

Table S2. Baseline characteristics of participants who underwent kidney transplant as the first modality of ESRD who were included versus excluded in the analysis

Variable	Exclude (n=32) ¹	Include (n=44) ²	P value ³
Age, year	52.42±12.29	46.90±12.33	0.0574
Men, N (%)	21 (65.63%)	23 (52.27%)	0.2444
Race, N (%)			
Non-Hispanic white	17 (53.13%)	24 (54.55%)	0.1794
Non-Hispanic black	5 (15.63%)	10 (22.73%)	.
Hispanic	5 (15.63%)	9 (20.45%)	.
Other	5 (15.63%)	1 (2.27%)	.
Current smoker, N (%)	4 (12.50%)	5 (11.36%)	0.8797
CRIC eGFR, ml/min/1.73m ²	32.80±9.18	30.62±7.14	0.2471
BMI, kg/m ²	28.37±5.46	28.64±7.87	0.8621
Systolic blood pressure, mmHg	132.08±24.42	121.86±16.96	0.0467
Diastolic blood pressure, mmHg	74.17±11.82	73.86±11.16	0.9098
LDL, mg/dl	93.78±37.26	94.50±35.62	0.9323
HDL, mg/dl	50.94±16.64	47.14±13.41	0.2741
Triglycerides, mg/dl	109.00 (79.50-177.00)	123.50 (82.50-195.50)	0.6187
Total cholesterol, mg/dl	181.53±56.11	178.75±41.06	0.8035
Hemoglobin, g/dl	12.03±1.31	12.00±1.52	0.9386
HbA1c hemoglobin, %	6.17±1.17	6.27±1.26	0.7301
Serum albumin, g/dl	3.80±0.45	3.86±0.47	0.6074
Hypertension, N (%)	28 (87.50%)	34 (77.27%)	0.2561
Cardiovascular disease, N (%)	7 (21.88%)	7 (15.91%)	0.5077
Chronic obstructive pulmonary disease, N (%)	0 (00.00%)	0 (00.00%)	.
Diabetes, N (%)	12 (37.50%)	18 (40.91%)	0.7640
β-blockers, N (%)	17 (53.13%)	12 (27.27%)	0.0220
Angiotensin converting enzyme inhibitors or angiotensin receptor blockers, N (%)	24 (75.00%)	35 (79.55%)	0.6387
Calcium channel blockers, N (%)	9 (28.13%)	13 (29.55%)	0.8928
Diuretics, N (%)	15 (46.88%)	22 (50.00%)	0.7878