

SUPPLEMENTARY TABLE S5. OR PER Gy AND CI FOR  
VARIOUS HISTOPATHOLOGIC AND INVASIVE PTC  
CHARACTERISTICS IN RADIOGENIC CASES

<i>Characteristics</i>	<i>OR per Gy</i>	<i>CI</i>	<i>p<sub>1</sub><sup>a</sup></i>	<i>p<sub>2</sub><sup>b</sup></i>
Tumor size ≤10 mm vs. >10 mm	1.20	1.06–1.39	0.004	NA
Dominant growth pattern <sup>c</sup>	1.18	0.98–1.63	0.103	0.081
Oxyphilic changes	1.09	0.88–1.26	0.341	0.423
Concomitant nodular pathology	1.24	1.08–1.45	0.003	0.052
Concomitant diffuse pathology	0.83	0.56–1.04	0.127	0.153
Intrathyroid spread	0.93	0.81–1.05	0.237	0.440
Extrathyroidal extension	0.87	0.74–0.99	0.031	0.106
Multifocality	1.04	0.87–1.18	0.581	0.773
Lymphatic/vascular invasion	0.87	0.76–0.98	0.220	0.083
Lymph node metastasis	0.93	0.83–1.04	0.197	0.553
Distant metastases	0.93	0.70–1.08	0.385	0.612

<sup>a</sup>Adjusted for age and sex.

<sup>b</sup>Additionally adjusted for tumor size.

<sup>c</sup>Follicular and solid-trabecular vs. papillary.

NA, not available.