

SUPPLEMENT TABLE 2, Participant Satisfaction, Confidence following Educational Intervention (n=16)

| Survey Statement | Response Score (%) | | | | | |
|--|--------------------|--------------|--------------|---------------|--------------|---------|
| | 1 | 2 | 3 | 4 | 5 | |
| 1) CERTAIN approach helped me feel better prepared during the emergency scenario | 0 (0%) | 1 (6.3%) | 0 (0%) | 8 (50%) | 7 (43.8%) | 4.3±0.8 |
| 2) CERTAIN software was easy to use | 0 (0%) | 0 (0%) | 2 (12.5%) | 10 (62.5%) | 4 (25%) | 4.1±0.6 |
| 3) I would use CERTAIN approach if I were treating decompensating patient in a real life | 0 (0%) | 0 (0%) | 5 (31.3%) | 8 (50%) | 3 (18.8%) | 3.9±0.7 |
| 4) CERTAIN did not disrupt clinical flow during simulated patient resuscitation | 0 (0%) | 1 (6.25%) | 5 (31.3%) | 8 (50%) | 2 (12.5%) | 3.7±0.8 |
| 5) If I were a patient, I would want CERTAIN to be used by my doctors | 0 (0%) | 1 (6.25%) | 2 (12.5%) | 7 (43.8%) | 6 (37.5%) | 4.1±0.9 |
| 6) The remote scenarios were realistic | 0 (0%) | 1 (6.25%) | 3 (18.8%) | 7 (43.8%) | 5 (31.2%) | 4.0±0.9 |
| 7) I enjoyed the remote training session. | 0 (0%) | 0 (0%) | 1 (6.25%) | 6 (37.5%) | 9 (56.3%) | 4.5±0.6 |
| 8) The knowledge gained will be helpful to me in my practice. | 0 (0%) | 0 (0%) | 2 (12.5%) | 7 (43.8%) | 7 (43.8%) | 4.3±0.7 |

| | | | | | | |
|--|------|---------|---------|---------|---------|---------|
| 9) I learned something new. | 0 | 0 | 2 | 3 | 11 | |
| | (0%) | (0%) | (12.5%) | (18.8%) | (68.8%) | 4.6±0.7 |
| 10) This training session should be taken by all ICU staff. | 0 | 1 | 2 | 3 | 10 | |
| | (0%) | (6.25%) | (12.5%) | (18.8%) | (62.5%) | 4.4±1.0 |
