Long-Term Survivors of Pancreatic Cancer A California Population-Based Study

SUPPLEMENTAL TABLE 1. Associations of Patient, Tumor, and Facility Characteristics With 5-Year Survival, Stratified by Surgical Status With Year of Diagnosis Dichotomized Before and After the Approval of Gemcitabine

| | OR (95% CI) Unresected | OR (95% CI) Surgically Resected |
|--|---------------------------|------------------------------------|
| Age (per year) | 0.97 (0.96-0.98) | 0.99 (0.98-1.00) |
| Sex | | |
| Female vs male | 0.97 (0.75-1.25) | 1.22 (1.04-1.43) |
| Race/Ethnicity | | |
| Asian/Pacific Islander vs Non-Hispanic White | 3.48 (2.51-4.82) | 0.97 (0.73-1.29) |
| Hispanic vs Non-Hispanic White | 1.82 (1.28-2.60) | 1.08 (0.84-1.38) |
| Non-Hispanic Black vs Non-Hispanic White | 1.02 (0.59-1.74) | 0.85 (0.60-1.21) |
| Degree of differentiation | | |
| Moderately vs poorly differentiated/anaplastic | 0.87 (0.55-1.37) | 1.66 (1.36-2.03) |
| Unknown differentiation vs poorly differentiated/anaplastic | 1.01 (0.71-1.44) | 2.10 (1.52-2.88) |
| Well-differentiated vs poorly differentiated/anaplastic | 2.53 (1.62-3.94) | 3.23 (2.51-4.16) |
| Anatomic location | | |
| Body of pancreas vs head of pancreas | 0.84 (0.54-1.33) | 0.94 (0.65-1.36) |
| Overlapping, other, NOS vs head of pancreas | 1.02 (0.75-1.38) | 1.05 (0.82-1.35) |
| Tail of pancreas vs head of pancreas | 0.82 (0.48-1.40) | 1.12 (0.81-1.55) |
| SEER summary stage | | |
| Localized vs distant | 3.73 (2.39-5.80) | 12.14 (7.65-19.29) |
| Regional vs distant | 1.91 (1.39-2.64) | 4.15 (2.66-6.48) |
| Unknown vs distant | 3.48 (2.34-5.18) | 6.56 (2.89-14.87) |
| Radiation | | |
| Yes vs no | 1.21 (0.86-1.71) | 0.97 (0.80-1.19) |
| Chemotherapy | | |
| Yes vs no | 1.06 (0.79-1.43) | 1.56 (1.27-1.91) |
| Quintile of neighborhood socioeconomic status | | |
| 2 vs 1 | 0.63 (0.40-0.97) | 1.16 (0.83-1.60) |
| 3 vs 1 | 0.75 (0.49-1.14) | 1.23 (0.90-1.69) |
| 4 vs 1 | 0.91 (0.60-1.37) | 1.41 (1.04-1.92) |
| 5 vs 1 | 0.96 (0.64-1.46) | 1.63 (1.20-2.20) |
| NCI cancer center | | |
| Yes vs no | 1.65 (1.20-2.27) | 1.38 (1.15-1.65) |
| Year of diagnosis | | |
| 1997-2009 vs 1988-1996 | 1.11 (0.84-1.48) | 1.68 (1.37-2.06) |
| Values in bold signify statistically significant association at $P < 0.05$. | | |