

Supplementary Information

Noninvasive assessment of characteristics of novel anti-HER2 antibodies by molecular imaging in a human gastric cancer xenograft-bearing mouse model

Wei-Ying Kuo¹, Jia-Jia Lin², Hung-Ju Hsu¹, Hong-Sen Chen¹, *An-Suei Yang¹, and *Chun-Yi Wu^{2,3}

¹Genomics Research Center, Academia Sinica, Taipei, Taiwan


²Department of Biomedical Imaging and Radiological Science, China Medical University, Taichung, Taiwan

³Master Program for Biomedical Engineering, China Medical University, Taichung, Taiwan

* Correspondence should be addressed to: Chun-Yi Wu, Department of Biomedical Imaging and Radiological Science, China Medical University, Taichung, Taiwan 40402. Phone: +886-4-2205-3366 ext.7808. email: cywu@mail.cmu.edu.tw

Supplementary Figure

a



b




Figure S1. (a) *In vivo* NIRF imaging (b) MicroSPECT/CT imaging of BT474 xenograft-bearing mice injected with 18.5 MBq (0.1 mg/mouse) of [^{111}In]labeled antibodies at 24 hr p.i..




Figure S2. MicroSPECT/CT imaging of N87 xenograft-bearing mice injected with 18.5 MBq (0.1 mg/mouse) of [^{123}I]labeled trastuzumab and 61 IgG at 12 and 36 hr p.i..