

eTable 1. Characteristics at celiac disease diagnosis in adults answering and not answering the study questionnaires.

	Questionnaires answered		P value
	Yes, n=237	No, n=322	
Current age, median (IQR), years	27.0 (22.0, 37.6)	25.9 (21.3, 34.7)	0.130
Age at diagnosis, median (IQR), years	9.7 (5.3, 13.6)	10.1 (6.4, 13.0)	0.529
Year of diagnosis, median (IQR)	1998 (1986, 2004)	1999 (1990, 2005)	0.075
Girls, No. (%)	164 (69.2)	167 (51.9)	<0.001
Main clinical presentation, No. (%)			0.259
Gastrointestinal	123 (52.1)	142 (45.1)	
Extra-intestinal	65 (27.5)	98 (31.1)	
Risk-group screening	48 (20.3)	75 (23.8)	
Symptoms ^a , No. (%)	172 (76.8)	226 (75.6)	0.750
In risk-group screened, No. (%)	21 (44.7)	43 (57.3)	0.173
Poor growth, No. (%)	97 (44.7)	115 (38.6)	0.164
Body mass index, median (IQR)	16.5 (15.2, 18.4) ^c	16.6 (15.4, 19.3) ^d	0.264
Anemia, No. (%)	63 (28.4)	63 (23.2)	0.194
Hemoglobin, median (IQR), g/l	124 (115, 131) ^e	127 (118, 134) ^f	0.051
Severity of villous atrophy, No. (%)			0.766
Partial	67 (31.5)	98 (33.6)	
Subtotal	84 (39.4)	106 (36.3)	
Total	62 (29.1)	88 (30.1)	
Type 1 diabetes, No. (%)	16 (8.8)	40 (15.9)	0.029
Celiac disease in the family ^b , No. (%)	75 (56.0) ^g	87 (44.2) ^h	0.035

^a Asymptomatic signs such as poor growth, anemia and other laboratory abnormalities excluded; ^b First degree relatives; ^{c-h} Data available only from ^c 160, ^d 223, ^e 190, ^f 223, ^g 134 and ^h 197 of patients. IQR, interquartile range.