

### Additional File 3. Treatment Outcome

	Clinic 1		Clinic 2		Clinic 3		Total	
	HIV	TB	HIV	TB	HIV	TB	HIV	TB
<b>Total TB cases on treatment for follow-up</b>	<b>122</b>		<b>67</b>		<b>22</b>		<b>211</b>	
<b>TB cases on Rx - identified by clinic (n, percentage of total TB cases)</b>	73 (59.8%)		35 (52.2%)		11 (50.0%)		119 (56.4%)	
<b>By provider (n, percentage of total TB cases by provider)</b>	24 (82.8%)	49 (52.7%)	12 (85.7%)	23 (43.4%)	7 (58.3%)	4 (40.0%)	43 (78.2%)	76 (48.7%)
<b>TB cases on treatment - transferred-in (n, percentage of total TB cases)</b>	49 (40.2%)		32 (47.8%)		11 (50.0%)		92 (43.6%)	
<b>By provider (n, percentage of total TB cases by provider)</b>	5 (17.2%)	44 (47.3%)	2 (14.3%)	30 (56.6%)	5 (41.7%)	6 (60.0%)	12 (21.8%)	80 (51.3%)
<b>By provider (n, percentage of total TB cases by clinic)</b>	29 (23.8%)	93 (76.2%)	14 (20.9%)	53 (79.1%)	12 (54.5%)	10 (45.5%)	55 (26.1%)	156 (73.9%)
	$X^2 = 10.522$ $p = 0.0052$							
<b>Outcome Intensive Phase<sup>1</sup></b>								
<b>Still on Intensive phase</b>	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (3.8%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (1.3%)
<b>Unknown (incomplete data)</b>	1 (3.4%)	4 (4.3%)	2 (14.2%)	2 (3.8%)	0 (0.0%)	0 (0.0%)	3 (5.5%)	6 (3.8%)
<b>Completed Intensive Phase</b>	23 (82.1%)	75 (84.3%)	10 (83.3%)	42 (85.7%)	5 (41.7%)	6 (60.0%)	38 (73.1%)	123 (83.1%)
	$X^2 = 2.4667$ $p = 0.1163$							
<b>Stopped</b>	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
<b>Defaulted</b>	2 (7.1%)	5 (5.6%)	0 (0.0%)	0 (0.0%)	6 (50.0%)	1 (10.0%)	8 (15.4%)	6 (4.1%)

	Clinic 1		Clinic 2		Clinic 3		Total	
	HIV	TB	HIV	TB	HIV	TB	HIV	TB
								p = 0.0104*
<b>Transferred out</b>	2 (7.1%)	4 (4.5%)	1 (8.3%)	0 (0.0%)	0 (0.0%)	1 (10.0%)	3 (5.8%)	5 (3.4%)
<b>Died</b>	1 (3.6%)	5 (5.6%)	1 (8.3%)	7 (14.3%)	1 (8.3%)	2 (20.0%)	3 (5.8%)	14 (9.5%)
								p = 0.5673*
								X <sup>2</sup> = 8.705 <b>p = 0.0335 (p = 0.0372)*</b>
<b>End-of-Intensive Phase test (completed IP only)</b>								
<b>&gt;=30 days and &lt;60 days of IP (n, percentage of cases completed IP)</b>	11 (47.8%)	48 (64.0%)	4 (40.0%)	19 (45.2%)	2 (40.0%)	1 (16.7%)	17 (44.7%)	68 (55.3%)
	X <sup>2</sup> = 1.9219 p = 0.1656		X <sup>2</sup> = 0.0898 p = 0.7644		X <sup>2</sup> = 0.7486 p = 0.3869 (p = 0.5455)*		X <sup>2</sup> = 1.296 p = 0.2549	
<b>Outcome Overall treatment<sup>2</sup></b>								
<b>Started TB Rx &gt;=180 days ago (n, percentage of total TB cases)</b>	93 (76.2%)		41 (61.2%)		14 (63.6%)		148 (70.1%)	
			X <sup>2</sup> = 5.1648 p = 0.0756					
<b>By provider (n, percentage of total TB cases by provider)</b>	22 (75.9%)	71 (76.3%)	3 (21.4%)	38 (71.7%)	9 (75.0%)	5 (50.0%)	34 (61.8%)	114 (73.1%)
	X <sup>2</sup> = 0.0028 p = 0.9575		X <sup>2</sup> = 11.785 <b>p = 0.0006 (p = 0.0009)*</b>		X <sup>2</sup> = 1.4732 p = 0.2248 (p = 0.3777)*		X <sup>2</sup> = 2.4612 p = 0.1167	
<b>Still on Continuation phase</b>	0 (0.0%)	0 (0.0%)	1 (33.3%)	9 (23.1%)	0 (0.0%)	0 (0.0%)	1 (1.5)	9 (7.9%)
<b>Unknown</b>	2 (9.1%)	3 (4.2%)	0 (0.0%)	1 (2.6%)	0 (0.0%)	0 (0.0%)	2 (2.9)	4 (3.5%)

	Clinic 1		Clinic 2		Clinic 3		Total	
	HIV	TB	HIV	TB	HIV	TB	HIV	TB
<b>Completed/cure</b>	14 (70.0%)	51 (75.0%)	0 (0.0%)	14 (50.0%)	2 (22.2%)	4 (80.0%)	16 (53.3%)	69 (68.3%)
<b>Stopped</b>	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
<b>Defaulted</b>	3 (15.0%)	9 (13.2%)	1 (50.0%)	1 (3.6%)	6 (66.6%)	0 (0.0%)	10 (33.3%)	10 (9.9%)
<b>Transferred out</b>	2 (10.0%)	4 (5.9%)	0 (0.0%)	6 (21.4%)	1 (11.1%)	0 (0.0%)	3 (10.0%)	10 (9.9%)
<b>Died</b>	1 (5.0%)	4 (5.9%)	1 (50.0%)	7 (25.0%)	0 (0.0%)	1 (20.0%)	2 (6.7%)	12 (11.9%)
	X <sup>2</sup> = 0.493 p = 0.9204 (p = 0.8640)*		X <sup>2</sup> = 7.902 p = 0.0481 (p = 0.0827)*		X <sup>2</sup> = 8.193 p = 0.0422 (p = 0.0210)*		X <sup>2</sup> = 9.513 p = 0.0232 (p = 0.0319)*	
<b>End-of-Continuation phase test (completed/cure only)</b>								
<b>&gt;=60 days of CP (n, percentage of completed/cure)</b>	11 (78.6%)	45 (88.2%)	-	4 (28.6%)	2 (100%)	3 (75.0%)	13 (81.3%)	52 (75.4%)
	p = 0.3915*		-	p = 1*		p = 0.7510*		

<sup>1</sup> Patients who started TB treatment at least 60 days before the end of the assessment; <sup>2</sup> Patients who started TB treatment at least 180 days before the end of the assessment; \* Fisher's Exact test (two tailed)