

Supplementary Material


Alterations in the abundance and co-occurrence of *Akkermansia muciniphila* and *Faecalibacterium prausnitzii* in the colonic mucosa of inflammatory bowel disease subjects

Mireia Lopez-Siles, Núria Enrich-Capó, Xavier Aldeguer, Miriam Sabat-Mir, Sylvia H. Duncan, L. Jesús Garcia-Gil*, Margarita Martinez-Medina


* Correspondence: L. Jesús Garcia-Gil, jesus.garcia@udg.edu

Supplementary Figure

A. UC location



B. CD location



Legend

Both species	Only <i>F. prausnitzii</i>	Only phylogroup I <i>A. muciniphila</i>	Only phylogroup II <i>A. muciniphila</i>	<i>A. muciniphila</i> and phylogroup II
Only <i>A. muciniphila</i>	None of the two species			

Figure S1. Prevalence of total *F. prausnitzii*, phylogroups and *A. muciniphila* (A) by disease subtype in patients with UC and (B) in patients with CD; UC, ulcerative colitis; CD, Crohn's disease; E1, ulcerative proctitis; E2, ulcerative left-sided colitis; E3, ulcerative pancolitis; IC-CD, ileocolonic CD, I-CD, ileal CD; C-CD, colonic CD.