

Table S1 Summary of ARIA events

	Low-dose SC cohort		High-dose IV cohort		All patients	
	(n = 39)		(n = 52)		(n = 91)	
	Placebo	Crenezumab	Placebo	Crenezumab	Placebo	Crenezumab
	(n = 13)	(n = 26)	(n = 16)	(n = 36)	(n = 29)	(n = 62)
Any ARIA-E, n (%)						
Symptomatic	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Asymptomatic	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Superficial siderosis, n (%)						
	0 (0.0)	0 (0.0)	1 (6.3)	0 (0.0)	1 (3.4)	0 (0.0)
Macrohemorrhage, n (%)						
	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
New ARIA-H, n (%)						
Symptomatic	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Asymptomatic	0 (0.0)	4 (15.4)	1 (6.3)	5 (13.9)	1 (3.4)	9 (14.5)
1 microhemorrhage	0 (0.0)	1 (3.8)	0 (0.0)	1 (2.8)	0 (0.0)	2 (3.2)
2–4 microhemorrhages	0 (0.0)	2 (7.7)	1 (6.3)	2 (5.6)	1 (3.4)	4 (6.5)
>4 microhemorrhages	0 (0.0)	1 (3.8)	0 (0.0)	2 (5.6)	0 (0.0)	3 (4.8)

ARIA-E amyloid-related imaging abnormalities: vasogenic edema or effusions; *ARIA-H* amyloid-related imaging abnormalities: microhemorrhage and siderosis; *IV* intravenous; *SC* subcutaneous