

THE LANCET

Supplementary appendix 2

This appendix formed part of the original submission and has been peer reviewed.
We post it as supplied by the authors.

Supplement to: GBD 2016 Alcohol Collaborators. Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2018; published online Aug 23. [http://dx.doi.org/10.1016/S0140-6736\(18\)31310-2](http://dx.doi.org/10.1016/S0140-6736(18)31310-2).

Supplemental Results Appendix for Alcohol use and burden: a systematic analysis from the Global Burden of Disease Study 2016 for 195 countries and territories, 1990–2016

Contents

Supplemental Results Table 1: Prevalence of current drinkers and abstainers, Population average of standard drinks daily, Percent of total attributable deaths, Percent of total attributable DALYs, Attributable deaths, and DALYs to alcohol use by location, year, sex, and age for GBD 2016	2
Supplemental Results Figure 1: Attributable DALY rate disaggregated by cause by location, sex, and age in 2016	3

Supplemental Results Table 1: Prevalence of current drinkers and abstainers, Population average of standard drinks daily, Percent of total attributable deaths, Percent of total attributable DALYs, Attributable deaths, and DALYs to alcohol use by location, year, sex, and age for GBD 2016

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	1990	Female	All Ages	26% (24-29%)	74% (71-76%)	0.76 (0.74-0.8)	1.3% (1.1-1.6%)	2% (1.4-2.8%)	450000 (300000-610000)	1500000 (1200000-1800000)
Global	1990	Female	15 to 19	21% (16-28%)	79% (72-84%)	0.56 (0.38-0.77)	1.1% (0.77-1.4%)	1.3% (0.84-2%)	4200 (2700-6300)	470000 (330000-640000)
Global	1990	Female	20 to 24	27% (20-34%)	73% (66-80%)	0.88 (0.64-1.2)	1.8% (1.4-2.4%)	2% (1.3-2.7%)	7700 (5200-11000)	890000 (650000-1200000)
Global	1990	Female	25 to 29	31% (23-38%)	69% (62-77%)	1 (0.77-1.3)	2.3% (1.8-2.8%)	2.7% (1.9-3.6%)	10000 (7100-13000)	1000000 (760000-1300000)
Global	1990	Female	30 to 34	34% (26-42%)	66% (58-74%)	1.1 (0.84-1.5)	2.6% (2-3.2%)	3.5% (2.6-4.4%)	13000 (9900-17000)	1100000 (850000-1300000)
Global	1990	Female	35 to 39	36% (28-43%)	64% (57-72%)	1.2 (0.9-1.5)	2.8% (2.3-3.5%)	4.2% (3.2-5.3%)	18000 (14000-23000)	1200000 (950000-1500000)
Global	1990	Female	40 to 44	38% (30-47%)	62% (53-70%)	1.3 (1-1.7)	3.2% (2.5-4%)	5% (3.8-6.2%)	22000 (17000-28000)	1200000 (980000-1500000)
Global	1990	Female	45 to 49	38% (28-45%)	62% (55-72%)	1.4 (1-1.8)	3.3% (2.5-4.3%)	5% (3.9-6.4%)	26000 (20000-34000)	1200000 (960000-1600000)
Global	1990	Female	50 to 54	39% (30-46%)	61% (54-70%)	1.4 (1.1-1.9)	3.5% (2.7-4.6%)	5.1% (3.8-6.5%)	37000 (28000-48000)	1500000 (1100000-1900000)
Global	1990	Female	55 to 59	38% (29-46%)	62% (54-71%)	1.5 (1.1-1.9)	3.4% (2.4-4.5%)	4.6% (3.4-6.1%)	44000 (32000-59000)	1500000 (1100000-2000000)
Global	1990	Female	60 to 64	40% (31-49%)	60% (51-69%)	1.5 (1.1-2.1)	3% (1.9-4.2%)	3.9% (2.5-5.5%)	50000 (32000-71000)	1500000 (940000-2000000)
Global	1990	Female	65 to 69	41% (32-49%)	59% (51-68%)	1.5 (1.1-1.9)	2.6% (1.6-3.6%)	3.3% (2.1-4.7%)	53000 (33000-75000)	1300000 (800000-1800000)
Global	1990	Female	70 to 74	38% (29-46%)	62% (54-71%)	1.2 (0.88-1.6)	1.9% (0.88-3%)	2.4% (1.1-3.7%)	44000 (20000-69000)	830000 (370000-1300000)
Global	1990	Female	75 to 79	39% (29-49%)	61% (51-71%)	1.3 (0.92-1.8)	1.7% (0.39-3.1%)	2% (0.42-3.6%)	43000 (9300-80000)	650000 (150000-1200000)
Global	1990	Female	80 to 84	40% (30-50%)	60% (50-70%)	1.4 (0.99-1.8)	1.6% (0.27-3%)	1.8% (0.26-3.5%)	37000 (5300-73000)	420000 (72000-800000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	1990	Female	85 to 89	41% (31-51%)	59% (49-69%)	1.4 (0.99-2)	1.4% (0.11-3%)	1.6% (-0.016-3.4%)	24000 (-250-52000)	200000 (15000-410000)
Global	1990	Female	90 to 94	44% (33-54%)	56% (46-67%)	1.5 (0.98-2.1)	1.3% (-0.35-3.2%)	1.3% (-0.62-3.7%)	9700 (-4500-27000)	57000 (-16000-150000)
Global	1990	Female	95 plus	47% (33-59%)	53% (41-67%)	1.5 (0.85-2.3)	1% (-1.1-3.5%)	1% (-1.5-4%)	2500 (-3700-10000)	10000 (-11000-35000)
Global	1990	Male	All Ages	39% (35-43%)	61% (57-65%)	1.4 (1.4-1.5)	4.5% (4-4.9%)	6.3% (5.4-7.2%)	1600000 (1300000-1800000)	5800000 (5200000-6400000)
Global	1990	Male	15 to 19	28% (21-36%)	72% (64-79%)	0.77 (0.54-1)	3.1% (2.3-4.2%)	4% (2.8-5.7%)	16000 (11000-23000)	1500000 (1000000-2000000)
Global	1990	Male	20 to 24	43% (33-52%)	57% (48-67%)	1.6 (1.2-2)	7.2% (5.6-8.9%)	8.6% (6.5-11%)	47000 (35000-60000)	3900000 (3100000-4900000)
Global	1990	Male	25 to 29	51% (40-60%)	49% (40-60%)	2.1 (1.6-2.6)	9% (7.3-11%)	11% (8.6-14%)	62000 (49000-76000)	4800000 (3800000-5700000)
Global	1990	Male	30 to 34	55% (44-64%)	45% (36-56%)	2.4 (1.9-3)	10% (8.3-12%)	13% (10-15%)	77000 (62000-91000)	5200000 (4300000-6100000)
Global	1990	Male	35 to 39	58% (48-66%)	42% (34-52%)	2.5 (1.9-3.2)	11% (8.8-13%)	14% (11-16%)	93000 (76000-110000)	5500000 (4500000-6500000)
Global	1990	Male	40 to 44	60% (50-68%)	40% (32-50%)	2.7 (2.1-3.4)	11% (9.1-13%)	14% (12-16%)	100000 (85000-120000)	5300000 (4400000-6200000)
Global	1990	Male	45 to 49	58% (48-67%)	42% (33-52%)	2.8 (2.1-3.6)	11% (9-13%)	14% (11-16%)	120000 (96000-140000)	5300000 (4400000-6200000)
Global	1990	Male	50 to 54	59% (49-67%)	41% (33-51%)	3 (2.3-3.8)	11% (8.9-13%)	13% (11-16%)	160000 (130000-190000)	6100000 (5000000-7200000)
Global	1990	Male	55 to 59	58% (48-67%)	42% (33-52%)	3 (2.3-3.7)	9.9% (8-12%)	12% (9.6-14%)	170000 (140000-210000)	5900000 (4800000-7000000)
Global	1990	Male	60 to 64	58% (48-67%)	42% (33-52%)	2.9 (2.3-3.7)	8.6% (6.7-11%)	9.9% (7.7-12%)	190000 (150000-240000)	5400000 (4200000-6700000)
Global	1990	Male	65 to 69	57% (47-66%)	43% (34-53%)	2.7 (2.2-3.3)	7.1% (5.5-8.9%)	8.2% (6.2-10%)	170000 (130000-210000)	4000000 (3100000-5000000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	1990	Male	70 to 74	54% (44-63%)	46% (37-56%)	2.1 (1.7-2.7)	5.6% (3.9-7.4%)	6.3% (4.4-8.4%)	130000 (94000-180000)	2600000 (1800000-3400000)
Global	1990	Male	75 to 79	54% (43-65%)	46% (35-57%)	2.3 (1.7-2.9)	4.7% (3.1-6.6%)	5.2% (3.3-7.4%)	110000 (70000-160000)	1600000 (1100000-2300000)
Global	1990	Male	80 to 84	54% (42-64%)	46% (36-58%)	2.3 (1.8-2.9)	4.1% (2.6-5.7%)	4.4% (2.8-6.3%)	69000 (43000-98000)	770000 (500000-1100000)
Global	1990	Male	85 to 89	53% (41-63%)	47% (37-59%)	2.1 (1.6-2.8)	3.3% (1.9-4.8%)	3.5% (1.9-5.3%)	32000 (18000-48000)	260000 (150000-390000)
Global	1990	Male	90 to 94	55% (43-65%)	45% (35-57%)	2 (1.4-2.7)	2.6% (1.1-4.2%)	2.7% (0.95-4.6%)	9200 (3300-16000)	53000 (22000-88000)
Global	1990	Male	95 plus	58% (44-69%)	42% (31-56%)	1.9 (1.2-2.7)	2% (0.19-3.9%)	1.9% (-0.18-4.1%)	1700 (-160-3800)	7100 (680-14000)
Global	1995	Female	All Ages	26% (24-28%)	74% (72-76%)	0.73 (0.7-0.75)	1.4% (1.1-1.7%)	2% (1.3-2.8%)	470000 (300000-650000)	1600000 (1300000-1900000)
Global	1995	Female	15 to 19	21% (16-27%)	79% (73-84%)	0.55 (0.38-0.76)	1.1% (0.8-1.5%)	1.4% (0.91-2.1%)	4300 (2800-6300)	48000 (33000-64000)
Global	1995	Female	20 to 24	27% (21-33%)	73% (67-79%)	0.85 (0.61-1.1)	1.8% (1.4-2.3%)	1.9% (1.3-2.7%)	7900 (5300-11000)	90000 (66000-120000)
Global	1995	Female	25 to 29	29% (23-36%)	71% (64-77%)	0.9 (0.68-1.2)	2.1% (1.6-2.5%)	2.4% (1.7-3.1%)	10000 (7500-14000)	100000 (78000-130000)
Global	1995	Female	30 to 34	33% (26-40%)	67% (60-74%)	1 (0.77-1.3)	2.4% (1.9-2.9%)	3.2% (2.4-4%)	14000 (11000-18000)	120000 (91000-140000)
Global	1995	Female	35 to 39	35% (29-42%)	65% (58-71%)	1.1 (0.88-1.5)	2.9% (2.3-3.5%)	4.3% (3.4-5.2%)	20000 (16000-25000)	130000 (110000-160000)
Global	1995	Female	40 to 44	36% (30-44%)	64% (56-70%)	1.2 (0.89-1.5)	3.2% (2.5-3.9%)	4.9% (3.9-6%)	27000 (22000-33000)	150000 (120000-180000)
Global	1995	Female	45 to 49	38% (31-45%)	62% (55-69%)	1.3 (0.99-1.7)	3.4% (2.7-4.3%)	5.2% (4.1-6.6%)	33000 (26000-41000)	150000 (120000-190000)
Global	1995	Female	50 to 54	37% (30-44%)	63% (56-70%)	1.3 (1-1.7)	3.4% (2.5-4.4%)	5% (3.7-6.3%)	37000 (28000-48000)	150000 (110000-190000)
Global	1995	Female	55 to 59	38% (31-45%)	62% (55-69%)	1.4 (1-1.9)	3.4% (2.5-4.6%)	4.8% (3.5-6.3%)	49000 (36000-65000)	170000 (120000-220000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	1995	Female	60 to 64	37% (30-45%)	63% (55-70%)	1.4 (1-1.8)	2.8% (1.8-4%)	3.7% (2.4-5.2%)	51000 (33000-72000)	1500000 (950000-2100000)
Global	1995	Female	65 to 69	39% (32-47%)	61% (53-68%)	1.3 (1-1.7)	2.5% (1.5-3.5%)	3.1% (1.9-4.5%)	56000 (34000-79000)	1300000 (800000-1900000)
Global	1995	Female	70 to 74	40% (32-48%)	60% (52-68%)	1.3 (0.93-1.7)	2% (0.85-3.2%)	2.4% (1-3.9%)	53000 (23000-84000)	1000000 (420000-1600000)
Global	1995	Female	75 to 79	37% (28-47%)	63% (53-72%)	1.1 (0.78-1.5)	1.4% (0.31-2.7%)	1.7% (0.32-3.2%)	37000 (7100-70000)	550000 (120000-1000000)
Global	1995	Female	80 to 84	39% (29-49%)	61% (51-71%)	1.3 (0.91-1.7)	1.3% (0.1-2.7%)	1.5% (0.066-3.1%)	34000 (1600-72000)	390000 (30000-790000)
Global	1995	Female	85 to 89	42% (32-52%)	58% (48-68%)	1.4 (0.94-1.9)	1.2% (-0.11-2.7%)	1.3% (-0.25-3%)	24000 (-4600-55000)	200000 (-180000-440000)
Global	1995	Female	90 to 94	45% (34-55%)	55% (45-66%)	1.4 (0.94-2)	1% (-0.57-2.9%)	1% (-0.85-3.3%)	9700 (-7900-30000)	58000 (-33000-170000)
Global	1995	Female	95 plus	49% (35-61%)	51% (39-65%)	1.5 (0.85-2.2)	0.84% (-1.2-3.3%)	0.78% (-1.6-3.7%)	2500 (-5300-12000)	11000 (-16000-42000)
Global	1995	Male	All Ages	39% (36-43%)	61% (57-64%)	1.5 (1.4-1.5)	5.2% (4.6-5.7%)	7% (6-8%)	1900000 (1600000-2100000)	6900000 (6200000-7700000)
Global	1995	Male	15 to 19	28% (21-35%)	72% (65-79%)	0.77 (0.54-1)	3.3% (2.4-4.3%)	4.3% (2.9-6%)	17000 (12000-24000)	1500000 (1100000-2000000)
Global	1995	Male	20 to 24	42% (33-50%)	58% (50-67%)	1.6 (1.2-2)	7.4% (5.8-9.2%)	8.9% (6.7-12%)	52000 (39000-67000)	4300000 (3300000-5400000)
Global	1995	Male	25 to 29	50% (41-59%)	50% (41-59%)	2 (1.5-2.5)	9.1% (7.3-11%)	11% (8.7-14%)	73000 (57000-90000)	5500000 (4300000-6600000)
Global	1995	Male	30 to 34	55% (45-63%)	45% (37-55%)	2.4 (1.9-3)	10% (8.5-12%)	13% (10-15%)	94000 (77000-110000)	6300000 (5200000-7300000)
Global	1995	Male	35 to 39	58% (49-65%)	42% (35-51%)	2.6 (2-3.3)	11% (9.4-13%)	14% (12-17%)	120000 (96000-140000)	6800000 (5600000-8000000)
Global	1995	Male	40 to 44	59% (51-67%)	41% (33-49%)	2.7 (2-3.3)	12% (10-14%)	15% (13-18%)	150000 (120000-170000)	7500000 (6400000-8600000)
Global	1995	Male	45 to 49	60% (51-68%)	40% (32-49%)	2.9 (2.2-3.8)	12% (10-14%)	15% (13-18%)	160000 (130000-190000)	7200000 (6000000-8400000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	1995	Male	50 to 54	59% (50-66%)	41% (34-50%)	2.9 (2.3-3.7)	11% (9.2-13%)	14% (11-16%)	160000 (130000-190000)	6400000 (5300000-7600000)
Global	1995	Male	55 to 59	59% (50-67%)	41% (33-50%)	3 (2.3-3.8)	11% (8.5-13%)	13% (10-15%)	210000 (170000-250000)	6900000 (5600000-8200000)
Global	1995	Male	60 to 64	57% (48-66%)	43% (34-52%)	2.8 (2.2-3.6)	8.9% (7-11%)	10% (8.1-13%)	210000 (170000-260000)	6100000 (4700000-7500000)
Global	1995	Male	65 to 69	57% (48-65%)	43% (35-52%)	2.6 (2.1-3.2)	7.4% (5.6-9.3%)	8.5% (6.5-11%)	200000 (160000-260000)	4800000 (3700000-6000000)
Global	1995	Male	70 to 74	55% (46-63%)	45% (37-54%)	2.3 (1.8-2.9)	5.9% (4.2-7.8%)	6.7% (4.8-8.9%)	160000 (120000-220000)	3100000 (2300000-4100000)
Global	1995	Male	75 to 79	52% (41-62%)	48% (38-59%)	1.9 (1.4-2.4)	4.6% (3.1-6.4%)	5.1% (3.3-7.2%)	110000 (73000-160000)	1700000 (1100000-2300000)
Global	1995	Male	80 to 84	53% (42-63%)	47% (37-58%)	2 (1.6-2.5)	4% (2.6-5.6%)	4.4% (2.8-6.2%)	80000 (51000-110000)	890000 (580000-1300000)
Global	1995	Male	85 to 89	54% (43-64%)	46% (36-57%)	2 (1.5-2.6)	3.3% (1.9-4.8%)	3.5% (1.9-5.3%)	39000 (21000-59000)	310000 (180000-470000)
Global	1995	Male	90 to 94	56% (44-66%)	44% (34-56%)	1.9 (1.4-2.6)	2.5% (1.4-4.2%)	2.6% (0.88-4.5%)	12000 (3900-20000)	67000 (27000-110000)
Global	1995	Male	95 plus	58% (45-70%)	42% (30-55%)	1.8 (1.2-2.6)	2% (0.19-4%)	2% (-0.11-4.3%)	2200 (-120-4800)	9000 (840-18000)
Global	2000	Female	All Ages	25% (23-27%)	75% (73-77%)	0.71 (0.68-0.74)	1.4% (1.2-1.7%)	2% (1.3-2.8%)	480000 (310000-660000)	17000000 (14000000-20000000)
Global	2000	Female	15 to 19	21% (16-27%)	79% (73-84%)	0.53 (0.37-0.74)	1.1% (0.82-1.5%)	1.5% (0.96-2.1%)	4600 (2900-6700)	510000 (360000-690000)
Global	2000	Female	20 to 24	27% (21-33%)	73% (67-79%)	0.87 (0.63-1.2)	1.9% (1.4-2.4%)	2.1% (1.4-2.8%)	8100 (5500-11000)	930000 (680000-1200000)
Global	2000	Female	25 to 29	28% (22-34%)	72% (66-78%)	0.89 (0.67-1.1)	2% (1.6-2.5%)	2.3% (1.7-3%)	11000 (7900-14000)	1100000 (820000-1300000)
Global	2000	Female	30 to 34	29% (23-36%)	71% (64-77%)	0.93 (0.69-1.2)	2.2% (1.7-2.6%)	2.8% (2.2-3.5%)	14000 (11000-18000)	1200000 (910000-1400000)
Global	2000	Female	35 to 39	32% (26-39%)	68% (61-74%)	1.1 (0.8-1.4)	2.6% (2.1-3.2%)	3.7% (3-4.6%)	20000 (16000-25000)	1400000 (1100000-1600000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2000	Female	40 to 44	34% (28-42%)	66% (58-72%)	1.2 (0.89-1.5)	3% (2.4-3.7%)	4.6% (3.7-5.6%)	28000 (22000-33000)	1500000 (1300000-1900000)
Global	2000	Female	45 to 49	35% (27-42%)	65% (58-73%)	1.2 (0.88-1.5)	3.2% (2.5-4%)	4.9% (3.9-6.2%)	36000 (28000-45000)	1700000 (1300000-2100000)
Global	2000	Female	50 to 54	36% (30-44%)	64% (56-70%)	1.3 (0.97-1.7)	3.4% (2.6-4.3%)	5% (3.8-6.4%)	42000 (32000-54000)	1700000 (1300000-2100000)
Global	2000	Female	55 to 59	35% (29-42%)	65% (58-71%)	1.3 (0.97-1.7)	3.2% (2.3-4.2%)	4.5% (3.3-5.9%)	46000 (34000-60000)	1600000 (1100000-2000000)
Global	2000	Female	60 to 64	36% (28-44%)	64% (56-72%)	1.3 (0.94-1.8)	2.8% (1.8-4%)	3.8% (2.4-5.3%)	54000 (35000-75000)	1500000 (1000000-2200000)
Global	2000	Female	65 to 69	35% (28-44%)	65% (56-72%)	1.2 (0.88-1.5)	2.3% (1.5-3.3%)	3% (1.9-4.2%)	56000 (36000-78000)	1300000 (830000-1800000)
Global	2000	Female	70 to 74	37% (29-45%)	63% (55-71%)	1.1 (0.81-1.5)	1.9% (0.83-3%)	2.3% (1-3.7%)	54000 (24000-86000)	1000000 (440000-1600000)
Global	2000	Female	75 to 79	37% (29-47%)	63% (53-71%)	1.1 (0.78-1.5)	1.4% (0.32-2.7%)	1.7% (0.36-3.2%)	43000 (9000-81000)	650000 (140000-1200000)
Global	2000	Female	80 to 84	37% (27-46%)	63% (54-73%)	1.1 (0.76-1.5)	1.1% (0.16-2.3%)	1.3% (0.16-2.7%)	30000 (3700-62000)	340000 (47000-680000)
Global	2000	Female	85 to 89	41% (30-51%)	59% (49-70%)	1.2 (0.86-1.7)	1% (-0.14-2.4%)	1.1% (-0.29-2.8%)	23000 (-5900-56000)	190000 (-26000-450000)
Global	2000	Female	90 to 94	46% (35-55%)	54% (45-65%)	1.4 (0.91-1.9)	0.9% (-0.63-2.8%)	0.93% (-0.91-3.2%)	10000 (-10000-36000)	64000 (-44000-200000)
Global	2000	Female	95 plus	50% (36-62%)	50% (38-64%)	1.4 (0.84-2.2)	0.74% (-1.3-3.2%)	0.67% (-1.7-3.5%)	2800 (-7000-14000)	12000 (-20000-52000)
Global	2000	Male	All Ages	40% (37-43%)	60% (57-63%)	1.5 (1.4-1.6)	5.5% (5-6.1%)	7.2% (6.2-8.2%)	2000000 (1700000-2300000)	75000000 (67000000-83000000)
Global	2000	Male	15 to 19	29% (23-36%)	71% (64-77%)	0.84 (0.59-1.1)	3.4% (2.5-4.5%)	4.6% (3.1-6.4%)	19000 (13000-26000)	1700000 (1200000-2300000)
Global	2000	Male	20 to 24	43% (34-50%)	57% (50-66%)	1.8 (1.3-2.4)	7.8% (6.2-9.7%)	9.5% (7.2-12%)	55000 (42000-71000)	4500000 (3500000-5600000)
Global	2000	Male	25 to 29	50% (41-58%)	50% (42-59%)	2.2 (1.7-2.7)	9.2% (7.4-11%)	11% (8.7-14%)	77000 (60000-94000)	5800000 (4600000-7000000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2000	Male	30 to 34	54% (44-62%)	46% (38-56%)	2.3 (1.8-3)	9.8% (8.2-12%)	12% (9.9-14%)	97000 (79000-120000)	650000 (540000-760000)
Global	2000	Male	35 to 39	57% (49-64%)	43% (36-51%)	2.5 (1.9-3.3)	11% (9-13%)	14% (11-16%)	120000 (100000-140000)	720000 (600000-840000)
Global	2000	Male	40 to 44	58% (51-66%)	42% (34-49%)	2.7 (2.1-3.4)	12% (9.8-14%)	15% (12-17%)	150000 (130000-180000)	790000 (670000-910000)
Global	2000	Male	45 to 49	59% (50-67%)	41% (33-50%)	2.8 (2.1-3.6)	12% (10-14%)	15% (13-18%)	190000 (160000-220000)	860000 (730000-1000000)
Global	2000	Male	50 to 54	60% (51-67%)	40% (33-49%)	2.9 (2.2-3.7)	12% (9.5-14%)	14% (12-17%)	190000 (160000-230000)	760000 (630000-900000)
Global	2000	Male	55 to 59	57% (49-65%)	43% (35-51%)	2.8 (2.2-3.6)	10% (8.2-12%)	12% (10-15%)	190000 (150000-220000)	640000 (520000-760000)
Global	2000	Male	60 to 64	57% (48-65%)	43% (35-52%)	2.7 (2.1-3.5)	8.9% (6.9-11%)	10% (8.1-13%)	220000 (170000-280000)	640000 (500000-790000)
Global	2000	Male	65 to 69	55% (46-64%)	45% (36-54%)	2.4 (1.9-2.9)	7.4% (5.7-9.3%)	8.6% (6.7-11%)	210000 (160000-260000)	500000 (390000-630000)
Global	2000	Male	70 to 74	55% (46-63%)	45% (37-54%)	2.2 (1.7-2.7)	5.9% (4.2-7.9%)	6.7% (4.7-9%)	190000 (130000-250000)	360000 (250000-470000)
Global	2000	Male	75 to 79	53% (43-63%)	47% (37-57%)	2 (1.5-2.5)	4.7% (3.1-6.5%)	5.2% (3.4-7.4%)	130000 (85000-180000)	200000 (130000-270000)
Global	2000	Male	80 to 84	51% (39-61%)	49% (39-61%)	1.7 (1.3-2.1)	3.8% (2.5-5.3%)	4.2% (2.6-5.9%)	78000 (49000-110000)	870000 (580000-1200000)
Global	2000	Male	85 to 89	53% (41-62%)	47% (38-59%)	1.8 (1.3-2.3)	3.1% (1.8-4.7%)	3.4% (1.9-5.2%)	44000 (24000-66000)	350000 (210000-530000)
Global	2000	Male	90 to 94	56% (45-66%)	44% (34-55%)	1.8 (1.3-2.4)	2.4% (0.93-4.1%)	2.5% (0.75-4.4%)	14000 (4100-24000)	80000 (30000-130000)
Global	2000	Male	95 plus	58% (45-70%)	42% (30-55%)	1.8 (1.1-2.5)	2% (0.11-4%)	1.9% (-0.2-4.3%)	2700 (-290-6000)	11000 (610-23000)
Global	2005	Female	All Ages	24% (23-27%)	76% (73-77%)	0.71 (0.69-0.74)	1.5% (1.3-1.8%)	2.1% (1.4-2.8%)	510000 (350000-670000)	1700000 (1400000-2000000)
Global	2005	Female	15 to 19	20% (16-25%)	80% (75-84%)	0.51 (0.36-0.7)	1.1% (0.8-1.5%)	1.5% (0.97-2.1%)	4400 (2900-6400)	510000 (360000-700000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2005	Female	20 to 24	26% (21-31%)	74% (69-79%)	0.87 (0.64-1.2)	1.9% (1.5-2.4%)	2.2% (1.6-3%)	8400 (5900-11000)	970000 (730000-1300000)
Global	2005	Female	25 to 29	28% (22-33%)	72% (67-78%)	0.94 (0.71-1.2)	2.1% (1.7-2.5%)	2.5% (1.8-3.1%)	11000 (8200-14000)	1100000 (840000-1400000)
Global	2005	Female	30 to 34	29% (23-34%)	71% (66-77%)	0.94 (0.71-1.2)	2.2% (1.8-2.6%)	2.8% (2.2-3.5%)	15000 (12000-18000)	1200000 (960000-1500000)
Global	2005	Female	35 to 39	30% (25-36%)	70% (64-75%)	0.98 (0.75-1.3)	2.4% (1.9-2.9%)	3.5% (2.8-4.3%)	20000 (16000-24000)	1300000 (1100000-1600000)
Global	2005	Female	40 to 44	33% (28-39%)	67% (61-72%)	1.1 (0.85-1.4)	2.8% (2.3-3.4%)	4.4% (3.6-5.3%)	28000 (23000-34000)	1600000 (1300000-1900000)
Global	2005	Female	45 to 49	34% (28-40%)	66% (60-72%)	1.2 (0.91-1.5)	3.2% (2.5-4%)	5% (4-6.2%)	38000 (30000-47000)	1800000 (1400000-2200000)
Global	2005	Female	50 to 54	34% (28-40%)	66% (60-72%)	1.2 (0.89-1.5)	3.3% (2.5-4.2%)	4.9% (3.8-6.2%)	48000 (37000-59000)	1900000 (1500000-2400000)
Global	2005	Female	55 to 59	35% (30-41%)	65% (59-70%)	1.3 (0.96-1.7)	3.2% (2.3-4.1%)	4.6% (3.4-6%)	52000 (39000-67000)	1800000 (1300000-2300000)
Global	2005	Female	60 to 64	34% (28-40%)	66% (60-72%)	1.2 (0.89-1.7)	2.6% (1.8-3.6%)	3.6% (2.5-4.9%)	49000 (34000-66000)	1400000 (950000-1900000)
Global	2005	Female	65 to 69	34% (27-41%)	66% (59-73%)	1.1 (0.86-1.5)	2.3% (1.5-3.2%)	3.1% (2-4.2%)	57000 (38000-78000)	1300000 (880000-1900000)
Global	2005	Female	70 to 74	33% (27-40%)	67% (60-73%)	1 (0.74-1.4)	1.8% (0.93-2.8%)	2.3% (1.2-3.5%)	53000 (28000-80000)	1000000 (510000-1500000)
Global	2005	Female	75 to 79	34% (27-42%)	66% (58-73%)	1 (0.7-1.4)	1.4% (0.41-2.5%)	1.7% (0.55-3.1%)	46000 (14000-81000)	690000 (200000-1200000)
Global	2005	Female	80 to 84	36% (27-45%)	64% (55-73%)	1.1 (0.78-1.5)	1.3% (0.32-2.4%)	1.5% (0.38-2.8%)	40000 (10000-76000)	440000 (110000-830000)
Global	2005	Female	85 to 89	37% (28-46%)	63% (54-72%)	1.1 (0.75-1.5)	1% (0.026-2.2%)	1.2% (-0.015-2.5%)	23000 (-310-51000)	190000 (4800-410000)
Global	2005	Female	90 to 94	43% (33-53%)	57% (47-67%)	1.3 (0.87-1.8)	0.93% (-0.46-2.6%)	0.99% (-0.67-3%)	13000 (-8700-39000)	76000 (-37000-210000)
Global	2005	Female	95 plus	49% (36-62%)	51% (38-64%)	1.4 (0.85-2.2)	0.86% (-1-3.2%)	0.83% (-1.4-3.6%)	4300 (-7300-18000)	18000 (-21000-66000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2005	Male	All Ages	41% (38-43%)	59% (57-62%)	1.6 (1.5-1.7)	5.9% (5.3-6.5%)	7.4% (6.5-8.4%)	2100000 (1800000-2400000)	79000000 (72000000-88000000)
Global	2005	Male	15 to 19	30% (25-37%)	70% (63-75%)	0.81 (0.58-1.1)	3.4% (2.5-4.4%)	4.6% (3.2-6.3%)	19000 (13000-26000)	1800000 (1300000-2300000)
Global	2005	Male	20 to 24	43% (37-49%)	57% (51-63%)	1.9 (1.4-2.4)	7.9% (6.3-9.7%)	9.8% (7.5-12%)	56000 (43000-71000)	4700000 (3700000-5800000)
Global	2005	Male	25 to 29	49% (43-56%)	51% (44-57%)	2.4 (1.8-3)	9.3% (7.6-11%)	11% (9.1-14%)	77000 (61000-93000)	5800000 (4700000-6900000)
Global	2005	Male	30 to 34	53% (46-60%)	47% (40-54%)	2.5 (1.9-3.1)	9.9% (8.3-12%)	12% (10-15%)	100000 (82000-120000)	6700000 (5600000-7800000)
Global	2005	Male	35 to 39	56% (50-62%)	44% (38-50%)	2.5 (1.8-3.2)	10% (8.7-12%)	13% (11-15%)	120000 (100000-150000)	7400000 (6200000-8600000)
Global	2005	Male	40 to 44	57% (51-64%)	43% (36-49%)	2.7 (2.1-3.4)	11% (9.7-13%)	15% (12-17%)	160000 (140000-190000)	8400000 (7200000-9600000)
Global	2005	Male	45 to 49	58% (52-65%)	42% (35-48%)	2.9 (2.2-3.7)	12% (10-14%)	15% (13-18%)	200000 (170000-230000)	9100000 (7800000-10000000)
Global	2005	Male	50 to 54	59% (52-65%)	41% (35-48%)	2.8 (2.1-3.5)	12% (9.9-14%)	15% (12-17%)	240000 (200000-280000)	9400000 (7900000-11000000)
Global	2005	Male	55 to 59	58% (52-65%)	42% (35-48%)	2.8 (2.2-3.6)	11% (8.6-12%)	13% (11-15%)	220000 (190000-270000)	7700000 (6400000-9000000)
Global	2005	Male	60 to 64	56% (49-63%)	44% (37-51%)	2.6 (2-3.4)	8.6% (6.7-11%)	10% (8-13%)	200000 (160000-240000)	5800000 (4500000-7100000)
Global	2005	Male	65 to 69	55% (48-62%)	45% (38-52%)	2.4 (1.9-2.9)	7.4% (5.7-9.2%)	8.7% (6.6-11%)	220000 (170000-270000)	5300000 (4100000-6500000)
Global	2005	Male	70 to 74	53% (45-60%)	47% (40-55%)	2.1 (1.6-2.6)	5.8% (4.1-7.7%)	6.7% (4.6-8.8%)	190000 (130000-250000)	3600000 (2600000-4800000)
Global	2005	Male	75 to 79	52% (44-61%)	48% (39-56%)	1.9 (1.5-2.4)	4.6% (2.9-6.4%)	5.1% (3.2-7.3%)	140000 (91000-200000)	2200000 (1400000-3000000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2005	Male	80 to 84	51% (40-61%)	49% (39-60%)	1.8 (1.4-2.2)	3.8% (2.5-5.3%)	4.2% (2.7-6%)	94000 (61000-130000)	1100000 (710000-1500000)
Global	2005	Male	85 to 89	50% (40-60%)	50% (40-60%)	1.6 (1.2-2)	2.9% (1.7-4.2%)	3.1% (1.7-4.7%)	42000 (23000-64000)	350000 (200000-520000)
Global	2005	Male	90 to 94	55% (44-64%)	45% (36-56%)	1.7 (1.3-2.3)	2.4% (0.87-3.9%)	2.4% (0.71-4.3%)	16000 (4700-28000)	93000 (35000-160000)
Global	2005	Male	95 plus	58% (46-70%)	42% (30-54%)	1.8 (1.1-2.5)	2% (0.12-4.2%)	1.9% (-0.22-4.3%)	3400 (-390-7700)	14000 (830-29000)
Global	2010	Female	All Ages	24% (23-26%)	76% (74-77%)	0.71 (0.68-0.74)	1.5% (1.3-1.8%)	2.1% (1.5-2.8%)	520000 (360000-680000)	17000000 (14000000-20000000)
Global	2010	Female	15 to 19	19% (15-24%)	81% (76-85%)	0.49 (0.34-0.7)	1% (0.75-1.4%)	1.4% (0.86-2%)	3600 (2300-5400)	450000 (320000-610000)
Global	2010	Female	20 to 24	25% (20-31%)	75% (69-80%)	0.82 (0.59-1.1)	1.9% (1.5-2.4%)	2.2% (1.5-3%)	7600 (5300-10000)	960000 (710000-1300000)
Global	2010	Female	25 to 29	27% (21-33%)	73% (67-79%)	0.93 (0.7-1.2)	2.1% (1.7-2.5%)	2.5% (1.8-3.2%)	10000 (7500-13000)	1100000 (830000-1400000)
Global	2010	Female	30 to 34	28% (22-34%)	72% (66-78%)	0.98 (0.73-1.3)	2.1% (1.7-2.6%)	2.8% (2.2-3.5%)	13000 (10000-16000)	1100000 (900000-1400000)
Global	2010	Female	35 to 39	30% (24-36%)	70% (64-76%)	0.98 (0.75-1.3)	2.3% (1.9-2.8%)	3.4% (2.7-4.2%)	18000 (14000-22000)	1300000 (1000000-1500000)
Global	2010	Female	40 to 44	31% (26-37%)	69% (63-74%)	1 (0.76-1.3)	2.5% (2-3%)	4% (3.2-4.9%)	24000 (20000-30000)	1400000 (1200000-1700000)
Global	2010	Female	45 to 49	33% (26-39%)	67% (61-74%)	1.1 (0.85-1.4)	2.9% (2.3-3.6%)	4.7% (3.7-5.9%)	35000 (28000-43000)	1700000 (1400000-2100000)
Global	2010	Female	50 to 54	33% (28-40%)	67% (60-72%)	1.2 (0.9-1.5)	3.2% (2.4-4%)	4.9% (3.8-6.1%)	46000 (36000-58000)	1900000 (1500000-2400000)
Global	2010	Female	55 to 59	33% (27-39%)	67% (61-73%)	1.2 (0.88-1.6)	3.1% (2.3-4%)	4.7% (3.5-5.9%)	56000 (42000-71000)	1900000 (1500000-2500000)
Global	2010	Female	60 to 64	34% (28-41%)	66% (59-72%)	1.2 (0.89-1.6)	2.7% (1.8-3.6%)	3.8% (2.6-5%)	55000 (38000-73000)	1600000 (1100000-2100000)
Global	2010	Female	65 to 69	34% (28-40%)	66% (60-72%)	1.1 (0.85-1.4)	2.2% (1.5-3%)	3.1% (2.1-4.1%)	52000 (36000-70000)	1200000 (830000-1700000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2010	Female	70 to 74	34% (27-41%)	66% (59-73%)	1 (0.75-1.4)	1.9% (0.97-2.8%)	2.4% (1.3-3.6%)	56000 (31000-84000)	1100000 (570000-1600000)
Global	2010	Female	75 to 79	33% (25-41%)	67% (59-75%)	0.95 (0.66-1.3)	1.4% (0.54-2.5%)	1.8% (0.71-3%)	47000 (18000-79000)	710000 (270000-1200000)
Global	2010	Female	80 to 84	35% (26-43%)	65% (57-74%)	1 (0.72-1.4)	1.3% (0.38-2.3%)	1.5% (0.48-2.8%)	44000 (14000-82000)	490000 (150000-910000)
Global	2010	Female	85 to 89	38% (29-48%)	62% (52-71%)	1.1 (0.77-1.5)	1.1% (0.16-2.3%)	1.3% (0.15-2.6%)	32000 (3700-65000)	260000 (36000-520000)
Global	2010	Female	90 to 94	42% (32-51%)	58% (49-68%)	1.2 (0.77-1.6)	0.85% (-0.31-2.3%)	0.93% (-0.46-2.7%)	13000 (-6300-36000)	74000 (-26000-200000)
Global	2010	Female	95 plus	49% (36-61%)	51% (39-64%)	1.4 (0.79-2.1)	0.84% (-0.93-3.1%)	0.83% (-1.3-3.6%)	5300 (-8400-23000)	22000 (-23000-82000)
Global	2010	Male	All Ages	40% (38-43%)	60% (57-62%)	1.6 (1.5-1.8)	5.9% (5.3-6.5%)	7.4% (6.4-8.4%)	2100000 (1900000-2400000)	7900000 (7100000-8800000)
Global	2010	Male	15 to 19	28% (22-34%)	72% (66-78%)	0.82 (0.56-1.1)	3.1% (2.3-4.1%)	4.1% (2.7-5.7%)	16000 (11000-23000)	150000 (110000-210000)
Global	2010	Male	20 to 24	42% (36-49%)	58% (51-64%)	1.8 (1.3-2.4)	7.5% (5.9-9.2%)	9.4% (7-12%)	54000 (40000-68000)	460000 (360000-570000)
Global	2010	Male	25 to 29	48% (41-54%)	52% (46-59%)	2.4 (1.8-3.1)	9.1% (7.4-11%)	11% (8.9-14%)	74000 (59000-90000)	570000 (460000-680000)
Global	2010	Male	30 to 34	51% (44-58%)	49% (42-56%)	2.6 (2-3.4)	9.6% (8.1-11%)	12% (9.9-14%)	91000 (75000-110000)	620000 (520000-720000)
Global	2010	Male	35 to 39	54% (48-61%)	46% (39-52%)	2.5 (1.9-3.3)	10% (8.4-12%)	13% (11-15%)	120000 (96000-140000)	700000 (590000-810000)
Global	2010	Male	40 to 44	56% (49-63%)	44% (37-51%)	2.5 (1.9-3.2)	11% (8.9-12%)	14% (12-16%)	150000 (120000-170000)	780000 (660000-900000)
Global	2010	Male	45 to 49	57% (50-63%)	43% (37-50%)	2.8 (2.1-3.5)	11% (9.5-13%)	15% (12-17%)	190000 (160000-220000)	870000 (740000-1000000)
Global	2010	Male	50 to 54	57% (50-63%)	43% (37-50%)	2.9 (2.2-3.6)	11% (9.3-13%)	14% (12-16%)	220000 (190000-260000)	900000 (760000-1000000)
Global	2010	Male	55 to 59	56% (49-63%)	44% (37-51%)	2.7 (2.1-3.5)	10% (8.5-12%)	13% (11-15%)	250000 (210000-300000)	870000 (730000-1000000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2010	Male	60 to 64	56% (49-64%)	44% (36-51%)	2.7 (2-3.4)	8.7% (6.9-11%)	11% (8.3-13%)	230000 (180000-280000)	6700000 (5400000-8100000)
Global	2010	Male	65 to 69	54% (46-61%)	46% (39-54%)	2.4 (1.9-2.9)	7.1% (5.6-8.7%)	8.4% (6.6-10%)	200000 (160000-240000)	4800000 (3800000-5900000)
Global	2010	Male	70 to 74	53% (45-60%)	47% (40-55%)	2.1 (1.6-2.7)	5.8% (4.2-7.7%)	6.8% (4.7-8.9%)	200000 (140000-260000)	3800000 (2700000-5000000)
Global	2010	Male	75 to 79	51% (43-60%)	49% (40-57%)	1.9 (1.4-2.4)	4.7% (3.1-6.4%)	5.3% (3.5-7.4%)	150000 (100000-210000)	2300000 (1600000-3200000)
Global	2010	Male	80 to 84	52% (41-61%)	48% (39-59%)	1.8 (1.4-2.2)	3.9% (2.6-5.4%)	4.3% (2.8-6.2%)	110000 (73000-160000)	1300000 (860000-1800000)
Global	2010	Male	85 to 89	53% (43-62%)	47% (38-57%)	1.8 (1.3-2.3)	3.1% (1.9-4.5%)	3.4% (1.9-5.1%)	58000 (32000-86000)	470000 (280000-690000)
Global	2010	Male	90 to 94	54% (44-63%)	46% (37-56%)	1.6 (1.2-2.2)	2.3% (0.92-3.8%)	2.4% (0.79-4.1%)	18000 (5800-31000)	100000 (42000-170000)
Global	2010	Male	95 plus	59% (47-70%)	41% (30-53%)	1.7 (1.1-2.5)	2.1% (0.25-4.1%)	2% (-0.076-4.4%)	4600 (-170-9900)	19000 (2300-38000)
Global	2016	Female	All Ages	25% (23-27%)	75% (73-77%)	0.74 (0.69-0.8)	1.7% (1.4-2%)	2.3% (1.6-3%)	560000 (390000-750000)	18000000 (15000000-22000000)
Global	2016	Female	15 to 19	19% (14-25%)	81% (75-86%)	0.52 (0.34-0.75)	1% (0.74-1.4%)	1.4% (0.84-2.1%)	3100 (1900-4700)	41000 (290000-560000)
Global	2016	Female	20 to 24	25% (19-32%)	75% (68-81%)	0.87 (0.62-1.2)	1.9% (1.5-2.5%)	2.3% (1.6-3.1%)	6400 (4400-8900)	870000 (640000-1200000)
Global	2016	Female	25 to 29	27% (21-34%)	73% (66-79%)	0.92 (0.67-1.2)	2.2% (1.7-2.7%)	2.8% (2-3.7%)	9700 (7000-13000)	1100000 (830000-1400000)
Global	2016	Female	30 to 34	29% (21-36%)	71% (64-79%)	0.99 (0.73-1.3)	2.3% (1.9-2.9%)	3.4% (2.6-4.4%)	14000 (10000-17000)	1200000 (960000-1500000)
Global	2016	Female	35 to 39	31% (23-38%)	69% (62-77%)	1.1 (0.78-1.4)	2.5% (2-3.2%)	4.2% (3.2-5.3%)	18000 (14000-24000)	1300000 (1000000-1600000)
Global	2016	Female	40 to 44	31% (24-39%)	69% (61-76%)	1.1 (0.78-1.4)	2.6% (2.1-3.3%)	4.7% (3.6-5.8%)	25000 (19000-31000)	1500000 (1200000-1800000)
Global	2016	Female	45 to 49	31% (23-38%)	69% (62-77%)	1.1 (0.76-1.4)	2.7% (2.1-3.5%)	4.8% (3.7-6%)	34000 (27000-43000)	1700000 (1300000-2100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2016	Female	50 to 54	32% (25-39%)	68% (61-75%)	1.1 (0.84-1.5)	3% (2.3-3.9%)	5% (3.8-6.2%)	48000 (37000-60000)	2000000 (1500000-2500000)
Global	2016	Female	55 to 59	33% (26-41%)	67% (59-74%)	1.2 (0.9-1.6)	3.1% (2.3-4%)	4.8% (3.6-6.2%)	58000 (44000-75000)	2000000 (1500000-2600000)
Global	2016	Female	60 to 64	33% (26-41%)	67% (59-74%)	1.2 (0.85-1.6)	2.7% (1.9-3.6%)	3.9% (2.8-5.3%)	65000 (46000-86000)	1900000 (1300000-2500000)
Global	2016	Female	65 to 69	35% (28-43%)	65% (57-72%)	1.1 (0.84-1.5)	2.3% (1.5-3.1%)	3.2% (2.3-4.3%)	63000 (43000-83000)	1500000 (1000000-2000000)
Global	2016	Female	70 to 74	34% (27-42%)	66% (58-73%)	1 (0.74-1.4)	1.8% (1-2.8%)	2.5% (1.4-3.7%)	54000 (31000-80000)	1000000 (580000-1500000)
Global	2016	Female	75 to 79	34% (26-44%)	66% (56-74%)	1 (0.68-1.4)	1.5% (0.55-2.6%)	1.9% (0.75-3.3%)	52000 (20000-89000)	780000 (290000-1300000)
Global	2016	Female	80 to 84	35% (26-45%)	65% (55-74%)	1 (0.71-1.4)	1.3% (0.44-2.3%)	1.6% (0.55-2.9%)	46000 (16000-84000)	510000 (180000-930000)
Global	2016	Female	85 to 89	38% (28-47%)	62% (53-72%)	1.1 (0.73-1.5)	1.1% (0.17-2.2%)	1.3% (0.15-2.7%)	37000 (4200-79000)	300000 (47000-620000)
Global	2016	Female	90 to 94	44% (33-53%)	56% (47-67%)	1.3 (0.81-1.8)	0.95% (-0.37-2.5%)	1% (-0.49-2.9%)	20000 (-9300-56000)	120000 (-45000-310000)
Global	2016	Female	95 plus	48% (35-61%)	52% (39-65%)	1.3 (0.72-1.9)	0.78% (-0.81-2.8%)	0.77% (-1.2-3.2%)	5900 (-9000-24000)	25000 (-26000-89000)
Global	2016	Male	All Ages	39% (36-43%)	61% (57-64%)	1.7 (1.5-1.9)	6.2% (5.6-6.9%)	7.5% (6.5-8.6%)	2300000 (1900000-2600000)	8100000 (7200000-9100000)
Global	2016	Male	15 to 19	25% (19-32%)	75% (68-81%)	0.79 (0.53-1.1)	2.9% (2.1-3.9%)	3.8% (2.6-5.4%)	14000 (9200-19000)	1400000 (990000-1800000)
Global	2016	Male	20 to 24	38% (30-46%)	62% (54-70%)	1.7 (1.3-2.3)	7% (5.5-8.8%)	8.7% (6.5-11%)	46000 (34000-59000)	4000000 (3100000-5100000)
Global	2016	Male	25 to 29	45% (36-54%)	55% (46-64%)	2.2 (1.6-2.9)	8.8% (7.2-11%)	11% (8.8-14%)	72000 (56000-89000)	5600000 (4600000-6800000)
Global	2016	Male	30 to 34	48% (38-57%)	52% (43-62%)	2.6 (1.9-3.5)	9.6% (8-11%)	13% (10-15%)	90000 (74000-110000)	6300000 (5200000-7400000)
Global	2016	Male	35 to 39	51% (42-59%)	49% (41-58%)	2.7 (2-3.7)	9.9% (8.3-12%)	13% (11-16%)	110000 (88000-130000)	6600000 (5500000-7700000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Global	2016	Male	40 to 44	53% (44-62%)	47% (38-56%)	2.7 (1.9-3.5)	10% (8.7-12%)	14% (12-16%)	140000 (110000-160000)	740000 (620000-860000)
Global	2016	Male	45 to 49	55% (45-64%)	45% (36-55%)	2.6 (1.9-3.5)	11% (9.1-13%)	14% (12-17%)	180000 (150000-210000)	850000 (720000-990000)
Global	2016	Male	50 to 54	56% (46-64%)	44% (36-54%)	2.7 (2-3.6)	11% (9.1-13%)	14% (12-16%)	230000 (190000-270000)	920000 (770000-1100000)
Global	2016	Male	55 to 59	55% (47-64%)	45% (36-53%)	2.8 (2.1-3.6)	10% (8.3-12%)	13% (10-15%)	250000 (210000-300000)	880000 (740000-1000000)
Global	2016	Male	60 to 64	55% (46-65%)	45% (35-54%)	2.6 (1.9-3.4)	8.9% (7.1-11%)	11% (8.7-13%)	280000 (230000-340000)	830000 (670000-1000000)
Global	2016	Male	65 to 69	55% (46-64%)	45% (36-54%)	2.4 (1.8-3)	7.4% (5.9-9.1%)	8.9% (7.1-11%)	250000 (200000-310000)	610000 (490000-740000)
Global	2016	Male	70 to 74	53% (43-61%)	47% (39-57%)	2.1 (1.6-2.7)	5.7% (4.2-7.5%)	6.7% (4.8-8.7%)	190000 (140000-250000)	380000 (280000-490000)
Global	2016	Male	75 to 79	51% (41-62%)	49% (38-59%)	1.9 (1.4-2.5)	4.6% (3.1-6.4%)	5.3% (3.4-7.4%)	170000 (110000-230000)	250000 (170000-350000)
Global	2016	Male	80 to 84	51% (40-61%)	49% (39-60%)	1.7 (1.3-2.2)	3.9% (2.6-5.3%)	4.4% (2.8-6.1%)	130000 (83000-180000)	140000 (97000-200000)
Global	2016	Male	85 to 89	52% (41-62%)	48% (38-59%)	1.7 (1.2-2.2)	3.1% (1.9-4.5%)	3.4% (1.9-5%)	76000 (43000-110000)	620000 (370000-900000)
Global	2016	Male	90 to 94	56% (45-66%)	44% (34-55%)	1.8 (1.3-2.4)	2.5% (1-4%)	2.6% (0.91-4.4%)	28000 (9700-47000)	160000 (68000-260000)
Global	2016	Male	95 plus	58% (45-69%)	42% (31-55%)	1.6 (0.98-2.3)	2% (0.29-3.8%)	1.9% (-0.027-4.1%)	5500 (-77-12000)	23000 (3400-45000)
High SDI	1990	Female	All Ages	73% (69-76%)	27% (24-31%)	2.3 (2-2.7)	3.3% (2-4.8%)	3.2% (0.62-6.1%)	140000 (33000-260000)	410000 (250000-590000)
High SDI	1990	Female	15 to 19	69% (59-78%)	31% (22-41%)	2 (1.4-2.8)	3% (1.9-4.4%)	9.3% (5-14%)	1100 (580-1600)	110000 (71000-160000)
High SDI	1990	Female	20 to 24	80% (72-86%)	20% (14-28%)	3.1 (2.2-4.1)	4.6% (3.3-6.1%)	12% (7.4-17%)	1800 (1100-2500)	210000 (150000-280000)
High SDI	1990	Female	25 to 29	83% (76-88%)	17% (12-24%)	2.9 (2.2-3.8)	4.2% (3-5.4%)	11% (7-15%)	2000 (1300-2800)	220000 (150000-290000)
High SDI	1990	Female	30 to 34	84% (76-89%)	16% (11-24%)	2.9 (2.1-3.7)	4% (2.9-5.3%)	11% (7.4-14%)	2600 (1800-3500)	230000 (160000-290000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	1990	Female	35 to 39	84% (77-89%)	16% (11-23%)	2.9 (2.2-3.9)	4.4% (3.2-5.8%)	12% (8.5-15%)	3800 (2800-4900)	260000 (190000-340000)
High SDI	1990	Female	40 to 44	83% (77-89%)	17% (11-23%)	3.1 (2.3-4)	4.9% (3.5-6.4%)	12% (8.9-15%)	5600 (4100-7100)	320000 (230000-410000)
High SDI	1990	Female	45 to 49	82% (75-88%)	18% (12-25%)	3.2 (2.3-4.3)	5.3% (3.7-7.2%)	12% (8.4-15%)	7000 (5100-9100)	330000 (240000-440000)
High SDI	1990	Female	50 to 54	81% (74-86%)	19% (14-26%)	3.3 (2.4-4.4)	5.6% (3.7-7.7%)	11% (7.6-14%)	9200 (6500-12000)	380000 (260000-510000)
High SDI	1990	Female	55 to 59	79% (71-85%)	21% (15-29%)	3.3 (2.4-4.4)	5.3% (3.3-7.5%)	9.2% (6.1-13%)	11000 (7400-16000)	400000 (250000-560000)
High SDI	1990	Female	60 to 64	76% (68-82%)	24% (18-32%)	3.1 (2.2-4.3)	4.5% (2.3-7.1%)	7.2% (3.9-11%)	13000 (7000-21000)	400000 (200000-630000)
High SDI	1990	Female	65 to 69	72% (63-80%)	28% (20-37%)	2.9 (2.1-3.8)	3.8% (1.6-6.2%)	5.5% (2.4-8.8%)	16000 (6200-26000)	390000 (150000-640000)
High SDI	1990	Female	70 to 74	68% (58-76%)	32% (24-42%)	2.4 (1.6-3.3)	2.5% (0.14-5.2%)	3.4% (0.23-6.9%)	12000 (140-25000)	240000 (5700-500000)
High SDI	1990	Female	75 to 79	66% (56-76%)	34% (24-44%)	2.4 (1.6-3.3)	2.3% (-0.54-5.5%)	2.8% (-0.79-6.9%)	17000 (-5300-41000)	260000 (-65000-630000)
High SDI	1990	Female	80 to 84	64% (52-73%)	36% (27-48%)	2.3 (1.6-3.2)	2.1% (-0.68-5.2%)	2.4% (-1.1-6.3%)	18000 (-7700-47000)	210000 (-66000-530000)
High SDI	1990	Female	85 to 89	61% (50-71%)	39% (29-50%)	2.2 (1.5-3.1)	1.7% (-1.1-4.9%)	1.8% (-1.6-5.8%)	13000 (-9700-39000)	110000 (-61000-310000)
High SDI	1990	Female	90 to 94	60% (49-70%)	40% (30-51%)	2.1 (1.3-3.1)	1.3% (-1.5-4.7%)	1.3% (-2.2-5.4%)	5300 (-7600-21000)	33000 (-34000-110000)
High SDI	1990	Female	95 plus	59% (44-72%)	41% (28-56%)	1.9 (1.1-3)	0.73% (-2.3-4.3%)	0.57% (-3.2-4.9%)	1100 (-4300-7500)	5400 (-13000-27000)
High SDI	1990	Male	All Ages	82% (79-85%)	18% (15-21%)	3.6 (3.1-4.1)	8.4% (6.5-10%)	8.2% (5.5-11%)	350000 (230000-460000)	12000000 (9200000-15000000)
High SDI	1990	Male	15 to 19	73% (64-81%)	27% (19-36%)	2.4 (1.7-3.2)	7.2% (4.7-10%)	13% (7.3-18%)	4000 (2400-5800)	350000 (230000-480000)
High SDI	1990	Male	20 to 24	86% (78-91%)	14% (9.3-22%)	4.5 (3.5-5.8)	14% (10-18%)	21% (14-28%)	9900 (6900-13000)	840000 (620000-1100000)
High SDI	1990	Male	25 to 29	89% (83-93%)	11% (7.3-17%)	4.7 (3.6-5.9)	13% (9.7-16%)	20% (14-26%)	10000 (7400-14000)	850000 (640000-1100000)
High SDI	1990	Male	30 to 34	91% (86-94%)	9.2% (5.6-14%)	4.6 (3.6-5.9)	12% (8.8-15%)	18% (13-23%)	11000 (8200-14000)	820000 (630000-1000000)
High SDI	1990	Male	35 to 39	92% (87-95%)	8.3% (5.1-13%)	4.4 (3.4-5.7)	11% (8.5-14%)	17% (12-21%)	13000 (9700-16000)	830000 (640000-1000000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	1990	Male	40 to 44	92% (88-95%)	7.7% (4.7-12%)	4.6 (3.5-5.8)	11% (8.4-14%)	16% (12-20%)	16000 (12000-21000)	910000 (700000-1100000)
High SDI	1990	Male	45 to 49	92% (88-95%)	7.8% (4.8-12%)	5 (3.8-6.5)	12% (8.8-15%)	16% (12-20%)	21000 (15000-26000)	970000 (740000-1200000)
High SDI	1990	Male	50 to 54	92% (88-95%)	8.1% (5.1-12%)	5.5 (4.2-7)	12% (8.8-15%)	15% (11-19%)	28000 (21000-36000)	1100000 (870000-1400000)
High SDI	1990	Male	55 to 59	91% (86-94%)	9.3% (6-14%)	5.7 (4.4-7.2)	11% (7.9-14%)	13% (9.6-17%)	35000 (26000-45000)	1200000 (900000-1600000)
High SDI	1990	Male	60 to 64	89% (84-93%)	11% (6.9-16%)	5.5 (4.2-7.1)	9.1% (6.2-12%)	11% (7.1-15%)	41000 (27000-56000)	1200000 (820000-1600000)
High SDI	1990	Male	65 to 69	88% (83-92%)	12% (7.9-17%)	5.1 (3.9-6.4)	7.6% (4.7-11%)	8.7% (5.3-12%)	41000 (25000-58000)	1000000 (630000-1400000)
High SDI	1990	Male	70 to 74	86% (80-91%)	14% (9.4-20%)	4.2 (3.1-5.4)	5.5% (2.7-8.4%)	6.1% (2.7-9.6%)	29000 (13000-47000)	590000 (280000-920000)
High SDI	1990	Male	75 to 79	85% (79-91%)	15% (9.4-21%)	4.1 (3-5.4)	5.2% (2.2-8.4%)	5.7% (2.1-9.4%)	35000 (13000-59000)	550000 (230000-900000)
High SDI	1990	Male	80 to 84	84% (76-90%)	16% (10-24%)	3.8 (2.8-5)	4.7% (1.7-7.8%)	5% (1.6-8.5%)	29000 (8900-50000)	330000 (120000-560000)
High SDI	1990	Male	85 to 89	82% (74-88%)	18% (12-26%)	3.5 (2.5-4.6)	3.8% (0.98-6.8%)	4% (0.66-7.4%)	15000 (2600-28000)	120000 (32000-220000)
High SDI	1990	Male	90 to 94	82% (74-88%)	18% (12-26%)	3.2 (2.2-4.4)	2.8% (-0.18-5.8%)	2.7% (-0.73-6.3%)	4200 (-900-9500)	25000 (-830-52000)
High SDI	1990	Male	95 plus	81% (71-89%)	19% (11-29%)	2.8 (1.7-4.1)	1.7% (-1.4-5%)	1.4% (-2.3-5.3%)	650 (-890-2200)	3000 (-2200-8500)
High SDI	1995	Female	All Ages	73% (70-76%)	27% (24-30%)	2.2 (1.9-2.6)	3.1% (1.8-4.4%)	2.8% (0.36-5.5%)	120000 (21000-240000)	3800000 (2300000-5500000)
High SDI	1995	Female	15 to 19	68% (59-77%)	32% (23-41%)	1.9 (1.3-2.7)	2.9% (1.8-4.2%)	8.9% (4.8-14%)	930 (510-1400)	100000 (65000-150000)
High SDI	1995	Female	20 to 24	80% (72-86%)	20% (14-28%)	3 (2.1-4)	4.4% (3.1-5.8%)	12% (7-17%)	1500 (940-2200)	190000 (130000-250000)
High SDI	1995	Female	25 to 29	83% (76-88%)	17% (12-24%)	2.8 (2.1-3.7)	3.9% (2.8-5.1%)	10% (6.4-14%)	1700 (1100-2400)	190000 (140000-250000)
High SDI	1995	Female	30 to 34	84% (77-89%)	16% (11-23%)	2.8 (2.1-3.7)	3.7% (2.7-4.9%)	10% (6.8-13%)	2400 (1600-3200)	210000 (160000-280000)
High SDI	1995	Female	35 to 39	83% (77-89%)	17% (11-23%)	2.8 (2.1-3.7)	4.2% (3-5.4%)	11% (7.9-14%)	3800 (2800-4800)	260000 (200000-340000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	1995	Female	40 to 44	83% (77-88%)	17% (12-23%)	2.9 (2.2-3.8)	4.7% (3.3-6.1%)	11% (8.5-14%)	5600 (4200-7000)	320000 (240000-410000)
High SDI	1995	Female	45 to 49	83% (76-88%)	17% (12-24%)	3.1 (2.2-4.1)	5% (3.5-6.7%)	11% (8.3-14%)	7700 (5700-9900)	370000 (270000-480000)
High SDI	1995	Female	50 to 54	81% (74-86%)	19% (14-26%)	3.1 (2.2-4.1)	5.2% (3.5-7.1%)	10% (7.4-13%)	8900 (6400-12000)	360000 (250000-490000)
High SDI	1995	Female	55 to 59	79% (72-85%)	21% (15-28%)	3.2 (2.3-4.2)	5% (3.2-7%)	9% (6.2-12%)	11000 (7200-15000)	380000 (240000-520000)
High SDI	1995	Female	60 to 64	76% (68-82%)	24% (18-32%)	3 (2.1-4.1)	4.2% (2.2-6.5%)	6.8% (3.9-10%)	12000 (6400-18000)	350000 (180000-540000)
High SDI	1995	Female	65 to 69	72% (64-80%)	28% (20-36%)	2.7 (2-3.5)	3.3% (1.3-5.5%)	5% (2.2-8%)	13000 (5400-22000)	330000 (130000-550000)
High SDI	1995	Female	70 to 74	69% (60-77%)	31% (23-40%)	2.4 (1.7-3.3)	2.6% (0.33-5.2%)	3.7% (0.56-7%)	15000 (1300-30000)	300000 (26000-590000)
High SDI	1995	Female	75 to 79	65% (55-75%)	35% (25-45%)	2.1 (1.4-3)	1.6% (-0.81-4.3%)	2% (-1.1-5.4%)	9700 (-5900-28000)	160000 (-82000-430000)
High SDI	1995	Female	80 to 84	64% (52-73%)	36% (27-48%)	2.1 (1.5-2.9)	1.6% (-0.89-4.3%)	1.8% (-1.3-5.3%)	13000 (-10000-39000)	160000 (-92000-450000)
High SDI	1995	Female	85 to 89	62% (51-71%)	38% (29-49%)	2.1 (1.4-2.9)	1.3% (-1.2-4.2%)	1.4% (-1.7-4.9%)	10000 (-13000-38000)	94000 (-84000-300000)
High SDI	1995	Female	90 to 94	61% (50-71%)	39% (29-50%)	2 (1.3-2.9)	0.9% (-1.7-4.1%)	0.83% (-2.3-4.7%)	4300 (-10000-22000)	29000 (-48000-120000)
High SDI	1995	Female	95 plus	60% (46-72%)	40% (28-54%)	1.9 (1-2.9)	0.5% (-2.4-4%)	0.3% (-3.3-4.5%)	810 (-5900-8800)	4700 (-18000-32000)
High SDI	1995	Male	All Ages	83% (80-85%)	17% (15-20%)	3.4 (3-3.9)	8.1% (6.4-9.9%)	7.8% (5.2-10%)	330000 (220000-450000)	12000000 (9100000-14000000)
High SDI	1995	Male	15 to 19	73% (64-80%)	27% (20-36%)	2.3 (1.7-3)	6.9% (4.5-9.6%)	12% (7-18%)	3500 (2100-5100)	310000 (210000-430000)
High SDI	1995	Male	20 to 24	86% (79-91%)	14% (9.1-21%)	4.3 (3.3-5.4)	13% (9.7-17%)	20% (13-27%)	8600 (5900-11000)	750000 (550000-950000)
High SDI	1995	Male	25 to 29	89% (84-93%)	11% (7.1-16%)	4.4 (3.4-5.6)	12% (9.1-15%)	19% (12-25%)	8800 (6000-12000)	740000 (550000-940000)
High SDI	1995	Male	30 to 34	91% (86-94%)	8.9% (5.5-14%)	4.4 (3.4-5.6)	11% (8-13%)	16% (11-21%)	10000 (7200-13000)	760000 (570000-960000)
High SDI	1995	Male	35 to 39	92% (88-95%)	8.2% (5.2-12%)	4.2 (3.3-5.4)	10% (8-13%)	16% (11-20%)	13000 (9600-16000)	840000 (650000-1000000)
High SDI	1995	Male	40 to 44	92% (88-95%)	7.8% (4.9-12%)	4.3 (3.3-5.5)	11% (8.3-13%)	16% (12-20%)	17000 (13000-22000)	960000 (750000-1200000)
High SDI	1995	Male	45 to 49	92% (88-95%)	7.6% (4.7-12%)	4.8 (3.6-6.1)	11% (8.5-14%)	16% (12-20%)	23000 (18000-29000)	1100000 (860000-1400000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	1995	Male	50 to 54	92% (88-95%)	8% (5.2-12%)	5 (3.8-6.4)	11% (8.3-14%)	15% (11-18%)	27000 (20000-33000)	1100000 (830000-1400000)
High SDI	1995	Male	55 to 59	91% (87-94%)	8.8% (5.8-13%)	5.4 (4.1-6.9)	11% (8-13%)	13% (10-17%)	34000 (26000-43000)	1200000 (910000-1500000)
High SDI	1995	Male	60 to 64	90% (85-93%)	10% (6.7-15%)	5.2 (3.9-6.8)	9% (6.3-12%)	11% (7.5-15%)	39000 (27000-52000)	1100000 (810000-1500000)
High SDI	1995	Male	65 to 69	88% (83-92%)	12% (7.8-17%)	4.7 (3.7-5.9)	7.4% (4.9-10%)	8.6% (5.5-12%)	42000 (26000-58000)	1000000 (670000-1400000)
High SDI	1995	Male	70 to 74	87% (81-91%)	13% (9.1-19%)	4.2 (3.1-5.4)	5.9% (3.2-8.7%)	6.6% (3.4-10%)	38000 (19000-58000)	760000 (400000-1100000)
High SDI	1995	Male	75 to 79	85% (78-90%)	15% (9.9-22%)	3.5 (2.5-4.6)	4% (1.4-6.9%)	4.3% (1.2-7.7%)	24000 (6200-42000)	380000 (120000-650000)
High SDI	1995	Male	80 to 84	84% (76-89%)	16% (11-24%)	3.5 (2.6-4.5)	4% (1.4-6.9%)	4.3% (1.2-7.6%)	26000 (6500-46000)	300000 (99000-520000)
High SDI	1995	Male	85 to 89	83% (75-88%)	17% (12-25%)	3.2 (2.3-4.3)	3.4% (0.7-6.2%)	3.5% (0.35-6.8%)	15000 (1500-29000)	130000 (26000-240000)
High SDI	1995	Male	90 to 94	83% (75-88%)	17% (12-25%)	3 (2-4.1)	2.5% (-0.41-5.5%)	2.4% (-1-6%)	4700 (-1800-11000)	29000 (-4300-64000)
High SDI	1995	Male	95 plus	82% (72-89%)	18% (11-28%)	2.7 (1.7-4)	1.7% (-1.4-5%)	1.5% (-2.2-5.3%)	790 (-1100-2700)	3600 (-2700-10000)
High SDI	2000	Female	All Ages	73% (69-76%)	27% (24-31%)	2.2 (1.9-2.5)	2.9% (1.7-4.1%)	2.5% (0.29-5%)	110000 (18000-220000)	3600000 (2100000-5100000)
High SDI	2000	Female	15 to 19	68% (58-76%)	32% (24-42%)	1.9 (1.3-2.6)	2.7% (1.7-3.9%)	8.9% (4.7-14%)	840 (450-1300)	95000 (62000-130000)
High SDI	2000	Female	20 to 24	79% (71-86%)	21% (14-29%)	2.9 (2.1-3.9)	4.1% (2.9-5.4%)	12% (7-17%)	1400 (830-1900)	160000 (120000-220000)
High SDI	2000	Female	25 to 29	83% (76-88%)	17% (12-24%)	2.8 (2.1-3.7)	3.7% (2.6-4.8%)	11% (6.6-15%)	1500 (930-2100)	170000 (120000-220000)
High SDI	2000	Female	30 to 34	84% (77-89%)	16% (11-23%)	2.7 (2-3.6)	3.5% (2.5-4.6%)	10% (7-14%)	2000 (1400-2700)	190000 (130000-250000)
High SDI	2000	Female	35 to 39	84% (78-89%)	16% (11-22%)	2.8 (2.1-3.7)	4% (2.9-5.2%)	11% (8.1-14%)	3500 (2600-4500)	250000 (190000-320000)
High SDI	2000	Female	40 to 44	83% (77-88%)	17% (12-23%)	2.9 (2.1-3.7)	4.5% (3.3-5.9%)	11% (8.6-14%)	5500 (4200-7000)	320000 (240000-410000)
High SDI	2000	Female	45 to 49	82% (75-87%)	18% (13-25%)	3 (2.2-3.9)	4.9% (3.5-6.5%)	11% (8.3-14%)	7700 (5800-9700)	370000 (270000-480000)
High SDI	2000	Female	50 to 54	81% (74-86%)	19% (14-26%)	3 (2.2-3.9)	4.9% (3.3-6.6%)	9.9% (7.2-13%)	9600 (7100-12000)	400000 (280000-530000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	2000	Female	55 to 59	79% (72-85%)	21% (15-28%)	3 (2.2-3.9)	4.6% (2.9-6.4%)	8.5% (6-11%)	10000 (7000-13000)	350000 (230000-490000)
High SDI	2000	Female	60 to 64	76% (68-82%)	24% (18-32%)	2.8 (2-3.9)	3.9% (2.1-6%)	6.7% (4-9.8%)	11000 (6400-16000)	330000 (180000-500000)
High SDI	2000	Female	65 to 69	71% (62-79%)	29% (21-38%)	2.5 (1.8-3.3)	3% (1.3-4.9%)	4.8% (2.3-7.4%)	11000 (5200-18000)	280000 (120000-460000)
High SDI	2000	Female	70 to 74	68% (59-76%)	32% (24-41%)	2.2 (1.5-3.1)	2.3% (0.27-4.5%)	3.3% (0.64-6.3%)	12000 (1800-24000)	240000 (23000-490000)
High SDI	2000	Female	75 to 79	66% (56-75%)	34% (25-44%)	2.1 (1.4-2.9)	1.6% (-0.55-4.1%)	2.1% (-0.71-5.3%)	12000 (-4700-31000)	190000 (-67000-490000)
High SDI	2000	Female	80 to 84	63% (51-73%)	37% (27-49%)	1.9 (1.3-2.6)	1% (-1-3.3%)	1.2% (-1.4-4.1%)	7800 (-8900-27000)	95000 (-91000-310000)
High SDI	2000	Female	85 to 89	62% (51-72%)	38% (28-49%)	1.9 (1.3-2.8)	1% (-1.2-3.6%)	1% (-1.7-4.2%)	8700 (-14000-34000)	79000 (-92000-280000)
High SDI	2000	Female	90 to 94	62% (51-72%)	38% (28-49%)	1.9 (1.2-2.9)	0.7% (-1.7-3.6%)	0.61% (-2.4-4.1%)	3800 (-13000-24000)	27000 (-61000-130000)
High SDI	2000	Female	95 plus	63% (49-75%)	37% (25-51%)	1.9 (1-2.9)	0.35% (-2.5-3.8%)	0.13% (-3.3-4.3%)	520 (-7800-11000)	4200 (-24000-38000)
High SDI	2000	Male	All Ages	83% (81-86%)	17% (14-19%)	3.4 (2.9-3.9)	8% (6.3-9.7%)	7.5% (5.1-10%)	320000 (220000-430000)	11000000 (8800000-14000000)
High SDI	2000	Male	15 to 19	72% (64-79%)	28% (21-36%)	2.2 (1.6-3)	6.5% (4.3-8.9%)	12% (6.9-18%)	2900 (1700-4200)	270000 (180000-370000)
High SDI	2000	Male	20 to 24	86% (79-91%)	14% (9.5-21%)	4.2 (3.3-5.4)	13% (9.2-16%)	20% (13-27%)	7300 (5000-9600)	640000 (470000-810000)
High SDI	2000	Male	25 to 29	89% (85-93%)	11% (7-15%)	4.3 (3.3-5.4)	12% (8.7-15%)	20% (13-26%)	7400 (5100-9800)	630000 (470000-800000)
High SDI	2000	Male	30 to 34	91% (87-95%)	8.7% (5.4-13%)	4.2 (3.3-5.4)	11% (8-14%)	18% (12-24%)	8300 (5800-11000)	640000 (480000-810000)
High SDI	2000	Male	35 to 39	92% (88-95%)	8% (5.1-12%)	4.1 (3.1-5.3)	11% (8-13%)	17% (12-22%)	12000 (8500-15000)	770000 (590000-950000)
High SDI	2000	Male	40 to 44	92% (89-95%)	7.7% (4.9-11%)	4.2 (3.3-5.4)	11% (8.6-14%)	17% (13-21%)	17000 (13000-21000)	960000 (750000-1200000)
High SDI	2000	Male	45 to 49	92% (88-95%)	7.6% (4.9-12%)	4.6 (3.5-5.9)	12% (8.9-14%)	17% (13-21%)	24000 (19000-30000)	1200000 (900000-1400000)
High SDI	2000	Male	50 to 54	92% (89-95%)	7.8% (5.1-11%)	4.8 (3.7-6.1)	11% (8.3-13%)	15% (11-18%)	30000 (23000-37000)	1200000 (940000-1500000)
High SDI	2000	Male	55 to 59	91% (88-94%)	8.6% (5.7-12%)	4.9 (3.8-6.3)	9.9% (7.5-12%)	13% (9.7-16%)	31000 (23000-39000)	1100000 (840000-1400000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	2000	Male	60 to 64	90% (86-94%)	9.7% (6.5-14%)	4.9 (3.7-6.4)	9% (6.4-12%)	11% (7.9-15%)	37000 (26000-48000)	1100000 (790000-1400000)
High SDI	2000	Male	65 to 69	89% (84-93%)	11% (7.5-16%)	4.4 (3.5-5.6)	7.3% (4.9-9.8%)	8.7% (5.8-12%)	38000 (25000-52000)	960000 (650000-1300000)
High SDI	2000	Male	70 to 74	87% (82-91%)	13% (8.8-18%)	4 (3-5.1)	5.7% (3.1-8.4%)	6.6% (3.5-9.8%)	38000 (20000-57000)	760000 (420000-1100000)
High SDI	2000	Male	75 to 79	86% (80-91%)	14% (9.4-20%)	3.5 (2.6-4.7)	4.4% (1.8-7.2%)	4.9% (1.7-8.2%)	31000 (11000-53000)	500000 (200000-820000)
High SDI	2000	Male	80 to 84	84% (76-89%)	16% (11-24%)	3 (2.2-4)	3.2% (0.79-5.7%)	3.3% (0.51-6.3%)	18000 (2700-34000)	220000 (53000-390000)
High SDI	2000	Male	85 to 89	83% (75-89%)	17% (11-25%)	3 (2.2-4)	3% (0.47-5.7%)	3% (0.076-6.2%)	14000 (390-29000)	130000 (19000-240000)
High SDI	2000	Male	90 to 94	83% (76-89%)	17% (11-24%)	2.9 (2-4)	2.3% (-0.47-5.3%)	2.2% (-1.1-5.7%)	5100 (-2400-13000)	32000 (-6600-74000)
High SDI	2000	Male	95 plus	83% (74-90%)	17% (9.9-26%)	2.6 (1.6-3.9)	1.6% (-1.4-4.9%)	1.3% (-2.3-5.2%)	870 (-1400-3400)	4100 (-3600-13000)
High SDI	2005	Female	All Ages	72% (69-75%)	28% (25-31%)	2.2 (1.9-2.5)	2.8% (1.8-3.9%)	2.5% (0.56-4.7%)	110000 (27000-200000)	3500000 (2300000-4900000)
High SDI	2005	Female	15 to 19	67% (58-75%)	33% (25-42%)	1.9 (1.3-2.6)	2.6% (1.7-3.6%)	8.8% (4.6-13%)	730 (400-1100)	88000 (58000-120000)
High SDI	2005	Female	20 to 24	79% (71-85%)	21% (15-29%)	3 (2.2-4)	4.1% (3-5.4%)	12% (7-17%)	1300 (830-1900)	170000 (120000-220000)
High SDI	2005	Female	25 to 29	82% (75-87%)	18% (13-25%)	2.8 (2.1-3.7)	3.7% (2.7-4.7%)	11% (6.6-15%)	1400 (880-1900)	160000 (120000-210000)
High SDI	2005	Female	30 to 34	83% (77-88%)	17% (12-23%)	2.7 (2-3.6)	3.4% (2.4-4.5%)	11% (6.8-14%)	1900 (1200-2500)	170000 (120000-230000)
High SDI	2005	Female	35 to 39	83% (78-88%)	17% (12-22%)	2.8 (2.1-3.7)	3.8% (2.7-4.9%)	11% (8-14%)	3100 (2300-4000)	230000 (170000-290000)
High SDI	2005	Female	40 to 44	83% (77-88%)	17% (12-23%)	2.9 (2.2-3.8)	4.3% (3.1-5.7%)	12% (8.6-15%)	5300 (4000-6700)	310000 (230000-400000)
High SDI	2005	Female	45 to 49	82% (75-87%)	18% (13-25%)	3 (2.2-3.9)	4.8% (3.5-6.4%)	11% (8.4-14%)	7900 (6100-9900)	390000 (290000-490000)
High SDI	2005	Female	50 to 54	80% (74-85%)	20% (15-26%)	2.9 (2.2-3.8)	4.8% (3.4-6.5%)	10% (7.5-13%)	9900 (7500-13000)	410000 (300000-540000)
High SDI	2005	Female	55 to 59	78% (72-84%)	22% (16-28%)	2.9 (2.1-3.8)	4.4% (2.9-6.1%)	8.4% (6.2-11%)	11000 (8300-15000)	400000 (280000-540000)
High SDI	2005	Female	60 to 64	75% (68-81%)	25% (19-32%)	2.7 (2-3.7)	3.7% (2.1-5.5%)	6.6% (4.3-9.1%)	11000 (6800-15000)	320000 (180000-460000)
High SDI	2005	Female	65 to 69	71% (62-78%)	29% (22-38%)	2.4 (1.8-3.2)	3% (1.5-4.6%)	4.9% (2.8-7.2%)	11000 (6200-16000)	270000 (130000-420000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	2005	Female	70 to 74	67% (59-74%)	33% (26-41%)	2.1 (1.5-2.9)	2.2% (0.5-4%)	3.4% (1.1-5.8%)	11000 (3400-20000)	220000 (48000-410000)
High SDI	2005	Female	75 to 79	65% (55-73%)	35% (27-45%)	1.9 (1.3-2.7)	1.5% (-0.27-3.6%)	2.1% (-0.23-4.7%)	11000 (-1500-26000)	170000 (-30000-410000)
High SDI	2005	Female	80 to 84	63% (51-72%)	37% (28-49%)	1.9 (1.3-2.6)	1.2% (-0.66-3.2%)	1.4% (-0.9-4%)	11000 (-6600-32000)	130000 (-68000-370000)
High SDI	2005	Female	85 to 89	61% (50-71%)	39% (29-50%)	1.8 (1.2-2.5)	0.76% (-1.1-2.9%)	0.81% (-1.5-3.4%)	6300 (-11000-25000)	57000 (-76000-210000)
High SDI	2005	Female	90 to 94	62% (52-72%)	38% (28-48%)	1.9 (1.2-2.7)	0.67% (-1.5-3.3%)	0.61% (-2.1-3.8%)	4200 (-13000-25000)	29000 (-61000-140000)
High SDI	2005	Female	95 plus	64% (50-76%)	36% (24-50%)	1.9 (1.1-3)	0.48% (-2.2-3.7%)	0.32% (-2.9-4.2%)	1200 (-8700-13000)	6800 (-27000-48000)
High SDI	2005	Male	All Ages	83% (81-86%)	17% (14-19%)	3.4 (2.9-3.9)	7.9% (6.3-9.6%)	7.6% (5.2-10%)	330000 (230000-430000)	11000000 (8900000-13000000)
High SDI	2005	Male	15 to 19	71% (63-78%)	29% (22-37%)	2.3 (1.6-3)	6.1% (4.2-8.3%)	12% (7.1-17%)	2600 (1600-3600)	250000 (170000-330000)
High SDI	2005	Male	20 to 24	86% (80-90%)	14% (9.6-20%)	4.4 (3.4-5.6)	12% (9.1-16%)	20% (13-27%)	7100 (5000-9300)	630000 (470000-800000)
High SDI	2005	Male	25 to 29	89% (85-93%)	11% (7.1-15%)	4.4 (3.4-5.5)	12% (8.7-15%)	20% (13-26%)	6900 (4800-9000)	590000 (450000-750000)
High SDI	2005	Male	30 to 34	91% (87-95%)	8.6% (5.5-13%)	4.3 (3.3-5.5)	10% (7.8-13%)	18% (12-24%)	7600 (5300-9900)	590000 (440000-740000)
High SDI	2005	Male	35 to 39	92% (88-95%)	7.9% (5-12%)	4.1 (3.1-5.3)	10% (7.6-13%)	17% (12-22%)	10000 (7300-13000)	680000 (520000-850000)
High SDI	2005	Male	40 to 44	92% (89-95%)	7.7% (4.9-11%)	4.2 (3.3-5.4)	11% (8.3-13%)	17% (13-21%)	16000 (12000-20000)	910000 (710000-1100000)
High SDI	2005	Male	45 to 49	92% (88-95%)	7.7% (5-12%)	4.7 (3.6-6)	12% (9-14%)	17% (13-21%)	25000 (19000-30000)	1200000 (930000-1400000)
High SDI	2005	Male	50 to 54	92% (88-95%)	8% (5.3-12%)	4.8 (3.7-6.1)	11% (8.8-14%)	16% (12-19%)	32000 (25000-39000)	1300000 (1000000-1600000)
High SDI	2005	Male	55 to 59	91% (88-94%)	8.7% (5.8-12%)	4.8 (3.7-6.1)	9.9% (7.7-12%)	13% (10-16%)	36000 (28000-44000)	1300000 (1000000-1600000)
High SDI	2005	Male	60 to 64	90% (86-93%)	9.8% (6.6-14%)	4.7 (3.5-6)	8.5% (6.2-11%)	11% (7.8-14%)	34000 (25000-44000)	1000000 (770000-1300000)
High SDI	2005	Male	65 to 69	89% (84-93%)	11% (7.4-16%)	4.4 (3.4-5.5)	7.3% (5.1-9.6%)	9% (6.2-12%)	37000 (26000-49000)	940000 (660000-1200000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	2005	Male	70 to 74	88% (83-91%)	12% (8.9-17%)	3.8 (2.9-5)	5.7% (3.3-8.2%)	6.7% (3.8-9.7%)	36000 (20000-52000)	730000 (430000-1000000)
High SDI	2005	Male	75 to 79	86% (80-91%)	14% (9.4-20%)	3.4 (2.5-4.5)	4.4% (1.9-7%)	4.9% (2-8%)	32000 (13000-53000)	520000 (230000-830000)
High SDI	2005	Male	80 to 84	84% (77-90%)	16% (10-23%)	3.1 (2.3-4.1)	3.5% (1.2-6%)	3.8% (1-6.7%)	25000 (7100-45000)	300000 (110000-520000)
High SDI	2005	Male	85 to 89	83% (76-88%)	17% (12-24%)	2.8 (2-3.7)	2.4% (0.26-4.9%)	2.5% (-0.15-5.3%)	11000 (-550-24000)	100000 (11000-200000)
High SDI	2005	Male	90 to 94	84% (77-89%)	16% (11-23%)	2.8 (1.9-3.8)	2.1% (-0.4-5%)	2% (-0.97-5.4%)	5600 (-2600-15000)	36000 (-6700-83000)
High SDI	2005	Male	95 plus	84% (75-91%)	16% (9.4-25%)	2.6 (1.6-3.9)	1.7% (-1.2-5%)	1.4% (-2-5.4%)	1200 (-1600-4500)	5600 (-3900-17000)
High SDI	2010	Female	All Ages	72% (68-75%)	28% (25-32%)	2.2 (1.8-2.6)	2.7% (1.8-3.8%)	2.5% (0.65-4.6%)	110000 (30000-200000)	350000 (230000-490000)
High SDI	2010	Female	15 to 19	61% (53-69%)	39% (31-47%)	1.6 (1.1-2.3)	2.2% (1.5-3.1%)	7.9% (4.1-12%)	540 (290-830)	73000 (48000-100000)
High SDI	2010	Female	20 to 24	79% (71-85%)	21% (15-29%)	3 (2.2-4)	4% (2.9-5.3%)	12% (7-17%)	1200 (770-1700)	160000 (110000-210000)
High SDI	2010	Female	25 to 29	82% (75-87%)	18% (13-25%)	2.8 (2.1-3.7)	3.7% (2.7-4.8%)	11% (6.5-15%)	1400 (900-2000)	170000 (120000-220000)
High SDI	2010	Female	30 to 34	83% (76-88%)	17% (12-24%)	2.7 (2-3.6)	3.4% (2.4-4.5%)	11% (6.9-14%)	1800 (1200-2400)	170000 (120000-220000)
High SDI	2010	Female	35 to 39	83% (77-88%)	17% (12-23%)	2.8 (2-3.7)	3.6% (2.6-4.8%)	11% (7.8-15%)	2800 (2000-3700)	210000 (150000-270000)
High SDI	2010	Female	40 to 44	82% (76-87%)	18% (13-24%)	2.9 (2.1-3.8)	4.1% (2.9-5.4%)	12% (8.5-15%)	4700 (3500-5900)	280000 (210000-360000)
High SDI	2010	Female	45 to 49	82% (75-87%)	18% (13-25%)	3 (2.2-3.9)	4.6% (3.3-6.1%)	11% (8.6-14%)	7600 (5900-9500)	380000 (280000-480000)
High SDI	2010	Female	50 to 54	80% (74-85%)	20% (15-26%)	2.9 (2.1-3.8)	4.8% (3.4-6.4%)	10% (7.6-13%)	10000 (7700-13000)	430000 (310000-550000)
High SDI	2010	Female	55 to 59	78% (71-84%)	22% (16-29%)	2.8 (2.1-3.8)	4.3% (2.9-5.9%)	8.5% (6.2-11%)	11000 (8500-15000)	410000 (280000-550000)
High SDI	2010	Female	60 to 64	75% (68-81%)	25% (19-32%)	2.6 (1.9-3.6)	3.5% (2-5.2%)	6.5% (4.3-8.9%)	12000 (8000-16000)	360000 (210000-520000)
High SDI	2010	Female	65 to 69	71% (63-78%)	29% (22-37%)	2.4 (1.7-3.1)	2.8% (1.4-4.3%)	4.8% (2.9-6.9%)	11000 (6500-16000)	260000 (140000-400000)
High SDI	2010	Female	70 to 74	68% (60-74%)	32% (26-40%)	2.1 (1.5-2.9)	2.2% (0.62-3.9%)	3.5% (1.4-5.8%)	11000 (4400-19000)	220000 (63000-390000)
High SDI	2010	Female	75 to 79	65% (55-73%)	35% (27-45%)	1.9 (1.3-2.7)	1.5% (-0.11-3.4%)	2.2% (0.097-4.6%)	11000 (280-22000)	160000 (-11000-360000)
High SDI	2010	Female	80 to 84	63% (51-72%)	37% (28-49%)	1.8 (1.2-2.5)	1.1% (-0.54-3%)	1.4% (-0.69-3.8%)	11000 (-4900-28000)	130000 (-56000-340000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	2010	Female	85 to 89	62% (51-71%)	38% (29-49%)	1.8 (1.2-2.6)	0.91% (-0.89-3%)	1% (-1.2-3.6%)	9900 (-11000-33000)	87000 (-77000-270000)
High SDI	2010	Female	90 to 94	62% (51-71%)	38% (29-49%)	1.7 (1.1-2.5)	0.5% (-1.4-2.8%)	0.44% (-1.9-3.2%)	3200 (-11000-21000)	22000 (-57000-120000)
High SDI	2010	Female	95 plus	64% (50-76%)	36% (24-50%)	1.8 (1-2.9)	0.47% (-2-3.5%)	0.33% (-2.7-4%)	1600 (-10000-16000)	8500 (-31000-58000)
High SDI	2010	Male	All Ages	83% (81-85%)	17% (15-19%)	3.4 (2.9-4)	7.7% (6-9.4%)	7.3% (5-9.7%)	320000 (220000-430000)	1100000 (850000-1300000)
High SDI	2010	Male	15 to 19	68% (60-76%)	32% (24-40%)	2.2 (1.5-2.9)	5.5% (3.8-7.4%)	11% (6.6-17%)	2000 (1200-2800)	200000 (140000-270000)
High SDI	2010	Male	20 to 24	86% (79-90%)	14% (9.6-21%)	4.4 (3.4-5.7)	12% (8.8-15%)	20% (13-27%)	6200 (4300-8100)	570000 (430000-720000)
High SDI	2010	Male	25 to 29	89% (84-93%)	11% (7.3-16%)	4.4 (3.4-5.7)	11% (8.4-14%)	19% (13-26%)	6500 (4500-8500)	570000 (430000-720000)
High SDI	2010	Male	30 to 34	91% (86-94%)	9.1% (5.7-14%)	4.2 (3.2-5.5)	10% (7.6-13%)	18% (12-24%)	6900 (4800-9000)	550000 (410000-690000)
High SDI	2010	Male	35 to 39	92% (88-95%)	8.2% (5.2-12%)	4 (3-5.3)	9.8% (7.3-12%)	17% (12-23%)	9000 (6400-12000)	610000 (460000-770000)
High SDI	2010	Male	40 to 44	92% (88-95%)	7.8% (4.9-12%)	4.2 (3.1-5.4)	10% (7.7-13%)	17% (13-22%)	14000 (10000-17000)	780000 (600000-970000)
High SDI	2010	Male	45 to 49	92% (88-95%)	7.9% (5-12%)	4.6 (3.4-5.9)	11% (8.6-14%)	17% (13-21%)	22000 (17000-28000)	1100000 (850000-1300000)
High SDI	2010	Male	50 to 54	92% (88-94%)	8.3% (5.5-12%)	4.8 (3.6-6.1)	11% (8.7-14%)	16% (12-20%)	32000 (25000-39000)	1300000 (1000000-1600000)
High SDI	2010	Male	55 to 59	91% (87-94%)	9.3% (6.2-13%)	4.8 (3.6-6.1)	10% (7.9-13%)	14% (11-17%)	38000 (29000-46000)	1300000 (1100000-1600000)
High SDI	2010	Male	60 to 64	90% (85-93%)	10% (6.9-15%)	4.5 (3.4-5.9)	8.4% (6.1-11%)	11% (8-14%)	39000 (28000-49000)	1200000 (880000-1500000)
High SDI	2010	Male	65 to 69	88% (84-92%)	12% (7.8-16%)	4.1 (3.2-5.2)	6.8% (4.8-9%)	8.6% (6-11%)	35000 (24000-45000)	880000 (620000-1200000)
High SDI	2010	Male	70 to 74	88% (83-91%)	12% (8.7-17%)	3.8 (2.8-4.9)	5.6% (3.4-8%)	6.8% (4-9.7%)	34000 (20000-49000)	710000 (430000-1000000)
High SDI	2010	Male	75 to 79	86% (81-91%)	14% (9.1-19%)	3.3 (2.4-4.5)	4.3% (2-6.7%)	4.9% (2.2-7.8%)	31000 (13000-49000)	500000 (230000-790000)
High SDI	2010	Male	80 to 84	85% (78-90%)	15% (9.9-22%)	3.1 (2.2-4)	3.4% (1.2-5.8%)	3.7% (1.1-6.5%)	27000 (7900-47000)	320000 (120000-550000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	2010	Male	85 to 89	84% (77-89%)	16% (11-23%)	2.9 (2.1-3.9)	2.7% (0.52-5.1%)	2.8% (0.16-5.6%)	17000 (1200-34000)	150000 (29000-280000)
High SDI	2010	Male	90 to 94	83% (76-89%)	17% (11-24%)	2.6 (1.8-3.6)	1.8% (-0.45-4.4%)	1.7% (-1-4.7%)	4900 (-2800-14000)	32000 (-8200-78000)
High SDI	2010	Male	95 plus	84% (75-91%)	16% (9.5-25%)	2.5 (1.5-3.8)	1.7% (-1-4.9%)	1.5% (-1.8-5.2%)	1600 (-1900-5700)	7400 (-4500-21000)
High SDI	2016	Female	All Ages	72% (69-75%)	28% (25-31%)	2.2 (1.6-2.8)	2.6% (1.5-4%)	2.4% (0.29-4.8%)	110000 (14000-240000)	350000 (200000-540000)
High SDI	2016	Female	15 to 19	61% (52-69%)	39% (31-48%)	1.6 (1-2.4)	2.1% (1.4-3%)	7.8% (4-12%)	470 (240-740)	65000 (43000-92000)
High SDI	2016	Female	20 to 24	79% (70-85%)	21% (15-30%)	3 (2-4.3)	3.9% (2.8-5.2%)	12% (6.8-17%)	1200 (720-1700)	160000 (110000-210000)
High SDI	2016	Female	25 to 29	83% (76-88%)	17% (12-24%)	2.9 (2-4)	3.6% (2.6-4.8%)	11% (6.4-16%)	1400 (850-2000)	160000 (120000-220000)
High SDI	2016	Female	30 to 34	83% (76-88%)	17% (12-24%)	2.7 (1.8-3.9)	3.4% (2.4-4.6%)	11% (6.7-15%)	1800 (1200-2500)	170000 (120000-230000)
High SDI	2016	Female	35 to 39	83% (76-88%)	17% (12-24%)	2.7 (1.8-3.9)	3.6% (2.4-4.9%)	11% (7.6-15%)	2600 (1800-3500)	200000 (140000-260000)
High SDI	2016	Female	40 to 44	83% (76-88%)	17% (12-24%)	2.8 (1.9-4)	3.9% (2.7-5.4%)	12% (8.2-15%)	4400 (3100-5800)	260000 (180000-350000)
High SDI	2016	Female	45 to 49	82% (75-87%)	18% (13-25%)	2.9 (2-4.2)	4.4% (3-6.1%)	11% (8.3-15%)	7000 (5100-9100)	350000 (240000-470000)
High SDI	2016	Female	50 to 54	80% (74-86%)	20% (14-26%)	2.9 (2-4.1)	4.6% (3.1-6.5%)	10% (7.4-13%)	10000 (7200-13000)	430000 (290000-580000)
High SDI	2016	Female	55 to 59	78% (71-84%)	22% (16-29%)	2.8 (1.9-3.9)	4.2% (2.7-6.1%)	8.4% (5.9-11%)	12000 (8100-16000)	420000 (270000-600000)
High SDI	2016	Female	60 to 64	75% (68-82%)	25% (18-32%)	2.6 (1.8-3.8)	3.5% (1.9-5.4%)	6.5% (4.1-9.3%)	13000 (7700-18000)	380000 (200000-580000)
High SDI	2016	Female	65 to 69	72% (63-79%)	28% (21-37%)	2.4 (1.6-3.3)	2.7% (1.2-4.4%)	4.8% (2.7-7.1%)	12000 (6800-19000)	310000 (140000-500000)
High SDI	2016	Female	70 to 74	68% (59-76%)	32% (24-41%)	2.1 (1.4-3.1)	2% (0.44-3.9%)	3.4% (1.2-5.9%)	11000 (3900-20000)	220000 (48000-430000)
High SDI	2016	Female	75 to 79	67% (57-76%)	33% (24-43%)	2 (1.2-2.9)	1.5% (-0.23-3.7%)	2.2% (-0.064-5%)	11000 (-490-25000)	170000 (-25000-410000)
High SDI	2016	Female	80 to 84	65% (53-74%)	35% (26-47%)	1.8 (1.2-2.7)	1.1% (-0.71-3.2%)	1.4% (-0.89-4.1%)	10000 (-6300-30000)	120000 (-76000-360000)
High SDI	2016	Female	85 to 89	63% (52-73%)	37% (27-48%)	1.8 (1.1-2.7)	0.84% (-1.1-3.1%)	0.95% (-1.4-3.7%)	9700 (-13000-37000)	86000 (-100000-310000)
High SDI	2016	Female	90 to 94	64% (53-73%)	36% (27-47%)	1.8 (1.1-2.8)	0.64% (-1.5-3.3%)	0.61% (-2-3.8%)	5900 (-18000-35000)	40000 (-87000-200000)
High SDI	2016	Female	95 plus	64% (50-76%)	36% (24-50%)	1.7 (0.91-2.8)	0.29% (-2.1-3.4%)	0.13% (-2.8-3.8%)	780 (-12000-16000)	5900 (-36000-59000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High SDI	2016	Male	All Ages	83% (80-85%)	17% (15-20%)	3.3 (2.5-4.4)	7.3% (5.4-9.4%)	6.8% (4.1-9.7%)	330000 (200000-480000)	11000000 (7800000-14000000)
High SDI	2016	Male	15 to 19	68% (59-76%)	32% (24-41%)	2.1 (1.4-3.1)	5.2% (3.5-7.3%)	11% (6.3-17%)	1700 (980-2500)	180000 (120000-240000)
High SDI	2016	Male	20 to 24	85% (78-90%)	15% (10-22%)	4.4 (3.1-6.1)	12% (8.5-15%)	20% (12-27%)	6000 (4000-8100)	550000 (410000-710000)
High SDI	2016	Male	25 to 29	89% (83-93%)	11% (7.4-17%)	4.4 (3.1-6.1)	11% (8.1-14%)	19% (12-27%)	6400 (4300-8700)	570000 (420000-730000)
High SDI	2016	Male	30 to 34	91% (85-94%)	9.4% (5.8-15%)	4.2 (2.9-5.9)	10% (7.4-13%)	18% (12-25%)	7000 (4800-9500)	560000 (410000-720000)
High SDI	2016	Male	35 to 39	91% (87-95%)	8.8% (5.4-13%)	4 (2.7-5.6)	9.7% (7.1-13%)	17% (12-23%)	8400 (5800-11000)	580000 (420000-740000)
High SDI	2016	Male	40 to 44	92% (88-95%)	8.1% (5-12%)	4 (2.8-5.6)	9.8% (7.2-13%)	17% (12-22%)	12000 (8600-16000)	720000 (530000-920000)
High SDI	2016	Male	45 to 49	92% (87-95%)	8.1% (5-13%)	4.4 (3-6.2)	11% (7.9-14%)	17% (12-22%)	20000 (14000-26000)	970000 (720000-1200000)
High SDI	2016	Male	50 to 54	91% (87-94%)	8.6% (5.6-13%)	4.6 (3.2-6.4)	11% (8.1-14%)	16% (12-20%)	30000 (22000-38000)	1300000 (940000-1600000)
High SDI	2016	Male	55 to 59	90% (86-94%)	9.6% (6.3-14%)	4.6 (3.2-6.4)	10% (7.4-13%)	14% (10-18%)	37000 (27000-48000)	1300000 (990000-1700000)
High SDI	2016	Male	60 to 64	89% (85-93%)	11% (7.1-15%)	4.5 (3.1-6.2)	8.5% (5.9-11%)	11% (7.8-15%)	41000 (28000-54000)	1200000 (870000-1700000)
High SDI	2016	Male	65 to 69	88% (83-92%)	12% (7.9-17%)	4 (2.8-5.4)	6.6% (4.3-9%)	8.4% (5.5-11%)	39000 (25000-54000)	1000000 (650000-1400000)
High SDI	2016	Male	70 to 74	87% (82-91%)	13% (8.9-18%)	3.6 (2.4-5)	5.2% (2.8-7.8%)	6.4% (3.4-9.5%)	34000 (18000-52000)	710000 (380000-1100000)
High SDI	2016	Male	75 to 79	87% (80-91%)	13% (8.7-20%)	3.3 (2.2-4.7)	4.1% (1.7-6.8%)	4.7% (1.9-8.1%)	31000 (12000-53000)	510000 (210000-860000)
High SDI	2016	Male	80 to 84	85% (77-90%)	15% (9.5-23%)	3 (2-4.2)	3.2% (0.91-5.9%)	3.5% (0.76-6.7%)	27000 (5800-51000)	330000 (92000-610000)
High SDI	2016	Male	85 to 89	83% (76-89%)	17% (11-24%)	2.7 (1.8-3.9)	2.5% (0.2-5.2%)	2.6% (-0.16-5.8%)	19000 (-960-43000)	170000 (13000-360000)
High SDI	2016	Male	90 to 94	83% (76-89%)	17% (11-24%)	2.6 (1.7-3.9)	2% (-0.5-4.9%)	1.9% (-1-5.4%)	8600 (-4500-24000)	55000 (-14000-140000)
High SDI	2016	Male	95 plus	83% (73-90%)	17% (10-27%)	2.3 (1.3-3.6)	1.3% (-1.2-4.4%)	1% (-2-4.7%)	1400 (-2500-6100)	6800 (-6500-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	1990	Female	All Ages	49% (45-54%)	51% (46-55%)	1.6 (1.3-1.9)	3.6% (2.2-5.2%)	3.7% (1.4-6.5%)	99000 (34000-170000)	3300000 (2000000-4700000)
High-middle SDI	1990	Female	15 to 19	46% (36-57%)	54% (43-64%)	1.5 (0.93-2.3)	3% (1.9-4.5%)	6.3% (3.3-10%)	680 (350-1100)	75000 (46000-110000)
High-middle SDI	1990	Female	20 to 24	54% (44-63%)	46% (37-56%)	2.1 (1.3-3.1)	4.6% (3.2-6.2%)	7.9% (4.6-12%)	1100 (630-1600)	130000 (89000-190000)
High-middle SDI	1990	Female	25 to 29	59% (49-67%)	41% (33-51%)	2.2 (1.5-3.1)	5.3% (3.9-6.8%)	8.8% (5.8-12%)	1500 (970-2100)	170000 (120000-230000)
High-middle SDI	1990	Female	30 to 34	63% (53-71%)	37% (29-47%)	2.4 (1.6-3.3)	5.9% (4.5-7.5%)	10% (7.3-14%)	2300 (1600-3100)	220000 (160000-280000)
High-middle SDI	1990	Female	35 to 39	64% (55-72%)	36% (28-45%)	2.6 (1.7-3.6)	6.7% (5-8.5%)	12% (8.6-16%)	3500 (2400-4700)	260000 (190000-340000)
High-middle SDI	1990	Female	40 to 44	65% (57-73%)	35% (27-43%)	2.7 (1.8-3.8)	7.3% (5.4-9.5%)	13% (9.2-17%)	4600 (3300-6200)	280000 (200000-370000)
High-middle SDI	1990	Female	45 to 49	64% (54-73%)	36% (27-46%)	2.8 (1.8-3.9)	7.3% (5.1-9.9%)	12% (8.5-17%)	5100 (3500-7100)	260000 (180000-350000)
High-middle SDI	1990	Female	50 to 54	65% (56-73%)	35% (27-44%)	2.5 (1.7-3.6)	7.4% (5-10%)	11% (7.5-16%)	9000 (5900-13000)	400000 (270000-540000)
High-middle SDI	1990	Female	55 to 59	60% (49-69%)	40% (31-51%)	2.4 (1.6-3.5)	6.5% (3.8-9.5%)	9.6% (5.6-14%)	9500 (5400-14000)	350000 (200000-510000)
High-middle SDI	1990	Female	60 to 64	56% (45-67%)	44% (33-55%)	2.2 (1.4-3.3)	5.1% (2.2-8.5%)	7% (3-12%)	12000 (4500-20000)	350000 (150000-600000)
High-middle SDI	1990	Female	65 to 69	53% (42-64%)	47% (36-58%)	2 (1.3-3)	4.3% (1.4-7.6%)	5.7% (1.8-10%)	11000 (2800-21000)	280000 (79000-510000)
High-middle SDI	1990	Female	70 to 74	48% (37-58%)	52% (42-63%)	1.8 (1.1-2.7)	3.3% (0.35-6.9%)	4.2% (0.45-8.7%)	8800 (-93-19000)	180000 (4800-380000)
High-middle SDI	1990	Female	75 to 79	40% (29-52%)	60% (48-71%)	1.5 (0.84-2.3)	2.4% (-0.41-6%)	2.9% (-0.6-7.3%)	11000 (-3500-28000)	160000 (-43000-420000)
High-middle SDI	1990	Female	80 to 84	36% (25-48%)	64% (52-75%)	1.4 (0.8-2.1)	2.2% (-0.4-5.4%)	2.5% (-0.57-6.4%)	9500 (-3200-25000)	110000 (-28000-280000)
High-middle SDI	1990	Female	85 to 89	35% (24-47%)	65% (53-76%)	1.4 (0.77-2.2)	2% (-0.58-5.3%)	2.3% (-0.82-6.3%)	6100 (-2900-17000)	49000 (-18000-130000)
High-middle SDI	1990	Female	90 to 94	37% (26-48%)	63% (52-74%)	1.4 (0.76-2.4)	2% (-0.8-5.7%)	2.3% (-1.1-6.7%)	2700 (-1500-8200)	15000 (-6300-43000)
High-middle SDI	1990	Female	95 plus	38% (24-54%)	62% (46-76%)	1.4 (0.65-2.6)	1.8% (-1.4-6.1%)	2% (-1.8-7.1%)	910 (-820-3200)	3400 (-2400-11000)
High-middle SDI	1990	Male	All Ages	57% (53-61%)	43% (39-47%)	3.1 (2.6-3.7)	9.3% (6.9-12%)	9.5% (6.3-13%)	290000 (190000-390000)	11000000 (8000000-14000000)
High-middle SDI	1990	Male	15 to 19	52% (41-63%)	48% (37-59%)	1.9 (1.2-2.9)	6.2% (3.9-9.1%)	9.4% (5.4-15%)	2400 (1400-3700)	210000 (130000-310000)
High-middle SDI	1990	Male	20 to 24	59% (50-67%)	41% (33-50%)	3.3 (2.2-4.7)	11% (7.8-16%)	15% (9.5-21%)	6600 (4200-9300)	540000 (370000-730000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	1990	Male	25 to 29	65% (58-72%)	35% (28-42%)	4.1 (2.8-5.6)	14% (10-18%)	18% (13-24%)	10000 (7200-14000)	770000 (570000-1000000)
High-middle SDI	1990	Male	30 to 34	69% (62-76%)	31% (24-38%)	4.7 (3.3-6.4)	15% (12-19%)	20% (15-26%)	15000 (11000-19000)	990000 (750000-1300000)
High-middle SDI	1990	Male	35 to 39	71% (64-78%)	29% (22-36%)	5.1 (3.4-7.1)	16% (12-21%)	21% (15-27%)	19000 (14000-25000)	1100000 (820000-1400000)
High-middle SDI	1990	Male	40 to 44	73% (66-79%)	27% (21-34%)	5.5 (3.8-7.6)	17% (12-21%)	22% (16-28%)	22000 (16000-28000)	1100000 (820000-1400000)
High-middle SDI	1990	Male	45 to 49	72% (64-78%)	28% (22-36%)	5.9 (4.1-8.3)	16% (12-21%)	20% (15-26%)	22000 (15000-29000)	980000 (710000-1300000)
High-middle SDI	1990	Male	50 to 54	76% (69-81%)	24% (19-31%)	6 (4-8.2)	16% (11-20%)	19% (13-25%)	38000 (25000-50000)	1500000 (1000000-1900000)
High-middle SDI	1990	Male	55 to 59	73% (65-80%)	27% (20-35%)	5.6 (3.9-7.7)	13% (8.4-18%)	15% (9.9-21%)	34000 (21000-47000)	1200000 (740000-1600000)
High-middle SDI	1990	Male	60 to 64	74% (65-81%)	26% (19-35%)	5.2 (3.5-7.3)	11% (6-16%)	12% (6.8-19%)	38000 (20000-57000)	1100000 (590000-1600000)
High-middle SDI	1990	Male	65 to 69	71% (62-78%)	29% (22-38%)	4.9 (3.5-6.6)	9% (4.6-14%)	10% (5.1-16%)	26000 (12000-42000)	640000 (310000-990000)
High-middle SDI	1990	Male	70 to 74	69% (60-76%)	31% (24-40%)	4.2 (2.8-5.8)	7% (2.9-11%)	8% (3.2-13%)	17000 (5900-29000)	330000 (120000-550000)
High-middle SDI	1990	Male	75 to 79	65% (55-74%)	35% (26-45%)	3.6 (2.3-5.2)	5.4% (1.4-10%)	6% (1.4-11%)	17000 (2800-33000)	260000 (53000-500000)
High-middle SDI	1990	Male	80 to 84	63% (52-73%)	37% (27-48%)	3.4 (2.2-4.7)	4.9% (1-9.3%)	5.4% (0.93-10%)	12000 (1300-24000)	140000 (22000-270000)
High-middle SDI	1990	Male	85 to 89	60% (47-70%)	40% (30-53%)	3 (1.9-4.4)	4.2% (0.7-8.2%)	4.6% (0.48-9.2%)	5800 (340-12000)	46000 (5900-93000)
High-middle SDI	1990	Male	90 to 94	58% (46-69%)	42% (31-54%)	2.7 (1.6-4.1)	3.4% (0.22-7.3%)	3.6% (-0.1-8.1%)	1700 (-87-3900)	9600 (440-21000)
High-middle SDI	1990	Male	95 plus	58% (43-72%)	42% (28-57%)	2.4 (1.2-4)	2.6% (-0.74-6.7%)	2.6% (-1.3-7.4%)	340 (-180-960)	1300 (-390-3500)
High-middle SDI	1995	Female	All Ages	49% (45-54%)	51% (46-55%)	1.5 (1.2-1.8)	3.7% (2.3-5.2%)	3.5% (1.2-6.1%)	110000 (34000-190000)	3800000 (2300000-5400000)
High-middle SDI	1995	Female	15 to 19	47% (38-57%)	53% (43-62%)	1.4 (0.89-2.2)	3.1% (1.9-4.5%)	6.2% (3.2-9.8%)	790 (430-1200)	83000 (53000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	1995	Female	20 to 24	54% (45-63%)	46% (37-55%)	2 (1.3-3)	4.7% (3.3-6.4%)	7.8% (4.6-12%)	1400 (850-2100)	160000 (110000-220000)
High-middle SDI	1995	Female	25 to 29	57% (48-66%)	43% (34-52%)	2.1 (1.3-2.9)	5.2% (3.9-6.8%)	8.4% (5.4-12%)	1800 (1200-2600)	180000 (130000-240000)
High-middle SDI	1995	Female	30 to 34	61% (52-69%)	39% (31-48%)	2.2 (1.5-3.1)	6.1% (4.6-7.7%)	10% (7.1-13%)	3100 (2200-4100)	260000 (190000-330000)
High-middle SDI	1995	Female	35 to 39	63% (55-71%)	37% (29-45%)	2.4 (1.6-3.4)	7.3% (5.6-9.2%)	12% (9-15%)	5400 (4000-7000)	360000 (270000-450000)
High-middle SDI	1995	Female	40 to 44	64% (56-72%)	36% (28-44%)	2.5 (1.7-3.4)	7.9% (6-10%)	13% (9.6-17%)	7700 (5700-10000)	430000 (320000-540000)
High-middle SDI	1995	Female	45 to 49	65% (56-73%)	35% (27-44%)	2.5 (1.7-3.6)	7.8% (5.6-10%)	12% (8.7-16%)	8700 (6100-12000)	420000 (300000-550000)
High-middle SDI	1995	Female	50 to 54	63% (54-71%)	37% (29-46%)	2.5 (1.6-3.5)	7.2% (4.8-10%)	11% (7.3-15%)	8200 (5400-11000)	340000 (230000-470000)
High-middle SDI	1995	Female	55 to 59	61% (51-70%)	39% (30-49%)	2.2 (1.4-3.3)	6.7% (3.9-9.8%)	9.4% (5.5-14%)	13000 (7800-20000)	470000 (280000-700000)
High-middle SDI	1995	Female	60 to 64	56% (46-67%)	44% (33-54%)	2.1 (1.3-3.2)	4.9% (2-8.4%)	6.8% (2.8-12%)	11000 (3900-19000)	330000 (120000-570000)
High-middle SDI	1995	Female	65 to 69	54% (43-64%)	46% (36-57%)	1.9 (1.2-2.7)	3.7% (0.81-7%)	4.8% (1.1-9.1%)	12000 (1500-25000)	310000 (49000-610000)
High-middle SDI	1995	Female	70 to 74	49% (38-59%)	51% (41-62%)	1.7 (1-2.6)	2.9% (-0.15-6.5%)	3.7% (-0.14-8.2%)	11000 (-2700-26000)	210000 (-41000-510000)
High-middle SDI	1995	Female	75 to 79	42% (31-54%)	58% (46-69%)	1.4 (0.82-2.3)	2.1% (-0.58-5.5%)	2.6% (-0.77-6.7%)	7100 (-4200-21000)	110000 (-54000-320000)
High-middle SDI	1995	Female	80 to 84	37% (27-49%)	63% (51-73%)	1.3 (0.75-2)	1.7% (-0.77-4.7%)	2% (-1-5.6%)	7500 (-6400-24000)	87000 (-61000-270000)
High-middle SDI	1995	Female	85 to 89	37% (26-48%)	63% (52-74%)	1.3 (0.74-2.1)	1.7% (-0.87-4.9%)	1.9% (-1.1-5.7%)	5500 (-5200-19000)	46000 (-35000-150000)
High-middle SDI	1995	Female	90 to 94	38% (27-50%)	62% (50-73%)	1.4 (0.74-2.3)	1.7% (-1-5.3%)	1.9% (-1.3-6.2%)	2600 (-2600-9700)	15000 (-12000-51000)
High-middle SDI	1995	Female	95 plus	40% (26-56%)	60% (44-74%)	1.5 (0.68-2.6)	1.8% (-1.5-6.2%)	2% (-1.9-7.2%)	930 (-1000-3500)	3400 (-3100-12000)
High-middle SDI	1995	Male	All Ages	57% (53-61%)	43% (39-47%)	3 (2.5-3.6)	10% (7.6-13%)	9.9% (6.7-13%)	390000 (260000-530000)	1500000 (1100000-1900000)
High-middle SDI	1995	Male	15 to 19	53% (43-63%)	47% (37-57%)	1.9 (1.1-2.8)	6.3% (4-9.3%)	9.2% (5.3-14%)	3000 (1700-4600)	260000 (160000-370000)
High-middle SDI	1995	Male	20 to 24	60% (51-67%)	40% (33-49%)	3.2 (2.1-4.5)	12% (8.1-16%)	15% (9.5-21%)	9400 (6100-13000)	730000 (510000-990000)
High-middle SDI	1995	Male	25 to 29	63% (55-70%)	37% (30-45%)	3.9 (2.6-5.4)	14% (10-18%)	17% (12-23%)	13000 (9200-18000)	950000 (690000-1200000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	1995	Male	30 to 34	67% (59-74%)	33% (26-41%)	4.4 (3-6)	15% (12-19%)	19% (14-24%)	21000 (16000-27000)	1300000 (1000000-1700000)
High-middle SDI	1995	Male	35 to 39	70% (63-76%)	30% (24-37%)	4.8 (3.3-6.6)	17% (13-21%)	21% (15-26%)	32000 (23000-41000)	1800000 (1300000-2300000)
High-middle SDI	1995	Male	40 to 44	72% (65-78%)	28% (22-35%)	5.2 (3.6-7.1)	18% (13-22%)	21% (16-27%)	41000 (30000-54000)	2000000 (1500000-2600000)
High-middle SDI	1995	Male	45 to 49	74% (66-80%)	26% (20-34%)	5.7 (4-7.9)	18% (13-22%)	21% (15-27%)	44000 (31000-57000)	1900000 (1400000-2500000)
High-middle SDI	1995	Male	50 to 54	72% (65-78%)	28% (22-35%)	5.7 (3.9-7.8)	15% (11-20%)	18% (12-24%)	37000 (25000-49000)	1400000 (960000-1800000)
High-middle SDI	1995	Male	55 to 59	74% (66-80%)	26% (20-34%)	5.2 (3.6-7.1)	14% (8.8-19%)	16% (10-22%)	54000 (33000-75000)	1800000 (1100000-2500000)
High-middle SDI	1995	Male	60 to 64	71% (63-79%)	29% (21-37%)	4.9 (3.4-7)	11% (5.9-16%)	12% (6.7-18%)	42000 (22000-64000)	1200000 (630000-1800000)
High-middle SDI	1995	Male	65 to 69	73% (64-80%)	27% (20-36%)	4.4 (3.1-6)	8.6% (4-13%)	9.7% (4.4-15%)	39000 (16000-65000)	940000 (400000-1500000)
High-middle SDI	1995	Male	70 to 74	69% (60-76%)	31% (24-40%)	4 (2.8-5.6)	7% (2.7-12%)	7.9% (3-13%)	24000 (7300-42000)	470000 (150000-810000)
High-middle SDI	1995	Male	75 to 79	65% (55-73%)	35% (27-45%)	3.4 (2.2-4.9)	5.2% (1.6-9.5%)	5.8% (1.6-11%)	14000 (2300-27000)	210000 (45000-400000)
High-middle SDI	1995	Male	80 to 84	64% (52-73%)	36% (27-48%)	3.1 (2.1-4.4)	4.5% (0.9-8.6%)	4.9% (0.79-9.6%)	12000 (410-24000)	130000 (13000-270000)
High-middle SDI	1995	Male	85 to 89	59% (48-69%)	41% (31-52%)	2.8 (1.8-4.1)	3.9% (0.56-7.8%)	4.2% (0.37-8.7%)	6300 (-180-14000)	51000 (2700-110000)
High-middle SDI	1995	Male	90 to 94	58% (47-69%)	42% (31-53%)	2.6 (1.6-4)	3.3% (0.21-7%)	3.5% (-0.048-7.9%)	2000 (-230-4700)	11000 (-35-25000)
High-middle SDI	1995	Male	95 plus	58% (43-71%)	42% (29-57%)	2.3 (1.2-3.9)	2.7% (-0.57-6.7%)	2.8% (-1.1-7.4%)	400 (-190-1100)	1600 (-430-4000)
High-middle SDI	2000	Female	All Ages	48% (44-52%)	52% (48-56%)	1.5 (1.2-1.8)	3.7% (2.3-5.1%)	3.4% (1.2-5.9%)	110000 (35000-190000)	3900000 (2400000-5400000)
High-middle SDI	2000	Female	15 to 19	47% (37-57%)	53% (43-63%)	1.4 (0.87-2.2)	3.1% (2-4.5%)	6.4% (3.4-10%)	800 (450-1200)	85000 (54000-120000)
High-middle SDI	2000	Female	20 to 24	52% (43-61%)	48% (39-57%)	2 (1.3-2.9)	4.8% (3.4-6.4%)	8.1% (4.9-12%)	1600 (980-2300)	170000 (120000-230000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2000	Female	25 to 29	56% (47-64%)	44% (36-53%)	2 (1.3-2.9)	5.3% (4-6.8%)	8.8% (5.8-12%)	2000 (1400-2800)	200000 (150000-270000)
High-middle SDI	2000	Female	30 to 34	57% (48-66%)	43% (34-52%)	2.1 (1.4-3)	5.7% (4.4-7.2%)	9.8% (7-13%)	2800 (2000-3700)	230000 (180000-300000)
High-middle SDI	2000	Female	35 to 39	60% (51-68%)	40% (32-49%)	2.3 (1.6-3.3)	6.9% (5.4-8.6%)	12% (9-15%)	4900 (3700-6300)	330000 (260000-420000)
High-middle SDI	2000	Female	40 to 44	62% (54-70%)	38% (30-46%)	2.5 (1.7-3.4)	7.9% (6.2-9.9%)	13% (10-17%)	7800 (5900-9900)	440000 (340000-560000)
High-middle SDI	2000	Female	45 to 49	63% (54-71%)	37% (29-46%)	2.4 (1.7-3.4)	7.8% (5.8-10%)	12% (9.1-16%)	9900 (7200-13000)	480000 (360000-620000)
High-middle SDI	2000	Female	50 to 54	61% (52-70%)	39% (30-48%)	2.3 (1.5-3.3)	7.4% (5.1-10%)	11% (7.7-15%)	10000 (7200-14000)	430000 (300000-590000)
High-middle SDI	2000	Female	55 to 59	56% (46-66%)	44% (34-54%)	2.2 (1.4-3.2)	6.3% (3.8-9.3%)	9.3% (5.6-14%)	9200 (5500-14000)	330000 (200000-480000)
High-middle SDI	2000	Female	60 to 64	55% (44-66%)	45% (34-56%)	1.9 (1.2-3)	4.9% (2.1-8.4%)	6.8% (2.8-12%)	13000 (4800-22000)	380000 (150000-660000)
High-middle SDI	2000	Female	65 to 69	51% (40-62%)	49% (38-60%)	1.8 (1.1-2.6)	3.7% (0.96-6.8%)	4.9% (1.3-9.1%)	10000 (1700-21000)	260000 (54000-510000)
High-middle SDI	2000	Female	70 to 74	46% (35-57%)	54% (43-65%)	1.5 (0.88-2.3)	2.6% (-0.21-5.9%)	3.3% (-0.28-7.5%)	11000 (-3000-28000)	220000 (-43000-540000)
High-middle SDI	2000	Female	75 to 79	39% (28-51%)	61% (49-72%)	1.3 (0.71-2)	2% (-0.65-5.3%)	2.4% (-0.82-6.5%)	9000 (-5300-27000)	140000 (-68000-410000)
High-middle SDI	2000	Female	80 to 84	37% (26-49%)	63% (51-74%)	1.2 (0.71-1.9)	1.8% (-0.6-4.7%)	2.1% (-0.79-5.7%)	6400 (-4200-20000)	74000 (-39000-220000)
High-middle SDI	2000	Female	85 to 89	37% (26-48%)	63% (52-74%)	1.2 (0.69-1.9)	1.5% (-0.9-4.6%)	1.6% (-1.2-5.3%)	5200 (-5700-19000)	44000 (-38000-150000)
High-middle SDI	2000	Female	90 to 94	38% (27-49%)	62% (51-73%)	1.3 (0.7-2.1)	1.5% (-1.1-5%)	1.7% (-1.4-5.9%)	2800 (-3300-11000)	16000 (-15000-57000)
High-middle SDI	2000	Female	95 plus	40% (26-55%)	60% (45-74%)	1.4 (0.65-2.5)	1.7% (-1.4-6.1%)	1.9% (-1.8-7.1%)	1100 (-1200-4200)	3900 (-3600-14000)
High-middle SDI	2000	Male	All Ages	57% (53-60%)	43% (40-47%)	3 (2.5-3.6)	10% (7.9-13%)	10% (6.9-13%)	410000 (280000-540000)	1600000 (1200000-1900000)
High-middle SDI	2000	Male	15 to 19	55% (45-64%)	45% (36-55%)	1.8 (1.1-2.7)	6.5% (4.1-9.4%)	9.5% (5.3-14%)	3000 (1700-4600)	260000 (160000-370000)
High-middle SDI	2000	Male	20 to 24	58% (50-65%)	42% (35-50%)	3 (2-4.3)	12% (8.1-15%)	15% (9.6-21%)	10000 (6800-14000)	790000 (560000-1100000)
High-middle SDI	2000	Male	25 to 29	61% (54-68%)	39% (32-46%)	3.7 (2.6-5.2)	14% (10-17%)	17% (12-23%)	15000 (10000-19000)	1000000 (760000-1300000)
High-middle SDI	2000	Male	30 to 34	64% (56-70%)	36% (30-44%)	4.2 (3-5.8)	14% (11-18%)	18% (13-23%)	18000 (13000-23000)	1200000 (880000-1400000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2000	Male	35 to 39	68% (61-74%)	32% (26-39%)	4.7 (3.2-6.4)	16% (13-20%)	20% (15-25%)	28000 (21000-36000)	1600000 (1200000-2000000)
High-middle SDI	2000	Male	40 to 44	70% (64-76%)	30% (24-36%)	5.2 (3.6-7)	18% (14-22%)	22% (17-27%)	42000 (32000-53000)	2100000 (1600000-2600000)
High-middle SDI	2000	Male	45 to 49	72% (64-77%)	28% (23-36%)	5.6 (4-7.6)	18% (14-22%)	21% (16-26%)	51000 (38000-64000)	2200000 (1700000-2800000)
High-middle SDI	2000	Male	50 to 54	73% (66-79%)	27% (21-34%)	5.7 (4-7.8)	16% (12-21%)	20% (14-25%)	49000 (35000-64000)	1900000 (1300000-2400000)
High-middle SDI	2000	Male	55 to 59	70% (63-77%)	30% (23-37%)	5.2 (3.7-7)	13% (8.9-18%)	16% (11-21%)	36000 (24000-49000)	1200000 (790000-1600000)
High-middle SDI	2000	Male	60 to 64	72% (64-79%)	28% (21-36%)	4.7 (3.2-6.7)	11% (6.5-16%)	13% (7.3-19%)	51000 (27000-75000)	1400000 (800000-2100000)
High-middle SDI	2000	Male	65 to 69	70% (61-77%)	30% (23-39%)	4.3 (3.1-5.8)	8.9% (4.6-13%)	10% (5.2-16%)	37000 (17000-58000)	890000 (420000-1400000)
High-middle SDI	2000	Male	70 to 74	70% (61-77%)	30% (23-39%)	3.7 (2.5-5.1)	6.8% (2.6-11%)	7.6% (2.8-13%)	32000 (9100-57000)	610000 (200000-1100000)
High-middle SDI	2000	Male	75 to 79	63% (54-72%)	37% (28-46%)	3.2 (2.1-4.6)	5.4% (1.7-9.6%)	6% (1.8-11%)	19000 (3700-36000)	280000 (66000-530000)
High-middle SDI	2000	Male	80 to 84	61% (50-71%)	39% (29-50%)	2.9 (1.9-4.1)	4.6% (1.3-8.3%)	5.1% (1.3-9.3%)	10000 (1500-20000)	110000 (23000-220000)
High-middle SDI	2000	Male	85 to 89	60% (48-70%)	40% (30-52%)	2.6 (1.7-3.8)	3.7% (0.66-7.3%)	4% (0.51-8.1%)	6100 (120-13000)	50000 (4600-100000)
High-middle SDI	2000	Male	90 to 94	58% (46-69%)	42% (31-54%)	2.4 (1.5-3.7)	3.2% (0.19-6.7%)	3.4% (-0.065-7.5%)	2200 (-270-5200)	13000 (-79-28000)
High-middle SDI	2000	Male	95 plus	58% (44-71%)	42% (29-56%)	2.2 (1.1-3.7)	2.7% (-0.46-6.6%)	2.8% (-0.95-7.3%)	490 (-190-1300)	1900 (-400-4700)
High-middle SDI	2005	Female	All Ages	47% (43-51%)	53% (49-57%)	1.6 (1.3-1.9)	4% (2.7-5.4%)	3.7% (1.6-6.1%)	130000 (58000-220000)	4500000 (3000000-6100000)
High-middle SDI	2005	Female	15 to 19	46% (37-55%)	54% (45-63%)	1.5 (0.91-2.2)	3.1% (2.1-4.5%)	6.7% (3.7-10%)	760 (430-1200)	83000 (54000-120000)
High-middle SDI	2005	Female	20 to 24	50% (41-59%)	50% (41-59%)	2 (1.3-2.9)	4.8% (3.5-6.3%)	8.3% (5.3-12%)	1700 (1100-2400)	180000 (130000-250000)
High-middle SDI	2005	Female	25 to 29	54% (45-62%)	46% (38-55%)	2 (1.4-2.9)	5.5% (4.2-6.9%)	9% (6.2-12%)	2600 (1800-3400)	240000 (180000-310000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2005	Female	30 to 34	56% (47-65%)	44% (35-53%)	2.2 (1.5-3.1)	6.1% (4.8-7.6%)	10% (7.5-13%)	3800 (2800-4900)	290000 (230000-370000)
High-middle SDI	2005	Female	35 to 39	57% (49-66%)	43% (34-51%)	2.4 (1.7-3.3)	6.9% (5.4-8.6%)	12% (8.9-15%)	5400 (4100-7000)	350000 (270000-440000)
High-middle SDI	2005	Female	40 to 44	60% (53-68%)	40% (32-47%)	2.6 (1.8-3.5)	8.1% (6.3-10%)	13% (10-17%)	8800 (6800-11000)	480000 (380000-610000)
High-middle SDI	2005	Female	45 to 49	62% (53-69%)	38% (31-47%)	2.5 (1.8-3.5)	8.3% (6.4-11%)	13% (9.9-17%)	12000 (9100-15000)	570000 (440000-720000)
High-middle SDI	2005	Female	50 to 54	59% (50-68%)	41% (32-50%)	2.3 (1.6-3.3)	7.9% (5.6-10%)	12% (8.4-16%)	14000 (9900-18000)	570000 (410000-750000)
High-middle SDI	2005	Female	55 to 59	56% (46-65%)	44% (35-54%)	2.2 (1.4-3.2)	6.9% (4.4-9.8%)	10% (6.4-14%)	14000 (8700-19000)	470000 (310000-670000)
High-middle SDI	2005	Female	60 to 64	51% (41-61%)	49% (39-59%)	2 (1.2-3)	5.1% (2.4-8.2%)	7.2% (3.6-12%)	9800 (4600-16000)	300000 (140000-480000)
High-middle SDI	2005	Female	65 to 69	49% (39-59%)	51% (41-61%)	1.7 (1.1-2.5)	4% (1.3-7%)	5.3% (1.8-9.3%)	14000 (3900-25000)	340000 (110000-600000)
High-middle SDI	2005	Female	70 to 74	42% (32-53%)	58% (47-68%)	1.4 (0.87-2.2)	2.9% (0.28-6%)	3.8% (0.4-7.7%)	11000 (-180-25000)	220000 (5200-480000)
High-middle SDI	2005	Female	75 to 79	36% (26-48%)	64% (52-74%)	1.2 (0.69-1.9)	2.1% (-0.3-5.2%)	2.6% (-0.41-6.4%)	12000 (-3600-32000)	190000 (-42000-480000)
High-middle SDI	2005	Female	80 to 84	35% (25-47%)	65% (53-75%)	1.2 (0.7-1.9)	2% (-0.33-5%)	2.4% (-0.46-5.9%)	10000 (-3500-28000)	120000 (-30000-310000)
High-middle SDI	2005	Female	85 to 89	36% (25-47%)	64% (53-75%)	1.2 (0.69-1.9)	1.8% (-0.41-4.8%)	2.1% (-0.57-5.7%)	6200 (-2500-18000)	50000 (-16000-140000)
High-middle SDI	2005	Female	90 to 94	37% (26-48%)	63% (52-74%)	1.3 (0.68-2.1)	1.7% (-0.8-5.1%)	1.9% (-1.1-6%)	3800 (-2600-13000)	21000 (-12000-66000)
High-middle SDI	2005	Female	95 plus	39% (26-54%)	61% (46-74%)	1.4 (0.64-2.4)	1.9% (-1.1-6.2%)	2.1% (-1.5-7.1%)	1500 (-1100-5100)	5500 (-3300-18000)
High-middle SDI	2005	Male	All Ages	56% (52-60%)	44% (40-48%)	3.2 (2.7-3.8)	11% (8.5-13%)	10% (7.5-13%)	470000 (340000-600000)	1700000 (1400000-2100000)
High-middle SDI	2005	Male	15 to 19	53% (44-62%)	47% (38-56%)	1.9 (1.2-2.8)	6.5% (4.2-9.2%)	9.9% (5.6-15%)	2800 (1600-4200)	240000 (150000-350000)
High-middle SDI	2005	Male	20 to 24	55% (48-62%)	45% (38-52%)	3.1 (2.1-4.3)	11% (8.2-15%)	15% (10-21%)	10000 (7000-14000)	790000 (570000-1000000)
High-middle SDI	2005	Male	25 to 29	59% (53-66%)	41% (34-47%)	3.9 (2.7-5.3)	13% (10-17%)	17% (12-23%)	17000 (12000-22000)	1200000 (880000-1500000)
High-middle SDI	2005	Male	30 to 34	61% (55-68%)	39% (32-45%)	4.4 (3.1-5.9)	14% (11-17%)	18% (14-23%)	22000 (17000-28000)	1400000 (1100000-1700000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2005	Male	35 to 39	63% (56-70%)	37% (30-44%)	4.7 (3.2-6.4)	15% (12-18%)	19% (14-24%)	28000 (21000-34000)	1500000 (1200000-1900000)
High-middle SDI	2005	Male	40 to 44	67% (61-73%)	33% (27-39%)	5.2 (3.6-7)	17% (14-20%)	21% (16-25%)	44000 (34000-54000)	2100000 (1700000-2600000)
High-middle SDI	2005	Male	45 to 49	70% (63-76%)	30% (24-37%)	5.8 (4.2-7.9)	18% (14-22%)	22% (17-26%)	60000 (46000-74000)	2600000 (2000000-3100000)
High-middle SDI	2005	Male	50 to 54	71% (64-76%)	29% (24-36%)	5.8 (4.1-7.9)	17% (13-21%)	20% (15-25%)	65000 (48000-81000)	2400000 (1800000-3100000)
High-middle SDI	2005	Male	55 to 59	71% (64-77%)	29% (23-36%)	5.5 (3.9-7.4)	15% (10-19%)	17% (12-22%)	54000 (37000-70000)	1800000 (1200000-2300000)
High-middle SDI	2005	Male	60 to 64	68% (61-75%)	32% (25-39%)	4.9 (3.3-6.9)	11% (6.8-16%)	13% (8-18%)	37000 (21000-53000)	1100000 (620000-1500000)
High-middle SDI	2005	Male	65 to 69	70% (62-77%)	30% (23-38%)	4.4 (3.1-5.9)	9.7% (5.2-14%)	11% (5.8-16%)	48000 (23000-72000)	1100000 (570000-1700000)
High-middle SDI	2005	Male	70 to 74	66% (58-73%)	34% (27-42%)	3.7 (2.6-5.1)	7.2% (3.2-11%)	8.2% (3.6-13%)	32000 (11000-54000)	620000 (230000-1000000)
High-middle SDI	2005	Male	75 to 79	64% (54-73%)	36% (27-46%)	3.2 (2.1-4.6)	5.5% (1.6-9.9%)	6.2% (1.7-11%)	25000 (4000-49000)	380000 (80000-730000)
High-middle SDI	2005	Male	80 to 84	61% (50-71%)	39% (29-50%)	2.9 (1.9-4.2)	4.9% (1.6-8.7%)	5.5% (1.6-9.9%)	14000 (3000-27000)	160000 (40000-300000)
High-middle SDI	2005	Male	85 to 89	57% (46-67%)	43% (33-54%)	2.5 (1.6-3.7)	3.9% (1.1-7.2%)	4.3% (1-8.1%)	6000 (950-12000)	48000 (11000-92000)
High-middle SDI	2005	Male	90 to 94	58% (47-68%)	42% (32-53%)	2.4 (1.4-3.6)	3.2% (0.36-6.5%)	3.4% (0.1-7.3%)	2400 (-100-5400)	14000 (790-29000)
High-middle SDI	2005	Male	95 plus	58% (43-71%)	42% (29-57%)	2.2 (1.1-3.7)	2.8% (-0.32-6.6%)	2.9% (-0.77-7.3%)	630 (-180-1600)	2400 (-300-5800)
High-middle SDI	2010	Female	All Ages	47% (43-51%)	53% (49-57%)	1.6 (1.3-2)	3.8% (2.5-5.2%)	3.6% (1.5-6.1%)	120000 (51000-210000)	4100000 (2700000-5600000)
High-middle SDI	2010	Female	15 to 19	44% (34-53%)	56% (47-66%)	1.4 (0.84-2.1)	2.7% (1.8-4%)	6.3% (3.3-9.9%)	520 (280-820)	61000 (38000-89000)
High-middle SDI	2010	Female	20 to 24	48% (39-58%)	52% (42-61%)	1.9 (1.2-2.8)	4.5% (3.3-6%)	7.9% (5-11%)	1300 (820-1800)	160000 (110000-220000)
High-middle SDI	2010	Female	25 to 29	51% (41-60%)	49% (40-59%)	2 (1.3-2.8)	5% (3.8-6.2%)	8.1% (5.6-11%)	2100 (1400-2700)	220000 (160000-280000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2010	Female	30 to 34	54% (45-63%)	46% (37-55%)	2.1 (1.4-3)	5.5% (4.4-6.9%)	9.4% (6.8-12%)	3300 (2400-4200)	280000 (210000-350000)
High-middle SDI	2010	Female	35 to 39	56% (48-65%)	44% (35-52%)	2.4 (1.6-3.4)	6.4% (5-7.9%)	11% (8.5-14%)	4800 (3600-6000)	330000 (250000-410000)
High-middle SDI	2010	Female	40 to 44	57% (49-66%)	43% (34-51%)	2.5 (1.7-3.5)	6.8% (5.3-8.6%)	12% (9.2-15%)	6200 (4700-7900)	360000 (280000-450000)
High-middle SDI	2010	Female	45 to 49	59% (50-67%)	41% (33-50%)	2.5 (1.7-3.6)	7.2% (5.5-9.3%)	12% (9.2-16%)	8800 (6600-11000)	440000 (330000-560000)
High-middle SDI	2010	Female	50 to 54	59% (50-67%)	41% (33-50%)	2.4 (1.6-3.4)	7.3% (5.3-9.7%)	11% (8.3-15%)	12000 (8500-16000)	500000 (360000-670000)
High-middle SDI	2010	Female	55 to 59	56% (45-66%)	44% (34-55%)	2.2 (1.4-3.3)	6.6% (4.3-9.3%)	9.9% (6.5-14%)	14000 (8800-19000)	490000 (320000-680000)
High-middle SDI	2010	Female	60 to 64	53% (42-64%)	47% (36-58%)	2.1 (1.3-3.2)	5.2% (2.7-8.3%)	7.5% (3.9-12%)	12000 (5700-19000)	360000 (180000-570000)
High-middle SDI	2010	Female	65 to 69	50% (39-61%)	50% (39-61%)	1.9 (1.2-2.7)	4% (1.6-7%)	5.7% (2.2-9.7%)	9400 (3300-17000)	240000 (86000-420000)
High-middle SDI	2010	Female	70 to 74	46% (35-57%)	54% (43-65%)	1.6 (0.95-2.5)	3.2% (0.39-6.6%)	4.1% (0.48-8.5%)	13000 (610-29000)	270000 (23000-580000)
High-middle SDI	2010	Female	75 to 79	39% (28-51%)	61% (49-72%)	1.3 (0.76-2.1)	2.4% (-0.15-5.6%)	3% (-0.25-7%)	11000 (-2100-28000)	170000 (-22000-430000)
High-middle SDI	2010	Female	80 to 84	36% (25-48%)	64% (52-75%)	1.2 (0.72-1.9)	2% (-0.3-5.1%)	2.4% (-0.45-6.2%)	13000 (-3400-33000)	140000 (-27000-370000)
High-middle SDI	2010	Female	85 to 89	36% (25-47%)	64% (53-75%)	1.2 (0.69-2)	1.9% (-0.4-5%)	2.2% (-0.59-5.9%)	8800 (-3300-25000)	71000 (-19000-190000)
High-middle SDI	2010	Female	90 to 94	37% (27-49%)	63% (51-73%)	1.3 (0.68-2.1)	1.7% (-0.59-4.9%)	2% (-0.83-5.8%)	3600 (-1900-11000)	20000 (-8300-60000)
High-middle SDI	2010	Female	95 plus	40% (26-54%)	60% (46-74%)	1.4 (0.64-2.4)	1.9% (-0.95-5.8%)	2.1% (-1.3-6.8%)	1700 (-1100-5700)	6300 (-3300-20000)
High-middle SDI	2010	Male	All Ages	54% (50-58%)	46% (42-50%)	3.3 (2.7-3.9)	9.9% (7.9-12%)	9.7% (7-12%)	400000 (290000-530000)	1500000 (1100000-1800000)
High-middle SDI	2010	Male	15 to 19	49% (39-58%)	51% (42-61%)	1.8 (1.1-2.7)	5.5% (3.5-8%)	9.2% (5.2-14%)	1700 (960-2600)	160000 (99000-230000)
High-middle SDI	2010	Male	20 to 24	53% (45-60%)	47% (40-55%)	3 (2-4.3)	11% (7.6-14%)	15% (9.6-20%)	7500 (5000-10000)	610000 (440000-810000)
High-middle SDI	2010	Male	25 to 29	56% (49-62%)	44% (38-51%)	3.8 (2.5-5.3)	12% (9.3-15%)	16% (12-22%)	13000 (9600-17000)	950000 (720000-1200000)
High-middle SDI	2010	Male	30 to 34	59% (52-66%)	41% (34-48%)	4.3 (3-6)	13% (10-16%)	17% (13-22%)	19000 (14000-23000)	1200000 (940000-1500000)
High-middle SDI	2010	Male	35 to 39	60% (53-67%)	40% (33-47%)	4.6 (3.1-6.5)	14% (11-17%)	18% (14-23%)	23000 (18000-29000)	1300000 (1000000-1600000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2010	Male	40 to 44	62% (55-69%)	38% (31-45%)	4.9 (3.4-6.8)	14% (11-17%)	19% (15-23%)	28000 (22000-35000)	1400000 (1100000-1700000)
High-middle SDI	2010	Male	45 to 49	66% (58-72%)	34% (28-42%)	5.6 (3.9-7.7)	16% (13-19%)	20% (16-25%)	41000 (32000-51000)	1800000 (1400000-2200000)
High-middle SDI	2010	Male	50 to 54	68% (61-74%)	32% (26-39%)	5.8 (3.9-8)	16% (12-20%)	20% (15-24%)	53000 (40000-66000)	2000000 (1500000-2500000)
High-middle SDI	2010	Male	55 to 59	68% (60-75%)	32% (25-40%)	5.3 (3.7-7.3)	14% (10-18%)	17% (12-22%)	53000 (38000-69000)	1800000 (1300000-2300000)
High-middle SDI	2010	Male	60 to 64	69% (61-76%)	31% (24-39%)	5 (3.4-7.1)	12% (7.5-16%)	14% (8.8-19%)	45000 (28000-62000)	1300000 (800000-1800000)
High-middle SDI	2010	Male	65 to 69	66% (58-73%)	34% (27-42%)	4.4 (3.1-5.9)	9.3% (5.5-13%)	11% (6.4-16%)	30000 (16000-44000)	730000 (410000-1100000)
High-middle SDI	2010	Male	70 to 74	66% (58-73%)	34% (27-42%)	3.7 (2.5-5.2)	7.4% (3.4-12%)	8.5% (3.8-14%)	35000 (14000-58000)	690000 (280000-1100000)
High-middle SDI	2010	Male	75 to 79	61% (51-70%)	39% (30-49%)	3.2 (2.1-4.7)	5.7% (2.1-9.8%)	6.5% (2.3-11%)	24000 (6400-43000)	360000 (110000-650000)
High-middle SDI	2010	Male	80 to 84	62% (50-71%)	38% (29-50%)	2.9 (1.9-4.2)	4.9% (1.5-8.9%)	5.5% (1.4-10%)	18000 (3300-35000)	210000 (48000-390000)
High-middle SDI	2010	Male	85 to 89	58% (47-68%)	42% (32-53%)	2.7 (1.7-3.9)	4.2% (1.2-7.7%)	4.7% (1.2-8.7%)	8800 (1600-17000)	71000 (17000-130000)
High-middle SDI	2010	Male	90 to 94	56% (46-67%)	44% (33-54%)	2.4 (1.4-3.7)	3.3% (0.57-6.5%)	3.6% (0.36-7.3%)	2600 (110-5500)	15000 (1900-30000)
High-middle SDI	2010	Male	95 plus	58% (44-71%)	42% (29-56%)	2.3 (1.2-3.8)	2.9% (-0.13-6.7%)	3.1% (-0.56-7.4%)	730 (-150-1800)	2800 (-170-6400)
High-middle SDI	2016	Female	All Ages	46% (42-51%)	54% (49-58%)	1.7 (1.2-2.2)	3.7% (2.3-5.4%)	3.5% (1.2-6.4%)	130000 (41000-250000)	4000000 (2200000-6400000)
High-middle SDI	2016	Female	15 to 19	42% (32-52%)	58% (48-68%)	1.4 (0.75-2.2)	2.4% (1.5-3.6%)	6% (3-9.8%)	380 (180-670)	47000 (28000-73000)
High-middle SDI	2016	Female	20 to 24	47% (38-57%)	53% (43-62%)	1.9 (1.1-3.1)	4.1% (2.9-5.6%)	7.7% (4.5-11%)	870 (450-1500)	120000 (75000-170000)
High-middle SDI	2016	Female	25 to 29	51% (41-60%)	49% (40-59%)	2.1 (1.2-3.1)	4.9% (3.7-6.4%)	8.3% (5.5-12%)	1900 (970-3100)	210000 (140000-300000)
High-middle SDI	2016	Female	30 to 34	51% (41-61%)	49% (39-59%)	2.1 (1.3-3.1)	5.2% (3.9-6.8%)	8.9% (6.3-12%)	3100 (1700-5300)	270000 (170000-410000)
High-middle SDI	2016	Female	35 to 39	53% (44-62%)	47% (38-56%)	2.3 (1.4-3.5)	5.7% (4.2-7.6%)	10% (7.6-14%)	4600 (2400-7900)	320000 (200000-500000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2016	Female	40 to 44	55% (46-64%)	45% (36-54%)	2.5 (1.5-3.7)	6.3% (4.5-8.4%)	12% (8.5-15%)	6000 (3300-9800)	350000 (220000-540000)
High-middle SDI	2016	Female	45 to 49	56% (46-65%)	44% (35-54%)	2.5 (1.5-3.7)	6.1% (4.3-8.5%)	11% (8-15%)	6800 (4000-11000)	350000 (220000-530000)
High-middle SDI	2016	Female	50 to 54	57% (47-66%)	43% (34-53%)	2.4 (1.5-3.7)	6.6% (4.5-9.2%)	11% (7.6-15%)	10000 (5900-16000)	440000 (270000-690000)
High-middle SDI	2016	Female	55 to 59	55% (45-66%)	45% (34-55%)	2.3 (1.4-3.6)	6.3% (3.9-9.2%)	9.8% (6.2-14%)	13000 (7300-22000)	480000 (280000-760000)
High-middle SDI	2016	Female	60 to 64	54% (42-65%)	46% (35-58%)	2.2 (1.2-3.4)	5.1% (2.5-8.4%)	7.6% (3.7-12%)	14000 (6100-24000)	420000 (200000-730000)
High-middle SDI	2016	Female	65 to 69	52% (41-63%)	48% (37-59%)	2 (1.2-3)	4.1% (1.4-7.3%)	5.8% (2.1-10%)	12000 (3700-23000)	300000 (100000-560000)
High-middle SDI	2016	Female	70 to 74	47% (36-58%)	53% (42-64%)	1.8 (0.98-2.8)	3.3% (0.44-6.9%)	4.5% (0.67-9.4%)	9800 (730-22000)	200000 (20000-450000)
High-middle SDI	2016	Female	75 to 79	42% (30-55%)	58% (45-70%)	1.5 (0.78-2.5)	2.5% (-0.32-6.3%)	3.1% (-0.51-8%)	13000 (-3300-38000)	210000 (-40000-580000)
High-middle SDI	2016	Female	80 to 84	38% (27-50%)	62% (50-73%)	1.4 (0.74-2.3)	2.3% (-0.21-5.7%)	2.8% (-0.31-7%)	12000 (-2700-33000)	140000 (-23000-370000)
High-middle SDI	2016	Female	85 to 89	37% (25-49%)	63% (51-75%)	1.3 (0.7-2.2)	1.9% (-0.5-5.4%)	2.2% (-0.72-6.5%)	11000 (-5200-34000)	91000 (-32000-260000)
High-middle SDI	2016	Female	90 to 94	38% (27-50%)	62% (50-73%)	1.4 (0.72-2.5)	2% (-0.62-5.8%)	2.2% (-0.9-6.9%)	6000 (-3100-19000)	34000 (-13000-100000)
High-middle SDI	2016	Female	95 plus	41% (27-56%)	59% (44-73%)	1.5 (0.67-2.8)	2.1% (-0.84-6.4%)	2.4% (-1.2-7.6%)	2300 (-1200-7400)	8400 (-3500-26000)
High-middle SDI	2016	Male	All Ages	52% (48-56%)	48% (44-52%)	3.1 (2.3-4.2)	9.3% (7-12%)	9.1% (6.1-12%)	380000 (220000-570000)	13000000 (8800000-19000000)
High-middle SDI	2016	Male	15 to 19	46% (36-55%)	54% (45-64%)	1.6 (0.91-2.6)	5% (3.1-7.5%)	8.6% (4.6-14%)	1300 (640-2200)	120000 (72000-190000)
High-middle SDI	2016	Male	20 to 24	51% (42-58%)	49% (42-58%)	2.8 (1.7-4.3)	9.8% (6.8-13%)	14% (8.6-20%)	5100 (2900-7900)	430000 (270000-630000)
High-middle SDI	2016	Male	25 to 29	55% (47-62%)	45% (38-53%)	3.7 (2.3-5.4)	12% (8.9-15%)	16% (11-22%)	12000 (7100-18000)	880000 (580000-1200000)
High-middle SDI	2016	Male	30 to 34	56% (47-63%)	44% (37-53%)	4 (2.6-5.9)	12% (9.3-15%)	16% (12-21%)	18000 (11000-27000)	1200000 (750000-1700000)
High-middle SDI	2016	Male	35 to 39	58% (50-65%)	42% (35-50%)	4.4 (2.7-6.5)	13% (9.5-16%)	17% (12-22%)	21000 (13000-32000)	1200000 (780000-1800000)
High-middle SDI	2016	Male	40 to 44	60% (52-67%)	40% (33-48%)	4.7 (3-6.9)	13% (10-17%)	18% (14-23%)	26000 (16000-39000)	1300000 (860000-1900000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
High-middle SDI	2016	Male	45 to 49	60% (52-67%)	40% (33-48%)	5 (3.2-7.3)	13% (10-17%)	18% (14-22%)	30000 (19000-45000)	1400000 (880000-1900000)
High-middle SDI	2016	Male	50 to 54	64% (56-71%)	36% (29-44%)	5.3 (3.3-7.8)	14% (11-18%)	18% (14-22%)	44000 (27000-65000)	1700000 (1100000-2500000)
High-middle SDI	2016	Male	55 to 59	65% (57-72%)	35% (28-43%)	5 (3.2-7.3)	13% (9.3-17%)	16% (12-21%)	50000 (31000-75000)	1700000 (1100000-2500000)
High-middle SDI	2016	Male	60 to 64	66% (57-73%)	34% (27-43%)	4.7 (2.9-7.1)	11% (7-15%)	13% (8.3-18%)	50000 (28000-76000)	1400000 (810000-2200000)
High-middle SDI	2016	Male	65 to 69	67% (58-74%)	33% (26-42%)	4.3 (2.8-6.3)	9.3% (5.3-13%)	11% (6.3-16%)	38000 (19000-61000)	930000 (480000-1400000)
High-middle SDI	2016	Male	70 to 74	63% (55-71%)	37% (29-45%)	3.7 (2.3-5.5)	7% (3.3-11%)	8.3% (3.9-13%)	24000 (9300-41000)	470000 (190000-780000)
High-middle SDI	2016	Male	75 to 79	63% (53-72%)	37% (28-47%)	3.2 (1.9-4.9)	5.6% (1.7-10%)	6.4% (1.8-11%)	26000 (5300-50000)	400000 (97000-750000)
High-middle SDI	2016	Male	80 to 84	61% (49-70%)	39% (30-51%)	3 (1.8-4.6)	5% (1.5-8.9%)	5.6% (1.5-10%)	18000 (3400-35000)	210000 (48000-390000)
High-middle SDI	2016	Male	85 to 89	58% (46-68%)	42% (32-54%)	2.6 (1.5-4.1)	3.9% (0.6-7.7%)	4.3% (0.44-8.8%)	11000 (360-24000)	94000 (10000-190000)
High-middle SDI	2016	Male	90 to 94	57% (46-68%)	43% (32-54%)	2.5 (1.4-4.1)	3.5% (0.42-7.1%)	3.8% (0.22-8.1%)	4200 (-26-9200)	24000 (1800-50000)
High-middle SDI	2016	Male	95 plus	57% (43-69%)	43% (31-57%)	2.2 (1.1-3.9)	2.9% (-0.15-6.6%)	3.1% (-0.57-7.4%)	930 (-190-2300)	3500 (-250-8100)
Middle SDI	1990	Female	All Ages	16% (14-19%)	84% (81-86%)	0.33 (0.26-0.4)	1.2% (0.84-1.5%)	1.8% (1.1-2.5%)	110000 (67000-160000)	3800000 (2800000-5000000)
Middle SDI	1990	Female	15 to 19	14% (9.1-20%)	86% (80-91%)	0.26 (0.14-0.45)	0.97% (0.59-1.5%)	1.4% (0.71-2.4%)	1300 (630-2200)	140000 (87000-220000)
Middle SDI	1990	Female	20 to 24	17% (11-25%)	83% (75-89%)	0.37 (0.2-0.62)	1.5% (0.99-2.2%)	1.8% (0.87-3%)	2300 (1100-3900)	280000 (180000-410000)
Middle SDI	1990	Female	25 to 29	19% (13-27%)	81% (73-87%)	0.45 (0.25-0.72)	1.8% (1.2-2.5%)	2.2% (1.1-3.5%)	2900 (1600-4600)	300000 (200000-420000)
Middle SDI	1990	Female	30 to 34	21% (13-30%)	79% (70-87%)	0.51 (0.28-0.82)	1.9% (1.3-2.6%)	2.5% (1.5-3.9%)	3500 (2100-5200)	290000 (200000-410000)
Middle SDI	1990	Female	35 to 39	22% (14-31%)	78% (69-86%)	0.52 (0.28-0.84)	1.9% (1.3-2.7%)	2.9% (1.8-4.3%)	4500 (2800-6600)	310000 (210000-430000)
Middle SDI	1990	Female	40 to 44	23% (15-32%)	77% (68-85%)	0.58 (0.31-0.93)	2.2% (1.5-3.1%)	3.5% (2.2-5%)	5200 (3300-7600)	290000 (200000-410000)
Middle SDI	1990	Female	45 to 49	22% (14-31%)	78% (69-86%)	0.62 (0.32-1)	2.4% (1.5-3.5%)	3.7% (2.3-5.5%)	6400 (4000-9600)	300000 (190000-430000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	1990	Female	50 to 54	22% (14-31%)	78% (69-86%)	0.63 (0.33-1.1)	2.5% (1.5-3.7%)	3.8% (2.3-5.6%)	8900 (5400-13000)	350000 (210000-510000)
Middle SDI	1990	Female	55 to 59	21% (13-30%)	79% (70-87%)	0.62 (0.33-1.1)	2.5% (1.5-3.8%)	3.6% (2.1-5.5%)	11000 (6800-17000)	370000 (220000-570000)
Middle SDI	1990	Female	60 to 64	21% (13-31%)	79% (69-87%)	0.63 (0.32-1.1)	2.3% (1.2-3.7%)	3.1% (1.7-5%)	13000 (7100-20000)	350000 (190000-570000)
Middle SDI	1990	Female	65 to 69	21% (13-29%)	79% (71-87%)	0.59 (0.32-0.98)	2.1% (1.1-3.3%)	2.7% (1.5-4.3%)	14000 (7600-22000)	310000 (170000-490000)
Middle SDI	1990	Female	70 to 74	20% (13-28%)	80% (72-87%)	0.5 (0.25-0.87)	1.6% (0.7-2.8%)	2% (0.89-3.5%)	13000 (6100-22000)	240000 (110000-410000)
Middle SDI	1990	Female	75 to 79	18% (11-27%)	82% (73-89%)	0.46 (0.22-0.82)	1.2% (0.35-2.4%)	1.5% (0.45-2.9%)	9900 (3100-19000)	140000 (42000-270000)
Middle SDI	1990	Female	80 to 84	17% (11-25%)	83% (75-89%)	0.46 (0.22-0.8)	1% (0.17-2.1%)	1.2% (0.21-2.5%)	6500 (1000-14000)	69000 (11000-140000)
Middle SDI	1990	Female	85 to 89	18% (11-27%)	82% (73-89%)	0.47 (0.23-0.85)	0.84% (0.019-2%)	0.99% (0.0038-2.4%)	3700 (-31-8800)	28000 (500-65000)
Middle SDI	1990	Female	90 to 94	20% (13-29%)	80% (71-87%)	0.54 (0.25-1)	0.71% (-0.19-2%)	0.81% (-0.26-2.4%)	1300 (-440-3800)	6900 (-1900-20000)
Middle SDI	1990	Female	95 plus	23% (13-36%)	77% (64-87%)	0.64 (0.24-1.3)	0.72% (-0.44-2.5%)	0.77% (-0.6-2.9%)	350 (-270-1300)	1300 (-790-4500)
Middle SDI	1990	Male	All Ages	40% (36-45%)	60% (55-64%)	1.6 (1.4-1.9)	5.7% (4.7-6.6%)	7.7% (6.3-9.3%)	610000 (490000-740000)	2200000 (1800000-2600000)
Middle SDI	1990	Male	15 to 19	29% (20-39%)	71% (61-80%)	0.63 (0.37-0.99)	2.9% (1.9-4.3%)	3.8% (2.1-5.9%)	6200 (3500-9700)	550000 (340000-810000)
Middle SDI	1990	Male	20 to 24	45% (33-56%)	55% (44-67%)	1.6 (1-2.2)	6.8% (4.8-9.2%)	8.3% (5.4-12%)	19000 (13000-27000)	1600000 (1100000-2100000)
Middle SDI	1990	Male	25 to 29	52% (40-63%)	48% (37-60%)	2.2 (1.5-3)	8.8% (6.4-11%)	11% (7.6-15%)	24000 (17000-33000)	1800000 (1300000-2300000)
Middle SDI	1990	Male	30 to 34	55% (43-66%)	45% (34-57%)	2.7 (1.9-3.6)	9.7% (7.4-12%)	12% (9.2-16%)	28000 (20000-35000)	1800000 (1400000-2300000)
Middle SDI	1990	Male	35 to 39	58% (46-68%)	42% (32-54%)	3.1 (2.2-4.3)	10% (8.1-13%)	13% (10-17%)	34000 (25000-43000)	2000000 (1500000-2500000)
Middle SDI	1990	Male	40 to 44	58% (47-69%)	42% (31-53%)	3.5 (2.5-4.6)	11% (8.7-14%)	14% (11-17%)	35000 (27000-43000)	1800000 (1400000-2200000)
Middle SDI	1990	Male	45 to 49	58% (46-69%)	42% (31-54%)	3.7 (2.6-5)	12% (9.2-14%)	15% (11-18%)	42000 (32000-52000)	1900000 (1500000-2300000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	1990	Male	50 to 54	57% (46-68%)	43% (32-54%)	3.8 (2.6-5.1)	12% (8.9-14%)	14% (11-17%)	55000 (42000-68000)	220000 (170000-270000)
Middle SDI	1990	Male	55 to 59	56% (44-67%)	44% (33-56%)	3.7 (2.6-5.1)	11% (8.4-14%)	13% (9.9-16%)	69000 (52000-86000)	230000 (180000-290000)
Middle SDI	1990	Male	60 to 64	55% (43-66%)	45% (34-57%)	3.4 (2.3-4.8)	10% (7.3-13%)	12% (8.4-15%)	77000 (56000-98000)	220000 (160000-280000)
Middle SDI	1990	Male	65 to 69	52% (40-63%)	48% (37-60%)	2.9 (2-4)	8.6% (6.2-11%)	9.8% (7.1-13%)	73000 (53000-95000)	170000 (120000-220000)
Middle SDI	1990	Male	70 to 74	48% (37-59%)	52% (41-63%)	2.3 (1.5-3.3)	7% (4.7-9.6%)	7.8% (5.2-11%)	68000 (46000-94000)	130000 (86000-170000)
Middle SDI	1990	Male	75 to 79	45% (33-57%)	55% (43-67%)	1.9 (1.2-2.9)	5.5% (3.4-8.1%)	6.1% (3.6-9.1%)	45000 (27000-66000)	65000 (40000-95000)
Middle SDI	1990	Male	80 to 84	41% (29-54%)	59% (46-71%)	1.7 (1-2.4)	4.4% (2.5-6.6%)	4.9% (2.8-7.4%)	23000 (13000-34000)	24000 (14000-36000)
Middle SDI	1990	Male	85 to 89	39% (28-51%)	61% (49-72%)	1.5 (0.88-2.3)	3.5% (1.8-5.5%)	3.8% (1.9-6.1%)	9600 (4900-15000)	74000 (39000-120000)
Middle SDI	1990	Male	90 to 94	40% (29-51%)	60% (49-71%)	1.5 (0.85-2.3)	2.8% (1.2-4.7%)	3% (1.2-5.2%)	2900 (1200-4900)	16000 (6800-26000)
Middle SDI	1990	Male	95 plus	43% (29-57%)	57% (43-71%)	1.5 (0.77-2.5)	2.6% (0.78-4.8%)	2.7% (0.62-5.2%)	620 (140-1200)	2300 (690-4300)
Middle SDI	1995	Female	All Ages	17% (14-20%)	83% (80-86%)	0.37 (0.3-0.45)	1.4% (1-1.8%)	2% (1.3-2.9%)	130000 (83000-190000)	440000 (320000-570000)
Middle SDI	1995	Female	15 to 19	15% (10-22%)	85% (78-90%)	0.33 (0.17-0.56)	1.1% (0.66-1.6%)	1.7% (0.83-2.8%)	1200 (600-2000)	14000 (8300-21000)
Middle SDI	1995	Female	20 to 24	18% (12-26%)	82% (74-88%)	0.43 (0.23-0.71)	1.6% (1-2.3%)	1.9% (0.96-3.2%)	2300 (1100-3800)	28000 (17000-41000)
Middle SDI	1995	Female	25 to 29	19% (13-27%)	81% (73-87%)	0.46 (0.26-0.75)	1.7% (1.1-2.4%)	2.1% (1.1-3.3%)	3100 (1600-4900)	32000 (21000-46000)
Middle SDI	1995	Female	30 to 34	21% (14-30%)	79% (70-86%)	0.54 (0.29-0.86)	1.8% (1.3-2.5%)	2.5% (1.5-3.8%)	3700 (2300-5600)	32000 (22000-45000)
Middle SDI	1995	Female	35 to 39	22% (15-31%)	78% (69-85%)	0.59 (0.32-0.96)	2% (1.4-2.8%)	3.1% (2-4.6%)	4700 (3000-6900)	32000 (22000-45000)
Middle SDI	1995	Female	40 to 44	22% (15-31%)	78% (69-85%)	0.59 (0.32-0.95)	2.2% (1.5-3%)	3.5% (2.3-5.1%)	6700 (4400-9700)	37000 (25000-52000)
Middle SDI	1995	Female	45 to 49	22% (13-30%)	78% (70-87%)	0.65 (0.34-1.1)	2.4% (1.6-3.5%)	3.9% (2.4-5.7%)	8000 (5000-12000)	37000 (24000-53000)
Middle SDI	1995	Female	50 to 54	21% (14-30%)	79% (70-86%)	0.68 (0.36-1.2)	2.7% (1.7-3.9%)	4.1% (2.5-5.9%)	9600 (6000-14000)	38000 (24000-55000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	1995	Female	55 to 59	21% (14-30%)	79% (70-86%)	0.7 (0.36-1.2)	2.7% (1.7-4.1%)	4% (2.5-6%)	13000 (7900-19000)	420000 (260000-630000)
Middle SDI	1995	Female	60 to 64	20% (13-30%)	80% (70-87%)	0.67 (0.35-1.2)	2.5% (1.4-3.9%)	3.4% (2-5.3%)	16000 (9100-24000)	430000 (250000-670000)
Middle SDI	1995	Female	65 to 69	21% (13-29%)	79% (71-87%)	0.64 (0.35-1.1)	2.3% (1.3-3.6%)	3.1% (1.7-4.7%)	17000 (9700-26000)	390000 (220000-600000)
Middle SDI	1995	Female	70 to 74	20% (13-28%)	80% (72-87%)	0.57 (0.3-1)	1.8% (0.84-3.2%)	2.4% (1.1-4%)	16000 (7800-27000)	290000 (140000-500000)
Middle SDI	1995	Female	75 to 79	19% (12-28%)	81% (72-88%)	0.52 (0.26-0.92)	1.4% (0.46-2.7%)	1.8% (0.59-3.3%)	13000 (4600-24000)	180000 (62000-350000)
Middle SDI	1995	Female	80 to 84	19% (12-27%)	81% (73-88%)	0.53 (0.27-0.9)	1.2% (0.29-2.4%)	1.4% (0.35-2.9%)	9600 (2300-19000)	100000 (24000-200000)
Middle SDI	1995	Female	85 to 89	19% (12-29%)	81% (71-88%)	0.56 (0.27-0.99)	1% (0.13-2.3%)	1.2% (0.14-2.8%)	5700 (580-13000)	43000 (4800-97000)
Middle SDI	1995	Female	90 to 94	21% (14-31%)	79% (69-86%)	0.63 (0.3-1.2)	0.9% (-0.12-2.4%)	1% (-0.17-2.8%)	2200 (-390-6000)	12000 (-1600-31000)
Middle SDI	1995	Female	95 plus	25% (14-38%)	75% (62-86%)	0.74 (0.29-1.5)	0.93% (-0.39-2.9%)	1% (-0.55-3.4%)	620 (-340-2100)	2300 (-980-7200)
Middle SDI	1995	Male	All Ages	41% (37-46%)	59% (54-63%)	1.9 (1.7-2.2)	6.8% (5.7-7.8%)	8.7% (7.1-10%)	740000 (610000-880000)	2700000 (2200000-3100000)
Middle SDI	1995	Male	15 to 19	30% (22-40%)	70% (60-78%)	0.73 (0.42-1.1)	3.3% (2.1-4.8%)	4.3% (2.5-6.7%)	6500 (3700-10000)	570000 (350000-840000)
Middle SDI	1995	Male	20 to 24	45% (34-56%)	55% (44-66%)	1.8 (1.2-2.5)	7.4% (5.3-10%)	9.1% (6-13%)	22000 (14000-30000)	170000 (120000-240000)
Middle SDI	1995	Male	25 to 29	53% (41-63%)	47% (37-59%)	2.5 (1.7-3.4)	9.3% (6.8-12%)	12% (8.1-15%)	31000 (22000-42000)	230000 (160000-290000)
Middle SDI	1995	Male	30 to 34	56% (43-66%)	44% (34-57%)	3 (2.2-4)	10% (7.9-13%)	13% (9.7-16%)	35000 (26000-45000)	230000 (180000-290000)
Middle SDI	1995	Male	35 to 39	57% (47-67%)	43% (33-53%)	3.5 (2.5-4.7)	11% (8.7-14%)	14% (11-18%)	39000 (30000-49000)	230000 (180000-280000)
Middle SDI	1995	Male	40 to 44	59% (48-69%)	41% (31-52%)	3.9 (2.9-5.2)	12% (9.4-14%)	15% (12-18%)	50000 (39000-61000)	260000 (210000-310000)
Middle SDI	1995	Male	45 to 49	59% (46-69%)	41% (31-54%)	4.2 (3-5.7)	12% (10-15%)	16% (12-19%)	54000 (42000-66000)	240000 (190000-300000)
Middle SDI	1995	Male	50 to 54	58% (46-68%)	42% (32-54%)	4.3 (3-5.7)	12% (9.7-15%)	15% (12-18%)	59000 (45000-72000)	230000 (180000-280000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	1995	Male	55 to 59	56% (44-67%)	44% (33-56%)	4.1 (2.9-5.6)	12% (9.1-14%)	14% (11-17%)	74000 (57000-91000)	250000 (190000-310000)
Middle SDI	1995	Male	60 to 64	54% (42-65%)	46% (35-58%)	3.7 (2.6-5.3)	11% (8.1-14%)	12% (9.2-16%)	91000 (68000-110000)	260000 (190000-320000)
Middle SDI	1995	Male	65 to 69	51% (40-62%)	49% (38-60%)	3.2 (2.2-4.3)	9.2% (6.8-12%)	11% (7.6-14%)	88000 (64000-110000)	210000 (150000-260000)
Middle SDI	1995	Male	70 to 74	48% (37-59%)	52% (41-63%)	2.6 (1.7-3.7)	7.6% (5.2-10%)	8.4% (5.8-11%)	77000 (53000-100000)	140000 (100000-190000)
Middle SDI	1995	Male	75 to 79	45% (33-58%)	55% (42-67%)	2.2 (1.4-3.3)	6.3% (4.1-9%)	7% (4.5-10%)	60000 (38000-85000)	860000 (560000-1200000)
Middle SDI	1995	Male	80 to 84	43% (31-55%)	57% (45-69%)	2 (1.3-2.9)	5.3% (3.2-7.6%)	5.8% (3.5-8.4%)	36000 (22000-51000)	380000 (240000-550000)
Middle SDI	1995	Male	85 to 89	40% (29-52%)	60% (48-71%)	1.7 (1-2.6)	4.1% (2.3-6.2%)	4.5% (2.5-6.9%)	15000 (8200-22000)	110000 (64000-170000)
Middle SDI	1995	Male	90 to 94	41% (30-52%)	59% (48-70%)	1.7 (0.98-2.6)	3.3% (1.6-5.3%)	3.5% (1.6-5.9%)	4300 (2000-7100)	23000 (11000-37000)
Middle SDI	1995	Male	95 plus	44% (30-57%)	56% (43-70%)	1.6 (0.85-2.7)	2.9% (0.93-5.3%)	3% (0.76-5.7%)	870 (220-1700)	3300 (1000-6000)
Middle SDI	2000	Female	All Ages	17% (15-20%)	83% (80-85%)	0.4 (0.33-0.48)	1.5% (1.1-1.9%)	2.1% (1.3-3%)	140000 (89000-210000)	470000 (350000-610000)
Middle SDI	2000	Female	15 to 19	17% (12-24%)	83% (76-88%)	0.38 (0.2-0.63)	1.1% (0.71-1.7%)	1.9% (0.95-3.2%)	1400 (690-2300)	150000 (93000-230000)
Middle SDI	2000	Female	20 to 24	20% (14-28%)	80% (72-86%)	0.52 (0.28-0.85)	1.7% (1.1-2.5%)	2.2% (1.2-3.6%)	2300 (1200-3800)	270000 (170000-400000)
Middle SDI	2000	Female	25 to 29	20% (13-27%)	80% (73-87%)	0.5 (0.28-0.8)	1.7% (1.2-2.4%)	2.1% (1.2-3.3%)	3300 (1800-5200)	34000 (22000-48000)
Middle SDI	2000	Female	30 to 34	20% (13-28%)	80% (72-87%)	0.51 (0.28-0.82)	1.7% (1.2-2.3%)	2.3% (1.4-3.5%)	4000 (2500-6100)	35000 (24000-50000)
Middle SDI	2000	Female	35 to 39	21% (14-29%)	79% (71-86%)	0.58 (0.32-0.94)	1.9% (1.3-2.6%)	2.9% (1.9-4.2%)	5000 (3200-7400)	36000 (24000-50000)
Middle SDI	2000	Female	40 to 44	21% (14-29%)	79% (71-86%)	0.62 (0.34-1)	2.1% (1.4-2.9%)	3.4% (2.2-4.9%)	6500 (4200-9400)	37000 (25000-52000)
Middle SDI	2000	Female	45 to 49	20% (13-28%)	80% (72-87%)	0.62 (0.32-1)	2.2% (1.4-3.1%)	3.6% (2.2-5.2%)	8900 (5600-13000)	42000 (27000-61000)
Middle SDI	2000	Female	50 to 54	20% (13-29%)	80% (71-87%)	0.68 (0.36-1.1)	2.5% (1.5-3.6%)	3.8% (2.4-5.6%)	11000 (6700-16000)	43000 (27000-62000)
Middle SDI	2000	Female	55 to 59	20% (14-29%)	80% (71-86%)	0.72 (0.38-1.2)	2.6% (1.6-4%)	3.9% (2.4-5.8%)	13000 (7800-19000)	42000 (26000-63000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2000	Female	60 to 64	20% (13-29%)	80% (71-87%)	0.7 (0.37-1.2)	2.4% (1.4-3.9%)	3.4% (1.9-5.3%)	16000 (9100-25000)	440000 (250000-710000)
Middle SDI	2000	Female	65 to 69	19% (12-27%)	81% (73-88%)	0.62 (0.34-1)	2.2% (1.2-3.5%)	3% (1.7-4.6%)	19000 (11000-29000)	430000 (240000-660000)
Middle SDI	2000	Female	70 to 74	19% (13-27%)	81% (73-87%)	0.57 (0.29-1)	1.8% (0.83-3.1%)	2.4% (1.1-4%)	18000 (8500-30000)	330000 (150000-560000)
Middle SDI	2000	Female	75 to 79	18% (12-27%)	82% (73-88%)	0.53 (0.26-0.94)	1.4% (0.44-2.6%)	1.8% (0.58-3.3%)	14000 (4600-26000)	200000 (63000-370000)
Middle SDI	2000	Female	80 to 84	18% (12-26%)	82% (74-88%)	0.53 (0.27-0.91)	1.2% (0.29-2.4%)	1.5% (0.38-2.9%)	11000 (2700-22000)	120000 (27000-230000)
Middle SDI	2000	Female	85 to 89	19% (12-28%)	81% (72-88%)	0.57 (0.28-0.99)	1.1% (0.15-2.4%)	1.3% (0.17-2.8%)	7200 (790-16000)	54000 (6700-120000)
Middle SDI	2000	Female	90 to 94	21% (14-30%)	79% (70-86%)	0.65 (0.31-1.2)	1% (-0.04-2.6%)	1.2% (-0.08-3%)	3200 (-270-8200)	17000 (-780-43000)
Middle SDI	2000	Female	95 plus	24% (14-37%)	76% (63-86%)	0.77 (0.3-1.6)	1.1% (-0.3-3.1%)	1.2% (-0.44-3.6%)	990 (-370-3000)	3600 (-1000-10000)
Middle SDI	2000	Male	All Ages	43% (39-48%)	57% (52-61%)	2.1 (1.8-2.4)	7.4% (6.3-8.6%)	9.1% (7.4-11%)	820000 (660000-980000)	29000000 (25000000-34000000)
Middle SDI	2000	Male	15 to 19	34% (26-43%)	66% (57-74%)	0.87 (0.52-1.3)	3.7% (2.4-5.4%)	5% (2.9-7.6%)	7600 (4300-12000)	650000 (400000-960000)
Middle SDI	2000	Male	20 to 24	46% (36-56%)	54% (44-64%)	1.9 (1.3-2.7)	8% (5.7-11%)	9.9% (6.5-14%)	21000 (14000-30000)	1700000 (1200000-2300000)
Middle SDI	2000	Male	25 to 29	52% (41-62%)	48% (38-59%)	2.5 (1.7-3.5)	9.4% (6.9-12%)	12% (8.2-16%)	30000 (20000-40000)	2200000 (1600000-2900000)
Middle SDI	2000	Male	30 to 34	56% (43-66%)	44% (34-57%)	3.1 (2.2-4.1)	10% (7.9-13%)	13% (9.7-17%)	38000 (28000-49000)	2500000 (1900000-3200000)
Middle SDI	2000	Male	35 to 39	58% (47-68%)	42% (32-53%)	3.6 (2.6-4.8)	11% (8.7-14%)	14% (11-18%)	44000 (33000-56000)	2600000 (2000000-3300000)
Middle SDI	2000	Male	40 to 44	58% (48-68%)	42% (32-52%)	4 (2.9-5.2)	12% (9.5-15%)	15% (12-19%)	52000 (40000-64000)	2700000 (2100000-3300000)
Middle SDI	2000	Male	45 to 49	59% (47-69%)	41% (31-53%)	4.3 (3.1-5.8)	13% (10-15%)	16% (12-19%)	69000 (54000-84000)	3200000 (2500000-3800000)
Middle SDI	2000	Male	50 to 54	58% (47-68%)	42% (32-53%)	4.4 (3.1-5.8)	12% (9.7-15%)	15% (12-18%)	69000 (53000-84000)	2800000 (2200000-3400000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2000	Male	55 to 59	56% (45-66%)	44% (34-55%)	4.2 (2.9-5.7)	12% (9-14%)	14% (11-17%)	75000 (57000-92000)	260000 (200000-320000)
Middle SDI	2000	Male	60 to 64	53% (42-65%)	47% (35-58%)	3.7 (2.6-5.3)	11% (7.9-13%)	12% (9.2-16%)	92000 (68000-120000)	260000 (200000-330000)
Middle SDI	2000	Male	65 to 69	51% (39-61%)	49% (39-61%)	3.2 (2.2-4.3)	9.3% (6.8-12%)	11% (7.7-14%)	100000 (73000-130000)	240000 (170000-300000)
Middle SDI	2000	Male	70 to 74	48% (37-58%)	52% (42-63%)	2.6 (1.7-3.7)	7.7% (5.3-10%)	8.6% (5.8-12%)	90000 (61000-120000)	170000 (120000-230000)
Middle SDI	2000	Male	75 to 79	45% (34-57%)	55% (43-66%)	2.2 (1.5-3.3)	6.3% (4-9%)	7% (4.4-10%)	64000 (41000-92000)	930000 (600000-1300000)
Middle SDI	2000	Male	80 to 84	43% (31-55%)	57% (45-69%)	2 (1.3-2.9)	5.4% (3.3-7.8%)	5.9% (3.6-8.7%)	42000 (26000-61000)	460000 (280000-660000)
Middle SDI	2000	Male	85 to 89	41% (30-52%)	59% (48-70%)	1.8 (1.1-2.7)	4.3% (2.5-6.5%)	4.8% (2.7-7.2%)	20000 (11000-30000)	150000 (88000-230000)
Middle SDI	2000	Male	90 to 94	41% (30-52%)	59% (48-70%)	1.7 (1-2.6)	3.4% (1.7-5.6%)	3.7% (1.7-6.1%)	5600 (2600-9400)	30000 (15000-50000)
Middle SDI	2000	Male	95 plus	44% (30-57%)	56% (43-70%)	1.7 (0.88-2.8)	3% (0.99-5.6%)	3.1% (0.83-6%)	1100 (290-2200)	4300 (1400-8100)
Middle SDI	2005	Female	All Ages	18% (16-20%)	82% (80-84%)	0.43 (0.35-0.52)	1.5% (1.1-1.9%)	2% (1.3-2.9%)	150000 (93000-210000)	480000 (350000-620000)
Middle SDI	2005	Female	15 to 19	18% (13-24%)	82% (76-87%)	0.38 (0.21-0.62)	1.1% (0.72-1.7%)	1.9% (1-3.2%)	1500 (770-2400)	170000 (100000-240000)
Middle SDI	2005	Female	20 to 24	22% (16-28%)	78% (72-84%)	0.57 (0.32-0.9)	1.8% (1.2-2.5%)	2.4% (1.3-3.8%)	2600 (1400-4000)	300000 (190000-430000)
Middle SDI	2005	Female	25 to 29	21% (16-28%)	79% (72-84%)	0.59 (0.36-0.91)	1.8% (1.3-2.4%)	2.4% (1.4-3.6%)	3100 (1800-4900)	320000 (210000-460000)
Middle SDI	2005	Female	30 to 34	20% (15-27%)	80% (73-85%)	0.54 (0.32-0.82)	1.6% (1.2-2.2%)	2.3% (1.5-3.4%)	4100 (2500-6100)	350000 (240000-500000)
Middle SDI	2005	Female	35 to 39	21% (16-28%)	79% (72-84%)	0.55 (0.33-0.85)	1.7% (1.2-2.3%)	2.7% (1.7-3.8%)	4900 (3000-7400)	360000 (240000-500000)
Middle SDI	2005	Female	40 to 44	22% (16-29%)	78% (71-84%)	0.62 (0.37-0.95)	1.9% (1.3-2.6%)	3.2% (2.1-4.5%)	6600 (4300-9500)	390000 (260000-540000)
Middle SDI	2005	Female	45 to 49	21% (15-27%)	79% (73-85%)	0.67 (0.39-1.1)	2% (1.3-3%)	3.5% (2.2-5.1%)	8400 (5300-13000)	410000 (270000-590000)
Middle SDI	2005	Female	50 to 54	20% (15-27%)	80% (73-85%)	0.64 (0.37-1)	2.2% (1.4-3.2%)	3.5% (2.2-5.1%)	12000 (7700-18000)	480000 (310000-700000)
Middle SDI	2005	Female	55 to 59	19% (14-25%)	81% (75-86%)	0.66 (0.38-1.1)	2.3% (1.4-3.5%)	3.5% (2.2-5.3%)	14000 (8400-20000)	460000 (280000-690000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2005	Female	60 to 64	18% (12-25%)	82% (75-88%)	0.65 (0.35-1.1)	2.2% (1.3-3.5%)	3.2% (1.9-4.9%)	14000 (8400-22000)	400000 (230000-620000)
Middle SDI	2005	Female	65 to 69	17% (12-24%)	83% (76-88%)	0.58 (0.32-0.94)	2% (1.2-3.2%)	2.8% (1.6-4.3%)	17000 (9900-27000)	400000 (220000-620000)
Middle SDI	2005	Female	70 to 74	16% (11-23%)	84% (77-89%)	0.52 (0.27-0.9)	1.7% (0.83-2.8%)	2.2% (1.1-3.6%)	18000 (8900-30000)	330000 (160000-550000)
Middle SDI	2005	Female	75 to 79	16% (10-23%)	84% (77-90%)	0.49 (0.24-0.87)	1.3% (0.48-2.4%)	1.7% (0.64-3%)	14000 (5300-26000)	210000 (72000-380000)
Middle SDI	2005	Female	80 to 84	15% (9.2-22%)	85% (78-91%)	0.49 (0.24-0.85)	1.2% (0.36-2.3%)	1.5% (0.46-2.8%)	12000 (3600-24000)	130000 (38000-250000)
Middle SDI	2005	Female	85 to 89	15% (9.3-22%)	85% (78-91%)	0.51 (0.25-0.9)	1.1% (0.22-2.2%)	1.3% (0.27-2.6%)	8100 (1500-17000)	62000 (12000-130000)
Middle SDI	2005	Female	90 to 94	17% (11-25%)	83% (75-89%)	0.58 (0.28-1.1)	1% (0.05-2.3%)	1.2% (0.032-2.8%)	3800 (86-9200)	20000 (1200-48000)
Middle SDI	2005	Female	95 plus	20% (12-31%)	80% (69-88%)	0.73 (0.28-1.5)	1.2% (-0.15-3%)	1.3% (-0.27-3.5%)	1400 (-280-3700)	5000 (-580-13000)
Middle SDI	2005	Male	All Ages	44% (41-48%)	56% (52-59%)	2.3 (2-2.6)	7.8% (6.6-9.1%)	9.1% (7.4-11%)	850000 (690000-1000000)	3000000 (2600000-3600000)
Middle SDI	2005	Male	15 to 19	36% (29-45%)	64% (55-71%)	0.93 (0.58-1.4)	3.8% (2.5-5.4%)	5.3% (3.2-7.9%)	8300 (4800-13000)	730000 (470000-1000000)
Middle SDI	2005	Male	20 to 24	49% (41-57%)	51% (43-59%)	2.1 (1.4-2.9)	8.4% (6.1-11%)	11% (7.1-15%)	21000 (14000-30000)	1700000 (1200000-2300000)
Middle SDI	2005	Male	25 to 29	53% (44-61%)	47% (39-56%)	2.7 (1.9-3.7)	9.7% (7.1-12%)	13% (8.8-17%)	26000 (17000-35000)	1900000 (1400000-2600000)
Middle SDI	2005	Male	30 to 34	56% (46-65%)	44% (35-54%)	3.2 (2.4-4.2)	10% (7.9-13%)	13% (9.9-17%)	35000 (26000-46000)	2400000 (1800000-3000000)
Middle SDI	2005	Male	35 to 39	58% (49-66%)	42% (34-51%)	3.7 (2.7-4.8)	11% (8.6-14%)	14% (11-18%)	46000 (35000-59000)	2800000 (2200000-3500000)
Middle SDI	2005	Male	40 to 44	58% (49-67%)	42% (33-51%)	4 (3-5.3)	12% (9.5-14%)	15% (12-19%)	58000 (45000-72000)	3100000 (2400000-3700000)
Middle SDI	2005	Male	45 to 49	58% (49-67%)	42% (33-51%)	4.3 (3.2-5.7)	12% (9.9-15%)	16% (13-19%)	66000 (52000-81000)	3100000 (2400000-3700000)
Middle SDI	2005	Male	50 to 54	59% (49-68%)	41% (32-51%)	4.5 (3.3-5.9)	12% (9.8-15%)	15% (12-18%)	89000 (70000-110000)	3600000 (2800000-4300000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2005	Male	55 to 59	56% (47-65%)	44% (35-53%)	4.2 (3.1-5.7)	12% (9-14%)	14% (11-17%)	86000 (67000-110000)	300000 (230000-360000)
Middle SDI	2005	Male	60 to 64	53% (44-63%)	47% (37-56%)	3.8 (2.7-5.3)	10% (7.7-13%)	12% (9.1-15%)	84000 (62000-110000)	240000 (180000-300000)
Middle SDI	2005	Male	65 to 69	50% (40-59%)	50% (41-60%)	3.1 (2.2-4.2)	8.8% (6.5-11%)	10% (7.5-13%)	96000 (70000-120000)	230000 (170000-290000)
Middle SDI	2005	Male	70 to 74	46% (36-56%)	54% (44-64%)	2.5 (1.7-3.6)	7.3% (5-9.6%)	8.2% (5.6-11%)	91000 (62000-120000)	170000 (120000-230000)
Middle SDI	2005	Male	75 to 79	43% (33-54%)	57% (46-67%)	2.1 (1.4-3.2)	5.8% (3.7-8.3%)	6.5% (4.1-9.3%)	67000 (42000-96000)	990000 (620000-1400000)
Middle SDI	2005	Male	80 to 84	40% (29-52%)	60% (48-71%)	1.9 (1.2-2.8)	5% (2.9-7.3%)	5.5% (3.2-8.1%)	45000 (26000-67000)	490000 (290000-730000)
Middle SDI	2005	Male	85 to 89	39% (28-50%)	61% (50-72%)	1.7 (1.1-2.6)	4% (2.3-6.2%)	4.4% (2.4-6.9%)	21000 (12000-33000)	170000 (94000-260000)
Middle SDI	2005	Male	90 to 94	40% (29-51%)	60% (49-71%)	1.6 (1-2.5)	3.2% (1.6-5.4%)	3.5% (1.6-6%)	6900 (3100-12000)	38000 (18000-64000)
Middle SDI	2005	Male	95 plus	42% (29-56%)	58% (44-71%)	1.6 (0.86-2.7)	2.9% (0.88-5.4%)	3% (0.72-5.9%)	1400 (320-2800)	5200 (1500-10000)
Middle SDI	2010	Female	All Ages	18% (16-20%)	82% (80-84%)	0.47 (0.38-0.57)	1.6% (1.2-2%)	2.1% (1.4-3%)	160000 (100000-220000)	500000 (370000-640000)
Middle SDI	2010	Female	15 to 19	18% (13-24%)	82% (76-87%)	0.41 (0.22-0.68)	1.1% (0.73-1.7%)	2% (1-3.3%)	1200 (640-2000)	140000 (91000-220000)
Middle SDI	2010	Female	20 to 24	21% (15-28%)	79% (72-85%)	0.54 (0.3-0.87)	1.8% (1.2-2.5%)	2.3% (1.2-3.7%)	2500 (1400-3800)	310000 (200000-440000)
Middle SDI	2010	Female	25 to 29	22% (16-28%)	78% (72-84%)	0.63 (0.37-0.99)	1.9% (1.3-2.5%)	2.5% (1.4-3.8%)	2900 (1700-4500)	320000 (210000-450000)
Middle SDI	2010	Female	30 to 34	22% (16-29%)	78% (71-84%)	0.63 (0.37-0.97)	1.8% (1.2-2.4%)	2.5% (1.6-3.7%)	3100 (1900-4700)	300000 (200000-420000)
Middle SDI	2010	Female	35 to 39	22% (16-28%)	78% (72-84%)	0.59 (0.35-0.93)	1.7% (1.2-2.3%)	2.8% (1.8-4%)	4300 (2700-6400)	330000 (220000-460000)
Middle SDI	2010	Female	40 to 44	21% (16-28%)	79% (72-84%)	0.61 (0.36-0.95)	1.8% (1.2-2.4%)	3.2% (2.1-4.4%)	6300 (4100-9100)	390000 (260000-540000)
Middle SDI	2010	Female	45 to 49	21% (15-28%)	79% (72-85%)	0.68 (0.39-1.1)	2% (1.3-2.9%)	3.6% (2.3-5.2%)	9100 (5800-13000)	450000 (290000-640000)
Middle SDI	2010	Female	50 to 54	20% (15-27%)	80% (73-85%)	0.72 (0.41-1.2)	2.3% (1.4-3.3%)	3.9% (2.5-5.6%)	12000 (7800-18000)	500000 (320000-730000)
Middle SDI	2010	Female	55 to 59	19% (14-25%)	81% (75-86%)	0.67 (0.38-1.1)	2.3% (1.4-3.5%)	3.8% (2.4-5.5%)	16000 (10000-24000)	550000 (340000-820000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2010	Female	60 to 64	17% (12-24%)	83% (76-88%)	0.64 (0.34-1.1)	2.2% (1.3-3.4%)	3.4% (2.1-5%)	17000 (10000-25000)	470000 (280000-720000)
Middle SDI	2010	Female	65 to 69	17% (12-24%)	83% (76-88%)	0.59 (0.32-0.95)	2.1% (1.2-3.1%)	3% (1.8-4.4%)	17000 (10000-26000)	390000 (230000-600000)
Middle SDI	2010	Female	70 to 74	16% (11-23%)	84% (77-89%)	0.54 (0.27-0.95)	1.7% (0.87-2.8%)	2.3% (1.2-3.8%)	18000 (9300-29000)	340000 (170000-550000)
Middle SDI	2010	Female	75 to 79	15% (9.8-22%)	85% (78-90%)	0.5 (0.24-0.89)	1.4% (0.53-2.4%)	1.8% (0.72-3.1%)	16000 (6300-28000)	230000 (88000-410000)
Middle SDI	2010	Female	80 to 84	15% (9-22%)	85% (78-91%)	0.52 (0.25-0.9)	1.2% (0.4-2.3%)	1.5% (0.51-2.8%)	14000 (4600-27000)	150000 (49000-290000)
Middle SDI	2010	Female	85 to 89	15% (9.4-23%)	85% (77-91%)	0.56 (0.27-1)	1.1% (0.25-2.3%)	1.4% (0.3-2.8%)	9900 (2000-20000)	76000 (17000-150000)
Middle SDI	2010	Female	90 to 94	17% (11-26%)	83% (74-89%)	0.64 (0.3-1.2)	1% (0.087-2.4%)	1.2% (0.04-2.9%)	4800 (180-11000)	26000 (2300-59000)
Middle SDI	2010	Female	95 plus	21% (12-32%)	79% (68-88%)	0.78 (0.3-1.6)	1.1% (-0.15-2.9%)	1.3% (-0.31-3.4%)	1800 (-370-4700)	6500 (-710-17000)
Middle SDI	2010	Male	All Ages	44% (41-48%)	56% (52-59%)	2.5 (2.2-2.9)	8.2% (6.9-9.5%)	9.3% (7.6-11%)	910000 (730000-1100000)	3200000 (2700000-3700000)
Middle SDI	2010	Male	15 to 19	34% (27-42%)	66% (58-73%)	0.93 (0.56-1.4)	3.8% (2.5-5.4%)	5.3% (3.2-8%)	7200 (4100-11000)	640000 (400000-930000)
Middle SDI	2010	Male	20 to 24	48% (39-57%)	52% (43-61%)	2.1 (1.5-3)	8.3% (5.9-11%)	11% (7.1-15%)	22000 (14000-31000)	180000 (130000-250000)
Middle SDI	2010	Male	25 to 29	52% (43-60%)	48% (40-57%)	2.8 (2-3.9)	9.7% (7.2-13%)	13% (9-17%)	26000 (17000-35000)	200000 (140000-260000)
Middle SDI	2010	Male	30 to 34	55% (45-63%)	45% (37-55%)	3.4 (2.4-4.5)	10% (7.9-13%)	14% (10-18%)	30000 (22000-40000)	210000 (160000-270000)
Middle SDI	2010	Male	35 to 39	57% (48-65%)	43% (35-52%)	3.9 (2.8-5.2)	11% (8.7-14%)	15% (11-18%)	42000 (31000-54000)	260000 (200000-330000)
Middle SDI	2010	Male	40 to 44	57% (48-67%)	43% (33-52%)	4.3 (3.2-5.6)	12% (9.5-14%)	15% (12-19%)	59000 (46000-74000)	320000 (260000-390000)
Middle SDI	2010	Male	45 to 49	58% (48-67%)	42% (33-52%)	4.6 (3.4-6.1)	12% (10-15%)	16% (13-19%)	73000 (57000-89000)	340000 (270000-420000)
Middle SDI	2010	Male	50 to 54	57% (47-66%)	43% (34-53%)	4.7 (3.4-6.1)	12% (9.7-15%)	15% (12-18%)	85000 (66000-100000)	350000 (270000-420000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2010	Male	55 to 59	56% (46-65%)	44% (35-54%)	4.6 (3.4-6.1)	12% (9.2-14%)	14% (11-17%)	110000 (83000-130000)	3700000 (2900000-4500000)
Middle SDI	2010	Male	60 to 64	53% (43-63%)	47% (37-57%)	4 (2.8-5.6)	10% (7.9-13%)	12% (9.4-15%)	98000 (75000-120000)	2900000 (2200000-3600000)
Middle SDI	2010	Male	65 to 69	49% (40-59%)	51% (41-60%)	3.3 (2.4-4.5)	8.9% (6.8-11%)	10% (7.9-13%)	92000 (70000-120000)	2200000 (1700000-2800000)
Middle SDI	2010	Male	70 to 74	46% (36-55%)	54% (45-64%)	2.7 (1.9-3.8)	7.4% (5.2-9.7%)	8.5% (5.9-11%)	93000 (65000-120000)	1800000 (1300000-2400000)
Middle SDI	2010	Male	75 to 79	43% (33-54%)	57% (46-67%)	2.3 (1.5-3.4)	6.1% (4-8.5%)	6.9% (4.4-9.7%)	78000 (50000-110000)	1200000 (760000-1600000)
Middle SDI	2010	Male	80 to 84	41% (30-53%)	59% (47-70%)	2.1 (1.3-3)	5.3% (3.3-7.6%)	5.9% (3.6-8.6%)	58000 (35000-84000)	640000 (400000-920000)
Middle SDI	2010	Male	85 to 89	39% (28-50%)	61% (50-72%)	1.9 (1.1-2.8)	4.3% (2.5-6.5%)	4.8% (2.7-7.2%)	28000 (16000-42000)	220000 (120000-330000)
Middle SDI	2010	Male	90 to 94	40% (29-51%)	60% (49-71%)	1.8 (1.1-2.8)	3.5% (1.7-5.7%)	3.8% (1.7-6.4%)	8900 (4000-15000)	49000 (24000-80000)
Middle SDI	2010	Male	95 plus	43% (29-57%)	57% (43-71%)	1.8 (0.97-3)	3.2% (1.1-5.9%)	3.3% (0.88-6.4%)	2000 (540-3900)	7800 (2600-14000)
Middle SDI	2016	Female	All Ages	19% (16-22%)	81% (78-84%)	0.52 (0.39-0.69)	1.7% (1.2-2.2%)	2.3% (1.4-3.4%)	170000 (110000-250000)	5300000 (3800000-7100000)
Middle SDI	2016	Female	15 to 19	18% (12-26%)	82% (74-88%)	0.46 (0.23-0.79)	1.1% (0.72-1.7%)	2.1% (1.1-3.6%)	880 (440-1500)	110000 (70000-170000)
Middle SDI	2016	Female	20 to 24	22% (15-30%)	78% (70-85%)	0.61 (0.32-1)	1.9% (1.3-2.6%)	2.5% (1.3-4.1%)	1700 (910-2800)	240000 (160000-360000)
Middle SDI	2016	Female	25 to 29	23% (16-31%)	77% (69-84%)	0.62 (0.34-1)	1.8% (1.3-2.5%)	2.4% (1.4-3.9%)	2500 (1400-3900)	310000 (210000-440000)
Middle SDI	2016	Female	30 to 34	24% (16-33%)	76% (67-84%)	0.67 (0.37-1.1)	1.8% (1.3-2.5%)	2.7% (1.7-4.1%)	3100 (1900-4700)	310000 (210000-440000)
Middle SDI	2016	Female	35 to 39	25% (17-34%)	75% (66-83%)	0.72 (0.38-1.2)	1.9% (1.3-2.6%)	3.3% (2.1-4.8%)	4100 (2500-6100)	320000 (210000-450000)
Middle SDI	2016	Female	40 to 44	23% (16-32%)	77% (68-84%)	0.69 (0.37-1.1)	1.9% (1.3-2.6%)	3.6% (2.3-5.1%)	6200 (4000-9200)	380000 (250000-550000)
Middle SDI	2016	Female	45 to 49	21% (13-30%)	79% (70-87%)	0.68 (0.36-1.1)	2% (1.3-2.8%)	3.8% (2.4-5.5%)	9700 (6100-14000)	480000 (310000-700000)
Middle SDI	2016	Female	50 to 54	20% (13-29%)	80% (71-87%)	0.73 (0.39-1.2)	2.3% (1.4-3.3%)	4.1% (2.6-5.8%)	14000 (8700-20000)	570000 (360000-830000)
Middle SDI	2016	Female	55 to 59	20% (13-29%)	80% (71-87%)	0.77 (0.4-1.3)	2.5% (1.6-3.7%)	4.3% (2.8-6.1%)	17000 (11000-26000)	590000 (360000-880000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2016	Female	60 to 64	19% (11-29%)	81% (71-89%)	0.69 (0.34-1.2)	2.3% (1.4-3.6%)	3.7% (2.3-5.5%)	21000 (13000-33000)	610000 (360000-940000)
Middle SDI	2016	Female	65 to 69	19% (12-27%)	81% (73-88%)	0.64 (0.33-1.1)	2.1% (1.2-3.3%)	3.2% (2-4.8%)	21000 (13000-32000)	490000 (280000-760000)
Middle SDI	2016	Female	70 to 74	18% (11-26%)	82% (74-89%)	0.59 (0.29-1.1)	1.8% (0.9-2.9%)	2.6% (1.4-4.1%)	18000 (9600-30000)	340000 (170000-580000)
Middle SDI	2016	Female	75 to 79	17% (10-26%)	83% (74-90%)	0.55 (0.25-1)	1.4% (0.55-2.6%)	1.9% (0.78-3.4%)	17000 (6600-30000)	240000 (91000-440000)
Middle SDI	2016	Female	80 to 84	16% (9.5-24%)	84% (76-90%)	0.57 (0.27-1)	1.3% (0.39-2.4%)	1.6% (0.51-3%)	16000 (4800-30000)	170000 (52000-320000)
Middle SDI	2016	Female	85 to 89	17% (9.9-26%)	83% (74-90%)	0.61 (0.28-1.1)	1.1% (0.22-2.3%)	1.4% (0.25-2.8%)	12000 (2200-25000)	95000 (19000-200000)
Middle SDI	2016	Female	90 to 94	19% (12-29%)	81% (71-88%)	0.71 (0.32-1.4)	1% (-0.0074-2.4%)	1.2% (-0.08-2.9%)	6300 (-320-15000)	35000 (350-81000)
Middle SDI	2016	Female	95 plus	23% (12-35%)	77% (65-88%)	0.85 (0.32-1.8)	1.1% (-0.27-3%)	1.2% (-0.46-3.5%)	2400 (-770-6600)	8800 (-1800-23000)
Middle SDI	2016	Male	All Ages	45% (41-49%)	55% (51-59%)	2.8 (2.3-3.3)	8.5% (7-10%)	9.4% (7.6-11%)	1000000 (800000-1200000)	3400000 (2800000-4000000)
Middle SDI	2016	Male	15 to 19	32% (24-42%)	68% (58-76%)	0.9 (0.51-1.5)	3.7% (2.3-5.3%)	5.2% (2.9-8.1%)	5800 (3100-9300)	520000 (320000-780000)
Middle SDI	2016	Male	20 to 24	45% (35-55%)	55% (45-65%)	2.1 (1.3-3.1)	8.2% (5.7-11%)	11% (6.8-15%)	18000 (11000-27000)	150000 (100000-210000)
Middle SDI	2016	Male	25 to 29	50% (39-60%)	50% (40-61%)	2.8 (1.9-4.1)	9.4% (6.9-12%)	13% (8.6-17%)	26000 (17000-36000)	210000 (150000-270000)
Middle SDI	2016	Male	30 to 34	53% (41-63%)	47% (37-59%)	3.4 (2.3-4.8)	10% (7.7-13%)	14% (10-18%)	30000 (21000-40000)	210000 (160000-270000)
Middle SDI	2016	Male	35 to 39	55% (45-65%)	45% (35-55%)	3.9 (2.6-5.5)	11% (8.2-13%)	15% (11-19%)	35000 (25000-46000)	220000 (160000-280000)
Middle SDI	2016	Male	40 to 44	57% (47-67%)	43% (33-53%)	4.4 (3.1-6)	11% (9.1-14%)	15% (12-19%)	51000 (39000-65000)	280000 (220000-350000)
Middle SDI	2016	Male	45 to 49	58% (47-68%)	42% (32-53%)	4.9 (3.5-6.6)	12% (9.8-15%)	16% (13-19%)	76000 (59000-94000)	370000 (290000-450000)
Middle SDI	2016	Male	50 to 54	58% (47-68%)	42% (32-53%)	5 (3.6-6.7)	12% (9.6-15%)	15% (12-19%)	94000 (73000-120000)	390000 (310000-480000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Middle SDI	2016	Male	55 to 59	57% (46-67%)	43% (33-54%)	4.8 (3.5-6.6)	12% (9.1-14%)	14% (11-17%)	100000 (80000-130000)	3700000 (2900000-4600000)
Middle SDI	2016	Male	60 to 64	55% (44-66%)	45% (34-56%)	4.5 (3.1-6.3)	11% (8.3-13%)	13% (10-16%)	130000 (100000-170000)	3900000 (3000000-4900000)
Middle SDI	2016	Male	65 to 69	52% (41-63%)	48% (37-59%)	3.7 (2.6-5.1)	9.5% (7.2-12%)	11% (8.4-14%)	120000 (92000-160000)	3000000 (2300000-3800000)
Middle SDI	2016	Male	70 to 74	48% (37-58%)	52% (42-63%)	2.9 (1.9-4.3)	7.6% (5.3-10%)	8.8% (6-12%)	97000 (67000-130000)	1900000 (1300000-2500000)
Middle SDI	2016	Male	75 to 79	44% (32-56%)	56% (44-68%)	2.5 (1.5-3.7)	6.3% (4-8.8%)	7.1% (4.5-10%)	84000 (53000-120000)	1300000 (810000-1800000)
Middle SDI	2016	Male	80 to 84	42% (30-54%)	58% (46-70%)	2.2 (1.4-3.3)	5.5% (3.4-7.8%)	6.1% (3.7-8.9%)	68000 (41000-98000)	750000 (460000-1100000)
Middle SDI	2016	Male	85 to 89	40% (29-51%)	60% (49-71%)	2 (1.2-3.1)	4.6% (2.7-6.9%)	5.1% (2.9-7.8%)	39000 (22000-59000)	310000 (180000-460000)
Middle SDI	2016	Male	90 to 94	41% (31-53%)	59% (47-69%)	1.9 (1.2-3)	3.7% (1.8-6%)	4.1% (1.8-6.7%)	13000 (5900-22000)	72000 (35000-120000)
Middle SDI	2016	Male	95 plus	44% (31-58%)	56% (42-69%)	1.9 (1-3.2)	3.3% (1.1-6.2%)	3.5% (0.87-6.7%)	2900 (720-5500)	11000 (3600-20000)
Low-middle SDI	1990	Female	All Ages	7.6% (6-9.5%)	92% (90-94%)	0.2 (0.15-0.26)	0.54% (0.38-0.73%)	0.88% (0.53-1.3%)	61000 (37000-93000)	2400000 (1700000-3300000)
Low-middle SDI	1990	Female	15 to 19	6.7% (4-10%)	93% (90-96%)	0.15 (0.074-0.28)	0.47% (0.25-0.79%)	0.49% (0.17-0.99%)	570 (180-1200)	74000 (39000-130000)
Low-middle SDI	1990	Female	20 to 24	7.8% (4.7-12%)	92% (88-95%)	0.24 (0.11-0.42)	0.91% (0.54-1.4%)	0.84% (0.35-1.6%)	1300 (530-2500)	160000 (90000-250000)
Low-middle SDI	1990	Female	25 to 29	8.5% (5.1-13%)	92% (87-95%)	0.3 (0.14-0.53)	1.2% (0.73-1.9%)	1.2% (0.57-2.3%)	2000 (890-3600)	190000 (110000-300000)
Low-middle SDI	1990	Female	30 to 34	9.5% (5.7-15%)	91% (85-94%)	0.36 (0.17-0.62)	1.4% (0.89-2.2%)	1.7% (0.91-2.9%)	2700 (1400-4500)	210000 (130000-320000)
Low-middle SDI	1990	Female	35 to 39	10% (6-15%)	90% (85-94%)	0.4 (0.19-0.69)	1.7% (1.1-2.6%)	2.2% (1.3-3.7%)	3600 (2000-5900)	230000 (140000-350000)
Low-middle SDI	1990	Female	40 to 44	11% (7-17%)	89% (83-93%)	0.42 (0.2-0.73)	1.8% (1.1-2.7%)	2.5% (1.5-4%)	4000 (2300-6400)	220000 (140000-330000)
Low-middle SDI	1990	Female	45 to 49	11% (6.7-17%)	89% (83-93%)	0.39 (0.17-0.71)	1.8% (1.1-2.8%)	2.4% (1.4-4%)	4700 (2600-7700)	220000 (130000-340000)
Low-middle SDI	1990	Female	50 to 54	12% (7.1-18%)	88% (82-93%)	0.41 (0.19-0.76)	1.8% (1.1-2.9%)	2.4% (1.4-4%)	6100 (3400-10000)	240000 (140000-390000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	1990	Female	55 to 59	12% (7-19%)	88% (81-93%)	0.48 (0.22-0.91)	1.9% (1-3.1%)	2.4% (1.3-4.1%)	7600 (3900-13000)	250000 (140000-420000)
Low-middle SDI	1990	Female	60 to 64	13% (7.3-20%)	87% (80-93%)	0.51 (0.23-0.99)	1.7% (0.8-2.9%)	2.1% (0.97-3.7%)	7800 (3700-14000)	220000 (110000-380000)
Low-middle SDI	1990	Female	65 to 69	13% (7.6-20%)	87% (80-92%)	0.48 (0.22-0.87)	1.4% (0.66-2.5%)	1.7% (0.78-3.1%)	7500 (3400-13000)	170000 (81000-310000)
Low-middle SDI	1990	Female	70 to 74	13% (7.5-19%)	87% (81-93%)	0.41 (0.18-0.78)	1.1% (0.4-2.2%)	1.3% (0.46-2.7%)	6100 (2200-12000)	110000 (42000-220000)
Low-middle SDI	1990	Female	75 to 79	12% (6.5-20%)	88% (80-93%)	0.35 (0.14-0.72)	0.79% (0.17-1.8%)	0.92% (0.18-2.1%)	3700 (830-8300)	53000 (13000-120000)
Low-middle SDI	1990	Female	80 to 84	11% (5.9-20%)	89% (80-94%)	0.33 (0.13-0.72)	0.61% (0.088-1.5%)	0.71% (0.086-1.7%)	2200 (330-5300)	23000 (4000-55000)
Low-middle SDI	1990	Female	85 to 89	12% (6.5-22%)	88% (78-93%)	0.36 (0.15-0.76)	0.5% (-0.01-1.3%)	0.56% (-0.042-1.5%)	990 (-41-2600)	7600 (71-20000)
Low-middle SDI	1990	Female	90 to 94	15% (7.8-26%)	85% (74-92%)	0.42 (0.16-0.9)	0.41% (-0.15-1.3%)	0.45% (-0.22-1.5%)	290 (-130-950)	1600 (-520-5000)
Low-middle SDI	1990	Female	95 plus	18% (8.5-31%)	82% (69-91%)	0.5 (0.17-1.1)	0.41% (-0.3-1.6%)	0.41% (-0.42-1.8%)	67 (-64-280)	260 (-180-980)
Low-middle SDI	1990	Male	All Ages	17% (15-20%)	83% (80-85%)	0.76 (0.6-0.96)	2.2% (1.7-2.7%)	3.1% (2.2-4.1%)	240000 (170000-320000)	10000000 (7900000-13000000)
Low-middle SDI	1990	Male	15 to 19	10% (6.9-15%)	90% (85-93%)	0.36 (0.19-0.61)	1.5% (0.93-2.2%)	1.6% (0.84-2.6%)	2100 (1100-3500)	220000 (140000-340000)
Low-middle SDI	1990	Male	20 to 24	18% (12-24%)	82% (76-88%)	0.85 (0.47-1.4)	4% (2.7-5.7%)	4.2% (2.5-6.5%)	7500 (4300-11000)	680000 (440000-970000)
Low-middle SDI	1990	Male	25 to 29	25% (18-33%)	75% (67-82%)	1.3 (0.79-2)	6% (4.2-8.2%)	6.7% (4.2-9.7%)	13000 (7800-18000)	1000000 (690000-1400000)
Low-middle SDI	1990	Male	30 to 34	31% (22-40%)	69% (60-78%)	1.6 (1-2.4)	7.5% (5.4-9.9%)	8.8% (5.9-12%)	18000 (12000-25000)	1200000 (870000-1600000)
Low-middle SDI	1990	Male	35 to 39	34% (25-42%)	66% (58-75%)	1.8 (1.1-2.7)	7.9% (5.7-11%)	9.6% (6.5-13%)	21000 (14000-29000)	1200000 (890000-1600000)
Low-middle SDI	1990	Male	40 to 44	34% (26-42%)	66% (58-74%)	1.9 (1.1-2.8)	8% (5.7-11%)	9.8% (6.7-13%)	22000 (15000-30000)	1100000 (800000-1500000)
Low-middle SDI	1990	Male	45 to 49	32% (23-40%)	68% (60-77%)	1.8 (1.1-2.9)	7.7% (5.3-11%)	9.5% (6.4-13%)	25000 (17000-35000)	1100000 (780000-1600000)
Low-middle SDI	1990	Male	50 to 54	30% (23-39%)	70% (61-77%)	1.7 (1-2.7)	6.9% (4.5-9.6%)	8.4% (5.4-12%)	27000 (17000-39000)	1100000 (690000-1500000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	1990	Male	55 to 59	29% (21-37%)	71% (63-79%)	1.6 (0.94-2.5)	6% (4-8.4%)	7.2% (4.8-10%)	28000 (18000-40000)	940000 (620000-1300000)
Low-middle SDI	1990	Male	60 to 64	27% (20-36%)	73% (64-80%)	1.4 (0.79-2.3)	4.8% (2.9-7.1%)	5.6% (3.4-8.5%)	26000 (15000-40000)	740000 (450000-1100000)
Low-middle SDI	1990	Male	65 to 69	26% (19-33%)	74% (67-81%)	1.2 (0.73-1.9)	3.8% (2.2-5.7%)	4.5% (2.6-6.7%)	22000 (12000-33000)	510000 (290000-760000)
Low-middle SDI	1990	Male	70 to 74	24% (17-31%)	76% (69-83%)	1 (0.57-1.6)	2.9% (1.5-4.6%)	3.3% (1.7-5.4%)	16000 (7900-26000)	300000 (150000-480000)
Low-middle SDI	1990	Male	75 to 79	21% (14-28%)	79% (72-86%)	0.77 (0.4-1.3)	2% (0.91-3.4%)	2.3% (0.98-3.9%)	8900 (3800-16000)	130000 (58000-220000)
Low-middle SDI	1990	Male	80 to 84	18% (12-25%)	82% (75-88%)	0.6 (0.33-0.98)	1.3% (0.55-2.4%)	1.5% (0.58-2.7%)	3800 (1500-7000)	42000 (17000-75000)
Low-middle SDI	1990	Male	85 to 89	18% (12-25%)	82% (75-88%)	0.52 (0.28-0.87)	0.95% (0.33-1.8%)	1% (0.3-2%)	1500 (430-2900)	12000 (4100-23000)
Low-middle SDI	1990	Male	90 to 94	19% (13-26%)	81% (74-87%)	0.5 (0.25-0.87)	0.67% (0.12-1.4%)	0.68% (0.042-1.6%)	360 (20-830)	2100 (380-4500)
Low-middle SDI	1990	Male	95 plus	21% (13-31%)	79% (69-87%)	0.49 (0.2-0.93)	0.59% (0.0079-1.4%)	0.54% (-0.14-1.5%)	66 (-17-190)	290 (3.4-710)
Low-middle SDI	1995	Female	All Ages	6.4% (5.4-7.6%)	94% (92-95%)	0.18 (0.14-0.23)	0.53% (0.39-0.7%)	0.86% (0.54-1.2%)	64000 (40000-93000)	2500000 (1800000-3300000)
Low-middle SDI	1995	Female	15 to 19	5.6% (3.7-8%)	94% (92-96%)	0.15 (0.073-0.26)	0.46% (0.26-0.75%)	0.46% (0.18-0.92%)	590 (220-1200)	80000 (43000-130000)
Low-middle SDI	1995	Female	20 to 24	6.9% (4.5-9.6%)	93% (90-95%)	0.23 (0.12-0.39)	0.84% (0.51-1.3%)	0.74% (0.32-1.3%)	1300 (560-2300)	160000 (95000-250000)
Low-middle SDI	1995	Female	25 to 29	7.4% (4.9-10%)	93% (90-95%)	0.27 (0.14-0.46)	1% (0.65-1.5%)	1% (0.5-1.7%)	1900 (930-3300)	190000 (120000-290000)
Low-middle SDI	1995	Female	30 to 34	8% (5.3-11%)	92% (89-95%)	0.31 (0.17-0.52)	1.2% (0.81-1.7%)	1.4% (0.8-2.2%)	2600 (1500-4100)	210000 (130000-300000)
Low-middle SDI	1995	Female	35 to 39	8.7% (5.9-12%)	91% (88-94%)	0.34 (0.18-0.56)	1.4% (0.97-2.1%)	1.8% (1.1-2.8%)	3500 (2100-5500)	230000 (150000-330000)
Low-middle SDI	1995	Female	40 to 44	9.1% (6.4-12%)	91% (88-94%)	0.35 (0.19-0.59)	1.6% (1-2.2%)	2.1% (1.3-3.1%)	4200 (2600-6400)	230000 (150000-340000)
Low-middle SDI	1995	Female	45 to 49	9.5% (6.4-13%)	90% (87-94%)	0.35 (0.17-0.61)	1.6% (1.1-2.4%)	2.2% (1.3-3.3%)	4700 (2800-7300)	230000 (140000-330000)
Low-middle SDI	1995	Female	50 to 54	9.9% (7-14%)	90% (86-93%)	0.37 (0.19-0.64)	1.7% (1.1-2.6%)	2.3% (1.4-3.5%)	6100 (3600-9500)	250000 (150000-370000)
Low-middle SDI	1995	Female	55 to 59	9.9% (6.9-14%)	90% (86-93%)	0.43 (0.21-0.74)	1.7% (1-2.7%)	2.2% (1.3-3.5%)	7600 (4300-12000)	260000 (150000-410000)
Low-middle SDI	1995	Female	60 to 64	9.8% (6.6-14%)	90% (86-93%)	0.43 (0.21-0.77)	1.5% (0.8-2.4%)	1.9% (0.97-3.1%)	7800 (4000-13000)	220000 (120000-370000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	1995	Female	65 to 69	10% (6.7-14%)	90% (86-93%)	0.4 (0.2-0.69)	1.3% (0.66-2.2%)	1.6% (0.78-2.7%)	7800 (3900-13000)	180000 (92000-300000)
Low-middle SDI	1995	Female	70 to 74	10% (6.9-15%)	90% (85-93%)	0.36 (0.18-0.65)	1.1% (0.42-2%)	1.3% (0.5-2.4%)	6600 (2600-12000)	120000 (49000-230000)
Low-middle SDI	1995	Female	75 to 79	10% (6.5-15%)	90% (85-93%)	0.34 (0.15-0.66)	0.8% (0.23-1.6%)	0.94% (0.26-2%)	4400 (1200-9100)	63000 (18000-130000)
Low-middle SDI	1995	Female	80 to 84	11% (6.5-16%)	89% (84-94%)	0.36 (0.16-0.7)	0.65% (0.14-1.4%)	0.75% (0.15-1.7%)	2700 (550-6000)	29000 (6500-63000)
Low-middle SDI	1995	Female	85 to 89	12% (7.3-19%)	88% (81-93%)	0.41 (0.18-0.81)	0.55% (0.049-1.3%)	0.63% (0.032-1.5%)	1300 (66-3300)	10000 (920-24000)
Low-middle SDI	1995	Female	90 to 94	14% (8.7-23%)	86% (77-91%)	0.49 (0.2-0.99)	0.48% (-0.067-1.4%)	0.53% (-0.12-1.6%)	430 (-99-1300)	2300 (-320-6600)
Low-middle SDI	1995	Female	95 plus	17% (9.6-28%)	83% (72-90%)	0.58 (0.2-1.3)	0.49% (-0.21-1.6%)	0.5% (-0.32-1.9%)	110 (-69-390)	410 (-180-1400)
Low-middle SDI	1995	Male	All Ages	17% (15-19%)	83% (81-85%)	0.91 (0.72-1.1)	2.6% (2.1-3.2%)	3.6% (2.7-4.6%)	300000 (220000-390000)	1300000 (1000000-1600000)
Low-middle SDI	1995	Male	15 to 19	8.5% (5.7-12%)	92% (88-94%)	0.34 (0.18-0.57)	1.4% (0.92-2.1%)	1.5% (0.85-2.5%)	2200 (1200-3500)	240000 (150000-360000)
Low-middle SDI	1995	Male	20 to 24	16% (12-21%)	84% (79-88%)	0.89 (0.51-1.4)	4% (2.7-5.5%)	4.2% (2.6-6.2%)	8300 (5000-13000)	760000 (500000-1100000)
Low-middle SDI	1995	Male	25 to 29	24% (18-30%)	76% (70-82%)	1.5 (0.92-2.2)	6% (4.3-8%)	6.7% (4.3-9.5%)	15000 (9400-21000)	1100000 (810000-1500000)
Low-middle SDI	1995	Male	30 to 34	29% (23-36%)	71% (64-77%)	1.9 (1.2-2.8)	7.6% (5.6-10%)	9% (6.2-12%)	21000 (15000-29000)	1500000 (1100000-1900000)
Low-middle SDI	1995	Male	35 to 39	33% (26-40%)	67% (60-74%)	2.2 (1.4-3.1)	8.2% (6.1-11%)	10% (7.1-13%)	26000 (18000-34000)	1500000 (1100000-2000000)
Low-middle SDI	1995	Male	40 to 44	34% (28-41%)	66% (59-72%)	2.3 (1.4-3.3)	8.5% (6.3-11%)	10% (7.5-14%)	30000 (21000-39000)	1500000 (1100000-2000000)
Low-middle SDI	1995	Male	45 to 49	33% (26-39%)	67% (61-74%)	2.3 (1.4-3.4)	8.3% (6-11%)	10% (7.1-14%)	31000 (22000-42000)	1400000 (1000000-1800000)
Low-middle SDI	1995	Male	50 to 54	31% (25-37%)	69% (63-75%)	2.2 (1.3-3.3)	7.5% (5.2-10%)	9.2% (6.2-12%)	33000 (22000-45000)	1300000 (870000-1700000)
Low-middle SDI	1995	Male	55 to 59	29% (23-36%)	71% (64-77%)	2 (1.2-3)	6.7% (4.6-9.1%)	8.1% (5.5-11%)	35000 (23000-49000)	1200000 (810000-1600000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	1995	Male	60 to 64	27% (21-34%)	73% (66-79%)	1.7 (1-2.7)	5.3% (3.5-7.6%)	6.2% (4-9%)	32000 (20000-47000)	920000 (600000-1300000)
Low-middle SDI	1995	Male	65 to 69	25% (19-31%)	75% (69-81%)	1.5 (0.88-2.2)	4.3% (2.7-6.1%)	5% (3.1-7.2%)	28000 (17000-40000)	650000 (410000-940000)
Low-middle SDI	1995	Male	70 to 74	23% (18-29%)	77% (71-82%)	1.2 (0.7-1.8)	3.2% (1.8-5%)	3.7% (2.1-5.8%)	20000 (11000-32000)	380000 (220000-590000)
Low-middle SDI	1995	Male	75 to 79	20% (15-26%)	80% (74-85%)	0.91 (0.51-1.5)	2.3% (1.1-3.7%)	2.6% (1.2-4.3%)	12000 (5600-20000)	170000 (86000-280000)
Low-middle SDI	1995	Male	80 to 84	18% (13-23%)	82% (77-87%)	0.72 (0.42-1.2)	1.6% (0.74-2.6%)	1.8% (0.8-3%)	5400 (2400-9100)	59000 (28000-98000)
Low-middle SDI	1995	Male	85 to 89	18% (13-24%)	82% (76-87%)	0.65 (0.37-1)	1.1% (0.48-2.1%)	1.3% (0.48-2.3%)	2200 (830-4100)	18000 (7300-31000)
Low-middle SDI	1995	Male	90 to 94	19% (13-25%)	81% (75-87%)	0.61 (0.33-1)	0.81% (0.24-1.6%)	0.84% (0.16-1.8%)	560 (110-1200)	3300 (970-6500)
Low-middle SDI	1995	Male	95 plus	21% (14-29%)	79% (71-86%)	0.59 (0.26-1.1)	0.7% (0.087-1.6%)	0.65% (-0.054-1.7%)	100 (-9.1-270)	450 (54-1000)
Low-middle SDI	2000	Female	All Ages	6.4% (5.4-7.5%)	94% (92-95%)	0.19 (0.14-0.24)	0.56% (0.41-0.73%)	0.87% (0.55-1.3%)	68000 (42000-100000)	2600000 (1900000-3600000)
Low-middle SDI	2000	Female	15 to 19	5.7% (3.9-8.1%)	94% (92-96%)	0.15 (0.079-0.27)	0.48% (0.27-0.76%)	0.47% (0.19-0.91%)	640 (250-1300)	89000 (50000-150000)
Low-middle SDI	2000	Female	20 to 24	6.9% (4.7-9.5%)	93% (91-95%)	0.23 (0.12-0.39)	0.84% (0.52-1.3%)	0.7% (0.32-1.3%)	1300 (600-2400)	180000 (100000-270000)
Low-middle SDI	2000	Female	25 to 29	7.4% (5.2-10%)	93% (90-95%)	0.28 (0.15-0.45)	0.98% (0.64-1.4%)	0.91% (0.46-1.5%)	1900 (990-3300)	200000 (130000-300000)
Low-middle SDI	2000	Female	30 to 34	7.8% (5.4-11%)	92% (89-95%)	0.31 (0.17-0.5)	1.1% (0.78-1.6%)	1.2% (0.75-1.9%)	2600 (1500-4200)	220000 (140000-320000)
Low-middle SDI	2000	Female	35 to 39	8.3% (5.8-11%)	92% (89-94%)	0.33 (0.18-0.53)	1.3% (0.93-1.9%)	1.7% (1.1-2.5%)	3700 (2300-5600)	250000 (170000-350000)
Low-middle SDI	2000	Female	40 to 44	8.7% (6.3-11%)	91% (89-94%)	0.34 (0.18-0.56)	1.4% (0.98-2%)	1.9% (1.2-2.8%)	4400 (2700-6600)	250000 (170000-360000)
Low-middle SDI	2000	Female	45 to 49	8.8% (6.1-12%)	91% (88-94%)	0.33 (0.17-0.57)	1.5% (0.99-2.1%)	2% (1.3-3%)	5200 (3100-8000)	250000 (160000-370000)
Low-middle SDI	2000	Female	50 to 54	9.4% (6.6-13%)	91% (87-93%)	0.36 (0.18-0.62)	1.6% (1-2.4%)	2.1% (1.3-3.2%)	6300 (3800-9900)	260000 (160000-390000)
Low-middle SDI	2000	Female	55 to 59	9.4% (6.4-13%)	91% (87-94%)	0.42 (0.21-0.73)	1.6% (0.98-2.5%)	2.1% (1.2-3.3%)	7800 (4300-12000)	260000 (150000-410000)
Low-middle SDI	2000	Female	60 to 64	9.3% (6.2-13%)	91% (87-94%)	0.42 (0.2-0.75)	1.4% (0.76-2.3%)	1.8% (0.92-2.9%)	8200 (4200-14000)	240000 (120000-390000)
Low-middle SDI	2000	Female	65 to 69	9.3% (6.2-13%)	91% (87-94%)	0.38 (0.19-0.66)	1.2% (0.65-2.1%)	1.5% (0.78-2.5%)	8500 (4300-14000)	200000 (100000-330000)
Low-middle SDI	2000	Female	70 to 74	9.6% (6.4-14%)	90% (86-94%)	0.35 (0.17-0.64)	1% (0.41-1.8%)	1.2% (0.48-2.2%)	7200 (2800-13000)	130000 (54000-250000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2000	Female	75 to 79	9.7% (6.1-15%)	90% (85-94%)	0.33 (0.15-0.64)	0.76% (0.22-1.5%)	0.9% (0.25-1.8%)	4800 (1300-9800)	69000 (20000-140000)
Low-middle SDI	2000	Female	80 to 84	9.9% (5.9-15%)	90% (85-94%)	0.34 (0.15-0.68)	0.62% (0.15-1.3%)	0.72% (0.16-1.6%)	3000 (660-6500)	32000 (7800-69000)
Low-middle SDI	2000	Female	85 to 89	11% (6.6-18%)	89% (82-93%)	0.39 (0.17-0.8)	0.53% (0.061-1.2%)	0.6% (0.051-1.4%)	1500 (120-3600)	12000 (1300-27000)
Low-middle SDI	2000	Female	90 to 94	13% (7.8-21%)	87% (79-92%)	0.47 (0.2-0.94)	0.46% (-0.048-1.3%)	0.51% (-0.099-1.5%)	500 (-100-1400)	2800 (-290-7600)
Low-middle SDI	2000	Female	95 plus	16% (8.6-27%)	84% (73-91%)	0.56 (0.2-1.2)	0.46% (-0.17-1.5%)	0.47% (-0.27-1.7%)	130 (-76-460)	490 (-190-1600)
Low-middle SDI	2000	Male	All Ages	18% (16-20%)	82% (80-84%)	1 (0.83-1.3)	2.9% (2.4-3.5%)	3.8% (2.9-4.8%)	340000 (250000-430000)	1500000 (1200000-1800000)
Low-middle SDI	2000	Male	15 to 19	11% (8.1-15%)	89% (85-92%)	0.48 (0.27-0.78)	1.8% (1.2-2.6%)	2% (1.1-3.1%)	3200 (1800-5100)	340000 (220000-500000)
Low-middle SDI	2000	Male	20 to 24	21% (16-26%)	79% (74-84%)	1.3 (0.77-1.9)	4.9% (3.5-6.7%)	5.3% (3.4-7.7%)	12000 (7400-17000)	1000000 (720000-1400000)
Low-middle SDI	2000	Male	25 to 29	28% (22-33%)	72% (67-78%)	1.9 (1.2-2.7)	6.7% (4.9-8.8%)	7.4% (5-10%)	19000 (13000-26000)	1500000 (1100000-1900000)
Low-middle SDI	2000	Male	30 to 34	31% (25-37%)	69% (63-75%)	2.1 (1.4-3.1)	7.7% (5.9-9.9%)	8.9% (6.4-12%)	25000 (18000-34000)	1700000 (1300000-2200000)
Low-middle SDI	2000	Male	35 to 39	33% (27-38%)	67% (62-73%)	2.3 (1.5-3.3)	8% (6-10%)	9.5% (6.9-12%)	30000 (21000-39000)	1800000 (1300000-2200000)
Low-middle SDI	2000	Male	40 to 44	33% (27-39%)	67% (61-73%)	2.4 (1.5-3.4)	8.1% (6.1-10%)	9.9% (7.2-13%)	34000 (24000-44000)	1800000 (1300000-2200000)
Low-middle SDI	2000	Male	45 to 49	32% (26-38%)	68% (62-74%)	2.4 (1.5-3.5)	8% (5.9-11%)	9.8% (7-13%)	38000 (27000-50000)	1700000 (1200000-2200000)
Low-middle SDI	2000	Male	50 to 54	30% (24-36%)	70% (64-76%)	2.2 (1.3-3.3)	7.3% (5.1-9.7%)	8.9% (6-12%)	36000 (24000-49000)	1400000 (970000-1900000)
Low-middle SDI	2000	Male	55 to 59	27% (21-33%)	73% (67-79%)	1.9 (1.2-3)	6.3% (4.4-8.5%)	7.6% (5.2-10%)	35000 (23000-49000)	1200000 (820000-1600000)
Low-middle SDI	2000	Male	60 to 64	24% (18-31%)	76% (69-82%)	1.6 (0.94-2.6)	5% (3.3-7.1%)	5.8% (3.8-8.4%)	33000 (21000-48000)	950000 (620000-1400000)
Low-middle SDI	2000	Male	65 to 69	22% (16-28%)	78% (72-84%)	1.4 (0.81-2.1)	4% (2.6-5.7%)	4.7% (3-6.7%)	29000 (18000-42000)	680000 (440000-980000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2000	Male	70 to 74	20% (15-26%)	80% (74-85%)	1.1 (0.64-1.8)	3% (1.8-4.6%)	3.5% (2-5.4%)	21000 (12000-34000)	410000 (240000-630000)
Low-middle SDI	2000	Male	75 to 79	17% (12-23%)	83% (77-88%)	0.82 (0.44-1.4)	2.1% (1.1-3.4%)	2.4% (1.2-3.9%)	13000 (6300-21000)	190000 (97000-300000)
Low-middle SDI	2000	Male	80 to 84	15% (9.9-20%)	85% (80-90%)	0.63 (0.35-1)	1.4% (0.73-2.4%)	1.6% (0.77-2.7%)	5900 (2800-10000)	65000 (32000-110000)
Low-middle SDI	2000	Male	85 to 89	15% (9.8-21%)	85% (79-90%)	0.57 (0.31-0.94)	1.1% (0.48-1.9%)	1.2% (0.48-2.1%)	2500 (990-4500)	20000 (8700-35000)
Low-middle SDI	2000	Male	90 to 94	16% (11-22%)	84% (78-89%)	0.54 (0.28-0.94)	0.77% (0.26-1.5%)	0.8% (0.2-1.7%)	650 (160-1400)	3800 (1300-7400)
Low-middle SDI	2000	Male	95 plus	18% (11-26%)	82% (74-89%)	0.53 (0.22-0.99)	0.67% (0.12-1.5%)	0.62% (-0.0097-1.6%)	130 (-3-320)	540 (95-1200)
Low-middle SDI	2005	Female	All Ages	6.5% (5.5-7.7%)	93% (92-95%)	0.2 (0.15-0.25)	0.57% (0.42-0.75%)	0.86% (0.54-1.2%)	67000 (41000-100000)	2600000 (1900000-3500000)
Low-middle SDI	2005	Female	15 to 19	5.8% (3.9-8.2%)	94% (92-96%)	0.16 (0.079-0.27)	0.45% (0.27-0.71%)	0.44% (0.18-0.83%)	570 (230-1100)	86000 (49000-140000)
Low-middle SDI	2005	Female	20 to 24	6.9% (4.8-9.6%)	93% (90-95%)	0.24 (0.12-0.41)	0.82% (0.52-1.2%)	0.68% (0.32-1.2%)	1200 (580-2200)	180000 (110000-270000)
Low-middle SDI	2005	Female	25 to 29	7.5% (5.3-10%)	92% (90-95%)	0.29 (0.16-0.47)	0.92% (0.62-1.3%)	0.85% (0.45-1.4%)	1800 (940-3100)	200000 (130000-290000)
Low-middle SDI	2005	Female	30 to 34	8.1% (5.6-11%)	92% (89-94%)	0.32 (0.18-0.52)	1% (0.71-1.4%)	1.1% (0.71-1.8%)	2500 (1400-3900)	210000 (140000-310000)
Low-middle SDI	2005	Female	35 to 39	8.4% (5.9-11%)	92% (89-94%)	0.34 (0.19-0.56)	1.2% (0.85-1.7%)	1.6% (1-2.3%)	3500 (2100-5400)	240000 (160000-340000)
Low-middle SDI	2005	Female	40 to 44	8.6% (6.2-11%)	91% (89-94%)	0.35 (0.18-0.57)	1.3% (0.88-1.8%)	1.7% (1.2-2.5%)	4300 (2700-6400)	250000 (160000-350000)
Low-middle SDI	2005	Female	45 to 49	8.8% (6.1-12%)	91% (88-94%)	0.33 (0.17-0.58)	1.3% (0.88-1.9%)	1.8% (1.2-2.7%)	5100 (3100-7800)	250000 (160000-360000)
Low-middle SDI	2005	Female	50 to 54	8.9% (6.3-12%)	91% (88-94%)	0.36 (0.18-0.62)	1.4% (0.92-2.1%)	1.9% (1.2-2.9%)	6300 (3700-9900)	260000 (160000-390000)
Low-middle SDI	2005	Female	55 to 59	9.3% (6.4-13%)	91% (87-94%)	0.43 (0.22-0.75)	1.5% (0.9-2.3%)	2% (1.1-3.1%)	7500 (4100-12000)	260000 (150000-410000)
Low-middle SDI	2005	Female	60 to 64	9.1% (6.1-13%)	91% (87-94%)	0.43 (0.21-0.77)	1.3% (0.71-2.1%)	1.7% (0.87-2.7%)	7800 (3900-13000)	230000 (120000-370000)
Low-middle SDI	2005	Female	65 to 69	9.2% (6.2-13%)	91% (87-94%)	0.39 (0.2-0.67)	1.1% (0.59-1.9%)	1.4% (0.71-2.4%)	8400 (4100-14000)	200000 (100000-330000)
Low-middle SDI	2005	Female	70 to 74	9.2% (6.2-13%)	91% (87-94%)	0.35 (0.17-0.64)	0.93% (0.37-1.7%)	1.1% (0.44-2.1%)	7400 (2800-14000)	140000 (55000-250000)
Low-middle SDI	2005	Female	75 to 79	9.2% (5.8-14%)	91% (86-94%)	0.33 (0.15-0.64)	0.71% (0.2-1.4%)	0.85% (0.23-1.7%)	5100 (1300-10000)	74000 (20000-150000)
Low-middle SDI	2005	Female	80 to 84	9.4% (5.6-15%)	91% (85-94%)	0.34 (0.15-0.66)	0.59% (0.14-1.2%)	0.69% (0.15-1.5%)	3300 (650-7200)	36000 (7800-77000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2005	Female	85 to 89	10% (6.2-17%)	90% (83-94%)	0.38 (0.17-0.78)	0.52% (0.066-1.2%)	0.59% (0.053-1.4%)	1800 (130-4200)	14000 (1600-32000)
Low-middle SDI	2005	Female	90 to 94	12% (7.3-20%)	88% (80-93%)	0.46 (0.19-0.91)	0.46% (-0.029-1.2%)	0.5% (-0.078-1.4%)	600 (-100-1700)	3300 (-210-8800)
Low-middle SDI	2005	Female	95 plus	15% (8-25%)	85% (75-92%)	0.55 (0.2-1.2)	0.47% (-0.15-1.5%)	0.48% (-0.26-1.7%)	150 (-85-540)	600 (-200-1900)
Low-middle SDI	2005	Male	All Ages	19% (17-21%)	81% (79-83%)	1.1 (0.89-1.4)	3.2% (2.6-3.9%)	4% (3.1-5%)	370000 (280000-460000)	1600000 (1300000-1900000)
Low-middle SDI	2005	Male	15 to 19	11% (8.1-15%)	89% (85-92%)	0.48 (0.27-0.79)	1.8% (1.2-2.5%)	1.9% (1.1-3%)	3200 (1800-5200)	35000 (230000-510000)
Low-middle SDI	2005	Male	20 to 24	21% (16-26%)	79% (74-84%)	1.3 (0.79-1.9)	5% (3.6-6.7%)	5.4% (3.5-7.8%)	13000 (8100-18000)	110000 (78000-150000)
Low-middle SDI	2005	Male	25 to 29	28% (22-33%)	72% (67-78%)	1.9 (1.2-2.7)	6.6% (4.9-8.6%)	7.3% (5-10%)	20000 (14000-27000)	160000 (110000-200000)
Low-middle SDI	2005	Male	30 to 34	31% (25-38%)	69% (62-75%)	2.2 (1.5-3.2)	7.5% (5.7-9.5%)	8.6% (6.3-11%)	27000 (19000-35000)	180000 (140000-230000)
Low-middle SDI	2005	Male	35 to 39	33% (27-39%)	67% (61-73%)	2.4 (1.6-3.4)	7.8% (6-9.9%)	9.2% (6.8-12%)	32000 (24000-42000)	190000 (150000-240000)
Low-middle SDI	2005	Male	40 to 44	33% (27-39%)	67% (61-73%)	2.4 (1.6-3.5)	7.9% (6-10%)	9.5% (7.1-12%)	36000 (27000-47000)	190000 (140000-240000)
Low-middle SDI	2005	Male	45 to 49	32% (26-38%)	68% (62-74%)	2.4 (1.5-3.6)	7.8% (5.8-10%)	9.5% (6.9-12%)	40000 (29000-54000)	180000 (140000-240000)
Low-middle SDI	2005	Male	50 to 54	30% (24-36%)	70% (64-76%)	2.3 (1.4-3.4)	7.2% (5.2-9.5%)	8.7% (6.1-12%)	42000 (29000-57000)	170000 (120000-220000)
Low-middle SDI	2005	Male	55 to 59	27% (22-34%)	73% (66-78%)	2 (1.3-3.1)	6.4% (4.5-8.5%)	7.7% (5.4-10%)	38000 (26000-52000)	130000 (91000-170000)
Low-middle SDI	2005	Male	60 to 64	25% (19-32%)	75% (68-81%)	1.7 (1-2.7)	5.1% (3.4-7.1%)	6% (3.9-8.4%)	35000 (22000-50000)	100000 (66000-140000)
Low-middle SDI	2005	Male	65 to 69	23% (17-29%)	77% (71-83%)	1.5 (0.88-2.2)	4% (2.6-5.7%)	4.7% (3-6.7%)	31000 (20000-46000)	74000 (48000-110000)
Low-middle SDI	2005	Male	70 to 74	21% (16-27%)	79% (73-84%)	1.2 (0.69-1.9)	3% (1.8-4.6%)	3.5% (2-5.4%)	24000 (13000-36000)	45000 (26000-69000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2005	Male	75 to 79	18% (13-24%)	82% (76-87%)	0.89 (0.48-1.4)	2.1% (1.1-3.4%)	2.4% (1.2-3.9%)	14000 (7100-24000)	210000 (110000-350000)
Low-middle SDI	2005	Male	80 to 84	15% (10-21%)	85% (79-90%)	0.66 (0.37-1.1)	1.5% (0.73-2.4%)	1.6% (0.77-2.8%)	7100 (3300-12000)	79000 (39000-130000)
Low-middle SDI	2005	Male	85 to 89	15% (9.7-21%)	85% (79-90%)	0.59 (0.32-0.99)	1.1% (0.49-1.9%)	1.2% (0.5-2.1%)	3000 (1200-5500)	24000 (11000-42000)
Low-middle SDI	2005	Male	90 to 94	16% (11-22%)	84% (78-89%)	0.57 (0.29-0.97)	0.8% (0.27-1.6%)	0.82% (0.2-1.7%)	800 (190-1700)	4700 (1600-9300)
Low-middle SDI	2005	Male	95 plus	18% (11-26%)	82% (74-89%)	0.56 (0.24-1)	0.63% (0.14-1.6%)	0.63% (-0.011-1.6%)	150 (-3.8-400)	670 (130-1500)
Low-middle SDI	2010	Female	All Ages	7.3% (6.1-8.7%)	93% (91-94%)	0.22 (0.16-0.29)	0.63% (0.47-0.83%)	0.91% (0.57-1.3%)	69000 (42000-100000)	2700000 (1900000-3600000)
Low-middle SDI	2010	Female	15 to 19	6.2% (4.1-8.9%)	94% (91-96%)	0.17 (0.078-0.31)	0.47% (0.28-0.73%)	0.43% (0.17-0.82%)	510 (200-1000)	85000 (49000-140000)
Low-middle SDI	2010	Female	20 to 24	7.2% (4.8-10%)	93% (90-95%)	0.25 (0.12-0.45)	0.88% (0.57-1.3%)	0.69% (0.33-1.2%)	1100 (510-2000)	180000 (110000-270000)
Low-middle SDI	2010	Female	25 to 29	7.8% (5.2-11%)	92% (89-95%)	0.31 (0.16-0.53)	0.99% (0.67-1.4%)	0.9% (0.48-1.5%)	1700 (890-2900)	210000 (140000-310000)
Low-middle SDI	2010	Female	30 to 34	8.2% (5.6-11%)	92% (89-94%)	0.34 (0.18-0.57)	1.1% (0.73-1.5%)	1.2% (0.7-1.8%)	2300 (1300-3700)	220000 (140000-310000)
Low-middle SDI	2010	Female	35 to 39	8.7% (5.9-12%)	91% (88-94%)	0.36 (0.19-0.62)	1.2% (0.83-1.6%)	1.5% (0.99-2.2%)	3300 (2000-5000)	240000 (160000-330000)
Low-middle SDI	2010	Female	40 to 44	8.7% (6.1-12%)	91% (88-94%)	0.36 (0.18-0.62)	1.2% (0.86-1.7%)	1.7% (1.1-2.5%)	4000 (2500-6100)	240000 (160000-350000)
Low-middle SDI	2010	Female	45 to 49	8.7% (5.8-12%)	91% (88-94%)	0.35 (0.17-0.62)	1.3% (0.87-1.8%)	1.8% (1.1-2.7%)	5000 (3100-7700)	250000 (160000-370000)
Low-middle SDI	2010	Female	50 to 54	9.1% (6.2-13%)	91% (87-94%)	0.38 (0.18-0.69)	1.4% (0.9-2.1%)	1.9% (1.2-2.9%)	6600 (3900-10000)	270000 (170000-410000)
Low-middle SDI	2010	Female	55 to 59	9.2% (6.2-13%)	91% (87-94%)	0.46 (0.22-0.83)	1.5% (0.86-2.3%)	2% (1.1-3.1%)	8000 (4400-13000)	280000 (160000-440000)
Low-middle SDI	2010	Female	60 to 64	9.6% (6.3-14%)	90% (86-94%)	0.48 (0.23-0.9)	1.3% (0.7-2.2%)	1.7% (0.87-2.9%)	8200 (4100-14000)	240000 (120000-400000)
Low-middle SDI	2010	Female	65 to 69	9.5% (6.2-14%)	90% (86-94%)	0.43 (0.21-0.77)	1.2% (0.57-1.9%)	1.5% (0.7-2.5%)	8500 (4100-15000)	200000 (100000-340000)
Low-middle SDI	2010	Female	70 to 74	9.1% (6-13%)	91% (87-94%)	0.37 (0.17-0.7)	0.91% (0.37-1.7%)	1.1% (0.44-2.1%)	7700 (3000-14000)	150000 (58000-270000)
Low-middle SDI	2010	Female	75 to 79	8.3% (5.1-13%)	92% (87-95%)	0.32 (0.14-0.64)	0.69% (0.21-1.4%)	0.83% (0.24-1.7%)	5500 (1600-11000)	81000 (24000-160000)
Low-middle SDI	2010	Female	80 to 84	7.8% (4.5-13%)	92% (87-96%)	0.3 (0.13-0.61)	0.57% (0.15-1.2%)	0.67% (0.16-1.4%)	3800 (850-8200)	41000 (10000-87000)
Low-middle SDI	2010	Female	85 to 89	8.4% (4.8-14%)	92% (86-95%)	0.32 (0.14-0.65)	0.47% (0.067-1.1%)	0.54% (0.052-1.3%)	2000 (160-4700)	16000 (2000-36000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2010	Female	90 to 94	9.5% (5.5-16%)	90% (84-95%)	0.37 (0.15-0.75)	0.42% (-0.0035-1.1%)	0.46% (-0.048-1.3%)	700 (-77-1900)	3900 (-41-10000)
Low-middle SDI	2010	Female	95 plus	11% (5.9-20%)	89% (80-94%)	0.43 (0.15-0.98)	0.43% (-0.098-1.3%)	0.44% (-0.19-1.4%)	180 (-79-590)	720 (-160-2100)
Low-middle SDI	2010	Male	All Ages	20% (18-22%)	80% (78-82%)	1.2 (0.93-1.6)	3.5% (2.8-4.2%)	4.2% (3.3-5.3%)	390000 (290000-490000)	1700000 (1300000-2000000)
Low-middle SDI	2010	Male	15 to 19	12% (8.5-16%)	88% (84-91%)	0.51 (0.28-0.85)	1.7% (1.1-2.5%)	1.9% (1.1-3.1%)	3100 (1700-5100)	34000 (22000-50000)
Low-middle SDI	2010	Male	20 to 24	22% (18-27%)	78% (73-82%)	1.4 (0.83-2.1)	5% (3.5-6.7%)	5.6% (3.6-8%)	12000 (7900-18000)	110000 (77000-150000)
Low-middle SDI	2010	Male	25 to 29	29% (24-34%)	71% (66-76%)	2 (1.3-3)	6.8% (5-8.8%)	7.8% (5.3-11%)	21000 (14000-28000)	160000 (120000-210000)
Low-middle SDI	2010	Male	30 to 34	32% (26-37%)	68% (63-74%)	2.3 (1.5-3.4)	7.5% (5.8-9.5%)	8.7% (6.4-11%)	26000 (19000-35000)	180000 (140000-230000)
Low-middle SDI	2010	Male	35 to 39	33% (28-39%)	67% (61-72%)	2.4 (1.6-3.5)	7.7% (5.9-9.7%)	9.2% (6.8-12%)	32000 (24000-42000)	190000 (150000-250000)
Low-middle SDI	2010	Male	40 to 44	33% (28-38%)	67% (62-72%)	2.5 (1.6-3.6)	7.6% (5.8-9.6%)	9.3% (6.9-12%)	36000 (26000-47000)	190000 (140000-240000)
Low-middle SDI	2010	Male	45 to 49	32% (26-37%)	68% (63-74%)	2.4 (1.5-3.7)	7.5% (5.6-9.7%)	9.2% (6.7-12%)	41000 (30000-55000)	190000 (140000-250000)
Low-middle SDI	2010	Male	50 to 54	30% (25-36%)	70% (64-75%)	2.3 (1.4-3.6)	7% (5-9.1%)	8.5% (6-11%)	44000 (31000-59000)	170000 (120000-230000)
Low-middle SDI	2010	Male	55 to 59	29% (23-35%)	71% (65-77%)	2.2 (1.3-3.4)	6.3% (4.4-8.3%)	7.6% (5.3-10%)	44000 (30000-60000)	150000 (110000-200000)
Low-middle SDI	2010	Male	60 to 64	28% (21-35%)	72% (65-79%)	1.9 (1.2-3.1)	5.1% (3.4-7.2%)	6.1% (3.9-8.6%)	38000 (24000-54000)	110000 (71000-150000)
Low-middle SDI	2010	Male	65 to 69	26% (20-33%)	74% (67-80%)	1.7 (1-2.6)	4.1% (2.6-5.9%)	4.8% (3-6.9%)	33000 (20000-48000)	79000 (50000-110000)
Low-middle SDI	2010	Male	70 to 74	24% (18-31%)	76% (69-82%)	1.4 (0.82-2.2)	3.1% (1.7-4.7%)	3.6% (2-5.6%)	27000 (14000-42000)	51000 (28000-79000)
Low-middle SDI	2010	Male	75 to 79	21% (15-28%)	79% (72-85%)	1.1 (0.6-1.8)	2.2% (1.1-3.6%)	2.5% (1.2-4.1%)	17000 (7700-28000)	25000 (12000-41000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2010	Male	80 to 84	18% (12-24%)	82% (76-88%)	0.83 (0.45-1.3)	1.5% (0.69-2.5%)	1.7% (0.73-2.9%)	8800 (3800-15000)	98000 (44000-170000)
Low-middle SDI	2010	Male	85 to 89	17% (11-25%)	83% (75-89%)	0.7 (0.37-1.2)	1.1% (0.45-2%)	1.2% (0.42-2.2%)	3800 (1300-7100)	31000 (12000-55000)
Low-middle SDI	2010	Male	90 to 94	18% (12-26%)	82% (74-88%)	0.68 (0.34-1.2)	0.79% (0.2-1.6%)	0.79% (0.1-1.8%)	990 (120-2200)	6000 (1500-12000)
Low-middle SDI	2010	Male	95 plus	22% (13-32%)	78% (68-87%)	0.69 (0.29-1.3)	0.68% (0.035-1.6%)	0.6% (-0.15-1.7%)	180 (-49-530)	830 (37-2000)
Low-middle SDI	2016	Female	All Ages	8% (6.6-9.7%)	92% (90-93%)	0.26 (0.17-0.4)	0.82% (0.59-1.1%)	1.1% (0.69-1.7%)	83000 (49000-130000)	320000 (230000-440000)
Low-middle SDI	2016	Female	15 to 19	7% (4.5-10%)	93% (90-96%)	0.19 (0.075-0.38)	0.51% (0.3-0.83%)	0.47% (0.17-0.99%)	520 (190-1100)	90000 (51000-150000)
Low-middle SDI	2016	Female	20 to 24	8.1% (5.1-12%)	92% (88-95%)	0.29 (0.12-0.57)	1% (0.64-1.5%)	0.82% (0.37-1.6%)	1200 (490-2200)	190000 (120000-300000)
Low-middle SDI	2016	Female	25 to 29	8.6% (5.5-12%)	91% (88-94%)	0.36 (0.16-0.68)	1.2% (0.78-1.7%)	1.1% (0.57-2%)	1800 (900-3400)	230000 (150000-350000)
Low-middle SDI	2016	Female	30 to 34	9% (5.8-13%)	91% (87-94%)	0.42 (0.19-0.78)	1.3% (0.86-1.9%)	1.5% (0.87-2.5%)	2700 (1500-4600)	260000 (170000-390000)
Low-middle SDI	2016	Female	35 to 39	9.4% (6.1-13%)	91% (87-94%)	0.45 (0.2-0.82)	1.4% (0.97-2%)	2% (1.2-3%)	3900 (2200-6200)	290000 (180000-420000)
Low-middle SDI	2016	Female	40 to 44	9.6% (6.5-13%)	90% (87-94%)	0.45 (0.2-0.82)	1.5% (0.97-2.1%)	2.1% (1.3-3.2%)	4800 (2800-7500)	290000 (190000-430000)
Low-middle SDI	2016	Female	45 to 49	9.5% (6.1-13%)	90% (87-94%)	0.42 (0.19-0.81)	1.5% (0.95-2.2%)	2.1% (1.3-3.3%)	5800 (3400-9300)	300000 (190000-440000)
Low-middle SDI	2016	Female	50 to 54	9.4% (6.2-13%)	91% (87-94%)	0.44 (0.19-0.84)	1.6% (0.98-2.4%)	2.2% (1.3-3.5%)	7800 (4600-13000)	330000 (200000-510000)
Low-middle SDI	2016	Female	55 to 59	9.6% (6.2-14%)	90% (86-94%)	0.51 (0.23-0.99)	1.6% (0.91-2.6%)	2.2% (1.2-3.6%)	9500 (5100-16000)	330000 (190000-540000)
Low-middle SDI	2016	Female	60 to 64	9.7% (6.1-15%)	90% (85-94%)	0.54 (0.24-1.1)	1.4% (0.72-2.4%)	1.9% (0.94-3.3%)	10000 (5000-18000)	300000 (150000-510000)
Low-middle SDI	2016	Female	65 to 69	9.8% (6.3-14%)	90% (86-94%)	0.49 (0.22-0.94)	1.3% (0.61-2.2%)	1.6% (0.77-2.8%)	10000 (4800-18000)	240000 (120000-420000)
Low-middle SDI	2016	Female	70 to 74	9.3% (6-14%)	91% (86-94%)	0.41 (0.18-0.83)	0.99% (0.4-1.9%)	1.2% (0.48-2.4%)	8700 (3400-17000)	170000 (66000-310000)
Low-middle SDI	2016	Female	75 to 79	8.4% (5.1-13%)	92% (87-95%)	0.35 (0.14-0.73)	0.74% (0.23-1.5%)	0.9% (0.27-1.9%)	6600 (1900-14000)	96000 (29000-200000)
Low-middle SDI	2016	Female	80 to 84	7.7% (4.3-13%)	92% (87-96%)	0.34 (0.14-0.7)	0.63% (0.18-1.3%)	0.75% (0.2-1.6%)	5000 (1300-11000)	54000 (15000-120000)
Low-middle SDI	2016	Female	85 to 89	8.2% (4.5-14%)	92% (86-95%)	0.36 (0.14-0.75)	0.55% (0.12-1.2%)	0.64% (0.11-1.5%)	3000 (480-6900)	23000 (4700-53000)
Low-middle SDI	2016	Female	90 to 94	9.3% (5.2-16%)	91% (84-95%)	0.4 (0.15-0.86)	0.49% (0.035-1.3%)	0.54% (-0.0036-1.5%)	1100 (-13-3100)	6300 (450-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2016	Female	95 plus	11% (5.5-19%)	89% (81-95%)	0.47 (0.15-1.1)	0.52% (-0.048-1.4%)	0.54% (-0.13-1.6%)	320 (-78-960)	1200 (-110-3400)
Low-middle SDI	2016	Male	All Ages	20% (17-22%)	80% (78-83%)	1.3 (0.94-1.7)	3.9% (3.2-4.7%)	4.6% (3.5-5.9%)	430000 (320000-550000)	1800000 (1400000-2200000)
Low-middle SDI	2016	Male	15 to 19	10% (6.9-15%)	90% (85-93%)	0.46 (0.22-0.85)	1.6% (0.99-2.4%)	1.7% (0.89-2.9%)	2700 (1300-4700)	300000 (190000-470000)
Low-middle SDI	2016	Male	20 to 24	18% (13-24%)	82% (76-87%)	1.2 (0.64-2)	4.4% (3-6.2%)	4.9% (2.9-7.4%)	11000 (6200-16000)	990000 (650000-1400000)
Low-middle SDI	2016	Male	25 to 29	26% (19-32%)	74% (68-81%)	1.9 (1.1-3)	6.3% (4.5-8.3%)	7.4% (4.9-10%)	19000 (12000-26000)	1500000 (1100000-2000000)
Low-middle SDI	2016	Male	30 to 34	29% (22-37%)	71% (63-78%)	2.3 (1.4-3.6)	7.3% (5.5-9.4%)	8.9% (6.4-12%)	26000 (18000-36000)	1800000 (1400000-2400000)
Low-middle SDI	2016	Male	35 to 39	32% (24-39%)	68% (61-76%)	2.5 (1.5-3.9)	7.7% (5.9-9.9%)	9.6% (7-12%)	33000 (24000-44000)	2000000 (1500000-2600000)
Low-middle SDI	2016	Male	40 to 44	32% (25-39%)	68% (61-75%)	2.6 (1.6-4)	7.8% (5.9-9.9%)	9.8% (7.2-13%)	37000 (27000-49000)	2000000 (1500000-2600000)
Low-middle SDI	2016	Male	45 to 49	31% (24-38%)	69% (62-76%)	2.6 (1.5-4.1)	7.7% (5.7-10%)	9.7% (7-13%)	44000 (31000-58000)	2000000 (1500000-2600000)
Low-middle SDI	2016	Male	50 to 54	30% (24-37%)	70% (63-76%)	2.5 (1.4-4)	7.1% (5.1-9.3%)	8.9% (6.2-12%)	48000 (33000-64000)	1900000 (1400000-2500000)
Low-middle SDI	2016	Male	55 to 59	29% (23-36%)	71% (64-77%)	2.4 (1.4-3.8)	6.5% (4.5-8.6%)	8% (5.5-11%)	51000 (35000-69000)	1800000 (1200000-2300000)
Low-middle SDI	2016	Male	60 to 64	28% (22-36%)	72% (64-78%)	2.2 (1.3-3.5)	5.3% (3.5-7.5%)	6.4% (4.1-9.1%)	47000 (30000-68000)	1400000 (890000-1900000)
Low-middle SDI	2016	Male	65 to 69	27% (21-34%)	73% (66-79%)	1.9 (1.2-3)	4.3% (2.7-6.2%)	5.1% (3.1-7.4%)	39000 (24000-58000)	940000 (580000-1400000)
Low-middle SDI	2016	Male	70 to 74	25% (19-32%)	75% (68-81%)	1.6 (0.89-2.5)	3.3% (1.8-5.1%)	3.8% (2-6%)	30000 (16000-48000)	580000 (320000-900000)
Low-middle SDI	2016	Male	75 to 79	22% (16-29%)	78% (71-84%)	1.2 (0.66-2)	2.3% (1.1-3.9%)	2.7% (1.2-4.5%)	20000 (9300-34000)	300000 (150000-500000)
Low-middle SDI	2016	Male	80 to 84	18% (12-25%)	82% (75-88%)	0.95 (0.52-1.6)	1.6% (0.75-2.8%)	1.8% (0.78-3.2%)	12000 (5000-21000)	130000 (60000-220000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low-middle SDI	2016	Male	85 to 89	18% (11-25%)	82% (75-89%)	0.81 (0.43-1.4)	1.2% (0.5-2.2%)	1.3% (0.48-2.4%)	5300 (2000-9900)	43000 (18000-77000)
Low-middle SDI	2016	Male	90 to 94	19% (13-26%)	81% (74-87%)	0.77 (0.39-1.4)	0.88% (0.22-1.8%)	0.91% (0.12-2%)	1500 (210-3400)	9100 (2200-19000)
Low-middle SDI	2016	Male	95 plus	22% (13-31%)	78% (69-87%)	0.77 (0.33-1.5)	0.77% (0.054-1.8%)	0.7% (-0.14-1.9%)	290 (-60-810)	1300 (87-3100)
Low SDI	1990	Female	All Ages	24% (20-29%)	76% (71-80%)	0.62 (0.47-0.81)	1% (0.59-1.5%)	1.9% (0.96-2.9%)	41000 (21000-64000)	1500000 (870000-2200000)
Low SDI	1990	Female	15 to 19	24% (15-35%)	76% (65-85%)	0.68 (0.35-1.2)	1.5% (0.65-2.5%)	1.8% (0.6-3.4%)	670 (210-1300)	61000 (27000-110000)
Low SDI	1990	Female	20 to 24	30% (19-43%)	70% (57-81%)	1 (0.53-1.7)	2.3% (1.1-3.9%)	2.5% (0.89-4.7%)	1200 (420-2300)	110000 (51000-180000)
Low SDI	1990	Female	25 to 29	33% (21-46%)	67% (54-79%)	1.2 (0.69-2)	3% (1.5-4.8%)	3.4% (1.4-5.9%)	1700 (720-3000)	130000 (66000-210000)
Low SDI	1990	Female	30 to 34	35% (23-49%)	65% (51-77%)	1.4 (0.75-2.2)	3.4% (1.8-5.4%)	4.1% (1.9-6.9%)	2100 (960-3600)	140000 (71000-220000)
Low SDI	1990	Female	35 to 39	38% (25-52%)	62% (48-75%)	1.3 (0.74-2.2)	3.6% (1.8-5.8%)	4.5% (2-7.5%)	2500 (1100-4100)	140000 (68000-220000)
Low SDI	1990	Female	40 to 44	39% (27-53%)	61% (47-73%)	1.3 (0.7-2.2)	3.7% (1.8-5.9%)	4.7% (2.2-7.7%)	2600 (1200-4300)	130000 (63000-210000)
Low SDI	1990	Female	45 to 49	39% (25-53%)	61% (47-75%)	1.3 (0.68-2.3)	4.1% (1.9-6.7%)	5.3% (2.4-8.8%)	3100 (1300-5300)	130000 (60000-220000)
Low SDI	1990	Female	50 to 54	37% (25-52%)	63% (48-75%)	1.3 (0.67-2.3)	4.6% (2.1-7.6%)	5.8% (2.5-9.7%)	3900 (1700-6700)	140000 (64000-250000)
Low SDI	1990	Female	55 to 59	36% (24-51%)	64% (49-76%)	1.3 (0.67-2.3)	4.7% (2-7.9%)	5.8% (2.4-9.7%)	4600 (1900-8000)	150000 (60000-250000)
Low SDI	1990	Female	60 to 64	37% (23-52%)	63% (48-77%)	1.3 (0.65-2.4)	4.2% (1.7-7.4%)	5.1% (2.1-8.9%)	4700 (1900-8300)	130000 (51000-220000)
Low SDI	1990	Female	65 to 69	37% (25-51%)	63% (49-75%)	1.2 (0.63-2.1)	3.9% (1.4-6.9%)	4.6% (1.7-8.1%)	5000 (1800-8900)	110000 (41000-200000)
Low SDI	1990	Female	70 to 74	36% (24-50%)	64% (50-76%)	1 (0.51-1.9)	3.2% (0.75-6.2%)	3.7% (0.84-7.2%)	4000 (870-7800)	71000 (16000-140000)
Low SDI	1990	Female	75 to 79	33% (21-48%)	67% (52-79%)	0.87 (0.39-1.6)	2.4% (0.28-5.1%)	2.8% (0.31-5.9%)	2600 (290-5500)	36000 (4100-76000)
Low SDI	1990	Female	80 to 84	32% (19-47%)	68% (53-81%)	0.79 (0.37-1.4)	1.9% (0.11-4.1%)	2.2% (0.11-4.8%)	1300 (59-2800)	13000 (690-29000)
Low SDI	1990	Female	85 to 89	31% (19-46%)	69% (54-81%)	0.76 (0.35-1.4)	1.4% (-0.095-3.5%)	1.7% (-0.15-4.1%)	510 (-43-1200)	3700 (-260-9100)
Low SDI	1990	Female	90 to 94	33% (21-48%)	67% (52-79%)	0.77 (0.34-1.5)	1.1% (-0.32-3.1%)	1.3% (-0.43-3.6%)	140 (-45-380)	710 (-200-1900)
Low SDI	1990	Female	95 plus	36% (21-55%)	64% (45-79%)	0.8 (0.3-1.7)	1.2% (-0.36-3.3%)	1.3% (-0.52-3.9%)	36 (-14-110)	130 (-39-360)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	1990	Male	All Ages	29% (24-35%)	71% (65-76%)	0.89 (0.68-1.1)	1.8% (1.2-2.5%)	3% (1.9-4.1%)	73000 (46000-100000)	3100000 (2100000-4100000)
Low SDI	1990	Male	15 to 19	26% (17-37%)	74% (63-83%)	0.81 (0.44-1.4)	2.6% (1.4-4.2%)	3% (1.4-5.2%)	1400 (590-2400)	120000 (64000-200000)
Low SDI	1990	Male	20 to 24	36% (24-49%)	64% (51-76%)	1.4 (0.76-2.2)	5% (2.8-7.6%)	5.3% (2.6-8.6%)	3300 (1600-5500)	270000 (150000-410000)
Low SDI	1990	Male	25 to 29	42% (28-55%)	58% (45-72%)	1.8 (1.1-2.8)	6.2% (3.7-9.3%)	6.8% (3.5-11%)	4700 (2500-7300)	330000 (200000-490000)
Low SDI	1990	Male	30 to 34	45% (31-59%)	55% (41-69%)	2.1 (1.2-3.2)	7.1% (4.3-10%)	8% (4.4-12%)	5600 (3100-8400)	350000 (210000-500000)
Low SDI	1990	Male	35 to 39	46% (33-60%)	54% (40-67%)	2.2 (1.3-3.5)	7.8% (4.7-11%)	8.9% (5.1-13%)	6400 (3600-9500)	350000 (210000-510000)
Low SDI	1990	Male	40 to 44	47% (34-61%)	53% (39-66%)	2.2 (1.2-3.5)	8% (4.8-12%)	9.4% (5.4-14%)	7000 (4000-10000)	340000 (200000-490000)
Low SDI	1990	Male	45 to 49	48% (34-62%)	52% (38-66%)	2.2 (1.2-3.5)	8.2% (4.9-12%)	9.7% (5.6-14%)	7400 (4200-11000)	310000 (180000-470000)
Low SDI	1990	Male	50 to 54	49% (35-62%)	51% (38-65%)	2.1 (1.2-3.5)	8.3% (4.7-12%)	9.9% (5.5-15%)	7800 (4200-12000)	290000 (160000-440000)
Low SDI	1990	Male	55 to 59	49% (35-62%)	51% (38-65%)	2.1 (1.1-3.4)	8% (4.5-12%)	9.4% (5.2-14%)	8100 (4400-12000)	260000 (140000-400000)
Low SDI	1990	Male	60 to 64	46% (32-61%)	54% (39-68%)	1.8 (0.97-3.2)	6.6% (3.5-10%)	7.6% (4-12%)	7100 (3700-11000)	190000 (100000-310000)
Low SDI	1990	Male	65 to 69	44% (30-58%)	56% (42-70%)	1.5 (0.82-2.5)	5.1% (2.5-8.1%)	5.8% (2.8-9.4%)	5600 (2700-9200)	130000 (63000-210000)
Low SDI	1990	Male	70 to 74	42% (28-56%)	58% (44-72%)	1.2 (0.64-2.1)	3.8% (1.5-6.7%)	4.4% (1.7-7.8%)	4200 (1600-7500)	76000 (30000-130000)
Low SDI	1990	Male	75 to 79	40% (27-55%)	60% (45-73%)	1 (0.5-1.8)	3% (0.92-5.7%)	3.4% (1-6.5%)	2600 (770-5100)	37000 (11000-70000)
Low SDI	1990	Male	80 to 84	38% (24-53%)	62% (47-76%)	0.86 (0.43-1.5)	2.3% (0.67-4.5%)	2.6% (0.71-5.1%)	1100 (300-2200)	12000 (3400-23000)
Low SDI	1990	Male	85 to 89	35% (22-49%)	65% (51-78%)	0.71 (0.35-1.3)	1.7% (0.46-3.3%)	1.9% (0.44-3.8%)	390 (91-790)	3000 (800-5900)
Low SDI	1990	Male	90 to 94	35% (22-49%)	65% (51-78%)	0.64 (0.3-1.2)	1.1% (0.12-2.5%)	1.2% (0.0068-2.8%)	83 (0.39-200)	460 (46-1000)
Low SDI	1990	Male	95 plus	37% (21-55%)	63% (45-79%)	0.62 (0.23-1.3)	1.1% (-0.038-2.7%)	1.1% (-0.24-3%)	18 (-3.9-48)	70 (-2.7-170)
Low SDI	1995	Female	All Ages	24% (20-29%)	76% (71-80%)	0.6 (0.46-0.78)	1% (0.57-1.5%)	1.8% (0.86-2.8%)	45000 (22000-71000)	1700000 (950000-2500000)
Low SDI	1995	Female	15 to 19	24% (16-35%)	76% (65-84%)	0.68 (0.36-1.1)	1.5% (0.67-2.6%)	1.8% (0.61-3.5%)	790 (260-1500)	72000 (33000-120000)
Low SDI	1995	Female	20 to 24	31% (20-43%)	69% (57-80%)	1 (0.54-1.6)	2.1% (1-3.6%)	2.2% (0.75-4.3%)	1400 (480-2600)	120000 (58000-200000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	1995	Female	25 to 29	33% (21-46%)	67% (54-79%)	1.2 (0.68-1.9)	2.5% (1.2-4%)	2.7% (1-4.8%)	1900 (820-3300)	140000 (74000-230000)
Low SDI	1995	Female	30 to 34	35% (23-50%)	65% (50-77%)	1.3 (0.73-2.1)	2.7% (1.4-4.4%)	3.1% (1.3-5.4%)	2300 (1000-4000)	150000 (78000-240000)
Low SDI	1995	Female	35 to 39	38% (25-52%)	62% (48-75%)	1.3 (0.71-2.1)	3% (1.5-4.9%)	3.6% (1.5-6.2%)	2700 (1200-4600)	150000 (75000-250000)
Low SDI	1995	Female	40 to 44	39% (27-53%)	61% (47-73%)	1.3 (0.68-2.1)	3.2% (1.6-5.3%)	4% (1.8-6.7%)	2900 (1400-4900)	140000 (71000-240000)
Low SDI	1995	Female	45 to 49	39% (25-53%)	61% (47-75%)	1.3 (0.66-2.2)	3.6% (1.6-6%)	4.5% (1.9-7.7%)	3500 (1500-6000)	150000 (67000-250000)
Low SDI	1995	Female	50 to 54	37% (25-52%)	63% (48-75%)	1.3 (0.65-2.2)	4.2% (1.8-7%)	5.1% (2.2-8.7%)	4500 (1800-7700)	160000 (69000-280000)
Low SDI	1995	Female	55 to 59	36% (23-51%)	64% (49-77%)	1.3 (0.64-2.2)	4.3% (1.7-7.3%)	5.2% (2.1-8.9%)	4900 (1900-8600)	160000 (62000-270000)
Low SDI	1995	Female	60 to 64	37% (23-52%)	63% (48-77%)	1.3 (0.62-2.3)	4% (1.5-7%)	4.7% (1.8-8.4%)	5200 (2000-9400)	140000 (54000-250000)
Low SDI	1995	Female	65 to 69	37% (25-51%)	63% (49-75%)	1.2 (0.61-2)	3.8% (1.3-6.7%)	4.4% (1.5-7.8%)	5400 (1900-9800)	120000 (43000-220000)
Low SDI	1995	Female	70 to 74	36% (24-50%)	64% (50-76%)	1 (0.48-1.8)	3% (0.62-5.8%)	3.4% (0.7-6.7%)	4400 (850-8800)	79000 (16000-160000)
Low SDI	1995	Female	75 to 79	33% (21-48%)	67% (52-79%)	0.83 (0.37-1.5)	2.2% (0.2-4.8%)	2.6% (0.21-5.5%)	2800 (250-6100)	39000 (3700-84000)
Low SDI	1995	Female	80 to 84	31% (19-46%)	69% (54-81%)	0.75 (0.35-1.4)	1.7% (0.0058-3.8%)	1.9% (-0.0058-4.4%)	1400 (-2.8-3100)	14000 (110-32000)
Low SDI	1995	Female	85 to 89	31% (19-45%)	69% (55-81%)	0.73 (0.34-1.3)	1.3% (-0.17-3.3%)	1.5% (-0.23-3.8%)	560 (-80-1400)	4200 (-520-10000)
Low SDI	1995	Female	90 to 94	33% (21-48%)	67% (52-79%)	0.74 (0.33-1.4)	1% (-0.37-2.9%)	1.2% (-0.49-3.5%)	150 (-62-450)	790 (-280-2300)
Low SDI	1995	Female	95 plus	36% (21-55%)	64% (45-79%)	0.75 (0.29-1.6)	1.1% (-0.4-3.1%)	1.2% (-0.57-3.6%)	41 (-19-130)	150 (-55-430)
Low SDI	1995	Male	All Ages	29% (24-35%)	71% (65-76%)	0.87 (0.67-1.1)	1.9% (1.3-2.5%)	2.9% (1.9-4.1%)	81000 (52000-110000)	350000 (230000-470000)
Low SDI	1995	Male	15 to 19	26% (18-38%)	74% (62-82%)	0.83 (0.45-1.4)	2.8% (1.5-4.4%)	3.2% (1.5-5.5%)	1700 (770-2900)	150000 (80000-240000)
Low SDI	1995	Male	20 to 24	37% (25-49%)	63% (51-75%)	1.3 (0.77-2.1)	5.1% (2.9-7.9%)	5.4% (2.5-8.9%)	3900 (1900-6400)	320000 (180000-480000)
Low SDI	1995	Male	25 to 29	42% (29-55%)	58% (45-71%)	1.8 (1.1-2.7)	5.9% (3.4-8.9%)	6.2% (3.2-10%)	5400 (2900-8300)	380000 (230000-560000)
Low SDI	1995	Male	30 to 34	45% (31-59%)	55% (41-69%)	2 (1.2-3.1)	6.1% (3.6-9.1%)	6.7% (3.6-10%)	6300 (3600-9500)	390000 (240000-570000)
Low SDI	1995	Male	35 to 39	46% (33-60%)	54% (40-67%)	2.1 (1.2-3.3)	6.6% (3.9-9.8%)	7.4% (4.1-11%)	7100 (4100-11000)	390000 (240000-570000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	1995	Male	40 to 44	47% (34-60%)	53% (40-66%)	2.1 (1.2-3.4)	6.9% (4-10%)	7.9% (4.4-12%)	7800 (4500-12000)	380000 (230000-550000)
Low SDI	1995	Male	45 to 49	48% (34-61%)	52% (39-66%)	2.1 (1.2-3.4)	7.1% (4.1-11%)	8.3% (4.6-12%)	8200 (4600-12000)	350000 (200000-520000)
Low SDI	1995	Male	50 to 54	48% (35-62%)	52% (38-65%)	2 (1.1-3.3)	7.5% (4.1-11%)	8.7% (4.7-13%)	8700 (4600-13000)	320000 (180000-490000)
Low SDI	1995	Male	55 to 59	48% (34-62%)	52% (38-66%)	2 (1.1-3.2)	7.3% (4.1-11%)	8.6% (4.7-13%)	8600 (4600-13000)	280000 (150000-420000)
Low SDI	1995	Male	60 to 64	46% (32-61%)	54% (39-68%)	1.8 (0.94-3)	6.1% (3.3-9.5%)	7.1% (3.7-11%)	7900 (4100-13000)	220000 (110000-340000)
Low SDI	1995	Male	65 to 69	44% (30-58%)	56% (42-70%)	1.5 (0.79-2.4)	4.7% (2.3-7.6%)	5.4% (2.6-8.8%)	6100 (2800-10000)	140000 (66000-220000)
Low SDI	1995	Male	70 to 74	41% (28-55%)	59% (45-72%)	1.2 (0.61-2)	3.6% (1.4-6.3%)	4.1% (1.5-7.3%)	4600 (1800-8200)	83000 (33000-150000)
Low SDI	1995	Male	75 to 79	39% (26-54%)	61% (46-74%)	0.96 (0.47-1.7)	2.7% (0.83-5.2%)	3.1% (0.9-6%)	3000 (880-5700)	42000 (13000-80000)
Low SDI	1995	Male	80 to 84	37% (23-51%)	63% (49-77%)	0.8 (0.4-1.4)	2.1% (0.58-4.1%)	2.4% (0.6-4.6%)	1300 (340-2600)	14000 (3900-27000)
Low SDI	1995	Male	85 to 89	34% (22-48%)	66% (52-78%)	0.67 (0.33-1.2)	1.6% (0.43-3.2%)	1.8% (0.41-3.6%)	450 (110-930)	3500 (950-7000)
Low SDI	1995	Male	90 to 94	34% (22-49%)	66% (51-78%)	0.6 (0.28-1.1)	1% (0.081-2.3%)	1.1% (-0.032-2.6%)	94 (-2.3-230)	530 (43-1200)
Low SDI	1995	Male	95 plus	37% (20-55%)	63% (45-80%)	0.57 (0.22-1.2)	1% (-0.066-2.6%)	1% (-0.27-2.8%)	22 (-5.6-59)	86 (-5.3-210)
Low SDI	2000	Female	All Ages	23% (19-28%)	77% (72-81%)	0.59 (0.45-0.76)	1.1% (0.63-1.6%)	1.9% (0.96-3%)	52000 (27000-82000)	1900000 (1100000-2800000)
Low SDI	2000	Female	15 to 19	24% (16-34%)	76% (66-84%)	0.68 (0.37-1.1)	1.5% (0.74-2.6%)	1.9% (0.73-3.6%)	910 (340-1700)	83000 (41000-140000)
Low SDI	2000	Female	20 to 24	29% (20-41%)	71% (59-80%)	0.98 (0.54-1.6)	2.2% (1.1-3.6%)	2.4% (0.88-4.4%)	1500 (590-2800)	140000 (70000-230000)
Low SDI	2000	Female	25 to 29	31% (21-43%)	69% (57-79%)	1.2 (0.67-1.8)	2.3% (1.2-3.8%)	2.4% (1-4.4%)	2100 (960-3700)	160000 (86000-260000)
Low SDI	2000	Female	30 to 34	33% (22-46%)	67% (54-78%)	1.2 (0.71-2)	2.4% (1.3-3.9%)	2.6% (1.2-4.6%)	2500 (1200-4200)	160000 (88000-260000)
Low SDI	2000	Female	35 to 39	36% (25-49%)	64% (51-75%)	1.2 (0.7-2)	2.6% (1.3-4.3%)	3% (1.4-5.3%)	2900 (1400-4900)	160000 (84000-260000)
Low SDI	2000	Female	40 to 44	37% (26-50%)	63% (50-74%)	1.2 (0.67-2)	2.9% (1.5-4.7%)	3.5% (1.6-5.9%)	3200 (1600-5400)	160000 (81000-260000)
Low SDI	2000	Female	45 to 49	37% (25-50%)	63% (50-75%)	1.3 (0.68-2.2)	3.5% (1.7-5.8%)	4.3% (1.9-7.3%)	4000 (1800-6900)	170000 (82000-290000)
Low SDI	2000	Female	50 to 54	36% (25-50%)	64% (50-75%)	1.2 (0.66-2.1)	4.2% (1.9-6.9%)	5.1% (2.3-8.6%)	5200 (2300-8900)	190000 (86000-320000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	2000	Female	55 to 59	35% (24-49%)	65% (51-76%)	1.3 (0.66-2.2)	4.5% (2-7.6%)	5.5% (2.4-9.3%)	6000 (2500-10000)	190000 (82000-320000)
Low SDI	2000	Female	60 to 64	36% (23-50%)	64% (50-77%)	1.2 (0.63-2.2)	4% (1.7-7%)	4.8% (2-8.5%)	5800 (2300-10000)	150000 (62000-280000)
Low SDI	2000	Female	65 to 69	37% (25-50%)	63% (50-75%)	1.2 (0.63-2)	4% (1.5-7%)	4.7% (1.8-8.2%)	6500 (2500-12000)	140000 (56000-260000)
Low SDI	2000	Female	70 to 74	35% (23-48%)	65% (52-77%)	1 (0.49-1.8)	3.1% (0.78-6.1%)	3.6% (0.87-7%)	5000 (1200-10000)	91000 (23000-180000)
Low SDI	2000	Female	75 to 79	32% (20-47%)	68% (53-80%)	0.85 (0.39-1.6)	2.5% (0.39-5.2%)	2.9% (0.43-6%)	3700 (580-7700)	51000 (8300-110000)
Low SDI	2000	Female	80 to 84	31% (19-46%)	69% (54-81%)	0.77 (0.36-1.4)	1.9% (0.15-4.1%)	2.2% (0.16-4.8%)	1900 (150-4100)	19000 (1500-42000)
Low SDI	2000	Female	85 to 89	31% (19-45%)	69% (55-81%)	0.72 (0.34-1.3)	1.4% (-0.11-3.4%)	1.6% (-0.16-3.9%)	690 (-57-1700)	5100 (-360-12000)
Low SDI	2000	Female	90 to 94	33% (21-47%)	67% (53-79%)	0.74 (0.34-1.4)	1.1% (-0.31-3.1%)	1.3% (-0.41-3.6%)	190 (-59-540)	1000 (-260-2700)
Low SDI	2000	Female	95 plus	36% (21-54%)	64% (46-79%)	0.77 (0.3-1.6)	1.2% (-0.34-3.4%)	1.3% (-0.51-3.9%)	53 (-20-160)	190 (-53-540)
Low SDI	2000	Male	All Ages	29% (24-34%)	71% (66-76%)	0.89 (0.69-1.1)	2% (1.4-2.7%)	3.1% (2.1-4.3%)	94000 (62000-130000)	400000 (280000-530000)
Low SDI	2000	Male	15 to 19	26% (18-36%)	74% (64-82%)	0.85 (0.47-1.4)	2.9% (1.7-4.5%)	3.4% (1.7-5.7%)	2000 (970-3400)	180000 (99000-280000)
Low SDI	2000	Male	20 to 24	36% (26-48%)	64% (52-74%)	1.4 (0.8-2.1)	5.5% (3.2-8.1%)	5.9% (3.1-9.3%)	4700 (2500-7600)	380000 (220000-570000)
Low SDI	2000	Male	25 to 29	42% (30-54%)	58% (46-70%)	1.8 (1.1-2.7)	6% (3.6-9%)	6.4% (3.4-10%)	6400 (3600-9700)	450000 (280000-650000)
Low SDI	2000	Male	30 to 34	45% (32-57%)	55% (43-68%)	2.1 (1.3-3.1)	6% (3.7-8.8%)	6.5% (3.7-9.9%)	7300 (4300-11000)	450000 (290000-650000)
Low SDI	2000	Male	35 to 39	46% (34-59%)	54% (41-66%)	2.2 (1.3-3.4)	6.2% (3.7-9%)	6.9% (3.9-10%)	8100 (4800-12000)	440000 (280000-640000)
Low SDI	2000	Male	40 to 44	47% (35-60%)	53% (40-65%)	2.2 (1.3-3.4)	6.6% (4-9.5%)	7.5% (4.3-11%)	8800 (5200-13000)	420000 (260000-610000)
Low SDI	2000	Male	45 to 49	48% (35-60%)	52% (40-65%)	2.2 (1.2-3.4)	7% (4.2-10%)	8.1% (4.7-12%)	9400 (5500-14000)	400000 (240000-590000)
Low SDI	2000	Male	50 to 54	49% (36-61%)	51% (39-64%)	2.1 (1.2-3.4)	7.6% (4.4-11%)	8.8% (5-13%)	9800 (5500-15000)	370000 (210000-550000)
Low SDI	2000	Male	55 to 59	48% (36-61%)	52% (39-64%)	2.1 (1.2-3.3)	7.7% (4.5-11%)	9% (5.1-13%)	10000 (5600-15000)	320000 (180000-480000)
Low SDI	2000	Male	60 to 64	46% (33-60%)	54% (40-67%)	1.8 (1-3.1)	6.3% (3.4-9.8%)	7.4% (3.9-11%)	8500 (4400-13000)	230000 (120000-360000)
Low SDI	2000	Male	65 to 69	44% (31-57%)	56% (43-69%)	1.5 (0.85-2.4)	5% (2.5-8.1%)	5.8% (2.9-9.4%)	7200 (3500-12000)	160000 (81000-260000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	2000	Male	70 to 74	42% (29-55%)	58% (45-71%)	1.2 (0.67-2)	3.8% (1.5-6.7%)	4.4% (1.7-7.7%)	5200 (2100-9200)	94000 (38000-170000)
Low SDI	2000	Male	75 to 79	40% (27-55%)	60% (45-73%)	1 (0.53-1.8)	3.1% (1-5.8%)	3.6% (1.1-6.7%)	3600 (1200-6900)	51000 (17000-96000)
Low SDI	2000	Male	80 to 84	37% (23-52%)	63% (48-77%)	0.85 (0.44-1.4)	2.3% (0.74-4.5%)	2.7% (0.79-5.1%)	1700 (520-3300)	18000 (5800-34000)
Low SDI	2000	Male	85 to 89	34% (22-48%)	66% (52-78%)	0.69 (0.35-1.2)	1.7% (0.49-3.3%)	1.9% (0.47-3.8%)	600 (150-1200)	4600 (1300-8900)
Low SDI	2000	Male	90 to 94	35% (23-49%)	65% (51-77%)	0.63 (0.3-1.2)	1.2% (0.17-2.5%)	1.3% (0.07-2.9%)	130 (7.6-300)	710 (100-1500)
Low SDI	2000	Male	95 plus	38% (21-56%)	62% (44-79%)	0.6 (0.23-1.2)	1.2% (0.027-2.9%)	1.3% (-0.16-3.2%)	31 (-3.9-79)	120 (2.8-280)
Low SDI	2005	Female	All Ages	23% (19-28%)	77% (72-81%)	0.6 (0.45-0.78)	1.1% (0.65-1.6%)	1.9% (0.96-3%)	53000 (27000-85000)	1900000 (1100000-2900000)
Low SDI	2005	Female	15 to 19	24% (16-34%)	76% (66-84%)	0.69 (0.38-1.1)	1.4% (0.71-2.4%)	1.7% (0.65-3.3%)	900 (340-1700)	86000 (42000-150000)
Low SDI	2005	Female	20 to 24	30% (20-41%)	70% (59-80%)	1 (0.55-1.6)	2.3% (1.2-3.7%)	2.4% (0.93-4.5%)	1600 (610-2900)	150000 (76000-240000)
Low SDI	2005	Female	25 to 29	31% (21-43%)	69% (57-79%)	1.2 (0.68-1.9)	2.3% (1.3-3.8%)	2.5% (1.1-4.6%)	2100 (980-3700)	170000 (93000-260000)
Low SDI	2005	Female	30 to 34	33% (23-46%)	67% (54-77%)	1.3 (0.73-2)	2.3% (1.2-3.7%)	2.5% (1.2-4.4%)	2500 (1200-4200)	170000 (93000-260000)
Low SDI	2005	Female	35 to 39	36% (25-49%)	64% (51-75%)	1.3 (0.71-2)	2.4% (1.3-3.9%)	2.8% (1.3-4.8%)	2900 (1400-4800)	160000 (87000-270000)
Low SDI	2005	Female	40 to 44	37% (26-50%)	63% (50-74%)	1.3 (0.69-2.1)	2.7% (1.4-4.3%)	3.2% (1.5-5.4%)	3200 (1500-5400)	160000 (82000-260000)
Low SDI	2005	Female	45 to 49	37% (25-50%)	63% (50-75%)	1.3 (0.68-2.2)	3.3% (1.6-5.5%)	4.1% (1.9-6.9%)	4100 (1900-7100)	180000 (85000-300000)
Low SDI	2005	Female	50 to 54	36% (24-50%)	64% (50-76%)	1.3 (0.66-2.2)	4.1% (1.9-6.7%)	5% (2.3-8.4%)	5300 (2300-9200)	200000 (88000-340000)
Low SDI	2005	Female	55 to 59	35% (23-49%)	65% (51-77%)	1.3 (0.65-2.2)	4.3% (1.9-7.3%)	5.3% (2.3-9.1%)	6100 (2500-11000)	190000 (81000-340000)
Low SDI	2005	Female	60 to 64	35% (23-50%)	65% (50-77%)	1.3 (0.63-2.3)	4% (1.6-7%)	4.8% (2-8.4%)	6100 (2400-11000)	160000 (64000-290000)
Low SDI	2005	Female	65 to 69	36% (24-50%)	64% (50-76%)	1.2 (0.61-2)	3.8% (1.4-6.7%)	4.5% (1.7-8%)	6300 (2200-11000)	140000 (50000-250000)
Low SDI	2005	Female	70 to 74	35% (23-48%)	65% (52-77%)	1 (0.5-1.8)	3.1% (0.77-6.1%)	3.7% (0.87-7.2%)	5400 (1200-11000)	98000 (23000-190000)
Low SDI	2005	Female	75 to 79	32% (20-47%)	68% (53-80%)	0.82 (0.38-1.5)	2.4% (0.31-5%)	2.7% (0.35-5.8%)	3500 (420-7500)	49000 (6000-100000)
Low SDI	2005	Female	80 to 84	31% (19-46%)	69% (54-81%)	0.78 (0.37-1.4)	1.9% (0.16-4.2%)	2.2% (0.19-4.8%)	2100 (160-4600)	22000 (1700-47000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	2005	Female	85 to 89	31% (19-45%)	69% (55-81%)	0.75 (0.35-1.4)	1.5% (-0.06-3.5%)	1.7% (-0.094-4.1%)	860 (-48-2100)	6400 (-270-15000)
Low SDI	2005	Female	90 to 94	32% (20-47%)	68% (53-80%)	0.74 (0.33-1.4)	1.2% (-0.28-3.1%)	1.3% (-0.39-3.7%)	230 (-66-630)	1200 (-280-3200)
Low SDI	2005	Female	95 plus	36% (21-54%)	64% (46-79%)	0.78 (0.3-1.6)	1.3% (-0.31-3.5%)	1.4% (-0.46-4.1%)	64 (-20-180)	220 (-54-620)
Low SDI	2005	Male	All Ages	29% (24-34%)	71% (66-76%)	0.9 (0.69-1.1)	2.2% (1.6-3%)	3.4% (2.2-4.7%)	99000 (65000-140000)	430000 (300000-570000)
Low SDI	2005	Male	15 to 19	27% (19-37%)	73% (63-81%)	0.88 (0.5-1.4)	3% (1.8-4.7%)	3.6% (1.8-5.9%)	2200 (1100-3600)	190000 (110000-300000)
Low SDI	2005	Male	20 to 24	37% (27-48%)	63% (52-73%)	1.4 (0.82-2.2)	6.1% (3.7-8.9%)	6.7% (3.6-11%)	5200 (2700-8300)	420000 (250000-630000)
Low SDI	2005	Male	25 to 29	41% (30-53%)	59% (47-70%)	1.8 (1.1-2.7)	6.8% (4.2-9.9%)	7.5% (4.1-12%)	7000 (3900-11000)	500000 (310000-720000)
Low SDI	2005	Male	30 to 34	44% (32-56%)	56% (44-68%)	2.1 (1.3-3.1)	6.5% (4.1-9.3%)	7.2% (4.1-11%)	7900 (4700-12000)	500000 (320000-710000)
Low SDI	2005	Male	35 to 39	45% (34-58%)	55% (42-66%)	2.2 (1.3-3.4)	6.3% (4-9.2%)	7.1% (4.2-11%)	8400 (5000-12000)	460000 (290000-660000)
Low SDI	2005	Male	40 to 44	46% (35-59%)	54% (41-65%)	2.2 (1.3-3.4)	6.5% (4-9.3%)	7.4% (4.4-11%)	9100 (5400-13000)	440000 (270000-640000)
Low SDI	2005	Male	45 to 49	48% (35-60%)	52% (40-65%)	2.2 (1.2-3.5)	7% (4.3-10%)	8.1% (4.8-12%)	9700 (5700-14000)	420000 (250000-610000)
Low SDI	2005	Male	50 to 54	48% (36-61%)	52% (39-64%)	2.1 (1.2-3.4)	7.5% (4.4-11%)	8.9% (5.1-13%)	10000 (5800-16000)	390000 (220000-570000)
Low SDI	2005	Male	55 to 59	48% (36-61%)	52% (39-64%)	2.1 (1.2-3.4)	7.7% (4.5-11%)	9.2% (5.2-14%)	10000 (5800-16000)	340000 (190000-510000)
Low SDI	2005	Male	60 to 64	46% (33-60%)	54% (40-67%)	1.8 (0.99-3.1)	6.4% (3.5-9.9%)	7.5% (4-12%)	9300 (4800-15000)	250000 (140000-400000)
Low SDI	2005	Male	65 to 69	43% (30-57%)	57% (43-70%)	1.5 (0.84-2.5)	5% (2.5-8.1%)	5.8% (2.8-9.5%)	7300 (3500-12000)	170000 (82000-270000)
Low SDI	2005	Male	70 to 74	42% (29-55%)	58% (45-71%)	1.3 (0.68-2.1)	3.9% (1.6-6.9%)	4.6% (1.8-8%)	5900 (2400-10000)	110000 (44000-190000)
Low SDI	2005	Male	75 to 79	40% (27-54%)	60% (46-73%)	1.1 (0.52-1.9)	3.1% (1-5.8%)	3.6% (1.1-6.8%)	3800 (1200-7200)	54000 (18000-100000)
Low SDI	2005	Male	80 to 84	38% (24-53%)	62% (47-76%)	0.88 (0.45-1.5)	2.4% (0.79-4.6%)	2.8% (0.84-5.3%)	1900 (590-3700)	20000 (6600-39000)
Low SDI	2005	Male	85 to 89	34% (22-48%)	66% (52-78%)	0.69 (0.35-1.2)	1.7% (0.52-3.3%)	1.9% (0.51-3.8%)	730 (190-1400)	5600 (1700-11000)
Low SDI	2005	Male	90 to 94	34% (22-49%)	66% (51-78%)	0.6 (0.29-1.1)	1.1% (0.16-2.5%)	1.2% (0.059-2.8%)	160 (7.7-360)	860 (120-1900)
Low SDI	2005	Male	95 plus	38% (21-56%)	62% (44-79%)	0.59 (0.23-1.2)	1.3% (0.061-3%)	1.3% (-0.12-3.4%)	38 (-3.6-97)	150 (6.6-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	2010	Female	All Ages	23% (19-28%)	77% (72-81%)	0.61 (0.44-0.84)	1.2% (0.65-1.8%)	2% (0.94-3.3%)	55000 (25000-90000)	200000 (1100000-3000000)
Low SDI	2010	Female	15 to 19	24% (16-34%)	76% (66-84%)	0.67 (0.34-1.2)	1.3% (0.62-2.2%)	1.5% (0.51-2.9%)	850 (300-1600)	86000 (42000-150000)
Low SDI	2010	Female	20 to 24	30% (20-42%)	70% (58-80%)	1 (0.53-1.7)	2.2% (1.2-3.6%)	2.3% (0.87-4.4%)	1500 (580-2800)	150000 (79000-240000)
Low SDI	2010	Female	25 to 29	32% (22-44%)	68% (56-78%)	1.2 (0.67-2)	2.4% (1.3-3.9%)	2.7% (1.1-4.8%)	2100 (960-3700)	170000 (96000-270000)
Low SDI	2010	Female	30 to 34	34% (23-47%)	66% (53-77%)	1.3 (0.73-2.2)	2.4% (1.3-3.9%)	2.8% (1.3-4.8%)	2600 (1200-4400)	180000 (97000-280000)
Low SDI	2010	Female	35 to 39	37% (25-50%)	63% (50-75%)	1.3 (0.7-2.2)	2.5% (1.3-4.1%)	3% (1.4-5.2%)	3000 (1400-5000)	170000 (90000-280000)
Low SDI	2010	Female	40 to 44	38% (27-51%)	62% (49-73%)	1.3 (0.67-2.2)	2.8% (1.4-4.5%)	3.4% (1.6-5.7%)	3300 (1500-5600)	170000 (83000-270000)
Low SDI	2010	Female	45 to 49	38% (25-51%)	62% (49-75%)	1.3 (0.66-2.3)	3.4% (1.6-5.6%)	4.3% (1.9-7.3%)	4200 (1800-7300)	180000 (82000-310000)
Low SDI	2010	Female	50 to 54	37% (25-52%)	63% (48-75%)	1.3 (0.64-2.3)	4.1% (1.8-6.8%)	5.2% (2.3-8.7%)	5400 (2300-9500)	200000 (87000-350000)
Low SDI	2010	Female	55 to 59	36% (23-51%)	64% (49-77%)	1.3 (0.64-2.3)	4.4% (1.8-7.5%)	5.5% (2.2-9.4%)	6300 (2400-11000)	200000 (78000-360000)
Low SDI	2010	Female	60 to 64	36% (23-51%)	64% (49-77%)	1.3 (0.6-2.3)	3.9% (1.4-6.9%)	4.7% (1.8-8.5%)	6200 (2100-11000)	170000 (58000-310000)
Low SDI	2010	Female	65 to 69	37% (25-51%)	63% (49-75%)	1.2 (0.59-2.1)	3.7% (1.3-6.7%)	4.5% (1.5-8.1%)	6700 (2100-12000)	150000 (48000-280000)
Low SDI	2010	Female	70 to 74	35% (24-49%)	65% (51-76%)	1 (0.48-1.8)	2.9% (0.56-6%)	3.4% (0.64-7.1%)	5300 (910-11000)	96000 (17000-200000)
Low SDI	2010	Female	75 to 79	33% (21-48%)	67% (52-79%)	0.86 (0.38-1.6)	2.4% (0.23-5.1%)	2.8% (0.26-6%)	4000 (310-8800)	56000 (4600-120000)
Low SDI	2010	Female	80 to 84	32% (19-47%)	68% (53-81%)	0.77 (0.35-1.4)	1.8% (0.04-4.1%)	2.1% (0.043-4.8%)	2200 (3.3-5200)	23000 (150-53000)
Low SDI	2010	Female	85 to 89	31% (19-45%)	69% (55-81%)	0.74 (0.34-1.4)	1.4% (-0.13-3.5%)	1.7% (-0.16-4.1%)	970 (-100-2400)	7200 (-700-18000)
Low SDI	2010	Female	90 to 94	33% (20-47%)	67% (53-80%)	0.75 (0.33-1.4)	1.1% (-0.34-3.2%)	1.3% (-0.47-3.8%)	280 (-99-790)	1400 (-450-4000)
Low SDI	2010	Female	95 plus	36% (20-53%)	64% (47-80%)	0.76 (0.29-1.6)	1.2% (-0.37-3.5%)	1.3% (-0.54-4%)	71 (-29-220)	250 (-79-740)
Low SDI	2010	Male	All Ages	28% (23-34%)	72% (66-77%)	0.89 (0.65-1.2)	2.4% (1.7-3.4%)	3.6% (2.3-5.1%)	110000 (66000-150000)	460000 (310000-630000)
Low SDI	2010	Male	15 to 19	26% (18-37%)	74% (63-82%)	0.82 (0.44-1.4)	2.7% (1.6-4.3%)	3% (1.4-5.2%)	2100 (990-3700)	200000 (110000-320000)
Low SDI	2010	Male	20 to 24	36% (26-48%)	64% (52-74%)	1.4 (0.76-2.2)	6% (3.6-8.8%)	6.4% (3.3-10%)	5400 (2700-8900)	460000 (270000-690000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	2010	Male	25 to 29	41% (29-53%)	59% (47-71%)	1.8 (1.1-2.8)	7.1% (4.4-10%)	7.9% (4.3-12%)	7600 (4100-12000)	550000 (340000-810000)
Low SDI	2010	Male	30 to 34	43% (31-57%)	57% (43-69%)	2 (1.2-3.2)	7% (4.4-10%)	8% (4.6-12%)	8600 (4900-13000)	550000 (340000-790000)
Low SDI	2010	Male	35 to 39	45% (33-58%)	55% (42-67%)	2.1 (1.2-3.5)	6.9% (4.3-9.9%)	7.9% (4.6-12%)	9100 (5300-14000)	510000 (320000-750000)
Low SDI	2010	Male	40 to 44	46% (34-59%)	54% (41-66%)	2.2 (1.2-3.6)	7% (4.3-10%)	8.2% (4.9-12%)	9700 (5600-15000)	480000 (290000-700000)
Low SDI	2010	Male	45 to 49	47% (34-60%)	53% (40-66%)	2.2 (1.2-3.6)	7.4% (4.5-11%)	8.7% (5.1-13%)	10000 (5900-16000)	440000 (260000-660000)
Low SDI	2010	Male	50 to 54	48% (35-61%)	52% (39-65%)	2.1 (1.1-3.6)	7.8% (4.5-12%)	9.3% (5.2-14%)	11000 (6000-17000)	410000 (230000-620000)
Low SDI	2010	Male	55 to 59	47% (35-61%)	53% (39-65%)	2 (1.1-3.4)	7.7% (4.3-11%)	9.2% (5.1-14%)	11000 (6000-17000)	360000 (200000-550000)
Low SDI	2010	Male	60 to 64	46% (32-61%)	54% (39-68%)	1.8 (0.94-3.2)	6.3% (3.3-9.9%)	7.5% (3.8-12%)	9800 (4900-16000)	270000 (140000-430000)
Low SDI	2010	Male	65 to 69	44% (30-58%)	56% (42-70%)	1.5 (0.79-2.5)	4.9% (2.3-8.1%)	5.8% (2.7-9.5%)	8000 (3600-14000)	180000 (85000-310000)
Low SDI	2010	Male	70 to 74	42% (28-55%)	58% (45-72%)	1.2 (0.63-2.2)	3.7% (1.4-6.7%)	4.3% (1.6-7.8%)	6000 (2200-11000)	110000 (41000-200000)
Low SDI	2010	Male	75 to 79	40% (26-55%)	60% (45-74%)	1 (0.5-1.9)	3% (0.9-5.8%)	3.4% (1-6.7%)	4200 (1200-8300)	59000 (18000-120000)
Low SDI	2010	Male	80 to 84	38% (24-53%)	62% (47-76%)	0.88 (0.42-1.6)	2.3% (0.69-4.5%)	2.7% (0.74-5.3%)	2200 (600-4400)	23000 (6800-46000)
Low SDI	2010	Male	85 to 89	34% (22-49%)	66% (51-78%)	0.7 (0.34-1.3)	1.7% (0.47-3.4%)	1.9% (0.45-3.9%)	820 (200-1700)	6300 (1700-13000)
Low SDI	2010	Male	90 to 94	34% (22-49%)	66% (51-78%)	0.6 (0.28-1.1)	1.1% (0.12-2.4%)	1.1% (0.0095-2.7%)	180 (1.8-440)	1000 (120-2300)
Low SDI	2010	Male	95 plus	37% (21-55%)	63% (45-79%)	0.56 (0.22-1.1)	1.2% (0.03-2.8%)	1.2% (-0.16-3.1%)	45 (-5.8-120)	170 (4.5-420)
Low SDI	2016	Female	All Ages	24% (20-29%)	76% (71-80%)	0.63 (0.36-1)	1.4% (0.68-2.3%)	2.4% (0.88-4.2%)	60000 (22000-110000)	2200000 (1100000-3600000)
Low SDI	2016	Female	15 to 19	24% (17-35%)	76% (65-83%)	0.67 (0.3-1.3)	1.3% (0.61-2.3%)	1.6% (0.5-3.3%)	860 (270-1800)	92000 (42000-160000)
Low SDI	2016	Female	20 to 24	30% (20-42%)	70% (58-80%)	1 (0.45-1.9)	2.3% (1.2-3.8%)	2.5% (0.87-4.8%)	1500 (520-3000)	160000 (82000-270000)
Low SDI	2016	Female	25 to 29	32% (21-44%)	68% (56-79%)	1.2 (0.57-2.3)	2.7% (1.5-4.3%)	3.1% (1.3-5.7%)	2100 (870-3900)	190000 (99000-300000)
Low SDI	2016	Female	30 to 34	34% (23-47%)	66% (53-77%)	1.3 (0.63-2.5)	2.9% (1.5-4.6%)	3.5% (1.5-6.2%)	2700 (1200-4900)	200000 (100000-320000)
Low SDI	2016	Female	35 to 39	37% (25-49%)	63% (51-75%)	1.3 (0.61-2.5)	3.1% (1.6-5%)	3.9% (1.8-6.8%)	3300 (1400-5800)	200000 (96000-330000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	2016	Female	40 to 44	38% (27-51%)	62% (49-73%)	1.3 (0.6-2.5)	3.3% (1.6-5.4%)	4.4% (2-7.4%)	3700 (1600-6500)	190000 (88000-320000)
Low SDI	2016	Female	45 to 49	38% (26-51%)	62% (49-74%)	1.3 (0.57-2.6)	3.8% (1.7-6.7%)	5.1% (2.1-9%)	4600 (1800-8300)	200000 (82000-360000)
Low SDI	2016	Female	50 to 54	37% (25-52%)	63% (48-75%)	1.3 (0.55-2.6)	4.3% (1.7-7.6%)	5.7% (2.2-10%)	5800 (2100-11000)	220000 (81000-390000)
Low SDI	2016	Female	55 to 59	36% (23-50%)	64% (50-77%)	1.3 (0.54-2.6)	4.4% (1.5-8.1%)	5.7% (2-10%)	6700 (2100-13000)	210000 (69000-400000)
Low SDI	2016	Female	60 to 64	36% (23-51%)	64% (49-77%)	1.3 (0.53-2.6)	3.9% (1.2-7.5%)	4.9% (1.5-9.3%)	6700 (1900-13000)	180000 (50000-360000)
Low SDI	2016	Female	65 to 69	37% (24-51%)	63% (49-76%)	1.2 (0.51-2.3)	3.6% (0.91-7.1%)	4.4% (1.1-8.7%)	7100 (1700-14000)	160000 (37000-320000)
Low SDI	2016	Female	70 to 74	36% (24-49%)	64% (51-76%)	1 (0.42-2.1)	2.9% (0.29-6.5%)	3.4% (0.33-7.7%)	5900 (510-14000)	110000 (9900-240000)
Low SDI	2016	Female	75 to 79	33% (21-48%)	67% (52-79%)	0.85 (0.33-1.8)	2.3% (0.0096-5.5%)	2.7% (0.017-6.5%)	4300 (-37-11000)	60000 (-580-150000)
Low SDI	2016	Female	80 to 84	32% (19-47%)	68% (53-81%)	0.78 (0.31-1.6)	1.8% (-0.15-4.5%)	2.1% (-0.17-5.2%)	2600 (-270-6600)	27000 (-2800-68000)
Low SDI	2016	Female	85 to 89	32% (20-46%)	68% (54-80%)	0.76 (0.3-1.6)	1.5% (-0.27-4%)	1.7% (-0.33-4.7%)	1200 (-250-3400)	9100 (-1800-25000)
Low SDI	2016	Female	90 to 94	33% (21-48%)	67% (52-79%)	0.76 (0.29-1.6)	1.1% (-0.47-3.6%)	1.3% (-0.6-4.2%)	350 (-160-1100)	1800 (-760-5800)
Low SDI	2016	Female	95 plus	36% (21-54%)	64% (46-79%)	0.81 (0.27-1.8)	1.3% (-0.41-4%)	1.5% (-0.58-4.7%)	100 (-42-330)	370 (-120-1100)
Low SDI	2016	Male	All Ages	29% (24-34%)	71% (66-76%)	0.91 (0.53-1.5)	3% (1.8-4.3%)	4.2% (2.3-6.4%)	120000 (62000-180000)	520000 (310000-760000)
Low SDI	2016	Male	15 to 19	26% (18-37%)	74% (63-82%)	0.82 (0.38-1.5)	2.8% (1.5-4.6%)	3.1% (1.3-5.7%)	2400 (950-4400)	230000 (120000-380000)
Low SDI	2016	Male	20 to 24	36% (25-48%)	64% (52-75%)	1.3 (0.64-2.4)	6.1% (3.5-9.4%)	6.5% (3-11%)	5900 (2600-10000)	520000 (280000-820000)
Low SDI	2016	Male	25 to 29	41% (29-53%)	59% (47-71%)	1.8 (0.89-3.2)	7.5% (4.5-11%)	8.4% (4.3-14%)	8300 (4000-14000)	620000 (350000-950000)
Low SDI	2016	Male	30 to 34	44% (31-57%)	56% (43-69%)	2.1 (1-3.7)	7.8% (4.6-11%)	9% (4.8-14%)	9500 (4800-15000)	620000 (350000-930000)
Low SDI	2016	Male	35 to 39	45% (33-57%)	55% (43-67%)	2.2 (1-3.9)	7.7% (4.5-11%)	9.1% (5-14%)	10000 (5300-16000)	580000 (330000-880000)
Low SDI	2016	Male	40 to 44	46% (34-59%)	54% (41-66%)	2.2 (1-4)	7.9% (4.5-12%)	9.5% (5.2-14%)	11000 (5600-17000)	540000 (300000-830000)
Low SDI	2016	Male	45 to 49	47% (34-60%)	53% (40-66%)	2.2 (1-4.1)	8.2% (4.5-12%)	9.9% (5.3-15%)	11000 (5700-18000)	500000 (260000-770000)
Low SDI	2016	Male	50 to 54	48% (35-61%)	52% (39-65%)	2.2 (0.98-4.1)	8.2% (4.3-13%)	10% (5-16%)	12000 (5700-19000)	450000 (220000-720000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Low SDI	2016	Male	55 to 59	47% (35-61%)	53% (39-65%)	2.1 (0.95-3.9)	7.8% (3.9-12%)	9.5% (4.7-15%)	12000 (5600-19000)	390000 (190000-630000)
Low SDI	2016	Male	60 to 64	46% (32-60%)	54% (40-68%)	1.8 (0.81-3.5)	6.3% (2.9-10%)	7.5% (3.3-13%)	11000 (4600-18000)	290000 (130000-500000)
Low SDI	2016	Male	65 to 69	43% (30-57%)	57% (43-70%)	1.5 (0.68-2.8)	4.8% (1.9-8.6%)	5.7% (2.2-10%)	8600 (3200-16000)	200000 (76000-360000)
Low SDI	2016	Male	70 to 74	42% (28-55%)	58% (45-72%)	1.2 (0.54-2.4)	3.6% (1.1-7.1%)	4.2% (1.3-8.4%)	6600 (1900-13000)	120000 (36000-240000)
Low SDI	2016	Male	75 to 79	40% (26-55%)	60% (45-74%)	1 (0.43-2.1)	2.9% (0.72-6.1%)	3.3% (0.8-7.2%)	4500 (1000-9900)	64000 (16000-140000)
Low SDI	2016	Male	80 to 84	38% (24-53%)	62% (47-76%)	0.88 (0.38-1.7)	2.3% (0.52-4.9%)	2.6% (0.54-5.7%)	2500 (500-5500)	27000 (5900-58000)
Low SDI	2016	Male	85 to 89	35% (22-50%)	65% (50-78%)	0.73 (0.31-1.5)	1.7% (0.37-3.9%)	1.9% (0.34-4.5%)	1000 (180-2400)	8000 (1700-18000)
Low SDI	2016	Male	90 to 94	35% (22-49%)	65% (51-78%)	0.63 (0.25-1.3)	1.2% (0.088-2.9%)	1.2% (-0.025-3.3%)	240 (-6.1-650)	1300 (97-3400)
Low SDI	2016	Male	95 plus	38% (21-55%)	62% (45-79%)	0.58 (0.2-1.3)	1.2% (0.021-3.1%)	1.2% (-0.17-3.5%)	60 (-8.7-170)	230 (2.7-610)
China	1990	Female	All Ages	12% (9.9-14%)	88% (86-90%)	0.14 (0.11-0.17)	0.85% (0.61-1.1%)	1.3% (0.81-1.8%)	49000 (31000-68000)	1500000 (1100000-2000000)
China	1990	Female	15 to 19	8.7% (5.4-13%)	91% (87-95%)	0.086 (0.046-0.15)	0.63% (0.42-0.92%)	0.72% (0.38-1.2%)	410 (220-710)	51000 (34000-74000)
China	1990	Female	20 to 24	11% (6.4-17%)	89% (83-94%)	0.095 (0.049-0.16)	0.91% (0.59-1.3%)	0.67% (0.23-1.3%)	560 (200-1100)	97000 (61000-140000)
China	1990	Female	25 to 29	12% (7.3-19%)	88% (81-93%)	0.12 (0.061-0.2)	1% (0.68-1.4%)	0.79% (0.3-1.5%)	540 (210-1000)	87000 (58000-120000)
China	1990	Female	30 to 34	14% (8.3-23%)	86% (77-92%)	0.16 (0.081-0.26)	1% (0.7-1.4%)	1% (0.54-1.7%)	720 (370-1200)	82000 (55000-120000)
China	1990	Female	35 to 39	17% (9.7-25%)	83% (75-90%)	0.21 (0.11-0.34)	1.2% (0.84-1.6%)	1.6% (0.99-2.3%)	1400 (870-2100)	110000 (75000-150000)
China	1990	Female	40 to 44	18% (11-28%)	82% (72-89%)	0.26 (0.15-0.41)	1.4% (0.98-1.9%)	2.1% (1.4-3%)	1800 (1200-2600)	100000 (76000-140000)
China	1990	Female	45 to 49	18% (11-27%)	82% (73-89%)	0.29 (0.16-0.44)	1.6% (1.1-2.2%)	2.5% (1.6-3.5%)	2400 (1600-3500)	110000 (75000-150000)
China	1990	Female	50 to 54	18% (11-27%)	82% (73-89%)	0.3 (0.17-0.47)	1.7% (1.2-2.5%)	2.6% (1.7-3.7%)	3700 (2400-5100)	140000 (92000-190000)
China	1990	Female	55 to 59	18% (11-26%)	82% (74-89%)	0.31 (0.18-0.49)	1.9% (1.2-2.7%)	2.7% (1.7-3.8%)	5400 (3500-7700)	170000 (110000-240000)
China	1990	Female	60 to 64	18% (11-27%)	82% (73-89%)	0.32 (0.18-0.54)	1.7% (1.1-2.6%)	2.3% (1.5-3.4%)	6400 (4000-9300)	170000 (110000-250000)
China	1990	Female	65 to 69	18% (12-26%)	82% (74-88%)	0.34 (0.2-0.52)	1.6% (1-2.4%)	2.1% (1.3-3%)	7500 (4700-11000)	160000 (100000-240000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	1990	Female	70 to 74	18% (11-25%)	82% (75-89%)	0.33 (0.19-0.54)	1.3% (0.75-2.1%)	1.7% (0.93-2.6%)	7900 (4500-12000)	140000 (77000-220000)
China	1990	Female	75 to 79	16% (10-24%)	84% (76-90%)	0.3 (0.16-0.51)	0.97% (0.42-1.7%)	1.2% (0.54-2%)	5600 (2500-9500)	76000 (32000-130000)
China	1990	Female	80 to 84	16% (9.7-23%)	84% (77-90%)	0.3 (0.17-0.48)	0.62% (0.12-1.2%)	0.73% (0.16-1.5%)	2700 (580-5400)	28000 (5600-55000)
China	1990	Female	85 to 89	16% (10-24%)	84% (76-90%)	0.31 (0.16-0.52)	0.43% (-0.011-1%)	0.51% (-0.019-1.2%)	1200 (-45-2800)	8800 (-230-21000)
China	1990	Female	90 to 94	18% (12-27%)	82% (73-88%)	0.35 (0.19-0.6)	0.3% (-0.19-1%)	0.33% (-0.23-1.1%)	310 (-210-1000)	1600 (-1000-5200)
China	1990	Female	95 plus	21% (12-32%)	79% (68-88%)	0.41 (0.18-0.76)	0.25% (-0.39-1.2%)	0.24% (-0.49-1.3%)	55 (-110-300)	220 (-350-1000)
China	1990	Male	All Ages	41% (37-47%)	59% (53-63%)	1.8 (1.6-2.1)	6.3% (5.4-7.2%)	8.8% (7.4-10%)	420000 (340000-500000)	1400000 (1200000-1600000)
China	1990	Male	15 to 19	24% (16-36%)	76% (64-84%)	0.46 (0.28-0.72)	2.3% (1.5-3.2%)	2.8% (1.7-4.4%)	2300 (1400-3600)	210000 (140000-310000)
China	1990	Male	20 to 24	43% (30-56%)	57% (44-70%)	1.4 (0.92-1.9)	5.8% (4.3-7.8%)	7.1% (4.8-9.7%)	8200 (5500-11000)	690000 (500000-910000)
China	1990	Male	25 to 29	52% (38-65%)	48% (35-62%)	2.1 (1.5-2.8)	7.8% (5.8-9.8%)	10% (7-13%)	9800 (6800-13000)	750000 (550000-960000)
China	1990	Male	30 to 34	57% (41-69%)	43% (31-59%)	2.7 (2-3.4)	9% (7-11%)	12% (8.9-15%)	12000 (9000-15000)	810000 (630000-990000)
China	1990	Male	35 to 39	59% (45-71%)	41% (29-55%)	3.3 (2.4-4.3)	9.9% (7.9-12%)	13% (10-16%)	17000 (13000-21000)	1100000 (840000-1300000)
China	1990	Male	40 to 44	61% (48-73%)	39% (27-52%)	3.8 (2.8-4.8)	11% (8.9-13%)	14% (11-17%)	19000 (15000-23000)	1000000 (820000-1200000)
China	1990	Male	45 to 49	62% (47-74%)	38% (26-53%)	4.2 (3.1-5.3)	12% (10-14%)	15% (12-18%)	25000 (20000-30000)	1100000 (930000-1400000)
China	1990	Male	50 to 54	62% (48-74%)	38% (26-52%)	4.3 (3.1-5.4)	13% (10-15%)	15% (12-18%)	37000 (29000-44000)	1400000 (1200000-1700000)
China	1990	Male	55 to 59	60% (47-72%)	40% (28-53%)	4.1 (3-5.4)	12% (9.8-15%)	15% (11-17%)	50000 (39000-60000)	1700000 (1300000-2000000)
China	1990	Male	60 to 64	59% (45-71%)	41% (29-55%)	3.8 (2.7-5.1)	12% (8.9-14%)	13% (10-16%)	59000 (45000-73000)	1700000 (1300000-2000000)
China	1990	Male	65 to 69	55% (42-67%)	45% (33-58%)	3.2 (2.4-4.3)	10% (7.8-13%)	11% (8.7-14%)	58000 (44000-73000)	1300000 (1000000-1700000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	1990	Male	70 to 74	51% (39-63%)	49% (37-61%)	2.7 (1.9-3.7)	8.3% (6-11%)	9.1% (6.5-12%)	57000 (41000-76000)	1100000 (760000-1400000)
China	1990	Male	75 to 79	48% (34-61%)	52% (39-66%)	2.3 (1.5-3.3)	6.9% (4.6-9.7%)	7.5% (4.9-11%)	37000 (25000-53000)	540000 (360000-750000)
China	1990	Male	80 to 84	44% (31-57%)	56% (43-69%)	2 (1.3-2.8)	5.7% (3.6-8%)	6.2% (3.9-8.7%)	18000 (11000-25000)	190000 (120000-270000)
China	1990	Male	85 to 89	42% (30-54%)	58% (46-70%)	1.8 (1.1-2.6)	4.6% (2.8-6.8%)	5% (3-7.4%)	7100 (4300-10000)	54000 (33000-79000)
China	1990	Male	90 to 94	43% (31-55%)	57% (45-69%)	1.8 (1.1-2.6)	3.8% (2.1-5.8%)	4.1% (2.2-6.3%)	2000 (1100-3000)	11000 (5800-16000)
China	1990	Male	95 plus	46% (31-60%)	54% (40-69%)	1.8 (0.99-2.8)	3.4% (1.5-5.7%)	3.5% (1.4-6.1%)	390 (160-680)	1400 (620-2400)
China	1995	Female	All Ages	12% (10-15%)	88% (85-90%)	0.16 (0.13-0.2)	1.1% (0.79-1.4%)	1.6% (1-2.2%)	62000 (40000-85000)	1900000 (1400000-2400000)
China	1995	Female	15 to 19	8.7% (5.5-13%)	91% (87-94%)	0.099 (0.052-0.17)	0.68% (0.46-0.96%)	0.79% (0.43-1.4%)	290 (150-490)	38000 (25000-55000)
China	1995	Female	20 to 24	11% (6.8-18%)	89% (82-93%)	0.11 (0.058-0.2)	0.98% (0.64-1.4%)	0.77% (0.33-1.5%)	570 (240-1100)	94000 (60000-140000)
China	1995	Female	25 to 29	13% (8.1-20%)	87% (80-92%)	0.14 (0.076-0.24)	1% (0.71-1.4%)	0.9% (0.37-1.7%)	800 (330-1500)	110000 (74000-160000)
China	1995	Female	30 to 34	15% (9-24%)	85% (76-91%)	0.18 (0.097-0.29)	1.1% (0.78-1.5%)	1.2% (0.69-2%)	980 (560-1600)	100000 (71000-150000)
China	1995	Female	35 to 39	17% (9.6-25%)	83% (75-90%)	0.23 (0.13-0.38)	1.3% (0.92-1.7%)	1.8% (1.1-2.6%)	1400 (910-2100)	110000 (75000-140000)
China	1995	Female	40 to 44	17% (11-26%)	83% (74-89%)	0.27 (0.16-0.44)	1.5% (1.1-2%)	2.4% (1.6-3.3%)	2700 (1900-3900)	150000 (110000-210000)
China	1995	Female	45 to 49	17% (10-26%)	83% (74-90%)	0.3 (0.17-0.47)	1.7% (1.2-2.4%)	2.8% (1.8-3.9%)	3400 (2200-4800)	150000 (100000-210000)
China	1995	Female	50 to 54	17% (11-26%)	83% (74-89%)	0.32 (0.19-0.51)	1.9% (1.3-2.7%)	3% (2-4.2%)	4000 (2700-5600)	150000 (100000-210000)
China	1995	Female	55 to 59	17% (11-25%)	83% (75-89%)	0.33 (0.19-0.52)	2.1% (1.4-2.9%)	3.1% (2.1-4.3%)	5800 (3900-8000)	190000 (130000-260000)
China	1995	Female	60 to 64	17% (10-26%)	83% (74-90%)	0.34 (0.19-0.57)	2% (1.3-2.8%)	2.7% (1.8-3.9%)	7800 (5300-11000)	210000 (140000-300000)
China	1995	Female	65 to 69	17% (11-25%)	83% (75-89%)	0.36 (0.21-0.56)	1.9% (1.2-2.6%)	2.5% (1.7-3.4%)	9000 (6100-12000)	200000 (130000-270000)
China	1995	Female	70 to 74	17% (11-25%)	83% (75-89%)	0.36 (0.21-0.6)	1.6% (0.92-2.4%)	2% (1.2-3%)	9400 (5600-15000)	170000 (97000-260000)
China	1995	Female	75 to 79	17% (11-26%)	83% (74-89%)	0.37 (0.2-0.61)	1.2% (0.54-2.1%)	1.5% (0.68-2.5%)	7800 (3700-13000)	110000 (48000-180000)
China	1995	Female	80 to 84	18% (11-26%)	82% (74-89%)	0.38 (0.22-0.61)	0.83% (0.25-1.6%)	0.99% (0.31-1.9%)	4600 (1400-8700)	47000 (14000-89000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	1995	Female	85 to 89	19% (12-28%)	81% (72-88%)	0.41 (0.22-0.67)	0.64% (0.074-1.4%)	0.75% (0.092-1.7%)	2200 (270-4900)	16000 (1900-36000)
China	1995	Female	90 to 94	21% (14-30%)	79% (70-86%)	0.46 (0.25-0.78)	0.51% (-0.13-1.4%)	0.57% (-0.16-1.6%)	700 (-200-2000)	3700 (-1000-10000)
China	1995	Female	95 plus	24% (14-36%)	76% (64-86%)	0.53 (0.24-0.96)	0.47% (-0.37-1.7%)	0.49% (-0.46-1.9%)	150 (-140-610)	570 (-450-2100)
China	1995	Male	All Ages	43% (38-48%)	57% (52-62%)	2.2 (2-2.5)	7.7% (6.6-8.8%)	10% (8.5-12%)	510000 (430000-600000)	1700000 (1400000-1900000)
China	1995	Male	15 to 19	23% (15-34%)	77% (66-85%)	0.51 (0.31-0.79)	2.5% (1.7-3.5%)	3.2% (2-4.9%)	1800 (1100-2800)	170000 (110000-250000)
China	1995	Male	20 to 24	42% (30-55%)	58% (45-70%)	1.6 (1.1-2.1)	6.6% (4.9-8.8%)	8.1% (5.5-11%)	8800 (5900-12000)	730000 (530000-980000)
China	1995	Male	25 to 29	53% (39-65%)	47% (35-61%)	2.4 (1.7-3.2)	8.8% (6.6-11%)	11% (8.1-15%)	15000 (11000-19000)	1100000 (810000-1400000)
China	1995	Male	30 to 34	57% (42-69%)	43% (31-58%)	3.1 (2.3-3.9)	9.9% (7.8-12%)	13% (9.8-16%)	17000 (13000-21000)	1100000 (880000-1400000)
China	1995	Male	35 to 39	59% (47-71%)	41% (29-53%)	3.8 (2.8-4.9)	11% (8.9-13%)	14% (11-18%)	19000 (15000-23000)	1100000 (920000-1400000)
China	1995	Male	40 to 44	61% (49-73%)	39% (27-51%)	4.4 (3.3-5.5)	12% (10-15%)	15% (13-18%)	30000 (25000-36000)	1600000 (1300000-1900000)
China	1995	Male	45 to 49	62% (48-74%)	38% (26-52%)	4.8 (3.6-6.1)	13% (11-16%)	16% (13-19%)	33000 (27000-39000)	1500000 (1300000-1800000)
China	1995	Male	50 to 54	62% (48-73%)	38% (27-52%)	4.9 (3.7-6.3)	14% (11-16%)	16% (13-19%)	38000 (31000-45000)	1500000 (1200000-1800000)
China	1995	Male	55 to 59	60% (46-72%)	40% (28-54%)	4.7 (3.4-6.1)	13% (11-16%)	16% (12-18%)	52000 (41000-61000)	1800000 (1400000-2100000)
China	1995	Male	60 to 64	57% (43-70%)	43% (30-57%)	4.2 (3-5.7)	12% (9.8-15%)	14% (11-17%)	69000 (54000-84000)	2000000 (1500000-2400000)
China	1995	Male	65 to 69	54% (41-66%)	46% (34-59%)	3.6 (2.6-4.8)	11% (8.5-14%)	12% (9.4-15%)	69000 (53000-86000)	1600000 (1200000-2000000)
China	1995	Male	70 to 74	51% (38-62%)	49% (38-62%)	3 (2.1-4.2)	9.2% (6.8-12%)	10% (7.4-13%)	63000 (46000-83000)	1200000 (860000-1500000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	1995	Male	75 to 79	48% (35-62%)	52% (38-65%)	2.7 (1.8-3.8)	7.9% (5.4-11%)	8.6% (5.9-12%)	51000 (35000-70000)	730000 (510000-1000000)
China	1995	Male	80 to 84	46% (33-59%)	54% (41-67%)	2.5 (1.6-3.4)	6.8% (4.5-9.3%)	7.4% (4.9-10%)	30000 (20000-41000)	320000 (210000-430000)
China	1995	Male	85 to 89	45% (32-57%)	55% (43-68%)	2.2 (1.4-3.2)	5.7% (3.6-8%)	6.1% (3.9-8.7%)	11000 (7200-16000)	87000 (55000-120000)
China	1995	Male	90 to 94	45% (34-57%)	55% (43-66%)	2.1 (1.4-3.2)	4.7% (2.8-6.9%)	5% (2.9-7.6%)	3100 (1800-4700)	17000 (9700-25000)
China	1995	Male	95 plus	48% (34-62%)	52% (38-66%)	2.1 (1.2-3.3)	4.2% (1.9-6.8%)	4.4% (1.9-7.2%)	590 (250-980)	2200 (950-3500)
China	2000	Female	All Ages	13% (11-16%)	87% (84-89%)	0.16 (0.14-0.2)	1.1% (0.77-1.4%)	1.4% (0.88-2%)	56000 (34000-80000)	1700000 (1300000-2200000)
China	2000	Female	15 to 19	12% (8-18%)	88% (82-92%)	0.14 (0.074-0.23)	0.76% (0.51-1.1%)	0.99% (0.54-1.7%)	320 (170-540)	42000 (28000-59000)
China	2000	Female	20 to 24	13% (8.4-20%)	87% (80-92%)	0.13 (0.068-0.22)	1.1% (0.74-1.5%)	0.87% (0.4-1.6%)	400 (190-720)	71000 (47000-100000)
China	2000	Female	25 to 29	14% (8.7-21%)	86% (79-91%)	0.15 (0.081-0.24)	1.1% (0.77-1.5%)	0.93% (0.46-1.6%)	620 (310-1100)	99000 (67000-140000)
China	2000	Female	30 to 34	15% (9-23%)	85% (77-91%)	0.18 (0.096-0.29)	1.1% (0.78-1.5%)	1.2% (0.68-1.8%)	900 (530-1400)	110000 (77000-150000)
China	2000	Female	35 to 39	16% (9.4-24%)	84% (76-91%)	0.22 (0.12-0.35)	1.2% (0.85-1.5%)	1.6% (1-2.3%)	1300 (830-1800)	110000 (77000-140000)
China	2000	Female	40 to 44	16% (9.8-24%)	84% (76-90%)	0.25 (0.14-0.39)	1.3% (0.94-1.7%)	2% (1.4-2.9%)	1900 (1300-2700)	120000 (85000-160000)
China	2000	Female	45 to 49	16% (9.2-23%)	84% (77-91%)	0.27 (0.15-0.42)	1.5% (0.98-2%)	2.4% (1.5-3.4%)	3400 (2200-4800)	160000 (110000-220000)
China	2000	Female	50 to 54	16% (10-24%)	84% (76-90%)	0.3 (0.18-0.47)	1.7% (1.1-2.3%)	2.7% (1.7-3.8%)	4300 (2700-6000)	160000 (110000-230000)
China	2000	Female	55 to 59	16% (10-24%)	84% (76-90%)	0.31 (0.18-0.49)	1.8% (1.2-2.7%)	2.8% (1.8-4%)	5100 (3300-7400)	160000 (110000-230000)
China	2000	Female	60 to 64	16% (9.6-24%)	84% (76-90%)	0.31 (0.18-0.53)	1.7% (1.1-2.5%)	2.4% (1.5-3.5%)	6700 (4300-9900)	180000 (120000-260000)
China	2000	Female	65 to 69	16% (9.8-23%)	84% (77-90%)	0.32 (0.19-0.5)	1.7% (1-2.4%)	2.2% (1.5-3.2%)	8900 (5700-13000)	200000 (120000-280000)
China	2000	Female	70 to 74	16% (10-24%)	84% (76-90%)	0.33 (0.19-0.54)	1.4% (0.74-2.1%)	1.7% (0.98-2.7%)	8800 (4900-14000)	150000 (84000-240000)
China	2000	Female	75 to 79	16% (10-24%)	84% (76-90%)	0.34 (0.18-0.56)	1% (0.38-1.7%)	1.3% (0.52-2.1%)	6600 (2800-11000)	90000 (35000-160000)
China	2000	Female	80 to 84	17% (11-24%)	83% (76-89%)	0.36 (0.2-0.57)	0.66% (0.089-1.4%)	0.81% (0.14-1.6%)	4000 (710-8200)	41000 (5600-84000)
China	2000	Female	85 to 89	18% (12-26%)	82% (74-88%)	0.39 (0.21-0.63)	0.5% (-0.091-1.3%)	0.58% (-0.1-1.5%)	2100 (-370-5400)	15000 (-2700-40000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2000	Female	90 to 94	20% (14-29%)	80% (71-86%)	0.44 (0.24-0.75)	0.38% (-0.27-1.3%)	0.42% (-0.32-1.5%)	650 (-490-2300)	3400 (-2400-12000)
China	2000	Female	95 plus	23% (14-34%)	77% (66-86%)	0.52 (0.23-0.93)	0.36% (-0.51-1.6%)	0.36% (-0.62-1.8%)	150 (-270-750)	600 (-850-2700)
China	2000	Male	All Ages	45% (40-50%)	55% (50-60%)	2.3 (2.1-2.6)	8.4% (7.2-9.6%)	10% (8.6-12%)	550000 (450000-640000)	1800000 (1500000-2000000)
China	2000	Male	15 to 19	29% (20-40%)	71% (60-80%)	0.64 (0.39-0.97)	3% (2.1-4.1%)	3.9% (2.4-5.8%)	2100 (1300-3100)	20000 (13000-28000)
China	2000	Male	20 to 24	43% (30-55%)	57% (45-70%)	1.6 (1.1-2.1)	7% (5.1-9.4%)	8.6% (5.8-12%)	6900 (4600-9600)	58000 (42000-79000)
China	2000	Male	25 to 29	52% (38-64%)	48% (36-62%)	2.3 (1.7-3.1)	8.8% (6.6-11%)	11% (8-15%)	13000 (8900-17000)	98000 (72000-120000)
China	2000	Male	30 to 34	56% (41-69%)	44% (31-59%)	3 (2.2-3.8)	9.9% (7.8-12%)	13% (9.7-16%)	19000 (14000-23000)	130000 (99000-160000)
China	2000	Male	35 to 39	59% (47-71%)	41% (29-53%)	3.7 (2.8-4.8)	11% (8.9-14%)	14% (11-18%)	22000 (17000-27000)	130000 (110000-160000)
China	2000	Male	40 to 44	61% (49-73%)	39% (27-51%)	4.4 (3.4-5.5)	12% (10-15%)	16% (13-19%)	28000 (23000-33000)	150000 (120000-180000)
China	2000	Male	45 to 49	62% (48-74%)	38% (26-52%)	4.8 (3.6-6.1)	13% (11-16%)	16% (13-19%)	43000 (35000-51000)	200000 (160000-230000)
China	2000	Male	50 to 54	62% (48-73%)	38% (27-52%)	4.9 (3.7-6.2)	13% (11-16%)	16% (13-19%)	44000 (35000-52000)	180000 (140000-210000)
China	2000	Male	55 to 59	60% (46-71%)	40% (29-54%)	4.6 (3.4-6)	13% (10-15%)	15% (12-18%)	50000 (40000-59000)	170000 (140000-200000)
China	2000	Male	60 to 64	56% (43-69%)	44% (31-57%)	4.1 (3-5.5)	12% (9.4-15%)	14% (11-17%)	67000 (52000-81000)	190000 (150000-230000)
China	2000	Male	65 to 69	53% (41-65%)	47% (35-59%)	3.5 (2.5-4.6)	11% (8.2-13%)	12% (9.2-15%)	77000 (59000-97000)	180000 (140000-230000)
China	2000	Male	70 to 74	51% (39-62%)	49% (38-61%)	3 (2.1-4.1)	9.1% (6.6-12%)	10% (7.2-13%)	72000 (52000-93000)	130000 (96000-170000)
China	2000	Male	75 to 79	48% (36-61%)	52% (39-64%)	2.7 (1.8-3.8)	7.8% (5.3-11%)	8.5% (5.8-12%)	52000 (35000-71000)	75000 (51000-100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2000	Male	80 to 84	46% (33-58%)	54% (42-67%)	2.4 (1.6-3.4)	6.6% (4.4-9.2%)	7.2% (4.8-10%)	34000 (22000-47000)	370000 (240000-510000)
China	2000	Male	85 to 89	44% (32-56%)	56% (44-68%)	2.2 (1.4-3.1)	5.5% (3.4-7.8%)	5.9% (3.7-8.5%)	16000 (9600-22000)	120000 (75000-170000)
China	2000	Male	90 to 94	45% (33-56%)	55% (44-67%)	2.1 (1.3-3.1)	4.5% (2.6-6.9%)	4.9% (2.7-7.5%)	4000 (2200-6200)	21000 (12000-33000)
China	2000	Male	95 plus	48% (34-61%)	52% (39-66%)	2.1 (1.2-3.1)	4.1% (1.7-6.9%)	4.2% (1.6-7.3%)	750 (280-1300)	2800 (1100-4700)
China	2005	Female	All Ages	15% (13-16%)	85% (84-87%)	0.18 (0.15-0.21)	0.98% (0.73-1.3%)	1.2% (0.76-1.7%)	47000 (29000-66000)	1500000 (1100000-1900000)
China	2005	Female	15 to 19	15% (11-20%)	85% (80-89%)	0.16 (0.09-0.25)	0.77% (0.52-1.1%)	1.1% (0.59-1.8%)	350 (190-570)	48000 (31000-67000)
China	2005	Female	20 to 24	17% (12-23%)	83% (77-88%)	0.16 (0.091-0.25)	1.1% (0.78-1.5%)	0.98% (0.44-1.7%)	300 (140-510)	63000 (42000-91000)
China	2005	Female	25 to 29	16% (11-21%)	84% (79-89%)	0.16 (0.096-0.24)	1.1% (0.81-1.5%)	0.99% (0.51-1.6%)	330 (170-540)	65000 (45000-91000)
China	2005	Female	30 to 34	16% (12-22%)	84% (78-88%)	0.18 (0.12-0.26)	1.1% (0.76-1.4%)	1.1% (0.68-1.7%)	620 (380-930)	88000 (62000-120000)
China	2005	Female	35 to 39	18% (13-24%)	82% (76-87%)	0.23 (0.15-0.34)	1.1% (0.8-1.4%)	1.5% (0.98-2.1%)	1200 (800-1700)	110000 (81000-150000)
China	2005	Female	40 to 44	18% (14-25%)	82% (75-86%)	0.27 (0.17-0.38)	1.1% (0.81-1.5%)	1.9% (1.3-2.6%)	1700 (1200-2400)	110000 (80000-150000)
China	2005	Female	45 to 49	17% (13-23%)	83% (77-87%)	0.28 (0.19-0.41)	1.2% (0.82-1.7%)	2.1% (1.4-3%)	2400 (1600-3500)	120000 (82000-160000)
China	2005	Female	50 to 54	17% (12-23%)	83% (77-88%)	0.3 (0.2-0.42)	1.4% (0.89-2%)	2.3% (1.5-3.3%)	4600 (3000-6500)	180000 (120000-250000)
China	2005	Female	55 to 59	16% (12-21%)	84% (79-88%)	0.29 (0.19-0.41)	1.5% (0.95-2.2%)	2.4% (1.6-3.5%)	5200 (3400-7700)	170000 (110000-240000)
China	2005	Female	60 to 64	14% (10-19%)	86% (81-90%)	0.27 (0.17-0.41)	1.4% (0.92-2.1%)	2.2% (1.4-3.1%)	5400 (3500-7600)	140000 (91000-200000)
China	2005	Female	65 to 69	14% (9.7-19%)	86% (81-90%)	0.27 (0.18-0.38)	1.4% (0.89-2%)	1.9% (1.3-2.7%)	7200 (4600-10000)	160000 (98000-220000)
China	2005	Female	70 to 74	13% (9.1-19%)	87% (81-91%)	0.26 (0.16-0.42)	1.1% (0.63-1.7%)	1.5% (0.87-2.2%)	7600 (4400-11000)	130000 (76000-200000)
China	2005	Female	75 to 79	13% (8.2-19%)	87% (81-92%)	0.25 (0.14-0.41)	0.76% (0.3-1.3%)	0.98% (0.42-1.6%)	5300 (2300-8700)	72000 (28000-120000)
China	2005	Female	80 to 84	12% (7.3-18%)	88% (82-93%)	0.25 (0.14-0.4)	0.44% (0.018-0.93%)	0.55% (0.044-1.1%)	2900 (230-5900)	29000 (1100-61000)
China	2005	Female	85 to 89	13% (7.8-19%)	87% (81-92%)	0.27 (0.14-0.44)	0.27% (-0.15-0.78%)	0.32% (-0.17-0.9%)	1300 (-690-3600)	9500 (-4900-27000)
China	2005	Female	90 to 94	14% (8.9-21%)	86% (79-91%)	0.31 (0.16-0.52)	0.16% (-0.32-0.77%)	0.15% (-0.38-0.84%)	290 (-720-1600)	1800 (-3500-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2005	Female	95 plus	17% (9.6-26%)	83% (74-90%)	0.37 (0.16-0.67)	0.13% (-0.51-0.95%)	0.087% (-0.65-1%)	46 (-350-550)	270 (-1100-2000)
China	2005	Male	All Ages	47% (44-51%)	53% (49-56%)	2.5 (2.2-2.8)	8.8% (7.5-10%)	10% (8.3-12%)	540000 (450000-640000)	1800000 (1500000-2000000)
China	2005	Male	15 to 19	35% (27-44%)	65% (56-73%)	0.75 (0.49-1.1)	3.2% (2.2-4.4%)	4.5% (2.9-6.6%)	2800 (1800-4200)	260000 (180000-360000)
China	2005	Male	20 to 24	48% (40-58%)	52% (42-60%)	1.8 (1.3-2.4)	7.4% (5.5-9.7%)	9.7% (6.6-14%)	6500 (4400-9000)	550000 (400000-730000)
China	2005	Male	25 to 29	54% (44-64%)	46% (36-56%)	2.4 (1.8-3.2)	9.1% (6.8-11%)	12% (8.8-16%)	8900 (6300-12000)	700000 (520000-890000)
China	2005	Male	30 to 34	57% (47-67%)	43% (33-53%)	3.1 (2.4-3.8)	9.9% (7.7-12%)	13% (9.8-16%)	16000 (12000-20000)	1100000 (860000-1400000)
China	2005	Male	35 to 39	59% (50-69%)	41% (31-50%)	3.7 (2.9-4.7)	11% (8.9-13%)	14% (11-18%)	24000 (19000-30000)	1500000 (1200000-1800000)
China	2005	Male	40 to 44	61% (50-71%)	39% (29-50%)	4.3 (3.4-5.3)	12% (10-15%)	16% (13-19%)	32000 (26000-38000)	1700000 (1400000-2000000)
China	2005	Male	45 to 49	62% (52-72%)	38% (28-48%)	4.7 (3.7-5.9)	13% (11-15%)	16% (13-19%)	37000 (30000-43000)	1700000 (1400000-2000000)
China	2005	Male	50 to 54	62% (52-72%)	38% (28-48%)	4.9 (3.8-6.1)	13% (11-15%)	16% (13-18%)	57000 (46000-67000)	2300000 (1900000-2700000)
China	2005	Male	55 to 59	60% (51-69%)	40% (31-49%)	4.6 (3.5-5.9)	13% (10-15%)	15% (12-18%)	57000 (46000-67000)	2000000 (1600000-2300000)
China	2005	Male	60 to 64	56% (46-66%)	44% (34-54%)	4 (3-5.3)	11% (9-14%)	13% (10-16%)	57000 (44000-69000)	1700000 (1300000-2000000)
China	2005	Male	65 to 69	52% (42-61%)	48% (39-58%)	3.3 (2.5-4.3)	9.9% (7.7-12%)	11% (8.6-14%)	70000 (54000-86000)	1700000 (1300000-2000000)
China	2005	Male	70 to 74	48% (38-58%)	52% (42-62%)	2.8 (2-3.7)	8.2% (5.9-10%)	9.1% (6.5-12%)	69000 (49000-89000)	1300000 (950000-1700000)
China	2005	Male	75 to 79	44% (34-55%)	56% (45-66%)	2.4 (1.6-3.3)	6.8% (4.6-9.2%)	7.4% (5-10%)	51000 (34000-71000)	750000 (510000-1000000)
China	2005	Male	80 to 84	42% (30-55%)	58% (45-70%)	2.1 (1.4-3)	5.7% (3.6-8.1%)	6.3% (3.9-8.8%)	34000 (21000-48000)	370000 (230000-520000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2005	Male	85 to 89	41% (29-52%)	59% (48-71%)	1.9 (1.3-2.8)	4.7% (2.9-6.9%)	5.1% (3.1-7.6%)	16000 (9400-23000)	120000 (74000-180000)
China	2005	Male	90 to 94	42% (31-53%)	58% (47-69%)	1.9 (1.2-2.7)	3.8% (2-6%)	4% (2.1-6.5%)	4700 (2400-7500)	25000 (13000-41000)
China	2005	Male	95 plus	45% (31-59%)	55% (41-69%)	1.8 (1.1-2.8)	3.4% (1.2-6.1%)	3.5% (1.1-6.5%)	810 (260-1500)	3000 (1100-5500)
China	2010	Female	All Ages	14% (13-16%)	86% (84-87%)	0.21 (0.17-0.24)	1.1% (0.8-1.3%)	1.3% (0.87-1.9%)	50000 (32000-70000)	160000 (120000-200000)
China	2010	Female	15 to 19	14% (10-19%)	86% (81-90%)	0.16 (0.092-0.26)	0.74% (0.51-1%)	1.1% (0.62-1.8%)	210 (120-340)	34000 (23000-48000)
China	2010	Female	20 to 24	16% (12-22%)	84% (78-88%)	0.17 (0.096-0.26)	1.1% (0.8-1.6%)	0.99% (0.48-1.7%)	310 (150-530)	75000 (50000-110000)
China	2010	Female	25 to 29	16% (11-22%)	84% (78-89%)	0.17 (0.1-0.27)	1.1% (0.86-1.5%)	1% (0.58-1.6%)	290 (170-470)	65000 (45000-91000)
China	2010	Female	30 to 34	16% (12-22%)	84% (78-88%)	0.2 (0.13-0.29)	1.1% (0.79-1.4%)	1.2% (0.78-1.8%)	420 (280-630)	67000 (46000-92000)
China	2010	Female	35 to 39	18% (13-23%)	82% (77-87%)	0.25 (0.16-0.37)	1.1% (0.81-1.4%)	1.6% (1.1-2.3%)	960 (650-1300)	96000 (70000-130000)
China	2010	Female	40 to 44	18% (13-24%)	82% (76-87%)	0.28 (0.19-0.41)	1.1% (0.82-1.5%)	2.1% (1.4-2.8%)	1900 (1300-2600)	130000 (92000-170000)
China	2010	Female	45 to 49	17% (12-23%)	83% (77-88%)	0.3 (0.2-0.43)	1.2% (0.86-1.7%)	2.4% (1.6-3.4%)	2700 (1800-3800)	140000 (95000-180000)
China	2010	Female	50 to 54	16% (12-22%)	84% (78-88%)	0.32 (0.21-0.45)	1.4% (0.96-2%)	2.7% (1.8-3.7%)	4000 (2600-5500)	160000 (110000-210000)
China	2010	Female	55 to 59	15% (11-20%)	85% (80-89%)	0.31 (0.21-0.44)	1.6% (1.1-2.3%)	2.9% (1.9-4%)	6600 (4300-9200)	210000 (140000-290000)
China	2010	Female	60 to 64	14% (9.8-19%)	86% (81-90%)	0.29 (0.19-0.44)	1.6% (1-2.2%)	2.5% (1.8-3.5%)	6900 (4700-9400)	190000 (130000-250000)
China	2010	Female	65 to 69	13% (9.5-18%)	87% (82-90%)	0.28 (0.18-0.41)	1.5% (0.99-2%)	2.2% (1.6-3%)	6900 (4800-9300)	150000 (100000-210000)
China	2010	Female	70 to 74	13% (8.8-18%)	87% (82-91%)	0.28 (0.17-0.45)	1.2% (0.69-1.7%)	1.6% (1-2.4%)	7300 (4500-11000)	130000 (78000-190000)
China	2010	Female	75 to 79	12% (8.1-18%)	88% (82-92%)	0.28 (0.16-0.44)	0.81% (0.32-1.4%)	1.1% (0.46-1.8%)	5900 (2500-9700)	82000 (32000-140000)
China	2010	Female	80 to 84	12% (7.3-18%)	88% (82-93%)	0.28 (0.15-0.44)	0.49% (0.049-0.97%)	0.61% (0.073-1.2%)	3400 (410-6800)	36000 (3600-72000)
China	2010	Female	85 to 89	13% (7.7-19%)	87% (81-92%)	0.29 (0.15-0.49)	0.31% (-0.15-0.84%)	0.35% (-0.19-1%)	1500 (-830-4300)	12000 (-5700-32000)
China	2010	Female	90 to 94	14% (8.7-21%)	86% (79-91%)	0.33 (0.18-0.57)	0.18% (-0.33-0.86%)	0.16% (-0.44-0.95%)	360 (-990-2100)	2400 (-4600-12000)
China	2010	Female	95 plus	16% (9.2-25%)	84% (75-91%)	0.4 (0.17-0.73)	0.16% (-0.53-1.1%)	0.1% (-0.73-1.2%)	72 (-510-850)	450 (-1500-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2010	Male	All Ages	47% (44-52%)	53% (48-56%)	3 (2.6-3.3)	9.4% (8-11%)	11% (8.8-12%)	580000 (480000-680000)	19000000 (16000000-21000000)
China	2010	Male	15 to 19	32% (25-41%)	68% (59-75%)	0.76 (0.5-1.1)	3.1% (2.2-4.3%)	4.7% (3.1-6.8%)	2000 (1300-2900)	190000 (130000-260000)
China	2010	Male	20 to 24	48% (39-58%)	52% (42-61%)	2 (1.4-2.6)	7.4% (5.5-9.7%)	10% (6.8-14%)	7400 (4900-10000)	650000 (480000-860000)
China	2010	Male	25 to 29	53% (44-63%)	47% (37-56%)	2.7 (2-3.5)	9.3% (7-12%)	13% (9.2-17%)	8400 (5900-11000)	680000 (500000-870000)
China	2010	Male	30 to 34	57% (47-67%)	43% (33-53%)	3.4 (2.6-4.3)	10% (8-13%)	14% (11-18%)	11000 (8600-14000)	830000 (650000-1000000)
China	2010	Male	35 to 39	59% (50-68%)	41% (32-50%)	4.1 (3.2-5.2)	11% (9.2-14%)	15% (12-19%)	21000 (16000-26000)	1300000 (1100000-1600000)
China	2010	Male	40 to 44	60% (50-71%)	40% (29-50%)	4.7 (3.7-5.8)	12% (10-15%)	16% (13-20%)	34000 (28000-41000)	1900000 (1600000-2300000)
China	2010	Male	45 to 49	61% (51-71%)	39% (29-49%)	5.2 (4.1-6.5)	13% (11-15%)	17% (14-20%)	41000 (34000-49000)	2000000 (1700000-2400000)
China	2010	Male	50 to 54	61% (51-71%)	39% (29-49%)	5.4 (4.2-6.6)	13% (11-16%)	16% (14-19%)	50000 (41000-58000)	2100000 (1700000-2400000)
China	2010	Male	55 to 59	59% (50-69%)	41% (31-50%)	5.1 (3.9-6.4)	13% (10-15%)	15% (13-18%)	69000 (57000-82000)	2500000 (2000000-2900000)
China	2010	Male	60 to 64	56% (46-66%)	44% (34-54%)	4.4 (3.3-5.8)	12% (9.3-14%)	14% (11-16%)	67000 (54000-81000)	2000000 (1600000-2400000)
China	2010	Male	65 to 69	51% (42-62%)	49% (38-58%)	3.6 (2.8-4.7)	10% (8.1-12%)	12% (9.3-14%)	66000 (53000-81000)	1600000 (1300000-2000000)
China	2010	Male	70 to 74	47% (38-57%)	53% (43-62%)	3 (2.2-4.1)	8.5% (6.2-11%)	9.6% (7-12%)	71000 (51000-90000)	1400000 (1000000-1700000)
China	2010	Male	75 to 79	44% (35-55%)	56% (45-65%)	2.6 (1.8-3.6)	7.1% (4.9-9.5%)	7.9% (5.4-11%)	60000 (41000-81000)	900000 (620000-1200000)
China	2010	Male	80 to 84	42% (31-55%)	58% (45-69%)	2.4 (1.6-3.3)	6.2% (4.1-8.6%)	6.8% (4.5-9.5%)	45000 (29000-62000)	490000 (330000-680000)
China	2010	Male	85 to 89	41% (30-52%)	59% (48-70%)	2.2 (1.4-3.1)	5.2% (3.2-7.6%)	5.7% (3.5-8.2%)	20000 (12000-29000)	160000 (96000-230000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2010	Male	90 to 94	42% (31-53%)	58% (47-69%)	2.1 (1.4-3)	4.3% (2.3-6.6%)	4.6% (2.3-7.2%)	6000 (3100-9500)	33000 (18000-51000)
China	2010	Male	95 plus	45% (31-59%)	55% (41-69%)	2.1 (1.2-3.2)	3.8% (1.5-6.6%)	3.9% (1.3-7%)	1300 (430-2300)	4800 (1800-8400)
China	2016	Female	All Ages	16% (13-19%)	84% (81-87%)	0.24 (0.2-0.3)	1.2% (0.9-1.5%)	1.6% (1-2.2%)	59000 (39000-81000)	1700000 (1300000-2200000)
China	2016	Female	15 to 19	14% (8.4-21%)	86% (79-92%)	0.17 (0.085-0.29)	0.72% (0.5-1%)	1.2% (0.66-2%)	120 (67-200)	21000 (15000-31000)
China	2016	Female	20 to 24	16% (9.4-25%)	84% (75-91%)	0.18 (0.086-0.3)	1.1% (0.79-1.6%)	1.1% (0.55-1.9%)	190 (97-340)	52000 (34000-77000)
China	2016	Female	25 to 29	18% (10-27%)	82% (73-90%)	0.2 (0.1-0.35)	1.2% (0.86-1.5%)	1.2% (0.65-1.9%)	340 (190-560)	78000 (54000-110000)
China	2016	Female	30 to 34	19% (11-30%)	81% (70-89%)	0.25 (0.13-0.41)	1.1% (0.83-1.5%)	1.4% (0.9-2.2%)	440 (280-680)	70000 (48000-100000)
China	2016	Female	35 to 39	20% (12-31%)	80% (69-88%)	0.3 (0.16-0.5)	1.1% (0.83-1.5%)	2% (1.3-2.9%)	680 (440-1000)	70000 (51000-96000)
China	2016	Female	40 to 44	19% (11-29%)	81% (71-89%)	0.33 (0.18-0.53)	1.2% (0.86-1.6%)	2.5% (1.7-3.5%)	1600 (1100-2300)	110000 (79000-150000)
China	2016	Female	45 to 49	18% (9.6-27%)	82% (73-90%)	0.33 (0.18-0.52)	1.3% (0.92-1.8%)	2.9% (1.9-4%)	3400 (2200-4600)	170000 (120000-220000)
China	2016	Female	50 to 54	16% (9.4-26%)	84% (74-91%)	0.33 (0.18-0.53)	1.5% (1.1-2.1%)	3.1% (2.1-4.2%)	4900 (3400-6600)	190000 (140000-260000)
China	2016	Female	55 to 59	15% (9.1-24%)	85% (76-91%)	0.33 (0.19-0.54)	1.8% (1.2-2.4%)	3.4% (2.4-4.5%)	6200 (4400-8300)	200000 (140000-270000)
China	2016	Female	60 to 64	15% (8.5-25%)	85% (75-92%)	0.34 (0.18-0.58)	1.8% (1.2-2.5%)	3.1% (2.2-4.2%)	9700 (6900-13000)	260000 (180000-360000)
China	2016	Female	65 to 69	15% (8.8-24%)	85% (76-91%)	0.35 (0.19-0.56)	1.6% (1.1-2.3%)	2.6% (1.8-3.6%)	9700 (6800-13000)	220000 (140000-300000)
China	2016	Female	70 to 74	15% (8.9-23%)	85% (77-91%)	0.34 (0.19-0.58)	1.3% (0.77-1.9%)	2% (1.2-2.9%)	7800 (4900-11000)	140000 (81000-210000)
China	2016	Female	75 to 79	14% (8-22%)	86% (78-92%)	0.33 (0.17-0.56)	0.96% (0.42-1.6%)	1.3% (0.63-2.1%)	6600 (3100-11000)	92000 (40000-150000)
China	2016	Female	80 to 84	14% (7.9-21%)	86% (79-92%)	0.33 (0.17-0.56)	0.6% (0.11-1.2%)	0.79% (0.16-1.5%)	4300 (860-8200)	45000 (8300-87000)
China	2016	Female	85 to 89	14% (8.4-23%)	86% (77-92%)	0.35 (0.18-0.6)	0.39% (-0.11-0.99%)	0.46% (-0.15-1.2%)	2400 (-800-6300)	19000 (-5300-48000)
China	2016	Female	90 to 94	16% (9.7-25%)	84% (75-90%)	0.4 (0.21-0.72)	0.23% (-0.38-1%)	0.22% (-0.53-1.1%)	620 (-1500-3200)	4100 (-6700-18000)
China	2016	Female	95 plus	19% (11-31%)	81% (69-89%)	0.49 (0.2-0.92)	0.18% (-0.69-1.3%)	0.11% (-0.91-1.4%)	110 (-900-1400)	740 (-2700-5300)
China	2016	Male	All Ages	48% (44-53%)	52% (47-56%)	3.3 (2.9-3.7)	9.8% (8.4-11%)	11% (9.1-13%)	65000 (54000-76000)	2000000 (1700000-2300000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2016	Male	15 to 19	28% (20-40%)	72% (60-80%)	0.72 (0.45-1.1)	3% (2.1-4.1%)	4.6% (2.9-6.8%)	1200 (770-1800)	120000 (87000-170000)
China	2016	Male	20 to 24	42% (30-54%)	58% (46-70%)	1.8 (1.2-2.5)	7.1% (5.2-9.5%)	9.9% (6.5-14%)	4900 (3200-7000)	460000 (330000-610000)
China	2016	Male	25 to 29	49% (36-62%)	51% (38-64%)	2.6 (1.9-3.6)	8.7% (6.5-11%)	12% (8.6-17%)	9500 (6600-13000)	790000 (590000-1000000)
China	2016	Male	30 to 34	54% (39-67%)	46% (33-61%)	3.4 (2.4-4.5)	9.8% (7.6-12%)	14% (10-18%)	11000 (8000-14000)	810000 (620000-1000000)
China	2016	Male	35 to 39	57% (45-70%)	43% (30-55%)	4.2 (3.1-5.6)	11% (8.6-13%)	15% (12-19%)	13000 (10000-16000)	880000 (690000-1100000)
China	2016	Male	40 to 44	60% (48-73%)	40% (27-52%)	4.9 (3.8-6.3)	12% (9.8-14%)	16% (13-20%)	26000 (21000-31000)	1500000 (1200000-1800000)
China	2016	Male	45 to 49	62% (49-74%)	38% (26-51%)	5.6 (4.3-7.1)	13% (11-15%)	17% (14-20%)	45000 (37000-53000)	2200000 (1800000-2600000)
China	2016	Male	50 to 54	63% (50-74%)	37% (26-50%)	5.9 (4.5-7.4)	13% (11-15%)	17% (14-20%)	57000 (47000-66000)	2400000 (2000000-2800000)
China	2016	Male	55 to 59	62% (50-73%)	38% (27-50%)	5.6 (4.3-7.2)	13% (11-15%)	16% (13-19%)	63000 (52000-74000)	2200000 (1900000-2700000)
China	2016	Male	60 to 64	59% (46-71%)	41% (29-54%)	5.1 (3.7-6.7)	12% (9.9-15%)	14% (12-17%)	94000 (76000-110000)	2800000 (2300000-3400000)
China	2016	Male	65 to 69	55% (43-67%)	45% (33-57%)	4.2 (3.1-5.5)	11% (8.7-13%)	13% (10-16%)	93000 (73000-110000)	2300000 (1800000-2800000)
China	2016	Male	70 to 74	50% (38-61%)	50% (39-62%)	3.4 (2.4-4.7)	9.1% (6.6-12%)	10% (7.5-13%)	74000 (54000-94000)	1400000 (1100000-1800000)
China	2016	Male	75 to 79	46% (33-59%)	54% (41-67%)	2.9 (1.9-4)	7.6% (5.1-10%)	8.4% (5.7-11%)	66000 (45000-89000)	1000000 (680000-1300000)
China	2016	Male	80 to 84	43% (31-56%)	57% (44-69%)	2.6 (1.7-3.6)	6.5% (4.3-9%)	7.2% (4.7-10%)	53000 (34000-73000)	580000 (390000-810000)
China	2016	Male	85 to 89	42% (31-53%)	58% (47-69%)	2.4 (1.6-3.4)	5.6% (3.7-8.1%)	6.1% (3.9-8.9%)	30000 (19000-43000)	230000 (150000-340000)
China	2016	Male	90 to 94	44% (33-55%)	56% (45-67%)	2.3 (1.5-3.4)	4.7% (2.6-7.1%)	5.1% (2.7-7.8%)	9300 (4900-14000)	51000 (28000-77000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
China	2016	Male	95 plus	47% (33-60%)	53% (40-67%)	2.3 (1.3-3.5)	4.2% (1.7-7.2%)	4.4% (1.5-7.6%)	1800 (640-3200)	6900 (2800-12000)
North Korea	1990	Female	All Ages	16% (13-20%)	84% (80-87%)	0.26 (0.22-0.31)	0.85% (0.5-1.2%)	1.2% (0.47-2.1%)	680 (270-1200)	24000 (15000-35000)
North Korea	1990	Female	15 to 19	16% (9.8-25%)	84% (75-90%)	0.25 (0.13-0.43)	0.92% (0.56-1.5%)	1.4% (0.63-2.6%)	8.6 (3.7-17)	1100 (680-1800)
North Korea	1990	Female	20 to 24	19% (12-29%)	81% (71-88%)	0.27 (0.13-0.47)	1.3% (0.8-1.9%)	1.3% (0.47-2.7%)	10 (3.6-20)	1800 (1100-2800)
North Korea	1990	Female	25 to 29	20% (11-31%)	80% (69-89%)	0.28 (0.15-0.48)	1.3% (0.81-1.8%)	1.3% (0.51-2.4%)	8.7 (3.4-16)	1400 (890-2100)
North Korea	1990	Female	30 to 34	21% (12-33%)	79% (67-88%)	0.32 (0.17-0.54)	1.2% (0.77-1.7%)	1.4% (0.66-2.5%)	13 (6.1-24)	1600 (1000-2400)
North Korea	1990	Female	35 to 39	22% (13-35%)	78% (65-87%)	0.4 (0.21-0.68)	1.2% (0.8-1.9%)	1.8% (0.91-3.1%)	13 (6.6-23)	1100 (660-1600)
North Korea	1990	Female	40 to 44	23% (14-36%)	77% (64-86%)	0.46 (0.25-0.75)	1.3% (0.77-2%)	2.1% (1.1-3.4%)	31 (16-51)	1900 (1100-2900)
North Korea	1990	Female	45 to 49	22% (12-35%)	78% (65-88%)	0.48 (0.25-0.81)	1.3% (0.69-2.2%)	2.1% (1-3.7%)	50 (24-86)	2400 (1300-4000)
North Korea	1990	Female	50 to 54	22% (13-34%)	78% (66-87%)	0.5 (0.29-0.82)	1.4% (0.68-2.5%)	2.2% (1-3.8%)	74 (33-130)	2900 (1400-5000)
North Korea	1990	Female	55 to 59	21% (12-32%)	79% (68-88%)	0.51 (0.25-0.85)	1.5% (0.52-2.7%)	2.2% (0.79-3.9%)	82 (30-150)	2700 (980-4900)
North Korea	1990	Female	60 to 64	19% (11-31%)	81% (69-89%)	0.49 (0.24-0.87)	1.3% (0.41-2.7%)	1.9% (0.65-3.7%)	86 (30-170)	2400 (750-4800)
North Korea	1990	Female	65 to 69	19% (11-29%)	81% (71-89%)	0.49 (0.25-0.83)	1.3% (0.34-2.5%)	1.7% (0.51-3.3%)	91 (27-180)	2100 (590-4100)
North Korea	1990	Female	70 to 74	18% (11-28%)	82% (72-89%)	0.47 (0.23-0.85)	1.1% (0.2-2.4%)	1.4% (0.29-3%)	85 (17-180)	1500 (280-3400)
North Korea	1990	Female	75 to 79	16% (9.9-27%)	84% (73-90%)	0.45 (0.22-0.8)	0.83% (-0.018-2%)	1% (0.0095-2.5%)	63 (0.59-150)	890 (-20-2200)
North Korea	1990	Female	80 to 84	16% (9.3-26%)	84% (74-91%)	0.43 (0.22-0.73)	0.58% (-0.18-1.7%)	0.69% (-0.2-2%)	36 (-10-110)	380 (-110-1100)
North Korea	1990	Female	85 to 89	16% (9.3-25%)	84% (75-91%)	0.44 (0.22-0.78)	0.45% (-0.3-1.5%)	0.51% (-0.38-1.8%)	22 (-16-74)	170 (-110-560)
North Korea	1990	Female	90 to 94	18% (11-27%)	82% (73-89%)	0.48 (0.23-0.86)	0.36% (-0.44-1.6%)	0.38% (-0.55-1.8%)	7.8 (-11-38)	45 (-55-200)
North Korea	1990	Female	95 plus	20% (11-33%)	80% (67-89%)	0.54 (0.23-1)	0.36% (-0.62-1.9%)	0.32% (-0.8-2.1%)	2.2 (-5.4-14)	9.3 (-16-50)
North Korea	1990	Male	All Ages	37% (31-45%)	63% (55-69%)	1.5 (1.4-1.7)	5.6% (4.4-6.8%)	7.9% (6-9.9%)	4100 (3000-5200)	170000 (130000-200000)
North Korea	1990	Male	15 to 19	30% (19-45%)	70% (55-81%)	0.54 (0.27-0.97)	2.6% (1.6-4.1%)	3.3% (1.7-5.6%)	44 (22-76)	4100 (2400-6600)
North Korea	1990	Male	20 to 24	42% (28-58%)	58% (42-72%)	1.3 (0.73-2)	5.8% (3.7-8.5%)	6.9% (3.7-11%)	130 (71-220)	12000 (7100-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	1990	Male	25 to 29	48% (31-65%)	52% (35-69%)	1.9 (1.1-2.9)	7.8% (4.9-11%)	9.8% (5.5-15%)	150 (81-220)	11000 (6900-17000)
North Korea	1990	Male	30 to 34	51% (34-68%)	49% (32-66%)	2.4 (1.5-3.5)	8.7% (5.8-12%)	11% (7.1-16%)	210 (130-310)	15000 (9600-20000)
North Korea	1990	Male	35 to 39	54% (37-70%)	46% (30-63%)	2.9 (1.8-4.4)	10% (6.7-14%)	13% (8.1-18%)	160 (98-230)	9600 (6200-13000)
North Korea	1990	Male	40 to 44	55% (39-72%)	45% (28-61%)	3.4 (2.1-4.9)	11% (7.4-14%)	14% (9.1-19%)	310 (200-430)	17000 (11000-23000)
North Korea	1990	Male	45 to 49	56% (37-73%)	44% (27-63%)	3.8 (2.4-5.5)	11% (7.9-15%)	14% (9.4-19%)	490 (330-660)	23000 (16000-30000)
North Korea	1990	Male	50 to 54	55% (37-73%)	45% (27-63%)	3.9 (2.5-5.6)	11% (7.5-15%)	14% (8.9-18%)	630 (390-860)	25000 (16000-34000)
North Korea	1990	Male	55 to 59	54% (36-71%)	46% (29-64%)	3.6 (2.2-5.6)	11% (6.8-15%)	13% (7.8-17%)	550 (320-780)	19000 (11000-26000)
North Korea	1990	Male	60 to 64	51% (33-69%)	49% (31-67%)	3.2 (1.8-5.2)	9.2% (5.1-14%)	11% (5.7-16%)	430 (220-670)	13000 (6700-19000)
North Korea	1990	Male	65 to 69	48% (32-65%)	52% (35-68%)	2.7 (1.5-4.1)	7.9% (4.2-12%)	9% (4.6-14%)	370 (190-600)	8700 (4600-14000)
North Korea	1990	Male	70 to 74	44% (29-61%)	56% (39-71%)	2.3 (1.2-3.6)	6.3% (2.9-10%)	7% (3.1-12%)	280 (130-490)	5500 (2600-9200)
North Korea	1990	Male	75 to 79	41% (27-59%)	59% (41-73%)	1.9 (0.99-3.3)	5.2% (2.1-9.4%)	5.8% (2.2-11%)	190 (74-360)	2900 (1100-5300)
North Korea	1990	Male	80 to 84	39% (24-56%)	61% (44-76%)	1.8 (0.9-2.9)	4.6% (1.9-8.3%)	5% (2-9.1%)	77 (29-140)	850 (340-1500)
North Korea	1990	Male	85 to 89	38% (24-53%)	62% (47-76%)	1.6 (0.85-2.7)	3.8% (1.4-7.1%)	4% (1.4-7.8%)	35 (12-68)	280 (100-530)
North Korea	1990	Male	90 to 94	39% (27-55%)	61% (45-73%)	1.6 (0.82-2.7)	3.1% (0.88-6.6%)	3.3% (0.77-7.2%)	12 (2.8-27)	66 (19-140)
North Korea	1990	Male	95 plus	43% (26-62%)	57% (38-74%)	1.7 (0.74-3.1)	2.8% (0.24-6.9%)	2.9% (-0.041-7.3%)	2.8 (-0.042-7.2)	11 (0.92-26)
North Korea	1995	Female	All Ages	16% (13-20%)	84% (80-87%)	0.28 (0.24-0.34)	0.55% (0.3-0.91%)	0.87% (0.37-1.6%)	930 (400-1600)	31000 (18000-46000)
North Korea	1995	Female	15 to 19	16% (10-25%)	84% (75-90%)	0.26 (0.14-0.47)	0.76% (0.46-1.2%)	0.97% (0.39-1.9%)	6.2 (2.7-12)	790 (490-1300)
North Korea	1995	Female	20 to 24	20% (12-30%)	80% (70-88%)	0.29 (0.14-0.51)	1.1% (0.65-1.7%)	0.92% (0.33-1.9%)	11 (4.1-22)	1900 (1100-3000)
North Korea	1995	Female	25 to 29	20% (12-31%)	80% (69-88%)	0.3 (0.16-0.49)	1% (0.64-1.5%)	0.93% (0.35-1.8%)	13 (5.1-24)	2000 (1200-2900)
North Korea	1995	Female	30 to 34	21% (12-33%)	79% (67-88%)	0.34 (0.19-0.56)	1% (0.64-1.5%)	1.1% (0.48-1.9%)	13 (6.4-24)	1600 (990-2300)
North Korea	1995	Female	35 to 39	23% (13-35%)	77% (65-87%)	0.42 (0.23-0.71)	1.1% (0.65-1.6%)	1.4% (0.69-2.5%)	26 (13-46)	2100 (1300-3200)
North Korea	1995	Female	40 to 44	23% (14-36%)	77% (64-86%)	0.48 (0.26-0.79)	1.2% (0.69-1.9%)	1.7% (0.86-3%)	24 (13-41)	1400 (840-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	1995	Female	45 to 49	22% (13-35%)	78% (65-87%)	0.52 (0.26-0.88)	1.2% (0.66-2.2%)	1.9% (0.87-3.4%)	53 (25-95)	2500 (1300-4300)
North Korea	1995	Female	50 to 54	22% (13-34%)	78% (66-87%)	0.54 (0.31-0.9)	1.4% (0.66-2.4%)	2% (0.96-3.5%)	89 (41-150)	3500 (1700-5900)
North Korea	1995	Female	55 to 59	21% (13-32%)	79% (68-87%)	0.54 (0.29-0.9)	1.4% (0.55-2.6%)	2% (0.82-3.6%)	120 (45-210)	3900 (1500-7000)
North Korea	1995	Female	60 to 64	20% (11-32%)	80% (68-89%)	0.53 (0.27-0.91)	1.3% (0.45-2.5%)	1.8% (0.64-3.3%)	120 (44-230)	3300 (1100-6400)
North Korea	1995	Female	65 to 69	19% (11-30%)	81% (70-89%)	0.52 (0.27-0.87)	1.3% (0.43-2.4%)	1.6% (0.61-3.1%)	130 (47-250)	2900 (1000-5700)
North Korea	1995	Female	70 to 74	18% (11-28%)	82% (72-89%)	0.5 (0.26-0.88)	1.1% (0.24-2.3%)	1.3% (0.33-2.8%)	130 (30-260)	2300 (490-4800)
North Korea	1995	Female	75 to 79	17% (10-28%)	83% (72-90%)	0.48 (0.23-0.85)	0.81% (0.042-2%)	0.96% (0.077-2.3%)	96 (8.2-230)	1400 (72-3300)
North Korea	1995	Female	80 to 84	16% (9.6-26%)	84% (74-90%)	0.46 (0.24-0.78)	0.57% (-0.097-1.5%)	0.64% (-0.1-1.7%)	54 (-9.5-140)	570 (-100-1500)
North Korea	1995	Female	85 to 89	17% (9.9-25%)	83% (75-90%)	0.48 (0.24-0.82)	0.44% (-0.17-1.3%)	0.47% (-0.2-1.5%)	32 (-14-99)	250 (-93-750)
North Korea	1995	Female	90 to 94	18% (11-27%)	82% (73-89%)	0.52 (0.25-0.95)	0.34% (-0.27-1.3%)	0.33% (-0.32-1.4%)	14 (-12-55)	76 (-58-290)
North Korea	1995	Female	95 plus	21% (11-34%)	79% (66-89%)	0.59 (0.24-1.2)	0.3% (-0.34-1.4%)	0.27% (-0.41-1.4%)	4.2 (-6.8-22)	18 (-21-81)
North Korea	1995	Male	All Ages	38% (31-46%)	62% (54-69%)	1.7 (1.5-2)	3.5% (2.3-5.4%)	5.4% (3.5-7.9%)	5400 (4000-6800)	210000 (170000-250000)
North Korea	1995	Male	15 to 19	30% (19-45%)	70% (55-81%)	0.57 (0.29-1)	2.3% (1.4-3.7%)	2.7% (1.4-4.9%)	32 (16-58)	3000 (1800-4900)
North Korea	1995	Male	20 to 24	42% (28-58%)	58% (42-72%)	1.4 (0.77-2.2)	5.3% (3.3-7.9%)	6% (3.2-9.6%)	150 (80-240)	12000 (7600-19000)
North Korea	1995	Male	25 to 29	48% (31-65%)	52% (35-69%)	2 (1.2-3.1)	6.8% (4.3-10%)	8% (4.3-13%)	220 (130-340)	17000 (10000-24000)
North Korea	1995	Male	30 to 34	51% (33-69%)	49% (31-67%)	2.6 (1.6-3.8)	7.9% (5.2-11%)	9.4% (5.7-14%)	220 (130-320)	15000 (9800-21000)
North Korea	1995	Male	35 to 39	54% (37-71%)	46% (29-63%)	3.1 (1.9-4.8)	8.6% (5.6-12%)	10% (6.3-15%)	330 (200-480)	20000 (13000-28000)
North Korea	1995	Male	40 to 44	55% (39-73%)	45% (27-61%)	3.6 (2.2-5.3)	9.6% (6.3-13%)	11% (7.1-16%)	250 (160-350)	13000 (9200-18000)
North Korea	1995	Male	45 to 49	56% (37-72%)	44% (28-63%)	4 (2.5-5.9)	10% (6.8-14%)	12% (7.7-16%)	510 (330-690)	23000 (16000-31000)
North Korea	1995	Male	50 to 54	55% (38-73%)	45% (27-62%)	4.1 (2.6-6.1)	10% (6.8-14%)	12% (7.7-16%)	750 (490-1000)	30000 (20000-40000)
North Korea	1995	Male	55 to 59	54% (36-71%)	46% (29-64%)	3.9 (2.3-5.7)	9.7% (6.3-13%)	11% (7-15%)	910 (550-1300)	31000 (20000-43000)
North Korea	1995	Male	60 to 64	51% (33-69%)	49% (31-67%)	3.4 (1.9-5.5)	8.5% (4.8-12%)	9.5% (5.1-14%)	710 (380-1100)	20000 (11000-31000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	1995	Male	65 to 69	48% (31-65%)	52% (35-69%)	2.9 (1.6-4.3)	7.3% (4.1-11%)	8% (4.4-12%)	500 (270-770)	12000 (6400-18000)
North Korea	1995	Male	70 to 74	44% (29-61%)	56% (39-71%)	2.4 (1.3-3.8)	5.6% (2.7-9.2%)	6.1% (2.8-10%)	350 (170-590)	6800 (3300-11000)
North Korea	1995	Male	75 to 79	41% (26-59%)	59% (41-74%)	2 (1-3.5)	4.6% (1.9-8.1%)	4.9% (1.9-8.8%)	240 (95-440)	3600 (1500-6400)
North Korea	1995	Male	80 to 84	39% (24-56%)	61% (44-76%)	1.8 (0.97-3.1)	3.8% (1.5-7%)	3.9% (1.5-7.5%)	110 (43-190)	1200 (510-2100)
North Korea	1995	Male	85 to 89	38% (24-53%)	62% (47-76%)	1.7 (0.89-2.8)	3% (1.1-5.8%)	3.1% (1-6.3%)	57 (20-110)	450 (170-820)
North Korea	1995	Male	90 to 94	40% (26-55%)	60% (45-74%)	1.7 (0.87-2.9)	2.4% (0.62-5.2%)	2.4% (0.52-5.4%)	17 (4.2-38)	95 (26-200)
North Korea	1995	Male	95 plus	43% (26-62%)	57% (38-74%)	1.8 (0.78-3.3)	1.8% (0.16-4.6%)	1.7% (0.048-4.7%)	4.2 (0.16-10)	17 (1.9-39)
North Korea	2000	Female	All Ages	16% (13-20%)	84% (80-87%)	0.27 (0.22-0.32)	0.52% (0.27-0.89%)	0.77% (0.26-1.5%)	970 (320-1800)	32000 (17000-50000)
North Korea	2000	Female	15 to 19	16% (9.9-25%)	84% (75-90%)	0.25 (0.13-0.44)	0.74% (0.44-1.2%)	0.94% (0.39-1.9%)	8 (3.4-16)	990 (580-1600)
North Korea	2000	Female	20 to 24	20% (12-30%)	80% (70-88%)	0.27 (0.13-0.49)	1% (0.63-1.6%)	0.91% (0.33-1.9%)	7.9 (2.9-16)	1300 (780-2000)
North Korea	2000	Female	25 to 29	20% (12-32%)	80% (68-88%)	0.28 (0.14-0.46)	0.99% (0.63-1.4%)	0.87% (0.32-1.7%)	13 (5-25)	2000 (1200-3000)
North Korea	2000	Female	30 to 34	21% (12-34%)	79% (66-88%)	0.31 (0.17-0.52)	0.94% (0.61-1.4%)	0.96% (0.43-1.8%)	19 (8.5-34)	2200 (1400-3300)
North Korea	2000	Female	35 to 39	23% (14-35%)	77% (65-86%)	0.39 (0.21-0.65)	1% (0.62-1.5%)	1.3% (0.61-2.2%)	24 (12-43)	1900 (1200-2900)
North Korea	2000	Female	40 to 44	23% (14-36%)	77% (64-86%)	0.45 (0.25-0.73)	1.1% (0.59-1.7%)	1.5% (0.73-2.6%)	44 (22-75)	2600 (1500-4100)
North Korea	2000	Female	45 to 49	23% (13-36%)	77% (64-87%)	0.48 (0.24-0.82)	1.1% (0.53-2.1%)	1.7% (0.73-3.1%)	37 (17-68)	1800 (870-3100)
North Korea	2000	Female	50 to 54	22% (13-35%)	78% (65-87%)	0.5 (0.27-0.85)	1.3% (0.55-2.3%)	1.8% (0.77-3.3%)	82 (35-150)	3200 (1400-5800)
North Korea	2000	Female	55 to 59	21% (13-32%)	79% (68-87%)	0.5 (0.26-0.83)	1.3% (0.38-2.4%)	1.7% (0.54-3.3%)	120 (36-230)	3900 (1200-7400)
North Korea	2000	Female	60 to 64	20% (11-32%)	80% (68-89%)	0.48 (0.25-0.81)	1.1% (0.28-2.2%)	1.5% (0.39-3%)	140 (34-280)	3900 (860-7600)
North Korea	2000	Female	65 to 69	19% (11-30%)	81% (70-89%)	0.48 (0.26-0.8)	1.1% (0.24-2.2%)	1.4% (0.36-2.8%)	140 (36-290)	3200 (730-6600)
North Korea	2000	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.47 (0.25-0.84)	0.9% (0.07-2.1%)	1.1% (0.13-2.6%)	140 (15-300)	2400 (170-5600)
North Korea	2000	Female	75 to 79	17% (10-27%)	83% (73-90%)	0.44 (0.22-0.81)	0.67% (-0.098-1.8%)	0.79% (-0.096-2.2%)	100 (-12-270)	1400 (-210-3800)
North Korea	2000	Female	80 to 84	17% (9.7-26%)	83% (74-90%)	0.43 (0.22-0.75)	0.43% (-0.24-1.4%)	0.49% (-0.28-1.6%)	51 (-30-170)	540 (-300-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	2000	Female	85 to 89	17% (10-26%)	83% (74-90%)	0.44 (0.23-0.77)	0.31% (-0.3-1.2%)	0.33% (-0.35-1.3%)	28 (-29-110)	210 (-200-830)
North Korea	2000	Female	90 to 94	18% (11-28%)	82% (72-89%)	0.48 (0.23-0.85)	0.21% (-0.37-1.1%)	0.2% (-0.44-1.2%)	9 (-19-52)	53 (-93-270)
North Korea	2000	Female	95 plus	21% (11-35%)	79% (65-89%)	0.54 (0.22-1.1)	0.17% (-0.47-1.2%)	0.14% (-0.56-1.2%)	2.6 (-10-23)	12 (-35-87)
North Korea	2000	Male	All Ages	38% (32-46%)	62% (54-68%)	1.6 (1.4-1.9)	3.7% (2.5-5.5%)	5.5% (3.5-8%)	6300 (4500-8100)	240000 (190000-300000)
North Korea	2000	Male	15 to 19	30% (19-44%)	70% (56-81%)	0.52 (0.26-0.93)	2.3% (1.3-3.6%)	2.7% (1.3-4.7%)	41 (20-73)	3700 (2200-6100)
North Korea	2000	Male	20 to 24	42% (28-58%)	58% (42-72%)	1.2 (0.69-2)	5.4% (3.4-8.2%)	6.1% (3.3-9.8%)	110 (60-180)	9200 (5600-14000)
North Korea	2000	Male	25 to 29	48% (32-65%)	52% (35-68%)	1.9 (1.1-2.9)	6.8% (4.2-10%)	7.8% (4.4-12%)	240 (130-380)	18000 (11000-26000)
North Korea	2000	Male	30 to 34	51% (34-68%)	49% (32-66%)	2.4 (1.5-3.5)	7.6% (5.1-11%)	8.9% (5.5-13%)	330 (200-490)	22000 (14000-31000)
North Korea	2000	Male	35 to 39	54% (38-70%)	46% (30-62%)	2.8 (1.7-4.2)	8.5% (5.5-12%)	10% (6-15%)	330 (200-500)	20000 (12000-28000)
North Korea	2000	Male	40 to 44	55% (39-73%)	45% (27-61%)	3.3 (2-4.8)	9.2% (6-13%)	11% (6.6-15%)	510 (320-730)	27000 (17000-37000)
North Korea	2000	Male	45 to 49	56% (37-72%)	44% (28-63%)	3.6 (2.2-5.4)	10% (6.5-14%)	12% (7.2-16%)	390 (250-540)	18000 (12000-24000)
North Korea	2000	Male	50 to 54	56% (37-73%)	44% (27-63%)	3.7 (2.3-5.6)	10% (6.3-14%)	11% (7-16%)	750 (450-1000)	30000 (19000-40000)
North Korea	2000	Male	55 to 59	54% (36-73%)	46% (27-64%)	3.5 (2.1-5.3)	9.4% (5.6-13%)	11% (6.1-15%)	1000 (590-1500)	35000 (21000-49000)
North Korea	2000	Male	60 to 64	51% (33-70%)	49% (30-67%)	3.1 (1.7-4.9)	8.2% (4.5-12%)	9% (4.8-14%)	1100 (560-1600)	31000 (17000-46000)
North Korea	2000	Male	65 to 69	48% (32-65%)	52% (35-68%)	2.6 (1.6-3.9)	7% (3.7-10%)	7.6% (3.9-12%)	690 (360-1100)	16000 (8900-25000)
North Korea	2000	Male	70 to 74	45% (29-62%)	55% (38-71%)	2.2 (1.2-3.6)	5.3% (2.4-8.9%)	5.7% (2.5-9.7%)	390 (180-670)	7500 (3400-13000)
North Korea	2000	Male	75 to 79	42% (27-60%)	58% (40-73%)	1.9 (0.97-3.2)	4.3% (1.6-7.9%)	4.5% (1.7-8.5%)	240 (88-450)	3500 (1300-6500)
North Korea	2000	Male	80 to 84	39% (24-56%)	61% (44-76%)	1.7 (0.9-2.8)	3.4% (1.3-6.6%)	3.6% (1.2-7%)	100 (39-200)	1100 (450-2100)
North Korea	2000	Male	85 to 89	38% (25-54%)	62% (46-75%)	1.6 (0.81-2.6)	2.7% (0.9-5.5%)	2.7% (0.8-5.8%)	65 (21-130)	510 (180-1000)
North Korea	2000	Male	90 to 94	40% (26-55%)	60% (45-74%)	1.5 (0.79-2.7)	2% (0.43-4.7%)	2% (0.33-4.9%)	21 (4.1-49)	120 (29-270)
North Korea	2000	Male	95 plus	43% (26-62%)	57% (38-74%)	1.6 (0.7-3)	1.6% (0.093-4.5%)	1.5% (-0.042-4.5%)	4.4 (-0.14-12)	18 (1.2-45)
North Korea	2005	Female	All Ages	17% (14-20%)	83% (80-86%)	0.24 (0.2-0.28)	0.63% (0.27-1.1%)	0.73% (0.043-1.6%)	730 (45-1600)	26000 (11000-45000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	2005	Female	15 to 19	16% (10-26%)	84% (74-90%)	0.21 (0.11-0.36)	0.83% (0.5-1.3%)	1.2% (0.5-2.2%)	7.4 (2.9-14)	940 (570-1500)
North Korea	2005	Female	20 to 24	20% (12-30%)	80% (70-88%)	0.23 (0.11-0.42)	1.1% (0.7-1.8%)	1.1% (0.39-2.2%)	8.2 (2.8-17)	1500 (870-2300)
North Korea	2005	Female	25 to 29	20% (12-31%)	80% (69-88%)	0.24 (0.12-0.41)	1.1% (0.72-1.5%)	1% (0.4-2%)	7.9 (2.9-16)	1300 (790-1900)
North Korea	2005	Female	30 to 34	21% (12-34%)	79% (66-88%)	0.27 (0.14-0.45)	1% (0.64-1.5%)	1.1% (0.45-1.9%)	16 (6.6-29)	2000 (1200-3000)
North Korea	2005	Female	35 to 39	23% (13-35%)	77% (65-87%)	0.33 (0.18-0.54)	1% (0.62-1.5%)	1.3% (0.58-2.3%)	28 (12-49)	2400 (1400-3600)
North Korea	2005	Female	40 to 44	23% (14-36%)	77% (64-86%)	0.38 (0.21-0.61)	1% (0.57-1.6%)	1.5% (0.68-2.6%)	34 (15-57)	2100 (1200-3200)
North Korea	2005	Female	45 to 49	23% (12-35%)	77% (65-88%)	0.41 (0.21-0.69)	1% (0.48-1.8%)	1.6% (0.64-2.9%)	56 (22-100)	2700 (1300-4800)
North Korea	2005	Female	50 to 54	22% (13-35%)	78% (65-87%)	0.43 (0.23-0.7)	1.1% (0.41-2%)	1.6% (0.6-3.1%)	46 (16-86)	1800 (670-3300)
North Korea	2005	Female	55 to 59	21% (13-33%)	79% (67-87%)	0.43 (0.22-0.71)	1% (0.22-2.1%)	1.5% (0.3-3%)	86 (18-180)	2800 (560-5800)
North Korea	2005	Female	60 to 64	20% (11-32%)	80% (68-89%)	0.41 (0.21-0.71)	0.88% (0.04-2%)	1.2% (0.12-2.7%)	110 (9.6-240)	2900 (130-6600)
North Korea	2005	Female	65 to 69	19% (11-30%)	81% (70-89%)	0.41 (0.22-0.67)	0.8% (-0.023-1.9%)	1.1% (0.031-2.5%)	130 (3.7-290)	2800 (-75-6600)
North Korea	2005	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.4 (0.21-0.72)	0.65% (-0.16-1.7%)	0.84% (-0.17-2.2%)	110 (-21-280)	1900 (-460-5200)
North Korea	2005	Female	75 to 79	17% (11-27%)	83% (73-89%)	0.38 (0.19-0.68)	0.42% (-0.35-1.5%)	0.53% (-0.39-1.9%)	70 (-51-250)	960 (-790-3600)
North Korea	2005	Female	80 to 84	17% (9.6-26%)	83% (74-90%)	0.37 (0.19-0.63)	0.2% (-0.53-1.2%)	0.24% (-0.65-1.4%)	25 (-66-150)	260 (-690-1500)
North Korea	2005	Female	85 to 89	17% (10-26%)	83% (74-90%)	0.38 (0.19-0.67)	0.089% (-0.62-1.1%)	0.083% (-0.75-1.3%)	5.9 (-54-90)	55 (-390-680)
North Korea	2005	Female	90 to 94	18% (11-28%)	82% (72-89%)	0.41 (0.2-0.73)	-0.0014% (-0.75-1.1%)	-0.047% (-0.96-1.2%)	-1.7 (-33-43)	-0.27 (-160-230)
North Korea	2005	Female	95 plus	21% (11-34%)	79% (66-89%)	0.47 (0.19-0.94)	-0.054% (-1-1.4%)	-0.14% (-1.3-1.5%)	-1.6 (-14-17)	-2.3 (-45-60)
North Korea	2005	Male	All Ages	39% (33-47%)	61% (53-67%)	1.4 (1.3-1.6)	5.4% (4-6.7%)	6.9% (4.8-9.1%)	6200 (4200-8200)	240000 (180000-300000)
North Korea	2005	Male	15 to 19	30% (19-45%)	70% (55-81%)	0.44 (0.21-0.81)	2.5% (1.4-4%)	3% (1.4-5.4%)	40 (19-72)	3700 (2100-5900)
North Korea	2005	Male	20 to 24	42% (28-59%)	58% (41-72%)	1.1 (0.57-1.7)	5.5% (3.4-8.3%)	6.2% (3.4-10%)	120 (66-200)	10000 (6200-16000)
North Korea	2005	Male	25 to 29	48% (31-64%)	52% (36-69%)	1.6 (0.88-2.5)	7.2% (4.4-10%)	8.6% (4.7-13%)	160 (81-240)	12000 (6900-17000)
North Korea	2005	Male	30 to 34	51% (33-68%)	49% (32-67%)	2 (1.2-2.9)	7.8% (5.1-11%)	9.3% (5.7-14%)	290 (170-440)	20000 (13000-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	2005	Male	35 to 39	54% (37-71%)	46% (29-63%)	2.4 (1.5-3.5)	8.6% (5.7-12%)	10% (6.5-15%)	420 (240-620)	25000 (16000-36000)
North Korea	2005	Male	40 to 44	55% (39-73%)	45% (27-61%)	2.8 (1.8-4)	9.4% (6.5-13%)	11% (7.5-16%)	430 (270-600)	23000 (15000-31000)
North Korea	2005	Male	45 to 49	56% (37-72%)	44% (28-63%)	3.1 (1.8-4.5)	10% (6.4-14%)	12% (7.4-17%)	680 (420-990)	31000 (20000-45000)
North Korea	2005	Male	50 to 54	55% (37-72%)	45% (28-63%)	3.2 (1.9-4.7)	10% (5.9-14%)	12% (6.8-17%)	490 (280-700)	19000 (12000-27000)
North Korea	2005	Male	55 to 59	53% (36-71%)	47% (29-64%)	3 (1.7-4.5)	9.2% (5-14%)	11% (5.4-16%)	850 (440-1300)	29000 (16000-43000)
North Korea	2005	Male	60 to 64	51% (33-70%)	49% (30-67%)	2.6 (1.5-4.3)	7.9% (3.8-13%)	8.9% (4.2-14%)	1000 (470-1700)	29000 (14000-47000)
North Korea	2005	Male	65 to 69	48% (32-66%)	52% (34-68%)	2.2 (1.3-3.3)	6.5% (3.2-10%)	7.3% (3.4-12%)	880 (420-1400)	21000 (10000-34000)
North Korea	2005	Male	70 to 74	45% (29-62%)	55% (38-71%)	1.8 (1-3)	5% (2-8.9%)	5.5% (2.2-10%)	460 (180-830)	8700 (3500-16000)
North Korea	2005	Male	75 to 79	41% (27-58%)	59% (42-73%)	1.6 (0.8-2.7)	3.9% (1.2-7.6%)	4.2% (1.3-8.4%)	220 (67-440)	3200 (1000-6400)
North Korea	2005	Male	80 to 84	39% (24-56%)	61% (44-76%)	1.4 (0.73-2.4)	3.3% (1.1-6.7%)	3.5% (1.1-7.4%)	87 (26-180)	970 (320-2000)
North Korea	2005	Male	85 to 89	38% (24-53%)	62% (47-76%)	1.3 (0.7-2.2)	2.6% (0.69-5.4%)	2.7% (0.61-5.8%)	43 (9.4-93)	350 (94-730)
North Korea	2005	Male	90 to 94	39% (26-54%)	61% (46-74%)	1.3 (0.68-2.2)	2.1% (0.26-5.2%)	2.1% (0.08-5.6%)	17 (0.66-46)	98 (12-250)
North Korea	2005	Male	95 plus	43% (26-61%)	57% (39-74%)	1.3 (0.59-2.5)	1.9% (-0.19-5.4%)	1.8% (-0.54-5.7%)	3.8 (-1.1-12)	15 (-1.5-44)
North Korea	2010	Female	All Ages	17% (14-21%)	83% (79-86%)	0.24 (0.19-0.29)	0.6% (0.22-1.1%)	0.63% (-0.094-1.5%)	700 (-100-1700)	26000 (8900-46000)
North Korea	2010	Female	15 to 19	16% (10-26%)	84% (74-90%)	0.2 (0.1-0.36)	0.82% (0.49-1.3%)	1.2% (0.49-2.1%)	7.4 (3-14)	960 (540-1500)
North Korea	2010	Female	20 to 24	20% (12-30%)	80% (70-88%)	0.22 (0.1-0.4)	1.1% (0.72-1.7%)	1% (0.33-2.1%)	7.7 (2.3-15)	1500 (880-2300)
North Korea	2010	Female	25 to 29	20% (12-32%)	80% (68-88%)	0.23 (0.11-0.4)	1.1% (0.71-1.6%)	0.99% (0.35-1.9%)	8.8 (3.1-17)	1500 (940-2200)
North Korea	2010	Female	30 to 34	21% (12-33%)	79% (67-88%)	0.26 (0.13-0.43)	1% (0.62-1.5%)	1% (0.43-1.9%)	9.9 (3.9-18)	1300 (780-1900)
North Korea	2010	Female	35 to 39	23% (14-35%)	77% (65-86%)	0.32 (0.17-0.52)	0.98% (0.61-1.5%)	1.3% (0.56-2.3%)	26 (11-45)	2200 (1300-3300)
North Korea	2010	Female	40 to 44	23% (14-37%)	77% (63-86%)	0.36 (0.2-0.59)	0.98% (0.55-1.5%)	1.4% (0.64-2.5%)	43 (19-75)	2700 (1500-4200)
North Korea	2010	Female	45 to 49	23% (12-36%)	77% (64-88%)	0.39 (0.2-0.65)	0.98% (0.44-1.7%)	1.5% (0.59-2.7%)	49 (18-93)	2400 (1100-4100)
North Korea	2010	Female	50 to 54	22% (14-35%)	78% (65-86%)	0.41 (0.22-0.67)	1% (0.35-1.9%)	1.5% (0.51-2.8%)	80 (27-150)	3100 (1100-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	2010	Female	55 to 59	21% (13-33%)	79% (67-87%)	0.41 (0.22-0.68)	0.95% (0.18-2%)	1.4% (0.28-2.9%)	57 (12-120)	1800 (370-3900)
North Korea	2010	Female	60 to 64	20% (11-32%)	80% (68-89%)	0.4 (0.2-0.7)	0.81% (0.024-1.9%)	1.1% (0.083-2.6%)	96 (7.1-210)	2600 (78-6000)
North Korea	2010	Female	65 to 69	19% (12-29%)	81% (71-88%)	0.39 (0.21-0.66)	0.7% (-0.075-1.6%)	0.97% (-0.055-2.2%)	120 (-7.5-270)	2600 (-280-6100)
North Korea	2010	Female	70 to 74	18% (12-29%)	82% (71-88%)	0.38 (0.2-0.69)	0.56% (-0.22-1.7%)	0.74% (-0.24-2.1%)	120 (-40-350)	2100 (-840-6300)
North Korea	2010	Female	75 to 79	17% (10-27%)	83% (73-90%)	0.37 (0.18-0.66)	0.33% (-0.42-1.4%)	0.42% (-0.5-1.7%)	67 (-81-280)	910 (-1200-3900)
North Korea	2010	Female	80 to 84	17% (10-26%)	83% (74-90%)	0.36 (0.18-0.61)	0.11% (-0.63-1.1%)	0.13% (-0.75-1.3%)	17 (-98-170)	180 (-1000-1700)
North Korea	2010	Female	85 to 89	17% (10-26%)	83% (74-90%)	0.37 (0.19-0.64)	0.0026% (-0.74-0.98%)	-0.018% (-0.88-1.1%)	-1.7 (-79-98)	1.9 (-560-740)
North Korea	2010	Female	90 to 94	19% (11-28%)	81% (72-89%)	0.4 (0.19-0.72)	-0.087% (-0.86-1%)	-0.15% (-1.1-1%)	-6 (-42-45)	-21 (-200-230)
North Korea	2010	Female	95 plus	21% (12-35%)	79% (65-88%)	0.45 (0.18-0.9)	-0.14% (-1.2-1.3%)	-0.24% (-1.4-1.4%)	-3.2 (-19-19)	-7.1 (-59-65)
North Korea	2010	Male	All Ages	40% (34-47%)	60% (53-66%)	1.4 (1.2-1.7)	5.4% (4-6.9%)	6.6% (4.4-9%)	6500 (4300-9100)	250000 (180000-320000)
North Korea	2010	Male	15 to 19	30% (19-45%)	70% (55-81%)	0.42 (0.2-0.77)	2.4% (1.4-3.9%)	3% (1.4-5.3%)	42 (20-73)	3900 (2200-6100)
North Korea	2010	Male	20 to 24	43% (28-58%)	57% (42-72%)	1 (0.55-1.7)	5.4% (3.3-8.1%)	6.1% (3.2-10%)	120 (65-210)	10000 (6100-16000)
North Korea	2010	Male	25 to 29	48% (31-64%)	52% (36-69%)	1.5 (0.82-2.4)	6.9% (4.2-10%)	8.1% (4.3-13%)	180 (91-280)	13000 (7900-20000)
North Korea	2010	Male	30 to 34	51% (33-68%)	49% (32-67%)	1.9 (1.2-2.9)	7.7% (5-11%)	9.2% (5.4-14%)	190 (110-300)	13000 (8400-19000)
North Korea	2010	Male	35 to 39	53% (37-70%)	47% (30-63%)	2.3 (1.4-3.3)	8.2% (5.3-12%)	9.9% (5.9-15%)	390 (220-600)	24000 (15000-35000)
North Korea	2010	Male	40 to 44	55% (38-72%)	45% (28-62%)	2.6 (1.7-3.8)	8.8% (5.8-12%)	11% (6.6-15%)	550 (330-820)	29000 (19000-42000)
North Korea	2010	Male	45 to 49	56% (37-72%)	44% (28-63%)	2.9 (1.8-4.4)	9.4% (6-13%)	11% (6.8-16%)	590 (350-880)	27000 (17000-40000)
North Korea	2010	Male	50 to 54	56% (38-72%)	44% (28-62%)	3 (1.9-4.3)	9.4% (5.5-13%)	11% (6.1-16%)	900 (500-1300)	36000 (21000-52000)
North Korea	2010	Male	55 to 59	54% (36-72%)	46% (28-64%)	2.8 (1.7-4.4)	8.7% (4.9-13%)	9.9% (5.4-15%)	580 (310-880)	20000 (11000-30000)
North Korea	2010	Male	60 to 64	51% (33-70%)	49% (30-67%)	2.5 (1.4-4.1)	7.5% (3.6-12%)	8.4% (3.9-14%)	920 (430-1500)	27000 (13000-44000)
North Korea	2010	Male	65 to 69	48% (32-66%)	52% (34-68%)	2.1 (1.2-3.3)	6.1% (2.9-10%)	6.8% (3.2-12%)	930 (430-1600)	22000 (11000-37000)
North Korea	2010	Male	70 to 74	45% (29-62%)	55% (38-71%)	1.7 (0.95-2.8)	4.7% (1.8-8.8%)	5.2% (1.9-9.8%)	690 (250-1300)	13000 (4900-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	2010	Male	75 to 79	41% (27-59%)	59% (41-73%)	1.5 (0.74-2.5)	3.6% (1-7.5%)	3.9% (1.1-8.2%)	300 (80-620)	4400 (1300-9000)
North Korea	2010	Male	80 to 84	39% (24-56%)	61% (44-76%)	1.3 (0.7-2.3)	3% (0.91-6.3%)	3.2% (0.85-7%)	100 (28-230)	1200 (350-2500)
North Korea	2010	Male	85 to 89	38% (24-53%)	62% (47-76%)	1.3 (0.65-2.1)	2.4% (0.51-5.1%)	2.5% (0.39-5.6%)	43 (6.7-100)	340 (74-770)
North Korea	2010	Male	90 to 94	40% (26-55%)	60% (45-74%)	1.2 (0.65-2.1)	1.8% (0.11-4.8%)	1.9% (-0.069-5.2%)	14 (-0.51-38)	78 (4.8-200)
North Korea	2010	Male	95 plus	43% (26-62%)	57% (38-74%)	1.3 (0.56-2.4)	1.7% (-0.34-5.2%)	1.5% (-0.69-5.5%)	3.7 (-1.7-13)	15 (-3.2-47)
North Korea	2016	Female	All Ages	18% (15-21%)	82% (79-85%)	0.26 (0.17-0.36)	0.68% (0.19-1.3%)	0.71% (-0.13-1.8%)	870 (-150-2300)	30000 (8000-57000)
North Korea	2016	Female	15 to 19	17% (11-26%)	83% (74-89%)	0.21 (0.096-0.4)	0.87% (0.52-1.3%)	1.2% (0.5-2.3%)	6.9 (2.8-13)	920 (540-1400)
North Korea	2016	Female	20 to 24	20% (12-30%)	80% (70-88%)	0.23 (0.098-0.44)	1.2% (0.74-1.8%)	1.1% (0.36-2.1%)	7.9 (2.5-16)	1500 (880-2300)
North Korea	2016	Female	25 to 29	20% (12-32%)	80% (68-88%)	0.23 (0.11-0.43)	1.1% (0.72-1.6%)	0.99% (0.33-2%)	8.6 (2.8-18)	1500 (910-2400)
North Korea	2016	Female	30 to 34	21% (13-33%)	79% (67-87%)	0.26 (0.13-0.49)	1% (0.65-1.5%)	1% (0.41-2%)	12 (4.6-23)	1600 (940-2300)
North Korea	2016	Female	35 to 39	23% (13-36%)	77% (64-87%)	0.33 (0.16-0.59)	1% (0.62-1.6%)	1.3% (0.59-2.4%)	17 (7.1-31)	1500 (840-2300)
North Korea	2016	Female	40 to 44	23% (14-37%)	77% (63-86%)	0.38 (0.19-0.67)	1% (0.56-1.8%)	1.5% (0.69-2.8%)	40 (18-79)	2500 (1300-4400)
North Korea	2016	Female	45 to 49	23% (13-37%)	77% (63-87%)	0.4 (0.2-0.72)	1% (0.46-2%)	1.6% (0.62-3.1%)	71 (27-140)	3500 (1500-6600)
North Korea	2016	Female	50 to 54	22% (14-35%)	78% (65-86%)	0.42 (0.21-0.76)	1.1% (0.39-2.1%)	1.6% (0.56-3.2%)	81 (29-160)	3200 (1200-6000)
North Korea	2016	Female	55 to 59	22% (13-33%)	78% (67-87%)	0.42 (0.21-0.72)	1% (0.2-2.2%)	1.5% (0.31-3.3%)	120 (24-250)	3800 (720-8200)
North Korea	2016	Female	60 to 64	20% (12-32%)	80% (68-88%)	0.41 (0.21-0.72)	0.9% (0.06-2.2%)	1.3% (0.14-3%)	84 (8.7-190)	2200 (150-5300)
North Korea	2016	Female	65 to 69	19% (12-30%)	81% (70-88%)	0.4 (0.21-0.7)	0.82% (-0.074-2%)	1.1% (-0.035-2.7%)	120 (-3.6-290)	2700 (-240-6500)
North Korea	2016	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.39 (0.19-0.73)	0.67% (-0.24-2%)	0.88% (-0.23-2.5%)	150 (-40-420)	2600 (-890-7600)
North Korea	2016	Female	75 to 79	17% (11-27%)	83% (73-89%)	0.38 (0.18-0.7)	0.43% (-0.38-1.7%)	0.53% (-0.44-2.1%)	110 (-90-440)	1500 (-1300-6300)
North Korea	2016	Female	80 to 84	17% (10-27%)	83% (73-90%)	0.37 (0.17-0.66)	0.2% (-0.63-1.4%)	0.23% (-0.75-1.7%)	39 (-120-280)	410 (-1300-3000)
North Korea	2016	Female	85 to 89	18% (10-27%)	82% (73-90%)	0.39 (0.18-0.71)	0.069% (-0.75-1.3%)	0.058% (-0.9-1.5%)	6.7 (-100-170)	69 (-740-1300)
North Korea	2016	Female	90 to 94	19% (12-29%)	81% (71-88%)	0.42 (0.18-0.79)	-0.027% (-0.85-1.3%)	-0.079% (-1.1-1.5%)	-4.2 (-55-81)	-8.1 (-270-410)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
North Korea	2016	Female	95 plus	22% (12-35%)	78% (65-88%)	0.48 (0.18-0.99)	-0.068% (-1.2-1.7%)	-0.16% (-1.4-1.9%)	-2.5 (-22-30)	-4 (-70-110)
North Korea	2016	Male	All Ages	41% (35-49%)	59% (51-65%)	1.5 (1.1-2.1)	5.7% (3.7-7.9%)	6.8% (3.7-10%)	7500 (4200-11000)	270000 (170000-380000)
North Korea	2016	Male	15 to 19	30% (19-45%)	70% (55-81%)	0.44 (0.19-0.84)	2.5% (1.3-4.1%)	3% (1.3-5.6%)	42 (18-78)	3900 (2100-6400)
North Korea	2016	Male	20 to 24	43% (28-59%)	57% (41-72%)	1.1 (0.51-1.8)	5.4% (3.2-8.6%)	6.2% (3.1-11%)	140 (64-230)	11000 (6300-18000)
North Korea	2016	Male	25 to 29	49% (31-65%)	51% (35-69%)	1.6 (0.77-2.7)	6.9% (4.1-11%)	8.2% (4.3-13%)	180 (89-300)	14000 (8000-22000)
North Korea	2016	Male	30 to 34	52% (34-69%)	48% (31-66%)	2 (1-3.2)	7.6% (4.7-11%)	9.1% (5.1-14%)	230 (130-380)	16000 (9600-24000)
North Korea	2016	Male	35 to 39	54% (38-71%)	46% (29-62%)	2.3 (1.2-3.9)	8.5% (5-13%)	10% (5.5-16%)	260 (140-440)	16000 (9400-25000)
North Korea	2016	Male	40 to 44	55% (39-73%)	45% (27-61%)	2.7 (1.5-4.3)	9% (5.5-13%)	11% (6.2-17%)	520 (290-850)	28000 (16000-44000)
North Korea	2016	Male	45 to 49	56% (37-73%)	44% (27-63%)	3 (1.6-4.7)	9.3% (5.4-14%)	11% (6-17%)	830 (440-1300)	39000 (22000-59000)
North Korea	2016	Male	50 to 54	56% (38-73%)	44% (27-62%)	3.1 (1.8-4.8)	9.4% (5.1-14%)	11% (5.7-17%)	870 (440-1300)	35000 (19000-52000)
North Korea	2016	Male	55 to 59	54% (36-72%)	46% (28-64%)	2.9 (1.6-4.8)	8.7% (4.6-13%)	10% (5.1-16%)	1200 (580-1800)	40000 (21000-62000)
North Korea	2016	Male	60 to 64	52% (34-70%)	48% (30-66%)	2.6 (1.4-4.3)	7.5% (3.2-12%)	8.4% (3.5-14%)	750 (310-1300)	22000 (9400-36000)
North Korea	2016	Male	65 to 69	48% (33-66%)	52% (34-67%)	2.2 (1.2-3.5)	6.3% (2.8-11%)	7.1% (3-12%)	870 (360-1500)	21000 (9000-35000)
North Korea	2016	Male	70 to 74	45% (29-62%)	55% (38-71%)	1.8 (0.91-3.2)	4.9% (1.6-9.7%)	5.4% (1.6-11%)	830 (240-1700)	16000 (4900-32000)
North Korea	2016	Male	75 to 79	42% (27-60%)	58% (40-73%)	1.5 (0.72-2.7)	3.9% (0.95-7.9%)	4.2% (0.94-8.7%)	540 (120-1100)	8000 (2000-16000)
North Korea	2016	Male	80 to 84	40% (25-57%)	60% (43-75%)	1.4 (0.68-2.4)	3.2% (0.64-7%)	3.4% (0.58-7.7%)	190 (32-420)	2100 (420-4500)
North Korea	2016	Male	85 to 89	39% (25-54%)	61% (46-75%)	1.3 (0.63-2.3)	2.5% (0.33-6%)	2.6% (0.21-6.6%)	64 (5.2-160)	510 (69-1200)
North Korea	2016	Male	90 to 94	40% (26-56%)	60% (44-74%)	1.3 (0.62-2.4)	1.9% (-0.064-5.6%)	2% (-0.27-6.2%)	17 (-2.3-53)	96 (-3.2-280)
North Korea	2016	Male	95 plus	43% (26-62%)	57% (38-74%)	1.3 (0.54-2.6)	1.7% (-0.42-5.8%)	1.6% (-0.77-6.3%)	3.5 (-1.7-13)	14 (-3.4-47)
Taiwan	1990	Female	All Ages	16% (13-20%)	84% (80-87%)	0.3 (0.17-0.5)	1.2% (0.55-2.2%)	1.6% (0.31-3.6%)	630 (120-1500)	23000 (11000-43000)
Taiwan	1990	Female	15 to 19	15% (9.4-24%)	85% (76-91%)	0.28 (0.12-0.58)	0.97% (0.52-1.6%)	1.6% (0.56-3.4%)	7.5 (2.5-16)	930 (490-1600)
Taiwan	1990	Female	20 to 24	19% (11-29%)	81% (71-89%)	0.3 (0.12-0.61)	1.4% (0.87-2.1%)	1.8% (0.69-3.5%)	9.8 (3.9-19)	1600 (950-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	1990	Female	25 to 29	19% (11-30%)	81% (70-89%)	0.31 (0.13-0.63)	1.4% (0.91-2.1%)	1.9% (0.83-3.5%)	13 (5.6-24)	1800 (1100-2800)
Taiwan	1990	Female	30 to 34	20% (12-32%)	80% (68-88%)	0.36 (0.16-0.71)	1.4% (0.94-2.2%)	2.3% (1.2-4.2%)	18 (9.7-33)	1900 (1200-3000)
Taiwan	1990	Female	35 to 39	22% (13-34%)	78% (66-87%)	0.45 (0.19-0.86)	1.6% (1.1-2.6%)	3.2% (1.9-5.3%)	30 (18-50)	2200 (1500-3400)
Taiwan	1990	Female	40 to 44	22% (13-35%)	78% (65-87%)	0.51 (0.22-0.99)	1.8% (1.1-2.9%)	3.6% (2.1-6%)	32 (19-54)	1800 (1200-3000)
Taiwan	1990	Female	45 to 49	22% (12-34%)	78% (66-88%)	0.55 (0.22-1.1)	1.8% (0.92-3.2%)	3.5% (1.9-6.3%)	41 (22-76)	1900 (1000-3500)
Taiwan	1990	Female	50 to 54	21% (13-33%)	79% (67-87%)	0.57 (0.24-1.2)	1.8% (0.77-3.5%)	3.3% (1.6-6.2%)	57 (28-110)	2200 (980-4300)
Taiwan	1990	Female	55 to 59	20% (12-31%)	80% (69-88%)	0.57 (0.23-1.1)	1.6% (0.5-3.5%)	2.8% (1.2-5.6%)	66 (29-130)	2100 (670-4500)
Taiwan	1990	Female	60 to 64	19% (11-30%)	81% (70-89%)	0.55 (0.21-1.1)	1.4% (0.18-3.3%)	2.2% (0.61-4.9%)	72 (20-160)	1900 (260-4600)
Taiwan	1990	Female	65 to 69	18% (11-28%)	82% (72-89%)	0.54 (0.23-1)	1.2% (-0.075-3.3%)	1.8% (0.13-4.5%)	77 (5.5-190)	1700 (-99-4600)
Taiwan	1990	Female	70 to 74	17% (11-27%)	83% (73-89%)	0.52 (0.21-1)	1% (-0.29-3.3%)	1.4% (-0.23-4.2%)	70 (-11-210)	1300 (-350-4100)
Taiwan	1990	Female	75 to 79	16% (9.6-26%)	84% (74-90%)	0.5 (0.2-1)	0.83% (-0.57-3.2%)	1% (-0.58-3.7%)	66 (-38-240)	940 (-590-3500)
Taiwan	1990	Female	80 to 84	15% (8.8-25%)	85% (75-91%)	0.48 (0.19-0.92)	0.67% (-0.59-2.8%)	0.8% (-0.66-3.3%)	42 (-35-170)	450 (-380-1900)
Taiwan	1990	Female	85 to 89	16% (9.1-24%)	84% (76-91%)	0.49 (0.19-0.98)	0.58% (-0.61-2.6%)	0.66% (-0.72-3%)	21 (-23-95)	160 (-180-730)
Taiwan	1990	Female	90 to 94	17% (10-26%)	83% (74-90%)	0.52 (0.21-1.1)	0.49% (-0.68-2.5%)	0.52% (-0.82-2.8%)	6.8 (-11-37)	39 (-53-200)
Taiwan	1990	Female	95 plus	19% (10-32%)	81% (68-90%)	0.6 (0.19-1.4)	0.55% (-0.81-3.3%)	0.54% (-1-3.6%)	2 (-3.7-13)	7.9 (-12-47)
Taiwan	1990	Male	All Ages	39% (32-46%)	61% (54-68%)	2 (1.1-3.2)	8.6% (5.4-12%)	10% (5.6-15%)	6800 (3700-9600)	240000 (150000-320000)
Taiwan	1990	Male	15 to 19	29% (18-43%)	71% (57-82%)	0.61 (0.25-1.3)	3.1% (1.5-5.5%)	4.1% (1.6-7.8%)	53 (21-100)	4500 (2200-8200)
Taiwan	1990	Male	20 to 24	41% (27-57%)	59% (43-73%)	1.5 (0.67-2.8)	6.7% (3.7-11%)	8.7% (4-16%)	130 (59-230)	11000 (5700-18000)
Taiwan	1990	Male	25 to 29	47% (31-64%)	53% (36-69%)	2.2 (0.99-4)	8.7% (4.6-14%)	12% (5.4-20%)	210 (97-350)	16000 (8100-25000)
Taiwan	1990	Male	30 to 34	50% (33-67%)	50% (33-67%)	2.8 (1.3-5)	10% (5.7-15%)	14% (7.2-22%)	270 (140-420)	18000 (9900-27000)
Taiwan	1990	Male	35 to 39	53% (36-69%)	47% (31-64%)	3.3 (1.5-6.2)	11% (6.9-16%)	16% (9-23%)	360 (200-530)	21000 (13000-30000)
Taiwan	1990	Male	40 to 44	54% (37-72%)	46% (28-63%)	3.9 (1.8-7)	13% (8.2-18%)	17% (10-24%)	380 (230-530)	20000 (12000-27000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	1990	Male	45 to 49	55% (36-72%)	45% (28-64%)	4.3 (2-8.3)	14% (8.8-19%)	18% (11-24%)	480 (300-640)	22000 (14000-29000)
Taiwan	1990	Male	50 to 54	54% (36-72%)	46% (28-64%)	4.4 (2-8.3)	14% (8.2-18%)	17% (10-23%)	600 (360-810)	24000 (15000-32000)
Taiwan	1990	Male	55 to 59	53% (35-70%)	47% (30-65%)	4.1 (2-7.7)	12% (7.1-17%)	15% (8.5-20%)	740 (430-1000)	26000 (15000-35000)
Taiwan	1990	Male	60 to 64	50% (33-68%)	50% (32-67%)	3.7 (1.7-6.9)	10% (5.3-15%)	12% (6.3-18%)	930 (470-1300)	27000 (14000-40000)
Taiwan	1990	Male	65 to 69	47% (31-64%)	53% (36-69%)	3.1 (1.4-5.7)	9.1% (4-14%)	11% (4.6-16%)	890 (390-1400)	22000 (9500-33000)
Taiwan	1990	Male	70 to 74	43% (28-60%)	57% (40-72%)	2.5 (1.1-4.9)	7.4% (2.7-12%)	8.5% (3-14%)	680 (240-1100)	13000 (4900-22000)
Taiwan	1990	Male	75 to 79	40% (25-58%)	60% (42-75%)	2.2 (0.91-4.2)	6.3% (1.8-11%)	7% (2-13%)	580 (160-1000)	8500 (2500-15000)
Taiwan	1990	Male	80 to 84	38% (23-54%)	62% (46-77%)	2 (0.79-3.7)	5.7% (1.4-11%)	6.3% (1.4-12%)	310 (71-590)	3400 (850-6500)
Taiwan	1990	Male	85 to 89	37% (24-52%)	63% (48-76%)	1.8 (0.75-3.6)	4.8% (1.1-9.8%)	5.3% (1.1-11%)	130 (25-260)	990 (220-2000)
Taiwan	1990	Male	90 to 94	39% (25-54%)	61% (46-75%)	1.8 (0.76-3.6)	4% (0.52-8.5%)	4.3% (0.45-9.4%)	32 (3.4-70)	170 (22-370)
Taiwan	1990	Male	95 plus	42% (26-61%)	58% (39-74%)	1.9 (0.65-4)	3.6% (0.072-8.6%)	3.7% (-0.19-9.3%)	6.5 (-0.33-16)	24 (0.46-57)
Taiwan	1995	Female	All Ages	16% (13-20%)	84% (80-87%)	0.34 (0.19-0.55)	1.4% (0.71-2.5%)	2% (0.58-4.1%)	880 (260-1800)	29000 (15000-50000)
Taiwan	1995	Female	15 to 19	15% (9.4-24%)	85% (76-91%)	0.3 (0.12-0.61)	1.8% (0.57-1.8%)	1.8% (0.63-3.7%)	8.2 (2.9-17)	1000 (540-1800)
Taiwan	1995	Female	20 to 24	19% (11-28%)	81% (72-89%)	0.32 (0.13-0.69)	1.5% (0.92-2.3%)	1.9% (0.75-3.7%)	9.2 (3.7-18)	1500 (890-2400)
Taiwan	1995	Female	25 to 29	19% (11-29%)	81% (71-89%)	0.33 (0.14-0.69)	1.5% (1-2.3%)	2% (0.93-3.6%)	12 (5.3-21)	1700 (1100-2600)
Taiwan	1995	Female	30 to 34	20% (12-31%)	80% (69-88%)	0.38 (0.15-0.77)	1.5% (1-2.3%)	2.5% (1.4-4.3%)	19 (11-34)	2000 (1300-3100)
Taiwan	1995	Female	35 to 39	22% (13-34%)	78% (66-87%)	0.47 (0.2-0.9)	1.7% (1.2-2.6%)	3.4% (2.1-5.5%)	34 (21-55)	2500 (1700-3800)
Taiwan	1995	Female	40 to 44	22% (13-35%)	78% (65-87%)	0.54 (0.24-1)	1.9% (1.1-3%)	3.8% (2.3-6.3%)	49 (29-81)	2900 (1800-4600)
Taiwan	1995	Female	45 to 49	21% (12-34%)	79% (66-88%)	0.58 (0.23-1.2)	1.9% (1-3.3%)	3.8% (2.2-6.6%)	49 (27-85)	2300 (1300-4000)
Taiwan	1995	Female	50 to 54	21% (12-33%)	79% (67-88%)	0.6 (0.25-1.2)	2% (0.95-3.7%)	3.8% (2.1-6.7%)	62 (33-110)	2400 (1200-4400)
Taiwan	1995	Female	55 to 59	20% (12-31%)	80% (69-88%)	0.6 (0.25-1.2)	2% (0.81-3.9%)	3.5% (1.8-6.4%)	85 (42-160)	2700 (1200-5300)
Taiwan	1995	Female	60 to 64	19% (11-30%)	81% (70-89%)	0.58 (0.23-1.2)	1.8% (0.56-3.8%)	3% (1.2-5.6%)	100 (42-190)	2700 (860-5600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	1995	Female	65 to 69	18% (11-29%)	82% (71-89%)	0.57 (0.24-1.1)	1.7% (0.33-3.8%)	2.6% (0.8-5.3%)	120 (37-240)	2700 (500-5900)
Taiwan	1995	Female	70 to 74	17% (11-27%)	83% (73-89%)	0.55 (0.22-1.1)	1.5% (0.077-3.8%)	2% (0.27-4.9%)	120 (15-290)	2200 (120-5500)
Taiwan	1995	Female	75 to 79	16% (9.6-26%)	84% (74-90%)	0.53 (0.2-1)	1.1% (-0.22-3.5%)	1.5% (-0.16-4.3%)	91 (-11-270)	1300 (-250-4000)
Taiwan	1995	Female	80 to 84	15% (9-25%)	85% (75-91%)	0.51 (0.2-1)	0.88% (-0.39-3%)	1.1% (-0.42-3.6%)	69 (-27-230)	740 (-330-2500)
Taiwan	1995	Female	85 to 89	16% (9.5-24%)	84% (76-91%)	0.53 (0.21-1.1)	0.77% (-0.49-2.8%)	0.89% (-0.53-3.3%)	39 (-23-140)	300 (-180-1100)
Taiwan	1995	Female	90 to 94	17% (10-26%)	83% (74-90%)	0.57 (0.22-1.2)	0.65% (-0.56-2.6%)	0.71% (-0.68-3%)	13 (-12-55)	74 (-63-300)
Taiwan	1995	Female	95 plus	20% (11-33%)	80% (67-89%)	0.65 (0.21-1.5)	0.73% (-0.79-3.5%)	0.74% (-0.99-3.9%)	4.1 (-5.4-21)	16 (-17-75)
Taiwan	1995	Male	All Ages	39% (33-47%)	61% (53-67%)	2.2 (1.2-3.5)	9.1% (5.8-12%)	11% (5.7-14%)	7800 (4400-11000)	270000 (170000-360000)
Taiwan	1995	Male	15 to 19	28% (18-43%)	72% (57-82%)	0.65 (0.25-1.4)	3.2% (1.6-5.8%)	4.3% (1.7-8.5%)	56 (22-110)	4800 (2400-8700)
Taiwan	1995	Male	20 to 24	41% (27-57%)	59% (43-73%)	1.5 (0.71-3)	7.2% (3.9-12%)	9.3% (4.3-17%)	120 (56-220)	10000 (5500-17000)
Taiwan	1995	Male	25 to 29	47% (30-64%)	53% (36-70%)	2.3 (1-4.4)	9.3% (5-14%)	13% (5.8-21%)	200 (90-320)	15000 (7800-23000)
Taiwan	1995	Male	30 to 34	50% (32-68%)	50% (32-68%)	3 (1.3-5.4)	11% (5.9-16%)	15% (7.3-22%)	300 (150-460)	20000 (11000-30000)
Taiwan	1995	Male	35 to 39	53% (36-70%)	47% (30-64%)	3.6 (1.6-6.5)	12% (7.4-17%)	16% (9.5-24%)	430 (250-620)	25000 (15000-36000)
Taiwan	1995	Male	40 to 44	54% (37-72%)	46% (28-63%)	4.1 (1.9-7.4)	14% (8.8-19%)	18% (11-25%)	590 (360-800)	31000 (19000-41000)
Taiwan	1995	Male	45 to 49	54% (36-72%)	46% (28-64%)	4.6 (2.1-8.6)	15% (9.4-19%)	19% (12-25%)	580 (360-760)	26000 (17000-35000)
Taiwan	1995	Male	50 to 54	54% (36-72%)	46% (28-64%)	4.7 (2.1-8.9)	14% (9.1-19%)	18% (11-23%)	630 (390-830)	25000 (16000-33000)
Taiwan	1995	Male	55 to 59	52% (34-70%)	48% (30-66%)	4.4 (2.1-7.8)	13% (7.9-17%)	16% (9.4-21%)	740 (450-970)	26000 (15000-34000)
Taiwan	1995	Male	60 to 64	50% (32-68%)	50% (32-68%)	3.9 (1.8-7.2)	11% (6-15%)	13% (7.1-18%)	870 (470-1200)	26000 (14000-36000)
Taiwan	1995	Male	65 to 69	47% (30-65%)	53% (35-70%)	3.3 (1.5-6.1)	9.3% (4.4-14%)	11% (5-16%)	1100 (490-1600)	26000 (12000-39000)
Taiwan	1995	Male	70 to 74	43% (27-60%)	57% (40-73%)	2.7 (1.1-5.1)	7.6% (2.9-12%)	8.7% (3.3-14%)	910 (350-1500)	18000 (6900-28000)
Taiwan	1995	Male	75 to 79	40% (25-57%)	60% (43-75%)	2.3 (0.94-4.4)	6.2% (2-11%)	7% (2.3-12%)	640 (200-1100)	9700 (3100-16000)
Taiwan	1995	Male	80 to 84	37% (23-54%)	63% (46-77%)	2.1 (0.86-4)	5.7% (1.6-10%)	6.3% (1.7-11%)	470 (120-830)	5100 (1400-9200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	1995	Male	85 to 89	37% (23-51%)	63% (49-77%)	1.9 (0.82-3.8)	4.9% (1.2-9.5%)	5.4% (1.2-11%)	190 (43-370)	1500 (360-2900)
Taiwan	1995	Male	90 to 94	38% (25-54%)	62% (46-75%)	1.9 (0.8-3.7)	4.2% (0.67-8.6%)	4.5% (0.57-9.5%)	52 (6.4-110)	280 (46-570)
Taiwan	1995	Male	95 plus	42% (25-60%)	58% (40-75%)	2 (0.66-4.2)	3.8% (0.078-8.7%)	3.9% (-0.14-9.5%)	10 (-0.37-25)	38 (0.81-89)
Taiwan	2000	Female	All Ages	17% (14-20%)	83% (80-86%)	0.38 (0.21-0.62)	1.6% (0.81-2.7%)	2.2% (0.84-4.4%)	1100 (400-2100)	34000 (17000-58000)
Taiwan	2000	Female	15 to 19	15% (9.5-24%)	85% (76-91%)	0.31 (0.13-0.65)	1% (0.59-1.7%)	1.8% (0.67-3.7%)	6.7 (2.5-14)	910 (490-1500)
Taiwan	2000	Female	20 to 24	19% (11-29%)	81% (71-89%)	0.34 (0.14-0.71)	1.5% (0.96-2.3%)	2% (0.82-3.9%)	8.8 (3.7-18)	1600 (980-2500)
Taiwan	2000	Female	25 to 29	19% (11-30%)	81% (70-89%)	0.35 (0.14-0.72)	1.6% (1-2.3%)	2.2% (0.99-4.1%)	9.9 (4.5-18)	1600 (1000-2400)
Taiwan	2000	Female	30 to 34	20% (12-32%)	80% (68-88%)	0.4 (0.16-0.82)	1.6% (1-2.3%)	2.6% (1.4-4.5%)	17 (9.3-30)	1900 (1200-3000)
Taiwan	2000	Female	35 to 39	21% (13-34%)	79% (66-87%)	0.5 (0.21-0.98)	1.7% (1.2-2.7%)	3.5% (2.2-5.8%)	34 (21-55)	2600 (1800-3900)
Taiwan	2000	Female	40 to 44	22% (13-35%)	78% (65-87%)	0.57 (0.25-1.1)	1.9% (1.1-3.1%)	3.9% (2.3-6.6%)	52 (30-88)	3100 (1900-5100)
Taiwan	2000	Female	45 to 49	21% (12-34%)	79% (66-88%)	0.61 (0.25-1.2)	1.9% (1-3.2%)	3.9% (2.3-6.7%)	68 (38-120)	3300 (1900-5600)
Taiwan	2000	Female	50 to 54	21% (12-33%)	79% (67-88%)	0.63 (0.26-1.3)	2% (0.97-3.6%)	3.9% (2.2-6.7%)	68 (37-120)	2700 (1300-4800)
Taiwan	2000	Female	55 to 59	20% (12-31%)	80% (69-88%)	0.63 (0.26-1.3)	2.1% (0.9-4.1%)	3.8% (2-6.8%)	86 (44-160)	2800 (1200-5500)
Taiwan	2000	Female	60 to 64	19% (11-30%)	81% (70-89%)	0.62 (0.23-1.2)	2% (0.73-4.2%)	3.5% (1.6-6.4%)	120 (53-220)	3200 (1100-6300)
Taiwan	2000	Female	65 to 69	18% (11-29%)	82% (71-89%)	0.61 (0.26-1.2)	2% (0.6-4.1%)	3.2% (1.2-5.9%)	150 (57-270)	3300 (990-6700)
Taiwan	2000	Female	70 to 74	17% (10-27%)	83% (73-90%)	0.6 (0.24-1.2)	1.8% (0.3-4.2%)	2.6% (0.65-5.6%)	160 (40-350)	3000 (480-7000)
Taiwan	2000	Female	75 to 79	16% (9.6-26%)	84% (74-90%)	0.56 (0.22-1.1)	1.4% (0.014-3.5%)	1.9% (0.16-4.4%)	130 (12-310)	1900 (19-4800)
Taiwan	2000	Female	80 to 84	15% (9-24%)	85% (76-91%)	0.54 (0.22-1)	1.1% (-0.19-3.2%)	1.3% (-0.18-3.9%)	89 (-11-260)	970 (-170-2900)
Taiwan	2000	Female	85 to 89	16% (9.2-25%)	84% (75-91%)	0.56 (0.23-1.1)	0.93% (-0.33-3%)	1.1% (-0.35-3.5%)	61 (-20-190)	480 (-170-1500)
Taiwan	2000	Female	90 to 94	17% (10-27%)	83% (73-90%)	0.6 (0.23-1.2)	0.8% (-0.47-3%)	0.88% (-0.57-3.4%)	23 (-15-88)	130 (-74-490)
Taiwan	2000	Female	95 plus	20% (11-32%)	80% (68-89%)	0.69 (0.22-1.6)	0.88% (-0.67-3.8%)	0.91% (-0.85-4.2%)	7.8 (-7.3-37)	30 (-22-130)
Taiwan	2000	Male	All Ages	40% (34-48%)	60% (52-66%)	2.5 (1.4-4)	9.7% (6.2-13%)	11% (6.3-15%)	8500 (5000-12000)	290000 (190000-380000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	2000	Male	15 to 19	28% (18-43%)	72% (57-82%)	0.67 (0.27-1.4)	3.1% (1.6-5.5%)	4.5% (1.9-8.7%)	40 (17-77)	3700 (1900-6500)
Taiwan	2000	Male	20 to 24	41% (26-56%)	59% (44-74%)	1.6 (0.72-3.2)	7.1% (3.9-12%)	9.6% (4.5-18%)	110 (49-200)	9400 (5100-16000)
Taiwan	2000	Male	25 to 29	47% (30-64%)	53% (36-70%)	2.5 (1.1-4.6)	9.5% (5.2-15%)	14% (6.4-22%)	160 (77-270)	13000 (6700-20000)
Taiwan	2000	Male	30 to 34	50% (33-67%)	50% (33-67%)	3.2 (1.4-5.7)	11% (6.5-16%)	16% (8.1-24%)	280 (140-420)	19000 (11000-27000)
Taiwan	2000	Male	35 to 39	52% (36-69%)	48% (31-64%)	3.7 (1.7-7)	13% (7.9-18%)	17% (10-24%)	450 (270-650)	27000 (16000-37000)
Taiwan	2000	Male	40 to 44	53% (37-71%)	47% (29-63%)	4.3 (2-7.9)	15% (9.5-19%)	19% (12-26%)	680 (420-910)	35000 (23000-47000)
Taiwan	2000	Male	45 to 49	54% (35-72%)	46% (28-65%)	4.8 (2.2-8.8)	15% (10-20%)	20% (13-26%)	850 (560-1100)	39000 (26000-50000)
Taiwan	2000	Male	50 to 54	54% (36-72%)	46% (28-64%)	4.9 (2.3-9.1)	15% (9.9-19%)	19% (12-24%)	730 (500-960)	30000 (20000-39000)
Taiwan	2000	Male	55 to 59	52% (35-71%)	48% (29-65%)	4.6 (2.1-8.5)	13% (8.3-17%)	16% (10-21%)	740 (460-970)	26000 (16000-34000)
Taiwan	2000	Male	60 to 64	49% (31-68%)	51% (32-69%)	4.1 (1.7-7.6)	11% (6.1-15%)	13% (7.4-18%)	800 (450-1100)	24000 (13000-33000)
Taiwan	2000	Male	65 to 69	46% (31-63%)	54% (37-69%)	3.4 (1.5-6.4)	9.3% (4.7-13%)	11% (5.5-16%)	900 (460-1300)	22000 (11000-32000)
Taiwan	2000	Male	70 to 74	43% (27-60%)	57% (40-73%)	2.8 (1.2-5.4)	7.5% (3.1-12%)	8.7% (3.6-14%)	1000 (440-1600)	20000 (8800-32000)
Taiwan	2000	Male	75 to 79	40% (25-58%)	60% (42-75%)	2.4 (0.98-4.7)	6.2% (2.3-10%)	7.1% (2.5-12%)	820 (290-1400)	13000 (4500-21000)
Taiwan	2000	Male	80 to 84	37% (23-54%)	63% (46-77%)	2.2 (0.94-4.2)	5.7% (1.9-9.8%)	6.4% (2-11%)	530 (170-920)	5900 (1900-10000)
Taiwan	2000	Male	85 to 89	36% (23-52%)	64% (48-77%)	2 (0.87-3.9)	5% (1.3-9.3%)	5.6% (1.4-10%)	290 (73-540)	2300 (610-4200)
Taiwan	2000	Male	90 to 94	38% (25-54%)	62% (46-75%)	2 (0.83-4)	4.3% (0.8-8.4%)	4.7% (0.75-9.4%)	80 (13-160)	440 (81-850)
Taiwan	2000	Male	95 plus	42% (25-60%)	58% (40-75%)	2.1 (0.71-4.5)	4.1% (0.27-8.9%)	4.2% (0.025-9.8%)	18 (0.11-41)	67 (4.6-150)
Taiwan	2005	Female	All Ages	17% (14-21%)	83% (79-86%)	0.41 (0.22-0.69)	1.5% (0.76-2.7%)	2.1% (0.8-4.2%)	1100 (420-2200)	34000 (17000-60000)
Taiwan	2005	Female	15 to 19	15% (9.4-25%)	85% (75-91%)	0.31 (0.13-0.66)	1% (0.59-1.7%)	1.9% (0.71-4.1%)	4.6 (1.7-9.7)	690 (400-1100)
Taiwan	2005	Female	20 to 24	19% (11-29%)	81% (71-89%)	0.34 (0.14-0.74)	1.6% (1-2.4%)	2.1% (0.84-4.4%)	8 (3.1-17)	1500 (930-2400)
Taiwan	2005	Female	25 to 29	19% (11-30%)	81% (70-89%)	0.35 (0.14-0.73)	1.6% (1-2.4%)	2.3% (0.91-4.6%)	11 (4.3-22)	1800 (1100-2700)
Taiwan	2005	Female	30 to 34	20% (12-31%)	80% (69-88%)	0.4 (0.16-0.81)	1.6% (1-2.4%)	2.7% (1.3-4.9%)	15 (7.7-28)	1800 (1100-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	2005	Female	35 to 39	21% (13-34%)	79% (66-87%)	0.5 (0.21-0.99)	1.7% (1.1-2.7%)	3.5% (2-5.9%)	29 (16-49)	2300 (1500-3700)
Taiwan	2005	Female	40 to 44	22% (13-35%)	78% (65-87%)	0.57 (0.24-1.2)	1.8% (1.1-3%)	3.8% (2.1-6.5%)	47 (27-80)	2900 (1800-4900)
Taiwan	2005	Female	45 to 49	21% (12-33%)	79% (67-88%)	0.61 (0.25-1.2)	1.7% (0.87-3%)	3.7% (2.1-6.5%)	64 (35-110)	3200 (1700-5600)
Taiwan	2005	Female	50 to 54	21% (13-33%)	79% (67-87%)	0.64 (0.26-1.2)	1.8% (0.8-3.2%)	3.7% (1.9-6.4%)	86 (44-150)	3500 (1600-6300)
Taiwan	2005	Female	55 to 59	20% (12-31%)	80% (69-88%)	0.64 (0.25-1.3)	1.8% (0.76-3.6%)	3.5% (1.8-6.4%)	83 (42-150)	2800 (1200-5400)
Taiwan	2005	Female	60 to 64	19% (11-30%)	81% (70-89%)	0.62 (0.24-1.2)	1.9% (0.67-3.9%)	3.4% (1.6-6.3%)	100 (49-180)	2800 (1000-5600)
Taiwan	2005	Female	65 to 69	18% (11-28%)	82% (72-89%)	0.61 (0.26-1.2)	1.9% (0.67-3.9%)	3.2% (1.4-5.9%)	150 (65-270)	3300 (1100-6700)
Taiwan	2005	Female	70 to 74	17% (11-27%)	83% (73-89%)	0.6 (0.24-1.3)	1.8% (0.39-4%)	2.6% (0.77-5.5%)	170 (46-340)	3100 (670-7100)
Taiwan	2005	Female	75 to 79	16% (9.6-26%)	84% (74-90%)	0.56 (0.22-1.2)	1.3% (0.11-3.4%)	1.8% (0.32-4.3%)	140 (26-350)	2100 (180-5400)
Taiwan	2005	Female	80 to 84	15% (9-25%)	85% (75-91%)	0.55 (0.23-1.1)	1% (-0.11-2.9%)	1.3% (-0.056-3.5%)	110 (-4.6-290)	1200 (-120-3300)
Taiwan	2005	Female	85 to 89	16% (9.5-25%)	84% (75-91%)	0.57 (0.23-1.1)	0.88% (-0.23-2.8%)	1% (-0.23-3.2%)	66 (-14-210)	530 (-140-1600)
Taiwan	2005	Female	90 to 94	17% (10-27%)	83% (73-90%)	0.62 (0.24-1.3)	0.77% (-0.37-2.9%)	0.85% (-0.46-3.3%)	31 (-17-120)	180 (-87-680)
Taiwan	2005	Female	95 plus	20% (11-33%)	80% (67-89%)	0.71 (0.23-1.6)	0.88% (-0.58-3.8%)	0.91% (-0.76-4.2%)	12 (-10-56)	46 (-31-200)
Taiwan	2005	Male	All Ages	41% (35-49%)	59% (51-65%)	2.6 (1.4-4.3)	10% (6.6-13%)	11% (6.3-15%)	9100 (5300-13000)	310000 (200000-410000)
Taiwan	2005	Male	15 to 19	28% (17-43%)	72% (57-83%)	0.67 (0.27-1.4)	3% (1.6-5.2%)	4.6% (1.9-9%)	26 (11-51)	2600 (1400-4400)
Taiwan	2005	Male	20 to 24	40% (26-57%)	60% (43-74%)	1.6 (0.68-3.3)	6.9% (3.9-11%)	9.8% (4.3-18%)	86 (37-160)	7900 (4500-13000)
Taiwan	2005	Male	25 to 29	46% (29-63%)	54% (37-71%)	2.4 (1-4.7)	9.4% (5.2-14%)	14% (6.4-22%)	170 (81-290)	14000 (7300-21000)
Taiwan	2005	Male	30 to 34	50% (32-67%)	50% (33-68%)	3.1 (1.4-5.8)	11% (6.7-17%)	16% (8.7-24%)	260 (140-400)	18000 (10000-25000)
Taiwan	2005	Male	35 to 39	52% (36-70%)	48% (30-64%)	3.7 (1.7-7)	13% (8.2-18%)	18% (11-26%)	450 (260-630)	27000 (16000-37000)
Taiwan	2005	Male	40 to 44	54% (37-71%)	46% (29-63%)	4.3 (2-8)	15% (10-20%)	20% (13-27%)	750 (460-990)	39000 (25000-51000)
Taiwan	2005	Male	45 to 49	54% (36-71%)	46% (29-64%)	4.8 (2.2-9)	16% (10-20%)	21% (13-27%)	970 (620-1300)	44000 (29000-57000)
Taiwan	2005	Male	50 to 54	54% (35-71%)	46% (29-65%)	4.9 (2.2-9)	15% (9.6-19%)	19% (12-25%)	1100 (690-1400)	43000 (28000-56000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	2005	Male	55 to 59	52% (34-70%)	48% (30-66%)	4.6 (2.1-8.3)	13% (8.2-17%)	16% (10-21%)	830 (510-1100)	29000 (18000-38000)
Taiwan	2005	Male	60 to 64	49% (31-68%)	51% (32-69%)	4.1 (1.8-7.8)	11% (6.1-15%)	13% (7.3-18%)	770 (430-1100)	23000 (13000-32000)
Taiwan	2005	Male	65 to 69	46% (30-65%)	54% (35-70%)	3.4 (1.5-6.6)	9% (4.7-13%)	11% (5.5-16%)	810 (410-1200)	20000 (10000-29000)
Taiwan	2005	Male	70 to 74	43% (28-61%)	57% (39-72%)	2.8 (1.2-5.5)	7.1% (3.2-11%)	8.4% (3.6-13%)	840 (370-1300)	17000 (7400-27000)
Taiwan	2005	Male	75 to 79	39% (25-56%)	61% (44-75%)	2.4 (0.96-4.8)	5.8% (2.1-9.8%)	6.6% (2.3-11%)	910 (320-1600)	14000 (5100-24000)
Taiwan	2005	Male	80 to 84	37% (22-54%)	63% (46-78%)	2.1 (0.92-4.1)	5.1% (1.7-9.1%)	5.8% (1.9-10%)	640 (210-1200)	7400 (2500-13000)
Taiwan	2005	Male	85 to 89	36% (22-52%)	64% (48-78%)	2 (0.87-3.8)	4.6% (1.2-8.7%)	5.1% (1.3-9.8%)	330 (82-640)	2700 (730-5100)
Taiwan	2005	Male	90 to 94	38% (25-53%)	62% (47-75%)	2 (0.81-4)	4.1% (0.83-8.1%)	4.4% (0.77-9.1%)	130 (22-270)	730 (150-1400)
Taiwan	2005	Male	95 plus	41% (25-59%)	59% (41-75%)	2 (0.71-4.6)	3.9% (0.43-8.7%)	4.1% (0.26-9.7%)	30 (1.8-70)	110 (12-250)
Taiwan	2010	Female	All Ages	18% (15-22%)	82% (78-85%)	0.44 (0.24-0.73)	1.5% (0.76-2.6%)	2.1% (0.85-4.2%)	1200 (500-2400)	35000 (18000-61000)
Taiwan	2010	Female	15 to 19	16% (9.7-25%)	84% (75-90%)	0.33 (0.13-0.69)	0.98% (0.57-1.6%)	2% (0.8-4.2%)	3.7 (1.4-7.5)	630 (360-1000)
Taiwan	2010	Female	20 to 24	19% (12-29%)	81% (71-88%)	0.35 (0.14-0.8)	1.6% (0.98-2.4%)	2.4% (0.94-5%)	5.6 (2.2-12)	1200 (710-1800)
Taiwan	2010	Female	25 to 29	19% (11-30%)	81% (70-89%)	0.37 (0.15-0.74)	1.6% (1.1-2.4%)	2.5% (0.89-5.1%)	9.7 (3.6-20)	1700 (1000-2500)
Taiwan	2010	Female	30 to 34	20% (12-31%)	80% (69-88%)	0.42 (0.17-0.88)	1.6% (1-2.4%)	2.8% (1.3-5.4%)	16 (7.4-31)	1900 (1200-3000)
Taiwan	2010	Female	35 to 39	22% (13-34%)	78% (66-87%)	0.52 (0.22-1)	1.7% (1-2.6%)	3.6% (2-6.3%)	25 (14-45)	2100 (1400-3300)
Taiwan	2010	Female	40 to 44	22% (14-35%)	78% (65-86%)	0.59 (0.25-1.1)	1.7% (1-2.8%)	3.9% (2.2-6.6%)	43 (24-74)	2700 (1700-4600)
Taiwan	2010	Female	45 to 49	22% (12-34%)	78% (66-88%)	0.62 (0.26-1.2)	1.6% (0.88-2.8%)	3.8% (2.1-6.4%)	63 (35-110)	3200 (1700-5500)
Taiwan	2010	Female	50 to 54	21% (13-33%)	79% (67-87%)	0.65 (0.28-1.3)	1.7% (0.77-3%)	3.7% (2-6.2%)	85 (46-140)	3500 (1700-6400)
Taiwan	2010	Female	55 to 59	20% (12-32%)	80% (68-88%)	0.66 (0.27-1.3)	1.7% (0.7-3.4%)	3.6% (1.9-6.3%)	110 (58-190)	3700 (1500-7500)
Taiwan	2010	Female	60 to 64	19% (11-31%)	81% (69-89%)	0.64 (0.25-1.3)	1.8% (0.68-3.6%)	3.4% (1.8-6.1%)	110 (54-190)	3000 (1100-6100)
Taiwan	2010	Female	65 to 69	18% (11-29%)	82% (71-89%)	0.63 (0.27-1.3)	1.9% (0.75-3.6%)	3.4% (1.6-5.9%)	130 (62-230)	3100 (1200-6100)
Taiwan	2010	Female	70 to 74	17% (11-28%)	83% (72-89%)	0.62 (0.27-1.3)	1.8% (0.52-3.8%)	2.8% (1.1-5.6%)	170 (67-340)	3300 (1000-7200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	2010	Female	75 to 79	16% (9.8-27%)	84% (73-90%)	0.59 (0.24-1.3)	1.4% (0.23-3.4%)	2% (0.48-4.5%)	160 (39-380)	2400 (410-6100)
Taiwan	2010	Female	80 to 84	16% (9.5-25%)	84% (75-91%)	0.57 (0.25-1.1)	1.1% (-0.022-3%)	1.4% (0.069-3.7%)	130 (6.7-370)	1500 (-29-4300)
Taiwan	2010	Female	85 to 89	16% (9.7-25%)	84% (75-90%)	0.59 (0.25-1.2)	0.92% (-0.13-2.8%)	1.1% (-0.13-3.4%)	95 (-11-300)	780 (-110-2500)
Taiwan	2010	Female	90 to 94	18% (11-27%)	82% (73-89%)	0.64 (0.24-1.3)	0.82% (-0.32-2.8%)	0.91% (-0.4-3.3%)	44 (-19-160)	260 (-100-900)
Taiwan	2010	Female	95 plus	20% (11-33%)	80% (67-89%)	0.73 (0.26-1.6)	0.94% (-0.46-3.9%)	0.99% (-0.63-4.4%)	21 (-13-94)	79 (-40-330)
Taiwan	2010	Male	All Ages	43% (36-51%)	57% (49-64%)	2.8 (1.5-4.7)	10% (6.9-13%)	11% (6.3-14%)	9500 (5600-13000)	320000 (210000-410000)
Taiwan	2010	Male	15 to 19	28% (18-43%)	72% (57-82%)	0.68 (0.27-1.5)	2.8% (1.6-4.8%)	4.6% (2-9%)	20 (8.7-39)	2200 (1300-3700)
Taiwan	2010	Male	20 to 24	41% (27-57%)	59% (43-73%)	1.7 (0.7-3.4)	6.8% (4-11%)	10% (4.6-20%)	60 (26-110)	5900 (3400-9700)
Taiwan	2010	Male	25 to 29	47% (30-64%)	53% (36-70%)	2.5 (1.1-4.8)	9% (5.2-13%)	14% (6.5-22%)	130 (59-210)	11000 (6100-16000)
Taiwan	2010	Male	30 to 34	50% (32-67%)	50% (33-68%)	3.2 (1.5-6)	11% (6.5-16%)	16% (9-24%)	230 (130-340)	17000 (9800-24000)
Taiwan	2010	Male	35 to 39	52% (36-69%)	48% (31-64%)	3.8 (1.7-7.3)	13% (8.2-18%)	19% (11-26%)	380 (230-530)	23000 (14000-32000)
Taiwan	2010	Male	40 to 44	53% (37-71%)	47% (29-63%)	4.3 (2-8)	15% (10-20%)	21% (13-27%)	680 (430-900)	36000 (23000-47000)
Taiwan	2010	Male	45 to 49	54% (36-71%)	46% (29-64%)	4.8 (2.3-8.5)	16% (10-20%)	21% (14-27%)	980 (650-1300)	46000 (30000-58000)
Taiwan	2010	Male	50 to 54	54% (36-71%)	46% (29-64%)	4.9 (2.3-8.8)	15% (9.9-19%)	20% (13-24%)	1100 (740-1400)	46000 (30000-59000)
Taiwan	2010	Male	55 to 59	52% (35-71%)	48% (29-65%)	4.7 (2.2-8.1)	13% (8.5-17%)	17% (11-22%)	1200 (750-1500)	41000 (27000-54000)
Taiwan	2010	Male	60 to 64	50% (32-69%)	50% (31-68%)	4.1 (1.8-7.6)	11% (6.2-15%)	14% (7.6-19%)	840 (470-1100)	26000 (14000-35000)
Taiwan	2010	Male	65 to 69	47% (31-64%)	53% (36-69%)	3.5 (1.6-6.4)	9% (4.7-13%)	11% (5.8-16%)	770 (400-1100)	20000 (10000-28000)
Taiwan	2010	Male	70 to 74	43% (27-60%)	57% (40-73%)	2.9 (1.3-5.6)	7.1% (3.2-11%)	8.4% (3.8-13%)	770 (340-1200)	16000 (7000-24000)
Taiwan	2010	Male	75 to 79	40% (25-58%)	60% (42-75%)	2.4 (1.1-4.9)	5.7% (2.2-9.8%)	6.6% (2.5-11%)	780 (290-1300)	12000 (4600-21000)
Taiwan	2010	Male	80 to 84	38% (23-54%)	62% (46-77%)	2.2 (0.97-4.4)	5% (1.7-9%)	5.7% (1.9-10%)	820 (270-1500)	9500 (3300-17000)
Taiwan	2010	Male	85 to 89	37% (23-52%)	63% (48-77%)	2.1 (0.89-4.1)	4.5% (1.4-8.5%)	5% (1.5-9.7%)	480 (140-920)	4000 (1300-7600)
Taiwan	2010	Male	90 to 94	38% (25-53%)	62% (47-75%)	2 (0.84-4.1)	4% (1-8%)	4.4% (0.96-9.2%)	180 (38-370)	1000 (260-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	2010	Male	95 plus	42% (25-61%)	58% (39-75%)	2.1 (0.8-4.5)	4% (0.71-8.9%)	4.2% (0.49-9.9%)	61 (7-140)	220 (40-500)
Taiwan	2016	Female	All Ages	18% (15-22%)	82% (78-85%)	0.47 (0.26-0.79)	1.6% (0.79-2.7%)	2.2% (0.89-4.2%)	1600 (610-3000)	40000 (20000-71000)
Taiwan	2016	Female	15 to 19	16% (9.8-25%)	84% (75-90%)	0.34 (0.14-0.7)	0.96% (0.58-1.6%)	2% (0.77-4.1%)	3.1 (1.2-6.6)	560 (320-930)
Taiwan	2016	Female	20 to 24	19% (12-29%)	81% (71-88%)	0.36 (0.14-0.79)	1.6% (1-2.4%)	2.4% (0.95-4.9%)	5.2 (2-11)	1100 (700-1800)
Taiwan	2016	Female	25 to 29	19% (12-30%)	81% (70-88%)	0.37 (0.15-0.78)	1.6% (1-2.4%)	2.5% (0.98-5%)	6.9 (2.6-14)	1300 (800-1900)
Taiwan	2016	Female	30 to 34	20% (12-32%)	80% (68-88%)	0.42 (0.18-0.86)	1.5% (0.97-2.4%)	2.8% (1.3-5.5%)	15 (6.3-28)	1800 (1100-2900)
Taiwan	2016	Female	35 to 39	22% (13-34%)	78% (66-87%)	0.52 (0.22-1)	1.7% (1.1-2.7%)	3.7% (2-6.6%)	28 (14-51)	2400 (1500-3900)
Taiwan	2016	Female	40 to 44	22% (13-35%)	78% (65-87%)	0.59 (0.26-1.2)	1.7% (1-2.9%)	3.9% (2.2-6.6%)	40 (21-70)	2600 (1500-4300)
Taiwan	2016	Female	45 to 49	22% (12-34%)	78% (66-88%)	0.64 (0.27-1.2)	1.7% (0.91-2.9%)	3.9% (2.1-6.5%)	60 (32-110)	3100 (1600-5500)
Taiwan	2016	Female	50 to 54	21% (13-33%)	79% (67-87%)	0.67 (0.3-1.3)	1.7% (0.83-3.2%)	3.8% (2.2-6.4%)	93 (50-150)	3900 (1800-7300)
Taiwan	2016	Female	55 to 59	20% (12-32%)	80% (68-88%)	0.67 (0.29-1.3)	1.7% (0.72-3.3%)	3.7% (2-6.2%)	120 (64-220)	4200 (1800-8200)
Taiwan	2016	Female	60 to 64	19% (11-31%)	81% (69-89%)	0.65 (0.26-1.3)	1.8% (0.72-3.6%)	3.6% (1.9-6.3%)	170 (89-320)	4900 (1900-9900)
Taiwan	2016	Female	65 to 69	18% (11-29%)	82% (71-89%)	0.64 (0.28-1.2)	2% (0.84-3.7%)	3.5% (1.8-6%)	160 (79-270)	3700 (1600-7100)
Taiwan	2016	Female	70 to 74	17% (11-27%)	83% (73-89%)	0.62 (0.26-1.3)	1.9% (0.66-3.9%)	3.1% (1.3-5.8%)	200 (84-390)	3800 (1300-7900)
Taiwan	2016	Female	75 to 79	16% (9.9-26%)	84% (74-90%)	0.6 (0.24-1.2)	1.5% (0.34-3.5%)	2.2% (0.68-4.7%)	210 (63-440)	3100 (680-7200)
Taiwan	2016	Female	80 to 84	16% (9.3-26%)	84% (74-91%)	0.59 (0.25-1.2)	1.2% (0.095-3.1%)	1.5% (0.2-3.8%)	180 (23-450)	2000 (160-5300)
Taiwan	2016	Female	85 to 89	17% (9.7-26%)	83% (74-90%)	0.61 (0.26-1.2)	0.99% (-0.085-2.9%)	1.2% (-0.11-3.5%)	140 (-13-420)	1200 (-95-3500)
Taiwan	2016	Female	90 to 94	18% (11-29%)	82% (71-89%)	0.66 (0.27-1.4)	0.9% (-0.24-3.1%)	1% (-0.32-3.6%)	81 (-26-290)	470 (-120-1600)
Taiwan	2016	Female	95 plus	21% (11-34%)	79% (66-89%)	0.76 (0.27-1.7)	1.1% (-0.45-4%)	1.1% (-0.61-4.5%)	38 (-21-160)	140 (-60-540)
Taiwan	2016	Male	All Ages	44% (37-52%)	56% (48-63%)	3 (1.7-5)	10% (6.9-13%)	10% (6.1-14%)	11000 (6100-15000)	340000 (220000-450000)
Taiwan	2016	Male	15 to 19	29% (18-43%)	71% (57-82%)	0.7 (0.28-1.5)	2.7% (1.6-4.5%)	4.6% (2-8.7%)	17 (7.3-33)	1900 (1100-3300)
Taiwan	2016	Male	20 to 24	41% (27-57%)	59% (43-73%)	1.7 (0.72-3.4)	6.7% (4-11%)	10% (4.7-19%)	56 (25-110)	5800 (3400-9400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Taiwan	2016	Male	25 to 29	47% (30-64%)	53% (36-70%)	2.5 (1.1-4.9)	8.8% (5.2-13%)	14% (7-23%)	94 (45-160)	8300 (4800-13000)
Taiwan	2016	Male	30 to 34	50% (33-68%)	50% (32-67%)	3.2 (1.5-6.2)	11% (6.6-15%)	16% (9-24%)	210 (110-320)	15000 (9300-22000)
Taiwan	2016	Male	35 to 39	53% (37-70%)	47% (30-63%)	3.9 (1.7-7.6)	13% (8.1-18%)	19% (11-25%)	410 (240-590)	25000 (15000-35000)
Taiwan	2016	Male	40 to 44	54% (38-72%)	46% (28-62%)	4.4 (2-8.2)	15% (9.6-20%)	21% (13-27%)	590 (360-820)	31000 (20000-43000)
Taiwan	2016	Male	45 to 49	55% (36-71%)	45% (29-64%)	4.9 (2.3-8.8)	16% (11-20%)	21% (14-27%)	910 (610-1200)	43000 (29000-57000)
Taiwan	2016	Male	50 to 54	54% (36-72%)	46% (28-64%)	5 (2.5-9.2)	15% (10-19%)	20% (14-25%)	1200 (790-1600)	49000 (33000-66000)
Taiwan	2016	Male	55 to 59	53% (35-71%)	47% (29-65%)	4.7 (2.3-8.6)	13% (8.8-17%)	17% (11-22%)	1300 (840-1800)	46000 (30000-63000)
Taiwan	2016	Male	60 to 64	50% (33-69%)	50% (31-67%)	4.2 (1.9-7.8)	11% (6.5-15%)	14% (8.1-19%)	1300 (770-1900)	41000 (24000-57000)
Taiwan	2016	Male	65 to 69	47% (32-65%)	53% (35-68%)	3.6 (1.6-6.4)	9.1% (5-13%)	11% (6.1-16%)	900 (480-1300)	23000 (12000-33000)
Taiwan	2016	Male	70 to 74	43% (28-61%)	57% (39-72%)	2.9 (1.3-5.7)	7.3% (3.3-11%)	8.7% (3.8-13%)	850 (390-1300)	17000 (8000-27000)
Taiwan	2016	Male	75 to 79	40% (26-58%)	60% (42-74%)	2.5 (1.1-5.1)	5.9% (2.3-9.8%)	6.8% (2.5-11%)	780 (290-1300)	12000 (4700-21000)
Taiwan	2016	Male	80 to 84	38% (24-55%)	62% (45-76%)	2.3 (1-4.3)	5.2% (1.7-9.1%)	5.9% (1.9-11%)	780 (260-1400)	9100 (3100-16000)
Taiwan	2016	Male	85 to 89	37% (23-52%)	63% (48-77%)	2.1 (0.93-4.1)	4.6% (1.4-8.5%)	5.1% (1.5-9.6%)	750 (230-1400)	6200 (2000-11000)
Taiwan	2016	Male	90 to 94	38% (25-54%)	62% (46-75%)	2.1 (0.91-4.2)	4.1% (1.1-8.1%)	4.5% (1.1-9.1%)	330 (80-670)	1900 (520-3800)
Taiwan	2016	Male	95 plus	42% (25-61%)	58% (39-75%)	2.2 (0.8-4.6)	4.1% (0.76-8.9%)	4.4% (0.58-9.9%)	110 (15-260)	420 (80-910)
Cambodia	1990	Female	All Ages	8.5% (5.9-12%)	92% (88-94%)	0.24 (0.15-0.36)	1.1% (0.63-1.6%)	2.1% (1.1-3.3%)	890 (450-1500)	28000 (17000-42000)
Cambodia	1990	Female	15 to 19	8.3% (3.7-16%)	92% (84-96%)	0.22 (0.069-0.49)	0.72% (0.34-1.4%)	0.9% (0.31-1.9%)	6.9 (2.4-15)	700 (320-1400)
Cambodia	1990	Female	20 to 24	10% (4.1-20%)	90% (80-96%)	0.29 (0.099-0.64)	1.4% (0.63-2.7%)	1.6% (0.5-3.5%)	16 (5.1-36)	1500 (670-2900)
Cambodia	1990	Female	25 to 29	11% (4.6-21%)	89% (79-95%)	0.34 (0.11-0.72)	1.6% (0.75-3.2%)	2% (0.65-4.4%)	20 (6.7-46)	1700 (730-3200)
Cambodia	1990	Female	30 to 34	12% (4.8-22%)	88% (78-95%)	0.39 (0.13-0.81)	1.9% (0.92-3.5%)	2.4% (1-4.9%)	27 (11-55)	1900 (900-3500)
Cambodia	1990	Female	35 to 39	13% (5.6-24%)	87% (76-94%)	0.43 (0.15-0.91)	2.4% (1.2-4.4%)	3.3% (1.5-6%)	42 (19-77)	2400 (1200-4300)
Cambodia	1990	Female	40 to 44	13% (5.9-24%)	87% (76-94%)	0.48 (0.17-1)	2.6% (1.3-4.8%)	3.6% (1.7-6.5%)	43 (20-78)	2200 (1100-3900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	1990	Female	45 to 49	13% (5.7-24%)	87% (76-94%)	0.54 (0.18-1.2)	2.9% (1.4-5.5%)	3.9% (1.8-7.5%)	53 (24-100)	2300 (1100-4400)
Cambodia	1990	Female	50 to 54	14% (6.1-26%)	86% (74-94%)	0.61 (0.2-1.4)	3.6% (1.6-6.8%)	4.6% (2-8.7%)	78 (34-150)	2900 (1300-5500)
Cambodia	1990	Female	55 to 59	14% (6.5-26%)	86% (74-94%)	0.66 (0.21-1.5)	3.7% (1.5-7.1%)	4.5% (1.8-8.7%)	94 (38-190)	3100 (1200-6000)
Cambodia	1990	Female	60 to 64	14% (6.2-26%)	86% (74-94%)	0.66 (0.21-1.6)	3.7% (1.3-7.4%)	4.4% (1.5-9%)	110 (39-230)	3100 (1100-6300)
Cambodia	1990	Female	65 to 69	13% (6.2-24%)	87% (76-94%)	0.61 (0.19-1.4)	3.4% (1.1-6.8%)	4% (1.3-8%)	120 (40-240)	2800 (920-5500)
Cambodia	1990	Female	70 to 74	13% (6-23%)	87% (77-94%)	0.55 (0.17-1.3)	3% (0.89-6.7%)	3.5% (1-7.7%)	110 (32-250)	2000 (600-4500)
Cambodia	1990	Female	75 to 79	12% (5.5-23%)	88% (77-94%)	0.51 (0.14-1.3)	2.5% (0.6-5.7%)	2.8% (0.68-6.5%)	77 (18-180)	1100 (260-2500)
Cambodia	1990	Female	80 to 84	12% (5.4-23%)	88% (77-95%)	0.51 (0.15-1.2)	2.5% (0.67-5.6%)	2.9% (0.77-6.4%)	52 (14-120)	540 (150-1200)
Cambodia	1990	Female	85 to 89	13% (5.7-24%)	87% (76-94%)	0.55 (0.16-1.4)	2.3% (0.58-5.6%)	2.7% (0.67-6.5%)	25 (6.2-62)	190 (47-450)
Cambodia	1990	Female	90 to 94	15% (6.5-28%)	85% (72-94%)	0.62 (0.16-1.6)	2.1% (0.42-5.3%)	2.5% (0.48-6.3%)	8.5 (1.6-21)	44 (8.6-110)
Cambodia	1990	Female	95 plus	19% (7.1-37%)	81% (63-93%)	0.76 (0.15-2.1)	2.3% (0.3-6%)	2.6% (0.33-7%)	2.4 (0.31-6.6)	8.4 (1.1-22)
Cambodia	1990	Male	All Ages	25% (19-32%)	75% (68-81%)	0.72 (0.49-0.98)	2.5% (1.7-3.6%)	4.1% (2.6-6%)	1800 (1100-2700)	76000 (49000-110000)
Cambodia	1990	Male	15 to 19	23% (13-38%)	77% (62-87%)	0.39 (0.16-0.74)	1.5% (0.75-2.6%)	1.8% (0.71-3.3%)	19 (7.6-35)	1700 (810-2900)
Cambodia	1990	Male	20 to 24	35% (20-53%)	65% (47-80%)	0.92 (0.43-1.6)	4.4% (2.2-6.9%)	5.1% (2.3-8.5%)	72 (32-120)	5500 (2700-8800)
Cambodia	1990	Male	25 to 29	41% (22-59%)	59% (41-78%)	1.3 (0.62-2.1)	6.6% (3.4-11%)	8% (3.7-13%)	100 (47-170)	6900 (3600-11000)
Cambodia	1990	Male	30 to 34	44% (24-63%)	56% (37-76%)	1.6 (0.83-2.6)	8.1% (4.7-12%)	10% (5.6-16%)	130 (73-210)	8100 (4700-12000)
Cambodia	1990	Male	35 to 39	46% (28-66%)	54% (34-72%)	1.8 (0.88-3.1)	9.4% (5.5-14%)	12% (6.6-18%)	180 (97-270)	9500 (5500-15000)
Cambodia	1990	Male	40 to 44	46% (28-67%)	54% (33-72%)	1.9 (0.96-3.2)	9.6% (5.7-14%)	12% (6.8-18%)	170 (96-260)	8200 (4700-13000)
Cambodia	1990	Male	45 to 49	44% (25-64%)	56% (36-75%)	1.9 (0.89-3.3)	9.5% (5.4-15%)	11% (6.5-18%)	190 (110-300)	8300 (4700-13000)
Cambodia	1990	Male	50 to 54	42% (24-62%)	58% (38-76%)	1.8 (0.83-3.1)	8.7% (4.6-14%)	10% (5.4-17%)	220 (110-350)	8200 (4400-13000)
Cambodia	1990	Male	55 to 59	40% (23-60%)	60% (40-77%)	1.6 (0.74-2.9)	7.4% (3.7-13%)	8.7% (4.2-15%)	220 (100-370)	7100 (3400-12000)
Cambodia	1990	Male	60 to 64	37% (21-57%)	63% (43-79%)	1.4 (0.62-2.7)	6.5% (2.8-11%)	7.4% (3.2-13%)	200 (87-360)	5600 (2400-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	1990	Male	65 to 69	33% (19-52%)	67% (48-81%)	1.2 (0.5-2.1)	5.1% (2.3-9.3%)	5.9% (2.6-11%)	150 (65-280)	3500 (1500-6400)
Cambodia	1990	Male	70 to 74	30% (17-47%)	70% (53-83%)	0.95 (0.41-1.8)	4.3% (1.6-8.7%)	4.9% (1.7-9.9%)	110 (39-220)	2000 (740-4100)
Cambodia	1990	Male	75 to 79	27% (15-44%)	73% (56-85%)	0.81 (0.33-1.6)	3.3% (1.1-6.8%)	3.7% (1.2-7.7%)	54 (17-110)	770 (250-1600)
Cambodia	1990	Male	80 to 84	25% (13-42%)	75% (58-87%)	0.7 (0.27-1.4)	2.9% (0.96-5.9%)	3.3% (1.1-6.7%)	23 (7.4-46)	240 (80-490)
Cambodia	1990	Male	85 to 89	24% (13-39%)	76% (61-87%)	0.62 (0.25-1.3)	2.1% (0.69-4.5%)	2.4% (0.75-5.2%)	6.9 (2.2-15)	53 (17-110)
Cambodia	1990	Male	90 to 94	25% (13-42%)	75% (58-87%)	0.6 (0.23-1.2)	1.6% (0.38-3.7%)	1.8% (0.4-4.3%)	2.3 (0.49-5.5)	12 (2.9-28)
Cambodia	1990	Male	95 plus	29% (13-48%)	71% (52-87%)	0.63 (0.21-1.4)	1.4% (0.27-3.7%)	1.5% (0.22-4.1%)	0.55 (0.077-1.5)	2 (0.37-5.2)
Cambodia	1995	Female	All Ages	8.5% (6.1-12%)	92% (88-94%)	0.41 (0.26-0.6)	1.7% (1.2-2.3%)	3.3% (2.1-4.6%)	1600 (1000-2200)	49000 (35000-67000)
Cambodia	1995	Female	15 to 19	8.3% (3.9-15%)	92% (85-96%)	0.38 (0.13-0.85)	1.2% (0.52-2.2%)	1.6% (0.53-3.4%)	11 (3.7-23)	1000 (440-2000)
Cambodia	1995	Female	20 to 24	10% (4.4-19%)	90% (81-96%)	0.52 (0.18-1.1)	2.1% (0.96-4%)	2.8% (0.96-5.6%)	30 (11-62)	2500 (1100-4700)
Cambodia	1995	Female	25 to 29	11% (4.8-21%)	89% (79-95%)	0.6 (0.19-1.3)	2.4% (1.1-4.4%)	3.2% (1.2-6.2%)	37 (14-72)	2800 (1300-5100)
Cambodia	1995	Female	30 to 34	12% (4.9-23%)	88% (77-95%)	0.67 (0.23-1.4)	2.8% (1.4-4.9%)	3.8% (1.7-6.9%)	48 (21-86)	3200 (1500-5500)
Cambodia	1995	Female	35 to 39	13% (5.3-24%)	87% (76-95%)	0.74 (0.25-1.5)	3.5% (1.8-6%)	4.9% (2.4-8.5%)	70 (34-120)	3900 (2000-6600)
Cambodia	1995	Female	40 to 44	13% (6-24%)	87% (76-94%)	0.83 (0.29-1.8)	3.9% (2-6.5%)	5.3% (2.7-9%)	86 (43-140)	4300 (2200-7200)
Cambodia	1995	Female	45 to 49	13% (5.8-24%)	87% (76-94%)	0.94 (0.31-2.1)	4.3% (2.2-7.1%)	5.7% (2.9-9.8%)	92 (46-160)	4000 (2000-6800)
Cambodia	1995	Female	50 to 54	14% (6.1-25%)	86% (75-94%)	1.1 (0.34-2.4)	5.2% (2.7-8.7%)	6.7% (3.3-11%)	130 (65-220)	4900 (2500-8100)
Cambodia	1995	Female	55 to 59	14% (6.2-25%)	86% (75-94%)	1.2 (0.36-2.6)	5.3% (2.6-9.1%)	6.6% (3.2-11%)	160 (74-270)	5100 (2500-8800)
Cambodia	1995	Female	60 to 64	14% (5.9-26%)	86% (74-94%)	1.2 (0.34-2.7)	5.3% (2.4-9.4%)	6.4% (3-12%)	180 (83-320)	5100 (2300-8900)
Cambodia	1995	Female	65 to 69	13% (6-24%)	87% (76-94%)	1.1 (0.33-2.4)	5.3% (2.2-9.1%)	6.2% (2.7-11%)	210 (90-360)	4800 (2100-8200)
Cambodia	1995	Female	70 to 74	13% (5.8-23%)	87% (77-94%)	0.97 (0.29-2.3)	4.7% (1.8-8.7%)	5.5% (2.1-10%)	200 (78-370)	3700 (1400-6800)
Cambodia	1995	Female	75 to 79	12% (5.3-23%)	88% (77-95%)	0.9 (0.25-2.3)	4% (1.3-7.6%)	4.6% (1.5-8.7%)	150 (49-290)	2200 (710-4100)
Cambodia	1995	Female	80 to 84	12% (5.1-23%)	88% (77-95%)	0.89 (0.26-2.1)	4.1% (1.5-7.6%)	4.7% (1.7-8.8%)	100 (36-190)	1100 (370-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	1995	Female	85 to 89	13% (5.9-24%)	87% (76-94%)	0.96 (0.28-2.3)	3.8% (1.4-7.6%)	4.4% (1.6-8.9%)	47 (16-95)	350 (120-700)
Cambodia	1995	Female	90 to 94	15% (6.6-28%)	85% (72-93%)	1.1 (0.27-2.8)	3.5% (1.1-7.1%)	4.1% (1.3-8.3%)	15 (4.8-31)	78 (25-160)
Cambodia	1995	Female	95 plus	19% (7.8-36%)	81% (64-92%)	1.3 (0.26-3.6)	3.6% (0.84-7.8%)	4.1% (0.95-9%)	4.1 (0.92-9)	14 (3.3-31)
Cambodia	1995	Male	All Ages	25% (19-32%)	75% (68-81%)	1.2 (0.83-1.6)	4.1% (2.9-5.2%)	6.7% (4.6-8.8%)	3400 (2300-4500)	130000 (97000-170000)
Cambodia	1995	Male	15 to 19	23% (13-38%)	77% (62-87%)	0.67 (0.29-1.3)	2.3% (1.2-3.9%)	2.8% (1.3-5.1%)	27 (12-49)	2200 (1200-3900)
Cambodia	1995	Male	20 to 24	35% (19-54%)	65% (46-81%)	1.6 (0.76-2.7)	6.4% (3.6-9.9%)	7.6% (3.9-12%)	130 (69-210)	9700 (5300-15000)
Cambodia	1995	Male	25 to 29	41% (23-59%)	59% (41-77%)	2.2 (1.1-3.6)	9.4% (5.5-15%)	12% (6.4-18%)	200 (110-310)	13000 (7700-20000)
Cambodia	1995	Male	30 to 34	44% (25-63%)	56% (37-75%)	2.7 (1.4-4.5)	11% (7.1-17%)	14% (8.5-21%)	230 (140-330)	14000 (8500-20000)
Cambodia	1995	Male	35 to 39	46% (29-66%)	54% (34-71%)	3.1 (1.6-5.3)	13% (8.4-19%)	16% (10-23%)	290 (170-400)	15000 (9400-21000)
Cambodia	1995	Male	40 to 44	46% (29-66%)	54% (34-71%)	3.3 (1.6-5.5)	14% (9-19%)	17% (11-23%)	340 (220-480)	16000 (11000-23000)
Cambodia	1995	Male	45 to 49	44% (25-64%)	56% (36-75%)	3.2 (1.6-5.6)	14% (8.7-19%)	17% (11-23%)	320 (190-440)	14000 (8300-18000)
Cambodia	1995	Male	50 to 54	42% (24-62%)	58% (38-76%)	3.1 (1.4-5.3)	13% (7.6-18%)	16% (9.2-21%)	340 (200-470)	13000 (7500-17000)
Cambodia	1995	Male	55 to 59	40% (23-59%)	60% (41-77%)	2.8 (1.3-5)	12% (6.4-16%)	14% (7.6-20%)	370 (210-540)	12000 (6700-18000)
Cambodia	1995	Male	60 to 64	37% (21-57%)	63% (43-79%)	2.4 (1.1-4.6)	10% (5.4-16%)	12% (6.2-18%)	380 (190-580)	11000 (5400-16000)
Cambodia	1995	Male	65 to 69	34% (19-52%)	66% (48-81%)	2 (0.9-3.7)	8.8% (4.4-14%)	10% (5.1-16%)	320 (160-510)	7300 (3600-12000)
Cambodia	1995	Male	70 to 74	30% (17-46%)	70% (54-83%)	1.7 (0.73-3.2)	7.6% (3.4-13%)	8.7% (3.8-15%)	250 (110-430)	4600 (2000-7900)
Cambodia	1995	Male	75 to 79	27% (15-44%)	73% (56-85%)	1.4 (0.57-2.8)	6% (2.5-11%)	6.9% (2.8-12%)	140 (57-250)	2000 (820-3600)
Cambodia	1995	Male	80 to 84	25% (13-41%)	75% (59-87%)	1.2 (0.49-2.5)	5.4% (2.2-10%)	6.2% (2.5-12%)	62 (24-110)	650 (250-1200)
Cambodia	1995	Male	85 to 89	24% (13-39%)	76% (61-87%)	1.1 (0.45-2.2)	4.2% (1.7-8%)	4.8% (1.9-9.3%)	21 (8.1-40)	160 (62-300)
Cambodia	1995	Male	90 to 94	25% (14-41%)	75% (59-86%)	1.1 (0.43-2.2)	3.3% (1.2-6.8%)	3.8% (1.3-7.9%)	4.7 (1.6-9.8)	25 (8.8-51)
Cambodia	1995	Male	95 plus	29% (14-50%)	71% (50-86%)	1.1 (0.37-2.4)	3% (0.88-6.6%)	3.3% (0.88-7.5%)	1.2 (0.31-2.6)	4.2 (1.2-9.3)
Cambodia	2000	Female	All Ages	8.5% (6.1-12%)	92% (88-94%)	0.81 (0.53-1.2)	2.4% (1.8-3.1%)	4.6% (3.3-6.2%)	2300 (1600-3100)	72000 (54000-92000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	2000	Female	15 to 19	8.2% (3.9-15%)	92% (85-96%)	0.73 (0.26-1.6)	1.8% (0.85-3.2%)	2.7% (1.1-5.1%)	33 (13-61)	2800 (1300-5000)
Cambodia	2000	Female	20 to 24	9.9% (4.2-18%)	90% (82-96%)	0.98 (0.34-2.1)	2.9% (1.4-4.9%)	4% (1.7-7.3%)	34 (15-62)	2800 (1400-4700)
Cambodia	2000	Female	25 to 29	11% (4.5-20%)	89% (80-95%)	1.1 (0.38-2.5)	3.1% (1.6-5.1%)	4.2% (1.8-7.3%)	52 (24-91)	3900 (2000-6400)
Cambodia	2000	Female	30 to 34	12% (4.7-22%)	88% (78-95%)	1.3 (0.43-2.7)	3.6% (1.9-5.7%)	5% (2.4-8.1%)	72 (36-120)	4700 (2500-7400)
Cambodia	2000	Female	35 to 39	13% (5.4-23%)	87% (77-95%)	1.4 (0.48-2.9)	4.4% (2.4-6.9%)	6.2% (3.3-9.8%)	100 (56-160)	5900 (3300-9000)
Cambodia	2000	Female	40 to 44	13% (6.2-23%)	87% (77-94%)	1.6 (0.57-3.3)	4.9% (2.9-7.5%)	6.8% (3.9-10%)	120 (71-190)	6100 (3600-9100)
Cambodia	2000	Female	45 to 49	13% (5.9-24%)	87% (76-94%)	1.8 (0.6-4)	5.4% (3.1-8.3%)	7.3% (4.2-12%)	160 (91-250)	7000 (4100-11000)
Cambodia	2000	Female	50 to 54	14% (6.1-26%)	86% (74-94%)	2 (0.66-4.5)	6.5% (3.8-10%)	8.4% (5-13%)	190 (110-300)	7200 (4300-11000)
Cambodia	2000	Female	55 to 59	14% (6.4-26%)	86% (74-94%)	2.2 (0.73-4.9)	6.6% (3.7-11%)	8.2% (4.7-13%)	220 (130-360)	7200 (4100-12000)
Cambodia	2000	Female	60 to 64	14% (5.9-26%)	86% (74-94%)	2.2 (0.65-5.3)	6.5% (3.7-11%)	8% (4.6-13%)	260 (150-430)	7100 (4000-12000)
Cambodia	2000	Female	65 to 69	13% (6.2-24%)	87% (76-94%)	2.1 (0.64-4.5)	6.5% (3.5-11%)	7.8% (4.2-13%)	300 (160-490)	6900 (3700-11000)
Cambodia	2000	Female	70 to 74	13% (6.1-23%)	87% (77-94%)	1.8 (0.56-4.3)	6% (3.1-10%)	7.1% (3.7-12%)	270 (140-450)	5000 (2700-8400)
Cambodia	2000	Female	75 to 79	12% (5.4-23%)	88% (77-95%)	1.7 (0.47-4.3)	5.3% (2.6-9%)	6.1% (3-10%)	200 (99-350)	2900 (1400-4900)
Cambodia	2000	Female	80 to 84	12% (5.7-24%)	88% (76-94%)	1.8 (0.53-4)	5.4% (2.6-9.1%)	6.3% (3-11%)	150 (70-250)	1500 (730-2600)
Cambodia	2000	Female	85 to 89	13% (5.9-25%)	87% (75-94%)	1.9 (0.56-4.7)	5% (2.4-9.1%)	5.9% (2.8-11%)	73 (34-130)	540 (250-960)
Cambodia	2000	Female	90 to 94	16% (6.7-28%)	84% (72-93%)	2.2 (0.53-5.5)	4.7% (2.1-8.4%)	5.5% (2.5-9.7%)	22 (10-40)	120 (51-200)
Cambodia	2000	Female	95 plus	19% (7.1-38%)	81% (62-93%)	2.6 (0.54-7.4)	4.9% (2-9.4%)	5.6% (2.2-11%)	5.8 (2.3-11)	20 (7.9-38)
Cambodia	2000	Male	All Ages	25% (19-32%)	75% (68-81%)	2.3 (1.6-3.1)	6% (5-7.1%)	9.7% (7.9-12%)	5300 (4300-6400)	210000 (170000-250000)
Cambodia	2000	Male	15 to 19	23% (13-38%)	77% (62-87%)	1.3 (0.56-2.4)	3.8% (1.9-6.7%)	5% (2.4-9.1%)	88 (42-160)	7100 (3600-12000)
Cambodia	2000	Male	20 to 24	35% (20-53%)	65% (47-80%)	3.1 (1.5-5.2)	10% (6.3-15%)	13% (7.4-19%)	170 (100-260)	12000 (7400-18000)
Cambodia	2000	Male	25 to 29	41% (23-59%)	59% (41-77%)	4.3 (2.2-7.1)	12% (8.2-17%)	15% (9.5-21%)	330 (220-450)	21000 (15000-29000)
Cambodia	2000	Male	30 to 34	44% (25-63%)	56% (37-75%)	5.3 (2.8-8.7)	14% (9.4-18%)	17% (11-23%)	410 (290-530)	24000 (17000-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	2000	Male	35 to 39	46% (28-66%)	54% (34-72%)	6 (3-10)	15% (11-20%)	19% (13-25%)	470 (350-600)	25000 (19000-32000)
Cambodia	2000	Male	40 to 44	45% (28-65%)	55% (35-72%)	6.2 (3.1-11)	16% (12-20%)	19% (14-25%)	440 (320-550)	21000 (16000-26000)
Cambodia	2000	Male	45 to 49	44% (25-64%)	56% (36-75%)	6.2 (3-11)	16% (12-21%)	20% (14-25%)	520 (380-670)	22000 (16000-28000)
Cambodia	2000	Male	50 to 54	42% (23-62%)	58% (38-77%)	5.9 (2.8-10)	16% (11-20%)	19% (14-24%)	460 (330-590)	17000 (13000-22000)
Cambodia	2000	Male	55 to 59	39% (23-60%)	61% (40-77%)	5.3 (2.5-9.6)	15% (11-20%)	18% (13-23%)	480 (340-630)	15000 (11000-20000)
Cambodia	2000	Male	60 to 64	37% (20-56%)	63% (44-80%)	4.6 (2-8.7)	14% (9.4-18%)	16% (11-21%)	520 (350-710)	14000 (9800-20000)
Cambodia	2000	Male	65 to 69	33% (19-52%)	67% (48-81%)	3.9 (1.7-7.3)	12% (8.1-17%)	14% (9.5-20%)	510 (330-710)	12000 (7700-16000)
Cambodia	2000	Male	70 to 74	30% (17-46%)	70% (54-83%)	3.2 (1.4-6.1)	11% (6.9-16%)	13% (7.9-19%)	430 (260-630)	7900 (4800-11000)
Cambodia	2000	Male	75 to 79	27% (15-44%)	73% (56-85%)	2.7 (1.1-5.4)	9.3% (5.2-14%)	11% (6-16%)	260 (150-390)	3800 (2100-5600)
Cambodia	2000	Male	80 to 84	25% (13-41%)	75% (59-87%)	2.3 (0.92-4.6)	8.8% (4.6-13%)	10% (5.2-15%)	140 (70-200)	1400 (740-2200)
Cambodia	2000	Male	85 to 89	23% (12-38%)	77% (62-88%)	2 (0.83-4.2)	7.3% (3.7-11%)	8.4% (4.2-13%)	54 (27-85)	410 (210-640)
Cambodia	2000	Male	90 to 94	25% (13-40%)	75% (60-87%)	2 (0.8-4)	6.2% (2.9-9.8%)	7.1% (3.2-11%)	14 (6.1-22)	70 (32-110)
Cambodia	2000	Male	95 plus	28% (13-48%)	72% (52-87%)	2 (0.69-4.5)	5.7% (2.3-9.8%)	6.4% (2.5-11%)	2.3 (0.89-4)	8.1 (3.3-14)
Cambodia	2005	Female	All Ages	8.6% (6.2-12%)	91% (88-94%)	1.1 (0.73-1.6)	2.8% (2.2-3.7%)	5.3% (3.8-7.1%)	2500 (1800-3400)	78000 (60000-100000)
Cambodia	2005	Female	15 to 19	8.1% (3.6-15%)	92% (85-96%)	0.93 (0.31-2)	1.9% (0.95-3.2%)	3.2% (1.4-5.7%)	28 (12-50)	2500 (1300-4200)
Cambodia	2005	Female	20 to 24	9.8% (4.2-18%)	90% (82-96%)	1.3 (0.44-2.6)	2.9% (1.6-4.7%)	4.2% (1.9-7.3%)	53 (24-91)	4600 (2500-7300)
Cambodia	2005	Female	25 to 29	11% (4.4-19%)	89% (81-96%)	1.5 (0.49-3.2)	3.2% (1.7-5%)	4.6% (2.2-7.8%)	35 (17-59)	2700 (1500-4300)
Cambodia	2005	Female	30 to 34	12% (4.8-22%)	88% (78-95%)	1.6 (0.56-3.5)	3.6% (2.1-5.7%)	5.2% (2.7-8.4%)	64 (34-100)	4300 (2500-6700)
Cambodia	2005	Female	35 to 39	12% (5.2-23%)	88% (77-95%)	1.8 (0.62-3.8)	4.4% (2.5-6.7%)	6.4% (3.5-9.7%)	98 (54-150)	5700 (3300-8400)
Cambodia	2005	Female	40 to 44	13% (5.8-23%)	87% (77-94%)	2 (0.7-4.1)	4.9% (3-7.3%)	7.2% (4.4-11%)	130 (80-200)	6800 (4100-10000)
Cambodia	2005	Female	45 to 49	13% (5.6-24%)	87% (76-94%)	2.3 (0.76-5.2)	5.5% (3.3-8.3%)	7.8% (4.6-12%)	160 (95-250)	7300 (4300-11000)
Cambodia	2005	Female	50 to 54	13% (6.1-25%)	87% (75-94%)	2.6 (0.86-5.9)	6.5% (3.9-10%)	8.7% (5.3-13%)	240 (150-380)	9200 (5600-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	2005	Female	55 to 59	14% (6.3-26%)	86% (74-94%)	2.9 (0.93-6.6)	6.7% (3.9-11%)	8.6% (5-14%)	250 (140-410)	8200 (4700-13000)
Cambodia	2005	Female	60 to 64	14% (6.1-26%)	86% (74-94%)	3 (0.88-6.7)	6.6% (3.8-11%)	8.3% (4.8-14%)	280 (160-460)	7800 (4500-13000)
Cambodia	2005	Female	65 to 69	13% (6-23%)	87% (77-94%)	2.7 (0.85-5.8)	6.6% (3.6-11%)	8.1% (4.5-13%)	320 (180-530)	7400 (4100-12000)
Cambodia	2005	Female	70 to 74	13% (5.8-23%)	87% (77-94%)	2.4 (0.71-5.6)	6.1% (3.2-10%)	7.4% (3.9-12%)	320 (170-540)	6000 (3100-9900)
Cambodia	2005	Female	75 to 79	12% (5.2-23%)	88% (77-95%)	2.3 (0.62-5.6)	5.4% (2.7-9.1%)	6.4% (3.1-11%)	240 (120-400)	3400 (1700-5600)
Cambodia	2005	Female	80 to 84	12% (5.3-23%)	88% (77-95%)	2.3 (0.69-5.3)	5.5% (2.6-9%)	6.4% (3.1-11%)	160 (79-280)	1700 (840-2900)
Cambodia	2005	Female	85 to 89	13% (5.9-25%)	87% (75-94%)	2.5 (0.72-6)	5.2% (2.5-9.2%)	6.1% (2.9-11%)	89 (43-160)	660 (320-1200)
Cambodia	2005	Female	90 to 94	15% (6.8-28%)	85% (72-93%)	2.8 (0.7-7.1)	4.8% (2.4-8.4%)	5.7% (2.8-9.9%)	29 (15-51)	150 (76-260)
Cambodia	2005	Female	95 plus	19% (7.4-36%)	81% (64-93%)	3.3 (0.67-9.1)	5% (2.1-9.1%)	5.8% (2.5-11%)	7 (3-13)	24 (11-44)
Cambodia	2005	Male	All Ages	25% (20-33%)	75% (67-80%)	3.1 (2.3-4.2)	7.2% (5.9-8.5%)	11% (9.1-13%)	5900 (4900-7200)	23000 (19000-27000)
Cambodia	2005	Male	15 to 19	23% (13-38%)	77% (62-87%)	1.6 (0.7-3)	4.7% (2.4-7.9%)	6.4% (3.1-11%)	96 (45-170)	7700 (4000-13000)
Cambodia	2005	Male	20 to 24	34% (19-52%)	66% (48-81%)	3.9 (1.9-6.5)	12% (7.2-16%)	15% (8.9-21%)	350 (210-500)	25000 (16000-35000)
Cambodia	2005	Male	25 to 29	39% (22-57%)	61% (43-78%)	5.5 (2.7-9.4)	14% (9.5-18%)	17% (12-23%)	240 (160-330)	16000 (11000-21000)
Cambodia	2005	Male	30 to 34	43% (24-62%)	57% (38-76%)	6.8 (3.6-11)	14% (9.5-18%)	17% (11-22%)	390 (280-510)	24000 (17000-30000)
Cambodia	2005	Male	35 to 39	45% (27-65%)	55% (35-73%)	7.7 (3.9-13)	15% (11-19%)	18% (13-24%)	510 (380-650)	27000 (21000-35000)
Cambodia	2005	Male	40 to 44	45% (27-65%)	55% (35-73%)	8 (4-13)	15% (12-19%)	19% (14-23%)	560 (420-700)	27000 (20000-34000)
Cambodia	2005	Male	45 to 49	43% (24-63%)	57% (37-76%)	8 (3.9-13)	16% (12-20%)	20% (15-25%)	510 (380-640)	22000 (16000-27000)
Cambodia	2005	Male	50 to 54	41% (23-61%)	59% (39-77%)	7.5 (3.6-13)	16% (12-20%)	19% (15-24%)	570 (430-730)	22000 (16000-27000)
Cambodia	2005	Male	55 to 59	39% (23-59%)	61% (41-77%)	6.7 (3.3-12)	15% (11-20%)	18% (13-24%)	520 (380-680)	17000 (12000-22000)
Cambodia	2005	Male	60 to 64	36% (20-55%)	64% (45-80%)	5.8 (2.6-11)	14% (9.8-18%)	17% (12-22%)	520 (360-700)	15000 (10000-19000)
Cambodia	2005	Male	65 to 69	33% (19-51%)	67% (49-81%)	4.9 (2.2-9.2)	13% (8.8-17%)	15% (10-20%)	530 (360-730)	12000 (8400-17000)
Cambodia	2005	Male	70 to 74	29% (16-46%)	71% (54-84%)	4.1 (1.8-7.8)	12% (7.6-16%)	13% (8.9-19%)	500 (330-720)	9300 (6000-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	2005	Male	75 to 79	27% (15-43%)	73% (57-85%)	3.4 (1.4-6.9)	9.8% (6.3-14%)	11% (7.1-16%)	330 (200-480)	4700 (2900-6900)
Cambodia	2005	Male	80 to 84	24% (12-40%)	76% (60-88%)	3 (1.2-5.9)	9.3% (5.5-14%)	11% (6.3-16%)	180 (100-260)	1900 (1100-2800)
Cambodia	2005	Male	85 to 89	23% (12-37%)	77% (63-88%)	2.6 (1-5.3)	7.9% (4.5-12%)	9% (5.1-14%)	81 (46-120)	610 (350-940)
Cambodia	2005	Male	90 to 94	24% (13-40%)	76% (60-87%)	2.5 (1-5.1)	6.8% (3.7-10%)	7.8% (4.2-12%)	23 (12-34)	120 (66-180)
Cambodia	2005	Male	95 plus	27% (13-47%)	73% (53-87%)	2.6 (0.84-5.7)	6.4% (3-11%)	7.2% (3.3-12%)	4.1 (1.9-6.9)	15 (6.7-24)
Cambodia	2010	Female	All Ages	8.7% (6.2-12%)	91% (88-94%)	1.3 (0.83-2)	3.1% (2.3-4.1%)	5.7% (4.1-7.6%)	2600 (1800-3400)	77000 (59000-100000)
Cambodia	2010	Female	15 to 19	8% (3.5-16%)	92% (84-96%)	1 (0.34-2.3)	1.7% (0.91-3%)	3% (1.3-5.6%)	21 (9.2-40)	2000 (1100-3400)
Cambodia	2010	Female	20 to 24	9.6% (3.8-18%)	90% (82-96%)	1.4 (0.47-3.1)	2.7% (1.5-4.4%)	4.1% (1.9-7.1%)	31 (14-56)	2900 (1700-4700)
Cambodia	2010	Female	25 to 29	10% (4.4-20%)	90% (80-96%)	1.6 (0.52-3.6)	3.1% (1.8-4.9%)	4.9% (2.3-8%)	61 (29-100)	5000 (2800-7900)
Cambodia	2010	Female	30 to 34	11% (4.6-22%)	89% (78-95%)	1.8 (0.63-4)	3.7% (2.2-5.7%)	5.8% (3.2-9.3%)	40 (22-66)	2700 (1600-4300)
Cambodia	2010	Female	35 to 39	12% (5.2-23%)	88% (77-95%)	2 (0.66-4.3)	4.4% (2.6-6.6%)	6.9% (4.1-11%)	82 (47-130)	4900 (2900-7400)
Cambodia	2010	Female	40 to 44	13% (5.8-23%)	87% (77-94%)	2.3 (0.77-4.8)	4.9% (3.1-7.3%)	7.6% (4.8-11%)	120 (74-180)	6100 (3900-8900)
Cambodia	2010	Female	45 to 49	13% (5.2-24%)	87% (76-95%)	2.6 (0.8-5.7)	5.4% (3.2-8.2%)	8% (4.8-12%)	170 (100-260)	7500 (4600-11000)
Cambodia	2010	Female	50 to 54	13% (5.6-25%)	87% (75-94%)	2.8 (0.9-6.4)	6.3% (3.8-9.8%)	8.7% (5.3-13%)	230 (140-360)	8800 (5400-14000)
Cambodia	2010	Female	55 to 59	13% (6.2-25%)	87% (75-94%)	3.1 (0.94-7)	6.4% (3.7-10%)	8.4% (5-13%)	280 (170-450)	9400 (5600-15000)
Cambodia	2010	Female	60 to 64	13% (5.8-25%)	87% (75-94%)	3.2 (0.96-7.4)	6.3% (3.7-10%)	8% (4.8-13%)	280 (160-450)	7900 (4700-13000)
Cambodia	2010	Female	65 to 69	13% (5.8-24%)	87% (76-94%)	3 (0.94-6.5)	6.3% (3.4-10%)	7.9% (4.4-13%)	320 (180-520)	7400 (4100-12000)
Cambodia	2010	Female	70 to 74	12% (5.5-22%)	88% (78-95%)	2.7 (0.78-6.5)	5.8% (3.1-9.6%)	7.1% (3.8-12%)	310 (160-520)	5800 (3100-9700)
Cambodia	2010	Female	75 to 79	12% (5.1-22%)	88% (78-95%)	2.5 (0.66-6.2)	5.1% (2.6-8.5%)	6.1% (3.1-10%)	270 (140-440)	3900 (2000-6300)
Cambodia	2010	Female	80 to 84	12% (5.6-23%)	88% (77-94%)	2.5 (0.75-5.9)	5.2% (2.7-8.6%)	6.2% (3.2-10%)	190 (100-320)	2000 (1100-3400)
Cambodia	2010	Female	85 to 89	13% (5.6-24%)	87% (76-94%)	2.7 (0.79-6.7)	5% (2.5-8.7%)	5.9% (2.9-10%)	100 (50-180)	760 (380-1300)
Cambodia	2010	Female	90 to 94	15% (6.4-28%)	85% (72-94%)	3.2 (0.75-8.3)	4.8% (2.3-8.4%)	5.7% (2.7-9.9%)	37 (18-65)	190 (92-330)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	2010	Female	95 plus	19% (7.2-36%)	81% (64-93%)	3.9 (0.75-11)	5.1% (2.2-9.3%)	5.9% (2.5-11%)	9.5 (4-18)	33 (14-60)
Cambodia	2010	Male	All Ages	26% (20-33%)	74% (67-80%)	3.7 (2.6-5.1)	8.3% (6.8-9.7%)	12% (10-14%)	6400 (5200-7600)	250000 (200000-290000)
Cambodia	2010	Male	15 to 19	22% (12-37%)	78% (63-88%)	1.7 (0.72-3.3)	4.7% (2.4-7.8%)	6.4% (3-11%)	98 (46-170)	7900 (4000-13000)
Cambodia	2010	Male	20 to 24	34% (19-51%)	66% (49-81%)	4.2 (2-7.2)	12% (7.4-16%)	15% (9-21%)	270 (160-390)	19000 (12000-28000)
Cambodia	2010	Male	25 to 29	39% (22-58%)	61% (42-78%)	5.9 (2.9-10)	14% (9.8-18%)	18% (12-23%)	510 (340-690)	34000 (23000-45000)
Cambodia	2010	Male	30 to 34	42% (23-62%)	58% (38-77%)	7.4 (3.8-13)	15% (11-19%)	19% (13-24%)	270 (190-350)	16000 (12000-21000)
Cambodia	2010	Male	35 to 39	45% (27-63%)	55% (37-73%)	8.4 (4.2-15)	15% (12-19%)	19% (14-24%)	470 (350-590)	25000 (19000-32000)
Cambodia	2010	Male	40 to 44	44% (27-64%)	56% (36-73%)	8.8 (4.3-15)	16% (12-19%)	19% (15-24%)	550 (420-690)	27000 (21000-34000)
Cambodia	2010	Male	45 to 49	43% (25-62%)	57% (38-75%)	8.7 (4.2-15)	16% (12-20%)	20% (15-25%)	660 (500-840)	29000 (21000-36000)
Cambodia	2010	Male	50 to 54	41% (22-60%)	59% (40-78%)	8.2 (3.9-14)	16% (12-20%)	20% (15-24%)	540 (400-680)	21000 (15000-26000)
Cambodia	2010	Male	55 to 59	38% (23-58%)	62% (42-77%)	7.4 (3.5-14)	15% (11-19%)	18% (13-24%)	610 (440-800)	20000 (15000-26000)
Cambodia	2010	Male	60 to 64	36% (20-56%)	64% (44-80%)	6.4 (2.8-12)	14% (9.7-18%)	16% (12-22%)	560 (390-750)	16000 (11000-21000)
Cambodia	2010	Male	65 to 69	32% (18-50%)	68% (50-82%)	5.4 (2.4-10)	12% (8.7-17%)	15% (10-20%)	530 (370-740)	12000 (8600-17000)
Cambodia	2010	Male	70 to 74	29% (16-45%)	71% (55-84%)	4.4 (1.9-8.8)	11% (7.5-16%)	13% (8.8-18%)	520 (350-720)	9700 (6500-13000)
Cambodia	2010	Male	75 to 79	26% (15-42%)	74% (58-85%)	3.8 (1.5-7.5)	9.6% (6.3-14%)	11% (7.2-16%)	400 (250-560)	5700 (3600-8100)
Cambodia	2010	Male	80 to 84	24% (12-40%)	76% (60-88%)	3.2 (1.3-6.5)	9.2% (5.6-13%)	11% (6.4-15%)	230 (140-340)	2500 (1400-3600)
Cambodia	2010	Male	85 to 89	22% (12-36%)	78% (64-88%)	2.8 (1.1-5.8)	7.8% (4.6-11%)	8.9% (5.2-13%)	100 (59-160)	790 (450-1200)
Cambodia	2010	Male	90 to 94	24% (13-39%)	76% (61-87%)	2.7 (1.1-5.9)	6.7% (3.9-10%)	7.7% (4.5-12%)	32 (18-49)	170 (97-260)
Cambodia	2010	Male	95 plus	27% (12-47%)	73% (53-88%)	2.8 (0.92-6.4)	6.6% (3.2-11%)	7.3% (3.4-12%)	6.6 (3.1-11)	23 (11-38)
Cambodia	2016	Female	All Ages	8.8% (6.3-12%)	91% (88-94%)	1.4 (0.79-2.2)	3.3% (2.4-4.3%)	5.8% (4.1-7.7%)	2600 (1900-3500)	77000 (58000-100000)
Cambodia	2016	Female	15 to 19	7.9% (3.5-15%)	92% (85-96%)	1 (0.32-2.4)	1.5% (0.8-2.5%)	2.7% (1.2-4.8%)	14 (5.9-26)	1500 (770-2400)
Cambodia	2016	Female	20 to 24	9.4% (3.9-18%)	91% (82-96%)	1.4 (0.45-3.2)	2.3% (1.4-3.7%)	3.6% (1.7-6.1%)	25 (12-44)	2700 (1600-4200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	2016	Female	25 to 29	10% (4.4-19%)	90% (81-96%)	1.6 (0.53-3.5)	2.8% (1.7-4.3%)	4.5% (2.3-7.6%)	33 (16-55)	2900 (1700-4500)
Cambodia	2016	Female	30 to 34	11% (4.7-21%)	89% (79-95%)	1.8 (0.58-4.1)	3.3% (2-5.1%)	5.5% (3.1-8.5%)	73 (40-120)	5400 (3300-8100)
Cambodia	2016	Female	35 to 39	12% (5.4-22%)	88% (78-95%)	2 (0.71-4.5)	4% (2.5-6%)	6.7% (4-9.9%)	62 (36-94)	3800 (2400-5600)
Cambodia	2016	Female	40 to 44	12% (5.7-22%)	88% (78-94%)	2.3 (0.75-4.9)	4.6% (2.9-6.7%)	7.4% (4.7-11%)	98 (61-150)	5100 (3300-7500)
Cambodia	2016	Female	45 to 49	13% (5.6-24%)	87% (76-94%)	2.6 (0.8-6.1)	5% (3-7.6%)	7.7% (4.6-12%)	150 (91-230)	6900 (4100-10000)
Cambodia	2016	Female	50 to 54	13% (5.6-25%)	87% (75-94%)	2.9 (0.87-6.4)	5.8% (3.5-8.9%)	8.3% (5.1-13%)	230 (140-370)	9100 (5500-14000)
Cambodia	2016	Female	55 to 59	13% (5.9-25%)	87% (75-94%)	3.1 (0.94-7.3)	5.9% (3.4-9.4%)	8.1% (4.7-13%)	280 (160-440)	9200 (5200-15000)
Cambodia	2016	Female	60 to 64	13% (5.6-24%)	87% (76-94%)	3.2 (0.92-7.3)	5.8% (3.5-9.2%)	7.6% (4.6-12%)	320 (190-520)	9200 (5500-15000)
Cambodia	2016	Female	65 to 69	13% (5.7-24%)	87% (76-94%)	3 (0.87-6.8)	5.8% (3.2-9.2%)	7.4% (4.1-12%)	330 (190-540)	7800 (4400-13000)
Cambodia	2016	Female	70 to 74	12% (5.9-22%)	88% (78-94%)	2.7 (0.77-6.9)	5.4% (2.9-8.8%)	6.8% (3.7-11%)	320 (170-530)	6000 (3200-9900)
Cambodia	2016	Female	75 to 79	12% (5.2-23%)	88% (77-95%)	2.6 (0.68-6.6)	4.8% (2.5-8.1%)	5.9% (3-9.8%)	270 (140-460)	3900 (2000-6600)
Cambodia	2016	Female	80 to 84	12% (5.2-23%)	88% (77-95%)	2.6 (0.74-6.1)	4.9% (2.5-8.1%)	5.9% (3-9.7%)	220 (110-370)	2300 (1200-3900)
Cambodia	2016	Female	85 to 89	13% (5.7-24%)	87% (76-94%)	2.8 (0.76-7.1)	4.7% (2.3-8.3%)	5.6% (2.7-9.9%)	140 (69-250)	1100 (520-1900)
Cambodia	2016	Female	90 to 94	15% (6.3-27%)	85% (73-94%)	3.2 (0.75-8.7)	4.4% (2.1-7.8%)	5.3% (2.5-9.3%)	49 (23-86)	250 (120-440)
Cambodia	2016	Female	95 plus	18% (7-35%)	82% (65-93%)	3.8 (0.71-11)	4.7% (2-8.6%)	5.4% (2.3-10%)	13 (5.5-24)	44 (19-82)
Cambodia	2016	Male	All Ages	27% (21-34%)	73% (66-79%)	4 (2.5-6.1)	9% (7.4-11%)	13% (10-15%)	6700 (5400-8000)	25000 (21000-30000)
Cambodia	2016	Male	15 to 19	22% (12-35%)	78% (65-88%)	1.7 (0.67-3.5)	4.5% (2.3-7.7%)	6.3% (2.9-11%)	81 (36-140)	6600 (3200-11000)
Cambodia	2016	Male	20 to 24	33% (18-51%)	67% (49-82%)	4.2 (1.9-7.7)	11% (6.8-15%)	14% (8.4-20%)	300 (180-450)	22000 (13000-32000)
Cambodia	2016	Male	25 to 29	39% (21-57%)	61% (43-79%)	5.9 (2.8-11)	14% (9.7-18%)	18% (12-23%)	320 (210-450)	22000 (15000-29000)
Cambodia	2016	Male	30 to 34	42% (23-62%)	58% (38-77%)	7.3 (3.5-13)	14% (11-19%)	19% (14-25%)	550 (380-740)	34000 (24000-45000)
Cambodia	2016	Male	35 to 39	44% (27-63%)	56% (37-73%)	8.4 (3.8-16)	16% (12-20%)	20% (15-26%)	350 (260-450)	19000 (14000-24000)
Cambodia	2016	Male	40 to 44	44% (27-64%)	56% (36-73%)	8.8 (3.9-16)	15% (12-19%)	20% (16-25%)	450 (330-570)	22000 (16000-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cambodia	2016	Male	45 to 49	42% (24-61%)	58% (39-76%)	8.6 (3.8-16)	16% (12-20%)	20% (15-25%)	610 (460-780)	27000 (20000-34000)
Cambodia	2016	Male	50 to 54	40% (23-61%)	60% (39-77%)	8.1 (3.5-15)	15% (11-19%)	19% (14-24%)	730 (530-930)	28000 (20000-35000)
Cambodia	2016	Male	55 to 59	38% (22-59%)	62% (41-78%)	7.4 (3.3-14)	14% (10-18%)	17% (13-22%)	610 (440-810)	20000 (15000-27000)
Cambodia	2016	Male	60 to 64	35% (19-55%)	65% (45-81%)	6.4 (2.7-13)	13% (9.1-17%)	16% (11-20%)	650 (460-870)	18000 (13000-24000)
Cambodia	2016	Male	65 to 69	32% (18-51%)	68% (49-82%)	5.3 (2.2-11)	12% (8.1-16%)	14% (9.7-19%)	600 (420-810)	14000 (9600-19000)
Cambodia	2016	Male	70 to 74	28% (16-45%)	72% (55-84%)	4.4 (1.8-9)	10% (6.8-14%)	12% (7.9-17%)	520 (340-720)	9600 (6300-13000)
Cambodia	2016	Male	75 to 79	26% (14-42%)	74% (58-86%)	3.7 (1.4-7.8)	8.8% (5.6-12%)	10% (6.4-14%)	420 (260-590)	6000 (3800-8500)
Cambodia	2016	Male	80 to 84	24% (12-40%)	76% (60-88%)	3.2 (1.2-6.7)	8.3% (5-12%)	9.5% (5.8-14%)	280 (170-410)	3000 (1800-4400)
Cambodia	2016	Male	85 to 89	22% (11-37%)	78% (63-89%)	2.8 (1.1-6.1)	7% (4-10%)	8% (4.6-12%)	130 (75-200)	1000 (570-1500)
Cambodia	2016	Male	90 to 94	23% (12-38%)	77% (62-88%)	2.7 (1-5.7)	6.1% (3.3-9.3%)	7% (3.7-11%)	40 (22-62)	210 (120-320)
Cambodia	2016	Male	95 plus	27% (12-45%)	73% (55-88%)	2.8 (0.88-6.9)	6% (2.8-9.6%)	6.6% (3-11%)	8.8 (4.1-15)	31 (15-51)
Indonesia	1990	Female	All Ages	2.5% (2-3.1%)	98% (97-98%)	0.11 (0.1-0.12)	0.76% (0.61-0.93%)	1.3% (0.92-1.7%)	8400 (6200-11000)	290000 (230000-350000)
Indonesia	1990	Female	15 to 19	2.3% (1.3-3.8%)	98% (96-99%)	0.092 (0.046-0.16)	0.54% (0.35-0.83%)	0.61% (0.32-1.1%)	80 (41-140)	9000 (5700-14000)
Indonesia	1990	Female	20 to 24	2.8% (1.5-4.7%)	97% (95-99%)	0.12 (0.06-0.21)	0.96% (0.62-1.4%)	0.92% (0.46-1.5%)	150 (76-250)	17000 (11000-25000)
Indonesia	1990	Female	25 to 29	3.1% (1.7-4.9%)	97% (95-98%)	0.14 (0.073-0.23)	1.1% (0.78-1.6%)	1.2% (0.66-1.9%)	190 (100-300)	19000 (12000-27000)
Indonesia	1990	Female	30 to 34	3.4% (1.8-5.5%)	97% (94-98%)	0.16 (0.084-0.26)	1.2% (0.88-1.8%)	1.4% (0.92-2.2%)	240 (150-360)	19000 (13000-28000)
Indonesia	1990	Female	35 to 39	3.7% (1.9-5.9%)	96% (94-98%)	0.19 (0.096-0.3)	1.5% (1.1-2.2%)	2% (1.3-3%)	330 (210-480)	21000 (14000-29000)
Indonesia	1990	Female	40 to 44	3.7% (2.1-6.1%)	96% (94-98%)	0.22 (0.11-0.36)	1.7% (1.2-2.4%)	2.3% (1.6-3.4%)	400 (270-590)	22000 (15000-31000)
Indonesia	1990	Female	45 to 49	3.7% (1.9-5.8%)	96% (94-98%)	0.25 (0.12-0.42)	1.9% (1.3-2.8%)	2.5% (1.7-3.8%)	570 (370-860)	26000 (18000-39000)
Indonesia	1990	Female	50 to 54	3.7% (2.1-6.2%)	96% (94-98%)	0.25 (0.13-0.43)	2.1% (1.4-3%)	2.7% (1.8-4%)	830 (550-1200)	32000 (22000-48000)
Indonesia	1990	Female	55 to 59	3.7% (2.1-6%)	96% (94-98%)	0.24 (0.12-0.41)	1.8% (1.2-2.8%)	2.4% (1.5-3.6%)	900 (570-1400)	30000 (19000-45000)
Indonesia	1990	Female	60 to 64	3.7% (2-6.3%)	96% (94-98%)	0.23 (0.11-0.43)	1.8% (1.1-2.8%)	2.2% (1.3-3.5%)	1000 (600-1600)	28000 (17000-44000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	1990	Female	65 to 69	3.7% (2.1-5.8%)	96% (94-98%)	0.22 (0.11-0.39)	1.6% (0.95-2.6%)	2% (1.1-3.1%)	970 (580-1500)	22000 (13000-36000)
Indonesia	1990	Female	70 to 74	3.6% (2.1-5.8%)	96% (94-98%)	0.21 (0.098-0.37)	1.5% (0.82-2.4%)	1.7% (0.96-2.8%)	960 (540-1600)	18000 (10000-29000)
Indonesia	1990	Female	75 to 79	3.5% (2-5.9%)	96% (94-98%)	0.2 (0.087-0.37)	1.2% (0.68-2.1%)	1.4% (0.79-2.4%)	810 (440-1300)	12000 (6400-19000)
Indonesia	1990	Female	80 to 84	3.5% (1.9-5.8%)	96% (94-98%)	0.19 (0.091-0.34)	1.2% (0.65-2%)	1.4% (0.75-2.3%)	570 (300-930)	6000 (3200-9700)
Indonesia	1990	Female	85 to 89	3.8% (2.1-6.4%)	96% (94-98%)	0.2 (0.093-0.37)	1.1% (0.55-1.9%)	1.3% (0.65-2.2%)	300 (150-520)	2200 (1200-3900)
Indonesia	1990	Female	90 to 94	4.4% (2.5-7.2%)	96% (93-97%)	0.23 (0.1-0.43)	0.96% (0.46-1.8%)	1.1% (0.54-2.1%)	96 (46-180)	500 (240-900)
Indonesia	1990	Female	95 plus	5.3% (2.6-9.3%)	95% (91-97%)	0.26 (0.098-0.52)	1% (0.42-1.9%)	1.1% (0.46-2.2%)	24 (9.9-46)	84 (36-160)
Indonesia	1990	Male	All Ages	6.6% (5.4-8.1%)	93% (92-95%)	0.061 (0.055-0.068)	0.64% (0.44-0.81%)	0.87% (0.56-1.2%)	6200 (4000-8400)	270000 (190000-350000)
Indonesia	1990	Male	15 to 19	5.6% (3.3-9%)	94% (91-97%)	0.027 (0.013-0.047)	0.44% (0.3-0.61%)	0.32% (0.11-0.55%)	47 (16-81)	7600 (4800-11000)
Indonesia	1990	Male	20 to 24	8.8% (4.9-14%)	91% (86-95%)	0.068 (0.037-0.11)	0.64% (0.55-1.4%)	0.64% (0.18-1.2%)	130 (36-240)	19000 (11000-28000)
Indonesia	1990	Male	25 to 29	10% (5.8-16%)	90% (84-94%)	0.1 (0.054-0.16)	1.2% (0.72-1.8%)	1% (0.3-1.8%)	210 (60-360)	23000 (13000-35000)
Indonesia	1990	Male	30 to 34	11% (6.3-18%)	89% (82-94%)	0.13 (0.073-0.2)	1.7% (1-2.3%)	1.7% (0.84-2.6%)	360 (180-540)	29000 (17000-40000)
Indonesia	1990	Male	35 to 39	12% (6.9-19%)	88% (81-93%)	0.15 (0.082-0.24)	2.1% (1.4-2.9%)	2.5% (1.4-3.6%)	510 (290-750)	32000 (20000-44000)
Indonesia	1990	Male	40 to 44	12% (7-19%)	88% (81-93%)	0.15 (0.08-0.26)	2.3% (1.5-3%)	2.8% (1.8-3.9%)	570 (370-790)	30000 (20000-40000)
Indonesia	1990	Male	45 to 49	11% (6-17%)	89% (83-94%)	0.14 (0.076-0.24)	2.2% (1.5-3%)	2.8% (1.8-3.8%)	720 (460-990)	32000 (22000-43000)
Indonesia	1990	Male	50 to 54	10% (6.3-16%)	90% (84-94%)	0.13 (0.066-0.21)	1.9% (1.3-2.6%)	2.4% (1.6-3.3%)	820 (530-1100)	31000 (21000-42000)
Indonesia	1990	Male	55 to 59	9.8% (5.8-15%)	90% (85-94%)	0.11 (0.056-0.18)	1.5% (0.97-2%)	1.8% (1.1-2.6%)	730 (450-1000)	24000 (15000-33000)
Indonesia	1990	Male	60 to 64	9.2% (5.4-15%)	91% (85-95%)	0.091 (0.044-0.16)	1.2% (0.7-1.7%)	1.5% (0.81-2.1%)	690 (390-1000)	19000 (11000-28000)
Indonesia	1990	Male	65 to 69	8.6% (5.2-13%)	91% (87-95%)	0.08 (0.042-0.13)	0.88% (0.48-1.3%)	1% (0.54-1.5%)	510 (270-760)	12000 (6600-17000)
Indonesia	1990	Male	70 to 74	7.5% (4.7-12%)	92% (88-95%)	0.066 (0.033-0.12)	0.68% (0.35-1%)	0.78% (0.38-1.2%)	390 (190-600)	7300 (3800-11000)
Indonesia	1990	Male	75 to 79	6.5% (3.8-10%)	94% (90-96%)	0.054 (0.024-0.097)	0.57% (0.27-0.84%)	0.65% (0.29-0.99%)	300 (140-460)	4400 (2100-6500)
Indonesia	1990	Male	80 to 84	5.8% (3.3-9%)	94% (91-97%)	0.046 (0.022-0.08)	0.46% (0.2-0.69%)	0.52% (0.21-0.8%)	150 (61-230)	1600 (710-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	1990	Male	85 to 89	5.6% (3.2-8.9%)	94% (91-97%)	0.043 (0.021-0.075)	0.33% (0.13-0.52%)	0.37% (0.13-0.6%)	63 (22-100)	480 (190-760)
Indonesia	1990	Male	90 to 94	6% (3.6-9.6%)	94% (90-96%)	0.043 (0.02-0.08)	0.25% (0.083-0.42%)	0.28% (0.072-0.48%)	17 (4.5-30)	91 (30-160)
Indonesia	1990	Male	95 plus	7.1% (3.8-12%)	93% (88-96%)	0.048 (0.019-0.092)	0.22% (0.048-0.4%)	0.23% (0.021-0.44%)	3.5 (0.33-6.8)	14 (2.9-24)
Indonesia	1995	Female	All Ages	2.2% (1.8-2.8%)	98% (97-98%)	0.11 (0.1-0.12)	0.84% (0.69-1%)	1.4% (1-1.8%)	9400 (7000-13000)	300000 (250000-380000)
Indonesia	1995	Female	15 to 19	2.2% (1.2-3.6%)	98% (96-99%)	0.1 (0.05-0.18)	0.57% (0.37-0.84%)	0.66% (0.35-1.1%)	75 (40-130)	9000 (5800-13000)
Indonesia	1995	Female	20 to 24	2.5% (1.3-4.2%)	97% (96-99%)	0.13 (0.061-0.22)	0.98% (0.66-1.4%)	0.95% (0.51-1.5%)	130 (72-220)	17000 (11000-25000)
Indonesia	1995	Female	25 to 29	2.6% (1.4-4.3%)	97% (96-99%)	0.14 (0.072-0.22)	1.1% (0.77-1.5%)	1.2% (0.67-1.8%)	170 (100-270)	18000 (13000-26000)
Indonesia	1995	Female	30 to 34	2.7% (1.4-4.5%)	97% (96-99%)	0.15 (0.076-0.24)	1.2% (0.85-1.6%)	1.4% (0.9-2%)	220 (140-320)	19000 (14000-26000)
Indonesia	1995	Female	35 to 39	2.8% (1.4-4.6%)	97% (95-99%)	0.16 (0.082-0.27)	1.4% (1-1.9%)	1.9% (1.2-2.7%)	330 (220-480)	22000 (16000-30000)
Indonesia	1995	Female	40 to 44	2.9% (1.6-4.9%)	97% (95-98%)	0.2 (0.1-0.33)	1.6% (1.1-2.2%)	2.2% (1.5-3.1%)	410 (280-580)	23000 (16000-31000)
Indonesia	1995	Female	45 to 49	3.1% (1.5-4.9%)	97% (95-98%)	0.24 (0.12-0.42)	1.8% (1.2-2.6%)	2.4% (1.6-3.6%)	530 (350-790)	25000 (17000-36000)
Indonesia	1995	Female	50 to 54	3.1% (1.7-5.2%)	97% (95-98%)	0.25 (0.13-0.43)	2% (1.3-2.9%)	2.6% (1.8-3.8%)	800 (540-1200)	32000 (22000-46000)
Indonesia	1995	Female	55 to 59	3.1% (1.7-5.1%)	97% (95-98%)	0.24 (0.11-0.41)	1.8% (1.2-2.6%)	2.3% (1.5-3.4%)	950 (620-1400)	32000 (21000-47000)
Indonesia	1995	Female	60 to 64	3.2% (1.8-5.4%)	97% (95-98%)	0.23 (0.11-0.43)	1.7% (1.1-2.7%)	2.2% (1.4-3.4%)	1100 (720-1700)	32000 (20000-49000)
Indonesia	1995	Female	65 to 69	3.3% (1.9-5.2%)	97% (95-98%)	0.23 (0.11-0.41)	1.7% (1-2.6%)	2% (1.2-3.1%)	1200 (750-1900)	28000 (17000-44000)
Indonesia	1995	Female	70 to 74	3.4% (2-5.4%)	97% (95-98%)	0.23 (0.11-0.4)	1.5% (0.9-2.5%)	1.8% (1.1-2.9%)	1100 (670-1900)	21000 (12000-35000)
Indonesia	1995	Female	75 to 79	3.4% (1.9-5.8%)	97% (94-98%)	0.23 (0.1-0.42)	1.4% (0.77-2.2%)	1.6% (0.9-2.6%)	970 (540-1600)	14000 (7700-23000)
Indonesia	1995	Female	80 to 84	3.5% (1.9-5.8%)	96% (94-98%)	0.23 (0.1-0.4)	1.3% (0.74-2.2%)	1.6% (0.87-2.6%)	730 (400-1200)	7600 (4200-12000)
Indonesia	1995	Female	85 to 89	3.8% (2.1-6.5%)	96% (94-98%)	0.24 (0.11-0.43)	1.2% (0.63-2.1%)	1.4% (0.74-2.4%)	430 (230-730)	3200 (1700-5500)
Indonesia	1995	Female	90 to 94	4.4% (2.5-7%)	96% (93-98%)	0.26 (0.11-0.5)	1.1% (0.55-1.9%)	1.3% (0.64-2.3%)	150 (76-280)	790 (400-1400)
Indonesia	1995	Female	95 plus	5.3% (2.6-9.2%)	95% (91-97%)	0.3 (0.11-0.6)	1.1% (0.49-1.9%)	1.3% (0.56-2.3%)	39 (17-68)	130 (59-240)
Indonesia	1995	Male	All Ages	7.5% (6-9.3%)	93% (91-94%)	0.078 (0.07-0.086)	0.81% (0.61-0.99%)	1.1% (0.76-1.4%)	7700 (5300-9700)	330000 (250000-410000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	1995	Male	15 to 19	8% (4.8-13%)	92% (87-95%)	0.045 (0.022-0.078)	0.5% (0.33-0.72%)	0.42% (0.16-0.72%)	61 (22-100)	8700 (5400-13000)
Indonesia	1995	Male	20 to 24	13% (7.4-19%)	87% (81-93%)	0.11 (0.063-0.18)	1.1% (0.62-1.7%)	0.9% (0.27-1.6%)	180 (51-320)	22000 (12000-34000)
Indonesia	1995	Male	25 to 29	13% (7.5-19%)	87% (81-92%)	0.15 (0.081-0.23)	1.5% (0.83-2.2%)	1.4% (0.47-2.3%)	280 (96-480)	28000 (15000-42000)
Indonesia	1995	Male	30 to 34	12% (6.9-19%)	88% (81-93%)	0.16 (0.092-0.25)	1.9% (1.3-2.6%)	2.1% (1.2-3%)	450 (250-650)	35000 (23000-47000)
Indonesia	1995	Male	35 to 39	12% (6.8-18%)	88% (82-93%)	0.17 (0.095-0.28)	2.3% (1.6-3.1%)	2.9% (1.9-4%)	680 (440-950)	42000 (29000-56000)
Indonesia	1995	Male	40 to 44	11% (6.3-17%)	89% (83-94%)	0.16 (0.083-0.27)	2.5% (1.8-3.2%)	3.2% (2.2-4.2%)	760 (530-1000)	39000 (28000-51000)
Indonesia	1995	Male	45 to 49	9.3% (4.9-15%)	91% (85-95%)	0.14 (0.071-0.24)	2.4% (1.7-3.1%)	3.1% (2.2-4.1%)	790 (550-1000)	35000 (25000-45000)
Indonesia	1995	Male	50 to 54	8.2% (4.9-13%)	92% (87-95%)	0.12 (0.059-0.2)	2.1% (1.5-2.7%)	2.7% (1.9-3.4%)	870 (630-1200)	34000 (24000-43000)
Indonesia	1995	Male	55 to 59	7.5% (4.4-12%)	93% (88-96%)	0.094 (0.047-0.16)	1.6% (1.2-2.1%)	2% (1.4-2.7%)	870 (600-1100)	29000 (20000-38000)
Indonesia	1995	Male	60 to 64	7.3% (4.2-12%)	93% (88-96%)	0.083 (0.04-0.15)	1.3% (0.88-1.8%)	1.6% (1-2.2%)	850 (550-1200)	24000 (16000-33000)
Indonesia	1995	Male	65 to 69	7.3% (4.3-11%)	93% (89-96%)	0.078 (0.039-0.13)	1% (0.64-1.4%)	1.2% (0.73-1.6%)	680 (420-950)	16000 (10000-22000)
Indonesia	1995	Male	70 to 74	6.7% (4.1-10%)	93% (90-96%)	0.067 (0.034-0.12)	0.78% (0.48-1.1%)	0.91% (0.54-1.3%)	510 (310-750)	9600 (5800-14000)
Indonesia	1995	Male	75 to 79	5.9% (3.5-9.5%)	94% (90-97%)	0.056 (0.025-0.1)	0.66% (0.36-0.92%)	0.76% (0.39-1.1%)	360 (190-520)	5200 (2900-7400)
Indonesia	1995	Male	80 to 84	5.4% (3.1-8.3%)	95% (92-97%)	0.049 (0.023-0.085)	0.53% (0.26-0.76%)	0.61% (0.29-0.89%)	200 (93-290)	2100 (1000-3000)
Indonesia	1995	Male	85 to 89	5.5% (3-8.6%)	95% (91-97%)	0.048 (0.023-0.084)	0.38% (0.18-0.56%)	0.43% (0.19-0.66%)	91 (40-140)	690 (330-1000)
Indonesia	1995	Male	90 to 94	6% (3.5-9.3%)	94% (91-96%)	0.049 (0.023-0.094)	0.28% (0.12-0.45%)	0.32% (0.11-0.52%)	26 (9.4-43)	140 (56-220)
Indonesia	1995	Male	95 plus	7% (3.6-11%)	93% (89-96%)	0.053 (0.021-0.1)	0.27% (0.092-0.44%)	0.28% (0.065-0.49%)	5.3 (1.3-9.5)	20 (7.1-34)
Indonesia	2000	Female	All Ages	2.3% (1.9-2.8%)	98% (97-98%)	0.12 (0.12-0.13)	0.99% (0.81-1.2%)	1.6% (1.2-2.1%)	12000 (8600-15000)	360000 (290000-440000)
Indonesia	2000	Female	15 to 19	2.2% (1.2-3.6%)	98% (96-99%)	0.11 (0.053-0.19)	0.59% (0.38-0.86%)	0.73% (0.39-1.2%)	73 (39-120)	9000 (5900-13000)
Indonesia	2000	Female	20 to 24	2.6% (1.3-4.3%)	97% (96-99%)	0.14 (0.066-0.24)	1% (0.68-1.4%)	1% (0.56-1.7%)	130 (70-220)	17000 (11000-24000)
Indonesia	2000	Female	25 to 29	2.7% (1.5-4.4%)	97% (96-99%)	0.15 (0.076-0.24)	1.2% (0.82-1.6%)	1.3% (0.73-2%)	180 (100-270)	19000 (13000-27000)
Indonesia	2000	Female	30 to 34	2.7% (1.4-4.5%)	97% (96-99%)	0.16 (0.082-0.26)	1.2% (0.9-1.7%)	1.5% (0.97-2.2%)	240 (160-360)	21000 (16000-29000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	2000	Female	35 to 39	2.8% (1.4-4.6%)	97% (95-99%)	0.18 (0.092-0.29)	1.4% (1-1.9%)	1.9% (1.3-2.8%)	370 (250-540)	25000 (18000-34000)
Indonesia	2000	Female	40 to 44	3% (1.6-4.9%)	97% (95-98%)	0.21 (0.11-0.35)	1.6% (1.2-2.2%)	2.3% (1.6-3.2%)	520 (370-730)	29000 (21000-39000)
Indonesia	2000	Female	45 to 49	3.1% (1.5-4.9%)	97% (95-98%)	0.26 (0.13-0.45)	1.8% (1.3-2.6%)	2.5% (1.7-3.7%)	640 (440-940)	30000 (21000-43000)
Indonesia	2000	Female	50 to 54	3.2% (1.8-5.3%)	97% (95-98%)	0.27 (0.13-0.47)	2% (1.4-2.9%)	2.7% (1.8-3.9%)	850 (580-1200)	33000 (23000-48000)
Indonesia	2000	Female	55 to 59	3.2% (1.8-5.2%)	97% (95-98%)	0.26 (0.12-0.45)	1.9% (1.3-2.7%)	2.4% (1.6-3.6%)	1100 (700-1500)	35000 (24000-51000)
Indonesia	2000	Female	60 to 64	3.3% (1.7-5.6%)	97% (94-98%)	0.25 (0.12-0.47)	1.8% (1.2-2.8%)	2.3% (1.5-3.5%)	1300 (860-2000)	38000 (24000-58000)
Indonesia	2000	Female	65 to 69	3.4% (1.8-5.4%)	97% (95-98%)	0.25 (0.12-0.44)	1.8% (1.1-2.7%)	2.1% (1.3-3.3%)	1500 (960-2400)	36000 (22000-55000)
Indonesia	2000	Female	70 to 74	3.4% (2-5.5%)	97% (95-98%)	0.25 (0.12-0.45)	1.6% (0.98-2.6%)	2% (1.2-3.1%)	1500 (920-2400)	29000 (17000-45000)
Indonesia	2000	Female	75 to 79	3.5% (1.9-5.9%)	97% (94-98%)	0.24 (0.11-0.45)	1.5% (0.83-2.4%)	1.7% (0.97-2.8%)	1300 (720-2100)	18000 (10000-29000)
Indonesia	2000	Female	80 to 84	3.6% (1.9-6.1%)	96% (94-98%)	0.25 (0.11-0.44)	1.4% (0.82-2.3%)	1.7% (0.96-2.7%)	950 (540-1500)	9900 (5600-16000)
Indonesia	2000	Female	85 to 89	3.9% (2.2-6.6%)	96% (93-98%)	0.26 (0.12-0.47)	1.3% (0.71-2.2%)	1.5% (0.84-2.6%)	550 (300-910)	4100 (2300-6700)
Indonesia	2000	Female	90 to 94	4.5% (2.6-7.2%)	95% (93-97%)	0.29 (0.13-0.55)	1.2% (0.63-2%)	1.4% (0.74-2.4%)	210 (110-370)	1100 (580-1900)
Indonesia	2000	Female	95 plus	5.4% (2.7-9.7%)	95% (90-97%)	0.33 (0.13-0.66)	1.2% (0.57-2.1%)	1.4% (0.64-2.5%)	58 (27-110)	200 (97-360)
Indonesia	2000	Male	All Ages	7.7% (6.2-9.6%)	92% (90-94%)	0.085 (0.075-0.094)	0.97% (0.75-1.2%)	1.2% (0.92-1.5%)	9100 (6700-11000)	390000 (300000-470000)
Indonesia	2000	Male	15 to 19	8% (4.8-12%)	92% (88-95%)	0.046 (0.023-0.081)	0.53% (0.35-0.74%)	0.46% (0.19-0.77%)	68 (28-110)	9500 (6100-14000)
Indonesia	2000	Male	20 to 24	13% (7.3-19%)	87% (81-93%)	0.12 (0.065-0.19)	1.2% (0.69-1.8%)	0.99% (0.35-1.7%)	200 (76-360)	25000 (14000-37000)
Indonesia	2000	Male	25 to 29	13% (7.4-19%)	87% (81-93%)	0.15 (0.082-0.24)	1.5% (0.94-2.2%)	1.5% (0.58-2.4%)	320 (130-530)	31000 (18000-46000)
Indonesia	2000	Male	30 to 34	12% (6.8-19%)	88% (81-93%)	0.17 (0.094-0.26)	2% (1.4-2.6%)	2.2% (1.4-3.1%)	530 (330-750)	41000 (28000-54000)
Indonesia	2000	Male	35 to 39	12% (7-18%)	88% (82-93%)	0.18 (0.098-0.28)	2.5% (1.8-3.2%)	3.1% (2.1-4.1%)	810 (550-1100)	49000 (35000-65000)
Indonesia	2000	Male	40 to 44	11% (6.4-17%)	89% (83-94%)	0.17 (0.088-0.27)	2.6% (2-3.3%)	3.4% (2.5-4.4%)	990 (730-1300)	51000 (39000-66000)
Indonesia	2000	Male	45 to 49	9.3% (5-14%)	91% (86-95%)	0.14 (0.074-0.24)	2.5% (1.9-3.3%)	3.3% (2.3-4.3%)	1000 (750-1400)	46000 (34000-60000)
Indonesia	2000	Male	50 to 54	8.1% (4.8-13%)	92% (87-95%)	0.12 (0.06-0.21)	2.2% (1.6-2.8%)	2.8% (2.1-3.6%)	960 (700-1200)	37000 (27000-46000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	2000	Male	55 to 59	7.5% (4.4-11%)	93% (89-96%)	0.097 (0.049-0.17)	1.7% (1.3-2.2%)	2.2% (1.5-2.8%)	940 (680-1200)	31000 (23000-40000)
Indonesia	2000	Male	60 to 64	7.3% (4.2-12%)	93% (88-96%)	0.086 (0.041-0.16)	1.4% (0.97-1.9%)	1.7% (1.1-2.3%)	990 (660-1300)	28000 (19000-37000)
Indonesia	2000	Male	65 to 69	7.3% (4.3-11%)	93% (89-96%)	0.08 (0.042-0.13)	1% (0.68-1.4%)	1.3% (0.78-1.7%)	800 (510-1100)	19000 (12000-25000)
Indonesia	2000	Male	70 to 74	6.8% (4.1-11%)	93% (89-96%)	0.07 (0.035-0.12)	0.82% (0.53-1.2%)	0.96% (0.61-1.4%)	630 (390-890)	12000 (7600-17000)
Indonesia	2000	Male	75 to 79	5.9% (3.4-9.6%)	94% (90-97%)	0.058 (0.026-0.1)	0.69% (0.41-0.94%)	0.81% (0.47-1.1%)	430 (250-600)	6200 (3700-8500)
Indonesia	2000	Male	80 to 84	5.5% (3.1-8.6%)	94% (91-97%)	0.052 (0.024-0.089)	0.56% (0.3-0.78%)	0.64% (0.33-0.91%)	220 (110-320)	2400 (1300-3300)
Indonesia	2000	Male	85 to 89	5.5% (3.1-8.6%)	95% (91-97%)	0.05 (0.024-0.088)	0.41% (0.21-0.58%)	0.46% (0.23-0.69%)	110 (52-160)	810 (420-1200)
Indonesia	2000	Male	90 to 94	6% (3.5-9.4%)	94% (91-96%)	0.052 (0.023-0.096)	0.3% (0.14-0.45%)	0.34% (0.14-0.53%)	34 (14-53)	180 (84-280)
Indonesia	2000	Male	95 plus	7.1% (3.8-12%)	93% (88-96%)	0.056 (0.022-0.11)	0.28% (0.11-0.45%)	0.29% (0.078-0.5%)	7.2 (2-12)	28 (10-45)
Indonesia	2005	Female	All Ages	2.4% (1.9-3%)	98% (97-98%)	0.14 (0.13-0.15)	1.1% (0.89-1.3%)	1.7% (1.3-2.2%)	12000 (9000-16000)	370000 (300000-450000)
Indonesia	2005	Female	15 to 19	2.3% (1.3-3.7%)	98% (96-99%)	0.12 (0.056-0.2)	0.59% (0.39-0.85%)	0.75% (0.4-1.3%)	63 (34-110)	8300 (5500-12000)
Indonesia	2005	Female	20 to 24	2.6% (1.4-4.3%)	97% (96-99%)	0.15 (0.069-0.25)	1% (0.7-1.4%)	1% (0.55-1.7%)	110 (59-180)	16000 (10000-22000)
Indonesia	2005	Female	25 to 29	2.7% (1.4-4.4%)	97% (96-99%)	0.16 (0.083-0.26)	1.1% (0.83-1.5%)	1.2% (0.73-1.9%)	160 (94-250)	19000 (13000-26000)
Indonesia	2005	Female	30 to 34	2.8% (1.5-4.7%)	97% (95-99%)	0.17 (0.088-0.28)	1.2% (0.88-1.6%)	1.4% (0.96-2.1%)	230 (150-330)	21000 (15000-28000)
Indonesia	2005	Female	35 to 39	2.9% (1.5-4.8%)	97% (95-99%)	0.19 (0.098-0.31)	1.4% (1-1.8%)	1.9% (1.3-2.6%)	370 (260-520)	26000 (19000-34000)
Indonesia	2005	Female	40 to 44	3.1% (1.8-5%)	97% (95-98%)	0.23 (0.12-0.37)	1.6% (1.2-2.1%)	2.2% (1.6-3.1%)	550 (390-760)	31000 (23000-42000)
Indonesia	2005	Female	45 to 49	3.2% (1.6-5.1%)	97% (95-98%)	0.27 (0.13-0.47)	1.8% (1.3-2.5%)	2.5% (1.7-3.6%)	730 (500-1100)	35000 (24000-48000)
Indonesia	2005	Female	50 to 54	3.3% (1.8-5.5%)	97% (95-98%)	0.29 (0.14-0.49)	2% (1.4-2.8%)	2.6% (1.8-3.7%)	960 (660-1400)	38000 (26000-54000)
Indonesia	2005	Female	55 to 59	3.3% (1.8-5.4%)	97% (95-98%)	0.27 (0.13-0.48)	1.8% (1.2-2.7%)	2.4% (1.6-3.5%)	1100 (720-1500)	36000 (25000-52000)
Indonesia	2005	Female	60 to 64	3.4% (1.8-5.8%)	97% (94-98%)	0.27 (0.13-0.5)	1.8% (1.1-2.7%)	2.2% (1.4-3.4%)	1300 (810-1900)	36000 (23000-54000)
Indonesia	2005	Female	65 to 69	3.5% (2-5.6%)	96% (94-98%)	0.27 (0.13-0.47)	1.7% (1.1-2.7%)	2.1% (1.3-3.3%)	1500 (930-2300)	35000 (22000-54000)
Indonesia	2005	Female	70 to 74	3.6% (2.1-5.8%)	96% (94-98%)	0.27 (0.13-0.48)	1.6% (0.97-2.6%)	2% (1.2-3.1%)	1600 (960-2500)	30000 (18000-47000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	2005	Female	75 to 79	3.6% (2-6.1%)	96% (94-98%)	0.26 (0.12-0.48)	1.5% (0.84-2.4%)	1.7% (0.98-2.8%)	1300 (750-2100)	19000 (11000-31000)
Indonesia	2005	Female	80 to 84	3.7% (2-6.1%)	96% (94-98%)	0.26 (0.12-0.46)	1.4% (0.85-2.3%)	1.7% (1-2.7%)	1100 (650-1700)	12000 (6800-18000)
Indonesia	2005	Female	85 to 89	4% (2.2-6.7%)	96% (93-98%)	0.28 (0.13-0.51)	1.3% (0.75-2.2%)	1.6% (0.88-2.6%)	630 (360-1100)	4700 (2700-7900)
Indonesia	2005	Female	90 to 94	4.7% (2.7-7.6%)	95% (92-97%)	0.31 (0.14-0.59)	1.2% (0.66-2.1%)	1.4% (0.78-2.5%)	240 (130-420)	1200 (680-2100)
Indonesia	2005	Female	95 plus	5.6% (2.7-9.9%)	94% (90-97%)	0.35 (0.14-0.7)	1.2% (0.6-2.2%)	1.4% (0.67-2.5%)	72 (34-130)	250 (120-440)
Indonesia	2005	Male	All Ages	7.6% (6.2-9.4%)	92% (91-94%)	0.084 (0.074-0.094)	1.1% (0.88-1.3%)	1.4% (1-1.6%)	10000 (7900-12000)	440000 (350000-520000)
Indonesia	2005	Male	15 to 19	7.8% (4.6-12%)	92% (88-95%)	0.044 (0.021-0.078)	0.55% (0.38-0.76%)	0.5% (0.23-0.8%)	70 (32-110)	9600 (6400-14000)
Indonesia	2005	Male	20 to 24	12% (7.3-19%)	88% (81-93%)	0.11 (0.06-0.18)	1.2% (0.77-1.7%)	1% (0.45-1.7%)	210 (91-350)	25000 (15000-36000)
Indonesia	2005	Male	25 to 29	13% (7.4-19%)	87% (81-93%)	0.15 (0.08-0.24)	1.6% (1-2.1%)	1.5% (0.67-2.3%)	340 (150-530)	33000 (21000-47000)
Indonesia	2005	Male	30 to 34	12% (6.8-19%)	88% (81-93%)	0.16 (0.091-0.26)	2% (1.5-2.6%)	2.3% (1.5-3.1%)	580 (380-790)	44000 (31000-57000)
Indonesia	2005	Male	35 to 39	12% (6.8-18%)	88% (82-93%)	0.17 (0.095-0.27)	2.5% (1.9-3.2%)	3.1% (2.2-4.1%)	910 (650-1200)	56000 (41000-72000)
Indonesia	2005	Male	40 to 44	11% (6.2-17%)	89% (83-94%)	0.16 (0.085-0.26)	2.6% (2.1-3.3%)	3.4% (2.6-4.3%)	1100 (860-1400)	59000 (45000-73000)
Indonesia	2005	Male	45 to 49	9.1% (4.8-14%)	91% (86-95%)	0.14 (0.072-0.23)	2.6% (1.9-3.3%)	3.4% (2.5-4.2%)	1300 (940-1600)	56000 (42000-71000)
Indonesia	2005	Male	50 to 54	8% (4.7-13%)	92% (87-95%)	0.11 (0.059-0.19)	2.2% (1.7-2.8%)	2.9% (2.2-3.6%)	1200 (920-1500)	47000 (36000-59000)
Indonesia	2005	Male	55 to 59	7.4% (4.4-12%)	93% (88-96%)	0.095 (0.048-0.16)	1.7% (1.3-2.2%)	2.2% (1.6-2.8%)	1100 (820-1400)	37000 (28000-46000)
Indonesia	2005	Male	60 to 64	7.3% (4.2-12%)	93% (88-96%)	0.084 (0.04-0.15)	1.4% (1-1.8%)	1.7% (1.2-2.3%)	1000 (720-1300)	29000 (20000-37000)
Indonesia	2005	Male	65 to 69	7.2% (4.3-11%)	93% (89-96%)	0.078 (0.041-0.13)	1% (0.71-1.4%)	1.3% (0.84-1.6%)	830 (570-1100)	19000 (14000-26000)
Indonesia	2005	Male	70 to 74	6.6% (4.1-10%)	93% (90-96%)	0.067 (0.032-0.12)	0.81% (0.54-1.1%)	0.97% (0.63-1.3%)	670 (440-920)	12000 (8400-17000)
Indonesia	2005	Male	75 to 79	5.8% (3.3-9.4%)	94% (91-97%)	0.055 (0.025-0.1)	0.71% (0.44-0.94%)	0.84% (0.5-1.1%)	480 (280-640)	6800 (4200-9100)
Indonesia	2005	Male	80 to 84	5.3% (3-8.3%)	95% (92-97%)	0.048 (0.023-0.082)	0.59% (0.35-0.8%)	0.7% (0.39-0.94%)	280 (160-390)	3000 (1700-4100)
Indonesia	2005	Male	85 to 89	5.2% (3-8.2%)	95% (92-97%)	0.046 (0.022-0.081)	0.43% (0.23-0.58%)	0.49% (0.27-0.68%)	130 (69-180)	990 (540-1300)
Indonesia	2005	Male	90 to 94	5.7% (3.3-9%)	94% (91-97%)	0.047 (0.022-0.088)	0.32% (0.16-0.46%)	0.36% (0.17-0.53%)	41 (19-61)	220 (110-320)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	2005	Male	95 plus	6.8% (3.6-11%)	93% (89-96%)	0.052 (0.02-0.1)	0.29% (0.13-0.44%)	0.3% (0.1-0.47%)	9.7 (3.3-15)	37 (16-55)
Indonesia	2010	Female	All Ages	3% (2.4-3.7%)	97% (96-98%)	0.17 (0.15-0.18)	1.2% (0.95-1.4%)	1.7% (1.3-2.2%)	12000 (9300-16000)	390000 (320000-480000)
Indonesia	2010	Female	15 to 19	2.6% (1.5-4.3%)	97% (96-99%)	0.12 (0.06-0.23)	0.59% (0.39-0.86%)	0.76% (0.4-1.3%)	54 (29-96)	7600 (4900-11000)
Indonesia	2010	Female	20 to 24	3.2% (1.7-5.2%)	97% (95-98%)	0.17 (0.078-0.3)	1.1% (0.73-1.5%)	1.1% (0.58-1.9%)	98 (51-170)	15000 (9900-22000)
Indonesia	2010	Female	25 to 29	3.5% (1.9-5.5%)	97% (94-98%)	0.19 (0.095-0.31)	1.2% (0.86-1.6%)	1.3% (0.76-2.1%)	160 (90-250)	20000 (14000-28000)
Indonesia	2010	Female	30 to 34	3.7% (2-6.1%)	96% (94-98%)	0.21 (0.11-0.34)	1.3% (0.93-1.7%)	1.6% (1-2.3%)	230 (150-330)	23000 (16000-31000)
Indonesia	2010	Female	35 to 39	4.1% (2.1-6.6%)	96% (93-98%)	0.24 (0.13-0.39)	1.5% (1.1-2%)	2.1% (1.4-2.9%)	410 (280-580)	29000 (21000-40000)
Indonesia	2010	Female	40 to 44	4.1% (2.4-6.8%)	96% (93-98%)	0.28 (0.15-0.45)	1.7% (1.2-2.3%)	2.4% (1.7-3.3%)	630 (440-870)	36000 (26000-49000)
Indonesia	2010	Female	45 to 49	4.1% (2.1-6.4%)	96% (94-98%)	0.32 (0.16-0.53)	1.9% (1.3-2.6%)	2.6% (1.8-3.8%)	880 (600-1300)	42000 (29000-59000)
Indonesia	2010	Female	50 to 54	4.1% (2.3-6.8%)	96% (93-98%)	0.33 (0.17-0.55)	2% (1.4-2.8%)	2.7% (1.9-3.9%)	1100 (770-1600)	46000 (32000-65000)
Indonesia	2010	Female	55 to 59	4.1% (2.3-6.6%)	96% (93-98%)	0.32 (0.16-0.54)	1.9% (1.2-2.7%)	2.4% (1.6-3.6%)	1100 (730-1600)	38000 (25000-55000)
Indonesia	2010	Female	60 to 64	4.1% (2.3-7%)	96% (93-98%)	0.3 (0.14-0.55)	1.8% (1.1-2.7%)	2.3% (1.4-3.4%)	1200 (730-1900)	35000 (21000-53000)
Indonesia	2010	Female	65 to 69	4.1% (2.3-6.4%)	96% (94-98%)	0.29 (0.15-0.49)	1.7% (1-2.6%)	2.1% (1.3-3.2%)	1400 (840-2100)	32000 (20000-50000)
Indonesia	2010	Female	70 to 74	4% (2.4-6.4%)	96% (94-98%)	0.27 (0.13-0.48)	1.6% (0.92-2.4%)	1.9% (1.1-2.9%)	1600 (930-2400)	29000 (18000-45000)
Indonesia	2010	Female	75 to 79	3.9% (2.2-6.5%)	96% (94-98%)	0.26 (0.12-0.46)	1.3% (0.77-2.1%)	1.6% (0.91-2.6%)	1300 (720-2000)	18000 (10000-30000)
Indonesia	2010	Female	80 to 84	3.9% (2.1-6.3%)	96% (94-98%)	0.26 (0.12-0.44)	1.4% (0.78-2.1%)	1.6% (0.92-2.5%)	1100 (640-1800)	12000 (6700-19000)
Indonesia	2010	Female	85 to 89	4.2% (2.3-7%)	96% (93-98%)	0.27 (0.12-0.49)	1.2% (0.71-2.1%)	1.5% (0.83-2.5%)	710 (400-1200)	5300 (3000-9000)
Indonesia	2010	Female	90 to 94	4.9% (2.8-7.9%)	95% (92-97%)	0.3 (0.13-0.58)	1.1% (0.61-2%)	1.3% (0.72-2.4%)	270 (140-470)	1400 (750-2400)
Indonesia	2010	Female	95 plus	5.9% (2.9-11%)	94% (89-97%)	0.35 (0.13-0.7)	1.2% (0.55-2.1%)	1.4% (0.62-2.5%)	78 (35-140)	270 (120-470)
Indonesia	2010	Male	All Ages	7.1% (5.8-8.6%)	93% (91-94%)	0.073 (0.065-0.082)	1.2% (0.96-1.4%)	1.4% (1.1-1.7%)	11000 (8600-13000)	470000 (380000-550000)
Indonesia	2010	Male	15 to 19	5.3% (3.1-8.4%)	95% (92-97%)	0.026 (0.012-0.049)	0.55% (0.38-0.72%)	0.47% (0.21-0.7%)	60 (27-90)	8800 (5900-12000)
Indonesia	2010	Male	20 to 24	8.4% (4.7-13%)	92% (87-95%)	0.066 (0.034-0.11)	1.1% (0.81-1.5%)	0.85% (0.45-1.3%)	150 (83-240)	21000 (15000-30000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	2010	Male	25 to 29	9.9% (5.5-15%)	90% (85-94%)	0.1 (0.052-0.17)	1.4% (1-1.9%)	1.3% (0.66-1.8%)	290 (150-420)	32000 (22000-43000)
Indonesia	2010	Male	30 to 34	11% (6.2-17%)	89% (83-94%)	0.13 (0.072-0.2)	1.9% (1.4-2.4%)	2.1% (1.4-2.8%)	530 (360-700)	43000 (32000-55000)
Indonesia	2010	Male	35 to 39	12% (6.6-18%)	88% (82-93%)	0.15 (0.082-0.23)	2.4% (1.9-3%)	3% (2.2-3.8%)	920 (670-1200)	58000 (45000-72000)
Indonesia	2010	Male	40 to 44	11% (6.8-18%)	89% (82-93%)	0.15 (0.08-0.24)	2.5% (2-3%)	3.3% (2.6-4.1%)	1200 (930-1500)	63000 (50000-77000)
Indonesia	2010	Male	45 to 49	11% (5.8-16%)	89% (84-94%)	0.14 (0.076-0.23)	2.5% (1.9-3.1%)	3.3% (2.5-4.1%)	1500 (1100-1800)	66000 (49000-82000)
Indonesia	2010	Male	50 to 54	10% (6-15%)	90% (85-94%)	0.13 (0.068-0.21)	2.2% (1.6-2.7%)	2.8% (2.1-3.6%)	1500 (1100-2000)	59000 (45000-76000)
Indonesia	2010	Male	55 to 59	9.4% (5.6-15%)	91% (85-94%)	0.11 (0.056-0.17)	1.7% (1.3-2.2%)	2.2% (1.6-2.7%)	1300 (960-1700)	44000 (33000-55000)
Indonesia	2010	Male	60 to 64	8.9% (5.2-14%)	91% (86-95%)	0.09 (0.045-0.16)	1.4% (0.95-1.8%)	1.7% (1.2-2.2%)	1100 (720-1400)	30000 (20000-38000)
Indonesia	2010	Male	65 to 69	8.3% (4.9-13%)	92% (87-95%)	0.079 (0.04-0.13)	0.99% (0.7-1.3%)	1.2% (0.83-1.6%)	850 (590-1100)	20000 (14000-26000)
Indonesia	2010	Male	70 to 74	7.2% (4.5-11%)	93% (89-95%)	0.064 (0.031-0.11)	0.76% (0.52-1%)	0.91% (0.61-1.2%)	690 (460-920)	13000 (8800-17000)
Indonesia	2010	Male	75 to 79	6.1% (3.6-9.9%)	94% (90-96%)	0.051 (0.023-0.092)	0.67% (0.43-0.89%)	0.8% (0.5-1.1%)	500 (310-670)	7200 (4500-9600)
Indonesia	2010	Male	80 to 84	5.4% (3.1-8.4%)	95% (92-97%)	0.044 (0.02-0.075)	0.57% (0.34-0.74%)	0.67% (0.38-0.88%)	310 (180-420)	3300 (2000-4400)
Indonesia	2010	Male	85 to 89	5.3% (3-8.3%)	95% (92-97%)	0.041 (0.02-0.072)	0.41% (0.24-0.54%)	0.47% (0.27-0.64%)	160 (88-210)	1200 (700-1600)
Indonesia	2010	Male	90 to 94	5.7% (3.4-9%)	94% (91-97%)	0.041 (0.019-0.077)	0.3% (0.16-0.42%)	0.34% (0.17-0.49%)	48 (24-69)	250 (140-360)
Indonesia	2010	Male	95 plus	6.7% (3.5-11%)	93% (89-96%)	0.045 (0.018-0.091)	0.28% (0.13-0.41%)	0.29% (0.11-0.45%)	11 (4.2-17)	42 (19-60)
Indonesia	2016	Female	All Ages	3.1% (2.5-3.7%)	97% (96-97%)	0.2 (0.15-0.24)	1.3% (1-1.6%)	1.8% (1.4-2.5%)	13000 (9700-18000)	420000 (330000-540000)
Indonesia	2016	Female	15 to 19	2.6% (1.5-4.3%)	97% (96-98%)	0.14 (0.064-0.27)	0.6% (0.39-0.88%)	0.81% (0.41-1.5%)	54 (27-97)	7900 (5100-12000)
Indonesia	2016	Female	20 to 24	3.2% (1.7-5.3%)	97% (95-98%)	0.19 (0.082-0.37)	1.1% (0.76-1.6%)	1.1% (0.58-2%)	89 (44-150)	15000 (9900-21000)
Indonesia	2016	Female	25 to 29	3.5% (1.9-5.5%)	97% (95-98%)	0.21 (0.097-0.36)	1.2% (0.88-1.7%)	1.4% (0.74-2.2%)	130 (70-210)	18000 (12000-25000)
Indonesia	2016	Female	30 to 34	3.7% (2-6.2%)	96% (94-98%)	0.23 (0.11-0.4)	1.3% (0.92-1.7%)	1.6% (1-2.4%)	210 (130-320)	23000 (16000-31000)
Indonesia	2016	Female	35 to 39	4.1% (2.2-6.5%)	96% (93-98%)	0.27 (0.13-0.45)	1.5% (1-2%)	2.1% (1.4-3%)	380 (250-550)	29000 (21000-40000)
Indonesia	2016	Female	40 to 44	4.1% (2.4-6.8%)	96% (93-98%)	0.31 (0.16-0.52)	1.6% (1.2-2.2%)	2.4% (1.7-3.4%)	610 (420-870)	37000 (26000-50000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	2016	Female	45 to 49	4.1% (2.1-6.5%)	96% (93-98%)	0.35 (0.18-0.6)	1.8% (1.3-2.6%)	2.6% (1.8-3.9%)	910 (610-1300)	45000 (31000-64000)
Indonesia	2016	Female	50 to 54	4.1% (2.3-6.8%)	96% (93-98%)	0.37 (0.2-0.63)	1.9% (1.3-2.8%)	2.7% (1.8-3.9%)	1200 (820-1800)	51000 (35000-73000)
Indonesia	2016	Female	55 to 59	4.2% (2.3-6.7%)	96% (93-98%)	0.35 (0.18-0.6)	1.8% (1.2-2.7%)	2.5% (1.6-3.7%)	1400 (890-2000)	48000 (32000-71000)
Indonesia	2016	Female	60 to 64	4.2% (2.3-7%)	96% (93-98%)	0.34 (0.16-0.64)	1.8% (1.1-2.7%)	2.3% (1.4-3.6%)	1400 (860-2200)	41000 (25000-63000)
Indonesia	2016	Female	65 to 69	4.2% (2.4-6.5%)	96% (94-98%)	0.33 (0.16-0.6)	1.7% (1-2.7%)	2.2% (1.3-3.5%)	1500 (890-2300)	35000 (21000-55000)
Indonesia	2016	Female	70 to 74	4% (2.4-6.3%)	96% (94-98%)	0.31 (0.14-0.59)	1.6% (0.94-2.5%)	2% (1.2-3.2%)	1500 (910-2400)	29000 (17000-46000)
Indonesia	2016	Female	75 to 79	3.9% (2.2-6.6%)	96% (93-98%)	0.29 (0.13-0.56)	1.4% (0.76-2.3%)	1.7% (0.92-2.8%)	1500 (810-2500)	22000 (12000-36000)
Indonesia	2016	Female	80 to 84	4% (2.1-6.6%)	96% (93-98%)	0.29 (0.13-0.53)	1.4% (0.76-2.2%)	1.6% (0.92-2.6%)	1200 (660-1900)	13000 (7100-20000)
Indonesia	2016	Female	85 to 89	4.3% (2.4-7.1%)	96% (93-98%)	0.31 (0.14-0.58)	1.3% (0.7-2.1%)	1.5% (0.83-2.5%)	770 (420-1300)	5800 (3200-9700)
Indonesia	2016	Female	90 to 94	5% (2.9-7.9%)	95% (92-97%)	0.34 (0.15-0.65)	1.2% (0.6-2%)	1.4% (0.71-2.4%)	330 (170-560)	1700 (900-2900)
Indonesia	2016	Female	95 plus	6% (3-10%)	94% (90-97%)	0.4 (0.15-0.85)	1.2% (0.54-2.2%)	1.4% (0.61-2.5%)	97 (42-180)	340 (150-610)
Indonesia	2016	Male	All Ages	7.3% (6-8.8%)	93% (91-94%)	0.082 (0.062-0.1)	1.3% (1.1-1.5%)	1.5% (1.2-1.8%)	12000 (10000-15000)	520000 (420000-620000)
Indonesia	2016	Male	15 to 19	5.2% (3.1-8.6%)	95% (91-97%)	0.028 (0.013-0.057)	0.59% (0.4-0.79%)	0.54% (0.23-0.78%)	70 (30-100)	10000 (6700-14000)
Indonesia	2016	Male	20 to 24	8.4% (4.7-13%)	92% (87-95%)	0.073 (0.034-0.14)	1.2% (0.89-1.7%)	0.96% (0.54-1.5%)	170 (98-270)	24000 (17000-33000)
Indonesia	2016	Male	25 to 29	9.9% (5.7-15%)	90% (85-94%)	0.11 (0.052-0.19)	1.6% (1.2-2.1%)	1.4% (0.85-2.1%)	260 (160-380)	29000 (21000-40000)
Indonesia	2016	Male	30 to 34	11% (6-17%)	89% (83-94%)	0.14 (0.07-0.23)	2% (1.6-2.6%)	2.3% (1.6-3%)	530 (370-700)	44000 (33000-57000)
Indonesia	2016	Male	35 to 39	12% (6.7-18%)	88% (82-93%)	0.16 (0.082-0.27)	2.5% (1.9-3.1%)	3.2% (2.4-4%)	910 (670-1200)	59000 (45000-73000)
Indonesia	2016	Male	40 to 44	11% (6.7-18%)	89% (82-93%)	0.16 (0.084-0.27)	2.5% (2-3.1%)	3.4% (2.6-4.2%)	1200 (960-1500)	66000 (52000-81000)
Indonesia	2016	Male	45 to 49	11% (5.8-16%)	89% (84-94%)	0.15 (0.079-0.27)	2.5% (1.9-3.1%)	3.3% (2.5-4.1%)	1600 (1200-2000)	71000 (55000-90000)
Indonesia	2016	Male	50 to 54	10% (6-16%)	90% (84-94%)	0.14 (0.072-0.23)	2.1% (1.7-2.7%)	2.8% (2.1-3.6%)	1700 (1200-2200)	65000 (50000-84000)
Indonesia	2016	Male	55 to 59	9.4% (5.6-15%)	91% (85-94%)	0.11 (0.058-0.19)	1.7% (1.3-2.2%)	2.2% (1.6-2.8%)	1600 (1200-2100)	53000 (40000-68000)
Indonesia	2016	Male	60 to 64	8.8% (5.2-14%)	91% (86-95%)	0.097 (0.046-0.18)	1.4% (1-1.8%)	1.7% (1.2-2.3%)	1400 (990-1900)	41000 (28000-54000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Indonesia	2016	Male	65 to 69	8.3% (5.1-13%)	92% (87-95%)	0.085 (0.042-0.15)	1% (0.75-1.3%)	1.2% (0.89-1.6%)	980 (700-1300)	23000 (17000-30000)
Indonesia	2016	Male	70 to 74	7.2% (4.5-11%)	93% (89-95%)	0.07 (0.033-0.13)	0.8% (0.58-1.1%)	0.97% (0.68-1.3%)	780 (540-1100)	15000 (10000-20000)
Indonesia	2016	Male	75 to 79	6.1% (3.5-9.8%)	94% (90-96%)	0.056 (0.024-0.11)	0.71% (0.49-0.94%)	0.85% (0.58-1.1%)	630 (410-870)	9100 (6100-12000)
Indonesia	2016	Male	80 to 84	5.4% (3-8.4%)	95% (92-97%)	0.047 (0.021-0.085)	0.61% (0.39-0.79%)	0.72% (0.44-0.95%)	360 (220-480)	3800 (2500-5000)
Indonesia	2016	Male	85 to 89	5.2% (3-8.3%)	95% (92-97%)	0.044 (0.021-0.081)	0.44% (0.28-0.58%)	0.51% (0.32-0.68%)	170 (110-230)	1300 (840-1700)
Indonesia	2016	Male	90 to 94	5.6% (3.4-8.9%)	94% (91-97%)	0.045 (0.021-0.084)	0.33% (0.19-0.45%)	0.38% (0.21-0.53%)	59 (33-83)	310 (190-430)
Indonesia	2016	Male	95 plus	6.6% (3.6-11%)	93% (89-96%)	0.049 (0.018-0.1)	0.31% (0.15-0.44%)	0.33% (0.14-0.49%)	14 (5.8-21)	52 (26-75)
Laos	1990	Female	All Ages	24% (20-28%)	76% (72-80%)	0.63 (0.52-0.76)	1.2% (0.68-1.8%)	2.5% (1.3-4%)	710 (370-1100)	22000 (13000-32000)
Laos	1990	Female	15 to 19	24% (17-33%)	76% (67-83%)	0.77 (0.46-1.2)	1.9% (1.3-3.3%)	2.7% (1.3-4.6%)	13 (6.2-23)	1000 (550-1700)
Laos	1990	Female	20 to 24	31% (21-42%)	69% (58-79%)	1.1 (0.62-1.6)	3% (1.6-4.9%)	3.7% (1.8-6.4%)	22 (10-37)	1700 (890-2700)
Laos	1990	Female	25 to 29	33% (23-43%)	67% (57-77%)	1.1 (0.69-1.6)	3.3% (1.7-5.4%)	4.2% (2-7.4%)	24 (11-41)	1600 (840-2700)
Laos	1990	Female	30 to 34	34% (23-46%)	66% (54-77%)	1.1 (0.69-1.7)	3.2% (1.7-5.1%)	4.1% (2-6.9%)	22 (11-36)	1400 (720-2200)
Laos	1990	Female	35 to 39	36% (24-46%)	64% (54-76%)	1.2 (0.75-1.9)	4.3% (2.2-7.1%)	5.8% (2.9-9.5%)	31 (16-52)	1700 (870-2700)
Laos	1990	Female	40 to 44	35% (25-46%)	65% (54-75%)	1.3 (0.77-1.9)	4.5% (2.2-7.5%)	6.1% (2.9-10%)	34 (16-55)	1600 (790-2600)
Laos	1990	Female	45 to 49	34% (23-45%)	66% (55-77%)	1.2 (0.7-2)	4.9% (2.3-8.7%)	6.5% (2.9-11%)	40 (18-70)	1700 (770-2900)
Laos	1990	Female	50 to 54	34% (24-45%)	66% (55-76%)	1.2 (0.72-2)	4.9% (1.9-8.9%)	6.3% (2.4-11%)	51 (20-91)	1900 (730-3400)
Laos	1990	Female	55 to 59	34% (25-46%)	66% (54-75%)	1.3 (0.7-2.1)	4.7% (1.6-8.8%)	5.8% (1.9-11%)	61 (20-110)	1900 (660-3600)
Laos	1990	Female	60 to 64	32% (22-44%)	68% (56-78%)	1.2 (0.7-2.1)	4.3% (1.3-8.6%)	5.2% (1.5-10%)	73 (22-140)	2000 (590-3900)
Laos	1990	Female	65 to 69	28% (19-37%)	72% (63-81%)	1.2 (0.72-1.8)	4.3% (1.7-8.1%)	5.1% (2-9.6%)	85 (32-160)	1900 (730-3500)
Laos	1990	Female	70 to 74	27% (19-36%)	73% (64-81%)	1.2 (0.7-1.9)	4.6% (1.6-8.8%)	5.3% (1.9-10%)	96 (34-190)	1700 (620-3400)
Laos	1990	Female	75 to 79	28% (19-38%)	72% (62-81%)	1.3 (0.71-2)	4.3% (1.4-8.3%)	4.9% (1.6-9.7%)	78 (25-150)	1100 (360-2100)
Laos	1990	Female	80 to 84	29% (20-38%)	71% (62-80%)	1.2 (0.75-1.9)	4.2% (1.4-7.8%)	4.9% (1.7-9%)	53 (18-98)	550 (190-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	1990	Female	85 to 89	29% (20-40%)	71% (60-80%)	1.2 (0.73-1.9)	3.6% (1.2-6.7%)	4.1% (1.4-7.8%)	20 (6.8-38)	150 (52-280)
Laos	1990	Female	90 to 94	31% (21-42%)	69% (58-79%)	1.3 (0.72-2.1)	3.1% (0.74-6.4%)	3.7% (0.81-7.7%)	5.3 (1.2-11)	27 (6.4-56)
Laos	1990	Female	95 plus	34% (22-47%)	66% (53-78%)	1.4 (0.64-2.5)	3.3% (0.33-7.4%)	3.7% (0.29-8.5%)	1.2 (0.095-2.8)	4.2 (0.41-9.5)
Laos	1990	Male	All Ages	48% (41-54%)	52% (46-59%)	1.4 (1.2-1.6)	2.9% (2-3.8%)	4.9% (3.3-6.7%)	1500 (1000-2100)	61000 (45000-80000)
Laos	1990	Male	15 to 19	46% (34-58%)	54% (42-66%)	1 (0.62-1.5)	3.5% (1.9-5.5%)	4.4% (2.4-7.1%)	26 (14-42)	2000 (1100-3200)
Laos	1990	Male	20 to 24	66% (53-76%)	34% (24-47%)	2.4 (1.6-3.1)	9.9% (6.5-14%)	12% (7.8-17%)	89 (56-130)	6300 (4100-8700)
Laos	1990	Male	25 to 29	71% (60-79%)	29% (21-40%)	2.9 (2-3.7)	14% (8.7-19%)	17% (10-23%)	120 (72-160)	7500 (4800-10000)
Laos	1990	Male	30 to 34	75% (65-82%)	25% (18-35%)	3.2 (2.4-4.2)	15% (10-20%)	18% (12-25%)	120 (80-170)	7200 (4800-9800)
Laos	1990	Male	35 to 39	76% (67-83%)	24% (17-33%)	3.3 (2.4-4.3)	16% (11-21%)	19% (13-26%)	130 (88-180)	6900 (4700-9300)
Laos	1990	Male	40 to 44	75% (65-82%)	25% (18-35%)	3.2 (2.3-4.3)	15% (9.9-21%)	18% (12-26%)	120 (77-170)	5700 (3700-7900)
Laos	1990	Male	45 to 49	72% (60-79%)	28% (21-40%)	3.2 (2.2-4.3)	15% (9.6-21%)	18% (11-26%)	140 (83-190)	5700 (3500-7900)
Laos	1990	Male	50 to 54	69% (58-77%)	31% (23-42%)	2.9 (1.9-4)	14% (7.9-20%)	16% (9.2-24%)	150 (87-220)	5600 (3300-8200)
Laos	1990	Male	55 to 59	67% (55-77%)	33% (23-45%)	2.5 (1.7-3.5)	11% (5.9-17%)	13% (7-20%)	140 (77-220)	4500 (2500-7000)
Laos	1990	Male	60 to 64	63% (50-74%)	37% (26-50%)	2.1 (1.4-3.2)	9% (4.4-15%)	11% (5-17%)	140 (65-230)	3800 (1800-6300)
Laos	1990	Male	65 to 69	56% (43-67%)	44% (33-57%)	1.8 (1.2-2.6)	7.5% (3.6-13%)	8.7% (4.1-15%)	120 (57-200)	2700 (1300-4500)
Laos	1990	Male	70 to 74	52% (39-64%)	48% (36-61%)	1.7 (1-2.4)	6.7% (2.7-11%)	7.7% (3.1-13%)	110 (43-190)	1900 (810-3300)
Laos	1990	Male	75 to 79	52% (38-67%)	48% (33-62%)	1.6 (0.92-2.4)	5% (1.6-9.2%)	5.8% (1.8-11%)	64 (19-120)	890 (290-1600)
Laos	1990	Male	80 to 84	50% (34-65%)	50% (35-66%)	1.3 (0.8-1.9)	4.1% (1.2-7.9%)	4.8% (1.3-9.2%)	31 (8.4-59)	320 (93-610)
Laos	1990	Male	85 to 89	45% (31-59%)	55% (41-69%)	1.1 (0.62-1.7)	2.6% (0.52-5.3%)	2.9% (0.51-6.1%)	9.2 (1.6-19)	69 (14-140)
Laos	1990	Male	90 to 94	44% (30-58%)	56% (42-70%)	0.95 (0.54-1.5)	1.6% (-0.03-3.8%)	1.8% (-0.13-4.5%)	1.9 (-0.14-4.7)	9.9 (-0.19-24)
Laos	1990	Male	95 plus	45% (29-64%)	55% (36-71%)	0.86 (0.39-1.6)	1.1% (-0.49-3.5%)	1.1% (-0.72-3.9%)	0.26 (-0.17-0.88)	1 (-0.43-3.1)
Laos	1995	Female	All Ages	24% (20-28%)	76% (72-80%)	0.62 (0.52-0.75)	1.2% (0.73-1.8%)	2.5% (1.3-4%)	730 (390-1100)	23000 (14000-33000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	1995	Female	15 to 19	24% (16-33%)	76% (67-84%)	0.76 (0.47-1.2)	2.1% (1.1-3.3%)	2.9% (1.5-4.9%)	15 (7.5-25)	1200 (640-1900)
Laos	1995	Female	20 to 24	31% (21-42%)	69% (58-79%)	1.1 (0.62-1.6)	3% (1.6-5%)	3.9% (1.9-6.6%)	24 (12-41)	1900 (990-3000)
Laos	1995	Female	25 to 29	33% (23-44%)	67% (56-77%)	1.1 (0.68-1.6)	3.3% (1.8-5.2%)	4.3% (2.1-7.2%)	25 (12-43)	1800 (930-2900)
Laos	1995	Female	30 to 34	34% (23-46%)	66% (54-77%)	1.1 (0.66-1.7)	3.2% (1.7-5.1%)	4.2% (2.1-7%)	24 (12-40)	1500 (840-2400)
Laos	1995	Female	35 to 39	36% (24-46%)	64% (54-76%)	1.2 (0.73-1.9)	4.2% (2.2-6.9%)	5.8% (2.9-9.6%)	34 (17-57)	1900 (1000-3000)
Laos	1995	Female	40 to 44	35% (25-46%)	65% (54-75%)	1.3 (0.76-1.9)	4.5% (2.3-7.1%)	6.1% (3.1-9.6%)	37 (19-59)	1800 (930-2800)
Laos	1995	Female	45 to 49	34% (23-44%)	66% (56-77%)	1.2 (0.72-2)	4.7% (2.1-8.1%)	6.3% (2.8-11%)	42 (19-72)	1800 (820-3000)
Laos	1995	Female	50 to 54	34% (24-45%)	66% (55-76%)	1.2 (0.71-2)	4.7% (1.8-8.5%)	6.1% (2.3-11%)	49 (18-88)	1800 (680-3200)
Laos	1995	Female	55 to 59	34% (25-45%)	66% (55-75%)	1.2 (0.68-2.1)	4.4% (1.4-8.6%)	5.5% (1.8-11%)	58 (18-110)	1800 (590-3600)
Laos	1995	Female	60 to 64	32% (22-44%)	68% (56-78%)	1.2 (0.68-2)	4.1% (1.2-8.4%)	4.9% (1.4-9.8%)	67 (19-130)	1800 (510-3700)
Laos	1995	Female	65 to 69	29% (20-37%)	71% (63-80%)	1.2 (0.72-1.8)	4.1% (1.5-7.5%)	4.9% (1.8-8.9%)	85 (31-160)	1900 (700-3500)
Laos	1995	Female	70 to 74	27% (19-37%)	73% (63-81%)	1.2 (0.71-1.9)	4.4% (1.6-8.4%)	5.1% (1.8-9.8%)	97 (34-190)	1800 (630-3500)
Laos	1995	Female	75 to 79	28% (19-39%)	72% (61-81%)	1.2 (0.69-2)	4.1% (1.3-8.3%)	4.7% (1.5-9.6%)	82 (25-170)	1200 (350-2300)
Laos	1995	Female	80 to 84	29% (19-39%)	71% (61-81%)	1.2 (0.73-2)	4.1% (1.5-7.5%)	4.7% (1.7-8.7%)	58 (21-110)	600 (220-1100)
Laos	1995	Female	85 to 89	29% (20-40%)	71% (60-80%)	1.2 (0.72-2)	3.5% (1.2-6.9%)	4% (1.4-8%)	24 (8.1-47)	170 (61-340)
Laos	1995	Female	90 to 94	31% (22-42%)	69% (58-78%)	1.3 (0.7-2.1)	3% (0.74-6.3%)	3.5% (0.82-7.4%)	6.3 (1.5-13)	32 (8.2-67)
Laos	1995	Female	95 plus	34% (22-46%)	66% (54-78%)	1.3 (0.61-2.5)	3.2% (0.25-7.4%)	3.6% (0.2-8.5%)	1.5 (0.085-3.5)	5.1 (0.41-12)
Laos	1995	Male	All Ages	48% (41-54%)	52% (46-59%)	1.3 (1.2-1.5)	3% (2.1-4%)	4.9% (3.3-6.8%)	1600 (1100-2100)	65000 (48000-84000)
Laos	1995	Male	15 to 19	46% (34-58%)	54% (42-66%)	1 (0.63-1.4)	3.6% (2-5.5%)	4.7% (2.5-7.3%)	27 (15-43)	2200 (1200-3300)
Laos	1995	Male	20 to 24	66% (54-76%)	34% (24-46%)	2.3 (1.6-3.1)	10% (6.7-14%)	13% (8.1-18%)	97 (61-140)	6900 (4500-9400)
Laos	1995	Male	25 to 29	72% (61-79%)	28% (21-39%)	2.8 (2.1-3.7)	14% (9.2-19%)	17% (11-23%)	120 (80-180)	8100 (5400-11000)
Laos	1995	Male	30 to 34	75% (65-82%)	25% (18-35%)	3.2 (2.4-4.1)	15% (10-20%)	18% (12-25%)	130 (89-190)	8000 (5400-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	1995	Male	35 to 39	76% (67-83%)	24% (17-33%)	3.3 (2.4-4.2)	16% (11-21%)	19% (13-26%)	140 (95-200)	7600 (5100-10000)
Laos	1995	Male	40 to 44	75% (66-82%)	25% (18-34%)	3.2 (2.3-4.2)	15% (10-20%)	18% (12-25%)	140 (94-200)	6800 (4600-9400)
Laos	1995	Male	45 to 49	72% (60-79%)	28% (21-40%)	3.1 (2.2-4.2)	15% (9.6-21%)	18% (11-25%)	140 (88-200)	5900 (3700-8200)
Laos	1995	Male	50 to 54	69% (58-77%)	31% (23-42%)	2.9 (1.9-4)	13% (7.9-19%)	16% (9.2-23%)	140 (84-210)	5300 (3200-7600)
Laos	1995	Male	55 to 59	67% (56-77%)	33% (23-44%)	2.5 (1.7-3.5)	10% (5.8-16%)	12% (6.8-19%)	140 (77-210)	4500 (2500-6800)
Laos	1995	Male	60 to 64	63% (50-74%)	37% (26-50%)	2.1 (1.4-3.2)	8.6% (4.3-14%)	10% (5-17%)	140 (67-230)	3800 (1900-6200)
Laos	1995	Male	65 to 69	56% (43-67%)	44% (33-57%)	1.8 (1.3-2.6)	7.1% (3.4-12%)	8.2% (3.9-14%)	120 (59-210)	2800 (1400-4600)
Laos	1995	Male	70 to 74	52% (40-64%)	48% (36-60%)	1.6 (1-2.4)	6.3% (2.6-11%)	7.3% (3-13%)	110 (45-190)	2000 (840-3500)
Laos	1995	Male	75 to 79	53% (37-67%)	47% (33-63%)	1.5 (0.92-2.3)	4.7% (1.4-8.7%)	5.4% (1.5-10%)	68 (19-120)	960 (290-1700)
Laos	1995	Male	80 to 84	50% (34-65%)	50% (35-66%)	1.3 (0.76-1.9)	3.8% (1-7.4%)	4.4% (1.1-8.7%)	33 (8.3-65)	350 (92-680)
Laos	1995	Male	85 to 89	45% (31-60%)	55% (40-69%)	1 (0.6-1.7)	2.4% (0.43-5%)	2.7% (0.39-5.7%)	10 (1.5-22)	77 (14-160)
Laos	1995	Male	90 to 94	45% (31-59%)	55% (41-69%)	0.93 (0.53-1.5)	1.4% (-0.11-3.5%)	1.6% (-0.25-4.1%)	2 (-0.32-5.1)	11 (-0.83-26)
Laos	1995	Male	95 plus	46% (29-65%)	54% (35-71%)	0.85 (0.4-1.6)	1% (-0.53-3.2%)	1% (-0.8-3.6%)	0.29 (-0.23-1)	1.1 (-0.61-3.6)
Laos	2000	Female	All Ages	22% (19-25%)	78% (75-81%)	0.54 (0.45-0.65)	1.1% (0.69-1.7%)	2.2% (1.2-3.6%)	630 (330-1000)	20000 (12000-29000)
Laos	2000	Female	15 to 19	22% (16-29%)	78% (71-84%)	0.66 (0.41-1)	1.9% (1-3%)	2.7% (1.4-4.4%)	14 (7.1-23)	1100 (620-1800)
Laos	2000	Female	20 to 24	29% (22-37%)	71% (63-78%)	0.96 (0.6-1.5)	2.8% (1.5-4.5%)	3.5% (1.7-6.1%)	22 (11-36)	1700 (940-2700)
Laos	2000	Female	25 to 29	30% (23-38%)	70% (62-77%)	0.95 (0.65-1.4)	3% (1.7-4.6%)	3.9% (1.9-6.4%)	23 (11-38)	1600 (900-2600)
Laos	2000	Female	30 to 34	31% (24-41%)	69% (59-76%)	0.97 (0.64-1.4)	2.8% (1.6-4.4%)	3.8% (1.9-6.1%)	22 (11-35)	1400 (790-2200)
Laos	2000	Female	35 to 39	34% (26-42%)	66% (58-74%)	1.1 (0.73-1.6)	3.7% (2.1-6.1%)	5.2% (2.8-8.4%)	31 (17-52)	1700 (980-2800)
Laos	2000	Female	40 to 44	34% (26-42%)	66% (58-74%)	1.2 (0.76-1.7)	3.9% (2.1-6.3%)	5.5% (2.9-8.8%)	34 (18-55)	1700 (900-2600)
Laos	2000	Female	45 to 49	31% (23-38%)	69% (62-77%)	1.1 (0.67-1.7)	3.9% (1.7-6.7%)	5.4% (2.3-9.1%)	37 (16-63)	1600 (720-2700)
Laos	2000	Female	50 to 54	28% (20-36%)	72% (64-80%)	0.96 (0.57-1.6)	3.6% (1.3-6.6%)	4.7% (1.7-8.5%)	39 (14-71)	1500 (510-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	2000	Female	55 to 59	27% (19-36%)	73% (64-81%)	0.95 (0.54-1.6)	3.3% (1.1-6.4%)	4.2% (1.3-8.1%)	42 (13-81)	1300 (430-2600)
Laos	2000	Female	60 to 64	26% (18-35%)	74% (65-82%)	0.95 (0.54-1.6)	3.1% (0.87-6.5%)	3.8% (1.1-8%)	51 (14-110)	1400 (380-2900)
Laos	2000	Female	65 to 69	24% (17-32%)	76% (68-83%)	0.95 (0.58-1.5)	3.2% (1.2-6%)	3.8% (1.5-7.1%)	63 (23-120)	1400 (540-2700)
Laos	2000	Female	70 to 74	24% (16-32%)	76% (68-84%)	1 (0.59-1.7)	3.5% (1.2-7.1%)	4.2% (1.4-8.3%)	82 (28-160)	1500 (500-3000)
Laos	2000	Female	75 to 79	26% (18-36%)	74% (64-82%)	1.2 (0.68-1.9)	3.6% (1-7.4%)	4.2% (1.2-8.5%)	76 (21-160)	1100 (300-2200)
Laos	2000	Female	80 to 84	28% (19-38%)	72% (62-81%)	1.2 (0.71-1.9)	3.7% (1.2-7%)	4.3% (1.4-8.3%)	59 (20-110)	610 (210-1200)
Laos	2000	Female	85 to 89	28% (19-38%)	72% (62-81%)	1.2 (0.67-1.8)	3.1% (1-6.2%)	3.7% (1.1-7.3%)	25 (7.8-50)	190 (59-360)
Laos	2000	Female	90 to 94	30% (21-40%)	70% (60-79%)	1.2 (0.65-2)	2.7% (0.58-5.8%)	3.2% (0.64-6.8%)	7 (1.4-15)	36 (7.6-76)
Laos	2000	Female	95 plus	32% (21-45%)	68% (55-79%)	1.3 (0.59-2.3)	2.9% (0.15-6.8%)	3.2% (0.08-7.8%)	1.6 (0.04-3.9)	5.5 (0.3-13)
Laos	2000	Male	All Ages	51% (45-57%)	49% (43-55%)	1.4 (1.2-1.6)	3.3% (2.4-4.4%)	5.3% (3.7-7.3%)	1600 (1100-2200)	69000 (52000-88000)
Laos	2000	Male	15 to 19	49% (40-59%)	51% (41-60%)	1 (0.68-1.4)	3.7% (2.2-5.4%)	5% (2.9-7.5%)	30 (17-47)	2400 (1400-3600)
Laos	2000	Male	20 to 24	70% (61-77%)	30% (23-39%)	2.4 (1.7-3.1)	10% (6.9-14%)	13% (8.5-18%)	100 (67-140)	7200 (4900-9900)
Laos	2000	Male	25 to 29	76% (69-81%)	24% (19-31%)	3 (2.2-3.8)	14% (9.4-19%)	17% (11-24%)	130 (87-180)	8700 (5800-12000)
Laos	2000	Male	30 to 34	79% (73-83%)	21% (17-27%)	3.3 (2.5-4.2)	15% (11-20%)	19% (13-25%)	140 (97-190)	8400 (5900-11000)
Laos	2000	Male	35 to 39	80% (75-84%)	20% (16-25%)	3.4 (2.6-4.3)	15% (11-21%)	19% (13-26%)	150 (100-210)	8100 (5600-11000)
Laos	2000	Male	40 to 44	78% (73-83%)	22% (17-27%)	3.3 (2.5-4.3)	15% (10-20%)	18% (12-25%)	150 (100-210)	7300 (5000-10000)
Laos	2000	Male	45 to 49	76% (69-81%)	24% (19-31%)	3.3 (2.4-4.4)	15% (9.7-20%)	18% (12-25%)	170 (110-230)	7000 (4600-9700)
Laos	2000	Male	50 to 54	73% (66-78%)	27% (22-34%)	3 (2.1-4.1)	13% (8-18%)	16% (9.4-22%)	150 (90-220)	5500 (3400-8000)
Laos	2000	Male	55 to 59	71% (62-78%)	29% (22-38%)	2.5 (1.8-3.5)	10% (5.9-15%)	12% (6.8-18%)	130 (75-200)	4200 (2500-6200)
Laos	2000	Male	60 to 64	64% (54-73%)	36% (27-46%)	2.1 (1.4-3)	8% (4-13%)	9.5% (4.6-15%)	130 (64-220)	3600 (1800-5900)
Laos	2000	Male	65 to 69	53% (42-64%)	47% (36-58%)	1.7 (1.2-2.3)	6.3% (3-11%)	7.4% (3.5-13%)	110 (55-190)	2600 (1300-4400)
Laos	2000	Male	70 to 74	54% (43-64%)	46% (36-57%)	1.7 (1.1-2.4)	6% (2.4-10%)	7% (2.8-12%)	120 (47-200)	2100 (850-3600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	2000	Male	75 to 79	56% (43-68%)	44% (32-57%)	1.6 (1-2.4)	4.6% (1.3-8.6%)	5.3% (1.4-9.9%)	74 (20-140)	1100 (300-2000)
Laos	2000	Male	80 to 84	50% (34-65%)	50% (35-66%)	1.3 (0.75-1.9)	3.6% (0.84-6.9%)	4.1% (0.91-8%)	36 (8-71)	380 (89-730)
Laos	2000	Male	85 to 89	43% (30-58%)	57% (42-70%)	0.97 (0.57-1.5)	2.1% (0.36-4.6%)	2.4% (0.33-5.3%)	11 (1.5-24)	84 (14-180)
Laos	2000	Male	90 to 94	42% (29-57%)	58% (43-71%)	0.85 (0.47-1.4)	1.3% (-0.11-3.1%)	1.4% (-0.28-3.6%)	2.2 (-0.44-5.6)	12 (-1-29)
Laos	2000	Male	95 plus	44% (26-62%)	56% (38-74%)	0.77 (0.36-1.4)	0.92% (-0.53-3%)	0.9% (-0.76-3.4%)	0.31 (-0.26-1.2)	1.2 (-0.71-4.1)
Laos	2005	Female	All Ages	22% (19-25%)	78% (75-81%)	0.53 (0.45-0.64)	1.1% (0.67-1.6%)	2.1% (1.1-3.4%)	540 (280-880)	17000 (11000-25000)
Laos	2005	Female	15 to 19	21% (15-28%)	79% (72-85%)	0.62 (0.39-0.93)	1.7% (0.95-2.6%)	2.5% (1.3-4%)	13 (6.6-21)	1100 (620-1700)
Laos	2005	Female	20 to 24	28% (21-36%)	72% (64-79%)	0.91 (0.56-1.4)	2.5% (1.4-3.9%)	3.2% (1.6-5.5%)	19 (9.2-32)	1500 (890-2400)
Laos	2005	Female	25 to 29	30% (23-37%)	70% (63-77%)	0.91 (0.61-1.3)	2.6% (1.5-4.1%)	3.5% (1.7-5.9%)	19 (9.4-32)	1400 (790-2200)
Laos	2005	Female	30 to 34	30% (23-39%)	70% (61-77%)	0.93 (0.6-1.4)	2.5% (1.5-3.9%)	3.4% (1.8-5.4%)	18 (9.7-29)	1200 (740-1900)
Laos	2005	Female	35 to 39	33% (25-41%)	67% (59-75%)	1.1 (0.67-1.6)	3.2% (1.8-5%)	4.5% (2.4-7.2%)	26 (14-42)	1500 (830-2300)
Laos	2005	Female	40 to 44	33% (26-40%)	67% (60-74%)	1.1 (0.73-1.6)	3.4% (1.8-5.3%)	4.8% (2.5-7.7%)	30 (15-48)	1500 (800-2400)
Laos	2005	Female	45 to 49	30% (23-38%)	70% (62-77%)	1 (0.64-1.6)	3.3% (1.4-5.6%)	4.7% (2-7.9%)	32 (14-56)	1400 (610-2400)
Laos	2005	Female	50 to 54	27% (19-36%)	73% (64-81%)	0.92 (0.53-1.5)	3% (0.99-5.6%)	4% (1.3-7.4%)	34 (11-63)	1300 (420-2400)
Laos	2005	Female	55 to 59	27% (19-36%)	73% (64-81%)	0.91 (0.52-1.5)	2.7% (0.75-5.4%)	3.5% (1-7.1%)	36 (10-71)	1200 (320-2300)
Laos	2005	Female	60 to 64	25% (17-34%)	75% (66-83%)	0.91 (0.5-1.5)	2.6% (0.66-5.7%)	3.2% (0.82-7.2%)	40 (9.9-88)	1100 (270-2500)
Laos	2005	Female	65 to 69	23% (16-31%)	77% (69-84%)	0.9 (0.54-1.4)	2.7% (0.85-5.2%)	3.3% (1.1-6.3%)	51 (17-100)	1200 (370-2300)
Laos	2005	Female	70 to 74	23% (16-31%)	77% (69-84%)	0.97 (0.56-1.6)	3% (0.92-6.3%)	3.6% (1.1-7.4%)	68 (20-140)	1200 (370-2500)
Laos	2005	Female	75 to 79	26% (18-36%)	74% (64-82%)	1.1 (0.63-1.8)	3.1% (0.82-6.6%)	3.6% (0.92-7.7%)	68 (18-150)	960 (260-2100)
Laos	2005	Female	80 to 84	27% (19-38%)	73% (62-81%)	1.1 (0.68-1.8)	3.2% (1-6.1%)	3.8% (1.2-7.2%)	54 (17-110)	570 (180-1100)
Laos	2005	Female	85 to 89	28% (19-38%)	72% (62-81%)	1.1 (0.64-1.8)	2.7% (0.83-5.5%)	3.2% (0.95-6.5%)	25 (7.5-51)	190 (57-380)
Laos	2005	Female	90 to 94	29% (20-40%)	71% (60-80%)	1.1 (0.64-1.9)	2.4% (0.4-5.3%)	2.8% (0.39-6.2%)	7.4 (1-17)	38 (6.3-84)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	2005	Female	95 plus	32% (21-44%)	68% (56-79%)	1.2 (0.56-2.1)	2.5% (-0.053-6.1%)	2.8% (-0.17-7%)	1.7 (-0.11-4.3)	5.9 (-0.13-15)
Laos	2005	Male	All Ages	51% (46-58%)	49% (42-54%)	1.4 (1.2-1.6)	3.4% (2.5-4.6%)	5.2% (3.6-7.1%)	1500 (1100-2000)	64000 (48000-82000)
Laos	2005	Male	15 to 19	48% (39-58%)	52% (42-61%)	0.99 (0.65-1.3)	3.6% (2.2-5.3%)	4.9% (2.8-7.3%)	31 (18-47)	2500 (1500-3700)
Laos	2005	Male	20 to 24	69% (60-76%)	31% (24-40%)	2.3 (1.7-3)	9.7% (6.4-13%)	12% (8-17%)	97 (61-130)	6900 (4600-9400)
Laos	2005	Male	25 to 29	75% (69-80%)	25% (20-31%)	2.9 (2.1-3.7)	13% (8.2-17%)	16% (10-22%)	120 (75-160)	7800 (5000-10000)
Laos	2005	Male	30 to 34	79% (73-83%)	21% (17-27%)	3.3 (2.5-4.1)	14% (9.5-18%)	17% (12-23%)	130 (85-170)	7600 (5300-10000)
Laos	2005	Male	35 to 39	79% (74-84%)	21% (16-26%)	3.3 (2.5-4.2)	14% (9.9-19%)	17% (12-24%)	130 (92-180)	7200 (5000-9800)
Laos	2005	Male	40 to 44	78% (73-83%)	22% (17-27%)	3.2 (2.4-4.2)	13% (9.2-18%)	16% (11-23%)	140 (94-200)	6800 (4700-9300)
Laos	2005	Male	45 to 49	75% (69-81%)	25% (19-31%)	3.2 (2.3-4.1)	13% (8.8-18%)	16% (11-22%)	160 (100-220)	6800 (4500-9300)
Laos	2005	Male	50 to 54	72% (65-78%)	28% (22-35%)	2.9 (2-3.9)	11% (7.1-17%)	14% (8.5-20%)	160 (95-230)	5800 (3600-8400)
Laos	2005	Male	55 to 59	70% (61-78%)	30% (22-39%)	2.5 (1.7-3.4)	8.9% (5-13%)	11% (5.8-16%)	120 (66-180)	4000 (2200-5900)
Laos	2005	Male	60 to 64	63% (53-72%)	37% (28-47%)	2 (1.4-3)	6.9% (3.1-12%)	8.2% (3.7-14%)	110 (49-190)	3000 (1400-5100)
Laos	2005	Male	65 to 69	53% (42-63%)	47% (37-58%)	1.6 (1.1-2.2)	5.5% (2.6-9%)	6.4% (3-11%)	100 (49-170)	2300 (1100-3800)
Laos	2005	Male	70 to 74	53% (42-63%)	47% (37-58%)	1.6 (1-2.2)	5.1% (2.1-8.6%)	6% (2.4-9.9%)	100 (43-170)	1900 (780-3100)
Laos	2005	Male	75 to 79	55% (43-67%)	45% (33-57%)	1.5 (0.99-2.3)	3.9% (0.98-7.3%)	4.5% (1-8.4%)	69 (16-130)	990 (250-1800)
Laos	2005	Male	80 to 84	50% (35-63%)	50% (37-65%)	1.2 (0.74-1.8)	3% (0.67-5.7%)	3.5% (0.69-6.5%)	35 (6.8-66)	370 (81-690)
Laos	2005	Male	85 to 89	42% (28-57%)	58% (43-72%)	0.92 (0.53-1.4)	1.7% (0.18-3.8%)	2% (0.1-4.4%)	11 (0.55-24)	82 (8.8-180)
Laos	2005	Male	90 to 94	41% (28-55%)	59% (45-72%)	0.8 (0.45-1.3)	1% (-0.25-2.6%)	1.1% (-0.46-2.9%)	2.1 (-0.88-5.6)	12 (-2.9-30)
Laos	2005	Male	95 plus	42% (26-61%)	58% (39-74%)	0.72 (0.34-1.4)	0.71% (-0.92-2.5%)	0.64% (-0.92-2.7%)	0.28 (-0.4-1.2)	1.2 (-1-4.1)
Laos	2010	Female	All Ages	24% (20-27%)	76% (73-80%)	0.63 (0.51-0.76)	1.2% (0.71-1.9%)	2.4% (1.1-4%)	560 (280-920)	18000 (11000-26000)
Laos	2010	Female	15 to 19	23% (15-31%)	77% (69-85%)	0.71 (0.42-1.1)	1.7% (0.98-2.8%)	2.7% (1.4-4.5%)	13 (6.5-21)	1100 (630-1800)
Laos	2010	Female	20 to 24	30% (20-40%)	70% (60-80%)	0.98 (0.59-1.5)	2.4% (1.4-3.9%)	3.2% (1.5-5.5%)	18 (8.1-31)	1600 (860-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	2010	Female	25 to 29	31% (21-41%)	69% (59-79%)	0.98 (0.62-1.5)	2.5% (1.5-4%)	3.4% (1.7-5.8%)	17 (8.2-28)	1300 (750-2100)
Laos	2010	Female	30 to 34	32% (22-44%)	68% (56-78%)	1 (0.61-1.6)	2.4% (1.4-3.8%)	3.4% (1.7-5.6%)	16 (8.2-27)	1100 (670-1800)
Laos	2010	Female	35 to 39	34% (23-45%)	66% (55-77%)	1.1 (0.67-1.8)	3% (1.6-4.7%)	4.4% (2.3-7%)	23 (12-38)	1400 (740-2100)
Laos	2010	Female	40 to 44	33% (24-44%)	67% (56-76%)	1.2 (0.71-1.8)	3.1% (1.6-5.1%)	4.7% (2.4-7.6%)	27 (14-44)	1400 (740-2200)
Laos	2010	Female	45 to 49	32% (21-42%)	68% (58-79%)	1.1 (0.65-1.9)	3.1% (1.4-5.5%)	4.6% (2-8%)	32 (14-56)	1400 (610-2500)
Laos	2010	Female	50 to 54	32% (22-42%)	68% (58-78%)	1.1 (0.63-1.9)	3.2% (0.92-6.2%)	4.4% (1.3-8.4%)	37 (11-72)	1400 (420-2800)
Laos	2010	Female	55 to 59	32% (23-43%)	68% (57-77%)	1.2 (0.63-1.9)	2.9% (0.73-6.3%)	3.9% (0.92-8.4%)	41 (9.9-87)	1300 (340-2900)
Laos	2010	Female	60 to 64	30% (20-41%)	70% (59-80%)	1.1 (0.61-2)	2.7% (0.51-6.4%)	3.5% (0.67-8%)	44 (8.4-100)	1200 (230-2800)
Laos	2010	Female	65 to 69	27% (18-36%)	73% (64-82%)	1.1 (0.63-1.8)	2.9% (0.75-6%)	3.6% (0.93-7.4%)	53 (14-110)	1200 (320-2500)
Laos	2010	Female	70 to 74	25% (18-35%)	75% (65-82%)	1.1 (0.64-1.9)	3.2% (0.82-6.5%)	3.8% (1-7.9%)	70 (19-140)	1300 (330-2600)
Laos	2010	Female	75 to 79	26% (18-37%)	74% (63-82%)	1.2 (0.65-2)	3.1% (0.59-6.4%)	3.6% (0.67-7.8%)	66 (12-140)	950 (180-2000)
Laos	2010	Female	80 to 84	27% (18-37%)	73% (63-82%)	1.2 (0.68-1.9)	3.2% (0.99-6.3%)	3.8% (1.2-7.5%)	56 (17-110)	590 (180-1200)
Laos	2010	Female	85 to 89	28% (19-38%)	72% (62-81%)	1.2 (0.68-1.9)	2.9% (0.78-5.8%)	3.4% (0.89-6.8%)	32 (8.4-64)	240 (65-480)
Laos	2010	Female	90 to 94	30% (20-40%)	70% (60-80%)	1.2 (0.67-2.1)	2.5% (0.38-5.7%)	3% (0.39-6.7%)	10 (1.4-23)	53 (8.3-120)
Laos	2010	Female	95 plus	32% (20-45%)	68% (55-80%)	1.3 (0.58-2.4)	2.7% (-0.077-6.6%)	3% (-0.18-7.5%)	2.4 (-0.15-6.2)	8.5 (-0.24-21)
Laos	2010	Male	All Ages	48% (41-54%)	52% (46-59%)	1.4 (1.2-1.7)	3.6% (2.3-4.9%)	5.2% (3.2-7.1%)	1400 (940-1900)	61000 (42000-78000)
Laos	2010	Male	15 to 19	44% (32-56%)	56% (44-68%)	0.93 (0.57-1.3)	3.3% (2-5.1%)	4.7% (2.6-7.2%)	30 (17-47)	2500 (1500-3700)
Laos	2010	Male	20 to 24	64% (51-74%)	36% (26-49%)	2.2 (1.5-3)	8.8% (5.7-12%)	11% (7-16%)	96 (59-140)	7000 (4500-9700)
Laos	2010	Male	25 to 29	70% (58-79%)	30% (21-42%)	2.7 (1.9-3.6)	11% (6.1-15%)	14% (7.2-20%)	110 (56-150)	7300 (4000-10000)
Laos	2010	Male	30 to 34	73% (62-81%)	27% (19-38%)	3.1 (2.2-4)	12% (6.8-16%)	15% (8.3-21%)	110 (61-160)	6800 (3900-9400)
Laos	2010	Male	35 to 39	75% (64-82%)	25% (18-36%)	3.2 (2.3-4.3)	12% (7.8-17%)	16% (9.5-22%)	120 (72-170)	6600 (4100-9300)
Laos	2010	Male	40 to 44	74% (64-81%)	26% (19-36%)	3.1 (2.2-4.3)	12% (7.8-17%)	15% (9.6-21%)	120 (78-180)	6100 (4000-8600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	2010	Male	45 to 49	70% (57-79%)	30% (21-43%)	3 (2.1-4.2)	12% (7.5-17%)	15% (9.3-21%)	150 (92-210)	6300 (4100-9000)
Laos	2010	Male	50 to 54	68% (57-76%)	32% (24-43%)	2.8 (1.8-3.9)	10% (6.1-15%)	13% (7.4-19%)	150 (87-220)	5600 (3300-8300)
Laos	2010	Male	55 to 59	66% (53-76%)	34% (24-47%)	2.4 (1.5-3.4)	8% (4.1-12%)	9.6% (4.9-15%)	130 (66-200)	4300 (2200-6700)
Laos	2010	Male	60 to 64	61% (47-73%)	39% (27-53%)	2 (1.3-3.1)	6.3% (2.9-11%)	7.4% (3.4-13%)	100 (48-180)	2900 (1400-5100)
Laos	2010	Male	65 to 69	54% (41-66%)	46% (34-59%)	1.7 (1.1-2.5)	5.2% (2.5-8.6%)	6.1% (2.9-10%)	95 (45-160)	2200 (1100-3700)
Laos	2010	Male	70 to 74	50% (37-62%)	50% (38-63%)	1.5 (0.97-2.4)	4.6% (1.9-7.7%)	5.3% (2.2-8.9%)	93 (37-160)	1700 (730-2900)
Laos	2010	Male	75 to 79	50% (35-65%)	50% (35-65%)	1.4 (0.83-2.2)	3.3% (0.9-6.2%)	3.8% (0.92-7.2%)	61 (15-120)	890 (230-1700)
Laos	2010	Male	80 to 84	47% (31-62%)	53% (38-69%)	1.2 (0.71-1.8)	2.7% (0.67-5.1%)	3% (0.68-6%)	33 (7.4-65)	360 (90-690)
Laos	2010	Male	85 to 89	43% (29-57%)	57% (43-71%)	0.97 (0.56-1.5)	1.6% (0.15-3.4%)	1.8% (0.026-3.7%)	11 (0.17-24)	89 (8.6-190)
Laos	2010	Male	90 to 94	42% (28-57%)	58% (43-72%)	0.85 (0.47-1.4)	0.89% (-0.34-2.4%)	0.93% (-0.56-2.7%)	2.2 (-1.3-6.5)	13 (-4.8-35)
Laos	2010	Male	95 plus	43% (26-61%)	57% (39-74%)	0.76 (0.34-1.4)	0.59% (-0.66-2.4%)	0.49% (-1-2.6%)	0.27 (-0.55-1.4)	1.3 (-1.4-5.2)
Laos	2016	Female	All Ages	23% (20-27%)	77% (73-80%)	0.66 (0.45-0.92)	1.4% (0.59-2.4%)	2.6% (0.92-4.9%)	560 (200-1000)	18000 (8300-29000)
Laos	2016	Female	15 to 19	22% (15-30%)	78% (70-85%)	0.73 (0.36-1.2)	1.7% (0.88-2.7%)	2.8% (1.3-4.8%)	9 (3.9-16)	850 (450-1400)
Laos	2016	Female	20 to 24	29% (19-39%)	71% (61-81%)	1 (0.52-1.6)	2.3% (1.3-3.8%)	3.3% (1.4-5.8%)	15 (6.3-27)	1400 (770-2300)
Laos	2016	Female	25 to 29	30% (21-41%)	70% (59-79%)	1 (0.58-1.6)	2.3% (1.3-3.7%)	3.3% (1.5-5.8%)	15 (6.7-26)	1300 (710-2100)
Laos	2016	Female	30 to 34	31% (21-43%)	69% (57-79%)	1 (0.58-1.7)	2.3% (1.2-3.6%)	3.3% (1.5-5.8%)	14 (6.3-25)	1100 (580-1800)
Laos	2016	Female	35 to 39	33% (22-44%)	67% (56-78%)	1.2 (0.63-1.9)	2.7% (1.4-4.5%)	4.3% (1.9-7.2%)	20 (9-35)	1300 (630-2100)
Laos	2016	Female	40 to 44	33% (23-43%)	67% (57-77%)	1.2 (0.64-2)	3% (1.4-5.1%)	4.7% (2.1-8%)	26 (12-45)	1400 (640-2400)
Laos	2016	Female	45 to 49	31% (21-42%)	69% (58-79%)	1.2 (0.64-2)	3% (1.2-5.4%)	4.5% (1.8-8.4%)	31 (12-58)	1400 (570-2600)
Laos	2016	Female	50 to 54	31% (22-42%)	69% (58-78%)	1.2 (0.61-2.1)	3% (0.85-6.1%)	4.4% (1.2-8.8%)	39 (11-77)	1500 (410-3000)
Laos	2016	Female	55 to 59	32% (23-43%)	68% (57-77%)	1.2 (0.62-2.2)	2.9% (0.5-6.5%)	4% (0.74-9%)	44 (8.4-97)	1500 (250-3300)
Laos	2016	Female	60 to 64	30% (20-41%)	70% (59-80%)	1.2 (0.61-2.2)	2.7% (0.35-6.6%)	3.5% (0.47-8.5%)	47 (6.6-110)	1300 (170-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	2016	Female	65 to 69	26% (18-35%)	74% (65-82%)	1.1 (0.6-2)	2.8% (0.54-6.2%)	3.6% (0.72-7.9%)	56 (11-120)	1300 (260-2800)
Laos	2016	Female	70 to 74	25% (18-35%)	75% (65-82%)	1.2 (0.62-2.1)	3.1% (0.47-6.6%)	3.8% (0.57-8%)	66 (10-140)	1200 (180-2700)
Laos	2016	Female	75 to 79	26% (17-37%)	74% (63-83%)	1.2 (0.62-2.1)	3% (0.23-6.7%)	3.6% (0.2-8.1%)	65 (3.6-150)	950 (71-2200)
Laos	2016	Female	80 to 84	27% (18-37%)	73% (63-82%)	1.2 (0.67-2)	3.2% (0.65-6.7%)	3.8% (0.74-8.1%)	56 (11-120)	600 (120-1300)
Laos	2016	Female	85 to 89	28% (18-38%)	72% (62-82%)	1.2 (0.65-2.1)	2.9% (0.48-6.5%)	3.4% (0.49-7.6%)	37 (5.1-82)	280 (46-610)
Laos	2016	Female	90 to 94	29% (20-40%)	71% (60-80%)	1.3 (0.67-2.2)	2.6% (0.17-6.5%)	3.1% (0.15-7.7%)	15 (0.74-38)	78 (5.3-190)
Laos	2016	Female	95 plus	32% (20-45%)	68% (55-80%)	1.4 (0.57-2.6)	2.8% (-0.14-7.2%)	3.1% (-0.28-8%)	3.9 (-0.34-10)	14 (-0.67-35)
Laos	2016	Male	All Ages	49% (42-55%)	51% (45-58%)	1.5 (1.1-2.1)	3.9% (2.2-5.9%)	5.4% (2.9-8.3%)	1400 (800-2100)	62000 (38000-87000)
Laos	2016	Male	15 to 19	43% (31-56%)	57% (44-69%)	0.96 (0.52-1.5)	3.3% (1.9-5.2%)	4.8% (2.5-7.8%)	26 (14-42)	2100 (1200-3300)
Laos	2016	Male	20 to 24	64% (50-74%)	36% (26-50%)	2.3 (1.4-3.4)	8.5% (5.3-12%)	11% (6.6-17%)	94 (54-140)	7100 (4400-10000)
Laos	2016	Male	25 to 29	70% (58-79%)	30% (21-42%)	2.8 (1.9-4.3)	10% (5.8-15%)	14% (6.8-20%)	110 (57-170)	7900 (4300-11000)
Laos	2016	Male	30 to 34	73% (63-81%)	27% (19-37%)	3.2 (2.1-4.7)	11% (5.1-16%)	14% (5.8-21%)	110 (43-160)	7000 (3200-10000)
Laos	2016	Male	35 to 39	75% (63-82%)	25% (18-37%)	3.3 (2.1-4.9)	11% (5.9-17%)	15% (6.9-22%)	120 (55-180)	6600 (3400-9600)
Laos	2016	Male	40 to 44	74% (63-81%)	26% (19-37%)	3.3 (2-4.8)	11% (6.3-17%)	14% (7.5-21%)	120 (63-180)	6000 (3300-9100)
Laos	2016	Male	45 to 49	70% (57-79%)	30% (21-43%)	3.2 (2-4.8)	11% (6.4-16%)	14% (7.8-21%)	140 (76-210)	6100 (3400-9200)
Laos	2016	Male	50 to 54	68% (56-77%)	32% (23-44%)	2.9 (1.7-4.5)	9.7% (5.2-15%)	12% (6.4-19%)	150 (77-230)	5700 (3000-8900)
Laos	2016	Male	55 to 59	66% (54-76%)	34% (24-46%)	2.5 (1.5-3.9)	7.7% (3.5-13%)	9.3% (4.1-16%)	130 (60-230)	4400 (2000-7700)
Laos	2016	Male	60 to 64	61% (47-73%)	39% (27-53%)	2.1 (1.2-3.6)	6% (2.4-12%)	7.2% (2.7-14%)	120 (46-230)	3400 (1300-6600)
Laos	2016	Male	65 to 69	53% (40-65%)	47% (35-60%)	1.8 (1-3)	4.9% (2-9.6%)	5.8% (2.3-11%)	98 (39-190)	2300 (950-4400)
Laos	2016	Male	70 to 74	49% (37-62%)	51% (38-63%)	1.6 (0.93-2.6)	4.2% (1.5-8%)	4.9% (1.7-9.3%)	84 (30-160)	1600 (570-3100)
Laos	2016	Male	75 to 79	50% (34-64%)	50% (36-66%)	1.5 (0.79-2.5)	3.1% (0.67-6.7%)	3.5% (0.69-7.8%)	58 (11-130)	860 (180-1900)
Laos	2016	Male	80 to 84	47% (31-61%)	53% (39-69%)	1.2 (0.67-2)	2.4% (0.44-5.3%)	2.7% (0.41-6.2%)	32 (5-73)	350 (66-780)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Laos	2016	Male	85 to 89	42% (28-57%)	58% (43-72%)	1 (0.53-1.6)	1.5% (-0.0076-3.6%)	1.6% (-0.15-4.2%)	12 (-1.2-31)	96 (-0.5-240)
Laos	2016	Male	90 to 94	41% (28-55%)	59% (45-72%)	0.88 (0.46-1.5)	0.8% (-0.48-2.6%)	0.8% (-0.76-2.9%)	2.5 (-2.3-9)	15 (-9-49)
Laos	2016	Male	95 plus	42% (26-60%)	58% (40-74%)	0.79 (0.34-1.5)	0.53% (-0.78-2.4%)	0.39% (-1.1-2.5%)	0.29 (-0.82-1.8)	1.5 (-2.3-6.8)
Malaysia	1990	Female	All Ages	2.5% (2.1-3.1%)	97% (97-98%)	0.46 (0.32-0.62)	1.4% (1.1-1.9%)	2.6% (1.8-3.5%)	910 (640-1300)	27000 (21000-35000)
Malaysia	1990	Female	15 to 19	2.4% (1.4-3.7%)	98% (96-99%)	0.42 (0.2-0.77)	0.93% (0.62-1.4%)	1.9% (1.1-3%)	8.8 (5.2-14)	950 (630-1400)
Malaysia	1990	Female	20 to 24	2.9% (1.6-4.4%)	97% (96-98%)	0.56 (0.27-1)	1.3% (0.9-1.9%)	2.2% (1.3-3.4%)	12 (7.1-19)	1400 (930-2000)
Malaysia	1990	Female	25 to 29	3.1% (1.8-4.9%)	97% (95-98%)	0.66 (0.33-1.1)	1.5% (1-2.1%)	2.5% (1.5-3.8%)	15 (9-24)	1600 (1100-2300)
Malaysia	1990	Female	30 to 34	3.3% (1.8-5.5%)	97% (95-98%)	0.72 (0.37-1.3)	1.6% (1.1-2.3%)	2.7% (1.6-4.2%)	19 (12-29)	1600 (1100-2400)
Malaysia	1990	Female	35 to 39	3.5% (1.9-5.4%)	96% (95-98%)	0.73 (0.35-1.2)	1.7% (1.1-2.4%)	3% (1.8-4.4%)	25 (15-37)	1800 (1100-2500)
Malaysia	1990	Female	40 to 44	3.6% (2.1-5.5%)	96% (95-98%)	0.81 (0.39-1.4)	1.9% (1.2-2.7%)	3.2% (2.1-4.7%)	32 (20-47)	1800 (1200-2600)
Malaysia	1990	Female	45 to 49	3.5% (2-5.4%)	96% (95-98%)	0.97 (0.44-1.7)	2.1% (1.4-3.1%)	3.5% (2.2-5.2%)	39 (25-58)	1900 (1200-2700)
Malaysia	1990	Female	50 to 54	3.6% (2.2-5.5%)	96% (94-98%)	1.1 (0.49-2)	2.5% (1.6-3.7%)	3.7% (2.3-5.5%)	63 (39-95)	2600 (1600-3800)
Malaysia	1990	Female	55 to 59	3.7% (2.2-5.7%)	96% (94-98%)	1.1 (0.52-2)	2.6% (1.6-4.1%)	3.6% (2.3-5.7%)	76 (47-120)	2600 (1600-4000)
Malaysia	1990	Female	60 to 64	3.6% (2.1-5.8%)	96% (94-98%)	1.1 (0.48-2)	2.5% (1.6-3.9%)	3.3% (2-5.1%)	97 (60-150)	2800 (1700-4400)
Malaysia	1990	Female	65 to 69	3.4% (1.9-5%)	97% (95-98%)	1 (0.46-1.8)	2.5% (1.5-3.8%)	3.1% (1.8-4.7%)	110 (65-170)	2600 (1500-3900)
Malaysia	1990	Female	70 to 74	3.2% (2-5%)	97% (95-98%)	0.99 (0.43-1.9)	2.4% (1.4-3.9%)	2.8% (1.7-4.7%)	130 (75-210)	2400 (1400-3900)
Malaysia	1990	Female	75 to 79	3.3% (1.9-5.2%)	97% (95-98%)	1 (0.41-2)	2.2% (1.2-3.6%)	2.6% (1.5-4.3%)	110 (60-180)	1600 (870-2500)
Malaysia	1990	Female	80 to 84	3.4% (1.8-5.1%)	97% (95-98%)	1 (0.42-1.9)	2.2% (1.2-3.6%)	2.6% (1.4-4.3%)	88 (45-140)	930 (480-1500)
Malaysia	1990	Female	85 to 89	3.5% (2-5.5%)	96% (94-98%)	1 (0.47-1.8)	2.2% (1.2-3.6%)	2.5% (1.4-4.3%)	58 (30-97)	430 (230-730)
Malaysia	1990	Female	90 to 94	3.8% (2.3-6%)	96% (94-98%)	1.1 (0.45-2.1)	2.1% (1.2-3.6%)	2.5% (1.3-4.3%)	23 (12-39)	120 (65-200)
Malaysia	1990	Female	95 plus	4.3% (2.5-7.1%)	96% (93-98%)	1.2 (0.45-2.6)	2.3% (1.2-3.9%)	2.7% (1.4-4.6%)	8.4 (4.4-15)	29 (15-49)
Malaysia	1990	Male	All Ages	7.9% (6.5-9.5%)	92% (90-94%)	1.8 (1.3-2.3)	4.9% (3.9-6%)	6.5% (5.1-8%)	3100 (2400-3800)	120000 (92000-140000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	1990	Male	15 to 19	7% (4.3-10%)	93% (90-96%)	0.87 (0.4-1.6)	3.5% (2.2-5.6%)	5.4% (3.1-8.8%)	61 (36-98)	5100 (3100-7900)
Malaysia	1990	Male	20 to 24	11% (6.5-16%)	89% (84-94%)	2.1 (1.1-3.6)	7% (4.3-10%)	9.5% (5.3-15%)	130 (74-200)	10000 (6400-15000)
Malaysia	1990	Male	25 to 29	12% (7.5-18%)	88% (82-93%)	2.8 (1.5-4.7)	7.7% (5.1-11%)	11% (6.8-16%)	150 (91-210)	11000 (7300-15000)
Malaysia	1990	Male	30 to 34	13% (7.8-20%)	87% (80-92%)	3.4 (1.9-5.5)	7.8% (5.3-11%)	11% (7.5-16%)	150 (100-220)	10000 (7200-14000)
Malaysia	1990	Male	35 to 39	13% (8.2-19%)	87% (81-92%)	4 (2.1-6.5)	8% (5.7-11%)	12% (8.2-16%)	170 (110-230)	9900 (7100-13000)
Malaysia	1990	Male	40 to 44	13% (8.4-19%)	87% (81-92%)	4.1 (2.1-6.6)	8.3% (5.9-11%)	12% (8.2-15%)	190 (130-250)	10000 (7100-13000)
Malaysia	1990	Male	45 to 49	13% (7.6-19%)	87% (81-92%)	3.7 (1.9-6.4)	8.4% (6-11%)	11% (8.1-15%)	220 (160-290)	9900 (7100-13000)
Malaysia	1990	Male	50 to 54	13% (8-18%)	87% (82-92%)	3.7 (1.8-6.4)	8% (5.7-11%)	10% (7.2-14%)	290 (210-390)	11000 (8100-15000)
Malaysia	1990	Male	55 to 59	12% (7.6-18%)	88% (82-92%)	3.9 (1.9-6.8)	7.7% (5.3-11%)	9.3% (6.4-13%)	310 (210-440)	10000 (7200-14000)
Malaysia	1990	Male	60 to 64	11% (7-17%)	89% (83-93%)	3.4 (1.6-6.2)	6.8% (4.6-9.5%)	8% (5.4-11%)	340 (230-480)	9700 (6600-13000)
Malaysia	1990	Male	65 to 69	9.6% (6.1-14%)	90% (86-94%)	2.6 (1.2-4.3)	5.9% (3.9-8.4%)	6.8% (4.5-9.8%)	310 (210-440)	7200 (4800-10000)
Malaysia	1990	Male	70 to 74	8.2% (5.3-12%)	92% (88-95%)	1.9 (0.9-3.4)	4.9% (3.3-7.1%)	5.7% (3.7-8.1%)	290 (190-420)	5500 (3600-7900)
Malaysia	1990	Male	75 to 79	7.4% (4.6-12%)	93% (88-95%)	1.6 (0.67-3.1)	4% (2.6-6%)	4.6% (2.9-6.9%)	210 (130-320)	3000 (1900-4600)
Malaysia	1990	Male	80 to 84	6.7% (3.9-9.9%)	93% (90-96%)	1.3 (0.62-2.5)	3.6% (2.3-5.3%)	4.2% (2.6-6.2%)	140 (86-210)	1500 (930-2200)
Malaysia	1990	Male	85 to 89	6.4% (3.8-9.9%)	94% (90-96%)	1.2 (0.55-2.2)	3.2% (1.9-4.8%)	3.7% (2.2-5.6%)	79 (47-120)	600 (370-900)
Malaysia	1990	Male	90 to 94	6.8% (4.1-11%)	93% (89-96%)	1.2 (0.52-2.3)	2.9% (1.8-4.3%)	3.4% (2-5.1%)	27 (16-41)	140 (85-210)
Malaysia	1990	Male	95 plus	7.9% (4.3-13%)	92% (87-96%)	1.3 (0.49-2.7)	3.1% (1.7-4.9%)	3.4% (1.9-5.6%)	7.2 (3.9-12)	26 (14-42)
Malaysia	1995	Female	All Ages	2.6% (2.1-3.1%)	97% (97-98%)	0.38 (0.26-0.5)	1.4% (1.1-1.8%)	2.5% (1.8-3.5%)	980 (700-1400)	29000 (22000-37000)
Malaysia	1995	Female	15 to 19	2.4% (1.4-3.6%)	98% (96-99%)	0.34 (0.16-0.61)	0.87% (0.59-1.3%)	1.8% (1-2.9%)	8.4 (4.9-13)	950 (620-1400)
Malaysia	1995	Female	20 to 24	2.9% (1.6-4.5%)	97% (96-98%)	0.45 (0.21-0.81)	1.3% (0.84-1.8%)	2.1% (1.3-3.2%)	11 (6.6-17)	1400 (920-2000)
Malaysia	1995	Female	25 to 29	3.2% (1.8-4.9%)	97% (95-98%)	0.53 (0.27-0.91)	1.4% (0.97-2%)	2.4% (1.4-3.6%)	14 (8.2-21)	1500 (1000-2200)
Malaysia	1995	Female	30 to 34	3.3% (1.8-5.5%)	97% (94-98%)	0.57 (0.3-1)	1.5% (1-2.1%)	2.5% (1.6-3.9%)	18 (11-28)	1700 (1200-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	1995	Female	35 to 39	3.5% (1.8-5.4%)	96% (95-98%)	0.57 (0.28-0.98)	1.5% (0.98-2.1%)	2.7% (1.7-4.1%)	24 (15-36)	1800 (1200-2500)
Malaysia	1995	Female	40 to 44	3.5% (2.1-5.5%)	96% (95-98%)	0.63 (0.31-1.1)	1.6% (1.1-2.4%)	2.9% (1.9-4.2%)	32 (20-47)	2000 (1300-2800)
Malaysia	1995	Female	45 to 49	3.5% (2-5.3%)	96% (95-98%)	0.76 (0.35-1.3)	1.9% (1.2-2.8%)	3.1% (2-4.7%)	44 (28-67)	2200 (1400-3300)
Malaysia	1995	Female	50 to 54	3.6% (2.2-5.4%)	96% (95-98%)	0.86 (0.41-1.6)	2.3% (1.4-3.3%)	3.4% (2.1-5%)	57 (36-85)	2400 (1500-3400)
Malaysia	1995	Female	55 to 59	3.7% (2.2-5.6%)	96% (94-98%)	0.87 (0.41-1.6)	2.4% (1.5-3.6%)	3.4% (2.1-5.1%)	81 (51-120)	2800 (1800-4300)
Malaysia	1995	Female	60 to 64	3.6% (2.1-5.7%)	96% (94-98%)	0.84 (0.39-1.6)	2.4% (1.5-3.7%)	3.1% (1.9-4.8%)	98 (60-150)	2800 (1800-4400)
Malaysia	1995	Female	65 to 69	3.4% (1.9-5%)	97% (95-98%)	0.8 (0.37-1.4)	2.3% (1.4-3.5%)	2.9% (1.7-4.5%)	120 (71-180)	2800 (1700-4200)
Malaysia	1995	Female	70 to 74	3.2% (2-5%)	97% (95-98%)	0.78 (0.34-1.5)	2.2% (1.3-3.7%)	2.7% (1.6-4.5%)	130 (78-220)	2500 (1500-4000)
Malaysia	1995	Female	75 to 79	3.3% (1.9-5.2%)	97% (95-98%)	0.8 (0.33-1.6)	2.1% (1.1-3.5%)	2.5% (1.4-4.2%)	120 (67-200)	1800 (970-2900)
Malaysia	1995	Female	80 to 84	3.4% (1.8-4.9%)	97% (95-98%)	0.79 (0.33-1.4)	2.2% (1.1-3.5%)	2.6% (1.3-4.1%)	100 (51-160)	1100 (540-1700)
Malaysia	1995	Female	85 to 89	3.5% (2-5.5%)	97% (94-98%)	0.79 (0.37-1.4)	2.1% (1.1-3.6%)	2.5% (1.3-4.2%)	72 (38-120)	530 (290-900)
Malaysia	1995	Female	90 to 94	3.8% (2.3-5.9%)	96% (94-98%)	0.84 (0.36-1.6)	2.1% (1.1-3.6%)	2.5% (1.3-4.2%)	32 (17-54)	160 (88-280)
Malaysia	1995	Female	95 plus	4.4% (2.5-6.9%)	96% (93-98%)	0.94 (0.36-2.1)	2.3% (1.2-3.9%)	2.7% (1.4-4.5%)	9 (4.6-15)	31 (16-51)
Malaysia	1995	Male	All Ages	8.1% (6.7-9.7%)	92% (90-93%)	1.5 (1.1-1.9)	4.9% (3.9-5.9%)	6.2% (4.9-7.6%)	3300 (2600-4000)	130000 (100000-150000)
Malaysia	1995	Male	15 to 19	6.9% (4.2-10%)	93% (90-96%)	0.68 (0.31-1.2)	3.3% (2-5.4%)	4.8% (2.7-7.9%)	68 (37-110)	5700 (3300-8800)
Malaysia	1995	Male	20 to 24	11% (6.3-16%)	89% (84-94%)	1.7 (0.86-2.8)	6.7% (4.2-10%)	8.8% (5.1-14%)	150 (85-230)	12000 (7300-17000)
Malaysia	1995	Male	25 to 29	12% (7.4-18%)	88% (82-93%)	2.2 (1.1-3.7)	7.2% (4.8-10%)	9.9% (6.2-15%)	150 (95-220)	11000 (7600-16000)
Malaysia	1995	Male	30 to 34	13% (7.8-20%)	87% (80-92%)	2.7 (1.5-4.5)	7.2% (4.9-9.8%)	10% (6.7-14%)	170 (110-240)	12000 (8000-16000)
Malaysia	1995	Male	35 to 39	13% (8.3-20%)	87% (80-92%)	3.2 (1.7-5.1)	7.5% (5.4-10%)	11% (7.4-14%)	190 (130-260)	12000 (8500-16000)
Malaysia	1995	Male	40 to 44	13% (8.5-19%)	87% (81-92%)	3.2 (1.7-5.3)	7.6% (5.5-10%)	11% (7.6-14%)	220 (160-290)	12000 (8500-15000)
Malaysia	1995	Male	45 to 49	13% (7.6-18%)	87% (82-92%)	2.9 (1.5-4.9)	7.7% (5.5-10%)	10% (7.4-14%)	250 (180-340)	12000 (8300-15000)
Malaysia	1995	Male	50 to 54	13% (8.2-18%)	87% (82-92%)	2.9 (1.5-4.9)	7.4% (5.3-10%)	9.5% (6.7-13%)	290 (200-390)	11000 (8100-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	1995	Male	55 to 59	12% (8.1-18%)	88% (82-92%)	3.1 (1.5-5.4)	7.1% (4.9-9.7%)	8.7% (6.1-12%)	350 (240-490)	12000 (8300-16000)
Malaysia	1995	Male	60 to 64	11% (7-17%)	89% (83-93%)	2.7 (1.3-4.8)	6.3% (4.3-8.9%)	7.5% (5.1-11%)	350 (240-490)	9900 (6800-14000)
Malaysia	1995	Male	65 to 69	9.7% (6-14%)	90% (86-94%)	2 (1-3.5)	5.4% (3.6-7.6%)	6.3% (4.2-8.9%)	340 (230-480)	7900 (5300-11000)
Malaysia	1995	Male	70 to 74	8.2% (5.4-12%)	92% (88-95%)	1.5 (0.7-2.6)	4.5% (3-6.4%)	5.1% (3.4-7.4%)	280 (190-410)	5300 (3500-7600)
Malaysia	1995	Male	75 to 79	7.3% (4.5-11%)	93% (89-96%)	1.2 (0.53-2.4)	3.6% (2.2-5.4%)	4.1% (2.5-6.2%)	200 (130-310)	3000 (1900-4500)
Malaysia	1995	Male	80 to 84	6.6% (3.9-9.8%)	93% (90-96%)	1 (0.47-1.9)	3.3% (2-4.9%)	3.7% (2.3-5.6%)	140 (87-220)	1600 (950-2300)
Malaysia	1995	Male	85 to 89	6.4% (3.7-9.9%)	94% (90-96%)	0.93 (0.44-1.7)	3% (1.8-4.6%)	3.4% (2-5.3%)	86 (50-130)	650 (390-1000)
Malaysia	1995	Male	90 to 94	6.9% (4.2-11%)	93% (89-96%)	0.95 (0.42-1.9)	2.8% (1.6-4.3%)	3.2% (1.8-5%)	35 (20-55)	180 (110-280)
Malaysia	1995	Male	95 plus	8% (4.4-13%)	92% (87-96%)	1 (0.38-2.3)	3% (1.5-4.9%)	3.3% (1.6-5.6%)	8.3 (4.1-14)	30 (15-50)
Malaysia	2000	Female	All Ages	2.3% (1.9-2.8%)	98% (97-98%)	0.23 (0.16-0.3)	1.3% (0.97-1.7%)	2.3% (1.6-3.1%)	970 (680-1300)	27000 (21000-35000)
Malaysia	2000	Female	15 to 19	2.3% (1.4-3.4%)	98% (97-99%)	0.22 (0.11-0.39)	0.73% (0.48-1.1%)	1.4% (0.84-2.3%)	6.8 (3.9-11)	860 (560-1200)
Malaysia	2000	Female	20 to 24	2.5% (1.5-3.9%)	97% (96-99%)	0.26 (0.13-0.48)	1.1% (0.72-1.5%)	1.6% (0.93-2.4%)	8.1 (4.8-13)	1200 (810-1800)
Malaysia	2000	Female	25 to 29	2.5% (1.4-3.9%)	97% (96-99%)	0.29 (0.14-0.5)	1.1% (0.77-1.5%)	1.6% (0.98-2.5%)	8.8 (5.2-13)	1300 (840-1800)
Malaysia	2000	Female	30 to 34	2.5% (1.4-4.1%)	98% (96-99%)	0.29 (0.15-0.51)	1.1% (0.74-1.6%)	1.7% (1.1-2.6%)	12 (7.2-18)	1300 (900-1900)
Malaysia	2000	Female	35 to 39	2.7% (1.4-4.1%)	97% (96-99%)	0.3 (0.14-0.51)	1.1% (0.74-1.5%)	1.9% (1.1-2.8%)	17 (10-26)	1500 (990-2100)
Malaysia	2000	Female	40 to 44	3% (1.8-4.5%)	97% (96-98%)	0.36 (0.18-0.6)	1.3% (0.85-1.8%)	2.2% (1.4-3.2%)	27 (17-41)	1800 (1200-2600)
Malaysia	2000	Female	45 to 49	2.9% (1.7-4.4%)	97% (96-98%)	0.43 (0.2-0.77)	1.5% (0.99-2.2%)	2.4% (1.5-3.7%)	40 (25-61)	2100 (1400-3200)
Malaysia	2000	Female	50 to 54	3% (1.9-4.6%)	97% (95-98%)	0.48 (0.23-0.85)	1.8% (1.2-2.6%)	2.7% (1.7-4%)	58 (37-85)	2500 (1600-3600)
Malaysia	2000	Female	55 to 59	3.5% (2.2-5.1%)	97% (95-98%)	0.54 (0.27-0.98)	2% (1.3-3%)	2.9% (1.8-4.3%)	71 (45-110)	2500 (1600-3700)
Malaysia	2000	Female	60 to 64	3.5% (2.1-5.3%)	97% (95-98%)	0.55 (0.27-0.98)	2.1% (1.3-3.2%)	2.7% (1.7-4.2%)	100 (62-150)	2900 (1900-4500)
Malaysia	2000	Female	65 to 69	3.2% (1.9-4.8%)	97% (95-98%)	0.52 (0.25-0.94)	2.1% (1.3-3.1%)	2.6% (1.6-4%)	120 (71-180)	2800 (1700-4300)
Malaysia	2000	Female	70 to 74	3.3% (2.1-5%)	97% (95-98%)	0.54 (0.25-0.98)	2% (1.2-3.4%)	2.5% (1.5-4.1%)	140 (81-230)	2600 (1500-4200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	2000	Female	75 to 79	3.5% (2.1-5.4%)	97% (95-98%)	0.57 (0.25-1.1)	2% (1.1-3.4%)	2.4% (1.3-4%)	140 (74-230)	2000 (1100-3200)
Malaysia	2000	Female	80 to 84	3.4% (1.9-5.1%)	97% (95-98%)	0.55 (0.24-0.98)	2% (1-3.3%)	2.4% (1.2-3.9%)	100 (53-170)	1100 (570-1800)
Malaysia	2000	Female	85 to 89	3.5% (2-5.4%)	97% (95-98%)	0.54 (0.25-0.96)	1.9% (1-3.2%)	2.3% (1.2-3.9%)	74 (39-130)	560 (300-950)
Malaysia	2000	Female	90 to 94	3.8% (2.3-6%)	96% (94-98%)	0.57 (0.25-1.1)	1.9% (0.96-3.3%)	2.2% (1.1-4%)	35 (18-62)	180 (92-320)
Malaysia	2000	Female	95 plus	4.3% (2.5-6.8%)	96% (93-98%)	0.63 (0.26-1.3)	2% (0.94-3.6%)	2.3% (1.1-4.2%)	11 (5.1-20)	39 (18-68)
Malaysia	2000	Male	All Ages	8.2% (7.1-9.5%)	92% (90-93%)	1 (0.74-1.3)	4.8% (3.9-5.8%)	6% (4.7-7.3%)	3500 (2800-4400)	140000 (110000-170000)
Malaysia	2000	Male	15 to 19	6.5% (4.1-9.6%)	94% (90-96%)	0.42 (0.2-0.76)	2.6% (1.5-4%)	3.7% (1.9-6.1%)	53 (27-87)	4700 (2700-7200)
Malaysia	2000	Male	20 to 24	10% (6.8-14%)	90% (86-93%)	1.1 (0.58-1.8)	5.6% (3.7-8.2%)	7.4% (4.4-12%)	120 (72-200)	10000 (6500-15000)
Malaysia	2000	Male	25 to 29	11% (7.8-15%)	89% (85-92%)	1.4 (0.78-2.3)	5.9% (4.2-8.4%)	8% (5.3-12%)	130 (88-200)	10000 (7200-14000)
Malaysia	2000	Male	30 to 34	12% (8.2-17%)	88% (83-92%)	1.7 (0.97-2.7)	6.1% (4.4-8.3%)	8.2% (5.7-11%)	160 (110-220)	11000 (8100-15000)
Malaysia	2000	Male	35 to 39	12% (8.1-17%)	88% (83-92%)	1.9 (1.1-3.1)	6.3% (4.7-8.6%)	8.7% (6.1-12%)	200 (140-270)	12000 (9100-17000)
Malaysia	2000	Male	40 to 44	13% (9.1-17%)	87% (83-91%)	2.1 (1.1-3.3)	6.7% (5-8.8%)	9.2% (6.8-12%)	250 (190-330)	13000 (10000-17000)
Malaysia	2000	Male	45 to 49	14% (9.5-18%)	86% (82-91%)	2.1 (1.2-3.4)	7.3% (5.5-9.2%)	9.8% (7.3-12%)	310 (230-390)	14000 (11000-18000)
Malaysia	2000	Male	50 to 54	14% (9.5-19%)	86% (81-90%)	2.1 (1.1-3.5)	7.2% (5.3-9.5%)	9.2% (6.7-12%)	350 (260-470)	14000 (10000-18000)
Malaysia	2000	Male	55 to 59	15% (10-20%)	85% (80-90%)	2.5 (1.3-4.2)	7.5% (5.2-9.9%)	9.2% (6.5-12%)	410 (280-540)	14000 (9800-18000)
Malaysia	2000	Male	60 to 64	14% (9.5-20%)	86% (80-91%)	2.3 (1.2-3.8)	6.7% (4.7-9%)	8% (5.6-11%)	470 (330-640)	13000 (9600-18000)
Malaysia	2000	Male	65 to 69	10% (6.9-14%)	90% (86-93%)	1.4 (0.78-2.4)	5.2% (3.5-7.1%)	6% (4.1-8.3%)	360 (250-500)	8500 (5900-12000)
Malaysia	2000	Male	70 to 74	8.4% (5.6-12%)	92% (88-94%)	1 (0.51-1.8)	3.9% (2.4-5.7%)	4.5% (2.8-6.5%)	290 (180-420)	5500 (3400-7900)
Malaysia	2000	Male	75 to 79	7.6% (4.9-11%)	92% (89-95%)	0.84 (0.39-1.6)	3.1% (1.7-4.8%)	3.5% (1.8-5.5%)	190 (100-300)	2900 (1600-4500)
Malaysia	2000	Male	80 to 84	6.7% (4-9.7%)	93% (90-96%)	0.7 (0.33-1.3)	2.7% (1.5-4.1%)	3.1% (1.6-4.8%)	120 (64-190)	1300 (720-2000)
Malaysia	2000	Male	85 to 89	6.3% (3.6-9.8%)	94% (90-96%)	0.61 (0.28-1.1)	2.3% (1.2-3.8%)	2.7% (1.3-4.4%)	78 (39-130)	610 (310-1000)
Malaysia	2000	Male	90 to 94	6.7% (4.1-11%)	93% (89-96%)	0.62 (0.27-1.2)	2.2% (0.9-3.7%)	2.5% (0.99-4.3%)	32 (13-57)	170 (73-300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	2000	Male	95 plus	7.8% (4.3-13%)	92% (87-96%)	0.67 (0.25-1.4)	2.3% (0.83-4.1%)	2.5% (0.82-4.7%)	8.9 (3-17)	32 (12-60)
Malaysia	2005	Female	All Ages	2.3% (1.9-2.8%)	98% (97-98%)	0.2 (0.14-0.27)	1.2% (0.89-1.6%)	2.1% (1.5-3%)	1000 (700-1500)	29000 (21000-37000)
Malaysia	2005	Female	15 to 19	2.2% (1.4-3.4%)	98% (97-99%)	0.18 (0.085-0.34)	0.63% (0.4-0.94%)	1.3% (0.73-2.2%)	6.1 (3.5-10)	810 (520-1200)
Malaysia	2005	Female	20 to 24	2.5% (1.5-3.8%)	97% (96-99%)	0.22 (0.11-0.4)	0.95% (0.64-1.4%)	1.4% (0.79-2.2%)	7.5 (4.3-12)	1200 (790-1800)
Malaysia	2005	Female	25 to 29	2.5% (1.4-3.9%)	98% (96-99%)	0.24 (0.12-0.42)	0.99% (0.69-1.4%)	1.4% (0.84-2.2%)	8.2 (4.8-13)	1300 (850-1800)
Malaysia	2005	Female	30 to 34	2.4% (1.3-4.1%)	98% (96-99%)	0.24 (0.12-0.43)	0.98% (0.66-1.4%)	1.5% (0.92-2.4%)	10 (6.3-16)	1300 (830-1800)
Malaysia	2005	Female	35 to 39	2.6% (1.4-4.2%)	97% (96-99%)	0.25 (0.11-0.45)	0.96% (0.64-1.4%)	1.6% (0.92-2.5%)	15 (8.6-23)	1400 (910-2000)
Malaysia	2005	Female	40 to 44	2.9% (1.8-4.3%)	97% (96-98%)	0.3 (0.15-0.5)	1.1% (0.73-1.6%)	1.9% (1.2-2.9%)	26 (15-39)	1700 (1100-2500)
Malaysia	2005	Female	45 to 49	2.9% (1.7-4.3%)	97% (96-98%)	0.36 (0.18-0.63)	1.4% (0.88-2%)	2.2% (1.4-3.4%)	43 (27-66)	2300 (1500-3500)
Malaysia	2005	Female	50 to 54	3% (1.9-4.6%)	97% (95-98%)	0.41 (0.19-0.72)	1.6% (1-2.4%)	2.5% (1.5-3.7%)	63 (39-96)	2700 (1700-4100)
Malaysia	2005	Female	55 to 59	3.4% (2.1-5%)	97% (95-98%)	0.46 (0.23-0.82)	1.8% (1.1-2.7%)	2.6% (1.6-3.9%)	82 (50-120)	2900 (1800-4400)
Malaysia	2005	Female	60 to 64	3.4% (2.1-5.2%)	97% (95-98%)	0.46 (0.22-0.83)	1.8% (1.1-2.8%)	2.4% (1.5-3.8%)	95 (56-150)	2900 (1700-4500)
Malaysia	2005	Female	65 to 69	3.2% (2-4.8%)	97% (95-98%)	0.44 (0.22-0.78)	1.9% (1.1-2.8%)	2.4% (1.4-3.6%)	120 (72-190)	2900 (1700-4400)
Malaysia	2005	Female	70 to 74	3.3% (2.1-5%)	97% (95-98%)	0.46 (0.21-0.83)	1.8% (1-3%)	2.3% (1.3-3.7%)	140 (78-230)	2700 (1500-4400)
Malaysia	2005	Female	75 to 79	3.5% (2.1-5.4%)	97% (95-98%)	0.49 (0.21-0.93)	1.9% (0.94-3.2%)	2.2% (1.1-3.8%)	140 (71-240)	2100 (1100-3500)
Malaysia	2005	Female	80 to 84	3.5% (1.9-5.2%)	97% (95-98%)	0.47 (0.21-0.84)	1.9% (0.94-3.1%)	2.2% (1.1-3.6%)	120 (60-200)	1300 (640-2100)
Malaysia	2005	Female	85 to 89	3.4% (2-5.4%)	97% (95-98%)	0.45 (0.21-0.82)	1.8% (0.93-3.2%)	2.1% (1-3.7%)	89 (44-150)	660 (340-1200)
Malaysia	2005	Female	90 to 94	3.8% (2.2-5.8%)	96% (94-98%)	0.48 (0.21-0.93)	1.9% (0.87-3.3%)	2.1% (0.97-3.9%)	49 (22-88)	250 (120-450)
Malaysia	2005	Female	95 plus	4.3% (2.4-7%)	96% (93-98%)	0.54 (0.21-1.1)	2% (0.76-3.7%)	2.2% (0.84-4.1%)	15 (5.5-28)	49 (19-91)
Malaysia	2005	Male	All Ages	8.3% (7.1-9.6%)	92% (90-93%)	0.89 (0.63-1.2)	4.4% (3.5-5.5%)	5.4% (4.1-6.8%)	3600 (2800-4600)	140000 (110000-170000)
Malaysia	2005	Male	15 to 19	6.3% (4-9.2%)	94% (91-96%)	0.35 (0.16-0.63)	2.2% (1.3-3.3%)	3.3% (1.7-5.4%)	47 (24-76)	4200 (2500-6300)
Malaysia	2005	Male	20 to 24	9.9% (6.6-14%)	90% (86-93%)	0.89 (0.47-1.5)	4.8% (3.1-7.3%)	6.5% (3.7-10%)	110 (64-190)	9700 (6200-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	2005	Male	25 to 29	11% (7.5-16%)	89% (84-92%)	1.2 (0.64-2)	5.1% (3.5-7.3%)	7% (4.5-10%)	120 (79-180)	9900 (6800-14000)
Malaysia	2005	Male	30 to 34	12% (8-17%)	88% (83-92%)	1.4 (0.81-2.2)	5.4% (3.8-7.4%)	7.3% (4.9-10%)	150 (99-210)	10000 (7500-14000)
Malaysia	2005	Male	35 to 39	12% (7.9-17%)	88% (83-92%)	1.6 (0.88-2.6)	5.8% (4.1-7.9%)	7.9% (5.3-11%)	190 (120-260)	11000 (8000-15000)
Malaysia	2005	Male	40 to 44	12% (8.7-17%)	88% (83-91%)	1.7 (0.91-2.8)	6.1% (4.4-8.2%)	8.4% (5.9-11%)	240 (170-330)	13000 (9400-17000)
Malaysia	2005	Male	45 to 49	14% (9.2-18%)	86% (82-91%)	1.8 (0.98-2.9)	6.6% (4.8-8.7%)	8.9% (6.3-11%)	330 (230-420)	15000 (11000-19000)
Malaysia	2005	Male	50 to 54	13% (9.1-18%)	87% (82-91%)	1.8 (0.92-2.8)	6.5% (4.5-8.7%)	8.4% (5.9-11%)	380 (260-510)	15000 (11000-20000)
Malaysia	2005	Male	55 to 59	15% (10-19%)	85% (81-90%)	2 (1.1-3.5)	6.9% (4.7-9.3%)	8.5% (5.8-12%)	460 (310-620)	16000 (11000-21000)
Malaysia	2005	Male	60 to 64	14% (9.1-19%)	86% (81-91%)	1.9 (0.95-3.1)	6.1% (4.1-8.2%)	7.4% (5-10%)	470 (320-650)	14000 (9000-19000)
Malaysia	2005	Male	65 to 69	10% (6.8-14%)	90% (86-93%)	1.2 (0.63-2)	4.7% (2.9-6.5%)	5.5% (3.4-7.7%)	400 (250-560)	9500 (6000-13000)
Malaysia	2005	Male	70 to 74	8.2% (5.4-12%)	92% (88-95%)	0.85 (0.42-1.5)	3.4% (2-5.1%)	4% (2.2-6%)	300 (160-450)	5700 (3200-8500)
Malaysia	2005	Male	75 to 79	7.4% (4.7-11%)	93% (89-95%)	0.7 (0.31-1.3)	2.6% (1.3-4.3%)	3% (1.4-5%)	210 (93-330)	3100 (1400-4900)
Malaysia	2005	Male	80 to 84	6.5% (4-9.5%)	93% (90-96%)	0.58 (0.27-1)	2.3% (1.1-3.6%)	2.6% (1.2-4.2%)	120 (56-190)	1300 (640-2100)
Malaysia	2005	Male	85 to 89	6.1% (3.5-9.6%)	94% (90-97%)	0.5 (0.24-0.91)	2% (0.85-3.4%)	2.2% (0.93-3.9%)	75 (31-130)	580 (250-1000)
Malaysia	2005	Male	90 to 94	6.5% (3.9-9.8%)	94% (90-96%)	0.5 (0.22-0.95)	1.8% (0.69-3.4%)	2% (0.73-3.9%)	38 (14-72)	200 (78-380)
Malaysia	2005	Male	95 plus	7.5% (4.1-12%)	93% (88-96%)	0.54 (0.2-1.1)	1.9% (0.63-3.8%)	2% (0.61-4.1%)	9.9 (3-20)	36 (12-70)
Malaysia	2010	Female	All Ages	2.7% (2.2-3.2%)	97% (97-98%)	0.28 (0.19-0.39)	1.3% (0.99-1.7%)	2.2% (1.5-3.2%)	1200 (840-1700)	34000 (26000-45000)
Malaysia	2010	Female	15 to 19	2.3% (1.4-3.6%)	98% (96-99%)	0.22 (0.095-0.41)	0.64% (0.41-0.97%)	1.4% (0.77-2.4%)	6.7 (3.7-11)	930 (590-1400)
Malaysia	2010	Female	20 to 24	2.8% (1.6-4.4%)	97% (96-98%)	0.29 (0.13-0.52)	1% (0.7-1.5%)	1.6% (0.93-2.6%)	9.3 (5.3-15)	1500 (970-2200)
Malaysia	2010	Female	25 to 29	3.1% (1.7-4.8%)	97% (95-98%)	0.34 (0.17-0.61)	1.1% (0.77-1.6%)	1.8% (1.1-2.8%)	12 (7-18)	1700 (1100-2500)
Malaysia	2010	Female	30 to 34	3.2% (1.8-5.4%)	97% (95-98%)	0.37 (0.19-0.65)	1.2% (0.81-1.6%)	2% (1.3-3.1%)	14 (8.5-21)	1600 (1000-2300)
Malaysia	2010	Female	35 to 39	3.4% (1.9-5.3%)	97% (95-98%)	0.37 (0.18-0.65)	1.2% (0.76-1.7%)	2.1% (1.3-3.2%)	19 (11-29)	1700 (1100-2400)
Malaysia	2010	Female	40 to 44	3.4% (2.1-5.3%)	97% (95-98%)	0.41 (0.2-0.71)	1.3% (0.84-1.9%)	2.3% (1.5-3.6%)	31 (19-48)	2100 (1400-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	2010	Female	45 to 49	3.4% (1.9-5.4%)	97% (95-98%)	0.49 (0.22-0.88)	1.5% (0.97-2.3%)	2.6% (1.6-4.1%)	54 (33-85)	2900 (1800-4400)
Malaysia	2010	Female	50 to 54	3.5% (2.1-5.5%)	96% (95-98%)	0.55 (0.25-0.99)	1.8% (1.1-2.7%)	2.8% (1.7-4.2%)	82 (51-120)	3600 (2300-5400)
Malaysia	2010	Female	55 to 59	3.7% (2.2-5.6%)	96% (94-98%)	0.56 (0.26-1)	1.9% (1.2-2.9%)	2.8% (1.7-4.4%)	100 (63-160)	3800 (2400-5700)
Malaysia	2010	Female	60 to 64	3.5% (2-5.6%)	96% (94-98%)	0.54 (0.25-1)	1.8% (1.1-2.9%)	2.6% (1.5-4%)	110 (70-180)	3500 (2100-5500)
Malaysia	2010	Female	65 to 69	3.3% (1.9-5%)	97% (95-98%)	0.52 (0.23-0.93)	1.9% (1.1-2.9%)	2.5% (1.4-3.8%)	130 (74-200)	3100 (1800-4800)
Malaysia	2010	Female	70 to 74	3.2% (2-4.8%)	97% (95-98%)	0.51 (0.22-0.97)	1.8% (1-3.1%)	2.3% (1.3-3.9%)	160 (91-280)	3100 (1700-5200)
Malaysia	2010	Female	75 to 79	3.2% (1.8-5.1%)	97% (95-98%)	0.51 (0.21-1)	1.8% (0.93-3.1%)	2.1% (1.1-3.6%)	150 (76-250)	2200 (1100-3700)
Malaysia	2010	Female	80 to 84	3.3% (1.8-4.9%)	97% (95-98%)	0.51 (0.22-0.97)	1.9% (0.95-3%)	2.2% (1.1-3.6%)	140 (68-220)	1400 (730-2400)
Malaysia	2010	Female	85 to 89	3.5% (2-5.6%)	97% (94-98%)	0.53 (0.25-0.98)	1.9% (1-3.4%)	2.2% (1.2-3.9%)	110 (57-200)	840 (440-1400)
Malaysia	2010	Female	90 to 94	3.8% (2.3-6.1%)	96% (94-98%)	0.56 (0.24-1.1)	2% (1-3.5%)	2.3% (1.1-4%)	52 (26-93)	270 (140-470)
Malaysia	2010	Female	95 plus	4.3% (2.5-6.8%)	96% (93-98%)	0.62 (0.24-1.3)	2.1% (0.92-3.7%)	2.3% (1-4.2%)	13 (5.6-23)	44 (19-78)
Malaysia	2010	Male	All Ages	8.6% (7.2-10%)	91% (90-93%)	1 (0.72-1.4)	4.4% (3.5-5.4%)	5.3% (4-6.5%)	4100 (3100-5100)	150000 (120000-190000)
Malaysia	2010	Male	15 to 19	6.7% (4.1-9.8%)	93% (90-96%)	0.42 (0.19-0.78)	2.4% (1.4-3.7%)	3.7% (1.9-6.2%)	52 (27-87)	4600 (2700-7200)
Malaysia	2010	Male	20 to 24	10% (6.3-15%)	90% (85-94%)	1.1 (0.53-1.8)	5.1% (3.3-7.7%)	7.2% (4-12%)	130 (74-220)	11000 (7200-17000)
Malaysia	2010	Male	25 to 29	12% (7.2-17%)	88% (83-93%)	1.4 (0.73-2.4)	5.6% (3.8-7.9%)	8% (5-12%)	150 (93-220)	12000 (8100-17000)
Malaysia	2010	Male	30 to 34	13% (7.7-19%)	87% (81-92%)	1.7 (0.94-2.8)	5.8% (4-8%)	8% (5.3-12%)	170 (110-240)	12000 (8400-16000)
Malaysia	2010	Male	35 to 39	13% (7.9-19%)	87% (81-92%)	2 (1-3.4)	6.2% (4.3-8.5%)	8.6% (5.7-12%)	210 (140-290)	13000 (8800-18000)
Malaysia	2010	Male	40 to 44	13% (8.1-19%)	87% (81-92%)	2 (1-3.4)	6.3% (4.4-8.4%)	8.7% (5.9-12%)	260 (180-350)	14000 (9900-19000)
Malaysia	2010	Male	45 to 49	13% (7.4-18%)	87% (82-93%)	1.9 (0.94-3.1)	6.4% (4.5-8.7%)	8.6% (5.9-11%)	350 (240-460)	16000 (11000-21000)
Malaysia	2010	Male	50 to 54	12% (7.7-18%)	88% (82-92%)	1.8 (0.91-3.1)	6.2% (4.2-8.5%)	8% (5.4-11%)	420 (290-590)	17000 (12000-23000)
Malaysia	2010	Male	55 to 59	12% (7.5-17%)	88% (83-92%)	1.9 (0.91-3.5)	6% (4-8.3%)	7.5% (4.9-10%)	470 (320-670)	16000 (11000-23000)
Malaysia	2010	Male	60 to 64	11% (6.8-16%)	89% (84-93%)	1.7 (0.79-3)	5.3% (3.4-7.4%)	6.4% (4.1-9%)	490 (320-690)	14000 (9200-20000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	2010	Male	65 to 69	9.3% (5.8-13%)	91% (87-94%)	1.3 (0.6-2.1)	4.5% (2.9-6.4%)	5.3% (3.4-7.6%)	420 (270-610)	10000 (6500-14000)
Malaysia	2010	Male	70 to 74	8% (5.1-12%)	92% (88-95%)	0.94 (0.44-1.7)	3.6% (2.1-5.2%)	4.2% (2.5-6.1%)	390 (230-580)	7500 (4500-11000)
Malaysia	2010	Male	75 to 79	7.1% (4.3-11%)	93% (89-96%)	0.76 (0.33-1.5)	2.7% (1.4-4.3%)	3.2% (1.5-5.2%)	250 (120-410)	3800 (1900-6100)
Malaysia	2010	Male	80 to 84	6.4% (3.7-9.3%)	94% (91-96%)	0.65 (0.3-1.3)	2.4% (1.2-3.7%)	2.8% (1.4-4.4%)	170 (82-260)	1900 (960-2900)
Malaysia	2010	Male	85 to 89	6.1% (3.4-9.5%)	94% (91-97%)	0.57 (0.26-1.1)	2.1% (0.99-3.4%)	2.4% (1.1-4%)	100 (47-170)	800 (380-1300)
Malaysia	2010	Male	90 to 94	6.4% (3.9-9.9%)	94% (90-96%)	0.57 (0.24-1.1)	2% (0.8-3.5%)	2.2% (0.84-4%)	45 (17-82)	240 (98-420)
Malaysia	2010	Male	95 plus	7.4% (4-12%)	93% (88-96%)	0.61 (0.23-1.3)	2% (0.75-3.8%)	2.2% (0.71-4.2%)	12 (3.9-23)	43 (16-82)
Malaysia	2016	Female	All Ages	2.7% (2.2-3.3%)	97% (97-98%)	0.36 (0.22-0.56)	1.3% (1-1.8%)	2.3% (1.6-3.1%)	1300 (930-1900)	39000 (29000-51000)
Malaysia	2016	Female	15 to 19	2.3% (1.4-3.5%)	98% (96-99%)	0.27 (0.11-0.51)	0.66% (0.42-0.98%)	1.5% (0.87-2.4%)	6.3 (3.6-10)	920 (590-1300)
Malaysia	2016	Female	20 to 24	2.8% (1.6-4.3%)	97% (96-98%)	0.35 (0.15-0.66)	1% (0.67-1.4%)	1.7% (1-2.6%)	9 (5.4-14)	1500 (1000-2200)
Malaysia	2016	Female	25 to 29	3.1% (1.7-4.8%)	97% (95-98%)	0.41 (0.18-0.76)	1.1% (0.78-1.6%)	1.9% (1.1-3%)	11 (6.7-18)	1800 (1200-2600)
Malaysia	2016	Female	30 to 34	3.2% (1.8-5.3%)	97% (95-98%)	0.44 (0.2-0.84)	1.1% (0.79-1.6%)	2.1% (1.3-3.2%)	17 (9.9-26)	2000 (1300-2900)
Malaysia	2016	Female	35 to 39	3.4% (1.8-5.2%)	97% (95-98%)	0.44 (0.19-0.85)	1.1% (0.76-1.6%)	2.2% (1.3-3.2%)	20 (12-31)	1900 (1200-2700)
Malaysia	2016	Female	40 to 44	3.4% (2-5.3%)	97% (95-98%)	0.49 (0.23-0.9)	1.3% (0.83-1.8%)	2.4% (1.5-3.6%)	30 (19-46)	2200 (1400-3100)
Malaysia	2016	Female	45 to 49	3.4% (1.9-5.3%)	97% (95-98%)	0.6 (0.24-1.2)	1.5% (0.94-2.3%)	2.6% (1.6-4%)	52 (31-81)	2900 (1800-4300)
Malaysia	2016	Female	50 to 54	3.5% (2.1-5.3%)	96% (95-98%)	0.67 (0.28-1.3)	1.7% (1.1-2.6%)	2.8% (1.8-4.2%)	85 (54-130)	3900 (2500-5700)
Malaysia	2016	Female	55 to 59	3.6% (2.2-5.7%)	96% (94-98%)	0.68 (0.29-1.4)	1.8% (1.1-2.8%)	2.8% (1.7-4.2%)	120 (74-190)	4400 (2800-6800)
Malaysia	2016	Female	60 to 64	3.5% (2-5.5%)	97% (94-98%)	0.65 (0.28-1.3)	1.8% (1.1-2.8%)	2.5% (1.5-4%)	130 (81-210)	4200 (2500-6600)
Malaysia	2016	Female	65 to 69	3.2% (1.9-4.9%)	97% (95-98%)	0.62 (0.26-1.2)	1.8% (1.1-2.8%)	2.4% (1.5-3.8%)	170 (100-270)	4300 (2600-6500)
Malaysia	2016	Female	70 to 74	3.1% (1.9-4.8%)	97% (95-98%)	0.61 (0.24-1.3)	1.8% (1-2.9%)	2.3% (1.3-3.7%)	180 (100-290)	3400 (1900-5500)
Malaysia	2016	Female	75 to 79	3.2% (1.9-5%)	97% (95-98%)	0.62 (0.24-1.3)	1.7% (0.92-2.9%)	2.1% (1.1-3.5%)	180 (94-310)	2700 (1500-4600)
Malaysia	2016	Female	80 to 84	3.3% (1.8-4.9%)	97% (95-98%)	0.62 (0.25-1.2)	1.8% (0.95-2.9%)	2.2% (1.1-3.5%)	150 (77-240)	1600 (860-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malaysia	2016	Female	85 to 89	3.5% (2-5.6%)	97% (94-98%)	0.64 (0.27-1.2)	1.8% (0.96-3.2%)	2.2% (1.1-3.7%)	110 (56-190)	840 (430-1400)
Malaysia	2016	Female	90 to 94	3.8% (2.3-6%)	96% (94-98%)	0.69 (0.28-1.4)	1.9% (0.99-3.2%)	2.2% (1.1-3.7%)	58 (30-100)	300 (160-510)
Malaysia	2016	Female	95 plus	4.4% (2.5-7%)	96% (93-98%)	0.77 (0.28-1.7)	2% (0.98-3.5%)	2.3% (1.1-4.1%)	17 (7.7-29)	57 (27-97)
Malaysia	2016	Male	All Ages	8.9% (7.3-11%)	91% (89-93%)	1.3 (0.82-2.1)	4.6% (3.6-5.7%)	5.3% (4.1-6.6%)	4800 (3700-6000)	180000 (140000-220000)
Malaysia	2016	Male	15 to 19	6.5% (4-9.7%)	93% (90-96%)	0.5 (0.21-0.98)	2.5% (1.5-3.9%)	4.1% (2.1-6.8%)	49 (25-81)	4400 (2700-7000)
Malaysia	2016	Male	20 to 24	10% (6.3-15%)	90% (85-94%)	1.3 (0.59-2.4)	5.3% (3.5-7.9%)	7.6% (4.3-12%)	140 (79-210)	11000 (7200-17000)
Malaysia	2016	Male	25 to 29	12% (7-17%)	88% (83-93%)	1.7 (0.81-3.1)	5.9% (4-8.2%)	8.6% (5.4-12%)	170 (110-250)	14000 (9500-19000)
Malaysia	2016	Male	30 to 34	12% (7.5-19%)	88% (81-92%)	2.1 (1-3.7)	6.1% (4.2-8.3%)	8.7% (5.7-12%)	220 (140-310)	15000 (11000-21000)
Malaysia	2016	Male	35 to 39	13% (7.8-19%)	87% (81-92%)	2.4 (1.1-4.5)	6.3% (4.3-8.6%)	9% (6-13%)	230 (150-330)	14000 (10000-20000)
Malaysia	2016	Male	40 to 44	13% (8-18%)	87% (82-92%)	2.5 (1.2-4.5)	6.3% (4.4-8.4%)	8.9% (6.1-12%)	280 (190-370)	15000 (11000-20000)
Malaysia	2016	Male	45 to 49	12% (7.3-18%)	88% (82-93%)	2.3 (1-4.2)	6.4% (4.4-8.6%)	8.7% (5.9-12%)	360 (250-490)	17000 (12000-23000)
Malaysia	2016	Male	50 to 54	12% (7.7-18%)	88% (82-92%)	2.2 (0.99-4.1)	6.1% (4.2-8.3%)	8% (5.6-11%)	450 (320-620)	18000 (13000-25000)
Malaysia	2016	Male	55 to 59	12% (7.3-17%)	88% (83-93%)	2.3 (1-4.4)	5.8% (4-8%)	7.3% (5.1-10%)	550 (380-760)	19000 (13000-26000)
Malaysia	2016	Male	60 to 64	11% (6.6-17%)	89% (83-93%)	2 (0.9-4.2)	5.2% (3.5-7.3%)	6.3% (4.2-9%)	570 (380-790)	16000 (11000-23000)
Malaysia	2016	Male	65 to 69	9.2% (5.7-13%)	91% (87-94%)	1.5 (0.7-2.9)	4.4% (2.9-6.3%)	5.3% (3.4-7.6%)	580 (380-820)	14000 (9100-20000)
Malaysia	2016	Male	70 to 74	7.9% (5.1-12%)	92% (88-95%)	1.1 (0.5-2.2)	3.6% (2.2-5.3%)	4.2% (2.6-6.2%)	430 (270-640)	8200 (5200-12000)
Malaysia	2016	Male	75 to 79	7% (4.3-11%)	93% (89-96%)	0.92 (0.37-1.9)	2.8% (1.5-4.4%)	3.3% (1.8-5.3%)	320 (170-520)	4800 (2700-7600)
Malaysia	2016	Male	80 to 84	6.2% (3.7-9.3%)	94% (91-96%)	0.77 (0.32-1.5)	2.5% (1.4-3.7%)	2.9% (1.5-4.4%)	240 (130-370)	2600 (1400-4100)
Malaysia	2016	Male	85 to 89	5.9% (3.3-9.3%)	94% (91-97%)	0.68 (0.27-1.3)	2.2% (1.1-3.6%)	2.5% (1.2-4.2%)	160 (77-260)	1200 (620-2000)
Malaysia	2016	Male	90 to 94	6.4% (3.9-10%)	94% (90-96%)	0.69 (0.28-1.5)	2.1% (0.94-3.5%)	2.4% (1-4%)	72 (31-120)	380 (170-640)
Malaysia	2016	Male	95 plus	7.4% (4.1-12%)	93% (88-96%)	0.74 (0.27-1.6)	2.2% (0.91-3.9%)	2.4% (0.89-4.4%)	18 (6.7-33)	65 (26-120)
Maldives	1990	Female	All Ages	14% (9.8-18%)	86% (82-90%)	0.1 (0.061-0.15)	0.14% (-0.024-0.31%)	0.027% (-0.34-0.45%)	0.17 (-2.2-3)	61 (-10-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	1990	Female	15 to 19	13% (6.7-23%)	87% (77-93%)	0.099 (0.034-0.21)	0.36% (0.2-0.58%)	0.27% (0.034-0.63%)	0.039 (0.005-0.091)	7.4 (3.9-12)
Maldives	1990	Female	20 to 24	16% (7.5-29%)	84% (71-93%)	0.13 (0.049-0.28)	0.54% (0.29-0.88%)	0.24% (-0.044-0.72%)	0.045 (-0.0081-0.13)	12 (6.4-20)
Maldives	1990	Female	25 to 29	18% (8.1-31%)	82% (69-92%)	0.16 (0.052-0.32)	0.49% (0.23-0.83%)	0.21% (-0.13-0.67%)	0.049 (-0.029-0.15)	11 (5.1-19)
Maldives	1990	Female	30 to 34	19% (8.5-35%)	81% (65-92%)	0.18 (0.062-0.37)	0.46% (0.16-0.8%)	0.25% (-0.16-0.74%)	0.053 (-0.033-0.15)	8.6 (3-15)
Maldives	1990	Female	35 to 39	21% (9.4-35%)	79% (65-91%)	0.19 (0.065-0.39)	0.47% (0.043-0.92%)	0.39% (-0.21-1.1%)	0.068 (-0.041-0.19)	6.5 (0.59-13)
Maldives	1990	Female	40 to 44	21% (10-35%)	79% (65-90%)	0.22 (0.074-0.47)	0.57% (0.13-1.1%)	0.69% (0.014-1.5%)	0.08 (0.0015-0.18)	5.5 (1.2-11)
Maldives	1990	Female	45 to 49	21% (9.8-35%)	79% (65-90%)	0.24 (0.079-0.53)	0.55% (0.016-1.3%)	0.71% (-0.094-1.9%)	0.14 (-0.02-0.39)	7.5 (0.21-18)
Maldives	1990	Female	50 to 54	21% (10-36%)	79% (64-90%)	0.27 (0.092-0.57)	0.43% (-0.26-1.3%)	0.55% (-0.41-1.7%)	0.17 (-0.13-0.55)	7.2 (-4.2-21)
Maldives	1990	Female	55 to 59	21% (11-37%)	79% (63-89%)	0.29 (0.1-0.63)	0.24% (-0.6-1.3%)	0.31% (-0.77-1.7%)	0.12 (-0.3-0.63)	3.9 (-9.7-21)
Maldives	1990	Female	60 to 64	21% (11-36%)	79% (64-89%)	0.29 (0.1-0.67)	-0.065% (-1-1%)	-0.087% (-1.2-1.3%)	-0.037 (-0.52-0.53)	-0.93 (-14-15)
Maldives	1990	Female	65 to 69	21% (10-34%)	79% (66-90%)	0.27 (0.094-0.55)	-0.27% (-1.2-0.64%)	-0.33% (-1.4-0.81%)	-0.15 (-0.68-0.38)	-3.4 (-15-8.5)
Maldives	1990	Female	70 to 74	20% (9.9-34%)	80% (66-90%)	0.25 (0.08-0.54)	-0.24% (-1.1-0.71%)	-0.29% (-1.2-0.86%)	-0.12 (-0.51-0.34)	-2.2 (-9.4-6.2)
Maldives	1990	Female	75 to 79	19% (9.2-34%)	81% (66-91%)	0.23 (0.071-0.58)	-0.34% (-1.1-0.48%)	-0.41% (-1.4-0.59%)	-0.11 (-0.38-0.16)	-1.6 (-5.4-2.2)
Maldives	1990	Female	80 to 84	19% (9.1-34%)	81% (66-91%)	0.24 (0.071-0.55)	-0.49% (-1.4-0.29%)	-0.58% (-1.7-0.33%)	-0.1 (-0.29-0.059)	-1 (-2.9-0.63)
Maldives	1990	Female	85 to 89	21% (11-36%)	79% (64-89%)	0.25 (0.077-0.6)	-0.59% (-1.5-0.14%)	-0.72% (-1.8-0.14%)	-0.054 (-0.14-0.01)	-0.38 (-1-0.092)
Maldives	1990	Female	90 to 94	24% (12-41%)	76% (59-88%)	0.29 (0.079-0.7)	-0.55% (-1.5-0.31%)	-0.69% (-1.8-0.31%)	-0.015 (-0.039-0.0067)	-0.07 (-0.19-0.04)
Maldives	1990	Female	95 plus	29% (13-49%)	71% (51-87%)	0.34 (0.069-0.91)	-0.59% (-1.9-0.45%)	-0.77% (-2.3-0.42%)	-0.0039 (-0.011-0.0022)	-0.012 (-0.037-0.009)
Maldives	1990	Male	All Ages	35% (29-42%)	65% (58-71%)	0.26 (0.17-0.38)	0.29% (-0.2-0.77%)	0.018% (-1-0.95%)	0.13 (-7.5-7.1)	140 (-94-360)
Maldives	1990	Male	15 to 19	34% (21-49%)	66% (51-79%)	0.16 (0.071-0.3)	0.83% (0.47-1.4%)	0.82% (0.23-1.7%)	0.095 (0.028-0.2)	14 (7.7-23)
Maldives	1990	Male	20 to 24	47% (31-63%)	53% (37-69%)	0.35 (0.18-0.59)	1.8% (0.95-3%)	1.7% (0.24-3.5%)	0.21 (0.03-0.44)	29 (14-47)
Maldives	1990	Male	25 to 29	51% (34-68%)	49% (32-66%)	0.47 (0.24-0.77)	2% (0.87-3.3%)	1.8% (-0.16-4.1%)	0.22 (-0.019-0.5)	27 (12-47)
Maldives	1990	Male	30 to 34	53% (35-70%)	47% (30-65%)	0.55 (0.29-0.91)	1.7% (0.37-3.3%)	1.5% (-0.71-4%)	0.21 (-0.1-0.56)	23 (4.5-43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	1990	Male	35 to 39	55% (39-71%)	45% (29-61%)	0.62 (0.31-1)	1.7% (0.019-3.5%)	1.7% (-0.89-4.4%)	0.22 (-0.12-0.58)	18 (0.21-37)
Maldives	1990	Male	40 to 44	55% (40-73%)	45% (27-60%)	0.65 (0.32-1.1)	1.3% (-0.44-3.1%)	1.3% (-1.3-4%)	0.19 (-0.18-0.56)	13 (-4.4-31)
Maldives	1990	Male	45 to 49	54% (36-71%)	46% (29-64%)	0.65 (0.32-1.1)	1.1% (-0.71-2.9%)	1.1% (-1.6-3.7%)	0.23 (-0.32-0.79)	14 (-9.2-38)
Maldives	1990	Male	50 to 54	52% (34-70%)	48% (30-66%)	0.63 (0.31-1.1)	0.49% (-1.3-2.3%)	0.35% (-2.1-2.8%)	0.13 (-0.79-1)	9.5 (-25-43)
Maldives	1990	Male	55 to 59	51% (34-68%)	49% (32-66%)	0.58 (0.29-1)	0.33% (-1.4-2%)	0.22% (-2.1-2.4%)	0.1 (-1.1-1.1)	6.9 (-29-40)
Maldives	1990	Male	60 to 64	48% (31-67%)	52% (33-69%)	0.52 (0.25-0.95)	0.034% (-1.7-1.6%)	-0.1% (-2.3-1.9%)	-0.063 (-1.4-1.1)	0.71 (-36-33)
Maldives	1990	Male	65 to 69	45% (29-63%)	55% (37-71%)	0.45 (0.22-0.77)	-0.26% (-1.7-1.1%)	-0.44% (-2.3-1.3%)	-0.27 (-1.4-0.73)	-4.5 (-29-18)
Maldives	1990	Male	70 to 74	41% (26-59%)	59% (41-74%)	0.38 (0.18-0.69)	-0.49% (-1.8-0.57%)	-0.7% (-2.4-0.62%)	-0.39 (-1.4-0.36)	-6.1 (-23-7.3)
Maldives	1990	Male	75 to 79	38% (23-56%)	62% (44-77%)	0.33 (0.15-0.62)	-0.59% (-1.9-0.32%)	-0.81% (-2.4-0.32%)	-0.36 (-1.1-0.14)	-4.5 (-15-2.4)
Maldives	1990	Male	80 to 84	35% (20-53%)	65% (47-80%)	0.29 (0.13-0.53)	-0.65% (-1.9-0.24%)	-0.81% (-2.2-0.22%)	-0.25 (-0.69-0.069)	-2.4 (-6.8-0.85)
Maldives	1990	Male	85 to 89	34% (20-51%)	66% (49-80%)	0.26 (0.11-0.48)	-0.54% (-1.5-0.18%)	-0.68% (-1.8-0.17%)	-0.11 (-0.29-0.026)	-0.72 (-2-0.25)
Maldives	1990	Male	90 to 94	36% (22-51%)	64% (49-78%)	0.25 (0.11-0.48)	-0.46% (-1.2-0.14%)	-0.61% (-1.5-0.099%)	-0.034 (-0.086-0.0057)	-0.15 (-0.42-0.043)
Maldives	1990	Male	95 plus	39% (22-58%)	61% (42-78%)	0.25 (0.087-0.51)	-0.54% (-1.5-0.1%)	-0.74% (-1.9-0.012%)	-0.012 (-0.03-0.00019)	-0.033 (-0.093-0.0064)
Maldives	1995	Female	All Ages	13% (9.8-18%)	87% (82-90%)	0.15 (0.099-0.21)	0.29% (0.045-0.59%)	0.25% (-0.36-1%)	1.3 (-1.8-5.2)	100 (15-210)
Maldives	1995	Female	15 to 19	13% (6.6-23%)	87% (77-93%)	0.14 (0.049-0.3)	0.47% (0.27-0.74%)	0.46% (0.13-1.1%)	0.041 (0.012-0.09)	8.5 (4.8-14)
Maldives	1995	Female	20 to 24	16% (7.5-28%)	84% (72-92%)	0.19 (0.072-0.4)	0.77% (0.47-1.2%)	0.45% (0.062-1.2%)	0.053 (0.0073-0.14)	14 (8.3-22)
Maldives	1995	Female	25 to 29	17% (8.1-31%)	83% (69-92%)	0.22 (0.081-0.44)	0.74% (0.46-1.1%)	0.41% (0.027-0.98%)	0.056 (0.0035-0.14)	13 (7.7-21)
Maldives	1995	Female	30 to 34	19% (8.6-33%)	81% (67-91%)	0.25 (0.096-0.53)	0.67% (0.39-1%)	0.47% (0.019-1.2%)	0.068 (0.0028-0.17)	11 (6.2-18)
Maldives	1995	Female	35 to 39	20% (8.9-35%)	80% (65-91%)	0.28 (0.1-0.56)	0.66% (0.25-1.2%)	0.66% (-0.015-1.6%)	0.11 (-0.0026-0.27)	11 (4-20)
Maldives	1995	Female	40 to 44	21% (10-35%)	79% (65-90%)	0.31 (0.12-0.67)	0.77% (0.31-1.5%)	1.1% (0.27-2.4%)	0.12 (0.029-0.26)	7.9 (3.2-15)
Maldives	1995	Female	45 to 49	21% (10-34%)	79% (66-90%)	0.36 (0.12-0.74)	0.82% (0.21-1.8%)	1.2% (0.21-3%)	0.15 (0.026-0.36)	7.7 (1.9-17)
Maldives	1995	Female	50 to 54	21% (10-37%)	79% (63-90%)	0.39 (0.14-0.81)	0.78% (-0.1-2%)	1.1% (-0.17-2.8%)	0.29 (-0.048-0.77)	12 (-1.6-31)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	1995	Female	55 to 59	21% (11-37%)	79% (63-89%)	0.43 (0.16-0.88)	0.64% (-0.32-1.9%)	0.88% (-0.45-2.7%)	0.33 (-0.18-1)	11 (-5.9-34)
Maldives	1995	Female	60 to 64	21% (10-37%)	79% (63-90%)	0.43 (0.15-0.9)	0.32% (-0.84-1.8%)	0.4% (-1.1-2.3%)	0.2 (-0.55-1.1)	5.7 (-15-32)
Maldives	1995	Female	65 to 69	20% (9.9-34%)	80% (66-90%)	0.4 (0.14-0.79)	0.076% (-1.1-1.4%)	0.089% (-1.3-1.7%)	0.041 (-0.61-0.77)	1 (-14-18)
Maldives	1995	Female	70 to 74	20% (9.6-33%)	80% (67-90%)	0.36 (0.12-0.79)	0.067% (-0.89-1.5%)	0.081% (-1.1-1.8%)	0.037 (-0.51-0.79)	0.7 (-9.4-15)
Maldives	1995	Female	75 to 79	19% (8.9-32%)	81% (68-91%)	0.34 (0.1-0.81)	-0.11% (-1.1-1.1%)	-0.14% (-1.2-1.3%)	-0.05 (-0.45-0.44)	-0.64 (-6.1-6.4)
Maldives	1995	Female	80 to 84	19% (9.1-33%)	81% (67-91%)	0.34 (0.11-0.76)	-0.25% (-1.3-0.84%)	-0.32% (-1.5-0.91%)	-0.067 (-0.32-0.2)	-0.65 (-3.3-2.1)
Maldives	1995	Female	85 to 89	21% (9.8-36%)	79% (64-90%)	0.37 (0.11-0.82)	-0.3% (-1.2-0.73%)	-0.41% (-1.6-0.83%)	-0.035 (-0.13-0.07)	-0.23 (-0.95-0.57)
Maldives	1995	Female	90 to 94	24% (12-42%)	76% (58-88%)	0.42 (0.12-1)	-0.31% (-1.4-1%)	-0.42% (-1.6-1.1%)	-0.014 (-0.056-0.038)	-0.062 (-0.27-0.21)
Maldives	1995	Female	95 plus	28% (13-49%)	72% (51-87%)	0.49 (0.11-1.3)	-0.34% (-1.7-1.6%)	-0.48% (-2.2-1.7%)	-0.0033 (-0.014-0.012)	-0.0089 (-0.046-0.043)
Maldives	1995	Male	All Ages	34% (28-42%)	66% (58-72%)	0.36 (0.26-0.5)	0.52% (-0.1-1.1%)	0.18% (-1-1.4%)	1.2 (-7-9.4)	210 (-45-450)
Maldives	1995	Male	15 to 19	33% (21-49%)	67% (51-79%)	0.23 (0.11-0.41)	0.99% (0.57-1.6%)	1.1% (0.4-2.2%)	0.12 (0.041-0.24)	17 (9.6-28)
Maldives	1995	Male	20 to 24	46% (31-62%)	54% (38-69%)	0.5 (0.27-0.82)	2.3% (1.3-3.5%)	2.4% (0.8-4.5%)	0.28 (0.092-0.53)	35 (19-54)
Maldives	1995	Male	25 to 29	51% (33-68%)	49% (32-67%)	0.67 (0.37-1)	2.5% (1.4-3.8%)	2.8% (0.7-5.2%)	0.26 (0.068-0.5)	32 (17-49)
Maldives	1995	Male	30 to 34	53% (35-70%)	47% (30-65%)	0.8 (0.46-1.2)	2.3% (1.1-3.8%)	2.5% (0.31-5.1%)	0.28 (0.039-0.57)	30 (13-49)
Maldives	1995	Male	35 to 39	55% (39-72%)	45% (28-61%)	0.89 (0.48-1.4)	2.3% (0.81-3.9%)	2.7% (0.13-5.5%)	0.35 (0.017-0.72)	28 (9.2-48)
Maldives	1995	Male	40 to 44	55% (39-72%)	45% (28-61%)	0.94 (0.5-1.5)	2% (0.41-3.8%)	2.4% (-0.29-5.3%)	0.28 (-0.033-0.64)	19 (3.7-37)
Maldives	1995	Male	45 to 49	53% (36-70%)	47% (30-64%)	0.94 (0.48-1.6)	1.7% (-0.11-3.7%)	2.1% (-0.81-5%)	0.3 (-0.12-0.74)	17 (-0.95-36)
Maldives	1995	Male	50 to 54	52% (35-69%)	48% (31-65%)	0.91 (0.47-1.5)	1% (-0.89-3.1%)	1.1% (-1.7-3.9%)	0.33 (-0.5-1.2)	17 (-14-49)
Maldives	1995	Male	55 to 59	50% (34-67%)	50% (33-66%)	0.84 (0.43-1.4)	0.76% (-1.1-2.7%)	0.75% (-1.7-3.3%)	0.39 (-0.92-1.7)	17 (-24-61)
Maldives	1995	Male	60 to 64	48% (30-66%)	52% (34-70%)	0.75 (0.37-1.3)	0.39% (-1.3-2.5%)	0.32% (-1.9-2.9%)	0.24 (-1.4-2.1)	10 (-34-63)
Maldives	1995	Male	65 to 69	44% (29-62%)	56% (38-71%)	0.65 (0.34-1.1)	-0.039% (-1.5-1.5%)	-0.18% (-2.1-1.8%)	-0.13 (-1.5-1.3)	-0.74 (-32-31)
Maldives	1995	Male	70 to 74	41% (26-58%)	59% (42-74%)	0.56 (0.27-1)	-0.45% (-1.9-0.87%)	-0.66% (-2.4-0.93%)	-0.47 (-1.7-0.66)	-7 (-30-14)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	1995	Male	75 to 79	38% (23-56%)	62% (44-77%)	0.48 (0.21-0.89)	-0.64% (-2.1-0.42%)	-0.88% (-2.7-0.44%)	-0.49 (-1.5-0.24)	-6 (-19-4.1)
Maldives	1995	Male	80 to 84	35% (20-53%)	65% (47-80%)	0.42 (0.19-0.76)	-0.67% (-1.9-0.29%)	-0.87% (-2.4-0.28%)	-0.31 (-0.85-0.096)	-2.9 (-8.4-1.3)
Maldives	1995	Male	85 to 89	34% (19-51%)	66% (49-81%)	0.37 (0.17-0.7)	-0.54% (-1.5-0.19%)	-0.74% (-1.9-0.16%)	-0.13 (-0.35-0.027)	-0.87 (-2.5-0.29)
Maldives	1995	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.36 (0.16-0.67)	-0.46% (-1.2-0.15%)	-0.67% (-1.7-0.098%)	-0.041 (-0.1-0.0059)	-0.18 (-0.47-0.057)
Maldives	1995	Male	95 plus	39% (22-59%)	61% (41-78%)	0.36 (0.13-0.75)	-0.58% (-1.5-0.12%)	-0.86% (-2--0.014%)	-0.013 (-0.031--0.00021)	-0.036 (-0.096-0.0073)
Maldives	2000	Female	All Ages	13% (9.7-18%)	87% (82-90%)	0.19 (0.13-0.27)	0.48% (0.15-0.87%)	0.53% (-0.29-1.6%)	2.4 (-1.3-7.1)	140 (44-260)
Maldives	2000	Female	15 to 19	13% (6.3-22%)	87% (78-94%)	0.18 (0.064-0.36)	0.53% (0.31-0.84%)	0.63% (0.21-1.4%)	0.041 (0.014-0.09)	11 (5.9-17)
Maldives	2000	Female	20 to 24	16% (7.1-28%)	84% (72-93%)	0.24 (0.087-0.5)	0.95% (0.6-1.4%)	0.68% (0.19-1.6%)	0.046 (0.012-0.11)	15 (9.2-24)
Maldives	2000	Female	25 to 29	17% (7.4-30%)	83% (70-93%)	0.28 (0.097-0.57)	0.95% (0.6-1.3%)	0.63% (0.18-1.3%)	0.046 (0.013-0.099)	14 (8.6-21)
Maldives	2000	Female	30 to 34	18% (8.1-33%)	82% (67-92%)	0.31 (0.12-0.66)	0.85% (0.54-1.3%)	0.72% (0.25-1.5%)	0.061 (0.02-0.13)	13 (7.5-20)
Maldives	2000	Female	35 to 39	20% (8.7-34%)	80% (66-91%)	0.34 (0.12-0.68)	0.82% (0.45-1.4%)	0.95% (0.24-2.1%)	0.12 (0.031-0.27)	13 (7-22)
Maldives	2000	Female	40 to 44	20% (9.9-33%)	80% (67-90%)	0.38 (0.14-0.79)	0.9% (0.44-1.6%)	1.4% (0.53-2.8%)	0.18 (0.065-0.35)	13 (6.1-22)
Maldives	2000	Female	45 to 49	20% (9.6-34%)	80% (66-90%)	0.43 (0.15-0.89)	0.97% (0.38-2%)	1.6% (0.49-3.6%)	0.18 (0.054-0.41)	9.9 (3.7-21)
Maldives	2000	Female	50 to 54	21% (10-36%)	79% (64-90%)	0.48 (0.18-0.99)	1.1% (0.098-2.6%)	1.5% (0.11-3.6%)	0.29 (0.021-0.71)	12 (1.1-28)
Maldives	2000	Female	55 to 59	21% (10-35%)	79% (65-90%)	0.53 (0.19-1.1)	0.96% (-0.049-2.5%)	1.4% (-0.1-3.7%)	0.39 (-0.027-1)	14 (-0.77-35)
Maldives	2000	Female	60 to 64	21% (9.9-35%)	79% (65-90%)	0.53 (0.18-1.1)	0.7% (-0.56-2.4%)	0.89% (-0.74-3.1%)	0.49 (-0.4-1.7)	14 (-11-49)
Maldives	2000	Female	65 to 69	20% (9.6-34%)	80% (66-90%)	0.5 (0.18-1)	0.42% (-0.77-2%)	0.52% (-0.97-2.4%)	0.28 (-0.51-1.3)	6.7 (-12-31)
Maldives	2000	Female	70 to 74	19% (9.5-32%)	81% (68-91%)	0.45 (0.16-1)	0.35% (-0.68-2%)	0.43% (-0.84-2.5%)	0.21 (-0.41-1.2)	4 (-7.6-22)
Maldives	2000	Female	75 to 79	19% (8.6-33%)	81% (67-91%)	0.43 (0.13-0.96)	0.13% (-0.92-1.6%)	0.14% (-1.1-1.9%)	0.062 (-0.51-0.87)	1 (-7-13)
Maldives	2000	Female	80 to 84	19% (9-34%)	81% (66-91%)	0.43 (0.13-0.94)	0.0018% (-1-1.4%)	-0.026% (-1.3-1.6%)	-0.0071 (-0.36-0.44)	0.01 (-3.6-4.8)
Maldives	2000	Female	85 to 89	20% (9.6-36%)	80% (64-90%)	0.45 (0.14-1)	-0.062% (-0.99-1.2%)	-0.13% (-1.3-1.4%)	-0.016 (-0.16-0.17)	-0.068 (-1.1-1.4)
Maldives	2000	Female	90 to 94	23% (11-41%)	77% (59-89%)	0.51 (0.14-1.2)	-0.036% (-1.2-1.7%)	-0.11% (-1.5-1.9%)	-0.0062 (-0.084-0.11)	-0.012 (-0.39-0.58)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	2000	Female	95 plus	28% (12-49%)	72% (51-88%)	0.61 (0.14-1.6)	-0.033% (-1.5-2.3%)	-0.15% (-1.9-2.5%)	-0.0018 (-0.023-0.03)	-0.0015 (-0.072-0.11)
Maldives	2000	Male	All Ages	35% (28-42%)	65% (58-72%)	0.47 (0.35-0.61)	0.77% (0.17-1.4%)	0.32% (-0.94-1.6%)	2 (-5.9-10)	260 (54-490)
Maldives	2000	Male	15 to 19	33% (20-49%)	67% (51-80%)	0.28 (0.13-0.5)	1.1% (0.64-1.7%)	1.4% (0.51-2.6%)	0.16 (0.06-0.31)	23 (13-37)
Maldives	2000	Male	20 to 24	46% (30-62%)	54% (38-70%)	0.62 (0.33-1)	2.6% (1.6-3.8%)	3% (1.2-5.1%)	0.32 (0.13-0.57)	41 (25-63)
Maldives	2000	Male	25 to 29	50% (33-67%)	50% (33-67%)	0.83 (0.45-1.3)	2.9% (1.9-4.1%)	3.6% (1.5-6.1%)	0.26 (0.11-0.45)	35 (21-51)
Maldives	2000	Male	30 to 34	53% (35-70%)	47% (30-65%)	0.99 (0.56-1.5)	2.7% (1.6-3.9%)	3.3% (1.1-5.8%)	0.27 (0.092-0.48)	33 (19-48)
Maldives	2000	Male	35 to 39	54% (38-71%)	46% (29-62%)	1.1 (0.59-1.7)	2.7% (1.4-4.1%)	3.5% (1.1-6.1%)	0.39 (0.12-0.69)	34 (17-52)
Maldives	2000	Male	40 to 44	54% (38-72%)	46% (28-62%)	1.2 (0.62-1.8)	2.3% (0.98-3.9%)	3.1% (0.66-5.7%)	0.41 (0.083-0.76)	29 (11-48)
Maldives	2000	Male	45 to 49	53% (35-71%)	47% (29-65%)	1.2 (0.61-1.9)	2.1% (0.46-3.9%)	2.7% (-0.0068-5.6%)	0.36 (-0.00094-0.76)	21 (4.4-39)
Maldives	2000	Male	50 to 54	51% (35-69%)	49% (31-65%)	1.1 (0.6-1.9)	1.4% (-0.58-3.4%)	1.5% (-1.3-4.4%)	0.32 (-0.27-0.93)	16 (-6.9-40)
Maldives	2000	Male	55 to 59	50% (33-67%)	50% (33-67%)	1 (0.55-1.7)	1.1% (-0.69-3.1%)	1.2% (-1.2-3.8%)	0.49 (-0.5-1.6)	20 (-12-58)
Maldives	2000	Male	60 to 64	47% (30-67%)	53% (33-70%)	0.93 (0.47-1.6)	0.7% (-0.99-2.8%)	0.68% (-1.5-3.3%)	0.46 (-1-2.2)	17 (-23-66)
Maldives	2000	Male	65 to 69	44% (28-62%)	56% (38-72%)	0.81 (0.41-1.4)	0.22% (-1.3-1.9%)	0.11% (-1.8-2.2%)	0.093 (-1.6-1.8)	5.5 (-31-45)
Maldives	2000	Male	70 to 74	40% (26-58%)	60% (42-74%)	0.68 (0.34-1.2)	-0.29% (-1.7-1.1%)	-0.5% (-2.3-1.2%)	-0.41 (-1.8-0.98)	-5.3 (-31-21)
Maldives	2000	Male	75 to 79	37% (22-55%)	63% (45-78%)	0.59 (0.26-1.1)	-0.55% (-1.9-0.64%)	-0.79% (-2.5-0.62%)	-0.51 (-1.7-0.39)	-6 (-22-7.3)
Maldives	2000	Male	80 to 84	35% (20-52%)	65% (48-80%)	0.51 (0.22-0.92)	-0.58% (-1.8-0.43%)	-0.8% (-2.2-0.41%)	-0.36 (-1-0.19)	-3.3 (-9.9-2.4)
Maldives	2000	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.46 (0.22-0.85)	-0.45% (-1.4-0.29%)	-0.67% (-1.8-0.27%)	-0.15 (-0.4-0.06)	-0.92 (-2.8-0.6)
Maldives	2000	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.44 (0.2-0.81)	-0.47% (-1.3-0.18%)	-0.68% (-1.7-0.14%)	-0.068 (-0.17-0.013)	-0.29 (-0.78-0.11)
Maldives	2000	Male	95 plus	38% (20-58%)	62% (42-80%)	0.45 (0.16-0.9)	-0.64% (-1.6-0.12%)	-0.95% (-2.2-0.002%)	-0.023 (-0.054-0.000047)	-0.064 (-0.16-0.012)
Maldives	2005	Female	All Ages	14% (10-18%)	86% (82-90%)	0.24 (0.16-0.33)	0.61% (0.28-1.1%)	0.73% (-0.17-2.1%)	3 (-0.69-8.7)	160 (72-300)
Maldives	2005	Female	15 to 19	12% (6.1-22%)	88% (78-94%)	0.2 (0.07-0.41)	0.55% (0.33-0.86%)	0.71% (0.27-1.5%)	0.037 (0.014-0.081)	11 (6.5-18)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	2005	Female	20 to 24	15% (7.1-26%)	85% (74-93%)	0.27 (0.1-0.55)	1% (0.66-1.6%)	0.8% (0.27-1.8%)	0.043 (0.014-0.097)	20 (12-31)
Maldives	2005	Female	25 to 29	16% (7.5-29%)	84% (71-92%)	0.31 (0.1-0.61)	1.1% (0.7-1.5%)	0.78% (0.28-1.6%)	0.036 (0.013-0.075)	16 (9.9-24)
Maldives	2005	Female	30 to 34	18% (7.9-31%)	82% (69-92%)	0.34 (0.12-0.71)	0.93% (0.59-1.4%)	0.88% (0.35-1.8%)	0.053 (0.021-0.11)	14 (8.1-21)
Maldives	2005	Female	35 to 39	19% (8.5-33%)	81% (67-91%)	0.38 (0.14-0.77)	0.88% (0.53-1.4%)	1.2% (0.45-2.4%)	0.094 (0.035-0.19)	13 (7.6-21)
Maldives	2005	Female	40 to 44	19% (9.4-32%)	81% (68-91%)	0.42 (0.15-0.9)	0.93% (0.49-1.6%)	1.6% (0.69-3.3%)	0.17 (0.072-0.34)	14 (7.2-24)
Maldives	2005	Female	45 to 49	20% (8.9-34%)	80% (66-91%)	0.48 (0.17-1)	1% (0.44-2.1%)	1.9% (0.71-4.2%)	0.23 (0.085-0.5)	14 (6-28)
Maldives	2005	Female	50 to 54	20% (9.7-35%)	80% (65-90%)	0.54 (0.19-1.1)	1.1% (0.27-2.5%)	1.8% (0.4-4.2%)	0.3 (0.067-0.69)	13 (3.1-29)
Maldives	2005	Female	55 to 59	20% (10-35%)	80% (65-90%)	0.59 (0.2-1.2)	1.1% (0.13-2.7%)	1.7% (0.19-4.2%)	0.29 (0.033-0.71)	11 (1.2-25)
Maldives	2005	Female	60 to 64	20% (10-35%)	80% (65-90%)	0.6 (0.2-1.2)	0.92% (-0.3-2.9%)	1.3% (-0.41-4%)	0.49 (-0.16-1.5)	15 (-5.1-45)
Maldives	2005	Female	65 to 69	20% (9.5-32%)	80% (68-90%)	0.55 (0.2-1.2)	0.66% (-0.45-2.4%)	0.88% (-0.64-3.1%)	0.47 (-0.33-1.7)	11 (-7.6-41)
Maldives	2005	Female	70 to 74	19% (9.6-32%)	81% (68-90%)	0.5 (0.17-1.1)	0.56% (-0.5-2.5%)	0.72% (-0.62-3.2%)	0.41 (-0.36-1.8)	7.9 (-6.7-35)
Maldives	2005	Female	75 to 79	18% (8.5-32%)	82% (68-91%)	0.47 (0.14-1.1)	0.35% (-0.68-2%)	0.4% (-0.85-2.4%)	0.21 (-0.44-1.2)	3.1 (-6.3-18)
Maldives	2005	Female	80 to 84	18% (8.7-32%)	82% (68-91%)	0.47 (0.15-1)	0.23% (-0.76-1.9%)	0.25% (-0.99-2.2%)	0.089 (-0.35-0.83)	1.1 (-3.5-9)
Maldives	2005	Female	85 to 89	20% (9.8-34%)	80% (66-90%)	0.51 (0.16-1.1)	0.17% (-0.74-1.8%)	0.14% (-1-2.1%)	0.026 (-0.18-0.38)	0.3 (-1.3-3)
Maldives	2005	Female	90 to 94	23% (11-40%)	77% (60-89%)	0.58 (0.16-1.4)	0.23% (-0.83-2.3%)	0.2% (-1.1-2.7%)	0.017 (-0.096-0.23)	0.12 (-0.46-1.2)
Maldives	2005	Female	95 plus	27% (12-47%)	73% (53-88%)	0.68 (0.15-1.8)	0.26% (-1.2-2.7%)	0.18% (-1.5-3%)	0.0043 (-0.035-0.07)	0.025 (-0.11-0.26)
Maldives	2005	Male	All Ages	36% (29-44%)	64% (56-71%)	0.57 (0.42-0.76)	0.96% (0.31-1.7%)	0.39% (-0.82-1.8%)	2.3 (-5-11)	300 (91-520)
Maldives	2005	Male	15 to 19	32% (20-48%)	68% (52-80%)	0.3 (0.14-0.55)	1.1% (0.65-1.7%)	1.5% (0.58-2.9%)	0.15 (0.06-0.31)	25 (14-39)
Maldives	2005	Male	20 to 24	45% (28-62%)	55% (38-72%)	0.69 (0.36-1.1)	2.6% (1.7-3.8%)	3.2% (1.5-5.5%)	0.35 (0.17-0.61)	51 (32-76)
Maldives	2005	Male	25 to 29	50% (32-67%)	50% (33-68%)	0.92 (0.48-1.4)	3% (2-4.1%)	3.8% (1.7-6.2%)	0.29 (0.13-0.49)	41 (26-59)
Maldives	2005	Male	30 to 34	52% (34-70%)	48% (30-66%)	1.1 (0.61-1.7)	2.8% (1.8-3.9%)	3.7% (1.6-6.2%)	0.26 (0.11-0.44)	35 (22-50)
Maldives	2005	Male	35 to 39	54% (37-70%)	46% (30-63%)	1.2 (0.69-1.9)	2.8% (1.7-4%)	3.9% (1.6-6.4%)	0.37 (0.16-0.61)	35 (20-51)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	2005	Male	40 to 44	54% (37-71%)	46% (29-63%)	1.3 (0.69-2.1)	2.4% (1.2-3.9%)	3.4% (0.99-6%)	0.42 (0.13-0.73)	32 (15-51)
Maldives	2005	Male	45 to 49	52% (33-69%)	48% (31-67%)	1.3 (0.69-2.2)	2.2% (0.74-4%)	3% (0.39-6.1%)	0.5 (0.063-0.99)	30 (9.5-57)
Maldives	2005	Male	50 to 54	51% (32-68%)	49% (32-68%)	1.2 (0.65-2)	1.5% (-0.18-3.6%)	1.8% (-0.96-4.9%)	0.36 (-0.19-0.95)	19 (-1.9-44)
Maldives	2005	Male	55 to 59	49% (32-66%)	51% (34-68%)	1.1 (0.59-2)	1.3% (-0.35-3.3%)	1.5% (-0.95-4.3%)	0.39 (-0.25-1.1)	16 (-4.2-41)
Maldives	2005	Male	60 to 64	46% (30-64%)	54% (36-70%)	1 (0.5-1.8)	0.9% (-0.73-2.9%)	0.93% (-1.2-3.5%)	0.46 (-0.61-1.7)	17 (-13-53)
Maldives	2005	Male	65 to 69	43% (27-60%)	57% (40-73%)	0.88 (0.46-1.5)	0.41% (-0.96-2.1%)	0.33% (-1.5-2.5%)	0.26 (-1.2-1.9)	9.6 (-23-49)
Maldives	2005	Male	70 to 74	39% (25-57%)	61% (43-75%)	0.74 (0.35-1.4)	-0.13% (-1.5-1.3%)	-0.33% (-2.1-1.3%)	-0.32 (-2-1.3)	-2.9 (-34-28)
Maldives	2005	Male	75 to 79	36% (22-53%)	64% (47-78%)	0.64 (0.28-1.2)	-0.42% (-1.7-0.78%)	-0.66% (-2.3-0.79%)	-0.51 (-1.8-0.59)	-5.6 (-23-11)
Maldives	2005	Male	80 to 84	34% (19-51%)	66% (49-81%)	0.56 (0.25-1)	-0.46% (-1.6-0.53%)	-0.68% (-2.1-0.53%)	-0.37 (-1.1-0.28)	-3.2 (-11-3.5)
Maldives	2005	Male	85 to 89	32% (18-48%)	68% (52-82%)	0.49 (0.22-0.93)	-0.35% (-1.1-0.34%)	-0.56% (-1.6-0.33%)	-0.16 (-0.47-0.093)	-0.94 (-3.1-0.97)
Maldives	2005	Male	90 to 94	33% (20-49%)	67% (51-80%)	0.48 (0.21-0.94)	-0.37% (-1.1-0.26%)	-0.59% (-1.5-0.21%)	-0.08 (-0.2-0.028)	-0.33 (-0.95-0.22)
Maldives	2005	Male	95 plus	37% (20-58%)	63% (42-80%)	0.48 (0.18-1)	-0.56% (-1.5-0.19%)	-0.89% (-2.2-0.055%)	-0.033 (-0.079-0.0019)	-0.087 (-0.24-0.03)
Maldives	2010	Female	All Ages	14% (10-19%)	86% (81-90%)	0.24 (0.14-0.37)	0.61% (0.29-1.1%)	0.63% (-0.25-2.1%)	2.6 (-1-8.4)	170 (74-300)
Maldives	2010	Female	15 to 19	12% (6.2-22%)	88% (78-94%)	0.18 (0.06-0.41)	0.56% (0.34-0.9%)	0.69% (0.26-1.5%)	0.031 (0.012-0.066)	10 (6-17)
Maldives	2010	Female	20 to 24	15% (7-27%)	85% (73-93%)	0.25 (0.088-0.53)	1.1% (0.65-1.6%)	0.75% (0.25-1.6%)	0.041 (0.013-0.089)	23 (13-35)
Maldives	2010	Female	25 to 29	16% (7.2-29%)	84% (71-93%)	0.28 (0.091-0.59)	1.1% (0.75-1.5%)	0.74% (0.26-1.6%)	0.036 (0.013-0.076)	21 (13-31)
Maldives	2010	Female	30 to 34	18% (7.9-33%)	82% (67-92%)	0.32 (0.11-0.67)	0.97% (0.62-1.4%)	0.86% (0.36-1.7%)	0.046 (0.019-0.094)	16 (9.4-25)
Maldives	2010	Female	35 to 39	19% (8.3-33%)	81% (67-92%)	0.35 (0.12-0.73)	0.87% (0.54-1.3%)	1.2% (0.49-2.3%)	0.073 (0.03-0.15)	13 (7.6-20)
Maldives	2010	Female	40 to 44	19% (9.5-33%)	81% (67-91%)	0.39 (0.14-0.84)	0.85% (0.48-1.4%)	1.6% (0.74-3.3%)	0.14 (0.061-0.28)	13 (7.1-23)
Maldives	2010	Female	45 to 49	19% (8.8-32%)	81% (68-91%)	0.44 (0.15-1)	0.88% (0.41-1.8%)	1.8% (0.75-4.1%)	0.2 (0.08-0.44)	13 (6.4-29)
Maldives	2010	Female	50 to 54	19% (9.4-33%)	81% (67-91%)	0.49 (0.16-1.1)	0.96% (0.3-2.1%)	1.8% (0.58-4%)	0.3 (0.096-0.7)	15 (4.6-32)
Maldives	2010	Female	55 to 59	20% (9.8-34%)	80% (66-90%)	0.55 (0.17-1.2)	0.94% (0.15-2.3%)	1.7% (0.31-4.1%)	0.28 (0.052-0.66)	11 (1.8-26)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	2010	Female	60 to 64	20% (9.3-34%)	80% (66-91%)	0.55 (0.17-1.2)	0.81% (-0.21-2.5%)	1.3% (-0.24-3.8%)	0.28 (-0.054-0.85)	8.6 (-2.2-27)
Maldives	2010	Female	65 to 69	19% (9.4-32%)	81% (68-91%)	0.51 (0.17-1.1)	0.57% (-0.4-2.1%)	0.81% (-0.56-2.9%)	0.36 (-0.24-1.3)	8.9 (-6.3-33)
Maldives	2010	Female	70 to 74	19% (9-31%)	81% (69-91%)	0.47 (0.14-1.1)	0.49% (-0.45-2.3%)	0.64% (-0.56-2.9%)	0.45 (-0.39-2)	8.6 (-8-40)
Maldives	2010	Female	75 to 79	18% (8.2-31%)	82% (69-92%)	0.44 (0.12-1.1)	0.29% (-0.64-1.9%)	0.34% (-0.83-2.3%)	0.24 (-0.57-1.6)	3.7 (-8.2-24)
Maldives	2010	Female	80 to 84	18% (8.9-33%)	82% (67-91%)	0.44 (0.13-1)	0.17% (-0.79-1.6%)	0.18% (-1-1.9%)	0.082 (-0.47-0.87)	1.1 (-4.9-9.8)
Maldives	2010	Female	85 to 89	20% (9.3-34%)	80% (66-91%)	0.48 (0.14-1.2)	0.12% (-0.69-1.5%)	0.09% (-1-1.9%)	0.021 (-0.22-0.41)	0.29 (-1.6-3.5)
Maldives	2010	Female	90 to 94	23% (10-40%)	77% (60-90%)	0.55 (0.14-1.4)	0.18% (-0.68-1.8%)	0.15% (-0.95-2.2%)	0.017 (-0.11-0.25)	0.15 (-0.55-1.5)
Maldives	2010	Female	95 plus	27% (11-48%)	73% (52-89%)	0.64 (0.14-1.7)	0.22% (-1-2.4%)	0.12% (-1.4-2.7%)	0.0042 (-0.049-0.094)	0.033 (-0.15-0.37)
Maldives	2010	Male	All Ages	37% (30-45%)	63% (55-70%)	0.58 (0.37-0.85)	0.98% (0.27-1.8%)	0.24% (-0.99-1.6%)	1.5 (-6.2-10)	310 (81-550)
Maldives	2010	Male	15 to 19	31% (19-47%)	69% (53-81%)	0.28 (0.12-0.51)	1.1% (0.66-1.7%)	1.4% (0.55-2.8%)	0.13 (0.05-0.25)	22 (13-34)
Maldives	2010	Male	20 to 24	44% (27-61%)	56% (39-73%)	0.63 (0.31-1.1)	2.5% (1.7-3.7%)	3% (1.3-5.5%)	0.37 (0.16-0.68)	57 (36-86)
Maldives	2010	Male	25 to 29	49% (31-66%)	51% (34-69%)	0.85 (0.44-1.4)	2.8% (1.9-4%)	3.5% (1.5-5.9%)	0.34 (0.14-0.58)	53 (33-77)
Maldives	2010	Male	30 to 34	52% (33-68%)	48% (32-67%)	1 (0.56-1.7)	2.7% (1.7-3.7%)	3.4% (1.3-5.8%)	0.28 (0.11-0.49)	39 (24-56)
Maldives	2010	Male	35 to 39	53% (36-70%)	47% (30-64%)	1.1 (0.6-2)	2.6% (1.6-3.9%)	3.8% (1.6-6.3%)	0.3 (0.12-0.5)	34 (20-50)
Maldives	2010	Male	40 to 44	53% (36-72%)	47% (28-64%)	1.2 (0.61-2.1)	2.3% (1.1-3.6%)	3.3% (0.93-6.1%)	0.37 (0.1-0.68)	32 (15-51)
Maldives	2010	Male	45 to 49	52% (33-69%)	48% (31-67%)	1.2 (0.59-2.1)	2% (0.66-3.7%)	2.8% (0.26-6%)	0.49 (0.042-1.1)	32 (10-60)
Maldives	2010	Male	50 to 54	50% (32-68%)	50% (32-68%)	1.2 (0.56-2)	1.4% (-0.16-3.2%)	1.6% (-1-4.5%)	0.42 (-0.28-1.2)	24 (-2.8-58)
Maldives	2010	Male	55 to 59	48% (32-66%)	52% (34-68%)	1.1 (0.5-1.9)	1.2% (-0.25-3.2%)	1.4% (-0.84-4.2%)	0.41 (-0.26-1.2)	18 (-3.7-47)
Maldives	2010	Male	60 to 64	46% (29-64%)	54% (36-71%)	0.95 (0.42-1.7)	0.82% (-0.7-2.6%)	0.84% (-1.2-3.3%)	0.31 (-0.41-1.2)	12 (-9.5-38)
Maldives	2010	Male	65 to 69	42% (27-60%)	58% (40-73%)	0.82 (0.39-1.5)	0.32% (-1-2%)	0.22% (-1.5-2.4%)	0.15 (-1-1.5)	6.5 (-20-40)
Maldives	2010	Male	70 to 74	39% (24-57%)	61% (43-76%)	0.69 (0.31-1.4)	-0.2% (-1.5-1.2%)	-0.42% (-2-1.3%)	-0.44 (-2.1-1.4)	-4.8 (-36-30)
Maldives	2010	Male	75 to 79	35% (21-55%)	65% (45-79%)	0.59 (0.24-1.2)	-0.41% (-1.7-0.65%)	-0.67% (-2.3-0.63%)	-0.72 (-2.5-0.7)	-7.8 (-31-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	2010	Male	80 to 84	33% (19-50%)	67% (50-81%)	0.51 (0.2-1)	-0.48% (-1.6-0.43%)	-0.71% (-2.1-0.4%)	-0.56 (-1.7-0.31)	-4.8 (-16-4.3)
Maldives	2010	Male	85 to 89	31% (18-47%)	69% (53-82%)	0.46 (0.2-0.91)	-0.35% (-1.1-0.29%)	-0.56% (-1.6-0.25%)	-0.22 (-0.61-0.099)	-1.3 (-4.1-1.1)
Maldives	2010	Male	90 to 94	33% (19-49%)	67% (51-81%)	0.44 (0.18-0.89)	-0.29% (-0.91-0.23%)	-0.51% (-1.4-0.16%)	-0.08 (-0.21-0.025)	-0.31 (-0.99-0.25)
Maldives	2010	Male	95 plus	37% (20-57%)	63% (43-80%)	0.45 (0.16-0.98)	-0.42% (-1.3-0.24%)	-0.78% (-1.9-0.091%)	-0.035 (-0.087-0.0042)	-0.086 (-0.26-0.047)
Maldives	2016	Female	All Ages	14% (11-19%)	86% (81-89%)	0.23 (0.098-0.41)	0.62% (0.3-1.2%)	0.54% (-0.26-2.1%)	2.4 (-1.1-9.6)	180 (80-350)
Maldives	2016	Female	15 to 19	12% (6.2-21%)	88% (79-94%)	0.16 (0.045-0.39)	0.56% (0.33-0.89%)	0.66% (0.25-1.4%)	0.02 (0.0072-0.044)	8.1 (4.5-13)
Maldives	2016	Female	20 to 24	15% (6.9-26%)	85% (74-93%)	0.22 (0.062-0.52)	1.1% (0.69-1.6%)	0.72% (0.23-1.8%)	0.031 (0.0093-0.08)	20 (12-31)
Maldives	2016	Female	25 to 29	16% (7.1-28%)	84% (72-93%)	0.25 (0.07-0.58)	1.1% (0.73-1.6%)	0.69% (0.24-1.5%)	0.038 (0.012-0.088)	24 (15-36)
Maldives	2016	Female	30 to 34	17% (7.5-31%)	83% (69-92%)	0.27 (0.078-0.6)	0.97% (0.61-1.4%)	0.82% (0.35-1.7%)	0.059 (0.022-0.13)	22 (13-34)
Maldives	2016	Female	35 to 39	19% (8.3-33%)	81% (67-92%)	0.31 (0.086-0.72)	0.86% (0.52-1.3%)	1.1% (0.49-2.3%)	0.09 (0.033-0.2)	17 (9.6-27)
Maldives	2016	Female	40 to 44	19% (9.4-33%)	81% (67-91%)	0.34 (0.1-0.83)	0.82% (0.45-1.4%)	1.5% (0.7-3.2%)	0.14 (0.057-0.29)	15 (7.7-26)
Maldives	2016	Female	45 to 49	19% (8.7-33%)	81% (67-91%)	0.39 (0.11-0.95)	0.81% (0.37-1.6%)	1.7% (0.67-3.9%)	0.2 (0.077-0.47)	14 (6.4-29)
Maldives	2016	Female	50 to 54	20% (9.2-34%)	80% (66-91%)	0.44 (0.13-1.1)	0.86% (0.27-2%)	1.6% (0.48-3.9%)	0.33 (0.091-0.85)	16 (4.7-39)
Maldives	2016	Female	55 to 59	20% (9.9-34%)	80% (66-90%)	0.48 (0.13-1.2)	0.81% (0.11-2.2%)	1.5% (0.26-4%)	0.37 (0.065-1)	15 (1.8-40)
Maldives	2016	Female	60 to 64	20% (9.5-34%)	80% (66-90%)	0.48 (0.12-1.2)	0.68% (-0.15-2.4%)	1.1% (-0.17-3.7%)	0.34 (-0.051-1.2)	11 (-2.4-39)
Maldives	2016	Female	65 to 69	20% (9.5-33%)	80% (67-90%)	0.45 (0.12-1.1)	0.45% (-0.41-2%)	0.68% (-0.54-2.9%)	0.22 (-0.18-0.99)	5.5 (-5.1-26)
Maldives	2016	Female	70 to 74	19% (9.4-31%)	81% (69-91%)	0.41 (0.11-1)	0.39% (-0.39-1.9%)	0.55% (-0.49-2.7%)	0.3 (-0.27-1.5)	5.9 (-5.9-29)
Maldives	2016	Female	75 to 79	18% (8.8-32%)	82% (68-91%)	0.39 (0.093-0.99)	0.21% (-0.59-1.7%)	0.27% (-0.76-2.3%)	0.21 (-0.64-1.7)	3.4 (-9.5-27)
Maldives	2016	Female	80 to 84	18% (8.5-32%)	82% (68-91%)	0.39 (0.094-0.94)	0.083% (-0.74-1.5%)	0.076% (-0.94-1.9%)	0.047 (-0.64-1.2)	0.73 (-6.6-13)
Maldives	2016	Female	85 to 89	20% (9.5-34%)	80% (66-91%)	0.42 (0.1-1.1)	0.038% (-0.7-1.3%)	-0.012% (-0.97-1.7%)	-0.0051 (-0.36-0.59)	0.14 (-2.7-5.1)
Maldives	2016	Female	90 to 94	22% (11-39%)	78% (61-89%)	0.47 (0.11-1.3)	0.059% (-0.7-1.6%)	-0.004% (-0.97-2%)	-0.0012 (-0.2-0.38)	0.079 (-0.99-2.2)
Maldives	2016	Female	95 plus	27% (12-47%)	73% (53-88%)	0.56 (0.11-1.6)	0.09% (-0.94-2.3%)	-0.019% (-1.3-2.5%)	-0.0015 (-0.095-0.18)	0.028 (-0.3-0.69)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Maldives	2016	Male	All Ages	38% (32-46%)	62% (54-68%)	0.55 (0.25-0.93)	1.1% (0.4-2%)	0.28% (-0.84-1.6%)	1.9 (-5.7-11)	350 (120-670)
Maldives	2016	Male	15 to 19	31% (18-46%)	69% (54-82%)	0.24 (0.083-0.51)	1% (0.63-1.6%)	1.3% (0.48-2.6%)	0.075 (0.026-0.16)	16 (9.3-25)
Maldives	2016	Male	20 to 24	44% (28-60%)	56% (40-72%)	0.55 (0.21-1.1)	2.5% (1.6-3.7%)	2.7% (0.97-5.4%)	0.28 (0.091-0.57)	50 (30-78)
Maldives	2016	Male	25 to 29	49% (31-66%)	51% (34-69%)	0.74 (0.28-1.4)	2.8% (1.8-4.1%)	3.1% (1.1-5.8%)	0.35 (0.11-0.7)	61 (38-92)
Maldives	2016	Male	30 to 34	51% (33-68%)	49% (32-67%)	0.88 (0.36-1.7)	2.6% (1.6-3.9%)	2.9% (1-5.3%)	0.36 (0.12-0.72)	56 (33-86)
Maldives	2016	Male	35 to 39	53% (35-70%)	47% (30-65%)	0.98 (0.41-1.9)	2.5% (1.5-4%)	3.3% (1.2-6.5%)	0.37 (0.12-0.79)	43 (25-71)
Maldives	2016	Male	40 to 44	53% (37-71%)	47% (29-63%)	1 (0.43-2)	2.2% (0.98-3.7%)	2.9% (0.55-5.9%)	0.4 (0.074-0.93)	36 (16-66)
Maldives	2016	Male	45 to 49	52% (34-71%)	48% (29-66%)	1 (0.43-2.1)	1.9% (0.62-3.7%)	2.4% (0.019-5.8%)	0.49 (0.0033-1.2)	34 (11-68)
Maldives	2016	Male	50 to 54	50% (32-69%)	50% (31-68%)	1 (0.4-2)	1.2% (-0.13-3.2%)	1.4% (-0.91-4.3%)	0.44 (-0.29-1.4)	26 (-2.9-68)
Maldives	2016	Male	55 to 59	48% (31-67%)	52% (33-69%)	0.92 (0.34-1.9)	1.1% (-0.21-3.1%)	1.2% (-0.81-4.1%)	0.53 (-0.35-1.9)	25 (-5.1-73)
Maldives	2016	Male	60 to 64	45% (29-64%)	55% (36-71%)	0.82 (0.28-1.7)	0.76% (-0.55-2.7%)	0.78% (-1-3.4%)	0.39 (-0.52-1.7)	15 (-10-53)
Maldives	2016	Male	65 to 69	42% (26-59%)	58% (41-74%)	0.7 (0.26-1.4)	0.33% (-0.8-1.9%)	0.23% (-1.3-2.3%)	0.12 (-0.65-1.2)	5.3 (-12-31)
Maldives	2016	Male	70 to 74	38% (24-55%)	62% (45-76%)	0.59 (0.2-1.3)	-0.082% (-1.2-1.1%)	-0.28% (-1.8-1.3%)	-0.22 (-1.3-0.98)	-1.5 (-22-21)
Maldives	2016	Male	75 to 79	35% (21-53%)	65% (47-79%)	0.51 (0.16-1.1)	-0.28% (-1.4-0.62%)	-0.52% (-2-0.63%)	-0.55 (-2.1-0.67)	-5.5 (-26-12)
Maldives	2016	Male	80 to 84	32% (18-50%)	68% (50-82%)	0.44 (0.15-0.92)	-0.37% (-1.3-0.42%)	-0.57% (-1.8-0.4%)	-0.59 (-1.9-0.46)	-4.9 (-18-5.8)
Maldives	2016	Male	85 to 89	31% (17-47%)	69% (53-83%)	0.39 (0.13-0.88)	-0.28% (-0.99-0.27%)	-0.48% (-1.4-0.27%)	-0.32 (-0.95-0.16)	-1.8 (-6.4-1.7)
Maldives	2016	Male	90 to 94	32% (19-49%)	68% (51-81%)	0.38 (0.13-0.81)	-0.28% (-0.91-0.21%)	-0.48% (-1.4-0.16%)	-0.13 (-0.39-0.045)	-0.52 (-1.8-0.4)
Maldives	2016	Male	95 plus	36% (19-57%)	64% (43-81%)	0.38 (0.11-0.94)	-0.41% (-1.3-0.19%)	-0.72% (-1.9-0.049%)	-0.061 (-0.16-0.0041)	-0.15 (-0.47-0.074)
Myanmar	1990	Female	All Ages	2.1% (1.8-2.7%)	98% (97-98%)	0.025 (0.016-0.035)	0.41% (0.32-0.52%)	0.58% (0.43-0.77%)	1200 (880-1600)	45000 (36000-56000)
Myanmar	1990	Female	15 to 19	2% (1.2-3.1%)	98% (97-99%)	0.024 (0.011-0.045)	0.38% (0.24-0.55%)	0.3% (0.15-0.5%)	10 (5.1-18)	1700 (1000-2500)
Myanmar	1990	Female	20 to 24	2.5% (1.3-3.8%)	98% (96-99%)	0.032 (0.014-0.062)	0.64% (0.42-0.9%)	0.38% (0.18-0.67%)	17 (8.1-30)	3100 (2000-4500)
Myanmar	1990	Female	25 to 29	2.7% (1.5-4.3%)	97% (96-99%)	0.035 (0.015-0.067)	0.7% (0.47-0.97%)	0.46% (0.22-0.77%)	21 (9.8-36)	3100 (2000-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	1990	Female	30 to 34	2.9% (1.6-4.8%)	97% (95-98%)	0.038 (0.018-0.072)	0.81% (0.57-1.1%)	0.74% (0.44-1.1%)	37 (22-56)	3600 (2500-5000)
Myanmar	1990	Female	35 to 39	3% (1.7-4.7%)	97% (95-98%)	0.042 (0.019-0.076)	1% (0.72-1.4%)	1.1% (0.71-1.6%)	65 (41-97)	4500 (3200-6200)
Myanmar	1990	Female	40 to 44	3% (1.8-4.7%)	97% (95-98%)	0.044 (0.021-0.08)	1.2% (0.84-1.6%)	1.4% (0.96-2%)	74 (49-110)	4100 (2900-5600)
Myanmar	1990	Female	45 to 49	3.1% (1.7-4.6%)	97% (95-98%)	0.047 (0.021-0.088)	1.2% (0.81-1.6%)	1.4% (0.91-2%)	88 (57-130)	4100 (2800-5700)
Myanmar	1990	Female	50 to 54	3.1% (1.8-4.9%)	97% (95-98%)	0.052 (0.024-0.096)	1.1% (0.78-1.6%)	1.3% (0.86-2%)	110 (73-170)	4500 (3000-6600)
Myanmar	1990	Female	55 to 59	3.2% (1.8-4.8%)	97% (95-98%)	0.057 (0.026-0.11)	0.96% (0.66-1.4%)	1.1% (0.76-1.7%)	130 (84-190)	4400 (2900-6300)
Myanmar	1990	Female	60 to 64	3% (1.8-4.9%)	97% (95-98%)	0.057 (0.026-0.11)	0.82% (0.51-1.2%)	0.97% (0.6-1.5%)	150 (92-220)	4200 (2600-6100)
Myanmar	1990	Female	65 to 69	2.8% (1.6-4.3%)	97% (96-98%)	0.054 (0.025-0.097)	0.69% (0.44-1%)	0.81% (0.51-1.2%)	140 (87-210)	3200 (2000-4800)
Myanmar	1990	Female	70 to 74	2.7% (1.6-4.2%)	97% (96-98%)	0.053 (0.023-0.1)	0.6% (0.36-0.96%)	0.7% (0.41-1.1%)	140 (80-220)	2500 (1500-4100)
Myanmar	1990	Female	75 to 79	2.8% (1.6-4.5%)	97% (96-98%)	0.054 (0.024-0.11)	0.47% (0.26-0.78%)	0.55% (0.29-0.91%)	97 (52-160)	1400 (740-2300)
Myanmar	1990	Female	80 to 84	3% (1.7-4.5%)	97% (96-98%)	0.056 (0.025-0.1)	0.42% (0.22-0.7%)	0.49% (0.25-0.83%)	63 (32-110)	650 (330-1100)
Myanmar	1990	Female	85 to 89	3.4% (1.8-5.5%)	97% (95-98%)	0.062 (0.027-0.12)	0.35% (0.16-0.6%)	0.41% (0.19-0.71%)	28 (13-50)	210 (98-360)
Myanmar	1990	Female	90 to 94	4% (2.2-6.5%)	96% (93-98%)	0.07 (0.029-0.14)	0.29% (0.12-0.55%)	0.34% (0.13-0.66%)	7.9 (3-15)	40 (16-76)
Myanmar	1990	Female	95 plus	5% (2.3-9.4%)	95% (91-98%)	0.084 (0.027-0.18)	0.34% (0.11-0.7%)	0.37% (0.11-0.81%)	2 (0.6-4.4)	7 (2.3-15)
Myanmar	1990	Male	All Ages	19% (16-22%)	81% (78-84%)	0.48 (0.34-0.63)	3.3% (2-4.7%)	4.8% (2.8-7.1%)	11000 (6400-16000)	420000 (270000-590000)
Myanmar	1990	Male	15 to 19	16% (10-22%)	84% (78-90%)	0.21 (0.11-0.38)	1.2% (0.56-2.2%)	1.4% (0.4-2.7%)	71 (20-140)	6600 (2900-11000)
Myanmar	1990	Male	20 to 24	25% (17-34%)	75% (66-83%)	0.55 (0.3-0.9)	4% (1.8-7%)	4.6% (1.8-8.4%)	320 (130-610)	25000 (11000-44000)
Myanmar	1990	Male	25 to 29	30% (21-41%)	70% (59-79%)	0.79 (0.46-1.2)	6.7% (3.3-11%)	8.1% (3.6-14%)	560 (250-990)	38000 (19000-64000)
Myanmar	1990	Male	30 to 34	32% (22-44%)	68% (56-78%)	0.98 (0.55-1.5)	8.6% (4.7-13%)	11% (5.5-17%)	780 (400-1200)	47000 (25000-73000)
Myanmar	1990	Male	35 to 39	34% (22-44%)	66% (56-78%)	1.1 (0.62-1.8)	10% (5.6-15%)	12% (6.9-19%)	1000 (550-1500)	54000 (31000-81000)
Myanmar	1990	Male	40 to 44	33% (23-43%)	67% (57-77%)	1.2 (0.68-1.9)	11% (6.5-16%)	13% (7.8-19%)	1000 (590-1500)	47000 (28000-70000)
Myanmar	1990	Male	45 to 49	32% (21-42%)	68% (58-79%)	1.3 (0.68-2.1)	11% (6.5-17%)	14% (7.7-20%)	1200 (670-1800)	49000 (28000-75000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	1990	Male	50 to 54	29% (21-39%)	71% (61-79%)	1.2 (0.63-1.9)	9.7% (5.3-15%)	12% (6.4-18%)	1200 (660-1900)	45000 (24000-70000)
Myanmar	1990	Male	55 to 59	26% (18-35%)	74% (65-82%)	0.96 (0.51-1.6)	7.3% (4-12%)	8.7% (4.7-14%)	1100 (600-1900)	37000 (19000-59000)
Myanmar	1990	Male	60 to 64	24% (16-33%)	76% (67-84%)	0.81 (0.42-1.4)	5.8% (2.7-10%)	6.8% (3.2-12%)	1100 (510-1900)	30000 (14000-52000)
Myanmar	1990	Male	65 to 69	21% (14-29%)	79% (71-86%)	0.7 (0.36-1.2)	4.5% (2.1-8.1%)	5.3% (2.4-9.2%)	880 (400-1500)	20000 (9200-34000)
Myanmar	1990	Male	70 to 74	19% (13-26%)	81% (74-87%)	0.59 (0.28-1)	3.7% (1.5-7%)	4.3% (1.8-8%)	730 (300-1400)	13000 (5500-25000)
Myanmar	1990	Male	75 to 79	17% (11-24%)	83% (76-89%)	0.5 (0.24-0.93)	2.8% (1.1-5.3%)	3.2% (1.2-6.1%)	410 (150-810)	5800 (2100-11000)
Myanmar	1990	Male	80 to 84	15% (9.8-21%)	85% (79-90%)	0.42 (0.2-0.72)	2.3% (0.88-4.3%)	2.7% (0.97-5%)	210 (77-390)	2200 (810-4100)
Myanmar	1990	Male	85 to 89	15% (9.2-23%)	85% (77-91%)	0.37 (0.17-0.69)	1.7% (0.6-3.4%)	2% (0.67-3.9%)	80 (27-160)	590 (200-1100)
Myanmar	1990	Male	90 to 94	17% (11-24%)	83% (76-89%)	0.36 (0.16-0.67)	1.3% (0.31-2.8%)	1.5% (0.34-3.3%)	21 (4.6-44)	110 (24-220)
Myanmar	1990	Male	95 plus	19% (11-30%)	81% (70-89%)	0.37 (0.14-0.7)	1.2% (0.18-2.7%)	1.3% (0.13-3.1%)	3.8 (0.38-9.1)	13 (2.1-31)
Myanmar	1995	Female	All Ages	2.2% (1.8-2.7%)	98% (97-98%)	0.028 (0.018-0.04)	0.47% (0.37-0.59%)	0.64% (0.47-0.84%)	1400 (1000-1800)	52000 (41000-64000)
Myanmar	1995	Female	15 to 19	2% (1.2-3.1%)	98% (97-99%)	0.026 (0.012-0.049)	0.4% (0.25-0.58%)	0.32% (0.16-0.51%)	11 (5.8-19)	1800 (1100-2700)
Myanmar	1995	Female	20 to 24	2.5% (1.4-3.8%)	98% (96-99%)	0.034 (0.015-0.066)	0.68% (0.46-0.95%)	0.4% (0.21-0.7%)	18 (9.4-32)	3300 (2100-4800)
Myanmar	1995	Female	25 to 29	2.7% (1.5-4.2%)	97% (96-99%)	0.038 (0.017-0.072)	0.73% (0.5-1%)	0.49% (0.26-0.81%)	23 (12-38)	3500 (2300-4900)
Myanmar	1995	Female	30 to 34	2.9% (1.5-4.7%)	97% (95-98%)	0.041 (0.018-0.077)	0.85% (0.61-1.1%)	0.78% (0.5-1.1%)	40 (26-60)	4000 (2700-5400)
Myanmar	1995	Female	35 to 39	3% (1.6-4.6%)	97% (95-98%)	0.045 (0.021-0.081)	1% (0.76-1.4%)	1.2% (0.77-1.7%)	71 (47-100)	5000 (3500-6900)
Myanmar	1995	Female	40 to 44	3% (1.8-4.7%)	97% (95-98%)	0.048 (0.023-0.087)	1.2% (0.86-1.6%)	1.5% (1-2.1%)	100 (70-150)	5800 (4200-8100)
Myanmar	1995	Female	45 to 49	3.1% (1.7-4.7%)	97% (95-98%)	0.051 (0.023-0.095)	1.2% (0.85-1.6%)	1.5% (1-2.1%)	100 (68-150)	4700 (3300-6600)
Myanmar	1995	Female	50 to 54	3.1% (1.8-4.7%)	97% (95-98%)	0.056 (0.026-0.1)	1.2% (0.8-1.7%)	1.4% (0.94-2%)	120 (79-180)	4800 (3200-6700)
Myanmar	1995	Female	55 to 59	3.1% (1.8-4.8%)	97% (95-98%)	0.062 (0.029-0.11)	1% (0.67-1.4%)	1.2% (0.78-1.7%)	140 (87-190)	4600 (3000-6300)
Myanmar	1995	Female	60 to 64	3% (1.8-4.9%)	97% (95-98%)	0.061 (0.028-0.12)	0.87% (0.54-1.3%)	1% (0.62-1.6%)	160 (95-240)	4400 (2700-6600)
Myanmar	1995	Female	65 to 69	2.8% (1.6-4.2%)	97% (96-98%)	0.058 (0.028-0.1)	0.73% (0.48-1.1%)	0.87% (0.55-1.3%)	170 (110-250)	3900 (2500-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	1995	Female	70 to 74	2.7% (1.7-4.2%)	97% (96-98%)	0.057 (0.025-0.11)	0.65% (0.38-1%)	0.76% (0.43-1.2%)	160 (91-250)	2900 (1700-4600)
Myanmar	1995	Female	75 to 79	2.8% (1.7-4.5%)	97% (96-98%)	0.059 (0.026-0.12)	0.51% (0.28-0.83%)	0.6% (0.32-0.97%)	120 (64-200)	1700 (920-2700)
Myanmar	1995	Female	80 to 84	3% (1.6-4.5%)	97% (95-98%)	0.061 (0.027-0.11)	0.46% (0.25-0.77%)	0.53% (0.28-0.9%)	79 (42-130)	810 (440-1400)
Myanmar	1995	Female	85 to 89	3.4% (1.8-5.6%)	97% (94-98%)	0.067 (0.029-0.13)	0.38% (0.19-0.65%)	0.45% (0.22-0.76%)	35 (17-59)	250 (120-430)
Myanmar	1995	Female	90 to 94	4% (2.2-6.6%)	96% (93-98%)	0.077 (0.032-0.14)	0.32% (0.14-0.6%)	0.38% (0.16-0.71%)	10 (4.5-19)	53 (23-99)
Myanmar	1995	Female	95 plus	4.9% (2.3-9.2%)	95% (91-98%)	0.092 (0.03-0.2)	0.37% (0.13-0.72%)	0.41% (0.13-0.83%)	2.8 (0.92-5.7)	9.9 (3.5-19)
Myanmar	1995	Male	All Ages	19% (16-23%)	81% (77-84%)	0.55 (0.39-0.73)	3.8% (2.4-5.3%)	5.3% (3.3-7.6%)	12000 (7600-18000)	490000 (310000-680000)
Myanmar	1995	Male	15 to 19	16% (10-23%)	84% (77-90%)	0.23 (0.12-0.42)	1.3% (0.65-2.3%)	1.5% (0.53-2.8%)	82 (29-160)	7500 (3600-13000)
Myanmar	1995	Male	20 to 24	26% (17-34%)	74% (66-83%)	0.6 (0.32-1)	4.2% (2-6.9%)	4.8% (2-8.2%)	370 (150-640)	28000 (13000-47000)
Myanmar	1995	Male	25 to 29	30% (21-41%)	70% (59-79%)	0.87 (0.5-1.4)	6.9% (3.6-11%)	8.2% (4-14%)	660 (310-1100)	44000 (23000-72000)
Myanmar	1995	Male	30 to 34	33% (22-45%)	67% (55-78%)	1.1 (0.61-1.7)	8.8% (5-13%)	11% (5.9-17%)	900 (490-1400)	54000 (31000-82000)
Myanmar	1995	Male	35 to 39	34% (23-45%)	66% (55-77%)	1.2 (0.68-1.9)	10% (6.1-15%)	13% (7.4-19%)	1200 (680-1800)	63000 (37000-94000)
Myanmar	1995	Male	40 to 44	33% (23-44%)	67% (56-77%)	1.3 (0.75-2.1)	11% (6.7-16%)	13% (8.2-19%)	1400 (830-2000)	66000 (40000-95000)
Myanmar	1995	Male	45 to 49	32% (21-42%)	68% (58-79%)	1.4 (0.73-2.3)	11% (6.9-17%)	14% (8.4-20%)	1300 (790-2000)	56000 (33000-84000)
Myanmar	1995	Male	50 to 54	29% (21-40%)	71% (60-79%)	1.3 (0.67-2.1)	10% (5.6-15%)	12% (6.7-18%)	1300 (710-2100)	49000 (26000-75000)
Myanmar	1995	Male	55 to 59	27% (18-36%)	73% (64-82%)	1.1 (0.56-1.7)	7.7% (4.3-12%)	9.2% (5.1-14%)	1200 (650-1900)	39000 (21000-61000)
Myanmar	1995	Male	60 to 64	24% (16-34%)	76% (66-84%)	0.89 (0.46-1.6)	6.1% (3.1-10%)	7.1% (3.6-12%)	1200 (570-2000)	32000 (16000-55000)
Myanmar	1995	Male	65 to 69	21% (14-29%)	79% (71-86%)	0.77 (0.4-1.3)	4.8% (2.3-8.1%)	5.6% (2.7-9.4%)	1100 (510-1800)	24000 (11000-40000)
Myanmar	1995	Male	70 to 74	19% (13-26%)	81% (74-87%)	0.64 (0.31-1.1)	4% (1.7-7.3%)	4.6% (2-8.3%)	840 (350-1500)	15000 (6500-28000)
Myanmar	1995	Male	75 to 79	17% (11-24%)	83% (76-89%)	0.55 (0.26-0.99)	3% (1.2-5.5%)	3.4% (1.3-6.4%)	520 (200-980)	7300 (2900-14000)
Myanmar	1995	Male	80 to 84	15% (9.8-21%)	85% (79-90%)	0.46 (0.22-0.78)	2.5% (0.96-4.5%)	2.8% (1.1-5.2%)	260 (99-490)	2800 (1100-5100)
Myanmar	1995	Male	85 to 89	15% (9.1-22%)	85% (78-91%)	0.41 (0.18-0.72)	1.9% (0.68-3.6%)	2.1% (0.75-4.1%)	99 (36-190)	730 (270-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	1995	Male	90 to 94	17% (11-24%)	83% (76-89%)	0.4 (0.17-0.7)	1.4% (0.43-2.9%)	1.7% (0.48-3.4%)	27 (7.7-56)	140 (42-280)
Myanmar	1995	Male	95 plus	20% (11-30%)	80% (70-89%)	0.4 (0.15-0.76)	1.3% (0.29-2.9%)	1.4% (0.27-3.3%)	5.4 (1-12)	19 (4.3-43)
Myanmar	2000	Female	All Ages	2% (1.6-2.4%)	98% (98-98%)	0.029 (0.019-0.04)	0.52% (0.41-0.64%)	0.7% (0.53-0.92%)	1500 (1100-1900)	56000 (45000-68000)
Myanmar	2000	Female	15 to 19	1.9% (1.1-2.9%)	98% (97-99%)	0.027 (0.012-0.052)	0.4% (0.26-0.58%)	0.31% (0.17-0.5%)	11 (5.8-18)	1800 (1200-2700)
Myanmar	2000	Female	20 to 24	2.2% (1.2-3.3%)	98% (97-99%)	0.034 (0.015-0.066)	0.68% (0.46-0.97%)	0.39% (0.21-0.66%)	17 (9.3-29)	3400 (2200-5000)
Myanmar	2000	Female	25 to 29	2.2% (1.3-3.4%)	98% (97-99%)	0.034 (0.016-0.065)	0.74% (0.53-0.99%)	0.46% (0.28-0.73%)	21 (13-34)	3500 (2400-4900)
Myanmar	2000	Female	30 to 34	2.2% (1.2-3.5%)	98% (96-99%)	0.036 (0.017-0.066)	0.84% (0.63-1.1%)	0.74% (0.5-1.1%)	39 (26-57)	4100 (2900-5400)
Myanmar	2000	Female	35 to 39	2.4% (1.4-3.8%)	98% (96-99%)	0.041 (0.019-0.073)	1% (0.76-1.4%)	1.1% (0.77-1.7%)	70 (46-100)	5000 (3700-6900)
Myanmar	2000	Female	40 to 44	2.6% (1.5-3.8%)	97% (96-99%)	0.045 (0.023-0.08)	1.2% (0.87-1.6%)	1.5% (1-2%)	110 (73-150)	6000 (4300-8100)
Myanmar	2000	Female	45 to 49	2.6% (1.5-3.9%)	97% (96-99%)	0.049 (0.022-0.091)	1.2% (0.89-1.6%)	1.5% (1.1-2.1%)	130 (94-190)	6400 (4500-8600)
Myanmar	2000	Female	50 to 54	2.9% (1.8-4.2%)	97% (96-98%)	0.059 (0.027-0.1)	1.2% (0.84-1.7%)	1.5% (0.98-2.1%)	130 (88-190)	5200 (3600-7400)
Myanmar	2000	Female	55 to 59	3.1% (1.9-4.5%)	97% (95-98%)	0.068 (0.032-0.12)	1.1% (0.71-1.5%)	1.3% (0.81-1.8%)	140 (89-200)	4600 (3100-6700)
Myanmar	2000	Female	60 to 64	2.8% (1.7-4.5%)	97% (96-98%)	0.065 (0.03-0.12)	0.93% (0.6-1.3%)	1.1% (0.71-1.7%)	160 (100-240)	4500 (2900-6500)
Myanmar	2000	Female	65 to 69	2.6% (1.5-3.9%)	97% (96-98%)	0.061 (0.03-0.11)	0.78% (0.52-1.1%)	0.93% (0.61-1.4%)	180 (120-260)	4100 (2700-5900)
Myanmar	2000	Female	70 to 74	2.6% (1.6-4%)	97% (96-98%)	0.062 (0.028-0.12)	0.69% (0.44-1.1%)	0.82% (0.52-1.3%)	190 (120-290)	3400 (2200-5300)
Myanmar	2000	Female	75 to 79	2.8% (1.6-4.4%)	97% (96-98%)	0.065 (0.028-0.13)	0.56% (0.33-0.89%)	0.66% (0.38-1.1%)	140 (78-220)	1900 (1100-3100)
Myanmar	2000	Female	80 to 84	3% (1.6-4.4%)	97% (96-98%)	0.068 (0.03-0.13)	0.5% (0.27-0.8%)	0.59% (0.31-0.96%)	95 (51-150)	980 (540-1600)
Myanmar	2000	Female	85 to 89	3.3% (1.8-5.4%)	97% (95-98%)	0.074 (0.033-0.14)	0.42% (0.23-0.69%)	0.49% (0.26-0.82%)	45 (24-75)	330 (180-540)
Myanmar	2000	Female	90 to 94	3.9% (2.2-6.5%)	96% (93-98%)	0.084 (0.036-0.16)	0.35% (0.17-0.62%)	0.41% (0.19-0.73%)	13 (6.1-23)	67 (32-120)
Myanmar	2000	Female	95 plus	4.9% (2.4-9.2%)	95% (91-98%)	0.1 (0.034-0.22)	0.4% (0.16-0.76%)	0.45% (0.17-0.88%)	3.7 (1.4-7.2)	13 (5.3-25)
Myanmar	2000	Male	All Ages	21% (19-23%)	79% (77-81%)	0.69 (0.49-0.91)	4.3% (2.9-5.9%)	5.8% (3.8-7.9%)	14000 (8800-19000)	560000 (380000-760000)
Myanmar	2000	Male	15 to 19	15% (11-20%)	85% (80-89%)	0.25 (0.13-0.41)	1.4% (0.75-2.2%)	1.6% (0.7-2.8%)	87 (38-150)	8000 (4300-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	2000	Male	20 to 24	27% (21-33%)	73% (67-79%)	0.7 (0.41-1.1)	4.6% (2.6-7.2%)	5.2% (2.7-8.6%)	420 (200-690)	32000 (17000-50000)
Myanmar	2000	Male	25 to 29	34% (28-40%)	66% (60-72%)	1.1 (0.7-1.6)	7.6% (4.2-11%)	9% (4.7-14%)	780 (410-1200)	52000 (29000-79000)
Myanmar	2000	Male	30 to 34	37% (30-43%)	63% (57-70%)	1.3 (0.84-1.9)	9.3% (5.7-13%)	11% (6.6-17%)	1100 (640-1500)	64000 (39000-91000)
Myanmar	2000	Male	35 to 39	37% (30-43%)	63% (57-70%)	1.5 (0.89-2.2)	11% (6.6-15%)	13% (7.9-18%)	1300 (810-1900)	72000 (44000-100000)
Myanmar	2000	Male	40 to 44	37% (31-43%)	63% (57-69%)	1.6 (1-2.4)	11% (7.5-16%)	14% (9.1-19%)	1600 (1000-2300)	77000 (50000-110000)
Myanmar	2000	Male	45 to 49	35% (29-41%)	65% (59-71%)	1.7 (1-2.6)	12% (7.9-17%)	15% (9.5-21%)	1900 (1200-2700)	80000 (51000-110000)
Myanmar	2000	Male	50 to 54	32% (26-38%)	68% (62-74%)	1.6 (0.9-2.4)	11% (6.5-16%)	13% (7.8-19%)	1600 (900-2300)	57000 (34000-85000)
Myanmar	2000	Male	55 to 59	26% (21-33%)	74% (67-79%)	1.2 (0.68-1.8)	7.9% (4.6-12%)	9.4% (5.5-14%)	1200 (700-1900)	40000 (23000-62000)
Myanmar	2000	Male	60 to 64	20% (15-27%)	80% (73-85%)	0.85 (0.47-1.5)	5.8% (3.1-9.3%)	6.7% (3.6-11%)	1100 (570-1800)	30000 (16000-49000)
Myanmar	2000	Male	65 to 69	17% (12-22%)	83% (78-88%)	0.68 (0.38-1.1)	4.3% (2.4-7.1%)	5% (2.7-8.3%)	940 (500-1500)	21000 (11000-35000)
Myanmar	2000	Male	70 to 74	15% (11-21%)	85% (79-89%)	0.57 (0.28-0.98)	3.6% (1.8-6.3%)	4.1% (2-7.3%)	850 (410-1500)	15000 (7500-27000)
Myanmar	2000	Male	75 to 79	15% (9.6-21%)	85% (79-90%)	0.53 (0.25-0.93)	2.8% (1.3-5.1%)	3.2% (1.4-5.8%)	540 (230-1000)	7600 (3300-14000)
Myanmar	2000	Male	80 to 84	15% (9.5-20%)	85% (80-90%)	0.49 (0.24-0.83)	2.5% (1.1-4.4%)	2.9% (1.2-5.2%)	310 (130-560)	3200 (1400-5700)
Myanmar	2000	Male	85 to 89	15% (9.5-22%)	85% (78-91%)	0.46 (0.22-0.79)	2% (0.82-3.6%)	2.3% (0.93-4.2%)	130 (52-230)	950 (390-1700)
Myanmar	2000	Male	90 to 94	17% (11-25%)	83% (75-89%)	0.45 (0.21-0.81)	1.6% (0.54-3.1%)	1.8% (0.62-3.7%)	35 (11-69)	180 (61-360)
Myanmar	2000	Male	95 plus	20% (11-30%)	80% (70-89%)	0.45 (0.17-0.85)	1.4% (0.38-3%)	1.6% (0.37-3.4%)	7.2 (1.7-16)	25 (6.7-54)
Myanmar	2005	Female	All Ages	2% (1.6-2.4%)	98% (98-98%)	0.032 (0.021-0.045)	0.57% (0.45-0.69%)	0.76% (0.58-0.96%)	1500 (1200-1900)	55000 (45000-67000)
Myanmar	2005	Female	15 to 19	1.8% (1.1-2.9%)	98% (97-99%)	0.029 (0.013-0.055)	0.41% (0.26-0.58%)	0.3% (0.17-0.48%)	8 (4.4-13)	1600 (990-2300)
Myanmar	2005	Female	20 to 24	2.1% (1.1-3.2%)	98% (97-99%)	0.036 (0.016-0.071)	0.72% (0.49-1%)	0.38% (0.22-0.61%)	14 (8.2-23)	3300 (2200-4800)
Myanmar	2005	Female	25 to 29	2.1% (1.2-3.3%)	98% (97-99%)	0.037 (0.016-0.069)	0.76% (0.54-1%)	0.45% (0.29-0.66%)	18 (12-28)	3400 (2300-4800)
Myanmar	2005	Female	30 to 34	2.2% (1.2-3.5%)	98% (96-99%)	0.038 (0.018-0.073)	0.84% (0.63-1.1%)	0.72% (0.49-1%)	34 (23-49)	3800 (2700-5200)
Myanmar	2005	Female	35 to 39	2.4% (1.4-3.6%)	98% (96-99%)	0.044 (0.021-0.078)	1% (0.74-1.3%)	1.1% (0.75-1.6%)	64 (42-92)	4800 (3500-6400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	2005	Female	40 to 44	2.5% (1.5-3.7%)	97% (96-98%)	0.049 (0.025-0.085)	1.1% (0.84-1.5%)	1.4% (1-2%)	96 (67-140)	5600 (4100-7500)
Myanmar	2005	Female	45 to 49	2.6% (1.5-3.8%)	97% (96-99%)	0.053 (0.024-0.097)	1.2% (0.86-1.6%)	1.5% (1-2.1%)	130 (87-180)	6100 (4400-8200)
Myanmar	2005	Female	50 to 54	2.8% (1.8-4.1%)	97% (96-98%)	0.063 (0.03-0.11)	1.2% (0.83-1.6%)	1.5% (1-2.1%)	160 (110-240)	6600 (4600-9200)
Myanmar	2005	Female	55 to 59	3% (1.8-4.5%)	97% (95-98%)	0.073 (0.035-0.13)	1% (0.71-1.5%)	1.3% (0.85-1.9%)	140 (93-210)	4800 (3200-6900)
Myanmar	2005	Female	60 to 64	2.8% (1.7-4.3%)	97% (96-98%)	0.07 (0.032-0.13)	0.94% (0.62-1.4%)	1.2% (0.76-1.7%)	150 (99-220)	4300 (2800-6100)
Myanmar	2005	Female	65 to 69	2.6% (1.5-3.8%)	97% (96-99%)	0.065 (0.032-0.12)	0.8% (0.54-1.1%)	0.98% (0.64-1.4%)	170 (110-240)	3900 (2600-5500)
Myanmar	2005	Female	70 to 74	2.6% (1.6-4.1%)	97% (96-98%)	0.067 (0.03-0.13)	0.72% (0.46-1.1%)	0.87% (0.55-1.3%)	190 (120-280)	3400 (2200-5200)
Myanmar	2005	Female	75 to 79	2.7% (1.6-4.4%)	97% (96-98%)	0.071 (0.032-0.14)	0.57% (0.35-0.9%)	0.69% (0.41-1.1%)	160 (93-240)	2200 (1300-3400)
Myanmar	2005	Female	80 to 84	2.9% (1.6-4.4%)	97% (96-98%)	0.074 (0.032-0.14)	0.52% (0.3-0.81%)	0.62% (0.36-0.97%)	110 (62-170)	1100 (650-1700)
Myanmar	2005	Female	85 to 89	3.3% (1.8-5.3%)	97% (95-98%)	0.081 (0.036-0.16)	0.44% (0.24-0.72%)	0.52% (0.28-0.85%)	52 (28-87)	380 (210-630)
Myanmar	2005	Female	90 to 94	3.9% (2.2-6.4%)	96% (94-98%)	0.092 (0.039-0.17)	0.37% (0.19-0.64%)	0.43% (0.22-0.75%)	16 (8.1-28)	84 (43-150)
Myanmar	2005	Female	95 plus	4.8% (2.4-9%)	95% (91-98%)	0.11 (0.037-0.24)	0.42% (0.18-0.78%)	0.47% (0.18-0.9%)	4.4 (1.7-8.5)	16 (6.7-29)
Myanmar	2005	Male	All Ages	21% (19-24%)	79% (76-81%)	0.78 (0.56-1)	4.5% (3.2-6%)	5.9% (4-7.8%)	14000 (9100-20000)	55000 (38000-77000)
Myanmar	2005	Male	15 to 19	15% (11-20%)	85% (80-89%)	0.26 (0.14-0.44)	1.4% (0.84-2.2%)	1.7% (0.8-2.8%)	74 (35-120)	7000 (4100-11000)
Myanmar	2005	Male	20 to 24	26% (21-32%)	74% (68-79%)	0.75 (0.44-1.2)	4.4% (2.7-6.6%)	5.1% (2.8-8%)	380 (200-610)	30000 (17000-46000)
Myanmar	2005	Male	25 to 29	34% (28-39%)	66% (61-72%)	1.2 (0.74-1.8)	7% (4.3-10%)	8.3% (4.7-13%)	720 (400-1100)	49000 (29000-74000)
Myanmar	2005	Male	30 to 34	36% (30-42%)	64% (58-70%)	1.4 (0.91-2.1)	8.4% (5.4-12%)	10% (6.1-15%)	980 (590-1400)	60000 (37000-85000)
Myanmar	2005	Male	35 to 39	36% (30-42%)	64% (58-70%)	1.6 (0.98-2.4)	9.7% (6.3-14%)	12% (7.6-16%)	1300 (820-1900)	70000 (45000-100000)
Myanmar	2005	Male	40 to 44	36% (30-42%)	64% (58-70%)	1.7 (1.1-2.6)	11% (7.1-14%)	13% (8.6-18%)	1600 (1000-2300)	74000 (49000-110000)
Myanmar	2005	Male	45 to 49	35% (28-40%)	65% (60-72%)	1.8 (1.1-2.8)	11% (7.5-16%)	14% (9-19%)	1900 (1200-2700)	79000 (51000-110000)
Myanmar	2005	Male	50 to 54	32% (26-37%)	68% (63-74%)	1.7 (0.96-2.6)	10% (6.3-15%)	12% (7.6-18%)	1800 (1100-2800)	69000 (41000-100000)
Myanmar	2005	Male	55 to 59	26% (20-32%)	74% (68-80%)	1.2 (0.73-2)	7.5% (4.7-11%)	9.1% (5.5-13%)	1200 (730-2000)	40000 (24000-63000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	2005	Male	60 to 64	20% (15-26%)	80% (74-85%)	0.9 (0.49-1.5)	5.6% (3.1-8.9%)	6.6% (3.6-10%)	1000 (550-1700)	28000 (15000-46000)
Myanmar	2005	Male	65 to 69	17% (12-22%)	83% (78-88%)	0.72 (0.4-1.2)	4.2% (2.4-6.7%)	4.9% (2.8-8%)	870 (480-1400)	20000 (11000-32000)
Myanmar	2005	Male	70 to 74	15% (10-20%)	85% (80-90%)	0.6 (0.3-1)	3.5% (1.8-5.9%)	4% (2.1-6.9%)	800 (400-1400)	15000 (7400-25000)
Myanmar	2005	Male	75 to 79	14% (9.4-20%)	86% (80-91%)	0.56 (0.27-1)	2.7% (1.3-4.9%)	3.2% (1.5-5.6%)	580 (260-1100)	8200 (3800-15000)
Myanmar	2005	Male	80 to 84	14% (9.4-20%)	86% (80-91%)	0.52 (0.26-0.88)	2.5% (1.1-4.2%)	2.9% (1.3-5%)	340 (150-600)	3600 (1600-6300)
Myanmar	2005	Male	85 to 89	15% (9.3-22%)	85% (78-91%)	0.48 (0.22-0.85)	1.9% (0.88-3.5%)	2.2% (1-4%)	140 (62-260)	1100 (470-1900)
Myanmar	2005	Male	90 to 94	16% (11-24%)	84% (76-89%)	0.47 (0.22-0.84)	1.5% (0.58-2.9%)	1.8% (0.65-3.4%)	40 (15-78)	210 (78-400)
Myanmar	2005	Male	95 plus	19% (11-29%)	81% (71-89%)	0.47 (0.18-0.89)	1.4% (0.42-2.8%)	1.5% (0.41-3.2%)	8 (2.1-17)	28 (9-58)
Myanmar	2010	Female	All Ages	2.2% (1.9-2.7%)	98% (97-98%)	0.045 (0.029-0.064)	0.67% (0.54-0.82%)	0.88% (0.67-1.1%)	1600 (1200-2000)	57000 (45000-69000)
Myanmar	2010	Female	15 to 19	1.9% (1.1-3%)	98% (97-99%)	0.037 (0.017-0.073)	0.44% (0.28-0.65%)	0.32% (0.18-0.49%)	6 (3.5-9.1)	1400 (880-2100)
Myanmar	2010	Female	20 to 24	2.4% (1.3-3.7%)	98% (96-99%)	0.049 (0.021-0.097)	0.82% (0.54-1.2%)	0.41% (0.24-0.66%)	9.9 (5.6-16)	2800 (1800-4200)
Myanmar	2010	Female	25 to 29	2.6% (1.4-4.2%)	97% (96-99%)	0.054 (0.023-0.11)	0.85% (0.61-1.2%)	0.5% (0.32-0.76%)	15 (9.6-23)	3200 (2200-4600)
Myanmar	2010	Female	30 to 34	2.8% (1.5-4.5%)	97% (95-98%)	0.058 (0.026-0.11)	0.89% (0.66-1.2%)	0.78% (0.54-1.1%)	30 (20-43)	3800 (2700-5100)
Myanmar	2010	Female	35 to 39	2.9% (1.6-4.5%)	97% (95-98%)	0.063 (0.03-0.12)	1% (0.77-1.3%)	1.2% (0.83-1.7%)	58 (39-82)	4600 (3300-6200)
Myanmar	2010	Female	40 to 44	3% (1.8-4.5%)	97% (96-98%)	0.068 (0.033-0.12)	1.2% (0.88-1.5%)	1.5% (1.1-2.1%)	93 (65-130)	5600 (4200-7400)
Myanmar	2010	Female	45 to 49	3% (1.6-4.5%)	97% (95-98%)	0.072 (0.032-0.13)	1.2% (0.87-1.6%)	1.6% (1.1-2.2%)	120 (84-180)	6100 (4500-8200)
Myanmar	2010	Female	50 to 54	3% (1.8-4.7%)	97% (95-98%)	0.08 (0.036-0.15)	1.2% (0.84-1.7%)	1.6% (1.1-2.2%)	160 (110-240)	6600 (4700-9200)
Myanmar	2010	Female	55 to 59	3.1% (1.7-4.8%)	97% (95-98%)	0.087 (0.041-0.17)	1.1% (0.73-1.5%)	1.4% (0.91-2%)	180 (120-270)	6300 (4200-8900)
Myanmar	2010	Female	60 to 64	2.9% (1.7-4.8%)	97% (95-98%)	0.087 (0.039-0.16)	0.98% (0.64-1.4%)	1.3% (0.81-1.8%)	160 (100-230)	4600 (3000-6600)
Myanmar	2010	Female	65 to 69	2.7% (1.5-4.1%)	97% (96-98%)	0.083 (0.04-0.15)	0.86% (0.58-1.2%)	1.1% (0.72-1.6%)	170 (110-240)	3800 (2600-5600)
Myanmar	2010	Female	70 to 74	2.6% (1.6-4.1%)	97% (96-98%)	0.082 (0.035-0.16)	0.78% (0.49-1.2%)	0.96% (0.59-1.5%)	180 (120-270)	3300 (2100-5100)
Myanmar	2010	Female	75 to 79	2.7% (1.6-4.5%)	97% (96-98%)	0.084 (0.037-0.17)	0.62% (0.37-0.99%)	0.76% (0.45-1.2%)	160 (93-250)	2200 (1300-3600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	2010	Female	80 to 84	3% (1.6-4.5%)	97% (95-98%)	0.088 (0.038-0.16)	0.57% (0.34-0.91%)	0.69% (0.41-1.1%)	120 (72-190)	1300 (740-2000)
Myanmar	2010	Female	85 to 89	3.3% (1.8-5.5%)	97% (94-98%)	0.097 (0.041-0.19)	0.49% (0.28-0.81%)	0.58% (0.33-0.97%)	68 (38-110)	500 (290-840)
Myanmar	2010	Female	90 to 94	3.9% (2.2-6.5%)	96% (94-98%)	0.11 (0.046-0.21)	0.42% (0.22-0.69%)	0.49% (0.24-0.83%)	22 (11-37)	110 (58-190)
Myanmar	2010	Female	95 plus	4.8% (2.3-9.2%)	95% (91-98%)	0.13 (0.043-0.28)	0.47% (0.21-0.87%)	0.52% (0.22-0.97%)	6.2 (2.6-11)	22 (9.7-39)
Myanmar	2010	Male	All Ages	20% (17-23%)	80% (77-83%)	0.86 (0.61-1.2)	5% (3.5-6.6%)	6.4% (4.3-8.4%)	14000 (9300-20000)	540000 (380000-740000)
Myanmar	2010	Male	15 to 19	15% (9.9-21%)	85% (79-90%)	0.32 (0.16-0.56)	1.5% (0.95-2.4%)	1.9% (0.95-3.1%)	71 (34-120)	6700 (4000-10000)
Myanmar	2010	Male	20 to 24	25% (16-33%)	75% (67-84%)	0.83 (0.43-1.4)	4.5% (2.7-6.8%)	5.2% (2.9-8.3%)	300 (160-510)	24000 (15000-39000)
Myanmar	2010	Male	25 to 29	29% (19-39%)	71% (61-81%)	1.2 (0.69-1.9)	6.5% (4.1-9.5%)	7.8% (4.5-12%)	580 (340-930)	41000 (26000-63000)
Myanmar	2010	Male	30 to 34	31% (21-43%)	69% (57-79%)	1.5 (0.84-2.4)	7.7% (4.9-11%)	9.3% (5.6-14%)	840 (510-1300)	53000 (34000-76000)
Myanmar	2010	Male	35 to 39	33% (22-43%)	67% (57-78%)	1.7 (0.93-2.7)	8.9% (6-13%)	11% (7.2-16%)	1100 (720-1700)	63000 (41000-91000)
Myanmar	2010	Male	40 to 44	32% (22-42%)	68% (58-78%)	1.8 (1-2.9)	9.6% (6.5-13%)	12% (8.1-16%)	1400 (880-2100)	69000 (44000-100000)
Myanmar	2010	Male	45 to 49	30% (20-40%)	70% (60-80%)	1.9 (1-3.1)	10% (6.8-14%)	13% (8.6-18%)	1700 (1100-2600)	74000 (47000-110000)
Myanmar	2010	Male	50 to 54	28% (19-38%)	72% (62-81%)	1.7 (0.93-2.8)	9.7% (5.8-14%)	12% (7.3-17%)	1800 (1100-2700)	68000 (41000-100000)
Myanmar	2010	Male	55 to 59	25% (17-34%)	75% (66-83%)	1.4 (0.74-2.4)	7.8% (4.7-11%)	9.5% (5.8-14%)	1600 (950-2500)	52000 (32000-81000)
Myanmar	2010	Male	60 to 64	23% (15-32%)	77% (68-85%)	1.2 (0.62-2.1)	6.3% (3.5-9.9%)	7.5% (4.2-12%)	1200 (640-1900)	33000 (18000-53000)
Myanmar	2010	Male	65 to 69	20% (14-27%)	80% (73-86%)	1 (0.55-1.8)	5.1% (2.8-8.1%)	6% (3.3-9.8%)	1000 (550-1700)	24000 (13000-39000)
Myanmar	2010	Male	70 to 74	18% (12-24%)	82% (76-88%)	0.87 (0.42-1.5)	4.2% (2.1-7%)	4.9% (2.4-8.3%)	930 (450-1600)	17000 (8400-30000)
Myanmar	2010	Male	75 to 79	16% (10-23%)	84% (77-90%)	0.74 (0.34-1.4)	3.2% (1.5-5.7%)	3.7% (1.8-6.5%)	680 (310-1200)	9800 (4500-17000)
Myanmar	2010	Male	80 to 84	15% (9.1-21%)	85% (79-91%)	0.62 (0.29-1.1)	2.7% (1.3-4.7%)	3.1% (1.4-5.4%)	410 (190-720)	4300 (2000-7600)
Myanmar	2010	Male	85 to 89	14% (8.5-21%)	86% (79-91%)	0.54 (0.25-0.97)	2.1% (0.99-3.7%)	2.4% (1.1-4.3%)	180 (85-330)	1400 (660-2500)
Myanmar	2010	Male	90 to 94	15% (9.8-23%)	85% (77-90%)	0.52 (0.24-0.95)	1.6% (0.69-3.1%)	1.9% (0.77-3.6%)	51 (21-99)	270 (110-510)
Myanmar	2010	Male	95 plus	18% (9.8-28%)	82% (72-90%)	0.53 (0.2-1)	1.5% (0.53-3%)	1.6% (0.52-3.4%)	11 (3.4-23)	39 (14-80)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	2016	Female	All Ages	2.3% (1.9-2.7%)	98% (97-98%)	0.054 (0.032-0.082)	0.78% (0.63-0.96%)	0.99% (0.76-1.3%)	1600 (1200-2100)	58000 (46000-71000)
Myanmar	2016	Female	15 to 19	1.9% (1.1-3%)	98% (97-99%)	0.043 (0.018-0.088)	0.48% (0.3-0.69%)	0.35% (0.21-0.57%)	5.4 (3.3-8.7)	1400 (850-2100)
Myanmar	2016	Female	20 to 24	2.4% (1.3-3.6%)	98% (96-99%)	0.057 (0.023-0.12)	0.92% (0.62-1.3%)	0.45% (0.27-0.75%)	8.2 (4.8-13)	2800 (1800-4200)
Myanmar	2016	Female	25 to 29	2.6% (1.4-4.1%)	97% (96-99%)	0.062 (0.027-0.12)	0.96% (0.67-1.3%)	0.56% (0.36-0.85%)	12 (7.6-18)	3100 (2000-4300)
Myanmar	2016	Female	30 to 34	2.7% (1.4-4.6%)	97% (95-99%)	0.067 (0.028-0.13)	0.98% (0.72-1.3%)	0.87% (0.6-1.2%)	24 (16-34)	3500 (2400-4900)
Myanmar	2016	Female	35 to 39	2.9% (1.5-4.4%)	97% (96-98%)	0.072 (0.032-0.14)	1.1% (0.8-1.4%)	1.3% (0.92-1.8%)	50 (34-70)	4400 (3100-5700)
Myanmar	2016	Female	40 to 44	2.9% (1.7-4.5%)	97% (96-98%)	0.078 (0.036-0.15)	1.2% (0.87-1.6%)	1.6% (1.2-2.3%)	85 (60-120)	5400 (4000-7200)
Myanmar	2016	Female	45 to 49	2.9% (1.6-4.5%)	97% (96-98%)	0.084 (0.035-0.16)	1.2% (0.88-1.6%)	1.7% (1.2-2.3%)	120 (82-170)	6100 (4500-8300)
Myanmar	2016	Female	50 to 54	3% (1.8-4.6%)	97% (95-98%)	0.092 (0.041-0.18)	1.2% (0.83-1.7%)	1.6% (1.1-2.3%)	150 (100-220)	6500 (4500-8900)
Myanmar	2016	Female	55 to 59	3% (1.7-4.6%)	97% (95-98%)	0.1 (0.042-0.19)	1.1% (0.74-1.5%)	1.5% (0.95-2.1%)	180 (110-260)	6300 (4300-8900)
Myanmar	2016	Female	60 to 64	2.9% (1.7-4.7%)	97% (95-98%)	0.1 (0.044-0.19)	1% (0.64-1.4%)	1.3% (0.85-1.9%)	210 (130-310)	6100 (4000-8800)
Myanmar	2016	Female	65 to 69	2.7% (1.6-4%)	97% (96-98%)	0.096 (0.043-0.18)	0.9% (0.6-1.3%)	1.2% (0.77-1.7%)	190 (120-280)	4300 (2800-6400)
Myanmar	2016	Female	70 to 74	2.6% (1.6-4%)	97% (96-98%)	0.095 (0.039-0.19)	0.82% (0.51-1.3%)	1.1% (0.65-1.6%)	180 (110-270)	3300 (2000-5000)
Myanmar	2016	Female	75 to 79	2.7% (1.6-4.3%)	97% (96-98%)	0.098 (0.039-0.2)	0.67% (0.39-1.1%)	0.83% (0.49-1.3%)	160 (91-250)	2200 (1300-3600)
Myanmar	2016	Female	80 to 84	2.9% (1.6-4.4%)	97% (96-98%)	0.1 (0.041-0.19)	0.62% (0.36-1%)	0.76% (0.45-1.3%)	130 (73-210)	1300 (770-2200)
Myanmar	2016	Female	85 to 89	3.3% (1.8-5.3%)	97% (95-98%)	0.11 (0.046-0.22)	0.53% (0.29-0.93%)	0.64% (0.35-1.1%)	83 (45-140)	620 (330-1100)
Myanmar	2016	Female	90 to 94	3.9% (2.2-6.5%)	96% (93-98%)	0.13 (0.051-0.26)	0.47% (0.23-0.8%)	0.55% (0.26-0.94%)	32 (15-56)	170 (83-290)
Myanmar	2016	Female	95 plus	4.9% (2.4-9.2%)	95% (91-98%)	0.16 (0.05-0.34)	0.52% (0.22-1%)	0.59% (0.24-1.1%)	9.2 (3.7-18)	33 (14-63)
Myanmar	2016	Male	All Ages	20% (17-23%)	80% (77-83%)	1.1 (0.67-1.6)	5.8% (4.1-7.7%)	7.1% (4.7-9.5%)	15000 (9700-20000)	560000 (390000-750000)
Myanmar	2016	Male	15 to 19	15% (9.8-21%)	85% (79-90%)	0.37 (0.17-0.68)	1.7% (1-2.7%)	2.1% (1.1-3.7%)	78 (40-140)	7400 (4400-12000)
Myanmar	2016	Male	20 to 24	24% (16-34%)	76% (66-84%)	0.97 (0.47-1.7)	4.9% (2.9-7.7%)	5.7% (3-9.5%)	310 (160-520)	25000 (15000-40000)
Myanmar	2016	Male	25 to 29	29% (20-39%)	71% (61-80%)	1.4 (0.72-2.4)	7% (4.4-10%)	8.6% (5.1-13%)	500 (280-790)	36000 (22000-55000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Myanmar	2016	Male	30 to 34	31% (21-43%)	69% (57-79%)	1.7 (0.93-2.9)	8.3% (5.4-12%)	10% (6.4-15%)	740 (440-1100)	47000 (30000-69000)
Myanmar	2016	Male	35 to 39	32% (22-43%)	68% (57-78%)	1.9 (1-3.4)	9.5% (6.3-13%)	12% (8-17%)	1000 (660-1500)	58000 (39000-83000)
Myanmar	2016	Male	40 to 44	32% (22-42%)	68% (58-78%)	2.1 (1.1-3.6)	10% (6.9-14%)	13% (9-17%)	1300 (860-1800)	65000 (44000-89000)
Myanmar	2016	Male	45 to 49	30% (19-39%)	70% (61-81%)	2.2 (1.1-3.8)	11% (7.2-15%)	14% (9.3-19%)	1700 (1100-2400)	74000 (48000-100000)
Myanmar	2016	Male	50 to 54	28% (19-38%)	72% (62-81%)	2 (1-3.6)	10% (6.3-14%)	13% (8-17%)	1800 (1100-2600)	70000 (44000-97000)
Myanmar	2016	Male	55 to 59	25% (17-35%)	75% (65-83%)	1.7 (0.82-3)	8.3% (5-12%)	10% (6.1-15%)	1800 (1100-2600)	59000 (36000-86000)
Myanmar	2016	Male	60 to 64	23% (15-33%)	77% (67-85%)	1.4 (0.67-2.5)	6.7% (3.6-10%)	8% (4.4-12%)	1700 (920-2600)	47000 (26000-73000)
Myanmar	2016	Male	65 to 69	20% (13-27%)	80% (73-87%)	1.2 (0.6-2.2)	5.4% (2.9-8.6%)	6.5% (3.4-10%)	1300 (680-2100)	30000 (16000-48000)
Myanmar	2016	Male	70 to 74	18% (12-24%)	82% (76-88%)	1 (0.46-1.8)	4.5% (2.1-7.4%)	5.3% (2.4-8.8%)	1000 (480-1700)	19000 (9200-32000)
Myanmar	2016	Male	75 to 79	16% (10-23%)	84% (77-90%)	0.85 (0.38-1.6)	3.5% (1.6-6.1%)	4% (1.8-7.1%)	760 (340-1300)	11000 (4900-19000)
Myanmar	2016	Male	80 to 84	14% (9.1-20%)	86% (80-91%)	0.71 (0.31-1.3)	3% (1.3-5.1%)	3.4% (1.5-5.9%)	470 (210-810)	5000 (2200-8600)
Myanmar	2016	Male	85 to 89	14% (8.4-21%)	86% (79-92%)	0.62 (0.26-1.2)	2.3% (1-4.1%)	2.6% (1.2-4.7%)	230 (110-430)	1800 (810-3200)
Myanmar	2016	Male	90 to 94	15% (9.5-23%)	85% (77-90%)	0.6 (0.25-1.2)	1.8% (0.73-3.5%)	2.1% (0.84-4%)	76 (31-150)	400 (160-780)
Myanmar	2016	Male	95 plus	18% (10-28%)	82% (72-90%)	0.62 (0.22-1.3)	1.7% (0.57-3.4%)	1.8% (0.56-3.8%)	16 (4.9-34)	57 (20-120)
Philippines	1990	Female	All Ages	16% (13-19%)	84% (81-87%)	0.38 (0.32-0.47)	1.2% (0.78-1.7%)	2.1% (1.2-3.3%)	3300 (1800-5100)	120000 (79000-170000)
Philippines	1990	Female	15 to 19	16% (11-23%)	84% (77-89%)	0.4 (0.23-0.64)	1.1% (0.66-1.8%)	1.8% (0.88-3.1%)	52 (26-91)	5300 (3100-8400)
Philippines	1990	Female	20 to 24	20% (13-28%)	80% (72-87%)	0.55 (0.31-0.89)	1.8% (1.1-2.7%)	2.3% (1.1-4%)	77 (35-130)	8300 (4800-13000)
Philippines	1990	Female	25 to 29	22% (14-31%)	78% (69-86%)	0.62 (0.36-0.97)	2.1% (1.2-3.3%)	2.9% (1.4-5%)	110 (50-180)	9500 (5600-15000)
Philippines	1990	Female	30 to 34	23% (15-33%)	77% (67-85%)	0.68 (0.39-1)	2.3% (1.4-3.6%)	3.5% (1.7-5.7%)	130 (67-220)	10000 (5800-15000)
Philippines	1990	Female	35 to 39	24% (15-32%)	76% (68-85%)	0.74 (0.42-1.1)	2.7% (1.5-4.4%)	4.3% (2.1-7.1%)	180 (86-290)	11000 (6100-17000)
Philippines	1990	Female	40 to 44	23% (16-31%)	77% (69-84%)	0.74 (0.43-1.1)	2.8% (1.5-4.5%)	4.6% (2.4-7.5%)	180 (90-290)	9400 (5000-15000)
Philippines	1990	Female	45 to 49	22% (14-30%)	78% (70-86%)	0.71 (0.4-1.2)	2.9% (1.4-4.9%)	4.6% (2.1-7.8%)	200 (94-350)	9200 (4500-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	1990	Female	50 to 54	22% (15-31%)	78% (69-85%)	0.79 (0.45-1.3)	3.3% (1.6-5.8%)	5% (2.4-8.7%)	270 (130-480)	11000 (5100-18000)
Philippines	1990	Female	55 to 59	23% (15-31%)	77% (69-85%)	0.97 (0.56-1.5)	4.2% (2-7%)	6.2% (3-10%)	370 (180-620)	12000 (6000-21000)
Philippines	1990	Female	60 to 64	21% (14-30%)	79% (70-86%)	0.93 (0.54-1.5)	3.9% (1.9-6.8%)	5.5% (2.6-9.7%)	390 (180-690)	11000 (5300-19000)
Philippines	1990	Female	65 to 69	19% (13-27%)	81% (73-87%)	0.76 (0.44-1.2)	3.1% (1.3-5.4%)	4.3% (1.9-7.5%)	360 (160-640)	8400 (3800-15000)
Philippines	1990	Female	70 to 74	18% (13-26%)	82% (74-87%)	0.65 (0.35-1.1)	2.3% (0.64-4.7%)	3.1% (0.88-6.2%)	300 (86-610)	5700 (1500-11000)
Philippines	1990	Female	75 to 79	19% (12-27%)	81% (73-88%)	0.62 (0.32-1.1)	1.8% (0.19-4.2%)	2.2% (0.23-5.3%)	250 (27-600)	3700 (380-8800)
Philippines	1990	Female	80 to 84	20% (13-27%)	80% (73-87%)	0.65 (0.36-1.1)	1.6% (0.18-3.7%)	1.9% (0.19-4.3%)	210 (22-500)	2300 (250-5300)
Philippines	1990	Female	85 to 89	22% (14-31%)	78% (69-86%)	0.7 (0.37-1.2)	1.4% (0.039-3.3%)	1.6% (0.019-3.8%)	130 (1.5-310)	960 (28-2300)
Philippines	1990	Female	90 to 94	24% (16-34%)	76% (66-84%)	0.8 (0.42-1.4)	1.2% (-0.14-3.3%)	1.3% (-0.24-3.8%)	46 (-8.3-130)	250 (-30-690)
Philippines	1990	Female	95 plus	28% (16-42%)	72% (58-84%)	0.93 (0.42-1.7)	1.3% (-0.34-4.2%)	1.4% (-0.51-4.8%)	18 (-6.5-59)	65 (-16-200)
Philippines	1990	Male	All Ages	56% (50-62%)	44% (38-50%)	1.7 (1.5-1.9)	5.8% (4.7-6.9%)	8.2% (6.5-10%)	18000 (14000-23000)	780000 (630000-930000)
Philippines	1990	Male	15 to 19	54% (43-65%)	46% (35-57%)	1.3 (0.79-1.8)	4.3% (2.8-6%)	6.1% (3.8-8.6%)	320 (200-450)	27000 (17000-37000)
Philippines	1990	Male	20 to 24	74% (64-81%)	26% (19-36%)	2.8 (2-3.7)	11% (7.8-14%)	14% (9.7-18%)	1000 (730-1400)	79000 (57000-100000)
Philippines	1990	Male	25 to 29	77% (69-84%)	23% (16-31%)	3.1 (2.3-4)	12% (9.3-16%)	16% (12-20%)	1300 (940-1700)	89000 (67000-110000)
Philippines	1990	Male	30 to 34	79% (71-85%)	21% (15-29%)	3.4 (2.7-4.3)	13% (10-16%)	17% (13-22%)	1400 (1000-1800)	87000 (66000-110000)
Philippines	1990	Male	35 to 39	79% (71-85%)	21% (15-29%)	3.8 (2.7-4.9)	15% (11-18%)	19% (14-24%)	1600 (1200-2100)	90000 (67000-110000)
Philippines	1990	Male	40 to 44	78% (71-84%)	22% (16-29%)	3.9 (2.8-5.1)	15% (11-19%)	20% (15-25%)	1600 (1200-2100)	79000 (58000-100000)
Philippines	1990	Male	45 to 49	75% (66-82%)	25% (18-34%)	3.9 (2.7-5.3)	16% (11-21%)	20% (14-26%)	1800 (1300-2400)	80000 (55000-110000)
Philippines	1990	Male	50 to 54	71% (62-78%)	29% (22-38%)	3.6 (2.5-5)	15% (10-20%)	19% (13-25%)	1900 (1300-2600)	73000 (48000-97000)
Philippines	1990	Male	55 to 59	66% (55-75%)	34% (25-45%)	3.3 (2.3-4.6)	14% (9.3-19%)	17% (12-23%)	1900 (1300-2700)	63000 (42000-88000)
Philippines	1990	Male	60 to 64	62% (49-72%)	38% (28-51%)	2.8 (1.9-4.2)	12% (7.2-17%)	14% (8.5-21%)	1700 (1000-2500)	47000 (28000-69000)
Philippines	1990	Male	65 to 69	58% (45-70%)	42% (30-55%)	2.2 (1.5-3.2)	8.9% (5.2-14%)	11% (6.3-16%)	1300 (750-2000)	30000 (18000-46000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	1990	Male	70 to 74	53% (40-65%)	47% (35-60%)	1.7 (1.1-2.6)	6.3% (3.1-10%)	7.4% (3.6-12%)	920 (440-1500)	17000 (8400-29000)
Philippines	1990	Male	75 to 79	47% (33-62%)	53% (38-67%)	1.4 (0.81-2.3)	4.3% (1.5-8%)	4.9% (1.6-9.4%)	610 (190-1200)	8800 (2900-17000)
Philippines	1990	Male	80 to 84	42% (29-56%)	58% (44-71%)	1.2 (0.69-1.8)	3% (0.77-5.7%)	3.2% (0.76-6.4%)	350 (80-690)	3700 (950-7400)
Philippines	1990	Male	85 to 89	41% (28-55%)	59% (45-72%)	1 (0.58-1.6)	1.9% (0.29-3.9%)	2% (0.2-4.4%)	140 (14-310)	1100 (170-2400)
Philippines	1990	Male	90 to 94	43% (29-57%)	57% (43-71%)	0.99 (0.57-1.6)	1.2% (-0.21-3.1%)	1.2% (-0.42-3.4%)	29 (-10-83)	170 (-31-440)
Philippines	1990	Male	95 plus	46% (28-64%)	54% (36-72%)	0.99 (0.48-1.7)	0.99% (-0.51-3.3%)	0.9% (-0.84-3.4%)	5.3 (-4.9-20)	23 (-12-76)
Philippines	1995	Female	All Ages	16% (14-19%)	84% (81-86%)	0.39 (0.32-0.48)	1.2% (0.85-1.7%)	2.1% (1.2-3.3%)	3400 (1900-5300)	120000 (84000-170000)
Philippines	1995	Female	15 to 19	16% (11-22%)	84% (78-89%)	0.4 (0.23-0.62)	1.1% (0.66-1.6%)	1.7% (0.89-2.9%)	46 (24-78)	5100 (3100-7900)
Philippines	1995	Female	20 to 24	20% (13-29%)	80% (71-87%)	0.54 (0.29-0.88)	1.7% (1-2.6%)	2.3% (1-3.9%)	71 (33-120)	8200 (4900-12000)
Philippines	1995	Female	25 to 29	22% (14-30%)	78% (70-86%)	0.61 (0.36-0.97)	2% (1.2-3.1%)	2.8% (1.4-4.9%)	100 (50-170)	9500 (5800-15000)
Philippines	1995	Female	30 to 34	22% (15-33%)	78% (67-85%)	0.67 (0.38-1.1)	2.2% (1.3-3.2%)	3.3% (1.8-5.2%)	130 (68-200)	10000 (6100-15000)
Philippines	1995	Female	35 to 39	23% (15-32%)	77% (68-85%)	0.73 (0.42-1.1)	2.5% (1.5-3.8%)	4% (2.1-6.5%)	170 (90-270)	11000 (6500-17000)
Philippines	1995	Female	40 to 44	23% (16-31%)	77% (69-84%)	0.74 (0.44-1.1)	2.5% (1.4-3.9%)	4.3% (2.4-6.6%)	200 (110-310)	11000 (6300-17000)
Philippines	1995	Female	45 to 49	22% (14-30%)	78% (70-86%)	0.71 (0.39-1.1)	2.6% (1.4-4.4%)	4.2% (2.1-6.9%)	210 (100-340)	9600 (5000-16000)
Philippines	1995	Female	50 to 54	22% (15-30%)	78% (70-85%)	0.79 (0.45-1.3)	3% (1.5-5%)	4.6% (2.3-7.6%)	270 (140-450)	11000 (5400-18000)
Philippines	1995	Female	55 to 59	23% (15-31%)	77% (69-85%)	0.98 (0.55-1.6)	3.8% (1.9-6.2%)	5.6% (2.9-9.2%)	380 (200-620)	13000 (6800-21000)
Philippines	1995	Female	60 to 64	21% (14-31%)	79% (69-86%)	0.94 (0.53-1.5)	3.7% (1.7-6.3%)	5.1% (2.4-8.7%)	450 (200-760)	13000 (5800-22000)
Philippines	1995	Female	65 to 69	19% (13-27%)	81% (73-87%)	0.77 (0.46-1.2)	2.9% (1.3-4.9%)	3.9% (1.9-6.7%)	370 (180-640)	8700 (4100-15000)
Philippines	1995	Female	70 to 74	18% (13-25%)	82% (75-87%)	0.65 (0.36-1.1)	2.2% (0.63-4.2%)	2.9% (0.89-5.5%)	340 (100-650)	6300 (1800-12000)
Philippines	1995	Female	75 to 79	19% (12-26%)	81% (74-88%)	0.62 (0.32-1.1)	1.6% (0.16-3.8%)	2.1% (0.25-4.8%)	260 (30-600)	3700 (370-8700)
Philippines	1995	Female	80 to 84	20% (13-27%)	80% (73-87%)	0.65 (0.37-1.1)	1.4% (0.056-3.4%)	1.7% (0.021-3.9%)	230 (2.9-540)	2400 (94-5700)
Philippines	1995	Female	85 to 89	21% (14-30%)	79% (70-86%)	0.7 (0.39-1.2)	1.2% (-0.096-2.9%)	1.3% (-0.15-3.4%)	150 (-16-360)	1100 (-88-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	1995	Female	90 to 94	24% (16-34%)	76% (66-84%)	0.79 (0.42-1.4)	1% (-0.27-3.1%)	1.1% (-0.4-3.6%)	51 (-18-160)	280 (-73-850)
Philippines	1995	Female	95 plus	28% (16-42%)	72% (58-84%)	0.92 (0.41-1.7)	1.2% (-0.49-4.1%)	1.3% (-0.67-4.6%)	18 (-9.5-66)	68 (-27-230)
Philippines	1995	Male	All Ages	56% (50-62%)	44% (38-50%)	1.8 (1.6-2)	6% (4.9-7.1%)	8.2% (6.4-10%)	19000 (15000-24000)	810000 (660000-970000)
Philippines	1995	Male	15 to 19	54% (43-65%)	46% (35-57%)	1.3 (0.81-1.8)	4.1% (2.8-5.7%)	6.2% (4.1-8.6%)	300 (200-430)	26000 (18000-36000)
Philippines	1995	Male	20 to 24	74% (63-81%)	26% (19-37%)	2.8 (2-3.6)	10% (7.4-13%)	13% (9.7-17%)	900 (650-1200)	71000 (53000-90000)
Philippines	1995	Male	25 to 29	77% (69-84%)	23% (16-31%)	3.1 (2.3-3.9)	11% (8.6-14%)	15% (11-19%)	1200 (840-1500)	82000 (60000-100000)
Philippines	1995	Male	30 to 34	79% (71-85%)	21% (15-29%)	3.4 (2.6-4.3)	12% (9.1-15%)	16% (12-20%)	1300 (970-1700)	84000 (63000-110000)
Philippines	1995	Male	35 to 39	79% (71-85%)	21% (15-29%)	3.8 (2.7-4.9)	13% (10-17%)	18% (13-23%)	1600 (1200-2100)	90000 (67000-110000)
Philippines	1995	Male	40 to 44	78% (70-84%)	22% (16-30%)	3.9 (2.9-5)	14% (10-18%)	19% (14-24%)	1800 (1300-2400)	92000 (67000-120000)
Philippines	1995	Male	45 to 49	75% (65-82%)	25% (18-35%)	3.9 (2.7-5.3)	15% (10-19%)	19% (13-24%)	2000 (1400-2600)	87000 (61000-110000)
Philippines	1995	Male	50 to 54	71% (62-78%)	29% (22-38%)	3.6 (2.5-5)	14% (9.4-19%)	17% (11-23%)	2100 (1400-2900)	81000 (53000-110000)
Philippines	1995	Male	55 to 59	66% (55-75%)	34% (25-45%)	3.3 (2.3-4.6)	13% (8.6-18%)	16% (11-22%)	2200 (1400-3000)	71000 (47000-98000)
Philippines	1995	Male	60 to 64	62% (49-73%)	38% (27-51%)	2.9 (1.9-4.2)	11% (7-17%)	13% (8.2-20%)	2100 (1300-3100)	59000 (37000-86000)
Philippines	1995	Male	65 to 69	58% (45-70%)	42% (30-55%)	2.2 (1.5-3.2)	8.4% (4.9-13%)	9.8% (5.9-15%)	1500 (870-2200)	34000 (20000-51000)
Philippines	1995	Male	70 to 74	53% (41-65%)	47% (35-59%)	1.8 (1.1-2.6)	5.9% (2.9-9.7%)	6.8% (3.3-11%)	1100 (500-1700)	20000 (9500-33000)
Philippines	1995	Male	75 to 79	46% (32-61%)	54% (39-68%)	1.4 (0.81-2.3)	4% (1.3-7.6%)	4.5% (1.4-8.7%)	630 (190-1200)	9100 (3100-17000)
Philippines	1995	Male	80 to 84	42% (29-56%)	58% (44-71%)	1.2 (0.69-1.8)	2.6% (0.55-5.3%)	2.8% (0.45-5.9%)	350 (56-730)	3800 (810-7700)
Philippines	1995	Male	85 to 89	41% (28-56%)	59% (44-72%)	1 (0.59-1.6)	1.6% (-0.056-3.7%)	1.7% (-0.17-4%)	140 (-14-330)	1100 (-39-2600)
Philippines	1995	Male	90 to 94	43% (30-57%)	57% (43-70%)	1 (0.57-1.6)	0.92% (-0.49-2.8%)	0.9% (-0.76-3.1%)	29 (-25-100)	180 (-96-540)
Philippines	1995	Male	95 plus	47% (29-65%)	53% (35-71%)	1 (0.48-1.8)	0.81% (-0.78-2.9%)	0.66% (-1.2-3.1%)	5.3 (-9.5-25)	25 (-23-92)
Philippines	2000	Female	All Ages	13% (11-15%)	87% (85-89%)	0.33 (0.28-0.4)	1.2% (0.86-1.7%)	2.1% (1.2-3.2%)	3800 (2200-5700)	130000 (91000-180000)
Philippines	2000	Female	15 to 19	13% (9.1-17%)	87% (83-91%)	0.34 (0.2-0.51)	0.9% (0.57-1.3%)	1.3% (0.72-2.2%)	37 (20-61)	4700 (2800-7000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	2000	Female	20 to 24	15% (11-20%)	85% (80-89%)	0.41 (0.25-0.64)	1.5% (0.93-2.1%)	1.7% (0.87-2.9%)	59 (29-98)	7800 (4800-11000)
Philippines	2000	Female	25 to 29	15% (11-21%)	85% (79-89%)	0.46 (0.29-0.7)	1.7% (1.1-2.5%)	2.2% (1.2-3.6%)	82 (44-140)	8600 (5500-13000)
Philippines	2000	Female	30 to 34	18% (14-23%)	82% (77-86%)	0.55 (0.35-0.81)	1.9% (1.2-2.7%)	2.7% (1.5-4.2%)	120 (65-180)	9600 (6100-14000)
Philippines	2000	Female	35 to 39	19% (14-24%)	81% (76-86%)	0.61 (0.39-0.9)	2.1% (1.3-3.1%)	3.3% (1.9-5.2%)	160 (95-250)	11000 (6800-16000)
Philippines	2000	Female	40 to 44	18% (14-23%)	82% (77-86%)	0.6 (0.39-0.86)	2.2% (1.3-3.2%)	3.5% (2.2-5.4%)	200 (120-300)	11000 (6900-16000)
Philippines	2000	Female	45 to 49	18% (13-23%)	82% (77-87%)	0.59 (0.37-0.91)	2.2% (1.3-3.5%)	3.5% (2-5.6%)	240 (130-370)	11000 (6600-17000)
Philippines	2000	Female	50 to 54	18% (14-24%)	82% (76-86%)	0.68 (0.41-1)	2.6% (1.4-4.2%)	3.9% (2.1-6.2%)	290 (160-460)	11000 (6100-18000)
Philippines	2000	Female	55 to 59	20% (16-26%)	80% (74-84%)	0.92 (0.59-1.4)	3.5% (2-5.7%)	5.2% (2.9-8.3%)	430 (240-700)	15000 (8100-24000)
Philippines	2000	Female	60 to 64	20% (15-26%)	80% (74-85%)	0.91 (0.57-1.4)	3.5% (1.8-5.8%)	4.8% (2.4-7.9%)	510 (260-840)	15000 (7400-25000)
Philippines	2000	Female	65 to 69	16% (12-22%)	84% (78-88%)	0.67 (0.42-1)	2.6% (1.3-4.4%)	3.5% (1.7-5.8%)	440 (220-740)	10000 (5100-18000)
Philippines	2000	Female	70 to 74	14% (10-20%)	86% (80-90%)	0.53 (0.3-0.88)	1.9% (0.68-3.6%)	2.4% (0.87-4.7%)	340 (120-650)	6300 (2300-12000)
Philippines	2000	Female	75 to 79	15% (9.8-22%)	85% (78-90%)	0.52 (0.28-0.91)	1.5% (0.25-3.3%)	1.8% (0.33-4%)	290 (51-640)	4200 (720-9500)
Philippines	2000	Female	80 to 84	18% (11-25%)	82% (75-89%)	0.61 (0.34-1)	1.4% (0.24-3.3%)	1.7% (0.26-3.8%)	260 (40-590)	2700 (450-6200)
Philippines	2000	Female	85 to 89	20% (13-29%)	80% (71-87%)	0.7 (0.39-1.1)	1.3% (0.077-3.1%)	1.5% (0.035-3.5%)	200 (4-5-460)	1500 (86-3400)
Philippines	2000	Female	90 to 94	23% (15-33%)	77% (67-85%)	0.79 (0.42-1.4)	1.2% (-0.13-3.3%)	1.3% (-0.23-3.8%)	88 (-15-250)	470 (-49-1300)
Philippines	2000	Female	95 plus	27% (15-40%)	73% (60-85%)	0.93 (0.42-1.7)	1.5% (-0.28-4.5%)	1.6% (-0.41-5.1%)	34 (-8.6-110)	120 (-22-360)
Philippines	2000	Male	All Ages	60% (56-65%)	40% (35-44%)	2 (1.8-2.2)	6.7% (5.4-7.9%)	8.9% (7-11%)	23000 (18000-29000)	970000 (800000-1200000)
Philippines	2000	Male	15 to 19	63% (56-69%)	37% (31-44%)	1.6 (1.1-2.2)	4.5% (2.8-6.1%)	6.9% (4.3-9.4%)	360 (230-500)	31000 (20000-42000)
Philippines	2000	Male	20 to 24	79% (74-82%)	21% (18-26%)	3.3 (2.4-4.1)	10% (7.5-13%)	14% (9.9-18%)	1000 (740-1300)	79000 (59000-100000)
Philippines	2000	Male	25 to 29	80% (76-84%)	20% (16-24%)	3.4 (2.6-4.3)	11% (8.3-14%)	15% (11-19%)	1200 (870-1600)	86000 (64000-110000)
Philippines	2000	Male	30 to 34	82% (78-85%)	18% (15-22%)	3.8 (3-4.8)	12% (9.1-15%)	16% (12-21%)	1500 (1100-1900)	93000 (69000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	2000	Male	35 to 39	81% (78-85%)	19% (15-22%)	4.2 (3.1-5.4)	14% (9.8-18%)	18% (13-24%)	1900 (1300-2400)	110000 (76000-140000)
Philippines	2000	Male	40 to 44	80% (76-84%)	20% (16-24%)	4.3 (3.3-5.5)	14% (11-18%)	19% (14-24%)	2200 (1600-2800)	110000 (82000-140000)
Philippines	2000	Male	45 to 49	78% (73-82%)	22% (18-27%)	4.3 (3.2-5.7)	15% (11-19%)	19% (14-25%)	2700 (1900-3600)	120000 (84000-150000)
Philippines	2000	Male	50 to 54	73% (68-78%)	27% (22-32%)	3.9 (2.8-5.2)	14% (9.8-19%)	18% (12-23%)	2700 (1800-3700)	100000 (69000-140000)
Philippines	2000	Male	55 to 59	66% (58-74%)	34% (26-42%)	3.5 (2.5-4.7)	13% (8.7-18%)	16% (10-22%)	2700 (1800-3700)	89000 (60000-120000)
Philippines	2000	Male	60 to 64	60% (50-70%)	40% (30-50%)	2.9 (2-4.1)	11% (6.9-16%)	13% (8-19%)	2500 (1600-3700)	70000 (44000-100000)
Philippines	2000	Male	65 to 69	55% (44-66%)	45% (34-56%)	2.2 (1.5-3)	8.1% (5-12%)	9.4% (5.9-14%)	1900 (1200-2900)	45000 (28000-66000)
Philippines	2000	Male	70 to 74	51% (39-61%)	49% (39-61%)	1.7 (1.2-2.5)	5.8% (3-9.3%)	6.6% (3.4-11%)	1200 (610-2000)	23000 (12000-37000)
Philippines	2000	Male	75 to 79	44% (31-57%)	56% (43-69%)	1.4 (0.82-2.2)	4% (1.7-8.3%)	4.4% (1.7-8.3%)	770 (280-1400)	11000 (4400-21000)
Philippines	2000	Male	80 to 84	39% (26-53%)	61% (47-74%)	1.1 (0.67-1.8)	2.7% (0.87-5.3%)	3% (0.86-5.8%)	400 (120-780)	4400 (1400-8400)
Philippines	2000	Male	85 to 89	38% (26-52%)	62% (48-74%)	1 (0.57-1.6)	1.8% (0.21-3.8%)	1.8% (0.11-4.1%)	180 (10-400)	1400 (170-3000)
Philippines	2000	Male	90 to 94	40% (27-55%)	60% (45-73%)	0.96 (0.53-1.6)	1.1% (-0.34-3.2%)	1.1% (-0.54-3.4%)	46 (-22-140)	260 (-78-750)
Philippines	2000	Male	95 plus	44% (27-62%)	56% (38-73%)	0.95 (0.45-1.7)	1% (-0.58-3.6%)	0.93% (-0.85-3.7%)	11 (-9.8-43)	45 (-26-150)
Philippines	2005	Female	All Ages	13% (11-15%)	87% (85-89%)	0.34 (0.29-0.42)	1.3% (0.92-1.8%)	2.2% (1.3-3.4%)	4500 (2500-6800)	150000 (100000-210000)
Philippines	2005	Female	15 to 19	13% (8.8-18%)	87% (82-91%)	0.34 (0.2-0.52)	0.88% (0.57-1.3%)	1.3% (0.7-2.2%)	40 (21-68)	5100 (3200-7500)
Philippines	2005	Female	20 to 24	15% (10-20%)	85% (80-90%)	0.42 (0.25-0.66)	1.5% (0.95-2.1%)	1.7% (0.83-2.8%)	65 (31-110)	8400 (5400-13000)
Philippines	2005	Female	25 to 29	15% (11-20%)	85% (80-89%)	0.46 (0.29-0.71)	1.6% (1.1-2.4%)	2.1% (1.1-3.4%)	89 (48-150)	9400 (5900-14000)
Philippines	2005	Female	30 to 34	17% (13-23%)	83% (77-87%)	0.56 (0.36-0.82)	1.8% (1.2-2.6%)	2.6% (1.4-4.1%)	120 (68-200)	10000 (6500-15000)
Philippines	2005	Female	35 to 39	18% (14-23%)	82% (77-86%)	0.61 (0.39-0.91)	2% (1.3-3%)	3.2% (1.8-4.9%)	180 (100-280)	12000 (7300-18000)
Philippines	2005	Female	40 to 44	18% (13-23%)	82% (77-87%)	0.6 (0.39-0.89)	2.1% (1.2-3.1%)	3.4% (2-5.2%)	220 (130-350)	12000 (7600-19000)
Philippines	2005	Female	45 to 49	18% (13-22%)	82% (78-87%)	0.6 (0.38-0.93)	2.2% (1.2-3.5%)	3.4% (1.8-5.5%)	280 (150-460)	13000 (7000-21000)
Philippines	2005	Female	50 to 54	18% (13-23%)	82% (77-87%)	0.68 (0.42-1)	2.5% (1.3-4.1%)	3.8% (1.9-6.2%)	390 (190-630)	16000 (7600-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	2005	Female	55 to 59	20% (15-25%)	80% (75-85%)	0.92 (0.6-1.4)	3.5% (1.9-5.8%)	5.1% (2.8-8.2%)	530 (290-890)	18000 (9600-30000)
Philippines	2005	Female	60 to 64	19% (14-25%)	81% (75-86%)	0.91 (0.58-1.4)	3.4% (1.7-5.8%)	4.7% (2.3-7.9%)	590 (290-980)	17000 (8200-28000)
Philippines	2005	Female	65 to 69	16% (11-21%)	84% (79-89%)	0.67 (0.43-1)	2.5% (1.2-4.4%)	3.4% (1.7-5.9%)	520 (250-910)	12000 (5900-22000)
Philippines	2005	Female	70 to 74	14% (10-19%)	86% (81-90%)	0.54 (0.3-0.9)	1.9% (0.55-3.8%)	2.4% (0.74-4.8%)	420 (130-830)	7900 (2400-16000)
Philippines	2005	Female	75 to 79	15% (9.9-21%)	85% (79-90%)	0.53 (0.28-0.92)	1.5% (0.22-3.4%)	1.8% (0.3-4.2%)	330 (54-760)	4800 (710-11000)
Philippines	2005	Female	80 to 84	17% (11-25%)	83% (75-89%)	0.61 (0.35-1)	1.4% (0.18-3.5%)	1.7% (0.2-4%)	310 (37-750)	3300 (400-7800)
Philippines	2005	Female	85 to 89	20% (13-29%)	80% (71-87%)	0.72 (0.38-1.2)	1.4% (0.1-3.3%)	1.6% (0.082-3.7%)	250 (13-600)	1900 (140-4500)
Philippines	2005	Female	90 to 94	23% (15-32%)	77% (68-85%)	0.81 (0.42-1.4)	1.3% (-0.093-3.6%)	1.5% (-0.19-4.1%)	120 (-15-330)	610 (-44-1700)
Philippines	2005	Female	95 plus	27% (16-40%)	73% (60-84%)	0.95 (0.43-1.7)	1.6% (-0.17-4.7%)	1.7% (-0.29-5.2%)	50 (-8.4-150)	180 (-18-520)
Philippines	2005	Male	All Ages	61% (56-65%)	39% (35-44%)	2.1 (1.9-2.3)	7.3% (6-8.7%)	9.5% (7.5-12%)	28000 (22000-34000)	1100000 (930000-1400000)
Philippines	2005	Male	15 to 19	63% (56-69%)	37% (31-44%)	1.6 (1.1-2.2)	4.5% (3-6.2%)	7.1% (4.5-9.7%)	410 (260-580)	34000 (22000-47000)
Philippines	2005	Male	20 to 24	78% (74-82%)	22% (18-26%)	3.3 (2.4-4.2)	10% (7.7-13%)	14% (10-19%)	1100 (810-1500)	87000 (63000-110000)
Philippines	2005	Male	25 to 29	80% (76-83%)	20% (17-24%)	3.5 (2.6-4.4)	11% (8.3-14%)	15% (11-19%)	1300 (960-1700)	93000 (68000-120000)
Philippines	2005	Male	30 to 34	82% (78-85%)	18% (15-22%)	3.9 (3-4.9)	12% (9.1-16%)	16% (12-21%)	1600 (1100-2000)	99000 (73000-130000)
Philippines	2005	Male	35 to 39	81% (77-85%)	19% (15-23%)	4.2 (3.2-5.4)	14% (10-18%)	18% (13-24%)	2100 (1500-2700)	120000 (84000-150000)
Philippines	2005	Male	40 to 44	80% (76-84%)	20% (16-24%)	4.4 (3.3-5.6)	15% (11-19%)	19% (14-25%)	2600 (1900-3400)	130000 (95000-160000)
Philippines	2005	Male	45 to 49	78% (73-82%)	22% (18-27%)	4.4 (3.2-5.8)	15% (11-20%)	19% (14-25%)	3200 (2300-4300)	140000 (100000-190000)
Philippines	2005	Male	50 to 54	73% (67-77%)	27% (23-33%)	4 (2.8-5.3)	15% (9.9-19%)	18% (12-24%)	3700 (2500-4900)	140000 (96000-190000)
Philippines	2005	Male	55 to 59	66% (57-74%)	34% (26-43%)	3.5 (2.6-4.8)	13% (8.7-18%)	16% (10-22%)	3400 (2200-4700)	110000 (73000-160000)
Philippines	2005	Male	60 to 64	59% (50-68%)	41% (32-50%)	2.9 (2-4.1)	11% (6.7-16%)	13% (7.8-19%)	3000 (1800-4300)	83000 (51000-120000)
Philippines	2005	Male	65 to 69	55% (44-65%)	45% (35-56%)	2.2 (1.5-3.1)	8.1% (4.9-12%)	9.4% (5.8-14%)	2300 (1400-3400)	53000 (33000-78000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	2005	Male	70 to 74	50% (39-61%)	50% (39-61%)	1.8 (1.2-2.6)	5.8% (3-9.5%)	6.6% (3.4-11%)	1600 (800-2600)	29000 (15000-48000)
Philippines	2005	Male	75 to 79	43% (31-57%)	57% (43-69%)	1.4 (0.82-2.2)	3.9% (1.5-7.3%)	4.3% (1.6-8.2%)	850 (320-1600)	12000 (4700-23000)
Philippines	2005	Male	80 to 84	38% (25-52%)	62% (48-75%)	1.1 (0.67-1.7)	2.6% (0.9-5.1%)	2.9% (0.87-5.6%)	440 (140-880)	4800 (1600-9300)
Philippines	2005	Male	85 to 89	37% (25-51%)	63% (49-75%)	0.99 (0.56-1.5)	1.7% (0.28-3.7%)	1.8% (0.17-4%)	200 (18-440)	1600 (250-3300)
Philippines	2005	Male	90 to 94	39% (26-53%)	61% (47-74%)	0.95 (0.54-1.6)	1.2% (-0.22-3.1%)	1.2% (-0.43-3.3%)	51 (-19-140)	290 (-55-760)
Philippines	2005	Male	95 plus	43% (26-62%)	57% (38-74%)	0.95 (0.45-1.7)	1.1% (-0.52-3.4%)	1% (-0.78-3.5%)	13 (-9.8-46)	53 (-25-170)
Philippines	2010	Female	All Ages	15% (13-18%)	85% (82-87%)	0.41 (0.33-0.51)	1.5% (0.97-2%)	2.2% (1.1-3.6%)	5300 (2700-8400)	180000 (120000-250000)
Philippines	2010	Female	15 to 19	14% (9.4-21%)	86% (79-91%)	0.37 (0.21-0.6)	0.94% (0.58-1.5%)	1.4% (0.64-2.4%)	45 (22-82)	5700 (3500-9200)
Philippines	2010	Female	20 to 24	18% (12-27%)	82% (73-88%)	0.51 (0.27-0.83)	1.6% (1-2.4%)	1.9% (0.86-3.3%)	76 (34-130)	9600 (5800-14000)
Philippines	2010	Female	25 to 29	20% (13-28%)	80% (72-87%)	0.57 (0.33-0.92)	1.8% (1.1-2.7%)	2.3% (1.1-4%)	110 (50-180)	11000 (6500-17000)
Philippines	2010	Female	30 to 34	20% (13-29%)	80% (71-87%)	0.63 (0.36-0.99)	1.9% (1.2-2.8%)	2.6% (1.3-4.3%)	140 (71-230)	12000 (7000-18000)
Philippines	2010	Female	35 to 39	21% (14-29%)	79% (71-86%)	0.68 (0.39-1.1)	2.1% (1.3-3.3%)	3.2% (1.7-5.2%)	200 (110-330)	13000 (7900-21000)
Philippines	2010	Female	40 to 44	21% (14-29%)	79% (71-86%)	0.69 (0.42-1.1)	2.2% (1.2-3.5%)	3.5% (1.9-5.6%)	270 (150-440)	15000 (8900-23000)
Philippines	2010	Female	45 to 49	20% (12-28%)	80% (72-88%)	0.66 (0.36-1.1)	2.2% (1.1-3.7%)	3.4% (1.7-5.9%)	330 (160-560)	16000 (8000-26000)
Philippines	2010	Female	50 to 54	20% (14-28%)	80% (72-86%)	0.74 (0.43-1.2)	2.5% (1.2-4.5%)	3.8% (1.8-6.7%)	450 (210-780)	18000 (8400-31000)
Philippines	2010	Female	55 to 59	21% (14-29%)	79% (71-86%)	0.92 (0.53-1.4)	3.3% (1.7-5.6%)	4.7% (2.4-7.9%)	650 (330-1100)	22000 (11000-38000)
Philippines	2010	Female	60 to 64	20% (13-28%)	80% (72-87%)	0.88 (0.5-1.4)	3.1% (1.5-5.5%)	4.2% (2-7.5%)	700 (340-1300)	20000 (9500-36000)
Philippines	2010	Female	65 to 69	18% (12-25%)	82% (75-88%)	0.72 (0.43-1.1)	2.4% (1-4.4%)	3.2% (1.4-5.8%)	580 (260-1000)	14000 (6000-25000)
Philippines	2010	Female	70 to 74	17% (11-24%)	83% (76-89%)	0.61 (0.33-1)	1.9% (0.42-3.9%)	2.4% (0.62-5%)	560 (140-1200)	10000 (2400-22000)
Philippines	2010	Female	75 to 79	17% (11-25%)	83% (75-89%)	0.59 (0.31-1)	1.4% (0.046-3.5%)	1.8% (0.09-4.3%)	450 (23-1100)	6400 (190-16000)
Philippines	2010	Female	80 to 84	18% (12-25%)	82% (75-88%)	0.62 (0.34-1)	1.3% (0.0026-3.2%)	1.5% (-0.023-3.7%)	350 (-5.3-890)	3700 (7.6-9400)
Philippines	2010	Female	85 to 89	20% (13-28%)	80% (72-87%)	0.67 (0.35-1.1)	1.2% (-0.039-2.9%)	1.3% (-0.076-3.3%)	240 (-15-600)	1800 (-62-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	2010	Female	90 to 94	22% (14-32%)	78% (68-86%)	0.76 (0.4-1.3)	1.1% (-0.25-3.2%)	1.2% (-0.38-3.6%)	120 (-36-350)	620 (-140-1800)
Philippines	2010	Female	95 plus	26% (15-39%)	74% (61-85%)	0.88 (0.4-1.6)	1.4% (-0.37-4.3%)	1.4% (-0.5-4.7%)	48 (-17-160)	180 (-47-550)
Philippines	2010	Male	All Ages	56% (50-62%)	44% (38-50%)	1.9 (1.7-2.1)	6.9% (5.5-8.3%)	8.5% (6.6-11%)	29000 (21000-36000)	1100000 (920000-1400000)
Philippines	2010	Male	15 to 19	51% (40-63%)	49% (37-60%)	1.2 (0.74-1.8)	3.6% (2.3-5.2%)	5.4% (3.2-7.8%)	330 (200-490)	29000 (18000-41000)
Philippines	2010	Male	20 to 24	72% (61-80%)	28% (20-39%)	2.7 (1.9-3.7)	9.2% (6.5-12%)	12% (8.5-17%)	1000 (690-1400)	78000 (55000-100000)
Philippines	2010	Male	25 to 29	76% (66-83%)	24% (17-34%)	3 (2.2-4)	10% (7.3-13%)	13% (9.3-18%)	1200 (840-1600)	87000 (62000-110000)
Philippines	2010	Male	30 to 34	78% (69-84%)	22% (16-31%)	3.4 (2.6-4.4)	11% (7.9-14%)	14% (10-19%)	1500 (1000-2000)	95000 (68000-120000)
Philippines	2010	Male	35 to 39	78% (69-84%)	22% (16-31%)	3.8 (2.7-5)	12% (8.6-16%)	16% (11-21%)	2000 (1300-2600)	110000 (77000-150000)
Philippines	2010	Male	40 to 44	77% (68-83%)	23% (17-32%)	3.9 (2.8-5.1)	13% (9.3-17%)	17% (12-22%)	2600 (1800-3400)	130000 (92000-170000)
Philippines	2010	Male	45 to 49	74% (62-81%)	26% (19-38%)	3.8 (2.7-5.2)	13% (8.9-18%)	17% (11-23%)	3200 (2100-4400)	140000 (96000-190000)
Philippines	2010	Male	50 to 54	69% (59-77%)	31% (23-41%)	3.5 (2.4-4.9)	13% (8.2-18%)	16% (9.9-21%)	3700 (2300-5000)	140000 (91000-190000)
Philippines	2010	Male	55 to 59	64% (52-74%)	36% (26-48%)	3.2 (2.2-4.4)	12% (7.5-16%)	14% (8.9-20%)	3900 (2400-5500)	130000 (82000-180000)
Philippines	2010	Male	60 to 64	59% (46-70%)	41% (30-54%)	2.8 (1.8-4.1)	9.7% (5.6-15%)	11% (6.6-17%)	3400 (2000-5200)	96000 (56000-140000)
Philippines	2010	Male	65 to 69	55% (41-67%)	45% (33-59%)	2.1 (1.4-3.1)	7.2% (4.3-11%)	8.3% (4.8-13%)	2400 (1400-3700)	56000 (34000-85000)
Philippines	2010	Male	70 to 74	49% (36-62%)	51% (38-64%)	1.6 (1-2.5)	5% (2.5-8.5%)	5.7% (2.8-9.7%)	1700 (850-2900)	32000 (16000-54000)
Philippines	2010	Male	75 to 79	43% (30-58%)	57% (42-70%)	1.3 (0.76-2.1)	3.3% (1.2-6.3%)	3.7% (1.3-7.1%)	990 (340-1900)	15000 (5000-28000)
Philippines	2010	Male	80 to 84	39% (26-53%)	61% (47-74%)	1.1 (0.64-1.7)	2.3% (0.64-4.5%)	2.4% (0.61-5%)	470 (120-960)	5200 (1500-11000)
Philippines	2010	Male	85 to 89	38% (25-52%)	62% (48-75%)	0.95 (0.54-1.5)	1.4% (0.068-3.4%)	1.5% (-0.045-3.6%)	170 (-5.2-420)	1400 (66-3200)
Philippines	2010	Male	90 to 94	39% (27-54%)	61% (46-73%)	0.92 (0.51-1.5)	0.92% (-0.4-2.8%)	0.91% (-0.56-3%)	43 (-26-140)	250 (-110-730)
Philippines	2010	Male	95 plus	43% (26-62%)	57% (38-74%)	0.92 (0.44-1.7)	0.87% (-0.93-3.2%)	0.76% (-0.93-3.3%)	9.6 (-12-41)	41 (-31-150)
Philippines	2016	Female	All Ages	16% (13-19%)	84% (81-87%)	0.41 (0.3-0.58)	1.4% (0.78-2.2%)	2.1% (0.81-3.8%)	5400 (2000-10000)	180000 (98000-300000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	2016	Female	15 to 19	15% (9.8-21%)	85% (79-90%)	0.36 (0.19-0.61)	0.88% (0.52-1.4%)	1.2% (0.56-2.3%)	38 (16-73)	5200 (3000-8400)
Philippines	2016	Female	20 to 24	18% (12-26%)	82% (74-88%)	0.49 (0.25-0.81)	1.5% (0.9-2.3%)	1.7% (0.7-3.1%)	71 (27-140)	10000 (5600-16000)
Philippines	2016	Female	25 to 29	20% (13-28%)	80% (72-87%)	0.55 (0.29-0.92)	1.7% (1-2.6%)	2.2% (0.9-3.9%)	99 (40-180)	11000 (6000-17000)
Philippines	2016	Female	30 to 34	21% (14-30%)	79% (70-86%)	0.61 (0.32-1)	1.7% (0.98-2.7%)	2.3% (1.1-4.1%)	120 (52-220)	11000 (6000-18000)
Philippines	2016	Female	35 to 39	22% (14-30%)	78% (70-86%)	0.67 (0.36-1.1)	1.9% (1.1-3.2%)	3% (1.4-5.2%)	190 (86-340)	13000 (7100-22000)
Philippines	2016	Female	40 to 44	21% (14-29%)	79% (71-86%)	0.67 (0.38-1.1)	2% (1.1-3.5%)	3.3% (1.7-5.7%)	250 (120-460)	14000 (7400-24000)
Philippines	2016	Female	45 to 49	21% (13-28%)	79% (72-87%)	0.64 (0.33-1.1)	2% (0.96-3.6%)	3.2% (1.5-5.8%)	330 (150-620)	16000 (7300-29000)
Philippines	2016	Female	50 to 54	21% (14-29%)	79% (71-86%)	0.72 (0.39-1.2)	2.3% (0.96-4.2%)	3.5% (1.5-6.5%)	450 (190-850)	18000 (7700-34000)
Philippines	2016	Female	55 to 59	21% (14-29%)	79% (71-86%)	0.89 (0.49-1.5)	2.9% (1.2-5.4%)	4.3% (1.9-7.7%)	700 (300-1300)	24000 (10000-46000)
Philippines	2016	Female	60 to 64	20% (13-28%)	80% (72-87%)	0.85 (0.47-1.5)	2.7% (0.99-5.4%)	3.8% (1.5-7.4%)	710 (270-1400)	21000 (7500-40000)
Philippines	2016	Female	65 to 69	18% (12-25%)	82% (75-88%)	0.7 (0.36-1.1)	2.2% (0.72-4.2%)	2.9% (1-5.6%)	650 (220-1300)	15000 (5000-31000)
Philippines	2016	Female	70 to 74	17% (11-24%)	83% (76-89%)	0.6 (0.3-1.1)	1.7% (0.27-3.9%)	2.2% (0.44-5%)	510 (99-1200)	9700 (1600-24000)
Philippines	2016	Female	75 to 79	17% (11-26%)	83% (74-89%)	0.58 (0.26-1.1)	1.3% (-0.081-3.5%)	1.6% (-0.078-4.3%)	450 (-20-1200)	6500 (-360-18000)
Philippines	2016	Female	80 to 84	18% (12-26%)	82% (74-88%)	0.6 (0.3-1)	1.1% (-0.1-3.1%)	1.3% (-0.12-3.7%)	400 (-37-1100)	4300 (-400-12000)
Philippines	2016	Female	85 to 89	20% (13-29%)	80% (71-87%)	0.66 (0.34-1.1)	1% (-0.26-2.9%)	1.1% (-0.33-3.3%)	270 (-78-790)	2000 (-510-5800)
Philippines	2016	Female	90 to 94	23% (15-32%)	77% (68-85%)	0.74 (0.38-1.3)	0.97% (-0.44-3.2%)	1.1% (-0.56-3.6%)	100 (-56-350)	550 (-250-1800)
Philippines	2016	Female	95 plus	27% (15-40%)	73% (60-85%)	0.86 (0.4-1.6)	1.2% (-0.54-4%)	1.3% (-0.72-4.4%)	45 (-26-160)	170 (-75-560)
Philippines	2016	Male	All Ages	57% (51-63%)	43% (37-49%)	1.9 (1.4-2.5)	6.7% (4.6-9.1%)	7.9% (5-11%)	29000 (18000-43000)	120000 (78000-160000)
Philippines	2016	Male	15 to 19	51% (40-63%)	49% (37-60%)	1.2 (0.64-1.8)	3.3% (2-5.1%)	5% (2.7-7.8%)	300 (160-500)	26000 (15000-42000)
Philippines	2016	Male	20 to 24	72% (61-80%)	28% (20-39%)	2.6 (1.6-3.8)	8.6% (5.7-12%)	12% (7.3-17%)	1000 (630-1600)	81000 (51000-120000)
Philippines	2016	Male	25 to 29	76% (67-83%)	24% (17-33%)	2.9 (1.9-4.2)	9.6% (6.5-13%)	13% (8.1-18%)	1300 (780-1800)	90000 (59000-130000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Philippines	2016	Male	30 to 34	78% (69-84%)	22% (16-31%)	3.3 (2.2-4.6)	10% (6.8-14%)	14% (8.6-19%)	1400 (860-2100)	92000 (59000-130000)
Philippines	2016	Male	35 to 39	78% (69-84%)	22% (16-31%)	3.6 (2.5-5.2)	11% (7.4-16%)	15% (9.4-21%)	1900 (1100-2900)	110000 (69000-160000)
Philippines	2016	Male	40 to 44	77% (68-83%)	23% (17-32%)	3.7 (2.5-5.3)	12% (7.9-17%)	16% (10-22%)	2400 (1500-3600)	120000 (77000-180000)
Philippines	2016	Male	45 to 49	74% (63-81%)	26% (19-37%)	3.7 (2.4-5.3)	12% (7.4-18%)	15% (9.3-22%)	3200 (1900-4700)	140000 (84000-210000)
Philippines	2016	Male	50 to 54	70% (60-77%)	30% (23-40%)	3.4 (2.2-5.1)	12% (6.8-17%)	14% (8.3-21%)	3700 (2100-5600)	140000 (82000-210000)
Philippines	2016	Male	55 to 59	64% (52-75%)	36% (25-48%)	3.1 (2-4.7)	11% (6.2-16%)	13% (7.3-20%)	4100 (2300-6500)	140000 (77000-210000)
Philippines	2016	Male	60 to 64	59% (47-71%)	41% (29-53%)	2.7 (1.6-4.1)	8.9% (4.7-14%)	10% (5.6-17%)	3700 (1900-6200)	110000 (55000-170000)
Philippines	2016	Male	65 to 69	56% (42-67%)	44% (33-58%)	2.1 (1.2-3.1)	6.5% (3.2-11%)	7.5% (3.6-12%)	2800 (1300-4600)	65000 (31000-110000)
Philippines	2016	Male	70 to 74	50% (37-62%)	50% (38-63%)	1.6 (0.9-2.5)	4.5% (1.8-8.2%)	5.1% (2-9.4%)	1600 (570-3000)	31000 (11000-57000)
Philippines	2016	Male	75 to 79	43% (30-58%)	57% (42-70%)	1.3 (0.68-2.2)	2.9% (0.65-6.3%)	3.2% (0.65-7%)	980 (190-2200)	15000 (3300-32000)
Philippines	2016	Male	80 to 84	39% (26-53%)	61% (47-74%)	1 (0.56-1.7)	2% (0.33-4.4%)	2.1% (0.24-4.9%)	540 (54-1300)	6000 (1000-14000)
Philippines	2016	Male	85 to 89	38% (25-52%)	62% (48-75%)	0.91 (0.48-1.5)	1.2% (-0.17-3.2%)	1.3% (-0.28-3.5%)	210 (-46-590)	1700 (-230-4500)
Philippines	2016	Male	90 to 94	40% (27-53%)	60% (47-73%)	0.89 (0.49-1.5)	0.78% (-0.63-2.8%)	0.74% (-0.85-3.1%)	37 (-42-150)	220 (-180-770)
Philippines	2016	Male	95 plus	44% (26-62%)	56% (38-74%)	0.89 (0.42-1.7)	0.74% (-0.72-3%)	0.61% (-1-3%)	8.1 (-14-40)	37 (-36-150)
Sri Lanka	1990	Female	All Ages	5% (4.2-6%)	95% (94-96%)	0.11 (0.08-0.13)	0.51% (0.38-0.67%)	0.56% (0.27-0.91%)	210 (100-340)	11000 (7800-14000)
Sri Lanka	1990	Female	15 to 19	4.5% (2.9-6.7%)	95% (93-97%)	0.088 (0.046-0.15)	0.5% (0.29-0.8%)	0.47% (0.13-1%)	3.9 (1-8.6)	630 (350-1000)
Sri Lanka	1990	Female	20 to 24	5.7% (3.4-8.4%)	94% (92-97%)	0.12 (0.06-0.21)	0.9% (0.55-1.3%)	0.65% (0.17-1.4%)	6.3 (1.6-14)	1200 (710-1800)
Sri Lanka	1990	Female	25 to 29	6.2% (3.7-9.4%)	94% (91-96%)	0.14 (0.075-0.24)	0.98% (0.63-1.4%)	0.76% (0.32-1.5%)	6.6 (2.7-13)	1200 (740-1700)
Sri Lanka	1990	Female	30 to 34	6.6% (3.9-10%)	93% (90-96%)	0.15 (0.083-0.26)	0.99% (0.69-1.3%)	0.97% (0.56-1.6%)	8.5 (4.9-14)	1200 (770-1600)
Sri Lanka	1990	Female	35 to 39	6.9% (3.9-10%)	93% (90-96%)	0.16 (0.079-0.27)	0.93% (0.66-1.3%)	1.1% (0.69-1.7%)	9.1 (5.7-14)	950 (660-1300)
Sri Lanka	1990	Female	40 to 44	6.8% (4.1-10%)	93% (90-96%)	0.16 (0.085-0.29)	0.91% (0.64-1.2%)	1.3% (0.86-1.9%)	12 (7.5-17)	900 (630-1200)
Sri Lanka	1990	Female	45 to 49	6.7% (3.8-9.9%)	93% (90-96%)	0.18 (0.091-0.32)	0.95% (0.65-1.4%)	1.4% (0.9-2.2%)	15 (9.1-23)	890 (610-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	1990	Female	50 to 54	6.8% (4.2-10%)	93% (90-96%)	0.21 (0.11-0.36)	0.93% (0.62-1.4%)	1.4% (0.84-2.2%)	18 (11-28)	860 (570-1300)
Sri Lanka	1990	Female	55 to 59	6.9% (4.3-10%)	93% (90-96%)	0.23 (0.12-0.4)	0.87% (0.51-1.4%)	1.3% (0.7-2.1%)	20 (11-34)	790 (460-1300)
Sri Lanka	1990	Female	60 to 64	6.7% (4-10%)	93% (90-96%)	0.23 (0.12-0.4)	0.75% (0.36-1.3%)	1.1% (0.47-1.8%)	22 (10-39)	710 (350-1200)
Sri Lanka	1990	Female	65 to 69	6.1% (3.7-9%)	94% (91-96%)	0.21 (0.11-0.35)	0.6% (0.25-1.1%)	0.81% (0.32-1.4%)	23 (9-41)	590 (250-1000)
Sri Lanka	1990	Female	70 to 74	5.9% (3.8-8.9%)	94% (91-96%)	0.2 (0.1-0.34)	0.52% (0.16-1%)	0.67% (0.19-1.3%)	23 (6.2-45)	460 (140-900)
Sri Lanka	1990	Female	75 to 79	6.3% (3.8-9.5%)	94% (90-96%)	0.2 (0.097-0.37)	0.38% (0.0096-0.86%)	0.46% (-0.017-1.1%)	17 (-0.64-41)	270 (6.8-620)
Sri Lanka	1990	Female	80 to 84	6.7% (4-9.6%)	93% (90-96%)	0.21 (0.11-0.37)	0.31% (-0.04-0.79%)	0.35% (-0.074-0.92%)	14 (-3-37)	160 (-20-410)
Sri Lanka	1990	Female	85 to 89	7.5% (4.4-12%)	93% (88-96%)	0.23 (0.11-0.4)	0.24% (-0.11-0.71%)	0.25% (-0.15-0.8%)	9.3 (-5.3-30)	76 (-34-240)
Sri Lanka	1990	Female	90 to 94	8.6% (5.1-13%)	91% (87-95%)	0.25 (0.12-0.46)	0.17% (-0.2-0.71%)	0.16% (-0.27-0.79%)	2.7 (-4.6-13)	17 (-21-72)
Sri Lanka	1990	Female	95 plus	10% (5.2-17%)	90% (83-95%)	0.28 (0.12-0.57)	0.15% (-0.32-0.89%)	0.11% (-0.42-0.93%)	0.56 (-2.2-4.7)	3 (-6.2-18)
Sri Lanka	1990	Male	All Ages	33% (28-39%)	67% (61-72%)	0.48 (0.38-0.58)	2% (1.3-2.8%)	2% (1.1-2.9%)	1300 (660-1900)	63000 (38000-87000)
Sri Lanka	1990	Male	15 to 19	26% (16-37%)	74% (63-84%)	0.19 (0.095-0.33)	1% (0.39-2%)	1.1% (0.15-2.5%)	19 (2.5-43)	1900 (710-3700)
Sri Lanka	1990	Male	20 to 24	39% (25-53%)	61% (47-75%)	0.46 (0.27-0.73)	2.5% (0.99-4.6%)	2.6% (0.62-5.3%)	79 (19-160)	6500 (2500-12000)
Sri Lanka	1990	Male	25 to 29	46% (31-60%)	54% (40-69%)	0.66 (0.39-0.98)	3.4% (1.6-5.4%)	3.8% (1.4-6.5%)	100 (38-180)	7800 (3700-12000)
Sri Lanka	1990	Male	30 to 34	51% (34-66%)	49% (34-66%)	0.85 (0.52-1.2)	4.1% (2.3-6.2%)	4.7% (2.3-7.8%)	120 (59-200)	8300 (4600-12000)
Sri Lanka	1990	Male	35 to 39	54% (39-68%)	46% (32-61%)	1 (0.64-1.5)	4.7% (3-6.9%)	5.8% (3.4-8.5%)	130 (78-200)	8000 (5100-12000)
Sri Lanka	1990	Male	40 to 44	55% (41-70%)	45% (30-59%)	1.1 (0.69-1.6)	4.8% (3-6.6%)	5.9% (3.5-8.6%)	140 (83-210)	7700 (4800-11000)
Sri Lanka	1990	Male	45 to 49	54% (37-68%)	46% (32-63%)	1.1 (0.64-1.6)	4.5% (2.7-6.5%)	5.6% (3.2-8.1%)	140 (82-220)	6700 (4100-9700)
Sri Lanka	1990	Male	50 to 54	52% (37-67%)	48% (33-63%)	1 (0.58-1.5)	3.6% (2-5.4%)	4.4% (2.2-6.8%)	130 (63-200)	5300 (2900-8000)
Sri Lanka	1990	Male	55 to 59	50% (35-65%)	50% (35-65%)	0.87 (0.5-1.4)	2.6% (1.2-4.1%)	3.2% (1.3-5.2%)	110 (43-180)	3900 (1800-6200)
Sri Lanka	1990	Male	60 to 64	46% (31-63%)	54% (37-69%)	0.74 (0.4-1.2)	1.9% (0.68-3.1%)	2.2% (0.52-3.8%)	90 (21-160)	2800 (1000-4700)
Sri Lanka	1990	Male	65 to 69	41% (27-55%)	59% (45-73%)	0.6 (0.33-0.95)	1.4% (0.42-2.3%)	1.5% (0.34-2.7%)	76 (17-130)	2000 (610-3300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	1990	Male	70 to 74	36% (24-50%)	64% (50-76%)	0.49 (0.26-0.8)	0.98% (0.21-1.9%)	1.1% (0.12-2.2%)	59 (6.5-120)	1200 (260-2400)
Sri Lanka	1990	Male	75 to 79	33% (20-49%)	67% (51-80%)	0.43 (0.23-0.75)	0.51% (-0.17-1.2%)	0.49% (-0.32-1.4%)	27 (-17-75)	480 (-150-1200)
Sri Lanka	1990	Male	80 to 84	29% (17-44%)	71% (56-83%)	0.38 (0.2-0.63)	0.32% (-0.26-0.95%)	0.29% (-0.41-1%)	16 (-22-54)	220 (-180-630)
Sri Lanka	1990	Male	85 to 89	27% (16-40%)	73% (60-84%)	0.32 (0.17-0.57)	0.14% (-0.42-0.62%)	0.086% (-0.57-0.64%)	3.6 (-24-27)	50 (-150-220)
Sri Lanka	1990	Male	90 to 94	27% (17-39%)	73% (61-83%)	0.3 (0.15-0.54)	0.0016% (-0.51-0.46%)	-0.078% (-0.7-0.46%)	-1.2 (-11-7.4)	0.16 (-48-44)
Sri Lanka	1990	Male	95 plus	28% (16-44%)	72% (56-84%)	0.3 (0.13-0.58)	-0.066% (-0.68-0.42%)	-0.2% (-0.93-0.36%)	-0.73 (-3.3-1.3)	-0.9 (-9.4-5.8)
Sri Lanka	1995	Female	All Ages	5.4% (4.6-6.4%)	95% (94-95%)	0.11 (0.085-0.15)	0.59% (0.43-0.78%)	0.6% (0.29-0.99%)	250 (120-410)	13000 (8900-17000)
Sri Lanka	1995	Female	15 to 19	4.8% (3-7.2%)	95% (93-97%)	0.091 (0.046-0.16)	0.52% (0.29-0.86%)	0.51% (0.13-1.1%)	4.9 (1.3-11)	730 (400-1200)
Sri Lanka	1995	Female	20 to 24	6% (3.6-9%)	94% (91-96%)	0.12 (0.062-0.22)	0.92% (0.57-1.4%)	0.7% (0.19-1.5%)	7 (1.9-15)	1300 (730-2000)
Sri Lanka	1995	Female	25 to 29	6.5% (3.9-10%)	93% (90-96%)	0.14 (0.076-0.25)	0.99% (0.68-1.4%)	0.79% (0.32-1.5%)	7.3 (2.9-14)	1300 (810-1800)
Sri Lanka	1995	Female	30 to 34	6.9% (4-11%)	93% (89-96%)	0.15 (0.082-0.27)	1% (0.69-1.4%)	1% (0.58-1.7%)	9.1 (5.2-15)	1200 (820-1700)
Sri Lanka	1995	Female	35 to 39	7.2% (4.2-11%)	93% (89-96%)	0.16 (0.081-0.27)	0.95% (0.7-1.3%)	1.1% (0.71-1.8%)	11 (6.9-18)	1200 (820-1600)
Sri Lanka	1995	Female	40 to 44	7.2% (4.4-10%)	93% (90-96%)	0.17 (0.086-0.28)	0.94% (0.67-1.3%)	1.4% (0.9-2%)	14 (9.2-20)	1100 (740-1400)
Sri Lanka	1995	Female	45 to 49	7% (4-10%)	93% (90-96%)	0.19 (0.092-0.33)	1% (0.72-1.5%)	1.5% (0.98-2.3%)	21 (14-31)	1200 (830-1700)
Sri Lanka	1995	Female	50 to 54	7.1% (4.5-11%)	93% (89-95%)	0.21 (0.11-0.37)	1% (0.66-1.5%)	1.5% (0.91-2.2%)	24 (15-37)	1100 (730-1600)
Sri Lanka	1995	Female	55 to 59	7.2% (4.5-11%)	93% (89-95%)	0.23 (0.12-0.42)	0.9% (0.51-1.5%)	1.3% (0.72-2.2%)	25 (14-41)	970 (560-1500)
Sri Lanka	1995	Female	60 to 64	6.9% (4.1-10%)	93% (90-96%)	0.23 (0.11-0.4)	0.75% (0.34-1.3%)	1% (0.46-1.8%)	27 (12-48)	860 (390-1500)
Sri Lanka	1995	Female	65 to 69	6.4% (3.8-9.4%)	94% (91-96%)	0.21 (0.11-0.35)	0.6% (0.25-1.1%)	0.79% (0.31-1.5%)	29 (11-54)	740 (310-1300)
Sri Lanka	1995	Female	70 to 74	6.2% (3.9-9.3%)	94% (91-96%)	0.2 (0.1-0.35)	0.5% (0.13-1%)	0.62% (0.13-1.3%)	29 (6.7-59)	580 (150-1200)
Sri Lanka	1995	Female	75 to 79	6.6% (4-9.9%)	93% (90-96%)	0.21 (0.1-0.37)	0.36% (-0.038-0.87%)	0.43% (-0.066-1.1%)	19 (-2.8-46)	300 (-30-710)
Sri Lanka	1995	Female	80 to 84	7.1% (4.3-10%)	93% (90-96%)	0.22 (0.11-0.38)	0.25% (-0.11-0.77%)	0.28% (-0.17-0.9%)	12 (-7.1-37)	140 (-64-420)
Sri Lanka	1995	Female	85 to 89	7.9% (4.6-12%)	92% (88-95%)	0.23 (0.12-0.4)	0.17% (-0.18-0.65%)	0.17% (-0.25-0.71%)	6.3 (-9.7-28)	58 (-64-220)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	1995	Female	90 to 94	9% (5.4-14%)	91% (86-95%)	0.25 (0.12-0.46)	0.1% (-0.27-0.58%)	0.071% (-0.39-0.65%)	1.4 (-7.4-12)	12 (-33-69)
Sri Lanka	1995	Female	95 plus	11% (5.8-18%)	89% (82-94%)	0.29 (0.12-0.58)	0.086% (-0.4-0.79%)	0.018% (-0.57-0.86%)	0.11 (-3.5-5.2)	2.1 (-9.8-20)
Sri Lanka	1995	Male	All Ages	35% (30-42%)	65% (58-70%)	0.5 (0.39-0.61)	2.6% (1.7-3.6%)	2.5% (1.4-3.6%)	1700 (990-2500)	83000 (54000-110000)
Sri Lanka	1995	Male	15 to 19	27% (17-39%)	73% (61-83%)	0.19 (0.096-0.33)	1% (0.4-2%)	1.1% (0.19-2.5%)	21 (3.7-46)	2000 (760-3800)
Sri Lanka	1995	Male	20 to 24	40% (27-55%)	60% (45-73%)	0.46 (0.27-0.74)	2.6% (1-4.8%)	2.8% (0.74-5.5%)	91 (25-180)	7300 (2800-13000)
Sri Lanka	1995	Male	25 to 29	47% (32-62%)	53% (38-68%)	0.65 (0.39-0.98)	3.7% (1.8-5.9%)	4% (1.6-6.9%)	130 (50-220)	9400 (4600-15000)
Sri Lanka	1995	Male	30 to 34	52% (35-67%)	48% (33-65%)	0.85 (0.53-1.3)	4.5% (2.7-6.8%)	5.3% (2.8-8.4%)	150 (79-240)	9900 (5800-15000)
Sri Lanka	1995	Male	35 to 39	56% (40-70%)	44% (30-60%)	1 (0.64-1.5)	5.5% (3.6-7.8%)	6.8% (4.2-9.8%)	200 (120-290)	12000 (7500-16000)
Sri Lanka	1995	Male	40 to 44	57% (42-70%)	43% (30-58%)	1.1 (0.68-1.6)	5.6% (3.7-7.6%)	7% (4.4-9.7%)	200 (130-290)	11000 (6900-14000)
Sri Lanka	1995	Male	45 to 49	55% (38-69%)	45% (31-62%)	1.1 (0.66-1.6)	5.4% (3.3-7.6%)	6.7% (3.8-9.5%)	230 (130-340)	11000 (6400-15000)
Sri Lanka	1995	Male	50 to 54	53% (38-68%)	47% (32-62%)	0.99 (0.58-1.5)	4.3% (2.4-6.2%)	5.2% (2.7-7.7%)	190 (98-290)	7700 (4100-11000)
Sri Lanka	1995	Male	55 to 59	51% (35-67%)	49% (33-65%)	0.87 (0.49-1.4)	3.1% (1.5-4.7%)	3.7% (1.7-5.7%)	150 (68-240)	5300 (2700-8100)
Sri Lanka	1995	Male	60 to 64	48% (32-63%)	52% (37-68%)	0.74 (0.4-1.2)	2.1% (0.83-3.4%)	2.5% (0.78-4.1%)	120 (39-210)	3600 (1400-6000)
Sri Lanka	1995	Male	65 to 69	42% (28-57%)	58% (43-72%)	0.6 (0.33-0.96)	1.5% (0.52-2.5%)	1.7% (0.48-3%)	100 (29-180)	2600 (900-4300)
Sri Lanka	1995	Male	70 to 74	37% (24-51%)	63% (49-76%)	0.49 (0.26-0.8)	1.1% (0.26-1.9%)	1.2% (0.22-2.2%)	79 (14-150)	1600 (390-3000)
Sri Lanka	1995	Male	75 to 79	34% (21-50%)	66% (50-79%)	0.44 (0.23-0.76)	0.52% (-0.11-1.3%)	0.51% (-0.26-1.4%)	29 (-14-77)	520 (-110-1200)
Sri Lanka	1995	Male	80 to 84	30% (18-45%)	70% (55-82%)	0.38 (0.2-0.63)	0.33% (-0.24-0.91%)	0.3% (-0.42-0.99%)	15 (-21-50)	220 (-160-590)
Sri Lanka	1995	Male	85 to 89	28% (17-41%)	72% (59-83%)	0.33 (0.17-0.56)	0.15% (-0.38-0.61%)	0.082% (-0.56-0.62%)	3.1 (-21-24)	51 (-130-210)
Sri Lanka	1995	Male	90 to 94	28% (17-41%)	72% (59-83%)	0.31 (0.15-0.56)	0.014% (-0.49-0.42%)	-0.089% (-0.71-0.4%)	-1.5 (-12-6.7)	1.4 (-50-44)
Sri Lanka	1995	Male	95 plus	30% (17-46%)	70% (54-83%)	0.31 (0.13-0.59)	-0.00028% (-0.58-0.43%)	-0.18% (-0.9-0.35%)	-0.72 (-3.7-1.5)	0.0039 (-9.6-7.1)
Sri Lanka	2000	Female	All Ages	5% (4.3-5.9%)	95% (94-96%)	0.14 (0.1-0.18)	0.74% (0.54-1%)	0.87% (0.45-1.4%)	380 (190-620)	16000 (11000-22000)
Sri Lanka	2000	Female	15 to 19	4.5% (2.9-6.7%)	95% (93-97%)	0.11 (0.057-0.2)	0.6% (0.35-1%)	0.65% (0.2-1.4%)	5.6 (1.7-12)	790 (440-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	2000	Female	20 to 24	5.6% (3.5-8.1%)	94% (92-97%)	0.15 (0.073-0.25)	1% (0.66-1.6%)	0.86% (0.28-1.8%)	7.6 (2.5-16)	1300 (800-2100)
Sri Lanka	2000	Female	25 to 29	6.3% (3.9-9.3%)	94% (91-96%)	0.18 (0.092-0.3)	1.1% (0.76-1.6%)	0.99% (0.42-1.9%)	7.3 (3.1-14)	1200 (800-1800)
Sri Lanka	2000	Female	30 to 34	6.4% (4-9.7%)	94% (90-96%)	0.18 (0.097-0.33)	1.1% (0.76-1.6%)	1.2% (0.7-2%)	9.4 (5.3-17)	1300 (840-1800)
Sri Lanka	2000	Female	35 to 39	6.2% (3.7-9%)	94% (91-96%)	0.17 (0.091-0.3)	1% (0.73-1.4%)	1.3% (0.82-2.1%)	12 (7.3-19)	1200 (850-1700)
Sri Lanka	2000	Female	40 to 44	6.1% (3.9-8.7%)	94% (91-96%)	0.18 (0.096-0.3)	1% (0.72-1.4%)	1.5% (1-2.2%)	16 (11-23)	1300 (900-1700)
Sri Lanka	2000	Female	45 to 49	6% (3.6-8.6%)	94% (91-96%)	0.2 (0.1-0.35)	1.1% (0.75-1.6%)	1.7% (1.1-2.5%)	24 (16-35)	1400 (940-1900)
Sri Lanka	2000	Female	50 to 54	5.8% (3.8-8.5%)	94% (91-96%)	0.22 (0.11-0.37)	1.1% (0.72-1.6%)	1.6% (1-2.5%)	32 (20-50)	1500 (1000-2200)
Sri Lanka	2000	Female	55 to 59	6.2% (3.9-8.9%)	94% (91-96%)	0.25 (0.13-0.44)	1.1% (0.63-1.7%)	1.6% (0.92-2.6%)	34 (19-55)	1300 (790-2200)
Sri Lanka	2000	Female	60 to 64	6.2% (3.9-9.3%)	94% (91-96%)	0.27 (0.13-0.47)	0.96% (0.46-1.7%)	1.3% (0.65-2.3%)	38 (18-67)	1200 (590-2100)
Sri Lanka	2000	Female	65 to 69	5.9% (3.7-8.5%)	94% (92-96%)	0.25 (0.13-0.41)	0.84% (0.39-1.5%)	1.1% (0.51-2%)	44 (21-80)	1100 (530-2000)
Sri Lanka	2000	Female	70 to 74	6% (3.9-8.7%)	94% (91-96%)	0.25 (0.12-0.44)	0.75% (0.26-1.4%)	0.94% (0.31-1.9%)	50 (16-99)	980 (320-1900)
Sri Lanka	2000	Female	75 to 79	6.4% (3.9-9.7%)	94% (90-96%)	0.26 (0.12-0.47)	0.59% (0.089-1.3%)	0.72% (0.074-1.6%)	36 (3.8-82)	570 (80-1300)
Sri Lanka	2000	Female	80 to 84	6.8% (4.2-9.9%)	93% (90-96%)	0.27 (0.13-0.46)	0.5% (0.025-1.2%)	0.58% (0.0084-1.4%)	30 (0.45-74)	340 (17-820)
Sri Lanka	2000	Female	85 to 89	7.7% (4.4-12%)	92% (88-96%)	0.29 (0.14-0.5)	0.42% (-0.0014-1.1%)	0.48% (-0.032-1.3%)	18 (-1.2-47)	150 (-0.52-390)
Sri Lanka	2000	Female	90 to 94	8.8% (5.3-14%)	91% (86-95%)	0.32 (0.16-0.58)	0.36% (-0.099-1.1%)	0.39% (-0.17-1.3%)	8.4 (-3.9-28)	50 (-14-150)
Sri Lanka	2000	Female	95 plus	11% (5.6-18%)	89% (82-94%)	0.36 (0.15-0.73)	0.39% (-0.25-1.5%)	0.37% (-0.37-1.7%)	2.8 (-2.8-13)	12 (-7.6-47)
Sri Lanka	2000	Male	All Ages	38% (33-43%)	62% (57-67%)	0.72 (0.55-0.91)	3.5% (2.4-4.8%)	3.4% (2.1-4.8%)	2500 (1600-3600)	110000 (76000-150000)
Sri Lanka	2000	Male	15 to 19	27% (19-38%)	73% (62-81%)	0.25 (0.13-0.43)	1.3% (0.55-2.3%)	1.5% (0.36-3.2%)	23 (5.2-47)	2200 (930-4000)
Sri Lanka	2000	Male	20 to 24	40% (29-52%)	60% (48-71%)	0.6 (0.34-0.95)	3.3% (1.6-5.7%)	3.7% (1.3-6.8%)	100 (37-190)	8200 (3900-14000)
Sri Lanka	2000	Male	25 to 29	48% (36-60%)	52% (40-64%)	0.85 (0.52-1.3)	4.5% (2.5-7.2%)	5.3% (2.4-8.8%)	130 (59-220)	9500 (5200-15000)
Sri Lanka	2000	Male	30 to 34	54% (40-66%)	46% (34-60%)	1.1 (0.72-1.7)	5.6% (3.4-8.2%)	6.9% (3.8-11%)	170 (97-270)	11000 (6700-16000)
Sri Lanka	2000	Male	35 to 39	59% (47-71%)	41% (29-53%)	1.4 (0.89-2.1)	6.9% (4.6-9.7%)	8.7% (5.5-12%)	260 (170-370)	15000 (9900-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	2000	Male	40 to 44	60% (48-72%)	40% (28-52%)	1.5 (0.96-2.2)	6.9% (4.8-9.3%)	8.8% (5.8-12%)	310 (200-430)	16000 (11000-21000)
Sri Lanka	2000	Male	45 to 49	58% (44-70%)	42% (30-56%)	1.5 (0.95-2.2)	6.6% (4.2-9.1%)	8.2% (5.1-11%)	350 (210-490)	15000 (9700-21000)
Sri Lanka	2000	Male	50 to 54	57% (44-69%)	43% (31-56%)	1.4 (0.84-2.1)	5.4% (3.2-7.8%)	6.5% (3.7-9.5%)	340 (190-510)	13000 (8000-20000)
Sri Lanka	2000	Male	55 to 59	55% (42-68%)	45% (32-58%)	1.2 (0.75-2)	3.9% (2.1-6.2%)	4.8% (2.4-7.5%)	250 (120-390)	8500 (4600-13000)
Sri Lanka	2000	Male	60 to 64	50% (37-64%)	50% (36-63%)	1 (0.59-1.7)	2.8% (1.2-4.7%)	3.3% (1.3-5.7%)	190 (74-320)	5500 (2400-9400)
Sri Lanka	2000	Male	65 to 69	43% (30-56%)	57% (44-70%)	0.8 (0.46-1.2)	2% (0.87-3.4%)	2.3% (0.93-3.9%)	150 (60-260)	3800 (1600-6200)
Sri Lanka	2000	Male	70 to 74	37% (25-50%)	63% (50-75%)	0.65 (0.35-1)	1.5% (0.46-2.8%)	1.6% (0.44-3.2%)	130 (34-240)	2500 (790-4700)
Sri Lanka	2000	Male	75 to 79	36% (23-50%)	64% (50-77%)	0.6 (0.32-1)	0.8% (-0.022-1.9%)	0.84% (-0.17-2.1%)	54 (-11-140)	910 (-24-2100)
Sri Lanka	2000	Male	80 to 84	31% (18-46%)	69% (54-82%)	0.5 (0.26-0.85)	0.57% (-0.14-1.5%)	0.58% (-0.28-1.6%)	33 (-16-93)	420 (-100-1100)
Sri Lanka	2000	Male	85 to 89	26% (16-39%)	74% (61-84%)	0.41 (0.21-0.7)	0.39% (-0.11-0.99%)	0.37% (-0.26-1.1%)	14 (-9.4-41)	140 (-41-350)
Sri Lanka	2000	Male	90 to 94	26% (16-39%)	74% (61-84%)	0.38 (0.19-0.68)	0.22% (-0.29-0.74%)	0.14% (-0.5-0.8%)	2.7 (-8.9-15)	26 (-34-86)
Sri Lanka	2000	Male	95 plus	28% (15-43%)	72% (57-85%)	0.37 (0.16-0.73)	0.21% (-0.37-0.77%)	0.044% (-0.67-0.67%)	0.24 (-3.6-3.6)	4.6 (-7.9-17)
Sri Lanka	2005	Female	All Ages	4.8% (4.1-5.6%)	95% (94-96%)	0.21 (0.15-0.28)	1.5% (0.71-1.4%)	1.5% (0.81-2.3%)	680 (370-1100)	22000 (16000-32000)
Sri Lanka	2005	Female	15 to 19	4.2% (2.7-6.3%)	96% (94-97%)	0.16 (0.081-0.28)	0.75% (0.42-1.2%)	0.98% (0.37-2.1%)	6.2 (2.3-13)	810 (450-1400)
Sri Lanka	2005	Female	20 to 24	5.2% (3.1-7.6%)	95% (92-97%)	0.21 (0.1-0.38)	1.3% (0.8-2%)	1.3% (0.5-2.6%)	9.3 (3.5-19)	1500 (910-2400)
Sri Lanka	2005	Female	25 to 29	5.8% (3.7-8.6%)	94% (91-96%)	0.26 (0.13-0.45)	1.4% (0.88-2%)	1.5% (0.68-2.8%)	9.8 (4.4-19)	1500 (960-2300)
Sri Lanka	2005	Female	30 to 34	6% (3.7-9.2%)	94% (91-96%)	0.27 (0.14-0.47)	1.3% (0.91-1.9%)	1.7% (0.91-3%)	11 (5.8-19)	1400 (880-2000)
Sri Lanka	2005	Female	35 to 39	5.8% (3.5-8.4%)	94% (92-97%)	0.25 (0.13-0.44)	1.2% (0.82-1.7%)	1.7% (1-2.8%)	13 (7.6-22)	1400 (910-2000)
Sri Lanka	2005	Female	40 to 44	5.7% (3.6-8.2%)	94% (92-96%)	0.26 (0.14-0.46)	1.2% (0.82-1.7%)	1.9% (1.2-2.9%)	18 (11-28)	1400 (980-2000)
Sri Lanka	2005	Female	45 to 49	5.6% (3.3-8.1%)	94% (92-97%)	0.3 (0.15-0.53)	1.3% (0.83-1.9%)	2.1% (1.3-3.3%)	29 (18-45)	1700 (1100-2600)
Sri Lanka	2005	Female	50 to 54	5.5% (3.5-8%)	95% (92-97%)	0.32 (0.17-0.55)	1.4% (0.87-2.2%)	2.2% (1.3-3.5%)	40 (24-65)	1900 (1200-3000)
Sri Lanka	2005	Female	55 to 59	5.8% (3.7-8.3%)	94% (92-96%)	0.37 (0.19-0.64)	1.4% (0.82-2.4%)	2.3% (1.3-3.8%)	53 (30-88)	2100 (1200-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	2005	Female	60 to 64	5.8% (3.7-8.7%)	94% (91-96%)	0.39 (0.19-0.66)	1.4% (0.71-2.4%)	2.1% (1-3.6%)	61 (29-100)	2000 (970-3300)
Sri Lanka	2005	Female	65 to 69	5.5% (3.5-7.7%)	94% (92-96%)	0.36 (0.19-0.59)	1.4% (0.67-2.3%)	1.9% (0.93-3.1%)	74 (37-120)	1900 (950-3100)
Sri Lanka	2005	Female	70 to 74	5.6% (3.5-8.3%)	94% (92-97%)	0.36 (0.18-0.64)	1.3% (0.58-2.5%)	1.7% (0.73-3.2%)	95 (41-180)	1900 (810-3600)
Sri Lanka	2005	Female	75 to 79	6.1% (3.7-9.3%)	94% (91-96%)	0.38 (0.18-0.68)	1.2% (0.41-2.3%)	1.5% (0.49-2.8%)	83 (28-160)	1300 (440-2600)
Sri Lanka	2005	Female	80 to 84	6.6% (3.9-9.5%)	93% (91-96%)	0.4 (0.2-0.69)	1.1% (0.37-2.3%)	1.4% (0.43-2.8%)	82 (26-170)	920 (300-1800)
Sri Lanka	2005	Female	85 to 89	7.3% (4.2-11%)	93% (89-96%)	0.43 (0.21-0.77)	1% (0.3-2.1%)	1.2% (0.33-2.6%)	52 (14-110)	420 (130-880)
Sri Lanka	2005	Female	90 to 94	8.4% (5-13%)	92% (87-95%)	0.47 (0.23-0.89)	0.97% (0.21-2.3%)	1.1% (0.22-2.8%)	28 (5.4-70)	160 (36-380)
Sri Lanka	2005	Female	95 plus	10% (5.1-16%)	90% (84-95%)	0.54 (0.21-1.1)	1.1% (0.096-2.7%)	1.2% (0.039-3.2%)	12 (0.36-31)	43 (4-110)
Sri Lanka	2005	Male	All Ages	37% (32-43%)	63% (57-68%)	1.1 (0.86-1.4)	4.6% (3.1-6.2%)	4.8% (3.1-6.9%)	3500 (2300-5000)	140000 (100000-190000)
Sri Lanka	2005	Male	15 to 19	25% (17-36%)	75% (64-83%)	0.37 (0.19-0.64)	1.7% (0.92-2.9%)	2.3% (0.92-4.3%)	22 (8.6-43)	2200 (1100-3700)
Sri Lanka	2005	Male	20 to 24	39% (27-51%)	61% (49-73%)	0.9 (0.51-1.4)	4.5% (2.5-7.3%)	5.7% (2.7-9.7%)	94 (43-160)	7700 (4100-12000)
Sri Lanka	2005	Male	25 to 29	46% (34-58%)	54% (42-66%)	1.3 (0.76-2)	6% (3.4-8.9%)	7.8% (4-12%)	130 (66-210)	9700 (5600-14000)
Sri Lanka	2005	Male	30 to 34	53% (39-66%)	47% (34-61%)	1.7 (1.1-2.5)	7.2% (4.6-10%)	9.8% (5.9-15%)	160 (95-250)	11000 (6900-16000)
Sri Lanka	2005	Male	35 to 39	58% (46-69%)	42% (31-54%)	2.2 (1.4-3.2)	8.7% (5.8-12%)	12% (7.5-17%)	270 (170-380)	16000 (10000-22000)
Sri Lanka	2005	Male	40 to 44	58% (47-71%)	42% (29-53%)	2.3 (1.5-3.3)	9.1% (6.3-12%)	12% (8.1-16%)	370 (240-510)	19000 (13000-26000)
Sri Lanka	2005	Male	45 to 49	57% (43-69%)	43% (31-57%)	2.3 (1.5-3.3)	8.8% (5.8-12%)	11% (7.4-16%)	460 (290-670)	21000 (13000-30000)
Sri Lanka	2005	Male	50 to 54	56% (43-68%)	44% (32-57%)	2.1 (1.3-3.1)	7.5% (4.6-11%)	9.4% (5.6-14%)	470 (280-720)	19000 (11000-28000)
Sri Lanka	2005	Male	55 to 59	54% (41-66%)	46% (34-59%)	1.9 (1.1-3)	5.9% (3.3-9.2%)	7.2% (4.1-12%)	420 (230-670)	15000 (8200-23000)
Sri Lanka	2005	Male	60 to 64	49% (36-63%)	51% (37-64%)	1.6 (0.89-2.5)	4.4% (2.1-7.3%)	5.3% (2.4-8.8%)	320 (140-550)	9400 (4400-16000)
Sri Lanka	2005	Male	65 to 69	41% (29-55%)	59% (45-71%)	1.2 (0.69-1.9)	3.3% (1.5-5.5%)	3.8% (1.6-6.5%)	260 (120-450)	6400 (3100-11000)
Sri Lanka	2005	Male	70 to 74	36% (24-49%)	64% (51-76%)	0.98 (0.52-1.6)	2.5% (1-4.6%)	2.9% (1.1-5.3%)	230 (88-420)	4500 (1800-8100)
Sri Lanka	2005	Male	75 to 79	34% (22-49%)	66% (51-78%)	0.91 (0.47-1.6)	1.7% (0.4-3.7%)	1.9% (0.36-4.2%)	140 (26-310)	2200 (530-4700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	2005	Male	80 to 84	29% (17-44%)	71% (56-83%)	0.75 (0.38-1.3)	1.3% (0.24-2.8%)	1.4% (0.18-3.2%)	90 (11-200)	1100 (190-2300)
Sri Lanka	2005	Male	85 to 89	25% (15-37%)	75% (63-85%)	0.6 (0.3-1.1)	0.88% (0.15-1.9%)	0.94% (0.072-2.2%)	39 (3-91)	340 (61-750)
Sri Lanka	2005	Male	90 to 94	25% (15-37%)	75% (63-85%)	0.56 (0.28-1)	0.63% (0.0096-1.5%)	0.62% (-0.14-1.7%)	14 (-3.2-38)	90 (1.4-220)
Sri Lanka	2005	Male	95 plus	26% (14-42%)	74% (58-86%)	0.55 (0.22-1.1)	0.61% (-0.089-1.7%)	0.49% (-0.36-1.7%)	3.4 (-2.6-12)	18 (-2.6-48)
Sri Lanka	2010	Female	All Ages	5.2% (4.4-6.1%)	95% (94-96%)	0.34 (0.25-0.46)	1.5% (1-2%)	2.3% (1.4-3.6%)	1200 (690-1900)	34000 (23000-48000)
Sri Lanka	2010	Female	15 to 19	4.4% (2.7-6.5%)	96% (93-97%)	0.24 (0.12-0.45)	0.82% (0.46-1.4%)	1% (0.38-2.2%)	7.2 (2.6-15)	920 (500-1600)
Sri Lanka	2010	Female	20 to 24	5.5% (3.3-8.2%)	95% (92-97%)	0.33 (0.16-0.61)	1.4% (0.85-2.2%)	1.5% (0.54-2.9%)	10 (3.9-21)	1600 (940-2500)
Sri Lanka	2010	Female	25 to 29	6% (3.6-9.3%)	94% (91-96%)	0.38 (0.19-0.68)	1.5% (0.97-2.2%)	1.7% (0.77-3.2%)	12 (5.4-24)	1700 (1100-2700)
Sri Lanka	2010	Female	30 to 34	6.4% (3.7-10%)	94% (90-96%)	0.41 (0.21-0.75)	1.5% (0.98-2.1%)	1.9% (0.97-3.4%)	14 (7.4-27)	1800 (1100-2600)
Sri Lanka	2010	Female	35 to 39	6.6% (3.7-10%)	93% (90-96%)	0.42 (0.21-0.76)	1.5% (0.93-2.2%)	2.1% (1.2-3.7%)	17 (9.1-30)	1700 (1000-2500)
Sri Lanka	2010	Female	40 to 44	6.6% (4-9.9%)	93% (90-96%)	0.45 (0.22-0.78)	1.5% (0.93-2.2%)	2.4% (1.4-3.9%)	24 (14-38)	1800 (1100-2800)
Sri Lanka	2010	Female	45 to 49	6.4% (3.7-9.5%)	94% (90-96%)	0.49 (0.25-0.89)	1.7% (1-2.6%)	2.8% (1.6-4.6%)	40 (23-66)	2400 (1500-3700)
Sri Lanka	2010	Female	50 to 54	6.5% (4.1-9.7%)	93% (90-96%)	0.56 (0.29-0.99)	1.9% (1.2-3.1%)	3.1% (1.8-5%)	66 (38-110)	3100 (1900-4900)
Sri Lanka	2010	Female	55 to 59	6.7% (4.1-10%)	93% (90-96%)	0.62 (0.31-1.1)	2.1% (1.2-3.4%)	3.3% (1.8-5.3%)	88 (46-140)	3500 (1900-5600)
Sri Lanka	2010	Female	60 to 64	6.4% (3.9-9.9%)	94% (90-96%)	0.61 (0.31-1)	2.1% (1.1-3.5%)	3.1% (1.6-5.2%)	110 (59-190)	3700 (2000-6400)
Sri Lanka	2010	Female	65 to 69	5.9% (3.6-8.5%)	94% (92-96%)	0.56 (0.29-0.95)	2% (1.1-3.4%)	2.8% (1.4-4.7%)	130 (69-230)	3400 (1800-5800)
Sri Lanka	2010	Female	70 to 74	5.7% (3.7-8.6%)	94% (91-96%)	0.54 (0.28-0.95)	2% (0.97-3.7%)	2.6% (1.2-4.6%)	150 (70-270)	2900 (1400-5200)
Sri Lanka	2010	Female	75 to 79	6.1% (3.7-9.3%)	94% (91-96%)	0.56 (0.26-0.99)	1.8% (0.78-3.4%)	2.3% (0.96-4.3%)	140 (60-280)	2200 (960-4300)
Sri Lanka	2010	Female	80 to 84	6.6% (4-9.8%)	93% (90-96%)	0.59 (0.29-1)	1.9% (0.84-3.5%)	2.3% (1-4.3%)	160 (70-320)	1800 (800-3400)
Sri Lanka	2010	Female	85 to 89	7.3% (4.3-11%)	93% (89-96%)	0.63 (0.3-1.1)	1.8% (0.72-3.4%)	2.2% (0.85-4.2%)	120 (47-240)	950 (380-1800)
Sri Lanka	2010	Female	90 to 94	8.4% (5.1-13%)	92% (87-95%)	0.7 (0.34-1.4)	1.8% (0.64-3.7%)	2.1% (0.72-4.6%)	66 (22-140)	360 (130-770)
Sri Lanka	2010	Female	95 plus	10% (5.3-17%)	90% (83-95%)	0.81 (0.33-1.6)	1.9% (0.48-3.9%)	2.3% (0.5-4.8%)	28 (6.1-59)	100 (26-210)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	2010	Male	All Ages	35% (30-41%)	65% (59-70%)	1.6 (1.2-2)	5.6% (4-7.5%)	6.3% (4.1-9%)	4700 (3000-6900)	180000 (120000-240000)
Sri Lanka	2010	Male	15 to 19	25% (16-37%)	75% (63-84%)	0.52 (0.25-0.94)	1.5% (0.79-2.6%)	1.7% (0.71-3.4%)	22 (9.7-42)	2200 (1200-3600)
Sri Lanka	2010	Male	20 to 24	38% (25-52%)	62% (48-75%)	1.3 (0.68-2.2)	3.7% (2-6.2%)	4.1% (1.9-7.6%)	91 (43-160)	7400 (4000-12000)
Sri Lanka	2010	Male	25 to 29	45% (30-59%)	55% (41-70%)	1.9 (1-3)	5.4% (3.1-8.6%)	6.5% (3.3-11%)	140 (74-230)	10000 (6000-16000)
Sri Lanka	2010	Male	30 to 34	50% (33-65%)	50% (35-67%)	2.4 (1.4-3.7)	6.8% (4.3-10%)	8.7% (5-13%)	190 (110-290)	13000 (7900-19000)
Sri Lanka	2010	Male	35 to 39	53% (38-67%)	47% (33-62%)	2.9 (1.7-4.4)	8.8% (5.8-12%)	12% (7.3-17%)	270 (170-400)	16000 (10000-23000)
Sri Lanka	2010	Male	40 to 44	54% (39-68%)	46% (32-61%)	3.1 (1.8-4.6)	10% (6.8-14%)	13% (8.9-18%)	380 (250-540)	20000 (13000-27000)
Sri Lanka	2010	Male	45 to 49	53% (36-67%)	47% (33-64%)	3.1 (1.9-4.7)	10% (7.1-14%)	14% (9.1-19%)	530 (340-750)	24000 (16000-34000)
Sri Lanka	2010	Male	50 to 54	51% (36-65%)	49% (35-64%)	2.8 (1.7-4.4)	9.4% (5.9-13%)	12% (7.5-17%)	600 (360-860)	24000 (14000-34000)
Sri Lanka	2010	Male	55 to 59	49% (34-65%)	51% (35-66%)	2.5 (1.4-4)	7.9% (4.5-12%)	9.8% (5.7-15%)	580 (320-890)	20000 (12000-31000)
Sri Lanka	2010	Male	60 to 64	45% (31-62%)	55% (38-69%)	2.1 (1.1-3.5)	6.2% (3-9.8%)	7.7% (3.6-12%)	540 (250-890)	16000 (7600-26000)
Sri Lanka	2010	Male	65 to 69	40% (26-55%)	60% (45-74%)	1.7 (0.91-2.7)	5.1% (2.5-8.5%)	6.1% (2.9-10%)	450 (210-760)	11000 (5400-18000)
Sri Lanka	2010	Male	70 to 74	35% (22-49%)	65% (51-78%)	1.4 (0.73-2.4)	4.1% (1.8-7.4%)	4.8% (2-8.6%)	360 (150-650)	6900 (3000-12000)
Sri Lanka	2010	Male	75 to 79	32% (19-48%)	68% (52-81%)	1.2 (0.62-2.3)	3.1% (1.1-6.2%)	3.5% (1.2-7.2%)	250 (80-510)	3800 (1300-7600)
Sri Lanka	2010	Male	80 to 84	28% (16-43%)	72% (57-84%)	1.1 (0.51-1.9)	2.5% (0.86-5.2%)	2.8% (0.86-6%)	180 (54-380)	2000 (700-4100)
Sri Lanka	2010	Male	85 to 89	25% (15-38%)	75% (62-85%)	0.9 (0.46-1.6)	1.8% (0.54-3.9%)	2.1% (0.52-4.5%)	92 (23-200)	760 (230-1600)
Sri Lanka	2010	Male	90 to 94	25% (15-37%)	75% (63-85%)	0.84 (0.41-1.5)	1.4% (0.28-3.2%)	1.5% (0.18-3.6%)	37 (4.4-91)	220 (45-510)
Sri Lanka	2010	Male	95 plus	27% (15-43%)	73% (57-85%)	0.82 (0.32-1.7)	1.3% (0.16-3.4%)	1.3% (-0.049-3.7%)	13 (-0.47-37)	55 (6.4-140)
Sri Lanka	2016	Female	All Ages	5.3% (4.4-6.1%)	95% (94-96%)	0.43 (0.28-0.62)	1.8% (1.2-2.5%)	2.8% (1.7-4.3%)	1400 (800-2300)	39000 (25000-57000)
Sri Lanka	2016	Female	15 to 19	4.3% (2.7-6.6%)	96% (93-97%)	0.3 (0.14-0.54)	1.1% (0.58-1.8%)	1.9% (0.72-3.9%)	7.5 (2.6-16)	950 (500-1600)
Sri Lanka	2016	Female	20 to 24	5.4% (3.2-8.1%)	95% (92-97%)	0.4 (0.19-0.75)	1.8% (1-2.8%)	2.6% (0.97-5%)	10 (3.7-21)	1600 (890-2500)
Sri Lanka	2016	Female	25 to 29	6% (3.5-9.2%)	94% (91-96%)	0.47 (0.21-0.86)	1.8% (1.2-2.7%)	2.7% (1.2-4.8%)	11 (4.3-22)	1600 (960-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	2016	Female	30 to 34	6.3% (3.7-10%)	94% (90-96%)	0.5 (0.23-0.98)	1.8% (1.1-2.6%)	2.9% (1.5-5%)	14 (6.3-26)	1800 (1100-2700)
Sri Lanka	2016	Female	35 to 39	6.5% (3.6-9.6%)	93% (90-96%)	0.5 (0.24-0.92)	1.7% (1.1-2.5%)	3% (1.6-5%)	19 (9.1-35)	1900 (1200-3000)
Sri Lanka	2016	Female	40 to 44	6.5% (3.9-9.6%)	94% (90-96%)	0.54 (0.25-0.97)	1.7% (1.1-2.5%)	3.2% (1.8-5.1%)	24 (13-42)	1900 (1200-3000)
Sri Lanka	2016	Female	45 to 49	6.4% (3.6-9.5%)	94% (91-96%)	0.6 (0.29-1.1)	1.9% (1.1-3%)	3.5% (2-5.4%)	40 (20-69)	2500 (1400-3900)
Sri Lanka	2016	Female	50 to 54	6.5% (4-9.6%)	94% (90-96%)	0.68 (0.33-1.2)	2.2% (1.3-3.4%)	3.8% (2.2-5.8%)	67 (35-120)	3300 (1900-5400)
Sri Lanka	2016	Female	55 to 59	6.6% (4.1-9.6%)	93% (90-96%)	0.77 (0.36-1.3)	2.3% (1.3-3.6%)	3.8% (2.2-5.8%)	95 (49-160)	3900 (2100-6300)
Sri Lanka	2016	Female	60 to 64	6.3% (3.8-9.5%)	94% (91-96%)	0.76 (0.36-1.3)	2.3% (1.3-3.7%)	3.5% (2-5.6%)	120 (62-200)	4100 (2100-6600)
Sri Lanka	2016	Female	65 to 69	5.8% (3.5-8.5%)	94% (91-97%)	0.69 (0.34-1.2)	2.2% (1.2-3.7%)	3.2% (1.8-5.1%)	180 (92-310)	4800 (2500-8100)
Sri Lanka	2016	Female	70 to 74	5.7% (3.6-8.6%)	94% (91-96%)	0.66 (0.32-1.2)	2.2% (1.1-3.6%)	2.9% (1.5-4.9%)	220 (110-390)	4400 (2200-7700)
Sri Lanka	2016	Female	75 to 79	6.1% (3.7-9.2%)	94% (91-96%)	0.69 (0.32-1.3)	2% (0.97-3.7%)	2.6% (1.2-4.8%)	160 (67-290)	2500 (1100-4500)
Sri Lanka	2016	Female	80 to 84	6.6% (3.9-9.8%)	93% (90-96%)	0.72 (0.34-1.3)	2.2% (1-3.9%)	2.7% (1.3-4.8%)	190 (82-350)	2200 (960-3900)
Sri Lanka	2016	Female	85 to 89	7.4% (4.3-11%)	93% (89-96%)	0.78 (0.35-1.5)	2.1% (0.89-3.8%)	2.6% (1.1-4.7%)	150 (62-280)	1200 (530-2200)
Sri Lanka	2016	Female	90 to 94	8.4% (5.1-13%)	92% (87-95%)	0.87 (0.39-1.7)	2.1% (0.79-3.8%)	2.5% (0.93-4.8%)	89 (31-170)	500 (190-930)
Sri Lanka	2016	Female	95 plus	10% (5.3-17%)	90% (83-95%)	1 (0.38-2.1)	2.3% (0.69-4.3%)	2.7% (0.73-5.3%)	38 (10-75)	140 (41-260)
Sri Lanka	2016	Male	All Ages	35% (30-41%)	65% (59-70%)	1.9 (1.3-2.7)	7.7% (5.3-10%)	8.5% (5.4-12%)	6100 (3500-9400)	220000 (140000-310000)
Sri Lanka	2016	Male	15 to 19	24% (15-36%)	76% (64-85%)	0.64 (0.28-1.2)	2.3% (1.2-4.1%)	3.8% (1.6-7%)	23 (8.7-46)	2300 (1200-4000)
Sri Lanka	2016	Male	20 to 24	38% (25-51%)	62% (49-75%)	1.6 (0.78-2.7)	6.7% (3.6-11%)	9.6% (4.5-16%)	100 (44-190)	8100 (4200-14000)
Sri Lanka	2016	Male	25 to 29	45% (29-59%)	55% (41-71%)	2.3 (1.1-3.7)	8.9% (5-14%)	13% (6.8-21%)	140 (63-240)	10000 (5300-16000)
Sri Lanka	2016	Male	30 to 34	50% (33-65%)	50% (35-67%)	2.9 (1.6-4.8)	10% (6.3-16%)	16% (9-23%)	200 (100-310)	13000 (7400-20000)
Sri Lanka	2016	Male	35 to 39	53% (38-68%)	47% (32-62%)	3.5 (1.9-5.7)	12% (7.5-17%)	17% (11-24%)	330 (180-500)	19000 (11000-28000)
Sri Lanka	2016	Male	40 to 44	54% (39-68%)	46% (32-61%)	3.8 (2.1-5.9)	12% (8.3-17%)	18% (12-24%)	430 (260-660)	22000 (14000-33000)
Sri Lanka	2016	Male	45 to 49	53% (36-67%)	47% (33-64%)	3.7 (2.1-6)	13% (8.5-17%)	17% (12-23%)	590 (350-890)	27000 (17000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sri Lanka	2016	Male	50 to 54	51% (36-65%)	49% (35-64%)	3.4 (1.9-5.5)	11% (6.9-16%)	15% (9.5-20%)	710 (390-1100)	28000 (16000-42000)
Sri Lanka	2016	Male	55 to 59	49% (34-64%)	51% (36-66%)	3 (1.6-4.9)	9.6% (5.6-14%)	12% (7.1-18%)	730 (390-1100)	25000 (14000-39000)
Sri Lanka	2016	Male	60 to 64	45% (30-62%)	55% (38-70%)	2.6 (1.4-4.3)	7.8% (4-12%)	9.8% (5-15%)	680 (330-1100)	20000 (9900-32000)
Sri Lanka	2016	Male	65 to 69	40% (26-54%)	60% (46-74%)	2.1 (1.1-3.5)	6.5% (3.4-10%)	7.9% (4-12%)	770 (350-1300)	19000 (8600-30000)
Sri Lanka	2016	Male	70 to 74	35% (23-48%)	65% (52-77%)	1.7 (0.85-3)	5.3% (2.4-8.9%)	6.3% (2.8-10%)	640 (270-1100)	12000 (5300-22000)
Sri Lanka	2016	Male	75 to 79	32% (19-47%)	68% (53-81%)	1.5 (0.7-2.9)	4.1% (1.6-7.6%)	4.8% (1.8-8.9%)	350 (120-670)	5300 (2000-10000)
Sri Lanka	2016	Male	80 to 84	28% (16-42%)	72% (58-84%)	1.3 (0.61-2.4)	3.4% (1.3-6.6%)	4% (1.4-7.7%)	250 (81-500)	2800 (960-5500)
Sri Lanka	2016	Male	85 to 89	25% (15-38%)	75% (62-85%)	1.1 (0.51-2.1)	2.6% (0.9-5.2%)	3% (0.97-6.3%)	140 (42-290)	1100 (360-2300)
Sri Lanka	2016	Male	90 to 94	25% (16-37%)	75% (63-84%)	1 (0.47-2)	2% (0.56-4.4%)	2.3% (0.55-5.2%)	60 (14-140)	350 (96-780)
Sri Lanka	2016	Male	95 plus	27% (15-43%)	73% (57-85%)	1 (0.39-2.1)	2% (0.4-4.6%)	2% (0.23-5.1%)	21 (2.5-54)	87 (18-200)
Thailand	1990	Female	All Ages	10% (7.2-14%)	90% (86-93%)	0.62 (0.46-0.82)	3% (2.4-3.8%)	5.8% (4.2-7.8%)	7900 (5600-11000)	220000 (180000-280000)
Thailand	1990	Female	15 to 19	9% (4-17%)	91% (83-96%)	0.47 (0.16-1)	1.5% (0.76-2.8%)	2.9% (1.3-5.7%)	69 (30-140)	6300 (3100-12000)
Thailand	1990	Female	20 to 24	11% (4.5-21%)	89% (79-95%)	0.63 (0.22-1.3)	2.3% (1.2-3.8%)	3.9% (1.6-7.3%)	100 (43-190)	9400 (5000-16000)
Thailand	1990	Female	25 to 29	12% (5.1-22%)	88% (78-95%)	0.72 (0.26-1.5)	2.7% (1.5-4.5%)	4.8% (2.2-8.8%)	120 (56-230)	11000 (5900-18000)
Thailand	1990	Female	30 to 34	13% (5.5-24%)	87% (76-94%)	0.83 (0.32-1.7)	3% (1.8-4.8%)	5.5% (3.1-8.9%)	160 (88-260)	12000 (7200-18000)
Thailand	1990	Female	35 to 39	14% (6.4-26%)	86% (74-94%)	0.93 (0.37-1.8)	3.5% (2.2-5.4%)	6.5% (3.8-9.9%)	220 (130-340)	14000 (8500-20000)
Thailand	1990	Female	40 to 44	15% (6.7-26%)	85% (74-93%)	1 (0.38-2)	4.3% (2.7-6.5%)	7.4% (4.6-11%)	300 (180-460)	16000 (10000-24000)
Thailand	1990	Female	45 to 49	15% (6.5-27%)	85% (73-93%)	1.2 (0.39-2.5)	5.2% (3.2-7.8%)	8.5% (5.3-13%)	420 (260-640)	19000 (12000-29000)
Thailand	1990	Female	50 to 54	15% (6.8-28%)	85% (72-93%)	1.3 (0.48-2.8)	6% (3.7-8.9%)	9.2% (5.7-13%)	630 (390-920)	25000 (16000-36000)
Thailand	1990	Female	55 to 59	15% (7.2-28%)	85% (72-93%)	1.4 (0.51-2.9)	6% (3.7-9.1%)	8.9% (5.5-14%)	720 (440-1100)	24000 (15000-37000)
Thailand	1990	Female	60 to 64	15% (6.7-28%)	85% (72-93%)	1.4 (0.51-3.2)	6% (3.6-9.5%)	8.5% (5.1-13%)	820 (480-1300)	23000 (14000-37000)
Thailand	1990	Female	65 to 69	15% (6.8-26%)	85% (74-93%)	1.3 (0.49-2.8)	6.2% (3.5-9.5%)	8.5% (4.9-13%)	870 (510-1300)	20000 (12000-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	1990	Female	70 to 74	14% (6.7-25%)	86% (75-93%)	1.2 (0.4-2.8)	5.5% (2.8-9%)	7.2% (3.7-12%)	860 (440-1400)	16000 (8200-26000)
Thailand	1990	Female	75 to 79	14% (6.2-25%)	86% (75-94%)	1.1 (0.34-2.6)	4.7% (2.1-8.3%)	6.1% (2.7-11%)	760 (350-1300)	11000 (5000-19000)
Thailand	1990	Female	80 to 84	14% (6-26%)	86% (74-94%)	1.1 (0.34-2.6)	5% (2.2-8.6%)	6.4% (2.8-11%)	830 (360-1400)	8700 (3800-15000)
Thailand	1990	Female	85 to 89	15% (6.3-27%)	85% (73-94%)	1.2 (0.36-2.8)	4.7% (1.9-8.9%)	5.8% (2.4-11%)	620 (250-1200)	4600 (1900-8700)
Thailand	1990	Female	90 to 94	17% (7.3-31%)	83% (69-93%)	1.4 (0.35-3.4)	4.2% (1.5-8.1%)	5.1% (1.8-9.9%)	280 (98-550)	1400 (510-2800)
Thailand	1990	Female	95 plus	21% (8.2-40%)	79% (60-92%)	1.7 (0.34-4.4)	4.2% (1.1-8.9%)	4.9% (1.2-11%)	83 (20-180)	290 (72-600)
Thailand	1990	Male	All Ages	31% (24-39%)	69% (61-76%)	1.8 (1.5-2.1)	8.1% (6.7-9.7%)	11% (8.9-14%)	21000 (17000-26000)	800000 (670000-950000)
Thailand	1990	Male	15 to 19	25% (15-41%)	75% (59-85%)	0.82 (0.37-1.5)	3.7% (1.9-6.3%)	5.1% (2.5-9.2%)	300 (150-540)	23000 (12000-41000)
Thailand	1990	Male	20 to 24	37% (21-56%)	63% (44-79%)	1.9 (1-3.1)	8.8% (5.1-14%)	11% (6.2-18%)	920 (500-1500)	65000 (38000-100000)
Thailand	1990	Male	25 to 29	43% (24-61%)	57% (39-76%)	2.7 (1.5-4.1)	12% (7.6-17%)	16% (9.4-23%)	1200 (750-1800)	81000 (50000-120000)
Thailand	1990	Male	30 to 34	46% (26-65%)	54% (35-74%)	3.3 (1.9-4.9)	14% (9.4-18%)	18% (12-24%)	1400 (930-1900)	84000 (58000-110000)
Thailand	1990	Male	35 to 39	49% (30-68%)	51% (32-70%)	3.8 (2.1-5.8)	15% (11-20%)	20% (14-26%)	1500 (1100-2000)	84000 (59000-110000)
Thailand	1990	Male	40 to 44	48% (30-69%)	52% (31-70%)	3.9 (2.3-6.2)	15% (11-20%)	20% (14-26%)	1600 (1100-2000)	78000 (56000-99000)
Thailand	1990	Male	45 to 49	47% (27-66%)	53% (34-73%)	3.9 (2.1-6.1)	15% (11-20%)	20% (14-26%)	1700 (1200-2200)	76000 (53000-97000)
Thailand	1990	Male	50 to 54	45% (26-64%)	55% (36-74%)	3.7 (2-5.9)	15% (10-19%)	19% (13-24%)	2100 (1400-2700)	80000 (56000-100000)
Thailand	1990	Male	55 to 59	42% (25-62%)	58% (38-75%)	3.3 (1.7-5.5)	13% (8.9-17%)	17% (11-22%)	2100 (1500-2800)	70000 (48000-94000)
Thailand	1990	Male	60 to 64	39% (23-59%)	61% (41-77%)	2.9 (1.5-5.1)	12% (7.3-16%)	14% (9.2-20%)	2100 (1300-2900)	59000 (38000-81000)
Thailand	1990	Male	65 to 69	36% (21-54%)	64% (46-79%)	2.5 (1.2-4.2)	10% (6.1-15%)	13% (7.7-18%)	1800 (1100-2600)	42000 (25000-60000)
Thailand	1990	Male	70 to 74	32% (18-50%)	68% (50-82%)	2 (0.94-3.7)	9.1% (5-14%)	11% (6-17%)	1600 (860-2400)	29000 (16000-45000)
Thailand	1990	Male	75 to 79	29% (17-47%)	71% (53-83%)	1.7 (0.75-3.2)	7.1% (3.4-11%)	8.5% (4.1-14%)	1100 (540-1800)	16000 (7800-26000)
Thailand	1990	Male	80 to 84	27% (14-44%)	73% (56-86%)	1.5 (0.65-2.8)	6.6% (3.3-11%)	7.8% (3.8-13%)	860 (430-1400)	9100 (4500-15000)
Thailand	1990	Male	85 to 89	26% (13-41%)	74% (59-87%)	1.3 (0.59-2.5)	5.1% (2.3-9%)	5.9% (2.7-11%)	430 (200-780)	3300 (1500-5900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	1990	Male	90 to 94	27% (15-44%)	73% (56-85%)	1.3 (0.53-2.5)	3.9% (1.5-7.5%)	4.5% (1.7-8.7%)	150 (56-290)	770 (300-1500)
Thailand	1990	Male	95 plus	31% (15-50%)	69% (50-85%)	1.3 (0.46-2.9)	3.6% (1.1-7.6%)	3.9% (1.1-8.5%)	35 (9.7-76)	120 (39-270)
Thailand	1995	Female	All Ages	11% (7.9-14%)	89% (86-92%)	0.74 (0.56-0.98)	3.4% (2.7-4.3%)	6.1% (4.4-8%)	9000 (6600-12000)	260000 (210000-320000)
Thailand	1995	Female	15 to 19	9.2% (4.3-17%)	91% (83-96%)	0.53 (0.18-1.1)	1.6% (0.82-3%)	3.1% (1.4-5.9%)	71 (31-140)	6500 (3200-12000)
Thailand	1995	Female	20 to 24	11% (4.9-21%)	89% (79-95%)	0.72 (0.24-1.5)	2.4% (1.3-4.1%)	3.6% (1.5-6.7%)	120 (51-230)	11000 (5600-19000)
Thailand	1995	Female	25 to 29	12% (5.5-22%)	88% (78-94%)	0.83 (0.3-1.6)	2.8% (1.5-4.5%)	4.2% (1.8-7.5%)	160 (70-290)	13000 (7100-22000)
Thailand	1995	Female	30 to 34	13% (5.6-25%)	87% (75-94%)	0.94 (0.34-1.9)	3.1% (1.9-4.9%)	5.1% (2.8-8.3%)	200 (110-320)	14000 (8600-22000)
Thailand	1995	Female	35 to 39	14% (6.2-26%)	86% (74-94%)	1 (0.41-2)	3.6% (2.2-5.5%)	6.2% (3.7-9.5%)	260 (150-410)	17000 (10000-25000)
Thailand	1995	Female	40 to 44	15% (6.9-27%)	85% (73-93%)	1.2 (0.43-2.3)	4.2% (2.7-6.3%)	7.2% (4.5-11%)	370 (220-550)	20000 (12000-29000)
Thailand	1995	Female	45 to 49	15% (6.3-27%)	85% (73-94%)	1.3 (0.45-2.7)	5.1% (3.2-7.4%)	8.3% (5.3-12%)	480 (300-710)	22000 (14000-33000)
Thailand	1995	Female	50 to 54	15% (6.7-28%)	85% (72-93%)	1.4 (0.54-3)	5.9% (3.8-8.6%)	9.1% (5.9-13%)	620 (400-900)	25000 (16000-36000)
Thailand	1995	Female	55 to 59	15% (7.1-28%)	85% (72-93%)	1.6 (0.55-3.2)	5.9% (3.6-8.8%)	8.7% (5.5-13%)	780 (490-1200)	27000 (17000-41000)
Thailand	1995	Female	60 to 64	15% (6.8-28%)	85% (72-93%)	1.6 (0.55-3.5)	5.9% (3.6-9.3%)	8.4% (5-13%)	970 (590-1500)	28000 (17000-44000)
Thailand	1995	Female	65 to 69	15% (6.7-26%)	85% (74-93%)	1.5 (0.55-3)	5.9% (3.6-9%)	8.2% (5.1-12%)	1100 (650-1600)	25000 (15000-39000)
Thailand	1995	Female	70 to 74	14% (6.7-25%)	86% (75-93%)	1.3 (0.46-3.1)	5.3% (2.9-8.5%)	7.1% (4-11%)	970 (540-1500)	18000 (10000-29000)
Thailand	1995	Female	75 to 79	14% (5.9-25%)	86% (75-94%)	1.3 (0.38-2.8)	4.6% (2.2-7.8%)	5.9% (2.9-9.9%)	830 (400-1400)	12000 (5900-20000)
Thailand	1995	Female	80 to 84	14% (6-25%)	86% (75-94%)	1.2 (0.39-2.8)	4.9% (2.3-8.1%)	6.1% (2.8-10%)	910 (410-1500)	9700 (4400-16000)
Thailand	1995	Female	85 to 89	15% (6.7-27%)	85% (73-93%)	1.3 (0.42-3)	4.6% (2-8.5%)	5.6% (2.4-10%)	740 (320-1400)	5600 (2400-10000)
Thailand	1995	Female	90 to 94	17% (7.6-31%)	83% (69-92%)	1.5 (0.41-3.6)	4.1% (1.6-7.9%)	4.9% (1.9-9.5%)	350 (130-680)	1800 (720-3500)
Thailand	1995	Female	95 plus	21% (8.1-39%)	79% (61-92%)	1.8 (0.39-4.7)	4.1% (1.2-8.4%)	4.7% (1.3-9.9%)	110 (30-230)	390 (110-790)
Thailand	1995	Male	All Ages	33% (26-41%)	67% (59-74%)	2.2 (1.9-2.5)	9.6% (7.9-11%)	12% (9.9-14%)	27000 (22000-32000)	110000 (89000-130000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	1995	Male	15 to 19	26% (14-41%)	74% (59-86%)	0.91 (0.42-1.7)	4.3% (2.2-7.7%)	5.6% (2.7-10%)	440 (210-790)	33000 (16000-59000)
Thailand	1995	Male	20 to 24	38% (22-56%)	62% (44-78%)	2.2 (1.1-3.6)	9.6% (5.4-15%)	12% (6.4-19%)	1400 (740-2200)	95000 (53000-150000)
Thailand	1995	Male	25 to 29	44% (25-62%)	56% (38-75%)	3 (1.7-4.6)	12% (7.6-18%)	14% (8.6-21%)	2200 (1300-3300)	140000 (87000-210000)
Thailand	1995	Male	30 to 34	47% (28-66%)	53% (34-72%)	3.7 (2.2-5.4)	14% (9.6-19%)	16% (11-22%)	2400 (1600-3200)	140000 (96000-190000)
Thailand	1995	Male	35 to 39	49% (32-69%)	51% (31-68%)	4.2 (2.4-6.3)	15% (11-20%)	18% (13-24%)	2400 (1700-3100)	130000 (90000-170000)
Thailand	1995	Male	40 to 44	49% (31-68%)	51% (32-69%)	4.4 (2.6-6.6)	15% (11-19%)	19% (13-24%)	2200 (1500-2800)	110000 (77000-140000)
Thailand	1995	Male	45 to 49	47% (28-67%)	53% (33-72%)	4.3 (2.3-6.8)	15% (11-19%)	19% (14-24%)	2100 (1500-2800)	93000 (67000-120000)
Thailand	1995	Male	50 to 54	45% (26-64%)	55% (36-74%)	4.1 (2.2-6.5)	14% (10-18%)	18% (13-23%)	2100 (1500-2600)	80000 (57000-100000)
Thailand	1995	Male	55 to 59	43% (26-63%)	57% (37-74%)	3.7 (2-6.1)	13% (8.9-17%)	16% (11-21%)	2200 (1500-2900)	75000 (52000-99000)
Thailand	1995	Male	60 to 64	40% (23-60%)	60% (40-77%)	3.3 (1.6-5.6)	11% (7.5-15%)	14% (9.3-19%)	2300 (1500-3100)	66000 (43000-89000)
Thailand	1995	Male	65 to 69	37% (21-55%)	63% (45-79%)	2.8 (1.4-4.6)	10% (6.4-14%)	13% (8-18%)	2200 (1400-3000)	51000 (32000-70000)
Thailand	1995	Male	70 to 74	33% (19-50%)	67% (50-81%)	2.3 (1.1-4.1)	8.9% (5.2-13%)	11% (6.5-16%)	1800 (1000-2600)	33000 (19000-49000)
Thailand	1995	Male	75 to 79	30% (17-47%)	70% (53-83%)	1.9 (0.87-3.6)	7% (3.6-11%)	8.3% (4.3-13%)	1200 (610-1900)	18000 (9100-27000)
Thailand	1995	Male	80 to 84	27% (15-44%)	73% (56-85%)	1.7 (0.72-3.1)	6.3% (3.2-10%)	7.4% (3.7-12%)	940 (470-1500)	10000 (5100-16000)
Thailand	1995	Male	85 to 89	26% (14-41%)	74% (59-86%)	1.5 (0.69-2.7)	5% (2.5-8.5%)	5.7% (2.8-10%)	520 (250-910)	4000 (2000-6900)
Thailand	1995	Male	90 to 94	28% (15-45%)	72% (55-85%)	1.5 (0.65-2.8)	3.9% (1.6-7.1%)	4.4% (1.8-8.4%)	180 (73-340)	960 (400-1800)
Thailand	1995	Male	95 plus	31% (15-52%)	69% (48-85%)	1.5 (0.55-3.1)	3.5% (1.2-7.4%)	3.8% (1.1-8.1%)	42 (12-89)	150 (49-320)
Thailand	2000	Female	All Ages	11% (8.1-15%)	89% (85-92%)	0.89 (0.66-1.2)	3.6% (2.8-4.5%)	6.1% (4.4-8.1%)	10000 (7400-14000)	290000 (230000-370000)
Thailand	2000	Female	15 to 19	9.4% (4.4-17%)	91% (83-96%)	0.59 (0.2-1.3)	1.7% (0.83-3.1%)	3.4% (1.5-6.8%)	61 (26-120)	5700 (2800-10000)
Thailand	2000	Female	20 to 24	11% (5-21%)	89% (79-95%)	0.8 (0.27-1.7)	2.3% (1.2-3.8%)	3.5% (1.5-6.4%)	110 (45-190)	9600 (4900-16000)
Thailand	2000	Female	25 to 29	12% (5.1-22%)	88% (78-95%)	0.9 (0.32-1.9)	2.7% (1.4-4.4%)	3.6% (1.6-6.4%)	210 (95-370)	16000 (8300-27000)
Thailand	2000	Female	30 to 34	13% (5.4-24%)	87% (76-95%)	1 (0.36-2)	3% (1.8-4.8%)	4.2% (2.3-6.7%)	280 (150-460)	20000 (12000-31000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	2000	Female	35 to 39	14% (6-26%)	86% (74-94%)	1.1 (0.43-2.2)	3.3% (2-5%)	5.2% (3-7.9%)	320 (190-490)	20000 (12000-30000)
Thailand	2000	Female	40 to 44	15% (6.8-26%)	85% (74-93%)	1.3 (0.49-2.5)	3.9% (2.5-5.8%)	6.5% (4.1-9.5%)	430 (270-630)	23000 (15000-34000)
Thailand	2000	Female	45 to 49	15% (6.5-27%)	85% (73-94%)	1.4 (0.49-2.9)	4.6% (3-7%)	7.7% (4.9-11%)	550 (350-810)	26000 (17000-39000)
Thailand	2000	Female	50 to 54	15% (6.7-29%)	85% (71-93%)	1.6 (0.62-3.3)	5.6% (3.6-8.1%)	8.7% (5.7-13%)	670 (430-990)	27000 (18000-40000)
Thailand	2000	Female	55 to 59	16% (7.3-29%)	84% (71-93%)	1.8 (0.65-3.8)	5.8% (3.6-8.8%)	8.6% (5.4-13%)	770 (480-1100)	27000 (17000-41000)
Thailand	2000	Female	60 to 64	16% (7-29%)	84% (71-93%)	1.8 (0.63-4)	5.8% (3.6-9.1%)	8.3% (5.3-13%)	1000 (650-1600)	30000 (18000-49000)
Thailand	2000	Female	65 to 69	15% (6.9-27%)	85% (73-93%)	1.7 (0.61-3.3)	5.8% (3.6-8.9%)	8.2% (5-12%)	1200 (730-1900)	29000 (18000-45000)
Thailand	2000	Female	70 to 74	14% (6.8-26%)	86% (74-93%)	1.5 (0.51-3.4)	5.2% (2.9-8.4%)	7.1% (4-11%)	1200 (670-1900)	23000 (13000-37000)
Thailand	2000	Female	75 to 79	14% (6.2-26%)	86% (74-94%)	1.4 (0.4-3.2)	4.5% (2.1-7.7%)	5.9% (2.8-9.7%)	980 (460-1600)	15000 (6900-24000)
Thailand	2000	Female	80 to 84	14% (6.3-26%)	86% (74-94%)	1.4 (0.42-3.1)	4.8% (2.1-8.1%)	6% (2.7-10%)	1000 (460-1800)	11000 (4900-19000)
Thailand	2000	Female	85 to 89	15% (6.7-28%)	85% (72-93%)	1.5 (0.47-3.4)	4.6% (2-8.3%)	5.6% (2.5-10%)	800 (350-1500)	6100 (2600-11000)
Thailand	2000	Female	90 to 94	17% (7.7-31%)	83% (69-92%)	1.7 (0.47-4.1)	4.2% (1.6-8.2%)	5.1% (2-9.9%)	440 (170-840)	2300 (910-4400)
Thailand	2000	Female	95 plus	21% (8.3-40%)	79% (60-92%)	2 (0.42-5.6)	4.3% (1.2-8.7%)	4.9% (1.3-10%)	150 (40-310)	510 (150-1000)
Thailand	2000	Male	All Ages	34% (27-42%)	66% (58-73%)	2.6 (2.2-3)	9.9% (8.3-12%)	12% (10-14%)	29000 (24000-35000)	1100000 (950000-1400000)
Thailand	2000	Male	15 to 19	26% (15-41%)	74% (59-85%)	1 (0.45-1.9)	4.7% (2.4-8.3%)	6.3% (3.1-12%)	390 (190-710)	29000 (15000-52000)
Thailand	2000	Male	20 to 24	38% (22-56%)	62% (44-78%)	2.4 (1.2-4.1)	10% (5.8-16%)	13% (7.1-21%)	1100 (600-1800)	80000 (44000-130000)
Thailand	2000	Male	25 to 29	44% (25-62%)	56% (38-75%)	3.3 (1.8-5.2)	12% (7.2-17%)	14% (8.1-20%)	1800 (1100-2700)	120000 (72000-170000)
Thailand	2000	Male	30 to 34	47% (27-66%)	53% (34-73%)	4 (2.3-6)	12% (8.4-16%)	14% (9.5-19%)	2400 (1700-3300)	140000 (100000-200000)
Thailand	2000	Male	35 to 39	49% (31-69%)	51% (31-69%)	4.5 (2.7-6.8)	13% (9.3-17%)	15% (11-20%)	2500 (1800-3400)	140000 (99000-180000)
Thailand	2000	Male	40 to 44	49% (30-69%)	51% (31-70%)	4.7 (2.8-7)	13% (10-17%)	17% (12-21%)	2600 (1900-3400)	130000 (95000-170000)
Thailand	2000	Male	45 to 49	47% (27-67%)	53% (33-73%)	4.6 (2.6-7.2)	14% (11-18%)	18% (14-22%)	2600 (1900-3300)	110000 (86000-150000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	2000	Male	50 to 54	45% (26-65%)	55% (35-74%)	4.4 (2.4-6.9)	14% (10-17%)	18% (13-22%)	2400 (1800-3100)	94000 (69000-120000)
Thailand	2000	Male	55 to 59	43% (25-63%)	57% (37-75%)	4 (2.2-6.7)	13% (9.2-17%)	17% (12-21%)	2300 (1600-2900)	77000 (54000-99000)
Thailand	2000	Male	60 to 64	40% (23-61%)	60% (39-77%)	3.6 (1.7-6.1)	12% (7.7-15%)	15% (9.9-19%)	2600 (1700-3400)	73000 (48000-97000)
Thailand	2000	Male	65 to 69	36% (21-56%)	64% (44-79%)	3 (1.5-5.1)	10% (6.9-14%)	13% (8.7-18%)	2500 (1600-3400)	59000 (39000-81000)
Thailand	2000	Male	70 to 74	33% (19-49%)	67% (51-81%)	2.5 (1.2-4.4)	9% (5.5-13%)	11% (6.8-16%)	2300 (1400-3300)	43000 (26000-63000)
Thailand	2000	Male	75 to 79	30% (17-47%)	70% (53-83%)	2.1 (0.91-3.9)	7.1% (4-11%)	8.7% (4.9-13%)	1500 (830-2200)	22000 (12000-33000)
Thailand	2000	Male	80 to 84	28% (14-44%)	72% (56-86%)	1.8 (0.82-3.4)	6.4% (3.2-9.9%)	7.6% (3.8-12%)	1100 (570-1700)	12000 (6200-19000)
Thailand	2000	Male	85 to 89	26% (14-41%)	74% (59-86%)	1.6 (0.73-3)	5.2% (2.4-8.7%)	6.1% (2.8-10%)	620 (280-1000)	4700 (2200-8000)
Thailand	2000	Male	90 to 94	28% (16-44%)	72% (56-84%)	1.6 (0.73-3)	4.2% (1.8-7.6%)	4.8% (2-8.9%)	240 (98-450)	1300 (540-2400)
Thailand	2000	Male	95 plus	31% (15-52%)	69% (48-85%)	1.6 (0.57-3.4)	3.8% (1.2-8%)	4.1% (1.2-8.8%)	56 (16-120)	200 (68-420)
Thailand	2005	Female	All Ages	11% (8.2-15%)	89% (85-92%)	0.97 (0.73-1.3)	3.6% (2.8-4.6%)	6.1% (4.4-8.2%)	11000 (7700-14000)	280000 (220000-360000)
Thailand	2005	Female	15 to 19	9.1% (4.2-17%)	91% (83-96%)	0.61 (0.21-1.4)	1.5% (0.77-2.7%)	3.2% (1.4-6.4%)	46 (20-93)	4500 (2200-8100)
Thailand	2005	Female	20 to 24	11% (4.7-20%)	89% (80-95%)	0.82 (0.27-1.7)	2% (1.1-3.3%)	3.4% (1.5-6.3%)	69 (30-130)	6900 (3800-11000)
Thailand	2005	Female	25 to 29	12% (5.1-22%)	88% (78-95%)	0.94 (0.32-2)	2.2% (1.2-3.6%)	3.2% (1.4-5.7%)	120 (55-210)	10000 (5600-17000)
Thailand	2005	Female	30 to 34	13% (5.2-24%)	87% (76-95%)	1.1 (0.36-2.2)	2.5% (1.4-4%)	3.7% (1.9-6.2%)	180 (90-300)	14000 (7700-21000)
Thailand	2005	Female	35 to 39	14% (5.7-26%)	86% (74-94%)	1.2 (0.44-2.3)	2.8% (1.7-4.3%)	4.7% (2.6-7.2%)	250 (140-390)	16000 (9500-24000)
Thailand	2005	Female	40 to 44	14% (6.5-26%)	86% (74-93%)	1.3 (0.49-2.5)	3.5% (2.2-5.1%)	6% (3.7-8.8%)	370 (230-540)	21000 (13000-31000)
Thailand	2005	Female	45 to 49	14% (6.4-25%)	86% (75-94%)	1.5 (0.53-2.9)	4.1% (2.6-6.1%)	7.2% (4.5-11%)	520 (330-790)	25000 (16000-38000)
Thailand	2005	Female	50 to 54	15% (6.9-27%)	85% (73-93%)	1.7 (0.67-3.5)	5% (3.2-7.4%)	8.2% (5.3-12%)	720 (460-1100)	30000 (19000-44000)
Thailand	2005	Female	55 to 59	15% (7.2-29%)	85% (71-93%)	1.9 (0.69-3.8)	5.5% (3.3-8.5%)	8.4% (5.2-13%)	840 (530-1300)	29000 (18000-46000)
Thailand	2005	Female	60 to 64	15% (6.9-28%)	85% (72-93%)	1.9 (0.68-4)	5.5% (3.4-8.7%)	8% (5-13%)	950 (590-1500)	28000 (17000-45000)
Thailand	2005	Female	65 to 69	15% (6.9-27%)	85% (73-93%)	1.7 (0.64-3.4)	5.6% (3.4-8.5%)	8% (4.9-12%)	1300 (780-1900)	31000 (19000-48000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	2005	Female	70 to 74	14% (6.5-25%)	86% (75-93%)	1.6 (0.52-3.5)	5% (2.8-7.9%)	6.8% (3.9-11%)	1300 (760-2100)	26000 (15000-40000)
Thailand	2005	Female	75 to 79	13% (6.1-25%)	87% (75-94%)	1.4 (0.43-3.3)	4.3% (2.1-7.3%)	5.7% (2.9-9.4%)	1200 (610-2000)	18000 (9000-30000)
Thailand	2005	Female	80 to 84	14% (6.3-26%)	86% (74-94%)	1.4 (0.45-3.2)	4.5% (2-7.7%)	5.7% (2.6-9.8%)	1300 (570-2200)	14000 (6100-23000)
Thailand	2005	Female	85 to 89	15% (6.6-27%)	85% (73-93%)	1.6 (0.49-3.5)	4.4% (2-8.1%)	5.5% (2.4-10%)	920 (400-1700)	7000 (3100-13000)
Thailand	2005	Female	90 to 94	17% (7.5-32%)	83% (68-93%)	1.8 (0.48-4.3)	4.2% (1.7-7.7%)	5.1% (2-9.6%)	480 (190-910)	2500 (1000-4700)
Thailand	2005	Female	95 plus	21% (8.1-40%)	79% (60-92%)	2.1 (0.44-5.8)	4.2% (1.3-8.7%)	4.9% (1.4-10%)	180 (50-360)	610 (180-1200)
Thailand	2005	Male	All Ages	34% (28-43%)	66% (57-72%)	2.8 (2.4-3.2)	10% (8.6-12%)	12% (10-15%)	29000 (24000-35000)	1100000 (920000-1300000)
Thailand	2005	Male	15 to 19	25% (14-41%)	75% (59-86%)	1 (0.46-1.9)	4.6% (2.5-8.2%)	6.5% (3.3-12%)	330 (170-600)	25000 (14000-45000)
Thailand	2005	Male	20 to 24	37% (21-56%)	63% (44-79%)	2.4 (1.2-4.2)	10% (5.9-16%)	14% (7.5-22%)	890 (480-1400)	64000 (37000-100000)
Thailand	2005	Male	25 to 29	43% (25-61%)	57% (39-75%)	3.4 (1.8-5.5)	12% (7.6-17%)	15% (9.2-22%)	1300 (790-1900)	89000 (56000-130000)
Thailand	2005	Male	30 to 34	46% (26-65%)	54% (35-74%)	4.1 (2.4-6.3)	12% (8.4-16%)	15% (10-19%)	1700 (1200-2300)	110000 (74000-140000)
Thailand	2005	Male	35 to 39	48% (29-68%)	52% (32-71%)	4.7 (2.6-7)	12% (8.4-16%)	15% (10-19%)	2000 (1300-2600)	110000 (78000-140000)
Thailand	2005	Male	40 to 44	48% (31-67%)	52% (33-69%)	4.9 (3-7.3)	13% (10-17%)	17% (13-21%)	2400 (1800-3100)	120000 (92000-150000)
Thailand	2005	Male	45 to 49	47% (27-66%)	53% (34-73%)	4.9 (2.8-7.3)	14% (11-17%)	19% (14-23%)	2800 (2100-3400)	120000 (95000-150000)
Thailand	2005	Male	50 to 54	45% (26-64%)	55% (36-74%)	4.6 (2.6-7)	14% (11-18%)	19% (14-23%)	2900 (2200-3600)	110000 (85000-140000)
Thailand	2005	Male	55 to 59	42% (26-62%)	58% (38-74%)	4.2 (2.2-6.9)	13% (9.6-17%)	18% (13-22%)	2700 (1900-3500)	92000 (67000-120000)
Thailand	2005	Male	60 to 64	39% (22-59%)	61% (41-78%)	3.7 (1.8-6.3)	12% (8.1-16%)	15% (10-20%)	2500 (1700-3300)	73000 (50000-96000)
Thailand	2005	Male	65 to 69	36% (20-54%)	64% (46-80%)	3.1 (1.5-5.1)	10% (7.1-14%)	13% (9.1-18%)	2700 (1800-3600)	64000 (44000-87000)
Thailand	2005	Male	70 to 74	32% (18-50%)	68% (50-82%)	2.5 (1.2-4.7)	8.8% (5.5-13%)	11% (6.9-16%)	2500 (1500-3600)	47000 (29000-67000)
Thailand	2005	Male	75 to 79	29% (17-46%)	71% (54-83%)	2.1 (0.95-4.1)	7% (4-11%)	8.7% (5-13%)	1900 (1100-2800)	28000 (16000-41000)
Thailand	2005	Male	80 to 84	27% (15-43%)	73% (57-85%)	1.9 (0.84-3.5)	6.2% (3.1-9.7%)	7.4% (3.8-12%)	1400 (730-2300)	16000 (8000-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	2005	Male	85 to 89	25% (14-40%)	75% (60-86%)	1.6 (0.73-3)	5.1% (2.3-8.3%)	6% (2.7-9.9%)	750 (340-1300)	5800 (2600-9500)
Thailand	2005	Male	90 to 94	27% (15-43%)	73% (57-85%)	1.6 (0.72-3)	4.1% (1.7-7.5%)	4.8% (1.9-8.7%)	280 (110-520)	1500 (620-2700)
Thailand	2005	Male	95 plus	30% (15-50%)	70% (50-85%)	1.6 (0.58-3.5)	3.9% (1.2-7.9%)	4.2% (1.2-8.7%)	73 (20-150)	260 (80-530)
Thailand	2010	Female	All Ages	11% (8.3-15%)	89% (85-92%)	1 (0.76-1.4)	3.6% (2.8-4.6%)	6.1% (4.4-8.2%)	11000 (7600-14000)	270000 (220000-350000)
Thailand	2010	Female	15 to 19	9% (4.2-17%)	91% (83-96%)	0.63 (0.21-1.4)	1.5% (0.78-2.8%)	3.3% (1.4-6.5%)	41 (18-82)	4000 (2000-7400)
Thailand	2010	Female	20 to 24	11% (4.6-20%)	89% (80-95%)	0.84 (0.26-1.9)	1.9% (1.1-3.1%)	3.5% (1.5-6.4%)	48 (21-88)	5400 (3000-8900)
Thailand	2010	Female	25 to 29	12% (5.2-23%)	88% (77-95%)	0.95 (0.31-2)	2.1% (1.2-3.3%)	3.4% (1.5-6.2%)	71 (31-130)	7000 (3900-12000)
Thailand	2010	Female	30 to 34	13% (5.3-25%)	87% (75-95%)	1.1 (0.38-2.2)	2.3% (1.3-3.6%)	3.9% (1.8-6.6%)	120 (56-210)	9900 (5500-16000)
Thailand	2010	Female	35 to 39	14% (5.8-26%)	86% (74-94%)	1.2 (0.45-2.3)	2.7% (1.6-4.3%)	4.9% (2.6-8%)	200 (110-330)	14000 (7700-21000)
Thailand	2010	Female	40 to 44	14% (6.5-25%)	86% (75-94%)	1.3 (0.49-2.5)	3.3% (2.1-5%)	6.1% (3.8-9.2%)	340 (210-520)	19000 (13000-29000)
Thailand	2010	Female	45 to 49	14% (5.8-26%)	86% (74-94%)	1.4 (0.54-2.9)	3.9% (2.4-6%)	7.2% (4.6-11%)	510 (320-760)	25000 (16000-37000)
Thailand	2010	Female	50 to 54	15% (6.4-28%)	85% (72-94%)	1.6 (0.64-3.4)	4.7% (2.9-7%)	8.1% (5.1-12%)	740 (460-1100)	31000 (19000-46000)
Thailand	2010	Female	55 to 59	15% (6.8-28%)	85% (72-93%)	1.8 (0.65-3.7)	5% (3.1-7.8%)	8.2% (5.2-13%)	880 (550-1400)	31000 (20000-49000)
Thailand	2010	Female	60 to 64	15% (6.4-28%)	85% (72-94%)	1.8 (0.63-4)	5.1% (3.2-8.1%)	7.9% (5-12%)	970 (610-1500)	29000 (18000-46000)
Thailand	2010	Female	65 to 69	14% (6.1-26%)	86% (74-94%)	1.7 (0.62-3.3)	5.3% (3.3-7.9%)	7.8% (5-11%)	1100 (700-1700)	27000 (17000-42000)
Thailand	2010	Female	70 to 74	14% (6.3-25%)	86% (75-94%)	1.5 (0.52-3.6)	4.7% (2.7-7.4%)	6.7% (4-10%)	1400 (790-2100)	26000 (15000-41000)
Thailand	2010	Female	75 to 79	13% (5.9-25%)	87% (75-94%)	1.5 (0.44-3.4)	4.2% (2.1-6.9%)	5.7% (3-9.2%)	1300 (680-2200)	20000 (10000-33000)
Thailand	2010	Female	80 to 84	13% (5.9-26%)	87% (74-94%)	1.5 (0.43-3.2)	4.2% (1.9-7%)	5.5% (2.6-9.5%)	1300 (620-2300)	15000 (6800-25000)
Thailand	2010	Female	85 to 89	15% (6.4-28%)	85% (72-94%)	1.6 (0.45-3.6)	4.1% (1.8-7.6%)	5.2% (2.3-9.6%)	1000 (440-1900)	8000 (3400-15000)
Thailand	2010	Female	90 to 94	17% (7.8-32%)	83% (68-92%)	1.8 (0.47-4.3)	3.8% (1.4-7.1%)	4.8% (1.7-9.2%)	470 (180-900)	2500 (940-4800)
Thailand	2010	Female	95 plus	21% (8.1-39%)	79% (61-92%)	2.2 (0.45-5.8)	4% (1.2-8.1%)	4.8% (1.3-9.8%)	170 (46-340)	590 (170-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	2010	Male	All Ages	35% (28-43%)	65% (57-72%)	2.9 (2.4-3.5)	11% (9-13%)	13% (11-15%)	30000 (24000-35000)	1100000 (910000-1200000)
Thailand	2010	Male	15 to 19	24% (14-39%)	76% (61-86%)	1 (0.45-1.9)	4.6% (2.6-7.9%)	6.6% (3.6-12%)	260 (140-450)	20000 (11000-34000)
Thailand	2010	Male	20 to 24	36% (20-54%)	64% (46-80%)	2.4 (1.2-4.1)	9.9% (5.7-15%)	14% (7.5-22%)	610 (330-950)	45000 (26000-69000)
Thailand	2010	Male	25 to 29	42% (24-60%)	58% (40-76%)	3.3 (1.7-5.6)	12% (7.5-17%)	16% (9.7-23%)	920 (560-1300)	63000 (39000-89000)
Thailand	2010	Male	30 to 34	45% (26-64%)	55% (36-74%)	4.1 (2.2-6.5)	13% (8.8-17%)	17% (11-22%)	1400 (950-1900)	88000 (61000-120000)
Thailand	2010	Male	35 to 39	48% (30-67%)	52% (33-70%)	4.7 (2.7-7.2)	13% (9.4-17%)	17% (12-22%)	1900 (1300-2500)	110000 (75000-140000)
Thailand	2010	Male	40 to 44	48% (30-67%)	52% (33-70%)	4.9 (2.9-7.2)	14% (10-17%)	18% (13-23%)	2400 (1800-3100)	120000 (92000-160000)
Thailand	2010	Male	45 to 49	46% (27-65%)	54% (35-73%)	4.8 (2.6-7.2)	15% (11-18%)	20% (15-24%)	3000 (2300-3800)	140000 (100000-170000)
Thailand	2010	Male	50 to 54	44% (25-63%)	56% (37-75%)	4.5 (2.5-7)	14% (11-18%)	19% (15-24%)	3300 (2500-4100)	130000 (100000-160000)
Thailand	2010	Male	55 to 59	42% (25-62%)	58% (38-75%)	4.1 (2.2-6.6)	13% (9.6-17%)	18% (13-22%)	3200 (2300-4000)	110000 (80000-140000)
Thailand	2010	Male	60 to 64	39% (22-58%)	61% (42-78%)	3.6 (1.7-6.1)	12% (8.2-15%)	15% (11-20%)	2800 (2000-3700)	82000 (57000-110000)
Thailand	2010	Male	65 to 69	35% (20-53%)	65% (47-80%)	3 (1.5-5)	10% (7-14%)	13% (9.1-18%)	2500 (1700-3400)	61000 (41000-81000)
Thailand	2010	Male	70 to 74	31% (18-49%)	69% (51-82%)	2.5 (1.2-4.5)	8.6% (5.4-12%)	11% (6.9-15%)	2600 (1600-3600)	49000 (31000-68000)
Thailand	2010	Male	75 to 79	29% (16-46%)	71% (54-84%)	2.1 (0.93-4)	6.8% (3.8-10%)	8.6% (4.8-13%)	2000 (1100-3000)	30000 (17000-44000)
Thailand	2010	Male	80 to 84	26% (14-44%)	74% (56-86%)	1.8 (0.79-3.4)	5.7% (2.9-9.1%)	7.1% (3.6-11%)	1500 (740-2400)	17000 (8300-26000)
Thailand	2010	Male	85 to 89	25% (13-40%)	75% (60-87%)	1.6 (0.69-3.1)	4.5% (2.1-7.6%)	5.5% (2.5-9.5%)	790 (350-1400)	6300 (2900-11000)
Thailand	2010	Male	90 to 94	26% (14-44%)	74% (56-86%)	1.6 (0.66-2.9)	3.6% (1.5-6.9%)	4.4% (1.7-8.4%)	260 (100-510)	1500 (610-2700)
Thailand	2010	Male	95 plus	30% (14-51%)	70% (49-86%)	1.6 (0.56-3.4)	3.5% (1.1-7.3%)	3.9% (1-8.5%)	64 (17-140)	250 (76-520)
Thailand	2016	Female	All Ages	12% (8.6-15%)	88% (85-91%)	1.1 (0.73-1.6)	3.5% (2.7-4.7%)	5.9% (4.1-7.9%)	12000 (7900-16000)	280000 (210000-370000)
Thailand	2016	Female	15 to 19	8.9% (4.2-16%)	91% (84-96%)	0.62 (0.2-1.4)	1.5% (0.75-2.6%)	3.3% (1.4-6.6%)	33 (14-63)	3400 (1700-6200)
Thailand	2016	Female	20 to 24	11% (4.5-20%)	89% (80-95%)	0.82 (0.25-1.9)	1.9% (1.1-3%)	3.5% (1.6-6.4%)	39 (18-73)	4800 (2800-7800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	2016	Female	25 to 29	11% (4.8-21%)	89% (79-95%)	0.93 (0.29-2)	1.9% (1.1-3%)	3.3% (1.4-6%)	51 (21-95)	5700 (3300-9100)
Thailand	2016	Female	30 to 34	12% (5.2-23%)	88% (77-95%)	1 (0.34-2.2)	2.1% (1.2-3.3%)	3.6% (1.7-6.3%)	83 (37-150)	7600 (4300-12000)
Thailand	2016	Female	35 to 39	14% (5.7-25%)	86% (75-94%)	1.2 (0.41-2.4)	2.4% (1.4-3.8%)	4.5% (2.1-7.3%)	150 (71-260)	11000 (6100-18000)
Thailand	2016	Female	40 to 44	14% (6.4-24%)	86% (76-94%)	1.3 (0.47-2.7)	3.1% (1.9-4.7%)	5.7% (3.4-8.7%)	290 (170-450)	17000 (10000-26000)
Thailand	2016	Female	45 to 49	14% (6-26%)	86% (74-94%)	1.5 (0.5-3.1)	3.7% (2.3-5.6%)	6.8% (4.2-10%)	480 (290-740)	24000 (15000-36000)
Thailand	2016	Female	50 to 54	15% (6.3-28%)	85% (72-94%)	1.6 (0.63-3.6)	4.4% (2.7-6.7%)	7.9% (4.8-12%)	730 (440-1100)	31000 (19000-46000)
Thailand	2016	Female	55 to 59	15% (6.9-28%)	85% (72-93%)	1.8 (0.64-3.7)	4.7% (2.9-7.4%)	8.1% (5.1-12%)	940 (590-1500)	34000 (21000-54000)
Thailand	2016	Female	60 to 64	15% (6.5-28%)	85% (72-93%)	1.8 (0.63-3.9)	4.8% (3-7.8%)	7.7% (4.9-12%)	1200 (730-1800)	36000 (22000-57000)
Thailand	2016	Female	65 to 69	14% (6.6-27%)	86% (73-93%)	1.7 (0.61-3.5)	5% (3.1-7.6%)	7.6% (4.8-11%)	1300 (820-1900)	32000 (20000-48000)
Thailand	2016	Female	70 to 74	14% (6.6-25%)	86% (75-93%)	1.5 (0.51-3.5)	4.4% (2.6-6.9%)	6.5% (3.8-9.8%)	1300 (720-2000)	25000 (14000-39000)
Thailand	2016	Female	75 to 79	13% (5.9-25%)	87% (75-94%)	1.4 (0.4-3.3)	3.8% (2-6.4%)	5.3% (2.8-8.6%)	1400 (700-2200)	21000 (10000-34000)
Thailand	2016	Female	80 to 84	13% (6-25%)	87% (75-94%)	1.4 (0.43-3.4)	3.8% (1.7-6.4%)	5.1% (2.3-8.8%)	1500 (680-2500)	16000 (7500-28000)
Thailand	2016	Female	85 to 89	14% (6.8-27%)	86% (73-93%)	1.6 (0.46-3.7)	3.7% (1.5-6.7%)	4.9% (2-8.8%)	1300 (510-2300)	9800 (4000-18000)
Thailand	2016	Female	90 to 94	17% (7.3-30%)	83% (70-93%)	1.8 (0.46-4.6)	3.5% (1.2-6.6%)	4.5% (1.6-8.5%)	660 (230-1300)	3500 (1200-6700)
Thailand	2016	Female	95 plus	20% (7.8-39%)	80% (61-92%)	2.2 (0.43-6)	3.7% (1.1-7.5%)	4.5% (1.2-9.3%)	220 (57-450)	770 (210-1500)
Thailand	2016	Male	All Ages	35% (29-44%)	65% (56-71%)	3 (2.2-4.1)	11% (8.7-13%)	13% (9.7-15%)	32000 (24000-40000)	1100000 (880000-1400000)
Thailand	2016	Male	15 to 19	24% (14-38%)	76% (62-86%)	0.99 (0.41-2)	4.5% (2.4-7.7%)	6.8% (3.4-12%)	220 (110-390)	17000 (9100-29000)
Thailand	2016	Male	20 to 24	36% (20-55%)	64% (45-80%)	2.4 (1.1-4.5)	9.9% (5.7-15%)	14% (7.5-23%)	550 (290-910)	42000 (24000-65000)
Thailand	2016	Male	25 to 29	42% (24-60%)	58% (40-76%)	3.4 (1.5-6)	12% (7.2-17%)	17% (9.5-24%)	800 (450-1200)	56000 (33000-80000)
Thailand	2016	Male	30 to 34	45% (26-65%)	55% (35-74%)	4.1 (2.1-7)	13% (8.6-17%)	17% (11-24%)	1200 (760-1700)	76000 (50000-110000)
Thailand	2016	Male	35 to 39	47% (30-66%)	53% (34-70%)	4.6 (2.4-7.7)	14% (9.7-18%)	19% (13-24%)	1800 (1200-2500)	100000 (71000-140000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Thailand	2016	Male	40 to 44	47% (30-67%)	53% (33-70%)	4.8 (2.5-8)	14% (11-19%)	19% (14-24%)	2500 (1800-3300)	120000 (90000-160000)
Thailand	2016	Male	45 to 49	46% (27-65%)	54% (35-73%)	4.8 (2.4-7.8)	15% (11-19%)	20% (15-25%)	3200 (2300-4100)	140000 (100000-180000)
Thailand	2016	Male	50 to 54	44% (25-64%)	56% (36-75%)	4.5 (2.3-7.4)	14% (11-18%)	19% (14-24%)	3600 (2600-4700)	140000 (100000-180000)
Thailand	2016	Male	55 to 59	42% (24-62%)	58% (38-76%)	4.1 (2.1-6.8)	13% (9.2-17%)	17% (13-22%)	3700 (2700-4800)	130000 (90000-160000)
Thailand	2016	Male	60 to 64	38% (22-59%)	62% (41-78%)	3.6 (1.7-6.3)	11% (7.7-15%)	15% (11-20%)	3600 (2400-4900)	110000 (72000-140000)
Thailand	2016	Male	65 to 69	35% (20-54%)	65% (46-80%)	3 (1.4-5.4)	9.9% (6.6-13%)	13% (8.9-17%)	3200 (2100-4300)	76000 (50000-100000)
Thailand	2016	Male	70 to 74	31% (18-49%)	69% (51-82%)	2.4 (1.1-4.6)	8.2% (5-12%)	11% (6.3-15%)	2600 (1500-3800)	51000 (29000-73000)
Thailand	2016	Male	75 to 79	28% (16-45%)	72% (55-84%)	2.1 (0.84-4)	6.4% (3.4-9.8%)	8.1% (4.5-12%)	2200 (1100-3300)	33000 (17000-49000)
Thailand	2016	Male	80 to 84	26% (14-43%)	74% (57-86%)	1.8 (0.73-3.4)	5.3% (2.5-8.4%)	6.6% (3.2-11%)	1700 (800-2700)	19000 (9000-30000)
Thailand	2016	Male	85 to 89	24% (13-39%)	76% (61-87%)	1.6 (0.65-3.1)	4.2% (1.8-7.2%)	5.2% (2.1-9.1%)	920 (380-1700)	7400 (3100-13000)
Thailand	2016	Male	90 to 94	26% (14-42%)	74% (58-86%)	1.5 (0.62-3)	3.4% (1.2-6.5%)	4.2% (1.4-8.4%)	380 (130-750)	2100 (750-4100)
Thailand	2016	Male	95 plus	29% (14-50%)	71% (50-86%)	1.6 (0.51-3.5)	3.4% (0.92-7.2%)	3.7% (0.82-8.5%)	88 (19-200)	340 (93-730)
Timor-Leste	1990	Female	All Ages	8.4% (5.9-12%)	92% (88-94%)	0.024 (0.018-0.032)	0.12% (0.055-0.19%)	0.091% (-0.031-0.22%)	3.2 (-1.1-8)	300 (130-480)
Timor-Leste	1990	Female	15 to 19	8.3% (3.8-16%)	92% (84-96%)	0.022 (0.0077-0.047)	0.27% (0.16-0.4%)	0.16% (0.037-0.33%)	0.13 (0.03-0.28)	26 (15-40)
Timor-Leste	1990	Female	20 to 24	10% (4.2-19%)	90% (81-96%)	0.029 (0.01-0.061)	0.44% (0.25-0.66%)	0.15% (-0.028-0.37%)	0.15 (-0.029-0.37)	44 (24-68)
Timor-Leste	1990	Female	25 to 29	11% (4.7-21%)	89% (79-95%)	0.034 (0.013-0.068)	0.47% (0.27-0.72%)	0.17% (-0.048-0.47%)	0.17 (-0.049-0.46)	43 (23-68)
Timor-Leste	1990	Female	30 to 34	12% (4.7-23%)	88% (77-95%)	0.038 (0.014-0.076)	0.47% (0.25-0.7%)	0.26% (0.012-0.54%)	0.25 (0.012-0.53)	39 (20-61)
Timor-Leste	1990	Female	35 to 39	13% (5.3-24%)	87% (76-95%)	0.043 (0.016-0.082)	0.46% (0.24-0.69%)	0.34% (0.066-0.66%)	0.33 (0.064-0.67)	34 (17-53)
Timor-Leste	1990	Female	40 to 44	13% (5.9-24%)	87% (76-94%)	0.048 (0.018-0.099)	0.46% (0.23-0.71%)	0.43% (0.11-0.8%)	0.43 (0.11-0.81)	32 (15-51)
Timor-Leste	1990	Female	45 to 49	14% (6-25%)	86% (75-94%)	0.055 (0.019-0.12)	0.43% (0.16-0.71%)	0.42% (0.06-0.84%)	0.45 (0.061-0.92)	26 (9.4-46)
Timor-Leste	1990	Female	50 to 54	14% (6.1-26%)	86% (74-94%)	0.061 (0.021-0.14)	0.44% (0.086-0.82%)	0.48% (0.0077-1%)	0.56 (0.0082-1.2)	25 (5-48)
Timor-Leste	1990	Female	55 to 59	14% (6.3-26%)	86% (74-94%)	0.067 (0.022-0.14)	0.33% (-0.055-0.64%)	0.36% (-0.14-0.78%)	0.42 (-0.15-0.98)	16 (-2.8-34)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	1990	Female	60 to 64	14% (6.1-27%)	86% (73-94%)	0.068 (0.021-0.16)	0.19% (-0.24-0.54%)	0.2% (-0.33-0.64%)	0.28 (-0.47-0.95)	8.8 (-12-27)
Timor-Leste	1990	Female	65 to 69	14% (6-25%)	86% (75-94%)	0.064 (0.021-0.13)	0.091% (-0.36-0.42%)	0.096% (-0.44-0.52%)	0.13 (-0.59-0.66)	3.3 (-13-15)
Timor-Leste	1990	Female	70 to 74	13% (6.1-24%)	87% (76-94%)	0.058 (0.018-0.14)	0.054% (-0.38-0.35%)	0.056% (-0.46-0.41%)	0.073 (-0.59-0.57)	1.5 (-11-10)
Timor-Leste	1990	Female	75 to 79	12% (5.4-24%)	88% (76-95%)	0.054 (0.015-0.13)	-0.028% (-0.47-0.23%)	-0.039% (-0.56-0.26%)	-0.036 (-0.51-0.26)	-0.41 (-7-3.6)
Timor-Leste	1990	Female	80 to 84	12% (5.5-24%)	88% (76-94%)	0.053 (0.016-0.12)	-0.057% (-0.46-0.19%)	-0.074% (-0.55-0.22%)	-0.044 (-0.32-0.13)	-0.41 (-3.3-1.4)
Timor-Leste	1990	Female	85 to 89	13% (5.7-25%)	87% (75-94%)	0.056 (0.017-0.14)	-0.093% (-0.49-0.13%)	-0.12% (-0.59-0.14%)	-0.052 (-0.26-0.065)	-0.35 (-1.9-0.49)
Timor-Leste	1990	Female	90 to 94	15% (6.3-29%)	85% (71-94%)	0.065 (0.017-0.16)	-0.12% (-0.54-0.11%)	-0.16% (-0.67-0.12%)	-0.036 (-0.14-0.027)	-0.17 (-0.71-0.14)
Timor-Leste	1990	Female	95 plus	19% (7.1-37%)	81% (63-93%)	0.08 (0.016-0.22)	-0.13% (-0.67-0.15%)	-0.18% (-0.84-0.16%)	-0.0099 (-0.045-0.0088)	-0.029 (-0.14-0.033)
Timor-Leste	1990	Male	All Ages	26% (20-34%)	74% (66-80%)	0.079 (0.07-0.089)	0.31% (0.094-0.54%)	0.35% (-0.041-0.76%)	12 (-1.5-26)	850 (280-1400)
Timor-Leste	1990	Male	15 to 19	23% (14-38%)	77% (62-86%)	0.04 (0.019-0.074)	0.52% (0.34-0.79%)	0.4% (0.19-0.7%)	0.32 (0.15-0.58)	48 (30-73)
Timor-Leste	1990	Male	20 to 24	35% (19-53%)	65% (47-81%)	0.094 (0.049-0.15)	0.98% (0.56-1.5%)	0.64% (0.15-1.3%)	0.72 (0.16-1.4)	100 (59-160)
Timor-Leste	1990	Male	25 to 29	40% (22-59%)	60% (41-78%)	0.13 (0.074-0.2)	1.2% (0.45-2.1%)	0.89% (-0.19-2.1%)	0.92 (-0.19-2.2)	120 (42-200)
Timor-Leste	1990	Male	30 to 34	44% (25-63%)	56% (37-75%)	0.16 (0.098-0.24)	1.5% (0.48-2.5%)	1.3% (-0.11-2.9%)	1.2 (-0.11-2.7)	120 (40-210)
Timor-Leste	1990	Male	35 to 39	46% (28-66%)	54% (34-72%)	0.19 (0.1-0.29)	1.6% (0.49-2.8%)	1.7% (-0.049-3.6%)	1.5 (-0.039-3.2)	120 (33-210)
Timor-Leste	1990	Male	40 to 44	46% (28-66%)	54% (34-72%)	0.2 (0.11-0.31)	1.6% (0.36-2.9%)	1.8% (-0.068-3.9%)	1.7 (-0.052-3.6)	110 (24-190)
Timor-Leste	1990	Male	45 to 49	44% (25-64%)	56% (36-75%)	0.19 (0.1-0.32)	1.5% (0.2-2.9%)	1.8% (-0.21-3.8%)	1.7 (-0.2-3.7)	88 (11-170)
Timor-Leste	1990	Male	50 to 54	42% (24-62%)	58% (38-76%)	0.18 (0.094-0.3)	1.2% (-0.15-2.5%)	1.4% (-0.53-3.2%)	1.3 (-0.49-3.1)	60 (-7.1-120)
Timor-Leste	1990	Male	55 to 59	40% (23-60%)	60% (40-77%)	0.16 (0.079-0.29)	0.96% (-0.12-2%)	1.1% (-0.4-2.6%)	0.96 (-0.33-2.3)	37 (-4.2-79)
Timor-Leste	1990	Male	60 to 64	37% (21-56%)	63% (44-79%)	0.14 (0.067-0.26)	0.75% (-0.22-1.7%)	0.82% (-0.46-2.1%)	0.83 (-0.43-2.1)	27 (-7.3-62)
Timor-Leste	1990	Male	65 to 69	34% (19-52%)	66% (48-81%)	0.12 (0.057-0.21)	0.5% (-0.39-1.3%)	0.52% (-0.63-1.6%)	0.45 (-0.54-1.4)	12 (-9.5-33)
Timor-Leste	1990	Male	70 to 74	30% (17-47%)	70% (53-83%)	0.096 (0.043-0.18)	0.35% (-0.4-1.1%)	0.35% (-0.58-1.2%)	0.3 (-0.49-1.1)	6.6 (-7.2-20)
Timor-Leste	1990	Male	75 to 79	27% (15-45%)	73% (55-85%)	0.081 (0.033-0.16)	0.21% (-0.44-0.72%)	0.19% (-0.61-0.82%)	0.11 (-0.36-0.5)	2.2 (-4.5-7.3)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	1990	Male	80 to 84	25% (13-42%)	75% (58-87%)	0.07 (0.029-0.13)	0.16% (-0.37-0.6%)	0.13% (-0.52-0.67%)	0.044 (-0.17-0.23)	0.67 (-1.5-2.5)
Timor-Leste	1990	Male	85 to 89	24% (12-39%)	76% (61-88%)	0.062 (0.026-0.12)	0.11% (-0.28-0.42%)	0.079% (-0.39-0.48%)	0.02 (-0.1-0.12)	0.24 (-0.61-0.95)
Timor-Leste	1990	Male	90 to 94	25% (14-41%)	75% (59-86%)	0.06 (0.024-0.12)	0.05% (-0.29-0.31%)	0.015% (-0.41-0.33%)	0.0018 (-0.049-0.04)	0.038 (-0.22-0.23)
Timor-Leste	1990	Male	95 plus	29% (13-48%)	71% (52-87%)	0.063 (0.021-0.13)	0.069% (-0.33-0.34%)	0.0032% (-0.46-0.34%)	0.00011 (-0.016-0.012)	0.0095 (-0.044-0.047)
Timor-Leste	1995	Female	All Ages	8.4% (6-11%)	92% (89-94%)	0.027 (0.02-0.036)	0.14% (0.076-0.21%)	0.11% (-0.021-0.24%)	3.7 (-0.68-8.2)	340 (180-500)
Timor-Leste	1995	Female	15 to 19	8.2% (3.6-15%)	92% (85-96%)	0.025 (0.0085-0.055)	0.29% (0.17-0.44%)	0.17% (0.049-0.33%)	0.11 (0.03-0.21)	21 (12-34)
Timor-Leste	1995	Female	20 to 24	10% (4.3-18%)	90% (82-96%)	0.033 (0.011-0.069)	0.48% (0.28-0.71%)	0.15% (0.0066-0.34%)	0.15 (0.0065-0.34)	48 (27-75)
Timor-Leste	1995	Female	25 to 29	11% (4.7-20%)	89% (80-95%)	0.038 (0.013-0.075)	0.51% (0.31-0.74%)	0.18% (0.0035-0.41%)	0.17 (0.0036-0.41)	46 (27-71)
Timor-Leste	1995	Female	30 to 34	12% (4.8-22%)	88% (78-95%)	0.043 (0.017-0.087)	0.5% (0.31-0.72%)	0.27% (0.056-0.52%)	0.27 (0.056-0.54)	44 (26-66)
Timor-Leste	1995	Female	35 to 39	13% (5.3-24%)	87% (76-95%)	0.048 (0.019-0.093)	0.48% (0.29-0.71%)	0.35% (0.088-0.66%)	0.37 (0.089-0.7)	39 (22-60)
Timor-Leste	1995	Female	40 to 44	13% (6.1-24%)	87% (76-94%)	0.055 (0.02-0.11)	0.49% (0.28-0.71%)	0.45% (0.17-0.78%)	0.47 (0.18-0.82)	35 (20-53)
Timor-Leste	1995	Female	45 to 49	13% (5.7-25%)	87% (75-94%)	0.063 (0.021-0.14)	0.45% (0.22-0.69%)	0.45% (0.12-0.8%)	0.52 (0.14-0.95)	31 (15-49)
Timor-Leste	1995	Female	50 to 54	13% (5.8-26%)	87% (74-94%)	0.069 (0.024-0.15)	0.47% (0.14-0.82%)	0.52% (0.09-1%)	0.67 (0.12-1.3)	30 (9.6-55)
Timor-Leste	1995	Female	55 to 59	14% (6-25%)	86% (75-94%)	0.076 (0.025-0.16)	0.35% (-0.017-0.68%)	0.4% (-0.075-0.84%)	0.56 (-0.1-1.2)	21 (-0.99-42)
Timor-Leste	1995	Female	60 to 64	13% (6-25%)	87% (75-94%)	0.076 (0.024-0.18)	0.21% (-0.2-0.52%)	0.23% (-0.29-0.64%)	0.33 (-0.39-0.93)	10 (-9.7-27)
Timor-Leste	1995	Female	65 to 69	13% (5.8-24%)	87% (76-94%)	0.07 (0.024-0.15)	0.1% (-0.31-0.41%)	0.11% (-0.37-0.51%)	0.19 (-0.68-0.83)	4.8 (-15-19)
Timor-Leste	1995	Female	70 to 74	12% (5.9-23%)	88% (77-94%)	0.063 (0.02-0.15)	0.07% (-0.31-0.36%)	0.078% (-0.38-0.42%)	0.11 (-0.56-0.63)	2.2 (-9.8-11)
Timor-Leste	1995	Female	75 to 79	12% (5.2-23%)	88% (77-95%)	0.059 (0.017-0.14)	-0.024% (-0.44-0.23%)	-0.035% (-0.54-0.27%)	-0.041 (-0.63-0.31)	-0.47 (-8.8-4.4)
Timor-Leste	1995	Female	80 to 84	12% (5.4-23%)	88% (77-95%)	0.059 (0.018-0.13)	-0.061% (-0.44-0.17%)	-0.078% (-0.52-0.2%)	-0.05 (-0.34-0.13)	-0.48 (-3.5-1.4)
Timor-Leste	1995	Female	85 to 89	13% (5.9-24%)	87% (76-94%)	0.064 (0.02-0.15)	-0.1% (-0.5-0.13%)	-0.13% (-0.59-0.14%)	-0.051 (-0.24-0.054)	-0.35 (-1.7-0.41)
Timor-Leste	1995	Female	90 to 94	15% (6.8-28%)	85% (72-93%)	0.073 (0.02-0.17)	-0.14% (-0.57-0.1%)	-0.18% (-0.72-0.1%)	-0.034 (-0.14-0.02)	-0.16 (-0.66-0.12)
Timor-Leste	1995	Female	95 plus	18% (7.2-36%)	82% (64-93%)	0.09 (0.019-0.24)	-0.14% (-0.68-0.15%)	-0.2% (-0.85-0.16%)	-0.014 (-0.061-0.011)	-0.04 (-0.2-0.041)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	1995	Male	All Ages	26% (21-34%)	74% (66-79%)	0.092 (0.081-0.1)	0.39% (0.16-0.65%)	0.45% (0.023-0.93%)	15 (0.86-31)	1000 (410-1600)
Timor-Leste	1995	Male	15 to 19	23% (13-38%)	77% (62-87%)	0.045 (0.021-0.086)	0.52% (0.33-0.76%)	0.39% (0.18-0.7%)	0.27 (0.13-0.47)	39 (25-57)
Timor-Leste	1995	Male	20 to 24	35% (19-53%)	65% (47-81%)	0.11 (0.054-0.17)	0.92% (0.51-1.4%)	0.59% (0.14-1.1%)	0.82 (0.21-1.6)	120 (63-180)
Timor-Leste	1995	Male	25 to 29	40% (22-59%)	60% (41-78%)	0.15 (0.083-0.22)	1.2% (0.57-1.9%)	0.83% (0.051-1.7%)	1 (0.059-2.2)	130 (62-200)
Timor-Leste	1995	Male	30 to 34	44% (24-63%)	56% (37-76%)	0.18 (0.11-0.27)	1.4% (0.61-2.3%)	1.3% (0.1-2.5%)	1.5 (0.11-3.1)	140 (61-240)
Timor-Leste	1995	Male	35 to 39	46% (28-65%)	54% (35-72%)	0.21 (0.12-0.32)	1.6% (0.67-2.8%)	1.7% (0.19-3.5%)	1.9 (0.19-4)	140 (54-250)
Timor-Leste	1995	Male	40 to 44	46% (28-65%)	54% (35-72%)	0.22 (0.12-0.34)	1.7% (0.55-2.8%)	1.9% (0.16-3.7%)	2 (0.14-3.9)	120 (41-220)
Timor-Leste	1995	Male	45 to 49	44% (26-64%)	56% (36-74%)	0.22 (0.11-0.35)	1.6% (0.38-2.8%)	1.9% (0.16-3.7%)	2.2 (0.17-4.3)	110 (26-200)
Timor-Leste	1995	Male	50 to 54	42% (24-61%)	58% (39-76%)	0.21 (0.11-0.34)	1.3% (0.098-2.5%)	1.6% (-0.19-3.2%)	1.8 (-0.23-3.8)	80 (5.8-150)
Timor-Leste	1995	Male	55 to 59	39% (23-60%)	61% (40-77%)	0.19 (0.091-0.32)	1% (0.088-2%)	1.2% (-0.11-2.6%)	1.4 (-0.13-3)	53 (4.4-100)
Timor-Leste	1995	Male	60 to 64	37% (20-57%)	63% (43-80%)	0.16 (0.076-0.3)	0.81% (-0.036-1.6%)	0.91% (-0.24-2.1%)	1 (-0.28-2.4)	34 (-1.5-69)
Timor-Leste	1995	Male	65 to 69	33% (19-51%)	67% (49-81%)	0.13 (0.064-0.24)	0.54% (-0.25-1.3%)	0.58% (-0.46-1.6%)	0.69 (-0.51-1.9)	19 (-8.1-44)
Timor-Leste	1995	Male	70 to 74	30% (17-46%)	70% (54-83%)	0.11 (0.049-0.2)	0.38% (-0.31-1%)	0.39% (-0.47-1.2%)	0.39 (-0.47-1.2)	8.4 (-6.7-23)
Timor-Leste	1995	Male	75 to 79	27% (15-44%)	73% (56-85%)	0.092 (0.038-0.18)	0.23% (-0.34-0.73%)	0.21% (-0.5-0.84%)	0.17 (-0.39-0.68)	3.2 (-4.6-10)
Timor-Leste	1995	Male	80 to 84	25% (13-41%)	75% (59-87%)	0.079 (0.033-0.15)	0.17% (-0.3-0.59%)	0.15% (-0.44-0.69%)	0.057 (-0.17-0.26)	0.83 (-1.5-2.9)
Timor-Leste	1995	Male	85 to 89	23% (13-39%)	77% (61-87%)	0.07 (0.03-0.13)	0.11% (-0.26-0.44%)	0.09% (-0.37-0.49%)	0.026 (-0.1-0.14)	0.29 (-0.63-1.1)
Timor-Leste	1995	Male	90 to 94	25% (14-41%)	75% (59-86%)	0.068 (0.028-0.13)	0.05% (-0.28-0.32%)	0.014% (-0.39-0.35%)	0.0021 (-0.057-0.052)	0.045 (-0.24-0.29)
Timor-Leste	1995	Male	95 plus	28% (14-48%)	72% (52-86%)	0.07 (0.024-0.15)	0.074% (-0.3-0.34%)	0.0071% (-0.44-0.32%)	0.00032 (-0.019-0.014)	0.013 (-0.053-0.061)
Timor-Leste	2000	Female	All Ages	7.7% (5.5-11%)	92% (89-94%)	0.03 (0.023-0.04)	0.16% (0.085-0.24%)	0.13% (-0.043-0.3%)	3.3 (-1.1-7.5)	280 (150-410)
Timor-Leste	2000	Female	15 to 19	7.8% (3.4-15%)	92% (85-97%)	0.029 (0.01-0.064)	0.33% (0.19-0.5%)	0.2% (0.066-0.38%)	0.13 (0.039-0.25)	27 (16-41)
Timor-Leste	2000	Female	20 to 24	9.5% (3.9-18%)	91% (82-96%)	0.04 (0.013-0.086)	0.54% (0.32-0.79%)	0.18% (0.015-0.4%)	0.088 (0.0071-0.2)	29 (17-45)
Timor-Leste	2000	Female	25 to 29	10% (4.3-20%)	90% (80-96%)	0.046 (0.016-0.097)	0.57% (0.36-0.82%)	0.2% (0.016-0.45%)	0.11 (0.0085-0.25)	32 (19-48)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	2000	Female	30 to 34	11% (4.5-22%)	89% (78-95%)	0.052 (0.019-0.11)	0.55% (0.35-0.78%)	0.3% (0.075-0.55%)	0.17 (0.041-0.31)	29 (17-43)
Timor-Leste	2000	Female	35 to 39	12% (5.3-23%)	88% (77-95%)	0.058 (0.022-0.11)	0.52% (0.31-0.74%)	0.38% (0.1-0.69%)	0.27 (0.07-0.5)	31 (18-45)
Timor-Leste	2000	Female	40 to 44	13% (5.9-22%)	87% (78-94%)	0.066 (0.024-0.13)	0.51% (0.31-0.74%)	0.48% (0.17-0.8%)	0.37 (0.13-0.64)	29 (17-43)
Timor-Leste	2000	Female	45 to 49	13% (5.6-23%)	87% (77-94%)	0.075 (0.025-0.16)	0.47% (0.24-0.72%)	0.49% (0.17-0.85%)	0.44 (0.14-0.8)	27 (13-41)
Timor-Leste	2000	Female	50 to 54	13% (5.6-25%)	87% (75-94%)	0.083 (0.03-0.17)	0.49% (0.17-0.82%)	0.57% (0.14-1%)	0.65 (0.16-1.2)	29 (10-49)
Timor-Leste	2000	Female	55 to 59	13% (6.1-25%)	87% (75-94%)	0.091 (0.032-0.19)	0.38% (0.0095-0.73%)	0.45% (-0.043-0.92%)	0.6 (-0.049-1.3)	22 (0.53-43)
Timor-Leste	2000	Female	60 to 64	13% (5.8-25%)	87% (75-94%)	0.093 (0.031-0.2)	0.23% (-0.2-0.59%)	0.27% (-0.25-0.74%)	0.41 (-0.43-1.2)	12 (-11-33)
Timor-Leste	2000	Female	65 to 69	13% (5.8-23%)	87% (77-94%)	0.086 (0.029-0.18)	0.11% (-0.33-0.42%)	0.12% (-0.42-0.54%)	0.19 (-0.63-0.81)	4.8 (-13-19)
Timor-Leste	2000	Female	70 to 74	12% (5.7-23%)	88% (77-94%)	0.078 (0.025-0.18)	0.074% (-0.32-0.37%)	0.086% (-0.38-0.46%)	0.15 (-0.65-0.8)	2.8 (-12-14)
Timor-Leste	2000	Female	75 to 79	12% (5.4-22%)	88% (78-95%)	0.073 (0.022-0.18)	-0.028% (-0.47-0.24%)	-0.038% (-0.55-0.29%)	-0.047 (-0.68-0.36)	-0.57 (-9.6-5.1)
Timor-Leste	2000	Female	80 to 84	12% (5.5-23%)	88% (77-95%)	0.072 (0.022-0.16)	-0.066% (-0.47-0.18%)	-0.085% (-0.56-0.22%)	-0.065 (-0.46-0.17)	-0.63 (-4.5-1.8)
Timor-Leste	2000	Female	85 to 89	13% (5.7-23%)	87% (77-94%)	0.077 (0.024-0.18)	-0.11% (-0.52-0.13%)	-0.14% (-0.63-0.15%)	-0.067 (-0.3-0.071)	-0.46 (-2.2-0.53)
Timor-Leste	2000	Female	90 to 94	15% (6.3-27%)	85% (73-94%)	0.089 (0.024-0.21)	-0.15% (-0.61-0.11%)	-0.19% (-0.74-0.12%)	-0.036 (-0.14-0.023)	-0.16 (-0.67-0.12)
Timor-Leste	2000	Female	95 plus	18% (6.7-34%)	82% (66-93%)	0.11 (0.022-0.29)	-0.15% (-0.7-0.16%)	-0.21% (-0.88-0.17%)	-0.014 (-0.056-0.011)	-0.039 (-0.18-0.042)
Timor-Leste	2000	Male	All Ages	23% (18-30%)	77% (70-82%)	0.097 (0.085-0.11)	0.46% (0.19-0.74%)	0.56% (0.056-1.1%)	14 (1.4-28)	860 (360-1400)
Timor-Leste	2000	Male	15 to 19	22% (12-37%)	78% (63-88%)	0.054 (0.024-0.1)	0.65% (0.42-0.97%)	0.57% (0.28-1%)	0.36 (0.17-0.65)	53 (33-78)
Timor-Leste	2000	Male	20 to 24	34% (19-52%)	66% (48-81%)	0.13 (0.063-0.22)	1.3% (0.74-1.9%)	0.99% (0.29-1.8%)	0.53 (0.14-1)	71 (39-110)
Timor-Leste	2000	Male	25 to 29	39% (22-58%)	61% (42-78%)	0.18 (0.098-0.29)	1.5% (0.73-2.3%)	1.2% (0.087-2.5%)	0.75 (0.051-1.5)	92 (45-140)
Timor-Leste	2000	Male	30 to 34	43% (24-62%)	57% (38-76%)	0.22 (0.13-0.34)	1.7% (0.74-2.7%)	1.6% (0.12-3.2%)	1 (0.077-2)	97 (41-160)
Timor-Leste	2000	Male	35 to 39	45% (27-65%)	55% (35-73%)	0.26 (0.14-0.39)	1.8% (0.79-3%)	2% (0.21-4%)	1.5 (0.14-3.2)	120 (48-200)
Timor-Leste	2000	Male	40 to 44	44% (27-64%)	56% (36-73%)	0.27 (0.15-0.4)	1.8% (0.66-3%)	2.1% (0.23-4%)	1.7 (0.18-3.4)	110 (38-190)
Timor-Leste	2000	Male	45 to 49	43% (24-63%)	57% (37-76%)	0.26 (0.14-0.42)	1.7% (0.49-2.9%)	2.1% (0.24-4%)	1.9 (0.2-3.7)	99 (28-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	2000	Male	50 to 54	41% (23-61%)	59% (39-77%)	0.25 (0.13-0.4)	1.4% (0.27-2.5%)	1.7% (0.061-3.4%)	1.9 (0.065-3.8)	84 (16-160)
Timor-Leste	2000	Male	55 to 59	39% (22-59%)	61% (41-78%)	0.22 (0.11-0.37)	1.1% (0.21-2.1%)	1.3% (0.034-2.7%)	1.6 (0.035-3.3)	59 (9.7-110)
Timor-Leste	2000	Male	60 to 64	36% (20-56%)	64% (44-80%)	0.19 (0.094-0.34)	0.86% (0.023-1.7%)	0.98% (-0.12-2.1%)	1.3 (-0.18-2.8)	41 (1.2-82)
Timor-Leste	2000	Male	65 to 69	32% (18-51%)	68% (49-82%)	0.16 (0.079-0.27)	0.57% (-0.2-1.3%)	0.63% (-0.41-1.6%)	0.74 (-0.44-1.9)	20 (-6.1-45)
Timor-Leste	2000	Male	70 to 74	29% (16-45%)	71% (55-84%)	0.13 (0.06-0.24)	0.41% (-0.29-1%)	0.43% (-0.45-1.2%)	0.52 (-0.56-1.5)	11 (-8.1-29)
Timor-Leste	2000	Male	75 to 79	26% (15-43%)	74% (57-85%)	0.11 (0.048-0.21)	0.24% (-0.32-0.71%)	0.22% (-0.5-0.8%)	0.18 (-0.41-0.67)	3.4 (-4.8-10)
Timor-Leste	2000	Male	80 to 84	24% (12-40%)	76% (60-88%)	0.095 (0.041-0.17)	0.18% (-0.33-0.62%)	0.16% (-0.46-0.71%)	0.074 (-0.22-0.35)	1.1 (-2-3.9)
Timor-Leste	2000	Male	85 to 89	23% (12-37%)	77% (63-88%)	0.084 (0.036-0.16)	0.11% (-0.27-0.45%)	0.082% (-0.39-0.5%)	0.026 (-0.12-0.16)	0.31 (-0.78-1.3)
Timor-Leste	2000	Male	90 to 94	24% (13-41%)	76% (59-87%)	0.082 (0.034-0.16)	0.042% (-0.3-0.33%)	0.0032% (-0.43-0.35%)	0.00049 (-0.069-0.057)	0.041 (-0.3-0.32)
Timor-Leste	2000	Male	95 plus	28% (13-48%)	72% (52-87%)	0.084 (0.029-0.18)	0.064% (-0.34-0.35%)	-0.0078% (-0.51-0.34%)	-0.00042 (-0.027-0.018)	0.014 (-0.073-0.073)
Timor-Leste	2005	Female	All Ages	7.6% (5.4-10%)	92% (90-95%)	0.033 (0.023-0.046)	0.19% (0.11-0.27%)	0.14% (-0.03-0.32%)	3.6 (-0.73-7.9)	310 (180-440)
Timor-Leste	2005	Female	15 to 19	7.7% (3.5-15%)	92% (85-97%)	0.033 (0.011-0.071)	0.36% (0.22-0.52%)	0.21% (0.088-0.4%)	0.12 (0.046-0.23)	30 (17-44)
Timor-Leste	2005	Female	20 to 24	9.3% (4.1-18%)	91% (82-96%)	0.044 (0.015-0.096)	0.62% (0.38-0.94%)	0.19% (0.046-0.4%)	0.12 (0.028-0.27)	49 (29-77)
Timor-Leste	2005	Female	25 to 29	10% (4.3-19%)	90% (81-96%)	0.05 (0.017-0.11)	0.64% (0.43-0.93%)	0.22% (0.029-0.45%)	0.087 (0.012-0.18)	29 (18-44)
Timor-Leste	2005	Female	30 to 34	11% (4.4-21%)	89% (79-96%)	0.057 (0.02-0.12)	0.6% (0.39-0.84%)	0.3% (0.049-0.54%)	0.15 (0.026-0.27)	31 (19-46)
Timor-Leste	2005	Female	35 to 39	12% (5-22%)	88% (78-95%)	0.063 (0.022-0.12)	0.55% (0.37-0.74%)	0.39% (0.11-0.67%)	0.22 (0.066-0.39)	28 (18-40)
Timor-Leste	2005	Female	40 to 44	12% (5.4-22%)	88% (78-95%)	0.071 (0.025-0.15)	0.52% (0.32-0.73%)	0.49% (0.2-0.8%)	0.36 (0.13-0.59)	31 (18-44)
Timor-Leste	2005	Female	45 to 49	12% (5.4-22%)	88% (78-95%)	0.081 (0.027-0.18)	0.48% (0.27-0.71%)	0.51% (0.18-0.87%)	0.44 (0.16-0.79)	28 (15-42)
Timor-Leste	2005	Female	50 to 54	13% (5.5-23%)	87% (77-94%)	0.089 (0.032-0.19)	0.52% (0.22-0.85%)	0.62% (0.2-1.1%)	0.67 (0.21-1.3)	31 (12-52)
Timor-Leste	2005	Female	55 to 59	13% (5.9-24%)	87% (76-94%)	0.098 (0.033-0.22)	0.41% (0.052-0.73%)	0.5% (0.012-0.97%)	0.7 (0.017-1.3)	26 (3.3-46)
Timor-Leste	2005	Female	60 to 64	13% (5.7-25%)	87% (75-94%)	0.099 (0.032-0.22)	0.26% (-0.16-0.65%)	0.31% (-0.21-0.83%)	0.53 (-0.36-1.4)	16 (-9.8-39)
Timor-Leste	2005	Female	65 to 69	13% (5.6-23%)	87% (77-94%)	0.092 (0.03-0.19)	0.13% (-0.27-0.44%)	0.15% (-0.36-0.55%)	0.28 (-0.7-1)	7 (-16-24)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	2005	Female	70 to 74	12% (5.4-22%)	88% (78-95%)	0.083 (0.026-0.2)	0.09% (-0.27-0.37%)	0.11% (-0.34-0.46%)	0.19 (-0.62-0.83)	3.7 (-11-15)
Timor-Leste	2005	Female	75 to 79	11% (5.1-22%)	89% (78-95%)	0.077 (0.023-0.19)	-0.014% (-0.44-0.27%)	-0.023% (-0.55-0.32%)	-0.039 (-0.87-0.55)	-0.4 (-12-7.8)
Timor-Leste	2005	Female	80 to 84	11% (5.2-22%)	89% (78-95%)	0.078 (0.024-0.18)	-0.056% (-0.47-0.19%)	-0.074% (-0.57-0.22%)	-0.08 (-0.61-0.24)	-0.76 (-6.2-2.6)
Timor-Leste	2005	Female	85 to 89	12% (5.5-23%)	88% (77-95%)	0.084 (0.025-0.2)	-0.11% (-0.52-0.13%)	-0.14% (-0.64-0.15%)	-0.089 (-0.39-0.094)	-0.61 (-2.8-0.72)
Timor-Leste	2005	Female	90 to 94	14% (6.3-27%)	86% (73-94%)	0.099 (0.026-0.24)	-0.15% (-0.57-0.11%)	-0.2% (-0.71-0.13%)	-0.05 (-0.18-0.033)	-0.23 (-0.87-0.18)
Timor-Leste	2005	Female	95 plus	18% (6.9-36%)	82% (64-93%)	0.12 (0.024-0.35)	-0.16% (-0.69-0.22%)	-0.22% (-0.88-0.23%)	-0.015 (-0.059-0.015)	-0.042 (-0.19-0.06)
Timor-Leste	2005	Male	All Ages	23% (18-30%)	77% (70-82%)	0.1 (0.083-0.13)	0.56% (0.27-0.87%)	0.66% (0.15-1.2%)	17 (3.8-31)	1000 (510-1600)
Timor-Leste	2005	Male	15 to 19	22% (12-36%)	78% (64-88%)	0.058 (0.027-0.11)	0.74% (0.47-1.1%)	0.7% (0.35-1.2%)	0.43 (0.21-0.77)	62 (38-94)
Timor-Leste	2005	Male	20 to 24	33% (18-51%)	67% (49-82%)	0.14 (0.067-0.23)	1.5% (0.88-2.2%)	1.3% (0.38-2.3%)	0.92 (0.28-1.7)	130 (70-190)
Timor-Leste	2005	Male	25 to 29	38% (21-56%)	62% (44-79%)	0.19 (0.1-0.33)	1.7% (0.94-2.6%)	1.5% (0.18-3%)	0.71 (0.081-1.4)	86 (46-130)
Timor-Leste	2005	Male	30 to 34	42% (23-61%)	58% (39-77%)	0.24 (0.13-0.4)	1.9% (0.92-2.9%)	1.9% (0.21-3.6%)	1.1 (0.13-2.2)	110 (52-170)
Timor-Leste	2005	Male	35 to 39	44% (26-64%)	56% (36-74%)	0.28 (0.15-0.43)	2% (0.83-3.2%)	2.3% (0.29-4.4%)	1.5 (0.19-3)	110 (49-190)
Timor-Leste	2005	Male	40 to 44	44% (26-64%)	56% (36-74%)	0.29 (0.15-0.47)	1.9% (0.73-3%)	2.3% (0.3-4.2%)	2 (0.25-3.7)	120 (47-210)
Timor-Leste	2005	Male	45 to 49	42% (24-62%)	58% (38-76%)	0.29 (0.14-0.48)	1.8% (0.65-3%)	2.3% (0.43-4.2%)	2.2 (0.4-4.2)	120 (40-200)
Timor-Leste	2005	Male	50 to 54	40% (22-60%)	60% (40-78%)	0.27 (0.14-0.44)	1.5% (0.43-2.5%)	1.9% (0.27-3.4%)	2.2 (0.34-4.1)	94 (27-160)
Timor-Leste	2005	Male	55 to 59	38% (21-57%)	62% (43-79%)	0.24 (0.11-0.41)	1.1% (0.33-2.1%)	1.4% (0.19-2.7%)	2 (0.26-3.9)	75 (21-140)
Timor-Leste	2005	Male	60 to 64	35% (19-54%)	65% (46-81%)	0.21 (0.096-0.37)	0.9% (0.12-1.6%)	1% (0.0027-2.1%)	1.7 (0.0044-3.5)	54 (6.6-100)
Timor-Leste	2005	Male	65 to 69	32% (17-49%)	68% (51-83%)	0.17 (0.083-0.3)	0.59% (-0.037-1.2%)	0.66% (-0.18-1.5%)	1.1 (-0.3-2.4)	28 (-1.8-57)
Timor-Leste	2005	Male	70 to 74	28% (15-45%)	72% (55-85%)	0.14 (0.063-0.26)	0.43% (-0.18-0.95%)	0.47% (-0.33-1.2%)	0.66 (-0.47-1.6)	14 (-6.1-31)
Timor-Leste	2005	Male	75 to 79	26% (14-42%)	74% (58-86%)	0.12 (0.05-0.23)	0.25% (-0.27-0.67%)	0.24% (-0.41-0.76%)	0.3 (-0.49-0.92)	5.4 (-5.7-14)
Timor-Leste	2005	Male	80 to 84	24% (13-39%)	76% (61-87%)	0.1 (0.043-0.2)	0.19% (-0.24-0.59%)	0.17% (-0.36-0.66%)	0.12 (-0.27-0.48)	1.7 (-2.2-5.4)
Timor-Leste	2005	Male	85 to 89	22% (11-36%)	78% (64-89%)	0.089 (0.037-0.17)	0.11% (-0.24-0.41%)	0.086% (-0.34-0.45%)	0.038 (-0.15-0.2)	0.44 (-0.95-1.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	2005	Male	90 to 94	23% (12-39%)	77% (61-88%)	0.086 (0.035-0.18)	0.044% (-0.3-0.31%)	0.0062% (-0.42-0.32%)	0.0012 (-0.085-0.067)	0.056 (-0.38-0.38)
Timor-Leste	2005	Male	95 plus	27% (12-46%)	73% (54-88%)	0.088 (0.029-0.2)	0.065% (-0.34-0.33%)	-0.0099% (-0.51-0.32%)	-0.00061 (-0.032-0.019)	0.016 (-0.085-0.081)
Timor-Leste	2010	Female	All Ages	7.6% (5.5-10%)	92% (90-94%)	0.037 (0.024-0.056)	0.21% (0.12-0.31%)	0.14% (-0.051-0.32%)	3.8 (-1.3-8.8)	340 (200-500)
Timor-Leste	2010	Female	15 to 19	7.5% (3.4-14%)	92% (86-97%)	0.034 (0.011-0.081)	0.4% (0.25-0.61%)	0.22% (0.1-0.41%)	0.09 (0.037-0.17)	30 (17-47)
Timor-Leste	2010	Female	20 to 24	9.1% (3.8-17%)	91% (83-96%)	0.046 (0.015-0.1)	0.76% (0.47-1.1%)	0.21% (0.083-0.42%)	0.093 (0.035-0.2)	51 (31-79)
Timor-Leste	2010	Female	25 to 29	9.9% (4.2-19%)	90% (81-96%)	0.052 (0.018-0.12)	0.75% (0.5-1.1%)	0.23% (0.039-0.43%)	0.099 (0.017-0.19)	45 (28-68)
Timor-Leste	2010	Female	30 to 34	11% (4.4-21%)	89% (79-96%)	0.06 (0.02-0.13)	0.69% (0.46-0.97%)	0.33% (0.053-0.54%)	0.12 (0.02-0.21)	31 (20-48)
Timor-Leste	2010	Female	35 to 39	12% (4.9-22%)	88% (78-95%)	0.065 (0.023-0.13)	0.6% (0.4-0.83%)	0.42% (0.071-0.73%)	0.22 (0.037-0.4)	34 (21-49)
Timor-Leste	2010	Female	40 to 44	12% (5.5-22%)	88% (78-95%)	0.073 (0.025-0.15)	0.56% (0.37-0.77%)	0.53% (0.24-0.86%)	0.33 (0.14-0.56)	32 (20-46)
Timor-Leste	2010	Female	45 to 49	12% (5.2-23%)	88% (77-95%)	0.083 (0.026-0.18)	0.51% (0.3-0.74%)	0.55% (0.23-0.93%)	0.41 (0.16-0.72)	28 (16-42)
Timor-Leste	2010	Female	50 to 54	12% (5.3-24%)	88% (76-95%)	0.093 (0.031-0.21)	0.52% (0.24-0.83%)	0.65% (0.24-1.2%)	0.56 (0.19-1)	27 (12-46)
Timor-Leste	2010	Female	55 to 59	13% (5.7-23%)	87% (77-94%)	0.1 (0.034-0.23)	0.41% (0.098-0.71%)	0.52% (0.086-1%)	0.49 (0.085-0.94)	19 (4.4-34)
Timor-Leste	2010	Female	60 to 64	13% (5.4-23%)	87% (77-95%)	0.1 (0.033-0.23)	0.27% (-0.1-0.63%)	0.34% (-0.15-0.83%)	0.93 (-0.43-2.3)	29 (-11-69)
Timor-Leste	2010	Female	65 to 69	12% (5.3-22%)	88% (78-95%)	0.095 (0.031-0.2)	0.14% (-0.24-0.44%)	0.18% (-0.34-0.59%)	0.44 (-0.83-1.6)	11 (-18-37)
Timor-Leste	2010	Female	70 to 74	12% (5.5-21%)	88% (79-95%)	0.086 (0.028-0.2)	0.11% (-0.24-0.38%)	0.13% (-0.31-0.5%)	0.31 (-0.73-1.2)	6 (-14-22)
Timor-Leste	2010	Female	75 to 79	11% (5-22%)	89% (78-95%)	0.082 (0.023-0.2)	0.0032% (-0.39-0.29%)	-0.0022% (-0.48-0.36%)	-0.0044 (-1-0.78)	0.14 (-15-12)
Timor-Leste	2010	Female	80 to 84	11% (5-22%)	89% (78-95%)	0.082 (0.024-0.2)	-0.042% (-0.42-0.21%)	-0.058% (-0.52-0.24%)	-0.09 (-0.8-0.38)	-0.84 (-8.5-4.2)
Timor-Leste	2010	Female	85 to 89	12% (5.3-23%)	88% (77-95%)	0.089 (0.026-0.22)	-0.1% (-0.5-0.15%)	-0.14% (-0.62-0.17%)	-0.12 (-0.56-0.15)	-0.82 (-4.1-1.1)
Timor-Leste	2010	Female	90 to 94	14% (6-27%)	86% (73-94%)	0.1 (0.025-0.26)	-0.15% (-0.56-0.16%)	-0.2% (-0.72-0.17%)	-0.069 (-0.26-0.057)	-0.31 (-1.2-0.34)
Timor-Leste	2010	Female	95 plus	18% (6.9-36%)	82% (64-93%)	0.13 (0.024-0.37)	-0.16% (-0.68-0.29%)	-0.23% (-0.89-0.28%)	-0.022 (-0.088-0.027)	-0.061 (-0.28-0.11)
Timor-Leste	2010	Male	All Ages	23% (18-29%)	77% (71-82%)	0.11 (0.078-0.16)	0.62% (0.35-0.94%)	0.71% (0.24-1.2%)	20 (6.9-37)	1200 (670-1800)
Timor-Leste	2010	Male	15 to 19	21% (12-34%)	79% (66-88%)	0.059 (0.026-0.12)	0.79% (0.52-1.1%)	0.74% (0.36-1.3%)	0.4 (0.2-0.74)	64 (39-96)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	2010	Male	20 to 24	32% (17-49%)	68% (51-83%)	0.14 (0.066-0.25)	1.7% (1.1-2.3%)	1.4% (0.54-2.5%)	0.87 (0.33-1.6)	130 (82-190)
Timor-Leste	2010	Male	25 to 29	37% (21-55%)	63% (45-79%)	0.2 (0.1-0.36)	1.9% (1.1-2.7%)	1.6% (0.22-3%)	0.98 (0.13-1.9)	130 (76-200)
Timor-Leste	2010	Male	30 to 34	41% (23-61%)	59% (39-77%)	0.25 (0.13-0.44)	2% (1.1-3%)	2.1% (0.27-4%)	1 (0.13-2)	110 (54-160)
Timor-Leste	2010	Male	35 to 39	43% (26-62%)	57% (38-74%)	0.29 (0.14-0.48)	2.1% (0.98-3.4%)	2.6% (0.29-4.8%)	1.7 (0.18-3.5)	140 (62-230)
Timor-Leste	2010	Male	40 to 44	43% (26-63%)	57% (37-74%)	0.3 (0.15-0.51)	2% (0.88-3.2%)	2.7% (0.38-4.7%)	2 (0.29-3.9)	130 (55-220)
Timor-Leste	2010	Male	45 to 49	42% (23-61%)	58% (39-77%)	0.3 (0.14-0.53)	1.9% (0.75-3.2%)	2.6% (0.52-4.8%)	2.3 (0.46-4.4)	120 (46-210)
Timor-Leste	2010	Male	50 to 54	39% (22-58%)	61% (42-78%)	0.28 (0.13-0.49)	1.6% (0.61-2.6%)	2.1% (0.56-3.7%)	2.1 (0.53-3.9)	92 (35-160)
Timor-Leste	2010	Male	55 to 59	37% (21-56%)	63% (44-79%)	0.25 (0.11-0.44)	1.2% (0.55-2.1%)	1.6% (0.46-2.8%)	1.6 (0.45-3)	62 (26-110)
Timor-Leste	2010	Male	60 to 64	34% (18-54%)	66% (46-82%)	0.21 (0.096-0.4)	0.97% (0.32-1.6%)	1.2% (0.24-2.1%)	3.5 (0.73-6.8)	110 (35-200)
Timor-Leste	2010	Male	65 to 69	31% (17-48%)	69% (52-83%)	0.17 (0.079-0.32)	0.66% (0.14-1.2%)	0.76% (0.011-1.5%)	1.9 (0.024-4)	52 (11-99)
Timor-Leste	2010	Male	70 to 74	27% (15-44%)	73% (56-85%)	0.14 (0.059-0.29)	0.49% (0.046-0.95%)	0.56% (-0.04-1.2%)	1.2 (-0.083-2.6)	25 (2.3-51)
Timor-Leste	2010	Male	75 to 79	25% (14-40%)	75% (60-86%)	0.12 (0.048-0.24)	0.29% (-0.14-0.64%)	0.29% (-0.27-0.75%)	0.54 (-0.47-1.4)	9.8 (-4.5-22)
Timor-Leste	2010	Male	80 to 84	23% (12-38%)	77% (62-88%)	0.1 (0.042-0.21)	0.22% (-0.17-0.53%)	0.2% (-0.27-0.62%)	0.24 (-0.33-0.74)	3.3 (-2.7-8.3)
Timor-Leste	2010	Male	85 to 89	21% (11-35%)	79% (65-89%)	0.09 (0.036-0.18)	0.13% (-0.18-0.38%)	0.1% (-0.28-0.42%)	0.067 (-0.18-0.27)	0.75 (-1.1-2.3)
Timor-Leste	2010	Male	90 to 94	22% (12-38%)	78% (62-88%)	0.087 (0.034-0.18)	0.054% (-0.27-0.28%)	0.016% (-0.38-0.3%)	0.0044 (-0.11-0.079)	0.093 (-0.46-0.48)
Timor-Leste	2010	Male	95 plus	26% (11-45%)	74% (55-89%)	0.09 (0.026-0.2)	0.075% (-0.3-0.31%)	-0.0043% (-0.48-0.29%)	-0.00033 (-0.039-0.024)	0.025 (-0.099-0.1)
Timor-Leste	2016	Female	All Ages	7.8% (5.6-10%)	92% (90-94%)	0.04 (0.023-0.064)	0.26% (0.15-0.38%)	0.15% (-0.078-0.38%)	4.1 (-2.1-11)	380 (220-570)
Timor-Leste	2016	Female	15 to 19	7.6% (3.2-14%)	92% (86-97%)	0.036 (0.01-0.083)	0.44% (0.26-0.68%)	0.25% (0.11-0.46%)	0.091 (0.034-0.18)	32 (18-53)
Timor-Leste	2016	Female	20 to 24	9.1% (3.8-17%)	91% (83-96%)	0.047 (0.015-0.11)	0.84% (0.52-1.3%)	0.23% (0.089-0.44%)	0.091 (0.032-0.19)	58 (34-92)
Timor-Leste	2016	Female	25 to 29	9.8% (4.1-18%)	90% (82-96%)	0.054 (0.018-0.12)	0.83% (0.55-1.1%)	0.25% (0.033-0.47%)	0.097 (0.013-0.21)	50 (32-76)
Timor-Leste	2016	Female	30 to 34	11% (4.4-20%)	89% (80-96%)	0.061 (0.019-0.14)	0.75% (0.5-1.1%)	0.35% (0.045-0.6%)	0.15 (0.019-0.28)	44 (27-65)
Timor-Leste	2016	Female	35 to 39	12% (5-22%)	88% (78-95%)	0.069 (0.021-0.15)	0.66% (0.45-0.9%)	0.47% (0.11-0.79%)	0.19 (0.053-0.36)	30 (20-44)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	2016	Female	40 to 44	12% (5.5-22%)	88% (78-94%)	0.078 (0.025-0.18)	0.6% (0.39-0.83%)	0.59% (0.21-0.96%)	0.36 (0.11-0.62)	35 (21-51)
Timor-Leste	2016	Female	45 to 49	12% (5.3-22%)	88% (78-95%)	0.088 (0.028-0.2)	0.55% (0.35-0.79%)	0.62% (0.28-1%)	0.51 (0.2-0.91)	34 (20-52)
Timor-Leste	2016	Female	50 to 54	13% (5.6-25%)	87% (75-94%)	0.099 (0.031-0.23)	0.55% (0.27-0.88%)	0.69% (0.28-1.2%)	0.7 (0.26-1.3)	34 (15-57)
Timor-Leste	2016	Female	55 to 59	13% (5.5-25%)	87% (75-95%)	0.11 (0.033-0.26)	0.42% (0.093-0.78%)	0.54% (0.063-1.1%)	0.65 (0.061-1.4)	25 (5-49)
Timor-Leste	2016	Female	60 to 64	13% (5.6-25%)	87% (75-94%)	0.11 (0.032-0.27)	0.28% (-0.095-0.65%)	0.35% (-0.14-0.87%)	0.42 (-0.16-1.1)	13 (-4.8-32)
Timor-Leste	2016	Female	65 to 69	13% (5.6-23%)	87% (77-94%)	0.1 (0.031-0.23)	0.15% (-0.25-0.51%)	0.19% (-0.35-0.65%)	0.66 (-1.3-2.3)	17 (-28-55)
Timor-Leste	2016	Female	70 to 74	12% (5.6-22%)	88% (78-94%)	0.095 (0.027-0.23)	0.12% (-0.24-0.42%)	0.15% (-0.31-0.54%)	0.53 (-1.2-2.1)	10 (-22-39)
Timor-Leste	2016	Female	75 to 79	12% (5-22%)	88% (78-95%)	0.089 (0.024-0.22)	0.0087% (-0.37-0.3%)	0.0034% (-0.48-0.39%)	0.0094 (-1.2-1)	0.44 (-17-15)
Timor-Leste	2016	Female	80 to 84	12% (5-22%)	88% (78-95%)	0.089 (0.024-0.23)	-0.035% (-0.42-0.23%)	-0.051% (-0.53-0.29%)	-0.1 (-1.1-0.57)	-0.93 (-11-6.1)
Timor-Leste	2016	Female	85 to 89	13% (5.4-23%)	87% (77-95%)	0.096 (0.026-0.26)	-0.097% (-0.5-0.21%)	-0.13% (-0.63-0.22%)	-0.18 (-0.89-0.32)	-1.2 (-6.5-2.6)
Timor-Leste	2016	Female	90 to 94	15% (6-27%)	85% (73-94%)	0.11 (0.025-0.29)	-0.14% (-0.57-0.21%)	-0.19% (-0.73-0.22%)	-0.1 (-0.39-0.13)	-0.46 (-1.9-0.7)
Timor-Leste	2016	Female	95 plus	18% (6.8-35%)	82% (65-93%)	0.14 (0.026-0.41)	-0.15% (-0.69-0.32%)	-0.22% (-0.9-0.33%)	-0.029 (-0.12-0.047)	-0.079 (-0.38-0.18)
Timor-Leste	2016	Male	All Ages	23% (18-30%)	77% (70-82%)	0.12 (0.07-0.18)	0.79% (0.45-1.2%)	0.83% (0.31-1.4%)	24 (8.8-43)	1400 (780-2000)
Timor-Leste	2016	Male	15 to 19	21% (12-34%)	79% (66-88%)	0.062 (0.025-0.13)	0.83% (0.53-1.2%)	0.81% (0.41-1.5%)	0.46 (0.21-0.85)	73 (45-110)
Timor-Leste	2016	Male	20 to 24	32% (18-50%)	68% (50-82%)	0.15 (0.063-0.29)	1.7% (1.1-2.5%)	1.5% (0.62-2.8%)	1.1 (0.4-2)	150 (94-230)
Timor-Leste	2016	Male	25 to 29	38% (21-56%)	62% (44-79%)	0.21 (0.093-0.38)	2% (1.2-2.9%)	1.8% (0.25-3.3%)	1.2 (0.15-2.4)	150 (83-230)
Timor-Leste	2016	Male	30 to 34	41% (22-61%)	59% (39-78%)	0.26 (0.12-0.47)	2.2% (1.2-3.2%)	2.2% (0.27-4.2%)	1.5 (0.17-3)	150 (76-240)
Timor-Leste	2016	Male	35 to 39	43% (26-62%)	57% (38-74%)	0.3 (0.13-0.54)	2.3% (1.1-3.6%)	2.8% (0.35-5%)	1.6 (0.17-3.2)	120 (57-200)
Timor-Leste	2016	Male	40 to 44	43% (25-63%)	57% (37-75%)	0.31 (0.14-0.58)	2.2% (0.91-3.5%)	2.8% (0.34-4.9%)	2.3 (0.26-4.3)	140 (58-240)
Timor-Leste	2016	Male	45 to 49	42% (24-60%)	58% (40-76%)	0.31 (0.14-0.57)	2.1% (0.93-3.3%)	2.8% (0.77-4.9%)	2.9 (0.74-5.4)	150 (64-250)
Timor-Leste	2016	Male	50 to 54	40% (22-60%)	60% (40-78%)	0.29 (0.12-0.55)	1.7% (0.77-2.7%)	2.3% (0.8-3.8%)	2.7 (0.84-5)	120 (54-210)
Timor-Leste	2016	Male	55 to 59	37% (21-57%)	63% (43-79%)	0.26 (0.11-0.51)	1.3% (0.59-2.1%)	1.7% (0.56-2.9%)	2.4 (0.76-4.4)	90 (39-160)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Timor-Leste	2016	Male	60 to 64	34% (18-54%)	66% (46-82%)	0.22 (0.09-0.44)	1% (0.41-1.7%)	1.3% (0.36-2.2%)	1.7 (0.48-3.3)	55 (21-97)
Timor-Leste	2016	Male	65 to 69	31% (17-49%)	69% (51-83%)	0.19 (0.074-0.36)	0.71% (0.18-1.3%)	0.83% (0.085-1.6%)	3 (0.34-6.1)	80 (20-150)
Timor-Leste	2016	Male	70 to 74	28% (15-45%)	72% (55-85%)	0.15 (0.06-0.31)	0.53% (0.083-1%)	0.61% (-0.015-1.2%)	2.1 (-0.048-4.4)	45 (6.9-86)
Timor-Leste	2016	Male	75 to 79	25% (13-41%)	75% (59-87%)	0.13 (0.046-0.26)	0.33% (-0.12-0.68%)	0.34% (-0.23-0.8%)	0.76 (-0.53-1.8)	13 (-5-29)
Timor-Leste	2016	Male	80 to 84	23% (11-38%)	77% (62-89%)	0.11 (0.039-0.24)	0.24% (-0.14-0.55%)	0.23% (-0.25-0.62%)	0.35 (-0.37-0.96)	4.7 (-2.7-11)
Timor-Leste	2016	Male	85 to 89	21% (11-35%)	79% (65-89%)	0.094 (0.034-0.2)	0.14% (-0.18-0.38%)	0.12% (-0.27-0.41%)	0.13 (-0.3-0.45)	1.4 (-1.8-3.7)
Timor-Leste	2016	Male	90 to 94	22% (12-37%)	78% (63-88%)	0.092 (0.033-0.2)	0.061% (-0.27-0.29%)	0.023% (-0.39-0.3%)	0.0091 (-0.15-0.12)	0.15 (-0.67-0.68)
Timor-Leste	2016	Male	95 plus	26% (12-45%)	74% (55-88%)	0.095 (0.027-0.21)	0.084% (-0.32-0.32%)	0.0064% (-0.5-0.3%)	0.00064 (-0.047-0.029)	0.033 (-0.12-0.13)
Vietnam	1990	Female	All Ages	9.3% (7.8-11%)	91% (89-92%)	0.13 (0.084-0.18)	0.86% (0.5-1.3%)	0.8% (0.074-1.9%)	2000 (180-4700)	90000 (51000-140000)
Vietnam	1990	Female	15 to 19	9.3% (5.8-14%)	91% (86-94%)	0.13 (0.062-0.23)	1.1% (0.7-1.5%)	0.8% (0.26-1.6%)	22 (6.9-42)	5200 (3200-7800)
Vietnam	1990	Female	20 to 24	11% (6.6-15%)	89% (85-93%)	0.15 (0.075-0.28)	2.2% (1.4-3.2%)	1% (0.23-2.3%)	33 (6.9-74)	11000 (6400-17000)
Vietnam	1990	Female	25 to 29	11% (7-17%)	89% (83-93%)	0.16 (0.081-0.29)	2.2% (1.5-3.2%)	1.1% (0.15-2.6%)	40 (5-97)	11000 (6600-17000)
Vietnam	1990	Female	30 to 34	12% (7.4-18%)	88% (82-93%)	0.18 (0.089-0.31)	1.9% (1.2-2.7%)	1.1% (0.31-2.3%)	44 (12-93)	8900 (5400-14000)
Vietnam	1990	Female	35 to 39	12% (7.6-18%)	88% (82-92%)	0.2 (0.098-0.35)	1.7% (1.1-2.6%)	1.4% (0.44-2.8%)	56 (18-120)	7300 (4300-11000)
Vietnam	1990	Female	40 to 44	12% (7.7-17%)	88% (83-92%)	0.21 (0.11-0.37)	1.6% (1-2.5%)	1.6% (0.7-2.9%)	60 (27-110)	5400 (3200-8200)
Vietnam	1990	Female	45 to 49	12% (7.1-17%)	88% (83-93%)	0.23 (0.11-0.44)	1.5% (0.95-2.5%)	1.6% (0.7-3.2%)	76 (32-150)	5000 (2900-8400)
Vietnam	1990	Female	50 to 54	12% (8.1-18%)	88% (82-92%)	0.26 (0.13-0.47)	1.4% (0.78-2.4%)	1.6% (0.63-3.1%)	110 (44-230)	5700 (3000-10000)
Vietnam	1990	Female	55 to 59	13% (8.3-19%)	87% (81-92%)	0.29 (0.15-0.52)	1.3% (0.59-2.4%)	1.5% (0.51-3.1%)	170 (53-360)	6700 (2900-13000)
Vietnam	1990	Female	60 to 64	13% (7.9-19%)	87% (81-92%)	0.29 (0.14-0.51)	1.2% (0.4-2.4%)	1.5% (0.37-3.1%)	210 (50-460)	6600 (2100-14000)
Vietnam	1990	Female	65 to 69	11% (7.2-16%)	89% (84-93%)	0.26 (0.13-0.44)	0.92% (0.16-2.1%)	1.1% (0.11-2.6%)	220 (23-540)	5500 (970-12000)
Vietnam	1990	Female	70 to 74	11% (7.1-16%)	89% (84-93%)	0.25 (0.13-0.45)	0.78% (0.007-2.1%)	0.93% (-0.06-2.5%)	230 (-12-640)	4400 (30-12000)
Vietnam	1990	Female	75 to 79	11% (7.2-17%)	89% (83-93%)	0.25 (0.12-0.46)	0.61% (-0.19-2%)	0.71% (-0.27-2.4%)	220 (-78-750)	3300 (-950-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	1990	Female	80 to 84	12% (7.5-17%)	88% (83-93%)	0.27 (0.13-0.46)	0.65% (-0.2-2%)	0.77% (-0.28-2.4%)	260 (-94-820)	2800 (-840-8400)
Vietnam	1990	Female	85 to 89	13% (8.2-20%)	87% (80-92%)	0.3 (0.14-0.53)	0.55% (-0.27-1.8%)	0.64% (-0.34-2.2%)	170 (-90-600)	1300 (-590-4300)
Vietnam	1990	Female	90 to 94	16% (9.7-24%)	84% (76-90%)	0.34 (0.16-0.62)	0.41% (-0.44-1.8%)	0.44% (-0.58-2.1%)	60 (-77-280)	330 (-350-1400)
Vietnam	1990	Female	95 plus	19% (10-31%)	81% (69-90%)	0.41 (0.16-0.84)	0.4% (-0.66-2.1%)	0.38% (-0.88-2.4%)	16 (-37-100)	67 (-110-360)
Vietnam	1990	Male	All Ages	43% (37-49%)	57% (51-63%)	0.75 (0.52-1)	3.4% (2-5%)	4.2% (1.9-6.6%)	11000 (4900-17000)	420000 (250000-640000)
Vietnam	1990	Male	15 to 19	35% (25-46%)	65% (54-75%)	0.39 (0.2-0.61)	2.2% (1.1-3.7%)	2.8% (1.1-5%)	150 (54-280)	14000 (6900-23000)
Vietnam	1990	Male	20 to 24	52% (40-63%)	48% (37-60%)	0.89 (0.53-1.4)	5.2% (3-7.9%)	6% (2.9-9.7%)	450 (200-770)	38000 (21000-60000)
Vietnam	1990	Male	25 to 29	62% (51-71%)	38% (29-49%)	1.2 (0.73-1.7)	6.6% (3.6-10%)	8% (3.6-13%)	630 (280-1100)	46000 (24000-73000)
Vietnam	1990	Male	30 to 34	70% (58-78%)	30% (22-42%)	1.5 (0.97-2.1)	7.7% (4.4-12%)	9.7% (5-15%)	770 (370-1200)	50000 (27000-78000)
Vietnam	1990	Male	35 to 39	72% (62-79%)	28% (21-38%)	1.8 (1.2-2.7)	8.8% (5.2-13%)	11% (6.3-17%)	880 (460-1400)	50000 (28000-76000)
Vietnam	1990	Male	40 to 44	71% (61-79%)	29% (21-39%)	1.9 (1.2-2.8)	8.7% (5.3-12%)	11% (6.5-16%)	700 (390-1100)	35000 (20000-53000)
Vietnam	1990	Male	45 to 49	66% (53-75%)	34% (25-47%)	1.7 (1.1-2.5)	8% (4.8-12%)	10% (5.7-15%)	730 (410-1100)	32000 (18000-49000)
Vietnam	1990	Male	50 to 54	63% (51-72%)	37% (28-49%)	1.6 (0.99-2.4)	7.2% (4.1-11%)	8.9% (4.9-13%)	900 (480-1400)	34000 (19000-53000)
Vietnam	1990	Male	55 to 59	61% (48-71%)	39% (29-52%)	1.6 (1-2.4)	6.6% (3.3-10%)	8.1% (4-13%)	1200 (580-1900)	39000 (20000-63000)
Vietnam	1990	Male	60 to 64	57% (44-69%)	43% (31-56%)	1.5 (0.87-2.3)	5.2% (2.2-8.7%)	6.2% (2.5-10%)	1100 (440-1900)	31000 (13000-53000)
Vietnam	1990	Male	65 to 69	53% (41-65%)	47% (35-59%)	1.3 (0.74-1.9)	3.9% (1.4-6.9%)	4.5% (1.5-8.1%)	990 (320-1800)	23000 (8000-41000)
Vietnam	1990	Male	70 to 74	50% (38-63%)	50% (37-62%)	1.1 (0.61-1.8)	3.3% (0.67-6.5%)	3.8% (0.72-7.7%)	890 (150-1800)	16000 (3000-33000)
Vietnam	1990	Male	75 to 79	49% (35-64%)	51% (36-65%)	0.98 (0.51-1.7)	2.3% (-0.14-5.4%)	2.6% (-0.25-6.3%)	600 (-58-1500)	8700 (-500-21000)
Vietnam	1990	Male	80 to 84	46% (30-61%)	54% (39-70%)	0.82 (0.44-1.4)	1.9% (-0.36-4.8%)	2.2% (-0.5-5.6%)	400 (-91-1000)	4200 (-840-11000)
Vietnam	1990	Male	85 to 89	42% (28-57%)	58% (43-72%)	0.67 (0.35-1.2)	1% (-0.68-3.1%)	1.1% (-0.88-3.6%)	130 (-99-430)	1000 (-660-3200)
Vietnam	1990	Male	90 to 94	41% (28-56%)	59% (44-72%)	0.58 (0.29-1)	0.36% (-0.98-2.1%)	0.33% (-1.3-2.3%)	16 (-62-120)	100 (-280-610)
Vietnam	1990	Male	95 plus	43% (27-61%)	57% (39-73%)	0.52 (0.22-1)	0.075% (-1.2-1.5%)	-0.084% (-1.6-1.7%)	-0.99 (-19-20)	3.6 (-55-75)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	1995	Female	All Ages	9.6% (8-11%)	90% (89-92%)	0.19 (0.13-0.26)	1.3% (0.79-2%)	1.5% (0.47-3%)	3900 (1200-7900)	130000 (78000-200000)
Vietnam	1995	Female	15 to 19	9.3% (5.8-14%)	91% (86-94%)	0.19 (0.088-0.33)	1.2% (0.82-1.8%)	1.1% (0.45-2.1%)	26 (11-50)	5800 (3700-8600)
Vietnam	1995	Female	20 to 24	11% (6.5-15%)	89% (85-93%)	0.22 (0.11-0.39)	2.6% (1.7-3.7%)	1.5% (0.52-3.1%)	43 (14-88)	12000 (7700-19000)
Vietnam	1995	Female	25 to 29	11% (7.2-17%)	89% (83-93%)	0.23 (0.12-0.41)	2.6% (1.8-3.7%)	1.8% (0.6-3.6%)	56 (17-120)	13000 (7800-19000)
Vietnam	1995	Female	30 to 34	12% (7.3-19%)	88% (81-93%)	0.25 (0.13-0.44)	2.3% (1.5-3.2%)	1.7% (0.74-3.3%)	64 (25-120)	11000 (7000-16000)
Vietnam	1995	Female	35 to 39	12% (7.5-18%)	88% (82-93%)	0.28 (0.14-0.47)	2.1% (1.4-3.1%)	2.1% (0.91-3.8%)	97 (40-190)	11000 (6600-16000)
Vietnam	1995	Female	40 to 44	12% (7.9-17%)	88% (83-92%)	0.3 (0.16-0.5)	2% (1.3-3%)	2.2% (1.1-4%)	120 (55-210)	9500 (5800-14000)
Vietnam	1995	Female	45 to 49	12% (7-17%)	88% (83-93%)	0.33 (0.16-0.59)	2% (1.2-3.2%)	2.3% (1.1-4.3%)	120 (51-230)	7400 (4200-12000)
Vietnam	1995	Female	50 to 54	13% (8.2-19%)	87% (81-92%)	0.37 (0.19-0.65)	2% (1.1-3.4%)	2.4% (1-4.6%)	160 (60-290)	7400 (3900-13000)
Vietnam	1995	Female	55 to 59	13% (8.3-19%)	87% (81-92%)	0.42 (0.22-0.72)	1.9% (0.93-3.5%)	2.4% (0.97-4.7%)	230 (86-450)	8600 (4000-16000)
Vietnam	1995	Female	60 to 64	13% (8.1-19%)	87% (81-92%)	0.41 (0.2-0.74)	1.9% (0.79-3.6%)	2.4% (0.86-4.6%)	370 (130-740)	11000 (4500-21000)
Vietnam	1995	Female	65 to 69	11% (7.2-16%)	89% (84-93%)	0.37 (0.19-0.61)	1.6% (0.56-3.1%)	1.9% (0.6-4%)	390 (110-810)	9500 (3100-19000)
Vietnam	1995	Female	70 to 74	11% (7-15%)	89% (85-93%)	0.35 (0.18-0.62)	1.4% (0.36-3.1%)	1.7% (0.41-3.9%)	470 (100-1100)	9000 (2200-21000)
Vietnam	1995	Female	75 to 79	11% (7.2-17%)	89% (83-93%)	0.36 (0.18-0.65)	1.2% (0.15-3%)	1.5% (0.16-3.6%)	470 (49-1200)	6800 (860-17000)
Vietnam	1995	Female	80 to 84	12% (7.7-17%)	88% (83-92%)	0.38 (0.19-0.64)	1.3% (0.17-3.1%)	1.6% (0.19-3.7%)	610 (69-1500)	6400 (780-15000)
Vietnam	1995	Female	85 to 89	13% (8.1-20%)	87% (80-92%)	0.42 (0.21-0.73)	1.2% (0.069-2.8%)	1.4% (0.047-3.4%)	460 (15-1100)	3400 (190-8300)
Vietnam	1995	Female	90 to 94	16% (9.7-23%)	84% (77-90%)	0.48 (0.23-0.89)	1% (-0.14-2.9%)	1.2% (-0.21-3.5%)	200 (-37-580)	1100 (-140-3000)
Vietnam	1995	Female	95 plus	19% (10-31%)	81% (69-90%)	0.57 (0.22-1.2)	1.1% (-0.33-3.4%)	1.2% (-0.49-3.9%)	70 (-30-240)	250 (-76-820)
Vietnam	1995	Male	All Ages	44% (39-50%)	56% (50-61%)	1.1 (0.81-1.5)	5.1% (3.4-7%)	6.4% (3.8-9.2%)	17000 (9700-25000)	640000 (420000-900000)
Vietnam	1995	Male	15 to 19	36% (26-46%)	64% (54-74%)	0.55 (0.3-0.87)	2.8% (1.6-4.4%)	3.6% (1.7-6.2%)	200 (85-370)	18000 (9900-30000)
Vietnam	1995	Male	20 to 24	52% (39-63%)	48% (37-61%)	1.3 (0.78-1.8)	6.5% (4.2-9.5%)	7.8% (4.4-12%)	620 (320-1000)	50000 (30000-79000)
Vietnam	1995	Male	25 to 29	62% (51-72%)	38% (28-49%)	1.7 (1.1-2.4)	8.4% (5.2-12%)	11% (6.1-16%)	870 (460-1400)	62000 (38000-94000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	1995	Male	30 to 34	69% (58-78%)	31% (22-42%)	2.1 (1.4-3)	9.8% (6.4-14%)	13% (7.9-18%)	1100 (640-1700)	70000 (43000-100000)
Vietnam	1995	Male	35 to 39	72% (61-80%)	28% (20-39%)	2.6 (1.7-3.7)	11% (7.4-15%)	14% (9.4-20%)	1400 (830-2000)	77000 (49000-110000)
Vietnam	1995	Male	40 to 44	71% (61-79%)	29% (21-39%)	2.7 (1.7-3.8)	11% (7.3-15%)	14% (9.4-19%)	1300 (830-2000)	67000 (43000-96000)
Vietnam	1995	Male	45 to 49	66% (54-76%)	34% (24-46%)	2.4 (1.5-3.5)	10% (6.8-15%)	13% (8.5-19%)	1100 (670-1600)	48000 (31000-70000)
Vietnam	1995	Male	50 to 54	63% (52-72%)	37% (28-48%)	2.3 (1.4-3.4)	9.7% (5.9-14%)	12% (7.3-17%)	1200 (680-1700)	44000 (26000-66000)
Vietnam	1995	Male	55 to 59	61% (48-70%)	39% (30-52%)	2.3 (1.4-3.4)	9.2% (5.5-14%)	11% (6.7-17%)	1500 (870-2300)	50000 (29000-76000)
Vietnam	1995	Male	60 to 64	57% (44-68%)	43% (32-56%)	2.1 (1.3-3.1)	7.6% (4.2-12%)	9% (4.9-14%)	1900 (960-3000)	52000 (27000-82000)
Vietnam	1995	Male	65 to 69	53% (40-65%)	47% (35-60%)	1.8 (1.1-2.7)	5.9% (2.8-9.7%)	6.9% (3.2-11%)	1600 (750-2700)	38000 (18000-62000)
Vietnam	1995	Male	70 to 74	50% (38-62%)	50% (38-62%)	1.6 (0.88-2.4)	5.2% (2-9%)	6.1% (2.3-11%)	1700 (630-2900)	30000 (12000-53000)
Vietnam	1995	Male	75 to 79	49% (34-64%)	51% (36-66%)	1.4 (0.75-2.2)	3.9% (0.85-7.4%)	4.4% (0.86-8.5%)	1200 (220-2300)	17000 (3600-33000)
Vietnam	1995	Male	80 to 84	46% (30-61%)	54% (39-70%)	1.2 (0.63-1.8)	3.3% (0.54-6.7%)	3.8% (0.51-7.8%)	880 (120-1800)	9300 (1500-18000)
Vietnam	1995	Male	85 to 89	42% (28-58%)	58% (42-72%)	0.94 (0.51-1.5)	2% (-0.048-4.6%)	2.3% (-0.15-5.3%)	350 (-23-810)	2700 (-64-6000)
Vietnam	1995	Male	90 to 94	42% (28-57%)	58% (43-72%)	0.82 (0.42-1.4)	1% (-0.57-3.2%)	1.1% (-0.84-3.6%)	72 (-55-240)	400 (-220-1200)
Vietnam	1995	Male	95 plus	43% (26-62%)	57% (38-74%)	0.72 (0.31-1.4)	0.5% (-1-2.5%)	0.39% (-1.4-2.7%)	6.8 (-24-48)	34 (-69-170)
Vietnam	2000	Female	All Ages	8.8% (7.3-11%)	91% (89-93%)	0.29 (0.2-0.4)	2.1% (1.4-3.1%)	3.1% (1.5-5.1%)	8500 (3900-14000)	220000 (130000-310000)
Vietnam	2000	Female	15 to 19	9% (5.7-13%)	91% (87-94%)	0.3 (0.15-0.53)	1.5% (1-2.2%)	1.7% (0.74-3.2%)	39 (17-73)	7300 (4600-11000)
Vietnam	2000	Female	20 to 24	9.8% (6.3-14%)	90% (86-94%)	0.34 (0.17-0.6)	3% (2-4.2%)	2.3% (1-4.3%)	60 (25-120)	14000 (8900-22000)
Vietnam	2000	Female	25 to 29	10% (6.6-15%)	90% (85-93%)	0.35 (0.18-0.6)	3.1% (2.2-4.3%)	2.7% (1.3-5%)	80 (34-160)	15000 (9700-22000)
Vietnam	2000	Female	30 to 34	10% (6.9-16%)	90% (84-93%)	0.37 (0.2-0.61)	2.7% (1.8-3.7%)	2.6% (1.4-4.4%)	91 (45-170)	14000 (8600-20000)
Vietnam	2000	Female	35 to 39	10% (6.3-15%)	90% (85-94%)	0.38 (0.2-0.64)	2.7% (1.8-3.8%)	3% (1.6-5.3%)	140 (68-250)	14000 (9100-21000)
Vietnam	2000	Female	40 to 44	9.7% (6.3-14%)	90% (86-94%)	0.4 (0.21-0.65)	2.5% (1.7-3.7%)	3.1% (1.7-5.1%)	190 (100-320)	14000 (9000-21000)
Vietnam	2000	Female	45 to 49	9.9% (6.1-14%)	90% (86-94%)	0.44 (0.22-0.81)	2.6% (1.6-4.1%)	3.4% (1.8-5.8%)	240 (120-420)	14000 (8200-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	2000	Female	50 to 54	11% (7.6-16%)	89% (84-92%)	0.53 (0.28-0.93)	2.9% (1.7-4.6%)	3.8% (1.8-6.5%)	270 (130-470)	12000 (6800-20000)
Vietnam	2000	Female	55 to 59	12% (8.1-17%)	88% (83-92%)	0.64 (0.34-1.1)	3.1% (1.6-5.2%)	4.1% (2-7.1%)	360 (170-630)	13000 (7000-22000)
Vietnam	2000	Female	60 to 64	12% (7.7-17%)	88% (83-92%)	0.64 (0.32-1.1)	3.3% (1.6-5.6%)	4.2% (2-7.4%)	570 (260-990)	17000 (8300-29000)
Vietnam	2000	Female	65 to 69	11% (6.9-15%)	89% (85-93%)	0.57 (0.3-0.93)	2.9% (1.3-4.9%)	3.7% (1.7-6.3%)	800 (340-1400)	19000 (8700-34000)
Vietnam	2000	Female	70 to 74	9.9% (6.6-14%)	90% (86-93%)	0.53 (0.27-0.93)	2.6% (1.1-4.9%)	3.3% (1.4-6.1%)	920 (360-1800)	17000 (7000-33000)
Vietnam	2000	Female	75 to 79	11% (6.7-16%)	89% (84-93%)	0.56 (0.28-1)	2.5% (0.87-4.8%)	3.1% (1.1-5.9%)	1100 (340-2200)	16000 (5100-31000)
Vietnam	2000	Female	80 to 84	12% (7.7-17%)	88% (83-92%)	0.62 (0.32-1.1)	2.8% (1.1-5.1%)	3.4% (1.3-6.2%)	1500 (540-2800)	16000 (5900-29000)
Vietnam	2000	Female	85 to 89	14% (8.3-20%)	86% (80-92%)	0.71 (0.35-1.2)	2.7% (0.97-5.3%)	3.3% (1.1-6.3%)	1200 (420-2400)	9200 (3200-18000)
Vietnam	2000	Female	90 to 94	16% (9.8-24%)	84% (76-90%)	0.82 (0.39-1.5)	2.6% (0.71-5.4%)	3% (0.78-6.5%)	640 (160-1400)	3300 (900-7000)
Vietnam	2000	Female	95 plus	20% (10-32%)	80% (68-90%)	1 (0.39-2)	2.8% (0.43-6.5%)	3.2% (0.41-7.7%)	250 (33-590)	850 (130-2000)
Vietnam	2000	Male	All Ages	49% (45-54%)	51% (46-55%)	2.1 (1.5-2.7)	8.8% (6.4-11%)	11% (7.3-15%)	32000 (20000-45000)	1100000 (760000-1500000)
Vietnam	2000	Male	15 to 19	34% (26-44%)	66% (56-74%)	0.87 (0.5-1.3)	3.7% (2.3-5.7%)	5.1% (2.8-8.2%)	300 (150-510)	26000 (15000-41000)
Vietnam	2000	Male	20 to 24	53% (43-62%)	47% (38-57%)	2.1 (1.4-3.1)	9.1% (6-13%)	11% (7.1-17%)	930 (550-1400)	73000 (45000-110000)
Vietnam	2000	Male	25 to 29	67% (59-73%)	33% (27-41%)	3 (2-4.3)	12% (8.1-16%)	16% (10-22%)	1300 (790-1900)	90000 (59000-130000)
Vietnam	2000	Male	30 to 34	74% (67-80%)	26% (20-33%)	3.8 (2.6-5.3)	14% (9.7-18%)	18% (13-25%)	1600 (980-2400)	100000 (67000-150000)
Vietnam	2000	Male	35 to 39	76% (70-81%)	24% (19-30%)	4.6 (3.1-6.5)	15% (11-20%)	21% (15-27%)	2000 (1300-2900)	110000 (75000-160000)
Vietnam	2000	Male	40 to 44	75% (69-81%)	25% (19-31%)	4.8 (3.2-6.7)	16% (11-20%)	21% (15-26%)	2300 (1500-3300)	120000 (78000-160000)
Vietnam	2000	Male	45 to 49	71% (63-77%)	29% (23-37%)	4.2 (2.8-6)	15% (11-20%)	20% (14-26%)	2500 (1600-3400)	110000 (72000-150000)
Vietnam	2000	Male	50 to 54	68% (59-75%)	32% (25-41%)	4 (2.6-5.8)	15% (9.9-20%)	19% (12-25%)	2100 (1300-2900)	80000 (51000-110000)
Vietnam	2000	Male	55 to 59	66% (56-73%)	34% (27-44%)	4.1 (2.7-6)	15% (10-21%)	19% (12-25%)	2500 (1600-3400)	81000 (53000-110000)
Vietnam	2000	Male	60 to 64	62% (52-70%)	38% (30-48%)	3.8 (2.4-5.6)	13% (8.1-19%)	16% (9.5-22%)	3000 (1800-4300)	84000 (51000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	2000	Male	65 to 69	59% (49-68%)	41% (32-51%)	3.4 (2.1-4.9)	11% (6.2-16%)	13% (7.2-19%)	3700 (2000-5400)	84000 (45000-120000)
Vietnam	2000	Male	70 to 74	57% (46-67%)	43% (33-54%)	2.9 (1.7-4.4)	10% (5-16%)	12% (5.9-18%)	3600 (1700-5500)	66000 (32000-100000)
Vietnam	2000	Male	75 to 79	53% (40-67%)	47% (33-60%)	2.5 (1.4-4)	7.9% (3.2-13%)	9% (3.6-15%)	2900 (1100-4900)	41000 (17000-69000)
Vietnam	2000	Male	80 to 84	48% (32-63%)	52% (37-68%)	2 (1.1-3.1)	6.5% (2.6-11%)	7.5% (2.9-13%)	2100 (820-3700)	22000 (8900-39000)
Vietnam	2000	Male	85 to 89	43% (29-59%)	57% (41-71%)	1.6 (0.87-2.6)	4.5% (1.5-8.2%)	5.1% (1.6-9.3%)	960 (300-1800)	7300 (2300-13000)
Vietnam	2000	Male	90 to 94	43% (29-58%)	57% (42-71%)	1.4 (0.74-2.4)	2.8% (0.51-6%)	3.2% (0.43-6.8%)	270 (36-580)	1400 (250-3000)
Vietnam	2000	Male	95 plus	44% (28-62%)	56% (38-72%)	1.2 (0.53-2.4)	1.9% (-0.26-5.5%)	1.9% (-0.58-5.9%)	44 (-13-130)	170 (-24-480)
Vietnam	2005	Female	All Ages	9.1% (7.7-11%)	91% (89-92%)	0.47 (0.32-0.66)	3.1% (2.2-4.1%)	4.8% (2.8-7.1%)	14000 (7800-21000)	320000 (210000-440000)
Vietnam	2005	Female	15 to 19	9.2% (5.8-13%)	91% (87-94%)	0.47 (0.23-0.83)	1.8% (1.2-2.8%)	2.6% (1.2-4.8%)	58 (26-110)	9100 (5700-14000)
Vietnam	2005	Female	20 to 24	9.9% (6.3-14%)	90% (86-94%)	0.51 (0.25-0.94)	3.4% (2.3-4.8%)	3.3% (1.5-5.9%)	82 (38-150)	17000 (11000-25000)
Vietnam	2005	Female	25 to 29	10% (6.8-15%)	90% (85-93%)	0.53 (0.28-0.93)	3.5% (2.5-4.8%)	3.5% (1.7-6.1%)	98 (45-180)	17000 (11000-25000)
Vietnam	2005	Female	30 to 34	10% (6.8-16%)	90% (84-93%)	0.55 (0.3-0.92)	3.1% (2.2-4.4%)	3.4% (1.9-5.6%)	120 (57-210)	16000 (10000-24000)
Vietnam	2005	Female	35 to 39	10% (6.4-15%)	90% (85-94%)	0.58 (0.31-0.99)	3.2% (2.2-4.5%)	4.1% (2.3-6.6%)	180 (91-320)	17000 (11000-26000)
Vietnam	2005	Female	40 to 44	9.8% (6.2-14%)	90% (86-94%)	0.6 (0.32-1)	3.2% (2.1-4.6%)	4.3% (2.5-6.7%)	270 (150-440)	19000 (12000-29000)
Vietnam	2005	Female	45 to 49	9.9% (6.1-14%)	90% (86-94%)	0.67 (0.34-1.3)	3.5% (2.2-5.3%)	4.8% (2.7-7.8%)	410 (220-690)	23000 (14000-35000)
Vietnam	2005	Female	50 to 54	11% (7.7-16%)	89% (84-92%)	0.82 (0.44-1.5)	4% (2.4-6%)	5.5% (3.1-8.7%)	550 (310-910)	24000 (15000-38000)
Vietnam	2005	Female	55 to 59	12% (8.1-18%)	88% (82-92%)	0.99 (0.53-1.7)	4.4% (2.6-6.8%)	6% (3.3-9.3%)	590 (320-950)	21000 (12000-33000)
Vietnam	2005	Female	60 to 64	12% (7.8-17%)	88% (83-92%)	0.98 (0.49-1.7)	4.6% (2.5-7.2%)	6.1% (3.3-9.7%)	760 (400-1200)	22000 (12000-35000)
Vietnam	2005	Female	65 to 69	11% (6.9-15%)	89% (85-93%)	0.86 (0.45-1.5)	4.2% (2.3-6.5%)	5.4% (2.9-8.4%)	1000 (520-1700)	25000 (13000-39000)
Vietnam	2005	Female	70 to 74	9.9% (6.7-14%)	90% (86-93%)	0.8 (0.42-1.4)	3.9% (2-6.4%)	4.9% (2.4-8.1%)	1500 (710-2600)	28000 (14000-48000)
Vietnam	2005	Female	75 to 79	11% (6.9-16%)	89% (84-93%)	0.85 (0.43-1.5)	3.8% (1.8-6.4%)	4.7% (2.2-7.9%)	1700 (790-2900)	26000 (12000-43000)
Vietnam	2005	Female	80 to 84	12% (7.5-17%)	88% (83-93%)	0.97 (0.49-1.7)	4.3% (2.1-7%)	5.3% (2.6-8.6%)	2600 (1300-4200)	28000 (13000-45000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	2005	Female	85 to 89	14% (8.5-21%)	86% (79-92%)	1.1 (0.54-1.9)	4.4% (2-7.3%)	5.3% (2.4-8.9%)	2300 (1000-3900)	17000 (7800-29000)
Vietnam	2005	Female	90 to 94	17% (10-24%)	83% (76-90%)	1.3 (0.63-2.4)	4.3% (1.8-7.7%)	5.1% (2.1-9.4%)	1300 (530-2300)	6600 (2800-12000)
Vietnam	2005	Female	95 plus	20% (11-33%)	80% (67-89%)	1.6 (0.62-3.2)	4.7% (1.5-9.2%)	5.4% (1.7-11%)	540 (160-1100)	1900 (600-3600)
Vietnam	2005	Male	All Ages	51% (47-56%)	49% (44-53%)	3.5 (2.5-4.5)	12% (9.5-15%)	16% (12-19%)	47000 (33000-62000)	1500000 (1200000-2000000)
Vietnam	2005	Male	15 to 19	34% (26-44%)	66% (56-74%)	1.3 (0.75-2)	4.9% (3-7.5%)	6.9% (3.8-11%)	430 (230-700)	36000 (21000-56000)
Vietnam	2005	Male	20 to 24	53% (44-62%)	47% (38-56%)	3.3 (2.1-4.7)	12% (8-16%)	15% (10-22%)	1300 (760-2000)	98000 (64000-150000)
Vietnam	2005	Male	25 to 29	67% (58-73%)	33% (27-42%)	4.5 (3-6.5)	14% (9.9-19%)	19% (13-26%)	1600 (910-2300)	110000 (69000-160000)
Vietnam	2005	Male	30 to 34	74% (67-80%)	26% (20-33%)	5.7 (3.9-8)	16% (12-20%)	22% (16-28%)	1900 (1200-2800)	120000 (79000-170000)
Vietnam	2005	Male	35 to 39	76% (70-82%)	24% (18-30%)	7 (4.6-9.9)	18% (13-22%)	25% (19-30%)	2500 (1600-3500)	140000 (98000-190000)
Vietnam	2005	Male	40 to 44	75% (69-81%)	25% (19-31%)	7.3 (4.9-10)	19% (14-22%)	25% (20-29%)	3100 (2200-4300)	160000 (110000-210000)
Vietnam	2005	Male	45 to 49	71% (63-78%)	29% (22-37%)	6.5 (4.4-9.1)	19% (15-23%)	25% (20-30%)	3900 (2800-5300)	170000 (130000-230000)
Vietnam	2005	Male	50 to 54	68% (59-75%)	32% (25-41%)	6.1 (4-8.8)	19% (14-23%)	24% (18-29%)	4200 (3000-5500)	160000 (120000-210000)
Vietnam	2005	Male	55 to 59	66% (56-73%)	34% (27-44%)	6.2 (4.1-8.9)	20% (15-24%)	24% (19-29%)	3600 (2600-4700)	120000 (88000-150000)
Vietnam	2005	Male	60 to 64	62% (52-71%)	38% (29-48%)	5.8 (3.7-8.6)	18% (13-22%)	21% (15-26%)	3700 (2500-4800)	100000 (72000-130000)
Vietnam	2005	Male	65 to 69	59% (49-68%)	41% (32-51%)	5.1 (3.1-7.4)	16% (10-20%)	19% (12-23%)	4400 (2900-5800)	100000 (68000-130000)
Vietnam	2005	Male	70 to 74	57% (46-66%)	43% (34-54%)	4.5 (2.7-6.7)	15% (8.9-20%)	17% (10-23%)	5700 (3400-7800)	110000 (62000-140000)
Vietnam	2005	Male	75 to 79	53% (39-66%)	47% (34-61%)	3.7 (2.1-6)	12% (6.5-17%)	14% (7.4-20%)	4600 (2400-6700)	65000 (35000-95000)
Vietnam	2005	Male	80 to 84	48% (31-63%)	52% (37-69%)	3 (1.7-4.8)	10% (5.1-15%)	12% (5.8-18%)	3800 (1900-5900)	40000 (20000-62000)
Vietnam	2005	Male	85 to 89	43% (28-58%)	57% (42-72%)	2.4 (1.3-4)	7.5% (3.2-12%)	8.5% (3.7-14%)	1900 (810-3300)	15000 (6300-24000)
Vietnam	2005	Male	90 to 94	42% (29-58%)	58% (42-71%)	2.1 (1.1-3.6)	5.2% (1.8-10%)	5.8% (1.9-12%)	590 (200-1200)	3100 (1100-6100)
Vietnam	2005	Male	95 plus	44% (27-63%)	56% (37-73%)	1.8 (0.78-3.7)	3.8% (0.54-9%)	4% (0.37-9.9%)	120 (11-290)	430 (61-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	2010	Female	All Ages	10% (8.7-12%)	90% (88-91%)	0.76 (0.51-1.1)	4% (2.9-5.1%)	6.1% (4-8.4%)	19000 (12000-26000)	420000 (300000-550000)
Vietnam	2010	Female	15 to 19	9.3% (5.9-14%)	91% (86-94%)	0.65 (0.31-1.1)	2.2% (1.4-3.3%)	3.6% (1.6-6.2%)	72 (31-130)	10000 (6500-16000)
Vietnam	2010	Female	20 to 24	10% (6.5-15%)	90% (85-94%)	0.76 (0.36-1.4)	4% (2.7-5.5%)	4.4% (2.1-7.4%)	110 (48-190)	20000 (13000-29000)
Vietnam	2010	Female	25 to 29	11% (7.1-16%)	89% (84-93%)	0.81 (0.4-1.5)	4.1% (2.9-5.5%)	4.7% (2.4-7.8%)	120 (59-220)	20000 (13000-29000)
Vietnam	2010	Female	30 to 34	12% (7.3-18%)	88% (82-93%)	0.87 (0.43-1.5)	3.7% (2.5-5.1%)	4.7% (2.5-7.5%)	150 (73-260)	19000 (12000-27000)
Vietnam	2010	Female	35 to 39	12% (7.4-17%)	88% (83-93%)	0.95 (0.49-1.6)	3.9% (2.7-5.3%)	5.6% (3.3-8.6%)	240 (130-400)	22000 (14000-31000)
Vietnam	2010	Female	40 to 44	12% (7.7-17%)	88% (83-92%)	1 (0.53-1.8)	4.1% (2.7-5.7%)	6.2% (3.7-9.1%)	360 (200-570)	25000 (16000-36000)
Vietnam	2010	Female	45 to 49	12% (6.9-17%)	88% (83-93%)	1.1 (0.54-2.1)	4.5% (2.9-6.6%)	6.8% (4-10%)	590 (330-920)	32000 (20000-47000)
Vietnam	2010	Female	50 to 54	12% (8.1-18%)	88% (82-92%)	1.2 (0.65-2.3)	5% (3.3-7.5%)	7.3% (4.4-11%)	870 (500-1300)	38000 (24000-56000)
Vietnam	2010	Female	55 to 59	13% (8.1-19%)	87% (81-92%)	1.4 (0.73-2.5)	5.4% (3.4-8%)	7.6% (4.7-11%)	1000 (610-1600)	38000 (24000-58000)
Vietnam	2010	Female	60 to 64	13% (7.9-19%)	87% (81-92%)	1.4 (0.69-2.6)	5.5% (3.4-8.3%)	7.5% (4.5-11%)	1000 (590-1500)	30000 (18000-45000)
Vietnam	2010	Female	65 to 69	11% (7.2-16%)	89% (84-93%)	1.3 (0.64-2.2)	5.2% (3.1-7.7%)	6.8% (4-10%)	1100 (640-1700)	28000 (16000-42000)
Vietnam	2010	Female	70 to 74	11% (7.2-16%)	89% (84-93%)	1.2 (0.59-2.2)	4.9% (2.8-7.7%)	6.3% (3.7-9.8%)	1600 (860-2600)	30000 (17000-49000)
Vietnam	2010	Female	75 to 79	11% (7.2-17%)	89% (83-93%)	1.2 (0.59-2.3)	4.7% (2.6-7.2%)	5.9% (3.3-9.2%)	2300 (1200-3600)	34000 (18000-53000)
Vietnam	2010	Female	80 to 84	12% (7.5-17%)	88% (83-92%)	1.3 (0.67-2.3)	5% (2.9-7.6%)	6.2% (3.6-9.5%)	3300 (1800-5000)	35000 (20000-53000)
Vietnam	2010	Female	85 to 89	14% (8.3-20%)	86% (80-92%)	1.5 (0.74-2.6)	5.2% (2.8-8.2%)	6.4% (3.4-10%)	3100 (1600-4800)	23000 (12000-36000)
Vietnam	2010	Female	90 to 94	16% (9.6-24%)	84% (76-90%)	1.7 (0.81-3.2)	5.2% (2.7-8.7%)	6.3% (3.1-10%)	1900 (950-3100)	9800 (5100-16000)
Vietnam	2010	Female	95 plus	19% (10-31%)	81% (69-90%)	2 (0.81-4.2)	5.7% (2.3-10%)	6.6% (2.6-12%)	800 (310-1500)	2800 (1100-5000)
Vietnam	2010	Male	All Ages	49% (44-54%)	51% (46-56%)	4.8 (3.4-6.4)	14% (11-17%)	18% (14-21%)	56000 (41000-72000)	1800000 (1400000-2300000)
Vietnam	2010	Male	15 to 19	35% (25-46%)	65% (54-75%)	1.9 (1-2.9)	6.4% (3.8-9.9%)	9.2% (5.1-15%)	550 (280-910)	46000 (25000-72000)
Vietnam	2010	Male	20 to 24	51% (39-62%)	49% (38-61%)	4.4 (2.6-6.6)	14% (9.7-20%)	19% (12-28%)	1600 (950-2500)	120000 (78000-180000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	2010	Male	25 to 29	62% (51-71%)	38% (29-49%)	5.8 (3.6-8.7)	16% (12-21%)	22% (16-29%)	1800 (1100-2600)	130000 (82000-170000)
Vietnam	2010	Male	30 to 34	69% (58-78%)	31% (22-42%)	7.4 (4.7-11)	17% (13-21%)	24% (19-30%)	2000 (1300-2800)	130000 (91000-170000)
Vietnam	2010	Male	35 to 39	72% (61-79%)	28% (21-39%)	9 (5.7-13)	19% (14-22%)	26% (21-31%)	2500 (1800-3500)	150000 (110000-200000)
Vietnam	2010	Male	40 to 44	71% (61-78%)	29% (22-39%)	9.4 (5.9-14)	20% (16-23%)	27% (22-31%)	3400 (2500-4500)	170000 (130000-230000)
Vietnam	2010	Male	45 to 49	66% (53-75%)	34% (25-47%)	8.3 (5.4-12)	21% (17-24%)	27% (23-32%)	4600 (3400-6100)	200000 (150000-260000)
Vietnam	2010	Male	50 to 54	63% (52-72%)	37% (28-48%)	7.8 (5-12)	20% (17-24%)	26% (22-30%)	5600 (4200-7200)	210000 (160000-270000)
Vietnam	2010	Male	55 to 59	60% (48-70%)	40% (30-52%)	7.8 (5-12)	21% (17-24%)	25% (21-30%)	5900 (4400-7500)	190000 (150000-240000)
Vietnam	2010	Male	60 to 64	57% (44-68%)	43% (32-56%)	7.3 (4.3-11)	19% (15-22%)	22% (18-26%)	4300 (3200-5500)	120000 (90000-150000)
Vietnam	2010	Male	65 to 69	53% (40-65%)	47% (35-60%)	6.3 (3.7-9.6)	17% (13-20%)	20% (15-24%)	4200 (3000-5400)	98000 (72000-130000)
Vietnam	2010	Male	70 to 74	50% (37-63%)	50% (37-63%)	5.4 (3-8.4)	16% (11-20%)	18% (13-23%)	5000 (3500-6600)	94000 (66000-120000)
Vietnam	2010	Male	75 to 79	49% (34-64%)	51% (36-66%)	4.7 (2.6-8)	14% (8.7-18%)	16% (9.9-21%)	5500 (3400-7600)	80000 (49000-110000)
Vietnam	2010	Male	80 to 84	46% (30-61%)	54% (39-70%)	4 (2.1-6.5)	12% (7.4-17%)	14% (8.3-20%)	5100 (3000-7100)	54000 (32000-76000)
Vietnam	2010	Male	85 to 89	41% (28-56%)	59% (44-72%)	3.2 (1.7-5.4)	9.7% (5.2-14%)	11% (5.9-16%)	2900 (1500-4300)	22000 (11000-33000)
Vietnam	2010	Male	90 to 94	40% (27-56%)	60% (44-73%)	2.8 (1.4-4.7)	7.2% (3.2-12%)	8.2% (3.5-14%)	1000 (430-1700)	5400 (2400-8700)
Vietnam	2010	Male	95 plus	42% (25-61%)	58% (39-75%)	2.4 (1-4.8)	5.6% (1.5-11%)	6.1% (1.4-12%)	220 (50-440)	790 (210-1500)
Vietnam	2016	Female	All Ages	10% (8.8-12%)	90% (88-91%)	0.85 (0.51-1.3)	4% (2.9-5.1%)	5.9% (3.8-8.1%)	19000 (12000-27000)	430000 (310000-560000)
Vietnam	2016	Female	15 to 19	9.3% (5.8-14%)	91% (86-94%)	0.71 (0.31-1.3)	2.3% (1.5-3.3%)	3.7% (1.6-6.5%)	51 (22-93)	7700 (4800-11000)
Vietnam	2016	Female	20 to 24	10% (6.5-15%)	90% (85-93%)	0.82 (0.36-1.6)	4.1% (2.8-5.5%)	4.4% (2.1-7.5%)	91 (42-160)	19000 (12000-27000)
Vietnam	2016	Female	25 to 29	11% (6.8-17%)	89% (83-93%)	0.86 (0.41-1.7)	4.1% (2.9-5.5%)	4.4% (2.1-7.5%)	110 (50-200)	21000 (14000-30000)
Vietnam	2016	Female	30 to 34	12% (7.2-18%)	88% (82-93%)	0.92 (0.42-1.8)	3.7% (2.6-5.1%)	4.4% (2.3-7.4%)	140 (67-250)	20000 (13000-29000)
Vietnam	2016	Female	35 to 39	12% (7.3-17%)	88% (83-93%)	1 (0.49-1.9)	3.8% (2.6-5.3%)	5.4% (2.9-8.2%)	220 (110-370)	22000 (14000-31000)
Vietnam	2016	Female	40 to 44	12% (7.6-17%)	88% (83-92%)	1.1 (0.53-2)	3.9% (2.6-5.5%)	5.9% (3.4-8.8%)	330 (180-540)	25000 (16000-36000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	2016	Female	45 to 49	12% (7-17%)	88% (83-93%)	1.2 (0.54-2.1)	4.3% (2.8-6.2%)	6.6% (3.8-9.7%)	530 (300-840)	31000 (19000-45000)
Vietnam	2016	Female	50 to 54	12% (7.8-18%)	88% (82-92%)	1.3 (0.64-2.4)	4.8% (3.1-6.9%)	7% (4.3-10%)	830 (480-1300)	38000 (23000-56000)
Vietnam	2016	Female	55 to 59	13% (8.4-18%)	87% (82-92%)	1.5 (0.73-2.7)	5.1% (3.3-7.4%)	7.2% (4.4-11%)	1200 (710-1800)	45000 (28000-66000)
Vietnam	2016	Female	60 to 64	12% (8-18%)	88% (82-92%)	1.5 (0.71-2.8)	5.1% (3.1-7.6%)	7% (4.2-10%)	1400 (810-2100)	44000 (26000-65000)
Vietnam	2016	Female	65 to 69	11% (7.3-16%)	89% (84-93%)	1.4 (0.67-2.6)	4.9% (2.9-7.2%)	6.5% (3.8-9.7%)	1300 (750-1900)	32000 (19000-47000)
Vietnam	2016	Female	70 to 74	11% (7.2-16%)	89% (84-93%)	1.3 (0.6-2.4)	4.6% (2.6-7.1%)	5.9% (3.4-9.1%)	1300 (720-2000)	26000 (15000-40000)
Vietnam	2016	Female	75 to 79	11% (7.2-17%)	89% (83-93%)	1.3 (0.62-2.7)	4.4% (2.4-6.9%)	5.6% (3.1-8.6%)	1700 (940-2700)	26000 (14000-40000)
Vietnam	2016	Female	80 to 84	12% (7.7-17%)	88% (83-92%)	1.4 (0.69-2.5)	4.7% (2.6-7.1%)	5.8% (3.2-9%)	3000 (1700-4600)	33000 (19000-50000)
Vietnam	2016	Female	85 to 89	14% (8.3-20%)	86% (80-92%)	1.6 (0.7-2.9)	4.9% (2.6-7.6%)	5.9% (3.1-9.4%)	3500 (1800-5500)	26000 (14000-41000)
Vietnam	2016	Female	90 to 94	16% (9.7-24%)	84% (76-90%)	1.8 (0.81-3.5)	5% (2.4-8.1%)	5.9% (2.8-9.8%)	2200 (1100-3700)	12000 (5600-19000)
Vietnam	2016	Female	95 plus	19% (9.9-31%)	81% (69-90%)	2.2 (0.79-4.7)	5.5% (2.2-9.8%)	6.3% (2.4-12%)	1000 (390-1900)	3600 (1400-6600)
Vietnam	2016	Male	All Ages	50% (45-55%)	50% (45-55%)	5.4 (3.4-7.9)	14% (12-17%)	18% (14-21%)	60000 (44000-76000)	1900000 (1500000-2400000)
Vietnam	2016	Male	15 to 19	35% (24-46%)	65% (54-76%)	2 (1-3.3)	6.6% (3.8-11%)	9.7% (5.3-16%)	410 (210-710)	34000 (19000-56000)
Vietnam	2016	Male	20 to 24	51% (39-62%)	49% (38-61%)	4.7 (2.6-7.5)	14% (9.6-20%)	19% (12-28%)	1400 (800-2200)	110000 (70000-160000)
Vietnam	2016	Male	25 to 29	62% (50-71%)	38% (29-50%)	6.2 (3.5-9.7)	16% (11-21%)	22% (15-29%)	1800 (1000-2600)	130000 (82000-180000)
Vietnam	2016	Male	30 to 34	69% (57-78%)	31% (22-43%)	7.9 (4.6-12)	17% (13-22%)	24% (18-30%)	2000 (1200-3000)	130000 (87000-190000)
Vietnam	2016	Male	35 to 39	72% (61-79%)	28% (21-39%)	9.7 (5.7-16)	18% (14-22%)	26% (20-31%)	2400 (1600-3500)	140000 (97000-200000)
Vietnam	2016	Male	40 to 44	70% (61-79%)	30% (21-39%)	10 (5.9-16)	19% (15-22%)	26% (21-30%)	3100 (2200-4300)	160000 (120000-220000)
Vietnam	2016	Male	45 to 49	66% (53-75%)	34% (25-47%)	8.9 (5.3-14)	20% (16-24%)	27% (22-31%)	4400 (3200-6000)	200000 (150000-260000)
Vietnam	2016	Male	50 to 54	63% (51-72%)	37% (28-49%)	8.3 (4.8-13)	20% (16-23%)	25% (21-29%)	5800 (4300-7700)	230000 (170000-290000)
Vietnam	2016	Male	55 to 59	60% (48-71%)	40% (29-52%)	8.4 (4.9-13)	20% (16-24%)	25% (20-29%)	7400 (5500-9400)	250000 (190000-310000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vietnam	2016	Male	60 to 64	57% (44-68%)	43% (32-56%)	7.8 (4.3-12)	18% (14-21%)	22% (17-26%)	7100 (5400-9000)	200000 (150000-250000)
Vietnam	2016	Male	65 to 69	53% (39-64%)	47% (36-61%)	6.6 (3.6-11)	16% (12-19%)	19% (14-23%)	5100 (3800-6500)	120000 (89000-150000)
Vietnam	2016	Male	70 to 74	50% (37-62%)	50% (38-63%)	5.7 (2.9-9.5)	14% (9.9-18%)	17% (12-21%)	4400 (3000-5800)	83000 (57000-110000)
Vietnam	2016	Male	75 to 79	49% (34-64%)	51% (36-66%)	5 (2.5-8.8)	13% (7.8-17%)	15% (9-20%)	4400 (2700-6100)	64000 (40000-88000)
Vietnam	2016	Male	80 to 84	45% (30-60%)	55% (40-70%)	4.2 (2.1-7.3)	12% (6.7-16%)	13% (7.6-19%)	5000 (2900-7100)	54000 (31000-75000)
Vietnam	2016	Male	85 to 89	41% (26-56%)	59% (44-74%)	3.4 (1.6-5.9)	9.2% (4.7-14%)	11% (5.2-16%)	3400 (1700-5100)	26000 (13000-39000)
Vietnam	2016	Male	90 to 94	40% (27-56%)	60% (44-73%)	2.9 (1.4-5.4)	7.1% (2.8-12%)	8.1% (3-13%)	1300 (480-2100)	6700 (2700-11000)
Vietnam	2016	Male	95 plus	42% (25-61%)	58% (39-75%)	2.6 (1-5.1)	5.7% (1.3-11%)	6.1% (1.1-12%)	300 (56-620)	1100 (260-2200)
Fiji	1990	Female	All Ages	7.9% (6.1-10%)	92% (90-94%)	0.19 (0.14-0.25)	0.42% (0.22-0.66%)	0.24% (-0.051-0.62%)	4.5 (-0.9-13)	440 (220-730)
Fiji	1990	Female	15 to 19	9.2% (5.2-16%)	91% (84-95%)	0.25 (0.12-0.45)	0.77% (0.44-1.3%)	1% (0.4-2.1%)	0.31 (0.11-0.7)	39 (21-67)
Fiji	1990	Female	20 to 24	12% (6.5-20%)	88% (80-94%)	0.41 (0.21-0.69)	1.4% (0.88-2.2%)	1.6% (0.7-3%)	0.69 (0.26-1.4)	83 (47-140)
Fiji	1990	Female	25 to 29	12% (6.7-21%)	88% (79-93%)	0.36 (0.2-0.59)	1.4% (0.89-1.9%)	1.5% (0.77-2.5%)	0.89 (0.37-1.7)	94 (56-150)
Fiji	1990	Female	30 to 34	12% (6-20%)	88% (80-94%)	0.33 (0.17-0.57)	1.1% (0.68-1.7%)	1.1% (0.58-2%)	0.85 (0.36-1.6)	80 (45-130)
Fiji	1990	Female	35 to 39	11% (6-19%)	89% (81-94%)	0.34 (0.16-0.59)	1.1% (0.59-1.7%)	1.2% (0.57-2.1%)	0.98 (0.38-1.9)	71 (36-120)
Fiji	1990	Female	40 to 44	10% (5.6-19%)	90% (81-94%)	0.29 (0.14-0.53)	0.79% (0.35-1.4%)	0.92% (0.33-1.8%)	0.86 (0.27-1.9)	52 (22-100)
Fiji	1990	Female	45 to 49	10% (5-18%)	90% (82-95%)	0.22 (0.094-0.4)	0.37% (-0.04-0.92%)	0.42% (-0.088-1.2%)	0.47 (-0.11-1.4)	25 (-3.2-66)
Fiji	1990	Female	50 to 54	10% (5.4-18%)	90% (82-95%)	0.19 (0.077-0.37)	0.071% (-0.41-0.58%)	0.092% (-0.47-0.73%)	0.12 (-0.62-1)	4.7 (-27-40)
Fiji	1990	Female	55 to 59	9.3% (5.1-16%)	91% (84-95%)	0.18 (0.079-0.35)	0.036% (-0.43-0.59%)	0.079% (-0.46-0.76%)	0.12 (-0.72-1.2)	2.4 (-28-39)
Fiji	1990	Female	60 to 64	8.5% (4.3-15%)	91% (85-96%)	0.16 (0.07-0.33)	-0.05% (-0.52-0.5%)	-0.026% (-0.55-0.63%)	-0.037 (-0.86-1.1)	-2.6 (-29-30)
Fiji	1990	Female	65 to 69	8.9% (4.8-15%)	91% (85-95%)	0.16 (0.076-0.31)	-0.072% (-0.56-0.42%)	-0.044% (-0.59-0.54%)	-0.067 (-0.98-0.89)	-3.1 (-25-20)
Fiji	1990	Female	70 to 74	9.1% (5.2-15%)	91% (85-95%)	0.16 (0.074-0.31)	-0.15% (-0.64-0.36%)	-0.16% (-0.7-0.43%)	-0.27 (-1.2-0.84)	-6 (-25-15)
Fiji	1990	Female	75 to 79	8.9% (4.9-16%)	91% (84-95%)	0.15 (0.064-0.3)	-0.082% (-0.51-0.4%)	-0.073% (-0.55-0.47%)	-0.11 (-0.9-0.81)	-2.2 (-14-11)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	1990	Female	80 to 84	8.9% (4.8-15%)	91% (85-95%)	0.15 (0.067-0.28)	-0.13% (-0.52-0.26%)	-0.15% (-0.58-0.3%)	-0.16 (-0.67-0.33)	-1.8 (-7.2-3.6)
Fiji	1990	Female	85 to 89	9.1% (4.9-15%)	91% (85-95%)	0.15 (0.07-0.27)	-0.14% (-0.49-0.24%)	-0.16% (-0.56-0.26%)	-0.11 (-0.37-0.17)	-0.8 (-2.8-1.3)
Fiji	1990	Female	90 to 94	9.8% (5.5-16%)	90% (84-95%)	0.16 (0.07-0.3)	-0.1% (-0.4-0.26%)	-0.12% (-0.47-0.28%)	-0.031 (-0.12-0.068)	-0.15 (-0.6-0.39)
Fiji	1990	Female	95 plus	11% (5.6-20%)	89% (80-94%)	0.18 (0.07-0.37)	-0.089% (-0.53-0.41%)	-0.14% (-0.64-0.42%)	-0.0082 (-0.037-0.024)	-0.02 (-0.12-0.091)
Fiji	1990	Male	All Ages	37% (29-47%)	63% (53-71%)	1.2 (1-1.4)	3.1% (1.9-4.2%)	2.8% (1.4-4.4%)	76 (37-130)	4200 (2600-6300)
Fiji	1990	Male	15 to 19	34% (19-51%)	66% (49-81%)	0.79 (0.37-1.4)	2.3% (1.2-3.8%)	3% (1.4-5.4%)	1.8 (0.81-3.3)	160 (82-270)
Fiji	1990	Male	20 to 24	52% (33-70%)	48% (30-67%)	2.1 (1.2-3.2)	7.1% (4.2-10%)	8.8% (5-13%)	8.1 (4.2-13)	610 (360-950)
Fiji	1990	Male	25 to 29	59% (39-76%)	41% (24-61%)	2.8 (1.8-4)	8.6% (5.5-12%)	11% (6.4-15%)	12 (6.8-19)	820 (490-1200)
Fiji	1990	Male	30 to 34	60% (39-77%)	40% (23-61%)	3 (1.8-4.3)	7.6% (4.7-11%)	9.2% (5.3-14%)	12 (6-19)	760 (420-1200)
Fiji	1990	Male	35 to 39	58% (39-75%)	42% (25-61%)	2.7 (1.6-4.1)	6.5% (3.2-10%)	7.6% (3.4-12%)	10 (4-18)	580 (260-990)
Fiji	1990	Male	40 to 44	56% (37-73%)	44% (27-63%)	2.3 (1.2-3.6)	4.7% (1.5-8.7%)	5.2% (1.3-10%)	8 (1.9-17)	420 (130-830)
Fiji	1990	Male	45 to 49	52% (31-71%)	48% (29-69%)	1.7 (0.88-2.9)	2.8% (0.23-6.4%)	3% (-0.27-7.2%)	5.4 (-0.52-14)	270 (23-630)
Fiji	1990	Male	50 to 54	49% (30-68%)	51% (32-70%)	1.4 (0.68-2.3)	2.1% (-0.28-5.5%)	2.2% (-0.56-6.1%)	4.5 (-1.2-13)	190 (-24-530)
Fiji	1990	Male	55 to 59	47% (29-67%)	53% (33-71%)	1.2 (0.59-2.2)	1.8% (-0.24-4.9%)	1.8% (-0.46-5.5%)	4.1 (-1.1-12)	150 (-21-420)
Fiji	1990	Male	60 to 64	44% (26-64%)	56% (36-74%)	1 (0.5-1.9)	1.5% (-0.22-4.7%)	1.6% (-0.38-5.1%)	3.7 (-0.88-12)	120 (-17-350)
Fiji	1990	Male	65 to 69	43% (24-62%)	57% (38-76%)	0.93 (0.46-1.6)	1.2% (-0.28-3.5%)	1.3% (-0.36-3.8%)	2.8 (-0.77-8.5)	72 (-16-210)
Fiji	1990	Male	70 to 74	40% (23-57%)	60% (43-77%)	0.79 (0.38-1.4)	0.9% (-0.42-3%)	0.93% (-0.49-3.2%)	2.1 (-1.2-7.2)	42 (-20-140)
Fiji	1990	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.66 (0.3-1.3)	0.68% (-0.4-2.4%)	0.7% (-0.48-2.5%)	1.3 (-0.9-4.8)	20 (-12-73)
Fiji	1990	Male	80 to 84	33% (18-52%)	67% (48-82%)	0.58 (0.27-1.1)	0.54% (-0.29-1.8%)	0.53% (-0.39-1.9%)	0.6 (-0.41-2.1)	7.2 (-3.8-24)
Fiji	1990	Male	85 to 89	32% (19-48%)	68% (52-81%)	0.55 (0.26-0.97)	0.38% (-0.37-1.5%)	0.35% (-0.47-1.6%)	0.19 (-0.25-0.84)	1.7 (-1.7-6.9)
Fiji	1990	Male	90 to 94	33% (20-49%)	67% (51-80%)	0.56 (0.25-1)	0.21% (-0.46-1.2%)	0.15% (-0.61-1.3%)	0.026 (-0.11-0.23)	0.23 (-0.49-1.3)
Fiji	1990	Male	95 plus	36% (19-57%)	64% (43-81%)	0.61 (0.25-1.2)	0.11% (-0.77-1.3%)	-0.024% (-1-1.4%)	-0.0012 (-0.05-0.067)	0.021 (-0.14-0.25)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	1995	Female	All Ages	8.2% (6.4-11%)	92% (89-94%)	0.19 (0.14-0.25)	0.35% (0.13-0.6%)	0.12% (-0.19-0.52%)	2.8 (-4-12)	410 (160-710)
Fiji	1995	Female	15 to 19	9.4% (5.4-16%)	91% (84-95%)	0.25 (0.12-0.45)	0.79% (0.42-1.3%)	1% (0.4-2.1%)	0.39 (0.14-0.81)	46 (25-80)
Fiji	1995	Female	20 to 24	12% (6.5-20%)	88% (80-93%)	0.41 (0.21-0.7)	1.5% (0.87-2.3%)	1.6% (0.71-2.9%)	0.75 (0.31-1.5)	89 (51-140)
Fiji	1995	Female	25 to 29	12% (6.7-21%)	88% (79-93%)	0.35 (0.19-0.6)	1.4% (0.91-2%)	1.4% (0.78-2.5%)	0.89 (0.41-1.6)	91 (56-140)
Fiji	1995	Female	30 to 34	12% (6.1-20%)	88% (80-94%)	0.33 (0.17-0.56)	1.1% (0.64-1.7%)	1.1% (0.52-1.9%)	0.9 (0.4-1.7)	84 (48-140)
Fiji	1995	Female	35 to 39	11% (5.8-20%)	89% (80-94%)	0.33 (0.16-0.57)	0.99% (0.55-1.6%)	1.1% (0.48-2%)	1.2 (0.48-2.2)	85 (45-140)
Fiji	1995	Female	40 to 44	11% (5.8-18%)	89% (82-94%)	0.28 (0.14-0.51)	0.7% (0.24-1.4%)	0.79% (0.16-1.7%)	0.91 (0.19-2)	56 (20-110)
Fiji	1995	Female	45 to 49	10% (5.3-19%)	90% (81-95%)	0.22 (0.098-0.4)	0.24% (-0.2-0.8%)	0.27% (-0.27-0.98%)	0.38 (-0.4-1.4)	20 (-18-66)
Fiji	1995	Female	50 to 54	10% (5.6-18%)	90% (82-94%)	0.19 (0.081-0.36)	-0.067% (-0.57-0.48%)	-0.068% (-0.65-0.58%)	-0.12 (-1.1-0.95)	-5.8 (-49-39)
Fiji	1995	Female	55 to 59	9.3% (5.1-16%)	91% (84-95%)	0.17 (0.076-0.34)	-0.092% (-0.6-0.46%)	-0.071% (-0.64-0.59%)	-0.13 (-1.2-1.1)	-7.1 (-47-36)
Fiji	1995	Female	60 to 64	8.6% (4.5-15%)	91% (85-96%)	0.16 (0.07-0.32)	-0.18% (-0.7-0.36%)	-0.18% (-0.76-0.44%)	-0.35 (-1.5-0.84)	-12 (-48-24)
Fiji	1995	Female	65 to 69	9.1% (5.1-16%)	91% (84-95%)	0.16 (0.077-0.31)	-0.2% (-0.7-0.3%)	-0.19% (-0.78-0.38%)	-0.37 (-1.5-0.69)	-10 (-38-15)
Fiji	1995	Female	70 to 74	9.4% (5.3-16%)	91% (84-95%)	0.16 (0.074-0.31)	-0.28% (-0.78-0.22%)	-0.3% (-0.86-0.28%)	-0.57 (-1.7-0.54)	-12 (-34-9.8)
Fiji	1995	Female	75 to 79	9.2% (5.1-16%)	91% (84-95%)	0.15 (0.065-0.28)	-0.18% (-0.65-0.27%)	-0.19% (-0.7-0.34%)	-0.4 (-1.5-0.74)	-6.4 (-23-10)
Fiji	1995	Female	80 to 84	9% (4.9-16%)	91% (84-95%)	0.14 (0.068-0.27)	-0.23% (-0.64-0.15%)	-0.25% (-0.71-0.17%)	-0.36 (-1.1-0.25)	-4 (-12-2.6)
Fiji	1995	Female	85 to 89	9.1% (5.1-15%)	91% (85-95%)	0.15 (0.066-0.27)	-0.22% (-0.61-0.14%)	-0.25% (-0.7-0.15%)	-0.2 (-0.58-0.13)	-1.5 (-4.3-1.1)
Fiji	1995	Female	90 to 94	9.9% (5.4-16%)	90% (84-95%)	0.16 (0.067-0.29)	-0.17% (-0.51-0.21%)	-0.21% (-0.59-0.21%)	-0.062 (-0.18-0.064)	-0.31 (-0.93-0.37)
Fiji	1995	Female	95 plus	11% (5.7-20%)	89% (80-94%)	0.18 (0.067-0.35)	-0.17% (-0.64-0.36%)	-0.23% (-0.79-0.35%)	-0.018 (-0.059-0.028)	-0.05 (-0.18-0.11)
Fiji	1995	Male	All Ages	39% (30-48%)	61% (52-70%)	1.2 (1.1-1.4)	3% (1.9-4.3%)	2.7% (1.3-4.4%)	83 (40-140)	4600 (2800-6900)
Fiji	1995	Male	15 to 19	33% (19-51%)	67% (49-81%)	0.76 (0.35-1.3)	2.3% (1.2-3.9%)	3% (1.3-5.5%)	2.1 (0.93-3.9)	180 (94-320)
Fiji	1995	Male	20 to 24	51% (33-69%)	49% (31-67%)	2.1 (1.2-3.1)	7.2% (4.2-11%)	8.8% (4.9-14%)	8.8 (4.7-14)	660 (380-1000)
Fiji	1995	Male	25 to 29	59% (39-76%)	41% (24-61%)	2.8 (1.8-4)	8.6% (5.4-12%)	10% (6.3-15%)	12 (6.8-18)	810 (500-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	1995	Male	30 to 34	60% (39-77%)	40% (23-61%)	2.9 (1.8-4.2)	7.6% (4.7-11%)	9% (5.2-14%)	12 (6.5-20)	800 (460-1200)
Fiji	1995	Male	35 to 39	59% (40-75%)	41% (25-60%)	2.7 (1.6-4)	6.3% (3.1-9.9%)	7.2% (3.1-12%)	12 (4.8-22)	710 (330-1200)
Fiji	1995	Male	40 to 44	56% (38-74%)	44% (26-62%)	2.2 (1.3-3.4)	4.4% (1.3-8.4%)	4.9% (1.1-9.7%)	9.2 (1.7-20)	490 (140-1000)
Fiji	1995	Male	45 to 49	52% (31-70%)	48% (30-69%)	1.7 (0.89-2.8)	2.7% (0.063-6%)	2.7% (-0.41-6.7%)	6.1 (-0.74-16)	300 (8.3-740)
Fiji	1995	Male	50 to 54	49% (30-68%)	51% (32-70%)	1.3 (0.66-2.3)	1.9% (-0.37-5.1%)	2% (-0.67-5.6%)	5 (-1.7-14)	220 (-40-570)
Fiji	1995	Male	55 to 59	47% (28-66%)	53% (34-72%)	1.2 (0.58-2.1)	1.6% (-0.33-4.6%)	1.6% (-0.54-5%)	4.4 (-1.4-13)	160 (-34-460)
Fiji	1995	Male	60 to 64	45% (26-65%)	55% (35-74%)	1 (0.5-1.9)	1.4% (-0.32-4.4%)	1.4% (-0.44-4.8%)	3.9 (-1.3-14)	120 (-28-390)
Fiji	1995	Male	65 to 69	43% (25-62%)	57% (38-75%)	0.9 (0.45-1.6)	1.1% (-0.35-3.3%)	1.1% (-0.45-3.5%)	3 (-1.3-9.3)	76 (-25-220)
Fiji	1995	Male	70 to 74	40% (23-58%)	60% (42-77%)	0.77 (0.37-1.4)	0.8% (-0.5-2.8%)	0.82% (-0.58-3%)	1.9 (-1.3-6.7)	38 (-24-130)
Fiji	1995	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.65 (0.29-1.2)	0.6% (-0.47-2.3%)	0.61% (-0.58-2.4%)	1.3 (-1.2-5.4)	21 (-17-81)
Fiji	1995	Male	80 to 84	33% (18-51%)	67% (49-82%)	0.57 (0.28-1)	0.47% (-0.34-1.7%)	0.46% (-0.45-1.8%)	0.56 (-0.56-2.2)	6.9 (-5-25)
Fiji	1995	Male	85 to 89	32% (18-49%)	68% (51-82%)	0.54 (0.25-0.96)	0.32% (-0.42-1.5%)	0.28% (-0.54-1.5%)	0.19 (-0.36-1)	1.8 (-2.5-8.3)
Fiji	1995	Male	90 to 94	33% (20-50%)	67% (50-80%)	0.55 (0.25-1)	0.17% (-0.5-1.2%)	0.093% (-0.67-1.2%)	0.02 (-0.15-0.26)	0.22 (-0.65-1.5)
Fiji	1995	Male	95 plus	36% (20-57%)	64% (43-80%)	0.6 (0.25-1.1)	0.065% (-0.8-1.3%)	-0.076% (-1.1-1.3%)	-0.0039 (-0.057-0.07)	0.013 (-0.17-0.27)
Fiji	2000	Female	All Ages	7.6% (6-9.7%)	92% (90-94%)	0.17 (0.13-0.23)	0.28% (0.07-0.51%)	0.03% (-0.26-0.38%)	0.81 (-6.8-9.9)	360 (86-660)
Fiji	2000	Female	15 to 19	9.3% (5.8-15%)	91% (85-94%)	0.24 (0.13-0.41)	0.78% (0.44-1.3%)	0.99% (0.41-2%)	0.42 (0.17-0.83)	50 (29-84)
Fiji	2000	Female	20 to 24	11% (6.6-18%)	89% (82-93%)	0.37 (0.21-0.63)	1.4% (0.84-2.1%)	1.4% (0.68-2.7%)	0.77 (0.34-1.5)	94 (56-150)
Fiji	2000	Female	25 to 29	10% (5.9-18%)	90% (82-94%)	0.3 (0.16-0.49)	1.2% (0.85-1.8%)	1.3% (0.69-2.1%)	0.75 (0.39-1.3)	82 (53-120)
Fiji	2000	Female	30 to 34	9.9% (5.5-17%)	90% (83-94%)	0.27 (0.14-0.46)	0.96% (0.62-1.5%)	0.91% (0.47-1.6%)	0.66 (0.31-1.2)	67 (42-110)
Fiji	2000	Female	35 to 39	9.8% (5.2-18%)	90% (82-95%)	0.28 (0.14-0.51)	0.88% (0.48-1.4%)	0.93% (0.41-1.7%)	1 (0.42-1.9)	77 (41-130)
Fiji	2000	Female	40 to 44	9.3% (5.1-16%)	91% (84-95%)	0.25 (0.12-0.45)	0.62% (0.21-1.2%)	0.67% (0.14-1.4%)	0.93 (0.17-2.1)	59 (20-120)
Fiji	2000	Female	45 to 49	9.2% (4.7-16%)	91% (84-95%)	0.19 (0.082-0.35)	0.19% (-0.21-0.65%)	0.19% (-0.29-0.79%)	0.29 (-0.45-1.2)	17 (-20-60)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	2000	Female	50 to 54	9.1% (5.1-16%)	91% (84-95%)	0.16 (0.067-0.32)	-0.11% (-0.58-0.35%)	-0.13% (-0.66-0.43%)	-0.25 (-1.3-0.83)	-10 (-56-34)
Fiji	2000	Female	55 to 59	8.4% (4.5-15%)	92% (85-95%)	0.15 (0.067-0.3)	-0.11% (-0.57-0.37%)	-0.11% (-0.63-0.46%)	-0.25 (-1.4-1.1)	-11 (-56-36)
Fiji	2000	Female	60 to 64	8% (4.1-15%)	92% (85-96%)	0.14 (0.062-0.29)	-0.2% (-0.71-0.29%)	-0.21% (-0.79-0.34%)	-0.53 (-1.9-0.79)	-17 (-59-24)
Fiji	2000	Female	65 to 69	8.8% (4.8-15%)	91% (85-95%)	0.15 (0.074-0.28)	-0.24% (-0.74-0.22%)	-0.25% (-0.82-0.28%)	-0.61 (-2-0.72)	-16 (-51-15)
Fiji	2000	Female	70 to 74	9.3% (5.2-16%)	91% (84-95%)	0.15 (0.071-0.29)	-0.33% (-0.86-0.16%)	-0.36% (-0.96-0.18%)	-0.87 (-2.4-0.43)	-17 (-47-8)
Fiji	2000	Female	75 to 79	9.2% (5-16%)	91% (84-95%)	0.15 (0.062-0.28)	-0.24% (-0.7-0.2%)	-0.25% (-0.77-0.24%)	-0.54 (-1.7-0.49)	-8.4 (-26-7.1)
Fiji	2000	Female	80 to 84	9% (4.9-16%)	91% (84-95%)	0.14 (0.067-0.26)	-0.28% (-0.7-0.098%)	-0.31% (-0.77-0.11%)	-0.55 (-1.4-0.2)	-6 (-15-2)
Fiji	2000	Female	85 to 89	9.2% (4.9-15%)	91% (85-95%)	0.14 (0.068-0.27)	-0.26% (-0.67-0.11%)	-0.3% (-0.76-0.11%)	-0.3 (-0.78-0.11)	-2.3 (-5.9-0.94)
Fiji	2000	Female	90 to 94	10% (5.4-17%)	90% (83-95%)	0.16 (0.068-0.29)	-0.21% (-0.57-0.17%)	-0.25% (-0.66-0.18%)	-0.092 (-0.24-0.068)	-0.46 (-1.2-0.36)
Fiji	2000	Female	95 plus	12% (5.6-21%)	88% (79-94%)	0.18 (0.068-0.36)	-0.21% (-0.71-0.31%)	-0.28% (-0.86-0.31%)	-0.025 (-0.078-0.028)	-0.074 (-0.25-0.11)
Fiji	2000	Male	All Ages	42% (34-50%)	58% (50-66%)	1.3 (1.1-1.5)	3% (1.9-4.2%)	2.6% (1.2-4.2%)	87 (42-140)	4900 (3200-7000)
Fiji	2000	Male	15 to 19	40% (25-56%)	60% (44-75%)	0.92 (0.47-1.5)	2.7% (1.5-4.4%)	3.6% (1.8-6.3%)	2.8 (1.3-4.9)	240 (130-400)
Fiji	2000	Male	20 to 24	59% (42-73%)	41% (27-58%)	2.5 (1.7-3.4)	8.2% (5.2-12%)	10% (6.2-15%)	11 (7-17)	850 (540-1200)
Fiji	2000	Male	25 to 29	64% (48-78%)	36% (22-52%)	3 (2-4.2)	9.2% (6-13%)	11% (7.1-16%)	12 (7.4-18)	850 (540-1200)
Fiji	2000	Male	30 to 34	62% (45-77%)	38% (23-55%)	3 (2-4.2)	7.6% (4.7-11%)	9.1% (5.3-14%)	11 (6.4-17)	730 (440-1100)
Fiji	2000	Male	35 to 39	60% (44-75%)	40% (25-56%)	2.7 (1.7-3.8)	6.3% (3.1-9.9%)	7.3% (3.2-12%)	12 (5.1-20)	710 (350-1200)
Fiji	2000	Male	40 to 44	56% (40-72%)	44% (28-60%)	2.2 (1.3-3.3)	4.3% (1.1-8%)	4.7% (0.87-9.2%)	10 (1.9-21)	550 (150-1000)
Fiji	2000	Male	45 to 49	51% (33-67%)	49% (33-67%)	1.6 (0.94-2.5)	2.4% (0.1-5.5%)	2.5% (-0.23-6%)	6.1 (-0.62-16)	310 (13-720)
Fiji	2000	Male	50 to 54	47% (30-64%)	53% (36-70%)	1.2 (0.63-2)	1.7% (-0.37-4.5%)	1.7% (-0.7-4.9%)	4.9 (-2-15)	220 (-49-590)
Fiji	2000	Male	55 to 59	44% (28-61%)	56% (39-72%)	1.1 (0.55-1.8)	1.4% (-0.27-4.1%)	1.4% (-0.46-4.5%)	4.6 (-1.5-14)	180 (-34-510)
Fiji	2000	Male	60 to 64	42% (25-60%)	58% (40-75%)	0.91 (0.46-1.7)	1.2% (-0.32-3.8%)	1.2% (-0.43-4.1%)	4.1 (-1.4-14)	130 (-33-410)
Fiji	2000	Male	65 to 69	41% (24-60%)	59% (40-76%)	0.84 (0.43-1.5)	0.97% (-0.39-3%)	0.99% (-0.5-3.2%)	3.1 (-1.6-9.8)	78 (-33-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	2000	Male	70 to 74	40% (23-57%)	60% (43-77%)	0.74 (0.35-1.3)	0.69% (-0.55-2.4%)	0.69% (-0.66-2.6%)	1.9 (-1.9-7.6)	40 (-34-150)
Fiji	2000	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.63 (0.29-1.2)	0.52% (-0.51-2.1%)	0.51% (-0.62-2.2%)	1.1 (-1.3-4.7)	17 (-17-72)
Fiji	2000	Male	80 to 84	34% (18-52%)	66% (48-82%)	0.56 (0.27-1)	0.41% (-0.38-1.6%)	0.39% (-0.47-1.6%)	0.54 (-0.66-2.3)	6.7 (-6.3-26)
Fiji	2000	Male	85 to 89	32% (18-49%)	68% (51-82%)	0.53 (0.25-0.95)	0.27% (-0.46-1.4%)	0.22% (-0.6-1.5%)	0.17 (-0.48-1.1)	1.8 (-3.2-9.1)
Fiji	2000	Male	90 to 94	34% (20-51%)	66% (49-80%)	0.55 (0.25-1)	0.13% (-0.56-1.2%)	0.055% (-0.74-1.2%)	0.015 (-0.21-0.34)	0.22 (-0.95-2)
Fiji	2000	Male	95 plus	37% (20-57%)	63% (43-80%)	0.6 (0.24-1.2)	0.039% (-0.85-1.3%)	-0.1% (-1.1-1.3%)	-0.007 (-0.078-0.086)	0.01 (-0.23-0.35)
Fiji	2005	Female	All Ages	7.6% (6.1-9.7%)	92% (90-94%)	0.19 (0.14-0.25)	0.27% (0.045-0.53%)	0.028% (-0.3-0.41%)	0.83 (-8.7-12)	370 (58-740)
Fiji	2005	Female	15 to 19	9% (5.6-15%)	91% (85-94%)	0.25 (0.14-0.44)	0.81% (0.45-1.3%)	1.1% (0.44-2.1%)	0.38 (0.16-0.78)	45 (25-75)
Fiji	2005	Female	20 to 24	11% (6.6-18%)	89% (82-93%)	0.4 (0.21-0.65)	1.4% (0.89-2.2%)	1.6% (0.73-2.8%)	0.75 (0.33-1.4)	94 (56-150)
Fiji	2005	Female	25 to 29	10% (5.8-18%)	90% (82-94%)	0.32 (0.18-0.53)	1.3% (0.86-1.8%)	1.3% (0.71-2.2%)	0.75 (0.38-1.3)	86 (56-130)
Fiji	2005	Female	30 to 34	9.7% (5.4-17%)	90% (83-95%)	0.29 (0.15-0.5)	1% (0.64-1.5%)	0.95% (0.5-1.7%)	0.61 (0.3-1.1)	65 (40-100)
Fiji	2005	Female	35 to 39	9.6% (5-17%)	90% (83-95%)	0.3 (0.15-0.54)	0.91% (0.51-1.5%)	0.98% (0.45-1.8%)	0.89 (0.37-1.7)	72 (38-120)
Fiji	2005	Female	40 to 44	9% (5-15%)	91% (85-95%)	0.26 (0.13-0.47)	0.66% (0.23-1.2%)	0.73% (0.18-1.5%)	1.1 (0.26-2.3)	70 (24-140)
Fiji	2005	Female	45 to 49	9% (4.6-16%)	91% (84-95%)	0.2 (0.091-0.37)	0.23% (-0.21-0.72%)	0.24% (-0.27-0.84%)	0.46 (-0.51-1.6)	26 (-24-83)
Fiji	2005	Female	50 to 54	9% (5-16%)	91% (84-95%)	0.17 (0.073-0.35)	-0.074% (-0.58-0.45%)	-0.093% (-0.67-0.52%)	-0.21 (-1.6-1.2)	-8.2 (-66-49)
Fiji	2005	Female	55 to 59	8.3% (4.5-15%)	92% (85-95%)	0.17 (0.074-0.32)	-0.086% (-0.56-0.43%)	-0.08% (-0.63-0.51%)	-0.21 (-1.7-1.4)	-9.3 (-63-47)
Fiji	2005	Female	60 to 64	7.9% (4.1-15%)	92% (85-96%)	0.16 (0.068-0.31)	-0.18% (-0.72-0.37%)	-0.19% (-0.79-0.43%)	-0.6 (-2.5-1.4)	-19 (-76-41)
Fiji	2005	Female	65 to 69	8.8% (4.8-16%)	91% (84-95%)	0.17 (0.079-0.31)	-0.22% (-0.75-0.29%)	-0.23% (-0.82-0.34%)	-0.67 (-2.4-1)	-18 (-61-24)
Fiji	2005	Female	70 to 74	9.2% (5.2-16%)	91% (84-95%)	0.17 (0.078-0.33)	-0.3% (-0.84-0.24%)	-0.33% (-0.94-0.28%)	-0.96 (-2.7-0.81)	-19 (-52-15)
Fiji	2005	Female	75 to 79	9% (4.9-16%)	91% (84-95%)	0.16 (0.069-0.3)	-0.2% (-0.7-0.3%)	-0.21% (-0.77-0.35%)	-0.5 (-1.8-0.83)	-8 (-28-12)
Fiji	2005	Female	80 to 84	8.9% (4.8-16%)	91% (84-95%)	0.15 (0.072-0.28)	-0.25% (-0.66-0.19%)	-0.28% (-0.75-0.2%)	-0.52 (-1.4-0.37)	-5.6 (-15-4.4)
Fiji	2005	Female	85 to 89	9.1% (4.9-15%)	91% (85-95%)	0.16 (0.073-0.29)	-0.23% (-0.64-0.19%)	-0.27% (-0.72-0.2%)	-0.3 (-0.82-0.24)	-2.2 (-6.2-1.9)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	2005	Female	90 to 94	9.9% (5.5-16%)	90% (84-94%)	0.17 (0.074-0.33)	-0.19% (-0.58-0.25%)	-0.23% (-0.67-0.27%)	-0.098 (-0.29-0.11)	-0.48 (-1.5-0.61)
Fiji	2005	Female	95 plus	11% (5.6-20%)	89% (80-94%)	0.19 (0.076-0.39)	-0.19% (-0.74-0.37%)	-0.26% (-0.89-0.37%)	-0.027 (-0.09-0.038)	-0.075 (-0.29-0.15)
Fiji	2005	Male	All Ages	43% (36-51%)	57% (49-64%)	1.5 (1.3-1.8)	3.2% (2.1-4.5%)	2.7% (1.4-4.4%)	100 (48-170)	5500 (3400-8100)
Fiji	2005	Male	15 to 19	40% (25-56%)	60% (44-75%)	1 (0.52-1.7)	2.9% (1.6-4.8%)	4% (2-6.7%)	2.4 (1.2-4.3)	200 (110-340)
Fiji	2005	Male	20 to 24	59% (42-73%)	41% (27-58%)	2.7 (1.8-3.8)	8.7% (5.4-12%)	11% (6.4-16%)	11 (6.4-17)	830 (500-1200)
Fiji	2005	Male	25 to 29	64% (48-78%)	36% (22-52%)	3.3 (2.2-4.5)	9.8% (6.4-14%)	12% (7.8-18%)	12 (7.4-18)	880 (550-1300)
Fiji	2005	Male	30 to 34	62% (44-77%)	38% (23-56%)	3.3 (2.1-4.6)	8.4% (5.3-12%)	10% (6.2-15%)	11 (6.2-16)	710 (440-1000)
Fiji	2005	Male	35 to 39	60% (44-75%)	40% (25-56%)	2.9 (1.9-4.2)	7.1% (4-10%)	8.5% (4.5-13%)	12 (5.7-19)	700 (390-1100)
Fiji	2005	Male	40 to 44	56% (40-72%)	44% (28-60%)	2.4 (1.5-3.5)	5.1% (1.8-8.7%)	5.7% (1.5-10%)	12 (3.4-22)	660 (240-1100)
Fiji	2005	Male	45 to 49	50% (33-66%)	50% (34-67%)	1.7 (1-2.6)	3.1% (0.58-6.2%)	3.2% (0.25-6.9%)	9.5 (0.7-22)	480 (87-1000)
Fiji	2005	Male	50 to 54	46% (30-64%)	54% (36-70%)	1.3 (0.68-2.1)	2.2% (-0.045-5.1%)	2.2% (-0.3-5.7%)	7.3 (-1-20)	320 (-6.8-810)
Fiji	2005	Male	55 to 59	44% (28-61%)	56% (39-72%)	1.1 (0.6-1.9)	1.8% (-0.1-4.6%)	1.9% (-0.25-5%)	6.9 (-0.97-19)	260 (-15-680)
Fiji	2005	Male	60 to 64	42% (26-61%)	58% (39-74%)	0.99 (0.5-1.8)	1.6% (-0.16-4.1%)	1.6% (-0.27-4.5%)	6.7 (-1.2-20)	210 (-23-580)
Fiji	2005	Male	65 to 69	41% (24-60%)	59% (40-76%)	0.92 (0.47-1.6)	1.3% (-0.18-3.6%)	1.3% (-0.3-3.8%)	5.2 (-1.1-15)	130 (-18-360)
Fiji	2005	Male	70 to 74	39% (22-57%)	61% (43-78%)	0.8 (0.37-1.5)	0.94% (-0.45-3%)	0.96% (-0.56-3.1%)	3.3 (-2-11)	67 (-35-210)
Fiji	2005	Male	75 to 79	36% (21-54%)	64% (46-79%)	0.67 (0.3-1.3)	0.71% (-0.43-2.5%)	0.72% (-0.5-2.6%)	2 (-1.5-7.4)	32 (-19-110)
Fiji	2005	Male	80 to 84	33% (18-51%)	67% (49-82%)	0.6 (0.29-1.1)	0.56% (-0.31-1.9%)	0.55% (-0.38-1.9%)	0.9 (-0.61-3.2)	11 (-5.9-36)
Fiji	2005	Male	85 to 89	31% (18-48%)	69% (52-82%)	0.56 (0.26-1)	0.42% (-0.37-1.6%)	0.4% (-0.48-1.7%)	0.34 (-0.41-1.5)	3.1 (-2.7-12)
Fiji	2005	Male	90 to 94	32% (20-49%)	68% (51-80%)	0.57 (0.27-1)	0.27% (-0.44-1.4%)	0.2% (-0.6-1.5%)	0.066 (-0.19-0.47)	0.51 (-0.86-2.7)
Fiji	2005	Male	95 plus	36% (19-56%)	64% (44-81%)	0.62 (0.25-1.2)	0.2% (-0.71-1.6%)	0.078% (-0.94-1.6%)	0.0066 (-0.079-0.14)	0.066 (-0.23-0.53)
Fiji	2010	Female	All Ages	8.4% (6.7-11%)	92% (89-93%)	0.26 (0.19-0.35)	0.39% (0.082-0.79%)	0.18% (-0.26-0.79%)	5.6 (-8.2-25)	560 (120-1100)
Fiji	2010	Female	15 to 19	9% (5.2-15%)	91% (85-95%)	0.31 (0.15-0.56)	0.95% (0.52-1.7%)	1.3% (0.52-2.7%)	0.5 (0.2-1.1)	56 (30-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	2010	Female	20 to 24	12% (6.2-19%)	88% (81-94%)	0.51 (0.25-0.9)	1.7% (1-2.6%)	2.1% (0.92-3.9%)	0.84 (0.36-1.6)	100 (60-160)
Fiji	2010	Female	25 to 29	12% (6.4-20%)	88% (80-94%)	0.45 (0.24-0.74)	1.6% (1-2.3%)	1.8% (0.94-3%)	0.96 (0.5-1.7)	110 (67-160)
Fiji	2010	Female	30 to 34	11% (5.8-19%)	89% (81-94%)	0.41 (0.21-0.71)	1.3% (0.78-1.9%)	1.4% (0.69-2.5%)	0.87 (0.43-1.6)	88 (53-140)
Fiji	2010	Female	35 to 39	11% (5.5-19%)	89% (81-94%)	0.41 (0.2-0.74)	1.2% (0.68-1.9%)	1.4% (0.65-2.6%)	1 (0.46-2)	83 (45-140)
Fiji	2010	Female	40 to 44	10% (5.5-18%)	90% (82-95%)	0.36 (0.17-0.64)	0.93% (0.36-1.8%)	1.1% (0.33-2.3%)	1.3 (0.4-2.9)	84 (34-170)
Fiji	2010	Female	45 to 49	10% (4.9-18%)	90% (82-95%)	0.27 (0.12-0.52)	0.42% (-0.11-1.2%)	0.51% (-0.2-1.4%)	0.9 (-0.36-2.5)	48 (-13-130)
Fiji	2010	Female	50 to 54	10% (5.4-18%)	90% (82-95%)	0.24 (0.097-0.46)	0.072% (-0.56-0.88%)	0.093% (-0.64-1.1%)	0.23 (-1.8-3)	9.3 (-75-120)
Fiji	2010	Female	55 to 59	9.1% (5.1-16%)	91% (84-95%)	0.22 (0.098-0.43)	0.046% (-0.56-0.83%)	0.077% (-0.63-1%)	0.22 (-1.9-2.9)	5.6 (-72-100)
Fiji	2010	Female	60 to 64	8.4% (4.2-15%)	92% (85-96%)	0.2 (0.089-0.4)	-0.041% (-0.67-0.73%)	-0.035% (-0.75-0.85%)	-0.13 (-2.8-3)	-5.2 (-82-88)
Fiji	2010	Female	65 to 69	8.7% (4.7-15%)	91% (85-95%)	0.2 (0.092-0.38)	-0.058% (-0.66-0.64%)	-0.047% (-0.73-0.75%)	-0.18 (-2.7-2.8)	-6.1 (-68-67)
Fiji	2010	Female	70 to 74	9% (5.2-15%)	91% (85-95%)	0.2 (0.091-0.4)	-0.12% (-0.71-0.63%)	-0.13% (-0.8-0.73%)	-0.49 (-3-2.6)	-10 (-57-52)
Fiji	2010	Female	75 to 79	8.9% (4.9-15%)	91% (85-95%)	0.19 (0.08-0.37)	-0.028% (-0.56-0.71%)	-0.021% (-0.64-0.84%)	-0.069 (-2-2.4)	-1.5 (-30-37)
Fiji	2010	Female	80 to 84	8.7% (4.7-15%)	91% (85-95%)	0.19 (0.084-0.35)	-0.084% (-0.55-0.52%)	-0.098% (-0.63-0.58%)	-0.22 (-1.4-1.4)	-2.4 (-15-15)
Fiji	2010	Female	85 to 89	8.9% (4.7-15%)	91% (85-95%)	0.19 (0.085-0.36)	-0.079% (-0.52-0.46%)	-0.1% (-0.6-0.49%)	-0.12 (-0.73-0.61)	-0.83 (-5.5-4.8)
Fiji	2010	Female	90 to 94	9.7% (5.4-16%)	90% (84-95%)	0.2 (0.09-0.38)	-0.062% (-0.47-0.5%)	-0.093% (-0.55-0.53%)	-0.043 (-0.26-0.25)	-0.17 (-1.3-1.4)
Fiji	2010	Female	95 plus	11% (5.5-20%)	89% (80-95%)	0.23 (0.087-0.46)	-0.057% (-0.6-0.61%)	-0.12% (-0.73-0.62%)	-0.015 (-0.088-0.076)	-0.027 (-0.28-0.29)
Fiji	2010	Male	All Ages	41% (33-50%)	59% (50-67%)	1.7 (1.4-2.1)	3.7% (2.4-5.2%)	3.4% (1.6-5.5%)	140 (63-220)	6500 (4000-9300)
Fiji	2010	Male	15 to 19	32% (18-50%)	68% (50-82%)	0.96 (0.47-1.7)	2.7% (1.5-4.8%)	3.9% (1.8-7.4%)	2 (0.88-3.8)	180 (94-320)
Fiji	2010	Male	20 to 24	50% (32-68%)	50% (32-68%)	2.6 (1.5-4)	8.5% (5.1-12%)	11% (6.3-17%)	9.1 (4.8-15)	700 (400-1100)
Fiji	2010	Male	25 to 29	58% (38-75%)	42% (25-62%)	3.5 (2.1-5.1)	9.9% (6.6-13%)	13% (8.4-18%)	12 (7.1-17)	850 (550-1200)
Fiji	2010	Male	30 to 34	59% (37-77%)	41% (23-63%)	3.7 (2.2-5.4)	9.1% (5.9-12%)	12% (7.3-16%)	11 (6.6-17)	760 (480-1100)
Fiji	2010	Male	35 to 39	58% (39-76%)	42% (24-61%)	3.4 (1.9-5.1)	8.1% (4.7-12%)	10% (5.5-15%)	12 (6.1-18)	700 (410-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	2010	Male	40 to 44	55% (37-73%)	45% (27-63%)	2.8 (1.6-4.4)	6.4% (2.9-10%)	7.5% (3-13%)	13 (5-22)	690 (320-1100)
Fiji	2010	Male	45 to 49	51% (31-71%)	49% (29-69%)	2.1 (1.1-3.4)	4.5% (1.4-8.2%)	4.9% (1.2-9.4%)	13 (3.3-27)	650 (200-1200)
Fiji	2010	Male	50 to 54	48% (29-68%)	52% (32-71%)	1.7 (0.85-2.9)	3.4% (0.51-7.3%)	3.7% (0.28-8.4%)	14 (1.1-31)	580 (88-1200)
Fiji	2010	Male	55 to 59	46% (28-66%)	54% (34-72%)	1.5 (0.74-2.7)	3% (0.4-6.6%)	3.2% (0.26-7.3%)	14 (1.1-30)	480 (63-1000)
Fiji	2010	Male	60 to 64	43% (25-63%)	57% (37-75%)	1.3 (0.61-2.4)	2.5% (0.25-5.9%)	2.7% (0.17-6.5%)	13 (0.77-32)	390 (39-930)
Fiji	2010	Male	65 to 69	42% (24-61%)	58% (39-76%)	1.1 (0.56-1.9)	2.1% (0.15-4.9%)	2.2% (0.11-5.3%)	10 (0.49-24)	260 (17-580)
Fiji	2010	Male	70 to 74	39% (23-56%)	61% (44-77%)	0.96 (0.45-1.8)	1.5% (-0.18-4.3%)	1.6% (-0.26-4.6%)	6.8 (-1.1-19)	130 (-15-370)
Fiji	2010	Male	75 to 79	35% (21-53%)	65% (47-79%)	0.8 (0.35-1.5)	1.2% (-0.25-3.5%)	1.2% (-0.32-3.8%)	4.1 (-1.1-12)	63 (-14-180)
Fiji	2010	Male	80 to 84	33% (18-51%)	67% (49-82%)	0.72 (0.33-1.3)	0.97% (-0.12-2.8%)	1% (-0.19-3%)	2 (-0.37-5.9)	23 (-2.7-66)
Fiji	2010	Male	85 to 89	31% (18-47%)	69% (53-82%)	0.68 (0.31-1.3)	0.82% (-0.18-2.5%)	0.84% (-0.27-2.7%)	0.79 (-0.25-2.4)	6.6 (-1.5-19)
Fiji	2010	Male	90 to 94	32% (19-49%)	68% (51-81%)	0.69 (0.32-1.3)	0.59% (-0.27-2%)	0.57% (-0.43-2.2%)	0.19 (-0.13-0.7)	1.2 (-0.54-4)
Fiji	2010	Male	95 plus	35% (18-56%)	65% (44-82%)	0.75 (0.3-1.4)	0.55% (-0.5-2.4%)	0.47% (-0.7-2.5%)	0.042 (-0.063-0.21)	0.19 (-0.18-0.79)
Fiji	2016	Female	All Ages	8.4% (6.7-11%)	92% (89-93%)	0.28 (0.18-0.43)	0.47% (0.046-1.1%)	0.28% (-0.27-1.1%)	10 (-10-41)	730 (76-1700)
Fiji	2016	Female	15 to 19	9% (5.2-15%)	91% (85-95%)	0.34 (0.15-0.67)	1.1% (0.53-2%)	1.5% (0.54-3.3%)	0.53 (0.18-1.2)	57 (27-100)
Fiji	2016	Female	20 to 24	12% (6.4-19%)	88% (81-94%)	0.57 (0.26-1)	1.9% (1.1-3.1%)	2.4% (0.93-4.6%)	0.98 (0.34-2.1)	110 (58-190)
Fiji	2016	Female	25 to 29	12% (6.2-20%)	88% (80-94%)	0.49 (0.24-0.86)	1.7% (1-2.7%)	2% (0.98-3.7%)	1 (0.42-2.1)	110 (63-180)
Fiji	2016	Female	30 to 34	12% (5.9-20%)	88% (80-94%)	0.45 (0.21-0.82)	1.4% (0.81-2.3%)	1.6% (0.71-3%)	1.1 (0.41-2.4)	110 (57-190)
Fiji	2016	Female	35 to 39	11% (5.8-19%)	89% (81-94%)	0.45 (0.21-0.82)	1.4% (0.71-2.4%)	1.6% (0.69-3.2%)	1.5 (0.5-3.2)	110 (53-210)
Fiji	2016	Female	40 to 44	10% (5.5-18%)	90% (82-94%)	0.39 (0.17-0.78)	1.1% (0.38-2.3%)	1.3% (0.34-2.9%)	1.6 (0.35-3.7)	96 (33-210)
Fiji	2016	Female	45 to 49	10% (5.1-17%)	90% (83-95%)	0.3 (0.12-0.59)	0.55% (-0.075-1.6%)	0.66% (-0.16-2%)	1.1 (-0.24-3.5)	59 (-7.5-170)
Fiji	2016	Female	50 to 54	10% (5.5-18%)	90% (82-94%)	0.25 (0.1-0.51)	0.2% (-0.48-1.2%)	0.25% (-0.57-1.5%)	0.66 (-1.7-3.7)	28 (-71-160)
Fiji	2016	Female	55 to 59	9.1% (5.1-16%)	91% (84-95%)	0.24 (0.099-0.48)	0.17% (-0.52-1.1%)	0.22% (-0.57-1.4%)	0.81 (-2.2-5.1)	26 (-86-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	2016	Female	60 to 64	8.4% (4.3-15%)	92% (85-96%)	0.22 (0.087-0.46)	0.083% (-0.61-1%)	0.11% (-0.67-1.2%)	0.43 (-3.1-5.3)	12 (-92-160)
Fiji	2016	Female	65 to 69	8.9% (4.6-16%)	91% (84-95%)	0.22 (0.092-0.43)	0.058% (-0.61-0.96%)	0.085% (-0.67-1.1%)	0.35 (-3.1-4.6)	6.9 (-77-110)
Fiji	2016	Female	70 to 74	9.2% (5.1-16%)	91% (84-95%)	0.22 (0.089-0.46)	-0.016% (-0.66-0.97%)	-0.013% (-0.75-1.1%)	-0.088 (-3.9-5.3)	-2.1 (-76-100)
Fiji	2016	Female	75 to 79	8.9% (5-16%)	91% (84-95%)	0.21 (0.082-0.44)	0.072% (-0.56-0.94%)	0.09% (-0.6-1.1%)	0.36 (-2.6-4.6)	4.8 (-39-67)
Fiji	2016	Female	80 to 84	8.8% (4.7-15%)	91% (85-95%)	0.21 (0.089-0.4)	0.0092% (-0.53-0.84%)	0.005% (-0.61-0.95%)	0.0095 (-2-2.9)	0.3 (-22-32)
Fiji	2016	Female	85 to 89	9% (4.9-15%)	91% (85-95%)	0.21 (0.089-0.4)	0.0038% (-0.51-0.76%)	-0.012% (-0.58-0.84%)	-0.023 (-1.1-1.5)	0.041 (-7.7-11)
Fiji	2016	Female	90 to 94	9.7% (5.5-16%)	90% (84-95%)	0.22 (0.093-0.44)	0.017% (-0.45-0.78%)	-0.007% (-0.55-0.84%)	-0.0045 (-0.32-0.51)	0.058 (-1.6-2.7)
Fiji	2016	Female	95 plus	11% (5.7-20%)	89% (80-94%)	0.25 (0.093-0.54)	0.021% (-0.59-0.98%)	-0.033% (-0.76-1%)	-0.0051 (-0.12-0.16)	0.012 (-0.36-0.57)
Fiji	2016	Male	All Ages	41% (33-50%)	59% (50-67%)	1.9 (1.3-2.7)	4.3% (2.3-6.5%)	3.9% (1.6-6.6%)	180 (67-340)	8100 (4100-14000)
Fiji	2016	Male	15 to 19	33% (19-50%)	67% (50-81%)	1.1 (0.47-2)	3.2% (1.5-6%)	4.4% (1.9-8.8%)	2.4 (0.91-4.9)	210 (95-390)
Fiji	2016	Male	20 to 24	50% (32-69%)	50% (31-68%)	2.9 (1.5-4.8)	9.5% (5.3-14%)	12% (6.5-19%)	11 (5.2-19)	840 (440-1400)
Fiji	2016	Male	25 to 29	58% (37-74%)	42% (26-63%)	3.8 (2.2-6.1)	11% (6.8-15%)	14% (8.5-20%)	13 (6.7-21)	950 (510-1400)
Fiji	2016	Male	30 to 34	59% (38-76%)	41% (24-62%)	4.1 (2.3-6.4)	10% (6.3-14%)	13% (7.5-18%)	15 (7.6-25)	980 (540-1500)
Fiji	2016	Male	35 to 39	58% (40-76%)	42% (24-60%)	3.7 (2.1-6.2)	8.9% (5-13%)	11% (5.7-16%)	16 (6.8-28)	950 (450-1600)
Fiji	2016	Male	40 to 44	55% (37-73%)	45% (27-63%)	3.1 (1.6-5.4)	7.1% (3.3-11%)	8.3% (3.4-14%)	15 (5.4-28)	800 (340-1400)
Fiji	2016	Male	45 to 49	51% (29-70%)	49% (30-71%)	2.4 (1.1-4.2)	5.2% (1.6-9.5%)	5.8% (1.3-11%)	15 (2.9-31)	710 (200-1400)
Fiji	2016	Male	50 to 54	48% (28-68%)	52% (32-72%)	1.9 (0.87-3.4)	4.1% (0.71-8.5%)	4.4% (0.45-9.3%)	18 (1.9-41)	740 (140-1600)
Fiji	2016	Male	55 to 59	46% (28-66%)	54% (34-72%)	1.7 (0.77-3.1)	3.6% (0.55-7.5%)	3.8% (0.41-8.4%)	19 (1.7-47)	700 (100-1600)
Fiji	2016	Male	60 to 64	44% (26-64%)	56% (36-74%)	1.4 (0.61-2.7)	3% (0.27-6.7%)	3.2% (0.19-7.4%)	17 (1-40)	510 (47-1200)
Fiji	2016	Male	65 to 69	42% (24-61%)	58% (39-76%)	1.2 (0.56-2.2)	2.6% (0.28-5.9%)	2.8% (0.24-6.4%)	15 (1.2-36)	360 (39-850)
Fiji	2016	Male	70 to 74	39% (23-57%)	61% (43-77%)	1.1 (0.46-2)	2% (-0.058-5.2%)	2.1% (-0.11-5.6%)	11 (-0.65-29)	220 (-6.4-570)
Fiji	2016	Male	75 to 79	35% (20-54%)	65% (46-80%)	0.89 (0.36-1.8)	1.6% (-0.12-4.4%)	1.6% (-0.19-4.7%)	6.8 (-0.84-20)	100 (-8.5-300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Fiji	2016	Male	80 to 84	32% (17-50%)	68% (50-83%)	0.79 (0.34-1.5)	1.3% (-0.031-3.4%)	1.3% (-0.086-3.7%)	3.5 (-0.22-9.8)	40 (-0.92-110)
Fiji	2016	Male	85 to 89	31% (17-47%)	69% (53-83%)	0.75 (0.34-1.4)	1.1% (-0.1-3.2%)	1.2% (-0.17-3.4%)	1.4 (-0.2-4.1)	12 (-1-31)
Fiji	2016	Male	90 to 94	32% (19-50%)	68% (50-81%)	0.77 (0.34-1.4)	0.86% (-0.2-2.7%)	0.86% (-0.32-2.9%)	0.33 (-0.12-1.1)	2 (-0.49-6.2)
Fiji	2016	Male	95 plus	36% (18-56%)	64% (44-82%)	0.84 (0.32-1.8)	0.85% (-0.47-3.1%)	0.79% (-0.65-3.3%)	0.069 (-0.055-0.29)	0.29 (-0.15-1.1)
Kiribati	1990	Female	All Ages	7.4% (5.8-9.6%)	93% (90-94%)	0.14 (0.11-0.18)	0.39% (0.24-0.57%)	0.41% (0.16-0.71%)	1.5 (0.59-2.6)	84 (52-120)
Kiribati	1990	Female	15 to 19	9.1% (5.1-16%)	91% (84-95%)	0.18 (0.092-0.32)	0.95% (0.44-1.9%)	1.3% (0.42-2.7%)	0.06 (0.019-0.13)	5.7 (2.6-11)
Kiribati	1990	Female	20 to 24	12% (6.4-19%)	88% (81-94%)	0.31 (0.16-0.5)	1.7% (0.87-2.7%)	2% (0.8-3.6%)	0.16 (0.066-0.3)	15 (7.8-25)
Kiribati	1990	Female	25 to 29	12% (6.5-21%)	88% (79-94%)	0.3 (0.17-0.47)	1.5% (0.93-2.3%)	1.7% (0.9-2.9%)	0.15 (0.076-0.25)	13 (7.9-19)
Kiribati	1990	Female	30 to 34	12% (6.1-20%)	88% (80-94%)	0.29 (0.15-0.49)	1.4% (0.81-2.2%)	1.6% (0.79-2.7%)	0.16 (0.078-0.29)	12 (6.9-19)
Kiribati	1990	Female	35 to 39	11% (5.9-19%)	89% (81-94%)	0.27 (0.14-0.49)	1.3% (0.74-2.2%)	1.6% (0.76-2.8%)	0.16 (0.079-0.3)	10 (5.7-17)
Kiribati	1990	Female	40 to 44	10% (5.6-18%)	90% (82-94%)	0.24 (0.12-0.42)	1.3% (0.58-2.3%)	1.6% (0.64-2.9%)	0.18 (0.076-0.34)	9.7 (4.5-17)
Kiribati	1990	Female	45 to 49	10% (5-18%)	90% (82-95%)	0.21 (0.095-0.39)	0.72% (0.22-1.4%)	0.9% (0.23-1.9%)	0.11 (0.03-0.24)	5.2 (1.5-10)
Kiribati	1990	Female	50 to 54	10% (5.4-18%)	90% (82-95%)	0.19 (0.088-0.35)	0.55% (-0.045-1.4%)	0.71% (0.01-1.8%)	0.12 (0.0017-0.31)	4.5 (-0.34-12)
Kiribati	1990	Female	55 to 59	9.2% (5.1-16%)	91% (84-95%)	0.16 (0.073-0.3)	0.4% (-0.11-1.1%)	0.53% (-0.098-1.4%)	0.093 (-0.019-0.24)	2.8 (-0.81-7.5)
Kiribati	1990	Female	60 to 64	8.4% (4.3-15%)	92% (85-96%)	0.14 (0.062-0.27)	0.25% (-0.19-0.78%)	0.34% (-0.16-0.98%)	0.09 (-0.042-0.27)	2.2 (-1.6-7.1)
Kiribati	1990	Female	65 to 69	8.8% (4.8-15%)	91% (85-95%)	0.14 (0.064-0.25)	0.22% (-0.21-0.78%)	0.3% (-0.2-0.92%)	0.079 (-0.05-0.25)	1.5 (-1.4-5.4)
Kiribati	1990	Female	70 to 74	9.1% (5.3-15%)	91% (85-95%)	0.14 (0.063-0.26)	0.16% (-0.3-0.81%)	0.22% (-0.32-0.98%)	0.069 (-0.1-0.32)	1.1 (-2-5.4)
Kiribati	1990	Female	75 to 79	9% (4.9-16%)	91% (84-95%)	0.13 (0.059-0.24)	0.21% (-0.21-0.71%)	0.26% (-0.21-0.84%)	0.075 (-0.064-0.24)	0.96 (-0.96-3.3)
Kiribati	1990	Female	80 to 84	8.8% (4.8-15%)	91% (85-95%)	0.12 (0.06-0.22)	0.07% (-0.2-0.37%)	0.089% (-0.21-0.42%)	0.018 (-0.044-0.085)	0.17 (-0.5-0.87)
Kiribati	1990	Female	85 to 89	9% (4.8-15%)	91% (85-95%)	0.13 (0.058-0.23)	-0.0034% (-0.24-0.23%)	-0.00054% (-0.25-0.26%)	-0.00005 (-0.026-0.026)	-0.0028 (-0.2-0.2)
Kiribati	1990	Female	90 to 94	9.7% (5.4-16%)	90% (84-95%)	0.14 (0.06-0.25)	0.03% (-0.17-0.26%)	0.034% (-0.19-0.29%)	0.0014 (-0.0075-0.012)	0.007 (-0.041-0.062)
Kiribati	1990	Female	95 plus	11% (5.4-20%)	89% (80-95%)	0.15 (0.06-0.31)	0.11% (-0.18-0.44%)	0.098% (-0.23-0.46%)	0.0012 (-0.0029-0.0057)	0.0055 (-0.0084-0.021)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	1990	Male	All Ages	39% (31-48%)	61% (52-69%)	0.76 (0.68-0.84)	2.3% (1.6-3%)	2.7% (1.7-3.8%)	12 (7.5-17)	580 (400-750)
Kiribati	1990	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.51 (0.25-0.83)	2.1% (0.84-4%)	2.5% (0.85-5.2%)	0.2 (0.069-0.42)	17 (6.8-32)
Kiribati	1990	Male	20 to 24	56% (37-73%)	44% (27-63%)	1.4 (0.84-2)	7.2% (4-11%)	8.5% (4.4-13%)	1.2 (0.61-1.8)	85 (47-130)
Kiribati	1990	Male	25 to 29	63% (44-79%)	37% (21-56%)	1.9 (1.3-2.6)	9.1% (5.8-13%)	11% (6.7-15%)	1.5 (0.91-2.2)	97 (61-140)
Kiribati	1990	Male	30 to 34	64% (43-80%)	36% (20-57%)	2 (1.2-2.9)	8.8% (5.5-13%)	10% (6.4-15%)	1.5 (0.89-2.2)	90 (55-130)
Kiribati	1990	Male	35 to 39	62% (44-79%)	38% (21-56%)	1.8 (1.1-2.6)	6.3% (3.5-9.6%)	7.2% (3.9-11%)	1.1 (0.57-1.7)	60 (33-93)
Kiribati	1990	Male	40 to 44	60% (41-75%)	40% (25-59%)	1.5 (0.85-2.4)	5.8% (2.8-9.4%)	6.6% (3.1-11%)	1.2 (0.54-2)	58 (28-95)
Kiribati	1990	Male	45 to 49	56% (34-74%)	44% (26-66%)	1.3 (0.71-2.2)	5% (2.2-8.6%)	5.6% (2.4-9.7%)	1.2 (0.5-2.1)	52 (23-91)
Kiribati	1990	Male	50 to 54	53% (34-72%)	47% (28-66%)	1.1 (0.56-1.9)	3.8% (1.2-7.3%)	4.2% (1.3-8.1%)	1.1 (0.32-2.1)	41 (13-79)
Kiribati	1990	Male	55 to 59	51% (33-71%)	49% (29-67%)	0.9 (0.47-1.6)	2.6% (0.61-5.2%)	2.8% (0.6-5.8%)	0.76 (0.16-1.6)	25 (5.8-52)
Kiribati	1990	Male	60 to 64	48% (29-69%)	52% (31-71%)	0.75 (0.37-1.4)	1.9% (0.29-4.5%)	2.1% (0.28-4.9%)	0.68 (0.088-1.6)	19 (2.7-44)
Kiribati	1990	Male	65 to 69	46% (27-65%)	54% (35-73%)	0.68 (0.36-1.1)	2.2% (0.52-4.7%)	2.4% (0.58-5.2%)	0.73 (0.17-1.6)	16 (3.8-35)
Kiribati	1990	Male	70 to 74	44% (26-61%)	56% (39-74%)	0.6 (0.3-1)	1.5% (0.17-3.6%)	1.7% (0.17-4%)	0.49 (0.05-1.1)	8.9 (1-21)
Kiribati	1990	Male	75 to 79	40% (24-59%)	60% (41-76%)	0.51 (0.25-0.92)	1.3% (0.17-3%)	1.4% (0.18-3.3%)	0.32 (0.039-0.74)	4.5 (0.57-10)
Kiribati	1990	Male	80 to 84	37% (21-56%)	63% (44-79%)	0.46 (0.24-0.77)	0.84% (0.12-1.9%)	0.92% (0.13-2%)	0.12 (0.017-0.27)	1.3 (0.19-2.9)
Kiribati	1990	Male	85 to 89	36% (21-53%)	64% (47-79%)	0.43 (0.21-0.76)	0.6% (-0.018-1.5%)	0.65% (-0.044-1.7%)	0.04 (-0.0027-0.1)	0.31 (-0.0094-0.79)
Kiribati	1990	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.44 (0.21-0.79)	0.36% (-0.089-1%)	0.37% (-0.13-1.1%)	0.0079 (-0.0029-0.023)	0.045 (-0.011-0.13)
Kiribati	1990	Male	95 plus	40% (22-61%)	60% (39-78%)	0.48 (0.2-0.91)	0.016% (-0.54-0.68%)	-0.08% (-0.71-0.65%)	-0.00048 (-0.0042-0.0038)	0.00037 (-0.012-0.016)
Kiribati	1995	Female	All Ages	7.4% (5.8-9.5%)	93% (91-94%)	0.14 (0.11-0.17)	0.4% (0.24-0.58%)	0.4% (0.14-0.7%)	1.5 (0.52-2.6)	83 (50-120)
Kiribati	1995	Female	15 to 19	9.2% (5.1-16%)	91% (84-95%)	0.18 (0.093-0.33)	0.92% (0.42-1.7%)	1.2% (0.39-2.5%)	0.064 (0.02-0.14)	6.3 (2.9-12)
Kiribati	1995	Female	20 to 24	12% (6.3-20%)	88% (80-94%)	0.3 (0.16-0.51)	1.6% (0.87-2.6%)	1.8% (0.73-3.5%)	0.11 (0.047-0.22)	11 (5.8-18)
Kiribati	1995	Female	25 to 29	12% (6.6-20%)	88% (80-93%)	0.3 (0.17-0.46)	1.5% (0.94-2.2%)	1.6% (0.86-2.7%)	0.16 (0.082-0.27)	14 (8.6-21)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	1995	Female	30 to 34	12% (6.2-20%)	88% (80-94%)	0.29 (0.15-0.48)	1.3% (0.82-2.1%)	1.5% (0.79-2.7%)	0.16 (0.082-0.28)	12 (7.3-19)
Kiribati	1995	Female	35 to 39	11% (5.9-19%)	89% (81-94%)	0.28 (0.15-0.49)	1.3% (0.75-2.1%)	1.6% (0.77-2.7%)	0.19 (0.095-0.33)	12 (7-20)
Kiribati	1995	Female	40 to 44	10% (5.6-18%)	90% (82-94%)	0.24 (0.12-0.42)	1.2% (0.59-2.2%)	1.6% (0.67-2.8%)	0.19 (0.08-0.34)	9.9 (4.7-17)
Kiribati	1995	Female	45 to 49	10% (5.1-18%)	90% (82-95%)	0.21 (0.097-0.38)	0.68% (0.17-1.3%)	0.85% (0.21-1.7%)	0.12 (0.027-0.24)	5.5 (1.4-10)
Kiribati	1995	Female	50 to 54	10% (5.3-18%)	90% (82-95%)	0.19 (0.086-0.36)	0.49% (-0.11-1.3%)	0.64% (-0.092-1.6%)	0.11 (-0.016-0.29)	4 (-0.85-11)
Kiribati	1995	Female	55 to 59	9.1% (4.9-16%)	91% (84-95%)	0.16 (0.072-0.3)	0.35% (-0.14-0.97%)	0.48% (-0.11-1.3%)	0.098 (-0.022-0.27)	2.9 (-1.2-8.3)
Kiribati	1995	Female	60 to 64	8.3% (4.3-15%)	92% (85-96%)	0.13 (0.061-0.26)	0.21% (-0.21-0.74%)	0.3% (-0.18-0.92%)	0.073 (-0.045-0.22)	1.7 (-1.7-5.9)
Kiribati	1995	Female	65 to 69	8.9% (4.9-15%)	91% (85-95%)	0.14 (0.065-0.25)	0.17% (-0.27-0.68%)	0.25% (-0.27-0.83%)	0.07 (-0.073-0.24)	1.3 (-2.5)
Kiribati	1995	Female	70 to 74	9.3% (5.3-16%)	91% (84-95%)	0.14 (0.062-0.25)	0.12% (-0.34-0.7%)	0.18% (-0.35-0.87%)	0.058 (-0.12-0.28)	0.84 (-2.3-4.9)
Kiribati	1995	Female	75 to 79	9.1% (4.9-16%)	91% (84-95%)	0.13 (0.057-0.25)	0.18% (-0.23-0.65%)	0.23% (-0.23-0.76%)	0.069 (-0.071-0.23)	0.87 (-1.1-3.1)
Kiribati	1995	Female	80 to 84	8.8% (4.7-15%)	91% (85-95%)	0.12 (0.058-0.21)	0.049% (-0.22-0.33%)	0.066% (-0.23-0.38%)	0.014 (-0.049-0.083)	0.13 (-0.56-0.84)
Kiribati	1995	Female	85 to 89	8.9% (4.9-15%)	91% (85-95%)	0.12 (0.056-0.23)	-0.021% (-0.26-0.21%)	-0.02% (-0.3-0.23%)	-0.0024 (-0.034-0.027)	-0.02 (-0.26-0.2)
Kiribati	1995	Female	90 to 94	9.7% (5.3-16%)	90% (84-95%)	0.13 (0.059-0.25)	0.014% (-0.19-0.23%)	0.016% (-0.21-0.26%)	0.0006 (-0.008-0.01)	0.0031 (-0.043-0.052)
Kiribati	1995	Female	95 plus	11% (5.5-20%)	89% (80-95%)	0.15 (0.059-0.3)	0.1% (-0.22-0.43%)	0.079% (-0.27-0.44%)	0.00087 (-0.003-0.0049)	0.0043 (-0.0089-0.018)
Kiribati	1995	Male	All Ages	38% (30-48%)	62% (52-70%)	0.74 (0.67-0.81)	2.4% (1.7-3.2%)	2.8% (1.8-3.9%)	13 (8-18)	610 (430-800)
Kiribati	1995	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.5 (0.24-0.84)	2% (0.76-3.9%)	2.3% (0.7-4.8%)	0.23 (0.073-0.49)	19 (7.7-38)
Kiribati	1995	Male	20 to 24	55% (36-72%)	45% (28-64%)	1.3 (0.81-2)	6.6% (3.5-10%)	7.7% (3.8-12%)	0.94 (0.46-1.5)	67 (35-110)
Kiribati	1995	Male	25 to 29	62% (43-78%)	38% (22-57%)	1.9 (1.2-2.6)	8.5% (5.4-12%)	9.9% (6.1-14%)	1.6 (1-2.4)	110 (68-150)
Kiribati	1995	Male	30 to 34	63% (42-79%)	37% (21-58%)	2 (1.3-2.8)	8.2% (5.1-12%)	9.5% (5.7-14%)	1.7 (1-2.5)	100 (63-150)
Kiribati	1995	Male	35 to 39	62% (44-78%)	38% (22-56%)	1.8 (1.1-2.6)	6% (3.2-9.1%)	6.8% (3.5-10%)	1.3 (0.72-2.1)	74 (41-110)
Kiribati	1995	Male	40 to 44	59% (42-76%)	41% (24-58%)	1.5 (0.86-2.4)	5.6% (2.6-8.9%)	6.3% (2.8-10%)	1.3 (0.59-2.2)	65 (30-100)
Kiribati	1995	Male	45 to 49	55% (34-73%)	45% (27-66%)	1.3 (0.71-2.1)	4.7% (2-8.1%)	5.3% (2.2-9.1%)	1.4 (0.56-2.4)	59 (25-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	1995	Male	50 to 54	52% (33-71%)	48% (29-67%)	1.1 (0.55-1.9)	3.6% (1.1-6.9%)	4% (1.1-7.6%)	1.1 (0.31-2.1)	41 (12-79)
Kiribati	1995	Male	55 to 59	50% (32-70%)	50% (30-68%)	0.88 (0.44-1.5)	2.4% (0.55-5.1%)	2.7% (0.58-5.6%)	0.77 (0.16-1.6)	25 (5.9-53)
Kiribati	1995	Male	60 to 64	48% (29-68%)	52% (32-71%)	0.74 (0.37-1.3)	1.8% (0.16-4.3%)	2% (0.17-4.7%)	0.65 (0.054-1.5)	18 (1.5-42)
Kiribati	1995	Male	65 to 69	47% (28-65%)	53% (35-72%)	0.68 (0.36-1.1)	2.1% (0.4-4.5%)	2.3% (0.45-5%)	0.78 (0.15-1.7)	18 (3.2-37)
Kiribati	1995	Male	70 to 74	44% (27-63%)	56% (37-73%)	0.6 (0.3-1.1)	1.5% (0.093-3.5%)	1.6% (0.12-3.9%)	0.52 (0.038-1.3)	9.5 (0.62-23)
Kiribati	1995	Male	75 to 79	40% (24-59%)	60% (41-76%)	0.51 (0.24-0.91)	1.3% (0.13-3%)	1.4% (0.14-3.4%)	0.33 (0.034-0.79)	4.7 (0.51-11)
Kiribati	1995	Male	80 to 84	37% (21-55%)	63% (45-79%)	0.45 (0.23-0.76)	0.8% (0.077-1.8%)	0.88% (0.075-2%)	0.13 (0.011-0.29)	1.4 (0.13-3)
Kiribati	1995	Male	85 to 89	35% (20-52%)	65% (48-80%)	0.43 (0.2-0.76)	0.55% (-0.047-1.4%)	0.6% (-0.071-1.6%)	0.04 (-0.0048-0.1)	0.31 (-0.026-0.79)
Kiribati	1995	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.43 (0.2-0.79)	0.33% (-0.13-0.95%)	0.33% (-0.19-1%)	0.0068 (-0.0039-0.021)	0.039 (-0.015-0.12)
Kiribati	1995	Male	95 plus	40% (22-60%)	60% (40-78%)	0.47 (0.19-0.89)	-0.016% (-0.59-0.64%)	-0.12% (-0.77-0.59%)	-0.00066 (-0.0043-0.0033)	-0.00034 (-0.013-0.014)
Kiribati	2000	Female	All Ages	7.3% (5.8-9.3%)	93% (91-94%)	0.13 (0.11-0.17)	0.38% (0.24-0.55%)	0.35% (0.11-0.63%)	1.3 (0.41-2.4)	78 (47-110)
Kiribati	2000	Female	15 to 19	9.5% (5.4-16%)	91% (84-95%)	0.19 (0.097-0.33)	0.91% (0.44-1.7%)	1.2% (0.42-2.5%)	0.071 (0.024-0.15)	7.1 (3.3-13)
Kiribati	2000	Female	20 to 24	12% (6.5-20%)	88% (80-93%)	0.3 (0.16-0.51)	1.6% (0.86-2.5%)	1.8% (0.73-3.4%)	0.12 (0.047-0.22)	11 (6.3-19)
Kiribati	2000	Female	25 to 29	12% (6.7-19%)	88% (81-93%)	0.29 (0.17-0.45)	1.4% (0.94-2.1%)	1.6% (0.87-2.6%)	0.11 (0.063-0.19)	10 (6.7-16)
Kiribati	2000	Female	30 to 34	11% (6-18%)	89% (82-94%)	0.26 (0.15-0.44)	1.2% (0.79-1.8%)	1.3% (0.72-2.2%)	0.15 (0.077-0.25)	12 (7.5-18)
Kiribati	2000	Female	35 to 39	9.1% (4.8-17%)	91% (83-95%)	0.22 (0.11-0.4)	1.1% (0.68-1.8%)	1.3% (0.66-2.3%)	0.17 (0.084-0.3)	11 (6.7-18)
Kiribati	2000	Female	40 to 44	8.6% (4.7-15%)	91% (85-95%)	0.2 (0.097-0.35)	1.1% (0.58-1.7%)	1.3% (0.64-2.3%)	0.19 (0.092-0.34)	10 (5.6-18)
Kiribati	2000	Female	45 to 49	10% (5.5-17%)	90% (83-95%)	0.2 (0.1-0.36)	0.61% (0.13-1.2%)	0.76% (0.16-1.6%)	0.12 (0.024-0.24)	5.4 (1.2-10)
Kiribati	2000	Female	50 to 54	10% (6.1-17%)	90% (83-94%)	0.19 (0.095-0.34)	0.39% (-0.18-1.2%)	0.53% (-0.16-1.5%)	0.1 (-0.031-0.28)	3.7 (-1.6-10)
Kiribati	2000	Female	55 to 59	8.9% (5.1-16%)	91% (84-95%)	0.15 (0.071-0.28)	0.28% (-0.18-0.83%)	0.39% (-0.14-1.1%)	0.081 (-0.029-0.23)	2.4 (-1.6-7.1)
Kiribati	2000	Female	60 to 64	8.1% (4.2-14%)	92% (86-96%)	0.13 (0.059-0.24)	0.16% (-0.24-0.64%)	0.24% (-0.22-0.81%)	0.062 (-0.058-0.21)	1.4 (-2-5.4)
Kiribati	2000	Female	65 to 69	8.9% (4.8-15%)	91% (85-95%)	0.13 (0.064-0.24)	0.12% (-0.3-0.58%)	0.19% (-0.29-0.74%)	0.055 (-0.084-0.21)	0.92 (-2.4-4.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	2000	Female	70 to 74	9.4% (5.4-16%)	91% (84-95%)	0.14 (0.062-0.25)	0.046% (-0.39-0.54%)	0.094% (-0.41-0.69%)	0.032 (-0.14-0.24)	0.34 (-2.9-4.1)
Kiribati	2000	Female	75 to 79	9.3% (5.3-16%)	91% (84-95%)	0.13 (0.058-0.24)	0.12% (-0.27-0.54%)	0.17% (-0.27-0.65%)	0.052 (-0.084-0.21)	0.61 (-1.4-2.8)
Kiribati	2000	Female	80 to 84	9.2% (4.9-16%)	91% (84-95%)	0.13 (0.061-0.22)	0.019% (-0.26-0.3%)	0.034% (-0.28-0.35%)	0.0077 (-0.062-0.08)	0.052 (-0.71-0.82)
Kiribati	2000	Female	85 to 89	9.4% (5-15%)	91% (85-95%)	0.13 (0.06-0.24)	-0.046% (-0.32-0.21%)	-0.047% (-0.35-0.24%)	-0.0059 (-0.045-0.03)	-0.049 (-0.35-0.22)
Kiribati	2000	Female	90 to 94	10% (5.7-17%)	90% (83-94%)	0.14 (0.063-0.26)	-0.0019% (-0.22-0.22%)	-0.00095% (-0.24-0.25%)	-0.000044 (-0.01-0.011)	-0.00052 (-0.056-0.057)
Kiribati	2000	Female	95 plus	12% (5.7-21%)	88% (79-94%)	0.16 (0.063-0.31)	0.084% (-0.26-0.45%)	0.062% (-0.33-0.46%)	0.00065 (-0.0035-0.0049)	0.0035 (-0.011-0.018)
Kiribati	2000	Male	All Ages	40% (32-48%)	60% (52-68%)	0.76 (0.69-0.83)	2.9% (1.7-3.4%)	2.9% (1.7-4%)	13 (7.9-19)	640 (440-850)
Kiribati	2000	Male	15 to 19	38% (22-56%)	62% (44-78%)	0.51 (0.25-0.84)	2.1% (0.79-4%)	2.5% (0.75-5.1%)	0.28 (0.085-0.57)	23 (9-44)
Kiribati	2000	Male	20 to 24	56% (38-74%)	44% (26-62%)	1.4 (0.86-2)	6.9% (3.6-11%)	8.1% (4-13%)	1.1 (0.54-1.8)	79 (41-120)
Kiribati	2000	Male	25 to 29	65% (50-79%)	35% (21-50%)	2 (1.4-2.7)	8.9% (5.5-13%)	10% (6.3-15%)	1.3 (0.81-2)	88 (54-130)
Kiribati	2000	Male	30 to 34	66% (49-80%)	34% (20-51%)	2.1 (1.4-2.8)	8.1% (4.9-12%)	9.4% (5.4-14%)	1.9 (1.1-2.7)	110 (67-160)
Kiribati	2000	Male	35 to 39	62% (47-77%)	38% (23-53%)	1.7 (1.1-2.5)	5.7% (3.2-8.6%)	6.4% (3.5-9.9%)	1.5 (0.8-2.4)	82 (46-130)
Kiribati	2000	Male	40 to 44	59% (43-74%)	41% (26-57%)	1.5 (0.89-2.2)	5.1% (2.2-8.3%)	5.7% (2.4-9.3%)	1.5 (0.63-2.6)	74 (33-120)
Kiribati	2000	Male	45 to 49	57% (38-72%)	43% (28-62%)	1.4 (0.81-2.1)	4.4% (1.7-7.7%)	4.9% (1.9-8.7%)	1.4 (0.54-2.5)	63 (24-110)
Kiribati	2000	Male	50 to 54	55% (36-70%)	45% (30-64%)	1.1 (0.63-1.9)	3.4% (0.86-6.6%)	3.7% (0.84-7.4%)	1.3 (0.31-2.5)	48 (13-95)
Kiribati	2000	Male	55 to 59	51% (34-67%)	49% (33-66%)	0.88 (0.47-1.4)	2.2% (0.3-4.7%)	2.4% (0.27-5.2%)	0.76 (0.088-1.7)	25 (3.4-57)
Kiribati	2000	Male	60 to 64	48% (30-66%)	52% (34-70%)	0.73 (0.38-1.3)	1.6% (0.0088-4%)	1.8% (-0.0011-4.4%)	0.61 (-0.00037-1.5)	17 (0.091-42)
Kiribati	2000	Male	65 to 69	47% (29-65%)	53% (35-71%)	0.69 (0.37-1.2)	1.8% (0.24-4.2%)	2% (0.27-4.6%)	0.66 (0.088-1.5)	15 (1.9-34)
Kiribati	2000	Male	70 to 74	44% (27-62%)	56% (38-73%)	0.6 (0.31-1.1)	1.3% (0.0028-3.3%)	1.4% (0.016-3.6%)	0.44 (0.0049-1.1)	8.1 (0.017-21)
Kiribati	2000	Male	75 to 79	41% (24-59%)	59% (41-76%)	0.52 (0.25-0.91)	1.1% (0.077-2.9%)	1.3% (0.083-3.2%)	0.29 (0.019-0.73)	4.1 (0.28-10)
Kiribati	2000	Male	80 to 84	38% (21-57%)	62% (43-79%)	0.46 (0.24-0.78)	0.75% (0.022-1.8%)	0.83% (0.02-2%)	0.12 (0.0029-0.29)	1.3 (0.037-3.1)
Kiribati	2000	Male	85 to 89	37% (22-54%)	63% (46-78%)	0.44 (0.21-0.77)	0.48% (-0.13-1.4%)	0.52% (-0.16-1.6%)	0.036 (-0.011-0.11)	0.28 (-0.074-0.84)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	2000	Male	90 to 94	38% (23-55%)	62% (45-77%)	0.45 (0.21-0.81)	0.28% (-0.2-0.98%)	0.27% (-0.28-1.1%)	0.0058 (-0.0059-0.023)	0.035 (-0.026-0.12)
Kiribati	2000	Male	95 plus	41% (23-62%)	59% (38-77%)	0.49 (0.2-0.91)	-0.059% (-0.69-0.64%)	-0.17% (-0.89-0.6%)	-0.00092 (-0.0047-0.0032)	-0.0012 (-0.014-0.013)
Kiribati	2005	Female	All Ages	7.5% (6-9.7%)	92% (90-94%)	0.14 (0.12-0.18)	0.38% (0.23-0.55%)	0.34% (0.086-0.6%)	1.3 (0.35-2.4)	82 (49-120)
Kiribati	2005	Female	15 to 19	9.5% (5.4-16%)	90% (84-95%)	0.19 (0.099-0.33)	0.89% (0.45-1.6%)	1.2% (0.42-2.4%)	0.079 (0.028-0.17)	8 (3.9-15)
Kiribati	2005	Female	20 to 24	12% (6.7-20%)	88% (80-93%)	0.31 (0.16-0.52)	1.5% (0.87-2.5%)	1.8% (0.75-3.3%)	0.14 (0.057-0.26)	14 (7.6-23)
Kiribati	2005	Female	25 to 29	12% (6.9-20%)	88% (80-93%)	0.29 (0.17-0.46)	1.4% (0.91-2%)	1.5% (0.83-2.5%)	0.12 (0.066-0.21)	12 (7.3-17)
Kiribati	2005	Female	30 to 34	11% (6.1-18%)	89% (82-94%)	0.26 (0.15-0.45)	1.2% (0.77-1.8%)	1.3% (0.71-2.2%)	0.12 (0.061-0.2)	9.7 (6-15)
Kiribati	2005	Female	35 to 39	9.2% (5-17%)	91% (83-95%)	0.23 (0.12-0.41)	1.1% (0.69-1.7%)	1.3% (0.69-2.3%)	0.18 (0.092-0.32)	12 (7.6-20)
Kiribati	2005	Female	40 to 44	8.7% (4.8-15%)	91% (85-95%)	0.2 (0.098-0.36)	1% (0.58-1.7%)	1.3% (0.62-2.2%)	0.19 (0.097-0.34)	11 (5.9-18)
Kiribati	2005	Female	45 to 49	10% (5.6-16%)	90% (84-94%)	0.21 (0.1-0.35)	0.57% (0.12-1.1%)	0.73% (0.12-1.5%)	0.13 (0.02-0.29)	6.3 (1.2-13)
Kiribati	2005	Female	50 to 54	11% (6.3-17%)	89% (83-94%)	0.19 (0.096-0.33)	0.33% (-0.23-0.99%)	0.47% (-0.2-1.3%)	0.098 (-0.042-0.27)	3.4 (-2.4-10)
Kiribati	2005	Female	55 to 59	9.1% (4.9-16%)	91% (84-95%)	0.15 (0.073-0.29)	0.24% (-0.24-0.76%)	0.36% (-0.2-1%)	0.086 (-0.046-0.24)	2.4 (-2.3-7.4)
Kiribati	2005	Female	60 to 64	8.2% (4.2-15%)	92% (85-96%)	0.13 (0.061-0.25)	0.14% (-0.28-0.59%)	0.22% (-0.27-0.76%)	0.055 (-0.07-0.19)	1.2 (-2.4-5)
Kiribati	2005	Female	65 to 69	9% (4.8-16%)	91% (84-95%)	0.14 (0.066-0.25)	0.096% (-0.35-0.56%)	0.17% (-0.35-0.69%)	0.051 (-0.1-0.21)	0.81 (-3-4.6)
Kiribati	2005	Female	70 to 74	9.6% (5.4-16%)	90% (84-95%)	0.14 (0.065-0.27)	0.021% (-0.44-0.48%)	0.067% (-0.46-0.63%)	0.029 (-0.2-0.27)	0.2 (-4.1-4.6)
Kiribati	2005	Female	75 to 79	9.4% (5.2-16%)	91% (84-95%)	0.13 (0.06-0.25)	0.1% (-0.3-0.53%)	0.15% (-0.29-0.65%)	0.049 (-0.094-0.22)	0.55 (-1.6-2.9)
Kiribati	2005	Female	80 to 84	9.2% (4.8-16%)	91% (84-95%)	0.13 (0.062-0.23)	0.0058% (-0.3-0.29%)	0.019% (-0.32-0.35%)	0.0045 (-0.077-0.082)	0.016 (-0.88-0.82)
Kiribati	2005	Female	85 to 89	9.4% (5.1-15%)	91% (85-95%)	0.13 (0.061-0.24)	-0.059% (-0.36-0.21%)	-0.063% (-0.39-0.23%)	-0.0088 (-0.056-0.031)	-0.07 (-0.43-0.24)
Kiribati	2005	Female	90 to 94	10% (5.7-17%)	90% (83-94%)	0.14 (0.066-0.26)	-0.012% (-0.24-0.23%)	-0.014% (-0.27-0.24%)	-0.00072 (-0.013-0.012)	-0.0037 (-0.071-0.067)
Kiribati	2005	Female	95 plus	12% (5.8-21%)	88% (79-94%)	0.16 (0.066-0.32)	0.072% (-0.29-0.43%)	0.043% (-0.38-0.46%)	0.00054 (-0.0048-0.0058)	0.0035 (-0.014-0.021)
Kiribati	2005	Male	All Ages	41% (33-50%)	59% (50-67%)	0.79 (0.72-0.86)	2.6% (1.7-3.5%)	2.8% (1.7-4%)	14 (8.3-20)	690 (470-940)
Kiribati	2005	Male	15 to 19	38% (22-56%)	62% (44-78%)	0.51 (0.26-0.84)	2.1% (0.83-4%)	2.5% (0.83-5.1%)	0.33 (0.11-0.7)	28 (11-53)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	2005	Male	20 to 24	58% (39-74%)	42% (26-61%)	1.4 (0.89-2)	6.8% (3.7-10%)	8.1% (4-13%)	1.3 (0.64-2.1)	96 (50-150)
Kiribati	2005	Male	25 to 29	66% (50-80%)	34% (20-50%)	2 (1.4-2.7)	8.7% (5.5-12%)	10% (6.2-15%)	1.6 (0.97-2.3)	110 (65-150)
Kiribati	2005	Male	30 to 34	66% (50-80%)	34% (20-50%)	2.1 (1.4-3)	7.9% (4.9-12%)	9.2% (5.5-14%)	1.5 (0.86-2.2)	90 (54-130)
Kiribati	2005	Male	35 to 39	62% (46-77%)	38% (23-54%)	1.7 (1.1-2.5)	5.7% (3.1-8.6%)	6.4% (3.3-9.9%)	1.6 (0.85-2.6)	91 (51-140)
Kiribati	2005	Male	40 to 44	59% (44-75%)	41% (25-56%)	1.5 (0.9-2.2)	5% (2.2-8.3%)	5.6% (2.3-9.3%)	1.7 (0.68-2.9)	84 (35-140)
Kiribati	2005	Male	45 to 49	57% (39-73%)	43% (27-61%)	1.4 (0.8-2.1)	4.2% (1.5-7.6%)	4.7% (1.6-8.5%)	1.6 (0.56-2.9)	71 (25-120)
Kiribati	2005	Male	50 to 54	55% (37-71%)	45% (29-63%)	1.1 (0.63-1.9)	3.2% (0.65-6.3%)	3.5% (0.76-7%)	1.2 (0.26-2.5)	47 (11-96)
Kiribati	2005	Male	55 to 59	51% (34-68%)	49% (32-66%)	0.89 (0.49-1.5)	2% (0.21-4.6%)	2.2% (0.2-5.1%)	0.82 (0.074-1.9)	28 (3-61)
Kiribati	2005	Male	60 to 64	48% (30-67%)	52% (33-70%)	0.73 (0.38-1.3)	1.5% (0.016-3.9%)	1.7% (0.014-4.3%)	0.61 (0.0056-1.6)	17 (0.18-43)
Kiribati	2005	Male	65 to 69	47% (28-66%)	53% (34-72%)	0.69 (0.37-1.1)	1.7% (0.079-3.9%)	1.9% (0.094-4.3%)	0.63 (0.032-1.4)	14 (0.68-33)
Kiribati	2005	Male	70 to 74	45% (27-62%)	55% (38-73%)	0.61 (0.3-1.1)	1.2% (-0.066-3%)	1.3% (-0.1-3.4%)	0.52 (-0.039-1.3)	9.5 (-0.54-25)
Kiribati	2005	Male	75 to 79	41% (25-60%)	59% (40-75%)	0.52 (0.25-0.94)	1.1% (0.024-2.7%)	1.2% (0.047-2.9%)	0.29 (0.012-0.72)	4.2 (0.091-10)
Kiribati	2005	Male	80 to 84	38% (21-56%)	62% (44-79%)	0.46 (0.24-0.77)	0.72% (0.019-1.8%)	0.78% (0.011-1.9%)	0.12 (0.0017-0.29)	1.3 (0.034-3.1)
Kiribati	2005	Male	85 to 89	36% (21-54%)	64% (46-79%)	0.44 (0.21-0.76)	0.44% (-0.2-1.4%)	0.47% (-0.24-1.5%)	0.034 (-0.018-0.11)	0.27 (-0.12-0.84)
Kiribati	2005	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.44 (0.21-0.8)	0.24% (-0.26-0.94%)	0.23% (-0.36-1%)	0.0054 (-0.0084-0.024)	0.034 (-0.036-0.13)
Kiribati	2005	Male	95 plus	41% (22-61%)	59% (39-78%)	0.48 (0.2-0.93)	-0.083% (-0.71-0.65%)	-0.2% (-0.91-0.6%)	-0.0012 (-0.0054-0.0034)	-0.0019 (-0.016-0.015)
Kiribati	2010	Female	All Ages	7.8% (6.1-9.9%)	92% (90-94%)	0.15 (0.12-0.19)	0.41% (0.24-0.61%)	0.36% (0.11-0.65%)	1.5 (0.45-2.7)	92 (54-130)
Kiribati	2010	Female	15 to 19	9.6% (5.5-16%)	90% (84-94%)	0.19 (0.098-0.33)	0.88% (0.46-1.6%)	1.2% (0.43-2.3%)	0.075 (0.028-0.16)	7.8 (4-14)
Kiribati	2010	Female	20 to 24	12% (6.8-20%)	88% (80-93%)	0.31 (0.17-0.51)	1.5% (0.91-2.4%)	1.7% (0.78-3.2%)	0.15 (0.064-0.28)	16 (9-25)
Kiribati	2010	Female	25 to 29	12% (6.9-20%)	88% (80-93%)	0.29 (0.17-0.45)	1.4% (0.92-2%)	1.5% (0.83-2.5%)	0.14 (0.074-0.24)	14 (8.5-21)
Kiribati	2010	Female	30 to 34	11% (6-18%)	89% (82-94%)	0.26 (0.15-0.44)	1.2% (0.77-1.8%)	1.3% (0.72-2.1%)	0.13 (0.069-0.23)	11 (7-17)
Kiribati	2010	Female	35 to 39	9.2% (4.8-17%)	91% (83-95%)	0.23 (0.11-0.43)	1.1% (0.7-1.8%)	1.3% (0.7-2.3%)	0.16 (0.08-0.29)	11 (6.6-18)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	2010	Female	40 to 44	8.7% (4.8-16%)	91% (84-95%)	0.21 (0.099-0.38)	1% (0.58-1.7%)	1.3% (0.66-2.2%)	0.23 (0.11-0.4)	13 (7-21)
Kiribati	2010	Female	45 to 49	10% (5.4-17%)	90% (83-95%)	0.21 (0.11-0.36)	0.6% (0.14-1.2%)	0.76% (0.14-1.5%)	0.17 (0.032-0.33)	7.8 (1.8-15)
Kiribati	2010	Female	50 to 54	11% (6.2-18%)	89% (82-94%)	0.2 (0.096-0.35)	0.35% (-0.2-0.94%)	0.48% (-0.21-1.3%)	0.12 (-0.057-0.33)	4.4 (-2.6-12)
Kiribati	2010	Female	55 to 59	9.2% (5-16%)	91% (84-95%)	0.16 (0.075-0.3)	0.27% (-0.25-0.77%)	0.38% (-0.21-1%)	0.11 (-0.06-0.29)	3.1 (-2.8-9.2)
Kiribati	2010	Female	60 to 64	8.3% (4.4-15%)	92% (85-96%)	0.13 (0.062-0.26)	0.15% (-0.23-0.62%)	0.23% (-0.21-0.79%)	0.065 (-0.061-0.22)	1.5 (-2.2-5.9)
Kiribati	2010	Female	65 to 69	9% (5-15%)	91% (85-95%)	0.14 (0.067-0.24)	0.11% (-0.33-0.56%)	0.19% (-0.32-0.73%)	0.066 (-0.11-0.25)	1.1 (-3.3-5.4)
Kiribati	2010	Female	70 to 74	9.5% (5.5-16%)	90% (84-94%)	0.14 (0.065-0.27)	0.022% (-0.44-0.49%)	0.069% (-0.45-0.6%)	0.029 (-0.19-0.27)	0.21 (-4-4.5)
Kiribati	2010	Female	75 to 79	9.3% (5.1-16%)	91% (84-95%)	0.13 (0.059-0.26)	0.11% (-0.29-0.52%)	0.16% (-0.29-0.63%)	0.06 (-0.11-0.24)	0.69 (-1.8-3.2)
Kiribati	2010	Female	80 to 84	9.2% (4.9-16%)	91% (84-95%)	0.13 (0.061-0.23)	0.011% (-0.29-0.3%)	0.024% (-0.31-0.36%)	0.0068 (-0.086-0.097)	0.037 (-0.99-1)
Kiribati	2010	Female	85 to 89	9.5% (5.1-15%)	91% (85-95%)	0.13 (0.062-0.23)	-0.053% (-0.37-0.18%)	-0.057% (-0.41-0.21%)	-0.0086 (-0.063-0.03)	-0.069 (-0.48-0.22)
Kiribati	2010	Female	90 to 94	10% (5.7-17%)	90% (83-94%)	0.14 (0.065-0.27)	-0.0089% (-0.24-0.23%)	-0.012% (-0.27-0.25%)	-0.00066 (-0.015-0.014)	-0.003 (-0.08-0.078)
Kiribati	2010	Female	95 plus	12% (5.8-20%)	88% (80-94%)	0.16 (0.068-0.31)	0.081% (-0.29-0.44%)	0.049% (-0.37-0.45%)	0.00071 (-0.0053-0.0065)	0.0045 (-0.016-0.024)
Kiribati	2010	Male	All Ages	42% (35-51%)	58% (49-65%)	0.84 (0.75-0.94)	2.7% (1.8-3.7%)	2.9% (1.7-4.1%)	15 (8.8-23)	760 (500-1000)
Kiribati	2010	Male	15 to 19	38% (22-56%)	62% (44-78%)	0.51 (0.26-0.84)	2.1% (0.8-3.9%)	2.5% (0.71-5.1%)	0.33 (0.094-0.67)	28 (10-52)
Kiribati	2010	Male	20 to 24	57% (39-74%)	43% (26-61%)	1.4 (0.88-2)	6.7% (3.6-11%)	8% (3.9-13%)	1.6 (0.76-2.5)	110 (60-180)
Kiribati	2010	Male	25 to 29	66% (51-79%)	34% (21-49%)	2 (1.4-2.7)	8.6% (5.2-12%)	10% (5.9-15%)	1.9 (1.1-2.8)	130 (78-180)
Kiribati	2010	Male	30 to 34	66% (49-81%)	34% (19-51%)	2.1 (1.4-2.9)	7.7% (4.8-11%)	8.9% (5.3-13%)	1.7 (1-2.6)	110 (65-160)
Kiribati	2010	Male	35 to 39	62% (46-78%)	38% (22-54%)	1.7 (1.1-2.5)	5.6% (3-8.6%)	6.3% (3.3-9.9%)	1.4 (0.71-2.3)	78 (41-120)
Kiribati	2010	Male	40 to 44	60% (43-75%)	40% (25-57%)	1.5 (0.9-2.3)	4.9% (2-8.1%)	5.5% (2.1-9.1%)	1.9 (0.68-3.2)	92 (36-160)
Kiribati	2010	Male	45 to 49	58% (39-73%)	42% (27-61%)	1.4 (0.83-2.2)	4.1% (1.4-7.5%)	4.6% (1.5-8.3%)	1.9 (0.62-3.6)	84 (29-150)
Kiribati	2010	Male	50 to 54	55% (37-71%)	45% (29-63%)	1.1 (0.64-1.9)	3.1% (0.55-6%)	3.4% (0.51-6.8%)	1.4 (0.22-2.9)	54 (11-110)
Kiribati	2010	Male	55 to 59	51% (34-68%)	49% (32-66%)	0.89 (0.5-1.4)	2% (0.24-4.3%)	2.1% (0.2-4.8%)	0.93 (0.085-2)	31 (3.6-68)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	2010	Male	60 to 64	48% (30-66%)	52% (34-70%)	0.74 (0.4-1.3)	1.5% (0.0019-3.8%)	1.6% (-0.013-4.1%)	0.65 (-0.0055-1.7)	18 (0.022-48)
Kiribati	2010	Male	65 to 69	47% (28-66%)	53% (34-72%)	0.69 (0.37-1.2)	1.6% (0.13-3.7%)	1.8% (0.17-4.1%)	0.7 (0.067-1.7)	16 (1.3-37)
Kiribati	2010	Male	70 to 74	45% (28-63%)	55% (37-72%)	0.61 (0.3-1.1)	1.2% (-0.058-3%)	1.3% (-0.066-3.4%)	0.48 (-0.026-1.3)	8.8 (-0.45-23)
Kiribati	2010	Male	75 to 79	41% (25-60%)	59% (40-75%)	0.52 (0.26-0.94)	1.1% (0.016-2.6%)	1.2% (0.024-2.9%)	0.32 (0.0065-0.79)	4.5 (0.067-11)
Kiribati	2010	Male	80 to 84	38% (22-57%)	62% (43-78%)	0.46 (0.24-0.77)	0.71% (-0.01-1.7%)	0.78% (-0.024-1.8%)	0.14 (-0.0041-0.32)	1.4 (-0.021-3.4)
Kiribati	2010	Male	85 to 89	37% (22-53%)	63% (47-78%)	0.44 (0.21-0.76)	0.43% (-0.21-1.3%)	0.45% (-0.26-1.4%)	0.033 (-0.019-0.1)	0.26 (-0.13-0.8)
Kiribati	2010	Male	90 to 94	37% (23-55%)	63% (45-77%)	0.44 (0.21-0.79)	0.23% (-0.26-0.89%)	0.21% (-0.35-0.94%)	0.005 (-0.0085-0.023)	0.032 (-0.038-0.13)
Kiribati	2010	Male	95 plus	41% (22-61%)	59% (39-78%)	0.47 (0.2-0.91)	-0.091% (-0.71-0.62%)	-0.21% (-0.91-0.54%)	-0.0013 (-0.0057-0.0033)	-0.0022 (-0.017-0.015)
Kiribati	2016	Female	All Ages	8.4% (6.6-11%)	92% (89-93%)	0.17 (0.12-0.23)	0.47% (0.23-0.77%)	0.39% (0.076-0.82%)	1.8 (0.35-3.9)	110 (52-170)
Kiribati	2016	Female	15 to 19	9.5% (5.4-16%)	91% (84-95%)	0.19 (0.093-0.36)	0.88% (0.44-1.6%)	1.2% (0.4-2.5%)	0.074 (0.025-0.16)	8 (3.9-15)
Kiribati	2016	Female	20 to 24	12% (6.7-20%)	88% (80-93%)	0.32 (0.15-0.58)	1.5% (0.86-2.5%)	1.8% (0.72-3.6%)	0.15 (0.055-0.3)	16 (8.3-26)
Kiribati	2016	Female	25 to 29	13% (6.7-21%)	87% (79-93%)	0.32 (0.15-0.53)	1.5% (0.89-2.2%)	1.6% (0.81-2.8%)	0.16 (0.075-0.3)	16 (9.3-25)
Kiribati	2016	Female	30 to 34	12% (6.3-21%)	88% (79-94%)	0.31 (0.15-0.54)	1.3% (0.78-2%)	1.5% (0.71-2.5%)	0.17 (0.079-0.32)	15 (8.4-23)
Kiribati	2016	Female	35 to 39	12% (6.1-21%)	88% (79-94%)	0.29 (0.14-0.53)	1.3% (0.74-2.1%)	1.5% (0.79-2.8%)	0.21 (0.1-0.39)	14 (7.9-24)
Kiribati	2016	Female	40 to 44	11% (6-19%)	89% (81-94%)	0.26 (0.12-0.49)	1.1% (0.56-2.1%)	1.5% (0.64-2.8%)	0.23 (0.094-0.45)	13 (5.9-23)
Kiribati	2016	Female	45 to 49	11% (5.3-19%)	89% (81-95%)	0.22 (0.099-0.43)	0.63% (0.086-1.4%)	0.8% (0.14-1.8%)	0.19 (0.033-0.42)	9.1 (1.3-20)
Kiribati	2016	Female	50 to 54	11% (5.7-19%)	89% (81-94%)	0.2 (0.09-0.38)	0.38% (-0.17-1.1%)	0.53% (-0.18-1.5%)	0.17 (-0.06-0.48)	6.2 (-2.9-18)
Kiribati	2016	Female	55 to 59	9.6% (5.2-17%)	90% (83-95%)	0.16 (0.072-0.33)	0.28% (-0.25-0.94%)	0.41% (-0.21-1.2%)	0.14 (-0.076-0.44)	4.2 (-3.7-14)
Kiribati	2016	Female	60 to 64	8.8% (4.6-16%)	91% (84-95%)	0.14 (0.061-0.28)	0.16% (-0.27-0.67%)	0.25% (-0.24-0.87%)	0.099 (-0.096-0.34)	2.2 (-3.8-9)
Kiribati	2016	Female	65 to 69	9.4% (5-16%)	91% (84-95%)	0.15 (0.067-0.28)	0.13% (-0.34-0.67%)	0.21% (-0.33-0.83%)	0.079 (-0.13-0.31)	1.3 (-3.4-6.9)
Kiribati	2016	Female	70 to 74	9.7% (5.6-16%)	90% (84-94%)	0.15 (0.063-0.29)	0.039% (-0.44-0.61%)	0.088% (-0.45-0.76%)	0.038 (-0.19-0.33)	0.38 (-4.1-5.7)
Kiribati	2016	Female	75 to 79	9.5% (5.3-17%)	90% (83-95%)	0.14 (0.059-0.27)	0.13% (-0.29-0.6%)	0.18% (-0.29-0.72%)	0.086 (-0.14-0.35)	1 (-2.3-4.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kiribati	2016	Female	80 to 84	9.3% (4.9-16%)	91% (84-95%)	0.13 (0.061-0.25)	0.019% (-0.29-0.35%)	0.034% (-0.31-0.41%)	0.012 (-0.11-0.14)	0.081 (-1.3-1.5)
Kiribati	2016	Female	85 to 89	9.5% (5.1-16%)	91% (84-95%)	0.14 (0.061-0.26)	-0.038% (-0.34-0.24%)	-0.041% (-0.38-0.27%)	-0.0082 (-0.076-0.053)	-0.066 (-0.58-0.42)
Kiribati	2016	Female	90 to 94	10% (5.6-16%)	90% (84-94%)	0.15 (0.064-0.28)	0.0032% (-0.24-0.26%)	0.00096% (-0.27-0.29%)	0.000058 (-0.019-0.02)	0.0013 (-0.099-0.11)
Kiribati	2016	Female	95 plus	12% (5.9-21%)	88% (79-94%)	0.17 (0.065-0.34)	0.087% (-0.31-0.48%)	0.061% (-0.39-0.51%)	0.0011 (-0.0072-0.0092)	0.0061 (-0.022-0.033)
Kiribati	2016	Male	All Ages	43% (35-52%)	57% (48-65%)	0.88 (0.65-1.1)	3% (1.6-4.6%)	3.1% (1.5-5.1%)	18 (8.1-30)	870 (470-1400)
Kiribati	2016	Male	15 to 19	38% (23-56%)	62% (44-77%)	0.53 (0.25-0.95)	2.1% (0.79-4.3%)	2.6% (0.71-5.8%)	0.34 (0.09-0.74)	29 (11-58)
Kiribati	2016	Male	20 to 24	56% (37-73%)	44% (27-63%)	1.4 (0.81-2.3)	6.8% (3.3-11%)	8% (3.5-14%)	1.6 (0.67-2.8)	110 (54-190)
Kiribati	2016	Male	25 to 29	63% (43-78%)	37% (22-57%)	1.9 (1.2-2.9)	8.3% (4.7-13%)	9.8% (5.2-15%)	2.2 (1.1-3.5)	150 (81-230)
Kiribati	2016	Male	30 to 34	64% (43-80%)	36% (20-57%)	2.1 (1.2-3.2)	7.7% (4.3-12%)	8.9% (4.5-14%)	2.1 (1-3.4)	130 (66-200)
Kiribati	2016	Male	35 to 39	63% (46-80%)	37% (20-54%)	1.8 (1.1-2.9)	5.9% (2.9-9.7%)	6.6% (3-11%)	1.7 (0.71-2.9)	93 (43-160)
Kiribati	2016	Male	40 to 44	61% (43-78%)	39% (22-57%)	1.6 (0.86-2.7)	5.1% (2-9.2%)	5.7% (2.1-10%)	1.7 (0.53-3.1)	83 (29-150)
Kiribati	2016	Male	45 to 49	57% (34-75%)	43% (25-66%)	1.4 (0.7-2.4)	4.2% (1.3-7.9%)	4.6% (1.3-9%)	2 (0.59-4.1)	90 (28-180)
Kiribati	2016	Male	50 to 54	54% (34-73%)	46% (27-66%)	1.2 (0.58-2.1)	3.2% (0.56-6.7%)	3.5% (0.54-7.5%)	2 (0.29-4.3)	76 (13-170)
Kiribati	2016	Male	55 to 59	52% (33-71%)	48% (29-67%)	0.95 (0.46-1.7)	2.1% (0.15-5.3%)	2.4% (0.083-5.9%)	1.3 (0.046-3.2)	43 (3-110)
Kiribati	2016	Male	60 to 64	49% (30-69%)	51% (31-70%)	0.79 (0.36-1.5)	1.6% (-0.049-4.5%)	1.8% (-0.053-5%)	0.95 (-0.028-2.8)	27 (-0.79-77)
Kiribati	2016	Male	65 to 69	48% (30-67%)	52% (33-70%)	0.71 (0.35-1.2)	1.7% (0.093-4.3%)	1.9% (0.13-4.7%)	0.9 (0.062-2.3)	21 (1.1-54)
Kiribati	2016	Male	70 to 74	45% (28-62%)	55% (38-72%)	0.62 (0.3-1.1)	1.2% (-0.094-3.2%)	1.4% (-0.091-3.6%)	0.57 (-0.037-1.6)	11 (-0.81-29)
Kiribati	2016	Male	75 to 79	40% (24-59%)	60% (41-76%)	0.52 (0.24-0.97)	1.1% (-0.0042-2.7%)	1.2% (-0.00046-3.1%)	0.43 (-0.00017-1.1)	6.2 (-0.022-16)
Kiribati	2016	Male	80 to 84	37% (21-56%)	63% (44-79%)	0.47 (0.23-0.87)	0.77% (-0.019-2%)	0.84% (-0.019-2.1%)	0.18 (-0.0036-0.44)	1.9 (-0.046-4.8)
Kiribati	2016	Male	85 to 89	36% (22-52%)	64% (48-78%)	0.45 (0.21-0.82)	0.49% (-0.19-1.6%)	0.52% (-0.23-1.7%)	0.047 (-0.021-0.16)	0.37 (-0.15-1.2)
Kiribati	2016	Male	90 to 94	38% (23-55%)	62% (45-77%)	0.46 (0.21-0.85)	0.27% (-0.26-1%)	0.25% (-0.35-1.1%)	0.0067 (-0.0092-0.028)	0.042 (-0.041-0.16)
Kiribati	2016	Male	95 plus	41% (22-62%)	59% (38-78%)	0.5 (0.2-0.97)	-0.036% (-0.7-0.8%)	-0.16% (-0.91-0.78%)	-0.001 (-0.0059-0.0051)	-0.00091 (-0.018-0.021)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	1990	Female	All Ages	18% (13-25%)	82% (75-87%)	0.28 (0.15-0.48)	0.061% (-0.39-0.7%)	-0.39% (-1.2-0.7%)	-0.46 (-1.4-0.79)	4.5 (-27-53)
Marshall Islands	1990	Female	15 to 19	20% (9-36%)	80% (64-91%)	0.35 (0.13-0.82)	1% (0.4-2.1%)	1.5% (0.33-3.7%)	0.034 (0.0076-0.088)	3.6 (1.4-7.5)
Marshall Islands	1990	Female	20 to 24	24% (10-43%)	76% (57-90%)	0.55 (0.19-1.2)	1.7% (0.81-3.2%)	2.2% (0.63-5.2%)	0.056 (0.017-0.13)	5.8 (2.8-11)
Marshall Islands	1990	Female	25 to 29	26% (10-46%)	74% (54-90%)	0.57 (0.21-1.2)	1.6% (0.75-2.9%)	2% (0.58-4.2%)	0.055 (0.016-0.12)	5.2 (2.5-9.7)
Marshall Islands	1990	Female	30 to 34	27% (11-47%)	73% (53-89%)	0.59 (0.22-1.2)	1.3% (0.45-2.7%)	1.6% (0.24-3.9%)	0.05 (0.008-0.12)	4.2 (1.5-9)
Marshall Islands	1990	Female	35 to 39	27% (11-46%)	73% (54-89%)	0.6 (0.22-1.2)	0.97% (-0.034-2.6%)	1.3% (-0.24-3.8%)	0.049 (-0.0096-0.15)	3.3 (-0.13-8.9)
Marshall Islands	1990	Female	40 to 44	26% (13-44%)	74% (56-87%)	0.55 (0.21-1.1)	0.46% (-0.89-2.5%)	0.65% (-1.3-3.5%)	0.028 (-0.054-0.16)	1.5 (-2.9-8.2)
Marshall Islands	1990	Female	45 to 49	26% (12-44%)	74% (56-88%)	0.48 (0.16-1)	-0.22% (-1.6-1.6%)	-0.15% (-2-2.3%)	-0.008 (-0.1-0.12)	-0.74 (-5.6-5.4)
Marshall Islands	1990	Female	50 to 54	26% (12-44%)	74% (56-88%)	0.43 (0.13-0.95)	-0.78% (-2.6-0.91%)	-0.8% (-2.9-1.5%)	-0.05 (-0.19-0.093)	-2.6 (-8.6-3.1)
Marshall Islands	1990	Female	55 to 59	25% (12-45%)	75% (55-88%)	0.39 (0.12-0.95)	-1% (-3-0.6%)	-1% (-3.3-0.98%)	-0.074 (-0.24-0.069)	-3.2 (-9.7-1.9)
Marshall Islands	1990	Female	60 to 64	25% (12-46%)	75% (54-88%)	0.37 (0.11-0.87)	-1.3% (-3.4-0.27%)	-1.3% (-3.8-0.51%)	-0.11 (-0.31-0.043)	-3.8 (-10-0.8)
Marshall Islands	1990	Female	65 to 69	26% (12-43%)	74% (57-88%)	0.37 (0.12-0.82)	-1.3% (-3.1-0.23%)	-1.4% (-3.4-0.39%)	-0.13 (-0.32-0.034)	-3.5 (-8.1-0.58)
Marshall Islands	1990	Female	70 to 74	26% (12-43%)	74% (57-88%)	0.36 (0.11-0.83)	-1.3% (-2.9-0.2%)	-1.4% (-3.3-0.35%)	-0.13 (-0.32-0.035)	-2.7 (-6.4-0.43)
Marshall Islands	1990	Female	75 to 79	25% (11-44%)	75% (56-89%)	0.34 (0.093-0.85)	-1% (-2.7-0.29%)	-1.1% (-3-0.4%)	-0.094 (-0.26-0.035)	-1.5 (-4-0.43)
Marshall Islands	1990	Female	80 to 84	24% (10-43%)	76% (57-90%)	0.32 (0.086-0.77)	-1% (-2.5-0.1%)	-1.1% (-2.9-0.16%)	-0.07 (-0.18-0.0099)	-0.79 (-2-0.073)
Marshall Islands	1990	Female	85 to 89	25% (11-43%)	75% (57-89%)	0.33 (0.11-0.85)	-1% (-2.5-0.1%)	-1.2% (-2.9-0.12%)	-0.041 (-0.1-0.004)	-0.32 (-0.76-0.031)
Marshall Islands	1990	Female	90 to 94	28% (13-48%)	72% (52-87%)	0.37 (0.12-0.86)	-0.97% (-2.2-0.17%)	-1.1% (-2.5-0.19%)	-0.015 (-0.033-0.0025)	-0.077 (-0.17-0.013)
Marshall Islands	1990	Female	95 plus	33% (15-56%)	67% (44-85%)	0.43 (0.12-1.1)	-1.3% (-3-0.31%)	-1.5% (-3.4-0.33%)	-0.0048 (-0.011-0.0011)	-0.016 (-0.038-0.004)
Marshall Islands	1990	Male	All Ages	39% (31-49%)	61% (51-69%)	0.64 (0.37-1.1)	1.9% (0.86-3.5%)	2% (0.65-4.3%)	3.4 (1.1-7.2)	190 (85-350)
Marshall Islands	1990	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.43 (0.16-0.87)	1.9% (0.79-3.6%)	2.4% (0.76-4.9%)	0.11 (0.035-0.23)	9.8 (4-18)
Marshall Islands	1990	Male	20 to 24	55% (37-73%)	45% (27-63%)	1.2 (0.55-2.1)	5.4% (2.5-9.6%)	6.5% (2.7-12%)	0.4 (0.16-0.73)	30 (14-53)
Marshall Islands	1990	Male	25 to 29	63% (43-78%)	37% (22-57%)	1.6 (0.8-2.9)	6.8% (3.5-12%)	8.2% (3.9-15%)	0.51 (0.24-0.93)	35 (18-61)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	1990	Male	30 to 34	64% (43-80%)	36% (20-57%)	1.7 (0.84-3.2)	6.2% (3.2-11%)	7.3% (3.4-14%)	0.47 (0.22-0.89)	30 (16-54)
Marshall Islands	1990	Male	35 to 39	62% (44-79%)	38% (21-56%)	1.5 (0.71-2.9)	4.8% (2.1-9.3%)	5.6% (2.2-11%)	0.4 (0.16-0.8)	23 (10-45)
Marshall Islands	1990	Male	40 to 44	59% (42-76%)	41% (24-58%)	1.3 (0.58-2.6)	3.7% (1.1-8%)	4.1% (0.98-9.3%)	0.33 (0.079-0.76)	17 (5.3-38)
Marshall Islands	1990	Male	45 to 49	56% (34-74%)	44% (26-66%)	1.1 (0.48-2.3)	2.9% (0.61-6.7%)	3.2% (0.44-7.8%)	0.3 (0.039-0.74)	14 (2.9-33)
Marshall Islands	1990	Male	50 to 54	53% (34-71%)	47% (29-66%)	0.94 (0.38-1.9)	2.2% (0.14-6%)	2.4% (0.0055-6.8%)	0.27 (0.00058-0.75)	11 (0.75-29)
Marshall Islands	1990	Male	55 to 59	51% (33-70%)	49% (30-67%)	0.77 (0.31-1.6)	1.4% (-0.16-4.2%)	1.5% (-0.29-4.6%)	0.18 (-0.036-0.58)	6.5 (-0.81-20)
Marshall Islands	1990	Male	60 to 64	48% (29-68%)	52% (32-71%)	0.64 (0.26-1.4)	0.97% (-0.4-3.6%)	1% (-0.46-3.9%)	0.14 (-0.062-0.53)	4.2 (-1.7-16)
Marshall Islands	1990	Male	65 to 69	46% (28-65%)	54% (35-72%)	0.58 (0.23-1.2)	0.85% (-0.36-3.1%)	0.94% (-0.39-3.5%)	0.12 (-0.05-0.46)	2.9 (-1.3-11)
Marshall Islands	1990	Male	70 to 74	44% (26-61%)	56% (39-74%)	0.52 (0.2-1.1)	0.58% (-0.42-2.7%)	0.63% (-0.48-3%)	0.072 (-0.055-0.33)	1.4 (-1-6.3)
Marshall Islands	1990	Male	75 to 79	40% (24-59%)	60% (41-76%)	0.44 (0.16-0.98)	0.43% (-0.41-2.2%)	0.45% (-0.47-2.4%)	0.038 (-0.04-0.21)	0.59 (-0.55-3)
Marshall Islands	1990	Male	80 to 84	37% (21-55%)	63% (45-79%)	0.39 (0.15-0.86)	0.34% (-0.34-1.6%)	0.35% (-0.4-1.7%)	0.018 (-0.021-0.092)	0.21 (-0.21-1)
Marshall Islands	1990	Male	85 to 89	35% (21-52%)	65% (48-79%)	0.37 (0.15-0.82)	0.17% (-0.48-1.3%)	0.15% (-0.59-1.4%)	0.0034 (-0.013-0.03)	0.033 (-0.093-0.24)
Marshall Islands	1990	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.38 (0.15-0.82)	0.12% (-0.44-1.1%)	0.074% (-0.57-1.1%)	0.00051 (-0.0039-0.0076)	0.0047 (-0.018-0.044)
Marshall Islands	1990	Male	95 plus	40% (22-62%)	60% (38-78%)	0.41 (0.14-0.94)	-0.093% (-0.79-1%)	-0.22% (-1-0.98%)	-0.00034 (-0.0016-0.0015)	-0.00056 (-0.0049-0.0062)
Marshall Islands	1995	Female	All Ages	19% (13-26%)	81% (74-87%)	0.29 (0.15-0.49)	-0.091% (-0.65-0.71%)	-0.69% (-1.6-0.59%)	-0.95 (-2.2-0.84)	-7.3 (-51-55)
Marshall Islands	1995	Female	15 to 19	20% (9-35%)	80% (65-91%)	0.35 (0.12-0.78)	0.98% (0.38-2.1%)	1.5% (0.28-3.6%)	0.036 (0.0068-0.089)	3.7 (1.4-7.6)
Marshall Islands	1995	Female	20 to 24	24% (10-43%)	76% (57-90%)	0.55 (0.19-1.2)	1.7% (0.79-3.2%)	2.1% (0.59-4.8%)	0.068 (0.018-0.16)	7 (3.2-14)
Marshall Islands	1995	Female	25 to 29	26% (10-45%)	74% (55-90%)	0.56 (0.2-1.2)	1.4% (0.59-2.8%)	1.7% (0.41-4%)	0.062 (0.015-0.15)	5.8 (2.3-11)
Marshall Islands	1995	Female	30 to 34	27% (10-48%)	73% (52-90%)	0.59 (0.21-1.2)	1.1% (0.083-2.7%)	1.2% (-0.15-3.6%)	0.052 (-0.0065-0.16)	4.4 (0.36-11)
Marshall Islands	1995	Female	35 to 39	28% (11-47%)	72% (53-89%)	0.61 (0.22-1.2)	0.69% (-0.55-2.6%)	0.85% (-0.82-3.5%)	0.045 (-0.043-0.19)	3 (-2.4-11)
Marshall Islands	1995	Female	40 to 44	27% (13-44%)	73% (56-87%)	0.56 (0.2-1.2)	0.095% (-1.5-2.3%)	0.15% (-2.1-3.1%)	0.0091 (-0.12-0.18)	0.4 (-6.6-9.7)
Marshall Islands	1995	Female	45 to 49	26% (11-45%)	74% (55-89%)	0.48 (0.16-1)	-0.59% (-2.3-1.4%)	-0.63% (-2.8-2%)	-0.043 (-0.19-0.13)	-2.5 (-10-6.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	1995	Female	50 to 54	25% (11-46%)	75% (54-89%)	0.42 (0.13-0.95)	-1.1% (-3.3-0.54%)	-1.3% (-3.8-0.81%)	-0.1 (-0.3-0.067)	-4.8 (-14-2.3)
Marshall Islands	1995	Female	55 to 59	25% (11-45%)	75% (55-89%)	0.38 (0.12-0.93)	-1.3% (-3.5-0.25%)	-1.4% (-3.9-0.42%)	-0.13 (-0.35-0.037)	-5.1 (-14-1)
Marshall Islands	1995	Female	60 to 64	25% (11-46%)	75% (54-89%)	0.36 (0.11-0.88)	-1.6% (-3.9-0.13%)	-1.7% (-4.4-0.24%)	-0.18 (-0.45-0.024)	-5.7 (-14-0.49)
Marshall Islands	1995	Female	65 to 69	27% (12-47%)	73% (53-88%)	0.38 (0.12-0.86)	-1.7% (-3.8--0.054%)	-1.8% (-4.2-0.096%)	-0.21 (-0.47-0.011)	-5.3 (-12--0.17)
Marshall Islands	1995	Female	70 to 74	27% (13-45%)	73% (55-87%)	0.37 (0.11-0.84)	-1.6% (-3.5--0.058%)	-1.8% (-3.9--0.011%)	-0.21 (-0.45--0.0012)	-4.2 (-8.9--0.14)
Marshall Islands	1995	Female	75 to 79	26% (11-45%)	74% (55-89%)	0.34 (0.093-0.85)	-1.3% (-3.2-0.14%)	-1.4% (-3.5-0.26%)	-0.15 (-0.38-0.029)	-2.3 (-5.7-0.26)
Marshall Islands	1995	Female	80 to 84	25% (11-44%)	75% (56-89%)	0.32 (0.088-0.77)	-1.3% (-2.9--0.039%)	-1.4% (-3.2--0.037%)	-0.11 (-0.25--0.0029)	-1.2 (-2.7--0.039)
Marshall Islands	1995	Female	85 to 89	26% (11-44%)	74% (56-89%)	0.33 (0.1-0.81)	-1.3% (-3--0.078%)	-1.4% (-3.3--0.096%)	-0.065 (-0.15--0.0045)	-0.5 (-1.1--0.033)
Marshall Islands	1995	Female	90 to 94	29% (13-48%)	71% (52-87%)	0.37 (0.12-0.86)	-1.2% (-2.7-0.046%)	-1.4% (-3-0.045%)	-0.022 (-0.047-0.0007)	-0.11 (-0.25-0.0042)
Marshall Islands	1995	Female	95 plus	34% (14-57%)	66% (43-86%)	0.43 (0.11-1)	-1.6% (-3.6-0.31%)	-1.8% (-4.2-0.29%)	-0.0073 (-0.016-0.0011)	-0.024 (-0.055-0.0048)
Marshall Islands	1995	Male	All Ages	39% (31-48%)	61% (52-69%)	0.64 (0.36-1.1)	1.9% (0.82-3.6%)	1.9% (0.58-4.3%)	3.4 (1.1-7.8)	200 (89-390)
Marshall Islands	1995	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.43 (0.16-0.88)	1.8% (0.74-3.5%)	2.3% (0.69-4.6%)	0.11 (0.033-0.23)	9.8 (4-18)
Marshall Islands	1995	Male	20 to 24	55% (36-72%)	45% (28-64%)	1.1 (0.54-2.1)	5.2% (2.5-9.1%)	6.2% (2.6-11%)	0.44 (0.18-0.82)	34 (16-60)
Marshall Islands	1995	Male	25 to 29	62% (43-78%)	38% (22-57%)	1.6 (0.78-2.9)	6.5% (3.3-12%)	7.7% (3.6-14%)	0.54 (0.25-1)	38 (19-67)
Marshall Islands	1995	Male	30 to 34	63% (42-79%)	37% (21-58%)	1.7 (0.81-3.2)	5.8% (2.9-11%)	6.7% (3.1-13%)	0.5 (0.23-0.98)	32 (16-60)
Marshall Islands	1995	Male	35 to 39	62% (44-78%)	38% (22-56%)	1.5 (0.71-2.8)	4.5% (1.9-9%)	5.2% (1.9-11%)	0.43 (0.16-0.88)	25 (11-50)
Marshall Islands	1995	Male	40 to 44	59% (42-76%)	41% (24-58%)	1.3 (0.56-2.5)	3.3% (0.91-7.7%)	3.7% (0.78-9%)	0.36 (0.075-0.85)	19 (5.2-43)
Marshall Islands	1995	Male	45 to 49	55% (34-73%)	45% (27-66%)	1.1 (0.45-2.2)	2.5% (0.34-6.2%)	2.8% (0.19-7.2%)	0.3 (0.02-0.77)	14 (2-36)
Marshall Islands	1995	Male	50 to 54	52% (32-71%)	48% (29-68%)	0.91 (0.36-1.8)	1.9% (-0.093-5.4%)	2% (-0.26-6.2%)	0.26 (-0.033-0.78)	11 (-0.53-31)
Marshall Islands	1995	Male	55 to 59	50% (32-70%)	50% (30-68%)	0.75 (0.29-1.6)	1.1% (-0.37-4.3%)	1.2% (-0.44-4.6%)	0.16 (-0.064-0.62)	6 (-1.9-22)
Marshall Islands	1995	Male	60 to 64	48% (28-68%)	52% (32-72%)	0.62 (0.25-1.4)	0.78% (-0.51-3.5%)	0.83% (-0.62-3.9%)	0.12 (-0.094-0.56)	3.7 (-2.5-16)
Marshall Islands	1995	Male	65 to 69	47% (28-65%)	53% (35-72%)	0.57 (0.23-1.2)	0.66% (-0.51-2.9%)	0.72% (-0.59-3.1%)	0.1 (-0.084-0.45)	2.4 (-2-11)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	1995	Male	70 to 74	44% (27-62%)	56% (38-73%)	0.51 (0.19-1.1)	0.43% (-0.59-2.4%)	0.46% (-0.66-2.5%)	0.057 (-0.08-0.31)	1.1 (-1.5-6)
Marshall Islands	1995	Male	75 to 79	40% (24-58%)	60% (42-76%)	0.43 (0.16-0.97)	0.3% (-0.56-2%)	0.3% (-0.63-2.1%)	0.029 (-0.061-0.2)	0.46 (-0.86-2.9)
Marshall Islands	1995	Male	80 to 84	37% (21-55%)	63% (45-79%)	0.38 (0.15-0.84)	0.25% (-0.43-1.5%)	0.25% (-0.49-1.6%)	0.013 (-0.026-0.093)	0.16 (-0.28-1)
Marshall Islands	1995	Male	85 to 89	35% (21-52%)	65% (48-79%)	0.36 (0.14-0.77)	0.097% (-0.58-1.2%)	0.063% (-0.68-1.2%)	0.0017 (-0.018-0.033)	0.022 (-0.13-0.26)
Marshall Islands	1995	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.37 (0.14-0.83)	0.053% (-0.5-0.95%)	-0.0014% (-0.65-0.99%)	-0.00001 (-0.005-0.0076)	0.0024 (-0.024-0.044)
Marshall Islands	1995	Male	95 plus	41% (22-61%)	59% (39-78%)	0.41 (0.14-0.96)	-0.15% (-0.87-0.92%)	-0.29% (-1.1-0.89%)	-0.00048 (-0.0019-0.0015)	-0.00097 (-0.0057-0.006)
Marshall Islands	2000	Female	All Ages	19% (13-26%)	81% (74-87%)	0.29 (0.16-0.49)	-0.21% (-0.81-0.55%)	-0.86% (-1.8-0.35%)	-1.5 (-3.1-0.63)	-20 (-77-54)
Marshall Islands	2000	Female	15 to 19	20% (9.3-36%)	80% (64-91%)	0.35 (0.12-0.78)	0.98% (0.37-2.2%)	1.4% (0.26-3.6%)	0.041 (0.0076-0.1)	4.1 (1.5-9.1)
Marshall Islands	2000	Female	20 to 24	24% (11-43%)	76% (57-89%)	0.55 (0.2-1.2)	1.6% (0.76-3.2%)	2% (0.56-4.6%)	0.077 (0.021-0.18)	7.7 (3.5-15)
Marshall Islands	2000	Female	25 to 29	26% (11-45%)	74% (55-89%)	0.56 (0.2-1.2)	1.3% (0.38-2.7%)	1.5% (0.13-3.6%)	0.071 (0.0067-0.17)	6.6 (1.9-14)
Marshall Islands	2000	Female	30 to 34	27% (10-46%)	73% (54-90%)	0.58 (0.21-1.2)	0.88% (-0.19-2.3%)	0.95% (-0.54-3.1%)	0.05 (-0.028-0.16)	4.2 (-0.95-11)
Marshall Islands	2000	Female	35 to 39	27% (12-46%)	73% (54-88%)	0.6 (0.22-1.2)	0.43% (-1-2.6%)	0.51% (-1.3-3.4%)	0.033 (-0.086-0.22)	2.2 (-5.2-13)
Marshall Islands	2000	Female	40 to 44	26% (13-44%)	74% (56-87%)	0.55 (0.21-1.2)	-0.2% (-2-2.1%)	-0.19% (-2.4-2.7%)	-0.014 (-0.19-0.21)	-1.1 (-10-11)
Marshall Islands	2000	Female	45 to 49	26% (11-43%)	74% (57-89%)	0.47 (0.16-0.97)	-0.84% (-2.8-1.2%)	-0.92% (-3.3-1.6%)	-0.082 (-0.29-0.14)	-4.5 (-15-6.5)
Marshall Islands	2000	Female	50 to 54	25% (12-44%)	75% (56-88%)	0.41 (0.13-0.94)	-1.4% (-3.7-0.35%)	-1.5% (-4.2-0.51%)	-0.16 (-0.42-0.056)	-7 (-19-2)
Marshall Islands	2000	Female	55 to 59	25% (12-44%)	75% (56-88%)	0.37 (0.12-0.9)	-1.5% (-3.8-0.086%)	-1.7% (-4.2-0.16%)	-0.19 (-0.48-0.017)	-7.2 (-18-0.39)
Marshall Islands	2000	Female	60 to 64	25% (11-44%)	75% (56-89%)	0.36 (0.11-0.86)	-1.7% (-4-0.11%)	-1.9% (-4.5-0.095%)	-0.24 (-0.56-0.012)	-7.5 (-17-0.47)
Marshall Islands	2000	Female	65 to 69	26% (12-45%)	74% (55-88%)	0.36 (0.12-0.84)	-1.8% (-3.9-0.18%)	-2% (-4.4-0.12%)	-0.28 (-0.6-0.016)	-6.9 (-15-0.71)
Marshall Islands	2000	Female	70 to 74	26% (12-44%)	74% (56-88%)	0.35 (0.11-0.8)	-1.8% (-3.7-0.13%)	-2% (-4.3-0.12%)	-0.28 (-0.6-0.018)	-5.4 (-12-0.4)
Marshall Islands	2000	Female	75 to 79	25% (11-44%)	75% (56-89%)	0.33 (0.087-0.83)	-1.4% (-3.4-0.022%)	-1.6% (-3.8-0.026%)	-0.21 (-0.48-0.0034)	-3.1 (-7.2-0.048)
Marshall Islands	2000	Female	80 to 84	24% (10-43%)	76% (57-90%)	0.31 (0.085-0.76)	-1.4% (-3.2-0.076%)	-1.6% (-3.5-0.083%)	-0.15 (-0.36-0.008)	-1.7 (-3.8-0.086)
Marshall Islands	2000	Female	85 to 89	25% (11-43%)	75% (57-89%)	0.33 (0.098-0.78)	-1.4% (-3.2-0.093%)	-1.6% (-3.6-0.11%)	-0.093 (-0.21-0.0064)	-0.7 (-1.6-0.047)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	2000	Female	90 to 94	29% (13-49%)	71% (51-87%)	0.38 (0.11-0.89)	-1.4% (-2.9-0.071%)	-1.6% (-3.3-0.03%)	-0.033 (-0.07-0.00063)	-0.17 (-0.36-0.0088)
Marshall Islands	2000	Female	95 plus	34% (15-56%)	66% (44-85%)	0.44 (0.12-1.1)	-1.7% (-4-0.25%)	-2% (-4.5-0.2%)	-0.01 (-0.023-0.00097)	-0.034 (-0.079-0.0047)
Marshall Islands	2000	Male	All Ages	39% (30-49%)	61% (51-70%)	0.66 (0.36-1.1)	1.8% (0.77-3.5%)	1.8% (0.48-3.9%)	3.7 (1-8.2)	220 (91-410)
Marshall Islands	2000	Male	15 to 19	37% (21-55%)	63% (45-79%)	0.43 (0.16-0.87)	1.9% (0.8-3.6%)	2.3% (0.72-4.8%)	0.12 (0.038-0.25)	11 (4.5-20)
Marshall Islands	2000	Male	20 to 24	55% (35-72%)	45% (28-65%)	1.2 (0.52-2.1)	5.2% (2.5-9.2%)	6.2% (2.6-11%)	0.48 (0.2-0.88)	36 (17-64)
Marshall Islands	2000	Male	25 to 29	62% (43-78%)	38% (22-57%)	1.6 (0.78-2.8)	6.5% (3.3-11%)	7.7% (3.5-14%)	0.65 (0.3-1.2)	45 (23-78)
Marshall Islands	2000	Male	30 to 34	63% (42-79%)	37% (21-58%)	1.7 (0.84-3.1)	5.6% (2.7-10%)	6.5% (2.8-12%)	0.55 (0.24-1)	35 (17-63)
Marshall Islands	2000	Male	35 to 39	62% (44-78%)	38% (22-56%)	1.5 (0.69-2.8)	4.4% (1.7-8.8%)	4.9% (1.7-10%)	0.47 (0.16-0.98)	28 (11-55)
Marshall Islands	2000	Male	40 to 44	59% (42-76%)	41% (24-58%)	1.3 (0.57-2.5)	3.2% (0.75-7.2%)	3.5% (0.55-8.3%)	0.39 (0.06-0.95)	21 (4.7-48)
Marshall Islands	2000	Male	45 to 49	56% (34-73%)	44% (27-66%)	1.1 (0.47-2.3)	2.4% (0.13-6.1%)	2.5% (-0.028-6.9%)	0.33 (-0.0036-0.9)	16 (0.88-41)
Marshall Islands	2000	Male	50 to 54	52% (33-70%)	48% (30-67%)	0.93 (0.37-1.9)	1.7% (-0.33-5.4%)	1.8% (-0.58-6%)	0.27 (-0.08-0.87)	11 (-2.3-34)
Marshall Islands	2000	Male	55 to 59	50% (32-68%)	50% (32-68%)	0.76 (0.3-1.6)	1% (-0.51-3.9%)	1% (-0.64-4.3%)	0.16 (-0.097-0.68)	6 (-3-24)
Marshall Islands	2000	Male	60 to 64	48% (28-67%)	52% (33-72%)	0.64 (0.26-1.3)	0.67% (-0.69-3.2%)	0.69% (-0.8-3.5%)	0.11 (-0.13-0.57)	3.5 (-3.6-17)
Marshall Islands	2000	Male	65 to 69	46% (27-65%)	54% (35-73%)	0.57 (0.24-1.2)	0.55% (-0.62-2.7%)	0.59% (-0.69-3%)	0.093 (-0.11-0.46)	2.3 (-2.6-11)
Marshall Islands	2000	Male	70 to 74	43% (25-61%)	57% (39-75%)	0.51 (0.19-1.1)	0.33% (-0.7-2.1%)	0.34% (-0.79-2.4%)	0.047 (-0.11-0.31)	0.95 (-1.9-6.1)
Marshall Islands	2000	Male	75 to 79	40% (23-59%)	60% (41-77%)	0.43 (0.17-0.97)	0.22% (-0.66-1.7%)	0.21% (-0.75-1.9%)	0.022 (-0.08-0.2)	0.38 (-1.2-2.9)
Marshall Islands	2000	Male	80 to 84	37% (20-55%)	63% (45-80%)	0.39 (0.15-0.82)	0.18% (-0.54-1.4%)	0.17% (-0.63-1.4%)	0.011 (-0.04-0.094)	0.14 (-0.41-1)
Marshall Islands	2000	Male	85 to 89	35% (21-53%)	65% (47-79%)	0.37 (0.15-0.75)	0.035% (-0.68-1.1%)	-0.013% (-0.82-1.1%)	-0.00041 (-0.027-0.037)	0.0098 (-0.19-0.3)
Marshall Islands	2000	Male	90 to 94	37% (23-55%)	63% (45-77%)	0.38 (0.14-0.8)	-0.0041% (-0.6-0.9%)	-0.071% (-0.74-0.93%)	-0.00076 (-0.008-0.0099)	-0.00027 (-0.038-0.058)
Marshall Islands	2000	Male	95 plus	41% (22-62%)	59% (38-78%)	0.42 (0.14-0.94)	-0.2% (-0.95-0.94%)	-0.35% (-1.2-0.9%)	-0.00077 (-0.0027-0.0019)	-0.0017 (-0.0082-0.0083)
Marshall Islands	2005	Female	All Ages	19% (13-25%)	81% (75-87%)	0.31 (0.17-0.53)	-0.31% (-1.1-0.68%)	-1% (-2.2-0.39%)	-2 (-4.4-0.85)	-32 (-110-73)
Marshall Islands	2005	Female	15 to 19	20% (9.6-35%)	80% (65-90%)	0.37 (0.13-0.8)	1% (0.38-2.2%)	1.5% (0.31-3.6%)	0.046 (0.0099-0.11)	4.6 (1.7-9.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	2005	Female	20 to 24	24% (11-42%)	76% (58-89%)	0.58 (0.22-1.2)	1.7% (0.74-3.3%)	2.1% (0.58-4.8%)	0.092 (0.026-0.22)	8.9 (3.8-17)
Marshall Islands	2005	Female	25 to 29	26% (11-44%)	74% (56-89%)	0.59 (0.2-1.2)	1.3% (0.36-2.9%)	1.5% (0.084-3.9%)	0.083 (0.005-0.23)	7.5 (2-17)
Marshall Islands	2005	Female	30 to 34	27% (10-45%)	73% (55-90%)	0.6 (0.21-1.3)	0.83% (-0.35-2.6%)	0.88% (-0.74-3.3%)	0.061 (-0.052-0.23)	5.1 (-2.3-16)
Marshall Islands	2005	Female	35 to 39	27% (11-46%)	73% (54-89%)	0.63 (0.23-1.3)	0.38% (-1.2-2.5%)	0.44% (-1.6-3.2%)	0.037 (-0.13-0.28)	2.4 (-8-16)
Marshall Islands	2005	Female	40 to 44	26% (13-44%)	74% (56-87%)	0.57 (0.22-1.2)	-0.26% (-2.1-2.3%)	-0.27% (-2.6-2.9%)	-0.026 (-0.25-0.28)	-1.7 (-14-15)
Marshall Islands	2005	Female	45 to 49	26% (11-44%)	74% (56-89%)	0.49 (0.17-1.1)	-0.91% (-3.1-1.2%)	-1% (-3.6-1.5%)	-0.11 (-0.41-0.17)	-6.1 (-21-7.6)
Marshall Islands	2005	Female	50 to 54	26% (12-44%)	74% (56-88%)	0.44 (0.13-1)	-1.5% (-3.9-0.5%)	-1.7% (-4.5-0.73%)	-0.22 (-0.6-0.092)	-9.7 (-25-3.2)
Marshall Islands	2005	Female	55 to 59	25% (11-45%)	75% (55-89%)	0.4 (0.12-0.94)	-1.6% (-4-0.23%)	-1.8% (-4.5-0.36%)	-0.26 (-0.67-0.058)	-9.6 (-24-1.5)
Marshall Islands	2005	Female	60 to 64	25% (11-45%)	75% (55-89%)	0.38 (0.11-0.89)	-1.8% (-4.2--0.1%)	-2% (-4.7--0.071%)	-0.32 (-0.76--0.012)	-9.7 (-23--0.56)
Marshall Islands	2005	Female	65 to 69	27% (12-46%)	73% (54-88%)	0.39 (0.13-0.88)	-1.9% (-4.2--0.24%)	-2.1% (-4.7--0.23%)	-0.36 (-0.82--0.039)	-8.9 (-20--1.1)
Marshall Islands	2005	Female	70 to 74	27% (12-44%)	73% (56-88%)	0.38 (0.11-0.85)	-1.9% (-4--0.22%)	-2.1% (-4.5--0.23%)	-0.38 (-0.82--0.043)	-7.4 (-16--0.81)
Marshall Islands	2005	Female	75 to 79	25% (11-45%)	75% (55-89%)	0.35 (0.097-0.9)	-1.5% (-3.6-0.07%)	-1.7% (-4-0.09%)	-0.28 (-0.65-0.015)	-4.1 (-9.5-0.17)
Marshall Islands	2005	Female	80 to 84	24% (11-44%)	76% (56-89%)	0.33 (0.094-0.81)	-1.5% (-3.5--0.1%)	-1.7% (-3.9--0.12%)	-0.21 (-0.49--0.015)	-2.2 (-5.3--0.16)
Marshall Islands	2005	Female	85 to 89	25% (11-44%)	75% (56-89%)	0.34 (0.11-0.81)	-1.5% (-3.4--0.00053%)	-1.7% (-3.8--0.05%)	-0.12 (-0.27--0.0036)	-0.9 (-2--0.00033)
Marshall Islands	2005	Female	90 to 94	28% (13-48%)	72% (52-87%)	0.39 (0.12-0.94)	-1.4% (-3.1-0.14%)	-1.7% (-3.5-0.098%)	-0.044 (-0.093-0.0026)	-0.23 (-0.49-0.021)
Marshall Islands	2005	Female	95 plus	33% (14-57%)	67% (43-86%)	0.46 (0.12-1.1)	-1.8% (-4.2-0.2%)	-2.1% (-4.9-0.17%)	-0.014 (-0.033-0.0011)	-0.047 (-0.11-0.0052)
Marshall Islands	2005	Male	All Ages	40% (32-49%)	60% (51-68%)	0.7 (0.39-1.2)	2% (0.77-4%)	1.9% (0.41-4.3%)	4.5 (1-9)	260 (97-510)
Marshall Islands	2005	Male	15 to 19	36% (21-54%)	64% (46-79%)	0.44 (0.17-0.9)	1.9% (0.85-3.7%)	2.4% (0.81-4.9%)	0.13 (0.045-0.29)	12 (5-22)
Marshall Islands	2005	Male	20 to 24	55% (36-72%)	45% (28-64%)	1.2 (0.55-2.2)	5.4% (2.6-9.4%)	6.4% (2.7-12%)	0.54 (0.22-1)	41 (19-74)
Marshall Islands	2005	Male	25 to 29	62% (43-78%)	38% (22-57%)	1.6 (0.79-2.9)	6.6% (3.4-12%)	7.8% (3.5-14%)	0.73 (0.33-1.3)	51 (26-88)
Marshall Islands	2005	Male	30 to 34	63% (42-80%)	37% (20-58%)	1.8 (0.88-3.1)	5.7% (2.6-10%)	6.5% (2.7-12%)	0.71 (0.29-1.3)	46 (21-80)
Marshall Islands	2005	Male	35 to 39	62% (43-78%)	38% (22-57%)	1.6 (0.73-2.9)	4.5% (1.6-8.9%)	5% (1.6-10%)	0.59 (0.18-1.2)	35 (12-68)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	2005	Male	40 to 44	59% (41-76%)	41% (24-59%)	1.4 (0.58-2.7)	3.2% (0.64-7.4%)	3.5% (0.5-8.5%)	0.49 (0.071-1.2)	26 (5.6-61)
Marshall Islands	2005	Male	45 to 49	55% (34-73%)	45% (27-66%)	1.2 (0.48-2.3)	2.4% (0.014-6.5%)	2.5% (-0.2-7.2%)	0.42 (-0.029-1.2)	20 (0.12-56)
Marshall Islands	2005	Male	50 to 54	52% (33-71%)	48% (29-67%)	0.97 (0.39-1.9)	1.7% (-0.44-5.5%)	1.8% (-0.67-6.1%)	0.34 (-0.13-1.2)	14 (-3.9-46)
Marshall Islands	2005	Male	55 to 59	50% (32-69%)	50% (31-68%)	0.79 (0.31-1.6)	0.98% (-0.7-4%)	1% (-0.81-4.2%)	0.19 (-0.16-0.81)	7.2 (-5.1-29)
Marshall Islands	2005	Male	60 to 64	48% (28-68%)	52% (32-72%)	0.66 (0.25-1.4)	0.65% (-0.76-3.2%)	0.65% (-0.9-3.5%)	0.13 (-0.18-0.68)	4 (-4.8-20)
Marshall Islands	2005	Male	65 to 69	46% (28-64%)	54% (36-72%)	0.6 (0.24-1.2)	0.51% (-0.72-2.6%)	0.53% (-0.82-2.8%)	0.1 (-0.16-0.55)	2.5 (-3.6-13)
Marshall Islands	2005	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.53 (0.2-1.1)	0.3% (-0.77-2%)	0.29% (-0.86-2.2%)	0.049 (-0.14-0.37)	1 (-2.6-7.3)
Marshall Islands	2005	Male	75 to 79	39% (24-58%)	61% (42-76%)	0.44 (0.17-0.97)	0.19% (-0.74-1.6%)	0.17% (-0.85-1.8%)	0.022 (-0.11-0.22)	0.39 (-1.5-3.3)
Marshall Islands	2005	Male	80 to 84	36% (21-55%)	64% (45-79%)	0.39 (0.16-0.82)	0.16% (-0.59-1.3%)	0.13% (-0.69-1.4%)	0.01 (-0.053-0.1)	0.15 (-0.55-1.2)
Marshall Islands	2005	Male	85 to 89	35% (20-52%)	65% (48-80%)	0.37 (0.15-0.79)	0.014% (-0.71-1%)	-0.044% (-0.85-1.1%)	-0.0017 (-0.034-0.043)	0.0047 (-0.24-0.36)
Marshall Islands	2005	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.38 (0.14-0.8)	-0.029% (-0.66-0.84%)	-0.11% (-0.83-0.87%)	-0.0015 (-0.011-0.012)	-0.0024 (-0.054-0.067)
Marshall Islands	2005	Male	95 plus	40% (22-61%)	60% (39-78%)	0.41 (0.13-0.93)	-0.21% (-0.98-0.83%)	-0.37% (-1.3-0.74%)	-0.0012 (-0.004-0.0023)	-0.0026 (-0.012-0.01)
Marshall Islands	2010	Female	All Ages	19% (13-26%)	81% (74-87%)	0.33 (0.17-0.57)	-0.36% (-1.2-0.75%)	-1.1% (-2.4-0.43%)	-2.4 (-5.2-0.92)	-40 (-130-84)
Marshall Islands	2010	Female	15 to 19	20% (8.9-35%)	80% (65-91%)	0.38 (0.13-0.84)	1% (0.38-2.2%)	1.5% (0.33-3.7%)	0.05 (0.011-0.13)	5 (2-11)
Marshall Islands	2010	Female	20 to 24	24% (10-42%)	76% (58-90%)	0.59 (0.21-1.3)	1.7% (0.76-3.3%)	2.1% (0.55-4.9%)	0.093 (0.025-0.22)	9.2 (4-18)
Marshall Islands	2010	Female	25 to 29	25% (11-44%)	75% (56-89%)	0.61 (0.21-1.3)	1.4% (0.35-3%)	1.5% (0.051-4.2%)	0.09 (0.0029-0.25)	8.3 (2-19)
Marshall Islands	2010	Female	30 to 34	26% (10-46%)	74% (54-90%)	0.63 (0.22-1.3)	0.88% (-0.4-2.7%)	0.93% (-0.79-3.6%)	0.068 (-0.057-0.24)	5.8 (-2.6-17)
Marshall Islands	2010	Female	35 to 39	27% (11-45%)	73% (55-89%)	0.65 (0.24-1.4)	0.38% (-1.4-2.6%)	0.44% (-1.8-3.3%)	0.044 (-0.19-0.33)	2.9 (-11-20)
Marshall Islands	2010	Female	40 to 44	26% (13-44%)	74% (56-87%)	0.61 (0.23-1.3)	-0.29% (-2.3-2.3%)	-0.29% (-2.8-2.8%)	-0.031 (-0.31-0.32)	-2.2 (-18-17)
Marshall Islands	2010	Female	45 to 49	26% (12-44%)	74% (56-88%)	0.53 (0.18-1.1)	-0.94% (-3.2-1.2%)	-1% (-3.7-1.7%)	-0.13 (-0.49-0.22)	-7.2 (-25-9.8)
Marshall Islands	2010	Female	50 to 54	26% (12-46%)	74% (54-88%)	0.46 (0.15-1)	-1.5% (-4-0.77%)	-1.7% (-4.5-1%)	-0.26 (-0.71-0.16)	-11 (-31-6.2)
Marshall Islands	2010	Female	55 to 59	25% (11-44%)	75% (56-89%)	0.42 (0.13-0.97)	-1.6% (-4-0.31%)	-1.8% (-4.6-0.44%)	-0.32 (-0.85-0.079)	-12 (-30-2.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	2010	Female	60 to 64	25% (11-44%)	75% (56-89%)	0.4 (0.11-0.94)	-1.8% (-4.2-0.014%)	-2% (-4.7-0.094%)	-0.38 (-0.9-0.018)	-12 (-27-0.092)
Marshall Islands	2010	Female	65 to 69	26% (11-45%)	74% (55-89%)	0.41 (0.13-0.93)	-1.9% (-4.3--0.16%)	-2.1% (-4.8--0.14%)	-0.43 (-0.98--0.028)	-11 (-24--0.88)
Marshall Islands	2010	Female	70 to 74	26% (11-43%)	74% (57-89%)	0.39 (0.11-0.92)	-1.9% (-4--0.21%)	-2.1% (-4.5--0.23%)	-0.44 (-0.96--0.049)	-8.5 (-18--0.96)
Marshall Islands	2010	Female	75 to 79	25% (11-43%)	75% (57-89%)	0.36 (0.096-0.9)	-1.5% (-3.6-0.0049%)	-1.7% (-4-0.0012%)	-0.34 (-0.8-0.00027)	-4.9 (-12-0.018)
Marshall Islands	2010	Female	80 to 84	24% (10-43%)	76% (57-90%)	0.35 (0.1-0.86)	-1.5% (-3.5--0.15%)	-1.7% (-3.9--0.2%)	-0.26 (-0.6--0.032)	-2.8 (-6.5--0.28)
Marshall Islands	2010	Female	85 to 89	25% (11-43%)	75% (57-89%)	0.36 (0.12-0.89)	-1.5% (-3.4--0.12%)	-1.7% (-3.9--0.16%)	-0.14 (-0.31--0.013)	-1.1 (-2.4--0.077)
Marshall Islands	2010	Female	90 to 94	28% (13-48%)	72% (52-87%)	0.41 (0.13-0.99)	-1.5% (-3.2-0.13%)	-1.7% (-3.7-0.094%)	-0.051 (-0.11-0.0028)	-0.26 (-0.57-0.022)
Marshall Islands	2010	Female	95 plus	33% (14-56%)	67% (44-86%)	0.48 (0.14-1.1)	-1.8% (-4.2-0.27%)	-2.1% (-4.9-0.19%)	-0.017 (-0.039-0.0015)	-0.055 (-0.13-0.008)
Marshall Islands	2010	Male	All Ages	40% (32-50%)	60% (50-68%)	0.74 (0.41-1.3)	2.2% (0.8-4.3%)	2% (0.38-4.5%)	5.2 (1-12)	300 (110-590)
Marshall Islands	2010	Male	15 to 19	36% (21-55%)	64% (45-79%)	0.46 (0.17-0.93)	2% (0.9-3.8%)	2.5% (0.83-5.2%)	0.15 (0.048-0.32)	13 (5.5-25)
Marshall Islands	2010	Male	20 to 24	54% (36-72%)	46% (28-64%)	1.2 (0.58-2.2)	5.5% (2.7-9.8%)	6.6% (2.9-13%)	0.58 (0.24-1.1)	43 (20-81)
Marshall Islands	2010	Male	25 to 29	61% (42-77%)	39% (23-58%)	1.7 (0.82-3.1)	6.8% (3.4-12%)	8.1% (3.6-15%)	0.81 (0.36-1.5)	56 (28-98)
Marshall Islands	2010	Male	30 to 34	63% (41-79%)	37% (21-59%)	1.8 (0.88-3.3)	5.9% (2.7-11%)	6.7% (2.8-13%)	0.79 (0.32-1.5)	51 (23-92)
Marshall Islands	2010	Male	35 to 39	62% (43-78%)	38% (22-57%)	1.6 (0.75-3.1)	4.6% (1.6-9.1%)	5.2% (1.6-11%)	0.75 (0.22-1.5)	44 (15-86)
Marshall Islands	2010	Male	40 to 44	59% (41-76%)	41% (24-59%)	1.4 (0.61-2.7)	3.3% (0.57-7.9%)	3.6% (0.46-9%)	0.58 (0.076-1.4)	31 (5.5-72)
Marshall Islands	2010	Male	45 to 49	55% (34-73%)	45% (27-66%)	1.2 (0.51-2.4)	2.5% (-0.1-6.6%)	2.6% (-0.39-7.3%)	0.51 (-0.068-1.4)	24 (-1-63)
Marshall Islands	2010	Male	50 to 54	52% (33-71%)	48% (29-67%)	1 (0.4-2)	1.8% (-0.44-5.5%)	1.9% (-0.61-6%)	0.42 (-0.14-1.3)	18 (-4.5-53)
Marshall Islands	2010	Male	55 to 59	50% (32-69%)	50% (31-68%)	0.82 (0.33-1.7)	1.1% (-0.65-4.1%)	1.1% (-0.84-4.5%)	0.25 (-0.2-1)	9.3 (-6-36)
Marshall Islands	2010	Male	60 to 64	47% (28-67%)	53% (33-72%)	0.68 (0.25-1.5)	0.7% (-0.75-3.3%)	0.71% (-0.9-3.6%)	0.16 (-0.21-0.84)	5.1 (-5.6-25)
Marshall Islands	2010	Male	65 to 69	45% (27-64%)	55% (36-73%)	0.62 (0.24-1.3)	0.54% (-0.73-2.8%)	0.56% (-0.85-3%)	0.12 (-0.19-0.67)	3.1 (-4.2-16)
Marshall Islands	2010	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.55 (0.22-1.2)	0.32% (-0.78-2.3%)	0.32% (-0.89-2.5%)	0.061 (-0.17-0.49)	1.3 (-3.1-9.4)
Marshall Islands	2010	Male	75 to 79	39% (24-58%)	61% (42-76%)	0.47 (0.18-1)	0.21% (-0.76-2%)	0.19% (-0.85-2.1%)	0.028 (-0.13-0.32)	0.51 (-1.7-4.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	2010	Male	80 to 84	37% (21-56%)	63% (44-79%)	0.42 (0.17-0.89)	0.17% (-0.6-1.5%)	0.14% (-0.72-1.6%)	0.013 (-0.062-0.15)	0.19 (-0.65-1.7)
Marshall Islands	2010	Male	85 to 89	35% (20-52%)	65% (48-80%)	0.39 (0.16-0.85)	0.029% (-0.69-1.1%)	-0.032% (-0.85-1.2%)	-0.0014 (-0.038-0.053)	0.011 (-0.26-0.43)
Marshall Islands	2010	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.4 (0.16-0.87)	-0.021% (-0.68-0.9%)	-0.1% (-0.85-0.92%)	-0.0016 (-0.013-0.014)	-0.0019 (-0.061-0.082)
Marshall Islands	2010	Male	95 plus	40% (21-60%)	60% (40-79%)	0.43 (0.15-0.97)	-0.19% (-1-0.99%)	-0.35% (-1.3-0.98%)	-0.0013 (-0.0049-0.0037)	-0.0028 (-0.015-0.015)
Marshall Islands	2016	Female	All Ages	19% (14-26%)	81% (74-86%)	0.36 (0.2-0.61)	-0.4% (-1.3-0.89%)	-1.2% (-2.6-0.62%)	-2.9 (-6.5-1.6)	-48 (-170-110)
Marshall Islands	2016	Female	15 to 19	20% (8.8-35%)	80% (65-91%)	0.41 (0.13-0.94)	1% (0.41-2.4%)	1.6% (0.38-4%)	0.054 (0.013-0.15)	5.6 (2.2-13)
Marshall Islands	2016	Female	20 to 24	24% (9.9-42%)	76% (58-90%)	0.64 (0.23-1.4)	1.8% (0.78-3.5%)	2.2% (0.64-5.4%)	0.1 (0.027-0.25)	10 (4.6-20)
Marshall Islands	2016	Female	25 to 29	25% (11-44%)	75% (56-89%)	0.65 (0.23-1.3)	1.4% (0.29-3.2%)	1.6% (-0.087-4.3%)	0.091 (-0.0053-0.26)	8.7 (1.9-20)
Marshall Islands	2016	Female	30 to 34	26% (9.8-46%)	74% (54-90%)	0.67 (0.23-1.3)	0.92% (-0.39-2.9%)	0.97% (-0.84-4%)	0.072 (-0.062-0.27)	6.4 (-2.8-20)
Marshall Islands	2016	Female	35 to 39	27% (11-45%)	73% (55-89%)	0.69 (0.25-1.5)	0.41% (-1.4-2.8%)	0.45% (-1.9-3.6%)	0.046 (-0.2-0.38)	3.4 (-11-24)
Marshall Islands	2016	Female	40 to 44	26% (12-43%)	74% (57-88%)	0.63 (0.24-1.3)	-0.23% (-2.3-2.2%)	-0.28% (-3-2.9%)	-0.035 (-0.37-0.38)	-2 (-20-20)
Marshall Islands	2016	Female	45 to 49	26% (12-44%)	74% (56-88%)	0.54 (0.19-1.2)	-0.88% (-3.1-1.4%)	-1% (-3.7-1.7%)	-0.15 (-0.55-0.26)	-7.6 (-27-12)
Marshall Islands	2016	Female	50 to 54	25% (11-44%)	75% (56-89%)	0.48 (0.15-1)	-1.4% (-3.8-1.1%)	-1.6% (-4.4-1.2%)	-0.28 (-0.81-0.22)	-12 (-35-9.1)
Marshall Islands	2016	Female	55 to 59	25% (11-45%)	75% (55-89%)	0.44 (0.15-1)	-1.5% (-4.1-0.63%)	-1.8% (-4.6-0.78%)	-0.36 (-0.99-0.15)	-13 (-36-5.3)
Marshall Islands	2016	Female	60 to 64	25% (12-45%)	75% (55-88%)	0.43 (0.13-1)	-1.8% (-4.3-0.27%)	-2% (-4.8-0.36%)	-0.46 (-1.1-0.091)	-14 (-33-2.3)
Marshall Islands	2016	Female	65 to 69	26% (12-45%)	74% (55-88%)	0.44 (0.14-1)	-1.9% (-4.3-0.16%)	-2.2% (-4.8-0.19%)	-0.51 (-1.2-0.048)	-13 (-29-1.2)
Marshall Islands	2016	Female	70 to 74	26% (12-44%)	74% (56-88%)	0.42 (0.13-1)	-1.9% (-4.1-0.17%)	-2.1% (-4.7-0.19%)	-0.53 (-1.2-0.043)	-10 (-23-0.91)
Marshall Islands	2016	Female	75 to 79	25% (11-44%)	75% (56-89%)	0.39 (0.11-0.99)	-1.5% (-3.7-0.35%)	-1.7% (-4.2-0.4%)	-0.41 (-1-0.095)	-6 (-15-1.3)
Marshall Islands	2016	Female	80 to 84	24% (11-43%)	76% (57-89%)	0.37 (0.11-0.94)	-1.5% (-3.6-0.14%)	-1.7% (-4.1-0.14%)	-0.31 (-0.78-0.026)	-3.3 (-8.2-0.32)
Marshall Islands	2016	Female	85 to 89	25% (11-44%)	75% (56-89%)	0.39 (0.13-0.93)	-1.5% (-3.6-0.099%)	-1.7% (-4.1-0.071%)	-0.17 (-0.41-0.0065)	-1.3 (-3.1-0.078)
Marshall Islands	2016	Female	90 to 94	28% (13-48%)	72% (52-87%)	0.43 (0.13-1)	-1.5% (-3.4-0.25%)	-1.7% (-4-0.18%)	-0.061 (-0.14-0.0061)	-0.31 (-0.72-0.047)
Marshall Islands	2016	Female	95 plus	33% (14-56%)	67% (44-86%)	0.51 (0.14-1.2)	-1.8% (-4.5-0.33%)	-2.2% (-5.2-0.27%)	-0.019 (-0.045-0.0023)	-0.063 (-0.15-0.011)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Marshall Islands	2016	Male	All Ages	41% (32-50%)	59% (50-68%)	0.82 (0.45-1.4)	2.4% (0.83-4.9%)	2.1% (0.41-5.2%)	6.3 (1.2-15)	360 (120-730)
Marshall Islands	2016	Male	15 to 19	36% (21-55%)	64% (45-79%)	0.5 (0.19-0.99)	2.1% (0.93-4%)	2.7% (0.92-5.5%)	0.17 (0.058-0.37)	15 (6.6-30)
Marshall Islands	2016	Male	20 to 24	55% (36-72%)	45% (28-64%)	1.3 (0.63-2.5)	5.8% (2.8-11%)	7% (3-13%)	0.68 (0.28-1.3)	51 (25-94)
Marshall Islands	2016	Male	25 to 29	62% (42-77%)	38% (23-58%)	1.9 (0.86-3.4)	7.1% (3.5-12%)	8.5% (3.8-16%)	0.88 (0.39-1.7)	62 (29-110)
Marshall Islands	2016	Male	30 to 34	63% (42-79%)	37% (21-58%)	2 (0.96-3.7)	6.1% (2.8-11%)	7.1% (2.8-14%)	0.89 (0.35-1.7)	58 (26-110)
Marshall Islands	2016	Male	35 to 39	62% (44-79%)	38% (21-56%)	1.7 (0.85-3.2)	4.9% (1.7-9.5%)	5.6% (1.7-12%)	0.86 (0.27-1.7)	51 (18-98)
Marshall Islands	2016	Male	40 to 44	59% (41-76%)	41% (24-59%)	1.5 (0.65-2.9)	3.5% (0.67-7.9%)	3.8% (0.47-9.2%)	0.75 (0.095-1.8)	41 (7.4-94)
Marshall Islands	2016	Male	45 to 49	55% (33-74%)	45% (26-67%)	1.3 (0.56-2.5)	2.6% (-0.039-6.9%)	2.8% (-0.29-7.9%)	0.63 (-0.062-1.8)	31 (-0.45-81)
Marshall Islands	2016	Male	50 to 54	52% (32-72%)	48% (28-68%)	1.1 (0.46-2.2)	2% (-0.49-6%)	2.1% (-0.67-6.5%)	0.53 (-0.17-1.7)	23 (-5.6-67)
Marshall Islands	2016	Male	55 to 59	50% (32-69%)	50% (31-68%)	0.89 (0.34-1.9)	1.2% (-0.57-4.8%)	1.2% (-0.75-5.2%)	0.34 (-0.21-1.5)	13 (-6.2-51)
Marshall Islands	2016	Male	60 to 64	47% (28-67%)	53% (33-72%)	0.73 (0.27-1.6)	0.83% (-0.72-3.9%)	0.85% (-0.88-4.3%)	0.24 (-0.25-1.2)	7.5 (-6.8-37)
Marshall Islands	2016	Male	65 to 69	46% (27-65%)	54% (35-73%)	0.66 (0.26-1.3)	0.65% (-0.7-3.3%)	0.67% (-0.81-3.6%)	0.17 (-0.2-0.91)	4.5 (-4.8-22)
Marshall Islands	2016	Male	70 to 74	43% (26-60%)	57% (40-74%)	0.58 (0.23-1.2)	0.39% (-0.73-2.6%)	0.39% (-0.84-2.8%)	0.089 (-0.2-0.64)	1.9 (-3.6-12)
Marshall Islands	2016	Male	75 to 79	39% (23-58%)	61% (42-77%)	0.5 (0.19-1.1)	0.27% (-0.74-2.1%)	0.24% (-0.86-2.3%)	0.043 (-0.16-0.39)	0.78 (-2.2-5.9)
Marshall Islands	2016	Male	80 to 84	36% (20-54%)	64% (46-80%)	0.44 (0.18-0.94)	0.23% (-0.54-1.7%)	0.2% (-0.67-1.8%)	0.021 (-0.07-0.19)	0.29 (-0.68-2.2)
Marshall Islands	2016	Male	85 to 89	34% (21-51%)	66% (49-79%)	0.42 (0.17-0.89)	0.078% (-0.71-1.4%)	0.017% (-0.88-1.4%)	0.00088 (-0.047-0.079)	0.036 (-0.33-0.64)
Marshall Islands	2016	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.44 (0.16-0.94)	0.015% (-0.75-1.1%)	-0.072% (-0.94-1.2%)	-0.0013 (-0.016-0.02)	0.0015 (-0.076-0.11)
Marshall Islands	2016	Male	95 plus	40% (21-61%)	60% (39-79%)	0.48 (0.17-1.1)	-0.15% (-1.1-1.2%)	-0.32% (-1.4-1.2%)	-0.0014 (-0.0057-0.005)	-0.0025 (-0.018-0.02)
Federated States of Micronesia	1990	Female	All Ages	15% (12-19%)	85% (81-88%)	0.23 (0.15-0.33)	0.4% (0.14-0.76%)	0.29% (-0.13-0.9%)	0.96 (-0.42-3)	72 (26-140)
Federated States of Micronesia	1990	Female	15 to 19	19% (12-30%)	81% (70-88%)	0.33 (0.15-0.58)	0.97% (0.44-1.7%)	1.4% (0.43-2.9%)	0.081 (0.024-0.17)	8.2 (3.6-15)
Federated States of Micronesia	1990	Female	20 to 24	24% (14-36%)	76% (64-86%)	0.52 (0.26-0.9)	1.7% (0.91-2.8%)	2.1% (0.82-4.1%)	0.14 (0.057-0.28)	14 (7.5-24)
Federated States of Micronesia	1990	Female	25 to 29	24% (14-37%)	76% (63-86%)	0.48 (0.24-0.81)	1.5% (0.9-2.5%)	1.8% (0.82-3.4%)	0.13 (0.059-0.26)	12 (6.6-20)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	1990	Female	30 to 34	23% (13-37%)	77% (63-87%)	0.47 (0.23-0.81)	1.3% (0.66-2.2%)	1.5% (0.63-2.9%)	0.13 (0.051-0.27)	11 (5.3-18)
Federated States of Micronesia	1990	Female	35 to 39	22% (13-36%)	78% (64-87%)	0.49 (0.24-0.89)	1.4% (0.67-2.5%)	1.7% (0.75-3.4%)	0.19 (0.079-0.39)	12 (5.8-23)
Federated States of Micronesia	1990	Female	40 to 44	21% (12-34%)	79% (66-88%)	0.45 (0.21-0.84)	1.3% (0.52-2.5%)	1.7% (0.61-3.4%)	0.16 (0.056-0.34)	8.3 (3.3-17)
Federated States of Micronesia	1990	Female	45 to 49	21% (11-33%)	79% (67-89%)	0.39 (0.18-0.72)	0.78% (0.046-1.7%)	1% (0.084-2.4%)	0.12 (0.0098-0.27)	5.1 (0.32-12)
Federated States of Micronesia	1990	Female	50 to 54	21% (12-33%)	79% (67-88%)	0.35 (0.16-0.68)	0.47% (-0.22-1.5%)	0.68% (-0.23-2.2%)	0.086 (-0.027-0.27)	2.9 (-1.4-9.7)
Federated States of Micronesia	1990	Female	55 to 59	19% (11-30%)	81% (70-89%)	0.28 (0.13-0.54)	0.27% (-0.43-1.2%)	0.43% (-0.43-1.6%)	0.075 (-0.071-0.29)	1.9 (-2.9-8.6)
Federated States of Micronesia	1990	Female	60 to 64	17% (9.8-29%)	83% (71-90%)	0.24 (0.11-0.47)	0.0092% (-0.63-0.79%)	0.095% (-0.66-1%)	0.021 (-0.14-0.23)	0.074 (-4.6-5.8)
Federated States of Micronesia	1990	Female	65 to 69	18% (10-28%)	82% (72-90%)	0.24 (0.11-0.44)	-0.019% (-0.7-0.75%)	0.063% (-0.73-0.95%)	0.018 (-0.2-0.28)	-0.15 (-5.3-5.8)
Federated States of Micronesia	1990	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.24 (0.11-0.46)	-0.17% (-0.8-0.61%)	-0.13% (-0.86-0.78%)	-0.045 (-0.29-0.25)	-1.2 (-5.7-4.4)
Federated States of Micronesia	1990	Female	75 to 79	18% (11-30%)	82% (70-89%)	0.23 (0.11-0.44)	-0.13% (-0.74-0.62%)	-0.095% (-0.79-0.73%)	-0.03 (-0.25-0.23)	-0.65 (-3.8-3.1)
Federated States of Micronesia	1990	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.23 (0.1-0.41)	-0.28% (-0.88-0.3%)	-0.3% (-0.98-0.37%)	-0.069 (-0.22-0.086)	-0.78 (-2.4-0.84)
Federated States of Micronesia	1990	Female	85 to 89	18% (11-29%)	82% (71-89%)	0.23 (0.11-0.42)	-0.32% (-0.9-0.24%)	-0.36% (-1-0.29%)	-0.041 (-0.12-0.033)	-0.31 (-0.87-0.23)
Federated States of Micronesia	1990	Female	90 to 94	20% (12-30%)	80% (70-88%)	0.24 (0.1-0.46)	-0.26% (-0.76-0.28%)	-0.29% (-0.86-0.31%)	-0.013 (-0.037-0.014)	-0.067 (-0.2-0.071)
Federated States of Micronesia	1990	Female	95 plus	22% (12-36%)	78% (64-88%)	0.27 (0.1-0.53)	-0.29% (-0.98-0.4%)	-0.36% (-1.2-0.41%)	-0.0044 (-0.014-0.0049)	-0.013 (-0.046-0.018)
Federated States of Micronesia	1990	Male	All Ages	36% (27-45%)	64% (55-73%)	0.71 (0.5-0.98)	2.1% (1.3-3.2%)	2.2% (1.1-3.6%)	9 (4.5-15)	470 (290-700)
Federated States of Micronesia	1990	Male	15 to 19	36% (21-53%)	64% (47-79%)	0.46 (0.21-0.82)	1.9% (0.86-3.5%)	2.4% (0.83-4.8%)	0.25 (0.087-0.5)	22 (9.6-40)
Federated States of Micronesia	1990	Male	20 to 24	54% (35-72%)	46% (28-65%)	1.3 (0.75-2.1)	5.9% (3.3-9.4%)	7.1% (3.7-12%)	0.96 (0.48-1.6)	72 (40-110)
Federated States of Micronesia	1990	Male	25 to 29	61% (42-77%)	39% (23-58%)	2 (1.2-3.1)	8.2% (5-12%)	9.9% (5.9-15%)	1.3 (0.72-2)	86 (54-130)
Federated States of Micronesia	1990	Male	30 to 34	62% (41-79%)	38% (21-59%)	2.2 (1.2-3.4)	7.6% (4.5-12%)	9.2% (5.2-14%)	1.3 (0.67-2.1)	80 (46-120)
Federated States of Micronesia	1990	Male	35 to 39	61% (43-78%)	39% (22-57%)	1.8 (1-2.9)	5.8% (3.2-9.4%)	6.8% (3.5-11%)	1.1 (0.55-1.8)	62 (34-100)
Federated States of Micronesia	1990	Male	40 to 44	58% (40-75%)	42% (25-60%)	1.6 (0.89-2.6)	4.6% (2.2-7.9%)	5.2% (2.2-9.2%)	0.78 (0.34-1.4)	40 (20-71)
Federated States of Micronesia	1990	Male	45 to 49	54% (33-72%)	46% (28-67%)	1.4 (0.72-2.5)	4.1% (1.7-7.4%)	4.6% (1.7-8.5%)	0.81 (0.3-1.5)	37 (15-67)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	1990	Male	50 to 54	51% (32-70%)	49% (30-68%)	1.2 (0.58-2.1)	3.3% (1.1-6.4%)	3.6% (0.94-7.3%)	0.62 (0.16-1.3)	25 (7.9-50)
Federated States of Micronesia	1990	Male	55 to 59	49% (31-69%)	51% (31-69%)	0.88 (0.44-1.6)	1.9% (0.35-4.3%)	2% (0.25-4.9%)	0.48 (0.061-1.1)	17 (3.1-38)
Federated States of Micronesia	1990	Male	60 to 64	46% (27-66%)	54% (34-73%)	0.69 (0.32-1.3)	1.3% (0.012-3.6%)	1.4% (-0.1-3.9%)	0.41 (-0.029-1.2)	12 (0.12-33)
Federated States of Micronesia	1990	Male	65 to 69	45% (26-64%)	55% (36-74%)	0.62 (0.3-1.2)	1.2% (-0.02-2.8%)	1.3% (-0.061-3.2%)	0.42 (-0.023-1.1)	9.9 (-0.17-25)
Federated States of Micronesia	1990	Male	70 to 74	42% (25-59%)	58% (41-75%)	0.55 (0.26-1)	0.81% (-0.16-2.2%)	0.88% (-0.22-2.5%)	0.33 (-0.082-0.94)	6.3 (-1.3-17)
Federated States of Micronesia	1990	Male	75 to 79	38% (23-57%)	62% (43-77%)	0.46 (0.21-0.91)	0.58% (-0.22-1.9%)	0.61% (-0.27-2.1%)	0.21 (-0.092-0.72)	3.1 (-1.2-10)
Federated States of Micronesia	1990	Male	80 to 84	36% (20-54%)	64% (46-80%)	0.41 (0.19-0.75)	0.43% (-0.2-1.3%)	0.44% (-0.26-1.5%)	0.096 (-0.058-0.31)	1.1 (-0.54-3.4)
Federated States of Micronesia	1990	Male	85 to 89	34% (21-51%)	66% (49-79%)	0.39 (0.18-0.72)	0.26% (-0.32-1%)	0.25% (-0.42-1.1%)	0.025 (-0.042-0.11)	0.22 (-0.27-0.88)
Federated States of Micronesia	1990	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.4 (0.17-0.75)	0.17% (-0.32-0.81%)	0.13% (-0.44-0.85%)	0.0057 (-0.019-0.038)	0.042 (-0.081-0.21)
Federated States of Micronesia	1990	Male	95 plus	38% (21-59%)	62% (41-79%)	0.43 (0.16-0.84)	-0.081% (-0.75-0.69%)	-0.21% (-0.97-0.66%)	-0.0022 (-0.011-0.0072)	-0.0034 (-0.032-0.029)
Federated States of Micronesia	1995	Female	All Ages	16% (13-20%)	84% (80-87%)	0.24 (0.18-0.31)	0.43% (0.17-0.75%)	0.26% (-0.17-0.79%)	0.92 (-0.59-2.7)	77 (30-130)
Federated States of Micronesia	1995	Female	15 to 19	19% (12-30%)	81% (70-88%)	0.34 (0.18-0.57)	0.99% (0.49-1.8%)	1.4% (0.52-3%)	0.089 (0.029-0.18)	9.1 (4.3-16)
Federated States of Micronesia	1995	Female	20 to 24	24% (14-36%)	76% (64-86%)	0.54 (0.29-0.87)	1.7% (0.96-2.7%)	2.1% (0.89-3.9%)	0.14 (0.055-0.27)	14 (7.7-23)
Federated States of Micronesia	1995	Female	25 to 29	24% (14-37%)	76% (63-86%)	0.49 (0.28-0.8)	1.6% (0.97-2.4%)	1.9% (0.96-3.2%)	0.14 (0.066-0.25)	12 (7.3-20)
Federated States of Micronesia	1995	Female	30 to 34	23% (13-37%)	77% (63-87%)	0.48 (0.26-0.79)	1.3% (0.75-2%)	1.5% (0.72-2.6%)	0.14 (0.065-0.26)	11 (6.5-19)
Federated States of Micronesia	1995	Female	35 to 39	22% (12-35%)	78% (65-88%)	0.51 (0.26-0.85)	1.4% (0.71-2.3%)	1.7% (0.81-3.1%)	0.22 (0.095-0.42)	14 (7.2-24)
Federated States of Micronesia	1995	Female	40 to 44	21% (12-34%)	79% (66-88%)	0.47 (0.24-0.8)	1.2% (0.51-2.2%)	1.6% (0.63-3.1%)	0.23 (0.083-0.45)	12 (5-22)
Federated States of Micronesia	1995	Female	45 to 49	21% (11-34%)	79% (66-89%)	0.41 (0.2-0.72)	0.74% (0.054-1.6%)	1% (0.078-2.3%)	0.14 (0.011-0.31)	5.9 (0.39-13)
Federated States of Micronesia	1995	Female	50 to 54	21% (12-33%)	79% (67-88%)	0.36 (0.17-0.67)	0.41% (-0.27-1.4%)	0.62% (-0.23-1.9%)	0.089 (-0.034-0.27)	2.9 (-1.9-9.9)
Federated States of Micronesia	1995	Female	55 to 59	19% (11-30%)	81% (70-89%)	0.29 (0.14-0.54)	0.21% (-0.46-1%)	0.38% (-0.43-1.4%)	0.064 (-0.072-0.24)	1.5 (-3.3-7.3)
Federated States of Micronesia	1995	Female	60 to 64	17% (9.7-29%)	83% (71-90%)	0.25 (0.12-0.46)	-0.053% (-0.71-0.62%)	0.034% (-0.73-0.83%)	0.0098 (-0.2-0.23)	-0.48 (-6.8-5.8)
Federated States of Micronesia	1995	Female	65 to 69	18% (11-30%)	82% (70-89%)	0.25 (0.13-0.46)	-0.098% (-0.77-0.61%)	-0.017% (-0.79-0.83%)	-0.0045 (-0.21-0.22)	-0.7 (-5.6-4.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	1995	Female	70 to 74	19% (11-30%)	81% (70-89%)	0.25 (0.12-0.47)	-0.26% (-0.96-0.4%)	-0.23% (-1-0.53%)	-0.092 (-0.4-0.21)	-2.2 (-8.1-3.2)
Federated States of Micronesia	1995	Female	75 to 79	19% (12-30%)	81% (70-88%)	0.24 (0.12-0.44)	-0.18% (-0.82-0.5%)	-0.16% (-0.87-0.62%)	-0.063 (-0.36-0.24)	-1.2 (-5.5-3.2)
Federated States of Micronesia	1995	Female	80 to 84	18% (11-29%)	82% (71-89%)	0.23 (0.11-0.41)	-0.35% (-0.97-0.23%)	-0.37% (-1.1-0.29%)	-0.11 (-0.31-0.083)	-1.2 (-3.4-0.83)
Federated States of Micronesia	1995	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.24 (0.11-0.42)	-0.38% (-1-0.15%)	-0.42% (-1.1-0.19%)	-0.058 (-0.16-0.027)	-0.45 (-1.2-0.17)
Federated States of Micronesia	1995	Female	90 to 94	20% (12-30%)	80% (70-88%)	0.25 (0.12-0.45)	-0.31% (-0.82-0.23%)	-0.35% (-0.95-0.27%)	-0.016 (-0.042-0.012)	-0.081 (-0.22-0.064)
Federated States of Micronesia	1995	Female	95 plus	22% (12-37%)	78% (63-88%)	0.28 (0.11-0.54)	-0.34% (-1.1-0.32%)	-0.42% (-1.3-0.31%)	-0.0052 (-0.016-0.0038)	-0.016 (-0.051-0.015)
Federated States of Micronesia	1995	Male	All Ages	36% (28-46%)	64% (54-72%)	0.75 (0.6-0.91)	2.4% (1.5-3.2%)	2.3% (1.3-3.4%)	9.9 (5.5-15)	520 (340-730)
Federated States of Micronesia	1995	Male	15 to 19	36% (20-54%)	64% (46-80%)	0.47 (0.23-0.83)	2% (0.94-3.5%)	2.5% (0.94-4.9%)	0.3 (0.11-0.57)	26 (12-45)
Federated States of Micronesia	1995	Male	20 to 24	53% (34-71%)	47% (29-66%)	1.4 (0.84-2)	6% (3.4-9.3%)	7.3% (3.8-12%)	0.93 (0.47-1.5)	70 (39-110)
Federated States of Micronesia	1995	Male	25 to 29	61% (41-77%)	39% (23-59%)	2.1 (1.3-3)	8.3% (5.4-12%)	10% (6.4-15%)	1.1 (0.68-1.7)	77 (48-110)
Federated States of Micronesia	1995	Male	30 to 34	62% (41-78%)	38% (22-59%)	2.2 (1.4-3.3)	7.7% (4.9-11%)	9.2% (5.6-14%)	1.4 (0.84-2.3)	92 (55-140)
Federated States of Micronesia	1995	Male	35 to 39	61% (42-77%)	39% (23-58%)	1.9 (1.1-2.8)	5.9% (3.5-8.6%)	6.9% (3.9-11%)	1.3 (0.69-2)	72 (43-110)
Federated States of Micronesia	1995	Male	40 to 44	58% (40-75%)	42% (25-60%)	1.6 (0.97-2.5)	4.5% (2.5-7.2%)	5.1% (2.6-8.5%)	1.1 (0.54-1.9)	60 (31-99)
Federated States of Micronesia	1995	Male	45 to 49	54% (33-72%)	46% (28-67%)	1.5 (0.78-2.4)	4% (1.7-6.4%)	4.4% (1.8-7.4%)	1 (0.43-1.8)	48 (21-80)
Federated States of Micronesia	1995	Male	50 to 54	51% (32-70%)	49% (30-68%)	1.2 (0.62-2)	3.2% (1.1-5.9%)	3.5% (1-6.6%)	0.68 (0.19-1.3)	28 (9.4-51)
Federated States of Micronesia	1995	Male	55 to 59	49% (31-68%)	51% (32-69%)	0.9 (0.48-1.6)	1.8% (0.42-4%)	1.9% (0.31-4.5%)	0.41 (0.062-0.96)	14 (3.1-33)
Federated States of Micronesia	1995	Male	60 to 64	47% (27-66%)	53% (34-73%)	0.72 (0.35-1.3)	1.2% (0.05-3.2%)	1.3% (-0.063-3.5%)	0.54 (-0.025-1.5)	16 (0.6-42)
Federated States of Micronesia	1995	Male	65 to 69	45% (27-63%)	55% (37-73%)	0.65 (0.33-1.1)	1.1% (-0.0012-2.6%)	1.2% (-0.034-3%)	0.39 (-0.0096-0.94)	9.1 (-0.0093-22)
Federated States of Micronesia	1995	Male	70 to 74	43% (25-61%)	57% (39-75%)	0.57 (0.29-1.1)	0.77% (-0.19-2.1%)	0.83% (-0.24-2.3%)	0.33 (-0.095-0.92)	6.3 (-1.7-17)
Federated States of Micronesia	1995	Male	75 to 79	39% (23-58%)	61% (42-77%)	0.49 (0.24-0.89)	0.55% (-0.23-1.7%)	0.58% (-0.31-1.9%)	0.23 (-0.12-0.75)	3.5 (-1.4-11)
Federated States of Micronesia	1995	Male	80 to 84	36% (20-53%)	64% (47-80%)	0.43 (0.22-0.76)	0.41% (-0.2-1.2%)	0.42% (-0.29-1.3%)	0.098 (-0.067-0.31)	1.1 (-0.54-3.4)
Federated States of Micronesia	1995	Male	85 to 89	34% (20-51%)	66% (49-80%)	0.41 (0.19-0.7)	0.24% (-0.33-1%)	0.23% (-0.44-1.1%)	0.026 (-0.049-0.12)	0.24 (-0.32-0.97)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	1995	Male	90 to 94	35% (22-52%)	65% (48-78%)	0.41 (0.19-0.76)	0.15% (-0.33-0.76%)	0.11% (-0.47-0.78%)	0.0038 (-0.016-0.028)	0.032 (-0.071-0.16)
Federated States of Micronesia	1995	Male	95 plus	39% (21-59%)	61% (41-79%)	0.45 (0.18-0.86)	-0.093% (-0.73-0.66%)	-0.23% (-0.97-0.62%)	-0.0025 (-0.011-0.0067)	-0.004 (-0.031-0.029)
Federated States of Micronesia	2000	Female	All Ages	15% (12-19%)	85% (81-88%)	0.21 (0.17-0.27)	0.33% (0.1-0.61%)	0.082% (-0.3-0.5%)	0.29 (-1.1-1.8)	58 (17-100)
Federated States of Micronesia	2000	Female	15 to 19	19% (11-30%)	81% (70-89%)	0.3 (0.16-0.51)	0.86% (0.45-1.5%)	1.2% (0.42-2.5%)	0.075 (0.025-0.16)	8.3 (4.1-15)
Federated States of Micronesia	2000	Female	20 to 24	22% (13-34%)	78% (66-87%)	0.45 (0.24-0.72)	1.4% (0.81-2.2%)	1.6% (0.71-3%)	0.11 (0.044-0.21)	12 (6.9-20)
Federated States of Micronesia	2000	Female	25 to 29	21% (13-33%)	79% (67-87%)	0.38 (0.23-0.6)	1.3% (0.85-1.9%)	1.4% (0.77-2.3%)	0.11 (0.05-0.19)	11 (6.6-16)
Federated States of Micronesia	2000	Female	30 to 34	20% (11-32%)	80% (68-89%)	0.37 (0.2-0.6)	1.1% (0.63-1.6%)	1.1% (0.55-1.9%)	0.11 (0.051-0.2)	9.9 (5.6-15)
Federated States of Micronesia	2000	Female	35 to 39	20% (11-33%)	80% (67-89%)	0.41 (0.21-0.71)	1.1% (0.59-1.9%)	1.3% (0.61-2.4%)	0.17 (0.07-0.31)	11 (5.6-18)
Federated States of Micronesia	2000	Female	40 to 44	20% (12-32%)	80% (68-88%)	0.4 (0.2-0.69)	0.94% (0.35-1.7%)	1.2% (0.41-2.3%)	0.19 (0.059-0.36)	10 (3.6-18)
Federated States of Micronesia	2000	Female	45 to 49	20% (10-32%)	80% (68-90%)	0.35 (0.17-0.6)	0.49% (-0.11-1.2%)	0.69% (-0.12-1.7%)	0.12 (-0.02-0.31)	5.2 (-1.3-13)
Federated States of Micronesia	2000	Female	50 to 54	20% (11-31%)	80% (69-89%)	0.31 (0.15-0.58)	0.16% (-0.55-0.85%)	0.3% (-0.57-1.2%)	0.057 (-0.11-0.23)	1.5 (-5-8.1)
Federated States of Micronesia	2000	Female	55 to 59	18% (11-30%)	82% (70-89%)	0.25 (0.12-0.46)	0.012% (-0.64-0.6%)	0.13% (-0.66-0.87%)	0.024 (-0.12-0.16)	0.091 (-4.9-4.6)
Federated States of Micronesia	2000	Female	60 to 64	17% (9.3-28%)	83% (72-91%)	0.22 (0.1-0.4)	-0.2% (-0.89-0.35%)	-0.14% (-0.92-0.52%)	-0.029 (-0.19-0.099)	-1.4 (-6.1-2.3)
Federated States of Micronesia	2000	Female	65 to 69	18% (11-28%)	82% (72-89%)	0.22 (0.11-0.38)	-0.25% (-0.9-0.31%)	-0.2% (-0.93-0.47%)	-0.063 (-0.29-0.14)	-2.2 (-7.8-2.6)
Federated States of Micronesia	2000	Female	70 to 74	19% (11-30%)	81% (70-89%)	0.22 (0.11-0.4)	-0.42% (-1-0.11%)	-0.42% (-1.1-0.23%)	-0.12 (-0.31-0.059)	-2.5 (-6.3-0.7)
Federated States of Micronesia	2000	Female	75 to 79	18% (11-29%)	82% (71-89%)	0.21 (0.1-0.39)	-0.33% (-0.95-0.18%)	-0.32% (-1-0.28%)	-0.14 (-0.46-0.12)	-2.3 (-7-1.3)
Federated States of Micronesia	2000	Female	80 to 84	18% (10-30%)	82% (70-90%)	0.21 (0.1-0.36)	-0.47% (-1.1--0.015%)	-0.51% (-1.2-0.017%)	-0.19 (-0.46-0.0066)	-2.1 (-5--0.068)
Federated States of Micronesia	2000	Female	85 to 89	18% (11-28%)	82% (72-89%)	0.21 (0.11-0.37)	-0.48% (-1.1--0.043%)	-0.54% (-1.2--0.036%)	-0.1 (-0.24--0.007)	-0.79 (-1.8--0.071)
Federated States of Micronesia	2000	Female	90 to 94	20% (12-31%)	80% (69-88%)	0.23 (0.11-0.41)	-0.41% (-0.97-0.0079%)	-0.47% (-1.1-0.012%)	-0.028 (-0.067-0.00069)	-0.15 (-0.35-0.0026)
Federated States of Micronesia	2000	Female	95 plus	22% (12-36%)	78% (64-88%)	0.25 (0.1-0.48)	-0.45% (-1.2-0.12%)	-0.55% (-1.4-0.12%)	-0.0074 (-0.019-0.0017)	-0.023 (-0.063-0.0064)
Federated States of Micronesia	2000	Male	All Ages	39% (31-47%)	61% (53-69%)	0.76 (0.65-0.89)	2.3% (1.5-3.1%)	2.1% (1.2-3%)	8.6 (4.8-13)	480 (320-650)
Federated States of Micronesia	2000	Male	15 to 19	35% (21-53%)	65% (47-79%)	0.43 (0.22-0.73)	1.9% (0.91-3.2%)	2.4% (0.91-4.5%)	0.28 (0.11-0.53)	24 (12-42)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	2000	Male	20 to 24	54% (36-72%)	46% (28-64%)	1.3 (0.84-1.9)	5.6% (3.3-8.6%)	6.8% (3.6-11%)	0.93 (0.5-1.5)	70 (41-110)
Federated States of Micronesia	2000	Male	25 to 29	64% (47-78%)	36% (22-53%)	2.1 (1.4-2.8)	7.9% (5.2-11%)	9.6% (6.1-14%)	1.1 (0.69-1.7)	79 (51-110)
Federated States of Micronesia	2000	Male	30 to 34	66% (47-81%)	34% (19-53%)	2.3 (1.4-3.1)	7.2% (4.8-10%)	8.6% (5.4-13%)	1.2 (0.71-1.9)	78 (48-110)
Federated States of Micronesia	2000	Male	35 to 39	64% (47-79%)	36% (21-53%)	1.9 (1.2-2.6)	5.5% (3.3-8.1%)	6.4% (3.6-9.8%)	1.1 (0.6-1.8)	67 (37-100)
Federated States of Micronesia	2000	Male	40 to 44	60% (43-75%)	40% (25-57%)	1.6 (0.96-2.3)	4% (2-6.5%)	4.5% (1.9-7.5%)	0.92 (0.39-1.6)	49 (24-82)
Federated States of Micronesia	2000	Male	45 to 49	54% (35-70%)	46% (30-65%)	1.4 (0.83-2)	3.3% (1.5-5.4%)	3.6% (1.3-6.1%)	1 (0.38-1.8)	48 (21-81)
Federated States of Micronesia	2000	Male	50 to 54	50% (32-67%)	50% (33-68%)	1.1 (0.6-1.7)	2.5% (0.82-4.7%)	2.7% (0.76-5.2%)	0.74 (0.2-1.4)	31 (9.8-55)
Federated States of Micronesia	2000	Male	55 to 59	47% (30-66%)	53% (34-70%)	0.79 (0.43-1.4)	1.4% (0.19-3%)	1.4% (0.046-3.3%)	0.3 (0.0075-0.7)	11 (1.5-24)
Federated States of Micronesia	2000	Male	60 to 64	45% (26-65%)	55% (35-74%)	0.63 (0.31-1.1)	0.88% (-0.17-2.4%)	0.91% (-0.31-2.6%)	0.23 (-0.075-0.66)	7.1 (-1.3-20)
Federated States of Micronesia	2000	Male	65 to 69	44% (26-62%)	56% (38-74%)	0.58 (0.3-1)	0.78% (-0.14-2%)	0.85% (-0.21-2.2%)	0.32 (-0.083-0.88)	7.8 (-1.3-21)
Federated States of Micronesia	2000	Male	70 to 74	42% (24-59%)	58% (41-76%)	0.52 (0.26-0.92)	0.5% (-0.33-1.6%)	0.53% (-0.41-1.8%)	0.13 (-0.1-0.43)	2.5 (-1.6-8.1)
Federated States of Micronesia	2000	Male	75 to 79	38% (22-57%)	62% (43-78%)	0.44 (0.22-0.8)	0.35% (-0.35-1.3%)	0.36% (-0.42-1.5%)	0.15 (-0.18-0.61)	2.4 (-2.4-9.1)
Federated States of Micronesia	2000	Male	80 to 84	35% (19-54%)	65% (46-81%)	0.39 (0.2-0.66)	0.27% (-0.3-0.91%)	0.27% (-0.38-0.99%)	0.08 (-0.12-0.3)	0.98 (-1.1-3.3)
Federated States of Micronesia	2000	Male	85 to 89	34% (19-51%)	66% (49-81%)	0.37 (0.19-0.63)	0.12% (-0.4-0.68%)	0.088% (-0.52-0.71%)	0.013 (-0.074-0.1)	0.15 (-0.5-0.85)
Federated States of Micronesia	2000	Male	90 to 94	35% (21-53%)	65% (47-79%)	0.37 (0.18-0.68)	0.052% (-0.44-0.54%)	-0.0023% (-0.55-0.54%)	-0.00009 (-0.027-0.027)	0.015 (-0.13-0.16)
Federated States of Micronesia	2000	Male	95 plus	39% (21-59%)	61% (41-79%)	0.41 (0.16-0.77)	-0.18% (-0.84-0.42%)	-0.33% (-1.1-0.35%)	-0.0036 (-0.012-0.0038)	-0.0078 (-0.035-0.018)
Federated States of Micronesia	2005	Female	All Ages	15% (12-19%)	85% (81-88%)	0.2 (0.15-0.25)	0.24% (-0.053-0.52%)	-0.099% (-0.54-0.32%)	-0.36 (-2-1.1)	38 (-8.2-82)
Federated States of Micronesia	2005	Female	15 to 19	19% (11-29%)	81% (71-89%)	0.27 (0.15-0.45)	0.77% (0.38-1.3%)	1% (0.32-2%)	0.061 (0.018-0.13)	7 (3.5-12)
Federated States of Micronesia	2005	Female	20 to 24	22% (13-34%)	78% (66-87%)	0.4 (0.22-0.65)	1.3% (0.76-2%)	1.4% (0.57-2.6%)	0.11 (0.043-0.22)	13 (7.6-21)
Federated States of Micronesia	2005	Female	25 to 29	20% (12-33%)	80% (67-88%)	0.34 (0.2-0.56)	1.2% (0.74-1.7%)	1.2% (0.56-2%)	0.089 (0.038-0.16)	9.2 (5.5-13)
Federated States of Micronesia	2005	Female	30 to 34	20% (12-32%)	80% (68-88%)	0.33 (0.18-0.56)	0.95% (0.53-1.5%)	0.98% (0.4-1.7%)	0.077 (0.031-0.14)	6.9 (3.7-11)
Federated States of Micronesia	2005	Female	35 to 39	20% (12-33%)	80% (67-88%)	0.38 (0.2-0.65)	0.95% (0.44-1.6%)	1.1% (0.4-2.1%)	0.14 (0.047-0.26)	8.8 (4.1-15)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	2005	Female	40 to 44	20% (12-32%)	80% (68-88%)	0.36 (0.18-0.63)	0.77% (0.17-1.5%)	0.98% (0.14-2%)	0.14 (0.021-0.3)	7.7 (1.8-15)
Federated States of Micronesia	2005	Female	45 to 49	19% (10-31%)	81% (69-90%)	0.31 (0.15-0.53)	0.32% (-0.27-0.91%)	0.47% (-0.33-1.3%)	0.093 (-0.063-0.26)	3.8 (-3.2-11)
Federated States of Micronesia	2005	Female	50 to 54	20% (12-32%)	80% (68-88%)	0.28 (0.14-0.5)	-0.016% (-0.72-0.58%)	0.083% (-0.72-0.85%)	0.02 (-0.19-0.21)	-0.2 (-8.4-7.4)
Federated States of Micronesia	2005	Female	55 to 59	18% (11-30%)	82% (70-89%)	0.23 (0.12-0.42)	-0.14% (-0.85-0.4%)	-0.044% (-0.86-0.62%)	-0.01 (-0.21-0.16)	-1.4 (-8.4-4.2)
Federated States of Micronesia	2005	Female	60 to 64	17% (9.4-29%)	83% (71-91%)	0.2 (0.091-0.36)	-0.32% (-1-0.19%)	-0.28% (-1.1-0.33%)	-0.063 (-0.25-0.079)	-2.4 (-7.9-1.6)
Federated States of Micronesia	2005	Female	65 to 69	18% (11-29%)	82% (71-89%)	0.2 (0.098-0.35)	-0.37% (-1.1-0.13%)	-0.34% (-1.2-0.28%)	-0.092 (-0.31-0.071)	-2.8 (-7.9-0.93)
Federated States of Micronesia	2005	Female	70 to 74	19% (11-30%)	81% (70-89%)	0.2 (0.096-0.36)	-0.55% (-1.2--0.023%)	-0.56% (-1.3-0.03%)	-0.24 (-0.56-0.013)	-4.9 (-11--0.2)
Federated States of Micronesia	2005	Female	75 to 79	18% (11-29%)	82% (71-89%)	0.19 (0.092-0.35)	-0.44% (-1-0.044%)	-0.45% (-1.1-0.088%)	-0.17 (-0.45-0.034)	-2.8 (-6.8-0.24)
Federated States of Micronesia	2005	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.19 (0.095-0.32)	-0.56% (-1.2--0.12%)	-0.62% (-1.3--0.12%)	-0.26 (-0.58--0.051)	-2.9 (-6.2--0.61)
Federated States of Micronesia	2005	Female	85 to 89	18% (11-28%)	82% (72-89%)	0.19 (0.098-0.35)	-0.56% (-1.2--0.12%)	-0.62% (-1.3--0.13%)	-0.19 (-0.39--0.038)	-1.4 (-3--0.3)
Federated States of Micronesia	2005	Female	90 to 94	19% (12-30%)	81% (70-88%)	0.21 (0.1-0.38)	-0.49% (-1--0.074%)	-0.55% (-1.2--0.082%)	-0.054 (-0.12--0.0079)	-0.28 (-0.6--0.044)
Federated States of Micronesia	2005	Female	95 plus	22% (12-36%)	78% (64-88%)	0.23 (0.092-0.43)	-0.53% (-1.3-0.046%)	-0.63% (-1.5-0.029%)	-0.014 (-0.032-0.00063)	-0.043 (-0.1-0.0038)
Federated States of Micronesia	2005	Male	All Ages	39% (31-48%)	61% (52-69%)	0.68 (0.57-0.79)	2.2% (1.4-3%)	1.7% (0.86-2.6%)	6.7 (3.4-11)	400 (250-570)
Federated States of Micronesia	2005	Male	15 to 19	35% (20-53%)	65% (47-80%)	0.37 (0.19-0.62)	1.7% (0.82-2.9%)	2.1% (0.82-4.1%)	0.24 (0.089-0.48)	22 (11-38)
Federated States of Micronesia	2005	Male	20 to 24	55% (37-72%)	45% (28-63%)	1.1 (0.74-1.6)	5% (3-7.6%)	6% (3.2-9.7%)	0.94 (0.49-1.6)	73 (42-120)
Federated States of Micronesia	2005	Male	25 to 29	65% (48-79%)	35% (21-52%)	1.8 (1.3-2.6)	7% (4.4-10%)	8.5% (5-12%)	1.1 (0.62-1.6)	75 (46-110)
Federated States of Micronesia	2005	Male	30 to 34	66% (48-81%)	34% (19-52%)	2 (1.3-2.8)	6.3% (3.9-9.3%)	7.5% (4.3-11%)	0.81 (0.44-1.3)	53 (31-80)
Federated States of Micronesia	2005	Male	35 to 39	64% (48-79%)	36% (21-52%)	1.7 (1.1-2.5)	4.8% (2.7-7.2%)	5.5% (2.8-8.7%)	0.75 (0.37-1.2)	45 (24-72)
Federated States of Micronesia	2005	Male	40 to 44	60% (43-75%)	40% (25-57%)	1.4 (0.86-2.1)	3.3% (1.5-5.6%)	3.7% (1.4-6.5%)	0.73 (0.28-1.3)	40 (18-69)
Federated States of Micronesia	2005	Male	45 to 49	54% (35-70%)	46% (30-65%)	1.2 (0.71-1.8)	2.6% (1-4.5%)	2.8% (0.82-5%)	0.71 (0.21-1.3)	35 (13-63)
Federated States of Micronesia	2005	Male	50 to 54	50% (32-68%)	50% (32-68%)	0.95 (0.54-1.5)	2% (0.48-3.6%)	2.1% (0.37-4%)	0.71 (0.13-1.4)	30 (7.5-59)
Federated States of Micronesia	2005	Male	55 to 59	48% (30-66%)	52% (34-70%)	0.7 (0.38-1.2)	1% (-0.059-2.4%)	1% (-0.22-2.7%)	0.32 (-0.066-0.83)	12 (-0.77-30)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	2005	Male	60 to 64	45% (26-65%)	55% (35-74%)	0.56 (0.28-1)	0.59% (-0.34-1.9%)	0.59% (-0.48-2.1%)	0.14 (-0.12-0.51)	4.5 (-2.6-15)
Federated States of Micronesia	2005	Male	65 to 69	44% (25-63%)	56% (37-75%)	0.52 (0.26-0.89)	0.51% (-0.32-1.5%)	0.53% (-0.39-1.7%)	0.15 (-0.11-0.49)	3.7 (-2.3-12)
Federated States of Micronesia	2005	Male	70 to 74	41% (24-60%)	59% (40-76%)	0.45 (0.22-0.8)	0.27% (-0.45-1.2%)	0.27% (-0.56-1.3%)	0.11 (-0.22-0.53)	2.3 (-4-11)
Federated States of Micronesia	2005	Male	75 to 79	37% (22-57%)	63% (43-78%)	0.38 (0.19-0.69)	0.18% (-0.49-0.93%)	0.17% (-0.57-0.99%)	0.047 (-0.16-0.28)	0.84 (-2.2-4.3)
Federated States of Micronesia	2005	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.34 (0.17-0.6)	0.15% (-0.38-0.68%)	0.12% (-0.47-0.71%)	0.037 (-0.14-0.22)	0.53 (-1.4-2.4)
Federated States of Micronesia	2005	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.32 (0.16-0.56)	0.011% (-0.5-0.48%)	-0.037% (-0.61-0.5%)	-0.0084 (-0.14-0.11)	0.022 (-0.96-0.93)
Federated States of Micronesia	2005	Male	90 to 94	34% (21-51%)	66% (49-79%)	0.33 (0.15-0.59)	-0.031% (-0.48-0.36%)	-0.097% (-0.66-0.35%)	-0.0072 (-0.049-0.026)	-0.013 (-0.22-0.16)
Federated States of Micronesia	2005	Male	95 plus	37% (20-58%)	63% (42-80%)	0.35 (0.14-0.67)	-0.24% (-0.84-0.28%)	-0.39% (-1.1-0.19%)	-0.0069 (-0.02-0.0034)	-0.017 (-0.06-0.019)
Federated States of Micronesia	2010	Female	All Ages	16% (13-20%)	84% (80-87%)	0.2 (0.15-0.27)	0.18% (-0.13-0.46%)	-0.17% (-0.62-0.24%)	-0.66 (-2.4-0.94)	28 (-19-74)
Federated States of Micronesia	2010	Female	15 to 19	18% (11-29%)	82% (71-89%)	0.25 (0.12-0.43)	0.74% (0.37-1.2%)	0.93% (0.26-1.9%)	0.056 (0.015-0.12)	6.8 (3.3-12)
Federated States of Micronesia	2010	Female	20 to 24	23% (14-35%)	77% (65-86%)	0.4 (0.2-0.65)	1.3% (0.73-2%)	1.3% (0.51-2.7%)	0.098 (0.035-0.2)	12 (6.4-20)
Federated States of Micronesia	2010	Female	25 to 29	23% (13-36%)	77% (64-87%)	0.37 (0.21-0.61)	1.2% (0.73-1.8%)	1.3% (0.55-2.2%)	0.11 (0.047-0.21)	11 (6.7-18)
Federated States of Micronesia	2010	Female	30 to 34	23% (13-36%)	77% (64-87%)	0.36 (0.18-0.61)	0.8% (0.35-1.3%)	0.75% (0.073-1.6%)	0.055 (0.0051-0.12)	5.4 (2.3-9.5)
Federated States of Micronesia	2010	Female	35 to 39	22% (12-35%)	78% (65-88%)	0.38 (0.2-0.66)	0.91% (0.39-1.6%)	1.1% (0.32-2%)	0.096 (0.027-0.19)	6.3 (2.5-11)
Federated States of Micronesia	2010	Female	40 to 44	20% (12-33%)	80% (67-88%)	0.35 (0.17-0.62)	0.7% (0.08-1.4%)	0.86% (0.027-1.9%)	0.12 (0.004-0.26)	6.4 (0.78-13)
Federated States of Micronesia	2010	Female	45 to 49	20% (11-32%)	80% (68-89%)	0.3 (0.15-0.54)	0.27% (-0.38-0.86%)	0.37% (-0.42-1.2%)	0.068 (-0.077-0.22)	2.8 (-3.8-9.4)
Federated States of Micronesia	2010	Female	50 to 54	20% (11-33%)	80% (67-89%)	0.27 (0.13-0.47)	-0.088% (-0.83-0.49%)	-0.026% (-0.9-0.69%)	-0.0067 (-0.25-0.19)	-1.1 (-11-6.6)
Federated States of Micronesia	2010	Female	55 to 59	18% (11-30%)	82% (70-89%)	0.22 (0.11-0.4)	-0.19% (-0.94-0.33%)	-0.13% (-0.98-0.52%)	-0.041 (-0.31-0.18)	-2.6 (-12-4.6)
Federated States of Micronesia	2010	Female	60 to 64	17% (9.5-28%)	83% (72-91%)	0.19 (0.086-0.35)	-0.38% (-1.1-0.16%)	-0.36% (-1.2-0.26%)	-0.11 (-0.38-0.08)	-4 (-12-1.6)
Federated States of Micronesia	2010	Female	65 to 69	18% (11-28%)	82% (72-89%)	0.19 (0.09-0.33)	-0.41% (-1.1-0.089%)	-0.4% (-1.2-0.18%)	-0.11 (-0.35-0.052)	-3.3 (-8.9-0.7)
Federated States of Micronesia	2010	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.19 (0.087-0.34)	-0.59% (-1.2--0.068%)	-0.62% (-1.3--0.036%)	-0.21 (-0.47--0.012)	-4.3 (-9.2--0.54)
Federated States of Micronesia	2010	Female	75 to 79	18% (11-29%)	82% (71-89%)	0.18 (0.081-0.32)	-0.48% (-1.1--0.037%)	-0.5% (-1.2--0.0021%)	-0.23 (-0.58--0.00094)	-3.6 (-8.7--0.29)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	2010	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.17 (0.083-0.3)	-0.58% (-1.2--0.15%)	-0.64% (-1.3--0.14%)	-0.28 (-0.62--0.06)	-3.1 (-6.6--0.76)
Federated States of Micronesia	2010	Female	85 to 89	18% (11-28%)	82% (72-89%)	0.17 (0.087-0.3)	-0.58% (-1.2--0.13%)	-0.65% (-1.3--0.14%)	-0.17 (-0.35--0.037)	-1.3 (-2.7--0.3)
Federated States of Micronesia	2010	Female	90 to 94	19% (12-29%)	81% (71-88%)	0.19 (0.088-0.35)	-0.52% (-1--0.13%)	-0.59% (-1.2--0.15%)	-0.072 (-0.15--0.018)	-0.37 (-0.76--0.086)
Federated States of Micronesia	2010	Female	95 plus	22% (12-35%)	78% (65-88%)	0.21 (0.084-0.39)	-0.55% (-1.3-0.025%)	-0.67% (-1.5--0.011%)	-0.019 (-0.043--0.00029)	-0.06 (-0.14-0.0026)
Federated States of Micronesia	2010	Male	All Ages	38% (30-48%)	62% (52-70%)	0.62 (0.49-0.76)	1.9% (1.1-2.8%)	1.4% (0.56-2.4%)	5.9 (2.3-10)	360 (200-550)
Federated States of Micronesia	2010	Male	15 to 19	34% (20-52%)	66% (48-80%)	0.34 (0.16-0.58)	1.6% (0.81-2.8%)	2% (0.73-3.9%)	0.21 (0.076-0.44)	20 (10-36)
Federated States of Micronesia	2010	Male	20 to 24	52% (34-70%)	48% (30-66%)	1 (0.62-1.5)	4.6% (2.7-7.2%)	5.4% (2.8-8.8%)	0.89 (0.45-1.5)	69 (40-110)
Federated States of Micronesia	2010	Male	25 to 29	59% (40-76%)	41% (24-60%)	1.5 (0.99-2.2)	6.1% (3.7-8.8%)	7.2% (4-11%)	1.2 (0.64-1.8)	82 (50-120)
Federated States of Micronesia	2010	Male	30 to 34	61% (39-78%)	39% (22-61%)	1.6 (0.96-2.5)	5.1% (2.5-8%)	5.7% (2.3-9.4%)	0.72 (0.28-1.2)	48 (22-78)
Federated States of Micronesia	2010	Male	35 to 39	59% (41-77%)	41% (23-59%)	1.4 (0.8-2.3)	4.3% (2.2-7%)	4.8% (2.2-8.1%)	0.53 (0.24-0.96)	32 (16-54)
Federated States of Micronesia	2010	Male	40 to 44	57% (39-75%)	43% (25-61%)	1.2 (0.66-1.9)	3% (1.2-5.3%)	3.2% (0.99-5.8%)	0.52 (0.15-1.1)	28 (11-52)
Federated States of Micronesia	2010	Male	45 to 49	53% (32-72%)	47% (28-68%)	1.1 (0.6-1.8)	2.4% (0.76-4.2%)	2.4% (0.54-4.6%)	0.64 (0.14-1.3)	31 (10-59)
Federated States of Micronesia	2010	Male	50 to 54	50% (31-69%)	50% (31-69%)	0.89 (0.47-1.5)	1.7% (0.24-3.5%)	1.8% (0.027-3.8%)	0.6 (0.0097-1.4)	26 (3.5-55)
Federated States of Micronesia	2010	Male	55 to 59	48% (30-67%)	52% (33-70%)	0.66 (0.35-1.1)	0.81% (-0.25-2.2%)	0.78% (-0.44-2.3%)	0.36 (-0.21-1)	14 (-4.1-37)
Federated States of Micronesia	2010	Male	60 to 64	45% (26-65%)	55% (35-74%)	0.52 (0.25-0.96)	0.43% (-0.5-1.7%)	0.41% (-0.68-1.8%)	0.16 (-0.28-0.72)	5.5 (-6.8-21)
Federated States of Micronesia	2010	Male	65 to 69	43% (25-62%)	57% (38-75%)	0.47 (0.23-0.82)	0.35% (-0.41-1.3%)	0.35% (-0.52-1.4%)	0.098 (-0.14-0.39)	2.6 (-3-9.4)
Federated States of Micronesia	2010	Male	70 to 74	41% (24-58%)	59% (42-76%)	0.4 (0.19-0.73)	0.17% (-0.5-0.96%)	0.15% (-0.58-1.1%)	0.047 (-0.19-0.32)	1.1 (-3.3-6.2)
Federated States of Micronesia	2010	Male	75 to 79	37% (22-56%)	63% (44-78%)	0.34 (0.16-0.63)	0.099% (-0.52-0.74%)	0.069% (-0.61-0.77%)	0.027 (-0.24-0.31)	0.63 (-3.2-4.8)
Federated States of Micronesia	2010	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.31 (0.15-0.54)	0.088% (-0.43-0.59%)	0.057% (-0.53-0.6%)	0.017 (-0.16-0.19)	0.32 (-1.5-2.2)
Federated States of Micronesia	2010	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.29 (0.15-0.51)	-0.041% (-0.53-0.39%)	-0.098% (-0.66-0.39%)	-0.017 (-0.12-0.066)	-0.061 (-0.82-0.58)
Federated States of Micronesia	2010	Male	90 to 94	34% (21-51%)	66% (49-79%)	0.29 (0.14-0.52)	-0.066% (-0.5-0.28%)	-0.14% (-0.66-0.26%)	-0.012 (-0.058-0.023)	-0.035 (-0.26-0.14)
Federated States of Micronesia	2010	Male	95 plus	37% (20-57%)	63% (43-80%)	0.31 (0.13-0.61)	-0.26% (-0.86-0.19%)	-0.41% (-1.1-0.11%)	-0.0085 (-0.023-0.0023)	-0.021 (-0.069-0.015)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	2016	Female	All Ages	16% (13-21%)	84% (79-87%)	0.2 (0.13-0.3)	0.15% (-0.17-0.5%)	-0.21% (-0.67-0.25%)	-0.82 (-2.6-0.96)	24 (-26-81)
Federated States of Micronesia	2016	Female	15 to 19	18% (11-28%)	82% (72-89%)	0.24 (0.11-0.44)	0.72% (0.36-1.3%)	0.9% (0.22-2.1%)	0.051 (0.011-0.12)	6.4 (3-11)
Federated States of Micronesia	2016	Female	20 to 24	23% (14-35%)	77% (65-86%)	0.39 (0.18-0.7)	1.2% (0.69-2.1%)	1.3% (0.39-2.9%)	0.1 (0.03-0.23)	13 (6.8-22)
Federated States of Micronesia	2016	Female	25 to 29	23% (13-36%)	77% (64-87%)	0.36 (0.18-0.62)	1.2% (0.68-1.8%)	1.2% (0.48-2.3%)	0.11 (0.04-0.21)	11 (6.3-18)
Federated States of Micronesia	2016	Female	30 to 34	22% (12-36%)	78% (64-88%)	0.34 (0.17-0.61)	0.89% (0.46-1.5%)	0.87% (0.24-1.7%)	0.094 (0.024-0.2)	8.9 (4.4-15)
Federated States of Micronesia	2016	Female	35 to 39	22% (12-35%)	78% (65-88%)	0.36 (0.17-0.7)	0.74% (0.22-1.6%)	0.8% (0.068-1.9%)	0.08 (0.0061-0.21)	5.7 (1.7-13)
Federated States of Micronesia	2016	Female	40 to 44	20% (12-33%)	80% (67-88%)	0.34 (0.15-0.66)	0.53% (0.045-1.4%)	0.61% (0.016-1.8%)	0.06 (0.0013-0.17)	3.5 (0.28-8.8)
Federated States of Micronesia	2016	Female	45 to 49	20% (11-32%)	80% (68-89%)	0.29 (0.12-0.54)	0.23% (-0.45-0.92%)	0.32% (-0.54-1.3%)	0.053 (-0.086-0.22)	2.2 (-4.2-9)
Federated States of Micronesia	2016	Female	50 to 54	20% (11-33%)	80% (67-89%)	0.26 (0.11-0.49)	-0.12% (-0.86-0.55%)	-0.067% (-0.93-0.73%)	-0.015 (-0.23-0.18)	-1.4 (-9.7-5.8)
Federated States of Micronesia	2016	Female	55 to 59	18% (11-30%)	82% (70-89%)	0.21 (0.094-0.41)	-0.22% (-0.92-0.37%)	-0.16% (-1-0.52%)	-0.056 (-0.34-0.19)	-3.1 (-13-4.8)
Federated States of Micronesia	2016	Female	60 to 64	17% (9.5-28%)	83% (72-91%)	0.18 (0.076-0.35)	-0.38% (-1.1-0.13%)	-0.37% (-1.2-0.22%)	-0.16 (-0.52-0.096)	-5.6 (-16-1.9)
Federated States of Micronesia	2016	Female	65 to 69	17% (10-28%)	83% (72-90%)	0.18 (0.079-0.33)	-0.44% (-1.1-0.12%)	-0.43% (-1.2-0.17%)	-0.18 (-0.53-0.076)	-5.1 (-13-1.4)
Federated States of Micronesia	2016	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.18 (0.079-0.35)	-0.6% (-1.2--0.086%)	-0.64% (-1.4--0.058%)	-0.24 (-0.51--0.022)	-4.9 (-10--0.72)
Federated States of Micronesia	2016	Female	75 to 79	18% (11-28%)	82% (72-89%)	0.17 (0.07-0.32)	-0.49% (-1.1--0.028%)	-0.51% (-1.2-0.0093%)	-0.19 (-0.46-0.0037)	-2.9 (-6.9--0.18)
Federated States of Micronesia	2016	Female	80 to 84	18% (10-28%)	82% (72-90%)	0.17 (0.074-0.31)	-0.59% (-1.2--0.15%)	-0.65% (-1.3--0.15%)	-0.26 (-0.54--0.058)	-2.8 (-5.8--0.71)
Federated States of Micronesia	2016	Female	85 to 89	18% (11-27%)	82% (73-89%)	0.17 (0.077-0.32)	-0.58% (-1.2--0.15%)	-0.66% (-1.3--0.16%)	-0.19 (-0.38--0.044)	-1.4 (-2.9--0.33)
Federated States of Micronesia	2016	Female	90 to 94	19% (12-29%)	81% (71-88%)	0.18 (0.074-0.36)	-0.52% (-1--0.15%)	-0.6% (-1.2--0.17%)	-0.063 (-0.13--0.017)	-0.32 (-0.67--0.087)
Federated States of Micronesia	2016	Female	95 plus	22% (12-35%)	78% (65-88%)	0.2 (0.077-0.43)	-0.56% (-1.3--0.0013%)	-0.67% (-1.5--0.025%)	-0.023 (-0.053--0.00081)	-0.074 (-0.17--0.00016)
Federated States of Micronesia	2016	Male	All Ages	39% (31-49%)	61% (51-69%)	0.65 (0.43-0.92)	1.9% (0.98-3.2%)	1.4% (0.41-2.7%)	6.2 (1.7-13)	380 (190-660)
Federated States of Micronesia	2016	Male	15 to 19	34% (20-51%)	66% (49-80%)	0.33 (0.14-0.61)	1.6% (0.77-2.8%)	1.9% (0.69-3.9%)	0.19 (0.067-0.4)	18 (8.8-33)
Federated States of Micronesia	2016	Male	20 to 24	52% (33-70%)	48% (30-67%)	0.97 (0.52-1.6)	4.4% (2.4-7.3%)	5.2% (2.5-9.1%)	0.85 (0.38-1.5)	67 (36-110)
Federated States of Micronesia	2016	Male	25 to 29	59% (39-76%)	41% (24-61%)	1.5 (0.86-2.3)	5.8% (3.3-9%)	6.9% (3.5-11%)	1.2 (0.61-2.1)	89 (49-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Federated States of Micronesia	2016	Male	30 to 34	60% (39-78%)	40% (22-61%)	1.6 (0.86-2.6)	5.1% (2.7-8.3%)	5.8% (2.7-10%)	1.1 (0.51-2)	73 (38-120)
Federated States of Micronesia	2016	Male	35 to 39	59% (40-77%)	41% (23-60%)	1.4 (0.74-2.3)	3.3% (0.92-6.3%)	3.6% (0.43-7.3%)	0.57 (0.063-1.2)	35 (9.8-69)
Federated States of Micronesia	2016	Male	40 to 44	56% (38-74%)	44% (26-62%)	1.2 (0.59-2)	2.7% (0.83-5.5%)	2.9% (0.43-6.2%)	0.39 (0.058-0.86)	21 (6.4-44)
Federated States of Micronesia	2016	Male	45 to 49	52% (31-71%)	48% (29-69%)	1.1 (0.51-1.9)	2.3% (0.49-4.8%)	2.4% (0.11-5.1%)	0.47 (0.019-1.1)	23 (4.4-49)
Federated States of Micronesia	2016	Male	50 to 54	49% (29-69%)	51% (31-71%)	0.86 (0.42-1.6)	1.7% (0.00082-3.8%)	1.7% (-0.23-4.2%)	0.58 (-0.075-1.5)	25 (0.013-59)
Federated States of Micronesia	2016	Male	55 to 59	47% (29-67%)	53% (33-71%)	0.64 (0.3-1.2)	0.79% (-0.37-2.4%)	0.77% (-0.56-2.5%)	0.33 (-0.25-1.1)	13 (-5.8-39)
Federated States of Micronesia	2016	Male	60 to 64	45% (26-66%)	55% (34-74%)	0.5 (0.22-0.98)	0.42% (-0.51-1.7%)	0.39% (-0.71-1.8%)	0.23 (-0.42-1.1)	7.8 (-10-33)
Federated States of Micronesia	2016	Male	65 to 69	43% (25-62%)	57% (38-75%)	0.45 (0.21-0.82)	0.32% (-0.46-1.3%)	0.32% (-0.57-1.5%)	0.17 (-0.28-0.78)	4.4 (-6.2-19)
Federated States of Micronesia	2016	Male	70 to 74	40% (24-58%)	60% (42-76%)	0.39 (0.18-0.74)	0.16% (-0.53-1%)	0.14% (-0.64-1.1%)	0.046 (-0.22-0.37)	1.1 (-3.7-7.2)
Federated States of Micronesia	2016	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.33 (0.14-0.64)	0.11% (-0.49-0.81%)	0.073% (-0.59-0.87%)	0.021 (-0.17-0.25)	0.49 (-2.3-3.7)
Federated States of Micronesia	2016	Male	80 to 84	33% (18-51%)	67% (49-82%)	0.29 (0.13-0.56)	0.11% (-0.39-0.63%)	0.076% (-0.48-0.68%)	0.021 (-0.13-0.19)	0.35 (-1.3-2.1)
Federated States of Micronesia	2016	Male	85 to 89	32% (19-48%)	68% (52-81%)	0.28 (0.12-0.54)	-0.03% (-0.52-0.43%)	-0.087% (-0.66-0.42%)	-0.017 (-0.13-0.08)	-0.05 (-0.87-0.7)
Federated States of Micronesia	2016	Male	90 to 94	33% (20-51%)	67% (49-80%)	0.29 (0.12-0.57)	-0.061% (-0.51-0.33%)	-0.13% (-0.66-0.3%)	-0.0087 (-0.043-0.02)	-0.023 (-0.2-0.13)
Federated States of Micronesia	2016	Male	95 plus	37% (19-57%)	63% (43-81%)	0.32 (0.12-0.67)	-0.26% (-0.87-0.23%)	-0.41% (-1.1-0.16%)	-0.01 (-0.029-0.004)	-0.025 (-0.086-0.023)
Papua New Guinea	1990	Female	All Ages	20% (14-27%)	80% (73-86%)	1 (0.66-1.5)	2% (1.2-2.9%)	3.1% (1.6-4.9%)	630 (320-1000)	25000 (15000-37000)
Papua New Guinea	1990	Female	15 to 19	21% (9.7-37%)	79% (63-90%)	1.3 (0.5-2.5)	2.6% (1.1-4.9%)	3.9% (1.4-7.8%)	17 (6.1-36)	1400 (560-2700)
Papua New Guinea	1990	Female	20 to 24	26% (12-45%)	74% (55-88%)	2 (0.86-4)	4% (2.1-6.9%)	5.4% (2.5-9.4%)	28 (13-51)	2100 (1100-3700)
Papua New Guinea	1990	Female	25 to 29	27% (11-47%)	73% (53-89%)	2.1 (0.89-3.8)	4.7% (2.4-7.7%)	6.1% (3-10%)	36 (16-61)	2400 (1200-4000)
Papua New Guinea	1990	Female	30 to 34	28% (11-49%)	72% (51-89%)	2.1 (0.78-4.1)	4.5% (2.3-7.4%)	5.8% (2.7-9.5%)	38 (17-66)	2300 (1200-3900)
Papua New Guinea	1990	Female	35 to 39	28% (12-48%)	72% (52-88%)	2.2 (0.85-4.1)	5% (2.6-8.1%)	6.3% (3.2-10%)	51 (24-88)	2800 (1400-4700)
Papua New Guinea	1990	Female	40 to 44	28% (14-46%)	72% (54-86%)	2 (0.81-3.9)	5.3% (2.5-8.9%)	6.6% (3.1-11%)	55 (24-97)	2700 (1200-4600)
Papua New Guinea	1990	Female	45 to 49	27% (12-46%)	73% (54-88%)	1.7 (0.66-3.4)	4.9% (2.1-9%)	6.1% (2.6-11%)	55 (22-100)	2400 (960-4300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	1990	Female	50 to 54	27% (13-46%)	73% (54-87%)	1.5 (0.58-3.2)	4.6% (1.5-9%)	5.6% (1.9-11%)	59 (20-120)	2200 (730-4400)
Papua New Guinea	1990	Female	55 to 59	27% (13-47%)	73% (53-87%)	1.4 (0.53-3.1)	4% (1.2-8.4%)	4.7% (1.4-9.9%)	56 (16-120)	1800 (500-3800)
Papua New Guinea	1990	Female	60 to 64	27% (13-47%)	73% (53-87%)	1.3 (0.48-3.1)	3.3% (0.62-7.4%)	3.7% (0.74-8.4%)	48 (9.5-110)	1300 (250-3100)
Papua New Guinea	1990	Female	65 to 69	28% (13-46%)	72% (54-87%)	1.4 (0.48-2.9)	3.6% (0.79-7.6%)	4% (0.9-8.6%)	57 (14-120)	1300 (300-2800)
Papua New Guinea	1990	Female	70 to 74	27% (12-45%)	73% (55-88%)	1.3 (0.44-2.8)	3.1% (0.54-7.2%)	3.5% (0.62-8.1%)	49 (8.7-120)	890 (150-2200)
Papua New Guinea	1990	Female	75 to 79	26% (11-44%)	74% (56-89%)	1.2 (0.39-2.8)	3.1% (0.48-7.2%)	3.4% (0.5-8%)	40 (6.2-95)	560 (86-1300)
Papua New Guinea	1990	Female	80 to 84	26% (11-45%)	74% (55-89%)	1.2 (0.36-2.6)	2.6% (0.45-6.5%)	2.9% (0.47-7.2%)	24 (3.9-60)	250 (43-630)
Papua New Guinea	1990	Female	85 to 89	27% (12-47%)	73% (53-88%)	1.2 (0.44-2.7)	2.5% (0.42-6.4%)	2.8% (0.43-7.1%)	9.3 (1.5-24)	70 (12-180)
Papua New Guinea	1990	Female	90 to 94	30% (15-50%)	70% (50-85%)	1.4 (0.47-3)	2.3% (0.32-5.8%)	2.6% (0.32-6.4%)	2.5 (0.31-6.2)	13 (1.8-32)
Papua New Guinea	1990	Female	95 plus	35% (16-57%)	65% (43-84%)	1.6 (0.5-3.6)	2.5% (0.042-6.6%)	2.7% (-0.039-7.2%)	0.6 (-0.0085-1.6)	2.1 (0.035-5.4)
Papua New Guinea	1990	Male	All Ages	40% (32-50%)	60% (50-68%)	2.3 (1.6-3.1)	4.9% (3.4-6.2%)	6.6% (4.5-8.6%)	1500 (1000-2100)	70000 (49000-92000)
Papua New Guinea	1990	Male	15 to 19	39% (23-57%)	61% (43-77%)	1.6 (0.71-2.9)	4.7% (2.2-8.7%)	6.2% (2.7-12%)	41 (18-78)	3100 (1400-5800)
Papua New Guinea	1990	Male	20 to 24	57% (38-74%)	43% (26-62%)	4.2 (2.3-6.7)	14% (8.4-19%)	17% (10-24%)	130 (82-200)	9300 (5800-13000)
Papua New Guinea	1990	Male	25 to 29	64% (45-79%)	36% (21-55%)	5.8 (3.3-9.1)	16% (11-20%)	19% (13-26%)	150 (100-210)	9600 (6500-13000)
Papua New Guinea	1990	Male	30 to 34	65% (45-81%)	35% (19-55%)	6.2 (3.5-9.9)	15% (11-20%)	18% (13-24%)	150 (100-210)	8900 (6200-12000)
Papua New Guinea	1990	Male	35 to 39	64% (46-79%)	36% (21-54%)	5.5 (2.9-9.4)	14% (9.2-18%)	17% (11-22%)	160 (99-220)	8400 (5400-12000)
Papua New Guinea	1990	Male	40 to 44	61% (44-77%)	39% (23-56%)	4.8 (2.3-8.5)	12% (7.7-17%)	15% (8.7-20%)	160 (92-230)	7500 (4500-11000)
Papua New Guinea	1990	Male	45 to 49	58% (36-75%)	42% (25-64%)	4.2 (2-7.4)	11% (6.5-15%)	13% (7.3-18%)	160 (90-240)	6800 (3900-10000)
Papua New Guinea	1990	Male	50 to 54	55% (35-74%)	45% (26-65%)	3.5 (1.6-6.4)	9.4% (4.8-14%)	11% (5.4-16%)	150 (78-230)	5800 (3000-8600)
Papua New Guinea	1990	Male	55 to 59	53% (34-73%)	47% (27-66%)	2.8 (1.3-5.3)	7.3% (3.4-11%)	8% (3.7-13%)	120 (55-200)	4100 (1800-6400)
Papua New Guinea	1990	Male	60 to 64	50% (30-70%)	50% (30-70%)	2.4 (1.1-4.7)	6% (2.4-9.9%)	6.5% (2.6-11%)	100 (41-180)	2900 (1200-4900)
Papua New Guinea	1990	Male	65 to 69	48% (29-68%)	52% (32-71%)	2.2 (0.98-4.1)	5.5% (2.3-9.2%)	6% (2.5-10%)	84 (33-140)	1900 (780-3300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	1990	Male	70 to 74	46% (27-63%)	54% (37-73%)	1.9 (0.86-3.7)	4.6% (1.7-8.1%)	5% (1.8-8.9%)	56 (20-100)	1000 (380-1800)
Papua New Guinea	1990	Male	75 to 79	42% (26-60%)	58% (40-74%)	1.6 (0.7-3.1)	3.8% (1.3-7%)	4.1% (1.4-7.6%)	30 (9.9-57)	440 (150-810)
Papua New Guinea	1990	Male	80 to 84	39% (22-57%)	61% (43-78%)	1.4 (0.64-2.7)	3.4% (1.3-6.1%)	3.6% (1.3-6.6%)	14 (5.2-25)	150 (58-270)
Papua New Guinea	1990	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.4 (0.6-2.6)	3% (0.99-5.7%)	3.2% (0.99-6.2%)	5.3 (1.6-10)	41 (14-79)
Papua New Guinea	1990	Male	90 to 94	39% (24-56%)	61% (44-76%)	1.4 (0.59-2.8)	2.4% (0.75-4.8%)	2.6% (0.72-5.3%)	1.3 (0.35-2.7)	7.1 (2.2-14)
Papua New Guinea	1990	Male	95 plus	42% (24-63%)	58% (37-76%)	1.5 (0.55-3.2)	2.4% (0.44-5.3%)	2.5% (0.3-5.7%)	0.29 (0.035-0.67)	1.1 (0.19-2.5)
Papua New Guinea	1995	Female	All Ages	20% (14-27%)	80% (73-86%)	1.1 (0.73-1.7)	2.3% (1.4-3.3%)	3.4% (1.8-5.4%)	780 (410-1300)	30000 (18000-45000)
Papua New Guinea	1995	Female	15 to 19	21% (9.8-38%)	79% (62-90%)	1.4 (0.52-2.7)	2.8% (1.2-5.2%)	4.1% (1.5-8.2%)	19 (7-39)	1500 (640-2900)
Papua New Guinea	1995	Female	20 to 24	26% (11-44%)	74% (56-89%)	2.1 (0.87-4.1)	4.1% (2.1-7%)	5.6% (2.6-9.8%)	36 (16-64)	2700 (1400-4600)
Papua New Guinea	1995	Female	25 to 29	27% (12-46%)	73% (54-88%)	2.2 (0.95-4)	4.9% (2.6-8%)	6.4% (3.2-11%)	43 (21-73)	2900 (1500-4800)
Papua New Guinea	1995	Female	30 to 34	28% (11-49%)	72% (51-89%)	2.3 (0.87-4.4)	4.8% (2.4-7.7%)	6.1% (2.9-10%)	45 (21-78)	2800 (1400-4700)
Papua New Guinea	1995	Female	35 to 39	29% (13-48%)	71% (52-87%)	2.4 (0.98-4.5)	5.5% (2.9-8.6%)	6.9% (3.6-11%)	62 (30-100)	3400 (1700-5600)
Papua New Guinea	1995	Female	40 to 44	28% (14-47%)	72% (53-86%)	2.2 (0.91-4.4)	5.7% (2.8-9.4%)	7.2% (3.4-12%)	72 (33-120)	3500 (1700-5900)
Papua New Guinea	1995	Female	45 to 49	28% (12-47%)	72% (53-88%)	1.9 (0.71-3.7)	5.3% (2.2-9.5%)	6.6% (2.8-12%)	70 (29-130)	3000 (1200-5500)
Papua New Guinea	1995	Female	50 to 54	27% (12-47%)	73% (53-88%)	1.7 (0.62-3.3)	5% (1.7-9.6%)	6% (2.1-12%)	73 (26-140)	2700 (960-5400)
Papua New Guinea	1995	Female	55 to 59	26% (12-46%)	74% (54-88%)	1.5 (0.55-3.3)	4.3% (1.3-8.9%)	5.1% (1.6-10%)	69 (21-140)	2200 (670-4700)
Papua New Guinea	1995	Female	60 to 64	27% (12-47%)	73% (53-88%)	1.4 (0.5-3.3)	3.5% (0.66-7.7%)	4% (0.77-8.6%)	59 (12-130)	1600 (300-3600)
Papua New Guinea	1995	Female	65 to 69	28% (13-47%)	72% (53-87%)	1.5 (0.53-3)	3.9% (0.9-8.1%)	4.4% (1-9.2%)	71 (17-150)	1600 (360-3400)
Papua New Guinea	1995	Female	70 to 74	28% (14-47%)	72% (53-86%)	1.4 (0.51-2.9)	3.4% (0.67-7.7%)	3.8% (0.67-8.5%)	62 (11-150)	1100 (200-2700)
Papua New Guinea	1995	Female	75 to 79	26% (12-47%)	74% (53-88%)	1.3 (0.43-2.9)	3.4% (0.56-7.5%)	3.7% (0.62-8.5%)	51 (8.9-120)	720 (120-1600)
Papua New Guinea	1995	Female	80 to 84	26% (12-46%)	74% (54-88%)	1.3 (0.41-2.8)	2.9% (0.56-6.9%)	3.2% (0.59-7.7%)	32 (5.9-78)	340 (64-820)
Papua New Guinea	1995	Female	85 to 89	27% (12-45%)	73% (55-88%)	1.3 (0.48-2.9)	2.8% (0.52-6.9%)	3.1% (0.54-7.7%)	13 (2.3-32)	96 (19-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	1995	Female	90 to 94	30% (15-50%)	70% (50-85%)	1.5 (0.5-3.2)	2.5% (0.46-6.2%)	2.8% (0.49-6.9%)	3.2 (0.58-8)	17 (3.1-42)
Papua New Guinea	1995	Female	95 plus	35% (16-57%)	65% (43-84%)	1.7 (0.53-3.8)	2.8% (0.22-7%)	3% (0.16-7.8%)	0.82 (0.043-2.1)	2.9 (0.23-7.2)
Papua New Guinea	1995	Male	All Ages	41% (33-50%)	59% (50-67%)	2.5 (1.7-3.4)	5.2% (3.8-6.8%)	7% (4.9-9.1%)	1800 (1200-2400)	81000 (58000-110000)
Papua New Guinea	1995	Male	15 to 19	39% (23-57%)	61% (43-77%)	1.7 (0.75-3.1)	4.9% (2.3-8.8%)	6.4% (2.8-12%)	44 (19-84)	3400 (1600-6200)
Papua New Guinea	1995	Male	20 to 24	56% (38-73%)	44% (27-62%)	4.5 (2.5-7.1)	14% (8.7-19%)	17% (11-24%)	170 (100-240)	12000 (7400-16000)
Papua New Guinea	1995	Male	25 to 29	63% (44-79%)	37% (21-56%)	6.2 (3.5-9.4)	16% (11-21%)	19% (14-26%)	180 (120-240)	11000 (7800-15000)
Papua New Guinea	1995	Male	30 to 34	65% (44-80%)	35% (20-56%)	6.6 (3.7-10)	15% (11-20%)	18% (13-24%)	170 (120-230)	10000 (7100-13000)
Papua New Guinea	1995	Male	35 to 39	64% (45-79%)	36% (21-55%)	5.8 (3.2-10)	14% (9.9-18%)	17% (12-22%)	180 (110-240)	9400 (6200-13000)
Papua New Guinea	1995	Male	40 to 44	61% (43-77%)	39% (23-57%)	5.1 (2.5-9.1)	13% (8-17%)	15% (9.3-20%)	180 (110-260)	8800 (5400-12000)
Papua New Guinea	1995	Male	45 to 49	57% (36-74%)	43% (26-64%)	4.4 (2.1-8)	11% (6.9-15%)	13% (7.7-18%)	180 (100-260)	7500 (4500-11000)
Papua New Guinea	1995	Male	50 to 54	54% (34-72%)	46% (28-66%)	3.7 (1.7-6.7)	9.6% (5.3-14%)	11% (5.8-16%)	170 (92-260)	6400 (3500-9500)
Papua New Guinea	1995	Male	55 to 59	52% (34-71%)	48% (29-66%)	3 (1.4-5.7)	7.5% (3.6-11%)	8.3% (3.9-13%)	140 (65-230)	4600 (2200-7300)
Papua New Guinea	1995	Male	60 to 64	50% (30-70%)	50% (30-70%)	2.5 (1.1-4.9)	6.2% (2.7-10%)	6.8% (2.9-11%)	120 (50-190)	3300 (1400-5400)
Papua New Guinea	1995	Male	65 to 69	48% (29-67%)	52% (33-71%)	2.3 (1-4.3)	5.8% (2.6-9.4%)	6.4% (2.7-10%)	100 (41-170)	2300 (960-3800)
Papua New Guinea	1995	Male	70 to 74	46% (28-64%)	54% (36-72%)	2.1 (0.94-4)	4.9% (1.9-8.4%)	5.3% (2-9.2%)	68 (25-120)	1300 (480-2200)
Papua New Guinea	1995	Male	75 to 79	42% (26-60%)	58% (40-74%)	1.8 (0.77-3.3)	4.1% (1.4-7.1%)	4.4% (1.5-7.8%)	38 (13-68)	540 (190-970)
Papua New Guinea	1995	Male	80 to 84	39% (22-57%)	61% (43-78%)	1.6 (0.67-2.9)	3.6% (1.4-6.3%)	3.9% (1.5-6.9%)	18 (6.9-31)	200 (77-340)
Papua New Guinea	1995	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.5 (0.63-2.8)	3.2% (1.1-5.9%)	3.4% (1.1-6.4%)	6.8 (2.3-13)	53 (19-100)
Papua New Guinea	1995	Male	90 to 94	39% (24-56%)	61% (44-76%)	1.5 (0.63-3)	2.6% (0.83-5.1%)	2.8% (0.77-5.7%)	1.6 (0.46-3.3)	8.9 (2.9-18)
Papua New Guinea	1995	Male	95 plus	42% (24-63%)	58% (37-76%)	1.6 (0.58-3.4)	2.6% (0.54-5.8%)	2.7% (0.41-6.1%)	0.38 (0.057-0.85)	1.4 (0.28-3.1)
Papua New Guinea	2000	Female	All Ages	20% (13-27%)	80% (73-87%)	1.1 (0.68-1.5)	2.2% (1.2-3.3%)	3.2% (1.6-5.2%)	810 (390-1300)	32000 (18000-48000)
Papua New Guinea	2000	Female	15 to 19	21% (9.3-36%)	79% (64-91%)	1.3 (0.48-2.5)	2.2% (0.91-4.3%)	3.3% (1.1-6.7%)	16 (5.3-32)	1300 (540-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	2000	Female	20 to 24	25% (11-45%)	75% (55-89%)	2 (0.81-3.9)	3.7% (1.9-6.1%)	4.9% (2.2-8.8%)	33 (14-58)	2500 (1300-4300)
Papua New Guinea	2000	Female	25 to 29	27% (11-46%)	73% (54-89%)	2 (0.9-3.7)	4.4% (2.3-7%)	5.7% (2.8-9.5%)	49 (23-83)	3300 (1700-5500)
Papua New Guinea	2000	Female	30 to 34	27% (12-47%)	73% (53-88%)	2.1 (0.82-4)	4.2% (2.2-7.1%)	5.4% (2.5-9.2%)	47 (22-82)	2900 (1500-5000)
Papua New Guinea	2000	Female	35 to 39	28% (11-47%)	72% (53-89%)	2.2 (0.84-4.1)	4.9% (2.4-7.9%)	6.1% (3-10%)	64 (29-110)	3500 (1600-6000)
Papua New Guinea	2000	Female	40 to 44	27% (13-45%)	73% (55-87%)	2 (0.8-4)	5.2% (2.4-8.8%)	6.5% (2.9-11%)	74 (32-130)	3600 (1600-6400)
Papua New Guinea	2000	Female	45 to 49	27% (12-44%)	73% (56-88%)	1.7 (0.66-3.4)	4.8% (1.9-9%)	5.9% (2.4-11%)	78 (31-150)	3400 (1300-6400)
Papua New Guinea	2000	Female	50 to 54	26% (12-46%)	74% (54-88%)	1.5 (0.57-3)	4.4% (1.4-8.9%)	5.3% (1.7-11%)	78 (25-160)	2900 (930-5900)
Papua New Guinea	2000	Female	55 to 59	26% (12-45%)	74% (55-88%)	1.4 (0.49-3.1)	3.8% (1-8.2%)	4.5% (1.2-9.7%)	72 (19-160)	2300 (600-5200)
Papua New Guinea	2000	Female	60 to 64	26% (12-45%)	74% (55-88%)	1.3 (0.47-3)	3.1% (0.48-7.1%)	3.5% (0.58-8.2%)	61 (10-140)	1700 (250-4000)
Papua New Guinea	2000	Female	65 to 69	27% (13-46%)	73% (54-87%)	1.3 (0.49-2.8)	3.4% (0.61-7.5%)	3.9% (0.71-8.6%)	71 (14-160)	1600 (310-3700)
Papua New Guinea	2000	Female	70 to 74	27% (13-45%)	73% (55-87%)	1.3 (0.46-2.7)	3% (0.34-7.2%)	3.3% (0.41-8%)	62 (7.3-150)	1100 (120-2800)
Papua New Guinea	2000	Female	75 to 79	26% (11-44%)	74% (56-89%)	1.2 (0.39-2.7)	2.9% (0.4-7.1%)	3.2% (0.45-7.9%)	52 (7.1-130)	740 (99-1800)
Papua New Guinea	2000	Female	80 to 84	25% (11-45%)	75% (55-89%)	1.1 (0.35-2.6)	2.5% (0.4-6.6%)	2.8% (0.42-7.2%)	34 (5.1-89)	360 (56-940)
Papua New Guinea	2000	Female	85 to 89	26% (12-46%)	74% (54-88%)	1.2 (0.41-2.7)	2.4% (0.38-6.3%)	2.7% (0.39-7%)	14 (2-36)	100 (17-270)
Papua New Guinea	2000	Female	90 to 94	29% (13-49%)	71% (51-87%)	1.4 (0.46-3)	2.3% (0.33-5.8%)	2.5% (0.34-6.5%)	3.6 (0.49-9.3)	19 (2.8-49)
Papua New Guinea	2000	Female	95 plus	34% (15-58%)	66% (42-85%)	1.6 (0.47-3.6)	2.5% (0.05-6.7%)	2.7% (-0.011-7.6%)	0.89 (-0.0036-2.4)	3.2 (0.061-8.3)
Papua New Guinea	2000	Male	All Ages	41% (32-50%)	59% (50-68%)	2.4 (1.7-3.3)	5.1% (3.5-6.7%)	6.6% (4.5-8.8%)	1800 (1200-2500)	84000 (61000-110000)
Papua New Guinea	2000	Male	15 to 19	38% (22-56%)	62% (44-78%)	1.6 (0.69-2.9)	4.3% (1.9-7.8%)	5.7% (2.3-11%)	39 (16-76)	3100 (1400-5600)
Papua New Guinea	2000	Male	20 to 24	56% (37-73%)	44% (27-63%)	4.2 (2.3-6.6)	12% (7.7-18%)	16% (9.3-22%)	160 (95-240)	11000 (6700-16000)
Papua New Guinea	2000	Male	25 to 29	63% (43-79%)	37% (21-57%)	5.8 (3.3-8.9)	14% (9.5-19%)	17% (12-23%)	200 (130-280)	13000 (8700-17000)
Papua New Guinea	2000	Male	30 to 34	64% (43-80%)	36% (20-57%)	6.1 (3.4-9.7)	14% (9.3-18%)	16% (11-22%)	180 (120-250)	11000 (7100-15000)
Papua New Guinea	2000	Male	35 to 39	63% (45-79%)	37% (21-55%)	5.4 (2.9-9.4)	13% (8.4-17%)	15% (10-20%)	180 (110-260)	9700 (6300-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	2000	Male	40 to 44	61% (43-77%)	39% (23-57%)	4.8 (2.3-8.4)	12% (7.1-16%)	14% (8.1-19%)	190 (110-270)	9000 (5400-13000)
Papua New Guinea	2000	Male	45 to 49	57% (35-74%)	43% (26-65%)	4.1 (1.9-7.5)	10% (6.2-14%)	12% (6.9-17%)	190 (110-290)	8200 (4800-12000)
Papua New Guinea	2000	Male	50 to 54	54% (34-72%)	46% (28-66%)	3.5 (1.6-6.3)	8.9% (4.7-13%)	10% (5.1-15%)	180 (87-270)	6600 (3400-10000)
Papua New Guinea	2000	Male	55 to 59	52% (34-70%)	48% (30-66%)	2.8 (1.3-5.4)	6.9% (3.2-11%)	7.7% (3.5-12%)	140 (62-230)	4700 (2100-7500)
Papua New Guinea	2000	Male	60 to 64	49% (30-69%)	51% (31-70%)	2.4 (1.1-4.6)	5.7% (2.4-9.5%)	6.3% (2.5-11%)	120 (49-210)	3400 (1400-5700)
Papua New Guinea	2000	Male	65 to 69	47% (28-66%)	53% (34-72%)	2.1 (0.98-4)	5.3% (2.3-8.9%)	5.8% (2.4-9.8%)	100 (41-180)	2400 (980-4000)
Papua New Guinea	2000	Male	70 to 74	45% (27-63%)	55% (37-73%)	1.9 (0.84-3.6)	4.4% (1.5-7.8%)	4.8% (1.6-8.6%)	71 (25-130)	1300 (470-2400)
Papua New Guinea	2000	Male	75 to 79	41% (24-60%)	59% (40-76%)	1.6 (0.69-3)	3.6% (1.2-6.7%)	3.9% (1.2-7.3%)	40 (13-75)	580 (190-1100)
Papua New Guinea	2000	Male	80 to 84	38% (21-57%)	62% (43-79%)	1.4 (0.63-2.7)	3.3% (1.2-5.8%)	3.5% (1.2-6.3%)	20 (7-35)	210 (80-380)
Papua New Guinea	2000	Male	85 to 89	37% (21-54%)	63% (46-79%)	1.4 (0.62-2.7)	2.9% (0.97-5.4%)	3.1% (0.98-6%)	7.4 (2.4-15)	58 (20-110)
Papua New Guinea	2000	Male	90 to 94	38% (23-57%)	62% (43-77%)	1.4 (0.61-2.8)	2.3% (0.72-4.7%)	2.5% (0.67-5.2%)	1.8 (0.49-3.8)	9.9 (3-20)
Papua New Guinea	2000	Male	95 plus	42% (23-63%)	58% (37-77%)	1.5 (0.55-3.3)	2.4% (0.44-5.2%)	2.4% (0.3-5.6%)	0.4 (0.048-0.93)	1.5 (0.29-3.3)
Papua New Guinea	2005	Female	All Ages	20% (14-27%)	80% (73-86%)	0.97 (0.63-1.4)	2.1% (1.1-3.3%)	2.9% (1.3-5%)	850 (370-1500)	34000 (18000-53000)
Papua New Guinea	2005	Female	15 to 19	21% (9.7-37%)	79% (63-90%)	1.1 (0.45-2.3)	2.1% (0.86-4.2%)	3.1% (1.1-6.4%)	16 (5.4-33)	1300 (540-2600)
Papua New Guinea	2005	Female	20 to 24	25% (11-45%)	75% (55-89%)	1.8 (0.72-3.5)	3.5% (1.7-5.9%)	4.6% (2-8.3%)	34 (14-62)	2600 (1300-4500)
Papua New Guinea	2005	Female	25 to 29	27% (11-47%)	73% (53-89%)	1.8 (0.77-3.4)	4.1% (2.1-6.7%)	5.2% (2.4-8.9%)	51 (23-88)	3500 (1700-5800)
Papua New Guinea	2005	Female	30 to 34	28% (11-48%)	72% (52-89%)	1.9 (0.71-3.6)	4% (2-6.6%)	5% (2.3-8.4%)	58 (27-100)	3600 (1800-6200)
Papua New Guinea	2005	Female	35 to 39	28% (12-47%)	72% (53-88%)	2 (0.79-3.6)	4.6% (2.2-7.6%)	5.7% (2.6-9.5%)	73 (32-130)	4000 (1800-6900)
Papua New Guinea	2005	Female	40 to 44	27% (14-46%)	73% (54-86%)	1.8 (0.72-3.5)	4.7% (2-8.4%)	5.9% (2.5-11%)	80 (33-150)	3900 (1600-7100)
Papua New Guinea	2005	Female	45 to 49	27% (12-46%)	73% (54-88%)	1.5 (0.59-3)	4.3% (1.6-8.4%)	5.3% (1.9-10%)	82 (29-160)	3500 (1300-6900)
Papua New Guinea	2005	Female	50 to 54	27% (12-45%)	73% (55-88%)	1.4 (0.52-2.7)	3.8% (1-8.1%)	4.6% (1.2-9.8%)	85 (22-190)	3200 (810-7000)
Papua New Guinea	2005	Female	55 to 59	26% (12-46%)	74% (54-88%)	1.2 (0.46-2.7)	3.2% (0.58-7.5%)	3.8% (0.74-8.8%)	75 (14-180)	2400 (440-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	2005	Female	60 to 64	26% (12-46%)	74% (54-88%)	1.2 (0.43-2.7)	2.6% (0.11-6.6%)	3% (0.14-7.5%)	61 (2.7-150)	1700 (65-4300)
Papua New Guinea	2005	Female	65 to 69	27% (13-46%)	73% (54-87%)	1.2 (0.43-2.5)	2.8% (0.22-6.8%)	3.2% (0.32-7.7%)	70 (6.3-170)	1600 (130-3900)
Papua New Guinea	2005	Female	70 to 74	27% (13-44%)	73% (56-87%)	1.1 (0.41-2.3)	2.4% (-0.053-6.5%)	2.7% (-0.018-7.2%)	60 (-0.45-160)	1100 (-25-3000)
Papua New Guinea	2005	Female	75 to 79	26% (11-44%)	74% (56-89%)	1.1 (0.34-2.4)	2.4% (0.12-6.3%)	2.7% (0.15-7%)	51 (2.7-140)	720 (34-2000)
Papua New Guinea	2005	Female	80 to 84	25% (11-44%)	75% (56-89%)	1 (0.33-2.3)	2.1% (0.17-5.8%)	2.3% (0.17-6.4%)	34 (2.7-99)	360 (29-1000)
Papua New Guinea	2005	Female	85 to 89	26% (12-45%)	74% (55-88%)	1.1 (0.39-2.4)	2% (0.078-5.7%)	2.2% (0.051-6.4%)	14 (0.32-40)	110 (4.2-300)
Papua New Guinea	2005	Female	90 to 94	30% (14-49%)	70% (51-86%)	1.2 (0.43-2.8)	1.9% (0.07-5.3%)	2.1% (0.03-6%)	3.7 (0.053-11)	20 (0.73-56)
Papua New Guinea	2005	Female	95 plus	34% (15-57%)	66% (43-85%)	1.4 (0.44-3.2)	2.1% (-0.22-6.2%)	2.2% (-0.32-6.9%)	0.89 (-0.13-2.8)	3.2 (-0.35-9.5)
Papua New Guinea	2005	Male	All Ages	41% (33-50%)	59% (50-67%)	2.2 (1.5-2.9)	5% (3.3-6.7%)	6.3% (4-8.6%)	2000 (1200-2700)	91000 (62000-120000)
Papua New Guinea	2005	Male	15 to 19	38% (22-56%)	62% (44-78%)	1.4 (0.59-2.5)	3.9% (1.8-7.1%)	5.1% (2.2-9.6%)	38 (15-75)	3000 (1300-5600)
Papua New Guinea	2005	Male	20 to 24	56% (37-73%)	44% (27-63%)	3.7 (2-5.9)	12% (6.9-17%)	14% (8.3-21%)	160 (91-240)	11000 (6600-16000)
Papua New Guinea	2005	Male	25 to 29	63% (44-79%)	37% (21-56%)	5.1 (2.9-7.7)	13% (9-18%)	16% (11-22%)	210 (130-290)	13000 (8700-18000)
Papua New Guinea	2005	Male	30 to 34	64% (44-80%)	36% (20-56%)	5.4 (3.1-8.5)	13% (8.5-17%)	15% (9.9-21%)	220 (140-320)	13000 (8700-18000)
Papua New Guinea	2005	Male	35 to 39	63% (45-79%)	37% (21-55%)	4.8 (2.6-8.1)	12% (7.5-16%)	14% (8.8-19%)	210 (120-310)	11000 (6800-16000)
Papua New Guinea	2005	Male	40 to 44	60% (42-77%)	40% (23-58%)	4.2 (2-7.3)	11% (6.1-15%)	12% (6.8-18%)	210 (110-310)	9900 (5400-15000)
Papua New Guinea	2005	Male	45 to 49	56% (35-74%)	44% (26-65%)	3.6 (1.7-6.5)	9.4% (5.2-14%)	11% (5.8-16%)	200 (100-310)	8700 (4600-13000)
Papua New Guinea	2005	Male	50 to 54	53% (34-73%)	47% (27-66%)	3 (1.4-5.6)	8% (3.8-12%)	9% (4.1-14%)	190 (85-310)	7300 (3300-11000)
Papua New Guinea	2005	Male	55 to 59	51% (33-71%)	49% (29-67%)	2.4 (1.1-4.6)	6.1% (2.6-10%)	6.8% (2.7-11%)	150 (57-250)	4800 (2000-8200)
Papua New Guinea	2005	Male	60 to 64	49% (30-69%)	51% (31-70%)	2.1 (0.93-4.1)	5% (1.9-9%)	5.5% (2-9.9%)	120 (43-220)	3400 (1300-6100)
Papua New Guinea	2005	Male	65 to 69	47% (28-66%)	53% (34-72%)	1.9 (0.87-3.5)	4.6% (1.7-8.2%)	5.1% (1.9-9.1%)	100 (37-190)	2400 (880-4400)
Papua New Guinea	2005	Male	70 to 74	44% (27-62%)	56% (38-73%)	1.6 (0.73-3.1)	3.8% (1.2-7.3%)	4.1% (1.3-8%)	70 (22-140)	1300 (420-2500)
Papua New Guinea	2005	Male	75 to 79	40% (25-60%)	60% (40-75%)	1.4 (0.59-2.6)	3.1% (0.92-6.1%)	3.3% (0.94-6.6%)	40 (11-81)	590 (170-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	2005	Male	80 to 84	38% (21-56%)	62% (44-79%)	1.2 (0.53-2.3)	2.7% (0.91-5.2%)	2.9% (0.9-5.5%)	20 (6.3-39)	220 (74-420)
Papua New Guinea	2005	Male	85 to 89	36% (22-53%)	64% (47-78%)	1.2 (0.52-2.3)	2.3% (0.7-4.8%)	2.5% (0.69-5.2%)	7.6 (2.1-16)	60 (18-120)
Papua New Guinea	2005	Male	90 to 94	37% (23-55%)	63% (45-77%)	1.2 (0.52-2.4)	1.9% (0.48-4.2%)	2% (0.42-4.6%)	1.8 (0.38-4.2)	10 (2.5-23)
Papua New Guinea	2005	Male	95 plus	41% (22-62%)	59% (38-78%)	1.3 (0.46-2.7)	1.9% (0.22-4.5%)	1.9% (0.047-4.8%)	0.4 (0.0094-0.99)	1.5 (0.18-3.6)
Papua New Guinea	2010	Female	All Ages	20% (14-27%)	80% (73-86%)	0.9 (0.57-1.3)	2% (0.96-3.4%)	2.7% (1-5%)	870 (310-1600)	35000 (16000-57000)
Papua New Guinea	2010	Female	15 to 19	20% (9.7-36%)	80% (64-90%)	1 (0.39-2.1)	2.1% (0.83-4.1%)	3.1% (1-6.5%)	17 (5.2-36)	1400 (550-2800)
Papua New Guinea	2010	Female	20 to 24	25% (11-44%)	75% (56-89%)	1.6 (0.65-3.2)	3.4% (1.7-5.9%)	4.5% (2-8.3%)	33 (13-60)	2600 (1200-4500)
Papua New Guinea	2010	Female	25 to 29	26% (11-46%)	74% (54-89%)	1.7 (0.71-3.2)	3.9% (2-6.6%)	5.1% (2.3-8.7%)	49 (21-88)	3400 (1700-5800)
Papua New Guinea	2010	Female	30 to 34	27% (11-47%)	73% (53-89%)	1.7 (0.65-3.3)	3.8% (1.9-6.4%)	4.8% (2.2-8.4%)	57 (25-100)	3600 (1700-6300)
Papua New Guinea	2010	Female	35 to 39	28% (12-47%)	72% (53-88%)	1.8 (0.73-3.3)	4.4% (2.1-7.4%)	5.5% (2.5-9.5%)	84 (35-150)	4600 (2000-8200)
Papua New Guinea	2010	Female	40 to 44	27% (13-46%)	73% (54-87%)	1.7 (0.69-3.3)	4.4% (1.9-8.2%)	5.6% (2.3-10%)	86 (33-170)	4200 (1600-7900)
Papua New Guinea	2010	Female	45 to 49	27% (12-45%)	73% (55-88%)	1.4 (0.54-2.9)	3.9% (1.2-8%)	4.9% (1.5-10%)	83 (26-170)	3600 (1100-7500)
Papua New Guinea	2010	Female	50 to 54	27% (12-47%)	73% (53-88%)	1.3 (0.49-2.6)	3.4% (0.72-7.8%)	4.2% (0.95-9.3%)	85 (19-200)	3200 (680-7400)
Papua New Guinea	2010	Female	55 to 59	26% (12-47%)	74% (53-88%)	1.2 (0.43-2.6)	2.9% (0.31-7%)	3.4% (0.42-8.2%)	80 (9.1-200)	2600 (270-6600)
Papua New Guinea	2010	Female	60 to 64	26% (12-46%)	74% (54-88%)	1.1 (0.38-2.5)	2.2% (-0.21-6.1%)	2.6% (-0.19-7.1%)	62 (-4.5-170)	1700 (-160-4700)
Papua New Guinea	2010	Female	65 to 69	27% (13-46%)	73% (54-87%)	1.1 (0.41-2.3)	2.3% (-0.074-6.3%)	2.7% (-0.063-7.2%)	68 (-1.4-180)	1500 (-45-4100)
Papua New Guinea	2010	Female	70 to 74	27% (12-45%)	73% (55-88%)	1.1 (0.36-2.3)	2.1% (-0.28-6%)	2.4% (-0.26-6.6%)	61 (-6.9-170)	1100 (-150-3200)
Papua New Guinea	2010	Female	75 to 79	26% (11-45%)	74% (55-89%)	0.99 (0.32-2.3)	2.1% (-0.13-6.2%)	2.3% (-0.12-6.7%)	51 (-2.5-150)	730 (-43-2200)
Papua New Guinea	2010	Female	80 to 84	25% (11-45%)	75% (55-89%)	0.95 (0.3-2.1)	1.8% (-0.023-5.3%)	2% (-0.037-6%)	35 (-0.7-100)	380 (-4.7-1100)
Papua New Guinea	2010	Female	85 to 89	26% (12-45%)	74% (55-88%)	0.99 (0.35-2.2)	1.7% (-0.15-5.6%)	1.9% (-0.21-6.1%)	14 (-1.5-46)	110 (-10-340)
Papua New Guinea	2010	Female	90 to 94	29% (14-49%)	71% (51-86%)	1.1 (0.38-2.4)	1.6% (-0.16-4.9%)	1.8% (-0.23-5.4%)	3.8 (-0.49-12)	20 (-2.1-60)
Papua New Guinea	2010	Female	95 plus	34% (15-56%)	66% (44-85%)	1.3 (0.4-2.9)	1.7% (-0.52-6%)	1.8% (-0.7-6.6%)	0.88 (-0.34-3.3)	3.2 (-0.98-11)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	2010	Male	All Ages	41% (33-51%)	59% (49-67%)	2 (1.4-2.8)	5% (3.2-7%)	6.2% (3.7-8.7%)	2100 (1200-3000)	97000 (62000-130000)
Papua New Guinea	2010	Male	15 to 19	38% (23-56%)	62% (44-77%)	1.3 (0.55-2.4)	3.8% (1.8-7%)	5.1% (2.2-9.6%)	41 (17-81)	3200 (1400-6100)
Papua New Guinea	2010	Male	20 to 24	56% (38-73%)	44% (27-62%)	3.4 (1.8-5.6)	11% (6.6-16%)	14% (7.8-21%)	160 (88-240)	11000 (6500-17000)
Papua New Guinea	2010	Male	25 to 29	63% (43-79%)	37% (21-57%)	4.7 (2.7-7.3)	13% (8.4-18%)	16% (10-22%)	210 (130-300)	14000 (8500-19000)
Papua New Guinea	2010	Male	30 to 34	64% (43-80%)	36% (20-57%)	5 (2.7-7.9)	12% (8.1-17%)	15% (9.4-21%)	230 (140-330)	14000 (8700-19000)
Papua New Guinea	2010	Male	35 to 39	63% (44-79%)	37% (21-56%)	4.4 (2.3-7.4)	11% (6.8-16%)	14% (7.9-19%)	250 (130-380)	13000 (7600-20000)
Papua New Guinea	2010	Male	40 to 44	61% (43-77%)	39% (23-57%)	3.9 (1.9-6.9)	10% (5.5-15%)	12% (6.2-18%)	230 (120-360)	11000 (5800-17000)
Papua New Guinea	2010	Male	45 to 49	57% (35-75%)	43% (25-65%)	3.4 (1.6-6.1)	8.9% (4.5-14%)	10% (5.1-16%)	220 (110-350)	9300 (4700-15000)
Papua New Guinea	2010	Male	50 to 54	54% (34-72%)	46% (28-66%)	2.8 (1.3-5.1)	7.5% (3.4-12%)	8.5% (3.7-14%)	200 (85-340)	7600 (3400-13000)
Papua New Guinea	2010	Male	55 to 59	52% (33-71%)	48% (29-67%)	2.3 (1.1-4.3)	5.7% (2.3-9.7%)	6.3% (2.5-11%)	160 (60-290)	5300 (2100-9500)
Papua New Guinea	2010	Male	60 to 64	49% (29-68%)	51% (32-71%)	1.9 (0.89-3.7)	4.6% (1.6-8.6%)	5% (1.6-9.6%)	130 (40-240)	3500 (1200-6700)
Papua New Guinea	2010	Male	65 to 69	47% (28-66%)	53% (34-72%)	1.7 (0.78-3.2)	4.2% (1.3-7.7%)	4.5% (1.4-8.6%)	110 (31-210)	2500 (770-4800)
Papua New Guinea	2010	Male	70 to 74	44% (27-62%)	56% (38-73%)	1.5 (0.67-2.9)	3.4% (0.95-6.9%)	3.6% (0.98-7.6%)	73 (20-150)	1400 (400-2800)
Papua New Guinea	2010	Male	75 to 79	41% (25-59%)	59% (41-75%)	1.3 (0.56-2.5)	2.7% (0.73-5.6%)	2.9% (0.74-6.1%)	42 (10-89)	620 (160-1300)
Papua New Guinea	2010	Male	80 to 84	38% (22-57%)	62% (43-78%)	1.2 (0.51-2.2)	2.4% (0.72-4.9%)	2.6% (0.72-5.4%)	22 (5.9-46)	240 (70-500)
Papua New Guinea	2010	Male	85 to 89	36% (22-53%)	64% (47-78%)	1.1 (0.46-2.2)	2% (0.53-4.5%)	2.2% (0.5-4.9%)	8.2 (1.9-19)	65 (16-140)
Papua New Guinea	2010	Male	90 to 94	38% (23-56%)	62% (44-77%)	1.1 (0.46-2.2)	1.7% (0.34-3.8%)	1.7% (0.25-4.2%)	2 (0.28-4.7)	11 (2.2-26)
Papua New Guinea	2010	Male	95 plus	41% (22-62%)	59% (38-78%)	1.2 (0.44-2.5)	1.7% (0.082-4.2%)	1.7% (-0.097-4.5%)	0.42 (-0.025-1.2)	1.7 (0.082-4.2)
Papua New Guinea	2016	Female	All Ages	20% (14-27%)	80% (73-86%)	0.94 (0.54-1.5)	2.1% (0.93-3.7%)	2.8% (0.79-5.4%)	940 (250-1900)	37000 (15000-65000)
Papua New Guinea	2016	Female	15 to 19	21% (9.6-36%)	79% (64-90%)	1.1 (0.38-2.2)	2.1% (0.84-4.2%)	3.3% (1.1-6.9%)	18 (5.8-38)	1600 (620-3100)
Papua New Guinea	2016	Female	20 to 24	25% (11-44%)	75% (56-89%)	1.7 (0.66-3.3)	3.4% (1.7-6.1%)	4.8% (2.1-8.9%)	34 (15-64)	2800 (1400-4800)
Papua New Guinea	2016	Female	25 to 29	27% (11-46%)	73% (54-89%)	1.7 (0.69-3.6)	4% (2-6.8%)	5.4% (2.4-9.7%)	47 (20-86)	3400 (1600-5900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	2016	Female	30 to 34	27% (10-48%)	73% (52-90%)	1.8 (0.61-3.6)	3.8% (1.7-6.7%)	5% (2.1-9%)	55 (21-100)	3500 (1600-6400)
Papua New Guinea	2016	Female	35 to 39	28% (11-47%)	72% (53-89%)	1.8 (0.69-3.8)	4.3% (1.9-7.5%)	5.5% (2.3-9.7%)	81 (32-150)	4500 (1900-8200)
Papua New Guinea	2016	Female	40 to 44	28% (13-46%)	72% (54-87%)	1.7 (0.66-3.8)	4.4% (1.6-8.1%)	5.6% (2.1-10%)	100 (36-200)	4900 (1800-9400)
Papua New Guinea	2016	Female	45 to 49	27% (12-45%)	73% (55-88%)	1.4 (0.53-2.9)	3.8% (1.1-7.7%)	4.9% (1.4-9.8%)	94 (26-190)	4100 (1100-8500)
Papua New Guinea	2016	Female	50 to 54	27% (13-46%)	73% (54-87%)	1.3 (0.45-2.6)	3.3% (0.46-7.6%)	4.1% (0.67-9.3%)	89 (15-210)	3400 (520-7800)
Papua New Guinea	2016	Female	55 to 59	26% (12-45%)	74% (55-88%)	1.1 (0.41-2.5)	2.7% (0.29-7%)	3.3% (0.42-8.2%)	83 (11-210)	2700 (280-6900)
Papua New Guinea	2016	Female	60 to 64	26% (12-46%)	74% (54-88%)	1.1 (0.36-2.6)	2.1% (-0.29-6.2%)	2.5% (-0.3-7.1%)	71 (-8.1-210)	2000 (-280-5600)
Papua New Guinea	2016	Female	65 to 69	27% (12-47%)	73% (53-88%)	1.1 (0.38-2.4)	2.2% (-0.19-6.4%)	2.6% (-0.17-7.3%)	77 (-4.9-210)	1700 (-150-5000)
Papua New Guinea	2016	Female	70 to 74	27% (13-46%)	73% (54-87%)	1.1 (0.37-2.4)	2% (-0.43-6.6%)	2.3% (-0.42-7.4%)	68 (-12-220)	1200 (-250-4000)
Papua New Guinea	2016	Female	75 to 79	26% (11-45%)	74% (55-89%)	1 (0.31-2.4)	2% (-0.32-6.2%)	2.3% (-0.34-6.9%)	59 (-9.1-180)	840 (-130-2600)
Papua New Guinea	2016	Female	80 to 84	25% (11-45%)	75% (55-89%)	0.97 (0.29-2.3)	1.8% (-0.26-5.9%)	2% (-0.27-6.6%)	43 (-5.8-140)	460 (-68-1500)
Papua New Guinea	2016	Female	85 to 89	26% (12-46%)	74% (54-88%)	1 (0.35-2.5)	1.8% (-0.32-5.8%)	2% (-0.4-6.4%)	18 (-3.6-60)	130 (-24-450)
Papua New Guinea	2016	Female	90 to 94	30% (13-49%)	70% (51-87%)	1.2 (0.36-2.7)	1.7% (-0.28-5.5%)	1.9% (-0.34-6.1%)	4.8 (-0.88-15)	26 (-4.1-82)
Papua New Guinea	2016	Female	95 plus	34% (14-57%)	66% (43-86%)	1.3 (0.4-3.2)	1.7% (-0.69-6.3%)	1.8% (-0.86-6.9%)	1.1 (-0.51-3.9)	3.8 (-1.5-14)
Papua New Guinea	2016	Male	All Ages	42% (34-51%)	58% (49-66%)	2.1 (1.3-3.2)	5.3% (3.2-7.5%)	6.3% (3.5-9.2%)	2300 (1200-3300)	100000 (62000-150000)
Papua New Guinea	2016	Male	15 to 19	38% (22-56%)	62% (44-78%)	1.3 (0.52-2.6)	3.9% (1.8-7.6%)	5.3% (2.2-10%)	47 (20-97)	3700 (1700-7300)
Papua New Guinea	2016	Male	20 to 24	56% (37-73%)	44% (27-63%)	3.5 (1.7-6)	11% (6.4-16%)	14% (7.6-21%)	180 (92-270)	12000 (6800-19000)
Papua New Guinea	2016	Male	25 to 29	63% (43-79%)	37% (21-57%)	4.8 (2.5-8.2)	13% (8.1-18%)	16% (10-23%)	210 (120-300)	14000 (8400-20000)
Papua New Guinea	2016	Male	30 to 34	64% (43-81%)	36% (19-57%)	5.1 (2.7-8.7)	12% (7.8-17%)	15% (9.2-21%)	230 (130-320)	14000 (8200-19000)
Papua New Guinea	2016	Male	35 to 39	63% (45-80%)	37% (20-55%)	4.5 (2.2-8.2)	11% (6.5-16%)	14% (7.7-20%)	250 (130-370)	14000 (7400-20000)
Papua New Guinea	2016	Male	40 to 44	60% (42-77%)	40% (23-58%)	4 (1.8-7.5)	10% (5.1-15%)	12% (5.6-17%)	270 (130-420)	13000 (6600-20000)
Papua New Guinea	2016	Male	45 to 49	56% (35-74%)	44% (26-65%)	3.4 (1.6-6.3)	8.7% (4.2-13%)	10% (4.8-15%)	250 (110-400)	11000 (5100-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Papua New Guinea	2016	Male	50 to 54	53% (33-73%)	47% (27-67%)	2.8 (1.2-5.3)	7.4% (3.1-12%)	8.4% (3.4-14%)	220 (88-370)	8400 (3600-14000)
Papua New Guinea	2016	Male	55 to 59	51% (33-71%)	49% (29-67%)	2.3 (1-4.5)	5.6% (2-9.7%)	6.2% (2.2-11%)	170 (61-330)	5800 (2100-11000)
Papua New Guinea	2016	Male	60 to 64	49% (29-69%)	51% (31-71%)	1.9 (0.8-3.8)	4.4% (1.4-8.4%)	4.9% (1.4-9.3%)	140 (40-280)	4000 (1200-7700)
Papua New Guinea	2016	Male	65 to 69	47% (29-66%)	53% (34-71%)	1.7 (0.73-3.5)	4% (1.2-7.7%)	4.4% (1.2-8.6%)	120 (32-230)	2800 (790-5400)
Papua New Guinea	2016	Male	70 to 74	45% (27-62%)	55% (38-73%)	1.5 (0.63-3.1)	3.3% (0.87-6.7%)	3.6% (0.9-7.3%)	82 (20-170)	1500 (390-3200)
Papua New Guinea	2016	Male	75 to 79	40% (24-59%)	60% (41-76%)	1.3 (0.53-2.7)	2.7% (0.61-5.5%)	2.9% (0.61-6%)	50 (10-110)	740 (170-1500)
Papua New Guinea	2016	Male	80 to 84	37% (21-56%)	63% (44-79%)	1.2 (0.49-2.3)	2.4% (0.62-5.2%)	2.6% (0.61-5.6%)	27 (6.4-59)	300 (75-640)
Papua New Guinea	2016	Male	85 to 89	36% (21-52%)	64% (48-79%)	1.1 (0.46-2.4)	2.1% (0.43-4.7%)	2.2% (0.39-5.1%)	10 (1.9-24)	82 (17-190)
Papua New Guinea	2016	Male	90 to 94	38% (23-55%)	62% (45-77%)	1.2 (0.46-2.4)	1.7% (0.3-4.1%)	1.8% (0.19-4.4%)	2.6 (0.27-6.4)	15 (2.5-34)
Papua New Guinea	2016	Male	95 plus	42% (22-62%)	58% (38-78%)	1.3 (0.43-2.8)	1.8% (0.1-4.6%)	1.8% (-0.14-5%)	0.57 (-0.047-1.6)	2.2 (0.13-5.8)
Samoa	1990	Female	All Ages	18% (12-24%)	82% (76-88%)	0.42 (0.31-0.53)	0.8% (0.39-1.3%)	0.84% (0.039-1.9%)	3.1 (0.13-7)	160 (79-260)
Samoa	1990	Female	15 to 19	19% (7.9-34%)	81% (66-92%)	0.51 (0.22-0.96)	1.2% (0.62-2.2%)	2.2% (0.81-4.4%)	0.13 (0.049-0.27)	15 (7.3-26)
Samoa	1990	Female	20 to 24	23% (9.3-41%)	77% (59-91%)	0.81 (0.35-1.4)	2.1% (1.3-3.4%)	3.2% (1.3-6%)	0.2 (0.079-0.37)	22 (12-35)
Samoa	1990	Female	25 to 29	24% (10-43%)	76% (57-90%)	0.83 (0.37-1.5)	2.2% (1.3-3.3%)	3.2% (1.5-5.5%)	0.22 (0.1-0.4)	22 (13-35)
Samoa	1990	Female	30 to 34	25% (9.9-45%)	75% (55-90%)	0.87 (0.35-1.5)	2% (1.2-3.3%)	3% (1.4-5.3%)	0.2 (0.088-0.37)	17 (9.3-27)
Samoa	1990	Female	35 to 39	26% (11-44%)	74% (56-89%)	0.9 (0.38-1.6)	2.1% (1.1-3.5%)	3.1% (1.4-5.4%)	0.3 (0.13-0.55)	20 (10-33)
Samoa	1990	Female	40 to 44	25% (12-43%)	75% (57-88%)	0.83 (0.35-1.5)	2% (0.85-3.6%)	3% (1.2-5.7%)	0.3 (0.11-0.58)	16 (6.7-31)
Samoa	1990	Female	45 to 49	25% (11-43%)	75% (57-89%)	0.72 (0.28-1.4)	1.4% (0.39-3.1%)	2.1% (0.64-4.7%)	0.33 (0.093-0.73)	15 (3.9-33)
Samoa	1990	Female	50 to 54	25% (11-44%)	75% (56-89%)	0.63 (0.24-1.3)	1.1% (-0.0077-3%)	1.6% (0.15-4.3%)	0.28 (0.027-0.73)	10 (-0.07-28)
Samoa	1990	Female	55 to 59	24% (11-42%)	76% (58-89%)	0.56 (0.21-1.1)	0.8% (-0.27-2.3%)	1.3% (-0.13-3.3%)	0.32 (-0.035-0.86)	9.6 (-3.1-28)
Samoa	1990	Female	60 to 64	24% (11-44%)	76% (56-89%)	0.54 (0.2-1.1)	0.42% (-0.74-1.8%)	0.7% (-0.67-2.5%)	0.2 (-0.2-0.74)	4.6 (-8.1-20)
Samoa	1990	Female	65 to 69	25% (11-42%)	75% (58-89%)	0.54 (0.21-1.1)	0.38% (-0.78-1.7%)	0.65% (-0.73-2.4%)	0.23 (-0.28-0.86)	4.2 (-8.5-19)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	1990	Female	70 to 74	24% (11-40%)	76% (60-89%)	0.52 (0.19-1)	0.23% (-0.86-1.8%)	0.39% (-0.9-2.4%)	0.13 (-0.31-0.79)	1.8 (-6.7-14)
Samoa	1990	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.48 (0.16-1)	0.27% (-0.8-1.8%)	0.4% (-0.87-2.2%)	0.15 (-0.35-0.85)	1.8 (-5.6-12)
Samoa	1990	Female	80 to 84	23% (9.3-42%)	77% (58-91%)	0.46 (0.16-0.99)	0.12% (-0.81-1.4%)	0.19% (-0.89-1.7%)	0.06 (-0.29-0.53)	0.5 (-3.3-5.6)
Samoa	1990	Female	85 to 89	24% (11-42%)	76% (58-89%)	0.48 (0.18-1)	0.03% (-0.87-1.2%)	0.049% (-1-1.4%)	0.0098 (-0.2-0.27)	0.054 (-1.5-2)
Samoa	1990	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.56 (0.2-1.2)	0.088% (-0.82-1.5%)	0.11% (-0.94-1.7%)	0.011 (-0.095-0.17)	0.054 (-0.51-0.87)
Samoa	1990	Female	95 plus	32% (13-55%)	68% (45-87%)	0.65 (0.21-1.5)	0.039% (-1.3-1.8%)	-0.0092% (-1.5-2%)	-0.00024 (-0.048-0.062)	0.0047 (-0.15-0.22)
Samoa	1990	Male	All Ages	36% (28-46%)	64% (54-72%)	0.91 (0.76-1.1)	3.1% (2-4.3%)	3.3% (2-4.9%)	17 (11-26)	860 (590-1200)
Samoa	1990	Male	15 to 19	35% (20-53%)	65% (47-80%)	0.64 (0.32-1.1)	2.4% (1.2-4.2%)	3.3% (1.3-6.2%)	0.55 (0.22-1)	47 (23-82)
Samoa	1990	Male	20 to 24	53% (35-71%)	47% (29-65%)	1.7 (1.1-2.5)	7% (4.2-11%)	9.1% (5.1-14%)	1.9 (1-3.1)	140 (82-220)
Samoa	1990	Male	25 to 29	61% (41-77%)	39% (23-59%)	2.4 (1.5-3.4)	8.9% (5.6-13%)	12% (7-17%)	2.3 (1.3-3.5)	160 (98-230)
Samoa	1990	Male	30 to 34	62% (41-79%)	38% (21-59%)	2.6 (1.6-3.8)	8.3% (5.1-12%)	11% (6.2-16%)	1.6 (0.9-2.6)	100 (61-160)
Samoa	1990	Male	35 to 39	60% (42-77%)	40% (23-58%)	2.3 (1.3-3.5)	7% (4-11%)	8.7% (4.7-14%)	1.6 (0.85-2.6)	91 (52-150)
Samoa	1990	Male	40 to 44	57% (39-74%)	43% (26-61%)	2 (1-3.2)	5.7% (2.9-9.5%)	6.9% (3.3-12%)	1.1 (0.51-2)	59 (28-100)
Samoa	1990	Male	45 to 49	53% (32-72%)	47% (28-68%)	1.7 (0.83-3)	4.8% (2.2-8.4%)	5.6% (2.4-10%)	1.4 (0.58-2.5)	63 (28-110)
Samoa	1990	Male	50 to 54	51% (31-69%)	49% (31-69%)	1.4 (0.71-2.4)	4% (1.6-7.4%)	4.6% (1.7-8.9%)	1.5 (0.52-2.9)	59 (23-110)
Samoa	1990	Male	55 to 59	48% (30-69%)	52% (31-70%)	1.1 (0.57-2)	2.9% (0.94-6.1%)	3.3% (0.97-7.1%)	1.3 (0.36-2.8)	44 (14-94)
Samoa	1990	Male	60 to 64	46% (27-66%)	54% (34-73%)	0.95 (0.45-1.7)	2.3% (0.58-5.3%)	2.6% (0.6-6.1%)	1.1 (0.26-2.7)	33 (8.6-78)
Samoa	1990	Male	65 to 69	44% (26-64%)	56% (36-74%)	0.87 (0.43-1.5)	2% (0.48-4.1%)	2.2% (0.53-4.8%)	1.3 (0.31-2.9)	31 (8-68)
Samoa	1990	Male	70 to 74	42% (25-59%)	58% (41-75%)	0.76 (0.37-1.4)	1.5% (0.21-3.6%)	1.7% (0.18-4%)	0.85 (0.092-2.1)	16 (2.1-39)
Samoa	1990	Male	75 to 79	38% (22-57%)	62% (43-78%)	0.64 (0.31-1.2)	1.2% (0.11-3%)	1.3% (0.056-3.4%)	0.55 (0.027-1.5)	8.2 (0.73-22)
Samoa	1990	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.57 (0.28-1)	0.97% (0.13-2.2%)	1.1% (0.11-2.5%)	0.28 (0.031-0.67)	3.1 (0.44-7.2)
Samoa	1990	Male	85 to 89	34% (20-50%)	66% (50-80%)	0.54 (0.26-0.99)	0.74% (-0.0085-1.8%)	0.8% (-0.062-2.1%)	0.1 (-0.0079-0.28)	0.84 (-0.0096-2.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	1990	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.55 (0.25-1.1)	0.55% (-0.067-1.5%)	0.56% (-0.15-1.7%)	0.026 (-0.0066-0.078)	0.15 (-0.018-0.43)
Samoa	1990	Male	95 plus	38% (21-59%)	62% (41-79%)	0.61 (0.24-1.2)	0.36% (-0.47-1.6%)	0.28% (-0.68-1.7%)	0.0025 (-0.0061-0.016)	0.013 (-0.016-0.057)
Samoa	1995	Female	All Ages	18% (12-24%)	82% (76-88%)	0.43 (0.32-0.56)	0.85% (0.4-1.4%)	0.84% (-0.022-2%)	3.3 (-0.087-7.8)	170 (81-290)
Samoa	1995	Female	15 to 19	19% (8.5-33%)	81% (67-91%)	0.53 (0.23-0.98)	1.3% (0.65-2.2%)	2.3% (0.85-4.6%)	0.12 (0.045-0.24)	13 (6.6-23)
Samoa	1995	Female	20 to 24	23% (9.4-39%)	77% (61-91%)	0.83 (0.36-1.5)	2.2% (1.3-3.5%)	3.3% (1.4-6.2%)	0.2 (0.083-0.39)	22 (13-37)
Samoa	1995	Female	25 to 29	24% (9.7-42%)	76% (58-90%)	0.85 (0.37-1.5)	2.2% (1.4-3.4%)	3.3% (1.6-5.6%)	0.24 (0.11-0.43)	23 (14-37)
Samoa	1995	Female	30 to 34	25% (9.4-44%)	75% (56-91%)	0.89 (0.35-1.6)	2.1% (1.2-3.3%)	3% (1.4-5.3%)	0.28 (0.12-0.49)	22 (13-36)
Samoa	1995	Female	35 to 39	26% (11-44%)	74% (56-89%)	0.93 (0.38-1.6)	2.1% (1.1-3.6%)	3.2% (1.5-5.6%)	0.29 (0.12-0.54)	19 (9.8-33)
Samoa	1995	Female	40 to 44	25% (12-42%)	75% (58-88%)	0.86 (0.37-1.6)	2% (0.87-3.7%)	3% (1.2-5.6%)	0.41 (0.16-0.8)	22 (9.6-42)
Samoa	1995	Female	45 to 49	24% (11-42%)	76% (58-89%)	0.73 (0.29-1.4)	1.4% (0.42-3.1%)	2.2% (0.65-4.6%)	0.31 (0.089-0.69)	14 (3.7-30)
Samoa	1995	Female	50 to 54	24% (11-43%)	76% (57-89%)	0.64 (0.24-1.3)	1% (-0.067-2.8%)	1.6% (0.1-4.1%)	0.35 (0.023-0.94)	13 (-0.73-36)
Samoa	1995	Female	55 to 59	23% (11-41%)	77% (59-89%)	0.57 (0.22-1.1)	0.77% (-0.31-2.3%)	1.2% (-0.16-3.2%)	0.26 (-0.035-0.7)	7.6 (-3.2-23)
Samoa	1995	Female	60 to 64	23% (10-42%)	77% (58-90%)	0.55 (0.2-1.2)	0.4% (-0.73-1.8%)	0.69% (-0.7-2.6%)	0.23 (-0.24-0.89)	5.1 (-8.9-24)
Samoa	1995	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.56 (0.22-1.1)	0.36% (-0.87-1.9%)	0.63% (-0.85-2.6%)	0.24 (-0.32-0.93)	4.2 (-9.8-21)
Samoa	1995	Female	70 to 74	25% (11-42%)	75% (58-89%)	0.54 (0.2-1.1)	0.2% (-0.94-1.7%)	0.36% (-0.99-2.3%)	0.15 (-0.42-0.98)	2 (-9.3-17)
Samoa	1995	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.5 (0.16-1.1)	0.25% (-0.9-1.8%)	0.38% (-0.92-2.3%)	0.16 (-0.38-0.91)	1.8 (-6.3-13)
Samoa	1995	Female	80 to 84	23% (9.7-42%)	77% (58-90%)	0.47 (0.17-0.99)	0.12% (-0.84-1.4%)	0.18% (-0.95-1.6%)	0.072 (-0.37-0.67)	0.6 (-4.3-7.1)
Samoa	1995	Female	85 to 89	24% (11-42%)	76% (58-89%)	0.49 (0.18-1)	0.033% (-0.88-1.2%)	0.051% (-1-1.5%)	0.014 (-0.27-0.4)	0.083 (-2.1-3)
Samoa	1995	Female	90 to 94	27% (13-47%)	73% (53-87%)	0.56 (0.22-1.2)	0.091% (-0.87-1.5%)	0.11% (-1-1.7%)	0.012 (-0.11-0.19)	0.06 (-0.57-0.96)
Samoa	1995	Female	95 plus	32% (14-55%)	68% (45-86%)	0.66 (0.22-1.5)	0.05% (-1.3-1.8%)	-0.0024% (-1.6-2.1%)	-0.000039 (-0.056-0.074)	0.0074 (-0.18-0.26)
Samoa	1995	Male	All Ages	36% (28-46%)	64% (54-72%)	0.98 (0.81-1.2)	3.3% (2.3-4.6%)	3.4% (2-5%)	18 (11-27)	890 (620-1200)
Samoa	1995	Male	15 to 19	35% (20-53%)	65% (47-80%)	0.65 (0.32-1.1)	2.4% (1.2-4.2%)	3.4% (1.4-6.3%)	0.5 (0.21-0.96)	43 (21-76)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	1995	Male	20 to 24	53% (34-70%)	47% (30-66%)	1.8 (1.1-2.7)	7.2% (4.3-11%)	9.4% (5.3-15%)	1.8 (0.91-2.8)	130 (75-210)
Samoa	1995	Male	25 to 29	60% (40-76%)	40% (24-60%)	2.5 (1.6-3.5)	9% (5.9-13%)	12% (7.3-17%)	2.2 (1.3-3.3)	150 (96-220)
Samoa	1995	Male	30 to 34	61% (40-78%)	39% (22-60%)	2.7 (1.7-3.9)	8.2% (5.2-12%)	11% (6.4-16%)	2.1 (1.2-3.4)	140 (83-210)
Samoa	1995	Male	35 to 39	60% (42-77%)	40% (23-58%)	2.3 (1.4-3.7)	7% (4-11%)	8.9% (4.9-14%)	1.5 (0.78-2.4)	86 (48-140)
Samoa	1995	Male	40 to 44	57% (40-75%)	43% (25-60%)	2 (1.1-3.3)	5.7% (3-9.4%)	7% (3.3-12%)	1.6 (0.75-2.8)	83 (41-140)
Samoa	1995	Male	45 to 49	53% (32-71%)	47% (29-68%)	1.7 (0.91-3)	4.8% (2.2-8.7%)	5.7% (2.4-11%)	1.2 (0.51-2.3)	58 (26-100)
Samoa	1995	Male	50 to 54	50% (30-69%)	50% (31-70%)	1.4 (0.74-2.5)	4% (1.6-7.7%)	4.6% (1.7-9.2%)	1.5 (0.53-3)	60 (24-120)
Samoa	1995	Male	55 to 59	48% (30-68%)	52% (32-70%)	1.2 (0.58-2)	2.8% (0.92-5.9%)	3.2% (0.88-6.9%)	1.2 (0.34-2.7)	43 (15-92)
Samoa	1995	Male	60 to 64	46% (27-66%)	54% (34-73%)	0.97 (0.45-1.8)	2.2% (0.51-5.1%)	2.5% (0.52-5.9%)	1.1 (0.24-2.7)	34 (7.9-78)
Samoa	1995	Male	65 to 69	44% (26-63%)	56% (37-74%)	0.88 (0.44-1.5)	1.9% (0.41-4.2%)	2.2% (0.4-4.8%)	1.2 (0.22-2.6)	28 (5.6-61)
Samoa	1995	Male	70 to 74	42% (25-59%)	58% (41-75%)	0.78 (0.38-1.4)	1.5% (0.19-3.6%)	1.7% (0.17-4.1%)	1 (0.1-2.5)	20 (2.5-48)
Samoa	1995	Male	75 to 79	38% (22-57%)	62% (43-78%)	0.67 (0.32-1.3)	1.2% (0.095-3%)	1.3% (0.07-3.3%)	0.6 (0.036-1.6)	9 (0.78-23)
Samoa	1995	Male	80 to 84	35% (20-53%)	65% (47-80%)	0.59 (0.29-1.1)	0.98% (0.14-2.3%)	1.1% (0.11-2.6%)	0.34 (0.033-0.82)	3.7 (0.53-8.9)
Samoa	1995	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.56 (0.27-1)	0.75% (0.002-1.9%)	0.8% (-0.055-2.1%)	0.14 (-0.01-0.38)	1.2 (0.0031-3)
Samoa	1995	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.57 (0.26-1.1)	0.55% (-0.083-1.6%)	0.57% (-0.17-1.7%)	0.033 (-0.01-0.1)	0.2 (-0.03-0.54)
Samoa	1995	Male	95 plus	38% (20-59%)	62% (41-80%)	0.63 (0.25-1.3)	0.37% (-0.49-1.7%)	0.29% (-0.69-1.9%)	0.0041 (-0.0097-0.026)	0.021 (-0.028-0.097)
Samoa	2000	Female	All Ages	17% (12-24%)	83% (76-88%)	0.44 (0.32-0.56)	0.83% (0.37-1.5%)	0.74% (-0.13-1.9%)	3.1 (-0.53-8.1)	170 (70-290)
Samoa	2000	Female	15 to 19	18% (8-34%)	82% (66-92%)	0.54 (0.22-1)	1.2% (0.63-2.3%)	2.3% (0.83-4.9%)	0.11 (0.04-0.24)	12 (6.1-22)
Samoa	2000	Female	20 to 24	22% (9.4-41%)	78% (59-91%)	0.84 (0.35-1.5)	2.2% (1.2-3.5%)	3.3% (1.3-6.3%)	0.19 (0.072-0.36)	21 (11-34)
Samoa	2000	Female	25 to 29	24% (10-43%)	76% (57-90%)	0.87 (0.38-1.5)	2.3% (1.4-3.5%)	3.3% (1.6-5.7%)	0.24 (0.11-0.43)	23 (13-36)
Samoa	2000	Female	30 to 34	25% (9.6-44%)	75% (56-90%)	0.9 (0.35-1.6)	2.1% (1.2-3.4%)	3% (1.3-5.2%)	0.28 (0.12-0.5)	23 (13-37)
Samoa	2000	Female	35 to 39	25% (9.7-44%)	75% (56-90%)	0.93 (0.39-1.6)	2.1% (1.1-3.5%)	3.1% (1.4-5.5%)	0.36 (0.16-0.67)	24 (12-42)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	2000	Female	40 to 44	25% (12-43%)	75% (57-88%)	0.86 (0.36-1.6)	1.9% (0.79-3.5%)	2.8% (1.1-5.3%)	0.4 (0.15-0.81)	22 (9.2-42)
Samoa	2000	Female	45 to 49	24% (10-41%)	76% (59-90%)	0.73 (0.29-1.4)	1.3% (0.28-3%)	2% (0.51-4.6%)	0.33 (0.082-0.74)	15 (3.1-34)
Samoa	2000	Female	50 to 54	24% (10-42%)	76% (58-90%)	0.64 (0.25-1.3)	0.94% (-0.26-2.7%)	1.5% (-0.053-3.8%)	0.28 (-0.01-0.76)	10 (-2.8-29)
Samoa	2000	Female	55 to 59	23% (10-42%)	77% (58-90%)	0.57 (0.22-1.2)	0.66% (-0.45-2.3%)	1.1% (-0.33-3.2%)	0.26 (-0.076-0.76)	7.4 (-5.5-24)
Samoa	2000	Female	60 to 64	23% (10-40%)	77% (60-90%)	0.54 (0.2-1.2)	0.3% (-0.87-1.7%)	0.55% (-0.85-2.4%)	0.16 (-0.25-0.68)	3.3 (-9.6-18)
Samoa	2000	Female	65 to 69	24% (11-41%)	76% (59-89%)	0.56 (0.22-1.1)	0.25% (-1-1.8%)	0.48% (-1-2.4%)	0.17 (-0.38-0.83)	2.8 (-12-19)
Samoa	2000	Female	70 to 74	24% (11-40%)	76% (60-89%)	0.54 (0.2-1.1)	0.1% (-1-1.5%)	0.23% (-1.1-2%)	0.11 (-0.53-0.94)	1.1 (-12-17)
Samoa	2000	Female	75 to 79	23% (10-41%)	77% (59-90%)	0.51 (0.17-1.1)	0.17% (-0.99-1.7%)	0.29% (-1.1-2.1%)	0.15 (-0.54-1.1)	1.6 (-8.9-15)
Samoa	2000	Female	80 to 84	23% (9.1-41%)	77% (59-91%)	0.49 (0.17-1)	0.069% (-0.91-1.5%)	0.12% (-1-1.8%)	0.058 (-0.51-0.85)	0.43 (-5.7-9.1)
Samoa	2000	Female	85 to 89	23% (10-43%)	77% (57-90%)	0.51 (0.19-1.1)	0.008% (-0.99-1.3%)	0.017% (-1.1-1.5%)	0.006 (-0.38-0.53)	0.027 (-2.9-4)
Samoa	2000	Female	90 to 94	26% (12-46%)	74% (54-88%)	0.57 (0.21-1.2)	0.078% (-0.92-1.5%)	0.085% (-1.1-1.8%)	0.013 (-0.16-0.28)	0.073 (-0.83-1.4)
Samoa	2000	Female	95 plus	31% (13-55%)	69% (45-87%)	0.68 (0.22-1.5)	0.037% (-1.4-1.9%)	-0.025% (-1.7-2.1%)	-0.00092 (-0.065-0.085)	0.0059 (-0.21-0.29)
Samoa	2000	Male	All Ages	36% (28-46%)	64% (54-72%)	1 (0.84-1.2)	3.3% (2.2-4.6%)	3.3% (1.9-5%)	18 (10-27)	880 (590-1200)
Samoa	2000	Male	15 to 19	34% (19-52%)	66% (48-81%)	0.67 (0.31-1.2)	2.5% (1.3-4.4%)	3.4% (1.5-6.6%)	0.45 (0.19-0.88)	39 (20-70)
Samoa	2000	Male	20 to 24	52% (33-69%)	48% (31-67%)	1.8 (1.1-2.8)	7.2% (4.4-11%)	9.5% (5.3-15%)	1.5 (0.81-2.5)	120 (66-180)
Samoa	2000	Male	25 to 29	59% (39-76%)	41% (24-61%)	2.5 (1.6-3.7)	9% (5.6-13%)	12% (7.1-18%)	2 (1.1-3.1)	140 (84-210)
Samoa	2000	Male	30 to 34	61% (40-77%)	39% (23-60%)	2.7 (1.7-3.9)	8% (5.1-11%)	10% (6.3-16%)	2 (1.1-3.1)	130 (78-200)
Samoa	2000	Male	35 to 39	59% (41-76%)	41% (24-59%)	2.4 (1.4-3.6)	6.9% (4-10%)	8.8% (4.7-14%)	1.8 (0.89-3)	110 (58-170)
Samoa	2000	Male	40 to 44	57% (39-74%)	43% (26-61%)	2.1 (1.1-3.3)	5.6% (2.9-9%)	6.9% (3.3-11%)	1.7 (0.74-3)	89 (43-150)
Samoa	2000	Male	45 to 49	53% (32-70%)	47% (30-68%)	1.8 (0.89-3)	4.7% (2.2-8.4%)	5.6% (2.4-10%)	1.5 (0.62-2.9)	72 (32-130)
Samoa	2000	Male	50 to 54	50% (30-68%)	50% (32-70%)	1.5 (0.74-2.6)	3.9% (1.5-7.4%)	4.6% (1.5-9.1%)	1.3 (0.46-2.7)	56 (22-110)
Samoa	2000	Male	55 to 59	48% (30-67%)	52% (33-70%)	1.2 (0.59-2.1)	2.7% (0.86-5.7%)	3.1% (0.81-6.8%)	1.2 (0.3-2.7)	42 (13-91)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	2000	Male	60 to 64	45% (26-65%)	55% (35-74%)	1 (0.44-1.9)	2.2% (0.42-5%)	2.4% (0.38-5.8%)	1.1 (0.17-2.7)	33 (6.4-80)
Samoa	2000	Male	65 to 69	43% (25-62%)	57% (38-75%)	0.91 (0.44-1.6)	1.9% (0.29-4.1%)	2.1% (0.28-4.7%)	1.1 (0.15-2.5)	28 (4.5-61)
Samoa	2000	Male	70 to 74	41% (23-59%)	59% (41-77%)	0.79 (0.38-1.4)	1.4% (0.1-3.7%)	1.6% (0.074-4.1%)	0.89 (0.041-2.3)	17 (1.3-44)
Samoa	2000	Male	75 to 79	37% (21-55%)	63% (45-79%)	0.67 (0.32-1.3)	1.1% (0.038-3%)	1.2% (-0.019-3.4%)	0.69 (-0.01-1.9)	11 (0.37-29)
Samoa	2000	Male	80 to 84	34% (18-53%)	66% (47-82%)	0.6 (0.29-1.1)	0.94% (0.075-2.4%)	1% (0.05-2.6%)	0.39 (0.02-1)	4.4 (0.36-11)
Samoa	2000	Male	85 to 89	33% (18-50%)	67% (50-82%)	0.57 (0.28-1)	0.71% (-0.055-2%)	0.75% (-0.11-2.2%)	0.16 (-0.023-0.46)	1.3 (-0.095-3.6)
Samoa	2000	Male	90 to 94	34% (21-52%)	66% (48-79%)	0.58 (0.26-1.1)	0.53% (-0.16-1.5%)	0.53% (-0.28-1.7%)	0.043 (-0.023-0.14)	0.25 (-0.081-0.75)
Samoa	2000	Male	95 plus	38% (20-59%)	62% (41-80%)	0.64 (0.26-1.3)	0.35% (-0.53-1.8%)	0.27% (-0.76-1.9%)	0.0046 (-0.013-0.033)	0.024 (-0.037-0.12)
Samoa	2005	Female	All Ages	17% (12-23%)	83% (77-88%)	0.47 (0.35-0.61)	0.88% (0.35-1.5%)	0.78% (-0.17-2%)	3.6 (-0.9-9.7)	180 (73-330)
Samoa	2005	Female	15 to 19	18% (8.4-32%)	82% (68-92%)	0.58 (0.25-1.1)	1.3% (0.67-2.4%)	2.4% (0.92-5.2%)	0.12 (0.043-0.26)	13 (6.5-24)
Samoa	2005	Female	20 to 24	22% (9.3-41%)	78% (59-91%)	0.91 (0.38-1.7)	2.3% (1.3-3.8%)	3.6% (1.5-7.1%)	0.19 (0.075-0.37)	21 (12-35)
Samoa	2005	Female	25 to 29	24% (9.9-42%)	76% (58-90%)	0.92 (0.39-1.7)	2.4% (1.4-3.7%)	3.6% (1.6-6.3%)	0.24 (0.11-0.44)	23 (13-37)
Samoa	2005	Female	30 to 34	24% (9.5-43%)	76% (57-91%)	0.95 (0.38-1.7)	2.2% (1.2-3.5%)	3.2% (1.5-5.5%)	0.29 (0.13-0.51)	24 (13-39)
Samoa	2005	Female	35 to 39	25% (10-43%)	75% (57-90%)	0.98 (0.41-1.7)	2.2% (1.1-3.6%)	3.2% (1.5-5.6%)	0.4 (0.18-0.71)	26 (14-46)
Samoa	2005	Female	40 to 44	24% (12-41%)	76% (59-88%)	0.9 (0.39-1.6)	1.9% (0.84-3.5%)	2.9% (1.2-5.5%)	0.45 (0.18-0.89)	25 (10-47)
Samoa	2005	Female	45 to 49	24% (10-41%)	76% (59-90%)	0.77 (0.31-1.5)	1.4% (0.3-3.1%)	2.1% (0.6-4.6%)	0.39 (0.11-0.88)	18 (4-41)
Samoa	2005	Female	50 to 54	24% (11-41%)	76% (59-89%)	0.68 (0.27-1.3)	0.97% (-0.26-2.8%)	1.5% (-0.011-4%)	0.35 (-0.0025-0.93)	13 (-3.5-37)
Samoa	2005	Female	55 to 59	23% (10-42%)	77% (58-90%)	0.62 (0.23-1.3)	0.71% (-0.47-2.3%)	1.2% (-0.3-3.3%)	0.29 (-0.078-0.84)	8.3 (-5.5-27)
Samoa	2005	Female	60 to 64	23% (10-42%)	77% (58-90%)	0.6 (0.22-1.3)	0.35% (-0.99-2%)	0.64% (-0.96-2.6%)	0.19 (-0.29-0.83)	4 (-12-23)
Samoa	2005	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.62 (0.23-1.2)	0.29% (-1-2%)	0.54% (-1.1-2.7%)	0.21 (-0.43-1)	3.5 (-13-23)
Samoa	2005	Female	70 to 74	25% (11-42%)	75% (58-89%)	0.59 (0.21-1.2)	0.14% (-1.1-1.7%)	0.28% (-1.2-2.2%)	0.13 (-0.55-1.1)	1.5 (-12-20)
Samoa	2005	Female	75 to 79	23% (10-41%)	77% (59-90%)	0.55 (0.19-1.2)	0.22% (-1.1-1.9%)	0.34% (-1.1-2.3%)	0.21 (-0.69-1.4)	2.5 (-11-20)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	2005	Female	80 to 84	22% (9.8-40%)	78% (60-90%)	0.52 (0.18-1.1)	0.13% (-0.9-1.5%)	0.19% (-1-1.9%)	0.11 (-0.6-1.1)	1 (-7.1-12)
Samoa	2005	Female	85 to 89	23% (11-41%)	77% (59-89%)	0.54 (0.21-1.1)	0.068% (-0.91-1.4%)	0.083% (-1.1-1.7%)	0.035 (-0.46-0.7)	0.26 (-3.5-5.3)
Samoa	2005	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.62 (0.23-1.3)	0.14% (-0.93-1.7%)	0.15% (-1.1-2%)	0.028 (-0.21-0.38)	0.16 (-1-2)
Samoa	2005	Female	95 plus	31% (12-54%)	69% (46-88%)	0.74 (0.24-1.6)	0.094% (-1.4-2.3%)	0.032% (-1.7-2.6%)	0.0018 (-0.095-0.15)	0.021 (-0.31-0.51)
Samoa	2005	Male	All Ages	36% (29-45%)	64% (55-71%)	1.1 (0.91-1.3)	3.4% (2.3-4.7%)	3.3% (1.9-5.1%)	19 (11-29)	910 (610-1300)
Samoa	2005	Male	15 to 19	34% (19-51%)	66% (49-81%)	0.7 (0.33-1.2)	2.6% (1.3-4.5%)	3.6% (1.5-6.8%)	0.44 (0.18-0.87)	38 (19-70)
Samoa	2005	Male	20 to 24	52% (33-70%)	48% (30-67%)	1.9 (1.1-2.9)	7.5% (4.5-11%)	9.9% (5.3-16%)	1.5 (0.77-2.4)	110 (64-180)
Samoa	2005	Male	25 to 29	60% (40-76%)	40% (24-60%)	2.7 (1.7-3.9)	9.2% (5.9-13%)	12% (7.6-19%)	1.8 (1-2.8)	130 (78-190)
Samoa	2005	Male	30 to 34	61% (40-78%)	39% (22-60%)	2.9 (1.8-4.2)	8.3% (5.3-12%)	11% (6.5-16%)	1.8 (1-2.8)	120 (72-180)
Samoa	2005	Male	35 to 39	59% (41-76%)	41% (24-59%)	2.5 (1.5-3.8)	7.2% (4.2-11%)	9.3% (5.1-14%)	1.9 (0.95-3.1)	110 (62-180)
Samoa	2005	Male	40 to 44	57% (39-74%)	43% (26-61%)	2.2 (1.2-3.4)	5.8% (3.1-9.1%)	7.3% (3.5-12%)	1.8 (0.81-3)	95 (48-160)
Samoa	2005	Male	45 to 49	52% (32-71%)	48% (29-68%)	1.9 (0.96-3.2)	4.9% (2.3-8.5%)	5.9% (2.5-11%)	1.7 (0.74-3.2)	84 (39-150)
Samoa	2005	Male	50 to 54	49% (30-69%)	51% (31-70%)	1.5 (0.8-2.6)	4.1% (1.5-7.6%)	4.7% (1.6-9.3%)	1.6 (0.54-3.2)	69 (26-130)
Samoa	2005	Male	55 to 59	47% (30-66%)	53% (34-70%)	1.3 (0.63-2.2)	2.9% (0.92-5.9%)	3.3% (0.9-7.1%)	1.3 (0.35-2.8)	46 (15-99)
Samoa	2005	Male	60 to 64	45% (26-65%)	55% (35-74%)	1.1 (0.49-1.9)	2.3% (0.52-5.3%)	2.6% (0.5-6.2%)	1.2 (0.23-2.8)	35 (7.9-81)
Samoa	2005	Male	65 to 69	43% (25-62%)	57% (38-75%)	0.95 (0.48-1.7)	1.9% (0.4-4.2%)	2.2% (0.41-4.8%)	1.2 (0.22-2.6)	29 (6-63)
Samoa	2005	Male	70 to 74	40% (23-58%)	60% (42-77%)	0.83 (0.4-1.5)	1.5% (0.14-3.8%)	1.7% (0.13-4.3%)	0.95 (0.076-2.4)	19 (2-47)
Samoa	2005	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.7 (0.33-1.3)	1.2% (0.053-3.1%)	1.3% (-0.0023-3.5%)	0.7 (-0.0014-1.9)	11 (0.54-29)
Samoa	2005	Male	80 to 84	34% (19-52%)	66% (48-81%)	0.62 (0.3-1.1)	0.97% (0.14-2.5%)	1% (0.096-2.7%)	0.45 (0.041-1.2)	5.2 (0.74-13)
Samoa	2005	Male	85 to 89	32% (18-49%)	68% (51-82%)	0.59 (0.29-1.1)	0.74% (-0.02-1.9%)	0.78% (-0.085-2.1%)	0.22 (-0.024-0.62)	1.8 (-0.049-4.9)
Samoa	2005	Male	90 to 94	34% (21-51%)	66% (49-79%)	0.61 (0.28-1.2)	0.55% (-0.17-1.7%)	0.55% (-0.29-1.8%)	0.056 (-0.03-0.18)	0.34 (-0.1-1)
Samoa	2005	Male	95 plus	37% (19-58%)	63% (42-81%)	0.66 (0.26-1.4)	0.4% (-0.55-1.9%)	0.31% (-0.78-2.1%)	0.008 (-0.02-0.052)	0.041 (-0.055-0.19)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	2010	Female	All Ages	17% (12-23%)	83% (77-88%)	0.48 (0.35-0.64)	0.92% (0.36-1.7%)	0.81% (-0.2-2.1%)	4.2 (-1-11)	200 (82-380)
Samoa	2010	Female	15 to 19	18% (8.2-33%)	82% (67-92%)	0.58 (0.24-1.1)	1.3% (0.65-2.4%)	2.4% (0.9-5.2%)	0.13 (0.047-0.27)	14 (6.8-26)
Samoa	2010	Female	20 to 24	22% (8.9-39%)	78% (61-91%)	0.91 (0.37-1.7)	2.3% (1.2-3.8%)	3.5% (1.4-6.9%)	0.19 (0.075-0.38)	22 (12-37)
Samoa	2010	Female	25 to 29	23% (9.7-41%)	77% (59-90%)	0.92 (0.39-1.7)	2.4% (1.4-3.9%)	3.5% (1.6-6.4%)	0.24 (0.11-0.45)	23 (13-38)
Samoa	2010	Female	30 to 34	24% (9.2-43%)	76% (57-91%)	0.96 (0.36-1.8)	2.2% (1.2-3.6%)	3.2% (1.5-5.7%)	0.29 (0.13-0.53)	24 (13-40)
Samoa	2010	Female	35 to 39	25% (9.6-43%)	75% (57-90%)	1 (0.4-1.8)	2.2% (1.1-3.7%)	3.3% (1.5-5.6%)	0.4 (0.17-0.74)	27 (14-46)
Samoa	2010	Female	40 to 44	24% (12-41%)	76% (59-88%)	0.91 (0.4-1.6)	2% (0.87-3.7%)	3% (1.2-5.7%)	0.5 (0.19-0.96)	27 (12-51)
Samoa	2010	Female	45 to 49	24% (9.9-41%)	76% (59-90%)	0.78 (0.32-1.5)	1.4% (0.38-3.2%)	2.2% (0.6-4.5%)	0.46 (0.13-1)	21 (5.6-47)
Samoa	2010	Female	50 to 54	23% (10-42%)	77% (58-90%)	0.69 (0.26-1.4)	1% (-0.21-3%)	1.6% (0.048-4.1%)	0.42 (0.014-1.1)	16 (-3.4-44)
Samoa	2010	Female	55 to 59	23% (10-41%)	77% (59-90%)	0.63 (0.24-1.3)	0.78% (-0.46-2.5%)	1.2% (-0.28-3.5%)	0.37 (-0.081-1.1)	11 (-6.8-36)
Samoa	2010	Female	60 to 64	23% (10-41%)	77% (59-90%)	0.6 (0.22-1.3)	0.42% (-0.86-2.3%)	0.71% (-0.89-3.1%)	0.22 (-0.26-0.94)	5.1 (-11-28)
Samoa	2010	Female	65 to 69	24% (11-42%)	76% (58-89%)	0.61 (0.23-1.2)	0.36% (-1-2.1%)	0.62% (-1-2.8%)	0.26 (-0.43-1.2)	4.7 (-13-28)
Samoa	2010	Female	70 to 74	24% (10-41%)	76% (59-90%)	0.58 (0.21-1.2)	0.2% (-1-1.9%)	0.34% (-1.1-2.5%)	0.17 (-0.58-1.3)	2.5 (-13-24)
Samoa	2010	Female	75 to 79	23% (9.5-41%)	77% (59-90%)	0.55 (0.17-1.2)	0.27% (-0.94-2%)	0.4% (-1.1-2.5%)	0.24 (-0.67-1.5)	3 (-11-22)
Samoa	2010	Female	80 to 84	22% (8.8-42%)	78% (58-91%)	0.53 (0.17-1.1)	0.17% (-0.89-1.6%)	0.24% (-1-1.9%)	0.17 (-0.74-1.4)	1.6 (-8.6-15)
Samoa	2010	Female	85 to 89	23% (11-42%)	77% (58-89%)	0.56 (0.21-1.2)	0.1% (-0.9-1.5%)	0.12% (-1-1.8%)	0.069 (-0.59-1)	0.53 (-4.6-7.7)
Samoa	2010	Female	90 to 94	27% (12-47%)	73% (53-88%)	0.64 (0.23-1.3)	0.16% (-0.93-1.8%)	0.17% (-1.1-2%)	0.044 (-0.28-0.51)	0.25 (-1.4-2.7)
Samoa	2010	Female	95 plus	31% (13-55%)	69% (45-87%)	0.75 (0.25-1.6)	0.11% (-1.4-2.3%)	0.044% (-1.8-2.5%)	0.0033 (-0.13-0.19)	0.033 (-0.42-0.68)
Samoa	2010	Male	All Ages	37% (29-46%)	63% (54-71%)	1.1 (0.89-1.4)	3.5% (2.3-4.8%)	3.3% (1.9-5.2%)	19 (11-31)	930 (620-1300)
Samoa	2010	Male	15 to 19	34% (19-52%)	66% (48-81%)	0.71 (0.35-1.2)	2.5% (1.3-4.5%)	3.6% (1.6-6.9%)	0.46 (0.18-0.87)	40 (20-70)
Samoa	2010	Male	20 to 24	51% (33-69%)	49% (31-67%)	1.9 (1.1-3)	7.4% (4.5-11%)	10% (5.5-16%)	1.4 (0.71-2.4)	110 (60-180)
Samoa	2010	Male	25 to 29	59% (39-76%)	41% (24-61%)	2.7 (1.6-4)	9.2% (5.6-13%)	12% (7.2-18%)	1.7 (0.94-2.7)	120 (72-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	2010	Male	30 to 34	60% (39-77%)	40% (23-61%)	2.9 (1.7-4.3)	8.3% (5.2-12%)	11% (6.4-17%)	1.7 (0.91-2.6)	110 (67-170)
Samoa	2010	Male	35 to 39	59% (40-76%)	41% (24-60%)	2.5 (1.5-3.9)	7.1% (4.1-11%)	9.4% (5-15%)	1.8 (0.92-2.9)	110 (60-170)
Samoa	2010	Male	40 to 44	56% (39-74%)	44% (26-61%)	2.2 (1.2-3.5)	5.8% (2.9-9.2%)	7.3% (3.4-12%)	1.9 (0.85-3.3)	100 (50-170)
Samoa	2010	Male	45 to 49	53% (32-71%)	47% (29-68%)	1.9 (1-3.3)	4.9% (2.4-8.4%)	6% (2.6-11%)	1.9 (0.75-3.5)	91 (41-160)
Samoa	2010	Male	50 to 54	49% (31-68%)	51% (32-69%)	1.6 (0.8-2.7)	4.1% (1.6-7.8%)	4.9% (1.7-9.5%)	1.9 (0.65-3.9)	82 (30-160)
Samoa	2010	Male	55 to 59	47% (30-67%)	53% (33-70%)	1.3 (0.65-2.2)	3% (0.97-6.2%)	3.4% (0.97-7.3%)	1.5 (0.43-3.4)	55 (17-120)
Samoa	2010	Male	60 to 64	45% (26-64%)	55% (36-74%)	1.1 (0.51-2)	2.3% (0.55-5.4%)	2.6% (0.51-6.4%)	1.2 (0.23-3)	36 (8.7-86)
Samoa	2010	Male	65 to 69	43% (25-61%)	57% (39-75%)	0.96 (0.48-1.7)	1.9% (0.47-4.4%)	2.2% (0.49-5.1%)	1.2 (0.27-2.8)	29 (7.3-67)
Samoa	2010	Male	70 to 74	40% (24-58%)	60% (42-76%)	0.83 (0.41-1.5)	1.5% (0.18-3.8%)	1.7% (0.13-4.3%)	0.99 (0.079-2.5)	20 (2.5-49)
Samoa	2010	Male	75 to 79	37% (22-55%)	63% (45-78%)	0.71 (0.34-1.3)	1.2% (0.097-3.1%)	1.3% (0.03-3.5%)	0.7 (0.017-1.9)	11 (0.94-30)
Samoa	2010	Male	80 to 84	34% (19-53%)	66% (47-81%)	0.63 (0.31-1.1)	0.99% (0.16-2.4%)	1.1% (0.13-2.7%)	0.5 (0.056-1.3)	5.7 (0.9-14)
Samoa	2010	Male	85 to 89	32% (19-49%)	68% (51-81%)	0.6 (0.29-1.1)	0.75% (-0.014-2%)	0.8% (-0.074-2.3%)	0.27 (-0.024-0.73)	2.2 (-0.042-5.9)
Samoa	2010	Male	90 to 94	34% (20-50%)	66% (50-80%)	0.61 (0.28-1.2)	0.56% (-0.15-1.8%)	0.56% (-0.26-1.9%)	0.083 (-0.039-0.28)	0.5 (-0.13-1.6)
Samoa	2010	Male	95 plus	37% (20-58%)	63% (42-80%)	0.67 (0.26-1.3)	0.43% (-0.49-2.1%)	0.34% (-0.72-2.2%)	0.011 (-0.024-0.074)	0.057 (-0.065-0.27)
Samoa	2016	Female	All Ages	18% (13-24%)	82% (76-87%)	0.49 (0.31-0.71)	0.91% (0.2-2%)	0.79% (-0.37-2.5%)	4.6 (-2.16)	210 (44-450)
Samoa	2016	Female	15 to 19	18% (7.7-33%)	82% (67-92%)	0.58 (0.22-1.2)	1.3% (0.62-2.4%)	2.4% (0.76-5.3%)	0.13 (0.041-0.3)	14 (6.9-27)
Samoa	2016	Female	20 to 24	22% (8.9-39%)	78% (61-91%)	0.91 (0.35-1.7)	2.2% (1.3-3.7%)	3.4% (1.2-7%)	0.21 (0.074-0.43)	24 (13-42)
Samoa	2016	Female	25 to 29	23% (9.6-41%)	77% (59-90%)	0.93 (0.36-1.9)	2.3% (1.3-3.9%)	3.4% (1.3-6.9%)	0.22 (0.081-0.45)	22 (12-39)
Samoa	2016	Female	30 to 34	24% (9.1-44%)	76% (56-91%)	0.96 (0.35-1.9)	2.2% (1.1-3.8%)	3.2% (1.2-6.3%)	0.26 (0.097-0.51)	21 (11-38)
Samoa	2016	Female	35 to 39	25% (9.7-43%)	75% (57-90%)	1 (0.38-1.9)	2.2% (1.1-4%)	3.3% (1.4-6.1%)	0.38 (0.15-0.72)	25 (12-47)
Samoa	2016	Female	40 to 44	24% (11-41%)	76% (59-89%)	0.93 (0.38-1.8)	2% (0.77-3.9%)	3% (1.1-5.9%)	0.48 (0.16-0.99)	27 (10-53)
Samoa	2016	Female	45 to 49	24% (10-42%)	76% (58-90%)	0.79 (0.3-1.6)	1.5% (0.28-3.4%)	2.2% (0.54-5%)	0.49 (0.11-1.2)	23 (4.2-53)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	2016	Female	50 to 54	24% (11-42%)	76% (58-89%)	0.7 (0.27-1.4)	1.1% (-0.25-3.1%)	1.6% (-0.064-4.3%)	0.48 (-0.017-1.4)	18 (-4.2-59)
Samoa	2016	Female	55 to 59	23% (11-41%)	77% (59-89%)	0.64 (0.23-1.4)	0.79% (-0.53-2.9%)	1.2% (-0.43-4%)	0.45 (-0.16-1.5)	14 (-9.4-52)
Samoa	2016	Female	60 to 64	24% (11-43%)	76% (57-89%)	0.62 (0.21-1.3)	0.43% (-0.97-2.5%)	0.72% (-0.98-3.4%)	0.3 (-0.41-1.4)	6.9 (-16-41)
Samoa	2016	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.63 (0.22-1.3)	0.4% (-1.1-2.5%)	0.66% (-1.1-3.3%)	0.28 (-0.46-1.4)	5.3 (-14-33)
Samoa	2016	Female	70 to 74	24% (11-41%)	76% (59-89%)	0.6 (0.21-1.3)	0.23% (-1.1-2.2%)	0.37% (-1.2-2.8%)	0.21 (-0.73-1.6)	3.2 (-16-31)
Samoa	2016	Female	75 to 79	23% (9.8-41%)	77% (59-90%)	0.56 (0.17-1.3)	0.3% (-1.2-2.2%)	0.42% (-1.1-2.7%)	0.29 (-0.73-1.8)	3.7 (-12-27)
Samoa	2016	Female	80 to 84	22% (8.9-41%)	78% (59-91%)	0.54 (0.16-1.2)	0.2% (-0.98-1.9%)	0.26% (-1.1-2.2%)	0.21 (-0.96-1.9)	2.1 (-11-21)
Samoa	2016	Female	85 to 89	24% (10-42%)	76% (58-90%)	0.56 (0.19-1.2)	0.14% (-0.97-1.7%)	0.15% (-1.1-2.1%)	0.11 (-0.8-1.4)	0.84 (-6.2-11)
Samoa	2016	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.65 (0.22-1.4)	0.18% (-0.92-2%)	0.19% (-1.1-2.3%)	0.072 (-0.41-0.86)	0.42 (-2.1-4.6)
Samoa	2016	Female	95 plus	32% (12-54%)	68% (46-88%)	0.76 (0.24-1.7)	0.12% (-1.5-2.4%)	0.054% (-1.8-2.7%)	0.0058 (-0.19-0.29)	0.051 (-0.62-1)
Samoa	2016	Male	All Ages	37% (29-45%)	63% (55-71%)	1.1 (0.77-1.5)	3.5% (2.1-5.5%)	3.3% (1.5-5.8%)	21 (9.3-38)	980 (550-1500)
Samoa	2016	Male	15 to 19	34% (19-51%)	66% (49-81%)	0.72 (0.33-1.3)	2.5% (1.3-4.4%)	3.7% (1.6-7.2%)	0.45 (0.17-0.88)	41 (19-73)
Samoa	2016	Male	20 to 24	52% (34-69%)	48% (31-66%)	2 (1-3.2)	7.5% (4.2-12%)	10% (5.1-17%)	1.6 (0.78-2.9)	130 (66-210)
Samoa	2016	Male	25 to 29	59% (39-75%)	41% (25-61%)	2.7 (1.5-4.5)	9.1% (5.3-14%)	12% (6.7-20%)	1.7 (0.83-2.7)	120 (67-190)
Samoa	2016	Male	30 to 34	60% (39-78%)	40% (22-61%)	2.9 (1.5-4.8)	8.3% (4.9-13%)	11% (5.9-17%)	1.5 (0.76-2.6)	100 (57-160)
Samoa	2016	Male	35 to 39	59% (40-77%)	41% (23-60%)	2.6 (1.3-4.3)	7.3% (4-12%)	9.5% (4.7-16%)	1.6 (0.78-2.7)	98 (51-160)
Samoa	2016	Male	40 to 44	56% (38-74%)	44% (26-62%)	2.2 (1.1-3.7)	5.9% (2.7-9.8%)	7.4% (3-13%)	1.8 (0.72-3.3)	98 (42-170)
Samoa	2016	Male	45 to 49	52% (30-71%)	48% (29-70%)	1.9 (0.93-3.4)	5% (2.2-8.8%)	6% (2.3-11%)	2.1 (0.76-4)	100 (44-190)
Samoa	2016	Male	50 to 54	49% (30-69%)	51% (31-70%)	1.6 (0.74-2.8)	4.2% (1.5-7.9%)	5% (1.5-9.4%)	2.2 (0.65-4.4)	93 (30-180)
Samoa	2016	Male	55 to 59	47% (29-67%)	53% (33-71%)	1.3 (0.6-2.4)	3.1% (0.9-6.7%)	3.5% (0.89-7.8%)	1.9 (0.5-4.5)	70 (21-160)
Samoa	2016	Male	60 to 64	45% (26-65%)	55% (35-74%)	1.1 (0.47-2.1)	2.4% (0.47-5.7%)	2.7% (0.41-6.7%)	1.6 (0.23-4.2)	49 (9.6-120)
Samoa	2016	Male	65 to 69	43% (25-61%)	57% (39-75%)	0.96 (0.43-1.7)	2% (0.36-4.7%)	2.3% (0.38-5.5%)	1.3 (0.2-3.2)	32 (5.6-76)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Samoa	2016	Male	70 to 74	40% (24-58%)	60% (42-76%)	0.85 (0.39-1.6)	1.5% (0.18-4.2%)	1.7% (0.15-4.7%)	1 (0.093-2.8)	20 (2.3-55)
Samoa	2016	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.72 (0.3-1.4)	1.2% (0.068-3.4%)	1.3% (0.013-3.8%)	0.83 (0.0066-2.4)	13 (0.69-37)
Samoa	2016	Male	80 to 84	33% (18-51%)	67% (49-82%)	0.63 (0.29-1.2)	1% (0.12-2.7%)	1.1% (0.084-3%)	0.57 (0.04-1.6)	6.6 (0.69-18)
Samoa	2016	Male	85 to 89	32% (19-48%)	68% (52-81%)	0.6 (0.27-1.1)	0.8% (-0.051-2.1%)	0.85% (-0.14-2.4%)	0.31 (-0.054-0.86)	2.6 (-0.16-6.8)
Samoa	2016	Male	90 to 94	34% (20-51%)	66% (49-80%)	0.63 (0.26-1.2)	0.62% (-0.13-1.9%)	0.62% (-0.23-2.1%)	0.11 (-0.041-0.38)	0.66 (-0.13-2.1)
Samoa	2016	Male	95 plus	38% (19-59%)	62% (41-81%)	0.69 (0.25-1.5)	0.49% (-0.48-2.2%)	0.4% (-0.72-2.3%)	0.021 (-0.038-0.12)	0.099 (-0.097-0.44)
Solomon Islands	1990	Female	All Ages	18% (13-25%)	82% (75-87%)	0.14 (0.11-0.18)	0.064% (-0.2-0.3%)	-0.2% (-0.65-0.21%)	-2.2 (-7.3-2.4)	41 (-130-190)
Solomon Islands	1990	Female	15 to 19	20% (8.9-36%)	80% (64-91%)	0.19 (0.079-0.35)	0.51% (0.18-1%)	0.5% (-0.023-1.3%)	0.13 (-0.0057-0.34)	17 (5.8-33)
Solomon Islands	1990	Female	20 to 24	24% (10-43%)	76% (57-90%)	0.29 (0.13-0.53)	0.88% (0.43-1.5%)	0.76% (0.15-1.8%)	0.22 (0.045-0.52)	27 (13-49)
Solomon Islands	1990	Female	25 to 29	25% (11-45%)	75% (55-89%)	0.29 (0.13-0.51)	0.86% (0.47-1.4%)	0.77% (0.23-1.5%)	0.26 (0.075-0.53)	27 (14-45)
Solomon Islands	1990	Female	30 to 34	27% (11-47%)	73% (53-89%)	0.31 (0.12-0.55)	0.64% (0.22-1.1%)	0.55% (0.016-1.3%)	0.22 (0.0063-0.5)	20 (6.8-37)
Solomon Islands	1990	Female	35 to 39	27% (11-47%)	73% (53-89%)	0.32 (0.13-0.56)	0.57% (0.052-1.2%)	0.58% (-0.11-1.4%)	0.28 (-0.05-0.7)	19 (1.8-40)
Solomon Islands	1990	Female	40 to 44	27% (13-45%)	73% (55-87%)	0.3 (0.13-0.55)	0.33% (-0.31-0.99%)	0.36% (-0.49-1.2%)	0.2 (-0.28-0.71)	12 (-11-35)
Solomon Islands	1990	Female	45 to 49	26% (11-45%)	74% (55-89%)	0.26 (0.11-0.48)	-0.056% (-0.77-0.47%)	-0.062% (-0.91-0.6%)	-0.042 (-0.64-0.42)	-2 (-28-17)
Solomon Islands	1990	Female	50 to 54	26% (12-46%)	74% (54-88%)	0.23 (0.087-0.45)	-0.3% (-1.2-0.32%)	-0.3% (-1.3-0.45%)	-0.19 (-0.86-0.27)	-8.5 (-35-9.2)
Solomon Islands	1990	Female	55 to 59	25% (12-45%)	75% (55-88%)	0.2 (0.075-0.42)	-0.42% (-1.5-0.19%)	-0.41% (-1.6-0.28%)	-0.33 (-1.3-0.23)	-13 (-46-5.8)
Solomon Islands	1990	Female	60 to 64	25% (11-45%)	75% (55-89%)	0.19 (0.069-0.41)	-0.63% (-1.8-0.029%)	-0.65% (-1.9-0.076%)	-0.5 (-1.4-0.057)	-16 (-43-0.67)
Solomon Islands	1990	Female	65 to 69	26% (12-43%)	74% (57-88%)	0.2 (0.073-0.38)	-0.7% (-1.7--0.0095%)	-0.72% (-1.9-0.039%)	-0.68 (-1.7-0.034)	-17 (-43--0.25)
Solomon Islands	1990	Female	70 to 74	26% (12-44%)	74% (56-88%)	0.19 (0.07-0.37)	-0.73% (-1.6--0.058%)	-0.76% (-1.8--0.039%)	-0.65 (-1.5--0.032)	-13 (-30--1.1)
Solomon Islands	1990	Female	75 to 79	25% (11-43%)	75% (57-89%)	0.17 (0.055-0.39)	-0.62% (-1.7--0.0024%)	-0.65% (-1.8-0.025%)	-0.51 (-1.4-0.018)	-7.6 (-20--0.029)
Solomon Islands	1990	Female	80 to 84	24% (10-43%)	76% (57-90%)	0.16 (0.053-0.35)	-0.6% (-1.5--0.068%)	-0.65% (-1.7--0.068%)	-0.38 (-0.95--0.039)	-4.1 (-10--0.47)
Solomon Islands	1990	Female	85 to 89	25% (11-44%)	75% (56-89%)	0.17 (0.063-0.36)	-0.65% (-1.6--0.11%)	-0.72% (-1.8--0.12%)	-0.17 (-0.43--0.031)	-1.3 (-3.2--0.22)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	1990	Female	90 to 94	29% (13-48%)	71% (52-87%)	0.2 (0.068-0.41)	-0.56% (-1.3--0.1%)	-0.63% (-1.4--0.12%)	-0.047 (-0.11--0.0088)	-0.25 (-0.56--0.046)
Solomon Islands	1990	Female	95 plus	34% (15-55%)	66% (45-85%)	0.23 (0.075-0.47)	-0.77% (-1.9--0.06%)	-0.9% (-2.1--0.091%)	-0.017 (-0.039--0.0017)	-0.055 (-0.13--0.0044)
Solomon Islands	1990	Male	All Ages	37% (30-47%)	63% (53-70%)	0.3 (0.25-0.37)	0.87% (0.39-1.4%)	0.72% (0.085-1.4%)	11 (1.3-21)	680 (310-1100)
Solomon Islands	1990	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.23 (0.11-0.38)	1% (0.44-1.9%)	1% (0.26-2.3%)	0.43 (0.1-0.94)	42 (18-79)
Solomon Islands	1990	Male	20 to 24	56% (37-73%)	44% (27-63%)	0.6 (0.36-0.89)	2.8% (1.4-4.5%)	3% (1.2-5.2%)	1.5 (0.58-2.6)	120 (59-200)
Solomon Islands	1990	Male	25 to 29	63% (44-78%)	37% (22-56%)	0.83 (0.5-1.2)	3.4% (1.8-5.4%)	3.6% (1.6-6.2%)	1.9 (0.82-3.2)	140 (71-220)
Solomon Islands	1990	Male	30 to 34	64% (43-80%)	36% (20-57%)	0.89 (0.53-1.3)	3% (1.5-4.9%)	3.2% (1.3-5.5%)	1.7 (0.63-3)	110 (53-190)
Solomon Islands	1990	Male	35 to 39	62% (44-79%)	38% (21-56%)	0.78 (0.45-1.2)	2.2% (1-3.8%)	2.3% (0.72-4.2%)	1.2 (0.38-2.4)	76 (34-130)
Solomon Islands	1990	Male	40 to 44	60% (41-76%)	40% (24-59%)	0.68 (0.37-1.1)	1.6% (0.44-3.1%)	1.6% (0.14-3.3%)	1.1 (0.097-2.3)	60 (16-120)
Solomon Islands	1990	Male	45 to 49	56% (34-74%)	44% (26-66%)	0.58 (0.3-0.99)	1.2% (0.0075-2.6%)	1.2% (-0.29-2.8%)	1.1 (-0.27-2.5)	55 (0.34-120)
Solomon Islands	1990	Male	50 to 54	53% (34-72%)	47% (28-66%)	0.48 (0.24-0.83)	0.93% (-0.19-2.1%)	0.88% (-0.38-2.2%)	0.84 (-0.37-2.2)	37 (-7.5-88)
Solomon Islands	1990	Male	55 to 59	51% (33-70%)	49% (30-67%)	0.39 (0.19-0.7)	0.47% (-0.42-1.4%)	0.42% (-0.62-1.5%)	0.46 (-0.67-1.7)	19 (-17-57)
Solomon Islands	1990	Male	60 to 64	48% (29-68%)	52% (32-71%)	0.33 (0.16-0.59)	0.29% (-0.49-1.1%)	0.27% (-0.62-1.2%)	0.28 (-0.65-1.3)	9.3 (-16-36)
Solomon Islands	1990	Male	65 to 69	47% (28-66%)	53% (34-72%)	0.3 (0.15-0.51)	0.25% (-0.46-1%)	0.24% (-0.56-1.1%)	0.28 (-0.63-1.3)	7.1 (-13-30)
Solomon Islands	1990	Male	70 to 74	44% (27-62%)	56% (38-73%)	0.26 (0.13-0.47)	0.12% (-0.46-0.76%)	0.11% (-0.53-0.83%)	0.13 (-0.61-0.97)	2.8 (-11-18)
Solomon Islands	1990	Male	75 to 79	40% (25-59%)	60% (41-75%)	0.23 (0.1-0.43)	0.061% (-0.43-0.56%)	0.04% (-0.5-0.59%)	0.04 (-0.5-0.61)	0.96 (-6.6-8.9)
Solomon Islands	1990	Male	80 to 84	37% (21-56%)	63% (44-79%)	0.2 (0.099-0.35)	0.055% (-0.32-0.4%)	0.032% (-0.39-0.43%)	0.024 (-0.29-0.31)	0.48 (-2.7-3.6)
Solomon Islands	1990	Male	85 to 89	35% (21-52%)	65% (48-79%)	0.19 (0.088-0.33)	-0.034% (-0.42-0.29%)	-0.073% (-0.52-0.3%)	-0.028 (-0.2-0.12)	-0.11 (-1.4-0.97)
Solomon Islands	1990	Male	90 to 94	37% (22-54%)	63% (46-78%)	0.19 (0.089-0.36)	-0.04% (-0.4-0.24%)	-0.087% (-0.51-0.23%)	-0.012 (-0.072-0.033)	-0.033 (-0.32-0.2)
Solomon Islands	1990	Male	95 plus	41% (22-62%)	59% (38-78%)	0.22 (0.088-0.43)	-0.22% (-0.72-0.11%)	-0.34% (-0.94-0.055%)	-0.013 (-0.037-0.0022)	-0.034 (-0.11-0.017)
Solomon Islands	1995	Female	All Ages	19% (13-25%)	81% (75-87%)	0.13 (0.096-0.17)	0.015% (-0.25-0.24%)	-0.28% (-0.75-0.12%)	-3.6 (-9.9-1.4)	11 (-180-170)
Solomon Islands	1995	Female	15 to 19	20% (9.4-36%)	80% (64-91%)	0.16 (0.07-0.31)	0.49% (0.18-0.96%)	0.47% (-0.03-1.3%)	0.13 (-0.0084-0.35)	18 (6.8-35)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	1995	Female	20 to 24	25% (10-42%)	75% (58-90%)	0.26 (0.12-0.46)	0.82% (0.41-1.3%)	0.66% (0.11-1.5%)	0.24 (0.042-0.58)	33 (16-56)
Solomon Islands	1995	Female	25 to 29	26% (11-45%)	74% (55-89%)	0.26 (0.11-0.45)	0.8% (0.44-1.2%)	0.67% (0.19-1.3%)	0.26 (0.073-0.54)	29 (15-46)
Solomon Islands	1995	Female	30 to 34	27% (11-48%)	73% (52-89%)	0.27 (0.11-0.49)	0.55% (0.18-0.99%)	0.42% (-0.054-1%)	0.19 (-0.025-0.47)	20 (6.6-37)
Solomon Islands	1995	Female	35 to 39	27% (11-46%)	73% (54-89%)	0.29 (0.12-0.5)	0.44% (-0.069-0.91%)	0.4% (-0.21-1.1%)	0.23 (-0.11-0.63)	17 (-2.4-37)
Solomon Islands	1995	Female	40 to 44	27% (13-44%)	73% (56-87%)	0.26 (0.12-0.47)	0.2% (-0.45-0.73%)	0.18% (-0.57-0.89%)	0.11 (-0.39-0.55)	7.2 (-18-27)
Solomon Islands	1995	Female	45 to 49	26% (12-44%)	74% (56-88%)	0.23 (0.09-0.44)	-0.15% (-0.88-0.34%)	-0.18% (-1-0.44%)	-0.13 (-0.78-0.3)	-6 (-35-13)
Solomon Islands	1995	Female	50 to 54	26% (12-44%)	74% (56-88%)	0.2 (0.076-0.4)	-0.4% (-1.3-0.16%)	-0.42% (-1.4-0.23%)	-0.38 (-1.3-0.2)	-16 (-54-7)
Solomon Islands	1995	Female	55 to 59	25% (11-44%)	75% (56-89%)	0.18 (0.067-0.37)	-0.49% (-1.6-0.1%)	-0.5% (-1.7-0.18%)	-0.41 (-1.4-0.14)	-16 (-50-3.4)
Solomon Islands	1995	Female	60 to 64	25% (11-45%)	75% (55-89%)	0.17 (0.058-0.38)	-0.7% (-1.9--0.046%)	-0.72% (-2-0.0047%)	-0.73 (-2.1-0.0047)	-23 (-62--1.4)
Solomon Islands	1995	Female	65 to 69	27% (12-45%)	73% (55-88%)	0.18 (0.061-0.34)	-0.78% (-1.8--0.054%)	-0.81% (-2--0.024%)	-0.79 (-2--0.022)	-20 (-49--1.4)
Solomon Islands	1995	Female	70 to 74	27% (12-45%)	73% (55-88%)	0.17 (0.06-0.33)	-0.8% (-1.8--0.12%)	-0.85% (-1.9--0.1%)	-0.97 (-2.3--0.12)	-19 (-43--2.7)
Solomon Islands	1995	Female	75 to 79	25% (11-46%)	75% (54-89%)	0.15 (0.048-0.34)	-0.66% (-1.7--0.034%)	-0.69% (-1.8--0.02%)	-0.6 (-1.6--0.017)	-9 (-23--0.43)
Solomon Islands	1995	Female	80 to 84	24% (11-45%)	76% (55-89%)	0.14 (0.047-0.31)	-0.64% (-1.6--0.088%)	-0.7% (-1.7--0.1%)	-0.45 (-1.1--0.066)	-4.8 (-12--0.65)
Solomon Islands	1995	Female	85 to 89	25% (11-44%)	75% (56-89%)	0.15 (0.054-0.32)	-0.68% (-1.6--0.12%)	-0.75% (-1.8--0.13%)	-0.22 (-0.52--0.039)	-1.7 (-4--0.29)
Solomon Islands	1995	Female	90 to 94	29% (14-49%)	71% (51-86%)	0.17 (0.06-0.36)	-0.6% (-1.3--0.13%)	-0.68% (-1.5--0.14%)	-0.056 (-0.12--0.012)	-0.29 (-0.65--0.062)
Solomon Islands	1995	Female	95 plus	34% (15-56%)	66% (44-85%)	0.2 (0.067-0.43)	-0.81% (-1.9--0.088%)	-0.94% (-2.2--0.13%)	-0.018 (-0.043--0.0026)	-0.061 (-0.14--0.0066)
Solomon Islands	1995	Male	All Ages	38% (30-47%)	62% (53-70%)	0.28 (0.22-0.34)	0.81% (0.34-1.3%)	0.63% (0.022-1.2%)	10 (0.35-20)	690 (280-1100)
Solomon Islands	1995	Male	15 to 19	37% (22-56%)	63% (44-78%)	0.2 (0.095-0.33)	0.96% (0.45-1.8%)	0.99% (0.24-2.2%)	0.44 (0.11-0.99)	46 (21-85)
Solomon Islands	1995	Male	20 to 24	55% (37-72%)	45% (28-63%)	0.52 (0.31-0.77)	2.5% (1.3-4.2%)	2.6% (0.94-4.8%)	1.7 (0.59-3)	140 (66-240)
Solomon Islands	1995	Male	25 to 29	62% (43-78%)	38% (22-57%)	0.73 (0.46-1)	3% (1.6-4.8%)	3.2% (1.4-5.5%)	1.8 (0.79-3.3)	140 (72-230)
Solomon Islands	1995	Male	30 to 34	63% (43-79%)	37% (21-57%)	0.78 (0.47-1.2)	2.6% (1.3-4.2%)	2.6% (0.98-4.7%)	1.6 (0.59-3)	110 (55-190)
Solomon Islands	1995	Male	35 to 39	62% (44-78%)	38% (22-56%)	0.68 (0.39-1.1)	2% (0.87-3.3%)	1.9% (0.47-3.6%)	1.3 (0.32-2.5)	83 (34-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	1995	Male	40 to 44	60% (42-77%)	40% (23-58%)	0.59 (0.31-0.97)	1.4% (0.25-2.7%)	1.3% (-0.12-2.9%)	0.91 (-0.074-2)	54 (10-110)
Solomon Islands	1995	Male	45 to 49	56% (34-73%)	44% (27-66%)	0.51 (0.26-0.88)	1% (-0.1-2.2%)	0.95% (-0.42-2.3%)	0.83 (-0.38-2.1)	44 (-4.1-96)
Solomon Islands	1995	Male	50 to 54	53% (32-72%)	47% (28-68%)	0.42 (0.2-0.73)	0.75% (-0.29-1.7%)	0.67% (-0.55-1.8%)	0.77 (-0.61-2.2)	37 (-14-86)
Solomon Islands	1995	Male	55 to 59	51% (33-70%)	49% (30-67%)	0.34 (0.16-0.61)	0.36% (-0.5-1.1%)	0.29% (-0.7-1.2%)	0.33 (-0.81-1.4)	15 (-21-49)
Solomon Islands	1995	Male	60 to 64	48% (29-68%)	52% (32-71%)	0.29 (0.14-0.52)	0.22% (-0.51-0.88%)	0.18% (-0.67-0.95%)	0.23 (-0.84-1.2)	8.6 (-20-35)
Solomon Islands	1995	Male	65 to 69	47% (28-66%)	53% (34-72%)	0.26 (0.13-0.47)	0.16% (-0.46-0.81%)	0.14% (-0.59-0.88%)	0.16 (-0.66-1)	4.7 (-14-24)
Solomon Islands	1995	Male	70 to 74	44% (27-63%)	56% (37-73%)	0.23 (0.11-0.42)	0.055% (-0.47-0.61%)	0.035% (-0.55-0.66%)	0.041 (-0.67-0.79)	1.3 (-11-15)
Solomon Islands	1995	Male	75 to 79	40% (24-59%)	60% (41-76%)	0.2 (0.093-0.38)	0.011% (-0.47-0.42%)	-0.02% (-0.57-0.43%)	-0.02 (-0.57-0.44)	0.18 (-7.4-6.6)
Solomon Islands	1995	Male	80 to 84	37% (21-55%)	63% (45-79%)	0.18 (0.085-0.32)	0.032% (-0.33-0.34%)	0.004% (-0.39-0.34%)	0.0029 (-0.29-0.24)	0.27 (-2.8-2.9)
Solomon Islands	1995	Male	85 to 89	36% (21-52%)	64% (48-79%)	0.17 (0.077-0.3)	-0.057% (-0.45-0.23%)	-0.1% (-0.55-0.22%)	-0.039 (-0.21-0.085)	-0.18 (-1.5-0.74)
Solomon Islands	1995	Male	90 to 94	37% (23-55%)	63% (45-77%)	0.17 (0.076-0.32)	-0.053% (-0.42-0.2%)	-0.1% (-0.54-0.19%)	-0.014 (-0.072-0.026)	-0.042 (-0.33-0.16)
Solomon Islands	1995	Male	95 plus	41% (22-62%)	59% (38-78%)	0.19 (0.078-0.38)	-0.23% (-0.73-0.1%)	-0.35% (-0.96-0.043%)	-0.013 (-0.036-0.0017)	-0.034 (-0.11-0.015)
Solomon Islands	2000	Female	All Ages	19% (13-26%)	81% (74-87%)	0.14 (0.11-0.19)	0.013% (-0.29-0.25%)	-0.31% (-0.83-0.11%)	-4.6 (-13-1.6)	11 (-230-200)
Solomon Islands	2000	Female	15 to 19	20% (9.2-35%)	80% (65-91%)	0.18 (0.074-0.34)	0.53% (0.21-1%)	0.56% (0.034-1.4%)	0.17 (0.009-0.42)	21 (8.1-40)
Solomon Islands	2000	Female	20 to 24	24% (11-43%)	76% (57-89%)	0.28 (0.12-0.5)	0.9% (0.46-1.5%)	0.79% (0.2-1.7%)	0.34 (0.082-0.73)	42 (21-72)
Solomon Islands	2000	Female	25 to 29	26% (11-45%)	74% (55-89%)	0.29 (0.13-0.49)	0.88% (0.48-1.3%)	0.79% (0.27-1.5%)	0.41 (0.12-0.81)	43 (23-66)
Solomon Islands	2000	Female	30 to 34	27% (11-48%)	73% (52-89%)	0.3 (0.12-0.55)	0.59% (0.2-1%)	0.49% (-0.052-1.1%)	0.25 (-0.028-0.6)	25 (8.3-45)
Solomon Islands	2000	Female	35 to 39	28% (11-47%)	72% (53-89%)	0.31 (0.12-0.55)	0.45% (-0.076-0.94%)	0.44% (-0.18-1.1%)	0.29 (-0.13-0.74)	21 (-3.7-45)
Solomon Islands	2000	Female	40 to 44	27% (13-45%)	73% (55-87%)	0.28 (0.12-0.52)	0.18% (-0.52-0.71%)	0.17% (-0.66-0.87%)	0.13 (-0.49-0.64)	8.2 (-23-32)
Solomon Islands	2000	Female	45 to 49	26% (11-44%)	74% (56-89%)	0.24 (0.098-0.47)	-0.18% (-0.95-0.37%)	-0.2% (-1.1-0.46%)	-0.16 (-0.89-0.36)	-7.6 (-40-16)
Solomon Islands	2000	Female	50 to 54	26% (12-44%)	74% (56-88%)	0.22 (0.079-0.44)	-0.43% (-1.4-0.16%)	-0.45% (-1.5-0.23%)	-0.45 (-1.6-0.23)	-20 (-65-7.5)
Solomon Islands	2000	Female	55 to 59	25% (11-44%)	75% (56-89%)	0.19 (0.072-0.4)	-0.54% (-1.7-0.095%)	-0.53% (-1.8-0.18%)	-0.65 (-2.2-0.22)	-25 (-80-4.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	2000	Female	60 to 64	25% (11-44%)	75% (56-89%)	0.18 (0.062-0.39)	-0.74% (-2--0.047%)	-0.76% (-2.1--0.018%)	-0.81 (-2.3--0.017)	-25 (-68--1.5)
Solomon Islands	2000	Female	65 to 69	26% (12-44%)	74% (56-88%)	0.19 (0.065-0.37)	-0.82% (-1.9--0.077%)	-0.85% (-2.1--0.045%)	-1.1 (-2.8--0.052)	-28 (-68--2.7)
Solomon Islands	2000	Female	70 to 74	26% (12-44%)	74% (56-88%)	0.18 (0.063-0.35)	-0.86% (-1.9--0.15%)	-0.91% (-2.1--0.15%)	-1.1 (-2.5--0.17)	-22 (-47--3.6)
Solomon Islands	2000	Female	75 to 79	25% (11-44%)	75% (56-89%)	0.16 (0.053-0.35)	-0.71% (-1.8--0.047%)	-0.75% (-1.9--0.037%)	-0.89 (-2.2--0.044)	-13 (-33--0.82)
Solomon Islands	2000	Female	80 to 84	24% (10-44%)	76% (56-90%)	0.15 (0.05-0.33)	-0.7% (-1.7--0.12%)	-0.76% (-1.8--0.14%)	-0.63 (-1.5--0.11)	-6.8 (-16--1.2)
Solomon Islands	2000	Female	85 to 89	25% (12-44%)	75% (56-88%)	0.16 (0.057-0.34)	-0.74% (-1.8--0.16%)	-0.83% (-2--0.17%)	-0.27 (-0.63--0.057)	-2 (-4.8--0.44)
Solomon Islands	2000	Female	90 to 94	29% (13-48%)	71% (52-87%)	0.19 (0.065-0.39)	-0.67% (-1.5--0.14%)	-0.75% (-1.7--0.16%)	-0.074 (-0.16--0.016)	-0.38 (-0.87--0.079)
Solomon Islands	2000	Female	95 plus	34% (14-56%)	66% (44-86%)	0.22 (0.07-0.47)	-0.9% (-2.1--0.11%)	-1% (-2.4--0.15%)	-0.022 (-0.051--0.0033)	-0.073 (-0.17--0.0089)
Solomon Islands	2000	Male	All Ages	39% (30-48%)	61% (52-70%)	0.31 (0.25-0.39)	0.93% (0.43-1.5%)	0.72% (0.087-1.4%)	13 (1.6-25)	880 (410-1400)
Solomon Islands	2000	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.22 (0.1-0.38)	1.1% (0.51-1.9%)	1.2% (0.34-2.5%)	0.54 (0.15-1.1)	53 (24-97)
Solomon Islands	2000	Male	20 to 24	55% (36-72%)	45% (28-64%)	0.58 (0.34-0.86)	2.9% (1.5-4.6%)	3.1% (1.3-5.5%)	2.2 (0.89-3.9)	180 (89-290)
Solomon Islands	2000	Male	25 to 29	62% (43-78%)	38% (22-57%)	0.79 (0.51-1.1)	3.4% (1.9-5.1%)	3.6% (1.8-5.9%)	2.7 (1.3-4.6)	200 (110-320)
Solomon Islands	2000	Male	30 to 34	63% (42-79%)	37% (21-58%)	0.83 (0.5-1.3)	2.8% (1.5-4.4%)	2.9% (1.2-4.9%)	2 (0.86-3.7)	140 (72-240)
Solomon Islands	2000	Male	35 to 39	62% (44-78%)	38% (22-56%)	0.73 (0.42-1.2)	2.1% (0.89-3.5%)	2.1% (0.56-3.9%)	1.7 (0.41-3.1)	110 (44-180)
Solomon Islands	2000	Male	40 to 44	59% (42-76%)	41% (24-58%)	0.64 (0.34-1.1)	1.4% (0.32-2.8%)	1.3% (-0.049-3%)	1.2 (-0.042-2.7)	71 (16-140)
Solomon Islands	2000	Male	45 to 49	56% (35-73%)	44% (27-65%)	0.55 (0.27-0.96)	1.1% (-0.058-2.3%)	1% (-0.39-2.5%)	0.94 (-0.37-2.4)	51 (-2.7-110)
Solomon Islands	2000	Male	50 to 54	53% (33-71%)	47% (29-67%)	0.46 (0.22-0.81)	0.8% (-0.24-1.8%)	0.72% (-0.49-2%)	0.81 (-0.57-2.3)	38 (-12-90)
Solomon Islands	2000	Male	55 to 59	50% (32-69%)	50% (31-68%)	0.38 (0.18-0.68)	0.38% (-0.49-1.3%)	0.3% (-0.68-1.3%)	0.42 (-0.97-1.9)	19 (-26-64)
Solomon Islands	2000	Male	60 to 64	48% (28-68%)	52% (32-72%)	0.31 (0.15-0.58)	0.21% (-0.5-0.95%)	0.17% (-0.66-0.98%)	0.23 (-0.93-1.4)	9 (-22-40)
Solomon Islands	2000	Male	65 to 69	46% (27-65%)	54% (35-73%)	0.29 (0.14-0.5)	0.15% (-0.43-0.78%)	0.14% (-0.54-0.85%)	0.2 (-0.83-1.3)	5.7 (-16-29)
Solomon Islands	2000	Male	70 to 74	44% (26-62%)	56% (38-74%)	0.25 (0.12-0.47)	0.042% (-0.51-0.58%)	0.019% (-0.61-0.65%)	0.023 (-0.74-0.77)	1.1 (-13-15)
Solomon Islands	2000	Male	75 to 79	40% (24-59%)	60% (41-76%)	0.22 (0.1-0.41)	-0.01% (-0.5-0.43%)	-0.045% (-0.6-0.45%)	-0.049 (-0.67-0.5)	-0.17 (-8.7-7.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	2000	Male	80 to 84	37% (20-56%)	63% (44-80%)	0.19 (0.093-0.35)	0.021% (-0.37-0.36%)	-0.01% (-0.45-0.37%)	-0.0082 (-0.37-0.3)	0.2 (-3.6-3.4)
Solomon Islands	2000	Male	85 to 89	36% (20-53%)	64% (47-80%)	0.18 (0.082-0.33)	-0.076% (-0.49-0.23%)	-0.13% (-0.61-0.22%)	-0.051 (-0.24-0.089)	-0.26 (-1.7-0.77)
Solomon Islands	2000	Male	90 to 94	37% (23-55%)	63% (45-77%)	0.19 (0.083-0.36)	-0.075% (-0.46-0.2%)	-0.13% (-0.59-0.19%)	-0.019 (-0.084-0.026)	-0.062 (-0.39-0.17)
Solomon Islands	2000	Male	95 plus	41% (23-61%)	59% (39-77%)	0.21 (0.083-0.41)	-0.26% (-0.78-0.12%)	-0.39% (-0.99-0.046%)	-0.015 (-0.038-0.0017)	-0.039 (-0.12-0.018)
Solomon Islands	2005	Female	All Ages	19% (13-26%)	81% (74-87%)	0.19 (0.14-0.24)	0.024% (-0.33-0.34%)	-0.34% (-0.95-0.16%)	-6.2 (-18-2.9)	23 (-310-320)
Solomon Islands	2005	Female	15 to 19	20% (9.1-36%)	80% (64-91%)	0.23 (0.092-0.42)	0.64% (0.26-1.2%)	0.75% (0.13-1.8%)	0.25 (0.046-0.61)	28 (11-54)
Solomon Islands	2005	Female	20 to 24	25% (10-42%)	75% (58-90%)	0.35 (0.16-0.63)	1.1% (0.56-1.8%)	1.1% (0.35-2.3%)	0.51 (0.15-1.1)	55 (28-98)
Solomon Islands	2005	Female	25 to 29	26% (11-44%)	74% (56-89%)	0.36 (0.16-0.63)	1% (0.55-1.7%)	1% (0.36-1.9%)	0.66 (0.23-1.3)	60 (32-99)
Solomon Islands	2005	Female	30 to 34	27% (11-47%)	73% (53-89%)	0.38 (0.16-0.68)	0.73% (0.25-1.3%)	0.69% (0.054-1.5%)	0.5 (0.038-1.1)	42 (14-78)
Solomon Islands	2005	Female	35 to 39	28% (11-47%)	72% (53-89%)	0.39 (0.17-0.69)	0.61% (-0.016-1.3%)	0.65% (-0.14-1.6%)	0.55 (-0.13-1.3)	35 (-0.8-76)
Solomon Islands	2005	Female	40 to 44	27% (13-44%)	73% (56-87%)	0.36 (0.15-0.67)	0.27% (-0.53-1%)	0.3% (-0.7-1.3%)	0.28 (-0.65-1.3)	16 (-30-62)
Solomon Islands	2005	Female	45 to 49	26% (12-45%)	74% (55-88%)	0.31 (0.12-0.58)	-0.19% (-1.1-0.52%)	-0.19% (-1.3-0.65%)	-0.19 (-1.3-0.67)	-10 (-61-27)
Solomon Islands	2005	Female	50 to 54	26% (12-45%)	74% (55-88%)	0.27 (0.1-0.56)	-0.51% (-1.7-0.25%)	-0.51% (-1.8-0.38%)	-0.59 (-2.2-0.42)	-27 (-93-13)
Solomon Islands	2005	Female	55 to 59	25% (12-46%)	75% (54-88%)	0.25 (0.091-0.51)	-0.63% (-1.9-0.12%)	-0.62% (-2-0.22%)	-0.85 (-2.8-0.29)	-33 (-99-6)
Solomon Islands	2005	Female	60 to 64	26% (12-45%)	74% (55-88%)	0.24 (0.084-0.52)	-0.87% (-2.2--0.091%)	-0.89% (-2.3--0.041%)	-1.4 (-3.6--0.065)	-42 (-110--4)
Solomon Islands	2005	Female	65 to 69	27% (12-45%)	73% (55-88%)	0.24 (0.086-0.48)	-0.97% (-2.2--0.1%)	-1% (-2.3--0.048%)	-1.4 (-3.3--0.06)	-35 (-80--3.3)
Solomon Islands	2005	Female	70 to 74	27% (12-44%)	73% (56-88%)	0.23 (0.081-0.44)	-1% (-2.2--0.17%)	-1.1% (-2.3--0.16%)	-1.8 (-4.2--0.27)	-36 (-80--6.1)
Solomon Islands	2005	Female	75 to 79	25% (11-45%)	75% (55-89%)	0.21 (0.069-0.46)	-0.82% (-2--0.071%)	-0.87% (-2.2--0.055%)	-1.2 (-3--0.073)	-18 (-45--1.5)
Solomon Islands	2005	Female	80 to 84	24% (11-44%)	76% (56-89%)	0.2 (0.066-0.42)	-0.81% (-2--0.17%)	-0.88% (-2.2--0.16%)	-0.96 (-2.3--0.18)	-10 (-24--2.2)
Solomon Islands	2005	Female	85 to 89	25% (12-43%)	75% (57-88%)	0.21 (0.075-0.44)	-0.86% (-2--0.19%)	-0.95% (-2.2--0.21%)	-0.44 (-1--0.097)	-3.3 (-7.7--0.72)
Solomon Islands	2005	Female	90 to 94	29% (14-48%)	71% (52-86%)	0.24 (0.084-0.49)	-0.76% (-1.6--0.18%)	-0.86% (-1.8--0.2%)	-0.1 (-0.22--0.025)	-0.53 (-1.1--0.13)
Solomon Islands	2005	Female	95 plus	34% (15-57%)	66% (43-85%)	0.28 (0.094-0.59)	-1% (-2.3--0.15%)	-1.2% (-2.6--0.21%)	-0.033 (-0.073--0.0059)	-0.11 (-0.24--0.016)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	2005	Male	All Ages	39% (31-48%)	61% (52-69%)	0.4 (0.32-0.49)	1.1% (0.59-1.8%)	0.92% (0.24-1.7%)	20 (5.5-36)	1200 (640-1900)
Solomon Islands	2005	Male	15 to 19	37% (22-56%)	63% (44-78%)	0.27 (0.12-0.46)	1.2% (0.59-2.2%)	1.4% (0.44-2.8%)	0.72 (0.23-1.5)	68 (32-120)
Solomon Islands	2005	Male	20 to 24	55% (37-72%)	45% (28-63%)	0.71 (0.42-1.1)	3.4% (1.9-5.4%)	3.8% (1.8-6.4%)	2.8 (1.3-4.7)	220 (120-350)
Solomon Islands	2005	Male	25 to 29	63% (43-79%)	37% (21-57%)	1 (0.63-1.4)	4.1% (2.3-6.1%)	4.6% (2.4-7.1%)	3.7 (1.9-6.1)	270 (150-410)
Solomon Islands	2005	Male	30 to 34	64% (43-80%)	36% (20-57%)	1.1 (0.65-1.6)	3.5% (1.9-5.4%)	3.7% (1.7-6.2%)	3.3 (1.5-5.6)	220 (120-350)
Solomon Islands	2005	Male	35 to 39	62% (43-78%)	38% (22-57%)	0.93 (0.55-1.4)	2.6% (1.2-4.2%)	2.6% (0.97-4.7%)	2.5 (0.9-4.7)	150 (72-270)
Solomon Islands	2005	Male	40 to 44	59% (41-76%)	41% (24-59%)	0.81 (0.44-1.3)	1.8% (0.5-3.4%)	1.7% (0.24-3.7%)	2 (0.27-4.3)	110 (32-220)
Solomon Islands	2005	Male	45 to 49	56% (34-74%)	44% (26-66%)	0.69 (0.34-1.2)	1.3% (0.019-2.9%)	1.3% (-0.28-3.1%)	1.6 (-0.35-3.9)	80 (1.5-170)
Solomon Islands	2005	Male	50 to 54	53% (33-72%)	47% (28-67%)	0.57 (0.29-0.99)	1% (-0.21-2.3%)	0.96% (-0.47-2.5%)	1.3 (-0.58-3.5)	58 (-13-140)
Solomon Islands	2005	Male	55 to 59	51% (33-70%)	49% (30-67%)	0.47 (0.22-0.83)	0.5% (-0.44-1.6%)	0.45% (-0.65-1.7%)	0.69 (-0.99-2.7)	28 (-25-90)
Solomon Islands	2005	Male	60 to 64	48% (29-68%)	52% (32-71%)	0.39 (0.19-0.71)	0.3% (-0.46-1.2%)	0.27% (-0.58-1.3%)	0.48 (-1.1-2.4)	17 (-26-69)
Solomon Islands	2005	Male	65 to 69	46% (27-65%)	54% (35-73%)	0.36 (0.18-0.62)	0.25% (-0.44-1.1%)	0.24% (-0.53-1.1%)	0.4 (-0.89-1.9)	10 (-19-45)
Solomon Islands	2005	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.31 (0.15-0.56)	0.1% (-0.49-0.8%)	0.087% (-0.57-0.87%)	0.15 (-0.99-1.5)	3.6 (-17-29)
Solomon Islands	2005	Male	75 to 79	39% (23-59%)	61% (41-77%)	0.27 (0.12-0.49)	0.032% (-0.51-0.63%)	0.0023% (-0.62-0.65%)	0.0032 (-0.84-0.91)	0.71 (-11-13)
Solomon Islands	2005	Male	80 to 84	37% (21-55%)	63% (45-79%)	0.24 (0.12-0.42)	0.035% (-0.41-0.44%)	0.0037% (-0.5-0.45%)	0.0038 (-0.5-0.47)	0.42 (-4.9-5.3)
Solomon Islands	2005	Male	85 to 89	35% (20-52%)	65% (48-80%)	0.22 (0.11-0.4)	-0.065% (-0.5-0.31%)	-0.12% (-0.61-0.3%)	-0.058 (-0.31-0.15)	-0.27 (-2.1-1.3)
Solomon Islands	2005	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.23 (0.11-0.43)	-0.081% (-0.49-0.25%)	-0.15% (-0.64-0.23%)	-0.024 (-0.11-0.038)	-0.08 (-0.49-0.23)
Solomon Islands	2005	Male	95 plus	40% (21-61%)	60% (39-79%)	0.25 (0.1-0.51)	-0.27% (-0.83-0.13%)	-0.41% (-1.1-0.052%)	-0.018 (-0.048-0.0023)	-0.047 (-0.14-0.023)
Solomon Islands	2010	Female	All Ages	19% (13-26%)	81% (74-87%)	0.23 (0.17-0.3)	0.11% (-0.3-0.51%)	-0.29% (-0.97-0.34%)	-5.9 (-20-6.8)	110 (-300-530)
Solomon Islands	2010	Female	15 to 19	20% (9-36%)	80% (64-91%)	0.27 (0.11-0.53)	0.78% (0.32-1.5%)	1% (0.25-2.3%)	0.36 (0.086-0.88)	37 (15-75)
Solomon Islands	2010	Female	20 to 24	24% (10-42%)	76% (58-90%)	0.42 (0.18-0.78)	1.3% (0.68-2.2%)	1.4% (0.5-3%)	0.68 (0.23-1.5)	69 (35-120)
Solomon Islands	2010	Female	25 to 29	25% (11-44%)	75% (56-89%)	0.43 (0.19-0.76)	1.3% (0.68-2.1%)	1.4% (0.55-2.5%)	0.89 (0.34-1.7)	77 (41-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	2010	Female	30 to 34	26% (10-46%)	74% (54-90%)	0.45 (0.19-0.81)	0.96% (0.36-1.8%)	1% (0.19-2.2%)	0.85 (0.15-1.9)	66 (24-130)
Solomon Islands	2010	Female	35 to 39	27% (12-46%)	73% (54-88%)	0.47 (0.21-0.82)	0.86% (0.12-1.8%)	0.99% (0.043-2.2%)	1.1 (0.041-2.4)	65 (7.9-130)
Solomon Islands	2010	Female	40 to 44	26% (13-44%)	74% (56-87%)	0.44 (0.19-0.8)	0.49% (-0.41-1.5%)	0.58% (-0.52-1.9%)	0.6 (-0.54-1.9)	31 (-27-93)
Solomon Islands	2010	Female	45 to 49	26% (11-45%)	74% (55-89%)	0.38 (0.15-0.73)	-0.049% (-1.1-0.85%)	-0.022% (-1.2-1.2%)	-0.027 (-1.5-1.3)	-3 (-68-53)
Solomon Islands	2010	Female	50 to 54	26% (12-46%)	74% (54-88%)	0.34 (0.13-0.67)	-0.45% (-1.7-0.49%)	-0.45% (-1.9-0.66%)	-0.57 (-2.5-0.84)	-26 (-100-28)
Solomon Islands	2010	Female	55 to 59	25% (11-45%)	75% (55-89%)	0.3 (0.11-0.61)	-0.61% (-2.1-0.29%)	-0.6% (-2.2-0.48%)	-0.82 (-3.1-0.66)	-32 (-110-15)
Solomon Islands	2010	Female	60 to 64	25% (11-44%)	75% (56-89%)	0.29 (0.1-0.6)	-0.89% (-2.3-0.022%)	-0.92% (-2.5-0.11%)	-1.5 (-4.2-0.18)	-46 (-120-1.3)
Solomon Islands	2010	Female	65 to 69	26% (12-44%)	74% (56-88%)	0.29 (0.11-0.56)	-0.98% (-2.3--0.049%)	-1% (-2.5-0.027%)	-1.9 (-4.7-0.05)	-48 (-110--2.1)
Solomon Islands	2010	Female	70 to 74	26% (12-44%)	74% (56-88%)	0.28 (0.099-0.56)	-1% (-2.3--0.11%)	-1.1% (-2.5-0.099%)	-1.8 (-4.3--0.16)	-36 (-83--4.3)
Solomon Islands	2010	Female	75 to 79	25% (11-43%)	75% (57-89%)	0.26 (0.085-0.54)	-0.83% (-2.1--0.041%)	-0.88% (-2.2-0.011%)	-1.7 (-4.2-0.022)	-25 (-61--1.1)
Solomon Islands	2010	Female	80 to 84	24% (11-44%)	76% (56-89%)	0.25 (0.084-0.53)	-0.83% (-2--0.12%)	-0.91% (-2.2--0.13%)	-1.3 (-3.1--0.17)	-14 (-33--2.1)
Solomon Islands	2010	Female	85 to 89	25% (12-44%)	75% (56-88%)	0.26 (0.095-0.56)	-0.88% (-2--0.17%)	-0.98% (-2.3--0.2%)	-0.54 (-1.2--0.11)	-4 (-9.4-0.79)
Solomon Islands	2010	Female	90 to 94	29% (14-48%)	71% (52-86%)	0.29 (0.1-0.62)	-0.78% (-1.7--0.14%)	-0.88% (-2--0.15%)	-0.14 (-0.31--0.024)	-0.72 (-1.6--0.13)
Solomon Islands	2010	Female	95 plus	34% (15-56%)	66% (44-85%)	0.35 (0.11-0.73)	-1% (-2.4--0.099%)	-1.2% (-2.7--0.15%)	-0.038 (-0.085--0.0045)	-0.12 (-0.28--0.012)
Solomon Islands	2010	Male	All Ages	39% (31-49%)	61% (51-69%)	0.5 (0.4-0.62)	1.5% (0.83-2.2%)	1.2% (0.45-2.1%)	29 (11-50)	1700 (980-2600)
Solomon Islands	2010	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.33 (0.16-0.56)	1.5% (0.71-2.6%)	1.8% (0.65-3.5%)	0.95 (0.34-1.9)	86 (40-150)
Solomon Islands	2010	Male	20 to 24	55% (36-72%)	45% (28-64%)	0.89 (0.53-1.3)	4.1% (2.4-6.3%)	4.7% (2.4-7.6%)	3.4 (1.7-5.7)	260 (150-410)
Solomon Islands	2010	Male	25 to 29	62% (43-78%)	38% (22-57%)	1.2 (0.77-1.8)	5% (3-7.4%)	5.7% (3.1-8.8%)	4.6 (2.5-7.2)	320 (190-490)
Solomon Islands	2010	Male	30 to 34	63% (42-79%)	37% (21-58%)	1.3 (0.8-2)	4.2% (2.4-6.4%)	4.7% (2.4-7.4%)	4.6 (2.3-7.7)	300 (170-480)
Solomon Islands	2010	Male	35 to 39	62% (44-79%)	38% (21-56%)	1.1 (0.66-1.8)	3.2% (1.7-5.2%)	3.4% (1.5-5.9%)	4.1 (1.8-7.1)	240 (120-400)
Solomon Islands	2010	Male	40 to 44	59% (41-76%)	41% (24-59%)	1 (0.54-1.6)	2.3% (0.76-4.2%)	2.4% (0.49-4.6%)	3 (0.62-6.1)	160 (53-300)
Solomon Islands	2010	Male	45 to 49	56% (34-73%)	44% (27-66%)	0.86 (0.44-1.5)	1.8% (0.23-3.5%)	1.8% (-0.047-3.9%)	2.6 (-0.073-5.7)	130 (16-260)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	2010	Male	50 to 54	53% (34-71%)	47% (29-66%)	0.71 (0.36-1.3)	1.4% (0.005-3.1%)	1.4% (-0.27-3.3%)	2.2 (-0.45-5.4)	93 (0.35-220)
Solomon Islands	2010	Male	55 to 59	50% (32-70%)	50% (30-68%)	0.58 (0.28-1)	0.75% (-0.33-2%)	0.72% (-0.55-2.2%)	1.2 (-0.95-3.7)	45 (-21-130)
Solomon Islands	2010	Male	60 to 64	48% (28-67%)	52% (33-72%)	0.49 (0.23-0.9)	0.5% (-0.41-1.8%)	0.5% (-0.56-1.9%)	0.9 (-0.99-3.6)	28 (-23-110)
Solomon Islands	2010	Male	65 to 69	46% (28-64%)	54% (36-72%)	0.44 (0.21-0.77)	0.42% (-0.36-1.5%)	0.43% (-0.45-1.6%)	0.87 (-0.95-3.3)	22 (-19-77)
Solomon Islands	2010	Male	70 to 74	43% (27-61%)	57% (39-73%)	0.38 (0.18-0.7)	0.22% (-0.44-1%)	0.21% (-0.53-1.1%)	0.4 (-0.97-2.1)	8.4 (-17-41)
Solomon Islands	2010	Male	75 to 79	40% (24-58%)	60% (42-76%)	0.33 (0.15-0.61)	0.11% (-0.48-0.83%)	0.089% (-0.58-0.86%)	0.16 (-1.1-1.6)	3.3 (-14-24)
Solomon Islands	2010	Male	80 to 84	37% (21-55%)	63% (45-79%)	0.29 (0.14-0.51)	0.093% (-0.4-0.61%)	0.064% (-0.49-0.65%)	0.084 (-0.66-0.84)	1.4 (-6.3-9.4)
Solomon Islands	2010	Male	85 to 89	35% (20-52%)	65% (48-80%)	0.28 (0.13-0.5)	-0.017% (-0.46-0.43%)	-0.067% (-0.58-0.45%)	-0.037 (-0.32-0.25)	-0.081 (-2.2-2.1)
Solomon Islands	2010	Male	90 to 94	37% (22-54%)	63% (46-78%)	0.28 (0.13-0.53)	-0.058% (-0.51-0.31%)	-0.13% (-0.65-0.3%)	-0.023 (-0.12-0.056)	-0.063 (-0.55-0.35)
Solomon Islands	2010	Male	95 plus	40% (21-61%)	60% (39-79%)	0.31 (0.12-0.64)	-0.27% (-0.84-0.19%)	-0.41% (-1.1-0.12%)	-0.019 (-0.052-0.0053)	-0.048 (-0.15-0.036)
Solomon Islands	2016	Female	All Ages	19% (13-26%)	81% (74-87%)	0.27 (0.18-0.4)	0.16% (-0.35-0.72%)	-0.25% (-1.1-0.64%)	-5.5 (-23-13)	170 (-380-760)
Solomon Islands	2016	Female	15 to 19	20% (9-35%)	80% (65-91%)	0.31 (0.13-0.63)	0.87% (0.39-1.8%)	1.2% (0.33-2.8%)	0.44 (0.12-1.1)	45 (20-91)
Solomon Islands	2016	Female	20 to 24	24% (9.9-43%)	76% (57-90%)	0.49 (0.21-0.96)	1.5% (0.77-2.6%)	1.7% (0.58-3.6%)	0.81 (0.27-1.8)	82 (41-150)
Solomon Islands	2016	Female	25 to 29	26% (11-45%)	74% (55-89%)	0.51 (0.2-0.95)	1.5% (0.78-2.5%)	1.6% (0.64-3.2%)	0.98 (0.33-2.1)	84 (44-150)
Solomon Islands	2016	Female	30 to 34	27% (10-47%)	73% (53-90%)	0.53 (0.21-1)	1.1% (0.43-2.2%)	1.3% (0.29-2.9%)	0.98 (0.21-2.3)	74 (26-150)
Solomon Islands	2016	Female	35 to 39	27% (11-47%)	73% (53-89%)	0.54 (0.23-0.99)	1% (0.17-2.2%)	1.2% (0.083-2.7%)	1.4 (0.091-3.3)	84 (14-180)
Solomon Islands	2016	Female	40 to 44	26% (13-44%)	74% (56-87%)	0.5 (0.21-0.96)	0.63% (-0.42-1.9%)	0.78% (-0.52-2.5%)	1 (-0.72-3.3)	52 (-37-170)
Solomon Islands	2016	Female	45 to 49	26% (11-45%)	74% (55-89%)	0.43 (0.17-0.84)	0.06% (-1.1-1.2%)	0.13% (-1.3-1.6%)	0.17 (-1.7-2.2)	4.4 (-80-88)
Solomon Islands	2016	Female	50 to 54	26% (12-44%)	74% (56-88%)	0.38 (0.14-0.77)	-0.37% (-1.6-0.82%)	-0.34% (-1.8-1.1%)	-0.5 (-2.7-1.8)	-25 (-120-59)
Solomon Islands	2016	Female	55 to 59	25% (12-45%)	75% (55-88%)	0.34 (0.13-0.71)	-0.56% (-2-0.5%)	-0.53% (-2.2-0.74%)	-0.88 (-3.6-1.2)	-36 (-130-32)
Solomon Islands	2016	Female	60 to 64	25% (11-46%)	75% (54-89%)	0.33 (0.11-0.71)	-0.88% (-2.4-0.19%)	-0.9% (-2.6-0.31%)	-1.5 (-4.4-0.51)	-48 (-130-11)
Solomon Islands	2016	Female	65 to 69	27% (12-45%)	73% (55-88%)	0.34 (0.12-0.68)	-0.97% (-2.4-0.1%)	-1% (-2.7-0.22%)	-2 (-5.4-0.4)	-51 (-130-5.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	2016	Female	70 to 74	26% (12-44%)	74% (56-88%)	0.32 (0.11-0.67)	-1% (-2.3-0.11%)	-1.1% (-2.5-0.18%)	-2.4 (-5.9-0.37)	-48 (-110-4.6)
Solomon Islands	2016	Female	75 to 79	25% (11-44%)	75% (56-89%)	0.3 (0.093-0.67)	-0.79% (-2.1-0.17%)	-0.84% (-2.2-0.25%)	-1.5 (-4.2-0.43)	-23 (-62-5.2)
Solomon Islands	2016	Female	80 to 84	24% (11-43%)	76% (57-89%)	0.28 (0.091-0.62)	-0.8% (-2-0.03%)	-0.88% (-2.2-0.072%)	-1.4 (-3.5-0.11)	-15 (-37-0.57)
Solomon Islands	2016	Female	85 to 89	25% (12-45%)	75% (55-88%)	0.3 (0.1-0.64)	-0.85% (-2.1--0.066%)	-0.95% (-2.3--0.069%)	-0.79 (-1.9--0.058)	-5.9 (-14--0.47)
Solomon Islands	2016	Female	90 to 94	29% (13-48%)	71% (52-87%)	0.34 (0.12-0.71)	-0.76% (-1.7-0.005%)	-0.86% (-2-0.02%)	-0.18 (-0.41-0.0041)	-0.91 (-2.1-0.006)
Solomon Islands	2016	Female	95 plus	34% (14-56%)	66% (44-86%)	0.4 (0.12-0.87)	-1% (-2.5-0.03%)	-1.2% (-2.8-0.027%)	-0.056 (-0.13--0.0013)	-0.18 (-0.42-0.0051)
Solomon Islands	2016	Male	All Ages	39% (31-48%)	61% (52-69%)	0.58 (0.41-0.8)	1.8% (0.92-2.9%)	1.5% (0.51-2.8%)	38 (13-71)	2200 (1100-3500)
Solomon Islands	2016	Male	15 to 19	37% (22-55%)	63% (45-78%)	0.38 (0.17-0.67)	1.6% (0.79-2.9%)	2% (0.73-3.9%)	1.2 (0.42-2.3)	110 (50-180)
Solomon Islands	2016	Male	20 to 24	55% (36-72%)	45% (28-64%)	1 (0.56-1.7)	4.5% (2.5-7.1%)	5.2% (2.6-8.8%)	4.3 (2.1-7.5)	340 (190-560)
Solomon Islands	2016	Male	25 to 29	62% (42-77%)	38% (23-58%)	1.4 (0.8-2.2)	5.5% (3.1-8.5%)	6.4% (3.2-10%)	5.2 (2.6-8.8)	370 (200-590)
Solomon Islands	2016	Male	30 to 34	63% (42-79%)	37% (21-58%)	1.5 (0.85-2.4)	4.7% (2.6-7.8%)	5.3% (2.6-9.2%)	5 (2.5-8.5)	330 (170-540)
Solomon Islands	2016	Male	35 to 39	62% (44-79%)	38% (21-56%)	1.3 (0.72-2.1)	3.7% (1.8-6.2%)	4.1% (1.7-7.2%)	4.9 (2-9.1)	290 (140-510)
Solomon Islands	2016	Male	40 to 44	59% (41-76%)	41% (24-59%)	1.1 (0.61-2)	2.7% (0.94-5.4%)	2.8% (0.72-6.1%)	4.5 (1.1-9.7)	240 (83-490)
Solomon Islands	2016	Male	45 to 49	55% (33-73%)	45% (27-67%)	0.98 (0.47-1.8)	2.1% (0.38-4.4%)	2.2% (0.12-4.9%)	3.7 (0.19-8.6)	180 (33-390)
Solomon Islands	2016	Male	50 to 54	52% (32-71%)	48% (29-68%)	0.82 (0.38-1.5)	1.6% (0.12-3.8%)	1.7% (-0.16-4.1%)	3.1 (-0.29-8.1)	130 (9.1-320)
Solomon Islands	2016	Male	55 to 59	50% (31-69%)	50% (31-69%)	0.66 (0.31-1.2)	0.93% (-0.31-2.6%)	0.92% (-0.5-2.8%)	1.8 (-0.95-6)	69 (-21-210)
Solomon Islands	2016	Male	60 to 64	48% (28-68%)	52% (32-72%)	0.54 (0.25-1.1)	0.65% (-0.33-2.1%)	0.65% (-0.45-2.3%)	1.3 (-0.88-4.8)	41 (-21-140)
Solomon Islands	2016	Male	65 to 69	46% (27-65%)	54% (35-73%)	0.5 (0.23-0.91)	0.57% (-0.29-1.8%)	0.59% (-0.38-2%)	1.2 (-0.79-4.2)	30 (-15-100)
Solomon Islands	2016	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.44 (0.2-0.84)	0.34% (-0.43-1.5%)	0.35% (-0.52-1.6%)	0.76 (-1.1-3.4)	16 (-19-66)
Solomon Islands	2016	Male	75 to 79	39% (23-58%)	61% (42-77%)	0.38 (0.17-0.74)	0.22% (-0.41-1.1%)	0.2% (-0.51-1.2%)	0.36 (-0.9-2.2)	6.3 (-12-32)
Solomon Islands	2016	Male	80 to 84	36% (20-54%)	64% (46-80%)	0.33 (0.15-0.63)	0.17% (-0.35-0.8%)	0.15% (-0.43-0.85%)	0.21 (-0.59-1.2)	2.9 (-5.8-14)
Solomon Islands	2016	Male	85 to 89	35% (20-51%)	65% (49-80%)	0.32 (0.14-0.61)	0.048% (-0.45-0.63%)	0.0019% (-0.56-0.66%)	0.0014 (-0.43-0.5)	0.31 (-2.9-4.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Solomon Islands	2016	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.33 (0.14-0.64)	-0.019% (-0.51-0.45%)	-0.089% (-0.66-0.46%)	-0.019 (-0.14-0.097)	-0.024 (-0.65-0.58)
Solomon Islands	2016	Male	95 plus	40% (22-61%)	60% (39-78%)	0.37 (0.14-0.74)	-0.23% (-0.87-0.37%)	-0.38% (-1.2-0.28%)	-0.019 (-0.058-0.015)	-0.046 (-0.17-0.074)
Tonga	1990	Female	All Ages	17% (12-24%)	83% (76-88%)	0.14 (0.099-0.19)	-0.044% (-0.32-0.19%)	-0.42% (-0.91--0.014%)	-1.2 (-2.8--0.047)	-6.2 (-46-27)
Tonga	1990	Female	15 to 19	18% (8.2-33%)	82% (67-92%)	0.17 (0.072-0.34)	0.48% (0.25-0.77%)	0.4% (0.075-1%)	0.016 (0.0027-0.039)	3.3 (1.7-5.4)
Tonga	1990	Female	20 to 24	23% (9.7-41%)	77% (59-90%)	0.27 (0.12-0.49)	0.87% (0.53-1.3%)	0.58% (0.19-1.2%)	0.034 (0.011-0.074)	7.6 (4.5-12)
Tonga	1990	Female	25 to 29	24% (10-44%)	76% (56-90%)	0.28 (0.12-0.51)	0.84% (0.54-1.2%)	0.59% (0.21-1.2%)	0.03 (0.011-0.066)	5.4 (3.2-8.1)
Tonga	1990	Female	30 to 34	25% (9.9-45%)	75% (55-90%)	0.29 (0.11-0.53)	0.7% (0.41-1.1%)	0.58% (0.21-1.1%)	0.03 (0.011-0.06)	3.9 (2.1-5.9)
Tonga	1990	Female	35 to 39	25% (10-43%)	75% (57-90%)	0.3 (0.12-0.56)	0.58% (0.25-0.97%)	0.62% (0.19-1.2%)	0.046 (0.014-0.097)	3.8 (1.6-6.5)
Tonga	1990	Female	40 to 44	25% (12-42%)	75% (58-88%)	0.27 (0.11-0.52)	0.39% (-0.067-0.84%)	0.52% (-0.048-1.2%)	0.045 (-0.0037-0.11)	2.6 (-0.41-5.8)
Tonga	1990	Female	45 to 49	24% (11-42%)	76% (58-89%)	0.24 (0.088-0.47)	0.023% (-0.63-0.49%)	0.13% (-0.62-0.76%)	0.013 (-0.067-0.076)	0.15 (-4.1-3.3)
Tonga	1990	Female	50 to 54	24% (11-43%)	76% (57-89%)	0.21 (0.076-0.42)	-0.33% (-1.2-0.26%)	-0.27% (-1.3-0.4%)	-0.038 (-0.19-0.055)	-2.5 (-9.7-1.9)
Tonga	1990	Female	55 to 59	23% (10-42%)	77% (58-90%)	0.19 (0.067-0.41)	-0.52% (-1.6-0.12%)	-0.46% (-1.7-0.25%)	-0.091 (-0.35-0.051)	-4.5 (-14-1.1)
Tonga	1990	Female	60 to 64	24% (11-43%)	76% (57-89%)	0.18 (0.063-0.38)	-0.72% (-2-0.019%)	-0.71% (-2.1-0.09%)	-0.18 (-0.53-0.023)	-6.7 (-18-0.16)
Tonga	1990	Female	65 to 69	24% (11-41%)	76% (59-89%)	0.18 (0.065-0.37)	-0.87% (-2--0.11%)	-0.91% (-2.2--0.059%)	-0.25 (-0.63--0.017)	-7 (-17--0.9)
Tonga	1990	Female	70 to 74	24% (12-40%)	76% (60-88%)	0.17 (0.058-0.36)	-0.86% (-1.8--0.16%)	-0.92% (-2--0.15%)	-0.28 (-0.64--0.045)	-5.9 (-13--1.1)
Tonga	1990	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.16 (0.051-0.35)	-0.6% (-1.6--0.056%)	-0.62% (-1.7--0.027%)	-0.18 (-0.51--0.0074)	-3 (-8.1--0.28)
Tonga	1990	Female	80 to 84	22% (9.2-42%)	78% (58-91%)	0.15 (0.045-0.34)	-0.6% (-1.6--0.079%)	-0.64% (-1.7--0.076%)	-0.19 (-0.5--0.021)	-2.2 (-5.7--0.28)
Tonga	1990	Female	85 to 89	23% (10-42%)	77% (58-90%)	0.16 (0.053-0.35)	-0.63% (-1.5--0.13%)	-0.68% (-1.7--0.13%)	-0.13 (-0.32--0.027)	-1.1 (-2.5--0.22)
Tonga	1990	Female	90 to 94	27% (12-47%)	73% (53-88%)	0.18 (0.059-0.39)	-0.68% (-1.5--0.15%)	-0.75% (-1.7--0.16%)	-0.064 (-0.14--0.014)	-0.34 (-0.76--0.075)
Tonga	1990	Female	95 plus	32% (13-53%)	68% (47-87%)	0.22 (0.067-0.49)	-0.83% (-2--0.052%)	-0.97% (-2.3--0.08%)	-0.021 (-0.049--0.0017)	-0.069 (-0.17--0.0043)
Tonga	1990	Male	All Ages	36% (28-45%)	64% (55-72%)	0.3 (0.22-0.39)	0.88% (0.51-1.3%)	0.62% (0.18-1.1%)	2.2 (0.67-3.8)	150 (89-230)
Tonga	1990	Male	15 to 19	35% (20-52%)	65% (48-80%)	0.2 (0.099-0.35)	1.1% (0.59-1.8%)	1.1% (0.43-2.2%)	0.096 (0.037-0.19)	10 (5.7-17)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	1990	Male	20 to 24	53% (34-71%)	47% (29-66%)	0.55 (0.33-0.84)	2.7% (1.6-4%)	2.8% (1.4-4.8%)	0.4 (0.18-0.68)	36 (21-55)
Tonga	1990	Male	25 to 29	60% (40-77%)	40% (23-60%)	0.77 (0.46-1.2)	3.3% (2-4.8%)	3.6% (1.9-5.6%)	0.38 (0.18-0.61)	31 (18-45)
Tonga	1990	Male	30 to 34	61% (40-78%)	39% (22-60%)	0.83 (0.48-1.3)	2.7% (1.6-3.9%)	2.8% (1.4-4.5%)	0.22 (0.1-0.38)	17 (10-27)
Tonga	1990	Male	35 to 39	60% (42-76%)	40% (24-58%)	0.73 (0.38-1.2)	2.2% (1.2-3.4%)	2.3% (0.98-3.9%)	0.23 (0.097-0.4)	16 (8.6-25)
Tonga	1990	Male	40 to 44	57% (39-74%)	43% (26-61%)	0.63 (0.3-1.1)	1.6% (0.85-2.6%)	1.7% (0.67-2.9%)	0.18 (0.062-0.33)	11 (5.7-19)
Tonga	1990	Male	45 to 49	53% (32-72%)	47% (28-68%)	0.54 (0.24-0.97)	1.5% (0.69-2.2%)	1.5% (0.53-2.6%)	0.18 (0.06-0.32)	9.7 (4.5-16)
Tonga	1990	Male	50 to 54	50% (31-69%)	50% (31-69%)	0.44 (0.2-0.79)	1.1% (0.35-1.9%)	1.1% (0.19-2.1%)	0.19 (0.03-0.37)	8.8 (2.7-16)
Tonga	1990	Male	55 to 59	48% (29-68%)	52% (32-71%)	0.36 (0.17-0.66)	0.63% (-0.091-1.3%)	0.61% (-0.27-1.4%)	0.14 (-0.06-0.33)	5.6 (-0.8-11)
Tonga	1990	Male	60 to 64	45% (27-65%)	55% (35-73%)	0.3 (0.13-0.57)	0.34% (-0.26-0.89%)	0.33% (-0.37-0.95%)	0.1 (-0.12-0.31)	3.5 (-2.7-9.4)
Tonga	1990	Male	65 to 69	44% (26-62%)	56% (38-74%)	0.27 (0.13-0.49)	0.23% (-0.26-0.69%)	0.23% (-0.36-0.76%)	0.076 (-0.12-0.25)	2 (-2.2-6.1)
Tonga	1990	Male	70 to 74	41% (24-58%)	59% (42-76%)	0.24 (0.1-0.44)	0.084% (-0.35-0.48%)	0.067% (-0.43-0.53%)	0.023 (-0.15-0.18)	0.6 (-2.6-3.6)
Tonga	1990	Male	75 to 79	37% (22-56%)	63% (44-78%)	0.21 (0.089-0.39)	-0.025% (-0.45-0.29%)	-0.068% (-0.55-0.28%)	-0.021 (-0.17-0.089)	-0.13 (-2.2-1.5)
Tonga	1990	Male	80 to 84	34% (19-53%)	66% (47-81%)	0.18 (0.082-0.33)	-0.001% (-0.36-0.26%)	-0.038% (-0.45-0.26%)	-0.0093 (-0.11-0.065)	-0.004 (-1.1-0.77)
Tonga	1990	Male	85 to 89	33% (19-49%)	67% (51-81%)	0.17 (0.081-0.31)	-0.081% (-0.45-0.17%)	-0.13% (-0.56-0.16%)	-0.02 (-0.082-0.024)	-0.1 (-0.57-0.21)
Tonga	1990	Male	90 to 94	34% (21-52%)	66% (48-79%)	0.18 (0.077-0.33)	-0.076% (-0.44-0.18%)	-0.13% (-0.57-0.17%)	-0.0077 (-0.032-0.0094)	-0.026 (-0.15-0.061)
Tonga	1990	Male	95 plus	38% (20-60%)	62% (40-80%)	0.2 (0.077-0.41)	-0.24% (-0.72-0.071%)	-0.38% (-0.96-0.00049%)	-0.0049 (-0.012-0.000063)	-0.012 (-0.036-0.0035)
Tonga	1995	Female	All Ages	18% (12-24%)	82% (76-88%)	0.17 (0.12-0.22)	-0.029% (-0.33-0.22%)	-0.4% (-0.96-0.042%)	-1.2 (-3-0.13)	-4.1 (-45-32)
Tonga	1995	Female	15 to 19	18% (8.3-33%)	82% (67-92%)	0.2 (0.086-0.38)	0.55% (0.3-0.89%)	0.56% (0.13-1.3%)	0.02 (0.0047-0.046)	3.5 (1.9-5.8)
Tonga	1995	Female	20 to 24	23% (9.7-40%)	77% (60-90%)	0.32 (0.14-0.58)	0.96% (0.58-1.4%)	0.74% (0.26-1.5%)	0.031 (0.011-0.067)	6.2 (3.6-9.5)
Tonga	1995	Female	25 to 29	24% (9.9-42%)	76% (58-90%)	0.33 (0.13-0.59)	0.94% (0.6-1.4%)	0.75% (0.31-1.4%)	0.042 (0.017-0.08)	6.7 (4.1-10)
Tonga	1995	Female	30 to 34	25% (8.9-44%)	75% (56-91%)	0.34 (0.14-0.61)	0.8% (0.45-1.2%)	0.74% (0.33-1.4%)	0.039 (0.017-0.073)	4.6 (2.5-7.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	1995	Female	35 to 39	25% (11-43%)	75% (57-89%)	0.35 (0.15-0.63)	0.7% (0.34-1.2%)	0.82% (0.35-1.5%)	0.059 (0.025-0.12)	4.5 (2.3-7.5)
Tonga	1995	Female	40 to 44	24% (11-41%)	76% (59-89%)	0.32 (0.13-0.6)	0.54% (0.082-1%)	0.76% (0.14-1.6%)	0.072 (0.012-0.15)	4 (0.57-7.7)
Tonga	1995	Female	45 to 49	24% (10-41%)	76% (59-90%)	0.28 (0.1-0.55)	0.16% (-0.46-0.7%)	0.35% (-0.41-1%)	0.037 (-0.044-0.11)	1.1 (-3.1-4.9)
Tonga	1995	Female	50 to 54	24% (11-42%)	76% (58-89%)	0.25 (0.091-0.48)	-0.21% (-1.1-0.37%)	-0.078% (-1.1-0.61%)	-0.011 (-0.16-0.084)	-1.6 (-8.2-2.8)
Tonga	1995	Female	55 to 59	23% (10-41%)	77% (59-90%)	0.22 (0.083-0.46)	-0.41% (-1.5-0.22%)	-0.3% (-1.5-0.42%)	-0.06 (-0.3-0.083)	-3.6 (-13-1.8)
Tonga	1995	Female	60 to 64	23% (11-42%)	77% (58-89%)	0.21 (0.077-0.44)	-0.67% (-1.9-0.079%)	-0.63% (-2-0.18%)	-0.16 (-0.51-0.045)	-6.4 (-18-0.7)
Tonga	1995	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.22 (0.081-0.43)	-0.86% (-2.1--0.068%)	-0.88% (-2.2--0.021%)	-0.29 (-0.73--0.0073)	-8.3 (-20--0.64)
Tonga	1995	Female	70 to 74	25% (11-42%)	75% (58-89%)	0.21 (0.073-0.42)	-0.9% (-2--0.14%)	-0.97% (-2.2--0.13%)	-0.33 (-0.76--0.04)	-7.2 (-16--1.1)
Tonga	1995	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.19 (0.063-0.43)	-0.62% (-1.6--0.028%)	-0.64% (-1.7-0.011%)	-0.22 (-0.61-0.0039)	-3.8 (-9.7--0.17)
Tonga	1995	Female	80 to 84	22% (9.8-42%)	78% (58-90%)	0.18 (0.059-0.38)	-0.61% (-1.5--0.091%)	-0.65% (-1.7--0.072%)	-0.23 (-0.58--0.024)	-2.7 (-6.5--0.39)
Tonga	1995	Female	85 to 89	24% (10-41%)	76% (59-90%)	0.19 (0.066-0.4)	-0.65% (-1.5--0.12%)	-0.71% (-1.7--0.13%)	-0.15 (-0.36--0.027)	-1.2 (-2.8--0.22)
Tonga	1995	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.21 (0.075-0.46)	-0.67% (-1.4--0.14%)	-0.74% (-1.6--0.15%)	-0.065 (-0.14--0.013)	-0.35 (-0.75--0.075)
Tonga	1995	Female	95 plus	32% (13-55%)	68% (45-87%)	0.26 (0.082-0.55)	-0.82% (-1.9--0.051%)	-0.96% (-2.3--0.09%)	-0.025 (-0.06--0.0023)	-0.084 (-0.2--0.0051)
Tonga	1995	Male	All Ages	36% (28-45%)	64% (55-72%)	0.35 (0.27-0.45)	0.98% (0.6-1.4%)	0.69% (0.23-1.1%)	2.5 (0.86-4.3)	170 (100-240)
Tonga	1995	Male	15 to 19	34% (20-52%)	66% (48-80%)	0.24 (0.11-0.42)	1.2% (0.7-2%)	1.4% (0.59-2.6%)	0.11 (0.048-0.21)	11 (6.6-19)
Tonga	1995	Male	20 to 24	52% (33-70%)	48% (30-67%)	0.66 (0.39-1)	3.1% (1.9-4.6%)	3.5% (1.8-5.6%)	0.36 (0.19-0.6)	31 (18-47)
Tonga	1995	Male	25 to 29	60% (40-76%)	40% (24-60%)	0.92 (0.57-1.4)	3.7% (2.4-5.2%)	4.1% (2.4-6.3%)	0.46 (0.25-0.72)	36 (22-53)
Tonga	1995	Male	30 to 34	61% (40-78%)	39% (22-60%)	0.99 (0.57-1.5)	3% (1.9-4.4%)	3.2% (1.7-5.2%)	0.28 (0.14-0.47)	21 (13-32)
Tonga	1995	Male	35 to 39	59% (41-77%)	41% (23-59%)	0.87 (0.49-1.4)	2.5% (1.5-3.7%)	2.7% (1.3-4.3%)	0.26 (0.12-0.45)	18 (10-28)
Tonga	1995	Male	40 to 44	57% (39-74%)	43% (26-61%)	0.76 (0.39-1.3)	1.9% (1-2.8%)	1.9% (0.83-3.2%)	0.24 (0.1-0.42)	15 (8.2-24)
Tonga	1995	Male	45 to 49	53% (32-71%)	47% (29-68%)	0.64 (0.31-1.1)	1.7% (0.78-2.6%)	1.8% (0.66-2.9%)	0.22 (0.078-0.38)	12 (5.4-19)
Tonga	1995	Male	50 to 54	50% (30-69%)	50% (31-70%)	0.52 (0.25-0.92)	1.2% (0.38-2.1%)	1.3% (0.24-2.3%)	0.21 (0.038-0.39)	9.7 (2.9-17)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	1995	Male	55 to 59	48% (30-67%)	52% (33-70%)	0.42 (0.21-0.73)	0.71% (-0.066-1.4%)	0.71% (-0.23-1.6%)	0.17 (-0.059-0.37)	6.7 (-0.63-14)
Tonga	1995	Male	60 to 64	45% (27-66%)	55% (34-73%)	0.35 (0.17-0.65)	0.41% (-0.27-0.98%)	0.4% (-0.39-1.1%)	0.12 (-0.11-0.33)	4.1 (-2.6-11)
Tonga	1995	Male	65 to 69	44% (26-63%)	56% (37-74%)	0.32 (0.16-0.56)	0.28% (-0.26-0.78%)	0.29% (-0.32-0.87%)	0.12 (-0.13-0.36)	3 (-2.9-8.6)
Tonga	1995	Male	70 to 74	41% (25-60%)	59% (40-75%)	0.29 (0.14-0.52)	0.11% (-0.39-0.55%)	0.096% (-0.46-0.6%)	0.037 (-0.19-0.23)	0.9 (-3.4-4.4)
Tonga	1995	Male	75 to 79	37% (22-55%)	63% (45-78%)	0.24 (0.11-0.45)	-0.015% (-0.43-0.34%)	-0.056% (-0.54-0.33%)	-0.019 (-0.19-0.12)	-0.082 (-2.5-1.9)
Tonga	1995	Male	80 to 84	34% (19-52%)	66% (48-81%)	0.22 (0.1-0.39)	0.0045% (-0.37-0.3%)	-0.032% (-0.45-0.3%)	-0.0089 (-0.12-0.081)	0.014 (-1.2-0.96)
Tonga	1995	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.2 (0.096-0.37)	-0.081% (-0.48-0.19%)	-0.13% (-0.6-0.18%)	-0.023 (-0.1-0.03)	-0.12 (-0.69-0.28)
Tonga	1995	Male	90 to 94	34% (21-52%)	66% (48-79%)	0.21 (0.093-0.39)	-0.081% (-0.47-0.2%)	-0.14% (-0.62-0.18%)	-0.0096 (-0.041-0.012)	-0.032 (-0.19-0.077)
Tonga	1995	Male	95 plus	38% (21-59%)	62% (41-79%)	0.24 (0.092-0.48)	-0.25% (-0.74-0.087%)	-0.4% (-0.99--0.0034%)	-0.0069 (-0.017--0.000061)	-0.017 (-0.051-0.006)
Tonga	2000	Female	All Ages	18% (12-24%)	82% (76-88%)	0.22 (0.16-0.29)	0.021% (-0.3-0.33%)	-0.36% (-0.97-0.15%)	-1.2 (-3.2-0.51)	3 (-46-46)
Tonga	2000	Female	15 to 19	18% (8.5-33%)	82% (67-91%)	0.27 (0.11-0.52)	0.62% (0.34-0.98%)	0.71% (0.19-1.6%)	0.027 (0.0073-0.059)	4.3 (2.4-6.9)
Tonga	2000	Female	20 to 24	22% (9.7-39%)	78% (61-90%)	0.42 (0.18-0.78)	1.1% (0.68-1.6%)	0.94% (0.34-1.9%)	0.041 (0.015-0.088)	7.2 (4.4-11)
Tonga	2000	Female	25 to 29	24% (9.5-42%)	76% (58-91%)	0.43 (0.18-0.76)	1.1% (0.69-1.5%)	0.96% (0.4-1.8%)	0.044 (0.018-0.084)	6.3 (3.9-9.5)
Tonga	2000	Female	30 to 34	25% (9.2-44%)	75% (56-91%)	0.45 (0.17-0.81)	0.93% (0.55-1.4%)	0.95% (0.45-1.8%)	0.058 (0.026-0.11)	6.3 (3.5-9.6)
Tonga	2000	Female	35 to 39	25% (10-43%)	75% (57-90%)	0.46 (0.19-0.83)	0.89% (0.44-1.5%)	1.1% (0.46-2.1%)	0.072 (0.028-0.14)	5.2 (2.6-9)
Tonga	2000	Female	40 to 44	24% (11-42%)	76% (58-89%)	0.43 (0.18-0.8)	0.73% (0.15-1.4%)	1.1% (0.31-2.2%)	0.097 (0.026-0.2)	5.2 (1.1-10)
Tonga	2000	Female	45 to 49	24% (10-41%)	76% (59-90%)	0.37 (0.14-0.7)	0.33% (-0.36-1%)	0.62% (-0.24-1.6%)	0.074 (-0.029-0.19)	2.6 (-2.9-7.9)
Tonga	2000	Female	50 to 54	24% (11-41%)	76% (59-89%)	0.32 (0.12-0.62)	-0.086% (-1-0.62%)	0.1% (-0.93-1%)	0.013 (-0.13-0.13)	-0.65 (-7.9-4.7)
Tonga	2000	Female	55 to 59	23% (10-42%)	77% (58-90%)	0.29 (0.11-0.59)	-0.33% (-1.4-0.37%)	-0.18% (-1.3-0.67%)	-0.033 (-0.25-0.11)	-2.7 (-12-3.1)
Tonga	2000	Female	60 to 64	23% (11-43%)	77% (57-89%)	0.28 (0.099-0.6)	-0.63% (-1.9-0.16%)	-0.57% (-2-0.32%)	-0.14 (-0.51-0.081)	-5.7 (-18-1.4)
Tonga	2000	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.28 (0.1-0.56)	-0.83% (-2.1-0.019%)	-0.84% (-2.3-0.12%)	-0.27 (-0.76-0.039)	-7.9 (-21-0.17)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	2000	Female	70 to 74	24% (11-42%)	76% (58-89%)	0.27 (0.098-0.53)	-0.88% (-2--0.11%)	-0.95% (-2.2--0.071%)	-0.39 (-0.94--0.034)	-8.4 (-20--1)
Tonga	2000	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.25 (0.079-0.52)	-0.59% (-1.5-0.028%)	-0.6% (-1.7-0.078%)	-0.24 (-0.68-0.031)	-4.2 (-11-0.21)
Tonga	2000	Female	80 to 84	22% (8.6-42%)	78% (58-91%)	0.23 (0.072-0.48)	-0.58% (-1.5--0.014%)	-0.62% (-1.6-0.013%)	-0.25 (-0.64-0.0058)	-3 (-7.6--0.073)
Tonga	2000	Female	85 to 89	23% (10-41%)	77% (59-90%)	0.25 (0.087-0.53)	-0.63% (-1.6--0.046%)	-0.69% (-1.7--0.039%)	-0.18 (-0.46--0.01)	-1.5 (-3.6--0.11)
Tonga	2000	Female	90 to 94	27% (11-46%)	73% (54-89%)	0.29 (0.094-0.62)	-0.63% (-1.5--0.049%)	-0.7% (-1.7--0.05%)	-0.074 (-0.18--0.0054)	-0.4 (-0.94--0.029)
Tonga	2000	Female	95 plus	32% (13-55%)	68% (45-87%)	0.34 (0.11-0.74)	-0.78% (-2-0.061%)	-0.92% (-2.3-0.049%)	-0.027 (-0.068-0.0013)	-0.089 (-0.23-0.0069)
Tonga	2000	Male	All Ages	36% (28-45%)	64% (55-72%)	0.48 (0.38-0.6)	1.2% (0.79-1.7%)	0.86% (0.39-1.5%)	3.2 (1.4-5.3)	200 (130-290)
Tonga	2000	Male	15 to 19	34% (19-52%)	66% (48-81%)	0.33 (0.16-0.56)	1.4% (0.79-2.3%)	1.7% (0.74-3%)	0.15 (0.068-0.29)	15 (8.5-25)
Tonga	2000	Male	20 to 24	52% (33-70%)	48% (30-67%)	0.88 (0.53-1.3)	3.6% (2.2-5.4%)	4.1% (2.2-6.6%)	0.45 (0.24-0.74)	38 (23-57)
Tonga	2000	Male	25 to 29	59% (39-75%)	41% (25-61%)	1.2 (0.76-1.8)	4.2% (2.7-5.9%)	4.8% (2.8-7.2%)	0.45 (0.24-0.71)	35 (22-52)
Tonga	2000	Male	30 to 34	61% (39-77%)	39% (23-61%)	1.3 (0.75-2)	3.3% (2.1-4.9%)	3.6% (2-5.7%)	0.39 (0.2-0.65)	29 (17-46)
Tonga	2000	Male	35 to 39	59% (41-76%)	41% (24-59%)	1.1 (0.64-1.8)	2.9% (1.8-4.3%)	3.1% (1.7-5%)	0.29 (0.14-0.48)	19 (11-30)
Tonga	2000	Male	40 to 44	57% (39-74%)	43% (26-61%)	1 (0.52-1.7)	2.2% (1.2-3.5%)	2.4% (1.1-4%)	0.28 (0.12-0.49)	17 (8.5-28)
Tonga	2000	Male	45 to 49	53% (32-71%)	47% (29-68%)	0.86 (0.43-1.6)	2% (0.99-3.2%)	2.2% (0.88-3.6%)	0.34 (0.14-0.6)	18 (8.6-30)
Tonga	2000	Male	50 to 54	50% (31-68%)	50% (32-69%)	0.71 (0.34-1.2)	1.5% (0.5-2.7%)	1.6% (0.33-3%)	0.25 (0.049-0.48)	11 (3.6-20)
Tonga	2000	Male	55 to 59	48% (30-66%)	52% (34-70%)	0.57 (0.28-0.99)	0.91% (0.11-1.8%)	0.93% (-0.072-2%)	0.2 (-0.015-0.45)	7.9 (0.9-16)
Tonga	2000	Male	60 to 64	45% (26-66%)	55% (34-74%)	0.47 (0.22-0.87)	0.56% (-0.2-1.3%)	0.57% (-0.31-1.5%)	0.18 (-0.092-0.47)	5.7 (-1.8-14)
Tonga	2000	Male	65 to 69	43% (25-62%)	57% (38-75%)	0.43 (0.22-0.74)	0.42% (-0.16-1.1%)	0.45% (-0.23-1.2%)	0.17 (-0.083-0.45)	4.2 (-1.7-11)
Tonga	2000	Male	70 to 74	41% (24-59%)	59% (41-76%)	0.38 (0.19-0.69)	0.21% (-0.35-0.81%)	0.21% (-0.39-0.87%)	0.096 (-0.18-0.39)	2 (-3.1-7.6)
Tonga	2000	Male	75 to 79	37% (21-56%)	63% (44-79%)	0.33 (0.15-0.61)	0.056% (-0.42-0.53%)	0.02% (-0.5-0.56%)	0.0077 (-0.19-0.2)	0.34 (-2.5-3.2)
Tonga	2000	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.29 (0.13-0.51)	0.058% (-0.36-0.44%)	0.024% (-0.44-0.44%)	0.0071 (-0.13-0.13)	0.21 (-1.3-1.6)
Tonga	2000	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.27 (0.13-0.51)	-0.04% (-0.46-0.3%)	-0.094% (-0.58-0.29%)	-0.018 (-0.11-0.056)	-0.065 (-0.75-0.49)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	2000	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.28 (0.12-0.52)	-0.054% (-0.45-0.25%)	-0.12% (-0.6-0.23%)	-0.0092 (-0.045-0.017)	-0.025 (-0.21-0.11)
Tonga	2000	Male	95 plus	38% (21-59%)	62% (41-79%)	0.32 (0.13-0.61)	-0.23% (-0.74-0.16%)	-0.39% (-1-0.076%)	-0.0076 (-0.02-0.0015)	-0.018 (-0.057-0.013)
Tonga	2005	Female	All Ages	18% (13-24%)	82% (76-87%)	0.26 (0.19-0.34)	0.057% (-0.31-0.39%)	-0.34% (-1-0.22%)	-1.2 (-3.3-0.76)	8.2 (-44-57)
Tonga	2005	Female	15 to 19	18% (8.3-33%)	82% (67-92%)	0.3 (0.12-0.59)	0.66% (0.36-1.1%)	0.82% (0.24-1.9%)	0.025 (0.0072-0.056)	3.9 (2-6.7)
Tonga	2005	Female	20 to 24	22% (9.1-41%)	78% (59-91%)	0.47 (0.19-0.9)	1.2% (0.7-1.7%)	1.1% (0.39-2.2%)	0.045 (0.017-0.093)	8 (4.7-12)
Tonga	2005	Female	25 to 29	24% (9.6-43%)	76% (57-90%)	0.48 (0.2-0.88)	1.1% (0.72-1.7%)	1.1% (0.44-2.2%)	0.048 (0.019-0.095)	6.9 (4.2-11)
Tonga	2005	Female	30 to 34	25% (9.1-44%)	75% (56-91%)	0.5 (0.2-0.91)	1% (0.59-1.5%)	1.1% (0.49-2%)	0.062 (0.026-0.12)	6.7 (3.7-10)
Tonga	2005	Female	35 to 39	25% (10-43%)	75% (57-90%)	0.52 (0.21-0.91)	0.98% (0.51-1.6%)	1.3% (0.59-2.4%)	0.1 (0.045-0.2)	7.6 (3.9-13)
Tonga	2005	Female	40 to 44	24% (12-41%)	76% (59-88%)	0.48 (0.2-0.88)	0.82% (0.21-1.6%)	1.2% (0.41-2.4%)	0.11 (0.034-0.21)	5.7 (1.5-11)
Tonga	2005	Female	45 to 49	24% (10-41%)	76% (59-90%)	0.42 (0.15-0.79)	0.38% (-0.42-1.1%)	0.75% (-0.2-1.9%)	0.09 (-0.023-0.22)	3.1 (-3.5-9.5)
Tonga	2005	Female	50 to 54	24% (11-41%)	76% (59-89%)	0.36 (0.14-0.72)	-0.078% (-1.1-0.78%)	0.17% (-1.1-1.2%)	0.022 (-0.14-0.16)	-0.6 (-8.8-5.7)
Tonga	2005	Female	55 to 59	23% (10-42%)	77% (58-90%)	0.33 (0.12-0.68)	-0.31% (-1.4-0.49%)	-0.12% (-1.3-0.91%)	-0.019 (-0.21-0.13)	-2.3 (-10-3.4)
Tonga	2005	Female	60 to 64	23% (10-41%)	77% (59-90%)	0.32 (0.11-0.67)	-0.61% (-2-0.24%)	-0.53% (-2-0.47%)	-0.11 (-0.41-0.099)	-4.8 (-15-2)
Tonga	2005	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.32 (0.12-0.64)	-0.82% (-2.2-0.085%)	-0.82% (-2.4-0.24%)	-0.23 (-0.66-0.072)	-7.1 (-18-0.84)
Tonga	2005	Female	70 to 74	25% (11-41%)	75% (59-89%)	0.31 (0.11-0.61)	-0.88% (-2.1--0.071%)	-0.94% (-2.3--0.011%)	-0.37 (-0.9--0.0048)	-8.2 (-19--0.65)
Tonga	2005	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.29 (0.095-0.62)	-0.59% (-1.6-0.07%)	-0.6% (-1.8-0.14%)	-0.29 (-0.85-0.068)	-5.1 (-14-0.67)
Tonga	2005	Female	80 to 84	23% (9.7-41%)	77% (59-90%)	0.27 (0.094-0.56)	-0.58% (-1.5--0.0038%)	-0.62% (-1.7-0.052%)	-0.3 (-0.83-0.023)	-3.6 (-9.7--0.024)
Tonga	2005	Female	85 to 89	23% (11-41%)	77% (59-89%)	0.28 (0.11-0.59)	-0.62% (-1.6--0.033%)	-0.69% (-1.8--0.028%)	-0.22 (-0.57--0.0091)	-1.8 (-4.5--0.097)
Tonga	2005	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.32 (0.11-0.7)	-0.63% (-1.5--0.015%)	-0.7% (-1.7--0.012%)	-0.096 (-0.23--0.0015)	-0.52 (-1.2--0.013)
Tonga	2005	Female	95 plus	32% (13-54%)	68% (46-87%)	0.38 (0.13-0.84)	-0.79% (-2-0.12%)	-0.95% (-2.4-0.1%)	-0.034 (-0.088-0.0039)	-0.11 (-0.3-0.017)
Tonga	2005	Male	All Ages	37% (29-46%)	63% (54-71%)	0.56 (0.44-0.71)	1.3% (0.87-1.9%)	0.99% (0.46-1.6%)	3.9 (1.7-6.7)	230 (150-340)
Tonga	2005	Male	15 to 19	34% (19-52%)	66% (48-81%)	0.36 (0.16-0.63)	1.5% (0.85-2.5%)	1.8% (0.82-3.3%)	0.14 (0.063-0.26)	14 (7.4-22)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	2005	Male	20 to 24	52% (33-71%)	48% (29-67%)	0.99 (0.56-1.5)	3.8% (2.3-5.7%)	4.4% (2.4-7%)	0.5 (0.26-0.83)	41 (25-63)
Tonga	2005	Male	25 to 29	60% (40-76%)	40% (24-60%)	1.4 (0.86-2.1)	4.6% (2.9-6.5%)	5.2% (3-7.9%)	0.48 (0.25-0.76)	37 (22-55)
Tonga	2005	Male	30 to 34	61% (40-78%)	39% (22-60%)	1.5 (0.87-2.2)	3.7% (2.3-5.7%)	4% (2.3-6.6%)	0.41 (0.21-0.69)	31 (19-49)
Tonga	2005	Male	35 to 39	59% (41-76%)	41% (24-59%)	1.3 (0.73-2)	3.1% (1.9-4.8%)	3.5% (1.8-5.6%)	0.45 (0.22-0.75)	30 (17-47)
Tonga	2005	Male	40 to 44	57% (39-74%)	43% (26-61%)	1.1 (0.6-1.9)	2.4% (1.3-3.9%)	2.6% (1.2-4.4%)	0.34 (0.14-0.6)	20 (10-34)
Tonga	2005	Male	45 to 49	53% (32-71%)	47% (29-68%)	0.97 (0.47-1.7)	2.2% (1.1-3.6%)	2.4% (1-4.1%)	0.42 (0.17-0.74)	22 (10-36)
Tonga	2005	Male	50 to 54	50% (31-69%)	50% (31-69%)	0.78 (0.38-1.4)	1.7% (0.6-3.1%)	1.9% (0.51-3.5%)	0.33 (0.088-0.66)	14 (5-28)
Tonga	2005	Male	55 to 59	48% (29-67%)	52% (33-71%)	0.63 (0.31-1.1)	1.1% (0.19-2.2%)	1.1% (0.083-2.4%)	0.21 (0.015-0.47)	8.2 (1.5-17)
Tonga	2005	Male	60 to 64	45% (27-66%)	55% (34-73%)	0.53 (0.25-1)	0.68% (-0.13-1.7%)	0.72% (-0.18-1.9%)	0.2 (-0.05-0.54)	6.4 (-1.1-16)
Tonga	2005	Male	65 to 69	44% (25-62%)	56% (38-75%)	0.49 (0.24-0.85)	0.52% (-0.11-1.3%)	0.57% (-0.16-1.5%)	0.22 (-0.062-0.58)	5.4 (-1.2-14)
Tonga	2005	Male	70 to 74	41% (24-58%)	59% (42-76%)	0.43 (0.21-0.77)	0.29% (-0.32-1%)	0.3% (-0.39-1.1%)	0.13 (-0.17-0.48)	2.7 (-2.8-9.4)
Tonga	2005	Male	75 to 79	37% (21-56%)	63% (44-79%)	0.36 (0.16-0.68)	0.11% (-0.37-0.71%)	0.084% (-0.47-0.72%)	0.039 (-0.21-0.33)	0.86 (-2.8-5.2)
Tonga	2005	Male	80 to 84	34% (19-53%)	66% (47-81%)	0.32 (0.15-0.58)	0.1% (-0.33-0.53%)	0.069% (-0.43-0.56%)	0.024 (-0.15-0.2)	0.42 (-1.4-2.2)
Tonga	2005	Male	85 to 89	33% (19-49%)	67% (51-81%)	0.3 (0.14-0.55)	-0.0051% (-0.43-0.4%)	-0.056% (-0.56-0.39%)	-0.012 (-0.11-0.08)	-0.0085 (-0.78-0.7)
Tonga	2005	Male	90 to 94	34% (20-52%)	66% (48-80%)	0.31 (0.14-0.56)	-0.029% (-0.41-0.33%)	-0.096% (-0.55-0.3%)	-0.0083 (-0.048-0.027)	-0.015 (-0.21-0.17)
Tonga	2005	Male	95 plus	37% (20-58%)	63% (42-80%)	0.34 (0.13-0.66)	-0.2% (-0.71-0.22%)	-0.36% (-0.97-0.15%)	-0.0083 (-0.022-0.0033)	-0.019 (-0.065-0.019)
Tonga	2010	Female	All Ages	18% (13-24%)	82% (76-87%)	0.28 (0.19-0.39)	0.097% (-0.29-0.48%)	-0.31% (-0.95-0.31%)	-1.1 (-3.5-1.1)	14 (-40-67)
Tonga	2010	Female	15 to 19	18% (8.3-32%)	82% (68-92%)	0.32 (0.13-0.64)	0.68% (0.35-1.1%)	0.89% (0.26-2%)	0.029 (0.0082-0.067)	4.5 (2.3-7.6)
Tonga	2010	Female	20 to 24	22% (8.9-40%)	78% (60-91%)	0.49 (0.2-0.97)	1.2% (0.72-1.9%)	1.2% (0.41-2.5%)	0.041 (0.014-0.089)	7.2 (4.2-12)
Tonga	2010	Female	25 to 29	23% (10-42%)	77% (58-90%)	0.51 (0.22-0.95)	1.2% (0.74-1.8%)	1.2% (0.47-2.3%)	0.054 (0.021-0.11)	7.8 (4.6-12)
Tonga	2010	Female	30 to 34	25% (9.6-44%)	75% (56-90%)	0.54 (0.22-0.99)	1.1% (0.61-1.7%)	1.2% (0.53-2.2%)	0.07 (0.029-0.13)	7.5 (4.1-12)
Tonga	2010	Female	35 to 39	25% (9.6-43%)	75% (57-90%)	0.56 (0.23-1)	1.1% (0.54-1.7%)	1.5% (0.66-2.6%)	0.11 (0.045-0.21)	8.3 (4.3-14)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	2010	Female	40 to 44	25% (12-42%)	75% (58-88%)	0.52 (0.22-0.99)	0.89% (0.23-1.7%)	1.4% (0.42-2.7%)	0.13 (0.037-0.26)	6.9 (1.7-14)
Tonga	2010	Female	45 to 49	24% (10-42%)	76% (58-90%)	0.45 (0.17-0.87)	0.44% (-0.42-1.3%)	0.83% (-0.22-2.2%)	0.091 (-0.026-0.24)	3.4 (-3.3-11)
Tonga	2010	Female	50 to 54	24% (10-42%)	76% (58-90%)	0.39 (0.14-0.77)	-0.032% (-1.3-0.96%)	0.21% (-1.2-1.5%)	0.032 (-0.19-0.23)	-0.3 (-12-8.5)
Tonga	2010	Female	55 to 59	23% (11-42%)	77% (58-89%)	0.35 (0.13-0.75)	-0.29% (-1.5-0.6%)	-0.11% (-1.4-1.1%)	-0.017 (-0.24-0.16)	-2.2 (-11-4.3)
Tonga	2010	Female	60 to 64	23% (10-42%)	77% (58-90%)	0.34 (0.12-0.72)	-0.61% (-2-0.31%)	-0.53% (-2.2-0.6%)	-0.11 (-0.45-0.11)	-4.6 (-16-2.3)
Tonga	2010	Female	65 to 69	24% (11-42%)	76% (58-89%)	0.34 (0.12-0.69)	-0.81% (-2.2-0.15%)	-0.82% (-2.4-0.31%)	-0.2 (-0.6-0.072)	-6.1 (-17-1)
Tonga	2010	Female	70 to 74	24% (11-40%)	76% (60-89%)	0.32 (0.12-0.65)	-0.83% (-2-0.0067%)	-0.9% (-2.3-0.079%)	-0.31 (-0.78-0.023)	-6.9 (-17-0.061)
Tonga	2010	Female	75 to 79	23% (10-40%)	77% (60-90%)	0.29 (0.095-0.64)	-0.54% (-1.6-0.14%)	-0.55% (-1.7-0.24%)	-0.26 (-0.81-0.11)	-4.6 (-13-1.3)
Tonga	2010	Female	80 to 84	22% (9.2-41%)	78% (59-91%)	0.28 (0.094-0.6)	-0.54% (-1.4-0.057%)	-0.58% (-1.6-0.089%)	-0.33 (-0.92-0.052)	-4 (-11-0.43)
Tonga	2010	Female	85 to 89	23% (10-41%)	77% (59-90%)	0.3 (0.11-0.64)	-0.6% (-1.6-0.01%)	-0.66% (-1.8-0.029%)	-0.28 (-0.77-0.012)	-2.2 (-6.1-0.04)
Tonga	2010	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.35 (0.11-0.75)	-0.62% (-1.5-0.093%)	-0.7% (-1.7-0.1%)	-0.13 (-0.3-0.017)	-0.67 (-1.6-0.11)
Tonga	2010	Female	95 plus	31% (14-55%)	69% (45-86%)	0.41 (0.13-0.94)	-0.78% (-2.1-0.19%)	-0.94% (-2.5-0.16%)	-0.049 (-0.13-0.0082)	-0.16 (-0.44-0.039)
Tonga	2010	Male	All Ages	37% (29-46%)	63% (54-71%)	0.59 (0.44-0.8)	1.4% (0.89-2.1%)	1.1% (0.48-1.9%)	4.3 (1.8-7.8)	250 (150-380)
Tonga	2010	Male	15 to 19	34% (20-52%)	66% (48-80%)	0.38 (0.17-0.69)	1.5% (0.86-2.6%)	1.9% (0.84-3.5%)	0.16 (0.071-0.29)	15 (8.2-24)
Tonga	2010	Male	20 to 24	52% (33-69%)	48% (31-67%)	1 (0.56-1.7)	4% (2.3-6.2%)	4.7% (2.4-7.8%)	0.45 (0.22-0.78)	38 (21-59)
Tonga	2010	Male	25 to 29	59% (40-76%)	41% (24-60%)	1.5 (0.86-2.3)	4.7% (2.9-6.9%)	5.5% (3-8.5%)	0.55 (0.29-0.89)	42 (25-65)
Tonga	2010	Male	30 to 34	60% (39-77%)	40% (23-61%)	1.6 (0.88-2.4)	3.8% (2.4-5.9%)	4.3% (2.3-6.9%)	0.44 (0.22-0.76)	33 (19-53)
Tonga	2010	Male	35 to 39	59% (40-77%)	41% (23-60%)	1.4 (0.77-2.2)	3.3% (1.9-5.1%)	3.7% (2-6.2%)	0.47 (0.23-0.85)	32 (17-52)
Tonga	2010	Male	40 to 44	57% (39-74%)	43% (26-61%)	1.2 (0.6-2)	2.6% (1.3-4.3%)	2.8% (1.3-5%)	0.39 (0.16-0.72)	24 (11-41)
Tonga	2010	Male	45 to 49	53% (32-72%)	47% (28-68%)	1 (0.49-1.8)	2.3% (1.1-4%)	2.5% (1.1-4.5%)	0.41 (0.17-0.75)	21 (9.7-37)
Tonga	2010	Male	50 to 54	50% (31-68%)	50% (32-69%)	0.84 (0.4-1.5)	1.8% (0.68-3.4%)	2% (0.67-3.8%)	0.46 (0.15-0.95)	20 (7.3-41)
Tonga	2010	Male	55 to 59	48% (30-67%)	52% (33-70%)	0.68 (0.32-1.2)	1.2% (0.24-2.5%)	1.3% (0.16-2.8%)	0.27 (0.034-0.63)	10 (2.2-23)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	2010	Male	60 to 64	45% (26-65%)	55% (35-74%)	0.57 (0.25-1.1)	0.78% (-0.051-2%)	0.83% (-0.12-2.2%)	0.21 (-0.032-0.6)	6.8 (-0.39-18)
Tonga	2010	Male	65 to 69	43% (25-63%)	57% (37-75%)	0.51 (0.25-0.91)	0.6% (-0.087-1.6%)	0.65% (-0.12-1.8%)	0.23 (-0.043-0.63)	5.6 (-0.83-15)
Tonga	2010	Male	70 to 74	40% (25-58%)	60% (42-75%)	0.44 (0.22-0.84)	0.34% (-0.27-1.2%)	0.36% (-0.33-1.3%)	0.15 (-0.14-0.57)	3.2 (-2.4-11)
Tonga	2010	Male	75 to 79	37% (22-55%)	63% (45-78%)	0.38 (0.17-0.71)	0.16% (-0.34-0.81%)	0.13% (-0.43-0.84%)	0.056 (-0.19-0.37)	1.1 (-2.4-6.1)
Tonga	2010	Male	80 to 84	34% (19-52%)	66% (48-81%)	0.34 (0.15-0.61)	0.14% (-0.29-0.62%)	0.11% (-0.38-0.65%)	0.041 (-0.15-0.25)	0.64 (-1.4-2.9)
Tonga	2010	Male	85 to 89	33% (19-49%)	67% (51-81%)	0.32 (0.15-0.58)	0.028% (-0.39-0.46%)	-0.021% (-0.5-0.47%)	-0.0051 (-0.12-0.11)	0.06 (-0.84-0.99)
Tonga	2010	Male	90 to 94	34% (21-51%)	66% (49-79%)	0.33 (0.14-0.61)	0.00034% (-0.4-0.41%)	-0.063% (-0.54-0.4%)	-0.0061 (-0.052-0.039)	0.00021 (-0.23-0.24)
Tonga	2010	Male	95 plus	37% (19-58%)	63% (42-81%)	0.36 (0.13-0.73)	-0.17% (-0.7-0.32%)	-0.32% (-0.96-0.25%)	-0.0087 (-0.026-0.0068)	-0.019 (-0.076-0.035)
Tonga	2016	Female	All Ages	18% (13-24%)	82% (76-87%)	0.28 (0.16-0.44)	0.12% (-0.3-0.64%)	-0.27% (-0.96-0.55%)	-1 (-3.7-2)	18 (-42-93)
Tonga	2016	Female	15 to 19	18% (8.3-32%)	82% (68-92%)	0.32 (0.11-0.67)	0.71% (0.37-1.2%)	0.92% (0.25-2.1%)	0.03 (0.0078-0.07)	4.7 (2.4-8.3)
Tonga	2016	Female	20 to 24	22% (9.3-39%)	78% (61-91%)	0.5 (0.19-0.98)	1.2% (0.72-1.9%)	1.2% (0.39-2.7%)	0.047 (0.015-0.1)	8.3 (4.8-13)
Tonga	2016	Female	25 to 29	24% (10-42%)	76% (58-90%)	0.51 (0.2-0.99)	1.2% (0.74-1.9%)	1.2% (0.44-2.5%)	0.046 (0.015-0.099)	6.7 (3.9-11)
Tonga	2016	Female	30 to 34	25% (9-45%)	75% (55-91%)	0.53 (0.2-1.1)	1.1% (0.61-1.8%)	1.5% (0.52-2.5%)	0.063 (0.023-0.13)	6.8 (3.5-12)
Tonga	2016	Female	35 to 39	26% (9.7-44%)	74% (56-90%)	0.56 (0.2-1.1)	1.1% (0.5-2%)	1.5% (0.61-3%)	0.11 (0.039-0.22)	7.8 (3.5-15)
Tonga	2016	Female	40 to 44	25% (12-43%)	75% (57-88%)	0.52 (0.18-1.1)	0.94% (0.2-2%)	1.5% (0.42-3.1%)	0.15 (0.038-0.31)	8 (1.7-17)
Tonga	2016	Female	45 to 49	24% (11-42%)	76% (58-89%)	0.44 (0.15-0.91)	0.5% (-0.32-1.6%)	0.93% (-0.13-2.4%)	0.11 (-0.015-0.29)	4.3 (-2.9-13)
Tonga	2016	Female	50 to 54	24% (11-42%)	76% (58-89%)	0.38 (0.13-0.82)	0.052% (-1-1.2%)	0.34% (-0.99-1.8%)	0.049 (-0.15-0.28)	0.45 (-9.1-10)
Tonga	2016	Female	55 to 59	23% (10-42%)	77% (58-90%)	0.35 (0.12-0.77)	-0.2% (-1.4-0.76%)	0.018% (-1.3-1.3%)	0.0031 (-0.27-0.26)	-2 (-14-7.9)
Tonga	2016	Female	60 to 64	23% (10-41%)	77% (59-90%)	0.33 (0.11-0.74)	-0.48% (-1.8-0.46%)	-0.38% (-1.9-0.8%)	-0.084 (-0.43-0.16)	-4.2 (-16-4.2)
Tonga	2016	Female	65 to 69	24% (11-42%)	76% (58-89%)	0.34 (0.12-0.7)	-0.7% (-2-0.3%)	-0.69% (-2.3-0.45%)	-0.18 (-0.59-0.12)	-5.8 (-17-2.4)
Tonga	2016	Female	70 to 74	24% (11-42%)	76% (58-89%)	0.32 (0.11-0.72)	-0.76% (-1.9-0.18%)	-0.8% (-2.1-0.29%)	-0.27 (-0.68-0.099)	-6 (-15-1.4)
Tonga	2016	Female	75 to 79	23% (10-41%)	77% (59-90%)	0.3 (0.088-0.66)	-0.49% (-1.5-0.35%)	-0.48% (-1.6-0.46%)	-0.21 (-0.71-0.21)	-3.8 (-12-2.7)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tonga	2016	Female	80 to 84	23% (9.5-41%)	77% (59-91%)	0.29 (0.088-0.62)	-0.51% (-1.5-0.24%)	-0.53% (-1.6-0.33%)	-0.31 (-0.96-0.19)	-3.8 (-11-1.7)
Tonga	2016	Female	85 to 89	24% (10-42%)	76% (58-90%)	0.3 (0.095-0.68)	-0.55% (-1.5-0.19%)	-0.61% (-1.7-0.25%)	-0.33 (-0.92-0.13)	-2.6 (-7.3-0.89)
Tonga	2016	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.35 (0.11-0.78)	-0.56% (-1.4-0.2%)	-0.64% (-1.6-0.22%)	-0.18 (-0.46-0.066)	-0.96 (-2.4-0.34)
Tonga	2016	Female	95 plus	31% (13-54%)	69% (46-87%)	0.41 (0.12-0.98)	-0.72% (-2.1-0.27%)	-0.87% (-2.5-0.24%)	-0.069 (-0.2-0.019)	-0.22 (-0.65-0.089)
Tonga	2016	Male	All Ages	37% (29-46%)	63% (54-71%)	0.6 (0.37-0.9)	1.6% (0.86-2.5%)	1.2% (0.46-2.2%)	4.6 (1.8-9)	270 (150-440)
Tonga	2016	Male	15 to 19	34% (20-52%)	66% (48-80%)	0.39 (0.16-0.74)	1.5% (0.83-2.7%)	1.9% (0.8-3.7%)	0.17 (0.069-0.32)	16 (8.4-28)
Tonga	2016	Male	20 to 24	52% (34-69%)	48% (31-66%)	1.1 (0.51-1.8)	4% (2.3-6.4%)	4.7% (2.3-8.1%)	0.54 (0.25-0.97)	45 (25-74)
Tonga	2016	Male	25 to 29	59% (39-76%)	41% (24-61%)	1.5 (0.76-2.4)	4.7% (2.7-7.5%)	5.6% (2.8-9.6%)	0.51 (0.25-0.88)	39 (22-65)
Tonga	2016	Male	30 to 34	61% (40-78%)	39% (22-60%)	1.6 (0.78-2.7)	3.9% (2.2-6.5%)	4.4% (2.1-8.1%)	0.42 (0.19-0.81)	32 (17-55)
Tonga	2016	Male	35 to 39	59% (40-77%)	41% (23-60%)	1.4 (0.69-2.4)	3.4% (1.9-5.9%)	3.9% (1.9-7.1%)	0.43 (0.2-0.83)	29 (15-51)
Tonga	2016	Male	40 to 44	57% (38-75%)	43% (25-62%)	1.2 (0.54-2.1)	2.6% (1.3-4.8%)	2.9% (1.2-5.6%)	0.44 (0.18-0.93)	27 (12-51)
Tonga	2016	Male	45 to 49	52% (31-71%)	48% (29-69%)	1 (0.43-1.9)	2.4% (1-4.3%)	2.6% (1-4.8%)	0.48 (0.18-0.92)	25 (10-45)
Tonga	2016	Male	50 to 54	49% (30-69%)	51% (31-70%)	0.83 (0.36-1.6)	1.9% (0.71-3.8%)	2.1% (0.69-4.3%)	0.45 (0.14-0.96)	20 (7.2-41)
Tonga	2016	Male	55 to 59	47% (29-67%)	53% (33-71%)	0.68 (0.3-1.3)	1.2% (0.27-2.8%)	1.3% (0.16-3.2%)	0.41 (0.048-0.97)	16 (3.4-36)
Tonga	2016	Male	60 to 64	45% (26-66%)	55% (34-74%)	0.56 (0.22-1.1)	0.85% (-0.022-2.3%)	0.91% (-0.066-2.6%)	0.27 (-0.021-0.75)	8.5 (-0.21-23)
Tonga	2016	Male	65 to 69	43% (25-62%)	57% (38-75%)	0.51 (0.23-0.96)	0.67% (-0.05-1.8%)	0.73% (-0.091-2%)	0.23 (-0.026-0.62)	5.7 (-0.43-15)
Tonga	2016	Male	70 to 74	41% (24-58%)	59% (42-76%)	0.45 (0.2-0.9)	0.4% (-0.2-1.4%)	0.43% (-0.26-1.5%)	0.17 (-0.098-0.56)	3.4 (-1.6-11)
Tonga	2016	Male	75 to 79	37% (21-55%)	63% (45-79%)	0.38 (0.16-0.77)	0.2% (-0.33-1%)	0.18% (-0.42-1%)	0.079 (-0.19-0.45)	1.5 (-2.2-7.2)
Tonga	2016	Male	80 to 84	34% (19-52%)	66% (48-81%)	0.34 (0.15-0.65)	0.17% (-0.27-0.72%)	0.15% (-0.33-0.76%)	0.06 (-0.13-0.31)	0.85 (-1.3-3.6)
Tonga	2016	Male	85 to 89	32% (18-48%)	68% (52-82%)	0.32 (0.14-0.61)	0.061% (-0.39-0.56%)	0.016% (-0.5-0.58%)	0.0044 (-0.14-0.16)	0.15 (-0.91-1.4)
Tonga	2016	Male	90 to 94	34% (21-52%)	66% (48-79%)	0.33 (0.14-0.66)	0.026% (-0.39-0.52%)	-0.035% (-0.51-0.49%)	-0.0043 (-0.062-0.06)	0.019 (-0.28-0.38)
Tonga	2016	Male	95 plus	38% (19-59%)	62% (41-81%)	0.37 (0.13-0.82)	-0.15% (-0.68-0.48%)	-0.29% (-0.92-0.38%)	-0.009 (-0.028-0.012)	-0.018 (-0.085-0.058)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	1990	Female	All Ages	9.9% (7.8-12%)	90% (88-92%)	0.28 (0.22-0.35)	0.79% (0.48-1.2%)	0.88% (0.39-1.5%)	4.7 (2.1-8.1)	230 (140-340)
Vanuatu	1990	Female	15 to 19	13% (7.3-21%)	87% (79-93%)	0.39 (0.19-0.66)	1.1% (0.54-2.1%)	1.7% (0.6-3.4%)	0.15 (0.05-0.31)	14 (6.7-26)
Vanuatu	1990	Female	20 to 24	16% (9.3-26%)	84% (74-91%)	0.64 (0.33-1.1)	2.1% (1.2-3.4%)	2.7% (1.2-4.9%)	0.32 (0.14-0.6)	28 (16-48)
Vanuatu	1990	Female	25 to 29	16% (9.9-24%)	84% (76-90%)	0.63 (0.37-0.94)	2.1% (1.3-3.2%)	2.7% (1.4-4.2%)	0.42 (0.21-0.69)	33 (19-50)
Vanuatu	1990	Female	30 to 34	16% (9.4-24%)	84% (76-91%)	0.63 (0.36-1)	2% (1.2-3%)	2.4% (1.2-4.1%)	0.44 (0.21-0.75)	31 (18-49)
Vanuatu	1990	Female	35 to 39	14% (8.5-22%)	86% (78-91%)	0.6 (0.33-0.99)	2.1% (1.2-3.5%)	2.6% (1.3-4.5%)	0.56 (0.27-0.97)	33 (17-54)
Vanuatu	1990	Female	40 to 44	13% (7.9-20%)	87% (80-92%)	0.51 (0.25-0.89)	2% (0.97-3.4%)	2.5% (1.1-4.6%)	0.55 (0.24-1)	28 (13-49)
Vanuatu	1990	Female	45 to 49	13% (7.4-19%)	87% (81-93%)	0.42 (0.21-0.75)	1.4% (0.56-2.7%)	1.7% (0.65-3.4%)	0.43 (0.15-0.84)	19 (7.4-37)
Vanuatu	1990	Female	50 to 54	12% (7.7-19%)	88% (81-92%)	0.36 (0.18-0.66)	1.1% (0.3-2.5%)	1.4% (0.32-3.1%)	0.38 (0.089-0.85)	14 (3.8-32)
Vanuatu	1990	Female	55 to 59	11% (6.7-17%)	89% (83-93%)	0.31 (0.14-0.58)	0.94% (0.2-2.1%)	1.1% (0.26-2.5%)	0.32 (0.073-0.74)	10 (2.2-24)
Vanuatu	1990	Female	60 to 64	10% (6-16%)	90% (84-94%)	0.27 (0.13-0.51)	0.56% (-0.06-1.6%)	0.68% (-0.062-1.9%)	0.23 (-0.019-0.63)	6.3 (-0.67-17)
Vanuatu	1990	Female	65 to 69	12% (6.8-19%)	88% (81-93%)	0.28 (0.14-0.49)	0.56% (-0.1-1.5%)	0.68% (-0.12-1.7%)	0.28 (-0.05-0.73)	6.2 (-1-16)
Vanuatu	1990	Female	70 to 74	12% (7.5-20%)	88% (80-93%)	0.28 (0.13-0.51)	0.44% (-0.28-1.4%)	0.51% (-0.3-1.6%)	0.22 (-0.13-0.72)	3.9 (-2.5-13)
Vanuatu	1990	Female	75 to 79	12% (7-21%)	88% (79-93%)	0.26 (0.12-0.48)	0.43% (-0.22-1.3%)	0.49% (-0.23-1.5%)	0.23 (-0.11-0.74)	3.2 (-1.7-10)
Vanuatu	1990	Female	80 to 84	12% (6.6-21%)	88% (79-93%)	0.26 (0.11-0.46)	0.27% (-0.26-1%)	0.31% (-0.29-1.2%)	0.13 (-0.12-0.49)	1.4 (-1.3-5.2)
Vanuatu	1990	Female	85 to 89	13% (6.9-20%)	87% (80-93%)	0.26 (0.12-0.49)	0.19% (-0.26-0.9%)	0.21% (-0.32-0.99%)	0.04 (-0.059-0.19)	0.31 (-0.42-1.5)
Vanuatu	1990	Female	90 to 94	14% (7.9-22%)	86% (78-92%)	0.28 (0.12-0.52)	0.21% (-0.25-0.96%)	0.22% (-0.3-1.1%)	0.012 (-0.016-0.061)	0.067 (-0.082-0.32)
Vanuatu	1990	Female	95 plus	16% (7.7-27%)	84% (73-92%)	0.32 (0.12-0.65)	0.26% (-0.3-1.2%)	0.25% (-0.39-1.3%)	0.003 (-0.0048-0.016)	0.012 (-0.014-0.055)
Vanuatu	1990	Male	All Ages	29% (21-36%)	71% (64-79%)	0.69 (0.55-0.83)	2.1% (1.4-3%)	2.1% (1.2-3.1%)	16 (8.8-25)	800 (510-1100)
Vanuatu	1990	Male	15 to 19	28% (16-45%)	72% (55-84%)	0.4 (0.18-0.71)	1.5% (0.78-2.6%)	1.8% (0.72-3.4%)	0.27 (0.1-0.54)	25 (12-43)
Vanuatu	1990	Male	20 to 24	48% (29-67%)	52% (33-71%)	1.2 (0.72-1.9)	5.1% (2.9-8.1%)	6% (3.1-10%)	1.2 (0.6-2)	91 (51-150)
Vanuatu	1990	Male	25 to 29	58% (42-71%)	42% (29-58%)	2 (1.3-2.9)	7.5% (4.8-11%)	9% (5.3-13%)	2 (1.2-3.1)	140 (83-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	1990	Male	30 to 34	60% (43-74%)	40% (26-57%)	2.3 (1.4-3.3)	7.4% (4.4-11%)	8.8% (4.9-13%)	2.1 (1.1-3.3)	130 (74-190)
Vanuatu	1990	Male	35 to 39	57% (41-70%)	43% (30-59%)	1.8 (1.1-2.8)	5.6% (3.1-8.7%)	6.4% (3.3-10%)	1.7 (0.87-2.8)	98 (54-160)
Vanuatu	1990	Male	40 to 44	52% (37-65%)	48% (35-63%)	1.5 (0.78-2.4)	4.5% (2.2-7.5%)	5% (2.2-8.9%)	1.7 (0.71-3)	85 (40-150)
Vanuatu	1990	Male	45 to 49	46% (30-60%)	54% (40-70%)	1.2 (0.62-2.1)	3.6% (1.5-6.5%)	3.9% (1.4-7.5%)	1.5 (0.56-3.1)	69 (29-130)
Vanuatu	1990	Male	50 to 54	43% (28-57%)	57% (43-72%)	0.97 (0.5-1.7)	2.9% (1.1-5.8%)	3.1% (0.96-6.6%)	1.4 (0.43-2.9)	55 (19-110)
Vanuatu	1990	Male	55 to 59	44% (29-57%)	56% (43-71%)	0.84 (0.44-1.4)	2.1% (0.59-4.4%)	2.2% (0.49-4.9%)	0.97 (0.22-2.1)	33 (8.9-70)
Vanuatu	1990	Male	60 to 64	41% (26-56%)	59% (44-74%)	0.68 (0.35-1.2)	1.7% (0.38-3.7%)	1.8% (0.3-4.1%)	0.96 (0.17-2.3)	28 (6.5-63)
Vanuatu	1990	Male	65 to 69	38% (22-56%)	62% (44-78%)	0.57 (0.29-1)	1.4% (0.34-2.9%)	1.5% (0.32-3.2%)	0.94 (0.19-2)	22 (5.4-46)
Vanuatu	1990	Male	70 to 74	34% (20-51%)	66% (49-80%)	0.46 (0.23-0.81)	0.91% (0.047-2.1%)	0.98% (0.019-2.3%)	0.6 (0.011-1.4)	11 (0.55-27)
Vanuatu	1990	Male	75 to 79	30% (16-47%)	70% (53-84%)	0.38 (0.17-0.71)	0.64% (-0.034-1.7%)	0.68% (-0.08-1.8%)	0.46 (-0.058-1.2)	6.9 (-0.36-18)
Vanuatu	1990	Male	80 to 84	27% (14-43%)	73% (57-86%)	0.33 (0.16-0.6)	0.49% (0.012-1.2%)	0.51% (-0.025-1.3%)	0.3 (-0.015-0.76)	3.4 (0.078-8.4)
Vanuatu	1990	Male	85 to 89	26% (14-41%)	74% (59-86%)	0.32 (0.14-0.59)	0.36% (-0.052-0.94%)	0.36% (-0.11-1%)	0.1 (-0.03-0.28)	0.83 (-0.12-2.2)
Vanuatu	1990	Male	90 to 94	27% (15-43%)	73% (57-85%)	0.33 (0.15-0.61)	0.26% (-0.12-0.75%)	0.25% (-0.19-0.8%)	0.021 (-0.016-0.07)	0.13 (-0.06-0.39)
Vanuatu	1990	Male	95 plus	30% (15-49%)	70% (51-85%)	0.36 (0.15-0.71)	0.07% (-0.41-0.8%)	-0.015% (-0.58-0.8%)	-0.0003 (-0.011-0.015)	0.0055 (-0.032-0.061)
Vanuatu	1995	Female	All Ages	10% (7.9-13%)	90% (87-92%)	0.25 (0.2-0.32)	0.72% (0.43-1.1%)	0.73% (0.26-1.3%)	4.4 (1.5-7.9)	240 (140-360)
Vanuatu	1995	Female	15 to 19	13% (7.4-22%)	87% (78-93%)	0.35 (0.17-0.59)	1.1% (0.56-2%)	1.7% (0.62-3.3%)	0.17 (0.062-0.35)	16 (7.9-29)
Vanuatu	1995	Female	20 to 24	16% (9.4-26%)	84% (74-91%)	0.57 (0.3-0.95)	1.9% (1.1-3.2%)	2.5% (1.1-4.5%)	0.33 (0.15-0.61)	30 (17-49)
Vanuatu	1995	Female	25 to 29	16% (9.7-25%)	84% (75-90%)	0.56 (0.33-0.85)	2% (1.2-3%)	2.5% (1.3-3.9%)	0.44 (0.23-0.71)	35 (21-53)
Vanuatu	1995	Female	30 to 34	16% (9.1-23%)	84% (77-91%)	0.55 (0.33-0.87)	1.8% (1-2.7%)	2.1% (1.1-3.6%)	0.46 (0.22-0.78)	33 (19-52)
Vanuatu	1995	Female	35 to 39	14% (8.5-22%)	86% (78-92%)	0.53 (0.29-0.89)	1.9% (1-3.1%)	2.3% (1.2-4%)	0.59 (0.28-1)	35 (19-59)
Vanuatu	1995	Female	40 to 44	13% (8.1-20%)	87% (80-92%)	0.45 (0.23-0.78)	1.7% (0.85-3%)	2.1% (0.96-3.9%)	0.57 (0.25-1)	29 (14-51)
Vanuatu	1995	Female	45 to 49	13% (7.3-19%)	87% (81-93%)	0.37 (0.19-0.67)	1.2% (0.43-2.3%)	1.4% (0.47-3%)	0.41 (0.13-0.86)	18 (6.8-37)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	1995	Female	50 to 54	12% (7.5-18%)	88% (82-93%)	0.32 (0.16-0.59)	0.91% (0.16-2.1%)	1.1% (0.18-2.6%)	0.37 (0.057-0.9)	14 (2.5-34)
Vanuatu	1995	Female	55 to 59	11% (6.7-16%)	89% (84-93%)	0.28 (0.13-0.51)	0.73% (0.052-1.8%)	0.9% (0.064-2.2%)	0.32 (0.023-0.81)	10 (0.7-26)
Vanuatu	1995	Female	60 to 64	10% (6-16%)	90% (84-94%)	0.25 (0.11-0.47)	0.39% (-0.2-1.3%)	0.46% (-0.23-1.5%)	0.18 (-0.084-0.61)	5 (-2.5-17)
Vanuatu	1995	Female	65 to 69	12% (6.6-19%)	88% (81-93%)	0.25 (0.12-0.43)	0.36% (-0.25-1.2%)	0.44% (-0.29-1.4%)	0.2 (-0.12-0.63)	4.4 (-3.1-14)
Vanuatu	1995	Female	70 to 74	13% (7.3-20%)	87% (80-93%)	0.25 (0.12-0.45)	0.22% (-0.43-1.1%)	0.26% (-0.48-1.2%)	0.13 (-0.23-0.59)	2.2 (-4.3-11)
Vanuatu	1995	Female	75 to 79	12% (7.1-21%)	88% (79-93%)	0.23 (0.1-0.43)	0.24% (-0.34-1%)	0.28% (-0.39-1.2%)	0.14 (-0.19-0.6)	1.9 (-2.8-8.3)
Vanuatu	1995	Female	80 to 84	12% (6.7-21%)	88% (79-93%)	0.23 (0.1-0.41)	0.12% (-0.37-0.79%)	0.13% (-0.42-0.9%)	0.068 (-0.21-0.44)	0.73 (-2.2-4.7)
Vanuatu	1995	Female	85 to 89	13% (7-20%)	87% (80-93%)	0.23 (0.11-0.43)	0.067% (-0.38-0.71%)	0.065% (-0.44-0.78%)	0.016 (-0.11-0.18)	0.14 (-0.8-1.4)
Vanuatu	1995	Female	90 to 94	13% (7.6-22%)	87% (78-92%)	0.25 (0.11-0.47)	0.092% (-0.31-0.75%)	0.087% (-0.38-0.84%)	0.0062 (-0.027-0.058)	0.038 (-0.13-0.31)
Vanuatu	1995	Female	95 plus	15% (7.9-26%)	85% (74-92%)	0.28 (0.11-0.57)	0.13% (-0.39-0.97%)	0.1% (-0.48-1.1%)	0.0016 (-0.0074-0.017)	0.0078 (-0.023-0.058)
Vanuatu	1995	Male	All Ages	30% (22-37%)	70% (63-78%)	0.62 (0.5-0.77)	1.9% (1.2-2.7%)	1.8% (0.94-2.7%)	15 (7.9-24)	790 (490-1100)
Vanuatu	1995	Male	15 to 19	29% (15-45%)	71% (55-85%)	0.35 (0.16-0.63)	1.5% (0.75-2.6%)	1.7% (0.7-3.4%)	0.31 (0.12-0.62)	28 (14-49)
Vanuatu	1995	Male	20 to 24	49% (30-66%)	51% (34-70%)	1.1 (0.63-1.7)	4.8% (2.8-7.4%)	5.7% (2.9-9.1%)	1.3 (0.65-2.1)	97 (54-150)
Vanuatu	1995	Male	25 to 29	59% (41-72%)	41% (28-59%)	1.8 (1.2-2.5)	6.9% (4.3-9.9%)	8.2% (4.8-12%)	2 (1.2-3.1)	140 (86-210)
Vanuatu	1995	Male	30 to 34	60% (43-75%)	40% (25-57%)	2 (1.3-2.9)	6.7% (4.1-9.7%)	7.8% (4.4-12%)	2.2 (1.2-3.4)	140 (79-210)
Vanuatu	1995	Male	35 to 39	57% (41-70%)	43% (30-59%)	1.6 (0.97-2.5)	4.9% (2.7-7.7%)	5.5% (2.8-9%)	1.8 (0.9-2.9)	100 (57-160)
Vanuatu	1995	Male	40 to 44	52% (38-65%)	48% (35-62%)	1.3 (0.71-2)	3.8% (1.8-6.7%)	4.2% (1.7-7.6%)	1.5 (0.61-2.9)	80 (36-140)
Vanuatu	1995	Male	45 to 49	45% (30-60%)	55% (40-70%)	1.1 (0.56-1.9)	3% (1.1-5.7%)	3.2% (0.99-6.4%)	1.4 (0.43-3)	66 (23-130)
Vanuatu	1995	Male	50 to 54	43% (29-56%)	57% (44-71%)	0.85 (0.43-1.5)	2.4% (0.72-4.7%)	2.5% (0.55-5.2%)	1.3 (0.29-2.8)	51 (16-110)
Vanuatu	1995	Male	55 to 59	44% (29-58%)	56% (42-71%)	0.74 (0.38-1.3)	1.6% (0.3-3.5%)	1.7% (0.19-3.9%)	0.89 (0.096-2)	31 (5.8-69)
Vanuatu	1995	Male	60 to 64	41% (26-57%)	59% (43-74%)	0.6 (0.31-1.1)	1.3% (0.21-3.1%)	1.4% (0.12-3.4%)	0.81 (0.072-2)	24 (3.6-57)
Vanuatu	1995	Male	65 to 69	38% (22-55%)	62% (45-78%)	0.5 (0.25-0.86)	1.1% (0.13-2.4%)	1.1% (0.076-2.7%)	0.75 (0.049-1.8)	18 (2.2-41)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	1995	Male	70 to 74	34% (20-50%)	66% (50-80%)	0.41 (0.21-0.72)	0.66% (-0.08-1.8%)	0.7% (-0.13-2%)	0.47 (-0.09-1.3)	9.1 (-1.1-24)
Vanuatu	1995	Male	75 to 79	30% (17-47%)	70% (53-83%)	0.34 (0.16-0.62)	0.47% (-0.12-1.4%)	0.48% (-0.18-1.5%)	0.33 (-0.13-1)	5 (-1.3-15)
Vanuatu	1995	Male	80 to 84	27% (15-44%)	73% (56-85%)	0.3 (0.14-0.54)	0.37% (-0.064-0.96%)	0.37% (-0.12-1%)	0.24 (-0.079-0.68)	2.8 (-0.48-7.5)
Vanuatu	1995	Male	85 to 89	26% (14-41%)	74% (59-86%)	0.28 (0.13-0.51)	0.24% (-0.15-0.79%)	0.23% (-0.21-0.84%)	0.076 (-0.068-0.27)	0.68 (-0.42-2.2)
Vanuatu	1995	Male	90 to 94	27% (16-43%)	73% (57-84%)	0.29 (0.13-0.55)	0.17% (-0.16-0.64%)	0.15% (-0.25-0.67%)	0.015 (-0.026-0.069)	0.11 (-0.1-0.39)
Vanuatu	1995	Male	95 plus	30% (15-50%)	70% (50-85%)	0.32 (0.13-0.64)	-0.014% (-0.46-0.55%)	-0.11% (-0.65-0.49%)	-0.0027 (-0.016-0.012)	-0.0013 (-0.044-0.051)
Vanuatu	2000	Female	All Ages	10% (7.9-13%)	90% (87-92%)	0.22 (0.17-0.28)	0.56% (0.3-0.88%)	0.49% (0.089-0.99%)	3.2 (0.6-6.7)	200 (110-320)
Vanuatu	2000	Female	15 to 19	13% (7.3-21%)	87% (79-93%)	0.29 (0.14-0.49)	0.95% (0.48-1.7%)	1.3% (0.49-2.7%)	0.16 (0.06-0.34)	16 (8.3-30)
Vanuatu	2000	Female	20 to 24	16% (9.4-26%)	84% (74-91%)	0.49 (0.26-0.81)	1.7% (0.95-2.8%)	2% (0.88-3.8%)	0.3 (0.13-0.56)	29 (16-48)
Vanuatu	2000	Female	25 to 29	16% (9.9-24%)	84% (76-90%)	0.48 (0.27-0.73)	1.7% (1.1-2.6%)	2.1% (1.1-3.4%)	0.38 (0.19-0.65)	31 (19-49)
Vanuatu	2000	Female	30 to 34	16% (9.4-24%)	84% (76-91%)	0.47 (0.27-0.75)	1.5% (0.86-2.2%)	1.7% (0.85-2.9%)	0.39 (0.18-0.7)	29 (16-48)
Vanuatu	2000	Female	35 to 39	14% (8.6-22%)	86% (78-91%)	0.45 (0.24-0.78)	1.5% (0.82-2.6%)	1.8% (0.87-3.3%)	0.53 (0.24-1)	33 (17-57)
Vanuatu	2000	Female	40 to 44	13% (7.9-20%)	87% (80-92%)	0.39 (0.19-0.69)	1.3% (0.62-2.5%)	1.6% (0.67-3.2%)	0.5 (0.21-0.99)	27 (13-49)
Vanuatu	2000	Female	45 to 49	13% (7.1-19%)	87% (81-93%)	0.32 (0.16-0.58)	0.86% (0.26-1.8%)	1% (0.24-2.3%)	0.34 (0.078-0.79)	16 (4.7-35)
Vanuatu	2000	Female	50 to 54	12% (7.8-19%)	88% (81-92%)	0.28 (0.13-0.5)	0.6% (-0.048-1.5%)	0.72% (-0.077-1.9%)	0.26 (-0.029-0.72)	10 (-0.85-27)
Vanuatu	2000	Female	55 to 59	11% (6.7-17%)	89% (83-93%)	0.23 (0.11-0.42)	0.45% (-0.15-1.3%)	0.56% (-0.17-1.6%)	0.24 (-0.071-0.66)	7.6 (-2.5-21)
Vanuatu	2000	Female	60 to 64	10% (5.9-16%)	90% (84-94%)	0.21 (0.095-0.38)	0.16% (-0.36-0.84%)	0.2% (-0.41-1%)	0.089 (-0.18-0.44)	2.4 (-5.2-12)
Vanuatu	2000	Female	65 to 69	12% (6.5-19%)	88% (81-93%)	0.21 (0.1-0.37)	0.13% (-0.43-0.77%)	0.17% (-0.5-0.91%)	0.08 (-0.23-0.44)	1.6 (-5.6-9.8)
Vanuatu	2000	Female	70 to 74	13% (7.3-20%)	87% (80-93%)	0.21 (0.097-0.38)	-0.011% (-0.57-0.67%)	0.00039% (-0.65-0.76%)	0.00022 (-0.34-0.39)	-0.11 (-6.3-7.1)
Vanuatu	2000	Female	75 to 79	12% (7-21%)	88% (79-93%)	0.2 (0.087-0.37)	0.032% (-0.5-0.64%)	0.044% (-0.55-0.73%)	0.023 (-0.29-0.38)	0.27 (-4.2-5.4)
Vanuatu	2000	Female	80 to 84	12% (6.8-21%)	88% (79-93%)	0.19 (0.089-0.34)	-0.052% (-0.5-0.49%)	-0.061% (-0.57-0.54%)	-0.032 (-0.3-0.29)	-0.32 (-3.2-3)
Vanuatu	2000	Female	85 to 89	13% (7.1-20%)	87% (80-93%)	0.2 (0.088-0.36)	-0.078% (-0.48-0.45%)	-0.098% (-0.57-0.49%)	-0.028 (-0.16-0.14)	-0.19 (-1.2-1.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	2000	Female	90 to 94	13% (7.9-22%)	87% (78-92%)	0.21 (0.09-0.41)	-0.045% (-0.41-0.48%)	-0.068% (-0.48-0.53%)	-0.0061 (-0.045-0.047)	-0.024 (-0.22-0.24)
Vanuatu	2000	Female	95 plus	16% (7.9-27%)	84% (73-92%)	0.24 (0.096-0.49)	-0.028% (-0.55-0.62%)	-0.077% (-0.68-0.66%)	-0.0015 (-0.013-0.012)	-0.002 (-0.04-0.044)
Vanuatu	2000	Male	All Ages	30% (23-37%)	70% (63-77%)	0.55 (0.43-0.67)	1.6% (0.95-2.2%)	1.4% (0.63-2.2%)	13 (5.8-20)	710 (420-1000)
Vanuatu	2000	Male	15 to 19	29% (16-47%)	71% (53-84%)	0.3 (0.14-0.53)	1.4% (0.72-2.4%)	1.6% (0.64-3.1%)	0.32 (0.12-0.63)	30 (15-53)
Vanuatu	2000	Male	20 to 24	50% (29-68%)	50% (32-71%)	0.97 (0.55-1.5)	4.3% (2.5-6.7%)	5% (2.6-8.1%)	1.3 (0.67-2.2)	100 (58-160)
Vanuatu	2000	Male	25 to 29	59% (41-72%)	41% (28-59%)	1.6 (1-2.2)	6.1% (3.8-8.7%)	7.1% (4.2-10%)	1.8 (1-2.9)	130 (76-190)
Vanuatu	2000	Male	30 to 34	60% (43-73%)	40% (27-57%)	1.7 (1.1-2.5)	5.6% (3.3-8.3%)	6.4% (3.5-9.8%)	1.9 (0.99-3)	120 (70-190)
Vanuatu	2000	Male	35 to 39	56% (41-70%)	44% (30-59%)	1.4 (0.83-2.1)	4.1% (2.2-6.6%)	4.5% (2.1-7.6%)	1.6 (0.72-2.8)	95 (48-160)
Vanuatu	2000	Male	40 to 44	52% (38-65%)	48% (35-62%)	1.1 (0.6-1.8)	3% (1.3-5.3%)	3.2% (1.1-6%)	1.3 (0.44-2.5)	70 (29-130)
Vanuatu	2000	Male	45 to 49	46% (29-60%)	54% (40-71%)	0.91 (0.47-1.6)	2.3% (0.74-4.3%)	2.4% (0.55-4.8%)	1.1 (0.23-2.3)	53 (16-100)
Vanuatu	2000	Male	50 to 54	43% (28-57%)	57% (43-72%)	0.74 (0.38-1.3)	1.8% (0.4-3.6%)	1.8% (0.2-3.9%)	1 (0.095-2.2)	42 (9-87)
Vanuatu	2000	Male	55 to 59	44% (29-58%)	56% (42-71%)	0.63 (0.33-1.1)	1.1% (0.036-2.6%)	1.1% (-0.14-2.8%)	0.67 (-0.089-1.6)	25 (0.81-55)
Vanuatu	2000	Male	60 to 64	41% (26-56%)	59% (44-74%)	0.52 (0.27-0.92)	0.86% (-0.059-2.2%)	0.87% (-0.15-2.4%)	0.55 (-0.098-1.5)	17 (-1.2-44)
Vanuatu	2000	Male	65 to 69	38% (23-54%)	62% (46-77%)	0.44 (0.22-0.77)	0.7% (-0.065-1.8%)	0.72% (-0.13-1.9%)	0.48 (-0.085-1.3)	12 (-1.1-30)
Vanuatu	2000	Male	70 to 74	34% (20-49%)	66% (51-80%)	0.35 (0.18-0.63)	0.41% (-0.23-1.3%)	0.42% (-0.3-1.4%)	0.3 (-0.22-1)	6 (-3.4-19)
Vanuatu	2000	Male	75 to 79	30% (17-47%)	70% (53-83%)	0.29 (0.13-0.54)	0.27% (-0.25-1%)	0.26% (-0.34-1.1%)	0.19 (-0.24-0.78)	3.2 (-2.9-12)
Vanuatu	2000	Male	80 to 84	27% (14-43%)	73% (57-86%)	0.25 (0.12-0.45)	0.23% (-0.16-0.71%)	0.21% (-0.22-0.75%)	0.14 (-0.14-0.49)	1.8 (-1.2-5.6)
Vanuatu	2000	Male	85 to 89	26% (14-40%)	74% (60-86%)	0.24 (0.11-0.43)	0.13% (-0.23-0.55%)	0.098% (-0.31-0.57%)	0.035 (-0.11-0.2)	0.38 (-0.68-1.6)
Vanuatu	2000	Male	90 to 94	27% (16-42%)	73% (58-84%)	0.25 (0.11-0.48)	0.082% (-0.24-0.44%)	0.04% (-0.34-0.45%)	0.0048 (-0.041-0.055)	0.058 (-0.17-0.31)
Vanuatu	2000	Male	95 plus	30% (15-50%)	70% (50-85%)	0.28 (0.11-0.56)	-0.095% (-0.54-0.33%)	-0.2% (-0.76-0.28%)	-0.0058 (-0.021-0.0077)	-0.01 (-0.06-0.037)
Vanuatu	2005	Female	All Ages	10% (8.2-13%)	90% (87-92%)	0.18 (0.14-0.24)	0.4% (0.17-0.67%)	0.27% (-0.075-0.68%)	2.1 (-0.55-5.4)	170 (70-290)
Vanuatu	2005	Female	15 to 19	13% (7.3-21%)	87% (79-93%)	0.24 (0.11-0.4)	0.75% (0.39-1.3%)	0.98% (0.33-2.1%)	0.13 (0.043-0.28)	14 (7.2-26)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	2005	Female	20 to 24	16% (8.9-26%)	84% (74-91%)	0.38 (0.2-0.63)	1.3% (0.74-2.2%)	1.5% (0.59-2.8%)	0.27 (0.11-0.51)	28 (16-47)
Vanuatu	2005	Female	25 to 29	16% (9.9-24%)	84% (76-90%)	0.38 (0.22-0.59)	1.4% (0.87-2.2%)	1.6% (0.8-2.8%)	0.34 (0.16-0.62)	30 (17-48)
Vanuatu	2005	Female	30 to 34	16% (9.5-24%)	84% (76-91%)	0.39 (0.22-0.6)	1.2% (0.65-1.8%)	1.3% (0.54-2.2%)	0.31 (0.12-0.58)	25 (13-42)
Vanuatu	2005	Female	35 to 39	14% (8.6-22%)	86% (78-91%)	0.37 (0.2-0.63)	1.2% (0.61-2%)	1.4% (0.6-2.5%)	0.46 (0.19-0.89)	29 (15-52)
Vanuatu	2005	Female	40 to 44	13% (8.1-20%)	87% (80-92%)	0.31 (0.16-0.54)	0.97% (0.42-1.8%)	1.2% (0.41-2.3%)	0.44 (0.15-0.88)	24 (9.9-44)
Vanuatu	2005	Female	45 to 49	13% (7.1-19%)	87% (81-93%)	0.26 (0.13-0.45)	0.56% (0.045-1.3%)	0.66% (-0.02-1.6%)	0.28 (-0.0074-0.7)	13 (1.1-30)
Vanuatu	2005	Female	50 to 54	12% (7.8-18%)	88% (82-92%)	0.22 (0.11-0.39)	0.31% (-0.25-1%)	0.37% (-0.31-1.3%)	0.17 (-0.14-0.6)	6.7 (-5.1-23)
Vanuatu	2005	Female	55 to 59	11% (6.6-17%)	89% (83-93%)	0.19 (0.091-0.36)	0.22% (-0.33-0.86%)	0.28% (-0.38-1%)	0.14 (-0.19-0.54)	4.4 (-6.8-17)
Vanuatu	2005	Female	60 to 64	10% (6-16%)	90% (84-94%)	0.17 (0.078-0.32)	-0.0038% (-0.5-0.55%)	0.012% (-0.56-0.69%)	0.0069 (-0.32-0.39)	-0.06 (-9.4-10)
Vanuatu	2005	Female	65 to 69	12% (6.8-19%)	88% (81-93%)	0.17 (0.082-0.31)	-0.051% (-0.59-0.49%)	-0.033% (-0.65-0.58%)	-0.02 (-0.39-0.36)	-0.83 (-9.4-7.9)
Vanuatu	2005	Female	70 to 74	13% (7.6-21%)	87% (79-92%)	0.17 (0.08-0.32)	-0.19% (-0.71-0.33%)	-0.21% (-0.79-0.4%)	-0.13 (-0.49-0.26)	-2.6 (-9.4-4.5)
Vanuatu	2005	Female	75 to 79	13% (7.1-21%)	87% (79-93%)	0.16 (0.07-0.3)	-0.13% (-0.62-0.34%)	-0.14% (-0.7-0.4%)	-0.082 (-0.41-0.24)	-1.2 (-5.9-3.3)
Vanuatu	2005	Female	80 to 84	12% (6.8-21%)	88% (79-93%)	0.16 (0.071-0.28)	-0.19% (-0.61-0.2%)	-0.22% (-0.69-0.23%)	-0.12 (-0.38-0.13)	-1.2 (-4-1.3)
Vanuatu	2005	Female	85 to 89	13% (7.1-21%)	87% (79-93%)	0.16 (0.074-0.3)	-0.2% (-0.62-0.2%)	-0.24% (-0.71-0.22%)	-0.067 (-0.21-0.065)	-0.48 (-1.5-0.51)
Vanuatu	2005	Female	90 to 94	14% (7.8-22%)	86% (78-92%)	0.17 (0.075-0.35)	-0.16% (-0.52-0.24%)	-0.2% (-0.61-0.26%)	-0.02 (-0.061-0.026)	-0.097 (-0.3-0.14)
Vanuatu	2005	Female	95 plus	16% (7.9-27%)	84% (73-92%)	0.2 (0.079-0.42)	-0.16% (-0.67-0.38%)	-0.23% (-0.81-0.4%)	-0.0054 (-0.019-0.0094)	-0.015 (-0.06-0.034)
Vanuatu	2005	Male	All Ages	31% (23-38%)	69% (62-77%)	0.45 (0.34-0.56)	1.2% (0.72-1.8%)	1% (0.36-1.7%)	10 (3.8-18)	630 (360-930)
Vanuatu	2005	Male	15 to 19	28% (15-45%)	72% (55-85%)	0.24 (0.11-0.42)	1.2% (0.61-2%)	1.3% (0.49-2.6%)	0.29 (0.11-0.57)	28 (15-47)
Vanuatu	2005	Male	20 to 24	48% (30-66%)	52% (34-70%)	0.74 (0.42-1.1)	3.5% (2.1-5.5%)	3.9% (2-6.5%)	1.2 (0.61-2.2)	100 (59-160)
Vanuatu	2005	Male	25 to 29	58% (42-72%)	42% (28-58%)	1.2 (0.76-1.8)	4.9% (2.9-7.2%)	5.5% (3-8.5%)	1.7 (0.91-2.8)	120 (70-190)
Vanuatu	2005	Male	30 to 34	60% (43-74%)	40% (26-57%)	1.4 (0.79-2)	4.4% (2.5-6.9%)	4.9% (2.5-8%)	1.6 (0.77-2.7)	110 (59-170)
Vanuatu	2005	Male	35 to 39	56% (42-70%)	44% (30-58%)	1.1 (0.64-1.7)	3.2% (1.7-5%)	3.4% (1.4-5.7%)	1.3 (0.56-2.3)	82 (43-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	2005	Male	40 to 44	52% (37-65%)	48% (35-63%)	0.9 (0.5-1.5)	2.3% (0.92-4%)	2.3% (0.62-4.4%)	1.1 (0.29-2.2)	62 (23-110)
Vanuatu	2005	Male	45 to 49	46% (29-60%)	54% (40-71%)	0.72 (0.38-1.2)	1.7% (0.45-3.3%)	1.7% (0.16-3.6%)	0.94 (0.081-2)	48 (12-94)
Vanuatu	2005	Male	50 to 54	43% (28-57%)	57% (43-72%)	0.57 (0.29-0.97)	1.3% (0.12-2.6%)	1.2% (-0.15-2.9%)	0.78 (-0.085-1.8)	35 (3.2-72)
Vanuatu	2005	Male	55 to 59	44% (29-57%)	56% (43-71%)	0.49 (0.25-0.85)	0.7% (-0.18-1.8%)	0.64% (-0.41-1.9%)	0.45 (-0.29-1.4)	18 (-5.1-48)
Vanuatu	2005	Male	60 to 64	41% (25-56%)	59% (44-75%)	0.41 (0.2-0.74)	0.47% (-0.27-1.5%)	0.44% (-0.41-1.6%)	0.33 (-0.31-1.2)	11 (-6.3-36)
Vanuatu	2005	Male	65 to 69	38% (21-55%)	62% (45-79%)	0.34 (0.17-0.61)	0.38% (-0.24-1.2%)	0.37% (-0.34-1.3%)	0.29 (-0.26-1)	7.7 (-4.7-24)
Vanuatu	2005	Male	70 to 74	34% (20-50%)	66% (50-80%)	0.28 (0.14-0.51)	0.19% (-0.33-0.81%)	0.17% (-0.41-0.86%)	0.13 (-0.32-0.67)	3 (-5.3-13)
Vanuatu	2005	Male	75 to 79	30% (17-47%)	70% (53-83%)	0.22 (0.1-0.41)	0.11% (-0.34-0.61%)	0.083% (-0.42-0.62%)	0.06 (-0.31-0.45)	1.3 (-3.9-6.9)
Vanuatu	2005	Male	80 to 84	27% (14-43%)	73% (57-86%)	0.2 (0.094-0.35)	0.11% (-0.25-0.48%)	0.083% (-0.32-0.49%)	0.052 (-0.21-0.31)	0.83 (-1.9-3.5)
Vanuatu	2005	Male	85 to 89	26% (14-40%)	74% (60-86%)	0.19 (0.087-0.34)	0.017% (-0.32-0.32%)	-0.025% (-0.42-0.32%)	-0.0088 (-0.14-0.11)	0.051 (-0.95-0.94)
Vanuatu	2005	Male	90 to 94	27% (15-41%)	73% (59-85%)	0.19 (0.09-0.37)	-0.0014% (-0.33-0.27%)	-0.053% (-0.43-0.26%)	-0.0067 (-0.054-0.032)	-0.00078 (-0.24-0.2)
Vanuatu	2005	Male	95 plus	29% (14-49%)	71% (51-86%)	0.21 (0.086-0.42)	-0.16% (-0.55-0.15%)	-0.28% (-0.76-0.091%)	-0.0088 (-0.023-0.0028)	-0.02 (-0.069-0.019)
Vanuatu	2010	Female	All Ages	10% (8-12%)	90% (88-92%)	0.14 (0.097-0.19)	0.28% (0.081-0.5%)	0.089% (-0.2-0.41%)	0.8 (-1.7-3.7)	130 (38-240)
Vanuatu	2010	Female	15 to 19	13% (7.5-21%)	87% (79-92%)	0.18 (0.085-0.32)	0.6% (0.3-1%)	0.7% (0.19-1.5%)	0.092 (0.025-0.2)	12 (5.8-20)
Vanuatu	2010	Female	20 to 24	16% (9.2-25%)	84% (75-91%)	0.29 (0.15-0.5)	1% (0.6-1.7%)	1% (0.35-2.1%)	0.21 (0.074-0.43)	26 (14-42)
Vanuatu	2010	Female	25 to 29	16% (10-23%)	84% (77-90%)	0.29 (0.16-0.46)	1.1% (0.69-1.7%)	1.2% (0.53-2.2%)	0.28 (0.12-0.51)	28 (16-44)
Vanuatu	2010	Female	30 to 34	15% (9.4-23%)	85% (77-91%)	0.29 (0.16-0.47)	0.88% (0.46-1.4%)	0.85% (0.3-1.6%)	0.23 (0.077-0.44)	21 (11-36)
Vanuatu	2010	Female	35 to 39	13% (8.3-21%)	87% (79-92%)	0.27 (0.14-0.45)	0.85% (0.44-1.5%)	0.92% (0.35-1.8%)	0.32 (0.12-0.67)	23 (11-40)
Vanuatu	2010	Female	40 to 44	12% (7.6-18%)	88% (82-92%)	0.22 (0.11-0.4)	0.67% (0.26-1.3%)	0.74% (0.2-1.6%)	0.3 (0.079-0.67)	18 (6.6-35)
Vanuatu	2010	Female	45 to 49	11% (6.7-16%)	89% (84-93%)	0.17 (0.088-0.31)	0.35% (-0.047-0.84%)	0.38% (-0.14-1%)	0.21 (-0.072-0.59)	11 (-1.4-26)
Vanuatu	2010	Female	50 to 54	11% (6.8-16%)	89% (84-93%)	0.15 (0.07-0.27)	0.14% (-0.27-0.62%)	0.15% (-0.36-0.76%)	0.079 (-0.19-0.37)	3.6 (-6.6-15)
Vanuatu	2010	Female	55 to 59	9.3% (5.8-14%)	91% (86-94%)	0.13 (0.059-0.24)	0.09% (-0.33-0.51%)	0.11% (-0.37-0.62%)	0.077 (-0.26-0.42)	2.5 (-9.1-14)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	2010	Female	60 to 64	9% (5.3-15%)	91% (85-95%)	0.12 (0.055-0.22)	-0.076% (-0.48-0.3%)	-0.081% (-0.57-0.38%)	-0.049 (-0.34-0.23)	-1.5 (-9.7-6.1)
Vanuatu	2010	Female	65 to 69	11% (6.2-17%)	89% (83-94%)	0.13 (0.06-0.23)	-0.14% (-0.61-0.26%)	-0.14% (-0.69-0.32%)	-0.12 (-0.58-0.28)	-3.3 (-14-5.9)
Vanuatu	2010	Female	70 to 74	12% (7.5-20%)	88% (80-93%)	0.13 (0.059-0.24)	-0.3% (-0.74-0.1%)	-0.33% (-0.84-0.14%)	-0.33 (-0.86-0.14)	-6.3 (-16-2.2)
Vanuatu	2010	Female	75 to 79	12% (7-21%)	88% (79-93%)	0.12 (0.054-0.23)	-0.23% (-0.67-0.11%)	-0.25% (-0.76-0.14%)	-0.18 (-0.53-0.094)	-2.6 (-7.6-1.2)
Vanuatu	2010	Female	80 to 84	12% (6.8-21%)	88% (79-93%)	0.12 (0.053-0.22)	-0.29% (-0.67-0.006%)	-0.33% (-0.74-0.018%)	-0.19 (-0.45-0.01)	-2 (-4.7-0.037)
Vanuatu	2010	Female	85 to 89	13% (7.1-20%)	87% (80-93%)	0.12 (0.055-0.24)	-0.29% (-0.72-0.012%)	-0.34% (-0.83-0.011%)	-0.096 (-0.24-0.0032)	-0.7 (-1.8-0.031)
Vanuatu	2010	Female	90 to 94	14% (7.8-22%)	86% (78-92%)	0.13 (0.054-0.26)	-0.26% (-0.63-0.0092%)	-0.31% (-0.72-0.0013%)	-0.031 (-0.072-0.00013)	-0.15 (-0.36-0.0054)
Vanuatu	2010	Female	95 plus	16% (8-27%)	84% (73-92%)	0.15 (0.057-0.31)	-0.26% (-0.77-0.16%)	-0.34% (-0.93-0.14%)	-0.0089 (-0.024-0.0036)	-0.026 (-0.076-0.016)
Vanuatu	2010	Male	All Ages	34% (26-41%)	66% (59-74%)	0.39 (0.27-0.53)	1% (0.54-1.7%)	0.77% (0.17-1.4%)	8.6 (2.1-16)	570 (300-930)
Vanuatu	2010	Male	15 to 19	29% (15-47%)	71% (53-85%)	0.19 (0.078-0.34)	1% (0.54-1.7%)	1.1% (0.39-2.2%)	0.22 (0.079-0.45)	23 (12-39)
Vanuatu	2010	Male	20 to 24	51% (31-69%)	49% (31-69%)	0.62 (0.32-1)	3% (1.6-4.8%)	3.3% (1.4-5.7%)	1.1 (0.49-2)	93 (49-150)
Vanuatu	2010	Male	25 to 29	63% (46-74%)	37% (26-54%)	1.1 (0.64-1.6)	4.2% (2.5-6.3%)	4.7% (2.4-7.4%)	1.8 (0.88-2.9)	130 (75-210)
Vanuatu	2010	Male	30 to 34	65% (50-76%)	35% (24-50%)	1.2 (0.69-1.8)	3.6% (2-5.7%)	3.9% (1.8-6.5%)	1.5 (0.66-2.6)	100 (54-170)
Vanuatu	2010	Male	35 to 39	60% (47-71%)	40% (29-53%)	0.92 (0.54-1.5)	2.5% (1.2-4.3%)	2.6% (0.92-4.9%)	1.1 (0.39-2.2)	73 (34-130)
Vanuatu	2010	Male	40 to 44	53% (41-66%)	47% (34-59%)	0.72 (0.38-1.2)	1.7% (0.52-3.2%)	1.6% (0.2-3.4%)	0.83 (0.092-1.8)	51 (15-97)
Vanuatu	2010	Male	45 to 49	47% (32-59%)	53% (41-68%)	0.56 (0.31-0.99)	1.3% (0.2-2.5%)	1.1% (-0.16-2.7%)	0.77 (-0.11-1.8)	43 (6.4-87)
Vanuatu	2010	Male	50 to 54	42% (30-55%)	58% (45-70%)	0.42 (0.21-0.76)	0.94% (-0.02-2%)	0.84% (-0.33-2.1%)	0.58 (-0.21-1.5)	28 (-0.56-61)
Vanuatu	2010	Male	55 to 59	41% (27-54%)	59% (46-73%)	0.34 (0.17-0.6)	0.47% (-0.22-1.2%)	0.38% (-0.45-1.2%)	0.32 (-0.38-1.1)	15 (-6.9-40)
Vanuatu	2010	Male	60 to 64	39% (25-53%)	61% (47-75%)	0.28 (0.14-0.51)	0.26% (-0.34-0.94%)	0.2% (-0.49-0.94%)	0.16 (-0.39-0.76)	6.4 (-8.2-23)
Vanuatu	2010	Male	65 to 69	37% (22-52%)	63% (48-78%)	0.25 (0.12-0.46)	0.19% (-0.32-0.75%)	0.15% (-0.42-0.79%)	0.17 (-0.51-0.88)	5.5 (-9.3-22)
Vanuatu	2010	Male	70 to 74	34% (20-49%)	66% (51-80%)	0.21 (0.099-0.39)	0.061% (-0.38-0.53%)	0.026% (-0.48-0.57%)	0.024 (-0.45-0.54)	1.2 (-7.3-10)
Vanuatu	2010	Male	75 to 79	30% (17-46%)	70% (54-83%)	0.17 (0.076-0.33)	0.025% (-0.36-0.37%)	-0.016% (-0.45-0.37%)	-0.012 (-0.36-0.29)	0.31 (-4.6-4.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	2010	Male	80 to 84	27% (14-42%)	73% (58-86%)	0.15 (0.068-0.28)	0.055% (-0.27-0.32%)	0.019% (-0.34-0.32%)	0.012 (-0.21-0.19)	0.39 (-1.9-2.3)
Vanuatu	2010	Male	85 to 89	26% (14-40%)	74% (60-86%)	0.14 (0.067-0.28)	-0.033% (-0.35-0.2%)	-0.083% (-0.45-0.19%)	-0.028 (-0.15-0.063)	-0.092 (-0.99-0.59)
Vanuatu	2010	Male	90 to 94	27% (15-42%)	73% (58-85%)	0.15 (0.066-0.29)	-0.03% (-0.33-0.2%)	-0.085% (-0.44-0.18%)	-0.011 (-0.056-0.023)	-0.022 (-0.25-0.15)
Vanuatu	2010	Male	95 plus	29% (14-49%)	71% (51-86%)	0.16 (0.064-0.33)	-0.18% (-0.57-0.095%)	-0.29% (-0.77-0.044%)	-0.0096 (-0.025-0.0015)	-0.023 (-0.075-0.012)
Vanuatu	2016	Female	All Ages	10% (8.2-13%)	90% (87-92%)	0.14 (0.072-0.23)	0.28% (0.084-0.57%)	0.089% (-0.22-0.51%)	0.95 (-2.3-5.4)	150 (42-310)
Vanuatu	2016	Female	15 to 19	13% (7.4-21%)	87% (79-93%)	0.18 (0.067-0.36)	0.59% (0.28-1.1%)	0.68% (0.14-1.7%)	0.092 (0.02-0.23)	12 (5.7-23)
Vanuatu	2016	Female	20 to 24	16% (9-26%)	84% (74-91%)	0.29 (0.12-0.55)	1.1% (0.58-1.8%)	1% (0.28-2.3%)	0.19 (0.052-0.43)	25 (13-45)
Vanuatu	2016	Female	25 to 29	16% (10-23%)	84% (77-90%)	0.28 (0.13-0.5)	1.1% (0.67-1.8%)	1.1% (0.44-2.2%)	0.31 (0.12-0.61)	33 (18-52)
Vanuatu	2016	Female	30 to 34	15% (9.4-22%)	85% (78-91%)	0.27 (0.12-0.51)	0.87% (0.43-1.4%)	0.82% (0.24-1.8%)	0.26 (0.069-0.56)	26 (12-45)
Vanuatu	2016	Female	35 to 39	13% (8.2-21%)	87% (79-92%)	0.25 (0.11-0.49)	0.84% (0.42-1.6%)	0.91% (0.3-2%)	0.35 (0.11-0.79)	25 (12-49)
Vanuatu	2016	Female	40 to 44	12% (7.4-18%)	88% (82-93%)	0.21 (0.089-0.43)	0.67% (0.25-1.4%)	0.74% (0.18-1.8%)	0.34 (0.075-0.8)	20 (7.2-42)
Vanuatu	2016	Female	45 to 49	11% (6.5-17%)	89% (83-93%)	0.17 (0.072-0.34)	0.38% (-0.02-0.99%)	0.42% (-0.1-1.2%)	0.23 (-0.061-0.69)	12 (-0.59-32)
Vanuatu	2016	Female	50 to 54	11% (6.8-16%)	89% (84-93%)	0.14 (0.059-0.3)	0.17% (-0.24-0.72%)	0.19% (-0.32-0.86%)	0.14 (-0.24-0.66)	6.1 (-8-27)
Vanuatu	2016	Female	55 to 59	9.3% (5.8-14%)	91% (86-94%)	0.12 (0.048-0.25)	0.12% (-0.27-0.58%)	0.15% (-0.33-0.72%)	0.12 (-0.25-0.54)	3.8 (-8.3-18)
Vanuatu	2016	Female	60 to 64	9% (5.2-14%)	91% (86-95%)	0.11 (0.045-0.23)	-0.041% (-0.43-0.36%)	-0.039% (-0.49-0.45%)	-0.035 (-0.43-0.38)	-1.2 (-13-11)
Vanuatu	2016	Female	65 to 69	11% (6.5-17%)	89% (83-94%)	0.12 (0.05-0.25)	-0.1% (-0.53-0.33%)	-0.095% (-0.62-0.42%)	-0.085 (-0.54-0.35)	-2.5 (-13-7.7)
Vanuatu	2016	Female	70 to 74	13% (7.5-20%)	87% (80-93%)	0.13 (0.048-0.28)	-0.26% (-0.72-0.15%)	-0.29% (-0.8-0.2%)	-0.34 (-0.97-0.24)	-6.6 (-18-4.2)
Vanuatu	2016	Female	75 to 79	12% (7.2-21%)	88% (79-93%)	0.12 (0.046-0.25)	-0.2% (-0.65-0.21%)	-0.21% (-0.72-0.24%)	-0.25 (-0.89-0.29)	-3.8 (-13-3.9)
Vanuatu	2016	Female	80 to 84	12% (6.7-21%)	88% (79-93%)	0.12 (0.044-0.24)	-0.27% (-0.64-0.063%)	-0.3% (-0.75-0.074%)	-0.25 (-0.6-0.059)	-2.6 (-6.4-0.61)
Vanuatu	2016	Female	85 to 89	13% (6.9-20%)	87% (80-93%)	0.12 (0.048-0.25)	-0.26% (-0.7-0.08%)	-0.31% (-0.8-0.089%)	-0.093 (-0.24-0.026)	-0.68 (-1.8-0.2)
Vanuatu	2016	Female	90 to 94	14% (7.8-23%)	86% (77-92%)	0.13 (0.046-0.27)	-0.24% (-0.6-0.062%)	-0.29% (-0.71-0.056%)	-0.028 (-0.068-0.0057)	-0.14 (-0.34-0.036)
Vanuatu	2016	Female	95 plus	16% (7.8-27%)	84% (73-92%)	0.14 (0.048-0.33)	-0.25% (-0.76-0.17%)	-0.32% (-0.92-0.15%)	-0.0076 (-0.022-0.0037)	-0.023 (-0.07-0.016)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Vanuatu	2016	Male	All Ages	34% (27-41%)	66% (59-73%)	0.39 (0.2-0.63)	1.1% (0.49-2%)	0.78% (0.13-1.7%)	9.9 (1.7-22)	650 (290-1200)
Vanuatu	2016	Male	15 to 19	29% (16-45%)	71% (55-84%)	0.18 (0.065-0.36)	1% (0.51-1.8%)	1.1% (0.34-2.2%)	0.26 (0.085-0.55)	27 (14-48)
Vanuatu	2016	Male	20 to 24	50% (32-67%)	50% (33-68%)	0.58 (0.26-1.1)	2.9% (1.5-5.1%)	3.2% (1.2-6%)	1 (0.38-2)	87 (42-150)
Vanuatu	2016	Male	25 to 29	63% (48-73%)	37% (27-52%)	1 (0.51-1.7)	4.1% (2.1-6.7%)	4.5% (1.9-7.9%)	1.8 (0.73-3.3)	140 (66-230)
Vanuatu	2016	Male	30 to 34	66% (50-76%)	34% (24-50%)	1.2 (0.55-2.1)	3.6% (1.7-6.3%)	3.8% (1.3-7.4%)	1.8 (0.61-3.7)	130 (57-230)
Vanuatu	2016	Male	35 to 39	60% (47-72%)	40% (28-53%)	0.92 (0.44-1.7)	2.6% (1.1-4.9%)	2.6% (0.7-5.6%)	1.4 (0.34-3)	90 (37-180)
Vanuatu	2016	Male	40 to 44	53% (41-65%)	47% (35-59%)	0.69 (0.31-1.3)	1.7% (0.45-3.7%)	1.6% (0.099-4%)	0.97 (0.061-2.4)	60 (16-130)
Vanuatu	2016	Male	45 to 49	46% (32-58%)	54% (42-68%)	0.55 (0.24-1.1)	1.3% (0.2-2.9%)	1.2% (-0.14-3%)	0.85 (-0.11-2.2)	47 (7.3-110)
Vanuatu	2016	Male	50 to 54	42% (29-54%)	58% (46-71%)	0.42 (0.18-0.85)	0.96% (0.0078-2.4%)	0.87% (-0.27-2.5%)	0.79 (-0.25-2.4)	38 (0.37-95)
Vanuatu	2016	Male	55 to 59	41% (27-54%)	59% (46-73%)	0.33 (0.14-0.64)	0.49% (-0.22-1.3%)	0.39% (-0.44-1.3%)	0.37 (-0.42-1.3)	17 (-8-49)
Vanuatu	2016	Male	60 to 64	38% (24-52%)	62% (48-76%)	0.27 (0.12-0.51)	0.28% (-0.32-0.98%)	0.22% (-0.49-1%)	0.24 (-0.51-1.1)	9.6 (-10-33)
Vanuatu	2016	Male	65 to 69	36% (21-54%)	64% (46-79%)	0.24 (0.1-0.48)	0.23% (-0.26-0.79%)	0.19% (-0.35-0.86%)	0.21 (-0.39-0.95)	6.3 (-7.5-24)
Vanuatu	2016	Male	70 to 74	33% (20-50%)	67% (50-80%)	0.2 (0.081-0.42)	0.095% (-0.31-0.64%)	0.061% (-0.42-0.68%)	0.08 (-0.55-0.91)	2.6 (-8.7-18)
Vanuatu	2016	Male	75 to 79	29% (17-46%)	71% (54-83%)	0.17 (0.061-0.35)	0.052% (-0.33-0.47%)	0.012% (-0.44-0.47%)	0.013 (-0.46-0.5)	0.88 (-5.9-7.9)
Vanuatu	2016	Male	80 to 84	26% (14-42%)	74% (58-86%)	0.15 (0.055-0.29)	0.084% (-0.25-0.37%)	0.051% (-0.33-0.38%)	0.034 (-0.22-0.25)	0.67 (-1.9-3)
Vanuatu	2016	Male	85 to 89	25% (14-40%)	75% (60-86%)	0.14 (0.057-0.3)	-0.012% (-0.34-0.26%)	-0.061% (-0.43-0.23%)	-0.02 (-0.14-0.076)	-0.033 (-0.93-0.72)
Vanuatu	2016	Male	90 to 94	26% (15-42%)	74% (58-85%)	0.15 (0.055-0.31)	-0.013% (-0.31-0.22%)	-0.067% (-0.41-0.21%)	-0.0079 (-0.049-0.025)	-0.0088 (-0.22-0.16)
Vanuatu	2016	Male	95 plus	30% (15-49%)	70% (51-85%)	0.16 (0.055-0.35)	-0.17% (-0.56-0.12%)	-0.28% (-0.76-0.056%)	-0.0092 (-0.025-0.0018)	-0.021 (-0.072-0.016)
Armenia	1990	Female	All Ages	30% (26-35%)	70% (65-74%)	1 (0.78-1.3)	2.4% (1.6-3.4%)	3.8% (2-6.1%)	450 (230-730)	13000 (8200-18000)
Armenia	1990	Female	15 to 19	25% (18-34%)	75% (66-82%)	0.78 (0.45-1.2)	1.6% (1-2.3%)	3.2% (1.9-5%)	1.7 (0.97-2.6)	220 (140-320)
Armenia	1990	Female	20 to 24	36% (25-47%)	64% (53-75%)	1.2 (0.73-1.9)	2.8% (2-3.8%)	4% (2.4-6.2%)	3.5 (2.1-5.4)	470 (310-680)
Armenia	1990	Female	25 to 29	41% (29-52%)	59% (48-71%)	1.5 (0.89-2.2)	3.1% (2.2-4.1%)	4.6% (3-6.7%)	5.7 (3.6-8.3)	700 (480-960)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	1990	Female	30 to 34	44% (31-56%)	56% (44-69%)	1.6 (0.98-2.4)	3.2% (2.4-4.3%)	5.3% (3.5-7.6%)	8.3 (5.5-12)	820 (600-1100)
Armenia	1990	Female	35 to 39	44% (32-55%)	56% (45-68%)	1.7 (1-2.6)	3.4% (2.4-4.6%)	6% (3.9-8.3%)	9.9 (6.5-14)	770 (540-1000)
Armenia	1990	Female	40 to 44	43% (32-54%)	57% (46-68%)	1.8 (1.1-2.7)	3.7% (2.5-5.1%)	6.4% (4.1-9.1%)	11 (6.9-15)	660 (450-930)
Armenia	1990	Female	45 to 49	42% (28-53%)	58% (47-72%)	1.8 (0.99-2.8)	4% (2.5-5.9%)	6.6% (4.1-10%)	12 (7-18)	590 (380-880)
Armenia	1990	Female	50 to 54	39% (28-52%)	61% (48-72%)	1.7 (1-2.7)	4.2% (2.6-6.4%)	6.4% (3.9-9.9%)	31 (18-49)	1300 (820-2100)
Armenia	1990	Female	55 to 59	37% (22-53%)	63% (47-78%)	1.7 (0.91-2.8)	4.3% (2.3-7%)	6.2% (3.2-10%)	38 (19-64)	1400 (710-2200)
Armenia	1990	Female	60 to 64	34% (20-51%)	66% (49-80%)	1.6 (0.83-2.7)	4.3% (1.9-7.4%)	5.8% (2.5-10%)	62 (27-110)	1800 (810-3200)
Armenia	1990	Female	65 to 69	32% (20-47%)	68% (53-80%)	1.5 (0.8-2.4)	4.3% (2-7.4%)	5.6% (2.6-9.7%)	56 (26-95)	1400 (640-2300)
Armenia	1990	Female	70 to 74	28% (18-42%)	72% (58-82%)	1.3 (0.69-2.2)	3.8% (1.3-7.2%)	4.8% (1.6-9.1%)	43 (14-82)	820 (280-1500)
Armenia	1990	Female	75 to 79	24% (15-38%)	76% (62-85%)	1.1 (0.54-1.9)	3.2% (0.77-6.7%)	3.8% (0.87-8%)	59 (13-130)	880 (210-1800)
Armenia	1990	Female	80 to 84	22% (13-33%)	78% (67-87%)	0.99 (0.52-1.6)	2.8% (0.73-5.9%)	3.3% (0.77-7%)	51 (12-110)	550 (140-1200)
Armenia	1990	Female	85 to 89	21% (13-32%)	79% (68-87%)	0.98 (0.5-1.7)	2.5% (0.46-6%)	2.9% (0.4-7.2%)	37 (5.1-93)	280 (53-690)
Armenia	1990	Female	90 to 94	23% (14-34%)	77% (66-86%)	1 (0.52-2)	2.2% (0.13-6.2%)	2.6% (-0.034-7.4%)	14 (-0.19-41)	77 (4.3-210)
Armenia	1990	Female	95 plus	25% (14-40%)	75% (60-86%)	1.1 (0.46-2.2)	2.2% (-0.32-6.8%)	2.5% (-0.6-8.1%)	4.1 (-0.95-13)	15 (-2.1-46)
Armenia	1990	Male	All Ages	79% (76-83%)	21% (17-24%)	1.2 (0.95-1.6)	2.5% (1.4-3.7%)	1.2% (-0.63-3.2%)	160 (-83-420)	16000 (8900-25000)
Armenia	1990	Male	15 to 19	68% (58-76%)	32% (24-42%)	0.57 (0.31-0.95)	3.8% (2.2-5.8%)	4.4% (1.9-7.9%)	8.4 (3.5-15)	870 (500-1400)
Armenia	1990	Male	20 to 24	90% (84-94%)	10% (6.4-16%)	1.3 (0.8-2)	7.9% (5.3-11%)	8.4% (4.5-13%)	22 (11-34)	2100 (1400-3000)
Armenia	1990	Male	25 to 29	93% (89-95%)	7.4% (4.9-11%)	2 (1.2-3)	9.1% (6.4-12%)	10% (6-15%)	32 (19-50)	3000 (2000-4100)
Armenia	1990	Male	30 to 34	94% (91-96%)	5.8% (3.6-9.3%)	2.3 (1.5-3.3)	8.3% (5.6-11%)	9.3% (5.3-14%)	37 (21-56)	3000 (2000-4200)
Armenia	1990	Male	35 to 39	95% (93-97%)	4.9% (3.1-7.5%)	2.3 (1.4-3.4)	6.6% (4.1-9.5%)	7.3% (3.5-12%)	28 (13-44)	2100 (1200-3000)
Armenia	1990	Male	40 to 44	95% (93-97%)	4.8% (2.9-7.4%)	2.3 (1.4-3.5)	5.2% (2.8-8%)	5.5% (2.1-9.4%)	20 (7.5-35)	1300 (700-2100)
Armenia	1990	Male	45 to 49	95% (91-97%)	5.4% (3.3-8.8%)	2.5 (1.5-3.8)	4.7% (1.8-8%)	4.9% (0.96-9.4%)	18 (3.4-35)	990 (390-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	1990	Male	50 to 54	93% (90-96%)	6.5% (4-10%)	2.4 (1.5-3.5)	3.6% (0.87-6.9%)	3.6% (-0.089-7.8%)	35 (-0.85-77)	1700 (430-3300)
Armenia	1990	Male	55 to 59	92% (85-96%)	8.2% (4.1-15%)	2.1 (1.3-3.1)	2.4% (0.13-5.1%)	2.3% (-0.66-5.6%)	25 (-7.1-60)	1100 (57-2300)
Armenia	1990	Male	60 to 64	90% (82-95%)	9.9% (4.7-18%)	1.8 (1.1-2.7)	1.3% (-0.97-3.9%)	1% (-1.7-4.2%)	17 (-29-72)	720 (-580-2300)
Armenia	1990	Male	65 to 69	89% (78-95%)	11% (5-22%)	1.6 (0.99-2.3)	0.4% (-1.7-2.6%)	0.11% (-2.5-2.8%)	1.2 (-27-31)	120 (-510-800)
Armenia	1990	Male	70 to 74	87% (77-93%)	13% (6.9-23%)	1.3 (0.78-2.1)	-0.4% (-2.5-1.8%)	-0.77% (-3.3-1.9%)	-5.4 (-23-13)	-61 (-390-280)
Armenia	1990	Male	75 to 79	85% (74-92%)	15% (7.7-26%)	1.1 (0.64-1.8)	-1.1% (-3.1-0.93%)	-1.5% (-4-0.96%)	-17 (-44-10)	-190 (-580-180)
Armenia	1990	Male	80 to 84	83% (71-91%)	17% (9.1-29%)	1 (0.6-1.6)	-1.7% (-3.9-0.33%)	-2.3% (-5-0.2%)	-23 (-49-1.9)	-210 (-470-39)
Armenia	1990	Male	85 to 89	81% (70-90%)	19% (10-30%)	0.95 (0.56-1.5)	-2.1% (-4.3--0.17%)	-2.8% (-5.6--0.49%)	-22 (-43--3.7)	-140 (-300--12)
Armenia	1990	Male	90 to 94	81% (70-90%)	19% (10-30%)	0.89 (0.51-1.5)	-2.2% (-4.6--0.27%)	-3.1% (-6--0.64%)	-9.2 (-18--1.9)	-41 (-84--4.9)
Armenia	1990	Male	95 plus	80% (66-91%)	20% (8.9-34%)	0.81 (0.4-1.5)	-2.1% (-4.4--0.15%)	-3% (-6--0.65%)	-2.5 (-4.9--0.53)	-7 (-15--0.52)
Armenia	1995	Female	All Ages	27% (24-30%)	73% (70-76%)	0.94 (0.75-1.1)	2.8% (1.9-4%)	4.2% (2.3-6.8%)	510 (280-830)	13000 (8900-19000)
Armenia	1995	Female	15 to 19	21% (18-26%)	79% (74-82%)	0.65 (0.42-0.94)	1.4% (0.98-1.9%)	2.8% (1.8-4.1%)	1.3 (0.84-1.9)	190 (130-270)
Armenia	1995	Female	20 to 24	29% (23-35%)	71% (65-77%)	0.98 (0.65-1.4)	2.6% (1.9-3.5%)	3.6% (2.3-5.4%)	2.4 (1.6-3.7)	340 (240-480)
Armenia	1995	Female	25 to 29	35% (29-42%)	65% (58-71%)	1.2 (0.85-1.7)	3% (2.3-3.9%)	4.4% (3-6.2%)	3.6 (2.4-5.1)	450 (320-610)
Armenia	1995	Female	30 to 34	38% (32-44%)	62% (56-68%)	1.4 (0.95-1.9)	3.1% (2.4-4%)	5% (3.6-6.8%)	7 (4.8-9.4)	690 (510-910)
Armenia	1995	Female	35 to 39	37% (31-44%)	63% (56-69%)	1.4 (0.95-1.9)	3.2% (2.4-4.1%)	5.5% (4-7.4%)	11 (7.5-14)	810 (610-1100)
Armenia	1995	Female	40 to 44	37% (31-44%)	63% (56-69%)	1.5 (1-2)	3.5% (2.5-4.5%)	5.9% (4.3-7.9%)	13 (9.5-18)	800 (580-1100)
Armenia	1995	Female	45 to 49	37% (31-45%)	63% (55-69%)	1.6 (1-2.3)	3.8% (2.6-5.4%)	6.2% (4.1-9.1%)	14 (9.4-21)	720 (480-1000)
Armenia	1995	Female	50 to 54	37% (27-49%)	63% (51-73%)	1.6 (0.99-2.5)	4.3% (2.6-6.4%)	6.5% (4-9.9%)	16 (9.7-25)	680 (430-1000)
Armenia	1995	Female	55 to 59	37% (23-54%)	63% (46-77%)	1.7 (0.93-2.7)	4.6% (2.3-7.4%)	6.5% (3.3-11%)	49 (25-81)	1700 (900-2800)
Armenia	1995	Female	60 to 64	34% (21-53%)	66% (47-79%)	1.6 (0.87-2.7)	4.4% (2-7.6%)	5.9% (2.6-10%)	57 (25-96)	1700 (740-2800)
Armenia	1995	Female	65 to 69	32% (20-48%)	68% (52-80%)	1.5 (0.85-2.4)	4.5% (2.2-7.6%)	5.7% (2.8-9.6%)	96 (46-160)	2300 (1100-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	1995	Female	70 to 74	28% (18-42%)	72% (58-82%)	1.3 (0.72-2.1)	4% (1.5-7.6%)	4.9% (1.9-9.5%)	72 (27-140)	1400 (530-2600)
Armenia	1995	Female	75 to 79	24% (15-37%)	76% (63-85%)	1.1 (0.57-1.9)	3.3% (0.9-6.9%)	3.9% (0.98-8.4%)	48 (12-100)	710 (190-1500)
Armenia	1995	Female	80 to 84	22% (13-33%)	78% (67-87%)	1 (0.53-1.7)	2.9% (0.79-5.8%)	3.4% (0.82-7%)	48 (12-100)	520 (140-1100)
Armenia	1995	Female	85 to 89	22% (13-33%)	78% (67-87%)	1 (0.51-1.7)	2.6% (0.56-5.9%)	3% (0.53-7%)	45 (8.2-100)	340 (73-790)
Armenia	1995	Female	90 to 94	23% (14-33%)	77% (67-86%)	1 (0.53-1.9)	2.3% (0.21-6%)	2.6% (0.059-7.4%)	20 (0.44-55)	110 (9.9-280)
Armenia	1995	Female	95 plus	25% (14-40%)	75% (60-86%)	1.1 (0.5-2.2)	2.3% (-0.3-7.1%)	2.6% (-0.6-8.4%)	6.6 (-1.5-21)	24 (-3-72)
Armenia	1995	Male	All Ages	82% (79-84%)	18% (16-21%)	1.3 (1.1-1.6)	2.4% (1.1-3.7%)	0.93% (-1.1-3%)	140 (-160-440)	15000 (6700-23000)
Armenia	1995	Male	15 to 19	74% (67-79%)	26% (21-33%)	0.66 (0.38-1.1)	3.8% (2.2-5.8%)	4.4% (2.1-7.4%)	9.4 (4.3-16)	910 (520-1400)
Armenia	1995	Male	20 to 24	92% (88-94%)	8.1% (5.6-12%)	1.4 (0.87-2.1)	7.7% (5.1-11%)	8.2% (4.6-12%)	19 (11-29)	1800 (1200-2500)
Armenia	1995	Male	25 to 29	94% (91-96%)	6.3% (4.3-9.1%)	2.1 (1.3-3.1)	9% (6.5-12%)	9.9% (6.3-14%)	23 (15-34)	2000 (1400-2700)
Armenia	1995	Male	30 to 34	95% (93-97%)	5% (3.2-7.4%)	2.4 (1.7-3.4)	8.3% (5.7-11%)	9.2% (5.6-13%)	34 (21-51)	2700 (1800-3600)
Armenia	1995	Male	35 to 39	96% (94-97%)	4.2% (2.8-6%)	2.4 (1.5-3.4)	6.5% (3.9-9.2%)	7.1% (3.6-11%)	34 (16-53)	2400 (1400-3400)
Armenia	1995	Male	40 to 44	96% (94-97%)	4.1% (2.8-6%)	2.4 (1.5-3.4)	4.9% (2.4-7.7%)	5.2% (1.6-8.9%)	28 (8.4-47)	1700 (830-2700)
Armenia	1995	Male	45 to 49	95% (93-97%)	4.6% (2.9-6.9%)	2.6 (1.7-3.9)	4.4% (1.4-8.1%)	4.5% (0.69-9.4%)	24 (3.8-50)	1300 (420-2400)
Armenia	1995	Male	50 to 54	94% (92-96%)	5.7% (3.6-8.2%)	2.5 (1.6-3.7)	3.5% (0.53-7.1%)	3.5% (-0.27-8.1%)	19 (-1.4-43)	880 (130-1800)
Armenia	1995	Male	55 to 59	92% (86-96%)	7.7% (4.1-14%)	2.1 (1.4-2.9)	2.1% (-0.32-4.8%)	1.9% (-1.5-3%)	28 (-15-76)	1200 (-190-2800)
Armenia	1995	Male	60 to 64	90% (82-95%)	9.8% (4.7-18%)	1.8 (1.1-2.6)	1% (-1.3-3.6%)	0.8% (-1.9-3.9%)	14 (-34-68)	590 (-730-2100)
Armenia	1995	Male	65 to 69	89% (79-95%)	11% (5-21%)	1.6 (1-2.2)	0.12% (-2.1-2.4%)	-0.16% (-2.8-2.6%)	-3.9 (-68-63)	79 (-1400-1600)
Armenia	1995	Male	70 to 74	87% (77-93%)	13% (6.8-23%)	1.3 (0.82-2)	-0.68% (-2.9-1.6%)	-1% (-3.6-1.6%)	-15 (-53-23)	-210 (-870-460)
Armenia	1995	Male	75 to 79	85% (75-93%)	15% (7.3-25%)	1.1 (0.66-1.8)	-1.3% (-3.5-0.78%)	-1.7% (-4.3-0.71%)	-14 (-35-5.9)	-180 (-470-100)
Armenia	1995	Male	80 to 84	83% (71-91%)	17% (8.9-29%)	1 (0.62-1.6)	-1.8% (-4.2-0.25%)	-2.5% (-5.3-0.13%)	-20 (-43-1.1)	-190 (-420-25)
Armenia	1995	Male	85 to 89	82% (71-90%)	18% (10-29%)	0.95 (0.58-1.4)	-2.3% (-4.6--0.31%)	-3% (-5.9--0.54%)	-27 (-52--4.9)	-180 (-360--24)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	1995	Male	90 to 94	81% (70-90%)	19% (10-30%)	0.89 (0.52-1.4)	-2.3% (-4.7--0.35%)	-3.2% (-6.1--0.74%)	-12 (-24--2.9)	-56 (-110--8.1)
Armenia	1995	Male	95 plus	80% (66-91%)	20% (8.7-34%)	0.81 (0.4-1.4)	-2.2% (-4.7--0.22%)	-3.2% (-6.2--0.68%)	-3.4 (-6.5--0.73)	-9.7 (-20--0.93)
Armenia	2000	Female	All Ages	27% (24-31%)	73% (69-76%)	0.97 (0.78-1.1)	3% (1.9-4.3%)	4.1% (2.2-6.9%)	500 (260-830)	13000 (8500-19000)
Armenia	2000	Female	15 to 19	21% (17-26%)	79% (74-83%)	0.63 (0.42-0.91)	1.3% (0.92-1.8%)	2.8% (1.8-4%)	1.1 (0.73-1.6)	190 (120-260)
Armenia	2000	Female	20 to 24	29% (24-35%)	71% (65-76%)	0.97 (0.66-1.4)	2.4% (1.8-3.3%)	3.7% (2.4-5.4%)	1.8 (1.2-2.7)	320 (220-460)
Armenia	2000	Female	25 to 29	35% (30-41%)	65% (59-70%)	1.2 (0.85-1.7)	2.9% (2.2-3.8%)	4.5% (3.1-6.2%)	2.8 (1.9-3.9)	370 (260-490)
Armenia	2000	Female	30 to 34	38% (32-44%)	62% (56-68%)	1.4 (0.97-1.8)	3% (2.3-3.9%)	5.1% (3.7-6.9%)	4.1 (2.9-5.5)	430 (320-580)
Armenia	2000	Female	35 to 39	37% (32-43%)	63% (57-68%)	1.4 (0.97-1.9)	3.1% (2.3-3.9%)	5.5% (4.1-7.2%)	8.4 (6.2-11)	680 (520-880)
Armenia	2000	Female	40 to 44	37% (32-43%)	63% (57-68%)	1.4 (1-1.9)	3.3% (2.5-4.3%)	5.9% (4.5-7.7%)	14 (10-18)	870 (670-1100)
Armenia	2000	Female	45 to 49	37% (31-44%)	63% (56-69%)	1.5 (1-2.2)	3.7% (2.5-5.1%)	6.3% (4.3-8.7%)	18 (12-26)	930 (630-1300)
Armenia	2000	Female	50 to 54	37% (27-49%)	63% (51-73%)	1.6 (0.96-2.5)	4.1% (2.6-6.2%)	6.5% (4.1-9.7%)	20 (12-30)	830 (530-1200)
Armenia	2000	Female	55 to 59	36% (22-54%)	64% (46-78%)	1.6 (0.89-2.7)	4.4% (2.3-7.3%)	6.4% (3.3-10%)	22 (11-36)	770 (400-1300)
Armenia	2000	Female	60 to 64	34% (20-52%)	66% (48-80%)	1.6 (0.83-2.6)	4.2% (1.7-7.5%)	5.7% (2.4-10%)	61 (26-110)	1800 (790-3300)
Armenia	2000	Female	65 to 69	31% (19-48%)	69% (52-81%)	1.4 (0.84-2.3)	4.2% (2-7.2%)	5.5% (2.7-9.4%)	71 (34-120)	1700 (840-2900)
Armenia	2000	Female	70 to 74	27% (17-42%)	73% (58-83%)	1.2 (0.69-2.1)	3.9% (1.5-7.4%)	4.8% (1.8-9.2%)	110 (40-210)	2100 (800-4000)
Armenia	2000	Female	75 to 79	23% (15-37%)	77% (63-85%)	1.1 (0.56-1.8)	3.2% (0.84-6.6%)	3.8% (1-7.9%)	69 (17-150)	1000 (260-2100)
Armenia	2000	Female	80 to 84	21% (13-33%)	79% (67-87%)	0.97 (0.52-1.6)	2.7% (0.85-5.4%)	3.2% (0.94-6.6%)	43 (13-89)	480 (150-960)
Armenia	2000	Female	85 to 89	21% (13-32%)	79% (68-87%)	0.97 (0.5-1.7)	2.4% (0.52-5.5%)	2.8% (0.53-6.7%)	33 (6.3-78)	260 (57-590)
Armenia	2000	Female	90 to 94	22% (14-33%)	78% (67-86%)	1 (0.51-1.8)	2.1% (0.14-5.7%)	2.4% (-0.025-6.8%)	19 (-0.2-55)	100 (6.9-280)
Armenia	2000	Female	95 plus	25% (14-40%)	75% (60-86%)	1.1 (0.48-2.1)	2.1% (-0.47-6.8%)	2.4% (-0.78-8.1%)	7.2 (-2.4-25)	26 (-5.8-84)
Armenia	2000	Male	All Ages	83% (80-86%)	17% (14-20%)	1.3 (1.1-1.6)	2.1% (0.76-3.4%)	0.48% (-1.5-2.5%)	63 (-200-330)	10000 (3900-17000)
Armenia	2000	Male	15 to 19	76% (71-80%)	24% (20-29%)	0.68 (0.4-1.1)	3.5% (2.3-5%)	4.1% (2.1-6.9%)	6.1 (3.2-10)	700 (450-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	2000	Male	20 to 24	92% (89-95%)	7.5% (5.3-11%)	1.4 (0.9-2.1)	7.2% (5.1-9.4%)	8% (4.7-12%)	11 (6.3-16)	1200 (840-1600)
Armenia	2000	Male	25 to 29	94% (92-96%)	5.8% (4.2-8%)	2.1 (1.3-3.1)	8.4% (6.2-11%)	9.8% (6.1-14%)	13 (7.9-19)	1300 (920-1700)
Armenia	2000	Male	30 to 34	95% (93-97%)	4.6% (3.1-6.7%)	2.4 (1.7-3.5)	8.1% (5.8-11%)	9.6% (6-14%)	17 (11-26)	1400 (1000-1900)
Armenia	2000	Male	35 to 39	96% (95-97%)	4% (2.7-5.3%)	2.4 (1.5-3.4)	6.6% (4.2-9.1%)	7.5% (3.9-11%)	23 (12-36)	1700 (1100-2500)
Armenia	2000	Male	40 to 44	96% (95-97%)	3.8% (2.6-5.4%)	2.4 (1.5-3.4)	4.9% (2.5-7.6%)	5.4% (1.8-8.9%)	26 (8.5-44)	1700 (820-2600)
Armenia	2000	Male	45 to 49	96% (94-97%)	4.2% (2.8-6.2%)	2.6 (1.8-3.8)	4% (1.4-7.5%)	4.1% (0.54-8.8%)	24 (3-51)	1300 (460-2500)
Armenia	2000	Male	50 to 54	95% (92-97%)	5.2% (3.5-7.6%)	2.5 (1.6-3.6)	3.2% (0.4-6.6%)	3.1% (-0.51-7.5%)	18 (-3-45)	900 (120-1900)
Armenia	2000	Male	55 to 59	93% (87-96%)	7.3% (3.8-13%)	2.1 (1.4-2.9)	1.8% (-0.62-4.4%)	1.6% (-1.4-4.8%)	9.1 (-8.2-28)	420 (-150-1000)
Armenia	2000	Male	60 to 64	90% (82-95%)	9.8% (4.6-18%)	1.7 (1.1-2.6)	0.92% (-1.3-3.4%)	0.69% (-2-3.7%)	12 (-35-62)	520 (-710-1900)
Armenia	2000	Male	65 to 69	89% (79-95%)	11% (5.2-21%)	1.5 (1-2.2)	0.072% (-2-2.3%)	-0.21% (-2.7-2.4%)	-4 (-52-47)	38 (-1000-1200)
Armenia	2000	Male	70 to 74	87% (77-93%)	13% (6.6-23%)	1.3 (0.8-2)	-0.63% (-2.7-1.5%)	-0.97% (-3.5-1.4%)	-26 (-93-39)	-360 (-1600-830)
Armenia	2000	Male	75 to 79	85% (73-92%)	15% (7.7-27%)	1.1 (0.67-1.8)	-1.2% (-3.2-0.76%)	-1.6% (-4-0.73%)	-21 (-54-9.8)	-260 (-720-170)
Armenia	2000	Male	80 to 84	83% (71-91%)	17% (9-29%)	1 (0.61-1.5)	-1.6% (-3.8-0.31%)	-2.3% (-5.1-0.13%)	-15 (-33-0.86)	-140 (-320-27)
Armenia	2000	Male	85 to 89	82% (70-90%)	18% (10-30%)	0.94 (0.59-1.4)	-2.2% (-4.4--0.22%)	-3% (-5.8--0.54%)	-15 (-30--2.7)	-100 (-210--10)
Armenia	2000	Male	90 to 94	81% (70-90%)	19% (10-30%)	0.88 (0.51-1.4)	-2.4% (-4.7--0.41%)	-3.3% (-6.1--0.8%)	-12 (-22--2.8)	-52 (-100--8.9)
Armenia	2000	Male	95 plus	80% (66-91%)	20% (8.7-34%)	0.8 (0.4-1.4)	-2.3% (-4.7--0.33%)	-3.3% (-6.3--0.74%)	-3.8 (-7.3--0.86)	-11 (-22--1.6)
Armenia	2005	Female	All Ages	28% (26-32%)	72% (68-74%)	1.1 (0.86-1.3)	3.3% (2.2-4.7%)	4.5% (2.5-7.4%)	570 (310-940)	14000 (9200-20000)
Armenia	2005	Female	15 to 19	22% (18-27%)	78% (73-82%)	0.67 (0.43-0.94)	1.4% (0.98-1.9%)	2.9% (1.9-4.1%)	1.2 (0.76-1.7)	190 (130-270)
Armenia	2005	Female	20 to 24	29% (24-36%)	71% (64-76%)	1 (0.67-1.5)	2.5% (1.8-3.4%)	3.8% (2.5-5.5%)	1.9 (1.2-2.8)	350 (240-500)
Armenia	2005	Female	25 to 29	36% (31-42%)	64% (58-69%)	1.3 (0.9-1.7)	2.9% (2.2-3.7%)	4.6% (3.2-6.5%)	2.7 (1.8-3.8)	390 (280-530)
Armenia	2005	Female	30 to 34	38% (33-45%)	62% (55-67%)	1.4 (1-2)	3% (2.3-3.9%)	5.3% (3.9-7.2%)	3.6 (2.6-5)	390 (300-530)
Armenia	2005	Female	35 to 39	37% (32-43%)	63% (57-68%)	1.4 (1-2)	3.1% (2.4-4%)	5.7% (4.2-7.6%)	5.8 (4.3-7.8)	480 (370-630)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	2005	Female	40 to 44	38% (32-44%)	62% (56-68%)	1.5 (1.1-2)	3.4% (2.6-4.4%)	6.1% (4.6-7.9%)	13 (9.6-17)	820 (620-1100)
Armenia	2005	Female	45 to 49	38% (31-44%)	62% (56-69%)	1.6 (1.1-2.2)	3.7% (2.6-5.1%)	6.4% (4.5-9%)	21 (14-30)	1100 (760-1500)
Armenia	2005	Female	50 to 54	38% (27-50%)	62% (50-73%)	1.7 (1-2.5)	4.3% (2.8-6.3%)	6.7% (4.4-9.9%)	28 (18-42)	1200 (770-1800)
Armenia	2005	Female	55 to 59	37% (23-54%)	63% (46-77%)	1.7 (0.92-2.9)	4.7% (2.4-7.8%)	6.8% (3.6-11%)	29 (16-49)	1000 (560-1700)
Armenia	2005	Female	60 to 64	34% (21-52%)	66% (48-79%)	1.6 (0.87-2.8)	4.5% (2-7.9%)	6.2% (2.8-11%)	32 (15-56)	960 (440-1700)
Armenia	2005	Female	65 to 69	32% (20-48%)	68% (52-80%)	1.5 (0.87-2.4)	4.5% (2.2-7.6%)	5.9% (3-9.9%)	88 (45-150)	2100 (1100-3600)
Armenia	2005	Female	70 to 74	28% (17-42%)	72% (58-83%)	1.3 (0.71-2.2)	4.2% (1.8-7.8%)	5.2% (2.2-9.8%)	96 (40-180)	1800 (760-3400)
Armenia	2005	Female	75 to 79	24% (15-37%)	76% (63-85%)	1.1 (0.58-1.9)	3.5% (1.1-6.8%)	4.2% (1.4-8.3%)	110 (36-220)	1600 (540-3200)
Armenia	2005	Female	80 to 84	22% (13-33%)	78% (67-87%)	1 (0.55-1.7)	3.1% (1.2-5.9%)	3.7% (1.4-7.1%)	80 (29-150)	870 (330-1600)
Armenia	2005	Female	85 to 89	22% (13-33%)	78% (67-87%)	1 (0.54-1.8)	2.7% (0.89-5.8%)	3.2% (0.95-7.1%)	33 (10-75)	260 (85-580)
Armenia	2005	Female	90 to 94	23% (14-34%)	77% (66-86%)	1.1 (0.54-1.9)	2.4% (0.4-6.1%)	2.8% (0.34-7.6%)	17 (2-44)	89 (15-230)
Armenia	2005	Female	95 plus	26% (15-41%)	74% (59-85%)	1.2 (0.5-2.2)	2.4% (-0.2-6.9%)	2.8% (-0.47-8.2%)	7.5 (-1.3-22)	27 (-2.3-77)
Armenia	2005	Male	All Ages	84% (82-87%)	16% (13-18%)	1.5 (1.2-1.7)	2.3% (1-3.7%)	0.77% (-1.3-2.9%)	110 (-180-400)	12000 (5200-19000)
Armenia	2005	Male	15 to 19	76% (69-80%)	24% (20-31%)	0.69 (0.4-1.1)	3.3% (2.2-4.6%)	4.1% (2.2-6.4%)	5 (2.6-8.1)	640 (420-910)
Armenia	2005	Male	20 to 24	92% (89-95%)	7.6% (5.2-11%)	1.5 (0.94-2.2)	7.1% (5.1-9.2%)	8.1% (4.9-12%)	11 (6.6-16)	1300 (910-1800)
Armenia	2005	Male	25 to 29	94% (92-96%)	5.9% (4.2-8.3%)	2.2 (1.4-3.2)	8.3% (6-11%)	9.9% (6-14%)	14 (8.3-20)	1400 (1000-1900)
Armenia	2005	Male	30 to 34	95% (93-97%)	4.7% (3.1-6.7%)	2.5 (1.7-3.6)	8.2% (5.9-11%)	10% (6.1-14%)	17 (10-25)	1400 (990-1900)
Armenia	2005	Male	35 to 39	96% (94-97%)	4% (2.7-5.8%)	2.4 (1.6-3.6)	7.1% (4.6-10%)	8.6% (4.5-13%)	20 (10-30)	1400 (900-2000)
Armenia	2005	Male	40 to 44	96% (94-97%)	3.9% (2.6-5.7%)	2.5 (1.6-3.5)	5.5% (3-8.2%)	6.2% (2.5-10%)	28 (11-46)	1800 (930-2600)
Armenia	2005	Male	45 to 49	96% (94-97%)	4.3% (2.9-6.4%)	2.7 (1.8-3.8)	4.6% (2-7.8%)	4.9% (1.3-9.5%)	34 (8.6-66)	1900 (780-3200)
Armenia	2005	Male	50 to 54	95% (92-97%)	5.3% (3.5-7.8%)	2.6 (1.7-3.6)	3.5% (0.7-6.9%)	3.6% (-0.15-8%)	30 (-1.2-68)	1400 (300-2900)
Armenia	2005	Male	55 to 59	93% (87-96%)	7.4% (4-13%)	2.2 (1.4-3)	2.1% (-0.33-5%)	2% (-1.1-5.5%)	16 (-8.5-45)	700 (-100-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	2005	Male	60 to 64	90% (82-95%)	9.8% (4.7-18%)	1.8 (1.1-2.7)	1.3% (-0.9-3.8%)	1.1% (-1.6-4.2%)	9.5 (-14-36)	360 (-260-1100)
Armenia	2005	Male	65 to 69	89% (79-95%)	11% (5.1-21%)	1.6 (1.1-2.3)	0.44% (-1.6-2.7%)	0.24% (-2.2-2.9%)	5.2 (-47-63)	260 (-910-1600)
Armenia	2005	Male	70 to 74	87% (77-93%)	13% (6.9-23%)	1.4 (0.85-2.1)	-0.25% (-2.3-1.9%)	-0.51% (-3-2%)	-12 (-69-48)	-120 (-1100-970)
Armenia	2005	Male	75 to 79	85% (74-92%)	15% (7.6-26%)	1.2 (0.7-1.8)	-0.79% (-2.7-1.2%)	-1.1% (-3.5-1.2%)	-29 (-90-32)	-340 (-1200-520)
Armenia	2005	Male	80 to 84	83% (71-91%)	17% (8.9-29%)	1 (0.65-1.6)	-1.3% (-3.3-0.59%)	-1.8% (-4.4-0.52%)	-23 (-56-6.9)	-200 (-540-93)
Armenia	2005	Male	85 to 89	82% (70-90%)	18% (10-30%)	0.97 (0.6-1.4)	-1.6% (-3.8-0.16%)	-2.3% (-5--0.027%)	-9.1 (-20--0.1)	-59 (-130-5.9)
Armenia	2005	Male	90 to 94	81% (70-90%)	19% (10-30%)	0.9 (0.55-1.4)	-2.1% (-4.3--0.22%)	-2.9% (-5.6--0.53%)	-5.7 (-11--1.1)	-25 (-52--2.6)
Armenia	2005	Male	95 plus	80% (66-91%)	20% (9.4-34%)	0.82 (0.43-1.4)	-2.1% (-4.5--0.18%)	-3% (-5.8--0.57%)	-2.6 (-5.1--0.49)	-7.4 (-16--0.62)
Armenia	2010	Female	All Ages	32% (28-37%)	68% (63-72%)	1.3 (1.1-1.7)	3.7% (2.5-5.3%)	5.1% (2.9-8.2%)	670 (390-1100)	15000 (10000-22000)
Armenia	2010	Female	15 to 19	25% (17-35%)	75% (65-83%)	0.87 (0.51-1.3)	1.6% (1.1-2.2%)	3.4% (2.1-5.1%)	1.1 (0.71-1.7)	170 (110-240)
Armenia	2010	Female	20 to 24	36% (25-47%)	64% (53-75%)	1.4 (0.84-2.1)	2.9% (2.1-4%)	4.7% (2.9-7.2%)	2.3 (1.4-3.6)	390 (260-550)
Armenia	2010	Female	25 to 29	41% (29-53%)	59% (47-71%)	1.6 (0.99-2.4)	3.1% (2.3-4%)	5.3% (3.4-7.5%)	2.8 (1.8-4)	420 (290-580)
Armenia	2010	Female	30 to 34	43% (31-57%)	57% (43-69%)	1.8 (1.1-2.7)	3.2% (2.4-4.3%)	6.1% (4.1-8.5%)	4 (2.7-5.7)	450 (320-610)
Armenia	2010	Female	35 to 39	44% (31-56%)	56% (44-69%)	1.9 (1.2-2.9)	3.5% (2.5-4.8%)	6.7% (4.6-9.4%)	5.8 (3.9-8.2)	470 (340-650)
Armenia	2010	Female	40 to 44	43% (32-54%)	57% (46-68%)	2 (1.2-2.9)	3.9% (2.7-5.3%)	7% (4.7-9.6%)	11 (7.3-15)	680 (480-940)
Armenia	2010	Female	45 to 49	42% (29-54%)	58% (46-71%)	2 (1.2-3)	4.3% (2.9-6.2%)	7.5% (4.9-11%)	22 (14-32)	1100 (770-1600)
Armenia	2010	Female	50 to 54	40% (27-53%)	60% (47-73%)	1.9 (1.1-3)	4.7% (3-7%)	7.5% (4.8-11%)	34 (22-51)	1500 (970-2200)
Armenia	2010	Female	55 to 59	37% (23-54%)	63% (46-77%)	1.9 (1-3.2)	5% (2.8-8.1%)	7.3% (4.2-12%)	40 (23-64)	1400 (820-2300)
Armenia	2010	Female	60 to 64	34% (20-52%)	66% (48-80%)	1.8 (0.97-3)	4.9% (2.4-8.2%)	6.7% (3.4-11%)	40 (19-67)	1200 (580-2000)
Armenia	2010	Female	65 to 69	32% (20-49%)	68% (51-80%)	1.7 (0.94-2.7)	5% (2.7-8.2%)	6.6% (3.5-11%)	50 (26-83)	1200 (640-2000)
Armenia	2010	Female	70 to 74	28% (18-43%)	72% (57-82%)	1.4 (0.78-2.4)	4.7% (2.1-8.7%)	5.9% (2.7-11%)	120 (55-230)	2300 (1100-4300)
Armenia	2010	Female	75 to 79	24% (15-37%)	76% (63-85%)	1.2 (0.61-2.1)	4.1% (1.6-7.5%)	4.9% (1.9-9.1%)	110 (43-210)	1600 (640-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	2010	Female	80 to 84	22% (14-33%)	78% (67-86%)	1.1 (0.59-1.9)	3.9% (1.7-6.9%)	4.5% (2-8.2%)	120 (53-220)	1300 (590-2400)
Armenia	2010	Female	85 to 89	21% (13-32%)	79% (68-87%)	1.1 (0.57-1.9)	3.4% (1.3-6.6%)	4.1% (1.6-8.1%)	73 (28-150)	570 (230-1100)
Armenia	2010	Female	90 to 94	22% (14-34%)	78% (66-86%)	1.2 (0.57-2.2)	2.9% (0.78-6.6%)	3.5% (0.85-8.3%)	22 (5.3-52)	120 (32-270)
Armenia	2010	Female	95 plus	25% (15-40%)	75% (60-85%)	1.3 (0.56-2.4)	2.9% (0.27-7.6%)	3.4% (0.086-9.2%)	7.8 (0.2-21)	29 (2.5-74)
Armenia	2010	Male	All Ages	83% (80-85%)	17% (15-20%)	1.6 (1.3-1.9)	2.9% (1.5-4.4%)	1.3% (-0.78-3.5%)	190 (-110-490)	14000 (7300-22000)
Armenia	2010	Male	15 to 19	69% (59-76%)	31% (24-41%)	0.64 (0.35-1)	3.2% (2.2-4.5%)	3.9% (2-6.3%)	4.1 (2.1-6.6)	540 (350-770)
Armenia	2010	Male	20 to 24	90% (84-94%)	10% (6.5-16%)	1.5 (0.92-2.2)	7% (5.1-9.2%)	8% (4.7-12%)	12 (7.3-18)	1500 (1000-2000)
Armenia	2010	Male	25 to 29	93% (88-95%)	7.4% (4.8-12%)	2.2 (1.4-3.3)	8.3% (6.1-11%)	10% (6.1-15%)	16 (9.6-24)	1700 (1200-2300)
Armenia	2010	Male	30 to 34	94% (91-96%)	5.8% (3.5-9.5%)	2.6 (1.7-3.8)	8.3% (6-11%)	10% (6.5-15%)	18 (11-26)	1600 (1100-2100)
Armenia	2010	Male	35 to 39	95% (92-97%)	4.9% (3.1-7.5%)	2.5 (1.5-3.8)	7.4% (4.9-10%)	9.1% (4.9-14%)	20 (10-29)	1400 (920-2000)
Armenia	2010	Male	40 to 44	95% (93-97%)	4.8% (2.8-7.2%)	2.6 (1.5-3.8)	6.1% (3.5-9.2%)	7.2% (3.3-12%)	25 (12-41)	1500 (890-2300)
Armenia	2010	Male	45 to 49	95% (91-97%)	5.4% (3.2-8.5%)	2.7 (1.8-4)	5.5% (2.8-8.9%)	6.3% (2.4-11%)	42 (16-73)	2200 (1100-3500)
Armenia	2010	Male	50 to 54	94% (90-96%)	6.5% (3.9-10%)	2.7 (1.7-3.8)	4.2% (1.3-7.7%)	4.5% (0.69-9%)	44 (6.5-89)	2000 (640-3800)
Armenia	2010	Male	55 to 59	92% (85-96%)	8.2% (4.2-15%)	2.3 (1.5-3.3)	2.8% (0.3-5.8%)	2.8% (-0.35-6.7%)	31 (-3.8-73)	1200 (140-2700)
Armenia	2010	Male	60 to 64	90% (82-95%)	9.8% (4.6-18%)	2 (1.2-3)	1.7% (-0.67-4.7%)	1.6% (-1.3-5.1%)	17 (-14-54)	610 (-240-1700)
Armenia	2010	Male	65 to 69	89% (79-95%)	11% (5.2-21%)	1.8 (1.2-2.6)	1% (-1-3.3%)	0.89% (-1.6-3.7%)	10 (-17-41)	310 (-320-1000)
Armenia	2010	Male	70 to 74	87% (77-93%)	13% (6.8-23%)	1.5 (0.92-2.3)	0.26% (-1.8-2.5%)	0.078% (-2.4-2.8%)	1.9 (-60-69)	140 (-960-1400)
Armenia	2010	Male	75 to 79	85% (73-93%)	15% (7.5-27%)	1.3 (0.75-2)	-0.26% (-2.2-1.7%)	-0.52% (-2.9-1.8%)	-12 (-67-43)	-100 (-850-680)
Armenia	2010	Male	80 to 84	83% (71-91%)	17% (8.7-29%)	1.2 (0.7-1.7)	-0.86% (-3-0.98%)	-1.2% (-3.7-1%)	-24 (-73-19)	-210 (-710-230)
Armenia	2010	Male	85 to 89	82% (70-90%)	18% (10-30%)	1.1 (0.66-1.6)	-1% (-2.9-0.65%)	-1.6% (-4.1-0.54%)	-13 (-35-4.6)	-80 (-230-51)
Armenia	2010	Male	90 to 94	81% (69-90%)	19% (10-31%)	1 (0.59-1.6)	-1.4% (-3.4-0.22%)	-2.2% (-4.8--0.052%)	-3.9 (-8.6--0.09)	-16 (-39-2.6)
Armenia	2010	Male	95 plus	80% (66-91%)	20% (9.1-34%)	0.91 (0.45-1.6)	-1.5% (-3.5-0.27%)	-2.4% (-5--0.15%)	-1.1 (-2.4-0.071)	-3 (-7.2-0.56)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	2016	Female	All Ages	32% (28-37%)	68% (63-72%)	1.6 (1.2-2.2)	4.1% (2.7-5.9%)	5.9% (3.3-9.5%)	780 (430-1300)	17000 (11000-25000)
Armenia	2016	Female	15 to 19	25% (17-34%)	75% (66-83%)	1 (0.55-1.6)	1.7% (1.1-2.5%)	4% (2.3-6.2%)	0.72 (0.42-1.1)	120 (74-180)
Armenia	2016	Female	20 to 24	36% (25-47%)	64% (53-75%)	1.6 (0.95-2.5)	3.1% (2.1-4.3%)	5.4% (3.3-8.4%)	2 (1.2-3.1)	370 (240-530)
Armenia	2016	Female	25 to 29	41% (29-52%)	59% (48-71%)	1.9 (1.1-3)	3.2% (2.4-4.3%)	5.9% (3.7-8.7%)	2.9 (1.8-4.4)	510 (350-720)
Armenia	2016	Female	30 to 34	43% (31-56%)	57% (44-69%)	2.1 (1.2-3.3)	3.4% (2.4-4.5%)	6.8% (4.4-10%)	4.5 (2.9-6.8)	570 (400-780)
Armenia	2016	Female	35 to 39	44% (31-55%)	56% (45-69%)	2.2 (1.3-3.4)	3.5% (2.5-4.9%)	7.5% (4.8-11%)	6.3 (3.9-9.2)	580 (400-810)
Armenia	2016	Female	40 to 44	43% (31-55%)	57% (45-69%)	2.3 (1.3-3.4)	3.9% (2.7-5.5%)	7.8% (5.1-11%)	9.4 (5.9-14)	650 (440-930)
Armenia	2016	Female	45 to 49	42% (28-53%)	58% (47-72%)	2.4 (1.3-3.7)	4.4% (2.9-6.6%)	8.3% (5.4-13%)	15 (9.1-23)	820 (540-1200)
Armenia	2016	Female	50 to 54	39% (27-52%)	61% (48-73%)	2.3 (1.3-3.6)	4.8% (3.1-7.3%)	8.3% (5.3-13%)	32 (20-50)	1500 (940-2300)
Armenia	2016	Female	55 to 59	36% (22-54%)	64% (46-78%)	2.2 (1.2-3.6)	5.1% (3-8.1%)	8.2% (4.8-13%)	52 (29-84)	2000 (1200-3200)
Armenia	2016	Female	60 to 64	34% (21-53%)	66% (47-79%)	2.1 (1.1-3.5)	5.2% (2.8-8.5%)	7.6% (4.2-12%)	68 (36-110)	2200 (1200-3600)
Armenia	2016	Female	65 to 69	32% (20-47%)	68% (53-80%)	1.9 (1.1-3.2)	5.4% (2.9-9%)	7.6% (4.2-13%)	68 (36-120)	1700 (920-2900)
Armenia	2016	Female	70 to 74	28% (17-42%)	72% (58-83%)	1.7 (0.87-2.9)	5.1% (2.4-9.3%)	6.7% (3.1-12%)	56 (26-100)	1100 (510-2000)
Armenia	2016	Female	75 to 79	23% (15-37%)	77% (63-85%)	1.4 (0.7-2.6)	4.4% (1.8-8.3%)	5.6% (2.3-10%)	130 (51-240)	1900 (810-3600)
Armenia	2016	Female	80 to 84	21% (13-33%)	79% (67-87%)	1.3 (0.68-2.2)	4.4% (1.9-8.2%)	5.4% (2.3-10%)	170 (71-320)	1800 (790-3400)
Armenia	2016	Female	85 to 89	21% (13-32%)	79% (68-87%)	1.3 (0.67-2.3)	4.1% (1.6-8.1%)	5.1% (1.8-10%)	95 (35-190)	740 (290-1500)
Armenia	2016	Female	90 to 94	23% (14-33%)	77% (67-86%)	1.4 (0.68-2.6)	3.8% (1.1-8.2%)	4.6% (1.2-10%)	57 (15-130)	310 (89-670)
Armenia	2016	Female	95 plus	25% (14-40%)	75% (60-86%)	1.5 (0.63-3)	3.8% (0.64-8.9%)	4.5% (0.56-11%)	16 (1.9-39)	57 (9.6-140)
Armenia	2016	Male	All Ages	84% (81-86%)	16% (14-19%)	1.9 (1.4-2.6)	3.5% (1.9-5.5%)	1.9% (-0.34-4.5%)	260 (-46-630)	17000 (8500-26000)
Armenia	2016	Male	15 to 19	68% (58-76%)	32% (24-42%)	0.76 (0.4-1.3)	3.5% (2.4-5.1%)	4.5% (2.4-7.4%)	2.7 (1.4-4.5)	340 (230-500)
Armenia	2016	Male	20 to 24	90% (84-94%)	10% (6.4-16%)	1.7 (1-2.8)	7.8% (5.6-11%)	9.3% (5.3-14%)	10 (5.8-16)	1200 (810-1700)
Armenia	2016	Male	25 to 29	93% (88-95%)	7.5% (4.8-12%)	2.6 (1.6-4.1)	9.2% (6.7-12%)	11% (6.6-17%)	17 (9.1-26)	1700 (1200-2400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Armenia	2016	Male	30 to 34	94% (90-96%)	5.9% (3.6-9.7%)	3.1 (2-4.5)	9.1% (6.3-12%)	12% (6.8-17%)	21 (12-33)	1900 (1300-2600)
Armenia	2016	Male	35 to 39	95% (92-97%)	5% (3.1-7.6%)	3 (1.7-4.6)	8% (5.1-11%)	10% (5.3-16%)	21 (10-34)	1600 (980-2300)
Armenia	2016	Male	40 to 44	95% (93-97%)	4.8% (2.9-7.3%)	3 (1.7-4.5)	7% (4.1-11%)	8.6% (4.1-14%)	23 (10-39)	1400 (800-2200)
Armenia	2016	Male	45 to 49	95% (91-97%)	5.4% (3.2-8.8%)	3.2 (2-5)	6.9% (3.7-11%)	8.2% (3.6-15%)	32 (14-58)	1700 (840-2800)
Armenia	2016	Male	50 to 54	93% (90-96%)	6.6% (3.9-10%)	3.2 (2-4.8)	5.7% (2.2-10%)	6.5% (1.9-12%)	54 (16-100)	2400 (950-4400)
Armenia	2016	Male	55 to 59	92% (85-96%)	8.3% (4.2-15%)	2.7 (1.8-4)	3.8% (0.99-7.8%)	4.2% (0.45-9.1%)	54 (5.8-120)	2100 (560-4100)
Armenia	2016	Male	60 to 64	90% (82-95%)	9.9% (4.6-18%)	2.3 (1.4-3.6)	2.6% (-0.067-6.5%)	2.7% (-0.52-7.2%)	42 (-8.4-110)	1400 (-37-3300)
Armenia	2016	Male	65 to 69	89% (79-95%)	11% (5.3-21%)	2.1 (1.3-3.1)	1.7% (-0.49-4.5%)	1.7% (-1-5.1%)	23 (-14-70)	670 (-200-1800)
Armenia	2016	Male	70 to 74	87% (76-93%)	13% (7-24%)	1.8 (1-2.7)	0.89% (-1.3-3.7%)	0.78% (-1.8-4.1%)	7.8 (-18-40)	200 (-290-800)
Armenia	2016	Male	75 to 79	85% (73-92%)	15% (7.6-27%)	1.5 (0.82-2.4)	0.25% (-1.8-2.7%)	0.033% (-2.4-2.9%)	0.66 (-51-62)	92 (-630-990)
Armenia	2016	Male	80 to 84	83% (71-91%)	17% (9.1-29%)	1.3 (0.76-2.1)	-0.47% (-2.7-1.7%)	-0.82% (-3.6-1.7%)	-18 (-80-37)	-130 (-760-470)
Armenia	2016	Male	85 to 89	81% (69-90%)	19% (10-31%)	1.2 (0.73-1.9)	-0.81% (-2.8-1.1%)	-1.3% (-3.9-1.1%)	-15 (-45-12)	-85 (-310-110)
Armenia	2016	Male	90 to 94	81% (69-89%)	19% (11-31%)	1.2 (0.66-1.9)	-1.3% (-3.4-0.74%)	-2% (-4.8-0.49%)	-9.9 (-24-2.6)	-40 (-110-24)
Armenia	2016	Male	95 plus	80% (65-91%)	20% (9.1-35%)	1 (0.5-1.9)	-1.3% (-3.5-0.65%)	-2.2% (-5-0.23%)	-1.7 (-3.8-0.18)	-4.3 (-11-2.1)
Azerbaijan	1990	Female	All Ages	55% (48-62%)	45% (38-52%)	0.98 (0.73-1.3)	0.94% (0.021-2%)	0.47% (-2-3.2%)	120 (-510-810)	13000 (260-27000)
Azerbaijan	1990	Female	15 to 19	55% (43-70%)	45% (30-57%)	0.94 (0.54-1.5)	1.7% (1-2.7%)	3% (1.4-5.2%)	5.1 (2.4-8.8)	630 (370-980)
Azerbaijan	1990	Female	20 to 24	62% (49-75%)	38% (25-51%)	1.3 (0.75-2.1)	3% (1.8-4.4%)	4.3% (1.9-7.4%)	11 (4.9-19)	1300 (810-2000)
Azerbaijan	1990	Female	25 to 29	65% (51-77%)	35% (23-49%)	1.5 (0.91-2.2)	3.3% (2.1-4.7%)	4.8% (2.1-7.8%)	16 (7-27)	1700 (1000-2500)
Azerbaijan	1990	Female	30 to 34	66% (52-79%)	34% (21-48%)	1.6 (0.99-2.3)	3.4% (2.1-4.9%)	5.5% (2.6-8.4%)	19 (9.1-31)	1700 (1000-2400)
Azerbaijan	1990	Female	35 to 39	66% (53-80%)	34% (20-47%)	1.6 (0.97-2.5)	3.4% (2-5.1%)	5.7% (2.8-9.1%)	20 (9.7-32)	1400 (810-2100)
Azerbaijan	1990	Female	40 to 44	66% (52-79%)	34% (21-48%)	1.6 (0.99-2.6)	3.3% (1.7-5.3%)	5.8% (2.5-9.6%)	17 (7.3-29)	970 (480-1600)
Azerbaijan	1990	Female	45 to 49	66% (52-79%)	34% (21-48%)	1.7 (0.98-2.6)	3% (0.95-5.6%)	5.1% (1.4-9.7%)	24 (6.2-46)	1100 (370-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	1990	Female	50 to 54	65% (50-77%)	35% (23-50%)	1.6 (0.96-2.6)	2.6% (0.16-5.9%)	3.9% (0.067-8.8%)	37 (0.58-82)	1500 (94-3400)
Azerbaijan	1990	Female	55 to 59	63% (49-76%)	37% (24-51%)	1.6 (0.95-2.6)	2.2% (-0.73-5.7%)	3% (-1.4-8%)	36 (-17-99)	1300 (-460-3500)
Azerbaijan	1990	Female	60 to 64	61% (47-76%)	39% (24-53%)	1.5 (0.89-2.6)	1.7% (-1.4-5.8%)	2.1% (-2.3-7.6%)	35 (-39-130)	1100 (-970-3800)
Azerbaijan	1990	Female	65 to 69	59% (47-74%)	41% (26-53%)	1.5 (0.93-2.3)	1.2% (-2-4.9%)	1.6% (-2.7-6.5%)	25 (-42-100)	600 (-930-2400)
Azerbaijan	1990	Female	70 to 74	57% (45-69%)	43% (31-55%)	1.4 (0.85-2.2)	0.16% (-3.5-4.2%)	0.15% (-4.4-5.3%)	2.9 (-83-100)	72 (-1500-1900)
Azerbaijan	1990	Female	75 to 79	54% (44-67%)	46% (33-56%)	1.4 (0.77-2.1)	-0.63% (-4.1-3.6%)	-0.87% (-5.2-4.2%)	-25 (-150-120)	-310 (-2000-1800)
Azerbaijan	1990	Female	80 to 84	52% (40-66%)	48% (34-60%)	1.3 (0.78-2)	-0.83% (-4.3-3%)	-1.1% (-5.4-3.4%)	-24 (-120-74)	-220 (-1200-810)
Azerbaijan	1990	Female	85 to 89	52% (41-63%)	48% (37-59%)	1.3 (0.77-1.9)	-1.6% (-5-2%)	-2% (-6-2.3%)	-42 (-130-48)	-290 (-910-360)
Azerbaijan	1990	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.3 (0.76-2.1)	-2.2% (-5.6-1.8%)	-2.7% (-6.8-2%)	-28 (-70-20)	-130 (-340-110)
Azerbaijan	1990	Female	95 plus	53% (40-68%)	47% (32-60%)	1.2 (0.62-2.2)	-2.5% (-6-1.9%)	-3.1% (-7.4-2.1%)	-9.6 (-23-6.5)	-30 (-74-23)
Azerbaijan	1990	Male	All Ages	48% (39-58%)	52% (42-61%)	1.9 (1.5-2.5)	5.3% (3.8-7%)	6.2% (3.7-9.1%)	1800 (1100-2700)	88000 (62000-120000)
Azerbaijan	1990	Male	15 to 19	36% (22-53%)	64% (47-78%)	0.47 (0.24-0.8)	4.2% (2.7-5.8%)	4.8% (2.3-7.3%)	20 (9.1-30)	2300 (1400-3200)
Azerbaijan	1990	Male	20 to 24	57% (38-74%)	43% (26-62%)	1.6 (0.98-2.4)	11% (8-14%)	11% (7.3-16%)	65 (41-93)	6400 (4600-8500)
Azerbaijan	1990	Male	25 to 29	68% (50-82%)	32% (18-50%)	3.5 (2.4-4.9)	16% (12-20%)	19% (14-25%)	140 (95-190)	11000 (8300-14000)
Azerbaijan	1990	Male	30 to 34	75% (56-87%)	25% (13-44%)	4.5 (3.1-6.3)	17% (13-22%)	21% (16-28%)	170 (120-230)	12000 (8800-15000)
Azerbaijan	1990	Male	35 to 39	77% (61-89%)	23% (11-39%)	4.1 (2.8-5.8)	15% (11-20%)	19% (13-26%)	150 (100-210)	9400 (6600-12000)
Azerbaijan	1990	Male	40 to 44	77% (62-89%)	23% (11-38%)	4.1 (2.7-5.9)	14% (9.9-20%)	17% (11-25%)	120 (74-170)	6300 (4300-8700)
Azerbaijan	1990	Male	45 to 49	76% (58-88%)	24% (12-42%)	4.6 (2.9-6.7)	15% (9.9-20%)	18% (11-25%)	180 (110-270)	8600 (5600-12000)
Azerbaijan	1990	Male	50 to 54	75% (58-87%)	25% (13-42%)	4.4 (2.8-6.3)	13% (8.1-18%)	15% (8.8-22%)	290 (160-420)	12000 (7100-16000)
Azerbaijan	1990	Male	55 to 59	72% (54-86%)	28% (14-46%)	3.4 (2.2-5.1)	9.4% (4.7-15%)	11% (4.7-17%)	260 (110-420)	8900 (4300-14000)
Azerbaijan	1990	Male	60 to 64	69% (49-85%)	31% (15-51%)	2.7 (1.6-4.4)	6.5% (2.3-12%)	7.1% (2.1-14%)	200 (59-400)	6000 (2100-11000)
Azerbaijan	1990	Male	65 to 69	66% (47-82%)	34% (18-53%)	2.4 (1.4-3.7)	4.9% (1.2-9.6%)	5.2% (0.95-11%)	96 (18-200)	2400 (580-4700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	1990	Male	70 to 74	62% (43-78%)	38% (22-57%)	1.9 (1.1-3.2)	3.4% (0.082-8.2%)	3.5% (-0.26-9.2%)	59 (-4.3-150)	1200 (27-2900)
Azerbaijan	1990	Male	75 to 79	57% (38-76%)	43% (24-62%)	1.6 (0.82-2.7)	2.6% (-0.33-6.9%)	2.7% (-0.66-7.6%)	58 (-14-160)	900 (-110-2300)
Azerbaijan	1990	Male	80 to 84	54% (35-71%)	46% (29-65%)	1.4 (0.76-2.3)	2% (-0.84-5.4%)	1.9% (-1.4-5.8%)	26 (-19-78)	310 (-140-860)
Azerbaijan	1990	Male	85 to 89	52% (34-69%)	48% (31-66%)	1.3 (0.68-2.1)	1.3% (-1.6-4.2%)	1.2% (-2.2-4.6%)	10 (-20-41)	100 (-120-320)
Azerbaijan	1990	Male	90 to 94	51% (34-68%)	49% (32-66%)	1.2 (0.61-1.9)	0.68% (-2.1-3.5%)	0.38% (-2.9-3.7%)	1.4 (-11-14)	15 (-47-79)
Azerbaijan	1990	Male	95 plus	52% (30-72%)	48% (28-70%)	1.1 (0.46-2)	0.68% (-2.1-3.5%)	0.28% (-3.1-3.4%)	0.31 (-3.5-3.8)	2.9 (-9.2-15)
Azerbaijan	1995	Female	All Ages	55% (49-62%)	45% (38-51%)	1 (0.78-1.3)	1.2% (0.075-2.3%)	0.81% (-1.8-3.6%)	210 (-470-920)	16000 (1000-32000)
Azerbaijan	1995	Female	15 to 19	55% (43-70%)	45% (30-57%)	0.96 (0.57-1.5)	1.8% (1-2.9%)	3.2% (1.3-5.4%)	6.7 (2.7-12)	740 (400-1200)
Azerbaijan	1995	Female	20 to 24	62% (49-74%)	38% (26-51%)	1.3 (0.78-2.1)	3.2% (2-5%)	4.7% (2-8.1%)	14 (6-24)	1500 (860-2300)
Azerbaijan	1995	Female	25 to 29	64% (51-77%)	36% (23-49%)	1.5 (0.91-2.2)	3.6% (2.2-5.2%)	5.2% (2.2-8.5%)	20 (8.8-34)	1900 (1100-2900)
Azerbaijan	1995	Female	30 to 34	66% (53-79%)	34% (21-47%)	1.6 (1-2.3)	3.7% (2.2-5.3%)	5.8% (2.9-9%)	30 (15-46)	2400 (1400-3400)
Azerbaijan	1995	Female	35 to 39	66% (53-80%)	34% (20-47%)	1.6 (1-2.4)	3.6% (2.1-5.5%)	6% (3-9.4%)	34 (17-54)	2200 (1300-3300)
Azerbaijan	1995	Female	40 to 44	66% (52-80%)	34% (20-48%)	1.6 (1-2.4)	3.6% (1.9-5.7%)	6.1% (3-9.7%)	33 (15-54)	1800 (930-2800)
Azerbaijan	1995	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.7 (1-2.6)	3.2% (0.97-6%)	5.1% (1.3-9.7%)	26 (6.8-49)	1200 (360-2200)
Azerbaijan	1995	Female	50 to 54	65% (51-78%)	35% (22-49%)	1.6 (0.98-2.6)	2.7% (0.16-5.7%)	4% (0.064-9%)	28 (0.44-62)	1100 (60-2500)
Azerbaijan	1995	Female	55 to 59	63% (49-76%)	37% (24-51%)	1.6 (0.98-2.6)	2.2% (-0.8-5.8%)	2.9% (-1.4-8.1%)	46 (-23-130)	1600 (-610-4400)
Azerbaijan	1995	Female	60 to 64	61% (48-76%)	39% (24-52%)	1.6 (0.92-2.5)	1.7% (-1.7-5.9%)	2% (-2.5-7.6%)	44 (-53-160)	1300 (-1400-4700)
Azerbaijan	1995	Female	65 to 69	59% (47-74%)	41% (26-53%)	1.5 (0.96-2.3)	1.1% (-2.4-4.9%)	1.4% (-3.1-6.3%)	39 (-87-170)	920 (-1900-4000)
Azerbaijan	1995	Female	70 to 74	57% (45-70%)	43% (30-55%)	1.4 (0.85-2.2)	0.028% (-3.8-4.3%)	-0.0021% (-4.8-5.2%)	-0.096 (-100-120)	14 (-1900-2200)
Azerbaijan	1995	Female	75 to 79	54% (44-67%)	46% (33-56%)	1.4 (0.78-2.2)	-0.69% (-4.2-3.6%)	-0.92% (-5.2-4.1%)	-22 (-120-96)	-270 (-1700-1400)
Azerbaijan	1995	Female	80 to 84	53% (41-66%)	47% (34-59%)	1.3 (0.81-2)	-0.83% (-4.3-2.8%)	-1.1% (-5.2-3.3%)	-22 (-110-66)	-210 (-1100-700)
Azerbaijan	1995	Female	85 to 89	52% (41-64%)	48% (36-59%)	1.3 (0.77-1.9)	-1.6% (-5.1-1.9%)	-2% (-6.1-2.3%)	-31 (-97-35)	-220 (-690-260)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	1995	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.3 (0.75-2.1)	-2.2% (-5.8-1.8%)	-2.8% (-7-2.1%)	-27 (-68-20)	-130 (-330-100)
Azerbaijan	1995	Female	95 plus	53% (41-68%)	47% (32-59%)	1.3 (0.62-2.2)	-2.5% (-6.2-1.6%)	-3.1% (-7.4-1.8%)	-10 (-24-5.7)	-32 (-78-20)
Azerbaijan	1995	Male	All Ages	50% (41-59%)	50% (41-59%)	2 (1.6-2.5)	6.6% (4.9-8.6%)	7.5% (4.9-11%)	2500 (1600-3500)	120000 (86000-150000)
Azerbaijan	1995	Male	15 to 19	36% (22-53%)	64% (47-78%)	0.48 (0.26-0.84)	4.3% (2.7-6.2%)	4.8% (2.4-7.4%)	25 (12-39)	2600 (1600-3700)
Azerbaijan	1995	Male	20 to 24	58% (38-74%)	42% (26-62%)	1.6 (1-2.4)	11% (8.3-15%)	12% (8-17%)	99 (63-140)	8800 (6300-12000)
Azerbaijan	1995	Male	25 to 29	68% (50-82%)	32% (18-50%)	3.5 (2.4-4.9)	18% (14-22%)	21% (15-26%)	190 (130-250)	14000 (10000-18000)
Azerbaijan	1995	Male	30 to 34	75% (56-87%)	25% (13-44%)	4.5 (3.1-6.2)	19% (15-24%)	23% (17-29%)	280 (200-360)	19000 (14000-23000)
Azerbaijan	1995	Male	35 to 39	77% (62-88%)	23% (12-38%)	4.1 (3-5.7)	17% (12-21%)	20% (14-26%)	270 (190-360)	16000 (12000-21000)
Azerbaijan	1995	Male	40 to 44	78% (62-89%)	22% (11-38%)	4.1 (2.8-5.7)	15% (10-20%)	18% (12-24%)	240 (150-340)	13000 (8600-17000)
Azerbaijan	1995	Male	45 to 49	77% (58-88%)	23% (12-42%)	4.6 (3-6.7)	16% (11-21%)	18% (12-25%)	200 (130-290)	9300 (6300-13000)
Azerbaijan	1995	Male	50 to 54	75% (58-87%)	25% (13-42%)	4.4 (2.9-6.3)	14% (8.8-19%)	16% (9.6-22%)	250 (150-360)	10000 (6300-14000)
Azerbaijan	1995	Male	55 to 59	73% (55-86%)	27% (14-45%)	3.4 (2.2-5)	9.7% (5-15%)	11% (5.2-17%)	300 (140-490)	10000 (5400-16000)
Azerbaijan	1995	Male	60 to 64	69% (50-84%)	31% (16-50%)	2.8 (1.6-4.3)	6.6% (2.2-12%)	7.1% (2.1-14%)	250 (75-480)	7400 (2600-14000)
Azerbaijan	1995	Male	65 to 69	66% (48-82%)	34% (18-52%)	2.4 (1.5-3.7)	4.9% (1.3-9.7%)	5.3% (1.1-11%)	190 (40-410)	4700 (1300-9500)
Azerbaijan	1995	Male	70 to 74	62% (44-78%)	38% (22-56%)	2 (1.1-3.3)	3.3% (0.024-8.2%)	3.5% (-0.39-9.1%)	74 (-7.8-190)	1500 (10-3600)
Azerbaijan	1995	Male	75 to 79	58% (40-75%)	42% (25-60%)	1.6 (0.86-2.7)	2.6% (-0.37-6.8%)	2.7% (-0.79-7.5%)	40 (-11-120)	630 (-89-1700)
Azerbaijan	1995	Male	80 to 84	54% (35-71%)	46% (29-65%)	1.4 (0.79-2.3)	2% (-0.98-5.4%)	1.9% (-1.6-5.8%)	21 (-16-64)	250 (-120-700)
Azerbaijan	1995	Male	85 to 89	52% (34-68%)	48% (32-66%)	1.3 (0.69-2.1)	1.3% (-1.6-4.3%)	1.1% (-2.3-4.6%)	7.7 (-16-31)	74 (-91-250)
Azerbaijan	1995	Male	90 to 94	51% (35-68%)	49% (32-65%)	1.2 (0.64-2)	0.65% (-2.2-3.5%)	0.34% (-2.9-3.6%)	1.1 (-9.2-11)	12 (-39-64)
Azerbaijan	1995	Male	95 plus	52% (31-72%)	48% (28-69%)	1.1 (0.48-2)	0.62% (-2.3-3.6%)	0.2% (-3.4-3.6%)	0.17 (-2.9-3.1)	2.1 (-7.6-12)
Azerbaijan	2000	Female	All Ages	56% (50-63%)	44% (37-50%)	1.1 (0.85-1.4)	1.5% (0.14-2.9%)	1.1% (-2.4-3%)	300 (-510-1100)	18000 (1700-36000)
Azerbaijan	2000	Female	15 to 19	55% (43-69%)	45% (31-57%)	1 (0.59-1.6)	1.7% (0.94-2.7%)	3.3% (1.2-5.6%)	6.2 (2.3-10)	740 (390-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	2000	Female	20 to 24	62% (49-74%)	38% (26-51%)	1.4 (0.8-2.2)	3.3% (1.9-5%)	5.2% (1.9-9%)	13 (4.7-22)	1400 (790-2200)
Azerbaijan	2000	Female	25 to 29	65% (51-77%)	35% (23-49%)	1.6 (0.94-2.4)	3.7% (2.2-5.4%)	5.9% (2.5-9.6%)	18 (7.6-29)	1700 (1000-2600)
Azerbaijan	2000	Female	30 to 34	66% (52-78%)	34% (22-48%)	1.7 (1.1-2.5)	3.8% (2.4-5.5%)	6.6% (3.3-10%)	27 (13-42)	2200 (1300-3100)
Azerbaijan	2000	Female	35 to 39	66% (52-79%)	34% (21-48%)	1.7 (1.1-2.5)	3.7% (2.2-5.5%)	6.7% (3.6-10%)	38 (20-60)	2600 (1500-3800)
Azerbaijan	2000	Female	40 to 44	66% (52-79%)	34% (21-48%)	1.7 (1.1-2.5)	3.7% (2.1-5.7%)	6.7% (3.6-10%)	43 (23-68)	2400 (1300-3700)
Azerbaijan	2000	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.7 (1.1-2.7)	3.4% (1.4-5.9%)	5.9% (2.3-10%)	41 (15-70)	1900 (730-3200)
Azerbaijan	2000	Female	50 to 54	65% (51-78%)	35% (22-49%)	1.7 (1-2.7)	3.1% (0.39-6.3%)	4.6% (0.48-9.4%)	30 (3.1-63)	1200 (160-2400)
Azerbaijan	2000	Female	55 to 59	63% (49-77%)	37% (23-51%)	1.7 (1-2.7)	2.6% (-0.52-6.1%)	3.5% (-0.98-8.5%)	31 (-8.5-75)	1100 (-220-2600)
Azerbaijan	2000	Female	60 to 64	61% (48-76%)	39% (24-52%)	1.6 (0.93-2.6)	2.1% (-1.5-6.5%)	2.5% (-2.3-8.1%)	62 (-57-200)	1900 (-1400-5700)
Azerbaijan	2000	Female	65 to 69	59% (47-73%)	41% (27-53%)	1.6 (1-2.4)	1.6% (-2.1-5.6%)	2% (-2.7-6.9%)	62 (-85-230)	1500 (-1900-5300)
Azerbaijan	2000	Female	70 to 74	57% (45-70%)	43% (30-55%)	1.5 (0.89-2.3)	0.49% (-3.4-4.9%)	0.58% (-4.2-6%)	21 (-160-220)	400 (-2800-4000)
Azerbaijan	2000	Female	75 to 79	54% (44-67%)	46% (33-56%)	1.4 (0.81-2.2)	-0.3% (-4.1-4.1%)	-0.44% (-5-4.8%)	-12 (-130-130)	-140 (-1800-1900)
Azerbaijan	2000	Female	80 to 84	53% (41-66%)	47% (34-59%)	1.4 (0.82-2.1)	-0.53% (-4.1-3.4%)	-0.72% (-4.9-3.9%)	-14 (-96-76)	-130 (-960-840)
Azerbaijan	2000	Female	85 to 89	52% (40-64%)	48% (36-60%)	1.3 (0.81-2)	-1.3% (-4.9-2.6%)	-1.6% (-5.7-3%)	-32 (-120-59)	-220 (-830-450)
Azerbaijan	2000	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.3 (0.77-2.1)	-2% (-5.6-2%)	-2.4% (-6.8-2.4%)	-24 (-68-23)	-110 (-330-120)
Azerbaijan	2000	Female	95 plus	53% (40-67%)	47% (33-60%)	1.3 (0.64-2.3)	-2.3% (-6.1-2.2%)	-2.9% (-7.4-2.5%)	-12 (-31-10)	-38 (-99-35)
Azerbaijan	2000	Male	All Ages	51% (42-61%)	49% (39-58%)	2.2 (1.7-2.7)	7% (5.1-9%)	7.7% (4.6-11%)	2400 (1400-3600)	110000 (79000-140000)
Azerbaijan	2000	Male	15 to 19	36% (22-54%)	64% (46-78%)	0.49 (0.27-0.85)	4.2% (2.9-5.8%)	4.8% (2.7-7.2%)	20 (11-31)	2300 (1600-3300)
Azerbaijan	2000	Male	20 to 24	57% (38-74%)	43% (26-62%)	1.6 (1-2.5)	12% (8.7-15%)	13% (9.1-18%)	67 (44-94)	6500 (4700-8600)
Azerbaijan	2000	Male	25 to 29	68% (51-82%)	32% (18-49%)	3.7 (2.5-5.1)	18% (14-23%)	23% (17-30%)	130 (98-170)	11000 (8200-13000)
Azerbaijan	2000	Male	30 to 34	75% (56-88%)	25% (12-44%)	4.7 (3.2-6.5)	20% (16-24%)	26% (20-32%)	190 (140-250)	13000 (10000-17000)
Azerbaijan	2000	Male	35 to 39	77% (62-89%)	23% (11-38%)	4.3 (3-5.7)	17% (13-21%)	22% (16-28%)	240 (170-320)	15000 (11000-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	2000	Male	40 to 44	78% (61-89%)	22% (11-39%)	4.3 (3-5.8)	15% (11-20%)	19% (13-25%)	260 (170-360)	14000 (10000-19000)
Azerbaijan	2000	Male	45 to 49	77% (58-89%)	23% (11-42%)	4.8 (3.2-6.8)	16% (11-21%)	19% (13-26%)	270 (180-380)	13000 (8900-17000)
Azerbaijan	2000	Male	50 to 54	75% (57-87%)	25% (13-43%)	4.6 (3-6.5)	14% (9.3-20%)	16% (10-23%)	220 (130-320)	8900 (5600-12000)
Azerbaijan	2000	Male	55 to 59	73% (55-86%)	27% (14-45%)	3.6 (2.3-5.2)	10% (5.2-16%)	12% (5.4-18%)	190 (88-300)	6500 (3400-9800)
Azerbaijan	2000	Male	60 to 64	69% (50-85%)	31% (15-50%)	2.9 (1.7-4.5)	7.2% (2.4-13%)	7.8% (2.2-15%)	290 (85-560)	8400 (2900-15000)
Azerbaijan	2000	Male	65 to 69	67% (49-83%)	33% (17-51%)	2.5 (1.5-3.8)	5.5% (1.6-11%)	6% (1.4-12%)	260 (61-520)	6200 (1700-12000)
Azerbaijan	2000	Male	70 to 74	62% (44-78%)	38% (22-56%)	2.1 (1.2-3.4)	3.8% (0.24-9.3%)	4.1% (-0.046-10%)	170 (-1.8-400)	3200 (200-7500)
Azerbaijan	2000	Male	75 to 79	58% (39-76%)	42% (24-61%)	1.7 (0.9-2.9)	3% (-0.37-7.8%)	3.2% (-0.65-8.6%)	60 (-12-160)	910 (-110-2400)
Azerbaijan	2000	Male	80 to 84	55% (35-72%)	45% (28-65%)	1.5 (0.82-2.4)	2.2% (-0.95-5.9%)	2.2% (-1.5-6.4%)	21 (-14-62)	250 (-110-670)
Azerbaijan	2000	Male	85 to 89	52% (35-69%)	48% (31-65%)	1.3 (0.74-2.2)	1.5% (-1.5-4.5%)	1.4% (-2.1-4.9%)	10 (-16-37)	95 (-93-290)
Azerbaijan	2000	Male	90 to 94	51% (35-69%)	49% (31-65%)	1.2 (0.66-2.1)	0.78% (-2.2-3.7%)	0.5% (-3-4%)	1.6 (-9.7-13)	15 (-41-70)
Azerbaijan	2000	Male	95 plus	52% (31-72%)	48% (28-69%)	1.1 (0.5-2.1)	0.72% (-2.3-3.9%)	0.33% (-3.3-4%)	0.32 (-3.1-3.9)	2.7 (-8.6-15)
Azerbaijan	2005	Female	All Ages	57% (51-65%)	43% (35-49%)	1.2 (0.93-1.6)	1.8% (0.39-3.4%)	1.8% (-1.5-5.2%)	520 (-430-1500)	24000 (5200-44000)
Azerbaijan	2005	Female	15 to 19	55% (43-69%)	45% (31-57%)	1.1 (0.6-1.7)	1.8% (0.99-2.8%)	3.3% (1.3-5.7%)	7.6 (3.1-13)	900 (480-1400)
Azerbaijan	2005	Female	20 to 24	62% (49-74%)	38% (26-51%)	1.5 (0.85-2.4)	3.3% (1.9-4.9%)	5.1% (2.1-8.7%)	14 (5.6-26)	1700 (940-2600)
Azerbaijan	2005	Female	25 to 29	65% (51-77%)	35% (23-49%)	1.7 (1-2.6)	3.6% (2.2-5.3%)	5.8% (2.7-9.4%)	17 (7.9-29)	1700 (1000-2600)
Azerbaijan	2005	Female	30 to 34	66% (52-79%)	34% (21-48%)	1.8 (1.1-2.8)	3.8% (2.4-5.5%)	6.4% (3.3-10%)	23 (12-38)	2000 (1200-2800)
Azerbaijan	2005	Female	35 to 39	66% (53-81%)	34% (19-47%)	1.8 (1.1-2.7)	3.9% (2.2-5.8%)	6.8% (3.5-11%)	37 (19-58)	2500 (1400-3700)
Azerbaijan	2005	Female	40 to 44	66% (52-79%)	34% (21-48%)	1.8 (1.1-2.7)	3.8% (2.1-5.9%)	6.9% (3.6-11%)	57 (28-90)	3100 (1700-4800)
Azerbaijan	2005	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.9 (1.1-2.8)	3.7% (1.6-6.3%)	6.4% (2.7-11%)	66 (27-110)	3000 (1300-5100)
Azerbaijan	2005	Female	50 to 54	64% (50-78%)	36% (22-50%)	1.8 (1.1-2.8)	3.5% (0.69-6.8%)	5.3% (1.1-10%)	61 (12-120)	2400 (470-4600)
Azerbaijan	2005	Female	55 to 59	63% (50-76%)	37% (24-50%)	1.8 (1.1-2.8)	3.1% (-0.21-6.8%)	4.3% (-0.38-9.3%)	44 (-3.8-97)	1500 (-91-3300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	2005	Female	60 to 64	61% (48-75%)	39% (25-52%)	1.7 (0.99-2.8)	2.8% (-0.93-7%)	3.5% (-1.4-9.1%)	52 (-20-140)	1500 (-490-4000)
Azerbaijan	2005	Female	65 to 69	59% (48-73%)	41% (27-52%)	1.7 (1.1-2.5)	2.3% (-1.6-6.4%)	2.9% (-1.9-8%)	100 (-65-280)	2300 (-1500-6400)
Azerbaijan	2005	Female	70 to 74	57% (46-70%)	43% (30-54%)	1.6 (0.96-2.5)	1.2% (-3.1-5.9%)	1.5% (-3.8-7.1%)	66 (-160-330)	1200 (-2900-6000)
Azerbaijan	2005	Female	75 to 79	54% (44-67%)	46% (33-56%)	1.5 (0.88-2.4)	0.38% (-3.6-5.1%)	0.39% (-4.3-5.9%)	15 (-160-230)	240 (-2300-3300)
Azerbaijan	2005	Female	80 to 84	53% (40-66%)	47% (34-60%)	1.5 (0.9-2.3)	0.084% (-3.6-4.2%)	0.017% (-4.4-4.8%)	0.43 (-130-140)	30 (-1300-1500)
Azerbaijan	2005	Female	85 to 89	52% (41-63%)	48% (37-59%)	1.4 (0.86-2.2)	-0.61% (-4.3-3.6%)	-0.81% (-5.1-4.3%)	-13 (-85-70)	-84 (-600-520)
Azerbaijan	2005	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.4 (0.82-2.3)	-1.3% (-5.1-3.2%)	-1.7% (-6.2-3.7%)	-18 (-66-39)	-82 (-320-200)
Azerbaijan	2005	Female	95 plus	53% (40-68%)	47% (32-60%)	1.4 (0.71-2.4)	-1.7% (-5.6-3.1%)	-2.2% (-6.7-3.3%)	-7.9 (-25-13)	-24 (-79-43)
Azerbaijan	2005	Male	All Ages	53% (45-62%)	47% (38-55%)	2.5 (2-3.2)	8.2% (5.9-11%)	8.8% (5.4-13%)	3100 (1900-4600)	130000 (96000-180000)
Azerbaijan	2005	Male	15 to 19	29% (18-45%)	71% (55-82%)	0.42 (0.23-0.74)	4% (3-5.2%)	4.5% (2.9-6.4%)	22 (14-33)	2600 (1900-3600)
Azerbaijan	2005	Male	20 to 24	54% (37-69%)	46% (31-63%)	1.6 (1.1-2.4)	11% (8.8-14%)	13% (8.9-17%)	82 (54-120)	8100 (5900-11000)
Azerbaijan	2005	Male	25 to 29	71% (57-83%)	29% (17-43%)	4.2 (2.9-5.7)	19% (15-23%)	24% (18-30%)	160 (110-210)	12000 (9300-16000)
Azerbaijan	2005	Male	30 to 34	78% (64-88%)	22% (12-36%)	5.4 (3.8-7.2)	21% (16-25%)	27% (20-33%)	210 (150-270)	14000 (11000-18000)
Azerbaijan	2005	Male	35 to 39	79% (68-89%)	21% (11-32%)	4.8 (3.5-6.4)	18% (13-23%)	23% (16-30%)	230 (160-330)	14000 (10000-19000)
Azerbaijan	2005	Male	40 to 44	79% (66-89%)	21% (11-34%)	4.8 (3.5-6.4)	16% (11-21%)	20% (13-27%)	340 (220-490)	18000 (13000-25000)
Azerbaijan	2005	Male	45 to 49	80% (66-89%)	20% (11-34%)	5.5 (3.9-7.4)	17% (12-22%)	20% (14-28%)	420 (270-580)	19000 (13000-26000)
Azerbaijan	2005	Male	50 to 54	79% (64-88%)	21% (12-36%)	5.3 (3.6-7.3)	15% (10-21%)	18% (11-25%)	410 (250-570)	16000 (10000-22000)
Azerbaijan	2005	Male	55 to 59	75% (59-86%)	25% (14-41%)	4 (2.7-5.8)	12% (6-18%)	13% (6.3-21%)	230 (110-370)	8000 (4100-12000)
Azerbaijan	2005	Male	60 to 64	70% (51-84%)	30% (16-49%)	3.2 (1.9-4.8)	8.3% (3.1-14%)	9.2% (3-16%)	200 (62-370)	5900 (2100-10000)
Azerbaijan	2005	Male	65 to 69	67% (47-82%)	33% (18-53%)	2.7 (1.6-4.1)	6.6% (2-12%)	7.2% (2-13%)	310 (83-590)	7400 (2200-14000)
Azerbaijan	2005	Male	70 to 74	63% (44-78%)	37% (22-56%)	2.3 (1.3-3.7)	4.8% (0.75-10%)	5.2% (0.61-12%)	270 (31-590)	5100 (790-11000)
Azerbaijan	2005	Male	75 to 79	58% (39-75%)	42% (25-61%)	1.9 (0.99-3)	3.8% (0.24-8.8%)	4.1% (-0.059-9.7%)	150 (-2.1-360)	2200 (140-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	2005	Male	80 to 84	55% (36-72%)	45% (28-64%)	1.6 (0.9-2.6)	2.9% (-0.38-7.1%)	3% (-0.87-7.9%)	56 (-17-150)	640 (-89-1600)
Azerbaijan	2005	Male	85 to 89	52% (34-69%)	48% (31-66%)	1.5 (0.8-2.4)	2.1% (-0.7-5.7%)	2.1% (-1.3-6.2%)	13 (-7.8-39)	110 (-39-300)
Azerbaijan	2005	Male	90 to 94	51% (35-69%)	49% (31-65%)	1.3 (0.72-2.3)	1.2% (-1.7-4.1%)	1.1% (-2.4-4.4%)	3.5 (-8-14)	24 (-32-78)
Azerbaijan	2005	Male	95 plus	52% (31-72%)	48% (28-69%)	1.2 (0.56-2.3)	1% (-2-4.4%)	0.72% (-2.9-4.6%)	0.67 (-2.7-4.3)	3.7 (-7.1-16)
Azerbaijan	2010	Female	All Ages	57% (51-64%)	43% (36-49%)	1.4 (1-1.9)	2.2% (0.47-4.1%)	2.4% (-1.4-6.5%)	760 (-440-2100)	30000 (6200-55000)
Azerbaijan	2010	Female	15 to 19	55% (43-70%)	45% (30-57%)	1.2 (0.65-1.9)	1.9% (1-3.1%)	3.5% (1.5-6.2%)	6.7 (2.9-12)	860 (470-1400)
Azerbaijan	2010	Female	20 to 24	62% (49-74%)	38% (26-51%)	1.7 (0.94-2.7)	3.3% (2-4.9%)	5.3% (2.3-9.1%)	15 (6.2-26)	1900 (1100-2900)
Azerbaijan	2010	Female	25 to 29	64% (51-77%)	36% (23-49%)	1.9 (1.1-2.9)	3.6% (2.3-5.2%)	6% (3-9.6%)	18 (8.7-29)	2000 (1200-2800)
Azerbaijan	2010	Female	30 to 34	66% (52-78%)	34% (22-48%)	2 (1.2-3.1)	3.8% (2.6-5.5%)	6.7% (3.8-10%)	22 (12-35)	2000 (1300-2800)
Azerbaijan	2010	Female	35 to 39	66% (53-79%)	34% (21-47%)	2 (1.2-3.1)	3.9% (2.4-5.8%)	7.3% (4-11%)	31 (17-50)	2200 (1300-3300)
Azerbaijan	2010	Female	40 to 44	66% (52-79%)	34% (21-48%)	2 (1.2-3.2)	4% (2.2-6.2%)	7.4% (3.9-12%)	50 (26-81)	2800 (1600-4500)
Azerbaijan	2010	Female	45 to 49	66% (51-79%)	34% (21-49%)	2.1 (1.2-3.1)	4% (1.7-6.7%)	7.1% (3.1-12%)	79 (34-130)	3700 (1700-6300)
Azerbaijan	2010	Female	50 to 54	64% (51-78%)	36% (22-49%)	2 (1.2-3.1)	4% (1.2-7.4%)	6.2% (1.9-11%)	93 (28-180)	3700 (1100-6900)
Azerbaijan	2010	Female	55 to 59	63% (50-76%)	37% (24-50%)	2 (1.2-3.2)	3.7% (0.35-7.5%)	5.2% (0.41-11%)	85 (6.5-180)	3000 (240-6200)
Azerbaijan	2010	Female	60 to 64	61% (47-76%)	39% (24-53%)	1.9 (1.1-3.2)	3.6% (-0.55-8.6%)	4.5% (-0.96-11%)	72 (-15-180)	2100 (-320-5100)
Azerbaijan	2010	Female	65 to 69	59% (48-74%)	41% (26-52%)	1.9 (1.1-2.9)	3.1% (-0.97-7.5%)	4% (-1.2-9.7%)	77 (-23-190)	1800 (-490-4400)
Azerbaijan	2010	Female	70 to 74	57% (45-70%)	43% (30-55%)	1.8 (1-2.8)	2% (-2.7-7.3%)	2.4% (-3.3-8.9%)	120 (-160-440)	2300 (-2900-8200)
Azerbaijan	2010	Female	75 to 79	54% (44-67%)	46% (33-56%)	1.7 (0.94-2.7)	1.2% (-3.3-6.5%)	1.3% (-4-7.5%)	68 (-210-390)	1000 (-2800-5600)
Azerbaijan	2010	Female	80 to 84	53% (40-66%)	47% (34-60%)	1.6 (0.95-2.6)	0.81% (-3.4-5.6%)	0.86% (-4-6.5%)	36 (-170-280)	420 (-1700-2900)
Azerbaijan	2010	Female	85 to 89	52% (40-64%)	48% (36-60%)	1.6 (0.95-2.5)	0.063% (-3.9-4.8%)	-0.057% (-4.8-5.5%)	-1.3 (-120-140)	14 (-830-1000)
Azerbaijan	2010	Female	90 to 94	53% (42-65%)	47% (35-58%)	1.6 (0.91-2.6)	-0.65% (-4.9-4.7%)	-0.89% (-5.8-5.3%)	-9.2 (-62-56)	-39 (-300-290)
Azerbaijan	2010	Female	95 plus	53% (40-68%)	47% (32-60%)	1.6 (0.78-2.7)	-1.1% (-5.4-5%)	-1.4% (-6.6-5.5%)	-6.1 (-27-23)	-17 (-87-81)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	2010	Male	All Ages	55% (46-63%)	45% (37-54%)	2.9 (2.2-3.8)	8.8% (6.1-12%)	9.4% (5.5-14%)	3600 (2100-5400)	150000 (100000-200000)
Azerbaijan	2010	Male	15 to 19	29% (18-45%)	71% (55-82%)	0.46 (0.24-0.82)	4.2% (3.1-5.5%)	4.7% (3.1-6.8%)	20 (13-30)	2500 (1800-3500)
Azerbaijan	2010	Male	20 to 24	53% (37-69%)	47% (31-63%)	1.8 (1.1-2.7)	11% (8.8-14%)	13% (8.8-17%)	73 (49-100)	8200 (6000-11000)
Azerbaijan	2010	Male	25 to 29	71% (57-82%)	29% (18-43%)	4.6 (3-6.4)	18% (14-22%)	23% (17-30%)	150 (100-200)	13000 (9700-16000)
Azerbaijan	2010	Male	30 to 34	78% (63-88%)	22% (12-37%)	5.9 (4.1-8.1)	20% (16-25%)	26% (20-35%)	190 (140-260)	14000 (11000-17000)
Azerbaijan	2010	Male	35 to 39	79% (67-88%)	21% (12-33%)	5.3 (3.7-7.5)	18% (13-24%)	23% (16-32%)	210 (140-290)	13000 (9200-18000)
Azerbaijan	2010	Male	40 to 44	79% (67-89%)	21% (11-33%)	5.3 (3.7-7.3)	17% (12-23%)	21% (14-30%)	280 (180-400)	15000 (10000-21000)
Azerbaijan	2010	Male	45 to 49	80% (64-89%)	20% (11-36%)	6.1 (4.2-8.5)	17% (12-23%)	21% (14-29%)	460 (310-660)	22000 (15000-30000)
Azerbaijan	2010	Male	50 to 54	78% (64-88%)	22% (12-36%)	5.9 (4-8.2)	16% (11-22%)	19% (13-27%)	560 (360-800)	22000 (15000-31000)
Azerbaijan	2010	Male	55 to 59	74% (60-86%)	26% (14-40%)	4.5 (2.9-6.6)	13% (6.9-19%)	15% (7.4-22%)	450 (220-700)	15000 (8000-23000)
Azerbaijan	2010	Male	60 to 64	70% (52-85%)	30% (15-48%)	3.5 (2.1-5.4)	9.4% (3.8-16%)	10% (3.7-18%)	260 (90-450)	7500 (2800-13000)
Azerbaijan	2010	Male	65 to 69	66% (48-82%)	34% (18-52%)	3 (1.8-4.6)	7.7% (2.8-13%)	8.5% (2.8-15%)	230 (72-410)	5400 (1900-9400)
Azerbaijan	2010	Male	70 to 74	62% (44-78%)	38% (22-56%)	2.5 (1.4-4.2)	5.8% (1.1-12%)	6.4% (1-14%)	340 (53-730)	6500 (1200-13000)
Azerbaijan	2010	Male	75 to 79	58% (39-76%)	42% (24-61%)	2.1 (1-3.4)	4.7% (0.6-10%)	5% (0.46-12%)	240 (21-560)	3600 (490-8200)
Azerbaijan	2010	Male	80 to 84	55% (36-72%)	45% (28-64%)	1.8 (0.97-2.9)	3.5% (-0.016-8.6%)	3.8% (-0.31-9.4%)	120 (-11-310)	1400 (-6.5-3300)
Azerbaijan	2010	Male	85 to 89	52% (34-69%)	48% (31-66%)	1.6 (0.86-2.7)	2.7% (-0.44-7%)	2.8% (-0.87-7.7%)	39 (-13-110)	330 (-50-840)
Azerbaijan	2010	Male	90 to 94	51% (35-69%)	49% (31-65%)	1.5 (0.77-2.6)	1.6% (-1.6-5.7%)	1.5% (-2.2-6%)	6.5 (-9.2-26)	39 (-40-140)
Azerbaijan	2010	Male	95 plus	52% (31-72%)	48% (28-69%)	1.4 (0.61-2.6)	1.2% (-2.2-5.8%)	0.98% (-2.9-6.1%)	0.95 (-2.8-5.9)	4.3 (-8-21)
Azerbaijan	2016	Female	All Ages	58% (51-65%)	42% (35-49%)	1.5 (0.99-2.1)	2.3% (0.42-4.3%)	2.4% (-1.4-6.9%)	710 (-400-2100)	29000 (5200-59000)
Azerbaijan	2016	Female	15 to 19	55% (43-69%)	45% (31-57%)	1.2 (0.62-2)	1.8% (1-2.9%)	3.5% (1.6-6.5%)	4.3 (1.9-7.8)	570 (310-930)
Azerbaijan	2016	Female	20 to 24	62% (49-74%)	38% (26-51%)	1.7 (0.89-2.8)	3.2% (2-4.8%)	5.2% (2.3-9.1%)	11 (4.8-20)	1500 (870-2300)
Azerbaijan	2016	Female	25 to 29	64% (51-78%)	36% (22-49%)	1.9 (1-3)	3.5% (2.2-5%)	5.7% (2.7-9.6%)	17 (7.6-31)	2100 (1300-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	2016	Female	30 to 34	66% (52-79%)	34% (21-48%)	2 (1.2-3.1)	3.7% (2.4-5.3%)	6.7% (3.7-11%)	25 (13-42)	2400 (1500-3600)
Azerbaijan	2016	Female	35 to 39	66% (53-80%)	34% (20-47%)	2 (1.1-3.4)	3.7% (2.2-5.7%)	7.3% (4-12%)	28 (14-47)	2100 (1200-3300)
Azerbaijan	2016	Female	40 to 44	66% (52-79%)	34% (21-48%)	2.1 (1.2-3.4)	3.8% (2-6.1%)	7.4% (3.5-12%)	41 (17-70)	2400 (1200-4000)
Azerbaijan	2016	Female	45 to 49	66% (50-79%)	34% (21-50%)	2.1 (1.2-3.4)	3.8% (1.6-6.7%)	7% (3-12%)	59 (24-110)	2900 (1200-5200)
Azerbaijan	2016	Female	50 to 54	64% (50-77%)	36% (23-50%)	2 (1.1-3.2)	3.7% (0.93-7.3%)	6.1% (1.5-12%)	100 (24-210)	4100 (1000-8400)
Azerbaijan	2016	Female	55 to 59	62% (50-75%)	38% (25-50%)	2 (1.1-3.3)	3.4% (0.18-7.6%)	5.1% (0.12-12%)	100 (2.4-230)	3600 (200-8200)
Azerbaijan	2016	Female	60 to 64	61% (47-75%)	39% (25-53%)	2 (1.1-3.3)	3.3% (-0.5-8.1%)	4.5% (-0.77-11%)	130 (-21-320)	3800 (-560-9600)
Azerbaijan	2016	Female	65 to 69	59% (47-74%)	41% (26-53%)	1.9 (1.1-3.1)	2.9% (-0.96-7.6%)	4% (-1.4-10%)	67 (-24-180)	1600 (-540-4300)
Azerbaijan	2016	Female	70 to 74	57% (46-70%)	43% (30-54%)	1.8 (0.95-3)	1.9% (-2.4-7.2%)	2.5% (-3.1-9.3%)	53 (-70-200)	1000 (-1300-3800)
Azerbaijan	2016	Female	75 to 79	54% (44-67%)	46% (33-56%)	1.7 (0.89-2.9)	1.2% (-2.9-6.9%)	1.4% (-3.9-8.5%)	64 (-180-410)	1000 (-2600-6000)
Azerbaijan	2016	Female	80 to 84	52% (40-66%)	48% (34-60%)	1.6 (0.9-2.7)	0.8% (-3.1-6%)	0.87% (-3.9-7.2%)	38 (-180-310)	460 (-1900-3400)
Azerbaijan	2016	Female	85 to 89	52% (40-63%)	48% (37-60%)	1.6 (0.89-2.6)	0.051% (-4-5.2%)	-0.094% (-5.1-6.1%)	-3 (-150-180)	12 (-1100-1400)
Azerbaijan	2016	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.6 (0.87-2.7)	-0.61% (-4.8-5%)	-0.92% (-5.8-5.6%)	-13 (-87-80)	-54 (-420-420)
Azerbaijan	2016	Female	95 plus	53% (40-68%)	47% (32-60%)	1.6 (0.73-2.8)	-1% (-5.2-5.6%)	-1.4% (-6.5-6.2%)	-5 (-23-21)	-14 (-72-74)
Azerbaijan	2016	Male	All Ages	56% (47-64%)	44% (36-53%)	3.1 (2.1-4.2)	9.3% (6.2-13%)	9.4% (5.1-14%)	3800 (2000-6200)	160000 (100000-230000)
Azerbaijan	2016	Male	15 to 19	36% (22-54%)	64% (46-78%)	0.59 (0.29-1.1)	4.5% (3.2-6.1%)	5.2% (3.2-7.8%)	19 (11-31)	2200 (1500-3200)
Azerbaijan	2016	Male	20 to 24	57% (37-74%)	43% (26-63%)	2 (1.1-3.2)	12% (8.5-15%)	13% (8.5-18%)	90 (52-140)	9000 (6000-13000)
Azerbaijan	2016	Male	25 to 29	68% (49-82%)	32% (18-51%)	4.4 (2.8-6.7)	17% (13-22%)	22% (15-29%)	190 (120-290)	16000 (11000-21000)
Azerbaijan	2016	Male	30 to 34	75% (56-87%)	25% (13-44%)	5.7 (3.6-8.4)	19% (14-24%)	25% (18-32%)	250 (160-370)	18000 (13000-25000)
Azerbaijan	2016	Male	35 to 39	77% (62-88%)	23% (12-38%)	5.2 (3.3-7.9)	17% (12-23%)	22% (14-31%)	210 (130-330)	14000 (9000-20000)
Azerbaijan	2016	Male	40 to 44	77% (62-89%)	23% (11-38%)	5.2 (3.3-7.7)	16% (11-22%)	20% (12-29%)	250 (140-400)	14000 (8700-21000)
Azerbaijan	2016	Male	45 to 49	76% (57-88%)	24% (12-43%)	5.7 (3.6-8.8)	16% (11-22%)	20% (13-28%)	340 (200-520)	16000 (10000-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Azerbaijan	2016	Male	50 to 54	74% (57-87%)	26% (13-43%)	5.5 (3.4-8.3)	15% (9.6-21%)	18% (11-26%)	600 (340-930)	25000 (15000-36000)
Azerbaijan	2016	Male	55 to 59	72% (55-86%)	28% (14-45%)	4.3 (2.6-6.7)	12% (6.1-18%)	14% (6.6-22%)	530 (240-890)	19000 (9600-30000)
Azerbaijan	2016	Male	60 to 64	69% (49-85%)	31% (15-51%)	3.5 (1.9-5.6)	9% (2.8-15%)	10% (2.9-18%)	500 (150-910)	15000 (4900-26000)
Azerbaijan	2016	Male	65 to 69	66% (47-82%)	34% (18-53%)	3 (1.7-4.8)	7.3% (2.5-13%)	8.3% (2.5-15%)	220 (61-420)	5300 (1700-9900)
Azerbaijan	2016	Male	70 to 74	61% (43-78%)	39% (22-57%)	2.5 (1.3-4.2)	5.6% (1-12%)	6.2% (0.72-14%)	170 (21-370)	3200 (570-7200)
Azerbaijan	2016	Male	75 to 79	57% (37-74%)	43% (26-63%)	2 (0.98-3.5)	4.5% (0.52-10%)	4.9% (0.15-12%)	220 (6.5-540)	3400 (350-7800)
Azerbaijan	2016	Male	80 to 84	53% (34-70%)	47% (30-66%)	1.7 (0.9-3)	3.4% (-0.19-8.7%)	3.7% (-0.61-9.7%)	160 (-28-440)	1900 (-110-4800)
Azerbaijan	2016	Male	85 to 89	51% (33-68%)	49% (32-67%)	1.6 (0.8-2.7)	2.6% (-0.55-7.5%)	2.7% (-1.1-8.3%)	68 (-26-210)	570 (-120-1600)
Azerbaijan	2016	Male	90 to 94	50% (34-67%)	50% (33-66%)	1.5 (0.72-2.6)	1.7% (-1.4-5.9%)	1.6% (-2.1-6.5%)	15 (-18-57)	92 (-74-300)
Azerbaijan	2016	Male	95 plus	51% (30-70%)	49% (30-70%)	1.4 (0.57-2.6)	1.5% (-1.8-5.6%)	1.2% (-2.7-5.9%)	1.3 (-2.8-6.4)	6 (-7.2-23)
Georgia	1990	Female	All Ages	53% (49-58%)	47% (42-51%)	0.41 (0.31-0.55)	-0.63% (-1.7-0.35%)	-2.4% (-4.5--0.46%)	-650 (-1200--130)	-5800 (-16000-3200)
Georgia	1990	Female	15 to 19	45% (33-57%)	55% (43-67%)	0.33 (0.19-0.52)	0.89% (0.5-1.4%)	0.99% (0.13-2.1%)	0.81 (0.11-1.7)	180 (99-290)
Georgia	1990	Female	20 to 24	61% (48-71%)	39% (29-52%)	0.52 (0.31-0.81)	1.6% (0.94-2.4%)	1.1% (-0.2-2.6%)	1.4 (-0.28-3.3)	380 (210-590)
Georgia	1990	Female	25 to 29	65% (54-73%)	35% (27-46%)	0.58 (0.36-0.85)	1.7% (1-2.5%)	1.1% (-0.23-2.5%)	2 (-0.41-4.5)	540 (300-810)
Georgia	1990	Female	30 to 34	66% (55-75%)	34% (25-45%)	0.6 (0.38-0.89)	1.6% (0.94-2.3%)	1.2% (-0.24-2.6%)	2.6 (-0.53-5.7)	540 (290-830)
Georgia	1990	Female	35 to 39	66% (55-74%)	34% (26-45%)	0.58 (0.37-0.91)	1.4% (0.68-2.1%)	1.4% (-0.12-2.9%)	3.4 (-0.28-7.5)	460 (220-730)
Georgia	1990	Female	40 to 44	65% (55-74%)	35% (26-45%)	0.58 (0.37-0.94)	1.2% (0.41-2%)	1.4% (-0.22-3.1%)	4.4 (-0.69-10)	390 (140-690)
Georgia	1990	Female	45 to 49	65% (53-74%)	35% (26-47%)	0.59 (0.35-0.92)	0.72% (-0.25-1.7%)	0.73% (-1.2-2.6%)	2.9 (-4.5-10)	240 (-74-540)
Georgia	1990	Female	50 to 54	63% (53-72%)	37% (28-47%)	0.57 (0.35-0.89)	0.39% (-0.83-1.5%)	0.33% (-1.7-2.2%)	3.1 (-16-21)	230 (-500-900)
Georgia	1990	Female	55 to 59	60% (48-70%)	40% (30-52%)	0.54 (0.32-0.83)	-0.19% (-1.6-1%)	-0.43% (-2.6-1.5%)	-5.1 (-31-17)	-120 (-970-620)
Georgia	1990	Female	60 to 64	57% (45-67%)	43% (33-55%)	0.54 (0.32-0.82)	-1.1% (-2.8-0.33%)	-1.7% (-4.1-0.49%)	-36 (-89-11)	-970 (-2400-280)
Georgia	1990	Female	65 to 69	55% (44-65%)	45% (35-56%)	0.55 (0.35-0.79)	-1.8% (-3.8-0.054%)	-2.4% (-5.1-0.016%)	-58 (-120-0.36)	-1400 (-2900--41)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	1990	Female	70 to 74	52% (40-62%)	48% (38-60%)	0.54 (0.33-0.8)	-2.3% (-4.5--0.38%)	-2.9% (-5.8--0.39%)	-80 (-160--11)	-1500 (-3000--250)
Georgia	1990	Female	75 to 79	49% (36-62%)	51% (38-64%)	0.5 (0.29-0.76)	-2.7% (-4.9--0.84%)	-3.4% (-6.2--1%)	-150 (-270--44)	-2100 (-3800--620)
Georgia	1990	Female	80 to 84	47% (33-59%)	53% (41-67%)	0.46 (0.27-0.68)	-2.9% (-5.2--0.96%)	-3.6% (-6.5--1.2%)	-150 (-260--48)	-1500 (-2700--500)
Georgia	1990	Female	85 to 89	45% (32-58%)	55% (42-68%)	0.44 (0.25-0.68)	-3% (-5.3--1.2%)	-3.7% (-6.5--1.4%)	-120 (-210--46)	-870 (-1500--340)
Georgia	1990	Female	90 to 94	45% (33-58%)	55% (42-67%)	0.44 (0.24-0.71)	-3.2% (-5.5--1.2%)	-3.8% (-6.6--1.5%)	-62 (-110--24)	-310 (-520--120)
Georgia	1990	Female	95 plus	47% (31-63%)	53% (37-69%)	0.46 (0.22-0.8)	-3.4% (-5.9--1.1%)	-4.1% (-7.2--1.3%)	-21 (-36--6.5)	-66 (-110--21)
Georgia	1990	Male	All Ages	80% (76-83%)	20% (17-24%)	3 (2.4-3.8)	8.5% (5.7-12%)	8.9% (4.8-14%)	2400 (1300-3800)	94000 (61000-130000)
Georgia	1990	Male	15 to 19	70% (59-80%)	30% (20-41%)	1.4 (0.94-2)	5% (3.6-7%)	7% (4.5-10%)	15 (9.4-22)	1500 (1000-2000)
Georgia	1990	Male	20 to 24	85% (78-90%)	15% (10-22%)	3 (2.2-4)	11% (8.1-15%)	14% (9.4-19%)	49 (33-69)	4200 (3000-5600)
Georgia	1990	Male	25 to 29	88% (83-92%)	12% (8.3-17%)	4.5 (3.2-5.9)	15% (11-20%)	19% (14-26%)	99 (68-130)	7300 (5300-9700)
Georgia	1990	Male	30 to 34	90% (85-93%)	10% (7-15%)	5.2 (3.8-6.9)	16% (12-21%)	21% (15-28%)	130 (90-170)	8500 (6300-11000)
Georgia	1990	Male	35 to 39	91% (87-94%)	9.3% (6.4-13%)	4.9 (3.5-6.5)	15% (10-19%)	19% (12-25%)	130 (84-180)	7700 (5300-10000)
Georgia	1990	Male	40 to 44	91% (87-94%)	9.1% (6.3-13%)	4.8 (3.6-6.5)	14% (9.8-19%)	18% (12-24%)	140 (91-200)	7400 (5000-9800)
Georgia	1990	Male	45 to 49	90% (85-93%)	9.5% (6.6-15%)	5.4 (3.9-7.3)	15% (11-20%)	19% (13-25%)	180 (120-240)	7900 (5400-11000)
Georgia	1990	Male	50 to 54	91% (87-93%)	9.4% (6.6-13%)	5.4 (4-7.2)	14% (9.4-19%)	17% (11-23%)	360 (220-490)	14000 (9000-19000)
Georgia	1990	Male	55 to 59	91% (87-94%)	8.7% (5.8-13%)	4.6 (3.4-6.1)	12% (7-17%)	14% (8.1-20%)	340 (190-500)	11000 (6600-16000)
Georgia	1990	Male	60 to 64	90% (85-93%)	10% (7-15%)	4.1 (2.9-5.6)	10% (4.8-16%)	12% (5.3-18%)	420 (190-670)	12000 (5900-19000)
Georgia	1990	Male	65 to 69	86% (80-90%)	14% (9.6-20%)	3.8 (2.7-5)	8.5% (3.4-14%)	9.7% (3.6-16%)	260 (98-450)	6200 (2500-10000)
Georgia	1990	Male	70 to 74	83% (77-88%)	17% (12-23%)	3.3 (2.4-4.4)	6.6% (1.7-12%)	7.3% (1.6-14%)	160 (35-300)	3000 (770-5600)
Georgia	1990	Male	75 to 79	82% (73-88%)	18% (12-27%)	2.9 (2-3.9)	4.3% (-0.099-9.8%)	4.6% (-0.53-11%)	130 (-16-330)	2100 (-52-4800)
Georgia	1990	Male	80 to 84	80% (70-87%)	20% (13-30%)	2.4 (1.6-3.2)	2.1% (-1.5-6.7%)	2% (-2.2-7.5%)	49 (-54-180)	610 (-450-2000)
Georgia	1990	Male	85 to 89	79% (69-86%)	21% (14-31%)	2 (1.3-2.8)	0.48% (-2.7-4.2%)	0.16% (-3.6-4.6%)	2.5 (-56-71)	64 (-360-580)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	1990	Male	90 to 94	79% (69-86%)	21% (14-31%)	1.8 (1.1-2.7)	-0.89% (-4-2.7%)	-1.5% (-5.3-2.7%)	-9.9 (-35-19)	-35 (-160-110)
Georgia	1990	Male	95 plus	78% (65-88%)	22% (12-35%)	1.5 (0.77-2.5)	-1.7% (-4.8-2%)	-2.5% (-6.3-1.8%)	-4.5 (-11-3.2)	-12 (-34-14)
Georgia	1995	Female	All Ages	55% (51-58%)	45% (42-49%)	0.37 (0.29-0.47)	-0.92% (-2.1-0.097%)	-3% (-5.3--0.86%)	-730 (-1300--210)	-7200 (-16000-760)
Georgia	1995	Female	15 to 19	53% (45-61%)	47% (39-55%)	0.34 (0.22-0.49)	0.78% (0.42-1.2%)	0.62% (-0.27-1.6%)	0.41 (-0.18-1)	140 (74-230)
Georgia	1995	Female	20 to 24	65% (58-71%)	35% (29-42%)	0.48 (0.33-0.7)	1.4% (0.77-2.2%)	0.5% (-0.8-1.9%)	0.47 (-0.76-1.8)	280 (140-450)
Georgia	1995	Female	25 to 29	67% (61-73%)	33% (27-39%)	0.52 (0.36-0.72)	1.3% (0.63-2.1%)	0.23% (-1.2-1.7%)	0.31 (-1.7-2.3)	310 (140-500)
Georgia	1995	Female	30 to 34	68% (62-73%)	32% (27-38%)	0.53 (0.37-0.74)	1.3% (0.55-2%)	0.4% (-1.2-2%)	0.77 (-2.3-3.8)	400 (160-670)
Georgia	1995	Female	35 to 39	66% (60-72%)	34% (28-40%)	0.5 (0.34-0.74)	1.1% (0.36-1.7%)	0.6% (-0.96-2.1%)	1.4 (-2.2-5.2)	350 (110-590)
Georgia	1995	Female	40 to 44	65% (59-72%)	35% (28-41%)	0.5 (0.34-0.74)	0.93% (0.22-1.7%)	0.9% (-0.65-2.5%)	2.8 (-2-7.7)	310 (78-570)
Georgia	1995	Female	45 to 49	65% (59-72%)	35% (28-41%)	0.51 (0.34-0.75)	0.42% (-0.49-1.3%)	0.15% (-1.6-2%)	0.57 (-6.1-7.2)	140 (-150-420)
Georgia	1995	Female	50 to 54	64% (57-70%)	36% (30-43%)	0.49 (0.32-0.72)	0.073% (-1-1.1%)	-0.21% (-2.2-1.6%)	-0.99 (-10-7.8)	24 (-330-360)
Georgia	1995	Female	55 to 59	59% (52-65%)	41% (35-48%)	0.44 (0.29-0.65)	-0.39% (-1.7-0.73%)	-0.78% (-2.8-1.1%)	-8.9 (-32-12)	-240 (-1000-450)
Georgia	1995	Female	60 to 64	55% (49-62%)	45% (38-51%)	0.44 (0.29-0.63)	-1.1% (-2.6-0.17%)	-1.7% (-3.9-0.25%)	-26 (-58-3.7)	-700 (-1600-120)
Georgia	1995	Female	65 to 69	55% (49-61%)	45% (39-51%)	0.47 (0.34-0.65)	-1.9% (-3.8--0.28%)	-2.6% (-5.3--0.3%)	-68 (-140--8.3)	-1600 (-3200--250)
Georgia	1995	Female	70 to 74	53% (46-60%)	47% (40-54%)	0.47 (0.33-0.67)	-2.3% (-4.4--0.59%)	-3% (-5.7--0.66%)	-89 (-170--19)	-1700 (-3100--430)
Georgia	1995	Female	75 to 79	53% (44-61%)	47% (39-56%)	0.47 (0.31-0.66)	-3.1% (-5.2--1.2%)	-3.9% (-6.6--1.4%)	-120 (-200--43)	-1700 (-2800--620)
Georgia	1995	Female	80 to 84	53% (40-65%)	47% (35-60%)	0.45 (0.29-0.67)	-3.5% (-5.9--1.3%)	-4.3% (-7.4--1.6%)	-160 (-270--60)	-1600 (-2800--630)
Georgia	1995	Female	85 to 89	51% (39-62%)	49% (38-61%)	0.43 (0.28-0.62)	-3.5% (-6.1--1.5%)	-4.3% (-7.4--1.7%)	-150 (-270--63)	-1100 (-1900--460)
Georgia	1995	Female	90 to 94	51% (39-62%)	49% (38-61%)	0.43 (0.25-0.66)	-3.6% (-6.1--1.4%)	-4.4% (-7.4--1.7%)	-85 (-140--32)	-420 (-700--160)
Georgia	1995	Female	95 plus	52% (36-68%)	48% (32-64%)	0.44 (0.22-0.73)	-3.8% (-6.4--1.2%)	-4.6% (-7.6--1.5%)	-29 (-49--9.6)	-95 (-160--32)
Georgia	1995	Male	All Ages	84% (82-87%)	16% (13-18%)	2.9 (2.4-3.4)	7.2% (4.5-9.9%)	6.6% (2.7-11%)	1600 (650-2600)	68000 (42000-97000)
Georgia	1995	Male	15 to 19	80% (74-85%)	20% (15-26%)	1.5 (1.1-2)	4.8% (3.4-6.4%)	6.9% (4.4-9.6%)	12 (7.4-17)	1200 (810-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	1995	Male	20 to 24	89% (86-92%)	11% (8.5-14%)	2.9 (2.3-3.6)	10% (7.6-13%)	12% (8.7-17%)	41 (28-56)	3500 (2500-4600)
Georgia	1995	Male	25 to 29	91% (89-93%)	9.1% (7.3-11%)	4.2 (3.3-5.3)	14% (10-17%)	17% (12-23%)	70 (49-94)	5200 (3800-6700)
Georgia	1995	Male	30 to 34	92% (90-94%)	8% (6.2-10%)	4.8 (3.8-6)	14% (11-18%)	19% (13-24%)	100 (72-140)	6900 (5100-8900)
Georgia	1995	Male	35 to 39	93% (91-94%)	7.4% (5.7-9.3%)	4.5 (3.5-5.6)	13% (9-17%)	16% (10-21%)	110 (72-150)	6700 (4700-8900)
Georgia	1995	Male	40 to 44	93% (91-95%)	7.1% (5.5-9%)	4.5 (3.6-5.6)	12% (7.9-16%)	15% (9.1-21%)	120 (73-170)	6500 (4200-8900)
Georgia	1995	Male	45 to 49	93% (91-94%)	7.4% (5.8-9.2%)	5.1 (3.9-6.4)	13% (8.6-18%)	16% (10-22%)	150 (91-210)	6800 (4500-9300)
Georgia	1995	Male	50 to 54	93% (91-94%)	7.2% (5.6-8.9%)	5.1 (3.9-6.4)	13% (7.6-17%)	15% (8.6-21%)	160 (89-230)	6300 (3800-9000)
Georgia	1995	Male	55 to 59	93% (91-95%)	6.5% (4.8-9%)	4.4 (3.4-5.4)	10% (5.2-15%)	12% (5.7-18%)	270 (130-410)	9300 (4800-14000)
Georgia	1995	Male	60 to 64	92% (90-94%)	7.8% (6-10%)	3.9 (3-4.9)	7.6% (2.7-13%)	8.7% (2.5-15%)	230 (66-400)	6700 (2300-12000)
Georgia	1995	Male	65 to 69	89% (86-92%)	11% (8.5-14%)	3.5 (2.8-4.4)	6% (1.4-11%)	6.8% (1.1-13%)	250 (42-470)	6100 (1500-11000)
Georgia	1995	Male	70 to 74	87% (83-89%)	13% (11-17%)	3.1 (2.4-4)	4.3% (-0.11-9%)	4.7% (-0.66-10%)	120 (-16-270)	2500 (-58-5200)
Georgia	1995	Male	75 to 79	86% (81-90%)	14% (10-19%)	2.8 (2.1-3.6)	2.3% (-1.9-7%)	2.2% (-2.8-7.7%)	41 (-51-140)	670 (-590-2000)
Georgia	1995	Male	80 to 84	84% (78-89%)	16% (11-22%)	2.3 (1.7-2.9)	0.38% (-3.2-4.2%)	0.059% (-4.2-4.6%)	1.1 (-84-93)	91 (-780-1000)
Georgia	1995	Male	85 to 89	83% (77-88%)	17% (12-23%)	1.9 (1.4-2.6)	-0.89% (-4.1-2.6%)	-1.4% (-5.4-2.7%)	-25 (-94-48)	-130 (-620-390)
Georgia	1995	Male	90 to 94	82% (75-88%)	18% (12-25%)	1.7 (1.1-2.4)	-2% (-5.2-1.5%)	-2.8% (-6.7-1.3%)	-21 (-50-9.6)	-88 (-230-66)
Georgia	1995	Male	95 plus	81% (71-90%)	19% (10-29%)	1.4 (0.78-2.3)	-2.5% (-5.6-0.81%)	-3.4% (-7.1-0.51%)	-7.1 (-15-1)	-20 (-46-6.6)
Georgia	2000	Female	All Ages	51% (47-55%)	49% (45-53%)	0.33 (0.26-0.42)	-0.74% (-1.7-0.13%)	-2.5% (-4.4--0.72%)	-580 (-1000--170)	-5300 (-12000-970)
Georgia	2000	Female	15 to 19	44% (33-54%)	56% (46-67%)	0.27 (0.17-0.4)	0.78% (0.46-1.2%)	0.66% (0.037-1.4%)	0.41 (0.02-0.88)	140 (78-220)
Georgia	2000	Female	20 to 24	58% (47-69%)	42% (31-53%)	0.41 (0.26-0.62)	1.5% (0.93-2.2%)	0.77% (-0.33-1.9%)	0.59 (-0.26-1.5)	280 (160-450)
Georgia	2000	Female	25 to 29	62% (52-71%)	38% (29-48%)	0.46 (0.3-0.65)	1.6% (0.99-2.3%)	0.69% (-0.51-1.9%)	0.64 (-0.48-1.8)	310 (170-470)
Georgia	2000	Female	30 to 34	63% (52-73%)	37% (27-48%)	0.47 (0.31-0.67)	1.4% (0.71-2.1%)	0.63% (-0.87-2%)	0.79 (-1-2.6)	310 (150-480)
Georgia	2000	Female	35 to 39	63% (52-72%)	37% (28-48%)	0.45 (0.29-0.67)	1.1% (0.47-1.7%)	0.64% (-0.8-2%)	1.3 (-1.6-4.1)	350 (140-550)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	2000	Female	40 to 44	62% (52-71%)	38% (29-48%)	0.45 (0.29-0.68)	1% (0.41-1.6%)	1.1% (-0.26-2.4%)	2.9 (-0.68-6.6)	340 (130-550)
Georgia	2000	Female	45 to 49	62% (51-71%)	38% (29-49%)	0.46 (0.29-0.68)	0.6% (-0.12-1.3%)	0.52% (-1.1-2%)	1.8 (-3.5-6.7)	200 (-36-420)
Georgia	2000	Female	50 to 54	60% (50-69%)	40% (31-50%)	0.44 (0.27-0.67)	0.27% (-0.66-1.1%)	0.1% (-1.6-1.7%)	0.44 (-7.4-7.1)	91 (-210-350)
Georgia	2000	Female	55 to 59	56% (46-66%)	44% (34-54%)	0.41 (0.25-0.65)	-0.22% (-1.3-0.75%)	-0.54% (-2.3-1.1%)	-3.2 (-14-6.7)	-72 (-420-240)
Georgia	2000	Female	60 to 64	53% (41-64%)	47% (36-59%)	0.4 (0.26-0.61)	-0.99% (-2.4-0.18%)	-1.5% (-3.7-0.32%)	-25 (-60-4.9)	-660 (-1600-140)
Georgia	2000	Female	65 to 69	50% (39-60%)	50% (40-61%)	0.41 (0.28-0.58)	-1.6% (-3.3--0.2%)	-2.2% (-4.5--0.2%)	-46 (-97--3.9)	-1100 (-2300--150)
Georgia	2000	Female	70 to 74	47% (37-57%)	53% (43-63%)	0.4 (0.26-0.59)	-1.9% (-3.7--0.38%)	-2.4% (-4.8--0.41%)	-87 (-170--14)	-1600 (-3200--360)
Georgia	2000	Female	75 to 79	46% (35-58%)	54% (42-65%)	0.38 (0.23-0.57)	-2.3% (-4.1--0.74%)	-2.9% (-5.4--0.88%)	-110 (-190--33)	-1500 (-2700--490)
Georgia	2000	Female	80 to 84	45% (32-57%)	55% (43-68%)	0.36 (0.22-0.54)	-2.6% (-4.7--0.93%)	-3.3% (-5.9--1.2%)	-100 (-190--37)	-1100 (-1900--390)
Georgia	2000	Female	85 to 89	44% (32-57%)	56% (43-68%)	0.35 (0.21-0.53)	-2.8% (-5.1--1.1%)	-3.5% (-6.3--1.4%)	-110 (-200--44)	-810 (-1500--320)
Georgia	2000	Female	90 to 94	45% (32-58%)	55% (42-68%)	0.36 (0.19-0.57)	-2.9% (-5--1.1%)	-3.6% (-6.1--1.3%)	-78 (-130--30)	-380 (-650--150)
Georgia	2000	Female	95 plus	47% (30-63%)	53% (37-70%)	0.37 (0.18-0.66)	-3.1% (-5.3--0.97%)	-3.7% (-6.5--1.2%)	-29 (-50--9.1)	-92 (-160--29)
Georgia	2000	Male	All Ages	83% (81-86%)	17% (14-19%)	2.8 (2.3-3.2)	6.6% (4-9.1%)	5.7% (2-9.7%)	1400 (490-2300)	56000 (34000-80000)
Georgia	2000	Male	15 to 19	77% (70-83%)	23% (17-30%)	1.4 (1-1.8)	4.3% (3.1-5.5%)	6.3% (4.1-8.8%)	7.9 (5.1-11)	890 (630-1200)
Georgia	2000	Male	20 to 24	88% (85-91%)	12% (9-15%)	2.7 (2.1-3.4)	9.8% (7.6-12%)	13% (9.1-17%)	28 (20-39)	2600 (1900-3400)
Georgia	2000	Male	25 to 29	90% (87-92%)	10% (7.7-13%)	3.9 (3.1-4.9)	13% (10-16%)	17% (13-22%)	52 (37-68)	4000 (3100-5100)
Georgia	2000	Male	30 to 34	92% (89-94%)	8.5% (6.4-11%)	4.6 (3.6-5.6)	14% (10-18%)	18% (13-24%)	69 (49-92)	4700 (3500-6100)
Georgia	2000	Male	35 to 39	92% (90-94%)	8% (6.1-10%)	4.2 (3.3-5.2)	12% (8.8-16%)	15% (10-21%)	92 (61-130)	5600 (4000-7400)
Georgia	2000	Male	40 to 44	92% (90-94%)	7.8% (5.7-9.8%)	4.2 (3.3-5.2)	12% (7.8-15%)	14% (8.9-20%)	110 (68-160)	6100 (3900-8100)
Georgia	2000	Male	45 to 49	92% (90-94%)	8.1% (6.1-10%)	4.7 (3.7-5.9)	12% (8.3-17%)	15% (9.6-22%)	150 (89-210)	6800 (4400-9300)
Georgia	2000	Male	50 to 54	92% (90-94%)	7.8% (5.9-10%)	4.7 (3.7-5.9)	12% (6.9-16%)	14% (8-20%)	140 (80-210)	5900 (3500-8300)
Georgia	2000	Male	55 to 59	93% (90-95%)	7.2% (5.1-9.6%)	4.1 (3.2-5)	9.2% (4.7-14%)	11% (5-17%)	130 (59-200)	4500 (2300-6800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	2000	Male	60 to 64	91% (89-94%)	8.6% (6.4-11%)	3.6 (2.8-4.6)	6.9% (2.4-12%)	7.9% (2.4-14%)	210 (60-370)	6100 (2100-11000)
Georgia	2000	Male	65 to 69	88% (84-91%)	12% (9-16%)	3.3 (2.6-4.1)	5.4% (1.2-9.9%)	6% (0.96-11%)	200 (31-370)	4700 (1100-8700)
Georgia	2000	Male	70 to 74	86% (82-89%)	14% (11-18%)	2.9 (2.3-3.6)	3.7% (-0.3-8.1%)	4.1% (-0.77-9.2%)	180 (-34-400)	3500 (-290-7600)
Georgia	2000	Male	75 to 79	84% (78-89%)	16% (11-22%)	2.5 (1.9-3.2)	1.9% (-1.7-6.1%)	1.9% (-2.3-6.7%)	50 (-63-180)	840 (-790-2700)
Georgia	2000	Male	80 to 84	82% (73-88%)	18% (12-27%)	2.1 (1.5-2.7)	0.21% (-3.1-3.6%)	-0.1% (-4-3.9%)	-1.7 (-67-65)	42 (-610-730)
Georgia	2000	Male	85 to 89	81% (72-87%)	19% (13-28%)	1.7 (1.2-2.4)	-1% (-4-2.1%)	-1.5% (-5.1-2.1%)	-20 (-64-28)	-110 (-430-230)
Georgia	2000	Male	90 to 94	80% (71-87%)	20% (13-29%)	1.5 (0.99-2.2)	-1.8% (-4.7-1.2%)	-2.5% (-6.1-1%)	-20 (-49-7.9)	-85 (-230-56)
Georgia	2000	Male	95 plus	79% (67-89%)	21% (11-33%)	1.3 (0.68-2.1)	-2.2% (-4.9-0.77%)	-3% (-6.3-0.42%)	-6.3 (-14-0.87)	-18 (-42-6.5)
Georgia	2005	Female	All Ages	51% (47-55%)	49% (45-53%)	0.36 (0.28-0.46)	-0.56% (-1.5-0.3%)	-2% (-3.8--0.34%)	-480 (-930--82)	-4000 (-11000-2200)
Georgia	2005	Female	15 to 19	43% (33-54%)	57% (46-67%)	0.28 (0.17-0.43)	0.73% (0.43-1.1%)	0.5% (-0.087-1.2%)	0.32 (-0.057-0.78)	140 (74-220)
Georgia	2005	Female	20 to 24	58% (47-68%)	42% (32-53%)	0.44 (0.28-0.65)	1.5% (0.85-2.2%)	0.66% (-0.4-1.8%)	0.52 (-0.33-1.4)	280 (150-430)
Georgia	2005	Female	25 to 29	62% (51-70%)	38% (30-49%)	0.48 (0.32-0.69)	1.6% (1-2.1%)	0.88% (-0.21-2%)	0.82 (-0.21-1.9)	310 (190-450)
Georgia	2005	Female	30 to 34	63% (51-72%)	37% (28-49%)	0.49 (0.33-0.71)	1.4% (0.84-2%)	0.94% (-0.34-2.1%)	1.2 (-0.44-2.8)	310 (170-470)
Georgia	2005	Female	35 to 39	62% (51-72%)	38% (28-49%)	0.48 (0.3-0.72)	1.2% (0.64-1.8%)	1.1% (-0.089-2.3%)	1.8 (-0.15-3.8)	300 (150-440)
Georgia	2005	Female	40 to 44	61% (52-71%)	39% (29-48%)	0.47 (0.31-0.71)	1.1% (0.5-1.7%)	1.3% (0.11-2.5%)	4 (0.33-7.7)	390 (170-600)
Georgia	2005	Female	45 to 49	61% (50-71%)	39% (29-50%)	0.48 (0.3-0.72)	0.7% (0.0079-1.4%)	0.81% (-0.63-2.2%)	3.5 (-2.7-9.3)	280 (3.2-540)
Georgia	2005	Female	50 to 54	60% (50-69%)	40% (31-50%)	0.46 (0.29-0.72)	0.37% (-0.51-1.1%)	0.39% (-1.2-1.8%)	2.2 (-6.7-10)	150 (-210-450)
Georgia	2005	Female	55 to 59	56% (45-65%)	44% (35-55%)	0.43 (0.27-0.67)	-0.062% (-1-0.81%)	-0.19% (-1.8-1.3%)	-1.4 (-13-9.5)	-25 (-420-320)
Georgia	2005	Female	60 to 64	52% (41-62%)	48% (38-59%)	0.42 (0.27-0.65)	-0.74% (-2-0.33%)	-1.1% (-3.1-0.55%)	-11 (-30-4.9)	-300 (-830-130)
Georgia	2005	Female	65 to 69	50% (39-59%)	50% (41-61%)	0.43 (0.3-0.62)	-1.2% (-2.8-0.06%)	-1.6% (-3.7-0.21%)	-40 (-91-5.1)	-980 (-2100-45)
Georgia	2005	Female	70 to 74	47% (36-57%)	53% (43-64%)	0.42 (0.27-0.63)	-1.5% (-3.1--0.12%)	-1.9% (-3.9-0.014%)	-56 (-120-0.42)	-1100 (-2300--83)
Georgia	2005	Female	75 to 79	45% (34-57%)	55% (43-66%)	0.4 (0.25-0.62)	-1.9% (-3.6--0.47%)	-2.4% (-4.5--0.54%)	-110 (-200--25)	-1600 (-2900--400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	2005	Female	80 to 84	44% (31-57%)	56% (43-69%)	0.38 (0.23-0.58)	-2.2% (-4--0.65%)	-2.7% (-5--0.77%)	-110 (-200--31)	-1200 (-2100--360)
Georgia	2005	Female	85 to 89	44% (31-56%)	56% (44-69%)	0.37 (0.22-0.57)	-2.4% (-4.4--0.81%)	-2.9% (-5.4--0.97%)	-84 (-160--28)	-610 (-1100--210)
Georgia	2005	Female	90 to 94	44% (31-57%)	56% (43-69%)	0.38 (0.21-0.61)	-2.5% (-4.4--0.86%)	-3.1% (-5.3--1%)	-59 (-100--20)	-290 (-510--100)
Georgia	2005	Female	95 plus	46% (30-63%)	54% (37-70%)	0.39 (0.19-0.7)	-2.7% (-4.7--0.81%)	-3.3% (-5.8--1%)	-29 (-51--9)	-94 (-170--29)
Georgia	2005	Male	All Ages	84% (81-86%)	16% (14-19%)	3.1 (2.6-3.6)	7.7% (5.1-10%)	7% (3.3-11%)	1900 (890-2900)	68000 (45000-92000)
Georgia	2005	Male	15 to 19	77% (70-83%)	23% (17-30%)	1.5 (1.1-1.9)	4.4% (3.1-5.8%)	6.1% (3.7-8.6%)	9.8 (5.9-14)	1100 (740-1400)
Georgia	2005	Male	20 to 24	88% (84-91%)	12% (9.3-16%)	2.9 (2.3-3.6)	10% (7.7-13%)	13% (9.1-17%)	32 (22-45)	2900 (2100-3800)
Georgia	2005	Male	25 to 29	90% (87-92%)	10% (7.9-13%)	4.2 (3.3-5.3)	14% (10-17%)	17% (12-23%)	51 (35-67)	3900 (2900-5000)
Georgia	2005	Male	30 to 34	91% (89-93%)	8.7% (6.6-11%)	4.9 (3.9-6.2)	15% (11-18%)	19% (13-24%)	73 (52-96)	5000 (3700-6400)
Georgia	2005	Male	35 to 39	92% (90-94%)	8.2% (6.2-10%)	4.5 (3.5-5.7)	18% (9.8-17%)	17% (12-22%)	81 (54-110)	4900 (3500-6500)
Georgia	2005	Male	40 to 44	92% (90-94%)	8% (5.9-10%)	4.5 (3.5-5.6)	13% (9.1-16%)	16% (11-21%)	120 (82-170)	6600 (4600-8700)
Georgia	2005	Male	45 to 49	92% (89-94%)	8.3% (6.3-11%)	5.1 (4-6.3)	14% (9.8-18%)	17% (11-23%)	190 (130-250)	8600 (6100-11000)
Georgia	2005	Male	50 to 54	92% (90-94%)	8% (6-10%)	5.1 (4-6.3)	13% (8.8-17%)	16% (10-21%)	220 (140-290)	8600 (5700-11000)
Georgia	2005	Male	55 to 59	93% (90-95%)	7.3% (5.2-10%)	4.4 (3.4-5.4)	11% (6.2-15%)	12% (6.9-18%)	190 (110-280)	6600 (3800-9400)
Georgia	2005	Male	60 to 64	91% (88-93%)	8.8% (6.5-12%)	3.9 (2.9-5)	8.4% (3.7-13%)	9.6% (3.9-16%)	160 (66-270)	4700 (2100-7600)
Georgia	2005	Male	65 to 69	88% (84-91%)	12% (9.2-16%)	3.5 (2.8-4.4)	6.9% (2.4-11%)	7.8% (2.5-13%)	290 (93-490)	6900 (2500-12000)
Georgia	2005	Male	70 to 74	85% (81-88%)	15% (12-19%)	3.1 (2.4-3.9)	5.3% (1-9.7%)	5.9% (0.85-11%)	230 (34-440)	4500 (890-8400)
Georgia	2005	Male	75 to 79	84% (78-88%)	16% (12-22%)	2.7 (2.1-3.5)	3.4% (-0.45-8%)	3.7% (-0.87-9%)	180 (-43-440)	2700 (-370-6400)
Georgia	2005	Male	80 to 84	82% (73-88%)	18% (12-27%)	2.2 (1.6-2.9)	1.8% (-1.6-5.5%)	1.7% (-2.1-6.2%)	52 (-60-180)	620 (-550-1900)
Georgia	2005	Male	85 to 89	80% (72-87%)	20% (13-28%)	1.9 (1.3-2.6)	0.32% (-2.6-3.6%)	0.077% (-3.3-3.9%)	0.85 (-38-43)	30 (-250-350)
Georgia	2005	Male	90 to 94	80% (71-87%)	20% (13-29%)	1.7 (1.1-2.4)	-0.75% (-3.6-2.3%)	-1.2% (-4.6-2.3%)	-6.7 (-26-13)	-24 (-120-72)
Georgia	2005	Male	95 plus	79% (66-89%)	21% (11-34%)	1.4 (0.75-2.3)	-1.4% (-4.2-1.6%)	-2% (-5.3-1.5%)	-4.4 (-12-3.3)	-12 (-36-14)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	2010	Female	All Ages	54% (50-58%)	46% (42-50%)	0.48 (0.37-0.61)	-0.46% (-1.4-0.4%)	-2% (-3.8--0.25%)	-450 (-870--55)	-3000 (-9100-2700)
Georgia	2010	Female	15 to 19	44% (33-55%)	56% (45-67%)	0.35 (0.21-0.54)	0.81% (0.45-1.2%)	0.74% (0.023-1.7%)	0.43 (0.013-0.99)	130 (70-210)
Georgia	2010	Female	20 to 24	60% (48-70%)	40% (30-52%)	0.55 (0.33-0.86)	1.7% (1.1-2.5%)	1.2% (0.13-2.5%)	0.94 (0.1-2)	310 (190-490)
Georgia	2010	Female	25 to 29	64% (52-73%)	36% (27-48%)	0.62 (0.4-0.91)	1.8% (1.2-2.5%)	1.4% (0.27-2.7%)	1.2 (0.23-2.3)	340 (220-510)
Georgia	2010	Female	30 to 34	66% (54-74%)	34% (26-46%)	0.63 (0.41-0.93)	1.8% (1.2-2.4%)	1.8% (0.7-3.1%)	2 (0.82-3.4)	380 (240-530)
Georgia	2010	Female	35 to 39	65% (53-74%)	35% (26-47%)	0.62 (0.39-0.93)	1.5% (0.99-2.1%)	1.8% (0.6-3%)	2.6 (0.89-4.6)	340 (210-490)
Georgia	2010	Female	40 to 44	65% (55-73%)	35% (27-45%)	0.62 (0.39-0.93)	1.8% (0.69-2%)	1.8% (0.6-3.2%)	4.2 (1.3-7.5)	350 (180-560)
Georgia	2010	Female	45 to 49	64% (52-74%)	36% (26-48%)	0.63 (0.4-0.98)	0.91% (0.18-1.7%)	1.2% (-0.13-2.8%)	5.3 (-0.58-12)	350 (72-670)
Georgia	2010	Female	50 to 54	63% (51-71%)	37% (29-49%)	0.61 (0.37-0.94)	0.6% (-0.27-1.5%)	0.82% (-0.73-2.5%)	5.1 (-4.7-15)	260 (-120-680)
Georgia	2010	Female	55 to 59	59% (48-69%)	41% (31-52%)	0.57 (0.35-0.87)	0.056% (-0.99-1.1%)	0.027% (-1.7-1.7%)	0.22 (-14-14)	25 (-440-470)
Georgia	2010	Female	60 to 64	57% (45-67%)	43% (33-55%)	0.57 (0.36-0.89)	-0.73% (-2.1-0.47%)	-1.1% (-3.2-0.67%)	-12 (-35-7.5)	-340 (-980-230)
Georgia	2010	Female	65 to 69	54% (43-64%)	46% (36-57%)	0.59 (0.39-0.86)	-1.3% (-2.8-0.066%)	-1.7% (-3.9-0.27%)	-24 (-55-3.7)	-600 (-1300-31)
Georgia	2010	Female	70 to 74	51% (39-62%)	49% (38-61%)	0.57 (0.36-0.84)	-1.5% (-3.2--0.027%)	-1.9% (-4.2-0.11%)	-60 (-130-4)	-1200 (-2500--23)
Georgia	2010	Female	75 to 79	48% (35-61%)	52% (39-65%)	0.52 (0.32-0.82)	-1.8% (-3.6--0.37%)	-2.4% (-4.7--0.44%)	-75 (-150--13)	-1100 (-2200--220)
Georgia	2010	Female	80 to 84	46% (33-59%)	54% (41-67%)	0.48 (0.29-0.73)	-2% (-3.8--0.54%)	-2.6% (-4.9--0.65%)	-100 (-200--26)	-1100 (-2100--300)
Georgia	2010	Female	85 to 89	44% (32-58%)	56% (42-68%)	0.47 (0.27-0.71)	-2.2% (-4.2--0.74%)	-2.8% (-5.3--0.9%)	-110 (-210--35)	-810 (-1500--270)
Georgia	2010	Female	90 to 94	45% (32-58%)	55% (42-68%)	0.47 (0.26-0.76)	-2.4% (-4.3--0.79%)	-3% (-5.3--0.95%)	-58 (-100--19)	-290 (-520--95)
Georgia	2010	Female	95 plus	47% (31-64%)	53% (36-69%)	0.49 (0.22-0.86)	-2.6% (-4.7--0.77%)	-3.2% (-5.9--0.95%)	-27 (-49--8)	-87 (-160--27)
Georgia	2010	Male	All Ages	81% (78-84%)	19% (16-22%)	3.5 (2.9-4.2)	9.7% (6.9-13%)	9.4% (5.4-14%)	2500 (1400-3700)	84000 (59000-110000)
Georgia	2010	Male	15 to 19	69% (58-79%)	31% (21-42%)	1.5 (1-2.1)	4.6% (3.3-6.2%)	6.5% (4.1-9.4%)	8.9 (5.5-13)	940 (660-1300)
Georgia	2010	Male	20 to 24	85% (77-90%)	15% (10-23%)	3.2 (2.4-4.3)	11% (8.3-14%)	14% (9.6-19%)	38 (26-53)	3400 (2500-4500)
Georgia	2010	Male	25 to 29	88% (82-92%)	12% (8.2-18%)	4.8 (3.6-6.2)	15% (11-19%)	19% (14-25%)	59 (41-79)	4500 (3400-5800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	2010	Male	30 to 34	90% (84-93%)	10% (7-16%)	5.6 (4.2-7.4)	16% (12-19%)	21% (16-26%)	75 (55-96)	5100 (3900-6400)
Georgia	2010	Male	35 to 39	91% (86-93%)	9.5% (6.5-14%)	5.2 (3.9-6.7)	15% (11-18%)	19% (14-24%)	92 (68-120)	5500 (4200-7000)
Georgia	2010	Male	40 to 44	91% (87-94%)	9.3% (6.4-13%)	5.2 (3.9-6.6)	15% (11-18%)	18% (14-23%)	130 (91-160)	6500 (4800-8300)
Georgia	2010	Male	45 to 49	90% (86-93%)	9.6% (6.7-14%)	5.8 (4.4-7.6)	16% (12-20%)	19% (14-24%)	230 (170-290)	10000 (7700-13000)
Georgia	2010	Male	50 to 54	91% (87-94%)	9.5% (6.5-13%)	5.8 (4.3-7.5)	15% (11-19%)	18% (13-23%)	290 (200-380)	11000 (8100-15000)
Georgia	2010	Male	55 to 59	91% (87-94%)	8.8% (5.8-13%)	4.9 (3.7-6.5)	13% (8.5-17%)	15% (9.7-21%)	290 (180-400)	9700 (6300-13000)
Georgia	2010	Male	60 to 64	90% (84-93%)	10% (7-16%)	4.4 (3.2-5.9)	11% (6-16%)	12% (6.7-19%)	270 (140-400)	7600 (4200-11000)
Georgia	2010	Male	65 to 69	86% (80-90%)	14% (9.7-20%)	4 (3-5.2)	9.4% (5-14%)	11% (5.4-17%)	230 (120-360)	5500 (2900-8600)
Georgia	2010	Male	70 to 74	83% (77-88%)	17% (12-23%)	3.5 (2.6-4.6)	7.7% (2.9-12%)	8.7% (3-14%)	390 (140-660)	7500 (2800-12000)
Georgia	2010	Male	75 to 79	81% (73-88%)	19% (12-27%)	3 (2.2-4.1)	5.8% (1.3-11%)	6.4% (1.1-12%)	260 (45-480)	3800 (820-7000)
Georgia	2010	Male	80 to 84	80% (69-87%)	20% (13-31%)	2.5 (1.7-3.4)	3.8% (0.14-8.1%)	4.1% (-0.13-9.1%)	150 (-4.6-330)	1600 (58-3500)
Georgia	2010	Male	85 to 89	79% (69-86%)	21% (14-31%)	2.1 (1.4-3)	2% (-1.3-5.9%)	2% (-1.9-6.7%)	44 (-41-140)	370 (-250-1100)
Georgia	2010	Male	90 to 94	78% (68-86%)	22% (14-32%)	1.9 (1.2-2.7)	0.63% (-2.3-4.3%)	0.39% (-3.2-4.7%)	2.2 (-19-27)	21 (-81-150)
Georgia	2010	Male	95 plus	78% (63-88%)	22% (12-37%)	1.6 (0.84-2.6)	-0.28% (-3.3-3.4%)	-0.76% (-4.5-3.5%)	-1.3 (-7.7-6.1)	-1.9 (-23-23)
Georgia	2016	Female	All Ages	54% (50-58%)	46% (42-50%)	0.52 (0.36-0.7)	-0.35% (-1.3-0.51%)	-1.8% (-3.6--0.16%)	-410 (-800--37)	-2100 (-7300-3200)
Georgia	2016	Female	15 to 19	44% (33-56%)	56% (44-67%)	0.38 (0.21-0.61)	0.91% (0.52-1.4%)	1% (0.16-2.2%)	0.37 (0.058-0.81)	94 (51-150)
Georgia	2016	Female	20 to 24	60% (48-70%)	40% (30-52%)	0.6 (0.35-0.98)	1.8% (1.1-2.7%)	1.5% (0.31-3%)	0.9 (0.18-1.9)	270 (160-400)
Georgia	2016	Female	25 to 29	64% (53-73%)	36% (27-47%)	0.67 (0.41-1)	2% (1.4-2.7%)	1.7% (0.56-3.2%)	1.4 (0.43-2.6)	360 (230-530)
Georgia	2016	Female	30 to 34	66% (54-74%)	34% (26-46%)	0.69 (0.42-1)	2% (1.4-2.7%)	2.3% (1-3.7%)	2.2 (1-3.8)	380 (250-560)
Georgia	2016	Female	35 to 39	65% (54-73%)	35% (27-46%)	0.67 (0.41-1)	1.7% (1.1-2.3%)	2.2% (1.1-3.5%)	2.9 (1.3-5)	360 (230-520)
Georgia	2016	Female	40 to 44	65% (55-73%)	35% (27-45%)	0.68 (0.41-1)	1.5% (0.87-2.2%)	2.3% (1-3.8%)	4.4 (1.9-7.8)	360 (210-570)
Georgia	2016	Female	45 to 49	64% (52-74%)	36% (26-48%)	0.69 (0.41-1.1)	1.1% (0.41-2%)	1.8% (0.39-3.4%)	5.3 (1-11)	330 (120-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	2016	Female	50 to 54	63% (51-71%)	37% (29-49%)	0.66 (0.39-1)	0.85% (0.041-1.8%)	1.4% (-0.12-3.1%)	7.3 (-0.61-17)	340 (17-760)
Georgia	2016	Female	55 to 59	59% (48-69%)	41% (31-52%)	0.62 (0.36-0.98)	0.28% (-0.67-1.2%)	0.44% (-1.1-2.1%)	3.5 (-9.6-17)	130 (-330-590)
Georgia	2016	Female	60 to 64	57% (45-68%)	43% (32-55%)	0.62 (0.37-0.96)	-0.51% (-1.8-0.6%)	-0.73% (-2.7-0.99%)	-8.1 (-30-11)	-250 (-880-300)
Georgia	2016	Female	65 to 69	54% (43-64%)	46% (36-57%)	0.64 (0.39-0.94)	-1% (-2.4-0.27%)	-1.3% (-3.4-0.53%)	-17 (-45-7.2)	-460 (-1100-120)
Georgia	2016	Female	70 to 74	51% (39-62%)	49% (38-61%)	0.61 (0.36-0.94)	-1.3% (-2.9-0.12%)	-1.7% (-3.9-0.34%)	-31 (-73-5.8)	-640 (-1400-62)
Georgia	2016	Female	75 to 79	48% (35-61%)	52% (39-65%)	0.56 (0.32-0.87)	-1.6% (-3.3--0.15%)	-2.1% (-4.4-0.13%)	-75 (-150-4.4)	-1100 (-2300--110)
Georgia	2016	Female	80 to 84	46% (33-58%)	54% (42-67%)	0.52 (0.31-0.8)	-1.9% (-3.6--0.42%)	-2.4% (-4.6--0.49%)	-94 (-180--20)	-1000 (-1900--230)
Georgia	2016	Female	85 to 89	45% (32-58%)	55% (42-68%)	0.51 (0.29-0.8)	-2.1% (-4--0.57%)	-2.6% (-5--0.67%)	-100 (-200--27)	-750 (-1400--200)
Georgia	2016	Female	90 to 94	45% (32-59%)	55% (41-68%)	0.52 (0.27-0.87)	-2.2% (-4.1--0.58%)	-2.8% (-5.1--0.73%)	-82 (-150--22)	-410 (-760--110)
Georgia	2016	Female	95 plus	47% (31-64%)	53% (36-69%)	0.54 (0.23-0.99)	-2.4% (-4.6--0.51%)	-3% (-5.6--0.68%)	-29 (-55--6.6)	-92 (-170--19)
Georgia	2016	Male	All Ages	82% (78-84%)	18% (16-22%)	3.9 (2.9-5.3)	11% (7.2-14%)	10% (5.5-15%)	2700 (1400-4200)	85000 (54000-120000)
Georgia	2016	Male	15 to 19	69% (58-79%)	31% (21-42%)	1.6 (1-2.4)	5.4% (3.6-7.5%)	7.4% (4.6-11%)	9.1 (5.3-14)	880 (570-1300)
Georgia	2016	Male	20 to 24	85% (78-90%)	15% (10-22%)	3.6 (2.4-5.1)	12% (9-17%)	16% (10-22%)	39 (23-58)	3300 (2200-4600)
Georgia	2016	Male	25 to 29	88% (82-92%)	12% (8.3-18%)	5.3 (3.7-7.5)	16% (12-21%)	21% (15-28%)	67 (43-94)	5100 (3500-6900)
Georgia	2016	Male	30 to 34	90% (85-93%)	10% (7.1-15%)	6.2 (4.3-8.8)	17% (13-21%)	22% (16-28%)	82 (54-120)	5600 (3900-7700)
Georgia	2016	Male	35 to 39	91% (87-94%)	9.4% (6.5-13%)	5.8 (4-8)	16% (12-20%)	21% (15-26%)	95 (60-140)	5700 (3900-8200)
Georgia	2016	Male	40 to 44	91% (87-94%)	9.3% (6.4-13%)	5.8 (4-8)	15% (11-19%)	20% (14-25%)	120 (76-180)	6300 (4200-9000)
Georgia	2016	Male	45 to 49	90% (85-93%)	9.7% (6.8-15%)	6.4 (4.6-8.7)	16% (12-20%)	20% (15-25%)	180 (120-260)	8000 (5500-11000)
Georgia	2016	Male	50 to 54	90% (86-93%)	9.5% (6.6-14%)	6.4 (4.5-8.8)	16% (11-20%)	19% (14-24%)	270 (170-380)	11000 (7000-15000)
Georgia	2016	Male	55 to 59	91% (87-94%)	8.8% (5.9-13%)	5.5 (3.7-7.6)	14% (8.7-18%)	16% (10-22%)	310 (180-450)	11000 (6500-15000)
Georgia	2016	Male	60 to 64	89% (84-93%)	11% (7.2-16%)	4.9 (3.2-6.8)	12% (6.4-17%)	14% (7.2-20%)	320 (160-480)	9100 (4800-14000)
Georgia	2016	Male	65 to 69	86% (79-90%)	14% (10-21%)	4.4 (3-6.2)	11% (5.6-16%)	13% (6.2-19%)	280 (130-440)	6700 (3400-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Georgia	2016	Male	70 to 74	83% (77-88%)	17% (12-23%)	3.9 (2.6-5.5)	9.3% (4-15%)	11% (4.2-17%)	260 (100-430)	4900 (2000-8000)
Georgia	2016	Male	75 to 79	81% (73-88%)	19% (12-27%)	3.4 (2.1-4.9)	7.4% (2.2-13%)	8.4% (2.3-15%)	330 (85-620)	4900 (1400-9100)
Georgia	2016	Male	80 to 84	80% (69-87%)	20% (13-31%)	2.8 (1.8-4)	5.5% (1-11%)	6.1% (0.91-13%)	240 (34-510)	2600 (520-5400)
Georgia	2016	Male	85 to 89	79% (68-86%)	21% (14-32%)	2.3 (1.4-3.5)	3.3% (-0.39-8.4%)	3.6% (-0.79-9.4%)	92 (-19-250)	740 (-85-1900)
Georgia	2016	Male	90 to 94	78% (68-86%)	22% (14-32%)	2 (1.2-3.2)	1.8% (-1.9-6.3%)	1.7% (-2.7-7.2%)	21 (-31-87)	130 (-140-460)
Georgia	2016	Male	95 plus	77% (63-88%)	23% (12-37%)	1.7 (0.85-2.9)	0.61% (-2.9-4.9%)	0.28% (-3.9-5.5%)	0.59 (-8-11)	5 (-24-42)
Kazakhstan	1990	Female	All Ages	57% (51-63%)	43% (37-49%)	2.3 (1.7-3)	5.6% (3.3-7.9%)	8% (3.9-13%)	5100 (2400-8400)	150000 (90000-210000)
Kazakhstan	1990	Female	15 to 19	47% (34-59%)	53% (41-66%)	1.8 (0.96-2.9)	4% (2.2-6.4%)	8.1% (4.3-14%)	41 (21-67)	3600 (2000-5700)
Kazakhstan	1990	Female	20 to 24	67% (54-78%)	33% (22-46%)	3 (1.8-4.8)	6.8% (4.3-9.9%)	13% (7.6-19%)	78 (45-120)	6500 (4100-9200)
Kazakhstan	1990	Female	25 to 29	74% (62-82%)	26% (18-38%)	3.7 (2.4-5.6)	7.6% (5.3-11%)	15% (9.9-20%)	110 (75-160)	8800 (6100-12000)
Kazakhstan	1990	Female	30 to 34	76% (66-85%)	24% (15-34%)	4.1 (2.6-5.9)	7.8% (5.5-10%)	15% (10-20%)	140 (93-190)	9700 (6700-13000)
Kazakhstan	1990	Female	35 to 39	77% (67-84%)	23% (16-33%)	4.1 (2.6-5.9)	8.1% (5.5-11%)	15% (10-20%)	160 (100-220)	9700 (6500-13000)
Kazakhstan	1990	Female	40 to 44	76% (65-84%)	24% (16-35%)	4.1 (2.6-5.9)	8.7% (5.7-12%)	15% (10-21%)	180 (110-250)	9500 (6300-13000)
Kazakhstan	1990	Female	45 to 49	74% (61-83%)	26% (17-39%)	3.9 (2.4-6)	9.6% (6.1-13%)	15% (9.9-22%)	200 (120-280)	8900 (5700-13000)
Kazakhstan	1990	Female	50 to 54	72% (60-82%)	28% (18-40%)	3.7 (2.2-5.8)	9.9% (5.7-14%)	15% (8.3-21%)	460 (260-690)	18000 (10000-27000)
Kazakhstan	1990	Female	55 to 59	69% (56-81%)	31% (19-44%)	3.4 (2.1-5.3)	9.5% (5-14%)	13% (7.1-20%)	390 (210-600)	13000 (7200-20000)
Kazakhstan	1990	Female	60 to 64	67% (52-80%)	33% (20-48%)	3.1 (1.8-5)	8.5% (3.3-14%)	11% (4.5-19%)	640 (250-1100)	18000 (7500-31000)
Kazakhstan	1990	Female	65 to 69	66% (50-79%)	34% (21-50%)	2.9 (1.7-4.5)	7.9% (3.1-13%)	10% (4-17%)	590 (220-1000)	14000 (5400-24000)
Kazakhstan	1990	Female	70 to 74	58% (43-71%)	42% (29-57%)	2.4 (1.3-4)	6.4% (1.3-13%)	8% (1.6-16%)	470 (93-950)	9000 (1800-18000)
Kazakhstan	1990	Female	75 to 79	47% (33-63%)	53% (37-67%)	2 (1-3.5)	5.1% (0.65-12%)	6.2% (0.7-14%)	570 (65-1300)	8400 (1000-19000)
Kazakhstan	1990	Female	80 to 84	42% (28-57%)	58% (43-72%)	2 (1-3.3)	5.1% (0.97-12%)	6% (1-14%)	510 (85-1200)	5600 (1100-12000)
Kazakhstan	1990	Female	85 to 89	41% (27-56%)	59% (44-73%)	2.1 (1.1-3.5)	5.1% (0.79-12%)	6.1% (0.78-14%)	370 (47-860)	2900 (430-6400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	1990	Female	90 to 94	42% (28-57%)	58% (43-72%)	2.3 (1.1-3.9)	5.1% (0.57-12%)	6.1% (0.45-15%)	160 (11-390)	850 (96-2000)
Kazakhstan	1990	Female	95 plus	45% (28-64%)	55% (36-72%)	2.5 (1.1-4.8)	5.6% (-0.14-14%)	6.5% (-0.5-17%)	45 (-3.4-120)	160 (-3.9-390)
Kazakhstan	1990	Male	All Ages	60% (54-68%)	40% (32-46%)	3 (2.2-3.9)	10% (7.3-13%)	12% (7.6-16%)	8300 (5300-12000)	340000 (240000-450000)
Kazakhstan	1990	Male	15 to 19	47% (34-60%)	53% (40-66%)	1.3 (0.66-2.2)	5.5% (3.2-8.6%)	7.4% (3.7-12%)	79 (39-130)	7000 (4000-11000)
Kazakhstan	1990	Male	20 to 24	74% (61-84%)	26% (16-39%)	3.5 (2.1-5.3)	14% (9.5-20%)	18% (11-26%)	290 (180-430)	22000 (15000-31000)
Kazakhstan	1990	Male	25 to 29	82% (72-88%)	18% (12-28%)	5.3 (3.4-7.9)	19% (13-25%)	24% (16-33%)	560 (360-770)	38000 (26000-51000)
Kazakhstan	1990	Male	30 to 34	86% (77-91%)	14% (8.6-23%)	6.1 (4-8.6)	20% (14-25%)	26% (18-33%)	700 (480-930)	44000 (31000-57000)
Kazakhstan	1990	Male	35 to 39	85% (77-91%)	15% (8.9-23%)	5.6 (3.5-8)	18% (12-23%)	23% (15-30%)	660 (410-900)	37000 (25000-50000)
Kazakhstan	1990	Male	40 to 44	85% (77-91%)	15% (9.2-23%)	5.2 (3.2-7.6)	17% (11-22%)	20% (13-27%)	610 (370-850)	31000 (19000-42000)
Kazakhstan	1990	Male	45 to 49	84% (75-90%)	16% (9.7-25%)	5.4 (3.4-8.3)	17% (10-22%)	20% (12-26%)	600 (360-830)	26000 (17000-36000)
Kazakhstan	1990	Male	50 to 54	83% (74-89%)	17% (11-26%)	5.5 (3.2-8.5)	16% (9-21%)	18% (10-24%)	1300 (690-1800)	49000 (28000-68000)
Kazakhstan	1990	Male	55 to 59	81% (70-88%)	19% (12-30%)	4.7 (2.9-7.1)	13% (7.4-18%)	15% (8.1-21%)	910 (470-1300)	30000 (16000-43000)
Kazakhstan	1990	Male	60 to 64	79% (67-88%)	21% (12-33%)	4.1 (2.3-6.4)	11% (5-16%)	12% (5.5-19%)	1100 (470-1700)	30000 (14000-48000)
Kazakhstan	1990	Male	65 to 69	78% (65-88%)	22% (12-35%)	3.8 (2.3-5.8)	9.6% (4.2-15%)	11% (4.6-18%)	610 (260-1000)	14000 (6200-23000)
Kazakhstan	1990	Male	70 to 74	74% (61-84%)	26% (16-39%)	3.3 (1.8-5.1)	7.8% (2.4-13%)	8.7% (2.5-15%)	400 (120-700)	7500 (2300-13000)
Kazakhstan	1990	Male	75 to 79	68% (52-81%)	32% (19-48%)	2.6 (1.4-4.3)	5.6% (0.77-11%)	6.2% (0.69-12%)	330 (38-670)	4900 (710-9800)
Kazakhstan	1990	Male	80 to 84	64% (47-78%)	36% (22-53%)	2.3 (1.2-3.6)	4% (-0.22-9.1%)	4.3% (-0.53-10%)	170 (-23-420)	2000 (-120-4600)
Kazakhstan	1990	Male	85 to 89	61% (45-75%)	39% (25-55%)	2.1 (1.1-3.4)	2.4% (-1.1-7.6%)	2.4% (-1.7-8.5%)	57 (-40-200)	490 (-220-1600)
Kazakhstan	1990	Male	90 to 94	60% (44-75%)	40% (25-56%)	1.9 (0.98-3.2)	1.1% (-2-5.9%)	0.82% (-2.9-6.4%)	7.3 (-26-59)	57 (-110-320)
Kazakhstan	1990	Male	95 plus	60% (40-77%)	40% (23-60%)	1.6 (0.71-3.2)	0.24% (-2.7-5.4%)	-0.24% (-3.8-5.9%)	-0.56 (-8.8-14)	2.2 (-25-51)
Kazakhstan	1995	Female	All Ages	60% (54-65%)	40% (35-46%)	2.2 (1.7-2.7)	5.9% (3.5-8.5%)	7.4% (3.1-12%)	5700 (2400-9600)	170000 (100000-250000)
Kazakhstan	1995	Female	15 to 19	51% (42-60%)	49% (40-58%)	1.7 (1-2.7)	4.3% (2.4-6.6%)	8.1% (4.4-12%)	49 (27-78)	4200 (2400-6500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	1995	Female	20 to 24	69% (60-76%)	31% (24-40%)	2.8 (1.7-4.3)	7.6% (4.8-11%)	13% (7.8-19%)	110 (62-150)	8300 (5300-12000)
Kazakhstan	1995	Female	25 to 29	75% (68-81%)	25% (19-32%)	3.4 (2.3-4.9)	8.9% (6.2-12%)	15% (10-20%)	140 (90-190)	9800 (6800-13000)
Kazakhstan	1995	Female	30 to 34	77% (71-83%)	23% (17-29%)	3.7 (2.5-5.1)	9% (6.5-12%)	15% (11-20%)	190 (130-250)	12000 (8400-16000)
Kazakhstan	1995	Female	35 to 39	77% (71-83%)	23% (17-29%)	3.7 (2.5-5.2)	8.8% (6.1-12%)	15% (10-19%)	230 (150-310)	13000 (9200-18000)
Kazakhstan	1995	Female	40 to 44	76% (70-82%)	24% (18-30%)	3.6 (2.5-5.1)	9.2% (6.2-13%)	14% (9.8-20%)	280 (190-390)	14000 (9600-20000)
Kazakhstan	1995	Female	45 to 49	76% (67-82%)	24% (18-33%)	3.6 (2.3-5.4)	9.7% (6.1-14%)	14% (8.9-20%)	310 (190-460)	14000 (8600-20000)
Kazakhstan	1995	Female	50 to 54	74% (64-82%)	26% (18-36%)	3.4 (2.1-5.1)	10% (5.5-15%)	14% (7.7-21%)	330 (180-500)	12000 (7000-19000)
Kazakhstan	1995	Female	55 to 59	71% (59-82%)	29% (18-41%)	3.1 (1.9-4.8)	9.3% (4.6-14%)	12% (6-19%)	680 (330-1100)	23000 (11000-36000)
Kazakhstan	1995	Female	60 to 64	70% (56-82%)	30% (18-44%)	2.9 (1.7-4.6)	7.8% (2.5-14%)	9.8% (3.2-17%)	480 (160-860)	14000 (4400-24000)
Kazakhstan	1995	Female	65 to 69	68% (54-80%)	32% (20-46%)	2.7 (1.6-4)	6.9% (2-13%)	8.4% (2.3-15%)	800 (220-1500)	19000 (5100-34000)
Kazakhstan	1995	Female	70 to 74	61% (47-73%)	39% (27-53%)	2.2 (1.2-3.6)	5.4% (0.26-12%)	6.5% (0.25-14%)	590 (21-1300)	11000 (470-24000)
Kazakhstan	1995	Female	75 to 79	51% (35-66%)	49% (34-65%)	1.9 (0.97-3.2)	4.3% (-0.26-11%)	5% (-0.44-13%)	410 (-32-1100)	6000 (-350-15000)
Kazakhstan	1995	Female	80 to 84	45% (31-60%)	55% (40-69%)	1.9 (1-3.1)	4.2% (0.058-10%)	4.8% (-0.16-12%)	480 (-15-1200)	5200 (72-13000)
Kazakhstan	1995	Female	85 to 89	44% (30-58%)	56% (42-70%)	2 (1.1-3.2)	4.2% (-0.13-11%)	4.9% (-0.4-12%)	380 (-30-1000)	3000 (-85-7400)
Kazakhstan	1995	Female	90 to 94	45% (31-59%)	55% (41-69%)	2.1 (1.1-3.6)	4.3% (-0.43-11%)	5.1% (-0.79-14%)	170 (-27-460)	920 (-90-2400)
Kazakhstan	1995	Female	95 plus	47% (30-65%)	53% (35-70%)	2.4 (1.1-4.3)	4.8% (-0.91-13%)	5.6% (-1.3-16%)	61 (-15-170)	210 (-42-580)
Kazakhstan	1995	Male	All Ages	67% (61-73%)	33% (27-39%)	2.9 (2.3-3.6)	12% (8.5-15%)	12% (7.9-16%)	11000 (7000-15000)	470000 (340000-600000)
Kazakhstan	1995	Male	15 to 19	50% (40-60%)	50% (40-60%)	1.2 (0.67-2)	5.8% (3.4-9.1%)	7.6% (3.9-12%)	97 (49-160)	8200 (4400-13000)
Kazakhstan	1995	Male	20 to 24	79% (71-85%)	21% (15-29%)	3.4 (2.2-5)	16% (11-22%)	20% (13-28%)	450 (290-640)	33000 (22000-45000)
Kazakhstan	1995	Male	25 to 29	87% (82-90%)	13% (9.6-18%)	5.1 (3.4-7.3)	22% (16-28%)	26% (19-34%)	690 (470-910)	45000 (32000-59000)
Kazakhstan	1995	Male	30 to 34	89% (85-92%)	11% (7.5-15%)	5.7 (3.9-7.9)	22% (16-28%)	27% (20-34%)	960 (670-1200)	58000 (42000-74000)
Kazakhstan	1995	Male	35 to 39	89% (84-93%)	11% (7.4-16%)	5.3 (3.6-7.3)	20% (14-26%)	24% (16-31%)	1100 (720-1500)	60000 (40000-80000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	1995	Male	40 to 44	88% (84-92%)	12% (7.9-16%)	5.1 (3.3-7.1)	19% (12-24%)	21% (14-29%)	1100 (710-1600)	56000 (36000-76000)
Kazakhstan	1995	Male	45 to 49	88% (82-92%)	12% (8.1-18%)	5.3 (3.5-7.7)	18% (11-24%)	20% (12-27%)	1100 (680-1600)	48000 (30000-67000)
Kazakhstan	1995	Male	50 to 54	87% (82-91%)	13% (9.2-18%)	5.3 (3.3-7.8)	16% (9.1-23%)	18% (9.8-25%)	950 (510-1400)	36000 (19000-50000)
Kazakhstan	1995	Male	55 to 59	85% (78-90%)	15% (10-22%)	4.6 (2.9-6.6)	13% (6.7-19%)	14% (7.1-21%)	1700 (810-2600)	55000 (28000-84000)
Kazakhstan	1995	Male	60 to 64	81% (72-88%)	19% (12-28%)	3.8 (2.3-5.8)	10% (4.2-16%)	11% (4.4-18%)	980 (400-1600)	27000 (12000-45000)
Kazakhstan	1995	Male	65 to 69	79% (68-88%)	21% (12-32%)	3.4 (2.2-5)	8.3% (3.3-14%)	9% (3.4-15%)	1100 (410-1900)	26000 (10000-43000)
Kazakhstan	1995	Male	70 to 74	76% (63-85%)	24% (15-37%)	3 (1.7-4.4)	6.5% (1.6-12%)	7.1% (1.5-14%)	480 (100-940)	9100 (2300-17000)
Kazakhstan	1995	Male	75 to 79	70% (55-82%)	30% (18-45%)	2.4 (1.3-3.8)	4.3% (0.091-9.6%)	4.6% (-0.18-11%)	220 (-7.7-510)	3300 (66-7300)
Kazakhstan	1995	Male	80 to 84	66% (50-79%)	34% (21-50%)	2.1 (1.2-3.2)	2.6% (-0.97-7.5%)	2.7% (-1.4-8.2%)	120 (-61-360)	1400 (-490-3800)
Kazakhstan	1995	Male	85 to 89	63% (47-77%)	37% (23-53%)	1.9 (1.1-3)	1.2% (-2-5.6%)	1% (-2.8-6.1%)	28 (-78-170)	290 (-480-1300)
Kazakhstan	1995	Male	90 to 94	62% (47-76%)	38% (24-53%)	1.7 (0.93-2.8)	0.19% (-2.6-4.4%)	-0.21% (-3.6-4.7%)	-2 (-36-46)	12 (-160-260)
Kazakhstan	1995	Male	95 plus	62% (42-79%)	38% (21-58%)	1.5 (0.69-2.8)	-0.4% (-3.2-3.9%)	-1% (-4.4-4.2%)	-2.6 (-11-11)	-4.2 (-33-40)
Kazakhstan	2000	Female	All Ages	60% (55-65%)	40% (35-45%)	2.3 (1.8-2.8)	6.3% (4.1-8.8%)	7.4% (3.5-12%)	5300 (2500-8800)	170000 (110000-240000)
Kazakhstan	2000	Female	15 to 19	50% (41-60%)	50% (40-59%)	1.7 (1.1-2.6)	4.5% (2.6-6.8%)	8.7% (4.9-14%)	50 (28-78)	4200 (2500-6300)
Kazakhstan	2000	Female	20 to 24	69% (60-76%)	31% (24-40%)	2.9 (1.8-4.3)	8.4% (5.5-12%)	15% (9.4-21%)	120 (73-170)	9100 (6000-13000)
Kazakhstan	2000	Female	25 to 29	76% (69-81%)	24% (19-31%)	3.6 (2.4-5)	10% (7.3-14%)	17% (12-23%)	170 (120-230)	12000 (8400-16000)
Kazakhstan	2000	Female	30 to 34	78% (72-84%)	22% (16-28%)	3.9 (2.7-5.3)	10% (7.7-14%)	17% (13-23%)	190 (140-260)	12000 (8900-16000)
Kazakhstan	2000	Female	35 to 39	78% (72-83%)	22% (17-28%)	3.8 (2.6-5.1)	9.9% (7.1-13%)	16% (12-21%)	250 (170-340)	14000 (10000-19000)
Kazakhstan	2000	Female	40 to 44	77% (71-82%)	23% (18-29%)	3.7 (2.6-5.1)	10% (7.2-13%)	16% (11-21%)	320 (220-440)	16000 (11000-22000)
Kazakhstan	2000	Female	45 to 49	75% (67-82%)	25% (18-33%)	3.6 (2.4-5.3)	10% (6.5-14%)	15% (9.8-21%)	400 (250-560)	18000 (11000-25000)
Kazakhstan	2000	Female	50 to 54	71% (62-79%)	29% (21-38%)	3.2 (2.1-4.7)	9.8% (5.8-14%)	14% (8-20%)	390 (220-580)	15000 (8800-22000)
Kazakhstan	2000	Female	55 to 59	70% (59-80%)	30% (20-41%)	3.1 (1.9-4.5)	9.4% (4.9-14%)	12% (6.4-19%)	370 (190-560)	12000 (6300-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	2000	Female	60 to 64	69% (57-80%)	31% (20-43%)	2.9 (1.7-4.4)	7.8% (2.6-14%)	9.9% (3.3-17%)	700 (230-1200)	20000 (6600-35000)
Kazakhstan	2000	Female	65 to 69	65% (53-77%)	35% (23-47%)	2.5 (1.6-3.7)	6.6% (1.8-12%)	8.1% (2.1-15%)	470 (120-880)	11000 (3000-20000)
Kazakhstan	2000	Female	70 to 74	54% (42-65%)	46% (35-58%)	1.9 (1.1-3)	4.8% (0.64-10%)	5.7% (0.65-12%)	620 (72-1300)	12000 (1600-25000)
Kazakhstan	2000	Female	75 to 79	42% (29-58%)	58% (42-71%)	1.5 (0.81-2.7)	3.6% (0.088-9%)	4.2% (0.0076-11%)	380 (0.74-980)	5600 (140-14000)
Kazakhstan	2000	Female	80 to 84	42% (28-56%)	58% (44-72%)	1.7 (0.97-2.8)	3.8% (0.15-9.2%)	4.4% (-0.08-11%)	330 (-6-800)	3600 (150-8500)
Kazakhstan	2000	Female	85 to 89	43% (30-58%)	57% (42-70%)	2 (1.1-3.2)	4.1% (-0.058-10%)	4.7% (-0.26-12%)	300 (-17-760)	2300 (-34-5700)
Kazakhstan	2000	Female	90 to 94	44% (30-59%)	56% (41-70%)	2.1 (1.2-3.6)	4.3% (-0.31-11%)	4.9% (-0.85-13%)	180 (-30-480)	950 (-70-2500)
Kazakhstan	2000	Female	95 plus	46% (29-65%)	54% (35-71%)	2.4 (1.1-4.4)	4.7% (-0.87-13%)	5.5% (-1.5-16%)	64 (-17-190)	220 (-42-620)
Kazakhstan	2000	Male	All Ages	68% (62-73%)	32% (27-38%)	3.1 (2.5-3.7)	13% (10-16%)	13% (9.3-17%)	12000 (7900-16000)	500000 (370000-640000)
Kazakhstan	2000	Male	15 to 19	52% (43-62%)	48% (38-57%)	1.3 (0.74-2.1)	6.2% (3.6-9.6%)	8.3% (4.3-13%)	100 (54-170)	8700 (4900-14000)
Kazakhstan	2000	Male	20 to 24	80% (72-85%)	20% (15-28%)	3.6 (2.3-5.3)	17% (12-23%)	21% (14-29%)	480 (310-670)	34000 (23000-47000)
Kazakhstan	2000	Male	25 to 29	87% (82-91%)	13% (9.4-18%)	5.3 (3.6-7.4)	24% (17-30%)	28% (21-36%)	820 (580-1100)	53000 (38000-69000)
Kazakhstan	2000	Male	30 to 34	89% (85-93%)	11% (7.3-15%)	6 (4.2-8.1)	24% (18-30%)	29% (21-36%)	930 (660-1200)	56000 (40000-71000)
Kazakhstan	2000	Male	35 to 39	89% (85-92%)	11% (7.6-15%)	5.5 (3.8-7.4)	22% (16-28%)	26% (18-33%)	1100 (790-1500)	62000 (44000-81000)
Kazakhstan	2000	Male	40 to 44	88% (84-92%)	12% (8.2-16%)	5.2 (3.5-7.1)	20% (14-26%)	23% (15-30%)	1300 (860-1800)	64000 (43000-86000)
Kazakhstan	2000	Male	45 to 49	87% (82-92%)	13% (8.4-18%)	5.4 (3.7-7.7)	19% (13-25%)	22% (14-28%)	1500 (910-2000)	62000 (40000-85000)
Kazakhstan	2000	Male	50 to 54	87% (82-91%)	13% (9.3-18%)	5.5 (3.5-8)	18% (10-24%)	20% (11-27%)	1300 (700-1800)	49000 (28000-69000)
Kazakhstan	2000	Male	55 to 59	84% (78-89%)	16% (11-22%)	4.7 (3.1-6.8)	14% (7.6-20%)	16% (8-23%)	900 (450-1300)	29000 (16000-44000)
Kazakhstan	2000	Male	60 to 64	80% (72-88%)	20% (12-28%)	3.8 (2.3-5.8)	10% (4.5-17%)	11% (4.7-19%)	1400 (580-2300)	39000 (17000-64000)
Kazakhstan	2000	Male	65 to 69	79% (69-87%)	21% (13-31%)	3.5 (2.4-5)	8.6% (3.4-14%)	9.4% (3.5-16%)	790 (300-1300)	18000 (7200-30000)
Kazakhstan	2000	Male	70 to 74	74% (63-83%)	26% (17-37%)	2.9 (1.8-4.3)	6.5% (1.7-12%)	7.1% (1.7-13%)	740 (180-1400)	14000 (3600-26000)
Kazakhstan	2000	Male	75 to 79	66% (50-78%)	34% (22-50%)	2.2 (1.2-3.5)	4.1% (0.24-9.3%)	4.5% (0.0021-10%)	230 (0.11-550)	3500 (200-7900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	2000	Male	80 to 84	64% (48-78%)	36% (22-52%)	2 (1.2-3.1)	2.6% (-0.9-7.1%)	2.6% (-1.4-7.9%)	82 (-45-250)	970 (-330-2800)
Kazakhstan	2000	Male	85 to 89	64% (48-76%)	36% (24-52%)	2 (1.2-3)	1.1% (-2.2-5.5%)	0.9% (-3-6%)	18 (-61-120)	200 (-390-960)
Kazakhstan	2000	Male	90 to 94	63% (47-76%)	37% (24-53%)	1.8 (0.98-2.8)	0.063% (-3-4.2%)	-0.38% (-4.1-4.6%)	-3.9 (-42-46)	3.8 (-180-260)
Kazakhstan	2000	Male	95 plus	62% (42-78%)	38% (22-58%)	1.5 (0.73-2.8)	-0.53% (-3.5-3.8%)	-1.2% (-4.8-3.9%)	-3.1 (-13-10)	-5.5 (-37-40)
Kazakhstan	2005	Female	All Ages	61% (57-66%)	39% (34-43%)	2.5 (2.1-3.1)	7.2% (5-9.8%)	8.5% (4.5-13%)	6300 (3300-10000)	200000 (140000-270000)
Kazakhstan	2005	Female	15 to 19	48% (38-59%)	52% (41-62%)	1.8 (1.1-2.7)	4.7% (2.7-7.4%)	9.6% (5.4-15%)	59 (33-93)	5000 (2900-7500)
Kazakhstan	2005	Female	20 to 24	69% (59-77%)	31% (23-41%)	3.1 (2-4.5)	8.9% (6-13%)	16% (10-22%)	130 (83-190)	10000 (6900-14000)
Kazakhstan	2005	Female	25 to 29	77% (70-83%)	23% (17-30%)	3.9 (2.6-5.3)	11% (7.9-14%)	18% (13-24%)	190 (130-260)	13000 (9600-18000)
Kazakhstan	2005	Female	30 to 34	79% (73-85%)	21% (15-27%)	4.2 (3-5.7)	12% (8.6-15%)	19% (14-24%)	250 (180-330)	16000 (11000-20000)
Kazakhstan	2005	Female	35 to 39	78% (72-84%)	22% (16-28%)	4.1 (2.8-5.7)	11% (8.2-15%)	18% (13-23%)	280 (190-380)	16000 (11000-21000)
Kazakhstan	2005	Female	40 to 44	77% (70-83%)	23% (17-30%)	4 (2.7-5.6)	12% (8.2-15%)	18% (13-23%)	390 (270-530)	20000 (14000-27000)
Kazakhstan	2005	Female	45 to 49	75% (66-82%)	25% (18-34%)	3.9 (2.6-5.6)	11% (7.8-15%)	17% (11-23%)	490 (330-680)	22000 (15000-30000)
Kazakhstan	2005	Female	50 to 54	71% (62-79%)	29% (21-38%)	3.5 (2.3-5.3)	11% (7-16%)	15% (9.9-22%)	550 (350-800)	21000 (14000-31000)
Kazakhstan	2005	Female	55 to 59	71% (60-81%)	29% (19-40%)	3.4 (2.1-5)	10% (5.9-15%)	14% (7.9-21%)	540 (300-800)	18000 (10000-27000)
Kazakhstan	2005	Female	60 to 64	69% (57-80%)	31% (20-43%)	3.1 (1.9-4.8)	8.9% (3.7-15%)	11% (4.7-19%)	460 (190-770)	13000 (5500-22000)
Kazakhstan	2005	Female	65 to 69	66% (53-77%)	34% (23-47%)	2.7 (1.7-4.1)	7.5% (2.7-13%)	9.3% (3.5-16%)	810 (300-1400)	19000 (7200-33000)
Kazakhstan	2005	Female	70 to 74	55% (42-67%)	45% (33-58%)	2.1 (1.2-3.4)	5.5% (0.9-11%)	6.6% (1.1-13%)	510 (82-1100)	9700 (1600-20000)
Kazakhstan	2005	Female	75 to 79	44% (30-59%)	56% (41-70%)	1.7 (0.92-2.9)	4.2% (0.31-10%)	4.9% (0.26-12%)	590 (33-1400)	8600 (650-21000)
Kazakhstan	2005	Female	80 to 84	43% (29-58%)	57% (42-71%)	1.9 (1.1-3.1)	4.4% (0.33-10%)	5.1% (0.24-12%)	470 (24-1200)	5200 (410-12000)
Kazakhstan	2005	Female	85 to 89	44% (30-59%)	56% (41-70%)	2.2 (1.2-3.5)	4.8% (0.37-12%)	5.5% (0.14-14%)	270 (7.2-710)	2100 (170-5200)
Kazakhstan	2005	Female	90 to 94	45% (32-60%)	55% (40-68%)	2.4 (1.3-4.1)	5% (-0.22-13%)	5.8% (-0.61-16%)	170 (-19-470)	900 (-41-2400)
Kazakhstan	2005	Female	95 plus	47% (30-65%)	53% (35-70%)	2.6 (1.2-4.8)	5.6% (-0.95-15%)	6.4% (-1.4-18%)	78 (-18-210)	270 (-46-710)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	2005	Male	All Ages	69% (64-75%)	31% (25-36%)	3.4 (2.8-4.1)	14% (11-18%)	15% (11-19%)	14000 (9700-18000)	580000 (450000-730000)
Kazakhstan	2005	Male	15 to 19	54% (44-64%)	46% (36-56%)	1.5 (0.84-2.3)	6.8% (4-10%)	9.3% (4.9-15%)	130 (65-200)	10000 (5900-16000)
Kazakhstan	2005	Male	20 to 24	80% (72-86%)	20% (14-28%)	3.8 (2.5-5.5)	18% (12-24%)	21% (14-29%)	530 (340-760)	38000 (26000-53000)
Kazakhstan	2005	Male	25 to 29	87% (83-91%)	13% (9.4-17%)	5.7 (3.8-7.8)	24% (18-31%)	28% (20-36%)	950 (670-1300)	62000 (45000-81000)
Kazakhstan	2005	Male	30 to 34	90% (86-93%)	10% (7.1-14%)	6.4 (4.5-8.7)	25% (19-31%)	29% (22-36%)	1200 (850-1500)	70000 (51000-89000)
Kazakhstan	2005	Male	35 to 39	89% (84-93%)	11% (7.5-16%)	5.8 (4-8)	23% (17-29%)	27% (19-34%)	1200 (820-1600)	65000 (46000-85000)
Kazakhstan	2005	Male	40 to 44	87% (82-91%)	13% (8.6-18%)	5.4 (3.7-7.3)	21% (15-27%)	24% (17-31%)	1500 (1000-2100)	74000 (50000-98000)
Kazakhstan	2005	Male	45 to 49	87% (81-91%)	13% (8.5-19%)	5.6 (3.9-8)	21% (14-26%)	23% (16-29%)	1800 (1200-2400)	77000 (51000-100000)
Kazakhstan	2005	Male	50 to 54	87% (81-91%)	13% (9.4-19%)	5.8 (3.7-8.3)	19% (12-25%)	21% (13-27%)	1800 (1100-2400)	68000 (41000-89000)
Kazakhstan	2005	Male	55 to 59	83% (76-89%)	17% (11-24%)	4.9 (3.2-7.1)	15% (8.7-21%)	17% (9.1-24%)	1300 (720-1900)	44000 (25000-62000)
Kazakhstan	2005	Male	60 to 64	81% (72-88%)	19% (12-28%)	4.1 (2.5-6.2)	12% (5.4-18%)	13% (5.6-20%)	860 (380-1400)	24000 (11000-38000)
Kazakhstan	2005	Male	65 to 69	80% (70-88%)	20% (12-30%)	3.8 (2.6-5.3)	9.6% (4.1-15%)	10% (4.3-17%)	1300 (540-2100)	30000 (13000-49000)
Kazakhstan	2005	Male	70 to 74	75% (63-84%)	25% (16-37%)	3.2 (2-4.7)	7.3% (2.3-13%)	7.9% (2.3-14%)	660 (180-1200)	12000 (3800-22000)
Kazakhstan	2005	Male	75 to 79	66% (50-80%)	34% (20-50%)	2.4 (1.3-3.8)	4.7% (0.55-10%)	5.1% (0.37-11%)	430 (32-970)	6500 (730-14000)
Kazakhstan	2005	Male	80 to 84	65% (48-78%)	35% (22-52%)	2.2 (1.3-3.4)	3.1% (-0.55-8%)	3.2% (-1.1-8.8%)	130 (-42-350)	1500 (-260-3900)
Kazakhstan	2005	Male	85 to 89	64% (49-77%)	36% (23-51%)	2.2 (1.3-3.3)	1.5% (-2.1-6.4%)	1.3% (-2.9-6.9%)	19 (-43-100)	190 (-270-800)
Kazakhstan	2005	Male	90 to 94	63% (48-77%)	37% (23-52%)	1.9 (1.1-3.1)	0.28% (-3-5.1%)	-0.13% (-4.1-5.5%)	-0.93 (-29-39)	12 (-130-220)
Kazakhstan	2005	Male	95 plus	62% (44-79%)	38% (21-56%)	1.7 (0.77-3.1)	-0.37% (-3.8-4.9%)	-0.97% (-5-5.2%)	-2.3 (-12-13)	-3.4 (-36-47)
Kazakhstan	2010	Female	All Ages	64% (59-69%)	36% (31-41%)	2.7 (2.2-3.4)	7.5% (5.1-10%)	9.5% (4.9-15%)	6200 (3300-9900)	190000 (130000-260000)
Kazakhstan	2010	Female	15 to 19	52% (39-64%)	48% (36-61%)	2 (1.2-3.1)	4.7% (2.7-7.3%)	11% (5.9-17%)	48 (25-76)	4200 (2400-6400)
Kazakhstan	2010	Female	20 to 24	71% (59-80%)	29% (20-41%)	3.3 (2.1-4.9)	7.8% (5.2-11%)	16% (10-22%)	110 (70-160)	9100 (6000-13000)
Kazakhstan	2010	Female	25 to 29	76% (66-83%)	24% (17-34%)	4 (2.6-5.6)	9.5% (6.9-13%)	18% (13-23%)	150 (100-200)	11000 (7800-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	2010	Female	30 to 34	79% (69-85%)	21% (15-31%)	4.3 (2.9-6)	11% (7.8-13%)	19% (14-24%)	200 (140-260)	13000 (9600-17000)
Kazakhstan	2010	Female	35 to 39	79% (70-85%)	21% (15-30%)	4.2 (2.8-6.1)	11% (8-14%)	19% (14-24%)	250 (180-330)	15000 (11000-19000)
Kazakhstan	2010	Female	40 to 44	78% (69-85%)	22% (15-31%)	4.2 (2.8-6)	11% (8-15%)	19% (14-24%)	310 (220-400)	16000 (12000-21000)
Kazakhstan	2010	Female	45 to 49	76% (65-84%)	24% (16-35%)	4.2 (2.7-6.1)	11% (8-15%)	18% (13-24%)	440 (310-600)	20000 (14000-27000)
Kazakhstan	2010	Female	50 to 54	75% (64-83%)	25% (17-36%)	3.9 (2.5-5.9)	11% (7.6-16%)	17% (11-23%)	530 (350-740)	21000 (14000-29000)
Kazakhstan	2010	Female	55 to 59	72% (60-83%)	28% (17-40%)	3.6 (2.2-5.5)	11% (6.3-15%)	15% (8.9-22%)	580 (350-860)	20000 (12000-29000)
Kazakhstan	2010	Female	60 to 64	71% (57-83%)	29% (17-43%)	3.3 (2-5.2)	9.3% (4.3-15%)	12% (5.8-20%)	520 (240-850)	15000 (7000-25000)
Kazakhstan	2010	Female	65 to 69	70% (56-82%)	30% (18-44%)	3.1 (1.9-4.7)	8.3% (3.4-14%)	11% (4.3-18%)	500 (200-840)	12000 (4900-20000)
Kazakhstan	2010	Female	70 to 74	63% (49-75%)	37% (25-51%)	2.6 (1.5-4.3)	6.6% (1.3-13%)	8.3% (1.6-16%)	790 (140-1600)	15000 (2800-30000)
Kazakhstan	2010	Female	75 to 79	53% (38-68%)	47% (32-62%)	2.2 (1.2-3.9)	5.4% (0.58-12%)	6.4% (0.55-14%)	480 (39-1100)	7100 (740-16000)
Kazakhstan	2010	Female	80 to 84	48% (33-63%)	52% (37-67%)	2.3 (1.2-3.6)	5.5% (0.68-12%)	6.2% (0.53-14%)	610 (50-1400)	6600 (800-15000)
Kazakhstan	2010	Female	85 to 89	46% (32-61%)	54% (39-68%)	2.4 (1.3-3.8)	5.3% (0.66-12%)	6.2% (0.41-15%)	420 (27-1000)	3300 (400-7500)
Kazakhstan	2010	Female	90 to 94	47% (33-61%)	53% (39-67%)	2.5 (1.4-4.4)	5.4% (0.26-13%)	6.4% (-0.088-16%)	150 (-2.1-390)	830 (41-2000)
Kazakhstan	2010	Female	95 plus	49% (32-67%)	51% (33-68%)	2.8 (1.3-5)	5.9% (-0.59-15%)	6.8% (-1.2-18%)	64 (-11-170)	220 (-21-560)
Kazakhstan	2010	Male	All Ages	68% (61-73%)	32% (27-39%)	3.4 (2.7-4.2)	14% (11-17%)	15% (10-19%)	12000 (8100-16000)	490000 (380000-620000)
Kazakhstan	2010	Male	15 to 19	52% (39-64%)	48% (36-61%)	1.4 (0.78-2.4)	6.2% (3.7-9.4%)	9.1% (4.7-15%)	81 (41-130)	7100 (4200-11000)
Kazakhstan	2010	Male	20 to 24	77% (65-85%)	23% (15-35%)	3.6 (2.4-5.5)	16% (11-21%)	20% (13-29%)	390 (240-570)	30000 (20000-42000)
Kazakhstan	2010	Male	25 to 29	84% (75-89%)	16% (11-25%)	5.5 (3.6-7.9)	22% (16-28%)	27% (19-35%)	690 (480-920)	47000 (34000-61000)
Kazakhstan	2010	Male	30 to 34	87% (80-92%)	13% (7.8-20%)	6.2 (4.2-8.8)	24% (18-29%)	28% (21-35%)	930 (660-1200)	57000 (42000-72000)
Kazakhstan	2010	Male	35 to 39	87% (80-92%)	13% (8.2-20%)	5.7 (3.8-8)	22% (17-28%)	27% (19-33%)	1000 (710-1300)	56000 (41000-72000)
Kazakhstan	2010	Male	40 to 44	86% (80-91%)	14% (8.5-20%)	5.4 (3.5-7.4)	21% (15-26%)	25% (18-31%)	1100 (770-1400)	54000 (39000-70000)
Kazakhstan	2010	Male	45 to 49	86% (78-91%)	14% (8.9-22%)	5.6 (3.8-8.3)	20% (15-25%)	24% (17-30%)	1500 (1000-2000)	65000 (48000-85000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	2010	Male	50 to 54	85% (77-90%)	15% (10-23%)	5.7 (3.6-8.5)	19% (12-24%)	21% (14-28%)	1600 (1000-2100)	60000 (40000-80000)
Kazakhstan	2010	Male	55 to 59	82% (73-89%)	18% (11-27%)	4.9 (3.1-7.2)	15% (9.5-21%)	17% (10-24%)	1300 (800-1900)	45000 (28000-63000)
Kazakhstan	2010	Male	60 to 64	81% (71-89%)	19% (11-29%)	4.2 (2.5-6.4)	12% (6.2-18%)	13% (6.5-20%)	970 (470-1500)	28000 (14000-42000)
Kazakhstan	2010	Male	65 to 69	80% (69-89%)	20% (11-31%)	3.9 (2.5-5.6)	10% (5-16%)	11% (5.3-18%)	740 (330-1200)	17000 (8100-27000)
Kazakhstan	2010	Male	70 to 74	77% (65-86%)	23% (14-35%)	3.4 (2.1-5.1)	8% (2.9-13%)	8.9% (3-15%)	940 (300-1600)	18000 (6300-30000)
Kazakhstan	2010	Male	75 to 79	71% (57-83%)	29% (17-43%)	2.8 (1.6-4.4)	5.7% (0.96-11%)	6.2% (0.75-13%)	360 (44-740)	5400 (890-11000)
Kazakhstan	2010	Male	80 to 84	68% (52-80%)	32% (20-48%)	2.4 (1.4-3.7)	3.7% (-0.42-9.1%)	3.9% (-0.92-10%)	200 (-47-510)	2200 (-250-5500)
Kazakhstan	2010	Male	85 to 89	65% (49-78%)	35% (22-51%)	2.2 (1.3-3.5)	2% (-1.6-7%)	1.8% (-2.5-7.8%)	42 (-57-180)	410 (-330-1400)
Kazakhstan	2010	Male	90 to 94	64% (50-78%)	36% (22-50%)	2 (1.1-3.3)	0.72% (-2.4-5.4%)	0.27% (-3.6-5.9%)	1.5 (-20-33)	25 (-83-190)
Kazakhstan	2010	Male	95 plus	63% (44-80%)	37% (20-56%)	1.7 (0.81-3.2)	0.057% (-3.1-4.9%)	-0.65% (-4.6-5.2%)	-1.1 (-7.2-8.6)	0.4 (-20-34)
Kazakhstan	2016	Female	All Ages	64% (58-69%)	36% (31-42%)	2.7 (2-3.5)	7.1% (4.4-10%)	9.4% (4.4-15%)	5700 (2400-9800)	170000 (100000-250000)
Kazakhstan	2016	Female	15 to 19	52% (40-64%)	48% (36-60%)	2 (1.1-3.3)	4.1% (2.3-6.7%)	11% (5.5-18%)	26 (13-45)	2400 (1300-3900)
Kazakhstan	2016	Female	20 to 24	71% (59-80%)	29% (20-41%)	3.3 (1.9-5.2)	6.7% (4.3-9.8%)	15% (9.5-23%)	64 (37-99)	5800 (3700-8400)
Kazakhstan	2016	Female	25 to 29	76% (67-83%)	24% (17-33%)	3.9 (2.5-5.8)	8.2% (5.8-11%)	17% (12-24%)	120 (78-180)	9900 (6900-14000)
Kazakhstan	2016	Female	30 to 34	79% (69-85%)	21% (15-31%)	4.3 (2.7-6.3)	9.4% (6.8-13%)	19% (14-25%)	170 (110-240)	12000 (8100-16000)
Kazakhstan	2016	Female	35 to 39	79% (70-86%)	21% (14-30%)	4.2 (2.6-6.3)	10% (7-13%)	19% (14-24%)	200 (130-290)	12000 (8400-17000)
Kazakhstan	2016	Female	40 to 44	78% (70-85%)	22% (15-30%)	4.2 (2.6-6.3)	10% (6.9-14%)	19% (13-24%)	260 (170-380)	14000 (9500-20000)
Kazakhstan	2016	Female	45 to 49	77% (66-84%)	23% (16-34%)	4.1 (2.5-6.4)	10% (6.9-14%)	18% (12-24%)	310 (190-450)	14000 (9500-21000)
Kazakhstan	2016	Female	50 to 54	75% (64-83%)	25% (17-36%)	3.9 (2.4-6.1)	10% (6.7-15%)	17% (11-23%)	480 (300-720)	20000 (12000-29000)
Kazakhstan	2016	Female	55 to 59	73% (60-83%)	27% (17-40%)	3.6 (2.1-5.8)	9.9% (5.6-15%)	15% (8.7-22%)	560 (300-850)	19000 (11000-29000)
Kazakhstan	2016	Female	60 to 64	71% (57-82%)	29% (18-43%)	3.4 (1.9-5.5)	8.8% (3.8-15%)	12% (5.4-20%)	630 (270-1100)	19000 (8000-31000)
Kazakhstan	2016	Female	65 to 69	70% (55-82%)	30% (18-45%)	3.1 (1.8-4.8)	7.9% (2.7-13%)	11% (3.7-18%)	480 (160-860)	12000 (3900-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	2016	Female	70 to 74	63% (49-75%)	37% (25-51%)	2.6 (1.4-4.3)	6.5% (1.3-13%)	8.3% (1.6-16%)	470 (81-980)	9100 (1700-19000)
Kazakhstan	2016	Female	75 to 79	53% (37-68%)	47% (32-63%)	2.2 (1.2-4.1)	5.3% (0.55-12%)	6.5% (0.6-15%)	670 (56-1600)	10000 (1100-23000)
Kazakhstan	2016	Female	80 to 84	48% (33-63%)	52% (37-67%)	2.3 (1.2-3.8)	5.5% (0.42-13%)	6.4% (0.29-15%)	670 (30-1600)	7400 (550-17000)
Kazakhstan	2016	Female	85 to 89	47% (33-61%)	53% (39-67%)	2.4 (1.3-4.1)	5.5% (0.33-13%)	6.5% (0.095-16%)	340 (5.6-820)	2600 (170-6200)
Kazakhstan	2016	Female	90 to 94	48% (34-62%)	52% (38-66%)	2.6 (1.3-4.6)	5.6% (-0.093-14%)	6.6% (-0.47-17%)	200 (-16-520)	1100 (-18-2700)
Kazakhstan	2016	Female	95 plus	50% (33-68%)	50% (32-67%)	2.8 (1.4-5.2)	6% (-0.54-15%)	7.1% (-1.1-19%)	58 (-9-160)	200 (-18-520)
Kazakhstan	2016	Male	All Ages	67% (61-73%)	33% (27-39%)	3.3 (2.4-4.5)	14% (10-18%)	14% (9.3-19%)	11000 (6400-16000)	440000 (300000-600000)
Kazakhstan	2016	Male	15 to 19	52% (39-65%)	48% (35-61%)	1.4 (0.75-2.5)	5.9% (3.6-9.3%)	9.4% (5-16%)	45 (22-77)	4300 (2500-6600)
Kazakhstan	2016	Male	20 to 24	77% (65-85%)	23% (15-35%)	3.6 (2.2-5.5)	15% (10-20%)	20% (13-29%)	250 (140-380)	20000 (12000-29000)
Kazakhstan	2016	Male	25 to 29	84% (75-89%)	16% (11-25%)	5.4 (3.4-8.2)	21% (15-27%)	27% (19-35%)	580 (360-840)	42000 (29000-57000)
Kazakhstan	2016	Male	30 to 34	87% (80-92%)	13% (7.9-20%)	6.1 (3.9-9.1)	23% (17-28%)	28% (21-36%)	800 (540-1100)	51000 (36000-69000)
Kazakhstan	2016	Male	35 to 39	87% (80-92%)	13% (8.2-20%)	5.7 (3.6-8.4)	22% (16-27%)	27% (20-34%)	840 (550-1200)	48000 (33000-67000)
Kazakhstan	2016	Male	40 to 44	86% (80-91%)	14% (8.6-20%)	5.4 (3.3-7.8)	21% (15-26%)	26% (18-32%)	930 (590-1300)	47000 (32000-66000)
Kazakhstan	2016	Male	45 to 49	86% (78-91%)	14% (8.9-22%)	5.6 (3.5-8.8)	20% (14-25%)	24% (17-31%)	1000 (650-1500)	46000 (31000-65000)
Kazakhstan	2016	Male	50 to 54	85% (77-90%)	15% (10-23%)	5.6 (3.3-8.8)	19% (12-24%)	22% (14-29%)	1400 (850-2100)	57000 (35000-82000)
Kazakhstan	2016	Male	55 to 59	83% (74-89%)	17% (11-26%)	4.8 (2.9-7.3)	15% (9.3-21%)	18% (11-25%)	1300 (720-2000)	46000 (26000-68000)
Kazakhstan	2016	Male	60 to 64	81% (71-88%)	19% (12-29%)	4.2 (2.4-6.6)	12% (5.9-18%)	13% (6.5-21%)	1200 (540-1900)	34000 (16000-54000)
Kazakhstan	2016	Male	65 to 69	80% (70-89%)	20% (11-30%)	3.9 (2.4-5.8)	10% (4.9-16%)	11% (5.3-18%)	760 (320-1300)	18000 (8100-29000)
Kazakhstan	2016	Male	70 to 74	77% (65-85%)	23% (15-35%)	3.4 (1.9-5.4)	8% (2.7-14%)	9% (2.8-16%)	530 (160-980)	10000 (3300-18000)
Kazakhstan	2016	Male	75 to 79	72% (57-83%)	28% (17-43%)	2.8 (1.5-4.6)	5.7% (0.96-12%)	6.4% (0.87-13%)	510 (70-1100)	7800 (1200-16000)
Kazakhstan	2016	Male	80 to 84	68% (53-80%)	32% (20-47%)	2.4 (1.3-3.9)	3.9% (-0.33-9.9%)	4.1% (-0.71-11%)	260 (-45-700)	2900 (-250-7500)
Kazakhstan	2016	Male	85 to 89	65% (50-78%)	35% (22-50%)	2.2 (1.2-3.6)	2.1% (-1.5-7.5%)	2% (-2.3-8.3%)	40 (-47-170)	380 (-270-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kazakhstan	2016	Male	90 to 94	64% (50-78%)	36% (22-50%)	2 (1-3.4)	0.87% (-2.2-5.4%)	0.46% (-3.4-6.1%)	4.1 (-30-55)	48 (-120-310)
Kazakhstan	2016	Male	95 plus	63% (44-80%)	37% (20-56%)	1.7 (0.76-3.2)	0.21% (-2.8-5.3%)	-0.47% (-4.3-5.4%)	-0.65 (-6-7.9)	1.3 (-16-31)
Kyrgyzstan	1990	Female	All Ages	53% (47-61%)	47% (39-53%)	1 (0.78-1.3)	1.9% (0.96-3%)	1.8% (-0.63-4.6%)	300 (-110-790)	16000 (7900-26000)
Kyrgyzstan	1990	Female	15 to 19	56% (43-70%)	44% (30-57%)	1.1 (0.6-1.7)	2.2% (1.3-3.6%)	4.3% (2.1-7%)	6 (2.9-9.9)	590 (340-910)
Kyrgyzstan	1990	Female	20 to 24	62% (49-75%)	38% (25-51%)	1.5 (0.86-2.4)	3.6% (2.3-5.4%)	5.5% (2.9-8.9%)	9.9 (5.2-16)	980 (610-1500)
Kyrgyzstan	1990	Female	25 to 29	65% (51-77%)	35% (23-49%)	1.7 (1-2.5)	4.5% (3.1-6.3%)	6.9% (4.2-10%)	16 (9.8-24)	1500 (970-2000)
Kyrgyzstan	1990	Female	30 to 34	66% (52-79%)	34% (21-48%)	1.8 (1.1-2.7)	5.2% (3.8-7%)	8.5% (5.7-12%)	22 (15-30)	1700 (1200-2300)
Kyrgyzstan	1990	Female	35 to 39	66% (53-80%)	34% (20-47%)	1.8 (1.1-2.8)	5.7% (4.2-7.7%)	9.4% (6.5-13%)	25 (17-34)	1600 (1200-2200)
Kyrgyzstan	1990	Female	40 to 44	66% (53-79%)	34% (21-47%)	1.8 (1.1-2.8)	6.4% (4.5-8.8%)	10% (7.1-14%)	27 (18-38)	1500 (1000-2000)
Kyrgyzstan	1990	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.9 (1.1-2.9)	6.4% (4-9.3%)	9.8% (5.8-15%)	24 (15-36)	1100 (710-1700)
Kyrgyzstan	1990	Female	50 to 54	65% (51-77%)	35% (23-49%)	1.9 (1.1-2.8)	6.4% (3.5-9.7%)	9.1% (4.7-14%)	57 (30-91)	2300 (1300-3600)
Kyrgyzstan	1990	Female	55 to 59	63% (50-76%)	37% (24-50%)	1.8 (1.1-2.8)	5.3% (2.2-8.9%)	7.4% (3-13%)	55 (22-94)	1900 (820-3200)
Kyrgyzstan	1990	Female	60 to 64	61% (48-76%)	39% (24-52%)	1.8 (1-2.8)	3.8% (0.51-7.6%)	5% (0.57-10%)	61 (7-130)	1800 (240-3600)
Kyrgyzstan	1990	Female	65 to 69	60% (48-74%)	40% (26-52%)	1.7 (1.1-2.5)	2.4% (-0.64-6.1%)	3.2% (-0.97-8.1%)	40 (-12-100)	940 (-250-2300)
Kyrgyzstan	1990	Female	70 to 74	58% (46-70%)	42% (30-54%)	1.6 (0.95-2.5)	1.3% (-2.2-5.4%)	1.6% (-2.8-6.9%)	20 (-35-84)	380 (-650-1600)
Kyrgyzstan	1990	Female	75 to 79	55% (44-67%)	45% (33-56%)	1.5 (0.87-2.4)	0.23% (-3.2-4.5%)	0.21% (-4-5.5%)	3.8 (-73-98)	74 (-1000-1400)
Kyrgyzstan	1990	Female	80 to 84	53% (41-66%)	47% (34-59%)	1.5 (0.9-2.2)	-0.63% (-3.8-3.4%)	-0.87% (-4.7-4%)	-16 (-89-75)	-150 (-900-800)
Kyrgyzstan	1990	Female	85 to 89	52% (41-63%)	48% (37-59%)	1.4 (0.87-2.2)	-1.2% (-4.4-2.7%)	-1.6% (-5.5-3%)	-26 (-92-50)	-180 (-650-400)
Kyrgyzstan	1990	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.5 (0.88-2.4)	-1.6% (-4.8-3%)	-2% (-5.9-3.3%)	-17 (-50-28)	-79 (-240-150)
Kyrgyzstan	1990	Female	95 plus	53% (41-68%)	47% (32-59%)	1.4 (0.75-2.5)	-1.7% (-5.1-3.1%)	-2.3% (-6.2-3.5%)	-5.5 (-15-8.5)	-17 (-49-30)
Kyrgyzstan	1990	Male	All Ages	36% (31-41%)	64% (59-69%)	2.8 (2.2-3.4)	8.6% (7.2-10%)	11% (9-14%)	2200 (1700-2700)	90000 (75000-110000)
Kyrgyzstan	1990	Male	15 to 19	26% (18-35%)	74% (65-82%)	0.91 (0.52-1.5)	4.1% (2.8-6.1%)	5.4% (3.2-8.7%)	13 (7.6-21)	1300 (870-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	1990	Male	20 to 24	51% (38-64%)	49% (36-62%)	3 (1.9-4.3)	13% (8.9-17%)	16% (11-23%)	60 (39-85)	4900 (3500-6700)
Kyrgyzstan	1990	Male	25 to 29	64% (51-74%)	36% (26-49%)	5.2 (3.5-7.1)	19% (15-24%)	24% (18-31%)	130 (95-170)	9500 (7300-12000)
Kyrgyzstan	1990	Male	30 to 34	72% (60-82%)	28% (18-40%)	6.5 (4.5-8.7)	22% (18-26%)	27% (22-33%)	170 (140-220)	11000 (9000-14000)
Kyrgyzstan	1990	Male	35 to 39	75% (64-84%)	25% (16-36%)	6.3 (4.3-8.9)	22% (18-26%)	27% (22-33%)	170 (130-210)	9800 (7800-12000)
Kyrgyzstan	1990	Male	40 to 44	75% (64-84%)	25% (16-36%)	6.6 (4.5-9.1)	23% (19-27%)	28% (23-33%)	170 (140-200)	8600 (7100-10000)
Kyrgyzstan	1990	Male	45 to 49	72% (59-82%)	28% (18-41%)	7.4 (4.6-11)	23% (20-27%)	28% (23-33%)	160 (130-190)	7200 (6000-8500)
Kyrgyzstan	1990	Male	50 to 54	68% (54-80%)	32% (20-46%)	6.9 (4.4-9.9)	22% (18-25%)	25% (21-30%)	310 (260-380)	12000 (10000-15000)
Kyrgyzstan	1990	Male	55 to 59	64% (45-80%)	36% (20-55%)	5.1 (3-7.7)	18% (14-22%)	21% (16-25%)	270 (200-340)	9200 (7100-11000)
Kyrgyzstan	1990	Male	60 to 64	59% (39-77%)	41% (23-61%)	4 (2.2-6.2)	14% (9-18%)	15% (10-21%)	290 (180-380)	8200 (5300-11000)
Kyrgyzstan	1990	Male	65 to 69	55% (35-73%)	45% (27-65%)	3.4 (2-5.3)	11% (6.7-16%)	13% (7.5-18%)	160 (98-230)	3800 (2300-5400)
Kyrgyzstan	1990	Male	70 to 74	50% (32-68%)	50% (32-68%)	2.9 (1.6-4.8)	8.4% (3.9-13%)	9.4% (4.3-15%)	96 (44-150)	1800 (860-2800)
Kyrgyzstan	1990	Male	75 to 79	46% (29-65%)	54% (35-71%)	2.3 (1.2-4.1)	6% (1.9-10%)	6.6% (2.1-12%)	81 (25-150)	1200 (390-2100)
Kyrgyzstan	1990	Male	80 to 84	43% (26-60%)	57% (40-74%)	2 (1.1-3.4)	4.5% (1.1-8.9%)	5% (1-9.8%)	56 (11-110)	620 (140-1200)
Kyrgyzstan	1990	Male	85 to 89	41% (26-57%)	59% (43-74%)	1.8 (0.96-3)	3.3% (0.35-7.6%)	3.6% (0.13-8.4%)	27 (0.97-64)	220 (23-490)
Kyrgyzstan	1990	Male	90 to 94	40% (26-57%)	60% (43-74%)	1.7 (0.9-2.9)	2.2% (-0.39-6.4%)	2.3% (-0.77-7.1%)	8 (-2.7-25)	47 (-8.3-130)
Kyrgyzstan	1990	Male	95 plus	42% (24-62%)	58% (38-76%)	1.6 (0.68-3)	1.5% (-1.2-5.8%)	1.3% (-1.8-6.2%)	1.3 (-1.8-6.1)	5.9 (-4.5-22)
Kyrgyzstan	1995	Female	All Ages	54% (48-61%)	46% (39-52%)	0.92 (0.71-1.2)	2% (0.81-3.2%)	1.3% (-1.3-4.1%)	230 (-230-750)	16000 (6600-26000)
Kyrgyzstan	1995	Female	15 to 19	56% (44-70%)	44% (30-56%)	0.96 (0.55-1.5)	2.1% (1.2-3.2%)	3.8% (1.8-6.4%)	5.7 (2.8-9.8)	580 (320-910)
Kyrgyzstan	1995	Female	20 to 24	62% (49-75%)	38% (25-51%)	1.3 (0.77-2.1)	3.5% (2.1-5.4%)	5.2% (2.7-8.5%)	11 (5.7-19)	1100 (660-1700)
Kyrgyzstan	1995	Female	25 to 29	65% (51-77%)	35% (23-49%)	1.5 (0.91-2.3)	4.4% (3-6.3%)	6.6% (3.8-9.9%)	15 (8.6-23)	1300 (860-1800)
Kyrgyzstan	1995	Female	30 to 34	66% (52-79%)	34% (21-48%)	1.6 (0.97-2.4)	5.2% (3.7-7%)	7.9% (5.2-11%)	27 (17-38)	2000 (1400-2700)
Kyrgyzstan	1995	Female	35 to 39	66% (53-80%)	34% (20-47%)	1.6 (0.97-2.4)	5.7% (4-7.8%)	8.7% (5.9-12%)	35 (23-48)	2200 (1600-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	1995	Female	40 to 44	66% (53-80%)	34% (20-47%)	1.6 (0.99-2.4)	6.3% (4.3-8.6%)	9.6% (6.4-13%)	41 (27-58)	2200 (1500-2900)
Kyrgyzstan	1995	Female	45 to 49	66% (52-79%)	34% (21-48%)	1.7 (0.98-2.6)	6.1% (3.7-9.2%)	8.8% (5-13%)	40 (22-60)	1800 (1100-2700)
Kyrgyzstan	1995	Female	50 to 54	65% (51-78%)	35% (22-49%)	1.6 (1-2.6)	6% (3-9.5%)	7.9% (3.8-13%)	35 (16-57)	1400 (690-2100)
Kyrgyzstan	1995	Female	55 to 59	63% (50-76%)	37% (24-50%)	1.6 (0.96-2.4)	4.7% (1.6-8.4%)	6.1% (1.9-11%)	64 (20-120)	2100 (720-3900)
Kyrgyzstan	1995	Female	60 to 64	61% (48-76%)	39% (24-52%)	1.6 (0.91-2.5)	3% (-0.1-7%)	3.8% (-0.3-9%)	47 (-3.8-110)	1300 (-49-3100)
Kyrgyzstan	1995	Female	65 to 69	60% (48-74%)	40% (26-52%)	1.5 (0.95-2.2)	1.6% (-1.5-5.4%)	2% (-2-6.7%)	38 (-38-130)	890 (-840-3000)
Kyrgyzstan	1995	Female	70 to 74	58% (45-70%)	42% (30-55%)	1.5 (0.85-2.1)	0.43% (-2.9-4.3%)	0.52% (-3.6-5.4%)	9.7 (-67-99)	180 (-1200-1800)
Kyrgyzstan	1995	Female	75 to 79	55% (44-68%)	45% (32-56%)	1.4 (0.79-2.1)	-0.64% (-3.9-3.4%)	-0.82% (-4.7-3.9%)	-13 (-78-64)	-180 (-1100-920)
Kyrgyzstan	1995	Female	80 to 84	53% (41-66%)	47% (34-59%)	1.3 (0.82-2)	-1.6% (-4.8-2.1%)	-1.9% (-5.7-2.5%)	-37 (-110-48)	-370 (-1100-500)
Kyrgyzstan	1995	Female	85 to 89	53% (41-64%)	47% (36-59%)	1.3 (0.78-1.9)	-2.2% (-5.3-1.7%)	-2.7% (-6.3-1.9%)	-47 (-110-33)	-330 (-800-250)
Kyrgyzstan	1995	Female	90 to 94	53% (43-64%)	47% (36-57%)	1.3 (0.78-2.1)	-2.5% (-5.6-1.5%)	-3.1% (-6.7-1.7%)	-29 (-65-16)	-140 (-320-83)
Kyrgyzstan	1995	Female	95 plus	53% (41-68%)	47% (32-59%)	1.3 (0.66-2.2)	-2.7% (-6-1.6%)	-3.3% (-7.1-1.8%)	-11 (-24-6)	-35 (-77-21)
Kyrgyzstan	1995	Male	All Ages	36% (31-42%)	64% (58-69%)	2.5 (2-3)	10% (8.5-12%)	12% (9.6-15%)	2700 (2100-3300)	110000 (91000-130000)
Kyrgyzstan	1995	Male	15 to 19	26% (18-35%)	74% (65-82%)	0.81 (0.46-1.3)	3.9% (2.6-5.7%)	4.9% (2.9-7.9%)	14 (8.3-22)	1400 (920-2100)
Kyrgyzstan	1995	Male	20 to 24	52% (39-64%)	48% (36-61%)	2.7 (1.7-3.8)	12% (8.7-16%)	15% (10-20%)	68 (46-94)	5500 (3900-7300)
Kyrgyzstan	1995	Male	25 to 29	64% (52-75%)	36% (25-48%)	4.6 (3.2-6.3)	19% (15-23%)	23% (17-28%)	140 (100-170)	9600 (7400-12000)
Kyrgyzstan	1995	Male	30 to 34	73% (61-82%)	27% (18-39%)	5.8 (4-7.7)	22% (17-26%)	26% (20-31%)	210 (170-260)	13000 (11000-16000)
Kyrgyzstan	1995	Male	35 to 39	76% (65-84%)	24% (16-35%)	5.7 (3.9-7.7)	22% (17-26%)	25% (20-31%)	250 (190-300)	14000 (11000-17000)
Kyrgyzstan	1995	Male	40 to 44	75% (65-83%)	25% (17-35%)	5.9 (4.1-8)	23% (18-27%)	26% (21-31%)	290 (230-350)	14000 (12000-17000)
Kyrgyzstan	1995	Male	45 to 49	72% (59-82%)	28% (18-41%)	6.6 (4.2-9.3)	23% (19-28%)	27% (22-32%)	280 (230-340)	12000 (9900-14000)
Kyrgyzstan	1995	Male	50 to 54	69% (55-81%)	31% (19-45%)	6.1 (3.9-9)	22% (18-27%)	25% (20-30%)	230 (180-280)	8800 (7000-11000)
Kyrgyzstan	1995	Male	55 to 59	64% (46-81%)	36% (19-54%)	4.6 (2.8-6.9)	18% (13-22%)	20% (14-25%)	370 (260-460)	12000 (8800-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	1995	Male	60 to 64	59% (39-77%)	41% (23-61%)	3.5 (1.9-5.6)	13% (7.9-18%)	14% (8.6-20%)	300 (180-410)	8300 (5100-11000)
Kyrgyzstan	1995	Male	65 to 69	55% (35-73%)	45% (27-65%)	3.1 (1.8-4.7)	10% (5.7-15%)	11% (6.2-17%)	290 (160-430)	6700 (3700-9900)
Kyrgyzstan	1995	Male	70 to 74	51% (32-68%)	49% (32-68%)	2.5 (1.4-4.1)	7.5% (3.2-12%)	8.2% (3.4-13%)	130 (54-210)	2400 (1000-3900)
Kyrgyzstan	1995	Male	75 to 79	46% (29-65%)	54% (35-71%)	2.1 (1.1-3.5)	5% (1.3-9.6%)	5.4% (1.2-11%)	55 (13-110)	800 (210-1600)
Kyrgyzstan	1995	Male	80 to 84	43% (26-61%)	57% (39-74%)	1.8 (0.95-3)	3.5% (0.46-7.8%)	3.8% (0.3-8.5%)	39 (3.2-87)	430 (57-930)
Kyrgyzstan	1995	Male	85 to 89	41% (26-57%)	59% (43-74%)	1.6 (0.87-2.7)	2.4% (-0.23-6.1%)	2.5% (-0.53-6.7%)	19 (-4.1-51)	160 (-14-390)
Kyrgyzstan	1995	Male	90 to 94	41% (27-57%)	59% (43-73%)	1.5 (0.83-2.7)	1.5% (-0.96-5.3%)	1.4% (-1.4-5.8%)	4.6 (-4.7-19)	28 (-18-100)
Kyrgyzstan	1995	Male	95 plus	42% (25-62%)	58% (38-75%)	1.4 (0.62-2.8)	0.89% (-1.6-5%)	0.63% (-2.2-5.3%)	0.65 (-2.3-5.5)	3.6 (-6.5-20)
Kyrgyzstan	2000	Female	All Ages	55% (48-62%)	45% (38-52%)	0.95 (0.73-1.2)	2% (0.96-3.3%)	1.2% (-1.3-4.2%)	210 (-220-700)	16000 (7100-26000)
Kyrgyzstan	2000	Female	15 to 19	56% (44-70%)	44% (30-56%)	0.95 (0.54-1.5)	1.9% (1.1-3.1%)	3.8% (1.8-6.4%)	5.4 (2.5-9.2)	580 (320-910)
Kyrgyzstan	2000	Female	20 to 24	62% (50-74%)	38% (26-50%)	1.3 (0.76-2.1)	3.6% (2.2-5.5%)	5.8% (2.8-9.4%)	11 (5.6-19)	1100 (670-1700)
Kyrgyzstan	2000	Female	25 to 29	65% (51-77%)	35% (23-49%)	1.5 (0.91-2.3)	4.6% (3.1-6.3%)	7.3% (4.2-11%)	17 (9.6-25)	1500 (980-2100)
Kyrgyzstan	2000	Female	30 to 34	66% (52-79%)	34% (21-48%)	1.6 (0.95-2.4)	5.3% (3.9-7%)	8.6% (5.9-12%)	23 (16-32)	1700 (1300-2300)
Kyrgyzstan	2000	Female	35 to 39	66% (53-79%)	34% (21-47%)	1.6 (0.95-2.4)	5.7% (4.2-7.7%)	9.4% (6.5-13%)	35 (24-47)	2200 (1600-2900)
Kyrgyzstan	2000	Female	40 to 44	66% (52-79%)	34% (21-48%)	1.6 (1-2.4)	6.2% (4.3-8.2%)	10% (7.1-14%)	45 (31-61)	2400 (1700-3200)
Kyrgyzstan	2000	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.6 (0.97-2.5)	6% (3.6-8.7%)	9.3% (5.4-14%)	47 (27-69)	2200 (1400-3100)
Kyrgyzstan	2000	Female	50 to 54	65% (51-78%)	35% (22-49%)	1.6 (0.97-2.5)	5.7% (3-8.9%)	8.2% (4-13%)	43 (21-67)	1700 (880-2700)
Kyrgyzstan	2000	Female	55 to 59	63% (50-77%)	37% (23-50%)	1.6 (0.95-2.4)	4.7% (1.6-8.3%)	6.3% (1.9-11%)	33 (9.9-59)	1100 (380-2000)
Kyrgyzstan	2000	Female	60 to 64	61% (48-76%)	39% (24-52%)	1.5 (0.91-2.5)	3% (-0.24-7%)	3.8% (-0.38-8.8%)	48 (-4.9-110)	1400 (-100-3200)
Kyrgyzstan	2000	Female	65 to 69	60% (47-73%)	40% (27-53%)	1.5 (0.93-2.2)	1.6% (-1.6-5.3%)	2% (-2-6.7%)	30 (-30-100)	700 (-670-2400)
Kyrgyzstan	2000	Female	70 to 74	57% (45-70%)	43% (30-55%)	1.4 (0.85-2.1)	0.35% (-3-4.3%)	0.42% (-3.7-5.3%)	9.6 (-85-120)	190 (-1600-2300)
Kyrgyzstan	2000	Female	75 to 79	55% (44-67%)	45% (33-56%)	1.4 (0.79-2)	-0.82% (-4-3%)	-1.1% (-5-3.6%)	-22 (-100-71)	-290 (-1400-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	2000	Female	80 to 84	53% (41-66%)	47% (34-59%)	1.3 (0.8-1.9)	-1.9% (-5.1-1.7%)	-2.4% (-6.3-2%)	-38 (-100-33)	-380 (-1000-360)
Kyrgyzstan	2000	Female	85 to 89	52% (41-64%)	48% (36-59%)	1.3 (0.77-1.9)	-2.5% (-5.6-1.1%)	-3.1% (-6.8-1.1%)	-40 (-88-15)	-280 (-620-130)
Kyrgyzstan	2000	Female	90 to 94	53% (43-64%)	47% (36-57%)	1.3 (0.77-2)	-2.8% (-5.9-0.94%)	-3.4% (-7.2-0.94%)	-28 (-59-8)	-140 (-290-48)
Kyrgyzstan	2000	Female	95 plus	53% (41-68%)	47% (32-59%)	1.2 (0.65-2.1)	-2.9% (-6.2-1.1%)	-3.6% (-7.6-1.1%)	-11 (-23-3.5)	-35 (-74-13)
Kyrgyzstan	2000	Male	All Ages	37% (33-43%)	63% (57-67%)	2.6 (2.1-3.1)	11% (9.8-13%)	13% (11-16%)	2700 (2200-3300)	110000 (96000-130000)
Kyrgyzstan	2000	Male	15 to 19	26% (18-34%)	74% (66-82%)	0.79 (0.46-1.3)	3.8% (2.6-5.5%)	5.1% (3.1-8.2%)	13 (7.7-21)	1400 (910-2000)
Kyrgyzstan	2000	Male	20 to 24	51% (38-63%)	49% (37-62%)	2.6 (1.7-3.8)	13% (9.5-17%)	17% (12-22%)	73 (51-98)	5800 (4200-7700)
Kyrgyzstan	2000	Male	25 to 29	64% (52-74%)	36% (26-48%)	4.5 (3.1-6.2)	20% (16-24%)	25% (19-30%)	150 (120-190)	11000 (8600-13000)
Kyrgyzstan	2000	Male	30 to 34	72% (60-82%)	28% (18-40%)	5.6 (3.9-7.5)	22% (18-26%)	27% (22-32%)	200 (160-230)	13000 (10000-15000)
Kyrgyzstan	2000	Male	35 to 39	75% (64-84%)	25% (16-36%)	5.5 (3.9-7.6)	23% (19-27%)	27% (22-32%)	270 (210-320)	15000 (12000-18000)
Kyrgyzstan	2000	Male	40 to 44	75% (64-83%)	25% (17-36%)	5.8 (4.1-7.6)	24% (20-28%)	29% (24-33%)	330 (270-390)	17000 (14000-19000)
Kyrgyzstan	2000	Male	45 to 49	72% (58-82%)	28% (18-42%)	6.4 (4.1-9)	24% (20-28%)	29% (24-33%)	340 (280-400)	15000 (12000-18000)
Kyrgyzstan	2000	Male	50 to 54	68% (54-80%)	32% (20-46%)	6 (3.8-8.7)	23% (19-27%)	27% (21-32%)	290 (230-350)	11000 (9100-14000)
Kyrgyzstan	2000	Male	55 to 59	63% (44-80%)	37% (20-56%)	4.4 (2.6-6.9)	18% (13-23%)	21% (15-27%)	210 (150-260)	6800 (4900-8700)
Kyrgyzstan	2000	Male	60 to 64	59% (39-77%)	41% (23-61%)	3.5 (1.9-5.5)	13% (8.3-18%)	15% (9.2-21%)	300 (190-430)	8600 (5300-12000)
Kyrgyzstan	2000	Male	65 to 69	55% (35-74%)	45% (26-65%)	3 (1.8-4.7)	10% (5.8-15%)	12% (6.3-18%)	240 (130-370)	5700 (3200-8400)
Kyrgyzstan	2000	Male	70 to 74	51% (32-68%)	49% (32-68%)	2.5 (1.4-4)	7.4% (3.2-12%)	8.3% (3.4-14%)	200 (82-330)	3700 (1600-6200)
Kyrgyzstan	2000	Male	75 to 79	46% (29-65%)	54% (35-71%)	2.1 (1.1-3.5)	4.8% (1.2-9.3%)	5.3% (1.1-11%)	70 (15-140)	1000 (260-2000)
Kyrgyzstan	2000	Male	80 to 84	43% (26-60%)	57% (40-74%)	1.8 (0.96-3)	3.2% (0.28-7.5%)	3.4% (0.061-8.3%)	28 (0.5-67)	310 (27-720)
Kyrgyzstan	2000	Male	85 to 89	41% (26-57%)	59% (43-74%)	1.6 (0.86-2.6)	2% (-0.49-5.7%)	2% (-0.88-6.3%)	10 (-4.6-32)	86 (-21-250)
Kyrgyzstan	2000	Male	90 to 94	40% (26-57%)	60% (43-74%)	1.5 (0.79-2.6)	1.1% (-1.2-4.7%)	1% (-1.7-5.1%)	2.7 (-4.8-14)	18 (-19-74)
Kyrgyzstan	2000	Male	95 plus	42% (24-61%)	58% (39-76%)	1.4 (0.61-2.7)	0.69% (-1.7-4.7%)	0.36% (-2.4-5%)	0.29 (-1.9-4)	2.1 (-5.4-15)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	2005	Female	All Ages	55% (49-63%)	45% (37-51%)	1 (0.78-1.3)	2.4% (1.2-3.8%)	1.5% (-1.4-4.6%)	260 (-250-810)	19000 (9300-30000)
Kyrgyzstan	2005	Female	15 to 19	55% (43-69%)	45% (31-57%)	0.97 (0.55-1.5)	2% (1.1-3.1%)	4.2% (1.9-7.1%)	6 (2.8-10)	650 (370-1000)
Kyrgyzstan	2005	Female	20 to 24	62% (49-74%)	38% (26-51%)	1.4 (0.77-2.1)	3.7% (2.3-5.5%)	6.4% (3.3-10%)	13 (6.6-20)	1300 (780-1900)
Kyrgyzstan	2005	Female	25 to 29	65% (51-78%)	35% (22-49%)	1.5 (0.93-2.3)	4.9% (3.4-6.8%)	8.2% (5.1-12%)	20 (12-28)	1700 (1100-2300)
Kyrgyzstan	2005	Female	30 to 34	66% (52-80%)	34% (20-48%)	1.6 (0.96-2.4)	5.8% (4.4-7.6%)	9.7% (7-13%)	30 (21-40)	2100 (1600-2800)
Kyrgyzstan	2005	Female	35 to 39	66% (53-80%)	34% (20-47%)	1.6 (0.99-2.5)	6.3% (4.7-8.3%)	11% (7.6-14%)	40 (29-52)	2400 (1800-3100)
Kyrgyzstan	2005	Female	40 to 44	66% (52-79%)	34% (21-48%)	1.7 (1.1-2.4)	6.8% (5-9%)	11% (8.2-15%)	57 (41-76)	3000 (2200-3900)
Kyrgyzstan	2005	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.7 (1-2.5)	6.5% (4.3-9.2%)	10% (6.7-14%)	65 (42-93)	2900 (2000-4100)
Kyrgyzstan	2005	Female	50 to 54	64% (50-78%)	36% (22-50%)	1.7 (1-2.5)	6.4% (3.6-9.5%)	9.2% (5-14%)	67 (37-100)	2600 (1500-3900)
Kyrgyzstan	2005	Female	55 to 59	63% (50-76%)	37% (24-50%)	1.6 (0.99-2.5)	5.4% (2.3-9.1%)	7.4% (3-13%)	57 (23-96)	1900 (810-3200)
Kyrgyzstan	2005	Female	60 to 64	61% (48-75%)	39% (25-52%)	1.6 (0.92-2.5)	3.3% (-0.099-7.6%)	4.2% (-0.23-9.6%)	31 (-1.7-71)	870 (-26-2000)
Kyrgyzstan	2005	Female	65 to 69	59% (48-74%)	41% (26-52%)	1.5 (0.97-2.3)	1.6% (-1.6-5.3%)	2% (-2.2-6.7%)	34 (-37-110)	800 (-800-2600)
Kyrgyzstan	2005	Female	70 to 74	57% (46-70%)	43% (30-54%)	1.5 (0.88-2.2)	0.26% (-3.3-4.5%)	0.29% (-4.1-5.5%)	5.6 (-81-110)	120 (-1400-1900)
Kyrgyzstan	2005	Female	75 to 79	55% (44-67%)	45% (33-56%)	1.4 (0.8-2.1)	-0.96% (-4.4-2.9%)	-1.2% (-5.4-3.4%)	-34 (-150-94)	-450 (-2100-1400)
Kyrgyzstan	2005	Female	80 to 84	53% (41-66%)	47% (34-59%)	1.3 (0.84-2)	-2% (-5.5-1.8%)	-2.5% (-6.6-1.9%)	-59 (-160-46)	-590 (-1600-500)
Kyrgyzstan	2005	Female	85 to 89	52% (42-63%)	48% (37-58%)	1.3 (0.79-2)	-2.5% (-6-1.1%)	-3.1% (-7.2-1.1%)	-41 (-94-15)	-280 (-660-120)
Kyrgyzstan	2005	Female	90 to 94	53% (43-64%)	47% (36-57%)	1.3 (0.79-2.1)	-2.8% (-6-1.1%)	-3.4% (-7.3-1.2%)	-22 (-47-7.9)	-110 (-230-44)
Kyrgyzstan	2005	Female	95 plus	53% (41-68%)	47% (32-59%)	1.3 (0.67-2.2)	-2.8% (-6.4-1.7%)	-3.5% (-7.7-1.8%)	-9.7 (-22-5.1)	-30 (-69-18)
Kyrgyzstan	2005	Male	All Ages	39% (34-44%)	61% (56-66%)	2.8 (2.3-3.4)	13% (11-15%)	14% (12-17%)	3200 (2500-3800)	130000 (110000-150000)
Kyrgyzstan	2005	Male	15 to 19	26% (18-34%)	74% (66-82%)	0.81 (0.47-1.3)	3.9% (2.7-5.6%)	5.6% (3.4-8.7%)	13 (8.2-21)	1400 (990-2100)
Kyrgyzstan	2005	Male	20 to 24	51% (38-63%)	49% (37-62%)	2.7 (1.7-3.9)	13% (9.4-17%)	17% (12-23%)	70 (48-94)	5800 (4200-7500)
Kyrgyzstan	2005	Male	25 to 29	64% (51-75%)	36% (25-49%)	4.6 (3.2-6.3)	20% (16-24%)	26% (20-32%)	150 (120-190)	11000 (8600-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	2005	Male	30 to 34	72% (60-82%)	28% (18-40%)	5.8 (4.1-7.7)	23% (19-27%)	28% (23-33%)	240 (190-280)	15000 (12000-18000)
Kyrgyzstan	2005	Male	35 to 39	75% (64-84%)	25% (16-36%)	5.7 (4-7.8)	24% (20-28%)	29% (23-33%)	280 (230-330)	16000 (13000-18000)
Kyrgyzstan	2005	Male	40 to 44	75% (64-83%)	25% (17-36%)	5.9 (4.2-7.8)	25% (21-29%)	30% (25-34%)	390 (320-460)	19000 (16000-22000)
Kyrgyzstan	2005	Male	45 to 49	72% (58-81%)	28% (19-42%)	6.6 (4.4-9.3)	25% (21-29%)	30% (25-34%)	440 (370-520)	19000 (16000-23000)
Kyrgyzstan	2005	Male	50 to 54	68% (55-80%)	32% (20-45%)	6.2 (4-8.8)	24% (19-28%)	27% (22-32%)	420 (340-510)	16000 (13000-19000)
Kyrgyzstan	2005	Male	55 to 59	63% (44-80%)	37% (20-56%)	4.6 (2.7-7)	19% (14-24%)	22% (16-27%)	300 (220-380)	9900 (7400-12000)
Kyrgyzstan	2005	Male	60 to 64	59% (39-77%)	41% (23-61%)	3.5 (1.9-5.7)	14% (8.7-19%)	16% (9.5-21%)	190 (120-270)	5300 (3300-7400)
Kyrgyzstan	2005	Male	65 to 69	55% (35-73%)	45% (27-65%)	3.1 (1.8-4.9)	11% (6.2-16%)	12% (6.8-18%)	290 (160-420)	6600 (3800-9500)
Kyrgyzstan	2005	Male	70 to 74	50% (32-68%)	50% (32-68%)	2.6 (1.4-4.1)	7.6% (3.3-13%)	8.4% (3.5-14%)	190 (79-320)	3500 (1500-5900)
Kyrgyzstan	2005	Male	75 to 79	45% (28-64%)	55% (36-72%)	2.1 (1.1-3.5)	4.9% (1.2-9.4%)	5.4% (1.1-11%)	130 (27-250)	1900 (460-3700)
Kyrgyzstan	2005	Male	80 to 84	42% (26-60%)	58% (40-74%)	1.8 (1-3.1)	3.2% (0.12-7.3%)	3.4% (-0.071-8.2%)	51 (-1-120)	570 (22-1300)
Kyrgyzstan	2005	Male	85 to 89	40% (26-57%)	60% (43-74%)	1.6 (0.89-2.6)	2% (-0.72-5.8%)	2% (-1.1-6.4%)	8.9 (-5.2-29)	75 (-28-220)
Kyrgyzstan	2005	Male	90 to 94	40% (26-56%)	60% (44-74%)	1.5 (0.81-2.6)	1.2% (-1.3-4.9%)	1.1% (-1.8-5.4%)	2.1 (-3.4-10)	13 (-14-54)
Kyrgyzstan	2005	Male	95 plus	41% (24-61%)	59% (39-76%)	1.4 (0.62-2.7)	0.8% (-1.8-5%)	0.52% (-2.5-5.4%)	0.34 (-1.6-3.5)	2 (-4.5-13)
Kyrgyzstan	2010	Female	All Ages	55% (49-63%)	45% (37-51%)	1 (0.78-1.3)	2.2% (1.2-3.5%)	1.1% (-1.6-4%)	180 (-280-690)	17000 (8900-27000)
Kyrgyzstan	2010	Female	15 to 19	55% (42-69%)	45% (31-58%)	0.95 (0.54-1.5)	1.9% (1-3%)	3.8% (1.6-6.5%)	5.8 (2.5-10)	630 (350-980)
Kyrgyzstan	2010	Female	20 to 24	62% (48-74%)	38% (26-52%)	1.3 (0.77-2.1)	3.5% (2.2-5.1%)	5.9% (3.1-9.4%)	13 (6.9-22)	1400 (880-2100)
Kyrgyzstan	2010	Female	25 to 29	64% (51-77%)	36% (23-49%)	1.5 (0.91-2.3)	4.5% (3.1-6.1%)	7.7% (4.7-11%)	17 (10-25)	1600 (1000-2200)
Kyrgyzstan	2010	Female	30 to 34	66% (52-78%)	34% (22-48%)	1.6 (0.95-2.4)	5.6% (4.1-7.3%)	9.6% (6.8-13%)	27 (19-36)	2000 (1500-2600)
Kyrgyzstan	2010	Female	35 to 39	66% (53-80%)	34% (20-47%)	1.6 (0.98-2.5)	5.9% (4.4-7.8%)	10% (7.4-14%)	35 (25-47)	2200 (1600-2900)
Kyrgyzstan	2010	Female	40 to 44	66% (52-79%)	34% (21-48%)	1.6 (1-2.5)	6.5% (4.7-8.5%)	11% (8.1-14%)	48 (34-62)	2500 (1900-3300)
Kyrgyzstan	2010	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.7 (1-2.6)	5.9% (3.9-8.5%)	9.8% (6.4-14%)	59 (38-83)	2800 (1800-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	2010	Female	50 to 54	64% (50-78%)	36% (22-50%)	1.7 (1-2.5)	5.7% (3.4-8.7%)	8.7% (4.9-13%)	68 (38-100)	2700 (1600-4100)
Kyrgyzstan	2010	Female	55 to 59	62% (50-76%)	38% (24-50%)	1.6 (0.98-2.5)	4.8% (2.1-8%)	6.9% (2.8-12%)	60 (25-100)	2000 (900-3400)
Kyrgyzstan	2010	Female	60 to 64	61% (47-76%)	39% (24-53%)	1.6 (0.92-2.5)	2.9% (-0.063-6.3%)	3.8% (-0.3-8.4%)	33 (-2.6-74)	950 (-22-2100)
Kyrgyzstan	2010	Female	65 to 69	59% (47-74%)	41% (26-53%)	1.5 (0.96-2.2)	1.3% (-1.7-4.8%)	1.7% (-2.4-6.3%)	14 (-21-53)	340 (-450-1200)
Kyrgyzstan	2010	Female	70 to 74	57% (45-70%)	43% (30-55%)	1.5 (0.85-2.2)	-0.02% (-3.5-3.9%)	-0.097% (-4.6-4.7%)	-1.7 (-85-88)	-6.8 (-1500-1600)
Kyrgyzstan	2010	Female	75 to 79	54% (44-67%)	46% (33-56%)	1.4 (0.78-2.1)	-1.3% (-4.7-2.6%)	-1.7% (-5.8-3%)	-34 (-120-65)	-450 (-1700-930)
Kyrgyzstan	2010	Female	80 to 84	53% (40-66%)	47% (34-60%)	1.3 (0.81-2)	-2.3% (-5.7-1.4%)	-2.9% (-7-1.5%)	-63 (-150-33)	-610 (-1500-380)
Kyrgyzstan	2010	Female	85 to 89	52% (41-63%)	48% (37-59%)	1.3 (0.78-1.9)	-2.8% (-6.3-0.63%)	-3.5% (-7.7-0.55%)	-69 (-150-11)	-480 (-1100-110)
Kyrgyzstan	2010	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.3 (0.76-2.1)	-3% (-6.4-0.78%)	-3.7% (-7.8-0.65%)	-26 (-54-4.6)	-120 (-260-32)
Kyrgyzstan	2010	Female	95 plus	53% (40-68%)	47% (32-60%)	1.3 (0.65-2.2)	-3% (-6.6-1.2%)	-3.8% (-8-1.2%)	-8.7 (-19-3)	-27 (-60-11)
Kyrgyzstan	2010	Male	All Ages	39% (35-43%)	61% (57-65%)	2.8 (2.3-3.4)	12% (10-14%)	13% (11-16%)	2900 (2300-3600)	120000 (100000-150000)
Kyrgyzstan	2010	Male	15 to 19	19% (13-28%)	81% (72-87%)	0.6 (0.34-0.99)	3.1% (2.2-4.4%)	3.9% (2.3-6.4%)	10 (6.1-16)	1200 (820-1700)
Kyrgyzstan	2010	Male	20 to 24	48% (39-58%)	52% (42-61%)	2.5 (1.6-3.5)	10% (7.7-14%)	13% (9.3-19%)	63 (43-87)	5500 (4100-7300)
Kyrgyzstan	2010	Male	25 to 29	65% (55-72%)	35% (28-45%)	4.7 (3.3-6.3)	18% (14-22%)	23% (18-29%)	130 (100-170)	9800 (7800-12000)
Kyrgyzstan	2010	Male	30 to 34	73% (64-81%)	27% (19-36%)	5.9 (4.2-7.9)	22% (19-26%)	27% (22-32%)	210 (170-250)	13000 (11000-16000)
Kyrgyzstan	2010	Male	35 to 39	77% (69-83%)	23% (17-31%)	5.8 (4.1-7.8)	23% (19-27%)	28% (23-33%)	260 (210-300)	15000 (12000-17000)
Kyrgyzstan	2010	Male	40 to 44	77% (70-83%)	23% (17-30%)	6.1 (4.4-7.9)	24% (20-28%)	29% (25-34%)	320 (270-380)	16000 (14000-19000)
Kyrgyzstan	2010	Male	45 to 49	74% (64-82%)	26% (18-36%)	6.8 (4.8-9.3)	24% (21-28%)	29% (25-34%)	420 (350-490)	18000 (16000-22000)
Kyrgyzstan	2010	Male	50 to 54	69% (56-80%)	31% (20-44%)	6.1 (4-8.6)	23% (19-27%)	27% (22-32%)	450 (370-540)	17000 (14000-21000)
Kyrgyzstan	2010	Male	55 to 59	62% (43-79%)	38% (21-57%)	4.4 (2.6-6.9)	18% (13-23%)	21% (15-27%)	360 (260-460)	12000 (8600-15000)
Kyrgyzstan	2010	Male	60 to 64	57% (37-75%)	43% (25-63%)	3.4 (1.9-5.5)	13% (8.1-18%)	15% (8.8-21%)	220 (130-310)	6200 (3800-8700)
Kyrgyzstan	2010	Male	65 to 69	54% (34-72%)	46% (28-66%)	3 (1.7-4.8)	10% (5.7-15%)	12% (6.3-18%)	140 (76-210)	3300 (1800-4900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	2010	Male	70 to 74	49% (31-67%)	51% (33-69%)	2.5 (1.3-4)	7.1% (2.7-12%)	7.9% (3-14%)	170 (63-300)	3200 (1300-5500)
Kyrgyzstan	2010	Male	75 to 79	44% (28-63%)	56% (37-72%)	2 (1-3.4)	4.5% (0.82-9.2%)	4.9% (0.67-10%)	92 (13-190)	1400 (240-2800)
Kyrgyzstan	2010	Male	80 to 84	42% (25-59%)	58% (41-75%)	1.7 (0.92-2.9)	2.8% (-0.32-7.1%)	2.9% (-0.62-7.8%)	49 (-10-130)	550 (-64-1400)
Kyrgyzstan	2010	Male	85 to 89	40% (24-55%)	60% (45-76%)	1.6 (0.8-2.6)	1.6% (-1.1-5.3%)	1.5% (-1.6-5.7%)	17 (-18-64)	150 (-100-500)
Kyrgyzstan	2010	Male	90 to 94	39% (25-55%)	61% (45-75%)	1.4 (0.77-2.5)	0.92% (-1.5-4.6%)	0.75% (-2-5%)	1.7 (-4.6-11)	12 (-19-60)
Kyrgyzstan	2010	Male	95 plus	40% (23-61%)	60% (39-77%)	1.3 (0.59-2.6)	0.56% (-2-4.7%)	0.24% (-2.7-5%)	0.14 (-1.6-3)	1.3 (-4.6-11)
Kyrgyzstan	2016	Female	All Ages	55% (49-62%)	45% (38-51%)	0.98 (0.69-1.3)	2.1% (1-3.4%)	0.93% (-1.7-4.1%)	150 (-260-630)	16000 (7700-26000)
Kyrgyzstan	2016	Female	15 to 19	55% (42-69%)	45% (31-58%)	0.91 (0.51-1.4)	1.8% (0.99-2.8%)	4.1% (1.6-7.2%)	3.8 (1.4-6.7)	450 (250-710)
Kyrgyzstan	2016	Female	20 to 24	62% (49-74%)	38% (26-51%)	1.3 (0.72-2.1)	3.2% (2-4.7%)	5.9% (3-9.7%)	8.9 (4.5-15)	1000 (640-1600)
Kyrgyzstan	2016	Female	25 to 29	65% (51-77%)	35% (23-49%)	1.4 (0.8-2.3)	4% (2.8-5.5%)	7.4% (4.6-11%)	16 (9.7-24)	1600 (1100-2200)
Kyrgyzstan	2016	Female	30 to 34	66% (52-79%)	34% (21-48%)	1.5 (0.89-2.4)	4.9% (3.7-6.6%)	9.4% (6.7-13%)	25 (17-34)	1900 (1400-2500)
Kyrgyzstan	2016	Female	35 to 39	66% (53-80%)	34% (20-47%)	1.5 (0.88-2.4)	5.4% (4-7.1%)	10% (7.5-14%)	30 (21-41)	1900 (1400-2600)
Kyrgyzstan	2016	Female	40 to 44	66% (52-79%)	34% (21-48%)	1.6 (0.91-2.4)	5.8% (4.2-7.8%)	11% (7.9-15%)	41 (29-57)	2200 (1600-3000)
Kyrgyzstan	2016	Female	45 to 49	66% (51-79%)	34% (21-49%)	1.6 (0.92-2.5)	5.3% (3.5-7.5%)	9.6% (6.2-14%)	45 (29-65)	2100 (1400-3000)
Kyrgyzstan	2016	Female	50 to 54	64% (50-77%)	36% (23-50%)	1.6 (0.88-2.5)	5.1% (3-7.9%)	8.3% (4.6-13%)	66 (36-100)	2600 (1500-4000)
Kyrgyzstan	2016	Female	55 to 59	63% (50-75%)	37% (25-50%)	1.5 (0.88-2.4)	4.2% (1.8-7.3%)	6.4% (2.5-11%)	64 (25-110)	2200 (910-3800)
Kyrgyzstan	2016	Female	60 to 64	61% (48-75%)	39% (25-52%)	1.5 (0.83-2.4)	2.5% (-0.28-5.8%)	3.5% (-0.54-8.1%)	43 (-6.4-100)	1300 (-130-3000)
Kyrgyzstan	2016	Female	65 to 69	59% (47-74%)	41% (26-53%)	1.4 (0.84-2.2)	1.1% (-1.8-4.4%)	1.5% (-2.6-6.1%)	15 (-27-64)	380 (-580-1500)
Kyrgyzstan	2016	Female	70 to 74	57% (46-70%)	43% (30-54%)	1.4 (0.75-2.2)	-0.22% (-3.4-3.7%)	-0.36% (-4.7-4.7%)	-3.9 (-49-48)	-57 (-900-920)
Kyrgyzstan	2016	Female	75 to 79	54% (43-67%)	46% (33-57%)	1.3 (0.69-2.1)	-1.3% (-4.4-2.5%)	-1.8% (-5.8-3%)	-35 (-110-54)	-460 (-1500-820)
Kyrgyzstan	2016	Female	80 to 84	52% (40-66%)	48% (34-60%)	1.2 (0.7-2)	-2.3% (-5.5-1.3%)	-3% (-7-1.4%)	-64 (-150-29)	-630 (-1500-340)
Kyrgyzstan	2016	Female	85 to 89	52% (41-63%)	48% (37-59%)	1.2 (0.7-1.9)	-2.9% (-6.3-0.58%)	-3.8% (-8-0.46%)	-57 (-120-6.9)	-400 (-840-75)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kyrgyzstan	2016	Female	90 to 94	53% (42-64%)	47% (36-58%)	1.2 (0.7-2)	-3.1% (-6.3-0.47%)	-4% (-7.9-0.31%)	-39 (-78-3.1)	-190 (-370-28)
Kyrgyzstan	2016	Female	95 plus	53% (40-68%)	47% (32-60%)	1.2 (0.6-2.2)	-3.2% (-6.8-0.72%)	-4% (-8.4-0.62%)	-11 (-22-1.6)	-33 (-70-7.3)
Kyrgyzstan	2016	Male	All Ages	39% (35-44%)	61% (56-65%)	2.7 (2-3.6)	11% (9.3-14%)	13% (9.6-16%)	2600 (1900-3400)	110000 (90000-140000)
Kyrgyzstan	2016	Male	15 to 19	19% (13-27%)	81% (73-87%)	0.56 (0.3-0.94)	3.3% (2.3-4.6%)	4.6% (2.7-7.3%)	6.9 (4-11)	920 (640-1300)
Kyrgyzstan	2016	Male	20 to 24	48% (38-58%)	52% (42-62%)	2.3 (1.4-3.4)	11% (7.9-14%)	15% (10-22%)	45 (29-64)	4300 (3100-5800)
Kyrgyzstan	2016	Male	25 to 29	65% (55-72%)	35% (28-45%)	4.5 (3-6.3)	18% (14-22%)	24% (18-31%)	130 (90-170)	9900 (7400-13000)
Kyrgyzstan	2016	Male	30 to 34	73% (63-80%)	27% (20-37%)	5.6 (3.8-7.8)	21% (17-26%)	28% (22-33%)	190 (150-240)	13000 (10000-16000)
Kyrgyzstan	2016	Male	35 to 39	76% (68-83%)	24% (17-32%)	5.5 (3.6-8)	22% (18-26%)	28% (22-34%)	210 (160-270)	13000 (9800-15000)
Kyrgyzstan	2016	Male	40 to 44	77% (69-83%)	23% (17-31%)	5.8 (3.8-8.3)	23% (19-27%)	29% (24-34%)	270 (210-330)	14000 (11000-17000)
Kyrgyzstan	2016	Male	45 to 49	74% (63-82%)	26% (18-37%)	6.5 (4.3-9.4)	23% (19-26%)	29% (24-33%)	310 (250-380)	14000 (11000-17000)
Kyrgyzstan	2016	Male	50 to 54	68% (55-80%)	32% (20-45%)	5.8 (3.6-8.8)	21% (17-25%)	26% (21-31%)	430 (330-530)	17000 (13000-21000)
Kyrgyzstan	2016	Male	55 to 59	62% (42-79%)	38% (21-58%)	4.2 (2.4-6.8)	17% (12-21%)	20% (14-26%)	380 (250-500)	13000 (8700-17000)
Kyrgyzstan	2016	Male	60 to 64	57% (36-75%)	43% (25-64%)	3.2 (1.6-5.2)	12% (7.1-17%)	14% (7.8-20%)	290 (170-430)	8400 (4900-12000)
Kyrgyzstan	2016	Male	65 to 69	53% (33-73%)	47% (27-67%)	2.8 (1.5-4.5)	9.4% (4.7-14%)	11% (5.3-17%)	150 (73-230)	3500 (1800-5400)
Kyrgyzstan	2016	Male	70 to 74	48% (30-67%)	52% (33-70%)	2.3 (1.2-3.9)	6.4% (2.3-11%)	7.2% (2.4-13%)	87 (29-160)	1700 (580-3000)
Kyrgyzstan	2016	Male	75 to 79	44% (27-63%)	56% (37-73%)	1.9 (0.93-3.3)	3.9% (0.43-8.5%)	4.3% (0.22-9.6%)	70 (3.3-160)	1100 (110-2300)
Kyrgyzstan	2016	Male	80 to 84	41% (24-58%)	59% (42-76%)	1.6 (0.83-2.8)	2.3% (-0.72-6.5%)	2.4% (-1.2-7.4%)	39 (-19-120)	450 (-150-1300)
Kyrgyzstan	2016	Male	85 to 89	39% (24-55%)	61% (45-76%)	1.4 (0.72-2.5)	1.2% (-1.4-5.2%)	1.1% (-1.9-5.7%)	9.9 (-18-53)	95 (-100-410)
Kyrgyzstan	2016	Male	90 to 94	38% (24-55%)	62% (45-76%)	1.3 (0.66-2.4)	0.62% (-1.7-4.1%)	0.39% (-2.3-4.6%)	1.9 (-12-23)	19 (-50-120)
Kyrgyzstan	2016	Male	95 plus	40% (23-60%)	60% (40-77%)	1.2 (0.52-2.4)	0.29% (-2.2-4.1%)	-0.098% (-3-4.4%)	-0.073 (-2.2-3.2)	0.83 (-6.1-12)
Mongolia	1990	Female	All Ages	43% (35-51%)	57% (49-65%)	0.72 (0.54-0.95)	1.6% (0.96-2.4%)	2.6% (1.1-4.6%)	220 (93-410)	8100 (5000-12000)
Mongolia	1990	Female	15 to 19	41% (26-57%)	59% (43-74%)	0.65 (0.27-1.2)	1.6% (0.56-3.2%)	2.4% (0.37-5.2%)	3.1 (0.48-7)	300 (100-600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	1990	Female	20 to 24	54% (37-70%)	46% (30-63%)	1.1 (0.45-1.9)	3% (1.2-5.4%)	3.8% (0.98-7.7%)	6.4 (1.6-14)	600 (240-1100)
Mongolia	1990	Female	25 to 29	59% (39-75%)	41% (25-61%)	1.3 (0.57-2.3)	3.8% (1.8-6.4%)	5.1% (1.8-9.5%)	9 (3.2-17)	730 (340-1200)
Mongolia	1990	Female	30 to 34	61% (41-77%)	39% (23-59%)	1.4 (0.68-2.4)	4.2% (2.4-6.8%)	6% (2.8-11%)	9.7 (4.5-17)	690 (380-1100)
Mongolia	1990	Female	35 to 39	62% (45-78%)	38% (22-55%)	1.5 (0.66-2.8)	4.4% (2.2-7.5%)	6.3% (2.8-11%)	10 (4.6-19)	630 (320-1100)
Mongolia	1990	Female	40 to 44	62% (46-79%)	38% (21-54%)	1.5 (0.64-2.7)	4.9% (2.5-8.3%)	7.2% (3.2-12%)	12 (5.3-21)	610 (300-1000)
Mongolia	1990	Female	45 to 49	61% (41-77%)	39% (23-59%)	1.6 (0.7-3)	4.9% (2.2-9.4%)	7% (2.9-13%)	15 (6.3-29)	670 (290-1300)
Mongolia	1990	Female	50 to 54	59% (41-75%)	41% (25-59%)	1.5 (0.66-3)	5.2% (2-10%)	7.1% (2.6-14%)	21 (7.7-42)	820 (310-1600)
Mongolia	1990	Female	55 to 59	56% (39-74%)	44% (26-61%)	1.5 (0.6-3.1)	5.1% (1.7-9.9%)	6.7% (2.2-13%)	26 (8.2-51)	850 (280-1700)
Mongolia	1990	Female	60 to 64	53% (35-71%)	47% (29-65%)	1.4 (0.56-3.1)	4.4% (1.2-9.5%)	5.6% (1.5-12%)	28 (8-60)	770 (220-1700)
Mongolia	1990	Female	65 to 69	50% (34-68%)	50% (32-66%)	1.3 (0.56-2.5)	3.8% (0.95-8.2%)	4.8% (1.2-10%)	28 (7.4-61)	640 (160-1400)
Mongolia	1990	Female	70 to 74	45% (30-62%)	55% (38-70%)	1.1 (0.42-2.2)	2.9% (0.34-7.4%)	3.6% (0.45-8.7%)	27 (3.5-67)	480 (58-1200)
Mongolia	1990	Female	75 to 79	40% (26-57%)	60% (43-74%)	0.92 (0.34-1.9)	2% (-0.49-5.9%)	2.3% (-0.58-7.1%)	15 (-3.5-45)	210 (-49-640)
Mongolia	1990	Female	80 to 84	37% (23-54%)	63% (46-77%)	0.92 (0.34-1.9)	1.3% (-1.1-5.2%)	1.5% (-1.4-6%)	7.6 (-6.6-30)	82 (-67-310)
Mongolia	1990	Female	85 to 89	36% (24-52%)	64% (48-76%)	0.98 (0.37-2.1)	0.91% (-1.5-5%)	0.98% (-1.9-5.8%)	3.7 (-6.9-22)	29 (-49-160)
Mongolia	1990	Female	90 to 94	38% (25-53%)	62% (47-75%)	1.1 (0.34-2.4)	0.73% (-2-5.8%)	0.75% (-2.4-6.5%)	1.1 (-3.4-9.3)	6.2 (-16-48)
Mongolia	1990	Female	95 plus	41% (25-60%)	59% (40-75%)	1.2 (0.29-3)	0.76% (-2.6-7%)	0.75% (-3.1-8.1%)	0.31 (-1.3-3.3)	1.2 (-4.1-11)
Mongolia	1990	Male	All Ages	51% (41-61%)	49% (39-59%)	1.1 (0.83-1.3)	3.2% (2.4-4.2%)	4.2% (2.9-5.9%)	430 (290-630)	19000 (14000-26000)
Mongolia	1990	Male	15 to 19	42% (26-60%)	58% (40-74%)	0.48 (0.24-0.83)	2.9% (1.6-4.6%)	3.1% (1.1-6%)	5.8 (2.1-11)	610 (330-990)
Mongolia	1990	Male	20 to 24	63% (44-79%)	37% (21-56%)	1.2 (0.74-1.9)	7.1% (4.7-10%)	7.8% (4.2-13%)	17 (9.1-28)	1600 (990-2400)
Mongolia	1990	Male	25 to 29	73% (56-85%)	27% (15-44%)	2.1 (1.3-3.1)	11% (7.4-15%)	12% (7.7-18%)	32 (19-46)	2500 (1600-3400)
Mongolia	1990	Male	30 to 34	79% (62-90%)	21% (10-38%)	2.6 (1.6-3.7)	12% (8.4-16%)	14% (9-20%)	32 (21-48)	2200 (1600-3100)
Mongolia	1990	Male	35 to 39	81% (68-91%)	19% (9.1-32%)	2.6 (1.6-4)	10% (7.2-15%)	12% (7.8-18%)	29 (18-45)	1800 (1200-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	1990	Male	40 to 44	81% (67-91%)	19% (8.8-33%)	2.6 (1.5-4)	9.9% (6.6-15%)	11% (7.1-18%)	29 (18-45)	1500 (1000-2300)
Mongolia	1990	Male	45 to 49	80% (63-91%)	20% (9.4-37%)	2.7 (1.5-4.2)	10% (6.7-14%)	12% (7.3-17%)	40 (24-61)	1800 (1200-2700)
Mongolia	1990	Male	50 to 54	79% (63-90%)	21% (10-37%)	2.7 (1.5-4.1)	9.6% (5.9-14%)	11% (6.5-16%)	53 (31-80)	2100 (1300-3100)
Mongolia	1990	Male	55 to 59	77% (61-89%)	23% (11-39%)	2.3 (1.2-3.6)	7.9% (4.9-12%)	8.9% (5.3-14%)	52 (31-82)	1800 (1100-2700)
Mongolia	1990	Male	60 to 64	74% (56-87%)	26% (13-44%)	1.9 (0.99-3.1)	6.3% (3.7-10%)	7.1% (4-11%)	51 (28-85)	1400 (820-2400)
Mongolia	1990	Male	65 to 69	71% (53-85%)	29% (15-47%)	1.6 (0.91-2.7)	4.8% (2.6-7.9%)	5.5% (2.8-9.1%)	42 (22-70)	970 (510-1600)
Mongolia	1990	Male	70 to 74	67% (49-82%)	33% (18-51%)	1.3 (0.73-2.3)	3.3% (1.4-5.9%)	3.7% (1.5-6.6%)	30 (12-56)	560 (230-1000)
Mongolia	1990	Male	75 to 79	63% (45-80%)	37% (20-55%)	1.1 (0.54-1.9)	2% (0.27-4.3%)	2.2% (0.19-4.8%)	14 (1.3-32)	210 (30-470)
Mongolia	1990	Male	80 to 84	60% (41-77%)	40% (23-59%)	0.96 (0.48-1.6)	1% (-0.53-2.9%)	1.1% (-0.76-3.2%)	4.4 (-3.3-14)	52 (-26-150)
Mongolia	1990	Male	85 to 89	58% (40-74%)	42% (26-60%)	0.9 (0.45-1.6)	0.4% (-1-2.1%)	0.29% (-1.4-2.2%)	0.71 (-3.5-5.4)	8.4 (-22-44)
Mongolia	1990	Male	90 to 94	57% (40-74%)	43% (26-60%)	0.85 (0.41-1.6)	-0.087% (-1.6-1.5%)	-0.32% (-2.2-1.5%)	-0.29 (-2-1.3)	-0.46 (-8.6-8)
Mongolia	1990	Male	95 plus	58% (36-77%)	42% (23-64%)	0.82 (0.33-1.6)	-0.32% (-2-1.3%)	-0.68% (-2.7-1.2%)	-0.16 (-0.61-0.28)	-0.3 (-1.8-1.2)
Mongolia	1995	Female	All Ages	44% (37-53%)	56% (47-63%)	0.71 (0.54-0.93)	2.1% (1.2-3.1%)	2.9% (1.2-5.4%)	250 (100-460)	9300 (5600-14000)
Mongolia	1995	Female	15 to 19	41% (27-59%)	59% (41-73%)	0.61 (0.25-1.1)	1.6% (0.56-3.1%)	2.4% (0.34-5.3%)	2.7 (0.38-6.2)	280 (96-530)
Mongolia	1995	Female	20 to 24	54% (37-70%)	46% (30-63%)	1 (0.43-1.8)	3% (1.3-5.3%)	3.9% (0.89-7.9%)	6.7 (1.5-14)	630 (260-1100)
Mongolia	1995	Female	25 to 29	59% (41-75%)	41% (25-59%)	1.2 (0.57-2.1)	3.8% (1.9-6.5%)	5.1% (1.7-9.6%)	9.5 (3.3-18)	790 (370-1300)
Mongolia	1995	Female	30 to 34	61% (42-77%)	39% (23-58%)	1.3 (0.67-2.3)	4.3% (2.4-7%)	6% (2.9-10%)	13 (6.1-23)	920 (520-1500)
Mongolia	1995	Female	35 to 39	62% (45-78%)	38% (22-55%)	1.4 (0.66-2.6)	4.4% (2.3-7.4%)	6.1% (2.7-11%)	14 (5.9-24)	820 (420-1400)
Mongolia	1995	Female	40 to 44	62% (45-78%)	38% (22-55%)	1.4 (0.61-2.6)	4.9% (2.4-8.3%)	6.9% (3.1-12%)	16 (7.3-29)	830 (400-1400)
Mongolia	1995	Female	45 to 49	61% (43-76%)	39% (24-57%)	1.5 (0.65-3)	5% (2-10%)	6.6% (2.4-13%)	18 (6.5-38)	780 (310-1600)
Mongolia	1995	Female	50 to 54	59% (42-75%)	41% (25-58%)	1.5 (0.62-2.9)	5.2% (1.8-11%)	6.6% (2.2-14%)	25 (8.2-52)	940 (330-1900)
Mongolia	1995	Female	55 to 59	56% (39-73%)	44% (27-61%)	1.4 (0.57-2.9)	5.1% (1.5-10%)	6.3% (1.8-13%)	29 (8.1-60)	930 (270-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	1995	Female	60 to 64	53% (35-71%)	47% (29-65%)	1.4 (0.53-2.9)	4.3% (1.1-9.5%)	5.2% (1.3-12%)	32 (8-71)	880 (220-2000)
Mongolia	1995	Female	65 to 69	51% (34-68%)	49% (32-66%)	1.2 (0.52-2.3)	3.6% (0.64-8.2%)	4.3% (0.83-9.8%)	32 (5.9-73)	710 (120-1700)
Mongolia	1995	Female	70 to 74	46% (30-62%)	54% (38-70%)	1 (0.38-2.1)	2.7% (0.091-7.1%)	3.3% (0.16-8.3%)	25 (1.1-63)	440 (15-1200)
Mongolia	1995	Female	75 to 79	40% (27-57%)	60% (43-73%)	0.87 (0.31-1.9)	1.7% (-0.59-5.7%)	2% (-0.7-6.6%)	18 (-6.3-58)	250 (-86-820)
Mongolia	1995	Female	80 to 84	37% (24-54%)	63% (46-76%)	0.86 (0.32-1.8)	1.1% (-1.3-5%)	1.3% (-1.6-5.7%)	7.3 (-9.1-32)	78 (-91-340)
Mongolia	1995	Female	85 to 89	37% (24-51%)	63% (49-76%)	0.93 (0.35-2)	0.71% (-1.7-4.9%)	0.75% (-2.1-5.5%)	2.9 (-8.2-22)	24 (-58-160)
Mongolia	1995	Female	90 to 94	38% (25-53%)	62% (47-75%)	1 (0.32-2.4)	0.54% (-2.1-5.8%)	0.53% (-2.6-6.6%)	1 (-4.8-12)	6 (-23-63)
Mongolia	1995	Female	95 plus	41% (25-60%)	59% (40-75%)	1.2 (0.28-2.9)	0.58% (-2.7-6.8%)	0.54% (-3.3-7.8%)	0.26 (-1.6-3.8)	1.1 (-5.2-13)
Mongolia	1995	Male	All Ages	52% (42-63%)	48% (37-58%)	1 (0.81-1.3)	4.3% (3.3-5.6%)	4.9% (3.4-6.9%)	520 (350-720)	24000 (18000-31000)
Mongolia	1995	Male	15 to 19	42% (26-61%)	58% (39-74%)	0.46 (0.23-0.81)	2.9% (1.5-4.5%)	3.2% (1.1-5.9%)	5.9 (2.2-11)	620 (330-990)
Mongolia	1995	Male	20 to 24	63% (44-79%)	37% (21-56%)	1.2 (0.68-1.8)	7.1% (4.5-11%)	7.6% (4-13%)	23 (12-38)	2000 (1200-3000)
Mongolia	1995	Male	25 to 29	74% (57-85%)	26% (15-43%)	2 (1.2-2.9)	11% (7.4-15%)	12% (7.5-18%)	42 (26-61)	3200 (2200-4300)
Mongolia	1995	Male	30 to 34	79% (63-90%)	21% (9.6-37%)	2.4 (1.6-3.5)	12% (8.5-16%)	13% (9-19%)	53 (35-75)	3600 (2500-4900)
Mongolia	1995	Male	35 to 39	81% (68-91%)	19% (9-32%)	2.4 (1.4-3.6)	10% (7.4-15%)	12% (7.6-17%)	43 (27-64)	2600 (1800-3700)
Mongolia	1995	Male	40 to 44	82% (69-91%)	18% (8.6-31%)	2.4 (1.4-3.7)	9.8% (6.6-15%)	11% (6.9-17%)	43 (26-68)	2300 (1500-3400)
Mongolia	1995	Male	45 to 49	81% (64-91%)	19% (9.1-36%)	2.6 (1.4-4)	10% (6.4-15%)	11% (6.7-17%)	48 (28-72)	2200 (1300-3100)
Mongolia	1995	Male	50 to 54	79% (64-90%)	21% (10-36%)	2.5 (1.4-3.9)	9.4% (5.7-14%)	10% (6.1-16%)	57 (33-88)	2200 (1300-3300)
Mongolia	1995	Male	55 to 59	77% (61-89%)	23% (11-39%)	2.1 (1.1-3.4)	7.5% (4.7-12%)	8.3% (4.9-13%)	55 (32-87)	1900 (1100-2800)
Mongolia	1995	Male	60 to 64	74% (57-87%)	26% (13-43%)	1.8 (0.91-3)	6% (3.2-9.8%)	6.5% (3.4-11%)	56 (28-95)	1600 (830-2600)
Mongolia	1995	Male	65 to 69	72% (54-86%)	28% (14-46%)	1.5 (0.84-2.5)	4.5% (2.3-7.5%)	5% (2.5-8.4%)	44 (22-76)	1000 (510-1700)
Mongolia	1995	Male	70 to 74	68% (50-82%)	32% (18-50%)	1.3 (0.69-2.2)	3% (1.2-5.6%)	3.3% (1.2-6.2%)	28 (9.6-52)	520 (190-960)
Mongolia	1995	Male	75 to 79	64% (46-79%)	36% (21-54%)	1 (0.52-1.8)	1.8% (0.13-4%)	1.9% (0.033-4.4%)	15 (0.26-35)	230 (16-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	1995	Male	80 to 84	61% (42-77%)	39% (23-58%)	0.92 (0.46-1.5)	0.91% (-0.69-2.7%)	0.89% (-0.98-3%)	4.1 (-4.5-14)	49 (-38-150)
Mongolia	1995	Male	85 to 89	58% (40-75%)	42% (25-60%)	0.85 (0.43-1.5)	0.3% (-1.1-1.9%)	0.18% (-1.5-2%)	0.47 (-4.1-5.2)	6.8 (-26-42)
Mongolia	1995	Male	90 to 94	58% (40-74%)	42% (26-60%)	0.8 (0.38-1.4)	-0.14% (-1.7-1.4%)	-0.38% (-2.2-1.4%)	-0.39 (-2.3-1.4)	-0.87 (-10-8.3)
Mongolia	1995	Male	95 plus	59% (36-78%)	41% (22-64%)	0.78 (0.32-1.5)	-0.38% (-1.9-1.2%)	-0.73% (-2.6-1.1%)	-0.19 (-0.66-0.28)	-0.38 (-1.9-1.2)
Mongolia	2000	Female	All Ages	47% (39-56%)	53% (44-61%)	0.82 (0.62-1.1)	3.1% (1.9-4.5%)	4.2% (1.9-7.3%)	340 (160-600)	13000 (8000-18000)
Mongolia	2000	Female	15 to 19	43% (28-60%)	57% (40-72%)	0.67 (0.27-1.3)	1.8% (0.64-3.4%)	2.9% (0.43-6.3%)	3.1 (0.45-7)	310 (110-600)
Mongolia	2000	Female	20 to 24	56% (39-71%)	44% (29-61%)	1.1 (0.46-1.9)	3.5% (1.5-6.4%)	4.8% (1.3-9.6%)	8 (2.1-16)	730 (310-1300)
Mongolia	2000	Female	25 to 29	60% (42-76%)	40% (24-58%)	1.3 (0.58-2.3)	4.7% (2.4-7.5%)	6.3% (2.4-11%)	12 (4.5-21)	1000 (490-1600)
Mongolia	2000	Female	30 to 34	63% (42-78%)	37% (22-58%)	1.4 (0.75-2.4)	5.4% (3.3-8.2%)	7.5% (3.9-12%)	16 (8.5-27)	1200 (700-1800)
Mongolia	2000	Female	35 to 39	63% (47-79%)	37% (21-53%)	1.5 (0.71-2.7)	5.4% (3-8.8%)	7.5% (3.6-12%)	23 (11-38)	1400 (720-2200)
Mongolia	2000	Female	40 to 44	63% (47-79%)	37% (21-53%)	1.6 (0.67-2.7)	5.9% (3-9.7%)	8.1% (3.8-14%)	27 (12-46)	1400 (690-2300)
Mongolia	2000	Female	45 to 49	62% (44-78%)	38% (22-56%)	1.6 (0.7-3.2)	6% (2.5-11%)	7.8% (2.9-15%)	28 (10-54)	1200 (480-2300)
Mongolia	2000	Female	50 to 54	60% (42-75%)	40% (25-58%)	1.6 (0.68-3.2)	6.4% (2.3-13%)	8% (2.8-16%)	30 (11-60)	1100 (420-2200)
Mongolia	2000	Female	55 to 59	57% (40-74%)	43% (26-60%)	1.6 (0.61-3.2)	6.3% (1.9-12%)	7.8% (2.4-15%)	38 (11-76)	1200 (380-2400)
Mongolia	2000	Female	60 to 64	55% (37-72%)	45% (28-63%)	1.5 (0.57-3.1)	5.5% (1.6-12%)	6.6% (1.9-14%)	40 (11-85)	1100 (300-2300)
Mongolia	2000	Female	65 to 69	52% (36-70%)	48% (30-64%)	1.3 (0.58-2.6)	4.5% (1.1-9.9%)	5.4% (1.4-12%)	42 (11-92)	940 (240-2100)
Mongolia	2000	Female	70 to 74	47% (32-63%)	53% (37-68%)	1.1 (0.42-2.3)	3.5% (0.28-8.6%)	4.1% (0.4-10%)	34 (3.3-84)	600 (50-1500)
Mongolia	2000	Female	75 to 79	42% (28-59%)	58% (41-72%)	0.97 (0.34-2.1)	2.4% (-0.42-7.2%)	2.8% (-0.49-8.3%)	21 (-3.6-63)	290 (-52-880)
Mongolia	2000	Female	80 to 84	39% (25-55%)	61% (45-75%)	0.97 (0.37-2)	1.8% (-1.1-6.5%)	2% (-1.3-7.4%)	12 (-7.7-45)	130 (-78-460)
Mongolia	2000	Female	85 to 89	38% (26-54%)	62% (46-74%)	1 (0.39-2.2)	1.3% (-1.5-6.2%)	1.5% (-1.8-7.1%)	6 (-7.4-29)	46 (-53-210)
Mongolia	2000	Female	90 to 94	40% (26-55%)	60% (45-74%)	1.1 (0.37-2.7)	1.1% (-2.7-2%)	1.2% (-2.5-8.3%)	2.1 (-4.4-15)	11 (-21-75)
Mongolia	2000	Female	95 plus	43% (26-61%)	57% (39-74%)	1.3 (0.31-3.3)	1.1% (-2.7-7.9%)	1.1% (-3.3-8.9%)	0.66 (-2.5-4)	2.5 (-6.4-19)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	2000	Male	All Ages	56% (46-66%)	44% (34-54%)	1.2 (0.98-1.5)	6% (4.7-7.5%)	6.5% (4.6-8.7%)	660 (460-890)	31000 (24000-39000)
Mongolia	2000	Male	15 to 19	44% (28-63%)	56% (37-72%)	0.51 (0.26-0.91)	3.2% (1.9-5%)	3.7% (1.5-6.8%)	6.7 (2.7-12)	730 (420-1100)
Mongolia	2000	Male	20 to 24	65% (47-80%)	35% (20-53%)	1.3 (0.75-2)	8.1% (5.2-12%)	8.7% (4.7-14%)	25 (13-40)	2300 (1400-3300)
Mongolia	2000	Male	25 to 29	75% (59-86%)	25% (14-41%)	2.2 (1.3-3.2)	13% (8.5-17%)	14% (8.4-20%)	51 (31-73)	3900 (2700-5300)
Mongolia	2000	Male	30 to 34	80% (64-91%)	20% (9.1-36%)	2.7 (1.7-3.8)	14% (10-18%)	15% (11-21%)	63 (43-87)	4300 (3200-5700)
Mongolia	2000	Male	35 to 39	82% (69-92%)	18% (8.4-31%)	2.6 (1.6-3.8)	12% (8.9-16%)	14% (9.4-19%)	71 (48-100)	4300 (3100-5800)
Mongolia	2000	Male	40 to 44	82% (69-92%)	18% (8.3-31%)	2.6 (1.5-3.9)	12% (8-16%)	13% (8.3-18%)	69 (44-100)	3600 (2400-5000)
Mongolia	2000	Male	45 to 49	82% (65-91%)	18% (9-35%)	2.8 (1.6-4.4)	12% (7.6-16%)	13% (8-19%)	73 (46-110)	3300 (2100-4700)
Mongolia	2000	Male	50 to 54	80% (64-90%)	20% (9.8-36%)	2.8 (1.5-4.3)	11% (6.5-15%)	12% (6.8-17%)	67 (39-100)	2600 (1600-3800)
Mongolia	2000	Male	55 to 59	79% (63-90%)	21% (9.9-37%)	2.4 (1.2-3.8)	8.5% (5.1-13%)	9.3% (5.4-14%)	65 (37-100)	2200 (1300-3300)
Mongolia	2000	Male	60 to 64	76% (57-88%)	24% (12-43%)	2 (1-3.3)	6.7% (3.6-11%)	7.3% (3.7-13%)	64 (32-110)	1800 (920-3000)
Mongolia	2000	Male	65 to 69	73% (55-87%)	27% (13-45%)	1.7 (0.96-2.8)	5.1% (2.5-8.7%)	5.6% (2.6-9.8%)	53 (25-91)	1200 (600-2100)
Mongolia	2000	Male	70 to 74	70% (52-83%)	30% (17-48%)	1.4 (0.75-2.4)	3.5% (1.2-6.7%)	3.8% (1.2-7.4%)	33 (10-63)	610 (200-1200)
Mongolia	2000	Male	75 to 79	66% (47-81%)	34% (19-53%)	1.2 (0.58-2)	2.2% (0.2-5%)	2.4% (0.11-5.4%)	17 (0.82-38)	250 (22-550)
Mongolia	2000	Male	80 to 84	63% (44-78%)	37% (22-56%)	1 (0.51-1.7)	1.3% (-0.63-3.4%)	1.3% (-0.88-3.6%)	5.7 (-4-16)	67 (-33-180)
Mongolia	2000	Male	85 to 89	60% (42-76%)	40% (24-58%)	0.95 (0.48-1.6)	0.59% (-1-2.5%)	0.49% (-1.5-2.7%)	1.4 (-4-7.4)	14 (-25-58)
Mongolia	2000	Male	90 to 94	59% (41-76%)	41% (24-59%)	0.89 (0.42-1.6)	0.032% (-1.6-1.9%)	-0.21% (-2.2-1.9%)	-0.22 (-2.4-2.1)	0.2 (-11-12)
Mongolia	2000	Male	95 plus	60% (38-78%)	40% (22-62%)	0.85 (0.35-1.6)	-0.27% (-2-1.6%)	-0.64% (-2.6-1.5%)	-0.19 (-0.76-0.42)	-0.3 (-2.2-1.8)
Mongolia	2005	Female	All Ages	51% (44-60%)	49% (40-56%)	1.1 (0.83-1.4)	5% (3.4-6.8%)	7.2% (4-11%)	570 (310-890)	19000 (13000-27000)
Mongolia	2005	Female	15 to 19	46% (31-62%)	54% (38-69%)	0.83 (0.35-1.5)	2.2% (0.86-4%)	4.2% (0.94-8.8%)	4 (0.87-8.3)	400 (160-750)
Mongolia	2005	Female	20 to 24	58% (42-73%)	42% (27-58%)	1.3 (0.56-2.3)	4.5% (2.2-7.6%)	7.3% (2.6-14%)	9.3 (3.3-17)	860 (430-1500)
Mongolia	2005	Female	25 to 29	62% (44-77%)	38% (23-56%)	1.6 (0.69-2.8)	6.4% (3.6-10%)	9.8% (4.4-16%)	16 (7.1-27)	1300 (700-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	2005	Female	30 to 34	64% (45-79%)	36% (21-55%)	1.7 (0.89-2.9)	7.4% (4.8-11%)	12% (6.8-18%)	22 (13-34)	1600 (1000-2300)
Mongolia	2005	Female	35 to 39	65% (49-80%)	35% (20-51%)	1.8 (0.89-3.2)	7.6% (4.7-11%)	12% (6.7-18%)	30 (16-46)	1800 (1100-2700)
Mongolia	2005	Female	40 to 44	65% (49-80%)	35% (20-51%)	1.9 (0.86-3.2)	8.3% (4.8-13%)	12% (6.6-19%)	45 (25-72)	2300 (1300-3600)
Mongolia	2005	Female	45 to 49	64% (46-79%)	36% (21-54%)	1.9 (0.88-3.7)	8.4% (3.8-15%)	12% (4.9-20%)	50 (22-90)	2200 (1000-3900)
Mongolia	2005	Female	50 to 54	62% (45-77%)	38% (23-55%)	1.9 (0.86-3.8)	9.2% (3.9-16%)	12% (5-21%)	55 (22-94)	2000 (850-3500)
Mongolia	2005	Female	55 to 59	59% (43-75%)	41% (25-57%)	1.9 (0.77-3.8)	9.2% (3.5-16%)	12% (4.4-20%)	55 (21-95)	1800 (670-3100)
Mongolia	2005	Female	60 to 64	57% (41-73%)	43% (27-59%)	1.8 (0.71-3.9)	8.4% (3.3-15%)	10% (4.1-19%)	61 (24-110)	1700 (620-2900)
Mongolia	2005	Female	65 to 69	55% (39-71%)	45% (29-61%)	1.7 (0.73-3.3)	7% (2.6-13%)	8.6% (3.2-16%)	61 (23-110)	1400 (490-2600)
Mongolia	2005	Female	70 to 74	51% (35-67%)	49% (33-65%)	1.4 (0.53-3)	5.8% (1.5-12%)	7.1% (1.9-15%)	60 (16-120)	1100 (290-2200)
Mongolia	2005	Female	75 to 79	45% (31-63%)	55% (37-69%)	1.3 (0.44-2.7)	4.5% (0.67-11%)	5.4% (0.8-12%)	42 (6.6-97)	590 (86-1400)
Mongolia	2005	Female	80 to 84	43% (27-60%)	57% (40-73%)	1.3 (0.47-2.5)	4.1% (0.21-10%)	4.8% (0.26-12%)	29 (1.6-69)	300 (15-720)
Mongolia	2005	Female	85 to 89	42% (28-57%)	58% (43-72%)	1.3 (0.5-2.8)	3.6% (-0.48-10%)	4.2% (-0.57-12%)	18 (-2.5-51)	130 (-17-370)
Mongolia	2005	Female	90 to 94	43% (30-58%)	57% (42-70%)	1.5 (0.46-3.4)	3% (-1.4-10%)	3.4% (-1.6-12%)	6.5 (-3.1-23)	34 (-15-120)
Mongolia	2005	Female	95 plus	46% (30-63%)	54% (37-70%)	1.6 (0.4-4.1)	2.7% (-2.2-11%)	3% (-2.7-12%)	1.7 (-1.5-7.1)	6 (-4.9-24)
Mongolia	2005	Male	All Ages	62% (53-71%)	38% (29-47%)	1.7 (1.3-2)	10% (7.8-12%)	11% (8-14%)	1300 (890-1600)	57000 (43000-70000)
Mongolia	2005	Male	15 to 19	48% (31-65%)	52% (35-69%)	0.67 (0.33-1.1)	4.6% (2.8-6.9%)	5.8% (2.7-9.6%)	12 (5.4-20)	1200 (700-1800)
Mongolia	2005	Male	20 to 24	68% (50-83%)	32% (17-50%)	1.6 (0.97-2.5)	11% (7.3-15%)	12% (7.3-18%)	38 (22-57)	3300 (2200-4600)
Mongolia	2005	Male	25 to 29	77% (62-88%)	23% (12-38%)	2.7 (1.6-3.9)	16% (11-21%)	19% (12-25%)	76 (50-110)	5600 (4000-7400)
Mongolia	2005	Male	30 to 34	82% (68-91%)	18% (8.5-32%)	3.3 (2.1-4.6)	19% (14-24%)	21% (15-28%)	110 (77-150)	7400 (5300-9600)
Mongolia	2005	Male	35 to 39	84% (72-92%)	16% (7.9-28%)	3.2 (2-4.6)	18% (13-24%)	21% (14-27%)	140 (88-190)	7700 (5300-10000)
Mongolia	2005	Male	40 to 44	84% (72-92%)	16% (7.5-28%)	3.2 (1.9-4.7)	17% (12-22%)	19% (13-25%)	170 (110-230)	8300 (5600-11000)
Mongolia	2005	Male	45 to 49	83% (69-92%)	17% (7.8-31%)	3.5 (2-5.2)	17% (11-22%)	18% (12-25%)	170 (110-240)	7600 (5000-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	2005	Male	50 to 54	82% (68-91%)	18% (8.9-32%)	3.4 (1.9-5.2)	15% (9.2-21%)	16% (9.8-23%)	150 (87-210)	5500 (3400-7700)
Mongolia	2005	Male	55 to 59	81% (66-91%)	19% (9.2-34%)	2.9 (1.6-4.5)	12% (7.2-18%)	13% (7.6-20%)	110 (61-160)	3500 (2100-5200)
Mongolia	2005	Male	60 to 64	78% (62-90%)	22% (10-38%)	2.4 (1.3-4.1)	9.4% (4.9-15%)	10% (5.2-17%)	99 (51-160)	2800 (1500-4500)
Mongolia	2005	Male	65 to 69	75% (59-88%)	25% (12-41%)	2.1 (1.2-3.4)	7.2% (3.6-12%)	7.9% (3.8-13%)	82 (40-140)	1900 (940-3100)
Mongolia	2005	Male	70 to 74	72% (55-85%)	28% (15-45%)	1.8 (0.98-2.9)	5.4% (2.1-10%)	5.9% (2.3-11%)	57 (23-110)	1100 (430-2000)
Mongolia	2005	Male	75 to 79	68% (51-83%)	32% (17-49%)	1.5 (0.76-2.5)	4% (1.2-8.8%)	4.4% (1.2-8.8%)	32 (8.7-65)	470 (140-930)
Mongolia	2005	Male	80 to 84	65% (47-80%)	35% (20-53%)	1.3 (0.66-2.1)	3% (0.48-6.2%)	3.2% (0.33-6.8%)	14 (1.5-30)	160 (27-320)
Mongolia	2005	Male	85 to 89	63% (45-78%)	37% (22-55%)	1.2 (0.61-2)	2% (-0.31-5.1%)	2% (-0.58-5.6%)	5.6 (-1.6-15)	46 (-7.1-120)
Mongolia	2005	Male	90 to 94	62% (45-78%)	38% (22-55%)	1.1 (0.54-2)	1% (-1-3.7%)	0.89% (-1.5-3.9%)	0.99 (-1.6-4.4)	6.5 (-6.8-24)
Mongolia	2005	Male	95 plus	63% (41-80%)	37% (20-59%)	1.1 (0.44-2)	0.44% (-1.8-3.3%)	0.14% (-2.4-3.4%)	0.042 (-0.75-1.1)	0.53 (-2.1-4)
Mongolia	2010	Female	All Ages	53% (45-60%)	47% (40-55%)	1.3 (0.97-1.7)	6.2% (4.3-8.3%)	9.2% (5.6-13%)	730 (440-1100)	25000 (17000-33000)
Mongolia	2010	Female	15 to 19	47% (32-63%)	53% (37-68%)	0.97 (0.38-1.8)	2.5% (1-4.6%)	5.1% (1.3-10%)	4 (1.1-8.2)	390 (150-750)
Mongolia	2010	Female	20 to 24	59% (44-74%)	41% (26-56%)	1.5 (0.64-2.8)	5% (2.6-8.3%)	8.7% (3.4-15%)	12 (4.8-22)	1100 (590-1900)
Mongolia	2010	Female	25 to 29	63% (45-77%)	37% (23-55%)	1.8 (0.83-3.3)	7% (4.2-10%)	12% (5.9-19%)	17 (8.6-28)	1400 (840-2200)
Mongolia	2010	Female	30 to 34	65% (47-79%)	35% (21-53%)	2 (1-3.4)	8.2% (5.6-12%)	14% (8.7-20%)	25 (15-38)	1800 (1200-2600)
Mongolia	2010	Female	35 to 39	66% (50-80%)	34% (20-50%)	2 (0.97-3.6)	8.8% (5.6-13%)	14% (8.2-20%)	36 (21-55)	2200 (1300-3200)
Mongolia	2010	Female	40 to 44	65% (50-80%)	35% (20-50%)	2.1 (0.96-3.8)	9.8% (5.9-14%)	15% (8.6-22%)	52 (30-81)	2700 (1500-4000)
Mongolia	2010	Female	45 to 49	64% (46-79%)	36% (21-54%)	2.2 (1-4.2)	9.7% (5-15%)	14% (6.9-22%)	69 (33-120)	3100 (1600-5000)
Mongolia	2010	Female	50 to 54	62% (46-77%)	38% (23-54%)	2.2 (1-4.2)	11% (4.9-17%)	14% (6.5-23%)	80 (36-130)	3000 (1400-5000)
Mongolia	2010	Female	55 to 59	60% (43-75%)	40% (25-57%)	2.2 (0.85-4.3)	10% (4.4-17%)	14% (6-22%)	72 (31-120)	2300 (1000-3800)
Mongolia	2010	Female	60 to 64	58% (41-75%)	42% (25-59%)	2.1 (0.82-4.5)	9.8% (4.4-16%)	13% (5.8-21%)	68 (31-110)	1800 (830-3100)
Mongolia	2010	Female	65 to 69	56% (40-72%)	44% (28-60%)	1.9 (0.83-3.8)	8.6% (3.7-15%)	10% (4.6-19%)	84 (36-150)	1900 (800-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	2010	Female	70 to 74	51% (36-67%)	49% (33-64%)	1.6 (0.61-3.4)	7.2% (2.7-14%)	9% (3.3-17%)	71 (27-130)	1300 (460-2500)
Mongolia	2010	Female	75 to 79	46% (32-63%)	54% (37-68%)	1.4 (0.52-3.1)	5.8% (1.5-12%)	7.1% (1.9-15%)	57 (15-120)	800 (200-1600)
Mongolia	2010	Female	80 to 84	43% (29-61%)	57% (39-71%)	1.5 (0.54-3)	5.5% (1.2-12%)	6.7% (1.4-14%)	45 (9.6-95)	470 (97-990)
Mongolia	2010	Female	85 to 89	43% (29-57%)	57% (43-71%)	1.5 (0.56-3.3)	5% (0.44-13%)	5.9% (0.41-15%)	25 (1.7-63)	180 (16-460)
Mongolia	2010	Female	90 to 94	44% (30-59%)	56% (41-70%)	1.7 (0.53-3.9)	4.2% (-0.67-12%)	4.9% (-0.95-14%)	8.7 (-1.6-24)	44 (-7-120)
Mongolia	2010	Female	95 plus	47% (30-64%)	53% (36-70%)	1.9 (0.47-4.6)	3.7% (-1.9-12%)	4.2% (-2.3-14%)	2.4 (-1.2-7.9)	8.3 (-4-26)
Mongolia	2010	Male	All Ages	64% (56-73%)	36% (27-44%)	2 (1.6-2.5)	12% (9.2-15%)	14% (9.8-17%)	1600 (1100-2100)	71000 (53000-90000)
Mongolia	2010	Male	15 to 19	49% (32-67%)	51% (33-68%)	0.75 (0.37-1.3)	5.3% (3.2-7.9%)	6.9% (3.4-11%)	12 (5.6-20)	1200 (700-1800)
Mongolia	2010	Male	20 to 24	69% (51-83%)	31% (17-49%)	1.8 (1.1-2.9)	12% (8.3-16%)	14% (8.4-20%)	47 (27-70)	4100 (2700-5700)
Mongolia	2010	Male	25 to 29	78% (63-88%)	22% (12-37%)	3.1 (1.9-4.6)	18% (13-24%)	21% (14-29%)	84 (54-120)	6400 (4400-8400)
Mongolia	2010	Male	30 to 34	83% (68-92%)	17% (8.3-32%)	3.7 (2.4-5.2)	21% (15-27%)	25% (17-33%)	130 (84-180)	8500 (5900-11000)
Mongolia	2010	Male	35 to 39	84% (72-92%)	16% (7.7-28%)	3.6 (2.3-5.3)	22% (14-29%)	25% (16-34%)	170 (100-230)	9700 (6400-13000)
Mongolia	2010	Male	40 to 44	85% (73-92%)	15% (7.5-27%)	3.7 (2.2-5.4)	20% (14-26%)	23% (15-30%)	190 (120-260)	9600 (6500-13000)
Mongolia	2010	Male	45 to 49	84% (69-92%)	16% (8-31%)	3.9 (2.3-6)	20% (14-26%)	22% (15-29%)	240 (160-340)	11000 (7300-15000)
Mongolia	2010	Male	50 to 54	83% (69-92%)	17% (8.4-31%)	3.9 (2.1-5.8)	17% (11-23%)	19% (12-26%)	220 (140-310)	8400 (5400-12000)
Mongolia	2010	Male	55 to 59	81% (67-91%)	19% (9-33%)	3.3 (1.8-5.1)	14% (9.2-20%)	16% (10-22%)	150 (93-220)	5000 (3100-7200)
Mongolia	2010	Male	60 to 64	79% (62-90%)	21% (9.9-38%)	2.8 (1.5-4.6)	11% (6.3-17%)	13% (6.8-19%)	110 (58-170)	3000 (1700-4600)
Mongolia	2010	Male	65 to 69	76% (60-89%)	24% (11-40%)	2.5 (1.4-3.9)	8.8% (4.5-14%)	9.8% (4.9-16%)	100 (52-170)	2400 (1200-3900)
Mongolia	2010	Male	70 to 74	73% (57-85%)	27% (15-43%)	2 (1.1-3.4)	6.8% (3.1-12%)	7.7% (3.4-13%)	73 (33-130)	1300 (610-2300)
Mongolia	2010	Male	75 to 79	69% (52-83%)	31% (17-48%)	1.7 (0.85-2.9)	5.4% (2.2-9.9%)	6.1% (2.3-11%)	43 (16-79)	620 (250-1100)
Mongolia	2010	Male	80 to 84	67% (48-81%)	33% (19-52%)	1.5 (0.76-2.5)	4.3% (1.5-8.3%)	4.8% (1.6-9.3%)	22 (7.4-43)	240 (88-460)
Mongolia	2010	Male	85 to 89	64% (46-79%)	36% (21-54%)	1.4 (0.7-2.3)	3% (0.56-6.5%)	3.3% (0.37-7.4%)	8.4 (0.94-18)	66 (12-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	2010	Male	90 to 94	63% (46-78%)	37% (22-54%)	1.3 (0.61-2.3)	1.8% (-0.53-5.1%)	1.8% (-1.1-5.6%)	1.5 (-0.9-4.9)	9.1 (-2.7-26)
Mongolia	2010	Male	95 plus	63% (42-81%)	37% (19-58%)	1.2 (0.5-2.2)	1.1% (-1.4-4.6%)	0.82% (-2.1-4.9%)	0.22 (-0.56-1.3)	1.1 (-1.5-4.8)
Mongolia	2016	Female	All Ages	52% (45-60%)	48% (40-55%)	1.4 (0.93-1.9)	6.5% (4.4-8.8%)	9.7% (5.7-14%)	810 (460-1200)	26000 (17000-36000)
Mongolia	2016	Female	15 to 19	46% (31-62%)	54% (38-69%)	1 (0.38-2)	2.5% (1.1-4.9%)	5.3% (1.6-11%)	3.2 (0.95-6.7)	320 (130-590)
Mongolia	2016	Female	20 to 24	59% (42-73%)	41% (27-58%)	1.6 (0.65-3.1)	4.9% (2.6-8%)	8.9% (3.6-16%)	8.4 (3.2-15)	820 (420-1300)
Mongolia	2016	Female	25 to 29	63% (45-77%)	37% (23-55%)	1.9 (0.82-3.5)	6.6% (4-10%)	12% (5.8-19%)	17 (8-29)	1500 (850-2300)
Mongolia	2016	Female	30 to 34	65% (46-79%)	35% (21-54%)	2.1 (1-3.6)	7.8% (5.2-11%)	14% (8.4-21%)	24 (14-38)	1800 (1200-2700)
Mongolia	2016	Female	35 to 39	65% (50-79%)	35% (21-50%)	2.2 (0.99-4)	8.5% (5.4-12%)	14% (8.4-21%)	34 (18-54)	2100 (1300-3200)
Mongolia	2016	Female	40 to 44	65% (49-80%)	35% (20-51%)	2.2 (0.93-4.1)	9.4% (5.7-14%)	15% (8.7-22%)	50 (27-79)	2600 (1500-4000)
Mongolia	2016	Female	45 to 49	64% (46-79%)	36% (21-54%)	2.3 (1-4.5)	9.3% (4.9-15%)	14% (7.1-22%)	63 (31-100)	2800 (1500-4500)
Mongolia	2016	Female	50 to 54	62% (45-76%)	38% (24-55%)	2.3 (1-4.5)	10% (5-16%)	14% (6.9-23%)	92 (43-150)	3500 (1600-5600)
Mongolia	2016	Female	55 to 59	60% (43-76%)	40% (24-57%)	2.3 (0.9-4.4)	10% (4.7-16%)	14% (6.5-22%)	97 (43-160)	3200 (1400-5100)
Mongolia	2016	Female	60 to 64	58% (40-74%)	42% (26-60%)	2.2 (0.84-4.5)	9.8% (4.7-16%)	13% (6.5-21%)	88 (41-140)	2400 (1100-3900)
Mongolia	2016	Female	65 to 69	55% (39-72%)	45% (28-61%)	2 (0.85-4)	8.7% (4.2-14%)	12% (5.7-19%)	74 (36-120)	1700 (790-2800)
Mongolia	2016	Female	70 to 74	51% (36-67%)	49% (33-64%)	1.7 (0.63-3.7)	7.6% (3-14%)	9.8% (4.2-18%)	74 (31-130)	1300 (540-2500)
Mongolia	2016	Female	75 to 79	46% (31-63%)	54% (37-69%)	1.5 (0.55-3.4)	6.1% (1.7-12%)	7.7% (2.3-15%)	63 (19-130)	890 (270-1800)
Mongolia	2016	Female	80 to 84	43% (28-59%)	57% (41-72%)	1.6 (0.57-3.2)	5.8% (1.2-12%)	7.1% (1.6-15%)	61 (13-130)	640 (140-1300)
Mongolia	2016	Female	85 to 89	42% (28-56%)	58% (44-72%)	1.6 (0.59-3.6)	5.3% (0.58-12%)	6.3% (0.71-15%)	39 (4.6-92)	290 (33-680)
Mongolia	2016	Female	90 to 94	43% (30-59%)	57% (41-70%)	1.8 (0.56-4.3)	4.6% (-0.42-12%)	5.3% (-0.56-14%)	15 (-1.5-40)	76 (-6.9-200)
Mongolia	2016	Female	95 plus	46% (30-64%)	54% (36-70%)	2 (0.49-4.9)	4.1% (-1.7-12%)	4.6% (-2.1-14%)	3.4 (-1.5-10)	12 (-4.7-34)
Mongolia	2016	Male	All Ages	64% (56-73%)	36% (27-44%)	2.2 (1.6-3)	13% (9.9-16%)	14% (10-19%)	1900 (1300-2500)	81000 (58000-110000)
Mongolia	2016	Male	15 to 19	48% (31-66%)	52% (34-69%)	0.81 (0.36-1.5)	5.5% (3.4-8.4%)	7.2% (3.6-12%)	10 (4.9-17)	990 (590-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mongolia	2016	Male	20 to 24	69% (50-83%)	31% (17-50%)	2 (1.1-3.1)	13% (8.3-18%)	14% (8.4-21%)	42 (24-64)	3700 (2300-5300)
Mongolia	2016	Male	25 to 29	78% (63-88%)	22% (12-37%)	3.3 (1.9-5.1)	19% (13-25%)	22% (14-31%)	100 (62-140)	7600 (5200-10000)
Mongolia	2016	Male	30 to 34	83% (69-92%)	17% (8.3-31%)	4 (2.4-6)	22% (15-28%)	26% (17-34%)	140 (89-200)	9500 (6500-13000)
Mongolia	2016	Male	35 to 39	84% (72-92%)	16% (8-28%)	3.9 (2.3-6.1)	22% (15-29%)	26% (16-35%)	170 (110-250)	10000 (6700-14000)
Mongolia	2016	Male	40 to 44	84% (73-92%)	16% (7.6-27%)	3.9 (2.3-6.2)	21% (14-27%)	24% (16-32%)	210 (130-300)	11000 (7100-15000)
Mongolia	2016	Male	45 to 49	84% (69-92%)	16% (7.8-31%)	4.2 (2.4-6.7)	20% (14-26%)	23% (15-30%)	250 (150-360)	11000 (7300-15000)
Mongolia	2016	Male	50 to 54	82% (69-91%)	18% (8.7-31%)	4.1 (2.2-6.5)	18% (12-24%)	20% (13-27%)	270 (160-390)	11000 (6500-15000)
Mongolia	2016	Male	55 to 59	81% (67-91%)	19% (9.4-33%)	3.5 (1.9-5.6)	15% (9.6-21%)	17% (11-23%)	220 (130-330)	7300 (4500-11000)
Mongolia	2016	Male	60 to 64	78% (61-90%)	22% (10-39%)	3 (1.5-5)	12% (6.9-18%)	13% (7.6-21%)	160 (86-250)	4400 (2500-6800)
Mongolia	2016	Male	65 to 69	76% (59-89%)	24% (11-41%)	2.6 (1.4-4.3)	9.7% (5.3-15%)	11% (5.9-18%)	100 (55-170)	2400 (1300-4000)
Mongolia	2016	Male	70 to 74	72% (55-85%)	28% (15-45%)	2.2 (1.2-3.7)	7.6% (3.7-13%)	8.6% (4.3-15%)	84 (40-140)	1600 (760-2700)
Mongolia	2016	Male	75 to 79	68% (50-83%)	32% (17-50%)	1.8 (0.84-3.2)	6% (2.6-11%)	6.8% (2.9-13%)	61 (26-110)	880 (380-1600)
Mongolia	2016	Male	80 to 84	66% (47-80%)	34% (20-53%)	1.6 (0.77-2.8)	4.7% (1.7-9.3%)	5.3% (1.9-11%)	37 (12-74)	400 (150-790)
Mongolia	2016	Male	85 to 89	63% (45-78%)	37% (22-55%)	1.5 (0.7-2.6)	3.4% (0.82-7.5%)	3.8% (0.73-8.6%)	14 (2.6-31)	110 (24-230)
Mongolia	2016	Male	90 to 94	62% (45-78%)	38% (22-55%)	1.4 (0.63-2.5)	2.1% (-0.38-6%)	2.1% (-0.75-6.6%)	2.6 (-0.93-8.1)	15 (-2.7-44)
Mongolia	2016	Male	95 plus	63% (41-80%)	37% (20-59%)	1.3 (0.5-2.5)	1.4% (-1.1-5.7%)	1.2% (-1.8-6.1%)	0.33 (-0.52-1.7)	1.6 (-1.3-6.5)
Tajikistan	1990	Female	All Ages	49% (43-57%)	51% (43-57%)	0.53 (0.4-0.69)	0.37% (-0.055-0.82%)	-0.2% (-1.5-1.4%)	-41 (-320-280)	4800 (-720-11000)
Tajikistan	1990	Female	15 to 19	52% (39-68%)	48% (32-61%)	0.54 (0.23-1)	1.3% (0.65-2.2%)	1.8% (0.51-3.6%)	3.4 (1-6.9)	420 (210-740)
Tajikistan	1990	Female	20 to 24	61% (48-74%)	39% (26-52%)	0.79 (0.36-1.4)	2.1% (1.2-3.3%)	2.2% (0.69-4.4%)	6.3 (1.9-13)	790 (420-1300)
Tajikistan	1990	Female	25 to 29	64% (50-77%)	36% (23-50%)	0.92 (0.44-1.6)	2.5% (1.5-3.9%)	3% (1.2-5.4%)	9.3 (3.8-17)	990 (560-1600)
Tajikistan	1990	Female	30 to 34	65% (51-78%)	35% (22-49%)	1 (0.54-1.7)	2.7% (1.7-4.1%)	3.6% (1.8-6%)	11 (5.6-19)	950 (580-1500)
Tajikistan	1990	Female	35 to 39	65% (52-79%)	35% (21-48%)	1 (0.5-1.9)	2.7% (1.6-4.5%)	3.9% (1.8-6.9%)	12 (5.5-22)	800 (440-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	1990	Female	40 to 44	65% (51-78%)	35% (22-49%)	1.1 (0.48-1.9)	2.7% (1.3-4.8%)	4.1% (1.6-7.6%)	11 (4.4-20)	610 (290-1100)
Tajikistan	1990	Female	45 to 49	65% (50-78%)	35% (22-50%)	1.1 (0.51-2.1)	2.5% (0.73-5.3%)	3.7% (0.85-8.2%)	9.6 (2.4-22)	450 (140-1000)
Tajikistan	1990	Female	50 to 54	64% (49-77%)	36% (23-51%)	1.2 (0.53-2.1)	2.2% (0.11-5.5%)	3.1% (-0.025-8%)	17 (-0.15-45)	700 (34-1700)
Tajikistan	1990	Female	55 to 59	62% (47-76%)	38% (24-53%)	1.2 (0.5-2.1)	1.5% (-0.93-5%)	2.1% (-1.4-7.2%)	15 (-9.7-50)	510 (-300-1700)
Tajikistan	1990	Female	60 to 64	60% (45-75%)	40% (25-55%)	1.1 (0.47-2.2)	0.45% (-2.2-4.5%)	0.57% (-3.1-5.8%)	6 (-31-61)	180 (-830-1700)
Tajikistan	1990	Female	65 to 69	58% (45-73%)	42% (27-55%)	1 (0.49-1.9)	-0.47% (-3-2.7%)	-0.59% (-3.9-3.6%)	-6.3 (-43-42)	-150 (-990-960)
Tajikistan	1990	Female	70 to 74	55% (43-68%)	45% (32-57%)	0.93 (0.37-1.8)	-1.1% (-3.7-2.3%)	-1.4% (-4.7-2.8%)	-16 (-54-33)	-300 (-990-610)
Tajikistan	1990	Female	75 to 79	52% (40-65%)	48% (35-60%)	0.87 (0.32-1.7)	-1.7% (-4.3-1.7%)	-2.1% (-5.3-2.1%)	-29 (-75-30)	-410 (-1100-430)
Tajikistan	1990	Female	80 to 84	50% (37-64%)	50% (36-63%)	0.89 (0.37-1.8)	-2% (-4.8-1.4%)	-2.5% (-6-1.5%)	-37 (-88-23)	-380 (-900-260)
Tajikistan	1990	Female	85 to 89	49% (37-61%)	51% (39-63%)	0.94 (0.37-1.9)	-2.2% (-5-2%)	-2.7% (-6-2.2%)	-34 (-77-27)	-240 (-550-230)
Tajikistan	1990	Female	90 to 94	50% (38-62%)	50% (38-62%)	1 (0.34-2.3)	-2% (-5-2.9%)	-2.6% (-6.1-3.2%)	-16 (-37-20)	-74 (-180-100)
Tajikistan	1990	Female	95 plus	51% (38-66%)	49% (34-62%)	1.1 (0.28-2.7)	-1.7% (-4.9-4.7%)	-2.2% (-6-5.3%)	-3.7 (-10-8.9)	-11 (-32-30)
Tajikistan	1990	Male	All Ages	60% (51-69%)	40% (31-49%)	0.83 (0.66-1)	1.4% (0.85-1.9%)	0.98% (-0.14-2.2%)	230 (-31-520)	20000 (13000-29000)
Tajikistan	1990	Male	15 to 19	55% (39-71%)	45% (29-61%)	0.46 (0.26-0.76)	2.6% (1.7-3.6%)	2.6% (1.3-4.3%)	6.9 (3.4-12)	950 (610-1400)
Tajikistan	1990	Male	20 to 24	73% (57-85%)	27% (15-43%)	1 (0.65-1.5)	6% (4.2-7.9%)	6.1% (3.4-9%)	21 (12-32)	2300 (1600-3200)
Tajikistan	1990	Male	25 to 29	80% (68-89%)	20% (11-32%)	1.7 (1.1-2.4)	7.9% (5.6-10%)	8.6% (5.3-12%)	42 (25-59)	3700 (2600-4800)
Tajikistan	1990	Male	30 to 34	84% (73-92%)	16% (8-27%)	2 (1.4-2.7)	8.3% (6.1-11%)	9.4% (6.1-13%)	45 (29-63)	3500 (2400-4600)
Tajikistan	1990	Male	35 to 39	86% (76-93%)	14% (7.4-24%)	2 (1.3-2.9)	7.4% (5.2-10%)	8.5% (5.2-12%)	37 (22-55)	2500 (1700-3500)
Tajikistan	1990	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	2 (1.3-3)	6.6% (4.3-9.3%)	7.4% (4.4-11%)	30 (17-46)	1800 (1200-2600)
Tajikistan	1990	Male	45 to 49	85% (73-93%)	15% (7.3-27%)	2.2 (1.3-3.2)	6.1% (3.4-8.8%)	6.8% (3.3-11%)	32 (15-51)	1700 (910-2500)
Tajikistan	1990	Male	50 to 54	84% (73-92%)	16% (8-27%)	2.2 (1.3-3.2)	5% (2.3-7.9%)	5.5% (1.9-9.2%)	49 (16-86)	2200 (990-3600)
Tajikistan	1990	Male	55 to 59	83% (70-91%)	17% (8.7-30%)	1.9 (1.1-2.9)	3.4% (0.81-6.3%)	3.6% (0.35-7.3%)	37 (3.9-78)	1500 (370-2800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	1990	Male	60 to 64	81% (67-91%)	19% (9.4-33%)	1.6 (0.95-2.6)	1.9% (-0.69-4.9%)	1.8% (-1.3-5.4%)	27 (-19-77)	920 (-330-2300)
Tajikistan	1990	Male	65 to 69	79% (64-90%)	21% (10-36%)	1.5 (0.9-2.2)	0.68% (-1.7-3.3%)	0.51% (-2.4-3.6%)	5.3 (-25-38)	200 (-480-950)
Tajikistan	1990	Male	70 to 74	76% (62-86%)	24% (14-38%)	1.2 (0.74-1.9)	-0.45% (-2.8-2.2%)	-0.76% (-3.6-2.4%)	-6.7 (-33-21)	-86 (-560-430)
Tajikistan	1990	Male	75 to 79	73% (57-85%)	27% (15-43%)	1.1 (0.57-1.7)	-1.3% (-3.5-0.97%)	-1.7% (-4.4-0.95%)	-20 (-51-11)	-250 (-680-180)
Tajikistan	1990	Male	80 to 84	70% (53-83%)	30% (17-47%)	0.96 (0.54-1.5)	-1.9% (-4.2-0.16%)	-2.5% (-5.2-0.091%)	-30 (-64-1)	-280 (-630-23)
Tajikistan	1990	Male	85 to 89	68% (52-81%)	32% (19-48%)	0.89 (0.51-1.4)	-2.3% (-4.5--0.27%)	-2.9% (-5.6--0.46%)	-28 (-54--4.3)	-190 (-370--22)
Tajikistan	1990	Male	90 to 94	67% (52-80%)	33% (20-48%)	0.82 (0.46-1.4)	-2.4% (-4.7--0.53%)	-3.1% (-5.9--0.76%)	-14 (-27--3.5)	-66 (-130--14)
Tajikistan	1990	Male	95 plus	66% (46-82%)	34% (18-54%)	0.75 (0.33-1.4)	-2.3% (-4.5--0.39%)	-3% (-5.7--0.71%)	-3.4 (-6.4--0.81)	-10 (-20--1.7)
Tajikistan	1995	Female	All Ages	49% (43-57%)	51% (43-57%)	0.51 (0.4-0.64)	0.42% (-0.076-0.93%)	-0.24% (-1.8-1.4%)	-50 (-370-290)	5300 (-890-12000)
Tajikistan	1995	Female	15 to 19	53% (40-68%)	47% (32-60%)	0.54 (0.23-0.98)	1.2% (0.58-2.1%)	1.6% (0.46-3.3%)	4 (1.1-8.2)	480 (220-830)
Tajikistan	1995	Female	20 to 24	61% (48-73%)	39% (27-52%)	0.78 (0.34-1.4)	2.1% (1.1-3.5%)	2.2% (0.6-4.4%)	7.4 (2-15)	880 (430-1500)
Tajikistan	1995	Female	25 to 29	64% (50-76%)	36% (24-50%)	0.9 (0.44-1.5)	2.6% (1.4-4.1%)	3% (1-5.5%)	11 (3.7-21)	1100 (570-1700)
Tajikistan	1995	Female	30 to 34	65% (51-79%)	35% (21-49%)	0.98 (0.53-1.6)	2.7% (1.6-4.2%)	3.6% (1.7-6%)	16 (7.5-27)	1200 (740-1900)
Tajikistan	1995	Female	35 to 39	65% (52-80%)	35% (20-48%)	1 (0.51-1.8)	2.7% (1.5-4.5%)	3.8% (1.7-6.7%)	16 (7.1-29)	1100 (580-1800)
Tajikistan	1995	Female	40 to 44	65% (51-79%)	35% (21-49%)	1.1 (0.48-1.8)	2.7% (1.2-4.9%)	4% (1.5-7.6%)	16 (5.4-31)	850 (370-1600)
Tajikistan	1995	Female	45 to 49	65% (50-78%)	35% (22-50%)	1.1 (0.51-2.2)	2.3% (0.47-5.4%)	3.4% (0.39-8.1%)	12 (1.5-29)	570 (120-1300)
Tajikistan	1995	Female	50 to 54	64% (49-77%)	36% (23-51%)	1.2 (0.53-2.2)	2% (-0.25-5.7%)	2.8% (-0.56-7.8%)	10 (-2.1-29)	410 (-54-1100)
Tajikistan	1995	Female	55 to 59	62% (48-75%)	38% (25-52%)	1.2 (0.49-2.1)	1.3% (-1.3-5.2%)	1.7% (-1.9-6.9%)	14 (-15-59)	480 (-480-1900)
Tajikistan	1995	Female	60 to 64	60% (45-75%)	40% (25-55%)	1.1 (0.46-2.2)	0.27% (-2.6-4.8%)	0.31% (-3.4-6.2%)	3.2 (-36-65)	100 (-980-1800)
Tajikistan	1995	Female	65 to 69	58% (45-73%)	42% (27-55%)	1 (0.48-1.9)	-0.73% (-3.6-3%)	-0.92% (-4.5-3.7%)	-13 (-61-53)	-300 (-1400-1200)
Tajikistan	1995	Female	70 to 74	56% (43-69%)	44% (31-57%)	0.92 (0.36-1.8)	-1.3% (-4.1-2.5%)	-1.6% (-5.1-3.1%)	-23 (-73-44)	-420 (-1300-800)
Tajikistan	1995	Female	75 to 79	52% (41-66%)	48% (34-59%)	0.86 (0.32-1.7)	-1.8% (-4.6-1.5%)	-2.2% (-5.6-1.7%)	-28 (-70-22)	-400 (-980-310)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	1995	Female	80 to 84	50% (37-65%)	50% (35-63%)	0.88 (0.35-1.7)	-2.2% (-5.1-1.3%)	-2.7% (-6.1-1.4%)	-36 (-84-20)	-370 (-860-210)
Tajikistan	1995	Female	85 to 89	50% (38-62%)	50% (38-62%)	0.93 (0.36-1.9)	-2.3% (-5.2-1.7%)	-2.8% (-6.2-1.8%)	-36 (-80-24)	-260 (-580-190)
Tajikistan	1995	Female	90 to 94	51% (39-62%)	49% (38-61%)	1 (0.35-2.3)	-2.2% (-5.1-2.8%)	-2.7% (-6.2-3.3%)	-19 (-43-22)	-89 (-210-120)
Tajikistan	1995	Female	95 plus	51% (38-67%)	49% (33-62%)	1 (0.28-2.6)	-1.8% (-5-4.5%)	-2.3% (-6.1-5.3%)	-5.1 (-14-12)	-16 (-44-41)
Tajikistan	1995	Male	All Ages	61% (52-70%)	39% (30-48%)	0.82 (0.68-0.95)	1.7% (1.1-2.4%)	1.4% (0.11-2.7%)	330 (25-640)	26000 (17000-37000)
Tajikistan	1995	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.46 (0.26-0.74)	2.4% (1.5-3.4%)	2.3% (1.1-3.8%)	9.4 (4.5-16)	1200 (710-1700)
Tajikistan	1995	Male	20 to 24	73% (58-85%)	27% (15-42%)	1 (0.66-1.4)	5.6% (3.7-7.6%)	5.5% (3-8%)	33 (17-50)	3200 (2100-4600)
Tajikistan	1995	Male	25 to 29	80% (69-89%)	20% (11-31%)	1.7 (1.1-2.3)	8% (5.6-10%)	8.4% (5.4-12%)	57 (35-81)	4600 (3100-6100)
Tajikistan	1995	Male	30 to 34	84% (73-92%)	16% (8-27%)	2 (1.4-2.6)	8.6% (6.3-11%)	9.5% (6.4-13%)	73 (48-100)	5200 (3700-6800)
Tajikistan	1995	Male	35 to 39	86% (75-93%)	14% (7.2-25%)	2 (1.3-2.7)	7.8% (5.5-10%)	8.8% (5.5-12%)	64 (39-89)	4100 (2800-5500)
Tajikistan	1995	Male	40 to 44	86% (77-93%)	14% (7-23%)	2 (1.3-2.9)	6.7% (4.4-9.1%)	7.4% (4.4-11%)	50 (28-73)	2800 (1800-3900)
Tajikistan	1995	Male	45 to 49	86% (74-93%)	14% (7.2-26%)	2.2 (1.4-3.2)	6.1% (3.3-9.1%)	6.7% (3.1-11%)	41 (19-66)	2000 (1100-3100)
Tajikistan	1995	Male	50 to 54	85% (73-92%)	15% (8-27%)	2.1 (1.3-3.1)	4.9% (2.1-8.1%)	5.3% (1.6-9.3%)	35 (11-65)	1600 (660-2700)
Tajikistan	1995	Male	55 to 59	83% (71-91%)	17% (8.8-29%)	1.9 (1.1-2.7)	3.1% (0.32-6.1%)	3.2% (-0.23-7%)	40 (-2.9-87)	1600 (160-3000)
Tajikistan	1995	Male	60 to 64	81% (67-91%)	19% (9.4-33%)	1.6 (0.94-2.5)	1.5% (-1.2-4.6%)	1.4% (-1.9-5.1%)	20 (-27-74)	720 (-540-2200)
Tajikistan	1995	Male	65 to 69	79% (65-89%)	21% (11-35%)	1.4 (0.9-2.1)	0.36% (-2.2-3.1%)	0.12% (-3-3.4%)	2.1 (-54-61)	170 (-1100-1500)
Tajikistan	1995	Male	70 to 74	76% (62-86%)	24% (14-38%)	1.2 (0.71-1.9)	-0.64% (-3.1-1.9%)	-0.98% (-3.9-2%)	-12 (-48-25)	-170 (-810-510)
Tajikistan	1995	Male	75 to 79	73% (58-85%)	27% (15-42%)	1.1 (0.58-1.7)	-1.5% (-3.8-0.77%)	-1.9% (-4.6-0.8%)	-17 (-41-7.2)	-210 (-550-120)
Tajikistan	1995	Male	80 to 84	71% (54-83%)	29% (17-46%)	0.95 (0.54-1.4)	-2% (-4.4-0.14%)	-2.6% (-5.3-0.047%)	-24 (-51-0.45)	-230 (-510-17)
Tajikistan	1995	Male	85 to 89	69% (53-80%)	31% (20-47%)	0.87 (0.51-1.4)	-2.3% (-4.5--0.34%)	-2.9% (-5.5--0.54%)	-24 (-45--4.4)	-160 (-310--24)
Tajikistan	1995	Male	90 to 94	68% (53-80%)	32% (20-47%)	0.81 (0.45-1.3)	-2.4% (-4.7--0.5%)	-3.1% (-5.8--0.74%)	-13 (-25--3.1)	-62 (-120--13)
Tajikistan	1995	Male	95 plus	67% (48-82%)	33% (18-52%)	0.73 (0.33-1.3)	-2.3% (-4.4--0.4%)	-2.9% (-5.6--0.71%)	-4 (-7.7--0.98)	-12 (-24--2.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	2000	Female	All Ages	50% (44-59%)	50% (41-56%)	0.46 (0.37-0.58)	0.4% (-0.15-0.95%)	-0.47% (-2.1-1.2%)	-91 (-400-240)	4500 (-1700-11000)
Tajikistan	2000	Female	15 to 19	53% (40-68%)	47% (32-60%)	0.48 (0.22-0.85)	1.1% (0.55-1.9%)	1.5% (0.41-3.1%)	3.5 (0.91-7.1)	480 (230-820)
Tajikistan	2000	Female	20 to 24	61% (49-73%)	39% (27-51%)	0.69 (0.31-1.2)	2% (1-3.3%)	2.2% (0.5-4.6%)	6.7 (1.6-14)	840 (420-1400)
Tajikistan	2000	Female	25 to 29	64% (49-76%)	36% (24-51%)	0.8 (0.38-1.4)	2.4% (1.2-3.9%)	2.9% (0.64-5.5%)	10 (2.3-20)	1000 (490-1700)
Tajikistan	2000	Female	30 to 34	65% (50-78%)	35% (22-50%)	0.87 (0.46-1.5)	2.7% (1.5-4.1%)	3.5% (1.4-6%)	14 (5.7-24)	1200 (630-1800)
Tajikistan	2000	Female	35 to 39	65% (52-79%)	35% (21-48%)	0.9 (0.45-1.5)	2.6% (1.4-4.3%)	3.8% (1.7-6.5%)	18 (7.7-31)	1200 (630-1900)
Tajikistan	2000	Female	40 to 44	65% (51-79%)	35% (21-49%)	0.95 (0.43-1.6)	2.5% (1.1-4.4%)	3.8% (1.5-7%)	18 (6.7-34)	980 (440-1700)
Tajikistan	2000	Female	45 to 49	65% (49-79%)	35% (21-51%)	0.99 (0.45-1.9)	2% (0.26-4.5%)	3% (0.091-7%)	13 (0.42-32)	620 (80-1400)
Tajikistan	2000	Female	50 to 54	64% (49-78%)	36% (22-51%)	1 (0.46-1.9)	1.6% (-0.4-4.8%)	2.3% (-0.78-6.9%)	10 (-3.4-30)	400 (-110-1200)
Tajikistan	2000	Female	55 to 59	62% (48-76%)	38% (24-52%)	1 (0.44-1.9)	0.85% (-1.6-4.4%)	1.1% (-2.2-5.9%)	5.5 (-11-29)	190 (-340-920)
Tajikistan	2000	Female	60 to 64	60% (46-75%)	40% (25-54%)	0.99 (0.41-1.9)	-0.19% (-2.9-3.8%)	-0.28% (-3.8-4.8%)	-3.2 (-43-53)	-80 (-1100-1500)
Tajikistan	2000	Female	65 to 69	58% (45-73%)	42% (27-55%)	0.91 (0.42-1.7)	-1.2% (-3.9-2.4%)	-1.4% (-4.8-2.8%)	-19 (-67-38)	-450 (-1500-890)
Tajikistan	2000	Female	70 to 74	55% (43-68%)	45% (32-57%)	0.81 (0.32-1.6)	-1.6% (-4.4-1.8%)	-1.9% (-5.2-2.1%)	-34 (-94-36)	-620 (-1700-640)
Tajikistan	2000	Female	75 to 79	52% (41-66%)	48% (34-59%)	0.76 (0.28-1.5)	-2% (-4.6-0.89%)	-2.4% (-5.6-0.98%)	-38 (-88-15)	-540 (-1200-210)
Tajikistan	2000	Female	80 to 84	50% (37-65%)	50% (35-63%)	0.78 (0.31-1.5)	-2.4% (-5.1-0.69%)	-2.8% (-6.2-0.74%)	-36 (-79-9)	-370 (-810-110)
Tajikistan	2000	Female	85 to 89	50% (37-62%)	50% (38-63%)	0.83 (0.32-1.8)	-2.5% (-5.3-0.97%)	-3% (-6.2-1%)	-33 (-69-10)	-240 (-490-84)
Tajikistan	2000	Female	90 to 94	51% (39-62%)	49% (38-61%)	0.89 (0.3-2.1)	-2.4% (-5.1-2.3%)	-2.9% (-6.2-2.5%)	-19 (-41-16)	-93 (-200-86)
Tajikistan	2000	Female	95 plus	51% (38-66%)	49% (34-62%)	0.93 (0.24-2.4)	-2% (-4.9-4.1%)	-2.5% (-5.9-4.6%)	-6.1 (-14-11)	-19 (-47-38)
Tajikistan	2000	Male	All Ages	61% (52-70%)	39% (30-48%)	0.75 (0.62-0.85)	1.8% (1.1-2.5%)	1.1% (-0.24-2.5%)	250 (-53-570)	24000 (15000-33000)
Tajikistan	2000	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.41 (0.23-0.64)	2.3% (1.5-3.2%)	2.2% (1.1-3.7%)	8.1 (4.1-14)	1100 (740-1700)
Tajikistan	2000	Male	20 to 24	73% (57-85%)	27% (15-43%)	0.91 (0.59-1.3)	5.4% (3.6-7.3%)	5.3% (2.7-8.1%)	26 (13-42)	2800 (1800-4000)
Tajikistan	2000	Male	25 to 29	80% (69-89%)	20% (11-31%)	1.5 (0.98-2)	7.6% (5.4-10%)	8.2% (5-12%)	48 (28-71)	4100 (2800-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	2000	Male	30 to 34	85% (73-92%)	15% (7.7-27%)	1.8 (1.3-2.4)	8.2% (6.1-11%)	9.2% (6.1-13%)	59 (39-82)	4400 (3100-5800)
Tajikistan	2000	Male	35 to 39	86% (75-93%)	14% (7.1-25%)	1.8 (1.2-2.4)	7.5% (5.3-9.9%)	8.6% (5.4-12%)	61 (38-88)	4100 (2800-5600)
Tajikistan	2000	Male	40 to 44	86% (76-93%)	14% (7-24%)	1.8 (1.2-2.5)	6.4% (4.1-8.6%)	7.2% (4-10%)	51 (28-76)	3100 (1900-4200)
Tajikistan	2000	Male	45 to 49	86% (73-93%)	14% (7.1-27%)	1.9 (1.2-2.8)	5.6% (3-8.2%)	6.1% (2.5-9.7%)	42 (17-69)	2200 (1100-3400)
Tajikistan	2000	Male	50 to 54	85% (73-92%)	15% (8-27%)	1.9 (1.1-2.8)	4.3% (1.6-7.4%)	4.6% (1.1-8.5%)	31 (6.9-58)	1400 (520-2400)
Tajikistan	2000	Male	55 to 59	83% (71-92%)	17% (8.1-29%)	1.7 (1-2.5)	2.8% (0.19-5.6%)	2.8% (-0.63-6.4%)	22 (-4.8-52)	890 (58-1800)
Tajikistan	2000	Male	60 to 64	81% (68-91%)	19% (9.4-32%)	1.4 (0.83-2.3)	1.1% (-1.5-3.9%)	0.9% (-2.2-4.2%)	14 (-35-67)	560 (-770-1900)
Tajikistan	2000	Male	65 to 69	79% (65-89%)	21% (11-35%)	1.3 (0.79-1.9)	-0.075% (-2.6-2.6%)	-0.35% (-3.3-2.8%)	-6 (-58-48)	-34 (-1200-1200)
Tajikistan	2000	Male	70 to 74	76% (62-86%)	24% (14-38%)	1.1 (0.63-1.7)	-0.95% (-3.4-1.3%)	-1.3% (-4.2-1.4%)	-28 (-89-29)	-430 (-1500-620)
Tajikistan	2000	Male	75 to 79	73% (57-85%)	27% (15-43%)	0.94 (0.5-1.5)	-1.6% (-3.8-0.46%)	-2% (-4.7-0.4%)	-26 (-60-5.4)	-330 (-800-100)
Tajikistan	2000	Male	80 to 84	71% (54-84%)	29% (16-46%)	0.85 (0.46-1.3)	-2.1% (-4.5-0.033%)	-2.6% (-5.4--0.13%)	-20 (-41--0.94)	-190 (-400-3)
Tajikistan	2000	Male	85 to 89	69% (53-80%)	31% (20-47%)	0.78 (0.44-1.2)	-2.4% (-4.6--0.42%)	-2.9% (-5.6--0.58%)	-17 (-33--3.4)	-120 (-230--21)
Tajikistan	2000	Male	90 to 94	68% (53-80%)	32% (20-47%)	0.72 (0.39-1.2)	-2.4% (-4.7--0.53%)	-3.1% (-5.8--0.8%)	-10 (-20--2.7)	-48 (-94--11)
Tajikistan	2000	Male	95 plus	67% (48-82%)	33% (18-52%)	0.65 (0.3-1.2)	-2.2% (-4.5--0.38%)	-2.9% (-5.5--0.7%)	-3.3 (-6.5--0.77)	-9.8 (-20--1.7)
Tajikistan	2005	Female	All Ages	51% (45-60%)	49% (40-55%)	0.44 (0.35-0.56)	0.41% (-0.26-1%)	-0.74% (-2.4-1%)	-130 (-440-190)	4100 (-2300-11000)
Tajikistan	2005	Female	15 to 19	52% (40-67%)	48% (33-60%)	0.43 (0.18-0.76)	1.1% (0.54-1.8%)	1.5% (0.3-3.1%)	3.5 (0.68-7.1)	520 (260-870)
Tajikistan	2005	Female	20 to 24	61% (48-73%)	39% (27-52%)	0.62 (0.27-1.1)	2% (1-3.1%)	2.2% (0.35-4.6%)	6.1 (0.92-13)	890 (440-1400)
Tajikistan	2005	Female	25 to 29	64% (49-77%)	36% (23-51%)	0.72 (0.34-1.3)	2.4% (1.2-3.9%)	2.9% (0.52-5.6%)	8.2 (1.4-17)	960 (480-1500)
Tajikistan	2005	Female	30 to 34	65% (51-79%)	35% (21-49%)	0.79 (0.4-1.3)	2.6% (1.6-3.9%)	3.6% (1.5-6.1%)	12 (5.2-20)	1100 (630-1600)
Tajikistan	2005	Female	35 to 39	65% (52-80%)	35% (20-48%)	0.81 (0.39-1.4)	2.5% (1.5-3.9%)	4% (2-6.7%)	15 (7.3-25)	1100 (630-1600)
Tajikistan	2005	Female	40 to 44	65% (51-79%)	35% (21-49%)	0.84 (0.38-1.5)	2.4% (1.2-4%)	3.9% (1.6-6.9%)	19 (7.9-33)	1100 (540-1800)
Tajikistan	2005	Female	45 to 49	65% (49-78%)	35% (22-51%)	0.89 (0.41-1.6)	1.9% (0.47-3.8%)	2.9% (0.4-6.4%)	17 (2.3-37)	800 (190-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	2005	Female	50 to 54	63% (49-77%)	37% (23-51%)	0.91 (0.42-1.7)	1.4% (-0.33-3.9%)	2% (-0.58-5.8%)	12 (-3.5-34)	480 (-120-1300)
Tajikistan	2005	Female	55 to 59	62% (48-75%)	38% (25-52%)	0.92 (0.39-1.7)	0.64% (-1.5-3.8%)	0.86% (-2.2-5.1%)	5.7 (-14-35)	190 (-440-1100)
Tajikistan	2005	Female	60 to 64	60% (45-74%)	40% (26-55%)	0.89 (0.36-1.8)	-0.45% (-3-3.3%)	-0.6% (-3.8-4.2%)	-4.2 (-27-29)	-110 (-730-800)
Tajikistan	2005	Female	65 to 69	58% (46-73%)	42% (27-54%)	0.82 (0.37-1.5)	-1.3% (-3.7-1.6%)	-1.6% (-4.6-2.1%)	-24 (-71-32)	-560 (-1600-730)
Tajikistan	2005	Female	70 to 74	56% (43-69%)	44% (31-57%)	0.73 (0.29-1.4)	-1.8% (-4.4-1.1%)	-2.1% (-5.2-1.2%)	-40 (-99-26)	-740 (-1800-460)
Tajikistan	2005	Female	75 to 79	52% (41-65%)	48% (35-59%)	0.68 (0.25-1.3)	-2.2% (-4.7-0.25%)	-2.6% (-5.5-0.29%)	-49 (-110-5.5)	-690 (-1500-72)
Tajikistan	2005	Female	80 to 84	50% (37-64%)	50% (36-63%)	0.7 (0.28-1.3)	-2.5% (-5.3-0.072%)	-3% (-6.2-0.044%)	-52 (-110-0.76)	-540 (-1100-15)
Tajikistan	2005	Female	85 to 89	50% (38-62%)	50% (38-62%)	0.74 (0.27-1.5)	-2.7% (-5.3-0.08%)	-3.3% (-6.3-0.021%)	-37 (-71-0.23)	-260 (-510-7.9)
Tajikistan	2005	Female	90 to 94	51% (39-62%)	49% (38-61%)	0.79 (0.27-1.8)	-2.6% (-5.2-1.1%)	-3.2% (-6.3-1.1%)	-20 (-39-6.9)	-96 (-190-41)
Tajikistan	2005	Female	95 plus	51% (38-67%)	49% (33-62%)	0.83 (0.22-2.1)	-2.3% (-4.9-2.6%)	-2.8% (-5.8-2.9%)	-7.2 (-15-7.3)	-23 (-49-26)
Tajikistan	2005	Male	All Ages	63% (55-72%)	37% (28-45%)	0.71 (0.6-0.82)	1.9% (1.1-2.7%)	0.96% (-0.67-2.4%)	210 (-140-540)	24000 (14000-33000)
Tajikistan	2005	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.37 (0.21-0.58)	2.3% (1.6-3.1%)	2.1% (0.99-3.5%)	7.6 (3.7-13)	1200 (790-1800)
Tajikistan	2005	Male	20 to 24	73% (58-85%)	27% (15-42%)	0.82 (0.55-1.1)	5.3% (3.5-7.2%)	5.1% (2.3-7.8%)	23 (10-37)	2900 (1800-4100)
Tajikistan	2005	Male	25 to 29	80% (69-89%)	20% (11-31%)	1.3 (0.89-1.8)	7.4% (5.3-9.7%)	7.9% (4.6-11%)	38 (22-56)	3700 (2600-5000)
Tajikistan	2005	Male	30 to 34	85% (73-92%)	15% (7.9-27%)	1.6 (1.1-2.1)	8% (5.9-10%)	9% (5.8-13%)	52 (33-74)	4200 (3000-5700)
Tajikistan	2005	Male	35 to 39	86% (76-93%)	14% (7.3-24%)	1.6 (1-2.2)	7.3% (5.2-9.5%)	8.5% (5.4-12%)	54 (33-79)	3800 (2600-5100)
Tajikistan	2005	Male	40 to 44	86% (77-93%)	14% (7-23%)	1.6 (1.1-2.2)	6.3% (4.2-8.4%)	7.2% (4.2-10%)	58 (33-86)	3500 (2400-4900)
Tajikistan	2005	Male	45 to 49	86% (74-93%)	14% (7.3-26%)	1.7 (1.1-2.5)	5.4% (2.9-7.8%)	6% (2.6-9.4%)	52 (23-85)	2800 (1500-4200)
Tajikistan	2005	Male	50 to 54	85% (73-92%)	15% (7.7-27%)	1.7 (1-2.5)	4.1% (1.5-6.8%)	4.3% (0.82-7.9%)	40 (7.9-76)	1900 (670-3200)
Tajikistan	2005	Male	55 to 59	83% (71-92%)	17% (8.4-29%)	1.5 (0.9-2.2)	2.6% (0.079-5.3%)	2.6% (-0.68-6%)	23 (-6-54)	960 (29-2000)
Tajikistan	2005	Male	60 to 64	81% (68-91%)	19% (9.4-32%)	1.3 (0.71-2.1)	0.9% (-1.6-3.4%)	0.67% (-2.4-3.8%)	7.8 (-28-43)	340 (-610-1300)
Tajikistan	2005	Male	65 to 69	79% (64-89%)	21% (11-36%)	1.1 (0.7-1.7)	-0.17% (-2.7-2.3%)	-0.44% (-3.4-2.6%)	-9.1 (-71-51)	-92 (-1500-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	2005	Male	70 to 74	76% (62-87%)	24% (13-38%)	0.98 (0.57-1.5)	-1.1% (-3.7-1.1%)	-1.5% (-4.3-1.1%)	-33 (-100-27)	-540 (-1700-550)
Tajikistan	2005	Male	75 to 79	73% (58-85%)	27% (15-42%)	0.84 (0.45-1.3)	-1.6% (-3.9-0.38%)	-2.1% (-4.7-0.33%)	-47 (-110-6.9)	-620 (-1500-150)
Tajikistan	2005	Male	80 to 84	71% (55-83%)	29% (17-45%)	0.75 (0.42-1.1)	-2.1% (-4.5-0.011%)	-2.6% (-5.4--0.09%)	-32 (-66--1)	-310 (-660-1.6)
Tajikistan	2005	Male	85 to 89	69% (53-80%)	31% (20-47%)	0.69 (0.4-1.1)	-2.4% (-4.6--0.35%)	-2.9% (-5.6--0.52%)	-14 (-27--2.3)	-93 (-190--15)
Tajikistan	2005	Male	90 to 94	68% (53-80%)	32% (20-47%)	0.64 (0.36-1.1)	-2.4% (-4.6--0.5%)	-3% (-5.7--0.73%)	-7 (-13--1.7)	-32 (-63--6.6)
Tajikistan	2005	Male	95 plus	66% (48-82%)	34% (18-52%)	0.58 (0.26-1)	-2.2% (-4.4--0.38%)	-2.8% (-5.5--0.64%)	-2.5 (-4.9--0.59)	-7.5 (-15--1.4)
Tajikistan	2010	Female	All Ages	52% (45-60%)	48% (40-55%)	0.41 (0.32-0.52)	0.38% (-0.16-0.91%)	-0.86% (-2.5-0.65%)	-150 (-420-110)	3800 (-1600-9300)
Tajikistan	2010	Female	15 to 19	52% (40-67%)	48% (33-60%)	0.38 (0.17-0.68)	0.98% (0.55-1.5%)	1.3% (0.31-2.5%)	2.9 (0.7-6.1)	470 (250-800)
Tajikistan	2010	Female	20 to 24	61% (48-74%)	39% (26-52%)	0.55 (0.25-0.97)	1.8% (1-2.8%)	1.8% (0.31-3.6%)	5.9 (1.1-12)	1000 (540-1600)
Tajikistan	2010	Female	25 to 29	64% (49-76%)	36% (24-51%)	0.64 (0.3-1.1)	2.1% (1.3-3.2%)	2.3% (0.66-4.5%)	7.5 (2.1-14)	1000 (590-1600)
Tajikistan	2010	Female	30 to 34	65% (51-77%)	35% (23-49%)	0.69 (0.37-1.2)	2.3% (1.5-3.4%)	3.1% (1.5-5%)	10 (5-17)	1000 (650-1500)
Tajikistan	2010	Female	35 to 39	65% (52-79%)	35% (21-48%)	0.71 (0.35-1.3)	2.2% (1.4-3.4%)	3.4% (1.7-5.6%)	13 (6.6-22)	980 (610-1500)
Tajikistan	2010	Female	40 to 44	65% (51-79%)	35% (21-49%)	0.74 (0.34-1.3)	2% (1.1-3.3%)	3.3% (1.4-5.8%)	16 (6.4-28)	940 (480-1500)
Tajikistan	2010	Female	45 to 49	65% (49-78%)	35% (22-51%)	0.78 (0.36-1.5)	1.5% (0.33-3.3%)	2.3% (0.25-5.3%)	15 (1.6-37)	780 (170-1700)
Tajikistan	2010	Female	50 to 54	63% (49-78%)	37% (22-51%)	0.8 (0.37-1.5)	1% (-0.37-3%)	1.4% (-0.8-4.7%)	11 (-6.3-35)	490 (-180-1400)
Tajikistan	2010	Female	55 to 59	61% (48-76%)	39% (24-52%)	0.81 (0.33-1.6)	0.27% (-1.6-2.6%)	0.34% (-2.3-3.6%)	2.8 (-19-30)	110 (-580-1000)
Tajikistan	2010	Female	60 to 64	60% (46-75%)	40% (25-54%)	0.78 (0.31-1.6)	-0.72% (-2.8-2.1%)	-1% (-3.9-2.7%)	-8.3 (-32-23)	-230 (-870-630)
Tajikistan	2010	Female	65 to 69	58% (45-73%)	42% (27-55%)	0.72 (0.33-1.4)	-1.5% (-3.7-0.67%)	-1.9% (-4.8-0.89%)	-16 (-41-8)	-380 (-930-170)
Tajikistan	2010	Female	70 to 74	56% (43-69%)	44% (31-57%)	0.64 (0.25-1.3)	-1.9% (-4.1-0.47%)	-2.3% (-5.2-0.63%)	-39 (-89-10)	-730 (-1600-170)
Tajikistan	2010	Female	75 to 79	52% (40-66%)	48% (34-60%)	0.6 (0.22-1.2)	-2.3% (-4.5--0.25%)	-2.8% (-5.6--0.28%)	-49 (-98--5.2)	-690 (-1400--82)
Tajikistan	2010	Female	80 to 84	50% (37-64%)	50% (36-63%)	0.61 (0.24-1.2)	-2.6% (-5.1--0.53%)	-3.2% (-6.3--0.63%)	-51 (-98--10)	-520 (-1000--110)
Tajikistan	2010	Female	85 to 89	50% (37-62%)	50% (38-63%)	0.65 (0.25-1.3)	-2.8% (-5.2--0.55%)	-3.4% (-6.3--0.71%)	-47 (-86--9.3)	-340 (-620--65)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	2010	Female	90 to 94	51% (38-63%)	49% (37-62%)	0.7 (0.24-1.6)	-2.7% (-5-0.041%)	-3.4% (-6.2--0.056%)	-20 (-36--0.35)	-97 (-180-1.5)
Tajikistan	2010	Female	95 plus	51% (38-66%)	49% (34-62%)	0.73 (0.19-1.9)	-2.4% (-4.8-1%)	-3% (-5.9-0.95%)	-7.1 (-14-2.3)	-23 (-45-9.9)
Tajikistan	2010	Male	All Ages	64% (55-73%)	36% (27-45%)	0.66 (0.55-0.77)	1.9% (1.2-2.7%)	0.77% (-0.79-2.2%)	180 (-180-500)	25000 (15000-35000)
Tajikistan	2010	Male	15 to 19	56% (40-71%)	44% (29-60%)	0.32 (0.18-0.5)	2.2% (1.5-3%)	1.9% (0.99-3.1%)	7.5 (3.8-12)	1200 (820-1800)
Tajikistan	2010	Male	20 to 24	73% (58-85%)	27% (15-42%)	0.72 (0.48-0.99)	4.9% (3.4-6.6%)	4.4% (2.2-6.6%)	26 (13-40)	3400 (2300-4800)
Tajikistan	2010	Male	25 to 29	80% (69-89%)	20% (11-31%)	1.2 (0.78-1.6)	6.7% (5-8.5%)	6.7% (4.2-9.3%)	41 (24-58)	4200 (3000-5600)
Tajikistan	2010	Male	30 to 34	85% (73-92%)	15% (7.7-27%)	1.4 (0.98-1.9)	7.3% (5.6-9.4%)	7.9% (5.1-11%)	48 (31-69)	4100 (3000-5500)
Tajikistan	2010	Male	35 to 39	86% (76-93%)	14% (7.4-24%)	1.4 (0.9-2)	6.7% (4.9-8.7%)	7.4% (4.6-10%)	50 (31-72)	3700 (2500-4900)
Tajikistan	2010	Male	40 to 44	86% (76-93%)	14% (7-24%)	1.4 (0.92-2)	5.7% (3.7-7.7%)	6.4% (3.4-9.6%)	49 (25-74)	3200 (2000-4500)
Tajikistan	2010	Male	45 to 49	85% (73-93%)	15% (7.4-27%)	1.5 (0.95-2.2)	4.8% (2.7-7.1%)	5.2% (1.9-8.5%)	52 (19-85)	2900 (1600-4500)
Tajikistan	2010	Male	50 to 54	85% (72-92%)	15% (8.1-28%)	1.5 (0.87-2.2)	3.5% (1.2-6%)	3.6% (0.24-6.8%)	45 (3.2-87)	2200 (730-3800)
Tajikistan	2010	Male	55 to 59	83% (71-91%)	17% (8.5-29%)	1.3 (0.78-2)	2.2% (-0.13-4.4%)	2.1% (-0.95-5%)	25 (-12-60)	1100 (-66-2200)
Tajikistan	2010	Male	60 to 64	81% (68-91%)	19% (9.3-32%)	1.1 (0.64-1.8)	0.71% (-1.5-2.9%)	0.42% (-2.4-3.3%)	5.2 (-29-41)	300 (-630-1300)
Tajikistan	2010	Male	65 to 69	79% (65-89%)	21% (11-35%)	1 (0.62-1.5)	-0.26% (-2.5-1.9%)	-0.59% (-3.4-2.1%)	-7.7 (-44-26)	-91 (-890-660)
Tajikistan	2010	Male	70 to 74	76% (63-86%)	24% (14-37%)	0.86 (0.5-1.3)	-1.2% (-3.5-0.99%)	-1.6% (-4.3-1.1%)	-35 (-100-24)	-570 (-1700-490)
Tajikistan	2010	Male	75 to 79	73% (58-85%)	27% (15-42%)	0.73 (0.39-1.2)	-1.7% (-4-0.29%)	-2.2% (-4.8-0.18%)	-43 (-99-3.7)	-560 (-1300-95)
Tajikistan	2010	Male	80 to 84	71% (54-83%)	29% (17-46%)	0.66 (0.37-1)	-2.2% (-4.4--0.062%)	-2.7% (-5.4--0.19%)	-44 (-89--3)	-430 (-890--12)
Tajikistan	2010	Male	85 to 89	69% (53-80%)	31% (20-47%)	0.6 (0.35-0.94)	-2.4% (-4.7--0.39%)	-2.9% (-5.6--0.55%)	-30 (-59--5.8)	-210 (-410--34)
Tajikistan	2010	Male	90 to 94	67% (53-80%)	33% (20-47%)	0.56 (0.3-0.93)	-2.4% (-4.7--0.49%)	-3% (-5.7--0.7%)	-8.4 (-16--1.9)	-40 (-77--7.8)
Tajikistan	2010	Male	95 plus	66% (47-82%)	34% (18-53%)	0.5 (0.22-0.91)	-2.1% (-4.5--0.33%)	-2.7% (-5.5--0.59%)	-2.3 (-4.6--0.49)	-6.9 (-14--1)
Tajikistan	2016	Female	All Ages	52% (46-60%)	48% (40-54%)	0.4 (0.29-0.54)	0.4% (-0.14-0.92%)	-0.74% (-2.3-0.63%)	-130 (-380-110)	4300 (-1400-9800)
Tajikistan	2016	Female	15 to 19	52% (39-67%)	48% (33-61%)	0.36 (0.14-0.65)	0.97% (0.53-1.6%)	1.2% (0.34-2.5%)	2.5 (0.68-5.4)	440 (230-730)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	2016	Female	20 to 24	61% (48-74%)	39% (26-52%)	0.52 (0.23-0.95)	1.8% (1.1-2.7%)	1.7% (0.47-3.4%)	4.8 (1.2-9.4)	940 (520-1400)
Tajikistan	2016	Female	25 to 29	64% (50-77%)	36% (23-50%)	0.6 (0.28-1.1)	2.1% (1.4-3%)	2.2% (0.78-4.1%)	7.9 (2.9-15)	1200 (760-1800)
Tajikistan	2016	Female	30 to 34	65% (50-78%)	35% (22-50%)	0.65 (0.33-1.1)	2.2% (1.6-3.2%)	3% (1.6-4.8%)	11 (5.4-18)	1200 (770-1700)
Tajikistan	2016	Female	35 to 39	65% (52-79%)	35% (21-48%)	0.67 (0.33-1.2)	2.2% (1.4-3.2%)	3.4% (1.8-5.3%)	12 (6.4-20)	980 (610-1400)
Tajikistan	2016	Female	40 to 44	65% (52-79%)	35% (21-48%)	0.7 (0.31-1.3)	2% (1.1-3.1%)	3.2% (1.5-5.6%)	15 (7.2-27)	930 (510-1500)
Tajikistan	2016	Female	45 to 49	65% (49-78%)	35% (22-51%)	0.73 (0.33-1.4)	1.4% (0.35-2.9%)	2.2% (0.27-4.9%)	14 (1.7-31)	730 (180-1500)
Tajikistan	2016	Female	50 to 54	63% (49-77%)	37% (23-51%)	0.74 (0.32-1.4)	0.88% (-0.37-2.6%)	1.3% (-0.87-4.1%)	12 (-8.3-38)	540 (-240-1500)
Tajikistan	2016	Female	55 to 59	62% (48-75%)	38% (25-52%)	0.75 (0.31-1.4)	0.12% (-1.5-1.9%)	0.13% (-2.3-2.8%)	1.4 (-27-33)	67 (-860-1100)
Tajikistan	2016	Female	60 to 64	60% (46-74%)	40% (26-54%)	0.73 (0.29-1.5)	-0.79% (-2.7-1.5%)	-1.1% (-3.8-2%)	-14 (-45-23)	-380 (-1300-720)
Tajikistan	2016	Female	65 to 69	58% (46-73%)	42% (27-54%)	0.68 (0.3-1.3)	-1.4% (-3.5-0.49%)	-1.9% (-4.7-0.72%)	-18 (-45-6.9)	-430 (-1000-140)
Tajikistan	2016	Female	70 to 74	55% (43-69%)	45% (31-57%)	0.61 (0.23-1.2)	-1.7% (-3.9-0.32%)	-2.2% (-5-0.41%)	-22 (-50-3.9)	-420 (-930-78)
Tajikistan	2016	Female	75 to 79	52% (40-66%)	48% (34-60%)	0.57 (0.21-1.2)	-2.1% (-4.2--0.23%)	-2.6% (-5.4--0.25%)	-39 (-80--3.5)	-570 (-1100--61)
Tajikistan	2016	Female	80 to 84	50% (37-65%)	50% (35-63%)	0.58 (0.23-1.2)	-2.5% (-4.9--0.54%)	-3.2% (-6.1--0.64%)	-46 (-88--9.7)	-480 (-920--100)
Tajikistan	2016	Female	85 to 89	50% (37-62%)	50% (38-63%)	0.61 (0.24-1.3)	-2.8% (-5--0.7%)	-3.4% (-6.2--0.88%)	-40 (-72--10)	-290 (-520--73)
Tajikistan	2016	Female	90 to 94	51% (39-63%)	49% (37-61%)	0.66 (0.22-1.5)	-2.8% (-5--0.4%)	-3.4% (-6.2--0.61%)	-25 (-44--4.3)	-120 (-220--18)
Tajikistan	2016	Female	95 plus	51% (38-66%)	49% (34-62%)	0.69 (0.18-1.7)	-2.5% (-4.9-0.38%)	-3.2% (-6-0.26%)	-7.1 (-13-0.6)	-23 (-43-3.5)
Tajikistan	2016	Male	All Ages	65% (57-73%)	35% (27-43%)	0.65 (0.5-0.82)	2.1% (1.2-3%)	0.95% (-0.56-2.5%)	230 (-150-590)	29000 (18000-41000)
Tajikistan	2016	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.3 (0.16-0.49)	2.2% (1.5-3.1%)	1.9% (0.99-3.1%)	7.4 (3.8-12)	1300 (810-1800)
Tajikistan	2016	Male	20 to 24	73% (56-85%)	27% (15-44%)	0.67 (0.42-0.99)	5.1% (3.5-6.8%)	4.4% (2.2-6.6%)	26 (13-42)	3500 (2300-5000)
Tajikistan	2016	Male	25 to 29	80% (69-89%)	20% (11-31%)	1.1 (0.7-1.6)	6.7% (5.1-8.7%)	6.5% (4.1-9.5%)	50 (30-75)	5400 (3800-7300)
Tajikistan	2016	Male	30 to 34	85% (73-92%)	15% (8-27%)	1.3 (0.89-1.8)	7.3% (5.5-9.4%)	7.7% (5-11%)	57 (35-85)	5100 (3700-6800)
Tajikistan	2016	Male	35 to 39	86% (76-93%)	14% (7.3-24%)	1.3 (0.81-1.9)	6.7% (4.8-8.8%)	7.3% (4.4-11%)	51 (30-76)	4000 (2700-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tajikistan	2016	Male	40 to 44	86% (76-93%)	14% (7-24%)	1.3 (0.82-2)	5.6% (3.6-7.8%)	6.2% (3.3-9.5%)	50 (26-76)	3300 (2100-4700)
Tajikistan	2016	Male	45 to 49	85% (72-93%)	15% (7.5-28%)	1.4 (0.87-2.1)	4.7% (2.4-6.9%)	5% (1.7-8.3%)	50 (16-88)	2900 (1500-4500)
Tajikistan	2016	Male	50 to 54	84% (72-92%)	16% (8.1-28%)	1.4 (0.82-2.1)	3.5% (1.3-5.9%)	3.6% (0.6-6.9%)	51 (8-100)	2700 (1000-4500)
Tajikistan	2016	Male	55 to 59	83% (71-91%)	17% (8.6-29%)	1.2 (0.71-1.9)	2.2% (0.081-4.3%)	2.1% (-0.8-5.1%)	33 (-13-83)	1500 (54-3100)
Tajikistan	2016	Male	60 to 64	81% (67-91%)	19% (9.4-33%)	1.1 (0.58-1.7)	0.8% (-1.3-2.9%)	0.51% (-2.2-3.3%)	8.4 (-36-55)	470 (-750-1800)
Tajikistan	2016	Male	65 to 69	79% (64-90%)	21% (10-36%)	0.93 (0.54-1.4)	-0.13% (-2.3-1.9%)	-0.49% (-3.3-2.1%)	-6 (-40-25)	-46 (-810-660)
Tajikistan	2016	Male	70 to 74	76% (62-87%)	24% (13-38%)	0.8 (0.44-1.3)	-0.94% (-3.2-1%)	-1.4% (-4.1-1.1%)	-20 (-62-16)	-310 (-1000-330)
Tajikistan	2016	Male	75 to 79	73% (57-84%)	27% (16-43%)	0.68 (0.35-1.1)	-1.5% (-3.6-0.35%)	-2% (-4.6-0.33%)	-35 (-85-5.7)	-460 (-1200-120)
Tajikistan	2016	Male	80 to 84	70% (53-82%)	30% (18-47%)	0.61 (0.32-0.98)	-2% (-4.2--0.013%)	-2.5% (-5.3--0.083%)	-43 (-93--1.5)	-420 (-920--3.1)
Tajikistan	2016	Male	85 to 89	68% (52-80%)	32% (20-48%)	0.56 (0.31-0.89)	-2.2% (-4.4--0.27%)	-2.8% (-5.5--0.47%)	-34 (-67--5.7)	-230 (-470--28)
Tajikistan	2016	Male	90 to 94	67% (52-80%)	33% (20-48%)	0.52 (0.28-0.85)	-2.2% (-4.4--0.36%)	-2.8% (-5.5--0.56%)	-17 (-33--3.4)	-78 (-160--13)
Tajikistan	2016	Male	95 plus	66% (47-82%)	34% (18-53%)	0.47 (0.21-0.86)	-1.9% (-4.1--0.24%)	-2.6% (-5.2--0.5%)	-2.2 (-4.5--0.41)	-6.4 (-14--0.75)
Turkmenistan	1990	Female	All Ages	51% (44-59%)	49% (41-56%)	0.69 (0.48-0.95)	1% (0.24-2%)	1% (-1.1-4.1%)	150 (-160-580)	8500 (2000-17000)
Turkmenistan	1990	Female	15 to 19	53% (39-68%)	47% (32-61%)	0.68 (0.27-1.3)	1.8% (0.93-3.1%)	2.9% (1.2-5.7%)	4.4 (1.7-8.4)	450 (220-780)
Turkmenistan	1990	Female	20 to 24	61% (48-74%)	39% (26-52%)	0.99 (0.4-1.8)	2.9% (1.5-4.8%)	3.8% (1.3-7.4%)	7.6 (2.6-15)	770 (390-1300)
Turkmenistan	1990	Female	25 to 29	64% (49-77%)	36% (23-51%)	1.1 (0.51-2.1)	3.4% (2-5.5%)	4.5% (2.1-8.1%)	11 (5.2-20)	1000 (590-1600)
Turkmenistan	1990	Female	30 to 34	65% (51-78%)	35% (22-49%)	1.2 (0.62-2.2)	3.7% (2.3-5.6%)	5.3% (2.7-8.9%)	13 (6.7-23)	1000 (600-1600)
Turkmenistan	1990	Female	35 to 39	65% (51-79%)	35% (21-49%)	1.3 (0.6-2.4)	3.8% (2.2-6.1%)	5.7% (2.9-10%)	13 (6.6-24)	860 (500-1400)
Turkmenistan	1990	Female	40 to 44	65% (51-78%)	35% (22-49%)	1.3 (0.59-2.4)	3.8% (1.9-6.8%)	5.9% (2.5-11%)	13 (5.3-24)	680 (330-1200)
Turkmenistan	1990	Female	45 to 49	65% (50-78%)	35% (22-50%)	1.4 (0.61-2.8)	3.7% (1.1-8.3%)	5.4% (1.2-13%)	12 (2.7-28)	540 (150-1200)
Turkmenistan	1990	Female	50 to 54	64% (49-77%)	36% (23-51%)	1.4 (0.62-2.8)	4% (0.73-9.8%)	5.7% (0.94-14%)	27 (4.4-65)	1000 (200-2500)
Turkmenistan	1990	Female	55 to 59	62% (47-75%)	38% (25-53%)	1.4 (0.6-2.9)	3.7% (-0.033-10%)	4.9% (-0.22-14%)	29 (-1.3-80)	950 (-8.4-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	1990	Female	60 to 64	60% (45-75%)	40% (25-55%)	1.4 (0.55-3)	2.7% (-1.3-9.9%)	3.4% (-1.6-13%)	31 (-15-120)	870 (-400-3200)
Turkmenistan	1990	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.3 (0.55-2.6)	1.6% (-2.2-7.8%)	2% (-2.9-9.8%)	20 (-27-95)	450 (-620-2200)
Turkmenistan	1990	Female	70 to 74	55% (42-69%)	45% (31-58%)	1.2 (0.43-2.3)	0.67% (-2.9-6.2%)	0.81% (-3.7-7.8%)	8 (-37-77)	150 (-660-1400)
Turkmenistan	1990	Female	75 to 79	52% (40-66%)	48% (34-60%)	1.1 (0.39-2.2)	-0.25% (-3.7-5.2%)	-0.38% (-4.6-6.2%)	-4.5 (-55-74)	-52 (-760-1000)
Turkmenistan	1990	Female	80 to 84	50% (37-64%)	50% (36-63%)	1.1 (0.42-2.3)	-1.1% (-4.6-4.2%)	-1.5% (-5.8-4.7%)	-15 (-59-49)	-140 (-590-540)
Turkmenistan	1990	Female	85 to 89	49% (37-62%)	51% (38-63%)	1.2 (0.45-2.4)	-1.4% (-4.9-4.2%)	-1.8% (-6.1-4.8%)	-13 (-46-37)	-91 (-320-280)
Turkmenistan	1990	Female	90 to 94	50% (39-63%)	50% (37-61%)	1.2 (0.41-2.9)	-1.4% (-5-5.3%)	-1.9% (-6.3-6.3%)	-6 (-20-20)	-27 (-98-110)
Turkmenistan	1990	Female	95 plus	51% (38-66%)	49% (34-62%)	1.3 (0.34-3.2)	-1.2% (-5.1-6.6%)	-1.6% (-6.3-7.6%)	-1.5 (-5.8-7)	-4.3 (-19-24)
Turkmenistan	1990	Male	All Ages	61% (53-70%)	39% (30-47%)	1.1 (0.77-1.4)	2.5% (1.5-3.6%)	2.7% (0.74-4.7%)	430 (120-770)	25000 (15000-36000)
Turkmenistan	1990	Male	15 to 19	56% (40-71%)	44% (29-60%)	0.57 (0.29-0.98)	3.3% (2.1-4.9%)	3.7% (1.9-6.2%)	8.9 (4.4-15)	970 (610-1400)
Turkmenistan	1990	Male	20 to 24	73% (57-85%)	27% (15-43%)	1.3 (0.76-2)	7.8% (5.4-10%)	8.4% (5-12%)	27 (16-40)	2600 (1800-3500)
Turkmenistan	1990	Male	25 to 29	80% (68-89%)	20% (11-32%)	2 (1.3-3.1)	10% (7-13%)	11% (7.2-16%)	43 (28-61)	3500 (2400-4700)
Turkmenistan	1990	Male	30 to 34	84% (73-92%)	16% (7.9-27%)	2.5 (1.6-3.6)	9.8% (6.9-13%)	11% (7.4-16%)	47 (30-66)	3400 (2400-4600)
Turkmenistan	1990	Male	35 to 39	86% (76-93%)	14% (7.4-24%)	2.5 (1.4-3.8)	9.4% (6.2-13%)	11% (6.4-16%)	46 (27-70)	2900 (1900-4200)
Turkmenistan	1990	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	2.5 (1.5-3.9)	8.9% (5.3-13%)	10% (5.2-16%)	41 (21-65)	2200 (1300-3400)
Turkmenistan	1990	Male	45 to 49	85% (73-93%)	15% (7.4-27%)	2.7 (1.5-4.2)	8.6% (4.2-14%)	9.5% (4.1-16%)	42 (17-71)	2000 (980-3200)
Turkmenistan	1990	Male	50 to 54	84% (73-92%)	16% (8.1-27%)	2.6 (1.5-4.2)	7.7% (3-13%)	8.5% (2.7-15%)	73 (23-130)	3000 (1200-5100)
Turkmenistan	1990	Male	55 to 59	83% (70-91%)	17% (8.8-30%)	2.3 (1.4-3.6)	6% (1.8-11%)	6.6% (1.5-13%)	64 (14-130)	2200 (640-4300)
Turkmenistan	1990	Male	60 to 64	81% (67-91%)	19% (9.4-33%)	2 (1.1-3.3)	4.2% (0.055-9.4%)	4.5% (-0.3-11%)	58 (-3.9-140)	1700 (22-3900)
Turkmenistan	1990	Male	65 to 69	79% (64-89%)	21% (11-36%)	1.8 (1-2.8)	2.5% (-1.2-6.6%)	2.6% (-1.7-7.3%)	24 (-16-68)	600 (-300-1600)
Turkmenistan	1990	Male	70 to 74	76% (62-86%)	24% (14-38%)	1.5 (0.83-2.4)	0.86% (-2.4-4.6%)	0.78% (-3-5.1%)	5.8 (-23-38)	130 (-370-710)
Turkmenistan	1990	Male	75 to 79	73% (57-85%)	27% (15-43%)	1.3 (0.65-2.2)	-0.74% (-3.7-2.4%)	-1.1% (-4.5-2.7%)	-9.2 (-39-23)	-100 (-520-350)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	1990	Male	80 to 84	70% (54-83%)	30% (17-46%)	1.2 (0.64-1.9)	-1.8% (-4.6-0.93%)	-2.4% (-5.6-0.91%)	-15 (-36-5.7)	-140 (-350-74)
Turkmenistan	1990	Male	85 to 89	68% (53-81%)	32% (19-47%)	1.1 (0.57-1.7)	-2.4% (-5.1-0.0088%)	-3.2% (-6.3--0.2%)	-13 (-26--0.82)	-88 (-180-0.31)
Turkmenistan	1990	Male	90 to 94	67% (52-81%)	33% (19-48%)	0.99 (0.51-1.7)	-2.7% (-5.2--0.26%)	-3.5% (-6.6--0.51%)	-5.8 (-11--0.85)	-26 (-51--2.6)
Turkmenistan	1990	Male	95 plus	66% (46-82%)	34% (18-54%)	0.9 (0.39-1.7)	-2.6% (-5.2--0.3%)	-3.5% (-6.7--0.62%)	-1.4 (-2.7--0.25)	-4.1 (-8.3--0.48)
Turkmenistan	1995	Female	All Ages	51% (45-59%)	49% (41-55%)	0.84 (0.6-1.1)	1.4% (0.36-2.8%)	1.7% (-1.1-5.5%)	260 (-170-850)	12000 (3200-24000)
Turkmenistan	1995	Female	15 to 19	52% (40-67%)	48% (33-60%)	0.81 (0.34-1.5)	2.1% (1-3.6%)	3.3% (1.2-6.4%)	6.2 (2.3-12)	600 (290-1100)
Turkmenistan	1995	Female	20 to 24	61% (48-73%)	39% (27-52%)	1.2 (0.51-2.2)	3.4% (1.6-5.8%)	4.6% (1.6-8.8%)	11 (3.8-22)	1100 (490-1900)
Turkmenistan	1995	Female	25 to 29	64% (50-76%)	36% (24-50%)	1.4 (0.62-2.5)	3.9% (2.1-6.2%)	5.2% (2.3-9.4%)	14 (6.2-26)	1200 (650-2000)
Turkmenistan	1995	Female	30 to 34	65% (50-79%)	35% (21-50%)	1.5 (0.75-2.6)	4.1% (2.4-6.7%)	5.9% (2.8-10%)	19 (9.1-33)	1400 (790-2300)
Turkmenistan	1995	Female	35 to 39	65% (52-80%)	35% (20-48%)	1.5 (0.73-2.7)	4.2% (2.3-7%)	6.5% (3-11%)	21 (9.9-37)	1300 (710-2200)
Turkmenistan	1995	Female	40 to 44	65% (51-79%)	35% (21-49%)	1.6 (0.71-2.8)	4.4% (2-7.8%)	6.9% (2.8-13%)	23 (9.1-42)	1200 (530-2100)
Turkmenistan	1995	Female	45 to 49	65% (49-79%)	35% (21-51%)	1.7 (0.72-3.2)	4.2% (0.97-9.4%)	6.3% (1.1-14%)	20 (3.7-46)	920 (220-2000)
Turkmenistan	1995	Female	50 to 54	63% (48-77%)	37% (23-52%)	1.7 (0.76-3.3)	4.6% (0.53-12%)	6.4% (0.54-16%)	23 (1.9-57)	870 (110-2100)
Turkmenistan	1995	Female	55 to 59	61% (48-75%)	39% (25-52%)	1.7 (0.7-3.5)	4.4% (-0.39-12%)	5.7% (-0.65-16%)	44 (-5-120)	1400 (-130-4000)
Turkmenistan	1995	Female	60 to 64	60% (45-75%)	40% (25-55%)	1.7 (0.67-3.5)	3.3% (-1.5-12%)	4.2% (-2.1-15%)	39 (-19-140)	1100 (-510-3800)
Turkmenistan	1995	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.5 (0.69-3)	2.3% (-2.5-9.5%)	2.8% (-3-12%)	38 (-42-160)	870 (-930-3700)
Turkmenistan	1995	Female	70 to 74	55% (43-69%)	45% (31-57%)	1.4 (0.54-2.7)	1.2% (-3.3-7.9%)	1.4% (-4-9.5%)	19 (-55-130)	360 (-980-2400)
Turkmenistan	1995	Female	75 to 79	52% (40-66%)	48% (34-60%)	1.3 (0.48-2.6)	0.31% (-3.7-6.8%)	0.29% (-4.5-8%)	3.5 (-55-100)	64 (-760-1400)
Turkmenistan	1995	Female	80 to 84	50% (37-65%)	50% (35-63%)	1.3 (0.54-2.6)	-0.44% (-4.5-5.5%)	-0.68% (-5.6-6.4%)	-7.5 (-62-72)	-61 (-620-760)
Turkmenistan	1995	Female	85 to 89	50% (38-62%)	50% (38-62%)	1.4 (0.53-2.9)	-0.64% (-4.9-6.1%)	-0.9% (-6-7.2%)	-8.4 (-55-66)	-51 (-390-500)
Turkmenistan	1995	Female	90 to 94	51% (39-63%)	49% (37-61%)	1.5 (0.5-3.4)	-0.61% (-4.9-7.5%)	-0.91% (-6.1-8.8%)	-3.6 (-24-36)	-14 (-120-190)
Turkmenistan	1995	Female	95 plus	51% (38-67%)	49% (33-62%)	1.6 (0.4-3.9)	-0.37% (-5.2-8.6%)	-0.64% (-6.5-10%)	-0.81 (-8.2-13)	-1.8 (-26-43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	1995	Male	All Ages	62% (54-71%)	38% (29-46%)	1.3 (0.99-1.7)	3.4% (2.1-4.9%)	3.7% (1.2-6.6%)	680 (230-1200)	37000 (22000-52000)
Turkmenistan	1995	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.69 (0.36-1.2)	3.6% (2.2-5.3%)	4% (2-6.7%)	13 (6.3-21)	1300 (800-1900)
Turkmenistan	1995	Male	20 to 24	73% (57-85%)	27% (15-43%)	1.6 (0.95-2.3)	8.8% (6.1-12%)	9.6% (5.8-14%)	40 (24-57)	3600 (2400-4800)
Turkmenistan	1995	Male	25 to 29	80% (68-89%)	20% (11-32%)	2.5 (1.5-3.6)	11% (8-15%)	13% (8.3-18%)	61 (39-85)	4700 (3300-6200)
Turkmenistan	1995	Male	30 to 34	84% (72-92%)	16% (7.9-28%)	3 (2-4.2)	11% (8-15%)	13% (8.5-18%)	71 (47-98)	4900 (3500-6600)
Turkmenistan	1995	Male	35 to 39	86% (75-93%)	14% (7.3-25%)	3 (1.9-4.4)	11% (7.2-16%)	13% (7.8-19%)	83 (50-120)	5000 (3300-7000)
Turkmenistan	1995	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	3.1 (1.8-4.6)	11% (6.1-16%)	12% (6.2-19%)	82 (43-130)	4300 (2600-6600)
Turkmenistan	1995	Male	45 to 49	86% (74-93%)	14% (7.4-26%)	3.3 (2-5.1)	10% (5.2-17%)	12% (5.3-20%)	77 (35-130)	3600 (1800-5800)
Turkmenistan	1995	Male	50 to 54	85% (73-92%)	15% (7.9-27%)	3.2 (1.8-5)	9.1% (3.5-16%)	10% (3.3-18%)	68 (23-120)	2700 (1000-4800)
Turkmenistan	1995	Male	55 to 59	83% (71-91%)	17% (8.8-29%)	2.8 (1.6-4.3)	6.8% (1.8-13%)	7.4% (1.5-15%)	93 (19-190)	3200 (850-6300)
Turkmenistan	1995	Male	60 to 64	81% (67-90%)	19% (9.5-33%)	2.4 (1.3-3.9)	4.8% (-0.13-11%)	5.2% (-0.58-13%)	71 (-7.8-170)	2100 (-58-4800)
Turkmenistan	1995	Male	65 to 69	79% (65-89%)	21% (11-35%)	2.2 (1.2-3.3)	2.9% (-1.4-8.3%)	3.1% (-1.9-9.2%)	52 (-30-160)	1300 (-570-3600)
Turkmenistan	1995	Male	70 to 74	76% (62-86%)	24% (14-38%)	1.8 (1-3)	1.2% (-2.8-6.2%)	1.1% (-3.4-6.8%)	12 (-37-74)	260 (-620-1400)
Turkmenistan	1995	Male	75 to 79	73% (58-85%)	27% (15-42%)	1.6 (0.84-2.6)	-0.5% (-4.1-3.5%)	-0.82% (-5-3.8%)	-6.1 (-37-28)	-59 (-480-410)
Turkmenistan	1995	Male	80 to 84	71% (54-83%)	29% (17-46%)	1.4 (0.79-2.2)	-1.7% (-5-1.7%)	-2.3% (-6.3-1.7%)	-14 (-39-11)	-130 (-370-130)
Turkmenistan	1995	Male	85 to 89	68% (52-80%)	32% (20-48%)	1.3 (0.71-2.1)	-2.4% (-5.5-0.56%)	-3.1% (-6.7-0.36%)	-15 (-32-1.7)	-97 (-220-22)
Turkmenistan	1995	Male	90 to 94	67% (53-80%)	33% (20-47%)	1.2 (0.63-2)	-2.7% (-5.5-0.042%)	-3.5% (-6.9-0.15%)	-6.7 (-13--0.29)	-30 (-62-0.48)
Turkmenistan	1995	Male	95 plus	66% (47-82%)	34% (18-53%)	1.1 (0.47-2)	-2.7% (-5.5--0.016%)	-3.5% (-7--0.38%)	-1.9 (-3.7--0.2)	-5.6 (-12--0.032)
Turkmenistan	2000	Female	All Ages	52% (46-60%)	48% (40-54%)	1 (0.75-1.4)	2.2% (0.82-4%)	2.8% (-0.57-7.7%)	430 (-82-1200)	18000 (6700-33000)
Turkmenistan	2000	Female	15 to 19	52% (40-67%)	48% (33-60%)	0.97 (0.41-1.8)	2.5% (1.1-4.5%)	4.1% (1.5-7.8%)	9.1 (3.3-18)	830 (380-1500)
Turkmenistan	2000	Female	20 to 24	61% (49-73%)	39% (27-51%)	1.4 (0.61-2.6)	4.2% (2-7.2%)	6% (2.2-11%)	17 (6.2-31)	1500 (680-2600)
Turkmenistan	2000	Female	25 to 29	64% (49-76%)	36% (24-51%)	1.6 (0.72-3)	4.9% (2.6-8.2%)	7% (3.1-13%)	21 (9.3-38)	1700 (860-2800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	2000	Female	30 to 34	65% (50-78%)	35% (22-50%)	1.8 (0.89-3.1)	5.1% (2.9-8.2%)	7.7% (3.8-13%)	25 (12-42)	1800 (940-2800)
Turkmenistan	2000	Female	35 to 39	65% (52-79%)	35% (21-48%)	1.8 (0.85-3.2)	5% (2.6-8.2%)	8% (3.7-14%)	30 (14-52)	1800 (940-3000)
Turkmenistan	2000	Female	40 to 44	65% (51-79%)	35% (21-49%)	1.9 (0.87-3.4)	5.3% (2.4-9.2%)	8.5% (3.6-15%)	36 (15-62)	1900 (850-3200)
Turkmenistan	2000	Female	45 to 49	65% (49-79%)	35% (21-51%)	2 (0.87-3.8)	5.3% (1.5-11%)	7.9% (1.9-17%)	37 (9-77)	1700 (490-3400)
Turkmenistan	2000	Female	50 to 54	64% (49-77%)	36% (23-51%)	2 (0.87-3.9)	5.8% (0.95-13%)	8.1% (1.2-18%)	39 (6.2-86)	1500 (250-3300)
Turkmenistan	2000	Female	55 to 59	62% (48-76%)	38% (24-52%)	2.1 (0.85-4.2)	5.7% (-0.039-14%)	7.6% (-0.16-19%)	39 (-0.8-96)	1300 (-8.5-3100)
Turkmenistan	2000	Female	60 to 64	60% (46-75%)	40% (25-54%)	2 (0.8-4.2)	4.6% (-1.5-14%)	5.7% (-2-17%)	62 (-21-190)	1700 (-530-5100)
Turkmenistan	2000	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.9 (0.81-3.7)	3.4% (-2.2-12%)	4.2% (-2.7-15%)	51 (-34-180)	1200 (-750-4100)
Turkmenistan	2000	Female	70 to 74	55% (43-68%)	45% (32-57%)	1.6 (0.65-3.3)	2.1% (-3.2-10%)	2.4% (-4-12%)	42 (-68-210)	790 (-1200-3900)
Turkmenistan	2000	Female	75 to 79	52% (40-65%)	48% (35-60%)	1.5 (0.56-3.1)	1.1% (-3.7-8.8%)	1.2% (-4.7-10%)	18 (-69-150)	280 (-950-2100)
Turkmenistan	2000	Female	80 to 84	50% (37-65%)	50% (35-63%)	1.6 (0.62-3.2)	0.4% (-4.2-7.8%)	0.3% (-5.3-9.3%)	3.2 (-57-100)	55 (-550-1100)
Turkmenistan	2000	Female	85 to 89	50% (37-62%)	50% (38-63%)	1.7 (0.62-3.5)	0.3% (-4.5-8.6%)	0.17% (-5.7-10%)	1.4 (-47-85)	22 (-330-640)
Turkmenistan	2000	Female	90 to 94	50% (39-62%)	50% (38-61%)	1.8 (0.59-4.1)	0.35% (-4.8-9.7%)	0.21% (-6-12%)	0.89 (-26-49)	9.1 (-120-250)
Turkmenistan	2000	Female	95 plus	51% (37-66%)	49% (34-63%)	1.9 (0.45-4.8)	0.56% (-5-10%)	0.44% (-6.3-12%)	0.58 (-8.2-16)	3 (-26-54)
Turkmenistan	2000	Male	All Ages	64% (55-72%)	36% (28-45%)	1.7 (1.3-2.1)	5.1% (3.2-7.3%)	5.6% (2.5-9.4%)	1100 (470-1800)	55000 (35000-76000)
Turkmenistan	2000	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.83 (0.43-1.4)	4% (2.3-6.1%)	4.5% (2.2-7.8%)	17 (8.4-30)	1700 (990-2600)
Turkmenistan	2000	Male	20 to 24	73% (57-85%)	27% (15-43%)	1.8 (1.1-2.8)	10% (6.8-14%)	11% (6.9-16%)	58 (35-85)	4900 (3300-6700)
Turkmenistan	2000	Male	25 to 29	80% (69-89%)	20% (11-31%)	2.9 (1.8-4.4)	14% (9.5-18%)	16% (10-23%)	98 (62-140)	7100 (4800-9600)
Turkmenistan	2000	Male	30 to 34	84% (72-92%)	16% (7.7-28%)	3.5 (2.3-5.1)	14% (9.7-19%)	16% (11-23%)	110 (73-160)	7300 (5100-10000)
Turkmenistan	2000	Male	35 to 39	86% (75-93%)	14% (7.2-25%)	3.5 (2.2-5.3)	13% (9.1-19%)	16% (10-23%)	130 (81-180)	7400 (5000-10000)
Turkmenistan	2000	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	3.6 (2.2-5.4)	13% (7.9-19%)	15% (8.6-23%)	140 (78-220)	7300 (4300-11000)
Turkmenistan	2000	Male	45 to 49	86% (73-93%)	14% (7.3-27%)	3.9 (2.4-6)	13% (7-20%)	15% (7.5-23%)	140 (71-220)	6400 (3600-9800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	2000	Male	50 to 54	85% (73-92%)	15% (8-27%)	3.8 (2.2-5.9)	11% (4.6-19%)	13% (4.5-22%)	110 (42-200)	4500 (1800-7600)
Turkmenistan	2000	Male	55 to 59	83% (71-92%)	17% (8.3-29%)	3.3 (1.9-5.2)	8.5% (2.6-16%)	9.4% (2.4-18%)	81 (20-160)	2800 (820-5300)
Turkmenistan	2000	Male	60 to 64	81% (68-91%)	19% (9.3-32%)	2.9 (1.6-4.8)	6.3% (0.49-14%)	6.8% (0.12-16%)	110 (2-260)	3100 (240-7100)
Turkmenistan	2000	Male	65 to 69	79% (65-90%)	21% (10-35%)	2.6 (1.5-4)	4.1% (-1.1-11%)	4.4% (-1.6-12%)	69 (-25-190)	1600 (-440-4300)
Turkmenistan	2000	Male	70 to 74	76% (61-86%)	24% (14-39%)	2.2 (1.2-3.6)	1.8% (-2.9-7.8%)	1.9% (-3.5-8.7%)	33 (-61-160)	670 (-1000-2900)
Turkmenistan	2000	Male	75 to 79	73% (57-85%)	27% (15-43%)	1.9 (1-3.2)	0.21% (-4.1-5.2%)	-0.038% (-4.9-5.8%)	-0.34 (-47-56)	32 (-610-830)
Turkmenistan	2000	Male	80 to 84	71% (54-84%)	29% (16-46%)	1.7 (0.94-2.7)	-1.2% (-4.9-3.2%)	-1.8% (-6.2-3.5%)	-9.2 (-33-18)	-75 (-320-210)
Turkmenistan	2000	Male	85 to 89	69% (53-81%)	31% (19-47%)	1.6 (0.85-2.5)	-2% (-5.5-1.5%)	-2.7% (-6.7-1.5%)	-11 (-26-5.8)	-67 (-180-52)
Turkmenistan	2000	Male	90 to 94	68% (53-80%)	32% (20-47%)	1.4 (0.75-2.5)	-2.5% (-5.6-0.75%)	-3.3% (-7.1-0.54%)	-6.2 (-13-1)	-27 (-61-8.3)
Turkmenistan	2000	Male	95 plus	67% (48-82%)	33% (18-52%)	1.3 (0.58-2.4)	-2.5% (-5.6-0.57%)	-3.4% (-7.1-0.22%)	-1.8 (-3.7-0.11)	-5.1 (-11-1.2)
Turkmenistan	2005	Female	All Ages	53% (47-61%)	47% (39-53%)	1.2 (0.83-1.6)	2.7% (1.1-4.6%)	3.5% (-0.14-8.8%)	520 (-20-1300)	21000 (8800-36000)
Turkmenistan	2005	Female	15 to 19	52% (40-67%)	48% (33-60%)	1 (0.42-1.9)	2.6% (1.2-4.6%)	4.2% (1.5-8%)	10 (3.7-20)	950 (420-1700)
Turkmenistan	2005	Female	20 to 24	61% (48-73%)	39% (27-52%)	1.5 (0.61-2.6)	4.4% (2.1-7.7%)	6.5% (2.3-12%)	19 (6.7-36)	1700 (770-2900)
Turkmenistan	2005	Female	25 to 29	64% (49-77%)	36% (23-51%)	1.7 (0.76-3.2)	5.3% (2.9-8.7%)	7.8% (3.6-14%)	24 (11-43)	1900 (1000-3100)
Turkmenistan	2005	Female	30 to 34	65% (50-78%)	35% (22-50%)	1.8 (0.92-3.4)	5.5% (3.3-8.8%)	8.5% (4.4-15%)	28 (15-48)	2000 (1100-3100)
Turkmenistan	2005	Female	35 to 39	65% (52-80%)	35% (20-48%)	1.9 (0.89-3.3)	5.5% (2.9-8.9%)	9% (4.4-15%)	30 (15-51)	1900 (1000-3000)
Turkmenistan	2005	Female	40 to 44	65% (51-79%)	35% (21-49%)	2 (0.89-3.6)	5.8% (2.7-9.6%)	9.5% (4.3-16%)	43 (20-71)	2200 (1100-3600)
Turkmenistan	2005	Female	45 to 49	65% (49-78%)	35% (22-51%)	2.1 (0.92-4)	5.7% (1.9-12%)	8.9% (2.5-18%)	51 (15-100)	2300 (750-4500)
Turkmenistan	2005	Female	50 to 54	63% (48-77%)	37% (23-52%)	2.1 (0.93-4.1)	6.1% (1.4-13%)	8.9% (2.1-18%)	57 (13-120)	2200 (520-4500)
Turkmenistan	2005	Female	55 to 59	61% (48-75%)	39% (25-52%)	2.2 (0.9-4.3)	6.2% (0.45-14%)	8.4% (0.52-20%)	55 (3.5-130)	1800 (130-4300)
Turkmenistan	2005	Female	60 to 64	60% (45-74%)	40% (26-55%)	2.1 (0.82-4.3)	5.2% (-1.2-15%)	6.5% (-1.6-18%)	45 (-11-120)	1200 (-270-3400)
Turkmenistan	2005	Female	65 to 69	58% (46-73%)	42% (27-54%)	1.9 (0.87-3.8)	4% (-1.9-13%)	5% (-2.3-16%)	69 (-32-220)	1600 (-710-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	2005	Female	70 to 74	55% (43-69%)	45% (31-57%)	1.7 (0.69-3.4)	2.6% (-3.1-11%)	3.1% (-3.8-14%)	46 (-57-200)	860 (-1000-3700)
Turkmenistan	2005	Female	75 to 79	52% (40-65%)	48% (35-60%)	1.6 (0.62-3.2)	1.5% (-3.8-9.4%)	1.7% (-4.8-11%)	29 (-85-200)	440 (-1200-2800)
Turkmenistan	2005	Female	80 to 84	50% (37-64%)	50% (36-63%)	1.7 (0.66-3.3)	0.78% (-3.8-8.3%)	0.78% (-5-10%)	9.5 (-60-120)	120 (-590-1300)
Turkmenistan	2005	Female	85 to 89	49% (38-61%)	51% (39-62%)	1.7 (0.66-3.7)	0.66% (-4.3-8.8%)	0.6% (-5.5-11%)	4.7 (-43-85)	46 (-300-630)
Turkmenistan	2005	Female	90 to 94	50% (39-62%)	50% (38-61%)	1.9 (0.63-4.2)	0.65% (-4.6-10%)	0.57% (-5.9-12%)	2.1 (-21-46)	15 (-100-230)
Turkmenistan	2005	Female	95 plus	51% (38-66%)	49% (34-62%)	1.9 (0.49-4.9)	0.8% (-4.9-11%)	0.72% (-6.2-13%)	0.96 (-8.1-17)	4.3 (-26-56)
Turkmenistan	2005	Male	All Ages	65% (57-74%)	35% (26-43%)	1.9 (1.4-2.4)	6.1% (4-8.5%)	6.7% (3.2-11%)	1300 (630-2100)	64000 (42000-88000)
Turkmenistan	2005	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.87 (0.46-1.5)	4.1% (2.5-6.2%)	4.8% (2.5-7.9%)	20 (10-33)	2000 (1200-3000)
Turkmenistan	2005	Male	20 to 24	73% (58-85%)	27% (15-42%)	1.9 (1.2-2.9)	10% (7-14%)	12% (7.2-17%)	61 (37-88)	5200 (3600-7100)
Turkmenistan	2005	Male	25 to 29	80% (69-89%)	20% (11-31%)	3.1 (1.9-4.6)	14% (9.9-19%)	16% (11-23%)	110 (68-150)	7700 (5300-10000)
Turkmenistan	2005	Male	30 to 34	85% (73-92%)	15% (7.8-27%)	3.7 (2.4-5.3)	14% (10-19%)	17% (11-23%)	130 (87-180)	8200 (5900-11000)
Turkmenistan	2005	Male	35 to 39	86% (76-93%)	14% (7.3-24%)	3.7 (2.3-5.4)	14% (10-20%)	17% (11-24%)	140 (92-200)	8100 (5600-11000)
Turkmenistan	2005	Male	40 to 44	86% (76-93%)	14% (6.9-24%)	3.8 (2.3-5.5)	14% (8.8-20%)	16% (9.7-24%)	160 (96-240)	8300 (5300-12000)
Turkmenistan	2005	Male	45 to 49	86% (73-93%)	14% (7.4-27%)	4.1 (2.5-6.1)	14% (7.9-21%)	16% (8.5-25%)	190 (100-290)	8500 (4800-13000)
Turkmenistan	2005	Male	50 to 54	85% (73-92%)	15% (7.7-27%)	4 (2.3-6)	13% (5.8-20%)	14% (5.9-23%)	170 (70-280)	6600 (3000-11000)
Turkmenistan	2005	Male	55 to 59	83% (70-91%)	17% (8.5-30%)	3.5 (2.1-5.3)	9.7% (3.6-18%)	11% (3.5-21%)	120 (38-220)	4000 (1500-7300)
Turkmenistan	2005	Male	60 to 64	81% (67-90%)	19% (9.5-33%)	3 (1.6-5)	7.4% (1.3-16%)	8.2% (1.1-18%)	86 (11-190)	2400 (430-5200)
Turkmenistan	2005	Male	65 to 69	79% (64-89%)	21% (11-36%)	2.7 (1.6-4.2)	4.9% (-0.71-12%)	5.3% (-1.2-14%)	93 (-20-240)	2200 (-320-5600)
Turkmenistan	2005	Male	70 to 74	76% (62-86%)	24% (14-38%)	2.3 (1.3-3.8)	2.5% (-2.6-9.1%)	2.6% (-3.2-10%)	42 (-53-170)	840 (-870-3100)
Turkmenistan	2005	Male	75 to 79	73% (58-84%)	27% (16-42%)	2 (1.1-3.3)	0.69% (-3.6-6.5%)	0.52% (-4.5-7.4%)	7.9 (-69-110)	170 (-890-1600)
Turkmenistan	2005	Male	80 to 84	71% (55-83%)	29% (17-45%)	1.8 (0.98-2.8)	-0.77% (-4.7-3.6%)	-1.3% (-6.1-3.9%)	-9.4 (-44-28)	-68 (-420-320)
Turkmenistan	2005	Male	85 to 89	68% (53-80%)	32% (20-47%)	1.6 (0.91-2.6)	-1.7% (-5.1-2.3%)	-2.4% (-6.5-2.4%)	-7.1 (-19-7.1)	-43 (-130-59)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	2005	Male	90 to 94	67% (52-80%)	33% (20-48%)	1.5 (0.79-2.6)	-2.3% (-5.5-1.2%)	-3.1% (-6.9-1%)	-4.1 (-9.1-1.4)	-18 (-42-9)
Turkmenistan	2005	Male	95 plus	66% (47-82%)	34% (18-53%)	1.4 (0.61-2.6)	-2.3% (-5.5-1.2%)	-3.2% (-7.2-0.88%)	-1.4 (-3.1-0.37)	-3.9 (-9.3-1.9)
Turkmenistan	2010	Female	All Ages	54% (47-61%)	46% (39-53%)	1.2 (0.84-1.6)	2.8% (1.2-4.8%)	3.6% (-0.033-8.6%)	520 (-4.8-1200)	21000 (9400-36000)
Turkmenistan	2010	Female	15 to 19	52% (39-68%)	48% (32-61%)	0.96 (0.39-1.8)	2.3% (1-4.1%)	3.7% (1.2-7.5%)	8.4 (2.7-17)	810 (360-1500)
Turkmenistan	2010	Female	20 to 24	61% (47-74%)	39% (26-53%)	1.4 (0.58-2.6)	4.1% (2-7%)	6% (2.2-11%)	17 (6.1-32)	1600 (760-2800)
Turkmenistan	2010	Female	25 to 29	64% (49-77%)	36% (23-51%)	1.6 (0.76-3.1)	4.9% (2.8-7.8%)	7.6% (3.7-13%)	22 (11-38)	1800 (1000-2900)
Turkmenistan	2010	Female	30 to 34	65% (50-77%)	35% (23-50%)	1.8 (0.91-3.3)	5.3% (3.3-8.4%)	8.7% (4.9-14%)	27 (15-43)	1900 (1200-2900)
Turkmenistan	2010	Female	35 to 39	65% (52-79%)	35% (21-48%)	1.8 (0.85-3.3)	5.4% (3-8.5%)	9.2% (4.7-15%)	31 (16-49)	1900 (1100-2900)
Turkmenistan	2010	Female	40 to 44	65% (52-78%)	35% (22-48%)	1.9 (0.87-3.5)	5.6% (3-9.2%)	9.6% (4.7-16%)	39 (19-64)	2000 (1100-3300)
Turkmenistan	2010	Female	45 to 49	65% (49-79%)	35% (21-51%)	2 (0.9-3.8)	5.4% (2-10%)	8.7% (3-17%)	52 (18-100)	2400 (860-4500)
Turkmenistan	2010	Female	50 to 54	63% (49-77%)	37% (23-51%)	2.1 (0.91-3.9)	5.7% (1.4-12%)	8.5% (2-17%)	65 (15-130)	2500 (610-5100)
Turkmenistan	2010	Female	55 to 59	61% (48-75%)	39% (25-52%)	2.1 (0.88-4.1)	5.7% (0.35-13%)	7.9% (0.36-18%)	70 (3.1-160)	2300 (140-5300)
Turkmenistan	2010	Female	60 to 64	60% (46-75%)	40% (25-54%)	2 (0.8-4.1)	4.8% (-0.92-13%)	6.2% (-1.3-17%)	55 (-11-150)	1500 (-300-4200)
Turkmenistan	2010	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.9 (0.85-3.6)	3.8% (-1.7-12%)	4.8% (-2.2-15%)	41 (-19-130)	950 (-410-3100)
Turkmenistan	2010	Female	70 to 74	55% (42-69%)	45% (31-58%)	1.7 (0.65-3.2)	2.6% (-2.9-10%)	3.1% (-3.6-13%)	51 (-59-210)	940 (-1100-3800)
Turkmenistan	2010	Female	75 to 79	52% (40-65%)	48% (35-60%)	1.5 (0.57-3.2)	1.4% (-3.5-8.7%)	1.6% (-4.5-11%)	24 (-67-160)	370 (-930-2200)
Turkmenistan	2010	Female	80 to 84	50% (37-64%)	50% (36-63%)	1.6 (0.62-3.1)	0.73% (-3.7-7.6%)	0.73% (-4.9-9.6%)	8.8 (-58-110)	120 (-590-1200)
Turkmenistan	2010	Female	85 to 89	49% (37-62%)	51% (38-63%)	1.7 (0.63-3.5)	0.57% (-4.1-8.3%)	0.5% (-5.3-10%)	5 (-53-100)	53 (-380-810)
Turkmenistan	2010	Female	90 to 94	50% (38-62%)	50% (38-62%)	1.8 (0.61-4)	0.51% (-4.4-9.3%)	0.41% (-5.8-11%)	1.6 (-23-45)	13 (-110-240)
Turkmenistan	2010	Female	95 plus	51% (36-66%)	49% (34-64%)	1.9 (0.48-4.7)	0.63% (-4.7-9.7%)	0.51% (-6.1-12%)	0.63 (-7.6-15)	3.3 (-24-50)
Turkmenistan	2010	Male	All Ages	66% (58-74%)	34% (26-42%)	1.9 (1.4-2.4)	6.4% (4.2-8.8%)	6.8% (3.4-11%)	1200 (620-2000)	62000 (42000-85000)
Turkmenistan	2010	Male	15 to 19	55% (40-72%)	45% (28-60%)	0.81 (0.42-1.4)	3.8% (2.5-5.5%)	4.4% (2.3-7.1%)	15 (7.6-24)	1600 (990-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	2010	Male	20 to 24	73% (57-85%)	27% (15-43%)	1.8 (1.1-2.8)	9.6% (6.8-13%)	11% (7-15%)	55 (35-79)	5100 (3500-6900)
Turkmenistan	2010	Male	25 to 29	80% (69-89%)	20% (11-31%)	3 (1.8-4.4)	13% (9.3-17%)	15% (10-21%)	88 (59-120)	6800 (4800-8800)
Turkmenistan	2010	Male	30 to 34	84% (72-92%)	16% (7.9-28%)	3.6 (2.3-5.2)	13% (9.7-18%)	16% (11-22%)	110 (73-150)	7200 (5200-9600)
Turkmenistan	2010	Male	35 to 39	86% (75-93%)	14% (7.3-25%)	3.5 (2.2-5.3)	14% (9.9-20%)	17% (11-24%)	130 (88-190)	7700 (5500-10000)
Turkmenistan	2010	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	3.6 (2.2-5.4)	14% (9.1-19%)	16% (11-24%)	150 (95-210)	7700 (5200-11000)
Turkmenistan	2010	Male	45 to 49	85% (72-93%)	15% (7.5-28%)	3.9 (2.4-5.8)	13% (7.8-20%)	16% (8.7-24%)	180 (100-280)	8400 (5000-12000)
Turkmenistan	2010	Male	50 to 54	84% (73-92%)	16% (8.2-27%)	3.8 (2.2-5.7)	12% (5.9-19%)	14% (6.1-22%)	190 (87-320)	7800 (3800-12000)
Turkmenistan	2010	Male	55 to 59	83% (71-91%)	17% (8.6-29%)	3.3 (2-5.1)	9.3% (3.5-17%)	11% (3.5-20%)	150 (50-280)	5100 (1900-9200)
Turkmenistan	2010	Male	60 to 64	81% (67-91%)	19% (9.3-33%)	2.9 (1.6-4.8)	7.1% (1.4-15%)	7.9% (1.1-18%)	100 (14-220)	2900 (570-6100)
Turkmenistan	2010	Male	65 to 69	79% (65-89%)	21% (11-35%)	2.6 (1.4-4)	4.7% (-0.43-11%)	5.1% (-0.87-13%)	59 (-9.8-150)	1400 (-130-3400)
Turkmenistan	2010	Male	70 to 74	76% (62-86%)	24% (14-38%)	2.2 (1.3-3.6)	2.4% (-2.4-8.8%)	2.5% (-3.1-10%)	43 (-54-170)	870 (-860-3200)
Turkmenistan	2010	Male	75 to 79	73% (58-85%)	27% (15-42%)	1.9 (1-3.2)	0.75% (-3.4-6.2%)	0.59% (-4.3-7%)	8.1 (-60-97)	170 (-780-1400)
Turkmenistan	2010	Male	80 to 84	71% (54-83%)	29% (17-46%)	1.7 (0.94-2.7)	-0.79% (-4.5-3.3%)	-1.3% (-5.8-3.7%)	-11 (-51-32)	-86 (-490-370)
Turkmenistan	2010	Male	85 to 89	69% (53-80%)	31% (20-47%)	1.6 (0.87-2.5)	-1.7% (-5-2%)	-2.4% (-6.3-2%)	-12 (-32-10)	-74 (-220-86)
Turkmenistan	2010	Male	90 to 94	67% (52-80%)	33% (20-48%)	1.4 (0.76-2.4)	-2.2% (-5.1-1.1%)	-3% (-6.6-0.93%)	-4 (-8.7-1.2)	-17 (-40-8.5)
Turkmenistan	2010	Male	95 plus	66% (47-82%)	34% (18-53%)	1.3 (0.57-2.5)	-2.2% (-5.2-0.98%)	-3.2% (-6.8-0.73%)	-1.1 (-2.4-0.25)	-3.2 (-7.5-1.4)
Turkmenistan	2016	Female	All Ages	54% (48-62%)	46% (38-52%)	1.2 (0.8-1.7)	2.9% (1.2-5.1%)	3.6% (-0.26-8.9%)	520 (-36-1300)	21000 (8800-37000)
Turkmenistan	2016	Female	15 to 19	52% (39-67%)	48% (33-61%)	0.95 (0.36-1.9)	2.1% (0.98-3.7%)	3.6% (1.2-7.3%)	5.3 (1.8-11)	550 (260-1000)
Turkmenistan	2016	Female	20 to 24	61% (47-73%)	39% (27-53%)	1.4 (0.56-2.7)	3.7% (1.9-6.2%)	5.8% (2.1-11%)	13 (4.5-24)	1300 (680-2200)
Turkmenistan	2016	Female	25 to 29	64% (50-77%)	36% (23-50%)	1.6 (0.7-3)	4.5% (2.6-7%)	7.3% (3.6-13%)	20 (9.3-34)	1800 (1000-2800)
Turkmenistan	2016	Female	30 to 34	65% (51-78%)	35% (22-49%)	1.8 (0.85-3.2)	5% (3.2-7.6%)	8.6% (4.8-14%)	25 (14-40)	1900 (1200-2900)
Turkmenistan	2016	Female	35 to 39	65% (52-79%)	35% (21-48%)	1.8 (0.8-3.3)	5.1% (3-8%)	9.3% (5-15%)	30 (16-48)	1900 (1100-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	2016	Female	40 to 44	65% (51-79%)	35% (21-49%)	1.9 (0.79-3.5)	5.4% (2.9-8.8%)	9.6% (4.8-16%)	38 (19-64)	2000 (1100-3300)
Turkmenistan	2016	Female	45 to 49	65% (49-79%)	35% (21-51%)	2 (0.84-3.7)	5.1% (1.9-9.9%)	8.5% (3.1-16%)	45 (16-86)	2000 (800-3900)
Turkmenistan	2016	Female	50 to 54	63% (49-76%)	37% (24-51%)	2 (0.9-3.9)	5.2% (1.3-11%)	8.1% (2-17%)	65 (16-140)	2600 (610-5300)
Turkmenistan	2016	Female	55 to 59	61% (48-75%)	39% (25-52%)	2 (0.81-3.9)	5.2% (0.4-12%)	7.5% (0.54-17%)	81 (5.7-180)	2700 (210-6200)
Turkmenistan	2016	Female	60 to 64	60% (46-75%)	40% (25-54%)	2 (0.77-4)	4.3% (-1-12%)	5.8% (-1.4-16%)	72 (-18-200)	2000 (-470-5600)
Turkmenistan	2016	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.8 (0.81-3.5)	3.4% (-1.7-11%)	4.4% (-2.2-15%)	52 (-27-170)	1200 (-600-4000)
Turkmenistan	2016	Female	70 to 74	55% (43-69%)	45% (31-57%)	1.6 (0.62-3.2)	2.3% (-2.7-10%)	2.9% (-3.5-13%)	29 (-36-130)	550 (-630-2400)
Turkmenistan	2016	Female	75 to 79	52% (40-65%)	48% (35-60%)	1.5 (0.56-3.2)	1.3% (-3.4-8.6%)	1.5% (-4.4-11%)	23 (-66-160)	350 (-900-2300)
Turkmenistan	2016	Female	80 to 84	50% (37-64%)	50% (36-63%)	1.6 (0.61-3.2)	0.69% (-3.7-7.4%)	0.69% (-5-9.4%)	9 (-67-130)	120 (-680-1400)
Turkmenistan	2016	Female	85 to 89	49% (36-61%)	51% (39-64%)	1.7 (0.63-3.6)	0.56% (-4.2-8.5%)	0.5% (-5.4-11%)	5.2 (-57-110)	55 (-400-840)
Turkmenistan	2016	Female	90 to 94	50% (39-62%)	50% (38-61%)	1.8 (0.6-4.2)	0.54% (-4.4-9.4%)	0.44% (-5.8-11%)	2.7 (-35-69)	21 (-170-350)
Turkmenistan	2016	Female	95 plus	51% (37-66%)	49% (34-63%)	1.9 (0.47-4.9)	0.67% (-4.8-9.7%)	0.55% (-6.1-12%)	0.94 (-10-20)	4.8 (-33-69)
Turkmenistan	2016	Male	All Ages	67% (60-75%)	33% (25-40%)	2 (1.4-2.7)	6.9% (4.3-9.9%)	7% (3.1-12%)	1300 (590-2200)	65000 (42000-93000)
Turkmenistan	2016	Male	15 to 19	55% (39-71%)	45% (29-61%)	0.8 (0.38-1.5)	3.8% (2.5-5.7%)	4.4% (2.3-7.5%)	12 (6-19)	1300 (810-1900)
Turkmenistan	2016	Male	20 to 24	73% (56-85%)	27% (15-44%)	1.8 (0.99-2.9)	9.5% (6.7-13%)	11% (6.6-15%)	52 (32-77)	4900 (3400-6800)
Turkmenistan	2016	Male	25 to 29	80% (69-89%)	20% (11-31%)	2.9 (1.7-4.5)	13% (9-17%)	15% (9.6-21%)	95 (61-130)	7500 (5400-10000)
Turkmenistan	2016	Male	30 to 34	85% (73-92%)	15% (8.2-27%)	3.5 (2.2-5.3)	13% (9.6-17%)	16% (11-22%)	110 (73-150)	7600 (5400-10000)
Turkmenistan	2016	Male	35 to 39	86% (76-93%)	14% (7.3-24%)	3.5 (2.1-5.5)	14% (10-19%)	17% (11-24%)	130 (87-180)	7800 (5600-11000)
Turkmenistan	2016	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	3.6 (2.1-5.6)	14% (9.2-19%)	17% (11-24%)	140 (92-210)	7700 (5100-11000)
Turkmenistan	2016	Male	45 to 49	85% (72-92%)	15% (7.6-28%)	3.8 (2.2-6)	13% (7.8-20%)	16% (8.8-24%)	170 (94-270)	8000 (4700-12000)
Turkmenistan	2016	Male	50 to 54	84% (72-92%)	16% (8.3-28%)	3.8 (2.1-6)	12% (5.9-19%)	14% (6.3-22%)	200 (89-330)	8100 (4000-13000)
Turkmenistan	2016	Male	55 to 59	83% (71-91%)	17% (8.6-29%)	3.3 (1.8-5.3)	9.1% (3.5-17%)	10% (3.6-20%)	180 (61-340)	6200 (2400-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkmenistan	2016	Male	60 to 64	81% (67-90%)	19% (9.6-33%)	2.8 (1.4-4.8)	6.9% (1.1-15%)	7.8% (0.87-18%)	140 (16-310)	4000 (640-8600)
Turkmenistan	2016	Male	65 to 69	79% (64-89%)	21% (11-36%)	2.5 (1.4-4.1)	4.7% (-0.4-11%)	5.2% (-0.84-13%)	77 (-13-200)	1900 (-160-4500)
Turkmenistan	2016	Male	70 to 74	76% (62-87%)	24% (13-38%)	2.2 (1.2-3.7)	2.4% (-2.2-8.7%)	2.5% (-2.8-10%)	30 (-34-120)	600 (-550-2200)
Turkmenistan	2016	Male	75 to 79	73% (57-84%)	27% (16-43%)	1.9 (0.95-3.2)	0.72% (-3.3-6.2%)	0.54% (-4.3-7.1%)	7.7 (-62-100)	170 (-800-1500)
Turkmenistan	2016	Male	80 to 84	70% (54-83%)	30% (17-46%)	1.7 (0.86-2.7)	-0.91% (-4.6-3.2%)	-1.5% (-5.8-3.4%)	-15 (-61-36)	-120 (-590-420)
Turkmenistan	2016	Male	85 to 89	68% (52-80%)	32% (20-48%)	1.5 (0.79-2.6)	-1.8% (-5.2-1.8%)	-2.5% (-6.5-1.7%)	-16 (-42-11)	-100 (-280-100)
Turkmenistan	2016	Male	90 to 94	67% (52-80%)	33% (20-48%)	1.4 (0.7-2.4)	-2.3% (-5.2-0.9%)	-3.1% (-6.7-0.73%)	-8.5 (-18-2)	-37 (-85-15)
Turkmenistan	2016	Male	95 plus	66% (47-82%)	34% (18-53%)	1.3 (0.53-2.4)	-2.3% (-5.3-0.91%)	-3.2% (-6.9-0.57%)	-1.5 (-3.1-0.26)	-4.2 (-9.7-1.7)
Uzbekistan	1990	Female	All Ages	50% (44-58%)	50% (42-56%)	0.66 (0.46-0.92)	1% (0.06-2.2%)	0.68% (-1.8-4.1%)	440 (-1200-2700)	34000 (2100-75000)
Uzbekistan	1990	Female	15 to 19	52% (39-68%)	48% (32-61%)	0.67 (0.27-1.3)	1.3% (0.64-2.3%)	2.2% (0.6-4.4%)	15 (4-30)	1800 (830-3100)
Uzbekistan	1990	Female	20 to 24	61% (48-74%)	39% (26-52%)	0.97 (0.42-1.8)	2.2% (1.1-3.7%)	2.8% (0.72-5.6%)	26 (6.6-54)	3200 (1500-5400)
Uzbekistan	1990	Female	25 to 29	64% (49-77%)	36% (23-51%)	1.1 (0.51-2)	2.6% (1.5-4.4%)	3.6% (1.1-6.9%)	38 (12-74)	4000 (2100-6600)
Uzbekistan	1990	Female	30 to 34	65% (51-78%)	35% (22-49%)	1.2 (0.62-2.2)	2.9% (1.6-4.6%)	4.4% (1.9-7.8%)	50 (21-89)	4200 (2400-6600)
Uzbekistan	1990	Female	35 to 39	65% (52-79%)	35% (21-48%)	1.3 (0.55-2.4)	3% (1.7-5.3%)	5% (2.2-9.4%)	54 (24-100)	3600 (1900-6300)
Uzbekistan	1990	Female	40 to 44	65% (52-78%)	35% (22-48%)	1.3 (0.58-2.4)	3.2% (1.5-6.4%)	5.4% (2.2-10%)	55 (23-110)	3000 (1400-5500)
Uzbekistan	1990	Female	45 to 49	65% (50-78%)	35% (22-50%)	1.4 (0.6-2.7)	3.4% (1.2-7.2%)	5.6% (1.9-12%)	59 (19-130)	2700 (920-5600)
Uzbekistan	1990	Female	50 to 54	64% (49-77%)	36% (23-51%)	1.4 (0.61-2.7)	3.6% (0.84-8.3%)	5.5% (1.2-12%)	130 (28-290)	4900 (1200-11000)
Uzbekistan	1990	Female	55 to 59	62% (48-76%)	38% (24-52%)	1.4 (0.55-2.8)	3.3% (-0.011-8.5%)	4.7% (-0.075-12%)	130 (-2.1-350)	4400 (-15-12000)
Uzbekistan	1990	Female	60 to 64	60% (45-75%)	40% (25-55%)	1.4 (0.54-2.8)	2.1% (-1.4-8.2%)	2.9% (-1.9-11%)	130 (-87-480)	3500 (-2400-13000)
Uzbekistan	1990	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.2 (0.55-2.4)	1.1% (-2.3-6.7%)	1.5% (-3-8.8%)	70 (-140-390)	1600 (-3200-9000)
Uzbekistan	1990	Female	70 to 74	55% (43-68%)	45% (32-57%)	1.1 (0.42-2.3)	0.13% (-3.1-5.6%)	0.17% (-4.1-6.8%)	8 (-190-320)	150 (-3400-6000)
Uzbekistan	1990	Female	75 to 79	52% (39-65%)	48% (35-61%)	1 (0.37-2.1)	-0.6% (-3.9-4.7%)	-0.81% (-4.9-5.7%)	-50 (-300-360)	-660 (-4100-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	1990	Female	80 to 84	50% (36-64%)	50% (36-64%)	1.1 (0.42-2.1)	-1.2% (-4.5-3.9%)	-1.5% (-5.5-4.6%)	-110 (-380-310)	-1000 (-3900-3400)
Uzbekistan	1990	Female	85 to 89	49% (37-61%)	51% (39-63%)	1.1 (0.42-2.4)	-1.3% (-4.7-4.6%)	-1.7% (-5.8-5.3%)	-110 (-370-330)	-720 (-2600-2600)
Uzbekistan	1990	Female	90 to 94	50% (38-62%)	50% (38-62%)	1.2 (0.41-2.8)	-1.1% (-4.7-5.3%)	-1.5% (-5.8-6.2%)	-48 (-180-200)	-220 (-880-1000)
Uzbekistan	1990	Female	95 plus	51% (38-67%)	49% (33-62%)	1.3 (0.34-3.2)	-0.81% (-4.8-6.8%)	-1.2% (-6-7.9%)	-9.9 (-50-68)	-27 (-160-230)
Uzbekistan	1990	Male	All Ages	61% (52-70%)	39% (30-48%)	1 (0.76-1.4)	2.8% (1.7-4.2%)	2.7% (0.57-5%)	2000 (400-3700)	120000 (70000-170000)
Uzbekistan	1990	Male	15 to 19	55% (40-71%)	45% (29-60%)	0.56 (0.3-0.93)	2.8% (1.8-4.2%)	3.2% (1.4-5.3%)	34 (15-59)	4200 (2500-6400)
Uzbekistan	1990	Male	20 to 24	73% (57-85%)	27% (15-43%)	1.3 (0.75-1.9)	6.4% (4.4-8.8%)	7% (3.9-11%)	100 (55-160)	11000 (7000-15000)
Uzbekistan	1990	Male	25 to 29	80% (68-89%)	20% (11-32%)	2 (1.3-3)	8.7% (6.1-12%)	10% (6.2-15%)	190 (120-280)	16000 (11000-22000)
Uzbekistan	1990	Male	30 to 34	84% (73-92%)	16% (7.8-27%)	2.4 (1.6-3.5)	8.9% (6.3-12%)	11% (6.7-15%)	210 (140-310)	16000 (11000-22000)
Uzbekistan	1990	Male	35 to 39	86% (75-93%)	14% (7.3-25%)	2.4 (1.4-3.7)	8.1% (5.1-12%)	9.7% (5.4-15%)	190 (100-290)	12000 (7600-18000)
Uzbekistan	1990	Male	40 to 44	86% (76-93%)	14% (7-24%)	2.5 (1.5-3.9)	7.6% (4.3-12%)	9.1% (4.4-15%)	170 (79-280)	9300 (5200-14000)
Uzbekistan	1990	Male	45 to 49	85% (74-93%)	15% (7.2-26%)	2.7 (1.6-4.3)	7.9% (4.2-13%)	9.3% (4.4-15%)	190 (92-320)	9200 (5000-15000)
Uzbekistan	1990	Male	50 to 54	84% (73-92%)	16% (7.9-27%)	2.6 (1.5-4.1)	7.6% (3.6-13%)	8.8% (3.8-15%)	370 (160-650)	15000 (7100-25000)
Uzbekistan	1990	Male	55 to 59	83% (70-91%)	17% (8.8-30%)	2.3 (1.2-3.7)	6.2% (2.4-11%)	7.1% (2.4-13%)	330 (110-610)	11000 (4200-20000)
Uzbekistan	1990	Male	60 to 64	81% (67-91%)	19% (9.3-33%)	2 (1-3.4)	4.4% (0.68-9.4%)	5% (0.5-11%)	310 (31-700)	9000 (1400-20000)
Uzbekistan	1990	Male	65 to 69	79% (64-89%)	21% (11-36%)	1.7 (0.98-2.8)	2.9% (-0.49-7%)	3.2% (-0.87-8.2%)	140 (-37-350)	3300 (-570-8200)
Uzbekistan	1990	Male	70 to 74	76% (62-86%)	24% (14-38%)	1.5 (0.82-2.5)	1.3% (-1.8-5%)	1.3% (-2.3-5.7%)	45 (-80-200)	940 (-1300-3700)
Uzbekistan	1990	Male	75 to 79	73% (57-85%)	27% (15-43%)	1.3 (0.65-2.1)	-0.16% (-3-3.1%)	-0.39% (-3.7-3.4%)	-17 (-170-150)	-120 (-2200-2200)
Uzbekistan	1990	Male	80 to 84	70% (54-83%)	30% (17-46%)	1.1 (0.62-1.8)	-1.4% (-4.1-1.4%)	-1.8% (-5-1.4%)	-92 (-250-70)	-840 (-2400-820)
Uzbekistan	1990	Male	85 to 89	68% (52-81%)	32% (19-48%)	1 (0.56-1.7)	-2% (-4.7-0.36%)	-2.6% (-5.8-0.26%)	-110 (-250-11)	-730 (-1700-130)
Uzbekistan	1990	Male	90 to 94	67% (52-81%)	33% (19-48%)	0.97 (0.49-1.7)	-2.3% (-4.8-0.046%)	-3% (-6-0.17%)	-60 (-120-3.4)	-270 (-580-5.6)
Uzbekistan	1990	Male	95 plus	66% (46-82%)	34% (18-54%)	0.87 (0.37-1.6)	-2.3% (-4.6-0.023%)	-3% (-6-0.26%)	-15 (-29-1.3)	-43 (-92-0.43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	1995	Female	All Ages	51% (44-58%)	49% (42-56%)	0.75 (0.54-1)	1.4% (0.23-2.9%)	1.2% (-1.6-5.3%)	980 (-1300-4200)	53000 (9000-110000)
Uzbekistan	1995	Female	15 to 19	52% (40-67%)	48% (33-60%)	0.74 (0.31-1.4)	1.4% (0.66-2.6%)	2.3% (0.58-4.9%)	19 (4.9-40)	2200 (970-3900)
Uzbekistan	1995	Female	20 to 24	61% (48-74%)	39% (26-52%)	1.1 (0.45-2)	2.5% (1.2-4.6%)	3.3% (0.82-6.7%)	39 (9.7-80)	4200 (1900-7500)
Uzbekistan	1995	Female	25 to 29	64% (50-76%)	36% (24-50%)	1.3 (0.58-2.2)	3.1% (1.5-5.4%)	4.3% (1.4-8.4%)	56 (17-110)	5100 (2400-8600)
Uzbekistan	1995	Female	30 to 34	65% (50-79%)	35% (21-50%)	1.4 (0.7-2.4)	3.5% (2-5.6%)	5.3% (2.5-9.4%)	83 (38-150)	6300 (3400-10000)
Uzbekistan	1995	Female	35 to 39	66% (52-80%)	34% (20-48%)	1.4 (0.65-2.5)	3.7% (2-6.4%)	6% (2.9-11%)	100 (48-180)	6300 (3300-11000)
Uzbekistan	1995	Female	40 to 44	65% (52-79%)	35% (21-48%)	1.5 (0.66-2.6)	3.8% (1.8-7.1%)	6.3% (2.9-11%)	110 (49-190)	5600 (2700-9800)
Uzbekistan	1995	Female	45 to 49	65% (49-79%)	35% (21-51%)	1.5 (0.68-3.1)	4% (1.4-8.4%)	6.2% (2-13%)	110 (33-220)	4600 (1700-9600)
Uzbekistan	1995	Female	50 to 54	63% (49-77%)	37% (23-51%)	1.6 (0.7-3.1)	4.2% (0.89-9.7%)	6.1% (1.3-14%)	110 (23-260)	4300 (890-9900)
Uzbekistan	1995	Female	55 to 59	61% (48-75%)	39% (25-52%)	1.6 (0.66-3.1)	4% (0.027-10%)	5.4% (0.0017-14%)	230 (0.068-560)	7400 (48-18000)
Uzbekistan	1995	Female	60 to 64	60% (45-75%)	40% (25-55%)	1.5 (0.59-3.2)	2.7% (-1.6-9.9%)	3.4% (-2.1-12%)	180 (-110-620)	5000 (-3000-17000)
Uzbekistan	1995	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.4 (0.64-2.7)	1.5% (-2.4-7.7%)	1.9% (-3-9.5%)	140 (-230-710)	3200 (-5200-16000)
Uzbekistan	1995	Female	70 to 74	55% (42-69%)	45% (31-58%)	1.2 (0.49-2.4)	0.46% (-3.4-6.5%)	0.55% (-4.1-7.8%)	42 (-320-600)	780 (-5800-11000)
Uzbekistan	1995	Female	75 to 79	52% (40-66%)	48% (34-60%)	1.2 (0.42-2.3)	-0.36% (-4.2-5.6%)	-0.51% (-5-6.5%)	-35 (-350-460)	-430 (-4900-6500)
Uzbekistan	1995	Female	80 to 84	50% (37-65%)	50% (35-63%)	1.2 (0.47-2.3)	-0.74% (-4.5-5.1%)	-0.99% (-5.5-5.9%)	-78 (-450-470)	-720 (-4500-5100)
Uzbekistan	1995	Female	85 to 89	50% (37-63%)	50% (37-63%)	1.3 (0.49-2.6)	-0.81% (-4.8-5.9%)	-1.1% (-5.7-6.9%)	-77 (-400-500)	-490 (-2800-3700)
Uzbekistan	1995	Female	90 to 94	51% (39-62%)	49% (38-61%)	1.3 (0.46-3)	-0.63% (-4.8-7.2%)	-0.91% (-5.8-8.3%)	-37 (-240-340)	-150 (-1100-1700)
Uzbekistan	1995	Female	95 plus	51% (38-66%)	49% (34-62%)	1.4 (0.38-3.4)	-0.33% (-5.1-8.3%)	-0.58% (-6.1-9.5%)	-7.8 (-82-130)	-17 (-260-440)
Uzbekistan	1995	Male	All Ages	62% (54-71%)	38% (29-46%)	1.2 (0.9-1.5)	3.6% (2.3-5.1%)	3.7% (1.2-6.2%)	3100 (1000-5300)	160000 (100000-230000)
Uzbekistan	1995	Male	15 to 19	56% (40-72%)	44% (28-60%)	0.63 (0.34-1)	2.9% (1.8-4.2%)	3.2% (1.5-5.4%)	45 (21-76)	5200 (3100-7600)
Uzbekistan	1995	Male	20 to 24	73% (58-85%)	27% (15-42%)	1.4 (0.87-2.1)	6.9% (4.8-9.4%)	7.6% (4.5-11%)	140 (80-220)	14000 (9100-19000)
Uzbekistan	1995	Male	25 to 29	80% (68-89%)	20% (11-32%)	2.3 (1.4-3.4)	9.8% (6.9-13%)	11% (7.2-16%)	240 (150-350)	19000 (13000-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	1995	Male	30 to 34	84% (73-92%)	16% (7.9-27%)	2.7 (1.8-3.9)	10% (7.4-14%)	12% (8.1-17%)	320 (210-450)	23000 (16000-30000)
Uzbekistan	1995	Male	35 to 39	86% (75-93%)	14% (7.2-25%)	2.7 (1.7-3.9)	9.3% (6.2-13%)	11% (7-16%)	340 (200-490)	21000 (14000-29000)
Uzbekistan	1995	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	2.8 (1.7-4.1)	8.9% (5.3-13%)	11% (5.8-16%)	340 (180-520)	18000 (10000-26000)
Uzbekistan	1995	Male	45 to 49	86% (74-93%)	14% (7.3-26%)	3 (1.8-4.7)	9.1% (5-14%)	11% (5.3-17%)	340 (170-560)	15000 (8400-25000)
Uzbekistan	1995	Male	50 to 54	85% (73-92%)	15% (8.1-27%)	2.9 (1.7-4.6)	8.7% (4.3-14%)	10% (4.4-17%)	340 (150-600)	14000 (6600-23000)
Uzbekistan	1995	Male	55 to 59	83% (71-91%)	17% (8.7-29%)	2.5 (1.4-4)	6.9% (2.6-13%)	7.8% (2.7-15%)	520 (180-980)	17000 (6600-32000)
Uzbekistan	1995	Male	60 to 64	81% (67-90%)	19% (9.6-33%)	2.2 (1.2-3.7)	4.9% (0.66-11%)	5.4% (0.48-12%)	400 (35-900)	11000 (1500-25000)
Uzbekistan	1995	Male	65 to 69	79% (65-89%)	21% (11-35%)	2 (1.1-3)	3.1% (-0.55-7.7%)	3.4% (-0.89-8.7%)	300 (-79-760)	7200 (-1300-18000)
Uzbekistan	1995	Male	70 to 74	76% (62-86%)	24% (14-38%)	1.7 (0.96-2.7)	1.5% (-1.9-5.3%)	1.5% (-2.5-6%)	88 (-140-340)	1800 (-2300-6400)
Uzbekistan	1995	Male	75 to 79	73% (58-85%)	27% (15-42%)	1.4 (0.77-2.4)	-0.018% (-3.2-3.6%)	-0.23% (-4-3.9%)	-9 (-160-150)	-11 (-2100-2300)
Uzbekistan	1995	Male	80 to 84	71% (54-83%)	29% (17-46%)	1.3 (0.72-2.1)	-1.2% (-4.1-1.9%)	-1.6% (-5.1-1.9%)	-75 (-240-86)	-660 (-2300-1000)
Uzbekistan	1995	Male	85 to 89	68% (53-80%)	32% (20-47%)	1.2 (0.66-1.9)	-1.9% (-4.8-0.95%)	-2.4% (-5.7-0.9%)	-96 (-230-36)	-620 (-1600-310)
Uzbekistan	1995	Male	90 to 94	67% (53-80%)	33% (20-47%)	1.1 (0.58-1.9)	-2.2% (-4.9-0.52%)	-2.8% (-6-0.42%)	-63 (-130-9.2)	-280 (-630-68)
Uzbekistan	1995	Male	95 plus	66% (47-82%)	34% (18-53%)	1 (0.44-1.9)	-2.2% (-4.7-0.48%)	-2.9% (-6-0.16%)	-20 (-41-1.1)	-59 (-130-13)
Uzbekistan	2000	Female	All Ages	52% (45-59%)	48% (41-55%)	0.85 (0.62-1.1)	1.8% (0.46-3.6%)	1.8% (-1.5-6.4%)	1600 (-1300-5400)	72000 (18000-140000)
Uzbekistan	2000	Female	15 to 19	52% (40-67%)	48% (33-60%)	0.8 (0.33-1.5)	1.6% (0.67-3%)	2.7% (0.67-5.9%)	26 (6.4-56)	2800 (1200-5200)
Uzbekistan	2000	Female	20 to 24	61% (48-73%)	39% (27-52%)	1.2 (0.49-2.2)	3% (1.3-5.3%)	4.2% (1-8.3%)	56 (14-110)	5500 (2400-9600)
Uzbekistan	2000	Female	25 to 29	64% (50-76%)	36% (24-50%)	1.4 (0.62-2.4)	3.7% (1.7-6.3%)	5.4% (1.6-10%)	80 (24-150)	6700 (3100-12000)
Uzbekistan	2000	Female	30 to 34	65% (50-78%)	35% (22-50%)	1.5 (0.73-2.5)	4% (2.3-6.6%)	6.3% (2.9-11%)	100 (47-170)	7300 (4000-12000)
Uzbekistan	2000	Female	35 to 39	65% (52-79%)	35% (21-48%)	1.5 (0.7-2.7)	4.3% (2.3-7.1%)	7% (3.4-12%)	140 (67-230)	8500 (4400-14000)
Uzbekistan	2000	Female	40 to 44	65% (51-79%)	35% (21-49%)	1.6 (0.73-2.8)	4.4% (2.1-7.6%)	7.2% (3.3-13%)	170 (77-290)	8600 (4100-15000)
Uzbekistan	2000	Female	45 to 49	65% (49-79%)	35% (21-51%)	1.6 (0.73-3.2)	4.4% (1.6-9%)	6.9% (2.4-14%)	170 (60-350)	7600 (2700-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	2000	Female	50 to 54	63% (48-77%)	37% (23-52%)	1.7 (0.76-3.3)	4.6% (1-10%)	6.5% (1.4-15%)	170 (37-380)	6500 (1500-15000)
Uzbekistan	2000	Female	55 to 59	62% (47-76%)	38% (24-53%)	1.7 (0.68-3.3)	4.4% (0.14-11%)	5.9% (0.14-14%)	170 (3.9-420)	5600 (180-13000)
Uzbekistan	2000	Female	60 to 64	60% (45-75%)	40% (25-55%)	1.6 (0.63-3.4)	3.1% (-1.5-11%)	4% (-1.9-13%)	250 (-120-850)	6900 (-3300-24000)
Uzbekistan	2000	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.5 (0.67-2.9)	1.9% (-2.6-8.9%)	2.3% (-3.2-11%)	180 (-240-830)	4000 (-5500-19000)
Uzbekistan	2000	Female	70 to 74	55% (43-68%)	45% (32-57%)	1.3 (0.51-2.8)	0.81% (-3.4-7.6%)	0.95% (-4.2-9.1%)	99 (-430-940)	1900 (-7900-17000)
Uzbekistan	2000	Female	75 to 79	52% (40-65%)	48% (35-60%)	1.3 (0.45-2.5)	0.014% (-3.9-6.5%)	-0.065% (-4.8-7.4%)	-6.2 (-460-700)	24 (-6300-10000)
Uzbekistan	2000	Female	80 to 84	50% (37-65%)	50% (35-63%)	1.3 (0.51-2.4)	-0.13% (-4.2-6.4%)	-0.27% (-5.2-7.4%)	-24 (-480-680)	-150 (-4800-7200)
Uzbekistan	2000	Female	85 to 89	49% (37-61%)	51% (39-63%)	1.4 (0.52-2.8)	-0.16% (-4.5-7.4%)	-0.32% (-5.5-8.6%)	-22 (-370-570)	-92 (-2600-4200)
Uzbekistan	2000	Female	90 to 94	50% (39-62%)	50% (38-61%)	1.5 (0.5-3.4)	0.0077% (-4.6-8.5%)	-0.15% (-5.6-10%)	-5.9 (-220-380)	2 (-1100-1900)
Uzbekistan	2000	Female	95 plus	51% (37-66%)	49% (34-63%)	1.5 (0.4-3.7)	0.28% (-4.9-9.2%)	0.14% (-5.9-11%)	2.1 (-91-160)	17 (-290-550)
Uzbekistan	2000	Male	All Ages	63% (54-72%)	37% (28-46%)	1.3 (1-1.7)	4.6% (3-6.4%)	4.5% (1.7-7.5%)	4200 (1500-7000)	210000 (130000-300000)
Uzbekistan	2000	Male	15 to 19	55% (39-71%)	45% (29-61%)	0.67 (0.36-1.1)	3% (1.8-4.4%)	3.3% (1.5-5.6%)	57 (27-98)	6400 (3800-9500)
Uzbekistan	2000	Male	20 to 24	73% (57-85%)	27% (15-43%)	1.5 (0.91-2.2)	7.3% (4.9-10%)	8.1% (4.6-12%)	190 (100-280)	17000 (11000-24000)
Uzbekistan	2000	Male	25 to 29	80% (69-89%)	20% (11-31%)	2.4 (1.5-3.6)	11% (7.1-15%)	12% (7.6-18%)	340 (200-490)	26000 (17000-35000)
Uzbekistan	2000	Male	30 to 34	84% (72-92%)	16% (7.8-28%)	2.9 (1.9-4.2)	11% (8.1-15%)	14% (9.2-19%)	400 (260-560)	27000 (19000-36000)
Uzbekistan	2000	Male	35 to 39	86% (75-93%)	14% (7.3-25%)	2.9 (1.8-4.3)	11% (7.1-15%)	13% (8-18%)	470 (280-680)	28000 (18000-39000)
Uzbekistan	2000	Male	40 to 44	86% (76-93%)	14% (7.1-24%)	3 (1.9-4.4)	10% (6.2-15%)	12% (6.9-18%)	520 (290-790)	27000 (16000-39000)
Uzbekistan	2000	Male	45 to 49	85% (73-93%)	15% (7.2-27%)	3.2 (2-5)	10% (5.8-16%)	12% (6.4-19%)	550 (290-900)	25000 (14000-40000)
Uzbekistan	2000	Male	50 to 54	84% (73-92%)	16% (8.1-27%)	3.2 (1.8-4.9)	9.5% (4.8-16%)	11% (5-18%)	490 (220-830)	19000 (9300-32000)
Uzbekistan	2000	Male	55 to 59	83% (71-92%)	17% (8.2-29%)	2.7 (1.6-4.4)	7.4% (2.8-14%)	8.3% (2.8-16%)	390 (130-740)	13000 (5000-25000)
Uzbekistan	2000	Male	60 to 64	81% (67-91%)	19% (9.4-33%)	2.4 (1.3-3.9)	5.2% (0.64-12%)	5.7% (0.39-13%)	520 (35-1200)	15000 (1800-33000)
Uzbekistan	2000	Male	65 to 69	79% (65-89%)	21% (11-35%)	2.1 (1.2-3.3)	3.2% (-0.67-8.3%)	3.5% (-1.1-9.2%)	320 (-100-850)	7700 (-1700-20000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	2000	Male	70 to 74	76% (61-86%)	24% (14-39%)	1.8 (1-2.9)	1.5% (-2.3-5.5%)	1.5% (-3-6.1%)	160 (-320-650)	3200 (-5200-12000)
Uzbekistan	2000	Male	75 to 79	73% (57-85%)	27% (15-43%)	1.6 (0.85-2.5)	0.13% (-3.4-4.1%)	-0.071% (-4.2-4.5%)	-4.3 (-260-260)	130 (-3400-3900)
Uzbekistan	2000	Male	80 to 84	71% (54-83%)	29% (17-46%)	1.4 (0.78-2.2)	-0.9% (-4-2.4%)	-1.3% (-5-2.5%)	-59 (-230-110)	-490 (-2200-1300)
Uzbekistan	2000	Male	85 to 89	68% (53-80%)	32% (20-47%)	1.3 (0.7-2.1)	-1.6% (-4.6-1.4%)	-2.1% (-5.7-1.4%)	-61 (-160-41)	-390 (-1100-350)
Uzbekistan	2000	Male	90 to 94	67% (52-80%)	33% (20-48%)	1.2 (0.63-2)	-1.9% (-4.8-0.92%)	-2.6% (-5.9-0.79%)	-43 (-98-13)	-190 (-460-88)
Uzbekistan	2000	Male	95 plus	66% (47-82%)	34% (18-53%)	1.1 (0.47-2)	-2% (-4.8-0.81%)	-2.7% (-6-0.61%)	-17 (-37-3.7)	-48 (-110-20)
Uzbekistan	2005	Female	All Ages	52% (46-60%)	48% (40-54%)	0.97 (0.7-1.3)	2.2% (0.65-4.3%)	2.2% (-1.4-7.3%)	2100 (-1200-6700)	89000 (26000-170000)
Uzbekistan	2005	Female	15 to 19	52% (38-67%)	48% (33-62%)	0.83 (0.34-1.6)	1.7% (0.76-3%)	3% (0.78-6.3%)	28 (7.1-57)	3100 (1400-5600)
Uzbekistan	2005	Female	20 to 24	61% (47-73%)	39% (27-53%)	1.2 (0.49-2.3)	3.1% (1.5-5.4%)	4.7% (1.2-9.2%)	64 (16-130)	6400 (2900-11000)
Uzbekistan	2005	Female	25 to 29	63% (49-76%)	37% (24-51%)	1.4 (0.64-2.7)	3.8% (1.8-6.6%)	5.7% (1.9-11%)	87 (28-170)	7400 (3400-12000)
Uzbekistan	2005	Female	30 to 34	65% (50-79%)	35% (21-50%)	1.6 (0.79-2.8)	4.3% (2.3-7.1%)	6.8% (3.1-12%)	120 (52-200)	8300 (4400-14000)
Uzbekistan	2005	Female	35 to 39	65% (52-80%)	35% (20-48%)	1.6 (0.74-2.9)	4.6% (2.4-7.7%)	7.5% (3.7-13%)	150 (71-260)	9000 (4700-15000)
Uzbekistan	2005	Female	40 to 44	65% (51-79%)	35% (21-49%)	1.7 (0.77-3)	4.7% (2.3-8.1%)	7.7% (3.7-13%)	210 (96-360)	11000 (5200-18000)
Uzbekistan	2005	Female	45 to 49	65% (49-78%)	35% (22-51%)	1.7 (0.78-3.4)	4.7% (1.8-9.5%)	7.4% (2.7-15%)	250 (86-500)	11000 (3900-22000)
Uzbekistan	2005	Female	50 to 54	63% (48-77%)	37% (23-52%)	1.8 (0.83-3.4)	4.9% (1.2-11%)	7% (1.7-15%)	280 (64-600)	11000 (2400-23000)
Uzbekistan	2005	Female	55 to 59	61% (47-75%)	39% (25-53%)	1.8 (0.74-3.6)	4.7% (0.27-11%)	6.3% (0.41-15%)	260 (16-620)	8300 (470-20000)
Uzbekistan	2005	Female	60 to 64	59% (45-74%)	41% (26-55%)	1.7 (0.67-3.6)	3.5% (-1.4-12%)	4.3% (-1.9-14%)	190 (-85-640)	5300 (-2300-18000)
Uzbekistan	2005	Female	65 to 69	58% (45-73%)	42% (27-55%)	1.6 (0.71-3.1)	2.3% (-2.4-9.8%)	2.8% (-3-12%)	250 (-270-1100)	5800 (-6000-24000)
Uzbekistan	2005	Female	70 to 74	55% (42-68%)	45% (32-58%)	1.4 (0.54-2.9)	1.1% (-3.5-8.3%)	1.3% (-4.3-10%)	130 (-450-1100)	2500 (-8100-19000)
Uzbekistan	2005	Female	75 to 79	52% (39-65%)	48% (35-61%)	1.3 (0.48-2.7)	0.25% (-4.1-7.3%)	0.21% (-5-8.5%)	26 (-610-1000)	520 (-8400-15000)
Uzbekistan	2005	Female	80 to 84	50% (36-64%)	50% (36-64%)	1.4 (0.53-2.6)	0.21% (-4.2-7.6%)	0.14% (-5.1-8.7%)	17 (-600-1100)	320 (-6100-11000)
Uzbekistan	2005	Female	85 to 89	49% (37-61%)	51% (39-63%)	1.4 (0.55-3)	0.26% (-4.4-8.5%)	0.16% (-5.3-9.8%)	12 (-380-680)	160 (-2700-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	2005	Female	90 to 94	50% (38-61%)	50% (39-62%)	1.6 (0.52-3.6)	0.53% (-4.5-9.7%)	0.46% (-5.5-11%)	16 (-190-380)	110 (-890-1900)
Uzbekistan	2005	Female	95 plus	51% (36-66%)	49% (34-64%)	1.6 (0.44-4)	0.88% (-4.9-11%)	0.84% (-5.8-12%)	11 (-81-170)	47 (-260-570)
Uzbekistan	2005	Male	All Ages	64% (56-73%)	36% (27-44%)	1.5 (1.2-1.9)	5% (3.1-7.1%)	4.8% (1.5-8.2%)	5000 (1600-8500)	250000 (150000-350000)
Uzbekistan	2005	Male	15 to 19	55% (39-71%)	45% (29-61%)	0.72 (0.38-1.2)	3.2% (2-4.8%)	3.7% (1.7-6.2%)	65 (30-110)	7400 (4400-11000)
Uzbekistan	2005	Male	20 to 24	73% (58-85%)	27% (15-42%)	1.6 (0.98-2.4)	7.6% (5.2-10%)	8.7% (5.1-13%)	200 (110-300)	19000 (12000-26000)
Uzbekistan	2005	Male	25 to 29	80% (68-89%)	20% (11-32%)	2.6 (1.6-3.8)	11% (7.5-15%)	13% (7.9-18%)	350 (220-510)	27000 (18000-38000)
Uzbekistan	2005	Male	30 to 34	84% (73-92%)	16% (8.1-27%)	3.1 (2-4.4)	12% (8.3-16%)	14% (9.4-20%)	440 (290-620)	30000 (21000-41000)
Uzbekistan	2005	Male	35 to 39	86% (76-93%)	14% (7.5-24%)	3.1 (1.9-4.5)	11% (7.4-16%)	14% (8.4-20%)	480 (290-710)	28000 (18000-40000)
Uzbekistan	2005	Male	40 to 44	86% (76-93%)	14% (7-24%)	3.2 (2-4.6)	11% (6.3-16%)	13% (7.2-19%)	610 (350-930)	32000 (19000-47000)
Uzbekistan	2005	Male	45 to 49	85% (73-92%)	15% (7.5-27%)	3.4 (2.1-5.2)	10% (5.8-16%)	12% (6.4-20%)	730 (380-1200)	33000 (18000-52000)
Uzbekistan	2005	Male	50 to 54	84% (73-92%)	16% (7.8-27%)	3.3 (2-5.1)	9.8% (4.7-16%)	11% (4.9-19%)	740 (320-1300)	29000 (14000-49000)
Uzbekistan	2005	Male	55 to 59	83% (70-91%)	17% (8.6-30%)	2.9 (1.7-4.6)	7.7% (2.7-14%)	8.6% (2.8-16%)	520 (160-990)	17000 (6400-33000)
Uzbekistan	2005	Male	60 to 64	81% (67-90%)	19% (9.6-33%)	2.5 (1.3-4.2)	5.4% (0.39-12%)	5.9% (0.097-14%)	400 (6-930)	11000 (810-26000)
Uzbekistan	2005	Male	65 to 69	79% (63-89%)	21% (11-37%)	2.2 (1.3-3.5)	3.5% (-0.75-9%)	3.8% (-1.2-10%)	440 (-130-1100)	10000 (-2300-26000)
Uzbekistan	2005	Male	70 to 74	76% (61-86%)	24% (14-39%)	1.9 (1.1-3.1)	1.5% (-2.6-6%)	1.5% (-3.2-6.6%)	170 (-360-720)	3400 (-5900-14000)
Uzbekistan	2005	Male	75 to 79	73% (57-85%)	27% (15-43%)	1.6 (0.87-2.7)	0.06% (-3.7-4.3%)	-0.16% (-4.5-4.6%)	-17 (-480-510)	100 (-6400-7500)
Uzbekistan	2005	Male	80 to 84	70% (54-83%)	30% (17-46%)	1.5 (0.81-2.3)	-0.81% (-4.3-2.8%)	-1.2% (-5.2-3%)	-83 (-370-210)	-680 (-3600-2400)
Uzbekistan	2005	Male	85 to 89	68% (53-80%)	32% (20-47%)	1.4 (0.75-2.2)	-1.5% (-4.6-1.7%)	-2% (-5.6-1.6%)	-53 (-150-42)	-330 (-1000-370)
Uzbekistan	2005	Male	90 to 94	67% (52-80%)	33% (20-48%)	1.3 (0.66-2.2)	-1.8% (-4.8-1.3%)	-2.4% (-6-1.3%)	-26 (-63-13)	-110 (-300-84)
Uzbekistan	2005	Male	95 plus	66% (47-81%)	34% (19-53%)	1.1 (0.5-2.2)	-1.9% (-4.9-1.3%)	-2.6% (-6.1-1.2%)	-11 (-25-4.7)	-30 (-77-21)
Uzbekistan	2010	Female	All Ages	53% (46-60%)	47% (40-54%)	1.2 (0.85-1.6)	2.8% (1.1-5.1%)	3.5% (-0.63-9.2%)	3100 (-580-8200)	110000 (42000-210000)
Uzbekistan	2010	Female	15 to 19	51% (37-66%)	49% (34-63%)	0.96 (0.38-1.8)	1.9% (0.82-3.5%)	3.7% (0.83-7.7%)	32 (7.2-68)	3500 (1400-6400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	2010	Female	20 to 24	60% (46-73%)	40% (27-54%)	1.4 (0.57-2.7)	3.4% (1.6-5.9%)	5.4% (1.3-11%)	71 (18-140)	7300 (3300-13000)
Uzbekistan	2010	Female	25 to 29	63% (49-76%)	37% (24-51%)	1.7 (0.75-3.1)	3.9% (1.9-6.8%)	6.2% (2.1-12%)	89 (30-170)	8100 (3900-14000)
Uzbekistan	2010	Female	30 to 34	65% (49-78%)	35% (22-51%)	1.8 (0.91-3.3)	4.5% (2.5-7.3%)	7.7% (3.7-13%)	120 (57-210)	8800 (4900-14000)
Uzbekistan	2010	Female	35 to 39	65% (52-78%)	35% (22-48%)	1.9 (0.87-3.5)	4.9% (2.6-8.3%)	8.5% (4.2-15%)	160 (77-280)	9900 (5200-16000)
Uzbekistan	2010	Female	40 to 44	65% (51-78%)	35% (22-49%)	1.9 (0.9-3.6)	5% (2.3-8.8%)	8.4% (3.8-15%)	200 (93-360)	11000 (5000-18000)
Uzbekistan	2010	Female	45 to 49	64% (48-78%)	36% (22-52%)	2 (0.91-3.9)	5.1% (1.8-10%)	8.2% (2.9-16%)	300 (110-590)	13000 (4800-26000)
Uzbekistan	2010	Female	50 to 54	63% (47-77%)	37% (23-53%)	2.1 (0.93-4)	5.5% (1.4-11%)	8% (2-17%)	390 (100-790)	15000 (3800-30000)
Uzbekistan	2010	Female	55 to 59	61% (47-75%)	39% (25-53%)	2.1 (0.86-4.1)	5.5% (0.44-13%)	7.5% (0.61-17%)	400 (31-910)	13000 (1000-30000)
Uzbekistan	2010	Female	60 to 64	59% (45-74%)	41% (26-55%)	2 (0.79-4.3)	4.4% (-1.1-13%)	5.7% (-1.5-17%)	300 (-74-860)	8300 (-2000-24000)
Uzbekistan	2010	Female	65 to 69	57% (43-73%)	43% (27-57%)	1.9 (0.84-3.7)	3.5% (-1.8-12%)	4.4% (-2.4-15%)	220 (-130-760)	5200 (-2800-18000)
Uzbekistan	2010	Female	70 to 74	54% (41-68%)	46% (32-59%)	1.7 (0.62-3.4)	2.2% (-3.1-10%)	2.7% (-3.9-13%)	280 (-410-1300)	5300 (-7300-24000)
Uzbekistan	2010	Female	75 to 79	51% (39-64%)	49% (36-61%)	1.5 (0.55-3.2)	1.4% (-3.5-9.2%)	1.6% (-4.4-11%)	160 (-450-1100)	2500 (-6300-16000)
Uzbekistan	2010	Female	80 to 84	49% (35-64%)	51% (36-65%)	1.6 (0.6-3)	1.3% (-3.7-9.5%)	1.4% (-4.6-11%)	160 (-540-1300)	1900 (-5500-14000)
Uzbekistan	2010	Female	85 to 89	48% (35-61%)	52% (39-65%)	1.6 (0.62-3.5)	1.3% (-4.1-11%)	1.4% (-5-12%)	130 (-470-1100)	1100 (-3200-8600)
Uzbekistan	2010	Female	90 to 94	49% (36-62%)	51% (38-64%)	1.8 (0.59-4.2)	1.7% (-4-11%)	1.8% (-4.8-13%)	69 (-190-510)	380 (-900-2600)
Uzbekistan	2010	Female	95 plus	50% (36-65%)	50% (35-64%)	1.9 (0.49-4.7)	2% (-4.3-12%)	2.2% (-5.3-14%)	27 (-65-170)	98 (-210-580)
Uzbekistan	2010	Male	All Ages	65% (57-73%)	35% (27-43%)	1.9 (1.4-2.4)	6.2% (3.8-8.6%)	6.1% (2.5-10%)	6300 (2500-10000)	300000 (190000-420000)
Uzbekistan	2010	Male	15 to 19	53% (37-70%)	47% (30-63%)	0.8 (0.41-1.3)	3.5% (2-5.3%)	4.1% (1.8-6.8%)	82 (37-140)	8600 (4900-13000)
Uzbekistan	2010	Male	20 to 24	72% (55-85%)	28% (15-45%)	1.8 (1.1-2.8)	8% (5.5-11%)	9% (5.3-13%)	290 (170-440)	26000 (17000-36000)
Uzbekistan	2010	Male	25 to 29	79% (67-88%)	21% (12-33%)	3 (1.8-4.4)	11% (7.7-15%)	13% (8.3-19%)	430 (270-630)	33000 (22000-46000)
Uzbekistan	2010	Male	30 to 34	84% (71-92%)	16% (8.1-29%)	3.6 (2.3-5.2)	12% (8.9-17%)	15% (9.9-21%)	490 (320-690)	33000 (23000-45000)
Uzbekistan	2010	Male	35 to 39	85% (74-93%)	15% (7.4-26%)	3.6 (2.2-5.2)	12% (8.2-17%)	15% (9.5-22%)	550 (340-800)	33000 (21000-46000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	2010	Male	40 to 44	85% (75-93%)	15% (7.4-25%)	3.7 (2.3-5.3)	12% (7.2-17%)	15% (8.4-21%)	620 (350-930)	32000 (19000-47000)
Uzbekistan	2010	Male	45 to 49	85% (72-92%)	15% (7.5-28%)	3.9 (2.5-6)	12% (6.7-18%)	14% (7.6-22%)	870 (450-1400)	40000 (22000-61000)
Uzbekistan	2010	Male	50 to 54	84% (72-92%)	16% (8.5-28%)	3.8 (2.3-6)	11% (5.4-18%)	13% (5.9-21%)	1000 (450-1700)	39000 (19000-64000)
Uzbekistan	2010	Male	55 to 59	82% (70-91%)	18% (8.9-30%)	3.3 (1.9-5.3)	9% (3.4-16%)	10% (3.7-19%)	790 (270-1500)	27000 (9800-48000)
Uzbekistan	2010	Male	60 to 64	80% (67-90%)	20% (9.8-33%)	2.9 (1.6-4.8)	6.8% (1-14%)	7.6% (0.87-17%)	530 (62-1200)	15000 (2400-33000)
Uzbekistan	2010	Male	65 to 69	78% (63-89%)	22% (11-37%)	2.6 (1.5-4)	5.1% (0.12-11%)	5.6% (-0.15-13%)	370 (-10-870)	8900 (200-20000)
Uzbekistan	2010	Male	70 to 74	75% (60-86%)	25% (14-40%)	2.2 (1.2-3.6)	2.8% (-1.7-8.8%)	3% (-2.3-10%)	350 (-260-1100)	6800 (-4100-21000)
Uzbekistan	2010	Male	75 to 79	72% (56-84%)	28% (16-44%)	1.9 (1-3.1)	1% (-3.3-6.4%)	0.91% (-4.1-6.9%)	87 (-390-670)	1600 (-5000-9800)
Uzbekistan	2010	Male	80 to 84	70% (52-82%)	30% (18-48%)	1.7 (0.92-2.7)	-0.082% (-3.9-4.1%)	-0.38% (-4.8-4.4%)	-31 (-400-370)	-78 (-3800-4100)
Uzbekistan	2010	Male	85 to 89	67% (51-80%)	33% (20-49%)	1.6 (0.84-2.5)	-0.95% (-4.4-2.7%)	-1.4% (-5.4-2.8%)	-66 (-260-130)	-380 (-1700-1100)
Uzbekistan	2010	Male	90 to 94	66% (51-80%)	34% (20-49%)	1.4 (0.75-2.5)	-1.3% (-4.7-2.3%)	-1.8% (-5.8-2.2%)	-22 (-69-26)	-88 (-320-150)
Uzbekistan	2010	Male	95 plus	65% (46-81%)	35% (19-54%)	1.3 (0.57-2.4)	-1.4% (-4.7-2.5%)	-2% (-6-2.4%)	-6.2 (-19-7.4)	-16 (-57-30)
Uzbekistan	2016	Female	All Ages	53% (46-60%)	47% (40-54%)	1.3 (0.91-1.9)	3.5% (1.4-6.2%)	4.6% (-0.0042-11%)	4300 (-3.9-11000)	140000 (54000-260000)
Uzbekistan	2016	Female	15 to 19	49% (36-65%)	51% (35-64%)	1 (0.41-2)	2% (0.84-4%)	4.1% (0.91-8.9%)	29 (6.3-64)	3100 (1200-5900)
Uzbekistan	2016	Female	20 to 24	59% (46-73%)	41% (27-54%)	1.5 (0.6-3)	3.5% (1.6-6.2%)	5.8% (1.5-12%)	63 (16-130)	6700 (3100-13000)
Uzbekistan	2016	Female	25 to 29	63% (48-76%)	37% (24-52%)	1.8 (0.77-3.5)	4% (2.1-6.9%)	6.6% (2.3-13%)	98 (33-200)	9200 (4600-16000)
Uzbekistan	2016	Female	30 to 34	64% (50-78%)	36% (22-50%)	2 (0.95-3.7)	4.6% (2.7-7.6%)	8.2% (4.1-14%)	140 (66-250)	11000 (6000-18000)
Uzbekistan	2016	Female	35 to 39	65% (51-78%)	35% (22-49%)	2 (0.9-3.8)	5.1% (2.7-8.4%)	9.1% (4.6-16%)	180 (86-310)	11000 (5800-19000)
Uzbekistan	2016	Female	40 to 44	65% (50-78%)	35% (22-50%)	2.1 (0.95-3.9)	5.3% (2.5-9.3%)	9.2% (4.1-16%)	230 (100-430)	12000 (5600-22000)
Uzbekistan	2016	Female	45 to 49	64% (48-78%)	36% (22-52%)	2.2 (0.98-4.4)	5.6% (2.1-11%)	9% (3.5-18%)	300 (110-610)	14000 (5300-27000)
Uzbekistan	2016	Female	50 to 54	62% (47-76%)	38% (24-53%)	2.3 (1-4.3)	6% (1.8-12%)	9% (2.8-18%)	460 (140-970)	18000 (5100-37000)
Uzbekistan	2016	Female	55 to 59	60% (46-74%)	40% (26-54%)	2.3 (0.93-4.4)	6.1% (0.98-13%)	8.6% (1.4-18%)	580 (93-1300)	19000 (2900-43000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	2016	Female	60 to 64	58% (44-73%)	42% (27-56%)	2.2 (0.86-4.6)	5.1% (-0.55-13%)	6.9% (-0.84-18%)	510 (-64-1400)	14000 (-1700-39000)
Uzbekistan	2016	Female	65 to 69	57% (43-72%)	43% (28-57%)	2 (0.9-4)	4.4% (-1.2-13%)	5.7% (-1.6-17%)	400 (-110-1100)	9200 (-2500-27000)
Uzbekistan	2016	Female	70 to 74	54% (41-67%)	46% (33-59%)	1.8 (0.67-3.9)	3.2% (-2.4-12%)	4% (-3.1-15%)	250 (-190-960)	4700 (-3400-18000)
Uzbekistan	2016	Female	75 to 79	50% (37-64%)	50% (36-63%)	1.7 (0.59-3.6)	2.3% (-2.9-11%)	2.8% (-3.8-14%)	300 (-410-1500)	4400 (-5600-22000)
Uzbekistan	2016	Female	80 to 84	48% (34-62%)	52% (38-66%)	1.7 (0.63-3.5)	2.1% (-3-11%)	2.4% (-4-13%)	310 (-500-1800)	3400 (-5100-18000)
Uzbekistan	2016	Female	85 to 89	47% (34-60%)	53% (40-66%)	1.8 (0.68-3.8)	2.1% (-3.6-12%)	2.3% (-4.3-14%)	220 (-440-1400)	1800 (-3100-10000)
Uzbekistan	2016	Female	90 to 94	48% (36-61%)	52% (39-64%)	1.9 (0.67-4.4)	2.3% (-3.7-12%)	2.6% (-4.6-14%)	150 (-260-820)	780 (-1300-4200)
Uzbekistan	2016	Female	95 plus	50% (35-65%)	50% (35-65%)	2.1 (0.56-5)	2.6% (-4-13%)	2.9% (-5-15%)	46 (-80-240)	170 (-250-830)
Uzbekistan	2016	Male	All Ages	65% (57-74%)	35% (26-43%)	2.2 (1.5-2.9)	7.2% (4.5-11%)	7.2% (3.2-12%)	8000 (3500-14000)	370000 (230000-540000)
Uzbekistan	2016	Male	15 to 19	52% (35-69%)	48% (31-65%)	0.86 (0.43-1.5)	3.7% (2.1-5.9%)	4.4% (2-7.6%)	77 (32-140)	7900 (4500-13000)
Uzbekistan	2016	Male	20 to 24	71% (53-84%)	29% (16-47%)	2 (1.1-3.2)	8.6% (5.6-12%)	9.7% (5.4-15%)	300 (160-470)	27000 (17000-39000)
Uzbekistan	2016	Male	25 to 29	79% (66-88%)	21% (12-34%)	3.3 (2-5.1)	12% (8.2-17%)	14% (8.8-21%)	530 (320-810)	41000 (27000-58000)
Uzbekistan	2016	Male	30 to 34	84% (71-92%)	16% (8.4-29%)	3.9 (2.4-5.9)	13% (9.1-18%)	16% (11-23%)	600 (380-900)	41000 (28000-59000)
Uzbekistan	2016	Male	35 to 39	85% (74-92%)	15% (7.7-26%)	3.9 (2.3-5.8)	13% (8.4-18%)	16% (10-24%)	620 (360-940)	37000 (24000-54000)
Uzbekistan	2016	Male	40 to 44	85% (75-93%)	15% (7.4-25%)	4 (2.4-6)	12% (7.7-18%)	16% (9-23%)	710 (400-1100)	37000 (22000-56000)
Uzbekistan	2016	Male	45 to 49	84% (71-92%)	16% (7.9-29%)	4.3 (2.6-6.7)	13% (7.1-19%)	16% (8.4-24%)	870 (460-1400)	39000 (22000-62000)
Uzbekistan	2016	Male	50 to 54	83% (71-91%)	17% (8.6-29%)	4.2 (2.4-6.8)	12% (5.7-19%)	15% (6.6-24%)	1200 (520-2000)	46000 (22000-76000)
Uzbekistan	2016	Male	55 to 59	82% (69-91%)	18% (9-31%)	3.6 (2-5.8)	10% (3.9-18%)	12% (4.4-21%)	1200 (430-2200)	40000 (16000-73000)
Uzbekistan	2016	Male	60 to 64	80% (65-90%)	20% (9.9-35%)	3.2 (1.7-5.3)	7.9% (2-16%)	9% (1.9-18%)	940 (200-1900)	27000 (6600-54000)
Uzbekistan	2016	Male	65 to 69	77% (61-89%)	23% (11-39%)	2.8 (1.6-4.5)	6.1% (0.88-13%)	6.9% (0.7-15%)	590 (60-1300)	14000 (2100-30000)
Uzbekistan	2016	Male	70 to 74	74% (59-86%)	26% (14-41%)	2.4 (1.3-4.1)	3.9% (-1.2-11%)	4.2% (-1.7-13%)	310 (-120-930)	6100 (-1900-17000)
Uzbekistan	2016	Male	75 to 79	71% (55-83%)	29% (17-45%)	2 (1.1-3.6)	1.9% (-2.5-8%)	1.9% (-3.3-9.2%)	200 (-340-950)	3300 (-4300-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uzbekistan	2016	Male	80 to 84	68% (51-82%)	32% (18-49%)	1.8 (0.94-3)	0.51% (-3.4-5.4%)	0.28% (-4.3-6%)	27 (-430-610)	600 (-4000-6600)
Uzbekistan	2016	Male	85 to 89	66% (49-79%)	34% (21-51%)	1.7 (0.84-2.8)	-0.56% (-4.3-3.6%)	-1% (-5.4-3.8%)	-58 (-310-230)	-280 (-2100-1800)
Uzbekistan	2016	Male	90 to 94	65% (50-79%)	35% (21-50%)	1.5 (0.77-2.7)	-0.96% (-4.6-2.7%)	-1.5% (-5.7-2.8%)	-38 (-150-72)	-140 (-680-400)
Uzbekistan	2016	Male	95 plus	65% (44-81%)	35% (19-56%)	1.4 (0.57-2.6)	-1.1% (-4.7-3.2%)	-1.7% (-5.9-3.2%)	-7.1 (-25-14)	-18 (-76-51)
Albania	1990	Female	All Ages	25% (22-29%)	75% (71-78%)	0.51 (0.42-0.61)	1.3% (0.87-1.7%)	1.8% (0.75-3.1%)	130 (54-230)	5000 (3400-6900)
Albania	1990	Female	15 to 19	25% (18-34%)	75% (66-82%)	0.52 (0.33-0.76)	1.5% (0.98-2%)	1.7% (0.87-3%)	1 (0.52-1.8)	230 (150-320)
Albania	1990	Female	20 to 24	34% (24-44%)	66% (56-76%)	0.79 (0.52-1.1)	2.7% (1.9-3.7%)	2.4% (1.2-3.7%)	1.8 (0.93-2.8)	460 (300-660)
Albania	1990	Female	25 to 29	35% (25-46%)	65% (54-75%)	0.73 (0.49-1)	2.6% (2-3.5%)	2.1% (1.1-3.3%)	1.8 (0.95-2.8)	470 (330-670)
Albania	1990	Female	30 to 34	36% (25-48%)	64% (52-75%)	0.72 (0.45-1.1)	2.4% (1.7-3.2%)	2.2% (1.3-3.4%)	2.1 (1.2-3.2)	430 (290-600)
Albania	1990	Female	35 to 39	37% (26-48%)	63% (52-74%)	0.77 (0.49-1.1)	2.2% (1.6-2.9%)	2.6% (1.5-3.9%)	2.3 (1.3-3.5)	350 (240-470)
Albania	1990	Female	40 to 44	37% (27-47%)	63% (53-73%)	0.78 (0.51-1.1)	2.1% (1.4-2.7%)	2.8% (1.7-4.3%)	2.8 (1.7-4.2)	290 (200-390)
Albania	1990	Female	45 to 49	34% (23-44%)	66% (56-77%)	0.75 (0.44-1.1)	1.9% (1.3-2.6%)	2.8% (1.6-4.4%)	3.4 (1.9-5.4)	270 (180-370)
Albania	1990	Female	50 to 54	32% (23-43%)	68% (57-77%)	0.86 (0.54-1.3)	2.1% (1.3-3.1%)	3.3% (1.8-5.3%)	6 (3.1-9.4)	320 (200-470)
Albania	1990	Female	55 to 59	31% (20-46%)	69% (54-80%)	1.1 (0.62-1.8)	2.7% (1.5-4.4%)	4.5% (2.1-7.5%)	10 (4.7-17)	400 (210-640)
Albania	1990	Female	60 to 64	29% (19-44%)	71% (56-81%)	1.1 (0.64-1.8)	3% (1.4-4.9%)	4.6% (2.1-7.9%)	16 (7-26)	480 (230-800)
Albania	1990	Female	65 to 69	27% (18-39%)	73% (61-82%)	0.97 (0.59-1.5)	2.6% (1.1-4.4%)	3.9% (1.7-6.6%)	18 (7.8-31)	450 (200-740)
Albania	1990	Female	70 to 74	24% (16-34%)	76% (66-84%)	0.78 (0.42-1.3)	2.1% (0.59-4.1%)	3% (0.79-5.8%)	19 (5-38)	370 (100-710)
Albania	1990	Female	75 to 79	20% (13-29%)	80% (71-87%)	0.61 (0.32-1.1)	1.6% (0.19-3.6%)	2.1% (0.2-4.7%)	19 (1.8-42)	270 (32-600)
Albania	1990	Female	80 to 84	17% (11-25%)	83% (75-89%)	0.52 (0.3-0.91)	1.3% (0.087-3.2%)	1.6% (0.079-3.9%)	15 (0.71-35)	150 (10-360)
Albania	1990	Female	85 to 89	17% (10-24%)	83% (76-90%)	0.5 (0.27-0.85)	1.1% (0.067-2.7%)	1.4% (0.032-3.4%)	8.8 (0.2-22)	67 (4-160)
Albania	1990	Female	90 to 94	18% (11-26%)	82% (74-89%)	0.53 (0.27-0.91)	0.9% (-0.17-2.5%)	1.1% (-0.28-3.1%)	4.1 (-1.1-12)	22 (-4.2-62)
Albania	1990	Female	95 plus	20% (12-30%)	80% (70-88%)	0.58 (0.25-1.1)	0.81% (-0.48-2.8%)	0.9% (-0.67-3.3%)	1.5 (-1.2-5.6)	5.6 (-3.2-20)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	1990	Male	All Ages	45% (40-51%)	55% (49-60%)	1.2 (1-1.4)	2.6% (1.9-3.4%)	2.9% (1.3-4.6%)	280 (130-450)	13000 (9200-18000)
Albania	1990	Male	15 to 19	39% (29-49%)	61% (51-71%)	0.71 (0.44-1.1)	2.9% (2-4%)	3.4% (1.9-5.3%)	4.9 (2.7-7.8)	600 (410-860)
Albania	1990	Male	20 to 24	59% (48-69%)	41% (31-52%)	1.6 (1-2.1)	6.5% (4.6-8.6%)	7.3% (4.4-11%)	16 (9.3-23)	1600 (1100-2300)
Albania	1990	Male	25 to 29	65% (53-74%)	35% (26-47%)	1.7 (1.2-2.3)	6.6% (4.8-8.5%)	7.5% (4.6-11%)	16 (9.5-22)	1600 (1200-2200)
Albania	1990	Male	30 to 34	67% (56-76%)	33% (24-44%)	1.9 (1.3-2.5)	6.2% (4.6-8%)	7.5% (4.7-11%)	14 (8.8-20)	1400 (990-1800)
Albania	1990	Male	35 to 39	67% (56-75%)	33% (25-44%)	2.1 (1.4-2.8)	5.7% (4-7.5%)	7.6% (4.7-11%)	13 (8.3-19)	1100 (770-1500)
Albania	1990	Male	40 to 44	69% (57-76%)	31% (24-43%)	2.2 (1.5-3)	5.1% (3.6-6.7%)	6.9% (4.3-9.6%)	13 (7.9-18)	900 (620-1200)
Albania	1990	Male	45 to 49	72% (60-80%)	28% (20-40%)	2.3 (1.6-3.3)	4.4% (2.8-6.3%)	5.9% (3.1-9.1%)	14 (7.5-23)	870 (530-1300)
Albania	1990	Male	50 to 54	69% (57-79%)	31% (21-43%)	2.4 (1.6-3.3)	4.2% (2.3-6.3%)	5.4% (2.6-8.8%)	21 (9.8-34)	980 (550-1500)
Albania	1990	Male	55 to 59	62% (44-79%)	38% (21-56%)	2.4 (1.4-3.5)	4.2% (2.1-7%)	5.3% (2.3-9.4%)	28 (12-49)	1100 (530-1800)
Albania	1990	Male	60 to 64	58% (39-77%)	42% (23-61%)	2.2 (1.3-3.4)	4.1% (1.8-7.2%)	5% (1.9-9%)	38 (15-70)	1200 (500-2100)
Albania	1990	Male	65 to 69	56% (38-74%)	44% (26-62%)	2 (1.2-2.8)	3.4% (1.3-6.2%)	4% (1.4-7.4%)	33 (11-61)	820 (300-1500)
Albania	1990	Male	70 to 74	53% (36-69%)	47% (31-64%)	1.7 (1-2.6)	2.6% (0.3-5.3%)	3.1% (0.19-6.3%)	33 (2-69)	650 (76-1300)
Albania	1990	Male	75 to 79	48% (33-65%)	52% (35-67%)	1.4 (0.77-2.2)	1.9% (-0.23-4.6%)	2.2% (-0.46-5.4%)	24 (-5-58)	360 (-43-860)
Albania	1990	Male	80 to 84	45% (29-62%)	55% (38-71%)	1.2 (0.67-1.9)	1.3% (-0.68-3.4%)	1.4% (-0.92-3.9%)	12 (-7.6-32)	130 (-68-340)
Albania	1990	Male	85 to 89	44% (30-59%)	56% (41-70%)	1.1 (0.63-1.7)	0.77% (-0.92-2.6%)	0.77% (-1.3-2.9%)	3.8 (-6.1-14)	33 (-39-110)
Albania	1990	Male	90 to 94	45% (31-60%)	55% (40-69%)	0.99 (0.57-1.6)	0.2% (-1.2-1.9%)	0.065% (-1.7-2.1%)	0.16 (-4.2-5)	3.1 (-18-28)
Albania	1990	Male	95 plus	46% (30-63%)	54% (37-70%)	0.9 (0.43-1.6)	-0.093% (-1.6-1.6%)	-0.37% (-2.3-1.8%)	-0.29 (-1.7-1.3)	-0.29 (-5.1-5.1)
Albania	1995	Female	All Ages	26% (22-30%)	74% (70-78%)	0.61 (0.5-0.73)	1.6% (1.1-2.2%)	2.6% (1.3-4.3%)	190 (95-310)	6200 (4300-8100)
Albania	1995	Female	15 to 19	26% (18-34%)	74% (66-82%)	0.62 (0.39-0.9)	1.7% (1.1-2.4%)	2.4% (1.2-4%)	1.6 (0.75-2.6)	250 (160-370)
Albania	1995	Female	20 to 24	34% (24-45%)	66% (55-76%)	0.92 (0.6-1.3)	3% (2.1-4.1%)	3.2% (1.6-5%)	2.3 (1.2-3.7)	470 (320-650)
Albania	1995	Female	25 to 29	35% (25-46%)	65% (54-75%)	0.84 (0.56-1.2)	2.9% (2.1-3.7%)	2.8% (1.5-4.4%)	2.1 (1.2-3.3)	440 (300-610)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	1995	Female	30 to 34	37% (26-48%)	63% (52-74%)	0.84 (0.54-1.2)	2.6% (1.9-3.4%)	2.8% (1.6-4.2%)	2.6 (1.5-3.9)	450 (310-620)
Albania	1995	Female	35 to 39	38% (27-49%)	62% (51-73%)	0.9 (0.57-1.3)	2.4% (1.8-3.1%)	3.2% (1.9-4.6%)	3.2 (1.9-4.8)	420 (290-560)
Albania	1995	Female	40 to 44	37% (27-48%)	63% (52-73%)	0.91 (0.6-1.3)	2.3% (1.6-3%)	3.4% (2-5%)	3.9 (2.3-5.9)	370 (240-490)
Albania	1995	Female	45 to 49	35% (23-45%)	65% (55-77%)	0.89 (0.53-1.3)	2.1% (1.4-2.9%)	3.3% (1.9-5.1%)	4.2 (2.4-6.4)	300 (200-420)
Albania	1995	Female	50 to 54	33% (23-44%)	67% (56-77%)	1 (0.65-1.5)	2.4% (1.6-3.4%)	4% (2.2-6.1%)	7.1 (4-11)	370 (240-540)
Albania	1995	Female	55 to 59	32% (21-46%)	68% (54-79%)	1.3 (0.74-2.1)	3.1% (1.7-4.9%)	5.4% (2.7-8.6%)	13 (6.6-21)	530 (290-810)
Albania	1995	Female	60 to 64	30% (19-45%)	70% (55-81%)	1.3 (0.77-2.1)	3.5% (1.8-5.7%)	5.7% (2.7-9.3%)	20 (9.5-32)	610 (310-970)
Albania	1995	Female	65 to 69	28% (18-40%)	72% (60-82%)	1.1 (0.7-1.7)	3.3% (1.7-5.4%)	5.1% (2.6-8.4%)	25 (13-41)	610 (330-990)
Albania	1995	Female	70 to 74	24% (16-34%)	76% (66-84%)	0.91 (0.5-1.5)	2.8% (1.5-4%)	4% (1.4-7.6%)	29 (9.9-55)	550 (190-1000)
Albania	1995	Female	75 to 79	20% (13-30%)	80% (70-87%)	0.72 (0.38-1.3)	2.2% (0.56-4.6%)	3% (0.68-6.2%)	26 (6-54)	370 (94-780)
Albania	1995	Female	80 to 84	17% (11-25%)	83% (75-89%)	0.62 (0.35-1)	2% (0.45-4.1%)	2.4% (0.54-5.1%)	23 (4.9-48)	240 (55-510)
Albania	1995	Female	85 to 89	17% (10-25%)	83% (75-90%)	0.6 (0.32-1)	1.7% (0.36-3.6%)	2.1% (0.42-4.6%)	16 (3.1-35)	120 (25-260)
Albania	1995	Female	90 to 94	18% (12-27%)	82% (73-88%)	0.63 (0.32-1.1)	1.4% (0.022-3.6%)	1.7% (-0.018-4.4%)	7.1 (-0.072-18)	38 (0.57-94)
Albania	1995	Female	95 plus	21% (12-31%)	79% (69-88%)	0.69 (0.3-1.3)	1.3% (-0.32-3.9%)	1.5% (-0.45-4.7%)	2.6 (-0.76-7.9)	9.4 (-2.2-27)
Albania	1995	Male	All Ages	45% (40-50%)	55% (50-60%)	1.4 (1.2-1.6)	3.3% (2.4-4.3%)	3.9% (2.1-6%)	380 (200-580)	16000 (11000-21000)
Albania	1995	Male	15 to 19	40% (30-50%)	60% (50-70%)	0.85 (0.53-1.3)	3.3% (2.2-4.6%)	4% (2.2-6.2%)	6.4 (3.6-10)	690 (450-980)
Albania	1995	Male	20 to 24	60% (49-69%)	40% (31-51%)	1.8 (1.2-2.5)	7.3% (5.1-9.9%)	8.5% (5.2-12%)	20 (12-29)	1800 (1200-2400)
Albania	1995	Male	25 to 29	65% (53-74%)	35% (26-47%)	2 (1.4-2.7)	7.6% (5.3-10%)	8.9% (5.5-13%)	19 (12-27)	1600 (1100-2200)
Albania	1995	Male	30 to 34	67% (56-76%)	33% (24-44%)	2.2 (1.6-3)	7.1% (5.2-9.4%)	8.8% (5.9-12%)	19 (12-26)	1500 (1100-2000)
Albania	1995	Male	35 to 39	68% (56-75%)	32% (25-44%)	2.4 (1.6-3.3)	6.6% (4.7-8.7%)	8.7% (5.8-12%)	17 (12-25)	1300 (960-1700)
Albania	1995	Male	40 to 44	69% (59-77%)	31% (23-41%)	2.6 (1.7-3.5)	5.9% (4.2-7.9%)	8.1% (5.2-11%)	17 (11-24)	1100 (770-1500)
Albania	1995	Male	45 to 49	72% (60-80%)	28% (20-40%)	2.7 (1.9-3.8)	5.2% (3.3-7.6%)	7.2% (4-11%)	16 (9-25)	930 (590-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	1995	Male	50 to 54	69% (58-79%)	31% (21-42%)	2.8 (1.9-3.9)	5% (2.8-7.5%)	6.6% (3.3-11%)	23 (12-38)	1100 (630-1700)
Albania	1995	Male	55 to 59	63% (45-79%)	37% (21-55%)	2.8 (1.7-4)	5.1% (2.7-8.3%)	6.6% (3.1-11%)	37 (17-62)	1400 (730-2200)
Albania	1995	Male	60 to 64	59% (39-76%)	41% (24-61%)	2.6 (1.6-3.9)	5.1% (2.3-8.8%)	6.2% (2.6-11%)	49 (22-89)	1500 (680-2600)
Albania	1995	Male	65 to 69	57% (40-74%)	43% (26-60%)	2.3 (1.5-3.3)	4.3% (1.9-7.5%)	5.2% (2.1-9.3%)	51 (21-92)	1300 (550-2200)
Albania	1995	Male	70 to 74	53% (37-69%)	47% (31-63%)	2 (1.2-3)	3.5% (0.85-6.8%)	4.1% (0.89-8.1%)	43 (9.2-85)	820 (210-1600)
Albania	1995	Male	75 to 79	49% (34-66%)	51% (34-66%)	1.6 (0.9-2.6)	2.7% (0.15-6.1%)	3.1% (-0.004-7.1%)	34 (-0.045-79)	510 (28-1100)
Albania	1995	Male	80 to 84	46% (31-63%)	54% (37-69%)	1.4 (0.81-2.2)	2% (-0.24-4.6%)	2.2% (-0.46-5.3%)	20 (-4.2-49)	230 (-28-530)
Albania	1995	Male	85 to 89	45% (31-59%)	55% (41-69%)	1.3 (0.74-2)	1.4% (-0.52-3.6%)	1.5% (-0.86-4.2%)	8.2 (-4.7-23)	67 (-26-180)
Albania	1995	Male	90 to 94	45% (31-61%)	55% (39-69%)	1.2 (0.66-1.9)	0.69% (-1-2.9%)	0.63% (-1.5-3.2%)	1.5 (-3.5-7.6)	10 (-15-41)
Albania	1995	Male	95 plus	47% (30-65%)	53% (35-70%)	1.1 (0.5-2)	0.33% (-1.4-2.7%)	0.097% (-2-3%)	0.07 (-1.5-2.2)	0.97 (-4.2-8.1)
Albania	2000	Female	All Ages	26% (23-30%)	74% (70-77%)	0.78 (0.64-0.94)	2.4% (1.7-3.2%)	4.1% (2.2-6.3%)	300 (170-470)	8300 (5800-11000)
Albania	2000	Female	15 to 19	26% (18-34%)	74% (66-82%)	0.76 (0.49-1.1)	1.9% (1.3-2.9%)	3.1% (1.5-5.4%)	2.1 (1-3.6)	300 (190-440)
Albania	2000	Female	20 to 24	34% (24-44%)	66% (56-76%)	1.1 (0.72-1.6)	3.4% (2.3-4.5%)	4.1% (2.1-6.6%)	3 (1.5-4.7)	490 (330-700)
Albania	2000	Female	25 to 29	35% (25-47%)	65% (53-75%)	1 (0.69-1.5)	3.2% (2.3-4.2%)	3.6% (2-5.7%)	2.8 (1.5-4.4)	460 (320-640)
Albania	2000	Female	30 to 34	37% (25-49%)	63% (51-75%)	1 (0.66-1.5)	2.9% (2.1-3.8%)	3.6% (2.1-5.6%)	3 (1.7-4.7)	450 (310-630)
Albania	2000	Female	35 to 39	38% (27-49%)	62% (51-73%)	1.1 (0.71-1.6)	2.8% (2-3.6%)	4% (2.5-5.9%)	4.5 (2.8-6.7)	500 (360-670)
Albania	2000	Female	40 to 44	37% (27-48%)	63% (52-73%)	1.1 (0.74-1.6)	2.6% (1.9-3.5%)	4.3% (2.7-6.3%)	6.1 (3.8-8.9)	500 (340-660)
Albania	2000	Female	45 to 49	35% (24-45%)	65% (55-76%)	1.1 (0.66-1.7)	2.5% (1.7-3.5%)	4.2% (2.6-6.4%)	6.8 (4.1-10)	440 (300-600)
Albania	2000	Female	50 to 54	33% (23-44%)	67% (56-77%)	1.3 (0.8-1.9)	2.9% (2-4.3%)	5% (3.2-7.6%)	9.6 (6.1-15)	470 (320-680)
Albania	2000	Female	55 to 59	32% (21-46%)	68% (54-79%)	1.6 (0.91-2.6)	3.9% (2.2-6%)	6.7% (3.6-10%)	17 (8.9-26)	630 (370-950)
Albania	2000	Female	60 to 64	30% (19-45%)	70% (55-81%)	1.6 (0.95-2.6)	4.5% (2.4-7.1%)	7.2% (3.9-11%)	29 (15-45)	870 (480-1300)
Albania	2000	Female	65 to 69	28% (18-40%)	72% (60-82%)	1.4 (0.87-2.1)	4.4% (2.5-7%)	6.6% (3.7-11%)	36 (20-57)	870 (500-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	2000	Female	70 to 74	24% (16-35%)	76% (65-84%)	1.1 (0.61-1.8)	4% (1.7-7%)	5.5% (2.3-9.6%)	46 (20-81)	870 (390-1500)
Albania	2000	Female	75 to 79	20% (13-31%)	80% (69-87%)	0.89 (0.45-1.5)	3.3% (1.1-6.6%)	4.3% (1.4-8.7%)	44 (14-86)	630 (210-1300)
Albania	2000	Female	80 to 84	17% (11-26%)	83% (74-89%)	0.76 (0.42-1.3)	3.1% (1.1-5.8%)	3.8% (1.3-7.2%)	41 (14-78)	430 (150-810)
Albania	2000	Female	85 to 89	17% (10-25%)	83% (75-90%)	0.73 (0.38-1.2)	2.7% (0.88-5.3%)	3.4% (1-6.5%)	31 (9.6-60)	230 (73-450)
Albania	2000	Female	90 to 94	18% (12-26%)	82% (74-88%)	0.77 (0.39-1.3)	2.4% (0.53-5.2%)	3% (0.59-6.3%)	16 (3.3-35)	85 (18-180)
Albania	2000	Female	95 plus	20% (12-30%)	80% (70-88%)	0.85 (0.38-1.6)	2.4% (0.035-5.9%)	2.8% (-0.084-7.1%)	6.1 (-0.19-16)	22 (0.31-53)
Albania	2000	Male	All Ages	47% (42-52%)	53% (48-58%)	1.8 (1.5-2.2)	4.7% (3.4-6.2%)	5.8% (3.3-8.7%)	600 (340-910)	22000 (15000-29000)
Albania	2000	Male	15 to 19	40% (30-49%)	60% (51-70%)	1 (0.65-1.5)	3.6% (2.4-5.1%)	4.5% (2.5-6.9%)	7 (4-11)	720 (480-1000)
Albania	2000	Male	20 to 24	60% (48-70%)	40% (30-52%)	2.3 (1.5-3.1)	8% (5.6-11%)	9.6% (6-14%)	21 (13-29)	1800 (1200-2400)
Albania	2000	Male	25 to 29	65% (54-74%)	35% (26-46%)	2.5 (1.7-3.4)	8.3% (5.8-11%)	10% (6.3-14%)	22 (14-32)	1800 (1200-2400)
Albania	2000	Male	30 to 34	67% (56-76%)	33% (24-44%)	2.7 (1.9-3.7)	8% (5.7-11%)	10% (6.6-14%)	22 (15-31)	1700 (1200-2300)
Albania	2000	Male	35 to 39	68% (57-75%)	32% (25-43%)	3 (2-4.1)	7.5% (5.2-10%)	10% (6.6-15%)	25 (16-36)	1800 (1200-2400)
Albania	2000	Male	40 to 44	69% (59-77%)	31% (23-41%)	3.2 (2.2-4.3)	6.9% (4.6-9.3%)	9.7% (6-14%)	28 (17-40)	1800 (1200-2400)
Albania	2000	Male	45 to 49	72% (61-80%)	28% (20-39%)	3.4 (2.3-4.7)	6.3% (3.9-9.2%)	8.8% (4.9-14%)	28 (16-44)	1500 (970-2200)
Albania	2000	Male	50 to 54	69% (57-80%)	31% (20-43%)	3.4 (2.3-4.7)	6.1% (3.5-9.1%)	8.3% (4.1-13%)	33 (17-52)	1500 (860-2300)
Albania	2000	Male	55 to 59	63% (45-79%)	37% (21-55%)	3.4 (2.2-5)	6.5% (3.7-10%)	8.6% (4.6-14%)	48 (25-78)	1800 (1000-2800)
Albania	2000	Male	60 to 64	59% (40-77%)	41% (23-60%)	3.2 (2-4.8)	6.6% (3.2-11%)	8.4% (3.9-14%)	74 (34-120)	2200 (1100-3500)
Albania	2000	Male	65 to 69	57% (39-74%)	43% (26-61%)	2.9 (1.8-4.1)	6% (2.8-9.8%)	7.3% (3.4-12%)	82 (37-140)	2000 (960-3300)
Albania	2000	Male	70 to 74	53% (37-69%)	47% (31-63%)	2.4 (1.5-3.7)	5.1% (1.8-9.5%)	6.1% (1.9-12%)	84 (27-160)	1600 (570-3000)
Albania	2000	Male	75 to 79	49% (34-65%)	51% (35-66%)	2 (1.1-3.2)	4.2% (0.87-8.8%)	4.9% (0.79-10%)	58 (9.6-120)	850 (170-1800)
Albania	2000	Male	80 to 84	46% (30-63%)	54% (37-70%)	1.7 (0.98-2.8)	3.5% (0.57-7.6%)	3.9% (0.53-8.7%)	40 (5.4-90)	430 (74-940)
Albania	2000	Male	85 to 89	45% (31-59%)	55% (41-69%)	1.6 (0.89-2.4)	2.6% (0.19-6%)	2.9% (-0.029-7%)	19 (-0.19-45)	150 (11-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	2000	Male	90 to 94	45% (31-60%)	55% (40-69%)	1.4 (0.79-2.4)	1.6% (-0.52-4.7%)	1.7% (-0.85-5.4%)	5.2 (-2.5-16)	30 (-9.3-84)
Albania	2000	Male	95 plus	46% (29-65%)	54% (35-71%)	1.3 (0.62-2.4)	1.1% (-1.1-4.6%)	0.98% (-1.7-5.1%)	0.83 (-1.5-4.4)	3.8 (-3.7-16)
Albania	2005	Female	All Ages	25% (23-27%)	75% (73-77%)	0.94 (0.77-1.1)	3.3% (2.3-4.4%)	5.4% (3.1-8.2%)	480 (270-720)	11000 (7900-15000)
Albania	2005	Female	15 to 19	22% (19-27%)	78% (73-81%)	0.8 (0.58-1.1)	2.1% (1.4-2.9%)	3.6% (1.9-5.7%)	2.3 (1.2-3.7)	330 (210-470)
Albania	2005	Female	20 to 24	31% (26-36%)	69% (64-74%)	1.2 (0.9-1.6)	3.7% (2.7-4.9%)	5% (2.8-7.6%)	3.5 (2-5.4)	540 (370-760)
Albania	2005	Female	25 to 29	33% (29-38%)	67% (62-71%)	1.2 (0.89-1.6)	3.5% (2.6-4.5%)	4.5% (2.6-6.6%)	2.9 (1.7-4.2)	440 (300-580)
Albania	2005	Female	30 to 34	34% (29-40%)	66% (60-71%)	1.1 (0.84-1.5)	3.1% (2.3-4.1%)	4.2% (2.6-6.1%)	3.2 (2-4.7)	430 (300-570)
Albania	2005	Female	35 to 39	35% (31-39%)	65% (61-69%)	1.2 (0.91-1.6)	3% (2.2-3.8%)	4.6% (3-6.3%)	4.8 (3.1-6.7)	500 (360-650)
Albania	2005	Female	40 to 44	33% (29-38%)	67% (62-71%)	1.2 (0.92-1.6)	2.9% (2.1-3.7%)	4.8% (3.2-6.6%)	7 (4.8-9.7)	580 (420-740)
Albania	2005	Female	45 to 49	32% (27-36%)	68% (64-73%)	1.2 (0.86-1.6)	2.8% (2-3.7%)	4.6% (3.1-6.7%)	9.4 (6.3-13)	600 (430-800)
Albania	2005	Female	50 to 54	32% (24-42%)	68% (58-76%)	1.5 (0.98-2.1)	3.4% (2.4-4.7%)	5.7% (3.8-8.4%)	14 (9.4-21)	690 (490-960)
Albania	2005	Female	55 to 59	32% (21-47%)	68% (53-79%)	2 (1.1-3.1)	4.7% (2.7-7%)	8% (4.4-12%)	23 (12-34)	860 (510-1300)
Albania	2005	Female	60 to 64	31% (19-46%)	69% (54-81%)	2 (1.1-3.2)	5.5% (3.1-8.5%)	8.6% (4.8-13%)	37 (20-57)	1100 (640-1700)
Albania	2005	Female	65 to 69	28% (19-41%)	72% (59-81%)	1.8 (1.1-2.6)	5.5% (3.3-8.5%)	8.1% (4.8-13%)	54 (32-83)	1300 (790-2000)
Albania	2005	Female	70 to 74	25% (16-36%)	75% (64-84%)	1.4 (0.79-2.3)	5.1% (2.4-8.7%)	7% (3.2-12%)	69 (32-110)	1300 (610-2200)
Albania	2005	Female	75 to 79	21% (13-30%)	79% (70-87%)	1.1 (0.58-1.9)	4.4% (1.7-8%)	5.7% (2.1-11%)	74 (27-140)	1100 (400-2000)
Albania	2005	Female	80 to 84	18% (11-26%)	82% (74-89%)	0.95 (0.52-1.6)	4.3% (1.7-7.6%)	5.2% (2-9.2%)	75 (29-130)	780 (310-1400)
Albania	2005	Female	85 to 89	17% (11-25%)	83% (75-89%)	0.91 (0.49-1.5)	3.9% (1.5-7.5%)	4.8% (1.7-9.2%)	54 (20-100)	400 (150-760)
Albania	2005	Female	90 to 94	19% (12-27%)	81% (73-88%)	0.95 (0.48-1.6)	3.5% (1-7.1%)	4.3% (1.2-8.7%)	31 (8.6-64)	160 (48-330)
Albania	2005	Female	95 plus	21% (12-31%)	79% (69-88%)	1 (0.46-1.9)	3.4% (0.45-7.8%)	4% (0.43-9.3%)	13 (1.4-30)	45 (5.8-100)
Albania	2005	Male	All Ages	51% (47-54%)	49% (46-53%)	2.4 (2-2.8)	6.1% (4.3-7.9%)	7.6% (4.4-11%)	850 (490-1200)	27000 (19000-35000)
Albania	2005	Male	15 to 19	38% (33-44%)	62% (56-67%)	1.2 (0.76-1.7)	3.5% (2.4-4.8%)	4.5% (2.4-6.8%)	6.4 (3.5-9.7)	720 (480-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	2005	Male	20 to 24	62% (56-66%)	38% (34-44%)	2.8 (1.9-3.7)	8.5% (5.9-11%)	11% (6.2-15%)	18 (11-25)	1600 (1100-2200)
Albania	2005	Male	25 to 29	70% (65-74%)	30% (26-35%)	3.2 (2.2-4.3)	9% (6.2-12%)	11% (6.8-16%)	17 (10-25)	1500 (1000-2000)
Albania	2005	Male	30 to 34	71% (66-75%)	29% (25-34%)	3.5 (2.4-4.6)	8.5% (5.9-12%)	11% (6.8-16%)	19 (11-28)	1500 (1000-2100)
Albania	2005	Male	35 to 39	70% (66-74%)	30% (26-34%)	3.7 (2.5-5.1)	8.3% (5.5-11%)	12% (6.9-17%)	22 (13-32)	1600 (1100-2200)
Albania	2005	Male	40 to 44	73% (69-77%)	27% (23-31%)	4.1 (2.9-5.4)	8% (5.1-11%)	12% (6.6-17%)	32 (18-47)	2000 (1300-2700)
Albania	2005	Male	45 to 49	77% (72-80%)	23% (20-28%)	4.4 (3.2-6)	7.8% (4.8-11%)	11% (6.1-17%)	41 (22-63)	2200 (1300-3100)
Albania	2005	Male	50 to 54	72% (62-81%)	28% (19-38%)	4.3 (3-5.9)	7.7% (4.6-11%)	11% (5.5-16%)	52 (27-78)	2300 (1300-3300)
Albania	2005	Male	55 to 59	63% (46-79%)	37% (21-54%)	4.2 (2.6-6)	7.9% (4.6-12%)	11% (5.8-16%)	62 (34-93)	2300 (1300-3300)
Albania	2005	Male	60 to 64	59% (40-77%)	41% (23-60%)	3.9 (2.4-5.8)	8.1% (4.3-12%)	10% (5.4-16%)	94 (48-140)	2800 (1500-4200)
Albania	2005	Male	65 to 69	58% (40-75%)	42% (25-60%)	3.5 (2.2-4.9)	7.7% (4-12%)	9.5% (4.7-14%)	120 (60-180)	2900 (1500-4400)
Albania	2005	Male	70 to 74	54% (37-70%)	46% (30-63%)	3 (1.8-4.5)	6.9% (2.7-12%)	8.2% (3.1-14%)	130 (49-230)	2500 (1000-4200)
Albania	2005	Male	75 to 79	49% (35-65%)	51% (35-65%)	2.4 (1.4-3.9)	5.8% (1.6-11%)	6.8% (1.8-13%)	110 (30-210)	1600 (470-3100)
Albania	2005	Male	80 to 84	47% (31-64%)	53% (36-69%)	2.1 (1.2-3.3)	5.1% (1.4-10%)	5.8% (1.6-12%)	76 (20-160)	810 (230-1700)
Albania	2005	Male	85 to 89	46% (32-60%)	54% (40-68%)	1.9 (1.1-2.9)	4.1% (0.88-8.7%)	4.7% (0.84-10%)	33 (5.9-70)	250 (55-530)
Albania	2005	Male	90 to 94	46% (32-61%)	54% (39-68%)	1.7 (0.99-2.8)	2.9% (0.11-7.2%)	3.2% (-0.13-8.2%)	12 (-0.49-31)	65 (2.6-160)
Albania	2005	Male	95 plus	47% (31-66%)	53% (34-69%)	1.6 (0.76-2.9)	2.1% (-0.84-6.8%)	2.1% (-1.4-7.8%)	2.4 (-1.5-8.8)	9.4 (-3.8-31)
Albania	2010	Female	All Ages	26% (24-29%)	74% (71-76%)	1.1 (0.88-1.3)	3.6% (2.5-5%)	5.9% (3.3-9.1%)	540 (300-840)	12000 (8400-16000)
Albania	2010	Female	15 to 19	23% (20-27%)	77% (73-80%)	0.85 (0.62-1.1)	2.1% (1.4-3%)	3.8% (2-6.1%)	1.8 (0.94-3)	270 (170-390)
Albania	2010	Female	20 to 24	31% (27-36%)	69% (64-73%)	1.3 (0.93-1.8)	3.8% (2.7-5.1%)	5.4% (3.1-8.4%)	2.9 (1.6-4.5)	470 (320-660)
Albania	2010	Female	25 to 29	34% (29-39%)	66% (61-71%)	1.3 (0.93-1.7)	3.6% (2.7-4.7%)	4.8% (2.8-7.5%)	2.5 (1.4-3.7)	400 (280-550)
Albania	2010	Female	30 to 34	34% (30-39%)	66% (61-70%)	1.2 (0.88-1.7)	3.2% (2.3-4.3%)	4.5% (2.8-6.8%)	2.6 (1.6-3.9)	370 (260-510)
Albania	2010	Female	35 to 39	35% (31-40%)	65% (60-69%)	1.3 (0.93-1.7)	3.1% (2.3-3.9%)	4.8% (3.1-6.8%)	3.9 (2.5-5.6)	430 (320-560)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	2010	Female	40 to 44	34% (30-39%)	66% (61-70%)	1.3 (0.95-1.8)	2.9% (2.2-3.9%)	5.1% (3.5-7%)	5.8 (4-8)	490 (360-650)
Albania	2010	Female	45 to 49	32% (28-37%)	68% (63-72%)	1.3 (0.9-1.8)	2.9% (2.1-3.9%)	5% (3.3-7.3%)	9.8 (6.5-14)	630 (460-860)
Albania	2010	Female	50 to 54	33% (25-42%)	67% (58-75%)	1.6 (1.1-2.3)	3.5% (2.4-4.9%)	6.1% (4.1-8.7%)	17 (11-24)	830 (590-1100)
Albania	2010	Female	55 to 59	33% (22-47%)	67% (53-78%)	2.1 (1.2-3.4)	4.8% (2.9-7.1%)	8.4% (4.8-12%)	28 (16-43)	1100 (670-1600)
Albania	2010	Female	60 to 64	31% (20-46%)	69% (54-80%)	2.2 (1.2-3.3)	5.5% (3.2-8.3%)	9% (5.2-14%)	39 (23-62)	1200 (730-1900)
Albania	2010	Female	65 to 69	29% (19-40%)	71% (60-81%)	1.9 (1.2-2.8)	5.6% (3.2-8.6%)	8.5% (4.9-13%)	56 (32-85)	1400 (820-2100)
Albania	2010	Female	70 to 74	25% (17-35%)	75% (65-83%)	1.5 (0.83-2.4)	5.2% (2.4-8.6%)	7.2% (3.4-12%)	79 (37-130)	1500 (710-2500)
Albania	2010	Female	75 to 79	21% (13-31%)	79% (69-87%)	1.2 (0.6-2)	4.5% (1.7-8.2%)	5.9% (2.1-11%)	80 (29-150)	1200 (430-2200)
Albania	2010	Female	80 to 84	18% (12-27%)	82% (73-88%)	1 (0.55-1.7)	4.4% (1.7-7.8%)	5.5% (2.1-9.8%)	87 (33-150)	920 (360-1600)
Albania	2010	Female	85 to 89	18% (11-25%)	82% (75-89%)	0.97 (0.53-1.6)	4.2% (1.6-7.9%)	5.1% (1.9-9.7%)	68 (26-130)	510 (200-950)
Albania	2010	Female	90 to 94	19% (12-27%)	81% (73-88%)	1 (0.53-1.7)	3.9% (1.2-8%)	4.7% (1.4-9.7%)	40 (12-83)	210 (67-420)
Albania	2010	Female	95 plus	21% (12-32%)	79% (68-88%)	1.1 (0.49-2.1)	3.8% (0.63-8.8%)	4.5% (0.64-11%)	18 (2.6-42)	63 (10-140)
Albania	2010	Male	All Ages	53% (48-56%)	47% (44-52%)	2.7 (2.2-3.3)	6.6% (4.7-8.7%)	8.2% (4.7-12%)	960 (560-1400)	28000 (20000-37000)
Albania	2010	Male	15 to 19	38% (34-44%)	62% (56-66%)	1.2 (0.8-1.8)	3.5% (2.4-5%)	4.5% (2.4-7.2%)	5.4 (2.9-8.6)	640 (420-910)
Albania	2010	Male	20 to 24	62% (57-67%)	38% (33-43%)	2.9 (2-4.1)	8.6% (6-12%)	11% (6.3-16%)	18 (10-26)	1700 (1200-2400)
Albania	2010	Male	25 to 29	70% (65-74%)	30% (26-35%)	3.4 (2.3-4.6)	9.1% (6-12%)	12% (6.8-17%)	17 (9.6-25)	1500 (1000-2000)
Albania	2010	Male	30 to 34	71% (66-75%)	29% (25-34%)	3.7 (2.5-5.1)	8.8% (6.1-12%)	12% (6.9-18%)	15 (8.9-23)	1200 (820-1700)
Albania	2010	Male	35 to 39	70% (66-74%)	30% (26-34%)	3.9 (2.7-5.5)	8.5% (5.5-12%)	12% (7-18%)	19 (10-28)	1400 (890-1900)
Albania	2010	Male	40 to 44	73% (69-77%)	27% (23-31%)	4.3 (3.1-5.9)	8.4% (5.3-12%)	12% (7.1-18%)	27 (15-40)	1700 (1100-2300)
Albania	2010	Male	45 to 49	77% (72-80%)	23% (20-28%)	4.6 (3.3-6.4)	8.2% (5.1-12%)	12% (6.6-17%)	41 (23-61)	2200 (1300-3100)
Albania	2010	Male	50 to 54	72% (61-81%)	28% (19-39%)	4.5 (3.1-6.3)	8.2% (4.9-12%)	11% (6.2-17%)	59 (32-87)	2600 (1600-3700)
Albania	2010	Male	55 to 59	64% (45-80%)	36% (20-55%)	4.4 (2.9-6.4)	8.3% (4.9-12%)	11% (6.3-17%)	72 (41-110)	2700 (1600-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	2010	Male	60 to 64	59% (40-78%)	41% (22-60%)	4.2 (2.6-6.1)	8.5% (4.8-12%)	11% (5.9-16%)	95 (51-140)	2900 (1600-4200)
Albania	2010	Male	65 to 69	58% (40-75%)	42% (25-60%)	3.7 (2.4-5.3)	8.1% (4.6-12%)	10% (5.6-15%)	120 (69-190)	3100 (1700-4500)
Albania	2010	Male	70 to 74	54% (38-70%)	46% (30-62%)	3.1 (1.9-4.7)	7.3% (3.1-12%)	8.8% (3.5-15%)	160 (63-270)	3100 (1300-5100)
Albania	2010	Male	75 to 79	50% (34-67%)	50% (33-66%)	2.6 (1.4-4.1)	6.2% (1.9-12%)	7.4% (2.1-14%)	140 (41-270)	2100 (640-3900)
Albania	2010	Male	80 to 84	47% (31-63%)	53% (37-69%)	2.2 (1.3-3.6)	5.5% (1.7-11%)	6.3% (1.8-12%)	99 (29-190)	1100 (320-2100)
Albania	2010	Male	85 to 89	46% (32-60%)	54% (40-68%)	2 (1.2-3.2)	4.6% (1.2-9.5%)	5.2% (1.2-11%)	48 (11-100)	370 (93-780)
Albania	2010	Male	90 to 94	46% (32-61%)	54% (39-68%)	1.9 (1.1-3.1)	3.3% (0.2-8.2%)	3.7% (-0.0075-9.5%)	14 (-0.029-37)	78 (4.8-190)
Albania	2010	Male	95 plus	47% (31-65%)	53% (35-69%)	1.7 (0.81-3.1)	2.5% (-0.79-7.8%)	2.6% (-1.3-8.9%)	3.5 (-1.8-12)	13 (-4.4-42)
Albania	2016	Female	All Ages	29% (26-33%)	71% (67-74%)	1.2 (0.92-1.7)	3.6% (2.4-5.2%)	5.6% (2.9-9.1%)	540 (270-900)	12000 (7900-18000)
Albania	2016	Female	15 to 19	26% (19-35%)	74% (65-81%)	0.99 (0.58-1.5)	2.1% (1.4-3.1%)	4.3% (1.8-7.4%)	1.5 (0.65-2.6)	240 (140-350)
Albania	2016	Female	20 to 24	36% (25-46%)	64% (54-75%)	1.5 (0.9-2.2)	3.9% (2.7-5.3%)	5.9% (3-9.7%)	2.9 (1.5-4.9)	520 (350-740)
Albania	2016	Female	25 to 29	37% (26-48%)	63% (52-74%)	1.4 (0.85-2.2)	3.6% (2.6-4.8%)	5% (2.6-8.1%)	2.5 (1.3-4.1)	470 (310-660)
Albania	2016	Female	30 to 34	37% (26-50%)	63% (50-74%)	1.3 (0.8-2.2)	3.3% (2.3-4.5%)	4.8% (2.7-7.8%)	2.5 (1.4-4)	390 (260-560)
Albania	2016	Female	35 to 39	39% (27-50%)	61% (50-73%)	1.4 (0.82-2.2)	3.2% (2.2-4.2%)	5.2% (3-7.7%)	3.2 (1.8-5)	380 (260-520)
Albania	2016	Female	40 to 44	38% (27-49%)	62% (51-73%)	1.5 (0.88-2.2)	3.1% (2.1-4.3%)	5.6% (3.5-8.2%)	5.3 (3.2-8.2)	460 (310-640)
Albania	2016	Female	45 to 49	36% (24-46%)	64% (54-76%)	1.4 (0.81-2.2)	3% (2.1-4.3%)	5.4% (3.3-8.1%)	8.4 (5-14)	570 (380-820)
Albania	2016	Female	50 to 54	34% (24-45%)	66% (55-76%)	1.6 (1-2.5)	3.5% (2.4-5.2%)	6.2% (3.9-9.4%)	16 (10-26)	850 (560-1200)
Albania	2016	Female	55 to 59	33% (22-48%)	67% (52-78%)	2.1 (1.2-3.4)	4.6% (2.7-7%)	8.2% (4.6-13%)	31 (16-49)	1200 (720-1900)
Albania	2016	Female	60 to 64	31% (20-46%)	69% (54-80%)	2.1 (1.2-3.4)	5.3% (2.9-8%)	8.8% (4.8-13%)	46 (25-74)	1500 (830-2300)
Albania	2016	Female	65 to 69	29% (19-41%)	71% (59-81%)	1.9 (1.1-2.9)	5.2% (2.8-8.2%)	8.2% (4.3-13%)	55 (30-89)	1400 (790-2200)
Albania	2016	Female	70 to 74	25% (17-35%)	75% (65-83%)	1.5 (0.77-2.5)	4.7% (2.1-8.2%)	6.9% (2.9-12%)	67 (28-120)	1300 (560-2200)
Albania	2016	Female	75 to 79	20% (14-30%)	80% (70-86%)	1.2 (0.57-2)	4% (1.4-7.6%)	5.5% (1.8-10%)	79 (26-150)	1200 (400-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Albania	2016	Female	80 to 84	18% (12-26%)	82% (74-88%)	0.99 (0.52-1.7)	3.9% (1.3-7.3%)	5% (1.7-9.6%)	86 (30-170)	920 (330-1800)
Albania	2016	Female	85 to 89	17% (11-25%)	83% (75-89%)	0.96 (0.49-1.7)	3.8% (1.3-7.5%)	4.8% (1.6-9.4%)	71 (24-140)	530 (190-1000)
Albania	2016	Female	90 to 94	19% (12-27%)	81% (73-88%)	1 (0.51-1.8)	3.6% (1-7.8%)	4.5% (1.2-9.7%)	44 (12-94)	230 (64-480)
Albania	2016	Female	95 plus	21% (13-32%)	79% (68-87%)	1.1 (0.48-2.2)	3.7% (0.48-8.6%)	4.4% (0.45-11%)	18 (1.8-45)	63 (8-150)
Albania	2016	Male	All Ages	53% (47-58%)	47% (42-53%)	2.8 (2.1-3.6)	6.7% (4.4-9.5%)	7.9% (4-13%)	1000 (510-1700)	29000 (18000-42000)
Albania	2016	Male	15 to 19	41% (31-50%)	59% (50-69%)	1.3 (0.79-2.1)	3.7% (2.4-5.4%)	4.9% (2.5-8.3%)	4.5 (2.3-7.6)	540 (340-810)
Albania	2016	Male	20 to 24	61% (49-70%)	39% (30-51%)	2.9 (1.8-4.3)	8.6% (5.9-12%)	11% (6.1-16%)	17 (9.4-27)	1700 (1100-2400)
Albania	2016	Male	25 to 29	66% (55-75%)	34% (25-45%)	3.2 (2-4.7)	8.8% (5.9-12%)	11% (6-17%)	18 (9.2-27)	1700 (1100-2400)
Albania	2016	Male	30 to 34	68% (58-77%)	32% (23-42%)	3.5 (2.2-5.1)	8.6% (5.7-12%)	12% (6.4-18%)	14 (7.8-22)	1300 (830-1800)
Albania	2016	Male	35 to 39	68% (58-76%)	32% (24-42%)	3.8 (2.3-5.8)	8.6% (5.5-12%)	12% (6.6-19%)	14 (7.3-22)	1100 (670-1500)
Albania	2016	Male	40 to 44	70% (60-77%)	30% (23-40%)	4.1 (2.7-6.1)	8.3% (5.2-12%)	12% (6.8-19%)	21 (11-32)	1300 (820-1900)
Albania	2016	Male	45 to 49	73% (62-80%)	27% (20-38%)	4.3 (2.8-6.5)	8% (4.8-12%)	12% (6.2-18%)	32 (16-51)	1800 (1000-2600)
Albania	2016	Male	50 to 54	70% (58-80%)	30% (20-42%)	4.4 (2.9-6.5)	8% (4.5-12%)	11% (5.8-17%)	55 (28-86)	2500 (1400-3700)
Albania	2016	Male	55 to 59	64% (46-80%)	36% (20-54%)	4.4 (2.7-6.6)	8.3% (4.8-12%)	11% (6.1-17%)	84 (43-130)	3200 (1700-4700)
Albania	2016	Male	60 to 64	59% (41-77%)	41% (23-59%)	4.1 (2.6-6.4)	8.4% (4.5-12%)	11% (5.4-16%)	110 (58-180)	3500 (1900-5400)
Albania	2016	Male	65 to 69	57% (40-75%)	43% (25-60%)	3.7 (2.3-5.6)	7.9% (4.2-12%)	9.9% (5.1-16%)	130 (64-210)	3200 (1700-5100)
Albania	2016	Male	70 to 74	54% (37-70%)	46% (30-63%)	3.1 (1.8-5)	7.1% (2.8-12%)	8.7% (3.3-15%)	150 (56-260)	2900 (1100-5100)
Albania	2016	Male	75 to 79	49% (35-66%)	51% (34-65%)	2.6 (1.3-4.4)	6% (1.6-11%)	7.2% (1.8-14%)	150 (39-300)	2300 (620-4400)
Albania	2016	Male	80 to 84	47% (31-63%)	53% (37-69%)	2.2 (1.2-3.8)	5.3% (1.4-11%)	6.2% (1.4-13%)	120 (25-240)	1300 (300-2600)
Albania	2016	Male	85 to 89	46% (32-61%)	54% (39-68%)	2 (1.1-3.4)	4.4% (0.97-9.9%)	5.1% (0.92-12%)	67 (12-160)	510 (110-1200)
Albania	2016	Male	90 to 94	46% (33-61%)	54% (39-67%)	1.9 (1-3.2)	3.2% (0.018-8.2%)	3.7% (-0.26-9.6%)	22 (-1.5-58)	120 (0.63-300)
Albania	2016	Male	95 plus	47% (31-65%)	53% (35-69%)	1.7 (0.78-3.2)	2.4% (-0.78-7.8%)	2.5% (-1.3-8.9%)	3.7 (-1.9-13)	14 (-4.7-46)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	1990	Female	All Ages	12% (10-14%)	88% (86-90%)	1.9 (1.4-2.5)	4.9% (3.7-6.4%)	6.6% (4.5-9.2%)	990 (650-1400)	29000 (21000-37000)
Bosnia and Herzegovina	1990	Female	15 to 19	12% (7.4-17%)	88% (83-93%)	1.3 (0.64-2.3)	3.3% (2.2-4.7%)	6.9% (3.9-11%)	4.6 (2.6-7.4)	620 (400-860)
Bosnia and Herzegovina	1990	Female	20 to 24	15% (9.8-23%)	85% (77-90%)	2.1 (1-3.7)	5.4% (4-7.1%)	8.2% (5-12%)	6.6 (3.8-9.8)	1100 (760-1500)
Bosnia and Herzegovina	1990	Female	25 to 29	16% (10-23%)	84% (77-90%)	2.1 (1.1-3.7)	5.2% (3.9-6.7%)	8.2% (5.3-11%)	8.2 (5-12)	1300 (880-1700)
Bosnia and Herzegovina	1990	Female	30 to 34	16% (10-25%)	84% (75-90%)	2.3 (1.2-4.1)	5% (3.7-6.6%)	9.3% (6.2-13%)	11 (7-16)	1300 (910-1700)
Bosnia and Herzegovina	1990	Female	35 to 39	17% (10-24%)	83% (76-90%)	2.7 (1.4-4.7)	5.2% (3.9-7%)	9.7% (6.7-13%)	17 (11-25)	1500 (1100-2000)
Bosnia and Herzegovina	1990	Female	40 to 44	16% (11-23%)	84% (77-89%)	3.1 (1.6-5)	5.6% (4.2-7.5%)	11% (7.4-14%)	25 (16-36)	1600 (1200-2200)
Bosnia and Herzegovina	1990	Female	45 to 49	16% (9.4-22%)	84% (78-91%)	3.4 (1.7-6.1)	6.7% (4.8-9.2%)	11% (7.6-16%)	38 (25-56)	1900 (1300-2700)
Bosnia and Herzegovina	1990	Female	50 to 54	15% (9.8-21%)	85% (79-90%)	3.4 (1.8-5.7)	6.7% (4.7-9.3%)	10% (7.1-15%)	69 (45-100)	2900 (2000-4100)
Bosnia and Herzegovina	1990	Female	55 to 59	14% (8.8-20%)	86% (80-91%)	3.1 (1.6-5.2)	6.6% (4.4-9.3%)	9.5% (6.2-14%)	94 (60-140)	3400 (2300-4800)
Bosnia and Herzegovina	1990	Female	60 to 64	13% (7.9-19%)	87% (81-92%)	2.8 (1.4-4.9)	6.4% (4.3-9.3%)	8.6% (5.6-13%)	130 (82-190)	3800 (2500-5400)
Bosnia and Herzegovina	1990	Female	65 to 69	12% (7.6-16%)	88% (84-92%)	2.5 (1.3-4.2)	6.3% (4.1-9.3%)	8% (5.1-12%)	150 (92-230)	3500 (2200-5300)
Bosnia and Herzegovina	1990	Female	70 to 74	10% (6.7-14%)	90% (86-93%)	2 (1-3.7)	5.5% (3.4-8.4%)	6.8% (4.1-11%)	110 (63-170)	2000 (1200-3100)
Bosnia and Herzegovina	1990	Female	75 to 79	8.8% (5.5-13%)	91% (87-95%)	1.7 (0.76-3.2)	4.6% (2.6-7.4%)	5.5% (3.1-9.1%)	130 (69-210)	1800 (1000-3000)
Bosnia and Herzegovina	1990	Female	80 to 84	7.8% (4.6-11%)	92% (89-95%)	1.5 (0.68-2.6)	4% (2.1-6.6%)	4.8% (2.4-8%)	97 (48-170)	1000 (530-1800)
Bosnia and Herzegovina	1990	Female	85 to 89	7.5% (4.4-12%)	93% (88-96%)	1.4 (0.68-2.7)	3.8% (1.8-6.7%)	4.5% (2.1-8.1%)	69 (31-120)	520 (250-910)
Bosnia and Herzegovina	1990	Female	90 to 94	8% (4.9-12%)	92% (88-95%)	1.5 (0.71-3)	3.7% (1.8-6.9%)	4.4% (2-8.1%)	31 (14-58)	160 (76-290)
Bosnia and Herzegovina	1990	Female	95 plus	9.3% (5.2-15%)	91% (85-95%)	1.8 (0.69-3.6)	4% (1.7-7.3%)	4.6% (1.9-8.5%)	9.4 (3.8-18)	32 (14-59)
Bosnia and Herzegovina	1990	Male	All Ages	37% (32-42%)	63% (58-68%)	5 (3.8-6.6)	13% (10-15%)	15% (12-19%)	2600 (2000-3400)	90000 (71000-110000)
Bosnia and Herzegovina	1990	Male	15 to 19	34% (23-48%)	66% (52-77%)	2.4 (1.2-4.3)	6.8% (4.2-10%)	11% (5.5-18%)	16 (7.8-27)	1700 (1000-2500)
Bosnia and Herzegovina	1990	Male	20 to 24	44% (30-57%)	56% (43-70%)	4.7 (2.5-8)	14% (9.6-18%)	19% (12-27%)	47 (29-68)	4300 (3000-5800)
Bosnia and Herzegovina	1990	Male	25 to 29	41% (29-53%)	59% (47-71%)	5.3 (2.8-8.6)	14% (11-17%)	20% (14-26%)	58 (40-79)	5000 (3700-6400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	1990	Male	30 to 34	41% (29-54%)	59% (46-71%)	6 (3.3-9.6)	15% (12-18%)	22% (17-28%)	74 (54-96)	5500 (4200-6900)
Bosnia and Herzegovina	1990	Male	35 to 39	49% (34-62%)	51% (38-66%)	7 (3.9-11)	17% (14-21%)	25% (20-30%)	100 (78-130)	6700 (5200-8400)
Bosnia and Herzegovina	1990	Male	40 to 44	52% (38-66%)	48% (34-62%)	8.3 (4.8-13)	18% (15-22%)	26% (21-31%)	130 (100-170)	7300 (5800-9200)
Bosnia and Herzegovina	1990	Male	45 to 49	52% (37-66%)	48% (34-63%)	9.6 (5.5-15)	19% (15-23%)	24% (19-29%)	160 (120-210)	7500 (5800-9600)
Bosnia and Herzegovina	1990	Male	50 to 54	50% (36-63%)	50% (37-64%)	10 (6-15)	18% (14-22%)	22% (17-28%)	310 (230-410)	12000 (9300-16000)
Bosnia and Herzegovina	1990	Male	55 to 59	48% (34-60%)	52% (40-66%)	10 (5.7-16)	16% (12-20%)	19% (14-24%)	370 (270-500)	13000 (9300-17000)
Bosnia and Herzegovina	1990	Male	60 to 64	46% (32-60%)	54% (40-68%)	9.6 (5.1-16)	14% (10-18%)	16% (12-21%)	370 (260-500)	11000 (7700-14000)
Bosnia and Herzegovina	1990	Male	65 to 69	44% (32-56%)	56% (44-68%)	8.5 (4.9-13)	14% (9.5-18%)	15% (10-20%)	290 (200-400)	6800 (4700-9100)
Bosnia and Herzegovina	1990	Male	70 to 74	41% (30-53%)	59% (47-70%)	7 (3.7-11)	12% (8.4-17%)	14% (9.6-19%)	180 (120-250)	3400 (2300-4700)
Bosnia and Herzegovina	1990	Male	75 to 79	36% (25-48%)	64% (52-75%)	5.5 (2.7-9.8)	10% (6.6-14%)	12% (7.3-16%)	210 (130-300)	3000 (1900-4300)
Bosnia and Herzegovina	1990	Male	80 to 84	34% (24-46%)	66% (54-76%)	4.7 (2.5-7.9)	9.2% (5.5-14%)	10% (6.1-16%)	170 (100-270)	1800 (1100-2800)
Bosnia and Herzegovina	1990	Male	85 to 89	34% (23-46%)	66% (54-77%)	4.2 (2.2-7.1)	8.5% (4.8-13%)	9.6% (5.3-15%)	110 (58-170)	810 (450-1300)
Bosnia and Herzegovina	1990	Male	90 to 94	35% (24-47%)	65% (53-76%)	3.9 (2-6.9)	7.8% (4.1-12%)	8.7% (4.4-14%)	34 (17-54)	180 (95-280)
Bosnia and Herzegovina	1990	Male	95 plus	37% (24-52%)	63% (48-76%)	3.6 (1.5-6.9)	7% (2.6-12%)	7.7% (2.6-14%)	6.2 (2.2-11)	22 (8.4-37)
Bosnia and Herzegovina	1995	Female	All Ages	13% (11-15%)	87% (85-89%)	1.9 (1.4-2.4)	4.5% (3.4-5.9%)	6.1% (4.1-8.7%)	1000 (650-1500)	28000 (20000-37000)
Bosnia and Herzegovina	1995	Female	15 to 19	12% (7.7-17%)	88% (83-92%)	1.3 (0.66-2.2)	2.1% (1.4-2.9%)	2% (1.1-3.4%)	3.6 (1.9-6)	480 (310-690)
Bosnia and Herzegovina	1995	Female	20 to 24	16% (9.7-23%)	84% (77-90%)	2 (0.96-3.6)	3.7% (2.6-4.9%)	3% (1.7-4.5%)	5.8 (3.4-8.9)	930 (630-1300)
Bosnia and Herzegovina	1995	Female	25 to 29	16% (10-24%)	84% (76-90%)	2 (1-3.5)	4.1% (3.1-5.2%)	3.8% (2.3-5.7%)	6.1 (3.6-9.3)	910 (640-1200)
Bosnia and Herzegovina	1995	Female	30 to 34	17% (10-25%)	83% (75-90%)	2.2 (1.2-3.9)	4.2% (3.1-5.4%)	4.9% (3.2-7.1%)	11 (6.9-16)	1200 (850-1600)
Bosnia and Herzegovina	1995	Female	35 to 39	17% (11-24%)	83% (76-89%)	2.6 (1.4-4.3)	4.5% (3.4-6.1%)	6.3% (4.2-8.9%)	17 (11-25)	1400 (960-1800)
Bosnia and Herzegovina	1995	Female	40 to 44	17% (11-24%)	83% (76-89%)	2.9 (1.6-4.5)	5.2% (3.8-7.1%)	8.1% (5.6-12%)	28 (18-42)	1800 (1200-2400)
Bosnia and Herzegovina	1995	Female	45 to 49	16% (9.6-23%)	84% (77-90%)	3.2 (1.7-5.5)	6.1% (4.4-8.5%)	9.4% (6.4-13%)	47 (30-70)	2300 (1600-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	1995	Female	50 to 54	15% (9.9-22%)	85% (78-90%)	3.2 (1.7-5.3)	6.3% (4.4-8.8%)	9.2% (6.3-13%)	45 (29-66)	1900 (1300-2700)
Bosnia and Herzegovina	1995	Female	55 to 59	14% (9-20%)	86% (80-91%)	3 (1.6-4.8)	6.3% (4.2-9%)	8.8% (5.8-13%)	87 (55-130)	3100 (2100-4500)
Bosnia and Herzegovina	1995	Female	60 to 64	13% (8.1-19%)	87% (81-92%)	2.7 (1.4-4.5)	6.3% (4.2-9.1%)	8.3% (5.4-12%)	130 (83-190)	3800 (2500-5600)
Bosnia and Herzegovina	1995	Female	65 to 69	12% (7.9-17%)	88% (83-92%)	2.4 (1.3-3.8)	6.2% (4-9.2%)	7.9% (4.9-12%)	170 (100-260)	4000 (2500-6000)
Bosnia and Herzegovina	1995	Female	70 to 74	10% (6.9-15%)	90% (85-93%)	2 (1-3.4)	5.5% (3.3-8.5%)	6.8% (4-11%)	150 (91-250)	2900 (1800-4500)
Bosnia and Herzegovina	1995	Female	75 to 79	8.9% (5.5-13%)	91% (87-94%)	1.6 (0.77-2.9)	4.6% (2.6-7.5%)	5.5% (3.1-9.1%)	96 (52-160)	1400 (780-2300)
Bosnia and Herzegovina	1995	Female	80 to 84	7.9% (4.7-11%)	92% (89-95%)	1.4 (0.68-2.4)	4.1% (2.1-6.7%)	4.8% (2.4-8.1%)	89 (43-150)	950 (470-1600)
Bosnia and Herzegovina	1995	Female	85 to 89	7.7% (4.4-12%)	92% (88-96%)	1.4 (0.65-2.6)	3.8% (1.8-6.6%)	4.5% (2.1-8.1%)	77 (35-140)	580 (280-1000)
Bosnia and Herzegovina	1995	Female	90 to 94	8.3% (5.1-13%)	92% (87-95%)	1.5 (0.7-2.9)	3.7% (1.8-6.8%)	4.4% (2-8.2%)	36 (16-68)	190 (90-350)
Bosnia and Herzegovina	1995	Female	95 plus	9.6% (5.4-15%)	90% (85-95%)	1.7 (0.69-3.5)	3.9% (1.7-6.9%)	4.5% (1.9-8.4%)	12 (4.8-22)	40 (18-73)
Bosnia and Herzegovina	1995	Male	All Ages	38% (33-43%)	62% (57-67%)	4.9 (3.8-6)	9.8% (7.8-12%)	13% (9.5-16%)	2600 (1900-3400)	85000 (66000-110000)
Bosnia and Herzegovina	1995	Male	15 to 19	34% (23-49%)	66% (51-77%)	2.2 (1.1-4)	3.2% (1.9-4.9%)	3% (1.5-5.2%)	14 (6.5-23)	1300 (770-2100)
Bosnia and Herzegovina	1995	Male	20 to 24	44% (30-57%)	56% (43-70%)	4.4 (2.2-7.3)	4.9% (3.2-6.6%)	4.2% (2.5-6.3%)	38 (23-57)	3400 (2300-4700)
Bosnia and Herzegovina	1995	Male	25 to 29	41% (28-53%)	59% (47-72%)	5 (2.5-8)	6.2% (4.5-7.9%)	5.7% (3.8-7.8%)	43 (28-59)	3600 (2600-4700)
Bosnia and Herzegovina	1995	Male	30 to 34	42% (29-54%)	58% (46-71%)	5.5 (3.1-8.9)	8% (6.2-10%)	8.2% (5.9-11%)	65 (46-85)	4800 (3600-6100)
Bosnia and Herzegovina	1995	Male	35 to 39	48% (35-62%)	52% (38-65%)	6.4 (3.7-9.9)	11% (8.6-14%)	12% (9.4-16%)	93 (67-120)	5900 (4600-7600)
Bosnia and Herzegovina	1995	Male	40 to 44	52% (38-66%)	48% (34-62%)	7.6 (4.5-11)	14% (11-17%)	17% (13-21%)	140 (100-180)	7500 (5800-9700)
Bosnia and Herzegovina	1995	Male	45 to 49	52% (36-65%)	48% (35-64%)	8.7 (5.1-13)	16% (13-20%)	20% (15-24%)	200 (140-260)	9200 (7000-12000)
Bosnia and Herzegovina	1995	Male	50 to 54	50% (36-63%)	50% (37-64%)	9.2 (5.6-14)	16% (13-21%)	20% (15-25%)	180 (130-250)	7400 (5500-9800)
Bosnia and Herzegovina	1995	Male	55 to 59	47% (34-60%)	53% (40-66%)	9.3 (5.5-14)	15% (11-19%)	17% (13-22%)	320 (220-440)	11000 (7800-15000)
Bosnia and Herzegovina	1995	Male	60 to 64	46% (32-59%)	54% (41-68%)	8.8 (4.8-14)	13% (9.8-18%)	15% (11-20%)	380 (260-520)	11000 (7800-15000)
Bosnia and Herzegovina	1995	Male	65 to 69	44% (32-56%)	56% (44-68%)	7.7 (4.6-12)	13% (9-17%)	15% (10-20%)	440 (290-600)	10000 (6900-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	1995	Male	70 to 74	41% (29-53%)	59% (47-71%)	6.4 (3.6-10)	12% (8.2-16%)	14% (9.2-19%)	270 (170-370)	5000 (3300-7000)
Bosnia and Herzegovina	1995	Male	75 to 79	37% (25-50%)	63% (50-75%)	5 (2.7-8.7)	10% (6.4-14%)	11% (7.2-16%)	130 (78-190)	1900 (1200-2700)
Bosnia and Herzegovina	1995	Male	80 to 84	34% (24-47%)	66% (53-76%)	4.3 (2.3-7.1)	8.9% (5.3-13%)	10% (6-15%)	120 (71-190)	1300 (790-2000)
Bosnia and Herzegovina	1995	Male	85 to 89	34% (23-46%)	66% (54-77%)	3.9 (2.1-6.4)	8.1% (4.5-12%)	9.2% (4.9-14%)	110 (57-170)	810 (450-1300)
Bosnia and Herzegovina	1995	Male	90 to 94	35% (24-47%)	65% (53-76%)	3.5 (1.8-6.2)	7.3% (3.8-12%)	8.2% (4.1-14%)	44 (22-72)	230 (120-370)
Bosnia and Herzegovina	1995	Male	95 plus	37% (24-52%)	63% (48-76%)	3.3 (1.4-6.1)	6.4% (2.1-11%)	7.1% (2.2-13%)	8.7 (2.5-16)	31 (10-55)
Bosnia and Herzegovina	2000	Female	All Ages	11% (9.7-13%)	89% (87-90%)	1.9 (1.4-2.4)	4.6% (3.4-6.1%)	6% (3.9-8.5%)	840 (550-1200)	23000 (17000-31000)
Bosnia and Herzegovina	2000	Female	15 to 19	11% (7.1-15%)	89% (85-93%)	1.2 (0.63-2.2)	2.9% (2-4%)	6.3% (3.6-9.9%)	2.4 (1.3-4)	400 (260-560)
Bosnia and Herzegovina	2000	Female	20 to 24	13% (8.3-20%)	87% (80-92%)	1.8 (0.88-3.4)	5% (3.7-6.6%)	7.4% (4.5-11%)	3.4 (2.5-1)	680 (460-940)
Bosnia and Herzegovina	2000	Female	25 to 29	14% (8.6-20%)	86% (80-91%)	1.8 (0.89-3.3)	4.9% (3.7-6.2%)	7.1% (4.4-10%)	3.6 (2.2-5.5)	730 (510-990)
Bosnia and Herzegovina	2000	Female	30 to 34	14% (9.1-22%)	86% (78-91%)	2.1 (1.1-3.7)	4.7% (3.6-6.1%)	8.1% (5.5-12%)	5.7 (3.7-8.6)	780 (560-1000)
Bosnia and Herzegovina	2000	Female	35 to 39	15% (9.4-22%)	85% (78-91%)	2.5 (1.3-4.1)	4.7% (3.6-6.2%)	8.7% (5.9-12%)	11 (7.1-17)	1100 (760-1400)
Bosnia and Herzegovina	2000	Female	40 to 44	16% (11-22%)	84% (78-89%)	3 (1.7-4.7)	5.1% (3.9-6.9%)	9.8% (6.7-14%)	20 (13-29)	1400 (960-1900)
Bosnia and Herzegovina	2000	Female	45 to 49	15% (9.1-20%)	85% (80-91%)	3.1 (1.8-5.2)	5.6% (4.1-7.7%)	9.8% (6.8-14%)	35 (23-51)	1900 (1300-2600)
Bosnia and Herzegovina	2000	Female	50 to 54	13% (8.6-19%)	87% (81-91%)	3 (1.6-4.8)	5.6% (3.9-7.7%)	9% (6.1-13%)	43 (28-64)	1900 (1400-2700)
Bosnia and Herzegovina	2000	Female	55 to 59	12% (8-18%)	88% (82-92%)	2.7 (1.5-4.5)	5.4% (3.7-7.7%)	8.3% (5.6-12%)	41 (27-61)	1500 (1100-2200)
Bosnia and Herzegovina	2000	Female	60 to 64	11% (7.3-16%)	89% (84-93%)	2.5 (1.3-4.2)	5.4% (3.7-7.8%)	7.5% (5-11%)	96 (61-140)	2900 (1900-4100)
Bosnia and Herzegovina	2000	Female	65 to 69	10% (6.5-14%)	90% (86-93%)	2.1 (1.2-3.4)	5.3% (3.4-7.8%)	7% (4.4-10%)	140 (86-210)	3300 (2100-5000)
Bosnia and Herzegovina	2000	Female	70 to 74	9% (5.9-13%)	91% (87-94%)	1.8 (0.95-3.1)	4.8% (2.9-7.5%)	6.1% (3.6-9.5%)	150 (89-240)	2900 (1700-4500)
Bosnia and Herzegovina	2000	Female	75 to 79	8.2% (5.1-12%)	92% (88-95%)	1.6 (0.75-2.8)	4.2% (2.4-6.8%)	5.1% (2.9-8.4%)	130 (71-210)	1900 (1100-3100)
Bosnia and Herzegovina	2000	Female	80 to 84	7.5% (4.5-11%)	93% (89-96%)	1.4 (0.7-2.5)	3.8% (1.9-6.2%)	4.6% (2.3-7.8%)	82 (40-140)	890 (440-1500)
Bosnia and Herzegovina	2000	Female	85 to 89	7.2% (4.2-11%)	93% (89-96%)	1.4 (0.65-2.5)	3.3% (1.7-5.8%)	4.2% (2-7.4%)	46 (21-81)	350 (170-620)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	2000	Female	90 to 94	7.8% (4.9-12%)	92% (88-95%)	1.5 (0.7-2.9)	3.2% (1.5-5.9%)	4% (1.8-7.5%)	31 (14-57)	160 (79-300)
Bosnia and Herzegovina	2000	Female	95 plus	9% (5.1-14%)	91% (86-95%)	1.7 (0.66-3.5)	3.4% (1.5-6.2%)	4.2% (1.7-7.7%)	11 (4.4-20)	39 (17-71)
Bosnia and Herzegovina	2000	Male	All Ages	40% (35-44%)	60% (56-65%)	5.7 (4.5-6.9)	13% (10-16%)	15% (11-19%)	2400 (1700-3100)	76000 (58000-96000)
Bosnia and Herzegovina	2000	Male	15 to 19	36% (26-49%)	64% (51-74%)	2.5 (1.3-4.3)	7.3% (4.6-11%)	13% (6.8-20%)	12 (6.4-21)	1300 (810-2000)
Bosnia and Herzegovina	2000	Male	20 to 24	44% (32-56%)	56% (44-68%)	4.7 (2.6-7.9)	14% (10-17%)	21% (14-28%)	26 (16-37)	2600 (1800-3400)
Bosnia and Herzegovina	2000	Male	25 to 29	37% (27-48%)	63% (52-73%)	4.8 (2.6-7.8)	13% (10-16%)	20% (15-27%)	31 (21-43)	2900 (2100-3700)
Bosnia and Herzegovina	2000	Male	30 to 34	38% (26-50%)	62% (50-74%)	5.4 (3-8.5)	13% (11-17%)	22% (17-28%)	37 (26-50)	3100 (2300-3900)
Bosnia and Herzegovina	2000	Male	35 to 39	49% (38-61%)	51% (39-62%)	7 (4.1-10)	16% (13-20%)	26% (20-31%)	72 (52-97)	4900 (3800-6300)
Bosnia and Herzegovina	2000	Male	40 to 44	57% (45-69%)	43% (31-55%)	9.1 (5.8-13)	18% (15-22%)	27% (22-33%)	110 (83-150)	6500 (5100-8300)
Bosnia and Herzegovina	2000	Male	45 to 49	58% (44-70%)	42% (30-56%)	11 (6.6-16)	18% (15-22%)	26% (20-31%)	180 (130-240)	8700 (6700-11000)
Bosnia and Herzegovina	2000	Male	50 to 54	53% (39-64%)	47% (36-61%)	11 (6.4-15)	17% (14-22%)	22% (17-28%)	220 (160-290)	9100 (6900-12000)
Bosnia and Herzegovina	2000	Male	55 to 59	47% (35-59%)	53% (41-65%)	10 (6-15)	15% (11-19%)	19% (14-24%)	170 (120-230)	6100 (4400-8000)
Bosnia and Herzegovina	2000	Male	60 to 64	44% (31-57%)	56% (43-69%)	9 (5-14)	13% (9.8-17%)	16% (11-20%)	330 (230-450)	9600 (6900-13000)
Bosnia and Herzegovina	2000	Male	65 to 69	41% (31-52%)	59% (48-69%)	7.6 (4.6-11)	12% (8.5-16%)	14% (9.9-19%)	390 (260-540)	9400 (6400-13000)
Bosnia and Herzegovina	2000	Male	70 to 74	38% (28-50%)	62% (50-72%)	6.4 (3.6-10)	11% (7.7-15%)	13% (8.8-18%)	380 (240-540)	7300 (4800-10000)
Bosnia and Herzegovina	2000	Male	75 to 79	35% (25-47%)	65% (53-75%)	5.2 (2.6-8.8)	9.6% (6-13%)	11% (6.9-16%)	180 (110-270)	2700 (1700-4000)
Bosnia and Herzegovina	2000	Male	80 to 84	33% (23-45%)	67% (55-77%)	4.4 (2.4-7.3)	8.6% (5.2-13%)	9.9% (5.7-15%)	98 (56-150)	1100 (650-1700)
Bosnia and Herzegovina	2000	Male	85 to 89	33% (22-45%)	67% (55-78%)	4 (2.1-6.5)	7.7% (4.3-12%)	8.9% (4.8-14%)	57 (30-92)	450 (250-710)
Bosnia and Herzegovina	2000	Male	90 to 94	34% (23-46%)	66% (54-77%)	3.7 (1.9-6.3)	7.2% (3.9-12%)	8.2% (4.3-13%)	39 (20-65)	210 (120-340)
Bosnia and Herzegovina	2000	Male	95 plus	36% (23-51%)	64% (49-77%)	3.4 (1.5-6.6)	6.5% (2.5-11%)	7.3% (2.4-13%)	10 (3.5-18)	38 (15-65)
Bosnia and Herzegovina	2005	Female	All Ages	10% (8.9-12%)	90% (88-91%)	1.9 (1.4-2.5)	4.1% (3-5.4%)	5.2% (3.4-7.4%)	820 (530-1200)	21000 (16000-28000)
Bosnia and Herzegovina	2005	Female	15 to 19	9.8% (6.3-14%)	90% (86-94%)	1.2 (0.63-2.2)	2.6% (1.8-3.5%)	5.4% (3.2-8.5%)	1.6 (0.94-2.5)	320 (210-450)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	2005	Female	20 to 24	12% (7.7-18%)	88% (82-92%)	1.8 (0.86-3.3)	4.7% (3.4-6.1%)	6.2% (3.8-9.3%)	2.8 (1.7-4.3)	700 (470-980)
Bosnia and Herzegovina	2005	Female	25 to 29	13% (7.9-19%)	87% (81-92%)	1.8 (0.89-3.3)	4.6% (3.5-5.9%)	6.1% (3.9-8.6%)	2.6 (1.6-3.9)	650 (450-890)
Bosnia and Herzegovina	2005	Female	30 to 34	13% (8-20%)	87% (80-92%)	2 (1-3.6)	4.4% (3.3-5.7%)	6.9% (4.5-10%)	4.7 (3-7)	760 (540-1000)
Bosnia and Herzegovina	2005	Female	35 to 39	14% (8.5-20%)	86% (80-91%)	2.4 (1.2-4.2)	4.3% (3.3-5.7%)	7.5% (4.9-11%)	7.1 (4.5-11)	810 (570-1100)
Bosnia and Herzegovina	2005	Female	40 to 44	14% (9.8-20%)	86% (80-90%)	2.9 (1.6-4.7)	4.6% (3.4-6.2%)	8.4% (5.6-12%)	16 (9.9-23)	1200 (870-1700)
Bosnia and Herzegovina	2005	Female	45 to 49	13% (8.4-19%)	87% (81-92%)	3.1 (1.7-5.3)	4.8% (3.5-6.7%)	8.3% (5.6-12%)	28 (18-42)	1600 (1100-2200)
Bosnia and Herzegovina	2005	Female	50 to 54	12% (7.9-17%)	88% (83-92%)	2.9 (1.6-4.9)	4.7% (3.3-6.5%)	7.6% (5.1-11%)	38 (24-57)	1800 (1200-2600)
Bosnia and Herzegovina	2005	Female	55 to 59	11% (7.2-16%)	89% (84-93%)	2.7 (1.5-4.3)	4.6% (3.1-6.4%)	7.1% (4.7-10%)	46 (30-68)	1800 (1200-2600)
Bosnia and Herzegovina	2005	Female	60 to 64	10% (6.5-15%)	90% (85-93%)	2.4 (1.2-4.1)	4.7% (3.1-6.8%)	6.5% (4.2-9.5%)	57 (37-84)	1800 (1200-2500)
Bosnia and Herzegovina	2005	Female	65 to 69	9.2% (5.9-13%)	91% (87-94%)	2.1 (1.1-3.5)	4.7% (3-6.9%)	6.1% (3.8-9.1%)	120 (75-190)	3000 (1900-4600)
Bosnia and Herzegovina	2005	Female	70 to 74	8.2% (5.4-12%)	92% (88-95%)	1.8 (0.92-3.1)	4.3% (2.6-6.6%)	5.4% (3.3-8.3%)	160 (94-250)	3000 (1800-4700)
Bosnia and Herzegovina	2005	Female	75 to 79	7.5% (4.6-11%)	93% (89-95%)	1.6 (0.73-2.7)	3.7% (2.2-6%)	4.6% (2.6-7.3%)	150 (87-250)	2200 (1300-3600)
Bosnia and Herzegovina	2005	Female	80 to 84	6.8% (4.2-9.6%)	93% (90-96%)	1.4 (0.69-2.4)	3.4% (1.8-5.5%)	4.1% (2.1-6.7%)	110 (57-180)	1200 (630-1900)
Bosnia and Herzegovina	2005	Female	85 to 89	6.6% (3.8-10%)	93% (90-96%)	1.4 (0.62-2.5)	2.9% (1.4-4.9%)	3.7% (1.8-6.4%)	43 (20-75)	340 (170-570)
Bosnia and Herzegovina	2005	Female	90 to 94	7.1% (4.3-11%)	93% (89-96%)	1.5 (0.66-2.8)	2.8% (1.4-4.9%)	3.6% (1.6-6.5%)	21 (9.7-37)	110 (56-200)
Bosnia and Herzegovina	2005	Female	95 plus	8.2% (4.7-13%)	92% (87-95%)	1.7 (0.63-3.5)	3% (1.4-5.3%)	3.8% (1.6-7%)	11 (4.5-20)	41 (18-72)
Bosnia and Herzegovina	2005	Male	All Ages	38% (34-43%)	62% (57-66%)	6 (4.7-7.6)	12% (9.1-14%)	13% (9.6-17%)	2300 (1600-3000)	72000 (55000-90000)
Bosnia and Herzegovina	2005	Male	15 to 19	34% (24-46%)	66% (54-76%)	2.5 (1.3-4.4)	6.8% (4.4-9.9%)	12% (6.4-18%)	9.1 (5-15)	1100 (700-1500)
Bosnia and Herzegovina	2005	Male	20 to 24	42% (30-54%)	58% (46-70%)	4.7 (2.5-8)	13% (9.6-16%)	19% (13-26%)	22 (14-31)	2500 (1800-3300)
Bosnia and Herzegovina	2005	Male	25 to 29	34% (25-45%)	66% (55-75%)	4.8 (2.6-8)	12% (9.6-15%)	18% (13-24%)	23 (16-32)	2400 (1800-3100)
Bosnia and Herzegovina	2005	Male	30 to 34	35% (25-48%)	65% (52-75%)	5.4 (2.9-8.9)	12% (9.2-14%)	20% (15-25%)	28 (20-37)	2600 (2000-3200)
Bosnia and Herzegovina	2005	Male	35 to 39	47% (35-59%)	53% (41-65%)	7.1 (3.9-11)	14% (11-17%)	23% (18-29%)	46 (34-62)	3400 (2600-4300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	2005	Male	40 to 44	55% (43-67%)	45% (33-57%)	9.3 (5.6-13)	16% (13-20%)	25% (20-31%)	95 (71-130)	5800 (4500-7200)
Bosnia and Herzegovina	2005	Male	45 to 49	56% (42-67%)	44% (33-58%)	11 (6.7-16)	17% (13-20%)	23% (18-29%)	150 (110-200)	7700 (5800-9900)
Bosnia and Herzegovina	2005	Male	50 to 54	50% (37-61%)	50% (39-63%)	11 (6.6-15)	15% (12-19%)	20% (15-25%)	210 (150-280)	9100 (6800-12000)
Bosnia and Herzegovina	2005	Male	55 to 59	44% (33-56%)	56% (44-67%)	10 (6-15)	13% (9.9-17%)	17% (12-21%)	210 (150-290)	7800 (5700-10000)
Bosnia and Herzegovina	2005	Male	60 to 64	41% (30-54%)	59% (46-70%)	9 (4.9-15)	12% (8.8-16%)	14% (10-19%)	210 (140-270)	6100 (4300-8100)
Bosnia and Herzegovina	2005	Male	65 to 69	39% (29-49%)	61% (51-71%)	7.7 (4.6-12)	11% (7.7-15%)	13% (9.1-17%)	370 (250-510)	9100 (6100-12000)
Bosnia and Herzegovina	2005	Male	70 to 74	36% (26-47%)	64% (53-74%)	6.4 (3.7-10)	10% (6.9-14%)	12% (7.8-16%)	390 (250-560)	7600 (4900-11000)
Bosnia and Herzegovina	2005	Male	75 to 79	32% (23-44%)	68% (56-77%)	5.1 (2.7-8.6)	8.7% (5.6-12%)	10% (6.3-14%)	290 (180-430)	4400 (2800-6300)
Bosnia and Herzegovina	2005	Male	80 to 84	31% (21-42%)	69% (58-79%)	4.3 (2.4-7.2)	7.9% (4.7-12%)	9.2% (5.4-14%)	150 (84-220)	1600 (970-2500)
Bosnia and Herzegovina	2005	Male	85 to 89	31% (21-42%)	69% (58-79%)	3.9 (2.1-6.6)	7.2% (4.1-11%)	8.3% (4.6-13%)	44 (24-70)	360 (200-550)
Bosnia and Herzegovina	2005	Male	90 to 94	32% (21-44%)	68% (56-79%)	3.7 (1.9-6.5)	6.8% (3.8-11%)	7.8% (4.1-13%)	22 (12-36)	120 (67-190)
Bosnia and Herzegovina	2005	Male	95 plus	33% (21-48%)	67% (52-79%)	3.4 (1.4-6.8)	6.3% (2.8-10%)	7% (2.9-12%)	9 (3.7-15)	34 (15-56)
Bosnia and Herzegovina	2010	Female	All Ages	11% (9.4-13%)	89% (87-91%)	2.1 (1.6-2.8)	4.1% (3.3-4.4%)	5% (3.3-7.2%)	850 (550-1300)	22000 (16000-29000)
Bosnia and Herzegovina	2010	Female	15 to 19	9.9% (6.3-15%)	90% (85-94%)	1.3 (0.64-2.4)	2.7% (1.9-3.6%)	5.2% (3.1-8.1%)	1.1 (0.69-1.8)	260 (170-370)
Bosnia and Herzegovina	2010	Female	20 to 24	13% (8.1-19%)	87% (81-92%)	2 (0.95-3.7)	5.1% (3.6-6.7%)	6.2% (3.7-9.2%)	2.3 (1.3-3.5)	670 (450-970)
Bosnia and Herzegovina	2010	Female	25 to 29	14% (8.6-20%)	86% (80-91%)	2.1 (1-3.8)	4.9% (3.7-6.2%)	6.2% (4-8.9%)	2.7 (1.6-4)	770 (540-1100)
Bosnia and Herzegovina	2010	Female	30 to 34	14% (8.7-22%)	86% (78-91%)	2.3 (1.1-4.2)	4.5% (3.4-5.8%)	7.1% (4.6-10%)	4.2 (2.6-6.5)	740 (510-1000)
Bosnia and Herzegovina	2010	Female	35 to 39	15% (9.1-22%)	85% (78-91%)	2.7 (1.3-4.7)	4.3% (3.3-5.7%)	7.4% (4.8-11%)	6.9 (4.2-10)	840 (600-1100)
Bosnia and Herzegovina	2010	Female	40 to 44	14% (9.1-21%)	86% (79-91%)	3 (1.6-5.1)	4.4% (3.2-6%)	7.9% (5.1-12%)	12 (7.1-18)	990 (690-1400)
Bosnia and Herzegovina	2010	Female	45 to 49	14% (8.1-20%)	86% (80-92%)	3.3 (1.7-5.8)	4.7% (3.3-6.4%)	8% (5.2-12%)	25 (16-38)	1500 (1100-2100)
Bosnia and Herzegovina	2010	Female	50 to 54	13% (8.3-19%)	87% (81-92%)	3.3 (1.8-5.6)	4.7% (3.3-6.5%)	7.6% (5-11%)	37 (23-56)	1800 (1200-2600)
Bosnia and Herzegovina	2010	Female	55 to 59	12% (7.7-17%)	88% (83-92%)	3 (1.6-5.1)	4.6% (3.1-6.5%)	7.3% (4.8-11%)	51 (32-75)	2000 (1400-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	2010	Female	60 to 64	11% (6.8-17%)	89% (83-93%)	2.8 (1.3-4.7)	4.6% (3-6.7%)	6.6% (4.2-9.8%)	73 (45-110)	2300 (1500-3400)
Bosnia and Herzegovina	2010	Female	65 to 69	10% (6.5-15%)	90% (85-93%)	2.4 (1.3-4.1)	4.7% (2.9-7%)	6.4% (3.9-9.6%)	86 (52-130)	2100 (1300-3200)
Bosnia and Herzegovina	2010	Female	70 to 74	8.8% (5.7-13%)	91% (87-94%)	2 (1-3.6)	4.3% (2.6-6.7%)	5.6% (3.4-8.6%)	150 (93-240)	3000 (1800-4700)
Bosnia and Herzegovina	2010	Female	75 to 79	7.6% (4.7-12%)	92% (88-95%)	1.7 (0.76-3)	3.7% (2.1-5.9%)	4.6% (2.6-7.3%)	160 (95-270)	2400 (1400-4000)
Bosnia and Herzegovina	2010	Female	80 to 84	6.7% (4.1-9.6%)	93% (90-96%)	1.4 (0.69-2.5)	3.3% (1.8-5.3%)	4% (2.1-6.5%)	130 (68-210)	1400 (760-2300)
Bosnia and Herzegovina	2010	Female	85 to 89	6.5% (3.7-9.9%)	94% (90-96%)	1.4 (0.66-2.6)	2.8% (1.4-4.8%)	3.6% (1.8-6.2%)	68 (32-120)	530 (270-900)
Bosnia and Herzegovina	2010	Female	90 to 94	7% (4.2-11%)	93% (89-96%)	1.5 (0.68-2.9)	2.7% (1.3-4.8%)	3.5% (1.6-6.4%)	26 (12-49)	140 (73-260)
Bosnia and Herzegovina	2010	Female	95 plus	8% (4.5-13%)	92% (87-96%)	1.7 (0.62-3.7)	2.9% (1.3-5.2%)	3.7% (1.5-6.8%)	10 (4.3-19)	37 (17-65)
Bosnia and Herzegovina	2010	Male	All Ages	36% (31-40%)	64% (60-69%)	6.2 (4.7-8)	11% (8.4-13%)	12% (8.9-16%)	2200 (1600-2900)	68000 (52000-84000)
Bosnia and Herzegovina	2010	Male	15 to 19	30% (20-42%)	70% (58-80%)	2.3 (1.1-4.2)	6.3% (4.2-8.9%)	11% (5.5-17%)	6.1 (3.1-10)	770 (500-1100)
Bosnia and Herzegovina	2010	Male	20 to 24	39% (25-52%)	61% (48-75%)	4.7 (2.3-7.9)	12% (9.2-16%)	18% (12-25%)	18 (12-27)	2200 (1600-3000)
Bosnia and Herzegovina	2010	Male	25 to 29	36% (24-48%)	64% (52-76%)	5.3 (2.8-8.9)	12% (9.2-15%)	18% (13-25%)	24 (16-33)	2600 (1900-3300)
Bosnia and Herzegovina	2010	Male	30 to 34	37% (25-50%)	63% (50-75%)	6 (3.1-9.9)	12% (9.1-14%)	20% (14-26%)	25 (18-34)	2400 (1800-3000)
Bosnia and Herzegovina	2010	Male	35 to 39	44% (30-57%)	56% (43-70%)	7.1 (3.8-11)	13% (10-17%)	22% (16-29%)	42 (30-57)	3200 (2400-4100)
Bosnia and Herzegovina	2010	Male	40 to 44	47% (33-62%)	53% (38-67%)	8.4 (4.9-13)	14% (11-18%)	22% (17-29%)	67 (49-91)	4200 (3200-5500)
Bosnia and Herzegovina	2010	Male	45 to 49	47% (31-61%)	53% (39-69%)	9.6 (5.5-15)	14% (11-18%)	20% (15-26%)	130 (92-170)	6600 (5000-8500)
Bosnia and Herzegovina	2010	Male	50 to 54	45% (31-59%)	55% (41-69%)	10 (6.1-15)	14% (11-18%)	18% (14-23%)	190 (130-250)	8100 (5900-11000)
Bosnia and Herzegovina	2010	Male	55 to 59	42% (29-56%)	58% (44-71%)	10 (5.8-16)	12% (9.1-16%)	16% (12-20%)	220 (160-300)	8200 (6000-11000)
Bosnia and Herzegovina	2010	Male	60 to 64	41% (27-55%)	59% (45-73%)	9.5 (5-15)	11% (8.2-15%)	14% (9.9-18%)	270 (180-360)	8100 (5700-11000)
Bosnia and Herzegovina	2010	Male	65 to 69	40% (27-53%)	60% (47-73%)	8.4 (4.8-13)	11% (7.5-15%)	13% (8.7-17%)	240 (160-340)	6000 (4000-8200)
Bosnia and Herzegovina	2010	Male	70 to 74	36% (26-48%)	64% (52-74%)	6.9 (3.8-11)	10% (6.6-14%)	12% (7.6-16%)	380 (250-540)	7500 (4900-11000)
Bosnia and Herzegovina	2010	Male	75 to 79	32% (22-45%)	68% (55-78%)	5.3 (2.7-9.2)	8.3% (5.3-12%)	9.6% (6.1-14%)	310 (190-450)	4800 (3000-6800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	2010	Male	80 to 84	29% (20-41%)	71% (59-80%)	4.4 (2.3-7.3)	7.6% (4.7-11%)	8.8% (5.3-13%)	200 (120-300)	2200 (1300-3300)
Bosnia and Herzegovina	2010	Male	85 to 89	29% (19-41%)	71% (59-81%)	4 (2.1-6.7)	6.8% (3.9-10%)	8% (4.4-13%)	85 (46-130)	690 (390-1100)
Bosnia and Herzegovina	2010	Male	90 to 94	31% (20-42%)	69% (58-80%)	3.8 (1.9-6.6)	6.6% (3.8-10%)	7.6% (4.1-12%)	25 (13-39)	130 (76-210)
Bosnia and Herzegovina	2010	Male	95 plus	33% (20-48%)	67% (52-80%)	3.5 (1.4-7)	6.2% (2.8-10%)	7% (2.9-12%)	6.5 (2.7-11)	24 (11-40)
Bosnia and Herzegovina	2016	Female	All Ages	11% (9-12%)	89% (88-91%)	2.1 (1.4-3)	3.9% (2.9-5.1%)	4.7% (3.1-6.7%)	860 (540-1300)	21000 (15000-29000)
Bosnia and Herzegovina	2016	Female	15 to 19	9.7% (6.2-15%)	90% (85-94%)	1.3 (0.59-2.4)	2.7% (1.9-3.6%)	5.2% (3-8.1%)	1 (0.57-1.7)	230 (160-330)
Bosnia and Herzegovina	2016	Female	20 to 24	13% (8.1-19%)	87% (81-92%)	2 (0.89-3.8)	5.2% (3.7-6.9%)	6.2% (3.8-9.3%)	1.7 (0.97-2.5)	530 (350-760)
Bosnia and Herzegovina	2016	Female	25 to 29	13% (8.4-20%)	87% (80-92%)	2.1 (0.93-3.7)	4.9% (3.7-6.2%)	6.2% (3.9-8.8%)	2.3 (1.3-3.5)	690 (470-940)
Bosnia and Herzegovina	2016	Female	30 to 34	14% (8.3-21%)	86% (79-92%)	2.3 (1.1-4.2)	4.5% (3.3-5.8%)	6.9% (4.4-10%)	4.4 (2.6-6.9)	830 (580-1100)
Bosnia and Herzegovina	2016	Female	35 to 39	14% (8.5-21%)	86% (79-92%)	2.7 (1.3-4.9)	4.3% (3.3-5.7%)	7.2% (4.7-11%)	6.3 (3.7-10)	830 (580-1100)
Bosnia and Herzegovina	2016	Female	40 to 44	14% (8.8-20%)	86% (80-91%)	3 (1.5-5.2)	4.4% (3.3-6%)	7.6% (5-11%)	11 (6.7-18)	1000 (690-1400)
Bosnia and Herzegovina	2016	Female	45 to 49	13% (7.6-19%)	87% (81-92%)	3.3 (1.6-5.7)	4.6% (3.3-6.4%)	7.7% (5.1-11%)	19 (12-30)	1200 (860-1700)
Bosnia and Herzegovina	2016	Female	50 to 54	12% (8-18%)	88% (82-92%)	3.2 (1.6-5.8)	4.5% (3.2-6.3%)	7.3% (4.8-11%)	33 (20-51)	1700 (1200-2400)
Bosnia and Herzegovina	2016	Female	55 to 59	11% (7.2-17%)	89% (83-93%)	2.9 (1.5-5.2)	4.3% (3-6.2%)	6.8% (4.5-10%)	47 (29-71)	1900 (1300-2900)
Bosnia and Herzegovina	2016	Female	60 to 64	11% (6.9-16%)	89% (84-93%)	2.7 (1.3-4.7)	4.3% (2.9-6.2%)	6.3% (4.1-9.3%)	73 (46-110)	2400 (1500-3600)
Bosnia and Herzegovina	2016	Female	65 to 69	9.8% (6.3-14%)	90% (86-94%)	2.4 (1.2-4.1)	4.4% (2.8-6.5%)	6.1% (3.9-9.1%)	110 (68-170)	2800 (1800-4200)
Bosnia and Herzegovina	2016	Female	70 to 74	8.5% (5.5-13%)	91% (87-94%)	2 (0.93-3.6)	4.1% (2.5-6.3%)	5.4% (3.3-8.3%)	100 (62-160)	2000 (1200-3100)
Bosnia and Herzegovina	2016	Female	75 to 79	7.3% (4.4-11%)	93% (89-96%)	1.6 (0.74-3)	3.4% (2-5.4%)	4.3% (2.6-7%)	140 (81-230)	2100 (1200-3400)
Bosnia and Herzegovina	2016	Female	80 to 84	6.4% (3.8-9.4%)	94% (91-96%)	1.4 (0.64-2.5)	3% (1.7-4.9%)	3.8% (2-6.1%)	140 (75-240)	1600 (840-2600)
Bosnia and Herzegovina	2016	Female	85 to 89	6.3% (3.6-9.7%)	94% (90-96%)	1.4 (0.66-2.6)	2.7% (1.4-4.6%)	3.5% (1.7-6%)	94 (43-170)	740 (360-1300)
Bosnia and Herzegovina	2016	Female	90 to 94	6.8% (4.2-11%)	93% (89-96%)	1.5 (0.7-3)	2.7% (1.3-4.7%)	3.4% (1.6-6.1%)	50 (22-89)	270 (130-470)
Bosnia and Herzegovina	2016	Female	95 plus	8% (4.6-13%)	92% (87-95%)	1.7 (0.64-3.7)	2.9% (1.3-5.1%)	3.6% (1.5-6.6%)	18 (7.3-33)	65 (30-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bosnia and Herzegovina	2016	Male	All Ages	36% (31-40%)	64% (60-69%)	6.5 (4.5-9)	11% (8.4-13%)	12% (8.6-15%)	2400 (1700-3200)	70000 (53000-88000)
Bosnia and Herzegovina	2016	Male	15 to 19	30% (19-42%)	70% (58-81%)	2.4 (1.1-4.4)	6.5% (4.2-9.2%)	11% (5.6-17%)	5.5 (2.6-8.9)	710 (460-1000)
Bosnia and Herzegovina	2016	Male	20 to 24	39% (25-52%)	61% (48-75%)	4.8 (2.2-8.1)	13% (9.5-17%)	19% (12-26%)	15 (8.9-21)	1800 (1300-2400)
Bosnia and Herzegovina	2016	Male	25 to 29	35% (24-48%)	65% (52-76%)	5.4 (2.5-9)	13% (9.7-16%)	19% (13-25%)	21 (14-30)	2400 (1700-3100)
Bosnia and Herzegovina	2016	Male	30 to 34	36% (24-50%)	64% (50-76%)	6.1 (3-10)	12% (9.3-15%)	20% (14-25%)	28 (19-39)	2800 (2100-3500)
Bosnia and Herzegovina	2016	Male	35 to 39	43% (30-57%)	57% (43-70%)	7.3 (3.8-12)	13% (10-17%)	22% (17-29%)	41 (28-57)	3200 (2400-4100)
Bosnia and Herzegovina	2016	Male	40 to 44	47% (33-62%)	53% (38-67%)	8.7 (4.7-14)	14% (11-18%)	23% (17-29%)	68 (49-95)	4400 (3300-5700)
Bosnia and Herzegovina	2016	Male	45 to 49	47% (31-60%)	53% (40-69%)	9.9 (5.5-17)	14% (11-18%)	21% (16-27%)	100 (75-140)	5500 (4200-7200)
Bosnia and Herzegovina	2016	Male	50 to 54	45% (31-58%)	55% (42-69%)	10 (6.1-17)	14% (10-17%)	18% (14-23%)	180 (130-250)	7900 (5700-10000)
Bosnia and Herzegovina	2016	Male	55 to 59	42% (29-56%)	58% (44-71%)	10 (5.6-17)	12% (9-15%)	15% (12-20%)	220 (150-300)	8200 (6000-11000)
Bosnia and Herzegovina	2016	Male	60 to 64	40% (27-55%)	60% (45-73%)	9.7 (5-16)	11% (8.1-14%)	14% (9.8-18%)	300 (200-400)	9100 (6500-12000)
Bosnia and Herzegovina	2016	Male	65 to 69	39% (28-52%)	61% (48-72%)	8.6 (4.7-14)	11% (7.3-14%)	13% (8.6-17%)	360 (240-500)	9100 (6200-12000)
Bosnia and Herzegovina	2016	Male	70 to 74	36% (25-47%)	64% (53-75%)	7 (3.7-12)	9.7% (6.5-13%)	11% (7.7-16%)	270 (180-380)	5400 (3600-7600)
Bosnia and Herzegovina	2016	Male	75 to 79	32% (21-44%)	68% (56-79%)	5.4 (2.6-9.4)	8.2% (5.3-12%)	9.5% (6.1-13%)	310 (190-440)	4700 (3000-6800)
Bosnia and Herzegovina	2016	Male	80 to 84	29% (20-41%)	71% (59-80%)	4.5 (2.3-7.8)	7.4% (4.5-11%)	8.6% (5.2-13%)	250 (150-370)	2800 (1700-4200)
Bosnia and Herzegovina	2016	Male	85 to 89	30% (19-41%)	70% (59-81%)	4.1 (2.1-7.1)	6.7% (3.8-10%)	7.8% (4.3-12%)	140 (77-230)	1200 (660-1800)
Bosnia and Herzegovina	2016	Male	90 to 94	31% (21-42%)	69% (58-79%)	3.8 (1.9-7.3)	6.5% (3.7-10%)	7.5% (4.1-12%)	65 (34-100)	360 (200-550)
Bosnia and Herzegovina	2016	Male	95 plus	33% (20-47%)	67% (53-80%)	3.6 (1.4-7.4)	6.1% (2.7-10%)	6.9% (2.8-12%)	11 (4.4-18)	39 (18-65)
Bulgaria	1990	Female	All Ages	76% (70-81%)	24% (19-30%)	2 (1.7-2.3)	4% (1.9-6.5%)	4.3% (0.45-8.2%)	2100 (220-4100)	58000 (28000-89000)
Bulgaria	1990	Female	15 to 19	76% (65-85%)	24% (15-35%)	1.9 (1.3-2.6)	3.4% (2.1-4.9%)	7.3% (3.7-11%)	11 (5.3-16)	1100 (710-1600)
Bulgaria	1990	Female	20 to 24	82% (70-90%)	18% (9.7-30%)	2.3 (1.6-3.3)	4.6% (3.1-6.2%)	7.6% (4-12%)	12 (6-18)	1500 (1000-2100)
Bulgaria	1990	Female	25 to 29	83% (71-91%)	17% (8.7-29%)	2.2 (1.5-3)	4.3% (2.9-5.7%)	6.9% (3.7-10%)	12 (6.5-18)	1600 (1000-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	1990	Female	30 to 34	84% (71-92%)	16% (8.3-29%)	2.3 (1.6-3.1)	4.2% (2.9-5.7%)	7.3% (4.4-10%)	19 (11-27)	1900 (1300-2600)
Bulgaria	1990	Female	35 to 39	84% (72-92%)	16% (8.4-28%)	2.6 (1.8-3.7)	4.6% (3.1-6.4%)	8.5% (5.2-12%)	33 (20-47)	2500 (1700-3400)
Bulgaria	1990	Female	40 to 44	83% (71-91%)	17% (8.7-29%)	2.8 (2-3.9)	5.1% (3.3-7.2%)	9.4% (5.9-13%)	57 (36-81)	3400 (2200-4800)
Bulgaria	1990	Female	45 to 49	82% (67-91%)	18% (9.1-33%)	2.9 (2-3.9)	5.6% (3.4-8%)	9.7% (6.1-14%)	78 (48-110)	3800 (2400-5500)
Bulgaria	1990	Female	50 to 54	80% (66-90%)	20% (10-34%)	2.9 (2-3.8)	5.8% (3.3-8.6%)	9.6% (5.5-14%)	120 (66-170)	4800 (2700-6900)
Bulgaria	1990	Female	55 to 59	79% (65-89%)	21% (11-35%)	2.8 (2-3.8)	6% (3.1-9%)	9.1% (4.9-14%)	200 (110-300)	6800 (3700-10000)
Bulgaria	1990	Female	60 to 64	76% (61-88%)	24% (12-39%)	2.7 (1.8-3.7)	5.5% (1.8-9.2%)	7.7% (2.7-13%)	270 (94-450)	7800 (2700-13000)
Bulgaria	1990	Female	65 to 69	73% (55-86%)	27% (14-45%)	2.5 (1.7-3.4)	5.1% (1.6-8.9%)	6.8% (2.3-12%)	390 (130-680)	9200 (2900-16000)
Bulgaria	1990	Female	70 to 74	68% (50-81%)	32% (19-50%)	2.3 (1.5-3.3)	4.5% (0.32-8.9%)	5.6% (0.3-11%)	290 (16-580)	5500 (420-11000)
Bulgaria	1990	Female	75 to 79	62% (44-79%)	38% (21-56%)	1.9 (1.2-3)	3.1% (-1.1-7.9%)	3.7% (-1.4-9.5%)	340 (-130-880)	5100 (-1700-13000)
Bulgaria	1990	Female	80 to 84	57% (38-74%)	43% (26-62%)	1.7 (1-2.6)	2% (-1.7-6.1%)	2.2% (-2.2-7%)	220 (-220-680)	2400 (-2000-7300)
Bulgaria	1990	Female	85 to 89	55% (37-70%)	45% (30-63%)	1.6 (0.96-2.5)	0.97% (-2.7-5%)	0.94% (-3.4-5.6%)	56 (-210-340)	500 (-1400-2600)
Bulgaria	1990	Female	90 to 94	56% (40-72%)	44% (28-60%)	1.6 (0.91-2.5)	0.48% (-3.4-5%)	0.35% (-4.4-5.6%)	7 (-87-110)	57 (-420-600)
Bulgaria	1990	Female	95 plus	59% (38-77%)	41% (23-62%)	1.6 (0.79-2.9)	0.16% (-4.3-5.9%)	-0.071% (-5.2-6.4%)	-0.4 (-30-37)	3.5 (-97-130)
Bulgaria	1990	Male	All Ages	86% (82-89%)	14% (11-18%)	5 (4.5-5.6)	15% (12-18%)	17% (13-21%)	10000 (7900-13000)	280000 (230000-330000)
Bulgaria	1990	Male	15 to 19	80% (70-87%)	20% (13-30%)	3 (2-4.1)	7.7% (5-11%)	12% (7.7-18%)	40 (25-57)	3400 (2200-4800)
Bulgaria	1990	Male	20 to 24	88% (78-94%)	12% (6.2-22%)	5.1 (3.6-6.9)	14% (10-19%)	21% (14-28%)	86 (59-120)	7000 (4900-9200)
Bulgaria	1990	Male	25 to 29	90% (82-95%)	10% (5-18%)	5.5 (3.9-7.3)	15% (11-19%)	22% (15-29%)	98 (67-130)	7600 (5500-9700)
Bulgaria	1990	Male	30 to 34	91% (83-96%)	8.8% (4-17%)	5.7 (4.1-7.4)	15% (11-19%)	22% (16-28%)	130 (96-180)	9300 (6900-12000)
Bulgaria	1990	Male	35 to 39	92% (85-96%)	8.1% (3.8-15%)	5.7 (4.1-7.6)	15% (11-20%)	21% (15-28%)	200 (140-260)	12000 (8500-15000)
Bulgaria	1990	Male	40 to 44	92% (85-96%)	7.8% (3.7-15%)	6.7 (5-8.5)	17% (13-21%)	23% (17-29%)	360 (270-440)	19000 (14000-23000)
Bulgaria	1990	Male	45 to 49	92% (83-96%)	8% (3.7-17%)	8.3 (6-11)	20% (16-24%)	26% (20-31%)	510 (400-610)	23000 (18000-27000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	1990	Male	50 to 54	91% (83-96%)	8.6% (3.8-17%)	8.9 (6.5-11)	20% (16-24%)	25% (20-30%)	720 (570-870)	28000 (22000-34000)
Bulgaria	1990	Male	55 to 59	91% (83-96%)	9.4% (4.3-17%)	8.7 (6.6-11)	19% (15-24%)	24% (18-29%)	1100 (840-1300)	37000 (29000-45000)
Bulgaria	1990	Male	60 to 64	90% (81-95%)	10% (4.5-19%)	8.1 (5.9-11)	18% (14-23%)	22% (17-28%)	1400 (1100-1800)	41000 (31000-51000)
Bulgaria	1990	Male	65 to 69	89% (80-95%)	11% (4.7-20%)	7.2 (5.5-9)	18% (14-22%)	21% (16-26%)	1800 (1400-2200)	42000 (32000-52000)
Bulgaria	1990	Male	70 to 74	88% (78-94%)	12% (6.1-22%)	6.1 (4.4-7.9)	16% (11-21%)	18% (12-24%)	1200 (780-1500)	22000 (15000-29000)
Bulgaria	1990	Male	75 to 79	86% (76-93%)	14% (7-24%)	5.1 (3.4-7.1)	13% (6.2-19%)	14% (6.8-22%)	1400 (640-2000)	20000 (9600-29000)
Bulgaria	1990	Male	80 to 84	84% (73-92%)	16% (7.7-27%)	4.4 (3.1-6)	10% (4.2-16%)	11% (4.3-18%)	900 (350-1500)	9800 (4100-16000)
Bulgaria	1990	Male	85 to 89	83% (72-91%)	17% (9.3-28%)	3.9 (2.7-5.4)	6.9% (1-14%)	7.4% (0.53-15%)	300 (22-620)	2400 (370-4700)
Bulgaria	1990	Male	90 to 94	83% (72-91%)	17% (9-28%)	3.4 (2.2-5)	4.1% (-1.5-11%)	4.2% (-2.4-13%)	53 (-31-160)	310 (-110-870)
Bulgaria	1990	Male	95 plus	83% (70-92%)	17% (7.7-30%)	2.9 (1.5-4.8)	2.1% (-3.2-10%)	1.9% (-4.3-11%)	5.1 (-12-31)	23 (-35-110)
Bulgaria	1995	Female	All Ages	76% (71-81%)	24% (19-29%)	1.7 (1.5-2)	2.6% (0.56-4.8%)	2.1% (-1.3-5.7%)	1100 (-700-2900)	37000 (8200-68000)
Bulgaria	1995	Female	15 to 19	77% (66-85%)	23% (15-34%)	1.6 (1.1-2.2)	3% (1.9-4.3%)	6.2% (3.3-9.5%)	8.6 (4.6-13)	950 (590-1400)
Bulgaria	1995	Female	20 to 24	82% (70-90%)	18% (9.6-30%)	1.9 (1.3-2.7)	4.1% (2.7-5.7%)	6.3% (3-10%)	9.8 (4.7-16)	1400 (880-1900)
Bulgaria	1995	Female	25 to 29	83% (72-91%)	17% (8.6-28%)	1.9 (1.3-2.5)	3.9% (2.7-5.2%)	5.6% (2.8-8.6%)	10 (5-16)	1400 (910-1800)
Bulgaria	1995	Female	30 to 34	84% (71-92%)	16% (8.3-29%)	1.9 (1.3-2.6)	3.6% (2.4-5%)	5.7% (3-8.8%)	14 (7-21)	1500 (960-2000)
Bulgaria	1995	Female	35 to 39	84% (73-92%)	16% (8.3-27%)	2.2 (1.5-3)	3.9% (2.5-5.4%)	6.8% (3.8-10%)	25 (14-37)	1900 (1200-2700)
Bulgaria	1995	Female	40 to 44	83% (72-92%)	17% (8.5-28%)	2.4 (1.7-3.2)	4.3% (2.6-6.2%)	7.7% (4.3-11%)	45 (25-66)	2700 (1700-3800)
Bulgaria	1995	Female	45 to 49	82% (69-91%)	18% (8.9-31%)	2.5 (1.8-3.3)	4.5% (2.6-6.7%)	7.7% (4.3-11%)	74 (42-110)	3600 (2100-5200)
Bulgaria	1995	Female	50 to 54	81% (68-90%)	19% (10-32%)	2.4 (1.7-3.2)	4.5% (2.3-7%)	7.2% (3.6-11%)	87 (43-130)	3600 (1800-5600)
Bulgaria	1995	Female	55 to 59	79% (65-90%)	21% (10-35%)	2.3 (1.6-3.2)	4.2% (1.7-7%)	6.4% (2.6-10%)	120 (50-200)	4200 (1600-7000)
Bulgaria	1995	Female	60 to 64	77% (62-88%)	23% (12-38%)	2.2 (1.6-3)	3.4% (0.44-6.7%)	4.9% (0.57-9.5%)	160 (20-330)	4800 (610-9400)
Bulgaria	1995	Female	65 to 69	74% (57-87%)	26% (13-43%)	2.1 (1.5-2.8)	3% (-0.097-6.2%)	4% (-0.1-8.2%)	210 (-5.6-450)	5100 (-170-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	1995	Female	70 to 74	69% (51-82%)	31% (18-49%)	1.9 (1.3-2.7)	2.4% (-1.4-6.3%)	3% (-1.7-7.9%)	260 (-150-660)	4900 (-2700-12000)
Bulgaria	1995	Female	75 to 79	62% (44-78%)	38% (22-56%)	1.6 (1-2.5)	1.3% (-2.4-5.1%)	1.5% (-3.1-6%)	100 (-220-420)	1600 (-2900-6200)
Bulgaria	1995	Female	80 to 84	57% (38-75%)	43% (25-62%)	1.4 (0.86-2.1)	0.4% (-3-3.9%)	0.34% (-3.6-4.3%)	35 (-380-450)	510 (-3800-4900)
Bulgaria	1995	Female	85 to 89	55% (38-71%)	45% (29-62%)	1.4 (0.77-2.1)	-0.43% (-3.5-3.2%)	-0.66% (-4.3-3.5%)	-48 (-310-250)	-270 (-2200-1900)
Bulgaria	1995	Female	90 to 94	56% (40-72%)	44% (28-60%)	1.4 (0.75-2.2)	-0.85% (-4.4-3.2%)	-1.2% (-5.2-3.4%)	-32 (-140-93)	-140 (-690-490)
Bulgaria	1995	Female	95 plus	59% (38-77%)	41% (23-62%)	1.4 (0.67-2.4)	-1.2% (-5.1-3.7%)	-1.6% (-6.1-4.1%)	-9.5 (-37-25)	-28 (-120-91)
Bulgaria	1995	Male	All Ages	86% (83-89%)	14% (11-17%)	4.4 (3.9-4.9)	14% (11-16%)	14% (10-18%)	9100 (6500-12000)	260000 (210000-310000)
Bulgaria	1995	Male	15 to 19	80% (70-88%)	20% (12-30%)	2.6 (1.7-3.5)	6.7% (4.4-9.5%)	11% (6.5-16%)	32 (19-46)	2800 (1900-3900)
Bulgaria	1995	Male	20 to 24	88% (78-94%)	12% (6.3-22%)	4.3 (3-5.8)	13% (9-17%)	18% (12-24%)	78 (51-110)	6300 (4500-8400)
Bulgaria	1995	Male	25 to 29	90% (81-95%)	10% (4.9-19%)	4.6 (3.3-6.1)	13% (9.5-17%)	19% (13-25%)	84 (57-110)	6500 (4700-8400)
Bulgaria	1995	Male	30 to 34	91% (83-96%)	8.7% (4-17%)	4.8 (3.5-6.3)	13% (9.5-17%)	18% (13-25%)	110 (75-150)	7600 (5400-9800)
Bulgaria	1995	Male	35 to 39	92% (85-96%)	8% (3.7-15%)	4.8 (3.4-6.4)	13% (9.4-18%)	18% (12-25%)	170 (110-240)	10000 (7200-14000)
Bulgaria	1995	Male	40 to 44	92% (86-96%)	7.8% (3.6-14%)	5.6 (4.2-7.2)	16% (11-20%)	20% (14-26%)	330 (230-420)	17000 (12000-21000)
Bulgaria	1995	Male	45 to 49	92% (84-96%)	8% (3.7-16%)	7 (5.2-9.2)	19% (14-23%)	23% (17-28%)	610 (450-760)	27000 (21000-34000)
Bulgaria	1995	Male	50 to 54	91% (83-96%)	8.7% (4-17%)	7.5 (5.4-9.5)	19% (15-23%)	23% (18-28%)	740 (570-910)	29000 (22000-35000)
Bulgaria	1995	Male	55 to 59	91% (82-96%)	9.5% (4.3-18%)	7.3 (5.6-9.3)	18% (14-22%)	22% (17-27%)	960 (740-1200)	32000 (25000-39000)
Bulgaria	1995	Male	60 to 64	90% (81-95%)	10% (4.6-19%)	6.8 (5-9)	17% (12-21%)	20% (14-25%)	1300 (950-1700)	38000 (27000-48000)
Bulgaria	1995	Male	65 to 69	89% (79-95%)	11% (4.9-21%)	6.1 (4.7-7.7)	16% (11-20%)	18% (13-23%)	1600 (1100-2000)	37000 (26000-47000)
Bulgaria	1995	Male	70 to 74	88% (78-94%)	12% (6.3-22%)	5.1 (3.8-6.7)	13% (7.5-18%)	15% (8.6-21%)	1600 (880-2200)	29000 (17000-41000)
Bulgaria	1995	Male	75 to 79	86% (76-93%)	14% (7-24%)	4.3 (3-6)	9.7% (3.5-16%)	11% (3.6-18%)	750 (250-1300)	11000 (4000-19000)
Bulgaria	1995	Male	80 to 84	84% (73-92%)	16% (7.6-27%)	3.7 (2.6-5)	6.8% (1.4-13%)	7.4% (1.1-15%)	620 (95-1200)	6900 (1500-13000)
Bulgaria	1995	Male	85 to 89	83% (73-91%)	17% (8.8-27%)	3.3 (2.3-4.4)	3.6% (-1.4-9.4%)	3.6% (-2-10%)	180 (-100-520)	1500 (-580-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	1995	Male	90 to 94	83% (73-91%)	17% (8.7-27%)	2.8 (1.8-4.1)	1.3% (-3.3-7.5%)	1% (-4.3-8.2%)	16 (-69-130)	120 (-300-690)
Bulgaria	1995	Male	95 plus	83% (69-93%)	17% (7.4-31%)	2.4 (1.3-3.8)	-0.056% (-4.4-6.4%)	-0.61% (-5.6-6.8%)	-1.9 (-17-21)	-0.7 (-53-77)
Bulgaria	2000	Female	All Ages	76% (70-81%)	24% (19-30%)	1.5 (1.4-1.8)	1.7% (-0.14-3.7%)	0.91% (-2.2-4%)	490 (-1200-2200)	25000 (-2100-52000)
Bulgaria	2000	Female	15 to 19	76% (64-85%)	24% (15-36%)	1.4 (0.95-1.9)	2.6% (1.7-3.7%)	5.3% (2.7-8.2%)	5.7 (2.9-8.8)	690 (440-990)
Bulgaria	2000	Female	20 to 24	82% (69-90%)	18% (9.8-31%)	1.7 (1.1-2.4)	3.8% (2.6-5.1%)	5.3% (2.4-8.5%)	7.5 (3.4-12)	1200 (770-1700)
Bulgaria	2000	Female	25 to 29	83% (70-91%)	17% (8.8-30%)	1.6 (1.1-2.2)	3.5% (2.5-4.7%)	4.6% (2.2-7.2%)	7.2 (3.4-11)	1200 (780-1600)
Bulgaria	2000	Female	30 to 34	84% (71-92%)	16% (8.1-29%)	1.7 (1.2-2.3)	3.3% (2.2-4.6%)	4.7% (2.3-7.4%)	11 (5.3-17)	1300 (820-1800)
Bulgaria	2000	Female	35 to 39	84% (72-92%)	16% (8.3-28%)	1.9 (1.3-2.7)	3.4% (2.2-4.8%)	5.6% (3-8.5%)	19 (10-29)	1600 (990-2200)
Bulgaria	2000	Female	40 to 44	83% (71-91%)	17% (9-29%)	2.1 (1.5-2.8)	3.6% (2-5.1%)	6.1% (3.1-9.2%)	34 (17-52)	2100 (1200-3000)
Bulgaria	2000	Female	45 to 49	82% (67-91%)	18% (9.5-33%)	2.2 (1.6-2.9)	3.7% (1.9-5.7%)	6.2% (3-9.6%)	56 (27-85)	2800 (1400-4200)
Bulgaria	2000	Female	50 to 54	80% (67-90%)	20% (10-33%)	2.1 (1.6-2.8)	3.6% (1.6-5.8%)	5.7% (2.5-9.2%)	80 (34-130)	3300 (1500-5300)
Bulgaria	2000	Female	55 to 59	78% (64-89%)	22% (11-36%)	2 (1.4-2.7)	3.2% (0.85-5.6%)	4.8% (1.3-8.4%)	88 (24-150)	3000 (810-5300)
Bulgaria	2000	Female	60 to 64	76% (59-88%)	24% (12-41%)	1.9 (1.3-2.7)	2.3% (-0.25-5.3%)	3.3% (-0.38-7.6%)	95 (-10-210)	2700 (-300-6300)
Bulgaria	2000	Female	65 to 69	73% (56-86%)	27% (14-44%)	1.8 (1.3-2.5)	1.8% (-0.94-4.6%)	2.5% (-1.2-6.3%)	130 (-63-320)	3000 (-1400-7600)
Bulgaria	2000	Female	70 to 74	68% (50-82%)	32% (18-50%)	1.6 (1.1-2.3)	1.3% (-1.9-4.6%)	1.6% (-2.4-5.7%)	130 (-200-470)	2500 (-3700-8700)
Bulgaria	2000	Female	75 to 79	61% (42-78%)	39% (22-58%)	1.4 (0.86-2.1)	0.34% (-2.8-3.5%)	0.34% (-3.5-4.3%)	39 (-400-490)	680 (-5600-6900)
Bulgaria	2000	Female	80 to 84	56% (36-73%)	44% (27-64%)	1.2 (0.72-1.8)	-0.36% (-3.3-2.4%)	-0.52% (-3.9-2.8%)	-41 (-310-220)	-360 (-3200-2300)
Bulgaria	2000	Female	85 to 89	53% (36-69%)	47% (31-64%)	1.1 (0.66-1.8)	-1.1% (-3.9-1.9%)	-1.4% (-4.6-2%)	-100 (-350-150)	-700 (-2500-1200)
Bulgaria	2000	Female	90 to 94	55% (38-71%)	45% (29-62%)	1.1 (0.64-1.8)	-1.5% (-4.6-1.9%)	-1.8% (-5.5-2%)	-56 (-170-61)	-260 (-820-340)
Bulgaria	2000	Female	95 plus	58% (36-78%)	42% (22-64%)	1.1 (0.54-2)	-1.8% (-5.3-2.3%)	-2.3% (-6.4-2.3%)	-14 (-41-15)	-45 (-130-56)
Bulgaria	2000	Male	All Ages	86% (83-89%)	14% (11-17%)	3.9 (3.5-4.3)	11% (8.7-14%)	11% (7.5-15%)	6900 (4600-9400)	200000 (160000-250000)
Bulgaria	2000	Male	15 to 19	79% (69-87%)	21% (13-31%)	2.2 (1.5-3)	5.7% (3.8-7.9%)	9.3% (5.5-14%)	20 (12-30)	1900 (1300-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	2000	Male	20 to 24	87% (77-94%)	13% (6.3-23%)	3.7 (2.7-5)	11% (7.5-14%)	15% (9.9-21%)	54 (35-75)	4800 (3300-6400)
Bulgaria	2000	Male	25 to 29	90% (81-95%)	10% (5.1-19%)	4 (2.9-5.2)	11% (8-14%)	16% (11-22%)	60 (39-82)	5100 (3600-6600)
Bulgaria	2000	Male	30 to 34	91% (82-96%)	9.1% (4.2-18%)	4.1 (3-5.4)	11% (7.9-15%)	15% (10-21%)	77 (51-110)	5700 (4000-7500)
Bulgaria	2000	Male	35 to 39	92% (84-96%)	8.2% (3.9-16%)	4.2 (3-5.7)	11% (7.5-15%)	15% (9.4-21%)	110 (69-150)	6900 (4700-9400)
Bulgaria	2000	Male	40 to 44	92% (85-96%)	7.9% (3.8-15%)	4.9 (3.7-6.3)	13% (8.8-17%)	17% (11-22%)	210 (140-290)	11000 (7800-15000)
Bulgaria	2000	Male	45 to 49	92% (84-96%)	8.2% (3.7-16%)	6.1 (4.5-7.8)	16% (11-20%)	20% (14-25%)	430 (300-560)	19000 (14000-25000)
Bulgaria	2000	Male	50 to 54	91% (83-96%)	8.9% (4.1-17%)	6.5 (4.9-8.2)	16% (12-20%)	20% (15-25%)	700 (510-880)	27000 (20000-34000)
Bulgaria	2000	Male	55 to 59	90% (81-95%)	9.8% (4.5-19%)	6.3 (4.8-8.1)	16% (12-20%)	19% (14-24%)	780 (560-990)	26000 (19000-33000)
Bulgaria	2000	Male	60 to 64	89% (80-95%)	11% (4.7-20%)	5.9 (4.3-7.8)	15% (9.7-19%)	17% (11-22%)	930 (610-1200)	27000 (18000-35000)
Bulgaria	2000	Male	65 to 69	89% (80-95%)	11% (4.8-20%)	5.2 (4.1-6.6)	13% (8.2-17%)	15% (9.4-20%)	1200 (750-1600)	28000 (18000-38000)
Bulgaria	2000	Male	70 to 74	87% (78-94%)	13% (6.5-22%)	4.5 (3.2-5.9)	10% (4.9-16%)	12% (5.4-18%)	1100 (540-1800)	22000 (11000-33000)
Bulgaria	2000	Male	75 to 79	86% (76-92%)	14% (7.5-24%)	3.7 (2.5-5.2)	7.3% (1.8-13%)	8.1% (1.9-15%)	860 (200-1600)	13000 (3300-24000)
Bulgaria	2000	Male	80 to 84	84% (72-92%)	16% (8.2-28%)	3.2 (2.3-4.3)	4.6% (0.11-10%)	4.9% (-0.28-11%)	290 (-16-660)	3300 (79-7200)
Bulgaria	2000	Male	85 to 89	83% (72-91%)	17% (9.4-28%)	2.8 (1.9-3.8)	1.9% (-2.1-6.8%)	1.8% (-2.8-7.3%)	88 (-140-350)	810 (-860-2800)
Bulgaria	2000	Male	90 to 94	83% (72-91%)	17% (9.2-28%)	2.5 (1.6-3.6)	0.19% (-3.8-5.2%)	-0.23% (-4.8-5.4%)	-4 (-81-91)	19 (-370-510)
Bulgaria	2000	Male	95 plus	82% (68-92%)	18% (8.2-32%)	2.1 (1.1-3.4)	-0.79% (-4.5-4.2%)	-1.4% (-5.8-4.4%)	-3.9 (-16-12)	-8.6 (-50-46)
Bulgaria	2005	Female	All Ages	73% (67-78%)	27% (22-33%)	1.5 (1.3-1.7)	1.6% (0.041-3.2%)	0.89% (-1.8-3.4%)	470 (-940-1800)	23000 (540-44000)
Bulgaria	2005	Female	15 to 19	75% (64-83%)	25% (17-36%)	1.3 (0.92-1.8)	2.4% (1.6-3.4%)	5.1% (2.6-8.1%)	4.3 (2.2-6.9)	580 (370-830)
Bulgaria	2005	Female	20 to 24	81% (69-89%)	19% (11-31%)	1.7 (1.1-2.3)	3.7% (2.5-4.9%)	5.1% (2.4-8.2%)	5.4 (2.5-8.7)	970 (630-1400)
Bulgaria	2005	Female	25 to 29	82% (71-90%)	18% (9.7-29%)	1.6 (1.1-2.1)	3.4% (2.4-4.4%)	4.2% (2.1-6.7%)	5.9 (2.9-9.6)	1100 (730-1500)
Bulgaria	2005	Female	30 to 34	82% (71-91%)	18% (9.3-29%)	1.6 (1.2-2.2)	3.1% (2.2-4.2%)	4.4% (2.3-6.6%)	8.6 (4.6-13)	1200 (790-1600)
Bulgaria	2005	Female	35 to 39	83% (73-91%)	17% (9.5-27%)	1.9 (1.3-2.6)	3.3% (2.1-4.5%)	5.2% (2.7-8.1%)	16 (8.5-25)	1400 (890-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	2005	Female	40 to 44	83% (73-91%)	17% (9.1-27%)	2.1 (1.5-2.8)	3.4% (2-5%)	5.8% (2.9-8.9%)	28 (14-43)	1800 (1000-2600)
Bulgaria	2005	Female	45 to 49	81% (67-90%)	19% (10-33%)	2.1 (1.6-2.8)	3.6% (1.9-5.6%)	6% (2.9-9.4%)	51 (24-79)	2500 (1300-3800)
Bulgaria	2005	Female	50 to 54	79% (63-88%)	21% (12-37%)	2 (1.5-2.7)	3.5% (1.6-5.6%)	5.5% (2.4-8.9%)	71 (32-110)	3000 (1400-4600)
Bulgaria	2005	Female	55 to 59	76% (61-87%)	24% (13-39%)	1.9 (1.4-2.5)	3% (1.1-5.3%)	4.7% (1.7-7.8%)	92 (35-150)	3200 (1200-5500)
Bulgaria	2005	Female	60 to 64	73% (57-85%)	27% (15-43%)	1.8 (1.2-2.4)	2.3% (0.076-4.8%)	3.3% (0.13-7%)	85 (3.4-180)	2500 (88-5300)
Bulgaria	2005	Female	65 to 69	66% (49-82%)	34% (18-51%)	1.5 (1.1-2)	1.8% (-0.42-4.2%)	2.5% (-0.65-5.7%)	99 (-26-230)	2400 (-590-5400)
Bulgaria	2005	Female	70 to 74	58% (41-73%)	42% (27-59%)	1.3 (0.81-1.8)	1.3% (-1.1-3.7%)	1.6% (-1.4-5%)	120 (-100-360)	2200 (-1900-6800)
Bulgaria	2005	Female	75 to 79	55% (38-72%)	45% (28-62%)	1.2 (0.77-1.8)	0.43% (-2.1-3.1%)	0.48% (-2.7-3.8%)	51 (-280-410)	820 (-3900-5900)
Bulgaria	2005	Female	80 to 84	53% (34-71%)	47% (29-66%)	1.1 (0.66-1.7)	-0.21% (-2.8-2.3%)	-0.32% (-3.4-2.7%)	-39 (-420-330)	-320 (-4400-3600)
Bulgaria	2005	Female	85 to 89	51% (33-68%)	49% (32-67%)	1.1 (0.61-1.6)	-0.83% (-3.4-1.7%)	-1.1% (-4-1.9%)	-63 (-230-110)	-420 (-1700-850)
Bulgaria	2005	Female	90 to 94	52% (35-69%)	48% (31-65%)	1.1 (0.59-1.7)	-1.1% (-4-1.7%)	-1.5% (-4.8-1.9%)	-48 (-160-63)	-220 (-760-350)
Bulgaria	2005	Female	95 plus	55% (34-75%)	45% (25-66%)	1.1 (0.5-1.9)	-1.4% (-4.7-2.1%)	-1.8% (-5.6-2.2%)	-15 (-46-18)	-47 (-150-68)
Bulgaria	2005	Male	All Ages	89% (86-92%)	11% (8.5-14%)	4.1 (3.7-4.5)	11% (8.7-14%)	11% (7.5-15%)	6700 (4500-9300)	190000 (150000-240000)
Bulgaria	2005	Male	15 to 19	82% (74-89%)	18% (11-26%)	2.3 (1.5-3.1)	5.5% (3.7-7.4%)	9.5% (5.6-14%)	15 (9-22)	1600 (1100-2200)
Bulgaria	2005	Male	20 to 24	89% (81-94%)	11% (5.6-19%)	3.9 (2.7-5.1)	11% (7.6-14%)	15% (9.8-21%)	47 (30-64)	4200 (2900-5600)
Bulgaria	2005	Male	25 to 29	92% (86-96%)	8.2% (4.2-14%)	4.1 (3-5.4)	11% (7.8-14%)	16% (10-21%)	58 (37-78)	4900 (3500-6400)
Bulgaria	2005	Male	30 to 34	93% (88-97%)	6.9% (3.5-12%)	4.3 (3.2-5.6)	10% (7.3-14%)	15% (9.8-21%)	69 (44-97)	5200 (3600-7000)
Bulgaria	2005	Male	35 to 39	93% (88-97%)	6.7% (3.2-12%)	4.3 (3.1-5.7)	10% (6.9-14%)	14% (8.8-21%)	93 (57-140)	6000 (4100-8400)
Bulgaria	2005	Male	40 to 44	93% (88-97%)	6.7% (3.3-12%)	4.9 (3.7-6.3)	12% (8-16%)	16% (10-22%)	180 (110-240)	9500 (6400-13000)
Bulgaria	2005	Male	45 to 49	94% (87-97%)	6.5% (3.2-13%)	6.2 (4.7-8)	15% (11-19%)	19% (13-25%)	380 (260-490)	17000 (12000-22000)
Bulgaria	2005	Male	50 to 54	93% (87-97%)	6.9% (3.3-13%)	6.7 (5.1-8.3)	17% (12-20%)	20% (15-25%)	660 (480-820)	26000 (19000-32000)
Bulgaria	2005	Male	55 to 59	92% (86-96%)	7.6% (3.7-14%)	6.6 (5.2-8.1)	16% (12-20%)	19% (14-24%)	940 (690-1200)	31000 (23000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	2005	Male	60 to 64	92% (84-96%)	8.4% (4-16%)	6.1 (4.5-7.9)	15% (9.7-19%)	17% (11-22%)	920 (600-1200)	26000 (17000-35000)
Bulgaria	2005	Male	65 to 69	91% (83-95%)	9.5% (4.6-17%)	5.4 (4.3-6.5)	13% (8.2-17%)	15% (9.3-20%)	990 (620-1300)	23000 (15000-31000)
Bulgaria	2005	Male	70 to 74	90% (82-95%)	10% (5.4-18%)	4.6 (3.4-5.9)	10% (5.1-15%)	12% (5.6-18%)	1100 (510-1600)	20000 (10000-31000)
Bulgaria	2005	Male	75 to 79	87% (79-93%)	13% (6.6-21%)	3.8 (2.7-5.1)	7.2% (1.8-13%)	8% (1.7-15%)	810 (180-1500)	12000 (3100-22000)
Bulgaria	2005	Male	80 to 84	84% (72-91%)	16% (8.5-28%)	3.1 (2.2-4.1)	4.5% (0.3-9.6%)	4.9% (0.078-11%)	450 (7.3-990)	5100 (350-11000)
Bulgaria	2005	Male	85 to 89	82% (70-90%)	18% (10-30%)	2.7 (1.8-3.7)	2% (-1.6-6.5%)	1.9% (-2.3-7.1%)	71 (-83-260)	640 (-510-2100)
Bulgaria	2005	Male	90 to 94	82% (71-90%)	18% (9.8-29%)	2.3 (1.5-3.4)	0.43% (-3.2-4.9%)	0.066% (-4.2-5.3%)	1.2 (-76-96)	46 (-340-530)
Bulgaria	2005	Male	95 plus	81% (66-92%)	19% (8.5-34%)	2 (1.1-3.2)	-0.51% (-4.1-4.3%)	-1.1% (-5.4-4.5%)	-4.1 (-20-17)	-7.6 (-60-63)
Bulgaria	2010	Female	All Ages	72% (66-78%)	28% (22-34%)	1.5 (1.3-1.7)	1.8% (0.26-3.4%)	1.1% (-1.5-3.6%)	570 (-760-1900)	24000 (3400-43000)
Bulgaria	2010	Female	15 to 19	74% (62-83%)	26% (17-38%)	1.4 (0.94-1.9)	2.5% (1.6-3.5%)	5.1% (2.6-8%)	3.4 (1.7-5.4)	460 (300-670)
Bulgaria	2010	Female	20 to 24	81% (68-89%)	19% (11-32%)	1.7 (1.2-2.4)	3.7% (2.6-5%)	5.2% (2.5-8.4%)	4.9 (2.4-7.7)	920 (610-1300)
Bulgaria	2010	Female	25 to 29	82% (71-90%)	18% (9.8-29%)	1.7 (1.1-2.2)	3.5% (2.5-4.5%)	4.3% (2.2-6.6%)	4.8 (2.4-7.4)	940 (640-1300)
Bulgaria	2010	Female	30 to 34	82% (71-90%)	18% (9.6-29%)	1.7 (1.2-2.3)	3.3% (2.3-4.3%)	4.7% (2.7-7.1%)	8.5 (4.8-13)	1200 (780-1600)
Bulgaria	2010	Female	35 to 39	82% (71-90%)	18% (9.8-29%)	1.9 (1.4-2.6)	3.4% (2.3-4.5%)	5.5% (3.1-8.1%)	16 (9.3-24)	1500 (940-2000)
Bulgaria	2010	Female	40 to 44	82% (71-91%)	18% (9.3-29%)	2.2 (1.6-2.8)	3.5% (2.3-5%)	6% (3.4-8.7%)	27 (15-39)	1700 (1100-2400)
Bulgaria	2010	Female	45 to 49	81% (67-90%)	19% (9.9-33%)	2.2 (1.6-2.9)	3.6% (2.1-5.4%)	6.1% (3.3-9.2%)	43 (23-65)	2200 (1300-3300)
Bulgaria	2010	Female	50 to 54	78% (64-88%)	22% (12-36%)	2.1 (1.6-2.8)	3.6% (1.9-5.7%)	5.8% (2.9-8.9%)	66 (34-100)	2800 (1500-4300)
Bulgaria	2010	Female	55 to 59	75% (61-87%)	25% (13-39%)	2 (1.4-2.6)	3.2% (1.4-5.2%)	5% (2.2-7.9%)	88 (39-140)	3200 (1400-5100)
Bulgaria	2010	Female	60 to 64	72% (56-85%)	28% (15-44%)	1.8 (1.3-2.5)	2.4% (0.36-4.9%)	3.7% (0.61-7.3%)	100 (17-200)	3000 (470-6000)
Bulgaria	2010	Female	65 to 69	65% (48-80%)	35% (20-52%)	1.6 (1.1-2.1)	2% (0.02-4.2%)	2.8% (0.043-5.8%)	100 (1.6-210)	2500 (26-5300)
Bulgaria	2010	Female	70 to 74	57% (40-73%)	43% (27-60%)	1.3 (0.83-1.9)	1.5% (-0.73-4%)	2% (-0.97-5.3%)	110 (-54-300)	2200 (-1000-5700)
Bulgaria	2010	Female	75 to 79	54% (38-71%)	46% (29-62%)	1.2 (0.79-1.8)	0.77% (-1.8-3.4%)	0.9% (-2.3-4.2%)	87 (-220-400)	1400 (-3100-6000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	2010	Female	80 to 84	52% (34-70%)	48% (30-66%)	1.1 (0.68-1.7)	0.14% (-2.3-2.6%)	0.088% (-3-3%)	10 (-350-350)	210 (-3500-3800)
Bulgaria	2010	Female	85 to 89	50% (33-66%)	50% (34-67%)	1.1 (0.63-1.6)	-0.48% (-3-2.1%)	-0.68% (-3.6-2.4%)	-64 (-340-220)	-400 (-2500-1700)
Bulgaria	2010	Female	90 to 94	51% (35-67%)	49% (33-65%)	1.1 (0.6-1.7)	-0.83% (-3.7-2.1%)	-1.1% (-4.4-2.3%)	-31 (-120-64)	-140 (-610-340)
Bulgaria	2010	Female	95 plus	54% (33-73%)	46% (27-67%)	1.1 (0.51-1.9)	-1.1% (-4.3-2.5%)	-1.5% (-5.2-2.7%)	-15 (-53-28)	-44 (-170-100)
Bulgaria	2010	Male	All Ages	89% (86-91%)	11% (8.9-14%)	4.3 (3.9-4.8)	11% (8.7-14%)	11% (7.7-15%)	6500 (4400-8800)	180000 (140000-230000)
Bulgaria	2010	Male	15 to 19	81% (73-88%)	19% (12-27%)	2.4 (1.6-3.3)	5.7% (3.9-7.9%)	9.9% (5.8-15%)	13 (7.6-19)	1300 (850-1800)
Bulgaria	2010	Male	20 to 24	89% (81-94%)	11% (6-19%)	4 (2.8-5.5)	11% (7.5-14%)	16% (9.9-22%)	40 (25-56)	3700 (2500-4900)
Bulgaria	2010	Male	25 to 29	91% (86-95%)	8.6% (4.7-14%)	4.3 (3.1-5.6)	11% (7.8-14%)	16% (10-22%)	49 (31-67)	4200 (3000-5600)
Bulgaria	2010	Male	30 to 34	93% (87-97%)	7.2% (3.4-13%)	4.4 (3.3-5.9)	10% (7.4-14%)	15% (9.8-21%)	62 (40-85)	4900 (3500-6500)
Bulgaria	2010	Male	35 to 39	93% (88-97%)	7% (3.4-12%)	4.4 (3.2-5.9)	10% (6.9-14%)	15% (9.3-21%)	85 (54-120)	5700 (3900-7800)
Bulgaria	2010	Male	40 to 44	93% (87-97%)	7.2% (3.4-13%)	5.1 (3.8-6.6)	12% (8-16%)	16% (10-22%)	160 (100-220)	8600 (5900-12000)
Bulgaria	2010	Male	45 to 49	93% (86-97%)	6.8% (3.2-14%)	6.5 (4.8-8.3)	15% (11-19%)	20% (14-25%)	330 (220-420)	15000 (11000-19000)
Bulgaria	2010	Male	50 to 54	93% (86-97%)	7.3% (3.5-14%)	6.9 (5.3-8.7)	17% (12-20%)	21% (15-26%)	570 (420-700)	22000 (17000-27000)
Bulgaria	2010	Male	55 to 59	92% (85-96%)	8% (4-15%)	6.7 (5.3-8.2)	16% (12-20%)	19% (14-24%)	830 (610-1000)	28000 (21000-35000)
Bulgaria	2010	Male	60 to 64	91% (84-96%)	8.8% (4.2-16%)	6.3 (4.7-8.1)	15% (10-19%)	17% (11-23%)	1000 (680-1300)	30000 (20000-38000)
Bulgaria	2010	Male	65 to 69	90% (82-95%)	10% (4.6-18%)	5.5 (4.4-6.7)	13% (8.6-17%)	15% (9.8-20%)	950 (610-1200)	23000 (15000-30000)
Bulgaria	2010	Male	70 to 74	89% (82-94%)	11% (5.5-18%)	4.7 (3.6-6)	11% (5.6-15%)	12% (6.3-18%)	900 (460-1300)	17000 (9100-25000)
Bulgaria	2010	Male	75 to 79	87% (77-93%)	13% (7-23%)	3.9 (2.8-5.3)	7.9% (2.5-13%)	8.9% (2.6-15%)	820 (240-1400)	12000 (4100-21000)
Bulgaria	2010	Male	80 to 84	83% (70-91%)	17% (8.9-30%)	3.2 (2.3-4.3)	5.3% (1.1-11%)	5.9% (0.86-12%)	520 (76-1000)	5800 (1300-11000)
Bulgaria	2010	Male	85 to 89	81% (69-90%)	19% (10-31%)	2.8 (1.9-3.8)	2.9% (-0.8-7.7%)	3% (-1.4-8.4%)	170 (-82-480)	1400 (-400-3800)
Bulgaria	2010	Male	90 to 94	81% (70-90%)	19% (9.9-30%)	2.4 (1.5-3.6)	1.2% (-2.6-6.2%)	0.95% (-3.4-6.7%)	14 (-51-98)	100 (-230-540)
Bulgaria	2010	Male	95 plus	81% (66-92%)	19% (8.3-34%)	2.1 (1.1-3.5)	0.12% (-3.6-5.6%)	-0.36% (-4.8-6.2%)	-1.9 (-25-32)	2.5 (-74-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	2016	Female	All Ages	74% (68-80%)	26% (20-32%)	1.7 (1.4-2.1)	2.1% (0.14-4.2%)	1.4% (-1.7-4.9%)	770 (-910-2600)	27000 (1700-54000)
Bulgaria	2016	Female	15 to 19	75% (62-84%)	25% (16-38%)	1.5 (0.94-2.2)	2.4% (1.6-3.5%)	5.3% (2.8-8.7%)	2 (1.1-3.4)	300 (190-440)
Bulgaria	2016	Female	20 to 24	81% (69-90%)	19% (10-31%)	1.9 (1.2-2.8)	3.7% (2.6-5%)	5.5% (2.7-9%)	4 (1.9-6.8)	840 (540-1200)
Bulgaria	2016	Female	25 to 29	83% (70-91%)	17% (9-30%)	1.8 (1.2-2.5)	3.5% (2.6-4.7%)	4.6% (2.3-7.2%)	3.9 (1.9-6.5)	820 (550-1200)
Bulgaria	2016	Female	30 to 34	83% (70-91%)	17% (8.6-30%)	1.9 (1.2-2.6)	3.4% (2.4-4.6%)	5.1% (2.9-7.7%)	7 (3.7-11)	1000 (700-1500)
Bulgaria	2016	Female	35 to 39	83% (71-91%)	17% (8.5-29%)	2.1 (1.4-3.1)	3.6% (2.4-4.9%)	6.1% (3.4-9.2%)	17 (8.8-28)	1500 (980-2200)
Bulgaria	2016	Female	40 to 44	83% (70-91%)	17% (8.9-30%)	2.3 (1.6-3.3)	3.7% (2.3-5.5%)	6.6% (3.7-9.8%)	27 (14-43)	1800 (1100-2800)
Bulgaria	2016	Female	45 to 49	81% (66-91%)	19% (9.1-34%)	2.4 (1.6-3.4)	3.9% (2.3-6%)	6.7% (3.5-10%)	38 (19-62)	2100 (1100-3200)
Bulgaria	2016	Female	50 to 54	80% (65-90%)	20% (10-35%)	2.3 (1.6-3.3)	4% (2.1-6.2%)	6.5% (3.3-10%)	62 (30-100)	2700 (1400-4300)
Bulgaria	2016	Female	55 to 59	78% (63-89%)	22% (11-37%)	2.2 (1.5-3.1)	3.6% (1.6-6%)	5.7% (2.6-9.4%)	87 (37-150)	3200 (1400-5300)
Bulgaria	2016	Female	60 to 64	75% (59-88%)	25% (12-41%)	2.1 (1.4-3.1)	2.8% (0.57-5.6%)	4.4% (0.94-8.6%)	130 (27-250)	3900 (790-7700)
Bulgaria	2016	Female	65 to 69	72% (55-85%)	28% (15-45%)	2 (1.3-2.8)	2.5% (-0.007-5.5%)	3.6% (0.023-7.7%)	130 (0.76-290)	3300 (-9.3-6900)
Bulgaria	2016	Female	70 to 74	66% (50-81%)	34% (19-50%)	1.8 (1.1-2.6)	2.1% (-0.85-5.4%)	2.9% (-1.1-7.2%)	130 (-54-350)	2600 (-1100-6700)
Bulgaria	2016	Female	75 to 79	59% (41-77%)	41% (23-59%)	1.5 (0.88-2.4)	1.2% (-1.8-4.6%)	1.5% (-2.4-5.8%)	140 (-210-530)	2200 (-3100-8000)
Bulgaria	2016	Female	80 to 84	54% (35-72%)	46% (28-65%)	1.3 (0.75-2)	0.56% (-2.2-3.6%)	0.59% (-2.8-4.3%)	65 (-310-490)	810 (-3200-5300)
Bulgaria	2016	Female	85 to 89	52% (35-69%)	48% (31-65%)	1.2 (0.66-2)	-0.087% (-2.9-3.1%)	-0.24% (-3.5-3.7%)	-29 (-420-420)	-99 (-3000-3300)
Bulgaria	2016	Female	90 to 94	53% (37-71%)	47% (29-63%)	1.2 (0.66-2)	-0.53% (-3.6-3.4%)	-0.77% (-4.5-3.8%)	-32 (-180-160)	-130 (-870-820)
Bulgaria	2016	Female	95 plus	57% (36-76%)	43% (24-64%)	1.2 (0.58-2.2)	-0.81% (-4.2-3.8%)	-1.2% (-5.2-4.2%)	-15 (-68-54)	-42 (-220-200)
Bulgaria	2016	Male	All Ages	87% (83-90%)	13% (10-17%)	4.4 (3.5-5.4)	12% (8.1-15%)	12% (6.9-16%)	6500 (3700-9700)	180000 (120000-240000)
Bulgaria	2016	Male	15 to 19	79% (68-86%)	21% (14-32%)	2.3 (1.5-3.5)	5.6% (3.7-8%)	9.9% (5.5-15%)	7.9 (4.2-13)	830 (520-1200)
Bulgaria	2016	Male	20 to 24	87% (76-93%)	13% (6.7-24%)	4 (2.6-5.8)	11% (7.3-14%)	16% (9.4-23%)	34 (20-52)	3300 (2200-4700)
Bulgaria	2016	Male	25 to 29	89% (81-95%)	11% (5.3-19%)	4.3 (2.9-6.1)	11% (7.7-14%)	16% (10-23%)	38 (23-56)	3600 (2500-5000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bulgaria	2016	Male	30 to 34	91% (82-96%)	9.3% (4.4-18%)	4.5 (3.1-6.2)	11% (7.5-14%)	16% (10-23%)	55 (33-82)	4500 (3100-6200)
Bulgaria	2016	Male	35 to 39	92% (84-96%)	8.4% (4-16%)	4.5 (3-6.4)	10% (6.9-14%)	16% (9.3-22%)	86 (49-130)	5900 (3700-8300)
Bulgaria	2016	Male	40 to 44	92% (85-96%)	8.1% (3.8-15%)	5.3 (3.8-7.2)	12% (8-17%)	17% (11-24%)	160 (93-230)	8800 (5500-12000)
Bulgaria	2016	Male	45 to 49	92% (83-96%)	8.5% (4-17%)	6.6 (4.5-9)	15% (11-19%)	20% (14-26%)	280 (180-400)	13000 (8700-18000)
Bulgaria	2016	Male	50 to 54	91% (83-96%)	9.1% (4.4-17%)	7 (4.9-9.5)	16% (12-21%)	21% (15-26%)	500 (340-680)	20000 (14000-26000)
Bulgaria	2016	Male	55 to 59	90% (82-95%)	10% (4.6-18%)	6.8 (4.8-9)	16% (12-20%)	20% (14-25%)	730 (510-1000)	25000 (18000-34000)
Bulgaria	2016	Male	60 to 64	89% (80-95%)	11% (4.8-20%)	6.3 (4.4-8.5)	15% (9.9-19%)	18% (12-23%)	1100 (710-1600)	32000 (21000-45000)
Bulgaria	2016	Male	65 to 69	88% (78-95%)	12% (4.9-22%)	5.6 (4.1-7.4)	13% (8.4-18%)	16% (9.9-21%)	1000 (620-1400)	24000 (15000-34000)
Bulgaria	2016	Male	70 to 74	87% (78-94%)	13% (6.1-22%)	4.8 (3.2-6.5)	11% (5.5-16%)	13% (6.2-19%)	770 (340-1200)	15000 (7200-23000)
Bulgaria	2016	Male	75 to 79	85% (74-93%)	15% (7.3-26%)	4 (2.6-5.8)	8.3% (2.6-14%)	9.5% (2.8-17%)	830 (240-1500)	13000 (4100-22000)
Bulgaria	2016	Male	80 to 84	83% (71-91%)	17% (8.6-29%)	3.4 (2.3-4.8)	6.2% (1.1-12%)	6.9% (0.83-14%)	590 (65-1200)	6700 (1100-13000)
Bulgaria	2016	Male	85 to 89	82% (70-90%)	18% (9.6-30%)	3 (1.9-4.3)	3.8% (-0.56-9.7%)	4% (-1.1-11%)	290 (-81-780)	2400 (-340-6100)
Bulgaria	2016	Male	90 to 94	82% (71-90%)	18% (9.7-29%)	2.7 (1.7-4)	1.8% (-2.5-7.6%)	1.7% (-3.4-8.3%)	34 (-70-170)	220 (-300-920)
Bulgaria	2016	Male	95 plus	82% (68-92%)	18% (7.9-32%)	2.2 (1.1-3.7)	0.54% (-3.5-6.3%)	0.089% (-4.8-7%)	0.63 (-33-48)	15 (-100-180)
Croatia	1990	Female	All Ages	63% (57-68%)	37% (32-43%)	1.8 (1.5-2.2)	5.6% (3.7-7.6%)	6.5% (3.5-10%)	1700 (900-2600)	42000 (28000-58000)
Croatia	1990	Female	15 to 19	65% (49-78%)	35% (22-51%)	1.3 (0.82-2)	3.1% (2-4.4%)	7.3% (4-11%)	4.4 (2.5-7)	500 (310-720)
Croatia	1990	Female	20 to 24	74% (60-84%)	26% (16-40%)	1.6 (1.1-2.3)	4.5% (3-6.1%)	8% (4.4-13%)	5.5 (3-8.6)	760 (510-1100)
Croatia	1990	Female	25 to 29	74% (61-84%)	26% (16-39%)	1.6 (1.1-2.2)	4.3% (3-5.7%)	7.4% (4.1-11%)	6.7 (3.7-10)	880 (590-1200)
Croatia	1990	Female	30 to 34	72% (59-81%)	28% (19-41%)	1.8 (1.2-2.5)	4.3% (3.1-5.7%)	8.2% (5.2-12%)	11 (6.6-15)	1100 (750-1500)
Croatia	1990	Female	35 to 39	69% (57-79%)	31% (21-43%)	2.1 (1.4-3)	5.1% (3.6-6.7%)	10% (6.9-14%)	21 (14-30)	1600 (1100-2100)
Croatia	1990	Female	40 to 44	68% (57-79%)	32% (21-43%)	2.4 (1.7-3.3)	6.3% (4.7-8.2%)	13% (9.4-16%)	37 (27-48)	2100 (1600-2700)
Croatia	1990	Female	45 to 49	68% (56-78%)	32% (22-44%)	2.8 (1.9-3.9)	8.1% (5.9-11%)	14% (10-18%)	57 (42-76)	2700 (2000-3500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	1990	Female	50 to 54	68% (56-78%)	32% (22-44%)	3.1 (2.2-4.1)	8.7% (6.3-11%)	14% (10-18%)	110 (76-140)	4300 (3200-5700)
Croatia	1990	Female	55 to 59	68% (56-78%)	32% (22-44%)	3.1 (2.2-4.3)	8.8% (6.1-12%)	14% (9.6-18%)	160 (110-220)	5600 (3900-7700)
Croatia	1990	Female	60 to 64	66% (52-77%)	34% (23-48%)	3 (2.1-4.2)	7.8% (4.7-11%)	11% (6.7-16%)	190 (120-290)	5800 (3500-8700)
Croatia	1990	Female	65 to 69	62% (47-74%)	38% (26-53%)	2.7 (1.8-3.7)	6.9% (3.9-11%)	9.3% (5.3-14%)	250 (140-380)	6100 (3400-9100)
Croatia	1990	Female	70 to 74	55% (41-66%)	45% (34-59%)	2.2 (1.4-3.3)	5.7% (2-10%)	7.2% (2.5-13%)	180 (65-320)	3500 (1300-6200)
Croatia	1990	Female	75 to 79	46% (33-61%)	54% (39-67%)	1.8 (1.1-2.8)	4.1% (0.71-8.4%)	4.9% (0.69-10%)	230 (34-480)	3500 (580-7000)
Croatia	1990	Female	80 to 84	41% (27-55%)	59% (45-73%)	1.7 (0.98-2.5)	3.6% (0.7-7.4%)	4.1% (0.65-8.6%)	220 (34-470)	2400 (470-5100)
Croatia	1990	Female	85 to 89	40% (27-54%)	60% (46-73%)	1.7 (0.97-2.6)	3.3% (0.58-7.1%)	3.8% (0.45-8.3%)	130 (14-280)	1000 (180-2200)
Croatia	1990	Female	90 to 94	41% (28-56%)	59% (44-72%)	1.8 (1-2.9)	3.5% (0.39-8.1%)	4.1% (0.25-9.7%)	53 (3.3-130)	290 (30-660)
Croatia	1990	Female	95 plus	45% (28-63%)	55% (37-72%)	2 (0.99-3.5)	4.2% (0.092-10%)	4.9% (-0.17-12%)	18 (-0.6-45)	64 (1.5-160)
Croatia	1990	Male	All Ages	74% (69-79%)	26% (21-31%)	5.4 (4.6-6.4)	18% (15-21%)	21% (17-25%)	5700 (4600-7000)	170000 (140000-200000)
Croatia	1990	Male	15 to 19	66% (51-80%)	34% (20-49%)	3.1 (1.9-4.6)	8.9% (5.7-13%)	15% (8.9-22%)	23 (13-35)	2000 (1200-2800)
Croatia	1990	Male	20 to 24	78% (65-87%)	22% (13-35%)	5.2 (3.5-7.4)	18% (12-24%)	25% (17-35%)	68 (45-94)	5200 (3600-7000)
Croatia	1990	Male	25 to 29	81% (69-89%)	19% (11-31%)	5.1 (3.5-6.8)	17% (12-22%)	25% (18-33%)	72 (50-97)	5400 (3900-7000)
Croatia	1990	Male	30 to 34	82% (70-90%)	18% (9.8-30%)	5.5 (3.8-7.3)	18% (13-22%)	26% (20-33%)	100 (76-130)	6900 (5200-8800)
Croatia	1990	Male	35 to 39	82% (73-90%)	18% (9.7-27%)	6.5 (4.5-8.9)	20% (15-24%)	29% (22-35%)	160 (120-200)	9600 (7500-12000)
Croatia	1990	Male	40 to 44	83% (74-91%)	17% (9.5-26%)	7.8 (5.8-10)	22% (18-26%)	31% (26-36%)	230 (190-280)	12000 (10000-14000)
Croatia	1990	Male	45 to 49	84% (74-91%)	16% (9.1-26%)	9.5 (6.7-13)	24% (20-28%)	31% (27-36%)	310 (260-370)	14000 (11000-16000)
Croatia	1990	Male	50 to 54	84% (74-91%)	16% (8.7-26%)	9.6 (6.8-13)	23% (19-27%)	28% (24-33%)	540 (430-650)	21000 (17000-25000)
Croatia	1990	Male	55 to 59	83% (72-91%)	17% (8.8-28%)	8.8 (6.5-12)	21% (18-25%)	25% (21-30%)	720 (580-890)	24000 (20000-29000)
Croatia	1990	Male	60 to 64	82% (70-90%)	18% (9.9-30%)	8.5 (6.1-12)	19% (16-24%)	23% (18-28%)	760 (600-940)	22000 (17000-27000)
Croatia	1990	Male	65 to 69	81% (69-89%)	19% (11-31%)	8.6 (6.4-11)	19% (14-23%)	22% (17-27%)	700 (530-880)	17000 (13000-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	1990	Male	70 to 74	79% (68-87%)	21% (13-32%)	8.3 (5.9-11)	18% (13-22%)	21% (15-26%)	520 (390-650)	9800 (7400-12000)
Croatia	1990	Male	75 to 79	76% (63-87%)	24% (13-37%)	7.4 (4.8-10)	16% (11-21%)	18% (13-24%)	680 (480-910)	10000 (7000-13000)
Croatia	1990	Male	80 to 84	74% (59-85%)	26% (15-41%)	6.1 (4.1-8.7)	14% (9.5-19%)	16% (10-22%)	550 (360-760)	5900 (4000-8100)
Croatia	1990	Male	85 to 89	72% (57-83%)	28% (17-43%)	5.1 (3.3-7.4)	11% (6.3-16%)	12% (6.9-19%)	210 (110-320)	1600 (950-2500)
Croatia	1990	Male	90 to 94	71% (56-82%)	29% (18-44%)	4.3 (2.6-6.5)	8.1% (2.9-14%)	9.2% (2.9-16%)	48 (15-82)	270 (99-450)
Croatia	1990	Male	95 plus	69% (51-83%)	31% (17-49%)	3.4 (1.7-5.8)	5.5% (0.079-12%)	5.9% (-0.71-14%)	6.5 (-0.77-15)	26 (0.37-55)
Croatia	1995	Female	All Ages	62% (57-67%)	38% (33-43%)	1.9 (1.6-2.2)	5.5% (3.9-7.4%)	6.6% (3.8-10%)	1700 (970-2600)	42000 (29000-56000)
Croatia	1995	Female	15 to 19	65% (50-79%)	35% (21-50%)	1.3 (0.83-1.9)	2.8% (1.8-4.1%)	5.2% (2.7-8.7%)	4.3 (2.3-7)	470 (300-670)
Croatia	1995	Female	20 to 24	74% (60-84%)	26% (16-40%)	1.6 (1-2.3)	4% (2.7-5.4%)	5.7% (2.8-9.5%)	4.9 (2.6-7.6)	680 (450-950)
Croatia	1995	Female	25 to 29	74% (61-84%)	26% (16-39%)	1.6 (1.1-2.2)	3.9% (2.9-5.1%)	5.8% (2.9-9.1%)	4.9 (2.6-7.4)	690 (470-940)
Croatia	1995	Female	30 to 34	72% (59-81%)	28% (19-41%)	1.8 (1.2-2.4)	3.9% (2.8-5.1%)	6.6% (4-9.7%)	8.3 (5-12)	900 (620-1200)
Croatia	1995	Female	35 to 39	69% (58-79%)	31% (21-42%)	2.1 (1.4-2.9)	4.7% (3.4-6.1%)	9% (6-12%)	19 (12-25)	1400 (960-1800)
Croatia	1995	Female	40 to 44	67% (56-79%)	33% (21-44%)	2.4 (1.7-3.2)	5.8% (4.4-7.5%)	11% (8.6-14%)	35 (26-45)	2000 (1600-2600)
Croatia	1995	Female	45 to 49	68% (55-78%)	32% (22-45%)	2.8 (1.9-3.7)	7.1% (5.3-9.3%)	13% (9.6-16%)	59 (44-76)	2800 (2100-3600)
Croatia	1995	Female	50 to 54	68% (56-78%)	32% (22-44%)	3.1 (2.2-4)	8.5% (6.3-11%)	14% (10-18%)	82 (60-110)	3300 (2400-4200)
Croatia	1995	Female	55 to 59	68% (55-78%)	32% (22-45%)	3.1 (2.2-4.2)	8.9% (6.4-12%)	14% (9.9-18%)	150 (110-190)	5100 (3600-6700)
Croatia	1995	Female	60 to 64	66% (51-77%)	34% (23-49%)	3 (2.1-4.1)	7.9% (5-11%)	11% (7.2-16%)	200 (130-280)	5900 (3800-8400)
Croatia	1995	Female	65 to 69	61% (47-73%)	39% (27-53%)	2.7 (1.9-3.6)	7% (4.2-10%)	9.5% (5.8-14%)	260 (160-380)	6200 (3800-9300)
Croatia	1995	Female	70 to 74	54% (41-66%)	46% (34-59%)	2.2 (1.4-3.2)	5.9% (2.4-9.7%)	7.5% (3.1-12%)	310 (120-510)	5900 (2400-9800)
Croatia	1995	Female	75 to 79	46% (32-60%)	54% (40-68%)	1.8 (1.1-2.7)	4.1% (0.85-8%)	4.9% (0.91-9.7%)	170 (31-340)	2600 (520-5100)
Croatia	1995	Female	80 to 84	40% (27-55%)	60% (45-73%)	1.6 (0.97-2.4)	3.5% (0.77-7.2%)	4% (0.81-8.3%)	190 (37-390)	2100 (450-4200)
Croatia	1995	Female	85 to 89	39% (26-53%)	61% (47-74%)	1.7 (0.97-2.6)	3.4% (0.62-7.1%)	3.9% (0.52-8.6%)	150 (20-320)	1200 (230-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	1995	Female	90 to 94	41% (28-55%)	59% (45-72%)	1.8 (0.98-2.8)	3.6% (0.48-8%)	4.2% (0.25-9.8%)	73 (4.6-170)	400 (55-880)
Croatia	1995	Female	95 plus	44% (27-62%)	56% (38-73%)	2 (0.99-3.4)	4.2% (0.019-10%)	5% (-0.18-13%)	25 (-0.92-61)	89 (0.38-220)
Croatia	1995	Male	All Ages	75% (69-80%)	25% (20-31%)	5.6 (4.8-6.3)	17% (14-20%)	20% (17-24%)	5600 (4500-6900)	160000 (130000-190000)
Croatia	1995	Male	15 to 19	67% (51-80%)	33% (20-49%)	3.1 (1.9-4.6)	7.7% (4.8-11%)	11% (6.3-17%)	24 (14-36)	2000 (1300-2900)
Croatia	1995	Male	20 to 24	78% (64-88%)	22% (12-36%)	5.2 (3.5-7.2)	14% (8.9-19%)	17% (10-27%)	62 (41-86)	4700 (3300-6300)
Croatia	1995	Male	25 to 29	81% (69-89%)	19% (11-31%)	5.1 (3.6-6.9)	14% (9.7-19%)	19% (12-27%)	60 (42-80)	4400 (3200-5800)
Croatia	1995	Male	30 to 34	82% (70-90%)	18% (10-30%)	5.5 (3.9-7.4)	15% (11-19%)	21% (14-28%)	81 (59-100)	5400 (4100-6900)
Croatia	1995	Male	35 to 39	82% (73-90%)	18% (10-27%)	6.5 (4.6-8.8)	18% (14-22%)	25% (19-31%)	140 (100-170)	8100 (6200-9800)
Croatia	1995	Male	40 to 44	83% (73-91%)	17% (9.4-27%)	7.8 (5.9-9.9)	21% (17-24%)	28% (24-33%)	230 (190-280)	12000 (10000-14000)
Croatia	1995	Male	45 to 49	84% (73-91%)	16% (9-27%)	9.5 (6.9-12)	23% (19-26%)	29% (25-34%)	330 (270-400)	15000 (12000-18000)
Croatia	1995	Male	50 to 54	84% (73-91%)	16% (9.1-27%)	9.6 (6.9-12)	23% (19-27%)	28% (23-33%)	390 (310-460)	15000 (12000-18000)
Croatia	1995	Male	55 to 59	83% (72-91%)	17% (8.9-28%)	8.7 (6.4-11)	21% (17-25%)	25% (21-30%)	630 (500-760)	21000 (17000-25000)
Croatia	1995	Male	60 to 64	82% (70-90%)	18% (10-30%)	8.5 (6.2-11)	19% (15-23%)	23% (18-27%)	810 (630-1000)	23000 (18000-28000)
Croatia	1995	Male	65 to 69	80% (69-88%)	20% (12-31%)	8.5 (6.3-11)	19% (14-23%)	22% (16-26%)	880 (670-1100)	21000 (16000-26000)
Croatia	1995	Male	70 to 74	78% (67-86%)	22% (14-33%)	8.2 (5.9-11)	18% (13-23%)	21% (15-26%)	760 (560-950)	14000 (11000-18000)
Croatia	1995	Male	75 to 79	76% (63-86%)	24% (14-37%)	7.3 (4.9-10)	16% (11-21%)	18% (13-24%)	470 (330-610)	6900 (4900-9000)
Croatia	1995	Male	80 to 84	74% (59-85%)	26% (15-41%)	6.1 (4.1-8.6)	14% (9.9-20%)	16% (11-22%)	470 (310-640)	5100 (3400-6800)
Croatia	1995	Male	85 to 89	71% (57-82%)	29% (18-43%)	5.1 (3.4-7.2)	11% (6.7-16%)	13% (7.5-19%)	230 (140-350)	1900 (1100-2700)
Croatia	1995	Male	90 to 94	70% (56-82%)	30% (18-44%)	4.2 (2.6-6.4)	8.2% (3.3-13%)	9.3% (3.3-16%)	68 (24-110)	380 (150-620)
Croatia	1995	Male	95 plus	69% (50-84%)	31% (16-50%)	3.3 (1.6-5.7)	5.6% (0.12-12%)	6% (-0.56-14%)	9.8 (-0.87-23)	38 (0.79-82)
Croatia	2000	Female	All Ages	60% (54-64%)	40% (36-46%)	1.9 (1.6-2.2)	5.5% (4-7.3%)	6.7% (4-9.8%)	1700 (1000-2600)	39000 (28000-53000)
Croatia	2000	Female	15 to 19	63% (47-77%)	37% (23-53%)	1.3 (0.84-1.9)	2.9% (1.8-4.1%)	7% (3.8-11%)	3.3 (1.7-5.2)	390 (240-560)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	2000	Female	20 to 24	72% (58-83%)	28% (17-42%)	1.6 (1.1-2.3)	4.2% (2.9-5.7%)	7.4% (4.1-12%)	3.7 (2-5.8)	590 (400-830)
Croatia	2000	Female	25 to 29	71% (58-82%)	29% (18-42%)	1.6 (1.1-2.1)	3.9% (2.8-5%)	6.8% (3.9-10%)	3.7 (2.1-5.5)	610 (420-830)
Croatia	2000	Female	30 to 34	68% (54-79%)	32% (21-46%)	1.7 (1.1-2.3)	3.9% (2.9-5.1%)	7.6% (4.9-11%)	6.2 (4-8.7)	720 (510-960)
Croatia	2000	Female	35 to 39	65% (54-76%)	35% (24-46%)	2 (1.3-2.9)	4.5% (3.3-5.9%)	9.6% (6.6-13%)	14 (9.4-18)	1100 (780-1400)
Croatia	2000	Female	40 to 44	65% (55-76%)	35% (24-45%)	2.4 (1.8-3.2)	5.5% (4.2-7.1%)	12% (8.9-15%)	29 (22-37)	1700 (1300-2200)
Croatia	2000	Female	45 to 49	66% (55-76%)	34% (24-45%)	2.8 (2-3.7)	6.8% (4.9-8.9%)	13% (9.7-16%)	52 (39-67)	2600 (1900-3300)
Croatia	2000	Female	50 to 54	66% (56-75%)	34% (25-44%)	3.1 (2.2-4)	8% (6-10%)	14% (10-17%)	80 (60-100)	3300 (2500-4300)
Croatia	2000	Female	55 to 59	67% (56-77%)	33% (23-44%)	3.2 (2.3-4.2)	8.8% (6.4-12%)	14% (10-18%)	100 (76-140)	3600 (2600-4800)
Croatia	2000	Female	60 to 64	65% (53-76%)	35% (24-47%)	3.1 (2.2-4.2)	7.9% (5.2-11%)	12% (7.8-16%)	180 (120-240)	5200 (3400-7200)
Croatia	2000	Female	65 to 69	60% (48-71%)	40% (29-52%)	2.7 (2-3.5)	7.1% (4.5-9.9%)	9.8% (6.3-14%)	250 (160-350)	6000 (3900-8500)
Croatia	2000	Female	70 to 74	51% (39-62%)	49% (38-61%)	2.1 (1.4-3)	6% (3-9.3%)	7.7% (3.8-12%)	290 (150-460)	5700 (2900-8800)
Croatia	2000	Female	75 to 79	41% (30-56%)	59% (44-70%)	1.7 (1-2.5)	4.4% (1.6-7.8%)	5.3% (1.9-9.5%)	280 (100-500)	4200 (1500-7500)
Croatia	2000	Female	80 to 84	38% (26-52%)	62% (48-74%)	1.6 (0.98-2.3)	3.7% (1.4-6.9%)	4.4% (1.6-8.4%)	180 (65-340)	2000 (750-3800)
Croatia	2000	Female	85 to 89	39% (26-52%)	61% (48-74%)	1.7 (0.99-2.6)	3.5% (0.94-6.9%)	4.1% (0.84-8.1%)	130 (27-250)	1000 (280-2000)
Croatia	2000	Female	90 to 94	41% (28-54%)	59% (46-72%)	1.9 (1-2.9)	3.8% (0.57-8.2%)	4.3% (0.42-9.7%)	100 (9.8-230)	540 (81-1200)
Croatia	2000	Female	95 plus	44% (28-62%)	56% (38-72%)	2.1 (1-3.4)	4.5% (0.17-10%)	5.2% (0.046-12%)	40 (0.37-94)	140 (5.4-320)
Croatia	2000	Male	All Ages	77% (72-82%)	23% (18-28%)	6.2 (5.6-6.9)	18% (15-21%)	21% (17-25%)	5700 (4600-7000)	150000 (130000-180000)
Croatia	2000	Male	15 to 19	70% (56-82%)	30% (18-44%)	3.4 (2.2-4.8)	9.5% (6.3-14%)	17% (10-24%)	22 (13-32)	1800 (1200-2600)
Croatia	2000	Male	20 to 24	80% (67-88%)	20% (12-33%)	5.6 (3.9-7.6)	17% (12-23%)	27% (18-36%)	51 (35-72)	4000 (2900-5300)
Croatia	2000	Male	25 to 29	82% (72-90%)	18% (9.9-28%)	5.5 (4-7.2)	16% (12-20%)	26% (19-34%)	50 (36-66)	3800 (2800-4900)
Croatia	2000	Male	30 to 34	84% (73-90%)	16% (9.5-27%)	5.9 (4.3-7.9)	16% (12-20%)	27% (21-34%)	60 (45-77)	4200 (3200-5200)
Croatia	2000	Male	35 to 39	84% (76-91%)	16% (9.3-24%)	6.9 (4.9-9.3)	19% (14-23%)	30% (23-36%)	110 (80-130)	6400 (4900-7700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	2000	Male	40 to 44	84% (76-91%)	16% (9.2-24%)	8.4 (6.5-11)	22% (18-25%)	32% (27-37%)	200 (160-240)	10000 (8600-12000)
Croatia	2000	Male	45 to 49	85% (76-91%)	15% (8.8-24%)	10 (7.6-13)	23% (19-27%)	31% (27-35%)	330 (280-390)	15000 (12000-17000)
Croatia	2000	Male	50 to 54	85% (75-92%)	15% (8.5-25%)	10 (7.6-13)	22% (19-27%)	29% (24-34%)	410 (330-490)	16000 (13000-19000)
Croatia	2000	Male	55 to 59	85% (76-92%)	15% (8.3-24%)	9.6 (7.4-12)	22% (18-26%)	27% (22-32%)	450 (360-550)	15000 (12000-18000)
Croatia	2000	Male	60 to 64	84% (74-91%)	16% (9.3-26%)	9.3 (6.8-12)	20% (16-24%)	24% (19-29%)	750 (590-920)	21000 (17000-26000)
Croatia	2000	Male	65 to 69	82% (72-89%)	18% (11-28%)	9.1 (7.1-11)	19% (15-23%)	22% (17-27%)	950 (720-1200)	22000 (17000-28000)
Croatia	2000	Male	70 to 74	80% (71-87%)	20% (13-29%)	8.9 (6.6-12)	18% (13-23%)	21% (15-26%)	950 (700-1200)	18000 (13000-22000)
Croatia	2000	Male	75 to 79	78% (65-87%)	22% (13-35%)	7.9 (5.5-11)	17% (12-22%)	19% (14-25%)	700 (510-910)	10000 (7600-13000)
Croatia	2000	Male	80 to 84	75% (61-86%)	25% (14-39%)	6.5 (4.7-8.8)	15% (10-19%)	17% (12-22%)	390 (270-530)	4300 (3000-5700)
Croatia	2000	Male	85 to 89	73% (59-83%)	27% (17-41%)	5.5 (3.8-7.6)	12% (7.7-17%)	13% (8.6-19%)	210 (130-300)	1600 (1100-2300)
Croatia	2000	Male	90 to 94	72% (58-83%)	28% (17-42%)	4.6 (3-6.8)	9.5% (4.6-14%)	11% (4.9-17%)	100 (46-160)	540 (270-830)
Croatia	2000	Male	95 plus	70% (52-84%)	30% (16-48%)	3.6 (1.8-6)	6.7% (0.97-13%)	7.2% (0.56-15%)	19 (1.4-40)	71 (10-140)
Croatia	2005	Female	All Ages	59% (55-64%)	41% (36-45%)	1.8 (1.5-2)	4.6% (3.3-6%)	5.3% (3-7.9%)	1400 (780-2100)	32000 (22000-42000)
Croatia	2005	Female	15 to 19	68% (54-80%)	32% (20-46%)	1.4 (0.92-1.9)	2.7% (1.8-3.9%)	7.1% (3.9-11%)	2.7 (1.5-4.2)	340 (220-490)
Croatia	2005	Female	20 to 24	75% (62-83%)	25% (17-38%)	1.6 (1.1-2.2)	4% (2.8-5.4%)	7% (3.8-11%)	3.3 (1.8-5.1)	560 (370-790)
Croatia	2005	Female	25 to 29	72% (60-81%)	28% (19-40%)	1.5 (1-2)	3.7% (2.8-4.8%)	6.2% (3.6-9.1%)	2.9 (1.6-4.3)	560 (390-760)
Croatia	2005	Female	30 to 34	68% (57-78%)	32% (22-43%)	1.6 (1.1-2.2)	3.6% (2.7-4.8%)	7.1% (4.6-9.9%)	4.7 (3.1-6.7)	640 (460-870)
Croatia	2005	Female	35 to 39	65% (56-76%)	35% (24-44%)	1.9 (1.3-2.7)	4.1% (3-5.3%)	9% (6.2-12%)	10 (7-14)	860 (620-1100)
Croatia	2005	Female	40 to 44	63% (53-74%)	37% (26-47%)	2.2 (1.6-2.9)	4.8% (3.5-6.2%)	11% (8.1-14%)	21 (16-27)	1300 (1000-1700)
Croatia	2005	Female	45 to 49	64% (54-73%)	36% (27-46%)	2.5 (1.9-3.3)	5.8% (4.3-7.4%)	11% (8.7-14%)	42 (32-54)	2100 (1600-2700)
Croatia	2005	Female	50 to 54	65% (55-74%)	35% (26-45%)	2.8 (2.1-3.6)	6.8% (5-8.7%)	12% (9.1-15%)	69 (52-89)	2900 (2200-3800)
Croatia	2005	Female	55 to 59	65% (55-75%)	35% (25-45%)	2.9 (2.2-3.7)	7.2% (5.3-9.4%)	12% (9.1-16%)	91 (68-120)	3300 (2500-4300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	2005	Female	60 to 64	64% (54-74%)	36% (26-46%)	2.8 (2.1-3.8)	6.9% (4.6-9.6%)	10% (7.2-14%)	100 (71-150)	3100 (2100-4400)
Croatia	2005	Female	65 to 69	59% (47-69%)	41% (31-53%)	2.5 (1.8-3.2)	6% (3.9-8.5%)	8.7% (5.8-12%)	180 (120-250)	4500 (2900-6200)
Croatia	2005	Female	70 to 74	49% (39-59%)	51% (41-61%)	1.9 (1.3-2.6)	4.7% (2.1-7.4%)	6.2% (2.9-9.8%)	220 (99-350)	4300 (2000-6800)
Croatia	2005	Female	75 to 79	38% (28-51%)	62% (49-72%)	1.4 (0.9-2.1)	3.3% (1.1-6%)	4.1% (1.3-7.5%)	210 (68-390)	3200 (1100-6000)
Croatia	2005	Female	80 to 84	36% (24-49%)	64% (51-76%)	1.4 (0.86-2)	3.1% (1-5.8%)	3.6% (1.1-6.7%)	210 (66-410)	2400 (750-4500)
Croatia	2005	Female	85 to 89	37% (25-50%)	63% (50-75%)	1.5 (0.91-2.3)	2.9% (0.68-5.8%)	3.3% (0.58-6.9%)	110 (19-240)	930 (220-1900)
Croatia	2005	Female	90 to 94	39% (27-52%)	61% (48-73%)	1.7 (0.94-2.5)	2.9% (0.12-6.7%)	3.3% (-0.048-8%)	58 (-0.85-140)	330 (14-750)
Croatia	2005	Female	95 plus	42% (27-60%)	58% (40-73%)	1.9 (0.94-3)	3.4% (-0.28-8.3%)	3.8% (-0.66-9.8%)	32 (-5.4-80)	120 (-9.7-280)
Croatia	2005	Male	All Ages	79% (74-83%)	21% (17-26%)	6.1 (5.5-6.7)	17% (15-20%)	21% (17-24%)	5500 (4400-6600)	140000 (120000-170000)
Croatia	2005	Male	15 to 19	72% (59-83%)	28% (17-41%)	3.3 (2.3-4.6)	8.8% (5.9-12%)	16% (10-23%)	17 (10-24)	1500 (960-2000)
Croatia	2005	Male	20 to 24	82% (71-89%)	18% (11-29%)	5.5 (3.9-7.4)	16% (11-21%)	26% (17-35%)	44 (30-61)	3600 (2500-4700)
Croatia	2005	Male	25 to 29	84% (75-90%)	16% (9.6-25%)	5.3 (3.9-6.8)	14% (11-19%)	24% (17-32%)	42 (30-56)	3300 (2500-4300)
Croatia	2005	Male	30 to 34	85% (77-91%)	15% (9-23%)	5.7 (4.2-7.5)	14% (11-18%)	25% (19-32%)	48 (35-62)	3500 (2600-4500)
Croatia	2005	Male	35 to 39	85% (78-91%)	15% (8.7-22%)	6.7 (4.8-8.9)	17% (13-21%)	29% (22-35%)	75 (57-94)	4700 (3600-5800)
Croatia	2005	Male	40 to 44	85% (78-92%)	15% (8.4-22%)	8 (6.2-10)	20% (17-23%)	31% (26-36%)	150 (130-180)	8200 (6800-9700)
Croatia	2005	Male	45 to 49	86% (79-92%)	14% (8.2-21%)	9.7 (7.3-12)	22% (19-26%)	31% (26-35%)	280 (230-330)	13000 (11000-15000)
Croatia	2005	Male	50 to 54	86% (77-92%)	14% (8-23%)	9.8 (7.4-12)	22% (18-26%)	28% (24-33%)	420 (350-500)	17000 (14000-20000)
Croatia	2005	Male	55 to 59	86% (76-92%)	14% (7.6-24%)	9 (7-11)	21% (17-25%)	26% (22-31%)	490 (400-580)	17000 (14000-20000)
Croatia	2005	Male	60 to 64	85% (76-92%)	15% (8.4-24%)	8.9 (6.7-12)	20% (16-24%)	24% (19-28%)	500 (400-610)	14000 (12000-18000)
Croatia	2005	Male	65 to 69	82% (74-89%)	18% (11-26%)	8.7 (6.9-11)	19% (15-22%)	22% (17-27%)	810 (640-990)	19000 (15000-24000)
Croatia	2005	Male	70 to 74	80% (71-88%)	20% (12-29%)	8.4 (6.4-11)	17% (13-22%)	20% (16-25%)	960 (730-1200)	18000 (14000-23000)
Croatia	2005	Male	75 to 79	79% (67-87%)	21% (13-33%)	7.6 (5.4-10)	16% (12-20%)	18% (13-23%)	840 (610-1100)	12000 (9200-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	2005	Male	80 to 84	77% (63-86%)	23% (14-37%)	6.3 (4.6-8.4)	14% (10-19%)	16% (11-21%)	530 (380-700)	5800 (4200-7600)
Croatia	2005	Male	85 to 89	75% (63-84%)	25% (16-37%)	5.3 (3.8-7.2)	11% (7.5-16%)	13% (8.2-18%)	180 (120-260)	1500 (960-2000)
Croatia	2005	Male	90 to 94	73% (61-84%)	27% (16-39%)	4.5 (3-6.5)	8.6% (3.8-13%)	9.5% (3.7-15%)	66 (26-110)	360 (160-570)
Croatia	2005	Male	95 plus	71% (55-85%)	29% (15-45%)	3.5 (1.8-5.7)	5.7% (0.16-12%)	6% (-0.37-14%)	15 (-0.93-35)	59 (1.7-120)
Croatia	2010	Female	All Ages	60% (55-64%)	40% (36-45%)	1.6 (1.4-1.8)	3.6% (2.4-4.8%)	3.8% (1.8-6.1%)	1000 (480-1700)	24000 (16000-33000)
Croatia	2010	Female	15 to 19	68% (53-80%)	32% (20-47%)	1.2 (0.8-1.7)	2.4% (1.6-3.3%)	6.2% (3.3-9.7%)	1.7 (0.89-2.7)	240 (150-340)
Croatia	2010	Female	20 to 24	75% (62-84%)	25% (16-38%)	1.4 (0.96-2.1)	3.8% (2.7-5%)	6.2% (3.4-9.6%)	2.2 (1.2-3.3)	460 (310-650)
Croatia	2010	Female	25 to 29	72% (61-82%)	28% (18-39%)	1.3 (0.93-1.8)	3.5% (2.6-4.7%)	5.2% (2.9-7.7%)	2.3 (1.3-3.4)	520 (360-730)
Croatia	2010	Female	30 to 34	70% (58-79%)	30% (21-42%)	1.5 (1-2)	3.4% (2.4-4.4%)	6.1% (3.8-8.6%)	3.4 (2.1-4.9)	570 (390-790)
Croatia	2010	Female	35 to 39	67% (58-77%)	33% (23-42%)	1.8 (1.2-2.5)	3.7% (2.7-4.8%)	7.9% (5.5-11%)	7.7 (5.1-10)	740 (530-960)
Croatia	2010	Female	40 to 44	64% (53-75%)	36% (25-47%)	1.9 (1.4-2.6)	4.2% (3.2-5.4%)	9.4% (7-12%)	15 (11-20)	1000 (750-1300)
Croatia	2010	Female	45 to 49	64% (54-75%)	36% (25-46%)	2.2 (1.6-3)	4.9% (3.7-6.4%)	10% (7.5-13%)	32 (24-41)	1700 (1300-2200)
Croatia	2010	Female	50 to 54	65% (54-75%)	35% (25-46%)	2.5 (1.8-3.2)	5.6% (4.2-7.4%)	10% (7.7-13%)	55 (42-71)	2400 (1800-3100)
Croatia	2010	Female	55 to 59	66% (54-76%)	34% (24-46%)	2.6 (1.9-3.4)	5.9% (4.3-7.7%)	10% (7.6-13%)	80 (60-100)	2900 (2200-3800)
Croatia	2010	Female	60 to 64	64% (52-75%)	36% (25-48%)	2.5 (1.8-3.3)	5.3% (3.5-7.3%)	8.5% (5.9-12%)	91 (63-130)	2800 (1900-4000)
Croatia	2010	Female	65 to 69	59% (47-70%)	41% (30-53%)	2.2 (1.6-2.8)	4.7% (2.9-6.6%)	7% (4.4-9.7%)	100 (64-150)	2600 (1600-3700)
Croatia	2010	Female	70 to 74	50% (39-60%)	50% (40-61%)	1.7 (1.2-2.4)	3.5% (1.3-5.8%)	4.8% (1.8-7.8%)	140 (54-230)	2900 (1100-4700)
Croatia	2010	Female	75 to 79	40% (29-54%)	60% (46-71%)	1.3 (0.85-2)	2.5% (0.51-4.8%)	3.1% (0.58-6.1%)	160 (30-300)	2400 (490-4700)
Croatia	2010	Female	80 to 84	37% (24-51%)	63% (49-76%)	1.3 (0.79-1.9)	2.2% (0.32-5.5%)	2.5% (0.32-5.3%)	160 (20-340)	1800 (310-3700)
Croatia	2010	Female	85 to 89	37% (25-51%)	63% (49-75%)	1.4 (0.79-2.1)	2% (-0.04-4.4%)	2.2% (-0.27-5.3%)	120 (-15-280)	960 (-19-2200)
Croatia	2010	Female	90 to 94	39% (27-53%)	61% (47-73%)	1.5 (0.83-2.3)	1.8% (-0.32-4.7%)	2% (-0.82-5.8%)	39 (-16-110)	230 (-43-610)
Croatia	2010	Female	95 plus	42% (26-60%)	58% (40-74%)	1.6 (0.82-2.8)	2% (-1.6-2.2%)	2.2% (-1.6-7.5%)	14 (-10-49)	56 (-28-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	2010	Male	All Ages	78% (74-82%)	22% (18-26%)	5.4 (4.9-6)	16% (14-18%)	18% (15-22%)	4900 (4000-5900)	120000 (110000-150000)
Croatia	2010	Male	15 to 19	72% (58-83%)	28% (17-42%)	2.9 (2-4.1)	7.5% (5-10%)	14% (8.7-21%)	10 (6.1-15)	960 (650-1300)
Croatia	2010	Male	20 to 24	82% (71-90%)	18% (10-29%)	4.9 (3.4-6.6)	14% (10-19%)	24% (16-32%)	31 (20-42)	2600 (1900-3500)
Croatia	2010	Male	25 to 29	84% (76-90%)	16% (9.6-24%)	4.7 (3.4-6)	13% (9.6-17%)	22% (16-30%)	31 (22-42)	2700 (1900-3500)
Croatia	2010	Male	30 to 34	85% (77-91%)	15% (9-23%)	5.1 (3.7-6.7)	13% (9.7-16%)	23% (17-30%)	37 (27-49)	2900 (2200-3700)
Croatia	2010	Male	35 to 39	85% (78-91%)	15% (8.9-22%)	6 (4.3-8)	15% (11-18%)	26% (20-32%)	60 (45-75)	3900 (3000-4900)
Croatia	2010	Male	40 to 44	85% (77-91%)	15% (8.9-23%)	7.1 (5.3-9)	18% (15-21%)	29% (24-33%)	110 (89-130)	6000 (4800-7200)
Croatia	2010	Male	45 to 49	86% (78-91%)	14% (8.6-22%)	8.5 (6.4-11)	20% (17-24%)	29% (25-34%)	220 (180-260)	10000 (8500-12000)
Croatia	2010	Male	50 to 54	85% (77-91%)	15% (8.7-23%)	8.6 (6.5-11)	20% (17-24%)	27% (23-31%)	350 (300-420)	14000 (12000-17000)
Croatia	2010	Male	55 to 59	84% (73-91%)	16% (8.8-27%)	7.7 (6-9.8)	19% (16-23%)	24% (21-29%)	480 (400-570)	17000 (14000-20000)
Croatia	2010	Male	60 to 64	84% (74-91%)	16% (9.4-26%)	7.7 (5.8-10)	18% (15-21%)	22% (18-26%)	530 (430-640)	15000 (13000-19000)
Croatia	2010	Male	65 to 69	82% (73-89%)	18% (11-27%)	7.7 (6.2-9.4)	17% (14-21%)	21% (17-25%)	550 (450-670)	13000 (11000-16000)
Croatia	2010	Male	70 to 74	80% (70-87%)	20% (13-30%)	7.4 (5.5-9.5)	16% (13-20%)	19% (15-23%)	790 (620-980)	15000 (12000-19000)
Croatia	2010	Male	75 to 79	79% (67-87%)	21% (13-33%)	6.6 (4.6-8.9)	14% (11-18%)	16% (12-21%)	810 (590-1100)	12000 (8900-16000)
Croatia	2010	Male	80 to 84	75% (61-85%)	25% (15-39%)	5.4 (3.8-7.2)	12% (8.6-16%)	14% (9.6-19%)	570 (390-760)	6300 (4400-8300)
Croatia	2010	Male	85 to 89	72% (59-83%)	28% (17-41%)	4.5 (3.1-6.1)	9.4% (5.4-13%)	11% (5.9-15%)	230 (130-340)	1800 (1100-2700)
Croatia	2010	Male	90 to 94	71% (57-82%)	29% (18-43%)	3.8 (2.5-5.5)	6.3% (2.1-11%)	6.9% (1.8-12%)	46 (12-84)	260 (87-450)
Croatia	2010	Male	95 plus	70% (53-83%)	30% (17-47%)	3 (1.5-5)	3.9% (-0.67-9.8%)	3.9% (-1.5-11%)	7.1 (-2.9-20)	29 (-5.1-74)
Croatia	2016	Female	All Ages	61% (56-66%)	39% (34-44%)	1.6 (1.3-2.1)	3.4% (1.9-5.1%)	3.5% (0.97-6.6%)	1000 (280-1900)	23000 (13000-34000)
Croatia	2016	Female	15 to 19	65% (49-78%)	35% (22-51%)	1.1 (0.66-1.7)	2.2% (1.4-3.1%)	5.7% (3-9.2%)	1.3 (0.68-2.2)	210 (140-300)
Croatia	2016	Female	20 to 24	74% (60-84%)	26% (16-40%)	1.3 (0.84-2.1)	3.7% (2.7-4.9%)	6% (3.2-9.6%)	1.6 (0.82-2.6)	380 (250-540)
Croatia	2016	Female	25 to 29	74% (61-84%)	26% (16-39%)	1.3 (0.88-2)	3.6% (2.7-4.7%)	5.3% (2.9-8%)	1.8 (0.95-2.9)	460 (300-650)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	2016	Female	30 to 34	72% (59-81%)	28% (19-41%)	1.5 (0.96-2.2)	3.4% (2.5-4.5%)	6.1% (3.8-8.9%)	3.1 (1.8-4.7)	550 (370-770)
Croatia	2016	Female	35 to 39	68% (57-79%)	32% (21-43%)	1.8 (1.1-2.6)	3.6% (2.6-4.8%)	7.9% (5.1-11%)	6.8 (4.1-10)	700 (480-960)
Croatia	2016	Female	40 to 44	67% (55-78%)	33% (22-45%)	2 (1.3-2.8)	4.1% (2.9-5.4%)	9.4% (6.8-12%)	14 (9.3-20)	980 (670-1300)
Croatia	2016	Female	45 to 49	68% (55-78%)	32% (22-45%)	2.4 (1.6-3.3)	5% (3.6-6.7%)	10% (7.5-14%)	27 (19-37)	1400 (990-1900)
Croatia	2016	Female	50 to 54	68% (55-78%)	32% (22-45%)	2.6 (1.8-3.6)	5.5% (3.9-7.6%)	10% (7.4-13%)	48 (33-67)	2100 (1500-2900)
Croatia	2016	Female	55 to 59	68% (55-78%)	32% (22-45%)	2.6 (1.8-3.7)	5.6% (3.8-7.8%)	9.8% (6.9-13%)	73 (49-100)	2700 (1800-3800)
Croatia	2016	Female	60 to 64	66% (51-77%)	34% (23-49%)	2.5 (1.7-3.5)	4.9% (2.9-7.1%)	8.1% (5.2-12%)	88 (54-130)	2800 (1600-4100)
Croatia	2016	Female	65 to 69	61% (47-73%)	39% (27-53%)	2.2 (1.4-3.1)	4.4% (2.4-6.7%)	6.6% (3.9-9.8%)	110 (63-180)	2900 (1600-4500)
Croatia	2016	Female	70 to 74	54% (40-65%)	46% (35-60%)	1.9 (1.1-2.8)	3.4% (1.1-6.4%)	4.7% (1.7-8.6%)	110 (36-190)	2100 (730-3900)
Croatia	2016	Female	75 to 79	46% (32-60%)	54% (40-68%)	1.5 (0.89-2.4)	2.6% (0.23-5.5%)	3.3% (0.25-7.3%)	140 (11-310)	2300 (170-4900)
Croatia	2016	Female	80 to 84	40% (26-55%)	60% (45-74%)	1.4 (0.8-2.1)	2.2% (0.056-5%)	2.5% (-0.05-6%)	170 (-3.4-400)	2000 (56-4500)
Croatia	2016	Female	85 to 89	39% (26-53%)	61% (47-74%)	1.4 (0.77-2.2)	1.9% (-0.42-4.7%)	2.1% (-0.73-5.7%)	130 (-48-360)	1100 (-260-2800)
Croatia	2016	Female	90 to 94	41% (28-54%)	59% (46-72%)	1.5 (0.8-2.5)	1.9% (-0.83-5.2%)	1.9% (-1.3-6.3%)	59 (-40-200)	360 (-170-1100)
Croatia	2016	Female	95 plus	44% (28-61%)	56% (39-72%)	1.7 (0.8-2.9)	2% (-1.3-6.7%)	2.2% (-2-8.3%)	23 (-22-87)	90 (-59-300)
Croatia	2016	Male	All Ages	76% (70-80%)	24% (20-30%)	5 (4-6.2)	15% (12-17%)	16% (13-20%)	4400 (3300-5600)	110000 (87000-140000)
Croatia	2016	Male	15 to 19	66% (49-80%)	34% (20-51%)	2.5 (1.5-3.8)	6.4% (4-9.2%)	13% (7.3-19%)	7.6 (4.3-12)	770 (490-1100)
Croatia	2016	Male	20 to 24	78% (64-88%)	22% (12-36%)	4.3 (2.8-6.3)	13% (9-18%)	22% (14-31%)	21 (13-31)	1900 (1300-2700)
Croatia	2016	Male	25 to 29	81% (69-89%)	19% (11-31%)	4.2 (2.8-5.9)	12% (8.8-16%)	21% (14-29%)	23 (14-33)	2100 (1500-2800)
Croatia	2016	Male	30 to 34	82% (70-90%)	18% (10-30%)	4.6 (3.1-6.4)	12% (8.8-16%)	21% (15-29%)	30 (20-41)	2500 (1800-3300)
Croatia	2016	Male	35 to 39	82% (72-90%)	18% (10-28%)	5.4 (3.6-7.7)	13% (10-17%)	24% (18-31%)	49 (33-66)	3400 (2500-4400)
Croatia	2016	Male	40 to 44	83% (73-90%)	17% (9.7-27%)	6.5 (4.6-8.9)	16% (13-20%)	28% (22-33%)	93 (69-120)	5300 (4000-6600)
Croatia	2016	Male	45 to 49	84% (73-91%)	16% (9.2-27%)	7.9 (5.4-11)	19% (16-23%)	28% (24-33%)	160 (120-210)	7800 (6000-9800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Croatia	2016	Male	50 to 54	84% (73-91%)	16% (9-27%)	8 (5.5-11)	19% (16-23%)	26% (22-31%)	280 (220-360)	11000 (9000-14000)
Croatia	2016	Male	55 to 59	83% (71-91%)	17% (8.6-29%)	7.3 (5.1-9.8)	18% (15-22%)	24% (20-28%)	410 (310-510)	14000 (11000-18000)
Croatia	2016	Male	60 to 64	82% (70-90%)	18% (9.7-30%)	7.1 (4.9-9.7)	17% (13-20%)	21% (17-25%)	520 (390-650)	15000 (12000-19000)
Croatia	2016	Male	65 to 69	80% (69-89%)	20% (11-31%)	7.1 (5.2-9.5)	16% (13-19%)	20% (16-24%)	630 (480-790)	15000 (12000-19000)
Croatia	2016	Male	70 to 74	78% (66-86%)	22% (14-34%)	6.8 (4.8-9.5)	15% (11-18%)	18% (14-22%)	550 (410-710)	11000 (8100-14000)
Croatia	2016	Male	75 to 79	76% (62-86%)	24% (14-38%)	6 (3.9-8.6)	13% (9.2-17%)	15% (11-20%)	660 (450-880)	10000 (6900-13000)
Croatia	2016	Male	80 to 84	73% (58-85%)	27% (15-42%)	5 (3.2-7.2)	11% (6.9-16%)	13% (7.8-18%)	610 (370-850)	6700 (4100-9300)
Croatia	2016	Male	85 to 89	71% (55-82%)	29% (18-45%)	4.2 (2.7-6.3)	8.5% (4.2-13%)	9.7% (4.5-15%)	300 (140-470)	2400 (1200-3700)
Croatia	2016	Male	90 to 94	70% (55-82%)	30% (18-45%)	3.5 (2.1-5.5)	5.7% (1.2-11%)	6.3% (0.88-12%)	79 (11-160)	450 (99-840)
Croatia	2016	Male	95 plus	69% (50-84%)	31% (16-50%)	2.8 (1.4-5)	3.5% (-0.88-9.5%)	3.5% (-1.8-11%)	9.4 (-4.8-29)	39 (-9.6-110)
Czech Republic	1990	Female	All Ages	77% (74-80%)	23% (20-26%)	2.3 (2-2.5)	3.8% (2.1-5.8%)	3.7% (0.67-7.3%)	2300 (420-4600)	66000 (36000-100000)
Czech Republic	1990	Female	15 to 19	79% (69-88%)	21% (12-31%)	2 (1.4-2.7)	3.5% (2.3-5.1%)	9% (5-14%)	13 (7.4-21)	1400 (900-2100)
Czech Republic	1990	Female	20 to 24	88% (81-93%)	12% (7.5-19%)	2.3 (1.5-3.2)	4.6% (3.1-6.3%)	9.5% (5.1-14%)	12 (6.6-19)	1600 (1100-2300)
Czech Republic	1990	Female	25 to 29	89% (82-93%)	11% (6.8-18%)	2 (1.4-2.8)	4.1% (2.9-5.6%)	8.2% (4.4-13%)	11 (6.2-18)	1600 (1000-2200)
Czech Republic	1990	Female	30 to 34	89% (82-93%)	11% (7.1-18%)	2.3 (1.7-3.2)	4.3% (2.9-5.7%)	8.9% (5.4-13%)	21 (12-29)	2100 (1400-2800)
Czech Republic	1990	Female	35 to 39	87% (80-91%)	13% (8.5-20%)	3.1 (2.2-4.1)	5.1% (3.5-6.7%)	11% (7.4-15%)	45 (30-60)	3400 (2300-4500)
Czech Republic	1990	Female	40 to 44	85% (78-89%)	15% (11-22%)	3.5 (2.6-4.5)	5.9% (4.2-7.7%)	12% (8.4-15%)	85 (62-110)	5000 (3700-6600)
Czech Republic	1990	Female	45 to 49	83% (75-88%)	17% (12-25%)	3.6 (2.6-4.8)	6.3% (4.3-8.8%)	11% (7.7-15%)	110 (77-150)	5500 (3900-7500)
Czech Republic	1990	Female	50 to 54	82% (75-87%)	18% (13-25%)	3.7 (2.7-4.9)	6.6% (4-9.3%)	11% (6.7-15%)	130 (80-180)	5500 (3500-7700)
Czech Republic	1990	Female	55 to 59	82% (75-87%)	18% (13-25%)	3.8 (2.8-5)	6.8% (3.9-10%)	10% (5.7-15%)	220 (120-310)	7800 (4300-11000)
Czech Republic	1990	Female	60 to 64	80% (73-85%)	20% (15-27%)	3.5 (2.5-4.6)	5.9% (2.3-9.5%)	8.1% (3.3-13%)	300 (120-490)	9100 (3600-15000)
Czech Republic	1990	Female	65 to 69	74% (66-81%)	26% (19-34%)	2.9 (2.2-3.8)	4.5% (1.4-7.9%)	5.9% (1.8-10%)	380 (120-680)	9500 (3000-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	1990	Female	70 to 74	66% (57-74%)	34% (26-43%)	2.4 (1.6-3.3)	3% (-0.52-6.7%)	3.7% (-0.74-8.5%)	210 (-42-490)	4200 (-740-9500)
Czech Republic	1990	Female	75 to 79	57% (45-68%)	43% (32-55%)	1.9 (1.2-2.7)	1.8% (-1.5-5.7%)	2.2% (-2-6.8%)	270 (-250-830)	4100 (-3400-12000)
Czech Republic	1990	Female	80 to 84	52% (39-63%)	48% (37-61%)	1.8 (1.2-2.5)	1.7% (-1.5-5.1%)	1.9% (-1.5-5.9%)	270 (-210-820)	3000 (-2000-8900)
Czech Republic	1990	Female	85 to 89	51% (38-63%)	49% (37-62%)	1.8 (1.2-2.8)	1.6% (-1.7-5.4%)	1.8% (-2.3-6.1%)	160 (-210-570)	1300 (-1400-4500)
Czech Republic	1990	Female	90 to 94	53% (41-66%)	47% (34-59%)	2 (1.2-3)	1.7% (-2.1-6.5%)	1.8% (-2.8-7.6%)	64 (-97-270)	370 (-470-1400)
Czech Republic	1990	Female	95 plus	57% (41-72%)	43% (28-59%)	2.2 (1.1-3.5)	1.9% (-3.2-8%)	2.1% (-4-9.4%)	14 (-27-63)	54 (-88-220)
Czech Republic	1990	Male	All Ages	88% (85-90%)	12% (10-15%)	4.7 (4.1-5.1)	13% (11-16%)	15% (11-19%)	9800 (7200-12000)	270000 (220000-330000)
Czech Republic	1990	Male	15 to 19	85% (76-92%)	15% (7.9-24%)	2.3 (1.6-3.3)	6.5% (4-9.4%)	12% (6.4-18%)	39 (22-60)	3400 (2100-5000)
Czech Republic	1990	Male	20 to 24	92% (87-95%)	8.1% (4.8-13%)	4.1 (3-5.4)	12% (8.4-17%)	20% (12-27%)	82 (51-110)	6600 (4500-8900)
Czech Republic	1990	Male	25 to 29	94% (90-96%)	6.3% (3.8-10%)	5 (3.5-6.6)	14% (9.5-18%)	23% (15-31%)	97 (64-130)	7500 (5300-9800)
Czech Republic	1990	Male	30 to 34	95% (91-97%)	5.3% (2.7-9.3%)	5.5 (4-7.3)	14% (10-18%)	23% (16-30%)	130 (91-170)	9100 (6600-12000)
Czech Republic	1990	Male	35 to 39	95% (92-97%)	4.9% (2.5-8.3%)	5.6 (4-7.7)	14% (10-19%)	21% (15-28%)	220 (150-300)	14000 (9700-18000)
Czech Republic	1990	Male	40 to 44	94% (90-96%)	6.2% (3.9-9.6%)	6.5 (4.9-8.2)	16% (12-19%)	22% (16-27%)	390 (280-480)	20000 (15000-25000)
Czech Republic	1990	Male	45 to 49	91% (87-94%)	9% (5.9-13%)	8 (6-10)	18% (14-22%)	23% (18-28%)	570 (450-690)	26000 (20000-31000)
Czech Republic	1990	Male	50 to 54	90% (85-93%)	10% (7.1-15%)	8.5 (6.4-11)	17% (13-21%)	21% (16-26%)	630 (490-780)	25000 (19000-31000)
Czech Republic	1990	Male	55 to 59	90% (86-93%)	9.7% (6.9-14%)	8.5 (6.6-11)	16% (12-21%)	19% (14-25%)	910 (670-1200)	31000 (23000-39000)
Czech Republic	1990	Male	60 to 64	90% (87-93%)	9.6% (6.9-13%)	8.4 (6.4-11)	16% (11-21%)	18% (12-24%)	1300 (900-1700)	37000 (26000-49000)
Czech Republic	1990	Male	65 to 69	90% (86-93%)	10% (7.3-14%)	7.8 (6.1-9.4)	15% (11-20%)	17% (12-23%)	1700 (1200-2200)	40000 (28000-52000)
Czech Republic	1990	Male	70 to 74	88% (84-91%)	12% (9.1-16%)	6.7 (5.1-8.6)	14% (9-19%)	16% (10-21%)	1000 (660-1400)	19000 (13000-26000)
Czech Republic	1990	Male	75 to 79	84% (77-89%)	16% (11-23%)	5.5 (4-7.5)	11% (5.8-17%)	13% (6.3-19%)	1400 (680-2000)	20000 (10000-29000)
Czech Republic	1990	Male	80 to 84	81% (73-87%)	19% (13-27%)	4.8 (3.4-6.4)	9.8% (4.6-15%)	11% (4.8-17%)	940 (420-1500)	10000 (4900-16000)
Czech Republic	1990	Male	85 to 89	79% (70-85%)	21% (15-30%)	4.3 (3-5.8)	7.8% (2.3-13%)	8.5% (2.2-15%)	340 (89-600)	2700 (790-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	1990	Male	90 to 94	78% (69-85%)	22% (15-31%)	3.8 (2.6-5.4)	5.2% (-0.33-12%)	5.6% (-0.94-13%)	59 (-10-140)	330 (-20-750)
Czech Republic	1990	Male	95 plus	77% (64-86%)	23% (14-36%)	3.2 (1.9-5.2)	3.1% (-2.6-11%)	3.2% (-3.6-12%)	4.7 (-5.3-18)	19 (-16-66)
Czech Republic	1995	Female	All Ages	78% (74-81%)	22% (19-26%)	2.4 (2.1-2.7)	4% (2.3-6.1%)	4% (0.93-7.8%)	2300 (540-4500)	65000 (37000-97000)
Czech Republic	1995	Female	15 to 19	79% (69-87%)	21% (13-31%)	2 (1.4-2.8)	3.4% (2.2-4.9%)	9.3% (5.2-14%)	13 (7.1-20)	1400 (850-2000)
Czech Republic	1995	Female	20 to 24	87% (80-92%)	13% (7.7-20%)	2.3 (1.6-3.3)	4.5% (3.1-6.2%)	9.5% (5.3-14%)	14 (7.9-22)	1900 (1300-2700)
Czech Republic	1995	Female	25 to 29	89% (82-93%)	11% (6.8-18%)	2.1 (1.4-3)	4.1% (2.8-5.5%)	8.5% (5-13%)	11 (6.4-16)	1500 (1000-2100)
Czech Republic	1995	Female	30 to 34	89% (83-93%)	11% (7-17%)	2.4 (1.7-3.3)	4.1% (2.9-5.6%)	9.2% (5.9-13%)	17 (11-24)	1800 (1300-2500)
Czech Republic	1995	Female	35 to 39	87% (80-91%)	13% (8.6-20%)	3.2 (2.2-4.4)	5.1% (3.6-6.8%)	12% (8-15%)	37 (26-49)	2800 (2000-3700)
Czech Republic	1995	Female	40 to 44	84% (78-90%)	16% (10-22%)	3.6 (2.7-4.7)	6% (4.4-7.8%)	12% (9.3-16%)	77 (58-99)	4700 (3500-6200)
Czech Republic	1995	Female	45 to 49	83% (75-88%)	17% (12-25%)	3.8 (2.7-4.9)	6.3% (4.3-8.5%)	12% (8.4-15%)	120 (90-160)	6300 (4500-8400)
Czech Republic	1995	Female	50 to 54	82% (76-87%)	18% (13-24%)	3.9 (2.8-5)	6.7% (4.4-9.2%)	11% (7.5-15%)	150 (100-210)	6700 (4500-9000)
Czech Republic	1995	Female	55 to 59	82% (76-87%)	18% (13-24%)	3.9 (2.9-5.2)	6.9% (4.4-9.8%)	10% (6.5-15%)	180 (110-260)	6600 (4100-9400)
Czech Republic	1995	Female	60 to 64	80% (72-85%)	20% (15-28%)	3.6 (2.6-4.8)	6% (2.9-9.5%)	8.5% (4.1-14%)	250 (120-400)	7600 (3600-12000)
Czech Republic	1995	Female	65 to 69	74% (66-81%)	26% (19-34%)	3 (2.2-3.9)	4.7% (1.8-7.8%)	6.3% (2.2-11%)	320 (120-550)	8100 (3000-13000)
Czech Republic	1995	Female	70 to 74	66% (57-75%)	34% (25-43%)	2.5 (1.7-3.4)	3.2% (0.027-6.9%)	4.1% (-0.24-8.7%)	350 (-21-740)	6900 (58-15000)
Czech Republic	1995	Female	75 to 79	57% (45-69%)	43% (31-55%)	1.9 (1.2-2.9)	2.1% (-1.1-6.1%)	2.5% (-1.5-7.3%)	170 (-110-510)	2700 (-1400-7800)
Czech Republic	1995	Female	80 to 84	52% (39-63%)	48% (37-61%)	1.8 (1.2-2.7)	1.9% (-0.89-5.3%)	2.2% (-1.2-6.4%)	280 (-150-800)	3200 (-1500-9000)
Czech Republic	1995	Female	85 to 89	51% (38-63%)	49% (37-62%)	1.9 (1.2-2.8)	1.9% (-1.3-5.8%)	2.1% (-1.8-6.8%)	220 (-180-700)	1800 (-1200-5200)
Czech Republic	1995	Female	90 to 94	53% (41-66%)	47% (34-59%)	2.1 (1.2-3.1)	1.9% (-1.9-6.8%)	2.1% (-2.6-7.9%)	89 (-110-330)	510 (-510-1800)
Czech Republic	1995	Female	95 plus	57% (40-71%)	43% (29-60%)	2.2 (1.2-3.6)	2.1% (-3.1-8.4%)	2.3% (-3.9-9.9%)	20 (-34-88)	77 (-110-310)
Czech Republic	1995	Male	All Ages	88% (85-90%)	12% (9.8-15%)	4.9 (4.3-5.6)	14% (11-16%)	15% (11-20%)	9000 (6600-11000)	260000 (210000-310000)
Czech Republic	1995	Male	15 to 19	85% (75-92%)	15% (8.4-25%)	2.4 (1.6-3.3)	6.8% (4.1-10%)	12% (6.6-18%)	44 (24-67)	3800 (2300-5500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	1995	Male	20 to 24	92% (87-95%)	8.2% (4.9-13%)	4.1 (3-5.5)	12% (8.4-17%)	20% (13-28%)	95 (60-130)	7700 (5200-10000)
Czech Republic	1995	Male	25 to 29	94% (89-96%)	6.4% (3.8-11%)	5.1 (3.5-6.8)	14% (9.7-18%)	24% (16-32%)	99 (64-130)	7600 (5300-9900)
Czech Republic	1995	Male	30 to 34	95% (91-97%)	5.3% (2.7-9.1%)	5.6 (4-7.5)	14% (10-18%)	24% (17-31%)	120 (84-160)	8600 (6200-11000)
Czech Republic	1995	Male	35 to 39	95% (91-97%)	5% (2.7-8.6%)	5.7 (4-7.8)	14% (10-18%)	22% (15-29%)	180 (120-230)	11000 (7700-14000)
Czech Republic	1995	Male	40 to 44	94% (90-96%)	6.3% (3.8-10%)	6.6 (4.9-8.5)	16% (12-20%)	23% (17-28%)	340 (260-420)	18000 (14000-22000)
Czech Republic	1995	Male	45 to 49	91% (86-94%)	9.1% (6.2-14%)	8.1 (6.1-10)	18% (14-22%)	24% (19-29%)	620 (490-740)	28000 (23000-34000)
Czech Republic	1995	Male	50 to 54	90% (86-93%)	10% (7.3-14%)	8.6 (6.5-11)	18% (14-21%)	22% (18-27%)	730 (580-900)	29000 (23000-35000)
Czech Republic	1995	Male	55 to 59	90% (86-93%)	9.7% (7-14%)	8.7 (6.7-11)	17% (13-21%)	20% (15-26%)	760 (570-970)	26000 (20000-33000)
Czech Republic	1995	Male	60 to 64	90% (86-93%)	9.6% (6.8-14%)	8.5 (6.4-11)	15% (11-20%)	18% (13-24%)	1000 (730-1400)	30000 (22000-40000)
Czech Republic	1995	Male	65 to 69	90% (86-93%)	10% (7.3-14%)	7.8 (6.1-9.6)	15% (11-19%)	17% (12-23%)	1400 (950-1800)	33000 (23000-43000)
Czech Republic	1995	Male	70 to 74	88% (83-91%)	12% (9.1-17%)	6.8 (5.1-8.8)	13% (8.6-18%)	15% (9.6-21%)	1500 (950-2100)	29000 (19000-39000)
Czech Republic	1995	Male	75 to 79	84% (77-89%)	16% (11-23%)	5.5 (3.9-7.6)	11% (5.8-16%)	13% (6.2-18%)	750 (370-1100)	11000 (5800-16000)
Czech Republic	1995	Male	80 to 84	81% (73-86%)	19% (14-27%)	4.8 (3.4-6.4)	9.7% (4.7-15%)	11% (4.8-17%)	860 (390-1400)	9600 (4600-15000)
Czech Republic	1995	Male	85 to 89	79% (70-85%)	21% (15-30%)	4.4 (3.1-5.9)	7.8% (2.7-13%)	8.6% (2.6-15%)	410 (120-710)	3200 (1100-5500)
Czech Republic	1995	Male	90 to 94	78% (69-84%)	22% (16-31%)	3.9 (2.5-5.5)	5.3% (-0.41-12%)	5.7% (-1.1-13%)	74 (-14-170)	410 (-32-910)
Czech Republic	1995	Male	95 plus	76% (64-85%)	24% (15-36%)	3.3 (1.9-5.2)	3.2% (-2.5-11%)	3.2% (-3.7-12%)	6.1 (-7-23)	25 (-19-83)
Czech Republic	2000	Female	All Ages	79% (76-81%)	21% (19-24%)	2.6 (2.3-2.8)	4.2% (2.8-5.9%)	4.3% (1.6-7.5%)	2300 (860-4100)	64000 (42000-90000)
Czech Republic	2000	Female	15 to 19	82% (75-89%)	18% (11-25%)	2.2 (1.5-3)	3.4% (2.1-4.9%)	10% (5.4-15%)	9.7 (5.3-15)	1100 (670-1500)
Czech Republic	2000	Female	20 to 24	89% (83-93%)	11% (7.3-17%)	2.5 (1.7-3.4)	4.4% (3.1-5.9%)	9.8% (5.6-15%)	13 (7.6-20)	1900 (1200-2600)
Czech Republic	2000	Female	25 to 29	90% (85-93%)	10% (6.7-15%)	2.2 (1.5-3)	3.9% (2.9-5.1%)	8.7% (5.3-13%)	12 (7-17)	1800 (1200-2400)
Czech Republic	2000	Female	30 to 34	90% (85-93%)	10% (6.9-15%)	2.5 (1.9-3.4)	4.1% (2.9-5.3%)	9.9% (6.8-13%)	15 (10-20)	1700 (1200-2200)
Czech Republic	2000	Female	35 to 39	88% (83-92%)	12% (8.1-17%)	3.4 (2.4-4.5)	5.1% (3.8-6.6%)	12% (9.2-16%)	31 (23-40)	2500 (1800-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	2000	Female	40 to 44	86% (82-90%)	14% (10-18%)	3.9 (2.9-4.9)	6.2% (4.8-8%)	14% (11-17%)	64 (51-79)	4000 (3100-5000)
Czech Republic	2000	Female	45 to 49	84% (80-88%)	16% (12-20%)	4 (3-5.1)	6.7% (4.8-8.9%)	13% (9.6-16%)	120 (91-150)	6200 (4600-8000)
Czech Republic	2000	Female	50 to 54	83% (78-87%)	17% (13-22%)	4 (3-5.1)	7% (4.9-9.3%)	12% (8.7-15%)	180 (130-240)	8000 (5700-10000)
Czech Republic	2000	Female	55 to 59	82% (78-86%)	18% (14-22%)	4.1 (3.1-5.2)	7.1% (4.8-9.6%)	11% (7.6-15%)	210 (150-290)	8000 (5400-11000)
Czech Republic	2000	Female	60 to 64	78% (73-83%)	22% (17-27%)	3.5 (2.6-4.7)	6% (3.3-8.9%)	8.8% (4.9-13%)	210 (110-310)	6500 (3500-9600)
Czech Republic	2000	Female	65 to 69	71% (63-79%)	29% (21-37%)	3 (2.1-3.8)	4.8% (2.4-7.4%)	6.6% (3.2-10%)	260 (130-410)	6800 (3300-11000)
Czech Republic	2000	Female	70 to 74	63% (54-72%)	37% (28-46%)	2.3 (1.6-3.2)	3.4% (0.57-6.6%)	4.4% (0.7-8.5%)	300 (49-590)	6100 (1100-12000)
Czech Republic	2000	Female	75 to 79	53% (40-64%)	47% (36-60%)	1.8 (1.1-2.7)	2.3% (-0.42-5.5%)	2.7% (-0.66-6.8%)	300 (-70-730)	4600 (-870-11000)
Czech Republic	2000	Female	80 to 84	49% (37-61%)	51% (39-63%)	1.7 (1.2-2.6)	2.1% (-0.46-5.1%)	2.5% (-0.7-6.2%)	190 (-54-470)	2200 (-460-5300)
Czech Republic	2000	Female	85 to 89	50% (37-62%)	50% (38-63%)	1.9 (1.2-2.8)	2.2% (-0.89-5.9%)	2.5% (-1.3-7%)	260 (-140-720)	2100 (-820-5600)
Czech Republic	2000	Female	90 to 94	52% (40-65%)	48% (35-60%)	2 (1.2-3.1)	2.2% (-1.5-6.9%)	2.4% (-2-8.3%)	130 (-110-430)	720 (-480-2300)
Czech Republic	2000	Female	95 plus	56% (39-71%)	44% (29-61%)	2.2 (1.2-3.6)	2.3% (-2.9-8.6%)	2.5% (-3.6-9.9%)	29 (-42-110)	110 (-130-400)
Czech Republic	2000	Male	All Ages	90% (88-92%)	10% (8.4-12%)	5.4 (4.9-5.9)	14% (12-17%)	16% (13-20%)	8800 (6800-11000)	25000 (21000-30000)
Czech Republic	2000	Male	15 to 19	88% (80-93%)	12% (7.1-20%)	2.6 (1.8-3.6)	6.7% (4.1-9.8%)	13% (7.3-20%)	31 (17-46)	2700 (1700-3900)
Czech Republic	2000	Male	20 to 24	93% (90-96%)	6.6% (4.3-9.7%)	4.4 (3.1-5.9)	12% (8.6-17%)	21% (13-28%)	95 (60-130)	7800 (5300-10000)
Czech Republic	2000	Male	25 to 29	95% (92-97%)	5.2% (3.4-7.7%)	5.3 (3.8-7.1)	13% (9.6-17%)	25% (17-33%)	110 (73-140)	8400 (6100-11000)
Czech Republic	2000	Male	30 to 34	96% (93-98%)	4.3% (2.5-7.1%)	5.8 (4.3-7.7)	14% (10-18%)	25% (18-32%)	110 (78-140)	7800 (5800-9900)
Czech Republic	2000	Male	35 to 39	96% (94-98%)	4.1% (2.3-6.4%)	5.9 (4.3-8.1)	14% (10-18%)	24% (17-31%)	150 (100-190)	9400 (6900-12000)
Czech Republic	2000	Male	40 to 44	95% (92-97%)	5.2% (3.4-7.8%)	6.8 (5.2-8.8)	16% (13-20%)	25% (19-30%)	280 (220-340)	15000 (12000-18000)
Czech Republic	2000	Male	45 to 49	92% (89-95%)	7.8% (5.4-11%)	8.5 (6.5-11)	19% (15-22%)	26% (22-30%)	570 (470-660)	26000 (21000-30000)
Czech Republic	2000	Male	50 to 54	91% (88-93%)	9.1% (6.7-12%)	9 (7-11)	18% (15-22%)	24% (20-28%)	840 (690-1000)	34000 (28000-40000)
Czech Republic	2000	Male	55 to 59	92% (89-93%)	8.5% (6.6-11%)	9 (7.2-11)	17% (14-21%)	21% (17-26%)	910 (710-1100)	31000 (25000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	2000	Male	60 to 64	92% (90-94%)	8.2% (6.4-10%)	8.9 (6.9-11)	16% (12-20%)	19% (14-24%)	900 (670-1100)	26000 (20000-33000)
Czech Republic	2000	Male	65 to 69	91% (89-93%)	8.5% (6.6-11%)	8.2 (6.7-9.9)	15% (11-19%)	18% (13-23%)	1200 (850-1500)	28000 (21000-35000)
Czech Republic	2000	Male	70 to 74	90% (87-92%)	10% (8.3-13%)	7.2 (5.6-9)	14% (9.2-18%)	16% (11-21%)	1300 (890-1700)	26000 (18000-34000)
Czech Republic	2000	Male	75 to 79	86% (80-90%)	14% (10-20%)	5.9 (4.3-7.8)	11% (6.5-16%)	13% (7.1-19%)	1200 (650-1700)	18000 (10000-26000)
Czech Republic	2000	Male	80 to 84	82% (75-87%)	18% (13-25%)	5 (3.6-6.6)	10% (5.1-15%)	11% (5.4-17%)	550 (270-840)	6200 (3100-9200)
Czech Republic	2000	Male	85 to 89	80% (72-85%)	20% (15-28%)	4.5 (3.3-6)	8.2% (3.3-14%)	9.1% (3.3-15%)	450 (160-760)	3600 (1400-5900)
Czech Republic	2000	Male	90 to 94	78% (70-85%)	22% (15-30%)	4 (2.7-5.6)	5.8% (0.12-12%)	6.3% (-0.49-13%)	120 (-9.3-250)	660 (14-1400)
Czech Republic	2000	Male	95 plus	77% (66-86%)	23% (14-34%)	3.4 (2-5.3)	3.6% (-2.4-11%)	3.7% (-3.3-12%)	9.9 (-8.8-32)	40 (-25-120)
Czech Republic	2005	Female	All Ages	79% (77-81%)	21% (19-23%)	2.6 (2.4-2.9)	4.2% (2.8-5.6%)	4.2% (1.6-7%)	2200 (840-3700)	64000 (43000-85000)
Czech Republic	2005	Female	15 to 19	83% (75-89%)	17% (11-25%)	2.2 (1.5-3)	3.2% (2-4.5%)	9.9% (5.6-15%)	7.8 (4.4-12)	940 (600-1300)
Czech Republic	2005	Female	20 to 24	89% (83-93%)	11% (6.9-17%)	2.5 (1.7-3.6)	4.3% (3-5.7%)	10% (5.9-15%)	9.3 (5.4-14)	1400 (980-2000)
Czech Republic	2005	Female	25 to 29	90% (86-94%)	9.8% (6.4-14%)	2.2 (1.6-3.1)	3.9% (2.8-5.1%)	9% (5.7-13%)	11 (7.2-16)	1800 (1300-2500)
Czech Republic	2005	Female	30 to 34	90% (86-93%)	9.9% (6.7-14%)	2.6 (2-3.5)	4.1% (3-5.4%)	10% (7.2-14%)	18 (13-24)	2100 (1500-2800)
Czech Republic	2005	Female	35 to 39	88% (84-92%)	12% (8.1-16%)	3.4 (2.5-4.6)	5.1% (3.9-6.5%)	13% (9.8-16%)	31 (23-39)	2500 (1900-3200)
Czech Republic	2005	Female	40 to 44	86% (82-90%)	14% (10-18%)	3.9 (3-5)	6.3% (4.9-7.9%)	14% (11-17%)	58 (47-71)	3700 (3000-4700)
Czech Republic	2005	Female	45 to 49	84% (80-88%)	16% (12-20%)	4.1 (3-5.2)	7% (5.2-9.1%)	14% (11-17%)	100 (81-130)	5400 (4100-6800)
Czech Republic	2005	Female	50 to 54	83% (79-86%)	17% (14-21%)	4.1 (3.1-5.3)	7.3% (5.3-9.5%)	13% (9.8-16%)	170 (130-220)	7700 (5800-9900)
Czech Republic	2005	Female	55 to 59	83% (79-86%)	17% (14-21%)	4.2 (3.2-5.2)	7.2% (5-9.4%)	11% (8.4-15%)	250 (180-330)	9500 (6800-13000)
Czech Republic	2005	Female	60 to 64	79% (75-83%)	21% (17-25%)	3.7 (2.8-4.7)	6.2% (3.7-8.8%)	9.3% (5.8-13%)	250 (160-360)	8100 (4900-12000)
Czech Republic	2005	Female	65 to 69	71% (65-77%)	29% (23-35%)	3 (2.3-3.7)	4.7% (2.5-7%)	6.7% (3.7-10%)	220 (120-330)	5800 (3200-8700)
Czech Republic	2005	Female	70 to 74	60% (53-66%)	40% (34-47%)	2.2 (1.6-3)	3.2% (0.73-5.7%)	4.2% (1.1-7.5%)	230 (60-410)	4800 (1100-8600)
Czech Republic	2005	Female	75 to 79	51% (42-61%)	49% (39-58%)	1.8 (1.2-2.5)	2.2% (-0.19-5.1%)	2.7% (-0.36-6.6%)	250 (-33-590)	4000 (-330-9300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	2005	Female	80 to 84	48% (36-58%)	52% (42-64%)	1.7 (1.1-2.4)	2% (-0.37-4.7%)	2.4% (-0.64-5.8%)	300 (-83-730)	3600 (-670-8200)
Czech Republic	2005	Female	85 to 89	47% (35-58%)	53% (42-65%)	1.8 (1.2-2.6)	2% (-0.75-5.1%)	2.2% (-1.2-6.1%)	150 (-79-420)	1300 (-470-3400)
Czech Republic	2005	Female	90 to 94	49% (38-61%)	51% (39-62%)	1.9 (1.2-2.9)	2% (-1.4-6.1%)	2.2% (-1.9-7.2%)	120 (-110-410)	710 (-510-2200)
Czech Republic	2005	Female	95 plus	53% (37-69%)	47% (31-63%)	2.1 (1.1-3.3)	2.1% (-2.7-7.6%)	2.3% (-3.4-8.9%)	37 (-55-140)	140 (-170-500)
Czech Republic	2005	Male	All Ages	91% (89-92%)	9.2% (7.9-11%)	5.6 (5.2-6)	14% (12-16%)	16% (13-20%)	8600 (6700-11000)	250000 (210000-290000)
Czech Republic	2005	Male	15 to 19	88% (82-93%)	12% (6.9-18%)	2.6 (1.8-3.6)	6.4% (3.9-9%)	13% (7.5-20%)	25 (14-38)	2300 (1400-3300)
Czech Republic	2005	Male	20 to 24	94% (91-96%)	6.2% (4.1-9.3%)	4.4 (3.2-5.9)	12% (8.6-16%)	21% (14-29%)	74 (47-100)	6200 (4300-8100)
Czech Republic	2005	Male	25 to 29	95% (93-97%)	4.9% (3.2-7.2%)	5.4 (3.8-6.9)	13% (9.4-17%)	25% (17-32%)	100 (71-140)	8200 (6100-11000)
Czech Republic	2005	Male	30 to 34	96% (93-98%)	4.1% (2.4-6.6%)	5.9 (4.3-7.5)	13% (10-17%)	25% (18-32%)	120 (89-160)	9000 (6900-11000)
Czech Republic	2005	Male	35 to 39	96% (94-98%)	3.9% (2.2-6.1%)	5.9 (4.3-7.9)	14% (10-18%)	25% (17-32%)	140 (99-180)	9100 (6700-11000)
Czech Republic	2005	Male	40 to 44	95% (93-97%)	4.9% (3.3-7.2%)	6.9 (5.3-8.6)	16% (13-20%)	26% (21-32%)	240 (190-290)	13000 (10000-15000)
Czech Republic	2005	Male	45 to 49	93% (90-95%)	7.2% (5.1-9.7%)	8.5 (6.6-11)	19% (15-22%)	28% (23-32%)	460 (380-540)	21000 (18000-25000)
Czech Republic	2005	Male	50 to 54	92% (90-94%)	8.3% (6.2-10%)	9.1 (7.2-11)	19% (16-22%)	25% (21-29%)	780 (660-920)	32000 (27000-37000)
Czech Republic	2005	Male	55 to 59	92% (91-94%)	7.8% (6.3-9.4%)	9.2 (7.4-11)	18% (14-21%)	22% (18-27%)	1100 (880-1300)	38000 (31000-45000)
Czech Republic	2005	Male	60 to 64	92% (91-94%)	7.7% (6.2-9.4%)	9 (7-11)	16% (12-20%)	19% (15-24%)	1100 (830-1300)	32000 (25000-39000)
Czech Republic	2005	Male	65 to 69	92% (90-94%)	8.2% (6.5-10%)	8.3 (6.9-9.9)	15% (11-18%)	18% (13-22%)	980 (750-1200)	24000 (19000-30000)
Czech Republic	2005	Male	70 to 74	90% (88-92%)	9.9% (8-12%)	7.3 (5.7-9.1)	13% (9-17%)	15% (10-20%)	1100 (750-1500)	22000 (15000-29000)
Czech Republic	2005	Male	75 to 79	86% (82-90%)	14% (10-18%)	6 (4.4-7.8)	11% (6.1-16%)	13% (6.5-18%)	1100 (560-1500)	17000 (9400-23000)
Czech Republic	2005	Male	80 to 84	81% (75-87%)	19% (13-25%)	5 (3.7-6.5)	9.4% (4.8-14%)	11% (5.1-16%)	880 (410-1300)	9900 (5000-15000)
Czech Republic	2005	Male	85 to 89	79% (71-85%)	21% (15-29%)	4.5 (3.2-5.9)	7.7% (2.9-13%)	8.6% (2.9-14%)	290 (98-490)	2400 (890-3900)
Czech Republic	2005	Male	90 to 94	78% (69-84%)	22% (16-31%)	4 (2.7-5.6)	5.4% (0.067-12%)	5.8% (-0.55-13%)	120 (-11-270)	690 (8.1-1500)
Czech Republic	2005	Male	95 plus	76% (66-85%)	24% (15-34%)	3.4 (1.9-5.2)	3.5% (-2.3-11%)	3.5% (-3.3-12%)	15 (-14-51)	60 (-39-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	2010	Female	All Ages	78% (76-81%)	22% (19-24%)	2.6 (2.3-2.8)	3.8% (2.5-5.1%)	3.4% (0.75-6.2%)	1800 (390-3300)	57000 (36000-78000)
Czech Republic	2010	Female	15 to 19	81% (72-88%)	19% (12-28%)	2 (1.4-2.8)	2.9% (1.9-4.1%)	8.8% (5-14%)	5.6 (3.2-8.6)	760 (490-1100)
Czech Republic	2010	Female	20 to 24	89% (82-93%)	11% (7.2-18%)	2.4 (1.6-3.4)	4.2% (3-5.7%)	9.2% (5.3-13%)	8.2 (4.7-12)	1400 (920-1900)
Czech Republic	2010	Female	25 to 29	90% (85-93%)	10% (6.6-15%)	2.1 (1.5-2.9)	3.9% (2.9-5.1%)	8.6% (5.6-12%)	7.9 (5.2-11)	1500 (1000-2000)
Czech Republic	2010	Female	30 to 34	89% (85-93%)	11% (7-15%)	2.5 (1.8-3.3)	4% (2.9-5.2%)	9.8% (7.1-13%)	17 (12-22)	2100 (1500-2800)
Czech Republic	2010	Female	35 to 39	88% (82-92%)	12% (8.2-18%)	3.3 (2.4-4.4)	4.7% (3.5-6%)	12% (9.4-15%)	31 (24-38)	2800 (2100-3600)
Czech Republic	2010	Female	40 to 44	85% (80-89%)	15% (11-20%)	3.7 (2.8-4.7)	5.8% (4.5-7.4%)	14% (11-16%)	51 (41-61)	3400 (2700-4300)
Czech Republic	2010	Female	45 to 49	83% (79-87%)	17% (13-21%)	3.8 (2.8-5)	6.4% (4.7-8.3%)	13% (11-17%)	84 (67-110)	4600 (3500-5900)
Czech Republic	2010	Female	50 to 54	83% (79-87%)	17% (13-21%)	4 (2.9-5.1)	6.8% (5-8.9%)	13% (9.6-16%)	130 (100-160)	6000 (4500-7800)
Czech Republic	2010	Female	55 to 59	83% (79-86%)	17% (14-21%)	4 (3.1-5)	6.6% (4.6-8.6%)	11% (8.1-14%)	210 (150-270)	8100 (5800-11000)
Czech Republic	2010	Female	60 to 64	80% (76-84%)	20% (16-24%)	3.6 (2.7-4.6)	5.6% (3.4-7.9%)	8.5% (5.5-12%)	260 (170-360)	8500 (5300-12000)
Czech Republic	2010	Female	65 to 69	73% (68-79%)	27% (21-32%)	3 (2.3-3.7)	4.3% (2.3-6.3%)	6.1% (3.4-9%)	230 (130-340)	6300 (3400-9300)
Czech Republic	2010	Female	70 to 74	63% (56-69%)	37% (31-44%)	2.3 (1.7-3.1)	2.8% (0.51-5.1%)	3.8% (0.72-6.9%)	170 (32-310)	3700 (720-7000)
Czech Republic	2010	Female	75 to 79	55% (46-63%)	45% (37-54%)	1.8 (1.2-2.6)	1.8% (-0.46-4.6%)	2.2% (-0.71-5.7%)	170 (-53-430)	2800 (-740-7100)
Czech Republic	2010	Female	80 to 84	50% (39-60%)	50% (40-61%)	1.7 (1.2-2.4)	1.5% (-0.84-4.1%)	1.7% (-1.2-4.9%)	200 (-140-550)	2500 (-1300-6600)
Czech Republic	2010	Female	85 to 89	49% (37-60%)	51% (40-63%)	1.8 (1.2-2.6)	1.4% (-1.3-4.5%)	1.5% (-1.9-5.3%)	180 (-230-640)	1600 (-1500-5100)
Czech Republic	2010	Female	90 to 94	51% (39-62%)	49% (38-61%)	1.9 (1.2-2.9)	1.3% (-2-5.3%)	1.3% (-2.8-6.4%)	55 (-120-270)	360 (-550-1500)
Czech Republic	2010	Female	95 plus	55% (39-71%)	45% (29-61%)	2.1 (1.1-3.4)	1.4% (-3.2-7%)	1.4% (-4.1-8.1%)	27 (-82-160)	120 (-260-580)
Czech Republic	2010	Male	All Ages	91% (89-92%)	9.5% (8.1-11%)	5.4 (5-5.8)	13% (11-15%)	14% (11-18%)	7900 (6100-9800)	230000 (190000-270000)
Czech Republic	2010	Male	15 to 19	87% (80-93%)	13% (7.4-20%)	2.4 (1.7-3.4)	5.7% (3.5-8%)	12% (6.9-19%)	18 (10-27)	1800 (1100-2500)
Czech Republic	2010	Male	20 to 24	93% (90-96%)	6.6% (4.4-9.9%)	4.1 (3-5.5)	11% (7.9-15%)	20% (13-28%)	59 (37-80)	5100 (3500-6800)
Czech Republic	2010	Male	25 to 29	95% (92-97%)	5.1% (3.4-7.6%)	5.1 (3.5-6.6)	12% (9.1-16%)	24% (16-32%)	76 (52-100)	6300 (4700-8200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	2010	Male	30 to 34	96% (93-97%)	4.3% (2.5-6.7%)	5.6 (4.1-7.1)	12% (9.3-16%)	24% (17-31%)	110 (76-140)	8400 (6200-11000)
Czech Republic	2010	Male	35 to 39	96% (94-98%)	4.1% (2.3-6.3%)	5.6 (4.1-7.5)	13% (9.4-16%)	24% (17-30%)	150 (100-190)	9900 (7300-12000)
Czech Republic	2010	Male	40 to 44	95% (93-97%)	5% (3.4-7.2%)	6.5 (5-8.2)	15% (12-18%)	26% (20-31%)	210 (170-250)	12000 (9600-14000)
Czech Republic	2010	Male	45 to 49	93% (91-95%)	7.2% (5.2-9.4%)	8.1 (6.2-10)	17% (14-21%)	27% (23-32%)	360 (300-420)	18000 (15000-21000)
Czech Republic	2010	Male	50 to 54	92% (90-94%)	8.2% (6.4-10%)	8.7 (6.9-11)	18% (15-21%)	25% (21-29%)	600 (510-700)	25000 (21000-29000)
Czech Republic	2010	Male	55 to 59	92% (91-94%)	7.7% (6.3-9.2%)	8.7 (7.1-11)	17% (14-20%)	22% (19-26%)	950 (790-1100)	33000 (28000-39000)
Czech Republic	2010	Male	60 to 64	92% (90-94%)	7.8% (6.3-9.6%)	8.5 (6.7-10)	15% (12-19%)	19% (15-23%)	1200 (950-1500)	36000 (29000-44000)
Czech Republic	2010	Male	65 to 69	91% (89-93%)	8.6% (6.8-11%)	7.8 (6.5-9.2)	14% (11-17%)	17% (13-21%)	1100 (850-1400)	28000 (22000-34000)
Czech Republic	2010	Male	70 to 74	90% (88-92%)	10% (8.2-12%)	6.9 (5.4-8.5)	12% (8.2-16%)	14% (9.4-19%)	880 (580-1200)	18000 (12000-24000)
Czech Republic	2010	Male	75 to 79	86% (82-90%)	14% (10-18%)	5.7 (4.2-7.4)	9.8% (5.1-14%)	11% (5.5-16%)	870 (430-1300)	14000 (7200-20000)
Czech Republic	2010	Male	80 to 84	81% (75-86%)	19% (14-25%)	4.7 (3.5-6)	8% (3.7-12%)	9.1% (3.9-15%)	740 (310-1200)	8700 (4000-13000)
Czech Republic	2010	Male	85 to 89	78% (71-84%)	22% (16-29%)	4.2 (3.1-5.5)	6.3% (1.9-11%)	6.9% (1.7-12%)	420 (100-750)	3500 (1100-6000)
Czech Republic	2010	Male	90 to 94	77% (68-84%)	23% (16-32%)	3.7 (2.5-5.1)	4.1% (-0.59-9.6%)	4.3% (-1.6-11%)	69 (-25-180)	410 (-60-960)
Czech Republic	2010	Male	95 plus	76% (64-85%)	24% (15-36%)	3.2 (1.9-4.9)	2.4% (-2.6-9.1%)	2.2% (-3.8-10%)	12 (-20-54)	55 (-56-200)
Czech Republic	2016	Female	All Ages	78% (76-81%)	22% (19-24%)	2.5 (2-2.9)	3.2% (1.8-4.9%)	2.7% (-0.087-5.8%)	1500 (-49-3200)	48000 (26000-71000)
Czech Republic	2016	Female	15 to 19	80% (70-88%)	20% (12-30%)	1.9 (1.2-2.7)	2.6% (1.7-3.7%)	8.5% (4.9-13%)	3.2 (1.8-4.9)	480 (310-700)
Czech Republic	2016	Female	20 to 24	88% (81-93%)	12% (7.5-19%)	2.2 (1.4-3.3)	3.9% (2.7-5.2%)	8.5% (4.7-13%)	5.4 (3-8.2)	1100 (710-1600)
Czech Republic	2016	Female	25 to 29	89% (83-93%)	11% (6.6-17%)	2 (1.3-2.9)	3.8% (2.8-4.9%)	7.9% (4.9-12%)	6.5 (4-9.5)	1300 (890-1800)
Czech Republic	2016	Female	30 to 34	89% (82-93%)	11% (6.9-18%)	2.3 (1.6-3.2)	3.8% (2.8-4.9%)	9.4% (6.6-13%)	11 (7.4-14)	1600 (1100-2100)
Czech Republic	2016	Female	35 to 39	87% (80-92%)	13% (8.3-20%)	3 (2.1-4.2)	4.2% (3.1-5.5%)	11% (8.5-15%)	27 (20-35)	2600 (1900-3500)
Czech Republic	2016	Female	40 to 44	85% (80-90%)	15% (10-20%)	3.5 (2.5-4.7)	4.9% (3.6-6.5%)	12% (9.7-15%)	48 (37-61)	3500 (2600-4600)
Czech Republic	2016	Female	45 to 49	84% (78-88%)	16% (12-22%)	3.6 (2.5-4.9)	5.6% (4-7.7%)	13% (9.6-16%)	68 (51-88)	3900 (2800-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	2016	Female	50 to 54	83% (78-87%)	17% (13-22%)	3.7 (2.7-4.9)	5.9% (4.2-8%)	12% (9-15%)	100 (76-130)	4800 (3400-6500)
Czech Republic	2016	Female	55 to 59	84% (81-88%)	16% (12-19%)	3.9 (2.9-5.1)	5.9% (4.1-8.2%)	10% (7.7-14%)	150 (110-190)	5900 (3900-8000)
Czech Republic	2016	Female	60 to 64	82% (79-86%)	18% (14-21%)	3.6 (2.6-4.9)	5% (2.9-7.6%)	8.1% (5-12%)	210 (130-300)	7000 (4000-10000)
Czech Republic	2016	Female	65 to 69	77% (72-81%)	23% (19-28%)	3.1 (2.3-4)	3.8% (1.9-6%)	5.7% (2.9-8.7%)	250 (130-380)	6700 (3300-11000)
Czech Republic	2016	Female	70 to 74	69% (63-74%)	31% (26-37%)	2.5 (1.7-3.3)	2.4% (0.2-4.7%)	3.5% (0.33-6.9%)	180 (16-370)	4000 (340-8100)
Czech Republic	2016	Female	75 to 79	59% (51-68%)	41% (32-49%)	1.9 (1.3-2.7)	1.4% (-0.83-4%)	1.8% (-1.2-5.2%)	110 (-71-310)	1900 (-1100-5500)
Czech Republic	2016	Female	80 to 84	53% (42-64%)	47% (36-58%)	1.7 (1.2-2.5)	1.1% (-1.2-3.8%)	1.2% (-1.7-4.6%)	120 (-170-450)	1600 (-1700-5500)
Czech Republic	2016	Female	85 to 89	53% (40-64%)	47% (36-60%)	1.8 (1.2-2.7)	0.92% (-1.8-4.4%)	0.88% (-2.5-5.3%)	100 (-310-620)	1000 (-2100-4900)
Czech Republic	2016	Female	90 to 94	55% (43-66%)	45% (34-57%)	2 (1.2-3)	0.79% (-2.7-5.1%)	0.66% (-3.6-6%)	59 (-330-540)	450 (-1500-2900)
Czech Republic	2016	Female	95 plus	59% (43-72%)	41% (28-57%)	2.1 (1.1-3.4)	0.8% (-3.8-6.7%)	0.61% (-5-7.9%)	9 (-76-120)	49 (-230-410)
Czech Republic	2016	Male	All Ages	89% (87-91%)	11% (9-13%)	5 (4.2-5.9)	12% (9-14%)	13% (8.7-17%)	6800 (4600-9000)	20000 (15000-24000)
Czech Republic	2016	Male	15 to 19	86% (77-92%)	14% (7.7-23%)	2.2 (1.4-3.2)	5.1% (3.1-7.3%)	12% (6.2-17%)	11 (6-17)	1100 (710-1600)
Czech Republic	2016	Male	20 to 24	92% (87-95%)	7.9% (4.8-13%)	3.7 (2.6-5.2)	9.9% (6.8-13%)	9.9% (12-27%)	39 (24-56)	3700 (2500-5000)
Czech Republic	2016	Male	25 to 29	94% (90-96%)	6.1% (3.8-10%)	4.6 (3.1-6.2)	11% (7.9-14%)	22% (14-30%)	54 (34-75)	4900 (3400-6500)
Czech Republic	2016	Male	30 to 34	95% (91-97%)	5.2% (2.9-9%)	5.1 (3.5-6.8)	11% (8.2-14%)	22% (16-30%)	68 (46-92)	5700 (4100-7400)
Czech Republic	2016	Male	35 to 39	95% (92-97%)	4.9% (2.7-8.2%)	5.1 (3.6-7.1)	11% (8.2-15%)	22% (15-29%)	120 (83-160)	8700 (6300-11000)
Czech Republic	2016	Male	40 to 44	94% (90-96%)	6.2% (3.8-9.8%)	5.9 (4.3-7.8)	13% (9.8-16%)	24% (18-30%)	200 (150-250)	12000 (8900-15000)
Czech Republic	2016	Male	45 to 49	91% (87-94%)	8.5% (5.8-13%)	7.3 (5.3-9.8)	16% (12-19%)	26% (20-30%)	290 (220-350)	14000 (11000-18000)
Czech Republic	2016	Male	50 to 54	91% (89-93%)	8.9% (6.8-11%)	7.9 (5.8-10)	16% (13-20%)	25% (20-29%)	460 (360-540)	19000 (16000-23000)
Czech Republic	2016	Male	55 to 59	92% (90-94%)	8.1% (6.4-10%)	8 (6-10)	16% (13-19%)	22% (17-26%)	660 (520-790)	24000 (19000-28000)
Czech Republic	2016	Male	60 to 64	92% (90-94%)	8% (6.4-10%)	7.9 (5.9-10)	14% (11-18%)	18% (14-22%)	970 (710-1200)	30000 (22000-37000)
Czech Republic	2016	Male	65 to 69	91% (89-93%)	8.6% (6.8-11%)	7.3 (5.7-9)	13% (8.8-16%)	16% (11-20%)	1200 (840-1500)	31000 (22000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Czech Republic	2016	Male	70 to 74	90% (87-92%)	10% (8.4-13%)	6.4 (4.7-8.3)	10% (6.5-14%)	13% (7.5-18%)	980 (580-1400)	20000 (13000-28000)
Czech Republic	2016	Male	75 to 79	86% (82-90%)	14% (10-18%)	5.3 (3.7-7.1)	8.4% (3.6-13%)	9.8% (3.8-16%)	650 (260-1000)	11000 (4600-17000)
Czech Republic	2016	Male	80 to 84	82% (76-87%)	18% (13-24%)	4.5 (3.2-6.1)	6.8% (2.3-12%)	7.7% (2.1-14%)	610 (160-1100)	7200 (2500-12000)
Czech Republic	2016	Male	85 to 89	80% (73-86%)	20% (14-27%)	4.1 (2.9-5.5)	5.1% (0.71-10%)	5.5% (0.2-12%)	370 (14-800)	3100 (450-6300)
Czech Republic	2016	Male	90 to 94	79% (71-85%)	21% (15-29%)	3.6 (2.4-5.1)	3.1% (-1.5-9%)	3.1% (-2.5-10%)	100 (-88-360)	660 (-330-2000)
Czech Republic	2016	Male	95 plus	77% (67-85%)	23% (15-33%)	3.1 (1.7-4.9)	1.6% (-3.3-8.3%)	1.2% (-4.7-9.4%)	4.8 (-19-39)	26 (-57-140)
Hungary	1990	Female	All Ages	73% (69-77%)	27% (23-31%)	2.7 (2.4-3)	8.9% (3.9-14%)	11% (3.7-20%)	7700 (2600-13000)	180000 (77000-280000)
Hungary	1990	Female	15 to 19	77% (64-87%)	23% (13-36%)	2.1 (1.4-3)	3.8% (1.1-6.6%)	9.2% (1.1-17%)	14 (1.7-26)	1500 (460-2600)
Hungary	1990	Female	20 to 24	85% (77-90%)	15% (9.9-23%)	2.5 (1.6-3.6)	5.4% (2.3-8.7%)	10% (1.4-20%)	18 (2.5-35)	2100 (900-3400)
Hungary	1990	Female	25 to 29	84% (76-90%)	16% (9.7-24%)	2.6 (1.7-3.6)	5.8% (2.7-8.9%)	11% (3.1-19%)	23 (6.2-39)	2300 (1100-3500)
Hungary	1990	Female	30 to 34	84% (76-90%)	16% (10-24%)	2.7 (1.8-3.7)	7.1% (3.8-10%)	13% (5.9-20%)	61 (28-92)	4600 (2500-6700)
Hungary	1990	Female	35 to 39	84% (76-90%)	16% (9.9-24%)	2.9 (2-4)	9.1% (5.3-13%)	16% (8.7-23%)	140 (73-200)	8300 (4600-12000)
Hungary	1990	Female	40 to 44	83% (75-89%)	17% (11-25%)	3.4 (2.5-4.6)	11% (6.6-15%)	18% (11-25%)	200 (120-280)	11000 (6400-15000)
Hungary	1990	Female	45 to 49	80% (72-86%)	20% (14-28%)	4.2 (3-5.7)	12% (7.6-17%)	20% (12-27%)	300 (190-420)	14000 (8500-19000)
Hungary	1990	Female	50 to 54	77% (68-83%)	23% (17-32%)	4.3 (3-5.8)	13% (7.4-18%)	19% (11-27%)	400 (230-560)	16000 (9200-23000)
Hungary	1990	Female	55 to 59	73% (64-79%)	27% (21-36%)	4.2 (2.9-5.7)	12% (6.3-18%)	17% (9.4-25%)	530 (290-800)	18000 (9600-28000)
Hungary	1990	Female	60 to 64	71% (61-78%)	29% (22-39%)	4.1 (2.9-5.7)	11% (4.6-18%)	15% (6.8-24%)	710 (320-1100)	21000 (8900-34000)
Hungary	1990	Female	65 to 69	70% (60-79%)	30% (21-40%)	4 (2.8-5.3)	11% (3.9-18%)	14% (5.4-24%)	990 (380-1700)	24000 (8900-41000)
Hungary	1990	Female	70 to 74	65% (54-74%)	35% (26-46%)	3.7 (2.4-5.1)	9.9% (2.5-18%)	13% (3.4-23%)	820 (220-1500)	16000 (4100-29000)
Hungary	1990	Female	75 to 79	55% (42-68%)	45% (32-58%)	3.1 (1.9-4.5)	8.6% (1.5-17%)	10% (1.8-21%)	1300 (230-2600)	20000 (3600-38000)
Hungary	1990	Female	80 to 84	50% (36-62%)	50% (38-64%)	2.6 (1.7-3.8)	7.2% (1.2-15%)	8.6% (1.4-18%)	1200 (190-2400)	13000 (2200-26000)
Hungary	1990	Female	85 to 89	49% (35-62%)	51% (38-65%)	2.5 (1.5-3.8)	6.5% (0.63-14%)	7.7% (0.65-17%)	680 (56-1500)	5200 (520-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	1990	Female	90 to 94	50% (37-64%)	50% (36-63%)	2.6 (1.5-4.1)	6% (-0.29-15%)	7.1% (-0.52-18%)	260 (-20-680)	1400 (-71-3600)
Hungary	1990	Female	95 plus	54% (36-70%)	46% (30-64%)	2.6 (1.3-4.6)	5.6% (-1.6-16%)	6.6% (-2.1-19%)	54 (-17-160)	190 (-53-540)
Hungary	1990	Male	All Ages	87% (84-90%)	13% (10-16%)	6.6 (5.9-7.2)	20% (9.9-29%)	22% (9.5-33%)	17000 (7100-25000)	490000 (240000-700000)
Hungary	1990	Male	15 to 19	84% (74-91%)	16% (8.6-26%)	2.4 (1.5-3.6)	7.6% (2.1-13%)	12% (2.1-23%)	48 (8-87)	4100 (1200-7000)
Hungary	1990	Male	20 to 24	92% (87-95%)	8.3% (5.3-13%)	4.6 (3.3-6.2)	15% (7.1-24%)	23% (8.6-36%)	120 (44-190)	9200 (4300-14000)
Hungary	1990	Male	25 to 29	93% (88-96%)	7.3% (4.5-12%)	6.7 (4.6-8.9)	22% (13-30%)	32% (17-45%)	190 (100-280)	14000 (8200-19000)
Hungary	1990	Male	30 to 34	93% (88-96%)	7.2% (4.4-12%)	7.8 (5.7-10)	25% (16-33%)	34% (22-46%)	400 (250-530)	25000 (16000-33000)
Hungary	1990	Male	35 to 39	92% (88-95%)	7.8% (4.7-12%)	7.9 (5.6-11)	26% (18-35%)	34% (23-45%)	710 (480-930)	39000 (26000-51000)
Hungary	1990	Male	40 to 44	92% (88-95%)	7.9% (4.8-12%)	9.2 (6.8-12)	28% (19-36%)	35% (24-45%)	930 (650-1200)	46000 (31000-59000)
Hungary	1990	Male	45 to 49	93% (89-95%)	7.4% (4.7-11%)	11 (8.7-14)	28% (17-37%)	34% (21-44%)	1300 (780-1700)	54000 (34000-72000)
Hungary	1990	Male	50 to 54	93% (90-96%)	6.9% (4.4-10%)	12 (9.3-16)	26% (14-35%)	30% (16-42%)	1400 (770-2000)	54000 (29000-75000)
Hungary	1990	Male	55 to 59	93% (90-96%)	6.5% (4.3-9.9%)	12 (9.6-15)	24% (11-34%)	27% (13-39%)	1800 (850-2600)	59000 (27000-85000)
Hungary	1990	Male	60 to 64	91% (87-94%)	8.7% (6-13%)	12 (8.9-15)	21% (8.7-32%)	24% (10-36%)	2100 (890-3200)	60000 (25000-89000)
Hungary	1990	Male	65 to 69	87% (82-91%)	13% (9-18%)	10 (8.2-13)	19% (6.5-31%)	22% (7.5-35%)	2300 (750-3500)	53000 (17000-83000)
Hungary	1990	Male	70 to 74	85% (79-89%)	15% (11-21%)	9.1 (6.7-12)	17% (5.2-28%)	20% (6.1-32%)	1400 (430-2300)	27000 (8000-44000)
Hungary	1990	Male	75 to 79	84% (77-90%)	16% (10-23%)	7.9 (5.5-10)	16% (4.5-26%)	18% (5.1-29%)	1900 (550-3200)	29000 (8100-47000)
Hungary	1990	Male	80 to 84	84% (75-90%)	16% (10-25%)	6.9 (4.9-9.1)	14% (4.3-24%)	16% (4.7-28%)	1400 (410-2400)	16000 (4400-27000)
Hungary	1990	Male	85 to 89	84% (76-90%)	16% (10-24%)	6.2 (4.3-8.4)	12% (3.5-23%)	14% (3.8-26%)	640 (170-1200)	5000 (1400-8900)
Hungary	1990	Male	90 to 94	85% (77-91%)	15% (9.4-23%)	5.5 (3.6-7.8)	10% (2.4-20%)	12% (2.5-23%)	170 (37-350)	940 (220-1800)
Hungary	1990	Male	95 plus	85% (75-93%)	15% (7.4-25%)	4.7 (2.7-7.4)	7.8% (-0.93-18%)	8.7% (-1.7-21%)	20 (-3.8-47)	73 (-8.6-170)
Hungary	1995	Female	All Ages	74% (70-77%)	26% (23-30%)	2.5 (2.2-2.8)	8.3% (3.7-13%)	10% (3-18%)	6800 (2000-12000)	160000 (72000-250000)
Hungary	1995	Female	15 to 19	77% (66-87%)	23% (13-34%)	1.9 (1.2-2.7)	3.1% (1.1-5.2%)	8.3% (0.93-16%)	10 (1.1-19)	1200 (410-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	1995	Female	20 to 24	85% (77-90%)	15% (9.5-23%)	2.3 (1.5-3.2)	4.6% (2.2-7.2%)	9.2% (1-17%)	14 (1.5-26)	1900 (910-3000)
Hungary	1995	Female	25 to 29	85% (77-90%)	15% (9.8-23%)	2.3 (1.6-3.2)	5.2% (2.7-7.8%)	10% (2.7-17%)	19 (5-32)	2100 (1100-3200)
Hungary	1995	Female	30 to 34	84% (76-90%)	16% (10-24%)	2.4 (1.7-3.4)	6.5% (3.7-9.4%)	13% (6.4-19%)	40 (20-60)	3100 (1800-4500)
Hungary	1995	Female	35 to 39	84% (77-90%)	16% (10-23%)	2.6 (1.8-3.7)	8.9% (5.5-13%)	16% (9.4-22%)	120 (69-170)	7100 (4200-9900)
Hungary	1995	Female	40 to 44	83% (76-88%)	17% (12-24%)	3.1 (2.3-4.2)	11% (6.8-15%)	18% (11-23%)	240 (150-320)	12000 (7700-16000)
Hungary	1995	Female	45 to 49	80% (72-86%)	20% (14-28%)	3.8 (2.7-5.2)	12% (7.4-17%)	19% (12-25%)	310 (200-430)	14000 (8900-20000)
Hungary	1995	Female	50 to 54	77% (69-83%)	23% (17-31%)	4 (2.8-5.2)	12% (7.5-17%)	18% (12-25%)	400 (250-550)	16000 (9600-22000)
Hungary	1995	Female	55 to 59	73% (64-79%)	27% (21-36%)	3.8 (2.7-5.1)	11% (6.3-17%)	16% (9.4-24%)	480 (270-710)	16000 (9000-24000)
Hungary	1995	Female	60 to 64	71% (61-78%)	29% (22-39%)	3.8 (2.6-5.2)	10% (4.4-16%)	14% (6.7-22%)	600 (280-970)	18000 (8000-29000)
Hungary	1995	Female	65 to 69	71% (61-79%)	29% (21-39%)	3.7 (2.6-4.8)	9.4% (3.2-16%)	12% (4.5-21%)	810 (290-1400)	20000 (7000-34000)
Hungary	1995	Female	70 to 74	66% (55-74%)	34% (26-45%)	3.3 (2.2-4.6)	8.7% (2-16%)	11% (2.6-21%)	1100 (270-2000)	21000 (4800-39000)
Hungary	1995	Female	75 to 79	56% (43-69%)	44% (31-57%)	2.8 (1.8-4.2)	7.4% (1-15%)	9% (1.2-18%)	740 (96-1500)	11000 (1600-23000)
Hungary	1995	Female	80 to 84	50% (36-63%)	50% (37-64%)	2.4 (1.6-3.5)	6.1% (0.34-13%)	7.3% (0.28-16%)	980 (39-2100)	11000 (630-23000)
Hungary	1995	Female	85 to 89	49% (36-63%)	51% (37-64%)	2.3 (1.4-3.6)	5.5% (0.04-12%)	6.5% (-0.063-14%)	670 (-6.3-1500)	5200 (39-12000)
Hungary	1995	Female	90 to 94	51% (38-65%)	49% (35-62%)	2.4 (1.4-3.8)	5% (-0.89-13%)	5.8% (-1.1-16%)	250 (-48-670)	1300 (-220-3600)
Hungary	1995	Female	95 plus	54% (37-71%)	46% (29-63%)	2.5 (1.2-4.2)	4.7% (-2.2-14%)	5.4% (-3-17%)	59 (-32-190)	210 (-98-640)
Hungary	1995	Male	All Ages	88% (85-90%)	12% (10-15%)	6.2 (5.6-6.8)	21% (11-29%)	22% (11-33%)	17000 (8200-25000)	510000 (270000-720000)
Hungary	1995	Male	15 to 19	84% (74-92%)	16% (8.3-26%)	2.2 (1.4-3.2)	6.5% (2-11%)	12% (1.8-21%)	36 (5.4-65)	3300 (1000-5500)
Hungary	1995	Male	20 to 24	92% (87-95%)	8.2% (5.2-13%)	4.2 (3-5.6)	14% (6.4-21%)	21% (7.6-35%)	94 (33-150)	7900 (3700-12000)
Hungary	1995	Male	25 to 29	93% (89-96%)	7.2% (4.4-11%)	6.2 (4.3-8.4)	19% (12-27%)	30% (16-43%)	160 (86-230)	12000 (7000-16000)
Hungary	1995	Male	30 to 34	93% (88-96%)	7% (4.3-12%)	7.2 (5.2-9.3)	24% (16-32%)	34% (22-44%)	290 (190-390)	18000 (13000-24000)
Hungary	1995	Male	35 to 39	92% (88-95%)	7.6% (4.7-12%)	7.3 (5.1-9.8)	27% (19-35%)	34% (24-44%)	650 (460-850)	36000 (25000-46000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	1995	Male	40 to 44	92% (88-95%)	7.7% (4.6-12%)	8.4 (6.3-11)	29% (21-36%)	35% (26-44%)	1200 (880-1600)	59000 (43000-75000)
Hungary	1995	Male	45 to 49	93% (89-95%)	7.3% (4.6-11%)	11 (8-13)	29% (19-37%)	34% (23-44%)	1500 (1000-1900)	64000 (42000-83000)
Hungary	1995	Male	50 to 54	93% (90-96%)	6.9% (4.5-10%)	11 (8.5-14)	27% (16-36%)	31% (19-42%)	1700 (1000-2300)	65000 (38000-88000)
Hungary	1995	Male	55 to 59	93% (90-96%)	6.5% (4.4-9.6%)	11 (8.9-14)	24% (13-34%)	28% (15-39%)	1800 (950-2500)	58000 (31000-82000)
Hungary	1995	Male	60 to 64	91% (88-94%)	8.5% (5.9-12%)	11 (8.1-14)	21% (9.8-32%)	25% (12-36%)	2000 (930-3000)	56000 (25000-84000)
Hungary	1995	Male	65 to 69	87% (82-91%)	13% (9-18%)	9.5 (7.4-12)	19% (7.2-30%)	22% (8.4-34%)	2200 (810-3400)	51000 (19000-80000)
Hungary	1995	Male	70 to 74	85% (80-89%)	15% (11-20%)	8.4 (6.2-11)	17% (5.8-27%)	20% (6.7-31%)	2200 (750-3400)	41000 (14000-66000)
Hungary	1995	Male	75 to 79	84% (77-90%)	16% (10-23%)	7.2 (5-9.5)	15% (5.1-25%)	17% (5.8-28%)	1200 (420-2000)	18000 (6200-30000)
Hungary	1995	Male	80 to 84	84% (76-90%)	16% (10-24%)	6.3 (4.5-8.4)	14% (4.5-23%)	15% (5-26%)	1300 (440-2300)	15000 (4900-25000)
Hungary	1995	Male	85 to 89	84% (77-90%)	16% (10-23%)	5.7 (3.9-7.6)	12% (3.4-21%)	13% (3.8-24%)	670 (190-1200)	5200 (1600-9500)
Hungary	1995	Male	90 to 94	85% (78-91%)	15% (9.5-22%)	5 (3.2-7.2)	9.3% (1.4-19%)	10% (1.2-22%)	170 (19-350)	930 (140-1900)
Hungary	1995	Male	95 plus	86% (75-93%)	14% (7.2-25%)	4.3 (2.4-6.8)	6.7% (-1.9-17%)	7.3% (-2.9-19%)	22 (-9-59)	84 (-26-220)
Hungary	2000	Female	All Ages	73% (69-76%)	27% (24-31%)	2.3 (2-2.6)	7.2% (3.1-11%)	8.3% (2.3-15%)	5400 (1500-9600)	130000 (58000-210000)
Hungary	2000	Female	15 to 19	79% (67-88%)	21% (12-33%)	1.8 (1.2-2.6)	2.9% (1-4.9%)	7.8% (0.58-15%)	6.8 (0.5-13)	870 (300-1400)
Hungary	2000	Female	20 to 24	85% (77-90%)	15% (9.7-23%)	2.2 (1.4-3.1)	4.3% (2.2-6.5%)	8.3% (0.88-16%)	11 (1.1-21)	1700 (880-2700)
Hungary	2000	Female	25 to 29	84% (77-90%)	16% (10-23%)	2.2 (1.5-3)	4.5% (2.4-6.5%)	8.8% (1.9-16%)	13 (2.9-24)	1900 (1000-2800)
Hungary	2000	Female	30 to 34	84% (77-89%)	16% (11-23%)	2.3 (1.6-3.1)	5.1% (2.9-7.5%)	11% (4.5-17%)	25 (11-40)	2400 (1300-3500)
Hungary	2000	Female	35 to 39	83% (77-89%)	17% (11-23%)	2.4 (1.6-3.5)	7.1% (4.2-10%)	13% (7.3-19%)	62 (33-92)	4000 (2300-5800)
Hungary	2000	Female	40 to 44	82% (75-88%)	18% (12-25%)	2.9 (2.1-3.9)	9.3% (5.8-13%)	16% (10-21%)	160 (100-220)	8500 (5500-12000)
Hungary	2000	Female	45 to 49	80% (73-85%)	20% (15-27%)	3.5 (2.5-4.7)	11% (6.8-15%)	17% (11-23%)	310 (200-420)	14000 (9000-19000)
Hungary	2000	Female	50 to 54	75% (68-82%)	25% (18-32%)	3.6 (2.6-4.7)	11% (6.7-15%)	16% (11-22%)	350 (230-480)	14000 (8800-20000)
Hungary	2000	Female	55 to 59	70% (61-77%)	30% (23-39%)	3.3 (2.4-4.5)	10% (5.8-15%)	15% (9-21%)	410 (250-590)	14000 (8200-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	2000	Female	60 to 64	68% (59-75%)	32% (25-41%)	3.3 (2.3-4.6)	8.8% (4.1-14%)	12% (6.3-20%)	460 (230-730)	14000 (6500-22000)
Hungary	2000	Female	65 to 69	68% (59-76%)	32% (24-41%)	3.2 (2.4-4.3)	8.2% (2.9-14%)	11% (4.2-18%)	620 (230-1000)	15000 (5500-26000)
Hungary	2000	Female	70 to 74	62% (52-72%)	38% (28-48%)	2.9 (1.9-4)	7.3% (1.5-14%)	9.3% (2.1-17%)	800 (180-1500)	16000 (3400-29000)
Hungary	2000	Female	75 to 79	52% (41-65%)	48% (35-59%)	2.4 (1.5-3.5)	5.9% (0.44-12%)	7.2% (0.4-15%)	880 (48-1800)	14000 (980-28000)
Hungary	2000	Female	80 to 84	47% (34-59%)	53% (41-66%)	2.1 (1.3-3)	4.8% (-0.07-11%)	5.7% (-0.19-13%)	500 (-17-1100)	5600 (-88-12000)
Hungary	2000	Female	85 to 89	46% (33-59%)	54% (41-67%)	2 (1.2-3.1)	4.3% (-0.44-10%)	5% (-0.63-12%)	530 (-67-1300)	4200 (-450-9800)
Hungary	2000	Female	90 to 94	48% (35-61%)	52% (39-65%)	2 (1.1-3.3)	3.9% (-1.4-11%)	4.4% (-1.9-14%)	230 (-96-700)	1300 (-430-3700)
Hungary	2000	Female	95 plus	51% (34-69%)	49% (31-66%)	2.2 (1-3.7)	3.7% (-2.5-12%)	4.2% (-3.3-15%)	54 (-43-190)	200 (-140-660)
Hungary	2000	Male	All Ages	90% (88-92%)	10% (8.5-12%)	6.3 (5.7-6.9)	20% (11-28%)	22% (11-32%)	16000 (8100-22000)	450000 (250000-620000)
Hungary	2000	Male	15 to 19	88% (80-93%)	12% (6.9-20%)	2.3 (1.5-3.3)	6% (1.9-10%)	12% (1.9-22%)	22 (3.4-42)	2200 (700-3700)
Hungary	2000	Male	20 to 24	93% (90-95%)	6.7% (4.5-9.7%)	4.2 (3-5.4)	13% (6-19%)	21% (7.3-33%)	81 (29-130)	7100 (3300-11000)
Hungary	2000	Male	25 to 29	94% (91-96%)	5.9% (3.8-8.6%)	6.1 (4.3-8.1)	17% (10-24%)	29% (15-42%)	130 (71-190)	10000 (6100-14000)
Hungary	2000	Male	30 to 34	94% (91-96%)	5.8% (3.8-8.5%)	7.1 (5.2-9.1)	21% (14-28%)	33% (21-43%)	220 (140-300)	15000 (9600-19000)
Hungary	2000	Male	35 to 39	94% (91-96%)	6.4% (4.1-8.9%)	7.1 (5.1-9.6)	24% (17-32%)	34% (24-44%)	380 (260-490)	21000 (15000-27000)
Hungary	2000	Male	40 to 44	94% (91-96%)	6.4% (4.2-9.1%)	8.3 (6.2-11)	28% (20-35%)	35% (26-44%)	900 (660-1100)	44000 (32000-55000)
Hungary	2000	Male	45 to 49	94% (92-96%)	5.9% (4-8.2%)	10 (8-13)	29% (19-36%)	35% (24-43%)	1500 (1000-1900)	65000 (44000-83000)
Hungary	2000	Male	50 to 54	94% (92-96%)	5.6% (3.9-7.7%)	11 (8.7-14)	27% (17-35%)	32% (20-42%)	1600 (1000-2200)	62000 (38000-82000)
Hungary	2000	Male	55 to 59	95% (93-96%)	5.4% (3.9-7.3%)	11 (8.9-14)	24% (14-33%)	29% (17-39%)	1700 (960-2400)	57000 (31000-78000)
Hungary	2000	Male	60 to 64	93% (90-94%)	7.4% (5.7-9.7%)	11 (8.1-13)	22% (11-31%)	25% (13-36%)	1700 (880-2500)	49000 (24000-70000)
Hungary	2000	Male	65 to 69	89% (85-91%)	11% (8.7-15%)	9.3 (7.4-11)	19% (8-29%)	22% (9.3-33%)	1900 (800-2800)	45000 (18000-67000)
Hungary	2000	Male	70 to 74	86% (82-89%)	14% (11-18%)	8.2 (6.1-10)	17% (6.5-27%)	19% (7.7-30%)	1900 (770-3000)	37000 (15000-58000)
Hungary	2000	Male	75 to 79	85% (79-90%)	15% (10-21%)	6.9 (5-9)	15% (5.2-24%)	17% (6.1-27%)	1700 (620-2800)	25000 (9300-41000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	2000	Male	80 to 84	84% (75-90%)	16% (10-25%)	5.9 (4.1-7.8)	13% (4.7-22%)	15% (5.3-25%)	800 (290-1300)	8900 (3300-15000)
Hungary	2000	Male	85 to 89	84% (76-90%)	16% (10-24%)	5.3 (3.7-7.1)	11% (3.3-20%)	12% (3.6-22%)	640 (190-1200)	5100 (1500-9000)
Hungary	2000	Male	90 to 94	85% (77-90%)	15% (9.7-23%)	4.7 (3-6.7)	8.6% (0.46-17%)	9.5% (0.071-20%)	190 (1.4-400)	1000 (57-2100)
Hungary	2000	Male	95 plus	86% (76-93%)	14% (7.4-24%)	4 (2.2-6.4)	6% (-2.5-16%)	6.4% (-3.7-18%)	22 (-13-63)	85 (-37-230)
Hungary	2005	Female	All Ages	70% (67-73%)	30% (27-33%)	2.4 (2.1-2.7)	6.8% (3-11%)	8% (2.5-14%)	5100 (1600-9100)	120000 (55000-190000)
Hungary	2005	Female	15 to 19	79% (68-87%)	21% (13-32%)	2 (1.3-2.8)	2.9% (1-4.9%)	7.9% (0.86-15%)	6.1 (0.68-12)	800 (280-1300)
Hungary	2005	Female	20 to 24	85% (77-90%)	15% (10-23%)	2.3 (1.5-3.3)	4.1% (2.3-6.2%)	8.4% (1-16%)	7.4 (0.85-14)	1300 (710-2000)
Hungary	2005	Female	25 to 29	83% (76-88%)	17% (12-24%)	2.2 (1.5-3.1)	4.3% (2.4-6.2%)	8.6% (2-15%)	13 (3-23)	1900 (1000-2900)
Hungary	2005	Female	30 to 34	83% (76-88%)	17% (12-24%)	2.3 (1.6-3.2)	4.5% (2.5-6.6%)	10% (3.9-16%)	22 (8.5-34)	2300 (1300-3300)
Hungary	2005	Female	35 to 39	82% (76-88%)	18% (12-24%)	2.5 (1.7-3.6)	5.9% (3.4-8.7%)	12% (6.5-19%)	48 (25-73)	3400 (1900-4900)
Hungary	2005	Female	40 to 44	81% (74-87%)	19% (13-26%)	3 (2.1-4.1)	8% (4.9-11%)	15% (8.9-20%)	100 (62-140)	5500 (3300-7700)
Hungary	2005	Female	45 to 49	78% (71-84%)	22% (16-29%)	3.6 (2.6-4.9)	10% (6.2-14%)	16% (11-22%)	240 (150-320)	11000 (6900-15000)
Hungary	2005	Female	50 to 54	73% (67-79%)	27% (21-33%)	3.7 (2.6-4.8)	10% (6.4-14%)	16% (10-21%)	380 (250-520)	15000 (9500-21000)
Hungary	2005	Female	55 to 59	66% (60-73%)	34% (27-40%)	3.3 (2.4-4.5)	9.5% (5.5-14%)	14% (8.7-20%)	390 (240-570)	14000 (8100-20000)
Hungary	2005	Female	60 to 64	64% (57-71%)	36% (29-43%)	3.2 (2.3-4.5)	8.5% (4.1-14%)	12% (6.4-19%)	440 (230-690)	13000 (6400-21000)
Hungary	2005	Female	65 to 69	64% (56-72%)	36% (28-44%)	3.2 (2.3-4.1)	8% (3.2-13%)	11% (4.6-18%)	540 (230-870)	13000 (5300-22000)
Hungary	2005	Female	70 to 74	56% (47-64%)	44% (36-53%)	2.7 (1.8-3.7)	6.9% (1.8-13%)	9% (2.5-16%)	670 (190-1200)	13000 (3500-24000)
Hungary	2005	Female	75 to 79	47% (37-58%)	53% (42-63%)	2.2 (1.5-3.2)	5.8% (0.81-12%)	7.1% (0.94-14%)	790 (100-1600)	12000 (1800-25000)
Hungary	2005	Female	80 to 84	43% (30-56%)	57% (44-70%)	2 (1.3-2.9)	4.8% (0.2-10%)	5.7% (0.14-12%)	780 (19-1700)	8800 (390-19000)
Hungary	2005	Female	85 to 89	42% (29-55%)	58% (45-71%)	1.9 (1.1-2.9)	4.2% (-0.11-9.8%)	5% (-0.34-12%)	370 (-26-900)	3000 (-76-7100)
Hungary	2005	Female	90 to 94	43% (31-57%)	57% (43-69%)	1.9 (1.1-3.2)	4% (-1-11%)	4.6% (-1.4-13%)	260 (-84-770)	1400 (-370-4100)
Hungary	2005	Female	95 plus	47% (30-65%)	53% (35-70%)	2 (0.97-3.6)	3.9% (-1.8-12%)	4.4% (-2.5-15%)	74 (-42-240)	270 (-130-850)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	2005	Male	All Ages	90% (88-92%)	10% (8.4-12%)	6.8 (6.2-7.5)	20% (11-28%)	22% (10-32%)	15000 (7200-22000)	420000 (230000-600000)
Hungary	2005	Male	15 to 19	88% (81-93%)	12% (6.8-19%)	2.5 (1.7-3.5)	6.2% (2-10%)	12% (2.2-23%)	21 (3.6-39)	2100 (720-3500)
Hungary	2005	Male	20 to 24	93% (91-95%)	6.7% (4.7-9.5%)	4.4 (3.2-5.9)	13% (6.1-19%)	22% (8.1-36%)	61 (22-99)	5500 (2700-8300)
Hungary	2005	Male	25 to 29	94% (91-96%)	5.9% (4-8.6%)	6.5 (4.6-8.6)	17% (9.8-24%)	29% (15-43%)	130 (69-190)	10000 (6000-14000)
Hungary	2005	Male	30 to 34	94% (92-96%)	5.8% (3.9-8.4%)	7.5 (5.6-9.7)	19% (12-26%)	33% (20-45%)	180 (110-250)	12000 (8100-17000)
Hungary	2005	Male	35 to 39	93% (91-96%)	6.7% (4.3-9.4%)	7.5 (5.4-10)	22% (15-30%)	34% (23-44%)	310 (210-420)	18000 (12000-24000)
Hungary	2005	Male	40 to 44	93% (91-96%)	6.7% (4.4-9.5%)	8.7 (6.6-11)	26% (19-34%)	35% (25-44%)	580 (420-760)	29000 (21000-37000)
Hungary	2005	Male	45 to 49	94% (92-96%)	6% (4.2-8.3%)	11 (8.5-14)	28% (18-36%)	35% (24-44%)	1200 (810-1600)	53000 (35000-69000)
Hungary	2005	Male	50 to 54	94% (92-96%)	5.8% (4-7.8%)	12 (9.4-15)	26% (16-35%)	32% (19-42%)	1800 (1100-2400)	68000 (41000-91000)
Hungary	2005	Male	55 to 59	94% (92-96%)	5.6% (4.1-7.6%)	12 (9.4-15)	24% (13-33%)	28% (16-39%)	1800 (970-2400)	58000 (31000-81000)
Hungary	2005	Male	60 to 64	92% (90-94%)	7.6% (5.8-9.7%)	11 (8.6-14)	21% (10-31%)	25% (12-36%)	1800 (880-2600)	50000 (24000-73000)
Hungary	2005	Male	65 to 69	88% (85-91%)	12% (8.8-15%)	9.9 (8-12)	19% (7.6-29%)	22% (9.1-33%)	1700 (710-2600)	40000 (16000-61000)
Hungary	2005	Male	70 to 74	86% (83-89%)	14% (11-17%)	8.7 (6.6-11)	17% (5.8-26%)	19% (7.1-30%)	1800 (640-2800)	34000 (12000-54000)
Hungary	2005	Male	75 to 79	84% (79-89%)	16% (11-21%)	7.3 (5.3-9.5)	15% (4.5-24%)	17% (5.3-27%)	1600 (510-2700)	24000 (7600-40000)
Hungary	2005	Male	80 to 84	83% (76-89%)	17% (11-24%)	6.2 (4.4-8.3)	13% (4.2-22%)	15% (4.7-25%)	1300 (400-2100)	14000 (4600-24000)
Hungary	2005	Male	85 to 89	84% (77-90%)	16% (10-23%)	5.7 (4-7.6)	11% (3.2-20%)	13% (3.4-23%)	450 (120-810)	3600 (1000-6300)
Hungary	2005	Male	90 to 94	85% (78-90%)	15% (9.6-22%)	5.1 (3.3-7.2)	9.2% (1.4-18%)	10% (1.1-21%)	220 (23-450)	1200 (180-2400)
Hungary	2005	Male	95 plus	86% (75-92%)	14% (7.6-25%)	4.3 (2.4-6.8)	6.7% (-1.8-16%)	7.4% (-3-19%)	34 (-14-87)	130 (-37-310)
Hungary	2010	Female	All Ages	68% (65-71%)	32% (29-35%)	2 (1.8-2.3)	5.4% (2.3-8.6%)	6% (1.3-11%)	3900 (800-7300)	93000 (40000-150000)
Hungary	2010	Female	15 to 19	77% (66-87%)	23% (13-34%)	1.7 (1.1-2.4)	2.5% (0.98-4.2%)	6.5% (0.21-13%)	4.2 (0.13-8.3)	640 (250-1100)
Hungary	2010	Female	20 to 24	84% (76-89%)	16% (11-24%)	2 (1.3-2.9)	3.9% (2.2-5.7%)	7.1% (0.55-14%)	5 (0.41-9.7)	1100 (610-1700)
Hungary	2010	Female	25 to 29	81% (73-87%)	19% (13-27%)	1.9 (1.3-2.6)	3.9% (2.3-5.6%)	7.2% (1.4-13%)	7.3 (1.4-14)	1400 (770-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	2010	Female	30 to 34	81% (74-87%)	19% (13-26%)	2 (1.4-2.8)	4% (2.3-5.7%)	8.8% (3.4-14%)	16 (6-26)	2000 (1100-2900)
Hungary	2010	Female	35 to 39	81% (74-87%)	19% (13-26%)	2.2 (1.5-3.1)	4.7% (2.6-7%)	11% (5.4-16%)	34 (17-52)	2700 (1500-4000)
Hungary	2010	Female	40 to 44	79% (72-86%)	21% (14-28%)	2.6 (1.8-3.5)	5.9% (3.3-8.7%)	12% (6.5-17%)	67 (37-96)	3900 (2100-5600)
Hungary	2010	Female	45 to 49	76% (69-83%)	24% (17-31%)	3.1 (2.2-4.2)	8.1% (4.8-12%)	14% (8.6-19%)	140 (83-190)	6500 (3900-9100)
Hungary	2010	Female	50 to 54	73% (67-78%)	27% (22-33%)	3.2 (2.3-4.3)	8.8% (5.2-12%)	14% (8.5-18%)	260 (160-360)	11000 (6500-15000)
Hungary	2010	Female	55 to 59	66% (60-73%)	34% (27-40%)	2.9 (2.2-3.9)	8.1% (4.6-12%)	12% (7.2-17%)	370 (220-530)	13000 (7400-19000)
Hungary	2010	Female	60 to 64	62% (56-69%)	38% (31-44%)	2.7 (2-3.7)	7.1% (3.4-11%)	10% (5.4-15%)	370 (190-570)	11000 (5500-18000)
Hungary	2010	Female	65 to 69	63% (56-70%)	37% (30-44%)	2.7 (2-3.5)	6.4% (2.6-11%)	9% (3.9-15%)	420 (180-680)	11000 (4300-18000)
Hungary	2010	Female	70 to 74	54% (46-61%)	46% (39-54%)	2.2 (1.5-3.1)	5.3% (1.3-9.9%)	7% (1.8-13%)	460 (120-850)	9600 (2200-18000)
Hungary	2010	Female	75 to 79	45% (37-56%)	55% (44-63%)	1.9 (1.2-2.7)	4.4% (0.36-9.3%)	5.5% (0.44-12%)	530 (40-1100)	8500 (650-18000)
Hungary	2010	Female	80 to 84	42% (30-54%)	58% (46-70%)	1.7 (1.1-2.5)	3.6% (-0.35-8.2%)	4.3% (-0.52-10%)	560 (-67-1300)	6500 (-590-15000)
Hungary	2010	Female	85 to 89	40% (28-52%)	60% (48-72%)	1.6 (0.95-2.5)	3.1% (-0.67-7.5%)	3.5% (-0.98-9%)	440 (-120-1100)	3600 (-800-9000)
Hungary	2010	Female	90 to 94	42% (30-55%)	58% (45-70%)	1.6 (0.92-2.7)	2.8% (-1.3-8.6%)	3.2% (-1.9-10%)	150 (-86-500)	860 (-420-2700)
Hungary	2010	Female	95 plus	45% (29-62%)	55% (38-71%)	1.7 (0.82-3)	2.7% (-2.2-9.6%)	3.1% (-2.9-11%)	65 (-62-240)	240 (-200-860)
Hungary	2010	Male	All Ages	89% (87-91%)	11% (9.1-13%)	6.1 (5.5-6.8)	18% (10-25%)	20% (10-29%)	13000 (6600-19000)	360000 (200000-500000)
Hungary	2010	Male	15 to 19	87% (80-92%)	13% (7.6-20%)	2.1 (1.4-3.1)	5.3% (1.8-8.9%)	11% (1.3-20%)	15 (1.8-27)	1600 (570-2600)
Hungary	2010	Male	20 to 24	93% (90-95%)	7.2% (4.9-10%)	3.9 (2.8-5.2)	11% (5.6-17%)	20% (6.9-33%)	42 (14-70)	4200 (2100-6300)
Hungary	2010	Male	25 to 29	93% (90-96%)	6.6% (4.3-9.7%)	5.7 (3.9-7.6)	14% (8.4-20%)	27% (14-41%)	75 (38-110)	6400 (3900-9200)
Hungary	2010	Male	30 to 34	94% (91-96%)	6.3% (4.3-9.2%)	6.6 (4.9-8.6)	16% (10-22%)	30% (18-42%)	140 (82-190)	10000 (6500-14000)
Hungary	2010	Male	35 to 39	93% (90-95%)	7.2% (4.6-10%)	6.6 (4.8-9)	18% (12-25%)	31% (20-41%)	220 (150-300)	14000 (8900-19000)
Hungary	2010	Male	40 to 44	92% (89-95%)	7.7% (4.8-11%)	7.6 (5.7-9.8)	22% (15-29%)	33% (23-41%)	430 (300-550)	22000 (15000-28000)
Hungary	2010	Male	45 to 49	93% (91-96%)	6.6% (4.5-9%)	9.7 (7.5-12)	26% (17-32%)	33% (23-42%)	750 (510-940)	33000 (22000-42000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	2010	Male	50 to 54	94% (92-96%)	6% (4.2-8.3%)	11 (8.1-13)	25% (16-33%)	31% (19-40%)	1300 (830-1700)	51000 (31000-67000)
Hungary	2010	Male	55 to 59	94% (92-96%)	5.8% (4.2-7.7%)	11 (8.3-13)	23% (13-31%)	27% (16-37%)	1800 (1100-2500)	60000 (34000-83000)
Hungary	2010	Male	60 to 64	92% (90-94%)	7.7% (5.9-10%)	9.9 (7.7-13)	20% (10-29%)	24% (13-34%)	1700 (910-2400)	49000 (25000-69000)
Hungary	2010	Male	65 to 69	88% (85-91%)	12% (9.1-15%)	8.8 (7-11)	18% (8.7-27%)	21% (11-31%)	1600 (810-2400)	39000 (19000-58000)
Hungary	2010	Male	70 to 74	86% (82-89%)	14% (11-18%)	7.7 (5.7-9.7)	16% (6.5-24%)	18% (7.9-28%)	1500 (630-2200)	28000 (12000-43000)
Hungary	2010	Male	75 to 79	85% (79-89%)	15% (11-21%)	6.5 (4.6-8.5)	13% (5.1-22%)	16% (6-25%)	1400 (530-2200)	21000 (8100-33000)
Hungary	2010	Male	80 to 84	84% (76-90%)	16% (10-24%)	5.6 (4-7.5)	12% (4.2-20%)	14% (4.9-22%)	1100 (400-1900)	13000 (4600-21000)
Hungary	2010	Male	85 to 89	84% (77-90%)	16% (10-23%)	5.1 (3.6-6.9)	10% (2.8-18%)	11% (2.9-21%)	650 (160-1200)	5200 (1300-9200)
Hungary	2010	Male	90 to 94	85% (78-91%)	15% (9.5-22%)	4.6 (2.9-6.5)	7.8% (-0.11-16%)	8.7% (-0.58-19%)	150 (-9.9-330)	850 (-12-1800)
Hungary	2010	Male	95 plus	86% (75-93%)	14% (7.2-25%)	3.9 (2.2-6.3)	5.5% (-3-15%)	5.9% (-4.3-17%)	34 (-25-100)	130 (-73-370)
Hungary	2016	Female	All Ages	70% (67-74%)	30% (26-33%)	2 (1.6-2.5)	4.9% (1.8-8.4%)	5.2% (0.56-11%)	3500 (370-7200)	83000 (29000-140000)
Hungary	2016	Female	15 to 19	75% (61-86%)	25% (14-39%)	1.5 (0.88-2.2)	2.2% (0.91-3.7%)	5.7% (0.12-12%)	2.5 (0.052-5.2)	450 (190-750)
Hungary	2016	Female	20 to 24	84% (75-90%)	16% (10-25%)	1.8 (1.1-2.7)	3.6% (2.2-5.4%)	6.3% (0.21-13%)	3.9 (0.12-8.1)	1000 (560-1500)
Hungary	2016	Female	25 to 29	84% (74-90%)	16% (10-26%)	1.8 (1.1-2.8)	3.8% (2.2-5.5%)	6.8% (0.89-13%)	6 (0.71-12)	1200 (660-1800)
Hungary	2016	Female	30 to 34	83% (74-89%)	17% (11-26%)	1.9 (1.2-2.8)	3.8% (2.2-5.6%)	8.5% (3.1-14%)	10 (3.8-18)	1400 (770-2100)
Hungary	2016	Female	35 to 39	83% (75-90%)	17% (10-25%)	2.1 (1.4-3)	4.3% (2.3-6.5%)	10% (5-16%)	33 (16-52)	2700 (1400-4100)
Hungary	2016	Female	40 to 44	82% (74-88%)	18% (12-26%)	2.5 (1.7-3.5)	5.2% (2.7-8%)	11% (5.7-17%)	61 (31-94)	3700 (1900-5600)
Hungary	2016	Female	45 to 49	79% (70-85%)	21% (15-30%)	3 (2-4.3)	7.1% (3.8-11%)	13% (7.1-18%)	130 (72-190)	6200 (3400-9200)
Hungary	2016	Female	50 to 54	76% (68-82%)	24% (18-32%)	3.1 (2.2-4.4)	8% (4.5-12%)	13% (7.5-18%)	180 (110-260)	7500 (4200-11000)
Hungary	2016	Female	55 to 59	70% (62-77%)	30% (23-38%)	2.9 (2-4.1)	7.5% (4.1-12%)	11% (6.8-17%)	300 (170-460)	11000 (5800-17000)
Hungary	2016	Female	60 to 64	66% (57-74%)	34% (26-43%)	2.7 (1.8-3.9)	6.5% (2.9-11%)	9.5% (4.7-15%)	400 (200-660)	12000 (5400-20000)
Hungary	2016	Female	65 to 69	66% (57-74%)	34% (26-43%)	2.7 (1.8-3.6)	5.9% (2.1-10%)	8.4% (3.3-14%)	410 (150-690)	10000 (3500-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	2016	Female	70 to 74	59% (49-69%)	41% (31-51%)	2.3 (1.5-3.4)	5% (0.84-10%)	6.7% (1.3-13%)	450 (91-910)	9300 (1700-19000)
Hungary	2016	Female	75 to 79	49% (38-61%)	51% (39-62%)	1.9 (1.2-2.9)	3.9% (-0.2-8.9%)	4.9% (-0.37-11%)	450 (-34-1000)	7100 (-340-17000)
Hungary	2016	Female	80 to 84	44% (32-57%)	56% (43-68%)	1.7 (0.99-2.6)	3% (-0.71-7.7%)	3.6% (-0.96-9.3%)	460 (-120-1200)	5300 (-1200-14000)
Hungary	2016	Female	85 to 89	43% (30-56%)	57% (44-70%)	1.6 (0.89-2.5)	2.5% (-1.3-7.2%)	2.9% (-1.7-8.5%)	360 (-210-1000)	3000 (-1500-8500)
Hungary	2016	Female	90 to 94	45% (33-58%)	55% (42-67%)	1.6 (0.89-2.7)	2.3% (-1.8-8%)	2.6% (-2.5-9.7%)	230 (-210-860)	1300 (-1100-4600)
Hungary	2016	Female	95 plus	49% (32-66%)	51% (34-68%)	1.7 (0.79-3.1)	2.2% (-2.8-9.2%)	2.5% (-3.6-11%)	46 (-68-200)	170 (-220-700)
Hungary	2016	Male	All Ages	88% (86-90%)	12% (9.7-14%)	5.5 (4.4-6.7)	17% (9.7-23%)	18% (9.7-27%)	12000 (6000-17000)	320000 (180000-450000)
Hungary	2016	Male	15 to 19	82% (71-91%)	18% (9.4-29%)	1.7 (1-2.6)	4.5% (1.8-7.6%)	9% (0.88-17%)	8.8 (0.81-18)	1100 (420-1800)
Hungary	2016	Male	20 to 24	91% (86-95%)	9% (5.5-14%)	3.3 (2.1-4.7)	9.9% (5.2-15%)	17% (4.9-30%)	33 (9.7-58)	3500 (1800-5500)
Hungary	2016	Male	25 to 29	92% (87-95%)	7.8% (4.5-13%)	4.9 (3.2-7)	13% (7.5-18%)	25% (12-38%)	54 (25-85)	4900 (2900-7200)
Hungary	2016	Male	30 to 34	92% (87-95%)	7.7% (4.6-13%)	5.7 (3.9-7.8)	14% (8.8-20%)	28% (16-40%)	78 (43-110)	6000 (3800-8500)
Hungary	2016	Male	35 to 39	92% (87-95%)	8.3% (4.8-13%)	5.8 (3.9-8.1)	16% (10-22%)	29% (18-39%)	190 (120-270)	12000 (7800-17000)
Hungary	2016	Male	40 to 44	92% (87-95%)	8.4% (5.1-13%)	6.7 (4.8-9)	20% (13-26%)	31% (22-40%)	360 (240-480)	19000 (13000-25000)
Hungary	2016	Male	45 to 49	93% (89-95%)	7.4% (4.7-11%)	8.5 (6-11)	23% (16-30%)	32% (23-40%)	680 (470-910)	30000 (21000-41000)
Hungary	2016	Male	50 to 54	94% (91-96%)	6.3% (4.2-9.3%)	9.5 (6.8-13)	24% (15-31%)	30% (20-39%)	930 (600-1200)	36000 (23000-48000)
Hungary	2016	Male	55 to 59	94% (92-96%)	5.7% (4-7.8%)	9.5 (7.1-13)	22% (13-30%)	27% (17-36%)	1500 (870-2000)	49000 (29000-68000)
Hungary	2016	Male	60 to 64	92% (90-94%)	7.7% (5.8-10%)	9 (6.5-12)	19% (11-28%)	23% (13-33%)	1800 (1100-2600)	53000 (30000-76000)
Hungary	2016	Male	65 to 69	88% (84-91%)	12% (9-16%)	7.9 (5.8-10)	17% (9.2-25%)	20% (11-29%)	1600 (860-2300)	38000 (20000-57000)
Hungary	2016	Male	70 to 74	85% (81-89%)	15% (11-19%)	6.9 (4.8-9.1)	15% (7.1-22%)	18% (8.6-26%)	1500 (720-2300)	30000 (14000-46000)
Hungary	2016	Male	75 to 79	85% (79-90%)	15% (10-21%)	5.9 (4-8.1)	12% (4.7-20%)	15% (5.6-23%)	1100 (430-1900)	17000 (6500-28000)
Hungary	2016	Male	80 to 84	84% (75-90%)	16% (10-25%)	5.1 (3.4-7)	11% (3.6-18%)	12% (4-21%)	1000 (330-1700)	11000 (3800-19000)
Hungary	2016	Male	85 to 89	84% (76-90%)	16% (10-24%)	4.5 (2.8-6.4)	8.4% (1.2-16%)	9.5% (1.1-19%)	540 (56-1100)	4400 (600-8400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Hungary	2016	Male	90 to 94	85% (77-90%)	15% (9.6-23%)	4 (2.5-6)	6.2% (-1.3-14%)	6.8% (-2.1-16%)	220 (-66-540)	1300 (-260-3000)
Hungary	2016	Male	95 plus	85% (75-92%)	15% (7.5-25%)	3.4 (1.9-5.7)	4.2% (-3.6-13%)	4.4% (-5.1-15%)	22 (-26-73)	89 (-77-260)
Macedonia	1990	Female	All Ages	54% (44-63%)	46% (37-56%)	0.98 (0.73-1.3)	2.8% (1.5-4.5%)	4.2% (1.8-7.7%)	300 (130-550)	7800 (4200-12000)
Macedonia	1990	Female	15 to 19	53% (33-72%)	47% (28-67%)	0.76 (0.33-1.5)	2% (1.2-3.1%)	3.8% (1.6-7%)	1 (0.42-1.9)	150 (94-240)
Macedonia	1990	Female	20 to 24	64% (44-80%)	36% (20-56%)	1.1 (0.48-2)	3.5% (2.2-5.1%)	4.6% (1.5-9%)	1.7 (0.58-3.4)	310 (170-480)
Macedonia	1990	Female	25 to 29	65% (44-81%)	35% (19-56%)	1.2 (0.53-2.2)	3.5% (2.3-5.1%)	4.7% (1.8-8.7%)	2 (0.77-3.8)	330 (200-490)
Macedonia	1990	Female	30 to 34	66% (45-82%)	34% (18-55%)	1.3 (0.63-2.2)	3.4% (2.1-5%)	5.3% (2.3-9.6%)	3 (1.3-5.3)	360 (220-550)
Macedonia	1990	Female	35 to 39	67% (48-83%)	33% (17-52%)	1.4 (0.65-2.4)	3.5% (2.1-5.3%)	5.9% (2.5-10%)	5.1 (2.1-8.9)	450 (260-690)
Macedonia	1990	Female	40 to 44	66% (47-82%)	34% (18-53%)	1.6 (0.76-2.7)	3.6% (1.8-5.8%)	6.3% (2.5-11%)	7.1 (2.8-12)	460 (230-740)
Macedonia	1990	Female	45 to 49	64% (41-81%)	36% (19-59%)	1.7 (0.82-3.1)	4% (1.7-7.3%)	6.7% (2.4-12%)	10 (3.6-19)	520 (210-940)
Macedonia	1990	Female	50 to 54	61% (40-78%)	39% (22-60%)	1.7 (0.87-3.1)	4.4% (1.6-8.2%)	6.9% (2.5-13%)	18 (6.2-32)	750 (280-1300)
Macedonia	1990	Female	55 to 59	57% (36-76%)	43% (24-64%)	1.7 (0.79-3)	4.5% (1.2-8.6%)	6.7% (1.8-12%)	26 (7.4-49)	910 (260-1700)
Macedonia	1990	Female	60 to 64	54% (33-74%)	46% (26-67%)	1.5 (0.68-2.9)	3.9% (0.42-8.5%)	5.4% (0.75-12%)	32 (4.4-69)	920 (110-2000)
Macedonia	1990	Female	65 to 69	49% (30-70%)	51% (30-70%)	1.3 (0.65-2.3)	3.4% (0.41-7.7%)	4.6% (0.69-10%)	34 (5.2-75)	780 (95-1800)
Macedonia	1990	Female	70 to 74	43% (26-62%)	57% (38-74%)	1.1 (0.43-2.1)	3.3% (0.043-8.2%)	4.1% (0.21-9.9%)	31 (1.6-77)	570 (7.7-1400)
Macedonia	1990	Female	75 to 79	36% (22-55%)	64% (45-78%)	0.84 (0.31-1.7)	3.1% (0.081-7.5%)	3.7% (0.12-8.8%)	46 (1.4-110)	630 (16-1500)
Macedonia	1990	Female	80 to 84	31% (18-50%)	69% (50-82%)	0.75 (0.31-1.4)	3.2% (0.6-7.3%)	3.7% (0.76-8.5%)	42 (8.4-96)	430 (84-970)
Macedonia	1990	Female	85 to 89	30% (18-45%)	70% (55-82%)	0.75 (0.3-1.5)	3.5% (0.75-7.6%)	4.3% (1-9.4%)	24 (5.5-51)	170 (37-370)
Macedonia	1990	Female	90 to 94	32% (19-49%)	68% (51-81%)	0.81 (0.28-1.8)	3.9% (1.2-8.2%)	5% (1.5-11%)	12 (3.7-25)	57 (17-120)
Macedonia	1990	Female	95 plus	35% (19-55%)	65% (45-81%)	0.89 (0.25-2.1)	4.5% (1.2-9.6%)	5.6% (1.5-12%)	4.3 (1.1-9.1)	14 (3.8-31)
Macedonia	1990	Male	All Ages	61% (50-73%)	39% (27-50%)	1.5 (1.2-2)	4.7% (2.8-7%)	5.6% (2.4-9.2%)	500 (220-850)	16000 (9700-24000)
Macedonia	1990	Male	15 to 19	53% (31-76%)	47% (24-69%)	0.74 (0.34-1.3)	3.2% (2-4.8%)	4.3% (2.1-7.4%)	2.6 (1.3-4.5)	320 (200-480)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	1990	Male	20 to 24	66% (43-84%)	34% (16-57%)	1.4 (0.71-2.3)	6.4% (4.4-8.7%)	7.6% (3.9-12%)	5.7 (2.9-8.8)	710 (470-1000)
Macedonia	1990	Male	25 to 29	70% (49-88%)	30% (12-51%)	1.8 (0.94-2.9)	7.1% (5.1-9.5%)	8.9% (5.1-14%)	7.5 (4.2-12)	810 (560-1200)
Macedonia	1990	Male	30 to 34	73% (51-90%)	27% (9.7-49%)	2 (1.2-3.2)	6.9% (4.9-9.7%)	9.3% (5.4-14%)	10 (5.8-16)	910 (620-1300)
Macedonia	1990	Male	35 to 39	75% (55-90%)	25% (10-45%)	2.2 (1.2-3.5)	6.6% (4.2-9.8%)	8.9% (4.5-14%)	14 (7.2-24)	1100 (660-1600)
Macedonia	1990	Male	40 to 44	76% (54-90%)	24% (9.9-46%)	2.5 (1.4-3.9)	6.7% (3.9-10%)	8.9% (4.5-15%)	19 (9.5-31)	1100 (640-1700)
Macedonia	1990	Male	45 to 49	76% (53-90%)	24% (9.9-47%)	2.9 (1.6-4.5)	7.2% (3.7-12%)	9.1% (4.2-16%)	27 (12-47)	1300 (660-2200)
Macedonia	1990	Male	50 to 54	75% (53-89%)	25% (11-47%)	3 (1.8-4.8)	7.2% (3.3-12%)	8.8% (3.7-15%)	40 (16-70)	1700 (750-2800)
Macedonia	1990	Male	55 to 59	73% (51-89%)	27% (11-49%)	3.1 (1.7-5)	7.4% (3.1-13%)	8.8% (3.4-15%)	62 (24-110)	2200 (920-3700)
Macedonia	1990	Male	60 to 64	71% (47-89%)	29% (11-53%)	3 (1.5-5)	6.8% (2.4-12%)	7.8% (2.6-15%)	70 (24-130)	2100 (730-3800)
Macedonia	1990	Male	65 to 69	70% (46-88%)	30% (12-54%)	2.7 (1.5-4.2)	5.9% (1.7-11%)	6.7% (1.7-13%)	72 (18-140)	1700 (490-3300)
Macedonia	1990	Male	70 to 74	67% (45-85%)	33% (15-55%)	2.3 (1.2-3.9)	5% (0.93-10%)	5.6% (0.92-12%)	49 (7.9-100)	930 (170-1900)
Macedonia	1990	Male	75 to 79	64% (41-83%)	36% (17-59%)	1.9 (0.87-3.3)	4.1% (0.43-9%)	4.5% (0.41-10%)	65 (5.8-150)	950 (100-2100)
Macedonia	1990	Male	80 to 84	62% (37-82%)	38% (18-63%)	1.7 (0.79-2.8)	3.4% (0.063-7.7%)	3.7% (-0.033-8.6%)	42 (-0.36-98)	460 (8.4-1000)
Macedonia	1990	Male	85 to 89	60% (37-81%)	40% (19-63%)	1.5 (0.75-2.7)	2.3% (-0.41-5.6%)	2.5% (-0.65-6.5%)	12 (-3.2-32)	99 (-18-240)
Macedonia	1990	Male	90 to 94	60% (38-82%)	40% (18-62%)	1.4 (0.67-2.7)	1.8% (-0.88-5%)	2% (-1.3-5.8%)	3.4 (-2.3-10)	19 (-9.5-53)
Macedonia	1990	Male	95 plus	62% (34-86%)	38% (14-66%)	1.3 (0.49-2.7)	1.9% (-1.3-5.5%)	2.1% (-2.1-6.3%)	0.87 (-0.87-2.6)	3.3 (-2.3-9.4)
Macedonia	1995	Female	All Ages	54% (45-63%)	46% (37-55%)	0.97 (0.77-1.2)	2.9% (1.5-4.6%)	4.1% (1.7-7.5%)	300 (130-540)	7700 (4200-12000)
Macedonia	1995	Female	15 to 19	54% (34-72%)	46% (28-66%)	0.75 (0.33-1.4)	1.9% (1.1-2.9%)	3.6% (1.4-6.5%)	0.88 (0.37-1.6)	140 (83-210)
Macedonia	1995	Female	20 to 24	64% (45-80%)	36% (20-55%)	1.1 (0.46-2)	3.4% (2.2-4.9%)	4.3% (1.4-8.7%)	1.4 (0.47-2.8)	270 (160-420)
Macedonia	1995	Female	25 to 29	65% (44-81%)	35% (19-56%)	1.1 (0.52-2)	3.4% (2.3-4.7%)	4.3% (1.6-8.1%)	1.6 (0.62-3.1)	300 (190-430)
Macedonia	1995	Female	30 to 34	66% (45-82%)	34% (18-55%)	1.2 (0.63-2.1)	3.3% (2.1-4.9%)	4.9% (2-8.8%)	2.8 (1.1-5)	340 (200-510)
Macedonia	1995	Female	35 to 39	67% (49-83%)	33% (17-51%)	1.4 (0.66-2.4)	3.3% (1.9-5.1%)	5.5% (2.3-9.7%)	4.4 (1.9-7.8)	390 (230-600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	1995	Female	40 to 44	66% (48-82%)	34% (18-52%)	1.5 (0.77-2.5)	3.4% (1.7-5.5%)	5.9% (2.4-10%)	7.8 (3.1-13)	500 (260-800)
Macedonia	1995	Female	45 to 49	64% (41-81%)	36% (19-59%)	1.7 (0.81-3)	3.8% (1.6-7%)	6.3% (2.2-12%)	11 (3.7-20)	540 (220-980)
Macedonia	1995	Female	50 to 54	61% (41-79%)	39% (21-59%)	1.7 (0.87-3)	4.2% (1.5-7.8%)	6.6% (2.4-12%)	16 (5.9-29)	670 (250-1200)
Macedonia	1995	Female	55 to 59	58% (38-77%)	42% (23-62%)	1.6 (0.79-2.9)	4.3% (1.2-8.1%)	6.4% (1.8-12%)	25 (7-47)	850 (240-1600)
Macedonia	1995	Female	60 to 64	54% (33-74%)	46% (26-67%)	1.5 (0.69-2.8)	3.7% (0.36-8.1%)	5.1% (0.65-11%)	31 (3.9-67)	870 (88-1900)
Macedonia	1995	Female	65 to 69	50% (30-70%)	50% (30-70%)	1.3 (0.65-2.2)	3.2% (0.22-7.2%)	4.3% (0.41-9.4%)	37 (3.5-82)	860 (62-1900)
Macedonia	1995	Female	70 to 74	43% (26-62%)	57% (38-74%)	1 (0.43-2)	3.1% (-0.0098-7.5%)	3.9% (0.1-9.3%)	45 (1.2-110)	810 (-2.6-2000)
Macedonia	1995	Female	75 to 79	36% (22-56%)	64% (44-78%)	0.83 (0.31-1.6)	2.8% (0.0074-7%)	3.4% (0.019-8.2%)	31 (0.18-76)	430 (1.2-1100)
Macedonia	1995	Female	80 to 84	32% (18-50%)	68% (50-82%)	0.73 (0.3-1.4)	3% (0.49-7%)	3.5% (0.55-8.2%)	38 (6-89)	390 (60-900)
Macedonia	1995	Female	85 to 89	30% (18-46%)	70% (54-82%)	0.73 (0.3-1.4)	3.4% (0.65-7.4%)	4.1% (0.84-9%)	30 (6-65)	210 (41-460)
Macedonia	1995	Female	90 to 94	32% (20-49%)	68% (51-80%)	0.79 (0.29-1.7)	3.7% (1-8.1%)	4.7% (1.3-10%)	13 (3.8-29)	66 (18-140)
Macedonia	1995	Female	95 plus	36% (19-56%)	64% (44-81%)	0.88 (0.24-2.1)	4.2% (1.1-9.2%)	5.3% (1.4-12%)	4.8 (1.3-11)	16 (4-35)
Macedonia	1995	Male	All Ages	62% (50-73%)	38% (27-50%)	1.5 (1.3-1.8)	5% (3.1-7%)	5.6% (2.5-8.9%)	510 (230-820)	16000 (10000-24000)
Macedonia	1995	Male	15 to 19	54% (32-76%)	46% (24-68%)	0.73 (0.35-1.3)	3.1% (2-4.6%)	4.2% (2-7.1%)	2.4 (1.1-4)	300 (180-440)
Macedonia	1995	Male	20 to 24	66% (42-85%)	34% (15-58%)	1.3 (0.66-2.2)	6.2% (4.4-8.6%)	7.4% (3.9-11%)	5.2 (2.7-8.2)	640 (430-910)
Macedonia	1995	Male	25 to 29	70% (49-87%)	30% (13-51%)	1.7 (0.97-2.8)	6.9% (5-9.3%)	8.7% (5.1-13%)	6.9 (4.1-10)	770 (540-1100)
Macedonia	1995	Male	30 to 34	73% (50-90%)	27% (10-50%)	2 (1.1-3.1)	6.8% (4.9-9.5%)	9.2% (5.7-14%)	9.1 (5.5-14)	830 (580-1200)
Macedonia	1995	Male	35 to 39	75% (54-90%)	25% (10-46%)	2.1 (1.2-3.3)	6.5% (4.2-9.4%)	8.7% (4.5-14%)	13 (6.8-20)	970 (600-1400)
Macedonia	1995	Male	40 to 44	76% (54-91%)	24% (9.3-46%)	2.4 (1.4-3.7)	6.6% (4.1-10%)	8.7% (4.6-14%)	22 (12-36)	1300 (790-1900)
Macedonia	1995	Male	45 to 49	76% (51-90%)	24% (10-49%)	2.8 (1.6-4.3)	7% (3.7-11%)	8.8% (4.2-15%)	29 (14-49)	1400 (740-2300)
Macedonia	1995	Male	50 to 54	75% (53-89%)	25% (11-47%)	3 (1.8-4.6)	7.1% (3.2-11%)	8.5% (3.7-14%)	38 (16-65)	1600 (740-2600)
Macedonia	1995	Male	55 to 59	73% (51-89%)	27% (11-49%)	3 (1.7-4.7)	7.2% (3.2-12%)	8.5% (3.5-15%)	59 (24-100)	2100 (900-3500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	1995	Male	60 to 64	71% (46-89%)	29% (11-54%)	2.9 (1.5-4.7)	6.5% (2-12%)	7.5% (2.2-14%)	73 (22-140)	2100 (710-3900)
Macedonia	1995	Male	65 to 69	70% (45-88%)	30% (12-55%)	2.6 (1.5-4)	5.7% (1.7-11%)	6.5% (1.8-12%)	76 (21-140)	1800 (530-3400)
Macedonia	1995	Male	70 to 74	68% (44-85%)	32% (15-56%)	2.2 (1.2-3.7)	4.8% (0.84-10%)	5.5% (0.86-11%)	73 (11-160)	1400 (230-2900)
Macedonia	1995	Male	75 to 79	65% (41-84%)	35% (16-59%)	1.8 (0.87-3.3)	4% (0.34-8.7%)	4.4% (0.31-9.7%)	40 (2.9-89)	580 (51-1300)
Macedonia	1995	Male	80 to 84	62% (37-82%)	38% (18-63%)	1.6 (0.83-2.7)	3.2% (-0.0044-7.3%)	3.5% (-0.034-8.1%)	37 (-0.37-85)	400 (-0.53-900)
Macedonia	1995	Male	85 to 89	60% (39-81%)	40% (19-61%)	1.5 (0.74-2.5)	2.2% (-0.43-5.4%)	2.4% (-0.66-6%)	17 (-4.7-42)	130 (-26-320)
Macedonia	1995	Male	90 to 94	60% (39-82%)	40% (18-61%)	1.4 (0.65-2.5)	1.7% (-0.98-4.5%)	1.9% (-1.4-5.2%)	4.3 (-3.2-12)	24 (-14-62)
Macedonia	1995	Male	95 plus	61% (34-86%)	39% (14-66%)	1.3 (0.47-2.5)	1.9% (-1.2-5%)	2.1% (-1.8-5.8%)	1.2 (-0.98-3.3)	4.4 (-2.8-12)
Macedonia	2000	Female	All Ages	55% (46-64%)	45% (36-54%)	0.92 (0.76-1.1)	2.6% (1.2-4.2%)	3.4% (1.1-6.5%)	280 (89-530)	7000 (3200-11000)
Macedonia	2000	Female	15 to 19	54% (34-72%)	46% (28-66%)	0.67 (0.3-1.3)	1.8% (1.1-2.6%)	3.1% (1.2-5.8%)	0.77 (0.31-1.4)	130 (81-200)
Macedonia	2000	Female	20 to 24	64% (44-80%)	36% (20-56%)	0.97 (0.41-1.8)	3.1% (2-4.6%)	3.6% (1.1-7.2%)	1.1 (0.34-2.1)	250 (150-380)
Macedonia	2000	Female	25 to 29	66% (45-82%)	34% (18-55%)	1 (0.49-1.8)	3.2% (2.2-4.4%)	3.7% (1.4-7%)	1.1 (0.42-2.2)	260 (170-380)
Macedonia	2000	Female	30 to 34	67% (46-82%)	33% (18-54%)	1.1 (0.57-1.9)	3% (2-4.4%)	4.2% (1.7-7.6%)	2.1 (0.86-3.9)	310 (200-460)
Macedonia	2000	Female	35 to 39	67% (49-83%)	33% (17-51%)	1.3 (0.58-2.1)	3% (1.8-4.5%)	4.7% (1.9-8.5%)	3.6 (1.5-6.5)	340 (200-520)
Macedonia	2000	Female	40 to 44	67% (48-82%)	33% (18-52%)	1.4 (0.68-2.3)	3% (1.5-5%)	5% (1.8-8.8%)	6.3 (2.3-12)	430 (210-690)
Macedonia	2000	Female	45 to 49	65% (42-81%)	35% (19-58%)	1.5 (0.76-2.7)	3.3% (1.4-5.9%)	5.4% (1.9-10%)	11 (3.9-21)	590 (240-1100)
Macedonia	2000	Female	50 to 54	62% (41-79%)	38% (21-59%)	1.6 (0.8-2.7)	3.6% (1.2-6.7%)	5.6% (1.8-10%)	15 (4.6-27)	610 (210-1100)
Macedonia	2000	Female	55 to 59	58% (38-77%)	42% (23-62%)	1.5 (0.73-2.6)	3.6% (0.85-7.1%)	5.5% (1.4-10%)	20 (5-38)	680 (160-1300)
Macedonia	2000	Female	60 to 64	55% (33-75%)	45% (25-67%)	1.4 (0.64-2.5)	3% (0.053-7%)	4.3% (0.17-9.8%)	26 (1-60)	730 (12-1700)
Macedonia	2000	Female	65 to 69	50% (30-71%)	50% (29-70%)	1.2 (0.61-1.9)	2.5% (-0.29-6%)	3.4% (-0.31-8%)	32 (-2.9-76)	730 (-81-1800)
Macedonia	2000	Female	70 to 74	43% (26-62%)	57% (38-74%)	0.94 (0.4-1.8)	2.4% (-0.36-6.3%)	3% (-0.39-7.8%)	41 (-5.6-110)	740 (-110-2000)
Macedonia	2000	Female	75 to 79	36% (22-56%)	64% (44-78%)	0.75 (0.29-1.5)	2.2% (-0.41-5.8%)	2.6% (-0.4-7%)	42 (-6-110)	580 (-100-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	2000	Female	80 to 84	32% (18-50%)	68% (50-82%)	0.67 (0.27-1.2)	2.4% (0.17-5.9%)	2.8% (0.2-6.9%)	32 (2.2-80)	330 (22-830)
Macedonia	2000	Female	85 to 89	31% (19-46%)	69% (54-81%)	0.67 (0.27-1.3)	2.9% (0.27-7.6%)	3.4% (0.36-9%)	25 (2.6-68)	180 (16-490)
Macedonia	2000	Female	90 to 94	32% (19-48%)	68% (52-81%)	0.72 (0.27-1.5)	3% (0.71-7%)	3.8% (0.94-9.1%)	12 (3.1-29)	61 (14-140)
Macedonia	2000	Female	95 plus	36% (19-57%)	64% (43-81%)	0.8 (0.22-1.9)	3.4% (0.79-8.1%)	4.3% (0.97-10%)	4.5 (1-11)	15 (3.5-35)
Macedonia	2000	Male	All Ages	63% (51-75%)	37% (25-49%)	1.5 (1.2-1.7)	4.7% (2.9-6.8%)	4.9% (2.1-8.1%)	480 (210-800)	16000 (9700-22000)
Macedonia	2000	Male	15 to 19	54% (32-77%)	46% (23-68%)	0.67 (0.33-1.2)	3% (1.9-4.4%)	3.9% (1.9-6.7%)	2.2 (1-3.6)	290 (170-420)
Macedonia	2000	Male	20 to 24	66% (43-84%)	34% (16-57%)	1.2 (0.62-2)	6% (4-8.3%)	6.9% (3.6-11%)	5.2 (2.7-8.5)	650 (420-920)
Macedonia	2000	Male	25 to 29	70% (48-88%)	30% (12-52%)	1.6 (0.85-2.5)	6.7% (4.8-9%)	8.2% (4.7-12%)	6.5 (3.8-10)	740 (510-1000)
Macedonia	2000	Male	30 to 34	73% (51-89%)	27% (11-49%)	1.8 (1.1-2.8)	6.5% (4.7-8.8%)	8.6% (5.2-13%)	8.1 (4.9-12)	790 (540-1100)
Macedonia	2000	Male	35 to 39	75% (54-90%)	25% (9.9-46%)	2 (1.1-3.1)	6.2% (4.1-8.9%)	8.3% (4.5-13%)	12 (6.3-18)	880 (570-1200)
Macedonia	2000	Male	40 to 44	76% (55-91%)	24% (9.3-45%)	2.2 (1.4-3.3)	6.2% (4-9%)	8.2% (4.7-13%)	20 (11-30)	1200 (730-1700)
Macedonia	2000	Male	45 to 49	76% (52-90%)	24% (10-48%)	2.5 (1.5-3.9)	6.3% (3.5-10%)	7.9% (4-13%)	32 (16-52)	1600 (880-2500)
Macedonia	2000	Male	50 to 54	75% (52-90%)	25% (10-48%)	2.7 (1.7-4)	6.2% (3-10%)	7.4% (3.1-13%)	38 (17-64)	1600 (790-2600)
Macedonia	2000	Male	55 to 59	73% (51-90%)	27% (10-49%)	2.8 (1.6-4.2)	6.4% (2.7-11%)	7.5% (3-13%)	49 (20-85)	1700 (750-2900)
Macedonia	2000	Male	60 to 64	72% (46-89%)	28% (11-54%)	2.6 (1.4-4.2)	5.8% (1.8-10%)	6.6% (1.9-12%)	65 (19-120)	1900 (600-3500)
Macedonia	2000	Male	65 to 69	70% (46-89%)	30% (11-54%)	2.4 (1.4-3.6)	4.9% (1.4-9.3%)	5.6% (1.3-11%)	76 (19-140)	1800 (510-3400)
Macedonia	2000	Male	70 to 74	68% (45-85%)	32% (15-55%)	2 (1.1-3.2)	4.1% (0.65-8.5%)	4.6% (0.65-9.7%)	68 (9.8-140)	1300 (200-2700)
Macedonia	2000	Male	75 to 79	65% (42-84%)	35% (16-58%)	1.7 (0.82-2.9)	3.3% (0.01-7.6%)	3.7% (-0.03-8.3%)	56 (-0.44-130)	820 (2.6-1900)
Macedonia	2000	Male	80 to 84	63% (39-83%)	37% (17-61%)	1.5 (0.75-2.5)	2.6% (-0.31-6.2%)	2.9% (-0.47-6.8%)	29 (-4.7-68)	310 (-36-740)
Macedonia	2000	Male	85 to 89	61% (39-82%)	39% (18-61%)	1.4 (0.68-2.4)	1.8% (-1-4.9%)	2% (-1.4-5.4%)	13 (-8.6-35)	100 (-56-260)
Macedonia	2000	Male	90 to 94	61% (38-83%)	39% (17-62%)	1.3 (0.61-2.3)	1.3% (-1.3-4.1%)	1.5% (-1.8-4.7%)	4 (-4.9-13)	22 (-22-68)
Macedonia	2000	Male	95 plus	62% (35-87%)	38% (13-65%)	1.2 (0.44-2.3)	1.7% (-1.4-4.9%)	1.9% (-1.9-5.8%)	1.3 (-1.3-4)	4.8 (-3.8-14)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	2005	Female	All Ages	55% (46-64%)	45% (36-54%)	0.8 (0.65-0.96)	1.9% (0.72-3.3%)	2.3% (0.43-5.2%)	210 (41-480)	5300 (2000-9400)
Macedonia	2005	Female	15 to 19	53% (34-71%)	47% (29-66%)	0.58 (0.26-1.1)	1.6% (1-2.4%)	2.7% (0.97-5%)	0.59 (0.21-1.1)	110 (70-170)
Macedonia	2005	Female	20 to 24	63% (45-80%)	37% (20-55%)	0.82 (0.35-1.5)	3% (2-4.2%)	2.9% (0.73-6.1%)	0.72 (0.18-1.5)	230 (140-360)
Macedonia	2005	Female	25 to 29	65% (45-81%)	35% (19-55%)	0.88 (0.4-1.6)	3% (2.1-4.1%)	2.8% (0.84-5.6%)	0.81 (0.24-1.6)	250 (160-360)
Macedonia	2005	Female	30 to 34	66% (45-82%)	34% (18-55%)	0.95 (0.47-1.6)	2.7% (1.8-3.8%)	3.2% (1.2-6%)	1.3 (0.46-2.3)	260 (160-370)
Macedonia	2005	Female	35 to 39	67% (49-83%)	33% (17-51%)	1 (0.48-1.8)	2.6% (1.7-3.9%)	3.8% (1.5-6.8%)	2.6 (1-4.7)	300 (180-450)
Macedonia	2005	Female	40 to 44	66% (48-82%)	34% (18-52%)	1.2 (0.59-1.9)	2.4% (1.2-3.9%)	3.8% (1.2-6.9%)	4.1 (1.3-7.8)	320 (160-530)
Macedonia	2005	Female	45 to 49	64% (42-81%)	36% (19-58%)	1.3 (0.65-2.3)	2.5% (0.95-4.8%)	4% (1-7.9%)	7.8 (2-16)	430 (160-800)
Macedonia	2005	Female	50 to 54	61% (40-79%)	39% (21-60%)	1.3 (0.69-2.2)	2.7% (0.81-5.2%)	4.2% (1-8.3%)	13 (3.2-25)	560 (160-1100)
Macedonia	2005	Female	55 to 59	58% (37-77%)	42% (23-63%)	1.3 (0.63-2.1)	2.6% (0.43-5.3%)	4% (0.62-8.1%)	16 (2.3-32)	540 (84-1100)
Macedonia	2005	Female	60 to 64	54% (32-74%)	46% (26-68%)	1.2 (0.55-2.1)	2% (-0.5-5.3%)	2.9% (-0.59-7.4%)	16 (-3-42)	450 (-110-1200)
Macedonia	2005	Female	65 to 69	50% (30-71%)	50% (29-70%)	1 (0.51-1.6)	1.5% (-0.82-4.5%)	2.1% (-0.84-5.9%)	20 (-7.9-56)	450 (-250-1300)
Macedonia	2005	Female	70 to 74	43% (26-62%)	57% (38-74%)	0.79 (0.34-1.5)	1.4% (-0.78-4.6%)	1.9% (-0.87-6%)	27 (-12-86)	470 (-260-1600)
Macedonia	2005	Female	75 to 79	36% (22-56%)	64% (44-78%)	0.62 (0.24-1.2)	1.4% (-0.67-4.5%)	1.7% (-0.73-5.5%)	32 (-14-100)	430 (-210-1500)
Macedonia	2005	Female	80 to 84	32% (18-49%)	68% (51-82%)	0.56 (0.22-1.1)	1.6% (-0.29-5.2%)	1.9% (-0.24-6.5%)	33 (-4-110)	330 (-56-1100)
Macedonia	2005	Female	85 to 89	30% (18-46%)	70% (54-82%)	0.56 (0.22-1.1)	2.1% (0.092-6.8%)	2.5% (0.18-8.3%)	20 (1.4-66)	140 (6.2-470)
Macedonia	2005	Female	90 to 94	32% (19-48%)	68% (52-81%)	0.6 (0.22-1.3)	2.2% (0.17-6.9%)	2.7% (0.27-8.2%)	8 (0.8-25)	39 (3.1-120)
Macedonia	2005	Female	95 plus	35% (19-56%)	65% (44-81%)	0.67 (0.18-1.6)	2.5% (0.2-7.4%)	3.1% (0.31-8.8%)	4.1 (0.39-12)	13 (1-38)
Macedonia	2005	Male	All Ages	64% (52-75%)	36% (25-48%)	1.3 (1.1-1.5)	3.8% (2.3-5.4%)	3.6% (1.2-6.1%)	380 (130-640)	13000 (7600-19000)
Macedonia	2005	Male	15 to 19	54% (32-77%)	46% (23-68%)	0.57 (0.27-1)	2.8% (1.8-4%)	3.6% (1.5-6.1%)	1.6 (0.68-2.8)	240 (150-340)
Macedonia	2005	Male	20 to 24	66% (43-85%)	34% (15-57%)	1 (0.54-1.7)	5.6% (3.8-7.7%)	6% (2.8-10%)	4.2 (1.9-6.9)	590 (380-850)
Macedonia	2005	Male	25 to 29	70% (48-88%)	30% (12-52%)	1.3 (0.71-2.2)	6.1% (4.4-8.2%)	7% (3.8-11%)	5.3 (2.9-8.5)	670 (460-930)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	2005	Male	30 to 34	73% (50-90%)	27% (10-50%)	1.5 (0.86-2.4)	5.7% (4-7.7%)	7.1% (3.9-11%)	5.9 (3.3-9.3)	660 (450-920)
Macedonia	2005	Male	35 to 39	75% (54-90%)	25% (10-46%)	1.7 (0.95-2.6)	5.3% (3.4-7.3%)	6.9% (3.4-11%)	8.3 (4.2-13)	720 (450-1000)
Macedonia	2005	Male	40 to 44	76% (54-90%)	24% (9.7-46%)	1.9 (1.1-2.8)	5% (3-7.4%)	6.5% (3.3-11%)	13 (6.7-21)	850 (510-1200)
Macedonia	2005	Male	45 to 49	76% (52-90%)	24% (9.7-48%)	2.1 (1.2-3.3)	5.1% (2.8-8.2%)	6.4% (2.9-11%)	24 (11-40)	1200 (680-2000)
Macedonia	2005	Male	50 to 54	75% (52-90%)	25% (10-48%)	2.3 (1.4-3.4)	5% (2.4-8.1%)	5.9% (2.4-9.9%)	37 (15-63)	1600 (750-2600)
Macedonia	2005	Male	55 to 59	73% (51-89%)	27% (11-49%)	2.3 (1.3-3.5)	5% (2.2-8.7%)	5.9% (2.3-11%)	42 (16-76)	1500 (640-2700)
Macedonia	2005	Male	60 to 64	71% (47-89%)	29% (11-53%)	2.2 (1.2-3.6)	4.4% (1.3-8.7%)	5.1% (1.3-10%)	46 (12-93)	1400 (400-2700)
Macedonia	2005	Male	65 to 69	70% (46-88%)	30% (12-54%)	2 (1.2-3)	3.6% (0.83-7.1%)	4.1% (0.87-8.1%)	54 (11-110)	1300 (310-2500)
Macedonia	2005	Male	70 to 74	67% (44-85%)	33% (15-56%)	1.7 (0.89-2.8)	3% (0.24-6.5%)	3.3% (0.17-7.3%)	56 (2.8-120)	1100 (86-2400)
Macedonia	2005	Male	75 to 79	64% (40-84%)	36% (16-60%)	1.4 (0.7-2.4)	2.3% (-0.34-5.9%)	2.5% (-0.53-6.5%)	42 (-9-110)	620 (-91-1600)
Macedonia	2005	Male	80 to 84	62% (38-81%)	38% (19-62%)	1.3 (0.63-2.1)	1.8% (-0.77-4.9%)	2% (-0.97-5.5%)	26 (-13-74)	290 (-120-780)
Macedonia	2005	Male	85 to 89	60% (37-81%)	40% (19-63%)	1.1 (0.56-2)	1.3% (-1.4-3.9%)	1.3% (-1.7-4.4%)	8.5 (-11-28)	67 (-74-210)
Macedonia	2005	Male	90 to 94	60% (38-82%)	40% (18-62%)	1.1 (0.51-1.9)	0.86% (-1.8-3.2%)	0.9% (-2.3-3.6%)	2.1 (-5.3-8.6)	12 (-25-44)
Macedonia	2005	Male	95 plus	61% (35-86%)	39% (14-65%)	0.98 (0.37-2)	1.2% (-1.9-3.9%)	1.3% (-2.5-4.4%)	1.2 (-2.3-4.2)	4.4 (-7.1-14)
Macedonia	2010	Female	All Ages	54% (45-63%)	46% (37-55%)	0.73 (0.58-0.9)	1.7% (0.66-3.1%)	2% (0.37-5.2%)	190 (35-500)	4800 (1800-8800)
Macedonia	2010	Female	15 to 19	52% (32-71%)	48% (29-68%)	0.51 (0.22-0.97)	1.6% (1-2.3%)	2.6% (0.95-4.7%)	0.46 (0.17-0.84)	100 (64-160)
Macedonia	2010	Female	20 to 24	63% (44-80%)	37% (20-56%)	0.74 (0.31-1.4)	3% (2-4.1%)	2.7% (0.8-5.4%)	0.49 (0.14-0.97)	210 (130-320)
Macedonia	2010	Female	25 to 29	65% (44-81%)	35% (19-56%)	0.79 (0.34-1.4)	2.9% (2-4%)	2.5% (0.7-4.9%)	0.64 (0.18-1.3)	250 (160-360)
Macedonia	2010	Female	30 to 34	66% (44-81%)	34% (19-56%)	0.85 (0.4-1.4)	2.6% (1.8-3.6%)	2.8% (1.1-5.2%)	1 (0.4-1.9)	250 (160-370)
Macedonia	2010	Female	35 to 39	66% (48-82%)	34% (18-52%)	0.94 (0.42-1.7)	2.5% (1.6-3.6%)	3.4% (1.5-6.3%)	2.1 (0.92-3.8)	270 (170-410)
Macedonia	2010	Female	40 to 44	65% (47-81%)	35% (19-53%)	1.1 (0.51-1.8)	2.2% (1.2-3.6%)	3.5% (1.2-6.6%)	3.4 (1.2-6.5)	300 (160-490)
Macedonia	2010	Female	45 to 49	63% (40-81%)	37% (19-60%)	1.2 (0.58-2.1)	2.3% (0.96-4.3%)	3.7% (1.1-7.5%)	6.4 (1.9-13)	380 (160-680)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	2010	Female	50 to 54	60% (40-78%)	40% (22-60%)	1.2 (0.61-2)	2.3% (0.71-4.4%)	3.6% (0.9-7.2%)	10 (2.6-20)	460 (150-860)
Macedonia	2010	Female	55 to 59	57% (37-75%)	43% (25-63%)	1.1 (0.53-1.9)	2.2% (0.35-4.5%)	3.5% (0.64-7%)	16 (2.9-33)	550 (86-1100)
Macedonia	2010	Female	60 to 64	53% (32-73%)	47% (27-68%)	1 (0.48-1.9)	1.6% (-0.35-4.5%)	2.5% (-0.35-6.6%)	14 (-2-38)	390 (-83-1100)
Macedonia	2010	Female	65 to 69	49% (29-70%)	51% (30-71%)	0.88 (0.45-1.5)	1.2% (-0.71-3.8%)	1.7% (-0.73-5.3%)	15 (-6.4-44)	320 (-190-1000)
Macedonia	2010	Female	70 to 74	42% (25-61%)	58% (39-75%)	0.7 (0.29-1.4)	1.1% (-0.71-3.8%)	1.6% (-0.77-5%)	22 (-11-72)	380 (-240-1300)
Macedonia	2010	Female	75 to 79	35% (21-55%)	65% (45-79%)	0.55 (0.21-1.1)	1.1% (-0.72-4.2%)	1.4% (-0.85-5.2%)	27 (-16-99)	360 (-250-1400)
Macedonia	2010	Female	80 to 84	31% (18-48%)	69% (52-82%)	0.49 (0.2-0.93)	1.4% (-0.35-4.8%)	1.7% (-0.36-6.2%)	32 (-7.2-120)	330 (-84-1200)
Macedonia	2010	Female	85 to 89	29% (17-45%)	71% (55-83%)	0.49 (0.2-0.99)	1.8% (0.082-6%)	2.2% (0.19-7.5%)	28 (2.4-96)	190 (8.8-670)
Macedonia	2010	Female	90 to 94	31% (18-47%)	69% (53-82%)	0.53 (0.19-1.2)	1.8% (0.15-5.5%)	2.4% (0.25-7.3%)	8.3 (0.86-26)	40 (3.2-120)
Macedonia	2010	Female	95 plus	34% (18-55%)	66% (45-82%)	0.59 (0.16-1.4)	2.1% (0.22-6.2%)	2.7% (0.34-7.7%)	2.6 (0.31-7.3)	8.3 (0.84-24)
Macedonia	2010	Male	All Ages	64% (53-75%)	36% (25-47%)	1.2 (0.99-1.5)	3.4% (2-5%)	3.2% (1-5.4%)	350 (110-610)	12000 (6900-17000)
Macedonia	2010	Male	15 to 19	53% (31-75%)	47% (25-69%)	0.51 (0.24-0.91)	2.7% (1.7-3.8%)	3.3% (1.3-5.8%)	1.3 (0.51-2.3)	210 (140-310)
Macedonia	2010	Male	20 to 24	65% (42-84%)	35% (16-58%)	0.94 (0.48-1.6)	5.4% (3.7-7.5%)	5.6% (2.4-9.3%)	3.2 (1.4-5.4)	510 (330-730)
Macedonia	2010	Male	25 to 29	69% (49-87%)	31% (13-51%)	1.2 (0.65-2)	5.8% (4.3-7.7%)	6.5% (3.3-10%)	4.2 (2.2-6.9)	630 (430-870)
Macedonia	2010	Male	30 to 34	72% (50-89%)	28% (11-50%)	1.4 (0.81-2.3)	5.4% (3.8-7.2%)	6.4% (3.2-10%)	4.8 (2.4-7.8)	610 (420-850)
Macedonia	2010	Male	35 to 39	75% (54-89%)	25% (11-46%)	1.5 (0.84-2.4)	4.9% (3.2-6.8%)	6.2% (2.9-10%)	6.7 (3.2-11)	630 (400-900)
Macedonia	2010	Male	40 to 44	76% (53-90%)	24% (9.7-47%)	1.7 (0.99-2.6)	4.4% (2.6-6.6%)	5.8% (2.7-9.5%)	11 (4.9-18)	740 (430-1100)
Macedonia	2010	Male	45 to 49	75% (49-90%)	25% (10-51%)	1.9 (1.1-3)	4.6% (2.4-7.3%)	5.7% (2.6-9.6%)	20 (8.9-33)	1000 (560-1600)
Macedonia	2010	Male	50 to 54	74% (50-89%)	26% (11-50%)	2 (1.3-3.1)	4.5% (2.1-7.3%)	5.4% (2.2-9.2%)	32 (13-53)	1400 (650-2200)
Macedonia	2010	Male	55 to 59	72% (49-89%)	28% (11-51%)	2.1 (1.2-3.1)	4.4% (1.8-7.6%)	5.2% (2-9.2%)	46 (17-81)	1700 (650-2900)
Macedonia	2010	Male	60 to 64	70% (45-89%)	30% (11-55%)	2 (1.1-3.2)	3.8% (1.2-7.3%)	4.4% (1.3-8.5%)	44 (12-85)	1300 (410-2500)
Macedonia	2010	Male	65 to 69	69% (44-88%)	31% (12-56%)	1.8 (1-2.7)	3.1% (0.59-6.1%)	3.6% (0.62-7%)	44 (7.5-84)	1100 (210-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	2010	Male	70 to 74	67% (43-84%)	33% (16-57%)	1.5 (0.8-2.5)	2.6% (0.18-5.9%)	2.9% (0.13-6.8%)	49 (2.1-110)	930 (63-2200)
Macedonia	2010	Male	75 to 79	64% (40-84%)	36% (16-60%)	1.3 (0.6-2.2)	2% (-0.35-5.2%)	2.3% (-0.55-5.8%)	44 (-9.9-110)	640 (-110-1600)
Macedonia	2010	Male	80 to 84	61% (35-82%)	39% (18-65%)	1.1 (0.54-1.9)	1.6% (-0.78-4.1%)	1.8% (-1-4.6%)	28 (-15-73)	300 (-140-780)
Macedonia	2010	Male	85 to 89	59% (36-81%)	41% (19-64%)	1 (0.51-1.7)	1.2% (-1.4-3.5%)	1.3% (-1.7-4%)	12 (-15-35)	90 (-100-270)
Macedonia	2010	Male	90 to 94	59% (38-82%)	41% (18-62%)	0.95 (0.44-1.7)	0.88% (-1.7-3.3%)	0.97% (-2.1-3.8%)	2.5 (-5.5-9.8)	14 (-26-51)
Macedonia	2010	Male	95 plus	61% (34-85%)	39% (15-66%)	0.89 (0.32-1.8)	1.2% (-1.7-3.5%)	1.3% (-2.3-4.1%)	0.79 (-1.4-2.5)	2.9 (-4.1-8.6)
Macedonia	2016	Female	All Ages	54% (45-62%)	46% (38-55%)	0.76 (0.56-1)	1.7% (0.68-3.1%)	2% (0.44-4.9%)	200 (44-510)	5000 (2000-9300)
Macedonia	2016	Female	15 to 19	52% (32-71%)	48% (29-68%)	0.53 (0.21-1)	1.6% (0.99-2.3%)	2.7% (0.96-5.2%)	0.42 (0.15-0.79)	92 (56-150)
Macedonia	2016	Female	20 to 24	63% (44-79%)	37% (21-56%)	0.76 (0.3-1.5)	3% (2-4.2%)	2.8% (0.76-5.9%)	0.5 (0.13-1.1)	200 (120-300)
Macedonia	2016	Female	25 to 29	64% (44-80%)	36% (20-56%)	0.8 (0.35-1.5)	3% (2.1-4.1%)	2.6% (0.84-5.1%)	0.61 (0.19-1.2)	230 (150-340)
Macedonia	2016	Female	30 to 34	65% (44-81%)	35% (19-56%)	0.87 (0.41-1.6)	2.7% (1.9-3.8%)	2.9% (1.1-5.4%)	1 (0.4-2)	270 (170-400)
Macedonia	2016	Female	35 to 39	66% (47-82%)	34% (18-53%)	0.95 (0.43-1.7)	2.5% (1.7-3.7%)	3.4% (1.5-6.4%)	2 (0.86-3.8)	280 (180-420)
Macedonia	2016	Female	40 to 44	65% (47-81%)	35% (19-53%)	1.1 (0.49-1.8)	2.3% (1.3-3.7%)	3.6% (1.3-6.8%)	3.3 (1.2-6.2)	300 (160-470)
Macedonia	2016	Female	45 to 49	63% (40-80%)	37% (20-60%)	1.2 (0.54-2.1)	2.4% (1.1-4.5%)	3.9% (1.3-7.7%)	6.4 (2.1-13)	390 (170-720)
Macedonia	2016	Female	50 to 54	60% (38-78%)	40% (22-62%)	1.2 (0.58-2.1)	2.4% (0.78-4.7%)	3.8% (1.1-7.8%)	9.8 (2.8-20)	460 (150-880)
Macedonia	2016	Female	55 to 59	56% (36-76%)	44% (24-64%)	1.1 (0.52-2)	2.2% (0.42-4.4%)	3.5% (0.7-7.1%)	14 (2.7-29)	510 (95-1000)
Macedonia	2016	Female	60 to 64	53% (32-73%)	47% (27-68%)	1 (0.47-1.9)	1.7% (-0.22-4.5%)	2.6% (-0.12-6.8%)	17 (-0.82-45)	500 (-63-1300)
Macedonia	2016	Female	65 to 69	48% (29-70%)	52% (30-71%)	0.89 (0.44-1.5)	1.3% (-0.49-3.8%)	1.9% (-0.49-5.3%)	17 (-4.4-49)	370 (-150-1100)
Macedonia	2016	Female	70 to 74	42% (25-61%)	58% (39-75%)	0.7 (0.29-1.4)	1.2% (-0.66-3.9%)	1.7% (-0.69-5.2%)	21 (-8.6-64)	360 (-200-1200)
Macedonia	2016	Female	75 to 79	35% (21-53%)	65% (47-79%)	0.56 (0.22-1.2)	1.2% (-0.63-4%)	1.5% (-0.69-5.1%)	28 (-13-98)	380 (-190-1400)
Macedonia	2016	Female	80 to 84	30% (17-48%)	70% (52-83%)	0.5 (0.21-0.97)	1.3% (-0.34-4.3%)	1.6% (-0.32-5.2%)	33 (-6.3-110)	330 (-76-1100)
Macedonia	2016	Female	85 to 89	29% (18-45%)	71% (55-82%)	0.5 (0.19-1)	1.6% (-0.07-5%)	2.1% (-0.053-6.3%)	31 (-0.8-94)	220 (-10-650)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Macedonia	2016	Female	90 to 94	31% (19-46%)	69% (54-81%)	0.54 (0.19-1.2)	1.8% (0.092-5.1%)	2.3% (0.23-7.1%)	15 (1.4-44)	70 (3.5-210)
Macedonia	2016	Female	95 plus	34% (18-54%)	66% (46-82%)	0.61 (0.17-1.5)	2.1% (0.28-6.1%)	2.7% (0.39-7.9%)	4.4 (0.63-13)	14 (2-43)
Macedonia	2016	Male	All Ages	64% (53-75%)	36% (25-47%)	1.3 (0.96-1.7)	3.6% (2-5.5%)	3.4% (1.1-6.3%)	390 (140-720)	13000 (7200-20000)
Macedonia	2016	Male	15 to 19	53% (31-75%)	47% (25-69%)	0.52 (0.23-0.96)	2.7% (1.7-4%)	3.4% (1.4-6.1%)	1.2 (0.47-2.1)	190 (120-280)
Macedonia	2016	Male	20 to 24	65% (42-85%)	35% (15-58%)	0.96 (0.46-1.7)	5.6% (3.8-7.8%)	5.7% (2.4-9.6%)	3.2 (1.3-5.3)	510 (330-730)
Macedonia	2016	Male	25 to 29	69% (47-87%)	31% (13-53%)	1.2 (0.63-2.2)	6.2% (4.5-8.4%)	6.7% (3.1-11%)	4.2 (2-7)	630 (430-900)
Macedonia	2016	Male	30 to 34	72% (49-89%)	28% (11-51%)	1.4 (0.79-2.4)	5.7% (4-7.8%)	6.5% (3.2-11%)	5.2 (2.6-8.9)	690 (470-970)
Macedonia	2016	Male	35 to 39	74% (53-89%)	26% (11-47%)	1.5 (0.81-2.5)	5.2% (3.4-7.5%)	6.5% (3.1-11%)	7.3 (3.4-13)	700 (450-1000)
Macedonia	2016	Male	40 to 44	76% (54-90%)	24% (10-46%)	1.8 (0.99-2.7)	4.7% (2.8-7.2%)	6.1% (2.7-10%)	11 (4.7-19)	770 (450-1200)
Macedonia	2016	Male	45 to 49	75% (52-90%)	25% (10-48%)	2 (1.1-3.2)	4.8% (2.5-8.1%)	6.1% (2.6-11%)	20 (8.6-36)	1100 (560-1800)
Macedonia	2016	Male	50 to 54	74% (52-89%)	26% (11-48%)	2.1 (1.2-3.3)	4.8% (2.3-8.1%)	5.9% (2.5-10%)	31 (13-54)	1400 (670-2300)
Macedonia	2016	Male	55 to 59	72% (50-89%)	28% (11-50%)	2.1 (1.2-3.3)	4.6% (2.1-8.2%)	5.6% (2.2-10%)	45 (18-80)	1600 (740-2900)
Macedonia	2016	Male	60 to 64	70% (44-89%)	30% (11-56%)	2 (1.1-3.3)	4.1% (1.3-7.8%)	4.8% (1.4-9.2%)	58 (18-110)	1800 (590-3400)
Macedonia	2016	Male	65 to 69	69% (44-88%)	31% (12-56%)	1.8 (1-2.9)	3.3% (0.84-6.6%)	3.9% (0.89-7.9%)	53 (12-110)	1300 (330-2600)
Macedonia	2016	Male	70 to 74	67% (43-84%)	33% (16-57%)	1.6 (0.78-2.7)	2.7% (0.35-6.4%)	3.2% (0.39-7.3%)	48 (6-110)	920 (120-2100)
Macedonia	2016	Male	75 to 79	63% (39-83%)	37% (17-61%)	1.3 (0.6-2.4)	2.2% (-0.2-5.9%)	2.5% (-0.34-6.6%)	48 (-6.2-130)	700 (-64-1800)
Macedonia	2016	Male	80 to 84	61% (35-82%)	39% (18-65%)	1.1 (0.53-2)	1.8% (-0.64-4.7%)	2% (-0.95-5.3%)	34 (-16-91)	370 (-140-960)
Macedonia	2016	Male	85 to 89	59% (37-81%)	41% (19-63%)	1 (0.49-1.9)	1.2% (-1.4-3.7%)	1.3% (-1.7-4.2%)	15 (-18-46)	110 (-120-350)
Macedonia	2016	Male	90 to 94	59% (38-81%)	41% (19-62%)	0.97 (0.44-1.9)	0.92% (-1.6-3.3%)	1% (-2-3.8%)	3.9 (-7.6-14)	21 (-36-74)
Macedonia	2016	Male	95 plus	61% (34-85%)	39% (15-66%)	0.9 (0.32-1.9)	1.3% (-1.5-3.8%)	1.5% (-2-4.5%)	1.3 (-1.9-4)	4.8 (-5.6-14)
Montenegro	1990	Female	All Ages	57% (49-66%)	43% (34-51%)	2.1 (1.3-3.1)	7.1% (4.2-10%)	11% (5.5-17%)	240 (120-380)	5300 (3200-7600)
Montenegro	1990	Female	15 to 19	58% (38-75%)	42% (25-62%)	1.6 (0.64-3.3)	3.3% (1.9-5.3%)	8.6% (4-15%)	0.66 (0.31-1.2)	76 (45-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	1990	Female	20 to 24	68% (49-82%)	32% (18-51%)	2.3 (0.92-4.5)	5.4% (3.3-8.2%)	11% (4.6-19%)	1.3 (0.56-2.3)	140 (85-220)
Montenegro	1990	Female	25 to 29	69% (50-84%)	31% (16-50%)	2.4 (1-4.6)	5.5% (3.4-8.1%)	12% (5.2-19%)	1.5 (0.69-2.6)	160 (97-240)
Montenegro	1990	Female	30 to 34	70% (49-84%)	30% (16-51%)	2.6 (1.2-4.8)	5.9% (3.6-8.8%)	13% (6.6-20%)	2.4 (1.2-3.8)	200 (120-300)
Montenegro	1990	Female	35 to 39	70% (53-84%)	30% (16-47%)	2.9 (1.3-5.4)	7% (3.8-11%)	14% (6.9-22%)	4.3 (2.1-6.7)	280 (150-430)
Montenegro	1990	Female	40 to 44	69% (51-84%)	31% (16-49%)	3.2 (1.4-6.1)	8.8% (4.4-13%)	16% (7.5-24%)	6.7 (3.2-11)	360 (180-550)
Montenegro	1990	Female	45 to 49	68% (45-83%)	32% (17-55%)	3.6 (1.5-7.2)	8.8% (4.2-14%)	16% (7.9-24%)	7.3 (3.6-11)	350 (170-520)
Montenegro	1990	Female	50 to 54	65% (44-81%)	35% (19-56%)	3.7 (1.6-7.2)	9% (4.5-13%)	16% (8.1-24%)	11 (5.4-16)	450 (230-670)
Montenegro	1990	Female	55 to 59	62% (41-79%)	38% (21-59%)	3.6 (1.6-6.7)	9.3% (4.5-14%)	16% (8-23%)	14 (7.3-22)	510 (250-790)
Montenegro	1990	Female	60 to 64	58% (37-77%)	42% (23-63%)	3.3 (1.4-6.4)	10% (4.3-16%)	14% (6.2-23%)	23 (9.8-37)	650 (280-1100)
Montenegro	1990	Female	65 to 69	54% (34-73%)	46% (27-66%)	2.8 (1.3-5.4)	9.8% (3.8-17%)	13% (5.3-22%)	22 (8.4-36)	530 (200-870)
Montenegro	1990	Female	70 to 74	47% (30-65%)	53% (35-70%)	2.3 (0.89-4.7)	8.5% (2.4-15%)	12% (3.3-20%)	19 (5.2-34)	360 (100-640)
Montenegro	1990	Female	75 to 79	40% (25-59%)	60% (41-75%)	1.8 (0.67-4.1)	7.7% (2.1-14%)	10% (2.7-19%)	31 (8.2-58)	460 (120-860)
Montenegro	1990	Female	80 to 84	36% (21-54%)	64% (46-79%)	1.7 (0.66-3.4)	7.8% (2.4-14%)	9.9% (2.9-18%)	34 (9.9-62)	370 (120-670)
Montenegro	1990	Female	85 to 89	34% (21-50%)	66% (50-79%)	1.7 (0.6-3.7)	8.1% (2.8-15%)	10% (3.4-19%)	35 (12-64)	260 (90-480)
Montenegro	1990	Female	90 to 94	36% (23-53%)	64% (47-77%)	1.8 (0.61-4.3)	8% (2.3-16%)	9.8% (2.7-19%)	20 (5.7-39)	100 (29-200)
Montenegro	1990	Female	95 plus	40% (22-60%)	60% (40-78%)	2 (0.52-5.1)	7.9% (1.6-16%)	9.7% (2.1-19%)	7.1 (1.5-14)	23 (4.6-46)
Montenegro	1990	Male	All Ages	65% (54-75%)	35% (25-46%)	3.1 (2.1-4.7)	11% (7.3-14%)	14% (8.4-20%)	310 (180-430)	9300 (6400-12000)
Montenegro	1990	Male	15 to 19	58% (36-80%)	42% (20-64%)	1.6 (0.7-2.9)	5% (3-7.9%)	7.9% (4-14%)	1.7 (0.87-3)	170 (99-270)
Montenegro	1990	Male	20 to 24	70% (48-87%)	30% (13-52%)	2.8 (1.3-5.1)	9% (6-13%)	13% (7.4-22%)	3.2 (1.8-5.3)	310 (200-470)
Montenegro	1990	Male	25 to 29	73% (54-89%)	27% (11-46%)	3.6 (1.8-6.4)	11% (6.8-15%)	16% (9-24%)	5.6 (3.1-8.7)	470 (290-670)
Montenegro	1990	Male	30 to 34	76% (55-91%)	24% (8.8-45%)	4.2 (2.1-7.1)	11% (7.4-15%)	18% (11-26%)	6.2 (3.6-8.9)	470 (300-650)
Montenegro	1990	Male	35 to 39	78% (58-91%)	22% (8.6-42%)	4.4 (2.2-7.4)	12% (7.6-17%)	19% (11-27%)	9 (5.2-13)	580 (350-810)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	1990	Male	40 to 44	79% (59-91%)	21% (8.6-41%)	5 (2.5-8.4)	14% (7.9-19%)	20% (10-27%)	14 (7.3-20)	720 (410-1000)
Montenegro	1990	Male	45 to 49	79% (57-91%)	21% (8.8-43%)	5.8 (2.9-10)	14% (8.5-19%)	20% (11-27%)	15 (8.6-21)	700 (430-950)
Montenegro	1990	Male	50 to 54	77% (57-90%)	23% (9.6-43%)	6.2 (3.2-11)	15% (9.1-19%)	19% (12-25%)	26 (15-36)	1100 (660-1400)
Montenegro	1990	Male	55 to 59	76% (55-90%)	24% (9.6-45%)	6.3 (3.3-11)	15% (9.1-20%)	18% (11-24%)	39 (24-52)	1300 (830-1800)
Montenegro	1990	Male	60 to 64	74% (51-90%)	26% (9.6-49%)	6 (3-11)	14% (7.6-19%)	17% (9.2-23%)	42 (23-58)	1200 (690-1700)
Montenegro	1990	Male	65 to 69	73% (50-90%)	27% (10-50%)	5.4 (2.8-9.2)	13% (6.9-18%)	16% (8.5-23%)	36 (19-51)	860 (460-1200)
Montenegro	1990	Male	70 to 74	71% (49-86%)	29% (14-51%)	4.6 (2.2-8.8)	11% (5-17%)	14% (5.9-22%)	22 (9-33)	420 (190-640)
Montenegro	1990	Male	75 to 79	68% (45-85%)	32% (15-55%)	4 (1.7-8)	11% (3.6-18%)	13% (3.9-21%)	32 (9.9-53)	470 (150-770)
Montenegro	1990	Male	80 to 84	66% (42-84%)	34% (16-58%)	3.5 (1.6-6.7)	10% (3.3-17%)	12% (3.8-21%)	27 (8.7-49)	300 (99-520)
Montenegro	1990	Male	85 to 89	64% (43-83%)	36% (17-57%)	3.2 (1.5-6.1)	8.9% (2.2-16%)	10% (2.4-19%)	20 (4.9-38)	160 (40-290)
Montenegro	1990	Male	90 to 94	64% (42-85%)	36% (15-58%)	3 (1.3-6)	7.1% (0.86-15%)	8.3% (0.86-17%)	7.4 (0.73-15)	39 (4.7-80)
Montenegro	1990	Male	95 plus	65% (38-88%)	35% (12-62%)	2.7 (0.92-6)	5.9% (-0.32-14%)	6.8% (-0.67-16%)	1.6 (-0.16-4)	5.8 (-0.3-14)
Montenegro	1995	Female	All Ages	58% (50-67%)	42% (33-50%)	2 (1.3-3)	7.3% (4.2-11%)	11% (5.2-17%)	280 (130-440)	5900 (3400-8700)
Montenegro	1995	Female	15 to 19	58% (39-75%)	42% (25-61%)	1.6 (0.64-3)	3.2% (1.9-5.2%)	8.3% (4-15%)	0.61 (0.29-1.1)	70 (41-110)
Montenegro	1995	Female	20 to 24	68% (50-83%)	32% (17-50%)	2.2 (0.87-4.4)	5.3% (3.2-8.1%)	10% (4.5-18%)	1.2 (0.52-2.1)	140 (81-220)
Montenegro	1995	Female	25 to 29	69% (50-84%)	31% (16-50%)	2.3 (0.96-4.5)	5.5% (3.4-8.2%)	11% (5.1-19%)	1.5 (0.65-2.5)	150 (91-220)
Montenegro	1995	Female	30 to 34	70% (50-84%)	30% (16-50%)	2.5 (1.2-4.5)	5.7% (3.5-8.5%)	12% (6.4-20%)	2.3 (1.2-3.6)	190 (120-280)
Montenegro	1995	Female	35 to 39	71% (54-85%)	29% (15-46%)	2.8 (1.2-5.2)	6.6% (3.6-9.9%)	14% (6.7-21%)	3.8 (1.8-6)	260 (140-390)
Montenegro	1995	Female	40 to 44	70% (52-84%)	30% (16-48%)	3.1 (1.3-5.7)	7.6% (3.9-12%)	15% (7.4-23%)	6.3 (3.1-9.8)	350 (180-540)
Montenegro	1995	Female	45 to 49	68% (47-83%)	32% (17-53%)	3.4 (1.5-6.7)	9% (4.3-14%)	16% (7.8-24%)	8.7 (4.2-13)	410 (200-630)
Montenegro	1995	Female	50 to 54	65% (46-81%)	35% (19-54%)	3.5 (1.6-6.6)	9.5% (4.7-14%)	16% (8-24%)	11 (5.3-16)	440 (220-650)
Montenegro	1995	Female	55 to 59	62% (42-79%)	38% (21-58%)	3.4 (1.5-6.6)	9.9% (4.8-15%)	16% (7.7-24%)	18 (8.7-28)	620 (300-960)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	1995	Female	60 to 64	58% (38-78%)	42% (22-62%)	3.1 (1.4-6.5)	9.4% (3.8-15%)	14% (5.7-23%)	24 (9.7-39)	690 (280-1100)
Montenegro	1995	Female	65 to 69	54% (35-74%)	46% (26-65%)	2.7 (1.3-5.1)	9.5% (3.7-16%)	13% (5.2-22%)	30 (12-50)	710 (280-1200)
Montenegro	1995	Female	70 to 74	48% (30-66%)	52% (34-70%)	2.2 (0.87-4.5)	9% (2.3-17%)	12% (3-21%)	32 (8.2-58)	590 (160-1100)
Montenegro	1995	Female	75 to 79	41% (26-60%)	59% (40-74%)	1.8 (0.62-3.7)	8% (1.9-15%)	10% (2.3-19%)	29 (6.4-55)	420 (98-790)
Montenegro	1995	Female	80 to 84	36% (21-56%)	64% (44-79%)	1.6 (0.59-3.2)	7.7% (2.3-14%)	9.7% (2.7-18%)	32 (8.7-61)	350 (100-650)
Montenegro	1995	Female	85 to 89	35% (22-51%)	65% (49-78%)	1.6 (0.57-3.4)	8% (2.5-15%)	9.9% (3.1-19%)	41 (13-78)	310 (100-580)
Montenegro	1995	Female	90 to 94	36% (23-53%)	64% (47-77%)	1.7 (0.58-4)	7.9% (2.1-15%)	9.6% (2.6-19%)	26 (7.1-51)	130 (36-250)
Montenegro	1995	Female	95 plus	40% (22-61%)	60% (39-78%)	1.9 (0.52-5)	7.8% (1.5-16%)	9.5% (1.9-20%)	9.5 (2-19)	31 (6.2-64)
Montenegro	1995	Male	All Ages	66% (55-77%)	34% (23-45%)	3.1 (2.1-4.5)	11% (7.3-14%)	14% (8.1-20%)	370 (210-530)	11000 (7100-14000)
Montenegro	1995	Male	15 to 19	59% (37-80%)	41% (20-63%)	1.5 (0.7-2.8)	5.1% (3.1-8%)	7.8% (4-13%)	1.7 (0.87-3)	160 (97-260)
Montenegro	1995	Male	20 to 24	69% (47-87%)	31% (13-53%)	2.7 (1.2-4.8)	9.4% (5.9-14%)	13% (6.9-21%)	4.2 (2.3-7.1)	370 (230-580)
Montenegro	1995	Male	25 to 29	73% (53-89%)	27% (11-47%)	3.4 (1.6-6.2)	11% (6.9-16%)	16% (8.7-24%)	5.4 (3-8.4)	440 (270-630)
Montenegro	1995	Male	30 to 34	76% (54-90%)	24% (9.5-46%)	3.9 (2-6.7)	11% (7.4-16%)	17% (10-26%)	7.1 (4.1-10)	520 (330-730)
Montenegro	1995	Male	35 to 39	78% (59-91%)	22% (8.7-41%)	4.2 (2.1-7)	12% (7.5-17%)	18% (10-26%)	9.9 (5.6-14)	620 (380-880)
Montenegro	1995	Male	40 to 44	79% (58-92%)	21% (8.2-42%)	4.7 (2.5-7.9)	13% (7.7-19%)	19% (10-27%)	16 (8.9-24)	870 (510-1200)
Montenegro	1995	Male	45 to 49	78% (55-91%)	22% (8.6-45%)	5.5 (2.7-9.4)	14% (8-20%)	19% (11-27%)	19 (10-26)	870 (500-1200)
Montenegro	1995	Male	50 to 54	77% (56-91%)	23% (9.4-44%)	5.8 (3-9.9)	15% (8.7-19%)	19% (11-25%)	22 (13-30)	900 (540-1200)
Montenegro	1995	Male	55 to 59	76% (54-91%)	24% (9.4-46%)	5.9 (3.2-10)	15% (9.1-19%)	18% (11-24%)	40 (24-55)	1400 (840-1900)
Montenegro	1995	Male	60 to 64	75% (50-91%)	25% (9.4-50%)	5.7 (2.9-10)	14% (7.6-19%)	17% (9.3-23%)	52 (28-73)	1500 (840-2100)
Montenegro	1995	Male	65 to 69	74% (49-90%)	26% (10-51%)	5.2 (2.7-8.6)	13% (6.9-19%)	16% (8.1-23%)	58 (30-84)	1400 (720-2000)
Montenegro	1995	Male	70 to 74	71% (49-87%)	29% (13-51%)	4.5 (2.2-8.2)	12% (5.1-19%)	14% (5.9-22%)	45 (19-69)	860 (370-1300)
Montenegro	1995	Male	75 to 79	68% (46-86%)	32% (14-54%)	3.8 (1.7-7.5)	11% (3.2-18%)	12% (3.6-21%)	30 (8.7-51)	440 (140-750)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	1995	Male	80 to 84	66% (42-84%)	34% (16-58%)	3.4 (1.5-6.4)	10% (3.1-18%)	12% (3.5-21%)	26 (7.9-47)	280 (90-500)
Montenegro	1995	Male	85 to 89	65% (43-83%)	35% (17-57%)	3.1 (1.4-5.7)	8.7% (2.1-17%)	10% (2.3-19%)	23 (5-43)	170 (43-330)
Montenegro	1995	Male	90 to 94	64% (44-85%)	36% (15-56%)	2.8 (1.2-5.4)	6.9% (0.71-15%)	7.9% (0.66-17%)	9.9 (0.83-21)	52 (5.2-110)
Montenegro	1995	Male	95 plus	65% (39-88%)	35% (12-61%)	2.6 (0.89-5.7)	5.7% (-0.3-14%)	6.5% (-0.71-16%)	2.2 (-0.24-5.5)	7.7 (-0.41-19)
Montenegro	2000	Female	All Ages	58% (50-66%)	42% (34-50%)	2.1 (1.5-3)	7.5% (4.6-11%)	11% (5.8-17%)	300 (150-460)	6400 (3900-9200)
Montenegro	2000	Female	15 to 19	58% (38-74%)	42% (26-62%)	1.6 (0.65-3)	3.3% (1.9-5.4%)	8.3% (4.1-15%)	0.68 (0.33-1.2)	75 (44-120)
Montenegro	2000	Female	20 to 24	67% (49-83%)	33% (17-51%)	2.2 (0.91-4.3)	5.2% (3.2-8%)	10% (4.6-18%)	1.1 (0.48-1.9)	130 (76-200)
Montenegro	2000	Female	25 to 29	69% (50-84%)	31% (16-50%)	2.4 (1-4.5)	5.5% (3.4-8.1%)	11% (5.2-19%)	1.4 (0.65-2.3)	150 (89-220)
Montenegro	2000	Female	30 to 34	70% (49-84%)	30% (16-51%)	2.5 (1.2-4.5)	5.8% (3.6-8.5%)	13% (6.5-20%)	2 (1-3.1)	170 (100-250)
Montenegro	2000	Female	35 to 39	70% (52-84%)	30% (16-48%)	2.8 (1.3-5.2)	6.5% (3.7-9.8%)	14% (6.8-21%)	3.7 (1.8-5.7)	250 (140-370)
Montenegro	2000	Female	40 to 44	70% (52-84%)	30% (16-48%)	3.1 (1.5-5.6)	7.6% (4.1-11%)	15% (8-23%)	6.2 (3.1-9.5)	340 (180-520)
Montenegro	2000	Female	45 to 49	68% (46-84%)	32% (16-54%)	3.4 (1.7-6.5)	8.8% (4.5-13%)	16% (8.3-24%)	11 (5.4-16)	500 (270-760)
Montenegro	2000	Female	50 to 54	65% (45-81%)	35% (19-55%)	3.5 (1.7-6.2)	9.7% (4.9-14%)	16% (8.5-24%)	13 (6.6-19)	520 (270-770)
Montenegro	2000	Female	55 to 59	62% (42-80%)	38% (20-58%)	3.5 (1.6-6.5)	10% (5-16%)	16% (8.1-24%)	16 (7.9-24)	540 (270-830)
Montenegro	2000	Female	60 to 64	59% (38-78%)	41% (22-62%)	3.2 (1.4-6.3)	9.6% (4.1-15%)	14% (6.1-23%)	25 (11-39)	720 (310-1100)
Montenegro	2000	Female	65 to 69	54% (34-73%)	46% (27-66%)	2.7 (1.4-5.1)	9.6% (4.2-16%)	13% (5.8-22%)	35 (15-58)	840 (370-1400)
Montenegro	2000	Female	70 to 74	48% (30-66%)	52% (34-70%)	2.2 (0.9-4.3)	9% (2.7-16%)	12% (3.5-21%)	41 (12-73)	780 (240-1400)
Montenegro	2000	Female	75 to 79	41% (25-60%)	59% (40-75%)	1.8 (0.67-3.6)	8.1% (2.2-15%)	10% (2.6-19%)	37 (9.4-67)	530 (140-970)
Montenegro	2000	Female	80 to 84	36% (21-54%)	64% (46-79%)	1.6 (0.61-3.1)	8.2% (2.8-15%)	10% (3.3-18%)	33 (11-59)	350 (120-630)
Montenegro	2000	Female	85 to 89	34% (21-50%)	66% (50-79%)	1.6 (0.61-3.3)	8.5% (3-15%)	10% (3.6-19%)	38 (13-68)	280 (95-510)
Montenegro	2000	Female	90 to 94	36% (23-53%)	64% (47-77%)	1.7 (0.59-3.8)	8.3% (2.3-15%)	10% (2.8-19%)	29 (8.1-55)	150 (43-280)
Montenegro	2000	Female	95 plus	40% (23-61%)	60% (39-77%)	1.9 (0.53-4.8)	8.2% (1.7-17%)	10% (2.1-20%)	12 (2.6-24)	39 (8.3-78)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	2000	Male	All Ages	66% (55-77%)	34% (23-45%)	3.3 (2.3-4.6)	11% (7.9-15%)	15% (8.7-20%)	430 (250-600)	12000 (8100-16000)
Montenegro	2000	Male	15 to 19	58% (36-79%)	42% (21-64%)	1.5 (0.71-2.7)	4.7% (2.9-7.3%)	7.7% (4-13%)	1.4 (0.71-2.3)	140 (84-210)
Montenegro	2000	Male	20 to 24	69% (47-86%)	31% (14-53%)	2.7 (1.3-4.8)	9.1% (5.8-13%)	13% (7.2-21%)	3.2 (1.7-5)	300 (190-440)
Montenegro	2000	Male	25 to 29	73% (53-89%)	27% (11-47%)	3.5 (1.7-6.3)	11% (7-15%)	16% (9.1-24%)	4.5 (2.6-6.8)	380 (240-540)
Montenegro	2000	Male	30 to 34	76% (55-91%)	24% (9.5-45%)	4 (2.1-7)	12% (7.8-16%)	18% (11-26%)	6 (3.5-8.7)	440 (280-610)
Montenegro	2000	Male	35 to 39	78% (58-91%)	22% (9.1-42%)	4.2 (2.3-7.2)	12% (7.6-17%)	18% (11-26%)	9.6 (5.5-14)	600 (380-850)
Montenegro	2000	Male	40 to 44	79% (58-92%)	21% (8.3-42%)	4.8 (2.7-7.7)	13% (8-18%)	19% (11-27%)	15 (8.8-22)	820 (500-1200)
Montenegro	2000	Male	45 to 49	79% (56-91%)	21% (8.7-44%)	5.6 (3-9.2)	14% (8.7-19%)	20% (11-26%)	26 (15-35)	1200 (710-1600)
Montenegro	2000	Male	50 to 54	77% (56-90%)	23% (9.6-44%)	5.9 (3.3-9.4)	15% (9.2-20%)	19% (12-25%)	29 (17-38)	1100 (710-1500)
Montenegro	2000	Male	55 to 59	76% (54-91%)	24% (9-46%)	6 (3.3-10)	15% (9.6-20%)	19% (12-25%)	32 (20-43)	1100 (700-1500)
Montenegro	2000	Male	60 to 64	74% (51-91%)	26% (9.4-49%)	5.8 (3-10)	14% (8.2-19%)	17% (9.9-24%)	52 (31-73)	1500 (880-2100)
Montenegro	2000	Male	65 to 69	73% (50-90%)	27% (10-50%)	5.2 (2.9-8.4)	14% (7.5-19%)	17% (8.9-23%)	67 (37-95)	1600 (900-2200)
Montenegro	2000	Male	70 to 74	71% (49-87%)	29% (13-51%)	4.5 (2.3-8)	12% (5.4-19%)	15% (6.4-22%)	67 (29-100)	1300 (560-1900)
Montenegro	2000	Male	75 to 79	68% (46-86%)	32% (14-54%)	3.8 (1.8-7.1)	11% (3.7-18%)	13% (4.1-21%)	52 (16-87)	760 (250-1300)
Montenegro	2000	Male	80 to 84	67% (43-84%)	33% (16-57%)	3.4 (1.6-6)	11% (3.6-18%)	12% (4.1-21%)	33 (11-58)	350 (120-610)
Montenegro	2000	Male	85 to 89	65% (43-83%)	35% (17-57%)	3.1 (1.5-5.6)	9.2% (2.5-17%)	11% (2.8-20%)	18 (4.6-34)	140 (37-250)
Montenegro	2000	Male	90 to 94	65% (43-85%)	35% (15-57%)	2.9 (1.3-5.4)	7.3% (1.2-15%)	8.4% (1.2-18%)	9.1 (1.3-19)	48 (7.6-99)
Montenegro	2000	Male	95 plus	65% (38-88%)	35% (12-62%)	2.6 (0.92-5.7)	6% (0.042-14%)	6.8% (-0.29-16%)	2.6 (-0.11-6.2)	9.1 (0.064-21)
Montenegro	2005	Female	All Ages	58% (50-66%)	42% (34-50%)	2.3 (1.7-3)	7.9% (5-11%)	12% (6.7-18%)	310 (170-460)	6700 (4300-9200)
Montenegro	2005	Female	15 to 19	58% (38-75%)	42% (25-62%)	1.7 (0.72-3.2)	3.4% (2-5.4%)	8.7% (4.2-15%)	0.65 (0.32-1.1)	72 (41-110)
Montenegro	2005	Female	20 to 24	67% (50-82%)	33% (18-50%)	2.3 (0.99-4.4)	5.4% (3.2-8.1%)	11% (4.8-18%)	1.2 (0.53-2)	140 (81-210)
Montenegro	2005	Female	25 to 29	69% (50-83%)	31% (17-50%)	2.5 (1.1-4.6)	5.6% (3.5-8.1%)	12% (5.5-19%)	1.4 (0.65-2.3)	150 (91-220)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	2005	Female	30 to 34	70% (50-84%)	30% (16-50%)	2.7 (1.3-4.8)	5.7% (3.6-8.4%)	13% (6.9-20%)	2 (1.1-3.1)	170 (100-250)
Montenegro	2005	Female	35 to 39	70% (53-84%)	30% (16-47%)	2.9 (1.4-5.3)	6.5% (3.7-9.6%)	14% (7.2-21%)	3.1 (1.6-4.8)	210 (120-310)
Montenegro	2005	Female	40 to 44	69% (52-84%)	31% (16-48%)	3.3 (1.6-5.6)	7.6% (4.4-11%)	15% (8.3-23%)	5.7 (3.1-8.6)	320 (180-470)
Montenegro	2005	Female	45 to 49	68% (45-83%)	32% (17-55%)	3.6 (1.8-6.8)	8.8% (5-13%)	16% (9.1-24%)	9.5 (5.4-14)	460 (260-670)
Montenegro	2005	Female	50 to 54	65% (45-81%)	35% (19-55%)	3.7 (1.9-6.4)	9.5% (5.2-14%)	16% (8.9-23%)	15 (8.6-22)	630 (360-910)
Montenegro	2005	Female	55 to 59	62% (41-79%)	38% (21-59%)	3.6 (1.8-6.4)	10% (5.3-15%)	16% (8.6-23%)	17 (9.6-26)	610 (330-910)
Montenegro	2005	Female	60 to 64	58% (37-78%)	42% (22-63%)	3.3 (1.5-6.3)	9.6% (4.5-15%)	14% (6.9-22%)	19 (9.2-30)	570 (270-910)
Montenegro	2005	Female	65 to 69	54% (34-74%)	46% (26-66%)	2.9 (1.4-5.1)	9.8% (4.4-15%)	14% (6.3-22%)	36 (16-56)	860 (390-1400)
Montenegro	2005	Female	70 to 74	47% (29-66%)	53% (34-71%)	2.3 (0.97-4.3)	9.3% (3-16%)	12% (4-21%)	48 (16-84)	920 (300-1600)
Montenegro	2005	Female	75 to 79	40% (26-60%)	60% (40-74%)	1.9 (0.73-3.7)	8.5% (2.5-15%)	11% (3.1-19%)	50 (14-89)	740 (210-1300)
Montenegro	2005	Female	80 to 84	36% (21-54%)	64% (46-79%)	1.7 (0.66-3.2)	8.6% (3.1-15%)	11% (4-19%)	41 (15-71)	440 (160-760)
Montenegro	2005	Female	85 to 89	34% (21-51%)	66% (49-79%)	1.7 (0.64-3.4)	9% (3.5-16%)	11% (4.2-19%)	30 (12-52)	230 (88-390)
Montenegro	2005	Female	90 to 94	36% (23-53%)	64% (47-77%)	1.8 (0.64-3.9)	8.8% (2.9-16%)	11% (3.5-19%)	22 (7.1-39)	110 (36-200)
Montenegro	2005	Female	95 plus	40% (23-61%)	60% (39-77%)	2 (0.56-4.7)	8.7% (1.9-17%)	11% (2.5-20%)	11 (2.7-21)	37 (8.1-71)
Montenegro	2005	Male	All Ages	67% (56-78%)	33% (22-44%)	3.6 (2.7-4.7)	12% (8.9-15%)	15% (10-21%)	460 (290-620)	12000 (8900-15000)
Montenegro	2005	Male	15 to 19	59% (36-80%)	41% (20-64%)	1.6 (0.79-2.9)	4.5% (2.9-6.8%)	8.2% (4.4-14%)	0.96 (0.52-1.6)	110 (69-170)
Montenegro	2005	Male	20 to 24	69% (48-86%)	31% (14-52%)	2.9 (1.4-4.9)	9.6% (6.3-14%)	14% (7.8-22%)	3.8 (2.1-6)	350 (220-510)
Montenegro	2005	Male	25 to 29	73% (52-89%)	27% (11-48%)	3.7 (1.9-6.4)	11% (7.4-16%)	17% (10-26%)	4.7 (2.7-7.1)	390 (260-560)
Montenegro	2005	Male	30 to 34	76% (55-91%)	24% (9.3-45%)	4.2 (2.3-7)	11% (7.9-16%)	19% (12-27%)	4.8 (3-6.9)	370 (250-510)
Montenegro	2005	Male	35 to 39	78% (59-91%)	22% (9.3-41%)	4.5 (2.4-7.3)	12% (7.9-17%)	19% (12-27%)	6.9 (4.1-9.8)	450 (290-610)
Montenegro	2005	Male	40 to 44	79% (58-91%)	21% (8.5-42%)	5 (2.8-7.8)	13% (8.3-18%)	20% (12-27%)	12 (7.5-17)	680 (440-920)
Montenegro	2005	Male	45 to 49	78% (57-91%)	22% (8.6-43%)	5.8 (3.3-9)	15% (9.4-19%)	20% (13-26%)	22 (14-30)	1000 (690-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	2005	Male	50 to 54	77% (56-91%)	23% (9.4-44%)	6.2 (3.7-9.4)	15% (10-19%)	20% (13-25%)	34 (22-45)	1400 (940-1800)
Montenegro	2005	Male	55 to 59	76% (54-91%)	24% (9.2-46%)	6.4 (3.7-10)	16% (10-20%)	19% (13-25%)	44 (30-58)	1500 (1000-2000)
Montenegro	2005	Male	60 to 64	74% (52-91%)	26% (9.1-48%)	6.2 (3.3-10)	15% (9.1-20%)	18% (11-24%)	44 (27-59)	1300 (780-1700)
Montenegro	2005	Male	65 to 69	73% (50-90%)	27% (10-50%)	5.5 (3.2-8.8)	14% (8.4-19%)	17% (10-23%)	70 (40-95)	1700 (980-2300)
Montenegro	2005	Male	70 to 74	71% (50-86%)	29% (14-50%)	4.7 (2.5-8)	13% (6-19%)	15% (7.1-22%)	71 (34-100)	1400 (660-2000)
Montenegro	2005	Male	75 to 79	68% (44-86%)	32% (14-56%)	4 (1.9-7.2)	12% (4.4-18%)	14% (5-21%)	66 (24-110)	960 (370-1500)
Montenegro	2005	Male	80 to 84	66% (42-84%)	34% (16-58%)	3.6 (1.7-6.1)	11% (4.2-18%)	13% (4.7-21%)	42 (15-70)	450 (170-740)
Montenegro	2005	Male	85 to 89	64% (42-82%)	36% (18-58%)	3.3 (1.6-5.6)	10% (3.5-17%)	12% (3.9-20%)	20 (6.6-35)	150 (52-260)
Montenegro	2005	Male	90 to 94	64% (42-84%)	36% (16-58%)	3 (1.4-5.5)	8.1% (1.6-16%)	9.4% (1.8-18%)	6.8 (1.3-13)	35 (7.2-68)
Montenegro	2005	Male	95 plus	64% (38-87%)	36% (13-62%)	2.8 (0.99-5.7)	6.8% (0.29-14%)	7.8% (0.096-17%)	2.3 (0.028-5)	8 (0.36-17)
Montenegro	2010	Female	All Ages	58% (50-66%)	42% (34-50%)	2.4 (1.8-3.1)	8.1% (5.4-11%)	12% (7.4-18%)	310 (180-460)	6600 (4400-8900)
Montenegro	2010	Female	15 to 19	57% (37-75%)	43% (25-63%)	1.7 (0.74-3.3)	3.2% (1.9-5.1%)	9% (4.3-16%)	0.51 (0.25-0.89)	61 (36-98)
Montenegro	2010	Female	20 to 24	68% (50-83%)	32% (17-50%)	2.5 (1-4.6)	5.3% (3.2-7.9%)	11% (5-19%)	0.86 (0.39-1.5)	110 (66-170)
Montenegro	2010	Female	25 to 29	69% (49-84%)	31% (16-51%)	2.6 (1.2-4.9)	5.5% (3.5-7.8%)	12% (6-19%)	1.3 (0.61-2.1)	150 (88-210)
Montenegro	2010	Female	30 to 34	70% (50-84%)	30% (16-50%)	2.8 (1.4-5.1)	5.6% (3.7-8.1%)	13% (7.4-20%)	1.8 (0.99-2.9)	170 (110-240)
Montenegro	2010	Female	35 to 39	70% (52-85%)	30% (15-48%)	3.1 (1.5-5.5)	6.4% (3.8-9.3%)	14% (7.6-21%)	3 (1.6-4.7)	220 (130-310)
Montenegro	2010	Female	40 to 44	69% (52-84%)	31% (16-48%)	3.5 (1.6-5.9)	7.6% (4.5-11%)	16% (8.9-23%)	5.1 (2.8-7.6)	290 (170-420)
Montenegro	2010	Female	45 to 49	68% (46-83%)	32% (17-54%)	3.8 (1.9-6.9)	8.6% (5-13%)	16% (9.5-23%)	9.1 (5.4-13)	450 (270-640)
Montenegro	2010	Female	50 to 54	65% (45-81%)	35% (19-55%)	3.9 (2-6.8)	9.4% (5.5-14%)	16% (9.9-23%)	13 (7.7-19)	560 (330-790)
Montenegro	2010	Female	55 to 59	62% (42-78%)	38% (22-58%)	3.8 (1.9-6.7)	9.8% (5.5-14%)	16% (9.1-23%)	21 (12-30)	740 (410-1100)
Montenegro	2010	Female	60 to 64	58% (37-77%)	42% (23-63%)	3.5 (1.7-6.7)	9.6% (4.6-15%)	15% (6.9-22%)	22 (10-33)	650 (310-1000)
Montenegro	2010	Female	65 to 69	54% (34-75%)	46% (25-66%)	3 (1.5-5.2)	10% (4.9-16%)	15% (7-22%)	28 (14-43)	690 (330-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	2010	Female	70 to 74	47% (30-66%)	53% (34-70%)	2.4 (1.1-4.7)	9.4% (3.4-16%)	13% (4.7-22%)	46 (17-79)	890 (320-1500)
Montenegro	2010	Female	75 to 79	40% (26-60%)	60% (40-74%)	2 (0.8-3.9)	8.6% (2.9-15%)	11% (3.6-20%)	48 (15-85)	710 (220-1300)
Montenegro	2010	Female	80 to 84	36% (21-54%)	64% (46-79%)	1.8 (0.74-3.3)	8.8% (3.4-15%)	11% (4.3-19%)	46 (17-77)	500 (190-830)
Montenegro	2010	Female	85 to 89	34% (21-50%)	66% (50-79%)	1.8 (0.7-3.5)	9.1% (3.6-15%)	11% (4.5-19%)	40 (16-69)	300 (120-510)
Montenegro	2010	Female	90 to 94	36% (22-53%)	64% (47-78%)	1.9 (0.7-3.8)	8.9% (3.2-16%)	11% (4-19%)	19 (6.7-34)	96 (34-170)
Montenegro	2010	Female	95 plus	39% (22-60%)	61% (40-78%)	2.1 (0.61-4.6)	8.8% (2.3-16%)	11% (2.8-20%)	8.5 (2.3-16)	28 (7.3-53)
Montenegro	2010	Male	All Ages	67% (56-77%)	33% (23-44%)	3.8 (3-4.9)	12% (9.4-15%)	16% (10-21%)	460 (310-610)	12000 (9200-15000)
Montenegro	2010	Male	15 to 19	58% (36-79%)	42% (21-64%)	1.7 (0.83-3)	4.9% (3.1-7.2%)	8.7% (4.7-14%)	1.1 (0.56-1.8)	110 (71-180)
Montenegro	2010	Male	20 to 24	69% (48-87%)	31% (13-52%)	3.1 (1.6-5.3)	10% (6.6-15%)	15% (8.5-23%)	3.4 (1.9-5.4)	310 (200-460)
Montenegro	2010	Male	25 to 29	73% (53-89%)	27% (11-47%)	3.9 (2-6.6)	12% (7.7-16%)	18% (11-27%)	5.1 (3.1-7.7)	430 (280-610)
Montenegro	2010	Male	30 to 34	76% (54-91%)	24% (9.4-46%)	4.5 (2.6-7.5)	12% (8.4-16%)	20% (13-28%)	5.4 (3.5-7.6)	420 (280-560)
Montenegro	2010	Male	35 to 39	78% (59-91%)	22% (9-41%)	4.8 (2.6-7.6)	13% (8.4-17%)	21% (13-28%)	7.8 (4.8-11)	510 (330-680)
Montenegro	2010	Male	40 to 44	79% (58-92%)	21% (8.4-42%)	5.4 (3.1-8.2)	14% (9.3-18%)	21% (14-28%)	12 (7.9-17)	680 (450-890)
Montenegro	2010	Male	45 to 49	78% (54-91%)	22% (8.7-46%)	6.2 (3.6-9.6)	15% (10-19%)	21% (14-27%)	21 (14-28)	1000 (700-1300)
Montenegro	2010	Male	50 to 54	77% (55-91%)	23% (9.3-45%)	6.5 (4-9.7)	15% (11-19%)	20% (14-25%)	30 (21-39)	1200 (860-1600)
Montenegro	2010	Male	55 to 59	75% (54-91%)	25% (9.3-46%)	6.7 (3.9-10)	15% (11-20%)	20% (14-25%)	51 (36-68)	1800 (1300-2300)
Montenegro	2010	Male	60 to 64	74% (50-90%)	26% (9.5-50%)	6.5 (3.5-10)	15% (9.6-19%)	18% (12-24%)	48 (32-64)	1400 (930-1900)
Montenegro	2010	Male	65 to 69	73% (49-89%)	27% (11-51%)	5.8 (3.3-8.9)	14% (8.7-19%)	18% (11-23%)	48 (29-64)	1200 (710-1500)
Montenegro	2010	Male	70 to 74	71% (48-87%)	29% (13-52%)	5 (2.7-8.2)	13% (6.6-18%)	16% (7.8-22%)	68 (34-98)	1300 (680-1900)
Montenegro	2010	Male	75 to 79	68% (45-86%)	32% (14-55%)	4.2 (2.1-7.5)	12% (5.1-18%)	14% (5.8-21%)	66 (28-100)	980 (410-1500)
Montenegro	2010	Male	80 to 84	66% (41-85%)	34% (15-59%)	3.8 (1.8-6.3)	11% (4.9-18%)	14% (5.6-21%)	49 (20-77)	530 (220-840)
Montenegro	2010	Male	85 to 89	64% (42-83%)	36% (17-58%)	3.4 (1.7-5.9)	10% (3.6-17%)	12% (4.1-21%)	30 (10-52)	230 (82-390)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	2010	Male	90 to 94	64% (44-84%)	36% (16-56%)	3.2 (1.5-5.7)	8.3% (1.9-15%)	9.7% (2.1-18%)	9 (2-17)	47 (11-88)
Montenegro	2010	Male	95 plus	65% (39-87%)	35% (13-61%)	2.9 (1.1-5.7)	7% (0.67-14%)	8.1% (0.54-17%)	1.9 (0.13-3.9)	6.7 (0.66-13)
Montenegro	2016	Female	All Ages	57% (49-66%)	43% (34-51%)	2.6 (1.8-3.5)	8.2% (5.1-11%)	12% (7.2-18%)	350 (190-520)	6900 (4300-9500)
Montenegro	2016	Female	15 to 19	57% (38-75%)	43% (25-62%)	1.8 (0.74-3.5)	3.2% (1.8-5.2%)	9.3% (4.4-16%)	0.46 (0.22-0.84)	57 (32-94)
Montenegro	2016	Female	20 to 24	67% (48-82%)	33% (18-52%)	2.6 (1-4.9)	5.1% (3.2-7.7%)	11% (5-19%)	0.74 (0.32-1.3)	100 (61-160)
Montenegro	2016	Female	25 to 29	69% (50-83%)	31% (17-50%)	2.8 (1.2-5.3)	5.3% (3.4-7.6%)	12% (5.7-20%)	0.91 (0.43-1.5)	120 (70-170)
Montenegro	2016	Female	30 to 34	70% (49-83%)	30% (17-51%)	3 (1.4-5.4)	5.4% (3.5-7.6%)	13% (7.2-20%)	1.6 (0.83-2.4)	160 (97-220)
Montenegro	2016	Female	35 to 39	70% (52-85%)	30% (15-48%)	3.2 (1.4-5.7)	6.1% (3.6-8.8%)	14% (7.6-21%)	2.7 (1.4-4.1)	200 (120-290)
Montenegro	2016	Female	40 to 44	69% (51-83%)	31% (17-49%)	3.6 (1.7-6.3)	7.3% (4.2-11%)	16% (8.4-23%)	4.8 (2.6-7.2)	290 (160-420)
Montenegro	2016	Female	45 to 49	67% (46-82%)	33% (18-54%)	4 (1.8-7.6)	8.5% (4.9-12%)	16% (9.5-23%)	7.7 (4.4-11)	390 (220-560)
Montenegro	2016	Female	50 to 54	64% (44-81%)	36% (19-56%)	4.1 (2-7.3)	9.4% (5.6-14%)	16% (9.8-23%)	13 (7.7-19)	560 (320-810)
Montenegro	2016	Female	55 to 59	61% (42-79%)	39% (21-58%)	4 (1.9-7)	9.8% (5.4-14%)	16% (9.1-23%)	18 (10-27)	660 (360-970)
Montenegro	2016	Female	60 to 64	58% (37-77%)	42% (23-63%)	3.7 (1.7-6.9)	9.5% (4.7-15%)	15% (7.5-23%)	27 (13-42)	830 (410-1300)
Montenegro	2016	Female	65 to 69	54% (34-74%)	46% (26-66%)	3.2 (1.5-5.6)	10% (4.9-15%)	15% (7.2-22%)	34 (16-53)	840 (400-1300)
Montenegro	2016	Female	70 to 74	47% (30-65%)	53% (35-70%)	2.6 (1.1-4.9)	9.4% (3.3-15%)	13% (4.9-21%)	36 (13-62)	710 (240-1200)
Montenegro	2016	Female	75 to 79	40% (25-58%)	60% (42-75%)	2.1 (0.79-4.1)	8.6% (2.7-15%)	11% (3.6-19%)	54 (17-94)	800 (250-1400)
Montenegro	2016	Female	80 to 84	35% (21-54%)	65% (46-79%)	1.9 (0.77-3.6)	8.9% (3.5-15%)	11% (4.4-19%)	52 (19-87)	570 (210-950)
Montenegro	2016	Female	85 to 89	34% (21-50%)	66% (50-79%)	1.9 (0.73-3.7)	9.2% (3.4-15%)	11% (4.2-19%)	51 (19-88)	390 (150-660)
Montenegro	2016	Female	90 to 94	36% (22-52%)	64% (48-78%)	2 (0.72-4.4)	9.1% (3.4-16%)	11% (4.1-19%)	32 (12-58)	160 (61-290)
Montenegro	2016	Female	95 plus	39% (22-60%)	61% (40-78%)	2.2 (0.62-5)	9% (2.5-16%)	11% (3.1-21%)	9.7 (2.8-18)	32 (8.9-59)
Montenegro	2016	Male	All Ages	67% (57-77%)	33% (23-43%)	4.1 (2.9-5.6)	13% (9.5-16%)	16% (11-21%)	500 (330-680)	13000 (9400-16000)
Montenegro	2016	Male	15 to 19	58% (36-79%)	42% (21-64%)	1.8 (0.84-3.3)	5% (3.1-7.8%)	9.1% (4.7-16%)	1 (0.53-1.8)	110 (69-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Montenegro	2016	Male	20 to 24	69% (47-87%)	31% (13-53%)	3.2 (1.6-5.7)	10% (6.8-15%)	16% (8.8-25%)	3.3 (1.8-5.2)	310 (190-460)
Montenegro	2016	Male	25 to 29	73% (53-89%)	27% (11-47%)	4.1 (2.1-7.4)	12% (8-17%)	19% (11-29%)	4.2 (2.5-6.5)	370 (240-520)
Montenegro	2016	Male	30 to 34	75% (54-91%)	25% (9.4-46%)	4.7 (2.5-8)	12% (8.3-16%)	21% (13-29%)	5.8 (3.6-8)	450 (300-610)
Montenegro	2016	Male	35 to 39	78% (58-90%)	22% (9.6-42%)	5 (2.6-8.4)	13% (8.6-17%)	21% (13-29%)	8.2 (5-11)	540 (360-710)
Montenegro	2016	Male	40 to 44	79% (59-92%)	21% (8.5-41%)	5.7 (3.1-9.2)	14% (9.4-19%)	22% (14-29%)	13 (8-17)	700 (460-900)
Montenegro	2016	Male	45 to 49	78% (57-91%)	22% (8.8-43%)	6.5 (3.5-11)	15% (10-19%)	22% (14-27%)	19 (12-24)	900 (600-1100)
Montenegro	2016	Male	50 to 54	77% (57-91%)	23% (9.4-43%)	7 (4-11)	15% (11-20%)	21% (15-26%)	30 (21-40)	1200 (870-1600)
Montenegro	2016	Male	55 to 59	75% (54-90%)	25% (9.7-46%)	7.1 (3.9-11)	15% (11-20%)	20% (14-26%)	44 (31-60)	1600 (1100-2100)
Montenegro	2016	Male	60 to 64	74% (50-90%)	26% (9.8-50%)	6.8 (3.6-11)	15% (9.8-19%)	18% (12-24%)	63 (41-85)	1900 (1200-2500)
Montenegro	2016	Male	65 to 69	73% (49-89%)	27% (11-51%)	6.2 (3.3-9.5)	14% (9-19%)	18% (11-24%)	61 (38-85)	1500 (970-2000)
Montenegro	2016	Male	70 to 74	70% (48-86%)	30% (14-52%)	5.3 (2.7-8.9)	13% (6.8-18%)	16% (8.3-23%)	51 (26-73)	990 (510-1400)
Montenegro	2016	Male	75 to 79	68% (44-85%)	32% (15-56%)	4.5 (2.1-7.9)	12% (4.8-18%)	14% (5.8-22%)	70 (29-110)	1100 (450-1600)
Montenegro	2016	Male	80 to 84	66% (41-84%)	34% (16-59%)	4 (2-6.8)	12% (5.1-19%)	14% (5.9-22%)	63 (27-100)	680 (300-1100)
Montenegro	2016	Male	85 to 89	64% (41-83%)	36% (17-59%)	3.7 (1.8-6.4)	11% (3.9-17%)	13% (4.3-21%)	43 (15-73)	330 (120-550)
Montenegro	2016	Male	90 to 94	64% (43-84%)	36% (16-57%)	3.4 (1.6-6.6)	8.9% (2.4-16%)	10% (2.8-19%)	18 (4.6-32)	93 (25-170)
Montenegro	2016	Male	95 plus	65% (39-87%)	35% (13-61%)	3.1 (1.1-6.3)	7.6% (0.78-15%)	8.8% (0.74-18%)	3.6 (0.3-7.1)	13 (1.3-24)
Poland	1990	Female	All Ages	76% (71-81%)	24% (19-29%)	1.5 (1.4-1.7)	2% (0.66-3.4%)	1.1% (-1.4-3.6%)	2000 (-2600-6600)	120000 (36000-190000)
Poland	1990	Female	15 to 19	76% (63-87%)	24% (13-37%)	1.4 (0.89-2.1)	2.5% (1.5-3.7%)	6.3% (3.2-9.8%)	30 (15-47)	3300 (2000-4900)
Poland	1990	Female	20 to 24	83% (71-90%)	17% (9.8-29%)	1.8 (1.1-2.5)	3.2% (2.2-4.5%)	6.3% (3.2-10%)	29 (15-46)	4000 (2700-5700)
Poland	1990	Female	25 to 29	84% (71-91%)	16% (8.7-29%)	1.7 (1.2-2.4)	3.3% (2.3-4.5%)	6% (3.4-9%)	38 (22-57)	5000 (3400-6900)
Poland	1990	Female	30 to 34	84% (71-92%)	16% (8.1-29%)	1.8 (1.2-2.5)	3.4% (2.4-4.7%)	6.5% (3.8-9.5%)	81 (47-120)	7900 (5300-11000)
Poland	1990	Female	35 to 39	84% (73-92%)	16% (8.2-27%)	1.9 (1.3-2.6)	3.9% (2.6-5.3%)	7.2% (4.5-10%)	150 (91-210)	11000 (7400-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	1990	Female	40 to 44	84% (75-91%)	16% (8.6-25%)	2.2 (1.5-3)	4.3% (2.7-6%)	7.6% (4.7-11%)	220 (140-310)	13000 (8300-18000)
Poland	1990	Female	45 to 49	85% (77-90%)	15% (9.7-23%)	2.6 (1.8-3.6)	4.9% (3-7.1%)	8.3% (5.2-12%)	250 (160-350)	12000 (7500-17000)
Poland	1990	Female	50 to 54	85% (78-90%)	15% (10-22%)	2.8 (1.9-3.7)	4.8% (2.8-7%)	7.7% (4.5-11%)	380 (230-550)	16000 (9200-23000)
Poland	1990	Female	55 to 59	84% (78-89%)	16% (11-22%)	2.7 (1.9-3.6)	4.2% (2.1-6.5%)	6.3% (3.2-9.5%)	510 (260-780)	18000 (8900-28000)
Poland	1990	Female	60 to 64	81% (74-86%)	19% (14-26%)	2.5 (1.7-3.4)	3.1% (0.52-6%)	4.3% (0.72-8.1%)	550 (90-1100)	16000 (2700-32000)
Poland	1990	Female	65 to 69	75% (66-82%)	25% (18-34%)	2.2 (1.5-3)	1.9% (-0.64-4.4%)	2.5% (-0.9-5.9%)	460 (-170-1100)	11000 (-3700-26000)
Poland	1990	Female	70 to 74	67% (57-75%)	33% (25-43%)	1.8 (1.2-2.6)	0.92% (-1.8-3.7%)	1.1% (-2.5-4.5%)	180 (-420-760)	3800 (-7700-15000)
Poland	1990	Female	75 to 79	58% (46-70%)	42% (30-54%)	1.5 (0.92-2.2)	0.11% (-2.6-2.9%)	0.017% (-3.4-3.5%)	5.4 (-1000-1100)	610 (-14000-16000)
Poland	1990	Female	80 to 84	53% (40-64%)	47% (36-60%)	1.3 (0.83-1.9)	-0.52% (-3-2%)	-0.77% (-3.8-2.3%)	-270 (-1300-800)	-2400 (-13000-8900)
Poland	1990	Female	85 to 89	52% (39-64%)	48% (36-61%)	1.2 (0.74-1.9)	-1.1% (-3.7-1.4%)	-1.5% (-4.6-1.5%)	-370 (-1200-400)	-2500 (-8400-3200)
Poland	1990	Female	90 to 94	54% (41-67%)	46% (33-59%)	1.3 (0.73-2)	-1.5% (-4.2-1.4%)	-2% (-5.3-1.4%)	-210 (-570-150)	-990 (-2700-880)
Poland	1990	Female	95 plus	58% (41-73%)	42% (27-59%)	1.3 (0.66-2.2)	-1.9% (-5.1-1.6%)	-2.5% (-6.5-1.7%)	-67 (-170-45)	-210 (-560-180)
Poland	1990	Male	All Ages	92% (89-94%)	8% (5.7-11%)	2.9 (2.6-3.2)	10% (7.9-12%)	10% (6.8-14%)	21000 (14000-29000)	760000 (600000-920000)
Poland	1990	Male	15 to 19	88% (78-95%)	12% (5.4-22%)	1.5 (0.99-2.3)	5.5% (3.3-8.2%)	9% (4.7-14%)	130 (69-200)	11000 (6700-16000)
Poland	1990	Male	20 to 24	93% (86-97%)	6.9% (3.1-14%)	2.5 (1.7-3.6)	10% (6.6-14%)	14% (8.3-21%)	290 (170-430)	23000 (15000-32000)
Poland	1990	Male	25 to 29	94% (89-97%)	5.6% (2.5-11%)	3.1 (2.1-4.3)	12% (8.9-16%)	17% (11-23%)	420 (280-580)	33000 (23000-43000)
Poland	1990	Male	30 to 34	95% (90-98%)	4.8% (2.1-9.7%)	3.3 (2.4-4.6)	13% (9.6-16%)	17% (12-23%)	730 (510-970)	51000 (38000-66000)
Poland	1990	Male	35 to 39	96% (91-98%)	4.3% (1.8-8.8%)	3.2 (2.1-4.3)	12% (8.5-15%)	15% (10-20%)	920 (630-1200)	57000 (42000-74000)
Poland	1990	Male	40 to 44	96% (92-98%)	3.8% (1.5-8.2%)	3.9 (2.8-5.1)	12% (8.7-16%)	15% (10-21%)	1200 (800-1600)	64000 (45000-85000)
Poland	1990	Male	45 to 49	97% (93-99%)	3.1% (1.4-6.6%)	5.5 (3.7-7.3)	16% (10-21%)	19% (12-25%)	1500 (950-2000)	69000 (45000-90000)
Poland	1990	Male	50 to 54	97% (94-98%)	3.3% (1.8-6.2%)	6.2 (4.4-8)	16% (11-20%)	19% (13-24%)	2500 (1700-3200)	99000 (69000-120000)
Poland	1990	Male	55 to 59	96% (93-97%)	4.5% (2.6-7.4%)	6.4 (4.8-8.2)	15% (11-19%)	17% (12-22%)	3300 (2300-4300)	110000 (80000-140000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	1990	Male	60 to 64	95% (92-97%)	5.1% (3-8.4%)	6.1 (4.5-8.2)	13% (8-17%)	14% (9-20%)	3600 (2200-4900)	100000 (65000-140000)
Poland	1990	Male	65 to 69	95% (91-97%)	5.3% (3.1-8.5%)	5.5 (4.1-7.1)	10% (6-15%)	12% (6.6-17%)	3100 (1700-4400)	73000 (42000-100000)
Poland	1990	Male	70 to 74	94% (91-96%)	5.8% (3.6-9.2%)	4.7 (3.3-6.3)	7.6% (2.7-12%)	8.4% (2.7-14%)	1700 (530-2800)	32000 (12000-53000)
Poland	1990	Male	75 to 79	93% (89-96%)	6.7% (3.8-11%)	3.9 (2.6-5.5)	4.9% (-0.023-10%)	5.2% (-0.5-12%)	1400 (-130-3200)	22000 (-100-47000)
Poland	1990	Male	80 to 84	93% (87-96%)	7.5% (4.1-13%)	3.4 (2.3-4.7)	2.6% (-1.7-7.8%)	2.6% (-2.5-8.7%)	610 (-590-2100)	7500 (-4800-23000)
Poland	1990	Male	85 to 89	92% (86-96%)	8% (4.5-14%)	3 (2-4.3)	0.4% (-3.6-5.3%)	-0.017% (-4.8-5.8%)	-2.2 (-580-700)	420 (-3700-5600)
Poland	1990	Male	90 to 94	92% (86-96%)	8.2% (4.3-14%)	2.7 (1.7-3.9)	-1.2% (-5.4-3.6%)	-2% (-6.9-3.8%)	-73 (-250-140)	-270 (-1200-790)
Poland	1990	Male	95 plus	92% (84-97%)	8.4% (3.5-16%)	2.3 (1.2-3.7)	-2.1% (-6-3.1%)	-3.1% (-7.9-3.1%)	-25 (-65-25)	-69 (-200-110)
Poland	1995	Female	All Ages	77% (72-81%)	23% (19-28%)	1.6 (1.4-1.8)	2.2% (0.86-3.6%)	1.2% (-1.2-3.7%)	2200 (-2300-6800)	120000 (48000-200000)
Poland	1995	Female	15 to 19	77% (63-87%)	23% (13-37%)	1.4 (0.9-2)	2.5% (1.5-3.8%)	6.6% (3.3-10%)	35 (17-55)	3800 (2300-5600)
Poland	1995	Female	20 to 24	83% (71-90%)	17% (9.7-29%)	1.7 (1.1-2.5)	3.3% (2.2-4.6%)	6.7% (3.6-10%)	32 (17-50)	4500 (3000-6400)
Poland	1995	Female	25 to 29	84% (72-91%)	16% (8.6-28%)	1.7 (1.2-2.4)	3.4% (2.4-4.5%)	6.6% (3.7-9.7%)	34 (19-50)	4500 (3100-6000)
Poland	1995	Female	30 to 34	84% (71-92%)	16% (8.4-29%)	1.8 (1.2-2.5)	3.6% (2.6-4.8%)	7.5% (4.8-11%)	64 (41-91)	6300 (4500-8400)
Poland	1995	Female	35 to 39	84% (73-92%)	16% (8.4-27%)	1.9 (1.2-2.6)	4.1% (2.9-5.5%)	8.2% (5.4-11%)	150 (98-210)	11000 (7700-15000)
Poland	1995	Female	40 to 44	84% (74-91%)	16% (8.8-26%)	2.2 (1.5-2.9)	4.6% (3-6.3%)	8.4% (5.5-11%)	270 (170-360)	15000 (11000-20000)
Poland	1995	Female	45 to 49	85% (77-91%)	15% (9.4-23%)	2.6 (1.8-3.5)	5.2% (3.4-7.2%)	8.9% (5.9-12%)	380 (250-530)	18000 (12000-25000)
Poland	1995	Female	50 to 54	85% (79-90%)	15% (10-21%)	2.8 (1.9-3.6)	5.2% (3.3-7.3%)	8.4% (5.3-12%)	350 (220-480)	14000 (9000-20000)
Poland	1995	Female	55 to 59	85% (78-89%)	15% (11-22%)	2.7 (1.9-3.6)	4.4% (2.4-6.7%)	6.8% (3.8-10%)	460 (260-690)	16000 (9000-25000)
Poland	1995	Female	60 to 64	81% (73-86%)	19% (14-27%)	2.5 (1.7-3.4)	3.2% (0.83-6%)	4.6% (1.2-8.4%)	520 (140-920)	15000 (4200-28000)
Poland	1995	Female	65 to 69	75% (66-81%)	25% (19-34%)	2.2 (1.5-2.9)	2.1% (-0.44-4.6%)	2.8% (-0.6-6.1%)	500 (-110-1100)	12000 (-2500-26000)
Poland	1995	Female	70 to 74	68% (58-75%)	32% (25-42%)	1.8 (1.2-2.6)	1% (-1.6-3.7%)	1.3% (-2.2-4.7%)	310 (-530-1200)	6300 (-10000-22000)
Poland	1995	Female	75 to 79	59% (46-70%)	41% (30-54%)	1.5 (0.93-2.2)	0.22% (-2.4-3%)	0.15% (-3.1-3.6%)	34 (-700-810)	880 (-10000-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	1995	Female	80 to 84	53% (40-64%)	47% (36-60%)	1.3 (0.82-1.9)	-0.41% (-2.8-1.9%)	-0.64% (-3.6-2.2%)	-220 (-1200-760)	-1800 (-13000-8500)
Poland	1995	Female	85 to 89	53% (40-64%)	47% (36-60%)	1.2 (0.76-1.9)	-0.98% (-3.6-1.4%)	-1.3% (-4.4-1.5%)	-400 (-1300-460)	-2600 (-9100-3700)
Poland	1995	Female	90 to 94	55% (43-67%)	45% (33-57%)	1.3 (0.75-2)	-1.4% (-4.1-1.4%)	-1.9% (-5.2-1.5%)	-250 (-700-210)	-1200 (-3400-1100)
Poland	1995	Female	95 plus	58% (42-73%)	42% (27-58%)	1.3 (0.69-2.2)	-1.8% (-5.1-1.8%)	-2.4% (-6.4-1.9%)	-81 (-220-64)	-250 (-700-250)
Poland	1995	Male	All Ages	93% (90-95%)	7.5% (5.3-10%)	3 (2.7-3.3)	11% (8.9-13%)	11% (7.7-14%)	22000 (16000-29000)	800000 (650000-960000)
Poland	1995	Male	15 to 19	89% (80-95%)	11% (5.1-20%)	1.6 (0.98-2.2)	5.5% (3.2-8%)	9.1% (4.8-14%)	140 (73-210)	12000 (7000-17000)
Poland	1995	Male	20 to 24	93% (87-97%)	6.7% (3.1-13%)	2.5 (1.7-3.6)	10% (6.6-14%)	15% (8.4-21%)	290 (170-420)	24000 (15000-33000)
Poland	1995	Male	25 to 29	95% (89-98%)	5.5% (2.4-11%)	3.1 (2.1-4.4)	13% (9.1-16%)	18% (11-24%)	360 (230-500)	28000 (20000-38000)
Poland	1995	Male	30 to 34	95% (90-98%)	4.8% (1.9-10%)	3.3 (2.4-4.6)	14% (11-18%)	19% (13-25%)	570 (410-770)	40000 (31000-52000)
Poland	1995	Male	35 to 39	96% (91-98%)	4.3% (1.7-8.6%)	3.2 (2.2-4.4)	13% (10-17%)	17% (12-22%)	1000 (720-1300)	61000 (46000-77000)
Poland	1995	Male	40 to 44	96% (92-99%)	3.7% (1.5-7.9%)	3.9 (2.8-5.1)	14% (10-18%)	18% (13-23%)	1600 (1100-2100)	84000 (62000-110000)
Poland	1995	Male	45 to 49	97% (93-99%)	3.1% (1.3-6.6%)	5.5 (3.8-7.3)	17% (11-22%)	21% (14-27%)	2300 (1500-3000)	110000 (71000-130000)
Poland	1995	Male	50 to 54	97% (94-98%)	3.3% (1.7-6.2%)	6.2 (4.4-8)	17% (12-21%)	21% (15-25%)	2200 (1600-2800)	86000 (63000-110000)
Poland	1995	Male	55 to 59	96% (93-97%)	4.4% (2.5-7.2%)	6.4 (4.8-8.3)	15% (11-20%)	18% (13-23%)	3100 (2300-3900)	100000 (77000-130000)
Poland	1995	Male	60 to 64	95% (91-97%)	5% (3-8.5%)	6.1 (4.3-8.3)	13% (8.8-17%)	15% (9.9-20%)	3500 (2300-4700)	100000 (67000-130000)
Poland	1995	Male	65 to 69	95% (91-97%)	5.2% (3.1-8.6%)	5.5 (4.1-7.1)	11% (6.4-15%)	12% (7-17%)	3400 (2000-4900)	81000 (49000-120000)
Poland	1995	Male	70 to 74	94% (91-96%)	5.8% (3.6-9.4%)	4.7 (3.3-6.3)	7.7% (3.1-13%)	8.6% (3.1-15%)	2400 (850-4000)	47000 (18000-77000)
Poland	1995	Male	75 to 79	93% (89-96%)	6.6% (3.9-11%)	3.9 (2.6-5.5)	5.1% (0.4-11%)	5.4% (-0.04-12%)	1000 (-7.7-2300)	16000 (1300-34000)
Poland	1995	Male	80 to 84	93% (87-96%)	7.3% (4.1-13%)	3.4 (2.3-4.7)	2.8% (-1.3-7.9%)	2.8% (-2-8.9%)	600 (-430-1900)	7400 (-3400-21000)
Poland	1995	Male	85 to 89	92% (87-96%)	7.8% (4.4-13%)	3 (2.1-4.2)	0.59% (-3.2-5.1%)	0.19% (-4.2-5.5%)	26 (-590-770)	720 (-3800-6200)
Poland	1995	Male	90 to 94	92% (87-96%)	8% (4.5-13%)	2.6 (1.7-3.8)	-1.1% (-5-3.7%)	-1.8% (-6.5-3.8%)	-85 (-300-180)	-310 (-1400-1100)
Poland	1995	Male	95 plus	92% (84-96%)	8.2% (3.7-16%)	2.2 (1.2-3.6)	-2% (-5.9-3%)	-3% (-7.6-3%)	-27 (-69-27)	-74 (-220-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	2000	Female	All Ages	77% (73-81%)	23% (19-27%)	1.8 (1.6-2)	2.9% (1.6-4.2%)	2.1% (-0.19-4.6%)	3600 (-330-7900)	150000 (83000-220000)
Poland	2000	Female	15 to 19	77% (62-87%)	23% (13-38%)	1.5 (0.95-2.1)	2.6% (1.5-3.8%)	7.3% (3.8-11%)	35 (18-55)	3800 (2300-5600)
Poland	2000	Female	20 to 24	83% (71-91%)	17% (9.4-29%)	1.9 (1.2-2.6)	3.5% (2.3-4.8%)	7.7% (4.2-12%)	37 (20-57)	5200 (3300-7400)
Poland	2000	Female	25 to 29	84% (72-92%)	16% (8.4-28%)	1.8 (1.2-2.6)	3.6% (2.6-4.7%)	7.8% (4.6-11%)	40 (24-57)	5200 (3600-7000)
Poland	2000	Female	30 to 34	84% (72-92%)	16% (7.9-28%)	1.9 (1.3-2.7)	3.7% (2.7-5%)	8.9% (6-12%)	56 (38-77)	5700 (4000-7500)
Poland	2000	Female	35 to 39	84% (73-92%)	16% (8.2-27%)	2 (1.3-2.9)	4.5% (3.2-5.9%)	9.8% (6.7-13%)	120 (81-160)	8700 (6300-11000)
Poland	2000	Female	40 to 44	84% (74-92%)	16% (8.3-26%)	2.3 (1.6-3.1)	5.2% (3.7-6.9%)	10% (7.1-13%)	270 (190-360)	16000 (11000-21000)
Poland	2000	Female	45 to 49	85% (78-91%)	15% (9.5-22%)	2.8 (2-3.7)	5.8% (4.1-7.6%)	10% (7.4-13%)	460 (330-610)	22000 (16000-29000)
Poland	2000	Female	50 to 54	85% (79-90%)	15% (9.7-21%)	2.9 (2-3.8)	5.9% (4.1-8%)	9.7% (6.9-13%)	560 (400-730)	23000 (16000-30000)
Poland	2000	Female	55 to 59	85% (79-89%)	15% (11-21%)	2.8 (2-3.9)	5.3% (3.4-7.5%)	8.3% (5.5-12%)	470 (310-650)	16000 (11000-23000)
Poland	2000	Female	60 to 64	81% (74-86%)	19% (14-26%)	2.6 (1.8-3.7)	4% (1.8-6.8%)	6% (2.9-9.8%)	560 (260-910)	17000 (7500-28000)
Poland	2000	Female	65 to 69	75% (66-82%)	25% (18-34%)	2.3 (1.6-3.1)	2.9% (0.67-5.3%)	4.1% (1-7.4%)	610 (150-1100)	15000 (3400-27000)
Poland	2000	Female	70 to 74	67% (59-76%)	33% (24-41%)	1.9 (1.3-2.7)	1.8% (-0.7-4.5%)	2.3% (-0.95-5.8%)	540 (-220-1300)	11000 (-4400-26000)
Poland	2000	Female	75 to 79	59% (46-70%)	41% (30-54%)	1.6 (1-2.3)	0.84% (-1.7-3.5%)	0.99% (-2.3-4.3%)	300 (-700-1300)	4900 (-9900-20000)
Poland	2000	Female	80 to 84	54% (41-65%)	46% (35-59%)	1.4 (0.89-2)	0.19% (-2.1-2.4%)	0.11% (-2.7-2.9%)	26 (-660-700)	620 (-6600-7900)
Poland	2000	Female	85 to 89	53% (40-65%)	47% (35-60%)	1.3 (0.81-2)	-0.37% (-2.7-1.9%)	-0.61% (-3.4-2.2%)	-170 (-960-620)	-950 (-6800-4800)
Poland	2000	Female	90 to 94	55% (43-68%)	45% (32-57%)	1.3 (0.78-2.1)	-0.9% (-3.5-1.8%)	-1.3% (-4.4-2%)	-200 (-680-320)	-860 (-3400-1700)
Poland	2000	Female	95 plus	59% (43-73%)	41% (27-57%)	1.4 (0.7-2.3)	-1.4% (-4.6-2.4%)	-1.9% (-5.8-2.6%)	-80 (-250-110)	-240 (-800-410)
Poland	2000	Male	All Ages	93% (90-95%)	7.1% (5-9.8%)	3.4 (3-3.7)	12% (11-15%)	13% (9.7-16%)	25000 (19000-32000)	850000 (710000-1000000)
Poland	2000	Male	15 to 19	89% (80-95%)	11% (5.3-20%)	1.6 (1.1-2.4)	5.5% (3.2-8.1%)	9.7% (5.2-15%)	130 (67-200)	11000 (6700-16000)
Poland	2000	Male	20 to 24	93% (87-97%)	6.7% (3.1-13%)	2.7 (1.8-3.8)	10% (6.7-15%)	16% (8.9-22%)	310 (170-440)	25000 (16000-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	2000	Male	25 to 29	95% (90-98%)	5.5% (2.4-10%)	3.4 (2.3-4.6)	13% (9.4-17%)	19% (13-27%)	370 (240-510)	30000 (21000-39000)
Poland	2000	Male	30 to 34	95% (90-98%)	4.7% (1.9-9.8%)	3.6 (2.5-5)	15% (11-18%)	21% (16-28%)	490 (360-640)	35000 (27000-45000)
Poland	2000	Male	35 to 39	96% (92-98%)	4.3% (1.7-8.5%)	3.4 (2.2-4.7)	15% (11-19%)	20% (15-26%)	780 (580-1000)	48000 (37000-60000)
Poland	2000	Male	40 to 44	96% (92-99%)	3.7% (1.4-7.6%)	4.1 (2.9-5.4)	16% (12-20%)	21% (16-26%)	1600 (1200-2000)	84000 (65000-100000)
Poland	2000	Male	45 to 49	97% (94-99%)	3% (1.3-6.4%)	5.8 (4-7.7)	19% (14-23%)	24% (18-29%)	2800 (2000-3400)	130000 (93000-150000)
Poland	2000	Male	50 to 54	97% (94-98%)	3.3% (1.8-6.1%)	6.6 (4.8-8.4)	19% (15-22%)	23% (18-27%)	3200 (2500-3800)	130000 (97000-150000)
Poland	2000	Male	55 to 59	96% (93-97%)	4.4% (2.5-6.9%)	6.7 (5-8.8)	17% (13-21%)	20% (16-25%)	2600 (2000-3200)	87000 (67000-110000)
Poland	2000	Male	60 to 64	95% (92-97%)	5% (3-8.1%)	6.5 (4.6-8.7)	14% (10-18%)	17% (12-21%)	3300 (2300-4200)	94000 (67000-120000)
Poland	2000	Male	65 to 69	95% (92-97%)	5.2% (3-8.2%)	5.8 (4.3-7.5)	12% (8-16%)	14% (9.1-19%)	3500 (2400-4800)	85000 (57000-110000)
Poland	2000	Male	70 to 74	94% (91-96%)	5.7% (3.6-9.1%)	5 (3.6-6.7)	9% (4.4-14%)	10% (4.8-16%)	3000 (1400-4600)	58000 (28000-88000)
Poland	2000	Male	75 to 79	93% (89-96%)	6.5% (3.7-11%)	4.2 (2.8-6)	6.4% (1.7-12%)	7.1% (1.5-13%)	1800 (410-3500)	28000 (7300-52000)
Poland	2000	Male	80 to 84	93% (87-96%)	7.2% (4.1-13%)	3.6 (2.5-5)	4.3% (0.28-9.3%)	4.6% (-0.24-11%)	700 (-36-1600)	8200 (520-18000)
Poland	2000	Male	85 to 89	92% (87-96%)	7.7% (4.4-13%)	3.2 (2.2-4.5)	2.1% (-1.6-6.8%)	2% (-2.4-7.5%)	260 (-320-970)	2400 (-1800-7700)
Poland	2000	Male	90 to 94	92% (87-96%)	7.9% (4.4-13%)	2.8 (1.8-4.1)	0.24% (-3.6-5%)	-0.29% (-5-5.5%)	-16 (-270-300)	79 (-1200-1700)
Poland	2000	Male	95 plus	92% (85-97%)	8.1% (3.5-15%)	2.4 (1.2-3.8)	-0.92% (-4.6-4.3%)	-1.7% (-6.3-4.5%)	-20 (-72-52)	-43 (-220-200)
Poland	2005	Female	All Ages	76% (72-80%)	24% (20-28%)	1.9 (1.7-2.1)	3.2% (2.3-4.3%)	2.6% (0.87-4.6%)	4500 (1500-8000)	170000 (120000-220000)
Poland	2005	Female	15 to 19	76% (62-86%)	24% (14-38%)	1.6 (0.99-2.2)	2.5% (1.5-3.6%)	7.7% (4-12%)	27 (14-43)	3100 (1900-4500)
Poland	2005	Female	20 to 24	83% (72-90%)	17% (9.8-28%)	2 (1.3-2.9)	3.4% (2.3-4.6%)	8.3% (4.6-13%)	35 (20-54)	5100 (3400-7200)
Poland	2005	Female	25 to 29	84% (73-91%)	16% (8.8-27%)	1.9 (1.3-2.7)	3.5% (2.5-4.5%)	8.6% (5.3-12%)	41 (26-58)	5600 (3800-7400)
Poland	2005	Female	30 to 34	84% (73-91%)	16% (8.6-27%)	2 (1.4-2.8)	3.8% (2.8-4.9%)	9.9% (6.9-13%)	63 (44-85)	6300 (4700-8300)
Poland	2005	Female	35 to 39	85% (76-92%)	15% (8.5-24%)	2.1 (1.4-3)	4.7% (3.4-6.2%)	11% (8.3-15%)	110 (80-140)	8100 (6100-10000)
Poland	2005	Female	40 to 44	85% (76-91%)	15% (9-24%)	2.5 (1.7-3.3)	5.6% (4.1-7.3%)	12% (8.9-15%)	220 (160-280)	13000 (9800-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	2005	Female	45 to 49	86% (80-90%)	14% (9.8-20%)	3 (2.2-4)	6.5% (4.8-8.4%)	12% (9.2-15%)	490 (370-610)	24000 (18000-30000)
Poland	2005	Female	50 to 54	85% (80-89%)	15% (11-20%)	3.1 (2.2-4)	6.4% (4.7-8.3%)	11% (8.3-14%)	680 (520-870)	28000 (21000-37000)
Poland	2005	Female	55 to 59	83% (79-86%)	17% (14-21%)	2.9 (2.1-3.8)	5.8% (4.1-7.8%)	9.2% (6.6-12%)	720 (520-950)	26000 (18000-35000)
Poland	2005	Female	60 to 64	77% (72-82%)	23% (18-28%)	2.5 (1.8-3.5)	4.5% (2.5-6.8%)	6.9% (4-10%)	500 (290-740)	15000 (8700-23000)
Poland	2005	Female	65 to 69	68% (62-75%)	32% (25-38%)	2.1 (1.5-2.8)	3.1% (1.5-5.1%)	4.6% (2.3-7.4%)	560 (280-900)	14000 (6800-23000)
Poland	2005	Female	70 to 74	59% (53-65%)	41% (35-47%)	1.7 (1.1-2.3)	2% (0.21-4.2%)	2.8% (0.38-5.6%)	550 (74-1100)	11000 (1200-22000)
Poland	2005	Female	75 to 79	47% (38-55%)	53% (45-62%)	1.2 (0.8-1.8)	1.1% (-0.57-3.1%)	1.4% (-0.82-4%)	410 (-240-1200)	6600 (-3300-18000)
Poland	2005	Female	80 to 84	42% (32-53%)	58% (47-68%)	1.1 (0.68-1.5)	0.58% (-0.96-2.3%)	0.62% (-1.4-2.8%)	220 (-470-1000)	2800 (-4500-12000)
Poland	2005	Female	85 to 89	44% (33-55%)	56% (45-67%)	1.1 (0.7-1.7)	0.13% (-1.6-2.1%)	0.015% (-2.1-2.4%)	3.5 (-460-520)	270 (-3300-4200)
Poland	2005	Female	90 to 94	46% (35-57%)	54% (43-65%)	1.1 (0.66-1.8)	-0.29% (-2.3-2%)	-0.53% (-3-2.3%)	-87 (-500-380)	-310 (-2400-2100)
Poland	2005	Female	95 plus	50% (34-66%)	50% (34-66%)	1.2 (0.59-2)	-0.63% (-3.2-2.3%)	-0.97% (-4-2.6%)	-56 (-240-150)	-150 (-770-560)
Poland	2005	Male	All Ages	95% (93-96%)	5.4% (3.9-7.4%)	4 (3.6-4.4)	14% (12-16%)	14% (11-18%)	28000 (22000-35000)	950000 (810000-1100000)
Poland	2005	Male	15 to 19	91% (84-95%)	9.5% (5-16%)	1.8 (1.2-2.6)	5.6% (3.4-8.1%)	11% (5.7-16%)	100 (55-160)	9400 (5800-14000)
Poland	2005	Male	20 to 24	95% (91-98%)	4.8% (2.4-9.1%)	3.1 (2.1-4.3)	11% (7.3-15%)	18% (10-25%)	320 (190-450)	27000 (17000-37000)
Poland	2005	Male	25 to 29	96% (93-98%)	4% (2-7.1%)	3.8 (2.7-5.1)	14% (9.8-17%)	22% (15-29%)	420 (280-560)	34000 (24000-44000)
Poland	2005	Male	30 to 34	97% (93-98%)	3.5% (1.6-6.6%)	4 (2.9-5.5)	15% (12-19%)	24% (18-31%)	550 (410-710)	39000 (30000-48000)
Poland	2005	Male	35 to 39	97% (94-99%)	3.2% (1.4-5.6%)	3.8 (2.5-5.2)	16% (12-20%)	24% (18-30%)	750 (560-960)	46000 (35000-57000)
Poland	2005	Male	40 to 44	97% (95-99%)	2.8% (1.2-5.2%)	4.6 (3.3-6.1)	18% (14-22%)	25% (19-31%)	1400 (1100-1700)	72000 (56000-88000)
Poland	2005	Male	45 to 49	98% (96-99%)	2.3% (1.1-4.4%)	6.4 (4.6-8.5)	21% (16-25%)	28% (21-33%)	3000 (2300-3700)	140000 (110000-160000)
Poland	2005	Male	50 to 54	97% (96-99%)	2.6% (1.5-4.4%)	7.3 (5.4-9.2)	21% (17-24%)	26% (21-30%)	4200 (3400-4900)	160000 (130000-190000)
Poland	2005	Male	55 to 59	96% (95-98%)	3.5% (2.3-5.2%)	7.5 (5.8-9.7)	19% (15-22%)	23% (19-27%)	4200 (3400-4900)	140000 (110000-170000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	2005	Male	60 to 64	96% (94-97%)	4% (2.6-5.8%)	7.3 (5.4-9.6)	16% (12-20%)	19% (15-23%)	3000 (2300-3700)	87000 (66000-110000)
Poland	2005	Male	65 to 69	95% (93-97%)	4.7% (3-7%)	6.4 (5-8.1)	13% (9.6-17%)	16% (11-20%)	3400 (2400-4400)	83000 (60000-100000)
Poland	2005	Male	70 to 74	94% (91-96%)	6% (4-8.7%)	5.2 (3.8-7)	9.9% (5.6-14%)	11% (6.3-17%)	3100 (1700-4500)	60000 (34000-87000)
Poland	2005	Male	75 to 79	93% (89-96%)	6.8% (4.2-11%)	4.4 (3-6.1)	7.1% (2.5-12%)	8% (2.5-14%)	2200 (700-3800)	34000 (12000-58000)
Poland	2005	Male	80 to 84	93% (87-96%)	7.3% (4.1-13%)	3.8 (2.7-5.3)	5.1% (0.93-9.9%)	5.6% (0.59-11%)	1200 (130-2400)	14000 (2500-27000)
Poland	2005	Male	85 to 89	92% (87-95%)	7.9% (4.6-13%)	3.4 (2.4-4.7)	3% (-0.75-7.8%)	3% (-1.5-8.7%)	310 (-150-910)	2800 (-710-7300)
Poland	2005	Male	90 to 94	92% (87-96%)	8% (4.4-13%)	3 (2-4.2)	1% (-2.7-5.8%)	0.61% (-4.1-6.4%)	36 (-230-360)	370 (-970-2100)
Poland	2005	Male	95 plus	92% (84-96%)	8.3% (3.6-16%)	2.5 (1.3-4)	-0.27% (-3.9-5%)	-1% (-5.6-5.3%)	-17 (-91-87)	-18 (-260-350)
Poland	2010	Female	All Ages	77% (73-81%)	23% (19-27%)	1.9 (1.7-2.2)	3.2% (2.2-4.4%)	2.4% (0.58-4.6%)	4400 (1000-8200)	170000 (120000-230000)
Poland	2010	Female	15 to 19	76% (63-87%)	24% (13-37%)	1.6 (1-2.3)	2.4% (1.5-3.6%)	7.7% (4-12%)	21 (11-33)	2500 (1500-3700)
Poland	2010	Female	20 to 24	83% (73-91%)	17% (9.2-27%)	2 (1.3-2.9)	3.4% (2.3-4.5%)	8.4% (4.7-13%)	28 (15-43)	4300 (2800-6100)
Poland	2010	Female	25 to 29	84% (73-91%)	16% (8.7-27%)	1.9 (1.3-2.7)	3.5% (2.6-4.6%)	8.9% (5.7-12%)	40 (26-56)	5700 (4000-7600)
Poland	2010	Female	30 to 34	84% (74-92%)	16% (8.5-26%)	2 (1.3-2.8)	3.8% (2.8-5%)	11% (7.8-14%)	66 (48-86)	6900 (5200-9000)
Poland	2010	Female	35 to 39	85% (76-92%)	15% (8.3-24%)	2.2 (1.4-3.1)	4.7% (3.5-6.1%)	12% (9-16%)	120 (86-150)	8900 (6600-11000)
Poland	2010	Female	40 to 44	85% (77-91%)	15% (8.5-23%)	2.5 (1.7-3.4)	5.6% (4.2-7.3%)	13% (9.8-16%)	190 (140-240)	11000 (8700-15000)
Poland	2010	Female	45 to 49	86% (80-91%)	14% (9.2-20%)	3 (2.2-4.1)	6.5% (4.9-8.4%)	13% (10-16%)	360 (280-450)	18000 (14000-23000)
Poland	2010	Female	50 to 54	85% (80-89%)	15% (11-20%)	3.1 (2.2-4.1)	6.6% (4.9-8.5%)	12% (9-14%)	660 (500-830)	28000 (21000-36000)
Poland	2010	Female	55 to 59	84% (79-87%)	16% (13-21%)	2.9 (2.2-3.8)	5.9% (4.2-7.7%)	9.6% (7.2-12%)	830 (630-1100)	30000 (22000-39000)
Poland	2010	Female	60 to 64	80% (75-84%)	20% (16-25%)	2.7 (1.9-3.7)	4.6% (2.7-6.8%)	7.1% (4.4-10%)	750 (460-1100)	23000 (14000-34000)
Poland	2010	Female	65 to 69	72% (66-78%)	28% (22-34%)	2.3 (1.7-3)	3.3% (1.7-5.2%)	4.9% (2.6-7.6%)	480 (250-760)	12000 (6200-20000)
Poland	2010	Female	70 to 74	62% (55-68%)	38% (32-45%)	1.8 (1.2-2.5)	2% (0.29-4.2%)	2.9% (0.46-5.8%)	470 (76-930)	9700 (1400-19000)
Poland	2010	Female	75 to 79	50% (40-60%)	50% (40-60%)	1.3 (0.85-1.9)	1.1% (-0.56-3.2%)	1.4% (-0.87-4.1%)	370 (-220-1100)	6100 (-3100-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	2010	Female	80 to 84	46% (35-57%)	54% (43-65%)	1.2 (0.76-1.7)	0.52% (-1.1-2.4%)	0.54% (-1.5-2.9%)	200 (-570-1100)	2700 (-5600-12000)
Poland	2010	Female	85 to 89	49% (37-59%)	51% (41-63%)	1.3 (0.79-1.9)	0.028% (-1.9-2.1%)	-0.14% (-2.6-2.5%)	-49 (-890-880)	92 (-6400-7300)
Poland	2010	Female	90 to 94	51% (39-62%)	49% (38-61%)	1.3 (0.75-2)	-0.4% (-2.7-2.3%)	-0.69% (-3.5-2.6%)	-100 (-510-380)	-380 (-2500-2200)
Poland	2010	Female	95 plus	54% (38-69%)	46% (31-62%)	1.3 (0.68-2.2)	-0.72% (-3.6-2.8%)	-1.1% (-4.7-3.2%)	-80 (-340-230)	-210 (-1100-830)
Poland	2010	Male	All Ages	95% (93-96%)	5.2% (3.8-6.9%)	4.1 (3.7-4.6)	14% (12-16%)	14% (11-18%)	29000 (23000-36000)	960000 (820000-1100000)
Poland	2010	Male	15 to 19	91% (84-95%)	9.5% (4.8-16%)	1.9 (1.2-2.7)	5.6% (3.3-8.2%)	11% (5.5-17%)	89 (45-140)	8000 (4900-12000)
Poland	2010	Male	20 to 24	95% (91-98%)	4.8% (2.3-9%)	3.1 (2.1-4.4)	11% (7.2-15%)	18% (10-26%)	280 (160-400)	23000 (15000-32000)
Poland	2010	Male	25 to 29	96% (93-98%)	3.9% (1.8-6.9%)	3.8 (2.6-5.1)	13% (9.8-17%)	22% (15-29%)	410 (280-540)	33000 (24000-44000)
Poland	2010	Male	30 to 34	97% (94-98%)	3.4% (1.6-6.2%)	4 (2.9-5.4)	15% (12-19%)	25% (19-32%)	590 (440-740)	42000 (33000-52000)
Poland	2010	Male	35 to 39	97% (94-99%)	3.2% (1.4-6.1%)	3.8 (2.5-5.3)	17% (13-21%)	25% (19-32%)	810 (620-1000)	50000 (39000-61000)
Poland	2010	Male	40 to 44	97% (95-99%)	2.7% (1.2-5.4%)	4.6 (3.2-6.2)	18% (14-23%)	26% (21-32%)	1200 (930-1500)	64000 (50000-78000)
Poland	2010	Male	45 to 49	98% (96-99%)	2.2% (1.1-4.3%)	6.5 (4.5-8.6)	21% (17-26%)	29% (22-34%)	2200 (1700-2700)	100000 (79000-120000)
Poland	2010	Male	50 to 54	97% (96-98%)	2.5% (1.5-4.3%)	7.4 (5.4-9.3)	21% (17-25%)	27% (22-32%)	4000 (3300-4700)	160000 (130000-190000)
Poland	2010	Male	55 to 59	97% (95-98%)	3.5% (2.2-5%)	7.6 (5.8-9.6)	19% (16-23%)	24% (20-28%)	4900 (4000-5800)	170000 (140000-200000)
Poland	2010	Male	60 to 64	96% (94-97%)	3.9% (2.6-5.6%)	7.3 (5.5-9.6)	16% (13-20%)	20% (15-24%)	4400 (3400-5300)	130000 (99000-160000)
Poland	2010	Male	65 to 69	96% (93-97%)	4.5% (2.9-6.6%)	6.4 (5-8.2)	13% (9.9-17%)	16% (12-20%)	2900 (2100-3700)	71000 (52000-89000)
Poland	2010	Male	70 to 74	94% (92-96%)	5.8% (3.9-8.4%)	5.3 (3.8-7.1)	10% (5.6-14%)	12% (6.4-17%)	2800 (1500-4000)	55000 (32000-78000)
Poland	2010	Male	75 to 79	94% (90-96%)	6.5% (4.1-10%)	4.5 (3-6.2)	7.1% (2.8-12%)	8.1% (2.7-14%)	2200 (760-3800)	35000 (14000-59000)
Poland	2010	Male	80 to 84	93% (88-96%)	7% (4-12%)	3.9 (2.7-5.4)	5.1% (0.96-9.8%)	5.6% (0.58-11%)	1400 (150-2900)	17000 (3200-33000)
Poland	2010	Male	85 to 89	92% (87-95%)	7.9% (4.6-13%)	3.4 (2.4-4.7)	2.9% (-0.69-7.5%)	2.9% (-1.5-8.4%)	470 (-240-1400)	4300 (-1100-11000)
Poland	2010	Male	90 to 94	92% (87-95%)	8% (4.6-13%)	3 (2-4.4)	1% (-2.6-6%)	0.6% (-3.9-6.7%)	33 (-210-370)	340 (-890-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	2010	Male	95 plus	92% (85-96%)	8.1% (3.6-15%)	2.5 (1.4-4)	-0.21% (-3.9-5%)	-0.92% (-5.5-5.4%)	-18 (-110-100)	-17 (-310-400)
Poland	2016	Female	All Ages	78% (74-81%)	22% (19-26%)	2.1 (1.6-2.6)	3.2% (1.8-4.8%)	2.4% (-0.22-5.4%)	4500 (-400-10000)	170000 (92000-250000)
Poland	2016	Female	15 to 19	77% (62-88%)	23% (12-38%)	1.6 (0.97-2.5)	2.4% (1.5-3.5%)	7.8% (3.9-13%)	15 (8-25)	1900 (1100-2800)
Poland	2016	Female	20 to 24	83% (72-91%)	17% (9.4-28%)	2 (1.2-3.1)	3.3% (2.3-4.6%)	8.4% (4.6-13%)	21 (12-34)	3500 (2300-5000)
Poland	2016	Female	25 to 29	84% (72-92%)	16% (8.4-28%)	2 (1.3-3)	3.4% (2.5-4.5%)	9.2% (5.7-13%)	31 (19-45)	4700 (3200-6500)
Poland	2016	Female	30 to 34	84% (72-92%)	16% (7.9-28%)	2 (1.3-3)	3.7% (2.7-5%)	11% (7.7-14%)	62 (43-85)	6900 (5000-9200)
Poland	2016	Female	35 to 39	84% (74-92%)	16% (8.3-26%)	2.2 (1.3-3.2)	4.4% (3.3-5.8%)	12% (9.2-16%)	120 (83-160)	9400 (6900-12000)
Poland	2016	Female	40 to 44	85% (74-92%)	15% (8.4-26%)	2.5 (1.6-3.6)	5.3% (3.8-7%)	13% (10-17%)	190 (130-250)	12000 (8400-15000)
Poland	2016	Female	45 to 49	86% (78-91%)	14% (9.4-22%)	3.1 (2-4.4)	6.3% (4.6-8.6%)	14% (10-18%)	290 (220-380)	15000 (11000-20000)
Poland	2016	Female	50 to 54	86% (80-90%)	14% (9.7-20%)	3.2 (2.2-4.5)	6.6% (4.6-8.9%)	12% (9-16%)	470 (340-630)	21000 (15000-27000)
Poland	2016	Female	55 to 59	85% (80-89%)	15% (11-20%)	3.2 (2.2-4.3)	6% (4.1-8.4%)	10% (7.4-13%)	800 (560-1100)	29000 (20000-40000)
Poland	2016	Female	60 to 64	83% (78-87%)	17% (13-22%)	3 (2-4.2)	4.9% (2.7-7.5%)	7.7% (4.6-11%)	920 (530-1400)	28000 (15000-43000)
Poland	2016	Female	65 to 69	78% (71-83%)	22% (17-29%)	2.7 (1.8-3.6)	3.6% (1.4-5.9%)	5.5% (2.5-8.7%)	770 (350-1200)	20000 (7800-32000)
Poland	2016	Female	70 to 74	70% (63-77%)	30% (23-37%)	2.2 (1.5-3.2)	2.3% (0.087-5.1%)	3.4% (0.4-7.2%)	430 (51-910)	9000 (330-19000)
Poland	2016	Female	75 to 79	61% (49-71%)	39% (29-51%)	1.8 (1.1-2.7)	1.4% (-0.91-4.3%)	1.8% (-1.3-5.7%)	410 (-280-1300)	6900 (-4400-21000)
Poland	2016	Female	80 to 84	56% (43-67%)	44% (33-57%)	1.6 (0.95-2.3)	0.7% (-1.4-3.2%)	0.75% (-2.1-4%)	260 (-750-1400)	3700 (-7600-17000)
Poland	2016	Female	85 to 89	55% (43-67%)	45% (33-57%)	1.5 (0.91-2.3)	0.15% (-2.1-3%)	-0.039% (-3.1-3.4%)	-17 (-1300-1400)	570 (-8600-11000)
Poland	2016	Female	90 to 94	57% (45-69%)	43% (31-55%)	1.6 (0.89-2.4)	-0.34% (-3.2-3.3%)	-0.67% (-4.2-3.7%)	-190 (-1200-1000)	-620 (-5900-5700)
Poland	2016	Female	95 plus	61% (44-75%)	39% (25-56%)	1.6 (0.79-2.6)	-0.67% (-4.1-3.9%)	-1.1% (-5.2-4.5%)	-77 (-360-310)	-190 (-1100-1100)
Poland	2016	Male	All Ages	94% (92-95%)	6.2% (4.5-8.2%)	4.2 (3.2-5.1)	14% (11-16%)	14% (10-18%)	28000 (19000-36000)	890000 (700000-1100000)
Poland	2016	Male	15 to 19	89% (80-95%)	11% (5.1-20%)	1.8 (1.1-2.8)	5.2% (3.1-7.8%)	11% (5.5-17%)	56 (28-90)	5300 (3200-7900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Poland	2016	Male	20 to 24	93% (88-97%)	6.6% (3-12%)	2.9 (1.9-4.3)	11% (7-15%)	18% (10-26%)	190 (110-300)	17000 (11000-24000)
Poland	2016	Male	25 to 29	95% (90-98%)	5.4% (2.4-10%)	3.6 (2.4-5.3)	13% (9.3-17%)	22% (15-30%)	310 (210-430)	26000 (19000-35000)
Poland	2016	Male	30 to 34	95% (90-98%)	4.6% (1.9-9.6%)	3.9 (2.6-5.6)	15% (11-19%)	25% (19-33%)	530 (390-720)	40000 (31000-51000)
Poland	2016	Male	35 to 39	96% (92-98%)	4.2% (1.7-8.5%)	3.7 (2.3-5.2)	16% (12-20%)	26% (20-33%)	800 (590-1000)	51000 (39000-64000)
Poland	2016	Male	40 to 44	96% (92-99%)	3.7% (1.5-7.7%)	4.5 (2.9-6.3)	18% (14-23%)	28% (22-34%)	1200 (900-1500)	64000 (49000-81000)
Poland	2016	Male	45 to 49	97% (94-99%)	3% (1.2-6.3%)	6.3 (4-8.9)	21% (16-25%)	30% (22-36%)	1700 (1300-2200)	79000 (60000-98000)
Poland	2016	Male	50 to 54	97% (94-98%)	3.3% (1.7-5.9%)	7.2 (4.8-9.7)	21% (16-25%)	28% (22-33%)	2700 (2000-3300)	110000 (83000-130000)
Poland	2016	Male	55 to 59	96% (94-98%)	4.1% (2.4-6.5%)	7.5 (5.2-10)	19% (15-23%)	24% (20-29%)	4400 (3400-5400)	150000 (120000-180000)
Poland	2016	Male	60 to 64	96% (93-97%)	4.3% (2.7-6.7%)	7.4 (5-10)	16% (12-20%)	20% (15-25%)	4900 (3600-6200)	140000 (110000-180000)
Poland	2016	Male	65 to 69	95% (93-97%)	4.6% (2.8-6.9%)	6.6 (4.8-8.6)	16% (9.4-17%)	16% (11-21%)	4200 (2900-5500)	100000 (72000-130000)
Poland	2016	Male	70 to 74	95% (92-97%)	5.2% (3.4-8.1%)	5.6 (3.8-7.8)	10% (5.6-14%)	12% (6.5-17%)	2300 (1200-3300)	46000 (25000-65000)
Poland	2016	Male	75 to 79	94% (90-97%)	6% (3.5-9.9%)	4.7 (3-6.9)	7.3% (2.8-12%)	8.5% (2.8-14%)	2100 (680-3600)	33000 (13000-55000)
Poland	2016	Male	80 to 84	93% (88-96%)	6.9% (3.8-12%)	4 (2.6-5.7)	5.2% (0.91-10%)	5.8% (0.56-12%)	1600 (150-3200)	19000 (3200-36000)
Poland	2016	Male	85 to 89	92% (87-96%)	7.7% (4.3-13%)	3.5 (2.3-5)	3.1% (-0.75-8.2%)	3.2% (-1.5-9.5%)	670 (-310-2000)	6100 (-1500-16000)
Poland	2016	Male	90 to 94	92% (87-96%)	7.8% (4.2-13%)	3.1 (1.9-4.7)	1.2% (-2.5-6.8%)	0.88% (-3.8-7.6%)	89 (-390-790)	790 (-1600-4400)
Poland	2016	Male	95 plus	92% (85-97%)	8% (3.4-15%)	2.6 (1.4-4.4)	0.0045% (-3.9-5.7%)	-0.7% (-5.6-6.3%)	-13 (-110-120)	0.29 (-310-460)
Romania	1990	Female	All Ages	64% (55-71%)	36% (29-45%)	2.6 (2.1-3.2)	8.4% (3.3-14%)	12% (3.6-22%)	15000 (4200-26000)	330000 (130000-530000)
Romania	1990	Female	15 to 19	65% (46-80%)	35% (20-54%)	2.3 (1.4-3.4)	3.7% (1.4-6.6%)	8.3% (1.8-16%)	37 (8.1-70)	3800 (1400-6800)
Romania	1990	Female	20 to 24	73% (56-85%)	27% (15-44%)	3 (1.9-4.3)	5.3% (2.6-8.4%)	9.5% (2.8-17%)	54 (15-97)	6000 (2900-9600)
Romania	1990	Female	25 to 29	74% (57-86%)	26% (14-43%)	2.9 (1.9-4.2)	5.7% (3.1-8.5%)	10% (3.9-17%)	54 (20-90)	5300 (2900-7800)
Romania	1990	Female	30 to 34	74% (57-86%)	26% (14-43%)	3 (1.9-4.4)	6.2% (3.5-9%)	11% (5.1-17%)	110 (49-170)	8600 (4800-13000)
Romania	1990	Female	35 to 39	75% (59-86%)	25% (14-41%)	3.3 (2.1-4.8)	7.3% (4.1-11%)	13% (6.8-20%)	180 (93-280)	12000 (6700-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	1990	Female	40 to 44	74% (58-86%)	26% (14-42%)	3.6 (2.4-5.1)	8.9% (4.9-13%)	16% (8.7-23%)	270 (150-400)	15000 (8700-22000)
Romania	1990	Female	45 to 49	73% (52-86%)	27% (14-48%)	3.9 (2.6-5.7)	10% (5.6-15%)	17% (9.5-25%)	370 (200-540)	17000 (9800-26000)
Romania	1990	Female	50 to 54	70% (52-84%)	30% (16-48%)	4 (2.7-5.7)	11% (6.1-17%)	17% (9.4-26%)	710 (390-1000)	29000 (16000-43000)
Romania	1990	Female	55 to 59	68% (49-83%)	32% (17-51%)	4 (2.7-5.8)	11% (5.6-18%)	17% (8.4-26%)	960 (480-1500)	33000 (16000-51000)
Romania	1990	Female	60 to 64	65% (45-82%)	35% (18-55%)	4 (2.6-5.9)	11% (4.9-19%)	16% (6.9-26%)	1400 (590-2200)	40000 (17000-65000)
Romania	1990	Female	65 to 69	61% (42-79%)	39% (21-58%)	4 (2.6-5.8)	12% (4.4-21%)	16% (6.1-27%)	2000 (730-3300)	47000 (17000-79000)
Romania	1990	Female	70 to 74	56% (38-73%)	44% (27-62%)	3.8 (2.3-5.9)	12% (3.4-22%)	15% (4.2-28%)	1700 (470-3100)	32000 (9000-60000)
Romania	1990	Female	75 to 79	49% (33-68%)	51% (32-67%)	3.3 (1.9-5.3)	10% (1.9-21%)	12% (2.4-25%)	2800 (530-5700)	40000 (7800-82000)
Romania	1990	Female	80 to 84	45% (29-63%)	55% (37-71%)	2.9 (1.7-4.5)	9.3% (1.4-20%)	11% (1.6-23%)	2400 (340-5000)	25000 (3900-54000)
Romania	1990	Female	85 to 89	43% (28-59%)	57% (41-72%)	2.8 (1.6-4.5)	8.1% (0.85-18%)	9.4% (0.88-21%)	1100 (110-2500)	8400 (870-19000)
Romania	1990	Female	90 to 94	45% (30-61%)	55% (39-70%)	2.9 (1.5-4.7)	7.7% (0.2-18%)	8.9% (0.023-22%)	440 (1.2-1100)	2300 (58-5500)
Romania	1990	Female	95 plus	48% (29-68%)	52% (32-71%)	3 (1.4-5.4)	7.6% (-0.56-20%)	8.7% (-0.87-23%)	130 (-13-350)	450 (-32-1200)
Romania	1990	Male	All Ages	70% (60-79%)	30% (21-40%)	8.1 (6.9-9.5)	16% (6.8-24%)	21% (6.8-32%)	28000 (9000-42000)	800000 (340000-1200000)
Romania	1990	Male	15 to 19	64% (43-83%)	36% (17-57%)	4.2 (2.5-6.4)	9.1% (4.1-16%)	16% (6-29%)	140 (51-260)	12000 (5300-20000)
Romania	1990	Male	20 to 24	74% (55-88%)	26% (12-45%)	7.4 (4.8-10)	17% (8.7-27%)	27% (12-43%)	340 (160-550)	27000 (14000-42000)
Romania	1990	Male	25 to 29	77% (60-91%)	23% (9.5-40%)	8.8 (6-12)	20% (11-30%)	31% (15-47%)	360 (180-540)	26000 (14000-38000)
Romania	1990	Male	30 to 34	79% (61-92%)	21% (7.8-39%)	10 (7.1-14)	22% (12-31%)	32% (17-47%)	690 (360-1000)	44000 (25000-62000)
Romania	1990	Male	35 to 39	81% (62-92%)	19% (7.6-38%)	12 (8.3-17)	23% (13-32%)	33% (18-46%)	1000 (580-1500)	59000 (34000-82000)
Romania	1990	Male	40 to 44	82% (64-92%)	18% (7.6-36%)	14 (9.7-18)	24% (13-33%)	32% (18-45%)	1300 (710-1800)	64000 (36000-88000)
Romania	1990	Male	45 to 49	81% (62-92%)	19% (7.8-38%)	14 (10-20)	23% (13-32%)	30% (16-42%)	1500 (790-2100)	65000 (35000-91000)
Romania	1990	Male	50 to 54	80% (62-92%)	20% (8.4-38%)	15 (10-20)	22% (11-32%)	27% (13-39%)	2400 (1200-3400)	91000 (45000-130000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	1990	Male	55 to 59	79% (60-92%)	21% (8.2-40%)	14 (10-19)	21% (8.9-30%)	25% (11-36%)	2900 (1300-4300)	98000 (43000-140000)
Romania	1990	Male	60 to 64	78% (57-92%)	22% (8.3-43%)	14 (9.2-19)	19% (6.6-29%)	23% (8.3-35%)	3400 (1200-5100)	97000 (34000-150000)
Romania	1990	Male	65 to 69	76% (55-91%)	24% (8.9-45%)	12 (8.4-17)	19% (5.4-29%)	22% (6.7-35%)	3500 (1100-5500)	83000 (25000-130000)
Romania	1990	Male	70 to 74	74% (54-88%)	26% (12-46%)	11 (7-15)	18% (4.6-28%)	20% (5.4-33%)	2300 (600-3600)	43000 (11000-69000)
Romania	1990	Male	75 to 79	72% (52-87%)	28% (13-48%)	9.1 (5.6-14)	16% (3.2-27%)	18% (3.5-31%)	3300 (640-5700)	49000 (9400-84000)
Romania	1990	Male	80 to 84	71% (48-86%)	29% (14-52%)	8.2 (5-12)	15% (2.3-26%)	17% (2.4-30%)	2800 (390-5000)	30000 (4500-53000)
Romania	1990	Male	85 to 89	69% (49-85%)	31% (15-51%)	7.5 (4.6-11)	13% (1.7-24%)	15% (1.7-27%)	1200 (130-2200)	9200 (1200-17000)
Romania	1990	Male	90 to 94	69% (49-87%)	31% (13-51%)	6.8 (4-11)	12% (1.9-22%)	13% (1.9-26%)	370 (52-700)	2000 (320-3700)
Romania	1990	Male	95 plus	69% (43-89%)	31% (11-57%)	6.1 (2.9-11)	10% (1.3-22%)	12% (0.95-24%)	76 (6.3-160)	280 (34-560)
Romania	1995	Female	All Ages	65% (57-72%)	35% (28-43%)	2.5 (2-3)	8.7% (3.5-14%)	12% (3.6-22%)	15000 (4500-27000)	340000 (130000-540000)
Romania	1995	Female	15 to 19	65% (47-80%)	35% (20-53%)	2.1 (1.3-3.1)	3.4% (1.2-6.1%)	7.9% (1.4-15%)	33 (5.9-65)	3500 (1300-6200)
Romania	1995	Female	20 to 24	73% (57-85%)	27% (15-43%)	2.7 (1.7-3.9)	5.1% (2.4-8.2%)	9.6% (2.1-18%)	48 (11-87)	5400 (2500-8800)
Romania	1995	Female	25 to 29	74% (58-86%)	26% (14-42%)	2.6 (1.7-3.9)	5.3% (2.8-8.2%)	10% (3.2-18%)	61 (19-110)	6300 (3300-9700)
Romania	1995	Female	30 to 34	75% (57-87%)	25% (13-43%)	2.8 (1.8-4)	6% (3.2-9%)	11% (4.7-18%)	73 (30-120)	5900 (3100-8900)
Romania	1995	Female	35 to 39	75% (60-87%)	25% (13-40%)	3 (1.9-4.4)	7.5% (4.1-11%)	13% (6.6-20%)	190 (93-290)	12000 (6400-18000)
Romania	1995	Female	40 to 44	74% (59-86%)	26% (14-41%)	3.3 (2.2-4.7)	8.9% (4.9-13%)	15% (8.1-22%)	340 (180-500)	18000 (10000-27000)
Romania	1995	Female	45 to 49	73% (53-85%)	27% (15-47%)	3.6 (2.3-5.1)	10% (5.7-15%)	17% (9.3-25%)	460 (260-690)	21000 (12000-32000)
Romania	1995	Female	50 to 54	71% (53-84%)	29% (16-47%)	3.7 (2.5-5.1)	11% (6.3-17%)	17% (9.4-26%)	590 (320-880)	24000 (13000-35000)
Romania	1995	Female	55 to 59	68% (49-83%)	32% (17-51%)	3.7 (2.4-5.3)	11% (5.5-18%)	17% (8.2-25%)	1100 (530-1600)	36000 (18000-56000)
Romania	1995	Female	60 to 64	65% (46-82%)	35% (18-54%)	3.7 (2.4-5.3)	11% (4.6-18%)	16% (6.6-25%)	1400 (610-2300)	40000 (17000-66000)
Romania	1995	Female	65 to 69	62% (43-80%)	38% (20-57%)	3.7 (2.4-5.2)	12% (4.1-20%)	15% (5.9-26%)	2100 (800-3500)	49000 (19000-83000)
Romania	1995	Female	70 to 74	57% (38-73%)	43% (27-62%)	3.5 (2.1-5.4)	11% (3.1-21%)	14% (4-26%)	2700 (730-5000)	51000 (14000-95000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	1995	Female	75 to 79	50% (34-69%)	50% (31-66%)	3.1 (1.8-4.9)	9.9% (1.8-20%)	12% (2.1-24%)	1800 (310-3900)	27000 (4800-56000)
Romania	1995	Female	80 to 84	46% (30-64%)	54% (36-70%)	2.7 (1.6-4.3)	8.8% (1.3-19%)	10% (1.2-22%)	2200 (270-4700)	23000 (3200-50000)
Romania	1995	Female	85 to 89	44% (29-61%)	56% (39-71%)	2.6 (1.4-4.1)	7.5% (0.59-17%)	8.6% (0.51-20%)	1400 (84-3200)	11000 (900-25000)
Romania	1995	Female	90 to 94	45% (30-62%)	55% (38-70%)	2.6 (1.5-4.3)	6.7% (-0.066-17%)	7.9% (-0.33-20%)	460 (-20-1200)	2400 (-24-6100)
Romania	1995	Female	95 plus	49% (30-69%)	51% (31-70%)	2.7 (1.3-5)	6.6% (-0.91-18%)	7.7% (-1.4-22%)	130 (-23-360)	450 (-63-1200)
Romania	1995	Male	All Ages	71% (60-80%)	29% (20-40%)	7.8 (6.6-9.1)	19% (8.1-28%)	22% (7.9-34%)	34000 (12000-52000)	980000 (430000-1500000)
Romania	1995	Male	15 to 19	65% (44-84%)	35% (16-56%)	3.9 (2.4-5.9)	8.8% (3.5-15%)	15% (5.4-27%)	140 (50-260)	12000 (5000-20000)
Romania	1995	Male	20 to 24	74% (53-89%)	26% (11-47%)	6.8 (4.4-9.6)	17% (8.8-26%)	26% (12-42%)	350 (160-560)	27000 (14000-42000)
Romania	1995	Male	25 to 29	77% (59-90%)	23% (9.8-41%)	8.1 (5.5-11)	20% (11-29%)	30% (16-45%)	490 (250-750)	35000 (20000-51000)
Romania	1995	Male	30 to 34	79% (60-92%)	21% (8.1-40%)	9.6 (6.5-13)	23% (13-33%)	33% (18-48%)	590 (320-850)	37000 (21000-52000)
Romania	1995	Male	35 to 39	81% (62-92%)	19% (7.7-38%)	11 (7.6-16)	26% (15-36%)	34% (19-47%)	1300 (750-1800)	71000 (42000-98000)
Romania	1995	Male	40 to 44	82% (64-93%)	18% (7-36%)	12 (8.8-17)	26% (15-36%)	34% (19-46%)	2000 (1100-2800)	98000 (55000-140000)
Romania	1995	Male	45 to 49	81% (60-93%)	19% (7.4-40%)	13 (9.1-18)	25% (14-35%)	31% (17-43%)	2300 (1200-3200)	97000 (53000-140000)
Romania	1995	Male	50 to 54	81% (62-92%)	19% (8.3-38%)	13 (9.4-18)	24% (12-35%)	29% (14-41%)	2400 (1200-3500)	91000 (45000-130000)
Romania	1995	Male	55 to 59	79% (59-92%)	21% (7.9-41%)	13 (9.3-18)	23% (9.7-33%)	27% (12-39%)	3700 (1600-5400)	120000 (52000-180000)
Romania	1995	Male	60 to 64	78% (56-92%)	22% (8.1-44%)	13 (8.5-18)	21% (7.5-31%)	24% (9.1-37%)	4200 (1500-6300)	120000 (42000-180000)
Romania	1995	Male	65 to 69	77% (55-91%)	23% (8.7-45%)	11 (7.8-15)	20% (6.1-31%)	23% (7.2-36%)	4800 (1500-7500)	110000 (34000-170000)
Romania	1995	Male	70 to 74	75% (55-89%)	25% (11-45%)	9.8 (6.6-14)	19% (5.1-30%)	22% (5.8-34%)	4500 (1200-7200)	83000 (23000-140000)
Romania	1995	Male	75 to 79	73% (52-88%)	27% (12-48%)	8.5 (5.3-12)	16% (3.7-28%)	19% (4.1-32%)	2400 (530-4000)	34000 (7900-59000)
Romania	1995	Male	80 to 84	71% (49-87%)	29% (13-51%)	7.6 (4.7-11)	16% (2.7-28%)	17% (2.9-31%)	2600 (450-4700)	28000 (5000-51000)
Romania	1995	Male	85 to 89	70% (50-85%)	30% (15-50%)	6.8 (4.3-10)	13% (2.1-24%)	15% (2.2-28%)	1600 (240-3000)	13000 (2100-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	1995	Male	90 to 94	69% (50-87%)	31% (13-50%)	6.2 (3.7-9.7)	12% (2.3-22%)	13% (2.4-26%)	480 (83-920)	2600 (520-4900)
Romania	1995	Male	95 plus	69% (45-89%)	31% (11-55%)	5.5 (2.7-9.7)	9.9% (1.1-21%)	11% (0.78-24%)	95 (6.5-210)	350 (37-720)
Romania	2000	Female	All Ages	64% (56-72%)	36% (28-44%)	2.3 (1.9-2.8)	7.8% (3.1-13%)	11% (3.1-20%)	13000 (3700-24000)	290000 (110000-480000)
Romania	2000	Female	15 to 19	64% (45-80%)	36% (20-55%)	1.9 (1.1-2.8)	3% (1.5-4%)	7.3% (1.1-14%)	24 (3.3-46)	2600 (900-4600)
Romania	2000	Female	20 to 24	72% (56-84%)	28% (16-44%)	2.4 (1.5-3.5)	4.5% (2.2-7.2%)	8.7% (1.6-17%)	35 (6.1-66)	4400 (2100-7000)
Romania	2000	Female	25 to 29	73% (57-86%)	27% (14-43%)	2.4 (1.6-3.4)	4.7% (2.4-7.3%)	9.2% (2.5-17%)	43 (12-78)	4800 (2500-7400)
Romania	2000	Female	30 to 34	74% (57-86%)	26% (14-43%)	2.5 (1.6-3.6)	5.1% (2.7-7.8%)	10% (3.9-17%)	72 (27-120)	6300 (3300-9500)
Romania	2000	Female	35 to 39	74% (59-87%)	26% (13-41%)	2.7 (1.7-4)	6.3% (3.3-9.6%)	12% (5.7-19%)	100 (48-160)	6800 (3400-10000)
Romania	2000	Female	40 to 44	74% (58-86%)	26% (14-42%)	2.9 (2-4.2)	7.7% (4.1-12%)	14% (7.3-21%)	260 (130-390)	14000 (7500-21000)
Romania	2000	Female	45 to 49	72% (52-85%)	28% (15-48%)	3.2 (2.1-4.6)	9% (5.1-14%)	15% (8.2-23%)	440 (240-680)	21000 (12000-32000)
Romania	2000	Female	50 to 54	70% (51-84%)	30% (16-49%)	3.3 (2.2-4.6)	10% (5.3-15%)	16% (8.4-24%)	570 (300-860)	23000 (12000-35000)
Romania	2000	Female	55 to 59	68% (49-83%)	32% (17-51%)	3.3 (2.1-4.8)	10% (4.9-16%)	15% (7.6-24%)	680 (330-1100)	23000 (11000-37000)
Romania	2000	Female	60 to 64	65% (45-82%)	35% (18-55%)	3.3 (2.1-4.8)	9.9% (4-16%)	14% (5.9-23%)	1200 (500-2000)	34000 (14000-57000)
Romania	2000	Female	65 to 69	61% (42-79%)	39% (21-58%)	3.3 (2.1-4.7)	10% (3.6-18%)	14% (5.5-24%)	1700 (650-2900)	41000 (15000-69000)
Romania	2000	Female	70 to 74	56% (37-72%)	44% (28-63%)	3.1 (1.9-4.8)	10% (2.6-19%)	13% (3.6-24%)	2300 (640-4400)	45000 (12000-84000)
Romania	2000	Female	75 to 79	49% (33-68%)	51% (32-67%)	2.7 (1.6-4.3)	8.7% (1.5-18%)	11% (1.8-22%)	2400 (430-5100)	35000 (6500-74000)
Romania	2000	Female	80 to 84	44% (28-62%)	56% (38-72%)	2.4 (1.4-3.7)	7.4% (0.99-16%)	8.7% (1.1-19%)	1600 (190-3400)	17000 (2300-36000)
Romania	2000	Female	85 to 89	42% (28-58%)	58% (42-72%)	2.3 (1.3-3.6)	6.5% (0.14-15%)	7.4% (-0.00032-17%)	1100 (-0.057-2600)	8400 (180-19000)
Romania	2000	Female	90 to 94	44% (29-61%)	56% (39-71%)	2.3 (1.3-3.8)	5.6% (-0.63-14%)	6.5% (-1-18%)	480 (-75-1300)	2600 (-280-6900)
Romania	2000	Female	95 plus	48% (29-68%)	52% (32-71%)	2.4 (1.1-4.4)	5.4% (-1.7-16%)	6.2% (-2.3-19%)	130 (-49-400)	450 (-150-1400)
Romania	2000	Male	All Ages	71% (61-80%)	29% (20-39%)	7.3 (6.2-8.4)	18% (8.5-27%)	22% (8.7-33%)	30000 (12000-46000)	850000 (400000-1200000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	2000	Male	15 to 19	63% (42-83%)	37% (17-58%)	3.5 (2.1-5.2)	7.5% (3.1-13%)	14% (4.4-25%)	88 (29-160)	7800 (3400-13000)
Romania	2000	Male	20 to 24	73% (53-88%)	27% (12-47%)	6.1 (3.9-8.6)	15% (8.2-23%)	25% (12-40%)	250 (120-410)	20000 (11000-31000)
Romania	2000	Male	25 to 29	76% (59-90%)	24% (9.9-41%)	7.3 (5-10)	18% (10-26%)	29% (16-43%)	350 (190-530)	26000 (15000-38000)
Romania	2000	Male	30 to 34	79% (60-92%)	21% (8.2-40%)	8.7 (5.9-12)	21% (13-29%)	32% (18-45%)	570 (330-810)	37000 (23000-51000)
Romania	2000	Male	35 to 39	80% (62-92%)	20% (7.9-38%)	10 (6.8-14)	24% (15-33%)	34% (20-46%)	730 (440-1000)	41000 (25000-56000)
Romania	2000	Male	40 to 44	81% (63-93%)	19% (7.5-37%)	11 (7.9-15)	26% (15-35%)	34% (20-45%)	1600 (930-2100)	77000 (46000-100000)
Romania	2000	Male	45 to 49	81% (60-92%)	19% (7.6-40%)	12 (8.3-16)	25% (14-34%)	32% (19-43%)	2200 (1300-3100)	97000 (56000-130000)
Romania	2000	Male	50 to 54	80% (60-92%)	20% (8.5-40%)	12 (8.4-16)	24% (13-33%)	29% (16-41%)	2400 (1300-3300)	91000 (47000-130000)
Romania	2000	Male	55 to 59	79% (59-92%)	21% (7.9-41%)	12 (8.3-16)	22% (10-32%)	27% (13-39%)	2500 (1200-3600)	82000 (38000-120000)
Romania	2000	Male	60 to 64	77% (55-92%)	23% (8.2-45%)	11 (7.6-16)	21% (8.4-31%)	24% (10-36%)	3600 (1500-5400)	100000 (42000-150000)
Romania	2000	Male	65 to 69	76% (54-91%)	24% (8.7-46%)	10 (6.9-14)	20% (6.9-30%)	23% (8.3-36%)	4200 (1500-6500)	98000 (35000-150000)
Romania	2000	Male	70 to 74	74% (55-89%)	26% (11-45%)	8.8 (5.8-13)	18% (6.1-29%)	21% (7.2-34%)	4500 (1500-7100)	85000 (27000-130000)
Romania	2000	Male	75 to 79	73% (52-88%)	27% (12-48%)	7.6 (4.7-11)	16% (4.4-27%)	19% (5-31%)	3500 (950-5900)	52000 (14000-88000)
Romania	2000	Male	80 to 84	71% (50-87%)	29% (13-50%)	6.8 (4.2-9.9)	15% (3.8-25%)	17% (4.3-29%)	2000 (490-3400)	22000 (5600-37000)
Romania	2000	Male	85 to 89	69% (49-86%)	31% (14-51%)	6.1 (3.8-9.1)	13% (2.9-24%)	15% (3.2-28%)	1300 (270-2300)	9700 (2100-17000)
Romania	2000	Male	90 to 94	69% (49-87%)	31% (13-51%)	5.6 (3.2-8.9)	11% (2.2-21%)	13% (2.2-25%)	520 (89-1000)	2800 (540-5500)
Romania	2000	Male	95 plus	69% (44-89%)	31% (11-56%)	5 (2.3-8.9)	9.2% (0.49-20%)	10% (-0.044-23%)	110 (-0.47-240)	390 (21-850)
Romania	2005	Female	All Ages	61% (54-68%)	39% (32-46%)	2.1 (1.7-2.5)	6.9% (2.6-11%)	9.4% (2.7-17%)	12000 (3400-21000)	250000 (98000-410000)
Romania	2005	Female	15 to 19	61% (43-77%)	39% (23-57%)	1.7 (1-2.5)	2.8% (0.96-5.1%)	6.5% (0.79-13%)	21 (2.6-42)	2400 (830-4200)
Romania	2005	Female	20 to 24	70% (54-83%)	30% (17-46%)	2.2 (1.4-3.2)	4.2% (2-6.6%)	8.2% (1.2-16%)	24 (3.6-47)	3400 (1700-5300)
Romania	2005	Female	25 to 29	72% (58-85%)	28% (15-42%)	2.2 (1.5-3.1)	4.3% (2.3-6.6%)	8.6% (1.8-16%)	34 (6.7-62)	4200 (2200-6400)
Romania	2005	Female	30 to 34	73% (58-85%)	27% (15-42%)	2.3 (1.5-3.3)	4.5% (2.3-7%)	9.4% (3.1-16%)	53 (17-91)	5000 (2400-7800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	2005	Female	35 to 39	73% (58-85%)	27% (15-42%)	2.4 (1.6-3.6)	5.3% (2.7-8.3%)	11% (4.6-17%)	100 (44-170)	7300 (3600-11000)
Romania	2005	Female	40 to 44	72% (57-84%)	28% (16-43%)	2.7 (1.8-3.8)	6.7% (3.5-10%)	13% (6.4-19%)	150 (75-230)	8300 (4400-13000)
Romania	2005	Female	45 to 49	71% (53-84%)	29% (16-47%)	2.9 (2-4.2)	8% (4.3-12%)	14% (7.5-21%)	360 (190-540)	17000 (8800-25000)
Romania	2005	Female	50 to 54	68% (50-81%)	32% (19-50%)	2.9 (2-4)	8.8% (4.5-13%)	14% (7.3-21%)	550 (280-830)	22000 (12000-34000)
Romania	2005	Female	55 to 59	64% (45-79%)	36% (21-55%)	2.8 (1.9-4.1)	8.9% (4.2-14%)	14% (6.7-21%)	660 (330-1000)	23000 (11000-36000)
Romania	2005	Female	60 to 64	61% (44-78%)	39% (22-56%)	2.9 (1.9-4.2)	8.8% (3.5-14%)	13% (5.4-20%)	750 (310-1200)	22000 (8900-36000)
Romania	2005	Female	65 to 69	54% (37-73%)	46% (27-63%)	2.6 (1.7-3.8)	8.8% (3-15%)	12% (4.6-20%)	1400 (530-2300)	34000 (12000-57000)
Romania	2005	Female	70 to 74	46% (30-63%)	54% (37-70%)	2.3 (1.4-3.6)	8.2% (2.2-15%)	11% (3-20%)	1800 (500-3300)	34000 (9300-64000)
Romania	2005	Female	75 to 79	45% (32-62%)	55% (38-68%)	2.3 (1.4-3.6)	7.7% (1.5-16%)	9.5% (1.8-20%)	2200 (430-4500)	33000 (6300-67000)
Romania	2005	Female	80 to 84	43% (27-61%)	57% (39-73%)	2.1 (1.3-3.3)	6.8% (0.69-14%)	8.1% (0.76-17%)	1900 (190-4000)	21000 (2200-44000)
Romania	2005	Female	85 to 89	40% (26-57%)	60% (43-74%)	2 (1.1-3.2)	5.6% (0.057-13%)	6.5% (-0.073-15%)	890 (-10-2100)	6900 (71-16000)
Romania	2005	Female	90 to 94	41% (27-58%)	59% (42-73%)	2 (1.1-3.4)	5.1% (-1-13%)	5.8% (-1.4-16%)	400 (-99-1100)	2100 (-450-5600)
Romania	2005	Female	95 plus	45% (26-66%)	55% (34-74%)	2.1 (0.98-3.8)	4.9% (-2-15%)	5.5% (-2.6-17%)	170 (-79-530)	600 (-250-1900)
Romania	2005	Male	All Ages	75% (65-83%)	25% (17-35%)	7.6 (6.4-8.8)	19% (9-27%)	22% (9.4-33%)	30000 (13000-44000)	810000 (390000-1200000)
Romania	2005	Male	15 to 19	69% (50-85%)	31% (15-50%)	3.6 (2.3-5.4)	7.6% (3-13%)	14% (4.4-25%)	89 (29-160)	7900 (3400-13000)
Romania	2005	Male	20 to 24	78% (61-90%)	22% (9.6-39%)	6.4 (4.3-8.8)	15% (8.1-23%)	26% (13-41%)	210 (100-330)	17000 (9200-25000)
Romania	2005	Male	25 to 29	80% (65-92%)	20% (8.5-35%)	7.5 (5.2-10)	17% (10-25%)	29% (15-44%)	310 (160-460)	23000 (13000-33000)
Romania	2005	Male	30 to 34	82% (67-93%)	18% (6.9-33%)	8.8 (6.2-12)	20% (12-27%)	32% (19-45%)	480 (280-660)	31000 (19000-42000)
Romania	2005	Male	35 to 39	83% (67-93%)	17% (6.9-33%)	10 (6.8-14)	22% (14-30%)	34% (21-46%)	810 (500-1100)	46000 (29000-62000)
Romania	2005	Male	40 to 44	84% (67-93%)	16% (6.7-33%)	11 (8.1-15)	25% (15-33%)	34% (20-46%)	1000 (620-1400)	50000 (30000-67000)
Romania	2005	Male	45 to 49	84% (66-93%)	16% (6.8-34%)	12 (8.6-16)	25% (14-34%)	32% (19-43%)	2000 (1200-2700)	87000 (50000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	2005	Male	50 to 54	82% (66-93%)	18% (7.4-34%)	12 (8.7-16)	24% (13-33%)	29% (16-40%)	2700 (1500-3700)	100000 (56000-140000)
Romania	2005	Male	55 to 59	82% (64-93%)	18% (7.3-36%)	12 (8.6-16)	22% (11-32%)	27% (14-39%)	2800 (1400-4000)	94000 (46000-130000)
Romania	2005	Male	60 to 64	79% (61-92%)	21% (7.7-39%)	11 (7.5-16)	21% (8.7-30%)	25% (11-36%)	2800 (1200-4000)	78000 (33000-110000)
Romania	2005	Male	65 to 69	77% (58-92%)	23% (8.3-42%)	9.9 (7-13)	20% (7.2-30%)	23% (8.7-35%)	4000 (1500-6000)	94000 (35000-140000)
Romania	2005	Male	70 to 74	77% (60-90%)	23% (10-40%)	8.8 (6-12)	18% (6.7-29%)	22% (7.9-34%)	4400 (1600-6700)	83000 (30000-130000)
Romania	2005	Male	75 to 79	73% (53-88%)	27% (12-47%)	7.3 (4.7-11)	16% (4.8-26%)	19% (5.8-31%)	4000 (1300-6600)	59000 (19000-96000)
Romania	2005	Male	80 to 84	70% (46-86%)	30% (14-54%)	6.3 (3.9-9.1)	15% (4.3-25%)	17% (4.9-29%)	2700 (800-4600)	30000 (8700-51000)
Romania	2005	Male	85 to 89	69% (48-85%)	31% (15-52%)	5.8 (3.6-8.6)	13% (3.7-23%)	15% (4.1-26%)	1100 (300-1900)	8600 (2400-15000)
Romania	2005	Male	90 to 94	68% (48-86%)	32% (14-52%)	5.3 (3.1-8.3)	11% (2.4-22%)	13% (2.5-25%)	430 (82-820)	2300 (480-4400)
Romania	2005	Male	95 plus	68% (44-88%)	32% (12-56%)	4.7 (2.2-8.3)	9.3% (0.094-21%)	10% (-0.35-23%)	140 (-4.9-320)	510 (5-1100)
Romania	2010	Female	All Ages	60% (53-67%)	40% (33-47%)	2.1 (1.7-2.6)	6.8% (2.7-11%)	9.4% (3-17%)	11000 (3600-21000)	230000 (92000-380000)
Romania	2010	Female	15 to 19	61% (42-76%)	39% (24-58%)	1.7 (0.99-2.6)	2.8% (0.99-5.1%)	6.9% (0.72-13%)	14 (1.4-27)	1600 (520-2700)
Romania	2010	Female	20 to 24	69% (52-82%)	31% (18-48%)	2.2 (1.3-3.3)	4.3% (2.1-6.8%)	8.2% (1.1-16%)	22 (2.9-42)	3100 (1500-4900)
Romania	2010	Female	25 to 29	72% (57-84%)	28% (16-43%)	2.2 (1.5-3.2)	4.3% (2.3-6.5%)	8.5% (1.9-15%)	24 (5.2-44)	3100 (1600-4800)
Romania	2010	Female	30 to 34	73% (57-85%)	27% (15-43%)	2.3 (1.5-3.3)	4.3% (2.2-6.6%)	9.2% (2.9-15%)	44 (14-75)	4500 (2300-6900)
Romania	2010	Female	35 to 39	72% (58-85%)	28% (15-42%)	2.5 (1.6-3.6)	5% (2.6-7.7%)	11% (4.9-17%)	82 (37-130)	6100 (3100-9200)
Romania	2010	Female	40 to 44	72% (57-84%)	28% (16-43%)	2.7 (1.9-3.9)	6% (3.1-9.3%)	12% (6.2-18%)	160 (84-250)	9600 (5000-14000)
Romania	2010	Female	45 to 49	70% (53-83%)	30% (17-47%)	3 (2-4.3)	7.4% (4-11%)	13% (7.4-20%)	220 (120-330)	11000 (5800-16000)
Romania	2010	Female	50 to 54	67% (48-80%)	33% (20-52%)	3 (2-4.1)	8.2% (4.3-13%)	14% (7.4-20%)	460 (250-680)	19000 (10000-28000)
Romania	2010	Female	55 to 59	63% (45-78%)	37% (22-55%)	2.9 (1.9-4.2)	8.4% (4-13%)	13% (6.7-20%)	680 (350-1000)	24000 (12000-37000)
Romania	2010	Female	60 to 64	60% (43-77%)	40% (23-57%)	2.9 (1.9-4.2)	8.4% (3.5-14%)	13% (5.8-20%)	800 (370-1300)	24000 (10000-39000)
Romania	2010	Female	65 to 69	53% (37-72%)	47% (28-63%)	2.7 (1.7-3.8)	8.6% (3.3-14%)	12% (5-20%)	960 (390-1600)	23000 (9100-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	2010	Female	70 to 74	46% (30-63%)	54% (37-70%)	2.3 (1.4-3.7)	8.1% (2.7-15%)	11% (3.8-20%)	1700 (580-3100)	33000 (11000-61000)
Romania	2010	Female	75 to 79	44% (31-61%)	56% (39-69%)	2.3 (1.4-3.7)	7.7% (1.8-15%)	9.8% (2.3-19%)	2100 (510-4200)	31000 (7600-62000)
Romania	2010	Female	80 to 84	42% (26-60%)	58% (40-74%)	2.2 (1.3-3.4)	7.1% (1.2-14%)	8.4% (1.4-17%)	2200 (350-4400)	23000 (4000-48000)
Romania	2010	Female	85 to 89	39% (25-55%)	61% (45-75%)	2 (1.1-3.3)	5.9% (0.44-13%)	7% (0.44-15%)	1400 (84-3000)	11000 (780-23000)
Romania	2010	Female	90 to 94	40% (27-58%)	60% (42-73%)	2 (1.1-3.4)	5.1% (-0.47-13%)	6.1% (-0.8-16%)	440 (-57-1100)	2400 (-240-6000)
Romania	2010	Female	95 plus	44% (27-64%)	56% (36-73%)	2.1 (0.98-3.9)	4.9% (-1.4-14%)	5.7% (-1.9-17%)	130 (-42-380)	460 (-130-1300)
Romania	2010	Male	All Ages	75% (66-83%)	25% (17-34%)	8 (6.7-9.6)	19% (8.8-27%)	22% (9.2-32%)	30000 (13000-44000)	780000 (380000-1100000)
Romania	2010	Male	15 to 19	68% (49-84%)	32% (16-51%)	3.7 (2.3-5.6)	8.5% (3.3-14%)	16% (5.3-27%)	69 (22-120)	5900 (2400-9900)
Romania	2010	Male	20 to 24	78% (60-91%)	22% (9.5-40%)	6.6 (4.3-9.2)	16% (8.3-24%)	26% (13-41%)	210 (100-340)	17000 (9000-25000)
Romania	2010	Male	25 to 29	79% (65-91%)	21% (8.8-35%)	7.7 (5.3-11)	17% (10-25%)	30% (16-44%)	240 (120-350)	18000 (9900-25000)
Romania	2010	Male	30 to 34	82% (68-93%)	18% (7.2-32%)	9 (6.4-12)	19% (11-27%)	32% (19-45%)	410 (240-580)	27000 (16000-37000)
Romania	2010	Male	35 to 39	83% (68-93%)	17% (6.9-32%)	11 (7-14)	21% (13-29%)	34% (20-45%)	670 (400-900)	38000 (23000-52000)
Romania	2010	Male	40 to 44	84% (67-93%)	16% (6.5-33%)	12 (8.2-16)	23% (14-31%)	34% (20-45%)	1200 (710-1600)	59000 (35000-79000)
Romania	2010	Male	45 to 49	83% (64-93%)	17% (7.1-36%)	12 (8.7-17)	24% (13-33%)	31% (18-42%)	1300 (780-1900)	59000 (34000-81000)
Romania	2010	Male	50 to 54	82% (64-92%)	18% (7.8-36%)	12 (8.7-17)	23% (13-32%)	29% (16-40%)	2500 (1400-3600)	97000 (53000-140000)
Romania	2010	Male	55 to 59	81% (63-92%)	19% (7.7-37%)	12 (8.9-17)	22% (11-31%)	27% (14-38%)	3300 (1700-4700)	110000 (54000-150000)
Romania	2010	Male	60 to 64	79% (60-92%)	21% (8.1-40%)	11 (7.9-16)	20% (8.8-30%)	24% (11-35%)	3200 (1400-4700)	90000 (39000-130000)
Romania	2010	Male	65 to 69	77% (57-91%)	23% (8.8-43%)	10 (7.2-14)	19% (7.4-29%)	23% (9-35%)	3100 (1200-4600)	72000 (28000-110000)
Romania	2010	Male	70 to 74	76% (59-90%)	24% (10-41%)	9 (6.2-13)	18% (6.8-28%)	21% (8.1-33%)	4200 (1600-6500)	80000 (29000-120000)
Romania	2010	Male	75 to 79	73% (52-88%)	27% (12-48%)	7.5 (4.8-11)	16% (5.1-26%)	19% (6-30%)	4000 (1300-6500)	58000 (19000-95000)
Romania	2010	Male	80 to 84	69% (46-86%)	31% (14-54%)	6.4 (4-9.2)	15% (4.5-25%)	17% (5.3-29%)	3300 (1000-5500)	36000 (11000-60000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	2010	Male	85 to 89	68% (47-85%)	32% (15-53%)	5.9 (3.7-8.7)	13% (3.6-22%)	15% (4.3-26%)	1700 (470-3000)	14000 (3700-23000)
Romania	2010	Male	90 to 94	68% (49-86%)	32% (14-51%)	5.4 (3.2-8.5)	11% (2.7-21%)	13% (2.9-24%)	450 (100-850)	2500 (600-4500)
Romania	2010	Male	95 plus	68% (43-88%)	32% (12-57%)	4.8 (2.3-8.5)	9.2% (0.64-20%)	10% (0.26-23%)	84 (2.1-180)	310 (21-640)
Romania	2016	Female	All Ages	62% (54-70%)	38% (30-46%)	2.4 (1.8-3.2)	7.2% (2.8-12%)	9.9% (2.9-19%)	12000 (3400-23000)	230000 (84000-390000)
Romania	2016	Female	15 to 19	62% (44-78%)	38% (22-56%)	1.9 (1.1-3)	2.8% (1.1-5.1%)	7.4% (0.9-15%)	11 (1.2-22)	1300 (510-2400)
Romania	2016	Female	20 to 24	71% (53-84%)	29% (16-47%)	2.4 (1.4-3.8)	4.4% (2.2-7%)	8.8% (1.4-17%)	14 (2.1-27)	2100 (1000-3300)
Romania	2016	Female	25 to 29	72% (55-85%)	28% (15-45%)	2.4 (1.4-3.8)	4.4% (2.4-6.7%)	9% (2.4-16%)	22 (5.6-39)	2900 (1500-4500)
Romania	2016	Female	30 to 34	73% (55-86%)	27% (14-45%)	2.5 (1.5-3.8)	4.4% (2.4-6.7%)	9.7% (3.6-17%)	32 (11-54)	3400 (1800-5300)
Romania	2016	Female	35 to 39	73% (57-86%)	27% (14-43%)	2.7 (1.6-4.1)	4.9% (2.5-7.6%)	11% (5-18%)	69 (29-110)	5400 (2800-8400)
Romania	2016	Female	40 to 44	73% (57-85%)	27% (15-43%)	3 (1.9-4.3)	5.8% (2.9-9.1%)	13% (6.2-19%)	140 (65-220)	8200 (4100-13000)
Romania	2016	Female	45 to 49	71% (51-84%)	29% (16-49%)	3.2 (2-4.8)	7% (3.6-11%)	14% (7.3-21%)	260 (140-410)	13000 (6800-20000)
Romania	2016	Female	50 to 54	69% (50-83%)	31% (17-50%)	3.3 (2.1-4.9)	8.3% (4.3-13%)	14% (7.9-21%)	340 (190-510)	14000 (7400-21000)
Romania	2016	Female	55 to 59	66% (47-82%)	34% (18-53%)	3.2 (2-5)	8.5% (3.9-13%)	14% (6.8-21%)	560 (280-880)	20000 (9500-32000)
Romania	2016	Female	60 to 64	63% (43-80%)	37% (20-57%)	3.2 (1.9-5.1)	8.5% (3.6-14%)	13% (5.9-22%)	890 (390-1500)	27000 (11000-46000)
Romania	2016	Female	65 to 69	59% (40-78%)	41% (22-60%)	3.2 (2-4.8)	9.1% (3.6-16%)	13% (5.7-23%)	1200 (490-2000)	29000 (12000-50000)
Romania	2016	Female	70 to 74	53% (35-71%)	47% (29-65%)	3 (1.7-4.9)	9.1% (2.7-17%)	12% (4.1-23%)	1400 (440-2500)	26000 (8100-49000)
Romania	2016	Female	75 to 79	46% (31-65%)	54% (35-69%)	2.6 (1.4-4.3)	8% (1.9-17%)	10% (2.6-21%)	2100 (520-4200)	31000 (7800-63000)
Romania	2016	Female	80 to 84	41% (25-60%)	59% (40-75%)	2.2 (1.2-3.7)	7.3% (1.3-15%)	8.9% (1.6-19%)	2400 (410-5100)	26000 (4500-55000)
Romania	2016	Female	85 to 89	40% (26-57%)	60% (43-74%)	2.2 (1.2-3.6)	6.5% (0.71-15%)	7.7% (0.76-18%)	1800 (160-4000)	13000 (1500-31000)
Romania	2016	Female	90 to 94	41% (27-58%)	59% (42-73%)	2.2 (1.1-3.8)	5.7% (-0.44-15%)	6.9% (-0.64-18%)	770 (-73-2000)	4100 (-300-11000)
Romania	2016	Female	95 plus	45% (27-65%)	55% (35-73%)	2.3 (1-4.3)	5.5% (-1.5-16%)	6.5% (-2-19%)	230 (-71-670)	820 (-210-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Romania	2016	Male	All Ages	71% (62-80%)	29% (20-38%)	8.2 (6.1-11)	18% (8.6-25%)	21% (8.7-31%)	27000 (12000-41000)	680000 (330000-1000000)
Romania	2016	Male	15 to 19	62% (40-81%)	38% (19-60%)	3.5 (1.9-5.7)	7.7% (3.1-13%)	15% (5.2-27%)	50 (17-92)	4500 (1900-7700)
Romania	2016	Male	20 to 24	72% (51-88%)	28% (12-49%)	6.3 (3.7-10)	15% (8-23%)	26% (13-41%)	120 (59-200)	10000 (5600-16000)
Romania	2016	Male	25 to 29	75% (57-90%)	25% (10-43%)	7.6 (4.7-12)	16% (9.3-23%)	29% (15-43%)	190 (96-280)	15000 (8300-21000)
Romania	2016	Male	30 to 34	78% (58-91%)	22% (8.7-42%)	8.9 (5.7-13)	18% (11-25%)	31% (17-44%)	270 (150-390)	18000 (11000-26000)
Romania	2016	Male	35 to 39	80% (61-91%)	20% (8.8-39%)	10 (6.5-16)	19% (12-27%)	32% (19-44%)	490 (300-690)	29000 (18000-41000)
Romania	2016	Male	40 to 44	81% (63-92%)	19% (7.8-37%)	12 (7.4-17)	21% (13-29%)	32% (20-44%)	880 (540-1200)	45000 (27000-62000)
Romania	2016	Male	45 to 49	81% (60-92%)	19% (8-40%)	12 (8-18)	22% (13-30%)	30% (18-41%)	1500 (880-2100)	67000 (39000-93000)
Romania	2016	Male	50 to 54	79% (60-92%)	21% (8.3-40%)	13 (8.2-18)	22% (13-31%)	28% (16-39%)	1700 (970-2400)	66000 (37000-93000)
Romania	2016	Male	55 to 59	78% (57-91%)	22% (8.7-43%)	12 (8-18)	21% (11-30%)	26% (14-37%)	2500 (1300-3600)	84000 (44000-120000)
Romania	2016	Male	60 to 64	76% (53-91%)	24% (8.9-47%)	12 (7.3-18)	19% (9.1-28%)	24% (12-34%)	3300 (1600-4900)	93000 (44000-140000)
Romania	2016	Male	65 to 69	75% (52-90%)	25% (9.7-48%)	10 (6.7-15)	18% (7.6-28%)	22% (9.4-34%)	3400 (1500-5200)	81000 (34000-120000)
Romania	2016	Male	70 to 74	73% (52-88%)	27% (12-48%)	9.1 (5.7-14)	17% (6.6-27%)	21% (8-32%)	3100 (1200-4800)	59000 (22000-92000)
Romania	2016	Male	75 to 79	71% (50-87%)	29% (13-50%)	7.7 (4.5-12)	15% (5-25%)	18% (5.9-29%)	3400 (1100-5500)	50000 (16000-81000)
Romania	2016	Male	80 to 84	69% (45-86%)	31% (14-55%)	6.8 (4-10)	14% (3.9-24%)	17% (4.4-28%)	3300 (890-5600)	37000 (9800-61000)
Romania	2016	Male	85 to 89	68% (45-85%)	32% (15-55%)	6.2 (3.7-9.6)	13% (3.4-23%)	15% (3.9-26%)	2100 (540-3600)	16000 (4300-28000)
Romania	2016	Male	90 to 94	68% (48-86%)	32% (14-52%)	5.7 (3.2-9.3)	11% (2.4-21%)	13% (2.7-25%)	740 (150-1400)	4000 (890-7500)
Romania	2016	Male	95 plus	68% (42-88%)	32% (12-58%)	5 (2.3-9.1)	9.2% (0.36-19%)	11% (-0.021-22%)	140 (-0.28-300)	520 (20-1100)
Serbia	1990	Female	All Ages	59% (50-67%)	41% (33-50%)	1.8 (1.2-2.6)	5.1% (1.2-10%)	7% (0.95-15%)	2800 (380-6200)	69000 (15000-140000)
Serbia	1990	Female	15 to 19	59% (39-76%)	41% (24-61%)	1.4 (0.56-2.8)	2.9% (0.8-5.8%)	6.7% (0.043-15%)	8.8 (0.056-21)	1000 (280-2100)
Serbia	1990	Female	20 to 24	68% (50-83%)	32% (17-50%)	2 (0.74-3.9)	4.6% (1.8-8.3%)	8.4% (-0.058-19%)	12 (-0.077-28)	1700 (630-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	1990	Female	25 to 29	70% (50-84%)	30% (16-50%)	2.1 (0.86-3.9)	4.6% (1.8-8.1%)	8.3% (-0.16-18%)	15 (-0.28-33)	1800 (710-3200)
Serbia	1990	Female	30 to 34	70% (50-84%)	30% (16-50%)	2.2 (1-3.9)	4.5% (1.6-7.9%)	8.6% (0.63-18%)	21 (1.5-46)	2100 (720-3800)
Serbia	1990	Female	35 to 39	71% (54-85%)	29% (15-46%)	2.5 (1.1-4.5)	4.8% (1.4-9.3%)	9.3% (1.1-20%)	41 (5-88)	3100 (860-5900)
Serbia	1990	Female	40 to 44	70% (53-84%)	30% (16-47%)	2.8 (1.3-4.9)	5.4% (1.3-11%)	10% (2.2-21%)	63 (13-130)	3800 (1000-7200)
Serbia	1990	Female	45 to 49	68% (47-83%)	32% (17-53%)	3.1 (1.4-5.7)	6.4% (1.7-13%)	12% (3-23%)	79 (22-150)	3900 (1100-7500)
Serbia	1990	Female	50 to 54	66% (45-81%)	34% (19-55%)	3.1 (1.5-5.8)	7.2% (2-14%)	12% (3.2-23%)	150 (43-290)	6300 (1800-12000)
Serbia	1990	Female	55 to 59	63% (42-80%)	37% (20-58%)	3.1 (1.4-5.3)	7.5% (1.7-15%)	11% (2.7-22%)	250 (59-490)	8800 (2000-17000)
Serbia	1990	Female	60 to 64	59% (39-78%)	41% (22-61%)	2.8 (1.3-5.4)	6.9% (0.59-15%)	9.8% (0.95-21%)	340 (33-730)	9900 (820-22000)
Serbia	1990	Female	65 to 69	55% (35-74%)	45% (26-65%)	2.4 (1.1-4.5)	6.6% (0.28-15%)	8.8% (0.54-20%)	400 (24-920)	9600 (400-22000)
Serbia	1990	Female	70 to 74	48% (31-67%)	52% (33-69%)	2 (0.79-3.9)	5.6% (-0.72-14%)	7.2% (-0.68-18%)	270 (-28-690)	5100 (-700-13000)
Serbia	1990	Female	75 to 79	41% (26-61%)	59% (39-74%)	1.6 (0.58-3.4)	4.6% (-0.78-12%)	5.7% (-0.88-15%)	340 (-54-930)	5000 (-840-14000)
Serbia	1990	Female	80 to 84	37% (22-55%)	63% (45-78%)	1.4 (0.56-2.8)	4.5% (-0.44-12%)	5.4% (-0.48-14%)	340 (-32-890)	3600 (-380-9500)
Serbia	1990	Female	85 to 89	35% (22-51%)	65% (49-78%)	1.4 (0.55-3.1)	4.5% (-0.37-12%)	5.4% (-0.4-15%)	290 (-22-830)	2200 (-180-6200)
Serbia	1990	Female	90 to 94	37% (23-54%)	63% (46-77%)	1.5 (0.56-3.6)	4.6% (-0.35-13%)	5.5% (-0.33-16%)	150 (-8.8-460)	780 (-55-2300)
Serbia	1990	Female	95 plus	41% (23-62%)	59% (38-77%)	1.7 (0.46-4.2)	4.9% (-0.12-14%)	5.8% (-0.11-17%)	51 (-0.96-150)	170 (-4.5-490)
Serbia	1990	Male	All Ages	67% (56-77%)	33% (23-44%)	2.9 (2-3.9)	9.7% (4-16%)	11% (3.4-20%)	5200 (1600-9000)	16000 (64000-260000)
Serbia	1990	Male	15 to 19	59% (37-80%)	41% (20-63%)	1.3 (0.58-2.6)	5% (1.1-9.9%)	7.5% (0.32-17%)	26 (1.1-58)	2400 (540-4800)
Serbia	1990	Male	20 to 24	70% (49-87%)	30% (13-51%)	2.4 (1.1-4.4)	9.9% (3-18%)	14% (1.8-27%)	65 (9-130)	5600 (1700-10000)
Serbia	1990	Male	25 to 29	74% (54-90%)	26% (10-46%)	3.1 (1.5-5.4)	11% (4-19%)	16% (3.3-30%)	77 (15-150)	6200 (2200-11000)
Serbia	1990	Male	30 to 34	76% (56-91%)	24% (8.7-44%)	3.6 (1.8-6)	12% (4.4-19%)	17% (4.3-31%)	100 (26-190)	7400 (2700-13000)
Serbia	1990	Male	35 to 39	78% (59-91%)	22% (9-41%)	3.8 (1.9-6.5)	12% (4.4-20%)	17% (4.7-31%)	160 (41-290)	10000 (3600-17000)
Serbia	1990	Male	40 to 44	79% (59-92%)	21% (8.4-41%)	4.2 (2.3-7.1)	12% (5.2-21%)	18% (6.4-31%)	210 (78-370)	12000 (4800-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	1990	Male	45 to 49	79% (58-91%)	21% (8.7-42%)	4.9 (2.5-8.1)	14% (6.2-22%)	18% (7.7-30%)	240 (100-390)	11000 (5000-18000)
Serbia	1990	Male	50 to 54	78% (57-91%)	22% (9.2-43%)	5.2 (2.9-8.4)	14% (6.6-22%)	18% (8.2-28%)	440 (200-680)	18000 (8400-27000)
Serbia	1990	Male	55 to 59	76% (56-91%)	24% (9.3-44%)	5.3 (2.9-8.5)	14% (5.9-22%)	17% (7.2-27%)	690 (290-1100)	24000 (10000-38000)
Serbia	1990	Male	60 to 64	75% (53-91%)	25% (9.5-47%)	5.1 (2.7-8.8)	12% (4.6-21%)	15% (5.3-25%)	820 (300-1400)	24000 (8900-41000)
Serbia	1990	Male	65 to 69	73% (51-90%)	27% (10-49%)	4.6 (2.5-7.5)	11% (3.4-20%)	13% (3.8-23%)	750 (220-1300)	18000 (5300-33000)
Serbia	1990	Male	70 to 74	71% (50-87%)	29% (13-50%)	4 (2-7)	9.5% (1.6-18%)	11% (1.9-21%)	440 (73-870)	8500 (1400-17000)
Serbia	1990	Male	75 to 79	69% (47-86%)	31% (14-53%)	3.4 (1.5-6.3)	7.7% (0.042-17%)	8.6% (-0.026-19%)	480 (-1.5-1100)	7100 (37-16000)
Serbia	1990	Male	80 to 84	67% (43-84%)	33% (16-57%)	3 (1.4-5.5)	6.4% (-0.47-15%)	7.1% (-0.67-17%)	370 (-36-870)	4100 (-320-9500)
Serbia	1990	Male	85 to 89	65% (43-84%)	35% (16-57%)	2.8 (1.3-5.1)	4.8% (-1.5-13%)	5.3% (-1.8-15%)	210 (-72-580)	1700 (-500-4500)
Serbia	1990	Male	90 to 94	65% (43-85%)	35% (15-57%)	2.6 (1.2-4.9)	3.6% (-1.9-12%)	3.9% (-2.3-13%)	65 (-38-210)	360 (-180-1100)
Serbia	1990	Male	95 plus	66% (38-88%)	34% (12-62%)	2.3 (0.82-5)	3.3% (-2-11%)	3.5% (-2.4-12%)	14 (-10-49)	53 (-31-170)
Serbia	1995	Female	All Ages	59% (51-67%)	41% (33-49%)	1.9 (1.3-2.5)	5.2% (1.2-10%)	7% (0.89-15%)	3500 (430-7700)	82000 (18000-160000)
Serbia	1995	Female	15 to 19	59% (40-75%)	41% (25-60%)	1.4 (0.59-2.6)	2.4% (0.62-4.6%)	3.7% (0.042-8.7%)	9.1 (0.11-22)	1100 (290-2100)
Serbia	1995	Female	20 to 24	68% (51-83%)	32% (17-49%)	2 (0.76-3.9)	3.9% (1.6-7%)	5.2% (-0.069-12%)	13 (-0.18-30)	1800 (710-3200)
Serbia	1995	Female	25 to 29	70% (51-84%)	30% (16-49%)	2.1 (0.84-3.9)	4.2% (1.7-7.2%)	6% (-0.12-13%)	15 (-0.3-34)	1900 (760-3300)
Serbia	1995	Female	30 to 34	71% (51-84%)	29% (16-49%)	2.2 (1-4)	4.1% (1.5-7.3%)	6.7% (0.43-14%)	21 (1.4-46)	2100 (720-3800)
Serbia	1995	Female	35 to 39	71% (55-85%)	29% (15-45%)	2.5 (1.1-4.4)	4.6% (1.3-9%)	8% (0.71-16%)	39 (3.3-79)	2800 (780-5200)
Serbia	1995	Female	40 to 44	71% (53-85%)	29% (15-47%)	2.8 (1.3-4.7)	5.4% (1.4-10%)	9.5% (1.9-19%)	80 (15-160)	4600 (1200-9000)
Serbia	1995	Female	45 to 49	69% (47-83%)	31% (17-53%)	3.1 (1.4-5.3)	6.3% (1.8-13%)	11% (2.8-21%)	120 (31-220)	5600 (1600-11000)
Serbia	1995	Female	50 to 54	66% (47-82%)	34% (18-53%)	3.1 (1.5-5.4)	7.3% (1.9-14%)	11% (3.2-22%)	150 (41-280)	5900 (1600-11000)
Serbia	1995	Female	55 to 59	63% (43-80%)	37% (20-57%)	3 (1.5-5.3)	7.7% (1.6-15%)	11% (2.4-22%)	260 (56-510)	9000 (1800-18000)
Serbia	1995	Female	60 to 64	60% (39-79%)	40% (21-61%)	2.8 (1.3-5.2)	7.1% (0.43-15%)	9.7% (0.66-21%)	390 (27-850)	11000 (650-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	1995	Female	65 to 69	55% (36-75%)	45% (25-64%)	2.4 (1.2-4.2)	6.7% (0.018-15%)	8.8% (0.26-19%)	550 (15-1200)	13000 (35-30000)
Serbia	1995	Female	70 to 74	49% (31-67%)	51% (33-69%)	2 (0.81-3.8)	5.8% (-1.1-15%)	7.2% (-1.2-18%)	580 (-94-1400)	11000 (-2000-27000)
Serbia	1995	Female	75 to 79	42% (26-62%)	58% (38-74%)	1.6 (0.58-3.3)	4.8% (-0.66-13%)	5.9% (-0.74-16%)	330 (-42-880)	4800 (-700-13000)
Serbia	1995	Female	80 to 84	37% (22-56%)	63% (44-78%)	1.4 (0.57-2.7)	4.7% (-0.47-12%)	5.6% (-0.56-14%)	360 (-36-940)	3800 (-390-9900)
Serbia	1995	Female	85 to 89	36% (22-52%)	64% (48-78%)	1.4 (0.57-3)	4.8% (-0.3-13%)	5.6% (-0.34-15%)	370 (-22-1000)	2700 (-170-7600)
Serbia	1995	Female	90 to 94	37% (23-54%)	63% (46-77%)	1.5 (0.56-3.4)	4.8% (-0.18-14%)	5.6% (-0.16-17%)	210 (-5.9-620)	1000 (-38-3100)
Serbia	1995	Female	95 plus	41% (23-62%)	59% (38-77%)	1.7 (0.47-4.2)	5% (-0.43-15%)	5.8% (-0.48-17%)	74 (-6.1-220)	250 (-22-730)
Serbia	1995	Male	All Ages	67% (56-77%)	33% (23-44%)	2.9 (2.1-3.9)	9.2% (3.7-15%)	11% (3.2-19%)	6200 (1800-11000)	180000 (73000-300000)
Serbia	1995	Male	15 to 19	59% (38-80%)	41% (20-62%)	1.4 (0.61-2.6)	3.4% (0.86-6.6%)	3.9% (0.12-8.6%)	23 (0.69-51)	2300 (570-4300)
Serbia	1995	Male	20 to 24	70% (47-87%)	30% (13-53%)	2.4 (1.1-4.2)	5.7% (2.1-10%)	5.8% (0.97-12%)	55 (9.4-110)	5100 (1800-8900)
Serbia	1995	Male	25 to 29	74% (54-89%)	26% (11-46%)	3.1 (1.5-5.3)	7.3% (2.9-12%)	8.3% (1.8-16%)	68 (15-130)	5700 (2200-9700)
Serbia	1995	Male	30 to 34	76% (56-91%)	24% (9.4-44%)	3.5 (1.8-5.8)	8.6% (3.3-14%)	11% (2.7-20%)	88 (22-160)	6500 (2500-11000)
Serbia	1995	Male	35 to 39	78% (59-91%)	22% (8.5-41%)	3.7 (1.9-6.3)	10% (3.6-17%)	13% (3.5-24%)	140 (36-250)	8800 (3200-15000)
Serbia	1995	Male	40 to 44	79% (59-92%)	21% (7.8-41%)	4.2 (2.3-6.7)	12% (4.8-20%)	15% (5.5-27%)	260 (93-450)	14000 (5700-24000)
Serbia	1995	Male	45 to 49	79% (56-92%)	21% (8.4-44%)	4.8 (2.6-7.8)	13% (5.7-21%)	17% (7-28%)	360 (150-590)	17000 (7300-27000)
Serbia	1995	Male	50 to 54	78% (57-91%)	22% (9.3-43%)	5.1 (2.8-8)	14% (6.4-21%)	17% (7.5-27%)	400 (180-640)	16000 (7400-25000)
Serbia	1995	Male	55 to 59	76% (55-91%)	24% (9-45%)	5.3 (2.9-8.2)	14% (5.8-22%)	16% (6.8-26%)	700 (290-1100)	24000 (10000-38000)
Serbia	1995	Male	60 to 64	75% (51-91%)	25% (9.3-49%)	5.1 (2.6-8.5)	12% (4.1-21%)	14% (4.6-24%)	970 (320-1700)	28000 (9200-48000)
Serbia	1995	Male	65 to 69	74% (50-90%)	26% (10-50%)	4.6 (2.6-7.3)	11% (3.2-20%)	13% (3.6-23%)	1100 (310-2000)	26000 (7500-47000)
Serbia	1995	Male	70 to 74	72% (50-87%)	28% (13-50%)	3.9 (2.1-6.7)	9.6% (1.4-19%)	11% (1.6-22%)	880 (130-1700)	17000 (2400-33000)
Serbia	1995	Male	75 to 79	69% (47-87%)	31% (13-53%)	3.3 (1.6-6.1)	7.6% (0.24-16%)	8.6% (0.18-19%)	420 (9.1-900)	6200 (200-13000)
Serbia	1995	Male	80 to 84	67% (44-84%)	33% (16-56%)	3 (1.5-5.1)	6.5% (-0.82-15%)	7.1% (-1-16%)	360 (-50-800)	3900 (-490-8700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	1995	Male	85 to 89	65% (44-83%)	35% (17-56%)	2.7 (1.3-4.8)	4.9% (-1.3-13%)	5.3% (-1.6-14%)	230 (-68-600)	1800 (-460-4700)
Serbia	1995	Male	90 to 94	65% (44-85%)	35% (15-56%)	2.5 (1.2-4.8)	3.6% (-1.7-11%)	3.9% (-2-12%)	86 (-45-280)	470 (-220-1500)
Serbia	1995	Male	95 plus	65% (39-88%)	35% (12-61%)	2.3 (0.83-4.9)	3.2% (-1.9-10%)	3.4% (-2.4-12%)	20 (-14-69)	75 (-45-250)
Serbia	2000	Female	All Ages	59% (50-67%)	41% (33-50%)	2 (1.5-2.6)	5.7% (1.3-11%)	7.5% (1.1-16%)	4000 (560-8800)	90000 (21000-180000)
Serbia	2000	Female	15 to 19	59% (39-75%)	41% (25-61%)	1.4 (0.63-2.7)	2.7% (0.85-5.1%)	6.1% (0.23-14%)	7.2 (0.28-16)	880 (270-1700)
Serbia	2000	Female	20 to 24	68% (50-83%)	32% (17-50%)	2 (0.8-3.9)	4.5% (1.9-7.7%)	7.9% (0.17-17%)	12 (0.25-26)	1600 (690-2900)
Serbia	2000	Female	25 to 29	70% (51-84%)	30% (16-49%)	2.2 (0.93-4.1)	4.5% (1.9-7.5%)	8.1% (0.48-18%)	13 (0.77-29)	1700 (730-3000)
Serbia	2000	Female	30 to 34	70% (51-84%)	30% (16-49%)	2.3 (1.1-4.2)	4.4% (1.7-7.8%)	8.5% (0.81-18%)	20 (1.8-41)	2000 (760-3400)
Serbia	2000	Female	35 to 39	71% (53-85%)	29% (15-47%)	2.5 (1.1-4.4)	4.8% (1.4-8.9%)	9.3% (1.3-19%)	33 (4.6-68)	2400 (730-4500)
Serbia	2000	Female	40 to 44	70% (53-84%)	30% (16-47%)	2.8 (1.4-4.8)	5.5% (1.5-11%)	10% (2.4-20%)	62 (14-120)	3600 (970-6800)
Serbia	2000	Female	45 to 49	69% (47-84%)	31% (16-53%)	3.1 (1.6-5.4)	6.7% (1.9-13%)	12% (3.2-22%)	140 (38-260)	6500 (1900-12000)
Serbia	2000	Female	50 to 54	66% (46-82%)	34% (18-54%)	3.2 (1.6-5.3)	7.3% (2.2-14%)	12% (3.9-22%)	180 (59-340)	7500 (2300-14000)
Serbia	2000	Female	55 to 59	63% (43-81%)	37% (19-57%)	3.2 (1.5-5.4)	8% (2-15%)	12% (3-22%)	220 (56-420)	7500 (1900-14000)
Serbia	2000	Female	60 to 64	59% (39-78%)	41% (22-61%)	2.9 (1.4-5.3)	7.4% (0.88-16%)	10% (1.4-21%)	370 (48-770)	11000 (1300-22000)
Serbia	2000	Female	65 to 69	55% (35-74%)	45% (26-65%)	2.5 (1.3-4.3)	7.1% (0.6-15%)	9.3% (0.9-20%)	590 (57-1300)	14000 (1100-31000)
Serbia	2000	Female	70 to 74	49% (31-66%)	51% (34-69%)	2 (0.89-3.8)	6.1% (-0.55-15%)	7.6% (-0.61-19%)	710 (-58-1700)	13000 (-1300-33000)
Serbia	2000	Female	75 to 79	42% (26-61%)	58% (39-74%)	1.6 (0.66-3.2)	5.2% (-0.7-13%)	6.2% (-0.73-16%)	670 (-81-1700)	9700 (-1300-26000)
Serbia	2000	Female	80 to 84	37% (22-55%)	63% (45-78%)	1.5 (0.61-2.7)	5% (-0.19-13%)	6% (-0.23-15%)	470 (-18-1200)	5000 (-180-13000)
Serbia	2000	Female	85 to 89	35% (22-51%)	65% (49-78%)	1.5 (0.6-2.9)	5.1% (0.21-13%)	6% (0.31-16%)	290 (14-760)	2200 (91-5600)
Serbia	2000	Female	90 to 94	37% (24-54%)	63% (46-76%)	1.6 (0.59-3.4)	5% (0.065-14%)	5.9% (0.097-16%)	190 (3.1-530)	960 (13-2700)
Serbia	2000	Female	95 plus	41% (24-62%)	59% (38-76%)	1.8 (0.48-4.1)	5.1% (-0.26-15%)	6% (-0.23-18%)	74 (-2.9-230)	250 (-12-750)
Serbia	2000	Male	All Ages	68% (57-78%)	32% (22-43%)	3.1 (2.4-3.9)	11% (4.7-17%)	12% (4.2-21%)	6900 (2400-12000)	190000 (82000-310000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	2000	Male	15 to 19	59% (36-80%)	41% (20-64%)	1.4 (0.66-2.5)	4.5% (1.2-8.6%)	6.9% (0.36-15%)	19 (0.98-42)	1900 (530-3600)
Serbia	2000	Male	20 to 24	70% (48-87%)	30% (13-52%)	2.5 (1.2-4.2)	8.7% (3.4-15%)	12% (2.2-23%)	52 (9.2-100)	4700 (1800-8400)
Serbia	2000	Male	25 to 29	74% (53-89%)	26% (11-47%)	3.2 (1.7-5.5)	9.9% (4.3-17%)	15% (3.9-28%)	62 (15-120)	5300 (2300-8900)
Serbia	2000	Male	30 to 34	76% (56-91%)	24% (9.4-44%)	3.7 (2-6)	10% (4.4-17%)	16% (4.8-29%)	76 (22-140)	5800 (2400-9500)
Serbia	2000	Male	35 to 39	78% (59-91%)	22% (8.9-41%)	3.9 (2-6.4)	11% (4.7-20%)	17% (5.1-30%)	110 (35-200)	7300 (2900-12000)
Serbia	2000	Male	40 to 44	79% (59-92%)	21% (8.3-41%)	4.4 (2.5-6.9)	13% (5.3-21%)	18% (6.4-31%)	210 (77-350)	11000 (4800-19000)
Serbia	2000	Male	45 to 49	79% (57-92%)	21% (8.3-43%)	5.1 (2.9-7.8)	14% (6.6-22%)	19% (8-30%)	420 (180-690)	19000 (9100-31000)
Serbia	2000	Male	50 to 54	78% (57-91%)	22% (9.1-43%)	5.4 (3.3-8)	14% (7.2-22%)	18% (8.8-28%)	550 (270-860)	22000 (11000-34000)
Serbia	2000	Male	55 to 59	76% (55-91%)	24% (8.6-45%)	5.5 (3.2-8.4)	14% (6.8-22%)	17% (8.2-27%)	590 (280-930)	20000 (9600-31000)
Serbia	2000	Male	60 to 64	75% (51-91%)	25% (9.2-49%)	5.3 (2.8-8.4)	13% (5-22%)	15% (5.9-26%)	880 (340-1500)	25000 (9900-43000)
Serbia	2000	Male	65 to 69	74% (50-90%)	26% (9.7-50%)	4.8 (2.8-7.2)	12% (4.2-21%)	14% (4.8-24%)	1200 (420-2100)	29000 (9800-50000)
Serbia	2000	Male	70 to 74	72% (50-87%)	28% (13-50%)	4.1 (2.2-6.9)	10% (2.4-19%)	12% (2.7-22%)	1200 (270-2200)	23000 (5100-43000)
Serbia	2000	Male	75 to 79	69% (47-86%)	31% (14-53%)	3.5 (1.7-6)	8.3% (0.89-17%)	9.4% (0.96-19%)	810 (82-1700)	12000 (1300-24000)
Serbia	2000	Male	80 to 84	67% (44-85%)	33% (15-56%)	3.1 (1.6-5.2)	7.1% (0.13-15%)	7.9% (0.032-17%)	420 (1.7-900)	4600 (87-9800)
Serbia	2000	Male	85 to 89	66% (43-83%)	34% (17-57%)	2.8 (1.4-5)	5.5% (-0.57-13%)	6.1% (-0.74-14%)	170 (-20-390)	1300 (-120-3000)
Serbia	2000	Male	90 to 94	65% (44-86%)	35% (14-56%)	2.6 (1.3-4.9)	4.2% (-0.99-11%)	4.6% (-1.4-13%)	74 (-22-210)	400 (-98-1100)
Serbia	2000	Male	95 plus	66% (39-88%)	34% (12-61%)	2.4 (0.9-4.8)	3.6% (-1.5-11%)	3.9% (-1.9-12%)	22 (-11-69)	79 (-33-240)
Serbia	2005	Female	All Ages	59% (51-67%)	41% (33-49%)	2.2 (1.8-2.7)	6% (1.5-11%)	7.8% (1.1-16%)	4300 (630-8800)	91000 (23000-170000)
Serbia	2005	Female	15 to 19	59% (39-75%)	41% (25-61%)	1.6 (0.71-2.9)	2.7% (0.96-4.9%)	7.1% (0.43-16%)	5.9 (0.37-13)	800 (290-1500)
Serbia	2005	Female	20 to 24	68% (51-83%)	32% (17-49%)	2.2 (0.95-4.3)	4.6% (2.1-7.7%)	9% (0.67-19%)	10 (0.75-22)	1500 (690-2500)
Serbia	2005	Female	25 to 29	70% (51-84%)	30% (16-49%)	2.4 (1-4.4)	4.6% (2.2-7.5%)	9% (1.2-19%)	12 (1.6-26)	1700 (800-2900)
Serbia	2005	Female	30 to 34	71% (51-84%)	29% (16-49%)	2.6 (1.2-4.6)	4.6% (2.1-7.9%)	9.4% (1.7-19%)	19 (3.3-37)	2000 (840-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	2005	Female	35 to 39	71% (55-85%)	29% (15-45%)	2.8 (1.3-4.9)	4.8% (1.8-8.8%)	10% (2.1-20%)	28 (5.9-55)	2200 (780-4000)
Serbia	2005	Female	40 to 44	70% (53-84%)	30% (16-47%)	3.1 (1.5-5.2)	5.8% (1.9-11%)	11% (3-20%)	57 (16-110)	3400 (1100-6200)
Serbia	2005	Female	45 to 49	69% (47-83%)	31% (17-53%)	3.5 (1.7-6.1)	6.9% (2.3-13%)	12% (3.9-22%)	110 (37-200)	5300 (1800-9700)
Serbia	2005	Female	50 to 54	66% (47-82%)	34% (18-53%)	3.6 (1.9-5.8)	7.5% (2.5-14%)	12% (4.4-21%)	200 (75-370)	8400 (3000-15000)
Serbia	2005	Female	55 to 59	63% (43-80%)	37% (20-57%)	3.5 (1.8-5.9)	7.8% (2.5-15%)	12% (4.1-22%)	260 (95-480)	9300 (3100-17000)
Serbia	2005	Female	60 to 64	60% (39-78%)	40% (22-61%)	3.2 (1.6-5.8)	7.7% (1.5-15%)	11% (2.5-21%)	280 (65-560)	8300 (1600-17000)
Serbia	2005	Female	65 to 69	55% (35-75%)	45% (25-65%)	2.8 (1.4-4.6)	7.5% (1.1-16%)	9.8% (1.6-21%)	500 (84-1000)	12000 (1800-25000)
Serbia	2005	Female	70 to 74	49% (31-67%)	51% (33-69%)	2.3 (1-4.2)	6.4% (-0.36-15%)	8.1% (-0.29-19%)	710 (-25-1600)	13000 (-770-32000)
Serbia	2005	Female	75 to 79	42% (26-62%)	58% (38-74%)	1.8 (0.74-3.4)	5.5% (-0.61-14%)	6.6% (-0.65-16%)	790 (-78-2000)	11000 (-1200-28000)
Serbia	2005	Female	80 to 84	37% (22-55%)	63% (45-78%)	1.6 (0.68-2.9)	5.4% (-0.061-13%)	6.4% (0.01-15%)	660 (0.99-1600)	7100 (-82-17000)
Serbia	2005	Female	85 to 89	36% (22-52%)	64% (48-78%)	1.6 (0.66-3.2)	5.5% (0.45-14%)	6.6% (0.51-17%)	380 (31-980)	2800 (230-7200)
Serbia	2005	Female	90 to 94	37% (24-55%)	63% (45-76%)	1.8 (0.65-3.9)	5.5% (0.36-15%)	6.5% (0.48-18%)	150 (11-400)	750 (47-2000)
Serbia	2005	Female	95 plus	41% (24-62%)	59% (38-76%)	2 (0.55-4.6)	5.5% (-0.19-16%)	6.5% (-0.17-19%)	66 (-1.7-200)	220 (-7.2-650)
Serbia	2005	Male	All Ages	68% (58-79%)	32% (21-42%)	3.5 (2.8-4.3)	11% (5.1-17%)	13% (4.9-21%)	7100 (2700-12000)	190000 (89000-300000)
Serbia	2005	Male	15 to 19	60% (37-81%)	40% (19-63%)	1.6 (0.74-2.7)	4.8% (1.5-8.9%)	8.2% (0.92-18%)	17 (1.9-37)	1800 (570-3300)
Serbia	2005	Male	20 to 24	70% (49-87%)	30% (13-51%)	2.8 (1.4-4.6)	9.8% (3.9-17%)	15% (3-28%)	48 (10-94)	4400 (1800-7600)
Serbia	2005	Male	25 to 29	74% (54-90%)	26% (10-46%)	3.5 (1.9-5.9)	11% (5-18%)	17% (5-32%)	64 (18-120)	5400 (2400-9100)
Serbia	2005	Male	30 to 34	76% (55-91%)	24% (9.3-45%)	4 (2.2-6.4)	11% (4.9-18%)	19% (6.2-33%)	77 (26-140)	5900 (2600-9600)
Serbia	2005	Male	35 to 39	78% (59-91%)	22% (9-41%)	4.3 (2.3-6.8)	12% (5.5-20%)	19% (7.1-33%)	100 (38-180)	6800 (3100-11000)
Serbia	2005	Male	40 to 44	79% (59-91%)	21% (8.5-41%)	4.8 (2.8-7.5)	13% (5.8-22%)	20% (7.8-32%)	180 (72-300)	9900 (4300-16000)
Serbia	2005	Male	45 to 49	79% (57-92%)	21% (8.3-43%)	5.6 (3.2-8.5)	15% (7.4-23%)	20% (9.8-31%)	350 (160-550)	16000 (7800-25000)
Serbia	2005	Male	50 to 54	78% (58-91%)	22% (9-42%)	5.9 (3.7-8.7)	15% (8.1-23%)	19% (10-29%)	630 (340-960)	25000 (14000-38000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	2005	Male	55 to 59	76% (56-91%)	24% (8.8-44%)	6.1 (3.6-9.1)	14% (7.5-22%)	18% (9.3-28%)	740 (390-1200)	26000 (13000-40000)
Serbia	2005	Male	60 to 64	75% (53-91%)	25% (9.1-47%)	5.8 (3.2-9.3)	13% (5.7-22%)	16% (6.8-26%)	700 (300-1100)	20000 (8600-33000)
Serbia	2005	Male	65 to 69	74% (50-90%)	26% (10-50%)	5.3 (3.2-7.9)	13% (5.4-21%)	15% (6.2-24%)	1100 (440-1800)	25000 (11000-42000)
Serbia	2005	Male	70 to 74	72% (51-87%)	28% (13-49%)	4.5 (2.6-7.5)	11% (3.1-20%)	12% (3.6-22%)	1200 (370-2200)	24000 (7000-42000)
Serbia	2005	Male	75 to 79	69% (47-86%)	31% (14-53%)	3.8 (2-6.5)	8.9% (1.5-17%)	10% (1.7-19%)	1000 (170-2000)	15000 (2400-30000)
Serbia	2005	Male	80 to 84	67% (42-84%)	33% (16-58%)	3.4 (1.8-5.6)	7.7% (0.89-15%)	8.7% (1.1-18%)	610 (76-1200)	6700 (770-13000)
Serbia	2005	Male	85 to 89	65% (43-83%)	35% (17-57%)	3.1 (1.6-5.2)	6.4% (0.26-14%)	7.2% (0.16-16%)	230 (5.1-490)	1800 (74-3800)
Serbia	2005	Male	90 to 94	65% (44-84%)	35% (16-56%)	2.9 (1.4-5.1)	5.1% (-0.39-13%)	5.6% (-0.59-15%)	59 (-6.3-150)	320 (-25-800)
Serbia	2005	Male	95 plus	65% (39-88%)	35% (12-61%)	2.6 (1-5.3)	4.3% (-1.2-12%)	4.7% (-1.6-14%)	19 (-6.5-56)	68 (-19-200)
Serbia	2010	Female	All Ages	59% (51-67%)	41% (33-49%)	2.5 (2-3.1)	5.9% (1.6-11%)	7.7% (1.3-16%)	4100 (730-8600)	87000 (25000-160000)
Serbia	2010	Female	15 to 19	59% (39-76%)	41% (24-61%)	1.7 (0.74-3.2)	2.8% (1-5%)	7.4% (0.63-16%)	5.1 (0.36-11)	710 (260-1200)
Serbia	2010	Female	20 to 24	68% (50-83%)	32% (17-50%)	2.5 (1-4.6)	4.7% (2.1-7.8%)	9.4% (1.1-20%)	9 (1.1-19)	1400 (650-2300)
Serbia	2010	Female	25 to 29	70% (50-84%)	30% (16-50%)	2.6 (1.1-4.7)	4.9% (2.3-7.9%)	9.5% (1.4-20%)	12 (1.7-24)	1600 (740-2600)
Serbia	2010	Female	30 to 34	71% (50-84%)	29% (16-50%)	2.8 (1.3-4.8)	4.7% (2.2-7.9%)	9.8% (2-19%)	18 (3.8-35)	1900 (870-3200)
Serbia	2010	Female	35 to 39	71% (54-85%)	29% (15-46%)	3.1 (1.4-5.4)	4.9% (2-8.7%)	10% (2.6-20%)	28 (6.9-55)	2400 (970-4100)
Serbia	2010	Female	40 to 44	70% (53-84%)	30% (16-47%)	3.4 (1.7-5.8)	5.7% (2.1-10%)	11% (3.8-20%)	51 (17-94)	3200 (1200-5700)
Serbia	2010	Female	45 to 49	69% (47-83%)	31% (17-53%)	3.8 (1.9-6.8)	6.7% (2.5-12%)	12% (4.6-22%)	93 (35-170)	4700 (1700-8500)
Serbia	2010	Female	50 to 54	66% (47-81%)	34% (19-53%)	3.9 (2.1-6.6)	7.4% (2.8-13%)	12% (4.8-21%)	150 (63-260)	6500 (2500-11000)
Serbia	2010	Female	55 to 59	63% (44-79%)	37% (21-56%)	3.8 (1.9-6.3)	7.7% (2.6-14%)	12% (4.5-21%)	270 (100-490)	10000 (3500-18000)
Serbia	2010	Female	60 to 64	59% (38-78%)	41% (22-62%)	3.5 (1.7-6.2)	7.2% (1.9-14%)	11% (3.2-20%)	320 (95-620)	10000 (2600-19000)
Serbia	2010	Female	65 to 69	55% (36-75%)	45% (25-64%)	3 (1.6-4.9)	7.3% (1.5-15%)	9.9% (2-20%)	380 (75-770)	9300 (1700-19000)
Serbia	2010	Female	70 to 74	49% (31-67%)	51% (33-69%)	2.5 (1.1-4.6)	6.3% (0.034-15%)	8.1% (0.21-19%)	570 (15-1300)	11000 (62-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	2010	Female	75 to 79	42% (26-60%)	58% (40-74%)	2 (0.79-3.8)	5.5% (-0.45-14%)	6.7% (-0.52-16%)	740 (-54-1800)	11000 (-910-27000)
Serbia	2010	Female	80 to 84	37% (22-55%)	63% (45-78%)	1.8 (0.75-3.2)	5.4% (0.023-13%)	6.4% (0.0082-16%)	700 (0.91-1700)	7600 (36-18000)
Serbia	2010	Female	85 to 89	35% (22-51%)	65% (49-78%)	1.8 (0.73-3.5)	5.5% (0.42-14%)	6.5% (0.52-17%)	540 (44-1400)	4100 (300-11000)
Serbia	2010	Female	90 to 94	37% (23-54%)	63% (46-77%)	1.9 (0.72-4)	5.4% (0.39-15%)	6.4% (0.47-18%)	200 (14-550)	1000 (73-2800)
Serbia	2010	Female	95 plus	40% (23-61%)	60% (39-77%)	2.1 (0.56-4.9)	5.4% (-0.15-15%)	6.4% (-0.15-18%)	52 (-1.2-150)	180 (-4.7-500)
Serbia	2010	Male	All Ages	69% (59-78%)	31% (22-41%)	4 (3.2-4.9)	11% (5.3-18%)	13% (5-21%)	7200 (2900-12000)	190000 (90000-300000)
Serbia	2010	Male	15 to 19	59% (38-80%)	41% (20-62%)	1.7 (0.81-3)	4.9% (1.7-9%)	8.9% (1.1-19%)	14 (1.8-30)	1500 (550-2800)
Serbia	2010	Male	20 to 24	70% (49-87%)	30% (13-51%)	3.1 (1.6-5.2)	10% (4.3-17%)	16% (3.6-30%)	41 (9.4-81)	4000 (1700-6800)
Serbia	2010	Male	25 to 29	74% (54-89%)	26% (11-46%)	3.9 (2-6.4)	12% (5.3-19%)	19% (6-35%)	60 (18-110)	5100 (2300-8400)
Serbia	2010	Male	30 to 34	76% (56-90%)	24% (9.5-44%)	4.4 (2.4-7)	12% (5.5-19%)	20% (7-35%)	81 (27-140)	6100 (2800-10000)
Serbia	2010	Male	35 to 39	79% (61-91%)	21% (8.7-39%)	4.7 (2.6-7.4)	12% (5.6-20%)	20% (8.2-35%)	100 (42-180)	7000 (3400-11000)
Serbia	2010	Male	40 to 44	79% (59-92%)	21% (8.3-41%)	5.3 (3.1-8.1)	13% (6.3-21%)	21% (9.1-34%)	170 (73-270)	9500 (4500-15000)
Serbia	2010	Male	45 to 49	79% (55-92%)	21% (8.4-45%)	6.1 (3.5-9.6)	15% (8.2-23%)	21% (11-32%)	300 (160-470)	14000 (7600-22000)
Serbia	2010	Male	50 to 54	78% (57-91%)	22% (9.3-43%)	6.4 (3.9-9.5)	15% (8.2-23%)	19% (10-29%)	500 (270-750)	20000 (11000-31000)
Serbia	2010	Male	55 to 59	76% (54-91%)	24% (9.3-46%)	6.6 (3.9-9.9)	15% (7.5-23%)	18% (9.3-27%)	810 (440-1300)	28000 (15000-44000)
Serbia	2010	Male	60 to 64	75% (51-91%)	25% (9.2-49%)	6.3 (3.5-10)	13% (6.1-21%)	16% (7.5-25%)	860 (410-1400)	26000 (12000-41000)
Serbia	2010	Male	65 to 69	74% (50-90%)	26% (10-50%)	5.7 (3.5-8.3)	13% (5.5-21%)	15% (6.6-24%)	830 (360-1400)	20000 (8700-34000)
Serbia	2010	Male	70 to 74	71% (49-87%)	29% (13-51%)	4.9 (2.8-8.2)	11% (3.9-20%)	13% (4.4-23%)	1100 (370-1900)	21000 (7100-37000)
Serbia	2010	Male	75 to 79	69% (46-86%)	31% (14-54%)	4.2 (2.2-7.4)	9.2% (2-18%)	10% (2.4-20%)	1100 (250-2100)	16000 (3600-31000)
Serbia	2010	Male	80 to 84	67% (42-85%)	33% (15-58%)	3.8 (2-6.1)	8.2% (1.4-16%)	9.2% (1.5-19%)	750 (130-1500)	8400 (1500-17000)
Serbia	2010	Male	85 to 89	65% (43-83%)	35% (17-57%)	3.4 (1.7-5.7)	6.9% (0.92-15%)	7.7% (0.88-17%)	380 (44-840)	3000 (400-6500)
Serbia	2010	Male	90 to 94	65% (44-85%)	35% (15-56%)	3.2 (1.5-5.6)	5.6% (-0.29-14%)	6.2% (-0.51-16%)	94 (-7.6-240)	510 (-27-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	2010	Male	95 plus	66% (40-88%)	34% (12-60%)	2.9 (1.1-5.6)	4.8% (-0.79-13%)	5.2% (-1-15%)	17 (-3.3-46)	61 (-9.8-160)
Serbia	2016	Female	All Ages	58% (50-67%)	42% (33-50%)	2.6 (1.9-3.4)	5.6% (1.5-11%)	7.1% (1.1-16%)	3700 (550-8100)	78000 (21000-150000)
Serbia	2016	Female	15 to 19	58% (39-76%)	42% (24-61%)	1.8 (0.72-3.4)	3% (1.1-5.4%)	7.7% (0.8-16%)	5.5 (0.59-12)	720 (260-1300)
Serbia	2016	Female	20 to 24	68% (50-82%)	32% (18-50%)	2.6 (1-4.9)	4.9% (2.2-8.4%)	9.7% (1.1-20%)	9.1 (1.1-19)	1300 (620-2200)
Serbia	2016	Female	25 to 29	70% (50-83%)	30% (17-50%)	2.7 (1.2-5)	5% (2.5-8.1%)	9.8% (1.8-20%)	12 (1.9-24)	1600 (770-2600)
Serbia	2016	Female	30 to 34	70% (50-84%)	30% (16-50%)	2.9 (1.3-5.1)	4.9% (2.4-8.2%)	10% (2.3-20%)	17 (3.7-33)	1800 (870-3000)
Serbia	2016	Female	35 to 39	71% (54-84%)	29% (16-46%)	3.2 (1.5-5.8)	5.1% (2.2-9.3%)	11% (3.1-21%)	29 (7.4-57)	2400 (990-4200)
Serbia	2016	Female	40 to 44	70% (53-84%)	30% (16-47%)	3.6 (1.7-6.1)	5.9% (2.2-11%)	12% (3.7-22%)	54 (17-100)	3500 (1300-6200)
Serbia	2016	Female	45 to 49	68% (46-83%)	32% (17-54%)	3.9 (1.9-7.2)	6.5% (2.6-12%)	12% (4.7-22%)	87 (34-160)	4600 (1800-8200)
Serbia	2016	Female	50 to 54	65% (45-81%)	35% (19-55%)	4 (2-7)	7.1% (2.7-13%)	12% (4.8-21%)	130 (53-240)	5800 (2200-11000)
Serbia	2016	Female	55 to 59	62% (43-80%)	38% (20-57%)	3.9 (1.9-6.6)	7.3% (2.4-13%)	11% (4.2-21%)	190 (70-340)	7100 (2400-13000)
Serbia	2016	Female	60 to 64	59% (38-78%)	41% (22-62%)	3.6 (1.7-6.6)	6.9% (1.7-13%)	10% (2.9-20%)	310 (88-590)	9800 (2500-19000)
Serbia	2016	Female	65 to 69	55% (35-74%)	45% (26-65%)	3.2 (1.6-5.2)	6.8% (1.3-14%)	9.5% (2.1-20%)	440 (96-910)	11000 (2100-23000)
Serbia	2016	Female	70 to 74	48% (30-67%)	52% (33-70%)	2.6 (1.1-4.7)	6% (0.1-14%)	7.9% (0.28-19%)	420 (15-960)	8400 (160-19000)
Serbia	2016	Female	75 to 79	41% (26-60%)	59% (40-74%)	2.1 (0.79-4)	5.1% (-0.26-13%)	6.4% (-0.26-17%)	510 (-21-1300)	7800 (-370-20000)
Serbia	2016	Female	80 to 84	36% (21-55%)	64% (45-79%)	1.9 (0.75-3.4)	4.8% (0.0076-12%)	5.7% (-0.069-15%)	580 (-7.2-1500)	6600 (12-17000)
Serbia	2016	Female	85 to 89	35% (22-51%)	65% (49-78%)	1.8 (0.75-3.6)	4.6% (-0.044-12%)	5.5% (-0.034-15%)	510 (-3.1-1400)	4000 (-38-11000)
Serbia	2016	Female	90 to 94	37% (23-53%)	63% (47-77%)	2 (0.75-4.3)	4.6% (-0.21-14%)	5.5% (-0.3-17%)	280 (-15-850)	1500 (-67-4400)
Serbia	2016	Female	95 plus	40% (23-61%)	60% (39-77%)	2.2 (0.6-5.2)	4.8% (-0.34-14%)	5.7% (-0.44-17%)	82 (-6.3-250)	280 (-20-860)
Serbia	2016	Male	All Ages	69% (59-79%)	31% (21-41%)	4.3 (3.1-5.7)	11% (5.5-18%)	13% (4.9-22%)	6800 (2700-12000)	180000 (88000-290000)
Serbia	2016	Male	15 to 19	59% (37-80%)	41% (20-63%)	1.8 (0.81-3.3)	5.4% (1.7-10%)	9.1% (0.97-19%)	17 (1.9-37)	1700 (550-3200)
Serbia	2016	Male	20 to 24	70% (49-87%)	30% (13-51%)	3.2 (1.5-5.5)	11% (4.3-20%)	16% (3.4-32%)	48 (10-95)	4400 (1700-7700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Serbia	2016	Male	25 to 29	73% (54-89%)	27% (11-46%)	4.1 (2.1-6.9)	12% (5.6-21%)	19% (6.1-36%)	69 (21-130)	5700 (2600-9500)
Serbia	2016	Male	30 to 34	76% (55-91%)	24% (9.3-45%)	4.7 (2.4-7.7)	13% (6.1-21%)	21% (7.5-36%)	84 (30-150)	6300 (2900-10000)
Serbia	2016	Male	35 to 39	78% (59-91%)	22% (9.4-41%)	4.9 (2.6-8.2)	13% (6-21%)	21% (8-35%)	120 (46-200)	7800 (3600-12000)
Serbia	2016	Male	40 to 44	79% (60-92%)	21% (8.3-40%)	5.6 (3.2-9)	14% (6.8-22%)	22% (9.8-35%)	190 (85-300)	11000 (5000-16000)
Serbia	2016	Male	45 to 49	79% (58-91%)	21% (8.7-42%)	6.4 (3.6-11)	15% (8.4-23%)	21% (11-33%)	300 (160-460)	14000 (7800-22000)
Serbia	2016	Male	50 to 54	78% (57-91%)	22% (9.3-43%)	6.9 (4.1-11)	15% (8.5-23%)	20% (11-30%)	430 (240-650)	18000 (9700-27000)
Serbia	2016	Male	55 to 59	76% (55-91%)	24% (9.3-45%)	7 (4.1-11)	15% (7.7-23%)	19% (9.7-28%)	590 (310-900)	21000 (11000-32000)
Serbia	2016	Male	60 to 64	75% (51-90%)	25% (9.5-49%)	6.8 (3.7-11)	13% (6-21%)	16% (7.5-26%)	860 (410-1400)	26000 (12000-42000)
Serbia	2016	Male	65 to 69	73% (49-90%)	27% (10-51%)	6.1 (3.6-9.3)	12% (5.8-20%)	15% (6.9-24%)	1100 (490-1700)	26000 (12000-43000)
Serbia	2016	Male	70 to 74	71% (49-87%)	29% (13-51%)	5.2 (2.9-8.4)	11% (4-19%)	13% (4.9-23%)	880 (330-1500)	17000 (6400-30000)
Serbia	2016	Male	75 to 79	69% (46-85%)	31% (15-54%)	4.4 (2.2-8)	9.1% (1.9-18%)	10% (2.1-20%)	830 (170-1600)	13000 (2700-24000)
Serbia	2016	Male	80 to 84	67% (43-85%)	33% (15-57%)	3.9 (1.9-6.7)	7.9% (1-16%)	8.9% (1-19%)	730 (86-1600)	8300 (1000-17000)
Serbia	2016	Male	85 to 89	65% (43-84%)	35% (16-57%)	3.6 (1.8-6.2)	6.7% (0.33-16%)	7.4% (0.055-18%)	450 (3.4-1100)	3700 (190-8400)
Serbia	2016	Male	90 to 94	65% (44-85%)	35% (15-56%)	3.3 (1.6-6.1)	5.4% (-0.64-15%)	6% (-1-16%)	160 (-28-430)	870 (-100-2300)
Serbia	2016	Male	95 plus	65% (40-88%)	35% (12-60%)	3 (1.1-6.1)	4.9% (-0.69-13%)	5.4% (-0.96-15%)	30 (-5.5-87)	110 (-15-310)
Slovakia	1990	Female	All Ages	87% (84-89%)	13% (11-16%)	1.9 (1.6-2.2)	2.6% (0.82-4.6%)	1.9% (-1.7-5.7%)	460 (-410-1400)	20000 (6400-35000)
Slovakia	1990	Female	15 to 19	88% (81-93%)	12% (6.9-19%)	2.3 (1.6-3.1)	3.8% (2.4-5.5%)	9.2% (4.9-14%)	7 (3.6-11)	780 (490-1100)
Slovakia	1990	Female	20 to 24	92% (87-95%)	8.3% (5.1-13%)	2.9 (2-4)	5.3% (3.7-7.1%)	10% (6.1-15%)	7.1 (4.1-11)	990 (670-1400)
Slovakia	1990	Female	25 to 29	93% (89-96%)	6.9% (3.9-11%)	2.4 (1.7-3.3)	4.8% (3.4-6.3%)	8.6% (5.2-12%)	8.1 (4.9-11)	1100 (740-1500)
Slovakia	1990	Female	30 to 34	93% (89-96%)	6.6% (3.9-11%)	2.3 (1.7-3.1)	4.6% (3.3-6.1%)	8.9% (6-12%)	13 (8.6-17)	1300 (940-1800)
Slovakia	1990	Female	35 to 39	92% (88-96%)	7.6% (4.5-12%)	2.7 (1.9-3.6)	5.4% (3.8-7.2%)	11% (7.3-14%)	27 (19-36)	1900 (1400-2600)
Slovakia	1990	Female	40 to 44	92% (87-95%)	8.1% (4.9-13%)	2.9 (2.1-3.8)	6.1% (4.4-8.3%)	11% (8.1-15%)	39 (27-52)	2300 (1600-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	1990	Female	45 to 49	92% (87-95%)	8% (4.7-13%)	3 (2.1-4.1)	6.6% (4.5-9.4%)	11% (7.7-16%)	52 (36-73)	2500 (1800-3500)
Slovakia	1990	Female	50 to 54	91% (87-95%)	8.6% (5.2-13%)	2.8 (2-3.9)	6.1% (3.7-9.3%)	9.6% (5.8-14%)	64 (38-93)	2600 (1600-3900)
Slovakia	1990	Female	55 to 59	90% (85-94%)	9.8% (6-15%)	2.5 (1.8-3.4)	4.3% (1.8-7.5%)	6.3% (2.6-11%)	68 (28-120)	2400 (970-4100)
Slovakia	1990	Female	60 to 64	89% (82-93%)	11% (6.8-18%)	2.4 (1.6-3.3)	2.4% (-0.6-6%)	3.2% (-1-8.2%)	54 (-17-140)	1600 (-400-4100)
Slovakia	1990	Female	65 to 69	86% (79-92%)	14% (7.8-21%)	2.4 (1.7-3.2)	1.4% (-2.1-4.9%)	1.7% (-2.9-6.3%)	47 (-81-170)	1200 (-1800-4000)
Slovakia	1990	Female	70 to 74	81% (73-88%)	19% (12-27%)	2.1 (1.4-3)	0.88% (-3-4.7%)	0.96% (-4-5.9%)	21 (-88-130)	480 (-1600-2600)
Slovakia	1990	Female	75 to 79	73% (61-83%)	27% (17-39%)	1.8 (1.2-2.8)	0.52% (-2.9-4.3%)	0.45% (-3.8-5.1%)	19 (-160-220)	400 (-2300-3300)
Slovakia	1990	Female	80 to 84	69% (54-80%)	31% (20-46%)	1.8 (1.2-2.6)	0.46% (-2.9-4.3%)	0.34% (-3.7-5%)	17 (-180-240)	280 (-1800-2700)
Slovakia	1990	Female	85 to 89	68% (54-79%)	32% (21-46%)	2 (1.2-2.9)	0.45% (-3.5-4.9%)	0.29% (-4.5-5.7%)	9.5 (-150-190)	130 (-1000-1400)
Slovakia	1990	Female	90 to 94	70% (56-80%)	30% (20-44%)	2.2 (1.3-3.3)	0.67% (-4.1-6.7%)	0.51% (-5.2-7.6%)	6.9 (-70-100)	55 (-340-540)
Slovakia	1990	Female	95 plus	72% (55-85%)	28% (15-45%)	2.3 (1.2-3.7)	1.1% (-4.5-8.3%)	0.96% (-5.6-9.5%)	2.5 (-15-25)	12 (-46-87)
Slovakia	1990	Male	All Ages	84% (80-88%)	16% (12-20%)	5.3 (4.6-6.2)	17% (14-20%)	18% (14-23%)	5500 (4200-7100)	170000 (140000-200000)
Slovakia	1990	Male	15 to 19	78% (66-87%)	22% (13-34%)	3 (1.9-4.1)	8.4% (5.8-12%)	14% (8.7-21%)	25 (15-37)	2300 (1600-3200)
Slovakia	1990	Male	20 to 24	89% (82-94%)	11% (6-18%)	5.5 (4.1-7.2)	17% (13-22%)	25% (18-33%)	60 (41-80)	5100 (3700-6600)
Slovakia	1990	Male	25 to 29	92% (87-96%)	7.6% (4.3-13%)	7.2 (5.2-9.3)	20% (15-24%)	30% (22-38%)	88 (66-110)	6800 (5200-8300)
Slovakia	1990	Male	30 to 34	94% (89-97%)	6% (2.9-11%)	8 (6-10)	21% (16-25%)	31% (24-37%)	140 (110-170)	9500 (7600-11000)
Slovakia	1990	Male	35 to 39	94% (90-97%)	5.8% (3-10%)	7.9 (5.8-10)	21% (17-26%)	30% (23-36%)	220 (170-260)	13000 (10000-15000)
Slovakia	1990	Male	40 to 44	94% (90-97%)	6.1% (3.4-10%)	8.2 (6.1-10)	23% (18-27%)	30% (24-35%)	300 (240-360)	15000 (12000-18000)
Slovakia	1990	Male	45 to 49	93% (87-96%)	7% (3.6-13%)	8.7 (6.4-12)	23% (19-28%)	29% (24-34%)	370 (300-450)	16000 (13000-20000)
Slovakia	1990	Male	50 to 54	92% (86-96%)	8% (4.3-14%)	9.3 (6.8-12)	22% (18-27%)	26% (21-32%)	460 (360-560)	18000 (14000-21000)
Slovakia	1990	Male	55 to 59	91% (84-95%)	8.9% (4.5-16%)	9.8 (7.3-13)	21% (16-26%)	24% (18-30%)	600 (450-760)	20000 (15000-25000)
Slovakia	1990	Male	60 to 64	90% (82-95%)	10% (5.1-18%)	9.2 (6.7-12)	19% (13-24%)	21% (15-28%)	740 (520-990)	21000 (15000-27000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	1990	Male	65 to 69	88% (80-95%)	12% (5.4-20%)	8 (6.1-10)	17% (12-22%)	19% (14-25%)	820 (570-1100)	19000 (14000-26000)
Slovakia	1990	Male	70 to 74	87% (78-92%)	13% (7.6-22%)	6.9 (4.9-9.2)	15% (9.2-20%)	17% (10-23%)	460 (290-640)	8800 (5600-12000)
Slovakia	1990	Male	75 to 79	85% (74-92%)	15% (8.2-26%)	5.9 (3.9-8.5)	12% (5.2-19%)	14% (5.5-21%)	580 (230-900)	8500 (3600-13000)
Slovakia	1990	Male	80 to 84	83% (72-91%)	17% (8.7-28%)	5.3 (3.6-7.4)	11% (4.2-17%)	12% (4.2-19%)	430 (150-690)	4700 (1800-7500)
Slovakia	1990	Male	85 to 89	83% (72-90%)	17% (10-28%)	4.9 (3.4-6.7)	8.8% (2.1-15%)	9.7% (1.7-17%)	180 (31-310)	1400 (330-2400)
Slovakia	1990	Male	90 to 94	83% (73-90%)	17% (9.6-27%)	4.5 (2.9-6.4)	6.6% (-0.4-14%)	7.1% (-1.2-16%)	41 (-7-90)	220 (-14-470)
Slovakia	1990	Male	95 plus	83% (71-92%)	17% (8.1-29%)	4 (2.2-6.3)	4.6% (-2.7-13%)	4.7% (-3.9-14%)	4.6 (-3.8-14)	18 (-11-50)
Slovakia	1995	Female	All Ages	89% (86-91%)	11% (9.3-14%)	1.9 (1.6-2.2)	2.2% (0.25-4.1%)	1.1% (-2.4-4.6%)	280 (-570-1100)	16000 (1800-30000)
Slovakia	1995	Female	15 to 19	89% (83-94%)	11% (6.2-17%)	2.2 (1.5-3)	3.4% (2.2-4.8%)	8.8% (4.6-13%)	6.3 (3.3-9.6)	740 (460-1100)
Slovakia	1995	Female	20 to 24	93% (89-95%)	7.3% (4.6-11%)	2.7 (1.9-3.7)	4.8% (3.3-6.4%)	9.7% (5.6-14%)	5.9 (3.4-8.6)	960 (650-1400)
Slovakia	1995	Female	25 to 29	94% (91-96%)	5.8% (3.5-9%)	2.3 (1.6-3.1)	4.4% (3.3-5.8%)	8.3% (5.1-12%)	5.7 (3.5-8.2)	880 (610-1200)
Slovakia	1995	Female	30 to 34	94% (91-96%)	5.7% (3.5-8.7%)	2.3 (1.6-3.1)	4.3% (3.1-5.7%)	8.7% (6-12%)	10 (7.2-14)	1100 (800-1500)
Slovakia	1995	Female	35 to 39	93% (90-96%)	6.7% (4-10%)	2.6 (1.8-3.5)	4.8% (3.5-6.3%)	10% (7.1-13%)	21 (15-28)	1600 (1100-2100)
Slovakia	1995	Female	40 to 44	93% (89-95%)	7.1% (4.6-11%)	2.8 (2-3.7)	5.3% (3.8-7.1%)	11% (7.7-14%)	37 (26-49)	2200 (1600-2900)
Slovakia	1995	Female	45 to 49	93% (89-96%)	7% (4.3-11%)	2.9 (2.1-3.8)	5.8% (3.9-8%)	11% (7.3-15%)	49 (34-68)	2400 (1700-3300)
Slovakia	1995	Female	50 to 54	92% (89-95%)	7.5% (4.9-11%)	2.7 (1.9-3.7)	5.5% (3.2-8.1%)	8.9% (5.2-13%)	57 (33-84)	2400 (1400-3400)
Slovakia	1995	Female	55 to 59	91% (87-95%)	8.6% (5.4-13%)	2.4 (1.8-3.2)	3.8% (1.4-6.6%)	5.8% (2.1-10%)	53 (19-90)	1800 (700-3200)
Slovakia	1995	Female	60 to 64	90% (85-94%)	9.9% (6.2-15%)	2.3 (1.6-3.2)	1.9% (-0.96-5.2%)	2.7% (-1.4-7.3%)	41 (-20-110)	1200 (-590-3300)
Slovakia	1995	Female	65 to 69	88% (82-93%)	12% (7.2-18%)	2.3 (1.7-3)	0.88% (-2.5-4.1%)	1.1% (-3.5-5.5%)	27 (-83-130)	690 (-2000-3200)
Slovakia	1995	Female	70 to 74	83% (76-88%)	17% (12-24%)	2.1 (1.4-2.8)	0.29% (-3.4-3.9%)	0.25% (-4.4-4.9%)	9.2 (-170-180)	270 (-3000-3600)
Slovakia	1995	Female	75 to 79	74% (63-83%)	26% (17-37%)	1.7 (1.1-2.5)	0.025% (-3.2-3.4%)	-0.15% (-4.1-4%)	-4.1 (-120-110)	14 (-1700-1800)
Slovakia	1995	Female	80 to 84	68% (53-80%)	32% (20-47%)	1.6 (1.1-2.4)	-0.093% (-3.2-3.1%)	-0.3% (-4.2-3.7%)	-14 (-200-180)	-55 (-1900-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	1995	Female	85 to 89	67% (53-78%)	33% (22-47%)	1.8 (1.1-2.7)	-0.23% (-3.8-3.9%)	-0.52% (-5-4.5%)	-20 (-190-170)	-78 (-1300-1300)
Slovakia	1995	Female	90 to 94	69% (55-80%)	31% (20-45%)	2 (1.2-3)	-0.11% (-4.5-5.1%)	-0.42% (-5.5-5.7%)	-6.8 (-89-94)	-11 (-430-500)
Slovakia	1995	Female	95 plus	72% (54-85%)	28% (15-46%)	2.1 (1.1-3.4)	0.19% (-5-6.3%)	-0.11% (-6.3-7.2%)	-0.39 (-23-27)	2.8 (-75-97)
Slovakia	1995	Male	All Ages	87% (83-90%)	13% (10-17%)	5.2 (4.5-5.9)	16% (13-19%)	17% (13-22%)	4900 (3700-6200)	150000 (130000-180000)
Slovakia	1995	Male	15 to 19	81% (72-88%)	19% (12-28%)	2.9 (2-3.9)	7.9% (5.4-11%)	13% (8.2-19%)	24 (15-34)	2300 (1500-3200)
Slovakia	1995	Male	20 to 24	91% (85-95%)	9.1% (5.5-15%)	5.3 (3.9-6.8)	16% (12-20%)	24% (16-31%)	57 (39-76)	5000 (3700-6400)
Slovakia	1995	Male	25 to 29	93% (89-96%)	6.7% (4-11%)	6.8 (4.9-8.8)	18% (14-23%)	29% (21-37%)	70 (50-90)	5600 (4200-6900)
Slovakia	1995	Male	30 to 34	95% (91-97%)	5.4% (2.8-9.4%)	7.5 (5.7-9.6)	19% (15-23%)	30% (23-36%)	100 (78-120)	7100 (5700-8500)
Slovakia	1995	Male	35 to 39	95% (91-97%)	5.1% (2.8-8.6%)	7.4 (5.5-9.6)	20% (16-24%)	29% (22-35%)	170 (130-200)	10000 (8100-12000)
Slovakia	1995	Male	40 to 44	95% (91-97%)	5.3% (3-8.7%)	7.7 (5.7-9.8)	21% (17-26%)	29% (23-34%)	300 (230-350)	15000 (12000-18000)
Slovakia	1995	Male	45 to 49	94% (90-97%)	5.9% (3.3-10%)	8.3 (6.1-11)	22% (18-26%)	28% (23-33%)	370 (300-440)	16000 (13000-19000)
Slovakia	1995	Male	50 to 54	93% (89-96%)	7% (4-11%)	8.8 (6.4-11)	22% (18-26%)	26% (22-32%)	420 (340-510)	16000 (13000-20000)
Slovakia	1995	Male	55 to 59	92% (86-96%)	8.1% (4.2-14%)	9.2 (6.9-12)	20% (16-25%)	24% (19-30%)	490 (380-610)	16000 (13000-21000)
Slovakia	1995	Male	60 to 64	91% (84-95%)	9.1% (4.8-16%)	8.7 (6.5-11)	18% (14-23%)	21% (16-27%)	630 (460-810)	18000 (13000-23000)
Slovakia	1995	Male	65 to 69	90% (83-95%)	10% (5.2-17%)	7.6 (5.8-9.7)	16% (12-21%)	19% (13-24%)	690 (490-910)	16000 (12000-21000)
Slovakia	1995	Male	70 to 74	88% (80-93%)	12% (7.3-20%)	6.5 (4.6-8.6)	14% (8.2-19%)	16% (9-22%)	690 (400-970)	13000 (7800-18000)
Slovakia	1995	Male	75 to 79	85% (75-92%)	15% (8.5-25%)	5.5 (3.7-7.7)	11% (4.1-17%)	12% (4.3-20%)	320 (110-510)	4800 (1700-7500)
Slovakia	1995	Male	80 to 84	83% (71-91%)	17% (9-29%)	4.9 (3.4-6.7)	9.2% (3-16%)	10% (3-18%)	340 (100-590)	3800 (1300-6300)
Slovakia	1995	Male	85 to 89	82% (71-90%)	18% (10-29%)	4.4 (3.1-6.1)	7% (0.92-14%)	7.6% (0.41-16%)	160 (8.4-320)	1200 (160-2500)
Slovakia	1995	Male	90 to 94	83% (72-90%)	17% (9.9-28%)	4.1 (2.7-5.8)	4.9% (-1.3-12%)	5.1% (-2.3-14%)	34 (-16-92)	200 (-54-480)
Slovakia	1995	Male	95 plus	83% (70-92%)	17% (8.1-30%)	3.6 (2-5.9)	3.1% (-3.5-11%)	2.9% (-4.9-13%)	3.2 (-5.3-14)	14 (-15-51)
Slovakia	2000	Female	All Ages	89% (87-91%)	11% (9.4-13%)	1.9 (1.7-2.1)	2.1% (0.41-3.7%)	0.92% (-2.2-4%)	220 (-550-980)	15000 (3100-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	2000	Female	15 to 19	89% (83-94%)	11% (6.3-17%)	2.1 (1.5-2.9)	3.3% (2.1-4.7%)	8.6% (4.5-13%)	5.7 (3.1-8.7)	680 (430-970)
Slovakia	2000	Female	20 to 24	93% (88-95%)	7.2% (4.6-12%)	2.6 (1.9-3.6)	4.6% (3.2-6.2%)	9.3% (5.4-13%)	6.1 (3.5-8.9)	1000 (680-1500)
Slovakia	2000	Female	25 to 29	94% (91-96%)	5.9% (3.6-9%)	2.2 (1.6-2.9)	4.3% (3.1-5.8%)	7.9% (4.8-11%)	6.2 (3.8-9)	980 (680-1300)
Slovakia	2000	Female	30 to 34	94% (91-96%)	5.8% (3.5-9.1%)	2.1 (1.6-2.9)	3.9% (2.9-5.1%)	8.3% (5.7-11%)	7 (4.8-9.5)	890 (640-1200)
Slovakia	2000	Female	35 to 39	93% (90-96%)	6.8% (4.3-10%)	2.5 (1.8-3.3)	4.5% (3.2-5.8%)	9.8% (6.8-13%)	17 (12-22)	1300 (950-1700)
Slovakia	2000	Female	40 to 44	93% (89-95%)	7.2% (4.7-11%)	2.7 (2-3.5)	5.1% (3.6-6.7%)	11% (7.7-14%)	33 (24-42)	1900 (1400-2500)
Slovakia	2000	Female	45 to 49	93% (89-95%)	7.2% (4.5-11%)	2.7 (1.9-3.6)	5.6% (4-7.6%)	10% (7.6-14%)	56 (41-73)	2700 (2000-3600)
Slovakia	2000	Female	50 to 54	92% (89-95%)	7.8% (5.1-11%)	2.6 (1.8-3.4)	5.2% (3.3-7.4%)	9% (5.7-12%)	62 (39-85)	2600 (1600-3500)
Slovakia	2000	Female	55 to 59	91% (87-94%)	9% (5.7-13%)	2.3 (1.7-3.1)	3.8% (1.7-6.2%)	6.2% (2.8-10%)	55 (25-87)	1900 (850-3000)
Slovakia	2000	Female	60 to 64	90% (85-94%)	10% (6.3-15%)	2.2 (1.6-3)	2% (-0.56-4.9%)	3% (-0.66-7.2%)	38 (-8.3-93)	1100 (-300-2800)
Slovakia	2000	Female	65 to 69	88% (82-93%)	12% (7.2-18%)	2.2 (1.6-2.9)	0.92% (-2-3.8%)	1.2% (-2.8-5.3%)	26 (-58-110)	640 (-1400-2800)
Slovakia	2000	Female	70 to 74	82% (75-88%)	18% (12-25%)	1.9 (1.3-2.7)	0.15% (-3-3.3%)	0.084% (-4.1-4.2%)	2.7 (-130-140)	130 (-2500-2700)
Slovakia	2000	Female	75 to 79	72% (61-82%)	28% (18-39%)	1.6 (1-2.2)	-0.15% (-3.1-2.8%)	-0.35% (-4-3.3%)	-17 (-190-160)	-140 (-2700-2500)
Slovakia	2000	Female	80 to 84	65% (50-78%)	35% (22-50%)	1.5 (0.97-2.1)	-0.32% (-3.1-2.4%)	-0.58% (-4.1-2.7%)	-19 (-140-88)	-140 (-1400-1000)
Slovakia	2000	Female	85 to 89	64% (50-77%)	36% (23-50%)	1.6 (0.98-2.4)	-0.53% (-3.8-3%)	-0.88% (-4.8-3.3%)	-34 (-190-130)	-190 (-1300-1000)
Slovakia	2000	Female	90 to 94	67% (53-79%)	33% (21-47%)	1.8 (1-2.7)	-0.47% (-4.4-4.1%)	-0.84% (-5.4-4.5%)	-17 (-110-89)	-57 (-530-490)
Slovakia	2000	Female	95 plus	70% (52-84%)	30% (16-48%)	1.9 (1-3.1)	-0.23% (-5-5.7%)	-0.6% (-6.2-6.3%)	-3 (-31-31)	-4.5 (-100-110)
Slovakia	2000	Male	All Ages	87% (84-90%)	13% (9.6-16%)	5.3 (4.7-5.8)	16% (13-19%)	17% (13-21%)	4800 (3700-6000)	150000 (130000-180000)
Slovakia	2000	Male	15 to 19	80% (71-88%)	20% (12-29%)	2.7 (1.9-3.8)	7.2% (5-9.8%)	13% (7.7-19%)	17 (11-25)	1800 (1200-2400)
Slovakia	2000	Male	20 to 24	91% (85-95%)	9.2% (5.4-15%)	5.1 (3.8-6.6)	15% (11-19%)	23% (16-30%)	58 (41-78)	5200 (3900-6700)
Slovakia	2000	Male	25 to 29	93% (89-96%)	6.6% (3.8-11%)	6.5 (4.8-8.4)	17% (13-22%)	28% (20-35%)	74 (54-95)	6000 (4600-7400)
Slovakia	2000	Male	30 to 34	95% (91-97%)	5.3% (2.8-9.2%)	7.2 (5.5-9.1)	19% (15-23%)	29% (23-36%)	90 (70-110)	6400 (5000-7600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	2000	Male	35 to 39	95% (91-97%)	5.1% (2.8-8.6%)	7.2 (5.4-9.3)	19% (15-23%)	28% (22-34%)	140 (110-160)	8400 (6700-10000)
Slovakia	2000	Male	40 to 44	95% (91-97%)	5.3% (3.1-9%)	7.4 (5.6-9.4)	21% (17-25%)	29% (23-34%)	260 (210-310)	14000 (11000-16000)
Slovakia	2000	Male	45 to 49	94% (89-97%)	6% (3.2-11%)	7.9 (6-10)	22% (18-26%)	28% (23-33%)	420 (340-500)	19000 (16000-22000)
Slovakia	2000	Male	50 to 54	93% (89-96%)	6.8% (3.9-11%)	8.5 (6.3-11)	22% (18-25%)	27% (22-32%)	510 (410-600)	20000 (16000-23000)
Slovakia	2000	Male	55 to 59	92% (87-96%)	7.7% (4-13%)	8.9 (6.9-11)	20% (16-25%)	24% (19-29%)	510 (400-620)	17000 (14000-21000)
Slovakia	2000	Male	60 to 64	91% (85-95%)	9.2% (4.7-15%)	8.4 (6.3-11)	18% (14-23%)	21% (16-27%)	570 (430-710)	16000 (12000-20000)
Slovakia	2000	Male	65 to 69	89% (82-94%)	11% (5.5-18%)	7.3 (5.6-9.1)	16% (12-20%)	18% (13-24%)	630 (450-820)	15000 (11000-19000)
Slovakia	2000	Male	70 to 74	87% (81-92%)	13% (7.5-19%)	6.3 (4.6-8.3)	13% (7.7-19%)	15% (8.6-21%)	620 (350-870)	12000 (6900-16000)
Slovakia	2000	Male	75 to 79	86% (77-92%)	14% (7.8-23%)	5.5 (3.8-7.6)	11% (4.1-17%)	12% (4.4-19%)	500 (180-800)	7500 (2900-12000)
Slovakia	2000	Male	80 to 84	84% (73-92%)	16% (8-27%)	4.9 (3.5-6.5)	8.7% (2.7-14%)	9.7% (2.7-17%)	220 (61-380)	2400 (760-4100)
Slovakia	2000	Male	85 to 89	84% (73-91%)	16% (9.5-27%)	4.4 (3.1-5.9)	6.5% (0.36-13%)	7% (-0.32-14%)	150 (-6.9-310)	1200 (68-2400)
Slovakia	2000	Male	90 to 94	84% (74-91%)	16% (9.1-26%)	4 (2.7-5.7)	4.4% (-1.7-11%)	4.5% (-2.8-13%)	37 (-23-110)	220 (-85-570)
Slovakia	2000	Male	95 plus	84% (71-93%)	16% (7.4-29%)	3.6 (2-5.7)	2.5% (-3.6-11%)	2.5% (-5-12%)	3.6 (-7.5-18)	16 (-22-67)
Slovakia	2005	Female	All Ages	88% (85-90%)	12% (10-15%)	1.8 (1.6-2)	1.9% (0.48-3.3%)	0.61% (-2.3-3.3%)	150 (-550-840)	14000 (3500-25000)
Slovakia	2005	Female	15 to 19	88% (82-93%)	12% (7-18%)	1.9 (1.4-2.6)	2.9% (1.9-4.1%)	7.7% (4.1-12%)	4 (2.1-6.1)	530 (350-750)
Slovakia	2005	Female	20 to 24	92% (88-95%)	7.8% (4.9-12%)	2.4 (1.7-3.3)	4.5% (3.1-5.9%)	8.4% (5.1-12%)	4.8 (2.9-7)	910 (610-1300)
Slovakia	2005	Female	25 to 29	94% (90-96%)	6.5% (4-10%)	2 (1.5-2.7)	4.1% (3-5.3%)	7.1% (4.4-9.9%)	5.2 (3.2-7.3)	970 (670-1300)
Slovakia	2005	Female	30 to 34	94% (90-96%)	6.4% (3.8-10%)	2 (1.4-2.6)	3.7% (2.8-4.8%)	7.5% (5.2-9.9%)	7.1 (4.9-9.6)	960 (690-1300)
Slovakia	2005	Female	35 to 39	93% (89-96%)	7.4% (4.4-11%)	2.3 (1.6-3.1)	4.2% (3-5.4%)	9.2% (6.3-12%)	13 (9.2-18)	1100 (810-1500)
Slovakia	2005	Female	40 to 44	92% (88-95%)	7.8% (4.8-12%)	2.5 (1.8-3.2)	4.8% (3.4-6.2%)	10% (7.3-13%)	28 (20-36)	1700 (1200-2200)
Slovakia	2005	Female	45 to 49	92% (88-95%)	7.7% (4.7-12%)	2.5 (1.8-3.2)	5.3% (3.8-7.1%)	10% (7.3-13%)	51 (37-67)	2500 (1800-3300)
Slovakia	2005	Female	50 to 54	91% (87-95%)	8.5% (5.5-13%)	2.3 (1.7-3.1)	4.9% (3.1-6.8%)	8.6% (5.7-11%)	68 (45-91)	2800 (1800-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	2005	Female	55 to 59	90% (84-94%)	10% (6.2-16%)	2 (1.5-2.7)	3.6% (1.8-5.6%)	6% (3.1-9%)	61 (32-92)	2100 (1100-3200)
Slovakia	2005	Female	60 to 64	88% (82-93%)	12% (7.4-18%)	1.9 (1.4-2.7)	2% (-0.089-4.3%)	3.2% (0.12-6.6%)	40 (1.5-82)	1100 (-52-2400)
Slovakia	2005	Female	65 to 69	85% (78-91%)	15% (8.8-22%)	1.9 (1.4-2.5)	0.88% (-1.5-3.2%)	1.3% (-2.2-4.6%)	23 (-40-83)	560 (-950-2000)
Slovakia	2005	Female	70 to 74	78% (70-85%)	22% (15-30%)	1.6 (1.1-2.3)	0.068% (-2.6-2.6%)	0.016% (-3.5-3.4%)	0.49 (-100-100)	53 (-1900-2000)
Slovakia	2005	Female	75 to 79	69% (56-80%)	31% (20-44%)	1.4 (0.89-2)	-0.29% (-2.7-2.2%)	-0.5% (-3.6-2.7%)	-22 (-160-120)	-240 (-2200-1800)
Slovakia	2005	Female	80 to 84	63% (49-76%)	37% (24-51%)	1.3 (0.88-1.9)	-0.59% (-3.1-1.8%)	-0.88% (-4-2%)	-51 (-230-120)	-440 (-2400-1300)
Slovakia	2005	Female	85 to 89	65% (51-76%)	35% (24-49%)	1.5 (0.94-2.2)	-0.94% (-4-2.2%)	-1.3% (-5.1-2.4%)	-39 (-150-70)	-240 (-1100-560)
Slovakia	2005	Female	90 to 94	67% (54-78%)	33% (22-46%)	1.7 (0.99-2.5)	-0.99% (-4.7-3.2%)	-1.4% (-5.8-3.6%)	-31 (-130-77)	-130 (-610-420)
Slovakia	2005	Female	95 plus	70% (52-84%)	30% (16-48%)	1.8 (0.94-2.9)	-0.86% (-5.2-4.6%)	-1.3% (-6.5-5.3%)	-8.4 (-42-33)	-22 (-140-120)
Slovakia	2005	Male	All Ages	89% (86-91%)	11% (8.8-14%)	5.3 (4.8-5.8)	16% (13-18%)	16% (12-21%)	4600 (3400-5800)	150000 (120000-170000)
Slovakia	2005	Male	15 to 19	81% (71-88%)	19% (12-29%)	2.6 (1.8-3.6)	6.9% (4.8-9.5%)	12% (7.5-18%)	14 (8.6-21)	1500 (1100-2100)
Slovakia	2005	Male	20 to 24	91% (86-95%)	8.8% (5.3-14%)	4.9 (3.7-6.3)	14% (11-18%)	22% (15-29%)	46 (33-61)	4400 (3300-5600)
Slovakia	2005	Male	25 to 29	94% (90-96%)	6.2% (3.7-9.7%)	6.3 (4.8-7.9)	16% (13-20%)	26% (19-33%)	64 (47-81)	5500 (4300-6800)
Slovakia	2005	Male	30 to 34	95% (92-97%)	5% (2.6-8.3%)	6.9 (5.3-8.7)	17% (14-20%)	28% (22-34%)	83 (65-100)	6200 (4900-7500)
Slovakia	2005	Male	35 to 39	95% (92-97%)	4.8% (2.6-7.9%)	6.9 (5.1-9.1)	18% (14-22%)	27% (21-33%)	110 (88-140)	7200 (5600-8600)
Slovakia	2005	Male	40 to 44	95% (92-97%)	5% (2.9-8.2%)	7.1 (5.5-9.1)	19% (15-23%)	28% (22-33%)	210 (160-250)	11000 (8700-13000)
Slovakia	2005	Male	45 to 49	94% (90-97%)	5.7% (3.1-9.6%)	7.6 (5.8-9.6)	21% (17-25%)	28% (23-33%)	380 (310-450)	17000 (14000-20000)
Slovakia	2005	Male	50 to 54	94% (89-96%)	6.4% (3.7-11%)	8.1 (6.2-10)	21% (17-25%)	27% (22-31%)	570 (470-670)	22000 (18000-26000)
Slovakia	2005	Male	55 to 59	93% (87-96%)	7.4% (4-13%)	8.6 (6.8-11)	20% (16-24%)	25% (20-29%)	600 (490-730)	20000 (17000-24000)
Slovakia	2005	Male	60 to 64	91% (85-95%)	8.9% (4.6-15%)	8 (6.1-10)	18% (14-22%)	22% (17-26%)	570 (430-700)	16000 (13000-20000)
Slovakia	2005	Male	65 to 69	90% (83-95%)	10% (5.2-17%)	7 (5.4-8.6)	16% (11-20%)	18% (13-23%)	540 (390-690)	13000 (9500-16000)
Slovakia	2005	Male	70 to 74	88% (81-93%)	12% (7-19%)	6.1 (4.5-7.9)	13% (7.1-18%)	15% (8-21%)	530 (290-770)	10000 (5900-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	2005	Male	75 to 79	86% (78-93%)	14% (7.3-22%)	5.2 (3.6-7.2)	9.7% (3.2-16%)	11% (3.3-18%)	440 (130-730)	6600 (2200-11000)
Slovakia	2005	Male	80 to 84	85% (75-92%)	15% (8.1-25%)	4.6 (3.4-6.2)	7.7% (1.9-14%)	8.5% (1.7-16%)	320 (61-580)	3500 (880-6300)
Slovakia	2005	Male	85 to 89	84% (75-90%)	16% (9.6-25%)	4.2 (3-5.6)	5.4% (-0.14-12%)	5.8% (-0.83-13%)	88 (-13-200)	720 (-18-1500)
Slovakia	2005	Male	90 to 94	84% (75-91%)	16% (9-25%)	3.9 (2.6-5.4)	3.3% (-2.4-10%)	3.3% (-3.6-12%)	31 (-34-110)	190 (-140-590)
Slovakia	2005	Male	95 plus	84% (72-93%)	16% (7.2-28%)	3.4 (1.9-5.4)	1.8% (-4.3-9.7%)	1.4% (-5.7-11%)	2.8 (-12-23)	15 (-35-83)
Slovakia	2010	Female	All Ages	87% (85-89%)	13% (11-15%)	1.7 (1.5-1.9)	1.8% (0.58-3.1%)	0.41% (-2.3-3%)	100 (-570-780)	13000 (4600-23000)
Slovakia	2010	Female	15 to 19	87% (80-92%)	13% (8-20%)	1.7 (1.2-2.4)	2.7% (1.8-3.8%)	7.1% (3.8-11%)	2.6 (1.4-4)	410 (260-580)
Slovakia	2010	Female	20 to 24	91% (87-95%)	8.5% (5.4-13%)	2.3 (1.6-3.1)	4.3% (3-5.8%)	7.7% (4.6-11%)	3.8 (2.2-5.5)	810 (540-1200)
Slovakia	2010	Female	25 to 29	93% (89-96%)	7% (4.2-11%)	1.9 (1.4-2.5)	3.9% (2.9-5.2%)	6.7% (4.3-9.1%)	4 (2.5-5.4)	890 (610-1200)
Slovakia	2010	Female	30 to 34	93% (90-96%)	6.7% (3.9-10%)	1.9 (1.3-2.5)	3.7% (2.7-4.9%)	7.4% (5.3-9.8%)	6.7 (4.7-8.9)	1000 (710-1400)
Slovakia	2010	Female	35 to 39	93% (89-95%)	7.4% (4.5-11%)	2.2 (1.5-2.9)	4.1% (3-5.2%)	9% (6.4-12%)	14 (9.7-18)	1200 (870-1600)
Slovakia	2010	Female	40 to 44	92% (88-95%)	8.1% (5.3-12%)	2.3 (1.7-3)	4.6% (3.4-5.9%)	10% (7.5-13%)	23 (17-30)	1500 (1100-1900)
Slovakia	2010	Female	45 to 49	92% (88-95%)	7.7% (4.9-12%)	2.4 (1.7-3.2)	4.9% (3.5-6.5%)	10% (7.5-13%)	40 (30-52)	2000 (1500-2600)
Slovakia	2010	Female	50 to 54	92% (88-94%)	8.4% (5.5-12%)	2.2 (1.6-3)	4.7% (3.1-6.5%)	8.5% (5.8-11%)	63 (43-83)	2600 (1700-3500)
Slovakia	2010	Female	55 to 59	90% (85-94%)	10% (6.4-15%)	2 (1.4-2.6)	3.5% (1.9-5.2%)	6.1% (3.5-8.9%)	67 (38-97)	2300 (1300-3400)
Slovakia	2010	Female	60 to 64	87% (81-92%)	13% (8-19%)	1.8 (1.3-2.5)	2.1% (0.35-4.2%)	3.5% (0.79-6.5%)	49 (11-89)	1400 (240-2700)
Slovakia	2010	Female	65 to 69	83% (75-90%)	17% (10-25%)	1.7 (1.2-2.3)	1% (-0.87-2.9%)	1.7% (-1.3-4.4%)	28 (-22-73)	660 (-560-1800)
Slovakia	2010	Female	70 to 74	75% (67-83%)	25% (17-33%)	1.4 (0.97-2)	0.21% (-1.8-2.2%)	0.25% (-2.6-3%)	6.3 (-65-75)	140 (-1200-1500)
Slovakia	2010	Female	75 to 79	67% (56-78%)	33% (22-44%)	1.3 (0.82-1.8)	-0.28% (-2.3-1.9%)	-0.48% (-3.2-2.3%)	-19 (-130-89)	-210 (-1800-1400)
Slovakia	2010	Female	80 to 84	65% (51-77%)	35% (23-49%)	1.3 (0.9-1.9)	-0.74% (-3.2-1.6%)	-1.1% (-4.2-1.9%)	-59 (-230-100)	-540 (-2300-1200)
Slovakia	2010	Female	85 to 89	66% (52-78%)	34% (22-48%)	1.5 (0.95-2.2)	-1.1% (-4.2-1.9%)	-1.6% (-5.4-2.1%)	-82 (-270-110)	-530 (-1900-890)
Slovakia	2010	Female	90 to 94	68% (55-79%)	32% (21-45%)	1.7 (0.98-2.5)	-1.3% (-4.8-2.8%)	-1.8% (-6.1-3%)	-31 (-100-51)	-130 (-490-290)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	2010	Female	95 plus	70% (53-84%)	30% (16-47%)	1.8 (0.94-2.9)	-1.3% (-5.6-3.9%)	-1.8% (-7-4.3%)	-13 (-50-30)	-36 (-160-110)
Slovakia	2010	Male	All Ages	89% (86-91%)	11% (9-14%)	5.2 (4.6-5.8)	15% (12-17%)	16% (12-20%)	4300 (3200-5500)	140000 (110000-160000)
Slovakia	2010	Male	15 to 19	80% (71-88%)	20% (12-29%)	2.5 (1.7-3.4)	6.5% (4.6-8.8%)	12% (7-17%)	11 (6.3-15)	1200 (820-1600)
Slovakia	2010	Male	20 to 24	91% (86-95%)	9.1% (5.3-14%)	4.6 (3.4-6)	14% (10-18%)	21% (14-28%)	38 (26-51)	3700 (2800-4800)
Slovakia	2010	Male	25 to 29	94% (90-96%)	6.5% (3.7-10%)	5.9 (4.4-7.6)	15% (12-19%)	25% (18-32%)	54 (40-69)	4800 (3700-6000)
Slovakia	2010	Male	30 to 34	95% (91-97%)	5.4% (2.8-9.2%)	6.5 (5-8.1)	15% (13-19%)	26% (21-32%)	75 (58-91)	5800 (4600-7100)
Slovakia	2010	Male	35 to 39	95% (92-97%)	5.2% (2.8-8.4%)	6.5 (4.8-8.4)	17% (13-20%)	26% (20-32%)	110 (82-130)	7000 (5400-8400)
Slovakia	2010	Male	40 to 44	95% (92-97%)	5.2% (3.1-8.4%)	6.8 (5.1-8.6)	18% (14-22%)	27% (21-32%)	160 (130-190)	8700 (6900-10000)
Slovakia	2010	Male	45 to 49	94% (90-97%)	5.9% (3.3-9.9%)	7.2 (5.4-9.2)	20% (16-24%)	27% (22-32%)	290 (230-350)	14000 (11000-16000)
Slovakia	2010	Male	50 to 54	93% (89-96%)	6.7% (3.9-11%)	7.7 (5.8-9.8)	21% (17-24%)	27% (22-31%)	500 (400-580)	20000 (16000-23000)
Slovakia	2010	Male	55 to 59	92% (86-96%)	8.1% (4.3-14%)	8 (6.3-10)	20% (16-23%)	25% (20-29%)	660 (530-800)	23000 (18000-27000)
Slovakia	2010	Male	60 to 64	90% (84-95%)	9.5% (5.1-16%)	7.5 (5.7-9.8)	17% (14-22%)	21% (16-26%)	650 (500-800)	19000 (15000-23000)
Slovakia	2010	Male	65 to 69	90% (83-94%)	10% (5.6-17%)	6.6 (5.2-8.5)	15% (11-19%)	18% (13-23%)	530 (390-680)	13000 (9500-16000)
Slovakia	2010	Male	70 to 74	88% (82-93%)	12% (7.3-18%)	5.8 (4.2-7.7)	12% (6.3-17%)	14% (7.2-20%)	440 (230-630)	8600 (4700-12000)
Slovakia	2010	Male	75 to 79	86% (78-92%)	14% (8.1-22%)	4.9 (3.4-6.9)	8.8% (2.9-15%)	10% (3-17%)	370 (110-630)	5600 (1900-9500)
Slovakia	2010	Male	80 to 84	83% (72-91%)	17% (9.3-28%)	4.3 (3-5.9)	6.7% (1.2-12%)	7.3% (0.8-14%)	260 (28-500)	2900 (520-5400)
Slovakia	2010	Male	85 to 89	82% (69-89%)	18% (11-31%)	3.8 (2.6-5.1)	4.5% (-0.39-10%)	4.7% (-1.1-11%)	110 (-28-280)	970 (-84-2200)
Slovakia	2010	Male	90 to 94	82% (72-90%)	18% (10-28%)	3.5 (2.3-5)	2.5% (-2.6-8.9%)	2.3% (-3.7-9.9%)	16 (-26-68)	110 (-110-380)
Slovakia	2010	Male	95 plus	82% (69-92%)	18% (8.3-31%)	3.1 (1.7-5)	1.1% (-4.1-8.5%)	0.57% (-5.8-9.4%)	1.2 (-12-19)	9.4 (-35-71)
Slovakia	2016	Female	All Ages	87% (85-90%)	13% (10-15%)	1.7 (1.4-2.1)	1.8% (0.44-3.3%)	0.53% (-2.4-3.6%)	140 (-630-940)	14000 (3200-25000)
Slovakia	2016	Female	15 to 19	88% (81-93%)	12% (7.1-19%)	1.8 (1.2-2.6)	2.7% (1.8-3.8%)	7.4% (3.8-11%)	2.1 (1.1-3.4)	320 (200-470)
Slovakia	2016	Female	20 to 24	92% (87-95%)	8.3% (5.1-13%)	2.3 (1.5-3.3)	4.3% (3-5.9%)	8% (4.6-12%)	3.1 (1.7-4.6)	690 (460-990)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	2016	Female	25 to 29	93% (89-96%)	6.9% (4-11%)	1.9 (1.3-2.7)	4% (3-5.3%)	6.9% (4.4-9.7%)	3.3 (2-4.9)	810 (550-1200)
Slovakia	2016	Female	30 to 34	93% (89-96%)	6.7% (3.9-11%)	1.9 (1.3-2.6)	3.9% (2.8-5.1%)	7.8% (5.5-10%)	6.3 (4.1-8.7)	970 (690-1400)
Slovakia	2016	Female	35 to 39	92% (88-95%)	7.7% (4.5-12%)	2.2 (1.4-3.1)	4.1% (3-5.4%)	9.4% (6.5-12%)	15 (9.8-21)	1400 (970-1800)
Slovakia	2016	Female	40 to 44	92% (87-95%)	8.2% (5-13%)	2.4 (1.6-3.3)	4.4% (3.2-5.9%)	10% (7.5-14%)	23 (16-32)	1500 (1100-2100)
Slovakia	2016	Female	45 to 49	92% (86-95%)	8.1% (4.8-14%)	2.4 (1.6-3.4)	4.8% (3.4-6.5%)	10% (7.5-13%)	34 (24-47)	1800 (1200-2400)
Slovakia	2016	Female	50 to 54	91% (86-95%)	8.8% (5.3-14%)	2.2 (1.5-3.1)	4.6% (3.1-6.4%)	8.9% (6.2-12%)	51 (33-71)	2200 (1400-3100)
Slovakia	2016	Female	55 to 59	90% (84-94%)	10% (6.2-16%)	2 (1.4-2.7)	3.7% (2.1-5.5%)	6.5% (4-9.4%)	64 (38-96)	2200 (1300-3400)
Slovakia	2016	Female	60 to 64	88% (82-93%)	12% (7-18%)	1.9 (1.3-2.7)	2.3% (0.5-4.3%)	4% (1.2-6.9%)	63 (19-110)	1800 (390-3400)
Slovakia	2016	Female	65 to 69	86% (77-92%)	14% (7.8-23%)	1.9 (1.3-2.6)	1.3% (-0.85-3.2%)	2.1% (-1-5.1%)	40 (-20-96)	920 (-630-2500)
Slovakia	2016	Female	70 to 74	80% (71-87%)	20% (13-29%)	1.7 (1.1-2.5)	0.41% (-1.8-2.9%)	0.59% (-2.7-4%)	15 (-64-99)	290 (-1300-2000)
Slovakia	2016	Female	75 to 79	72% (59-83%)	28% (17-41%)	1.4 (0.89-2.2)	-0.12% (-2.4-2.4%)	-0.3% (-3.4-2.9%)	-10 (-120-100)	-85 (-1700-1600)
Slovakia	2016	Female	80 to 84	67% (52-79%)	33% (21-48%)	1.4 (0.9-2.1)	-0.51% (-3.1-2.1%)	-0.82% (-4.1-2.6%)	-43 (-210-140)	-360 (-2200-1500)
Slovakia	2016	Female	85 to 89	66% (52-78%)	34% (22-48%)	1.5 (0.92-2.3)	-0.82% (-3.9-2.4%)	-1.2% (-5-2.7%)	-62 (-250-130)	-380 (-1800-1100)
Slovakia	2016	Female	90 to 94	68% (55-80%)	32% (20-45%)	1.7 (0.95-2.6)	-0.99% (-4.4-3.2%)	-1.5% (-5.8-3.5%)	-56 (-220-130)	-230 (-1000-730)
Slovakia	2016	Female	95 plus	71% (54-85%)	29% (15-46%)	1.8 (0.92-3)	-0.98% (-5.3-4.5%)	-1.5% (-6.8-5.1%)	-8.5 (-38-28)	-22 (-120-100)
Slovakia	2016	Male	All Ages	87% (84-90%)	13% (10-16%)	5.1 (4.1-6.3)	14% (11-17%)	15% (10-20%)	4100 (2700-5600)	130000 (100000-160000)
Slovakia	2016	Male	15 to 19	77% (66-87%)	23% (13-34%)	2.3 (1.4-3.4)	6.3% (4.3-8.6%)	12% (6.6-17%)	7.3 (4.2-11)	880 (590-1200)
Slovakia	2016	Male	20 to 24	89% (82-94%)	11% (6.2-18%)	4.4 (3.1-6)	13% (9.8-17%)	21% (13-28%)	27 (17-38)	2900 (2100-3800)
Slovakia	2016	Male	25 to 29	92% (86-96%)	8% (4.3-14%)	5.7 (3.9-7.6)	15% (11-18%)	25% (17-32%)	42 (29-57)	4000 (3000-5200)
Slovakia	2016	Male	30 to 34	94% (88-97%)	6.3% (3-12%)	6.4 (4.5-8.3)	15% (12-18%)	26% (20-33%)	61 (44-80)	5000 (3800-6400)
Slovakia	2016	Male	35 to 39	94% (89-97%)	6% (3.1-11%)	6.4 (4.5-8.7)	16% (12-20%)	27% (20-33%)	110 (76-140)	7200 (5400-9100)
Slovakia	2016	Male	40 to 44	94% (89-96%)	6.3% (3.6-11%)	6.5 (4.6-9)	17% (13-21%)	27% (21-33%)	160 (110-200)	8800 (6500-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovakia	2016	Male	45 to 49	93% (87-96%)	7.2% (3.6-13%)	7 (4.9-9.5)	19% (15-23%)	27% (21-33%)	230 (170-290)	11000 (8200-14000)
Slovakia	2016	Male	50 to 54	92% (85-96%)	8.3% (4.4-15%)	7.4 (5.2-10)	20% (15-24%)	26% (21-31%)	390 (290-490)	16000 (12000-19000)
Slovakia	2016	Male	55 to 59	91% (83-95%)	9.2% (4.6-17%)	7.8 (5.5-10)	19% (15-23%)	24% (20-29%)	580 (440-730)	20000 (16000-25000)
Slovakia	2016	Male	60 to 64	90% (82-95%)	10% (5.1-18%)	7.3 (5-10)	17% (13-21%)	21% (16-26%)	730 (550-950)	21000 (16000-27000)
Slovakia	2016	Male	65 to 69	88% (80-94%)	12% (5.8-20%)	6.4 (4.7-8.5)	14% (10-18%)	17% (12-22%)	600 (410-800)	15000 (10000-19000)
Slovakia	2016	Male	70 to 74	86% (78-92%)	14% (8.1-22%)	5.5 (3.7-7.6)	11% (5.6-16%)	13% (6.4-19%)	450 (210-670)	8900 (4400-13000)
Slovakia	2016	Male	75 to 79	84% (74-91%)	16% (8.9-26%)	4.7 (3-6.8)	8.5% (2.5-14%)	9.8% (2.6-17%)	320 (79-560)	4900 (1400-8400)
Slovakia	2016	Male	80 to 84	82% (71-91%)	18% (9.3-29%)	4.2 (2.7-6)	6.6% (1.1-13%)	7.4% (0.84-14%)	250 (28-510)	2900 (500-5600)
Slovakia	2016	Male	85 to 89	82% (71-90%)	18% (10-29%)	3.8 (2.5-5.5)	4.7% (-0.44-11%)	5% (-1.1-12%)	120 (-27-300)	1000 (-95-2300)
Slovakia	2016	Male	90 to 94	82% (71-90%)	18% (10-29%)	3.5 (2.2-5.3)	2.8% (-2.4-9.4%)	2.6% (-3.5-11%)	39 (-53-150)	250 (-210-840)
Slovakia	2016	Male	95 plus	82% (69-92%)	18% (8.2-31%)	3.2 (1.7-5.2)	1.4% (-3.8-8.9%)	0.9% (-5.4-10%)	1.6 (-9.3-18)	10 (-26-64)
Slovenia	1990	Female	All Ages	67% (62-71%)	33% (29-38%)	2.5 (2.1-2.9)	7.3% (5.6-9.2%)	9.8% (6.8-14%)	920 (630-1300)	21000 (16000-27000)
Slovenia	1990	Female	15 to 19	68% (53-80%)	32% (20-47%)	2.3 (1.5-3.3)	4.6% (2.9-6.7%)	13% (7-19%)	3.1 (1.7-4.9)	320 (190-460)
Slovenia	1990	Female	20 to 24	77% (64-86%)	23% (14-36%)	3 (1.9-4.2)	6.4% (4.4-8.9%)	15% (8.6-21%)	4.9 (2.9-7.3)	540 (370-740)
Slovenia	1990	Female	25 to 29	79% (68-87%)	21% (13-32%)	2.5 (1.7-3.5)	5.3% (3.8-6.9%)	12% (7.3-17%)	4.1 (2.5-5.9)	490 (330-650)
Slovenia	1990	Female	30 to 34	80% (68-88%)	20% (12-32%)	2.3 (1.6-3.2)	4.9% (3.6-6.5%)	12% (7.7-16%)	5.5 (3.5-7.8)	540 (370-710)
Slovenia	1990	Female	35 to 39	79% (67-87%)	21% (13-33%)	2.5 (1.6-3.7)	5.9% (4.2-7.9%)	14% (9.4-18%)	11 (7.3-15)	790 (560-1000)
Slovenia	1990	Female	40 to 44	76% (64-85%)	24% (15-36%)	3.1 (2.2-4.3)	7.4% (5.5-9.7%)	16% (12-20%)	17 (12-22)	980 (720-1200)
Slovenia	1990	Female	45 to 49	72% (61-81%)	28% (19-39%)	3.9 (2.7-5.3)	9.5% (7.1-12%)	18% (14-23%)	30 (23-38)	1500 (1100-1900)
Slovenia	1990	Female	50 to 54	68% (59-76%)	32% (24-41%)	4 (2.8-5.3)	10% (8-14%)	18% (14-23%)	45 (34-57)	1900 (1400-2400)
Slovenia	1990	Female	55 to 59	66% (56-74%)	34% (26-44%)	3.8 (2.6-5)	10% (7.4-13%)	16% (12-21%)	62 (47-81)	2200 (1700-2900)
Slovenia	1990	Female	60 to 64	65% (54-73%)	35% (27-46%)	3.6 (2.5-5)	9.1% (6.4-12%)	14% (9.8-18%)	79 (56-110)	2400 (1700-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	1990	Female	65 to 69	64% (54-72%)	36% (28-46%)	3.6 (2.6-4.7)	8.8% (6.1-12%)	13% (8.7-17%)	100 (71-140)	2600 (1800-3500)
Slovenia	1990	Female	70 to 74	61% (50-70%)	39% (30-50%)	3.4 (2.3-4.7)	8.6% (5.1-12%)	11% (6.8-16%)	81 (48-120)	1600 (960-2300)
Slovenia	1990	Female	75 to 79	56% (42-67%)	44% (33-58%)	2.9 (1.9-4.2)	7.4% (3.9-12%)	9.2% (4.8-15%)	160 (81-250)	2400 (1200-3800)
Slovenia	1990	Female	80 to 84	51% (37-64%)	49% (36-63%)	2.5 (1.6-3.7)	6.4% (3.5-11%)	7.8% (4.1-13%)	150 (80-250)	1700 (910-2700)
Slovenia	1990	Female	85 to 89	48% (35-61%)	52% (39-65%)	2.4 (1.5-3.6)	5.9% (2.7-10%)	6.9% (3.1-12%)	100 (45-180)	790 (360-1400)
Slovenia	1990	Female	90 to 94	48% (35-61%)	52% (39-65%)	2.4 (1.4-3.7)	6% (2.3-11%)	7.1% (2.6-13%)	50 (18-94)	260 (100-490)
Slovenia	1990	Female	95 plus	50% (33-66%)	50% (34-67%)	2.5 (1.3-4.2)	6.2% (1.6-13%)	7.3% (1.7-15%)	13 (3.1-26)	44 (12-89)
Slovenia	1990	Male	All Ages	72% (66-78%)	28% (22-34%)	5.2 (4.5-6)	17% (14-20%)	20% (16-24%)	2000 (1500-2400)	59000 (49000-70000)
Slovenia	1990	Male	15 to 19	62% (46-77%)	38% (23-54%)	3 (2-4.3)	9.4% (5.9-14%)	16% (9.3-24%)	11 (6.2-16)	900 (560-1300)
Slovenia	1990	Male	20 to 24	77% (63-87%)	23% (13-37%)	5.4 (3.8-7.1)	19% (13-25%)	29% (19-39%)	34 (22-46)	2500 (1800-3400)
Slovenia	1990	Male	25 to 29	82% (70-90%)	18% (10-30%)	5.5 (4.1-7.3)	19% (14-24%)	30% (21-38%)	35 (24-46)	2500 (1900-3300)
Slovenia	1990	Male	30 to 34	85% (73-92%)	15% (8-27%)	5.7 (4.2-7.5)	18% (14-23%)	29% (22-37%)	43 (31-56)	2900 (2200-3700)
Slovenia	1990	Male	35 to 39	86% (76-92%)	14% (7.8-24%)	5.8 (4.3-7.5)	19% (14-23%)	29% (22-35%)	65 (48-81)	3900 (2900-4700)
Slovenia	1990	Male	40 to 44	85% (76-92%)	15% (8.3-24%)	6.4 (4.8-8.2)	20% (16-24%)	30% (24-35%)	81 (64-99)	4200 (3300-5100)
Slovenia	1990	Male	45 to 49	83% (74-90%)	17% (9.9-26%)	7.5 (5.5-9.7)	22% (19-26%)	30% (25-35%)	120 (100-150)	5500 (4600-6600)
Slovenia	1990	Male	50 to 54	80% (72-86%)	20% (14-28%)	8.4 (6.4-11)	22% (18-25%)	28% (24-33%)	170 (140-210)	6700 (5600-8100)
Slovenia	1990	Male	55 to 59	77% (69-83%)	23% (17-31%)	9.4 (7.2-12)	21% (17-25%)	25% (21-30%)	250 (200-300)	8300 (6700-10000)
Slovenia	1990	Male	60 to 64	77% (69-82%)	23% (18-31%)	10 (7.6-13)	18% (14-22%)	22% (17-27%)	240 (180-300)	6900 (5300-8500)
Slovenia	1990	Male	65 to 69	79% (72-85%)	21% (15-28%)	10 (7.8-13)	17% (13-21%)	20% (15-25%)	220 (160-280)	5200 (3800-6700)
Slovenia	1990	Male	70 to 74	77% (69-83%)	23% (17-31%)	9.4 (7.1-12)	16% (11-20%)	19% (13-24%)	140 (94-180)	2700 (1900-3400)
Slovenia	1990	Male	75 to 79	71% (61-79%)	29% (21-39%)	8 (5.8-10)	14% (9.6-19%)	16% (11-21%)	250 (160-330)	3700 (2500-4800)
Slovenia	1990	Male	80 to 84	68% (56-77%)	32% (23-44%)	6.9 (5-9)	13% (8.5-17%)	14% (9.5-20%)	190 (120-270)	2100 (1400-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	1990	Male	85 to 89	69% (57-77%)	31% (23-43%)	6.3 (4.5-8.7)	12% (7.5-17%)	13% (8.4-19%)	90 (57-130)	710 (470-1000)
Slovenia	1990	Male	90 to 94	70% (59-79%)	30% (21-41%)	5.9 (3.9-8.6)	11% (7.3-16%)	13% (8-19%)	30 (19-43)	160 (100-230)
Slovenia	1990	Male	95 plus	71% (56-83%)	29% (17-44%)	5.3 (3-8.1)	11% (5.2-16%)	12% (5.5-18%)	5.4 (2.5-8.1)	19 (9.4-29)
Slovenia	1995	Female	All Ages	67% (62-71%)	33% (29-38%)	2.2 (1.9-2.6)	6.5% (5-8.2%)	8.4% (5.4-12%)	790 (520-1100)	18000 (14000-23000)
Slovenia	1995	Female	15 to 19	68% (53-81%)	32% (19-47%)	2.1 (1.3-2.9)	4.2% (2.6-6.1%)	11% (6-17%)	2.9 (1.6-4.5)	300 (180-440)
Slovenia	1995	Female	20 to 24	77% (64-86%)	23% (14-36%)	2.6 (1.7-3.7)	5.7% (3.9-7.9%)	13% (7.1-19%)	3.5 (2-5.3)	420 (280-580)
Slovenia	1995	Female	25 to 29	79% (68-87%)	21% (13-32%)	2.2 (1.5-3)	4.9% (3.5-6.4%)	11% (6.3-15%)	3.4 (2-4.9)	430 (290-570)
Slovenia	1995	Female	30 to 34	80% (67-88%)	20% (12-33%)	2 (1.3-2.9)	4.6% (3.3-6%)	10% (6.7-15%)	4.6 (2.9-6.6)	470 (330-630)
Slovenia	1995	Female	35 to 39	79% (66-88%)	21% (12-34%)	2.2 (1.4-3.2)	5.4% (3.9-7.2%)	12% (8.6-16%)	9.5 (6.6-13)	680 (500-900)
Slovenia	1995	Female	40 to 44	76% (65-85%)	24% (15-35%)	2.7 (1.9-3.7)	6.7% (4.9-8.6%)	14% (11-18%)	18 (14-24)	1100 (800-1400)
Slovenia	1995	Female	45 to 49	72% (62-81%)	28% (19-38%)	3.4 (2.3-4.6)	8.7% (6.4-11%)	16% (13-21%)	28 (21-36)	1400 (1000-1700)
Slovenia	1995	Female	50 to 54	68% (59-76%)	32% (24-41%)	3.5 (2.5-4.7)	9.4% (7.1-12%)	16% (13-20%)	40 (31-51)	1700 (1300-2100)
Slovenia	1995	Female	55 to 59	66% (56-74%)	34% (26-44%)	3.3 (2.3-4.4)	8.7% (6.5-11%)	15% (11-18%)	47 (36-61)	1700 (1300-2200)
Slovenia	1995	Female	60 to 64	65% (55-74%)	35% (26-45%)	3.2 (2.2-4.4)	8% (5.6-11%)	12% (8.9-16%)	69 (48-92)	2100 (1500-2800)
Slovenia	1995	Female	65 to 69	65% (55-73%)	35% (27-45%)	3.2 (2.3-4.1)	7.5% (5.2-10%)	11% (7.6-14%)	90 (63-120)	2300 (1600-3100)
Slovenia	1995	Female	70 to 74	62% (51-70%)	38% (30-49%)	3 (2-4.1)	7.2% (4.1-11%)	9.7% (5.6-14%)	120 (69-180)	2400 (1400-3600)
Slovenia	1995	Female	75 to 79	56% (43-68%)	44% (32-57%)	2.6 (1.7-3.7)	6.1% (2.9-9.9%)	7.7% (3.7-13%)	70 (33-120)	1100 (510-1800)
Slovenia	1995	Female	80 to 84	52% (37-64%)	48% (36-63%)	2.2 (1.4-3.2)	5.3% (2.6-8.8%)	6.4% (3-11%)	100 (50-170)	1200 (570-1900)
Slovenia	1995	Female	85 to 89	49% (36-62%)	51% (38-64%)	2.1 (1.3-3.1)	4.8% (2-8.6%)	5.7% (2.2-10%)	110 (43-200)	880 (360-1500)
Slovenia	1995	Female	90 to 94	49% (36-62%)	51% (38-64%)	2.2 (1.3-3.3)	4.8% (1.5-9.5%)	5.7% (1.6-11%)	53 (15-100)	280 (84-540)
Slovenia	1995	Female	95 plus	50% (35-66%)	50% (34-65%)	2.2 (1.1-3.6)	5% (0.66-11%)	5.9% (0.6-13%)	15 (1.6-34)	54 (7.2-120)
Slovenia	1995	Male	All Ages	73% (67-78%)	27% (22-33%)	4.8 (4.2-5.4)	16% (14-19%)	19% (16-23%)	1900 (1500-2300)	56000 (47000-66000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	1995	Male	15 to 19	63% (47-78%)	37% (22-53%)	2.7 (1.8-3.8)	8.8% (5.5-13%)	15% (8.5-22%)	11 (6.2-17)	930 (560-1300)
Slovenia	1995	Male	20 to 24	77% (63-87%)	23% (13-37%)	4.8 (3.4-6.3)	17% (12-23%)	27% (18-36%)	28 (18-37)	2100 (1500-2800)
Slovenia	1995	Male	25 to 29	82% (71-90%)	18% (10-29%)	4.9 (3.6-6.3)	17% (12-22%)	27% (19-35%)	28 (19-38)	2100 (1500-2800)
Slovenia	1995	Male	30 to 34	85% (75-92%)	15% (7.9-25%)	5 (3.7-6.5)	17% (13-21%)	27% (20-34%)	35 (26-46)	2400 (1800-3100)
Slovenia	1995	Male	35 to 39	86% (76-92%)	14% (7.8-24%)	5.1 (3.7-6.6)	17% (13-21%)	26% (20-33%)	50 (37-63)	3000 (2300-3800)
Slovenia	1995	Male	40 to 44	85% (77-91%)	15% (8.6-23%)	5.6 (4.3-7.2)	19% (15-23%)	27% (22-32%)	90 (70-110)	4700 (3700-5700)
Slovenia	1995	Male	45 to 49	83% (74-90%)	17% (9.7-26%)	6.6 (4.9-8.4)	21% (17-24%)	28% (24-33%)	110 (91-140)	5100 (4200-6100)
Slovenia	1995	Male	50 to 54	81% (73-86%)	19% (14-27%)	7.4 (5.6-9.2)	21% (18-24%)	27% (23-31%)	150 (130-190)	6100 (5100-7300)
Slovenia	1995	Male	55 to 59	77% (70-83%)	23% (17-30%)	8.2 (6.4-10)	20% (17-24%)	25% (21-29%)	200 (160-240)	6800 (5600-8200)
Slovenia	1995	Male	60 to 64	77% (69-83%)	23% (17-31%)	8.7 (6.7-11)	18% (15-22%)	22% (18-26%)	260 (210-320)	7500 (6000-9200)
Slovenia	1995	Male	65 to 69	79% (72-85%)	21% (15-28%)	8.8 (6.9-11)	17% (13-21%)	20% (15-24%)	260 (200-330)	6300 (4800-7800)
Slovenia	1995	Male	70 to 74	77% (69-83%)	23% (17-31%)	8.2 (6.3-10)	15% (12-19%)	18% (13-22%)	230 (160-290)	4400 (3200-5600)
Slovenia	1995	Male	75 to 79	71% (61-79%)	29% (21-39%)	6.9 (5.1-8.9)	14% (10-18%)	16% (12-20%)	120 (85-150)	1800 (1300-2300)
Slovenia	1995	Male	80 to 84	69% (57-77%)	31% (23-43%)	6 (4.4-7.8)	12% (8.8-16%)	14% (10-19%)	150 (100-200)	1600 (1200-2200)
Slovenia	1995	Male	85 to 89	69% (58-77%)	31% (23-42%)	5.5 (3.9-7.4)	11% (7.8-16%)	13% (8.6-18%)	130 (84-180)	1000 (680-1400)
Slovenia	1995	Male	90 to 94	70% (59-79%)	30% (21-41%)	5.1 (3.4-7.2)	11% (6.6-15%)	12% (7.3-18%)	41 (25-59)	220 (130-310)
Slovenia	1995	Male	95 plus	71% (56-83%)	29% (17-44%)	4.7 (2.6-7.1)	9.7% (3.9-15%)	11% (4.1-17%)	7.8 (2.9-12)	28 (11-43)
Slovenia	2000	Female	All Ages	66% (61-70%)	34% (30-39%)	2 (1.8-2.3)	5.6% (4.4-7%)	7.1% (4.8-9.6%)	680 (450-940)	16000 (12000-20000)
Slovenia	2000	Female	15 to 19	67% (51-81%)	33% (19-49%)	1.8 (1.2-2.6)	3.6% (2.2-5.3%)	9.9% (5.1-15%)	2.1 (1.1-3.3)	230 (140-340)
Slovenia	2000	Female	20 to 24	76% (62-85%)	24% (15-38%)	2.3 (1.5-3.2)	4.9% (3.3-6.7%)	11% (5.9-17%)	2.6 (1.3-3.9)	360 (240-500)
Slovenia	2000	Female	25 to 29	77% (65-86%)	23% (14-35%)	1.9 (1.3-2.6)	4.3% (3.1-5.5%)	9.4% (5.5-14%)	2.2 (1.2-3.2)	330 (220-440)
Slovenia	2000	Female	30 to 34	78% (64-87%)	22% (13-36%)	1.7 (1.1-2.4)	4% (2.9-5.1%)	9.1% (5.9-13%)	3.3 (2.1-4.5)	380 (270-520)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	2000	Female	35 to 39	77% (65-87%)	23% (13-35%)	2 (1.3-2.9)	4.8% (3.5-6.2%)	12% (8.2-15%)	7 (5-9.4)	550 (410-710)
Slovenia	2000	Female	40 to 44	76% (64-85%)	24% (15-36%)	2.5 (1.8-3.5)	5.9% (4.4-7.6%)	13% (10-16%)	14 (11-18)	850 (650-1100)
Slovenia	2000	Female	45 to 49	72% (62-80%)	28% (20-38%)	3.1 (2.2-4.3)	7.7% (5.7-9.9%)	15% (12-19%)	29 (22-37)	1400 (1100-1800)
Slovenia	2000	Female	50 to 54	67% (59-76%)	33% (24-41%)	3.1 (2.2-4.1)	8.4% (6.4-11%)	15% (12-19%)	35 (27-44)	1500 (1100-1800)
Slovenia	2000	Female	55 to 59	64% (54-72%)	36% (28-46%)	2.9 (2.1-3.8)	7.7% (5.8-9.8%)	13% (10-17%)	40 (31-51)	1400 (1100-1800)
Slovenia	2000	Female	60 to 64	62% (52-71%)	38% (29-48%)	2.7 (1.9-3.9)	6.9% (4.8-9.4%)	11% (8.2-15%)	48 (35-65)	1500 (1100-2000)
Slovenia	2000	Female	65 to 69	62% (51-71%)	38% (29-49%)	2.7 (1.9-3.6)	6.4% (4.5-8.6%)	9.6% (6.9-13%)	70 (50-95)	1800 (1300-2500)
Slovenia	2000	Female	70 to 74	60% (50-69%)	40% (31-50%)	2.7 (1.9-3.7)	6% (3.7-8.8%)	8.3% (5.2-12%)	95 (58-140)	1900 (1200-2900)
Slovenia	2000	Female	75 to 79	56% (43-68%)	44% (32-57%)	2.3 (1.6-3.4)	5% (2.5-8.1%)	6.5% (3.3-11%)	110 (52-180)	1700 (830-2800)
Slovenia	2000	Female	80 to 84	51% (37-64%)	49% (36-63%)	2 (1.3-2.9)	4.4% (2.2-7%)	5.4% (2.6-8.9%)	86 (42-140)	970 (480-1600)
Slovenia	2000	Female	85 to 89	49% (36-61%)	51% (39-64%)	1.9 (1.2-2.8)	4% (1.6-6.9%)	4.7% (1.8-8.5%)	68 (25-120)	540 (210-940)
Slovenia	2000	Female	90 to 94	49% (37-61%)	51% (39-63%)	1.9 (1.2-3)	3.9% (1-7.7%)	4.6% (1.1-9.3%)	53 (13-110)	290 (79-570)
Slovenia	2000	Female	95 plus	50% (35-65%)	50% (35-65%)	2 (1.1-3.3)	4.1% (0.57-9.2%)	4.7% (0.5-11%)	16 (1.7-38)	57 (8-130)
Slovenia	2000	Male	All Ages	76% (71-80%)	24% (20-29%)	4.7 (4.1-5.2)	16% (13-18%)	19% (16-22%)	1900 (1500-2200)	53000 (45000-62000)
Slovenia	2000	Male	15 to 19	66% (51-80%)	34% (20-49%)	2.6 (1.8-3.6)	7.9% (4.9-11%)	14% (8-21%)	7.9 (4.3-12)	690 (420-980)
Slovenia	2000	Male	20 to 24	79% (66-87%)	21% (13-34%)	4.5 (3.3-5.9)	16% (11-21%)	25% (16-33%)	24 (15-32)	1900 (1300-2500)
Slovenia	2000	Male	25 to 29	83% (74-90%)	17% (9.7-26%)	4.6 (3.4-5.8)	15% (11-19%)	25% (18-33%)	21 (14-28)	1700 (1200-2200)
Slovenia	2000	Male	30 to 34	86% (77-93%)	14% (7.4-23%)	4.7 (3.5-6)	15% (11-19%)	25% (18-33%)	27 (19-35)	2000 (1400-2500)
Slovenia	2000	Male	35 to 39	87% (78-92%)	13% (7.8-22%)	4.7 (3.5-6.1)	15% (12-19%)	25% (19-31%)	40 (30-51)	2500 (1900-3100)
Slovenia	2000	Male	40 to 44	86% (78-92%)	14% (8.3-22%)	5.2 (4-6.6)	17% (14-21%)	26% (21-31%)	69 (52-84)	3700 (2900-4500)
Slovenia	2000	Male	45 to 49	84% (76-90%)	16% (9.8-24%)	6.1 (4.7-7.7)	19% (16-23%)	27% (23-32%)	130 (100-150)	5800 (4700-6900)
Slovenia	2000	Male	50 to 54	82% (75-87%)	18% (13-25%)	6.9 (5.4-8.5)	20% (17-23%)	27% (23-30%)	140 (120-160)	5500 (4600-6500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	2000	Male	55 to 59	79% (72-83%)	21% (17-28%)	7.7 (6.1-9.3)	19% (16-22%)	25% (21-28%)	180 (140-210)	6100 (5000-7300)
Slovenia	2000	Male	60 to 64	78% (72-83%)	22% (17-28%)	8.2 (6.4-10)	18% (15-21%)	22% (18-26%)	220 (170-260)	6400 (5100-7700)
Slovenia	2000	Male	65 to 69	81% (75-86%)	19% (14-25%)	8.3 (6.7-10)	17% (14-20%)	20% (17-24%)	270 (220-330)	6600 (5200-7900)
Slovenia	2000	Male	70 to 74	79% (73-84%)	21% (16-27%)	7.8 (6.2-9.4)	15% (12-18%)	18% (14-22%)	260 (200-320)	5100 (3900-6200)
Slovenia	2000	Male	75 to 79	73% (65-80%)	27% (20-35%)	6.6 (5.1-8.3)	13% (10-16%)	15% (12-19%)	200 (150-250)	3000 (2300-3700)
Slovenia	2000	Male	80 to 84	70% (59-78%)	30% (22-41%)	5.6 (4.2-7.2)	12% (8.9-15%)	14% (10-17%)	130 (97-170)	1500 (1100-1900)
Slovenia	2000	Male	85 to 89	70% (60-78%)	30% (22-40%)	5.1 (3.6-6.7)	11% (7.5-15%)	12% (8.4-17%)	85 (59-120)	670 (470-900)
Slovenia	2000	Male	90 to 94	71% (61-80%)	29% (20-39%)	4.8 (3.2-6.5)	9.8% (5.6-14%)	11% (6.2-16%)	50 (28-73)	270 (150-390)
Slovenia	2000	Male	95 plus	72% (59-83%)	28% (17-41%)	4.3 (2.4-6.5)	8.8% (3.2-14%)	9.9% (3.2-16%)	9 (2.9-14)	33 (12-51)
Slovenia	2005	Female	All Ages	68% (64-71%)	32% (29-36%)	2 (1.7-2.2)	4.7% (3.6-5.9%)	5.8% (3.8-8.2%)	540 (350-780)	13000 (9900-17000)
Slovenia	2005	Female	15 to 19	71% (58-83%)	29% (17-42%)	1.8 (1.2-2.6)	2.9% (1.9-4.2%)	9.2% (4.8-14%)	1.2 (0.61-1.9)	160 (100-230)
Slovenia	2005	Female	20 to 24	79% (67-87%)	21% (13-33%)	2.2 (1.5-3.1)	4.6% (3.2-6.1%)	10% (5.4-15%)	2.2 (1.2-3.4)	320 (210-450)
Slovenia	2005	Female	25 to 29	80% (68-87%)	20% (13-32%)	1.8 (1.3-2.5)	3.9% (2.9-5.1%)	8.4% (4.9-12%)	1.7 (0.94-2.5)	310 (210-420)
Slovenia	2005	Female	30 to 34	80% (70-88%)	20% (12-30%)	1.7 (1.2-2.4)	3.8% (2.8-4.9%)	8.5% (5.5-12%)	2.7 (1.7-3.8)	340 (240-450)
Slovenia	2005	Female	35 to 39	79% (69-88%)	21% (12-31%)	1.8 (1.2-2.6)	4.1% (3.1-5.2%)	10% (7.5-13%)	5.1 (3.6-6.8)	450 (330-580)
Slovenia	2005	Female	40 to 44	75% (65-84%)	25% (16-35%)	2.2 (1.6-3)	4.9% (3.7-6.4%)	12% (8.9-15%)	10 (7.6-13)	660 (500-840)
Slovenia	2005	Female	45 to 49	71% (63-80%)	29% (20-37%)	2.8 (2-3.7)	6.3% (4.8-8%)	13% (10-16%)	20 (16-26)	1000 (800-1300)
Slovenia	2005	Female	50 to 54	70% (63-76%)	30% (24-37%)	3 (2.2-3.9)	7.1% (5.4-9%)	13% (11-16%)	34 (26-42)	1500 (1100-1800)
Slovenia	2005	Female	55 to 59	66% (60-73%)	34% (27-40%)	2.7 (2-3.6)	6.7% (5-8.4%)	12% (9.3-15%)	33 (25-42)	1200 (930-1600)
Slovenia	2005	Female	60 to 64	64% (57-71%)	36% (29-43%)	2.6 (1.9-3.5)	5.8% (4.2-7.8%)	10% (7.5-13%)	38 (28-50)	1200 (860-1600)
Slovenia	2005	Female	65 to 69	63% (57-70%)	37% (30-43%)	2.5 (2-3.2)	5.3% (3.7-6.9%)	8.5% (6.2-11%)	47 (35-62)	1200 (890-1700)
Slovenia	2005	Female	70 to 74	61% (53-67%)	39% (33-47%)	2.4 (1.8-3.2)	4.8% (2.8-7.1%)	7% (4.4-10%)	70 (43-100)	1500 (850-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	2005	Female	75 to 79	59% (49-68%)	41% (32-51%)	2.3 (1.6-3)	4.2% (2-6.8%)	5.6% (2.7-8.9%)	85 (40-140)	1400 (630-2200)
Slovenia	2005	Female	80 to 84	54% (41-66%)	46% (34-59%)	1.9 (1.3-2.7)	3.6% (1.6-5.9%)	4.5% (1.9-7.6%)	90 (39-150)	1000 (480-1800)
Slovenia	2005	Female	85 to 89	49% (37-61%)	51% (39-63%)	1.8 (1.1-2.6)	3% (0.93-5.5%)	3.6% (1-6.7%)	56 (16-100)	460 (140-830)
Slovenia	2005	Female	90 to 94	50% (37-60%)	50% (40-63%)	1.8 (1.1-2.6)	3% (0.52-6.1%)	3.5% (0.45-7.4%)	31 (4.1-65)	170 (29-350)
Slovenia	2005	Female	95 plus	51% (36-66%)	49% (34-64%)	1.8 (0.96-2.9)	3.1% (-0.022-7.5%)	3.6% (-0.26-9.1%)	18 (-1.3-46)	64 (-0.44-160)
Slovenia	2005	Male	All Ages	78% (74-82%)	22% (18-26%)	4.6 (4-5.1)	14% (12-16%)	17% (15-20%)	1700 (1400-2000)	47000 (40000-56000)
Slovenia	2005	Male	15 to 19	71% (58-83%)	29% (17-42%)	2.6 (1.8-3.5)	7% (4.3-9.7%)	13% (7.6-20%)	5.3 (3-7.9)	500 (310-700)
Slovenia	2005	Male	20 to 24	81% (69-89%)	19% (11-31%)	4.3 (3.2-5.6)	14% (10-19%)	23% (15-31%)	18 (12-25)	1500 (1100-2000)
Slovenia	2005	Male	25 to 29	86% (78-91%)	14% (8.8-22%)	4.3 (3.3-5.5)	14% (9.9-18%)	23% (15-30%)	21 (14-29)	1700 (1200-2200)
Slovenia	2005	Male	30 to 34	88% (81-93%)	12% (6.8-19%)	4.4 (3.4-5.6)	13% (9.7-16%)	23% (17-30%)	19 (14-25)	1500 (1100-1900)
Slovenia	2005	Male	35 to 39	88% (81-93%)	12% (7.2-19%)	4.4 (3.3-5.6)	13% (10-17%)	23% (17-29%)	28 (20-36)	1900 (1400-2300)
Slovenia	2005	Male	40 to 44	87% (81-92%)	13% (7.8-19%)	4.9 (3.8-6.1)	15% (12-19%)	24% (19-29%)	53 (40-67)	2900 (2300-3600)
Slovenia	2005	Male	45 to 49	86% (80-91%)	14% (9.2-20%)	5.7 (4.5-7.1)	17% (14-20%)	26% (21-30%)	95 (75-120)	4500 (3600-5400)
Slovenia	2005	Male	50 to 54	83% (78-87%)	17% (13-22%)	6.4 (5.1-7.8)	18% (15-21%)	25% (21-28%)	150 (120-170)	6000 (4900-7100)
Slovenia	2005	Male	55 to 59	80% (75-83%)	20% (17-25%)	7.2 (5.8-8.6)	17% (15-20%)	23% (20-26%)	150 (120-180)	5300 (4400-6300)
Slovenia	2005	Male	60 to 64	80% (76-84%)	20% (16-24%)	7.7 (6.1-9.4)	17% (14-20%)	21% (18-25%)	180 (150-220)	5500 (4500-6600)
Slovenia	2005	Male	65 to 69	83% (78-86%)	17% (14-22%)	7.8 (6.4-9.3)	16% (13-18%)	19% (16-22%)	210 (170-260)	5300 (4400-6300)
Slovenia	2005	Male	70 to 74	81% (76-84%)	19% (16-24%)	7.3 (5.9-8.9)	14% (12-17%)	17% (14-20%)	250 (200-300)	4900 (4000-5900)
Slovenia	2005	Male	75 to 79	76% (69-81%)	24% (19-31%)	6.3 (4.8-7.8)	12% (9.7-15%)	14% (11-17%)	210 (160-260)	3300 (2600-4100)
Slovenia	2005	Male	80 to 84	73% (65-79%)	27% (21-35%)	5.4 (4.2-6.7)	11% (8.3-14%)	13% (9.5-16%)	160 (120-200)	1800 (1300-2300)
Slovenia	2005	Male	85 to 89	73% (65-79%)	27% (21-35%)	5 (3.7-6.4)	9.9% (6.8-13%)	11% (7.5-15%)	74 (48-100)	590 (400-810)
Slovenia	2005	Male	90 to 94	74% (65-81%)	26% (19-35%)	4.6 (3.2-6.3)	8.9% (4.6-13%)	10% (4.8-15%)	31 (15-46)	170 (85-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	2005	Male	95 plus	74% (60-85%)	26% (15-40%)	4.1 (2.3-6.2)	7.8% (2.3-13%)	8.7% (2.2-15%)	11 (2.6-18)	40 (12-66)
Slovenia	2010	Female	All Ages	67% (63-71%)	33% (29-37%)	1.8 (1.5-2.1)	3.9% (2.9-5%)	4.7% (2.8-7%)	450 (270-680)	11000 (7900-14000)
Slovenia	2010	Female	15 to 19	71% (57-82%)	29% (18-43%)	1.7 (1.1-2.4)	2.5% (1.7-3.5%)	8% (4.3-12%)	0.7 (0.37-1.1)	110 (71-160)
Slovenia	2010	Female	20 to 24	79% (67-87%)	21% (13-33%)	2.1 (1.4-2.9)	3.9% (2.8-5.2%)	8.6% (4.7-13%)	1 (0.54-1.5)	230 (150-320)
Slovenia	2010	Female	25 to 29	80% (69-87%)	20% (13-31%)	1.7 (1.2-2.4)	3.6% (2.7-4.7%)	7.4% (4.4-10%)	0.92 (0.54-1.3)	250 (170-360)
Slovenia	2010	Female	30 to 34	81% (71-88%)	19% (12-29%)	1.6 (1.1-2.3)	3.4% (2.5-4.4%)	7.8% (5.2-11%)	1.8 (1.2-2.5)	300 (220-420)
Slovenia	2010	Female	35 to 39	80% (70-88%)	20% (12-30%)	1.7 (1.2-2.6)	3.8% (2.8-4.9%)	10% (7.3-13%)	3.7 (2.7-4.9)	370 (270-480)
Slovenia	2010	Female	40 to 44	76% (66-85%)	24% (15-34%)	2.1 (1.5-2.9)	4.4% (3.3-5.6%)	11% (8.6-14%)	7.8 (5.9-10)	540 (410-700)
Slovenia	2010	Female	45 to 49	71% (63-80%)	29% (20-37%)	2.6 (1.8-3.5)	5.4% (4-6.9%)	12% (9.6-15%)	15 (12-19)	830 (630-1100)
Slovenia	2010	Female	50 to 54	69% (62-76%)	31% (24-38%)	2.7 (2-3.6)	5.9% (4.5-7.5%)	12% (9.4-14%)	24 (19-30)	1100 (860-1400)
Slovenia	2010	Female	55 to 59	65% (58-72%)	35% (28-42%)	2.5 (1.8-3.3)	5.7% (4.2-7.2%)	11% (8.5-13%)	33 (25-41)	1300 (960-1600)
Slovenia	2010	Female	60 to 64	63% (55-70%)	37% (30-45%)	2.4 (1.7-3.2)	5% (3.6-6.7%)	9.1% (7-12%)	30 (23-39)	980 (720-1300)
Slovenia	2010	Female	65 to 69	62% (55-70%)	38% (30-45%)	2.3 (1.8-3)	4.6% (3.3-6.1%)	7.8% (5.9-10%)	38 (28-50)	1000 (740-1400)
Slovenia	2010	Female	70 to 74	60% (52-67%)	40% (33-48%)	2.2 (1.6-2.9)	4.1% (2.5-6%)	6.3% (4.1-8.9%)	47 (29-67)	1000 (630-1500)
Slovenia	2010	Female	75 to 79	57% (47-66%)	43% (34-53%)	2 (1.4-2.8)	3.5% (1.6-5.6%)	4.8% (2.4-7.9%)	67 (32-110)	1100 (520-1800)
Slovenia	2010	Female	80 to 84	52% (40-64%)	48% (36-60%)	1.7 (1.2-2.4)	2.8% (1.2-5%)	3.7% (1.6-6.5%)	71 (30-130)	840 (360-1500)
Slovenia	2010	Female	85 to 89	48% (36-59%)	52% (41-64%)	1.6 (0.96-2.3)	2.4% (0.64-4.7%)	2.9% (0.63-5.9%)	67 (15-140)	560 (140-1100)
Slovenia	2010	Female	90 to 94	48% (36-59%)	52% (41-64%)	1.6 (0.94-2.4)	2.3% (0.11-5.2%)	2.7% (-0.053-6.3%)	33 (-0.65-78)	190 (9.3-420)
Slovenia	2010	Female	95 plus	50% (34-66%)	50% (34-66%)	1.6 (0.84-2.7)	2.4% (-0.44-6.3%)	2.8% (-0.75-7.6%)	12 (-3.4-34)	46 (-8.3-120)
Slovenia	2010	Male	All Ages	78% (74-82%)	22% (18-26%)	4.3 (3.7-5)	13% (11-15%)	16% (14-19%)	1500 (1200-1800)	42000 (36000-50000)
Slovenia	2010	Male	15 to 19	69% (55-81%)	31% (19-45%)	2.3 (1.6-3.2)	5.6% (3.7-7.8%)	12% (6.6-18%)	2.7 (1.4-4.1)	290 (190-420)
Slovenia	2010	Male	20 to 24	81% (70-89%)	19% (11-30%)	4 (2.9-5.3)	12% (8.5-16%)	21% (14-29%)	12 (7.4-16)	1100 (740-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	2010	Male	25 to 29	86% (78-91%)	14% (8.7-22%)	4 (3-5.2)	11% (8.3-14%)	21% (14-28%)	12 (7.7-17)	1100 (830-1500)
Slovenia	2010	Male	30 to 34	88% (81-93%)	12% (6.9-19%)	4.1 (3-5.3)	11% (8.5-14%)	21% (15-28%)	17 (12-22)	1400 (1000-1800)
Slovenia	2010	Male	35 to 39	88% (81-93%)	12% (7-19%)	4.1 (3.1-5.3)	11% (8.7-15%)	22% (16-28%)	20 (14-26)	1500 (1100-1900)
Slovenia	2010	Male	40 to 44	87% (80-93%)	13% (7.3-20%)	4.5 (3.4-5.7)	14% (10-17%)	24% (18-29%)	41 (31-52)	2400 (1800-2900)
Slovenia	2010	Male	45 to 49	86% (80-91%)	14% (8.9-20%)	5.3 (4.1-6.7)	16% (13-19%)	25% (20-30%)	69 (54-86)	3400 (2700-4200)
Slovenia	2010	Male	50 to 54	83% (77-87%)	17% (13-23%)	5.8 (4.6-7.3)	17% (14-20%)	24% (20-28%)	110 (92-140)	4800 (3900-5800)
Slovenia	2010	Male	55 to 59	79% (75-83%)	21% (17-25%)	6.5 (5.2-8.1)	16% (14-19%)	23% (19-26%)	170 (140-210)	6200 (5100-7300)
Slovenia	2010	Male	60 to 64	79% (75-83%)	21% (17-25%)	7 (5.5-8.6)	16% (13-18%)	21% (17-24%)	160 (130-190)	4900 (4100-5900)
Slovenia	2010	Male	65 to 69	81% (77-85%)	19% (15-23%)	7 (5.7-8.5)	14% (12-17%)	19% (16-22%)	180 (150-210)	4600 (3800-5400)
Slovenia	2010	Male	70 to 74	80% (75-84%)	20% (16-25%)	6.7 (5.3-8.3)	13% (11-16%)	16% (13-19%)	200 (170-250)	4200 (3400-5100)
Slovenia	2010	Male	75 to 79	75% (68-81%)	25% (19-32%)	5.8 (4.4-7.3)	11% (9-14%)	14% (11-17%)	200 (160-260)	3300 (2600-4100)
Slovenia	2010	Male	80 to 84	72% (64-79%)	28% (21-36%)	5 (3.8-6.3)	9.9% (7.3-13%)	12% (8.5-15%)	170 (120-220)	1900 (1400-2500)
Slovenia	2010	Male	85 to 89	73% (66-79%)	27% (21-34%)	4.6 (3.4-6.1)	8.9% (5.6-12%)	10% (6.3-14%)	99 (60-140)	810 (500-1100)
Slovenia	2010	Male	90 to 94	74% (66-81%)	26% (19-34%)	4.3 (2.9-5.9)	7.8% (3.7-12%)	8.9% (3.9-14%)	32 (14-50)	180 (84-270)
Slovenia	2010	Male	95 plus	74% (62-85%)	26% (15-38%)	3.8 (2.2-5.6)	6.7% (1.7-12%)	7.5% (1.4-14%)	7.5 (1.4-13)	28 (7-48)
Slovenia	2016	Female	All Ages	64% (59-68%)	36% (32-41%)	1.7 (1.3-2.2)	3.6% (2.5-5%)	4.3% (2.1-7%)	460 (220-760)	10000 (7000-14000)
Slovenia	2016	Female	15 to 19	67% (53-80%)	33% (20-47%)	1.6 (0.96-2.4)	2.4% (1.6-3.5%)	7.6% (3.8-12%)	0.53 (0.27-0.84)	90 (58-130)
Slovenia	2016	Female	20 to 24	77% (64-86%)	23% (14-36%)	2 (1.2-3.1)	3.9% (2.8-5.3%)	8.5% (4.5-13%)	0.78 (0.41-1.2)	180 (120-260)
Slovenia	2016	Female	25 to 29	79% (68-87%)	21% (13-32%)	1.7 (1.1-2.6)	3.6% (2.7-4.7%)	7.7% (4.6-11%)	0.79 (0.46-1.2)	220 (150-310)
Slovenia	2016	Female	30 to 34	80% (68-88%)	20% (12-32%)	1.6 (0.97-2.4)	3.3% (2.4-4.4%)	7.8% (5.1-11%)	1.4 (0.88-2.1)	270 (180-380)
Slovenia	2016	Female	35 to 39	79% (67-88%)	21% (12-33%)	1.7 (1.1-2.7)	3.6% (2.7-4.7%)	10% (7.3-13%)	3.4 (2.3-4.8)	360 (270-490)
Slovenia	2016	Female	40 to 44	75% (63-85%)	25% (15-37%)	2.1 (1.4-3.1)	4.2% (3.1-5.5%)	11% (8.8-14%)	6.2 (4.4-8.5)	460 (340-620)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	2016	Female	45 to 49	70% (58-79%)	30% (21-42%)	2.6 (1.7-3.7)	5.1% (3.7-6.9%)	12% (9.4-16%)	13 (9-17)	730 (540-980)
Slovenia	2016	Female	50 to 54	65% (56-74%)	35% (26-44%)	2.5 (1.7-3.6)	5.4% (4-7.1%)	11% (8.9-14%)	20 (14-26)	940 (700-1200)
Slovenia	2016	Female	55 to 59	61% (53-71%)	39% (29-47%)	2.3 (1.6-3.3)	5.2% (3.7-6.7%)	10% (7.7-13%)	27 (20-36)	1100 (790-1400)
Slovenia	2016	Female	60 to 64	59% (50-70%)	41% (30-50%)	2.2 (1.5-3.2)	4.6% (3.2-6.3%)	8.6% (6.4-12%)	34 (25-47)	1100 (800-1600)
Slovenia	2016	Female	65 to 69	59% (50-69%)	41% (31-50%)	2.2 (1.5-3.1)	4.2% (2.9-5.8%)	7.4% (5.4-9.9%)	33 (23-46)	920 (620-1300)
Slovenia	2016	Female	70 to 74	57% (47-67%)	43% (33-53%)	2.1 (1.4-3.1)	3.8% (2.2-5.9%)	6.2% (3.8-9%)	43 (25-65)	950 (540-1500)
Slovenia	2016	Female	75 to 79	52% (41-64%)	48% (36-59%)	1.8 (1.2-2.8)	3.2% (1.5-5.5%)	4.6% (2.3-8%)	53 (24-93)	890 (410-1600)
Slovenia	2016	Female	80 to 84	48% (35-60%)	52% (40-65%)	1.6 (0.96-2.4)	2.7% (1-5%)	3.6% (1.3-6.6%)	66 (24-120)	790 (300-1500)
Slovenia	2016	Female	85 to 89	46% (33-57%)	54% (43-67%)	1.5 (0.87-2.4)	2.4% (0.6-4.8%)	3% (0.63-6.1%)	75 (15-160)	620 (160-1300)
Slovenia	2016	Female	90 to 94	46% (34-59%)	54% (41-66%)	1.6 (0.87-2.5)	2.3% (0.098-5.3%)	2.8% (-0.081-6.6%)	56 (-1.7-130)	310 (13-720)
Slovenia	2016	Female	95 plus	48% (33-64%)	52% (36-67%)	1.6 (0.79-2.8)	2.4% (-0.48-6.4%)	2.8% (-0.77-7.8%)	25 (-6.8-69)	90 (-18-240)
Slovenia	2016	Male	All Ages	74% (70-79%)	26% (21-30%)	4.2 (3.3-5.4)	12% (10-15%)	15% (12-18%)	1500 (1200-2000)	40000 (32000-49000)
Slovenia	2016	Male	15 to 19	62% (45-77%)	38% (23-55%)	2.1 (1.2-3.1)	5% (3.3-7.1%)	11% (5.7-17%)	1.9 (0.97-3)	220 (140-310)
Slovenia	2016	Male	20 to 24	76% (62-87%)	24% (13-38%)	3.7 (2.4-5.3)	11% (8.1-15%)	21% (13-30%)	6.9 (4-10)	690 (460-930)
Slovenia	2016	Male	25 to 29	82% (71-90%)	18% (10-29%)	3.8 (2.6-5.4)	11% (8-14%)	21% (13-29%)	8.8 (5.4-13)	880 (620-1200)
Slovenia	2016	Male	30 to 34	85% (74-92%)	15% (7.8-26%)	3.9 (2.6-5.5)	11% (7.9-14%)	21% (14-29%)	13 (8.4-18)	1100 (820-1500)
Slovenia	2016	Male	35 to 39	86% (76-92%)	14% (8-24%)	3.9 (2.7-5.6)	11% (8.6-15%)	23% (16-30%)	20 (14-28)	1500 (1100-2000)
Slovenia	2016	Male	40 to 44	85% (76-91%)	15% (8.6-24%)	4.4 (3-5.9)	13% (9.8-16%)	25% (19-31%)	32 (22-43)	1900 (1400-2500)
Slovenia	2016	Male	45 to 49	84% (75-90%)	16% (9.9-25%)	5.1 (3.6-7.1)	15% (12-19%)	26% (21-32%)	57 (42-77)	2900 (2200-3700)
Slovenia	2016	Male	50 to 54	81% (75-86%)	19% (14-25%)	5.8 (4.2-7.9)	16% (13-20%)	26% (21-30%)	97 (74-130)	4100 (3200-5200)
Slovenia	2016	Male	55 to 59	78% (73-83%)	22% (17-27%)	6.5 (4.8-8.7)	16% (14-19%)	24% (20-27%)	140 (110-180)	5200 (4100-6400)
Slovenia	2016	Male	60 to 64	77% (72-83%)	23% (17-28%)	6.9 (5-9)	15% (13-18%)	21% (17-24%)	200 (160-250)	6100 (4900-7600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Slovenia	2016	Male	65 to 69	80% (74-84%)	20% (16-26%)	6.9 (5.1-9.1)	14% (11-17%)	19% (15-22%)	180 (140-230)	4600 (3600-5800)
Slovenia	2016	Male	70 to 74	77% (71-82%)	23% (18-29%)	6.5 (4.6-8.7)	12% (10-15%)	16% (13-19%)	190 (140-250)	4000 (3000-5000)
Slovenia	2016	Male	75 to 79	71% (62-78%)	29% (22-38%)	5.4 (3.7-7.4)	10% (7.7-13%)	13% (9.5-16%)	180 (130-250)	3000 (2200-3900)
Slovenia	2016	Male	80 to 84	67% (57-76%)	33% (24-43%)	4.6 (3.1-6.3)	9% (6.1-12%)	11% (7.1-15%)	170 (110-250)	2000 (1400-2800)
Slovenia	2016	Male	85 to 89	69% (58-77%)	31% (23-42%)	4.3 (2.8-6.2)	8.1% (4.7-12%)	9.5% (5.4-14%)	130 (70-200)	1100 (590-1600)
Slovenia	2016	Male	90 to 94	70% (59-79%)	30% (21-41%)	4 (2.5-6)	7.3% (3.3-11%)	8.4% (3.4-13%)	59 (24-98)	320 (150-520)
Slovenia	2016	Male	95 plus	71% (57-83%)	29% (17-43%)	3.7 (2-5.7)	6.5% (1.7-11%)	7.4% (1.5-13%)	13 (2.6-24)	48 (12-85)
Belarus	1990	Female	All Ages	69% (63-75%)	31% (25-37%)	2.8 (2.3-3.3)	6.5% (4.2-9%)	5.9% (1.8-11%)	3300 (1000-6000)	110000 (72000-160000)
Belarus	1990	Female	15 to 19	63% (48-77%)	37% (23-52%)	2.2 (1.2-3.7)	5.2% (3.5-7.5%)	9.9% (5.5-16%)	18 (10-30)	2100 (1400-3100)
Belarus	1990	Female	20 to 24	75% (60-86%)	25% (14-40%)	3.4 (1.8-5.5)	8.5% (6.1-11%)	13% (7.6-19%)	27 (16-39)	3700 (2600-5000)
Belarus	1990	Female	25 to 29	80% (67-88%)	20% (12-33%)	3.8 (2.2-5.9)	9.2% (7.1-12%)	15% (9.7-21%)	40 (26-57)	5200 (3700-6800)
Belarus	1990	Female	30 to 34	83% (69-91%)	17% (9.1-31%)	4.2 (2.6-6.4)	9.8% (7.6-12%)	17% (12-23%)	66 (47-89)	6800 (5000-8800)
Belarus	1990	Female	35 to 39	84% (73-92%)	16% (8.5-27%)	4.4 (2.5-6.7)	11% (8.3-13%)	18% (13-24%)	92 (65-120)	7500 (5700-9500)
Belarus	1990	Female	40 to 44	84% (73-92%)	16% (8.5-27%)	4.4 (2.5-6.8)	12% (8.6-15%)	19% (13-25%)	130 (87-170)	8300 (6100-11000)
Belarus	1990	Female	45 to 49	83% (69-91%)	17% (8.6-31%)	4.3 (2.6-6.8)	12% (8.3-16%)	18% (12-25%)	150 (100-210)	8300 (5900-11000)
Belarus	1990	Female	50 to 54	80% (66-89%)	20% (11-34%)	4.1 (2.4-6.5)	11% (7.4-15%)	16% (10-23%)	300 (180-420)	14000 (8900-19000)
Belarus	1990	Female	55 to 59	75% (59-87%)	25% (13-41%)	3.8 (2.2-5.8)	10% (5.8-15%)	14% (7.7-21%)	370 (200-540)	14000 (7900-20000)
Belarus	1990	Female	60 to 64	71% (53-85%)	29% (15-47%)	3.4 (1.9-5.5)	8.4% (3.4-14%)	11% (3.9-19%)	470 (170-790)	15000 (5800-25000)
Belarus	1990	Female	65 to 69	66% (47-82%)	34% (18-53%)	2.9 (1.7-4.4)	6.6% (1.8-12%)	8.2% (1.8-15%)	450 (97-840)	12000 (3100-21000)
Belarus	1990	Female	70 to 74	59% (41-75%)	41% (25-59%)	2.3 (1.2-4.1)	4.5% (-0.45-11%)	5.3% (-0.93-13%)	270 (-49-670)	5600 (-580-13000)
Belarus	1990	Female	75 to 79	49% (32-68%)	51% (32-68%)	1.8 (0.88-3.2)	3% (-1.5-8.8%)	3.4% (-2.1-11%)	310 (-190-970)	5000 (-2400-15000)
Belarus	1990	Female	80 to 84	43% (26-61%)	57% (39-74%)	1.6 (0.81-2.7)	2.5% (-1.1-7.8%)	2.7% (-1.7-9.1%)	270 (-170-910)	3200 (-1400-9800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	1990	Female	85 to 89	40% (24-57%)	60% (43-76%)	1.6 (0.79-2.8)	2.3% (-1.2-7.9%)	2.5% (-1.8-9.2%)	190 (-130-700)	1600 (-850-5300)
Belarus	1990	Female	90 to 94	41% (27-58%)	59% (42-73%)	1.7 (0.8-3.1)	2.3% (-1.5-8.4%)	2.5% (-2.2-9.9%)	84 (-73-320)	490 (-320-1700)
Belarus	1990	Female	95 plus	45% (26-65%)	55% (35-74%)	1.8 (0.7-3.7)	2.5% (-2.1-9.8%)	2.7% (-2.9-12%)	62 (-67-270)	230 (-180-900)
Belarus	1990	Male	All Ages	79% (73-85%)	21% (15-27%)	4.8 (4-5.7)	15% (12-18%)	15% (11-19%)	8000 (5700-10000)	310000 (250000-370000)
Belarus	1990	Male	15 to 19	69% (53-83%)	31% (17-47%)	2.8 (1.5-4.4)	9% (5.7-13%)	14% (8-21%)	61 (36-93)	5000 (3100-7400)
Belarus	1990	Male	20 to 24	82% (69-91%)	18% (9.4-31%)	5.1 (3.1-7.5)	18% (12-25%)	23% (15-33%)	180 (120-250)	14000 (9300-19000)
Belarus	1990	Male	25 to 29	87% (79-93%)	13% (6.7-21%)	6.3 (4-9)	21% (16-27%)	28% (20-36%)	300 (210-390)	22000 (16000-27000)
Belarus	1990	Male	30 to 34	90% (82-96%)	9.7% (4.3-18%)	7.1 (4.9-9.9)	23% (18-28%)	30% (22-36%)	420 (310-530)	28000 (21000-34000)
Belarus	1990	Male	35 to 39	91% (85-96%)	8.7% (4-15%)	7.5 (4.9-11)	23% (18-29%)	30% (22-37%)	500 (370-620)	29000 (22000-35000)
Belarus	1990	Male	40 to 44	92% (85-96%)	8.5% (3.9-15%)	8 (5.1-11)	24% (18-29%)	30% (22-36%)	600 (450-740)	30000 (23000-37000)
Belarus	1990	Male	45 to 49	91% (82-96%)	8.9% (4-18%)	8.5 (5.5-12)	24% (18-28%)	28% (22-34%)	650 (500-790)	29000 (22000-35000)
Belarus	1990	Male	50 to 54	90% (81-95%)	10% (5-19%)	8.2 (5.3-12)	21% (16-25%)	25% (19-30%)	1100 (850-1400)	43000 (33000-53000)
Belarus	1990	Male	55 to 59	88% (78-94%)	12% (6-22%)	7.2 (4.8-10)	18% (13-22%)	21% (15-26%)	1100 (810-1400)	37000 (28000-47000)
Belarus	1990	Male	60 to 64	86% (74-93%)	14% (6.6-26%)	6.3 (3.9-9.4)	14% (8.8-19%)	16% (9.8-22%)	1200 (700-1600)	34000 (21000-46000)
Belarus	1990	Male	65 to 69	85% (72-93%)	15% (7.2-28%)	5.3 (3.5-7.7)	12% (6.4-17%)	13% (6.9-19%)	750 (390-1100)	18000 (9700-26000)
Belarus	1990	Male	70 to 74	82% (69-90%)	18% (9.9-31%)	4.3 (2.5-6.6)	8.5% (2.5-14%)	9.4% (2.4-16%)	370 (99-640)	7300 (2100-12000)
Belarus	1990	Male	75 to 79	78% (63-89%)	22% (11-37%)	3.6 (2.1-5.8)	6% (0.049-12%)	6.4% (-0.42-14%)	350 (-24-760)	5500 (45-11000)
Belarus	1990	Male	80 to 84	74% (57-85%)	26% (15-43%)	3.3 (1.9-4.9)	4.7% (-0.73-11%)	4.9% (-1.3-12%)	250 (-69-620)	2900 (-460-6800)
Belarus	1990	Male	85 to 89	70% (54-82%)	30% (18-46%)	3.1 (1.8-4.9)	3.6% (-1.1-9.9%)	3.7% (-1.8-11%)	110 (-57-340)	990 (-290-2700)
Belarus	1990	Male	90 to 94	69% (54-81%)	31% (19-46%)	2.9 (1.6-4.7)	2.7% (-1.9-8.5%)	2.6% (-2.9-9.5%)	30 (-34-110)	190 (-140-600)
Belarus	1990	Male	95 plus	67% (48-83%)	33% (17-52%)	2.7 (1.3-4.8)	2.1% (-2.8-8.3%)	1.8% (-4.9-1%)	8.7 (-19-44)	41 (-54-160)
Belarus	1995	Female	All Ages	70% (63-76%)	30% (24-37%)	2.7 (2.3-3.2)	6.9% (4.3-9.8%)	6% (1.9-11%)	3900 (1200-7200)	130000 (82000-180000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	1995	Female	15 to 19	64% (48-79%)	36% (21-52%)	2.2 (1.2-3.5)	5.5% (3.6-7.9%)	10% (5.9-16%)	21 (12-33)	2300 (1500-3300)
Belarus	1995	Female	20 to 24	76% (61-87%)	24% (13-39%)	3.3 (1.8-5.3)	8.9% (6.3-12%)	14% (8-20%)	32 (19-47)	4100 (2800-5600)
Belarus	1995	Female	25 to 29	80% (68-89%)	20% (11-32%)	3.7 (2.2-5.6)	9.9% (7.4-13%)	15% (9.8-22%)	44 (27-62)	5000 (3600-6600)
Belarus	1995	Female	30 to 34	83% (70-91%)	17% (8.8-30%)	4 (2.5-6.1)	10% (7.9-13%)	17% (12-24%)	79 (53-110)	7400 (5400-9600)
Belarus	1995	Female	35 to 39	84% (73-92%)	16% (8.2-27%)	4.2 (2.5-6.4)	11% (8.6-14%)	19% (13-24%)	130 (92-180)	10000 (7500-13000)
Belarus	1995	Female	40 to 44	84% (74-92%)	16% (7.8-26%)	4.3 (2.6-6.5)	12% (9-16%)	19% (14-25%)	190 (130-250)	11000 (8500-15000)
Belarus	1995	Female	45 to 49	83% (68-92%)	17% (8.4-32%)	4.2 (2.5-6.7)	13% (8.8-17%)	19% (13-26%)	240 (160-330)	12000 (8500-16000)
Belarus	1995	Female	50 to 54	80% (67-89%)	20% (11-33%)	4 (2.3-6.4)	12% (7.8-17%)	17% (11-24%)	260 (160-380)	11000 (7400-16000)
Belarus	1995	Female	55 to 59	76% (61-88%)	24% (12-39%)	3.7 (2.1-5.6)	11% (6.1-16%)	15% (7.8-22%)	490 (260-750)	18000 (9900-27000)
Belarus	1995	Female	60 to 64	71% (54-85%)	29% (15-46%)	3.4 (1.9-5.5)	8.8% (3.3-15%)	11% (3.7-19%)	500 (160-870)	15000 (5600-26000)
Belarus	1995	Female	65 to 69	67% (48-82%)	33% (18-52%)	2.9 (1.7-4.4)	6.5% (1.4-12%)	8% (1.4-15%)	580 (98-1100)	15000 (3300-27000)
Belarus	1995	Female	70 to 74	60% (41-76%)	40% (24-59%)	2.3 (1.2-4)	4.4% (-0.85-11%)	5.1% (-1.5-13%)	450 (-130-1200)	9200 (-1800-23000)
Belarus	1995	Female	75 to 79	50% (33-70%)	50% (30-67%)	1.8 (0.87-3.2)	2.9% (-1.8-8.8%)	3.2% (-2.5-10%)	240 (-190-790)	3800 (-2300-11000)
Belarus	1995	Female	80 to 84	44% (26-62%)	56% (38-74%)	1.6 (0.8-2.7)	2.3% (-1.4-7.4%)	2.5% (-2-8.4%)	290 (-230-980)	3400 (-2100-11000)
Belarus	1995	Female	85 to 89	41% (26-58%)	59% (42-74%)	1.6 (0.8-2.7)	2.2% (-1.5-7.8%)	2.4% (-2-8.8%)	220 (-180-810)	1800 (-1200-6200)
Belarus	1995	Female	90 to 94	42% (27-59%)	58% (41-73%)	1.7 (0.8-3.1)	2.3% (-2-8.8%)	2.4% (-2.6-10%)	110 (-120-460)	610 (-520-2400)
Belarus	1995	Female	95 plus	46% (26-66%)	54% (34-74%)	1.9 (0.71-3.6)	2.5% (-2.6-10%)	2.6% (-3.4-12%)	55 (-72-250)	210 (-210-850)
Belarus	1995	Male	All Ages	80% (74-85%)	20% (15-26%)	4.7 (4.1-5.5)	16% (13-19%)	16% (11-20%)	11000 (7900-14000)	400000 (330000-480000)
Belarus	1995	Male	15 to 19	69% (54-82%)	31% (18-46%)	2.7 (1.6-4.3)	9.4% (5.7-14%)	14% (8-21%)	71 (41-110)	5800 (3600-8400)
Belarus	1995	Male	20 to 24	82% (70-91%)	18% (9.4-30%)	5 (3-7.3)	19% (13-25%)	23% (15-32%)	220 (140-300)	16000 (11000-22000)
Belarus	1995	Male	25 to 29	87% (79-93%)	13% (6.9-21%)	6.2 (3.9-8.7)	23% (16-28%)	28% (20-36%)	330 (230-430)	23000 (16000-29000)
Belarus	1995	Male	30 to 34	90% (82-95%)	9.6% (4.5-18%)	7 (4.6-9.6)	24% (19-29%)	30% (22-36%)	540 (400-670)	34000 (26000-41000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	1995	Male	35 to 39	91% (85-96%)	8.6% (4.2-15%)	7.3 (4.9-10)	24% (19-29%)	29% (22-36%)	790 (580-980)	44000 (33000-53000)
Belarus	1995	Male	40 to 44	92% (85-96%)	8.4% (4.1-15%)	7.7 (5.2-11)	25% (19-30%)	30% (23-35%)	970 (730-1200)	47000 (37000-57000)
Belarus	1995	Male	45 to 49	91% (82-96%)	9% (4-18%)	8.2 (5.4-12)	24% (19-29%)	28% (22-34%)	1100 (870-1400)	48000 (38000-58000)
Belarus	1995	Male	50 to 54	90% (81-95%)	10% (4.6-19%)	7.9 (5.1-11)	22% (17-27%)	25% (19-31%)	1100 (790-1300)	40000 (30000-49000)
Belarus	1995	Male	55 to 59	88% (79-94%)	12% (5.6-21%)	7 (4.6-9.9)	19% (13-24%)	21% (15-27%)	1600 (1100-2000)	52000 (37000-67000)
Belarus	1995	Male	60 to 64	87% (76-94%)	13% (6.4-24%)	6.1 (3.9-9.2)	15% (8.6-20%)	16% (9.3-23%)	1400 (800-1900)	39000 (23000-54000)
Belarus	1995	Male	65 to 69	85% (73-93%)	15% (7-27%)	5.2 (3.4-7.5)	12% (5.9-17%)	13% (6.1-19%)	1300 (630-2000)	32000 (15000-46000)
Belarus	1995	Male	70 to 74	82% (70-90%)	18% (9.7-30%)	4.2 (2.6-6.5)	8.4% (2-15%)	9.1% (1.7-16%)	670 (130-1200)	13000 (3100-23000)
Belarus	1995	Male	75 to 79	79% (64-89%)	21% (11-36%)	3.6 (2-5.6)	5.9% (-0.32-12%)	6.3% (-0.82-14%)	280 (-37-600)	4300 (-220-8900)
Belarus	1995	Male	80 to 84	75% (59-86%)	25% (14-41%)	3.3 (1.9-4.9)	4.5% (-1-11%)	4.6% (-1.7-12%)	240 (-87-620)	2800 (-630-6800)
Belarus	1995	Male	85 to 89	71% (55-82%)	29% (18-45%)	3 (1.8-4.7)	3.4% (-1.4-10%)	3.4% (-2.2-11%)	120 (-79-390)	1000 (-430-3000)
Belarus	1995	Male	90 to 94	69% (54-82%)	31% (18-46%)	2.9 (1.6-4.6)	2.4% (-2.2-8.6%)	2.2% (-3.2-9.6%)	31 (-45-130)	200 (-190-710)
Belarus	1995	Male	95 plus	68% (50-84%)	32% (16-50%)	2.7 (1.2-4.7)	1.7% (-3.3-8.3%)	1.4% (-4.5-9.2%)	6 (-19-40)	30 (-57-140)
Belarus	2000	Female	All Ages	71% (64-77%)	29% (23-36%)	2.9 (2.4-3.4)	7.2% (4.5-10%)	6.1% (1.7-11%)	4000 (1100-7600)	130000 (83000-190000)
Belarus	2000	Female	15 to 19	64% (48-79%)	36% (21-52%)	2.2 (1.2-3.6)	5.4% (3.6-7.7%)	11% (5.9-17%)	21 (12-33)	2300 (1600-3400)
Belarus	2000	Female	20 to 24	76% (61-86%)	24% (14-39%)	3.3 (1.8-5.2)	9.1% (6.4-12%)	14% (8.2-21%)	34 (20-50)	4200 (2900-5700)
Belarus	2000	Female	25 to 29	80% (68-89%)	20% (11-32%)	3.8 (2.1-5.8)	10% (7.7-13%)	16% (9.9-23%)	50 (31-72)	5500 (3800-7200)
Belarus	2000	Female	30 to 34	83% (70-91%)	17% (8.7-30%)	4.1 (2.5-6.3)	11% (8.4-14%)	18% (12-25%)	75 (51-100)	6800 (5000-8700)
Belarus	2000	Female	35 to 39	84% (74-92%)	16% (8.2-26%)	4.3 (2.6-6.5)	12% (9-15%)	20% (14-26%)	130 (91-170)	9800 (7200-12000)
Belarus	2000	Female	40 to 44	84% (73-92%)	16% (7.9-27%)	4.4 (2.7-6.4)	13% (9.6-16%)	20% (14-27%)	210 (150-290)	13000 (9900-17000)
Belarus	2000	Female	45 to 49	83% (69-92%)	17% (8.2-31%)	4.3 (2.6-6.4)	13% (9.1-17%)	19% (13-26%)	280 (190-380)	14000 (10000-19000)
Belarus	2000	Female	50 to 54	80% (66-89%)	20% (11-34%)	4 (2.3-6.3)	13% (8.3-18%)	18% (11-25%)	340 (210-480)	14000 (9700-20000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	2000	Female	55 to 59	76% (60-88%)	24% (12-40%)	3.7 (2.1-5.8)	12% (6.6-17%)	16% (8.1-23%)	350 (190-510)	12000 (7000-18000)
Belarus	2000	Female	60 to 64	72% (55-85%)	28% (15-45%)	3.4 (1.9-5.6)	9.1% (3.4-15%)	12% (3.9-20%)	550 (180-950)	17000 (6300-28000)
Belarus	2000	Female	65 to 69	67% (49-83%)	33% (17-51%)	2.9 (1.7-4.4)	6.6% (1.3-12%)	8.1% (1.1-16%)	510 (76-1000)	13000 (2500-25000)
Belarus	2000	Female	70 to 74	60% (43-76%)	40% (24-57%)	2.3 (1.3-4)	4.3% (-0.92-11%)	5.1% (-1.6-13%)	530 (-160-1400)	11000 (-2300-26000)
Belarus	2000	Female	75 to 79	51% (33-69%)	49% (31-67%)	1.9 (0.9-3.2)	2.8% (-1.9-8.9%)	3.2% (-2.6-11%)	370 (-310-1200)	5900 (-4000-18000)
Belarus	2000	Female	80 to 84	44% (27-62%)	56% (38-73%)	1.6 (0.79-2.7)	2.3% (-1.6-7.4%)	2.5% (-2.2-8.4%)	200 (-180-690)	2400 (-1600-7600)
Belarus	2000	Female	85 to 89	41% (26-58%)	59% (42-74%)	1.6 (0.78-2.7)	2.3% (-1.5-7.9%)	2.4% (-2-9.1%)	220 (-180-820)	1800 (-1200-6200)
Belarus	2000	Female	90 to 94	43% (27-60%)	57% (40-73%)	1.7 (0.81-3)	2.3% (-2-9.1%)	2.5% (-2.7-10%)	110 (-120-470)	630 (-540-2400)
Belarus	2000	Female	95 plus	47% (26-68%)	53% (32-74%)	1.9 (0.7-3.7)	2.6% (-2.9-11%)	2.7% (-3.8-12%)	52 (-74-240)	190 (-220-810)
Belarus	2000	Male	All Ages	81% (75-86%)	19% (14-25%)	5 (4.3-5.7)	17% (14-20%)	16% (12-21%)	12000 (8500-15000)	430000 (350000-520000)
Belarus	2000	Male	15 to 19	70% (54-83%)	30% (17-46%)	2.7 (1.5-4.4)	9.1% (5.6-14%)	14% (8.1-22%)	68 (39-100)	5500 (3400-8100)
Belarus	2000	Male	20 to 24	82% (70-91%)	18% (9.3-30%)	5 (3.1-7.4)	19% (13-26%)	24% (15-33%)	240 (160-340)	18000 (12000-24000)
Belarus	2000	Male	25 to 29	87% (78-93%)	13% (6.8-22%)	6.2 (3.9-9)	23% (17-29%)	28% (20-36%)	370 (250-480)	25000 (18000-32000)
Belarus	2000	Male	30 to 34	90% (82-96%)	9.5% (4.5-18%)	7 (4.6-9.8)	25% (19-30%)	30% (23-37%)	510 (370-630)	32000 (24000-38000)
Belarus	2000	Male	35 to 39	92% (85-96%)	8.4% (4.2-15%)	7.4 (4.9-10)	25% (19-30%)	30% (23-37%)	780 (580-960)	43000 (33000-52000)
Belarus	2000	Male	40 to 44	92% (85-96%)	8.2% (3.9-15%)	7.8 (5.3-11)	26% (20-31%)	30% (24-36%)	1100 (870-1400)	55000 (43000-66000)
Belarus	2000	Male	45 to 49	91% (82-96%)	8.7% (3.9-18%)	8.3 (5.6-12)	25% (20-30%)	29% (23-35%)	1300 (1000-1600)	57000 (45000-69000)
Belarus	2000	Male	50 to 54	90% (82-95%)	9.9% (4.6-18%)	8 (5.2-11)	23% (17-28%)	26% (20-32%)	1400 (1000-1700)	51000 (39000-63000)
Belarus	2000	Male	55 to 59	88% (78-95%)	12% (5.4-22%)	7.1 (4.7-10)	19% (14-25%)	22% (16-28%)	1100 (810-1500)	37000 (27000-48000)
Belarus	2000	Male	60 to 64	87% (75-94%)	13% (6.3-25%)	6.2 (3.7-9.4)	15% (8.7-21%)	17% (9.4-24%)	1600 (890-2300)	45000 (25000-63000)
Belarus	2000	Male	65 to 69	85% (72-93%)	15% (7-28%)	5.3 (3.4-7.7)	12% (6-18%)	13% (6.2-20%)	1300 (590-2000)	30000 (15000-45000)
Belarus	2000	Male	70 to 74	82% (70-90%)	18% (9.7-30%)	4.3 (2.5-6.6)	8.5% (1.8-15%)	9.3% (1.6-17%)	1000 (180-1900)	20000 (4200-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	2000	Male	75 to 79	79% (65-89%)	21% (11-35%)	3.6 (2-5.8)	5.8% (-0.64-13%)	6.2% (-1.1-14%)	440 (-82-990)	6600 (-690-14000)
Belarus	2000	Male	80 to 84	75% (60-86%)	25% (14-40%)	3.3 (2-4.9)	4.6% (-1-11%)	4.7% (-1.7-12%)	170 (-60-440)	2000 (-460-4800)
Belarus	2000	Male	85 to 89	71% (56-83%)	29% (17-44%)	3.1 (1.8-4.8)	3.5% (-1.7-10%)	3.5% (-2.5-11%)	100 (-72-320)	850 (-420-2500)
Belarus	2000	Male	90 to 94	69% (56-82%)	31% (18-44%)	2.9 (1.6-4.7)	2.4% (-2.5-8.6%)	2.2% (-3.5-9.5%)	28 (-45-120)	180 (-190-650)
Belarus	2000	Male	95 plus	68% (50-83%)	32% (17-50%)	2.7 (1.3-4.6)	1.7% (-3.5-8.5%)	1.3% (-4.7-9.3%)	5.7 (-21-41)	29 (-60-150)
Belarus	2005	Female	All Ages	71% (65-77%)	29% (23-35%)	3.1 (2.6-3.6)	7.7% (4.8-11%)	6.6% (2-12%)	4300 (1300-8200)	140000 (88000-200000)
Belarus	2005	Female	15 to 19	64% (48-78%)	36% (22-52%)	2.3 (1.2-3.7)	5.1% (3.4-7.2%)	11% (6.4-17%)	16 (9.4-26)	2000 (1400-2900)
Belarus	2005	Female	20 to 24	75% (61-86%)	25% (14-39%)	3.5 (1.9-5.5)	9.2% (6.6-12%)	15% (8.7-22%)	35 (20-51)	4400 (3100-5900)
Belarus	2005	Female	25 to 29	80% (68-89%)	20% (11-32%)	3.9 (2.2-6.2)	11% (8-14%)	17% (11-24%)	55 (34-77)	5800 (4100-7700)
Belarus	2005	Female	30 to 34	83% (71-91%)	17% (8.6-29%)	4.3 (2.6-6.4)	12% (9-15%)	19% (13-25%)	86 (59-120)	7400 (5500-9600)
Belarus	2005	Female	35 to 39	84% (74-92%)	16% (7.9-26%)	4.5 (2.6-6.6)	13% (9.7-16%)	21% (15-27%)	130 (91-170)	9200 (6800-12000)
Belarus	2005	Female	40 to 44	84% (74-92%)	16% (8-26%)	4.5 (2.7-6.8)	13% (9.9-17%)	21% (15-28%)	210 (150-280)	13000 (9800-17000)
Belarus	2005	Female	45 to 49	83% (69-91%)	17% (8.5-31%)	4.4 (2.8-6.7)	13% (9.6-17%)	20% (14-27%)	320 (220-430)	17000 (12000-22000)
Belarus	2005	Female	50 to 54	80% (67-89%)	20% (11-33%)	4.2 (2.5-6.4)	13% (9.1-18%)	19% (13-26%)	400 (270-550)	17000 (12000-23000)
Belarus	2005	Female	55 to 59	76% (60-87%)	24% (13-40%)	3.9 (2.2-6)	13% (7.4-18%)	17% (9.6-24%)	480 (280-700)	17000 (10000-25000)
Belarus	2005	Female	60 to 64	71% (54-85%)	29% (15-46%)	3.6 (2-5.8)	9.9% (4.2-16%)	13% (5-21%)	390 (150-650)	12000 (5000-19000)
Belarus	2005	Female	65 to 69	67% (49-82%)	33% (18-51%)	3 (1.8-4.6)	7% (1.7-13%)	8.7% (1.7-17%)	550 (110-1100)	14000 (3400-26000)
Belarus	2005	Female	70 to 74	61% (43-76%)	39% (24-57%)	2.4 (1.3-4.1)	4.6% (-1-11%)	5.4% (-1.8-14%)	460 (-150-1200)	9500 (-2100-23000)
Belarus	2005	Female	75 to 79	51% (34-69%)	49% (31-66%)	1.9 (0.92-3.3)	3% (-2-9.4%)	3.4% (-2.8-11%)	450 (-370-1500)	7200 (-4700-22000)
Belarus	2005	Female	80 to 84	44% (27-62%)	56% (38-73%)	1.7 (0.86-2.8)	2.7% (-1.4-8.5%)	2.9% (-1.9-9.6%)	380 (-250-1300)	4400 (-2200-14000)
Belarus	2005	Female	85 to 89	42% (25-58%)	58% (42-75%)	1.7 (0.85-2.9)	2.7% (-1.4-8.7%)	2.9% (-2-10%)	190 (-130-650)	1500 (-810-4800)
Belarus	2005	Female	90 to 94	43% (28-60%)	57% (40-72%)	1.8 (0.84-3.2)	2.9% (-1.9-10%)	3.1% (-2.5-12%)	130 (-110-500)	740 (-500-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	2005	Female	95 plus	47% (27-67%)	53% (33-73%)	2 (0.77-3.9)	3.1% (-2.5-12%)	3.4% (-3.2-14%)	55 (-51-220)	200 (-150-740)
Belarus	2005	Male	All Ages	82% (76-87%)	18% (13-24%)	5.4 (4.6-6.2)	18% (14-21%)	17% (12-22%)	12000 (8900-16000)	450000 (370000-550000)
Belarus	2005	Male	15 to 19	70% (54-83%)	30% (17-46%)	2.8 (1.6-4.5)	9.2% (5.6-13%)	15% (8.3-22%)	62 (35-96)	5200 (3100-7600)
Belarus	2005	Male	20 to 24	82% (70-90%)	18% (9.5-30%)	5.2 (3.2-7.6)	19% (13-25%)	25% (16-34%)	230 (150-310)	17000 (11000-22000)
Belarus	2005	Male	25 to 29	87% (79-93%)	13% (6.7-21%)	6.4 (4.1-9.3)	23% (17-29%)	29% (21-37%)	360 (260-460)	25000 (18000-31000)
Belarus	2005	Male	30 to 34	90% (82-95%)	9.5% (4.6-18%)	7.3 (4.8-10)	25% (19-30%)	31% (23-37%)	530 (400-650)	32000 (25000-39000)
Belarus	2005	Male	35 to 39	92% (85-96%)	8.5% (4.2-15%)	7.7 (5-11)	26% (20-31%)	31% (23-37%)	690 (520-840)	38000 (29000-46000)
Belarus	2005	Male	40 to 44	92% (85-96%)	8.2% (3.9-15%)	8.1 (5.5-11)	27% (21-31%)	31% (25-37%)	1100 (870-1400)	55000 (43000-66000)
Belarus	2005	Male	45 to 49	91% (82-96%)	8.8% (3.9-18%)	8.6 (5.8-12)	26% (20-31%)	30% (24-36%)	1600 (1300-2000)	68000 (55000-83000)
Belarus	2005	Male	50 to 54	90% (81-95%)	9.9% (4.5-19%)	8.3 (5.4-12)	24% (18-28%)	27% (21-33%)	1600 (1200-2000)	62000 (48000-76000)
Belarus	2005	Male	55 to 59	88% (78-95%)	12% (5.3-22%)	7.4 (4.8-11)	20% (15-26%)	23% (16-29%)	1500 (1100-2000)	50000 (36000-64000)
Belarus	2005	Male	60 to 64	87% (76-94%)	13% (6.3-24%)	6.5 (3.9-9.9)	16% (9.4-22%)	18% (10-25%)	1200 (660-1600)	33000 (19000-45000)
Belarus	2005	Male	65 to 69	85% (72-93%)	15% (6.9-28%)	5.5 (3.6-8)	13% (6.5-19%)	14% (6.9-22%)	1500 (720-2200)	35000 (18000-51000)
Belarus	2005	Male	70 to 74	82% (70-90%)	18% (9.6-30%)	4.5 (2.7-6.7)	9.1% (2.3-16%)	9.9% (2.2-18%)	970 (210-1700)	19000 (4600-32000)
Belarus	2005	Male	75 to 79	79% (66-89%)	21% (11-34%)	3.8 (2.1-6)	6.3% (-0.47-13%)	6.7% (-0.99-15%)	670 (-100-1500)	10000 (-780-21000)
Belarus	2005	Male	80 to 84	75% (61-86%)	25% (14-39%)	3.5 (2-5.1)	5.1% (-0.92-12%)	5.3% (-1.6-13%)	280 (-85-720)	3200 (-600-7800)
Belarus	2005	Male	85 to 89	72% (57-83%)	28% (17-43%)	3.2 (1.9-4.8)	3.9% (-1.7-11%)	4% (-2.5-12%)	83 (-53-250)	700 (-310-1900)
Belarus	2005	Male	90 to 94	70% (56-82%)	30% (18-44%)	3 (1.7-4.8)	2.7% (-2.6-9.4%)	2.6% (-3.7-10%)	28 (-40-110)	170 (-170-600)
Belarus	2005	Male	95 plus	68% (50-83%)	32% (17-50%)	2.8 (1.3-4.8)	1.9% (-3.5-8.9%)	1.6% (-4.8-9.7%)	5.4 (-17-33)	26 (-48-120)
Belarus	2010	Female	All Ages	71% (65-77%)	29% (23-35%)	3.3 (2.8-3.9)	8.1% (5.3-11%)	7.1% (2.3-13%)	4600 (1500-8800)	140000 (91000-200000)
Belarus	2010	Female	15 to 19	64% (49-78%)	36% (22-51%)	2.4 (1.3-4)	5.2% (3.5-7.2%)	12% (6.7-18%)	12 (6.9-19)	1500 (1000-2100)
Belarus	2010	Female	20 to 24	76% (62-86%)	24% (14-38%)	3.8 (2.1-6)	8.9% (6.6-12%)	15% (9.5-22%)	26 (16-38)	3900 (2700-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	2010	Female	25 to 29	80% (68-89%)	20% (11-32%)	4.2 (2.4-6.5)	10% (7.9-13%)	17% (11-24%)	48 (31-67)	5600 (4000-7400)
Belarus	2010	Female	30 to 34	83% (71-91%)	17% (8.8-29%)	4.5 (2.8-7)	12% (9.1-15%)	19% (14-25%)	82 (59-110)	7200 (5400-9200)
Belarus	2010	Female	35 to 39	85% (74-92%)	15% (8.2-26%)	4.8 (2.8-7.4)	13% (9.8-16%)	21% (15-27%)	120 (87-160)	8900 (6700-11000)
Belarus	2010	Female	40 to 44	85% (74-92%)	15% (7.9-26%)	4.9 (2.9-7.3)	14% (10-17%)	23% (16-29%)	180 (130-240)	11000 (8200-14000)
Belarus	2010	Female	45 to 49	83% (70-92%)	17% (8.4-30%)	4.7 (2.9-7.3)	14% (10-18%)	22% (16-28%)	280 (200-370)	15000 (11000-19000)
Belarus	2010	Female	50 to 54	80% (67-90%)	20% (10-33%)	4.5 (2.7-6.9)	14% (9.8-18%)	20% (14-27%)	430 (290-570)	19000 (13000-25000)
Belarus	2010	Female	55 to 59	76% (60-88%)	24% (12-40%)	4.2 (2.4-6.4)	13% (8.3-18%)	18% (11-26%)	540 (340-760)	20000 (12000-27000)
Belarus	2010	Female	60 to 64	72% (55-85%)	28% (15-45%)	3.8 (2.1-6.1)	11% (5.1-16%)	14% (6.5-22%)	480 (220-770)	15000 (7100-23000)
Belarus	2010	Female	65 to 69	67% (49-82%)	33% (18-51%)	3.3 (1.9-4.9)	7.7% (2.3-14%)	9.8% (2.4-19%)	360 (90-690)	9300 (2800-17000)
Belarus	2010	Female	70 to 74	60% (43-76%)	40% (24-57%)	2.6 (1.3-4.4)	5.2% (-0.44-12%)	6.3% (-1.1-16%)	510 (-93-1300)	11000 (-870-25000)
Belarus	2010	Female	75 to 79	51% (34-69%)	49% (31-66%)	2.1 (1-3.7)	3.6% (-1.6-11%)	4.1% (-2.4-13%)	440 (-260-1400)	7100 (-3000-21000)
Belarus	2010	Female	80 to 84	44% (27-62%)	56% (38-73%)	1.8 (0.93-3)	3.2% (-1-9.4%)	3.6% (-1.5-11%)	520 (-220-1600)	6100 (-1900-18000)
Belarus	2010	Female	85 to 89	42% (26-58%)	58% (42-74%)	1.8 (0.9-3.1)	3.2% (-1.3-9.9%)	3.5% (-1.8-12%)	370 (-190-1200)	3000 (-1200-9100)
Belarus	2010	Female	90 to 94	43% (28-60%)	57% (40-72%)	1.9 (0.9-3.5)	3.3% (-1.4-11%)	3.6% (-2.1-12%)	120 (-71-430)	680 (-300-2200)
Belarus	2010	Female	95 plus	47% (28-67%)	53% (33-72%)	2.1 (0.81-4.1)	3.6% (-2.3-12%)	3.9% (-3-15%)	57 (-45-220)	210 (-140-740)
Belarus	2010	Male	All Ages	82% (77-87%)	18% (13-23%)	5.8 (4.8-6.9)	18% (15-22%)	17% (12-22%)	12000 (8700-16000)	430000 (350000-520000)
Belarus	2010	Male	15 to 19	70% (54-83%)	30% (17-46%)	3 (1.7-4.7)	9.1% (5.6-13%)	15% (8.6-23%)	44 (25-67)	3700 (2300-5500)
Belarus	2010	Male	20 to 24	82% (70-91%)	18% (9.5-30%)	5.5 (3.4-8.3)	19% (13-24%)	26% (17-35%)	190 (120-260)	15000 (10000-19000)
Belarus	2010	Male	25 to 29	87% (79-94%)	13% (6.5-21%)	6.9 (4.4-10)	23% (17-28%)	29% (22-37%)	320 (230-400)	22000 (17000-28000)
Belarus	2010	Male	30 to 34	90% (82-96%)	9.5% (4.5-18%)	7.8 (5.2-11)	25% (20-30%)	31% (24-37%)	470 (360-560)	29000 (23000-35000)
Belarus	2010	Male	35 to 39	91% (85-96%)	8.5% (4-15%)	8.2 (5.1-12)	26% (20-31%)	32% (24-38%)	630 (480-750)	35000 (27000-42000)
Belarus	2010	Male	40 to 44	92% (85-96%)	8.2% (4.1-15%)	8.6 (5.8-12)	27% (22-32%)	32% (26-38%)	880 (700-1100)	43000 (35000-52000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	2010	Male	45 to 49	91% (83-96%)	8.7% (4-17%)	9.1 (6.2-13)	26% (21-31%)	31% (25-37%)	1400 (1100-1600)	59000 (47000-70000)
Belarus	2010	Male	50 to 54	90% (82-95%)	9.9% (4.8-18%)	8.8 (5.8-12)	24% (19-29%)	28% (22-34%)	1800 (1400-2200)	67000 (52000-81000)
Belarus	2010	Male	55 to 59	88% (79-94%)	12% (5.8-21%)	7.8 (5.2-11)	21% (16-26%)	24% (18-30%)	1700 (1300-2100)	56000 (42000-70000)
Belarus	2010	Male	60 to 64	87% (76-94%)	13% (6.4-24%)	6.9 (4.3-10)	17% (11-24%)	19% (12-27%)	1500 (930-2100)	43000 (27000-59000)
Belarus	2010	Male	65 to 69	85% (73-93%)	15% (6.7-27%)	5.9 (3.8-8.6)	14% (7.5-20%)	16% (8.1-23%)	970 (510-1400)	23000 (13000-33000)
Belarus	2010	Male	70 to 74	82% (70-91%)	18% (9.1-30%)	4.8 (2.8-7.2)	10% (3.1-17%)	11% (2.9-19%)	1100 (280-1900)	21000 (6100-35000)
Belarus	2010	Male	75 to 79	79% (65-89%)	21% (11-35%)	4 (2.3-6.4)	7% (-0.21-14%)	7.6% (-0.84-15%)	640 (-71-1300)	9700 (-290-19000)
Belarus	2010	Male	80 to 84	75% (60-86%)	25% (14-40%)	3.7 (2.2-5.5)	5.9% (-0.59-13%)	6.3% (-1.4-15%)	470 (-100-1100)	5300 (-570-12000)
Belarus	2010	Male	85 to 89	72% (56-83%)	28% (17-44%)	3.4 (2.1-5.3)	4.7% (-1.1-12%)	4.9% (-2-13%)	160 (-65-430)	1300 (-320-3200)
Belarus	2010	Male	90 to 94	70% (55-82%)	30% (18-45%)	3.2 (1.8-5.2)	3.5% (-2.3-11%)	3.5% (-3.5-12%)	31 (-31-100)	190 (-130-560)
Belarus	2010	Male	95 plus	68% (50-83%)	32% (17-50%)	3 (1.5-5.2)	2.6% (-3.4-10%)	2.3% (-4.8-11%)	6.7 (-14-32)	30 (-39-110)
Belarus	2016	Female	All Ages	71% (64-77%)	29% (23-36%)	3.4 (2.6-4.4)	7.7% (4.8-12%)	6.9% (1.8-14%)	4200 (1100-8600)	130000 (75000-190000)
Belarus	2016	Female	15 to 19	64% (48-78%)	36% (22-52%)	2.5 (1.3-4.3)	4.8% (3.3-6.9%)	12% (6.8-19%)	6.4 (3.4-11)	950 (630-1400)
Belarus	2016	Female	20 to 24	75% (61-86%)	25% (14-39%)	3.8 (1.9-6.5)	8.7% (6.5-11%)	15% (9.2-22%)	16 (9-25)	2700 (1900-3800)
Belarus	2016	Female	25 to 29	80% (67-88%)	20% (12-33%)	4.3 (2.4-7.1)	9.7% (7.5-12%)	17% (11-23%)	34 (21-52)	4800 (3400-6600)
Belarus	2016	Female	30 to 34	83% (70-91%)	17% (8.9-30%)	4.7 (2.7-7.4)	11% (8.2-13%)	19% (13-25%)	64 (41-95)	6500 (4700-8700)
Belarus	2016	Female	35 to 39	84% (73-92%)	16% (8.4-27%)	4.9 (2.7-7.8)	12% (8.8-15%)	21% (15-27%)	93 (58-140)	7500 (5300-10000)
Belarus	2016	Female	40 to 44	84% (73-92%)	16% (8-27%)	5 (2.8-8.1)	13% (9.2-16%)	22% (16-29%)	140 (88-200)	9300 (6400-13000)
Belarus	2016	Female	45 to 49	83% (69-91%)	17% (8.7-31%)	4.9 (2.7-7.9)	13% (9.1-17%)	22% (15-28%)	180 (120-270)	10000 (7100-14000)
Belarus	2016	Female	50 to 54	80% (67-89%)	20% (11-33%)	4.6 (2.6-7.5)	13% (8.7-17%)	20% (14-27%)	320 (200-470)	15000 (9900-21000)
Belarus	2016	Female	55 to 59	76% (60-87%)	24% (13-40%)	4.3 (2.4-6.6)	12% (7.7-17%)	18% (11-25%)	490 (280-730)	19000 (12000-27000)
Belarus	2016	Female	60 to 64	71% (55-85%)	29% (15-45%)	3.9 (2-6.5)	10% (4.9-16%)	14% (6.5-23%)	470 (210-800)	15000 (7300-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	2016	Female	65 to 69	67% (48-83%)	33% (17-52%)	3.3 (1.8-5.3)	7.6% (2.4-14%)	10% (2.6-19%)	410 (98-820)	11000 (3200-21000)
Belarus	2016	Female	70 to 74	60% (42-76%)	40% (24-58%)	2.6 (1.3-4.7)	5.2% (-0.31-12%)	6.6% (-0.97-17%)	270 (-40-720)	5800 (-350-14000)
Belarus	2016	Female	75 to 79	50% (33-69%)	50% (31-67%)	2.1 (0.98-3.9)	3.7% (-1.3-11%)	4.4% (-2.1-14%)	470 (-230-1500)	7700 (-2600-23000)
Belarus	2016	Female	80 to 84	43% (26-62%)	57% (38-74%)	1.9 (0.92-3.3)	3.4% (-0.84-10%)	3.9% (-1.4-12%)	440 (-160-1400)	5100 (-1300-15000)
Belarus	2016	Female	85 to 89	41% (25-58%)	59% (42-75%)	1.8 (0.86-3.3)	3.4% (-0.92-10%)	3.9% (-1.4-12%)	490 (-180-1500)	4000 (-1100-12000)
Belarus	2016	Female	90 to 94	42% (27-60%)	58% (40-73%)	2 (0.9-3.8)	3.6% (-1.4-11%)	4% (-2-13%)	250 (-130-830)	1400 (-550-4300)
Belarus	2016	Female	95 plus	46% (27-68%)	54% (32-73%)	2.2 (0.81-4.3)	3.9% (-2.2-13%)	4.3% (-2.9-15%)	53 (-35-190)	190 (-100-630)
Belarus	2016	Male	All Ages	82% (77-87%)	18% (13-23%)	6 (4.6-7.8)	17% (13-21%)	17% (11-22%)	10000 (6300-14000)	350000 (260000-460000)
Belarus	2016	Male	15 to 19	70% (53-83%)	30% (17-47%)	3.1 (1.7-5.1)	8.4% (5.3-13%)	16% (9-24%)	23 (12-37)	2100 (1300-3200)
Belarus	2016	Male	20 to 24	82% (70-91%)	18% (9.5-30%)	5.7 (3.3-9)	18% (12-23%)	26% (17-36%)	110 (65-150)	8600 (5700-12000)
Belarus	2016	Male	25 to 29	87% (79-93%)	13% (6.8-21%)	7.1 (4.3-11)	21% (16-26%)	30% (21-37%)	240 (160-330)	18000 (13000-24000)
Belarus	2016	Male	30 to 34	90% (82-96%)	9.6% (4.5-18%)	8 (5.2-12)	24% (19-28%)	31% (24-38%)	380 (270-510)	25000 (19000-33000)
Belarus	2016	Male	35 to 39	91% (85-96%)	8.5% (3.9-15%)	8.5 (5.3-13)	25% (19-30%)	32% (24-38%)	480 (340-640)	27000 (20000-36000)
Belarus	2016	Male	40 to 44	92% (85-96%)	8.2% (4-15%)	8.9 (5.5-13)	26% (20-31%)	33% (25-39%)	640 (440-860)	32000 (23000-42000)
Belarus	2016	Male	45 to 49	91% (82-96%)	8.7% (4-18%)	9.4 (6-14)	25% (20-30%)	32% (25-38%)	810 (580-1100)	36000 (26000-47000)
Belarus	2016	Male	50 to 54	90% (82-95%)	10% (4.6-18%)	9.1 (5.7-13)	23% (18-28%)	28% (22-35%)	1300 (920-1700)	50000 (36000-66000)
Belarus	2016	Male	55 to 59	88% (78-94%)	12% (5.7-22%)	8.1 (5.1-12)	20% (15-26%)	24% (18-31%)	1600 (1100-2200)	53000 (37000-71000)
Belarus	2016	Male	60 to 64	86% (76-93%)	14% (6.5-24%)	7.1 (4.3-11)	17% (11-23%)	19% (12-27%)	1500 (870-2100)	42000 (26000-59000)
Belarus	2016	Male	65 to 69	85% (71-93%)	15% (7-29%)	6 (3.7-9.1)	14% (7.4-20%)	16% (8.2-23%)	1100 (580-1700)	27000 (15000-40000)
Belarus	2016	Male	70 to 74	82% (69-90%)	18% (9.7-31%)	4.9 (2.9-7.7)	10% (3.1-16%)	11% (2.9-19%)	560 (140-980)	11000 (3300-18000)
Belarus	2016	Male	75 to 79	78% (64-89%)	22% (11-36%)	4.1 (2.2-6.8)	7.1% (0.26-14%)	7.8% (-0.31-16%)	660 (-27-1400)	10000 (340-21000)
Belarus	2016	Male	80 to 84	74% (58-86%)	26% (14-42%)	3.8 (2.2-5.9)	6% (-0.37-13%)	6.5% (-0.96-15%)	370 (-59-850)	4200 (-260-9300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belarus	2016	Male	85 to 89	71% (55-83%)	29% (17-45%)	3.5 (2-5.6)	4.9% (-1.3-12%)	5.2% (-2-14%)	250 (-93-640)	2100 (-530-5000)
Belarus	2016	Male	90 to 94	69% (55-82%)	31% (18-45%)	3.3 (1.8-5.5)	3.7% (-2.2-11%)	3.8% (-3.3-12%)	53 (-46-170)	320 (-190-910)
Belarus	2016	Male	95 plus	68% (49-84%)	32% (16-51%)	3.1 (1.4-5.6)	2.8% (-3.2-10%)	2.6% (-4.7-11%)	6.2 (-11-27)	27 (-31-99)
Estonia	1990	Female	All Ages	80% (77-83%)	20% (17-23%)	1 (0.79-1.3)	1.5% (0.16-3%)	-1.8% (-4-0.69%)	-180 (-390-66)	4200 (440-8600)
Estonia	1990	Female	15 to 19	78% (66-87%)	22% (13-34%)	0.89 (0.55-1.3)	3.9% (2.6-5.2%)	6.3% (3.3-9.5%)	1.9 (0.98-2.8)	250 (160-340)
Estonia	1990	Female	20 to 24	87% (80-92%)	13% (7.9-20%)	1.2 (0.75-1.8)	6.1% (4.4-8%)	6.6% (3.5-10%)	2.1 (1.1-3.2)	410 (290-590)
Estonia	1990	Female	25 to 29	89% (82-93%)	11% (7.1-18%)	1.2 (0.79-1.8)	6.7% (5.3-8.3%)	7.3% (4.4-11%)	3 (1.8-4.4)	560 (400-760)
Estonia	1990	Female	30 to 34	89% (83-93%)	11% (6.5-17%)	1.3 (0.9-1.9)	7.4% (6-9.1%)	10% (7.2-14%)	5.4 (3.8-7.2)	730 (550-950)
Estonia	1990	Female	35 to 39	90% (85-94%)	10% (6.5-15%)	1.5 (0.98-2.1)	8.9% (7.3-11%)	13% (10-17%)	11 (8.2-14)	1000 (810-1300)
Estonia	1990	Female	40 to 44	90% (85-94%)	9.9% (6.1-15%)	1.6 (1.1-2.3)	8.8% (7.2-11%)	12% (9.1-16%)	15 (11-19)	1100 (870-1400)
Estonia	1990	Female	45 to 49	90% (86-93%)	9.8% (7-14%)	1.6 (1.1-2.3)	7.9% (6.1-9.9%)	11% (7.9-15%)	17 (12-23)	1100 (800-1300)
Estonia	1990	Female	50 to 54	88% (84-91%)	12% (8.8-16%)	1.5 (1-2.2)	5.9% (4.2-7.9%)	7.6% (4.7-11%)	21 (13-31)	1100 (790-1500)
Estonia	1990	Female	55 to 59	85% (80-88%)	15% (12-20%)	1.5 (1-2.1)	3.7% (1.9-5.7%)	4.4% (1.7-7.4%)	17 (6.4-29)	760 (380-1200)
Estonia	1990	Female	60 to 64	82% (76-86%)	18% (14-24%)	1.4 (0.94-2)	0.91% (-1.1-3.3%)	0.64% (-2.1-4%)	4.2 (-14-27)	250 (-290-900)
Estonia	1990	Female	65 to 69	79% (72-84%)	21% (16-28%)	1.2 (0.8-1.6)	-1.3% (-3.4-1.1%)	-2.1% (-5.1-0.99%)	-19 (-45-8.9)	-370 (-970-310)
Estonia	1990	Female	70 to 74	71% (63-78%)	29% (22-37%)	0.96 (0.59-1.4)	-2.5% (-4.9--0.16%)	-3.6% (-6.5--0.42%)	-32 (-59--3.7)	-560 (-1100--35)
Estonia	1990	Female	75 to 79	59% (48-70%)	41% (30-52%)	0.78 (0.46-1.2)	-2.9% (-5.1--0.67%)	-3.8% (-6.6--0.99%)	-62 (-110--17)	-850 (-1500--200)
Estonia	1990	Female	80 to 84	51% (40-62%)	49% (38-60%)	0.69 (0.41-1.1)	-3% (-5.4--0.92%)	-3.8% (-6.6--1.2%)	-73 (-130--23)	-750 (-1300--220)
Estonia	1990	Female	85 to 89	50% (38-62%)	50% (38-62%)	0.69 (0.4-1.1)	-3% (-5.5--0.89%)	-3.7% (-6.6--1.2%)	-55 (-100--17)	-400 (-720--120)
Estonia	1990	Female	90 to 94	52% (40-64%)	48% (36-60%)	0.76 (0.41-1.2)	-3% (-5.4--0.68%)	-3.8% (-6.7--0.99%)	-26 (-45--6.8)	-120 (-220--28)
Estonia	1990	Female	95 plus	56% (41-70%)	44% (30-59%)	0.84 (0.41-1.4)	-3.2% (-6--0.51%)	-4% (-7.5--0.87%)	-7.9 (-15--1.7)	-25 (-47--3.8)
Estonia	1990	Male	All Ages	87% (83-90%)	13% (9.8-17%)	2.4 (1.9-3)	9% (6.4-12%)	6.7% (3.2-10%)	620 (300-960)	31000 (22000-41000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	1990	Male	15 to 19	78% (66-88%)	22% (12-34%)	1.5 (0.88-2.2)	7.1% (4.5-10%)	9.5% (5.5-14%)	8.5 (4.9-13)	720 (450-1000)
Estonia	1990	Male	20 to 24	88% (80-93%)	12% (6.7-20%)	2.6 (1.6-3.8)	13% (9.1-18%)	16% (10-22%)	21 (14-30)	1700 (1200-2300)
Estonia	1990	Male	25 to 29	91% (85-95%)	8.8% (4.8-15%)	2.9 (1.9-4.2)	15% (11-19%)	18% (12-24%)	29 (19-40)	2300 (1600-3000)
Estonia	1990	Male	30 to 34	93% (87-97%)	6.7% (3.4-13%)	3.2 (2.2-4.6)	16% (12-20%)	19% (14-25%)	42 (31-56)	2900 (2200-3700)
Estonia	1990	Male	35 to 39	94% (89-97%)	6% (2.9-11%)	3.5 (2.3-5)	17% (13-22%)	21% (15-27%)	58 (42-77)	3500 (2700-4500)
Estonia	1990	Male	40 to 44	95% (90-98%)	5.3% (2.4-9.9%)	3.7 (2.5-5.3)	17% (12-22%)	20% (14-26%)	72 (50-96)	3800 (2800-4900)
Estonia	1990	Male	45 to 49	95% (91-98%)	4.8% (2.5-9.1%)	4.1 (2.7-5.8)	16% (11-21%)	19% (13-25%)	83 (57-110)	3800 (2700-5000)
Estonia	1990	Male	50 to 54	95% (91-97%)	4.9% (2.8-8.8%)	4.3 (2.8-6)	14% (9.2-19%)	16% (10-22%)	120 (72-160)	4600 (3000-6400)
Estonia	1990	Male	55 to 59	94% (91-97%)	5.6% (3.4-8.9%)	4.1 (2.8-5.6)	11% (6.1-15%)	12% (6.5-17%)	110 (58-160)	3700 (2200-5400)
Estonia	1990	Male	60 to 64	93% (90-96%)	6.5% (4.1-10%)	3.7 (2.4-5.2)	7.9% (2.8-13%)	7.9% (2.7-14%)	93 (31-160)	2800 (1100-4800)
Estonia	1990	Male	65 to 69	92% (88-95%)	7.9% (4.9-12%)	3.1 (2.1-4.2)	3.9% (0.043-8.3%)	3.9% (-0.62-9.2%)	38 (-6-89)	1000 (11-2200)
Estonia	1990	Male	70 to 74	90% (86-93%)	9.9% (6.8-14%)	2.5 (1.6-3.6)	1.3% (-2.4-5.7%)	0.98% (-3.3-6%)	8.2 (-28-51)	230 (-440-1000)
Estonia	1990	Male	75 to 79	87% (80-92%)	13% (7.6-20%)	2.1 (1.3-3.1)	-0.43% (-4-3.9%)	-0.97% (-5.1-4%)	-10 (-53-42)	-77 (-690-680)
Estonia	1990	Male	80 to 84	84% (75-90%)	16% (9.6-25%)	1.8 (1.2-2.7)	-1.7% (-5-2.1%)	-2.4% (-6.4-1.9%)	-21 (-58-17)	-180 (-560-230)
Estonia	1990	Male	85 to 89	82% (73-88%)	18% (12-27%)	1.7 (1.1-2.5)	-2.3% (-5.6-1.2%)	-3.1% (-7-0.89%)	-16 (-37-4.8)	-110 (-260-55)
Estonia	1990	Male	90 to 94	80% (71-88%)	20% (12-29%)	1.6 (0.93-2.5)	-2.7% (-5.9-0.55%)	-3.6% (-7.4-0.19%)	-5.6 (-11-0.29)	-25 (-54-5.1)
Estonia	1990	Male	95 plus	79% (67-89%)	21% (11-33%)	1.5 (0.73-2.6)	-2.7% (-5.8-0.6%)	-3.7% (-7.5-0.16%)	-1.3 (-2.6-0.055)	-3.7 (-8-0.84)
Estonia	1995	Female	All Ages	80% (77-83%)	20% (17-23%)	1.2 (0.99-1.4)	2.3% (0.87-3.9%)	-0.93% (-3.2-1.7%)	-94 (-330-170)	6500 (2300-11000)
Estonia	1995	Female	15 to 19	78% (66-88%)	22% (12-34%)	1 (0.65-1.4)	4.3% (3-5.7%)	7.4% (4.2-11%)	2.1 (1.2-3.1)	260 (170-350)
Estonia	1995	Female	20 to 24	87% (79-92%)	13% (7.8-21%)	1.4 (0.89-2)	6.6% (4.9-8.4%)	7.6% (4.3-11%)	2.9 (1.7-4.4)	460 (320-620)
Estonia	1995	Female	25 to 29	89% (82-93%)	11% (7.1-18%)	1.4 (0.94-2)	7.4% (5.8-9.2%)	8.7% (5.6-12%)	3.5 (2.2-4.9)	530 (390-700)
Estonia	1995	Female	30 to 34	90% (84-93%)	10% (6.5-16%)	1.5 (1.1-2.1)	8.5% (6.9-10%)	12% (9.4-16%)	6.9 (5.3-9)	760 (600-960)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	1995	Female	35 to 39	90% (85-94%)	10% (6.4-15%)	1.7 (1.2-2.3)	11% (8.6-13%)	15% (12-19%)	17 (13-21)	1300 (1100-1600)
Estonia	1995	Female	40 to 44	90% (85-94%)	9.8% (6.2-15%)	1.8 (1.3-2.5)	10% (8.3-12%)	14% (11-18%)	23 (17-28)	1500 (1200-1800)
Estonia	1995	Female	45 to 49	90% (86-93%)	9.7% (6.9-14%)	1.8 (1.3-2.5)	9.2% (7.2-11%)	13% (9.4-16%)	28 (21-36)	1500 (1200-1900)
Estonia	1995	Female	50 to 54	89% (84-91%)	11% (8.6-16%)	1.8 (1.2-2.3)	7.2% (5.2-9.5%)	9.4% (6.4-13%)	26 (18-35)	1200 (880-1600)
Estonia	1995	Female	55 to 59	85% (80-89%)	15% (11-20%)	1.7 (1.2-2.3)	4.8% (2.7-7%)	5.9% (2.8-9.1%)	28 (13-44)	1100 (630-1700)
Estonia	1995	Female	60 to 64	82% (76-86%)	18% (14-24%)	1.6 (1.1-2.2)	1.8% (-0.42-4.4%)	1.9% (-1.2-5.5%)	12 (-7.5-35)	460 (-110-1100)
Estonia	1995	Female	65 to 69	79% (72-84%)	21% (16-28%)	1.4 (0.96-1.8)	-0.66% (-3.1-1.9%)	-1.3% (-4.3-2.1%)	-13 (-44-21)	-210 (-940-610)
Estonia	1995	Female	70 to 74	72% (64-78%)	28% (22-36%)	1.1 (0.71-1.6)	-2.1% (-4.4-0.5%)	-2.9% (-5.9-0.36%)	-38 (-77-4.9)	-650 (-1400-160)
Estonia	1995	Female	75 to 79	60% (49-70%)	40% (30-51%)	0.9 (0.55-1.4)	-2.5% (-4.8--0.22%)	-3.3% (-6.1--0.44%)	-39 (-73--5.2)	-530 (-1000--49)
Estonia	1995	Female	80 to 84	52% (41-63%)	48% (37-59%)	0.8 (0.5-1.2)	-2.7% (-5-0.37%)	-3.4% (-6.2--0.63%)	-64 (-120--12)	-650 (-1200--87)
Estonia	1995	Female	85 to 89	51% (39-63%)	49% (37-61%)	0.8 (0.48-1.2)	-2.7% (-5.1--0.41%)	-3.4% (-6.3--0.66%)	-56 (-100--11)	-390 (-730--61)
Estonia	1995	Female	90 to 94	53% (41-65%)	47% (35-59%)	0.89 (0.49-1.4)	-2.8% (-5.3--0.31%)	-3.5% (-6.5--0.53%)	-27 (-50--4.1)	-130 (-250--14)
Estonia	1995	Female	95 plus	57% (41-70%)	43% (30-59%)	1 (0.5-1.7)	-2.9% (-6-0.14%)	-3.7% (-7.3--0.071%)	-7.3 (-14-0.14)	-23 (-47-1.1)
Estonia	1995	Male	All Ages	87% (84-91%)	13% (9.4-16%)	2.8 (2.3-3.3)	12% (9-15%)	9.6% (6-13%)	1100 (660-1500)	46000 (35000-57000)
Estonia	1995	Male	15 to 19	79% (67-88%)	21% (12-33%)	1.7 (1.1-2.5)	8.1% (5.4-12%)	11% (6.6-16%)	9.7 (5.8-14)	800 (510-1100)
Estonia	1995	Male	20 to 24	88% (80-93%)	12% (6.6-20%)	3 (2-4.2)	15% (11-20%)	18% (12-24%)	29 (19-39)	2200 (1500-2900)
Estonia	1995	Male	25 to 29	91% (85-95%)	8.8% (4.7-15%)	3.3 (2.2-4.6)	17% (13-22%)	20% (14-27%)	36 (25-48)	2600 (1900-3300)
Estonia	1995	Male	30 to 34	93% (87-97%)	6.7% (3.2-13%)	3.6 (2.5-5)	19% (14-23%)	22% (16-28%)	56 (42-73)	3600 (2700-4600)
Estonia	1995	Male	35 to 39	94% (90-97%)	5.8% (2.9-10%)	4 (2.7-5.6)	21% (16-26%)	24% (18-31%)	96 (73-120)	5400 (4200-6700)
Estonia	1995	Male	40 to 44	95% (90-98%)	5.2% (2.5-9.8%)	4.3 (3-5.8)	20% (16-26%)	23% (17-29%)	130 (93-160)	6300 (4700-7800)
Estonia	1995	Male	45 to 49	95% (91-98%)	4.8% (2.4-9.2%)	4.7 (3.3-6.5)	20% (15-25%)	22% (16-28%)	150 (110-190)	6500 (4900-8300)
Estonia	1995	Male	50 to 54	95% (91-97%)	4.9% (2.8-8.6%)	4.9 (3.5-6.6)	18% (12-23%)	20% (13-26%)	160 (110-210)	6000 (4200-7800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	1995	Male	55 to 59	95% (91-97%)	5.5% (3.4-8.6%)	4.7 (3.4-6.1)	13% (8.5-19%)	15% (9.1-21%)	180 (110-250)	5900 (3700-8200)
Estonia	1995	Male	60 to 64	94% (90-96%)	6.4% (4-10%)	4.2 (2.9-5.7)	9.5% (4.4-15%)	10% (4.4-17%)	140 (60-230)	4000 (1800-6500)
Estonia	1995	Male	65 to 69	92% (88-95%)	7.7% (4.9-12%)	3.5 (2.6-4.6)	5.6% (1.3-10%)	5.8% (0.92-11%)	87 (14-170)	2200 (500-4000)
Estonia	1995	Male	70 to 74	90% (86-93%)	9.8% (6.9-14%)	2.9 (2-4)	2.6% (-1.2-7.6%)	2.5% (-1.8-8.1%)	29 (-21-92)	630 (-280-1800)
Estonia	1995	Male	75 to 79	88% (81-92%)	12% (7.6-19%)	2.4 (1.5-3.5)	0.73% (-3-5.6%)	0.39% (-3.9-5.9%)	3.3 (-33-50)	100 (-420-780)
Estonia	1995	Male	80 to 84	85% (76-90%)	15% (9.5-24%)	2.1 (1.4-3.1)	-0.73% (-4.2-3.3%)	-1.3% (-5.5-3.4%)	-12 (-49-31)	-80 (-470-360)
Estonia	1995	Male	85 to 89	82% (73-89%)	18% (11-27%)	1.9 (1.3-2.8)	-1.7% (-5.2-2.1%)	-2.4% (-6.5-2%)	-14 (-37-11)	-82 (-250-100)
Estonia	1995	Male	90 to 94	81% (71-88%)	19% (12-29%)	1.8 (1.1-2.7)	-2.2% (-5.7-1.4%)	-3% (-7.1-1.1%)	-6 (-14-2.2)	-25 (-66-16)
Estonia	1995	Male	95 plus	80% (67-89%)	20% (11-33%)	1.7 (0.86-2.8)	-2.2% (-5.6-1.5%)	-3.2% (-7.3-1.3%)	-1 (-2.3-0.4)	-2.8 (-7-2)
Estonia	2000	Female	All Ages	79% (76-82%)	21% (18-24%)	1.5 (1.3-1.8)	3.6% (2.3-5.1%)	0.87% (-1.5-3.3%)	82 (-140-310)	9400 (5900-14000)
Estonia	2000	Female	15 to 19	79% (68-87%)	21% (13-32%)	1.3 (0.89-1.9)	4.3% (3.1-5.7%)	8.4% (5.2-12%)	1.8 (1.1-2.6)	230 (160-310)
Estonia	2000	Female	20 to 24	87% (81-92%)	13% (7.9-19%)	1.8 (1.2-2.6)	7% (5.2-8.9%)	8.9% (5.4-13%)	2.3 (1.4-3.4)	420 (290-570)
Estonia	2000	Female	25 to 29	89% (82-93%)	11% (7.2-18%)	1.8 (1.2-2.5)	7.8% (6.1-9.6%)	10% (6.6-13%)	3 (2-4.1)	520 (390-680)
Estonia	2000	Female	30 to 34	89% (83-93%)	11% (6.8-17%)	1.9 (1.4-2.6)	9.1% (7.4-11%)	14% (11-17%)	6.6 (5-8.3)	720 (560-900)
Estonia	2000	Female	35 to 39	90% (85-93%)	10% (6.9-15%)	2.2 (1.6-2.9)	11% (9-13%)	17% (14-20%)	14 (11-17)	1100 (890-1300)
Estonia	2000	Female	40 to 44	90% (86-94%)	9.7% (6.2-14%)	2.4 (1.8-3.2)	11% (9.1-13%)	17% (14-20%)	21 (17-26)	1400 (1100-1700)
Estonia	2000	Female	45 to 49	91% (87-93%)	9.5% (7-13%)	2.4 (1.7-3.2)	11% (8.4-13%)	15% (12-18%)	32 (25-39)	1700 (1400-2100)
Estonia	2000	Female	50 to 54	88% (84-91%)	12% (9.2-16%)	2.2 (1.6-3)	9% (7-11%)	12% (9.5-16%)	35 (26-44)	1600 (1200-2000)
Estonia	2000	Female	55 to 59	83% (80-87%)	17% (13-20%)	2.1 (1.5-2.8)	7% (4.9-9.2%)	9.3% (6.3-12%)	34 (22-45)	1300 (900-1700)
Estonia	2000	Female	60 to 64	79% (72-84%)	21% (16-28%)	1.9 (1.3-2.7)	3.9% (1.7-6.6%)	5.1% (1.9-8.6%)	28 (11-48)	930 (400-1500)
Estonia	2000	Female	65 to 69	74% (66-81%)	26% (19-34%)	1.5 (1.1-2.1)	1.4% (-0.77-3.7%)	1.4% (-1.5-4.6%)	11 (-11-35)	350 (-200-970)
Estonia	2000	Female	70 to 74	64% (54-73%)	36% (27-46%)	1.2 (0.73-1.7)	-0.2% (-2.4-2.1%)	-0.56% (-3.3-2.4%)	-7 (-43-30)	-61 (-730-660)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	2000	Female	75 to 79	53% (43-64%)	47% (36-57%)	0.96 (0.59-1.5)	-0.88% (-2.9-1.3%)	-1.3% (-4-1.4%)	-21 (-63-22)	-250 (-860-390)
Estonia	2000	Female	80 to 84	49% (37-59%)	51% (41-63%)	0.94 (0.6-1.4)	-1.3% (-3.4-0.83%)	-1.8% (-4.5-0.84%)	-23 (-57-11)	-220 (-570-150)
Estonia	2000	Female	85 to 89	49% (37-61%)	51% (39-63%)	1 (0.6-1.5)	-1.5% (-4-1%)	-2.1% (-5.1-0.93%)	-32 (-79-14)	-210 (-550-140)
Estonia	2000	Female	90 to 94	51% (40-63%)	49% (37-60%)	1.1 (0.62-1.7)	-1.6% (-4.2-1.2%)	-2.1% (-5.4-1.2%)	-18 (-46-11)	-81 (-220-63)
Estonia	2000	Female	95 plus	55% (40-69%)	45% (31-60%)	1.2 (0.63-2.1)	-1.6% (-4.8-1.9%)	-2.2% (-6-1.9%)	-5.4 (-15-4.7)	-15 (-46-18)
Estonia	2000	Male	All Ages	89% (86-92%)	11% (8.3-14%)	3.9 (3.4-4.6)	15% (12-18%)	13% (9.5-17%)	1200 (880-1600)	48000 (39000-57000)
Estonia	2000	Male	15 to 19	80% (70-89%)	20% (11-30%)	2.3 (1.5-3.3)	8.6% (5.8-12%)	13% (7.8-19%)	7.9 (4.8-11)	680 (440-920)
Estonia	2000	Male	20 to 24	89% (82-94%)	11% (5.9-18%)	4 (2.7-5.6)	17% (13-22%)	21% (15-28%)	25 (18-34)	2000 (1500-2600)
Estonia	2000	Male	25 to 29	92% (88-96%)	7.7% (4.2-12%)	4.4 (3-5.9)	19% (14-23%)	24% (17-31%)	31 (23-40)	2300 (1800-2900)
Estonia	2000	Male	30 to 34	94% (90-97%)	5.8% (3-10%)	4.9 (3.4-6.7)	21% (17-26%)	27% (21-34%)	43 (33-54)	2900 (2300-3500)
Estonia	2000	Male	35 to 39	95% (91-97%)	5% (2.7-8.6%)	5.3 (3.7-7.4)	24% (19-29%)	29% (23-36%)	73 (56-89)	4200 (3400-5100)
Estonia	2000	Male	40 to 44	95% (91-98%)	4.7% (2.3-8.5%)	5.7 (4.1-7.7)	25% (20-29%)	29% (23-35%)	120 (96-150)	6100 (4900-7200)
Estonia	2000	Male	45 to 49	96% (92-98%)	4.4% (2.3-8.1%)	6.2 (4.5-8.2)	24% (19-28%)	28% (22-33%)	160 (120-190)	6900 (5500-8200)
Estonia	2000	Male	50 to 54	95% (93-97%)	4.5% (2.6-7.4%)	6.5 (4.6-8.7)	22% (17-26%)	25% (19-30%)	170 (130-200)	6600 (5100-7800)
Estonia	2000	Male	55 to 59	95% (93-97%)	4.9% (3.1-7.4%)	6.3 (4.6-8.2)	18% (14-23%)	21% (15-26%)	160 (110-200)	5400 (3900-6600)
Estonia	2000	Male	60 to 64	94% (92-96%)	5.7% (3.9-8.4%)	5.7 (4-7.7)	14% (8.6-20%)	16% (9.3-22%)	180 (110-250)	5300 (3100-7100)
Estonia	2000	Male	65 to 69	93% (89-95%)	7.3% (4.9-11%)	4.7 (3.5-6.1)	9.9% (4.9-15%)	11% (4.9-17%)	130 (60-210)	3200 (1600-4900)
Estonia	2000	Male	70 to 74	91% (87-93%)	9.5% (6.8-13%)	3.8 (2.6-5.3)	6.2% (1.2-12%)	6.6% (0.8-13%)	89 (11-180)	1800 (360-3400)
Estonia	2000	Male	75 to 79	88% (81-92%)	12% (7.7-19%)	3.1 (2-4.6)	3.6% (-0.88-9.5%)	3.7% (-1.6-11%)	34 (-15-97)	570 (-130-1500)
Estonia	2000	Male	80 to 84	84% (75-91%)	16% (9.5-25%)	2.7 (1.8-3.9)	1.9% (-2.1-6.9%)	1.6% (-3.1-7.4%)	9.7 (-18-45)	140 (-160-520)
Estonia	2000	Male	85 to 89	81% (72-88%)	19% (12-28%)	2.5 (1.6-3.5)	0.56% (-3.2-5.4%)	0.087% (-4.4-5.7%)	0.46 (-23-31)	26 (-150-250)
Estonia	2000	Male	90 to 94	80% (70-88%)	20% (12-30%)	2.3 (1.4-3.5)	-0.29% (-4-4.4%)	-0.94% (-5.5-4.4%)	-2 (-12-9.6)	-3.7 (-52-57)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	2000	Male	95 plus	79% (65-89%)	21% (11-35%)	2.1 (1.1-3.4)	-0.66% (-4.4-4.6%)	-1.5% (-6.1-4.7%)	-0.7 (-2.9-2.3)	-1.3 (-8.5-8.5)
Estonia	2005	Female	All Ages	81% (79-84%)	19% (16-21%)	2.1 (1.8-2.3)	4.8% (3.4-6.3%)	2.6% (0.13-5.2%)	230 (12-450)	11000 (8000-15000)
Estonia	2005	Female	15 to 19	79% (69-88%)	21% (12-31%)	1.7 (1.1-2.4)	4.5% (3.2-5.8%)	9.8% (6.3-14%)	1.6 (1-2.3)	220 (160-300)
Estonia	2005	Female	20 to 24	88% (82-93%)	12% (7.4-18%)	2.3 (1.6-3.3)	7.3% (5.5-9.3%)	9.8% (6.2-14%)	2.5 (1.6-3.6)	430 (300-590)
Estonia	2005	Female	25 to 29	90% (85-93%)	10% (6.5-15%)	2.3 (1.6-3.2)	8% (6.4-9.7%)	11% (7.4-14%)	2.4 (1.7-3.2)	480 (350-640)
Estonia	2005	Female	30 to 34	90% (86-94%)	9.6% (6.2-14%)	2.5 (1.8-3.4)	9.5% (7.8-11%)	16% (13-20%)	6 (4.7-7.4)	700 (540-870)
Estonia	2005	Female	35 to 39	91% (87-94%)	9.3% (6.1-13%)	2.8 (2-3.7)	11% (9-13%)	19% (16-23%)	10 (8.3-12)	870 (720-1100)
Estonia	2005	Female	40 to 44	91% (87-94%)	8.8% (5.7-13%)	3 (2.2-3.9)	12% (9.7-14%)	20% (17-23%)	19 (16-22)	1300 (1000-1500)
Estonia	2005	Female	45 to 49	91% (89-93%)	8.8% (6.6-11%)	3.1 (2.2-4)	11% (9.3-13%)	18% (15-21%)	29 (24-35)	1600 (1300-1900)
Estonia	2005	Female	50 to 54	89% (87-91%)	11% (8.8-13%)	2.9 (2.1-3.8)	11% (8.6-13%)	16% (13-19%)	40 (32-49)	1800 (1500-2200)
Estonia	2005	Female	55 to 59	86% (83-88%)	14% (12-17%)	2.8 (2-3.7)	9.3% (7-12%)	13% (10-17%)	46 (35-57)	1700 (1300-2200)
Estonia	2005	Female	60 to 64	82% (80-85%)	18% (15-20%)	2.6 (1.8-3.5)	6.3% (3.9-9.1%)	8.8% (5.5-13%)	34 (21-49)	1100 (670-1600)
Estonia	2005	Female	65 to 69	79% (76-82%)	21% (18-24%)	2.2 (1.7-2.9)	3.5% (1.1-6%)	4.5% (1.2-8.1%)	30 (7.7-54)	830 (260-1400)
Estonia	2005	Female	70 to 74	73% (69-76%)	27% (24-31%)	1.8 (1.3-2.5)	1.4% (-1.4-4.2%)	1.6% (-2.1-5.2%)	15 (-20-49)	350 (-340-1100)
Estonia	2005	Female	75 to 79	61% (55-66%)	39% (34-45%)	1.5 (0.99-2.1)	0.4% (-2.2-3.3%)	0.25% (-3.1-3.9%)	3.7 (-47-58)	110 (-640-950)
Estonia	2005	Female	80 to 84	51% (42-60%)	49% (40-58%)	1.3 (0.84-1.8)	0.05% (-2.3-2.5%)	-0.17% (-3.2-2.9%)	-2.9 (-56-50)	12 (-550-590)
Estonia	2005	Female	85 to 89	47% (37-57%)	53% (43-63%)	1.2 (0.72-1.7)	-0.15% (-2.4-2.5%)	-0.43% (-3.2-2.8%)	-4.8 (-36-32)	-15 (-250-260)
Estonia	2005	Female	90 to 94	49% (39-59%)	51% (41-61%)	1.3 (0.75-2)	-0.12% (-2.9-3.2%)	-0.41% (-3.8-3.6%)	-3.7 (-34-32)	-6.4 (-160-170)
Estonia	2005	Female	95 plus	53% (39-66%)	47% (34-61%)	1.4 (0.74-2.4)	0.021% (-3.2-4.4%)	-0.31% (-4.2-4.8%)	-0.97 (-13-15)	0.28 (-40-54)
Estonia	2005	Male	All Ages	91% (88-93%)	9.5% (7.3-12%)	5.1 (4.5-5.8)	17% (15-20%)	16% (13-20%)	1400 (1100-1800)	51000 (43000-58000)
Estonia	2005	Male	15 to 19	83% (75-90%)	17% (10-25%)	3 (2-4.2)	9.6% (6.6-13%)	15% (9.7-22%)	8.5 (5.4-12)	720 (490-970)
Estonia	2005	Male	20 to 24	91% (85-95%)	9.4% (5.4-15%)	5 (3.5-7)	18% (13-22%)	23% (17-30%)	24 (17-32)	1900 (1400-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	2005	Male	25 to 29	93% (89-96%)	6.8% (4-11%)	5.6 (3.9-7.5)	19% (15-23%)	24% (18-31%)	30 (23-38)	2200 (1800-2800)
Estonia	2005	Male	30 to 34	95% (91-97%)	5.2% (2.8-8.9%)	6.1 (4.5-8.5)	22% (18-26%)	29% (24-35%)	43 (35-52)	2900 (2300-3400)
Estonia	2005	Male	35 to 39	95% (92-98%)	4.7% (2.3-8.4%)	6.7 (4.7-9.1)	25% (21-29%)	32% (26-38%)	58 (47-69)	3400 (2800-4000)
Estonia	2005	Male	40 to 44	96% (92-98%)	4.3% (2.1-7.9%)	7.2 (5.3-9.4)	27% (23-31%)	34% (28-39%)	100 (85-120)	5200 (4400-5900)
Estonia	2005	Male	45 to 49	96% (93-98%)	3.9% (2-7%)	8 (5.9-10)	27% (22-30%)	32% (27-37%)	150 (130-170)	6600 (5600-7700)
Estonia	2005	Male	50 to 54	96% (94-98%)	3.8% (2.3-6.3%)	8.4 (6.1-11)	25% (21-29%)	30% (25-34%)	190 (160-220)	7300 (6200-8400)
Estonia	2005	Male	55 to 59	96% (94-97%)	4.3% (2.9-6.4%)	8 (5.9-10)	22% (18-26%)	25% (21-30%)	190 (160-230)	6400 (5300-7600)
Estonia	2005	Male	60 to 64	95% (92-96%)	5.2% (3.6-7.6%)	7.1 (5.1-9.6)	18% (13-22%)	20% (15-26%)	180 (130-220)	5000 (3700-6300)
Estonia	2005	Male	65 to 69	94% (91-96%)	6.5% (4.4-9.3%)	6 (4.5-7.7)	14% (8.9-18%)	15% (9.9-20%)	180 (110-240)	4400 (2900-5700)
Estonia	2005	Male	70 to 74	92% (89-94%)	8.1% (6.1-11%)	5 (3.6-6.6)	9.9% (4.5-15%)	11% (4.7-17%)	130 (57-200)	2600 (1200-3900)
Estonia	2005	Male	75 to 79	90% (85-93%)	10% (6.7-15%)	4.2 (2.8-5.9)	6.9% (1.4-13%)	7.5% (1.1-14%)	91 (13-170)	1400 (290-2600)
Estonia	2005	Male	80 to 84	87% (80-92%)	13% (8.5-20%)	3.6 (2.5-5)	4.9% (0.02-11%)	5.2% (-0.63-12%)	40 (-4.9-91)	480 (1.9-1000)
Estonia	2005	Male	85 to 89	84% (76-90%)	16% (10-24%)	3.3 (2.2-4.7)	3.3% (-1.4-9%)	3.3% (-2.3-9.9%)	13 (-9.2-40)	120 (-49-320)
Estonia	2005	Male	90 to 94	83% (74-89%)	17% (11-26%)	3.1 (2-4.6)	2.1% (-2.9-8.3%)	1.8% (-4.1-9.2%)	4.5 (-10-23)	32 (-43-120)
Estonia	2005	Male	95 plus	81% (69-90%)	19% (9.8-31%)	2.8 (1.5-4.7)	1.4% (-3.7-8.6%)	0.92% (-5.2-9.5%)	0.56 (-3.2-5.8)	3.5 (-9.1-21)
Estonia	2010	Female	All Ages	80% (78-82%)	20% (18-22%)	2.3 (1.9-2.7)	5.2% (3.8-6.8%)	3.9% (0.97-6.8%)	300 (75-520)	11000 (7800-15000)
Estonia	2010	Female	15 to 19	78% (67-87%)	22% (13-33%)	1.8 (1.2-2.6)	4.2% (3.1-5.5%)	10% (6.5-14%)	0.9 (0.58-1.3)	150 (110-200)
Estonia	2010	Female	20 to 24	88% (82-92%)	12% (7.8-18%)	2.6 (1.7-3.7)	7.1% (5.2-9.1%)	9.4% (6.2-13%)	1.4 (0.93-2)	370 (250-510)
Estonia	2010	Female	25 to 29	89% (84-93%)	11% (6.9-16%)	2.6 (1.8-3.7)	7.9% (6.4-9.7%)	11% (7.8-14%)	2.4 (1.7-3.1)	470 (340-620)
Estonia	2010	Female	30 to 34	90% (85-94%)	9.9% (6.4-15%)	2.8 (2-3.8)	8.8% (7.1-11%)	16% (13-20%)	4.1 (3.3-5)	550 (420-700)
Estonia	2010	Female	35 to 39	90% (87-94%)	9.7% (6.3-13%)	3.2 (2.2-4.3)	9.6% (8-11%)	19% (16-22%)	7.4 (6.1-9)	730 (580-900)
Estonia	2010	Female	40 to 44	91% (87-94%)	9.2% (6.1-13%)	3.4 (2.5-4.6)	11% (8.8-12%)	20% (17-24%)	12 (10-14)	900 (730-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	2010	Female	45 to 49	91% (88-93%)	9.4% (7-12%)	3.5 (2.5-4.7)	11% (8.9-13%)	19% (16-22%)	20 (17-23)	1200 (980-1400)
Estonia	2010	Female	50 to 54	89% (86-91%)	11% (9.1-14%)	3.3 (2.3-4.4)	10% (8.3-12%)	17% (14-20%)	29 (24-35)	1400 (1200-1700)
Estonia	2010	Female	55 to 59	85% (83-88%)	15% (12-17%)	3.2 (2.3-4.2)	9.3% (7.2-12%)	15% (12-18%)	36 (29-44)	1400 (1100-1800)
Estonia	2010	Female	60 to 64	82% (79-84%)	18% (16-21%)	2.9 (2-4)	7.4% (5.1-10%)	11% (7.8-15%)	40 (27-53)	1300 (880-1800)
Estonia	2010	Female	65 to 69	79% (75-82%)	21% (18-25%)	2.5 (1.9-3.3)	4.9% (2.7-7.3%)	7.1% (3.9-10%)	32 (17-48)	860 (460-1300)
Estonia	2010	Female	70 to 74	72% (68-75%)	28% (25-32%)	2 (1.4-2.8)	2.9% (0.21-5.5%)	3.9% (0.012-7.7%)	31 (0.099-61)	680 (52-1300)
Estonia	2010	Female	75 to 79	60% (55-65%)	40% (35-45%)	1.7 (1.1-2.4)	1.9% (-0.81-4.7%)	2.3% (-1.5-6.2%)	27 (-17-71)	450 (-200-1100)
Estonia	2010	Female	80 to 84	50% (41-59%)	50% (41-59%)	1.4 (0.93-2)	1.4% (-1-4.1%)	1.6% (-1.6-5%)	27 (-27-83)	340 (-250-960)
Estonia	2010	Female	85 to 89	46% (36-55%)	54% (45-64%)	1.3 (0.78-2)	1.1% (-1.3-3.8%)	1.2% (-2-4.5%)	18 (-31-70)	170 (-200-570)
Estonia	2010	Female	90 to 94	48% (37-58%)	52% (42-63%)	1.4 (0.83-2.2)	1.2% (-1.8-4.9%)	1.2% (-2.5-5.6%)	8.1 (-17-38)	52 (-76-210)
Estonia	2010	Female	95 plus	52% (38-65%)	48% (35-62%)	1.6 (0.81-2.6)	1.4% (-2.4-6.4%)	1.4% (-3.3-7.4%)	4.7 (-11-25)	19 (-33-88)
Estonia	2010	Male	All Ages	90% (88-93%)	9.7% (7.4-12%)	5.9 (5-6.9)	18% (15-20%)	18% (13-22%)	1300 (1000-1600)	43000 (37000-50000)
Estonia	2010	Male	15 to 19	82% (74-89%)	18% (11-26%)	3.4 (2.2-4.8)	8.4% (5.8-11%)	16% (10-23%)	3.9 (2.5-5.5)	370 (260-490)
Estonia	2010	Male	20 to 24	90% (84-94%)	9.8% (5.6-16%)	5.7 (3.7-7.9)	16% (12-20%)	23% (17-30%)	16 (11-21)	1400 (1000-1700)
Estonia	2010	Male	25 to 29	93% (89-96%)	7.1% (4.1-11%)	6.3 (4.3-8.7)	18% (14-21%)	23% (17-29%)	24 (18-31)	1900 (1500-2300)
Estonia	2010	Male	30 to 34	95% (91-97%)	5.5% (2.8-9.3%)	7 (5-9.6)	20% (17-24%)	28% (22-33%)	32 (26-39)	2300 (1800-2700)
Estonia	2010	Male	35 to 39	95% (92-97%)	5.1% (2.5-8.5%)	7.6 (5.2-11)	23% (19-27%)	33% (27-38%)	42 (34-50)	2600 (2200-3100)
Estonia	2010	Male	40 to 44	95% (92-98%)	4.6% (2.2-8.1%)	8.1 (5.8-11)	26% (22-30%)	36% (31-41%)	62 (52-72)	3300 (2800-3900)
Estonia	2010	Male	45 to 49	96% (93-98%)	4.1% (2.2-7.3%)	9 (6.6-12)	27% (23-30%)	36% (31-40%)	100 (89-120)	4800 (4100-5400)
Estonia	2010	Male	50 to 54	96% (94-97%)	4% (2.5-6.5%)	9.5 (6.7-13)	26% (22-30%)	34% (29-38%)	150 (130-170)	5900 (5100-6800)
Estonia	2010	Male	55 to 59	95% (93-97%)	4.5% (3-6.8%)	9 (6.6-12)	24% (20-28%)	29% (24-34%)	180 (150-210)	6100 (5100-7100)
Estonia	2010	Male	60 to 64	95% (92-96%)	5.5% (3.8-8.3%)	8 (5.8-11)	20% (15-24%)	24% (18-29%)	180 (140-220)	5200 (4000-6200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	2010	Male	65 to 69	93% (90-95%)	6.7% (4.6-9.9%)	6.7 (5.1-8.8)	16% (12-20%)	19% (14-23%)	150 (110-190)	3600 (2700-4500)
Estonia	2010	Male	70 to 74	92% (89-94%)	8.3% (6.2-11%)	5.6 (4-7.6)	12% (7-17%)	14% (7.8-20%)	150 (87-210)	3000 (1800-4200)
Estonia	2010	Male	75 to 79	89% (84-93%)	11% (6.9-16%)	4.7 (3.1-6.8)	9.1% (3.6-14%)	10% (3.5-17%)	110 (36-170)	1700 (640-2600)
Estonia	2010	Male	80 to 84	86% (79-91%)	14% (8.9-21%)	4.1 (2.8-5.7)	7.1% (1.9-12%)	7.9% (1.6-14%)	78 (16-140)	900 (230-1600)
Estonia	2010	Male	85 to 89	83% (76-89%)	17% (11-24%)	3.7 (2.5-5.3)	5.7% (0.5-11%)	6.1% (-0.078-13%)	32 (-0.41-67)	270 (23-540)
Estonia	2010	Male	90 to 94	82% (73-89%)	18% (11-27%)	3.5 (2.2-5.3)	4.4% (-1.2-11%)	4.5% (-2.3-12%)	8 (-4-22)	47 (-13-120)
Estonia	2010	Male	95 plus	81% (69-90%)	19% (10-31%)	3.2 (1.7-5.3)	3.4% (-2.4-11%)	3.3% (-3.9-12%)	2.6 (-3-9.6)	11 (-7.7-34)
Estonia	2016	Female	All Ages	79% (77-81%)	21% (19-23%)	2.4 (1.8-3.1)	5.2% (3.4-7.3%)	4.1% (0.65-7.9%)	350 (53-690)	11000 (6900-16000)
Estonia	2016	Female	15 to 19	77% (65-87%)	23% (13-35%)	1.9 (1.2-3)	4.1% (3-5.5%)	10% (6.4-14%)	0.57 (0.35-0.83)	99 (69-140)
Estonia	2016	Female	20 to 24	87% (79-92%)	13% (8.1-21%)	2.7 (1.6-4.2)	7.3% (5.3-9.4%)	9.8% (6.2-14%)	1.1 (0.66-1.6)	290 (190-410)
Estonia	2016	Female	25 to 29	88% (82-93%)	12% (7.1-18%)	2.7 (1.7-4)	7.8% (6.1-9.7%)	11% (7.7-15%)	1.9 (1.2-2.7)	440 (310-610)
Estonia	2016	Female	30 to 34	89% (82-93%)	11% (6.6-18%)	2.9 (1.9-4.2)	8.4% (6.6-10%)	16% (12-19%)	3.3 (2.3-4.6)	500 (360-670)
Estonia	2016	Female	35 to 39	90% (85-94%)	10% (6.4-15%)	3.3 (2.1-4.8)	9.4% (7.6-11%)	19% (15-23%)	6.3 (4.5-8.6)	640 (480-820)
Estonia	2016	Female	40 to 44	90% (85-94%)	9.9% (6.2-15%)	3.5 (2.3-5.1)	10% (8.2-13%)	20% (17-24%)	12 (8.6-15)	880 (670-1100)
Estonia	2016	Female	45 to 49	91% (87-93%)	9.4% (6.9-13%)	3.6 (2.4-5.3)	10% (8.4-13%)	19% (16-23%)	16 (12-21)	960 (740-1200)
Estonia	2016	Female	50 to 54	89% (87-91%)	11% (8.7-13%)	3.5 (2.4-5)	10% (8.1-13%)	18% (15-22%)	26 (19-34)	1300 (980-1700)
Estonia	2016	Female	55 to 59	86% (83-88%)	14% (12-17%)	3.4 (2.3-4.8)	9.5% (7-12%)	15% (12-20%)	36 (25-47)	1400 (1000-1900)
Estonia	2016	Female	60 to 64	83% (80-86%)	17% (14-20%)	3.2 (2.1-4.6)	7.6% (5-11%)	12% (8.2-16%)	41 (27-59)	1400 (890-2000)
Estonia	2016	Female	65 to 69	79% (76-83%)	21% (17-24%)	2.7 (1.9-3.8)	5.2% (2.7-8.1%)	7.6% (3.9-12%)	33 (16-52)	910 (450-1500)
Estonia	2016	Female	70 to 74	72% (67-76%)	28% (24-33%)	2.1 (1.4-3.2)	3.3% (0.55-6.6%)	4.6% (0.54-9.1%)	31 (3.6-63)	680 (110-1400)
Estonia	2016	Female	75 to 79	59% (53-65%)	41% (35-47%)	1.7 (1.1-2.6)	2.3% (-0.48-5.7%)	3% (-0.95-7.4%)	36 (-12-91)	600 (-130-1500)
Estonia	2016	Female	80 to 84	50% (39-59%)	50% (41-61%)	1.5 (0.9-2.3)	1.9% (-0.75-5.1%)	2.2% (-1.5-6.5%)	37 (-23-110)	460 (-190-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Estonia	2016	Female	85 to 89	47% (36-58%)	53% (42-64%)	1.4 (0.82-2.3)	1.7% (-1-5.1%)	1.8% (-1.7-6.3%)	36 (-33-120)	320 (-190-980)
Estonia	2016	Female	90 to 94	50% (39-60%)	50% (40-61%)	1.6 (0.86-2.6)	1.8% (-1.7-6.2%)	1.9% (-2.4-7.3%)	24 (-30-93)	140 (-140-500)
Estonia	2016	Female	95 plus	54% (39-68%)	46% (32-61%)	1.8 (0.9-3.1)	2% (-2.5-8%)	2.1% (-3.3-9.3%)	6.5 (-11-29)	25 (-31-98)
Estonia	2016	Male	All Ages	89% (86-92%)	11% (8.3-14%)	6 (4.6-7.8)	17% (14-21%)	17% (13-22%)	1300 (900-1700)	40000 (32000-50000)
Estonia	2016	Male	15 to 19	77% (65-87%)	23% (13-35%)	3.2 (1.9-5)	7.7% (5.2-11%)	16% (9.6-23%)	2.3 (1.4-3.6)	220 (150-320)
Estonia	2016	Male	20 to 24	87% (79-93%)	13% (6.9-21%)	5.6 (3.4-8.5)	15% (11-20%)	23% (16-30%)	11 (7.2-15)	930 (680-1200)
Estonia	2016	Male	25 to 29	91% (85-95%)	9% (4.9-15%)	6.3 (4-9.3)	16% (12-20%)	23% (16-29%)	18 (13-24)	1500 (1100-2000)
Estonia	2016	Male	30 to 34	93% (87-97%)	6.8% (3.3-13%)	7 (4.5-10)	19% (15-22%)	27% (21-33%)	25 (19-34)	1900 (1400-2300)
Estonia	2016	Male	35 to 39	94% (89-97%)	6% (2.9-11%)	7.6 (4.8-11)	22% (18-26%)	33% (27-39%)	37 (28-49)	2300 (1800-2900)
Estonia	2016	Male	40 to 44	95% (90-97%)	5.4% (2.6-9.9%)	8.1 (5.5-12)	25% (21-29%)	36% (30-41%)	64 (49-81)	3400 (2700-4200)
Estonia	2016	Male	45 to 49	95% (92-98%)	4.6% (2.5-8.5%)	9.1 (6.2-13)	26% (22-30%)	36% (30-40%)	83 (65-100)	3800 (3100-4600)
Estonia	2016	Male	50 to 54	96% (93-97%)	4.3% (2.5-7%)	9.8 (6.7-14)	26% (22-31%)	35% (30-39%)	140 (110-170)	5400 (4300-6500)
Estonia	2016	Male	55 to 59	95% (93-97%)	4.8% (3.1-6.9%)	9.4 (6.6-13)	24% (20-28%)	30% (25-35%)	170 (130-210)	5800 (4600-7000)
Estonia	2016	Male	60 to 64	94% (92-96%)	5.6% (3.9-8.2%)	8.4 (5.7-12)	20% (16-24%)	25% (19-30%)	190 (140-240)	5500 (4200-6900)
Estonia	2016	Male	65 to 69	93% (90-95%)	6.8% (4.6-9.8%)	7.1 (5-9.6)	16% (12-21%)	19% (14-25%)	150 (110-200)	3600 (2700-4700)
Estonia	2016	Male	70 to 74	91% (88-94%)	8.6% (6.3-12%)	5.8 (3.9-8.2)	12% (7.1-17%)	15% (8-21%)	130 (70-190)	2600 (1500-3600)
Estonia	2016	Male	75 to 79	89% (83-93%)	11% (7.1-17%)	4.8 (3.1-7.2)	9.4% (3.9-15%)	11% (4.1-17%)	110 (42-180)	1800 (730-2800)
Estonia	2016	Male	80 to 84	85% (78-91%)	15% (9.1-22%)	4.2 (2.7-6.2)	7.5% (2.1-13%)	8.5% (1.9-16%)	86 (19-160)	1000 (280-1800)
Estonia	2016	Male	85 to 89	83% (75-89%)	17% (11-25%)	3.9 (2.4-5.9)	6.2% (0.89-12%)	6.9% (0.47-14%)	55 (4-120)	450 (65-910)
Estonia	2016	Male	90 to 94	82% (73-89%)	18% (11-27%)	3.6 (2.1-5.7)	4.9% (-0.89-12%)	5.3% (-1.7-14%)	15 (-5-40)	88 (-16-220)
Estonia	2016	Male	95 plus	80% (68-90%)	20% (10-32%)	3.3 (1.7-5.8)	4% (-2-12%)	4.1% (-3.2-13%)	2.4 (-2-8.1)	9.8 (-4.9-28)
Latvia	1990	Female	All Ages	77% (73-81%)	23% (19-27%)	1.5 (1.3-1.8)	2.4% (0.85-4%)	0.081% (-2.5-3.1%)	14 (-450-530)	12000 (4200-20000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	1990	Female	15 to 19	73% (60-84%)	27% (16-40%)	1.6 (1-2.3)	4.6% (3.1-6.3%)	7.7% (4.2-12%)	4.2 (2.3-6.7)	490 (330-700)
Latvia	1990	Female	20 to 24	83% (72-90%)	17% (9.9-28%)	2.2 (1.5-3.2)	7.4% (5.5-9.5%)	9.9% (6.1-14%)	5.6 (3.4-8)	860 (620-1200)
Latvia	1990	Female	25 to 29	86% (79-92%)	14% (8.1-21%)	2.3 (1.7-3.1)	7.9% (6.2-9.5%)	11% (7.2-14%)	9 (6.1-12)	1200 (910-1600)
Latvia	1990	Female	30 to 34	88% (80-93%)	12% (7.1-20%)	2.4 (1.8-3.2)	7.9% (6.3-9.6%)	12% (8.4-15%)	12 (8.9-16)	1400 (1000-1800)
Latvia	1990	Female	35 to 39	87% (80-93%)	13% (7.3-20%)	2.3 (1.7-3.2)	8.4% (6.8-10%)	12% (9.5-15%)	18 (14-23)	1600 (1200-2000)
Latvia	1990	Female	40 to 44	87% (80-92%)	13% (7.9-20%)	2.2 (1.6-3)	8.1% (6.3-9.9%)	11% (8.2-14%)	24 (17-31)	1700 (1300-2100)
Latvia	1990	Female	45 to 49	86% (78-92%)	14% (8.4-22%)	2 (1.5-2.7)	6.9% (5.2-8.8%)	9.3% (6.6-12%)	34 (24-45)	2000 (1500-2600)
Latvia	1990	Female	50 to 54	84% (76-90%)	16% (10-24%)	1.8 (1.3-2.5)	5.6% (3.8-7.5%)	7.1% (4.3-10%)	37 (23-53)	1900 (1300-2600)
Latvia	1990	Female	55 to 59	81% (73-88%)	19% (12-27%)	1.8 (1.2-2.4)	4.2% (2.1-6.4%)	5.1% (2.2-8.4%)	38 (16-62)	1600 (810-2500)
Latvia	1990	Female	60 to 64	79% (69-86%)	21% (14-31%)	1.7 (1.2-2.3)	2.2% (-0.16-4.8%)	2.4% (-0.95-5.9%)	28 (-12-71)	1100 (-78-2300)
Latvia	1990	Female	65 to 69	77% (66-85%)	23% (15-34%)	1.5 (1-2)	0.26% (-2.4-2.9%)	-0.054% (-3.5-3.5%)	-0.91 (-55-54)	130 (-1100-1500)
Latvia	1990	Female	70 to 74	72% (60-80%)	28% (20-40%)	1.3 (0.82-1.9)	-0.89% (-3.7-2.1%)	-1.4% (-5-2.4%)	-22 (-78-38)	-350 (-1400-790)
Latvia	1990	Female	75 to 79	63% (48-75%)	37% (25-52%)	1.1 (0.65-1.7)	-1.4% (-4.2-1.6%)	-2% (-5.3-1.7%)	-58 (-160-50)	-760 (-2200-850)
Latvia	1990	Female	80 to 84	55% (39-69%)	45% (31-61%)	1.1 (0.62-1.6)	-1.4% (-4-1.7%)	-1.8% (-5.1-1.7%)	-60 (-170-57)	-580 (-1700-690)
Latvia	1990	Female	85 to 89	51% (36-65%)	49% (35-64%)	1.1 (0.6-1.8)	-1% (-3.8-2.2%)	-1.4% (-4.7-2.4%)	-37 (-120-61)	-240 (-880-510)
Latvia	1990	Female	90 to 94	50% (35-65%)	50% (35-65%)	1.2 (0.64-2)	-0.75% (-3.7-3.1%)	-1.1% (-4.8-3.4%)	-13 (-57-40)	-54 (-270-220)
Latvia	1990	Female	95 plus	52% (33-71%)	48% (29-67%)	1.3 (0.63-2.4)	-0.52% (-4.1-4.1%)	-0.9% (-5.1-4.7%)	-3.7 (-21-19)	-8.5 (-66-67)
Latvia	1990	Male	All Ages	82% (77-86%)	18% (14-23%)	4.1 (3.4-4.8)	13% (11-16%)	12% (8-15%)	1900 (1300-2600)	79000 (63000-96000)
Latvia	1990	Male	15 to 19	72% (59-83%)	28% (17-41%)	2.6 (1.7-3.9)	9.4% (6-13%)	13% (7.8-19%)	22 (13-32)	1700 (1100-2400)
Latvia	1990	Male	20 to 24	85% (75-92%)	15% (8.2-25%)	5 (3.5-6.9)	18% (13-24%)	24% (16-32%)	55 (38-73)	4100 (3000-5300)
Latvia	1990	Male	25 to 29	89% (81-94%)	11% (5.9-19%)	5.9 (4.1-8)	22% (17-27%)	27% (20-35%)	90 (66-120)	6400 (4800-7900)
Latvia	1990	Male	30 to 34	91% (84-96%)	8.6% (4.2-16%)	6.2 (4.5-8.4)	22% (18-27%)	28% (21-34%)	110 (84-140)	7200 (5700-8700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	1990	Male	35 to 39	92% (86-96%)	7.9% (4-14%)	6.5 (4.6-8.7)	22% (17-27%)	28% (21-34%)	130 (94-150)	7300 (5700-8900)
Latvia	1990	Male	40 to 44	92% (86-96%)	7.9% (4-14%)	6.6 (4.6-8.8)	22% (17-26%)	26% (20-32%)	160 (120-190)	8000 (6100-9700)
Latvia	1990	Male	45 to 49	91% (82-96%)	8.6% (4-18%)	6.1 (4.4-8.3)	19% (15-24%)	23% (17-29%)	210 (150-270)	9500 (7200-12000)
Latvia	1990	Male	50 to 54	91% (82-95%)	9.4% (4.5-18%)	5.9 (4.1-8.1)	17% (13-22%)	20% (14-26%)	270 (180-350)	10000 (7400-13000)
Latvia	1990	Male	55 to 59	90% (82-94%)	10% (5.5-18%)	5.7 (4.2-7.5)	16% (11-20%)	18% (12-23%)	290 (200-380)	9900 (7000-13000)
Latvia	1990	Male	60 to 64	89% (81-94%)	11% (5.7-19%)	5.1 (3.6-7.1)	13% (7.7-18%)	14% (8-21%)	280 (150-400)	8000 (4600-11000)
Latvia	1990	Male	65 to 69	88% (79-94%)	12% (6-21%)	4 (2.9-5.3)	9% (4.3-14%)	9.8% (4.5-16%)	160 (72-250)	3900 (1900-6000)
Latvia	1990	Male	70 to 74	86% (76-92%)	14% (7.8-24%)	3 (2-4.3)	4.9% (0.57-10%)	5.1% (0.17-11%)	73 (2.4-160)	1500 (170-3100)
Latvia	1990	Male	75 to 79	82% (70-91%)	18% (9-30%)	2.5 (1.5-3.8)	2.6% (-1.4-8.2%)	2.6% (-2.1-9.1%)	49 (-40-170)	830 (-450-2600)
Latvia	1990	Male	80 to 84	77% (63-86%)	23% (14-37%)	2.3 (1.5-3.3)	1.7% (-2-6.8%)	1.5% (-2.9-7.2%)	26 (-50-130)	360 (-420-1400)
Latvia	1990	Male	85 to 89	72% (58-82%)	28% (18-42%)	2.2 (1.4-3.3)	1.3% (-2.3-6%)	1% (-3.1-6.5%)	11 (-33-69)	110 (-210-550)
Latvia	1990	Male	90 to 94	69% (55-81%)	31% (19-45%)	2.1 (1.2-3.5)	0.87% (-2.8-6.2%)	0.52% (-3.8-6.8%)	1.7 (-12-21)	16 (-53-120)
Latvia	1990	Male	95 plus	67% (48-82%)	33% (18-52%)	2.1 (0.99-3.6)	0.74% (-3.1-6.9%)	0.28% (-4.3-7.6%)	0.29 (-4.4-7.8)	3 (-13-29)
Latvia	1995	Female	All Ages	78% (73-82%)	22% (18-27%)	1.4 (1.2-1.7)	2.4% (0.77-4.1%)	0.0049% (-2.6-2.9%)	1 (-490-550)	12000 (4000-22000)
Latvia	1995	Female	15 to 19	74% (61-85%)	26% (15-39%)	1.5 (0.98-2.1)	4.7% (3.1-6.5%)	7.8% (4.5-12%)	4.1 (2.3-6.3)	460 (300-640)
Latvia	1995	Female	20 to 24	83% (72-90%)	17% (9.7-28%)	2.1 (1.4-2.9)	7.5% (5.5-9.4%)	9.8% (6.1-14%)	6.1 (3.8-8.6)	850 (600-1100)
Latvia	1995	Female	25 to 29	86% (79-92%)	14% (8-21%)	2.2 (1.6-2.9)	8.1% (6.2-9.9%)	10% (7.2-14%)	8.5 (5.8-11)	1100 (800-1400)
Latvia	1995	Female	30 to 34	88% (80-93%)	12% (6.9-20%)	2.2 (1.7-2.9)	8.2% (6.5-9.9%)	11% (8.4-15%)	14 (10-18)	1400 (1100-1800)
Latvia	1995	Female	35 to 39	87% (81-93%)	13% (7-19%)	2.1 (1.6-2.9)	8.8% (7.1-11%)	12% (9.4-15%)	27 (21-33)	2000 (1600-2400)
Latvia	1995	Female	40 to 44	87% (80-92%)	13% (7.8-20%)	2 (1.5-2.8)	8.2% (6.3-10%)	11% (7.8-14%)	35 (25-46)	2200 (1600-2700)
Latvia	1995	Female	45 to 49	86% (78-92%)	14% (8-22%)	1.9 (1.4-2.5)	7.1% (5.1-9.4%)	9% (6-12%)	42 (28-58)	2200 (1600-2900)
Latvia	1995	Female	50 to 54	84% (76-90%)	16% (10-24%)	1.7 (1.2-2.3)	5.6% (3.5-7.9%)	6.7% (3.7-10%)	43 (24-65)	2000 (1200-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	1995	Female	55 to 59	82% (73-88%)	18% (12-27%)	1.7 (1.2-2.2)	4% (1.9-6.5%)	4.7% (1.6-8.2%)	47 (17-81)	1900 (850-3000)
Latvia	1995	Female	60 to 64	79% (69-86%)	21% (14-31%)	1.6 (1.1-2.2)	1.8% (-0.48-4.5%)	1.9% (-1.2-5.4%)	23 (-15-65)	860 (-210-2100)
Latvia	1995	Female	65 to 69	77% (66-85%)	23% (15-34%)	1.4 (1-1.9)	-0.017% (-2.5-2.5%)	-0.38% (-3.7-2.8%)	-7 (-68-53)	-9.2 (-1400-1400)
Latvia	1995	Female	70 to 74	72% (59-81%)	28% (19-41%)	1.2 (0.76-1.8)	-1.1% (-3.8-1.6%)	-1.7% (-5-1.9%)	-40 (-120-45)	-650 (-2100-930)
Latvia	1995	Female	75 to 79	64% (48-76%)	36% (24-52%)	1.1 (0.63-1.6)	-1.7% (-4.2-1.1%)	-2.3% (-5.5-1.1%)	-48 (-120-25)	-640 (-1600-420)
Latvia	1995	Female	80 to 84	56% (40-69%)	44% (31-60%)	1 (0.6-1.5)	-1.7% (-4.4-1.3%)	-2.2% (-5.4-1.4%)	-72 (-180-43)	-700 (-1800-530)
Latvia	1995	Female	85 to 89	51% (36-66%)	49% (34-64%)	1 (0.58-1.7)	-1.5% (-4.1-1.6%)	-1.9% (-5-1.7%)	-54 (-140-48)	-360 (-1000-400)
Latvia	1995	Female	90 to 94	51% (37-66%)	49% (34-63%)	1.1 (0.62-1.9)	-1.2% (-4.4-2.4%)	-1.6% (-5.4-2.7%)	-23 (-76-38)	-100 (-370-210)
Latvia	1995	Female	95 plus	53% (34-72%)	47% (28-66%)	1.3 (0.6-2.3)	-0.99% (-4.5-3.7%)	-1.4% (-5.6-4.1%)	-5.7 (-23-17)	-16 (-74-60)
Latvia	1995	Male	All Ages	83% (78-87%)	17% (13-22%)	3.8 (3.2-4.4)	14% (11-17%)	12% (8.4-16%)	2400 (1600-3100)	97000 (75000-120000)
Latvia	1995	Male	15 to 19	73% (59-84%)	27% (16-41%)	2.4 (1.5-3.5)	9.5% (6-13%)	13% (7.8-19%)	20 (12-29)	1600 (990-2200)
Latvia	1995	Male	20 to 24	85% (75-92%)	15% (8.2-25%)	4.6 (3.3-6.2)	18% (14-24%)	23% (16-29%)	64 (45-85)	4700 (3400-6000)
Latvia	1995	Male	25 to 29	89% (81-94%)	11% (5.7-19%)	5.5 (3.9-7.3)	22% (17-28%)	26% (19-33%)	100 (73-130)	6800 (5100-8600)
Latvia	1995	Male	30 to 34	91% (85-96%)	8.5% (4.1-15%)	5.8 (4.3-7.8)	23% (17-28%)	26% (20-33%)	140 (100-180)	8500 (6600-11000)
Latvia	1995	Male	35 to 39	92% (87-96%)	7.9% (4-13%)	6 (4.2-8)	22% (17-28%)	26% (19-32%)	190 (140-240)	11000 (7900-13000)
Latvia	1995	Male	40 to 44	92% (86-96%)	7.9% (4-14%)	6 (4.3-8.1)	22% (16-27%)	25% (18-31%)	240 (170-310)	12000 (8600-15000)
Latvia	1995	Male	45 to 49	92% (84-96%)	8.3% (3.9-16%)	5.7 (4.1-7.7)	19% (14-25%)	22% (16-28%)	270 (190-350)	12000 (8300-15000)
Latvia	1995	Male	50 to 54	91% (84-96%)	9% (4.4-16%)	5.5 (3.8-7.6)	17% (12-23%)	19% (12-26%)	320 (210-430)	12000 (8100-16000)
Latvia	1995	Male	55 to 59	90% (83-95%)	10% (5.4-17%)	5.3 (3.9-7.1)	16% (10-21%)	17% (11-23%)	380 (240-510)	13000 (8100-17000)
Latvia	1995	Male	60 to 64	89% (81-95%)	11% (5.4-19%)	4.8 (3.4-6.5)	12% (6.7-18%)	13% (7-20%)	320 (170-480)	9200 (4900-13000)
Latvia	1995	Male	65 to 69	89% (80-94%)	11% (5.9-20%)	3.7 (2.7-4.9)	8% (3.5-13%)	8.6% (3.4-14%)	220 (85-360)	5200 (2300-8500)
Latvia	1995	Male	70 to 74	86% (77-92%)	14% (7.7-23%)	2.8 (1.9-4)	4% (0.082-9.1%)	4.2% (-0.28-9.8%)	80 (-5.5-190)	1600 (33-3600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	1995	Male	75 to 79	82% (70-91%)	18% (9.5-30%)	2.3 (1.4-3.5)	2% (-1.8-7.1%)	1.9% (-2.5-7.7%)	26 (-34-110)	450 (-410-1600)
Latvia	1995	Male	80 to 84	77% (63-87%)	23% (13-37%)	2.1 (1.4-3.1)	1% (-2.6-5.4%)	0.78% (-3.4-5.9%)	12 (-53-92)	190 (-480-1000)
Latvia	1995	Male	85 to 89	73% (59-83%)	27% (17-41%)	2.1 (1.3-3.1)	0.53% (-3-4.9%)	0.18% (-3.8-5.1%)	1.9 (-40-54)	48 (-260-440)
Latvia	1995	Male	90 to 94	70% (55-81%)	30% (19-45%)	2 (1.2-3.1)	0.2% (-3.3-5%)	-0.22% (-4.4-5.2%)	-0.91 (-18-22)	4.8 (-80-120)
Latvia	1995	Male	95 plus	67% (49-82%)	33% (18-51%)	1.9 (0.93-3.3)	0.13% (-3.5-6%)	-0.38% (-4.7-6.4%)	-0.27 (-3.3-4.4)	0.37 (-9.7-16)
Latvia	2000	Female	All Ages	77% (72-81%)	23% (19-28%)	1.5 (1.3-1.7)	2.7% (1.4-4.2%)	0.51% (-1.7-3%)	80 (-260-470)	12000 (6000-19000)
Latvia	2000	Female	15 to 19	74% (61-85%)	26% (15-39%)	1.6 (1.1-2.2)	4.4% (3-6%)	8.1% (4.6-12%)	3.3 (1.9-5.1)	420 (290-590)
Latvia	2000	Female	20 to 24	83% (72-90%)	17% (9.7-28%)	2.2 (1.5-3)	7.5% (5.6-9.5%)	10% (6.6-14%)	5 (3.2-6.9)	750 (530-1000)
Latvia	2000	Female	25 to 29	87% (79-92%)	13% (8-21%)	2.3 (1.7-3.1)	8% (6.3-9.7%)	11% (7.9-15%)	6.5 (4.7-8.5)	920 (690-1200)
Latvia	2000	Female	30 to 34	88% (81-93%)	12% (6.8-19%)	2.4 (1.8-3.2)	8.2% (6.7-10%)	12% (9.2-15%)	10 (7.9-13)	1100 (860-1400)
Latvia	2000	Female	35 to 39	87% (80-93%)	13% (7.2-20%)	2.2 (1.7-3)	8.6% (7.1-10%)	13% (10-16%)	19 (15-24)	1600 (1200-1900)
Latvia	2000	Female	40 to 44	86% (79-92%)	14% (8.2-21%)	2.1 (1.5-2.8)	8.3% (6.6-10%)	12% (9.4-15%)	28 (21-35)	1900 (1500-2300)
Latvia	2000	Female	45 to 49	86% (78-92%)	14% (8.3-22%)	2 (1.5-2.6)	7.5% (5.8-9.5%)	11% (7.8-14%)	35 (26-45)	1900 (1500-2400)
Latvia	2000	Female	50 to 54	84% (76-90%)	16% (10-24%)	1.8 (1.3-2.4)	6.3% (4.6-8.3%)	8.5% (5.9-12%)	39 (26-53)	1800 (1300-2400)
Latvia	2000	Female	55 to 59	80% (71-87%)	20% (13-29%)	1.7 (1.2-2.3)	4.7% (2.8-7%)	6.1% (3.1-9.4%)	39 (20-61)	1600 (900-2300)
Latvia	2000	Female	60 to 64	77% (66-85%)	23% (15-34%)	1.6 (1.1-2.2)	2.6% (0.57-4.9%)	3.1% (0.21-6.3%)	31 (2.1-62)	1100 (240-2000)
Latvia	2000	Female	65 to 69	73% (60-83%)	27% (17-40%)	1.3 (0.9-1.8)	0.68% (-1.4-2.8%)	0.55% (-2.3-3.5%)	7.3 (-30-46)	300 (-610-1300)
Latvia	2000	Female	70 to 74	64% (50-76%)	36% (24-50%)	1 (0.61-1.5)	-0.34% (-2.3-1.8%)	-0.68% (-3.4-2.2%)	-14 (-72-46)	-180 (-1300-940)
Latvia	2000	Female	75 to 79	56% (40-70%)	44% (30-60%)	0.89 (0.53-1.4)	-0.91% (-2.9-1.3%)	-1.3% (-3.8-1.5%)	-36 (-100-39)	-450 (-1400-640)
Latvia	2000	Female	80 to 84	52% (36-66%)	48% (34-64%)	0.95 (0.57-1.4)	-1.2% (-3.5-1.3%)	-1.6% (-4.4-1.3%)	-34 (-94-28)	-320 (-960-370)
Latvia	2000	Female	85 to 89	50% (35-65%)	50% (35-65%)	1 (0.59-1.7)	-1.1% (-3.6-1.7%)	-1.5% (-4.5-1.8%)	-39 (-110-45)	-250 (-800-380)
Latvia	2000	Female	90 to 94	49% (34-64%)	51% (36-66%)	1.1 (0.61-1.9)	-0.84% (-3.7-2.6%)	-1.2% (-4.7-2.9%)	-17 (-64-41)	-71 (-310-220)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	2000	Female	95 plus	51% (33-71%)	49% (29-67%)	1.3 (0.61-2.3)	-0.56% (-4.1-3.9%)	-0.91% (-5-4.4%)	-3.8 (-21-18)	-9.4 (-68-66)
Latvia	2000	Male	All Ages	84% (80-88%)	16% (12-20%)	4.2 (3.7-4.8)	14% (12-17%)	12% (9-16%)	1900 (1400-2500)	77000 (62000-92000)
Latvia	2000	Male	15 to 19	75% (63-85%)	25% (15-37%)	2.7 (1.8-3.8)	9.2% (6.1-13%)	14% (8.6-20%)	16 (9.9-23)	1300 (860-1800)
Latvia	2000	Male	20 to 24	87% (78-93%)	13% (7.2-22%)	5 (3.6-6.7)	19% (14-24%)	24% (18-31%)	49 (35-64)	3600 (2700-4600)
Latvia	2000	Male	25 to 29	90% (84-95%)	9.7% (5.4-16%)	5.9 (4.2-7.7)	22% (17-27%)	28% (21-35%)	69 (52-88)	4900 (3800-6000)
Latvia	2000	Male	30 to 34	92% (86-96%)	8% (4-14%)	6.2 (4.6-8.1)	23% (18-27%)	28% (22-34%)	88 (69-110)	5600 (4500-6800)
Latvia	2000	Male	35 to 39	92% (87-96%)	7.6% (3.8-13%)	6.4 (4.7-8.5)	23% (18-27%)	28% (22-34%)	130 (100-160)	7500 (6000-9000)
Latvia	2000	Male	40 to 44	93% (88-96%)	7.4% (3.9-12%)	6.4 (4.7-8.4)	23% (18-27%)	27% (21-33%)	190 (150-230)	9400 (7400-11000)
Latvia	2000	Male	45 to 49	92% (85-96%)	7.7% (3.8-15%)	6.1 (4.4-8.1)	21% (16-26%)	25% (19-30%)	210 (160-270)	9400 (7200-12000)
Latvia	2000	Male	50 to 54	92% (85-96%)	8.3% (4.3-15%)	5.8 (4.2-7.9)	19% (13-23%)	21% (15-27%)	250 (170-310)	9400 (6800-12000)
Latvia	2000	Male	55 to 59	91% (84-95%)	9.3% (5.1-16%)	5.8 (4.3-7.4)	17% (12-22%)	19% (13-25%)	270 (190-350)	9200 (6300-12000)
Latvia	2000	Male	60 to 64	90% (82-95%)	10% (5.5-18%)	5.1 (3.7-7)	13% (8-19%)	15% (8.9-21%)	300 (180-430)	8700 (5300-12000)
Latvia	2000	Male	65 to 69	89% (80-94%)	11% (5.7-20%)	3.9 (2.9-5.1)	8.8% (4.5-14%)	9.7% (4.6-15%)	200 (94-320)	4900 (2400-7600)
Latvia	2000	Male	70 to 74	85% (75-92%)	15% (8.2-25%)	2.8 (1.9-3.9)	4.4% (0.65-9.1%)	4.7% (0.27-10%)	100 (5.9-220)	2100 (310-4300)
Latvia	2000	Male	75 to 79	81% (68-90%)	19% (9.8-32%)	2.3 (1.4-3.5)	2.3% (-1.2-6.8%)	2.3% (-1.9-7.6%)	34 (-28-110)	580 (-300-1700)
Latvia	2000	Male	80 to 84	76% (62-86%)	24% (14-38%)	2.2 (1.4-3.1)	1.4% (-2.1-5.7%)	1.2% (-2.9-6.2%)	11 (-28-58)	170 (-240-680)
Latvia	2000	Male	85 to 89	71% (58-82%)	29% (18-42%)	2.1 (1.3-3.1)	0.84% (-2.6-4.9%)	0.5% (-3.5-5.3%)	4.3 (-30-46)	62 (-190-370)
Latvia	2000	Male	90 to 94	69% (55-80%)	31% (20-45%)	2 (1.2-3.2)	0.58% (-2.9-5.2%)	0.2% (-3.8-5.7%)	0.83 (-16-23)	14 (-72-130)
Latvia	2000	Male	95 plus	66% (48-82%)	34% (18-52%)	2 (0.96-3.4)	0.58% (-2.9-6%)	0.12% (-4-6.4%)	0.12 (-3.9-6.2)	2.2 (-11-23)
Latvia	2005	Female	All Ages	78% (75-82%)	22% (18-25%)	1.7 (1.5-2)	3% (1.6-4.5%)	1% (-1.3-3.5%)	160 (-210-560)	13000 (6900-20000)
Latvia	2005	Female	15 to 19	77% (66-86%)	23% (14-34%)	1.8 (1.3-2.5)	4.5% (3.1-5.9%)	8.4% (4.9-13%)	3.1 (1.8-4.8)	410 (280-560)
Latvia	2005	Female	20 to 24	85% (75-91%)	15% (8.6-25%)	2.5 (1.8-3.4)	7.5% (5.7-9.5%)	11% (7.2-15%)	4.4 (2.9-6)	740 (530-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	2005	Female	25 to 29	87% (80-92%)	13% (7.8-20%)	2.6 (1.9-3.4)	8% (6.5-9.7%)	12% (8.5-15%)	5.4 (3.9-7.1)	820 (610-1100)
Latvia	2005	Female	30 to 34	88% (82-93%)	12% (6.9-18%)	2.6 (2-3.5)	8.3% (6.7-9.9%)	13% (9.9-16%)	9.9 (7.6-12)	1100 (830-1300)
Latvia	2005	Female	35 to 39	88% (82-93%)	12% (7.1-18%)	2.5 (1.9-3.3)	8.9% (7.4-11%)	14% (11-17%)	17 (13-20)	1400 (1100-1700)
Latvia	2005	Female	40 to 44	88% (82-92%)	12% (7.6-18%)	2.4 (1.8-3.2)	9.2% (7.4-11%)	14% (11-17%)	30 (24-36)	1900 (1500-2300)
Latvia	2005	Female	45 to 49	87% (80-92%)	13% (7.9-20%)	2.2 (1.7-2.9)	8.1% (6.4-10%)	12% (9.3-15%)	39 (31-49)	2100 (1700-2600)
Latvia	2005	Female	50 to 54	85% (78-90%)	15% (10-22%)	2 (1.5-2.6)	7% (5.2-9.1%)	9.7% (7-13%)	44 (32-58)	2000 (1500-2600)
Latvia	2005	Female	55 to 59	81% (73-87%)	19% (13-27%)	1.9 (1.4-2.5)	5.4% (3.4-7.6%)	7% (4.1-10%)	43 (26-65)	1700 (1000-2400)
Latvia	2005	Female	60 to 64	79% (70-86%)	21% (14-30%)	1.8 (1.3-2.5)	3.4% (1.2-5.9%)	4.3% (1.2-7.8%)	37 (10-67)	1200 (420-2100)
Latvia	2005	Female	65 to 69	76% (67-83%)	24% (17-33%)	1.6 (1.2-2.1)	1.2% (-0.97-3.5%)	1.3% (-1.8-4.5%)	18 (-24-60)	540 (-430-1600)
Latvia	2005	Female	70 to 74	67% (58-75%)	33% (25-42%)	1.2 (0.81-1.7)	-0.0026% (-2.2-2.5%)	-0.2% (-3.2-3%)	-3.8 (-60-56)	-1.2 (-1100-1200)
Latvia	2005	Female	75 to 79	55% (41-68%)	45% (32-59%)	0.97 (0.58-1.4)	-0.5% (-2.5-1.8%)	-0.82% (-3.3-2%)	-23 (-96-57)	-260 (-1300-940)
Latvia	2005	Female	80 to 84	50% (35-65%)	50% (35-65%)	1 (0.59-1.5)	-0.66% (-2.9-1.8%)	-0.98% (-3.7-1.9%)	-32 (-120-59)	-280 (-1200-750)
Latvia	2005	Female	85 to 89	49% (34-64%)	51% (36-66%)	1.1 (0.63-1.8)	-0.66% (-3.2-2.2%)	-0.99% (-4-2.4%)	-19 (-76-46)	-110 (-560-390)
Latvia	2005	Female	90 to 94	49% (35-64%)	51% (36-65%)	1.2 (0.67-2.1)	-0.37% (-3.4-3.4%)	-0.66% (-4.2-3.8%)	-9.3 (-60-54)	-32 (-290-280)
Latvia	2005	Female	95 plus	51% (32-70%)	49% (30-68%)	1.4 (0.65-2.5)	-0.017% (-3.8-4.9%)	-0.28% (-4.7-5.5%)	-1.4 (-24-27)	-0.35 (-76-93)
Latvia	2005	Male	All Ages	86% (83-90%)	14% (10-17%)	4.9 (4.3-5.5)	15% (12-18%)	13% (9.5-17%)	2100 (1500-2800)	81000 (66000-96000)
Latvia	2005	Male	15 to 19	79% (69-87%)	21% (13-31%)	3.1 (2.2-4.4)	9.2% (6.3-13%)	15% (9.9-21%)	13 (8.5-18)	1100 (760-1500)
Latvia	2005	Male	20 to 24	88% (80-93%)	12% (6.7-20%)	5.7 (4.1-7.4)	19% (15-25%)	27% (20-34%)	43 (32-56)	3300 (2500-4100)
Latvia	2005	Male	25 to 29	91% (86-95%)	8.7% (4.8-14%)	6.6 (4.8-8.7)	23% (18-28%)	30% (23-37%)	61 (47-75)	4300 (3400-5200)
Latvia	2005	Male	30 to 34	93% (88-96%)	7.1% (3.6-12%)	7 (5.2-9.2)	24% (20-28%)	30% (25-36%)	89 (71-110)	5700 (4600-6700)
Latvia	2005	Male	35 to 39	93% (89-96%)	6.9% (3.6-11%)	7.2 (5.3-9.5)	24% (20-29%)	30% (24-35%)	120 (98-150)	7000 (5600-8300)
Latvia	2005	Male	40 to 44	93% (89-96%)	6.8% (3.6-11%)	7.3 (5.3-9.6)	24% (20-28%)	29% (23-34%)	190 (150-220)	9200 (7300-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	2005	Male	45 to 49	93% (87-97%)	6.9% (3.5-13%)	6.9 (5.1-9.1)	22% (17-27%)	26% (21-32%)	240 (180-290)	10000 (8200-13000)
Latvia	2005	Male	50 to 54	93% (87-96%)	7.2% (3.8-13%)	6.7 (4.8-8.9)	20% (15-25%)	23% (18-29%)	270 (200-340)	10000 (7800-13000)
Latvia	2005	Male	55 to 59	92% (86-96%)	8.2% (4.4-14%)	6.5 (4.9-8.5)	18% (14-23%)	21% (15-26%)	290 (210-370)	9600 (7100-12000)
Latvia	2005	Male	60 to 64	91% (85-95%)	8.9% (4.7-15%)	5.9 (4.2-8)	15% (9.6-20%)	17% (10-23%)	310 (180-420)	8700 (5500-12000)
Latvia	2005	Male	65 to 69	90% (83-95%)	9.7% (5-17%)	4.6 (3.5-6)	11% (5.6-15%)	12% (5.9-17%)	260 (130-390)	6400 (3400-9200)
Latvia	2005	Male	70 to 74	88% (80-93%)	12% (6.9-20%)	3.4 (2.4-4.6)	5.8% (1.2-11%)	6.2% (0.89-12%)	140 (20-270)	2800 (560-5200)
Latvia	2005	Male	75 to 79	82% (71-91%)	18% (9.2-29%)	2.6 (1.7-3.9)	3.2% (-0.85-8.2%)	3.3% (-1.3-8.9%)	70 (-28-190)	1100 (-300-2900)
Latvia	2005	Male	80 to 84	75% (60-86%)	25% (14-40%)	2.3 (1.5-3.3)	2.1% (-1.4-6.3%)	1.9% (-2.2-6.8%)	25 (-28-86)	320 (-230-980)
Latvia	2005	Male	85 to 89	71% (57-82%)	29% (18-43%)	2.3 (1.4-3.4)	1.4% (-2.5-5.8%)	1.2% (-2.8-6.3%)	7.4 (-18-40)	79 (-110-320)
Latvia	2005	Male	90 to 94	68% (54-80%)	32% (20-46%)	2.2 (1.3-3.5)	1% (-2.5-6.2%)	0.72% (-3.5-6.7%)	2.7 (-13-25)	23 (-56-140)
Latvia	2005	Male	95 plus	66% (48-81%)	34% (19-52%)	2.1 (1-3.7)	0.96% (-2.9-6.7%)	0.55% (-4.7-3%)	0.61 (-4.4-8.1)	4.3 (-13-30)
Latvia	2010	Female	All Ages	78% (74-81%)	22% (19-26%)	1.9 (1.6-2.2)	3.4% (1.9-4.9%)	1.6% (-1-4.2%)	230 (-150-630)	13000 (7100-20000)
Latvia	2010	Female	15 to 19	76% (65-86%)	24% (14-35%)	2 (1.4-2.7)	4.1% (2.9-5.4%)	8.2% (4.9-13%)	1.5 (0.89-2.4)	250 (170-350)
Latvia	2010	Female	20 to 24	84% (74-91%)	16% (8.9-26%)	2.7 (1.9-3.8)	7.2% (5.4-9.1%)	11% (7.2-15%)	2.9 (1.9-4)	620 (430-870)
Latvia	2010	Female	25 to 29	87% (79-92%)	13% (8.2-21%)	2.7 (2-3.6)	7.7% (6.1-9.4%)	11% (8.4-14%)	4.2 (3.1-5.5)	730 (530-960)
Latvia	2010	Female	30 to 34	88% (81-93%)	12% (7.2-19%)	2.8 (2-3.6)	8.1% (6.6-9.8%)	13% (11-16%)	7.3 (5.7-9.1)	860 (670-1100)
Latvia	2010	Female	35 to 39	88% (82-93%)	12% (7-18%)	2.7 (2-3.6)	8.6% (7-10%)	14% (11-17%)	14 (12-18)	1200 (970-1500)
Latvia	2010	Female	40 to 44	88% (82-92%)	12% (7.5-18%)	2.6 (1.9-3.5)	9.7% (7.8-12%)	17% (13-20%)	26 (20-31)	1600 (1300-1900)
Latvia	2010	Female	45 to 49	87% (80-92%)	13% (8.2-20%)	2.4 (1.8-3.1)	8.9% (7.1-11%)	15% (11-17%)	37 (29-45)	2000 (1500-2400)
Latvia	2010	Female	50 to 54	84% (78-90%)	16% (10-22%)	2.1 (1.6-2.8)	7.8% (5.9-10%)	12% (8.9-15%)	47 (35-60)	2100 (1600-2700)
Latvia	2010	Female	55 to 59	81% (73-87%)	19% (13-27%)	2 (1.5-2.7)	6.4% (4.3-8.5%)	9.1% (6-12%)	49 (32-66)	1800 (1200-2500)
Latvia	2010	Female	60 to 64	79% (70-86%)	21% (14-30%)	1.9 (1.4-2.7)	4.5% (2.2-6.9%)	6.2% (3-9.7%)	46 (22-71)	1400 (720-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	2010	Female	65 to 69	76% (67-84%)	24% (16-33%)	1.7 (1.3-2.2)	2% (-0.14-4.3%)	2.6% (-0.5-5.7%)	28 (-5.5-61)	760 (-54-1600)
Latvia	2010	Female	70 to 74	69% (59-77%)	31% (23-41%)	1.4 (0.91-1.9)	0.54% (-1.7-3%)	0.55% (-2.4-3.8%)	9.4 (-43-67)	250 (-790-1400)
Latvia	2010	Female	75 to 79	61% (47-73%)	39% (27-53%)	1.2 (0.76-1.8)	-0.24% (-2.6-2.4%)	-0.54% (-3.6-2.8%)	-13 (-84-65)	-110 (-1200-1100)
Latvia	2010	Female	80 to 84	54% (37-68%)	46% (32-63%)	1.2 (0.72-1.8)	-0.27% (-2.7-2.3%)	-0.56% (-3.5-2.7%)	-18 (-110-86)	-120 (-1200-1000)
Latvia	2010	Female	85 to 89	49% (34-64%)	51% (36-66%)	1.2 (0.68-1.9)	-0.12% (-2.6-3%)	-0.39% (-3.5-3.5%)	-10 (-92-91)	-28 (-650-730)
Latvia	2010	Female	90 to 94	48% (34-64%)	52% (36-66%)	1.3 (0.71-2.3)	0.17% (-2.8-4.1%)	-0.066% (-3.6-4.6%)	-0.68 (-36-46)	11 (-180-250)
Latvia	2010	Female	95 plus	50% (31-70%)	50% (30-69%)	1.5 (0.68-2.7)	0.56% (-3.1-5.6%)	0.35% (-4.2-6.5%)	1.6 (-19-29)	10 (-60-100)
Latvia	2010	Male	All Ages	86% (83-89%)	14% (11-17%)	5.3 (4.5-6.1)	16% (14-19%)	15% (11-18%)	2000 (1500-2600)	74000 (61000-86000)
Latvia	2010	Male	15 to 19	78% (69-87%)	22% (13-31%)	3.4 (2.3-4.7)	9.3% (6.3-13%)	16% (10-22%)	8.9 (5.7-12)	760 (520-1000)
Latvia	2010	Male	20 to 24	88% (80-93%)	12% (7-20%)	6.1 (4.3-8)	19% (14-24%)	29% (22-37%)	32 (23-41)	2500 (1900-3100)
Latvia	2010	Male	25 to 29	91% (86-95%)	9% (5-14%)	7.1 (5-9.3)	23% (18-28%)	33% (26-40%)	50 (39-61)	3600 (2900-4400)
Latvia	2010	Male	30 to 34	93% (88-96%)	7.2% (3.7-12%)	7.5 (5.5-9.8)	25% (21-30%)	35% (28-41%)	69 (55-82)	4500 (3700-5300)
Latvia	2010	Male	35 to 39	93% (89-96%)	7% (3.7-11%)	7.7 (5.6-10)	26% (21-31%)	35% (29-40%)	100 (83-120)	5800 (4700-6800)
Latvia	2010	Male	40 to 44	93% (88-96%)	7% (3.8-12%)	7.7 (5.6-10)	26% (22-31%)	34% (28-39%)	140 (120-170)	7200 (5800-8300)
Latvia	2010	Male	45 to 49	93% (86-96%)	7.2% (3.7-14%)	7.3 (5.4-9.9)	26% (21-30%)	32% (26-37%)	220 (180-260)	9700 (7800-11000)
Latvia	2010	Male	50 to 54	92% (86-96%)	7.6% (4.2-14%)	7.1 (5.1-9.7)	23% (18-27%)	28% (22-33%)	270 (210-330)	10000 (8100-12000)
Latvia	2010	Male	55 to 59	91% (86-95%)	8.6% (4.9-14%)	7 (5.2-9.1)	21% (16-25%)	24% (19-29%)	290 (230-360)	9700 (7600-12000)
Latvia	2010	Male	60 to 64	91% (84-95%)	9.3% (5.1-16%)	6.3 (4.4-8.5)	17% (12-22%)	20% (14-26%)	290 (200-370)	8200 (5800-10000)
Latvia	2010	Male	65 to 69	90% (83-95%)	9.9% (5.1-17%)	4.9 (3.7-6.4)	12% (7.7-17%)	14% (8.5-19%)	250 (150-340)	6000 (3700-8100)
Latvia	2010	Male	70 to 74	88% (81-93%)	12% (6.9-19%)	3.7 (2.6-5)	7.2% (2.6-12%)	8% (2.5-14%)	170 (53-290)	3300 (1200-5600)
Latvia	2010	Male	75 to 79	83% (72-91%)	17% (9.3-28%)	2.9 (1.9-4.3)	4.2% (-0.076-9.4%)	4.4% (-0.44-10%)	84 (-8.7-200)	1400 (-25-3000)
Latvia	2010	Male	80 to 84	76% (61-86%)	24% (14-39%)	2.6 (1.6-3.7)	2.9% (-0.53-7.4%)	2.9% (-1.2-8.1%)	46 (-19-130)	580 (-100-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	2010	Male	85 to 89	71% (56-82%)	29% (18-44%)	2.5 (1.5-3.7)	2.3% (-1.2-6.9%)	2.2% (-2.1-7.7%)	16 (-16-58)	150 (-82-470)
Latvia	2010	Male	90 to 94	68% (54-80%)	32% (20-46%)	2.4 (1.4-3.9)	1.7% (-2-7.3%)	1.5% (-2.9-8.1%)	3.7 (-7.4-20)	27 (-31-110)
Latvia	2010	Male	95 plus	66% (48-81%)	34% (19-52%)	2.3 (1.1-4)	1.6% (-2.5-7.6%)	1.2% (-3.8-8.4%)	1 (-3.2-7.2)	5.6 (-8.9-26)
Latvia	2016	Female	All Ages	76% (72-80%)	24% (20-28%)	1.9 (1.4-2.4)	3.4% (1.7-5.4%)	1.8% (-1.1-5.5%)	270 (-170-820)	13000 (6100-21000)
Latvia	2016	Female	15 to 19	72% (58-84%)	28% (16-42%)	1.9 (1.2-2.9)	3.9% (2.7-5.2%)	8.2% (4.6-13%)	0.62 (0.34-0.99)	120 (83-180)
Latvia	2016	Female	20 to 24	82% (70-90%)	18% (10-30%)	2.7 (1.7-4.2)	7.3% (5.5-9.5%)	11% (6.9-15%)	1.9 (1.2-2.8)	460 (310-640)
Latvia	2016	Female	25 to 29	86% (77-92%)	14% (8.4-23%)	2.8 (1.9-4)	7.8% (6.1-9.8%)	11% (8.1-15%)	3.6 (2.3-5.1)	700 (500-930)
Latvia	2016	Female	30 to 34	87% (78-93%)	13% (7-22%)	2.9 (1.9-4)	7.9% (6.3-9.8%)	13% (9.8-16%)	5.9 (4.1-8.1)	760 (570-1000)
Latvia	2016	Female	35 to 39	87% (79-93%)	13% (7.2-21%)	2.8 (1.9-4)	8.7% (6.9-11%)	15% (12-19%)	11 (7.8-15)	960 (730-1200)
Latvia	2016	Female	40 to 44	86% (79-92%)	14% (8-21%)	2.7 (1.8-3.9)	9.5% (7.5-12%)	17% (14-21%)	22 (16-28)	1400 (1100-1800)
Latvia	2016	Female	45 to 49	86% (77-92%)	14% (8.3-23%)	2.4 (1.7-3.5)	8.9% (6.9-11%)	15% (12-19%)	29 (21-38)	1500 (1100-1900)
Latvia	2016	Female	50 to 54	84% (75-90%)	16% (10-25%)	2.2 (1.5-3.1)	8% (5.9-10%)	13% (9.6-17%)	43 (30-58)	1900 (1400-2500)
Latvia	2016	Female	55 to 59	81% (72-88%)	19% (12-28%)	2.1 (1.4-3.1)	6.7% (4.5-9.1%)	10% (6.8-14%)	49 (32-70)	1900 (1200-2600)
Latvia	2016	Female	60 to 64	78% (68-86%)	22% (14-32%)	2 (1.3-3)	4.8% (2.4-7.4%)	6.9% (3.5-11%)	47 (23-76)	1500 (760-2400)
Latvia	2016	Female	65 to 69	76% (64-84%)	24% (16-36%)	1.8 (1.2-2.6)	2.4% (0.088-5%)	3.4% (-0.023-6.9%)	26 (-0.18-55)	720 (23-1500)
Latvia	2016	Female	70 to 74	70% (58-80%)	30% (20-42%)	1.5 (0.92-2.3)	0.93% (-1.4-3.6%)	1.1% (-2.2-4.9%)	16 (-31-70)	380 (-580-1500)
Latvia	2016	Female	75 to 79	61% (45-75%)	39% (25-55%)	1.3 (0.75-2)	0.14% (-2.2-3%)	-0.036% (-3.1-3.6%)	-0.72 (-70-80)	66 (-1000-1400)
Latvia	2016	Female	80 to 84	53% (37-67%)	47% (33-63%)	1.2 (0.71-2)	0.13% (-2.4-3.2%)	-0.064% (-3.2-3.7%)	-1.8 (-91-110)	53 (-950-1300)
Latvia	2016	Female	85 to 89	49% (33-64%)	51% (36-67%)	1.3 (0.69-2.1)	0.3% (-2.4-4%)	0.12% (-3.2-4.6%)	4.1 (-110-150)	97 (-780-1300)
Latvia	2016	Female	90 to 94	49% (34-64%)	51% (36-66%)	1.4 (0.73-2.5)	0.62% (-2.7-5%)	0.49% (-3.5-5.9%)	9.1 (-64-110)	73 (-310-590)
Latvia	2016	Female	95 plus	51% (31-70%)	49% (30-69%)	1.6 (0.69-3.1)	1.1% (-3-7.5%)	1% (-3.7-8.4%)	4.2 (-16-35)	18 (-49-120)
Latvia	2016	Male	All Ages	84% (80-87%)	16% (13-20%)	5.2 (3.9-6.7)	16% (13-19%)	14% (9.6-18%)	1900 (1300-2600)	65000 (49000-82000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Latvia	2016	Male	15 to 19	71% (57-82%)	29% (18-43%)	3.1 (1.9-4.7)	8.1% (5.4-11%)	16% (9.6-23%)	3.6 (2.1-5.4)	330 (220-470)
Latvia	2016	Male	20 to 24	84% (74-91%)	16% (8.5-26%)	5.9 (3.9-8.7)	18% (13-23%)	29% (21-38%)	19 (13-26)	1600 (1200-2000)
Latvia	2016	Male	25 to 29	89% (81-94%)	11% (6-19%)	7 (4.7-10)	22% (18-27%)	33% (25-40%)	43 (30-55)	3200 (2400-4000)
Latvia	2016	Male	30 to 34	91% (83-96%)	8.9% (4.3-17%)	7.5 (5-10)	25% (20-30%)	35% (27-41%)	62 (44-79)	4000 (3000-5000)
Latvia	2016	Male	35 to 39	92% (86-96%)	8.2% (4.1-14%)	7.8 (5.1-11)	26% (21-31%)	35% (28-41%)	79 (57-100)	4500 (3400-5600)
Latvia	2016	Male	40 to 44	92% (86-96%)	8.1% (4-14%)	7.8 (5.1-11)	26% (21-31%)	35% (28-40%)	120 (90-160)	6200 (4700-7700)
Latvia	2016	Male	45 to 49	91% (83-96%)	8.6% (4.2-17%)	7.4 (4.9-11)	26% (20-31%)	33% (26-39%)	160 (120-210)	7200 (5400-9200)
Latvia	2016	Male	50 to 54	91% (82-95%)	9.4% (4.8-18%)	7 (4.6-10)	23% (18-28%)	29% (22-34%)	250 (180-320)	9400 (7000-12000)
Latvia	2016	Male	55 to 59	89% (82-95%)	11% (5.3-18%)	6.9 (4.7-9.5)	21% (16-25%)	25% (20-30%)	290 (210-380)	9800 (7200-13000)
Latvia	2016	Male	60 to 64	89% (80-94%)	11% (5.9-20%)	6.1 (4.1-9.1)	17% (12-22%)	20% (14-26%)	280 (180-380)	8100 (5300-11000)
Latvia	2016	Male	65 to 69	88% (79-94%)	12% (6-21%)	4.8 (3.3-6.7)	13% (7.9-18%)	15% (8.8-20%)	190 (110-280)	4700 (2900-6600)
Latvia	2016	Male	70 to 74	85% (75-92%)	15% (8-25%)	3.5 (2.3-5)	7.6% (3-12%)	8.7% (3.1-14%)	150 (55-270)	3100 (1200-5100)
Latvia	2016	Male	75 to 79	81% (68-90%)	19% (9.9-32%)	2.9 (1.7-4.4)	4.7% (0.51-9.8%)	5.2% (0.2-11%)	100 (3.7-220)	1600 (170-3400)
Latvia	2016	Male	80 to 84	76% (61-86%)	24% (14-39%)	2.7 (1.6-4.1)	3.7% (-0.41-8.8%)	3.8% (-1.1-9.9%)	65 (-18-170)	790 (-88-2000)
Latvia	2016	Male	85 to 89	71% (55-82%)	29% (18-45%)	2.6 (1.5-4.1)	3.1% (-1-8.2%)	3.1% (-1.7-9.3%)	38 (-22-120)	340 (-110-930)
Latvia	2016	Male	90 to 94	68% (53-80%)	32% (20-47%)	2.6 (1.4-4.3)	2.4% (-1.9-8.4%)	2.4% (-2.8-9.4%)	9.6 (-11-39)	61 (-46-210)
Latvia	2016	Male	95 plus	66% (48-81%)	34% (19-52%)	2.5 (1.2-4.5)	2.2% (-2.4-8.5%)	1.9% (-3.6-9.5%)	1.7 (-3.1-8.3)	7.6 (-8.5-30)
Lithuania	1990	Female	All Ages	73% (66-78%)	27% (22-34%)	1.6 (1.3-1.9)	3.8% (2.4-5.1%)	0.6% (-2-3.2%)	110 (-390-600)	22000 (14000-31000)
Lithuania	1990	Female	15 to 19	68% (54-81%)	32% (19-46%)	1.7 (1.1-2.5)	4.8% (3.3-6.6%)	8.7% (5-13%)	6.3 (3.6-9.4)	750 (500-1000)
Lithuania	1990	Female	20 to 24	78% (64-87%)	22% (13-36%)	2.3 (1.5-3.2)	7.6% (5.7-9.7%)	11% (6.7-15%)	7.6 (4.8-11)	1300 (870-1700)
Lithuania	1990	Female	25 to 29	81% (71-89%)	19% (11-29%)	2.2 (1.5-2.9)	8% (6.3-9.7%)	11% (7.5-14%)	10 (7-14)	1600 (1200-2100)
Lithuania	1990	Female	30 to 34	84% (73-91%)	16% (8.6-27%)	2.2 (1.6-3)	8.7% (7.1-10%)	13% (9.8-17%)	18 (13-23)	2100 (1600-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	1990	Female	35 to 39	85% (76-92%)	15% (8-24%)	2.5 (1.8-3.3)	11% (8.9-13%)	16% (12-20%)	31 (24-38)	2700 (2200-3300)
Lithuania	1990	Female	40 to 44	85% (75-92%)	15% (7.8-25%)	2.5 (1.8-3.4)	12% (9.5-14%)	17% (13-20%)	49 (37-60)	3400 (2700-4100)
Lithuania	1990	Female	45 to 49	84% (71-92%)	16% (7.9-29%)	2.2 (1.6-3)	10% (8-12%)	14% (11-18%)	59 (46-76)	3500 (2700-4300)
Lithuania	1990	Female	50 to 54	82% (70-90%)	18% (9.9-30%)	2.1 (1.4-2.8)	8.1% (6-10%)	11% (7.7-14%)	64 (45-84)	3200 (2400-4100)
Lithuania	1990	Female	55 to 59	78% (63-88%)	22% (12-37%)	2.1 (1.4-2.9)	6.7% (4.5-9.1%)	8.8% (5.6-12%)	78 (49-110)	3200 (2200-4300)
Lithuania	1990	Female	60 to 64	74% (58-86%)	26% (14-42%)	1.9 (1.3-2.8)	4.1% (1.8-6.6%)	5% (1.7-8.6%)	64 (21-110)	2300 (1000-3700)
Lithuania	1990	Female	65 to 69	70% (54-84%)	30% (16-46%)	1.7 (1.1-2.4)	1.4% (-1.1-3.7%)	1.3% (-2.1-4.5%)	21 (-36-75)	790 (-590-2100)
Lithuania	1990	Female	70 to 74	65% (48-79%)	35% (21-52%)	1.4 (0.86-2.1)	-0.42% (-3.2-2.3%)	-0.99% (-4.6-2.5%)	-16 (-73-41)	-170 (-1300-960)
Lithuania	1990	Female	75 to 79	57% (41-74%)	43% (26-59%)	1.2 (0.67-1.9)	-1.5% (-4-1.2%)	-2.2% (-5.5-1.1%)	-61 (-150-30)	-760 (-2100-620)
Lithuania	1990	Female	80 to 84	51% (34-69%)	49% (31-66%)	1.1 (0.62-1.7)	-1.9% (-4.7-0.54%)	-2.6% (-6.1-0.44%)	-89 (-210-15)	-850 (-2100-240)
Lithuania	1990	Female	85 to 89	49% (32-65%)	51% (35-68%)	1.1 (0.62-1.7)	-1.9% (-4.6-0.46%)	-2.6% (-5.9-0.35%)	-72 (-160-9.7)	-480 (-1200-110)
Lithuania	1990	Female	90 to 94	50% (34-67%)	50% (33-66%)	1.2 (0.63-1.8)	-1.9% (-4.7-1%)	-2.6% (-6.3-0.94%)	-37 (-89-13)	-170 (-420-91)
Lithuania	1990	Female	95 plus	53% (34-72%)	47% (28-66%)	1.3 (0.62-2.1)	-2% (-5.3-2%)	-2.7% (-6.8-1.9%)	-21 (-52-15)	-60 (-160-58)
Lithuania	1990	Male	All Ages	80% (75-85%)	20% (15-25%)	4 (3.4-4.7)	15% (13-18%)	13% (9.5-17%)	2700 (2000-3400)	110000 (95000-130000)
Lithuania	1990	Male	15 to 19	71% (55-84%)	29% (16-45%)	2.4 (1.5-3.4)	9.1% (6-12%)	13% (8.2-19%)	25 (16-35)	2100 (1400-2800)
Lithuania	1990	Male	20 to 24	83% (71-91%)	17% (9.2-29%)	4.2 (3.1-5.7)	18% (13-23%)	23% (16-29%)	73 (52-95)	5700 (4200-7300)
Lithuania	1990	Male	25 to 29	88% (80-94%)	12% (6.5-20%)	4.6 (3.4-6.2)	21% (16-26%)	26% (19-33%)	110 (81-140)	8200 (6500-10000)
Lithuania	1990	Male	30 to 34	91% (84-96%)	9.2% (4.3-16%)	5.4 (4-7)	24% (19-29%)	30% (23-36%)	160 (120-190)	10000 (8400-13000)
Lithuania	1990	Male	35 to 39	92% (86-96%)	8.3% (3.9-14%)	6.5 (4.6-8.6)	27% (22-32%)	33% (27-40%)	210 (170-250)	12000 (9900-14000)
Lithuania	1990	Male	40 to 44	92% (85-96%)	8% (3.7-15%)	7.6 (5.7-9.8)	28% (24-33%)	35% (29-40%)	280 (230-320)	14000 (12000-16000)
Lithuania	1990	Male	45 to 49	92% (83-96%)	8.5% (3.9-17%)	8.7 (6.2-11)	28% (23-33%)	34% (28-39%)	370 (310-440)	17000 (14000-19000)
Lithuania	1990	Male	50 to 54	90% (82-95%)	9.7% (4.7-18%)	7.9 (5.5-11)	24% (19-28%)	28% (23-34%)	410 (340-500)	16000 (13000-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	1990	Male	55 to 59	89% (79-94%)	11% (5.7-21%)	5.4 (4-7.1)	17% (12-21%)	19% (14-24%)	360 (260-450)	12000 (9200-15000)
Lithuania	1990	Male	60 to 64	87% (76-94%)	13% (6.4-24%)	4.2 (2.8-5.7)	11% (6.3-16%)	12% (6.7-18%)	270 (150-390)	7900 (4600-11000)
Lithuania	1990	Male	65 to 69	85% (74-93%)	15% (6.8-26%)	3.9 (2.8-5.2)	8.1% (3.9-13%)	9% (3.9-14%)	190 (82-290)	4600 (2200-7100)
Lithuania	1990	Male	70 to 74	83% (71-91%)	17% (9.4-29%)	3.6 (2.3-4.9)	5.9% (1.3-11%)	6.3% (0.85-12%)	99 (13-190)	2000 (440-3700)
Lithuania	1990	Male	75 to 79	79% (66-89%)	21% (11-34%)	3.2 (2-4.7)	4% (-0.56-9.3%)	4.1% (-1.2-10%)	81 (-24-200)	1300 (-190-3100)
Lithuania	1990	Male	80 to 84	76% (61-86%)	24% (14-39%)	2.9 (1.9-4)	2.2% (-2.1-7.2%)	2% (-3.1-7.9%)	50 (-76-200)	690 (-640-2200)
Lithuania	1990	Male	85 to 89	72% (57-84%)	28% (16-43%)	2.6 (1.7-3.8)	1% (-2.7-5.7%)	0.59% (-3.9-6.1%)	8.6 (-56-88)	130 (-350-730)
Lithuania	1990	Male	90 to 94	71% (57-83%)	29% (17-43%)	2.4 (1.5-3.7)	0.19% (-3.7-5.4%)	-0.46% (-5.2-5.8%)	-2.6 (-29-32)	6.7 (-120-190)
Lithuania	1990	Male	95 plus	69% (51-84%)	31% (16-49%)	2.2 (1.1-3.8)	-0.23% (-4.5-8.8%)	-1% (-5.7-6.1%)	-3.2 (-18-19)	-2.9 (-52-73)
Lithuania	1995	Female	All Ages	73% (67-79%)	27% (21-33%)	1.5 (1.3-1.7)	4% (2.5-5.4%)	0.75% (-1.8-3.3%)	160 (-390-710)	25000 (16000-35000)
Lithuania	1995	Female	15 to 19	69% (55-81%)	31% (19-45%)	1.5 (0.99-2.3)	4.8% (3.3-6.6%)	8.7% (5.1-13%)	5.9 (3.4-8.9)	680 (470-950)
Lithuania	1995	Female	20 to 24	78% (65-88%)	22% (12-35%)	2.1 (1.4-2.9)	7.8% (5.8-9.8%)	11% (6.8-15%)	9.3 (5.8-13)	1300 (960-1800)
Lithuania	1995	Female	25 to 29	82% (70-90%)	18% (10-30%)	2 (1.4-2.7)	8.4% (6.7-10%)	11% (7.7-15%)	12 (8.2-16)	1600 (1200-2100)
Lithuania	1995	Female	30 to 34	84% (73-92%)	16% (8.1-27%)	2 (1.5-2.7)	9.4% (7.6-11%)	14% (11-18%)	23 (18-29)	2400 (1800-2900)
Lithuania	1995	Female	35 to 39	85% (75-92%)	15% (7.9-25%)	2.3 (1.6-3)	11% (9.3-13%)	17% (13-21%)	47 (36-57)	3500 (2800-4200)
Lithuania	1995	Female	40 to 44	86% (76-92%)	14% (7.5-24%)	2.3 (1.6-3)	12% (10-15%)	18% (14-21%)	67 (53-81)	4000 (3300-4800)
Lithuania	1995	Female	45 to 49	85% (71-92%)	15% (7.9-29%)	2 (1.4-2.7)	11% (8.6-13%)	15% (12-19%)	82 (62-100)	4100 (3300-5000)
Lithuania	1995	Female	50 to 54	82% (70-90%)	18% (9.8-30%)	1.9 (1.3-2.5)	9.1% (7.1-11%)	13% (9.6-16%)	89 (68-110)	3900 (3100-4800)
Lithuania	1995	Female	55 to 59	78% (65-89%)	22% (11-35%)	1.9 (1.3-2.7)	7.4% (5.3-9.6%)	10% (7-13%)	100 (70-130)	3700 (2700-4900)
Lithuania	1995	Female	60 to 64	75% (59-86%)	25% (14-41%)	1.8 (1.2-2.6)	4.2% (1.9-6.5%)	5.3% (2.2-8.6%)	74 (30-120)	2400 (1100-3800)
Lithuania	1995	Female	65 to 69	71% (54-85%)	29% (15-46%)	1.6 (1.1-2.2)	1.2% (-0.96-3.6%)	1.2% (-1.9-4.5%)	24 (-38-89)	820 (-670-2400)
Lithuania	1995	Female	70 to 74	66% (49-79%)	34% (21-51%)	1.3 (0.81-1.9)	-0.67% (-3.2-1.9%)	-1.2% (-4.4-2.1%)	-31 (-120-55)	-430 (-2100-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	1995	Female	75 to 79	59% (42-75%)	41% (25-58%)	1.1 (0.63-1.8)	-1.7% (-4.3-0.81%)	-2.4% (-5.6-0.79%)	-56 (-130-18)	-740 (-1800-330)
Lithuania	1995	Female	80 to 84	53% (34-71%)	47% (29-66%)	1 (0.59-1.6)	-2.2% (-4.9-0.27%)	-2.9% (-6.3-0.19%)	-100 (-230-6.5)	-1000 (-2300-130)
Lithuania	1995	Female	85 to 89	50% (33-66%)	50% (34-67%)	1 (0.58-1.6)	-2.4% (-5.1-0.15%)	-3% (-6.3-0.061%)	-100 (-220-2)	-720 (-1500-43)
Lithuania	1995	Female	90 to 94	51% (35-68%)	49% (32-65%)	1.1 (0.6-1.7)	-2.4% (-5.4-0.54%)	-3.1% (-6.7-0.4%)	-53 (-120-6.8)	-250 (-560-56)
Lithuania	1995	Female	95 plus	54% (35-72%)	46% (28-65%)	1.2 (0.59-2)	-2.5% (-6.1-1.3%)	-3.2% (-7.5-1.2%)	-26 (-59-9.8)	-79 (-190-39)
Lithuania	1995	Male	All Ages	81% (76-86%)	19% (14-24%)	3.7 (3.2-4.3)	16% (14-19%)	14% (10-17%)	3300 (2500-4100)	140000 (120000-160000)
Lithuania	1995	Male	15 to 19	71% (56-84%)	29% (16-44%)	2.2 (1.4-3.1)	9.2% (6-13%)	13% (8-19%)	25 (16-37)	2100 (1400-2900)
Lithuania	1995	Male	20 to 24	83% (72-91%)	17% (9-28%)	3.9 (2.8-5.1)	18% (14-24%)	22% (16-29%)	89 (63-120)	6600 (4900-8600)
Lithuania	1995	Male	25 to 29	88% (80-93%)	12% (6.5-20%)	4.2 (3.1-5.6)	21% (16-26%)	26% (19-32%)	120 (93-160)	8800 (6700-11000)
Lithuania	1995	Male	30 to 34	91% (83-96%)	9.2% (4.3-17%)	4.9 (3.7-6.4)	24% (20-29%)	29% (23-36%)	210 (160-250)	13000 (10000-16000)
Lithuania	1995	Male	35 to 39	92% (86-96%)	8.2% (4-14%)	5.9 (4.2-7.8)	28% (23-33%)	33% (27-40%)	310 (240-370)	17000 (14000-20000)
Lithuania	1995	Male	40 to 44	92% (85-96%)	8% (3.9-15%)	6.9 (5.2-8.9)	30% (25-34%)	35% (29-40%)	410 (340-480)	20000 (17000-23000)
Lithuania	1995	Male	45 to 49	91% (83-96%)	8.5% (3.8-17%)	7.8 (5.8-10)	29% (25-34%)	34% (29-39%)	490 (420-570)	21000 (18000-24000)
Lithuania	1995	Male	50 to 54	90% (82-96%)	9.6% (4.4-18%)	7.1 (5-9.5)	25% (20-30%)	28% (23-34%)	540 (430-640)	20000 (16000-24000)
Lithuania	1995	Male	55 to 59	89% (80-95%)	11% (5.4-20%)	5 (3.7-6.5)	17% (12-21%)	19% (14-24%)	430 (310-550)	14000 (11000-18000)
Lithuania	1995	Male	60 to 64	87% (76-94%)	13% (6.4-24%)	3.8 (2.6-5.3)	10% (5.7-15%)	11% (6.2-17%)	300 (170-460)	8700 (4900-13000)
Lithuania	1995	Male	65 to 69	86% (75-93%)	14% (6.9-25%)	3.6 (2.6-4.7)	7.1% (3.1-11%)	7.7% (3-13%)	230 (88-370)	5500 (2400-8900)
Lithuania	1995	Male	70 to 74	83% (71-91%)	17% (9.4-29%)	3.2 (2.2-4.4)	4.7% (0.57-9.2%)	5% (0.15-10%)	130 (3.8-260)	2600 (320-5100)
Lithuania	1995	Male	75 to 79	80% (66-89%)	20% (11-34%)	2.9 (1.9-4.3)	3% (-1.2-7.8%)	3% (-1.9-8.6%)	51 (-33-150)	860 (-330-2300)
Lithuania	1995	Male	80 to 84	76% (62-87%)	24% (13-38%)	2.6 (1.8-3.7)	1.2% (-2.7-5.8%)	0.9% (-3.6-6.1%)	17 (-67-110)	280 (-610-1300)
Lithuania	1995	Male	85 to 89	73% (59-84%)	27% (16-41%)	2.4 (1.6-3.5)	0.041% (-3.4-4.6%)	-0.53% (-4.7-4.8%)	-9.6 (-85-87)	6.7 (-540-710)
Lithuania	1995	Male	90 to 94	71% (57-83%)	29% (17-43%)	2.2 (1.4-3.4)	-0.65% (-4.3-3.9%)	-1.4% (-5.8-4.1%)	-10 (-41-29)	-28 (-190-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	1995	Male	95 plus	69% (53-84%)	31% (16-47%)	2 (1-3.5)	-0.95% (-4.7-4.2%)	-1.8% (-6.4-4.4%)	-4.4 (-15-11)	-9.1 (-45-41)
Lithuania	2000	Female	All Ages	72% (66-78%)	28% (22-34%)	1.7 (1.5-2)	3.9% (2.7-5.3%)	1.2% (-1.2-3.8%)	230 (-230-700)	22000 (15000-30000)
Lithuania	2000	Female	15 to 19	67% (53-81%)	33% (19-47%)	1.7 (1.1-2.5)	4.7% (3.2-6.5%)	9.2% (5.1-14%)	5.3 (2.9-8.1)	650 (430-920)
Lithuania	2000	Female	20 to 24	77% (63-87%)	23% (13-37%)	2.4 (1.6-3.3)	7.8% (6-9.8%)	11% (7.2-16%)	7.4 (4.6-11)	1100 (800-1500)
Lithuania	2000	Female	25 to 29	81% (70-89%)	19% (11-30%)	2.2 (1.6-3)	8.1% (6.5-10%)	11% (7.6-15%)	9.3 (6.2-12)	1400 (1000-1800)
Lithuania	2000	Female	30 to 34	84% (72-92%)	16% (8.2-28%)	2.3 (1.7-3.1)	8.8% (7.1-11%)	13% (9.7-17%)	16 (12-20)	1800 (1400-2300)
Lithuania	2000	Female	35 to 39	85% (76-92%)	15% (7.8-24%)	2.6 (1.9-3.5)	10% (8.2-12%)	16% (12-19%)	31 (24-39)	2700 (2100-3300)
Lithuania	2000	Female	40 to 44	85% (76-92%)	15% (7.7-24%)	2.6 (1.9-3.5)	10% (8.3-12%)	16% (13-19%)	46 (36-56)	3100 (2500-3800)
Lithuania	2000	Female	45 to 49	84% (71-92%)	16% (8-29%)	2.3 (1.6-3.1)	9.4% (7.5-11%)	14% (11-17%)	57 (44-69)	3100 (2500-3800)
Lithuania	2000	Female	50 to 54	81% (69-90%)	19% (10-31%)	2.1 (1.5-2.9)	8.6% (6.7-11%)	12% (9.6-15%)	65 (51-81)	3000 (2400-3700)
Lithuania	2000	Female	55 to 59	77% (63-88%)	23% (12-37%)	2.2 (1.5-3.1)	7.4% (5.5-9.6%)	11% (7.6-14%)	79 (57-100)	3000 (2200-4000)
Lithuania	2000	Female	60 to 64	74% (58-86%)	26% (14-42%)	2 (1.3-2.9)	4.8% (2.7-7.2%)	6.6% (3.5-10%)	70 (38-110)	2300 (1300-3500)
Lithuania	2000	Female	65 to 69	70% (53-84%)	30% (16-47%)	1.8 (1.2-2.5)	2.2% (0.011-4.4%)	2.6% (-0.5-5.8%)	40 (-7.6-90)	1200 (6.5-2400)
Lithuania	2000	Female	70 to 74	64% (47-79%)	36% (21-53%)	1.5 (0.89-2.2)	0.3% (-2.2-2.9%)	0.047% (-3.2-3.5%)	1.1 (-77-84)	190 (-1400-1800)
Lithuania	2000	Female	75 to 79	56% (39-73%)	44% (27-61%)	1.2 (0.69-2)	-0.77% (-3.2-1.8%)	-1.3% (-4.4-2%)	-39 (-130-59)	-450 (-1800-1000)
Lithuania	2000	Female	80 to 84	50% (31-67%)	50% (33-69%)	1.1 (0.63-1.7)	-1.2% (-3.7-1.1%)	-1.8% (-5-1.2%)	-45 (-120-29)	-420 (-1200-390)
Lithuania	2000	Female	85 to 89	47% (30-64%)	53% (36-70%)	1.1 (0.61-1.7)	-1.5% (-4-0.92%)	-2% (-5.2-0.92%)	-59 (-150-27)	-390 (-1100-250)
Lithuania	2000	Female	90 to 94	48% (32-65%)	52% (35-68%)	1.2 (0.65-1.9)	-1.6% (-4.5-1.4%)	-2.2% (-5.7-1.4%)	-39 (-100-24)	-170 (-480-140)
Lithuania	2000	Female	95 plus	52% (31-72%)	48% (28-69%)	1.3 (0.61-2.2)	-1.7% (-5.3-2.3%)	-2.3% (-6.5-2.5%)	-15 (-44-17)	-44 (-140-61)
Lithuania	2000	Male	All Ages	81% (75-86%)	19% (14-25%)	4.4 (3.9-5)	16% (14-19%)	15% (11-18%)	3000 (2300-3700)	120000 (100000-140000)
Lithuania	2000	Male	15 to 19	70% (54-83%)	30% (17-46%)	2.5 (1.6-3.5)	9.5% (6.1-13%)	14% (8.4-20%)	24 (15-36)	2000 (1300-2800)
Lithuania	2000	Male	20 to 24	82% (69-91%)	18% (9.3-31%)	4.5 (3.2-5.8)	20% (14-25%)	25% (17-33%)	73 (51-98)	5500 (4000-7200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	2000	Male	25 to 29	87% (79-93%)	13% (6.6-21%)	4.9 (3.6-6.5)	22% (17-27%)	27% (20-35%)	98 (72-130)	7000 (5400-8800)
Lithuania	2000	Male	30 to 34	90% (82-96%)	9.5% (4.5-18%)	5.7 (4.3-7.4)	24% (20-29%)	30% (24-37%)	150 (110-180)	9400 (7500-11000)
Lithuania	2000	Male	35 to 39	92% (85-96%)	8.4% (4.1-15%)	6.9 (5-9.1)	27% (22-32%)	35% (28-41%)	230 (190-270)	13000 (11000-15000)
Lithuania	2000	Male	40 to 44	92% (85-96%)	8.2% (3.9-15%)	8 (6.1-10)	29% (24-33%)	36% (31-41%)	320 (270-370)	16000 (14000-18000)
Lithuania	2000	Male	45 to 49	91% (82-96%)	8.8% (3.9-18%)	9.1 (6.8-12)	28% (24-33%)	34% (29-39%)	380 (330-450)	17000 (14000-19000)
Lithuania	2000	Male	50 to 54	90% (81-95%)	10% (4.6-19%)	8.3 (5.7-11)	25% (21-29%)	30% (25-34%)	400 (340-480)	15000 (13000-18000)
Lithuania	2000	Male	55 to 59	88% (78-94%)	12% (5.5-22%)	5.7 (4.4-7.4)	18% (14-22%)	21% (17-26%)	370 (280-450)	12000 (9600-15000)
Lithuania	2000	Male	60 to 64	87% (75-94%)	13% (6.3-25%)	4.4 (3-6.2)	12% (7.5-17%)	14% (8.2-19%)	300 (180-420)	8600 (5300-12000)
Lithuania	2000	Male	65 to 69	85% (73-93%)	15% (7.2-27%)	4.1 (3-5.5)	9.1% (5.1-13%)	10% (5.4-15%)	260 (140-390)	6400 (3500-9300)
Lithuania	2000	Male	70 to 74	82% (69-90%)	18% (10-31%)	3.7 (2.5-5.2)	6.7% (2.3-11%)	7.4% (2.2-13%)	200 (58-340)	4000 (1400-6800)
Lithuania	2000	Male	75 to 79	79% (64-89%)	21% (11-36%)	3.3 (2.2-4.9)	5% (0.51-9.9%)	5.3% (0.073-11%)	120 (1.6-250)	1900 (190-3700)
Lithuania	2000	Male	80 to 84	75% (58-86%)	25% (14-42%)	3 (2-4.3)	3.3% (-0.73-8.2%)	3.3% (-1.4-9.2%)	45 (-20-130)	560 (-120-1400)
Lithuania	2000	Male	85 to 89	71% (55-83%)	29% (17-45%)	2.7 (1.8-4)	1.9% (-1.9-6.7%)	1.6% (-2.9-7.3%)	18 (-32-82)	190 (-190-660)
Lithuania	2000	Male	90 to 94	69% (55-82%)	31% (18-45%)	2.5 (1.5-3.8)	0.79% (-3.1-5.9%)	0.29% (-4.4-6.5%)	2.4 (-35-53)	40 (-150-300)
Lithuania	2000	Male	95 plus	68% (50-83%)	32% (17-50%)	2.3 (1.2-3.9)	0.16% (-3.9-6.2%)	-0.54% (-5.5-6.7%)	-1.5 (-15-19)	1.8 (-43-70)
Lithuania	2005	Female	All Ages	74% (69-79%)	26% (21-31%)	2.1 (1.8-2.5)	4.8% (3.3-6.4%)	2.5% (0.018-5.3%)	520 (3.6-1100)	28000 (19000-37000)
Lithuania	2005	Female	15 to 19	71% (58-82%)	29% (18-42%)	2.1 (1.5-3.1)	5.2% (3.7-7.1%)	11% (6.2-16%)	5.7 (3.3-8.5)	680 (470-940)
Lithuania	2005	Female	20 to 24	80% (69-88%)	20% (12-31%)	3 (2-4.1)	8.3% (6.2-11%)	13% (8.6-19%)	6.8 (4.4-9.5)	1100 (780-1500)
Lithuania	2005	Female	25 to 29	83% (74-90%)	17% (9.5-26%)	2.8 (2-3.7)	8.8% (7-11%)	13% (9.3-18%)	10 (7.2-14)	1300 (1000-1700)
Lithuania	2005	Female	30 to 34	86% (76-93%)	14% (7.5-24%)	2.8 (2-3.8)	9.2% (7.4-11%)	15% (11-19%)	17 (13-22)	1700 (1400-2200)
Lithuania	2005	Female	35 to 39	86% (78-93%)	14% (7.2-22%)	3.1 (2.3-4.1)	10% (8.6-12%)	17% (14-21%)	33 (26-40)	2500 (2000-3000)
Lithuania	2005	Female	40 to 44	87% (77-93%)	13% (7.1-23%)	3.1 (2.3-4.2)	11% (9.3-13%)	18% (15-21%)	57 (47-68)	3500 (2900-4200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	2005	Female	45 to 49	86% (75-93%)	14% (7.3-25%)	2.8 (2.1-3.7)	10% (8.5-13%)	16% (13-19%)	77 (62-93)	4000 (3300-4800)
Lithuania	2005	Female	50 to 54	84% (73-90%)	16% (9.5-27%)	2.7 (1.9-3.6)	10% (7.9-12%)	15% (12-18%)	88 (70-110)	3800 (3100-4700)
Lithuania	2005	Female	55 to 59	79% (66-89%)	21% (11-34%)	2.6 (1.9-3.7)	9% (6.7-12%)	13% (9.6-17%)	100 (77-130)	3700 (2800-4800)
Lithuania	2005	Female	60 to 64	75% (61-86%)	25% (14-39%)	2.5 (1.7-3.5)	6.5% (4-9.4%)	9.1% (5.4-13%)	98 (58-140)	3100 (1900-4500)
Lithuania	2005	Female	65 to 69	69% (55-84%)	31% (16-45%)	2 (1.4-2.8)	3.3% (0.95-5.8%)	4.4% (1.1-7.9%)	67 (16-120)	1900 (540-3300)
Lithuania	2005	Female	70 to 74	63% (48-77%)	37% (23-52%)	1.6 (1.1-2.4)	1.3% (-1.2-4.3%)	1.5% (-1.9-5.3%)	34 (-46-120)	830 (-740-2700)
Lithuania	2005	Female	75 to 79	55% (40-69%)	45% (31-60%)	1.4 (0.86-2.2)	0.086% (-2.5-3%)	-0.17% (-3.4-3.5%)	-6 (-120-120)	60 (-1700-2000)
Lithuania	2005	Female	80 to 84	46% (29-63%)	54% (37-71%)	1.2 (0.68-1.8)	-0.33% (-2.5-2.2%)	-0.67% (-3.6-2.5%)	-27 (-140-99)	-180 (-1400-1200)
Lithuania	2005	Female	85 to 89	43% (27-59%)	57% (41-73%)	1.1 (0.65-1.8)	-0.49% (-2.9-2.1%)	-0.86% (-3.7-2.3%)	-22 (-94-58)	-110 (-660-480)
Lithuania	2005	Female	90 to 94	44% (29-61%)	56% (39-71%)	1.2 (0.68-2)	-0.62% (-3.4-2.7%)	-1% (-4.4-2.9%)	-19 (-81-53)	-69 (-390-300)
Lithuania	2005	Female	95 plus	48% (28-68%)	52% (32-72%)	1.4 (0.66-2.4)	-0.56% (-4.3-3.9%)	-0.98% (-5.3-4.3%)	-7 (-38-30)	-16 (-120-110)
Lithuania	2005	Male	All Ages	84% (79-88%)	16% (12-21%)	5.8 (5.1-6.5)	19% (17-22%)	18% (14-22%)	4200 (3300-5100)	150000 (130000-180000)
Lithuania	2005	Male	15 to 19	74% (62-85%)	26% (15-38%)	3.2 (2.2-4.4)	11% (7.4-15%)	17% (11-24%)	28 (18-40)	2200 (1500-3100)
Lithuania	2005	Male	20 to 24	86% (77-92%)	14% (7.7-23%)	5.7 (4.2-7.4)	22% (17-28%)	29% (21-37%)	84 (61-110)	6100 (4600-7700)
Lithuania	2005	Male	25 to 29	90% (84-94%)	10% (5.8-16%)	6.1 (4.6-8)	25% (20-30%)	31% (24-39%)	110 (83-140)	7400 (5800-9100)
Lithuania	2005	Male	30 to 34	92% (87-96%)	7.6% (3.8-13%)	7.1 (5.4-9.2)	28% (23-33%)	35% (28-41%)	170 (140-200)	11000 (8700-12000)
Lithuania	2005	Male	35 to 39	93% (88-97%)	6.9% (3.4-12%)	8.5 (6.3-11)	31% (26-35%)	38% (32-43%)	260 (220-290)	14000 (12000-16000)
Lithuania	2005	Male	40 to 44	93% (88-97%)	6.9% (3.5-12%)	9.9 (7.6-12)	32% (28-36%)	39% (34-43%)	410 (350-460)	20000 (17000-22000)
Lithuania	2005	Male	45 to 49	93% (86-97%)	7.2% (3.5-14%)	11 (8.5-14)	30% (26-35%)	36% (31-41%)	520 (440-600)	22000 (19000-26000)
Lithuania	2005	Male	50 to 54	92% (84-96%)	8.1% (4-16%)	10 (7.5-13)	28% (23-32%)	32% (27-37%)	540 (450-640)	20000 (17000-24000)
Lithuania	2005	Male	55 to 59	91% (83-95%)	9.5% (4.6-17%)	7.3 (5.6-9.2)	22% (18-26%)	25% (20-30%)	480 (390-580)	16000 (13000-19000)
Lithuania	2005	Male	60 to 64	89% (80-95%)	11% (5.4-20%)	5.6 (4.1-7.7)	15% (9.9-20%)	17% (11-23%)	410 (260-550)	12000 (7600-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	2005	Male	65 to 69	87% (78-94%)	13% (6.1-22%)	5.2 (3.9-6.8)	12% (7.6-17%)	14% (8.4-19%)	390 (230-540)	9400 (5800-13000)
Lithuania	2005	Male	70 to 74	84% (74-91%)	16% (9-26%)	4.6 (3.3-6.4)	9.5% (5-14%)	11% (5.2-16%)	320 (160-480)	6200 (3200-9300)
Lithuania	2005	Male	75 to 79	81% (69-90%)	19% (10-31%)	4.2 (2.8-5.8)	7.5% (2.5-13%)	8.3% (2.4-14%)	240 (69-420)	3700 (1300-6300)
Lithuania	2005	Male	80 to 84	78% (65-87%)	22% (13-35%)	3.8 (2.6-5.3)	5.8% (0.93-11%)	6.3% (0.44-13%)	130 (9.3-260)	1500 (240-2900)
Lithuania	2005	Male	85 to 89	75% (64-84%)	25% (16-36%)	3.5 (2.4-4.9)	4.3% (-0.52-9.7%)	4.4% (-1.3-11%)	43 (-13-110)	380 (-47-840)
Lithuania	2005	Male	90 to 94	73% (60-84%)	27% (16-40%)	3.2 (2.1-4.8)	2.7% (-2.4-8.7%)	2.5% (-3.7-9.7%)	14 (-21-55)	93 (-82-300)
Lithuania	2005	Male	95 plus	71% (54-85%)	29% (15-46%)	2.9 (1.5-4.9)	1.7% (-3.5-8.6%)	1.2% (-4.9-9.5%)	3.9 (-16-31)	22 (-45-110)
Lithuania	2010	Female	All Ages	73% (67-79%)	27% (21-33%)	2.4 (2-2.8)	5% (3.6-6.7%)	3.1% (0.49-6.1%)	610 (100-1200)	27000 (19000-37000)
Lithuania	2010	Female	15 to 19	70% (57-81%)	30% (19-43%)	2.4 (1.6-3.4)	4.9% (3.5-6.5%)	11% (6.6-17%)	3.5 (2.1-5.2)	510 (350-690)
Lithuania	2010	Female	20 to 24	80% (69-88%)	20% (12-31%)	3.3 (2.3-4.6)	8.4% (6.4-11%)	14% (9.1-19%)	6 (3.9-8.2)	1000 (730-1400)
Lithuania	2010	Female	25 to 29	83% (74-90%)	17% (10-26%)	3.1 (2.2-4.1)	8.6% (6.9-10%)	13% (9.4-18%)	7.4 (5.2-10)	1100 (820-1500)
Lithuania	2010	Female	30 to 34	85% (76-92%)	15% (7.9-24%)	3.2 (2.3-4.3)	9.2% (7.5-11%)	16% (12-19%)	13 (10-17)	1400 (1100-1800)
Lithuania	2010	Female	35 to 39	86% (77-93%)	14% (7.4-23%)	3.5 (2.6-4.8)	10% (8.5-12%)	18% (15-22%)	26 (21-32)	2100 (1700-2500)
Lithuania	2010	Female	40 to 44	86% (77-93%)	14% (7.4-23%)	3.5 (2.6-4.7)	11% (9-13%)	19% (15-22%)	45 (37-54)	2800 (2300-3400)
Lithuania	2010	Female	45 to 49	86% (76-92%)	14% (7.5-24%)	3.2 (2.3-4.1)	10% (8.3-12%)	16% (14-19%)	67 (55-81)	3600 (2900-4300)
Lithuania	2010	Female	50 to 54	83% (73-91%)	17% (9.3-27%)	3 (2.1-3.9)	9.7% (7.7-12%)	15% (12-18%)	88 (70-110)	3900 (3200-4800)
Lithuania	2010	Female	55 to 59	79% (65-88%)	21% (12-35%)	3 (2.1-4.1)	9.3% (6.9-12%)	14% (10-17%)	99 (76-130)	3700 (2900-4800)
Lithuania	2010	Female	60 to 64	74% (60-86%)	26% (14-40%)	2.8 (1.8-3.9)	7.2% (4.7-10%)	10% (6.7-15%)	95 (61-140)	3000 (2000-4300)
Lithuania	2010	Female	65 to 69	68% (53-83%)	32% (17-47%)	2.3 (1.6-3.1)	4.1% (1.8-6.9%)	5.5% (2.2-9.4%)	71 (27-120)	2000 (870-3300)
Lithuania	2010	Female	70 to 74	62% (47-76%)	38% (24-53%)	1.8 (1.2-2.7)	2.1% (-0.37-5.1%)	2.6% (-0.91-6.6%)	51 (-18-130)	1200 (-220-2900)
Lithuania	2010	Female	75 to 79	54% (39-69%)	46% (31-61%)	1.5 (0.93-2.3)	0.9% (-1.6-3.8%)	0.87% (-2.5-4.8%)	26 (-76-150)	560 (-1000-2400)
Lithuania	2010	Female	80 to 84	45% (28-62%)	55% (38-72%)	1.3 (0.75-2)	0.44% (-1.7-3.2%)	0.27% (-2.5-3.7%)	11 (-110-160)	260 (-1000-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	2010	Female	85 to 89	42% (26-57%)	58% (43-74%)	1.3 (0.72-2)	0.29% (-2.1-3.1%)	0.063% (-2.9-3.5%)	2.5 (-110-140)	110 (-750-1100)
Lithuania	2010	Female	90 to 94	43% (29-60%)	57% (40-71%)	1.4 (0.77-2.2)	0.25% (-2.6-4.2%)	-0.01% (-3.5-4.8%)	-0.15 (-56-75)	25 (-260-410)
Lithuania	2010	Female	95 plus	47% (28-67%)	53% (33-72%)	1.5 (0.76-2.7)	0.44% (-3.5-5.9%)	0.18% (-4.5-6.7%)	1.1 (-28-43)	11 (-85-150)
Lithuania	2010	Male	All Ages	84% (79-88%)	16% (12-21%)	6.7 (5.9-7.5)	20% (17-23%)	18% (15-22%)	3900 (3200-4800)	130000 (120000-160000)
Lithuania	2010	Male	15 to 19	74% (61-84%)	26% (16-39%)	3.6 (2.5-4.9)	11% (7.3-15%)	18% (12-25%)	21 (14-29)	1700 (1200-2400)
Lithuania	2010	Male	20 to 24	86% (76-92%)	14% (7.8-24%)	6.4 (4.7-8.2)	23% (17-28%)	32% (23-40%)	67 (49-85)	5000 (3800-6200)
Lithuania	2010	Male	25 to 29	90% (84-94%)	10% (5.8-16%)	6.9 (5.1-9.1)	24% (19-29%)	33% (25-40%)	80 (62-99)	5700 (4500-6900)
Lithuania	2010	Male	30 to 34	92% (86-96%)	7.8% (3.8-14%)	8 (6.1-10)	27% (23-32%)	36% (30-41%)	120 (100-140)	7800 (6500-9100)
Lithuania	2010	Male	35 to 39	93% (88-96%)	7.1% (3.5-12%)	9.6 (7-13)	30% (26-35%)	39% (34-44%)	200 (170-230)	11000 (9600-13000)
Lithuania	2010	Male	40 to 44	93% (88-96%)	7.1% (3.6-12%)	11 (8.6-14)	31% (27-35%)	39% (35-43%)	290 (260-330)	15000 (13000-16000)
Lithuania	2010	Male	45 to 49	93% (86-96%)	7.4% (3.6-14%)	13 (9.5-16)	30% (25-34%)	36% (31-41%)	420 (360-480)	18000 (16000-21000)
Lithuania	2010	Male	50 to 54	92% (85-96%)	8.2% (4.2-15%)	12 (8.4-15)	27% (23-32%)	33% (28-38%)	510 (420-590)	19000 (16000-23000)
Lithuania	2010	Male	55 to 59	90% (82-95%)	9.8% (4.9-18%)	8.2 (6.4-10)	23% (19-27%)	26% (22-31%)	470 (400-570)	16000 (13000-19000)
Lithuania	2010	Male	60 to 64	89% (81-95%)	11% (5.5-19%)	6.4 (4.5-8.8)	17% (12-21%)	20% (14-25%)	410 (290-520)	12000 (8300-15000)
Lithuania	2010	Male	65 to 69	87% (78-94%)	13% (6.1-22%)	5.9 (4.4-7.7)	14% (10-18%)	16% (11-21%)	390 (270-510)	9300 (6500-12000)
Lithuania	2010	Male	70 to 74	83% (73-91%)	17% (8.6-27%)	5.2 (3.7-7.1)	11% (6.3-15%)	13% (7.2-18%)	340 (190-480)	6600 (3900-9200)
Lithuania	2010	Male	75 to 79	80% (68-89%)	20% (11-32%)	4.7 (3.1-6.5)	8.9% (4-14%)	10% (4.2-16%)	280 (120-440)	4300 (1900-6600)
Lithuania	2010	Male	80 to 84	77% (64-87%)	23% (13-36%)	4.3 (2.9-5.9)	7.5% (2.9-13%)	8.4% (2.9-15%)	210 (69-370)	2400 (920-4100)
Lithuania	2010	Male	85 to 89	75% (62-84%)	25% (16-38%)	3.9 (2.6-5.5)	6% (0.87-11%)	6.5% (0.27-13%)	93 (3.7-180)	770 (110-1400)
Lithuania	2010	Male	90 to 94	73% (59-84%)	27% (16-41%)	3.6 (2.3-5.4)	4.4% (-1.2-10%)	4.6% (-2.2-12%)	24 (-11-61)	140 (-40-330)
Lithuania	2010	Male	95 plus	71% (54-85%)	29% (15-46%)	3.2 (1.7-5.5)	3.1% (-2.9-10%)	2.9% (-4.2-11%)	5.6 (-8.2-21)	24 (-23-79)
Lithuania	2016	Female	All Ages	71% (65-77%)	29% (23-35%)	2.5 (2-3.1)	5% (3.2-7%)	3.3% (0.066-7.2%)	670 (13-1400)	26000 (16000-36000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	2016	Female	15 to 19	66% (51-80%)	34% (20-49%)	2.4 (1.5-3.7)	4.7% (3.3-6.4%)	11% (6.3-16%)	2.2 (1.2-3.2)	330 (230-460)
Lithuania	2016	Female	20 to 24	77% (63-87%)	23% (13-37%)	3.4 (2.2-5)	8.3% (6.2-11%)	14% (8.7-20%)	4.3 (2.6-6.2)	890 (620-1200)
Lithuania	2016	Female	25 to 29	81% (70-89%)	19% (11-30%)	3.2 (2.2-4.5)	8.7% (6.8-11%)	14% (9.2-18%)	6.2 (4.1-8.4)	1000 (740-1300)
Lithuania	2016	Female	30 to 34	84% (71-91%)	16% (8.6-29%)	3.3 (2.2-4.7)	9.2% (7.4-11%)	16% (12-20%)	10 (7.5-13)	1100 (870-1500)
Lithuania	2016	Female	35 to 39	85% (75-92%)	15% (8.1-25%)	3.7 (2.6-5.2)	10% (8.4-12%)	19% (15-23%)	19 (14-24)	1500 (1200-1900)
Lithuania	2016	Female	40 to 44	85% (75-92%)	15% (7.7-25%)	3.7 (2.6-5.2)	11% (8.4-13%)	19% (15-23%)	35 (27-44)	2300 (1800-2800)
Lithuania	2016	Female	45 to 49	84% (72-92%)	16% (8.2-28%)	3.3 (2.3-4.7)	10% (7.9-12%)	17% (14-21%)	50 (38-62)	2700 (2100-3300)
Lithuania	2016	Female	50 to 54	81% (69-90%)	19% (10-31%)	3.1 (2.1-4.4)	9.6% (7.5-12%)	16% (12-19%)	81 (63-100)	3700 (2900-4600)
Lithuania	2016	Female	55 to 59	77% (63-88%)	23% (12-37%)	3.1 (2.1-4.6)	9.1% (6.8-12%)	14% (10-18%)	95 (70-120)	3600 (2700-4800)
Lithuania	2016	Female	60 to 64	73% (56-86%)	27% (14-44%)	2.9 (1.8-4.4)	7.4% (4.7-11%)	11% (6.9-15%)	95 (59-140)	3100 (2000-4400)
Lithuania	2016	Female	65 to 69	69% (51-85%)	31% (15-49%)	2.5 (1.7-3.6)	4.7% (2.2-7.6%)	6.4% (2.7-11%)	63 (26-110)	1800 (840-2900)
Lithuania	2016	Female	70 to 74	63% (45-78%)	37% (22-55%)	2.1 (1.2-3.2)	2.7% (-0.089-6.1%)	3.5% (-0.49-8.2%)	57 (-8-130)	1300 (-44-3000)
Lithuania	2016	Female	75 to 79	55% (37-71%)	45% (29-63%)	1.7 (0.93-2.8)	1.5% (-1.1-4.9%)	1.7% (-2-6.4%)	46 (-53-170)	890 (-620-2900)
Lithuania	2016	Female	80 to 84	48% (31-66%)	52% (34-69%)	1.5 (0.86-2.5)	1.1% (-1.5-4.4%)	1% (-2.4-5.2%)	42 (-98-210)	630 (-900-2600)
Lithuania	2016	Female	85 to 89	45% (29-62%)	55% (38-71%)	1.5 (0.85-2.5)	0.91% (-1.9-4.5%)	0.79% (-2.8-5.2%)	37 (-130-240)	400 (-850-1900)
Lithuania	2016	Female	90 to 94	47% (31-64%)	53% (36-69%)	1.6 (0.86-2.7)	0.92% (-2.6-5.7%)	0.76% (-3.7-6.8%)	21 (-100-190)	160 (-460-1000)
Lithuania	2016	Female	95 plus	51% (30-71%)	49% (29-70%)	1.8 (0.86-3.3)	1.2% (-3.4-8.2%)	1% (-4.4-9.2%)	5 (-22-45)	23 (-65-160)
Lithuania	2016	Male	All Ages	82% (77-86%)	18% (14-23%)	7 (5.6-8.6)	19% (16-22%)	18% (14-22%)	3500 (2700-4400)	110000 (95000-130000)
Lithuania	2016	Male	15 to 19	68% (52-82%)	32% (18-48%)	3.5 (2.2-5.2)	10% (6.7-15%)	18% (11-26%)	12 (7.3-18)	1000 (670-1500)
Lithuania	2016	Male	20 to 24	82% (69-90%)	18% (9.8-31%)	6.5 (4.4-9)	21% (16-27%)	32% (23-41%)	52 (36-68)	4000 (3000-5100)
Lithuania	2016	Male	25 to 29	87% (78-93%)	13% (6.9-22%)	7.2 (4.8-10)	24% (18-29%)	33% (24-41%)	68 (49-86)	4900 (3800-6100)
Lithuania	2016	Male	30 to 34	90% (81-95%)	9.8% (4.5-19%)	8.3 (5.9-11)	26% (22-31%)	36% (29-42%)	92 (73-110)	5900 (4900-7000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lithuania	2016	Male	35 to 39	91% (84-96%)	8.7% (4-16%)	10 (6.8-14)	29% (25-33%)	38% (33-43%)	140 (110-160)	7700 (6400-8900)
Lithuania	2016	Male	40 to 44	92% (85-96%)	8.4% (4-15%)	12 (8.3-16)	29% (26-34%)	39% (34-43%)	220 (180-250)	11000 (9200-13000)
Lithuania	2016	Male	45 to 49	91% (81-96%)	8.9% (4.1-19%)	13 (9.4-18)	28% (24-33%)	36% (31-41%)	290 (240-340)	13000 (11000-15000)
Lithuania	2016	Male	50 to 54	90% (81-95%)	10% (4.7-19%)	12 (8.3-16)	26% (22-31%)	32% (28-37%)	440 (360-530)	17000 (14000-20000)
Lithuania	2016	Male	55 to 59	88% (78-94%)	12% (5.8-22%)	8.5 (6.1-11)	22% (18-26%)	26% (22-31%)	450 (360-550)	15000 (12000-18000)
Lithuania	2016	Male	60 to 64	86% (75-94%)	14% (6.5-25%)	6.6 (4.3-9.4)	17% (13-21%)	20% (15-25%)	400 (290-520)	12000 (8500-15000)
Lithuania	2016	Male	65 to 69	84% (72-93%)	16% (7.2-28%)	6.1 (4.1-8.5)	14% (10-19%)	17% (12-22%)	310 (220-410)	7500 (5300-9700)
Lithuania	2016	Male	70 to 74	82% (68-90%)	18% (10-32%)	5.4 (3.6-7.8)	11% (6.9-16%)	13% (7.8-19%)	300 (180-420)	5900 (3600-8300)
Lithuania	2016	Male	75 to 79	78% (63-89%)	22% (11-37%)	4.8 (2.9-7.1)	9.2% (4.5-14%)	11% (4.8-17%)	270 (120-420)	4200 (2100-6300)
Lithuania	2016	Male	80 to 84	74% (58-85%)	26% (15-42%)	4.3 (2.7-6.4)	7.9% (3.2-13%)	8.9% (3.2-15%)	240 (89-400)	2800 (1100-4500)
Lithuania	2016	Male	85 to 89	70% (54-83%)	30% (17-46%)	3.9 (2.4-5.9)	6.6% (1.6-12%)	7.3% (1.1-14%)	130 (20-250)	1100 (260-2000)
Lithuania	2016	Male	90 to 94	68% (54-82%)	32% (18-46%)	3.6 (2.1-5.5)	5.1% (-0.1-11%)	5.5% (-0.88-13%)	45 (-7.3-100)	260 (-5.3-580)
Lithuania	2016	Male	95 plus	67% (48-84%)	33% (16-52%)	3.3 (1.6-5.7)	3.8% (-2.2-11%)	3.8% (-3.5-12%)	6 (-5.5-19)	24 (-14-70)
Moldova	1990	Female	All Ages	66% (59-73%)	34% (27-41%)	2.6 (1.6-3.7)	8.7% (5.5-12%)	10% (5.1-17%)	2300 (1100-3700)	71000 (45000-100000)
Moldova	1990	Female	15 to 19	59% (43-74%)	41% (26-57%)	1.7 (0.93-2.7)	4.5% (3-6.7%)	7.5% (4-12%)	8.1 (4.2-13)	950 (610-1400)
Moldova	1990	Female	20 to 24	73% (56-85%)	27% (15-44%)	2.7 (1.6-4.4)	7.6% (5.4-10%)	10% (6-15%)	11 (6.6-18)	1500 (1100-2100)
Moldova	1990	Female	25 to 29	78% (63-88%)	22% (12-37%)	3.5 (2.1-5.3)	8.7% (6.6-11%)	13% (8.5-18%)	19 (12-28)	2300 (1700-3100)
Moldova	1990	Female	30 to 34	81% (67-90%)	19% (9.7-33%)	3.9 (2.3-6)	9.6% (7.3-12%)	16% (11-21%)	36 (25-50)	3300 (2400-4300)
Moldova	1990	Female	35 to 39	83% (71-91%)	17% (9.1-29%)	4 (2.4-6.2)	11% (8.6-14%)	18% (13-23%)	64 (46-87)	4500 (3300-5700)
Moldova	1990	Female	40 to 44	83% (71-91%)	17% (8.8-29%)	4.2 (2.5-6.4)	13% (9.7-17%)	20% (15-26%)	84 (61-110)	4800 (3600-6100)
Moldova	1990	Female	45 to 49	81% (65-91%)	19% (9.1-35%)	4.4 (2.6-6.9)	15% (11-19%)	21% (16-27%)	130 (91-170)	6000 (4400-7700)
Moldova	1990	Female	50 to 54	78% (63-88%)	22% (12-37%)	4.4 (2.4-6.9)	15% (11-20%)	21% (15-27%)	230 (160-300)	9300 (6700-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	1990	Female	55 to 59	72% (55-86%)	28% (14-45%)	4.2 (2.2-7.1)	15% (10-20%)	20% (13-27%)	300 (200-410)	10000 (6700-14000)
Moldova	1990	Female	60 to 64	67% (48-83%)	33% (17-52%)	3.8 (2-6.3)	13% (7.2-19%)	17% (9.2-25%)	360 (200-550)	10000 (5700-15000)
Moldova	1990	Female	65 to 69	62% (42-79%)	38% (21-58%)	3.2 (1.7-5.3)	11% (4.8-18%)	13% (5.8-22%)	370 (160-620)	8600 (3700-14000)
Moldova	1990	Female	70 to 74	54% (36-71%)	46% (29-64%)	2.5 (1.2-4.4)	7.6% (1.8-15%)	9.1% (1.9-18%)	220 (48-440)	4200 (950-8200)
Moldova	1990	Female	75 to 79	44% (28-63%)	56% (37-72%)	1.9 (0.92-3.5)	5.1% (0.044-12%)	5.9% (-0.088-14%)	200 (-2.9-510)	3000 (26-7300)
Moldova	1990	Female	80 to 84	38% (22-56%)	62% (44-78%)	1.7 (0.78-3)	4% (-0.27-10%)	4.5% (-0.56-12%)	130 (-16-360)	1400 (-100-3800)
Moldova	1990	Female	85 to 89	35% (20-52%)	65% (48-80%)	1.6 (0.72-3)	3.5% (-0.69-10%)	3.9% (-0.98-12%)	73 (-18-220)	570 (-110-1600)
Moldova	1990	Female	90 to 94	37% (22-53%)	63% (47-78%)	1.7 (0.77-3.2)	3.4% (-1.2-10%)	3.8% (-1.6-12%)	30 (-13-95)	160 (-54-490)
Moldova	1990	Female	95 plus	40% (22-60%)	60% (40-78%)	1.9 (0.72-4)	3.6% (-1.6-13%)	3.9% (-2.2-14%)	8.2 (-4.5-30)	29 (-13-100)
Moldova	1990	Male	All Ages	78% (71-84%)	22% (16-29%)	5.6 (3.6-8.1)	16% (12-19%)	17% (12-22%)	4000 (2700-5200)	150000 (110000-180000)
Moldova	1990	Male	15 to 19	69% (54-83%)	31% (17-46%)	4 (2.3-6.3)	11% (6.7-16%)	17% (9.7-25%)	37 (21-55)	3000 (1800-4300)
Moldova	1990	Male	20 to 24	82% (69-91%)	18% (9.4-31%)	7.3 (4.4-11)	22% (16-29%)	29% (20-38%)	110 (73-150)	7800 (5500-10000)
Moldova	1990	Male	25 to 29	87% (79-93%)	13% (6.8-21%)	8.4 (5.1-13)	24% (18-30%)	31% (23-39%)	150 (110-190)	10000 (7700-13000)
Moldova	1990	Male	30 to 34	90% (82-96%)	9.6% (4.4-18%)	9.3 (5.6-14)	24% (19-29%)	32% (25-38%)	210 (160-260)	13000 (10000-16000)
Moldova	1990	Male	35 to 39	91% (85-96%)	8.5% (4-15%)	9.9 (6.2-15)	25% (21-29%)	33% (27-38%)	270 (210-320)	15000 (12000-18000)
Moldova	1990	Male	40 to 44	92% (85-96%)	8.3% (3.9-15%)	10 (6.4-15)	26% (22-30%)	33% (27-38%)	270 (220-330)	13000 (11000-16000)
Moldova	1990	Male	45 to 49	91% (82-96%)	8.9% (4.1-18%)	10 (6-15)	25% (21-29%)	31% (26-35%)	310 (250-370)	13000 (11000-16000)
Moldova	1990	Male	50 to 54	90% (81-95%)	10% (4.9-19%)	9.4 (5.7-14)	23% (19-27%)	27% (23-31%)	470 (380-570)	18000 (14000-21000)
Moldova	1990	Male	55 to 59	88% (78-94%)	12% (5.9-22%)	8.5 (5.1-13)	21% (16-24%)	24% (19-29%)	490 (380-590)	16000 (12000-20000)
Moldova	1990	Male	60 to 64	86% (75-93%)	14% (6.6-25%)	7.4 (4.4-11)	18% (13-23%)	21% (15-26%)	570 (400-730)	16000 (11000-20000)
Moldova	1990	Male	65 to 69	85% (73-93%)	15% (7-27%)	6.1 (3.6-9.2)	15% (9.3-21%)	17% (10-24%)	470 (280-640)	11000 (6500-15000)
Moldova	1990	Male	70 to 74	82% (69-90%)	18% (9.9-31%)	4.7 (2.6-7.6)	11% (3.6-17%)	12% (3.8-20%)	260 (81-430)	4900 (1600-8000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	1990	Male	75 to 79	78% (64-89%)	22% (11-36%)	3.8 (2-6.5)	7.2% (0.26-15%)	7.9% (-0.12-17%)	200 (-3.2-430)	3000 (110-6200)
Moldova	1990	Male	80 to 84	74% (58-86%)	26% (14-42%)	3.5 (1.9-5.8)	5.5% (-1.2-13%)	5.9% (-1.9-15%)	100 (-33-270)	1200 (-270-2900)
Moldova	1990	Male	85 to 89	71% (55-83%)	29% (17-45%)	3.2 (1.7-5.4)	4.2% (-2.3-12%)	4.3% (-3.1-14%)	39 (-29-130)	330 (-180-960)
Moldova	1990	Male	90 to 94	69% (54-82%)	31% (18-46%)	3.1 (1.6-5.4)	3% (-3-11%)	2.9% (-4-13%)	9.6 (-13-40)	58 (-58-210)
Moldova	1990	Male	95 plus	68% (49-83%)	32% (17-51%)	2.8 (1.2-5.5)	2.2% (-3.5-10%)	2% (-4.8-11%)	1.4 (-3.3-8.1)	6.2 (-9.7-28)
Moldova	1995	Female	All Ages	67% (60-74%)	33% (26-40%)	2.6 (1.7-3.6)	9.6% (6.1-14%)	11% (5.2-17%)	2900 (1400-4700)	86000 (53000-120000)
Moldova	1995	Female	15 to 19	60% (44-76%)	40% (24-56%)	1.7 (1-2.8)	4.4% (2.9-6.2%)	7.7% (4.1-12%)	7.8 (4.2-12)	950 (620-1400)
Moldova	1995	Female	20 to 24	73% (58-85%)	27% (15-42%)	2.8 (1.7-4.2)	7.8% (5.7-10%)	11% (6.4-15%)	13 (7.7-19)	1700 (1200-2300)
Moldova	1995	Female	25 to 29	78% (64-88%)	22% (12-36%)	3.5 (2.1-5.2)	9.4% (7-12%)	14% (9.1-19%)	20 (13-28)	2100 (1500-2800)
Moldova	1995	Female	30 to 34	81% (67-90%)	19% (9.6-33%)	3.9 (2.4-5.8)	11% (8.1-13%)	17% (12-22%)	43 (30-57)	3500 (2600-4500)
Moldova	1995	Female	35 to 39	83% (71-91%)	17% (8.8-29%)	4 (2.5-5.9)	12% (9.1-15%)	19% (14-24%)	82 (60-110)	5300 (4100-6700)
Moldova	1995	Female	40 to 44	83% (71-92%)	17% (8.3-29%)	4.2 (2.6-6.2)	14% (11-18%)	21% (16-27%)	140 (100-180)	7500 (5800-9500)
Moldova	1995	Female	45 to 49	82% (66-91%)	18% (9-34%)	4.4 (2.6-6.8)	15% (11-20%)	17% (16-28%)	170 (120-220)	7600 (5700-9900)
Moldova	1995	Female	50 to 54	78% (64-89%)	22% (11-36%)	4.4 (2.5-6.7)	17% (12-22%)	22% (16-28%)	230 (160-300)	8800 (6200-12000)
Moldova	1995	Female	55 to 59	73% (57-87%)	27% (13-43%)	4.2 (2.3-7)	16% (10-22%)	20% (13-28%)	390 (250-530)	13000 (8300-18000)
Moldova	1995	Female	60 to 64	68% (50-84%)	32% (16-50%)	3.9 (2-6.4)	14% (7.7-21%)	17% (9.2-25%)	410 (220-620)	11000 (6300-17000)
Moldova	1995	Female	65 to 69	63% (43-81%)	37% (19-57%)	3.3 (1.7-5.3)	12% (5-19%)	14% (5.9-23%)	480 (210-780)	11000 (4900-18000)
Moldova	1995	Female	70 to 74	55% (37-72%)	45% (28-63%)	2.6 (1.3-4.4)	8.2% (1.9-16%)	9.5% (2-19%)	400 (85-780)	7500 (1700-14000)
Moldova	1995	Female	75 to 79	45% (28-65%)	55% (35-72%)	2 (0.95-3.5)	5.3% (-0.11-13%)	6% (-0.27-15%)	200 (-8.6-500)	2900 (-58-7000)
Moldova	1995	Female	80 to 84	39% (23-57%)	61% (43-77%)	1.7 (0.84-2.9)	4% (-0.62-11%)	4.4% (-0.91-12%)	140 (-29-390)	1500 (-230-4100)
Moldova	1995	Female	85 to 89	36% (21-54%)	64% (46-79%)	1.6 (0.75-2.9)	3.4% (-0.87-10%)	3.8% (-1.3-12%)	96 (-33-290)	740 (-190-2200)
Moldova	1995	Female	90 to 94	38% (23-55%)	62% (45-77%)	1.7 (0.81-3.2)	3.4% (-1.1-11%)	3.7% (-1.6-12%)	38 (-16-120)	200 (-68-620)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	1995	Female	95 plus	41% (22-61%)	59% (39-78%)	1.9 (0.75-3.9)	3.7% (-1.7-13%)	4% (-2.2-14%)	12 (-6.5-43)	42 (-20-140)
Moldova	1995	Male	All Ages	79% (72-84%)	21% (16-28%)	5.7 (3.8-8.1)	18% (14-21%)	19% (13-24%)	5200 (3700-6800)	190000 (150000-230000)
Moldova	1995	Male	15 to 19	70% (54-82%)	30% (18-46%)	3.9 (2.3-6)	11% (7.1-16%)	17% (10-24%)	43 (25-62)	3400 (2100-4800)
Moldova	1995	Male	20 to 24	82% (70-90%)	18% (9.5-30%)	7.3 (4.6-11)	22% (17-29%)	30% (22-38%)	110 (79-150)	8300 (5900-11000)
Moldova	1995	Male	25 to 29	87% (79-93%)	13% (6.7-21%)	8.4 (5.3-12)	25% (20-31%)	33% (25-40%)	150 (110-190)	9900 (7700-12000)
Moldova	1995	Male	30 to 34	90% (82-96%)	9.6% (4.5-18%)	9.2 (5.7-13)	26% (21-31%)	33% (26-38%)	230 (180-280)	14000 (11000-17000)
Moldova	1995	Male	35 to 39	91% (85-96%)	8.6% (4.2-15%)	9.8 (6.2-14)	27% (23-32%)	34% (28-39%)	370 (300-440)	20000 (17000-23000)
Moldova	1995	Male	40 to 44	92% (85-96%)	8.4% (4.1-15%)	10 (6.4-14)	28% (24-32%)	34% (29-38%)	530 (430-620)	25000 (21000-30000)
Moldova	1995	Male	45 to 49	91% (82-96%)	8.9% (4.1-18%)	9.9 (6.3-14)	27% (23-31%)	32% (27-37%)	500 (410-600)	21000 (18000-25000)
Moldova	1995	Male	50 to 54	90% (81-95%)	10% (4.6-19%)	9.4 (5.9-14)	25% (21-29%)	29% (24-33%)	510 (420-620)	19000 (16000-23000)
Moldova	1995	Male	55 to 59	88% (79-94%)	12% (5.6-21%)	8.5 (5.2-12)	23% (18-27%)	26% (20-31%)	700 (540-850)	23000 (17000-27000)
Moldova	1995	Male	60 to 64	87% (76-94%)	13% (6.4-24%)	7.4 (4.6-11)	19% (13-25%)	22% (15-27%)	680 (470-870)	19000 (13000-24000)
Moldova	1995	Male	65 to 69	85% (73-93%)	15% (7-27%)	6 (3.7-8.9)	16% (9.7-22%)	18% (11-24%)	670 (390-930)	15000 (9200-21000)
Moldova	1995	Male	70 to 74	82% (70-90%)	18% (9.6-30%)	4.7 (2.7-7.4)	11% (3.5-18%)	12% (3.7-21%)	420 (120-700)	7700 (2400-13000)
Moldova	1995	Male	75 to 79	79% (65-89%)	21% (11-35%)	3.8 (2.1-6.3)	7.2% (-0.26-15%)	7.8% (-0.71-17%)	180 (-16-390)	2700 (-94-5600)
Moldova	1995	Male	80 to 84	75% (60-86%)	25% (14-40%)	3.5 (1.9-5.5)	5.2% (-2-13%)	5.4% (-2.6-15%)	99 (-49-280)	1100 (-430-2900)
Moldova	1995	Male	85 to 89	71% (55-82%)	29% (18-45%)	3.2 (1.7-5.1)	3.6% (-3.1-11%)	3.7% (-3.9-12%)	45 (-49-150)	380 (-320-1200)
Moldova	1995	Male	90 to 94	70% (55-82%)	30% (18-45%)	3 (1.6-5.1)	2.5% (-3.6-11%)	2.4% (-4.7-12%)	10 (-20-49)	62 (-86-260)
Moldova	1995	Male	95 plus	68% (50-84%)	32% (16-50%)	2.8 (1.2-5.3)	2% (-4-10%)	1.7% (-5.2-11%)	1.7 (-5.3-11)	7.6 (-15-40)
Moldova	2000	Female	All Ages	66% (59-72%)	34% (28-41%)	2.2 (1.5-3.1)	8% (5.1-12%)	8.4% (3.9-14%)	1900 (900-3200)	60000 (38000-87000)
Moldova	2000	Female	15 to 19	57% (42-71%)	43% (29-58%)	1.4 (0.83-2.1)	3.8% (2.6-5.1%)	6.4% (3.6-9.8%)	5.5 (3.2-8.6)	810 (550-1100)
Moldova	2000	Female	20 to 24	70% (55-82%)	30% (18-45%)	2.2 (1.4-3.2)	7% (5.1-9.2%)	9.6% (5.8-14%)	8.8 (5.2-13)	1500 (1000-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	2000	Female	25 to 29	76% (65-86%)	24% (14-35%)	2.9 (1.8-4.1)	8.3% (6.3-10%)	12% (8.1-17%)	15 (9.6-21)	1800 (1300-2400)
Moldova	2000	Female	30 to 34	80% (68-89%)	20% (11-32%)	3.2 (2-4.6)	8.9% (6.9-11%)	14% (10-19%)	22 (15-30)	2000 (1500-2600)
Moldova	2000	Female	35 to 39	82% (71-90%)	18% (9.9-29%)	3.3 (2.1-4.8)	10% (7.9-13%)	17% (13-22%)	49 (36-64)	3500 (2700-4400)
Moldova	2000	Female	40 to 44	82% (71-90%)	18% (9.6-29%)	3.5 (2.2-4.9)	12% (8.9-15%)	19% (14-24%)	92 (69-120)	5300 (4000-6700)
Moldova	2000	Female	45 to 49	81% (69-90%)	19% (9.9-31%)	3.7 (2.3-5.4)	13% (9.5-17%)	20% (15-25%)	150 (110-200)	7100 (5300-9200)
Moldova	2000	Female	50 to 54	78% (63-88%)	22% (12-37%)	3.7 (2.2-5.6)	14% (10-19%)	20% (14-26%)	160 (120-220)	6600 (4700-8800)
Moldova	2000	Female	55 to 59	72% (55-86%)	28% (14-45%)	3.5 (1.9-5.7)	14% (8.6-20%)	18% (11-26%)	210 (130-310)	7200 (4500-10000)
Moldova	2000	Female	60 to 64	67% (49-83%)	33% (17-51%)	3.2 (1.7-5.3)	12% (6.4-18%)	15% (8.1-23%)	300 (160-470)	8500 (4500-13000)
Moldova	2000	Female	65 to 69	62% (43-80%)	38% (20-57%)	2.7 (1.5-4.4)	9.6% (4.4-16%)	12% (5.3-20%)	300 (130-520)	7100 (3200-12000)
Moldova	2000	Female	70 to 74	54% (36-72%)	46% (28-64%)	2.1 (1.1-3.6)	6.5% (1.4-13%)	7.7% (1.6-16%)	270 (55-560)	5100 (1100-10000)
Moldova	2000	Female	75 to 79	44% (28-64%)	56% (36-72%)	1.6 (0.83-2.8)	3.9% (-0.39-9.8%)	4.4% (-0.59-12%)	170 (-24-450)	2600 (-260-6400)
Moldova	2000	Female	80 to 84	37% (21-56%)	63% (44-79%)	1.4 (0.71-2.4)	2.5% (-0.97-7.4%)	2.7% (-1.2-8.5%)	75 (-33-240)	830 (-310-2500)
Moldova	2000	Female	85 to 89	35% (21-52%)	65% (48-79%)	1.3 (0.64-2.4)	1.8% (-1.4-6.6%)	2% (-1.8-7.7%)	35 (-33-140)	280 (-210-1000)
Moldova	2000	Female	90 to 94	36% (21-54%)	64% (46-79%)	1.4 (0.67-2.5)	1.8% (-1.6-7.4%)	1.9% (-2.3-8.6%)	19 (-23-87)	110 (-99-450)
Moldova	2000	Female	95 plus	40% (21-62%)	60% (38-79%)	1.5 (0.62-3.2)	2.1% (-2.3-9.2%)	2.2% (-2.9-11%)	6.2 (-8.3-30)	23 (-25-100)
Moldova	2000	Male	All Ages	81% (75-86%)	19% (14-25%)	5.2 (3.5-7.1)	17% (12-20%)	17% (11-22%)	4000 (2600-5300)	150000 (110000-180000)
Moldova	2000	Male	15 to 19	72% (58-83%)	28% (17-42%)	3.5 (2.2-5.2)	9.5% (6.2-13%)	16% (9.6-22%)	31 (19-44)	2600 (1700-3600)
Moldova	2000	Male	20 to 24	84% (72-91%)	16% (8.7-28%)	6.4 (4.2-9.2)	20% (14-25%)	29% (20-37%)	84 (58-110)	6400 (4600-8400)
Moldova	2000	Male	25 to 29	89% (82-94%)	11% (6-18%)	7.4 (4.8-11)	23% (17-28%)	31% (23-38%)	110 (77-130)	7400 (5400-9200)
Moldova	2000	Male	30 to 34	92% (86-96%)	8% (3.9-14%)	8.2 (5.3-12)	23% (18-28%)	31% (24-37%)	120 (95-150)	7800 (6100-9500)
Moldova	2000	Male	35 to 39	93% (88-96%)	7.2% (3.6-12%)	8.6 (5.7-12)	26% (21-30%)	33% (27-38%)	240 (190-290)	13000 (11000-16000)
Moldova	2000	Male	40 to 44	93% (88-97%)	6.9% (3.5-12%)	8.9 (5.9-13)	27% (22-31%)	34% (28-39%)	400 (320-480)	20000 (16000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	2000	Male	45 to 49	93% (86-96%)	7.3% (3.5-14%)	8.8 (5.8-13)	27% (22-31%)	32% (27-37%)	530 (430-650)	23000 (18000-28000)
Moldova	2000	Male	50 to 54	91% (83-95%)	9.3% (4.5-17%)	8.1 (5.4-11)	25% (20-29%)	29% (24-34%)	450 (350-540)	17000 (13000-20000)
Moldova	2000	Male	55 to 59	88% (79-94%)	12% (5.6-21%)	7.2 (4.6-10)	22% (16-26%)	26% (19-31%)	450 (330-550)	15000 (11000-18000)
Moldova	2000	Male	60 to 64	87% (76-94%)	13% (6.4-24%)	6.3 (3.9-9.4)	18% (12-23%)	21% (13-27%)	540 (340-710)	15000 (9500-20000)
Moldova	2000	Male	65 to 69	85% (72-93%)	15% (7.4-28%)	5.1 (3.2-7.5)	14% (7.5-20%)	16% (8.3-23%)	480 (240-690)	11000 (5700-16000)
Moldova	2000	Male	70 to 74	82% (69-90%)	18% (10-31%)	3.9 (2.3-6.2)	9.1% (2.3-16%)	10% (2.4-19%)	320 (76-600)	6100 (1600-11000)
Moldova	2000	Male	75 to 79	78% (64-89%)	22% (11-36%)	3.2 (1.8-5.1)	5.1% (-1.3-13%)	5.5% (-1.8-14%)	150 (-51-400)	2300 (-590-5800)
Moldova	2000	Male	80 to 84	74% (58-86%)	26% (14-42%)	2.9 (1.6-4.5)	2.9% (-2.9-10%)	2.9% (-3.8-11%)	47 (-62-180)	560 (-570-1900)
Moldova	2000	Male	85 to 89	71% (54-83%)	29% (17-46%)	2.7 (1.5-4.2)	1.4% (-3.9-8.4%)	1.2% (-5.1-9.2%)	11 (-45-83)	110 (-290-640)
Moldova	2000	Male	90 to 94	69% (54-82%)	31% (18-46%)	2.5 (1.3-4.3)	0.71% (-4.3-8.2%)	0.37% (-5.4-9.1%)	1.6 (-24-41)	19 (-110-220)
Moldova	2000	Male	95 plus	68% (49-83%)	32% (17-51%)	2.4 (1.1-4.5)	0.49% (-4.6-8.2%)	0.052% (-5.8-9.1%)	0.056 (-6-9.2)	1.9 (-18-33)
Moldova	2005	Female	All Ages	67% (61-74%)	33% (26-39%)	2 (1.4-2.7)	7.6% (4.9-11%)	7.2% (3.3-12%)	1700 (740-2800)	56000 (36000-78000)
Moldova	2005	Female	15 to 19	57% (43-71%)	43% (29-57%)	1.2 (0.75-1.8)	3.5% (2.4-4.8%)	5.9% (3.3-9.1%)	4.1 (2.3-6.3)	720 (490-1000)
Moldova	2005	Female	20 to 24	70% (56-82%)	30% (18-44%)	1.9 (1.2-2.8)	6.7% (4.9-8.6%)	8.4% (5.1-12%)	7.4 (4.5-11)	1500 (1000-2100)
Moldova	2005	Female	25 to 29	77% (65-87%)	23% (13-35%)	2.5 (1.6-3.5)	7.5% (5.8-9.3%)	11% (7.5-15%)	12 (8-17)	1800 (1300-2300)
Moldova	2005	Female	30 to 34	81% (69-90%)	19% (10-31%)	2.8 (1.8-4.1)	8.2% (6.5-10%)	13% (9.7-17%)	21 (15-28)	2100 (1600-2600)
Moldova	2005	Female	35 to 39	82% (71-90%)	18% (9.6-29%)	2.9 (1.9-4.2)	9.8% (7.8-12%)	16% (12-20%)	37 (28-48)	2600 (2000-3300)
Moldova	2005	Female	40 to 44	83% (72-91%)	17% (9.3-28%)	3 (2-4.2)	11% (8.8-14%)	18% (14-22%)	82 (63-100)	4700 (3600-5800)
Moldova	2005	Female	45 to 49	82% (68-90%)	18% (9.8-32%)	3.2 (2.1-4.6)	12% (9.4-16%)	19% (14-24%)	150 (110-190)	7000 (5300-8800)
Moldova	2005	Female	50 to 54	78% (65-88%)	22% (12-35%)	3.2 (2-4.7)	13% (9.6-17%)	19% (14-24%)	220 (160-290)	8700 (6300-11000)
Moldova	2005	Female	55 to 59	73% (56-86%)	27% (14-44%)	3.1 (1.7-5)	13% (8.3-18%)	17% (11-24%)	210 (130-310)	7100 (4500-10000)
Moldova	2005	Female	60 to 64	68% (49-83%)	32% (17-51%)	2.8 (1.5-4.5)	11% (6-17%)	14% (7.7-21%)	240 (130-370)	6800 (3600-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	2005	Female	65 to 69	63% (43-80%)	37% (20-57%)	2.4 (1.3-3.8)	8.2% (3.6-14%)	10% (4.3-17%)	260 (110-450)	6100 (2700-10000)
Moldova	2005	Female	70 to 74	55% (37-72%)	45% (28-63%)	1.8 (1-3.1)	5.2% (0.93-11%)	6.1% (0.98-13%)	200 (31-420)	3800 (680-7900)
Moldova	2005	Female	75 to 79	45% (28-64%)	55% (36-72%)	1.4 (0.74-2.5)	2.8% (-0.72-7.5%)	3.2% (-1-8.9%)	130 (-40-360)	2000 (-490-5300)
Moldova	2005	Female	80 to 84	38% (22-56%)	62% (44-78%)	1.2 (0.63-2.1)	1.6% (-1.3-5.9%)	1.7% (-1.7-6.7%)	61 (-60-240)	690 (-560-2600)
Moldova	2005	Female	85 to 89	36% (22-52%)	64% (48-78%)	1.2 (0.56-2.1)	1% (-1.7-5.1%)	1% (-2.3-6%)	20 (-44-110)	170 (-290-860)
Moldova	2005	Female	90 to 94	37% (23-55%)	63% (45-77%)	1.2 (0.62-2.3)	0.93% (-2-5.6%)	0.9% (-2.7-6.4%)	7.5 (-22-55)	47 (-110-280)
Moldova	2005	Female	95 plus	41% (23-62%)	59% (38-77%)	1.4 (0.56-2.8)	1.1% (-2.6-7.2%)	1.1% (-3.4-8.5%)	3.6 (-12-28)	15 (-34-97)
Moldova	2005	Male	All Ages	82% (76-87%)	18% (13-24%)	4.8 (3.4-6.5)	17% (13-21%)	16% (10-21%)	4000 (2600-5500)	150000 (110000-190000)
Moldova	2005	Male	15 to 19	72% (58-84%)	28% (16-42%)	3.1 (2-4.6)	8.3% (5.4-12%)	14% (8.8-20%)	24 (15-35)	2100 (1400-2900)
Moldova	2005	Male	20 to 24	84% (73-91%)	16% (8.7-27%)	5.7 (3.8-8)	18% (13-23%)	27% (19-35%)	70 (47-95)	5500 (4000-7200)
Moldova	2005	Male	25 to 29	89% (82-94%)	11% (6.1-18%)	6.5 (4.3-9.2)	20% (15-25%)	29% (21-36%)	98 (70-120)	7000 (5300-8800)
Moldova	2005	Male	30 to 34	92% (85-96%)	8.3% (4-15%)	7.2 (4.7-10)	22% (17-27%)	29% (23-36%)	130 (100-170)	8500 (6500-10000)
Moldova	2005	Male	35 to 39	93% (87-96%)	7.4% (3.7-13%)	7.6 (5.1-11)	25% (20-29%)	31% (25-37%)	200 (150-240)	11000 (8600-13000)
Moldova	2005	Male	40 to 44	93% (87-97%)	7.1% (3.4-13%)	7.8 (5.3-11)	27% (22-32%)	33% (26-38%)	400 (310-480)	19000 (15000-23000)
Moldova	2005	Male	45 to 49	92% (85-96%)	7.5% (3.6-15%)	7.7 (5.2-11)	26% (21-31%)	31% (25-37%)	610 (480-740)	26000 (20000-31000)
Moldova	2005	Male	50 to 54	91% (83-96%)	9.1% (4.3-17%)	7.2 (4.8-10)	24% (19-29%)	29% (22-34%)	710 (540-880)	27000 (21000-33000)
Moldova	2005	Male	55 to 59	89% (80-95%)	11% (5.4-20%)	6.4 (4.1-9)	21% (15-26%)	25% (18-30%)	500 (350-630)	16000 (11000-20000)
Moldova	2005	Male	60 to 64	87% (76-94%)	13% (6.2-24%)	5.5 (3.4-8.1)	17% (10-23%)	19% (12-26%)	430 (260-590)	12000 (7300-16000)
Moldova	2005	Male	65 to 69	85% (73-93%)	15% (6.8-27%)	4.5 (2.9-6.5)	13% (6.3-19%)	14% (6.9-21%)	440 (210-680)	10000 (4900-15000)
Moldova	2005	Male	70 to 74	82% (70-90%)	18% (9.6-30%)	3.4 (2-5.2)	7.4% (1.3-15%)	8.2% (1.2-16%)	260 (39-520)	4900 (800-9600)
Moldova	2005	Male	75 to 79	79% (65-89%)	21% (11-35%)	2.8 (1.6-4.5)	3.6% (-1.7-10%)	3.8% (-2.2-12%)	120 (-70-360)	1800 (-820-5200)
Moldova	2005	Male	80 to 84	75% (60-86%)	25% (14-40%)	2.5 (1.5-3.9)	1.4% (-3.6-7.6%)	1.3% (-4.6-8.2%)	26 (-97-170)	360 (-910-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	2005	Male	85 to 89	71% (55-83%)	29% (17-45%)	2.4 (1.3-3.7)	0.12% (-4.5-6.3%)	-0.31% (-5.7-6.7%)	-2.9 (-53-63)	9.1 (-360-500)
Moldova	2005	Male	90 to 94	69% (55-82%)	31% (18-45%)	2.2 (1.2-3.8)	-0.46% (-4.5-5.9%)	-0.99% (-5.9-6.3%)	-3.5 (-21-23)	-9.6 (-96-120)
Moldova	2005	Male	95 plus	68% (49-83%)	32% (17-51%)	2.1 (0.95-3.9)	-0.55% (-4.9-6.1%)	-1.1% (-6.4-6.7%)	-1.3 (-7.4-7.6)	-2.5 (-22-28)
Moldova	2010	Female	All Ages	69% (62-75%)	31% (25-38%)	2 (1.3-2.7)	7% (4.7-9.8%)	6.3% (2.7-11%)	1400 (610-2400)	49000 (33000-69000)
Moldova	2010	Female	15 to 19	58% (44-72%)	42% (28-56%)	1.1 (0.67-1.7)	3.4% (2.4-4.6%)	5.6% (3.2-8.7%)	3.1 (1.7-4.7)	560 (370-790)
Moldova	2010	Female	20 to 24	71% (57-82%)	29% (18-43%)	1.8 (1.1-2.6)	6.5% (4.7-8.5%)	7.7% (4.7-11%)	5.8 (3.5-8.6)	1500 (990-2100)
Moldova	2010	Female	25 to 29	78% (66-87%)	22% (13-34%)	2.3 (1.5-3.3)	7.3% (5.7-9%)	10% (7.1-14%)	10 (7.1-15)	1700 (1200-2300)
Moldova	2010	Female	30 to 34	81% (70-90%)	19% (10-30%)	2.6 (1.6-3.9)	7.8% (6.2-9.7%)	12% (9.5-16%)	18 (13-24)	2000 (1500-2500)
Moldova	2010	Female	35 to 39	83% (71-91%)	17% (9.4-29%)	2.7 (1.7-4)	9.5% (7.6-12%)	16% (13-20%)	35 (27-45)	2600 (2100-3300)
Moldova	2010	Female	40 to 44	83% (72-91%)	17% (9.3-28%)	2.8 (1.8-4.1)	11% (8.4-13%)	18% (14-22%)	58 (44-73)	3400 (2700-4300)
Moldova	2010	Female	45 to 49	82% (70-90%)	18% (9.8-30%)	3 (1.9-4.5)	12% (8.9-15%)	18% (15-23%)	120 (90-150)	5700 (4400-7200)
Moldova	2010	Female	50 to 54	79% (66-89%)	21% (11-34%)	3 (1.8-4.5)	12% (9.1-16%)	18% (14-23%)	180 (140-240)	7400 (5500-9700)
Moldova	2010	Female	55 to 59	73% (57-87%)	27% (13-43%)	2.9 (1.6-4.6)	16% (7.8-16%)	16% (11-23%)	250 (160-350)	8400 (5600-12000)
Moldova	2010	Female	60 to 64	68% (50-83%)	32% (17-50%)	2.6 (1.4-4.2)	10% (5.9-16%)	13% (7.6-20%)	210 (120-310)	5900 (3300-9000)
Moldova	2010	Female	65 to 69	63% (44-80%)	37% (20-56%)	2.2 (1.3-3.5)	7.3% (3-12%)	9.1% (3.8-16%)	180 (77-320)	4400 (1800-7500)
Moldova	2010	Female	70 to 74	56% (38-73%)	44% (27-62%)	1.8 (0.95-2.9)	4.6% (0.41-9.8%)	5.5% (0.43-12%)	170 (14-370)	3200 (290-7000)
Moldova	2010	Female	75 to 79	46% (29-64%)	54% (36-71%)	1.4 (0.7-2.4)	2.4% (-0.92-7.2%)	2.8% (-1.2-8.6%)	100 (-45-310)	1500 (-590-4500)
Moldova	2010	Female	80 to 84	39% (22-57%)	61% (43-78%)	1.2 (0.59-2)	1.3% (-1.4-5.2%)	1.3% (-1.9-6.1%)	50 (-69-220)	580 (-640-2400)
Moldova	2010	Female	85 to 89	36% (22-53%)	64% (47-78%)	1.1 (0.51-2)	0.74% (-1.8-4.4%)	0.68% (-2.4-5.1%)	19 (-66-140)	180 (-440-1000)
Moldova	2010	Female	90 to 94	38% (23-55%)	62% (45-77%)	1.2 (0.57-2.1)	0.59% (-2.1-4.7%)	0.47% (-2.7-5.4%)	4.6 (-28-57)	36 (-130-300)
Moldova	2010	Female	95 plus	42% (23-62%)	58% (38-77%)	1.3 (0.5-2.5)	0.72% (-2.8-6%)	0.56% (-3.7-7.1%)	1.6 (-11-20)	8.2 (-32-72)
Moldova	2010	Male	All Ages	83% (78-88%)	17% (12-22%)	4.7 (3.2-6.4)	16% (12-20%)	15% (9.7-20%)	3700 (2400-5200)	140000 (100000-180000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	2010	Male	15 to 19	73% (58-84%)	27% (16-42%)	2.9 (1.8-4.3)	8% (5.3-11%)	14% (8.6-20%)	18 (11-27)	1600 (1100-2300)
Moldova	2010	Male	20 to 24	84% (74-91%)	16% (8.6-26%)	5.3 (3.5-7.6)	17% (12-22%)	25% (18-34%)	67 (45-93)	5300 (3900-7100)
Moldova	2010	Male	25 to 29	89% (82-94%)	11% (6-18%)	6.1 (3.9-8.8)	19% (14-24%)	28% (20-35%)	87 (63-110)	6400 (4800-8100)
Moldova	2010	Male	30 to 34	92% (85-96%)	8.3% (4-15%)	6.7 (4.4-9.6)	21% (16-26%)	28% (21-35%)	130 (96-160)	8300 (6400-10000)
Moldova	2010	Male	35 to 39	93% (87-96%)	7.4% (3.7-13%)	7.1 (4.6-10)	24% (19-28%)	30% (24-36%)	200 (150-240)	11000 (8400-13000)
Moldova	2010	Male	40 to 44	93% (88-96%)	7.1% (3.6-12%)	7.3 (4.8-10)	26% (21-30%)	32% (25-37%)	300 (230-370)	15000 (11000-18000)
Moldova	2010	Male	45 to 49	93% (85-97%)	7.5% (3.4-15%)	7.3 (4.8-10)	26% (20-30%)	31% (24-36%)	500 (380-610)	21000 (16000-26000)
Moldova	2010	Male	50 to 54	91% (84-96%)	9% (4.5-16%)	6.8 (4.5-9.6)	24% (18-29%)	28% (22-34%)	660 (490-820)	25000 (19000-31000)
Moldova	2010	Male	55 to 59	89% (81-94%)	11% (5.6-19%)	6 (3.9-8.5)	21% (14-26%)	24% (17-30%)	690 (470-880)	23000 (15000-29000)
Moldova	2010	Male	60 to 64	87% (77-94%)	13% (6.4-23%)	5.2 (3.3-7.6)	16% (9.9-22%)	19% (11-25%)	450 (260-610)	12000 (7400-17000)
Moldova	2010	Male	65 to 69	85% (73-93%)	15% (6.8-27%)	4.2 (2.6-6)	12% (5.8-18%)	13% (6.5-21%)	330 (160-510)	7700 (3700-12000)
Moldova	2010	Male	70 to 74	82% (70-91%)	18% (9.1-30%)	3.2 (1.9-4.9)	6.7% (1-14%)	7.4% (0.94-15%)	230 (28-470)	4300 (650-8700)
Moldova	2010	Male	75 to 79	79% (65-89%)	21% (11-35%)	2.6 (1.4-4.3)	3.1% (-1.8-9.7%)	3.2% (-2.5-11%)	94 (-72-320)	1500 (-860-4700)
Moldova	2010	Male	80 to 84	75% (59-86%)	25% (14-41%)	2.4 (1.4-3.8)	0.98% (-3.5-7%)	0.73% (-4.6-7.6%)	17 (-100-170)	270 (-960-1900)
Moldova	2010	Male	85 to 89	72% (56-82%)	28% (18-44%)	2.2 (1.2-3.5)	-0.25% (-4.4-5.3%)	-0.76% (-5.7-5.7%)	-9.9 (-76-78)	-29 (-510-600)
Moldova	2010	Male	90 to 94	70% (55-82%)	30% (18-45%)	2.1 (1.1-3.5)	-0.82% (-4.7-4.8%)	-1.5% (-6.3-5.2%)	-6.2 (-26-23)	-21 (-120-130)
Moldova	2010	Male	95 plus	68% (50-83%)	32% (17-50%)	2 (0.88-3.5)	-0.92% (-4.8-5.3%)	-1.7% (-6.6-5.6%)	-1.7 (-6.6-5.6)	-3.7 (-20-21)
Moldova	2016	Female	All Ages	69% (62-76%)	31% (24-38%)	2.1 (1.3-3)	7% (4.5-10%)	6.5% (2.7-11%)	1300 (560-2300)	46000 (29000-66000)
Moldova	2016	Female	15 to 19	59% (42-75%)	41% (25-58%)	1.2 (0.64-1.9)	3.3% (2.3-4.6%)	5.7% (3.2-9%)	1.6 (0.89-2.7)	360 (240-510)
Moldova	2016	Female	20 to 24	73% (57-85%)	27% (15-43%)	1.9 (1.1-2.9)	6.7% (4.9-8.9%)	7.9% (4.7-12%)	3.7 (2.1-5.8)	1100 (720-1600)
Moldova	2016	Female	25 to 29	78% (63-87%)	22% (13-37%)	2.4 (1.4-3.7)	7.3% (5.7-9.1%)	10% (6.6-14%)	8.9 (5.5-13)	1900 (1300-2500)
Moldova	2016	Female	30 to 34	81% (66-90%)	19% (9.6-34%)	2.7 (1.6-4.2)	7.7% (6-9.6%)	13% (9.2-17%)	16 (11-23)	2100 (1500-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	2016	Female	35 to 39	83% (71-91%)	17% (8.9-29%)	2.8 (1.6-4.3)	9.2% (7.4-12%)	16% (12-21%)	33 (23-44)	2700 (2000-3400)
Moldova	2016	Female	40 to 44	83% (72-92%)	17% (8.4-28%)	2.9 (1.7-4.5)	10% (7.8-13%)	18% (14-22%)	50 (36-67)	3100 (2400-4000)
Moldova	2016	Female	45 to 49	82% (66-91%)	18% (9.1-34%)	3 (1.8-4.8)	11% (8.2-14%)	18% (14-24%)	73 (53-99)	3700 (2700-4800)
Moldova	2016	Female	50 to 54	78% (64-88%)	22% (12-36%)	3 (1.7-4.7)	11% (8.5-15%)	18% (14-23%)	140 (100-190)	5900 (4300-8000)
Moldova	2016	Female	55 to 59	73% (56-86%)	27% (14-44%)	2.9 (1.5-4.8)	11% (7.6-16%)	17% (11-23%)	200 (140-290)	7200 (4800-10000)
Moldova	2016	Female	60 to 64	68% (49-83%)	32% (17-51%)	2.7 (1.4-4.5)	9.9% (5.5-15%)	13% (7.8-20%)	290 (160-450)	8400 (4800-13000)
Moldova	2016	Female	65 to 69	63% (43-81%)	37% (19-57%)	2.3 (1.2-3.7)	7.4% (3.3-13%)	9.5% (4-17%)	180 (78-310)	4200 (1800-7400)
Moldova	2016	Female	70 to 74	55% (37-72%)	45% (28-63%)	1.8 (0.89-3.1)	4.8% (0.81-10%)	5.9% (0.94-13%)	130 (21-280)	2600 (440-5400)
Moldova	2016	Female	75 to 79	45% (28-64%)	55% (36-72%)	1.4 (0.67-2.5)	2.7% (-0.66-7.7%)	3.2% (-0.94-9.4%)	100 (-33-310)	1600 (-400-4500)
Moldova	2016	Female	80 to 84	38% (22-57%)	62% (43-78%)	1.2 (0.55-2.1)	1.6% (-1.2-5.7%)	1.7% (-1.7-6.8%)	62 (-64-230)	710 (-570-2500)
Moldova	2016	Female	85 to 89	36% (21-52%)	64% (48-79%)	1.1 (0.5-2)	0.97% (-1.6-4.9%)	0.97% (-2.3-5.7%)	25 (-59-150)	230 (-390-1100)
Moldova	2016	Female	90 to 94	37% (23-54%)	63% (46-77%)	1.2 (0.55-2.2)	0.77% (-1.7-4.9%)	0.69% (-2.6-6.1%)	7.8 (-29-70)	58 (-130-370)
Moldova	2016	Female	95 plus	41% (22-63%)	59% (37-78%)	1.3 (0.49-2.6)	0.9% (-2.4-6.5%)	0.77% (-3.4-7.8%)	2.7 (-12-27)	14 (-37-98)
Moldova	2016	Male	All Ages	82% (77-87%)	18% (13-23%)	4.8 (3.1-7.1)	16% (11-20%)	15% (9.3-21%)	3400 (2000-4900)	130000 (87000-170000)
Moldova	2016	Male	15 to 19	70% (53-83%)	30% (17-47%)	2.9 (1.6-4.6)	7.6% (5.1-11%)	14% (8.3-21%)	11 (6.1-17)	980 (640-1400)
Moldova	2016	Male	20 to 24	82% (70-91%)	18% (9.4-30%)	5.3 (3.3-8.1)	15% (11-21%)	25% (17-34%)	38 (24-55)	3200 (2300-4400)
Moldova	2016	Male	25 to 29	87% (79-93%)	13% (6.8-21%)	6.1 (3.7-9.3)	17% (13-22%)	27% (19-35%)	75 (52-100)	5900 (4300-7600)
Moldova	2016	Male	30 to 34	90% (82-96%)	9.5% (4.4-18%)	6.6 (4.1-10)	19% (15-24%)	28% (21-34%)	110 (80-150)	7700 (5700-9700)
Moldova	2016	Male	35 to 39	92% (85-96%)	8.4% (4-15%)	7.1 (4.4-11)	22% (18-27%)	30% (23-36%)	170 (130-230)	10000 (7500-13000)
Moldova	2016	Male	40 to 44	92% (85-96%)	8.2% (3.8-15%)	7.3 (4.5-11)	25% (19-29%)	32% (25-37%)	250 (180-330)	13000 (9400-16000)
Moldova	2016	Male	45 to 49	91% (82-96%)	8.7% (4-18%)	7.2 (4.4-11)	25% (19-29%)	30% (24-36%)	320 (230-420)	14000 (10000-18000)
Moldova	2016	Male	50 to 54	90% (82-95%)	9.9% (4.6-18%)	6.8 (4.2-10)	23% (18-28%)	28% (21-34%)	520 (370-680)	20000 (14000-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Moldova	2016	Male	55 to 59	88% (79-94%)	12% (5.7-21%)	6 (3.7-9)	20% (14-25%)	24% (17-30%)	590 (400-780)	19000 (13000-25000)
Moldova	2016	Male	60 to 64	87% (76-94%)	13% (6.4-24%)	5.2 (3.1-8.1)	16% (10-22%)	19% (12-26%)	660 (390-930)	18000 (11000-26000)
Moldova	2016	Male	65 to 69	85% (72-93%)	15% (7-28%)	4.2 (2.5-6.5)	12% (5.8-19%)	14% (6.4-22%)	350 (160-550)	8100 (3800-13000)
Moldova	2016	Male	70 to 74	82% (70-91%)	18% (9.5-30%)	3.2 (1.8-5.2)	7.1% (1.3-14%)	8% (1.2-16%)	180 (29-380)	3500 (630-7000)
Moldova	2016	Male	75 to 79	78% (64-89%)	22% (11-36%)	2.6 (1.4-4.4)	3.5% (-1.5-10%)	3.7% (-2.1-12%)	91 (-51-290)	1400 (-600-4300)
Moldova	2016	Male	80 to 84	75% (59-86%)	25% (14-41%)	2.4 (1.3-3.9)	1.4% (-3.2-8.1%)	1.3% (-4.2-9%)	28 (-92-190)	380 (-850-2100)
Moldova	2016	Male	85 to 89	71% (55-83%)	29% (17-45%)	2.3 (1.2-3.7)	0.27% (-3.8-6.1%)	-0.21% (-5.3-6.8%)	-2.5 (-63-80)	29 (-410-640)
Moldova	2016	Male	90 to 94	69% (56-82%)	31% (18-44%)	2.1 (1.1-3.6)	-0.26% (-3.8-5.2%)	-0.99% (-5.8-6%)	-4.5 (-26-27)	-7.9 (-120-160)
Moldova	2016	Male	95 plus	68% (50-84%)	32% (16-50%)	2 (0.83-3.7)	-0.39% (-4-5.3%)	-1.2% (-6.6-1%)	-1.4 (-6.6-6.7)	-1.9 (-19-26)
Russian Federation	1990	Female	All Ages	61% (56-65%)	39% (35-44%)	1.2 (0.95-1.5)	3.5% (2.4-4.7%)	1.6% (0.12-3.3%)	14000 (1000-29000)	910000 (640000-1300000)
Russian Federation	1990	Female	15 to 19	47% (37-58%)	53% (42-63%)	0.93 (0.57-1.4)	3.4% (2.4-4.6%)	5.3% (3.1-8.3%)	170 (100-260)	21000 (14000-29000)
Russian Federation	1990	Female	20 to 24	71% (62-78%)	29% (22-38%)	1.9 (1.2-2.8)	6.6% (5-8.5%)	9.2% (5.9-13%)	310 (200-450)	42000 (30000-56000)
Russian Federation	1990	Female	25 to 29	78% (71-83%)	22% (17-29%)	1.9 (1.3-2.7)	8.2% (6.7-9.8%)	12% (8.5-15%)	540 (390-730)	70000 (52000-89000)
Russian Federation	1990	Female	30 to 34	81% (74-86%)	19% (14-26%)	2 (1.4-2.8)	9.2% (7.6-11%)	14% (11-17%)	980 (760-1200)	100000 (78000-130000)
Russian Federation	1990	Female	35 to 39	81% (76-86%)	19% (14-24%)	2.2 (1.6-3)	10% (8.4-12%)	16% (13-20%)	1600 (1200-2000)	120000 (96000-150000)
Russian Federation	1990	Female	40 to 44	80% (75-86%)	20% (14-25%)	2.1 (1.5-2.9)	10% (8.4-13%)	16% (13-20%)	2000 (1500-2500)	120000 (97000-150000)
Russian Federation	1990	Female	45 to 49	79% (74-84%)	21% (16-26%)	1.8 (1.3-2.5)	9.4% (7.4-12%)	14% (11-18%)	1700 (1300-2200)	89000 (67000-110000)
Russian Federation	1990	Female	50 to 54	75% (68-80%)	25% (20-32%)	1.5 (1-2.1)	7.7% (5.7-10%)	11% (8.1-14%)	3400 (2400-4500)	150000 (110000-200000)
Russian Federation	1990	Female	55 to 59	66% (59-73%)	34% (27-41%)	1.3 (0.84-1.8)	5.7% (3.9-7.9%)	7.8% (5-11%)	2900 (1800-4100)	110000 (72000-150000)
Russian Federation	1990	Female	60 to 64	58% (50-66%)	42% (34-50%)	1.1 (0.69-1.6)	2.9% (1.2-5.1%)	3.7% (1.2-6.6%)	2600 (860-4700)	83000 (34000-150000)
Russian Federation	1990	Female	65 to 69	50% (43-59%)	50% (41-57%)	0.91 (0.59-1.3)	1.2% (-0.45-3.1%)	1.3% (-0.81-3.8%)	1100 (-690-3100)	30000 (-11000-80000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	1990	Female	70 to 74	42% (35-51%)	58% (49-65%)	0.75 (0.46-1.1)	0.15% (-1.5-2.2%)	0.038% (-2.1-2.5%)	34 (-1800-2300)	3300 (-31000-47000)
Russian Federation	1990	Female	75 to 79	33% (25-44%)	67% (56-75%)	0.56 (0.32-0.87)	-0.27% (-1.6-1.4%)	-0.45% (-2.2-1.6%)	-700 (-3500-2600)	-7500 (-48000-40000)
Russian Federation	1990	Female	80 to 84	28% (19-39%)	72% (61-81%)	0.48 (0.28-0.76)	-0.56% (-1.9-0.78%)	-0.77% (-2.3-0.84%)	-1200 (-3600-1400)	-11000 (-37000-16000)
Russian Federation	1990	Female	85 to 89	27% (17-37%)	73% (63-83%)	0.47 (0.26-0.78)	-0.77% (-2.2-0.64%)	-1% (-2.7-0.64%)	-1000 (-2800-680)	-7100 (-20000-5900)
Russian Federation	1990	Female	90 to 94	27% (18-36%)	73% (64-82%)	0.48 (0.25-0.83)	-0.76% (-2.1-0.69%)	-1% (-2.6-0.7%)	-390 (-1000-270)	-1800 (-5100-1700)
Russian Federation	1990	Female	95 plus	28% (17-42%)	72% (58-83%)	0.5 (0.22-0.93)	-0.72% (-2.3-0.93%)	-0.95% (-2.9-0.99%)	-140 (-440-150)	-440 (-1400-560)
Russian Federation	1990	Male	All Ages	80% (75-84%)	20% (16-25%)	3.3 (2.6-4)	14% (11-17%)	12% (9.3-16%)	100000 (73000-130000)	450000 (360000-550000)
Russian Federation	1990	Male	15 to 19	64% (51-76%)	36% (24-49%)	1.5 (0.92-2.2)	7.3% (5-10%)	9.5% (6.2-14%)	790 (490-1200)	69000 (46000-97000)
Russian Federation	1990	Male	20 to 24	83% (76-89%)	17% (11-24%)	3.3 (2.2-4.8)	16% (12-21%)	19% (13-25%)	2400 (1600-3300)	190000 (140000-260000)
Russian Federation	1990	Male	25 to 29	90% (85-93%)	10% (6.6-15%)	4.3 (3-5.9)	20% (16-25%)	24% (18-31%)	4900 (3500-6300)	360000 (280000-450000)
Russian Federation	1990	Male	30 to 34	92% (88-95%)	7.9% (4.9-12%)	5.1 (3.7-6.7)	23% (19-28%)	28% (22-34%)	7900 (6000-9900)	520000 (410000-640000)
Russian Federation	1990	Male	35 to 39	92% (89-95%)	8.1% (5.1-11%)	5.7 (3.9-7.7)	25% (20-30%)	30% (24-36%)	10000 (7700-13000)	590000 (460000-740000)
Russian Federation	1990	Male	40 to 44	91% (88-94%)	8.6% (5.7-12%)	5.7 (3.9-7.7)	24% (19-29%)	29% (23-35%)	11000 (8300-14000)	560000 (430000-700000)
Russian Federation	1990	Male	45 to 49	91% (86-94%)	9.2% (5.9-14%)	5.7 (4-8)	23% (18-28%)	27% (21-33%)	9100 (6800-12000)	400000 (310000-510000)
Russian Federation	1990	Male	50 to 54	90% (86-93%)	10% (7-14%)	5.5 (3.7-7.7)	20% (14-25%)	23% (16-29%)	18000 (13000-24000)	700000 (500000-910000)
Russian Federation	1990	Male	55 to 59	87% (81-91%)	13% (8.7-19%)	4.5 (3.2-6.2)	15% (10-20%)	17% (12-23%)	14000 (9000-19000)	460000 (320000-630000)
Russian Federation	1990	Male	60 to 64	85% (79-91%)	15% (9.4-21%)	3.7 (2.5-5.4)	11% (6.1-16%)	12% (6.5-18%)	14000 (7800-23000)	420000 (240000-650000)
Russian Federation	1990	Male	65 to 69	84% (77-89%)	16% (11-23%)	3.1 (2.2-4.3)	6.7% (2.9-12%)	7.2% (2.7-13%)	5500 (2100-10000)	140000 (58000-240000)
Russian Federation	1990	Male	70 to 74	80% (73-85%)	20% (15-27%)	2.3 (1.5-3.3)	3.2% (-0.2-7.5%)	3.3% (-0.62-8.2%)	2000 (-380-5100)	42000 (-2700-100000)
Russian Federation	1990	Male	75 to 79	77% (67-84%)	23% (16-33%)	1.9 (1.2-2.9)	1.2% (-2-5.5%)	1% (-2.7-5.8%)	820 (-2200-4700)	16000 (-27000-73000)
Russian Federation	1990	Male	80 to 84	70% (58-80%)	30% (20-42%)	1.6 (1-2.4)	0.075% (-2.9-3.5%)	-0.26% (-3.6-3.6%)	-150 (-2100-2000)	510 (-20000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	1990	Male	85 to 89	61% (47-73%)	39% (27-53%)	1.3 (0.83-2)	-0.45% (-3-2.3%)	-0.86% (-3.8-2.3%)	-220 (-990-600)	-1000 (-6700-5100)
Russian Federation	1990	Male	90 to 94	57% (44-70%)	43% (30-56%)	1.2 (0.68-1.9)	-0.75% (-2.9-1.7%)	-1.2% (-3.8-1.7%)	-96 (-300-130)	-350 (-1400-800)
Russian Federation	1990	Male	95 plus	55% (38-72%)	45% (28-62%)	1.1 (0.53-1.9)	-0.7% (-3-1.8%)	-1.3% (-4.1-1.7%)	-32 (-100-43)	-71 (-300-180)
Russian Federation	1995	Female	All Ages	64% (60-68%)	36% (32-40%)	1.3 (1.1-1.6)	4.5% (3.2-5.9%)	2.3% (0.52-4.3%)	24000 (5500-45000)	140000 (98000-180000)
Russian Federation	1995	Female	15 to 19	50% (41-61%)	50% (39-59%)	1 (0.65-1.6)	4.1% (2.8-5.6%)	6.4% (3.9-9.5%)	300 (180-450)	30000 (21000-42000)
Russian Federation	1995	Female	20 to 24	73% (65-80%)	27% (20-35%)	2 (1.3-3)	7.8% (5.8-10%)	11% (7.3-16%)	590 (390-850)	61000 (45000-82000)
Russian Federation	1995	Female	25 to 29	80% (74-85%)	20% (15-26%)	2.1 (1.4-2.9)	9.9% (7.9-12%)	14% (10-18%)	860 (610-1100)	82000 (63000-100000)
Russian Federation	1995	Female	30 to 34	83% (77-87%)	17% (13-23%)	2.2 (1.5-3)	11% (9.5-14%)	17% (14-21%)	1700 (1300-2200)	140000 (110000-170000)
Russian Federation	1995	Female	35 to 39	83% (78-88%)	17% (12-22%)	2.4 (1.7-3.2)	13% (11-15%)	19% (16-23%)	3200 (2500-3900)	200000 (170000-250000)
Russian Federation	1995	Female	40 to 44	82% (77-87%)	18% (13-23%)	2.2 (1.6-3)	13% (11-15%)	19% (16-23%)	4400 (3400-5400)	240000 (190000-290000)
Russian Federation	1995	Female	45 to 49	81% (76-86%)	19% (14-24%)	1.9 (1.4-2.6)	11% (9.1-14%)	16% (13-19%)	4500 (3400-5700)	210000 (160000-260000)
Russian Federation	1995	Female	50 to 54	77% (72-82%)	23% (18-28%)	1.6 (1.1-2.3)	9.4% (7.3-12%)	13% (9.8-16%)	3300 (2400-4300)	130000 (100000-170000)
Russian Federation	1995	Female	55 to 59	69% (62-76%)	31% (24-38%)	1.4 (0.94-2)	6.8% (4.6-9.1%)	8.9% (6-12%)	5600 (3700-7800)	190000 (130000-270000)
Russian Federation	1995	Female	60 to 64	61% (53-69%)	39% (31-47%)	1.2 (0.77-1.7)	3.3% (1.2-5.9%)	4.1% (1.4-7.6%)	2900 (930-5200)	86000 (32000-150000)
Russian Federation	1995	Female	65 to 69	53% (46-62%)	47% (38-54%)	1 (0.67-1.4)	1.2% (-0.67-3.5%)	1.4% (-1.1-4.3%)	1700 (-1400-5500)	47000 (-25000-140000)
Russian Federation	1995	Female	70 to 74	45% (38-53%)	55% (47-62%)	0.84 (0.52-1.3)	0.078% (-1.9-2.3%)	-0.049% (-2.5-2.7%)	-66 (-3400-3700)	2400 (-59000-73000)
Russian Federation	1995	Female	75 to 79	36% (27-46%)	64% (54-73%)	0.63 (0.37-1)	-0.38% (-2-1.6%)	-0.57% (-2.6-1.7%)	-730 (-3300-2300)	-8400 (-45000-36000)
Russian Federation	1995	Female	80 to 84	31% (21-41%)	69% (59-79%)	0.54 (0.32-0.87)	-0.68% (-2.2-0.99%)	-0.89% (-2.7-1%)	-1700 (-5100-2000)	-16000 (-53000-23000)
Russian Federation	1995	Female	85 to 89	29% (19-39%)	71% (61-81%)	0.54 (0.31-0.88)	-0.92% (-2.5-0.77%)	-1.2% (-3-0.8%)	-1600 (-4100-1100)	-11000 (-29000-9000)
Russian Federation	1995	Female	90 to 94	29% (20-39%)	71% (61-80%)	0.54 (0.28-0.91)	-0.93% (-2.5-0.88%)	-1.2% (-3-0.91%)	-650 (-1600-530)	-3100 (-8100-2900)
Russian Federation	1995	Female	95 plus	31% (18-44%)	69% (56-82%)	0.57 (0.26-1)	-0.89% (-2.8-1.3%)	-1.1% (-3.4-1.3%)	-170 (-500-200)	-540 (-1700-760)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	1995	Male	All Ages	82% (78-86%)	18% (14-22%)	3.5 (2.9-4.2)	17% (14-21%)	15% (12-19%)	180000 (130000-240000)	7800000 (6200000-9700000)
Russian Federation	1995	Male	15 to 19	67% (55-79%)	33% (21-45%)	1.6 (1-2.5)	8.7% (5.8-12%)	11% (6.8-16%)	1400 (850-2100)	110000 (75000-160000)
Russian Federation	1995	Male	20 to 24	85% (78-90%)	15% (9.7-22%)	3.5 (2.4-5)	18% (13-23%)	20% (14-27%)	4700 (3200-6500)	350000 (250000-470000)
Russian Federation	1995	Male	25 to 29	91% (87-94%)	9.1% (6-13%)	4.5 (3-6.3)	24% (19-29%)	26% (20-33%)	7100 (5100-9200)	500000 (380000-630000)
Russian Federation	1995	Male	30 to 34	93% (89-96%)	7.2% (4.4-11%)	5.3 (3.8-7.2)	27% (22-32%)	30% (24-36%)	13000 (10000-16000)	810000 (640000-1000000)
Russian Federation	1995	Male	35 to 39	93% (89-95%)	7.4% (4.6-11%)	5.9 (4.2-7.8)	29% (23-34%)	32% (25-38%)	21000 (16000-26000)	1100000 (880000-1400000)
Russian Federation	1995	Male	40 to 44	92% (89-95%)	7.8% (5-11%)	6 (4.3-8)	29% (23-34%)	32% (25-38%)	26000 (20000-33000)	1300000 (970000-1600000)
Russian Federation	1995	Male	45 to 49	92% (88-95%)	8.3% (5.3-12%)	6 (4.2-8.3)	27% (21-32%)	29% (23-35%)	26000 (19000-34000)	1100000 (830000-1400000)
Russian Federation	1995	Male	50 to 54	91% (87-94%)	9.5% (6.5-13%)	5.8 (3.9-8)	23% (17-29%)	25% (18-32%)	18000 (13000-24000)	680000 (490000-890000)
Russian Federation	1995	Male	55 to 59	88% (84-92%)	12% (8-16%)	4.8 (3.4-6.6)	17% (12-23%)	19% (13-26%)	28000 (18000-39000)	930000 (610000-1300000)
Russian Federation	1995	Male	60 to 64	87% (81-91%)	13% (8.8-19%)	4 (2.7-5.7)	12% (6.9-19%)	13% (7.2-21%)	18000 (9700-28000)	510000 (280000-790000)
Russian Federation	1995	Male	65 to 69	86% (80-90%)	14% (9.6-20%)	3.4 (2.4-4.6)	7.4% (2.9-13%)	7.9% (2.7-14%)	14000 (4700-25000)	330000 (120000-580000)
Russian Federation	1995	Male	70 to 74	82% (76-88%)	18% (12-24%)	2.5 (1.6-3.5)	3.5% (-0.2-8.2%)	3.6% (-0.62-8.8%)	3500 (-600-8400)	71000 (-4200-160000)
Russian Federation	1995	Male	75 to 79	79% (70-86%)	21% (14-30%)	2.1 (1.3-3.2)	1.4% (-2.3-6%)	1.2% (-3-6.4%)	800 (-2000-4300)	15000 (-25000-65000)
Russian Federation	1995	Male	80 to 84	72% (61-82%)	28% (18-39%)	1.8 (1.1-2.7)	0.18% (-3.1-4%)	-0.13% (-3.9-4.2%)	-86 (-2700-2900)	1500 (-26000-33000)
Russian Federation	1995	Male	85 to 89	64% (51-75%)	36% (25-49%)	1.5 (0.94-2.2)	-0.42% (-3.4-2.7%)	-0.8% (-4.1-2.8%)	-280 (-1400-970)	-1200 (-9900-8000)
Russian Federation	1995	Male	90 to 94	60% (47-72%)	40% (28-53%)	1.3 (0.77-2)	-0.81% (-3.3-2.2%)	-1.3% (-4.1-2.1%)	-120 (-410-210)	-470 (-1900-1300)
Russian Federation	1995	Male	95 plus	58% (41-73%)	42% (27-59%)	1.2 (0.59-2.1)	-0.79% (-3.3-2.2%)	-1.4% (-4.3-2.1%)	-28 (-89-44)	-65 (-270-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2000	Female	All Ages	64% (59-68%)	36% (32-41%)	1.5 (1.2-1.8)	5.2% (3.8-6.8%)	2.9% (0.85-5.3%)	31000 (9100-56000)	1600000 (1100000-2100000)
Russian Federation	2000	Female	15 to 19	52% (41-64%)	48% (36-59%)	1.2 (0.75-1.9)	4.4% (3-6.2%)	7.4% (4.5-11%)	360 (220-540)	35000 (24000-50000)
Russian Federation	2000	Female	20 to 24	74% (63-81%)	26% (19-37%)	2.3 (1.5-3.4)	8.8% (6.7-11%)	13% (8.7-17%)	780 (520-1100)	77000 (57000-100000)
Russian Federation	2000	Female	25 to 29	78% (70-84%)	22% (16-30%)	2.3 (1.5-3.1)	11% (8.8-13%)	16% (12-19%)	1100 (800-1400)	100000 (78000-120000)
Russian Federation	2000	Female	30 to 34	80% (72-86%)	20% (14-28%)	2.3 (1.6-3.2)	12% (10-14%)	18% (15-22%)	1500 (1200-1900)	120000 (99000-150000)
Russian Federation	2000	Female	35 to 39	81% (74-87%)	19% (13-26%)	2.6 (1.9-3.5)	14% (12-16%)	21% (18-24%)	2900 (2300-3500)	190000 (160000-230000)
Russian Federation	2000	Female	40 to 44	81% (74-87%)	19% (13-26%)	2.5 (1.8-3.3)	14% (12-16%)	21% (18-24%)	4600 (3700-5600)	250000 (210000-310000)
Russian Federation	2000	Female	45 to 49	80% (72-85%)	20% (15-28%)	2.1 (1.5-2.8)	12% (10-15%)	18% (15-21%)	5400 (4300-6600)	260000 (210000-310000)
Russian Federation	2000	Female	50 to 54	74% (66-81%)	26% (19-34%)	1.8 (1.2-2.5)	10% (8.1-13%)	14% (11-17%)	5200 (4000-6700)	210000 (160000-270000)
Russian Federation	2000	Female	55 to 59	68% (59-76%)	32% (24-41%)	1.6 (1-2.3)	8.1% (5.7-11%)	11% (7.3-14%)	3600 (2500-4800)	120000 (88000-170000)
Russian Federation	2000	Female	60 to 64	63% (53-73%)	37% (27-47%)	1.4 (0.93-2.2)	4.4% (1.8-7.4%)	5.5% (2.1-9.6%)	4600 (1800-8000)	140000 (58000-240000)
Russian Federation	2000	Female	65 to 69	57% (47-67%)	43% (33-53%)	1.3 (0.81-1.8)	2% (-0.39-4.8%)	2.4% (-0.87-5.9%)	2200 (-800-5500)	57000 (-12000-130000)
Russian Federation	2000	Female	70 to 74	47% (36-58%)	53% (42-64%)	1 (0.59-1.6)	0.74% (-1.6-3.4%)	0.75% (-2.3-4%)	1300 (-3700-6700)	29000 (-63000-130000)
Russian Federation	2000	Female	75 to 79	37% (26-50%)	63% (50-74%)	0.76 (0.42-1.2)	0.11% (-2-2.5%)	0.0018% (-2.5-2.9%)	0.41 (-4100-4800)	3200 (-56000-75000)
Russian Federation	2000	Female	80 to 84	33% (23-45%)	67% (55-77%)	0.69 (0.39-1.1)	-0.29% (-2.1-1.9%)	-0.46% (-2.6-2.1%)	-610 (-3400-2800)	-4900 (-34000-31000)
Russian Federation	2000	Female	85 to 89	32% (21-43%)	68% (57-79%)	0.69 (0.37-1.1)	-0.58% (-2.4-1.5%)	-0.78% (-3-1.7%)	-1200 (-4500-2500)	-7400 (-32000-20000)
Russian Federation	2000	Female	90 to 94	32% (22-43%)	68% (57-78%)	0.7 (0.36-1.2)	-0.67% (-2.7-1.7%)	-0.88% (-3.3-1.8%)	-560 (-2100-1200)	-2500 (-10000-6200)
Russian Federation	2000	Female	95 plus	34% (21-49%)	66% (51-79%)	0.73 (0.34-1.3)	-0.66% (-3.1-2.3%)	-0.88% (-3.7-2.5%)	-140 (-600-410)	-430 (-2000-1500)
Russian Federation	2000	Male	All Ages	82% (77-86%)	18% (14-23%)	4.1 (3.4-4.8)	19% (16-22%)	17% (13-21%)	200000 (150000-250000)	840000 (680000-1000000)
Russian Federation	2000	Male	15 to 19	73% (60-83%)	27% (17-40%)	2.1 (1.3-3.1)	9.8% (6.4-14%)	12% (7.6-18%)	1600 (950-2400)	130000 (83000-180000)
Russian Federation	2000	Male	20 to 24	85% (78-90%)	15% (9.6-22%)	4 (2.7-5.6)	20% (15-25%)	22% (16-29%)	5700 (4000-7700)	420000 (310000-560000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2000	Male	25 to 29	90% (85-93%)	10% (6.7-15%)	5 (3.4-7)	25% (20-31%)	28% (21-35%)	8500 (6200-11000)	590000 (440000-740000)
Russian Federation	2000	Male	30 to 34	92% (88-95%)	7.8% (4.8-12%)	6 (4.3-8)	28% (23-33%)	32% (25-38%)	11000 (8200-13000)	670000 (530000-820000)
Russian Federation	2000	Male	35 to 39	92% (88-95%)	8% (4.9-12%)	6.6 (4.7-8.9)	31% (25-35%)	35% (28-40%)	18000 (14000-22000)	1000000 (800000-1200000)
Russian Federation	2000	Male	40 to 44	91% (87-94%)	8.7% (5.7-13%)	6.7 (4.7-8.7)	31% (25-35%)	35% (28-40%)	27000 (21000-33000)	1300000 (1000000-1600000)
Russian Federation	2000	Male	45 to 49	91% (85-94%)	9.3% (5.9-15%)	6.7 (4.8-9)	29% (24-34%)	33% (27-38%)	32000 (24000-39000)	1300000 (1000000-1700000)
Russian Federation	2000	Male	50 to 54	89% (85-93%)	11% (7.1-15%)	6.4 (4.4-8.9)	25% (19-31%)	28% (21-34%)	29000 (21000-37000)	1100000 (800000-1400000)
Russian Federation	2000	Male	55 to 59	87% (81-92%)	13% (8.3-19%)	5.3 (3.8-7.3)	20% (14-26%)	22% (15-29%)	17000 (12000-23000)	560000 (390000-730000)
Russian Federation	2000	Male	60 to 64	86% (79-91%)	14% (9-21%)	4.5 (3-6.6)	15% (8.6-21%)	16% (9.3-23%)	26000 (15000-38000)	740000 (440000-1100000)
Russian Federation	2000	Male	65 to 69	85% (78-90%)	15% (10-22%)	3.8 (2.7-5.3)	9.7% (4.6-15%)	10% (4.7-17%)	14000 (6200-23000)	340000 (150000-540000)
Russian Federation	2000	Male	70 to 74	82% (75-88%)	18% (12-25%)	2.9 (1.9-4)	5.1% (0.85-10%)	5.4% (0.55-11%)	9100 (960-19000)	180000 (30000-370000)
Russian Federation	2000	Male	75 to 79	79% (69-87%)	21% (13-31%)	2.4 (1.4-3.6)	2.5% (-1.5-7.8%)	2.5% (-2.1-8.4%)	2100 (-1800-7000)	34000 (-21000-100000)
Russian Federation	2000	Male	80 to 84	73% (60-82%)	27% (18-40%)	2.1 (1.3-3)	1.2% (-2.4-5.5%)	0.93% (-3.2-5.9%)	450 (-1500-2900)	6700 (-14000-32000)
Russian Federation	2000	Male	85 to 89	66% (53-77%)	34% (23-47%)	1.8 (1.1-2.7)	0.29% (-3-4.1%)	-0.052% (-3.9-4.3%)	-19 (-1400-1600)	930 (-9700-13000)
Russian Federation	2000	Male	90 to 94	63% (48-74%)	37% (26-52%)	1.6 (0.93-2.5)	-0.26% (-3.2-3.4%)	-0.68% (-4.1-3.5%)	-82 (-500-430)	-180 (-2300-2400)
Russian Federation	2000	Male	95 plus	60% (43-74%)	40% (26-57%)	1.4 (0.71-2.4)	-0.31% (-3.2-3.5%)	-0.85% (-4.2-3.5%)	-19 (-94-79)	-27 (-280-310)
Russian Federation	2005	Female	All Ages	64% (60-68%)	36% (32-40%)	1.8 (1.5-2.1)	6.8% (5.2-8.4%)	4.5% (2.5-6.7%)	48000 (26000-72000)	2100000 (1600000-2700000)
Russian Federation	2005	Female	15 to 19	52% (41-64%)	48% (36-59%)	1.4 (0.9-2.2)	4.8% (3.3-6.7%)	8.8% (5.5-13%)	350 (210-530)	35000 (24000-50000)
Russian Federation	2005	Female	20 to 24	73% (63-80%)	27% (20-37%)	2.6 (1.7-3.8)	9.7% (7.4-12%)	15% (11-20%)	930 (650-1300)	90000 (69000-120000)
Russian Federation	2005	Female	25 to 29	77% (70-84%)	23% (16-30%)	2.6 (1.7-3.6)	12% (10-15%)	18% (14-22%)	1600 (1200-2000)	130000 (100000-160000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2005	Female	30 to 34	80% (73-86%)	20% (14-27%)	2.7 (1.9-3.7)	15% (12-17%)	21% (17-25%)	2400 (1800-3000)	170000 (140000-210000)
Russian Federation	2005	Female	35 to 39	82% (77-87%)	18% (13-23%)	3.1 (2.2-4.1)	16% (14-19%)	24% (20-28%)	3300 (2600-4100)	200000 (160000-250000)
Russian Federation	2005	Female	40 to 44	82% (77-87%)	18% (13-23%)	2.9 (2.1-3.9)	16% (14-19%)	24% (20-28%)	5500 (4400-6700)	290000 (240000-350000)
Russian Federation	2005	Female	45 to 49	80% (72-85%)	20% (15-28%)	2.4 (1.8-3.2)	14% (12-17%)	20% (17-24%)	7000 (5700-8600)	330000 (270000-390000)
Russian Federation	2005	Female	50 to 54	74% (66-80%)	26% (20-34%)	2 (1.4-2.8)	12% (9.6-15%)	17% (13-20%)	7600 (5900-9600)	310000 (240000-380000)
Russian Federation	2005	Female	55 to 59	67% (58-75%)	33% (25-42%)	1.7 (1.2-2.5)	9.6% (7-12%)	13% (9.2-16%)	6700 (4800-8900)	230000 (170000-300000)
Russian Federation	2005	Female	60 to 64	60% (51-69%)	40% (31-49%)	1.5 (1-2.3)	5.8% (3.2-8.7%)	7.4% (4.1-11%)	3400 (1800-5100)	99000 (54000-150000)
Russian Federation	2005	Female	65 to 69	54% (45-63%)	46% (37-55%)	1.3 (0.86-1.9)	3.1% (0.67-5.8%)	3.8% (0.63-7.1%)	4300 (700-8100)	110000 (22000-200000)
Russian Federation	2005	Female	70 to 74	43% (34-53%)	57% (47-66%)	1 (0.63-1.6)	1.5% (-0.75-4.1%)	1.7% (-1.2-4.9%)	2200 (-1400-6100)	45000 (-23000-120000)
Russian Federation	2005	Female	75 to 79	32% (22-43%)	68% (57-78%)	0.74 (0.42-1.2)	0.79% (-1-2.9%)	0.82% (-1.4-3.3%)	1700 (-2900-7000)	29000 (-37000-110000)
Russian Federation	2005	Female	80 to 84	30% (21-41%)	70% (59-79%)	0.71 (0.4-1.1)	0.42% (-1.2-2.4%)	0.39% (-1.6-2.7%)	680 (-2800-4800)	9200 (-27000-52000)
Russian Federation	2005	Female	85 to 89	30% (21-41%)	70% (59-79%)	0.75 (0.42-1.2)	0.2% (-1.7-2.5%)	0.13% (-2-2.8%)	130 (-2000-2800)	1800 (-15000-22000)
Russian Federation	2005	Female	90 to 94	30% (21-40%)	70% (60-79%)	0.76 (0.4-1.2)	0.11% (-2-2.6%)	0.027% (-2.5-2.9%)	19 (-1700-2000)	460 (-8400-11000)
Russian Federation	2005	Female	95 plus	32% (19-46%)	68% (54-81%)	0.79 (0.37-1.4)	0.21% (-2.4-3.2%)	0.13% (-2.9-3.6%)	24 (-520-620)	150 (-1700-2200)
Russian Federation	2005	Male	All Ages	83% (79-86%)	17% (14-21%)	4.8 (4-5.6)	22% (19-25%)	20% (16-24%)	240000 (190000-300000)	980000 (810000-1200000)
Russian Federation	2005	Male	15 to 19	73% (60-83%)	27% (17-40%)	2.4 (1.5-3.5)	11% (7.1-15%)	14% (9-21%)	1400 (900-2100)	120000 (78000-170000)
Russian Federation	2005	Male	20 to 24	85% (78-91%)	15% (9.1-22%)	4.6 (3.2-6.3)	22% (16-27%)	25% (18-33%)	5900 (4200-7900)	440000 (320000-570000)
Russian Federation	2005	Male	25 to 29	90% (86-94%)	9.6% (6.3-14%)	5.8 (4-8)	28% (22-33%)	31% (24-38%)	11000 (8100-14000)	710000 (550000-890000)
Russian Federation	2005	Male	30 to 34	93% (89-96%)	7.4% (4.5-11%)	6.9 (5.1-9.2)	31% (26-36%)	35% (29-41%)	14000 (11000-17000)	870000 (700000-1100000)
Russian Federation	2005	Male	35 to 39	92% (89-95%)	7.8% (4.6-11%)	7.6 (5.2-10)	34% (28-39%)	38% (31-44%)	18000 (14000-21000)	960000 (770000-1100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2005	Male	40 to 44	92% (88-95%)	8.4% (5.3-12%)	7.6 (5.4-10)	34% (29-39%)	38% (32-44%)	29000 (23000-35000)	1400000 (1100000-1600000)
Russian Federation	2005	Male	45 to 49	91% (86-94%)	9% (5.7-14%)	7.7 (5.6-10)	32% (27-37%)	36% (30-41%)	38000 (31000-47000)	1600000 (1300000-1900000)
Russian Federation	2005	Male	50 to 54	90% (85-93%)	10% (7.1-15%)	7.4 (5.1-9.9)	29% (23-34%)	32% (25-37%)	40000 (31000-50000)	1500000 (1200000-1800000)
Russian Federation	2005	Male	55 to 59	87% (81-92%)	13% (8.3-19%)	6.1 (4.4-8.2)	23% (17-28%)	25% (19-31%)	31000 (22000-39000)	990000 (730000-1200000)
Russian Federation	2005	Male	60 to 64	86% (79-91%)	14% (9.3-21%)	5.1 (3.4-7.4)	17% (11-23%)	19% (11-26%)	17000 (9800-23000)	460000 (280000-630000)
Russian Federation	2005	Male	65 to 69	85% (79-90%)	15% (9.7-21%)	4.4 (3.1-6)	12% (6.2-18%)	13% (6.5-20%)	22000 (11000-33000)	520000 (250000-760000)
Russian Federation	2005	Male	70 to 74	82% (75-88%)	18% (12-25%)	3.3 (2.2-4.5)	6.9% (1.8-13%)	7.3% (1.6-14%)	9600 (2100-18000)	190000 (49000-340000)
Russian Federation	2005	Male	75 to 79	78% (67-86%)	22% (14-33%)	2.6 (1.7-3.9)	3.7% (-0.8-9.5%)	3.8% (-1.3-10%)	5400 (-1900-15000)	85000 (-18000-220000)
Russian Federation	2005	Male	80 to 84	72% (59-82%)	28% (18-41%)	2.3 (1.5-3.3)	2.2% (-1.7-7%)	2.1% (-2.3-7.6%)	1200 (-1300-4500)	15000 (-12000-49000)
Russian Federation	2005	Male	85 to 89	66% (52-77%)	34% (23-48%)	2 (1.3-3)	1.1% (-2.3-5.2%)	0.83% (-3.1-5.7%)	210 (-800-1400)	2400 (-5100-11000)
Russian Federation	2005	Male	90 to 94	62% (48-74%)	38% (26-52%)	1.8 (1.1-2.8)	0.42% (-2.8-4.3%)	0.06% (-3.8-4.6%)	8 (-500-600)	330 (-2200-3300)
Russian Federation	2005	Male	95 plus	59% (42-75%)	41% (25-58%)	1.6 (0.8-2.7)	0.4% (-2.7-4.6%)	-0.071% (-3.8-4.7%)	-1.8 (-99-120)	42 (-280-480)
Russian Federation	2010	Female	All Ages	64% (60-68%)	36% (32-40%)	1.9 (1.6-2.3)	6.6% (5-8.2%)	4.5% (2.2-6.9%)	43000 (21000-67000)	1800000 (1400000-2300000)
Russian Federation	2010	Female	15 to 19	51% (39-63%)	49% (37-61%)	1.5 (0.93-2.3)	4.6% (3.2-6.4%)	9.1% (5.4-14%)	200 (120-310)	22000 (15000-31000)
Russian Federation	2010	Female	20 to 24	71% (58-80%)	29% (20-42%)	2.7 (1.7-3.9)	9.2% (7.1-12%)	15% (10-20%)	680 (460-930)	76000 (56000-99000)
Russian Federation	2010	Female	25 to 29	76% (65-83%)	24% (17-35%)	2.7 (1.8-3.8)	12% (9.5-14%)	16% (13-20%)	1200 (920-1600)	120000 (94000-150000)
Russian Federation	2010	Female	30 to 34	80% (71-86%)	20% (14-29%)	2.9 (2-3.9)	14% (12-16%)	20% (16-23%)	2100 (1600-2600)	160000 (130000-200000)
Russian Federation	2010	Female	35 to 39	82% (75-87%)	18% (13-25%)	3.3 (2.4-4.5)	16% (13-18%)	24% (20-28%)	2900 (2300-3600)	190000 (150000-230000)
Russian Federation	2010	Female	40 to 44	81% (75-87%)	19% (13-25%)	3.1 (2.3-4.3)	15% (13-18%)	24% (20-28%)	3500 (2800-4300)	200000 (160000-240000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2010	Female	45 to 49	79% (70-85%)	21% (15-30%)	2.6 (1.8-3.5)	13% (11-16%)	20% (17-24%)	4800 (3800-5800)	230000 (190000-280000)
Russian Federation	2010	Female	50 to 54	74% (65-81%)	26% (19-35%)	2.2 (1.5-3)	11% (9.3-14%)	17% (14-20%)	6100 (4800-7700)	260000 (200000-320000)
Russian Federation	2010	Female	55 to 59	67% (57-76%)	33% (24-43%)	1.9 (1.3-2.7)	9.4% (7-12%)	13% (9.7-17%)	6600 (4800-8600)	230000 (170000-300000)
Russian Federation	2010	Female	60 to 64	61% (49-71%)	39% (29-51%)	1.7 (1.1-2.5)	6.1% (3.6-9%)	8.1% (4.7-12%)	4600 (2600-6900)	140000 (80000-210000)
Russian Federation	2010	Female	65 to 69	56% (45-66%)	44% (34-55%)	1.5 (0.98-2.2)	3.7% (1.2-6.4%)	4.7% (1.3-8.3%)	2400 (670-4100)	61000 (20000-100000)
Russian Federation	2010	Female	70 to 74	47% (36-59%)	53% (41-64%)	1.3 (0.75-2)	2.2% (-0.33-4.9%)	2.6% (-0.63-6.1%)	3400 (-830-8200)	72000 (-11000-160000)
Russian Federation	2010	Female	75 to 79	38% (26-52%)	62% (48-74%)	0.97 (0.54-1.5)	1.3% (-0.91-3.8%)	1.4% (-1.4-4.5%)	2000 (-1800-6400)	33000 (-23000-97000)
Russian Federation	2010	Female	80 to 84	33% (23-45%)	67% (55-77%)	0.86 (0.5-1.4)	0.83% (-1.1-3.1%)	0.86% (-1.5-3.6%)	1800 (-2900-7200)	22000 (-28000-83000)
Russian Federation	2010	Female	85 to 89	32% (22-42%)	68% (58-78%)	0.86 (0.48-1.4)	0.55% (-1.4-2.9%)	0.52% (-1.8-3.2%)	680 (-2500-4400)	6500 (-16000-34000)
Russian Federation	2010	Female	90 to 94	32% (22-43%)	68% (57-78%)	0.87 (0.46-1.5)	0.4% (-1.8-3.1%)	0.34% (-2.2-3.4%)	170 (-1100-1700)	1200 (-5300-9100)
Russian Federation	2010	Female	95 plus	34% (20-48%)	66% (52-80%)	0.93 (0.43-1.7)	0.63% (-2.1-4.1%)	0.61% (-2.6-4.5%)	130 (-540-910)	520 (-1700-3300)
Russian Federation	2010	Male	All Ages	82% (78-85%)	18% (15-22%)	5.2 (4.4-6.1)	21% (18-24%)	19% (15-23%)	200000 (150000-240000)	780000 (640000-930000)
Russian Federation	2010	Male	15 to 19	69% (54-81%)	31% (19-46%)	2.4 (1.5-3.6)	9.6% (6.5-13%)	15% (8.9-21%)	740 (440-1100)	62000 (41000-88000)
Russian Federation	2010	Male	20 to 24	84% (75-90%)	16% (9.9-25%)	4.8 (3.3-6.8)	20% (15-26%)	26% (18-34%)	4100 (2900-5500)	310000 (230000-410000)
Russian Federation	2010	Male	25 to 29	90% (84-93%)	10% (6.6-16%)	6.2 (4.3-8.7)	26% (21-31%)	30% (24-37%)	8100 (6200-10000)	560000 (440000-690000)
Russian Federation	2010	Male	30 to 34	92% (87-95%)	8% (4.7-13%)	7.4 (5.4-10)	29% (25-34%)	34% (28-39%)	12000 (9600-15000)	740000 (600000-900000)
Russian Federation	2010	Male	35 to 39	92% (88-95%)	8.4% (5.1-12%)	8.2 (5.6-11)	38% (27-37%)	38% (32-43%)	15000 (12000-18000)	810000 (660000-970000)
Russian Federation	2010	Male	40 to 44	91% (87-94%)	8.9% (5.7-13%)	8.2 (5.8-11)	33% (28-37%)	39% (33-44%)	17000 (13000-20000)	820000 (660000-980000)
Russian Federation	2010	Male	45 to 49	90% (84-94%)	9.8% (6.2-16%)	8.2 (6-11)	31% (27-35%)	37% (32-41%)	25000 (20000-29000)	1100000 (870000-1300000)
Russian Federation	2010	Male	50 to 54	88% (83-92%)	12% (7.7-17%)	7.8 (5.3-11)	28% (23-33%)	32% (27-37%)	31000 (24000-38000)	1200000 (910000-1400000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2010	Male	55 to 59	86% (80-91%)	14% (8.9-20%)	6.4 (4.6-8.7)	23% (18-28%)	26% (20-32%)	29000 (22000-37000)	960000 (740000-1200000)
Russian Federation	2010	Male	60 to 64	85% (77-90%)	15% (9.6-23%)	5.4 (3.7-7.9)	18% (12-23%)	20% (13-26%)	23000 (15000-31000)	640000 (420000-860000)
Russian Federation	2010	Male	65 to 69	84% (77-90%)	16% (9.9-23%)	4.7 (3.3-6.4)	13% (7.6-19%)	15% (8.1-21%)	11000 (5900-16000)	260000 (150000-370000)
Russian Federation	2010	Male	70 to 74	81% (73-87%)	19% (13-27%)	3.5 (2.3-4.9)	7.9% (2.8-14%)	8.6% (2.8-15%)	12000 (4000-22000)	240000 (85000-420000)
Russian Federation	2010	Male	75 to 79	75% (63-84%)	25% (16-37%)	2.7 (1.7-4)	4.6% (0.27-10%)	4.8% (-0.067-11%)	4900 (-71-12000)	77000 (4000-170000)
Russian Federation	2010	Male	80 to 84	70% (57-81%)	30% (19-43%)	2.4 (1.5-3.5)	3% (-0.76-8%)	3.1% (-1.3-8.8%)	3000 (-1200-8500)	36000 (-8800-94000)
Russian Federation	2010	Male	85 to 89	65% (51-76%)	35% (24-49%)	2.2 (1.4-3.2)	1.8% (-1.7-6.2%)	1.6% (-2.4-6.8%)	510 (-770-2200)	4900 (-4700-18000)
Russian Federation	2010	Male	90 to 94	61% (47-73%)	39% (27-53%)	1.9 (1.2-3)	0.93% (-2.5-5.1%)	0.63% (-3.6-5.5%)	63 (-350-550)	560 (-1500-3100)
Russian Federation	2010	Male	95 plus	59% (42-74%)	41% (26-58%)	1.7 (0.87-2.9)	0.93% (-2.4-5.6%)	0.56% (-3.5-6%)	18 (-110-200)	120 (-320-720)
Russian Federation	2016	Female	All Ages	64% (60-68%)	36% (32-40%)	1.9 (1.4-2.5)	6.2% (4.5-8.6%)	4.3% (1.8-7.2%)	43000 (16000-86000)	1700000 (1100000-2800000)
Russian Federation	2016	Female	15 to 19	51% (39-63%)	49% (37-61%)	1.5 (0.88-2.4)	4.4% (2.9-6.3%)	9.1% (5.3-14%)	130 (67-230)	15000 (9400-23000)
Russian Federation	2016	Female	20 to 24	71% (59-80%)	29% (20-41%)	2.8 (1.7-4.3)	8.7% (6.5-11%)	15% (9.8-21%)	400 (210-690)	46000 (30000-66000)
Russian Federation	2016	Female	25 to 29	77% (66-84%)	23% (16-34%)	2.8 (1.8-4.1)	11% (8.6-14%)	16% (12-21%)	1100 (570-1900)	110000 (73000-170000)
Russian Federation	2016	Female	30 to 34	80% (70-86%)	20% (14-30%)	3 (1.9-4.2)	13% (9.9-16%)	19% (15-23%)	2000 (1000-3600)	160000 (100000-250000)
Russian Federation	2016	Female	35 to 39	80% (72-86%)	20% (14-28%)	3.2 (2.1-4.6)	14% (11-18%)	22% (18-27%)	2900 (1400-5200)	190000 (120000-300000)
Russian Federation	2016	Female	40 to 44	80% (72-86%)	20% (14-28%)	3.1 (2.1-4.5)	14% (11-18%)	23% (19-27%)	3500 (1900-5900)	200000 (120000-310000)
Russian Federation	2016	Female	45 to 49	79% (69-85%)	21% (15-31%)	2.6 (1.8-3.8)	13% (9.9-16%)	20% (17-24%)	3400 (2000-5800)	170000 (110000-260000)
Russian Federation	2016	Female	50 to 54	75% (65-82%)	25% (18-35%)	2.3 (1.4-3.4)	11% (8.4-14%)	17% (13-21%)	5100 (2900-8400)	220000 (140000-340000)
Russian Federation	2016	Female	55 to 59	69% (58-78%)	31% (22-42%)	2 (1.3-3.1)	9.2% (6.6-13%)	13% (9.7-18%)	6600 (3800-11000)	230000 (150000-370000)
Russian Federation	2016	Female	60 to 64	64% (53-74%)	36% (26-47%)	1.9 (1.1-2.8)	6.1% (3.4-9.5%)	8.5% (4.7-13%)	5600 (2800-10000)	170000 (89000-300000)
Russian Federation	2016	Female	65 to 69	60% (48-70%)	40% (30-52%)	1.7 (1.1-2.5)	3.9% (1.2-7%)	5.1% (1.4-9%)	4000 (1100-8200)	100000 (32000-200000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2016	Female	70 to 74	53% (42-64%)	47% (36-58%)	1.5 (0.88-2.3)	2.5% (-0.48-6%)	3% (-0.84-7.6%)	2000 (-480-5600)	42000 (-7600-110000)
Russian Federation	2016	Female	75 to 79	43% (31-56%)	57% (44-69%)	1.2 (0.64-1.9)	1.5% (-1.1-4.8%)	1.7% (-1.6-5.7%)	3400 (-2900-13000)	55000 (-40000-190000)
Russian Federation	2016	Female	80 to 84	36% (24-47%)	64% (53-76%)	0.96 (0.51-1.6)	0.93% (-1.2-3.6%)	0.98% (-1.6-4.2%)	1500 (-2500-7100)	18000 (-25000-79000)
Russian Federation	2016	Female	85 to 89	33% (22-44%)	67% (56-78%)	0.91 (0.5-1.6)	0.59% (-1.5-3.4%)	0.54% (-1.9-3.9%)	1100 (-4200-7600)	11000 (-29000-60000)
Russian Federation	2016	Female	90 to 94	33% (23-45%)	67% (55-77%)	0.93 (0.48-1.6)	0.4% (-1.8-3.5%)	0.31% (-2.4-3.9%)	250 (-1900-3200)	1900 (-9300-17000)
Russian Federation	2016	Female	95 plus	36% (21-52%)	64% (48-79%)	0.99 (0.45-1.9)	0.65% (-2.3-4.5%)	0.62% (-2.8-5%)	120 (-520-990)	480 (-1700-3400)
Russian Federation	2016	Male	All Ages	80% (76-84%)	20% (16-24%)	5.1 (3.8-6.5)	20% (16-24%)	18% (14-23%)	180000 (110000-270000)	710000 (450000-1000000)
Russian Federation	2016	Male	15 to 19	69% (54-81%)	31% (19-46%)	2.5 (1.4-3.8)	9.3% (6-14%)	15% (8.8-23%)	500 (250-840)	43000 (25000-68000)
Russian Federation	2016	Male	20 to 24	82% (72-89%)	18% (11-28%)	4.8 (3.1-7)	20% (15-26%)	26% (18-35%)	2500 (1400-3900)	190000 (120000-280000)
Russian Federation	2016	Male	25 to 29	88% (81-92%)	12% (7.6-19%)	6.1 (4-8.5)	25% (20-31%)	30% (23-38%)	7300 (4400-11000)	510000 (330000-730000)
Russian Federation	2016	Male	30 to 34	91% (84-95%)	9.5% (5.2-16%)	7.2 (5-9.9)	28% (22-33%)	33% (26-39%)	12000 (6800-18000)	730000 (460000-1100000)
Russian Federation	2016	Male	35 to 39	90% (85-95%)	9.6% (5.4-15%)	8.1 (5.2-12)	30% (24-35%)	36% (29-41%)	14000 (8200-21000)	770000 (480000-1100000)
Russian Federation	2016	Male	40 to 44	90% (85-94%)	10% (6-15%)	8.2 (5.4-12)	31% (26-36%)	38% (32-43%)	16000 (9600-24000)	780000 (490000-1100000)
Russian Federation	2016	Male	45 to 49	89% (82-94%)	11% (6.4-18%)	8.1 (5.5-12)	30% (25-35%)	36% (31-41%)	17000 (10000-25000)	730000 (460000-1100000)
Russian Federation	2016	Male	50 to 54	87% (81-92%)	13% (7.9-19%)	7.8 (5-11)	27% (22-32%)	33% (26-38%)	25000 (15000-38000)	950000 (590000-1400000)
Russian Federation	2016	Male	55 to 59	85% (77-91%)	15% (9-23%)	6.4 (4.2-9.1)	23% (17-28%)	26% (20-32%)	28000 (17000-42000)	940000 (580000-1400000)
Russian Federation	2016	Male	60 to 64	84% (75-90%)	16% (9.9-25%)	5.4 (3.5-8.1)	18% (12-23%)	20% (13-27%)	25000 (14000-40000)	720000 (420000-1100000)
Russian Federation	2016	Male	65 to 69	84% (75-90%)	16% (10-25%)	4.7 (3.1-6.5)	13% (7.3-19%)	15% (7.9-21%)	17000 (8400-29000)	420000 (210000-680000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Russian Federation	2016	Male	70 to 74	81% (71-87%)	19% (13-29%)	3.5 (2.2-5.1)	8.3% (2.8-14%)	9.2% (2.7-16%)	6200 (1800-12000)	120000 (39000-220000)
Russian Federation	2016	Male	75 to 79	77% (65-86%)	23% (14-35%)	2.8 (1.7-4.4)	5.2% (0.37-11%)	5.6% (-0.00041-12%)	7500 (-0.65-17000)	120000 (8500-260000)
Russian Federation	2016	Male	80 to 84	71% (58-81%)	29% (19-42%)	2.5 (1.5-3.8)	3.5% (-0.78-8.8%)	3.6% (-1.4-9.7%)	2600 (-1100-7200)	32000 (-7600-81000)
Russian Federation	2016	Male	85 to 89	65% (51-77%)	35% (23-49%)	2.2 (1.3-3.4)	2% (-1.7-6.8%)	1.9% (-2.4-7.6%)	1300 (-1600-5200)	12000 (-9800-41000)
Russian Federation	2016	Male	90 to 94	61% (47-74%)	39% (26-53%)	2 (1.1-3.2)	1.1% (-2.4-5.9%)	0.84% (-3.3-6.5%)	120 (-500-990)	1000 (-2200-5400)
Russian Federation	2016	Male	95 plus	59% (42-74%)	41% (26-58%)	1.7 (0.84-3)	1% (-2.3-6%)	0.61% (-3.5-6.4%)	20 (-110-210)	130 (-310-760)
Ukraine	1990	Female	All Ages	64% (60-68%)	36% (32-40%)	2.8 (2.2-3.4)	5.5% (3.4-8%)	4% (0.83-7.7%)	14000 (2700-26000)	540000 (320000-770000)
Ukraine	1990	Female	15 to 19	56% (45-67%)	44% (33-55%)	2.4 (1.7-3.4)	5.9% (4.2-7.8%)	10% (6.4-15%)	100 (62-150)	12000 (8500-17000)
Ukraine	1990	Female	20 to 24	74% (64-81%)	26% (19-36%)	4.3 (3.1-5.8)	10% (7.8-13%)	14% (10-19%)	150 (110-200)	22000 (16000-29000)
Ukraine	1990	Female	25 to 29	79% (71-85%)	21% (15-29%)	4.1 (3-5.4)	10% (7.9-12%)	14% (10-19%)	190 (140-260)	27000 (20000-35000)
Ukraine	1990	Female	30 to 34	83% (75-88%)	17% (12-25%)	4.3 (3.1-5.6)	9.9% (7.9-12%)	15% (12-20%)	300 (230-390)	33000 (25000-41000)
Ukraine	1990	Female	35 to 39	85% (79-89%)	15% (11-21%)	4.8 (3.6-6.3)	11% (9.1-13%)	18% (14-22%)	490 (370-610)	41000 (32000-51000)
Ukraine	1990	Female	40 to 44	86% (80-90%)	14% (10-20%)	4.8 (3.5-6.1)	12% (9.7-14%)	18% (15-23%)	680 (530-850)	46000 (36000-57000)
Ukraine	1990	Female	45 to 49	85% (77-90%)	15% (9.9-23%)	4.4 (3.2-5.8)	12% (9.1-15%)	17% (13-22%)	870 (650-1100)	48000 (37000-60000)
Ukraine	1990	Female	50 to 54	78% (69-85%)	22% (15-31%)	3.7 (2.6-5)	11% (7.8-13%)	15% (10-19%)	1700 (1200-2200)	80000 (59000-100000)
Ukraine	1990	Female	55 to 59	67% (54-78%)	33% (22-46%)	3.1 (1.9-4.6)	8.9% (5.4-13%)	12% (6.6-17%)	1400 (800-2100)	56000 (33000-81000)
Ukraine	1990	Female	60 to 64	60% (47-72%)	40% (28-53%)	2.7 (1.6-4.3)	6.7% (2.7-11%)	8.5% (2.9-15%)	2100 (730-3800)	69000 (28000-120000)
Ukraine	1990	Female	65 to 69	57% (44-68%)	43% (32-56%)	2.3 (1.4-3.3)	4.7% (0.88-9%)	5.6% (0.62-11%)	1900 (210-3700)	49000 (9500-91000)
Ukraine	1990	Female	70 to 74	50% (37-61%)	50% (39-63%)	1.8 (1.1-2.7)	2.7% (-0.94-7.2%)	3.1% (-1.6-8.7%)	1000 (-520-2900)	22000 (-7800-57000)
Ukraine	1990	Female	75 to 79	40% (27-54%)	60% (46-73%)	1.4 (0.78-2.3)	1.6% (-1.7-5.9%)	1.6% (-2.3-6.8%)	1000 (-1500-4400)	18000 (-19000-67000)
Ukraine	1990	Female	80 to 84	34% (22-47%)	66% (53-78%)	1.2 (0.66-1.9)	1.2% (-1.5-4.7%)	1.3% (-1.9-5.4%)	810 (-1200-3500)	10000 (-12000-38000)
Ukraine	1990	Female	85 to 89	33% (21-48%)	67% (52-79%)	1.2 (0.61-1.9)	1.1% (-1.4-4.6%)	1.1% (-1.8-5.1%)	480 (-770-2200)	4200 (-5000-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	1990	Female	90 to 94	36% (24-51%)	64% (49-76%)	1.3 (0.66-2.1)	1.1% (-1.9-5.1%)	1.1% (-2.4-5.8%)	190 (-400-960)	1200 (-1900-5000)
Ukraine	1990	Female	95 plus	41% (24-60%)	59% (40-76%)	1.5 (0.65-2.6)	1.3% (-2.4-6.7%)	1.3% (-3.1-7.5%)	89 (-220-520)	370 (-670-1800)
Ukraine	1990	Male	All Ages	75% (71-78%)	25% (22-29%)	4.5 (3.6-5.5)	14% (11-17%)	12% (8.5-17%)	37000 (26000-51000)	1500000 (1200000-1900000)
Ukraine	1990	Male	15 to 19	60% (49-69%)	40% (31-51%)	2.5 (1.6-3.7)	8.1% (5.6-12%)	12% (8-18%)	280 (180-410)	23000 (15000-33000)
Ukraine	1990	Male	20 to 24	83% (76-88%)	17% (12-24%)	5.6 (3.8-7.8)	19% (14-25%)	25% (17-33%)	930 (650-1200)	70000 (50000-91000)
Ukraine	1990	Male	25 to 29	89% (84-92%)	11% (7.8-16%)	6.9 (4.7-9.4)	22% (17-28%)	29% (21-37%)	1500 (1100-1900)	100000 (80000-130000)
Ukraine	1990	Male	30 to 34	92% (88-95%)	7.7% (5.1-12%)	7.3 (5.2-9.7)	23% (18-29%)	30% (23-38%)	2100 (1600-2700)	130000 (100000-170000)
Ukraine	1990	Male	35 to 39	93% (89-95%)	7.1% (4.8-11%)	7.4 (5.1-10)	24% (18-31%)	31% (23-41%)	2700 (1900-3500)	150000 (110000-200000)
Ukraine	1990	Male	40 to 44	93% (90-95%)	6.6% (4.5-9.6%)	7.7 (5.5-11)	25% (19-33%)	31% (23-41%)	3200 (2400-4300)	160000 (120000-210000)
Ukraine	1990	Male	45 to 49	94% (90-96%)	6.2% (4.1-9.5%)	7.8 (5.5-11)	24% (18-31%)	29% (22-38%)	3800 (2800-5100)	170000 (120000-220000)
Ukraine	1990	Male	50 to 54	90% (85-93%)	9.9% (6.7-15%)	6.6 (4.4-9)	20% (14-26%)	24% (17-31%)	6500 (4600-8700)	250000 (180000-330000)
Ukraine	1990	Male	55 to 59	83% (75-90%)	17% (10-25%)	5.1 (3.4-7.2)	15% (10-20%)	18% (12-24%)	4500 (2900-6200)	150000 (100000-210000)
Ukraine	1990	Male	60 to 64	80% (71-87%)	20% (13-29%)	4.4 (2.8-6.6)	11% (5.6-17%)	13% (6-19%)	5200 (2500-8000)	150000 (75000-230000)
Ukraine	1990	Male	65 to 69	80% (71-87%)	20% (13-29%)	3.9 (2.6-5.4)	8.3% (3.3-14%)	9.1% (3.4-15%)	3100 (1100-5100)	74000 (30000-120000)
Ukraine	1990	Male	70 to 74	76% (66-84%)	24% (16-34%)	3.3 (2.2-4.8)	5.7% (0.97-11%)	6.1% (0.75-13%)	1500 (190-3200)	30000 (5200-60000)
Ukraine	1990	Male	75 to 79	70% (57-82%)	30% (18-43%)	2.8 (1.7-4.4)	3.6% (-0.83-9.6%)	3.8% (-1.3-11%)	1400 (-470-3800)	22000 (-4700-57000)
Ukraine	1990	Male	80 to 84	64% (50-76%)	36% (24-50%)	2.4 (1.5-3.7)	2.1% (-1.7-7.5%)	2% (-2.4-8.2%)	570 (-680-2400)	7300 (-5700-26000)
Ukraine	1990	Male	85 to 89	59% (45-71%)	41% (29-55%)	2.1 (1.3-3.2)	0.91% (-2.5-5.3%)	0.64% (-3.3-5.7%)	98 (-500-870)	1200 (-3200-7000)
Ukraine	1990	Male	90 to 94	56% (42-69%)	44% (31-58%)	1.9 (1-2.9)	0.02% (-3.1-4.2%)	-0.4% (-4.1-4.5%)	-20 (-200-230)	6.1 (-930-1300)
Ukraine	1990	Male	95 plus	54% (37-70%)	46% (30-63%)	1.7 (0.8-2.9)	-0.4% (-3.6-3.8%)	-0.97% (-4.8-3.9%)	-15 (-73-60)	-24 (-220-230)
Ukraine	1995	Female	All Ages	65% (61-69%)	35% (31-39%)	2.7 (2.2-3.3)	5.9% (3.5-8.3%)	4% (0.74-7.7%)	16000 (3000-31000)	640000 (380000-920000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	1995	Female	15 to 19	57% (46-67%)	43% (33-54%)	2.4 (1.6-3.3)	6.1% (4.3-8.2%)	10% (6.4-15%)	120 (74-170)	13000 (9200-18000)
Ukraine	1995	Female	20 to 24	75% (66-82%)	25% (18-34%)	4.1 (2.8-5.6)	10% (8.1-13%)	15% (10-20%)	220 (150-290)	26000 (19000-34000)
Ukraine	1995	Female	25 to 29	80% (72-85%)	20% (15-28%)	4 (2.8-5.2)	11% (8.2-13%)	14% (10-19%)	260 (180-350)	29000 (22000-37000)
Ukraine	1995	Female	30 to 34	83% (76-88%)	17% (12-24%)	4.1 (3.1-5.4)	11% (8.5-13%)	15% (12-20%)	410 (310-530)	38000 (29000-48000)
Ukraine	1995	Female	35 to 39	85% (79-90%)	15% (10-21%)	4.7 (3.5-6.1)	12% (9.8-15%)	18% (14-22%)	760 (580-940)	55000 (44000-68000)
Ukraine	1995	Female	40 to 44	86% (81-90%)	14% (9.6-19%)	4.6 (3.4-5.9)	13% (10-16%)	19% (15-23%)	1100 (870-1400)	68000 (54000-83000)
Ukraine	1995	Female	45 to 49	85% (78-90%)	15% (10-22%)	4.2 (3-5.6)	13% (9.8-16%)	18% (13-22%)	1400 (1100-1800)	71000 (54000-89000)
Ukraine	1995	Female	50 to 54	78% (69-85%)	22% (15-31%)	3.6 (2.5-4.9)	11% (8.1-15%)	15% (10-20%)	1500 (1000-2000)	66000 (46000-85000)
Ukraine	1995	Female	55 to 59	67% (54-78%)	33% (22-46%)	3 (1.8-4.5)	9.3% (5.3-14%)	12% (6.3-19%)	2700 (1400-4200)	98000 (56000-150000)
Ukraine	1995	Female	60 to 64	61% (48-72%)	39% (28-52%)	2.7 (1.6-4.2)	6.9% (2.5-13%)	8.4% (2.6-16%)	1900 (580-3600)	59000 (21000-110000)
Ukraine	1995	Female	65 to 69	58% (45-69%)	42% (31-55%)	2.2 (1.5-3.2)	4.5% (0.55-9.2%)	5.3% (0.27-11%)	2400 (120-5000)	62000 (7500-120000)
Ukraine	1995	Female	70 to 74	51% (38-63%)	49% (37-62%)	1.8 (1.1-2.6)	2.4% (-1.4-7%)	2.6% (-2.1-8.1%)	1500 (-1200-4700)	31000 (-18000-91000)
Ukraine	1995	Female	75 to 79	41% (28-55%)	59% (45-72%)	1.4 (0.77-2.3)	1.2% (-2.1-5.5%)	1.2% (-2.8-6.4%)	620 (-1400-3300)	11000 (-19000-48000)
Ukraine	1995	Female	80 to 84	35% (23-49%)	65% (51-77%)	1.2 (0.66-1.9)	0.97% (-1.8-4.4%)	0.93% (-2.2-5%)	740 (-1800-4000)	9500 (-18000-43000)
Ukraine	1995	Female	85 to 89	34% (22-49%)	66% (51-78%)	1.2 (0.61-1.9)	0.88% (-1.7-4.3%)	0.83% (-2.2-4.8%)	470 (-1200-2600)	4200 (-8400-20000)
Ukraine	1995	Female	90 to 94	37% (24-52%)	63% (48-76%)	1.3 (0.64-2.1)	0.91% (-2.2-5%)	0.85% (-2.7-5.6%)	190 (-610-1300)	1200 (-2900-6600)
Ukraine	1995	Female	95 plus	42% (24-62%)	58% (38-76%)	1.4 (0.63-2.6)	1.1% (-2.8-6.5%)	0.98% (-3.5-7.2%)	59 (-210-430)	250 (-660-1500)
Ukraine	1995	Male	All Ages	76% (72-79%)	24% (21-28%)	4.4 (3.6-5.3)	15% (11-19%)	13% (9-17%)	52000 (35000-70000)	210000 (160000-260000)
Ukraine	1995	Male	15 to 19	60% (50-69%)	40% (31-50%)	2.5 (1.6-3.6)	8.6% (5.8-12%)	12% (7.9-18%)	350 (220-520)	28000 (19000-40000)
Ukraine	1995	Male	20 to 24	83% (77-88%)	17% (12-23%)	5.5 (3.8-7.7)	20% (14-25%)	25% (17-32%)	1300 (910-1700)	95000 (67000-120000)
Ukraine	1995	Male	25 to 29	89% (84-92%)	11% (7.9-16%)	6.7 (4.6-8.9)	23% (18-29%)	29% (21-36%)	1900 (1300-2400)	120000 (92000-160000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	1995	Male	30 to 34	92% (88-95%)	7.7% (5.3-12%)	7.1 (5.1-9.4)	24% (19-29%)	29% (22-36%)	2900 (2200-3700)	180000 (140000-220000)
Ukraine	1995	Male	35 to 39	93% (90-95%)	7% (4.9-10%)	7.3 (5.1-9.9)	25% (19-31%)	30% (22-37%)	4300 (3200-5500)	230000 (170000-290000)
Ukraine	1995	Male	40 to 44	93% (91-96%)	6.6% (4.5-9.4%)	7.5 (5.4-10)	26% (20-32%)	30% (23-37%)	5900 (4400-7500)	280000 (220000-350000)
Ukraine	1995	Male	45 to 49	94% (90-96%)	6.2% (4.1-9.9%)	7.5 (5.3-10)	25% (19-31%)	29% (21-36%)	6600 (4900-8200)	280000 (210000-350000)
Ukraine	1995	Male	50 to 54	90% (86-93%)	9.8% (6.6-14%)	6.4 (4.2-8.7)	21% (15-27%)	24% (16-31%)	6200 (4200-8000)	230000 (160000-290000)
Ukraine	1995	Male	55 to 59	84% (75-90%)	16% (10-25%)	5 (3.3-6.9)	15% (9.9-21%)	17% (11-24%)	8600 (5400-12000)	280000 (180000-380000)
Ukraine	1995	Male	60 to 64	81% (72-87%)	19% (13-28%)	4.3 (2.8-6.4)	11% (5-17%)	12% (5.2-19%)	5100 (2200-8100)	140000 (66000-230000)
Ukraine	1995	Male	65 to 69	80% (71-88%)	20% (12-29%)	3.8 (2.6-5.3)	7.8% (2.7-13%)	8.4% (2.7-15%)	5100 (1600-9000)	120000 (41000-210000)
Ukraine	1995	Male	70 to 74	77% (68-84%)	23% (16-32%)	3.2 (2.1-4.6)	5.1% (0.19-11%)	5.4% (-0.11-12%)	2300 (-53-5200)	45000 (1700-97000)
Ukraine	1995	Male	75 to 79	71% (57-81%)	29% (19-43%)	2.7 (1.7-4.2)	3.1% (-1.1-9.1%)	3.1% (-1.6-9.9%)	890 (-480-2800)	14000 (-5200-41000)
Ukraine	1995	Male	80 to 84	65% (51-76%)	35% (24-49%)	2.4 (1.5-3.6)	1.6% (-2.3-6.8%)	1.4% (-3-7.5%)	470 (-960-2400)	6200 (-8800-27000)
Ukraine	1995	Male	85 to 89	60% (46-72%)	40% (28-54%)	2.1 (1.2-3.1)	0.46% (-3-5%)	0.17% (-3.9-5.3%)	32 (-730-1000)	730 (-4800-7900)
Ukraine	1995	Male	90 to 94	57% (43-70%)	43% (30-57%)	1.8 (1-2.9)	-0.41% (-3.7-3.8%)	-0.83% (-4.6-4.1%)	-50 (-280-250)	-140 (-1300-1300)
Ukraine	1995	Male	95 plus	55% (39-71%)	45% (29-61%)	1.6 (0.8-2.9)	-0.79% (-4.1-3.4%)	-1.3% (-5.3-3.5%)	-18 (-71-47)	-42 (-220-180)
Ukraine	2000	Female	All Ages	63% (59-67%)	37% (33-41%)	3 (2.4-3.7)	6.4% (4.2-8.7%)	4.7% (1.4-8.4%)	17000 (5300-32000)	650000 (420000-910000)
Ukraine	2000	Female	15 to 19	57% (48-66%)	43% (34-52%)	2.7 (1.8-3.7)	6.3% (4.5-8.4%)	12% (7.7-16%)	120 (77-170)	14000 (9400-18000)
Ukraine	2000	Female	20 to 24	75% (67-81%)	25% (19-33%)	4.6 (3.3-6.2)	11% (8.8-14%)	17% (12-22%)	230 (160-310)	27000 (20000-35000)
Ukraine	2000	Female	25 to 29	80% (74-85%)	20% (15-26%)	4.6 (3.4-5.9)	11% (8.9-14%)	16% (12-21%)	290 (210-380)	31000 (24000-39000)
Ukraine	2000	Female	30 to 34	84% (79-88%)	16% (12-21%)	4.8 (3.6-6.2)	12% (9.4-14%)	17% (13-22%)	410 (310-520)	36000 (28000-45000)
Ukraine	2000	Female	35 to 39	85% (80-89%)	15% (11-20%)	5.3 (4-6.8)	13% (11-16%)	20% (16-25%)	730 (570-910)	53000 (43000-64000)
Ukraine	2000	Female	40 to 44	86% (81-90%)	14% (10-19%)	5.2 (3.8-6.6)	14% (11-17%)	21% (17-26%)	1200 (910-1500)	71000 (57000-87000)
Ukraine	2000	Female	45 to 49	85% (79-90%)	15% (10-21%)	4.7 (3.5-6.2)	14% (11-17%)	20% (15-24%)	1600 (1200-2000)	80000 (62000-100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	2000	Female	50 to 54	76% (68-83%)	24% (17-32%)	3.8 (2.7-5.2)	12% (9-16%)	17% (12-22%)	1700 (1200-2300)	76000 (55000-99000)
Ukraine	2000	Female	55 to 59	62% (50-73%)	38% (27-50%)	3 (1.9-4.6)	10% (6.2-15%)	13% (7.4-20%)	1700 (960-2600)	63000 (38000-93000)
Ukraine	2000	Female	60 to 64	52% (39-65%)	48% (35-61%)	2.4 (1.4-3.9)	7.1% (3.2-12%)	8.8% (3.6-16%)	2600 (1000-4800)	81000 (35000-140000)
Ukraine	2000	Female	65 to 69	46% (34-59%)	54% (41-66%)	1.9 (1.2-2.9)	4.7% (1.4-8.7%)	5.5% (1.3-11%)	1600 (350-3200)	42000 (13000-79000)
Ukraine	2000	Female	70 to 74	40% (29-52%)	60% (48-71%)	1.5 (0.88-2.3)	2.7% (-0.4-6.6%)	3% (-0.89-8%)	1800 (-530-4700)	7000 (-5500-91000)
Ukraine	2000	Female	75 to 79	35% (24-47%)	65% (53-76%)	1.3 (0.7-2.1)	1.7% (-1.4-5.7%)	1.8% (-1.9-6.7%)	1200 (-1300-4500)	20000 (-16000-67000)
Ukraine	2000	Female	80 to 84	33% (22-46%)	67% (54-78%)	1.3 (0.7-2)	1.6% (-1.4-5.4%)	1.6% (-1.8-6.2%)	830 (-920-3100)	10000 (-8700-34000)
Ukraine	2000	Female	85 to 89	34% (22-48%)	66% (52-78%)	1.3 (0.68-2.1)	1.6% (-1.3-5.6%)	1.6% (-1.8-6.3%)	920 (-1000-3500)	7700 (-6200-27000)
Ukraine	2000	Female	90 to 94	37% (24-52%)	63% (48-76%)	1.4 (0.73-2.4)	1.7% (-1.8-6.3%)	1.7% (-2.3-7.2%)	420 (-560-1800)	2400 (-2600-9100)
Ukraine	2000	Female	95 plus	42% (23-61%)	58% (39-77%)	1.6 (0.72-2.9)	2% (-2.6-8.6%)	2% (-3.2-9.7%)	120 (-190-590)	460 (-590-2000)
Ukraine	2000	Male	All Ages	79% (76-81%)	21% (19-24%)	5.2 (4.3-6.3)	16% (13-20%)	15% (10-19%)	56000 (39000-74000)	220000 (170000-270000)
Ukraine	2000	Male	15 to 19	65% (57-72%)	35% (28-43%)	3 (2-4.3)	9.5% (6.4-13%)	14% (9.3-20%)	370 (230-530)	30000 (19000-41000)
Ukraine	2000	Male	20 to 24	86% (81-89%)	14% (11-19%)	6.4 (4.5-8.8)	22% (16-28%)	28% (20-36%)	1400 (1000-1800)	100000 (73000-130000)
Ukraine	2000	Male	25 to 29	91% (87-93%)	9.4% (7.2-13%)	7.6 (5.3-10)	24% (19-30%)	30% (23-37%)	2000 (1500-2500)	130000 (100000-160000)
Ukraine	2000	Male	30 to 34	93% (91-95%)	6.7% (4.9-9.3%)	8.1 (5.8-11)	24% (20-29%)	29% (23-35%)	2500 (1900-3000)	150000 (120000-180000)
Ukraine	2000	Male	35 to 39	94% (92-95%)	6.1% (4.5-8%)	8.3 (5.8-11)	26% (20-31%)	31% (24-37%)	4000 (3100-4900)	220000 (170000-260000)
Ukraine	2000	Male	40 to 44	94% (92-96%)	5.6% (4.1-7.6%)	8.6 (6.2-12)	28% (22-33%)	32% (26-39%)	6200 (4900-7600)	300000 (240000-360000)
Ukraine	2000	Male	45 to 49	95% (92-96%)	5.4% (3.8-8%)	8.7 (6.1-12)	27% (21-32%)	31% (24-37%)	7700 (6000-9400)	330000 (260000-400000)
Ukraine	2000	Male	50 to 54	91% (88-94%)	9% (6.2-12%)	7.3 (5-9.9)	23% (17-28%)	27% (20-33%)	7200 (5300-9100)	270000 (200000-340000)
Ukraine	2000	Male	55 to 59	85% (78-90%)	15% (9.6-22%)	5.7 (3.9-7.9)	18% (12-23%)	20% (13-27%)	5700 (3700-7600)	190000 (120000-240000)
Ukraine	2000	Male	60 to 64	82% (76-88%)	18% (12-24%)	5 (3.2-7.4)	13% (6-19%)	14% (6.4-22%)	7800 (3500-12000)	220000 (100000-330000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	2000	Male	65 to 69	81% (73-87%)	19% (13-27%)	4.3 (3-6)	9.4% (3.7-15%)	10% (3.9-17%)	4500 (1700-7400)	110000 (43000-170000)
Ukraine	2000	Male	70 to 74	76% (66-83%)	24% (17-34%)	3.5 (2.3-5)	6.3% (0.47-13%)	6.9% (0.17-14%)	4000 (110-8200)	78000 (6100-150000)
Ukraine	2000	Male	75 to 79	70% (56-81%)	30% (19-44%)	3 (1.8-4.5)	4.1% (-1-10%)	4.3% (-1.6-11%)	1600 (-580-4300)	25000 (-6100-62000)
Ukraine	2000	Male	80 to 84	64% (49-75%)	36% (25-51%)	2.6 (1.6-3.8)	2.6% (-1.7-8.1%)	2.5% (-2.4-8.8%)	510 (-490-1800)	6300 (-4300-20000)
Ukraine	2000	Male	85 to 89	58% (44-71%)	42% (29-56%)	2.2 (1.3-3.3)	1.3% (-2.5-6.4%)	1% (-3.2-6.9%)	180 (-550-1200)	1800 (-3600-9100)
Ukraine	2000	Male	90 to 94	56% (41-69%)	44% (31-59%)	2 (1.1-3.1)	0.2% (-3.4-4.9%)	-0.16% (-4.2-5.2%)	-10 (-270-330)	72 (-1200-1800)
Ukraine	2000	Male	95 plus	54% (37-71%)	46% (29-63%)	1.8 (0.88-3.2)	-0.36% (-4-4.7%)	-0.87% (-5.2-4.8%)	-11 (-63-59)	-17 (-190-220)
Ukraine	2005	Female	All Ages	63% (59-66%)	37% (34-41%)	3.7 (3-4.5)	8.3% (5.8-11%)	6.9% (3.1-12%)	26000 (12000-44000)	850000 (590000-1200000)
Ukraine	2005	Female	15 to 19	56% (49-63%)	44% (37-51%)	3.2 (2.3-4.3)	6.9% (5.1-9%)	14% (9.5-19%)	120 (78-160)	13000 (9600-17000)
Ukraine	2005	Female	20 to 24	73% (67-79%)	27% (21-33%)	5.5 (4.1-7.3)	12% (9.5-15%)	19% (14-24%)	250 (190-320)	30000 (22000-37000)
Ukraine	2005	Female	25 to 29	79% (75-83%)	21% (17-25%)	5.4 (4.1-6.9)	12% (9.8-14%)	17% (13-21%)	370 (280-470)	36000 (28000-45000)
Ukraine	2005	Female	30 to 34	83% (78-86%)	17% (14-22%)	5.7 (4.3-7.3)	13% (11-15%)	18% (15-22%)	570 (450-720)	46000 (37000-56000)
Ukraine	2005	Female	35 to 39	85% (82-88%)	15% (12-18%)	6.3 (4.9-8.1)	15% (13-18%)	23% (19-27%)	920 (740-1100)	61000 (50000-72000)
Ukraine	2005	Female	40 to 44	86% (83-89%)	14% (11-17%)	6.3 (4.8-8)	17% (14-20%)	25% (20-29%)	1500 (1200-1800)	86000 (70000-100000)
Ukraine	2005	Female	45 to 49	85% (81-89%)	15% (11-19%)	5.8 (4.4-7.5)	16% (13-20%)	23% (19-28%)	2100 (1700-2600)	100000 (83000-130000)
Ukraine	2005	Female	50 to 54	76% (69-83%)	24% (17-31%)	4.7 (3.4-6.2)	15% (11-19%)	20% (15-26%)	2500 (1900-3200)	110000 (82000-130000)
Ukraine	2005	Female	55 to 59	61% (51-72%)	39% (28-49%)	3.6 (2.3-5.4)	12% (8-17%)	16% (9.9-23%)	2600 (1600-3700)	92000 (59000-130000)
Ukraine	2005	Female	60 to 64	50% (38-62%)	50% (38-62%)	2.8 (1.7-4.4)	9.1% (4.3-15%)	11% (5.1-19%)	2200 (960-3700)	66000 (32000-110000)
Ukraine	2005	Female	65 to 69	44% (33-55%)	56% (45-67%)	2.2 (1.4-3.2)	6.3% (2.6-11%)	7.6% (2.7-14%)	3200 (1200-5800)	81000 (34000-140000)
Ukraine	2005	Female	70 to 74	39% (28-50%)	61% (50-72%)	1.7 (1-2.6)	4.2% (0.57-9.1%)	4.9% (0.32-11%)	2000 (130-4700)	41000 (5800-90000)
Ukraine	2005	Female	75 to 79	34% (24-46%)	66% (54-76%)	1.5 (0.85-2.5)	3.2% (-0.51-8.1%)	3.5% (-0.95-9.5%)	2700 (-740-7300)	42000 (-6900-110000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	2005	Female	80 to 84	33% (21-47%)	67% (53-79%)	1.6 (0.87-2.5)	3.2% (-0.47-8.2%)	3.5% (-0.76-9.4%)	2500 (-540-6900)	29000 (-4100-74000)
Ukraine	2005	Female	85 to 89	35% (22-50%)	65% (50-78%)	1.6 (0.84-2.7)	3.3% (-0.3-8.7%)	3.7% (-0.61-10%)	1400 (-230-3900)	11000 (-990-29000)
Ukraine	2005	Female	90 to 94	37% (25-53%)	63% (47-75%)	1.8 (0.91-3)	3.6% (-0.73-10%)	4% (-1-12%)	1000 (-270-3100)	5600 (-1100-16000)
Ukraine	2005	Female	95 plus	43% (25-62%)	57% (38-75%)	2.1 (0.93-3.7)	4.2% (-1.5-12%)	4.6% (-2.1-14%)	300 (-140-930)	1100 (-390-3100)
Ukraine	2005	Male	All Ages	80% (78-83%)	20% (17-22%)	6.6 (5.4-8)	19% (16-23%)	18% (13-22%)	71000 (53000-90000)	2600000 (2100000-3100000)
Ukraine	2005	Male	15 to 19	64% (58-70%)	36% (30-42%)	3.6 (2.4-5.1)	10% (7.2-15%)	17% (11-24%)	360 (230-500)	28000 (19000-39000)
Ukraine	2005	Male	20 to 24	86% (82-89%)	14% (11-18%)	7.8 (5.4-11)	24% (18-30%)	32% (24-40%)	1500 (1100-1900)	100000 (80000-130000)
Ukraine	2005	Male	25 to 29	91% (89-93%)	8.7% (6.9-11%)	9.4 (6.6-12)	26% (21-30%)	32% (25-37%)	2300 (1800-2700)	150000 (120000-180000)
Ukraine	2005	Male	30 to 34	94% (92-95%)	6.2% (4.7-8.2%)	10 (7.2-13)	26% (22-30%)	30% (26-35%)	3100 (2600-3700)	190000 (150000-220000)
Ukraine	2005	Male	35 to 39	94% (92-95%)	6% (4.5-7.7%)	10 (7.2-14)	28% (23-32%)	33% (27-38%)	4200 (3400-5000)	220000 (180000-260000)
Ukraine	2005	Male	40 to 44	95% (93-96%)	5.5% (4-7.1%)	11 (7.5-14)	30% (26-35%)	35% (30-40%)	7000 (5800-8300)	330000 (280000-390000)
Ukraine	2005	Male	45 to 49	95% (93-96%)	5% (3.5-7.2%)	11 (7.5-14)	29% (25-35%)	34% (28-40%)	9400 (7700-11000)	400000 (320000-470000)
Ukraine	2005	Male	50 to 54	91% (88-94%)	8.6% (6.2-12%)	8.9 (6.2-12)	26% (21-31%)	29% (24-35%)	9900 (7700-12000)	370000 (290000-440000)
Ukraine	2005	Male	55 to 59	85% (79-91%)	15% (9.5-21%)	6.9 (4.7-9.6)	21% (15-26%)	23% (17-30%)	8100 (5800-10000)	260000 (190000-330000)
Ukraine	2005	Male	60 to 64	83% (76-88%)	17% (12-24%)	6.1 (3.9-8.9)	16% (8.8-22%)	18% (9.6-25%)	6300 (3400-9000)	170000 (96000-250000)
Ukraine	2005	Male	65 to 69	81% (73-88%)	19% (12-27%)	5.3 (3.6-7.2)	13% (6.1-18%)	14% (6.5-21%)	8500 (4000-13000)	200000 (96000-290000)
Ukraine	2005	Male	70 to 74	76% (66-83%)	24% (17-34%)	4.3 (2.8-6.1)	9% (2.3-15%)	9.9% (2.2-17%)	4500 (1000-7700)	85000 (22000-140000)
Ukraine	2005	Male	75 to 79	70% (56-80%)	30% (20-44%)	3.6 (2.2-5.5)	6.4% (-0.15-13%)	6.9% (-0.58-15%)	3800 (-330-7900)	57000 (-1300-110000)
Ukraine	2005	Male	80 to 84	64% (49-75%)	36% (25-51%)	3.1 (1.9-4.7)	4.8% (-0.63-11%)	5% (-1.2-12%)	1500 (-370-3600)	16000 (-2200-39000)
Ukraine	2005	Male	85 to 89	58% (44-70%)	42% (30-56%)	2.7 (1.6-4.1)	3.2% (-1.5-9.1%)	3.2% (-2.3-10%)	380 (-260-1100)	3200 (-1500-8800)
Ukraine	2005	Male	90 to 94	55% (41-69%)	45% (31-59%)	2.4 (1.4-3.8)	1.8% (-2.6-7.4%)	1.6% (-3.5-8.1%)	96 (-210-490)	620 (-930-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	2005	Male	95 plus	54% (37-71%)	46% (29-63%)	2.1 (1-3.8)	0.83% (-3.7-6.6%)	0.43% (-4.8-7.2%)	5.5 (-61-91)	42 (-180-330)
Ukraine	2010	Female	All Ages	65% (61-68%)	35% (32-39%)	4.2 (3.3-5.1)	8.8% (5.9-12%)	7.8% (3.3-14%)	27000 (11000-47000)	810000 (540000-1100000)
Ukraine	2010	Female	15 to 19	55% (48-63%)	45% (37-52%)	3.4 (2.4-4.6)	6.8% (5-8.9%)	15% (10-20%)	76 (50-110)	9400 (6800-12000)
Ukraine	2010	Female	20 to 24	72% (66-78%)	28% (22-34%)	5.9 (4.3-7.8)	12% (9.3-14%)	19% (15-24%)	180 (140-230)	25000 (19000-32000)
Ukraine	2010	Female	25 to 29	78% (73-82%)	22% (18-27%)	5.8 (4.4-7.4)	12% (9.3-14%)	17% (13-20%)	290 (220-370)	33000 (26000-41000)
Ukraine	2010	Female	30 to 34	82% (77-85%)	18% (15-23%)	6 (4.6-7.8)	12% (10-14%)	17% (14-21%)	450 (350-560)	40000 (31000-48000)
Ukraine	2010	Female	35 to 39	84% (81-88%)	16% (12-19%)	6.9 (5.2-8.9)	15% (12-17%)	22% (19-26%)	770 (620-920)	54000 (44000-64000)
Ukraine	2010	Female	40 to 44	86% (82-89%)	14% (11-18%)	6.8 (5.2-8.7)	16% (13-19%)	24% (20-29%)	1100 (870-1300)	64000 (52000-75000)
Ukraine	2010	Female	45 to 49	85% (80-89%)	15% (11-20%)	6.2 (4.6-8.1)	15% (13-19%)	24% (19-29%)	1600 (1200-1900)	82000 (65000-98000)
Ukraine	2010	Female	50 to 54	78% (70-84%)	22% (16-30%)	5.3 (3.7-7.1)	15% (11-18%)	22% (16-27%)	2100 (1600-2700)	95000 (73000-120000)
Ukraine	2010	Female	55 to 59	67% (54-78%)	33% (22-46%)	4.4 (2.7-6.7)	13% (8.6-18%)	18% (12-26%)	2500 (1600-3600)	94000 (63000-130000)
Ukraine	2010	Female	60 to 64	60% (47-72%)	40% (28-53%)	3.9 (2.3-6)	11% (5.6-17%)	14% (6.9-24%)	2500 (1200-4200)	79000 (40000-120000)
Ukraine	2010	Female	65 to 69	57% (44-68%)	43% (32-56%)	3.2 (2.1-4.7)	8.5% (3.7-15%)	11% (4.1-20%)	2400 (920-4300)	61000 (26000-110000)
Ukraine	2010	Female	70 to 74	50% (37-61%)	50% (39-63%)	2.6 (1.5-3.9)	5.9% (1.1-12%)	7.1% (0.78-15%)	3700 (410-8100)	75000 (14000-160000)
Ukraine	2010	Female	75 to 79	40% (28-55%)	60% (45-72%)	2 (1.1-3.3)	4.3% (-0.15-11%)	5% (-0.68-13%)	2400 (-330-6400)	38000 (-1300-94000)
Ukraine	2010	Female	80 to 84	34% (22-48%)	66% (52-78%)	1.7 (0.99-2.8)	3.9% (0.1-9.5%)	4.4% (-0.26-12%)	3400 (-190-8800)	38000 (1100-94000)
Ukraine	2010	Female	85 to 89	33% (21-48%)	67% (52-79%)	1.7 (0.89-2.8)	3.8% (0.076-9.9%)	4.3% (-0.33-11%)	2300 (-170-6300)	18000 (380-47000)
Ukraine	2010	Female	90 to 94	36% (23-51%)	64% (49-77%)	1.9 (0.93-3.2)	4.1% (-0.41-11%)	4.6% (-0.73-13%)	840 (-130-2300)	4500 (-400-12000)
Ukraine	2010	Female	95 plus	41% (23-61%)	59% (39-77%)	2.1 (0.95-3.9)	4.8% (-1.1-14%)	5.2% (-1.7-16%)	380 (-120-1200)	1300 (-320-3900)
Ukraine	2010	Male	All Ages	80% (77-82%)	20% (18-23%)	7.1 (5.7-8.6)	18% (15-22%)	17% (13-22%)	57000 (41000-73000)	200000 (160000-240000)
Ukraine	2010	Male	15 to 19	60% (53-66%)	40% (34-47%)	3.5 (2.3-5.1)	9.6% (6.5-13%)	17% (11-24%)	200 (130-290)	17000 (11000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	2010	Male	20 to 24	85% (80-88%)	15% (12-20%)	8.2 (5.6-11)	22% (17-28%)	32% (25-40%)	1000 (760-1300)	76000 (59000-95000)
Ukraine	2010	Male	25 to 29	91% (88-93%)	9% (7.1-12%)	10 (6.9-13)	24% (20-28%)	32% (27-37%)	1700 (1400-2100)	120000 (99000-140000)
Ukraine	2010	Male	30 to 34	94% (91-95%)	6.4% (4.8-8.6%)	11 (7.6-14)	24% (21-28%)	29% (25-33%)	2300 (1900-2700)	140000 (120000-170000)
Ukraine	2010	Male	35 to 39	94% (92-95%)	6.5% (5-8.2%)	11 (7.6-14)	26% (23-30%)	32% (28-37%)	3100 (2600-3700)	170000 (150000-200000)
Ukraine	2010	Male	40 to 44	94% (92-96%)	6% (4.4-7.8%)	11 (7.9-15)	29% (25-33%)	35% (30-40%)	4200 (3500-4900)	200000 (170000-240000)
Ukraine	2010	Male	45 to 49	95% (92-96%)	5.2% (3.6-7.5%)	11 (7.9-15)	29% (24-33%)	35% (29-40%)	6200 (5100-7300)	270000 (220000-310000)
Ukraine	2010	Male	50 to 54	91% (87-94%)	9% (6.3-13%)	9.5 (6.4-13)	26% (21-31%)	31% (25-36%)	7500 (6000-9100)	280000 (230000-340000)
Ukraine	2010	Male	55 to 59	83% (75-90%)	17% (10-25%)	7.2 (4.8-10)	21% (16-26%)	25% (19-31%)	7300 (5400-9200)	240000 (180000-300000)
Ukraine	2010	Male	60 to 64	80% (71-87%)	20% (13-29%)	6.3 (4-9.3)	17% (10-23%)	19% (12-26%)	6200 (3700-8600)	170000 (110000-240000)
Ukraine	2010	Male	65 to 69	80% (71-87%)	20% (13-29%)	5.5 (3.8-7.7)	14% (7.5-19%)	15% (8.2-22%)	4700 (2500-6800)	110000 (63000-160000)
Ukraine	2010	Male	70 to 74	76% (67-84%)	24% (16-33%)	4.6 (3.1-6.6)	10% (3.5-16%)	11% (3.5-18%)	6100 (1900-10000)	120000 (41000-190000)
Ukraine	2010	Male	75 to 79	70% (57-81%)	30% (19-43%)	4 (2.5-6.1)	7.4% (0.49-14%)	8.1% (0.0056-16%)	3000 (2.2-5900)	45000 (3000-87000)
Ukraine	2010	Male	80 to 84	64% (50-76%)	36% (24-50%)	3.4 (2.1-5.3)	5.7% (-0.39-12%)	6.1% (-1-14%)	2400 (-400-5300)	27000 (-1800-57000)
Ukraine	2010	Male	85 to 89	59% (44-71%)	41% (29-56%)	3 (1.8-4.5)	4.1% (-1.2-10%)	4.2% (-1.9-11%)	690 (-310-1800)	5700 (-1700-14000)
Ukraine	2010	Male	90 to 94	56% (42-69%)	44% (31-58%)	2.6 (1.5-4.1)	2.5% (-2.3-8.6%)	2.4% (-3.3-9.4%)	110 (-140-420)	650 (-610-2200)
Ukraine	2010	Male	95 plus	54% (37-71%)	46% (29-63%)	2.4 (1.2-4.2)	1.4% (-3.5-7.9%)	1.1% (-4.6-8.7%)	15 (-62-120)	76 (-180-420)
Ukraine	2016	Female	All Ages	66% (61-70%)	34% (30-39%)	4.2 (3-5.6)	8.5% (5.4-12%)	7.5% (2.8-14%)	25000 (8600-49000)	750000 (430000-1200000)
Ukraine	2016	Female	15 to 19	57% (45-67%)	43% (33-55%)	3.5 (2.1-5.2)	6.6% (4.6-8.8%)	15% (9.6-22%)	47 (26-75)	6300 (4300-9100)
Ukraine	2016	Female	20 to 24	75% (65-82%)	25% (18-35%)	6.1 (4-8.7)	12% (9-14%)	20% (14-25%)	120 (72-190)	18000 (13000-25000)
Ukraine	2016	Female	25 to 29	79% (71-85%)	21% (15-29%)	5.9 (3.9-8.1)	12% (9.1-14%)	17% (12-22%)	260 (140-430)	32000 (23000-44000)
Ukraine	2016	Female	30 to 34	83% (76-88%)	17% (12-24%)	6.2 (4.2-8.5)	12% (9.4-15%)	18% (14-23%)	430 (240-700)	40000 (27000-56000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	2016	Female	35 to 39	85% (79-89%)	15% (11-21%)	6.9 (4.8-9.5)	14% (11-18%)	23% (18-27%)	690 (370-1100)	50000 (32000-73000)
Ukraine	2016	Female	40 to 44	86% (81-90%)	14% (9.9-19%)	6.9 (4.8-9.5)	15% (12-19%)	24% (19-29%)	950 (550-1500)	59000 (39000-85000)
Ukraine	2016	Female	45 to 49	85% (78-90%)	15% (10-22%)	6.3 (4.1-8.8)	15% (11-19%)	24% (18-29%)	1100 (680-1700)	59000 (41000-83000)
Ukraine	2016	Female	50 to 54	78% (70-85%)	22% (15-30%)	5.4 (3.4-7.7)	14% (10-18%)	21% (15-27%)	1900 (1100-2900)	85000 (57000-120000)
Ukraine	2016	Female	55 to 59	67% (54-77%)	33% (23-46%)	4.5 (2.6-7.2)	13% (8.2-18%)	18% (11-26%)	2400 (1300-3800)	90000 (55000-140000)
Ukraine	2016	Female	60 to 64	60% (47-72%)	40% (28-53%)	3.9 (2.3-6.4)	11% (5.6-16%)	14% (7-24%)	2900 (1300-5200)	92000 (46000-150000)
Ukraine	2016	Female	65 to 69	57% (44-68%)	43% (32-56%)	3.3 (2-5)	8.3% (3.3-15%)	11% (3.7-20%)	2200 (750-4300)	58000 (23000-110000)
Ukraine	2016	Female	70 to 74	50% (38-62%)	50% (38-62%)	2.6 (1.4-4.1)	5.8% (1-13%)	7.2% (0.67-17%)	2400 (180-5600)	49000 (8400-110000)
Ukraine	2016	Female	75 to 79	40% (27-54%)	60% (46-73%)	2 (1.1-3.5)	4.2% (-0.18-11%)	4.9% (-0.72-14%)	3300 (-460-9300)	52000 (-2300-140000)
Ukraine	2016	Female	80 to 84	34% (22-48%)	66% (52-78%)	1.7 (0.9-2.9)	3.7% (-0.027-9.9%)	4.3% (-0.33-12%)	2200 (-170-6500)	26000 (-170-70000)
Ukraine	2016	Female	85 to 89	33% (21-47%)	67% (53-79%)	1.7 (0.88-2.8)	3.7% (-0.043-9.8%)	4.2% (-0.45-12%)	2900 (-300-8000)	23000 (-260-61000)
Ukraine	2016	Female	90 to 94	36% (23-50%)	64% (50-77%)	1.8 (0.92-3.2)	4% (-0.5-11%)	4.5% (-0.88-13%)	1300 (-240-3900)	6900 (-790-20000)
Ukraine	2016	Female	95 plus	41% (23-60%)	59% (40-77%)	2.1 (0.93-4.1)	4.5% (-1.3-14%)	5% (-1.8-16%)	320 (-110-1000)	1100 (-310-3500)
Ukraine	2016	Male	All Ages	79% (75-82%)	21% (18-25%)	7 (5.2-9.2)	18% (14-22%)	16% (11-22%)	53000 (32000-83000)	1900000 (1200000-2600000)
Ukraine	2016	Male	15 to 19	60% (49-69%)	40% (31-51%)	3.6 (2.1-5.5)	9.3% (6-14%)	18% (11-26%)	130 (71-220)	11000 (6800-18000)
Ukraine	2016	Male	20 to 24	83% (76-88%)	17% (12-24%)	8 (5.2-12)	21% (16-27%)	32% (24-40%)	670 (410-990)	51000 (34000-72000)
Ukraine	2016	Male	25 to 29	89% (84-92%)	11% (7.8-16%)	9.7 (6.3-14)	24% (19-29%)	33% (26-38%)	1600 (1100-2300)	110000 (80000-150000)
Ukraine	2016	Male	30 to 34	92% (88-95%)	7.7% (5.2-12%)	10 (7-15)	24% (20-29%)	31% (26-36%)	2400 (1500-3400)	150000 (100000-200000)
Ukraine	2016	Male	35 to 39	93% (90-95%)	7.1% (4.9-10%)	11 (7-15)	26% (21-31%)	33% (27-38%)	3000 (1900-4300)	170000 (110000-230000)
Ukraine	2016	Male	40 to 44	93% (90-96%)	6.6% (4.5-9.5%)	11 (7.2-15)	28% (23-33%)	35% (30-40%)	4000 (2500-5700)	200000 (130000-280000)
Ukraine	2016	Male	45 to 49	94% (90-96%)	6.2% (4-9.8%)	11 (7.2-16)	28% (23-33%)	34% (29-40%)	4600 (2900-6500)	200000 (130000-280000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ukraine	2016	Male	50 to 54	90% (85-93%)	9.9% (6.7-15%)	9.2 (5.9-13)	25% (20-30%)	30% (24-36%)	6800 (4300-9900)	260000 (170000-370000)
Ukraine	2016	Male	55 to 59	83% (75-90%)	17% (10-25%)	7.2 (4.6-11)	21% (15-26%)	24% (18-31%)	7000 (4400-10000)	230000 (150000-330000)
Ukraine	2016	Male	60 to 64	80% (72-87%)	20% (13-28%)	6.3 (3.8-9.7)	16% (10-23%)	19% (11-26%)	7300 (3900-11000)	210000 (110000-310000)
Ukraine	2016	Male	65 to 69	80% (71-87%)	20% (13-29%)	5.5 (3.5-8.2)	13% (6.4-19%)	15% (7-22%)	4700 (2000-7600)	110000 (50000-180000)
Ukraine	2016	Male	70 to 74	77% (66-84%)	23% (16-34%)	4.7 (2.9-6.9)	10% (3-16%)	11% (3-19%)	3700 (940-6800)	72000 (20000-130000)
Ukraine	2016	Male	75 to 79	70% (57-81%)	30% (19-43%)	4 (2.4-6.4)	7.3% (0.47-14%)	8.1% (-0.038-16%)	4200 (-19-8800)	64000 (3900-130000)
Ukraine	2016	Male	80 to 84	64% (50-76%)	36% (24-50%)	3.4 (2.1-5.5)	5.6% (-0.45-12%)	6.1% (-1.1-14%)	1900 (-320-4400)	21000 (-1600-47000)
Ukraine	2016	Male	85 to 89	59% (45-71%)	41% (29-55%)	3 (1.8-4.7)	4.1% (-1.2-11%)	4.3% (-1.9-12%)	1200 (-520-3300)	9700 (-2900-25000)
Ukraine	2016	Male	90 to 94	56% (42-69%)	44% (31-58%)	2.7 (1.5-4.3)	2.5% (-2.3-8.7%)	2.5% (-3.3-9.6%)	170 (-240-680)	1100 (-990-3700)
Ukraine	2016	Male	95 plus	54% (38-71%)	46% (29-62%)	2.4 (1.1-4.3)	1.5% (-3.3-8%)	1.2% (-4.6-8.7%)	12 (-46-86)	58 (-130-320)
Brunei	1990	Female	All Ages	73% (66-80%)	27% (20-34%)	0.36 (0.25-0.49)	-0.13% (-0.84-0.51%)	-0.9% (-2.2-0.19%)	-3.2 (-7.6-0.72)	-31 (-190-120)
Brunei	1990	Female	15 to 19	71% (56-83%)	29% (17-44%)	0.36 (0.2-0.58)	1.1% (0.45-2%)	2.7% (0.73-5%)	0.15 (0.04-0.28)	14 (6-24)
Brunei	1990	Female	20 to 24	80% (68-90%)	20% (10-32%)	0.62 (0.37-0.93)	1.3% (0.43-2.4%)	2.5% (0.22-5%)	0.21 (0.019-0.43)	23 (7.6-40)
Brunei	1990	Female	25 to 29	84% (74-91%)	16% (8.7-26%)	0.65 (0.42-0.93)	0.87% (-0.0062-1.8%)	1.7% (-0.2-3.9%)	0.2 (-0.024-0.46)	18 (-0.13-36)
Brunei	1990	Female	30 to 34	86% (75-93%)	14% (7.4-25%)	0.64 (0.41-0.92)	0.45% (-0.45-1.4%)	1.2% (-0.54-3.1%)	0.2 (-0.09-0.52)	10 (-11-31)
Brunei	1990	Female	35 to 39	86% (76-93%)	14% (7.2-24%)	0.59 (0.36-0.9)	0.29% (-0.7-1.2%)	1.2% (-0.51-2.7%)	0.22 (-0.099-0.53)	6.1 (-14-26)
Brunei	1990	Female	40 to 44	85% (76-92%)	15% (8.1-24%)	0.56 (0.32-0.9)	0.11% (-0.98-1.1%)	0.95% (-0.76-2.5%)	0.17 (-0.13-0.46)	1.7 (-16-18)
Brunei	1990	Female	45 to 49	84% (72-92%)	16% (8.2-28%)	0.53 (0.3-0.87)	-0.36% (-1.7-0.82%)	0.25% (-1.6-1.9%)	0.04 (-0.23-0.3)	-4.1 (-19-8.9)
Brunei	1990	Female	50 to 54	82% (71-90%)	18% (10-29%)	0.5 (0.28-0.84)	-0.85% (-2.4-0.43%)	-0.4% (-2.3-1.2%)	-0.077 (-0.45-0.24)	-9.8 (-29-4.9)
Brunei	1990	Female	55 to 59	80% (66-89%)	20% (11-34%)	0.47 (0.27-0.76)	-1.5% (-3.3--0.09%)	-1.2% (-3.3-0.46%)	-0.28 (-0.77-0.1)	-17 (-37--0.98)
Brunei	1990	Female	60 to 64	77% (62-88%)	23% (12-38%)	0.42 (0.23-0.71)	-2.2% (-4.2--0.56%)	-2.2% (-4.4--0.31%)	-0.47 (-0.98--0.068)	-19 (-35--4.7)
Brunei	1990	Female	65 to 69	74% (57-87%)	26% (13-43%)	0.35 (0.2-0.57)	-2.2% (-4.1--0.62%)	-2.3% (-4.4-0.46%)	-0.56 (-1.1--0.11)	-17 (-31--4.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	1990	Female	70 to 74	73% (57-84%)	27% (16-43%)	0.29 (0.15-0.47)	-1.8% (-3.6--0.44%)	-1.9% (-4--0.35%)	-0.49 (-0.99--0.087)	-11 (-22--2.7)
Brunei	1990	Female	75 to 79	72% (56-84%)	28% (16-44%)	0.26 (0.13-0.44)	-1.8% (-3.7--0.47%)	-2.1% (-4.2--0.52%)	-0.63 (-1.3--0.16)	-10 (-21--2.6)
Brunei	1990	Female	80 to 84	70% (53-83%)	30% (17-47%)	0.28 (0.15-0.45)	-2.1% (-3.9--0.66%)	-2.4% (-4.5--0.74%)	-0.82 (-1.6--0.25)	-9.3 (-18--3)
Brunei	1990	Female	85 to 89	68% (52-80%)	32% (20-48%)	0.29 (0.16-0.48)	-2.1% (-3.7--0.78%)	-2.4% (-4.3--0.88%)	-0.62 (-1.1--0.22)	-4.9 (-8.6--1.8)
Brunei	1990	Female	90 to 94	68% (52-81%)	32% (19-48%)	0.31 (0.16-0.52)	-2.2% (-3.9--0.75%)	-2.5% (-4.6--0.87%)	-0.36 (-0.66--0.13)	-1.9 (-3.5--0.67)
Brunei	1990	Female	95 plus	68% (48-83%)	32% (17-52%)	0.33 (0.16-0.57)	-2.1% (-4--0.64%)	-2.6% (-4.9--0.78%)	-0.11 (-0.21--0.034)	-0.39 (-0.73--0.11)
Brunei	1990	Male	All Ages	88% (84-92%)	12% (8.2-16%)	0.36 (0.25-0.48)	1.5% (0.39-2.6%)	0.68% (-0.83-2%)	3.6 (-4.1-11)	430 (110-780)
Brunei	1990	Male	15 to 19	84% (73-92%)	16% (7.9-27%)	0.29 (0.16-0.46)	2.1% (0.88-3.8%)	2.8% (0.52-5.7%)	0.36 (0.064-0.76)	37 (15-67)
Brunei	1990	Male	20 to 24	91% (84-95%)	9.2% (4.7-16%)	0.52 (0.32-0.76)	4% (1.8-6.9%)	4.3% (0.84-8.5%)	0.88 (0.16-1.8)	91 (40-160)
Brunei	1990	Male	25 to 29	93% (88-97%)	6.9% (3.5-12%)	0.6 (0.39-0.87)	3.7% (1.8-5.9%)	3.8% (0.79-7.3%)	0.86 (0.18-1.7)	91 (43-150)
Brunei	1990	Male	30 to 34	94% (89-97%)	5.6% (2.6-11%)	0.62 (0.41-0.89)	2.8% (1.3-4.5%)	2.7% (0.35-5.6%)	0.71 (0.097-1.5)	74 (33-130)
Brunei	1990	Male	35 to 39	95% (91-98%)	5% (2.4-9.1%)	0.58 (0.35-0.89)	2.3% (0.97-3.8%)	2.4% (0.39-4.7%)	0.65 (0.097-1.4)	58 (23-100)
Brunei	1990	Male	40 to 44	95% (91-98%)	4.7% (2-8.7%)	0.57 (0.34-0.89)	2% (0.5-3.5%)	2.2% (0.016-4.3%)	0.57 (0.0044-1.2)	40 (10-73)
Brunei	1990	Male	45 to 49	95% (90-98%)	4.6% (2-9.8%)	0.59 (0.34-0.93)	1.5% (-0.39-3%)	1.6% (-0.79-3.6%)	0.37 (-0.2-0.86)	22 (-5.9-46)
Brunei	1990	Male	50 to 54	95% (90-98%)	4.9% (2.1-9.8%)	0.57 (0.31-0.94)	0.97% (-0.95-2.6%)	1% (-1.4-2.9%)	0.28 (-0.39-0.84)	14 (-14-36)
Brunei	1990	Male	55 to 59	95% (90-98%)	5.3% (2.5-10%)	0.53 (0.3-0.86)	0.64% (-1.2-2.1%)	0.66% (-1.5-2.4%)	0.2 (-0.45-0.73)	8.6 (-16-28)
Brunei	1990	Male	60 to 64	94% (88-98%)	5.9% (2.5-12%)	0.5 (0.27-0.82)	0.2% (-1.6-1.5%)	0.11% (-1.9-1.6%)	0.043 (-0.72-0.62)	2.6 (-20-21)
Brunei	1990	Male	65 to 69	94% (88-97%)	6.5% (2.6-12%)	0.46 (0.28-0.74)	-0.08% (-1.8-1.3%)	-0.21% (-2.1-1.3%)	-0.086 (-0.84-0.54)	-0.94 (-20-15)
Brunei	1990	Male	70 to 74	93% (87-97%)	7% (3.2-13%)	0.42 (0.24-0.67)	-0.31% (-1.9-0.9%)	-0.46% (-2.2-0.87%)	-0.2 (-0.97-0.39)	-3.1 (-19-9)
Brunei	1990	Male	75 to 79	93% (87-96%)	7.3% (3.5-13%)	0.39 (0.21-0.65)	-0.52% (-2-0.62%)	-0.73% (-2.4-0.54%)	-0.31 (-1-0.23)	-3.9 (-14-4.5)
Brunei	1990	Male	80 to 84	92% (85-96%)	7.9% (3.8-15%)	0.36 (0.21-0.58)	-0.61% (-1.9-0.48%)	-0.84% (-2.3-0.38%)	-0.36 (-1-0.16)	-3.3 (-11-2.5)
Brunei	1990	Male	85 to 89	91% (85-96%)	8.6% (4.4-15%)	0.33 (0.19-0.54)	-0.65% (-1.8-0.34%)	-0.92% (-2.2-0.2%)	-0.25 (-0.62-0.056)	-1.6 (-4.6-0.82)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	1990	Male	90 to 94	91% (84-96%)	9% (4.3-16%)	0.31 (0.18-0.5)	-0.68% (-1.9-0.25%)	-0.99% (-2.4-0.065%)	-0.13 (-0.31-0.0083)	-0.54 (-1.5-0.2)
Brunei	1990	Male	95 plus	91% (82-96%)	9.1% (3.5-18%)	0.29 (0.15-0.5)	-0.67% (-1.9-0.22%)	-1% (-2.6--0.00057%)	-0.035 (-0.086--0.000019)	-0.093 (-0.27-0.031)
Brunei	1995	Female	All Ages	73% (66-80%)	27% (20-34%)	0.18 (0.11-0.27)	0.23% (-0.28-0.66%)	-0.076% (-0.93-0.62%)	-0.29 (-3.5-2.5)	59 (-70-170)
Brunei	1995	Female	15 to 19	70% (55-83%)	30% (17-45%)	0.17 (0.085-0.29)	0.85% (0.47-1.4%)	1.8% (0.77-3.2%)	0.11 (0.048-0.2)	13 (7-20)
Brunei	1995	Female	20 to 24	80% (68-89%)	20% (11-32%)	0.3 (0.16-0.48)	1.1% (0.56-1.8%)	1.7% (0.32-3.2%)	0.15 (0.03-0.29)	20 (10-34)
Brunei	1995	Female	25 to 29	84% (74-92%)	16% (8.4-26%)	0.33 (0.18-0.5)	0.85% (0.27-1.5%)	1.4% (0.041-2.8%)	0.2 (0.0063-0.41)	22 (6.8-38)
Brunei	1995	Female	30 to 34	86% (76-93%)	14% (7.2-24%)	0.32 (0.18-0.5)	0.63% (0.011-1.2%)	1.2% (0.039-2.5%)	0.2 (0.0063-0.42)	14 (0.34-29)
Brunei	1995	Female	35 to 39	86% (76-93%)	14% (7.3-24%)	0.29 (0.15-0.47)	0.55% (-0.16-1.2%)	1.2% (0.16-2.2%)	0.26 (0.038-0.49)	13 (-3.8-28)
Brunei	1995	Female	40 to 44	86% (76-92%)	14% (7.7-24%)	0.27 (0.13-0.46)	0.46% (-0.38-1.2%)	1.2% (0.041-2.2%)	0.27 (0.011-0.53)	9.7 (-8.6-25)
Brunei	1995	Female	45 to 49	84% (72-92%)	16% (8.2-28%)	0.27 (0.14-0.45)	0.2% (-0.77-0.94%)	0.74% (-0.47-1.7%)	0.15 (-0.1-0.37)	3.1 (-12-14)
Brunei	1995	Female	50 to 54	82% (70-90%)	18% (9.8-30%)	0.25 (0.12-0.43)	-0.049% (-1.2-0.8%)	0.37% (-1-1.4%)	0.068 (-0.17-0.27)	-0.53 (-14-8.9)
Brunei	1995	Female	55 to 59	80% (66-89%)	20% (11-34%)	0.23 (0.12-0.39)	-0.37% (-1.7-0.53%)	-0.043% (-1.5-0.95%)	-0.0078 (-0.29-0.19)	-3.5 (-16-5.2)
Brunei	1995	Female	60 to 64	77% (61-88%)	23% (12-39%)	0.21 (0.1-0.37)	-0.77% (-2.2-0.21%)	-0.65% (-2.2-0.44%)	-0.19 (-0.63-0.13)	-9 (-26-2.6)
Brunei	1995	Female	65 to 69	74% (59-87%)	26% (13-41%)	0.19 (0.094-0.32)	-0.85% (-2.2-0.099%)	-0.81% (-2.3-0.24%)	-0.22 (-0.63-0.068)	-7.3 (-19-0.85)
Brunei	1995	Female	70 to 74	72% (57-84%)	28% (16-43%)	0.17 (0.081-0.28)	-0.78% (-2.1-0.091%)	-0.78% (-2.3-0.19%)	-0.24 (-0.69-0.057)	-6 (-16-0.7)
Brunei	1995	Female	75 to 79	71% (56-83%)	29% (17-44%)	0.15 (0.071-0.27)	-0.84% (-2.1-0.037%)	-0.92% (-2.4-0.12%)	-0.27 (-0.72-0.035)	-4.6 (-12-0.21)
Brunei	1995	Female	80 to 84	70% (52-83%)	30% (17-48%)	0.14 (0.072-0.25)	-0.85% (-2-0.017%)	-0.97% (-2.3-0.0032%)	-0.32 (-0.79-0.0011)	-3.7 (-9-0.073)
Brunei	1995	Female	85 to 89	68% (51-80%)	32% (20-49%)	0.13 (0.066-0.23)	-0.75% (-1.8--0.054%)	-0.85% (-2.1--0.051%)	-0.28 (-0.69--0.016)	-2.2 (-5.3--0.15)
Brunei	1995	Female	90 to 94	67% (51-81%)	33% (19-49%)	0.13 (0.064-0.23)	-0.69% (-1.7--0.043%)	-0.8% (-2--0.02%)	-0.13 (-0.32--0.0032)	-0.69 (-1.7--0.044)
Brunei	1995	Female	95 plus	68% (48-83%)	32% (17-52%)	0.12 (0.054-0.24)	-0.6% (-1.6-0.26%)	-0.72% (-1.9-0.3%)	-0.042 (-0.12-0.018)	-0.14 (-0.4-0.061)
Brunei	1995	Male	All Ages	88% (84-92%)	12% (7.9-16%)	0.17 (0.11-0.26)	1.4% (0.75-2.1%)	0.96% (0.11-1.8%)	5.6 (0.62-11)	460 (240-700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	1995	Male	15 to 19	84% (73-92%)	16% (8-27%)	0.14 (0.067-0.23)	1.4% (0.74-2.4%)	1.5% (0.36-3.2%)	0.22 (0.053-0.46)	28 (15-48)
Brunei	1995	Male	20 to 24	91% (83-95%)	9.3% (4.6-17%)	0.25 (0.13-0.4)	2.9% (1.6-4.5%)	2.5% (0.61-5.2%)	0.53 (0.13-1.1)	69 (38-110)
Brunei	1995	Male	25 to 29	93% (88-96%)	7% (3.6-12%)	0.3 (0.16-0.46)	2.8% (1.7-4.2%)	2.5% (0.75-4.7%)	0.63 (0.2-1.3)	78 (45-120)
Brunei	1995	Male	30 to 34	94% (89-97%)	5.6% (2.7-11%)	0.31 (0.17-0.48)	2.4% (1.4-3.6%)	2.2% (0.76-4.1%)	0.56 (0.19-1.1)	63 (35-98)
Brunei	1995	Male	35 to 39	95% (91-98%)	5% (2.3-9.5%)	0.28 (0.15-0.45)	2.2% (1.3-3.2%)	2.3% (0.97-3.7%)	0.75 (0.31-1.3)	65 (37-98)
Brunei	1995	Male	40 to 44	95% (91-98%)	4.7% (2.1-9%)	0.27 (0.14-0.45)	2% (1.1-3%)	2.3% (0.96-3.6%)	0.85 (0.34-1.5)	57 (30-88)
Brunei	1995	Male	45 to 49	95% (90-98%)	4.6% (2-9.6%)	0.29 (0.14-0.48)	1.7% (0.49-2.7%)	1.9% (0.44-3.2%)	0.69 (0.16-1.2)	39 (11-64)
Brunei	1995	Male	50 to 54	95% (91-98%)	4.8% (2.2-9.2%)	0.28 (0.14-0.47)	1.4% (0.2-2.4%)	1.6% (0.2-2.8%)	0.53 (0.072-0.94)	24 (3.3-42)
Brunei	1995	Male	55 to 59	95% (90-98%)	5.4% (2.4-9.8%)	0.26 (0.13-0.45)	1.1% (-0.097-2%)	1.2% (-0.061-2.3%)	0.44 (-0.021-0.83)	17 (-1.5-33)
Brunei	1995	Male	60 to 64	94% (88-97%)	5.9% (2.6-12%)	0.24 (0.12-0.42)	0.71% (-0.36-1.6%)	0.77% (-0.48-1.7%)	0.31 (-0.2-0.69)	10 (-5.4-23)
Brunei	1995	Male	65 to 69	94% (87-97%)	6.5% (2.7-13%)	0.23 (0.12-0.37)	0.43% (-0.55-1.2%)	0.45% (-0.62-1.3%)	0.22 (-0.3-0.62)	6.1 (-7.4-17)
Brunei	1995	Male	70 to 74	93% (87-97%)	6.9% (3.2-13%)	0.21 (0.11-0.36)	0.18% (-0.73-0.94%)	0.16% (-0.84-0.96%)	0.075 (-0.41-0.45)	2 (-8-10)
Brunei	1995	Male	75 to 79	93% (87-97%)	7.2% (3.5-13%)	0.2 (0.096-0.33)	-0.025% (-0.85-0.62%)	-0.093% (-1.1-0.63%)	-0.04 (-0.45-0.27)	-0.18 (-6.3-4.7)
Brunei	1995	Male	80 to 84	92% (86-96%)	7.8% (3.6-14%)	0.17 (0.091-0.29)	-0.12% (-0.86-0.45%)	-0.2% (-1-0.41%)	-0.076 (-0.38-0.15)	-0.56 (-4.1-2.1)
Brunei	1995	Male	85 to 89	91% (85-96%)	8.7% (4.4-15%)	0.15 (0.082-0.26)	-0.15% (-0.73-0.34%)	-0.25% (-0.95-0.29%)	-0.083 (-0.31-0.094)	-0.44 (-2.2-1)
Brunei	1995	Male	90 to 94	91% (84-96%)	9% (4.4-16%)	0.14 (0.069-0.23)	-0.17% (-0.74-0.26%)	-0.29% (-0.98-0.21%)	-0.042 (-0.15-0.031)	-0.15 (-0.68-0.23)
Brunei	1995	Male	95 plus	91% (82-96%)	9.3% (3.7-18%)	0.12 (0.053-0.22)	-0.13% (-0.74-0.26%)	-0.27% (-0.98-0.21%)	-0.011 (-0.041-0.0088)	-0.023 (-0.13-0.045)
Brunei	2000	Female	All Ages	74% (67-80%)	26% (20-33%)	0.065 (0.023-0.12)	0.57% (0.31-0.73%)	0.67% (0.16-0.91%)	2.5 (0.64-3.6)	160 (85-220)
Brunei	2000	Female	15 to 19	70% (54-83%)	30% (17-46%)	0.059 (0.019-0.12)	0.66% (0.43-0.99%)	1.2% (0.61-2.1%)	0.073 (0.037-0.13)	10 (6.4-15)
Brunei	2000	Female	20 to 24	80% (67-89%)	20% (11-33%)	0.1 (0.035-0.2)	1% (0.67-1.5%)	1.2% (0.61-2%)	0.11 (0.056-0.19)	22 (14-33)
Brunei	2000	Female	25 to 29	84% (74-91%)	16% (8.6-26%)	0.11 (0.038-0.21)	0.91% (0.59-1.2%)	1.2% (0.6-2%)	0.15 (0.068-0.25)	23 (14-32)
Brunei	2000	Female	30 to 34	86% (76-93%)	14% (7.2-24%)	0.11 (0.038-0.2)	0.8% (0.45-1.2%)	1.3% (0.73-2.1%)	0.21 (0.11-0.35)	22 (13-32)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	2000	Female	35 to 39	86% (76-93%)	14% (7-24%)	0.1 (0.032-0.19)	0.76% (0.43-1.1%)	1.4% (0.87-2%)	0.26 (0.15-0.4)	20 (11-28)
Brunei	2000	Female	40 to 44	85% (76-92%)	15% (7.9-24%)	0.094 (0.029-0.19)	0.75% (0.36-1.1%)	1.4% (0.84-2%)	0.31 (0.18-0.47)	18 (9-27)
Brunei	2000	Female	45 to 49	84% (70-92%)	16% (8.2-30%)	0.092 (0.03-0.19)	0.67% (0.21-1%)	1.1% (0.6-1.7%)	0.28 (0.14-0.43)	14 (4.3-22)
Brunei	2000	Female	50 to 54	82% (69-90%)	18% (9.9-31%)	0.087 (0.028-0.17)	0.63% (0.1-1%)	1% (0.38-1.5%)	0.24 (0.082-0.39)	9.8 (1.5-16)
Brunei	2000	Female	55 to 59	79% (64-89%)	21% (11-36%)	0.08 (0.027-0.17)	0.51% (-0.14-0.96%)	0.86% (0.13-1.4%)	0.17 (0.026-0.29)	5.4 (-1.4-10)
Brunei	2000	Female	60 to 64	76% (60-87%)	24% (13-40%)	0.072 (0.024-0.16)	0.31% (-0.49-0.78%)	0.51% (-0.34-1.1%)	0.14 (-0.093-0.29)	3.5 (-5.4-8.8)
Brunei	2000	Female	65 to 69	74% (58-86%)	26% (14-42%)	0.062 (0.022-0.13)	0.32% (-0.48-0.77%)	0.49% (-0.42-1%)	0.16 (-0.14-0.33)	3.5 (-5.2-8.2)
Brunei	2000	Female	70 to 74	72% (56-84%)	28% (16-44%)	0.053 (0.018-0.11)	0.35% (-0.43-0.73%)	0.49% (-0.42-0.96%)	0.089 (-0.079-0.17)	1.6 (-2.1-3.4)
Brunei	2000	Female	75 to 79	71% (56-83%)	29% (17-44%)	0.048 (0.016-0.1)	0.31% (-0.44-0.67%)	0.41% (-0.48-0.86%)	0.095 (-0.11-0.2)	1.4 (-2-3)
Brunei	2000	Female	80 to 84	69% (52-83%)	31% (17-48%)	0.047 (0.016-0.097)	0.26% (-0.45-0.59%)	0.31% (-0.53-0.72%)	0.097 (-0.17-0.23)	1.1 (-2-2.5)
Brunei	2000	Female	85 to 89	67% (51-80%)	33% (20-49%)	0.047 (0.016-0.095)	0.22% (-0.45-0.55%)	0.27% (-0.51-0.7%)	0.08 (-0.15-0.2)	0.59 (-1.2-1.5)
Brunei	2000	Female	90 to 94	67% (51-81%)	33% (19-49%)	0.047 (0.015-0.098)	0.18% (-0.46-0.52%)	0.23% (-0.52-0.64%)	0.046 (-0.1-0.13)	0.23 (-0.58-0.64)
Brunei	2000	Female	95 plus	67% (48-83%)	33% (17-52%)	0.047 (0.013-0.1)	0.18% (-0.48-0.54%)	0.24% (-0.58-0.7%)	0.015 (-0.036-0.045)	0.049 (-0.13-0.14)
Brunei	2000	Male	All Ages	88% (84-92%)	12% (8.3-16%)	0.061 (0.022-0.11)	1.3% (1-1.6%)	1% (0.72-1.4%)	5.7 (3.9-7.6)	420 (320-550)
Brunei	2000	Male	15 to 19	84% (73-92%)	16% (8.2-27%)	0.048 (0.015-0.097)	0.89% (0.59-1.4%)	0.52% (0.25-1.4%)	0.067 (0.031-0.19)	17 (10-27)
Brunei	2000	Male	20 to 24	91% (84-95%)	9.3% (4.8-16%)	0.087 (0.03-0.17)	2% (1.2-3.1%)	0.91% (0.31-2.4%)	0.19 (0.062-0.49)	48 (28-77)
Brunei	2000	Male	25 to 29	93% (88-96%)	7% (3.7-12%)	0.1 (0.034-0.19)	2.1% (1.4-3%)	1.2% (0.49-2.5%)	0.3 (0.12-0.61)	61 (38-89)
Brunei	2000	Male	30 to 34	94% (90-97%)	5.6% (2.7-10%)	0.1 (0.035-0.2)	2% (1.4-2.8%)	1.5% (0.83-2.5%)	0.39 (0.2-0.66)	57 (38-82)
Brunei	2000	Male	35 to 39	95% (91-98%)	5% (2.3-9.1%)	0.097 (0.031-0.19)	2% (1.5-2.6%)	2% (1.3-2.9%)	0.58 (0.35-0.89)	56 (39-77)
Brunei	2000	Male	40 to 44	95% (91-98%)	4.6% (2.1-8.8%)	0.096 (0.03-0.19)	1.9% (1.4-2.6%)	2.3% (1.6-3.2%)	0.84 (0.53-1.2)	57 (40-78)
Brunei	2000	Male	45 to 49	95% (91-98%)	4.5% (2-8.9%)	0.097 (0.031-0.19)	1.7% (1.2-2.3%)	2.1% (1.5-2.9%)	0.83 (0.53-1.2)	46 (32-63)
Brunei	2000	Male	50 to 54	95% (91-98%)	4.8% (2.2-8.8%)	0.095 (0.031-0.19)	1.5% (1-2%)	1.9% (1.2-2.5%)	0.69 (0.43-0.96)	31 (20-41)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	2000	Male	55 to 59	95% (90-98%)	5.4% (2.4-10%)	0.089 (0.03-0.19)	1.2% (0.73-1.7%)	1.5% (0.91-2.1%)	0.49 (0.28-0.69)	18 (11-26)
Brunei	2000	Male	60 to 64	94% (88-97%)	6% (2.6-12%)	0.082 (0.027-0.17)	0.94% (0.5-1.3%)	1.1% (0.63-1.6%)	0.43 (0.23-0.61)	13 (6.9-19)
Brunei	2000	Male	65 to 69	93% (87-97%)	6.6% (2.8-13%)	0.075 (0.027-0.15)	0.71% (0.33-0.98%)	0.84% (0.37-1.2%)	0.36 (0.16-0.51)	9.2 (4.1-13)
Brunei	2000	Male	70 to 74	93% (87-97%)	7.1% (3.3-13%)	0.067 (0.022-0.14)	0.48% (0.13-0.69%)	0.56% (0.14-0.83%)	0.19 (0.047-0.28)	3.8 (0.99-5.5)
Brunei	2000	Male	75 to 79	93% (86-97%)	7.5% (3.4-14%)	0.061 (0.02-0.12)	0.33% (-0.023-0.51%)	0.38% (-0.044-0.62%)	0.15 (-0.018-0.24)	2.3 (-0.16-3.6)
Brunei	2000	Male	80 to 84	92% (85-96%)	8.1% (3.9-15%)	0.057 (0.02-0.11)	0.22% (-0.098-0.36%)	0.24% (-0.14-0.42%)	0.1 (-0.056-0.17)	1.2 (-0.52-1.9)
Brunei	2000	Male	85 to 89	91% (84-95%)	8.9% (4.6-16%)	0.053 (0.02-0.11)	0.17% (-0.14-0.28%)	0.18% (-0.22-0.32%)	0.054 (-0.066-0.097)	0.46 (-0.39-0.78)
Brunei	2000	Male	90 to 94	91% (84-96%)	9.1% (4.4-16%)	0.051 (0.018-0.1)	0.13% (-0.19-0.24%)	0.13% (-0.25-0.27%)	0.024 (-0.046-0.048)	0.14 (-0.21-0.27)
Brunei	2000	Male	95 plus	91% (82-96%)	9.2% (3.7-18%)	0.049 (0.014-0.1)	0.12% (-0.2-0.23%)	0.12% (-0.31-0.24%)	0.0057 (-0.015-0.012)	0.025 (-0.04-0.047)
Brunei	2005	Female	All Ages	75% (68-81%)	25% (19-32%)	0.052 (0.014-0.1)	0.6% (0.37-0.74%)	0.71% (0.27-0.9%)	3.2 (1.2-4.2)	180 (110-240)
Brunei	2005	Female	15 to 19	70% (53-83%)	30% (17-47%)	0.046 (0.0094-0.1)	0.63% (0.4-0.91%)	1.1% (0.59-1.8%)	0.069 (0.037-0.12)	11 (6.8-16)
Brunei	2005	Female	20 to 24	80% (66-90%)	20% (10-34%)	0.081 (0.019-0.17)	0.99% (0.68-1.4%)	1.1% (0.6-1.8%)	0.096 (0.052-0.16)	22 (14-33)
Brunei	2005	Female	25 to 29	84% (74-91%)	16% (8.5-26%)	0.087 (0.022-0.18)	0.92% (0.63-1.2%)	1.2% (0.66-1.9%)	0.13 (0.069-0.21)	23 (15-32)
Brunei	2005	Female	30 to 34	86% (76-93%)	14% (7.4-24%)	0.085 (0.023-0.18)	0.81% (0.52-1.1%)	1.3% (0.78-2%)	0.18 (0.1-0.31)	22 (13-32)
Brunei	2005	Female	35 to 39	86% (76-93%)	14% (7-24%)	0.078 (0.02-0.17)	0.78% (0.5-1.1%)	1.3% (0.89-1.9%)	0.25 (0.15-0.36)	21 (13-29)
Brunei	2005	Female	40 to 44	86% (76-92%)	14% (7.8-24%)	0.073 (0.018-0.16)	0.77% (0.42-1.1%)	1.4% (0.89-2%)	0.33 (0.2-0.48)	21 (11-29)
Brunei	2005	Female	45 to 49	84% (72-92%)	16% (8.2-28%)	0.071 (0.017-0.15)	0.71% (0.31-1%)	1.2% (0.68-1.7%)	0.32 (0.17-0.47)	17 (7.9-25)
Brunei	2005	Female	50 to 54	82% (71-90%)	18% (9.9-29%)	0.067 (0.016-0.14)	0.68% (0.19-1%)	1.1% (0.52-1.5%)	0.32 (0.15-0.48)	14 (4.2-21)
Brunei	2005	Female	55 to 59	79% (65-89%)	21% (11-35%)	0.062 (0.013-0.14)	0.61% (-0.0052-0.94%)	0.96% (0.29-1.4%)	0.29 (0.096-0.44)	10 (-0.086-16)
Brunei	2005	Female	60 to 64	76% (61-88%)	24% (12-39%)	0.056 (0.011-0.13)	0.45% (-0.27-0.8%)	0.66% (-0.15-1.1%)	0.22 (-0.048-0.38)	6.4 (-4-11)
Brunei	2005	Female	65 to 69	73% (58-86%)	27% (14-42%)	0.048 (0.01-0.1)	0.46% (-0.23-0.77%)	0.64% (-0.14-1%)	0.22 (-0.05-0.36)	5.3 (-2.7-8.8)
Brunei	2005	Female	70 to 74	72% (57-83%)	28% (17-43%)	0.041 (0.0092-0.087)	0.44% (-0.25-0.71%)	0.59% (-0.19-0.95%)	0.2 (-0.066-0.33)	3.9 (-2.1-6.3)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	2005	Female	75 to 79	70% (55-82%)	30% (18-45%)	0.037 (0.0082-0.081)	0.39% (-0.27-0.64%)	0.5% (-0.26-0.83%)	0.19 (-0.1-0.32)	2.9 (-2-4.7)
Brunei	2005	Female	80 to 84	69% (51-83%)	31% (17-49%)	0.036 (0.0077-0.08)	0.33% (-0.29-0.55%)	0.39% (-0.33-0.68%)	0.14 (-0.12-0.24)	1.6 (-1.3-2.6)
Brunei	2005	Female	85 to 89	67% (49-80%)	33% (20-51%)	0.036 (0.0078-0.08)	0.29% (-0.3-0.53%)	0.35% (-0.34-0.67%)	0.13 (-0.13-0.25)	0.99 (-1.1-1.9)
Brunei	2005	Female	90 to 94	67% (51-80%)	33% (20-49%)	0.037 (0.0082-0.08)	0.25% (-0.35-0.48%)	0.31% (-0.4-0.61%)	0.072 (-0.092-0.14)	0.36 (-0.52-0.71)
Brunei	2005	Female	95 plus	67% (47-83%)	33% (17-53%)	0.037 (0.0073-0.085)	0.24% (-0.34-0.51%)	0.31% (-0.4-0.65%)	0.031 (-0.041-0.066)	0.1 (-0.15-0.22)
Brunei	2005	Male	All Ages	89% (85-92%)	11% (7.7-15%)	0.049 (0.013-0.098)	1.3% (1-1.5%)	0.98% (0.75-1.2%)	6.2 (4.6-8)	450 (350-570)
Brunei	2005	Male	15 to 19	84% (73-92%)	16% (8-27%)	0.036 (0.0077-0.078)	0.91% (0.62-1.3%)	0.5% (0.26-1.1%)	0.066 (0.033-0.14)	19 (12-28)
Brunei	2005	Male	20 to 24	91% (83-95%)	9.3% (4.8-17%)	0.066 (0.016-0.14)	2% (1.3-3%)	0.72% (0.33-1.8%)	0.14 (0.058-0.37)	50 (30-78)
Brunei	2005	Male	25 to 29	93% (88-96%)	7% (3.6-12%)	0.078 (0.019-0.16)	2.2% (1.5-3%)	1.1% (0.49-2.1%)	0.23 (0.097-0.45)	59 (38-86)
Brunei	2005	Male	30 to 34	94% (89-97%)	5.7% (2.7-11%)	0.081 (0.022-0.17)	2% (1.5-2.7%)	1.4% (0.87-2.2%)	0.31 (0.18-0.53)	55 (37-78)
Brunei	2005	Male	35 to 39	95% (91-98%)	5.1% (2.4-9.5%)	0.075 (0.019-0.16)	2% (1.5-2.6%)	2% (1.4-2.8%)	0.51 (0.32-0.75)	54 (39-72)
Brunei	2005	Male	40 to 44	95% (91-98%)	4.7% (2.1-9.1%)	0.073 (0.018-0.16)	1.9% (1.5-2.4%)	2.3% (1.7-3.2%)	0.77 (0.52-1.1)	56 (40-74)
Brunei	2005	Male	45 to 49	95% (91-98%)	4.6% (1.9-9.5%)	0.075 (0.017-0.16)	1.7% (1.3-2.2%)	2.1% (1.6-2.8%)	0.85 (0.58-1.2)	49 (36-66)
Brunei	2005	Male	50 to 54	95% (91-98%)	4.9% (2.1-9.3%)	0.073 (0.016-0.15)	1.5% (1.1-1.9%)	1.9% (1.4-2.5%)	0.86 (0.58-1.2)	39 (28-52)
Brunei	2005	Male	55 to 59	95% (90-97%)	5.4% (2.6-10%)	0.067 (0.014-0.15)	1.2% (0.8-1.6%)	1.5% (0.98-2%)	0.7 (0.46-0.98)	26 (18-36)
Brunei	2005	Male	60 to 64	94% (88-97%)	6% (2.5-12%)	0.063 (0.013-0.14)	0.94% (0.58-1.3%)	1.2% (0.74-1.6%)	0.55 (0.34-0.76)	17 (10-23)
Brunei	2005	Male	65 to 69	94% (88-97%)	6.5% (2.7-12%)	0.06 (0.013-0.12)	0.72% (0.42-0.95%)	0.87% (0.51-1.2%)	0.44 (0.26-0.59)	11 (6.4-14)
Brunei	2005	Male	70 to 74	93% (87-97%)	7% (3.3-13%)	0.056 (0.013-0.12)	0.5% (0.18-0.68%)	0.59% (0.22-0.84%)	0.3 (0.11-0.43)	5.9 (2.2-8.2)
Brunei	2005	Male	75 to 79	93% (87-96%)	7.4% (3.6-13%)	0.051 (0.011-0.12)	0.35% (0.056-0.51%)	0.41% (0.055-0.62%)	0.23 (0.031-0.35)	3.5 (0.59-5.1)
Brunei	2005	Male	80 to 84	92% (85-96%)	8.1% (3.9-15%)	0.045 (0.01-0.098)	0.24% (0.0036-0.35%)	0.27% (-0.011-0.42%)	0.14 (-0.0054-0.21)	1.5 (0.023-2.3)
Brunei	2005	Male	85 to 89	91% (84-95%)	9% (4.6-16%)	0.039 (0.0087-0.086)	0.19% (0.035-0.27%)	0.21% (-0.006-0.32%)	0.09 (-0.0025-0.13)	0.73 (0.13-1.1)
Brunei	2005	Male	90 to 94	91% (84-95%)	9.2% (4.7-16%)	0.035 (0.0074-0.075)	0.15% (0.098-0.24%)	0.18% (0.088-0.28%)	0.037 (0.018-0.058)	0.2 (0.13-0.31)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	2005	Male	95 plus	91% (82-96%)	9.3% (4-18%)	0.03 (0.0058-0.067)	0.15% (0.098-0.22%)	0.16% (0.1-0.26%)	0.012 (0.0076-0.019)	0.046 (0.03-0.07)
Brunei	2010	Female	All Ages	75% (69-81%)	25% (19-31%)	0.052 (0.0042-0.12)	0.58% (0.31-0.73%)	0.67% (0.2-0.87%)	3.5 (1-4.7)	200 (110-250)
Brunei	2010	Female	15 to 19	70% (55-84%)	30% (16-45%)	0.044 (0.0033-0.11)	0.63% (0.41-0.9%)	1.1% (0.63-1.9%)	0.073 (0.04-0.13)	11 (7.4-17)
Brunei	2010	Female	20 to 24	80% (68-89%)	20% (11-32%)	0.078 (0.0061-0.19)	1% (0.68-1.4%)	1.1% (0.62-1.8%)	0.094 (0.051-0.16)	22 (14-33)
Brunei	2010	Female	25 to 29	84% (74-91%)	16% (8.7-26%)	0.086 (0.0062-0.21)	0.91% (0.64-1.2%)	1.1% (0.65-1.8%)	0.12 (0.063-0.21)	22 (15-32)
Brunei	2010	Female	30 to 34	86% (76-93%)	14% (7.2-24%)	0.084 (0.0068-0.21)	0.79% (0.5-1.1%)	1.2% (0.74-1.9%)	0.16 (0.096-0.27)	21 (13-30)
Brunei	2010	Female	35 to 39	86% (76-93%)	14% (6.9-24%)	0.074 (0.006-0.19)	0.76% (0.49-1%)	1.3% (0.85-2%)	0.24 (0.15-0.37)	21 (14-30)
Brunei	2010	Female	40 to 44	85% (76-92%)	15% (7.6-24%)	0.07 (0.0059-0.18)	0.75% (0.4-1%)	1.4% (0.89-1.9%)	0.34 (0.21-0.5)	22 (12-31)
Brunei	2010	Female	45 to 49	84% (70-92%)	16% (8.3-30%)	0.07 (0.0056-0.18)	0.71% (0.35-0.99%)	1.2% (0.75-1.7%)	0.36 (0.22-0.52)	19 (9.7-27)
Brunei	2010	Female	50 to 54	82% (70-90%)	18% (9.8-30%)	0.066 (0.005-0.17)	0.67% (0.22-0.97%)	1.1% (0.63-1.5%)	0.4 (0.23-0.58)	17 (6-26)
Brunei	2010	Female	55 to 59	79% (66-89%)	21% (11-34%)	0.059 (0.0046-0.15)	0.59% (0.033-0.93%)	0.95% (0.37-1.4%)	0.39 (0.15-0.6)	14 (0.82-22)
Brunei	2010	Female	60 to 64	76% (61-88%)	24% (12-39%)	0.053 (0.0043-0.14)	0.44% (-0.26-0.76%)	0.66% (-0.095-1.1%)	0.26 (-0.038-0.45)	7.7 (-4.6-14)
Brunei	2010	Female	65 to 69	74% (58-86%)	26% (14-42%)	0.048 (0.0037-0.12)	0.42% (-0.3-0.74%)	0.6% (-0.19-1%)	0.22 (-0.065-0.37)	5.2 (-3.7-9.1)
Brunei	2010	Female	70 to 74	72% (56-83%)	28% (17-44%)	0.043 (0.0032-0.11)	0.38% (-0.35-0.69%)	0.53% (-0.31-0.92%)	0.25 (-0.15-0.44)	4.9 (-4.4-8.7)
Brunei	2010	Female	75 to 79	71% (55-83%)	29% (17-45%)	0.039 (0.0028-0.096)	0.32% (-0.38-0.61%)	0.42% (-0.38-0.79%)	0.22 (-0.2-0.42)	3.3 (-3.8-6.2)
Brunei	2010	Female	80 to 84	69% (51-83%)	31% (17-49%)	0.037 (0.0029-0.09)	0.27% (-0.39-0.52%)	0.33% (-0.45-0.64%)	0.17 (-0.24-0.33)	1.9 (-2.8-3.6)
Brunei	2010	Female	85 to 89	67% (50-80%)	33% (20-50%)	0.035 (0.0025-0.086)	0.24% (-0.38-0.47%)	0.29% (-0.43-0.59%)	0.11 (-0.16-0.22)	0.82 (-1.3-1.6)
Brunei	2010	Female	90 to 94	67% (52-80%)	33% (20-48%)	0.034 (0.0023-0.088)	0.22% (-0.36-0.44%)	0.27% (-0.42-0.55%)	0.071 (-0.11-0.14)	0.36 (-0.6-0.72)
Brunei	2010	Female	95 plus	67% (48-83%)	33% (17-52%)	0.032 (0.0021-0.085)	0.24% (-0.37-0.45%)	0.3% (-0.43-0.58%)	0.031 (-0.044-0.06)	0.1 (-0.15-0.2)
Brunei	2010	Male	All Ages	89% (86-93%)	11% (7.4-14%)	0.049 (0.0038-0.12)	1.3% (1-1.5%)	0.98% (0.72-1.3%)	7.3 (5.3-9.8)	510 (400-660)
Brunei	2010	Male	15 to 19	84% (73-92%)	16% (8.4-27%)	0.034 (0.0027-0.088)	0.94% (0.63-1.4%)	0.54% (0.27-1.2%)	0.078 (0.037-0.17)	21 (13-33)
Brunei	2010	Male	20 to 24	91% (84-95%)	9.4% (4.8-16%)	0.064 (0.005-0.16)	2% (1.3-3%)	0.77% (0.33-2.1%)	0.17 (0.066-0.47)	57 (35-89)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	2010	Male	25 to 29	93% (88-97%)	7% (3.5-12%)	0.076 (0.0056-0.18)	2.2% (1.6-3%)	1.1% (0.53-2.3%)	0.24 (0.11-0.53)	64 (42-94)
Brunei	2010	Male	30 to 34	94% (89-98%)	5.7% (2.4-11%)	0.078 (0.0063-0.19)	2% (1.5-2.8%)	1.4% (0.86-2.3%)	0.33 (0.19-0.56)	60 (40-84)
Brunei	2010	Male	35 to 39	95% (90-98%)	5% (2.3-9.7%)	0.072 (0.0055-0.18)	2% (1.5-2.6%)	2% (1.4-2.8%)	0.54 (0.35-0.82)	59 (43-79)
Brunei	2010	Male	40 to 44	95% (91-98%)	4.7% (2-9%)	0.07 (0.0057-0.17)	1.9% (1.4-2.5%)	2.3% (1.7-3.3%)	0.82 (0.54-1.2)	60 (43-81)
Brunei	2010	Male	45 to 49	95% (91-98%)	4.5% (2-9%)	0.072 (0.0057-0.18)	1.7% (1.3-2.2%)	2.1% (1.5-2.9%)	0.96 (0.66-1.4)	56 (40-75)
Brunei	2010	Male	50 to 54	95% (91-98%)	4.8% (2.1-8.9%)	0.071 (0.0053-0.17)	1.5% (1-2%)	1.9% (1.2-2.6%)	1.1 (0.69-1.5)	50 (34-67)
Brunei	2010	Male	55 to 59	95% (90-98%)	5.4% (2.5-10%)	0.066 (0.0053-0.17)	1.2% (0.75-1.6%)	1.5% (0.93-2.1%)	0.93 (0.58-1.3)	35 (22-49)
Brunei	2010	Male	60 to 64	94% (88-97%)	5.9% (2.6-12%)	0.061 (0.0048-0.16)	0.93% (0.56-1.3%)	1.2% (0.67-1.6%)	0.67 (0.38-0.95)	21 (12-29)
Brunei	2010	Male	65 to 69	94% (88-97%)	6.5% (2.8-12%)	0.057 (0.0046-0.14)	0.72% (0.41-0.99%)	0.87% (0.49-1.2%)	0.52 (0.29-0.73)	13 (7.3-18)
Brunei	2010	Male	70 to 74	93% (87-97%)	7.1% (3.3-13%)	0.052 (0.0041-0.13)	0.5% (0.19-0.69%)	0.6% (0.23-0.83%)	0.41 (0.15-0.57)	8.1 (3-11)
Brunei	2010	Male	75 to 79	93% (86-96%)	7.5% (3.7-14%)	0.047 (0.0035-0.12)	0.35% (0.076-0.5%)	0.41% (0.062-0.6%)	0.3 (0.046-0.44)	4.5 (0.97-6.6)
Brunei	2010	Male	80 to 84	92% (85-96%)	8.1% (3.9-15%)	0.042 (0.0033-0.1)	0.23% (-0.036-0.34%)	0.26% (-0.068-0.41%)	0.18 (-0.047-0.28)	2 (-0.31-3)
Brunei	2010	Male	85 to 89	91% (84-96%)	8.9% (4.4-16%)	0.039 (0.0029-0.093)	0.18% (-0.079-0.28%)	0.2% (-0.13-0.32%)	0.084 (-0.057-0.14)	0.69 (-0.31-1.1)
Brunei	2010	Male	90 to 94	91% (84-95%)	9.2% (4.5-16%)	0.036 (0.0025-0.09)	0.14% (-0.12-0.23%)	0.16% (-0.19-0.27%)	0.037 (-0.045-0.065)	0.21 (-0.18-0.34)
Brunei	2010	Male	95 plus	91% (81-96%)	9.4% (3.7-19%)	0.033 (0.0022-0.088)	0.14% (-0.11-0.22%)	0.14% (-0.2-0.26%)	0.01 (-0.014-0.018)	0.041 (-0.031-0.068)
Brunei	2016	Female	All Ages	76% (70-81%)	24% (19-30%)	0.069 (0.0075-0.15)	0.52% (0.22-0.7%)	0.56% (0.041-0.8%)	3.8 (0.3-6.1)	200 (83-290)
Brunei	2016	Female	15 to 19	70% (54-83%)	30% (17-46%)	0.057 (0.0057-0.13)	0.63% (0.41-0.92%)	1.1% (0.57-2%)	0.071 (0.031-0.13)	11 (6.8-17)
Brunei	2016	Female	20 to 24	80% (68-89%)	20% (11-32%)	0.1 (0.011-0.23)	1% (0.68-1.4%)	1.1% (0.58-2%)	0.086 (0.042-0.16)	21 (13-31)
Brunei	2016	Female	25 to 29	84% (74-91%)	16% (8.7-26%)	0.11 (0.011-0.24)	0.9% (0.61-1.2%)	1.1% (0.61-1.8%)	0.11 (0.052-0.19)	21 (14-31)
Brunei	2016	Female	30 to 34	86% (76-92%)	14% (7.5-24%)	0.11 (0.011-0.25)	0.76% (0.46-1.1%)	1.2% (0.68-1.9%)	0.16 (0.079-0.28)	21 (12-30)
Brunei	2016	Female	35 to 39	86% (77-93%)	14% (7.4-23%)	0.097 (0.011-0.23)	0.72% (0.44-1%)	1.3% (0.8-2%)	0.24 (0.13-0.41)	22 (13-32)
Brunei	2016	Female	40 to 44	85% (76-92%)	15% (7.9-24%)	0.092 (0.01-0.22)	0.69% (0.35-0.99%)	1.3% (0.82-2%)	0.35 (0.19-0.57)	23 (11-35)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	2016	Female	45 to 49	84% (72-92%)	16% (8.3-28%)	0.09 (0.0099-0.21)	0.66% (0.26-0.96%)	1.2% (0.71-1.8%)	0.43 (0.23-0.71)	23 (8.8-36)
Brunei	2016	Female	50 to 54	82% (70-90%)	18% (10-30%)	0.085 (0.0089-0.2)	0.61% (0.17-0.94%)	1% (0.51-1.5%)	0.47 (0.21-0.75)	20 (5.3-33)
Brunei	2016	Female	55 to 59	79% (65-89%)	21% (11-35%)	0.078 (0.0078-0.18)	0.5% (-0.077-0.88%)	0.86% (0.28-1.4%)	0.47 (0.15-0.83)	16 (-2.5-29)
Brunei	2016	Female	60 to 64	76% (60-88%)	24% (12-40%)	0.07 (0.0071-0.16)	0.33% (-0.4-0.75%)	0.55% (-0.21-1%)	0.35 (-0.12-0.69)	9.3 (-12-22)
Brunei	2016	Female	65 to 69	73% (57-86%)	27% (14-43%)	0.06 (0.0059-0.13)	0.32% (-0.39-0.71%)	0.5% (-0.32-0.97%)	0.32 (-0.21-0.64)	7.1 (-8.7-16)
Brunei	2016	Female	70 to 74	72% (56-83%)	28% (17-44%)	0.051 (0.0047-0.12)	0.32% (-0.41-0.68%)	0.45% (-0.38-0.89%)	0.26 (-0.22-0.52)	4.8 (-5.9-10)
Brunei	2016	Female	75 to 79	70% (56-83%)	30% (17-44%)	0.046 (0.0044-0.11)	0.25% (-0.48-0.56%)	0.33% (-0.53-0.73%)	0.22 (-0.36-0.5)	3.3 (-6.3-7.5)
Brunei	2016	Female	80 to 84	69% (51-82%)	31% (18-49%)	0.046 (0.0043-0.1)	0.17% (-0.51-0.48%)	0.2% (-0.6-0.58%)	0.15 (-0.45-0.43)	1.7 (-5.2-4.8)
Brunei	2016	Female	85 to 89	67% (50-80%)	33% (20-50%)	0.046 (0.0046-0.1)	0.11% (-0.55-0.43%)	0.14% (-0.62-0.53%)	0.084 (-0.39-0.33)	0.64 (-3.1-2.5)
Brunei	2016	Female	90 to 94	67% (52-80%)	33% (20-48%)	0.047 (0.0047-0.11)	0.089% (-0.57-0.38%)	0.11% (-0.65-0.47%)	0.035 (-0.2-0.15)	0.17 (-1.1-0.75)
Brunei	2016	Female	95 plus	68% (48-83%)	32% (17-52%)	0.048 (0.0047-0.12)	0.086% (-0.6-0.4%)	0.11% (-0.71-0.51%)	0.013 (-0.083-0.059)	0.041 (-0.29-0.2)
Brunei	2016	Male	All Ages	90% (87-93%)	9.8% (6.9-13%)	0.065 (0.007-0.14)	1.3% (1-1.6%)	1% (0.66-1.3%)	10 (6-15)	630 (460-850)
Brunei	2016	Male	15 to 19	84% (73-92%)	16% (8.4-27%)	0.043 (0.0046-0.1)	0.98% (0.62-1.5%)	0.097 (0.29-1.7%)	0.097 (0.038-0.27)	23 (14-38)
Brunei	2016	Male	20 to 24	91% (83-95%)	9.3% (4.8-17%)	0.08 (0.0088-0.18)	2.1% (1.3-3.2%)	0.95% (0.33-2.4%)	0.2 (0.059-0.56)	55 (33-91)
Brunei	2016	Male	25 to 29	93% (88-96%)	7% (3.6-12%)	0.096 (0.01-0.21)	2.3% (1.5-3.1%)	1.3% (0.53-2.5%)	0.29 (0.11-0.67)	66 (43-100)
Brunei	2016	Male	30 to 34	94% (90-97%)	5.6% (2.6-10%)	0.099 (0.0098-0.22)	2.1% (1.5-2.9%)	1.5% (0.87-2.5%)	0.41 (0.2-0.81)	69 (46-100)
Brunei	2016	Male	35 to 39	95% (91-98%)	5% (2.3-9.3%)	0.09 (0.0095-0.21)	2.1% (1.6-2.7%)	2% (1.3-2.9%)	0.67 (0.36-1.2)	71 (50-100)
Brunei	2016	Male	40 to 44	95% (91-98%)	4.7% (2-8.7%)	0.088 (0.0096-0.21)	2% (1.5-2.6%)	2.3% (1.6-3.2%)	0.99 (0.55-1.7)	73 (49-110)
Brunei	2016	Male	45 to 49	95% (91-98%)	4.6% (2-9.3%)	0.092 (0.0097-0.21)	1.8% (1.3-2.3%)	2.1% (1.4-3%)	1.2 (0.7-2)	71 (48-100)
Brunei	2016	Male	50 to 54	95% (91-98%)	4.9% (2.2-9.1%)	0.09 (0.009-0.21)	1.5% (1-2%)	1.9% (1.1-2.6%)	1.4 (0.78-2.3)	66 (41-97)
Brunei	2016	Male	55 to 59	95% (90-98%)	5.4% (2.4-10%)	0.082 (0.0085-0.19)	1.2% (0.73-1.7%)	1.5% (0.86-2.1%)	1.4 (0.77-2.2)	54 (32-82)
Brunei	2016	Male	60 to 64	94% (88-98%)	6% (2.4-12%)	0.077 (0.0074-0.18)	0.93% (0.49-1.3%)	1.1% (0.55-1.6%)	1.2 (0.58-1.9)	38 (19-57)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brunei	2016	Male	65 to 69	93% (87-97%)	6.5% (2.7-13%)	0.073 (0.0082-0.16)	0.71% (0.35-1%)	0.85% (0.4-1.2%)	0.85 (0.39-1.3)	21 (10-32)
Brunei	2016	Male	70 to 74	93% (87-97%)	7% (3.3-13%)	0.068 (0.0069-0.15)	0.48% (0.12-0.71%)	0.57% (0.12-0.86%)	0.51 (0.12-0.79)	10 (2.6-15)
Brunei	2016	Male	75 to 79	93% (87-96%)	7.4% (3.7-13%)	0.061 (0.0064-0.14)	0.33% (-0.022-0.51%)	0.38% (-0.063-0.6%)	0.34 (-0.059-0.56)	5.3 (-0.36-8.3)
Brunei	2016	Male	80 to 84	92% (85-96%)	8% (3.9-15%)	0.055 (0.0055-0.12)	0.2% (-0.12-0.35%)	0.23% (-0.17-0.41%)	0.19 (-0.15-0.35)	2.2 (-1.3-3.8)
Brunei	2016	Male	85 to 89	91% (84-96%)	8.9% (4.5-16%)	0.049 (0.005-0.11)	0.17% (-0.16-0.28%)	0.18% (-0.21-0.33%)	0.1 (-0.12-0.19)	0.88 (-0.83-1.5)
Brunei	2016	Male	90 to 94	91% (84-95%)	9.2% (4.5-16%)	0.044 (0.0044-0.1)	0.13% (-0.18-0.24%)	0.14% (-0.25-0.27%)	0.036 (-0.063-0.069)	0.21 (-0.29-0.37)
Brunei	2016	Male	95 plus	91% (82-96%)	9.3% (3.8-18%)	0.039 (0.0036-0.096)	0.13% (-0.13-0.24%)	0.13% (-0.23-0.26%)	0.0099 (-0.017-0.019)	0.044 (-0.042-0.074)
Japan	1990	Female	All Ages	79% (75-82%)	21% (18-25%)	1.9 (1.9-2)	0.99% (0.1-1.9%)	0.1% (-1.6-2%)	400 (-6100-7500)	130000 (13000-250000)
Japan	1990	Female	15 to 19	77% (68-85%)	23% (15-32%)	2 (1.3-2.9)	2% (1.1-3.1%)	8.7% (4.6-14%)	98 (52-150)	9100 (5000-14000)
Japan	1990	Female	20 to 24	87% (80-92%)	13% (8.2-20%)	3.2 (2.1-4.4)	3.3% (1.8-4.9%)	13% (6.8-20%)	180 (91-260)	15000 (9000-22000)
Japan	1990	Female	25 to 29	90% (85-94%)	9.9% (6.3-15%)	3.2 (2.2-4.3)	2.9% (1.6-4.3%)	12% (6.2-18%)	170 (86-250)	14000 (7300-20000)
Japan	1990	Female	30 to 34	92% (86-95%)	8.4% (4.9-14%)	3 (2.1-4.2)	2.5% (1.3-3.8%)	10% (5.5-15%)	180 (98-270)	13000 (6900-20000)
Japan	1990	Female	35 to 39	91% (86-95%)	8.9% (5-14%)	2.8 (2-3.8)	2.5% (1.3-3.9%)	9.3% (5.5-13%)	290 (170-400)	17000 (8600-25000)
Japan	1990	Female	40 to 44	89% (84-93%)	11% (6.6-16%)	2.7 (1.9-3.6)	2.7% (1.2-4.3%)	8.9% (5-12%)	500 (280-700)	24000 (12000-37000)
Japan	1990	Female	45 to 49	87% (79-92%)	13% (7.9-21%)	2.5 (1.7-3.4)	2.7% (1.1-4.5%)	7.9% (4.3-11%)	590 (330-860)	24000 (11000-38000)
Japan	1990	Female	50 to 54	82% (75-88%)	18% (12-25%)	2.3 (1.5-3.1)	2.6% (0.92-4.6%)	6.7% (3.5-10%)	700 (360-1000)	24000 (8600-41000)
Japan	1990	Female	55 to 59	78% (68-86%)	22% (14-32%)	2.2 (1.5-2.9)	2.3% (0.73-4.1%)	5.4% (2.8-8.3%)	800 (410-1200)	23000 (7400-41000)
Japan	1990	Female	60 to 64	73% (62-81%)	27% (19-38%)	1.9 (1.2-2.7)	1.5% (-0.14-3.5%)	3.4% (0.94-6.3%)	700 (190-1300)	16000 (-1500-36000)
Japan	1990	Female	65 to 69	68% (58-78%)	32% (22-42%)	1.5 (1-2)	0.4% (-0.92-1.9%)	1.4% (-0.52-3.5%)	390 (-150-970)	4300 (-9900-21000)
Japan	1990	Female	70 to 74	67% (57-76%)	33% (24-43%)	1.2 (0.74-1.7)	-0.58% (-1.9-0.89%)	-0.27% (-2.1-1.7%)	-100 (-810-650)	-6400 (-21000-9800)
Japan	1990	Female	75 to 79	70% (60-79%)	30% (21-40%)	1.1 (0.68-1.6)	-1.3% (-2.8-0.3%)	-1.5% (-3.4-0.6%)	-860 (-2000-360)	-16000 (-33000-3700)
Japan	1990	Female	80 to 84	72% (60-81%)	28% (19-40%)	1.2 (0.82-1.7)	-1.7% (-3.4-0.2%)	-1.9% (-4.1-0.27%)	-1400 (-3000-190)	-17000 (-34000-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	1990	Female	85 to 89	71% (60-79%)	29% (21-40%)	1.4 (0.89-1.9)	-1.5% (-3.5-0.81%)	-1.8% (-4.2-0.9%)	-1200 (-2800-590)	-9100 (-21000-4700)
Japan	1990	Female	90 to 94	71% (61-80%)	29% (20-39%)	1.5 (0.94-2.2)	-1.2% (-3.3-1.5%)	-1.5% (-4.1-1.6%)	-520 (-1400-560)	-2600 (-7300-3100)
Japan	1990	Female	95 plus	71% (58-82%)	29% (18-42%)	1.6 (0.91-2.5)	-0.73% (-3.3-2.5%)	-1% (-4.1-2.7%)	-110 (-440-300)	-320 (-1500-1100)
Japan	1990	Male	All Ages	89% (86-91%)	11% (9.1-14%)	1.8 (1.8-1.9)	3.5% (2.3-4.6%)	2.8% (1.1-4.4%)	12000 (4900-20000)	530000 (350000-700000)
Japan	1990	Male	15 to 19	80% (70-88%)	20% (12-30%)	1.2 (0.7-1.7)	3.7% (1.6-5.8%)	7.4% (2.6-12%)	240 (84-390)	21000 (9500-32000)
Japan	1990	Male	20 to 24	89% (81-94%)	11% (6.5-19%)	2.2 (1.5-3)	7.1% (4.3-10%)	12% (6.4-19%)	430 (220-650)	37000 (22000-53000)
Japan	1990	Male	25 to 29	92% (86-95%)	8.4% (4.8-14%)	2.7 (1.9-3.7)	7.1% (4.3-9.9%)	13% (6.8-20%)	400 (200-580)	33000 (20000-47000)
Japan	1990	Male	30 to 34	93% (88-96%)	6.7% (3.5-12%)	2.8 (2-3.9)	6.1% (3.7-8.7%)	12% (6.2-18%)	390 (200-580)	31000 (19000-44000)
Japan	1990	Male	35 to 39	94% (91-97%)	5.6% (2.9-9.5%)	2.6 (1.8-3.5)	5.5% (3.2-7.6%)	10% (5.3-15%)	540 (290-800)	37000 (22000-52000)
Japan	1990	Male	40 to 44	95% (91-97%)	5.2% (2.7-9%)	2.5 (1.7-3.4)	5.4% (3.2-7.7%)	9.3% (5-13%)	940 (510-1300)	52000 (31000-73000)
Japan	1990	Male	45 to 49	94% (90-97%)	5.5% (2.9-10%)	2.6 (1.8-3.6)	5.9% (3.7-8.4%)	9.2% (5.5-13%)	1300 (780-1800)	61000 (38000-85000)
Japan	1990	Male	50 to 54	93% (89-96%)	6.5% (3.6-11%)	2.5 (1.7-3.4)	5.6% (3.4-8%)	8.2% (5-12%)	1700 (1000-2400)	67000 (41000-96000)
Japan	1990	Male	55 to 59	92% (86-96%)	8.1% (4.4-14%)	2.3 (1.5-3.1)	4.6% (2.9-6.7%)	6.3% (3.9-9.1%)	2100 (1300-3000)	70000 (44000-100000)
Japan	1990	Male	60 to 64	90% (84-94%)	9.9% (5.5-16%)	2.1 (1.4-3)	3.4% (1.7-5.5%)	4.5% (2.4-7.1%)	1900 (1000-3000)	55000 (28000-89000)
Japan	1990	Male	65 to 69	88% (82-93%)	12% (6.8-18%)	2 (1.4-2.6)	2.5% (0.98-4.1%)	3.1% (1.3-5.2%)	1400 (570-2300)	33000 (13000-55000)
Japan	1990	Male	70 to 74	88% (80-93%)	12% (7.3-20%)	1.8 (1.2-2.5)	1.7% (0.074-3.6%)	2% (0.032-4.2%)	1000 (16-2200)	20000 (900-43000)
Japan	1990	Male	75 to 79	88% (82-93%)	12% (6.5-18%)	1.7 (1-2.4)	0.92% (-0.74-2.9%)	0.97% (-1-3.2%)	680 (-710-2200)	11000 (-9300-36000)
Japan	1990	Male	80 to 84	89% (81-94%)	11% (6.2-19%)	1.4 (0.95-2)	0.17% (-1.4-1.9%)	0.032% (-1.9-2%)	22 (-1300-1400)	1500 (-13000-16000)
Japan	1990	Male	85 to 89	88% (80-93%)	12% (6.9-20%)	1.2 (0.82-1.8)	-0.37% (-2-1.3%)	-0.67% (-2.5-1.1%)	-310 (-1200-520)	-1500 (-8000-5000)
Japan	1990	Male	90 to 94	88% (81-93%)	12% (6.9-19%)	1.1 (0.68-1.6)	-0.76% (-2.3-0.8%)	-1.2% (-3.1-0.6%)	-220 (-550-110)	-830 (-2500-890)
Japan	1990	Male	95 plus	89% (80-95%)	11% (5.5-20%)	0.95 (0.51-1.6)	-1.1% (-2.6-0.52%)	-1.7% (-3.8-0.17%)	-71 (-150-6.8)	-180 (-440-88)
Japan	1995	Female	All Ages	78% (75-82%)	22% (18-25%)	2 (2-2.1)	1.2% (0.28-2%)	0.5% (-1.1-2.3%)	2100 (-4700-9600)	160000 (39000-280000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	1995	Female	15 to 19	77% (67-85%)	23% (15-33%)	2 (1.3-2.9)	1.9% (1-2.9%)	8.1% (4.1-13%)	79 (40-120)	7300 (4000-11000)
Japan	1995	Female	20 to 24	87% (79-92%)	13% (8.3-21%)	3.2 (2.1-4.4)	3.1% (1.8-4.7%)	13% (6.8-19%)	190 (100-280)	17000 (9800-24000)
Japan	1995	Female	25 to 29	90% (84-94%)	10% (6.4-16%)	3.3 (2.3-4.5)	2.9% (1.6-4.3%)	12% (6.8-18%)	180 (99-270)	15000 (8400-21000)
Japan	1995	Female	30 to 34	91% (86-95%)	8.6% (5.1-14%)	3.2 (2.2-4.4)	2.7% (1.4-4.1%)	11% (6.2-16%)	200 (110-290)	14000 (7700-21000)
Japan	1995	Female	35 to 39	91% (86-95%)	8.9% (5.2-14%)	2.9 (2-3.9)	2.4% (1.2-3.8%)	9.5% (5.6-13%)	240 (140-330)	14000 (6800-21000)
Japan	1995	Female	40 to 44	89% (84-93%)	11% (6.5-16%)	2.7 (1.9-3.7)	2.6% (1.1-4.2%)	8.9% (5.1-12%)	410 (230-570)	20000 (8700-31000)
Japan	1995	Female	45 to 49	87% (79-92%)	13% (7.9-21%)	2.6 (1.8-3.6)	2.9% (1.3-4.6%)	8.3% (4.9-12%)	710 (410-1000)	29000 (13000-45000)
Japan	1995	Female	50 to 54	82% (75-88%)	18% (12-25%)	2.4 (1.6-3.3)	2.8% (1.1-4.8%)	7.4% (4-11%)	800 (440-1200)	28000 (11000-46000)
Japan	1995	Female	55 to 59	78% (68-86%)	22% (14-32%)	2.2 (1.5-3)	2.2% (0.72-3.9%)	5.6% (3-8.3%)	790 (420-1200)	22000 (7200-39000)
Japan	1995	Female	60 to 64	73% (62-81%)	27% (19-38%)	1.9 (1.2-2.8)	1.6% (-0.028-3.5%)	3.8% (1.3-6.6%)	790 (280-1400)	17000 (-320-39000)
Japan	1995	Female	65 to 69	69% (57-78%)	31% (22-43%)	1.6 (1.1-2.2)	0.81% (-0.49-2.3%)	2.2% (0.29-4.2%)	640 (85-1200)	9700 (-6100-27000)
Japan	1995	Female	70 to 74	68% (57-77%)	32% (23-43%)	1.4 (0.94-2)	0.15% (-1.2-1.8%)	0.87% (-1-3%)	360 (-420-1200)	1900 (-15000-22000)
Japan	1995	Female	75 to 79	71% (60-79%)	29% (21-40%)	1.4 (0.84-2)	-0.51% (-2-1.3%)	-0.32% (-2.3-2%)	-180 (-1300-1100)	-6000 (-24000-16000)
Japan	1995	Female	80 to 84	71% (59-81%)	29% (19-41%)	1.4 (0.9-1.9)	-1% (-2.6-0.71%)	-1.1% (-3.2-0.99%)	-910 (-2500-790)	-12000 (-30000-8000)
Japan	1995	Female	85 to 89	70% (59-79%)	30% (21-41%)	1.4 (0.91-1.9)	-1.2% (-2.9-0.68%)	-1.5% (-3.6-0.8%)	-1100 (-2800-620)	-9300 (-22000-5000)
Japan	1995	Female	90 to 94	71% (60-79%)	29% (21-40%)	1.4 (0.87-2)	-1.3% (-3.1-0.78%)	-1.6% (-3.7-0.89%)	-770 (-1800-410)	-4000 (-9200-2400)
Japan	1995	Female	95 plus	71% (58-82%)	29% (18-42%)	1.4 (0.75-2.2)	-1.3% (-3.4-1.3%)	-1.7% (-4.1-1.4%)	-300 (-730-240)	-950 (-2400-950)
Japan	1995	Male	All Ages	89% (87-91%)	11% (9-13%)	1.9 (1.8-2)	3.4% (2.3-4.5%)	2.8% (1.3-4.3%)	14000 (6300-22000)	56000 (37000-73000)
Japan	1995	Male	15 to 19	80% (69-88%)	20% (12-31%)	1.2 (0.7-1.7)	3.3% (1.5-5.1%)	6.7% (2.4-11%)	170 (62-280)	15000 (7600-24000)
Japan	1995	Male	20 to 24	88% (81-94%)	12% (6.5-19%)	2.3 (1.5-3.1)	6.5% (4-9.4%)	11% (5.7-18%)	430 (220-670)	38000 (23000-55000)
Japan	1995	Male	25 to 29	91% (86-95%)	8.5% (4.8-14%)	2.7 (1.9-3.7)	6.8% (4-9.5%)	13% (6.2-19%)	420 (210-620)	36000 (21000-51000)
Japan	1995	Male	30 to 34	93% (87-97%)	6.8% (3.4-13%)	2.9 (2-3.9)	6% (3.6-8.6%)	11% (5.9-18%)	400 (200-610)	31000 (19000-45000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	1995	Male	35 to 39	94% (90-97%)	5.7% (3.1-9.6%)	2.6 (1.8-3.6)	5.4% (3.2-7.7%)	10% (5.1-15%)	460 (240-690)	31000 (18000-44000)
Japan	1995	Male	40 to 44	95% (91-97%)	5.3% (2.8-9.1%)	2.6 (1.8-3.5)	5.3% (3.1-7.6%)	9.2% (5-13%)	770 (420-1100)	43000 (26000-61000)
Japan	1995	Male	45 to 49	94% (90-97%)	5.6% (2.8-10%)	2.6 (1.8-3.6)	5.7% (3.4-8.2%)	9.1% (5.2-13%)	1500 (830-2000)	68000 (41000-95000)
Japan	1995	Male	50 to 54	93% (89-96%)	6.5% (3.6-11%)	2.5 (1.7-3.4)	5.5% (3.3-7.8%)	8.2% (4.9-12%)	1800 (1100-2500)	72000 (43000-100000)
Japan	1995	Male	55 to 59	92% (87-96%)	8.1% (4.4-13%)	2.3 (1.6-3.2)	4.6% (2.9-6.7%)	6.5% (4.1-9.4%)	2000 (1300-2800)	67000 (43000-97000)
Japan	1995	Male	60 to 64	90% (84-94%)	9.9% (5.6-16%)	2.1 (1.4-3)	3.6% (2-5.7%)	4.9% (2.8-7.5%)	2300 (1300-3500)	65000 (35000-100000)
Japan	1995	Male	65 to 69	88% (81-93%)	12% (6.7-19%)	2 (1.4-2.6)	2.7% (1.3-4.2%)	3.5% (1.8-5.3%)	2100 (1100-3200)	49000 (24000-77000)
Japan	1995	Male	70 to 74	87% (80-93%)	13% (7.5-20%)	1.8 (1.3-2.5)	1.8% (0.36-3.5%)	2.3% (0.49-4.3%)	1400 (300-2600)	27000 (5100-51000)
Japan	1995	Male	75 to 79	88% (81-93%)	12% (6.7-19%)	1.7 (1.1-2.4)	1.1% (-0.45-2.9%)	1.2% (-0.61-3.3%)	810 (-410-2200)	13000 (-5600-35000)
Japan	1995	Male	80 to 84	88% (81-94%)	12% (6.4-19%)	1.5 (0.97-2)	0.37% (-1.1-1.9%)	0.29% (-1.4-2.1%)	220 (-1100-1600)	3600 (-11000-19000)
Japan	1995	Male	85 to 89	88% (80-93%)	12% (7-20%)	1.3 (0.84-1.8)	-0.12% (-1.6-1.4%)	-0.36% (-2.1-1.4%)	-200 (-1100-750)	-600 (-7800-7000)
Japan	1995	Male	90 to 94	88% (81-93%)	12% (6.8-19%)	1.1 (0.72-1.7)	-0.49% (-1.9-1%)	-0.9% (-2.7-0.94%)	-220 (-650-230)	-740 (-2900-1600)
Japan	1995	Male	95 plus	89% (81-95%)	11% (5.3-19%)	1 (0.55-1.6)	-0.76% (-2.2-0.82%)	-1.4% (-3.2-0.59%)	-86 (-200-38)	-200 (-590-220)
Japan	2000	Female	All Ages	78% (75-81%)	22% (19-25%)	2 (1.9-2.1)	1.2% (0.35-2%)	0.59% (-0.84-2.2%)	2600 (-3800-9700)	160000 (50000-280000)
Japan	2000	Female	15 to 19	78% (69-86%)	22% (14-31%)	2 (1.3-2.9)	1.8% (0.95-2.9%)	9.2% (4.4-14%)	65 (31-100)	6100 (3200-9400)
Japan	2000	Female	20 to 24	87% (80-92%)	13% (7.7-20%)	3.2 (2.1-4.4)	3.3% (1.8-5%)	15% (7.4-22%)	170 (85-260)	15000 (8300-21000)
Japan	2000	Female	25 to 29	91% (86-94%)	9.5% (5.9-14%)	3.3 (2.3-4.4)	3.1% (1.6-4.6%)	14% (7.1-21%)	220 (110-320)	17000 (9100-25000)
Japan	2000	Female	30 to 34	92% (87-95%)	8.1% (4.9-13%)	3.1 (2.2-4.3)	2.8% (1.4-4.2%)	12% (6.6-18%)	230 (120-340)	16000 (8200-24000)
Japan	2000	Female	35 to 39	91% (87-95%)	8.6% (4.8-13%)	2.9 (2-3.9)	2.6% (1.2-4%)	10% (6.2-15%)	250 (150-360)	15000 (7100-22000)
Japan	2000	Female	40 to 44	90% (85-94%)	10% (6.4-15%)	2.7 (1.8-3.8)	2.6% (1.1-4.2%)	9.6% (5.6-13%)	350 (200-490)	17000 (7400-26000)
Japan	2000	Female	45 to 49	87% (79-92%)	13% (7.8-21%)	2.6 (1.8-3.5)	2.8% (1.2-4.6%)	8.7% (5.3-12%)	580 (350-820)	24000 (11000-37000)
Japan	2000	Female	50 to 54	82% (75-88%)	18% (12-25%)	2.4 (1.6-3.2)	2.9% (1.1-4.8%)	7.6% (4.3-11%)	910 (520-1300)	31000 (13000-50000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	2000	Female	55 to 59	77% (67-85%)	23% (15-33%)	2.1 (1.5-2.9)	2.3% (0.84-4%)	6% (3.5-8.6%)	860 (500-1200)	24000 (9400-41000)
Japan	2000	Female	60 to 64	71% (60-81%)	29% (19-40%)	1.8 (1.2-2.6)	1.6% (0.11-3.3%)	4% (1.7-6.6%)	760 (320-1200)	17000 (1200-36000)
Japan	2000	Female	65 to 69	66% (54-77%)	34% (23-46%)	1.5 (1-2)	0.8% (-0.3-2.1%)	2.3% (0.68-4%)	660 (190-1200)	9800 (-4000-25000)
Japan	2000	Female	70 to 74	66% (55-75%)	34% (25-45%)	1.3 (0.85-1.8)	0.15% (-1-1.4%)	1% (-0.55-2.7%)	420 (-220-1100)	2000 (-14000-19000)
Japan	2000	Female	75 to 79	70% (60-78%)	30% (22-40%)	1.3 (0.77-1.8)	-0.44% (-1.7-1%)	-0.12% (-1.8-1.8%)	-68 (-1000-1000)	-5800 (-22000-13000)
Japan	2000	Female	80 to 84	71% (59-81%)	29% (19-41%)	1.3 (0.88-1.8)	-0.87% (-2.2-0.57%)	-0.9% (-2.6-0.87%)	-680 (-2000-660)	-10000 (-25000-6600)
Japan	2000	Female	85 to 89	70% (60-79%)	30% (21-40%)	1.3 (0.89-1.9)	-0.94% (-2.4-0.66%)	-1.1% (-2.8-0.84%)	-980 (-2500-750)	-8600 (-21000-6000)
Japan	2000	Female	90 to 94	71% (61-79%)	29% (21-39%)	1.4 (0.88-2)	-0.96% (-2.5-0.88%)	-1.2% (-3.1-0.95%)	-760 (-1900-600)	-4100 (-11000-3800)
Japan	2000	Female	95 plus	71% (57-81%)	29% (19-43%)	1.4 (0.79-2.3)	-0.88% (-2.8-1.5%)	-1.2% (-3.4-1.7%)	-340 (-970-480)	-1000 (-3300-1800)
Japan	2000	Male	All Ages	91% (89-92%)	9.5% (7.8-11%)	1.9 (1.8-2)	3.4% (2.1-4.6%)	2.9% (1.3-4.3%)	15000 (7200-23000)	570000 (360000-760000)
Japan	2000	Male	15 to 19	83% (75-90%)	17% (10-25%)	1.2 (0.73-1.7)	3.2% (1.5-5%)	7.5% (2.8-12%)	130 (48-220)	12000 (5900-19000)
Japan	2000	Male	20 to 24	91% (85-95%)	9% (5.3-15%)	2.3 (1.6-3.1)	6.8% (4-9.8%)	13% (5.9-20%)	360 (170-570)	32000 (19000-47000)
Japan	2000	Male	25 to 29	93% (88-96%)	7% (4-12%)	2.7 (1.9-3.7)	7% (4-10%)	14% (6-21%)	470 (200-720)	39000 (22000-57000)
Japan	2000	Male	30 to 34	94% (90-97%)	5.7% (3-10%)	2.8 (2-3.8)	6.3% (3.5-9.4%)	13% (5.4-20%)	470 (200-740)	36000 (20000-53000)
Japan	2000	Male	35 to 39	95% (92-97%)	4.8% (2.6-8%)	2.6 (1.8-3.6)	5.6% (3-8.3%)	11% (4.8-17%)	510 (230-780)	33000 (17000-49000)
Japan	2000	Male	40 to 44	96% (92-98%)	4.5% (2.4-7.7%)	2.5 (1.7-3.5)	5.4% (2.8-8%)	9.8% (4.7-15%)	680 (330-1000)	38000 (20000-55000)
Japan	2000	Male	45 to 49	95% (92-98%)	4.7% (2.5-8.2%)	2.6 (1.8-3.6)	5.7% (3.1-8.4%)	9.4% (4.8-13%)	1200 (630-1800)	58000 (32000-82000)
Japan	2000	Male	50 to 54	94% (91-97%)	5.6% (3.2-9.2%)	2.5 (1.7-3.4)	5.4% (3-7.9%)	8.2% (4.4-12%)	2000 (1100-3000)	80000 (44000-120000)
Japan	2000	Male	55 to 59	93% (88-96%)	7.1% (4-12%)	2.3 (1.6-3.1)	4.6% (2.6-6.8%)	6.6% (3.9-9.5%)	2100 (1300-3000)	71000 (42000-110000)
Japan	2000	Male	60 to 64	91% (86-95%)	8.8% (5-14%)	2.1 (1.4-3)	3.6% (1.9-5.8%)	5.1% (2.9-7.9%)	2200 (1200-3300)	61000 (33000-97000)
Japan	2000	Male	65 to 69	89% (82-94%)	11% (6.4-18%)	2 (1.4-2.6)	2.8% (1.4-4.3%)	3.7% (2-5.6%)	2300 (1200-3400)	53000 (27000-83000)
Japan	2000	Male	70 to 74	88% (81-93%)	12% (7.1-19%)	1.8 (1.2-2.5)	1.9% (0.55-3.5%)	2.5% (0.8-4.3%)	1900 (620-3400)	37000 (10000-66000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	2000	Male	75 to 79	88% (81-93%)	12% (6.9-19%)	1.6 (1-2.3)	1.2% (-0.22-2.7%)	1.4% (-0.27-3.2%)	1000 (-210-2500)	16000 (-3200-39000)
Japan	2000	Male	80 to 84	88% (80-94%)	12% (6.4-20%)	1.4 (0.94-1.9)	0.48% (-0.78-1.9%)	0.46% (-1.1-2.1%)	340 (-790-1600)	4700 (-7800-18000)
Japan	2000	Male	85 to 89	87% (80-93%)	13% (7.1-20%)	1.2 (0.82-1.7)	0.039% (-1.2-1.3%)	-0.17% (-1.7-1.4%)	-110 (-1100-870)	230 (-7100-8000)
Japan	2000	Male	90 to 94	88% (80-93%)	12% (7.3-20%)	1.1 (0.69-1.6)	-0.28% (-1.5-1%)	-0.66% (-2.2-0.92%)	-210 (-710-300)	-580 (-3100-2100)
Japan	2000	Male	95 plus	89% (80-94%)	11% (5.6-20%)	0.96 (0.53-1.6)	-0.5% (-1.8-0.87%)	-1% (-2.7-0.66%)	-96 (-250-60)	-190 (-680-330)
Japan	2005	Female	All Ages	78% (74-81%)	22% (19-26%)	1.9 (1.8-2)	1% (0.26-1.8%)	0.5% (-0.82-2%)	2500 (-4200-10000)	150000 (38000-260000)
Japan	2005	Female	15 to 19	78% (69-86%)	22% (14-31%)	2 (1.3-2.8)	1.8% (0.84-2.8%)	9.3% (3.9-15%)	55 (23-88)	5100 (2600-8100)
Japan	2005	Female	20 to 24	87% (80-92%)	13% (8-20%)	3.1 (2.1-4.4)	3.3% (1.7-5.3%)	15% (6.8-24%)	160 (69-240)	13000 (6900-20000)
Japan	2005	Female	25 to 29	90% (85-94%)	9.7% (6.2-15%)	3.1 (2.2-4.2)	3% (1.5-4.6%)	15% (6.5-22%)	190 (85-290)	15000 (7300-22000)
Japan	2005	Female	30 to 34	92% (87-95%)	8.3% (4.9-13%)	3 (2.1-4.1)	2.7% (1.3-4.1%)	12% (6-19%)	250 (120-380)	17000 (8300-26000)
Japan	2005	Female	35 to 39	91% (86-95%)	8.7% (5.1-14%)	2.8 (1.9-3.8)	2.6% (1.2-4%)	11% (6-15%)	280 (160-400)	16000 (7500-24000)
Japan	2005	Female	40 to 44	90% (85-93%)	10% (6.5-15%)	2.6 (1.8-3.6)	2.6% (1.1-4.2%)	9.9% (5.5-14%)	340 (190-480)	17000 (7300-26000)
Japan	2005	Female	45 to 49	87% (79-92%)	13% (7.8-21%)	2.5 (1.7-3.4)	2.7% (1.2-4.3%)	8.8% (5.2-12%)	460 (270-650)	19000 (8700-30000)
Japan	2005	Female	50 to 54	82% (74-88%)	18% (12-26%)	2.3 (1.5-3.1)	2.7% (1.1-4.5%)	7.6% (4.5-11%)	720 (420-1000)	25000 (10000-39000)
Japan	2005	Female	55 to 59	77% (67-86%)	23% (14-33%)	2.1 (1.5-2.8)	2.3% (0.92-3.8%)	6.1% (3.8-8.6%)	950 (580-1300)	27000 (11000-44000)
Japan	2005	Female	60 to 64	71% (60-81%)	29% (19-40%)	1.8 (1.2-2.5)	1.5% (0.17-3%)	4.1% (2-6.4%)	770 (370-1200)	17000 (2000-34000)
Japan	2005	Female	65 to 69	66% (54-77%)	34% (23-46%)	1.4 (0.99-1.9)	0.66% (-0.27-1.8%)	2.3% (0.9-3.9%)	600 (230-1000)	7800 (-3700-21000)
Japan	2005	Female	70 to 74	66% (55-76%)	34% (24-45%)	1.2 (0.77-1.6)	0.042% (-0.94-1.1%)	1% (-0.32-2.5%)	410 (-130-1000)	510 (-13000-15000)
Japan	2005	Female	75 to 79	70% (60-78%)	30% (22-40%)	1.1 (0.69-1.6)	-0.48% (-1.5-0.71%)	-0.042% (-1.4-1.5%)	-26 (-860-890)	-6900 (-21000-9900)
Japan	2005	Female	80 to 84	71% (59-81%)	29% (19-41%)	1.2 (0.82-1.7)	-0.85% (-2-0.37%)	-0.81% (-2.3-0.78%)	-700 (-2000-670)	-12000 (-27000-5300)
Japan	2005	Female	85 to 89	70% (59-79%)	30% (21-41%)	1.3 (0.87-1.8)	-0.76% (-2-0.61%)	-0.83% (-2.4-0.86%)	-800 (-2300-820)	-7700 (-20000-6300)
Japan	2005	Female	90 to 94	71% (60-79%)	29% (21-40%)	1.4 (0.89-2)	-0.7% (-2.1-1%)	-0.88% (-2.6-1.2%)	-750 (-2300-1000)	-4200 (-13000-5800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	2005	Female	95 plus	71% (57-82%)	29% (18-43%)	1.5 (0.84-2.3)	-0.54% (-2.4-1.8%)	-0.78% (-3-2.1%)	-380 (-1500-1000)	-1100 (-5000-3700)
Japan	2005	Male	All Ages	91% (89-92%)	9.4% (7.7-11%)	1.8 (1.8-1.9)	3% (1.7-4.2%)	2.5% (1-3.9%)	14000 (6000-23000)	520000 (300000-710000)
Japan	2005	Male	15 to 19	83% (75-90%)	17% (10-25%)	1.1 (0.68-1.7)	2.8% (1.3-4.3%)	7% (2.3-12%)	88 (29-150)	8700 (4200-13000)
Japan	2005	Male	20 to 24	91% (85-95%)	9.1% (5.3-15%)	2.2 (1.5-3)	6.3% (3.4-9.5%)	12% (5.1-20%)	290 (120-460)	26000 (14000-38000)
Japan	2005	Male	25 to 29	93% (89-96%)	7% (4.1-11%)	2.6 (1.8-3.5)	6.7% (3.5-9.8%)	14% (5.4-22%)	390 (150-620)	33000 (17000-48000)
Japan	2005	Male	30 to 34	94% (90-97%)	5.8% (3-10%)	2.7 (1.9-3.6)	6% (3-8.9%)	12% (4.3-20%)	490 (170-790)	37000 (20000-56000)
Japan	2005	Male	35 to 39	95% (92-97%)	4.8% (2.6-8%)	2.5 (1.7-3.4)	5.3% (2.7-8.1%)	11% (4.1-17%)	520 (200-840)	34000 (17000-52000)
Japan	2005	Male	40 to 44	96% (92-98%)	4.5% (2.4-7.6%)	2.4 (1.7-3.3)	5.1% (2.4-7.7%)	9.5% (3.9-15%)	650 (270-1000)	36000 (17000-54000)
Japan	2005	Male	45 to 49	95% (92-98%)	4.7% (2.5-8.3%)	2.5 (1.7-3.4)	5.4% (2.7-8.2%)	9.1% (4.3-14%)	970 (450-1400)	45000 (23000-65000)
Japan	2005	Male	50 to 54	94% (91-97%)	5.6% (3.2-9.1%)	2.4 (1.6-3.2)	5% (2.6-7.4%)	7.8% (3.9-12%)	1500 (760-2200)	60000 (31000-88000)
Japan	2005	Male	55 to 59	93% (89-96%)	7.1% (4-11%)	2.2 (1.6-2.9)	4.2% (2.3-6.2%)	6.2% (3.4-9.1%)	2100 (1200-3200)	72000 (40000-110000)
Japan	2005	Male	60 to 64	91% (86-95%)	8.7% (4.9-14%)	2 (1.3-2.8)	3.4% (1.7-5.3%)	4.8% (2.6-7.4%)	2100 (1100-3200)	59000 (31000-92000)
Japan	2005	Male	65 to 69	89% (83-94%)	11% (6.2-17%)	1.8 (1.3-2.4)	2.5% (1.3-3.9%)	3.6% (1.9-5.4%)	2000 (1100-3000)	46000 (24000-71000)
Japan	2005	Male	70 to 74	88% (81-93%)	12% (7.2-19%)	1.7 (1.1-2.3)	1.8% (0.58-3.2%)	2.4% (0.89-4.1%)	1900 (710-3300)	36000 (12000-63000)
Japan	2005	Male	75 to 79	88% (81-93%)	12% (6.9-19%)	1.5 (0.94-2.1)	1% (-0.15-2.4%)	1.3% (-0.18-2.9%)	1300 (-180-2900)	19000 (-2800-45000)
Japan	2005	Male	80 to 84	88% (80-93%)	12% (6.6-20%)	1.3 (0.88-1.8)	0.46% (-0.84-1.6%)	0.49% (-0.88-1.9%)	440 (-800-1800)	5600 (-7900-20000)
Japan	2005	Male	85 to 89	87% (80-93%)	13% (7.2-20%)	1.2 (0.8-1.6)	0.09% (-1-1.3%)	-0.06% (-1.4-1.4%)	-42 (-980-940)	590 (-6800-8300)
Japan	2005	Male	90 to 94	88% (81-93%)	12% (7-19%)	1.1 (0.7-1.6)	-0.18% (-1.3-1%)	-0.48% (-1.9-1%)	-210 (-810-440)	-480 (-3700-2900)
Japan	2005	Male	95 plus	89% (80-95%)	11% (5.4-20%)	1 (0.56-1.6)	-0.34% (-1.6-1%)	-0.81% (-2.4-0.92%)	-120 (-350-130)	-210 (-970-630)
Japan	2010	Female	All Ages	77% (74-81%)	23% (19-26%)	1.8 (1.7-2)	0.86% (0.16-1.6%)	0.37% (-0.87-1.8%)	2100 (-5000-10000)	130000 (24000-240000)
Japan	2010	Female	15 to 19	77% (67-84%)	23% (16-33%)	1.9 (1.2-2.7)	1.6% (0.74-2.7%)	9% (3.3-15%)	45 (17-74)	4300 (2000-6800)
Japan	2010	Female	20 to 24	87% (79-92%)	13% (8.4-21%)	3 (2-4.3)	3.2% (1.5-5.2%)	15% (6.4-24%)	130 (54-200)	11000 (5400-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	2010	Female	25 to 29	90% (85-94%)	10% (6.2-15%)	3.1 (2.2-4.2)	3% (1.4-4.7%)	15% (6.3-22%)	160 (70-250)	13000 (6000-19000)
Japan	2010	Female	30 to 34	91% (86-95%)	8.6% (4.8-14%)	3 (2.1-4.1)	2.6% (1.2-4.1%)	13% (6.1-19%)	210 (100-320)	14000 (6900-22000)
Japan	2010	Female	35 to 39	91% (86-95%)	8.9% (5.1-14%)	2.7 (1.9-3.7)	2.4% (1.1-3.9%)	11% (5.8-16%)	300 (160-430)	17000 (7900-26000)
Japan	2010	Female	40 to 44	89% (84-94%)	11% (6.5-16%)	2.6 (1.7-3.5)	2.4% (0.94-4%)	9.9% (5.4-14%)	360 (200-510)	17000 (7100-27000)
Japan	2010	Female	45 to 49	87% (79-92%)	13% (7.9-21%)	2.4 (1.6-3.4)	2.6% (1.1-4.3%)	8.9% (5.2-13%)	450 (270-640)	18000 (8400-29000)
Japan	2010	Female	50 to 54	82% (75-88%)	18% (12-25%)	2.2 (1.5-3.1)	2.5% (0.98-4.3%)	7.7% (4.6-11%)	570 (340-820)	20000 (8000-32000)
Japan	2010	Female	55 to 59	77% (68-86%)	23% (14-32%)	2 (1.4-2.7)	2.1% (0.75-3.6%)	6.1% (3.8-8.7%)	750 (460-1100)	21000 (8200-35000)
Japan	2010	Female	60 to 64	73% (63-81%)	27% (19-37%)	1.8 (1.2-2.5)	1.4% (0.11-2.9%)	4.2% (2.2-6.5%)	850 (440-1300)	18000 (1500-36000)
Japan	2010	Female	65 to 69	68% (58-78%)	32% (22-42%)	1.5 (1.1-2)	0.65% (-0.28-1.7%)	2.6% (1.2-4.1%)	660 (300-1000)	8200 (-3600-21000)
Japan	2010	Female	70 to 74	68% (57-76%)	32% (24-43%)	1.3 (0.84-1.7)	0.15% (-0.8-1.2%)	1.4% (0.094-2.8%)	510 (35-1000)	2000 (-11000-16000)
Japan	2010	Female	75 to 79	70% (60-79%)	30% (21-40%)	1.2 (0.74-1.7)	-0.27% (-1.2-0.84%)	0.36% (-0.89-1.8%)	220 (-550-1100)	-4100 (-18000-13000)
Japan	2010	Female	80 to 84	71% (59-81%)	29% (19-41%)	1.2 (0.83-1.7)	-0.69% (-1.7-0.45%)	-0.55% (-1.9-0.86%)	-510 (-1800-810)	-11000 (-27000-6800)
Japan	2010	Female	85 to 89	70% (59-79%)	30% (21-41%)	1.3 (0.83-1.8)	-0.67% (-1.8-0.54%)	-0.67% (-2-0.82%)	-800 (-2400-980)	-8800 (-23000-7000)
Japan	2010	Female	90 to 94	71% (60-79%)	29% (21-40%)	1.3 (0.83-1.9)	-0.77% (-2-0.7%)	-0.92% (-2.4-0.88%)	-960 (-2500-920)	-5700 (-15000-5100)
Japan	2010	Female	95 plus	71% (58-82%)	29% (18-42%)	1.3 (0.73-2.1)	-0.82% (-2.5-1.2%)	-1.1% (-3-1.4%)	-820 (-2300-1000)	-2600 (-7900-4000)
Japan	2010	Male	All Ages	89% (87-91%)	11% (8.7-13%)	1.7 (1.6-1.9)	2.7% (1.5-3.7%)	2.1% (0.81-3.4%)	14000 (5200-22000)	470000 (270000-650000)
Japan	2010	Male	15 to 19	80% (70-88%)	20% (12-30%)	1 (0.62-1.5)	2.4% (1.1-3.9%)	6.4% (1.7-12%)	64 (17-120)	6800 (3300-11000)
Japan	2010	Male	20 to 24	89% (81-94%)	11% (6.3-19%)	2 (1.4-2.8)	6.1% (3.1-9.4%)	12% (4.2-20%)	240 (81-400)	22000 (11000-33000)
Japan	2010	Male	25 to 29	92% (86-95%)	8.4% (4.7-14%)	2.5 (1.7-3.4)	6.4% (3.2-9.7%)	14% (4.5-22%)	320 (110-520)	27000 (14000-40000)
Japan	2010	Male	30 to 34	93% (88-97%)	6.7% (3.4-12%)	2.6 (1.8-3.5)	5.7% (2.8-8.6%)	12% (4-20%)	390 (130-640)	31000 (15000-46000)
Japan	2010	Male	35 to 39	94% (90-97%)	5.6% (3-9.6%)	2.4 (1.6-3.3)	4.9% (2.4-7.6%)	10% (3.6-17%)	510 (180-850)	35000 (17000-52000)
Japan	2010	Male	40 to 44	95% (91-97%)	5.2% (2.7-9.1%)	2.3 (1.6-3.2)	4.7% (2.2-7.2%)	9.2% (3.5-15%)	620 (240-1000)	35000 (17000-53000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	2010	Male	45 to 49	95% (90-97%)	5.5% (2.8-10%)	2.4 (1.6-3.3)	4.9% (2.4-7.6%)	8.8% (3.9-13%)	850 (380-1300)	40000 (20000-60000)
Japan	2010	Male	50 to 54	94% (89-96%)	6.4% (3.6-11%)	2.3 (1.6-3.2)	4.6% (2.4-7%)	7.6% (3.8-11%)	1100 (570-1700)	45000 (23000-68000)
Japan	2010	Male	55 to 59	92% (87-96%)	8% (4.5-13%)	2.1 (1.5-2.8)	3.8% (2.1-5.7%)	5.9% (3.3-8.7%)	1600 (880-2300)	54000 (30000-81000)
Japan	2010	Male	60 to 64	90% (84-94%)	9.8% (5.7-16%)	1.9 (1.3-2.6)	3.1% (1.6-4.8%)	4.5% (2.4-6.9%)	2100 (1100-3300)	61000 (31000-95000)
Japan	2010	Male	65 to 69	88% (81-93%)	12% (6.9-19%)	1.8 (1.3-2.3)	2.4% (1.3-3.7%)	3.5% (2-5.2%)	2000 (1100-3000)	48000 (25000-72000)
Japan	2010	Male	70 to 74	87% (80-92%)	13% (7.5-20%)	1.6 (1.1-2.2)	1.8% (0.67-3.1%)	2.5% (1.1-4.1%)	1900 (820-3100)	35000 (13000-60000)
Japan	2010	Male	75 to 79	88% (82-93%)	12% (6.8-18%)	1.5 (0.96-2.1)	1.1% (0.013-2.4%)	1.4% (0.069-3%)	1500 (72-3100)	23000 (270-48000)
Japan	2010	Male	80 to 84	88% (80-94%)	12% (6.3-20%)	1.3 (0.88-1.8)	0.49% (-0.54-1.5%)	0.56% (-0.7-1.9%)	680 (-850-2300)	8100 (-9000-26000)
Japan	2010	Male	85 to 89	88% (80-93%)	12% (7-20%)	1.1 (0.76-1.6)	0.081% (-0.93-1.2%)	-0.02% (-1.3-1.3%)	-18 (-1200-1200)	710 (-8200-10000)
Japan	2010	Male	90 to 94	88% (81-93%)	12% (6.8-19%)	1 (0.65-1.5)	-0.23% (-1.2-0.86%)	-0.48% (-1.8-0.88%)	-230 (-870-430)	-720 (-4000-2700)
Japan	2010	Male	95 plus	89% (81-95%)	11% (5.2-19%)	0.89 (0.48-1.4)	-0.4% (-1.5-0.81%)	-0.78% (-2.3-0.67%)	-160 (-470-140)	-350 (-1300-710)
Japan	2016	Female	All Ages	77% (73-80%)	23% (20-27%)	1.9 (1.5-2.3)	0.86% (0.062-1.8%)	0.36% (-1-2%)	2400 (-6800-13000)	140000 (10000-270000)
Japan	2016	Female	15 to 19	77% (67-85%)	23% (15-33%)	1.9 (1.2-3)	1.5% (0.72-2.4%)	9.2% (3.6-15%)	37 (15-61)	3800 (1900-6100)
Japan	2016	Female	20 to 24	87% (79-92%)	13% (8.1-21%)	3.1 (2-4.7)	2.9% (1.4-4.6%)	16% (6.5-25%)	100 (43-170)	9300 (4800-14000)
Japan	2016	Female	25 to 29	90% (84-94%)	10% (6.2-16%)	3.2 (2.1-4.7)	2.8% (1.3-4.4%)	15% (6.2-24%)	130 (51-200)	10000 (5000-16000)
Japan	2016	Female	30 to 34	91% (87-95%)	8.5% (4.9-13%)	3.1 (2-4.5)	2.5% (1.2-4%)	13% (5.8-21%)	160 (70-260)	12000 (5700-18000)
Japan	2016	Female	35 to 39	91% (86-95%)	8.8% (5-14%)	2.8 (1.9-4)	2.4% (1.1-3.8%)	11% (6-17%)	220 (120-330)	14000 (6300-21000)
Japan	2016	Female	40 to 44	89% (84-93%)	11% (6.6-16%)	2.7 (1.7-3.9)	2.4% (1-4%)	10% (5.6-15%)	380 (210-550)	19000 (8400-30000)
Japan	2016	Female	45 to 49	86% (78-92%)	14% (8-22%)	2.5 (1.7-3.7)	2.6% (1.2-4.3%)	9.2% (5.2-13%)	470 (270-690)	20000 (9200-32000)
Japan	2016	Female	50 to 54	82% (74-88%)	18% (12-26%)	2.3 (1.5-3.4)	2.6% (1.1-4.4%)	8% (4.7-12%)	550 (320-800)	20000 (8800-33000)
Japan	2016	Female	55 to 59	78% (68-86%)	22% (14-32%)	2.1 (1.4-3)	2.2% (0.86-3.9%)	6.4% (3.8-9.3%)	600 (360-860)	18000 (7200-30000)
Japan	2016	Female	60 to 64	73% (63-81%)	27% (19-37%)	1.9 (1.2-2.7)	1.6% (0.28-3.2%)	4.4% (2.2-7.1%)	710 (350-1100)	17000 (3100-33000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	2016	Female	65 to 69	68% (57-78%)	32% (22-43%)	1.5 (1.1-2.1)	0.85% (-0.11-2%)	2.8% (1.3-4.3%)	760 (350-1200)	12000 (-1700-27000)
Japan	2016	Female	70 to 74	67% (55-76%)	33% (24-45%)	1.3 (0.83-1.8)	0.31% (-0.6-1.4%)	1.5% (0.2-2.9%)	590 (79-1200)	4600 (-9500-20000)
Japan	2016	Female	75 to 79	70% (60-78%)	30% (22-40%)	1.2 (0.71-1.8)	-0.094% (-1-1.1%)	0.54% (-0.75-2%)	310 (-440-1200)	-1500 (-16000-16000)
Japan	2016	Female	80 to 84	71% (59-81%)	29% (19-41%)	1.3 (0.8-1.8)	-0.48% (-1.5-0.66%)	-0.34% (-1.7-1.2%)	-340 (-1700-1200)	-8200 (-26000-11000)
Japan	2016	Female	85 to 89	70% (59-79%)	30% (21-41%)	1.3 (0.85-1.9)	-0.44% (-1.5-0.84%)	-0.43% (-1.9-1.2%)	-620 (-2600-1600)	-7100 (-24000-13000)
Japan	2016	Female	90 to 94	71% (60-79%)	29% (21-40%)	1.4 (0.85-2.1)	-0.51% (-1.8-1.1%)	-0.66% (-2.2-1.3%)	-940 (-3100-1800)	-5200 (-18000-12000)
Japan	2016	Female	95 plus	71% (57-82%)	29% (18-43%)	1.4 (0.75-2.4)	-0.51% (-2.3-1.9%)	-0.75% (-2.8-2.2%)	-750 (-2800-2100)	-2100 (-9400-7600)
Japan	2016	Male	All Ages	89% (87-91%)	11% (8.7-13%)	1.8 (1.4-2.1)	2.5% (1.5-3.6%)	2% (0.63-3.3%)	14000 (4300-22000)	450000 (260000-630000)
Japan	2016	Male	15 to 19	80% (69-88%)	20% (12-31%)	1 (0.59-1.6)	2.2% (1.1-3.6%)	6.4% (1.6-12%)	52 (13-97)	5900 (2900-9400)
Japan	2016	Male	20 to 24	89% (81-94%)	11% (6.4-19%)	2.1 (1.3-3.1)	5.8% (2.8-8.9%)	12% (3.7-21%)	190 (57-320)	18000 (9100-28000)
Japan	2016	Male	25 to 29	92% (86-95%)	8.4% (4.8-14%)	2.5 (1.6-3.7)	6.2% (3.2-9.5%)	14% (4.5-23%)	250 (78-410)	22000 (11000-33000)
Japan	2016	Male	30 to 34	93% (87-97%)	6.8% (3.4-13%)	2.6 (1.7-3.8)	5.6% (2.8-8.3%)	13% (4.2-21%)	300 (97-490)	24000 (12000-37000)
Japan	2016	Male	35 to 39	94% (90-97%)	5.6% (2.9-9.6%)	2.4 (1.6-3.4)	4.8% (2.3-7.2%)	11% (3.5-18%)	370 (120-610)	27000 (13000-40000)
Japan	2016	Male	40 to 44	95% (91-97%)	5.2% (2.7-8.8%)	2.4 (1.5-3.4)	4.5% (2.1-7.1%)	9.5% (3.3-15%)	630 (210-1000)	37000 (18000-56000)
Japan	2016	Male	45 to 49	95% (90-97%)	5.4% (2.8-10%)	2.5 (1.6-3.5)	4.8% (2.4-7.4%)	9.2% (4-14%)	830 (370-1300)	41000 (21000-61000)
Japan	2016	Male	50 to 54	94% (89-96%)	6.4% (3.6-11%)	2.4 (1.5-3.4)	4.6% (2.4-7%)	8% (4.1-12%)	1000 (520-1500)	42000 (23000-63000)
Japan	2016	Male	55 to 59	92% (86-96%)	8% (4.3-14%)	2.1 (1.4-3)	3.9% (2.3-5.9%)	6.3% (3.6-9.3%)	1200 (690-1800)	42000 (24000-62000)
Japan	2016	Male	60 to 64	90% (84-94%)	9.9% (5.6-16%)	1.9 (1.2-2.8)	3.3% (1.8-5.2%)	5% (2.9-7.5%)	1800 (1000-2600)	51000 (29000-78000)
Japan	2016	Male	65 to 69	88% (81-93%)	12% (6.8-19%)	1.8 (1.3-2.5)	2.6% (1.4-3.9%)	3.9% (2.2-5.6%)	2400 (1400-3400)	56000 (31000-83000)
Japan	2016	Male	70 to 74	87% (80-93%)	13% (7.4-20%)	1.7 (1.1-2.4)	2% (0.81-3.2%)	2.8% (1.2-4.4%)	2200 (970-3400)	42000 (17000-69000)
Japan	2016	Male	75 to 79	88% (81-93%)	12% (6.9-19%)	1.5 (0.95-2.2)	1.3% (0.2-2.5%)	1.7% (0.3-3.3%)	1700 (290-3300)	26000 (4100-52000)
Japan	2016	Male	80 to 84	88% (80-94%)	12% (6.4-20%)	1.3 (0.87-1.8)	0.63% (-0.37-1.7%)	0.73% (-0.51-2.1%)	970 (-660-2700)	12000 (-6800-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Japan	2016	Male	85 to 89	88% (80-93%)	12% (7.2-20%)	1.1 (0.75-1.6)	0.2% (-0.81-1.3%)	0.11% (-1.1-1.5%)	140 (-1400-1900)	2600 (-10000-16000)
Japan	2016	Male	90 to 94	88% (81-93%)	12% (7.1-19%)	0.99 (0.61-1.5)	-0.12% (-1.1-0.95%)	-0.38% (-1.6-0.97%)	-250 (-1100-650)	-530 (-4900-4100)
Japan	2016	Male	95 plus	89% (80-95%)	11% (5.4-20%)	0.87 (0.45-1.5)	-0.31% (-1.4-0.85%)	-0.71% (-2.1-0.77%)	-170 (-520-190)	-320 (-1400-860)
South Korea	1990	Female	All Ages	76% (70-82%)	24% (18-30%)	1.9 (1.7-2.2)	2.9% (1.5-4.5%)	1.9% (-0.78-5%)	1900 (-700-4900)	140000 (71000-210000)
South Korea	1990	Female	15 to 19	70% (54-84%)	30% (16-46%)	1.7 (1-2.6)	3.3% (1.8-5%)	7.7% (3.5-13%)	85 (38-140)	8500 (4600-13000)
South Korea	1990	Female	20 to 24	81% (69-90%)	19% (10-31%)	3 (2.1-4.2)	6.3% (4.3-8.7%)	13% (7.6-19%)	190 (110-280)	19000 (13000-26000)
South Korea	1990	Female	25 to 29	85% (75-92%)	15% (7.7-25%)	3.2 (2.2-4.2)	6% (4.1-7.9%)	13% (8.5-18%)	220 (130-310)	19000 (13000-26000)
South Korea	1990	Female	30 to 34	87% (76-94%)	13% (6.5-24%)	3.1 (2.1-4)	5.5% (3.7-7.6%)	13% (8.2-17%)	250 (160-340)	19000 (13000-26000)
South Korea	1990	Female	35 to 39	87% (78-94%)	13% (6.3-22%)	2.9 (2-3.9)	5.4% (3.4-7.7%)	12% (7.5-17%)	250 (150-360)	16000 (9800-22000)
South Korea	1990	Female	40 to 44	86% (77-93%)	14% (6.9-23%)	2.7 (1.7-3.8)	5.6% (3.1-8.2%)	12% (6.6-17%)	280 (160-420)	15000 (8300-22000)
South Korea	1990	Female	45 to 49	85% (72-93%)	15% (7.4-28%)	2.6 (1.7-3.7)	5.3% (2.4-8.4%)	9.8% (4.6-15%)	340 (160-530)	15000 (7000-24000)
South Korea	1990	Female	50 to 54	83% (71-91%)	17% (9.2-29%)	2.4 (1.5-3.5)	5% (1.7-8.6%)	8.4% (3.1-14%)	400 (140-690)	15000 (5300-27000)
South Korea	1990	Female	55 to 59	80% (66-90%)	20% (10-34%)	2.3 (1.4-3.4)	4.1% (0.89-7.8%)	6.7% (1.7-12%)	390 (99-730)	13000 (2600-24000)
South Korea	1990	Female	60 to 64	77% (62-88%)	23% (12-38%)	2 (1.2-3.2)	2.4% (-0.99-6.7%)	3.7% (-1-9.5%)	270 (-72-690)	7100 (-2900-20000)
South Korea	1990	Female	65 to 69	75% (58-87%)	25% (13-42%)	1.7 (1.1-2.4)	0.66% (-2.3-4.4%)	1.1% (-2.8-6.1%)	110 (-280-570)	2000 (-7100-13000)
South Korea	1990	Female	70 to 74	73% (57-85%)	27% (15-43%)	1.3 (0.81-2)	-0.8% (-3.6-2.5%)	-0.86% (-4.4-3.3%)	-100 (-520-390)	-2300 (-10000-6800)
South Korea	1990	Female	75 to 79	72% (55-84%)	28% (16-45%)	1.2 (0.7-1.9)	-1.7% (-4.6-1.6%)	-2.1% (-5.5-1.9%)	-300 (-800-260)	-4400 (-12000-3800)
South Korea	1990	Female	80 to 84	70% (51-83%)	30% (17-49%)	1.3 (0.77-1.9)	-1.7% (-4.4-1.3%)	-2.1% (-5.3-1.3%)	-290 (-720-180)	-3000 (-7500-2100)
South Korea	1990	Female	85 to 89	67% (49-80%)	33% (20-51%)	1.3 (0.77-2)	-1.4% (-4-1.6%)	-1.8% (-4.9-1.8%)	-150 (-410-140)	-1000 (-2900-1200)
South Korea	1990	Female	90 to 94	67% (50-80%)	33% (20-50%)	1.4 (0.81-2.2)	-1.1% (-3.8-2%)	-1.6% (-4.7-2.2%)	-46 (-140-60)	-200 (-660-340)
South Korea	1990	Female	95 plus	67% (46-83%)	33% (17-54%)	1.5 (0.76-2.5)	-0.79% (-3.8-3%)	-1.3% (-4.9-3.2%)	-8.4 (-32-21)	-21 (-100-80)
South Korea	1990	Male	All Ages	89% (85-93%)	11% (7.4-15%)	1.9 (1.7-2.1)	8.2% (6.4-10%)	8.1% (5.4-11%)	11000 (7200-15000)	520000 (390000-660000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	1990	Male	15 to 19	83% (71-91%)	17% (8.5-29%)	1.4 (0.83-2)	5.6% (3-8.2%)	7.7% (3.2-12%)	200 (81-320)	20000 (11000-29000)
South Korea	1990	Male	20 to 24	91% (83-96%)	9.4% (4.4-17%)	2.5 (1.8-3.4)	11% (7.8-15%)	14% (7.7-21%)	440 (240-680)	44000 (29000-60000)
South Korea	1990	Male	25 to 29	93% (88-97%)	6.8% (3.3-12%)	2.9 (2.1-3.9)	12% (8.4-16%)	15% (8.5-21%)	610 (350-890)	52000 (35000-71000)
South Korea	1990	Male	30 to 34	95% (89-98%)	5.5% (2.4-11%)	3 (2.1-4.1)	12% (8.5-16%)	15% (9.4-21%)	780 (480-1100)	57000 (39000-75000)
South Korea	1990	Male	35 to 39	95% (90-98%)	4.8% (2.1-9.7%)	2.8 (2-3.9)	12% (9-16%)	16% (11-20%)	890 (600-1200)	55000 (39000-70000)
South Korea	1990	Male	40 to 44	96% (91-98%)	4.5% (1.9-9%)	2.8 (1.8-3.9)	13% (9.3-16%)	16% (11-20%)	1100 (770-1400)	57000 (41000-74000)
South Korea	1990	Male	45 to 49	96% (91-98%)	4.4% (2-9.2%)	2.8 (1.9-4)	13% (9.2-16%)	15% (11-20%)	1500 (1000-1900)	66000 (46000-85000)
South Korea	1990	Male	50 to 54	95% (90-98%)	4.7% (2.2-9.6%)	2.8 (1.8-3.9)	11% (7.8-15%)	13% (9.2-18%)	1700 (1100-2300)	64000 (44000-87000)
South Korea	1990	Male	55 to 59	95% (90-98%)	5.3% (2.3-10%)	2.6 (1.7-3.7)	9% (5.8-12%)	11% (6.7-15%)	1300 (830-1900)	45000 (28000-64000)
South Korea	1990	Male	60 to 64	94% (88-97%)	5.8% (2.5-12%)	2.4 (1.4-3.6)	7.6% (3.5-11%)	7.6% (3.9-12%)	980 (490-1600)	28000 (14000-44000)
South Korea	1990	Male	65 to 69	94% (88-97%)	6.3% (2.5-12%)	2.2 (1.5-3.1)	4.9% (1.8-8.3%)	5.5% (2-9.5%)	830 (290-1400)	20000 (7200-34000)
South Korea	1990	Male	70 to 74	93% (87-97%)	7% (3.3-13%)	2 (1.3-2.9)	3.3% (0.23-7.1%)	3.6% (0.095-7.9%)	540 (13-1200)	10000 (730-22000)
South Korea	1990	Male	75 to 79	92% (86-97%)	7.6% (3.5-14%)	1.8 (1.1-2.7)	1.9% (-1.1-5.7%)	1.9% (-1.4-6.2%)	240 (-180-780)	3800 (-2200-12000)
South Korea	1990	Male	80 to 84	92% (84-96%)	8.4% (3.9-16%)	1.6 (1.1-2.4)	0.84% (-1.9-4%)	0.7% (-2.4-4.3%)	59 (-200-360)	850 (-1900-4100)
South Korea	1990	Male	85 to 89	91% (83-95%)	9.4% (4.6-17%)	1.5 (0.99-2.2)	0.08% (-2.3-2.6%)	-0.28% (-3-2.5%)	-10 (-110-91)	25 (-740-800)
South Korea	1990	Male	90 to 94	90% (83-95%)	9.7% (4.7-17%)	1.4 (0.87-2.2)	-0.48% (-2.7-1.8%)	-1.1% (-3.8-1.6%)	-11 (-37-16)	-28 (-160-100)
South Korea	1990	Male	95 plus	90% (81-96%)	9.6% (3.8-19%)	1.3 (0.73-2.2)	-0.77% (-3.1-1.8%)	-1.6% (-4.4-1.5%)	-3.2 (-8.8-3)	-6.2 (-25-15)
South Korea	1995	Female	All Ages	77% (70-82%)	23% (18-30%)	2.2 (2-2.5)	3.3% (1.9-5%)	2.9% (0.068-6.1%)	3000 (71-6500)	150000 (87000-230000)
South Korea	1995	Female	15 to 19	69% (52-83%)	31% (17-48%)	1.8 (1.1-2.8)	3.4% (2-5.2%)	9.1% (4.5-15%)	75 (36-120)	7300 (4200-11000)
South Korea	1995	Female	20 to 24	81% (67-90%)	19% (10-33%)	3.3 (2.2-4.7)	6% (4.1-8.4%)	15% (8.6-21%)	170 (93-240)	17000 (11000-23000)
South Korea	1995	Female	25 to 29	85% (75-92%)	15% (8-25%)	3.6 (2.5-4.9)	6% (4-8.1%)	15% (9.4-21%)	190 (120-270)	18000 (12000-24000)
South Korea	1995	Female	30 to 34	87% (76-93%)	13% (6.6-24%)	3.5 (2.4-4.7)	5.4% (3.7-7.5%)	14% (9.3-19%)	220 (140-310)	17000 (12000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	1995	Female	35 to 39	87% (77-94%)	13% (6.4-23%)	3.1 (2.2-4.3)	5.1% (3.3-7.2%)	13% (8.4-17%)	280 (180-390)	18000 (12000-25000)
South Korea	1995	Female	40 to 44	86% (77-93%)	14% (7.3-23%)	3 (1.9-4.1)	5.1% (2.9-7.4%)	12% (7.5-17%)	280 (160-390)	15000 (8800-22000)
South Korea	1995	Female	45 to 49	85% (71-92%)	15% (7.7-29%)	2.9 (1.9-4.3)	5.3% (2.8-7.9%)	11% (6.4-16%)	310 (170-450)	15000 (7800-22000)
South Korea	1995	Female	50 to 54	83% (70-90%)	17% (9.6-30%)	2.8 (1.7-4)	5.5% (2.4-8.9%)	10% (5-15%)	390 (190-600)	15000 (6800-24000)
South Korea	1995	Female	55 to 59	80% (66-90%)	20% (10-34%)	2.5 (1.6-3.7)	4.5% (1.4-8.1%)	7.8% (2.9-13%)	430 (160-730)	14000 (4300-25000)
South Korea	1995	Female	60 to 64	77% (62-88%)	23% (12-38%)	2.2 (1.3-3.5)	3% (-0.19-7.2%)	4.9% (0.37-11%)	350 (27-770)	9600 (-620-23000)
South Korea	1995	Female	65 to 69	74% (57-87%)	26% (13-43%)	2 (1.3-2.9)	2% (-1.2-5.8%)	3% (-1.2-8.2%)	290 (-110-740)	6300 (-3600-18000)
South Korea	1995	Female	70 to 74	72% (55-85%)	28% (15-45%)	1.8 (1.1-2.7)	1.1% (-2.1-5.3%)	1.6% (-2.6-6.7%)	210 (-330-880)	3500 (-7000-17000)
South Korea	1995	Female	75 to 79	71% (55-84%)	29% (16-45%)	1.6 (0.95-2.6)	0.16% (-3.2-4.3%)	0.23% (-3.7-5.2%)	35 (-550-830)	430 (-8300-12000)
South Korea	1995	Female	80 to 84	69% (51-83%)	31% (17-49%)	1.5 (0.9-2.3)	-0.41% (-3.1-2.9%)	-0.53% (-3.8-3.4%)	-85 (-600-540)	-850 (-6500-5900)
South Korea	1995	Female	85 to 89	67% (49-80%)	33% (20-51%)	1.4 (0.82-2.1)	-0.58% (-3.1-2.5%)	-0.8% (-3.7-2.7%)	-94 (-430-320)	-620 (-3200-2600)
South Korea	1995	Female	90 to 94	66% (49-80%)	34% (20-51%)	1.3 (0.79-2.1)	-0.88% (-3.3-2.1%)	-1.2% (-4-2.2%)	-60 (-200-110)	-270 (-990-600)
South Korea	1995	Female	95 plus	67% (46-83%)	33% (17-54%)	1.3 (0.65-2.2)	-1.1% (-3.5-2.2%)	-1.5% (-4.5-2.3%)	-16 (-48-25)	-46 (-150-95)
South Korea	1995	Male	All Ages	90% (86-93%)	10% (6.9-14%)	2.1 (1.9-2.4)	9.2% (7.3-11%)	9.2% (6.7-12%)	13000 (9000-17000)	560000 (440000-690000)
South Korea	1995	Male	15 to 19	83% (70-91%)	17% (8.6-30%)	1.4 (0.86-2.1)	6.1% (3.4-8.9%)	9% (4.1-14%)	180 (81-290)	17000 (9900-25000)
South Korea	1995	Male	20 to 24	91% (83-96%)	9.4% (4.5-17%)	2.7 (1.9-3.7)	12% (8.2-16%)	15% (8.2-22%)	420 (220-640)	43000 (28000-59000)
South Korea	1995	Male	25 to 29	93% (88-97%)	6.8% (3.3-12%)	3.3 (2.3-4.4)	13% (8.9-17%)	17% (9.5-24%)	580 (320-840)	50000 (34000-68000)
South Korea	1995	Male	30 to 34	95% (89-98%)	5.5% (2.5-11%)	3.4 (2.4-4.5)	13% (9.1-17%)	17% (11-22%)	740 (460-1000)	55000 (39000-74000)
South Korea	1995	Male	35 to 39	95% (90-98%)	4.8% (2-9.7%)	3.1 (2.2-4.2)	13% (9.8-16%)	17% (12-22%)	1100 (740-1400)	66000 (49000-85000)
South Korea	1995	Male	40 to 44	95% (91-98%)	4.5% (1.9-8.6%)	3 (2-4.2)	14% (10-17%)	17% (13-22%)	1200 (870-1600)	64000 (47000-82000)
South Korea	1995	Male	45 to 49	96% (91-98%)	4.5% (2-9%)	3.1 (2.1-4.3)	14% (10-17%)	17% (13-21%)	1400 (1000-1800)	64000 (48000-83000)
South Korea	1995	Male	50 to 54	95% (91-98%)	4.7% (2.1-8.9%)	3 (1.9-4.4)	12% (8.9-16%)	15% (11-19%)	1600 (1100-2100)	62000 (44000-80000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	1995	Male	55 to 59	95% (90-98%)	5.2% (2.3-10%)	2.8 (1.8-4)	9.9% (6.7-13%)	12% (8-16%)	1600 (1100-2300)	55000 (37000-76000)
South Korea	1995	Male	60 to 64	94% (89-97%)	5.8% (2.6-11%)	2.6 (1.6-4)	7.5% (4.5-11%)	8.9% (5.3-13%)	1300 (740-1900)	36000 (21000-54000)
South Korea	1995	Male	65 to 69	94% (87-97%)	6.5% (2.7-13%)	2.5 (1.7-3.4)	6% (3.2-9.1%)	6.9% (3.7-11%)	960 (510-1500)	23000 (12000-35000)
South Korea	1995	Male	70 to 74	93% (86-97%)	7.2% (3.2-14%)	2.3 (1.5-3.3)	4.7% (1.8-8.2%)	5.3% (2-9.2%)	840 (320-1500)	16000 (6000-28000)
South Korea	1995	Male	75 to 79	92% (86-97%)	7.7% (3.4-14%)	2.1 (1.3-3.1)	3.2% (0.31-7%)	3.5% (0.21-7.8%)	480 (27-1100)	7200 (680-16000)
South Korea	1995	Male	80 to 84	92% (84-96%)	8.4% (3.8-16%)	1.8 (1.2-2.7)	2% (-0.59-5.2%)	2.1% (-0.89-5.6%)	200 (-87-550)	2400 (-740-6100)
South Korea	1995	Male	85 to 89	91% (83-95%)	9.4% (4.6-17%)	1.6 (1.1-2.4)	1% (-1.2-3.4%)	0.84% (-1.7-3.6%)	41 (-86-180)	440 (-520-1500)
South Korea	1995	Male	90 to 94	90% (83-95%)	9.7% (4.6-17%)	1.4 (0.89-2.2)	0.23% (-1.8-2.3%)	-0.24% (-2.6-2.2%)	-3.3 (-36-30)	20 (-150-200)
South Korea	1995	Male	95 plus	91% (81-96%)	9.4% (3.7-19%)	1.3 (0.69-2.1)	-0.33% (-2.4-2%)	-1.1% (-3.5-1.6%)	-2.7 (-8.5-3.8)	-3.3 (-23-20)
South Korea	2000	Female	All Ages	77% (71-82%)	23% (18-29%)	2.4 (2.1-2.6)	3.1% (1.7-4.7%)	2.7% (-0.059-6%)	3000 (-67-6700)	150000 (80000-220000)
South Korea	2000	Female	15 to 19	68% (52-83%)	32% (17-48%)	1.9 (1.1-2.9)	3.1% (1.7-4.8%)	9.9% (4.6-16%)	56 (26-93)	5800 (3300-8800)
South Korea	2000	Female	20 to 24	81% (67-90%)	19% (10-33%)	3.5 (2.3-4.9)	5.7% (3.7-8%)	17% (9.2-24%)	120 (67-190)	13000 (8300-18000)
South Korea	2000	Female	25 to 29	85% (74-92%)	15% (8-26%)	3.8 (2.6-5.1)	5.6% (3.7-7.7%)	17% (10-24%)	170 (98-240)	15000 (10000-21000)
South Korea	2000	Female	30 to 34	87% (75-93%)	13% (6.7-25%)	3.7 (2.5-4.9)	5.3% (3.5-7.2%)	15% (9.7-21%)	210 (130-290)	16000 (11000-22000)
South Korea	2000	Female	35 to 39	87% (77-93%)	13% (6.5-23%)	3.3 (2.3-4.5)	5.1% (3.2-7%)	14% (9.1-18%)	260 (170-360)	17000 (11000-24000)
South Korea	2000	Female	40 to 44	86% (76-93%)	14% (7-24%)	3.1 (2.1-4.4)	5.1% (3-7.3%)	13% (8.3-17%)	330 (200-460)	19000 (11000-26000)
South Korea	2000	Female	45 to 49	85% (71-92%)	15% (7.5-29%)	3.1 (2-4.3)	5% (2.8-7.4%)	12% (7.2-16%)	310 (190-440)	15000 (8600-22000)
South Korea	2000	Female	50 to 54	82% (70-91%)	18% (9.3-30%)	2.9 (1.8-4.1)	4.9% (2.3-7.8%)	10% (5.7-15%)	330 (180-490)	13000 (6600-21000)
South Korea	2000	Female	55 to 59	80% (65-89%)	20% (11-35%)	2.6 (1.7-3.8)	4.5% (1.7-7.6%)	8.2% (3.8-13%)	390 (180-620)	13000 (5000-22000)
South Korea	2000	Female	60 to 64	76% (61-88%)	24% (12-39%)	2.3 (1.4-3.7)	3.2% (0.03-6.9%)	5.5% (0.92-11%)	380 (61-760)	11000 (85-23000)
South Korea	2000	Female	65 to 69	73% (56-87%)	27% (13-44%)	2 (1.3-2.9)	1.8% (-0.99-5.3%)	3.1% (-0.83-7.7%)	300 (-79-770)	6500 (-3600-18000)
South Korea	2000	Female	70 to 74	71% (55-84%)	29% (16-45%)	1.7 (1-2.6)	0.74% (-2.2-4.4%)	1.2% (-2.7-5.8%)	170 (-360-800)	2600 (-7900-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	2000	Female	75 to 79	70% (53-83%)	30% (17-47%)	1.6 (0.89-2.5)	-0.071% (-3-3.5%)	-0.0035% (-3.6-4.5%)	-0.62 (-640-790)	-240 (-9800-12000)
South Korea	2000	Female	80 to 84	68% (49-83%)	32% (17-51%)	1.5 (0.92-2.3)	-0.23% (-2.8-3.1%)	-0.28% (-3.4-3.8%)	-51 (-620-690)	-540 (-7000-7600)
South Korea	2000	Female	85 to 89	66% (47-79%)	34% (21-53%)	1.5 (0.86-2.3)	-0.029% (-2.5-3.1%)	-0.11% (-3.1-3.6%)	-16 (-430-530)	-38 (-3300-4200)
South Korea	2000	Female	90 to 94	66% (49-80%)	34% (20-51%)	1.5 (0.88-2.4)	0.031% (-2.4-3%)	-0.12% (-3.1-3.4%)	-8.8 (-220-250)	14 (-1100-1300)
South Korea	2000	Female	95 plus	67% (45-83%)	33% (17-55%)	1.6 (0.77-2.6)	0.029% (-2.7-3.7%)	-0.26% (-3.5-4.1%)	-4.8 (-67-78)	2.4 (-210-290)
South Korea	2000	Male	All Ages	90% (87-93%)	9.7% (6.8-13%)	2.3 (2-2.5)	9% (7.1-11%)	8.9% (6.6-11%)	12000 (9100-16000)	540000 (420000-660000)
South Korea	2000	Male	15 to 19	82% (70-91%)	18% (8.6-30%)	1.5 (0.89-2.2)	5.7% (3.5-8.2%)	9.6% (4.5-15%)	120 (56-200)	13000 (7800-19000)
South Korea	2000	Male	20 to 24	90% (83-95%)	9.6% (4.6-17%)	2.9 (1.9-3.9)	12% (8.4-16%)	16% (8.8-24%)	290 (160-430)	32000 (22000-44000)
South Korea	2000	Male	25 to 29	93% (88-97%)	6.9% (3.4-12%)	3.3 (2.3-4.6)	13% (9.2-17%)	18% (10-25%)	420 (230-600)	40000 (27000-53000)
South Korea	2000	Male	30 to 34	94% (89-98%)	5.6% (2.4-11%)	3.4 (2.4-4.6)	12% (8.9-16%)	17% (11-24%)	540 (330-760)	43000 (30000-57000)
South Korea	2000	Male	35 to 39	95% (90-98%)	4.9% (2.1-9.6%)	3.2 (2.3-4.4)	13% (9.6-16%)	18% (13-23%)	880 (610-1200)	57000 (42000-74000)
South Korea	2000	Male	40 to 44	95% (91-98%)	4.6% (1.9-9%)	3.1 (2.1-4.3)	13% (10-17%)	18% (13-22%)	1400 (1000-1800)	74000 (56000-95000)
South Korea	2000	Male	45 to 49	95% (91-98%)	4.5% (1.9-9.5%)	3.2 (2.1-4.4)	13% (10-17%)	17% (13-21%)	1500 (1100-1900)	68000 (50000-86000)
South Korea	2000	Male	50 to 54	95% (90-98%)	4.8% (2.1-9.8%)	3.1 (2.1-4.4)	12% (8.8-15%)	15% (11-19%)	1400 (1000-1900)	57000 (41000-74000)
South Korea	2000	Male	55 to 59	95% (90-98%)	5.3% (2.4-10%)	2.9 (1.9-4.2)	10% (7.1-13%)	12% (8.7-16%)	1500 (1100-2100)	52000 (36000-71000)
South Korea	2000	Male	60 to 64	94% (88-97%)	5.8% (2.5-12%)	2.7 (1.7-4)	7.8% (4.9-12%)	9.4% (5.9-14%)	1500 (930-2300)	44000 (27000-65000)
South Korea	2000	Male	65 to 69	94% (87-97%)	6.5% (2.7-13%)	2.5 (1.6-3.4)	5.9% (3.3-8.9%)	7.1% (4.1-11%)	1100 (660-1700)	27000 (15000-42000)
South Korea	2000	Male	70 to 74	93% (86-97%)	7.3% (3.3-14%)	2.2 (1.4-3.2)	4.3% (1.9-7.4%)	5.1% (2.2-8.7%)	800 (340-1400)	15000 (6300-26000)
South Korea	2000	Male	75 to 79	92% (85-96%)	7.9% (3.6-15%)	2 (1.2-3)	3% (0.49-6.3%)	3.4% (0.52-7.1%)	550 (84-1200)	8200 (1400-17000)
South Korea	2000	Male	80 to 84	91% (83-96%)	8.8% (4-17%)	1.8 (1.2-2.7)	2.2% (-0.041-5.1%)	2.4% (-0.21-5.7%)	290 (-27-680)	3300 (-60-7700)
South Korea	2000	Male	85 to 89	90% (81-95%)	10% (4.6-19%)	1.7 (1.1-2.5)	1.7% (-0.45-4.1%)	1.7% (-0.75-4.4%)	110 (-49-280)	950 (-240-2300)
South Korea	2000	Male	90 to 94	90% (81-95%)	10% (4.8-19%)	1.6 (0.99-2.5)	1.2% (-0.79-3.6%)	0.99% (-1.3-3.8%)	22 (-29-84)	160 (-110-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	2000	Male	95 plus	90% (79-96%)	10% (3.9-21%)	1.6 (0.86-2.5)	0.71% (-1.4-3.4%)	0.16% (-2.3-3.3%)	0.71 (-9.9-15)	13 (-26-62)
South Korea	2005	Female	All Ages	77% (71-82%)	23% (18-29%)	2.4 (2.2-2.7)	3% (1.8-4.4%)	2.6% (0.23-5.4%)	2900 (280-6200)	150000 (84000-210000)
South Korea	2005	Female	15 to 19	67% (50-82%)	33% (18-50%)	1.9 (1.1-2.9)	2.6% (1.6-3.9%)	10% (4.2-17%)	35 (14-60)	4400 (2600-6500)
South Korea	2005	Female	20 to 24	80% (67-90%)	20% (10-33%)	3.5 (2.3-5.1)	6.3% (4-8.9%)	18% (9.3-28%)	120 (60-180)	13000 (8200-18000)
South Korea	2005	Female	25 to 29	85% (73-92%)	15% (8.2-27%)	3.8 (2.6-5.2)	6.1% (4-8.4%)	19% (10-28%)	160 (85-230)	15000 (9800-21000)
South Korea	2005	Female	30 to 34	86% (74-93%)	14% (6.8-26%)	3.7 (2.5-5.1)	5.5% (3.6-7.6%)	17% (9.7-24%)	190 (110-290)	16000 (10000-22000)
South Korea	2005	Female	35 to 39	87% (76-93%)	13% (6.9-24%)	3.4 (2.4-4.5)	5.1% (3.3-7.1%)	15% (9.1-20%)	250 (150-350)	17000 (11000-23000)
South Korea	2005	Female	40 to 44	86% (75-93%)	14% (7.1-25%)	3.2 (2.1-4.4)	5% (3.1-7.2%)	14% (8.6-18%)	320 (200-450)	18000 (11000-25000)
South Korea	2005	Female	45 to 49	84% (70-92%)	16% (7.7-30%)	3.1 (2-4.4)	5.1% (3.1-7.6%)	12% (7.8-17%)	380 (240-540)	18000 (11000-26000)
South Korea	2005	Female	50 to 54	82% (69-91%)	18% (9.3-31%)	2.9 (1.9-4.1)	4.4% (2.3-6.9%)	10% (6.4-15%)	350 (210-490)	15000 (7800-22000)
South Korea	2005	Female	55 to 59	79% (65-90%)	21% (10-35%)	2.6 (1.7-3.8)	3.8% (1.5-6.4%)	8.3% (4.7-12%)	320 (180-490)	11000 (4800-18000)
South Korea	2005	Female	60 to 64	76% (60-88%)	24% (12-40%)	2.4 (1.4-3.6)	3% (0.31-6.1%)	5.7% (1.9-10%)	340 (110-620)	9500 (1100-19000)
South Korea	2005	Female	65 to 69	73% (55-86%)	27% (14-45%)	2 (1.3-2.9)	1.8% (-0.56-4.7%)	3.4% (-0.054-7.3%)	310 (-5.2-670)	6500 (-2100-16000)
South Korea	2005	Female	70 to 74	71% (54-84%)	29% (16-46%)	1.7 (1-2.6)	0.77% (-1.7-3.7%)	1.5% (-1.8-5.3%)	200 (-250-730)	2900 (-6300-14000)
South Korea	2005	Female	75 to 79	69% (53-83%)	31% (17-47%)	1.6 (0.89-2.4)	-0.017% (-2.5-3%)	0.13% (-3-4.1%)	24 (-550-740)	-74 (-9400-11000)
South Korea	2005	Female	80 to 84	67% (48-82%)	33% (18-52%)	1.5 (0.89-2.3)	-0.19% (-2.4-2.6%)	-0.22% (-3-3.2%)	-45 (-610-650)	-550 (-6900-7400)
South Korea	2005	Female	85 to 89	64% (46-79%)	36% (21-54%)	1.5 (0.86-2.3)	0.016% (-2.2-2.8%)	-0.049% (-2.6-3.3%)	-8.5 (-440-540)	24 (-3300-4500)
South Korea	2005	Female	90 to 94	64% (47-79%)	36% (21-53%)	1.5 (0.86-2.3)	0.15% (-2.1-3%)	0.021% (-2.7-3.4%)	1.8 (-250-300)	88 (-1300-1700)
South Korea	2005	Female	95 plus	65% (43-82%)	35% (18-57%)	1.6 (0.77-2.6)	0.25% (-2.4-3.6%)	0.013% (-3.1-4.1%)	0.36 (-94-120)	33 (-300-460)
South Korea	2005	Male	All Ages	91% (87-93%)	9.4% (6.5-13%)	2.4 (2.1-2.6)	8.3% (6.6-10%)	8.3% (6-10%)	12000 (8200-15000)	490000 (370000-600000)
South Korea	2005	Male	15 to 19	82% (69-91%)	18% (9.1-31%)	1.5 (0.86-2.2)	4.9% (3.2-6.8%)	9.2% (4.1-15%)	65 (30-110)	9100 (5800-13000)
South Korea	2005	Male	20 to 24	90% (83-95%)	9.8% (4.7-17%)	2.8 (1.9-3.9)	12% (8.3-15%)	16% (8.3-24%)	190 (95-290)	25000 (17000-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	2005	Male	25 to 29	93% (87-97%)	7% (3.4-13%)	3.4 (2.3-4.6)	13% (9.1-16%)	18% (9.6-27%)	290 (150-430)	32000 (22000-43000)
South Korea	2005	Male	30 to 34	94% (88-98%)	5.7% (2.4-12%)	3.5 (2.4-4.7)	12% (8.3-16%)	18% (9.8-25%)	400 (230-580)	35000 (24000-48000)
South Korea	2005	Male	35 to 39	95% (90-98%)	5% (2.3-9.7%)	3.2 (2.3-4.4)	12% (8.3-15%)	17% (11-23%)	620 (400-850)	44000 (32000-57000)
South Korea	2005	Male	40 to 44	95% (91-98%)	4.7% (2-9.4%)	3.1 (2.1-4.3)	12% (9.3-16%)	17% (12-22%)	1100 (750-1400)	60000 (45000-78000)
South Korea	2005	Male	45 to 49	95% (90-98%)	4.6% (2-9.8%)	3.2 (2.2-4.5)	13% (9.5-16%)	17% (12-21%)	1500 (1100-2000)	72000 (53000-92000)
South Korea	2005	Male	50 to 54	95% (90-98%)	4.9% (2.2-10%)	3.1 (2.1-4.4)	11% (8.1-14%)	15% (11-19%)	1500 (1000-1900)	60000 (43000-79000)
South Korea	2005	Male	55 to 59	95% (90-98%)	5.4% (2.4-10%)	2.9 (1.9-4.1)	8.9% (6.3-12%)	12% (8.4-15%)	1300 (870-1700)	44000 (31000-60000)
South Korea	2005	Male	60 to 64	94% (88-97%)	6% (2.7-12%)	2.7 (1.7-4)	7.3% (4.6-11%)	9.2% (5.9-13%)	1300 (830-1900)	39000 (24000-57000)
South Korea	2005	Male	65 to 69	93% (87-97%)	6.7% (2.6-13%)	2.5 (1.8-3.5)	5.8% (3.5-8.7%)	7.2% (4.4-10%)	1300 (790-1900)	31000 (19000-46000)
South Korea	2005	Male	70 to 74	92% (85-97%)	7.5% (3.2-15%)	2.4 (1.5-3.4)	4.6% (2.1-7.4%)	5.6% (2.7-8.8%)	1000 (490-1600)	20000 (9300-33000)
South Korea	2005	Male	75 to 79	92% (84-96%)	8.3% (3.6-16%)	2.1 (1.3-3.2)	3.2% (1-6.1%)	3.8% (1.2-7.2%)	650 (210-1200)	10000 (3000-19000)
South Korea	2005	Male	80 to 84	91% (82-96%)	9.1% (4.2-18%)	1.9 (1.2-2.7)	2.1% (0.16-4.6%)	2.3% (0.032-5.3%)	320 (4.6-730)	3800 (270-8300)
South Korea	2005	Male	85 to 89	90% (81-95%)	10% (4.8-19%)	1.6 (1-2.3)	1.3% (-0.4-3.5%)	1.3% (-0.75-3.8%)	110 (-59-310)	1000 (-300-2600)
South Korea	2005	Male	90 to 94	89% (81-95%)	11% (5-19%)	1.4 (0.9-2.2)	0.78% (-0.87-2.8%)	0.51% (-1.5-2.8%)	16 (-45-85)	150 (-170-540)
South Korea	2005	Male	95 plus	90% (79-96%)	10% (4-21%)	1.3 (0.69-2.2)	0.28% (-1.4-2.3%)	-0.31% (-2.4-2.2%)	-2.1 (-16-15)	8 (-41-68)
South Korea	2010	Female	All Ages	77% (71-82%)	23% (18-29%)	2.3 (2-2.7)	2.8% (1.6-4.2%)	2.3% (0.17-4.7%)	2800 (210-5800)	140000 (78000-210000)
South Korea	2010	Female	15 to 19	67% (50-82%)	33% (18-50%)	1.8 (1-2.8)	2.5% (1.4-3.9%)	9.8% (3.3-18%)	32 (11-57)	4000 (2300-6300)
South Korea	2010	Female	20 to 24	80% (66-89%)	20% (11-34%)	3.3 (2.1-4.9)	5.8% (3.6-8.5%)	18% (7.9-28%)	92 (40-150)	11000 (6600-15000)
South Korea	2010	Female	25 to 29	85% (74-92%)	15% (8.1-26%)	3.6 (2.5-5.1)	6.4% (3.8-9.3%)	20% (9.1-30%)	170 (79-270)	15000 (9000-22000)
South Korea	2010	Female	30 to 34	86% (74-93%)	14% (6.8-26%)	3.6 (2.4-4.9)	5.5% (3.3-8.1%)	17% (9-26%)	190 (96-290)	15000 (8900-21000)
South Korea	2010	Female	35 to 39	86% (75-93%)	14% (6.9-25%)	3.2 (2.2-4.3)	4.9% (2.8-7%)	15% (8.3-21%)	240 (130-350)	16000 (9500-22000)
South Korea	2010	Female	40 to 44	86% (75-93%)	14% (7.2-25%)	3 (1.9-4.2)	4.6% (2.6-6.6%)	14% (7.9-19%)	290 (170-410)	16000 (9400-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	2010	Female	45 to 49	84% (71-92%)	16% (8.1-29%)	2.9 (1.9-4.2)	4.6% (2.6-6.7%)	12% (7.4-17%)	370 (230-520)	18000 (10000-26000)
South Korea	2010	Female	50 to 54	82% (69-90%)	18% (9.6-31%)	2.8 (1.7-4)	4.1% (2.1-6.5%)	10% (6.2-15%)	390 (230-560)	16000 (8600-25000)
South Korea	2010	Female	55 to 59	79% (64-89%)	21% (11-36%)	2.5 (1.6-3.6)	3.2% (1.3-5.4%)	7.9% (4.6-12%)	300 (170-450)	10000 (4300-17000)
South Korea	2010	Female	60 to 64	76% (59-88%)	24% (12-41%)	2.2 (1.3-3.5)	2.4% (0.32-5%)	5.5% (2.3-9.5%)	270 (110-460)	7300 (900-15000)
South Korea	2010	Female	65 to 69	72% (54-86%)	28% (14-46%)	2 (1.2-2.9)	1.7% (-0.2-4.1%)	3.7% (0.94-6.9%)	290 (73-550)	6000 (-650-14000)
South Korea	2010	Female	70 to 74	70% (53-83%)	30% (17-47%)	1.8 (1-2.6)	1.1% (-1.1-3.6%)	2.1% (-0.76-5.5%)	270 (-100-710)	4300 (-4200-15000)
South Korea	2010	Female	75 to 79	69% (52-83%)	31% (17-48%)	1.6 (0.89-2.5)	0.34% (-1.8-3.1%)	0.7% (-2-4.2%)	130 (-360-780)	1400 (-7200-12000)
South Korea	2010	Female	80 to 84	67% (48-82%)	33% (18-52%)	1.5 (0.84-2.2)	-0.19% (-2.1-2.3%)	-0.19% (-2.6-2.9%)	-44 (-580-630)	-640 (-7200-7900)
South Korea	2010	Female	85 to 89	64% (45-79%)	36% (21-55%)	1.4 (0.78-2.1)	-0.29% (-2.1-2%)	-0.42% (-2.7-2.3%)	-89 (-580-500)	-610 (-4500-4200)
South Korea	2010	Female	90 to 94	64% (47-79%)	36% (21-53%)	1.3 (0.75-2.1)	-0.38% (-2.3-1.9%)	-0.61% (-3-2.1%)	-78 (-380-270)	-320 (-1900-1600)
South Korea	2010	Female	95 plus	65% (43-83%)	35% (17-57%)	1.3 (0.62-2.2)	-0.39% (-2.6-2.1%)	-0.76% (-3.4-2.4%)	-34 (-150-110)	-76 (-490-430)
South Korea	2010	Male	All Ages	91% (88-94%)	9.1% (6.3-12%)	2.3 (1.9-2.6)	7.4% (5.5-9.1%)	7% (4.8-9.3%)	11000 (6900-14000)	440000 (320000-560000)
South Korea	2010	Male	15 to 19	81% (68-91%)	19% (9.5-32%)	1.4 (0.82-2.1)	4.8% (3-6.8%)	8.7% (3.6-15%)	61 (25-110)	8600 (5400-12000)
South Korea	2010	Male	20 to 24	90% (82-95%)	9.9% (4.8-18%)	2.7 (1.8-3.8)	11% (7.9-15%)	16% (7.3-25%)	140 (68-230)	21000 (14000-29000)
South Korea	2010	Male	25 to 29	93% (87-97%)	7.1% (3.4-13%)	3.2 (2.1-4.5)	12% (8.3-16%)	18% (8.1-27%)	270 (120-410)	29000 (19000-40000)
South Korea	2010	Male	30 to 34	94% (88-98%)	5.8% (2.4-12%)	3.3 (2.2-4.6)	11% (7.1-15%)	17% (8.4-26%)	320 (150-490)	29000 (19000-41000)
South Korea	2010	Male	35 to 39	95% (90-98%)	5.1% (2.2-9.9%)	3 (2.1-4.1)	10% (7.1-14%)	16% (9.5-23%)	500 (280-720)	37000 (25000-49000)
South Korea	2010	Male	40 to 44	95% (90-98%)	4.7% (1.9-9.5%)	3 (2-4.1)	11% (7.9-14%)	16% (11-22%)	820 (540-1200)	48000 (34000-65000)
South Korea	2010	Male	45 to 49	95% (90-98%)	4.7% (2-10%)	3 (2-4.1)	11% (8.3-15%)	16% (12-21%)	1300 (920-1800)	64000 (46000-84000)
South Korea	2010	Male	50 to 54	95% (90-98%)	4.9% (2.2-10%)	2.9 (1.9-4.1)	10% (7.3-13%)	14% (9.9-18%)	1600 (1100-2100)	64000 (45000-86000)
South Korea	2010	Male	55 to 59	95% (89-98%)	5.5% (2.3-11%)	2.7 (1.8-3.8)	8.2% (5.7-11%)	11% (7.7-15%)	1200 (830-1700)	44000 (30000-59000)
South Korea	2010	Male	60 to 64	94% (87-97%)	6.1% (2.5-13%)	2.5 (1.6-3.8)	6.5% (4.1-9.4%)	8.7% (5.6-12%)	1100 (690-1600)	32000 (20000-47000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	2010	Male	65 to 69	93% (86-97%)	6.9% (2.8-14%)	2.3 (1.6-3.2)	5.1% (3.1-7.5%)	6.6% (4.1-9.5%)	1100 (670-1600)	26000 (16000-40000)
South Korea	2010	Male	70 to 74	92% (85-97%)	7.7% (3.3-15%)	2.1 (1.4-3)	3.9% (1.9-6.3%)	4.8% (2.5-7.7%)	1100 (530-1700)	20000 (10000-34000)
South Korea	2010	Male	75 to 79	92% (84-96%)	8.4% (3.7-16%)	1.9 (1.2-2.9)	2.6% (0.74-5.1%)	3.1% (0.95-5.9%)	640 (190-1200)	9900 (2700-19000)
South Korea	2010	Male	80 to 84	91% (83-96%)	9.4% (4.2-17%)	1.7 (1.1-2.5)	1.6% (-0.047-3.8%)	1.8% (-0.23-4.3%)	300 (-37-720)	3700 (-110-8500)
South Korea	2010	Male	85 to 89	90% (81-95%)	10% (5-19%)	1.6 (1-2.3)	1.1% (-0.47-3%)	1% (-0.83-3.3%)	110 (-93-360)	1100 (-500-3100)
South Korea	2010	Male	90 to 94	89% (81-95%)	11% (5-19%)	1.4 (0.88-2.1)	0.66% (-0.85-2.4%)	0.39% (-1.4-2.5%)	18 (-62-110)	190 (-240-720)
South Korea	2010	Male	95 plus	90% (79-96%)	10% (4.1-21%)	1.3 (0.71-2.2)	0.33% (-1.4-2.3%)	-0.24% (-2.2-2.1%)	-2.6 (-24-22)	15 (-62-100)
South Korea	2016	Female	All Ages	77% (71-82%)	23% (18-29%)	2.2 (1.7-2.9)	2.2% (0.89-3.9%)	1.4% (-0.68-3.9%)	2200 (-1100-6700)	120000 (46000-210000)
South Korea	2016	Female	15 to 19	67% (50-82%)	33% (18-50%)	1.7 (0.89-2.8)	2.4% (1.3-3.9%)	9.6% (3.2-17%)	26 (7.9-52)	3400 (1800-5600)
South Korea	2016	Female	20 to 24	80% (66-89%)	20% (11-34%)	3.2 (1.9-5)	5.4% (3.2-8%)	17% (6.9-28%)	85 (31-150)	10000 (6000-16000)
South Korea	2016	Female	25 to 29	84% (73-92%)	16% (8.3-27%)	3.5 (2.2-5)	5.8% (3.3-8.6%)	19% (7.6-29%)	120 (42-210)	12000 (6400-18000)
South Korea	2016	Female	30 to 34	86% (74-93%)	14% (6.8-26%)	3.4 (2.2-5)	5.1% (2.8-7.8%)	17% (7.9-26%)	160 (64-280)	13000 (6700-20000)
South Korea	2016	Female	35 to 39	86% (76-93%)	14% (6.8-24%)	3.1 (1.9-4.5)	4.4% (2.3-7%)	15% (7.7-21%)	180 (74-330)	13000 (6400-21000)
South Korea	2016	Female	40 to 44	86% (76-93%)	14% (7.2-24%)	2.9 (1.7-4.3)	4.2% (2.1-6.6%)	13% (7.3-19%)	250 (110-440)	15000 (7000-24000)
South Korea	2016	Female	45 to 49	84% (71-92%)	16% (7.9-29%)	2.8 (1.7-4.2)	4% (2-6.4%)	12% (6.8-17%)	320 (160-550)	16000 (7700-27000)
South Korea	2016	Female	50 to 54	82% (69-91%)	18% (9.3-31%)	2.6 (1.6-3.9)	3.7% (1.7-6.3%)	9.7% (5.8-14%)	370 (180-630)	15000 (6800-28000)
South Korea	2016	Female	55 to 59	79% (65-90%)	21% (10-35%)	2.4 (1.5-3.6)	2.9% (1-5.3%)	7.5% (4.4-11%)	360 (180-620)	12000 (4400-23000)
South Korea	2016	Female	60 to 64	76% (60-88%)	24% (12-40%)	2.2 (1.3-3.4)	2.1% (-0.027-4.8%)	5.2% (2-9.6%)	300 (99-580)	8000 (-110-18000)
South Korea	2016	Female	65 to 69	73% (55-86%)	27% (14-45%)	1.9 (1.1-2.8)	1.3% (-0.62-3.9%)	3.2% (0.57-6.6%)	260 (37-600)	5000 (-2300-15000)
South Korea	2016	Female	70 to 74	70% (54-83%)	30% (17-46%)	1.6 (0.9-2.5)	0.58% (-1.3-3.2%)	1.5% (-1.1-5%)	190 (-140-670)	2400 (-5300-13000)
South Korea	2016	Female	75 to 79	69% (51-83%)	31% (17-49%)	1.4 (0.76-2.3)	-0.064% (-1.9-2.5%)	0.2% (-2.3-3.7%)	43 (-500-740)	-290 (-9400-12000)
South Korea	2016	Female	80 to 84	67% (48-82%)	33% (18-52%)	1.4 (0.75-2.2)	-0.44% (-2.4-1.8%)	-0.51% (-3-2.3%)	-140 (-860-660)	-1900 (-10000-8000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Korea	2016	Female	85 to 89	64% (45-79%)	36% (21-55%)	1.3 (0.73-2.1)	-0.33% (-2.2-2.1%)	-0.48% (-2.8-2.4%)	-150 (-870-730)	-1000 (-7000-6200)
South Korea	2016	Female	90 to 94	64% (47-79%)	36% (21-53%)	1.4 (0.71-2.2)	-0.25% (-2.3-2.3%)	-0.46% (-2.9-2.6%)	-100 (-630-550)	-360 (-3300-3300)
South Korea	2016	Female	95 plus	65% (43-83%)	35% (17-57%)	1.4 (0.67-2.5)	-0.075% (-2.2-3%)	-0.38% (-3-3.4%)	-32 (-260-270)	-28 (-830-1100)
South Korea	2016	Male	All Ages	91% (88-94%)	8.7% (6-12%)	2.2 (1.7-2.8)	6.7% (4.8-8.9%)	6% (3.7-8.6%)	11000 (5800-18000)	440000 (290000-640000)
South Korea	2016	Male	15 to 19	81% (68-91%)	19% (9.3-32%)	1.3 (0.72-2.1)	4.5% (2.8-6.4%)	8.4% (3-14%)	45 (16-83)	7000 (4200-10000)
South Korea	2016	Male	20 to 24	90% (82-95%)	10% (4.8-18%)	2.5 (1.5-3.7)	11% (7.4-14%)	15% (6.6-24%)	130 (56-240)	21000 (14000-30000)
South Korea	2016	Male	25 to 29	93% (87-97%)	7.2% (3.4-13%)	3 (1.9-4.3)	12% (7.9-16%)	17% (7.6-27%)	210 (85-370)	25000 (16000-36000)
South Korea	2016	Male	30 to 34	94% (88-98%)	5.8% (2.4-12%)	3.1 (2-4.6)	10% (6.6-14%)	17% (7.8-26%)	270 (110-470)	26000 (16000-38000)
South Korea	2016	Male	35 to 39	95% (90-98%)	5.1% (2.2-10%)	2.8 (1.8-4.1)	9.8% (6.5-13%)	16% (8.6-22%)	390 (190-650)	30000 (19000-44000)
South Korea	2016	Male	40 to 44	95% (90-98%)	4.7% (1.9-9.5%)	2.7 (1.7-4.1)	10% (7-14%)	16% (9.8-21%)	710 (380-1200)	42000 (27000-64000)
South Korea	2016	Male	45 to 49	95% (90-98%)	4.7% (2.1-9.8%)	2.8 (1.8-4.1)	11% (7.6-15%)	16% (11-21%)	1200 (690-1900)	58000 (36000-86000)
South Korea	2016	Male	50 to 54	95% (90-98%)	5% (2.2-9.8%)	2.8 (1.7-4.2)	9.7% (6.7-13%)	14% (9.6-18%)	1500 (890-2400)	64000 (40000-97000)
South Korea	2016	Male	55 to 59	94% (89-98%)	5.6% (2.4-11%)	2.6 (1.6-3.7)	7.7% (5.1-11%)	11% (7.3-15%)	1500 (870-2400)	54000 (32000-84000)
South Korea	2016	Male	60 to 64	94% (87-97%)	6.2% (2.7-13%)	2.4 (1.4-3.7)	6.1% (3.7-9.1%)	8.4% (5.4-12%)	1300 (650-2100)	38000 (21000-62000)
South Korea	2016	Male	65 to 69	93% (86-97%)	6.9% (2.8-14%)	2.2 (1.4-3.3)	4.9% (2.7-7.5%)	6.5% (3.8-9.6%)	1200 (640-2000)	29000 (15000-48000)
South Korea	2016	Male	70 to 74	92% (85-96%)	7.8% (3.6-15%)	2.1 (1.3-3.2)	3.8% (1.7-6.6%)	4.7% (2.3-7.9%)	1100 (500-2100)	22000 (9200-41000)
South Korea	2016	Male	75 to 79	91% (84-96%)	8.6% (4-16%)	1.9 (1-2.9)	2.5% (0.69-5.3%)	3% (0.91-6.2%)	830 (200-1800)	13000 (3200-28000)
South Korea	2016	Male	80 to 84	90% (82-96%)	9.6% (4.5-18%)	1.6 (0.99-2.5)	1.5% (-0.2-3.7%)	1.6% (-0.3-4.1%)	400 (-73-1000)	4900 (-640-12000)
South Korea	2016	Male	85 to 89	89% (80-95%)	11% (5.1-20%)	1.4 (0.85-2.2)	0.88% (-0.55-2.7%)	0.76% (-0.96-3%)	130 (-180-510)	1400 (-920-4400)
South Korea	2016	Male	90 to 94	89% (81-95%)	11% (5-19%)	1.3 (0.74-2)	0.45% (-0.91-2.2%)	0.09% (-1.5-2.2%)	7.2 (-130-170)	240 (-490-1100)
South Korea	2016	Male	95 plus	89% (78-96%)	11% (4.1-22%)	1.1 (0.58-2)	0.17% (-1.3-2%)	-0.52% (-2.4-1.5%)	-10 (-46-30)	15 (-110-170)
Singapore	1990	Female	All Ages	63% (55-71%)	37% (29-45%)	0.51 (0.44-0.59)	-0.24% (-0.82-0.28%)	-1.1% (-2.2--0.14%)	-62 (-130--8)	-640 (-2200-720)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	1990	Female	15 to 19	50% (32-69%)	50% (31-68%)	0.39 (0.2-0.63)	0.67% (0.24-1.3%)	1.9% (0.0049-4.5%)	0.58 (0.0015-1.4)	73 (26-130)
Singapore	1990	Female	20 to 24	68% (50-83%)	32% (17-50%)	0.78 (0.48-1.2)	1.4% (0.43-2.5%)	3.6% (-0.12-7.7%)	1.9 (-0.064-4.1)	200 (62-350)
Singapore	1990	Female	25 to 29	76% (59-87%)	24% (13-41%)	0.84 (0.56-1.2)	1.2% (0.35-2.1%)	3.4% (-0.099-7%)	2.4 (-0.069-5)	220 (67-390)
Singapore	1990	Female	30 to 34	78% (61-90%)	22% (10-39%)	0.81 (0.53-1.1)	0.99% (0.23-1.9%)	3.1% (0.4-6%)	2.9 (0.38-6)	210 (50-380)
Singapore	1990	Female	35 to 39	78% (64-90%)	22% (10-36%)	0.76 (0.5-1.1)	0.86% (0.2-1.6%)	2.8% (0.9-5.1%)	3.4 (1.1-6.3)	180 (41-320)
Singapore	1990	Female	40 to 44	77% (63-89%)	23% (11-37%)	0.72 (0.42-1.1)	0.58% (-0.073-1.4%)	2.1% (0.45-3.9%)	3.1 (0.64-6)	110 (-14-250)
Singapore	1990	Female	45 to 49	75% (56-88%)	25% (12-44%)	0.68 (0.4-1)	0.087% (-0.66-0.9%)	0.87% (-0.58-2.6%)	1.3 (-0.89-3.9)	12 (-94-130)
Singapore	1990	Female	50 to 54	71% (53-85%)	29% (15-47%)	0.62 (0.36-0.97)	-0.52% (-1.5-0.44%)	-0.31% (-1.9-1.2%)	-0.77 (-4.9-3.1)	-86 (-240-72)
Singapore	1990	Female	55 to 59	67% (48-83%)	33% (17-52%)	0.57 (0.33-0.89)	-1.2% (-2.3--0.2%)	-1.3% (-3-0.22%)	-4.6 (-11-0.85)	-210 (-430-36)
Singapore	1990	Female	60 to 64	62% (42-79%)	38% (21-58%)	0.48 (0.26-0.81)	-1.7% (-3.1--0.61%)	-2% (-3.9--0.56%)	-9.8 (-19--2.7)	-320 (-590--110)
Singapore	1990	Female	65 to 69	58% (38-76%)	42% (24-62%)	0.37 (0.21-0.59)	-1.7% (-3--0.66%)	-2% (-3.7--0.67%)	-12 (-22--4.1)	-310 (-560--120)
Singapore	1990	Female	70 to 74	55% (36-72%)	45% (28-64%)	0.28 (0.15-0.48)	-1.5% (-2.8--0.56%)	-1.7% (-3.2--0.59%)	-13 (-25--4.5)	-270 (-500--100)
Singapore	1990	Female	75 to 79	53% (34-70%)	47% (30-66%)	0.25 (0.13-0.42)	-1.4% (-2.6--0.51%)	-1.6% (-3.1--0.55%)	-14 (-28-5)	-220 (-420--82)
Singapore	1990	Female	80 to 84	50% (31-68%)	50% (32-69%)	0.25 (0.13-0.39)	-1.3% (-2.3--0.46%)	-1.5% (-2.8--0.51%)	-12 (-22--4.1)	-140 (-250--50)
Singapore	1990	Female	85 to 89	46% (30-63%)	54% (37-70%)	0.25 (0.13-0.41)	-1.2% (-2.2--0.44%)	-1.3% (-2.6--0.46%)	-7.8 (-15--2.7)	-62 (-110--23)
Singapore	1990	Female	90 to 94	45% (29-63%)	55% (37-71%)	0.26 (0.13-0.46)	-1.1% (-2.1--0.36%)	-1.3% (-2.5--0.41%)	-3 (-5.7--0.93)	-16 (-31--5.5)
Singapore	1990	Female	95 plus	47% (26-68%)	53% (32-74%)	0.29 (0.13-0.53)	-1.1% (-2.1--0.29%)	-1.3% (-2.6--0.33%)	-0.83 (-1.7--0.21)	-2.9 (-5.6--0.78)
Singapore	1990	Male	All Ages	84% (79-89%)	16% (11-21%)	0.58 (0.5-0.67)	0.66% (-0.42-1.7%)	-0.24% (-1.7-1.2%)	-18 (-130-89)	2100 (-1300-5600)
Singapore	1990	Male	15 to 19	71% (54-85%)	29% (15-46%)	0.36 (0.21-0.57)	1.6% (0.66-2.8%)	2.7% (0.45-5.5%)	1.9 (0.34-4)	210 (85-370)
Singapore	1990	Male	20 to 24	85% (72-93%)	15% (7.2-28%)	0.74 (0.49-1.1)	3.7% (1.4-6.3%)	5.3% (0.91-10%)	6.2 (0.96-12)	590 (230-1000)
Singapore	1990	Male	25 to 29	89% (79-95%)	11% (5.1-21%)	0.89 (0.63-1.2)	3.5% (1.4-5.8%)	5% (0.55-9.6%)	7.5 (0.85-15)	700 (280-1200)
Singapore	1990	Male	30 to 34	91% (82-96%)	8.9% (3.6-18%)	0.92 (0.66-1.2)	2.7% (1.1-4.6%)	3.8% (0.22-7.6%)	6.4 (0.36-13)	590 (230-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	1990	Male	35 to 39	92% (85-97%)	7.6% (3.1-15%)	0.86 (0.6-1.2)	2.2% (0.84-3.8%)	2.9% (0.12-6%)	5.7 (0.25-12)	470 (180-810)
Singapore	1990	Male	40 to 44	93% (86-97%)	7.1% (2.8-14%)	0.84 (0.55-1.2)	1.6% (0.21-3.1%)	1.9% (-0.38-4.4%)	4.9 (-0.95-11)	340 (43-660)
Singapore	1990	Male	45 to 49	93% (84-97%)	7.2% (2.9-16%)	0.85 (0.55-1.3)	0.98% (-0.56-2.5%)	1% (-1.3-3.4%)	2.8 (-3.6-9.1)	170 (-96-440)
Singapore	1990	Male	50 to 54	92% (84-97%)	7.6% (3.3-16%)	0.82 (0.49-1.2)	0.19% (-1.5-1.8%)	-0.0035% (-2.2-2.2%)	-0.038 (-9.7-9.3)	41 (-320-390)
Singapore	1990	Male	55 to 59	91% (83-96%)	8.5% (3.5-17%)	0.75 (0.46-1.1)	-0.5% (-2.4-1.1%)	-0.79% (-3.2-1.3%)	-5.1 (-21-8.4)	-130 (-640-300)
Singapore	1990	Male	60 to 64	90% (80-96%)	9.5% (3.9-20%)	0.68 (0.41-1)	-0.87% (-2.6-0.56%)	-1.2% (-3.2-0.62%)	-10 (-30-5.5)	-260 (-770-170)
Singapore	1990	Male	65 to 69	89% (80-96%)	11% (4.4-20%)	0.61 (0.38-0.92)	-0.93% (-2.5-0.38%)	-1.2% (-3-0.38%)	-11 (-28-4)	-240 (-630-100)
Singapore	1990	Male	70 to 74	88% (78-95%)	12% (5.3-22%)	0.52 (0.31-0.82)	-0.78% (-2.1-0.31%)	-0.96% (-2.5-0.3%)	-9.8 (-26-3.1)	-170 (-470-70)
Singapore	1990	Male	75 to 79	86% (76-94%)	14% (6.1-24%)	0.45 (0.26-0.72)	-0.67% (-1.8-0.26%)	-0.84% (-2.1-0.22%)	-8.2 (-21-2.2)	-110 (-290-43)
Singapore	1990	Male	80 to 84	85% (73-93%)	15% (7.2-27%)	0.39 (0.23-0.6)	-0.6% (-1.5-0.2%)	-0.78% (-1.9-0.16%)	-5.5 (-14-1.1)	-55 (-140-18)
Singapore	1990	Male	85 to 89	83% (71-92%)	17% (8.3-29%)	0.36 (0.21-0.56)	-0.57% (-1.5-0.18%)	-0.75% (-1.8-0.14%)	-2.7 (-6.6-0.52)	-19 (-48-5.7)
Singapore	1990	Male	90 to 94	83% (70-92%)	17% (8.2-30%)	0.33 (0.18-0.55)	-0.57% (-1.5-0.15%)	-0.81% (-1.9-0.056%)	-0.83 (-2-0.055)	-3.7 (-9.8-0.98)
Singapore	1990	Male	95 plus	83% (67-93%)	17% (6.8-33%)	0.32 (0.15-0.56)	-0.71% (-1.8-0.09%)	-1.1% (-2.4--0.07%)	-0.19 (-0.44--0.013)	-0.54 (-1.4-0.07)
Singapore	1995	Female	All Ages	62% (53-70%)	38% (30-47%)	0.58 (0.49-0.67)	-0.21% (-0.74-0.28%)	-1% (-2.1--0.12%)	-64 (-130--7.2)	-600 (-2100-810)
Singapore	1995	Female	15 to 19	49% (31-67%)	51% (33-69%)	0.41 (0.22-0.68)	0.67% (0.28-1.2%)	2% (0.11-4.7%)	0.48 (0.028-1.1)	63 (27-110)
Singapore	1995	Female	20 to 24	66% (47-81%)	34% (19-53%)	0.84 (0.5-1.3)	1.3% (0.57-2.2%)	3.9% (0.3-8.1%)	1.4 (0.091-2.9)	170 (75-280)
Singapore	1995	Female	25 to 29	74% (56-86%)	26% (14-44%)	0.95 (0.62-1.3)	1.1% (0.46-1.9%)	3.8% (0.7-7.3%)	1.8 (0.35-3.7)	180 (77-310)
Singapore	1995	Female	30 to 34	77% (59-89%)	23% (11-41%)	0.92 (0.59-1.3)	0.95% (0.35-1.7%)	3.6% (1.2-6.4%)	2.5 (0.79-4.7)	190 (70-330)
Singapore	1995	Female	35 to 39	77% (61-89%)	23% (11-39%)	0.83 (0.54-1.2)	0.87% (0.25-1.6%)	3.3% (1.4-5.5%)	3.7 (1.5-6.5)	200 (62-340)
Singapore	1995	Female	40 to 44	76% (61-88%)	24% (12-39%)	0.79 (0.47-1.1)	0.66% (0.064-1.4%)	2.6% (1-4.6%)	3.9 (1.5-7.1)	150 (14-310)
Singapore	1995	Female	45 to 49	74% (53-87%)	26% (13-47%)	0.77 (0.47-1.2)	0.33% (-0.41-1.2%)	1.6% (0.016-3.4%)	3.2 (0.028-7.1)	69 (-89-240)
Singapore	1995	Female	50 to 54	70% (51-83%)	30% (17-49%)	0.7 (0.42-1.1)	-0.19% (-1.1-0.79%)	0.32% (-1.3-2%)	0.72 (-2.7-4.6)	-32 (-180-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	1995	Female	55 to 59	65% (47-81%)	35% (19-53%)	0.6 (0.34-0.95)	-0.84% (-1.9-0.084%)	-0.81% (-2.4-0.65%)	-2.8 (-8.6-2.4)	-160 (-360-16)
Singapore	1995	Female	60 to 64	60% (40-78%)	40% (22-60%)	0.51 (0.28-0.86)	-1.5% (-2.7--0.42%)	-1.7% (-3.4--0.3%)	-8.4 (-17--1.4)	-290 (-550--88)
Singapore	1995	Female	65 to 69	56% (36-74%)	44% (26-64%)	0.44 (0.25-0.68)	-1.6% (-2.9--0.62%)	-2% (-3.6--0.63%)	-13 (-25--4)	-340 (-660--130)
Singapore	1995	Female	70 to 74	54% (35-71%)	46% (29-65%)	0.38 (0.2-0.64)	-1.6% (-2.8--0.58%)	-1.8% (-3.2--0.6%)	-14 (-26--4.7)	-290 (-520--100)
Singapore	1995	Female	75 to 79	52% (34-70%)	48% (30-66%)	0.33 (0.18-0.57)	-1.5% (-2.8--0.58%)	-1.8% (-3.3--0.63%)	-16 (-30--5.7)	-250 (-460--96)
Singapore	1995	Female	80 to 84	49% (30-67%)	51% (33-70%)	0.29 (0.16-0.48)	-1.3% (-2.3--0.46%)	-1.5% (-2.8--0.48%)	-13 (-25--4.1)	-160 (-290--56)
Singapore	1995	Female	85 to 89	45% (28-63%)	55% (37-72%)	0.26 (0.13-0.44)	-1.1% (-2--0.38%)	-1.2% (-2.4--0.4%)	-8.7 (-17--2.7)	-74 (-140--26)
Singapore	1995	Female	90 to 94	44% (28-63%)	56% (37-72%)	0.25 (0.12-0.44)	-0.98% (-1.9--0.31%)	-1.1% (-2.2--0.31%)	-4.1 (-8--1.1)	-23 (-45--7.3)
Singapore	1995	Female	95 plus	46% (25-67%)	54% (33-75%)	0.25 (0.11-0.46)	-0.9% (-1.8--0.2%)	-1.1% (-2.2--0.21%)	-1.1 (-2.4--0.23)	-4.1 (-8.1--0.9)
Singapore	1995	Male	All Ages	85% (80-90%)	15% (10-20%)	0.67 (0.58-0.76)	-0.28% (-0.43-1.6%)	-0.28% (-1.7-1.1%)	-22 (-140-89)	2000 (-1300-5300)
Singapore	1995	Male	15 to 19	72% (55-85%)	28% (15-45%)	0.4 (0.22-0.63)	1.6% (0.66-2.7%)	2.9% (0.51-6%)	1.5 (0.27-3)	170 (70-290)
Singapore	1995	Male	20 to 24	85% (72-93%)	15% (7.1-28%)	0.83 (0.54-1.2)	3.6% (1.6-5.9%)	5.7% (1.1-11%)	4.8 (0.94-9.2)	490 (220-810)
Singapore	1995	Male	25 to 29	89% (80-95%)	11% (5-20%)	1 (0.7-1.4)	3.4% (1.6-5.2%)	5.6% (0.94-10%)	4.6 (0.75-8.5)	500 (230-770)
Singapore	1995	Male	30 to 34	91% (82-96%)	8.6% (3.5-18%)	1.1 (0.75-1.4)	2.8% (1.2-4.4%)	4.3% (0.64-8.2%)	5.4 (0.76-10)	540 (250-860)
Singapore	1995	Male	35 to 39	93% (85-97%)	7.4% (3-15%)	0.96 (0.68-1.3)	2.2% (0.78-3.7%)	3.2% (0.11-6.5%)	6 (0.22-13)	500 (190-830)
Singapore	1995	Male	40 to 44	93% (86-97%)	6.9% (2.6-14%)	0.94 (0.62-1.3)	1.7% (0.45-3.1%)	2.4% (0.066-4.8%)	5.9 (0.14-13)	400 (110-720)
Singapore	1995	Male	45 to 49	93% (86-97%)	7% (2.9-14%)	0.97 (0.64-1.4)	1.2% (-0.29-2.7%)	1.5% (-0.99-3.8%)	5.2 (-3.4-14)	300 (-73-680)
Singapore	1995	Male	50 to 54	93% (85-97%)	7.3% (3.1-15%)	0.93 (0.58-1.4)	0.55% (-0.98-2.1%)	0.57% (-1.7-2.9%)	2.1 (-6.1-11)	110 (-200-430)
Singapore	1995	Male	55 to 59	92% (85-97%)	8.1% (3.4-15%)	0.84 (0.52-1.3)	-0.23% (-1.9-1.3%)	-0.43% (-2.7-1.7%)	-2.5 (-16-9.8)	-58 (-480-340)
Singapore	1995	Male	60 to 64	91% (81-96%)	9.3% (4-19%)	0.77 (0.46-1.2)	-0.76% (-2.5-0.75%)	-1% (-3.2-0.87%)	-8.6 (-27-7.3)	-220 (-700-210)
Singapore	1995	Male	65 to 69	89% (79-95%)	11% (4.6-21%)	0.71 (0.46-1)	-0.94% (-2.5-0.42%)	-1.2% (-3.1-0.45%)	-12 (-33-4.9)	-280 (-740-130)
Singapore	1995	Male	70 to 74	88% (77-95%)	12% (5.5-23%)	0.63 (0.37-0.96)	-0.87% (-2.2-0.33%)	-1.1% (-2.7-0.36%)	-11 (-29-3.7)	-200 (-520-81)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	1995	Male	75 to 79	86% (75-94%)	14% (6.1-25%)	0.53 (0.31-0.85)	-0.82% (-2-0.25%)	-0.99% (-2.4-0.25%)	-10 (-25-2.7)	-140 (-360-46)
Singapore	1995	Male	80 to 84	85% (72-93%)	15% (6.9-28%)	0.45 (0.26-0.69)	-0.67% (-1.7-0.2%)	-0.84% (-2.1-0.18%)	-7.1 (-18-1.6)	-73 (-190-22)
Singapore	1995	Male	85 to 89	83% (71-92%)	17% (8.3-29%)	0.39 (0.23-0.61)	-0.52% (-1.4-0.21%)	-0.71% (-1.7-0.17%)	-3.6 (-9-0.88)	-25 (-69-10)
Singapore	1995	Male	90 to 94	83% (71-92%)	17% (8.2-29%)	0.34 (0.19-0.58)	-0.55% (-1.5-0.15%)	-0.77% (-1.9-0.089%)	-1.4 (-3.4-0.16)	-6.5 (-17-1.8)
Singapore	1995	Male	95 plus	83% (68-93%)	17% (6.5-32%)	0.31 (0.14-0.55)	-0.59% (-1.6-0.14%)	-0.91% (-2.2--0.0055%)	-0.32 (-0.76--0.0019)	-0.89 (-2.3-0.22)
Singapore	2000	Female	All Ages	61% (53-69%)	39% (31-47%)	0.65 (0.57-0.75)	-0.098% (-0.61-0.39%)	-0.86% (-1.9-0.04%)	-56 (-120-2.9)	-290 (-1800-1200)
Singapore	2000	Female	15 to 19	48% (30-67%)	52% (33-70%)	0.46 (0.24-0.79)	0.71% (0.26-1.3%)	2.2% (0.033-5%)	0.56 (0.0084-1.3)	71 (26-130)
Singapore	2000	Female	20 to 24	66% (47-81%)	34% (19-53%)	0.97 (0.57-1.5)	1.3% (0.59-2.1%)	4.5% (0.75-8.9%)	1.2 (0.2-2.4)	150 (69-250)
Singapore	2000	Female	25 to 29	73% (56-86%)	27% (14-44%)	1.1 (0.69-1.6)	1.2% (0.54-1.9%)	4.5% (1-8.4%)	1.7 (0.38-3.4)	180 (83-290)
Singapore	2000	Female	30 to 34	76% (58-88%)	24% (12-42%)	1 (0.66-1.5)	1.1% (0.47-1.8%)	4.1% (1.6-7.4%)	2.7 (1-4.8)	210 (92-350)
Singapore	2000	Female	35 to 39	77% (61-89%)	23% (11-39%)	0.95 (0.62-1.3)	0.93% (0.34-1.6%)	3.8% (1.7-6.1%)	3.8 (1.7-6.5)	210 (78-370)
Singapore	2000	Female	40 to 44	76% (59-88%)	24% (12-41%)	0.91 (0.57-1.3)	0.74% (0.097-1.5%)	3.1% (1.3-5.3%)	4.6 (1.9-8.3)	190 (26-380)
Singapore	2000	Female	45 to 49	73% (52-87%)	27% (13-48%)	0.88 (0.55-1.3)	0.56% (-0.17-1.5%)	2.3% (0.54-4.4%)	4.9 (1.1-9.9)	140 (-43-360)
Singapore	2000	Female	50 to 54	69% (50-83%)	31% (17-50%)	0.8 (0.49-1.2)	0.16% (-0.65-1.2%)	1.1% (-0.46-3%)	3.2 (-1.3-8.7)	36 (-160-260)
Singapore	2000	Female	55 to 59	65% (46-81%)	35% (19-54%)	0.7 (0.4-1.1)	-0.48% (-1.5-0.44%)	-0.19% (-1.7-1.4%)	-0.53 (-5-4.3)	-84 (-250-77)
Singapore	2000	Female	60 to 64	59% (39-77%)	41% (23-61%)	0.59 (0.32-0.97)	-1.2% (-2.3--0.14%)	-1.4% (-3-0.13%)	-6.3 (-14-0.63)	-240 (-480--28)
Singapore	2000	Female	65 to 69	55% (35-74%)	45% (26-65%)	0.49 (0.28-0.76)	-1.5% (-2.7--0.55%)	-1.8% (-3.5--0.51%)	-12 (-24--3.5)	-340 (-620--120)
Singapore	2000	Female	70 to 74	52% (32-70%)	48% (30-68%)	0.4 (0.21-0.69)	-1.4% (-2.6--0.52%)	-1.7% (-3.1--0.56%)	-14 (-26--4.6)	-300 (-530--110)
Singapore	2000	Female	75 to 79	49% (31-68%)	51% (32-69%)	0.34 (0.18-0.59)	-1.4% (-2.5--0.52%)	-1.6% (-3--0.56%)	-15 (-28--4.9)	-240 (-440--84)
Singapore	2000	Female	80 to 84	47% (28-65%)	53% (35-72%)	0.32 (0.17-0.52)	-1.2% (-2.2--0.42%)	-1.4% (-2.7--0.47%)	-14 (-25--4.3)	-160 (-290--59)
Singapore	2000	Female	85 to 89	44% (27-61%)	56% (39-73%)	0.3 (0.16-0.5)	-1% (-1.9--0.37%)	-1.2% (-2.3--0.38%)	-10 (-20--3.2)	-89 (-170--32)
Singapore	2000	Female	90 to 94	43% (27-61%)	57% (39-73%)	0.3 (0.15-0.51)	-0.91% (-1.8--0.25%)	-1.1% (-2.1--0.23%)	-5.1 (-10--1.2)	-30 (-59--9)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	2000	Female	95 plus	45% (24-66%)	55% (34-76%)	0.31 (0.14-0.57)	-0.89% (-1.8--0.16%)	-1.1% (-2.3--0.16%)	-1.5 (-3.2--0.22)	-5.6 (-11--1.1)
Singapore	2000	Male	All Ages	85% (81-90%)	15% (10-19%)	0.76 (0.67-0.87)	0.73% (-0.35-1.7%)	-0.16% (-1.6-1.2%)	-13 (-130-99)	2400 (-1200-5600)
Singapore	2000	Male	15 to 19	71% (54-85%)	29% (15-46%)	0.46 (0.25-0.73)	1.5% (0.74-2.5%)	3.1% (0.62-6.4%)	1.2 (0.23-2.5)	150 (72-260)
Singapore	2000	Male	20 to 24	84% (72-93%)	16% (7.3-28%)	0.96 (0.61-1.4)	3.9% (1.8-6.2%)	6.3% (1.6-11%)	4.7 (1.2-8.4)	470 (220-750)
Singapore	2000	Male	25 to 29	89% (80-95%)	11% (5.2-20%)	1.1 (0.77-1.6)	3.5% (1.8-5.4%)	6% (1.3-11%)	4.7 (1-8.7)	500 (260-770)
Singapore	2000	Male	30 to 34	91% (82-96%)	8.7% (3.6-18%)	1.2 (0.83-1.6)	2.9% (1.3-4.6%)	4.8% (0.79-9%)	5.1 (0.81-9.8)	500 (230-790)
Singapore	2000	Male	35 to 39	93% (85-97%)	7.5% (3-15%)	1.1 (0.78-1.5)	2.4% (0.94-4%)	3.6% (0.33-7.2%)	6.6 (0.56-14)	540 (220-920)
Singapore	2000	Male	40 to 44	93% (86-97%)	7% (2.7-14%)	1.1 (0.73-1.5)	1.9% (0.59-3.3%)	2.8% (0.21-5.5%)	7.6 (0.59-15)	510 (150-880)
Singapore	2000	Male	45 to 49	93% (85-97%)	7% (2.9-15%)	1.1 (0.74-1.5)	1.3% (-0.17-2.7%)	1.8% (-0.79-4.2%)	6.2 (-2.9-15)	360 (-51-750)
Singapore	2000	Male	50 to 54	93% (85-97%)	7.3% (3.1-15%)	1 (0.67-1.5)	0.62% (-0.98-2.1%)	0.69% (-1.8-2.9%)	3.2 (-8.3-14)	180 (-270-580)
Singapore	2000	Male	55 to 59	92% (84-97%)	8.1% (3.3-16%)	0.98 (0.61-1.4)	0.043% (-1.5-1.5%)	-0.083% (-2.3-2%)	-0.4 (-12-9.9)	10 (-370-350)
Singapore	2000	Male	60 to 64	91% (81-96%)	9.3% (3.9-19%)	0.89 (0.53-1.4)	-0.49% (-2.1-0.97%)	-0.73% (-2.9-1.2%)	-5.7 (-23-9.4)	-140 (-610-280)
Singapore	2000	Male	65 to 69	89% (79-95%)	11% (4.5-21%)	0.78 (0.49-1.1)	-0.78% (-2.3-0.55%)	-1% (-2.8-0.64%)	-10 (-29-6.7)	-230 (-660-160)
Singapore	2000	Male	70 to 74	87% (76-94%)	13% (5.9-24%)	0.65 (0.39-0.99)	-0.78% (-2.1-0.36%)	-0.98% (-2.6-0.42%)	-11 (-30-4.3)	-200 (-550-100)
Singapore	2000	Male	75 to 79	86% (74-93%)	14% (6.6-26%)	0.54 (0.32-0.87)	-0.76% (-1.9-0.24%)	-0.95% (-2.3-0.22%)	-10 (-25-2.4)	-140 (-350-45)
Singapore	2000	Male	80 to 84	84% (71-92%)	16% (7.6-29%)	0.48 (0.28-0.73)	-0.68% (-1.7-0.17%)	-0.88% (-2.2-0.16%)	-7.6 (-19-1.4)	-79 (-200-21)
Singapore	2000	Male	85 to 89	82% (68-91%)	18% (8.9-32%)	0.44 (0.25-0.7)	-0.55% (-1.4-0.21%)	-0.76% (-1.9-0.17%)	-4.3 (-11-0.98)	-30 (-79-12)
Singapore	2000	Male	90 to 94	81% (68-91%)	19% (9.2-32%)	0.42 (0.23-0.7)	-0.52% (-1.4-0.21%)	-0.79% (-2-0.1%)	-2 (-4.9-0.27)	-9.1 (-25-3.6)
Singapore	2000	Male	95 plus	82% (64-93%)	18% (7.3-36%)	0.41 (0.19-0.71)	-0.55% (-1.6-0.2%)	-0.95% (-2.2-0.063%)	-0.48 (-1.1-0.031)	-1.3 (-3.5-0.46)
Singapore	2005	Female	All Ages	61% (52-68%)	39% (32-48%)	0.64 (0.55-0.74)	-0.0034% (-0.45-0.45%)	-0.67% (-1.6-0.13%)	-49 (-110-8.8)	-12 (-1500-1400)
Singapore	2005	Female	15 to 19	47% (29-66%)	53% (34-71%)	0.44 (0.23-0.75)	0.61% (0.26-1.1%)	2.1% (0.22-4.7%)	0.44 (0.048-0.99)	65 (28-110)
Singapore	2005	Female	20 to 24	65% (46-80%)	35% (20-54%)	0.94 (0.54-1.5)	1.2% (0.61-2.1%)	4.3% (0.75-8.5%)	1.2 (0.2-2.4)	150 (67-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	2005	Female	25 to 29	72% (55-85%)	28% (15-45%)	1.1 (0.67-1.6)	1.1% (0.54-1.8%)	4.3% (1-8.2%)	1.4 (0.33-2.7)	160 (80-260)
Singapore	2005	Female	30 to 34	75% (57-88%)	25% (12-43%)	1 (0.64-1.5)	1% (0.46-1.7%)	4% (1.5-7.2%)	2.3 (0.85-4.2)	190 (89-320)
Singapore	2005	Female	35 to 39	76% (59-88%)	24% (12-41%)	0.93 (0.59-1.3)	0.85% (0.34-1.5%)	3.7% (1.6-6%)	2.9 (1.3-5)	180 (78-300)
Singapore	2005	Female	40 to 44	75% (58-87%)	25% (13-42%)	0.88 (0.54-1.3)	0.78% (0.18-1.5%)	3% (1.2-5.2%)	4.4 (1.7-7.7)	200 (47-370)
Singapore	2005	Female	45 to 49	72% (52-86%)	28% (14-48%)	0.85 (0.53-1.3)	0.61% (-0.061-1.4%)	2.3% (0.66-4.3%)	5.1 (1.4-9.6)	170 (-16-380)
Singapore	2005	Female	50 to 54	68% (49-83%)	32% (17-51%)	0.77 (0.48-1.1)	0.38% (-0.34-1.2%)	1.5% (0.016-3.1%)	4.6 (0.045-9.9)	100 (-90-330)
Singapore	2005	Female	55 to 59	64% (44-81%)	36% (19-56%)	0.68 (0.39-1.1)	-0.16% (-1-0.68%)	0.27% (-1.2-1.7%)	1.1 (-4.6-6.7)	-42 (-260-170)
Singapore	2005	Female	60 to 64	58% (37-77%)	42% (23-63%)	0.57 (0.31-0.93)	-0.77% (-1.7-0.062%)	-0.86% (-2.4-0.43%)	-3.5 (-9.5-1.7)	-150 (-350-11)
Singapore	2005	Female	65 to 69	54% (33-74%)	46% (26-67%)	0.47 (0.26-0.73)	-1.1% (-2.1--0.32%)	-1.4% (-2.8--0.24%)	-8 (-16--1.5)	-240 (-460--68)
Singapore	2005	Female	70 to 74	51% (32-69%)	49% (31-68%)	0.38 (0.2-0.63)	-1.1% (-2.1--0.41%)	-1.4% (-2.6--0.4%)	-11 (-22--3.1)	-250 (-450--86)
Singapore	2005	Female	75 to 79	49% (31-69%)	51% (31-69%)	0.33 (0.18-0.56)	-1.2% (-2.1--0.42%)	-1.4% (-2.6--0.44%)	-14 (-27--4.7)	-230 (-440--83)
Singapore	2005	Female	80 to 84	46% (28-65%)	54% (35-72%)	0.3 (0.16-0.48)	-0.99% (-1.8--0.36%)	-1.2% (-2.3--0.38%)	-13 (-24--4.1)	-160 (-300--57)
Singapore	2005	Female	85 to 89	42% (26-59%)	58% (41-74%)	0.29 (0.15-0.47)	-0.91% (-1.7--0.3%)	-1.1% (-2.1--0.34%)	-11 (-22--3.5)	-96 (-180--32)
Singapore	2005	Female	90 to 94	42% (26-59%)	58% (41-74%)	0.29 (0.15-0.48)	-0.87% (-1.7--0.24%)	-1% (-2.1--0.24%)	-7.6 (-15--1.8)	-43 (-84--12)
Singapore	2005	Female	95 plus	44% (24-65%)	56% (35-76%)	0.3 (0.13-0.57)	-0.87% (-1.8--0.15%)	-1.1% (-2.2--0.15%)	-3.1 (-6.4--0.41)	-11 (-23--1.9)
Singapore	2005	Male	All Ages	86% (81-90%)	14% (10-19%)	0.76 (0.66-0.87)	0.61% (-0.4-1.6%)	-0.23% (-1.6-1.1%)	-20 (-140-94)	2100 (-1300-5400)
Singapore	2005	Male	15 to 19	70% (54-85%)	30% (15-46%)	0.44 (0.24-0.7)	1.4% (0.71-2.3%)	3% (0.6-6.1%)	1.1 (0.22-2.3)	150 (76-250)
Singapore	2005	Male	20 to 24	84% (72-92%)	16% (7.6-28%)	0.92 (0.58-1.4)	3.6% (1.8-5.7%)	6.1% (1.4-11%)	4.1 (0.9-7.4)	430 (210-670)
Singapore	2005	Male	25 to 29	89% (80-95%)	11% (5.3-20%)	1.1 (0.75-1.6)	3.4% (1.8-5.1%)	6% (1.3-11%)	3.6 (0.82-6.8)	410 (210-630)
Singapore	2005	Male	30 to 34	91% (82-96%)	8.8% (3.5-18%)	1.2 (0.81-1.6)	2.8% (1.3-4.4%)	4.8% (0.84-9.2%)	4.1 (0.69-8.1)	440 (220-690)
Singapore	2005	Male	35 to 39	92% (85-97%)	7.7% (3.4-15%)	1.1 (0.74-1.5)	2.1% (0.89-3.6%)	3.3% (0.13-6.8%)	4.5 (0.18-9.5)	420 (180-690)
Singapore	2005	Male	40 to 44	93% (85-97%)	7.2% (2.9-15%)	1 (0.7-1.4)	1.6% (0.38-3%)	2.3% (-0.36-5.2%)	5.7 (-0.92-13)	430 (98-790)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	2005	Male	45 to 49	93% (84-97%)	7.3% (3.1-16%)	1.1 (0.73-1.5)	1.2% (-0.25-2.5%)	1.4% (-1.2-3.8%)	5.4 (-4.5-14)	360 (-76-750)
Singapore	2005	Male	50 to 54	92% (84-97%)	7.6% (3.3-16%)	1 (0.67-1.5)	0.56% (-0.98-1.9%)	0.5% (-1.9-2.7%)	2.7 (-11-15)	190 (-330-650)
Singapore	2005	Male	55 to 59	92% (84-96%)	8.4% (3.6-16%)	0.93 (0.59-1.4)	-0.029% (-1.6-1.4%)	-0.26% (-2.6-1.8%)	-1.8 (-19-13)	-11 (-550-480)
Singapore	2005	Male	60 to 64	90% (81-96%)	9.6% (4.2-19%)	0.85 (0.51-1.3)	-0.37% (-1.8-0.96%)	-0.62% (-2.8-1.2%)	-4.2 (-18-8.2)	-95 (-460-250)
Singapore	2005	Male	65 to 69	89% (78-95%)	11% (4.6-22%)	0.78 (0.5-1.1)	-0.61% (-1.9-0.6%)	-0.84% (-2.6-0.71%)	-8.4 (-26-7.3)	-180 (-600-190)
Singapore	2005	Male	70 to 74	87% (75-94%)	13% (5.7-25%)	0.69 (0.41-1.1)	-0.61% (-1.8-0.48%)	-0.8% (-2.3-0.55%)	-9.1 (-26-6.2)	-160 (-480-130)
Singapore	2005	Male	75 to 79	85% (73-94%)	15% (6.5-27%)	0.58 (0.34-0.92)	-0.61% (-1.7-0.35%)	-0.81% (-2.1-0.36%)	-9.4 (-25-4.2)	-130 (-350-73)
Singapore	2005	Male	80 to 84	84% (70-92%)	16% (7.9-30%)	0.49 (0.29-0.73)	-0.58% (-1.5-0.21%)	-0.79% (-2-0.21%)	-8.4 (-21-2.2)	-86 (-230-30)
Singapore	2005	Male	85 to 89	82% (69-91%)	18% (8.8-31%)	0.42 (0.24-0.66)	-0.54% (-1.4-0.17%)	-0.75% (-1.8-0.14%)	-5.8 (-14-1.1)	-40 (-100-13)
Singapore	2005	Male	90 to 94	81% (68-91%)	19% (8.9-32%)	0.36 (0.2-0.59)	-0.54% (-1.4-0.16%)	-0.78% (-1.9-0.11%)	-3.3 (-7.9-0.48)	-15 (-39-4.5)
Singapore	2005	Male	95 plus	82% (65-93%)	18% (7-35%)	0.32 (0.15-0.57)	-0.49% (-1.4-0.19%)	-0.77% (-1.9-0.097%)	-1 (-2.5-0.13)	-2.8 (-8-1.1)
Singapore	2010	Female	All Ages	61% (53-68%)	39% (32-47%)	0.59 (0.51-0.68)	0.053% (-0.34-0.43%)	-0.4% (-1.2-0.29%)	-30 (-90-22)	180 (-1100-1500)
Singapore	2010	Female	15 to 19	47% (29-65%)	53% (35-71%)	0.4 (0.2-0.68)	0.55% (0.26-0.94%)	2.1% (0.16-4.6%)	0.36 (0.029-0.78)	63 (30-100)
Singapore	2010	Female	20 to 24	65% (46-80%)	35% (20-54%)	0.85 (0.51-1.3)	1.1% (0.55-1.9%)	3.9% (0.51-7.9%)	1 (0.13-2.2)	140 (66-240)
Singapore	2010	Female	25 to 29	73% (55-86%)	27% (14-45%)	0.98 (0.61-1.4)	0.97% (0.49-1.5%)	4% (1-7.2%)	1.1 (0.27-2.1)	150 (74-230)
Singapore	2010	Female	30 to 34	76% (57-88%)	24% (12-43%)	0.95 (0.6-1.4)	0.83% (0.41-1.4%)	3.7% (1.5-6.5%)	1.6 (0.64-2.8)	150 (73-240)
Singapore	2010	Female	35 to 39	76% (60-88%)	24% (12-40%)	0.85 (0.54-1.2)	0.76% (0.32-1.3%)	3.4% (1.5-5.6%)	2.6 (1.1-4.4)	160 (72-270)
Singapore	2010	Female	40 to 44	75% (58-87%)	25% (13-42%)	0.8 (0.49-1.1)	0.64% (0.14-1.3%)	2.9% (1.2-5%)	3.3 (1.3-5.7)	150 (33-280)
Singapore	2010	Female	45 to 49	72% (53-86%)	28% (14-47%)	0.79 (0.5-1.2)	0.54% (-0.017-1.2%)	2.3% (0.74-4%)	4.4 (1.4-8)	140 (-4.5-320)
Singapore	2010	Female	50 to 54	68% (50-82%)	32% (18-50%)	0.72 (0.45-1.1)	0.41% (-0.22-1.1%)	1.6% (0.29-3.2%)	5.1 (0.92-10)	120 (-65-330)
Singapore	2010	Female	55 to 59	63% (44-81%)	37% (19-56%)	0.62 (0.37-0.94)	0.065% (-0.63-0.75%)	0.77% (-0.45-2%)	3.2 (-1.9-8.5)	18 (-180-210)
Singapore	2010	Female	60 to 64	58% (37-76%)	42% (24-63%)	0.52 (0.29-0.83)	-0.39% (-1.2-0.28%)	-0.18% (-1.4-0.96%)	-0.93 (-7.1-4.9)	-110 (-320-72)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	2010	Female	65 to 69	54% (34-73%)	46% (27-66%)	0.44 (0.24-0.68)	-0.67% (-1.5--0.053%)	-0.7% (-1.8-0.22%)	-3.8 (-10-1.2)	-140 (-320--9.8)
Singapore	2010	Female	70 to 74	51% (32-69%)	49% (31-68%)	0.37 (0.19-0.63)	-0.79% (-1.6--0.18%)	-0.9% (-2--0.052%)	-7 (-16--0.4)	-180 (-350--40)
Singapore	2010	Female	75 to 79	49% (31-68%)	51% (32-69%)	0.32 (0.17-0.54)	-0.9% (-1.7--0.25%)	-1.1% (-2.1--0.22%)	-11 (-22--2.4)	-190 (-370--54)
Singapore	2010	Female	80 to 84	46% (27-65%)	54% (35-73%)	0.29 (0.15-0.46)	-0.8% (-1.5--0.24%)	-0.97% (-1.9--0.24%)	-12 (-23--2.8)	-160 (-300--49)
Singapore	2010	Female	85 to 89	43% (26-61%)	57% (39-74%)	0.26 (0.13-0.43)	-0.73% (-1.5--0.22%)	-0.86% (-1.8--0.19%)	-11 (-23--2.4)	-100 (-200--30)
Singapore	2010	Female	90 to 94	42% (26-60%)	58% (40-74%)	0.25 (0.12-0.42)	-0.65% (-1.3--0.16%)	-0.77% (-1.6--0.13%)	-5 (-11--0.89)	-33 (-66--7.7)
Singapore	2010	Female	95 plus	44% (24-66%)	56% (34-76%)	0.25 (0.11-0.47)	-0.65% (-1.3--0.11%)	-0.8% (-1.7--0.075%)	-2.2 (-4.8--0.21)	-8.9 (-18--1.5)
Singapore	2010	Male	All Ages	86% (81-90%)	14% (10-19%)	0.7 (0.61-0.79)	0.56% (-0.35-1.4%)	-0.17% (-1.5-1.1%)	-15 (-130-97)	2100 (-1400-5400)
Singapore	2010	Male	15 to 19	70% (53-84%)	30% (16-47%)	0.39 (0.22-0.63)	1.3% (0.68-2%)	2.7% (0.53-5.6%)	0.91 (0.18-1.9)	150 (76-240)
Singapore	2010	Male	20 to 24	84% (71-92%)	16% (7.5-29%)	0.83 (0.52-1.2)	3.2% (1.5-5.1%)	5.5% (1-10%)	3.2 (0.57-6.2)	390 (180-620)
Singapore	2010	Male	25 to 29	89% (79-95%)	11% (5.4-21%)	1 (0.67-1.4)	3.1% (1.7-4.8%)	5.6% (0.91-11%)	3.2 (0.52-6.1)	400 (220-620)
Singapore	2010	Male	30 to 34	91% (81-96%)	9.2% (3.7-19%)	1 (0.72-1.5)	2.5% (1.3-3.9%)	4.3% (0.46-8.5%)	3 (0.31-5.9)	380 (190-580)
Singapore	2010	Male	35 to 39	92% (84-97%)	8% (3.3-16%)	0.97 (0.65-1.3)	2% (0.85-3.2%)	3% (-0.078-6.2%)	3.5 (-0.086-7.4)	370 (160-610)
Singapore	2010	Male	40 to 44	93% (85-97%)	7.5% (2.8-15%)	0.94 (0.62-1.3)	1.5% (0.3-2.7%)	1.9% (-0.81-4.7%)	3.6 (-1.4-8.9)	330 (72-600)
Singapore	2010	Male	45 to 49	92% (83-97%)	7.6% (3.1-17%)	0.96 (0.65-1.3)	1% (-0.33-2.3%)	1.1% (-1.6-3.6%)	3.8 (-5.5-13)	300 (-100-670)
Singapore	2010	Male	50 to 54	92% (84-97%)	7.9% (3.3-16%)	0.92 (0.61-1.3)	0.51% (-0.89-1.8%)	0.36% (-2-2.5%)	2.1 (-11-15)	190 (-340-680)
Singapore	2010	Male	55 to 59	91% (83-96%)	8.7% (3.6-17%)	0.84 (0.55-1.2)	0.16% (-1.2-1.4%)	-0.014% (-2.1-1.9%)	-0.054 (-16-14)	61 (-490-540)
Singapore	2010	Male	60 to 64	90% (79-96%)	10% (4.2-21%)	0.76 (0.46-1.1)	-0.17% (-1.5-0.98%)	-0.38% (-2.4-1.3%)	-3.4 (-21-12)	-62 (-560-380)
Singapore	2010	Male	65 to 69	89% (77-95%)	11% (4.7-23%)	0.69 (0.44-1)	-0.39% (-1.6-0.72%)	-0.58% (-2.2-0.87%)	-5.2 (-20-8.3)	-110 (-460-210)
Singapore	2010	Male	70 to 74	87% (75-94%)	13% (5.7-25%)	0.59 (0.35-0.89)	-0.42% (-1.5-0.51%)	-0.58% (-2-0.61%)	-6.8 (-22-7.2)	-120 (-410-150)
Singapore	2010	Male	75 to 79	85% (74-93%)	15% (6.9-26%)	0.5 (0.3-0.79)	-0.47% (-1.4-0.39%)	-0.63% (-1.8-0.43%)	-8.2 (-23-5.8)	-110 (-330-92)
Singapore	2010	Male	80 to 84	84% (71-92%)	16% (7.6-29%)	0.43 (0.26-0.66)	-0.42% (-1.2-0.26%)	-0.61% (-1.6-0.26%)	-6.9 (-19-3)	-71 (-200-42)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	2010	Male	85 to 89	82% (69-91%)	18% (9.2-31%)	0.38 (0.22-0.6)	-0.38% (-1.1-0.24%)	-0.57% (-1.5-0.23%)	-5.3 (-14-2.1)	-36 (-100-22)
Singapore	2010	Male	90 to 94	81% (68-91%)	19% (8.9-32%)	0.35 (0.19-0.57)	-0.35% (-1.1-0.23%)	-0.58% (-1.5-0.17%)	-2.2 (-5.6-0.61)	-9.5 (-29-6.4)
Singapore	2010	Male	95 plus	82% (65-93%)	18% (7.1-35%)	0.33 (0.16-0.58)	-0.3% (-1.1-0.29%)	-0.59% (-1.6-0.18%)	-0.68 (-1.8-0.22)	-1.7 (-5.8-1.6)
Singapore	2016	Female	All Ages	62% (53-69%)	38% (31-47%)	0.59 (0.46-0.74)	0.08% (-0.31-0.46%)	-0.28% (-0.97-0.34%)	-25 (-90-32)	290 (-1100-1700)
Singapore	2016	Female	15 to 19	47% (30-66%)	53% (34-70%)	0.4 (0.2-0.71)	0.55% (0.26-0.93%)	2% (0.15-4.6%)	0.35 (0.03-0.83)	63 (29-110)
Singapore	2016	Female	20 to 24	65% (47-81%)	35% (19-53%)	0.86 (0.49-1.4)	1% (0.54-1.7%)	3.9% (0.48-8.2%)	0.82 (0.11-1.7)	140 (71-220)
Singapore	2016	Female	25 to 29	73% (55-86%)	27% (14-45%)	0.96 (0.59-1.5)	0.88% (0.47-1.4%)	4% (0.99-7.3%)	0.81 (0.2-1.6)	120 (65-190)
Singapore	2016	Female	30 to 34	76% (57-88%)	24% (12-43%)	0.94 (0.57-1.4)	0.79% (0.34-1.4%)	3.8% (1.4-6.8%)	1.5 (0.52-2.9)	140 (60-240)
Singapore	2016	Female	35 to 39	77% (62-89%)	23% (11-38%)	0.86 (0.54-1.3)	0.65% (0.24-1.1%)	3.5% (1.6-5.8%)	2 (0.87-3.7)	130 (51-230)
Singapore	2016	Female	40 to 44	76% (61-88%)	24% (12-39%)	0.82 (0.47-1.2)	0.65% (0.15-1.3%)	3.2% (1.3-5.5%)	3.2 (1.2-6)	150 (32-290)
Singapore	2016	Female	45 to 49	73% (54-87%)	27% (13-46%)	0.79 (0.48-1.2)	0.57% (0.014-1.3%)	2.5% (0.88-4.4%)	4.5 (1.5-8.6)	150 (3.5-350)
Singapore	2016	Female	50 to 54	69% (51-84%)	31% (16-49%)	0.72 (0.41-1.1)	0.47% (-0.13-1.2%)	1.9% (0.51-3.5%)	5.5 (1.4-11)	140 (-39-360)
Singapore	2016	Female	55 to 59	65% (46-82%)	35% (18-54%)	0.63 (0.36-1)	0.18% (-0.44-0.88%)	1% (-0.2-2.4%)	4.4 (-0.85-11)	57 (-140-270)
Singapore	2016	Female	60 to 64	59% (38-77%)	41% (23-62%)	0.53 (0.29-0.9)	-0.21% (-0.88-0.39%)	0.16% (-0.98-1.3%)	0.94 (-5.3-7.8)	-70 (-290-130)
Singapore	2016	Female	65 to 69	55% (34-74%)	45% (26-66%)	0.43 (0.24-0.69)	-0.49% (-1.2-0.083%)	-0.43% (-1.6-0.46%)	-2.3 (-8.3-2.4)	-100 (-260-19)
Singapore	2016	Female	70 to 74	51% (33-69%)	49% (31-67%)	0.34 (0.17-0.59)	-0.57% (-1.2--0.063%)	-0.58% (-1.5-0.14%)	-4.8 (-13-1.3)	-140 (-310--14)
Singapore	2016	Female	75 to 79	49% (31-68%)	51% (32-69%)	0.29 (0.15-0.51)	-0.67% (-1.4--0.13%)	-0.77% (-1.7--0.07%)	-8.1 (-19--0.75)	-150 (-310--28)
Singapore	2016	Female	80 to 84	46% (28-65%)	54% (35-72%)	0.28 (0.14-0.45)	-0.65% (-1.3--0.19%)	-0.78% (-1.6--0.14%)	-11 (-24--2.1)	-170 (-330--47)
Singapore	2016	Female	85 to 89	43% (26-61%)	57% (39-74%)	0.26 (0.13-0.46)	-0.59% (-1.2--0.14%)	-0.68% (-1.5--0.086%)	-13 (-28--1.5)	-120 (-250--27)
Singapore	2016	Female	90 to 94	42% (27-60%)	58% (40-73%)	0.27 (0.13-0.47)	-0.54% (-1.1--0.071%)	-0.62% (-1.4--0.0053%)	-7.1 (-16--0.056)	-48 (-99--6.9)
Singapore	2016	Female	95 plus	44% (24-65%)	56% (35-76%)	0.28 (0.12-0.54)	-0.54% (-1.1--0.035%)	-0.66% (-1.4--0.015%)	-2.9 (-6.2--0.068)	-12 (-25--0.68)
Singapore	2016	Male	All Ages	85% (80-90%)	15% (10-20%)	0.68 (0.52-0.85)	0.59% (-0.27-1.4%)	-0.077% (-1.2-1%)	-8 (-130-110)	2300 (-1000-5600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Singapore	2016	Male	15 to 19	69% (52-84%)	31% (16-48%)	0.38 (0.2-0.62)	1.2% (0.63-1.9%)	2.5% (0.39-5.6%)	0.79 (0.12-1.8)	140 (71-220)
Singapore	2016	Male	20 to 24	83% (70-92%)	17% (7.8-30%)	0.8 (0.48-1.2)	3% (1.5-4.8%)	5.1% (0.51-10%)	2.9 (0.28-6.2)	380 (190-620)
Singapore	2016	Male	25 to 29	88% (79-94%)	12% (5.5-21%)	0.98 (0.6-1.4)	3% (1.6-4.6%)	5.6% (0.62-11%)	2.6 (0.28-5.2)	360 (190-550)
Singapore	2016	Male	30 to 34	91% (81-96%)	9.5% (3.7-19%)	1 (0.65-1.5)	2.5% (1.3-3.8%)	4.4% (0.21-8.7%)	2.9 (0.14-6.2)	370 (190-590)
Singapore	2016	Male	35 to 39	92% (84-97%)	8.2% (3.3-16%)	0.93 (0.58-1.4)	1.9% (0.86-3.1%)	3% (-0.018-6.5%)	3 (-0.02-6.8)	350 (160-570)
Singapore	2016	Male	40 to 44	92% (85-97%)	7.7% (2.8-15%)	0.91 (0.56-1.4)	1.5% (0.36-2.7%)	2% (-0.73-5%)	3.4 (-1.2-8.7)	310 (80-580)
Singapore	2016	Male	45 to 49	92% (84-97%)	7.8% (3.2-16%)	0.93 (0.59-1.3)	1.2% (-0.0013-2.3%)	1.6% (-1-4.1%)	4.4 (-2.9-12)	330 (-0.38-660)
Singapore	2016	Male	50 to 54	92% (84-96%)	8.2% (3.6-16%)	0.89 (0.54-1.3)	0.73% (-0.53-1.9%)	0.76% (-1.4-2.9%)	3.7 (-8.2-15)	260 (-200-690)
Singapore	2016	Male	55 to 59	91% (82-96%)	9.2% (3.8-18%)	0.8 (0.48-1.2)	0.29% (-0.94-1.5%)	0.18% (-1.7-2.1%)	1.4 (-14-16)	120 (-390-610)
Singapore	2016	Male	60 to 64	90% (79-96%)	10% (4.2-21%)	0.73 (0.42-1.2)	-0.013% (-1.2-1.1%)	-0.18% (-1.9-1.4%)	-1.9 (-20-15)	-6.8 (-520-470)
Singapore	2016	Male	65 to 69	88% (77-95%)	12% (5-23%)	0.67 (0.41-1)	-0.28% (-1.4-0.78%)	-0.45% (-2-1%)	-4.2 (-20-9.1)	-82 (-440-220)
Singapore	2016	Male	70 to 74	87% (75-94%)	13% (6.1-25%)	0.59 (0.33-0.92)	-0.26% (-1.2-0.59%)	-0.39% (-1.7-0.76%)	-4.9 (-22-9.2)	-82 (-420-190)
Singapore	2016	Male	75 to 79	85% (73-93%)	15% (7-27%)	0.5 (0.28-0.81)	-0.35% (-1.2-0.4%)	-0.5% (-1.6-0.46%)	-6.6 (-22-6.2)	-87 (-300-100)
Singapore	2016	Male	80 to 84	83% (70-92%)	17% (8.1-30%)	0.42 (0.24-0.66)	-0.29% (-0.98-0.29%)	-0.46% (-1.3-0.33%)	-6.8 (-20-4.7)	-66 (-210-63)
Singapore	2016	Male	85 to 89	81% (68-91%)	19% (9.4-32%)	0.36 (0.2-0.59)	-0.24% (-0.87-0.3%)	-0.41% (-1.2-0.29%)	-5.2 (-16-3.6)	-32 (-120-40)
Singapore	2016	Male	90 to 94	81% (68-91%)	19% (9.1-32%)	0.32 (0.17-0.56)	-0.2% (-0.81-0.28%)	-0.41% (-1.2-0.21%)	-2.8 (-8-1.4)	-10 (-41-15)
Singapore	2016	Male	95 plus	82% (65-93%)	18% (7.3-35%)	0.29 (0.14-0.54)	-0.14% (-0.77-0.36%)	-0.41% (-1.3-0.21%)	-0.76 (-2.3-0.38)	-1.3 (-7.1-3.4)
Australia	1990	Female	All Ages	84% (81-86%)	16% (14-19%)	2.2 (1.9-2.5)	2.4% (1.2-3.7%)	1.4% (-1.5-4%)	750 (-820-2200)	48000 (23000-73000)
Australia	1990	Female	15 to 19	87% (81-92%)	13% (7.7-19%)	2.2 (1.7-2.8)	3.3% (2-4.7%)	11% (5.2-16%)	27 (13-41)	2700 (1600-3900)
Australia	1990	Female	20 to 24	92% (88-95%)	8.2% (5.5-12%)	3 (2.3-3.6)	4.5% (3.1-6%)	12% (7.1-17%)	37 (21-52)	4300 (2900-5700)
Australia	1990	Female	25 to 29	93% (90-95%)	7.3% (5.1-10%)	2.5 (2.1-2.9)	3.6% (2.5-4.7%)	10% (6-14%)	33 (19-47)	3700 (2500-4800)
Australia	1990	Female	30 to 34	93% (90-95%)	7.1% (5-10%)	2.5 (2.1-3)	3.2% (2.2-4.1%)	10% (6.4-14%)	38 (25-52)	3400 (2400-4400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	1990	Female	35 to 39	92% (89-95%)	7.8% (5.4-11%)	3.1 (2.6-3.6)	3.8% (2.8-4.8%)	12% (9.2-15%)	60 (46-74)	4200 (3200-5200)
Australia	1990	Female	40 to 44	91% (89-94%)	8.6% (6.1-11%)	3.3 (2.8-3.9)	4.4% (3.3-5.5%)	13% (10-15%)	90 (72-110)	5200 (4100-6300)
Australia	1990	Female	45 to 49	90% (87-93%)	9.5% (6.7-13%)	3.3 (2.8-3.8)	4.7% (3.6-6%)	11% (9-14%)	110 (83-130)	5100 (3900-6200)
Australia	1990	Female	50 to 54	87% (84-91%)	13% (9.5-16%)	3.2 (2.7-3.7)	4.9% (3.5-6.4%)	9.8% (7.2-12%)	130 (92-160)	5200 (3800-6500)
Australia	1990	Female	55 to 59	83% (78-87%)	17% (13-22%)	3.1 (2.6-3.6)	4.3% (2.8-6%)	7.7% (5.1-10%)	140 (91-180)	4800 (3200-6500)
Australia	1990	Female	60 to 64	79% (74-84%)	21% (16-26%)	2.9 (2.4-3.5)	3.5% (1.7-5.5%)	5.4% (2.7-8.5%)	160 (81-250)	4900 (2400-7600)
Australia	1990	Female	65 to 69	77% (71-82%)	23% (18-29%)	2.7 (2.3-3.2)	2.4% (0.39-4.5%)	3.4% (0.37-6.5%)	160 (17-300)	3900 (600-7200)
Australia	1990	Female	70 to 74	73% (66-78%)	27% (22-34%)	2.4 (2-2.9)	1.3% (-0.93-3.6%)	1.7% (-1.5-4.8%)	100 (-93-300)	2200 (-1500-5900)
Australia	1990	Female	75 to 79	67% (58-76%)	33% (24-42%)	2 (1.6-2.5)	0.33% (-2.2-2.7%)	0.21% (-3.2-3.3%)	18 (-280-290)	550 (-3600-4500)
Australia	1990	Female	80 to 84	65% (52-75%)	35% (25-48%)	1.8 (1.4-2.3)	-0.44% (-3.1-2%)	-0.77% (-4.1-2.4%)	-73 (-390-220)	-560 (-3900-2600)
Australia	1990	Female	85 to 89	64% (52-74%)	36% (26-48%)	1.7 (1.2-2.3)	-0.95% (-3.6-1.6%)	-1.4% (-4.7-1.8%)	-120 (-400-150)	-740 (-2800-1300)
Australia	1990	Female	90 to 94	64% (51-75%)	36% (25-49%)	1.6 (1-2.3)	-1.4% (-4-1.6%)	-1.9% (-5.1-1.6%)	-100 (-270-82)	-440 (-1300-510)
Australia	1990	Female	95 plus	65% (49-79%)	35% (21-51%)	1.5 (0.81-2.4)	-1.7% (-4.7-1.8%)	-2.4% (-6.1-1.8%)	-55 (-140-42)	-160 (-440-160)
Australia	1990	Male	All Ages	88% (85-91%)	12% (9.5-15%)	5.2 (4.6-5.9)	11% (8.8-13%)	12% (8.8-15%)	7700 (5700-9800)	260000 (210000-310000)
Australia	1990	Male	15 to 19	81% (72-88%)	19% (12-28%)	3.1 (2.3-3.9)	7.9% (4.7-11%)	16% (8.6-22%)	100 (56-140)	8400 (5100-12000)
Australia	1990	Male	20 to 24	90% (85-94%)	10% (6.5-15%)	5.4 (4.3-6.5)	15% (11-19%)	25% (17-33%)	230 (160-300)	18000 (13000-24000)
Australia	1990	Male	25 to 29	92% (89-94%)	8% (5.8-11%)	5.9 (5-6.9)	14% (10-18%)	25% (18-32%)	240 (170-300)	18000 (14000-22000)
Australia	1990	Male	30 to 34	94% (91-96%)	6.4% (4.3-9.1%)	6.6 (5.6-7.8)	13% (10-16%)	24% (18-30%)	230 (170-280)	16000 (13000-20000)
Australia	1990	Male	35 to 39	95% (94-97%)	4.6% (3-6.2%)	7.5 (6.4-8.8)	13% (10-16%)	25% (20-30%)	250 (200-300)	16000 (13000-19000)
Australia	1990	Male	40 to 44	96% (95-98%)	3.7% (2.4-5.2%)	8.1 (6.9-9.3)	13% (11-16%)	24% (20-29%)	320 (260-370)	18000 (15000-21000)
Australia	1990	Male	45 to 49	96% (94-97%)	4.1% (2.7-5.7%)	8.3 (7.1-9.7)	14% (11-17%)	23% (19-27%)	360 (290-430)	18000 (14000-21000)
Australia	1990	Male	50 to 54	95% (94-97%)	4.7% (3.3-6.3%)	8.6 (7.3-10)	15% (12-18%)	21% (17-25%)	480 (380-570)	20000 (16000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	1990	Male	55 to 59	94% (92-96%)	5.5% (3.9-7.5%)	9 (7.7-10)	15% (12-18%)	20% (16-25%)	680 (530-840)	24000 (19000-29000)
Australia	1990	Male	60 to 64	93% (91-95%)	6.9% (4.9-9.2%)	8.7 (7.3-10)	14% (11-18%)	18% (13-22%)	1000 (760-1300)	30000 (23000-38000)
Australia	1990	Male	65 to 69	91% (88-94%)	8.7% (6.2-12%)	7.6 (6.5-8.9)	13% (9.4-16%)	15% (11-19%)	1200 (890-1600)	30000 (22000-37000)
Australia	1990	Male	70 to 74	90% (86-93%)	9.8% (6.9-14%)	6.4 (5.3-7.7)	10% (6.6-14%)	12% (7.3-16%)	1100 (660-1500)	21000 (13000-28000)
Australia	1990	Male	75 to 79	90% (84-94%)	9.9% (6.3-16%)	5.4 (4.3-6.7)	7.7% (3.2-12%)	8.4% (3.1-13%)	880 (320-1400)	14000 (5700-21000)
Australia	1990	Male	80 to 84	89% (83-94%)	11% (6.4-17%)	4.6 (3.6-5.7)	5.4% (1.3-9.3%)	5.7% (0.95-10%)	490 (82-900)	5800 (1400-10000)
Australia	1990	Male	85 to 89	88% (81-92%)	12% (7.6-19%)	3.8 (2.9-4.8)	3% (-0.44-6.6%)	2.9% (-1.2-7%)	150 (-61-370)	1400 (-200-3000)
Australia	1990	Male	90 to 94	87% (79-92%)	13% (8-21%)	3.1 (2.2-4.2)	1% (-2.1-4.8%)	0.63% (-3-4.9%)	12 (-59-97)	120 (-250-560)
Australia	1990	Male	95 plus	86% (76-93%)	14% (7-24%)	2.4 (1.3-3.8)	-0.36% (-3.4-3.6%)	-0.99% (-4.6-3.7%)	-5.6 (-26-21)	-8.1 (-77-84)
Australia	1995	Female	All Ages	85% (82-87%)	15% (13-18%)	2 (1.7-2.2)	2% (0.87-3.1%)	0.48% (-2.1-3.2%)	280 (-1300-1900)	41000 (18000-64000)
Australia	1995	Female	15 to 19	87% (81-92%)	13% (7.8-19%)	1.9 (1.5-2.5)	2.8% (1.8-3.9%)	9.3% (4.8-14%)	19 (9.7-28)	2100 (1300-2900)
Australia	1995	Female	20 to 24	92% (88-95%)	8% (5.4-12%)	2.6 (2.1-3.2)	4% (2.8-5.2%)	11% (5.8-16%)	29 (16-42)	3900 (2700-5200)
Australia	1995	Female	25 to 29	93% (90-95%)	6.8% (4.7-9.6%)	2.2 (1.9-2.6)	3.2% (2.3-4.2%)	9.1% (5.1-13%)	26 (15-36)	3200 (2300-4300)
Australia	1995	Female	30 to 34	93% (91-95%)	6.5% (4.6-8.9%)	2.3 (1.9-2.7)	2.9% (2-3.8%)	8.8% (5.7-12%)	35 (22-48)	3200 (2300-4300)
Australia	1995	Female	35 to 39	93% (91-95%)	7.1% (4.9-9.4%)	2.8 (2.4-3.2)	3.4% (2.5-4.4%)	11% (8.1-14%)	58 (42-73)	4100 (3100-5200)
Australia	1995	Female	40 to 44	92% (90-94%)	7.8% (5.6-10%)	3 (2.6-3.5)	3.9% (2.9-5%)	12% (8.8-14%)	85 (64-100)	4900 (3800-6000)
Australia	1995	Female	45 to 49	91% (89-94%)	8.6% (6.1-11%)	3 (2.5-3.5)	4.2% (3.1-5.3%)	11% (8.3-13%)	110 (86-140)	5600 (4200-6900)
Australia	1995	Female	50 to 54	89% (86-92%)	11% (8.2-14%)	2.9 (2.5-3.4)	4.4% (3.1-5.7%)	9.5% (7.2-12%)	130 (95-160)	5200 (3800-6600)
Australia	1995	Female	55 to 59	85% (81-89%)	15% (11-19%)	2.9 (2.4-3.4)	4% (2.6-5.5%)	7.5% (5.3-9.8%)	130 (92-170)	4600 (3100-6100)
Australia	1995	Female	60 to 64	81% (76-86%)	19% (14-24%)	2.7 (2.2-3.2)	3% (1.5-4.7%)	5.1% (2.7-7.6%)	130 (70-190)	3800 (1900-5900)
Australia	1995	Female	65 to 69	80% (74-84%)	20% (16-26%)	2.5 (2.1-2.9)	1.9% (0.16-3.7%)	3% (0.25-5.6%)	120 (10-230)	3000 (220-5600)
Australia	1995	Female	70 to 74	76% (70-81%)	24% (19-30%)	2.2 (1.8-2.7)	0.85% (-1.3-3%)	1.2% (-1.8-4.2%)	77 (-110-270)	1500 (-2300-5300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	1995	Female	75 to 79	71% (63-79%)	29% (21-37%)	1.9 (1.5-2.4)	-0.22% (-2.5-2.2%)	-0.39% (-3.6-2.8%)	-32 (-290-230)	-350 (-4100-3500)
Australia	1995	Female	80 to 84	68% (56-77%)	32% (23-44%)	1.7 (1.3-2.1)	-1.1% (-3.5-1.4%)	-1.5% (-4.6-1.8%)	-160 (-510-200)	-1600 (-5200-2200)
Australia	1995	Female	85 to 89	66% (54-75%)	34% (25-46%)	1.5 (1.1-2)	-1.6% (-4-0.89%)	-2.1% (-5.1-0.96%)	-220 (-520-97)	-1500 (-3800-830)
Australia	1995	Female	90 to 94	66% (54-76%)	34% (24-46%)	1.4 (0.94-1.9)	-2% (-4.4-0.56%)	-2.6% (-5.7-0.55%)	-170 (-360-35)	-810 (-1800-230)
Australia	1995	Female	95 plus	66% (51-79%)	34% (21-49%)	1.3 (0.74-2.1)	-2.4% (-5-0.58%)	-3.1% (-6.3-0.44%)	-91 (-190-13)	-280 (-590-68)
Australia	1995	Male	All Ages	88% (85-90%)	12% (9.7-15%)	4.6 (4.1-5.2)	9.2% (7.4-11%)	9.5% (6.5-13%)	6300 (4300-8400)	220000 (180000-270000)
Australia	1995	Male	15 to 19	79% (68-87%)	21% (13-32%)	2.6 (1.9-3.4)	6.5% (3.9-9.2%)	13% (7.1-20%)	68 (36-99)	6000 (3600-8300)
Australia	1995	Male	20 to 24	88% (82-93%)	12% (7.1-18%)	4.6 (3.6-5.5)	13% (9-17%)	21% (14-29%)	190 (120-250)	16000 (11000-21000)
Australia	1995	Male	25 to 29	91% (87-94%)	9% (6-13%)	5.1 (4.3-6)	12% (8.9-15%)	21% (15-28%)	180 (130-240)	15000 (11000-18000)
Australia	1995	Male	30 to 34	93% (90-96%)	6.9% (4.4-10%)	5.8 (4.8-6.8)	11% (8.4-14%)	20% (15-26%)	210 (150-260)	15000 (11000-18000)
Australia	1995	Male	35 to 39	95% (93-97%)	4.8% (3.2-6.8%)	6.5 (5.5-7.6)	11% (8.7-14%)	21% (16-26%)	230 (180-290)	15000 (12000-18000)
Australia	1995	Male	40 to 44	96% (94-97%)	4% (2.5-5.9%)	7 (6-8.1)	12% (9.3-14%)	22% (18-26%)	280 (220-330)	16000 (13000-19000)
Australia	1995	Male	45 to 49	96% (94-97%)	4.3% (2.8-6.4%)	7.2 (6.2-8.4)	12% (9.7-15%)	21% (17-25%)	360 (290-430)	18000 (14000-21000)
Australia	1995	Male	50 to 54	95% (93-97%)	4.8% (3.3-6.8%)	7.5 (6.4-8.7)	13% (10-15%)	19% (16-23%)	430 (340-510)	18000 (15000-21000)
Australia	1995	Male	55 to 59	95% (92-96%)	5.5% (3.7-7.8%)	7.8 (6.7-9.1)	13% (11-16%)	18% (15-22%)	570 (450-680)	20000 (17000-24000)
Australia	1995	Male	60 to 64	93% (91-95%)	6.7% (4.6-9.5%)	7.6 (6.4-9)	13% (9.9-16%)	16% (12-20%)	760 (590-950)	23000 (18000-28000)
Australia	1995	Male	65 to 69	91% (88-94%)	8.5% (5.8-12%)	6.7 (5.6-7.8)	11% (7.8-14%)	13% (9-17%)	980 (670-1300)	24000 (17000-31000)
Australia	1995	Male	70 to 74	91% (87-93%)	9.3% (6.6-13%)	5.7 (4.6-6.8)	8.3% (4.7-12%)	9.6% (5.2-14%)	950 (520-1300)	19000 (11000-27000)
Australia	1995	Male	75 to 79	91% (87-94%)	9% (5.8-13%)	4.8 (3.8-6)	5.8% (1.8-9.5%)	6.2% (1.5-11%)	630 (150-1100)	10000 (3200-17000)
Australia	1995	Male	80 to 84	90% (84-94%)	10% (6.1-16%)	4.1 (3.2-5)	3.7% (0.24-7.3%)	3.8% (-0.35-8%)	390 (-36-820)	4700 (300-9300)
Australia	1995	Male	85 to 89	88% (82-93%)	12% (7.5-18%)	3.3 (2.6-4.2)	1.8% (-1.3-5.1%)	1.5% (-2.1-5.4%)	100 (-140-360)	1000 (-750-3000)
Australia	1995	Male	90 to 94	87% (80-92%)	13% (7.8-20%)	2.8 (2-3.8)	0.27% (-2.5-3.5%)	-0.2% (-3.5-3.7%)	-5.5 (-98-100)	46 (-430-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	1995	Male	95 plus	86% (76-93%)	14% (6.9-24%)	2.2 (1.2-3.4)	-0.8% (-3.5-2.5%)	-1.5% (-4.7-2.4%)	-10 (-34-17)	-23 (-100-72)
Australia	2000	Female	All Ages	85% (82-87%)	15% (13-18%)	1.9 (1.7-2.1)	1.8% (0.82-2.8%)	0.27% (-2.1-2.6%)	160 (-1300-1600)	39000 (17000-60000)
Australia	2000	Female	15 to 19	85% (77-92%)	15% (8.3-23%)	1.8 (1.3-2.2)	2.6% (1.6-3.6%)	8.6% (4.4-13%)	17 (8.9-27)	2000 (1200-2800)
Australia	2000	Female	20 to 24	91% (87-94%)	8.6% (5.6-13%)	2.4 (1.9-3)	3.7% (2.6-4.9%)	9.7% (5.2-14%)	23 (12-34)	3300 (2300-4400)
Australia	2000	Female	25 to 29	93% (90-95%)	6.9% (4.6-10%)	2.1 (1.7-2.4)	3.1% (2.2-4%)	8.3% (4.6-12%)	25 (14-36)	3200 (2200-4200)
Australia	2000	Female	30 to 34	94% (91-95%)	6.5% (4.5-9.1%)	2.1 (1.8-2.5)	2.7% (1.8-3.6%)	8.2% (5-11%)	30 (18-42)	2900 (2000-3900)
Australia	2000	Female	35 to 39	93% (91-95%)	7% (4.9-9.4%)	2.6 (2.2-3)	3.1% (2.2-4.2%)	10% (7-13%)	55 (38-71)	3900 (2900-5000)
Australia	2000	Female	40 to 44	92% (90-94%)	7.6% (5.6-10%)	2.8 (2.4-3.3)	3.6% (2.7-4.7%)	11% (8.3-14%)	84 (63-100)	4900 (3700-6100)
Australia	2000	Female	45 to 49	92% (89-94%)	8.4% (5.8-11%)	2.8 (2.4-3.2)	3.8% (2.7-4.9%)	10% (7.9-13%)	110 (83-130)	5300 (4000-6600)
Australia	2000	Female	50 to 54	89% (86-92%)	11% (7.9-14%)	2.7 (2.3-3.2)	3.9% (2.7-5.1%)	9.3% (6.9-11%)	140 (100-170)	5700 (4200-7200)
Australia	2000	Female	55 to 59	85% (81-89%)	15% (11-19%)	2.7 (2.3-3.2)	3.7% (2.4-5%)	7.4% (5.1-9.5%)	140 (96-170)	4700 (3200-6200)
Australia	2000	Female	60 to 64	82% (77-86%)	18% (14-23%)	2.5 (2.1-3)	2.8% (1.5-4.2%)	5.3% (2.9-7.4%)	120 (70-180)	3600 (1800-5300)
Australia	2000	Female	65 to 69	80% (75-85%)	20% (15-25%)	2.4 (2-2.7)	1.9% (0.36-3.3%)	3.2% (0.96-5.4%)	110 (33-180)	2500 (480-4500)
Australia	2000	Female	70 to 74	78% (72-82%)	22% (18-28%)	2.1 (1.8-2.6)	0.88% (-0.88-2.7%)	1.4% (-1.2-4.1%)	80 (-67-230)	1400 (-1500-4600)
Australia	2000	Female	75 to 79	73% (66-81%)	27% (19-34%)	1.9 (1.5-2.3)	-0.13% (-2-1.8%)	-0.19% (-2.8-2.5%)	-16 (-240-210)	-230 (-3500-3300)
Australia	2000	Female	80 to 84	70% (59-79%)	30% (21-41%)	1.6 (1.3-2.1)	-1% (-3.1-1%)	-1.4% (-4.2-1.3%)	-150 (-440-130)	-1600 (-4700-1500)
Australia	2000	Female	85 to 89	68% (57-76%)	32% (24-43%)	1.5 (1.1-1.9)	-1.6% (-3.8-0.59%)	-2.2% (-4.9-0.65%)	-260 (-590-78)	-1900 (-4300-660)
Australia	2000	Female	90 to 94	68% (57-78%)	32% (22-43%)	1.4 (0.93-1.9)	-2.1% (-4.3-0.34%)	-2.7% (-5.5-0.24%)	-220 (-450-20)	-1100 (-2200-180)
Australia	2000	Female	95 plus	68% (53-81%)	32% (19-47%)	1.3 (0.72-2)	-2.5% (-5-0.32%)	-3.2% (-6.3-0.14%)	-120 (-230-5)	-360 (-730-48)
Australia	2000	Male	All Ages	88% (85-91%)	12% (9.4-15%)	4.3 (3.8-4.9)	8.5% (6.8-10%)	8.3% (5.8-11%)	5600 (3900-7500)	200000 (160000-250000)
Australia	2000	Male	15 to 19	79% (68-87%)	21% (13-32%)	2.4 (1.8-3.1)	6.1% (3.6-8.7%)	13% (6.6-19%)	63 (32-92)	5600 (3400-7800)
Australia	2000	Male	20 to 24	88% (82-93%)	12% (7-18%)	4.2 (3.4-5.2)	12% (8.1-15%)	20% (13-27%)	140 (88-200)	12000 (8700-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	2000	Male	25 to 29	91% (87-94%)	9% (6-13%)	4.7 (4-5.5)	12% (8.4-15%)	20% (14-27%)	180 (120-240)	14000 (10000-18000)
Australia	2000	Male	30 to 34	93% (90-96%)	6.8% (4.3-10%)	5.3 (4.5-6.2)	11% (8.1-14%)	20% (15-26%)	190 (130-240)	13000 (10000-17000)
Australia	2000	Male	35 to 39	95% (93-97%)	4.8% (3-6.8%)	6 (5.1-7.1)	11% (8.4-13%)	21% (16-26%)	220 (170-280)	15000 (11000-18000)
Australia	2000	Male	40 to 44	96% (94-98%)	4% (2.5-5.8%)	6.5 (5.6-7.5)	11% (9.1-14%)	21% (17-26%)	280 (220-340)	16000 (13000-19000)
Australia	2000	Male	45 to 49	96% (94-97%)	4.3% (2.7-6.5%)	6.6 (5.7-7.7)	12% (9.3-14%)	20% (16-24%)	330 (260-400)	17000 (13000-20000)
Australia	2000	Male	50 to 54	95% (93-97%)	4.8% (3.2-6.8%)	6.9 (5.9-8)	12% (9.4-14%)	19% (15-22%)	450 (360-540)	19000 (16000-23000)
Australia	2000	Male	55 to 59	95% (92-96%)	5.5% (3.8-7.7%)	7.2 (6.1-8.5)	12% (9.8-15%)	18% (14-21%)	550 (430-660)	20000 (16000-24000)
Australia	2000	Male	60 to 64	93% (91-95%)	6.7% (4.7-9.4%)	7 (5.9-8.3)	12% (9.2-14%)	16% (12-19%)	660 (500-800)	20000 (16000-25000)
Australia	2000	Male	65 to 69	92% (88-94%)	8.5% (5.9-12%)	6.2 (5.3-7.2)	9.9% (7.2-13%)	12% (8.6-16%)	730 (520-940)	19000 (13000-24000)
Australia	2000	Male	70 to 74	91% (87-93%)	9.2% (6.6-13%)	5.2 (4.3-6.3)	7.5% (4.2-11%)	8.8% (4.8-13%)	800 (440-1200)	16000 (9500-24000)
Australia	2000	Male	75 to 79	91% (87-94%)	8.9% (5.8-13%)	4.5 (3.5-5.5)	5.1% (1.8-8.6%)	5.5% (1.6-9.7%)	620 (180-1100)	10000 (3600-17000)
Australia	2000	Male	80 to 84	90% (84-94%)	9.9% (6-16%)	3.8 (3-4.6)	3.1% (0.27-6.2%)	3.1% (-0.28-7%)	320 (-29-700)	4000 (340-8100)
Australia	2000	Male	85 to 89	88% (82-93%)	12% (7.2-18%)	3.1 (2.4-3.9)	1.3% (-1.1-4.3%)	1.1% (-1.9-4.7%)	89 (-160-380)	980 (-840-3200)
Australia	2000	Male	90 to 94	87% (81-92%)	13% (7.6-19%)	2.6 (1.8-3.5)	0.022% (-2.6-3%)	-0.45% (-3.6-3.1%)	-17 (-130-110)	5.2 (-590-680)
Australia	2000	Male	95 plus	86% (77-93%)	14% (6.8-23%)	2 (1.1-3.2)	-0.94% (-3.5-2.1%)	-1.6% (-4.6-2%)	-16 (-46-20)	-37 (-140-85)
Australia	2005	Female	All Ages	84% (81-86%)	16% (14-19%)	1.9 (1.7-2.2)	1.9% (1.1-2.8%)	0.73% (-1.2-2.9%)	480 (-790-1900)	42000 (23000-62000)
Australia	2005	Female	15 to 19	88% (81-92%)	12% (7.5-19%)	1.8 (1.4-2.3)	2.3% (1.5-3.2%)	9.2% (4.6-14%)	15 (7.6-23)	1800 (1100-2500)
Australia	2005	Female	20 to 24	92% (88-94%)	8.4% (5.7-12%)	2.5 (2-3)	3.3% (2.3-4.4%)	10% (5.4-15%)	22 (11-31)	3100 (2100-4200)
Australia	2005	Female	25 to 29	92% (89-95%)	7.8% (5.4-11%)	2.1 (1.7-2.4)	2.8% (2-3.7%)	8.7% (4.9-13%)	20 (11-29)	2600 (1800-3600)
Australia	2005	Female	30 to 34	93% (89-95%)	7.5% (5.2-11%)	2.1 (1.8-2.5)	2.5% (1.7-3.3%)	8.4% (5.2-12%)	27 (17-38)	2800 (1900-3700)
Australia	2005	Female	35 to 39	92% (90-95%)	7.8% (5.4-10%)	2.6 (2.2-3)	3% (2.1-3.9%)	10% (7.3-14%)	49 (35-64)	3600 (2600-4600)
Australia	2005	Female	40 to 44	92% (89-94%)	8.2% (5.9-11%)	2.8 (2.5-3.3)	3.4% (2.5-4.4%)	11% (8.4-14%)	82 (63-100)	4800 (3700-6000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	2005	Female	45 to 49	91% (88-94%)	9.2% (6.4-12%)	2.8 (2.4-3.3)	3.6% (2.6-4.7%)	10% (7.9-13%)	110 (85-140)	5500 (4100-6800)
Australia	2005	Female	50 to 54	88% (84-91%)	12% (9-16%)	2.7 (2.3-3.2)	3.8% (2.6-4.9%)	9.5% (7.3-11%)	140 (110-170)	5700 (4200-7100)
Australia	2005	Female	55 to 59	84% (80-88%)	16% (12-20%)	2.7 (2.3-3.2)	3.5% (2.5-4.7%)	7.8% (5.9-9.6%)	160 (120-200)	5500 (4000-7200)
Australia	2005	Female	60 to 64	82% (78-86%)	18% (14-22%)	2.6 (2.2-3.1)	3.1% (1.8-4.4%)	6% (4-7.8%)	150 (100-200)	4400 (2700-6100)
Australia	2005	Female	65 to 69	78% (73-84%)	22% (16-27%)	2.3 (2-2.7)	2.2% (1-3.5%)	4.1% (2.2-5.8%)	130 (73-190)	3100 (1400-4800)
Australia	2005	Female	70 to 74	74% (68-79%)	26% (21-32%)	2 (1.7-2.5)	1.3% (0.0017-2.7%)	2.3% (0.3-4.2%)	110 (14-200)	1900 (2.3-3900)
Australia	2005	Female	75 to 79	68% (59-77%)	32% (23-41%)	1.7 (1.3-2.2)	0.44% (-1-2%)	0.67% (-1.4-2.7%)	52 (-110-210)	730 (-1700-3400)
Australia	2005	Female	80 to 84	65% (53-75%)	35% (25-47%)	1.5 (1.1-2)	-0.39% (-2-1.3%)	-0.54% (-2.7-1.7%)	-63 (-320-200)	-670 (-3500-2200)
Australia	2005	Female	85 to 89	66% (55-76%)	34% (24-45%)	1.5 (1.1-1.9)	-1.1% (-2.9-0.89%)	-1.4% (-3.8-1.1%)	-180 (-470-130)	-1300 (-3500-1100)
Australia	2005	Female	90 to 94	66% (54-77%)	34% (23-46%)	1.4 (0.91-1.9)	-1.6% (-3.6-0.72%)	-2.1% (-4.7-0.79%)	-220 (-480-82)	-1100 (-2400-480)
Australia	2005	Female	95 plus	66% (51-80%)	34% (20-49%)	1.3 (0.71-2)	-2% (-4.3-0.71%)	-2.7% (-5.5-0.57%)	-130 (-270-27)	-390 (-840-130)
Australia	2005	Male	All Ages	89% (86-91%)	11% (8.6-14%)	4.5 (4-5.1)	8.4% (6.9-10%)	8.2% (6-11%)	5600 (4100-7400)	200000 (160000-240000)
Australia	2005	Male	15 to 19	83% (74-89%)	17% (11-26%)	2.6 (1.9-3.3)	5.5% (3.4-7.8%)	14% (7.3-20%)	49 (26-73)	4600 (2900-6400)
Australia	2005	Male	20 to 24	91% (85-94%)	9.5% (6-15%)	4.4 (3.5-5.4)	11% (7.9-14%)	21% (14-29%)	130 (82-170)	11000 (8000-15000)
Australia	2005	Male	25 to 29	92% (89-94%)	8.2% (5.8-11%)	4.8 (4.1-5.7)	11% (8.1-14%)	21% (15-28%)	130 (91-170)	11000 (8200-14000)
Australia	2005	Male	30 to 34	94% (91-96%)	6.5% (4.4-9.2%)	5.4 (4.6-6.3)	11% (8-13%)	22% (15-28%)	170 (120-220)	13000 (9700-16000)
Australia	2005	Male	35 to 39	96% (94-97%)	4.5% (3-6%)	6.1 (5.3-7.1)	11% (8.3-13%)	22% (17-27%)	190 (150-240)	13000 (10000-16000)
Australia	2005	Male	40 to 44	96% (95-98%)	3.7% (2.4-5.3%)	6.6 (5.7-7.6)	11% (9-14%)	22% (18-26%)	280 (220-330)	16000 (13000-19000)
Australia	2005	Male	45 to 49	96% (94-97%)	4% (2.6-5.6%)	6.8 (5.8-7.9)	12% (9.4-14%)	21% (17-25%)	360 (290-430)	18000 (15000-22000)
Australia	2005	Male	50 to 54	96% (94-97%)	4.4% (3.1-5.9%)	7.1 (6.1-8.2)	12% (9.5-14%)	19% (16-23%)	450 (360-530)	20000 (16000-23000)
Australia	2005	Male	55 to 59	95% (93-96%)	5.2% (3.6-7.3%)	7.4 (6.3-8.5)	12% (9.7-14%)	18% (15-21%)	610 (500-720)	23000 (19000-27000)
Australia	2005	Male	60 to 64	93% (91-95%)	6.7% (4.7-9.2%)	7.1 (6-8.3)	11% (9-14%)	16% (12-19%)	670 (530-790)	21000 (17000-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	2005	Male	65 to 69	91% (88-94%)	8.8% (5.9-12%)	6.2 (5.3-7.1)	9.7% (7.4-12%)	12% (9.4-15%)	700 (530-860)	18000 (14000-22000)
Australia	2005	Male	70 to 74	91% (88-93%)	9.2% (6.8-12%)	5.3 (4.4-6.3)	7.5% (4.9-10%)	9.1% (5.7-12%)	680 (420-920)	14000 (9300-19000)
Australia	2005	Male	75 to 79	91% (87-94%)	8.9% (5.9-13%)	4.5 (3.6-5.6)	5.4% (2.6-8.3%)	6.1% (2.7-9.7%)	650 (290-1000)	11000 (5400-17000)
Australia	2005	Male	80 to 84	90% (84-94%)	10% (6.3-16%)	3.8 (3-4.6)	3.5% (1.1-6.2%)	3.7% (0.74-7%)	440 (90-830)	5500 (1700-9800)
Australia	2005	Male	85 to 89	88% (81-92%)	12% (7.5-19%)	3.1 (2.4-4)	1.8% (-0.59-4.4%)	1.6% (-1.3-4.7%)	140 (-120-420)	1400 (-510-3500)
Australia	2005	Male	90 to 94	87% (80-92%)	13% (8-20%)	2.6 (1.9-3.5)	0.37% (-2-3.2%)	-0.042% (-2.9-3.3%)	-2.1 (-140-170)	110 (-620-990)
Australia	2005	Male	95 plus	86% (76-93%)	14% (6.9-24%)	2.1 (1.1-3.2)	-0.59% (-3.1-2.6%)	-1.2% (-4.1-2.6%)	-17 (-60-37)	-34 (-170-150)
Australia	2010	Female	All Ages	84% (82-86%)	16% (14-18%)	2 (1.7-2.3)	2.1% (1.3-3%)	1.2% (-0.73-3.1%)	830 (-510-2200)	50000 (31000-69000)
Australia	2010	Female	15 to 19	88% (82-93%)	12% (7.4-18%)	1.9 (1.4-2.4)	2.3% (1.5-3.2%)	9.4% (4.7-14%)	15 (7.5-23)	1900 (1200-2600)
Australia	2010	Female	20 to 24	92% (88-94%)	8.4% (5.7-12%)	2.6 (2.1-3.2)	3.4% (2.4-4.6%)	11% (5.6-16%)	23 (12-34)	3500 (2400-4700)
Australia	2010	Female	25 to 29	92% (89-95%)	7.6% (5.2-11%)	2.2 (1.8-2.6)	2.8% (2-3.7%)	9% (5-13%)	22 (12-32)	3100 (2200-4100)
Australia	2010	Female	30 to 34	93% (90-95%)	7.3% (5.1-10%)	2.2 (1.8-2.6)	2.5% (1.7-3.3%)	8.8% (5.4-12%)	27 (16-37)	2800 (1900-3700)
Australia	2010	Female	35 to 39	92% (90-95%)	7.7% (5.3-10%)	2.7 (2.2-3.2)	3% (2.1-4%)	11% (7.4-14%)	53 (36-69)	3900 (2900-5100)
Australia	2010	Female	40 to 44	92% (89-94%)	8.1% (5.9-11%)	2.9 (2.5-3.5)	3.4% (2.5-4.4%)	11% (8.7-14%)	81 (62-99)	4900 (3700-6100)
Australia	2010	Female	45 to 49	91% (88-94%)	9.1% (6.4-12%)	2.9 (2.4-3.4)	3.6% (2.6-4.7%)	11% (8.2-13%)	120 (92-140)	5900 (4300-7300)
Australia	2010	Female	50 to 54	88% (85-91%)	12% (9-15%)	2.8 (2.3-3.3)	3.8% (2.7-5%)	9.8% (7.6-12%)	150 (120-190)	6400 (4700-8000)
Australia	2010	Female	55 to 59	85% (80-88%)	15% (12-20%)	2.8 (2.3-3.3)	3.6% (2.5-4.8%)	8.2% (6.2-10%)	170 (130-210)	6000 (4300-7800)
Australia	2010	Female	60 to 64	82% (77-86%)	18% (14-23%)	2.7 (2.2-3.2)	3.1% (2-4.4%)	6.5% (4.7-8.3%)	190 (130-240)	5500 (3600-7400)
Australia	2010	Female	65 to 69	78% (72-84%)	22% (16-28%)	2.4 (2-2.9)	2.4% (1.3-3.6%)	4.7% (3-6.3%)	170 (100-220)	3900 (2100-5700)
Australia	2010	Female	70 to 74	74% (67-80%)	26% (20-33%)	2.1 (1.7-2.6)	1.7% (0.43-3%)	3% (1.2-4.8%)	150 (58-230)	2700 (710-4600)
Australia	2010	Female	75 to 79	68% (59-77%)	32% (23-41%)	1.8 (1.4-2.3)	0.89% (-0.41-2.3%)	1.4% (-0.46-3.2%)	97 (-31-220)	1400 (-640-3500)
Australia	2010	Female	80 to 84	65% (54-76%)	35% (24-46%)	1.6 (1.2-2.1)	0.16% (-1.3-1.6%)	0.22% (-1.7-2.1%)	26 (-200-250)	280 (-2300-2800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	2010	Female	85 to 89	66% (54-76%)	34% (24-46%)	1.5 (1.1-2)	-0.53% (-2.3-1.2%)	-0.77% (-3-1.5%)	-120 (-460-220)	-800 (-3500-1800)
Australia	2010	Female	90 to 94	66% (55-76%)	34% (24-45%)	1.4 (0.97-2)	-1.2% (-3.1-1.1%)	-1.6% (-4-1.2%)	-190 (-470-140)	-870 (-2400-840)
Australia	2010	Female	95 plus	67% (51-80%)	33% (20-49%)	1.4 (0.75-2.2)	-1.6% (-3.9-1.2%)	-2.2% (-5-1.1%)	-150 (-340-75)	-440 (-1100-320)
Australia	2010	Male	All Ages	89% (87-91%)	11% (8.6-13%)	4.7 (4.1-5.4)	8.5% (6.9-10%)	8.4% (6.2-11%)	6200 (4600-8000)	220000 (180000-260000)
Australia	2010	Male	15 to 19	82% (74-89%)	18% (11-26%)	2.7 (2-3.4)	5.5% (3.5-7.8%)	14% (7.4-20%)	48 (26-69)	4800 (3100-6500)
Australia	2010	Male	20 to 24	90% (85-94%)	9.7% (6.2-15%)	4.6 (3.6-5.7)	11% (7.8-14%)	22% (14-30%)	120 (75-160)	11000 (8200-15000)
Australia	2010	Male	25 to 29	92% (89-94%)	8.2% (5.8-11%)	5 (4.1-6)	10% (7.9-13%)	22% (15-28%)	130 (93-170)	12000 (9000-15000)
Australia	2010	Male	30 to 34	94% (91-96%)	6.4% (4.2-9.2%)	5.6 (4.7-6.7)	10% (8-13%)	22% (16-28%)	150 (110-190)	12000 (9200-15000)
Australia	2010	Male	35 to 39	96% (94-97%)	4.4% (3-5.9%)	6.3 (5.3-7.4)	11% (8.4-13%)	23% (18-28%)	210 (170-260)	14000 (11000-17000)
Australia	2010	Male	40 to 44	96% (95-98%)	3.7% (2.4-5%)	6.8 (5.8-7.9)	11% (8.9-14%)	23% (18-27%)	270 (220-320)	16000 (13000-19000)
Australia	2010	Male	45 to 49	96% (95-97%)	4% (2.6-5.5%)	7 (6-8.2)	12% (9.6-14%)	22% (18-26%)	380 (310-460)	19000 (16000-23000)
Australia	2010	Male	50 to 54	96% (94-97%)	4.4% (3.1-6%)	7.3 (6.2-8.5)	12% (9.8-14%)	20% (17-23%)	500 (410-590)	22000 (18000-26000)
Australia	2010	Male	55 to 59	95% (93-96%)	5.2% (3.6-7.2%)	7.6 (6.4-9)	12% (10-14%)	19% (16-22%)	640 (540-750)	24000 (20000-28000)
Australia	2010	Male	60 to 64	93% (91-95%)	6.7% (4.8-9.1%)	7.4 (6.1-8.7)	11% (9.2-13%)	16% (13-19%)	780 (640-920)	25000 (21000-30000)
Australia	2010	Male	65 to 69	91% (88-94%)	8.7% (6.1-12%)	6.4 (5.4-7.5)	9.7% (7.6-12%)	13% (10-16%)	770 (600-940)	21000 (16000-25000)
Australia	2010	Male	70 to 74	91% (88-93%)	9.1% (6.7-12%)	5.5 (4.4-6.6)	7.8% (5.3-10%)	9.8% (6.6-13%)	730 (500-970)	16000 (11000-21000)
Australia	2010	Male	75 to 79	91% (87-94%)	9% (6-13%)	4.7 (3.6-5.7)	5.8% (3-8.4%)	6.7% (3.3-9.9%)	650 (320-960)	11000 (5800-16000)
Australia	2010	Male	80 to 84	90% (84-94%)	10% (6.3-16%)	3.9 (3-4.8)	3.9% (1.5-6.5%)	4.2% (1.3-7.4%)	550 (170-950)	6900 (2700-11000)
Australia	2010	Male	85 to 89	88% (81-92%)	12% (7.6-19%)	3.2 (2.4-4.1)	2.3% (-0.074-4.7%)	2.2% (-0.57-5.2%)	260 (-68-620)	2500 (-86-5300)
Australia	2010	Male	90 to 94	87% (80-92%)	13% (7.8-20%)	2.7 (1.9-3.7)	0.85% (-1.5-3.7%)	0.52% (-2.3-3.9%)	33 (-150-250)	340 (-590-1500)
Australia	2010	Male	95 plus	86% (77-93%)	14% (7.1-23%)	2.1 (1.2-3.4)	-0.17% (-2.6-3%)	-0.69% (-3.6-3.1%)	-15 (-76-65)	-14 (-220-260)
Australia	2016	Female	All Ages	83% (80-86%)	17% (14-20%)	2.1 (1.6-2.7)	2.2% (1.2-3.3%)	1.4% (-0.81-3.6%)	1100 (-640-2900)	58000 (32000-85000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	2016	Female	15 to 19	85% (77-92%)	15% (8.2-23%)	2.1 (1.2-3.1)	2.4% (1.5-3.5%)	9.8% (4.5-16%)	15 (7.1-24)	1900 (1200-2700)
Australia	2016	Female	20 to 24	91% (86-94%)	9.1% (5.8-14%)	2.9 (1.7-4.3)	3.6% (2.4-4.9%)	12% (6.4-18%)	25 (13-37)	3900 (2600-5400)
Australia	2016	Female	25 to 29	92% (88-95%)	7.9% (5.1-12%)	2.4 (1.7-3.2)	2.9% (2.1-3.9%)	9.4% (5.3-14%)	25 (14-37)	3700 (2600-5100)
Australia	2016	Female	30 to 34	92% (89-95%)	7.5% (5-11%)	2.4 (1.8-3.2)	2.6% (1.8-3.6%)	9.3% (5.8-13%)	31 (19-45)	3200 (2300-4400)
Australia	2016	Female	35 to 39	92% (88-95%)	8.1% (5.4-12%)	3 (2.2-3.9)	3.2% (2.2-4.3%)	11% (7.8-15%)	53 (35-71)	4000 (2800-5300)
Australia	2016	Female	40 to 44	91% (88-94%)	8.8% (6.1-12%)	3.1 (2.3-4.1)	3.6% (2.5-4.7%)	12% (8.7-15%)	92 (67-120)	5600 (4100-7300)
Australia	2016	Female	45 to 49	90% (85-94%)	9.8% (6.4-15%)	2.9 (2.1-3.9)	3.7% (2.5-5%)	11% (8-14%)	110 (83-150)	5800 (4100-7700)
Australia	2016	Female	50 to 54	88% (83-91%)	12% (8.7-17%)	2.7 (2-3.6)	3.7% (2.4-5.1%)	9.5% (6.9-12%)	160 (120-210)	6900 (4600-9200)
Australia	2016	Female	55 to 59	84% (77-89%)	16% (11-23%)	2.7 (2-3.5)	3.6% (2.3-5%)	8.1% (5.8-10%)	180 (130-230)	6500 (4200-8800)
Australia	2016	Female	60 to 64	81% (74-86%)	19% (14-26%)	2.6 (1.9-3.5)	3.1% (1.8-4.6%)	6.4% (4.2-8.5%)	210 (140-290)	6300 (3600-9300)
Australia	2016	Female	65 to 69	79% (72-85%)	21% (15-28%)	2.4 (1.8-3.2)	2.5% (1.2-4%)	4.9% (2.9-6.7%)	210 (130-300)	5100 (2400-7900)
Australia	2016	Female	70 to 74	76% (69-82%)	24% (18-31%)	2.3 (1.6-3)	1.9% (0.41-3.5%)	3.4% (1.2-5.5%)	180 (64-300)	3400 (710-6300)
Australia	2016	Female	75 to 79	72% (61-80%)	28% (20-39%)	2 (1.4-2.7)	1.2% (-0.42-3%)	1.9% (-0.31-4.3%)	130 (-20-300)	1900 (-690-4800)
Australia	2016	Female	80 to 84	69% (57-78%)	31% (22-43%)	1.8 (1.2-2.4)	0.46% (-1.3-2.2%)	0.63% (-1.6-2.9%)	73 (-190-340)	820 (-2200-3900)
Australia	2016	Female	85 to 89	67% (56-76%)	33% (24-44%)	1.6 (1.1-2.3)	-0.16% (-2.1-1.9%)	-0.31% (-2.8-2.3%)	-56 (-500-400)	-290 (-3700-3300)
Australia	2016	Female	90 to 94	67% (56-77%)	33% (23-44%)	1.6 (0.96-2.3)	-0.78% (-2.9-1.6%)	-1.2% (-3.8-1.9%)	-170 (-550-270)	-740 (-2800-1500)
Australia	2016	Female	95 plus	68% (53-81%)	32% (19-47%)	1.5 (0.76-2.4)	-1.2% (-3.6-1.9%)	-1.7% (-4.8-2%)	-160 (-430-180)	-440 (-1300-710)
Australia	2016	Male	All Ages	88% (85-90%)	12% (9.8-15%)	4.8 (3.7-6.1)	8.4% (6.4-10%)	8.2% (5.4-11%)	7000 (4500-9800)	240000 (180000-290000)
Australia	2016	Male	15 to 19	79% (68-88%)	21% (12-32%)	2.2 (1.3-3.4)	4.6% (2.8-6.8%)	12% (5.8-19%)	36 (17-57)	3900 (2400-5600)
Australia	2016	Male	20 to 24	88% (81-93%)	12% (7.3-19%)	4 (2.6-6.1)	9.8% (6.7-13%)	20% (12-30%)	110 (64-160)	11000 (7600-15000)
Australia	2016	Male	25 to 29	90% (85-94%)	10% (6.3-15%)	5.4 (3.7-7.5)	10% (7.7-14%)	23% (15-31%)	150 (98-210)	14000 (10000-18000)
Australia	2016	Male	30 to 34	92% (87-95%)	7.8% (4.8-13%)	6.2 (4.4-8.3)	10% (7.8-14%)	24% (17-31%)	170 (120-220)	13000 (9700-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Australia	2016	Male	35 to 39	95% (92-97%)	5.4% (3.4-8.4%)	6.5 (4.7-8.7)	11% (7.8-14%)	23% (17-30%)	200 (140-270)	13000 (10000-17000)
Australia	2016	Male	40 to 44	96% (93-97%)	4.5% (2.6-7%)	6.7 (4.8-9)	11% (8.1-14%)	23% (17-28%)	300 (210-380)	17000 (13000-22000)
Australia	2016	Male	45 to 49	95% (92-97%)	4.9% (3-8%)	6.9 (5-9.4)	11% (8.5-15%)	21% (16-27%)	360 (260-460)	18000 (14000-23000)
Australia	2016	Male	50 to 54	95% (92-96%)	5.5% (3.6-8.2%)	7.3 (5.3-9.8)	12% (9-15%)	20% (15-24%)	540 (400-670)	24000 (18000-29000)
Australia	2016	Male	55 to 59	94% (91-96%)	6.2% (4-9.2%)	7.8 (5.6-10)	12% (9.6-15%)	19% (15-23%)	680 (530-830)	26000 (20000-31000)
Australia	2016	Male	60 to 64	92% (89-95%)	7.5% (4.9-11%)	7.6 (5.4-10)	11% (8.6-14%)	16% (13-20%)	880 (660-1100)	28000 (21000-35000)
Australia	2016	Male	65 to 69	91% (86-94%)	9.4% (6.2-14%)	6.6 (4.8-8.6)	9.8% (7-12%)	13% (9.2-17%)	960 (680-1200)	26000 (18000-33000)
Australia	2016	Male	70 to 74	90% (86-93%)	10% (7-14%)	5.4 (3.7-7.2)	7.7% (4.4-11%)	9.8% (5.5-13%)	840 (460-1200)	18000 (10000-25000)
Australia	2016	Male	75 to 79	90% (85-94%)	9.8% (6.3-15%)	4.6 (3.1-6.2)	5.7% (2.6-9%)	6.7% (2.8-11%)	660 (270-1100)	11000 (4900-17000)
Australia	2016	Male	80 to 84	89% (83-94%)	11% (6.4-17%)	3.9 (2.7-5.3)	4.2% (1.3-7.5%)	4.6% (1.1-8.7%)	640 (160-1200)	7900 (2500-14000)
Australia	2016	Male	85 to 89	88% (81-92%)	12% (7.7-19%)	3.3 (2.2-4.5)	2.6% (-0.027-5.7%)	2.6% (-0.48-6.4%)	390 (-68-970)	3600 (-38-8000)
Australia	2016	Male	90 to 94	87% (80-92%)	13% (7.9-20%)	2.7 (1.8-4)	1.2% (-1.3-4.4%)	0.98% (-2.1-4.7%)	91 (-190-440)	710 (-750-2500)
Australia	2016	Male	95 plus	86% (76-93%)	14% (7-24%)	2.2 (1.2-3.7)	0.21% (-2.3-3.9%)	-0.24% (-3.2-4.2%)	-7.4 (-99-130)	26 (-280-500)
New Zealand	1990	Female	All Ages	88% (86-91%)	12% (9-14%)	1.8 (1.6-2)	1.5% (0.16-2.7%)	-0.93% (-3.6-1.8%)	-120 (-450-230)	6300 (740-12000)
New Zealand	1990	Female	15 to 19	89% (80-95%)	11% (5.3-20%)	1.6 (0.97-2.3)	3.8% (2-5.6%)	9.7% (4.1-15%)	7.6 (3.2-12)	730 (400-1100)
New Zealand	1990	Female	20 to 24	93% (88-96%)	7.2% (3.9-12%)	2.4 (1.5-3.4)	5.1% (3.4-7%)	11% (5.9-18%)	8.8 (4.6-14)	1100 (690-1500)
New Zealand	1990	Female	25 to 29	93% (89-96%)	6.8% (3.8-11%)	2.4 (1.6-3.4)	4.4% (3.1-5.9%)	11% (6.2-16%)	8.7 (4.9-13)	960 (660-1300)
New Zealand	1990	Female	30 to 34	93% (89-96%)	6.5% (3.8-11%)	2.5 (1.7-3.5)	3.7% (2.5-5.1%)	9.9% (6.1-14%)	9.2 (5.8-13)	820 (560-1100)
New Zealand	1990	Female	35 to 39	94% (90-96%)	6.4% (3.6-10%)	2.6 (1.7-3.6)	3.8% (2.4-5.2%)	10% (6.3-14%)	12 (7.3-16)	830 (540-1100)
New Zealand	1990	Female	40 to 44	94% (90-96%)	6.3% (3.6-10%)	2.6 (1.7-3.7)	4.1% (2.5-5.8%)	9.5% (5.6-13%)	17 (10-24)	980 (610-1400)
New Zealand	1990	Female	45 to 49	94% (89-96%)	6.4% (3.5-11%)	2.7 (1.8-3.9)	4.3% (2.3-6.4%)	8.7% (4.6-13%)	21 (11-31)	1000 (560-1400)
New Zealand	1990	Female	50 to 54	92% (87-95%)	7.7% (4.5-13%)	2.7 (1.8-3.8)	4.1% (1.8-6.5%)	7% (3-11%)	24 (10-38)	970 (440-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	1990	Female	55 to 59	90% (84-94%)	9.9% (5.9-16%)	2.7 (1.9-3.7)	3.4% (1-5.8%)	5.3% (1.6-9.1%)	24 (7.4-42)	850 (250-1500)
New Zealand	1990	Female	60 to 64	88% (81-93%)	12% (7.2-19%)	2.5 (1.6-3.7)	2.1% (-0.48-5.1%)	3% (-0.85-7.3%)	21 (-6.1-52)	640 (-140-1500)
New Zealand	1990	Female	65 to 69	87% (78-92%)	13% (7.6-22%)	2.1 (1.5-2.9)	0.42% (-2.1-3.1%)	0.52% (-3.1-4.3%)	5.4 (-33-44)	150 (-720-1100)
New Zealand	1990	Female	70 to 74	85% (76-91%)	15% (9.3-24%)	1.8 (1.2-2.6)	-1.1% (-3.9-1.9%)	-1.6% (-5.3-2.4%)	-23 (-76-34)	-410 (-1400-660)
New Zealand	1990	Female	75 to 79	83% (72-91%)	17% (9.2-28%)	1.5 (0.93-2.3)	-2.2% (-4.8-0.88%)	-3% (-6.5-1%)	-55 (-120-19)	-770 (-1700-300)
New Zealand	1990	Female	80 to 84	81% (68-90%)	19% (10-32%)	1.4 (0.94-2)	-2.8% (-5.4-0.086%)	-3.5% (-6.8-0.092%)	-76 (-150-2)	-780 (-1500-24)
New Zealand	1990	Female	85 to 89	80% (67-89%)	20% (11-33%)	1.3 (0.87-1.9)	-2.9% (-5.4--0.4%)	-3.7% (-6.7--0.59%)	-65 (-120--10)	-470 (-870--67)
New Zealand	1990	Female	90 to 94	80% (68-88%)	20% (12-32%)	1.2 (0.77-1.9)	-3.1% (-5.5--0.65%)	-3.8% (-6.8--0.9%)	-40 (-70--9.2)	-200 (-360--43)
New Zealand	1990	Female	95 plus	80% (65-91%)	20% (9.3-35%)	1.2 (0.59-2)	-3% (-5.5--0.44%)	-3.8% (-6.9--0.67%)	-17 (-31--3)	-54 (-99--8)
New Zealand	1990	Male	All Ages	89% (85-92%)	11% (8.4-15%)	5 (4.4-5.5)	11% (8.3-13%)	11% (7.2-16%)	1600 (990-2100)	54000 (42000-66000)
New Zealand	1990	Male	15 to 19	81% (70-89%)	19% (11-30%)	3.1 (1.9-4.5)	10% (6.4-15%)	17% (9.7-25%)	34 (19-51)	2700 (1700-4000)
New Zealand	1990	Male	20 to 24	88% (80-93%)	12% (6.6-20%)	5.2 (3.5-7.1)	19% (13-25%)	26% (17-37%)	71 (45-100)	5600 (3800-7500)
New Zealand	1990	Male	25 to 29	92% (87-96%)	7.9% (4.4-13%)	6.2 (4.2-8.5)	18% (13-23%)	28% (19-38%)	58 (39-77)	4500 (3200-5700)
New Zealand	1990	Male	30 to 34	95% (90-97%)	5.4% (2.5-10%)	6.9 (4.9-9.4)	15% (11-19%)	26% (19-34%)	48 (35-62)	3400 (2600-4300)
New Zealand	1990	Male	35 to 39	95% (92-98%)	4.7% (2.3-8.3%)	7.3 (4.9-10)	14% (9.8-18%)	25% (17-32%)	48 (32-63)	3100 (2200-3900)
New Zealand	1990	Male	40 to 44	96% (92-98%)	4.5% (2.2-8.1%)	7.7 (5.4-10)	13% (9.4-17%)	22% (15-29%)	58 (39-75)	3200 (2300-4100)
New Zealand	1990	Male	45 to 49	95% (92-98%)	4.6% (2.3-8.5%)	8.2 (5.5-12)	14% (9.5-18%)	21% (14-28%)	70 (46-92)	3400 (2300-4400)
New Zealand	1990	Male	50 to 54	95% (92-98%)	4.9% (2.5-8.4%)	8.5 (5.7-12)	14% (9.5-19%)	19% (13-26%)	93 (60-120)	3800 (2600-5000)
New Zealand	1990	Male	55 to 59	95% (91-97%)	5.3% (2.7-9.2%)	8.5 (6-12)	14% (9.6-19%)	18% (12-24%)	130 (88-180)	4700 (3200-6100)
New Zealand	1990	Male	60 to 64	94% (89-97%)	6.3% (3.3-11%)	8.1 (5.3-11)	13% (7.5-18%)	16% (8.7-22%)	190 (110-270)	5600 (3300-7700)
New Zealand	1990	Male	65 to 69	92% (87-96%)	7.6% (3.8-13%)	7.1 (5-9.5)	12% (6.7-17%)	14% (7.5-19%)	230 (120-330)	5600 (3100-7800)
New Zealand	1990	Male	70 to 74	92% (86-96%)	8.3% (4.4-14%)	5.9 (4.1-8.1)	9.1% (3.7-14%)	10% (3.8-16%)	200 (73-320)	3900 (1600-6100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	1990	Male	75 to 79	92% (85-96%)	8.2% (4.1-15%)	4.9 (3.1-7)	6.7% (1-13%)	7.2% (0.6-14%)	160 (13-320)	2500 (370-4700)
New Zealand	1990	Male	80 to 84	91% (83-96%)	8.8% (4.1-17%)	4.3 (2.9-6)	5.4% (0.83-11%)	5.7% (0.44-12%)	110 (8.3-220)	1200 (190-2500)
New Zealand	1990	Male	85 to 89	90% (82-95%)	9.7% (4.9-18%)	3.7 (2.5-5.3)	3.9% (0.13-8.9%)	3.9% (-0.35-9.9%)	43 (-3.9-110)	370 (12-850)
New Zealand	1990	Male	90 to 94	90% (83-95%)	10% (5-17%)	3.3 (2.1-4.9)	2.5% (-0.84-7.3%)	2.4% (-1.5-7.8%)	9.4 (-5.8-30)	60 (-20-170)
New Zealand	1990	Male	95 plus	89% (79-96%)	11% (4.4-21%)	2.8 (1.5-4.8)	1.5% (-2.1-7.4%)	1.2% (-3.1-8.2%)	1.3 (-3.3-8.9)	6.6 (-9.2-32)
New Zealand	1995	Female	All Ages	88% (86-91%)	12% (9.1-14%)	1.3 (1.1-1.5)	0.67% (-0.36-1.7%)	-1.9% (-4.3-0.34%)	-260 (-570-45)	3000 (-1600-7400)
New Zealand	1995	Female	15 to 19	89% (80-95%)	11% (5.5-20%)	1.2 (0.71-1.7)	3.1% (1.5-4.8%)	7.8% (2.5-13%)	5.3 (1.7-9)	550 (270-830)
New Zealand	1995	Female	20 to 24	93% (87-96%)	7.3% (3.9-13%)	1.7 (1.1-2.6)	4.4% (2.9-6%)	8.9% (3.9-15%)	6.7 (3-11)	930 (610-1300)
New Zealand	1995	Female	25 to 29	93% (88-96%)	6.9% (3.9-12%)	1.8 (1.2-2.6)	3.7% (2.6-5%)	8.5% (4.3-13%)	6.2 (3.2-9.4)	800 (530-1100)
New Zealand	1995	Female	30 to 34	93% (89-96%)	6.6% (3.6-11%)	1.9 (1.3-2.7)	3.1% (2.1-4.2%)	7.7% (4.4-11%)	7.7 (4.5-11)	750 (500-1000)
New Zealand	1995	Female	35 to 39	94% (90-96%)	6.4% (3.6-10%)	1.9 (1.2-2.6)	2.9% (1.8-4%)	7.6% (4.5-11%)	9.4 (5.5-14)	720 (470-1000)
New Zealand	1995	Female	40 to 44	94% (89-96%)	6.4% (3.6-11%)	1.9 (1.2-2.6)	2.8% (1.7-4.2%)	7% (3.9-10%)	12 (6.6-18)	720 (450-1000)
New Zealand	1995	Female	45 to 49	94% (89-96%)	6.5% (3.6-11%)	1.9 (1.3-2.8)	3% (1.6-4.7%)	6.4% (3.4-9.8%)	17 (8.8-26)	830 (460-1200)
New Zealand	1995	Female	50 to 54	92% (87-95%)	7.7% (4.6-13%)	1.9 (1.3-2.7)	2.7% (1.1-4.5%)	4.9% (1.9-8.3%)	18 (6.8-30)	720 (300-1200)
New Zealand	1995	Female	55 to 59	90% (84-94%)	10% (5.8-16%)	1.9 (1.3-2.7)	1.9% (0.3-3.8%)	3.2% (0.54-6.2%)	15 (2.5-28)	490 (81-1000)
New Zealand	1995	Female	60 to 64	88% (81-93%)	12% (7.2-19%)	1.8 (1.1-2.6)	0.77% (-0.95-2.9%)	1.3% (-1.4-4.3%)	8 (-9-27)	220 (-270-790)
New Zealand	1995	Female	65 to 69	87% (78-92%)	13% (7.7-22%)	1.5 (1-2.1)	-0.55% (-2.3-1.2%)	-0.7% (-3.3-1.8%)	-6.8 (-32-18)	-180 (-770-400)
New Zealand	1995	Female	70 to 74	85% (76-91%)	15% (9.1-24%)	1.3 (0.81-1.8)	-1.7% (-3.7-0.26%)	-2.2% (-4.9-0.48%)	-32 (-70-7.1)	-630 (-1300-100)
New Zealand	1995	Female	75 to 79	83% (72-91%)	17% (9.3-28%)	1.1 (0.65-1.6)	-2.6% (-4.8--0.51%)	-3.4% (-6.3--0.71%)	-62 (-110-13)	-900 (-1700--180)
New Zealand	1995	Female	80 to 84	81% (68-90%)	19% (10-32%)	1 (0.66-1.4)	-3.2% (-5.5--1%)	-4% (-6.9--1.2%)	-93 (-160--29)	-990 (-1700--310)
New Zealand	1995	Female	85 to 89	79% (67-88%)	21% (12-33%)	0.93 (0.61-1.3)	-3.3% (-5.6--1.1%)	-4.1% (-7--1.5%)	-88 (-150--31)	-660 (-1100--230)
New Zealand	1995	Female	90 to 94	79% (67-88%)	21% (12-33%)	0.88 (0.53-1.3)	-3.4% (-5.8--1.2%)	-4.2% (-7.1--1.4%)	-56 (-95--19)	-280 (-480--98)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	1995	Female	95 plus	79% (64-90%)	21% (9.9-36%)	0.83 (0.42-1.5)	-3.3% (-5.8--0.99%)	-4.1% (-7--1.2%)	-24 (-41--7)	-76 (-130--23)
New Zealand	1995	Male	All Ages	89% (85-92%)	11% (8.2-15%)	4.8 (4.3-5.3)	10% (7.8-12%)	9.9% (6.1-14%)	1400 (870-2000)	50000 (39000-61000)
New Zealand	1995	Male	15 to 19	81% (70-90%)	19% (10-30%)	2.9 (1.8-4.4)	9.4% (5.9-13%)	16% (8.7-24%)	25 (14-37)	2100 (1300-3000)
New Zealand	1995	Male	20 to 24	88% (80-93%)	12% (6.6-20%)	4.9 (3.2-7)	17% (12-23%)	26% (16-36%)	57 (36-79)	4700 (3200-6300)
New Zealand	1995	Male	25 to 29	92% (86-95%)	8% (4.5-14%)	5.8 (3.9-8.1)	17% (12-22%)	27% (18-36%)	55 (36-73)	4200 (3000-5500)
New Zealand	1995	Male	30 to 34	95% (89-97%)	5.5% (2.6-11%)	6.5 (4.7-8.9)	14% (10-18%)	25% (17-33%)	48 (33-63)	3500 (2600-4400)
New Zealand	1995	Male	35 to 39	95% (92-98%)	4.7% (2.3-8.3%)	6.9 (4.7-9.6)	13% (9-16%)	24% (16-30%)	47 (32-61)	3100 (2200-4000)
New Zealand	1995	Male	40 to 44	96% (92-98%)	4.5% (2.2-8%)	7.3 (5.1-9.8)	12% (8.7-16%)	21% (14-28%)	53 (36-68)	3000 (2100-3900)
New Zealand	1995	Male	45 to 49	95% (91-98%)	4.6% (2.4-8.7%)	7.8 (5.2-11)	13% (8.6-17%)	20% (13-27%)	76 (48-98)	3700 (2500-4800)
New Zealand	1995	Male	50 to 54	95% (91-97%)	4.9% (2.5-9%)	8.1 (5.4-11)	13% (8.8-17%)	19% (12-24%)	94 (61-120)	3900 (2700-5000)
New Zealand	1995	Male	55 to 59	95% (90-97%)	5.4% (2.8-9.6%)	8.1 (5.6-11)	13% (9.1-18%)	18% (12-23%)	120 (80-160)	4300 (2900-5700)
New Zealand	1995	Male	60 to 64	94% (89-97%)	6.4% (3.2-11%)	7.6 (4.9-11)	12% (6.8-16%)	15% (8.1-20%)	150 (83-210)	4600 (2600-6200)
New Zealand	1995	Male	65 to 69	92% (87-96%)	7.7% (3.7-13%)	6.7 (4.7-9)	11% (5.8-15%)	13% (6.5-18%)	200 (110-290)	5000 (2700-7100)
New Zealand	1995	Male	70 to 74	92% (86-96%)	8.3% (4.4-14%)	5.5 (3.8-7.7)	8.1% (3.1-13%)	9.1% (3.2-15%)	190 (64-310)	3700 (1400-5900)
New Zealand	1995	Male	75 to 79	92% (86-96%)	8.2% (4.2-14%)	4.6 (3-6.7)	5.8% (0.77-11%)	6.2% (0.4-12%)	130 (8.5-270)	2100 (280-4000)
New Zealand	1995	Male	80 to 84	91% (84-96%)	8.7% (4.2-16%)	4.1 (2.7-5.6)	4.4% (0.43-9.4%)	4.6% (-0.11-10%)	96 (-2.2-210)	1100 (110-2400)
New Zealand	1995	Male	85 to 89	90% (82-95%)	9.7% (4.9-18%)	3.5 (2.4-5.1)	3.2% (-0.23-7.5%)	3.1% (-0.74-8.3%)	43 (-10-110)	380 (-28-920)
New Zealand	1995	Male	90 to 94	90% (82-95%)	10% (5-18%)	3.1 (2-4.7)	2.1% (-1-6.7%)	1.9% (-1.7-7.2%)	12 (-10-44)	76 (-38-250)
New Zealand	1995	Male	95 plus	89% (79-96%)	11% (4.4-21%)	2.7 (1.4-4.7)	1.2% (-2.2-7%)	0.91% (-3.1-7.5%)	1.3 (-4.5-11)	7.1 (-13-41)
New Zealand	2000	Female	All Ages	88% (86-91%)	12% (9-14%)	1.3 (1.1-1.5)	0.73% (-0.19-1.6%)	-1.8% (-4-0.34%)	-240 (-530-45)	3300 (-850-7400)
New Zealand	2000	Female	15 to 19	89% (80-95%)	11% (5.4-20%)	1.1 (0.69-1.7)	2.7% (1.3-4.2%)	7.5% (2.2-13%)	4.4 (1.2-7.5)	470 (230-720)
New Zealand	2000	Female	20 to 24	93% (87-96%)	7.4% (4-13%)	1.7 (1.1-2.6)	3.7% (2.5-5.1%)	8.7% (3.8-14%)	4.2 (1.8-6.8)	670 (440-940)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	2000	Female	25 to 29	93% (89-96%)	6.9% (3.8-11%)	1.8 (1.2-2.6)	3.4% (2.4-4.6%)	8.3% (3.8-13%)	5.4 (2.5-8.4)	700 (470-970)
New Zealand	2000	Female	30 to 34	93% (89-96%)	6.5% (3.6-11%)	1.9 (1.2-2.7)	2.8% (1.9-3.9%)	7.7% (4.2-11%)	6.5 (3.5-9.9)	670 (440-930)
New Zealand	2000	Female	35 to 39	94% (90-96%)	6.3% (3.6-10%)	1.9 (1.2-2.6)	2.6% (1.7-3.7%)	7.6% (4.3-11%)	8.8 (4.9-13)	700 (460-970)
New Zealand	2000	Female	40 to 44	94% (90-97%)	6.3% (3.5-10%)	1.9 (1.2-2.6)	2.7% (1.6-4%)	7% (3.9-10%)	13 (7.1-19)	770 (480-1100)
New Zealand	2000	Female	45 to 49	94% (89-97%)	6.5% (3.4-11%)	2 (1.3-2.8)	2.8% (1.5-4.3%)	6.5% (3.6-9.9%)	16 (9-25)	800 (450-1200)
New Zealand	2000	Female	50 to 54	92% (88-95%)	7.7% (4.5-12%)	1.9 (1.3-2.7)	2.6% (1.2-4.3%)	5.4% (2.5-8.5%)	20 (9.3-32)	810 (370-1300)
New Zealand	2000	Female	55 to 59	90% (84-94%)	9.9% (5.8-16%)	1.9 (1.3-2.7)	2.1% (0.69-3.8%)	3.8% (1.4-6.9%)	18 (6.4-32)	600 (200-1100)
New Zealand	2000	Female	60 to 64	88% (81-93%)	12% (7.1-19%)	1.8 (1.1-2.6)	1.2% (-0.34-3.1%)	2% (-0.4-5.1%)	13 (-2.5-32)	340 (-99-930)
New Zealand	2000	Female	65 to 69	87% (78-92%)	13% (7.6-22%)	1.5 (1-2.1)	-0.00076% (-1.5-1.6%)	0.19% (-2-2.5%)	1.5 (-17-20)	-0.55 (-440-460)
New Zealand	2000	Female	70 to 74	85% (76-91%)	15% (9.4-24%)	1.3 (0.81-1.8)	-1.4% (-2.8-0.54%)	-1.4% (-3.8-0.93%)	-18 (-47-11)	-380 (-920-180)
New Zealand	2000	Female	75 to 79	83% (71-91%)	17% (9.4-29%)	1.1 (0.65-1.7)	-2.1% (-3.9--0.23%)	-2.7% (-5.2--0.31%)	-49 (-94--5.5)	-740 (-1400--77)
New Zealand	2000	Female	80 to 84	81% (67-90%)	19% (10-33%)	1 (0.64-1.4)	-2.9% (-5.1--0.91%)	-3.7% (-6.5--1.1%)	-82 (-140--24)	-890 (-1500--280)
New Zealand	2000	Female	85 to 89	79% (67-88%)	21% (12-33%)	0.93 (0.6-1.4)	-3.3% (-5.5--1.2%)	-4.1% (-6.9--1.5%)	-99 (-170--37)	-750 (-1200--260)
New Zealand	2000	Female	90 to 94	79% (67-88%)	21% (12-33%)	0.89 (0.54-1.4)	-3.5% (-5.9--1.2%)	-4.3% (-7.2--1.6%)	-70 (-120--26)	-360 (-600--130)
New Zealand	2000	Female	95 plus	79% (64-90%)	21% (9.6-36%)	0.84 (0.42-1.5)	-3.5% (-6.1--1.1%)	-4.4% (-7.4--1.4%)	-32 (-54--10)	-100 (-170--33)
New Zealand	2000	Male	All Ages	89% (86-92%)	11% (8.2-14%)	4.3 (3.8-4.9)	8.5% (6.6-11%)	8.1% (4.9-12%)	1100 (680-1600)	42000 (33000-51000)
New Zealand	2000	Male	15 to 19	81% (70-89%)	19% (11-30%)	2.6 (1.6-4)	7.9% (4.8-11%)	15% (7.8-22%)	18 (9.8-28)	1600 (990-2300)
New Zealand	2000	Male	20 to 24	88% (80-93%)	12% (6.5-20%)	4.4 (2.9-6.3)	15% (10-20%)	24% (15-33%)	36 (22-52)	3200 (2200-4200)
New Zealand	2000	Male	25 to 29	92% (87-96%)	7.9% (4.4-13%)	5.2 (3.5-7.4)	15% (11-20%)	25% (16-34%)	41 (26-56)	3300 (2300-4300)
New Zealand	2000	Male	30 to 34	95% (90-97%)	5.4% (2.6-10%)	5.8 (4.1-7.9)	13% (9.1-17%)	24% (16-32%)	42 (28-57)	3100 (2200-4000)
New Zealand	2000	Male	35 to 39	95% (92-98%)	4.7% (2.3-8.4%)	6.1 (4.2-8.4)	11% (8-15%)	22% (14-29%)	45 (29-59)	3000 (2000-3800)
New Zealand	2000	Male	40 to 44	96% (92-98%)	4.5% (2.2-8.1%)	6.5 (4.5-8.7)	11% (7.5-14%)	20% (13-26%)	52 (33-68)	3000 (2000-3900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	2000	Male	45 to 49	95% (92-98%)	4.6% (2.3-8.4%)	6.9 (4.7-9.6)	11% (7.3-15%)	19% (12-25%)	65 (40-86)	3200 (2200-4300)
New Zealand	2000	Male	50 to 54	95% (91-98%)	4.9% (2.5-8.5%)	7.2 (4.9-9.8)	11% (7.3-15%)	17% (11-22%)	87 (54-110)	3700 (2400-4800)
New Zealand	2000	Male	55 to 59	95% (91-97%)	5.3% (2.7-9.5%)	7.2 (5-10)	12% (7.6-15%)	16% (11-21%)	110 (70-140)	3900 (2600-5100)
New Zealand	2000	Male	60 to 64	94% (89-97%)	6.3% (3.3-11%)	6.8 (4.4-10)	11% (6.1-15%)	14% (7.5-19%)	120 (68-170)	3800 (2200-5200)
New Zealand	2000	Male	65 to 69	92% (87-96%)	7.7% (3.8-13%)	6 (4.2-8.1)	9.2% (4.8-13%)	11% (5.5-16%)	150 (72-210)	3700 (2000-5300)
New Zealand	2000	Male	70 to 74	92% (86-96%)	8.3% (4.3-14%)	4.9 (3.3-6.9)	6.7% (2.5-11%)	7.7% (2.6-13%)	150 (49-240)	3000 (1100-4800)
New Zealand	2000	Male	75 to 79	92% (85-96%)	8.2% (4-15%)	4.2 (2.6-6.1)	4.6% (0.61-9.5%)	5% (0.23-11%)	110 (5.3-240)	1800 (250-3800)
New Zealand	2000	Male	80 to 84	91% (83-96%)	8.8% (4.2-17%)	3.6 (2.4-5)	3.2% (-0.2-7.5%)	3.3% (-0.73-8.3%)	66 (-15-170)	820 (-51-1900)
New Zealand	2000	Male	85 to 89	90% (82-95%)	9.8% (4.9-18%)	3.2 (2.1-4.5)	2% (-0.9-5.6%)	1.9% (-1.6-6%)	30 (-26-99)	290 (-130-810)
New Zealand	2000	Male	90 to 94	90% (82-95%)	10% (5-18%)	2.8 (1.8-4.2)	1.1% (-1.7-5.1%)	0.81% (-2.5-5.5%)	5.9 (-18-40)	49 (-72-230)
New Zealand	2000	Male	95 plus	89% (80-96%)	11% (4.5-20%)	2.4 (1.2-4.2)	0.34% (-2.7-5%)	-0.11% (-3.8-5.4%)	-0.22 (-7.5-11)	2.7 (-21-39)
New Zealand	2005	Female	All Ages	90% (87-92%)	10% (8.3-13%)	1.4 (1.2-1.6)	0.99% (0.14-1.8%)	-1.4% (-3.5-0.55%)	-200 (-500-77)	4700 (650-8500)
New Zealand	2005	Female	15 to 19	90% (82-95%)	10% (4.9-18%)	1.2 (0.74-1.8)	2.6% (1.3-4.1%)	7.5% (2.2-13%)	4.7 (1.4-8.1)	510 (260-780)
New Zealand	2005	Female	20 to 24	94% (90-96%)	6.3% (3.7-10%)	1.8 (1.2-2.7)	3.7% (2.5-5.1%)	9% (3.9-15%)	4.8 (2.1-7.8)	740 (480-1100)
New Zealand	2005	Female	25 to 29	94% (91-96%)	6% (3.6-9.4%)	1.9 (1.2-2.8)	3.2% (2.2-4.3%)	8.7% (4.1-14%)	4 (1.9-6.4)	600 (410-830)
New Zealand	2005	Female	30 to 34	94% (91-96%)	5.9% (3.7-9.2%)	1.9 (1.3-2.8)	2.9% (1.9-4%)	8% (4.4-12%)	6.6 (3.7-10)	680 (440-940)
New Zealand	2005	Female	35 to 39	94% (91-97%)	5.8% (3.5-8.7%)	2 (1.3-2.8)	2.7% (1.7-3.9%)	7.9% (4.5-12%)	9.7 (5.5-15)	750 (490-1100)
New Zealand	2005	Female	40 to 44	94% (92-97%)	5.6% (3.4-8.5%)	2 (1.3-2.8)	2.7% (1.6-4%)	7.4% (4.3-11%)	14 (8.1-21)	860 (530-1200)
New Zealand	2005	Female	45 to 49	94% (91-97%)	5.7% (3.2-9.1%)	2.1 (1.4-2.9)	2.9% (1.6-4.4%)	7.1% (4-10%)	20 (11-29)	970 (570-1400)
New Zealand	2005	Female	50 to 54	93% (90-96%)	6.9% (4.3-10%)	2.1 (1.4-2.9)	2.7% (1.4-4.2%)	6% (3-8.8%)	21 (11-32)	870 (450-1300)
New Zealand	2005	Female	55 to 59	91% (86-95%)	8.8% (5.4-14%)	2 (1.4-2.9)	2.3% (0.99-3.9%)	4.5% (2.1-7.3%)	23 (11-37)	780 (340-1300)
New Zealand	2005	Female	60 to 64	89% (84-93%)	11% (6.7-16%)	1.9 (1.2-2.8)	1.6% (0.19-3.4%)	2.9% (0.6-5.6%)	18 (3.7-35)	500 (61-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	2005	Female	65 to 69	87% (79-92%)	13% (7.6-21%)	1.6 (1.1-2.2)	0.62% (-0.72-2.1%)	1.2% (-0.81-3.4%)	9.8 (-6.6-28)	190 (-220-630)
New Zealand	2005	Female	70 to 74	85% (77-91%)	15% (9-23%)	1.3 (0.85-2)	-0.53% (-1.9-1%)	-0.6% (-2.6-1.7%)	-6.6 (-30-19)	-170 (-600-320)
New Zealand	2005	Female	75 to 79	83% (72-91%)	17% (9.4-28%)	1.2 (0.68-1.8)	-1.5% (-3.1-0.13%)	-2% (-4.2-0.23%)	-33 (-70-4)	-500 (-1000-47)
New Zealand	2005	Female	80 to 84	81% (68-90%)	19% (10-32%)	1.1 (0.68-1.5)	-2.4% (-4.4--0.6%)	-3.1% (-5.6--0.76%)	-76 (-140--18)	-820 (-1500--210)
New Zealand	2005	Female	85 to 89	80% (67-88%)	20% (12-33%)	0.99 (0.63-1.4)	-3% (-5.1--0.92%)	-3.8% (-6.4--1.2%)	-98 (-170--30)	-720 (-1200--220)
New Zealand	2005	Female	90 to 94	80% (68-89%)	20% (11-32%)	0.94 (0.57-1.4)	-3.3% (-5.7--1.1%)	-4.2% (-7.1--1.4%)	-84 (-140--29)	-420 (-710--140)
New Zealand	2005	Female	95 plus	80% (65-91%)	20% (9.3-35%)	0.89 (0.43-1.5)	-3.5% (-6.1--1%)	-4.3% (-7.5--1.4%)	-41 (-71--13)	-130 (-230--41)
New Zealand	2005	Male	All Ages	91% (88-93%)	9.2% (6.9-12%)	4.6 (4-5.2)	8.5% (6.7-10%)	8.2% (5.1-12%)	1100 (700-1600)	42000 (33000-51000)
New Zealand	2005	Male	15 to 19	85% (77-91%)	15% (8.8-23%)	2.8 (1.8-4.3)	8% (5-12%)	15% (8.4-23%)	21 (11-32)	1800 (1100-2600)
New Zealand	2005	Male	20 to 24	90% (83-94%)	10% (5.9-17%)	4.6 (3-6.6)	14% (9.7-19%)	24% (15-34%)	35 (21-49)	3200 (2200-4200)
New Zealand	2005	Male	25 to 29	93% (90-96%)	6.8% (4-10%)	5.4 (3.6-7.6)	14% (10-18%)	26% (17-35%)	32 (20-43)	2700 (1900-3500)
New Zealand	2005	Male	30 to 34	96% (92-98%)	4.4% (2.2-7.8%)	6 (4.2-8.2)	12% (8.8-16%)	24% (16-32%)	34 (23-45)	2600 (1900-3300)
New Zealand	2005	Male	35 to 39	96% (94-98%)	3.7% (1.9-6%)	6.4 (4.4-8.8)	11% (7.9-15%)	22% (15-30%)	43 (27-57)	2900 (1900-3700)
New Zealand	2005	Male	40 to 44	97% (94-98%)	3.5% (1.8-6%)	6.8 (4.7-9.1)	11% (7.6-14%)	21% (13-27%)	56 (36-73)	3200 (2200-4200)
New Zealand	2005	Male	45 to 49	96% (94-98%)	3.6% (1.9-6.1%)	7.2 (4.9-10)	11% (7.6-15%)	19% (12-25%)	70 (45-91)	3500 (2400-4500)
New Zealand	2005	Male	50 to 54	96% (94-98%)	3.9% (2-6.1%)	7.5 (5.1-10)	11% (7.6-15%)	18% (12-23%)	87 (56-110)	3800 (2600-4700)
New Zealand	2005	Male	55 to 59	96% (93-98%)	4.4% (2.4-7.4%)	7.6 (5.3-11)	12% (7.9-15%)	17% (11-21%)	120 (80-150)	4400 (3100-5600)
New Zealand	2005	Male	60 to 64	95% (91-97%)	5.4% (3-8.6%)	7.2 (4.7-10)	11% (6.4-14%)	14% (8.4-19%)	130 (75-160)	3900 (2400-5100)
New Zealand	2005	Male	65 to 69	93% (89-97%)	6.7% (3.4-11%)	6.2 (4.4-8.3)	9.2% (5.3-13%)	12% (6.4-16%)	130 (75-190)	3500 (2100-4800)
New Zealand	2005	Male	70 to 74	92% (88-96%)	7.9% (4.2-12%)	5.1 (3.4-7)	7% (3-11%)	8.3% (3.2-13%)	130 (50-200)	2700 (1200-4100)
New Zealand	2005	Male	75 to 79	92% (86-96%)	8.1% (4-14%)	4.2 (2.7-6.2)	5% (1.2-9.2%)	5.5% (0.81-11%)	120 (17-230)	1900 (470-3600)
New Zealand	2005	Male	80 to 84	91% (84-96%)	8.6% (4-16%)	3.7 (2.5-5.1)	3.5% (0.11-7.5%)	3.7% (-0.3-8.5%)	87 (-7.1-200)	1100 (35-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	2005	Male	85 to 89	90% (82-95%)	9.6% (4.9-18%)	3.2 (2.2-4.6)	2.2% (-0.74-5.9%)	2.1% (-1.4-6.5%)	36 (-24-110)	340 (-110-920)
New Zealand	2005	Male	90 to 94	90% (83-95%)	10% (4.9-17%)	2.8 (1.8-4.2)	1.2% (-1.7-5.1%)	0.86% (-2.6-5.4%)	7.8 (-23-49)	64 (-95-280)
New Zealand	2005	Male	95 plus	90% (79-96%)	10% (4.4-21%)	2.5 (1.3-4.2)	0.43% (-2.6-5.3%)	-0.029% (-3.7-5.7%)	-0.072 (-9.2-14)	4.3 (-26-53)
New Zealand	2010	Female	All Ages	90% (87-92%)	10% (8.3-13%)	1.6 (1.4-1.8)	1.4% (0.56-2.2%)	-0.78% (-2.8-1.1%)	-110 (-400-160)	6700 (2700-11000)
New Zealand	2010	Female	15 to 19	90% (83-95%)	10% (5-17%)	1.3 (0.81-1.9)	2.6% (1.4-4%)	8% (2.4-14%)	4.3 (1.3-7.4)	500 (270-740)
New Zealand	2010	Female	20 to 24	94% (90-96%)	6.3% (3.7-10%)	2 (1.2-3)	4% (2.8-5.4%)	9.6% (4.3-16%)	5.2 (2.3-8.4)	810 (540-1100)
New Zealand	2010	Female	25 to 29	94% (91-96%)	5.9% (3.6-9.3%)	2.1 (1.3-3)	3.6% (2.5-4.8%)	9.3% (4.4-14%)	5.2 (2.4-8.1)	710 (480-990)
New Zealand	2010	Female	30 to 34	94% (91-97%)	5.8% (3.5-9.1%)	2.1 (1.4-3.1)	3% (2.1-4.2%)	8.8% (5-13%)	6.4 (3.6-9.5)	660 (450-900)
New Zealand	2010	Female	35 to 39	94% (91-97%)	5.7% (3.3-8.6%)	2.2 (1.4-3.2)	3% (1.9-4.1%)	9% (5.6-13%)	9.8 (6-14)	780 (510-1100)
New Zealand	2010	Female	40 to 44	95% (92-97%)	5.5% (3.3-8.4%)	2.3 (1.4-3.2)	3.1% (1.9-4.4%)	8.5% (5.1-12%)	15 (9.1-22)	950 (600-1300)
New Zealand	2010	Female	45 to 49	94% (91-97%)	5.6% (3.3-8.9%)	2.3 (1.5-3.3)	3.3% (2-4.8%)	8.3% (4.9-12%)	23 (14-32)	1100 (700-1600)
New Zealand	2010	Female	50 to 54	93% (90-96%)	6.8% (4.3-10%)	2.3 (1.5-3.2)	3.1% (1.7-4.5%)	6.9% (3.9-9.8%)	26 (15-37)	1100 (620-1600)
New Zealand	2010	Female	55 to 59	91% (87-95%)	8.6% (5.4-13%)	2.3 (1.6-3.2)	2.7% (1.4-4.3%)	5.6% (3.2-8.2%)	26 (15-39)	920 (480-1400)
New Zealand	2010	Female	60 to 64	89% (84-93%)	11% (6.6-16%)	2.1 (1.4-3.1)	2.1% (0.64-3.8%)	4% (1.5-6.6%)	27 (10-45)	780 (230-1400)
New Zealand	2010	Female	65 to 69	88% (80-93%)	12% (7.2-20%)	1.8 (1.3-2.4)	1.2% (-0.1-2.6%)	2.2% (0.23-4.3%)	19 (1.9-38)	410 (-37-890)
New Zealand	2010	Female	70 to 74	85% (77-91%)	15% (8.9-23%)	1.5 (0.99-2.2)	0.083% (-1.3-1.7%)	0.36% (-1.6-2.7%)	4 (-19-30)	27 (-420-550)
New Zealand	2010	Female	75 to 79	83% (72-91%)	17% (9.4-28%)	1.3 (0.78-2)	-0.81% (-2.3-0.87%)	-1% (-3.1-1.3%)	-15 (-45-19)	-260 (-710-280)
New Zealand	2010	Female	80 to 84	81% (68-90%)	19% (10-32%)	1.2 (0.77-1.7)	-1.7% (-3.4-0.019%)	-2.2% (-4.5-0.025%)	-51 (-100-0.56)	-570 (-1100-6.6)
New Zealand	2010	Female	85 to 89	80% (67-88%)	20% (12-33%)	1.1 (0.72-1.6)	-2.4% (-4.4--0.53%)	-3.1% (-5.6--0.68%)	-91 (-160--20)	-670 (-1200--150)
New Zealand	2010	Female	90 to 94	79% (67-89%)	21% (11-33%)	1 (0.64-1.6)	-2.9% (-5.1--0.88%)	-3.7% (-6.4--1.2%)	-82 (-140--26)	-410 (-720--120)
New Zealand	2010	Female	95 plus	79% (64-90%)	21% (9.6-36%)	0.96 (0.5-1.6)	-3.2% (-5.7--0.97%)	-4% (-7.1--1.2%)	-46 (-80--14)	-150 (-260--44)
New Zealand	2010	Male	All Ages	91% (88-93%)	9.1% (6.8-12%)	4.7 (4.1-5.4)	8.5% (6.7-10%)	8.2% (5.2-11%)	1200 (760-1700)	43000 (34000-52000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	2010	Male	15 to 19	85% (78-91%)	15% (8.7-22%)	2.9 (1.8-4.3)	8% (4.8-11%)	16% (8.4-24%)	19 (10-29)	1700 (1100-2500)
New Zealand	2010	Male	20 to 24	90% (84-94%)	10% (5.6-16%)	4.7 (3-6.7)	14% (9.7-19%)	25% (15-35%)	35 (21-49)	3200 (2200-4300)
New Zealand	2010	Male	25 to 29	93% (90-96%)	6.8% (4-10%)	5.5 (3.6-7.9)	13% (9.6-17%)	26% (17-35%)	27 (18-37)	2500 (1800-3200)
New Zealand	2010	Male	30 to 34	96% (92-98%)	4.4% (2.2-7.8%)	6.2 (4.2-8.6)	12% (8.6-15%)	24% (17-32%)	28 (19-37)	2200 (1600-2900)
New Zealand	2010	Male	35 to 39	96% (94-98%)	3.7% (1.9-6.2%)	6.6 (4.4-9.2)	11% (8-15%)	23% (15-30%)	38 (25-50)	2600 (1900-3400)
New Zealand	2010	Male	40 to 44	96% (94-98%)	3.5% (1.9-5.9%)	6.9 (4.8-9.5)	11% (7.9-15%)	21% (14-28%)	52 (34-67)	3100 (2100-3900)
New Zealand	2010	Male	45 to 49	96% (94-98%)	3.7% (2-6.2%)	7.3 (5-10)	11% (8-15%)	20% (13-26%)	74 (51-96)	3800 (2700-4800)
New Zealand	2010	Male	50 to 54	96% (94-98%)	3.9% (2.1-6.4%)	7.6 (5.1-10)	11% (7.9-15%)	18% (12-23%)	92 (64-120)	4100 (2900-5200)
New Zealand	2010	Male	55 to 59	96% (93-98%)	4.4% (2.3-7.2%)	7.7 (5.4-10)	12% (7.8-15%)	17% (11-22%)	120 (79-150)	4400 (3100-5600)
New Zealand	2010	Male	60 to 64	95% (91-97%)	5.3% (2.9-8.5%)	7.2 (4.7-10)	14% (6.3-14%)	14% (8.4-18%)	140 (81-180)	4400 (2800-5700)
New Zealand	2010	Male	65 to 69	93% (89-97%)	6.6% (3.3-11%)	6.3 (4.5-8.3)	9.1% (5.6-12%)	12% (7-16%)	140 (84-190)	3700 (2300-5100)
New Zealand	2010	Male	70 to 74	92% (87-96%)	7.8% (4.2-13%)	5.2 (3.5-7.1)	7.3% (3.7-11%)	8.9% (4.2-13%)	140 (65-200)	2900 (1500-4300)
New Zealand	2010	Male	75 to 79	92% (86-96%)	8.1% (4.1-14%)	4.4 (2.8-6.4)	5.4% (1.7-9.5%)	6.2% (1.6-11%)	120 (31-220)	2000 (630-3500)
New Zealand	2010	Male	80 to 84	91% (84-96%)	8.6% (4.2-16%)	3.8 (2.5-5.4)	3.9% (0.66-7.8%)	4.3% (0.24-8.9%)	110 (6-220)	1300 (220-2600)
New Zealand	2010	Male	85 to 89	90% (82-95%)	9.6% (4.8-18%)	3.3 (2.2-4.8)	2.6% (-0.28-6.4%)	2.5% (-0.88-7%)	56 (-19-150)	520 (-58-1300)
New Zealand	2010	Male	90 to 94	90% (83-95%)	9.9% (4.7-17%)	2.9 (1.8-4.4)	1.5% (-1.4-5.3%)	1.2% (-2.2-5.7%)	13 (-23-61)	96 (-84-350)
New Zealand	2010	Male	95 plus	90% (80-96%)	10% (4.3-20%)	2.5 (1.3-4.3)	0.68% (-2.4-5.6%)	0.27% (-3.4-6%)	0.99 (-12-22)	9.8 (-35-81)
New Zealand	2016	Female	All Ages	89% (86-91%)	11% (9.1-14%)	1.7 (1.3-2.2)	1.6% (0.68-2.6%)	-0.31% (-2.2-1.9%)	-49 (-360-310)	8300 (3500-14000)
New Zealand	2016	Female	15 to 19	89% (80-95%)	11% (5.3-20%)	1.4 (0.79-2.2)	2.7% (1.4-4.1%)	8.3% (2.4-15%)	4 (1.2-7)	490 (270-750)
New Zealand	2016	Female	20 to 24	93% (87-96%)	7.3% (4-13%)	2.1 (1.2-3.3)	4.1% (2.8-5.6%)	10% (4.5-16%)	5.3 (2.3-9)	900 (590-1300)
New Zealand	2016	Female	25 to 29	93% (89-96%)	6.8% (3.8-11%)	2.2 (1.3-3.4)	3.5% (2.5-4.7%)	9.7% (4.7-15%)	4.4 (2.2-7.2)	710 (480-990)
New Zealand	2016	Female	30 to 34	93% (89-96%)	6.5% (3.8-11%)	2.3 (1.4-3.5)	3.1% (2.1-4.4%)	9.3% (5.3-14%)	6.4 (3.5-9.9)	690 (450-960)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	2016	Female	35 to 39	94% (90-96%)	6.3% (3.6-10%)	2.3 (1.4-3.6)	3.1% (2-4.3%)	9.7% (5.9-14%)	8.2 (4.7-12)	680 (450-950)
New Zealand	2016	Female	40 to 44	94% (90-96%)	6.3% (3.6-10%)	2.4 (1.5-3.7)	3.3% (2-4.7%)	9.3% (5.6-13%)	16 (8.9-23)	1000 (610-1400)
New Zealand	2016	Female	45 to 49	94% (89-96%)	6.5% (3.6-11%)	2.5 (1.5-3.8)	3.5% (2-5.2%)	9% (5.6-13%)	23 (13-33)	1200 (700-1700)
New Zealand	2016	Female	50 to 54	92% (88-95%)	7.7% (4.7-12%)	2.5 (1.5-3.7)	3.4% (1.9-5.2%)	7.6% (4.3-11%)	32 (18-48)	1400 (760-2100)
New Zealand	2016	Female	55 to 59	90% (84-94%)	10% (5.9-16%)	2.5 (1.6-3.5)	3% (1.5-4.7%)	6.3% (3.6-9.1%)	33 (18-48)	1200 (600-1800)
New Zealand	2016	Female	60 to 64	88% (82-93%)	12% (7.2-18%)	2.3 (1.4-3.4)	2.5% (0.89-4.3%)	4.7% (1.9-7.5%)	34 (14-55)	980 (350-1700)
New Zealand	2016	Female	65 to 69	87% (78-93%)	13% (7.3-22%)	2 (1.3-2.8)	1.6% (0.13-3.2%)	2.9% (0.67-5.3%)	31 (6.8-59)	690 (52-1500)
New Zealand	2016	Female	70 to 74	85% (76-91%)	15% (9.2-24%)	1.6 (1-2.5)	0.46% (-0.94-2.2%)	1% (-1.1-3.5%)	13 (-14-46)	180 (-390-890)
New Zealand	2016	Female	75 to 79	83% (72-91%)	17% (9.3-28%)	1.4 (0.79-2.2)	-0.44% (-1.9-1.5%)	-0.45% (-2.5-2.1%)	-7.1 (-40-33)	-150 (-680-470)
New Zealand	2016	Female	80 to 84	81% (67-90%)	19% (10-33%)	1.3 (0.78-2)	-1.3% (-3-0.65%)	-1.7% (-3.9-0.86%)	-38 (-91-18)	-430 (-1000-210)
New Zealand	2016	Female	85 to 89	79% (67-88%)	21% (12-33%)	1.2 (0.73-1.9)	-2% (-3.9-0.068%)	-2.6% (-4.9-0.0073%)	-79 (-150-0.22)	-590 (-1200-21)
New Zealand	2016	Female	90 to 94	79% (66-89%)	21% (11-34%)	1.1 (0.64-1.8)	-2.5% (-4.6--0.42%)	-3.2% (-5.9--0.61%)	-87 (-160--17)	-430 (-810--73)
New Zealand	2016	Female	95 plus	79% (64-90%)	21% (9.6-36%)	1.1 (0.51-1.9)	-2.8% (-5.2--0.34%)	-3.6% (-6.5--0.58%)	-48 (-88--7.8)	-150 (-280--18)
New Zealand	2016	Male	All Ages	89% (86-92%)	11% (7.9-14%)	4.7 (3.6-6)	8.2% (6-11%)	7.8% (4.3-12%)	1300 (710-2000)	45000 (32000-59000)
New Zealand	2016	Male	15 to 19	81% (71-89%)	19% (11-29%)	2.7 (1.6-4.2)	7.5% (4.6-11%)	15% (7.8-24%)	17 (8.9-27)	1600 (940-2300)
New Zealand	2016	Male	20 to 24	88% (80-94%)	12% (6.5-20%)	4.6 (2.7-7)	14% (9.5-19%)	24% (13-35%)	35 (19-50)	3400 (2300-4700)
New Zealand	2016	Male	25 to 29	92% (87-96%)	8% (4.4-13%)	5.5 (3.3-8.5)	13% (9.5-17%)	26% (17-35%)	28 (18-40)	2600 (1800-3500)
New Zealand	2016	Male	30 to 34	95% (90-97%)	5.5% (2.5-10%)	6.1 (3.9-9.3)	11% (8.3-15%)	24% (16-33%)	27 (17-37)	2200 (1600-2900)
New Zealand	2016	Male	35 to 39	95% (92-98%)	4.7% (2.3-8.4%)	6.5 (4-9.6)	11% (7.5-15%)	23% (14-30%)	30 (19-42)	2100 (1500-2800)
New Zealand	2016	Male	40 to 44	96% (92-98%)	4.5% (2.2-7.9%)	6.8 (4.4-9.9)	11% (7.5-15%)	21% (14-28%)	48 (31-66)	2900 (2000-3800)
New Zealand	2016	Male	45 to 49	95% (91-98%)	4.6% (2.3-8.5%)	7.2 (4.5-11)	11% (7.3-15%)	20% (12-26%)	66 (40-90)	3400 (2200-4500)
New Zealand	2016	Male	50 to 54	95% (91-97%)	4.9% (2.6-8.7%)	7.5 (4.6-11)	11% (6.9-14%)	17% (11-23%)	94 (57-130)	4200 (2700-5600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
New Zealand	2016	Male	55 to 59	95% (90-97%)	5.3% (2.8-9.5%)	7.5 (5-11)	11% (7.7-14%)	17% (11-21%)	120 (81-170)	4700 (3200-6200)
New Zealand	2016	Male	60 to 64	94% (89-97%)	6.3% (3.2-11%)	7.1 (4.3-10)	9.9% (6.2-13%)	14% (8.2-18%)	140 (82-190)	4500 (2800-6100)
New Zealand	2016	Male	65 to 69	92% (87-96%)	7.7% (3.8-13%)	6.2 (4.1-8.8)	8.8% (5.2-12%)	11% (6.5-16%)	180 (100-260)	4700 (2800-6600)
New Zealand	2016	Male	70 to 74	92% (86-96%)	8.2% (4.4-14%)	5.2 (3.3-7.6)	7.2% (3.3-11%)	8.9% (3.8-14%)	170 (73-270)	3600 (1700-5500)
New Zealand	2016	Male	75 to 79	92% (86-96%)	8.1% (4.2-14%)	4.4 (2.6-6.8)	5.5% (1.5-9.8%)	6.4% (1.4-12%)	130 (30-250)	2200 (590-4000)
New Zealand	2016	Male	80 to 84	91% (84-96%)	8.6% (4-16%)	3.8 (2.4-5.6)	4.1% (0.57-8.5%)	4.5% (0.29-9.7%)	120 (7.1-260)	1500 (200-3100)
New Zealand	2016	Male	85 to 89	90% (83-95%)	9.6% (4.9-17%)	3.3 (2-5)	2.8% (-0.32-7.4%)	2.8% (-0.81-8.2%)	80 (-25-230)	720 (-85-1900)
New Zealand	2016	Male	90 to 94	90% (82-95%)	10% (5-18%)	2.9 (1.7-4.6)	1.7% (-1.2-6.1%)	1.5% (-1.9-6.8%)	24 (-31-110)	170 (-120-610)
New Zealand	2016	Male	95 plus	90% (80-96%)	10% (4.3-20%)	2.5 (1.2-4.4)	0.91% (-2.3-6%)	0.56% (-3.1-6.5%)	2.9 (-16-34)	19 (-48-120)
Andorra	1990	Female	All Ages	73% (65-81%)	27% (19-35%)	4.6 (3-6.7)	6.7% (4-9.8%)	11% (5.1-17%)	22 (8.9-36)	420 (240-640)
Andorra	1990	Female	15 to 19	69% (49-85%)	31% (15-51%)	3.3 (1.5-6.2)	3% (1.6-5%)	12% (6-20%)	0.048 (0.022-0.086)	5.3 (2.9-8.8)
Andorra	1990	Female	20 to 24	75% (56-88%)	25% (12-44%)	4.3 (1.9-8.1)	4.1% (2.4-6.1%)	15% (7-23%)	0.087 (0.038-0.14)	9.9 (5.7-15)
Andorra	1990	Female	25 to 29	77% (60-89%)	23% (11-40%)	4.5 (2.1-8.3)	3.8% (2.2-5.6%)	15% (7.4-22%)	0.1 (0.047-0.16)	10 (5.9-15)
Andorra	1990	Female	30 to 34	79% (61-90%)	21% (9.7-39%)	4.8 (2.3-8.7)	3.9% (2.3-5.9%)	16% (9-23%)	0.13 (0.069-0.21)	11 (6.6-16)
Andorra	1990	Female	35 to 39	80% (65-91%)	20% (8.6-35%)	5.2 (2.4-10)	4.6% (2.7-7%)	18% (11-25%)	0.2 (0.1-0.31)	14 (7.7-20)
Andorra	1990	Female	40 to 44	81% (63-92%)	19% (8.3-37%)	5.9 (2.9-10)	5.7% (3.4-8.6%)	19% (12-26%)	0.33 (0.18-0.51)	19 (11-29)
Andorra	1990	Female	45 to 49	81% (60-91%)	19% (8.7-40%)	6.7 (3.1-12)	7% (4.4-10%)	19% (13-27%)	0.45 (0.26-0.72)	23 (13-35)
Andorra	1990	Female	50 to 54	80% (62-91%)	20% (8.7-38%)	7.1 (3.5-13)	7.8% (4.8-11%)	18% (12-26%)	0.66 (0.37-1)	29 (17-44)
Andorra	1990	Female	55 to 59	79% (62-91%)	21% (8.9-38%)	7.3 (3.8-12)	8.4% (5-12%)	17% (11-24%)	0.87 (0.48-1.4)	33 (19-50)
Andorra	1990	Female	60 to 64	78% (57-91%)	22% (8.9-43%)	7 (3.5-12)	8.3% (4.6-12%)	15% (8.9-22%)	1.2 (0.61-1.9)	38 (20-60)
Andorra	1990	Female	65 to 69	76% (55-90%)	24% (10-45%)	6.3 (3.5-10)	8.4% (4.7-12%)	14% (8.1-20%)	1.7 (0.83-2.9)	46 (23-75)
Andorra	1990	Female	70 to 74	73% (52-86%)	27% (14-48%)	5.4 (2.7-9.9)	8% (3.3-13%)	12% (5-19%)	1.8 (0.67-3.2)	37 (14-65)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	1990	Female	75 to 79	69% (49-85%)	31% (15-51%)	4.9 (2.2-9.2)	7.7% (2.1-13%)	10% (2.5-18%)	3 (0.72-5.6)	48 (12-87)
Andorra	1990	Female	80 to 84	66% (44-83%)	34% (17-56%)	4.6 (2.2-8.6)	8.3% (2.4-14%)	11% (3-19%)	3.8 (0.94-7.1)	44 (12-79)
Andorra	1990	Female	85 to 89	63% (43-80%)	37% (20-57%)	4.3 (2-8.1)	8.3% (2.3-15%)	10% (2.6-18%)	3.9 (0.96-7.3)	32 (8.4-58)
Andorra	1990	Female	90 to 94	62% (43-81%)	38% (19-57%)	4.2 (1.9-8.1)	7.6% (1.6-14%)	9.1% (1.7-17%)	2.4 (0.45-4.7)	13 (2.7-25)
Andorra	1990	Female	95 plus	63% (39-83%)	37% (17-61%)	4.2 (1.5-8.8)	7% (0.47-14%)	8.5% (0.4-17%)	1 (0.052-2.1)	3.6 (0.25-7.3)
Andorra	1990	Male	All Ages	80% (71-87%)	20% (13-29%)	5.6 (3.6-8.2)	11% (7.7-14%)	14% (8.1-19%)	27 (16-39)	720 (470-1000)
Andorra	1990	Male	15 to 19	72% (51-88%)	28% (12-49%)	3.6 (1.7-6.3)	7.4% (4.1-11%)	16% (7.9-25%)	0.17 (0.08-0.29)	16 (8.5-25)
Andorra	1990	Male	20 to 24	78% (58-90%)	22% (10-42%)	5.3 (2.6-9.3)	13% (8.4-19%)	23% (13-33%)	0.41 (0.2-0.67)	37 (22-54)
Andorra	1990	Male	25 to 29	81% (67-92%)	19% (8.3-33%)	5.9 (3-9.9)	12% (7.3-17%)	23% (11-34%)	0.43 (0.17-0.71)	36 (20-53)
Andorra	1990	Male	30 to 34	84% (68-94%)	16% (6.1-32%)	6.2 (3.2-11)	11% (5.9-16%)	22% (9.3-32%)	0.43 (0.17-0.74)	32 (17-50)
Andorra	1990	Male	35 to 39	87% (73-95%)	13% (5.1-27%)	6.3 (3.3-11)	11% (5.8-16%)	22% (9.7-31%)	0.53 (0.21-0.93)	34 (16-55)
Andorra	1990	Male	40 to 44	88% (74-96%)	12% (4.2-26%)	6.9 (3.7-11)	12% (6.7-17%)	22% (12-30%)	0.78 (0.36-1.3)	43 (22-68)
Andorra	1990	Male	45 to 49	89% (71-96%)	11% (4-29%)	8.2 (4.3-14)	13% (7.8-18%)	22% (12-29%)	1.1 (0.52-1.7)	51 (28-78)
Andorra	1990	Male	50 to 54	89% (74-96%)	11% (3.9-26%)	9.1 (4.8-15)	14% (9-18%)	21% (14-27%)	1.6 (0.92-2.4)	65 (40-95)
Andorra	1990	Male	55 to 59	89% (76-96%)	11% (3.8-24%)	9.5 (5.4-15)	13% (9.1-18%)	19% (13-25%)	2.1 (1.2-3)	73 (45-100)
Andorra	1990	Male	60 to 64	88% (73-96%)	12% (3.9-27%)	9.2 (4.9-15)	12% (7.9-17%)	16% (10-22%)	2.6 (1.5-3.8)	78 (46-110)
Andorra	1990	Male	65 to 69	88% (73-96%)	12% (4.2-27%)	8.3 (5-13)	12% (7.3-16%)	15% (8.8-20%)	3.2 (1.8-4.5)	79 (46-110)
Andorra	1990	Male	70 to 74	86% (72-95%)	14% (5.4-28%)	7.3 (4-12)	11% (5.1-15%)	13% (5.9-18%)	2.7 (1.2-4.1)	55 (26-82)
Andorra	1990	Male	75 to 79	85% (70-94%)	15% (5.8-30%)	6.6 (3.4-12)	9.6% (3.4-15%)	11% (3.6-17%)	3.6 (1.2-5.7)	57 (20-87)
Andorra	1990	Male	80 to 84	83% (67-93%)	17% (6.9-33%)	6.3 (3.4-11)	9.8% (3.7-15%)	11% (3.8-17%)	3.4 (1.2-5.4)	39 (15-61)
Andorra	1990	Male	85 to 89	81% (65-92%)	19% (7.6-35%)	6 (3.2-10)	9.7% (3.6-15%)	11% (3.9-17%)	2.6 (0.92-4.1)	20 (8-33)
Andorra	1990	Male	90 to 94	82% (66-93%)	18% (7-34%)	5.8 (3.1-10)	9.2% (3-15%)	10% (3-17%)	1.1 (0.33-1.8)	5.9 (2-9.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	1990	Male	95 plus	82% (64-94%)	18% (5.6-36%)	5.8 (2.6-11)	8.9% (2-15%)	10% (2-18%)	0.27 (0.054-0.47)	0.99 (0.23-1.7)
Andorra	1995	Female	All Ages	73% (65-80%)	27% (20-35%)	4.7 (3.3-6.5)	6.7% (4.3-9.5%)	11% (5.5-17%)	27 (12-45)	500 (310-750)
Andorra	1995	Female	15 to 19	69% (49-85%)	31% (15-51%)	3.4 (1.6-6.3)	2.8% (1.6-4.6%)	12% (6.2-20%)	0.047 (0.022-0.083)	5.4 (3-8.7)
Andorra	1995	Female	20 to 24	74% (56-88%)	26% (12-44%)	4.4 (2-8.1)	4% (2.4-5.8%)	15% (7.1-22%)	0.084 (0.039-0.13)	10 (5.9-15)
Andorra	1995	Female	25 to 29	77% (60-89%)	23% (11-40%)	4.6 (2.2-8.1)	3.6% (2.2-5.3%)	14% (6.9-22%)	0.11 (0.051-0.17)	11 (6.9-16)
Andorra	1995	Female	30 to 34	79% (61-91%)	21% (9.2-39%)	4.9 (2.5-8.4)	3.7% (2.2-5.4%)	15% (8.5-22%)	0.15 (0.081-0.24)	13 (8-19)
Andorra	1995	Female	35 to 39	80% (64-91%)	20% (9-36%)	5.3 (2.6-9.4)	4.5% (2.6-6.8%)	17% (10-24%)	0.24 (0.12-0.36)	16 (9.8-24)
Andorra	1995	Female	40 to 44	81% (64-92%)	19% (8.2-36%)	6 (3.1-10)	5.7% (3.6-8.6%)	19% (13-25%)	0.37 (0.22-0.61)	22 (14-34)
Andorra	1995	Female	45 to 49	81% (61-91%)	19% (8.6-39%)	6.9 (3.3-12)	7% (4.5-10%)	19% (14-27%)	0.6 (0.36-0.96)	30 (18-46)
Andorra	1995	Female	50 to 54	80% (62-91%)	20% (8.9-38%)	7.3 (3.8-12)	8% (5.1-11%)	19% (13-26%)	0.77 (0.45-1.2)	34 (21-51)
Andorra	1995	Female	55 to 59	79% (62-91%)	21% (8.7-38%)	7.5 (4-12)	8.5% (5.4-12%)	18% (12-24%)	1.1 (0.61-1.7)	41 (25-62)
Andorra	1995	Female	60 to 64	78% (57-91%)	22% (8.9-43%)	7.2 (3.8-12)	8.3% (5.1-12%)	15% (9.7-21%)	1.3 (0.74-2.2)	43 (25-68)
Andorra	1995	Female	65 to 69	76% (55-90%)	24% (10-45%)	6.4 (3.6-10)	8.3% (4.8-12%)	14% (8.5-19%)	1.8 (0.97-3.1)	50 (27-80)
Andorra	1995	Female	70 to 74	73% (54-87%)	27% (13-46%)	5.6 (2.8-9.6)	7.9% (3.4-12%)	12% (5.3-18%)	2.6 (1-4.6)	56 (22-94)
Andorra	1995	Female	75 to 79	69% (49-86%)	31% (14-51%)	5 (2.3-9.3)	7.6% (2.1-13%)	11% (3-17%)	2.6 (0.72-4.9)	43 (11-77)
Andorra	1995	Female	80 to 84	66% (44-83%)	34% (17-56%)	4.7 (2.2-8.6)	8.2% (2.7-14%)	11% (3.3-18%)	4.3 (1.3-7.8)	50 (15-90)
Andorra	1995	Female	85 to 89	63% (43-80%)	37% (20-57%)	4.5 (2.2-8.1)	8.4% (2.8-14%)	10% (3.1-18%)	5.5 (1.7-10)	44 (14-78)
Andorra	1995	Female	90 to 94	62% (42-80%)	38% (20-58%)	4.4 (2.1-8.1)	7.8% (2.1-14%)	9.4% (2.3-17%)	3.9 (0.93-7.5)	22 (5.6-40)
Andorra	1995	Female	95 plus	63% (38-83%)	37% (17-62%)	4.3 (1.6-8.8)	7.4% (0.86-14%)	8.9% (0.74-17%)	1.6 (0.14-3.3)	5.9 (0.66-11)
Andorra	1995	Male	All Ages	80% (72-87%)	20% (13-28%)	5.8 (4.1-8.1)	11% (8-14%)	14% (9.1-18%)	33 (21-47)	860 (600-1200)
Andorra	1995	Male	15 to 19	72% (52-87%)	28% (13-48%)	3.7 (1.8-6.3)	6.8% (4.2-10%)	16% (8.5-25%)	0.15 (0.071-0.26)	14 (8.3-23)
Andorra	1995	Male	20 to 24	77% (58-91%)	23% (9.4-42%)	5.4 (2.7-9.5)	13% (8.3-18%)	23% (13-33%)	0.38 (0.2-0.61)	36 (23-51)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	1995	Male	25 to 29	81% (66-93%)	19% (7.3-34%)	6 (3.2-9.7)	12% (7.3-16%)	23% (12-33%)	0.44 (0.21-0.72)	39 (23-55)
Andorra	1995	Male	30 to 34	84% (68-94%)	16% (5.9-32%)	6.4 (3.4-10)	10% (5.9-15%)	21% (8.7-31%)	0.5 (0.18-0.85)	38 (20-59)
Andorra	1995	Male	35 to 39	86% (72-95%)	14% (5-28%)	6.4 (3.6-10)	11% (5.3-16%)	21% (8.1-30%)	0.62 (0.19-1.1)	41 (18-64)
Andorra	1995	Male	40 to 44	88% (73-96%)	12% (4.1-27%)	7.1 (4-11)	12% (6.4-17%)	22% (11-30%)	0.9 (0.38-1.5)	50 (24-78)
Andorra	1995	Male	45 to 49	88% (71-96%)	12% (4.1-29%)	8.4 (4.6-14)	13% (8.2-18%)	23% (14-29%)	1.4 (0.76-2.2)	68 (40-100)
Andorra	1995	Male	50 to 54	89% (75-96%)	11% (4-25%)	9.3 (5.4-15)	14% (9.5-18%)	22% (15-27%)	1.9 (1.1-2.7)	77 (50-110)
Andorra	1995	Male	55 to 59	89% (76-96%)	11% (3.9-24%)	9.8 (5.5-15)	14% (9.8-18%)	20% (14-25%)	2.5 (1.6-3.6)	90 (59-120)
Andorra	1995	Male	60 to 64	88% (73-96%)	12% (4.2-27%)	9.5 (5.4-15)	13% (8.7-17%)	17% (12-22%)	3 (1.9-4.3)	90 (58-130)
Andorra	1995	Male	65 to 69	88% (73-96%)	12% (4.3-27%)	8.6 (5.2-13)	12% (8-16%)	15% (9.9-20%)	3.7 (2.3-5.4)	94 (60-130)
Andorra	1995	Male	70 to 74	87% (73-95%)	13% (5.3-27%)	7.5 (4.2-12)	11% (5.6-15%)	13% (6.6-18%)	4.1 (2-6.1)	83 (42-120)
Andorra	1995	Male	75 to 79	85% (71-94%)	15% (5.8-29%)	6.9 (3.5-11)	9.8% (4-14%)	11% (4.3-17%)	3.3 (1.3-5.1)	53 (22-80)
Andorra	1995	Male	80 to 84	84% (67-93%)	16% (6.7-33%)	6.5 (3.6-11)	10% (4.5-15%)	11% (4.7-17%)	4 (1.6-6.1)	46 (20-69)
Andorra	1995	Male	85 to 89	82% (64-93%)	18% (7.4-36%)	6.1 (3.4-10)	10% (4.5-16%)	11% (4.7-18%)	3.6 (1.5-5.8)	29 (13-46)
Andorra	1995	Male	90 to 94	82% (65-93%)	18% (6.8-35%)	6 (3.3-10)	9.4% (3.8-15%)	11% (3.9-17%)	1.8 (0.66-2.9)	9.9 (3.8-16)
Andorra	1995	Male	95 plus	82% (62-95%)	18% (5.4-38%)	5.9 (2.7-11)	9.2% (2.6-15%)	10% (2.6-18%)	0.48 (0.12-0.82)	1.7 (0.49-2.9)
Andorra	2000	Female	All Ages	73% (65-80%)	27% (20-35%)	4.5 (3.6-5.7)	6% (4.1-8.3%)	10% (5.4-15%)	23 (11-39)	430 (280-650)
Andorra	2000	Female	15 to 19	69% (50-85%)	31% (15-50%)	3.2 (1.6-5.7)	2.5% (1.4-3.9%)	12% (5.9-19%)	0.037 (0.018-0.063)	4.6 (2.7-7.4)
Andorra	2000	Female	20 to 24	75% (57-88%)	25% (12-43%)	4.2 (2-7.4)	3.7% (2.3-5.3%)	14% (7.1-22%)	0.066 (0.03-0.11)	8.4 (5.1-12)
Andorra	2000	Female	25 to 29	77% (60-89%)	23% (11-40%)	4.4 (2.2-7.6)	3.3% (2.1-4.9%)	14% (7-21%)	0.082 (0.04-0.13)	9.4 (5.8-14)
Andorra	2000	Female	30 to 34	79% (61-91%)	21% (9.4-39%)	4.7 (2.5-7.8)	3.4% (2.1-4.8%)	15% (9.2-22%)	0.13 (0.07-0.2)	12 (7.5-17)
Andorra	2000	Female	35 to 39	80% (64-91%)	20% (9.3-36%)	5.1 (2.6-8.5)	4.1% (2.5-6%)	17% (11-24%)	0.22 (0.12-0.33)	16 (9.8-23)
Andorra	2000	Female	40 to 44	81% (64-92%)	19% (8.3-36%)	5.7 (3.2-9)	5.1% (3.2-7.8%)	18% (13-25%)	0.33 (0.21-0.5)	21 (13-30)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	2000	Female	45 to 49	81% (61-92%)	19% (8.4-39%)	6.5 (3.5-11)	6.3% (4.1-9.4%)	19% (14-27%)	0.52 (0.32-0.81)	27 (17-41)
Andorra	2000	Female	50 to 54	80% (62-91%)	20% (8.9-38%)	7 (4-11)	7.2% (4.7-10%)	18% (13-25%)	0.74 (0.45-1.2)	34 (21-50)
Andorra	2000	Female	55 to 59	79% (63-91%)	21% (8.6-37%)	7.2 (4-11)	7.7% (5-11%)	17% (12-23%)	0.87 (0.52-1.4)	35 (22-52)
Andorra	2000	Female	60 to 64	78% (58-91%)	22% (8.9-42%)	6.9 (3.7-11)	7.5% (4.6-11%)	15% (9.7-20%)	1.1 (0.65-1.8)	38 (22-59)
Andorra	2000	Female	65 to 69	76% (54-90%)	24% (10-46%)	6.2 (3.6-9.4)	7.4% (4.5-10%)	13% (8.3-18%)	1.4 (0.8-2.3)	40 (23-63)
Andorra	2000	Female	70 to 74	73% (54-87%)	27% (13-46%)	5.4 (2.7-8.9)	7% (3.2-11%)	11% (5.2-17%)	1.9 (0.87-3.4)	44 (20-74)
Andorra	2000	Female	75 to 79	69% (49-86%)	31% (14-51%)	4.8 (2.3-8.5)	6.8% (2.3-11%)	9.6% (3.2-16%)	2.7 (0.88-5.1)	47 (15-84)
Andorra	2000	Female	80 to 84	66% (44-83%)	34% (17-56%)	4.5 (2.3-7.5)	7.3% (2.6-12%)	9.6% (3.1-16%)	3.2 (1-5.9)	39 (13-69)
Andorra	2000	Female	85 to 89	63% (43-80%)	37% (20-57%)	4.3 (2.2-7.4)	7.6% (2.6-13%)	9.5% (3.1-17%)	4.1 (1.3-7.6)	34 (11-61)
Andorra	2000	Female	90 to 94	62% (43-80%)	38% (20-57%)	4.2 (1.9-7.4)	7.1% (1.9-13%)	8.6% (2.2-16%)	3.7 (0.89-7)	21 (5.5-38)
Andorra	2000	Female	95 plus	63% (39-83%)	37% (17-61%)	4.1 (1.5-8.5)	6.7% (0.84-13%)	8.2% (0.69-16%)	1.8 (0.15-3.6)	6.6 (0.77-13)
Andorra	2000	Male	All Ages	81% (72-87%)	19% (13-28%)	5.7 (4.4-7.1)	11% (8.5-13%)	14% (9.4-18%)	31 (21-43)	830 (620-1100)
Andorra	2000	Male	15 to 19	72% (52-87%)	28% (13-48%)	3.5 (1.8-5.8)	6% (3.8-9%)	15% (8-24%)	0.11 (0.056-0.19)	12 (7-18)
Andorra	2000	Male	20 to 24	78% (58-90%)	22% (10-42%)	5.2 (2.8-8.5)	12% (8.3-17%)	23% (13-33%)	0.3 (0.16-0.47)	30 (20-42)
Andorra	2000	Male	25 to 29	81% (66-92%)	19% (8.3-34%)	5.8 (3.2-9)	11% (7.6-15%)	23% (13-32%)	0.36 (0.19-0.56)	33 (22-45)
Andorra	2000	Male	30 to 34	84% (67-94%)	16% (6.2-33%)	6.1 (3.5-9.2)	10% (6.6-14%)	23% (14-32%)	0.47 (0.26-0.73)	37 (23-52)
Andorra	2000	Male	35 to 39	87% (73-95%)	13% (5.2-27%)	6.2 (3.7-9.5)	10% (6.7-14%)	23% (13-31%)	0.64 (0.32-0.99)	42 (25-61)
Andorra	2000	Male	40 to 44	88% (74-96%)	12% (4.1-26%)	6.8 (4.2-10)	12% (7.5-16%)	24% (15-31%)	0.91 (0.48-1.4)	51 (30-74)
Andorra	2000	Male	45 to 49	89% (72-96%)	11% (4.1-28%)	8.1 (4.8-12)	13% (9.1-18%)	24% (16-30%)	1.4 (0.81-2)	65 (41-91)
Andorra	2000	Male	50 to 54	89% (75-96%)	11% (3.8-25%)	8.9 (5.6-13)	14% (10-18%)	23% (17-28%)	2 (1.3-2.9)	83 (56-110)
Andorra	2000	Male	55 to 59	89% (76-96%)	11% (3.7-24%)	9.4 (5.7-14)	14% (10-17%)	20% (15-25%)	2.3 (1.6-3.2)	84 (57-110)
Andorra	2000	Male	60 to 64	88% (74-96%)	12% (3.9-26%)	9.1 (5.4-14)	12% (8.7-16%)	17% (12-22%)	2.9 (1.9-4)	88 (61-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	2000	Male	65 to 69	87% (73-96%)	13% (4.3-27%)	8.2 (5.2-12)	12% (8.1-15%)	15% (11-20%)	3.3 (2.2-4.8)	86 (59-120)
Andorra	2000	Male	70 to 74	87% (72-95%)	13% (5.4-28%)	7.2 (4.1-11)	10% (5.8-14%)	13% (6.9-18%)	3.8 (2-5.6)	79 (46-110)
Andorra	2000	Male	75 to 79	85% (70-94%)	15% (5.8-30%)	6.5 (3.5-10)	9.3% (4.2-13%)	11% (4.5-16%)	4 (1.8-6.1)	65 (29-94)
Andorra	2000	Male	80 to 84	83% (67-93%)	17% (7-33%)	6.2 (3.6-9.6)	9.4% (4.5-14%)	11% (5-16%)	3.5 (1.6-5.3)	41 (19-60)
Andorra	2000	Male	85 to 89	82% (65-92%)	18% (7.9-35%)	5.9 (3.5-9.2)	9.5% (4.5-15%)	11% (4.9-17%)	3 (1.4-4.7)	25 (12-38)
Andorra	2000	Male	90 to 94	82% (67-93%)	18% (7.1-33%)	5.7 (3.2-9.5)	9% (3.9-14%)	10% (4-16%)	1.8 (0.72-3)	10 (4.3-16)
Andorra	2000	Male	95 plus	82% (63-94%)	18% (5.7-37%)	5.6 (2.7-10)	8.8% (2.7-15%)	10% (2.6-17%)	0.56 (0.15-0.98)	2.1 (0.64-3.5)
Andorra	2005	Female	All Ages	73% (65-81%)	27% (19-35%)	4.1 (3.4-4.7)	5.1% (3.7-7%)	8.5% (4.3-13%)	22 (11-36)	420 (290-600)
Andorra	2005	Female	15 to 19	69% (49-85%)	31% (15-51%)	2.8 (1.4-4.9)	2.2% (1.3-3.4%)	10% (5.1-17%)	0.033 (0.016-0.058)	4.5 (2.7-7)
Andorra	2005	Female	20 to 24	75% (56-88%)	25% (12-44%)	3.7 (1.8-6.3)	3.3% (2.1-4.9%)	13% (6.3-20%)	0.061 (0.027-0.1)	8.5 (5.2-13)
Andorra	2005	Female	25 to 29	77% (60-90%)	23% (10-40%)	3.9 (2.1-6.6)	3% (1.9-4.4%)	13% (6.9-20%)	0.074 (0.036-0.12)	9.1 (5.6-13)
Andorra	2005	Female	30 to 34	79% (61-90%)	21% (9.7-39%)	4.1 (2.3-6.8)	3% (1.9-4.3%)	14% (8.7-20%)	0.11 (0.063-0.18)	11 (6.9-16)
Andorra	2005	Female	35 to 39	80% (65-91%)	20% (8.8-35%)	4.5 (2.4-7.2)	3.6% (2.2-5.4%)	16% (11-22%)	0.21 (0.12-0.33)	16 (9.5-23)
Andorra	2005	Female	40 to 44	81% (63-92%)	19% (8.3-37%)	5 (2.9-7.7)	4.4% (2.8-6.7%)	17% (12-23%)	0.35 (0.22-0.51)	22 (13-32)
Andorra	2005	Female	45 to 49	81% (61-91%)	19% (8.7-39%)	5.7 (3.2-9.2)	5.6% (3.6-8.1%)	18% (13-25%)	0.52 (0.33-0.8)	28 (18-42)
Andorra	2005	Female	50 to 54	80% (62-91%)	20% (8.8-38%)	6.2 (3.6-9.3)	6.3% (4.2-9.1%)	17% (12-24%)	0.7 (0.45-1.1)	33 (22-51)
Andorra	2005	Female	55 to 59	79% (62-91%)	21% (8.7-38%)	6.3 (3.7-9.5)	6.7% (4.3-9.7%)	16% (11-21%)	0.91 (0.58-1.4)	37 (24-54)
Andorra	2005	Female	60 to 64	78% (57-91%)	22% (8.8-43%)	6.1 (3.3-9.7)	6.5% (3.9-9.5%)	13% (8.6-19%)	0.98 (0.6-1.5)	35 (21-51)
Andorra	2005	Female	65 to 69	76% (56-90%)	24% (10-44%)	5.4 (3.3-8.1)	6.2% (3.8-9.1%)	11% (7.4-16%)	1.3 (0.77-2)	38 (23-59)
Andorra	2005	Female	70 to 74	73% (53-86%)	27% (14-47%)	4.7 (2.5-7.6)	5.8% (2.8-9.3%)	9.5% (4.9-15%)	1.6 (0.78-2.8)	39 (18-66)
Andorra	2005	Female	75 to 79	69% (49-85%)	31% (15-51%)	4.2 (2.1-7.3)	5.6% (1.9-9.4%)	8% (2.9-14%)	2.3 (0.77-4.2)	41 (13-73)
Andorra	2005	Female	80 to 84	66% (44-83%)	34% (17-56%)	3.9 (2.1-6.5)	5.9% (2-10%)	7.9% (2.6-14%)	3.1 (1-5.7)	39 (13-69)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	2005	Female	85 to 89	63% (43-80%)	37% (20-57%)	3.7 (1.9-6.3)	6.2% (1.8-11%)	7.7% (2.2-14%)	4.1 (1.2-7.8)	35 (10-64)
Andorra	2005	Female	90 to 94	62% (43-80%)	38% (20-57%)	3.6 (1.7-6.5)	5.8% (1.3-11%)	7% (1.4-14%)	3.7 (0.75-7.2)	21 (4.9-40)
Andorra	2005	Female	95 plus	63% (39-83%)	37% (17-61%)	3.6 (1.3-7.3)	5.6% (0.32-12%)	6.8% (0.28-15%)	2.2 (0.091-4.8)	8.3 (0.52-17)
Andorra	2005	Male	All Ages	81% (73-88%)	19% (12-27%)	5.1 (4.3-6)	10% (7.9-12%)	12% (8.3-16%)	34 (23-47)	900 (680-1100)
Andorra	2005	Male	15 to 19	72% (51-88%)	28% (12-49%)	3.1 (1.7-4.9)	5.5% (3.4-8.1%)	14% (7.1-21%)	0.11 (0.054-0.18)	12 (7.2-18)
Andorra	2005	Male	20 to 24	78% (60-91%)	22% (9.4-40%)	4.6 (2.5-7.3)	11% (7.6-16%)	21% (12-32%)	0.3 (0.16-0.46)	31 (20-43)
Andorra	2005	Male	25 to 29	81% (66-92%)	19% (7.5-34%)	5.1 (3.1-7.7)	11% (7.5-14%)	22% (13-31%)	0.35 (0.19-0.53)	33 (22-45)
Andorra	2005	Male	30 to 34	84% (67-94%)	16% (6-33%)	5.4 (3.3-8)	9.6% (6.6-13%)	22% (13-30%)	0.43 (0.25-0.68)	35 (24-49)
Andorra	2005	Male	35 to 39	86% (72-95%)	14% (5.3-28%)	5.4 (3.4-8.2)	9.8% (6.4-13%)	22% (14-29%)	0.64 (0.37-0.99)	44 (28-62)
Andorra	2005	Male	40 to 44	88% (74-96%)	12% (4.3-26%)	6 (3.9-8.5)	11% (7.3-15%)	23% (16-29%)	1 (0.59-1.5)	57 (36-80)
Andorra	2005	Male	45 to 49	88% (73-96%)	12% (4-27%)	7.1 (4.5-10)	12% (8.5-16%)	23% (16-29%)	1.4 (0.89-2.1)	70 (46-97)
Andorra	2005	Male	50 to 54	89% (75-96%)	11% (4.1-25%)	7.9 (5.3-11)	13% (9.5-17%)	22% (16-27%)	2 (1.3-2.8)	85 (58-110)
Andorra	2005	Male	55 to 59	89% (77-96%)	11% (3.8-23%)	8.3 (5.5-12)	13% (9.5-17%)	20% (14-24%)	2.7 (1.9-3.7)	99 (69-130)
Andorra	2005	Male	60 to 64	88% (73-96%)	12% (3.8-27%)	8.1 (5-12)	12% (8.1-15%)	16% (11-21%)	2.9 (1.9-4)	91 (61-120)
Andorra	2005	Male	65 to 69	88% (73-96%)	12% (4-27%)	7.2 (4.9-10)	11% (7.3-14%)	14% (9.4-18%)	3.6 (2.4-5)	94 (64-130)
Andorra	2005	Male	70 to 74	87% (73-95%)	13% (5.4-27%)	6.3 (3.9-9.2)	9.2% (5-13%)	11% (6-16%)	3.9 (2.1-5.6)	83 (45-120)
Andorra	2005	Male	75 to 79	85% (71-94%)	15% (5.7-29%)	5.7 (3.3-8.9)	8.2% (3.4-13%)	9.6% (3.7-15%)	4.3 (1.7-6.8)	70 (29-110)
Andorra	2005	Male	80 to 84	84% (67-93%)	16% (6.7-33%)	5.4 (3.3-8)	8.2% (3.5-12%)	9.3% (3.7-14%)	4.2 (1.7-6.7)	50 (21-77)
Andorra	2005	Male	85 to 89	82% (65-93%)	18% (7.3-35%)	5.2 (3.1-7.9)	8.2% (3.6-13%)	9.2% (3.7-15%)	3.7 (1.4-6)	30 (13-48)
Andorra	2005	Male	90 to 94	82% (65-93%)	18% (6.7-35%)	5.1 (2.9-8.2)	7.8% (2.9-13%)	8.7% (2.9-15%)	2.1 (0.68-3.5)	11 (4.2-19)
Andorra	2005	Male	95 plus	83% (64-95%)	17% (5.3-36%)	5 (2.3-8.7)	7.7% (1.5-14%)	8.7% (1.4-16%)	0.77 (0.13-1.4)	2.9 (0.59-5.2)
Andorra	2010	Female	All Ages	73% (65-80%)	27% (20-35%)	3.6 (2.9-4.4)	4.5% (3-6.3%)	7% (3-12%)	22 (9.1-38)	420 (270-610)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	2010	Female	15 to 19	69% (50-85%)	31% (15-50%)	2.4 (1.2-4.3)	2% (1.2-3.1%)	9.1% (4.4-15%)	0.03 (0.014-0.053)	4.3 (2.5-6.8)
Andorra	2010	Female	20 to 24	74% (57-88%)	26% (12-43%)	3.2 (1.5-5.6)	3.1% (1.9-4.5%)	11% (5.2-19%)	0.057 (0.025-0.099)	8.4 (5.3-13)
Andorra	2010	Female	25 to 29	77% (61-89%)	23% (11-39%)	3.4 (1.8-5.9)	2.8% (1.7-4.1%)	12% (6.1-18%)	0.071 (0.035-0.12)	9.1 (5.6-13)
Andorra	2010	Female	30 to 34	79% (61-91%)	21% (9.2-39%)	3.6 (2-6.1)	2.8% (1.8-4.2%)	13% (8.2-19%)	0.11 (0.058-0.17)	10 (6.4-15)
Andorra	2010	Female	35 to 39	80% (64-91%)	20% (8.9-36%)	3.9 (2-6.3)	3.3% (1.9-4.9%)	15% (9.5-21%)	0.19 (0.11-0.3)	15 (8.7-21)
Andorra	2010	Female	40 to 44	81% (64-92%)	19% (8.1-36%)	4.4 (2.4-6.9)	4.1% (2.6-6.1%)	16% (11-22%)	0.35 (0.21-0.52)	22 (14-32)
Andorra	2010	Female	45 to 49	81% (61-92%)	19% (8.4-39%)	5 (2.8-8)	5.1% (3.2-7.6%)	16% (12-22%)	0.57 (0.35-0.89)	31 (19-46)
Andorra	2010	Female	50 to 54	80% (62-91%)	20% (8.9-38%)	5.4 (3.2-8.3)	5.7% (3.7-8.4%)	15% (11-21%)	0.74 (0.48-1.2)	35 (23-53)
Andorra	2010	Female	55 to 59	79% (62-91%)	21% (8.8-38%)	5.5 (3.2-8.4)	6.1% (3.8-8.9%)	14% (9.8-20%)	0.91 (0.57-1.4)	37 (22-57)
Andorra	2010	Female	60 to 64	78% (58-91%)	22% (8.8-42%)	5.3 (2.9-8.4)	5.8% (3.4-8.8%)	12% (7.6-17%)	1.1 (0.65-1.8)	39 (22-62)
Andorra	2010	Female	65 to 69	76% (55-90%)	24% (10-45%)	4.7 (2.8-7)	5.4% (3.2-8.1%)	9.9% (6.3-14%)	1.2 (0.68-2.1)	35 (19-56)
Andorra	2010	Female	70 to 74	73% (53-86%)	27% (14-47%)	4.1 (2.2-6.6)	5% (2.2-8.4%)	8.1% (4-13%)	1.6 (0.71-2.8)	37 (16-66)
Andorra	2010	Female	75 to 79	69% (48-86%)	31% (14-52%)	3.6 (1.8-6.2)	4.6% (1.4-8.3%)	6.6% (2.1-12%)	2 (0.65-4.1)	37 (11-72)
Andorra	2010	Female	80 to 84	66% (44-83%)	34% (17-56%)	3.4 (1.8-5.5)	4.8% (1.4-8.6%)	6.3% (1.7-12%)	2.9 (0.8-5.8)	37 (9.8-71)
Andorra	2010	Female	85 to 89	63% (44-80%)	37% (20-56%)	3.3 (1.6-5.3)	4.9% (1-9.7%)	6% (1.2-12%)	4 (0.8-8.4)	34 (7.3-69)
Andorra	2010	Female	90 to 94	62% (43-81%)	38% (19-57%)	3.2 (1.5-5.8)	4.7% (0.58-9.9%)	5.6% (0.5-12%)	3.7 (0.32-8.2)	21 (2.6-45)
Andorra	2010	Female	95 plus	63% (39-84%)	37% (16-61%)	3.1 (1.2-6.2)	4.5% (-0.11-10%)	5.5% (-0.29-13%)	2.2 (-0.12-5.1)	8.1 (-0.19-18)
Andorra	2010	Male	All Ages	81% (73-87%)	19% (13-27%)	4.5 (3.7-5.5)	8.8% (6.6-11%)	10% (6.3-14%)	34 (20-49)	890 (640-1200)
Andorra	2010	Male	15 to 19	72% (50-87%)	28% (13-50%)	2.7 (1.4-4.5)	5% (3.1-7.2%)	12% (6.1-20%)	0.097 (0.045-0.16)	11 (6.9-17)
Andorra	2010	Male	20 to 24	78% (59-90%)	22% (10-41%)	4 (2.2-6.6)	10% (6.9-14%)	19% (9.8-29%)	0.28 (0.14-0.44)	30 (19-43)
Andorra	2010	Male	25 to 29	81% (66-92%)	19% (8.4-34%)	4.5 (2.6-6.9)	9.8% (6.6-13%)	20% (11-29%)	0.32 (0.17-0.5)	32 (21-45)
Andorra	2010	Male	30 to 34	84% (68-94%)	16% (6.1-32%)	4.7 (2.8-7.2)	8.8% (5.8-12%)	20% (12-28%)	0.39 (0.22-0.61)	32 (21-46)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	2010	Male	35 to 39	87% (73-95%)	13% (5.1-27%)	4.7 (2.9-7.3)	8.7% (5.7-12%)	20% (12-27%)	0.55 (0.31-0.89)	39 (25-57)
Andorra	2010	Male	40 to 44	88% (74-96%)	12% (4.2-26%)	5.2 (3.4-7.8)	9.7% (6.3-14%)	21% (14-28%)	0.94 (0.54-1.5)	55 (35-81)
Andorra	2010	Male	45 to 49	89% (72-96%)	11% (4-28%)	6.2 (3.9-9.2)	11% (7.6-16%)	21% (14-27%)	1.5 (0.89-2.3)	74 (48-110)
Andorra	2010	Male	50 to 54	89% (74-96%)	11% (3.9-26%)	6.8 (4.5-9.8)	12% (8.1-16%)	20% (14-25%)	2 (1.2-3)	86 (56-120)
Andorra	2010	Male	55 to 59	89% (76-96%)	11% (3.7-24%)	7.1 (4.8-10)	12% (8-15%)	18% (12-23%)	2.6 (1.6-3.7)	95 (63-130)
Andorra	2010	Male	60 to 64	88% (73-96%)	12% (3.8-27%)	7 (4.3-10)	11% (6.5-14%)	15% (9.1-20%)	3.2 (1.8-4.5)	100 (60-140)
Andorra	2010	Male	65 to 69	88% (73-96%)	12% (4.2-27%)	6.3 (4.2-8.8)	9.5% (5.9-13%)	12% (7.8-17%)	3.3 (1.9-4.8)	88 (52-120)
Andorra	2010	Male	70 to 74	87% (73-95%)	13% (5.4-27%)	5.5 (3.4-8.3)	8% (3.9-12%)	9.9% (4.7-14%)	3.9 (1.7-6)	82 (38-120)
Andorra	2010	Male	75 to 79	85% (70-94%)	15% (5.8-30%)	5 (2.8-7.9)	6.9% (2.5-11%)	8.1% (2.7-13%)	4.1 (1.3-7.1)	68 (24-110)
Andorra	2010	Male	80 to 84	83% (67-93%)	17% (7-33%)	4.7 (2.8-7)	6.7% (2.4-11%)	7.6% (2.5-13%)	4.5 (1.4-7.6)	53 (19-89)
Andorra	2010	Male	85 to 89	82% (66-92%)	18% (7.8-34%)	4.5 (2.7-6.9)	6.6% (2.2-11%)	7.3% (2.2-13%)	3.8 (1.1-6.8)	31 (11-55)
Andorra	2010	Male	90 to 94	82% (66-93%)	18% (6.9-34%)	4.4 (2.5-7.2)	6.2% (1.6-11%)	6.9% (1.5-13%)	2.2 (0.46-4.1)	12 (3-22)
Andorra	2010	Male	95 plus	82% (63-94%)	18% (5.7-37%)	4.3 (2-7.7)	6.2% (0.96-12%)	6.9% (0.68-14%)	0.74 (0.075-1.5)	2.8 (0.44-5.4)
Andorra	2016	Female	All Ages	73% (65-81%)	27% (19-35%)	3.4 (2.5-4.6)	4.3% (2.6-6.4%)	6.3% (2.4-11%)	22 (7.7-41)	420 (240-680)
Andorra	2016	Female	15 to 19	69% (51-85%)	31% (15-49%)	2.3 (1.1-4.1)	1.9% (1.1-2.9%)	8.4% (3.9-14%)	0.026 (0.012-0.047)	3.8 (2.2-6)
Andorra	2016	Female	20 to 24	75% (57-88%)	25% (12-43%)	3 (1.3-5.4)	3% (1.9-4.4%)	11% (4.9-18%)	0.053 (0.023-0.093)	8 (4.9-12)
Andorra	2016	Female	25 to 29	77% (60-89%)	23% (11-40%)	3.2 (1.6-5.6)	2.7% (1.7-4%)	11% (5.8-18%)	0.065 (0.031-0.12)	8.5 (5.3-13)
Andorra	2016	Female	30 to 34	79% (61-91%)	21% (9.1-39%)	3.4 (1.8-6)	2.7% (1.7-4.1%)	12% (7.5-18%)	0.099 (0.052-0.17)	9.7 (5.9-15)
Andorra	2016	Female	35 to 39	80% (64-91%)	20% (8.9-36%)	3.7 (1.9-6.4)	3.2% (1.8-5%)	14% (9.3-20%)	0.17 (0.089-0.28)	13 (7.7-20)
Andorra	2016	Female	40 to 44	81% (64-92%)	19% (8.1-36%)	4.1 (2.2-7)	4% (2.4-6.1%)	15% (11-21%)	0.33 (0.2-0.51)	21 (13-32)
Andorra	2016	Female	45 to 49	81% (62-92%)	19% (8.3-38%)	4.7 (2.5-8.1)	5% (3-7.7%)	16% (11-22%)	0.6 (0.35-0.98)	32 (19-51)
Andorra	2016	Female	50 to 54	80% (62-91%)	20% (8.9-38%)	5.1 (2.8-8.3)	5.6% (3.5-8.5%)	15% (10-21%)	0.77 (0.47-1.3)	36 (22-58)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	2016	Female	55 to 59	79% (63-91%)	21% (8.7-37%)	5.2 (2.9-8)	5.9% (3.5-8.8%)	14% (8.9-19%)	0.89 (0.5-1.5)	36 (20-57)
Andorra	2016	Female	60 to 64	78% (58-91%)	22% (8.9-42%)	5 (2.6-7.9)	5.6% (3-8.6%)	11% (7-16%)	1.3 (0.69-2.2)	45 (23-73)
Andorra	2016	Female	65 to 69	76% (54-90%)	24% (10-46%)	4.4 (2.5-6.8)	5.1% (2.8-8.1%)	9.3% (5.4-14%)	1.1 (0.6-2)	33 (17-58)
Andorra	2016	Female	70 to 74	73% (54-87%)	27% (13-46%)	3.8 (2-6.6)	4.6% (1.8-8%)	7.5% (3.4-13%)	1.6 (0.67-3)	37 (14-69)
Andorra	2016	Female	75 to 79	70% (49-86%)	30% (14-51%)	3.4 (1.7-6.1)	4.2% (1.1-8.1%)	6.1% (1.9-11%)	1.9 (0.56-3.9)	34 (8.8-70)
Andorra	2016	Female	80 to 84	66% (44-83%)	34% (17-56%)	3.2 (1.6-5.5)	4.3% (0.98-8.6%)	5.6% (1.3-11%)	2.6 (0.57-5.7)	33 (7.3-70)
Andorra	2016	Female	85 to 89	63% (44-80%)	37% (20-56%)	3.1 (1.5-5.3)	4.3% (0.66-9%)	5.4% (0.75-11%)	4 (0.53-8.8)	34 (5.1-72)
Andorra	2016	Female	90 to 94	63% (43-81%)	37% (19-57%)	3 (1.3-5.6)	4.2% (0.22-9.3%)	5% (0.084-12%)	3.9 (0.069-9.1)	23 (1-50)
Andorra	2016	Female	95 plus	63% (38-83%)	37% (17-62%)	3 (1-6.3)	4.1% (-0.38-9.9%)	4.9% (-0.53-12%)	2.6 (-0.26-6.3)	9.5 (-0.86-23)
Andorra	2016	Male	All Ages	82% (74-88%)	18% (12-26%)	4.3 (3.1-5.8)	8.2% (5.5-11%)	9.1% (4.7-14%)	35 (17-55)	890 (560-1200)
Andorra	2016	Male	15 to 19	72% (51-88%)	28% (12-49%)	2.5 (1.3-4.3)	4.8% (3-7.2%)	12% (5.6-19%)	0.087 (0.039-0.15)	10 (6-15)
Andorra	2016	Male	20 to 24	78% (60-90%)	22% (9.6-40%)	3.8 (1.9-6.4)	10% (6.6-14%)	18% (8.8-28%)	0.26 (0.12-0.42)	29 (18-42)
Andorra	2016	Male	25 to 29	82% (67-92%)	18% (7.9-33%)	4.2 (2.3-6.8)	9.5% (6.3-13%)	19% (10-28%)	0.3 (0.15-0.49)	30 (19-43)
Andorra	2016	Male	30 to 34	84% (69-94%)	16% (6-31%)	4.4 (2.5-7.2)	8.5% (5.6-12%)	19% (11-27%)	0.37 (0.19-0.6)	31 (19-44)
Andorra	2016	Male	35 to 39	87% (74-95%)	13% (5.1-26%)	4.5 (2.5-7.2)	8.6% (5.4-12%)	19% (12-27%)	0.5 (0.25-0.85)	35 (21-53)
Andorra	2016	Male	40 to 44	88% (75-96%)	12% (4.3-25%)	5 (2.9-7.9)	9.4% (6.1-14%)	20% (13-27%)	0.89 (0.47-1.5)	52 (31-81)
Andorra	2016	Male	45 to 49	89% (74-96%)	11% (3.9-26%)	5.9 (3.5-9.4)	11% (6.8-15%)	20% (13-27%)	1.6 (0.85-2.6)	77 (46-120)
Andorra	2016	Male	50 to 54	89% (77-96%)	11% (4-23%)	6.4 (3.9-9.8)	11% (7.1-15%)	19% (12-24%)	2.1 (1.2-3.3)	88 (54-130)
Andorra	2016	Male	55 to 59	89% (77-97%)	11% (3.5-23%)	6.7 (4.1-10)	11% (7-15%)	17% (11-22%)	2.4 (1.4-3.6)	89 (53-130)
Andorra	2016	Male	60 to 64	88% (73-96%)	12% (3.8-27%)	6.5 (3.8-9.9)	10% (5.7-14%)	14% (7.7-19%)	3.6 (1.8-5.4)	110 (60-160)
Andorra	2016	Male	65 to 69	88% (72-96%)	12% (4.3-28%)	5.9 (3.7-8.9)	8.8% (4.9-12%)	12% (6.2-16%)	3.1 (1.6-4.7)	82 (42-120)
Andorra	2016	Male	70 to 74	87% (74-95%)	13% (5.2-26%)	5.2 (3-8.1)	7.3% (3.3-11%)	9% (3.8-14%)	3.9 (1.5-6.5)	82 (35-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Andorra	2016	Male	75 to 79	86% (72-95%)	14% (5.5-28%)	4.7 (2.6-7.9)	6.3% (2-11%)	7.3% (2.1-13%)	4 (1.1-7.3)	65 (20-120)
Andorra	2016	Male	80 to 84	84% (67-93%)	16% (6.8-33%)	4.5 (2.5-7.1)	6% (1.6-11%)	6.7% (1.5-12%)	4.2 (0.98-7.9)	51 (14-93)
Andorra	2016	Male	85 to 89	82% (65-92%)	18% (7.7-35%)	4.2 (2.3-6.9)	5.9% (1.4-11%)	6.5% (1.4-13%)	4.3 (0.82-8.3)	36 (8.7-67)
Andorra	2016	Male	90 to 94	82% (66-94%)	18% (6.5-34%)	4.1 (2.3-7.2)	5.6% (0.87-11%)	6.2% (0.71-13%)	2.5 (0.28-5.3)	14 (2.2-29)
Andorra	2016	Male	95 plus	82% (64-95%)	18% (5.2-36%)	4.1 (1.8-7.5)	5.6% (0.37-12%)	6.2% (-0.056-14%)	0.92 (-0.008-2)	3.4 (0.21-7.2)
Austria	1990	Female	All Ages	81% (78-84%)	19% (16-22%)	3 (2.7-3.4)	5.7% (3.8-7.7%)	6.6% (3-11%)	2900 (1300-4800)	67000 (45000-91000)
Austria	1990	Female	15 to 19	79% (71-85%)	21% (15-29%)	2.4 (1.7-3.3)	3.7% (2.2-5.3%)	12% (5.9-18%)	9.7 (4.8-15)	1000 (610-1500)
Austria	1990	Female	20 to 24	84% (75-91%)	16% (8.9-25%)	2.9 (2-4.1)	4.8% (3.3-6.4%)	13% (6.9-19%)	15 (8.2-23)	1900 (1300-2600)
Austria	1990	Female	25 to 29	86% (76-92%)	14% (7.7-24%)	2.8 (1.9-3.7)	4% (2.7-5.3%)	12% (6.1-17%)	15 (8.1-22)	1800 (1200-2400)
Austria	1990	Female	30 to 34	87% (77-93%)	13% (6.8-23%)	3 (2.1-4)	4.2% (2.9-5.6%)	13% (8.2-18%)	22 (14-30)	1900 (1300-2500)
Austria	1990	Female	35 to 39	87% (80-94%)	13% (6.5-20%)	3.6 (2.6-4.9)	5.5% (4-7.3%)	16% (12-21%)	37 (27-47)	2400 (1800-3100)
Austria	1990	Female	40 to 44	87% (80-92%)	13% (7.9-20%)	4 (3-5.2)	6.9% (5.2-8.8%)	17% (13-21%)	65 (50-80)	3600 (2800-4500)
Austria	1990	Female	45 to 49	86% (80-91%)	14% (9-20%)	4.2 (2.9-5.5)	7.8% (5.8-10%)	17% (13-21%)	100 (77-120)	4700 (3600-5900)
Austria	1990	Female	50 to 54	86% (81-91%)	14% (9-19%)	4.4 (3.2-5.8)	8.3% (6-11%)	16% (12-19%)	120 (91-150)	5000 (3800-6300)
Austria	1990	Female	55 to 59	87% (82-91%)	13% (8.9-18%)	4.8 (3.5-6.3)	8.5% (6-11%)	14% (10-18%)	150 (110-200)	5300 (3900-7000)
Austria	1990	Female	60 to 64	86% (80-91%)	14% (9.4-20%)	4.7 (3.4-6.3)	7.9% (4.8-11%)	12% (7.6-17%)	210 (130-300)	6500 (4100-9200)
Austria	1990	Female	65 to 69	84% (78-89%)	16% (11-22%)	4.4 (3.3-5.5)	7.2% (4.2-10%)	10% (5.9-15%)	340 (200-500)	8600 (5200-12000)
Austria	1990	Female	70 to 74	80% (73-86%)	20% (14-27%)	3.8 (2.8-5.1)	6.3% (2.8-10%)	8.1% (3.4-14%)	280 (120-470)	5700 (2500-9200)
Austria	1990	Female	75 to 79	75% (65-84%)	25% (16-35%)	3.4 (2.3-4.7)	5.4% (1.6-9.8%)	6.4% (1.7-12%)	480 (130-900)	7600 (2300-14000)
Austria	1990	Female	80 to 84	73% (61-83%)	27% (17-39%)	3.2 (2.3-4.3)	4.7% (1.1-8.9%)	5.4% (1-10%)	560 (110-1100)	6500 (1500-12000)
Austria	1990	Female	85 to 89	72% (60-82%)	28% (18-40%)	3 (2-4.3)	3.8% (0.27-8%)	4.1% (-0.13-9.2%)	370 (-11-830)	3100 (220-6600)
Austria	1990	Female	90 to 94	71% (59-82%)	29% (18-41%)	2.9 (1.8-4.3)	3.1% (-0.74-8%)	3.3% (-1.4-9.1%)	140 (-57-380)	790 (-190-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	1990	Female	95 plus	71% (55-85%)	29% (15-45%)	2.8 (1.5-4.5)	2.8% (-1.8-8.6%)	3% (-2.5-9.9%)	32 (-27-110)	130 (-80-390)
Austria	1990	Male	All Ages	88% (85-91%)	12% (9.3-15%)	5.2 (4.6-5.8)	15% (12-17%)	17% (13-20%)	6400 (4900-7900)	180000 (150000-210000)
Austria	1990	Male	15 to 19	82% (75-88%)	18% (12-25%)	3.4 (2.4-4.6)	10% (6.5-14%)	17% (9.8-25%)	46 (26-67)	3900 (2500-5400)
Austria	1990	Male	20 to 24	88% (79-94%)	12% (6.3-21%)	4.9 (3.5-6.7)	17% (12-22%)	24% (16-34%)	110 (70-150)	9300 (6700-12000)
Austria	1990	Male	25 to 29	90% (83-95%)	9.6% (5.1-17%)	5.1 (3.8-6.6)	16% (12-20%)	25% (17-33%)	100 (70-140)	8700 (6400-11000)
Austria	1990	Male	30 to 34	92% (85-96%)	8.1% (3.8-15%)	5.5 (4-7)	15% (11-19%)	25% (18-32%)	110 (78-140)	7800 (5800-9800)
Austria	1990	Male	35 to 39	93% (87-97%)	7.1% (3.1-13%)	6 (4.2-8)	16% (12-20%)	27% (19-33%)	130 (91-160)	7900 (5900-9700)
Austria	1990	Male	40 to 44	93% (88-97%)	6.8% (3.3-12%)	6.8 (5.1-8.7)	18% (14-22%)	28% (21-33%)	220 (170-260)	11000 (8900-14000)
Austria	1990	Male	45 to 49	93% (88-96%)	7.2% (3.8-12%)	8 (5.9-10)	20% (16-24%)	29% (23-34%)	350 (280-420)	16000 (13000-19000)
Austria	1990	Male	50 to 54	93% (89-96%)	7% (4-11%)	8.6 (6.3-11)	20% (16-23%)	27% (22-31%)	430 (350-500)	17000 (14000-20000)
Austria	1990	Male	55 to 59	94% (91-96%)	6.2% (3.9-9.3%)	8.7 (6.7-11)	19% (15-23%)	24% (19-29%)	540 (430-650)	18000 (15000-22000)
Austria	1990	Male	60 to 64	94% (91-97%)	5.8% (3.5-9.2%)	8.4 (6.4-11)	17% (13-21%)	21% (16-26%)	720 (550-890)	21000 (16000-25000)
Austria	1990	Male	65 to 69	94% (91-96%)	5.9% (3.6-9.4%)	7.7 (6.1-9.5)	15% (11-19%)	18% (13-22%)	790 (570-990)	19000 (14000-24000)
Austria	1990	Male	70 to 74	94% (90-96%)	6.1% (3.8-9.8%)	7 (5.3-9.1)	14% (8.7-18%)	16% (9.8-21%)	560 (360-760)	11000 (7100-15000)
Austria	1990	Male	75 to 79	93% (89-96%)	6.8% (4-11%)	6.4 (4.7-8.6)	12% (6.6-17%)	13% (7.2-19%)	860 (460-1200)	13000 (7200-18000)
Austria	1990	Male	80 to 84	92% (86-96%)	7.9% (4.3-14%)	6.1 (4.6-7.9)	11% (6.5-16%)	12% (6.8-18%)	800 (440-1100)	8900 (5100-13000)
Austria	1990	Male	85 to 89	91% (85-95%)	9.2% (5.2-15%)	5.8 (4.1-7.9)	10% (5.3-15%)	11% (5.4-17%)	450 (220-670)	3600 (1800-5300)
Austria	1990	Male	90 to 94	91% (84-95%)	9.5% (5.3-16%)	5.7 (3.8-8.2)	9.1% (3.6-15%)	9.9% (3.5-16%)	130 (46-220)	720 (280-1100)
Austria	1990	Male	95 plus	90% (82-96%)	9.8% (4.4-18%)	5.5 (3.2-8.7)	8.5% (1.5-15%)	9.1% (1.1-16%)	23 (2.9-42)	87 (16-150)
Austria	1995	Female	All Ages	82% (78-85%)	18% (15-22%)	2.9 (2.6-3.3)	5.2% (3.4-7.2%)	5.7% (1.9-9.9%)	2500 (850-4300)	59000 (38000-82000)
Austria	1995	Female	15 to 19	79% (70-85%)	21% (15-30%)	2.3 (1.6-3.2)	3.5% (2.1-5.1%)	11% (5.6-17%)	8 (4.1-12)	850 (510-1200)
Austria	1995	Female	20 to 24	85% (75-91%)	15% (8.9-25%)	2.8 (2-4)	4.4% (3-5.9%)	12% (6.5-18%)	10 (5.5-16)	1500 (980-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	1995	Female	25 to 29	86% (77-93%)	14% (7.3-23%)	2.7 (1.9-3.6)	3.8% (2.6-5%)	11% (6-16%)	14 (7.6-20)	1700 (1200-2300)
Austria	1995	Female	30 to 34	87% (78-93%)	13% (6.7-22%)	2.9 (2.1-3.9)	3.8% (2.6-5%)	12% (7.5-17%)	22 (13-30)	1900 (1300-2600)
Austria	1995	Female	35 to 39	88% (80-93%)	12% (6.6-20%)	3.5 (2.5-4.8)	5% (3.6-6.6%)	15% (11-20%)	37 (27-48)	2500 (1900-3300)
Austria	1995	Female	40 to 44	87% (80-92%)	13% (7.6-20%)	3.9 (2.8-5)	6.2% (4.7-8.1%)	16% (13-20%)	57 (43-70)	3200 (2500-4000)
Austria	1995	Female	45 to 49	86% (80-91%)	14% (8.8-20%)	4 (2.9-5.3)	7.6% (5.7-10%)	16% (13-20%)	95 (73-120)	4500 (3400-5600)
Austria	1995	Female	50 to 54	86% (81-91%)	14% (8.9-19%)	4.3 (3.1-5.6)	8.1% (5.9-10%)	15% (12-19%)	130 (100-170)	5500 (4100-6900)
Austria	1995	Female	55 to 59	87% (82-91%)	13% (8.7-18%)	4.6 (3.4-6)	8% (5.8-11%)	14% (10-18%)	140 (110-190)	5300 (3900-6900)
Austria	1995	Female	60 to 64	86% (81-91%)	14% (9.2-19%)	4.5 (3.3-6)	7.5% (4.8-11%)	12% (7.5-16%)	170 (110-240)	5200 (3400-7400)
Austria	1995	Female	65 to 69	85% (79-90%)	15% (10-21%)	4.2 (3.2-5.3)	6.6% (3.9-9.4%)	9.4% (5.4-13%)	240 (140-340)	6200 (3700-8800)
Austria	1995	Female	70 to 74	81% (74-86%)	19% (14-26%)	3.7 (2.7-5)	5.7% (2.4-9.5%)	7.4% (2.9-12%)	360 (140-610)	7500 (3100-12000)
Austria	1995	Female	75 to 79	76% (66-84%)	24% (16-34%)	3.3 (2.2-4.6)	4.7% (1-9%)	5.6% (0.88-11%)	270 (42-540)	4400 (930-8500)
Austria	1995	Female	80 to 84	74% (62-83%)	26% (17-38%)	3.1 (2.2-4.3)	4% (0.42-8.3%)	4.4% (0.079-9.8%)	420 (7.6-940)	5100 (500-11000)
Austria	1995	Female	85 to 89	72% (59-82%)	28% (18-41%)	3 (2-4.2)	3.2% (-0.48-7.5%)	3.3% (-1-8.4%)	320 (-99-830)	2800 (-450-6800)
Austria	1995	Female	90 to 94	71% (59-82%)	29% (18-41%)	2.8 (1.8-4.2)	2.5% (-1.5-7.5%)	2.5% (-2.1-8.6%)	130 (-110-450)	800 (-460-2500)
Austria	1995	Female	95 plus	72% (55-85%)	28% (15-45%)	2.7 (1.4-4.4)	2.3% (-2.5-8.1%)	2.2% (-3.3-9.3%)	35 (-52-150)	140 (-160-510)
Austria	1995	Male	All Ages	88% (86-91%)	12% (9.2-14%)	5 (4.4-5.6)	14% (11-17%)	16% (12-20%)	5900 (4400-7500)	170000 (140000-200000)
Austria	1995	Male	15 to 19	82% (75-88%)	18% (12-25%)	3.2 (2.2-4.3)	9.6% (5.8-14%)	16% (8.6-24%)	37 (20-54)	3100 (2000-4400)
Austria	1995	Male	20 to 24	88% (80-94%)	12% (6.3-20%)	4.7 (3.4-6.4)	16% (12-21%)	23% (15-32%)	82 (52-110)	7200 (5100-9600)
Austria	1995	Male	25 to 29	90% (84-95%)	9.6% (4.8-16%)	5 (3.6-6.5)	14% (11-18%)	23% (15-31%)	88 (58-120)	7700 (5600-9900)
Austria	1995	Male	30 to 34	92% (85-96%)	8.1% (3.8-15%)	5.3 (3.9-6.8)	13% (10-17%)	23% (16-30%)	100 (73-140)	7900 (5800-9900)
Austria	1995	Male	35 to 39	93% (87-97%)	7.2% (3.4-13%)	5.7 (4.1-7.6)	15% (11-18%)	25% (17-31%)	140 (97-170)	8600 (6400-11000)
Austria	1995	Male	40 to 44	93% (88-97%)	6.9% (3.3-12%)	6.5 (4.9-8.3)	17% (13-20%)	27% (21-32%)	190 (150-230)	10000 (8100-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	1995	Male	45 to 49	93% (88-96%)	7.2% (3.8-12%)	7.7 (5.7-10)	19% (15-23%)	28% (23-33%)	320 (250-370)	14000 (12000-17000)
Austria	1995	Male	50 to 54	93% (88-96%)	7% (4-12%)	8.3 (6.1-11)	20% (16-23%)	27% (22-31%)	460 (370-540)	18000 (15000-22000)
Austria	1995	Male	55 to 59	94% (91-96%)	6.2% (3.9-9.3%)	8.4 (6.5-11)	19% (15-22%)	24% (19-29%)	530 (430-630)	18000 (15000-22000)
Austria	1995	Male	60 to 64	94% (90-96%)	5.9% (3.5-9.6%)	8.1 (6.2-11)	17% (13-21%)	21% (16-26%)	630 (480-770)	18000 (14000-23000)
Austria	1995	Male	65 to 69	94% (90-96%)	6% (3.6-9.7%)	7.4 (5.9-9.1)	15% (11-19%)	17% (13-22%)	760 (540-970)	18000 (13000-23000)
Austria	1995	Male	70 to 74	94% (90-96%)	6.2% (3.9-9.8%)	6.7 (5-8.6)	13% (7.8-17%)	15% (8.7-20%)	760 (450-1000)	15000 (9000-20000)
Austria	1995	Male	75 to 79	93% (89-96%)	6.8% (3.8-11%)	6.2 (4.5-8.3)	11% (5.7-16%)	12% (5.9-18%)	500 (240-750)	7800 (4000-11000)
Austria	1995	Male	80 to 84	92% (86-96%)	7.9% (4.2-14%)	5.9 (4.4-7.7)	10% (5.3-15%)	11% (5.4-17%)	680 (330-1000)	7800 (4000-12000)
Austria	1995	Male	85 to 89	91% (84-95%)	9.3% (5.2-16%)	5.6 (4-7.6)	9.4% (4.4-15%)	10% (4.3-16%)	460 (200-740)	3700 (1700-5800)
Austria	1995	Male	90 to 94	90% (84-95%)	9.6% (5.3-16%)	5.5 (3.6-7.9)	8.4% (2.6-14%)	9% (2.3-16%)	160 (41-280)	910 (280-1500)
Austria	1995	Male	95 plus	90% (82-95%)	9.9% (4.5-18%)	5.3 (3.1-8.3)	7.8% (1.1-14%)	8.3% (0.45-16%)	30 (1.6-57)	110 (16-200)
Austria	2000	Female	All Ages	81% (77-84%)	19% (16-23%)	2.8 (2.5-3.2)	4.8% (3.2-6.5%)	4.9% (1.4-8.9%)	2100 (600-3700)	52000 (34000-72000)
Austria	2000	Female	15 to 19	79% (70-87%)	21% (13-30%)	2.2 (1.5-3.1)	3.3% (2-4.6%)	10% (5.2-16%)	7.3 (3.9-11)	820 (520-1200)
Austria	2000	Female	20 to 24	82% (70-91%)	18% (9.4-30%)	2.6 (1.7-3.7)	4.3% (3-5.7%)	11% (5.6-17%)	8 (4.2-12)	1200 (830-1700)
Austria	2000	Female	25 to 29	84% (72-92%)	16% (8-28%)	2.5 (1.7-3.4)	3.6% (2.5-4.6%)	10% (5.7-15%)	9.1 (5-13)	1300 (890-1700)
Austria	2000	Female	30 to 34	85% (73-92%)	15% (7.6-27%)	2.7 (1.9-3.8)	3.4% (2.5-4.6%)	12% (7.3-16%)	17 (11-23)	1600 (1200-2200)
Austria	2000	Female	35 to 39	86% (76-93%)	14% (7.3-24%)	3.3 (2.3-4.5)	4.6% (3.3-6%)	15% (11-19%)	35 (25-44)	2500 (1900-3200)
Austria	2000	Female	40 to 44	85% (76-91%)	15% (8.6-24%)	3.6 (2.7-4.8)	5.9% (4.5-7.7%)	16% (12-20%)	58 (45-72)	3400 (2600-4200)
Austria	2000	Female	45 to 49	84% (76-90%)	16% (9.7-24%)	3.8 (2.7-5)	6.9% (5.2-9%)	16% (12-20%)	80 (61-99)	3900 (3000-4800)
Austria	2000	Female	50 to 54	84% (77-90%)	16% (10-23%)	4 (2.9-5.2)	7.7% (5.7-9.8%)	15% (11-19%)	120 (89-140)	4800 (3700-6100)
Austria	2000	Female	55 to 59	86% (80-91%)	14% (9.2-20%)	4.3 (3.1-5.7)	7.7% (5.5-10%)	14% (10-17%)	150 (110-190)	5400 (4000-7000)
Austria	2000	Female	60 to 64	86% (80-91%)	14% (8.9-20%)	4.4 (3.2-5.9)	7.1% (4.7-10%)	12% (7.9-16%)	160 (110-220)	5000 (3300-7200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	2000	Female	65 to 69	85% (78-90%)	15% (9.8-22%)	4 (3.1-5.2)	6.1% (3.6-8.6%)	9% (5.4-13%)	170 (100-240)	4500 (2700-6400)
Austria	2000	Female	70 to 74	81% (73-88%)	19% (12-27%)	3.6 (2.6-4.8)	5.2% (2.3-8.7%)	7% (2.9-12%)	250 (100-410)	5300 (2300-8700)
Austria	2000	Female	75 to 79	78% (68-86%)	22% (14-32%)	3.3 (2.2-4.6)	4.3% (0.86-8.4%)	5.2% (0.77-10%)	350 (53-700)	5800 (1100-11000)
Austria	2000	Female	80 to 84	75% (63-85%)	25% (15-37%)	3.1 (2.2-4.2)	3.5% (0.084-7.1%)	3.9% (-0.46-8.5%)	250 (-29-540)	3000 (72-6300)
Austria	2000	Female	85 to 89	73% (60-83%)	27% (17-40%)	2.9 (1.9-4)	2.6% (-0.84-6.6%)	2.7% (-1.6-7.6%)	260 (-160-730)	2400 (-750-6000)
Austria	2000	Female	90 to 94	72% (60-83%)	28% (17-40%)	2.7 (1.7-4)	2% (-1.9-6.9%)	1.9% (-2.8-8%)	120 (-180-510)	770 (-740-2800)
Austria	2000	Female	95 plus	72% (55-85%)	28% (15-45%)	2.6 (1.4-4.3)	1.6% (-2.9-7.7%)	1.4% (-4-8.6%)	30 (-84-180)	130 (-250-640)
Austria	2000	Male	All Ages	88% (86-91%)	12% (9.3-14%)	4.9 (4.3-5.5)	13% (11-16%)	15% (11-19%)	5300 (3900-6700)	150000 (120000-180000)
Austria	2000	Male	15 to 19	78% (68-85%)	22% (15-32%)	2.9 (2-4)	8.5% (5.3-12%)	15% (7.7-22%)	29 (16-44)	2700 (1700-3800)
Austria	2000	Male	20 to 24	86% (77-93%)	14% (7.3-23%)	4.4 (3-6)	15% (11-20%)	21% (13-30%)	53 (33-75)	5200 (3700-6900)
Austria	2000	Male	25 to 29	90% (82-95%)	10% (5.2-18%)	4.7 (3.3-6.2)	14% (11-18%)	22% (14-30%)	61 (39-83)	5700 (4100-7400)
Austria	2000	Male	30 to 34	92% (85-96%)	8.2% (3.8-15%)	5 (3.7-6.5)	13% (9.4-16%)	23% (15-30%)	77 (52-100)	6400 (4700-8200)
Austria	2000	Male	35 to 39	93% (87-97%)	7.2% (3.2-13%)	5.4 (4-7.3)	14% (10-17%)	24% (17-31%)	120 (87-160)	8100 (6100-10000)
Austria	2000	Male	40 to 44	93% (87-97%)	6.9% (3.4-13%)	6.2 (4.7-7.8)	16% (12-19%)	26% (20-32%)	190 (150-230)	10000 (8200-12000)
Austria	2000	Male	45 to 49	93% (88-96%)	7.2% (3.8-12%)	7.3 (5.4-9.5)	18% (14-22%)	28% (22-33%)	260 (210-310)	12000 (9900-14000)
Austria	2000	Male	50 to 54	93% (89-96%)	7% (4.1-11%)	7.9 (5.9-10)	19% (15-22%)	26% (21-31%)	410 (330-470)	16000 (13000-19000)
Austria	2000	Male	55 to 59	94% (91-96%)	6.2% (3.9-9.1%)	8 (6.2-10)	18% (14-21%)	24% (19-28%)	530 (430-620)	18000 (15000-22000)
Austria	2000	Male	60 to 64	94% (91-97%)	5.7% (3.4-8.8%)	7.8 (5.9-10)	16% (12-20%)	21% (16-25%)	570 (440-700)	17000 (13000-21000)
Austria	2000	Male	65 to 69	94% (91-96%)	5.8% (3.7-8.8%)	7.1 (5.7-8.7)	14% (10-18%)	17% (13-22%)	620 (450-770)	15000 (11000-19000)
Austria	2000	Male	70 to 74	94% (90-96%)	6.2% (4-9.6%)	6.4 (4.7-8.3)	12% (7.3-16%)	14% (8.1-19%)	690 (400-950)	14000 (8400-19000)
Austria	2000	Male	75 to 79	93% (89-96%)	6.9% (4.1-11%)	5.8 (4.2-7.9)	10% (4.9-15%)	11% (5-17%)	630 (280-960)	10000 (5000-15000)
Austria	2000	Male	80 to 84	92% (86-95%)	8.1% (4.5-14%)	5.6 (4.1-7.3)	9.2% (4.6-14%)	10% (4.6-16%)	400 (180-620)	4600 (2300-7100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	2000	Male	85 to 89	91% (85-95%)	9.2% (5.2-15%)	5.4 (3.8-7.2)	8.4% (3.6-14%)	9% (3.3-15%)	410 (150-680)	3300 (1400-5400)
Austria	2000	Male	90 to 94	90% (84-95%)	9.5% (5.3-16%)	5.3 (3.5-7.5)	7.6% (1.9-13%)	8.1% (1.5-15%)	180 (33-330)	1000 (260-1800)
Austria	2000	Male	95 plus	90% (82-96%)	9.8% (4.3-18%)	5.1 (2.9-8)	7% (0.35-14%)	7.4% (-0.41-15%)	35 (-2-71)	130 (6.7-260)
Austria	2005	Female	All Ages	81% (78-84%)	19% (16-22%)	2.7 (2.4-3.1)	4.2% (2.8-5.8%)	4.2% (0.97-7.7%)	1700 (390-3100)	45000 (29000-62000)
Austria	2005	Female	15 to 19	79% (71-86%)	21% (14-29%)	2.1 (1.5-2.9)	2.9% (1.8-4%)	8.6% (4.5-13%)	5.8 (3-9.1)	710 (450-1000)
Austria	2005	Female	20 to 24	84% (73-91%)	16% (9.2-27%)	2.6 (1.7-3.7)	4% (2.8-5.3%)	9.6% (4.9-15%)	7.1 (3.7-11)	1200 (820-1700)
Austria	2005	Female	25 to 29	85% (75-92%)	15% (8-25%)	2.4 (1.6-3.4)	3.3% (2.4-4.3%)	9.5% (5.3-14%)	6.8 (3.8-9.8)	1100 (750-1500)
Austria	2005	Female	30 to 34	85% (74-93%)	15% (7.5-26%)	2.6 (1.8-3.6)	3.1% (2.2-4.1%)	11% (7.4-15%)	12 (7.8-16)	1300 (890-1700)
Austria	2005	Female	35 to 39	86% (78-93%)	14% (6.9-22%)	3.2 (2.3-4.2)	4% (2.9-5.2%)	14% (10-18%)	28 (20-35)	2100 (1600-2700)
Austria	2005	Female	40 to 44	86% (77-92%)	14% (8.5-23%)	3.5 (2.6-4.5)	5% (3.8-6.4%)	15% (12-18%)	50 (39-60)	3000 (2400-3700)
Austria	2005	Female	45 to 49	85% (78-90%)	15% (9.9-22%)	3.6 (2.6-4.8)	6% (4.6-7.7%)	15% (12-18%)	73 (58-89)	3700 (2900-4600)
Austria	2005	Female	50 to 54	85% (79-90%)	15% (9.7-21%)	3.9 (2.8-5)	6.7% (5-8.5%)	14% (11-17%)	94 (74-120)	4000 (3100-5000)
Austria	2005	Female	55 to 59	86% (80-90%)	14% (9.6-20%)	4.1 (3.1-5.5)	7.2% (5.3-9.3%)	13% (10-16%)	130 (100-170)	4900 (3800-6300)
Austria	2005	Female	60 to 64	86% (80-91%)	14% (9.4-20%)	4.1 (3-5.5)	6.6% (4.5-9.2%)	11% (8-15%)	160 (120-220)	5200 (3600-7000)
Austria	2005	Female	65 to 69	84% (78-89%)	16% (11-22%)	3.9 (2.9-4.9)	5.6% (3.6-7.8%)	8.6% (5.6-12%)	160 (110-230)	4400 (2800-6200)
Austria	2005	Female	70 to 74	80% (73-86%)	20% (14-27%)	3.4 (2.4-4.6)	4.6% (2.2-7.7%)	6.5% (3-10%)	180 (81-280)	3800 (1700-6300)
Austria	2005	Female	75 to 79	76% (66-84%)	24% (16-34%)	3.1 (2-4.4)	3.7% (0.72-7.2%)	4.6% (0.63-9.2%)	230 (32-470)	4000 (770-7700)
Austria	2005	Female	80 to 84	74% (62-84%)	26% (16-38%)	2.9 (2-3.9)	2.9% (-0.065-6%)	3.2% (-0.57-7.2%)	300 (-53-670)	3800 (-86-7900)
Austria	2005	Female	85 to 89	72% (61-82%)	28% (18-39%)	2.7 (1.8-3.8)	2.1% (-1.8-5.5%)	2.1% (-1.8-6.5%)	150 (-130-460)	1400 (-720-3800)
Austria	2005	Female	90 to 94	72% (60-82%)	28% (18-40%)	2.6 (1.6-3.8)	1.5% (-2-5.7%)	1.3% (-3-6.5%)	92 (-200-450)	640 (-870-2500)
Austria	2005	Female	95 plus	72% (56-86%)	28% (14-44%)	2.5 (1.3-4)	1.2% (-3-6.4%)	0.93% (-4.1-7.3%)	26 (-110-200)	130 (-320-710)
Austria	2005	Male	All Ages	89% (86-91%)	11% (8.9-14%)	4.8 (4.2-5.4)	12% (10-15%)	14% (10-17%)	4800 (3600-6100)	140000 (110000-160000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	2005	Male	15 to 19	79% (72-85%)	21% (15-28%)	2.8 (2-3.8)	7.7% (5-11%)	14% (7.3-20%)	24 (13-36)	2300 (1500-3200)
Austria	2005	Male	20 to 24	88% (80-93%)	12% (6.6-20%)	4.3 (3-5.8)	14% (10-18%)	20% (12-28%)	48 (29-67)	5100 (3700-6600)
Austria	2005	Male	25 to 29	91% (85-95%)	9% (4.7-15%)	4.5 (3.3-6)	13% (9.6-16%)	21% (13-29%)	42 (26-59)	4400 (3300-5700)
Austria	2005	Male	30 to 34	93% (87-96%)	7.5% (3.6-13%)	4.8 (3.6-6.3)	12% (8.8-15%)	22% (15-29%)	55 (37-74)	4700 (3500-6000)
Austria	2005	Male	35 to 39	93% (88-97%)	6.8% (3.1-12%)	5.2 (3.8-7.1)	12% (9.4-16%)	23% (17-30%)	98 (71-130)	6800 (5200-8600)
Austria	2005	Male	40 to 44	93% (88-97%)	6.6% (3.3-12%)	6 (4.6-7.6)	14% (11-17%)	26% (20-31%)	160 (130-200)	9300 (7400-11000)
Austria	2005	Male	45 to 49	93% (88-96%)	6.9% (3.7-12%)	7.1 (5.3-9.2)	17% (13-20%)	27% (22-32%)	260 (210-300)	12000 (10000-15000)
Austria	2005	Male	50 to 54	93% (89-96%)	6.7% (4-11%)	7.7 (5.8-9.8)	18% (15-21%)	26% (21-30%)	350 (280-410)	14000 (12000-17000)
Austria	2005	Male	55 to 59	94% (92-96%)	5.9% (3.8-8.5%)	7.8 (6-9.8)	17% (14-21%)	23% (19-28%)	480 (390-560)	17000 (14000-20000)
Austria	2005	Male	60 to 64	94% (91-97%)	5.5% (3.4-8.5%)	7.5 (5.7-9.8)	15% (12-19%)	20% (15-24%)	570 (440-680)	17000 (13000-21000)
Austria	2005	Male	65 to 69	94% (91-96%)	5.8% (3.6-8.9%)	6.8 (5.3-8.4)	13% (10-17%)	17% (13-21%)	580 (440-710)	15000 (11000-18000)
Austria	2005	Male	70 to 74	94% (91-96%)	6.2% (4.1-9.3%)	6.1 (4.5-8)	11% (7.1-15%)	13% (8.3-18%)	560 (340-760)	11000 (7300-15000)
Austria	2005	Male	75 to 79	93% (88-96%)	7.1% (4.1-12%)	5.6 (4-7.5)	9.1% (4.6-14%)	10% (5-16%)	590 (280-890)	9400 (4900-14000)
Austria	2005	Male	80 to 84	92% (86-96%)	8.3% (4.5-14%)	5.3 (3.9-6.8)	8.1% (3.7-13%)	8.9% (3.6-14%)	530 (220-850)	6200 (2800-9600)
Austria	2005	Male	85 to 89	91% (85-95%)	9.3% (5.3-15%)	5.1 (3.6-6.8)	7.4% (2.9-12%)	7.9% (2.6-13%)	260 (88-450)	2200 (850-3600)
Austria	2005	Male	90 to 94	90% (84-95%)	9.6% (5.3-16%)	5 (3.3-7.2)	6.7% (1.4-12%)	7.1% (0.8-14%)	180 (20-340)	1000 (210-1800)
Austria	2005	Male	95 plus	90% (82-95%)	10% (4.6-18%)	4.9 (2.8-7.8)	6.3% (0.067-13%)	6.5% (-0.87-14%)	43 (-5.8-96)	170 (1.8-340)
Austria	2010	Female	All Ages	83% (80-85%)	17% (15-20%)	2.8 (2.3-3.2)	3.9% (2.5-5.5%)	3.9% (0.61-7.3%)	1600 (250-3000)	42000 (27000-59000)
Austria	2010	Female	15 to 19	81% (73-87%)	19% (13-27%)	2.1 (1.4-2.9)	2.4% (1.6-3.4%)	8% (4.1-12%)	3.9 (2-6.2)	590 (380-840)
Austria	2010	Female	20 to 24	86% (77-92%)	14% (7.8-23%)	2.6 (1.7-3.7)	3.7% (2.7-5%)	8.9% (4.6-13%)	5.8 (3-8.8)	1100 (770-1600)
Austria	2010	Female	25 to 29	86% (77-93%)	14% (7.2-23%)	2.4 (1.6-3.2)	3.1% (2.3-4.1%)	8.8% (4.8-13%)	6.5 (3.6-9.3)	1100 (770-1500)
Austria	2010	Female	30 to 34	86% (76-92%)	14% (7.6-24%)	2.5 (1.7-3.4)	2.8% (2.1-3.8%)	10% (6.7-14%)	8.8 (5.7-12)	1000 (720-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	2010	Female	35 to 39	87% (79-93%)	13% (7.1-21%)	3 (2.1-4.2)	3.6% (2.6-4.7%)	13% (9.5-17%)	20 (14-25)	1600 (1200-2000)
Austria	2010	Female	40 to 44	86% (78-92%)	14% (8.1-22%)	3.4 (2.4-4.5)	4.4% (3.2-5.6%)	14% (11-17%)	40 (31-49)	2600 (2000-3200)
Austria	2010	Female	45 to 49	85% (79-91%)	15% (9.4-21%)	3.5 (2.5-4.7)	5.5% (4.1-7.1%)	14% (11-17%)	75 (58-91)	3800 (2900-4700)
Austria	2010	Female	50 to 54	87% (81-91%)	13% (8.9-19%)	3.8 (2.7-5)	6.3% (4.7-8%)	14% (11-17%)	100 (79-120)	4300 (3300-5400)
Austria	2010	Female	55 to 59	87% (83-91%)	13% (8.7-17%)	4.1 (3.1-5.4)	6.5% (4.8-8.3%)	12% (9.6-15%)	120 (91-140)	4300 (3300-5500)
Austria	2010	Female	60 to 64	87% (82-91%)	13% (8.6-18%)	4.1 (3-5.4)	6.3% (4.3-8.6%)	11% (7.7-14%)	150 (110-200)	4800 (3300-6400)
Austria	2010	Female	65 to 69	86% (81-91%)	14% (9.2-19%)	3.8 (2.9-5)	5.3% (3.3-7.4%)	8.4% (5.7-11%)	170 (120-240)	4600 (3000-6400)
Austria	2010	Female	70 to 74	84% (79-89%)	16% (11-21%)	3.5 (2.5-4.8)	4.3% (2.1-7.2%)	6.3% (3.1-10%)	170 (83-280)	3800 (1800-6100)
Austria	2010	Female	75 to 79	80% (72-87%)	20% (13-28%)	3.2 (2.2-4.5)	3.4% (0.65-6.5%)	4.4% (0.75-8.6%)	180 (30-340)	3100 (630-5900)
Austria	2010	Female	80 to 84	78% (67-86%)	22% (14-33%)	3 (2.1-4.1)	2.6% (-0.21-5.9%)	3% (-0.68-7.1%)	220 (-50-520)	2900 (-210-6200)
Austria	2010	Female	85 to 89	77% (67-85%)	23% (15-33%)	2.8 (1.9-3.9)	1.9% (-1.2-5.3%)	1.9% (-2.1-6.1%)	210 (-240-680)	2000 (-1300-5600)
Austria	2010	Female	90 to 94	76% (65-84%)	24% (16-35%)	2.7 (1.7-3.9)	1.3% (-2.2-5.6%)	1.1% (-3.2-6.4%)	63 (-180-360)	470 (-770-2000)
Austria	2010	Female	95 plus	76% (60-87%)	24% (13-40%)	2.6 (1.4-4.1)	1% (-3.2-6.2%)	0.78% (-4.4-7.1%)	26 (-140-230)	140 (-430-840)
Austria	2010	Male	All Ages	90% (88-92%)	10% (8.2-12%)	4.7 (4-5.5)	11% (9.3-14%)	13% (8.8-16%)	4600 (3200-5900)	130000 (100000-150000)
Austria	2010	Male	15 to 19	82% (76-87%)	18% (13-24%)	2.8 (1.9-3.7)	6.8% (4.4-9.3%)	13% (7-20%)	18 (9.4-26)	1900 (1200-2600)
Austria	2010	Male	20 to 24	89% (82-94%)	11% (5.9-18%)	4.2 (2.9-5.7)	13% (9.7-17%)	19% (11-27%)	40 (24-58)	4500 (3200-6000)
Austria	2010	Male	25 to 29	91% (85-95%)	8.8% (4.7-15%)	4.3 (3.1-5.8)	12% (8.8-15%)	19% (11-27%)	36 (21-51)	4200 (3100-5400)
Austria	2010	Male	30 to 34	93% (87-97%)	7.4% (3.4-13%)	4.6 (3.3-6.2)	11% (7.9-14%)	20% (13-28%)	41 (27-57)	3800 (2800-4900)
Austria	2010	Male	35 to 39	93% (88-97%)	6.7% (2.9-12%)	5 (3.5-6.7)	11% (8.2-14%)	22% (15-29%)	67 (45-88)	4900 (3600-6200)
Austria	2010	Male	40 to 44	94% (89-97%)	6.5% (3.2-11%)	5.7 (4.2-7.4)	12% (9.5-15%)	24% (18-30%)	130 (99-160)	7700 (6000-9400)
Austria	2010	Male	45 to 49	93% (89-96%)	6.8% (3.7-11%)	6.7 (5-8.7)	15% (12-18%)	25% (20-30%)	250 (200-300)	12000 (9900-15000)
Austria	2010	Male	50 to 54	94% (90-96%)	6.4% (3.8-10%)	7.2 (5.4-9.2)	17% (13-20%)	25% (20-29%)	360 (290-420)	15000 (12000-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	2010	Male	55 to 59	94% (92-96%)	5.6% (3.7-8%)	7.4 (5.6-9.3)	16% (13-19%)	22% (18-26%)	420 (340-480)	15000 (12000-17000)
Austria	2010	Male	60 to 64	95% (92-97%)	5.4% (3.4-8.2%)	7.1 (5.3-9.2)	15% (11-18%)	19% (15-23%)	530 (410-640)	16000 (12000-19000)
Austria	2010	Male	65 to 69	95% (92-96%)	5.5% (3.5-8.4%)	6.5 (5-8.1)	13% (9.4-16%)	16% (12-20%)	590 (440-730)	15000 (11000-18000)
Austria	2010	Male	70 to 74	94% (92-96%)	5.7% (3.8-8.5%)	5.9 (4.3-7.8)	10% (6.4-14%)	12% (7.7-17%)	540 (330-760)	11000 (7000-15000)
Austria	2010	Male	75 to 79	94% (90-96%)	6.1% (3.6-9.6%)	5.5 (3.9-7.4)	8.6% (4.4-13%)	10% (4.8-15%)	500 (240-770)	8000 (4100-12000)
Austria	2010	Male	80 to 84	93% (89-96%)	6.9% (4-11%)	5.3 (3.8-6.9)	7.3% (3-12%)	8.1% (2.7-13%)	510 (170-840)	6200 (2600-9700)
Austria	2010	Male	85 to 89	92% (88-95%)	7.8% (4.8-12%)	5.1 (3.6-6.8)	6.6% (2.1-11%)	7.1% (1.7-12%)	380 (91-660)	3200 (1000-5300)
Austria	2010	Male	90 to 94	92% (87-95%)	8.2% (4.8-13%)	5 (3.3-7.1)	6% (0.93-11%)	6.3% (0.34-12%)	130 (7-260)	760 (120-1500)
Austria	2010	Male	95 plus	91% (84-96%)	8.9% (4.1-16%)	4.8 (2.9-7.7)	5.7% (-0.42-12%)	5.9% (-1.2-14%)	50 (-11-120)	190 (-14-420)
Austria	2016	Female	All Ages	83% (80-86%)	17% (14-20%)	2.8 (2.1-3.7)	3.8% (2.1-5.9%)	3.7% (0.13-8%)	1600 (59-3500)	42000 (24000-65000)
Austria	2016	Female	15 to 19	82% (74-89%)	18% (11-26%)	2.1 (1.3-3.2)	2.3% (1.5-3.3%)	8% (4.1-13%)	3.1 (1.6-5)	490 (320-700)
Austria	2016	Female	20 to 24	85% (75-92%)	15% (8-25%)	2.5 (1.5-3.8)	3.5% (2.5-4.7%)	8.8% (4.3-14%)	5 (2.5-8)	1100 (730-1600)
Austria	2016	Female	25 to 29	86% (76-93%)	14% (6.9-24%)	2.3 (1.5-3.5)	3% (2.1-4%)	8.8% (4.4-13%)	5.9 (3-9.3)	1100 (720-1500)
Austria	2016	Female	30 to 34	86% (75-93%)	14% (7.1-25%)	2.5 (1.6-3.7)	2.8% (1.9-3.7%)	10% (6.6-15%)	10 (6.3-14)	1100 (800-1600)
Austria	2016	Female	35 to 39	86% (77-93%)	14% (7.1-23%)	3 (1.9-4.5)	3.3% (2.3-4.5%)	13% (9.2-17%)	16 (11-21)	1300 (950-1800)
Austria	2016	Female	40 to 44	86% (77-92%)	14% (8.1-23%)	3.3 (2.2-4.8)	4.1% (3-5.6%)	14% (11-18%)	32 (24-41)	2100 (1500-2700)
Austria	2016	Female	45 to 49	86% (79-91%)	14% (9-21%)	3.5 (2.4-5.1)	5.1% (3.7-6.9%)	14% (11-18%)	68 (52-87)	3500 (2600-4500)
Austria	2016	Female	50 to 54	86% (80-91%)	14% (9-20%)	3.7 (2.5-5.4)	5.8% (4.1-7.8%)	13% (10-17%)	110 (81-130)	4600 (3400-6000)
Austria	2016	Female	55 to 59	88% (84-92%)	12% (8-16%)	4.1 (2.8-5.9)	6.2% (4.3-8.4%)	12% (9.1-16%)	130 (97-170)	4900 (3400-6500)
Austria	2016	Female	60 to 64	88% (84-92%)	12% (7.6-16%)	4.1 (2.7-5.9)	6% (3.7-8.8%)	10% (7-14%)	140 (96-200)	4600 (2900-6600)
Austria	2016	Female	65 to 69	87% (82-92%)	13% (8.5-18%)	3.9 (2.7-5.4)	5.1% (2.9-7.8%)	8.2% (5.1-12%)	160 (100-230)	4200 (2500-6500)
Austria	2016	Female	70 to 74	85% (80-90%)	15% (10-20%)	3.6 (2.5-5)	4.2% (1.7-7.2%)	6.3% (2.7-11%)	210 (90-350)	4600 (1800-8100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	2016	Female	75 to 79	82% (74-88%)	18% (12-26%)	3.3 (2.1-4.8)	3.4% (0.47-7.1%)	4.4% (0.62-9.4%)	180 (24-380)	3100 (440-6500)
Austria	2016	Female	80 to 84	79% (68-86%)	21% (14-32%)	3 (1.9-4.3)	2.6% (-0.54-6.7%)	3% (-1-8.2%)	190 (-66-520)	2500 (-520-6400)
Austria	2016	Female	85 to 89	77% (66-85%)	23% (15-34%)	2.8 (1.8-4.1)	1.9% (-1.4-6%)	1.9% (-2.3-6.9%)	190 (-230-660)	1800 (-1400-5600)
Austria	2016	Female	90 to 94	76% (65-85%)	24% (15-35%)	2.7 (1.6-4.2)	1.4% (-2.3-6%)	1.2% (-3.4-6.8%)	120 (-360-710)	910 (-1600-4000)
Austria	2016	Female	95 plus	76% (61-87%)	24% (13-39%)	2.5 (1.3-4.2)	1.1% (-3.4-6.8%)	0.83% (-4.6-7.7%)	24 (-130-220)	130 (-400-780)
Austria	2016	Male	All Ages	89% (87-91%)	11% (8.7-13%)	4.7 (3.5-6.1)	11% (8-14%)	12% (7-16%)	4600 (2700-6300)	120000 (93000-160000)
Austria	2016	Male	15 to 19	81% (73-88%)	19% (12-27%)	2.7 (1.7-4.1)	6.2% (4-8.6%)	13% (6.4-20%)	13 (6.4-20)	1500 (940-2100)
Austria	2016	Male	20 to 24	87% (78-93%)	13% (6.8-22%)	4 (2.5-5.9)	12% (8.7-15%)	18% (10-27%)	31 (17-46)	4000 (2800-5300)
Austria	2016	Male	25 to 29	89% (82-95%)	11% (5.4-18%)	4.1 (2.7-6)	11% (7.8-14%)	18% (10-27%)	31 (17-46)	3800 (2700-5100)
Austria	2016	Male	30 to 34	91% (83-96%)	8.9% (4.2-17%)	4.4 (2.9-6.4)	9.6% (6.9-13%)	20% (12-28%)	39 (24-55)	3800 (2700-5000)
Austria	2016	Male	35 to 39	92% (85-96%)	7.9% (3.8-15%)	4.8 (3.1-7.1)	10% (7.1-13%)	22% (14-30%)	51 (34-70)	3900 (2800-5100)
Austria	2016	Male	40 to 44	92% (86-96%)	7.6% (3.7-14%)	5.5 (3.7-8)	12% (8.5-15%)	24% (17-31%)	93 (67-120)	5600 (4200-7200)
Austria	2016	Male	45 to 49	92% (87-96%)	7.6% (4.1-13%)	6.4 (4.5-9)	14% (11-18%)	25% (19-31%)	210 (160-270)	11000 (8200-13000)
Austria	2016	Male	50 to 54	93% (89-96%)	6.9% (4-11%)	7 (4.9-9.8)	16% (12-19%)	25% (19-30%)	370 (280-450)	16000 (12000-19000)
Austria	2016	Male	55 to 59	94% (91-96%)	5.9% (3.8-8.8%)	7.2 (5-9.8)	15% (12-19%)	22% (17-27%)	450 (340-540)	16000 (12000-19000)
Austria	2016	Male	60 to 64	95% (91-97%)	5.5% (3.4-8.6%)	7 (4.7-9.9)	14% (10-18%)	19% (14-23%)	470 (340-590)	14000 (11000-18000)
Austria	2016	Male	65 to 69	94% (91-97%)	5.5% (3.4-8.7%)	6.4 (4.5-8.7)	12% (8.5-16%)	16% (11-20%)	520 (360-670)	13000 (9200-17000)
Austria	2016	Male	70 to 74	94% (91-96%)	5.7% (3.7-8.8%)	5.8 (4-8.2)	9.8% (5.9-14%)	12% (7.1-17%)	610 (360-890)	13000 (7600-19000)
Austria	2016	Male	75 to 79	94% (90-96%)	6.2% (3.5-10%)	5.4 (3.6-7.8)	8% (3.8-12%)	9.3% (4.2-15%)	490 (220-790)	8200 (3900-13000)
Austria	2016	Male	80 to 84	93% (88-96%)	7.3% (3.9-12%)	5.1 (3.5-7.3)	7% (2.5-12%)	7.7% (2.4-14%)	480 (150-830)	5900 (2100-9700)
Austria	2016	Male	85 to 89	92% (86-95%)	8.5% (5-14%)	4.9 (3.2-7.3)	6.2% (1.3-11%)	6.6% (0.78-12%)	430 (51-810)	3700 (780-6600)
Austria	2016	Male	90 to 94	91% (86-95%)	8.9% (5-14%)	4.8 (2.9-7.5)	5.5% (0.26-11%)	5.8% (-0.4-12%)	230 (-15-470)	1300 (63-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Austria	2016	Male	95 plus	90% (83-96%)	9.5% (4.5-17%)	4.6 (2.6-7.8)	5.3% (-0.75-12%)	5.5% (-1.7-13%)	39 (-12-95)	150 (-22-340)
Belgium	1990	Female	All Ages	80% (77-83%)	20% (17-23%)	2 (1.9-2.2)	3.2% (1.8-4.6%)	2.9% (0.3-5.6%)	1500 (160-2900)	47000 (27000-67000)
Belgium	1990	Female	15 to 19	77% (68-84%)	23% (16-32%)	1.6 (1.2-2.1)	2.6% (1.5-4%)	8.3% (4-13%)	8.7 (4.2-14)	910 (540-1300)
Belgium	1990	Female	20 to 24	83% (73-91%)	17% (9.4-27%)	1.8 (1.3-2.6)	3.5% (2.2-4.9%)	8.7% (4.2-13%)	14 (7-22)	1600 (1000-2300)
Belgium	1990	Female	25 to 29	84% (73-92%)	16% (8.1-27%)	1.7 (1.2-2.3)	2.9% (1.8-4.1%)	8.3% (4.1-13%)	15 (7.3-23)	1600 (1000-2200)
Belgium	1990	Female	30 to 34	84% (73-92%)	16% (8.3-27%)	1.9 (1.4-2.5)	3.1% (2-4.4%)	9.4% (5.5-13%)	24 (14-35)	1900 (1200-2600)
Belgium	1990	Female	35 to 39	83% (75-91%)	17% (9.5-25%)	2.3 (1.7-2.9)	4.2% (2.8-5.7%)	12% (8.1-16%)	44 (29-58)	2700 (1900-3600)
Belgium	1990	Female	40 to 44	84% (75-90%)	16% (9.6-25%)	2.7 (2.1-3.3)	5.7% (4.1-7.4%)	14% (10-17%)	80 (58-99)	4200 (3100-5200)
Belgium	1990	Female	45 to 49	85% (79-90%)	15% (9.5-21%)	3.2 (2.5-4.1)	6.8% (5-8.7%)	15% (12-18%)	97 (75-120)	4400 (3400-5500)
Belgium	1990	Female	50 to 54	87% (81-92%)	13% (8.2-19%)	3.4 (2.6-4.3)	6.9% (5-8.8%)	14% (11-17%)	130 (100-160)	5200 (4000-6500)
Belgium	1990	Female	55 to 59	89% (84-93%)	11% (7.4-16%)	3.3 (2.6-4.1)	6.5% (4.7-8.2%)	12% (9-14%)	180 (140-220)	6100 (4600-7700)
Belgium	1990	Female	60 to 64	89% (83-92%)	11% (7.6-17%)	3.2 (2.4-4)	5.1% (3-7.2%)	8.2% (5.2-11%)	190 (120-270)	5600 (3400-8000)
Belgium	1990	Female	65 to 69	87% (81-91%)	13% (8.9-19%)	2.9 (2.3-3.5)	3.7% (1.6-5.9%)	5.6% (2.6-8.6%)	210 (98-320)	5000 (2100-7900)
Belgium	1990	Female	70 to 74	84% (77-88%)	16% (12-23%)	2.6 (1.9-3.3)	2.4% (-0.35-5%)	3.3% (-0.6-6.9%)	130 (-25-280)	2600 (-410-5400)
Belgium	1990	Female	75 to 79	80% (70-87%)	20% (13-30%)	2.4 (1.7-3.1)	1.3% (-1.6-4.2%)	1.6% (-2.2-5.3%)	130 (-180-440)	2100 (-2500-6600)
Belgium	1990	Female	80 to 84	76% (65-84%)	24% (16-35%)	2.3 (1.7-2.9)	1% (-1.6-3.6%)	1.1% (-2.2-4.3%)	120 (-240-460)	1500 (-2300-5300)
Belgium	1990	Female	85 to 89	73% (62-82%)	27% (18-38%)	2.2 (1.5-3)	1% (-1.4-3.5%)	1% (-2-4.1%)	100 (-210-420)	960 (-1400-3300)
Belgium	1990	Female	90 to 94	72% (60-82%)	28% (18-40%)	2.2 (1.4-3.1)	0.88% (-1.8-3.9%)	0.79% (-2.4-4.3%)	43 (-130-230)	300 (-590-1300)
Belgium	1990	Female	95 plus	71% (56-83%)	29% (17-44%)	2.2 (1.3-3.4)	0.95% (-2.3-4.7%)	0.81% (-3-5.3%)	14 (-50-89)	67 (-150-330)
Belgium	1990	Male	All Ages	88% (85-90%)	12% (9.8-15%)	3.5 (3.3-3.7)	7.9% (6.3-9.5%)	7.1% (4.7-9.4%)	3700 (2500-5000)	130000 (100000-160000)
Belgium	1990	Male	15 to 19	80% (71-86%)	20% (14-29%)	2.7 (2-3.4)	7.9% (4.7-11%)	14% (7.2-21%)	41 (21-60)	3500 (2100-5000)
Belgium	1990	Male	20 to 24	87% (78-93%)	13% (6.8-22%)	4 (3-5)	14% (9.4-19%)	21% (12-29%)	100 (61-140)	8500 (5600-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	1990	Male	25 to 29	90% (81-95%)	10% (5.2-19%)	3.2 (2.5-4)	11% (7.6-15%)	18% (10-25%)	89 (51-130)	7400 (4900-9900)
Belgium	1990	Male	30 to 34	91% (83-96%)	8.6% (4-17%)	3.3 (2.6-4.1)	10% (6.9-14%)	17% (9.9-24%)	93 (54-130)	6800 (4500-8900)
Belgium	1990	Male	35 to 39	93% (86-97%)	7.5% (3.1-14%)	4.2 (3.3-5.1)	12% (8.2-15%)	20% (14-26%)	130 (89-170)	8000 (5800-10000)
Belgium	1990	Male	40 to 44	93% (87-97%)	6.8% (3.1-13%)	4.9 (3.9-5.8)	12% (9.2-16%)	20% (15-25%)	190 (140-240)	10000 (7600-13000)
Belgium	1990	Male	45 to 49	94% (88-96%)	6.5% (3.6-12%)	5.3 (4.4-6.4)	13% (10-16%)	19% (15-24%)	220 (170-270)	10000 (7900-13000)
Belgium	1990	Male	50 to 54	94% (90-97%)	5.8% (3.3-9.8%)	5.7 (4.7-6.7)	12% (9.3-15%)	17% (13-21%)	290 (220-360)	12000 (9200-15000)
Belgium	1990	Male	55 to 59	95% (93-97%)	4.8% (2.9-7.5%)	5.9 (5-6.9)	12% (8.7-14%)	15% (11-18%)	440 (320-550)	15000 (12000-19000)
Belgium	1990	Male	60 to 64	95% (93-97%)	4.5% (2.5-7.4%)	5.7 (4.7-6.9)	9.6% (6.6-12%)	11% (7.6-15%)	540 (360-700)	16000 (11000-21000)
Belgium	1990	Male	65 to 69	95% (92-97%)	5% (2.9-7.9%)	5 (4.2-5.7)	7.1% (4.4-9.8%)	8.1% (4.7-11%)	560 (330-770)	14000 (8500-19000)
Belgium	1990	Male	70 to 74	94% (91-96%)	6% (3.8-9.3%)	4.2 (3.5-5.1)	4.9% (2-7.6%)	5.3% (2-8.5%)	320 (120-510)	6500 (2700-10000)
Belgium	1990	Male	75 to 79	92% (88-96%)	7.6% (4.3-12%)	3.7 (2.9-4.6)	3.5% (0.86-6.3%)	3.7% (0.54-6.8%)	330 (50-620)	5300 (1300-9600)
Belgium	1990	Male	80 to 84	91% (85-95%)	8.9% (5-15%)	3.5 (2.7-4.2)	2.9% (0.47-5.6%)	3% (0.21-6.1%)	250 (18-500)	3000 (500-5700)
Belgium	1990	Male	85 to 89	90% (84-94%)	9.6% (5.6-16%)	3.2 (2.5-4)	2.3% (-0.03-4.8%)	2.3% (-0.37-5.2%)	120 (-20-280)	1100 (-14-2300)
Belgium	1990	Male	90 to 94	91% (85-95%)	9.4% (5.4-15%)	3.1 (2.2-4)	1.6% (-0.87-4.4%)	1.5% (-1.5-4.6%)	31 (-31-96)	200 (-110-540)
Belgium	1990	Male	95 plus	91% (84-96%)	9% (4.2-16%)	2.9 (1.9-4.1)	1.2% (-1.6-4.7%)	0.98% (-2.3-5%)	4.9 (-11-25)	25 (-31-93)
Belgium	1995	Female	All Ages	81% (77-84%)	19% (16-23%)	1.9 (1.7-2.1)	3.1% (1.8-4.4%)	2.7% (0.38-5.1%)	1400 (200-2700)	44000 (27000-62000)
Belgium	1995	Female	15 to 19	77% (69-84%)	23% (16-31%)	1.5 (1.1-2)	2.6% (1.5-3.8%)	7.9% (3.9-12%)	8.1 (4-13)	840 (490-1200)
Belgium	1995	Female	20 to 24	84% (73-91%)	16% (9.2-27%)	1.7 (1.2-2.4)	3.3% (2.1-4.7%)	8.5% (4.2-13%)	11 (5.6-18)	1400 (860-2000)
Belgium	1995	Female	25 to 29	85% (73-92%)	15% (7.9-27%)	1.6 (1.1-2.2)	2.7% (1.7-3.8%)	8% (4-12%)	13 (6.2-20)	1400 (910-1900)
Belgium	1995	Female	30 to 34	84% (72-92%)	16% (8.2-28%)	1.7 (1.3-2.3)	2.8% (1.8-4%)	8.9% (5.1-13%)	22 (12-31)	1700 (1100-2400)
Belgium	1995	Female	35 to 39	83% (74-91%)	17% (9.1-26%)	2.1 (1.6-2.8)	3.9% (2.6-5.4%)	12% (7.8-16%)	42 (28-56)	2700 (1900-3500)
Belgium	1995	Female	40 to 44	84% (75-91%)	16% (9.5-25%)	2.5 (1.9-3.1)	5.4% (3.9-7.1%)	14% (10-17%)	77 (58-97)	4100 (3100-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	1995	Female	45 to 49	86% (78-91%)	14% (9.4-22%)	3 (2.3-3.8)	6.5% (4.8-8.3%)	15% (11-18%)	120 (89-140)	5300 (4100-6600)
Belgium	1995	Female	50 to 54	87% (81-92%)	13% (8.2-19%)	3.2 (2.5-4)	6.8% (5-8.5%)	14% (11-16%)	120 (96-150)	4900 (3800-6100)
Belgium	1995	Female	55 to 59	89% (84-92%)	11% (7.6-16%)	3.1 (2.5-3.9)	6.2% (4.5-8%)	11% (8.9-14%)	150 (120-190)	5300 (3900-6700)
Belgium	1995	Female	60 to 64	89% (83-92%)	11% (7.7-17%)	2.9 (2.3-3.7)	4.8% (2.9-6.7%)	8.2% (5.4-11%)	170 (110-230)	5100 (3100-7100)
Belgium	1995	Female	65 to 69	87% (81-91%)	13% (8.9-19%)	2.7 (2.1-3.3)	3.5% (1.5-5.3%)	5.5% (2.6-8%)	180 (86-270)	4400 (1900-6600)
Belgium	1995	Female	70 to 74	84% (77-88%)	16% (12-23%)	2.4 (1.8-3)	2.2% (-0.27-4.5%)	3.3% (-0.25-6.4%)	180 (-14-350)	3400 (-420-6800)
Belgium	1995	Female	75 to 79	80% (70-87%)	20% (13-30%)	2.2 (1.6-2.9)	1.2% (-1.3-3.6%)	1.6% (-1.6-4.6%)	91 (-90-260)	1400 (-1400-4100)
Belgium	1995	Female	80 to 84	76% (66-85%)	24% (15-34%)	2.1 (1.5-2.7)	0.83% (-1.5-3.1%)	0.94% (-1.9-3.8%)	99 (-200-400)	1200 (-2200-4500)
Belgium	1995	Female	85 to 89	73% (62-82%)	27% (18-38%)	2.1 (1.4-2.8)	0.73% (-1.5-3%)	0.73% (-2-3.4%)	81 (-220-370)	770 (-1600-3100)
Belgium	1995	Female	90 to 94	72% (60-82%)	28% (18-40%)	2.1 (1.3-2.9)	0.65% (-1.8-3.2%)	0.56% (-2.4-3.6%)	39 (-160-250)	280 (-780-1400)
Belgium	1995	Female	95 plus	71% (56-83%)	29% (17-44%)	2.1 (1.2-3.1)	0.71% (-2-3.8%)	0.57% (-2.6-4.1%)	13 (-58-95)	67 (-190-370)
Belgium	1995	Male	All Ages	88% (85-90%)	12% (9.5-15%)	3.3 (3-3.6)	7.6% (6-9.3%)	6.5% (4.5-8.8%)	3500 (2300-4700)	120000 (96000-150000)
Belgium	1995	Male	15 to 19	80% (71-86%)	20% (14-29%)	2.4 (1.8-3.1)	7.3% (4.2-10%)	13% (6.7-20%)	32 (16-47)	2800 (1700-4100)
Belgium	1995	Male	20 to 24	87% (78-93%)	13% (7-22%)	3.6 (2.8-4.7)	13% (9.1-18%)	20% (11-27%)	84 (50-120)	7300 (4900-9800)
Belgium	1995	Male	25 to 29	90% (81-95%)	10% (5-19%)	3 (2.3-3.8)	11% (7.1-15%)	17% (9-24%)	77 (40-110)	6500 (4300-8800)
Belgium	1995	Male	30 to 34	92% (83-96%)	8.5% (3.8-17%)	3 (2.3-3.9)	9.5% (6.2-13%)	16% (8.8-23%)	87 (47-120)	6500 (4200-8800)
Belgium	1995	Male	35 to 39	93% (86-97%)	7.5% (3.3-14%)	3.9 (3-4.8)	11% (7.8-14%)	19% (13-25%)	130 (90-180)	8400 (6100-11000)
Belgium	1995	Male	40 to 44	93% (87-97%)	6.8% (3.1-13%)	4.5 (3.6-5.5)	12% (9.3-15%)	20% (15-25%)	190 (140-250)	10000 (8000-13000)
Belgium	1995	Male	45 to 49	94% (88-96%)	6.5% (3.5-12%)	4.9 (4.1-5.9)	13% (9.9-16%)	19% (15-23%)	280 (210-340)	13000 (10000-16000)
Belgium	1995	Male	50 to 54	94% (90-97%)	5.8% (3.2-10%)	5.3 (4.3-6.2)	12% (9.6-15%)	17% (13-21%)	290 (230-360)	12000 (9400-15000)
Belgium	1995	Male	55 to 59	95% (93-97%)	4.8% (2.8-7.4%)	5.5 (4.6-6.4)	11% (8.4-14%)	14% (11-18%)	370 (280-470)	13000 (10000-16000)
Belgium	1995	Male	60 to 64	96% (93-98%)	4.5% (2.5-7.3%)	5.3 (4.3-6.4)	8.9% (6.2-12%)	11% (7.3-14%)	450 (300-600)	14000 (9400-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	1995	Male	65 to 69	95% (92-97%)	4.9% (2.8-8.3%)	4.6 (3.8-5.4)	6.5% (3.9-8.8%)	7.5% (4.3-10%)	460 (270-640)	12000 (7000-16000)
Belgium	1995	Male	70 to 74	94% (91-96%)	5.9% (3.7-9.2%)	3.9 (3.2-4.8)	4.4% (2-6.9%)	4.8% (2-7.8%)	410 (170-670)	8400 (3900-14000)
Belgium	1995	Male	75 to 79	93% (88-96%)	7.5% (4.2-12%)	3.5 (2.7-4.3)	3.1% (0.86-5.7%)	3.3% (0.63-6.2%)	220 (42-430)	3600 (1000-6600)
Belgium	1995	Male	80 to 84	91% (85-95%)	8.8% (4.9-15%)	3.2 (2.6-3.9)	2.5% (0.4-5%)	2.5% (0.15-5.4%)	220 (13-460)	2600 (420-5200)
Belgium	1995	Male	85 to 89	90% (84-94%)	9.5% (5.6-16%)	3 (2.3-3.8)	2% (-0.046-4.4%)	2% (-0.42-4.8%)	110 (-25-280)	1000 (-23-2300)
Belgium	1995	Male	90 to 94	91% (85-95%)	9.4% (5.3-15%)	2.9 (2-3.7)	1.3% (-0.89-3.9%)	1.2% (-1.5-4.1%)	29 (-36-100)	200 (-140-580)
Belgium	1995	Male	95 plus	91% (83-96%)	9.1% (4.2-17%)	2.7 (1.7-3.9)	1% (-1.4-4%)	0.72% (-2.1-4.2%)	4.4 (-13-26)	25 (-34-100)
Belgium	2000	Female	All Ages	80% (77-83%)	20% (17-23%)	1.9 (1.7-2.2)	3.1% (1.9-4.4%)	2.8% (0.6-5%)	1500 (320-2700)	45000 (28000-62000)
Belgium	2000	Female	15 to 19	74% (65-82%)	26% (18-35%)	1.4 (0.98-1.9)	2.4% (1.4-3.5%)	7.8% (4-12%)	7 (3.5-11)	750 (450-1100)
Belgium	2000	Female	20 to 24	81% (69-90%)	19% (10-31%)	1.6 (1.1-2.3)	3.2% (2.1-4.5%)	8.4% (4.1-13%)	10 (4.9-16)	1300 (780-1800)
Belgium	2000	Female	25 to 29	84% (71-92%)	16% (8.3-29%)	1.6 (1.1-2.2)	2.6% (1.6-3.6%)	8.3% (4.1-13%)	11 (5.4-17)	1200 (790-1700)
Belgium	2000	Female	30 to 34	84% (73-92%)	16% (8.3-27%)	1.7 (1.2-2.3)	2.7% (1.8-3.7%)	9.1% (5.3-13%)	18 (11-26)	1500 (1000-2100)
Belgium	2000	Female	35 to 39	84% (76-91%)	16% (8.9-24%)	2.1 (1.6-2.8)	3.7% (2.5-5.2%)	12% (7.8-16%)	40 (26-53)	2600 (1800-3400)
Belgium	2000	Female	40 to 44	84% (76-91%)	16% (9.2-24%)	2.5 (2-3.2)	5% (3.5-6.6%)	14% (10-17%)	74 (54-92)	3900 (2900-5000)
Belgium	2000	Female	45 to 49	86% (79-91%)	14% (9.5-21%)	3 (2.3-3.8)	6.4% (4.7-8.1%)	15% (12-18%)	120 (91-140)	5300 (4100-6500)
Belgium	2000	Female	50 to 54	88% (82-92%)	12% (8.2-18%)	3.2 (2.5-4.1)	6.9% (5.2-8.8%)	14% (11-17%)	160 (120-190)	6200 (4800-7600)
Belgium	2000	Female	55 to 59	89% (85-93%)	11% (7.5-15%)	3.1 (2.5-3.9)	6.3% (4.7-8%)	12% (9.3-14%)	150 (120-180)	5100 (3900-6400)
Belgium	2000	Female	60 to 64	88% (84-92%)	12% (7.9-16%)	2.9 (2.2-3.7)	5% (3.3-6.8%)	8.7% (6.2-11%)	160 (110-210)	4700 (3100-6300)
Belgium	2000	Female	65 to 69	86% (81-91%)	14% (9.4-19%)	2.6 (2.1-3.3)	3.6% (1.9-5.3%)	5.9% (3.4-8.3%)	170 (100-250)	4100 (2200-6100)
Belgium	2000	Female	70 to 74	83% (76-88%)	17% (12-24%)	2.3 (1.7-3)	2.4% (0.29-4.6%)	3.6% (0.56-6.6%)	180 (26-320)	3300 (400-6300)
Belgium	2000	Female	75 to 79	78% (70-86%)	22% (14-30%)	2.1 (1.5-2.8)	1.4% (-0.69-3.6%)	1.9% (-0.88-4.7%)	150 (-70-370)	2300 (-1100-5800)
Belgium	2000	Female	80 to 84	75% (65-84%)	25% (16-35%)	2 (1.5-2.8)	0.94% (-1.3-3.1%)	1.1% (-1.7-3.8%)	88 (-130-300)	1000 (-1400-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	2000	Female	85 to 89	73% (62-81%)	27% (19-38%)	2 (1.4-2.8)	0.77% (-1.4-3%)	0.8% (-1.8-3.5%)	91 (-210-390)	840 (-1500-3200)
Belgium	2000	Female	90 to 94	71% (59-81%)	29% (19-41%)	2 (1.3-2.9)	0.64% (-1.6-3.1%)	0.57% (-2.1-3.6%)	46 (-170-280)	330 (-840-1600)
Belgium	2000	Female	95 plus	71% (55-82%)	29% (18-45%)	2 (1.1-3.1)	0.71% (-1.8-3.8%)	0.6% (-2.4-4.2%)	19 (-75-140)	95 (-240-500)
Belgium	2000	Male	All Ages	89% (87-91%)	11% (8.8-13%)	3.3 (2.9-3.7)	7.8% (6-9.6%)	6.8% (4.4-9.2%)	3500 (2400-4800)	120000 (96000-150000)
Belgium	2000	Male	15 to 19	78% (70-85%)	22% (15-30%)	2.4 (1.7-3)	7% (4.2-10%)	13% (6.6-20%)	29 (15-44)	2700 (1600-3800)
Belgium	2000	Male	20 to 24	88% (80-93%)	12% (6.6-20%)	3.6 (2.7-4.7)	13% (8.9-18%)	20% (12-27%)	76 (44-110)	6600 (4400-8900)
Belgium	2000	Male	25 to 29	91% (84-95%)	9.1% (4.8-16%)	3 (2.3-3.8)	11% (7-14%)	17% (9.3-24%)	68 (37-97)	5900 (3900-7900)
Belgium	2000	Male	30 to 34	92% (86-96%)	7.6% (3.6-14%)	3 (2.3-3.9)	9.5% (6.2-13%)	17% (9.1-24%)	78 (43-110)	5900 (3800-7900)
Belgium	2000	Male	35 to 39	93% (88-97%)	6.6% (2.9-12%)	3.9 (2.9-4.9)	11% (7.8-14%)	20% (13-26%)	130 (83-170)	8200 (5800-11000)
Belgium	2000	Male	40 to 44	94% (89-97%)	5.9% (2.8-11%)	4.5 (3.6-5.5)	13% (9.4-16%)	21% (16-27%)	200 (150-250)	11000 (8200-13000)
Belgium	2000	Male	45 to 49	94% (91-97%)	5.6% (3.1-9.4%)	4.9 (4.1-6)	13% (10-17%)	20% (16-25%)	290 (230-360)	14000 (11000-17000)
Belgium	2000	Male	50 to 54	95% (92-97%)	4.9% (2.9-7.9%)	5.3 (4.3-6.4)	13% (9.8-16%)	18% (14-22%)	380 (290-470)	15000 (12000-19000)
Belgium	2000	Male	55 to 59	96% (94-97%)	4% (2.6-5.9%)	5.5 (4.5-6.6)	12% (9.1-15%)	16% (12-19%)	380 (290-480)	13000 (10000-17000)
Belgium	2000	Male	60 to 64	96% (94-98%)	3.9% (2.3-6%)	5.3 (4.3-6.5)	9.5% (6.7-12%)	12% (8.2-16%)	420 (290-550)	13000 (8900-16000)
Belgium	2000	Male	65 to 69	96% (93-97%)	4.4% (2.6-6.7%)	4.6 (3.8-5.5)	6.8% (4.3-9.3%)	8.1% (4.9-11%)	450 (270-630)	11000 (6900-16000)
Belgium	2000	Male	70 to 74	95% (92-97%)	5.2% (3.5-7.6%)	4 (3.2-4.9)	4.6% (2.2-7.1%)	5.1% (2.4-8.1%)	390 (180-630)	8100 (3900-13000)
Belgium	2000	Male	75 to 79	93% (90-96%)	6.6% (4-10%)	3.5 (2.8-4.4)	3.2% (0.88-5.9%)	3.4% (0.77-6.4%)	330 (73-620)	5300 (1500-9500)
Belgium	2000	Male	80 to 84	92% (87-95%)	8% (4.6-13%)	3.2 (2.6-4)	2.5% (0.28-5%)	2.5% (-0.042-5.4%)	170 (-2.7-360)	2100 (240-4200)
Belgium	2000	Male	85 to 89	91% (86-95%)	8.6% (5.1-14%)	3 (2.3-3.8)	1.9% (-0.29-4.4%)	1.9% (-0.69-4.7%)	120 (-43-290)	1100 (-170-2400)
Belgium	2000	Male	90 to 94	92% (86-95%)	8.5% (5.1-14%)	2.9 (2-3.8)	1.4% (-0.89-4%)	1.2% (-1.4-4.2%)	33 (-40-120)	230 (-150-660)
Belgium	2000	Male	95 plus	91% (85-96%)	8.5% (4-15%)	2.7 (1.7-3.9)	1.1% (-1.5-4.1%)	0.76% (-2.2-4.2%)	5.5 (-16-31)	31 (-43-120)
Belgium	2005	Female	All Ages	81% (78-83%)	19% (17-22%)	2.2 (1.9-2.4)	3.5% (2.4-4.7%)	3.7% (1.7-5.8%)	1900 (850-3000)	50000 (34000-66000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	2005	Female	15 to 19	77% (70-83%)	23% (17-30%)	1.6 (1.2-2.1)	2.2% (1.3-3.1%)	8.2% (4.2-12%)	5.6 (2.9-8.7)	660 (410-940)
Belgium	2005	Female	20 to 24	82% (71-90%)	18% (10-29%)	1.9 (1.3-2.6)	2.8% (1.9-3.8%)	8.9% (4.4-14%)	7.2 (3.6-11)	1100 (700-1500)
Belgium	2005	Female	25 to 29	84% (73-91%)	16% (8.6-27%)	1.7 (1.2-2.4)	2.5% (1.6-3.4%)	8.6% (4.4-13%)	9.4 (4.7-14)	1100 (730-1500)
Belgium	2005	Female	30 to 34	84% (75-91%)	16% (8.7-25%)	1.9 (1.4-2.5)	2.6% (1.7-3.6%)	9.7% (5.9-14%)	16 (9.7-22)	1400 (930-1900)
Belgium	2005	Female	35 to 39	84% (76-91%)	16% (9.4-24%)	2.3 (1.7-3.1)	3.5% (2.4-4.8%)	12% (8.1-16%)	33 (22-45)	2200 (1600-3000)
Belgium	2005	Female	40 to 44	84% (77-91%)	16% (9.4-23%)	2.8 (2.2-3.5)	4.8% (3.5-6.3%)	14% (11-18%)	68 (51-85)	3700 (2800-4700)
Belgium	2005	Female	45 to 49	86% (81-90%)	14% (9.5-19%)	3.3 (2.6-4.2)	6.2% (4.7-7.8%)	15% (12-18%)	110 (90-140)	5400 (4100-6700)
Belgium	2005	Female	50 to 54	88% (84-92%)	12% (7.9-16%)	3.6 (2.8-4.5)	7.1% (5.3-9%)	15% (12-17%)	160 (130-200)	6700 (5300-8100)
Belgium	2005	Female	55 to 59	90% (87-93%)	10% (7.3-13%)	3.5 (2.8-4.3)	6.6% (5-8.4%)	13% (10-15%)	190 (150-230)	6600 (5100-8100)
Belgium	2005	Female	60 to 64	89% (85-92%)	11% (7.8-15%)	3.2 (2.5-4.1)	5.6% (4-7.5%)	9.9% (7.6-12%)	170 (130-210)	4900 (3600-6400)
Belgium	2005	Female	65 to 69	87% (83-91%)	13% (9.1-17%)	3 (2.4-3.6)	4.3% (2.7-5.9%)	7.3% (5-9.4%)	180 (120-230)	4400 (2900-6000)
Belgium	2005	Female	70 to 74	84% (79-88%)	16% (12-21%)	2.6 (2-3.4)	3.2% (1.3-5.1%)	4.9% (2.3-7.5%)	210 (96-320)	4000 (1700-6500)
Belgium	2005	Female	75 to 79	78% (70-85%)	22% (15-30%)	2.3 (1.7-3.1)	2.2% (0.21-4.3%)	3% (0.47-5.7%)	210 (33-400)	3300 (310-6400)
Belgium	2005	Female	80 to 84	74% (64-83%)	26% (17-36%)	2.2 (1.6-2.9)	1.7% (-0.18-3.7%)	2.1% (-0.36-4.7%)	230 (-40-500)	2700 (-290-5800)
Belgium	2005	Female	85 to 89	72% (62-81%)	28% (19-38%)	2.2 (1.5-2.9)	1.5% (-0.54-3.6%)	1.7% (-0.74-4.3%)	150 (-64-360)	1300 (-450-3000)
Belgium	2005	Female	90 to 94	71% (59-80%)	29% (20-41%)	2.2 (1.4-3)	1.3% (-0.95-3.7%)	1.3% (-1.4-4.3%)	110 (-110-360)	680 (-510-2000)
Belgium	2005	Female	95 plus	70% (55-82%)	30% (18-45%)	2.2 (1.2-3.3)	1.3% (-1.3-4.4%)	1.3% (-1.8-5.1%)	49 (-67-190)	210 (-200-710)
Belgium	2005	Male	All Ages	90% (87-92%)	10% (8.4-13%)	3.7 (3.3-4.1)	8.6% (7.1-10%)	8% (5.8-10%)	4100 (3000-5300)	130000 (110000-160000)
Belgium	2005	Male	15 to 19	80% (74-85%)	20% (15-26%)	2.6 (2-3.4)	6.5% (4.1-9%)	14% (7.3-20%)	24 (12-35)	2300 (1400-3100)
Belgium	2005	Male	20 to 24	89% (83-94%)	11% (6.1-17%)	4.1 (3.1-5.2)	13% (9-17%)	21% (13-29%)	62 (38-87)	5700 (3900-7400)
Belgium	2005	Male	25 to 29	91% (86-95%)	8.7% (4.7-14%)	3.3 (2.5-4.2)	11% (7.2-14%)	18% (10-26%)	58 (33-82)	5100 (3400-6700)
Belgium	2005	Male	30 to 34	92% (86-96%)	7.6% (3.6-14%)	3.3 (2.6-4.2)	9.3% (6.3-12%)	18% (10-25%)	64 (36-91)	5100 (3400-6600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	2005	Male	35 to 39	94% (88-97%)	6.5% (2.9-12%)	4.3 (3.3-5.3)	11% (7.7-14%)	21% (14-28%)	110 (72-140)	7000 (5000-9000)
Belgium	2005	Male	40 to 44	94% (90-97%)	5.7% (2.7-10%)	5 (4-6)	13% (9.5-16%)	23% (17-28%)	190 (150-240)	11000 (8300-13000)
Belgium	2005	Male	45 to 49	95% (91-97%)	5.4% (3.1-8.9%)	5.4 (4.5-6.6)	14% (11-17%)	23% (18-27%)	300 (240-370)	14000 (12000-17000)
Belgium	2005	Male	50 to 54	95% (93-97%)	4.7% (2.8-7.3%)	5.8 (4.7-6.9)	14% (12-17%)	21% (17-25%)	400 (330-480)	17000 (14000-20000)
Belgium	2005	Male	55 to 59	96% (95-98%)	3.8% (2.4-5.3%)	6.1 (5.1-7.3)	13% (11-16%)	18% (15-22%)	520 (420-620)	18000 (15000-22000)
Belgium	2005	Male	60 to 64	96% (95-98%)	3.6% (2.2-5.5%)	5.9 (4.8-7.2)	11% (8.5-14%)	14% (11-18%)	460 (340-560)	14000 (11000-17000)
Belgium	2005	Male	65 to 69	96% (94-97%)	4.2% (2.5-6.3%)	5.1 (4.3-6)	8.4% (6.1-11%)	10% (7.5-13%)	460 (330-590)	12000 (8600-15000)
Belgium	2005	Male	70 to 74	95% (93-96%)	5.1% (3.5-7.3%)	4.4 (3.6-5.4)	5.9% (3.5-8.3%)	6.9% (4-9.8%)	460 (270-650)	9500 (5700-13000)
Belgium	2005	Male	75 to 79	93% (90-96%)	6.6% (4.2-9.9%)	3.8 (3.1-4.8)	4.2% (1.9-6.8%)	4.7% (1.9-7.7%)	410 (160-670)	6700 (3000-11000)
Belgium	2005	Male	80 to 84	92% (88-95%)	7.5% (4.5-12%)	3.6 (2.9-4.4)	3.4% (1.2-5.9%)	3.6% (0.99-6.5%)	350 (96-630)	4200 (1500-7300)
Belgium	2005	Male	85 to 89	92% (89-95%)	7.7% (4.8-11%)	3.5 (2.7-4.3)	2.7% (0.54-5.3%)	2.8% (0.18-5.8%)	140 (9.5-300)	1300 (260-2500)
Belgium	2005	Male	90 to 94	92% (88-95%)	7.7% (4.7-12%)	3.3 (2.4-4.2)	2.1% (-0.3-4.9%)	2.1% (-0.77-5.2%)	65 (-24-170)	420 (-60-960)
Belgium	2005	Male	95 plus	92% (86-96%)	8% (3.7-14%)	3.1 (2-4.2)	1.9% (-0.96-5.2%)	1.7% (-1.6-5.5%)	13 (-13-44)	61 (-32-170)
Belgium	2010	Female	All Ages	82% (79-84%)	18% (16-21%)	2.3 (2.1-2.6)	3.9% (2.8-5.1%)	4.3% (2.4-6.5%)	2300 (1300-3500)	57000 (41000-73000)
Belgium	2010	Female	15 to 19	80% (74-85%)	20% (15-26%)	1.8 (1.3-2.4)	2.1% (1.3-3.1%)	8.6% (4.5-13%)	5.6 (2.9-8.6)	670 (430-990)
Belgium	2010	Female	20 to 24	85% (76-91%)	15% (8.9-24%)	2 (1.4-2.9)	3% (2-4.1%)	9.4% (4.7-14%)	8.4 (4.2-13)	1200 (740-1700)
Belgium	2010	Female	25 to 29	84% (74-92%)	16% (8.3-26%)	1.9 (1.3-2.6)	2.6% (1.8-3.7%)	9% (4.6-14%)	11 (5.5-17)	1200 (800-1700)
Belgium	2010	Female	30 to 34	85% (75-92%)	15% (8.2-25%)	2 (1.5-2.7)	2.7% (1.8-3.7%)	10% (6.2-14%)	15 (9.5-22)	1400 (910-1900)
Belgium	2010	Female	35 to 39	84% (75-91%)	16% (9-25%)	2.4 (1.8-3.2)	3.5% (2.4-4.8%)	13% (8.8-17%)	32 (22-41)	2100 (1500-2800)
Belgium	2010	Female	40 to 44	84% (77-91%)	16% (9.4-23%)	3 (2.2-3.7)	4.7% (3.4-6.1%)	15% (11-18%)	63 (47-77)	3500 (2600-4300)
Belgium	2010	Female	45 to 49	87% (82-91%)	13% (9.1-18%)	3.6 (2.8-4.5)	6.2% (4.6-8%)	16% (12-19%)	120 (92-140)	5500 (4300-6900)
Belgium	2010	Female	50 to 54	89% (84-92%)	11% (7.7-16%)	3.8 (3-4.7)	7.1% (5.4-9%)	15% (12-18%)	170 (140-210)	7100 (5600-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	2010	Female	55 to 59	90% (87-93%)	9.7% (7-13%)	3.7 (3-4.6)	6.9% (5.3-8.7%)	13% (11-16%)	200 (170-240)	7200 (5700-8800)
Belgium	2010	Female	60 to 64	90% (86-93%)	10% (7.5-14%)	3.5 (2.7-4.5)	6% (4.3-7.8%)	11% (8.1-13%)	220 (170-280)	6700 (4900-8700)
Belgium	2010	Female	65 to 69	88% (84-91%)	12% (8.8-16%)	3.2 (2.6-4)	4.8% (3.4-6.4%)	8.1% (6-10%)	190 (140-240)	4800 (3300-6300)
Belgium	2010	Female	70 to 74	85% (81-89%)	15% (11-19%)	2.9 (2.2-3.7)	3.6% (1.9-5.5%)	5.8% (3.4-8.5%)	210 (120-310)	4200 (2200-6300)
Belgium	2010	Female	75 to 79	80% (73-86%)	20% (14-27%)	2.6 (1.9-3.4)	2.8% (0.89-4.9%)	4% (1.5-6.8%)	250 (94-420)	3900 (1300-6900)
Belgium	2010	Female	80 to 84	77% (67-84%)	23% (16-33%)	2.5 (1.9-3.2)	2.3% (0.53-4.2%)	3% (0.66-5.4%)	300 (66-560)	3500 (780-6500)
Belgium	2010	Female	85 to 89	74% (64-81%)	26% (19-36%)	2.4 (1.7-3.2)	2% (0.12-4.2%)	2.4% (-0.0015-5%)	300 (-0.2-650)	2600 (160-5300)
Belgium	2010	Female	90 to 94	72% (61-81%)	28% (19-39%)	2.4 (1.6-3.3)	1.7% (-0.46-4.4%)	1.9% (-0.66-5.1%)	130 (-46-360)	790 (-200-2100)
Belgium	2010	Female	95 plus	71% (56-83%)	29% (17-44%)	2.4 (1.4-3.6)	1.8% (-0.92-5%)	1.9% (-1.3-5.8%)	81 (-57-250)	320 (-170-910)
Belgium	2010	Male	All Ages	90% (88-92%)	10% (8.2-12%)	4 (3.5-4.5)	9.1% (7.5-11%)	8.7% (6.5-11%)	4600 (3400-5900)	140000 (120000-170000)
Belgium	2010	Male	15 to 19	82% (76-87%)	18% (13-24%)	2.9 (2.1-3.7)	6.2% (3.9-8.6%)	15% (8.1-21%)	20 (11-29)	2000 (1300-2800)
Belgium	2010	Male	20 to 24	90% (84-94%)	10% (5.7-16%)	4.3 (3.3-5.5)	13% (9.1-17%)	22% (14-30%)	57 (35-77)	5400 (3800-7100)
Belgium	2010	Male	25 to 29	91% (85-95%)	8.7% (4.6-15%)	3.5 (2.6-4.5)	11% (7.2-14%)	19% (10-27%)	56 (30-79)	5000 (3400-6700)
Belgium	2010	Male	30 to 34	92% (86-96%)	7.8% (3.9-14%)	3.5 (2.6-4.5)	9.4% (6.2-13%)	19% (11-26%)	60 (33-84)	4800 (3200-6400)
Belgium	2010	Male	35 to 39	93% (89-97%)	6.5% (3-11%)	4.5 (3.3-5.6)	11% (7.7-14%)	22% (15-29%)	100 (68-130)	6700 (4800-8600)
Belgium	2010	Male	40 to 44	94% (90-97%)	5.6% (2.6-10%)	5.2 (4.1-6.4)	12% (9.5-16%)	24% (18-30%)	170 (130-210)	9600 (7400-12000)
Belgium	2010	Male	45 to 49	95% (91-97%)	5.4% (3.1-8.9%)	5.7 (4.7-6.9)	14% (11-17%)	24% (19-29%)	290 (230-350)	14000 (11000-17000)
Belgium	2010	Male	50 to 54	95% (93-97%)	4.6% (2.8-7.3%)	6.1 (5-7.3)	15% (12-18%)	23% (19-26%)	430 (360-510)	18000 (15000-21000)
Belgium	2010	Male	55 to 59	96% (95-98%)	3.7% (2.4-5.3%)	6.5 (5.4-7.7)	14% (12-17%)	20% (17-23%)	550 (450-650)	19000 (16000-23000)
Belgium	2010	Male	60 to 64	97% (95-98%)	3.5% (2.1-5.4%)	6.2 (5.1-7.6)	12% (9.7-15%)	16% (13-19%)	620 (490-750)	19000 (15000-23000)
Belgium	2010	Male	65 to 69	96% (94-97%)	4% (2.5-6%)	5.4 (4.5-6.4)	9.4% (7.1-12%)	12% (9-15%)	480 (370-600)	12000 (9500-15000)
Belgium	2010	Male	70 to 74	95% (93-97%)	5% (3.4-6.9%)	4.6 (3.8-5.5)	6.7% (4.5-9.1%)	8.1% (5.4-11%)	470 (310-640)	9700 (6500-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	2010	Male	75 to 79	94% (90-96%)	6.4% (4-9.6%)	4.1 (3.2-5)	4.9% (2.7-7.3%)	5.6% (3-8.5%)	460 (240-700)	7500 (4000-11000)
Belgium	2010	Male	80 to 84	93% (89-96%)	7.1% (4.5-11%)	3.8 (3.1-4.6)	4% (1.8-6.4%)	4.4% (1.8-7.3%)	420 (170-690)	5000 (2300-8100)
Belgium	2010	Male	85 to 89	93% (89-95%)	7.3% (4.7-11%)	3.7 (2.8-4.5)	3.3% (1-5.8%)	3.4% (0.78-6.5%)	290 (66-540)	2500 (800-4500)
Belgium	2010	Male	90 to 94	93% (89-95%)	7.4% (4.6-11%)	3.5 (2.6-4.5)	2.6% (0.045-5.5%)	2.6% (-0.4-6%)	78 (-12-180)	480 (8.3-1000)
Belgium	2010	Male	95 plus	92% (87-96%)	7.7% (3.7-13%)	3.3 (2-4.5)	2.3% (-0.65-6%)	2.2% (-1.2-6.4%)	22 (-12-67)	95 (-27-250)
Belgium	2016	Female	All Ages	82% (79-85%)	18% (15-21%)	2.4 (1.8-3)	3.8% (2.4-5.4%)	4.3% (1.8-7.6%)	2400 (1000-4400)	57000 (37000-82000)
Belgium	2016	Female	15 to 19	78% (69-84%)	22% (16-31%)	1.7 (1.1-2.4)	1.9% (1.2-2.7%)	8.3% (4.4-13%)	4.4 (2.3-7)	580 (370-850)
Belgium	2016	Female	20 to 24	84% (74-91%)	16% (9.2-26%)	2 (1.3-2.9)	2.8% (1.9-3.8%)	9.3% (4.7-15%)	7 (3.5-11)	1100 (710-1500)
Belgium	2016	Female	25 to 29	85% (74-92%)	15% (7.9-26%)	1.9 (1.3-2.7)	2.5% (1.6-3.4%)	9.2% (4.7-14%)	9 (4.6-14)	1200 (750-1600)
Belgium	2016	Female	30 to 34	85% (73-92%)	15% (8.1-27%)	2 (1.3-2.9)	2.5% (1.6-3.5%)	10% (6.4-15%)	14 (8.6-21)	1300 (860-1800)
Belgium	2016	Female	35 to 39	84% (75-91%)	16% (9-25%)	2.5 (1.6-3.5)	3.4% (2.3-4.6%)	13% (9-17%)	29 (19-39)	2000 (1400-2600)
Belgium	2016	Female	40 to 44	84% (76-91%)	16% (9.1-24%)	3 (2.1-4.1)	4.5% (3.1-6.1%)	15% (11-19%)	54 (39-72)	3100 (2200-4100)
Belgium	2016	Female	45 to 49	87% (81-91%)	13% (9-19%)	3.6 (2.5-4.9)	5.9% (4.2-7.8%)	16% (12-20%)	100 (74-130)	4900 (3600-6400)
Belgium	2016	Female	50 to 54	89% (84-92%)	11% (7.7-16%)	3.8 (2.7-5.1)	6.7% (4.8-8.7%)	15% (12-19%)	160 (120-210)	6700 (5000-8700)
Belgium	2016	Female	55 to 59	90% (87-93%)	9.7% (7-13%)	3.7 (2.6-5)	6.8% (4.9-8.8%)	13% (10-16%)	210 (160-270)	7500 (5500-9700)
Belgium	2016	Female	60 to 64	90% (87-93%)	9.8% (6.9-13%)	3.6 (2.5-4.8)	6% (3.9-8.2%)	11% (7.8-14%)	230 (160-300)	6800 (4500-9400)
Belgium	2016	Female	65 to 69	89% (85-92%)	11% (8.2-15%)	3.3 (2.4-4.4)	4.8% (3-7%)	8.2% (5.8-11%)	240 (160-330)	5900 (3700-8500)
Belgium	2016	Female	70 to 74	86% (82-90%)	14% (10-18%)	2.9 (2.1-4.1)	3.7% (1.8-6.1%)	6% (3.4-9.2%)	200 (110-310)	4000 (1900-6600)
Belgium	2016	Female	75 to 79	83% (76-88%)	17% (12-24%)	2.7 (1.8-3.9)	2.9% (0.8-5.5%)	4.2% (1.6-7.7%)	230 (79-420)	3600 (1000-7000)
Belgium	2016	Female	80 to 84	79% (70-86%)	21% (14-30%)	2.6 (1.8-3.7)	2.4% (0.29-5.3%)	3.1% (0.45-6.9%)	300 (41-650)	3500 (460-7800)
Belgium	2016	Female	85 to 89	76% (67-83%)	24% (17-33%)	2.5 (1.6-3.5)	2.1% (-0.087-5.2%)	2.5% (-0.22-6.4%)	330 (-27-830)	2800 (-110-7000)
Belgium	2016	Female	90 to 94	74% (64-83%)	26% (17-36%)	2.5 (1.5-3.7)	1.8% (-0.67-5.1%)	2% (-1-6%)	240 (-120-710)	1400 (-510-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belgium	2016	Female	95 plus	73% (59-84%)	27% (16-41%)	2.5 (1.4-3.9)	1.9% (-1.5-5.9%)	2% (-1.5-6.9%)	93 (-68-320)	370 (-200-1100)
Belgium	2016	Male	All Ages	89% (87-91%)	11% (8.7-13%)	3.9 (3-5)	8.9% (6.6-11%)	8.4% (5.4-12%)	4700 (3000-6700)	140000 (100000-180000)
Belgium	2016	Male	15 to 19	80% (72-86%)	20% (14-28%)	2.8 (2-4)	5.6% (3.6-7.8%)	14% (7.6-22%)	16 (8-24)	1700 (1100-2500)
Belgium	2016	Male	20 to 24	87% (78-93%)	13% (6.6-22%)	4.2 (2.9-5.9)	12% (8.4-16%)	22% (13-31%)	50 (29-73)	5000 (3400-6800)
Belgium	2016	Male	25 to 29	90% (82-95%)	10% (5.1-18%)	3.4 (2.3-4.7)	10% (6.8-13%)	19% (9.4-27%)	46 (23-69)	4600 (3000-6200)
Belgium	2016	Male	30 to 34	92% (84-96%)	8.3% (3.8-16%)	3.4 (2.4-4.8)	9.1% (6.1-12%)	19% (10-26%)	57 (31-81)	4700 (3000-6300)
Belgium	2016	Male	35 to 39	93% (86-97%)	7.4% (3.2-14%)	4.4 (3-6.1)	10% (7.4-14%)	22% (14-30%)	90 (58-120)	6100 (4200-8100)
Belgium	2016	Male	40 to 44	93% (88-97%)	6.7% (3.1-12%)	5.1 (3.7-7)	12% (8.8-15%)	24% (18-31%)	150 (100-200)	8300 (6100-11000)
Belgium	2016	Male	45 to 49	94% (90-97%)	6.1% (3.3-10%)	5.7 (4.1-7.6)	13% (10-17%)	24% (19-30%)	250 (180-320)	12000 (9100-15000)
Belgium	2016	Male	50 to 54	95% (92-97%)	5% (2.8-8%)	6.1 (4.5-7.9)	14% (11-18%)	23% (18-28%)	400 (310-520)	17000 (13000-21000)
Belgium	2016	Male	55 to 59	96% (94-98%)	3.9% (2.5-5.8%)	6.4 (4.8-8.4)	14% (11-17%)	20% (16-24%)	560 (430-710)	20000 (15000-25000)
Belgium	2016	Male	60 to 64	96% (94-98%)	3.7% (2.2-5.6%)	6.2 (4.6-8.1)	12% (8.7-15%)	16% (12-20%)	600 (440-780)	18000 (13000-23000)
Belgium	2016	Male	65 to 69	96% (94-98%)	4.1% (2.5-6.4%)	5.4 (4.1-7)	9.3% (6.5-12%)	12% (8.2-16%)	610 (420-830)	16000 (11000-21000)
Belgium	2016	Male	70 to 74	95% (93-97%)	5% (3.3-7.4%)	4.6 (3.4-6.2)	6.8% (4.1-10%)	8.3% (5-12%)	440 (260-660)	9200 (5500-14000)
Belgium	2016	Male	75 to 79	94% (90-96%)	6.4% (3.8-9.9%)	4.1 (3-5.5)	5% (2.3-8.2%)	5.8% (2.6-9.8%)	430 (190-720)	7100 (3200-12000)
Belgium	2016	Male	80 to 84	92% (87-96%)	7.7% (4.4-13%)	3.8 (2.7-5.1)	4.1% (1.5-7.2%)	4.6% (1.5-8.2%)	450 (140-820)	5400 (2000-9500)
Belgium	2016	Male	85 to 89	92% (87-95%)	8.4% (5.1-13%)	3.6 (2.5-4.9)	3.4% (0.8-6.6%)	3.6% (0.57-7.3%)	350 (55-680)	3000 (690-5700)
Belgium	2016	Male	90 to 94	92% (86-95%)	8.4% (4.8-14%)	3.4 (2.3-4.7)	2.8% (-0.11-6.2%)	2.8% (-0.55-6.9%)	160 (-31-400)	990 (-38-2300)
Belgium	2016	Male	95 plus	92% (85-96%)	8.4% (3.9-15%)	3.2 (2-4.8)	2.5% (-0.76-6.6%)	2.5% (-1.3-7.4%)	31 (-16-94)	130 (-39-350)
Cyprus	1990	Female	All Ages	71% (62-79%)	29% (21-38%)	1 (0.86-1.2)	-0.18% (-1.4-1.1%)	-2% (-4.3-0.32%)	-59 (-120-9.6)	-150 (-1200-910)
Cyprus	1990	Female	15 to 19	69% (49-85%)	31% (15-51%)	1 (0.66-1.5)	1.7% (0.88-2.6%)	5.6% (2-10%)	0.4 (0.15-0.72)	51 (28-79)
Cyprus	1990	Female	20 to 24	75% (56-88%)	25% (12-44%)	1.4 (0.91-1.9)	2.3% (1.5-3.2%)	5.7% (2.7-9.1%)	0.45 (0.2-0.71)	73 (47-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	1990	Female	25 to 29	77% (60-90%)	23% (10-40%)	1.3 (0.94-1.8)	2.1% (1.3-2.8%)	5.1% (2.4-7.9%)	0.55 (0.27-0.87)	71 (45-99)
Cyprus	1990	Female	30 to 34	78% (60-90%)	22% (9.6-40%)	1.2 (0.82-1.7)	1.8% (1.1-2.5%)	4.8% (2.5-7.3%)	0.62 (0.32-0.94)	61 (38-87)
Cyprus	1990	Female	35 to 39	80% (64-91%)	20% (9-36%)	1.1 (0.72-1.6)	1.7% (0.96-2.4%)	4.8% (2.6-7.1%)	0.88 (0.47-1.3)	65 (39-94)
Cyprus	1990	Female	40 to 44	81% (63-92%)	19% (8.3-37%)	1.3 (0.87-1.8)	1.7% (0.9-2.7%)	5.1% (2.9-7.7%)	1.3 (0.72-1.9)	72 (38-110)
Cyprus	1990	Female	45 to 49	81% (61-91%)	19% (8.8-39%)	1.8 (1.2-2.5)	2.5% (1.2-3.9%)	6.4% (3.3-9.6%)	2.5 (1.3-3.8)	110 (53-180)
Cyprus	1990	Female	50 to 54	80% (61-91%)	20% (8.9-39%)	1.9 (1.3-2.6)	2.5% (1-4.1%)	5.6% (2.8-8.3%)	3.1 (1.5-4.6)	120 (45-180)
Cyprus	1990	Female	55 to 59	79% (62-91%)	21% (9.1-38%)	1.7 (1.1-2.3)	1.5% (-0.23-3.2%)	3.5% (0.71-6.3%)	2.7 (0.54-4.8)	71 (-10-150)
Cyprus	1990	Female	60 to 64	78% (58-91%)	22% (8.8-42%)	1.5 (0.94-2.1)	0.027% (-1.8-2%)	0.98% (-1.9-3.9%)	1.1 (-2.2-4.6)	1.1 (-100-110)
Cyprus	1990	Female	65 to 69	76% (55-90%)	24% (10-45%)	1.4 (0.89-1.9)	-1.2% (-3.5-1%)	-1.1% (-4-1.7%)	-2 (-7.2-3)	-73 (-200-60)
Cyprus	1990	Female	70 to 74	73% (53-86%)	27% (14-47%)	1.2 (0.78-1.8)	-2.2% (-4.6-0.24%)	-2.5% (-5.5-0.39%)	-7.8 (-18-1.2)	-160 (-350-17)
Cyprus	1990	Female	75 to 79	69% (49-85%)	31% (15-51%)	1.2 (0.71-1.8)	-3% (-5.8--0.23%)	-3.4% (-6.8--0.27%)	-16 (-32--1.3)	-230 (-450--18)
Cyprus	1990	Female	80 to 84	66% (44-83%)	34% (17-56%)	1.1 (0.7-1.7)	-2.9% (-5.7--0.54%)	-3.4% (-6.5--0.62%)	-22 (-43--4.1)	-240 (-450--43)
Cyprus	1990	Female	85 to 89	63% (43-80%)	37% (20-57%)	1.1 (0.65-1.7)	-2.5% (-5--0.33%)	-3% (-5.8--0.48%)	-15 (-30--2.4)	-110 (-220--15)
Cyprus	1990	Female	90 to 94	62% (43-80%)	38% (20-57%)	1.1 (0.61-1.8)	-2.1% (-4.4-0.13%)	-2.6% (-5.4-0.045%)	-6.6 (-14-0.11)	-32 (-67-1.9)
Cyprus	1990	Female	95 plus	63% (38-83%)	37% (17-62%)	1.2 (0.54-2)	-1.9% (-4.6-0.55%)	-2.4% (-5.7-0.47%)	-2.1 (-5-0.4)	-6.5 (-16-2)
Cyprus	1990	Male	All Ages	78% (68-86%)	22% (14-32%)	3.6 (3.1-4.1)	8% (5.6-10%)	8.4% (4.7-12%)	260 (140-380)	7500 (5300-9600)
Cyprus	1990	Male	15 to 19	73% (52-88%)	27% (12-48%)	3.4 (2.2-4.8)	7.9% (4.8-11%)	15% (8.2-23%)	3.4 (1.8-5.2)	300 (180-440)
Cyprus	1990	Male	20 to 24	78% (60-90%)	22% (9.6-40%)	5.3 (3.8-7.1)	15% (10-21%)	23% (14-33%)	7.6 (4.8-11)	620 (430-850)
Cyprus	1990	Male	25 to 29	82% (68-92%)	18% (8-32%)	4.8 (3.6-6.3)	13% (8.8-17%)	21% (13-29%)	5.9 (3.7-8.2)	470 (320-620)
Cyprus	1990	Male	30 to 34	85% (69-94%)	15% (6.1-31%)	4.4 (3.1-5.9)	10% (6.8-13%)	17% (11-24%)	4.6 (2.9-6.4)	340 (230-450)
Cyprus	1990	Male	35 to 39	87% (74-95%)	13% (4.9-26%)	4 (2.8-5.5)	8.3% (5.6-11%)	15% (9.2-21%)	4.5 (2.8-6.5)	310 (200-420)
Cyprus	1990	Male	40 to 44	88% (75-96%)	12% (4.1-25%)	4.5 (3.3-5.9)	8.4% (5.4-12%)	14% (8.1-20%)	6 (3.5-8.7)	350 (220-480)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	1990	Male	45 to 49	89% (72-96%)	11% (4-28%)	6 (4.4-8)	11% (7.4-15%)	17% (11-23%)	11 (6.6-14)	520 (340-690)
Cyprus	1990	Male	50 to 54	89% (75-96%)	11% (3.8-25%)	6.6 (4.7-8.7)	12% (7.5-16%)	17% (10-23%)	17 (9.9-23)	700 (430-930)
Cyprus	1990	Male	55 to 59	89% (77-96%)	11% (3.6-23%)	6.2 (4.6-8.1)	11% (6.4-15%)	14% (7.9-20%)	19 (11-27)	700 (420-970)
Cyprus	1990	Male	60 to 64	89% (75-96%)	11% (3.8-25%)	5.8 (4.1-7.9)	10% (5-15%)	12% (5.7-18%)	25 (12-38)	760 (390-1100)
Cyprus	1990	Male	65 to 69	88% (74-96%)	12% (4.1-26%)	5.3 (4-6.9)	8.9% (4.4-13%)	10% (4.8-16%)	28 (13-43)	690 (340-1000)
Cyprus	1990	Male	70 to 74	87% (74-95%)	13% (5.1-26%)	4.8 (3.5-6.4)	7.5% (3-12%)	8.3% (3-14%)	30 (11-51)	600 (250-990)
Cyprus	1990	Male	75 to 79	86% (72-94%)	14% (5.6-28%)	4.4 (3-6.1)	6.5% (1.6-12%)	7% (1.4-13%)	32 (6.5-60)	490 (120-880)
Cyprus	1990	Male	80 to 84	84% (69-93%)	16% (6.6-31%)	4.2 (2.9-5.6)	5.8% (1.5-10%)	6.1% (1.3-11%)	33 (6.7-61)	370 (95-670)
Cyprus	1990	Male	85 to 89	82% (67-93%)	18% (7.4-33%)	3.9 (2.6-5.5)	5% (1.1-9.6%)	5.3% (0.84-10%)	21 (3.3-41)	170 (36-330)
Cyprus	1990	Male	90 to 94	83% (68-93%)	17% (6.7-32%)	3.8 (2.3-5.7)	4.3% (0.15-9.1%)	4.4% (-0.29-9.9%)	8.7 (-0.57-20)	50 (1.8-110)
Cyprus	1990	Male	95 plus	83% (65-95%)	17% (5.3-35%)	3.8 (2.1-6.1)	4% (-0.5-9.5%)	4.1% (-1-10%)	2.5 (-0.6-6.3)	9.6 (-1.2-23)
Cyprus	1995	Female	All Ages	71% (63-79%)	29% (21-37%)	1.1 (0.92-1.3)	0.068% (-1.1-1.2%)	-1.6% (-3.8-0.51%)	-50 (-120-16)	60 (-910-1000)
Cyprus	1995	Female	15 to 19	69% (50-85%)	31% (15-50%)	1.1 (0.67-1.5)	1.9% (1-3.1%)	5.9% (2.1-10%)	0.52 (0.18-0.91)	59 (32-91)
Cyprus	1995	Female	20 to 24	74% (56-88%)	26% (12-44%)	1.4 (0.94-2)	2.6% (1.7-3.6%)	5.9% (2.5-9.5%)	0.62 (0.26-0.99)	84 (53-120)
Cyprus	1995	Female	25 to 29	77% (60-90%)	23% (10-40%)	1.4 (0.97-1.9)	2.3% (1.4-3.2%)	5.3% (2.3-8.3%)	0.74 (0.31-1.2)	83 (51-120)
Cyprus	1995	Female	30 to 34	79% (61-91%)	21% (9.3-39%)	1.3 (0.86-1.8)	1.9% (1.2-2.7%)	5% (2.7-7.5%)	0.76 (0.4-1.1)	71 (44-100)
Cyprus	1995	Female	35 to 39	80% (64-91%)	20% (9.1-36%)	1.1 (0.74-1.7)	1.8% (1.1-2.6%)	5% (2.7-7.5%)	1.1 (0.59-1.6)	77 (45-110)
Cyprus	1995	Female	40 to 44	81% (64-92%)	19% (8.2-36%)	1.3 (0.94-1.9)	1.8% (0.94-2.8%)	5.4% (3.1-8%)	1.4 (0.76-2)	77 (43-120)
Cyprus	1995	Female	45 to 49	81% (61-92%)	19% (8.4-39%)	1.9 (1.3-2.6)	2.8% (1.4-4.3%)	6.8% (3.8-10%)	2.9 (1.6-4.3)	130 (67-200)
Cyprus	1995	Female	50 to 54	80% (62-91%)	20% (9.1-38%)	2 (1.4-2.7)	2.6% (1.1-4.2%)	6% (3-8.8%)	3.3 (1.6-4.8)	130 (51-200)
Cyprus	1995	Female	55 to 59	79% (62-91%)	21% (8.8-38%)	1.7 (1.2-2.4)	1.7% (0.0076-3.4%)	3.8% (1.1-6.4%)	3.1 (0.88-5.3)	88 (0.46-180)
Cyprus	1995	Female	60 to 64	78% (57-91%)	22% (8.8-43%)	1.5 (0.97-2.2)	0.31% (-1.5-2.4%)	1.3% (-1.3-4.3%)	1.5 (-1.5-5)	17 (-83-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	1995	Female	65 to 69	76% (55-90%)	24% (10-45%)	1.4 (0.94-2)	-0.88% (-3-1.3%)	-0.71% (-3.5-2.1%)	-1.3 (-6.6-4.1)	-56 (-190-82)
Cyprus	1995	Female	70 to 74	73% (54-87%)	27% (13-46%)	1.3 (0.83-1.9)	-1.8% (-4.2-0.51%)	-2.1% (-5-0.85%)	-6.6 (-16-2.6)	-140 (-330-40)
Cyprus	1995	Female	75 to 79	69% (48-86%)	31% (14-52%)	1.2 (0.76-1.9)	-2.5% (-5--0.013%)	-2.9% (-6.1-0.037%)	-13 (-26-0.16)	-190 (-380--0.97)
Cyprus	1995	Female	80 to 84	66% (44-83%)	34% (17-56%)	1.2 (0.73-1.7)	-2.5% (-5--0.21%)	-3% (-6--0.27%)	-20 (-41--1.8)	-210 (-430--18)
Cyprus	1995	Female	85 to 89	63% (44-80%)	37% (20-56%)	1.2 (0.68-1.8)	-2.2% (-4.6--0.068%)	-2.7% (-5.4--0.13%)	-16 (-31--0.76)	-110 (-230--3.4)
Cyprus	1995	Female	90 to 94	62% (43-80%)	38% (20-57%)	1.2 (0.65-1.9)	-1.9% (-4.1-0.26%)	-2.4% (-5-0.13%)	-7.2 (-16-0.4)	-35 (-77-4.7)
Cyprus	1995	Female	95 plus	63% (39-83%)	37% (17-61%)	1.2 (0.59-2.1)	-1.6% (-4.2-0.75%)	-2.1% (-5.3-0.64%)	-2.4 (-5.9-0.71)	-7.3 (-19-3.4)
Cyprus	1995	Male	All Ages	79% (69-86%)	21% (14-31%)	3.8 (3.4-4.4)	8.9% (6.4-12%)	9.3% (5.5-13%)	300 (180-440)	8600 (6300-11000)
Cyprus	1995	Male	15 to 19	73% (53-88%)	27% (12-47%)	3.6 (2.3-5.1)	9.3% (5.4-14%)	16% (8.6-24%)	4.7 (2.6-7.2)	390 (230-570)
Cyprus	1995	Male	20 to 24	78% (60-91%)	22% (9.2-40%)	5.6 (3.9-7.7)	16% (11-23%)	24% (15-34%)	8.7 (5.5-12)	690 (480-940)
Cyprus	1995	Male	25 to 29	82% (68-93%)	18% (7.2-32%)	5.1 (3.7-6.6)	14% (9.4-18%)	22% (14-30%)	6.6 (4.1-9.2)	510 (350-680)
Cyprus	1995	Male	30 to 34	85% (69-94%)	15% (5.6-31%)	4.6 (3.2-6.2)	11% (7.4-15%)	18% (12-25%)	5.9 (3.7-8.1)	410 (280-560)
Cyprus	1995	Male	35 to 39	87% (73-95%)	13% (5-27%)	4.2 (2.9-5.8)	9% (6-12%)	15% (9.5-22%)	6 (3.7-8.6)	390 (260-540)
Cyprus	1995	Male	40 to 44	88% (74-96%)	12% (4-26%)	4.8 (3.4-6.3)	9.1% (5.8-12%)	15% (9.2-21%)	7.4 (4.5-10)	420 (280-590)
Cyprus	1995	Male	45 to 49	89% (72-96%)	11% (3.9-28%)	6.3 (4.5-8.3)	12% (8.1-16%)	18% (12-24%)	13 (8-17)	610 (410-800)
Cyprus	1995	Male	50 to 54	89% (75-96%)	11% (4-25%)	6.9 (5-8.9)	12% (8.4-16%)	18% (12-23%)	17 (11-22)	710 (470-930)
Cyprus	1995	Male	55 to 59	89% (77-96%)	11% (3.7-23%)	6.6 (5-8.6)	12% (7.4-16%)	15% (9.4-21%)	22 (13-31)	800 (500-1100)
Cyprus	1995	Male	60 to 64	89% (75-96%)	11% (3.9-25%)	6.1 (4.5-8.3)	11% (6-15%)	13% (7.1-19%)	25 (14-37)	780 (430-1100)
Cyprus	1995	Male	65 to 69	88% (75-96%)	12% (4-25%)	5.6 (4.2-7.3)	9.6% (5.4-14%)	11% (5.8-16%)	31 (16-45)	760 (430-1100)
Cyprus	1995	Male	70 to 74	87% (74-95%)	13% (5.2-26%)	5.1 (3.7-6.7)	8.3% (3.7-13%)	9.3% (3.7-15%)	37 (15-59)	730 (310-1200)
Cyprus	1995	Male	75 to 79	86% (72-95%)	14% (5.4-28%)	4.6 (3.2-6.5)	7.2% (2.4-12%)	7.8% (2.3-14%)	36 (10-64)	560 (190-960)
Cyprus	1995	Male	80 to 84	84% (68-94%)	16% (6.3-32%)	4.4 (3.2-6)	6.5% (2.2-11%)	6.9% (2-12%)	42 (12-73)	480 (170-820)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	1995	Male	85 to 89	82% (66-93%)	18% (7.1-34%)	4.1 (2.7-5.9)	5.7% (1.7-10%)	6% (1.5-11%)	27 (6.5-49)	220 (65-390)
Cyprus	1995	Male	90 to 94	82% (66-94%)	18% (6.5-34%)	4 (2.5-6)	4.9% (0.8-9.8%)	5.1% (0.47-11%)	11 (1-24)	65 (11-130)
Cyprus	1995	Male	95 plus	83% (63-95%)	17% (5.2-37%)	4 (2.2-6.3)	4.6% (-0.062-10%)	4.7% (-0.58-11%)	3.5 (-0.42-8)	13 (-0.19-29)
Cyprus	2000	Female	All Ages	72% (63-79%)	28% (21-37%)	1.1 (0.93-1.3)	0.15% (-0.89-1.2%)	-1.5% (-3.5-0.53%)	-45 (-100-15)	130 (-770-1100)
Cyprus	2000	Female	15 to 19	69% (50-85%)	31% (15-50%)	1 (0.65-1.5)	2% (0.99-3.1%)	5.8% (2-11%)	0.57 (0.2-1)	61 (31-97)
Cyprus	2000	Female	20 to 24	74% (56-88%)	26% (12-44%)	1.4 (0.92-2)	2.7% (1.6-3.8%)	5.8% (2.4-9.6%)	0.73 (0.3-1.2)	92 (57-130)
Cyprus	2000	Female	25 to 29	77% (60-89%)	23% (11-40%)	1.4 (0.94-1.8)	2.2% (1.4-3.1%)	5.3% (2.4-8.3%)	0.64 (0.29-1)	76 (48-110)
Cyprus	2000	Female	30 to 34	79% (61-91%)	21% (9.2-39%)	1.3 (0.84-1.8)	1.9% (1.2-2.8%)	4.9% (2.5-7.6%)	0.86 (0.44-1.3)	77 (47-110)
Cyprus	2000	Female	35 to 39	80% (63-91%)	20% (9-37%)	1.1 (0.73-1.6)	1.7% (1-2.5%)	5% (2.7-7.3%)	1 (0.56-1.5)	76 (45-110)
Cyprus	2000	Female	40 to 44	81% (64-92%)	19% (8.4-36%)	1.3 (0.91-1.8)	1.7% (0.96-2.6%)	5.5% (3.2-8%)	1.3 (0.77-1.9)	77 (45-110)
Cyprus	2000	Female	45 to 49	81% (61-91%)	19% (8.5-39%)	1.8 (1.3-2.5)	2.6% (1.4-3.9%)	6.8% (4-9.7%)	2.7 (1.6-3.8)	120 (68-180)
Cyprus	2000	Female	50 to 54	80% (62-91%)	20% (8.8-38%)	2 (1.4-2.6)	2.7% (1.2-4.2%)	6% (3.1-8.7%)	3.7 (1.9-5.4)	140 (64-210)
Cyprus	2000	Female	55 to 59	79% (62-91%)	21% (8.6-38%)	1.7 (1.1-2.3)	1.6% (0.16-3.2%)	3.9% (1.4-6.6%)	2.9 (1-5)	83 (7.8-160)
Cyprus	2000	Female	60 to 64	78% (58-91%)	22% (9-42%)	1.5 (0.95-2.2)	0.38% (-1.3-2.3%)	1.5% (-0.99-4.3%)	1.6 (-1.1-4.6)	20 (-69-120)
Cyprus	2000	Female	65 to 69	76% (55-89%)	24% (11-45%)	1.4 (0.93-1.9)	-0.75% (-2.7-1.2%)	-0.53% (-3.2-2.2%)	-0.99 (-6-4)	-49 (-170-81)
Cyprus	2000	Female	70 to 74	72% (53-87%)	28% (13-47%)	1.3 (0.8-1.9)	-1.7% (-3.8-0.48%)	-2% (-4.7-0.78%)	-6.1 (-14-2.3)	-130 (-290-38)
Cyprus	2000	Female	75 to 79	69% (48-86%)	31% (14-52%)	1.2 (0.72-1.8)	-2.2% (-4.4--0.043%)	-2.7% (-5.5--0.068%)	-11 (-23--0.27)	-170 (-340--3.3)
Cyprus	2000	Female	80 to 84	66% (44-83%)	34% (17-56%)	1.1 (0.73-1.7)	-2.3% (-4.4--0.24%)	-2.8% (-5.5--0.34%)	-17 (-33--2.1)	-180 (-350--19)
Cyprus	2000	Female	85 to 89	63% (44-80%)	37% (20-56%)	1.1 (0.67-1.7)	-2.2% (-4.3--0.11%)	-2.7% (-5.2--0.22%)	-16 (-31--1.3)	-120 (-230--6)
Cyprus	2000	Female	90 to 94	62% (42-80%)	38% (20-58%)	1.1 (0.62-1.8)	-1.8% (-4-0.12%)	-2.3% (-4.9--0.051%)	-7.5 (-16--0.17)	-36 (-81-2.3)
Cyprus	2000	Female	95 plus	63% (39-83%)	37% (17-61%)	1.2 (0.57-2)	-1.6% (-4.1-0.66%)	-2.1% (-5.1-0.53%)	-2.6 (-6.2-0.64)	-7.8 (-20-3.3)
Cyprus	2000	Male	All Ages	80% (70-87%)	20% (13-30%)	3.9 (3.4-4.5)	8.6% (6.3-11%)	8.8% (5.4-13%)	290 (180-420)	8400 (6200-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	2000	Male	15 to 19	73% (52-88%)	27% (12-48%)	3.5 (2.2-5.1)	9.1% (5.4-14%)	15% (8.4-24%)	4.7 (2.5-7.1)	380 (230-550)
Cyprus	2000	Male	20 to 24	78% (61-90%)	22% (9.9-39%)	5.5 (3.8-7.4)	16% (11-22%)	24% (15-34%)	8.4 (5.3-12)	670 (460-910)
Cyprus	2000	Male	25 to 29	82% (67-92%)	18% (8-33%)	5 (3.6-6.5)	14% (9.4-18%)	21% (13-29%)	6.9 (4.3-9.6)	520 (350-700)
Cyprus	2000	Male	30 to 34	85% (69-94%)	15% (6.1-31%)	4.5 (3.2-6)	10% (7.2-14%)	18% (11-24%)	5.7 (3.7-7.9)	410 (280-560)
Cyprus	2000	Male	35 to 39	87% (75-95%)	13% (5-25%)	4.1 (2.9-5.6)	8.9% (5.9-12%)	15% (9.1-21%)	6.3 (3.9-8.9)	410 (270-560)
Cyprus	2000	Male	40 to 44	88% (75-96%)	12% (4-25%)	4.7 (3.4-6.1)	8.4% (5.5-12%)	14% (8.6-20%)	7 (4.2-9.9)	410 (270-570)
Cyprus	2000	Male	45 to 49	89% (73-96%)	11% (3.9-27%)	6.2 (4.4-8.2)	11% (7.8-15%)	18% (12-23%)	12 (8.1-17)	610 (420-810)
Cyprus	2000	Male	50 to 54	89% (75-96%)	11% (3.7-25%)	6.7 (4.9-8.8)	12% (7.7-15%)	17% (11-23%)	16 (10-22)	690 (460-920)
Cyprus	2000	Male	55 to 59	89% (77-96%)	11% (3.6-23%)	6.4 (4.7-8.3)	11% (6.8-16%)	15% (8.6-21%)	24 (13-33)	850 (520-1200)
Cyprus	2000	Male	60 to 64	89% (74-96%)	11% (3.9-26%)	6 (4.3-8)	10% (5.4-15%)	13% (6.3-18%)	25 (12-37)	760 (410-1100)
Cyprus	2000	Male	65 to 69	88% (74-96%)	12% (4.1-26%)	5.4 (4.1-6.9)	9% (4.9-13%)	10% (5.2-16%)	29 (14-44)	720 (400-1100)
Cyprus	2000	Male	70 to 74	87% (74-95%)	13% (5.2-26%)	4.9 (3.5-6.5)	7.7% (3.2-13%)	8.6% (3.3-14%)	33 (13-55)	670 (280-1100)
Cyprus	2000	Male	75 to 79	86% (72-94%)	14% (5.5-28%)	4.5 (3.1-6.1)	6.6% (2-11%)	7.2% (1.9-13%)	33 (8.9-58)	520 (160-890)
Cyprus	2000	Male	80 to 84	84% (69-93%)	16% (6.6-31%)	4.3 (3-5.7)	5.9% (2.1-10%)	6.4% (1.9-11%)	36 (10-63)	420 (150-720)
Cyprus	2000	Male	85 to 89	83% (67-93%)	17% (7.4-33%)	4 (2.7-5.6)	5.4% (1.5-9.9%)	5.7% (1.2-11%)	27 (5.9-52)	220 (61-410)
Cyprus	2000	Male	90 to 94	83% (67-93%)	17% (6.7-33%)	3.9 (2.5-5.9)	4.7% (0.54-9.8%)	4.9% (0.11-11%)	11 (0.26-25)	65 (7.5-140)
Cyprus	2000	Male	95 plus	83% (64-95%)	17% (5.4-36%)	3.9 (2.1-6.4)	4.4% (-0.19-10%)	4.6% (-0.69-11%)	3.4 (-0.5-8.5)	13 (-0.55-30)
Cyprus	2005	Female	All Ages	70% (62-79%)	30% (21-38%)	1 (0.87-1.2)	0.27% (-0.63-1.2%)	-1.2% (-3-0.57%)	-37 (-89-17)	250 (-580-1100)
Cyprus	2005	Female	15 to 19	70% (51-85%)	30% (15-49%)	0.98 (0.63-1.4)	1.9% (0.88-3%)	5.8% (1.6-10%)	0.51 (0.15-0.92)	57 (28-91)
Cyprus	2005	Female	20 to 24	77% (60-89%)	23% (11-40%)	1.3 (0.89-1.9)	2.5% (1.6-3.6%)	5.8% (2.3-9.5%)	0.64 (0.25-1)	90 (56-130)
Cyprus	2005	Female	25 to 29	77% (62-89%)	23% (11-38%)	1.3 (0.9-1.7)	2.1% (1.3-2.9%)	5.2% (2.2-8.3%)	0.6 (0.27-0.96)	78 (51-110)
Cyprus	2005	Female	30 to 34	76% (58-89%)	24% (11-42%)	1.1 (0.73-1.6)	1.8% (1.1-2.5%)	4.7% (2.4-7.1%)	0.75 (0.39-1.1)	73 (47-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	2005	Female	35 to 39	76% (60-89%)	24% (11-40%)	0.98 (0.63-1.4)	1.5% (0.95-2.2%)	4.9% (2.8-7.1%)	0.85 (0.5-1.2)	67 (43-95)
Cyprus	2005	Female	40 to 44	78% (60-90%)	22% (9.9-40%)	1.2 (0.8-1.6)	1.7% (0.97-2.5%)	5.3% (3.3-7.6%)	1.4 (0.9-2.1)	85 (52-120)
Cyprus	2005	Female	45 to 49	78% (58-90%)	22% (10-42%)	1.6 (1.1-2.2)	2.2% (1.2-3.3%)	6.5% (4.1-9%)	2.6 (1.6-3.6)	120 (69-180)
Cyprus	2005	Female	50 to 54	77% (58-89%)	23% (11-42%)	1.7 (1.2-2.3)	2.4% (1.2-3.7%)	5.7% (3.3-8.2%)	3.5 (2-5.1)	130 (68-200)
Cyprus	2005	Female	55 to 59	76% (58-89%)	24% (11-42%)	1.5 (1-2.1)	1.6% (0.38-2.9%)	3.9% (1.7-6.1%)	3.3 (1.5-5.2)	94 (24-170)
Cyprus	2005	Female	60 to 64	74% (54-90%)	26% (10-46%)	1.3 (0.83-1.9)	0.56% (-0.86-2.1%)	1.8% (-0.46-4.1%)	2 (-0.51-4.6)	32 (-50-120)
Cyprus	2005	Female	65 to 69	70% (49-86%)	30% (14-51%)	1.2 (0.75-1.6)	-0.31% (-1.7-1.2%)	0.086% (-1.9-2.1%)	0.16 (-3.6-3.9)	-21 (-120-82)
Cyprus	2005	Female	70 to 74	68% (48-84%)	32% (16-52%)	1.1 (0.66-1.6)	-1.2% (-2.8-0.42%)	-1.3% (-3.4-0.78%)	-3.7 (-9.5-2.2)	-90 (-210-33)
Cyprus	2005	Female	75 to 79	67% (46-83%)	33% (17-54%)	1 (0.63-1.6)	-1.9% (-3.9--0.15%)	-2.3% (-4.8--0.072%)	-11 (-22--0.32)	-160 (-330--12)
Cyprus	2005	Female	80 to 84	64% (42-81%)	36% (19-58%)	1 (0.63-1.5)	-2.1% (-4--0.3%)	-2.6% (-4.8--0.38%)	-15 (-29--2.2)	-170 (-320--23)
Cyprus	2005	Female	85 to 89	61% (41-78%)	39% (22-59%)	1 (0.59-1.5)	-1.9% (-3.8--0.23%)	-2.4% (-4.8--0.29%)	-14 (-27--1.7)	-100 (-200--12)
Cyprus	2005	Female	90 to 94	60% (40-79%)	40% (21-60%)	1 (0.56-1.7)	-1.7% (-3.7--0.0055%)	-2.2% (-4.5--0.13%)	-7.4 (-16--0.46)	-36 (-77--0.12)
Cyprus	2005	Female	95 plus	61% (36-82%)	39% (18-64%)	1.1 (0.51-1.9)	-1.5% (-3.8-0.55%)	-2% (-4.7-0.38%)	-2.7 (-6.3-0.52)	-8.2 (-20-3)
Cyprus	2005	Male	All Ages	83% (74-89%)	17% (11-26%)	4 (3.4-4.5)	8.1% (5.9-10%)	8% (4.6-12%)	260 (150-390)	8200 (6100-11000)
Cyprus	2005	Male	15 to 19	76% (59-89%)	24% (11-41%)	3.4 (2.2-4.8)	8.5% (4.9-13%)	15% (8.1-23%)	4 (2.1-6.1)	340 (200-490)
Cyprus	2005	Male	20 to 24	82% (67-92%)	18% (7.9-33%)	5.4 (3.9-7.1)	15% (11-21%)	23% (14-33%)	9.2 (5.7-13)	750 (510-1000)
Cyprus	2005	Male	25 to 29	86% (75-94%)	14% (6-25%)	4.9 (3.7-6.3)	13% (9-17%)	21% (13-28%)	7.6 (4.7-11)	590 (390-780)
Cyprus	2005	Male	30 to 34	88% (77-95%)	12% (4.7-23%)	4.4 (3.2-5.9)	10% (7.1-14%)	17% (11-24%)	5.8 (3.7-8.1)	420 (290-570)
Cyprus	2005	Male	35 to 39	88% (77-95%)	12% (4.5-23%)	3.9 (2.8-5.3)	8.5% (5.8-12%)	14% (9.2-20%)	5.3 (3.3-7.4)	350 (240-480)
Cyprus	2005	Male	40 to 44	90% (78-96%)	10% (3.9-22%)	4.4 (3.3-5.8)	8% (5.2-11%)	13% (8-19%)	6.7 (4-9.6)	400 (260-560)
Cyprus	2005	Male	45 to 49	91% (80-97%)	9.1% (3.5-20%)	6 (4.4-7.8)	10% (6.8-14%)	17% (10-22%)	13 (7.9-17)	630 (400-840)
Cyprus	2005	Male	50 to 54	90% (79-97%)	9.5% (3.4-21%)	6.5 (4.8-8.3)	11% (7-15%)	16% (9.9-22%)	17 (10-23)	720 (480-970)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	2005	Male	55 to 59	90% (80-96%)	9.8% (3.5-20%)	6.1 (4.7-7.9)	10% (6.1-14%)	14% (7.8-19%)	23 (13-33)	850 (510-1200)
Cyprus	2005	Male	60 to 64	90% (79-97%)	9.8% (3.4-21%)	5.8 (4.2-7.6)	9.3% (4.6-14%)	11% (5.5-17%)	24 (11-36)	750 (370-1100)
Cyprus	2005	Male	65 to 69	89% (78-96%)	11% (3.5-22%)	5.3 (4-6.7)	8% (3.9-12%)	9.3% (4.3-14%)	28 (13-43)	730 (360-1100)
Cyprus	2005	Male	70 to 74	88% (77-95%)	12% (4.6-23%)	4.7 (3.4-6.2)	6.7% (2.6-11%)	7.5% (2.7-13%)	28 (9.9-47)	580 (230-960)
Cyprus	2005	Male	75 to 79	87% (75-95%)	13% (5-25%)	4.3 (3-5.9)	5.7% (1.7-10%)	6.2% (1.6-12%)	30 (7.4-54)	470 (140-840)
Cyprus	2005	Male	80 to 84	85% (70-94%)	15% (6.4-30%)	4 (2.9-5.4)	5% (1.6-8.7%)	5.3% (1.3-9.7%)	28 (7-51)	340 (110-600)
Cyprus	2005	Male	85 to 89	82% (67-93%)	18% (7.4-33%)	3.7 (2.5-5.3)	4.4% (1.2-8.5%)	4.6% (0.97-9.3%)	21 (4.3-42)	180 (48-340)
Cyprus	2005	Male	90 to 94	82% (67-93%)	18% (6.7-33%)	3.7 (2.3-5.4)	4% (0.14-8.9%)	4.1% (-0.29-9.5%)	11 (-0.76-25)	61 (2.1-140)
Cyprus	2005	Male	95 plus	83% (64-95%)	17% (5.3-36%)	3.6 (1.9-6)	3.8% (-0.41-9.4%)	3.8% (-0.92-10%)	2.9 (-0.71-7.9)	11 (-1.3-29)
Cyprus	2010	Female	All Ages	70% (62-78%)	30% (22-38%)	1 (0.85-1.2)	0.38% (-0.39-1.2%)	-0.94% (-2.6-0.68%)	-29 (-79-21)	370 (-380-1100)
Cyprus	2010	Female	15 to 19	69% (51-85%)	31% (15-49%)	0.96 (0.62-1.4)	1.5% (0.8-2.1%)	5.2% (1.5-9.5%)	0.3 (0.087-0.55)	44 (25-67)
Cyprus	2010	Female	20 to 24	76% (59-88%)	24% (12-41%)	1.3 (0.88-1.9)	2.1% (1.4-2.9%)	5.3% (2.3-8.3%)	0.41 (0.17-0.65)	84 (54-120)
Cyprus	2010	Female	25 to 29	77% (61-89%)	23% (11-39%)	1.3 (0.87-1.7)	1.8% (1.2-2.5%)	4.6% (2.3-7.2%)	0.52 (0.25-0.81)	85 (56-120)
Cyprus	2010	Female	30 to 34	75% (57-89%)	25% (11-43%)	1.1 (0.68-1.5)	1.5% (1-2.1%)	4.4% (2.5-6.4%)	0.6 (0.34-0.9)	72 (48-100)
Cyprus	2010	Female	35 to 39	76% (59-89%)	24% (11-41%)	0.95 (0.6-1.4)	1.4% (0.87-2%)	4.7% (2.9-6.7%)	0.75 (0.46-1.1)	67 (43-93)
Cyprus	2010	Female	40 to 44	77% (59-90%)	23% (10-41%)	1.1 (0.74-1.6)	1.6% (0.94-2.3%)	5.3% (3.4-7.4%)	1.3 (0.84-1.9)	82 (51-120)
Cyprus	2010	Female	45 to 49	78% (56-90%)	22% (10-44%)	1.6 (1.1-2.3)	2.1% (1.2-3.2%)	6.6% (4.3-9.3%)	2.5 (1.6-3.5)	120 (72-180)
Cyprus	2010	Female	50 to 54	76% (57-89%)	24% (11-43%)	1.7 (1.2-2.4)	2.3% (1.2-3.5%)	5.9% (3.6-8.3%)	3.6 (2.2-5.2)	140 (75-220)
Cyprus	2010	Female	55 to 59	76% (58-89%)	24% (11-42%)	1.4 (0.97-2.1)	1.5% (0.37-2.7%)	4.1% (2.1-6.4%)	3.2 (1.6-5)	94 (24-170)
Cyprus	2010	Female	60 to 64	74% (54-89%)	26% (11-46%)	1.3 (0.79-1.9)	0.62% (-0.61-2%)	2.1% (0.053-4.3%)	2.5 (0.064-5.1)	42 (-44-130)
Cyprus	2010	Female	65 to 69	69% (46-86%)	31% (14-54%)	1.1 (0.72-1.7)	-0.071% (-1.3-1.2%)	0.6% (-1.2-2.3%)	0.96 (-1.8-3.7)	-5.1 (-89-77)
Cyprus	2010	Female	70 to 74	67% (47-83%)	33% (17-53%)	1 (0.64-1.6)	-0.75% (-2.1-0.62%)	-0.68% (-2.6-1.2%)	-1.7 (-6.3-3)	-56 (-160-46)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	2010	Female	75 to 79	66% (44-83%)	34% (17-56%)	1 (0.6-1.6)	-1.4% (-3.1-0.1%)	-1.7% (-3.9-0.24%)	-7.2 (-16-1)	-120 (-260-8.8)
Cyprus	2010	Female	80 to 84	63% (41-81%)	37% (19-59%)	1 (0.61-1.5)	-1.8% (-3.5--0.17%)	-2.2% (-4.4--0.25%)	-14 (-27--1.5)	-150 (-300--14)
Cyprus	2010	Female	85 to 89	60% (40-78%)	40% (22-60%)	0.99 (0.56-1.6)	-1.6% (-3.3--0.039%)	-2% (-4.2--0.1%)	-13 (-26--0.64)	-95 (-190--2.4)
Cyprus	2010	Female	90 to 94	59% (39-78%)	41% (22-61%)	1 (0.54-1.6)	-1.3% (-3-0.13%)	-1.8% (-3.9-0.036%)	-7 (-15-0.14)	-34 (-76-3.3)
Cyprus	2010	Female	95 plus	61% (36-82%)	39% (18-64%)	1 (0.49-1.9)	-1.2% (-3.1-0.53%)	-1.7% (-4.1-0.42%)	-2.7 (-6.6-0.66)	-8 (-21-3.7)
Cyprus	2010	Male	All Ages	83% (75-89%)	17% (11-25%)	4.1 (3.4-4.8)	7.5% (5.4-9.8%)	7.4% (4.2-11%)	250 (140-380)	7900 (5600-10000)
Cyprus	2010	Male	15 to 19	76% (57-89%)	24% (11-43%)	3.4 (2.2-4.9)	6.8% (4.1-9.8%)	14% (7.8-22%)	2.6 (1.4-3.9)	240 (150-340)
Cyprus	2010	Male	20 to 24	82% (66-92%)	18% (8.3-34%)	5.4 (3.8-7.2)	14% (9.4-18%)	21% (14-30%)	7.3 (4.7-10)	650 (460-870)
Cyprus	2010	Male	25 to 29	86% (75-93%)	14% (6.5-25%)	4.9 (3.6-6.3)	11% (7.9-15%)	19% (12-26%)	6.4 (4.1-8.8)	550 (390-730)
Cyprus	2010	Male	30 to 34	88% (77-95%)	12% (5-23%)	4.4 (3.1-5.9)	9% (6.3-12%)	16% (10-21%)	5.3 (3.4-7.3)	420 (290-560)
Cyprus	2010	Male	35 to 39	88% (77-95%)	12% (4.6-23%)	3.9 (2.8-5.4)	7.5% (5.1-10%)	13% (8.7-19%)	4.5 (2.9-6.4)	320 (220-440)
Cyprus	2010	Male	40 to 44	90% (77-96%)	10% (3.8-23%)	4.4 (3.2-5.9)	7.6% (4.9-10%)	13% (7.6-19%)	5.9 (3.5-8.7)	360 (240-510)
Cyprus	2010	Male	45 to 49	91% (77-96%)	9.1% (3.6-23%)	5.9 (4.3-8)	9.7% (6.2-13%)	16% (9.5-21%)	11 (6.7-15)	580 (370-770)
Cyprus	2010	Male	50 to 54	90% (78-96%)	9.6% (3.6-22%)	6.4 (4.8-8.5)	10% (6.7-14%)	15% (9.5-20%)	16 (10-22)	730 (480-970)
Cyprus	2010	Male	55 to 59	90% (79-97%)	10% (3.4-21%)	6.1 (4.6-8)	9.5% (5.9-13%)	13% (7.5-18%)	20 (12-29)	780 (490-1100)
Cyprus	2010	Male	60 to 64	90% (78-97%)	10% (3.5-22%)	5.7 (4.2-7.8)	8.5% (4.5-13%)	11% (5.3-16%)	25 (13-38)	820 (450-1200)
Cyprus	2010	Male	65 to 69	89% (77-96%)	11% (3.9-23%)	5.2 (4-6.8)	7.4% (3.8-11%)	8.8% (4.1-14%)	26 (13-41)	710 (370-1100)
Cyprus	2010	Male	70 to 74	88% (76-95%)	12% (4.8-24%)	4.7 (3.3-6.3)	6.2% (2.6-10%)	7% (2.6-12%)	29 (11-49)	610 (260-1000)
Cyprus	2010	Male	75 to 79	87% (74-95%)	13% (5.3-26%)	4.2 (3-6)	5.4% (1.8-9.6%)	5.9% (1.7-11%)	28 (7.9-52)	460 (160-840)
Cyprus	2010	Male	80 to 84	84% (68-93%)	16% (6.8-32%)	3.9 (2.7-5.4)	4.7% (1.5-8.7%)	5.1% (1.3-9.9%)	28 (6.9-53)	340 (110-630)
Cyprus	2010	Male	85 to 89	82% (66-92%)	18% (7.9-34%)	3.7 (2.4-5.4)	4.3% (1.2-8.5%)	4.5% (0.94-9.4%)	23 (4.8-47)	190 (55-390)
Cyprus	2010	Male	90 to 94	82% (67-93%)	18% (7.1-33%)	3.6 (2.3-5.5)	3.9% (0.31-8.5%)	4% (-0.12-9.2%)	11 (-0.34-26)	66 (5.3-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	2010	Male	95 plus	82% (64-94%)	18% (5.6-36%)	3.5 (1.9-5.9)	3.8% (-0.3-9.2%)	3.9% (-0.95-10%)	3.2 (-0.79-8.5)	13 (-0.99-31)
Cyprus	2016	Female	All Ages	72% (64-80%)	28% (20-36%)	1.1 (0.77-1.5)	0.38% (-0.51-1.3%)	-0.93% (-2.7-0.86%)	-33 (-95-30)	400 (-510-1400)
Cyprus	2016	Female	15 to 19	68% (48-84%)	32% (16-52%)	0.92 (0.54-1.4)	1.4% (0.77-2.1%)	5% (1.4-9%)	0.25 (0.071-0.45)	40 (23-61)
Cyprus	2016	Female	20 to 24	73% (55-87%)	27% (13-45%)	1.3 (0.75-2)	2.1% (1.4-2.9%)	5.1% (2.1-8.1%)	0.4 (0.16-0.64)	87 (56-120)
Cyprus	2016	Female	25 to 29	76% (59-89%)	24% (11-41%)	1.2 (0.78-1.9)	1.8% (1.2-2.4%)	4.5% (2-7.3%)	0.52 (0.23-0.83)	92 (61-130)
Cyprus	2016	Female	30 to 34	78% (60-90%)	22% (9.6-40%)	1.1 (0.69-1.8)	1.5% (0.99-2.1%)	4.5% (2.4-6.8%)	0.66 (0.36-1)	83 (55-120)
Cyprus	2016	Female	35 to 39	79% (63-91%)	21% (9.3-37%)	1 (0.62-1.6)	1.4% (0.9-1.9%)	4.9% (3.1-7.2%)	0.81 (0.51-1.2)	75 (48-110)
Cyprus	2016	Female	40 to 44	80% (63-91%)	20% (8.5-37%)	1.2 (0.72-1.8)	1.6% (0.94-2.3%)	5.6% (3.5-8.1%)	1.4 (0.86-2)	88 (55-130)
Cyprus	2016	Female	45 to 49	80% (60-91%)	20% (8.6-40%)	1.7 (1-2.5)	2.1% (1.1-3.3%)	6.9% (4.3-10%)	2.6 (1.6-3.7)	120 (71-190)
Cyprus	2016	Female	50 to 54	79% (61-91%)	21% (9.2-39%)	1.8 (1.1-2.6)	2.3% (1.1-3.7%)	6.2% (3.6-9.1%)	3.9 (2.2-5.8)	150 (75-240)
Cyprus	2016	Female	55 to 59	78% (62-91%)	22% (8.8-38%)	1.5 (0.94-2.3)	1.6% (0.42-3.1%)	4.4% (2.1-7.1%)	3.5 (1.7-5.7)	100 (26-190)
Cyprus	2016	Female	60 to 64	77% (56-91%)	23% (9.2-44%)	1.4 (0.81-2.2)	0.69% (-0.64-2.3%)	2.3% (0.17-4.8%)	3 (0.21-6.2)	52 (-51-170)
Cyprus	2016	Female	65 to 69	75% (53-89%)	25% (11-47%)	1.3 (0.77-2)	-0.096% (-1.5-1.5%)	0.59% (-1.5-2.7%)	1 (-2.5-5)	-7.3 (-110-110)
Cyprus	2016	Female	70 to 74	71% (51-86%)	29% (14-49%)	1.1 (0.66-1.8)	-0.77% (-2.2-0.84%)	-0.69% (-2.8-1.5%)	-1.8 (-7.4-4)	-62 (-180-67)
Cyprus	2016	Female	75 to 79	68% (46-85%)	32% (15-54%)	1 (0.59-1.7)	-1.3% (-3-0.55%)	-1.6% (-3.7-0.74%)	-6.7 (-16-3.2)	-110 (-250-46)
Cyprus	2016	Female	80 to 84	64% (42-82%)	36% (18-58%)	1 (0.56-1.6)	-1.6% (-3.4-0.092%)	-2.1% (-4.3-0.1%)	-14 (-29-0.69)	-150 (-310-9)
Cyprus	2016	Female	85 to 89	62% (42-79%)	38% (21-58%)	1 (0.54-1.7)	-1.5% (-3.3-0.12%)	-1.9% (-4.1-0.087%)	-15 (-32-0.65)	-110 (-240-8.8)
Cyprus	2016	Female	90 to 94	61% (41-79%)	39% (21-59%)	1 (0.53-1.8)	-1.3% (-2.9-0.4%)	-1.7% (-3.8-0.3%)	-9.6 (-21-1.7)	-46 (-110-15)
Cyprus	2016	Female	95 plus	62% (37-83%)	38% (17-63%)	1.1 (0.5-1.9)	-1.1% (-3-0.66%)	-1.6% (-4-0.56%)	-4.2 (-10-1.5)	-12 (-33-7.2)
Cyprus	2016	Male	All Ages	81% (73-87%)	19% (13-27%)	4 (2.9-5.2)	7.1% (4.4-10%)	7.1% (3.2-12%)	260 (120-430)	7900 (4900-11000)
Cyprus	2016	Male	15 to 19	72% (52-88%)	28% (12-48%)	3.2 (1.9-5)	6.4% (4-9.3%)	14% (7.7-22%)	2.2 (1.2-3.4)	210 (140-320)
Cyprus	2016	Male	20 to 24	78% (60-90%)	22% (9.7-40%)	5 (3.3-7.4)	13% (8.8-17%)	20% (12-29%)	6.7 (4-9.8)	620 (420-860)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cyprus	2016	Male	25 to 29	82% (67-92%)	18% (7.7-33%)	4.6 (3.1-6.5)	11% (7.5-14%)	18% (12-26%)	6.1 (3.9-8.8)	560 (390-750)
Cyprus	2016	Male	30 to 34	84% (68-94%)	16% (6-32%)	4.2 (2.7-6.2)	8.4% (5.8-11%)	15% (9.2-22%)	5.3 (3.2-7.6)	440 (300-600)
Cyprus	2016	Male	35 to 39	87% (73-95%)	13% (5.2-27%)	3.8 (2.5-5.7)	7.2% (4.9-10%)	13% (7.9-19%)	4.5 (2.7-6.7)	330 (220-470)
Cyprus	2016	Male	40 to 44	88% (75-96%)	12% (4.2-25%)	4.3 (2.8-6.3)	7.3% (4.6-11%)	13% (7.3-19%)	5.6 (3.1-8.6)	350 (220-510)
Cyprus	2016	Male	45 to 49	89% (75-96%)	11% (4-25%)	5.7 (3.6-8.4)	9.2% (5.2-13%)	15% (8.3-22%)	10 (5.4-14)	530 (310-740)
Cyprus	2016	Male	50 to 54	89% (76-96%)	11% (4-24%)	6.2 (4.1-8.9)	9.5% (5.5-13%)	15% (8-20%)	16 (8.3-22)	710 (400-980)
Cyprus	2016	Male	55 to 59	89% (77-96%)	11% (3.6-23%)	5.9 (4-8.5)	9.1% (5-13%)	13% (6.5-18%)	19 (10-29)	750 (420-1100)
Cyprus	2016	Male	60 to 64	88% (73-96%)	12% (3.8-27%)	5.6 (3.7-8)	8.1% (3.8-12%)	10% (4.7-16%)	26 (12-40)	850 (400-1300)
Cyprus	2016	Male	65 to 69	88% (73-96%)	12% (4.2-27%)	5.1 (3.4-7.2)	7.2% (3.4-11%)	8.5% (3.7-14%)	27 (12-44)	730 (330-1100)
Cyprus	2016	Male	70 to 74	87% (74-95%)	13% (5.3-26%)	4.5 (2.9-6.6)	6% (2.3-10%)	6.9% (2.3-12%)	30 (10-52)	640 (240-1100)
Cyprus	2016	Male	75 to 79	86% (72-94%)	14% (5.6-28%)	4.1 (2.6-6.2)	5.2% (1.3-9.6%)	5.7% (1.2-11%)	29 (6.1-56)	480 (120-910)
Cyprus	2016	Male	80 to 84	84% (67-93%)	16% (6.8-33%)	3.9 (2.5-5.7)	4.6% (1.2-8.8%)	5% (0.96-9.9%)	30 (5.5-60)	370 (91-720)
Cyprus	2016	Male	85 to 89	82% (65-92%)	18% (7.6-35%)	3.7 (2.2-5.9)	4.3% (0.89-9.1%)	4.5% (0.59-10%)	26 (3.5-59)	220 (46-470)
Cyprus	2016	Male	90 to 94	82% (66-93%)	18% (6.6-34%)	3.6 (2-5.9)	3.8% (0.19-8.9%)	4% (-0.24-9.9%)	14 (-0.83-33)	79 (3.8-180)
Cyprus	2016	Male	95 plus	82% (64-95%)	18% (5.1-36%)	3.5 (1.8-6.2)	3.7% (-0.5-9.2%)	3.8% (-1.1-10%)	4.2 (-1.2-11)	16 (-2.2-41)
Denmark	1990	Female	All Ages	95% (94-96%)	4.9% (3.8-6.3%)	3.2 (2.8-3.6)	5.4% (3.3-7.5%)	5.8% (1.5-10%)	1700 (480-3100)	44000 (26000-62000)
Denmark	1990	Female	15 to 19	94% (88-97%)	6.4% (3.1-12%)	3 (2.2-3.9)	3.4% (2-4.9%)	13% (6.5-19%)	6.6 (3.4-9.7)	650 (390-910)
Denmark	1990	Female	20 to 24	95% (92-98%)	4.5% (2.3-8.4%)	3.5 (2.4-4.6)	4.2% (2.8-5.8%)	14% (7.7-20%)	9.4 (5.2-14)	1000 (670-1400)
Denmark	1990	Female	25 to 29	96% (94-98%)	3.6% (1.8-6.3%)	3 (2.2-3.8)	3.5% (2.1-4.8%)	12% (5.6-17%)	9.4 (4.4-14)	920 (560-1300)
Denmark	1990	Female	30 to 34	97% (94-98%)	3.3% (1.6-6%)	2.8 (2.1-3.7)	3.5% (2.2-5%)	11% (6.2-16%)	13 (6.9-18)	970 (590-1400)
Denmark	1990	Female	35 to 39	96% (94-98%)	3.6% (1.7-6%)	2.9 (2.1-4)	4.9% (3.1-6.8%)	14% (8.5-18%)	27 (17-37)	1600 (1100-2200)
Denmark	1990	Female	40 to 44	96% (94-98%)	3.6% (1.9-6%)	3.5 (2.7-4.4)	6.7% (4.8-8.7%)	16% (11-20%)	58 (42-73)	3000 (2200-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	1990	Female	45 to 49	97% (95-98%)	3.3% (2-5.2%)	4.4 (3.4-5.5)	8.4% (6.4-10%)	17% (13-21%)	90 (70-110)	4100 (3200-5000)
Denmark	1990	Female	50 to 54	97% (95-98%)	2.9% (1.8-4.5%)	4.9 (3.8-6.1)	9.5% (7.2-12%)	16% (13-20%)	120 (90-140)	4600 (3500-5700)
Denmark	1990	Female	55 to 59	97% (96-98%)	2.6% (1.7-3.8%)	5.1 (4-6.3)	9.1% (6.6-12%)	14% (10-17%)	140 (100-180)	4900 (3500-6300)
Denmark	1990	Female	60 to 64	97% (96-98%)	2.7% (1.7-4.1%)	5.1 (4-6.3)	7.8% (4.8-11%)	11% (6.8-15%)	160 (100-230)	4900 (3000-6800)
Denmark	1990	Female	65 to 69	97% (95-98%)	3.1% (1.9-4.9%)	4.8 (3.9-5.7)	6.4% (3.2-9.3%)	8.5% (4.2-12%)	190 (94-290)	4800 (2400-7100)
Denmark	1990	Female	70 to 74	96% (94-97%)	3.8% (2.5-5.8%)	4.3 (3.3-5.5)	5.4% (1.9-9.3%)	6.8% (1.9-12%)	210 (58-370)	4200 (1300-7200)
Denmark	1990	Female	75 to 79	95% (92-97%)	4.8% (2.7-8.3%)	4 (2.9-5.2)	4.3% (0.05-9.1%)	5.1% (-0.45-11%)	220 (-19-470)	3500 (43-7300)
Denmark	1990	Female	80 to 84	93% (88-96%)	7.2% (4.5-12%)	3.7 (2.9-4.8)	3.6% (-0.89-8.4%)	4.1% (-1.8-10%)	220 (-94-550)	2600 (-640-6100)
Denmark	1990	Female	85 to 89	89% (83-93%)	11% (7.2-17%)	3.6 (2.7-4.6)	2.8% (-1.7-7.7%)	3% (-2.7-9.2%)	150 (-140-470)	1300 (-800-3700)
Denmark	1990	Female	90 to 94	86% (79-91%)	14% (8.7-21%)	3.4 (2.3-4.6)	2% (-2.5-7.2%)	2% (-3.6-8.6%)	60 (-110-260)	390 (-470-1400)
Denmark	1990	Female	95 plus	83% (74-90%)	17% (9.7-26%)	3.2 (1.9-4.8)	1.5% (-3.6-7.8%)	1.3% (-4.9-9.1%)	14 (-52-98)	67 (-160-350)
Denmark	1990	Male	All Ages	97% (96-98%)	3.2% (2.2-4.5%)	4.6 (4.1-5.1)	11% (8.3-14%)	11% (7-16%)	3400 (2100-4800)	97000 (72000-120000)
Denmark	1990	Male	15 to 19	94% (89-97%)	5.9% (2.8-11%)	4 (3-5.1)	8.1% (5.3-11%)	18% (10-25%)	19 (11-27)	1700 (1100-2400)
Denmark	1990	Male	20 to 24	96% (92-98%)	4.2% (2-7.7%)	5.7 (4.4-7)	15% (11-19%)	26% (17-34%)	48 (32-64)	4100 (3000-5300)
Denmark	1990	Male	25 to 29	97% (94-99%)	3% (1.5-5.6%)	4.9 (3.9-6.1)	13% (9-16%)	22% (14-29%)	48 (30-65)	3900 (2800-5000)
Denmark	1990	Male	30 to 34	98% (95-99%)	2.2% (0.93-4.6%)	4.3 (3.3-5.5)	11% (7.7-15%)	19% (12-26%)	54 (33-75)	3800 (2500-5100)
Denmark	1990	Male	35 to 39	98% (96-99%)	1.8% (0.83-3.6%)	4 (3-5.2)	11% (8-15%)	19% (13-25%)	70 (46-93)	4400 (3100-5600)
Denmark	1990	Male	40 to 44	98% (97-99%)	1.6% (0.63-3.1%)	4.8 (3.8-6)	13% (10-17%)	21% (15-26%)	130 (92-160)	6900 (5200-8500)
Denmark	1990	Male	45 to 49	99% (97-99%)	1.4% (0.63-2.8%)	6.7 (5.4-8.1)	17% (13-20%)	25% (20-31%)	200 (150-250)	9300 (7300-11000)
Denmark	1990	Male	50 to 54	99% (98-99%)	1.2% (0.57-2.4%)	7.6 (6.2-9)	18% (14-21%)	24% (19-29%)	250 (190-300)	9800 (7600-12000)
Denmark	1990	Male	55 to 59	99% (98-100%)	1.1% (0.49-1.9%)	7.5 (6.1-8.9)	16% (12-20%)	20% (14-25%)	300 (210-380)	10000 (7400-13000)
Denmark	1990	Male	60 to 64	99% (98-100%)	1.2% (0.47-2.2%)	7.2 (5.9-8.7)	13% (9-18%)	16% (10-21%)	360 (230-490)	11000 (6900-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	1990	Male	65 to 69	99% (97-99%)	1.4% (0.54-3%)	6.8 (5.7-7.9)	11% (7-16%)	13% (7.7-18%)	450 (260-620)	11000 (6700-15000)
Denmark	1990	Male	70 to 74	98% (97-99%)	1.7% (0.74-3.4%)	6.4 (5.3-7.6)	9.6% (5-14%)	11% (5.3-16%)	460 (220-680)	9000 (4600-13000)
Denmark	1990	Male	75 to 79	98% (96-99%)	2% (0.84-3.8%)	5.9 (4.8-7.1)	8.2% (3.6-13%)	8.8% (3.4-14%)	450 (180-740)	7000 (3100-11000)
Denmark	1990	Male	80 to 84	98% (96-99%)	2.4% (1.1-4.5%)	5.4 (4.4-6.6)	6.9% (2.3-12%)	7.3% (1.8-13%)	340 (86-600)	3900 (1300-6700)
Denmark	1990	Male	85 to 89	97% (95-99%)	2.7% (1.3-4.8%)	5 (4-6.2)	5.3% (0.71-10%)	5.5% (0.12-11%)	170 (3.6-340)	1400 (190-2700)
Denmark	1990	Male	90 to 94	97% (95-99%)	2.7% (1.2-4.9%)	4.7 (3.5-6.1)	3.6% (-1.3-9.2%)	3.6% (-2.3-10%)	48 (-31-140)	300 (-110-760)
Denmark	1990	Male	95 plus	97% (95-99%)	2.7% (1-5.5%)	4.4 (2.9-6.1)	2.4% (-2.9-8.4%)	2.1% (-4.1-9.3%)	8.2 (-16-36)	38 (-46-130)
Denmark	1995	Female	All Ages	96% (94-97%)	4.5% (3.5-5.7%)	3.2 (2.9-3.6)	5.5% (3.6-7.5%)	6% (2.1-10%)	1900 (650-3200)	45000 (30000-62000)
Denmark	1995	Female	15 to 19	94% (90-97%)	5.7% (2.8-10%)	3 (2.2-4)	3.1% (1.9-4.4%)	12% (6.4-18%)	4.8 (2.6-7)	500 (310-710)
Denmark	1995	Female	20 to 24	96% (93-98%)	4% (2-7.4%)	3.5 (2.4-4.7)	3.9% (2.7-5.3%)	13% (7.4-18%)	7.8 (4.6-11)	890 (610-1200)
Denmark	1995	Female	25 to 29	97% (95-98%)	3.3% (1.7-5.4%)	3 (2.3-3.8)	3.2% (2.1-4.4%)	10% (5.5-15%)	8.2 (4.4-12)	850 (550-1200)
Denmark	1995	Female	30 to 34	97% (95-99%)	2.9% (1.4-4.7%)	2.8 (2.1-3.7)	3.3% (2.2-4.6%)	11% (6.6-15%)	13 (7.8-18)	1000 (650-1400)
Denmark	1995	Female	35 to 39	97% (95-98%)	3.1% (1.6-5%)	3 (2.1-4.1)	4.8% (3.2-6.4%)	14% (9.1-18%)	26 (18-35)	1600 (1100-2100)
Denmark	1995	Female	40 to 44	97% (95-98%)	3.2% (1.7-5.1%)	3.5 (2.7-4.5)	6.5% (4.8-8.5%)	16% (12-20%)	51 (38-64)	2700 (2000-3400)
Denmark	1995	Female	45 to 49	97% (96-98%)	2.8% (1.9-4%)	4.5 (3.5-5.7)	8.3% (6.5-10%)	17% (14-21%)	97 (78-120)	4400 (3500-5400)
Denmark	1995	Female	50 to 54	97% (97-98%)	2.5% (1.8-3.4%)	5 (3.9-6.3)	9.3% (7.2-11%)	17% (14-20%)	130 (110-160)	5300 (4200-6500)
Denmark	1995	Female	55 to 59	98% (97-98%)	2.3% (1.6-3.1%)	5.2 (4.1-6.5)	9% (6.7-11%)	14% (11-17%)	140 (110-170)	4900 (3700-6100)
Denmark	1995	Female	60 to 64	98% (97-98%)	2.3% (1.6-3.3%)	5.2 (4-6.5)	7.9% (5.4-11%)	11% (7.6-15%)	170 (110-220)	4900 (3300-6500)
Denmark	1995	Female	65 to 69	97% (96-98%)	2.7% (1.8-4%)	4.9 (4-5.8)	6.7% (4-9.3%)	8.7% (5.2-12%)	200 (120-280)	4800 (2800-6800)
Denmark	1995	Female	70 to 74	97% (95-98%)	3.4% (2.4-4.8%)	4.4 (3.5-5.6)	5.7% (2.4-9.1%)	7.2% (2.8-12%)	230 (95-380)	4700 (2000-7400)
Denmark	1995	Female	75 to 79	96% (93-97%)	4.5% (2.6-7.2%)	4 (3.1-5.2)	4.6% (0.7-8.7%)	5.4% (0.48-11%)	230 (21-440)	3600 (570-6800)
Denmark	1995	Female	80 to 84	93% (90-96%)	6.8% (4.3-10%)	3.8 (3-4.8)	3.9% (-0.14-8.2%)	4.4% (-0.75-9.9%)	240 (-39-560)	2900 (-100-6300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	1995	Female	85 to 89	90% (86-93%)	9.7% (6.7-14%)	3.7 (2.8-4.7)	3.2% (-1.1-8.1%)	3.4% (-1.9-9.5%)	190 (-110-540)	1700 (-580-4200)
Denmark	1995	Female	90 to 94	87% (82-92%)	13% (8.1-18%)	3.5 (2.5-4.8)	2.3% (-2.2-7.8%)	2.3% (-3.2-9.1%)	86 (-120-330)	540 (-500-1800)
Denmark	1995	Female	95 plus	85% (77-91%)	15% (9.1-23%)	3.3 (2-5)	1.8% (-3.5-8.3%)	1.7% (-4.7-9.7%)	22 (-63-130)	100 (-190-450)
Denmark	1995	Male	All Ages	97% (96-98%)	2.7% (1.9-3.7%)	4.7 (4.1-5.3)	11% (8.7-14%)	12% (7.4-16%)	3500 (2200-4900)	100000 (78000-120000)
Denmark	1995	Male	15 to 19	95% (92-98%)	4.6% (2.3-8.1%)	4.1 (3.2-5.2)	8.6% (5.5-12%)	18% (11-26%)	19 (11-27)	1700 (1100-2200)
Denmark	1995	Male	20 to 24	96% (94-98%)	3.5% (1.8-6.2%)	5.8 (4.4-7.2)	14% (10-18%)	25% (17-34%)	42 (28-55)	3700 (2700-4700)
Denmark	1995	Male	25 to 29	97% (95-99%)	2.7% (1.4-4.6%)	4.9 (4-6.1)	12% (8.5-15%)	20% (13-27%)	44 (28-60)	3700 (2600-4800)
Denmark	1995	Male	30 to 34	98% (96-99%)	1.9% (0.89-3.6%)	4.4 (3.4-5.5)	10% (7-13%)	17% (11-23%)	48 (30-67)	3600 (2500-4700)
Denmark	1995	Male	35 to 39	98% (97-99%)	1.5% (0.69-2.7%)	4.1 (3-5.4)	11% (7.9-13%)	18% (12-22%)	65 (44-85)	4200 (3100-5300)
Denmark	1995	Male	40 to 44	99% (98-99%)	1.3% (0.59-2.5%)	4.9 (3.8-6.1)	13% (10-17%)	21% (16-26%)	120 (88-150)	6400 (5000-7900)
Denmark	1995	Male	45 to 49	99% (98-99%)	1.2% (0.58-2.2%)	6.8 (5.5-8.2)	17% (14-20%)	26% (21-30%)	220 (170-260)	10000 (8200-12000)
Denmark	1995	Male	50 to 54	99% (98-100%)	1% (0.48-1.9%)	7.7 (6.3-9.2)	17% (14-21%)	25% (20-29%)	290 (230-350)	12000 (9400-14000)
Denmark	1995	Male	55 to 59	99% (99-100%)	0.9% (0.44-1.5%)	7.6 (6.3-9.2)	16% (12-19%)	21% (16-26%)	310 (240-390)	11000 (8500-13000)
Denmark	1995	Male	60 to 64	99% (98-100%)	0.98% (0.43-1.7%)	7.4 (6-9)	14% (10-18%)	17% (12-22%)	370 (250-470)	11000 (7600-14000)
Denmark	1995	Male	65 to 69	99% (98-100%)	1.2% (0.46-2.5%)	6.9 (5.9-8.1)	12% (7.8-15%)	14% (8.8-18%)	430 (270-570)	10000 (6800-14000)
Denmark	1995	Male	70 to 74	99% (97-99%)	1.5% (0.68-2.8%)	6.5 (5.3-7.7)	9.9% (5.6-14%)	11% (6-16%)	490 (260-710)	9600 (5300-14000)
Denmark	1995	Male	75 to 79	98% (97-99%)	1.8% (0.77-3.3%)	6 (4.9-7.3)	8.4% (3.8-13%)	9.1% (3.6-14%)	450 (180-710)	6900 (3100-11000)
Denmark	1995	Male	80 to 84	98% (96-99%)	2% (0.95-3.7%)	5.6 (4.6-6.7)	7.2% (2.6-12%)	7.6% (2.4-13%)	370 (110-640)	4300 (1600-7100)
Denmark	1995	Male	85 to 89	98% (97-99%)	2.1% (1.1-3.5%)	5.2 (4.2-6.4)	5.7% (0.82-10%)	5.9% (0.24-11%)	200 (8.5-400)	1700 (250-3200)
Denmark	1995	Male	90 to 94	98% (96-99%)	2.1% (1-3.6%)	4.9 (3.6-6.3)	4.1% (-1.2-9.6%)	4% (-2.1-11%)	62 (-32-160)	380 (-110-900)
Denmark	1995	Male	95 plus	98% (96-99%)	2.3% (0.84-4.4%)	4.5 (2.9-6.3)	2.8% (-3-9.3%)	2.5% (-4.3-10%)	10 (-17-41)	46 (-50-150)
Denmark	2000	Female	All Ages	96% (94-96%)	4.4% (3.5-5.6%)	3.4 (3-3.9)	6% (4.3-7.9%)	7% (3.4-11%)	2100 (980-3300)	48000 (34000-62000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	2000	Female	15 to 19	94% (90-97%)	5.7% (2.8-10%)	3.2 (2.3-4.2)	3.1% (1.9-4.5%)	12% (6.7-18%)	4.2 (2.4-6.2)	440 (280-630)
Denmark	2000	Female	20 to 24	96% (92-98%)	4% (2-7.6%)	3.7 (2.6-5.1)	3.7% (2.6-5%)	13% (7.8-18%)	5.7 (3.5-7.9)	710 (500-960)
Denmark	2000	Female	25 to 29	97% (95-98%)	3.2% (1.7-5.2%)	3.2 (2.3-4.1)	3.1% (2.1-4.1%)	11% (6.2-15%)	7 (3.9-9.9)	770 (520-1000)
Denmark	2000	Female	30 to 34	97% (95-99%)	2.9% (1.4-4.8%)	3 (2.2-3.9)	3.1% (2.2-4.3%)	11% (7.5-15%)	11 (7-15)	910 (620-1200)
Denmark	2000	Female	35 to 39	97% (95-98%)	3.1% (1.6-5%)	3.1 (2.3-4.2)	4.4% (3.1-6%)	14% (10-18%)	24 (16-31)	1500 (1100-2000)
Denmark	2000	Female	40 to 44	97% (95-98%)	3.1% (1.7-4.9%)	3.7 (2.9-4.8)	6.3% (4.7-8.2%)	17% (13-20%)	45 (35-56)	2400 (1900-3100)
Denmark	2000	Female	45 to 49	97% (96-98%)	2.8% (1.9-4%)	4.7 (3.6-5.9)	8.5% (6.7-11%)	18% (15-22%)	84 (68-100)	3900 (3200-4800)
Denmark	2000	Female	50 to 54	97% (97-98%)	2.5% (1.7-3.4%)	5.3 (4.1-6.6)	9.4% (7.4-12%)	17% (14-21%)	130 (110-160)	5500 (4400-6700)
Denmark	2000	Female	55 to 59	98% (97-98%)	2.2% (1.5-3%)	5.5 (4.4-6.8)	9.7% (7.7-12%)	16% (13-19%)	170 (140-210)	6000 (4800-7400)
Denmark	2000	Female	60 to 64	98% (97-98%)	2.3% (1.5-3.3%)	5.5 (4.3-6.8)	8.8% (6.5-11%)	13% (9.7-16%)	180 (130-230)	5300 (3900-6800)
Denmark	2000	Female	65 to 69	97% (96-98%)	2.7% (1.8-4%)	5.1 (4.1-6.2)	7.5% (5.1-9.7%)	9.9% (6.8-13%)	200 (140-260)	4900 (3300-6600)
Denmark	2000	Female	70 to 74	97% (95-98%)	3.3% (2.3-4.8%)	4.7 (3.6-5.9)	6.4% (3.2-9.4%)	8.1% (4.1-12%)	230 (120-340)	4600 (2400-6800)
Denmark	2000	Female	75 to 79	96% (93-97%)	4.4% (2.6-7.4%)	4.3 (3.2-5.6)	5.4% (1.8-9%)	6.5% (1.9-11%)	270 (79-450)	4200 (1400-7000)
Denmark	2000	Female	80 to 84	93% (89-96%)	7% (4.5-11%)	4 (3.1-5)	4.7% (0.83-8.6%)	5.5% (0.6-10%)	270 (30-510)	3200 (570-5800)
Denmark	2000	Female	85 to 89	90% (86-93%)	10% (6.8-14%)	3.9 (2.9-5)	4.1% (0.037-8.6%)	4.7% (-0.42-10%)	260 (-24-580)	2200 (18-4600)
Denmark	2000	Female	90 to 94	87% (81-92%)	13% (8.3-19%)	3.7 (2.6-5.1)	3.3% (-0.99-8.7%)	3.6% (-1.8-10%)	140 (-71-410)	860 (-250-2200)
Denmark	2000	Female	95 plus	84% (76-91%)	16% (9.2-24%)	3.5 (2.2-5.3)	2.9% (-2.1-9.2%)	3.1% (-2.9-11%)	49 (-46-170)	200 (-140-620)
Denmark	2000	Male	All Ages	97% (97-98%)	2.6% (1.7-3.5%)	5 (4.3-5.6)	13% (10-15%)	13% (9.8-17%)	3800 (2700-4800)	100000 (86000-120000)
Denmark	2000	Male	15 to 19	95% (92-98%)	4.7% (2.4-8.5%)	4.3 (3.2-5.6)	8.6% (5.6-12%)	19% (11-26%)	16 (9.5-22)	1400 (930-1900)
Denmark	2000	Male	20 to 24	97% (94-98%)	3.4% (1.8-6%)	6.1 (4.6-7.9)	14% (11-19%)	26% (18-34%)	35 (24-47)	3200 (2300-4000)
Denmark	2000	Male	25 to 29	98% (96-99%)	2.4% (1.3-4.1%)	5.2 (4.1-6.5)	12% (8.9-15%)	22% (15-29%)	35 (24-47)	3100 (2300-4000)
Denmark	2000	Male	30 to 34	98% (97-99%)	1.8% (0.8-3.3%)	4.6 (3.5-5.8)	10% (7.7-13%)	20% (14-27%)	42 (28-57)	3300 (2400-4200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	2000	Male	35 to 39	99% (97-99%)	1.5% (0.65-2.6%)	4.3 (3.2-5.5)	11% (8.6-14%)	21% (15-26%)	63 (45-80)	4200 (3200-5200)
Denmark	2000	Male	40 to 44	99% (98-99%)	1.2% (0.53-2.3%)	5.1 (4-6.4)	15% (12-18%)	25% (20-30%)	120 (91-140)	6400 (5200-7700)
Denmark	2000	Male	45 to 49	99% (98-99%)	1.1% (0.55-2.1%)	7.1 (5.8-8.7)	19% (16-22%)	30% (26-35%)	220 (180-260)	10000 (8500-12000)
Denmark	2000	Male	50 to 54	99% (98-100%)	0.97% (0.46-1.8%)	8.1 (6.6-9.8)	20% (17-23%)	29% (25-33%)	330 (280-390)	14000 (11000-16000)
Denmark	2000	Male	55 to 59	99% (99-100%)	0.85% (0.41-1.5%)	8 (6.6-9.5)	18% (15-21%)	25% (21-28%)	410 (340-490)	14000 (12000-17000)
Denmark	2000	Male	60 to 64	99% (98-100%)	0.91% (0.4-1.6%)	7.8 (6.3-9.3)	16% (12-19%)	20% (16-23%)	380 (300-460)	11000 (8900-14000)
Denmark	2000	Male	65 to 69	99% (98-100%)	1.2% (0.44-2.3%)	7.3 (6.2-8.6)	13% (10-16%)	16% (12-20%)	420 (310-520)	10000 (7600-13000)
Denmark	2000	Male	70 to 74	99% (97-99%)	1.4% (0.63-2.6%)	6.9 (5.6-8.3)	11% (7.4-15%)	13% (8.5-17%)	450 (300-610)	9000 (6100-12000)
Denmark	2000	Male	75 to 79	98% (97-99%)	1.6% (0.72-3%)	6.4 (5.2-7.8)	9.6% (5.6-14%)	11% (5.9-15%)	490 (280-710)	7600 (4500-11000)
Denmark	2000	Male	80 to 84	98% (96-99%)	1.9% (0.9-3.6%)	5.9 (4.9-7.1)	8.3% (4-12%)	9% (3.9-14%)	400 (170-620)	4600 (2200-6900)
Denmark	2000	Male	85 to 89	98% (96-99%)	2.1% (1.1-3.6%)	5.5 (4.4-6.7)	6.8% (2.3-11%)	7.2% (2.1-12%)	250 (73-440)	2100 (720-3500)
Denmark	2000	Male	90 to 94	98% (96-99%)	2.2% (1.1-4%)	5.1 (3.8-6.6)	5.2% (0.19-11%)	5.5% (-0.33-12%)	91 (-5.2-200)	530 (19-1100)
Denmark	2000	Male	95 plus	98% (95-99%)	2.4% (0.9-4.6%)	4.7 (3-6.6)	4% (-1.6-11%)	3.9% (-2.6-12%)	17 (-11-50)	69 (-28-180)
Denmark	2005	Female	All Ages	96% (95-97%)	4.3% (3.3-5.4%)	3.4 (3-3.9)	6.2% (4.6-7.9%)	7.6% (4.2-11%)	2200 (1200-3300)	48000 (35000-62000)
Denmark	2005	Female	15 to 19	94% (90-97%)	5.6% (2.7-9.7%)	3.2 (2.3-4.2)	2.7% (1.7-3.9%)	12% (6.4-17%)	3.6 (2-5.2)	410 (260-580)
Denmark	2005	Female	20 to 24	96% (93-98%)	3.9% (2-6.9%)	3.7 (2.6-5.1)	3.3% (2.4-4.5%)	13% (7.8-18%)	4 (2.4-5.6)	560 (390-780)
Denmark	2005	Female	25 to 29	97% (95-98%)	3.1% (1.6-5.2%)	3.2 (2.4-4.1)	2.9% (2-3.8%)	11% (6.4-15%)	5.6 (3.2-7.7)	650 (440-870)
Denmark	2005	Female	30 to 34	97% (95-99%)	2.8% (1.5-4.8%)	3 (2.2-4)	2.8% (2-3.8%)	12% (8-15%)	8.7 (5.9-12)	780 (550-1000)
Denmark	2005	Female	35 to 39	97% (95-98%)	3.1% (1.6-5%)	3.2 (2.3-4.3)	3.9% (2.7-5.2%)	14% (10-18%)	19 (14-25)	1300 (950-1700)
Denmark	2005	Female	40 to 44	97% (95-98%)	3% (1.6-4.7%)	3.7 (2.8-4.9)	5.7% (4.3-7.3%)	17% (14-20%)	41 (33-51)	2300 (1800-2800)
Denmark	2005	Female	45 to 49	97% (96-98%)	2.7% (1.8-3.9%)	4.7 (3.6-6.1)	8.4% (6.6-11%)	19% (16-23%)	80 (65-95)	3700 (3000-4500)
Denmark	2005	Female	50 to 54	98% (97-98%)	2.4% (1.7-3.3%)	5.3 (4.1-6.7)	9.7% (7.8-12%)	18% (16-21%)	120 (100-150)	5000 (4100-6100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	2005	Female	55 to 59	98% (97-98%)	2.2% (1.5-2.9%)	5.5 (4.3-6.8)	9.9% (7.8-12%)	17% (14-20%)	180 (150-210)	6300 (5100-7700)
Denmark	2005	Female	60 to 64	98% (97-98%)	2.2% (1.5-3.2%)	5.5 (4.2-6.9)	9% (6.9-11%)	14% (11-17%)	200 (150-250)	6000 (4600-7600)
Denmark	2005	Female	65 to 69	97% (96-98%)	2.6% (1.7-3.9%)	5.2 (4.2-6.2)	7.9% (5.8-10%)	11% (8.3-14%)	200 (150-250)	5000 (3600-6400)
Denmark	2005	Female	70 to 74	97% (96-98%)	3.2% (2.3-4.4%)	4.7 (3.6-6)	6.7% (3.9-9.6%)	8.8% (5.3-12%)	220 (130-310)	4400 (2600-6300)
Denmark	2005	Female	75 to 79	96% (93-98%)	4.1% (2.5-6.7%)	4.3 (3.2-5.6)	5.8% (2.4-9.1%)	7.1% (3-11%)	250 (100-400)	4000 (1700-6300)
Denmark	2005	Female	80 to 84	94% (90-96%)	6.5% (4.4-9.6%)	4.1 (3.1-5.2)	5.2% (1.8-8.7%)	6.3% (2-11%)	310 (97-520)	3600 (1200-5800)
Denmark	2005	Female	85 to 89	90% (86-93%)	9.7% (6.7-14%)	3.9 (2.9-5)	4.7% (1-8.7%)	5.4% (0.85-11%)	280 (44-560)	2300 (520-4400)
Denmark	2005	Female	90 to 94	88% (83-92%)	12% (8.4-17%)	3.7 (2.6-5.1)	3.8% (-0.35-8.5%)	4.3% (-0.77-10%)	180 (-32-420)	1000 (-91-2300)
Denmark	2005	Female	95 plus	85% (78-91%)	15% (9-22%)	3.5 (2.1-5.2)	3.3% (-1.8-9.8%)	3.5% (-2.5-11%)	68 (-48-220)	260 (-140-770)
Denmark	2005	Male	All Ages	98% (97-98%)	2.5% (1.7-3.4%)	5 (4.3-5.7)	13% (11-15%)	14% (11-18%)	3900 (3000-4900)	110000 (89000-130000)
Denmark	2005	Male	15 to 19	96% (92-98%)	4.5% (2.3-7.8%)	4.3 (3.2-5.6)	7.8% (5.2-11%)	18% (11-25%)	15 (8.7-21)	1300 (900-1800)
Denmark	2005	Male	20 to 24	97% (94-98%)	3.2% (1.7-5.6%)	6.1 (4.7-7.8)	13% (10-17%)	26% (18-34%)	26 (18-35)	2500 (1800-3100)
Denmark	2005	Male	25 to 29	98% (96-99%)	2.3% (1.2-3.9%)	5.3 (4.1-6.6)	11% (8.4-14%)	22% (15-30%)	28 (19-38)	2600 (1900-3300)
Denmark	2005	Male	30 to 34	98% (97-99%)	1.7% (0.78-3.2%)	4.6 (3.6-5.9)	9.8% (7.4-13%)	20% (14-27%)	35 (24-47)	2800 (2100-3600)
Denmark	2005	Male	35 to 39	99% (97-99%)	1.4% (0.65-2.6%)	4.3 (3.2-5.6)	11% (8.1-13%)	21% (15-26%)	54 (39-68)	3700 (2900-4600)
Denmark	2005	Male	40 to 44	99% (98-99%)	1.2% (0.55-2.3%)	5.1 (4-6.4)	14% (11-17%)	26% (21-31%)	110 (87-140)	6200 (5000-7500)
Denmark	2005	Male	45 to 49	99% (98-99%)	1.1% (0.54-2.1%)	7.1 (5.7-8.9)	20% (16-23%)	32% (27-36%)	210 (180-250)	10000 (8400-12000)
Denmark	2005	Male	50 to 54	99% (98-100%)	0.96% (0.45-1.8%)	8.1 (6.5-9.9)	21% (18-25%)	31% (28-35%)	330 (280-390)	13000 (12000-16000)
Denmark	2005	Male	55 to 59	99% (99-100%)	0.84% (0.41-1.4%)	8 (6.6-9.6)	19% (16-22%)	27% (23-30%)	440 (370-510)	15000 (13000-18000)
Denmark	2005	Male	60 to 64	99% (98-100%)	0.89% (0.39-1.6%)	7.8 (6.3-9.4)	17% (13-19%)	21% (17-25%)	470 (380-560)	14000 (11000-17000)
Denmark	2005	Male	65 to 69	99% (98-100%)	1.1% (0.44-2.2%)	7.3 (6.1-8.7)	14% (11-17%)	17% (13-20%)	440 (340-530)	11000 (8400-13000)
Denmark	2005	Male	70 to 74	99% (98-99%)	1.4% (0.64-2.5%)	6.8 (5.6-8.3)	11% (8-15%)	13% (9.4-17%)	430 (300-550)	8500 (6100-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	2005	Male	75 to 79	98% (97-99%)	1.6% (0.72-3%)	6.4 (5.1-7.8)	9.8% (6.4-13%)	11% (7-15%)	450 (280-610)	7000 (4600-9400)
Denmark	2005	Male	80 to 84	98% (97-99%)	1.8% (0.88-3.3%)	6 (4.8-7.2)	8.7% (4.8-12%)	9.5% (5.1-14%)	430 (230-620)	4900 (2700-7000)
Denmark	2005	Male	85 to 89	98% (97-99%)	1.9% (1-3.2%)	5.6 (4.4-6.8)	7.4% (3.3-12%)	8.1% (3.3-13%)	280 (110-450)	2300 (1000-3600)
Denmark	2005	Male	90 to 94	98% (96-99%)	2% (0.99-3.6%)	5.2 (3.8-6.7)	6% (1.3-11%)	6.4% (0.86-12%)	120 (16-230)	680 (140-1300)
Denmark	2005	Male	95 plus	98% (96-99%)	2.2% (0.8-4.2%)	4.8 (3.1-6.6)	4.8% (-0.75-11%)	4.9% (-1.5-12%)	26 (-8-64)	100 (-16-230)
Denmark	2010	Female	All Ages	96% (95-97%)	4.3% (3.4-5.5%)	3.3 (2.8-3.9)	5.7% (4.2-7.4%)	7.2% (4.1-11%)	1900 (1100-2800)	43000 (32000-55000)
Denmark	2010	Female	15 to 19	94% (90-97%)	5.7% (2.7-10%)	3 (2.1-4)	2.3% (1.5-3.3%)	11% (5.8-16%)	3.1 (1.7-4.6)	390 (250-560)
Denmark	2010	Female	20 to 24	96% (93-98%)	3.8% (1.9-7.2%)	3.6 (2.5-4.9)	3.2% (2.2-4.2%)	12% (7-16%)	4.3 (2.6-6)	610 (420-830)
Denmark	2010	Female	25 to 29	97% (95-98%)	3.1% (1.6-5.1%)	3 (2.2-4)	2.5% (1.7-3.3%)	10% (5.9-14%)	3.5 (2.1-5)	490 (340-670)
Denmark	2010	Female	30 to 34	97% (95-98%)	3% (1.5-5%)	2.8 (2-3.8)	2.6% (1.8-3.6%)	11% (7.7-15%)	7.1 (4.7-9.6)	660 (460-880)
Denmark	2010	Female	35 to 39	97% (95-98%)	3.2% (1.6-5.1%)	3 (2.1-4.1)	3.1% (2.2-4.2%)	13% (9.5-17%)	13 (9.3-17)	960 (680-1300)
Denmark	2010	Female	40 to 44	97% (95-98%)	3% (1.7-4.7%)	3.5 (2.6-4.6)	4.8% (3.6-6.3%)	16% (12-19%)	32 (25-41)	1900 (1400-2400)
Denmark	2010	Female	45 to 49	97% (96-98%)	2.8% (1.9-3.9%)	4.5 (3.3-5.8)	7.3% (5.8-9.4%)	19% (16-22%)	70 (57-85)	3400 (2700-4200)
Denmark	2010	Female	50 to 54	98% (97-98%)	2.4% (1.7-3.3%)	5 (3.8-6.5)	8.9% (7.1-11%)	18% (15-22%)	110 (87-130)	4500 (3600-5500)
Denmark	2010	Female	55 to 59	98% (97-99%)	2.2% (1.5-2.9%)	5.2 (4-6.6)	9.3% (7.3-11%)	17% (14-19%)	150 (120-180)	5300 (4200-6400)
Denmark	2010	Female	60 to 64	98% (97-98%)	2.3% (1.5-3.2%)	5.2 (3.9-6.6)	8.5% (6.3-11%)	14% (11-17%)	190 (150-240)	5900 (4500-7500)
Denmark	2010	Female	65 to 69	97% (96-98%)	2.6% (1.8-3.8%)	4.9 (3.9-6.1)	7.4% (5.5-9.4%)	11% (8.5-14%)	210 (160-270)	5300 (3900-6800)
Denmark	2010	Female	70 to 74	97% (96-98%)	3.1% (2.2-4.4%)	4.5 (3.4-5.8)	6.2% (3.7-8.8%)	8.5% (5.5-12%)	200 (130-280)	4000 (2400-5700)
Denmark	2010	Female	75 to 79	96% (94-98%)	4% (2.4-6.5%)	4.1 (3-5.4)	5.2% (2.3-8.4%)	6.6% (3.1-10%)	210 (97-340)	3400 (1500-5300)
Denmark	2010	Female	80 to 84	94% (90-96%)	6.4% (4.2-9.7%)	3.9 (2.9-5)	4.7% (1.8-8%)	5.7% (2.1-9.7%)	250 (89-420)	2900 (1100-4800)
Denmark	2010	Female	85 to 89	90% (86-93%)	9.8% (6.8-14%)	3.7 (2.8-4.8)	4.1% (0.95-7.7%)	4.9% (0.98-9.4%)	250 (49-480)	2100 (470-3900)
Denmark	2010	Female	90 to 94	88% (82-92%)	12% (8.1-18%)	3.5 (2.4-4.9)	3.5% (-0.27-7.8%)	4% (-0.51-9.3%)	160 (-20-370)	920 (-70-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	2010	Female	95 plus	85% (77-91%)	15% (9.1-23%)	3.3 (2-5.1)	3% (-1.6-8.9%)	3.3% (-2.2-10%)	71 (-48-220)	270 (-140-780)
Denmark	2010	Male	All Ages	98% (97-98%)	2.5% (1.7-3.3%)	4.7 (4-5.6)	12% (10-15%)	13% (10-17%)	3500 (2600-4500)	96000 (79000-120000)
Denmark	2010	Male	15 to 19	96% (92-98%)	4.4% (2.3-7.6%)	4.1 (3-5.3)	6.3% (4.2-8.5%)	17% (10-24%)	10 (6.1-15)	1100 (730-1500)
Denmark	2010	Male	20 to 24	97% (94-98%)	3.2% (1.7-5.6%)	5.7 (4.2-7.3)	12% (8.7-15%)	23% (15-31%)	21 (14-29)	2200 (1700-2900)
Denmark	2010	Male	25 to 29	98% (96-99%)	2.3% (1.2-4%)	4.9 (3.8-6.2)	9.9% (7.4-13%)	20% (13-28%)	19 (13-27)	2000 (1400-2500)
Denmark	2010	Male	30 to 34	98% (97-99%)	1.8% (0.82-3.4%)	4.3 (3.2-5.6)	8.5% (6.2-11%)	18% (12-25%)	23 (15-31)	2100 (1500-2700)
Denmark	2010	Male	35 to 39	99% (97-99%)	1.5% (0.68-2.7%)	4 (2.9-5.2)	9.3% (7.1-12%)	19% (14-25%)	40 (28-52)	3000 (2300-3700)
Denmark	2010	Male	40 to 44	99% (98-99%)	1.2% (0.52-2.3%)	4.8 (3.7-6.2)	12% (9.7-15%)	24% (19-29%)	85 (67-110)	5000 (4000-6100)
Denmark	2010	Male	45 to 49	99% (98-99%)	1.1% (0.54-2%)	6.7 (5.2-8.5)	17% (14-20%)	30% (26-35%)	180 (150-210)	8700 (7300-10000)
Denmark	2010	Male	50 to 54	99% (98-100%)	0.95% (0.44-1.8%)	7.6 (5.9-9.6)	20% (17-23%)	31% (26-35%)	290 (240-340)	12000 (10000-14000)
Denmark	2010	Male	55 to 59	99% (99-100%)	0.84% (0.41-1.4%)	7.5 (6.1-9.4)	19% (16-22%)	27% (23-31%)	380 (320-440)	13000 (11000-15000)
Denmark	2010	Male	60 to 64	99% (98-100%)	0.89% (0.39-1.6%)	7.3 (5.7-9.2)	16% (13-19%)	22% (18-25%)	470 (380-560)	14000 (11000-17000)
Denmark	2010	Male	65 to 69	99% (98-100%)	1.1% (0.43-2.2%)	6.9 (5.6-8.4)	13% (11-16%)	17% (14-20%)	460 (360-560)	12000 (9200-14000)
Denmark	2010	Male	70 to 74	99% (97-99%)	1.4% (0.63-2.6%)	6.4 (5.1-7.9)	11% (7.8-14%)	13% (9.4-17%)	410 (290-530)	8200 (5900-11000)
Denmark	2010	Male	75 to 79	98% (97-99%)	1.6% (0.72-3%)	5.9 (4.7-7.4)	8.9% (5.6-12%)	10% (6.5-14%)	370 (230-520)	5900 (3700-8200)
Denmark	2010	Male	80 to 84	98% (97-99%)	1.8% (0.85-3.3%)	5.6 (4.4-6.8)	7.6% (4.2-11%)	8.5% (4.5-13%)	350 (180-530)	4100 (2200-6100)
Denmark	2010	Male	85 to 89	98% (97-99%)	1.9% (1-3%)	5.2 (4.1-6.6)	6.5% (2.8-10%)	7% (2.8-12%)	260 (100-430)	2100 (940-3500)
Denmark	2010	Male	90 to 94	98% (97-99%)	2% (0.99-3.5%)	4.9 (3.5-6.4)	5.4% (1.1-10%)	5.8% (0.85-12%)	110 (16-220)	630 (130-1200)
Denmark	2010	Male	95 plus	98% (96-99%)	2.2% (0.84-4.2%)	4.5 (2.8-6.4)	4.4% (-0.6-10%)	4.6% (-1.3-12%)	29 (-15-74)	110 (-15-270)
Denmark	2016	Female	All Ages	95% (94-96%)	4.7% (3.7-6%)	3.2 (2.3-4.1)	5% (3.2-7%)	6.3% (3-10%)	1700 (750-2800)	37000 (24000-53000)
Denmark	2016	Female	15 to 19	94% (88-97%)	6.2% (3-12%)	2.8 (1.8-3.9)	2% (1.2-2.8%)	9.9% (5.1-15%)	2.2 (1.1-3.4)	320 (200-480)
Denmark	2016	Female	20 to 24	96% (92-98%)	4.5% (2.2-8.4%)	3.3 (2.1-4.8)	2.9% (2-3.9%)	11% (6.3-16%)	3.8 (2.1-5.7)	630 (420-880)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	2016	Female	25 to 29	96% (94-98%)	3.5% (1.8-6%)	2.8 (1.9-4)	2.3% (1.6-3.1%)	9.3% (5.2-13%)	3.2 (1.7-4.9)	500 (330-690)
Denmark	2016	Female	30 to 34	97% (94-98%)	3.2% (1.6-5.7%)	2.7 (1.8-3.8)	2.4% (1.6-3.3%)	11% (7.2-15%)	5.4 (3.4-7.7)	530 (360-720)
Denmark	2016	Female	35 to 39	96% (94-98%)	3.5% (1.7-5.8%)	2.8 (1.9-4)	2.8% (1.8-3.9%)	13% (8.8-17%)	10 (6.7-14)	770 (520-1000)
Denmark	2016	Female	40 to 44	97% (94-98%)	3.5% (1.8-5.8%)	3.3 (2.3-4.6)	4.1% (2.8-5.7%)	15% (12-19%)	25 (18-33)	1500 (1000-2000)
Denmark	2016	Female	45 to 49	97% (95-98%)	3.1% (2-4.7%)	4.2 (2.9-6)	6.3% (4.6-8.4%)	18% (14-22%)	56 (42-73)	2800 (2000-3600)
Denmark	2016	Female	50 to 54	97% (96-98%)	2.7% (1.8-3.9%)	4.7 (3.3-6.8)	7.8% (5.7-10%)	17% (14-21%)	94 (71-120)	4000 (3000-5200)
Denmark	2016	Female	55 to 59	98% (97-98%)	2.4% (1.6-3.3%)	4.9 (3.4-6.7)	8.2% (6-11%)	16% (12-19%)	120 (92-160)	4400 (3200-5700)
Denmark	2016	Female	60 to 64	97% (96-98%)	2.5% (1.7-3.8%)	4.9 (3.3-6.7)	7.6% (5.3-10%)	13% (9.9-16%)	150 (110-200)	4600 (3200-6100)
Denmark	2016	Female	65 to 69	97% (95-98%)	3.1% (1.9-4.7%)	4.5 (3.3-6.2)	6.5% (4.4-9%)	10% (7.6-14%)	200 (140-270)	5200 (3400-7200)
Denmark	2016	Female	70 to 74	96% (94-98%)	3.7% (2.5-5.5%)	4.1 (2.8-5.8)	5.3% (2.9-8.3%)	7.8% (4.7-12%)	200 (120-310)	4100 (2200-6400)
Denmark	2016	Female	75 to 79	95% (92-97%)	4.7% (2.6-8.1%)	3.8 (2.5-5.5)	4.5% (1.7-8%)	5.9% (2.5-10%)	190 (78-340)	3000 (1100-5400)
Denmark	2016	Female	80 to 84	93% (89-96%)	7% (4.4-11%)	3.6 (2.4-5.1)	4% (1-7.6%)	5% (1.3-9.6%)	190 (51-370)	2300 (570-4300)
Denmark	2016	Female	85 to 89	89% (84-93%)	11% (6.9-16%)	3.4 (2.3-4.8)	3.5% (0.27-7.4%)	4.1% (0.2-9%)	190 (9-420)	1600 (120-3400)
Denmark	2016	Female	90 to 94	87% (80-91%)	13% (8.5-20%)	3.3 (2.1-4.8)	2.9% (-0.77-7.7%)	3.3% (-1.3-8.9%)	150 (-54-400)	850 (-230-2200)
Denmark	2016	Female	95 plus	84% (76-91%)	16% (9.3-24%)	3.1 (1.8-4.9)	2.5% (-1.8-8.5%)	2.8% (-2.5-10%)	64 (-57-230)	240 (-170-790)
Denmark	2016	Male	All Ages	97% (96-98%)	2.9% (2-4%)	4.5 (3.4-5.8)	11% (8.3-14%)	12% (8-16%)	3200 (2100-4500)	87000 (64000-110000)
Denmark	2016	Male	15 to 19	94% (89-97%)	5.7% (2.6-11%)	3.8 (2.5-5.3)	5.4% (3.5-7.5%)	16% (8.7-23%)	7.8 (4.1-12)	900 (600-1300)
Denmark	2016	Male	20 to 24	96% (93-98%)	4.1% (2-7.5%)	5.4 (3.6-7.3)	10% (7.5-13%)	21% (13-31%)	18 (11-27)	2200 (1600-2900)
Denmark	2016	Male	25 to 29	97% (95-99%)	2.9% (1.4-5.4%)	4.6 (3.3-6.3)	8.8% (6.4-11%)	19% (11-27%)	17 (10-25)	1900 (1400-2600)
Denmark	2016	Male	30 to 34	98% (96-99%)	2.2% (0.99-4.3%)	4 (2.8-5.7)	7.9% (5.6-10%)	18% (11-25%)	18 (11-26)	1700 (1200-2300)
Denmark	2016	Male	35 to 39	98% (97-99%)	1.8% (0.8-3.5%)	3.8 (2.5-5.4)	8.5% (6.2-11%)	19% (12-25%)	30 (20-41)	2300 (1700-3000)
Denmark	2016	Male	40 to 44	98% (97-99%)	1.6% (0.61-3.2%)	4.5 (3.2-6.1)	11% (8.5-14%)	23% (18-29%)	65 (48-85)	4000 (3000-5000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Denmark	2016	Male	45 to 49	99% (97-99%)	1.4% (0.59-2.8%)	6.2 (4.4-8.5)	16% (12-19%)	29% (24-35%)	150 (120-190)	7500 (5900-9500)
Denmark	2016	Male	50 to 54	99% (98-99%)	1.2% (0.54-2.4%)	7.1 (5-9.6)	18% (15-22%)	29% (24-34%)	260 (200-320)	11000 (8500-13000)
Denmark	2016	Male	55 to 59	99% (98-100%)	1.1% (0.49-1.8%)	7 (5.1-9.5)	17% (14-21%)	26% (21-30%)	320 (250-400)	11000 (8800-14000)
Denmark	2016	Male	60 to 64	99% (98-100%)	1.1% (0.47-2.2%)	6.8 (4.9-9.1)	15% (11-18%)	21% (16-25%)	360 (270-460)	11000 (8200-14000)
Denmark	2016	Male	65 to 69	99% (97-99%)	1.4% (0.52-2.9%)	6.4 (4.7-8.4)	12% (8.8-15%)	16% (12-20%)	470 (340-600)	12000 (8500-15000)
Denmark	2016	Male	70 to 74	98% (97-99%)	1.7% (0.71-3.3%)	6 (4.2-8)	9.7% (6.2-13%)	12% (7.7-16%)	440 (270-610)	9000 (5600-12000)
Denmark	2016	Male	75 to 79	98% (96-99%)	1.9% (0.76-3.6%)	5.5 (3.9-7.6)	8% (4.3-12%)	9.3% (4.9-14%)	370 (190-560)	5900 (3100-9000)
Denmark	2016	Male	80 to 84	98% (96-99%)	2.2% (0.98-4%)	5.1 (3.6-6.9)	6.7% (2.9-11%)	7.5% (3.1-12%)	300 (130-510)	3600 (1500-5900)
Denmark	2016	Male	85 to 89	98% (96-99%)	2.4% (1.1-4.3%)	4.8 (3.4-6.4)	5.6% (1.7-11%)	6.1% (1.5-12%)	230 (59-450)	1900 (590-3600)
Denmark	2016	Male	90 to 94	98% (96-99%)	2.4% (1.1-4.3%)	4.5 (3-6.3)	4.5% (0.35-10%)	4.9% (-0.057-11%)	110 (-1.3-260)	640 (50-1400)
Denmark	2016	Male	95 plus	97% (95-99%)	2.5% (0.95-4.9%)	4.1 (2.5-6.3)	3.7% (-1.1-10%)	3.8% (-1.8-12%)	27 (-13-81)	100 (-31-290)
Finland	1990	Female	All Ages	74% (66-81%)	26% (19-34%)	2.2 (1.8-2.5)	2% (0.28-3.7%)	0.43% (-2.8-3.8%)	110 (-680-940)	14000 (2100-27000)
Finland	1990	Female	15 to 19	71% (51-86%)	29% (14-49%)	2.6 (1.7-3.6)	4.1% (2.6-5.8%)	13% (6.7-19%)	7.2 (3.8-11)	710 (450-1000)
Finland	1990	Female	20 to 24	76% (58-89%)	24% (11-42%)	3.3 (2.2-4.7)	5.7% (3.9-7.6%)	16% (8.8-23%)	12 (6.6-17)	1300 (920-1800)
Finland	1990	Female	25 to 29	78% (62-90%)	22% (10-38%)	2.7 (2-3.5)	4.1% (2.8-5.5%)	12% (6.6-18%)	9.5 (5.1-14)	1100 (750-1500)
Finland	1990	Female	30 to 34	80% (63-91%)	20% (9.2-37%)	2.4 (1.7-3.2)	3.7% (2.4-5.1%)	11% (6.3-16%)	13 (7.3-19)	1100 (730-1500)
Finland	1990	Female	35 to 39	81% (66-92%)	19% (8.1-34%)	2.5 (1.8-3.4)	4.3% (2.8-5.9%)	13% (8-17%)	23 (15-32)	1600 (1000-2100)
Finland	1990	Female	40 to 44	82% (65-92%)	18% (7.9-35%)	2.9 (2.1-3.8)	5.5% (3.8-7.5%)	14% (10-19%)	46 (33-60)	2500 (1800-3300)
Finland	1990	Female	45 to 49	82% (63-92%)	18% (8.4-37%)	3.5 (2.5-4.5)	6.8% (5-9.1%)	16% (11-20%)	54 (40-69)	2600 (1900-3300)
Finland	1990	Female	50 to 54	81% (64-91%)	19% (8.5-36%)	3.4 (2.5-4.5)	6.7% (4.5-9.1%)	13% (9.6-17%)	62 (44-80)	2600 (1800-3400)
Finland	1990	Female	55 to 59	80% (63-92%)	20% (8.4-37%)	2.8 (2-3.8)	4.6% (2.7-6.9%)	8.6% (5.4-12%)	53 (33-75)	1900 (1100-2700)
Finland	1990	Female	60 to 64	79% (59-92%)	21% (8.4-41%)	2.4 (1.6-3.3)	2.5% (0.19-5.3%)	4.1% (0.54-8.1%)	45 (6-88)	1300 (97-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	1990	Female	65 to 69	77% (57-91%)	23% (9.5-43%)	2.1 (1.4-2.9)	0.67% (-2.2-3.4%)	0.97% (-3-4.7%)	18 (-57-88)	440 (-1400-2200)
Finland	1990	Female	70 to 74	74% (55-87%)	26% (13-45%)	1.8 (1.2-2.6)	-0.72% (-3.7-2.4%)	-0.99% (-5-3.3%)	-27 (-140-87)	-500 (-2600-1600)
Finland	1990	Female	75 to 79	71% (51-86%)	29% (14-49%)	1.7 (1.1-2.5)	-1.3% (-4.4-2%)	-1.8% (-5.7-2.4%)	-78 (-250-110)	-1100 (-3600-1600)
Finland	1990	Female	80 to 84	68% (47-84%)	32% (16-53%)	1.8 (1.1-2.6)	-1.1% (-4.4-2.1%)	-1.5% (-5.4-2.4%)	-85 (-300-130)	-830 (-3200-1500)
Finland	1990	Female	85 to 89	65% (46-81%)	35% (19-54%)	1.8 (1.1-2.7)	-0.57% (-3.6-2.8%)	-0.88% (-4.5-3.2%)	-38 (-200-140)	-230 (-1400-1100)
Finland	1990	Female	90 to 94	64% (45-82%)	36% (18-55%)	1.9 (1.1-3)	-0.12% (-3.3-3.7%)	-0.4% (-4.1-4.2%)	-8.2 (-85-85)	-16 (-420-470)
Finland	1990	Female	95 plus	65% (41-84%)	35% (16-59%)	2 (1-3.4)	0.38% (-3.3-4.9%)	0.15% (-4.1-5.5%)	0.83 (-23-31)	8.7 (-76-110)
Finland	1990	Male	All Ages	80% (71-87%)	20% (13-29%)	4.6 (4-5.2)	12% (9.9-15%)	11% (7-15%)	2700 (1700-3700)	110000 (84000-130000)
Finland	1990	Male	15 to 19	72% (51-88%)	28% (12-49%)	3.3 (2.2-4.7)	10% (6.4-15%)	18% (10-27%)	28 (16-42)	2400 (1500-3400)
Finland	1990	Male	20 to 24	78% (59-90%)	22% (10-41%)	5.3 (3.6-7.1)	28% (14-25%)	28% (19-38%)	74 (49-100)	6100 (4400-8000)
Finland	1990	Male	25 to 29	82% (66-92%)	18% (8.2-34%)	5.2 (3.7-6.9)	18% (13-23%)	28% (19-37%)	80 (54-110)	6400 (4800-8100)
Finland	1990	Male	30 to 34	85% (68-94%)	15% (6-32%)	5.4 (4-7.1)	18% (14-23%)	28% (20-36%)	110 (80-140)	7600 (5900-9600)
Finland	1990	Male	35 to 39	87% (74-95%)	13% (5.2-26%)	6 (4.3-8)	20% (15-24%)	29% (22-37%)	170 (130-210)	10000 (8000-13000)
Finland	1990	Male	40 to 44	88% (75-96%)	12% (4.2-25%)	6.6 (5-8.6)	20% (16-25%)	30% (23-35%)	270 (210-320)	14000 (11000-17000)
Finland	1990	Male	45 to 49	89% (71-96%)	11% (4-29%)	7.4 (5.4-9.5)	21% (16-25%)	28% (22-34%)	250 (200-310)	12000 (9200-14000)
Finland	1990	Male	50 to 54	89% (75-96%)	11% (3.8-25%)	7.5 (5.5-9.9)	19% (14-23%)	25% (18-31%)	280 (210-360)	11000 (8700-14000)
Finland	1990	Male	55 to 59	89% (77-96%)	11% (3.7-23%)	7.2 (5.5-9.3)	16% (12-21%)	20% (14-26%)	330 (230-420)	11000 (8000-14000)
Finland	1990	Male	60 to 64	88% (73-96%)	12% (3.9-27%)	6.4 (4.6-8.6)	13% (7.6-18%)	15% (8.4-21%)	370 (210-520)	11000 (6400-15000)
Finland	1990	Male	65 to 69	88% (73-96%)	12% (4.2-27%)	5.3 (3.9-6.9)	8.8% (3.6-14%)	9.7% (3.5-16%)	290 (100-480)	7300 (3000-12000)
Finland	1990	Male	70 to 74	87% (73-95%)	13% (5.4-27%)	4.1 (2.9-5.4)	4.5% (-0.24-10%)	4.6% (-0.83-11%)	140 (-26-350)	3000 (-170-6900)
Finland	1990	Male	75 to 79	85% (71-94%)	15% (5.8-29%)	3.4 (2.3-5)	2.6% (-1.5-8.2%)	2.5% (-2.3-9%)	96 (-86-340)	1700 (-940-5300)
Finland	1990	Male	80 to 84	84% (67-93%)	16% (7.2-33%)	3.4 (2.3-4.7)	2.7% (-1.7-5.5%)	2.7% (-1.7-8.2%)	88 (-56-270)	1100 (-410-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	1990	Male	85 to 89	82% (66-92%)	18% (7.7-34%)	3.3 (2.2-4.7)	3.1% (-0.64-7.8%)	3.1% (-1.1-8.4%)	57 (-21-160)	500 (-110-1300)
Finland	1990	Male	90 to 94	82% (67-93%)	18% (7-33%)	3.4 (2.1-5)	3.2% (-1-8.9%)	3.3% (-1.6-9.9%)	22 (-11-66)	130 (-41-370)
Finland	1990	Male	95 plus	83% (64-94%)	17% (5.7-36%)	3.5 (1.9-5.6)	3.8% (-1.5-11%)	3.9% (-2.1-12%)	5.1 (-2.7-16)	20 (-7.7-56)
Finland	1995	Female	All Ages	75% (67-82%)	25% (18-33%)	2.2 (1.9-2.5)	2.3% (0.72-3.9%)	0.66% (-2.2-3.8%)	170 (-550-940)	16000 (4900-28000)
Finland	1995	Female	15 to 19	72% (55-86%)	28% (14-45%)	2.6 (1.8-3.7)	3.5% (2.2-4.8%)	13% (6.8-19%)	4.9 (2.6-7.4)	600 (400-850)
Finland	1995	Female	20 to 24	77% (62-89%)	23% (11-38%)	3.4 (2.3-4.8)	5.8% (4.1-7.6%)	16% (8.9-24%)	7.8 (4.4-12)	1100 (790-1500)
Finland	1995	Female	25 to 29	79% (66-90%)	21% (10-34%)	2.7 (2-3.5)	4.6% (3.2-5.9%)	13% (6.5-19%)	8.8 (4.4-13)	1100 (790-1500)
Finland	1995	Female	30 to 34	82% (69-92%)	18% (8.1-31%)	2.5 (1.8-3.2)	3.8% (2.5-5.1%)	12% (6.7-18%)	12 (6.5-17)	1100 (730-1500)
Finland	1995	Female	35 to 39	83% (68-93%)	17% (7.4-32%)	2.6 (1.8-3.4)	4.6% (3.1-6.2%)	14% (8.8-18%)	23 (15-30)	1600 (1100-2100)
Finland	1995	Female	40 to 44	83% (67-92%)	17% (7.9-33%)	2.9 (2.2-3.8)	5.9% (4.3-7.7%)	16% (12-20%)	42 (31-53)	2400 (1800-3100)
Finland	1995	Female	45 to 49	83% (67-92%)	17% (8-33%)	3.5 (2.7-4.6)	7.2% (5.3-9.4%)	17% (13-21%)	73 (55-92)	3600 (2700-4500)
Finland	1995	Female	50 to 54	83% (68-92%)	17% (7.8-32%)	3.5 (2.6-4.5)	7.1% (5.2-9.2%)	15% (11-18%)	68 (51-86)	2900 (2100-3700)
Finland	1995	Female	55 to 59	82% (67-92%)	18% (8.2-33%)	2.9 (2.1-3.7)	5.2% (3.4-7.3%)	9.9% (6.9-13%)	58 (40-78)	2100 (1400-2800)
Finland	1995	Female	60 to 64	79% (62-91%)	21% (9-38%)	2.3 (1.6-3.3)	3% (0.97-5.3%)	5.2% (1.9-8.7%)	44 (16-74)	1300 (410-2300)
Finland	1995	Female	65 to 69	77% (57-90%)	23% (9.6-43%)	2.1 (1.4-2.9)	1.1% (-1.2-3.4%)	1.8% (-1.7-5%)	27 (-26-78)	670 (-670-2000)
Finland	1995	Female	70 to 74	75% (57-88%)	25% (12-43%)	1.8 (1.2-2.6)	-0.29% (-2.8-2.5%)	-0.39% (-3.9-3.5%)	-10 (-100-90)	-210 (-2000-1700)
Finland	1995	Female	75 to 79	72% (51-87%)	28% (13-49%)	1.7 (1.1-2.5)	-1.1% (-3.9-1.9%)	-1.5% (-5.1-2.3%)	-54 (-190-84)	-760 (-2700-1300)
Finland	1995	Female	80 to 84	69% (48-85%)	31% (15-52%)	1.8 (1.1-2.6)	-1% (-4.1-2%)	-1.4% (-5.2-2.2%)	-76 (-280-110)	-750 (-3000-1400)
Finland	1995	Female	85 to 89	66% (47-82%)	34% (18-53%)	1.8 (1.1-2.7)	-0.62% (-3.5-2.6%)	-0.95% (-4.4-2.9%)	-49 (-230-150)	-300 (-1700-1200)
Finland	1995	Female	90 to 94	65% (46-82%)	35% (18-54%)	1.9 (1.1-2.9)	-0.19% (-3.3-3.4%)	-0.48% (-4.3-3.7%)	-13 (-110-97)	-31 (-550-540)
Finland	1995	Female	95 plus	66% (42-85%)	34% (15-58%)	2 (1-3.3)	0.29% (-3.1-4.6%)	0.051% (-4.5-2%)	0.41 (-32-42)	9.4 (-100-150)
Finland	1995	Male	All Ages	83% (75-89%)	17% (11-25%)	4.7 (4.1-5.3)	12% (9.9-15%)	11% (6.9-15%)	2500 (1600-3500)	100000 (81000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	1995	Male	15 to 19	77% (61-89%)	23% (11-39%)	3.5 (2.4-4.9)	9.8% (6.3-14%)	18% (11-27%)	24 (14-35)	2200 (1400-3100)
Finland	1995	Male	20 to 24	82% (67-93%)	18% (7.5-33%)	5.4 (3.9-7.3)	19% (14-24%)	29% (20-38%)	50 (34-66)	4600 (3400-5900)
Finland	1995	Male	25 to 29	85% (75-94%)	15% (6.2-25%)	5.3 (3.8-6.9)	18% (14-23%)	28% (19-37%)	64 (43-84)	5600 (4200-7100)
Finland	1995	Male	30 to 34	88% (77-95%)	12% (4.6-23%)	5.5 (4.1-7.2)	17% (13-22%)	29% (21-37%)	87 (63-110)	6400 (4900-8000)
Finland	1995	Male	35 to 39	89% (77-96%)	11% (4.1-23%)	6 (4.4-8.1)	20% (15-24%)	31% (24-38%)	140 (110-180)	8800 (6900-11000)
Finland	1995	Male	40 to 44	90% (78-96%)	10% (3.5-22%)	6.7 (5-8.5)	21% (17-25%)	32% (25-37%)	230 (180-270)	12000 (9600-14000)
Finland	1995	Male	45 to 49	91% (78-97%)	9.3% (3.4-22%)	7.4 (5.7-9.6)	22% (17-25%)	31% (25-36%)	340 (270-400)	16000 (13000-19000)
Finland	1995	Male	50 to 54	91% (80-97%)	9% (3.4-20%)	7.6 (5.7-9.7)	20% (15-23%)	26% (21-32%)	290 (230-350)	12000 (9400-14000)
Finland	1995	Male	55 to 59	91% (81-97%)	9.1% (3.4-19%)	7.3 (5.7-9.3)	17% (12-21%)	21% (15-27%)	300 (220-380)	11000 (8000-13000)
Finland	1995	Male	60 to 64	90% (79-97%)	9.7% (3.4-21%)	6.5 (4.8-8.5)	13% (8-18%)	16% (9.3-21%)	310 (190-420)	9200 (5800-13000)
Finland	1995	Male	65 to 69	89% (77-96%)	11% (3.7-23%)	5.2 (3.9-6.7)	8.7% (3.9-13%)	9.8% (3.8-16%)	290 (110-460)	7200 (3300-11000)
Finland	1995	Male	70 to 74	87% (75-95%)	13% (5.2-25%)	4 (2.8-5.3)	4.3% (0.12-9.3%)	4.5% (-0.52-10%)	160 (-18-360)	3300 (87-7200)
Finland	1995	Male	75 to 79	86% (72-95%)	14% (5.4-28%)	3.4 (2.3-4.9)	2.5% (-1.4-7.4%)	2.4% (-2.3-8.1%)	76 (-74-260)	1400 (-830-4200)
Finland	1995	Male	80 to 84	84% (68-94%)	16% (6.4-32%)	3.3 (2.3-4.7)	2.5% (-1.2-6.9%)	2.4% (-1.9-7.5%)	80 (-61-250)	1000 (-490-2900)
Finland	1995	Male	85 to 89	82% (65-93%)	18% (7.2-35%)	3.3 (2.2-4.6)	2.8% (-0.85-7.1%)	2.7% (-1.4-7.8%)	61 (-31-170)	550 (-170-1400)
Finland	1995	Male	90 to 94	82% (65-93%)	18% (6.6-35%)	3.3 (2-4.9)	3% (-1-8.5%)	3% (-1.6-9.5%)	25 (-14-78)	150 (-52-430)
Finland	1995	Male	95 plus	83% (63-95%)	17% (5.4-37%)	3.4 (1.8-5.4)	3.5% (-1.4-10%)	3.6% (-2.1-11%)	6.7 (-3.9-21)	27 (-10-77)
Finland	2000	Female	All Ages	75% (67-82%)	25% (18-33%)	2.3 (2-2.7)	3.2% (1.8-4.6%)	1.7% (-1.1-4.6%)	420 (-280-1100)	22000 (13000-32000)
Finland	2000	Female	15 to 19	70% (52-85%)	30% (15-48%)	2.6 (1.7-3.6)	3.7% (2.5-5.1%)	13% (7.3-19%)	5.7 (3.3-8.6)	660 (440-920)
Finland	2000	Female	20 to 24	76% (60-89%)	24% (11-40%)	3.5 (2.3-4.9)	6.1% (4.5-8%)	17% (10-24%)	9.7 (5.8-14)	1300 (940-1700)
Finland	2000	Female	25 to 29	80% (66-91%)	20% (9.2-34%)	2.9 (2.1-3.8)	4.9% (3.4-6.2%)	14% (8.1-20%)	8.1 (4.6-11)	1000 (710-1300)
Finland	2000	Female	30 to 34	82% (68-92%)	18% (7.9-32%)	2.6 (1.9-3.4)	4.4% (3.1-6%)	15% (9.3-20%)	14 (8.7-18)	1200 (820-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	2000	Female	35 to 39	83% (69-93%)	17% (7.3-31%)	2.7 (1.9-3.7)	5.2% (3.8-6.9%)	16% (11-21%)	24 (17-31)	1700 (1300-2200)
Finland	2000	Female	40 to 44	84% (69-93%)	16% (7.3-31%)	3.1 (2.3-4)	7% (5.5-9%)	19% (15-23%)	47 (38-57)	2700 (2100-3200)
Finland	2000	Female	45 to 49	84% (67-93%)	16% (7.5-33%)	3.7 (2.8-4.8)	8.5% (6.6-11%)	20% (16-24%)	79 (64-95)	3900 (3100-4700)
Finland	2000	Female	50 to 54	83% (68-92%)	17% (7.7-32%)	3.7 (2.7-4.7)	8.2% (6.2-10%)	17% (13-20%)	100 (83-130)	4400 (3400-5400)
Finland	2000	Female	55 to 59	82% (67-92%)	18% (7.8-33%)	3 (2.2-4)	6.4% (4.6-8.3%)	12% (9.7-15%)	75 (58-93)	2700 (2000-3400)
Finland	2000	Female	60 to 64	80% (62-92%)	20% (8.4-38%)	2.5 (1.7-3.4)	4.2% (2.4-6.4%)	7.4% (4.5-10%)	63 (38-89)	1900 (1100-2800)
Finland	2000	Female	65 to 69	76% (55-90%)	24% (10-45%)	2.1 (1.4-2.9)	2.3% (0.35-4.2%)	3.8% (0.9-6.5%)	46 (11-81)	1100 (180-2100)
Finland	2000	Female	70 to 74	73% (54-86%)	27% (14-46%)	1.8 (1.2-2.6)	0.72% (-1.3-2.9%)	1.1% (-1.8-4.1%)	25 (-40-93)	470 (-860-1900)
Finland	2000	Female	75 to 79	69% (48-85%)	31% (15-52%)	1.7 (1.1-2.5)	-0.2% (-2.6-2.3%)	-0.35% (-3.5-2.8%)	-13 (-130-110)	-150 (-2000-1700)
Finland	2000	Female	80 to 84	68% (47-84%)	32% (16-53%)	1.8 (1.1-2.6)	-0.47% (-3.2-2.2%)	-0.78% (-4.1-2.6%)	-37 (-190-120)	-310 (-2000-1400)
Finland	2000	Female	85 to 89	67% (48-82%)	33% (18-52%)	1.9 (1.2-2.8)	-0.29% (-3-2.8%)	-0.61% (-3.9-3.1%)	-33 (-210-160)	-150 (-1600-1400)
Finland	2000	Female	90 to 94	66% (48-83%)	34% (17-52%)	2 (1.2-3.1)	0.067% (-2.9-3.7%)	-0.21% (-3.9-4.1%)	-7.3 (-130-140)	14 (-630-790)
Finland	2000	Female	95 plus	66% (43-85%)	34% (15-57%)	2.1 (1.1-3.5)	0.62% (-2.9-5.4%)	0.42% (-3.8-6%)	4.5 (-41-64)	27 (-130-240)
Finland	2000	Male	All Ages	83% (76-89%)	17% (11-24%)	4.9 (4.2-5.5)	13% (11-16%)	12% (8.2-15%)	2800 (1900-3700)	110000 (89000-130000)
Finland	2000	Male	15 to 19	75% (55-89%)	25% (11-45%)	3.5 (2.3-4.8)	9.3% (6.1-13%)	18% (11-27%)	21 (12-31)	2000 (1300-2800)
Finland	2000	Male	20 to 24	81% (65-91%)	19% (8.7-35%)	5.5 (3.9-7.3)	19% (14-24%)	28% (20-37%)	53 (37-69)	5000 (3800-6400)
Finland	2000	Male	25 to 29	85% (74-93%)	15% (6.6-26%)	5.4 (3.9-7.1)	19% (14-23%)	29% (21-37%)	54 (38-69)	4800 (3700-6100)
Finland	2000	Male	30 to 34	88% (77-95%)	12% (4.9-23%)	5.6 (4.2-7.3)	19% (15-23%)	31% (24-39%)	80 (60-99)	6000 (4700-7300)
Finland	2000	Male	35 to 39	89% (78-96%)	11% (4-22%)	6.2 (4.4-8.2)	20% (17-25%)	35% (28-41%)	130 (100-150)	8100 (6600-9600)
Finland	2000	Male	40 to 44	90% (79-96%)	9.6% (3.6-21%)	6.8 (5.1-8.8)	23% (19-27%)	36% (30-42%)	220 (180-250)	11000 (9600-13000)
Finland	2000	Male	45 to 49	91% (77-97%)	9% (3.4-23%)	7.6 (5.8-9.6)	24% (20-28%)	35% (30-40%)	330 (270-370)	15000 (13000-17000)
Finland	2000	Male	50 to 54	91% (79-97%)	8.8% (3.2-21%)	7.7 (5.8-9.8)	22% (18-25%)	30% (24-35%)	440 (360-510)	18000 (15000-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	2000	Male	55 to 59	91% (81-97%)	9.1% (3.1-19%)	7.4 (5.8-9.4)	18% (14-22%)	24% (19-28%)	340 (270-410)	12000 (9600-14000)
Finland	2000	Male	60 to 64	90% (79-97%)	9.7% (3.3-21%)	6.6 (5-8.5)	14% (10-18%)	18% (12-23%)	320 (220-420)	9800 (6900-13000)
Finland	2000	Male	65 to 69	89% (78-96%)	11% (3.7-22%)	5.3 (4.1-6.8)	9.8% (5.6-14%)	11% (6.2-17%)	280 (150-410)	7100 (4100-10000)
Finland	2000	Male	70 to 74	88% (76-95%)	12% (4.7-24%)	4.1 (2.9-5.5)	5.2% (1.2-9.6%)	5.6% (0.89-11%)	190 (31-370)	4100 (1000-7600)
Finland	2000	Male	75 to 79	87% (75-95%)	13% (5-25%)	3.6 (2.4-5.1)	3% (-0.75-7.7%)	2.9% (-1.6-8.5%)	110 (-58-310)	1900 (-500-5000)
Finland	2000	Male	80 to 84	85% (71-94%)	15% (6.2-29%)	3.5 (2.4-4.8)	2.8% (-0.7-6.9%)	2.7% (-1.4-7.5%)	82 (-42-230)	1100 (-270-2800)
Finland	2000	Male	85 to 89	83% (68-93%)	17% (7.1-32%)	3.4 (2.3-4.8)	3% (-0.52-7.2%)	2.9% (-1.1-7.8%)	72 (-26-190)	660 (-120-1600)
Finland	2000	Male	90 to 94	83% (69-93%)	17% (6.7-31%)	3.5 (2.2-5.1)	3.3% (-0.7-8.7%)	3.3% (-1.3-9.7%)	37 (-14-110)	220 (-48-600)
Finland	2000	Male	95 plus	83% (66-95%)	17% (5.3-34%)	3.6 (2-5.7)	4% (-1.2-10%)	4.1% (-1.9-12%)	10 (-4.6-29)	41 (-12-110)
Finland	2005	Female	All Ages	76% (68-83%)	24% (17-32%)	2.4 (2.1-2.8)	3.7% (2.5-5%)	2.5% (0.13-5.1%)	610 (32-1200)	26000 (17000-35000)
Finland	2005	Female	15 to 19	71% (53-85%)	29% (15-47%)	2.7 (1.8-3.8)	3.7% (2.5-5.1%)	13% (7.5-20%)	5.7 (3.2-8.4)	640 (430-880)
Finland	2005	Female	20 to 24	78% (62-90%)	22% (10-38%)	3.7 (2.5-5.2)	6% (4.4-7.9%)	18% (11-25%)	9.6 (6-14)	1300 (920-1700)
Finland	2005	Female	25 to 29	81% (68-91%)	19% (8.9-32%)	3 (2.2-3.9)	4.6% (3.4-6%)	15% (8.5-21%)	8.1 (4.7-11)	1000 (740-1300)
Finland	2005	Female	30 to 34	82% (70-92%)	18% (7.9-30%)	2.7 (1.9-3.5)	4.2% (3-5.5%)	16% (10-20%)	11 (7.1-14)	950 (690-1200)
Finland	2005	Female	35 to 39	84% (71-93%)	16% (6.9-29%)	2.8 (2-3.8)	5.2% (3.9-6.7%)	17% (13-22%)	22 (17-28)	1500 (1100-1900)
Finland	2005	Female	40 to 44	84% (71-93%)	16% (7-29%)	3.2 (2.4-4.2)	7% (5.4-8.7%)	20% (17-25%)	44 (36-54)	2500 (2000-3100)
Finland	2005	Female	45 to 49	85% (70-93%)	15% (7.2-30%)	3.9 (3-5)	8.8% (6.9-11%)	21% (18-25%)	76 (63-90)	3700 (3000-4500)
Finland	2005	Female	50 to 54	84% (70-93%)	16% (7.5-30%)	3.9 (2.9-5)	9.2% (7.2-11%)	19% (16-22%)	110 (92-130)	4700 (3800-5600)
Finland	2005	Female	55 to 59	83% (69-92%)	17% (7.5-31%)	3.1 (2.4-4.2)	7.5% (5.8-9.5%)	14% (12-17%)	130 (110-150)	4500 (3600-5600)
Finland	2005	Female	60 to 64	80% (64-92%)	20% (8.4-36%)	2.6 (1.8-3.6)	5.1% (3.5-7.1%)	9.3% (6.9-12%)	81 (60-110)	2400 (1700-3400)
Finland	2005	Female	65 to 69	77% (59-90%)	23% (10-41%)	2.3 (1.6-3)	3.1% (1.5-4.9%)	5.5% (2.9-8.1%)	63 (34-93)	1500 (710-2300)
Finland	2005	Female	70 to 74	74% (57-87%)	26% (13-43%)	1.9 (1.3-2.8)	1.4% (-0.47-3.4%)	2.3% (-0.38-5.1%)	40 (-6.8-90)	760 (-260-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	2005	Female	75 to 79	68% (48-84%)	32% (16-52%)	1.7 (1.1-2.5)	0.42% (-1.5-2.6%)	0.55% (-2-3.5%)	17 (-64-110)	290 (-1100-1800)
Finland	2005	Female	80 to 84	65% (43-81%)	35% (19-57%)	1.7 (1.1-2.5)	0.038% (-2.3-2.3%)	-0.14% (-3-2.7%)	-7.1 (-150-130)	27 (-1600-1600)
Finland	2005	Female	85 to 89	64% (45-80%)	36% (20-55%)	1.9 (1.2-2.7)	0.046% (-2.5-2.7%)	-0.23% (-3.3-2.9%)	-11 (-160-140)	22 (-1200-1300)
Finland	2005	Female	90 to 94	63% (44-81%)	37% (19-56%)	2 (1.2-3)	0.28% (-2.6-3.4%)	0.0095% (-3.4-3.8%)	0.36 (-130-140)	67 (-620-820)
Finland	2005	Female	95 plus	64% (40-83%)	36% (17-60%)	2.1 (1.1-3.4)	0.71% (-2.9-4.8%)	0.5% (-3.8-5.3%)	7.2 (-54-76)	43 (-170-290)
Finland	2005	Male	All Ages	84% (77-90%)	16% (10-23%)	5.1 (4.5-5.7)	14% (12-16%)	13% (9.5-16%)	3000 (2300-3900)	110000 (97000-130000)
Finland	2005	Male	15 to 19	75% (57-89%)	25% (11-43%)	3.6 (2.4-5)	9.1% (6.3-12%)	18% (11-26%)	18 (11-25)	1800 (1200-2400)
Finland	2005	Male	20 to 24	82% (67-92%)	18% (7.8-33%)	5.7 (4.1-7.6)	19% (15-23%)	28% (20-37%)	49 (35-64)	5000 (3900-6300)
Finland	2005	Male	25 to 29	86% (76-94%)	14% (5.9-24%)	5.6 (4.1-7.5)	18% (15-22%)	29% (21-37%)	48 (35-61)	4800 (3700-6000)
Finland	2005	Male	30 to 34	89% (78-96%)	11% (4.5-22%)	5.8 (4.2-7.7)	18% (15-22%)	31% (24-39%)	61 (47-76)	4800 (3900-5900)
Finland	2005	Male	35 to 39	90% (79-96%)	10% (4-21%)	6.3 (4.6-8.5)	20% (17-24%)	35% (28-41%)	110 (91-130)	7200 (5900-8500)
Finland	2005	Male	40 to 44	91% (81-96%)	9.4% (3.6-19%)	7 (5.3-9)	23% (19-26%)	37% (31-42%)	190 (160-220)	10000 (8600-12000)
Finland	2005	Male	45 to 49	91% (80-97%)	8.7% (3.2-20%)	7.8 (6-10)	25% (21-28%)	37% (31-41%)	300 (260-340)	14000 (12000-16000)
Finland	2005	Male	50 to 54	92% (82-97%)	8.3% (3-18%)	8 (6-10)	24% (20-27%)	33% (28-37%)	430 (370-500)	18000 (15000-20000)
Finland	2005	Male	55 to 59	91% (82-97%)	8.6% (3-18%)	7.7 (6.1-9.5)	20% (17-24%)	26% (22-31%)	540 (440-630)	19000 (16000-22000)
Finland	2005	Male	60 to 64	91% (81-97%)	9% (3.1-19%)	6.9 (5.2-8.9)	16% (12-19%)	20% (15-25%)	390 (280-480)	12000 (8800-14000)
Finland	2005	Male	65 to 69	90% (79-97%)	9.9% (3.4-21%)	5.6 (4.4-7)	11% (7.4-15%)	13% (8.6-18%)	310 (200-420)	8000 (5300-11000)
Finland	2005	Male	70 to 74	89% (78-95%)	11% (4.7-22%)	4.3 (3.1-5.8)	6.2% (2.3-11%)	7% (2.2-12%)	200 (66-360)	4400 (1700-7500)
Finland	2005	Male	75 to 79	87% (74-95%)	13% (5-26%)	3.7 (2.5-5.3)	3.7% (0.12-8.5%)	3.9% (-0.47-9.5%)	150 (-18-370)	2600 (85-5900)
Finland	2005	Male	80 to 84	85% (70-94%)	15% (6.2-30%)	3.6 (2.5-4.9)	3.3% (0.13-7.3%)	3.2% (-0.55-8%)	120 (-20-290)	1600 (62-3500)
Finland	2005	Male	85 to 89	83% (68-93%)	17% (7.2-32%)	3.5 (2.4-4.9)	3.1% (-0.31-7.2%)	3% (-1-7.8%)	69 (-23-180)	660 (-67-1600)
Finland	2005	Male	90 to 94	83% (68-93%)	17% (6.5-32%)	3.6 (2.3-5.2)	3.4% (-0.69-8.6%)	3.3% (-1.4-9.6%)	42 (-17-120)	260 (-54-660)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	2005	Male	95 plus	83% (66-95%)	17% (5.1-34%)	3.6 (2-5.7)	3.9% (-1.1-10%)	4% (-1.9-12%)	13 (-6.2-37)	53 (-15-140)
Finland	2010	Female	All Ages	76% (68-82%)	24% (18-32%)	2.4 (2-2.8)	3.5% (2.3-4.7%)	2.5% (0.18-5.1%)	640 (47-1300)	25000 (17000-33000)
Finland	2010	Female	15 to 19	72% (56-86%)	28% (14-44%)	2.7 (1.9-3.7)	3.2% (2.1-4.4%)	13% (7.2-19%)	4.5 (2.5-6.7)	540 (350-750)
Finland	2010	Female	20 to 24	78% (63-89%)	22% (11-37%)	3.7 (2.5-5.1)	5.6% (4-7.5%)	17% (10-25%)	9.4 (5.5-14)	1100 (800-1600)
Finland	2010	Female	25 to 29	80% (67-91%)	20% (9.3-33%)	2.9 (2.1-3.9)	4.5% (3.2-5.9%)	14% (8.1-20%)	8.7 (5-12)	1000 (720-1400)
Finland	2010	Female	30 to 34	82% (69-92%)	18% (8-31%)	2.6 (1.8-3.5)	4.1% (3-5.4%)	15% (9.7-20%)	11 (6.7-14)	990 (710-1300)
Finland	2010	Female	35 to 39	83% (69-93%)	17% (7.2-31%)	2.8 (2-3.7)	4.7% (3.5-6.1%)	17% (12-21%)	15 (11-19)	1100 (870-1400)
Finland	2010	Female	40 to 44	84% (69-93%)	16% (7.3-31%)	3.2 (2.3-4.1)	6.2% (4.8-7.9%)	20% (16-23%)	33 (27-40)	2000 (1600-2500)
Finland	2010	Female	45 to 49	85% (70-93%)	15% (7.1-30%)	3.8 (2.9-5)	8.2% (6.4-10%)	21% (18-25%)	64 (53-76)	3200 (2600-3900)
Finland	2010	Female	50 to 54	84% (69-93%)	16% (7.4-31%)	3.8 (2.8-4.9)	8.7% (6.7-11%)	19% (16-22%)	93 (78-110)	4000 (3200-4800)
Finland	2010	Female	55 to 59	82% (69-92%)	18% (7.9-31%)	3.1 (2.3-4)	7.4% (5.7-9.4%)	15% (12-18%)	110 (93-130)	4000 (3200-4900)
Finland	2010	Female	60 to 64	80% (63-91%)	20% (8.6-37%)	2.5 (1.8-3.5)	5.2% (3.6-7.2%)	9.9% (7.6-13%)	120 (88-150)	3400 (2400-4500)
Finland	2010	Female	65 to 69	78% (60-90%)	22% (9.6-40%)	2.2 (1.6-3)	3.2% (1.7-5%)	6% (3.7-8.3%)	72 (45-100)	1700 (940-2500)
Finland	2010	Female	70 to 74	75% (59-87%)	25% (13-41%)	1.9 (1.3-2.7)	1.6% (-0.13-3.6%)	2.9% (0.48-5.4%)	49 (8.2-91)	880 (-74-1900)
Finland	2010	Female	75 to 79	68% (49-85%)	32% (15-51%)	1.7 (1.1-2.5)	0.64% (-1.1-2.7%)	1% (-1.4-3.7%)	28 (-37-99)	390 (-700-1600)
Finland	2010	Female	80 to 84	65% (42-82%)	35% (18-58%)	1.7 (1.1-2.4)	0.18% (-1.8-2.3%)	0.15% (-2.4-2.8%)	6.8 (-110-130)	130 (-1300-1600)
Finland	2010	Female	85 to 89	62% (44-79%)	38% (21-56%)	1.8 (1.1-2.6)	0.16% (-2.1-2.7%)	-0.014% (-2.8-3.1%)	-0.85 (-170-180)	97 (-1200-1600)
Finland	2010	Female	90 to 94	62% (42-80%)	38% (20-58%)	1.8 (1.1-2.9)	0.31% (-2.5-3.6%)	0.095% (-3.3-4%)	3.8 (-130-160)	82 (-650-910)
Finland	2010	Female	95 plus	62% (38-83%)	38% (17-62%)	2 (0.97-3.3)	0.7% (-2.9-4.7%)	0.53% (-3.7-5.4%)	10 (-73-100)	57 (-240-380)
Finland	2010	Male	All Ages	84% (77-89%)	16% (11-23%)	4.9 (4.3-5.7)	13% (11-15%)	11% (8.6-15%)	2900 (2200-3800)	110000 (90000-120000)
Finland	2010	Male	15 to 19	74% (54-88%)	26% (12-46%)	3.4 (2.3-4.8)	8.4% (5.6-11%)	18% (10-26%)	16 (9.2-23)	1600 (1100-2200)
Finland	2010	Male	20 to 24	82% (67-92%)	18% (8-33%)	5.6 (4-7.4)	18% (14-23%)	28% (19-36%)	47 (33-62)	4600 (3500-5800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	2010	Male	25 to 29	86% (76-94%)	14% (6.4-24%)	5.5 (4-7.2)	17% (14-21%)	27% (19-35%)	52 (37-67)	4900 (3700-6100)
Finland	2010	Male	30 to 34	89% (78-95%)	11% (4.7-22%)	5.7 (4.1-7.6)	17% (14-20%)	29% (22-36%)	60 (45-75)	4900 (3800-6000)
Finland	2010	Male	35 to 39	89% (79-96%)	11% (4.1-21%)	6.2 (4.4-8.1)	19% (15-22%)	33% (27-39%)	79 (63-94)	5400 (4400-6400)
Finland	2010	Male	40 to 44	90% (78-96%)	9.9% (3.6-22%)	6.8 (5-8.7)	21% (18-25%)	36% (30-41%)	140 (120-160)	8100 (6700-9300)
Finland	2010	Male	45 to 49	91% (79-97%)	8.6% (3.3-21%)	7.7 (5.9-9.9)	23% (19-27%)	36% (31-40%)	240 (210-280)	12000 (10000-13000)
Finland	2010	Male	50 to 54	92% (81-97%)	8.2% (2.9-19%)	7.9 (6-10)	23% (19-26%)	33% (28-38%)	350 (290-400)	15000 (12000-17000)
Finland	2010	Male	55 to 59	91% (82-97%)	8.8% (3.1-18%)	7.5 (5.9-9.3)	20% (17-24%)	27% (23-32%)	460 (380-530)	16000 (13000-19000)
Finland	2010	Male	60 to 64	91% (80-97%)	9.2% (3-20%)	6.7 (5-8.7)	16% (12-20%)	21% (16-25%)	520 (390-640)	16000 (12000-19000)
Finland	2010	Male	65 to 69	90% (80-96%)	10% (3.5-20%)	5.4 (4.3-6.9)	11% (7.8-15%)	14% (9.3-18%)	330 (220-440)	8600 (6000-11000)
Finland	2010	Male	70 to 74	88% (76-95%)	12% (4.9-24%)	4.1 (3-5.6)	6.3% (3-9.9%)	7.3% (3.1-12%)	210 (89-350)	4600 (2100-7400)
Finland	2010	Male	75 to 79	86% (73-94%)	14% (5.9-27%)	3.5 (2.4-4.9)	3.8% (0.56-7.6%)	4% (0.18-8.7%)	140 (6.4-310)	2500 (370-5000)
Finland	2010	Male	80 to 84	84% (68-93%)	16% (6.7-32%)	3.4 (2.4-4.7)	3.1% (0.16-6.7%)	3.1% (-0.54-7.4%)	130 (-23-310)	1800 (93-3800)
Finland	2010	Male	85 to 89	83% (68-93%)	17% (7.2-32%)	3.4 (2.4-4.8)	2.9% (-0.19-6.9%)	2.8% (-1-7.5%)	89 (-32-240)	860 (-60-2100)
Finland	2010	Male	90 to 94	83% (68-93%)	17% (6.7-32%)	3.5 (2.2-5.2)	3.1% (-0.67-8.2%)	3.1% (-1.5-9.2%)	44 (-22-130)	280 (-63-750)
Finland	2010	Male	95 plus	83% (66-95%)	17% (5.4-34%)	3.5 (2-5.5)	3.7% (-1.1-9.7%)	3.7% (-1.9-11%)	16 (-8.4-49)	67 (-20-180)
Finland	2016	Female	All Ages	75% (67-81%)	25% (19-33%)	2.3 (1.8-2.9)	2.9% (1.6-4.4%)	2.1% (-0.57-5.4%)	540 (-150-1400)	20000 (11000-31000)
Finland	2016	Female	15 to 19	72% (53-87%)	28% (13-47%)	2.6 (1.7-3.8)	3% (1.9-4.1%)	13% (6.9-19%)	3.6 (1.9-5.5)	450 (290-630)
Finland	2016	Female	20 to 24	76% (59-89%)	24% (11-41%)	3.5 (2.3-5)	4.6% (3.2-6.1%)	16% (9.5-24%)	6 (3.4-8.9)	930 (640-1300)
Finland	2016	Female	25 to 29	79% (64-90%)	21% (9.8-36%)	2.8 (1.9-4)	3.8% (2.7-4.9%)	13% (7.4-19%)	5.7 (3.1-8.2)	830 (580-1100)
Finland	2016	Female	30 to 34	80% (65-92%)	20% (8.4-35%)	2.5 (1.7-3.6)	3.6% (2.5-4.7%)	14% (8.9-19%)	8.7 (5.7-12)	890 (630-1200)
Finland	2016	Female	35 to 39	82% (66-92%)	18% (8.2-34%)	2.6 (1.7-3.9)	3.7% (2.7-4.9%)	15% (11-20%)	12 (8.4-16)	990 (710-1300)
Finland	2016	Female	40 to 44	82% (66-92%)	18% (7.8-34%)	3 (2-4.2)	4.9% (3.5-6.4%)	18% (14-22%)	20 (16-25)	1300 (970-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	2016	Female	45 to 49	82% (65-92%)	18% (7.8-35%)	3.6 (2.5-4.9)	6.8% (5.2-8.8%)	20% (16-24%)	44 (35-54)	2300 (1700-2900)
Finland	2016	Female	50 to 54	82% (65-92%)	18% (8.2-35%)	3.5 (2.5-4.9)	7.3% (5.5-9.4%)	18% (14-21%)	69 (54-84)	3100 (2400-3800)
Finland	2016	Female	55 to 59	81% (65-92%)	19% (8.1-35%)	2.9 (2-4.1)	6.4% (4.7-8.5%)	14% (12-17%)	83 (66-100)	3000 (2300-3900)
Finland	2016	Female	60 to 64	79% (60-92%)	21% (8.3-40%)	2.5 (1.6-3.6)	4.7% (3-6.9%)	9.5% (6.9-12%)	88 (63-120)	2600 (1700-3800)
Finland	2016	Female	65 to 69	78% (58-91%)	22% (9.4-42%)	2.2 (1.4-3.1)	3% (1.4-4.9%)	5.8% (3.4-8.4%)	92 (53-140)	2200 (1000-3500)
Finland	2016	Female	70 to 74	75% (57-88%)	25% (12-43%)	1.9 (1.2-2.8)	1.5% (-0.13-3.5%)	2.9% (0.61-5.6%)	47 (9.7-92)	840 (-71-1900)
Finland	2016	Female	75 to 79	72% (53-87%)	28% (13-47%)	1.8 (1.1-2.7)	0.64% (-1.2-2.9%)	1.2% (-1.4-4%)	29 (-35-100)	390 (-730-1800)
Finland	2016	Female	80 to 84	70% (49-85%)	30% (15-51%)	1.8 (1.1-2.8)	0.21% (-1.9-2.7%)	0.23% (-2.5-3.5%)	8.8 (-100-140)	130 (-1200-1700)
Finland	2016	Female	85 to 89	67% (49-83%)	33% (17-51%)	1.9 (1.1-2.9)	0.2% (-2.2-3.2%)	0.045% (-3-3.8%)	2.6 (-180-220)	120 (-1400-1900)
Finland	2016	Female	90 to 94	66% (48-83%)	34% (17-52%)	2 (1.1-3.1)	0.35% (-2.6-4.1%)	0.16% (-3.3-4.6%)	8.4 (-170-250)	120 (-870-1400)
Finland	2016	Female	95 plus	66% (43-85%)	34% (15-57%)	2.1 (1-3.5)	0.73% (-3.2-5.3%)	0.58% (-4.1-6.1%)	13 (-92-140)	68 (-300-490)
Finland	2016	Male	All Ages	82% (75-88%)	18% (12-25%)	4.6 (3.6-5.8)	11% (9.1-14%)	9.9% (6.4-14%)	2600 (1600-3700)	89000 (69000-110000)
Finland	2016	Male	15 to 19	74% (54-89%)	26% (11-46%)	3.3 (2.1-4.9)	7% (4.8-9.5%)	17% (9.2-24%)	9.2 (4.9-14)	1100 (750-1500)
Finland	2016	Male	20 to 24	79% (62-91%)	21% (9.2-38%)	5.2 (3.5-7.3)	16% (12-20%)	25% (17-35%)	32 (21-45)	3700 (2700-4800)
Finland	2016	Male	25 to 29	82% (68-93%)	18% (7.3-32%)	5.1 (3.4-7.2)	15% (12-19%)	25% (17-33%)	37 (24-51)	3900 (3000-5100)
Finland	2016	Male	30 to 34	85% (70-94%)	15% (5.8-30%)	5.3 (3.5-7.7)	15% (12-18%)	27% (19-35%)	47 (34-61)	4200 (3300-5300)
Finland	2016	Male	35 to 39	87% (75-95%)	13% (4.9-25%)	5.8 (3.8-8.4)	16% (13-20%)	31% (24-38%)	66 (51-82)	4800 (3900-5900)
Finland	2016	Male	40 to 44	89% (76-96%)	11% (4.1-24%)	6.5 (4.5-9)	18% (15-22%)	35% (28-40%)	92 (73-110)	5500 (4500-6600)
Finland	2016	Male	45 to 49	89% (76-96%)	11% (3.8-24%)	7.2 (5.1-9.8)	21% (17-24%)	35% (29-40%)	170 (140-200)	8400 (7000-10000)
Finland	2016	Male	50 to 54	90% (77-96%)	10% (3.7-23%)	7.4 (5.2-10)	20% (17-24%)	32% (27-37%)	250 (210-300)	11000 (9000-13000)
Finland	2016	Male	55 to 59	89% (78-97%)	11% (3.3-22%)	7 (5.1-9.4)	19% (15-22%)	27% (22-31%)	340 (270-410)	12000 (10000-15000)
Finland	2016	Male	60 to 64	89% (74-97%)	11% (3.5-26%)	6.3 (4.4-8.6)	15% (11-19%)	20% (15-25%)	390 (290-500)	12000 (8900-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Finland	2016	Male	65 to 69	88% (74-96%)	12% (4-26%)	5.2 (3.7-7)	11% (7.2-14%)	13% (8.9-18%)	420 (270-560)	11000 (7300-15000)
Finland	2016	Male	70 to 74	88% (75-95%)	12% (5-25%)	4 (2.7-5.6)	6.2% (3-10%)	7.5% (3.3-12%)	220 (94-370)	4700 (2200-7800)
Finland	2016	Male	75 to 79	87% (73-95%)	13% (5.2-27%)	3.4 (2.3-5.1)	3.8% (0.68-8%)	4.2% (0.47-9.5%)	150 (17-340)	2700 (460-5600)
Finland	2016	Male	80 to 84	85% (70-94%)	15% (6.3-30%)	3.4 (2.2-4.9)	3.2% (0.15-7.2%)	3.3% (-0.38-8%)	130 (-15-330)	1800 (76-4100)
Finland	2016	Male	85 to 89	83% (68-93%)	17% (6.9-32%)	3.4 (2.1-4.9)	2.9% (-0.29-7.6%)	2.9% (-0.85-8.3%)	120 (-35-330)	1100 (-110-2900)
Finland	2016	Male	90 to 94	83% (68-94%)	17% (6-32%)	3.4 (2-5.4)	3.1% (-0.72-8.5%)	3.1% (-1.3-9.5%)	67 (-29-200)	420 (-98-1100)
Finland	2016	Male	95 plus	83% (66-95%)	17% (4.9-34%)	3.5 (1.9-5.8)	3.5% (-1.3-10%)	3.7% (-2-12%)	21 (-11-66)	84 (-30-240)
France	1990	Female	All Ages	86% (83-89%)	14% (11-17%)	3.7 (3-4.4)	7.5% (5.6-9.5%)	10% (6.6-15%)	26000 (16000-37000)	550000 (420000-690000)
France	1990	Female	15 to 19	86% (77-92%)	14% (7.5-23%)	3.1 (2.1-4.3)	3.9% (2.4-5.7%)	14% (7.5-21%)	96 (53-140)	9300 (5600-13000)
France	1990	Female	20 to 24	90% (84-94%)	10% (5.7-16%)	3.6 (2.4-5.2)	5.1% (3.3-7.1%)	15% (8.8-22%)	140 (82-200)	14000 (9300-19000)
France	1990	Female	25 to 29	91% (85-95%)	8.9% (5-15%)	3.5 (2.4-4.8)	4.6% (3.1-6.3%)	14% (8.1-20%)	160 (93-230)	15000 (9700-20000)
France	1990	Female	30 to 34	90% (84-94%)	10% (5.7-16%)	3.6 (2.5-4.9)	4.9% (3.4-6.6%)	15% (10-20%)	220 (150-300)	17000 (12000-22000)
France	1990	Female	35 to 39	87% (80-92%)	13% (7.8-20%)	4 (2.8-5.5)	6.4% (4.7-8.3%)	19% (14-24%)	390 (290-480)	24000 (18000-30000)
France	1990	Female	40 to 44	86% (80-91%)	14% (8.7-20%)	4.5 (3.3-6.1)	8.1% (6.3-10%)	21% (17-26%)	630 (510-760)	34000 (27000-42000)
France	1990	Female	45 to 49	88% (82-92%)	12% (8-18%)	5.3 (3.9-7.1)	10% (7.7-13%)	23% (19-28%)	680 (560-820)	32000 (26000-39000)
France	1990	Female	50 to 54	90% (84-93%)	10% (6.6-16%)	5.9 (4.4-7.7)	11% (8.7-14%)	23% (19-27%)	1000 (820-1200)	41000 (34000-51000)
France	1990	Female	55 to 59	92% (88-95%)	8.3% (5.5-12%)	6.2 (4.6-7.9)	11% (8.8-14%)	21% (17-25%)	1400 (1200-1700)	51000 (41000-63000)
France	1990	Female	60 to 64	92% (88-95%)	7.8% (4.9-12%)	6.1 (4.5-8)	11% (7.9-14%)	18% (14-23%)	1800 (1400-2300)	56000 (42000-71000)
France	1990	Female	65 to 69	91% (86-94%)	9.1% (5.7-14%)	5.7 (4.4-7.2)	10% (7.3-13%)	15% (12-20%)	2300 (1700-2900)	59000 (44000-76000)
France	1990	Female	70 to 74	88% (83-92%)	12% (8-17%)	5.2 (3.8-6.9)	9% (5.6-13%)	13% (8.2-18%)	1800 (1100-2500)	37000 (24000-52000)
France	1990	Female	75 to 79	85% (77-91%)	15% (9.3-23%)	4.7 (3.3-6.4)	7.9% (3.9-12%)	10% (5-16%)	3400 (1600-5300)	54000 (27000-83000)
France	1990	Female	80 to 84	82% (72-89%)	18% (11-28%)	4.4 (3.2-5.8)	7.3% (3.6-12%)	8.9% (4.1-14%)	4500 (2100-7300)	52000 (25000-83000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	1990	Female	85 to 89	80% (70-87%)	20% (13-30%)	4.2 (2.9-5.7)	6.6% (2.9-11%)	7.7% (3.2-13%)	4100 (1700-7100)	34000 (15000-56000)
France	1990	Female	90 to 94	79% (69-87%)	21% (13-31%)	4.1 (2.7-5.8)	6% (1.9-11%)	6.8% (1.9-13%)	2300 (630-4200)	13000 (4100-23000)
France	1990	Female	95 plus	78% (64-88%)	22% (12-36%)	4 (2.3-6.2)	5.8% (0.68-11%)	6.5% (0.34-13%)	820 (44-1700)	3000 (340-6000)
France	1990	Male	All Ages	92% (90-94%)	7.6% (5.6-10%)	7.1 (5.8-8.4)	18% (16-21%)	21% (18-25%)	57000 (49000-65000)	160000 (140000-1800000)
France	1990	Male	15 to 19	89% (82-94%)	11% (5.7-18%)	5.2 (3.7-7)	12% (8-16%)	23% (15-31%)	420 (270-580)	34000 (23000-45000)
France	1990	Male	20 to 24	94% (89-97%)	6.2% (3.3-11%)	7.8 (5.8-10)	22% (16-28%)	33% (23-43%)	1000 (730-1400)	79000 (58000-100000)
France	1990	Male	25 to 29	95% (91-97%)	4.9% (2.6-8.6%)	7.6 (5.6-9.7)	19% (15-25%)	30% (22-38%)	1000 (740-1300)	74000 (56000-92000)
France	1990	Male	30 to 34	96% (92-98%)	4.3% (2.2-8%)	7.7 (5.9-9.9)	18% (14-23%)	29% (22-36%)	1100 (860-1400)	74000 (58000-90000)
France	1990	Male	35 to 39	96% (93-98%)	4.3% (2.1-7.4%)	8 (6.2-10)	20% (16-24%)	31% (25-37%)	1600 (1200-1800)	91000 (74000-110000)
France	1990	Male	40 to 44	96% (92-98%)	4.3% (2.2-7.5%)	9.1 (7.1-11)	22% (18-26%)	33% (28-37%)	2300 (2000-2700)	120000 (100000-140000)
France	1990	Male	45 to 49	96% (93-98%)	4.1% (2.2-7.4%)	11 (8.4-13)	24% (21-27%)	34% (30-38%)	2500 (2200-2800)	110000 (98000-120000)
France	1990	Male	50 to 54	96% (93-98%)	3.9% (2.2-6.7%)	12 (9.2-14)	25% (21-28%)	33% (29-36%)	3600 (3100-4000)	140000 (120000-160000)
France	1990	Male	55 to 59	96% (94-98%)	3.7% (2.2-5.9%)	12 (9.6-15)	24% (20-27%)	30% (26-33%)	5100 (4500-5700)	170000 (150000-190000)
France	1990	Male	60 to 64	96% (94-98%)	3.7% (2.1-6.1%)	12 (9.3-14)	22% (18-25%)	26% (23-30%)	6200 (5300-7100)	180000 (150000-210000)
France	1990	Male	65 to 69	96% (94-98%)	4% (2.3-6.3%)	11 (8.9-14)	19% (16-22%)	23% (19-27%)	6800 (5700-7900)	170000 (140000-190000)
France	1990	Male	70 to 74	96% (93-97%)	4.4% (2.7-7.1%)	10 (8-13)	17% (14-21%)	21% (17-24%)	4700 (3800-5500)	91000 (75000-110000)
France	1990	Male	75 to 79	95% (91-97%)	4.9% (2.5-8.5%)	9.4 (7.2-12)	18% (13-19%)	18% (14-22%)	7100 (5600-8600)	110000 (86000-130000)
France	1990	Male	80 to 84	94% (89-97%)	5.9% (3.1-11%)	8.5 (6.6-11)	15% (12-18%)	17% (13-20%)	7000 (5400-8500)	78000 (61000-95000)
France	1990	Male	85 to 89	93% (87-96%)	7.4% (4.1-13%)	7.8 (5.9-9.9)	13% (9.6-16%)	15% (11-19%)	4500 (3200-5600)	35000 (26000-44000)
France	1990	Male	90 to 94	92% (87-96%)	8.1% (4.3-13%)	7.2 (5.2-9.6)	12% (7.7-15%)	13% (8.4-17%)	1500 (1000-2000)	8400 (5500-11000)
France	1990	Male	95 plus	91% (84-96%)	8.7% (3.9-16%)	6.5 (4.4-9.3)	10% (4.8-15%)	11% (4.9-17%)	290 (130-420)	1100 (500-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	1995	Female	All Ages	87% (84-89%)	13% (11-16%)	3.2 (2.7-3.9)	6.4% (4.7-8.2%)	8.1% (4.7-12%)	20000 (12000-30000)	460000 (350000-590000)
France	1995	Female	15 to 19	87% (79-92%)	13% (7.6-21%)	2.7 (1.8-3.7)	3.4% (2-4.9%)	12% (6.1-18%)	67 (34-100)	6800 (4000-9900)
France	1995	Female	20 to 24	91% (86-95%)	9.2% (5.4-14%)	3.1 (2.1-4.5)	4.6% (2.9-6.4%)	13% (7.1-20%)	110 (60-170)	12000 (7800-17000)
France	1995	Female	25 to 29	92% (87-95%)	8.1% (4.8-13%)	3 (2.1-4.1)	4% (2.6-5.5%)	12% (6.5-17%)	130 (70-180)	12000 (8100-17000)
France	1995	Female	30 to 34	91% (86-94%)	9.3% (5.9-14%)	3.1 (2.2-4.1)	4.1% (2.8-5.7%)	12% (7.7-16%)	180 (120-250)	14000 (10000-19000)
France	1995	Female	35 to 39	88% (83-92%)	12% (7.6-17%)	3.5 (2.5-4.7)	5.5% (4-7.2%)	16% (12-20%)	330 (240-420)	21000 (15000-27000)
France	1995	Female	40 to 44	87% (83-92%)	13% (8.5-17%)	4 (2.9-5.2)	7.2% (5.5-9.3%)	19% (15-23%)	560 (450-680)	31000 (24000-38000)
France	1995	Female	45 to 49	89% (85-92%)	11% (7.7-15%)	4.7 (3.4-6.2)	9% (7-12%)	21% (17-25%)	880 (720-1100)	42000 (34000-51000)
France	1995	Female	50 to 54	91% (87-94%)	9.2% (6.1-13%)	5.2 (3.9-6.8)	10% (7.8-13%)	21% (17-25%)	850 (700-1000)	35000 (29000-43000)
France	1995	Female	55 to 59	92% (90-95%)	7.5% (5.2-10%)	5.4 (4.1-7)	10% (7.7-13%)	19% (15-23%)	1100 (890-1300)	39000 (31000-48000)
France	1995	Female	60 to 64	93% (90-95%)	7.2% (4.7-10%)	5.4 (4-7.1)	9.6% (7.1-12%)	16% (13-20%)	1500 (1200-1900)	46000 (35000-58000)
France	1995	Female	65 to 69	92% (88-95%)	8.1% (5.5-12%)	5.1 (3.9-6.4)	8.6% (6.3-11%)	13% (9.9-17%)	1800 (1400-2400)	48000 (35000-62000)
France	1995	Female	70 to 74	90% (86-93%)	10% (7.5-14%)	4.6 (3.4-6)	7.4% (4.4-11%)	11% (6.6-15%)	2300 (1400-3300)	49000 (29000-71000)
France	1995	Female	75 to 79	86% (80-91%)	14% (8.9-20%)	4.1 (2.9-5.6)	6.3% (2.8-10%)	8.1% (3.6-13%)	1700 (750-2700)	28000 (13000-44000)
France	1995	Female	80 to 84	82% (73-89%)	18% (11-27%)	3.8 (2.8-5)	5.5% (2.4-8.9%)	6.6% (2.6-11%)	3100 (1200-5100)	37000 (16000-60000)
France	1995	Female	85 to 89	79% (69-87%)	21% (13-31%)	3.6 (2.5-4.8)	4.8% (1.7-8.6%)	5.5% (1.7-10%)	3200 (990-5800)	27000 (9800-48000)
France	1995	Female	90 to 94	79% (68-87%)	21% (13-32%)	3.5 (2.3-4.9)	4.3% (0.76-8.6%)	4.7% (0.5-9.9%)	2000 (210-4000)	11000 (2000-22000)
France	1995	Female	95 plus	78% (64-88%)	22% (12-36%)	3.4 (1.9-5.2)	4.2% (-0.31-9.1%)	4.5% (-0.8-10%)	780 (-140-1800)	3000 (-220-6600)
France	1995	Male	All Ages	94% (92-95%)	6.1% (4.6-7.9%)	6.3 (5.3-7.4)	17% (14-19%)	19% (16-22%)	52000 (43000-60000)	140000 (120000-160000)
France	1995	Male	15 to 19	91% (86-95%)	8.9% (4.9-14%)	4.6 (3.4-6.1)	10% (6.7-14%)	21% (13-29%)	290 (180-400)	24000 (16000-32000)
France	1995	Male	20 to 24	95% (92-97%)	4.7% (2.7-7.7%)	6.8 (5.1-9)	19% (14-25%)	30% (21-40%)	810 (560-1100)	63000 (46000-81000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	1995	Male	25 to 29	96% (94-98%)	3.8% (2.2-6%)	6.6 (5-8.6)	17% (13-22%)	27% (19-36%)	850 (600-1100)	63000 (47000-79000)
France	1995	Male	30 to 34	97% (95-98%)	3.4% (1.9-5.4%)	6.7 (5.1-8.6)	15% (11-19%)	23% (17-29%)	910 (660-1200)	62000 (47000-77000)
France	1995	Male	35 to 39	96% (94-98%)	3.5% (1.9-5.6%)	7 (5.5-8.9)	17% (13-20%)	25% (20-31%)	1300 (990-1600)	76000 (61000-92000)
France	1995	Male	40 to 44	97% (94-98%)	3.4% (1.9-5.6%)	7.9 (6.3-10)	20% (17-24%)	30% (25-35%)	2100 (1800-2500)	110000 (92000-130000)
France	1995	Male	45 to 49	97% (95-98%)	3.2% (1.9-5%)	9.4 (7.5-12)	22% (19-25%)	31% (27-35%)	3200 (2800-3600)	140000 (130000-160000)
France	1995	Male	50 to 54	97% (95-98%)	3% (1.9-4.8%)	10 (8.1-13)	22% (19-25%)	30% (26-33%)	3000 (2600-3300)	120000 (110000-130000)
France	1995	Male	55 to 59	97% (96-98%)	2.9% (1.9-4.3%)	10 (8.5-13)	22% (19-24%)	28% (24-31%)	3900 (3500-4400)	130000 (120000-150000)
France	1995	Male	60 to 64	97% (95-98%)	3% (1.8-4.5%)	10 (8.3-13)	20% (17-23%)	25% (21-28%)	5400 (4700-6000)	160000 (140000-180000)
France	1995	Male	65 to 69	97% (95-98%)	3.3% (2-5%)	9.6 (7.8-12)	18% (15-21%)	22% (18-25%)	6200 (5300-7200)	150000 (130000-170000)
France	1995	Male	70 to 74	96% (95-98%)	3.5% (2.2-5.1%)	9 (7-11)	16% (13-19%)	19% (16-22%)	6900 (5700-8100)	140000 (110000-160000)
France	1995	Male	75 to 79	96% (94-98%)	3.8% (2.1-6.2%)	8.3 (6.5-10)	14% (11-17%)	16% (13-20%)	4400 (3400-5400)	68000 (54000-82000)
France	1995	Male	80 to 84	95% (91-97%)	5.2% (3-8.5%)	7.5 (5.8-9.1)	13% (9.8-16%)	15% (11-18%)	6100 (4500-7600)	69000 (52000-86000)
France	1995	Male	85 to 89	93% (87-96%)	7.4% (4.1-13%)	6.6 (5.1-8.3)	11% (7.7-15%)	13% (8.2-16%)	4300 (2900-5600)	35000 (24000-45000)
France	1995	Male	90 to 94	92% (86-96%)	8.1% (4.4-14%)	6.1 (4.5-8)	9.7% (5.7-13%)	11% (5.9-15%)	1700 (960-2500)	9700 (5600-14000)
France	1995	Male	95 plus	91% (84-96%)	8.7% (3.9-16%)	5.6 (3.8-7.8)	8.5% (3.5-13%)	9.1% (3.4-15%)	360 (130-570)	1300 (550-2100)
France	2000	Female	All Ages	86% (83-89%)	14% (11-17%)	2.9 (2.4-3.4)	5.4% (4-7%)	6.4% (3.8-9.8%)	17000 (9900-25000)	400000 (300000-520000)
France	2000	Female	15 to 19	85% (76-92%)	15% (8.5-24%)	2.3 (1.6-3.1)	2.9% (1.7-4.3%)	10% (5.1-16%)	57 (28-86)	6000 (3600-8700)
France	2000	Female	20 to 24	90% (84-94%)	10% (6-16%)	2.7 (1.8-3.9)	3.9% (2.6-5.4%)	12% (6-18%)	77 (39-110)	9000 (5800-12000)
France	2000	Female	25 to 29	91% (86-95%)	8.7% (5-14%)	2.6 (1.9-3.5)	3.4% (2.3-4.6%)	11% (6.2-17%)	89 (48-130)	9500 (6200-13000)
France	2000	Female	30 to 34	90% (85-94%)	9.8% (6.1-15%)	2.7 (1.9-3.7)	3.6% (2.4-5%)	12% (7.4-17%)	140 (86-200)	12000 (8000-16000)
France	2000	Female	35 to 39	87% (82-92%)	13% (7.9-18%)	3 (2.2-4.1)	4.9% (3.5-6.5%)	15% (11-19%)	290 (210-370)	19000 (14000-24000)
France	2000	Female	40 to 44	87% (81-91%)	13% (8.8-19%)	3.5 (2.6-4.6)	6.5% (4.9-8.4%)	17% (14-21%)	510 (400-620)	28000 (22000-34000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	2000	Female	45 to 49	88% (84-92%)	12% (8.1-16%)	4.1 (3.1-5.4)	8.2% (6.4-10%)	19% (15-22%)	820 (670-990)	39000 (31000-47000)
France	2000	Female	50 to 54	90% (86-94%)	9.8% (6.4-14%)	4.5 (3.4-5.9)	9% (7-11%)	19% (15-22%)	1100 (900-1300)	45000 (37000-55000)
France	2000	Female	55 to 59	92% (89-95%)	8% (5.4-11%)	4.7 (3.6-6.2)	9% (7-11%)	17% (14-20%)	940 (780-1100)	34000 (27000-42000)
France	2000	Female	60 to 64	92% (89-95%)	7.7% (4.9-11%)	4.7 (3.5-6.2)	8.3% (6-11%)	15% (12-18%)	1100 (890-1400)	35000 (26000-45000)
France	2000	Female	65 to 69	91% (86-94%)	9.1% (5.9-14%)	4.4 (3.4-5.5)	7.1% (5.1-9.3%)	12% (8.6-15%)	1400 (1100-1800)	38000 (27000-50000)
France	2000	Female	70 to 74	88% (84-91%)	12% (8.5-16%)	3.9 (2.8-5.1)	6% (3.5-8.7%)	8.9% (5.4-13%)	1800 (1100-2600)	38000 (22000-55000)
France	2000	Female	75 to 79	85% (78-90%)	15% (9.8-22%)	3.5 (2.5-4.8)	4.9% (2-8%)	6.5% (2.5-11%)	2200 (850-3600)	36000 (15000-59000)
France	2000	Female	80 to 84	81% (70-88%)	19% (12-30%)	3.3 (2.4-4.3)	4.1% (1.6-7%)	4.9% (1.8-8.7%)	1500 (570-2700)	19000 (7100-32000)
France	2000	Female	85 to 89	78% (68-86%)	22% (14-32%)	3.1 (2.2-4.2)	3.6% (0.88-6.8%)	4% (0.73-8%)	2300 (420-4700)	20000 (5000-39000)
France	2000	Female	90 to 94	77% (67-86%)	23% (14-33%)	3 (1.9-4.3)	3% (-0.088-6.8%)	3.2% (-0.37-7.8%)	1600 (-180-3800)	9600 (-270-22000)
France	2000	Female	95 plus	77% (63-88%)	23% (12-37%)	3 (1.6-4.6)	2.8% (-0.92-7.7%)	2.9% (-1.4-8.7%)	690 (-340-2000)	2800 (-880-7400)
France	2000	Male	All Ages	94% (92-95%)	6.4% (4.8-8.2%)	5.6 (4.7-6.6)	15% (13-18%)	17% (14-20%)	46000 (37000-54000)	1300000 (1100000-1500000)
France	2000	Male	15 to 19	89% (83-94%)	11% (6-17%)	4 (2.9-5.2)	9.2% (5.7-13%)	19% (10-26%)	250 (140-350)	22000 (14000-29000)
France	2000	Male	20 to 24	95% (91-97%)	5.2% (2.9-8.7%)	6 (4.4-7.9)	17% (12-22%)	28% (19-37%)	560 (370-750)	46000 (33000-59000)
France	2000	Male	25 to 29	96% (94-98%)	3.9% (2.2-6.2%)	5.8 (4.4-7.4)	16% (12-20%)	27% (18-36%)	630 (420-850)	49000 (36000-63000)
France	2000	Male	30 to 34	97% (94-98%)	3.5% (1.9-5.7%)	5.8 (4.4-7.5)	15% (11-19%)	26% (19-34%)	720 (510-940)	51000 (37000-64000)
France	2000	Male	35 to 39	96% (94-98%)	3.6% (1.9-5.7%)	6.2 (4.6-7.8)	16% (12-20%)	27% (20-34%)	1100 (800-1300)	66000 (51000-81000)
France	2000	Male	40 to 44	97% (94-98%)	3.5% (1.8-5.5%)	7 (5.4-8.7)	19% (15-23%)	29% (24-35%)	1800 (1500-2200)	96000 (78000-110000)
France	2000	Male	45 to 49	97% (95-98%)	3.3% (1.9-5.2%)	8.3 (6.6-10)	21% (18-25%)	30% (26-34%)	2900 (2500-3400)	130000 (120000-150000)
France	2000	Male	50 to 54	97% (95-98%)	3.1% (1.8-4.8%)	9 (7.2-11)	21% (18-24%)	28% (25-32%)	3800 (3300-4300)	150000 (130000-170000)
France	2000	Male	55 to 59	97% (95-98%)	3% (1.9-4.5%)	9.2 (7.4-11)	20% (17-22%)	26% (22-28%)	3300 (2900-3700)	120000 (100000-130000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	2000	Male	60 to 64	97% (95-98%)	3.2% (1.9-4.8%)	9 (7.1-11)	18% (15-21%)	22% (19-25%)	4000 (3400-4500)	120000 (100000-130000)
France	2000	Male	65 to 69	97% (95-98%)	3.4% (2.1-5.1%)	8.4 (6.7-10)	16% (14-19%)	20% (17-23%)	5200 (4400-5900)	130000 (110000-150000)
France	2000	Male	70 to 74	96% (95-98%)	3.6% (2.3-5.4%)	7.9 (6.2-9.7)	14% (11-17%)	17% (14-20%)	5900 (4700-7100)	120000 (94000-140000)
France	2000	Male	75 to 79	96% (93-98%)	4% (2.2-6.6%)	7.2 (5.7-9.1)	12% (9.3-16%)	14% (11-18%)	6200 (4600-7800)	97000 (72000-120000)
France	2000	Male	80 to 84	95% (91-97%)	5.3% (2.9-8.9%)	6.5 (5.1-8)	11% (7.6-14%)	12% (8.3-16%)	3700 (2400-4800)	42000 (29000-55000)
France	2000	Male	85 to 89	93% (88-96%)	6.9% (3.8-12%)	5.9 (4.6-7.4)	9.4% (5.9-13%)	10% (6.2-14%)	3700 (2200-5200)	30000 (19000-42000)
France	2000	Male	90 to 94	92% (88-96%)	7.6% (4.2-12%)	5.5 (4.1-7)	7.8% (3.9-12%)	8.4% (3.8-13%)	1700 (750-2600)	9400 (4600-14000)
France	2000	Male	95 plus	92% (85-96%)	8.5% (3.9-15%)	5 (3.3-7)	6.6% (2-11%)	6.9% (1.5-12%)	380 (82-680)	1500 (440-2500)
France	2005	Female	All Ages	85% (83-88%)	15% (12-17%)	2.7 (2.2-3.3)	4.9% (3.7-6.4%)	5.7% (3.3-8.8%)	15000 (8500-23000)	360000 (270000-460000)
France	2005	Female	15 to 19	82% (73-90%)	18% (9.8-27%)	2.1 (1.4-2.8)	2.3% (1.4-3.4%)	9.1% (4.6-14%)	38 (19-59)	4800 (2900-6900)
France	2005	Female	20 to 24	89% (82-94%)	11% (6.4-18%)	2.6 (1.6-3.7)	3.5% (2.3-4.7%)	11% (5.5-16%)	59 (30-88)	8100 (5300-11000)
France	2005	Female	25 to 29	91% (86-95%)	9.1% (5.4-14%)	2.5 (1.8-3.4)	3% (2-4.1%)	10% (5.4-15%)	60 (32-89)	7400 (5000-10000)
France	2005	Female	30 to 34	89% (84-93%)	11% (6.6-16%)	2.5 (1.8-3.5)	3.2% (2.1-4.3%)	11% (6.8-16%)	110 (65-150)	9600 (6600-13000)
France	2005	Female	35 to 39	87% (81-92%)	13% (8.1-19%)	2.9 (2-3.8)	4.3% (3-5.8%)	14% (9.5-18%)	230 (150-300)	15000 (11000-20000)
France	2005	Female	40 to 44	86% (81-91%)	14% (9-19%)	3.3 (2.4-4.3)	5.7% (4.2-7.3%)	16% (12-19%)	430 (320-530)	24000 (18000-30000)
France	2005	Female	45 to 49	88% (83-91%)	12% (8.7-17%)	3.9 (2.8-5.1)	7.4% (5.7-9.4%)	17% (14-21%)	720 (580-880)	34000 (27000-43000)
France	2005	Female	50 to 54	90% (85-93%)	10% (6.9-15%)	4.3 (3.2-5.6)	8.4% (6.5-10%)	17% (14-21%)	1000 (830-1200)	43000 (34000-52000)
France	2005	Female	55 to 59	92% (88-94%)	8.4% (5.8-12%)	4.5 (3.4-5.9)	8.3% (6.4-10%)	16% (13-19%)	1200 (1000-1500)	44000 (35000-55000)
France	2005	Female	60 to 64	92% (88-94%)	8.4% (5.6-12%)	4.4 (3.3-5.7)	7.6% (5.7-9.8%)	14% (11-17%)	1000 (790-1200)	31000 (24000-40000)
France	2005	Female	65 to 69	90% (86-93%)	10% (6.8-14%)	4.1 (3.2-5.2)	6.4% (4.6-8.4%)	11% (8.3-13%)	1100 (870-1400)	30000 (22000-39000)
France	2005	Female	70 to 74	87% (83-91%)	13% (9.3-17%)	3.6 (2.6-4.8)	5.3% (3.1-7.6%)	8.1% (5.2-11%)	1500 (940-2000)	31000 (18000-45000)
France	2005	Female	75 to 79	83% (76-89%)	17% (11-24%)	3.2 (2.3-4.3)	4.2% (1.9-7%)	5.8% (2.6-9.3%)	1700 (760-2800)	29000 (13000-48000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	2005	Female	80 to 84	79% (69-86%)	21% (14-31%)	2.9 (2.1-3.8)	3.5% (1.2-6.2%)	4.3% (1.4-7.8%)	2100 (650-3700)	26000 (9300-46000)
France	2005	Female	85 to 89	76% (67-84%)	24% (16-33%)	2.8 (2-3.8)	3.1% (0.77-5.9%)	3.5% (0.64-6.9%)	1400 (250-2700)	12000 (3000-23000)
France	2005	Female	90 to 94	75% (65-84%)	25% (16-35%)	2.8 (1.8-3.9)	2.6% (-0.017-6%)	2.8% (-0.37-6.9%)	1400 (-180-3400)	8500 (-56-20000)
France	2005	Female	95 plus	75% (60-86%)	25% (14-40%)	2.7 (1.6-4.2)	2.5% (-0.75-6.8%)	2.5% (-1.3-7.7%)	720 (-380-2200)	2900 (-910-8000)
France	2005	Male	All Ages	94% (92-95%)	6.3% (4.8-8.2%)	5.3 (4.4-6.4)	14% (12-16%)	16% (13-18%)	42000 (34000-50000)	1200000 (990000-1300000)
France	2005	Male	15 to 19	89% (83-94%)	11% (6-17%)	3.7 (2.7-5)	7.8% (5.1-11%)	17% (10-25%)	180 (110-260)	17000 (11000-23000)
France	2005	Male	20 to 24	95% (91-97%)	5.3% (3-8.9%)	5.6 (4.1-7.4)	15% (11-20%)	26% (17-35%)	450 (300-610)	39000 (28000-51000)
France	2005	Male	25 to 29	96% (94-98%)	4% (2.2-6.4%)	5.5 (4.1-7.1)	14% (11-18%)	26% (18-35%)	440 (290-590)	37000 (27000-47000)
France	2005	Male	30 to 34	96% (94-98%)	3.6% (2-5.9%)	5.5 (4.2-7.1)	14% (10-18%)	26% (17-33%)	590 (400-770)	43000 (32000-55000)
France	2005	Male	35 to 39	96% (94-98%)	3.7% (1.9-5.9%)	5.8 (4.4-7.4)	15% (11-18%)	26% (19-33%)	870 (630-1100)	55000 (42000-68000)
France	2005	Male	40 to 44	96% (94-98%)	3.5% (1.8-5.9%)	6.6 (5.1-8.3)	17% (14-21%)	28% (22-33%)	1500 (1200-1800)	80000 (64000-96000)
France	2005	Male	45 to 49	97% (95-98%)	3.3% (1.9-5.3%)	7.8 (6.2-9.7)	20% (17-23%)	29% (25-33%)	2600 (2200-2900)	120000 (100000-130000)
France	2005	Male	50 to 54	97% (95-98%)	3.2% (1.9-4.9%)	8.5 (6.7-10)	20% (17-23%)	28% (24-31%)	3600 (3100-4000)	140000 (130000-160000)
France	2005	Male	55 to 59	97% (96-98%)	3.1% (1.9-4.5%)	8.7 (7-11)	19% (16-21%)	24% (21-27%)	4300 (3800-4900)	150000 (130000-170000)
France	2005	Male	60 to 64	97% (95-98%)	3.2% (1.9-4.7%)	8.4 (6.7-11)	17% (14-19%)	21% (18-24%)	3400 (2900-3900)	100000 (88000-120000)
France	2005	Male	65 to 69	97% (95-98%)	3.4% (2.1-5%)	7.9 (6.5-9.6)	15% (12-17%)	18% (15-21%)	3900 (3300-4600)	99000 (83000-110000)
France	2005	Male	70 to 74	96% (95-98%)	3.6% (2.3-5.2%)	7.4 (5.9-9.1)	13% (10-16%)	16% (13-19%)	4900 (3900-6000)	100000 (79000-120000)
France	2005	Male	75 to 79	96% (93-98%)	4.1% (2.1-6.7%)	6.8 (5.3-8.6)	11% (8.2-14%)	13% (9.6-17%)	5300 (3900-6700)	84000 (62000-110000)
France	2005	Male	80 to 84	95% (91-97%)	5.2% (2.9-8.8%)	6.2 (4.9-7.7)	10% (6.7-13%)	11% (7.4-15%)	5300 (3500-7000)	62000 (42000-80000)
France	2005	Male	85 to 89	93% (90-96%)	6.5% (3.9-10%)	5.6 (4.4-7.1)	8.4% (5-12%)	9.2% (5.2-13%)	2500 (1400-3500)	20000 (12000-28000)
France	2005	Male	90 to 94	93% (88-96%)	7.3% (4-12%)	5.2 (3.9-6.7)	6.9% (3-11%)	7.4% (2.9-12%)	1600 (630-2500)	9100 (4100-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	2005	Male	95 plus	92% (86-96%)	8.1% (3.8-14%)	4.7 (3.2-6.6)	5.8% (1.5-11%)	6% (1.1-12%)	420 (76-820)	1600 (420-3000)
France	2010	Female	All Ages	86% (83-88%)	14% (12-17%)	2.6 (2.1-3.2)	4.4% (3.1-5.9%)	5.2% (2.7-8%)	14000 (7100-22000)	330000 (240000-440000)
France	2010	Female	15 to 19	85% (77-92%)	15% (8.3-23%)	2 (1.4-2.8)	2.1% (1.3-3%)	8.9% (4.3-14%)	29 (14-44)	4000 (2500-5800)
France	2010	Female	20 to 24	90% (84-94%)	10% (5.7-16%)	2.5 (1.5-3.5)	3.3% (2.3-4.4%)	10% (5.1-15%)	47 (24-71)	7400 (4900-10000)
France	2010	Female	25 to 29	91% (86-95%)	8.8% (5-14%)	2.4 (1.6-3.3)	2.9% (2-3.9%)	9.4% (5-14%)	54 (29-82)	7300 (4800-9900)
France	2010	Female	30 to 34	90% (84-94%)	10% (6.4-16%)	2.4 (1.6-3.3)	2.8% (1.9-3.8%)	11% (6.5-15%)	79 (47-110)	7700 (5300-10000)
France	2010	Female	35 to 39	87% (81-92%)	13% (8-19%)	2.7 (1.9-3.8)	3.7% (2.6-5%)	13% (9.1-17%)	180 (120-240)	13000 (9200-17000)
France	2010	Female	40 to 44	86% (81-91%)	14% (9.1-19%)	3.1 (2.2-4.3)	5% (3.6-6.5%)	15% (11-19%)	350 (260-440)	20000 (15000-25000)
France	2010	Female	45 to 49	88% (83-91%)	12% (8.5-17%)	3.6 (2.7-4.9)	6.4% (4.9-8.2%)	16% (13-19%)	620 (490-760)	30000 (24000-37000)
France	2010	Female	50 to 54	90% (86-93%)	10% (6.8-14%)	4 (3-5.4)	7.4% (5.7-9.4%)	16% (13-19%)	910 (730-1100)	38000 (30000-47000)
France	2010	Female	55 to 59	92% (89-94%)	7.9% (5.5-11%)	4.2 (3.1-5.7)	7.5% (5.6-9.5%)	15% (12-18%)	1200 (930-1400)	42000 (32000-52000)
France	2010	Female	60 to 64	92% (89-95%)	7.7% (5-11%)	4.2 (3-5.7)	6.9% (5.1-8.8%)	13% (10-16%)	1300 (1000-1600)	41000 (30000-53000)
France	2010	Female	65 to 69	91% (87-94%)	9.4% (6.2-13%)	3.9 (2.9-5.1)	5.8% (4.2-7.8%)	10% (7.8-13%)	1000 (760-1300)	26000 (19000-35000)
France	2010	Female	70 to 74	88% (84-91%)	12% (8.8-16%)	3.5 (2.5-4.8)	4.8% (2.8-7.1%)	7.8% (4.8-11%)	1100 (710-1600)	25000 (14000-36000)
France	2010	Female	75 to 79	84% (77-90%)	16% (10-23%)	3.1 (2.2-4.4)	3.8% (1.5-6.4%)	5.5% (2.5-8.8%)	1400 (630-2300)	24000 (9300-41000)
France	2010	Female	80 to 84	80% (71-87%)	20% (13-29%)	2.9 (2.1-3.9)	3.1% (0.87-5.6%)	4% (1-7.4%)	1800 (470-3300)	22000 (6000-41000)
France	2010	Female	85 to 89	78% (69-85%)	22% (15-31%)	2.8 (1.9-3.8)	2.8% (0.46-5.5%)	3.2% (0.39-6.7%)	2100 (250-4300)	18000 (3000-36000)
France	2010	Female	90 to 94	77% (67-86%)	23% (14-33%)	2.7 (1.7-3.9)	2.3% (-0.36-5.6%)	2.5% (-0.68-6.5%)	940 (-260-2500)	5900 (-920-15000)
France	2010	Female	95 plus	77% (63-88%)	23% (12-37%)	2.6 (1.5-4.2)	2.1% (-1.1-6.5%)	2.1% (-1.7-7.4%)	740 (-610-2600)	3000 (-1600-9400)
France	2010	Male	All Ages	94% (92-95%)	6.2% (4.7-8%)	5.1 (4.1-6.3)	13% (11-15%)	14% (11-17%)	39000 (30000-47000)	1100000 (870000-1300000)
France	2010	Male	15 to 19	91% (86-95%)	8.8% (5-14%)	3.6 (2.5-4.9)	7.1% (4.6-9.8%)	17% (9.5-24%)	140 (79-200)	14000 (9000-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	2010	Male	20 to 24	95% (92-97%)	5% (2.8-8.2%)	5.3 (3.9-7.1)	14% (10-19%)	25% (16-34%)	360 (230-490)	34000 (24000-43000)
France	2010	Male	25 to 29	96% (94-98%)	4.1% (2.3-6.4%)	5.2 (3.9-6.8)	13% (9.9-17%)	25% (16-33%)	400 (260-530)	35000 (25000-44000)
France	2010	Male	30 to 34	96% (94-98%)	3.8% (2-6.1%)	5.2 (3.9-6.8)	12% (9.2-16%)	25% (17-32%)	450 (310-590)	35000 (25000-44000)
France	2010	Male	35 to 39	96% (94-98%)	3.7% (2-5.8%)	5.4 (4-7.1)	13% (9.9-17%)	25% (18-32%)	750 (530-960)	49000 (37000-61000)
France	2010	Male	40 to 44	96% (94-98%)	3.6% (1.9-5.8%)	6.2 (4.6-8)	15% (12-19%)	27% (21-33%)	1200 (940-1500)	68000 (53000-82000)
France	2010	Male	45 to 49	97% (94-98%)	3.4% (2-5.5%)	7.3 (5.7-9.4)	18% (15-21%)	28% (23-33%)	2100 (1700-2400)	99000 (82000-110000)
France	2010	Male	50 to 54	97% (95-98%)	3.2% (1.9-5%)	8 (6.1-10)	19% (16-22%)	27% (22-30%)	3100 (2600-3600)	130000 (110000-140000)
France	2010	Male	55 to 59	97% (96-98%)	3% (1.9-4.4%)	8.2 (6.4-10)	18% (15-20%)	23% (20-27%)	4000 (3400-4600)	140000 (120000-160000)
France	2010	Male	60 to 64	97% (95-98%)	3.1% (1.9-4.6%)	8 (6.1-10)	16% (13-18%)	20% (17-23%)	4400 (3700-5100)	130000 (110000-150000)
France	2010	Male	65 to 69	97% (95-98%)	3.3% (2-4.9%)	7.5 (5.9-9.4)	14% (11-16%)	17% (14-20%)	3400 (2700-3900)	86000 (69000-100000)
France	2010	Male	70 to 74	96% (95-98%)	3.5% (2.2-5.2%)	7 (5.4-9)	12% (8.9-15%)	15% (11-18%)	3800 (2900-4600)	79000 (60000-97000)
France	2010	Male	75 to 79	96% (93-98%)	4% (2.1-6.6%)	6.4 (4.9-8.3)	10% (7.2-13%)	12% (8.4-16%)	4500 (3100-5700)	73000 (50000-94000)
France	2010	Male	80 to 84	95% (92-97%)	5% (2.9-8.2%)	5.9 (4.6-7.4)	8.9% (5.7-12%)	10% (6.2-14%)	4600 (2800-6300)	55000 (35000-75000)
France	2010	Male	85 to 89	94% (91-96%)	6.1% (3.7-9.3%)	5.4 (4.2-6.9)	7.5% (4.1-11%)	8.2% (4.2-12%)	3800 (1900-5600)	32000 (17000-46000)
France	2010	Male	90 to 94	93% (89-96%)	6.9% (3.8-11%)	4.9 (3.7-6.6)	6% (2.4-9.8%)	6.5% (2-11%)	1200 (370-2000)	7000 (2700-11000)
France	2010	Male	95 plus	92% (87-96%)	7.8% (3.6-13%)	4.5 (3-6.3)	5% (1.1-9.7%)	5.2% (0.51-11%)	470 (45-960)	1800 (380-3500)
France	2016	Female	All Ages	87% (84-89%)	13% (11-16%)	2.6 (1.9-3.5)	4.1% (2.6-5.8%)	4.6% (1.7-8.1%)	14000 (4900-24000)	320000 (200000-450000)
France	2016	Female	15 to 19	86% (76-92%)	14% (7.9-24%)	2 (1.2-2.9)	1.9% (1.3-2.8%)	8.6% (4.3-13%)	25 (12-39)	3700 (2300-5400)
France	2016	Female	20 to 24	90% (83-94%)	10% (5.6-17%)	2.4 (1.4-3.5)	3.1% (2.1-4.1%)	9.9% (5.1-15%)	36 (19-56)	6200 (4100-8800)
France	2016	Female	25 to 29	91% (85-95%)	9% (4.9-15%)	2.3 (1.4-3.3)	2.7% (1.8-3.7%)	9.3% (4.9-14%)	47 (25-73)	6600 (4400-9200)
France	2016	Female	30 to 34	90% (84-94%)	10% (6-16%)	2.3 (1.5-3.4)	2.6% (1.8-3.6%)	10% (6.3-15%)	74 (45-110)	7400 (5000-10000)
France	2016	Female	35 to 39	87% (81-92%)	13% (8-19%)	2.6 (1.6-3.8)	3.5% (2.4-4.8%)	13% (8.7-17%)	140 (96-190)	10000 (7200-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	2016	Female	40 to 44	86% (80-91%)	14% (8.8-20%)	3 (2-4.3)	4.6% (3.2-6.1%)	15% (11-19%)	310 (230-410)	18000 (13000-24000)
France	2016	Female	45 to 49	88% (83-92%)	12% (8.1-17%)	3.5 (2.4-5)	6% (4.3-8%)	16% (12-19%)	550 (420-690)	27000 (20000-34000)
France	2016	Female	50 to 54	90% (86-94%)	9.5% (6.3-14%)	3.9 (2.7-5.5)	6.9% (5.1-9.1%)	15% (12-19%)	840 (650-1100)	36000 (27000-46000)
France	2016	Female	55 to 59	93% (90-95%)	7.2% (5-9.9%)	4.2 (2.8-5.8)	7.1% (5-9.3%)	14% (11-18%)	1100 (810-1300)	39000 (28000-50000)
France	2016	Female	60 to 64	93% (91-96%)	6.7% (4.3-9.5%)	4.2 (2.8-5.9)	6.5% (4.4-8.9%)	12% (9.2-16%)	1200 (910-1600)	39000 (26000-53000)
France	2016	Female	65 to 69	92% (88-95%)	7.9% (5.3-12%)	3.9 (2.7-5.4)	5.4% (3.4-7.9%)	9.9% (7-13%)	1400 (980-1900)	36000 (23000-52000)
France	2016	Female	70 to 74	90% (86-93%)	10% (7.2-14%)	3.5 (2.3-5.1)	4.4% (2.2-7.1%)	7.5% (4.3-11%)	1000 (580-1500)	21000 (11000-34000)
France	2016	Female	75 to 79	87% (81-92%)	13% (8.4-19%)	3.2 (2-4.7)	3.5% (1-6.4%)	5.2% (2-9%)	1200 (450-2000)	20000 (5900-36000)
France	2016	Female	80 to 84	84% (76-90%)	16% (10-24%)	3 (2-4.3)	2.8% (0.36-5.8%)	3.7% (0.51-7.6%)	1500 (220-3100)	20000 (2400-40000)
France	2016	Female	85 to 89	82% (74-88%)	18% (12-26%)	2.9 (1.9-4.1)	2.5% (-0.086-5.5%)	3% (-0.24-6.7%)	1900 (-160-4300)	17000 (-590-37000)
France	2016	Female	90 to 94	81% (72-88%)	19% (12-28%)	2.8 (1.8-4.2)	2% (-0.73-5.5%)	2.2% (-1.2-6.4%)	1600 (-910-4800)	10000 (-3700-27000)
France	2016	Female	95 plus	80% (68-90%)	20% (10-32%)	2.7 (1.5-4.4)	1.9% (-1.4-6.4%)	1.9% (-2-7.2%)	570 (-600-2100)	2300 (-1700-7600)
France	2016	Male	All Ages	93% (91-95%)	6.8% (5.1-8.8%)	4.9 (3.5-6.6)	12% (9.3-15%)	13% (8.9-17%)	38000 (26000-50000)	1000000 (780000-1200000)
France	2016	Male	15 to 19	89% (81-94%)	11% (5.8-19%)	3.4 (2.1-4.9)	6.1% (4-8.6%)	16% (8.7-24%)	110 (60-170)	12000 (7400-17000)
France	2016	Male	20 to 24	94% (89-97%)	6.2% (3.5-11%)	5 (3.4-7.1)	13% (9.3-17%)	24% (15-34%)	280 (170-390)	27000 (19000-36000)
France	2016	Male	25 to 29	95% (91-97%)	5% (2.8-8.6%)	4.9 (3.4-6.8)	13% (9-17%)	24% (15-33%)	340 (210-470)	30000 (21000-40000)
France	2016	Male	30 to 34	96% (92-98%)	4.4% (2.3-8.2%)	4.9 (3.4-6.9)	12% (8.6-16%)	24% (15-33%)	410 (270-580)	32000 (23000-42000)
France	2016	Male	35 to 39	96% (92-98%)	4.4% (2.2-7.5%)	5.2 (3.5-7.3)	13% (9.2-17%)	25% (17-33%)	560 (380-740)	38000 (27000-48000)
France	2016	Male	40 to 44	96% (92-98%)	4.2% (2.1-7.5%)	5.8 (4-8.1)	15% (11-18%)	26% (20-33%)	1100 (790-1400)	61000 (46000-75000)
France	2016	Male	45 to 49	96% (93-98%)	3.8% (2.1-6.6%)	7 (5-9.7)	17% (14-21%)	28% (22-33%)	1800 (1400-2300)	88000 (69000-110000)
France	2016	Male	50 to 54	97% (95-98%)	3.4% (2-5.5%)	7.7 (5.4-11)	18% (15-21%)	26% (21-31%)	2800 (2300-3400)	120000 (94000-140000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
France	2016	Male	55 to 59	97% (95-98%)	3.2% (2-4.8%)	7.8 (5.6-11)	17% (14-20%)	23% (19-27%)	3600 (2800-4200)	130000 (100000-150000)
France	2016	Male	60 to 64	97% (95-98%)	3.2% (2-5%)	7.7 (5.5-11)	15% (12-18%)	20% (16-23%)	4100 (3300-4900)	120000 (99000-150000)
France	2016	Male	65 to 69	97% (95-98%)	3.5% (2.1-5.2%)	7.1 (5.1-9.9)	13% (9.6-16%)	17% (13-20%)	4700 (3600-5700)	120000 (91000-150000)
France	2016	Male	70 to 74	96% (94-98%)	3.8% (2.3-5.8%)	6.7 (4.7-9.2)	11% (7.9-14%)	14% (10-18%)	3400 (2400-4400)	72000 (52000-93000)
France	2016	Male	75 to 79	96% (93-98%)	4.2% (2.3-6.9%)	6.1 (4.3-8.4)	9.5% (6.1-13%)	11% (7.3-16%)	3800 (2400-5200)	62000 (39000-84000)
France	2016	Male	80 to 84	95% (91-97%)	5.2% (2.9-8.9%)	5.6 (4-7.8)	8.1% (4.5-12%)	9.3% (4.9-14%)	4400 (2300-6600)	53000 (29000-77000)
France	2016	Male	85 to 89	93% (89-96%)	6.6% (3.7-11%)	5.1 (3.5-7.1)	6.8% (3.1-11%)	7.6% (3.2-12%)	3700 (1600-6100)	32000 (14000-51000)
France	2016	Male	90 to 94	93% (88-96%)	7.3% (4-12%)	4.7 (3.2-6.6)	5.5% (1.8-9.7%)	5.9% (1.6-11%)	2300 (580-4200)	13000 (4200-24000)
France	2016	Male	95 plus	92% (85-96%)	8.1% (3.7-15%)	4.3 (2.7-6.5)	4.5% (0.62-9.6%)	4.7% (0.13-11%)	390 (10-900)	1500 (210-3300)
Germany	1990	Female	All Ages	90% (88-92%)	9.6% (8-12%)	3.6 (3.2-4)	6.8% (4.4-9.6%)	8.1% (3.1-14%)	40000 (15000-67000)	880000 (570000-1200000)
Germany	1990	Female	15 to 19	90% (84-95%)	9.6% (5.1-16%)	2.8 (2-3.8)	3.8% (2.2-5.3%)	12% (6.1-19%)	86 (42-130)	9000 (5400-13000)
Germany	1990	Female	20 to 24	93% (89-96%)	6.6% (3.9-11%)	3.1 (2-4.4)	4.5% (3.1-6.1%)	12% (6.2-17%)	140 (74-210)	18000 (12000-25000)
Germany	1990	Female	25 to 29	95% (91-97%)	5.4% (3.2-8.9%)	2.9 (2-3.8)	3.9% (2.7-5.2%)	11% (6.5-16%)	160 (94-230)	18000 (13000-25000)
Germany	1990	Female	30 to 34	95% (92-97%)	4.7% (2.5-8.1%)	3.1 (2.3-4.2)	4.6% (3.3-6.1%)	14% (9.8-18%)	270 (190-350)	21000 (15000-28000)
Germany	1990	Female	35 to 39	96% (93-98%)	4.4% (2.4-7.2%)	3.9 (2.7-5.3)	6.5% (4.8-8.3%)	18% (14-23%)	500 (370-630)	32000 (24000-41000)
Germany	1990	Female	40 to 44	95% (93-97%)	4.6% (2.7-7.3%)	4.5 (3.3-5.9)	8.1% (6.2-10%)	20% (16-24%)	730 (580-890)	40000 (32000-50000)
Germany	1990	Female	45 to 49	95% (92-97%)	5.3% (3.2-8.4%)	5 (3.6-6.6)	9.3% (7.1-12%)	20% (16-24%)	1300 (990-1500)	60000 (47000-73000)
Germany	1990	Female	50 to 54	94% (91-96%)	5.9% (3.9-8.8%)	5.3 (4-6.8)	10% (7.5-13%)	19% (15-23%)	2000 (1600-2500)	83000 (64000-100000)
Germany	1990	Female	55 to 59	94% (91-96%)	6.2% (4.1-9.4%)	5.7 (4.4-7.3)	10% (7.4-14%)	17% (13-23%)	2300 (1700-3000)	80000 (58000-110000)
Germany	1990	Female	60 to 64	92% (88-94%)	8.4% (5.6-12%)	5.6 (4.2-7.4)	9.8% (6.2-14%)	15% (9.4-21%)	3100 (1900-4400)	94000 (57000-130000)
Germany	1990	Female	65 to 69	88% (83-92%)	12% (7.9-17%)	5 (4-6.3)	8.6% (5-13%)	12% (6.8-17%)	4300 (2400-6300)	110000 (63000-160000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	1990	Female	70 to 74	85% (79-90%)	15% (10-21%)	4.5 (3.3-6)	7.5% (3-12%)	9.7% (3.6-16%)	3700 (1400-6300)	74000 (29000-130000)
Germany	1990	Female	75 to 79	83% (74-89%)	17% (11-26%)	4.2 (2.9-5.7)	6.3% (0.95-12%)	7.5% (0.84-15%)	6500 (720-13000)	100000 (15000-200000)
Germany	1990	Female	80 to 84	81% (71-88%)	19% (12-29%)	4 (3-5.3)	5.6% (0.68-11%)	6.3% (0.29-13%)	7200 (330-15000)	83000 (10000-170000)
Germany	1990	Female	85 to 89	80% (69-87%)	20% (13-31%)	3.9 (2.8-5.2)	4.8% (-0.3-11%)	5.3% (-0.89-12%)	5000 (-840-12000)	41000 (-2600-91000)
Germany	1990	Female	90 to 94	79% (69-87%)	21% (13-31%)	3.8 (2.6-5.4)	4.1% (-1.2-11%)	4.4% (-2-12%)	2000 (-920-5400)	11000 (-3400-29000)
Germany	1990	Female	95 plus	79% (66-89%)	21% (11-34%)	3.8 (2.2-5.7)	3.9% (-2.5-12%)	4.1% (-3.4-14%)	500 (-420-1700)	1900 (-1200-5800)
Germany	1990	Male	All Ages	94% (92-96%)	6% (4.5-7.8%)	5 (4.4-5.6)	15% (12-17%)	16% (12-20%)	67000 (50000-85000)	1900000 (1600000-2200000)
Germany	1990	Male	15 to 19	90% (83-95%)	10% (5.4-17%)	3.2 (2.2-4.2)	8.8% (5.5-12%)	16% (8.1-23%)	280 (150-410)	26000 (16000-36000)
Germany	1990	Male	20 to 24	94% (89-97%)	6.2% (3.5-11%)	4.5 (3.2-5.9)	15% (11-19%)	21% (13-29%)	740 (440-1000)	73000 (50000-97000)
Germany	1990	Male	25 to 29	96% (92-97%)	4.4% (2.5-7.5%)	4.6 (3.4-6.1)	14% (11-18%)	21% (14-28%)	840 (550-1100)	80000 (60000-100000)
Germany	1990	Male	30 to 34	97% (94-98%)	3.3% (1.7-6.1%)	4.7 (3.4-6.1)	14% (11-18%)	22% (16-29%)	990 (720-1300)	79000 (59000-98000)
Germany	1990	Male	35 to 39	97% (95-99%)	2.7% (1.4-4.8%)	4.5 (3.2-5.9)	15% (11-18%)	23% (17-28%)	1300 (960-1600)	85000 (66000-100000)
Germany	1990	Male	40 to 44	98% (96-99%)	2.3% (1.2-4.3%)	5 (3.7-6.4)	16% (12-19%)	24% (18-29%)	1700 (1300-2100)	94000 (72000-110000)
Germany	1990	Male	45 to 49	98% (96-99%)	2.2% (1-4.2%)	6.3 (4.6-8)	18% (14-22%)	26% (20-31%)	3200 (2400-4000)	150000 (120000-180000)
Germany	1990	Male	50 to 54	98% (96-99%)	2.4% (1.3-4.3%)	7.8 (6.2-9.5)	20% (16-23%)	26% (21-30%)	6000 (4800-7200)	240000 (190000-280000)
Germany	1990	Male	55 to 59	97% (95-98%)	3% (1.7-5.3%)	9.6 (7.7-12)	20% (16-24%)	25% (20-30%)	7200 (5800-8700)	250000 (200000-290000)
Germany	1990	Male	60 to 64	96% (94-98%)	3.8% (2.1-6.2%)	9.8 (7.7-12)	18% (14-23%)	22% (17-27%)	8600 (6600-11000)	250000 (190000-310000)
Germany	1990	Male	65 to 69	95% (92-98%)	4.7% (2.5-7.8%)	8.4 (6.8-10)	16% (12-20%)	19% (13-23%)	8300 (6000-10000)	200000 (150000-250000)
Germany	1990	Male	70 to 74	94% (91-97%)	5.6% (3.4-8.9%)	7.1 (5.6-8.8)	13% (8.6-18%)	15% (9.4-21%)	5900 (3700-8100)	120000 (75000-160000)
Germany	1990	Male	75 to 79	93% (88-97%)	6.6% (3.3-12%)	6.3 (4.7-8)	11% (5.5-17%)	12% (5.7-18%)	8600 (3900-13000)	130000 (63000-190000)
Germany	1990	Male	80 to 84	92% (85-96%)	8.1% (4.3-15%)	5.9 (4.5-7.4)	10% (4.8-16%)	11% (4.9-17%)	8000 (3500-12000)	89000 (42000-130000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	1990	Male	85 to 89	90% (84-94%)	10% (5.8-16%)	5.5 (4.1-7.1)	9.1% (3.5-14%)	9.8% (3.4-16%)	4400 (1500-7100)	35000 (14000-55000)
Germany	1990	Male	90 to 94	89% (82-94%)	11% (6.1-18%)	5.3 (3.6-7.2)	7.6% (1.3-14%)	8.2% (0.96-15%)	1200 (140-2200)	6600 (1200-12000)
Germany	1990	Male	95 plus	88% (80-95%)	12% (5.3-20%)	5 (2.9-7.3)	6.7% (-0.31-14%)	7.1% (-1.1-15%)	210 (-32-460)	800 (-37-1700)
Germany	1995	Female	All Ages	91% (90-93%)	8.6% (7.2-10%)	3.3 (3-3.7)	6% (3.8-8.4%)	6.5% (2.2-11%)	31000 (10000-54000)	740000 (470000-1000000)
Germany	1995	Female	15 to 19	92% (87-96%)	8.1% (4.4-13%)	2.6 (1.8-3.6)	3.7% (2.3-5.2%)	12% (5.7-18%)	78 (37-120)	8600 (5300-12000)
Germany	1995	Female	20 to 24	95% (91-97%)	5.5% (3.4-8.8%)	2.9 (1.9-4.2)	4.4% (3.1-5.8%)	12% (6.3-17%)	96 (51-140)	13000 (8900-18000)
Germany	1995	Female	25 to 29	95% (93-97%)	4.5% (2.8-6.9%)	2.7 (1.9-3.6)	3.6% (2.6-4.7%)	11% (6.2-15%)	130 (77-180)	16000 (11000-22000)
Germany	1995	Female	30 to 34	96% (94-98%)	3.9% (2.3-6.3%)	2.9 (2.1-3.8)	4% (2.9-5.2%)	13% (9.3-17%)	250 (170-330)	21000 (16000-28000)
Germany	1995	Female	35 to 39	96% (94-98%)	3.7% (2.2-5.8%)	3.6 (2.5-4.9)	5.8% (4.3-7.4%)	18% (13-22%)	480 (360-600)	31000 (24000-39000)
Germany	1995	Female	40 to 44	96% (94-98%)	3.9% (2.5-6.2%)	4.2 (3.1-5.5)	7.7% (5.8-9.6%)	20% (16-23%)	810 (650-980)	44000 (35000-54000)
Germany	1995	Female	45 to 49	95% (93-97%)	4.6% (2.9-7%)	4.7 (3.4-6)	8.8% (6.8-11%)	20% (16-23%)	1100 (860-1300)	50000 (40000-62000)
Germany	1995	Female	50 to 54	95% (93-97%)	5.1% (3.4-7.2%)	5 (3.8-6.3)	9.7% (7.4-12%)	19% (16-23%)	1700 (1400-2000)	69000 (55000-85000)
Germany	1995	Female	55 to 59	95% (92-96%)	5.4% (3.7-7.7%)	5.3 (4.1-6.7)	9.7% (7.2-12%)	17% (14-21%)	2500 (1900-3100)	89000 (68000-110000)
Germany	1995	Female	60 to 64	93% (90-95%)	7.5% (5.2-10%)	5.2 (3.9-6.8)	8.9% (5.7-12%)	14% (9.4-19%)	2500 (1700-3600)	77000 (51000-110000)
Germany	1995	Female	65 to 69	89% (84-92%)	11% (7.6-16%)	4.7 (3.7-5.8)	7.4% (4.2-11%)	11% (6.1-15%)	3100 (1800-4500)	79000 (45000-110000)
Germany	1995	Female	70 to 74	85% (79-90%)	15% (10-21%)	4.2 (3.1-5.5)	6.2% (2.2-10%)	8.2% (2.8-14%)	4100 (1400-7000)	84000 (30000-140000)
Germany	1995	Female	75 to 79	83% (74-89%)	17% (11-26%)	3.9 (2.7-5.2)	4.9% (0.22-10%)	5.9% (-0.048-13%)	3100 (-25-6600)	50000 (2300-100000)
Germany	1995	Female	80 to 84	81% (71-89%)	19% (11-29%)	3.7 (2.7-4.9)	4.2% (-0.22-9.2%)	4.8% (-0.76-11%)	5000 (-790-11000)	59000 (-2900-130000)
Germany	1995	Female	85 to 89	80% (69-87%)	20% (13-31%)	3.6 (2.6-4.8)	3.5% (-0.95-8.6%)	3.8% (-1.7-10%)	4000 (-1800-11000)	34000 (-9400-85000)
Germany	1995	Female	90 to 94	79% (68-87%)	21% (13-32%)	3.5 (2.4-5)	2.9% (-1.8-8.8%)	3% (-2.8-10%)	1700 (-1500-5700)	10000 (-6200-30000)
Germany	1995	Female	95 plus	79% (66-90%)	21% (10-34%)	3.5 (2.1-5.4)	2.7% (-3-10%)	2.7% (-4.1-11%)	460 (-690-1900)	1900 (-2000-6800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	1995	Male	All Ages	95% (94-96%)	4.7% (3.6-6%)	4.7 (4.2-5.2)	14% (12-17%)	15% (11-19%)	62000 (46000-77000)	1800000 (1500000-2100000)
Germany	1995	Male	15 to 19	92% (88-95%)	7.9% (4.6-12%)	3 (2.1-4)	8.7% (5.4-12%)	15% (7.3-22%)	250 (120-370)	24000 (15000-33000)
Germany	1995	Male	20 to 24	95% (93-97%)	4.7% (2.8-7.5%)	4.1 (3-5.5)	15% (11-19%)	20% (12-28%)	510 (290-710)	55000 (38000-72000)
Germany	1995	Male	25 to 29	97% (95-98%)	3.3% (2.1-5.2%)	4.3 (3.2-5.6)	14% (11-17%)	20% (13-27%)	680 (430-930)	72000 (53000-92000)
Germany	1995	Male	30 to 34	98% (96-99%)	2.4% (1.3-3.9%)	4.3 (3.2-5.7)	13% (10-16%)	21% (15-27%)	990 (710-1300)	81000 (61000-100000)
Germany	1995	Male	35 to 39	98% (97-99%)	2% (1.2-3.4%)	4.1 (3-5.5)	14% (11-18%)	23% (17-28%)	1400 (1100-1800)	92000 (72000-110000)
Germany	1995	Male	40 to 44	98% (97-99%)	1.8% (0.95-3.1%)	4.7 (3.4-6)	16% (13-19%)	24% (19-29%)	2000 (1600-2500)	110000 (91000-140000)
Germany	1995	Male	45 to 49	98% (97-99%)	1.6% (0.83-2.8%)	5.8 (4.3-7.6)	18% (14-22%)	26% (20-31%)	2800 (2100-3400)	130000 (100000-160000)
Germany	1995	Male	50 to 54	98% (97-99%)	1.8% (1-2.9%)	7.3 (5.7-9)	19% (16-23%)	26% (21-30%)	4800 (3900-5700)	190000 (160000-230000)
Germany	1995	Male	55 to 59	98% (96-99%)	2.3% (1.3-3.6%)	8.9 (7.2-11)	20% (17-23%)	26% (21-30%)	8200 (6800-9600)	280000 (240000-330000)
Germany	1995	Male	60 to 64	97% (96-98%)	2.9% (1.8-4.4%)	9.1 (7.2-11)	18% (14-22%)	22% (17-27%)	8500 (6700-10000)	250000 (200000-300000)
Germany	1995	Male	65 to 69	96% (94-98%)	4.1% (2.4-6.3%)	7.8 (6.4-9.4)	15% (11-19%)	18% (13-22%)	8900 (6600-11000)	220000 (160000-270000)
Germany	1995	Male	70 to 74	94% (91-97%)	5.5% (3.4-8.5%)	6.5 (5-8.2)	12% (7.1-16%)	14% (7.9-19%)	7200 (4100-9900)	140000 (83000-190000)
Germany	1995	Male	75 to 79	93% (89-97%)	6.5% (3.4-11%)	5.7 (4.2-7.4)	9.5% (3.7-14%)	10% (3.7-16%)	4400 (1600-6900)	68000 (27000-110000)
Germany	1995	Male	80 to 84	92% (86-96%)	8% (4.3-14%)	5.4 (4.1-6.7)	8.6% (3.3-13%)	9.3% (3.2-15%)	5900 (2000-9500)	67000 (26000-100000)
Germany	1995	Male	85 to 89	90% (83-94%)	9.9% (5.6-17%)	5 (3.8-6.6)	7.3% (2.3-12%)	7.8% (2-14%)	3700 (950-6300)	30000 (9500-49000)
Germany	1995	Male	90 to 94	89% (82-94%)	11% (5.9-18%)	4.8 (3.3-6.6)	6.1% (0.32-12%)	6.4% (-0.34-13%)	1200 (-64-2500)	6900 (360-13000)
Germany	1995	Male	95 plus	88% (80-95%)	12% (5.3-20%)	4.6 (2.7-6.8)	5.3% (-1.2-12%)	5.5% (-1.9-13%)	210 (-75-510)	800 (-1800-1800)
Germany	2000	Female	All Ages	90% (89-92%)	9.5% (7.9-11%)	3.1 (2.8-3.5)	5.2% (3.3-7.3%)	5.5% (1.5-10%)	25000 (7000-46000)	620000 (380000-870000)
Germany	2000	Female	15 to 19	90% (83-95%)	10% (5.4-17%)	2.3 (1.6-3.3)	3.2% (1.9-4.5%)	11% (5.2-17%)	66 (33-100)	7600 (4700-11000)
Germany	2000	Female	20 to 24	93% (89-96%)	6.7% (4.1-11%)	2.7 (1.7-3.9)	4.1% (3-5.4%)	11% (5.8-17%)	75 (40-110)	11000 (7500-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	2000	Female	25 to 29	95% (92-97%)	5.1% (3.1-8.2%)	2.5 (1.8-3.4)	3.4% (2.4-4.4%)	10% (6-15%)	81 (46-120)	11000 (7800-15000)
Germany	2000	Female	30 to 34	96% (93-98%)	4.3% (2.4-7%)	2.7 (2-3.6)	3.4% (2.5-4.5%)	12% (8.6-16%)	170 (120-230)	16000 (12000-21000)
Germany	2000	Female	35 to 39	96% (93-98%)	4.1% (2.3-6.7%)	3.4 (2.4-4.6)	5% (3.7-6.4%)	16% (12-20%)	420 (310-520)	29000 (21000-36000)
Germany	2000	Female	40 to 44	96% (93-97%)	4.4% (2.7-6.7%)	3.9 (2.9-5.1)	6.7% (5.1-8.4%)	18% (15-22%)	710 (570-860)	40000 (32000-49000)
Germany	2000	Female	45 to 49	95% (92-97%)	5.1% (3.1-7.9%)	4.3 (3.2-5.6)	8% (6.1-10%)	18% (15-22%)	1100 (860-1300)	51000 (40000-62000)
Germany	2000	Female	50 to 54	94% (92-96%)	5.6% (3.7-8.1%)	4.7 (3.5-6)	8.9% (6.7-11%)	18% (15-21%)	1300 (1100-1600)	55000 (43000-68000)
Germany	2000	Female	55 to 59	94% (92-96%)	5.9% (3.9-8.5%)	4.9 (3.8-6.2)	8.9% (6.7-11%)	16% (13-20%)	1900 (1500-2300)	68000 (52000-87000)
Germany	2000	Female	60 to 64	92% (89-94%)	8.1% (5.6-11%)	4.8 (3.6-6.2)	8% (5.4-11%)	13% (9.4-17%)	2600 (1900-3400)	80000 (55000-110000)
Germany	2000	Female	65 to 69	88% (83-92%)	12% (7.8-17%)	4.3 (3.4-5.4)	6.6% (3.9-9.4%)	9.9% (6.1-14%)	2500 (1500-3400)	63000 (37000-88000)
Germany	2000	Female	70 to 74	84% (78-89%)	16% (11-22%)	3.9 (2.8-5.1)	5.4% (2-8.8%)	7.3% (2.8-12%)	3000 (1100-4900)	62000 (22000-100000)
Germany	2000	Female	75 to 79	82% (73-89%)	18% (11-27%)	3.6 (2.5-4.9)	4.2% (0.068-8.5%)	5.1% (-0.15-11%)	3500 (-100-7400)	57000 (940-120000)
Germany	2000	Female	80 to 84	80% (69-88%)	20% (12-31%)	3.4 (2.5-4.5)	3.5% (-0.41-7.6%)	3.9% (-0.86-9.2%)	2600 (-570-6100)	32000 (-4000-70000)
Germany	2000	Female	85 to 89	79% (68-86%)	21% (14-32%)	3.3 (2.4-4.5)	2.8% (-1.2-7.3%)	2.9% (-1.9-8.4%)	3100 (-2000-8800)	27000 (-12000-72000)
Germany	2000	Female	90 to 94	79% (68-87%)	21% (13-32%)	3.3 (2.2-4.6)	2.2% (-2.2-7.3%)	2.1% (-3.3-8.4%)	1500 (-2200-5900)	9500 (-9400-32000)
Germany	2000	Female	95 plus	79% (65-89%)	21% (11-35%)	3.2 (1.9-4.9)	1.8% (-3.5-8.2%)	1.6% (-4.6-9.2%)	370 (-1100-2100)	1700 (-3300-7700)
Germany	2000	Male	All Ages	95% (93-96%)	5.3% (4-6.6%)	4.4 (3.9-4.9)	13% (11-15%)	14% (10-17%)	54000 (40000-69000)	160000 (130000-190000)
Germany	2000	Male	15 to 19	90% (84-95%)	10% (5.5-16%)	2.7 (1.8-3.7)	7.7% (4.9-11%)	14% (6.7-21%)	200 (98-300)	21000 (13000-29000)
Germany	2000	Male	20 to 24	94% (90-97%)	5.9% (3.4-9.6%)	3.8 (2.7-5.2)	14% (10-18%)	19% (10-27%)	390 (220-560)	44000 (31000-58000)
Germany	2000	Male	25 to 29	96% (94-98%)	3.9% (2.4-6.1%)	4 (2.9-5.3)	13% (9.9-16%)	20% (12-27%)	410 (240-560)	46000 (33000-59000)
Germany	2000	Male	30 to 34	97% (95-98%)	2.7% (1.5-4.6%)	4 (2.9-5.3)	12% (9.2-15%)	21% (14-27%)	680 (470-890)	61000 (45000-76000)
Germany	2000	Male	35 to 39	98% (96-99%)	2.3% (1.3-3.9%)	3.8 (2.7-5.1)	13% (10-16%)	22% (17-27%)	1200 (860-1500)	82000 (63000-100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	2000	Male	40 to 44	98% (96-99%)	2% (1.1-3.6%)	4.3 (3.2-5.5)	15% (12-18%)	24% (19-29%)	1900 (1500-2300)	110000 (85000-130000)
Germany	2000	Male	45 to 49	98% (97-99%)	1.8% (0.92-3.3%)	5.4 (4.1-7)	17% (13-21%)	25% (20-30%)	2800 (2200-3500)	130000 (110000-160000)
Germany	2000	Male	50 to 54	98% (97-99%)	2% (1.1-3.4%)	6.7 (5.3-8.3)	18% (15-21%)	25% (20-30%)	3700 (3000-4400)	150000 (120000-180000)
Germany	2000	Male	55 to 59	97% (96-99%)	2.5% (1.5-4.2%)	8.2 (6.7-9.9)	19% (16-22%)	25% (21-29%)	6000 (5000-7000)	210000 (180000-240000)
Germany	2000	Male	60 to 64	97% (95-98%)	3.2% (2-5.1%)	8.4 (6.7-10)	17% (14-21%)	22% (18-26%)	8900 (7100-11000)	260000 (210000-310000)
Germany	2000	Male	65 to 69	96% (93-98%)	4.2% (2.3-6.8%)	7.2 (5.9-8.7)	14% (11-18%)	17% (13-22%)	8000 (6000-10000)	200000 (150000-250000)
Germany	2000	Male	70 to 74	95% (92-97%)	5.2% (3.3-8.1%)	6 (4.7-7.6)	11% (6.2-15%)	12% (7-18%)	7300 (4100-10000)	140000 (84000-200000)
Germany	2000	Male	75 to 79	94% (89-97%)	6.2% (3.3-11%)	5.3 (4-6.9)	8.3% (3.2-13%)	9.2% (3.2-15%)	5200 (1800-8600)	82000 (32000-130000)
Germany	2000	Male	80 to 84	92% (86-96%)	7.9% (4.3-14%)	5 (3.9-6.3)	7.2% (2.6-12%)	7.8% (2.4-13%)	3200 (970-5300)	37000 (13000-60000)
Germany	2000	Male	85 to 89	90% (85-94%)	9.8% (5.8-15%)	4.7 (3.6-6)	6.2% (1.6-11%)	6.5% (1.2-12%)	3000 (550-5500)	25000 (6300-44000)
Germany	2000	Male	90 to 94	89% (83-94%)	11% (6.1-17%)	4.5 (3.1-6.1)	5% (-0.16-11%)	5.1% (-0.83-12%)	1100 (-180-2600)	6600 (-220-14000)
Germany	2000	Male	95 plus	88% (80-95%)	12% (5.3-20%)	4.3 (2.6-6.4)	4.3% (-1.7-11%)	4.2% (-2.7-12%)	210 (-140-630)	860 (-330-2300)
Germany	2005	Female	All Ages	90% (89-92%)	9.6% (8.1-11%)	3.1 (2.7-3.5)	4.8% (3.1-6.5%)	4.9% (1.5-8.5%)	22000 (6600-38000)	550000 (370000-750000)
Germany	2005	Female	15 to 19	90% (83-94%)	10% (5.6-17%)	2.2 (1.5-3.2)	2.6% (1.7-3.8%)	10% (5-16%)	48 (24-75)	6300 (4000-9300)
Germany	2005	Female	20 to 24	93% (89-96%)	6.6% (4-11%)	2.6 (1.6-3.8)	3.7% (2.7-5%)	10% (5.4-16%)	60 (32-90)	10000 (7100-15000)
Germany	2005	Female	25 to 29	95% (92-97%)	5% (3.1-7.6%)	2.4 (1.7-3.3)	3.1% (2.2-4%)	9.6% (5.5-14%)	58 (33-85)	9200 (6400-13000)
Germany	2005	Female	30 to 34	96% (93-98%)	4.3% (2.5-7%)	2.6 (1.9-3.6)	3% (2.2-4%)	11% (7.8-15%)	100 (71-140)	11000 (7500-14000)
Germany	2005	Female	35 to 39	96% (94-98%)	4.2% (2.4-6.4%)	3.2 (2.3-4.4)	4.1% (2.9-5.3%)	15% (11-19%)	290 (220-370)	21000 (16000-27000)
Germany	2005	Female	40 to 44	96% (93-97%)	4.3% (2.6-6.5%)	3.8 (2.8-4.9)	5.6% (4.2-7.2%)	17% (13-20%)	620 (500-750)	36000 (28000-45000)
Germany	2005	Female	45 to 49	95% (93-97%)	4.9% (3.2-7.2%)	4.2 (3.1-5.4)	7.2% (5.5-9.1%)	17% (14-21%)	1000 (810-1200)	48000 (38000-60000)
Germany	2005	Female	50 to 54	95% (92-96%)	5.4% (3.8-7.6%)	4.5 (3.4-5.7)	8% (6.2-10%)	17% (14-20%)	1300 (1100-1600)	57000 (45000-70000)
Germany	2005	Female	55 to 59	94% (92-96%)	5.8% (3.9-8.3%)	4.8 (3.6-6)	8.3% (6.3-11%)	15% (12-19%)	1600 (1200-1900)	57000 (44000-71000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	2005	Female	60 to 64	92% (89-94%)	7.9% (5.6-11%)	4.7 (3.5-6.1)	7.6% (5.3-10%)	13% (9.8-17%)	2000 (1500-2700)	64000 (44000-84000)
Germany	2005	Female	65 to 69	88% (83-92%)	12% (8.3-17%)	4.2 (3.3-5.2)	6.2% (4-8.6%)	9.9% (6.7-13%)	2600 (1800-3600)	69000 (45000-93000)
Germany	2005	Female	70 to 74	84% (79-88%)	16% (12-21%)	3.7 (2.7-4.8)	4.9% (2.2-7.8%)	7% (3.2-11%)	2400 (1100-3900)	51000 (22000-81000)
Germany	2005	Female	75 to 79	81% (73-87%)	19% (13-27%)	3.4 (2.4-4.5)	3.8% (0.59-7.4%)	4.8% (0.66-9.4%)	2800 (380-5600)	46000 (7200-90000)
Germany	2005	Female	80 to 84	79% (68-87%)	21% (13-32%)	3.2 (2.4-4.1)	3.1% (-0.12-6.6%)	3.5% (-0.51-7.8%)	3300 (-500-7300)	40000 (-1600-86000)
Germany	2005	Female	85 to 89	78% (67-86%)	22% (14-33%)	3.1 (2.3-4.2)	2.4% (-1.2-6.1%)	2.6% (-1.8-7.1%)	1900 (-1400-5200)	17000 (-8500-43000)
Germany	2005	Female	90 to 94	77% (66-86%)	23% (14-34%)	3.1 (2.1-4.4)	1.8% (-2.2-6.6%)	1.8% (-3.1-7.5%)	1400 (-2400-5800)	8900 (-11000-32000)
Germany	2005	Female	95 plus	77% (63-88%)	23% (12-37%)	3 (1.8-4.7)	1.5% (-3.2-7.5%)	1.4% (-4.4-8.5%)	440 (-1400-2800)	2000 (-4200-9900)
Germany	2005	Male	All Ages	95% (94-96%)	5% (3.8-6.2%)	4.3 (3.8-4.8)	12% (9.8-14%)	13% (9.2-16%)	49000 (36000-64000)	1400000 (1200000-1700000)
Germany	2005	Male	15 to 19	90% (85-94%)	9.7% (5.7-15%)	2.6 (1.8-3.6)	6.5% (4.2-8.9%)	13% (6.5-20%)	140 (71-220)	17000 (11000-23000)
Germany	2005	Male	20 to 24	94% (91-97%)	5.5% (3.3-9.1%)	3.6 (2.5-5.1)	12% (9-16%)	18% (10-26%)	290 (160-420)	36000 (26000-48000)
Germany	2005	Male	25 to 29	96% (94-98%)	3.6% (2.2-5.8%)	3.8 (2.8-5.1)	12% (8.8-14%)	19% (11-26%)	300 (170-410)	35000 (26000-46000)
Germany	2005	Male	30 to 34	97% (96-99%)	2.6% (1.5-4.3%)	3.8 (2.8-5.1)	10% (8.1-13%)	19% (13-26%)	390 (250-520)	37000 (27000-47000)
Germany	2005	Male	35 to 39	98% (96-99%)	2.2% (1.3-3.5%)	3.7 (2.6-5)	11% (8.5-13%)	20% (15-26%)	800 (570-1000)	59000 (46000-73000)
Germany	2005	Male	40 to 44	98% (97-99%)	1.9% (1-3.5%)	4.1 (3-5.3)	13% (10-16%)	22% (17-27%)	1600 (1300-2000)	93000 (74000-110000)
Germany	2005	Male	45 to 49	98% (97-99%)	1.8% (0.89-3.3%)	5.1 (3.8-6.7)	15% (12-19%)	24% (18-29%)	2600 (2000-3300)	130000 (100000-150000)
Germany	2005	Male	50 to 54	98% (97-99%)	1.9% (1.1-3.2%)	6.4 (5-8)	17% (14-20%)	24% (19-29%)	3900 (3100-4600)	160000 (130000-190000)
Germany	2005	Male	55 to 59	98% (96-99%)	2.4% (1.4-3.9%)	7.9 (6.2-9.7)	18% (15-21%)	24% (20-28%)	4800 (4000-5600)	170000 (140000-200000)
Germany	2005	Male	60 to 64	97% (95-98%)	3.1% (1.9-4.7%)	8 (6.2-10)	16% (13-20%)	21% (17-25%)	6600 (5300-7900)	200000 (160000-240000)
Germany	2005	Male	65 to 69	96% (94-98%)	4% (2.4-6%)	6.9 (5.7-8.3)	14% (10-17%)	17% (13-21%)	8500 (6500-11000)	210000 (160000-260000)
Germany	2005	Male	70 to 74	95% (93-97%)	4.8% (3.3-7%)	5.8 (4.6-7.2)	10% (6.4-14%)	12% (7.3-17%)	6700 (4000-9400)	130000 (84000-190000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	2005	Male	75 to 79	94% (91-97%)	5.7% (3.2-9.2%)	5.2 (3.8-6.6)	7.7% (3.3-12%)	8.6% (3.4-14%)	5700 (2200-9200)	90000 (40000-140000)
Germany	2005	Male	80 to 84	93% (88-96%)	7.4% (4.3-12%)	4.8 (3.8-6.1)	6.5% (2.3-11%)	7% (2.1-12%)	4100 (1300-6900)	48000 (17000-80000)
Germany	2005	Male	85 to 89	91% (86-94%)	9.2% (5.6-14%)	4.5 (3.4-5.7)	5.4% (1.3-9.7%)	5.6% (0.8-11%)	1800 (260-3500)	16000 (3900-28000)
Germany	2005	Male	90 to 94	90% (84-94%)	10% (5.8-16%)	4.3 (3-5.9)	4.3% (-0.36-9.5%)	4.3% (-1.1-10%)	1000 (-270-2500)	6200 (-490-14000)
Germany	2005	Male	95 plus	89% (80-95%)	11% (5.1-20%)	4.1 (2.5-6.1)	3.8% (-1.5-10%)	3.7% (-2.5-11%)	250 (-170-750)	1000 (-410-2700)
Germany	2010	Female	All Ages	90% (89-92%)	9.7% (8.2-11%)	3 (2.5-3.5)	4.3% (2.7-6%)	4.4% (1.2-7.9%)	20000 (5700-36000)	500000 (320000-680000)
Germany	2010	Female	15 to 19	90% (84-95%)	9.9% (5.4-16%)	2.1 (1.4-3)	2.5% (1.6-3.5%)	9.4% (4.9-15%)	34 (18-52)	5000 (3200-7100)
Germany	2010	Female	20 to 24	94% (90-96%)	6.1% (3.7-9.9%)	2.4 (1.6-3.6)	3.7% (2.7-4.9%)	9.7% (5-15%)	50 (27-77)	10000 (7000-14000)
Germany	2010	Female	25 to 29	95% (92-97%)	4.9% (3.1-7.5%)	2.3 (1.5-3.2)	3.1% (2.2-4%)	8.8% (4.8-13%)	53 (29-79)	9300 (6400-13000)
Germany	2010	Female	30 to 34	95% (92-97%)	4.7% (2.6-7.6%)	2.4 (1.7-3.4)	2.9% (2-3.7%)	11% (7.1-15%)	84 (55-120)	9300 (6500-12000)
Germany	2010	Female	35 to 39	95% (93-97%)	4.5% (2.6-7%)	3 (2.1-4.4)	3.7% (2.6-4.9%)	14% (10-18%)	180 (130-230)	14000 (10000-18000)
Germany	2010	Female	40 to 44	96% (93-97%)	4.3% (2.7-6.7%)	3.6 (2.6-4.7)	4.9% (3.5-6.4%)	16% (12-19%)	470 (360-580)	28000 (21000-36000)
Germany	2010	Female	45 to 49	95% (93-97%)	4.8% (3.1-7.3%)	4 (2.9-5.2)	6.3% (4.7-8.1%)	16% (13-20%)	920 (740-1100)	45000 (35000-57000)
Germany	2010	Female	50 to 54	94% (92-96%)	5.6% (3.8-8%)	4.2 (3.2-5.5)	7.3% (5.4-9.2%)	16% (13-19%)	1300 (1100-1500)	54000 (42000-68000)
Germany	2010	Female	55 to 59	94% (92-96%)	5.7% (3.8-8%)	4.5 (3.4-5.9)	7.5% (5.5-9.8%)	14% (11-17%)	1600 (1200-2000)	59000 (44000-75000)
Germany	2010	Female	60 to 64	92% (90-95%)	7.5% (5.2-10%)	4.5 (3.3-5.9)	7.1% (4.9-9.6%)	12% (9.3-16%)	1700 (1300-2200)	54000 (38000-72000)
Germany	2010	Female	65 to 69	88% (83-92%)	12% (8.2-17%)	3.9 (3.1-5)	5.8% (3.8-8%)	9.4% (6.7-12%)	2100 (1500-2800)	54000 (37000-74000)
Germany	2010	Female	70 to 74	85% (80-88%)	15% (12-20%)	3.5 (2.6-4.6)	4.5% (2.1-7.1%)	6.7% (3.5-10%)	2600 (1400-4000)	55000 (26000-86000)
Germany	2010	Female	75 to 79	80% (72-87%)	20% (13-28%)	3.2 (2.2-4.3)	3.4% (0.63-6.4%)	4.5% (0.87-8.2%)	2300 (450-4500)	38000 (7300-72000)
Germany	2010	Female	80 to 84	77% (66-86%)	23% (14-34%)	2.9 (2.1-3.9)	2.6% (-0.13-5.7%)	3% (-0.48-6.9%)	2500 (-400-5700)	31000 (-1400-68000)
Germany	2010	Female	85 to 89	77% (68-85%)	23% (15-32%)	2.9 (2.1-3.9)	2.1% (-1.1-5.7%)	2.2% (-1.7-6.5%)	2500 (-1900-7400)	22000 (-11000-59000)
Germany	2010	Female	90 to 94	77% (67-86%)	23% (14-33%)	2.9 (1.9-4.1)	1.6% (-2.1-5.9%)	1.6% (-2.9-6.8%)	930 (-1700-4000)	6100 (-8000-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	2010	Female	95 plus	77% (64-89%)	23% (11-36%)	2.8 (1.6-4.4)	1.4% (-3.3-7%)	1.2% (-4.6-7.8%)	470 (-1800-3000)	2200 (-5200-11000)
Germany	2010	Male	All Ages	95% (94-96%)	4.9% (3.7-6.1%)	4.1 (3.5-4.8)	11% (8.7-13%)	11% (7.7-15%)	45000 (31000-60000)	130000 (100000-160000)
Germany	2010	Male	15 to 19	91% (86-95%)	9% (5.1-14%)	2.4 (1.7-3.4)	5.9% (3.9-8.1%)	12% (6.2-19%)	95 (46-140)	12000 (8300-17000)
Germany	2010	Male	20 to 24	95% (92-97%)	5% (3-8%)	3.4 (2.3-4.7)	12% (8.9-15%)	18% (9.1-25%)	230 (120-340)	34000 (24000-44000)
Germany	2010	Male	25 to 29	97% (94-98%)	3.5% (2.2-5.5%)	3.6 (2.5-4.8)	11% (8.4-14%)	18% (9.9-25%)	260 (150-380)	34000 (24000-44000)
Germany	2010	Male	30 to 34	97% (96-99%)	2.7% (1.5-4.5%)	3.6 (2.5-4.8)	9.8% (7.3-12%)	18% (11-25%)	310 (200-430)	31000 (22000-39000)
Germany	2010	Male	35 to 39	98% (96-99%)	2.2% (1.2-3.8%)	3.5 (2.5-4.8)	10% (7.6-12%)	19% (13-25%)	480 (340-620)	38000 (29000-47000)
Germany	2010	Male	40 to 44	98% (96-99%)	2% (1-3.6%)	3.9 (2.9-5.1)	11% (8.7-14%)	21% (16-26%)	1100 (860-1400)	69000 (54000-85000)
Germany	2010	Male	45 to 49	98% (97-99%)	1.8% (0.92-3.2%)	4.8 (3.5-6.3)	14% (11-17%)	22% (17-28%)	2300 (1800-2900)	120000 (91000-140000)
Germany	2010	Male	50 to 54	98% (97-99%)	1.9% (1-3.1%)	6 (4.7-7.6)	16% (13-19%)	23% (19-28%)	3700 (2900-4400)	150000 (120000-180000)
Germany	2010	Male	55 to 59	98% (96-99%)	2.5% (1.5-4%)	7.4 (5.7-9.3)	17% (14-20%)	23% (19-27%)	5000 (4000-5900)	180000 (150000-210000)
Germany	2010	Male	60 to 64	97% (95-98%)	3.1% (1.9-4.8%)	7.5 (5.8-9.6)	15% (12-19%)	20% (16-24%)	5400 (4200-6400)	160000 (130000-200000)
Germany	2010	Male	65 to 69	96% (94-98%)	3.9% (2.3-6.1%)	6.5 (5.2-8)	13% (9.4-16%)	16% (12-20%)	6300 (4700-7900)	160000 (120000-200000)
Germany	2010	Male	70 to 74	95% (93-97%)	4.7% (3.1-6.9%)	5.5 (4.2-6.9)	9.4% (6-13%)	11% (6.9-16%)	7100 (4400-10000)	150000 (93000-200000)
Germany	2010	Male	75 to 79	94% (91-97%)	5.5% (3-9.2%)	4.9 (3.6-6.3)	7% (3.3-11%)	7.9% (3.3-13%)	5200 (2200-8500)	84000 (40000-130000)
Germany	2010	Male	80 to 84	93% (88-96%)	7.1% (4.1-12%)	4.6 (3.5-5.7)	5.5% (2.1-9.4%)	5.9% (1.7-11%)	4200 (1300-7600)	52000 (20000-90000)
Germany	2010	Male	85 to 89	91% (87-94%)	8.8% (5.6-13%)	4.3 (3.2-5.7)	4.5% (0.78-8.7%)	4.6% (0.18-9.6%)	2400 (92-5000)	21000 (3600-40000)
Germany	2010	Male	90 to 94	90% (85-94%)	9.9% (5.7-15%)	4.1 (2.8-5.7)	3.6% (-0.58-8.7%)	3.5% (-1.3-9.4%)	680 (-250-1800)	4300 (-670-10000)
Germany	2010	Male	95 plus	89% (81-95%)	11% (5.1-19%)	3.9 (2.4-5.9)	3.3% (-1.8-9.6%)	3.1% (-2.8-10%)	240 (-220-820)	1000 (-560-3000)
Germany	2016	Female	All Ages	90% (88-92%)	9.3% (8.3-12%)	2.9 (2.2-3.8)	4% (2.1-6.2%)	4% (0.45-8%)	19000 (2000-38000)	470000 (250000-720000)
Germany	2016	Female	15 to 19	90% (83-95%)	9.8% (5-17%)	2 (1.3-3)	2.2% (1.4-3.2%)	9% (4.3-14%)	27 (13-42)	4300 (2700-6100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	2016	Female	20 to 24	93% (89-96%)	6.6% (3.9-11%)	2.3 (1.4-3.6)	3.3% (2.3-4.4%)	9.4% (4.8-15%)	40 (20-65)	8400 (5500-12000)
Germany	2016	Female	25 to 29	95% (91-97%)	5.4% (3.1-9.2%)	2.2 (1.4-3.2)	2.7% (1.9-3.6%)	8.5% (4.4-13%)	47 (24-72)	8500 (5600-12000)
Germany	2016	Female	30 to 34	95% (92-98%)	4.7% (2.5-8%)	2.4 (1.5-3.5)	2.6% (1.8-3.6%)	10% (6.6-14%)	83 (52-120)	9200 (6300-13000)
Germany	2016	Female	35 to 39	96% (92-98%)	4.4% (2.4-7.5%)	2.9 (1.9-4.4)	3.3% (2.2-4.6%)	14% (9.5-18%)	150 (100-200)	12000 (7900-16000)
Germany	2016	Female	40 to 44	95% (92-97%)	4.7% (2.8-7.7%)	3.4 (2.3-4.8)	4.5% (3.1-6.1%)	16% (12-19%)	320 (230-410)	19000 (14000-26000)
Germany	2016	Female	45 to 49	95% (91-97%)	5.4% (3.2-8.9%)	3.8 (2.6-5.3)	5.7% (3.9-7.6%)	16% (12-19%)	790 (600-1000)	39000 (28000-52000)
Germany	2016	Female	50 to 54	94% (91-96%)	5.9% (3.8-9.2%)	4.1 (2.7-5.6)	6.7% (4.7-8.9%)	15% (12-18%)	1300 (1000-1700)	57000 (41000-74000)
Germany	2016	Female	55 to 59	94% (91-96%)	6% (3.9-8.8%)	4.4 (3-6.1)	6.9% (4.7-9.5%)	14% (10-17%)	1600 (1100-2100)	57000 (39000-78000)
Germany	2016	Female	60 to 64	92% (88-95%)	8.2% (5.4-12%)	4.2 (2.9-6.1)	6.5% (4-9.3%)	11% (8-16%)	2000 (1400-2800)	62000 (38000-90000)
Germany	2016	Female	65 to 69	88% (83-92%)	12% (7.6-17%)	3.8 (2.7-5.3)	5.4% (3.2-8%)	8.9% (5.9-13%)	1600 (1000-2400)	42000 (24000-64000)
Germany	2016	Female	70 to 74	86% (80-90%)	14% (10-20%)	3.5 (2.4-4.8)	4.2% (1.6-7.1%)	6.4% (3-10%)	2300 (1000-3800)	48000 (18000-83000)
Germany	2016	Female	75 to 79	83% (74-89%)	17% (11-26%)	3.2 (2.1-4.6)	3.2% (0.25-6.5%)	4.3% (0.44-8.5%)	2700 (270-5400)	45000 (3400-92000)
Germany	2016	Female	80 to 84	80% (69-88%)	20% (12-31%)	3 (2.1-4.2)	2.4% (-0.62-6%)	2.8% (-1.1-7.4%)	2100 (-770-5500)	26000 (-6500-65000)
Germany	2016	Female	85 to 89	79% (68-86%)	21% (14-32%)	2.9 (1.9-4.2)	1.9% (-1.6-5.8%)	2% (-2.3-6.8%)	2100 (-2400-7300)	19000 (-16000-59000)
Germany	2016	Female	90 to 94	78% (68-87%)	22% (13-32%)	2.8 (1.8-4.4)	1.4% (-2.3-6.1%)	1.3% (-3.3-7.1%)	1300 (-3400-7000)	9300 (-15000-39000)
Germany	2016	Female	95 plus	79% (65-89%)	21% (11-35%)	2.8 (1.5-4.6)	1.2% (-3.5-7.1%)	0.97% (-4.6-7.9%)	240 (-1100-1900)	1200 (-3300-6700)
Germany	2016	Male	All Ages	94% (93-96%)	5.7% (4.3-7.2%)	4 (3-5.2)	9.9% (7.2-13%)	9.7% (5.6-14%)	43000 (25000-64000)	1200000 (890000-1600000)
Germany	2016	Male	15 to 19	90% (82-95%)	10% (5.5-18%)	2.3 (1.5-3.4)	5.2% (3.4-7.3%)	12% (5.8-18%)	78 (37-120)	10000 (6700-15000)
Germany	2016	Male	20 to 24	94% (89-97%)	6.3% (3.4-11%)	3.2 (2.1-4.7)	11% (7.5-14%)	17% (8.2-24%)	190 (93-290)	27000 (19000-37000)
Germany	2016	Male	25 to 29	95% (92-98%)	4.5% (2.5-7.6%)	3.4 (2.2-4.8)	9.9% (7.3-13%)	17% (9-24%)	240 (120-360)	31000 (22000-41000)
Germany	2016	Male	30 to 34	97% (94-98%)	3.4% (1.7-6%)	3.4 (2.3-5)	8.9% (6.5-11%)	17% (10-24%)	300 (180-440)	31000 (22000-41000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Germany	2016	Male	35 to 39	97% (95-99%)	2.8% (1.4-5.1%)	3.3 (2.1-4.8)	9.2% (6.9-12%)	19% (13-24%)	400 (270-530)	31000 (23000-41000)
Germany	2016	Male	40 to 44	98% (95-99%)	2.4% (1.2-4.8%)	3.7 (2.5-5.2)	11% (8-14%)	21% (15-26%)	760 (540-970)	46000 (35000-58000)
Germany	2016	Male	45 to 49	98% (96-99%)	2.2% (1.1-4.5%)	4.6 (3.1-6.5)	13% (9.7-16%)	22% (17-28%)	2000 (1500-2600)	100000 (77000-130000)
Germany	2016	Male	50 to 54	98% (95-99%)	2.5% (1.3-4.5%)	5.7 (3.9-7.8)	15% (11-18%)	23% (17-28%)	3700 (2800-4700)	160000 (120000-200000)
Germany	2016	Male	55 to 59	97% (95-98%)	3% (1.7-5%)	7 (5-9.5)	16% (12-19%)	22% (17-27%)	4900 (3600-6100)	180000 (130000-220000)
Germany	2016	Male	60 to 64	96% (94-98%)	3.6% (2.1-5.7%)	7.1 (5.1-9.8)	15% (11-18%)	19% (14-24%)	6100 (4400-7600)	190000 (130000-230000)
Germany	2016	Male	65 to 69	96% (93-98%)	4.4% (2.5-7.2%)	6.1 (4.5-8.4)	12% (8-15%)	15% (10-19%)	4400 (2900-5800)	110000 (75000-150000)
Germany	2016	Male	70 to 74	95% (92-97%)	5.3% (3.3-8.1%)	5.2 (3.7-7.1)	8.7% (4.9-13%)	11% (5.8-15%)	6200 (3400-9300)	130000 (72000-190000)
Germany	2016	Male	75 to 79	94% (89-97%)	6.4% (3.3-11%)	4.6 (3.1-6.3)	6.4% (2.7-11%)	7.3% (2.9-13%)	6200 (2400-11000)	100000 (44000-180000)
Germany	2016	Male	80 to 84	92% (86-96%)	8.2% (4.4-14%)	4.3 (3-5.8)	5% (1.3-9.3%)	5.3% (0.97-10%)	4000 (760-8000)	49000 (12000-94000)
Germany	2016	Male	85 to 89	90% (83-94%)	10% (5.7-17%)	4 (2.7-5.7)	4% (0.086-8.4%)	4.1% (-0.4-9.2%)	2800 (-270-6400)	25000 (530-53000)
Germany	2016	Male	90 to 94	89% (82-94%)	11% (5.8-18%)	3.9 (2.4-5.7)	3.2% (-1.1-8.4%)	3.1% (-1.7-9.3%)	1100 (-620-3300)	7000 (-2400-19000)
Germany	2016	Male	95 plus	88% (79-95%)	12% (5.2-21%)	3.7 (2.1-5.9)	2.8% (-2.2-9.2%)	2.7% (-3.1-10%)	140 (-160-530)	600 (-440-1900)
Greece	1990	Female	All Ages	78% (73-82%)	22% (18-27%)	2.5 (2.2-2.9)	4.2% (2.3-6.1%)	5.2% (1.1-9.5%)	2300 (490-4200)	55000 (31000-82000)
Greece	1990	Female	15 to 19	74% (59-86%)	26% (14-41%)	1.9 (1.3-2.7)	2.4% (1.3-3.6%)	8.3% (3.7-14%)	8.3 (3.6-14)	920 (540-1400)
Greece	1990	Female	20 to 24	79% (66-89%)	21% (11-34%)	2.3 (1.5-3.3)	3.2% (2.1-4.5%)	8.7% (4.1-14%)	12 (5.9-20)	1500 (950-2200)
Greece	1990	Female	25 to 29	81% (69-90%)	19% (10-31%)	2.2 (1.5-3.2)	2.7% (1.7-3.8%)	7.9% (3.8-12%)	12 (5.8-19)	1300 (820-1900)
Greece	1990	Female	30 to 34	82% (69-90%)	18% (10-31%)	2.5 (1.8-3.5)	2.6% (1.7-3.7%)	8.5% (5.1-12%)	16 (9.6-23)	1400 (930-1900)
Greece	1990	Female	35 to 39	81% (72-88%)	19% (12-28%)	3.2 (2.2-4.5)	3.3% (2.3-4.6%)	11% (7.5-15%)	25 (17-34)	1800 (1200-2400)
Greece	1990	Female	40 to 44	82% (75-88%)	18% (12-25%)	3.6 (2.6-4.8)	4.2% (2.9-5.7%)	12% (8.8-16%)	44 (31-57)	2600 (1900-3400)
Greece	1990	Female	45 to 49	85% (79-89%)	15% (11-21%)	3.8 (2.7-5)	4.9% (3.4-6.8%)	12% (8.9-16%)	57 (40-74)	2900 (2000-3800)
Greece	1990	Female	50 to 54	87% (82-90%)	13% (9.9-18%)	3.9 (2.8-5.1)	5.5% (3.6-7.6%)	12% (8.5-16%)	100 (71-140)	4400 (3000-6000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	1990	Female	55 to 59	86% (81-90%)	14% (10-19%)	4 (2.9-5.1)	6% (3.7-8.4%)	11% (7.6-15%)	150 (100-210)	5600 (3600-7700)
Greece	1990	Female	60 to 64	84% (78-88%)	16% (12-22%)	3.8 (2.8-5.1)	6% (3-8.9%)	9.9% (5.3-15%)	230 (120-330)	6900 (3500-10000)
Greece	1990	Female	65 to 69	80% (73-85%)	20% (15-27%)	3.5 (2.6-4.4)	5.8% (2.8-8.8%)	8.6% (4.1-13%)	260 (120-390)	6400 (3100-9600)
Greece	1990	Female	70 to 74	78% (72-83%)	22% (17-28%)	3.2 (2.2-4.2)	5.7% (1.9-9.8%)	7.6% (2.4-13%)	370 (110-630)	7000 (2300-12000)
Greece	1990	Female	75 to 79	79% (71-85%)	21% (15-29%)	2.9 (2-4.2)	4.6% (-0.023-9.8%)	5.7% (-0.14-12%)	450 (-11-960)	6700 (-33-14000)
Greece	1990	Female	80 to 84	77% (68-84%)	23% (16-32%)	2.7 (1.9-3.7)	3.5% (-0.84-7.9%)	4.1% (-1.2-9.5%)	370 (-110-850)	4100 (-970-9300)
Greece	1990	Female	85 to 89	74% (63-81%)	26% (19-37%)	2.4 (1.6-3.4)	2.2% (-1.5-6.3%)	2.5% (-2-7.3%)	190 (-150-560)	1500 (-1000-4400)
Greece	1990	Female	90 to 94	71% (59-80%)	29% (20-41%)	2.1 (1.3-3.2)	1.1% (-2.5-5.3%)	1.1% (-3.2-6.1%)	47 (-140-260)	290 (-660-1400)
Greece	1990	Female	95 plus	69% (56-80%)	31% (20-44%)	1.9 (1-3.2)	0.47% (-3.3-5.3%)	0.3% (-4.2-6.1%)	4.5 (-63-92)	28 (-200-320)
Greece	1990	Male	All Ages	90% (86-93%)	10% (7.3-14%)	5 (4.3-5.6)	12% (9.2-14%)	15% (11-19%)	7400 (5600-9500)	170000 (140000-210000)
Greece	1990	Male	15 to 19	84% (72-92%)	16% (8.3-28%)	2.8 (1.9-3.9)	7.1% (3.9-10%)	13% (6.1-20%)	39 (18-61)	3400 (2000-5100)
Greece	1990	Male	20 to 24	89% (79-94%)	11% (5.6-21%)	4.1 (2.9-5.7)	13% (8.3-18%)	19% (10-27%)	93 (52-140)	8000 (5100-11000)
Greece	1990	Male	25 to 29	91% (83-96%)	9.1% (4.4-17%)	4.9 (3.5-6.6)	12% (8.1-17%)	20% (12-29%)	82 (48-120)	6600 (4300-8900)
Greece	1990	Male	30 to 34	93% (85-97%)	7.4% (3.2-15%)	5.3 (3.9-6.9)	10% (6.9-14%)	18% (11-26%)	76 (47-110)	5700 (3700-7700)
Greece	1990	Male	35 to 39	94% (88-97%)	6.3% (2.7-12%)	5.3 (3.8-7.2)	9.1% (5.8-13%)	16% (9-23%)	77 (44-110)	5200 (3300-7300)
Greece	1990	Male	40 to 44	95% (89-98%)	5.3% (2.2-11%)	6.2 (4.6-7.9)	10% (6.6-14%)	16% (10-23%)	120 (73-170)	6800 (4500-9100)
Greece	1990	Male	45 to 49	96% (92-98%)	4.2% (2.2-7.9%)	8 (5.9-10)	13% (9-16%)	19% (13-25%)	180 (120-230)	8600 (6200-11000)
Greece	1990	Male	50 to 54	96% (93-98%)	3.7% (2-6.6%)	8.8 (6.6-11)	14% (10-17%)	20% (15-25%)	320 (250-410)	13000 (10000-17000)
Greece	1990	Male	55 to 59	96% (93-98%)	4.1% (2.4-6.6%)	8.8 (6.9-11)	14% (11-18%)	19% (14-24%)	520 (380-660)	18000 (14000-23000)
Greece	1990	Male	60 to 64	95% (92-97%)	5% (3.1-7.8%)	8.4 (6.3-11)	14% (10-18%)	17% (12-23%)	760 (540-1000)	23000 (16000-29000)
Greece	1990	Male	65 to 69	93% (90-96%)	6.6% (4-9.8%)	7.7 (5.9-9.5)	14% (10-18%)	17% (12-21%)	830 (590-1100)	20000 (15000-25000)
Greece	1990	Male	70 to 74	93% (90-95%)	6.9% (4.7-10%)	7 (5.2-9)	14% (9.3-18%)	16% (11-21%)	930 (630-1200)	18000 (12000-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	1990	Male	75 to 79	94% (90-96%)	6.2% (3.6-10%)	6.4 (4.6-8.7)	13% (8-17%)	15% (9.1-20%)	1200 (740-1700)	18000 (11000-25000)
Greece	1990	Male	80 to 84	93% (88-96%)	6.9% (3.8-12%)	5.9 (4.4-7.8)	12% (7.6-17%)	14% (8.5-19%)	1100 (680-1500)	12000 (7700-17000)
Greece	1990	Male	85 to 89	91% (85-95%)	8.8% (5-15%)	5.4 (3.8-7.3)	11% (5.9-16%)	12% (6.5-18%)	700 (380-1000)	5500 (3000-8000)
Greece	1990	Male	90 to 94	90% (84-94%)	9.8% (5.6-16%)	5 (3.3-7.1)	8.9% (3.2-15%)	10% (3.3-17%)	280 (92-470)	1500 (550-2500)
Greece	1990	Male	95 plus	90% (82-95%)	10% (5.1-18%)	4.7 (2.6-7.3)	7.9% (1.2-15%)	8.7% (0.9-16%)	74 (7.8-140)	270 (39-490)
Greece	1995	Female	All Ages	78% (74-82%)	22% (18-26%)	2.4 (2.1-2.8)	3.4% (1.8-5.4%)	3.9% (0.16-7.9%)	1800 (74-3700)	47000 (24000-73000)
Greece	1995	Female	15 to 19	74% (59-85%)	26% (15-41%)	1.8 (1.2-2.5)	2.3% (1.3-3.4%)	7.8% (3.5-13%)	7.9 (3.5-13)	890 (510-1300)
Greece	1995	Female	20 to 24	79% (66-88%)	21% (12-34%)	2.1 (1.4-3)	2.9% (1.8-4.1%)	8% (3.7-13%)	9.9 (4.7-16)	1400 (830-1900)
Greece	1995	Female	25 to 29	81% (69-90%)	19% (10-31%)	2.1 (1.4-2.9)	2.4% (1.6-3.5%)	7.3% (3.6-11%)	11 (5.5-17)	1300 (840-1900)
Greece	1995	Female	30 to 34	82% (69-89%)	18% (11-31%)	2.4 (1.7-3.3)	2.4% (1.5-3.4%)	7.6% (4.6-11%)	14 (8.1-20)	1300 (850-1900)
Greece	1995	Female	35 to 39	81% (73-87%)	19% (13-27%)	3 (2.1-4.3)	3% (2-4.3%)	9.6% (6.3-13%)	24 (16-34)	1800 (1200-2500)
Greece	1995	Female	40 to 44	82% (74-88%)	18% (12-26%)	3.4 (2.4-4.5)	3.8% (2.6-5.2%)	11% (7.7-15%)	39 (27-51)	2400 (1700-3200)
Greece	1995	Female	45 to 49	85% (78-89%)	15% (11-22%)	3.6 (2.5-4.8)	4.3% (2.9-6.1%)	11% (7.8-15%)	60 (41-81)	3100 (2100-4200)
Greece	1995	Female	50 to 54	86% (81-90%)	14% (9.8-19%)	3.7 (2.6-4.9)	4.9% (3.2-6.9%)	11% (7.5-15%)	74 (50-100)	3200 (2100-4500)
Greece	1995	Female	55 to 59	86% (81-90%)	14% (10-19%)	3.7 (2.7-4.9)	5.2% (3.2-7.3%)	10% (6.5-14%)	130 (79-170)	4700 (2900-6600)
Greece	1995	Female	60 to 64	84% (78-88%)	16% (12-22%)	3.6 (2.5-4.8)	5% (2.5-7.7%)	8.7% (4.5-13%)	180 (91-260)	5500 (2700-8300)
Greece	1995	Female	65 to 69	80% (73-85%)	20% (15-27%)	3.2 (2.4-4.2)	4.8% (2.2-7.5%)	7.4% (3.4-11%)	260 (120-410)	6600 (3000-10000)
Greece	1995	Female	70 to 74	78% (71-84%)	22% (16-29%)	2.9 (2.1-4)	4.6% (1.1-8.4%)	6.3% (1.5-11%)	300 (73-540)	5800 (1500-11000)
Greece	1995	Female	75 to 79	79% (70-85%)	21% (15-30%)	2.7 (1.9-3.9)	3.6% (-0.75-8.3%)	4.5% (-1.1-11%)	310 (-74-730)	4700 (-950-11000)
Greece	1995	Female	80 to 84	77% (68-84%)	23% (16-32%)	2.5 (1.7-3.5)	2.6% (-1.2-6.6%)	3% (-1.9-8%)	270 (-170-730)	3100 (-1500-8000)
Greece	1995	Female	85 to 89	73% (63-81%)	27% (19-37%)	2.2 (1.5-3.1)	1.4% (-2.5-3%)	1.5% (-2.6-6.1%)	140 (-230-540)	1200 (-1700-4200)
Greece	1995	Female	90 to 94	71% (60-80%)	29% (20-40%)	2 (1.2-3)	0.51% (-2.8-4%)	0.37% (-3.8-4.7%)	19 (-190-240)	160 (-920-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	1995	Female	95 plus	69% (55-80%)	31% (20-45%)	1.8 (0.92-3)	-0.079% (-3.6-4.1%)	-0.37% (-4.5-4.6%)	-8 (-98-100)	-6.8 (-310-350)
Greece	1995	Male	All Ages	90% (86-93%)	10% (7.4-14%)	4.8 (4.2-5.5)	11% (8.5-13%)	14% (9.8-18%)	7200 (5100-9400)	170000 (140000-210000)
Greece	1995	Male	15 to 19	83% (72-92%)	17% (8.3-28%)	2.6 (1.8-3.7)	6.8% (3.7-10%)	12% (5.6-19%)	38 (17-59)	3400 (1900-5000)
Greece	1995	Male	20 to 24	88% (79-94%)	12% (5.8-21%)	3.9 (2.7-5.5)	12% (7.7-17%)	17% (9.5-26%)	87 (48-130)	7700 (4900-11000)
Greece	1995	Male	25 to 29	91% (83-95%)	9.2% (4.5-17%)	4.7 (3.3-6.3)	11% (7.5-15%)	18% (11-26%)	93 (54-130)	7600 (4900-10000)
Greece	1995	Male	30 to 34	93% (85-97%)	7.5% (3.3-15%)	5 (3.7-6.6)	9.8% (6.4-13%)	17% (9.8-24%)	82 (49-120)	6100 (3900-8300)
Greece	1995	Male	35 to 39	94% (88-97%)	6.4% (2.7-12%)	5.1 (3.7-7)	8.6% (5.3-12%)	14% (8-22%)	84 (47-130)	5700 (3600-8100)
Greece	1995	Male	40 to 44	95% (90-98%)	5.4% (2.2-10%)	5.9 (4.4-7.8)	9.4% (6.1-13%)	15% (9.1-21%)	120 (69-170)	6800 (4300-9300)
Greece	1995	Male	45 to 49	96% (92-98%)	4.3% (2.1-8.1%)	7.6 (5.6-9.9)	12% (8.2-16%)	18% (12-24%)	210 (140-280)	10000 (7100-13000)
Greece	1995	Male	50 to 54	96% (93-98%)	3.8% (2.1-6.8%)	8.4 (6.3-11)	13% (9.6-17%)	19% (13-24%)	270 (200-350)	11000 (8400-14000)
Greece	1995	Male	55 to 59	96% (93-98%)	4.1% (2.4-6.6%)	8.4 (6.4-11)	13% (9.7-17%)	18% (13-23%)	460 (340-590)	16000 (12000-21000)
Greece	1995	Male	60 to 64	95% (92-97%)	5.1% (3.2-7.8%)	8 (5.9-10)	13% (8.9-17%)	16% (11-21%)	690 (470-890)	21000 (14000-26000)
Greece	1995	Male	65 to 69	93% (90-96%)	6.6% (4-10%)	7.3 (5.7-9.2)	13% (9.1-16%)	16% (11-20%)	970 (680-1300)	24000 (17000-31000)
Greece	1995	Male	70 to 74	93% (89-95%)	7% (4.6-11%)	6.6 (4.9-8.6)	12% (8.3-17%)	15% (9.3-20%)	950 (600-1300)	19000 (12000-25000)
Greece	1995	Male	75 to 79	94% (89-96%)	6.4% (3.5-11%)	6.1 (4.3-8.2)	12% (6.6-16%)	13% (7.5-19%)	930 (520-1300)	14000 (8100-20000)
Greece	1995	Male	80 to 84	93% (88-96%)	7.1% (3.7-12%)	5.6 (4.1-7.5)	11% (6.7-16%)	13% (7.5-18%)	1000 (600-1500)	11000 (6600-16000)
Greece	1995	Male	85 to 89	91% (85-95%)	9% (5.1-15%)	5.1 (3.6-6.9)	9.6% (4.9-15%)	11% (5.4-17%)	770 (390-1200)	6000 (3100-9300)
Greece	1995	Male	90 to 94	90% (84-94%)	10% (5.6-16%)	4.8 (3.1-6.8)	8% (2.8-13%)	8.9% (2.8-15%)	310 (97-540)	1700 (600-2900)
Greece	1995	Male	95 plus	89% (82-95%)	11% (5.2-18%)	4.4 (2.4-6.9)	7.1% (0.86-14%)	7.7% (0.4-16%)	94 (4.8-190)	340 (41-660)
Greece	2000	Female	All Ages	77% (72-81%)	23% (19-28%)	2.2 (1.9-2.5)	2.7% (1.2-4.4%)	2.7% (-0.41-6%)	1300 (-200-2900)	38000 (18000-60000)
Greece	2000	Female	15 to 19	71% (56-84%)	29% (16-44%)	1.5 (0.99-2.2)	2.1% (1.1-3.2%)	7.1% (3.2-12%)	6.2 (2.8-11)	730 (420-1100)
Greece	2000	Female	20 to 24	77% (63-87%)	23% (13-37%)	1.9 (1.2-2.7)	2.7% (1.8-3.7%)	7.1% (3.2-11%)	8.6 (3.8-14)	1300 (810-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	2000	Female	25 to 29	81% (69-90%)	19% (10-31%)	1.9 (1.3-2.7)	2.2% (1.5-3.1%)	6.5% (3.1-10%)	8.7 (4.2-14)	1200 (770-1700)
Greece	2000	Female	30 to 34	82% (69-89%)	18% (11-31%)	2.2 (1.5-3)	2.1% (1.3-3%)	6.7% (3.8-10%)	12 (6.8-18)	1300 (800-1800)
Greece	2000	Female	35 to 39	80% (72-87%)	20% (13-28%)	2.7 (1.9-3.9)	2.6% (1.7-3.7%)	8.4% (5.1-12%)	20 (12-29)	1600 (1000-2200)
Greece	2000	Female	40 to 44	81% (73-87%)	19% (13-27%)	3.1 (2.2-4.2)	3.2% (2.1-4.5%)	9.6% (6.3-13%)	35 (23-48)	2200 (1500-3100)
Greece	2000	Female	45 to 49	84% (78-88%)	16% (12-22%)	3.2 (2.3-4.4)	3.7% (2.3-5.4%)	9.6% (6.2-13%)	52 (33-72)	2700 (1700-3800)
Greece	2000	Female	50 to 54	85% (80-89%)	15% (11-20%)	3.3 (2.3-4.4)	4.1% (2.4-6%)	9.3% (5.9-13%)	74 (47-100)	3300 (2000-4600)
Greece	2000	Female	55 to 59	84% (79-88%)	16% (12-21%)	3.3 (2.4-4.4)	4.3% (2.4-6.2%)	8.7% (5.2-12%)	84 (51-120)	3200 (1900-4600)
Greece	2000	Female	60 to 64	81% (75-86%)	19% (14-25%)	3.1 (2.2-4.3)	4.1% (2-6.5%)	7.4% (3.7-11%)	130 (68-210)	4300 (2100-6600)
Greece	2000	Female	65 to 69	76% (69-83%)	24% (17-31%)	2.8 (2.1-3.6)	3.8% (1.7-6.1%)	6% (2.7-9.6%)	190 (87-310)	4800 (2100-7800)
Greece	2000	Female	70 to 74	74% (66-81%)	26% (19-34%)	2.5 (1.7-3.4)	3.5% (0.81-6.4%)	5% (1.2-8.9%)	280 (70-510)	5600 (1300-10000)
Greece	2000	Female	75 to 79	75% (67-82%)	25% (18-33%)	2.3 (1.6-3.3)	2.6% (-0.72-6.2%)	3.3% (-1.1-7.9%)	230 (-77-550)	3600 (-1000-8300)
Greece	2000	Female	80 to 84	74% (64-82%)	26% (18-36%)	2.1 (1.5-3)	1.6% (-1.6-5%)	1.9% (-2.3-6%)	170 (-210-540)	1900 (-1900-6000)
Greece	2000	Female	85 to 89	71% (59-79%)	29% (21-41%)	1.9 (1.3-2.7)	0.67% (-2.4-3.8%)	0.63% (-3.1-4.4%)	58 (-280-410)	570 (-2000-3200)
Greece	2000	Female	90 to 94	68% (57-78%)	32% (22-43%)	1.7 (1.1-2.6)	-0.1% (-3-3.1%)	-0.35% (-3.8-3.5%)	-22 (-230-210)	-39 (-1100-1200)
Greece	2000	Female	95 plus	67% (53-79%)	33% (21-47%)	1.5 (0.8-2.6)	-0.66% (-3.6-2.9%)	-1% (-4.6-3.2%)	-27 (-120-85)	-69 (-380-300)
Greece	2000	Male	All Ages	91% (87-93%)	9.2% (6.8-13%)	4.6 (4-5.3)	10% (7.5-13%)	12% (8-16%)	6600 (4400-8800)	160000 (120000-200000)
Greece	2000	Male	15 to 19	83% (72-91%)	17% (8.8-28%)	2.4 (1.6-3.3)	6.3% (3.5-9.3%)	12% (5.1-18%)	31 (14-49)	2800 (1600-4100)
Greece	2000	Male	20 to 24	89% (80-94%)	11% (5.7-20%)	3.6 (2.5-5)	11% (6.9-15%)	15% (8-23%)	77 (40-120)	7100 (4600-9700)
Greece	2000	Male	25 to 29	92% (86-96%)	8.4% (4.1-14%)	4.4 (3.1-5.9)	10% (6.8-14%)	16% (8.7-23%)	79 (43-120)	6800 (4300-9400)
Greece	2000	Male	30 to 34	93% (87-97%)	6.6% (3-13%)	4.7 (3.5-6.2)	8.7% (5.6-12%)	14% (8.1-21%)	74 (42-110)	5900 (3700-8100)
Greece	2000	Male	35 to 39	94% (89-97%)	5.8% (2.6-11%)	4.7 (3.4-6.6)	7.7% (4.6-11%)	13% (6.5-20%)	72 (37-110)	5100 (3100-7400)
Greece	2000	Male	40 to 44	95% (90-98%)	4.9% (2.1-9.5%)	5.5 (4.2-7.4)	8.5% (5.1-12%)	14% (7.1-20%)	110 (59-170)	6600 (3800-9500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	2000	Male	45 to 49	96% (93-98%)	3.9% (2.1-7.2%)	7.1 (5.2-9.2)	11% (7.1-15%)	17% (10-22%)	200 (130-270)	9900 (6400-13000)
Greece	2000	Male	50 to 54	97% (94-98%)	3.4% (2-5.7%)	7.9 (5.9-10)	12% (8.7-16%)	17% (12-22%)	310 (220-410)	13000 (9400-17000)
Greece	2000	Male	55 to 59	96% (95-98%)	3.6% (2.3-5.4%)	7.9 (6-10)	12% (8.7-16%)	17% (12-21%)	360 (250-460)	13000 (9200-17000)
Greece	2000	Male	60 to 64	95% (93-97%)	4.6% (3-6.7%)	7.5 (5.5-9.8)	12% (7.7-16%)	15% (9.4-20%)	570 (360-750)	17000 (11000-22000)
Greece	2000	Male	65 to 69	94% (91-96%)	6.2% (4-8.8%)	6.8 (5.3-8.6)	12% (7.8-15%)	14% (9.5-19%)	830 (560-1100)	21000 (14000-27000)
Greece	2000	Male	70 to 74	93% (90-95%)	6.7% (4.6-9.8%)	6.2 (4.7-7.9)	11% (6.8-15%)	13% (7.9-18%)	1100 (660-1500)	21000 (13000-29000)
Greece	2000	Male	75 to 79	94% (90-96%)	6.2% (3.7-9.8%)	5.6 (4.1-7.5)	10% (5.3-15%)	12% (5.9-17%)	930 (470-1400)	14000 (7300-20000)
Greece	2000	Male	80 to 84	93% (88-96%)	7% (3.9-12%)	5.2 (3.8-6.8)	9.4% (5-14%)	11% (5.4-16%)	840 (430-1200)	9400 (5000-14000)
Greece	2000	Male	85 to 89	91% (85-95%)	9.1% (5.3-15%)	4.7 (3.3-6.4)	8.1% (3.8-13%)	9.1% (4-15%)	620 (270-1000)	4900 (2300-7900)
Greece	2000	Male	90 to 94	90% (84-94%)	10% (5.8-16%)	4.4 (2.9-6.3)	6.7% (1.8-12%)	7.4% (1.5-14%)	310 (67-580)	1700 (450-3100)
Greece	2000	Male	95 plus	89% (82-95%)	11% (5.3-18%)	4 (2.3-6.4)	5.9% (0.28-13%)	6.4% (-0.3-14%)	92 (-4.4-210)	340 (16-720)
Greece	2005	Female	All Ages	77% (73-80%)	23% (20-27%)	2.1 (1.8-2.5)	2.3% (1-3.8%)	2.2% (-0.59-5%)	1100 (-290-2500)	33000 (14000-54000)
Greece	2005	Female	15 to 19	73% (58-85%)	27% (15-42%)	1.5 (1-2.1)	1.9% (1.1-2.7%)	7.1% (3.2-12%)	4.5 (2.1-7.5)	570 (340-830)
Greece	2005	Female	20 to 24	80% (66-88%)	20% (12-34%)	1.9 (1.2-2.8)	2.5% (1.6-3.4%)	6.8% (3-11%)	6.7 (3-11)	1100 (700-1600)
Greece	2005	Female	25 to 29	83% (72-90%)	17% (9.7-28%)	1.9 (1.3-2.7)	2.1% (1.4-2.9%)	6.1% (3-9.6%)	7.8 (3.8-12)	1100 (710-1600)
Greece	2005	Female	30 to 34	83% (73-90%)	17% (10-27%)	2.2 (1.5-3)	2% (1.3-2.8%)	6.4% (3.6-9.9%)	10 (5.8-16)	1100 (730-1600)
Greece	2005	Female	35 to 39	81% (74-88%)	19% (12-26%)	2.7 (1.9-3.8)	2.4% (1.6-3.4%)	8% (4.8-12%)	19 (12-28)	1600 (1000-2300)
Greece	2005	Female	40 to 44	82% (76-87%)	18% (13-24%)	3.1 (2.2-4.1)	3% (1.9-4.2%)	9% (5.9-13%)	32 (21-45)	2100 (1400-2900)
Greece	2005	Female	45 to 49	85% (81-88%)	15% (12-19%)	3.2 (2.3-4.2)	3.4% (2-4.9%)	8.8% (5.5-12%)	50 (31-70)	2600 (1600-3700)
Greece	2005	Female	50 to 54	85% (82-89%)	15% (11-18%)	3.2 (2.3-4.3)	3.7% (2.1-5.4%)	8.4% (5-12%)	63 (38-88)	2800 (1700-4000)
Greece	2005	Female	55 to 59	84% (80-88%)	16% (12-20%)	3.2 (2.3-4.2)	3.9% (2.2-5.7%)	7.9% (4.7-11%)	89 (53-130)	3400 (2000-5000)
Greece	2005	Female	60 to 64	80% (76-84%)	20% (16-24%)	3 (2.1-4)	3.5% (1.5-5.6%)	6.5% (3.1-10%)	88 (42-140)	2800 (1300-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	2005	Female	65 to 69	74% (68-79%)	26% (21-32%)	2.6 (1.9-3.3)	3.2% (1.4-5%)	5.4% (2.4-8.5%)	150 (65-230)	3800 (1600-5900)
Greece	2005	Female	70 to 74	70% (64-76%)	30% (24-36%)	2.3 (1.6-3)	2.9% (0.68-5.3%)	4.4% (1.1-7.8%)	210 (53-380)	4300 (1000-7600)
Greece	2005	Female	75 to 79	72% (63-78%)	28% (22-37%)	2.1 (1.4-2.9)	2.1% (-0.56-5.1%)	2.8% (-0.84-6.7%)	240 (-75-590)	3800 (-980-9100)
Greece	2005	Female	80 to 84	69% (58-78%)	31% (22-42%)	1.9 (1.3-2.7)	1.3% (-1.4-4%)	1.5% (-2.1-5%)	160 (-210-530)	1900 (-2100-5900)
Greece	2005	Female	85 to 89	65% (53-74%)	35% (26-47%)	1.6 (1.1-2.3)	0.52% (-2-3.1%)	0.47% (-2.6-3.7%)	42 (-240-340)	440 (-1700-2600)
Greece	2005	Female	90 to 94	62% (50-72%)	38% (28-50%)	1.5 (0.91-2.2)	-0.17% (-2.5-2.5%)	-0.41% (-3.4-2.9%)	-26 (-210-180)	-67 (-1000-1000)
Greece	2005	Female	95 plus	61% (47-74%)	39% (26-53%)	1.3 (0.67-2.2)	-0.65% (-3.3-2.3%)	-1% (-4-2.5%)	-31 (-130-78)	-80 (-400-280)
Greece	2005	Male	All Ages	92% (89-94%)	8.2% (6.2-11%)	4.6 (4-5.3)	9.4% (7-12%)	11% (7.1-15%)	6100 (3900-8500)	150000 (110000-190000)
Greece	2005	Male	15 to 19	85% (75-92%)	15% (8.1-25%)	2.4 (1.6-3.4)	6.1% (3.5-9.1%)	12% (5.2-18%)	23 (10-36)	2200 (1200-3200)
Greece	2005	Male	20 to 24	90% (83-95%)	9.8% (5.2-17%)	3.6 (2.5-5)	11% (6.8-15%)	15% (7.8-22%)	65 (34-99)	6100 (4000-8500)
Greece	2005	Male	25 to 29	93% (87-96%)	7.4% (3.9-13%)	4.3 (3.1-5.7)	9.8% (6.3-13%)	15% (7.8-22%)	72 (38-100)	6300 (4100-8700)
Greece	2005	Male	30 to 34	94% (89-97%)	6.1% (2.8-11%)	4.6 (3.5-6)	8.2% (5.4-11%)	14% (7.9-20%)	64 (36-93)	5300 (3300-7200)
Greece	2005	Male	35 to 39	95% (90-98%)	5.3% (2.4-9.6%)	4.6 (3.3-6.4)	7.3% (4.5-11%)	12% (6.1-19%)	72 (37-110)	5200 (3200-7700)
Greece	2005	Male	40 to 44	95% (91-98%)	4.5% (2-8.5%)	5.4 (4-7.2)	8.1% (4.8-12%)	13% (6.6-20%)	100 (52-160)	6200 (3700-9100)
Greece	2005	Male	45 to 49	96% (94-98%)	3.5% (2-6.3%)	6.9 (5.1-8.9)	11% (6.8-14%)	16% (9.5-22%)	200 (110-270)	9800 (6100-13000)
Greece	2005	Male	50 to 54	97% (95-98%)	3% (1.8-5.1%)	7.7 (5.7-9.7)	12% (8-15%)	17% (11-22%)	290 (190-380)	12000 (8300-16000)
Greece	2005	Male	55 to 59	97% (95-98%)	3.3% (2.2-4.9%)	7.7 (5.9-9.8)	12% (7.9-15%)	16% (10-21%)	420 (270-550)	15000 (10000-20000)
Greece	2005	Male	60 to 64	96% (94-97%)	4.3% (3-6.1%)	7.3 (5.4-9.4)	11% (6.9-15%)	14% (8.6-19%)	440 (270-590)	14000 (8500-18000)
Greece	2005	Male	65 to 69	94% (92-96%)	5.8% (3.9-8.1%)	6.7 (5.2-8.3)	11% (7-14%)	13% (8.5-18%)	680 (430-910)	17000 (11000-23000)
Greece	2005	Male	70 to 74	94% (91-96%)	6.3% (4.5-8.6%)	6 (4.5-7.7)	10% (6-14%)	12% (7-17%)	940 (540-1300)	19000 (11000-26000)
Greece	2005	Male	75 to 79	94% (91-97%)	5.8% (3.4-9.2%)	5.5 (4-7.2)	9.1% (4.5-14%)	11% (5.1-16%)	1100 (520-1600)	17000 (8300-25000)
Greece	2005	Male	80 to 84	93% (88-96%)	6.7% (3.7-12%)	5 (3.7-6.6)	8.4% (4.2-13%)	9.5% (4.3-15%)	870 (400-1300)	10000 (4900-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	2005	Male	85 to 89	91% (86-95%)	8.8% (5.1-14%)	4.6 (3.2-6.1)	7.2% (3.2-12%)	8.1% (3.3-13%)	500 (210-840)	4100 (1800-6700)
Greece	2005	Male	90 to 94	90% (85-94%)	9.8% (5.7-15%)	4.2 (2.8-6.1)	6% (1.4-11%)	6.5% (1.1-13%)	250 (43-490)	1400 (330-2600)
Greece	2005	Male	95 plus	89% (82-95%)	11% (5.2-18%)	3.9 (2.3-6.2)	5.2% (-0.15-12%)	5.5% (-0.71-13%)	86 (-11-210)	320 (-9.5-740)
Greece	2010	Female	All Ages	78% (74-81%)	22% (19-26%)	2.1 (1.8-2.5)	2.1% (0.76-3.4%)	1.7% (-1.1-4.5%)	940 (-580-2400)	30000 (11000-50000)
Greece	2010	Female	15 to 19	76% (64-86%)	24% (14-36%)	1.6 (1.1-2.3)	1.8% (1.1-2.5%)	6.9% (3.3-11%)	3.7 (1.7-5.9)	500 (310-740)
Greece	2010	Female	20 to 24	82% (70-90%)	18% (10-30%)	2 (1.3-2.9)	2.5% (1.6-3.4%)	6.7% (3.1-11%)	5.5 (2.5-8.9)	920 (580-1300)
Greece	2010	Female	25 to 29	84% (74-91%)	16% (9.5-26%)	1.9 (1.3-2.7)	2% (1.3-2.7%)	5.8% (2.8-9.1%)	6.1 (3-9.7)	990 (620-1400)
Greece	2010	Female	30 to 34	84% (75-90%)	16% (9.6-25%)	2.2 (1.5-3.1)	1.9% (1.2-2.7%)	6% (3.1-9.5%)	9.4 (4.9-15)	1100 (680-1600)
Greece	2010	Female	35 to 39	82% (76-87%)	18% (13-24%)	2.7 (1.8-3.9)	2.3% (1.5-3.3%)	7.8% (4.4-11%)	17 (9.8-25)	1500 (920-2200)
Greece	2010	Female	40 to 44	82% (77-87%)	18% (13-23%)	3 (2.1-4.1)	2.8% (1.8-4%)	8.6% (5.3-12%)	31 (19-44)	2100 (1300-3000)
Greece	2010	Female	45 to 49	85% (81-88%)	15% (12-19%)	3.2 (2.3-4.2)	3.2% (1.9-4.7%)	8.4% (5.2-12%)	46 (29-65)	2500 (1500-3600)
Greece	2010	Female	50 to 54	86% (83-89%)	14% (11-17%)	3.2 (2.3-4.2)	3.5% (2-5.2%)	7.9% (4.7-11%)	66 (39-95)	3000 (1700-4400)
Greece	2010	Female	55 to 59	85% (82-88%)	15% (12-18%)	3.2 (2.3-4.3)	3.7% (1.9-5.4%)	7.4% (4.2-11%)	81 (46-120)	3100 (1700-4600)
Greece	2010	Female	60 to 64	80% (76-83%)	20% (17-24%)	2.9 (2.1-3.9)	3.2% (1.3-5.1%)	6% (2.7-9.3%)	97 (44-150)	3200 (1300-5100)
Greece	2010	Female	65 to 69	74% (68-78%)	26% (22-32%)	2.5 (1.8-3.3)	2.9% (1.1-4.7%)	5% (2.2-7.9%)	100 (44-160)	2700 (1100-4300)
Greece	2010	Female	70 to 74	70% (64-75%)	30% (25-36%)	2.2 (1.5-3)	2.5% (0.49-4.6%)	4% (0.87-7.2%)	170 (37-300)	3400 (680-6300)
Greece	2010	Female	75 to 79	71% (64-77%)	29% (23-36%)	2 (1.4-2.9)	1.8% (-0.59-4.6%)	2.5% (-0.88-6.1%)	190 (-69-490)	3100 (-990-7800)
Greece	2010	Female	80 to 84	68% (58-77%)	32% (23-42%)	1.8 (1.2-2.5)	1.1% (-1.5-3.7%)	1.2% (-2.1-4.6%)	150 (-250-550)	1800 (-2600-6300)
Greece	2010	Female	85 to 89	63% (52-72%)	37% (28-48%)	1.5 (0.98-2.2)	0.36% (-2-2.9%)	0.29% (-2.7-3.4%)	34 (-320-410)	410 (-2300-3200)
Greece	2010	Female	90 to 94	60% (49-70%)	40% (30-51%)	1.4 (0.84-2.1)	-0.28% (-2.4-2.2%)	-0.54% (-3.1-2.5%)	-37 (-210-170)	-120 (-1000-930)
Greece	2010	Female	95 plus	59% (45-73%)	41% (27-55%)	1.2 (0.62-2.1)	-0.74% (-3.2-2.2%)	-1.1% (-4.1-2.4%)	-39 (-140-81)	-100 (-440-300)
Greece	2010	Male	All Ages	92% (89-94%)	8% (6-11%)	4.6 (3.9-5.3)	8.7% (6.3-12%)	10% (5.9-14%)	5800 (3500-8300)	140000 (99000-180000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	2010	Male	15 to 19	86% (77-92%)	14% (7.7-23%)	2.4 (1.6-3.4)	5.7% (3.4-8.2%)	12% (5.5-18%)	18 (8.3-27)	1800 (1100-2500)
Greece	2010	Male	20 to 24	91% (84-95%)	9% (4.6-16%)	3.6 (2.5-5.1)	10% (6.7-14%)	16% (8.4-23%)	47 (25-68)	4600 (3000-6300)
Greece	2010	Male	25 to 29	93% (88-96%)	7.3% (3.7-12%)	4.2 (3-5.7)	9.3% (6-13%)	15% (7.9-21%)	61 (32-90)	5600 (3500-7600)
Greece	2010	Male	30 to 34	94% (89-97%)	6% (2.7-11%)	4.5 (3.3-6.1)	7.7% (5-11%)	13% (7.1-19%)	60 (34-91)	5100 (3300-7000)
Greece	2010	Male	35 to 39	95% (90-98%)	5.2% (2.3-9.9%)	4.5 (3.1-6.5)	6.7% (4.1-10%)	11% (5.3-19%)	60 (28-98)	4600 (2800-6900)
Greece	2010	Male	40 to 44	96% (92-98%)	4.3% (1.9-8%)	5.3 (3.8-7.1)	7.7% (4.5-11%)	13% (6.2-20%)	100 (51-160)	6300 (3800-9300)
Greece	2010	Male	45 to 49	97% (94-98%)	3.3% (1.8-5.7%)	6.8 (5-8.7)	11% (6.6-14%)	16% (9.3-22%)	190 (110-260)	9600 (5900-13000)
Greece	2010	Male	50 to 54	97% (95-98%)	2.9% (1.7-4.6%)	7.6 (5.7-9.5)	11% (7.7-15%)	16% (10-22%)	300 (190-390)	13000 (8300-16000)
Greece	2010	Male	55 to 59	97% (95-98%)	3.3% (2.2-4.7%)	7.5 (5.6-9.8)	11% (7.6-15%)	15% (9.6-20%)	380 (240-510)	14000 (9200-18000)
Greece	2010	Male	60 to 64	96% (94-97%)	4.4% (3-6.2%)	7.1 (5.2-9.4)	11% (6.4-14%)	13% (7.8-18%)	480 (280-650)	15000 (9200-20000)
Greece	2010	Male	65 to 69	94% (92-96%)	5.7% (3.9-7.7%)	6.5 (5-8.2)	9.8% (6.3-13%)	12% (7.6-17%)	500 (310-680)	13000 (8300-18000)
Greece	2010	Male	70 to 74	94% (92-96%)	6.2% (4.5-8.3%)	5.9 (4.4-7.7)	9.2% (5.2-13%)	11% (6-16%)	750 (410-1100)	16000 (8800-23000)
Greece	2010	Male	75 to 79	94% (91-96%)	5.7% (3.5-8.9%)	5.4 (3.9-7.3)	8.3% (4.1-13%)	9.7% (4.6-15%)	950 (450-1500)	15000 (7400-23000)
Greece	2010	Male	80 to 84	94% (89-96%)	6.4% (3.7-11%)	5 (3.6-6.7)	7.7% (3.6-12%)	8.8% (3.7-14%)	960 (400-1600)	11000 (5200-18000)
Greece	2010	Male	85 to 89	91% (86-95%)	8.9% (5.4-14%)	4.5 (3.1-6)	6.5% (2.2-11%)	7.3% (2.2-13%)	610 (180-1100)	5000 (1800-8500)
Greece	2010	Male	90 to 94	90% (85-94%)	9.9% (5.8-15%)	4.2 (2.7-5.9)	5.4% (0.84-11%)	5.9% (0.39-12%)	250 (17-500)	1400 (220-2800)
Greece	2010	Male	95 plus	89% (82-95%)	11% (5.1-18%)	3.9 (2.2-6.2)	4.7% (-0.53-11%)	5% (-1.2-12%)	81 (-20-200)	310 (-34-720)
Greece	2016	Female	All Ages	78% (75-82%)	22% (18-25%)	2.2 (1.6-2.9)	1.9% (0.15-3.9%)	1.3% (-2.3-5.2%)	810 (-1400-3200)	29000 (1900-59000)
Greece	2016	Female	15 to 19	74% (59-85%)	26% (15-41%)	1.5 (0.91-2.3)	1.6% (1-2.5%)	6.6% (3-11%)	2.8 (1.3-4.6)	400 (250-610)
Greece	2016	Female	20 to 24	79% (65-88%)	21% (12-35%)	1.9 (1.1-3)	2.4% (1.6-3.4%)	6.5% (3.2-11%)	4.4 (2.1-7.4)	800 (520-1100)
Greece	2016	Female	25 to 29	81% (69-90%)	19% (10-31%)	1.8 (1.1-2.7)	2% (1.3-2.8%)	5.9% (2.9-9.6%)	4.2 (2.1-6.6)	700 (440-990)
Greece	2016	Female	30 to 34	82% (69-89%)	18% (11-31%)	2.1 (1.3-3.1)	1.8% (1.1-2.6%)	5.8% (2.8-9.3%)	7.6 (3.6-12)	920 (540-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	2016	Female	35 to 39	81% (72-87%)	19% (13-28%)	2.6 (1.6-4)	2.2% (1.3-3.3%)	7.7% (3.8-12%)	15 (7.3-23)	1300 (760-2000)
Greece	2016	Female	40 to 44	82% (74-88%)	18% (12-26%)	3 (1.8-4.3)	2.7% (1.6-4.1%)	8.6% (4.7-12%)	28 (15-42)	1900 (1100-2800)
Greece	2016	Female	45 to 49	85% (78-89%)	15% (11-22%)	3.1 (2-4.5)	3.1% (1.7-4.8%)	8.3% (4.7-12%)	45 (26-68)	2500 (1400-3700)
Greece	2016	Female	50 to 54	86% (81-90%)	14% (9.9-19%)	3.2 (2.1-4.6)	3.4% (1.7-5.2%)	7.8% (4.3-12%)	64 (34-96)	2900 (1600-4500)
Greece	2016	Female	55 to 59	86% (80-89%)	14% (11-20%)	3.2 (2.1-4.6)	3.5% (1.6-5.7%)	7.2% (3.6-11%)	85 (42-130)	3300 (1600-5300)
Greece	2016	Female	60 to 64	84% (78-88%)	16% (12-22%)	3.1 (2-4.5)	3.2% (1-5.6%)	6% (2.2-10%)	89 (33-150)	2900 (950-5200)
Greece	2016	Female	65 to 69	80% (72-85%)	20% (15-28%)	2.8 (1.8-3.9)	2.9% (0.83-5.4%)	5.1% (1.7-9.2%)	130 (44-230)	3400 (1000-6300)
Greece	2016	Female	70 to 74	78% (71-83%)	22% (17-29%)	2.5 (1.6-3.7)	2.6% (0.13-5.4%)	4.1% (0.27-8.2%)	140 (9.4-280)	2900 (140-6000)
Greece	2016	Female	75 to 79	78% (70-85%)	22% (15-30%)	2.3 (1.5-3.5)	1.9% (-1.2-5.5%)	2.5% (-1.8-7.4%)	190 (-140-560)	3100 (-1900-8800)
Greece	2016	Female	80 to 84	77% (67-84%)	23% (16-33%)	2.1 (1.3-3.1)	1% (-2.2-4.5%)	1.2% (-3-5.5%)	160 (-410-750)	1900 (-4100-8400)
Greece	2016	Female	85 to 89	73% (62-81%)	27% (19-38%)	1.9 (1.1-2.9)	0.24% (-2.9-3.6%)	0.099% (-3.8-4.3%)	15 (-600-680)	360 (-4300-5400)
Greece	2016	Female	90 to 94	70% (59-79%)	30% (21-41%)	1.7 (1-2.7)	-0.51% (-3.4-2.7%)	-0.87% (-4.3-3%)	-95 (-480-330)	-350 (-2300-1900)
Greece	2016	Female	95 plus	69% (56-80%)	31% (20-44%)	1.5 (0.74-2.6)	-1.1% (-4-2.6%)	-1.6% (-5.1-2.8%)	-71 (-240-120)	-190 (-730-480)
Greece	2016	Male	All Ages	91% (88-93%)	9.2% (7-12%)	4.5 (3.3-5.9)	8.1% (4.8-11%)	9% (4-14%)	5800 (2600-9300)	130000 (74000-190000)
Greece	2016	Male	15 to 19	84% (72-92%)	16% (8.3-28%)	2.3 (1.4-3.5)	5.3% (3.2-7.7%)	11% (5.5-18%)	13 (6.3-21)	1400 (850-2000)
Greece	2016	Male	20 to 24	88% (79-94%)	12% (5.8-21%)	3.4 (2.1-5.1)	9.9% (6.5-14%)	15% (7.9-23%)	38 (20-58)	3900 (2500-5500)
Greece	2016	Male	25 to 29	91% (83-95%)	9.2% (4.5-17%)	4 (2.5-5.7)	9.2% (5.9-13%)	15% (8-22%)	40 (22-62)	3800 (2500-5300)
Greece	2016	Male	30 to 34	92% (85-97%)	7.5% (3.3-15%)	4.3 (2.8-6.1)	7.4% (4.6-10%)	13% (6.5-20%)	46 (23-74)	4100 (2500-5900)
Greece	2016	Male	35 to 39	94% (88-97%)	6.5% (2.9-12%)	4.4 (2.7-6.6)	6.6% (3.8-10%)	11% (4.9-19%)	51 (22-89)	4100 (2400-6400)
Greece	2016	Male	40 to 44	95% (89-98%)	5.4% (2.3-11%)	5.1 (3.3-7.4)	7.5% (4-12%)	12% (5.2-21%)	89 (36-150)	5700 (2900-8900)
Greece	2016	Male	45 to 49	96% (92-98%)	4.3% (2.2-7.9%)	6.5 (4.4-9.2)	9.9% (5.7-14%)	16% (8-23%)	180 (94-270)	9300 (5100-13000)
Greece	2016	Male	50 to 54	96% (93-98%)	3.8% (2.1-7%)	7.2 (5-10)	11% (6.3-15%)	16% (8.7-22%)	260 (150-370)	11000 (6700-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greece	2016	Male	55 to 59	96% (93-98%)	4.2% (2.5-6.7%)	7.2 (4.9-10)	11% (5.9-15%)	14% (7.6-20%)	360 (190-510)	13000 (7600-18000)
Greece	2016	Male	60 to 64	95% (92-97%)	5.1% (3.2-7.9%)	6.8 (4.6-9.7)	9.8% (5.4-14%)	13% (6.6-18%)	390 (200-570)	12000 (6500-18000)
Greece	2016	Male	65 to 69	93% (90-96%)	6.7% (4.1-10%)	6.2 (4.3-8.6)	9.1% (5-13%)	11% (6.2-16%)	550 (290-800)	14000 (7800-21000)
Greece	2016	Male	70 to 74	93% (89-95%)	7.1% (4.6-11%)	5.6 (3.7-8)	8.5% (4.2-13%)	10% (5-16%)	590 (270-920)	12000 (6000-19000)
Greece	2016	Male	75 to 79	94% (89-96%)	6.4% (3.5-11%)	5.2 (3.3-7.6)	7.7% (3.2-12%)	9.1% (3.6-15%)	820 (320-1400)	13000 (5500-22000)
Greece	2016	Male	80 to 84	93% (88-96%)	7.1% (3.8-12%)	4.8 (3.1-7)	7.2% (2.5-12%)	8.2% (2.6-14%)	1000 (300-1800)	12000 (4200-20000)
Greece	2016	Male	85 to 89	91% (85-95%)	9% (5.2-15%)	4.4 (2.8-6.5)	6.1% (1.5-11%)	6.9% (1.3-14%)	820 (160-1600)	6700 (1600-13000)
Greece	2016	Male	90 to 94	90% (84-94%)	10% (5.7-16%)	4.1 (2.5-6.1)	5.1% (0.25-11%)	5.6% (-0.22-12%)	410 (-16-880)	2300 (110-4800)
Greece	2016	Male	95 plus	89% (82-95%)	11% (5.3-18%)	3.8 (2-6.2)	4.4% (-1.2-11%)	4.6% (-1.9-12%)	120 (-52-320)	460 (-120-1100)
Iceland	1990	Female	All Ages	85% (81-88%)	15% (12-19%)	1.4 (1.2-1.6)	1.2% (0.023-2.4%)	-0.29% (-2.9-2.5%)	-2.4 (-24-20)	330 (5.6-670)
Iceland	1990	Female	15 to 19	85% (76-92%)	15% (8.1-24%)	1.2 (0.73-2)	1.9% (0.96-2.9%)	6.3% (2.1-11%)	0.2 (0.065-0.35)	22 (11-34)
Iceland	1990	Female	20 to 24	90% (83-95%)	9.5% (5-17%)	1.6 (0.84-2.4)	2.3% (1.5-3.3%)	6.8% (2.6-11%)	0.17 (0.065-0.28)	28 (17-41)
Iceland	1990	Female	25 to 29	92% (86-96%)	7.8% (4.2-14%)	1.6 (0.94-2.5)	2.1% (1.2-3%)	5.9% (1.9-9.9%)	0.23 (0.074-0.4)	30 (18-45)
Iceland	1990	Female	30 to 34	93% (88-97%)	6.9% (3.4-12%)	1.7 (1.1-2.5)	1.8% (1.1-2.6%)	6.2% (2.9-10%)	0.26 (0.12-0.42)	27 (17-39)
Iceland	1990	Female	35 to 39	93% (88-97%)	6.7% (3.3-12%)	1.8 (0.98-2.8)	2.3% (1.3-3.5%)	7.1% (3.6-11%)	0.51 (0.25-0.79)	36 (21-54)
Iceland	1990	Female	40 to 44	93% (88-96%)	6.9% (3.6-12%)	2 (1.2-3)	2.5% (1.4-3.9%)	7.8% (4.3-12%)	0.62 (0.34-0.94)	36 (21-54)
Iceland	1990	Female	45 to 49	93% (86-96%)	7.4% (3.7-14%)	2.3 (1.3-3.7)	3.6% (1.9-5.9%)	8.3% (4.4-13%)	1.2 (0.6-1.9)	54 (27-90)
Iceland	1990	Female	50 to 54	91% (84-94%)	9.3% (5.5-16%)	2.5 (1.4-3.9)	3.9% (1.9-6.3%)	7.7% (4-12%)	1.4 (0.7-2.1)	54 (26-86)
Iceland	1990	Female	55 to 59	88% (81-93%)	12% (7-19%)	2.4 (1.4-3.7)	3.5% (1.4-6%)	6.2% (2.6-10%)	1.8 (0.77-3)	61 (24-100)
Iceland	1990	Female	60 to 64	84% (76-91%)	16% (9.2-24%)	2.3 (1.2-3.6)	2.3% (0.0082-5.2%)	3.8% (0.19-8%)	1.6 (0.08-3.5)	47 (0.16-100)
Iceland	1990	Female	65 to 69	79% (70-88%)	21% (12-30%)	2 (1.3-2.9)	0.98% (-1.1-3.4%)	1.5% (-1.6-5.1%)	0.86 (-0.94-2.9)	20 (-23-73)
Iceland	1990	Female	70 to 74	77% (66-85%)	23% (15-34%)	1.8 (0.99-2.8)	-0.087% (-2.5-2.9%)	-0.13% (-3.4-3.9%)	-0.11 (-2.8-3.1)	-1.8 (-55-62)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	1990	Female	75 to 79	76% (64-85%)	24% (15-36%)	1.6 (0.86-2.7)	-1% (-3.7-2.1%)	-1.5% (-4.8-2.7%)	-1.7 (-5.7-3.1)	-23 (-81-50)
Iceland	1990	Female	80 to 84	75% (62-85%)	25% (15-38%)	1.6 (0.9-2.5)	-1.5% (-4.1-1.5%)	-2.1% (-5.3-1.6%)	-2.8 (-7.1-2.1)	-28 (-74-28)
Iceland	1990	Female	85 to 89	74% (61-84%)	26% (16-39%)	1.5 (0.86-2.5)	-1.7% (-4.3-1.4%)	-2.3% (-5.3-1.4%)	-3.5 (-8.4-2.2)	-25 (-61-20)
Iceland	1990	Female	90 to 94	75% (62-85%)	25% (15-38%)	1.5 (0.78-2.7)	-1.6% (-4.1-1.6%)	-2.2% (-5.3-1.7%)	-2 (-4.9-1.6)	-9.7 (-24-9.3)
Iceland	1990	Female	95 plus	76% (60-88%)	24% (12-40%)	1.5 (0.64-2.8)	-1.6% (-4.5-2.6%)	-2.2% (-5.8-2.7%)	-1 (-2.8-1.3)	-3.1 (-9-5)
Iceland	1990	Male	All Ages	89% (86-92%)	11% (7.8-14%)	1.6 (1.4-1.8)	2.4% (0.79-4.1%)	0.22% (-2.6-3.1%)	2 (-24-28)	750 (240-1300)
Iceland	1990	Male	15 to 19	86% (77-92%)	14% (7.8-23%)	1.3 (0.77-1.9)	5% (2.6-7.7%)	7.9% (2.9-13%)	0.88 (0.33-1.5)	80 (39-120)
Iceland	1990	Male	20 to 24	90% (83-94%)	9.8% (5.6-17%)	1.8 (1.1-2.7)	8.3% (5.1-12%)	11% (5-18%)	1.3 (0.59-2.1)	130 (81-190)
Iceland	1990	Male	25 to 29	93% (88-96%)	7.3% (4.1-12%)	2 (1.3-2.8)	7.4% (4.9-10%)	11% (5.1-17%)	1.3 (0.61-2)	120 (79-170)
Iceland	1990	Male	30 to 34	94% (89-97%)	5.6% (2.9-11%)	2.1 (1.4-2.9)	5.9% (3.7-8.1%)	9.8% (4.6-15%)	0.96 (0.46-1.5)	90 (57-130)
Iceland	1990	Male	35 to 39	95% (92-98%)	4.7% (2.5-8.4%)	2.1 (1.3-2.9)	5.3% (3.4-7.5%)	9% (4.2-14%)	1 (0.49-1.7)	83 (54-120)
Iceland	1990	Male	40 to 44	96% (93-98%)	4.2% (2-7.5%)	2.2 (1.4-3.1)	5.1% (3.1-7.4%)	8.4% (3.9-13%)	1.3 (0.6-2)	82 (49-120)
Iceland	1990	Male	45 to 49	96% (92-98%)	3.9% (1.9-7.5%)	2.6 (1.6-3.8)	4.9% (2.4-7.8%)	7% (2.7-12%)	1.4 (0.52-2.3)	75 (38-120)
Iceland	1990	Male	50 to 54	96% (92-98%)	4.4% (2.2-7.7%)	2.8 (1.8-4.1)	3.7% (1.1-6.8%)	4.7% (0.26-9.5%)	1.2 (0.066-2.5)	59 (17-110)
Iceland	1990	Male	55 to 59	94% (90-97%)	5.6% (2.9-9.8%)	2.9 (1.9-4.1)	2.7% (-0.37-6.3%)	2.9% (-1.7-8.1%)	1.3 (-0.69-3.6)	57 (-8.5-130)
Iceland	1990	Male	60 to 64	93% (89-97%)	6.6% (3.4-11%)	2.8 (1.7-4.2)	1.7% (-1.5-5.7%)	1.6% (-2.8-6.8%)	1 (-1.8-4.4)	42 (-38-140)
Iceland	1990	Male	65 to 69	93% (88-97%)	7.1% (3.4-12%)	2.5 (1.7-3.5)	0.29% (-2.6-3.4%)	-0.12% (-3.9-3.8%)	-0.11 (-3.6-3.7)	8.2 (-76-95)
Iceland	1990	Male	70 to 74	92% (87-96%)	7.9% (4.3-13%)	2.2 (1.4-3.3)	-0.46% (-3.3-2.5%)	-0.94% (-4.5-2.7%)	-1.1 (-5.1-2.9)	-12 (-86-67)
Iceland	1990	Male	75 to 79	91% (85-95%)	9.1% (4.8-15%)	2 (1.1-3.1)	-1% (-3.8-2.1%)	-1.6% (-4.9-2.1%)	-2.1 (-6.6-2.8)	-24 (-87-48)
Iceland	1990	Male	80 to 84	90% (83-95%)	10% (5.1-17%)	1.9 (1.2-2.8)	-1.3% (-3.9-1.4%)	-1.9% (-5-1.5%)	-2.6 (-6.7-2)	-23 (-67-25)
Iceland	1990	Male	85 to 89	89% (82-94%)	11% (5.6-18%)	1.8 (1.1-2.8)	-1.3% (-3.6-1.2%)	-1.8% (-4.6-1.1%)	-2.2 (-5.4-1.3)	-14 (-38-12)
Iceland	1990	Male	90 to 94	90% (83-95%)	10% (4.9-17%)	1.8 (1-2.9)	-1.4% (-3.6-1.2%)	-2% (-4.8-1.1%)	-1.1 (-2.7-0.62)	-4.8 (-13-4.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	1990	Male	95 plus	91% (82-96%)	9.5% (3.9-18%)	1.7 (0.85-3.1)	-1.4% (-3.9-1.9%)	-2.1% (-5.1-1.7%)	-0.41 (-1-0.33)	-1.1 (-3.1-1.5)
Iceland	1995	Female	All Ages	85% (81-88%)	15% (12-19%)	1.3 (1.2-1.5)	1.1% (0.18-2.2%)	-0.19% (-2.5-2.5%)	-1.7 (-22-23)	340 (55-660)
Iceland	1995	Female	15 to 19	85% (76-92%)	15% (8.3-24%)	1.2 (0.69-1.9)	1.7% (0.84-2.7%)	4.8% (1.4-8.1%)	0.18 (0.055-0.3)	20 (10-31)
Iceland	1995	Female	20 to 24	90% (83-95%)	9.6% (5-17%)	1.5 (0.82-2.4)	2.2% (1.4-3.2%)	5.1% (1.8-8.8%)	0.18 (0.066-0.31)	29 (18-43)
Iceland	1995	Female	25 to 29	92% (86-96%)	8% (4.3-14%)	1.5 (0.89-2.4)	1.8% (1.2-2.6%)	4.6% (1.7-7.8%)	0.17 (0.059-0.28)	25 (16-38)
Iceland	1995	Female	30 to 34	93% (88-96%)	7% (3.6-12%)	1.6 (1-2.4)	1.7% (1-2.3%)	4.9% (2.3-8.1%)	0.23 (0.11-0.38)	26 (16-37)
Iceland	1995	Female	35 to 39	93% (88-97%)	6.7% (3.4-12%)	1.7 (0.96-2.6)	1.9% (1.2-2.8%)	6.1% (3.4-9.2%)	0.42 (0.23-0.65)	32 (20-48)
Iceland	1995	Female	40 to 44	93% (88-96%)	7% (3.7-12%)	1.9 (1.1-2.9)	2.3% (1.3-3.5%)	6.9% (3.9-10%)	0.66 (0.38-0.97)	39 (23-59)
Iceland	1995	Female	45 to 49	92% (86-96%)	7.6% (3.8-14%)	2.2 (1.3-3.5)	3.2% (1.7-5.1%)	7.7% (4.3-12%)	1.2 (0.68-1.9)	57 (31-91)
Iceland	1995	Female	50 to 54	91% (84-94%)	9.4% (5.7-16%)	2.3 (1.4-3.6)	3.7% (2-5.8%)	7.3% (4.1-11%)	1.6 (0.9-2.5)	64 (34-100)
Iceland	1995	Female	55 to 59	88% (81-93%)	12% (6.9-19%)	2.3 (1.4-3.5)	3.5% (1.6-5.6%)	6.3% (3.2-9.6%)	1.7 (0.86-2.6)	56 (26-92)
Iceland	1995	Female	60 to 64	84% (76-90%)	16% (9.6-24%)	2.2 (1.2-3.5)	2.4% (0.35-4.8%)	4% (0.83-7.7%)	1.5 (0.32-3)	45 (7-90)
Iceland	1995	Female	65 to 69	79% (69-88%)	21% (12-31%)	1.9 (1.2-2.9)	1.2% (-0.66-3.4%)	1.9% (-0.83-4.9%)	1.1 (-0.51-3)	26 (-14-75)
Iceland	1995	Female	70 to 74	77% (66-85%)	23% (15-34%)	1.7 (0.96-2.7)	0.13% (-1.9-2.6%)	0.24% (-2.7-3.7%)	0.21 (-2.3-3.2)	3.2 (-46-63)
Iceland	1995	Female	75 to 79	76% (64-85%)	24% (15-36%)	1.5 (0.82-2.6)	-0.75% (-3.1-2.2%)	-1% (-4-2.7%)	-1.2 (-4.8-3.3)	-18 (-69-50)
Iceland	1995	Female	80 to 84	75% (61-86%)	25% (14-39%)	1.5 (0.85-2.4)	-1.4% (-3.9-1.5%)	-1.9% (-5-1.7%)	-2.9 (-7.5-2.6)	-30 (-79-32)
Iceland	1995	Female	85 to 89	74% (60-84%)	26% (16-40%)	1.5 (0.81-2.4)	-1.6% (-4-1.4%)	-2.1% (-5-1.5%)	-3.5 (-8.3-2.4)	-25 (-61-22)
Iceland	1995	Female	90 to 94	75% (62-85%)	25% (15-38%)	1.5 (0.76-2.6)	-1.6% (-3.9-1.9%)	-2.1% (-5-2.1%)	-2.4 (-5.7-2.4)	-12 (-29-14)
Iceland	1995	Female	95 plus	76% (59-88%)	24% (12-41%)	1.5 (0.59-2.9)	-1.4% (-4.2-3%)	-1.9% (-5.2-3.2%)	-0.98 (-2.7-1.7)	-3 (-8.7-6.1)
Iceland	1995	Male	All Ages	90% (87-92%)	10% (7.7-13%)	1.5 (1.3-1.8)	2.2% (0.77-3.6%)	0.16% (-2.4-2.7%)	1.5 (-23-26)	700 (230-1200)
Iceland	1995	Male	15 to 19	86% (77-92%)	14% (7.6-23%)	1.2 (0.74-1.8)	4.1% (2.1-6.2%)	6.2% (2.3-10%)	0.61 (0.22-1)	61 (31-91)
Iceland	1995	Male	20 to 24	90% (83-94%)	9.8% (5.6-17%)	1.8 (1.1-2.6)	7.1% (4.6-9.9%)	8.8% (3.6-14%)	0.92 (0.38-1.5)	110 (66-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	1995	Male	25 to 29	93% (87-96%)	7.4% (4.2-13%)	1.9 (1.2-2.8)	6.4% (4.2-8.8%)	8.5% (3.8-14%)	0.85 (0.37-1.4)	94 (61-130)
Iceland	1995	Male	30 to 34	94% (89-97%)	5.7% (2.9-11%)	2 (1.3-2.9)	5.2% (3.3-7.2%)	7.8% (3.4-13%)	0.86 (0.39-1.4)	84 (53-120)
Iceland	1995	Male	35 to 39	95% (92-98%)	4.7% (2.5-8.4%)	1.9 (1.2-2.8)	4.8% (3-6.7%)	7.5% (3.3-11%)	0.99 (0.45-1.5)	82 (53-110)
Iceland	1995	Male	40 to 44	96% (92-98%)	4.2% (2-7.8%)	2.1 (1.4-2.9)	4.8% (2.9-7%)	7.4% (3.6-12%)	1.3 (0.64-2.1)	88 (53-130)
Iceland	1995	Male	45 to 49	96% (92-98%)	4% (1.9-7.7%)	2.5 (1.6-3.6)	4.5% (2.4-6.8%)	6.6% (2.7-11%)	1.5 (0.57-2.4)	83 (45-120)
Iceland	1995	Male	50 to 54	96% (92-98%)	4.5% (2.3-8.4%)	2.7 (1.7-3.8)	3.7% (1.5-6.3%)	4.9% (1.3-9.2%)	1.3 (0.34-2.3)	64 (25-110)
Iceland	1995	Male	55 to 59	94% (90-97%)	5.7% (2.9-9.7%)	2.8 (1.8-4)	3% (0.35-5.9%)	3.5% (-0.42-7.7%)	1.3 (-0.17-2.9)	56 (6.7-110)
Iceland	1995	Male	60 to 64	93% (89-97%)	6.6% (3.4-11%)	2.7 (1.6-4)	1.9% (-0.81-5%)	1.9% (-1.9-6.2%)	1.2 (-1.2-3.7)	44 (-20-120)
Iceland	1995	Male	65 to 69	93% (87-97%)	7.1% (3.4-13%)	2.4 (1.6-3.4)	0.56% (-1.9-3.3%)	0.26% (-3.1-3.8%)	0.24 (-2.8-3.4)	16 (-55-90)
Iceland	1995	Male	70 to 74	92% (87-96%)	8% (4.4-13%)	2.1 (1.3-3.1)	-0.32% (-2.7-2.4%)	-0.74% (-3.8-2.6%)	-0.92 (-4.8-3.3)	-9.2 (-82-69)
Iceland	1995	Male	75 to 79	91% (85-95%)	9% (4.8-15%)	1.9 (1.1-2.9)	-0.88% (-3.3-1.8%)	-1.4% (-4.3-1.9%)	-2 (-6.1-2.7)	-22 (-82-44)
Iceland	1995	Male	80 to 84	90% (82-95%)	10% (5-18%)	1.8 (1.1-2.6)	-1.3% (-3.7-1.4%)	-1.9% (-4.6-1.4%)	-2.6 (-6.6-2)	-24 (-66-24)
Iceland	1995	Male	85 to 89	90% (81-95%)	10% (5.4-19%)	1.7 (1.1-2.6)	-1.3% (-3.5-1.1%)	-1.9% (-4.5-1.1%)	-2.2 (-5.3-1.3)	-14 (-37-12)
Iceland	1995	Male	90 to 94	90% (83-95%)	10% (4.9-17%)	1.7 (0.98-2.7)	-1.3% (-3.4-1.1%)	-1.9% (-4.4-0.93%)	-1.3 (-3.1-0.65)	-5.7 (-15-4.8)
Iceland	1995	Male	95 plus	90% (82-96%)	9.5% (3.9-18%)	1.7 (0.82-2.9)	-1.3% (-3.7-1.7%)	-1.9% (-4.8-1.4%)	-0.5 (-1.3-0.38)	-1.4 (-3.9-1.8)
Iceland	2000	Female	All Ages	86% (83-89%)	14% (11-17%)	1.4 (1.2-1.6)	1.3% (0.41-2.3%)	0.18% (-2-2.8%)	1.6 (-18-25)	390 (120-690)
Iceland	2000	Female	15 to 19	87% (79-92%)	13% (7.6-21%)	1.2 (0.71-1.9)	1.8% (0.88-2.7%)	5.9% (1.8-9.9%)	0.17 (0.056-0.29)	20 (10-30)
Iceland	2000	Female	20 to 24	92% (85-95%)	8.3% (4.5-15%)	1.6 (0.86-2.5)	2.4% (1.5-3.4%)	6% (2.2-10%)	0.22 (0.079-0.37)	32 (19-47)
Iceland	2000	Female	25 to 29	93% (89-96%)	6.9% (3.8-11%)	1.6 (0.92-2.5)	1.9% (1.2-2.7%)	5.8% (2.1-9.7%)	0.18 (0.067-0.31)	27 (16-38)
Iceland	2000	Female	30 to 34	94% (90-97%)	6.2% (3.2-10%)	1.7 (1-2.6)	1.8% (1.2-2.6%)	6.3% (3.1-10%)	0.25 (0.13-0.4)	26 (16-39)
Iceland	2000	Female	35 to 39	94% (90-97%)	6% (3.2-9.9%)	1.8 (0.97-2.8)	1.8% (1.1-2.6%)	7.1% (4.1-10%)	0.35 (0.2-0.52)	29 (17-42)
Iceland	2000	Female	40 to 44	94% (90-97%)	6.1% (3.4-9.8%)	2 (1.2-2.9)	2.5% (1.4-3.8%)	7.9% (4.7-11%)	0.8 (0.48-1.1)	47 (28-68)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	2000	Female	45 to 49	93% (88-97%)	6.5% (3.4-12%)	2.3 (1.4-3.7)	3% (1.6-4.6%)	8.3% (4.8-12%)	1.1 (0.66-1.7)	55 (31-85)
Iceland	2000	Female	50 to 54	91% (86-95%)	8.6% (5.4-14%)	2.4 (1.5-3.8)	3.4% (1.8-5.2%)	7.6% (4.5-11%)	1.6 (0.93-2.3)	63 (35-96)
Iceland	2000	Female	55 to 59	88% (82-93%)	12% (7.3-18%)	2.4 (1.4-3.5)	3.4% (1.5-5.3%)	6.3% (3.3-9.5%)	1.8 (0.95-2.8)	63 (30-99)
Iceland	2000	Female	60 to 64	84% (78-90%)	16% (9.7-22%)	2.3 (1.2-3.5)	2.4% (0.51-4.8%)	4.2% (1.1-7.9%)	1.4 (0.38-2.7)	42 (8.7-82)
Iceland	2000	Female	65 to 69	77% (68-87%)	23% (13-32%)	1.9 (1.2-2.8)	1.5% (-0.092-3.4%)	2.5% (0.00052-5.3%)	1.4 (0.00029-2.8)	32 (-2.1-71)
Iceland	2000	Female	70 to 74	73% (63-83%)	27% (17-37%)	1.6 (0.91-2.6)	0.64% (-1.1-2.9%)	1% (-1.5-4%)	0.87 (-1.2-3.4)	15 (-27-68)
Iceland	2000	Female	75 to 79	75% (63-84%)	25% (16-37%)	1.6 (0.8-2.6)	-0.27% (-2.3-2.5%)	-0.38% (-3.1-3.2%)	-0.43 (-3.5-3.7)	-6.6 (-53-60)
Iceland	2000	Female	80 to 84	74% (61-85%)	26% (15-39%)	1.5 (0.86-2.4)	-0.93% (-3.1-1.8%)	-1.2% (-4-2.2%)	-1.9 (-6.2-3.6)	-20 (-67-41)
Iceland	2000	Female	85 to 89	74% (60-84%)	26% (16-40%)	1.5 (0.83-2.4)	-1.3% (-3.6-1.3%)	-1.8% (-4.5-1.4%)	-2.9 (-7.4-2.5)	-21 (-56-20)
Iceland	2000	Female	90 to 94	75% (62-85%)	25% (15-38%)	1.5 (0.77-2.6)	-1.5% (-3.8-1.6%)	-2% (-4.9-1.7%)	-2.3 (-5.6-2)	-11 (-28-12)
Iceland	2000	Female	95 plus	76% (59-88%)	24% (12-41%)	1.5 (0.6-2.7)	-1.2% (-3.9-3%)	-1.7% (-4.8-3.3%)	-1 (-2.9-2)	-3 (-9.5-7.3)
Iceland	2000	Male	All Ages	91% (89-93%)	8.7% (6.5-11%)	1.7 (1.5-1.9)	2.6% (1.2-4.2%)	0.71% (-1.8-3.2%)	6.7 (-17-30)	830 (370-1300)
Iceland	2000	Male	15 to 19	89% (82-93%)	11% (6.7-18%)	1.3 (0.79-1.9)	4.3% (2.3-6.4%)	7.7% (2.8-13%)	0.52 (0.19-0.87)	55 (29-81)
Iceland	2000	Male	20 to 24	92% (87-95%)	7.9% (4.8-13%)	1.9 (1.1-2.8)	8% (5-11%)	11% (4.5-17%)	1.1 (0.45-1.8)	120 (73-170)
Iceland	2000	Male	25 to 29	94% (91-96%)	5.8% (3.6-8.8%)	2 (1.3-3)	7.1% (4.6-9.9%)	11% (4.8-17%)	0.99 (0.45-1.6)	100 (68-150)
Iceland	2000	Male	30 to 34	95% (93-98%)	4.5% (2.5-7.5%)	2.1 (1.3-3)	5.7% (3.7-7.9%)	10% (4.5-16%)	0.82 (0.38-1.3)	80 (51-110)
Iceland	2000	Male	35 to 39	96% (94-98%)	3.8% (2.2-6.1%)	2 (1.3-2.9)	5% (3.2-7%)	9.1% (4.2-14%)	0.93 (0.43-1.4)	79 (51-110)
Iceland	2000	Male	40 to 44	97% (94-98%)	3.4% (1.7-5.7%)	2.2 (1.5-3)	4.9% (3.1-6.9%)	8.5% (4.3-13%)	1.4 (0.7-2.1)	92 (58-130)
Iceland	2000	Male	45 to 49	97% (94-98%)	3.3% (1.7-5.9%)	2.6 (1.7-3.7)	4.8% (2.7-7.3%)	7.8% (3.6-12%)	1.7 (0.75-2.7)	94 (54-140)
Iceland	2000	Male	50 to 54	96% (94-98%)	3.6% (2-6%)	2.9 (1.9-4)	4.2% (2.1-6.7%)	6.2% (2.5-10%)	1.8 (0.73-3.1)	89 (44-140)
Iceland	2000	Male	55 to 59	95% (92-97%)	4.9% (2.7-8%)	2.9 (1.9-4.1)	3.5% (0.99-6.4%)	4.5% (0.57-8.6%)	1.9 (0.23-3.6)	75 (21-130)
Iceland	2000	Male	60 to 64	94% (90-97%)	6.1% (3.4-9.6%)	2.8 (1.8-4.1)	2.6% (-0.015-5.7%)	3% (-0.81-7.3%)	1.5 (-0.39-3.6)	52 (-0.33-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	2000	Male	65 to 69	93% (89-97%)	6.6% (3.5-11%)	2.5 (1.7-3.5)	1.1% (-1.3-3.5%)	1% (-2.3-4.2%)	0.79 (-1.8-3.3)	28 (-32-93)
Iceland	2000	Male	70 to 74	92% (88-96%)	7.7% (4.5-12%)	2.2 (1.3-3.1)	0.26% (-2.1-2.8%)	-0.0088% (-3-3.2%)	-0.011 (-3.6-3.8)	7.7 (-61-81)
Iceland	2000	Male	75 to 79	91% (85-95%)	9.2% (5.1-15%)	1.9 (1.2-3)	-0.32% (-2.5-2.2%)	-0.71% (-3.4-2.4%)	-1.1 (-5.1-3.5)	-8.8 (-68-60)
Iceland	2000	Male	80 to 84	90% (82-95%)	10% (5.1-18%)	1.8 (1.2-2.7)	-0.82% (-2.9-1.7%)	-1.3% (-4-1.7%)	-1.9 (-5.9-2.5)	-16 (-59-33)
Iceland	2000	Male	85 to 89	90% (82-94%)	10% (5.6-18%)	1.8 (1.1-2.7)	-1.1% (-3.1-1.3%)	-1.6% (-4.1-1.1%)	-1.9 (-4.9-1.4)	-11 (-34-14)
Iceland	2000	Male	90 to 94	90% (83-95%)	10% (4.9-17%)	1.7 (1-2.8)	-1.2% (-3.5-1.3%)	-1.8% (-4.4-1.2%)	-1.2 (-3.1-0.79)	-5.2 (-15-5.4)
Iceland	2000	Male	95 plus	90% (82-96%)	9.6% (3.9-18%)	1.7 (0.88-2.9)	-1.3% (-3.8-1.7%)	-1.9% (-4.9-1.6%)	-0.58 (-1.5-0.47)	-1.6 (-4.6-2.1)
Iceland	2005	Female	All Ages	86% (83-89%)	14% (11-17%)	1.6 (1.4-1.8)	1.6% (0.76-2.6%)	0.93% (-1.3-3.5%)	8.4 (-12-33)	500 (230-790)
Iceland	2005	Female	15 to 19	87% (79-92%)	13% (7.6-21%)	1.4 (0.77-2.1)	1.6% (0.88-2.5%)	6% (2.2-9.9%)	0.13 (0.047-0.22)	17 (9.4-26)
Iceland	2005	Female	20 to 24	92% (86-95%)	8.2% (4.6-14%)	1.7 (0.94-2.8)	2.3% (1.5-3.3%)	6.2% (2.4-10%)	0.18 (0.069-0.3)	29 (18-42)
Iceland	2005	Female	25 to 29	93% (89-96%)	6.8% (3.7-11%)	1.8 (1-2.9)	1.8% (1.2-2.6%)	6.1% (2.7-9.9%)	0.16 (0.068-0.26)	26 (16-36)
Iceland	2005	Female	30 to 34	94% (90-97%)	6.1% (3.2-10%)	1.9 (1.1-2.9)	1.9% (1.2-2.8%)	6.9% (3.7-11%)	0.27 (0.14-0.42)	28 (18-40)
Iceland	2005	Female	35 to 39	94% (90-97%)	6% (3.1-10%)	2 (1.1-3.2)	2% (1.2-3%)	8% (4.7-12%)	0.38 (0.22-0.57)	31 (20-45)
Iceland	2005	Female	40 to 44	94% (90-97%)	6.1% (3.3-10%)	2.2 (1.4-3.3)	2.6% (1.5-3.8%)	8.8% (5.5-12%)	0.82 (0.5-1.1)	49 (30-70)
Iceland	2005	Female	45 to 49	94% (88-97%)	6.4% (3.4-12%)	2.6 (1.6-4)	3.1% (1.8-4.7%)	9.3% (6-13%)	1.3 (0.8-1.8)	62 (37-92)
Iceland	2005	Female	50 to 54	91% (86-95%)	8.5% (5.4-14%)	2.7 (1.7-4)	3.6% (2.1-5.4%)	8.7% (5.6-12%)	1.8 (1.2-2.5)	75 (44-110)
Iceland	2005	Female	55 to 59	88% (81-93%)	12% (7.3-19%)	2.7 (1.6-3.9)	3.7% (2-5.6%)	7.4% (4.6-10%)	2.3 (1.4-3.2)	79 (45-120)
Iceland	2005	Female	60 to 64	84% (77-90%)	16% (9.7-23%)	2.5 (1.4-3.9)	2.8% (0.99-5.2%)	5.3% (2.3-8.8%)	1.8 (0.81-3.1)	55 (19-96)
Iceland	2005	Female	65 to 69	77% (67-86%)	23% (14-33%)	2.1 (1.3-3.1)	2.1% (0.53-3.9%)	3.5% (1.1-6.1%)	1.6 (0.54-2.9)	39 (10-74)
Iceland	2005	Female	70 to 74	73% (62-83%)	27% (17-38%)	1.8 (1-2.8)	1.2% (-0.55-3.5%)	1.9% (-0.58-5%)	1.5 (-0.44-3.8)	27 (-13-81)
Iceland	2005	Female	75 to 79	74% (63-84%)	26% (16-37%)	1.7 (0.88-2.9)	0.31% (-1.7-3.1%)	0.45% (-2.3-4.1%)	0.54 (-2.7-5)	8 (-44-79)
Iceland	2005	Female	80 to 84	74% (60-85%)	26% (15-40%)	1.7 (0.94-2.6)	-0.22% (-2.3-2.5%)	-0.31% (-3-3.1%)	-0.51 (-4.9-5)	-5.3 (-54-59)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	2005	Female	85 to 89	74% (61-84%)	26% (16-39%)	1.7 (0.94-2.6)	-0.65% (-2.9-2.1%)	-0.92% (-3.7-2.4%)	-1.6 (-6.4-4.3)	-11 (-49-35)
Iceland	2005	Female	90 to 94	75% (62-85%)	25% (15-38%)	1.7 (0.86-2.9)	-0.85% (-3.3-2.6%)	-1.2% (-4.2-2.9%)	-1.5 (-5.3-3.6)	-6.9 (-27-22)
Iceland	2005	Female	95 plus	76% (60-88%)	24% (12-40%)	1.7 (0.68-3.1)	-0.66% (-3.5-3.9%)	-0.99% (-4.3-4.4%)	-0.67 (-2.9-3)	-1.8 (-9.7-11)
Iceland	2005	Male	All Ages	91% (89-93%)	8.6% (6.5-11%)	1.9 (1.7-2.2)	3.2% (1.9-4.6%)	1.6% (-0.91-4%)	15 (-8.6-38)	1000 (600-1500)
Iceland	2005	Male	15 to 19	89% (82-93%)	11% (7-18%)	1.4 (0.88-2.2)	4.2% (2.4-6%)	8.1% (3.1-13%)	0.43 (0.16-0.71)	49 (28-71)
Iceland	2005	Male	20 to 24	92% (87-95%)	8% (4.9-13%)	2.1 (1.3-3.1)	8% (5.1-11%)	11% (5-18%)	1 (0.46-1.7)	120 (73-160)
Iceland	2005	Male	25 to 29	94% (91-96%)	5.8% (3.6-9%)	2.2 (1.4-3.3)	6.9% (4.6-9.5%)	11% (5.3-18%)	0.88 (0.42-1.4)	99 (65-140)
Iceland	2005	Male	30 to 34	95% (92-97%)	4.6% (2.6-7.7%)	2.3 (1.5-3.4)	5.7% (3.7-7.8%)	11% (4.7-17%)	0.77 (0.34-1.2)	81 (52-110)
Iceland	2005	Male	35 to 39	96% (94-98%)	3.9% (2.2-6.2%)	2.3 (1.5-3.2)	5.1% (3.5-6.9%)	10% (5-15%)	0.79 (0.38-1.2)	73 (49-98)
Iceland	2005	Male	40 to 44	97% (94-98%)	3.4% (1.8-6%)	2.5 (1.7-3.4)	5.2% (3.4-7.2%)	10% (5.6-15%)	1.4 (0.75-2)	93 (62-130)
Iceland	2005	Male	45 to 49	97% (94-98%)	3.3% (1.7-6.2%)	2.9 (1.8-4)	5.3% (3.2-7.8%)	9.2% (4.7-14%)	2 (1-3.1)	110 (69-170)
Iceland	2005	Male	50 to 54	96% (94-98%)	3.7% (2-6.2%)	3.2 (2.1-4.2)	5.1% (2.9-7.4%)	8% (3.9-12%)	2.4 (1.2-3.6)	110 (64-170)
Iceland	2005	Male	55 to 59	95% (92-97%)	5% (2.8-8.2%)	3.3 (2.2-4.5)	4.5% (2.2-7.4%)	6.4% (2.5-11%)	2.7 (1.1-4.7)	110 (54-180)
Iceland	2005	Male	60 to 64	94% (90-97%)	6.2% (3.5-9.8%)	3.1 (1.9-4.5)	3.5% (0.84-6.9%)	4.4% (0.56-9.1%)	2.4 (0.31-5.1)	83 (19-160)
Iceland	2005	Male	65 to 69	93% (89-96%)	6.7% (3.5-11%)	2.8 (1.9-3.8)	2% (-0.19-4.6%)	2.3% (-0.81-5.7%)	1.5 (-0.52-3.6)	45 (-4.5-100)
Iceland	2005	Male	70 to 74	92% (88-95%)	7.8% (4.5-12%)	2.4 (1.5-3.5)	1.2% (-1-3.9%)	1.2% (-1.6-4.6%)	1.3 (-1.7-5)	33 (-26-110)
Iceland	2005	Male	75 to 79	91% (85-95%)	9.3% (5-15%)	2.1 (1.3-3.3)	0.52% (-1.6-3.2%)	0.33% (-2.4-3.5%)	0.48 (-3.5-5.2)	14 (-45-87)
Iceland	2005	Male	80 to 84	90% (82-95%)	10% (5.2-18%)	2.1 (1.3-3)	-0.14% (-2.4-2.3%)	-0.5% (-3.3-2.3%)	-0.85 (-5.5-4)	-3.1 (-54-52)
Iceland	2005	Male	85 to 89	89% (81-95%)	11% (5.4-19%)	2 (1.2-3)	-0.46% (-2.5-2%)	-0.89% (-3.4-2%)	-1.2 (-4.8-2.8)	-5.8 (-33-25)
Iceland	2005	Male	90 to 94	90% (83-95%)	10% (5.2-17%)	2 (1.1-3.1)	-0.8% (-3.2-2%)	-1.3% (-4.3-1.8%)	-1 (-3.3-1.4)	-3.8 (-16-9.2)
Iceland	2005	Male	95 plus	90% (82-96%)	9.6% (4.1-18%)	1.9 (0.99-3.4)	-0.66% (-3.3-2.9%)	-1.2% (-4.3-3%)	-0.29 (-1.1-0.72)	-0.65 (-3.3-2.9)
Iceland	2010	Female	All Ages	85% (82-88%)	15% (12-18%)	1.8 (1.5-2.2)	1.9% (1-3.1%)	1.7% (-0.64-4.6%)	16 (-6.3-44)	620 (320-970)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	2010	Female	15 to 19	85% (75-92%)	15% (8.3-25%)	1.5 (0.82-2.4)	1.5% (0.92-2.2%)	5.9% (2.3-9.6%)	0.12 (0.048-0.2)	17 (10-26)
Iceland	2010	Female	20 to 24	90% (83-95%)	9.7% (5.2-17%)	1.9 (1-3.1)	2.3% (1.5-3.2%)	6.3% (2.6-11%)	0.17 (0.074-0.3)	31 (20-46)
Iceland	2010	Female	25 to 29	92% (86-96%)	8.1% (4.3-14%)	2 (1.1-3.2)	1.8% (1.2-2.5%)	6.3% (3-10%)	0.14 (0.066-0.22)	27 (17-38)
Iceland	2010	Female	30 to 34	93% (88-96%)	7.1% (3.5-12%)	2 (1.2-3.3)	1.7% (1.1-2.4%)	7.2% (4.2-11%)	0.2 (0.11-0.31)	25 (17-36)
Iceland	2010	Female	35 to 39	93% (88-97%)	6.8% (3.5-12%)	2.2 (1.2-3.6)	2.2% (1.4-3.3%)	8.6% (5.3-13%)	0.45 (0.27-0.66)	36 (22-52)
Iceland	2010	Female	40 to 44	93% (88-96%)	7% (3.7-12%)	2.4 (1.4-3.8)	2.7% (1.6-3.9%)	9.4% (6-13%)	0.76 (0.48-1.1)	48 (29-69)
Iceland	2010	Female	45 to 49	92% (85-96%)	7.6% (3.8-15%)	2.8 (1.6-4.4)	3.2% (1.8-4.8%)	10% (6.6-14%)	1.3 (0.83-1.8)	65 (39-97)
Iceland	2010	Female	50 to 54	91% (84-94%)	9.5% (5.7-16%)	2.9 (1.8-4.5)	3.5% (1.9-5.2%)	9.3% (6.1-13%)	1.7 (1.1-2.3)	75 (44-110)
Iceland	2010	Female	55 to 59	88% (81-93%)	12% (7.1-19%)	3 (1.8-4.4)	3.7% (2.1-5.6%)	8.1% (5.3-11%)	2.3 (1.5-3.3)	85 (48-130)
Iceland	2010	Female	60 to 64	84% (76-91%)	16% (9.5-24%)	2.8 (1.5-4.5)	3.3% (1.4-5.9%)	6.2% (3.2-9.8%)	2.6 (1.3-4)	78 (33-130)
Iceland	2010	Female	65 to 69	79% (70-88%)	21% (12-30%)	2.4 (1.5-3.7)	2.5% (0.91-4.5%)	4.4% (2-7.2%)	2.3 (1-3.7)	56 (21-100)
Iceland	2010	Female	70 to 74	77% (66-85%)	23% (15-34%)	2.1 (1.2-3.4)	1.8% (-0.076-4.2%)	2.9% (0.29-6.1%)	2 (0.2-4.1)	38 (-1.7-87)
Iceland	2010	Female	75 to 79	75% (63-85%)	25% (15-37%)	1.9 (0.99-3.3)	1.5% (-1.1-3.9%)	1.5% (-1.3-5.3%)	1.7 (-1.5-6)	26 (-27-98)
Iceland	2010	Female	80 to 84	74% (60-85%)	26% (15-40%)	1.9 (1.1-3)	0.51% (-1.7-3.5%)	0.67% (-2.1-4.4%)	1.1 (-3.7-7.3)	13 (-43-88)
Iceland	2010	Female	85 to 89	74% (59-84%)	26% (16-41%)	1.8 (1-3.1)	0.076% (-2.3-3.2%)	-0.022% (-2.8-3.7%)	-0.043 (-5.7-7.4)	1.5 (-44-62)
Iceland	2010	Female	90 to 94	74% (62-85%)	26% (15-38%)	1.9 (0.94-3.3)	-0.2% (-2.9-3.7%)	-0.43% (-3.6-4.2%)	-0.62 (-5.3-6.2)	-1.8 (-27-36)
Iceland	2010	Female	95 plus	75% (59-88%)	25% (12-41%)	1.9 (0.74-3.6)	-0.12% (-3.2-5.1%)	-0.39% (-4.1-5.9%)	-0.29 (-3.1-4.4)	-0.38 (-10-16)
Iceland	2010	Male	All Ages	90% (87-93%)	9.8% (7.4-13%)	2.1 (1.7-2.5)	3.8% (2.4-5.3%)	2.4% (-0.08-5.1%)	24 (-0.8-51)	1300 (800-1800)
Iceland	2010	Male	15 to 19	86% (77-92%)	14% (7.8-23%)	1.5 (0.9-2.4)	3.9% (2.4-5.6%)	8.1% (3.1-13%)	0.38 (0.15-0.64)	47 (29-67)
Iceland	2010	Male	20 to 24	90% (83-94%)	10% (5.7-17%)	2.2 (1.3-3.4)	7.7% (5.2-11%)	11% (5.1-19%)	0.8 (0.36-1.3)	110 (69-150)
Iceland	2010	Male	25 to 29	93% (87-96%)	7.4% (4.2-13%)	2.4 (1.5-3.6)	6.9% (4.6-9.5%)	12% (5.2-19%)	0.9 (0.41-1.5)	110 (69-150)
Iceland	2010	Male	30 to 34	94% (89-97%)	5.7% (2.9-11%)	2.5 (1.5-3.7)	6.5% (4.1-9.2%)	12% (5.4-18%)	1.2 (0.55-1.9)	110 (67-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	2010	Male	35 to 39	95% (92-97%)	4.8% (2.5-8.4%)	2.5 (1.6-3.6)	5.6% (3.8-7.7%)	11% (5.6-17%)	1 (0.51-1.5)	88 (59-120)
Iceland	2010	Male	40 to 44	96% (92-98%)	4.2% (2-7.6%)	2.7 (1.7-3.8)	5.9% (3.9-8.1%)	11% (6.6-17%)	1.5 (0.89-2.2)	100 (68-140)
Iceland	2010	Male	45 to 49	96% (92-98%)	4% (1.9-7.6%)	3.1 (1.9-4.5)	6% (3.6-8.6%)	11% (5.9-16%)	2.2 (1.2-3.4)	130 (78-180)
Iceland	2010	Male	50 to 54	96% (92-98%)	4.4% (2.2-7.9%)	3.4 (2.3-4.7)	5.9% (3.7-8.6%)	9.8% (5.7-14%)	3.1 (1.8-4.6)	150 (91-210)
Iceland	2010	Male	55 to 59	94% (90-97%)	5.7% (3-9.6%)	3.6 (2.3-4.9)	5.4% (3.1-8.3%)	8.1% (4.2-13%)	3.4 (1.8-5.5)	140 (82-220)
Iceland	2010	Male	60 to 64	93% (89-97%)	6.6% (3.4-11%)	3.4 (2.1-5)	4.7% (2-8%)	6.4% (2.5-11%)	3.9 (1.5-7)	130 (57-230)
Iceland	2010	Male	65 to 69	93% (88-97%)	7.1% (3.4-12%)	3.1 (2-4.3)	3% (0.75-5.6%)	3.7% (0.56-7.4%)	2.8 (0.43-5.4)	79 (19-150)
Iceland	2010	Male	70 to 74	92% (86-96%)	8% (4.4-14%)	2.6 (1.6-3.9)	2.1% (-0.21-5%)	2.4% (-0.51-6.1%)	2.2 (-0.5-5.6)	51 (-5.2-120)
Iceland	2010	Male	75 to 79	91% (85-95%)	9.1% (4.8-15%)	2.4 (1.4-3.7)	1.3% (-1.1-4.2%)	1.3% (-1.7-4.8%)	1.8 (-2.5-7.1)	35 (-30-120)
Iceland	2010	Male	80 to 84	90% (82-95%)	10% (5.1-18%)	2.3 (1.4-3.4)	0.51% (-1.9-3.1%)	0.24% (-2.7-3.3%)	0.44 (-5-6.2)	13 (-47-79)
Iceland	2010	Male	85 to 89	89% (82-94%)	11% (5.6-18%)	2.2 (1.3-3.4)	0.096% (-2.2-2.8%)	-0.27% (-3.1-2.9%)	-0.45 (-5.1-4.9)	1.5 (-34-45)
Iceland	2010	Male	90 to 94	90% (83-95%)	10% (5-17%)	2.2 (1.3-3.6)	-0.32% (-2.9-3%)	-0.79% (-3.8-2.8%)	-0.75 (-3.6-2.7)	-1.9 (-17-17)
Iceland	2010	Male	95 plus	90% (81-96%)	9.7% (4-19%)	2.1 (1.1-3.7)	-0.33% (-3.2-3.9%)	-0.87% (-4.3-4%)	-0.22 (-1.1-1)	-0.33 (-3.2-4)
Iceland	2016	Female	All Ages	85% (81-88%)	15% (12-19%)	2 (1.4-2.6)	2.3% (1.2-3.9%)	2.4% (-0.24-6.3%)	26 (-2.7-68)	820 (400-1300)
Iceland	2016	Female	15 to 19	85% (75-92%)	15% (8.4-25%)	1.5 (0.79-2.7)	1.5% (0.9-2.3%)	5.8% (2.3-10%)	0.11 (0.043-0.19)	16 (9.5-25)
Iceland	2016	Female	20 to 24	90% (83-95%)	9.7% (5.1-17%)	2 (1-3.4)	2.4% (1.6-3.4%)	6.4% (2.8-11%)	0.19 (0.083-0.33)	34 (22-51)
Iceland	2016	Female	25 to 29	92% (86-96%)	8.1% (4.4-14%)	2.1 (1.1-3.6)	2% (1.3-2.8%)	6.7% (3.3-11%)	0.15 (0.07-0.24)	27 (17-39)
Iceland	2016	Female	30 to 34	93% (87-96%)	7.1% (3.6-13%)	2.2 (1.1-3.6)	1.8% (1.2-2.6%)	7.9% (4.6-12%)	0.24 (0.14-0.36)	29 (19-42)
Iceland	2016	Female	35 to 39	93% (88-96%)	6.8% (3.5-12%)	2.3 (1.2-4.1)	2.3% (1.4-3.5%)	9.3% (5.6-13%)	0.47 (0.28-0.68)	37 (23-55)
Iceland	2016	Female	40 to 44	93% (88-96%)	7% (3.7-12%)	2.6 (1.4-4.4)	3% (1.8-4.5%)	10% (6.8-14%)	0.9 (0.58-1.3)	55 (35-82)
Iceland	2016	Female	45 to 49	92% (86-96%)	7.6% (3.8-14%)	3 (1.6-5)	3.4% (1.8-5.3%)	11% (7.2-15%)	1.3 (0.82-1.8)	68 (39-100)
Iceland	2016	Female	50 to 54	90% (84-94%)	9.5% (5.7-16%)	3.1 (1.7-5)	3.9% (2.1-5.9%)	10% (6.8-14%)	2.1 (1.4-3)	93 (54-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	2016	Female	55 to 59	88% (81-93%)	12% (7.1-19%)	3.1 (1.8-5)	4.1% (2.3-6.4%)	8.9% (5.9-13%)	3 (2-4.4)	110 (64-170)
Iceland	2016	Female	60 to 64	84% (75-91%)	16% (9.4-25%)	2.9 (1.6-5)	3.8% (1.8-6.7%)	7.1% (3.9-11%)	3.6 (2-5.6)	110 (52-180)
Iceland	2016	Female	65 to 69	79% (69-88%)	21% (12-31%)	2.6 (1.5-4.1)	3% (1.2-5.5%)	5.2% (2.6-8.4%)	3.5 (1.8-5.7)	86 (37-160)
Iceland	2016	Female	70 to 74	77% (66-85%)	23% (15-34%)	2.3 (1.2-3.8)	2.3% (0.27-5.1%)	3.7% (0.91-7.1%)	2.9 (0.71-5.8)	58 (6.7-130)
Iceland	2016	Female	75 to 79	75% (64-85%)	25% (15-36%)	2.1 (0.98-3.6)	1.6% (-0.67-5%)	2.3% (-0.72-6.6%)	2.4 (-0.76-7)	37 (-16-110)
Iceland	2016	Female	80 to 84	74% (60-86%)	26% (14-40%)	2 (1-3.4)	1.2% (-1.2-4.8%)	1.5% (-1.5-5.9%)	2.7 (-2.8-11)	31 (-34-130)
Iceland	2016	Female	85 to 89	74% (60-84%)	26% (16-40%)	2 (1-3.4)	0.73% (-1.9-4.8%)	0.79% (-2.4-5.7%)	1.8 (-5.6-13)	17 (-43-110)
Iceland	2016	Female	90 to 94	74% (61-85%)	26% (15-39%)	2 (0.94-3.7)	0.47% (-2.5-5.4%)	0.37% (-3.2-6.1%)	0.74 (-6.6-12)	6.1 (-34-66)
Iceland	2016	Female	95 plus	75% (59-88%)	25% (12-41%)	2 (0.76-4.1)	0.44% (-3-6.8%)	0.27% (-3.9-8%)	0.23 (-3.3-6.7)	1.5 (-10-24)
Iceland	2016	Male	All Ages	90% (88-93%)	9.7% (7.3-12%)	2.3 (1.6-3.1)	4.4% (2.7-6.5%)	3.3% (0.46-7%)	38 (5.6-80)	1600 (970-2300)
Iceland	2016	Male	15 to 19	86% (77-92%)	14% (8.1-23%)	1.6 (0.87-2.6)	3.9% (2.4-5.6%)	8.5% (3.4-14%)	0.33 (0.13-0.56)	42 (26-61)
Iceland	2016	Male	20 to 24	90% (83-94%)	10% (5.6-17%)	2.4 (1.3-3.8)	7.8% (5.3-11%)	12% (5.2-20%)	0.83 (0.35-1.4)	110 (74-160)
Iceland	2016	Male	25 to 29	93% (87-96%)	7.4% (4.2-13%)	2.6 (1.5-4.2)	7.5% (5-11%)	12% (5.5-20%)	0.9 (0.4-1.5)	110 (70-150)
Iceland	2016	Male	30 to 34	94% (89-97%)	5.7% (2.8-11%)	2.7 (1.5-4.3)	7% (4.5-9.9%)	13% (6.1-20%)	1.4 (0.66-2.2)	120 (76-180)
Iceland	2016	Male	35 to 39	95% (92-97%)	4.8% (2.5-8.5%)	2.7 (1.5-4.2)	6.1% (4.1-8.5%)	13% (6.4-19%)	1.1 (0.53-1.7)	90 (61-130)
Iceland	2016	Male	40 to 44	96% (92-98%)	4.2% (2.1-7.7%)	2.9 (1.7-4.5)	6.5% (4.3-9.2%)	13% (7.7-19%)	1.8 (1-2.6)	120 (76-160)
Iceland	2016	Male	45 to 49	96% (92-98%)	3.9% (1.9-7.9%)	3.3 (1.9-5.2)	6.9% (4.4-10%)	13% (7.4-20%)	2.4 (1.3-3.6)	130 (82-190)
Iceland	2016	Male	50 to 54	96% (92-98%)	4.5% (2.3-8%)	3.7 (2.2-5.4)	6.9% (4.2-10%)	12% (6.9-17%)	3.7 (2.1-5.5)	170 (100-250)
Iceland	2016	Male	55 to 59	94% (90-97%)	5.7% (2.9-9.7%)	3.8 (2.4-5.7)	6.4% (3.6-9.9%)	9.9% (5.5-15%)	4.9 (2.7-7.8)	190 (110-300)
Iceland	2016	Male	60 to 64	93% (89-97%)	6.6% (3.4-11%)	3.6 (2.1-5.7)	5.6% (2.6-9.6%)	8% (3.7-14%)	5.4 (2.5-9.4)	180 (87-300)
Iceland	2016	Male	65 to 69	93% (87-97%)	7.2% (3.4-13%)	3.3 (2-5)	4% (1.5-7.4%)	5.2% (1.8-9.8%)	5.3 (1.8-10)	150 (55-270)
Iceland	2016	Male	70 to 74	92% (87-95%)	8% (4.5-13%)	2.8 (1.6-4.4)	2.9% (0.48-6.3%)	3.6% (0.44-7.7%)	3.9 (0.5-8.4)	85 (14-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iceland	2016	Male	75 to 79	91% (85-95%)	9% (4.6-15%)	2.6 (1.3-4.2)	1.9% (-0.67-5.8%)	2.1% (-1.1-6.6%)	2.8 (-1.5-9.2)	51 (-17-150)
Iceland	2016	Male	80 to 84	90% (82-95%)	10% (5-18%)	2.4 (1.3-3.9)	1.2% (-1.3-4.7%)	1.1% (-2-5.2%)	2.2 (-4.1-11)	34 (-37-130)
Iceland	2016	Male	85 to 89	90% (82-94%)	10% (5.8-18%)	2.3 (1.3-3.9)	0.74% (-1.8-4.5%)	0.51% (-2.6-5%)	1 (-5.4-10)	14 (-35-88)
Iceland	2016	Male	90 to 94	90% (83-95%)	10% (5-17%)	2.3 (1.2-4)	0.37% (-2.5-5%)	0.034% (-3.3-5.3%)	0.04 (-4.4-6.9)	2.9 (-20-39)
Iceland	2016	Male	95 plus	90% (82-96%)	9.5% (3.9-18%)	2.3 (1.1-4.3)	0.2% (-3.1-5.6%)	-0.25% (-4.2-6.3%)	-0.1 (-1.7-2.5)	0.31 (-5-9.2)
Ireland	1990	Female	All Ages	66% (61-71%)	34% (29-39%)	2.3 (2.1-2.6)	3.7% (2.2-5.2%)	4.1% (1.3-7.3%)	610 (190-1100)	17000 (10000-24000)
Ireland	1990	Female	15 to 19	69% (55-81%)	31% (19-45%)	2.6 (1.8-3.6)	3.3% (2.2-4.6%)	10% (5.8-16%)	4.8 (2.7-7.3)	590 (390-820)
Ireland	1990	Female	20 to 24	79% (67-87%)	21% (13-33%)	3.6 (2.5-4.8)	5.2% (3.8-6.8%)	14% (8.6-19%)	6.1 (3.9-8.5)	870 (610-1200)
Ireland	1990	Female	25 to 29	78% (67-87%)	22% (13-33%)	3.1 (2.2-4.1)	4.3% (3-5.5%)	12% (7.1-17%)	5.8 (3.4-8.2)	750 (530-1000)
Ireland	1990	Female	30 to 34	78% (67-85%)	22% (15-33%)	2.9 (2.1-4.1)	3.7% (2.6-5%)	11% (7.1-15%)	7.2 (4.6-10)	690 (480-940)
Ireland	1990	Female	35 to 39	77% (66-84%)	23% (16-34%)	3 (2.1-4.3)	4.2% (3-5.6%)	12% (8.8-16%)	12 (8.2-15)	850 (600-1100)
Ireland	1990	Female	40 to 44	76% (67-84%)	24% (16-33%)	3.3 (2.4-4.4)	5.2% (3.7-6.8%)	13% (10-17%)	19 (14-25)	1100 (840-1400)
Ireland	1990	Female	45 to 49	77% (66-84%)	23% (16-34%)	3.8 (2.7-5.1)	6.5% (4.8-8.8%)	14% (10-18%)	29 (21-37)	1400 (1000-1800)
Ireland	1990	Female	50 to 54	76% (67-83%)	24% (17-33%)	4 (2.9-5.2)	7% (4.8-9.6%)	12% (8.8-17%)	37 (26-51)	1500 (1100-2100)
Ireland	1990	Female	55 to 59	75% (66-82%)	25% (18-34%)	3.9 (2.9-5.2)	6.7% (4.2-9.6%)	10% (6.6-15%)	49 (31-68)	1700 (1100-2400)
Ireland	1990	Female	60 to 64	72% (62-80%)	28% (20-38%)	3.6 (2.5-4.9)	5.5% (2.5-9.2%)	7.7% (3.4-13%)	60 (27-99)	1800 (830-2900)
Ireland	1990	Female	65 to 69	68% (58-76%)	32% (24-42%)	3 (2.2-4)	3.9% (1.4-6.8%)	5.1% (1.5-8.9%)	66 (20-120)	1600 (570-2800)
Ireland	1990	Female	70 to 74	63% (53-72%)	37% (28-47%)	2.5 (1.7-3.4)	2.7% (-0.23-5.9%)	3.3% (-0.41-7.3%)	61 (-7.3-140)	1200 (-100-2600)
Ireland	1990	Female	75 to 79	59% (45-71%)	41% (29-55%)	2.2 (1.4-3.2)	2.1% (-0.99-5.5%)	2.4% (-1.4-6.5%)	62 (-34-170)	970 (-440-2600)
Ireland	1990	Female	80 to 84	55% (41-67%)	45% (33-59%)	2.2 (1.5-3.1)	2.4% (-0.34-5.6%)	2.7% (-0.57-6.6%)	73 (-15-180)	840 (-120-2000)
Ireland	1990	Female	85 to 89	52% (38-64%)	48% (36-62%)	2.2 (1.4-3.3)	2.7% (-0.13-6.3%)	3% (-0.36-7.1%)	70 (-8.2-160)	560 (-26-1300)
Ireland	1990	Female	90 to 94	50% (37-62%)	50% (38-63%)	2.3 (1.4-3.6)	2.9% (-0.33-7.4%)	3.2% (-0.58-8.4%)	35 (-6.6-94)	200 (-23-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	1990	Female	95 plus	51% (34-66%)	49% (34-66%)	2.5 (1.3-4.1)	3.4% (-0.7-8.9%)	3.8% (-0.93-10%)	15 (-3.7-41)	55 (-11-140)
Ireland	1990	Male	All Ages	68% (61-74%)	32% (26-39%)	3.3 (2.9-3.7)	7.9% (5.8-10%)	8% (4.4-12%)	1300 (740-2000)	41000 (30000-54000)
Ireland	1990	Male	15 to 19	65% (49-79%)	35% (21-51%)	2.8 (1.8-3.9)	7.5% (5.1-11%)	14% (7.9-21%)	17 (9.5-25)	1600 (1100-2300)
Ireland	1990	Male	20 to 24	76% (63-86%)	24% (14-37%)	4.7 (3.2-6.2)	15% (11-20%)	23% (15-31%)	34 (22-47)	3200 (2300-4200)
Ireland	1990	Male	25 to 29	79% (69-87%)	21% (13-31%)	4.5 (3.2-5.8)	14% (10-17%)	22% (14-29%)	29 (19-39)	2600 (1900-3400)
Ireland	1990	Male	30 to 34	81% (70-89%)	19% (11-30%)	4.2 (2.9-5.6)	11% (7.6-14%)	19% (12-25%)	25 (16-34)	2000 (1500-2700)
Ireland	1990	Male	35 to 39	83% (73-90%)	17% (10-27%)	3.9 (2.7-5.3)	8.7% (6.1-12%)	15% (9.4-21%)	24 (15-34)	1700 (1200-2300)
Ireland	1990	Male	40 to 44	84% (74-91%)	16% (9.3-26%)	4.4 (3.2-5.7)	8.4% (5.3-12%)	13% (7.6-20%)	31 (17-45)	1900 (1200-2600)
Ireland	1990	Male	45 to 49	84% (74-91%)	16% (9.4-26%)	5.7 (4.1-7.5)	11% (6.9-15%)	16% (9.4-22%)	53 (31-73)	2600 (1700-3500)
Ireland	1990	Male	50 to 54	83% (75-89%)	17% (11-25%)	6.4 (4.7-8.1)	12% (7.8-16%)	16% (10-22%)	80 (50-110)	3300 (2100-4300)
Ireland	1990	Male	55 to 59	82% (73-88%)	18% (12-27%)	6.4 (4.9-8.3)	12% (8.1-16%)	15% (9.7-21%)	120 (79-170)	4300 (2800-5800)
Ireland	1990	Male	60 to 64	79% (69-85%)	21% (15-31%)	5.9 (4.3-8)	11% (5.8-15%)	13% (6.6-18%)	170 (86-240)	4900 (2600-6900)
Ireland	1990	Male	65 to 69	76% (66-83%)	24% (17-34%)	5 (3.7-6.4)	8.6% (4.4-13%)	9.7% (4.6-15%)	200 (95-310)	4900 (2400-7400)
Ireland	1990	Male	70 to 74	74% (64-81%)	26% (19-36%)	4.1 (3.1-5.4)	6.2% (1.8-11%)	6.8% (1.7-12%)	180 (45-340)	3600 (1000-6300)
Ireland	1990	Male	75 to 79	73% (62-82%)	27% (18-38%)	3.7 (2.5-5.1)	4.8% (0.67-10%)	5.1% (0.46-11%)	160 (15-350)	2400 (340-5100)
Ireland	1990	Male	80 to 84	72% (58-82%)	28% (18-42%)	3.5 (2.4-4.8)	4.6% (0.95-9.1%)	4.9% (0.72-9.9%)	120 (18-250)	1400 (290-2800)
Ireland	1990	Male	85 to 89	71% (57-81%)	29% (19-43%)	3.4 (2.2-4.6)	4.4% (0.59-8.7%)	4.6% (0.32-9.5%)	65 (4.5-140)	530 (74-1100)
Ireland	1990	Male	90 to 94	73% (60-83%)	27% (17-40%)	3.4 (2.1-4.9)	4% (0.089-8.9%)	4.3% (-0.2-9.9%)	23 (-1.1-53)	130 (2.8-280)
Ireland	1990	Male	95 plus	75% (60-87%)	25% (13-40%)	3.5 (2-5.3)	4.2% (-0.48-10%)	4.4% (-0.93-11%)	5.1 (-1.1-13)	19 (-2.2-47)
Ireland	1995	Female	All Ages	68% (63-73%)	32% (27-37%)	2.8 (2.5-3.2)	4.7% (3.2-6.4%)	5.8% (2.6-9.4%)	870 (380-1400)	21000 (14000-29000)
Ireland	1995	Female	15 to 19	71% (57-82%)	29% (18-43%)	3.1 (2.1-4.1)	3.5% (2.4-4.8%)	11% (6.8-17%)	5.3 (3.1-7.8)	640 (430-870)
Ireland	1995	Female	20 to 24	80% (69-87%)	20% (13-31%)	4.1 (2.9-5.5)	5.6% (4.1-7.2%)	15% (10-21%)	7.4 (4.7-10)	1000 (730-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	1995	Female	25 to 29	79% (69-87%)	21% (13-31%)	3.6 (2.6-4.8)	4.7% (3.4-6%)	13% (8.6-18%)	6.7 (4.2-9.2)	820 (580-1100)
Ireland	1995	Female	30 to 34	78% (68-86%)	22% (14-32%)	3.4 (2.4-4.8)	4% (2.9-5.3%)	12% (8.4-17%)	8.4 (5.9-12)	800 (580-1100)
Ireland	1995	Female	35 to 39	77% (68-85%)	23% (15-32%)	3.5 (2.4-4.8)	4.6% (3.4-6.1%)	14% (10-18%)	14 (9.9-18)	1000 (740-1300)
Ireland	1995	Female	40 to 44	77% (68-84%)	23% (16-32%)	3.8 (2.8-5.1)	5.7% (4.3-7.3%)	15% (12-18%)	22 (17-28)	1300 (1000-1700)
Ireland	1995	Female	45 to 49	78% (68-85%)	22% (15-32%)	4.4 (3.1-5.9)	7.2% (5.4-9.5%)	16% (12-20%)	37 (28-48)	1800 (1400-2300)
Ireland	1995	Female	50 to 54	77% (69-84%)	23% (16-31%)	4.6 (3.3-6.2)	8% (5.8-11%)	15% (11-19%)	46 (35-61)	1900 (1500-2600)
Ireland	1995	Female	55 to 59	76% (67-83%)	24% (17-33%)	4.6 (3.3-6)	7.9% (5.4-11%)	13% (8.9-18%)	55 (38-75)	2000 (1400-2700)
Ireland	1995	Female	60 to 64	73% (64-80%)	27% (20-36%)	4.2 (3-5.7)	6.8% (3.8-10%)	9.9% (5.4-15%)	66 (36-99)	2000 (1100-3000)
Ireland	1995	Female	65 to 69	69% (60-77%)	31% (23-40%)	3.6 (2.6-4.6)	5.3% (2.6-8.4%)	7% (3.2-11%)	78 (36-130)	2000 (940-3100)
Ireland	1995	Female	70 to 74	65% (55-73%)	35% (27-45%)	3 (2-4.1)	3.9% (0.95-7.3%)	4.9% (1-9.3%)	88 (18-170)	1800 (440-3300)
Ireland	1995	Female	75 to 79	60% (47-72%)	40% (28-53%)	2.7 (1.7-3.8)	3.4% (0.15-7.1%)	3.9% (-0.019-8.7%)	95 (-0.47-210)	1500 (66-3200)
Ireland	1995	Female	80 to 84	56% (43-68%)	44% (32-57%)	2.6 (1.7-3.7)	3.7% (0.61-7.4%)	4.3% (0.53-8.7%)	130 (17-270)	1500 (230-3000)
Ireland	1995	Female	85 to 89	53% (41-65%)	47% (35-59%)	2.7 (1.7-3.9)	4.1% (0.69-8.4%)	4.6% (0.54-9.7%)	120 (13-240)	930 (160-1900)
Ireland	1995	Female	90 to 94	52% (39-64%)	48% (36-61%)	2.8 (1.7-4.2)	4.3% (0.44-9.6%)	4.8% (0.22-11%)	64 (3-150)	360 (36-760)
Ireland	1995	Female	95 plus	52% (35-69%)	48% (31-65%)	3 (1.5-4.8)	5% (0.063-11%)	5.7% (-0.13-14%)	26 (-0.62-63)	95 (1.1-220)
Ireland	1995	Male	All Ages	71% (64-76%)	29% (24-36%)	4 (3.5-4.5)	9.7% (7.6-12%)	10% (6.4-14%)	1700 (1000-2400)	50000 (38000-63000)
Ireland	1995	Male	15 to 19	67% (50-80%)	33% (20-50%)	3.3 (2.2-4.5)	8.3% (5.5-12%)	16% (9.2-23%)	19 (11-29)	1800 (1200-2600)
Ireland	1995	Male	20 to 24	78% (65-87%)	22% (13-35%)	5.4 (3.8-7.1)	17% (12-22%)	25% (17-34%)	47 (31-65)	4200 (3000-5400)
Ireland	1995	Male	25 to 29	80% (70-88%)	20% (12-30%)	5.1 (3.6-6.8)	15% (12-19%)	24% (17-32%)	36 (25-48)	3100 (2300-4000)
Ireland	1995	Male	30 to 34	82% (71-89%)	18% (11-29%)	4.8 (3.4-6.5)	12% (8.9-15%)	20% (14-27%)	31 (21-42)	2500 (1800-3200)
Ireland	1995	Male	35 to 39	84% (75-90%)	16% (9.7-25%)	4.4 (3.1-6.1)	10% (7.3-13%)	18% (12-24%)	30 (20-41)	2200 (1500-2800)
Ireland	1995	Male	40 to 44	85% (75-91%)	15% (8.7-25%)	5 (3.7-6.5)	10% (7-13%)	17% (11-23%)	40 (26-54)	2400 (1700-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	1995	Male	45 to 49	85% (76-91%)	15% (9-24%)	6.6 (4.7-8.6)	13% (9.1-16%)	20% (14-25%)	72 (49-93)	3600 (2600-4500)
Ireland	1995	Male	50 to 54	84% (76-90%)	16% (10-24%)	7.3 (5.4-9.4)	14% (10-17%)	19% (14-25%)	97 (69-130)	4100 (3000-5100)
Ireland	1995	Male	55 to 59	82% (74-88%)	18% (12-26%)	7.4 (5.6-9.5)	14% (9.9-18%)	17% (12-23%)	130 (95-170)	4700 (3400-6000)
Ireland	1995	Male	60 to 64	80% (72-86%)	20% (14-28%)	6.8 (4.9-9.1)	12% (8.2-17%)	15% (9.6-20%)	180 (110-240)	5200 (3400-7100)
Ireland	1995	Male	65 to 69	77% (68-84%)	23% (16-32%)	5.8 (4.3-7.4)	10% (6.3-15%)	12% (7-17%)	220 (130-310)	5300 (3200-7500)
Ireland	1995	Male	70 to 74	75% (66-82%)	25% (18-34%)	4.8 (3.5-6.4)	8.1% (3.6-13%)	9% (3.8-14%)	230 (99-370)	4500 (2000-7100)
Ireland	1995	Male	75 to 79	74% (63-83%)	26% (17-37%)	4.3 (2.8-5.9)	6.7% (1.8-12%)	7.2% (1.7-13%)	210 (51-380)	3100 (870-5500)
Ireland	1995	Male	80 to 84	73% (61-83%)	27% (17-39%)	4.1 (2.8-5.5)	6.5% (2.1-11%)	6.9% (2.1-12%)	190 (57-340)	2200 (750-3700)
Ireland	1995	Male	85 to 89	72% (60-82%)	28% (18-40%)	3.9 (2.6-5.4)	6.1% (1.8-11%)	6.5% (1.6-12%)	100 (27-190)	830 (240-1500)
Ireland	1995	Male	90 to 94	74% (62-84%)	26% (16-38%)	4 (2.6-5.7)	5.8% (1.1-11%)	6.2% (0.96-12%)	37 (5.8-73)	200 (39-390)
Ireland	1995	Male	95 plus	76% (61-87%)	24% (13-39%)	4 (2.2-6.2)	5.9% (0.39-12%)	6.3% (-0.0027-14%)	9.4 (-0.0042-20)	35 (2.3-72)
Ireland	2000	Female	All Ages	70% (65-74%)	30% (26-35%)	3.4 (3-3.9)	5.6% (4.1-7.5%)	7.4% (4.2-12%)	1100 (620-1800)	26000 (19000-35000)
Ireland	2000	Female	15 to 19	70% (57-81%)	30% (19-43%)	3.4 (2.3-4.7)	4% (2.8-5.5%)	13% (7.6-18%)	6.3 (3.8-9)	710 (480-970)
Ireland	2000	Female	20 to 24	82% (73-88%)	18% (12-27%)	4.9 (3.4-6.7)	6.3% (4.7-8%)	17% (12-23%)	9.8 (6.5-13)	1200 (910-1600)
Ireland	2000	Female	25 to 29	82% (75-88%)	18% (12-25%)	4.3 (3.1-5.7)	5% (3.8-6.3%)	15% (11-20%)	7.9 (5.4-11)	980 (710-1300)
Ireland	2000	Female	30 to 34	80% (74-86%)	20% (14-26%)	4 (3-5.5)	4.5% (3.4-5.9%)	15% (11-19%)	10 (7.5-13)	950 (700-1200)
Ireland	2000	Female	35 to 39	79% (73-85%)	21% (15-27%)	4.1 (2.9-5.6)	5.5% (4.1-7.1%)	16% (12-20%)	19 (14-24)	1300 (1000-1700)
Ireland	2000	Female	40 to 44	78% (71-85%)	22% (15-29%)	4.4 (3.2-5.8)	6.4% (4.8-8.2%)	17% (13-21%)	27 (21-34)	1600 (1300-2100)
Ireland	2000	Female	45 to 49	79% (72-85%)	21% (15-28%)	5.1 (3.7-6.7)	7.9% (5.9-10%)	17% (14-23%)	42 (33-54)	2100 (1600-2800)
Ireland	2000	Female	50 to 54	78% (72-84%)	22% (16-28%)	5.3 (3.9-7)	9% (6.8-12%)	17% (13-22%)	62 (48-83)	2600 (2000-3500)
Ireland	2000	Female	55 to 59	75% (69-81%)	25% (19-31%)	5.1 (3.8-6.7)	8.7% (6.2-12%)	15% (10-19%)	65 (47-88)	2400 (1700-3200)
Ireland	2000	Female	60 to 64	70% (63-77%)	30% (23-37%)	4.6 (3.2-6.2)	7.8% (5-11%)	12% (7.5-17%)	74 (47-110)	2300 (1400-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	2000	Female	65 to 69	65% (58-73%)	35% (27-42%)	3.8 (2.8-5)	6.3% (3.7-9.1%)	8.6% (5-13%)	82 (47-120)	2100 (1200-3100)
Ireland	2000	Female	70 to 74	62% (52-71%)	38% (29-48%)	3.2 (2.2-4.4)	5% (2.1-8.4%)	6.4% (2.5-11%)	99 (39-170)	2000 (820-3400)
Ireland	2000	Female	75 to 79	61% (50-71%)	39% (29-50%)	3.1 (2-4.4)	4.7% (1.3-9.1%)	5.6% (1.4-11%)	130 (34-260)	2100 (570-4100)
Ireland	2000	Female	80 to 84	57% (44-68%)	43% (32-56%)	3 (2-4.3)	5.1% (1.7-9.2%)	5.9% (1.8-11%)	170 (53-320)	1900 (650-3500)
Ireland	2000	Female	85 to 89	53% (40-64%)	47% (36-60%)	3 (1.9-4.4)	5.4% (1.9-10%)	6.2% (2-12%)	170 (56-340)	1400 (500-2600)
Ireland	2000	Female	90 to 94	51% (39-64%)	49% (36-61%)	3.1 (1.8-4.9)	5.7% (1.5-11%)	6.5% (1.5-13%)	97 (22-200)	530 (130-1100)
Ireland	2000	Female	95 plus	51% (34-68%)	49% (32-66%)	3.3 (1.7-5.5)	6.4% (1.2-13%)	7.4% (1-15%)	41 (5.7-85)	140 (26-300)
Ireland	2000	Male	All Ages	74% (69-79%)	26% (21-31%)	4.9 (4.3-5.5)	12% (9.4-14%)	13% (9-16%)	2000 (1500-2700)	60000 (50000-73000)
Ireland	2000	Male	15 to 19	69% (55-81%)	31% (19-45%)	3.9 (2.6-5.4)	9.8% (6.5-13%)	18% (11-26%)	24 (14-35)	2200 (1400-2900)
Ireland	2000	Male	20 to 24	81% (71-88%)	19% (12-29%)	6.5 (4.7-8.7)	29% (14-24%)	29% (20-37%)	57 (39-75)	4900 (3700-6300)
Ireland	2000	Male	25 to 29	84% (77-89%)	16% (11-23%)	6.2 (4.6-8)	17% (13-21%)	27% (20-35%)	49 (35-64)	4100 (3100-5100)
Ireland	2000	Male	30 to 34	85% (78-90%)	15% (9.8-22%)	5.7 (4.2-7.6)	14% (11-18%)	25% (18-32%)	42 (30-54)	3200 (2500-4000)
Ireland	2000	Male	35 to 39	86% (79-91%)	14% (8.9-21%)	5.3 (3.8-7.2)	12% (9.2-16%)	22% (16-29%)	43 (31-55)	2900 (2200-3700)
Ireland	2000	Male	40 to 44	86% (80-92%)	14% (8.5-20%)	6 (4.3-7.8)	13% (9.6-16%)	23% (16-28%)	60 (42-76)	3400 (2600-4300)
Ireland	2000	Male	45 to 49	87% (80-92%)	13% (8.5-20%)	7.7 (5.5-10)	15% (12-19%)	24% (18-29%)	91 (69-110)	4400 (3500-5400)
Ireland	2000	Male	50 to 54	86% (80-90%)	14% (9.5-20%)	8.6 (6.4-11)	15% (12-19%)	22% (18-28%)	130 (100-170)	5500 (4300-6800)
Ireland	2000	Male	55 to 59	83% (78-88%)	17% (12-22%)	8.6 (6.4-11)	15% (11-19%)	20% (15-25%)	160 (120-200)	5500 (4200-7100)
Ireland	2000	Male	60 to 64	79% (72-85%)	21% (15-28%)	7.8 (5.7-10)	13% (9.9-18%)	17% (12-22%)	180 (130-250)	5500 (4000-7300)
Ireland	2000	Male	65 to 69	76% (68-83%)	24% (17-32%)	6.5 (5-8.3)	12% (8.4-16%)	14% (9.8-18%)	230 (160-300)	5600 (4000-7400)
Ireland	2000	Male	70 to 74	74% (66-81%)	26% (19-34%)	5.4 (3.9-7.4)	9.8% (5.4-14%)	11% (5.9-16%)	250 (130-360)	4800 (2800-6900)
Ireland	2000	Male	75 to 79	74% (64-82%)	26% (18-36%)	4.9 (3.3-6.8)	8.4% (3.6-13%)	9.2% (3.8-14%)	260 (100-410)	3900 (1700-6000)
Ireland	2000	Male	80 to 84	75% (63-84%)	25% (16-37%)	4.8 (3.3-6.5)	8.4% (4-13%)	9.2% (4.1-14%)	240 (110-370)	2600 (1300-4100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	2000	Male	85 to 89	75% (64-83%)	25% (17-36%)	4.7 (3.3-6.5)	8.1% (3.6-13%)	8.8% (3.7-14%)	160 (67-260)	1300 (550-2000)
Ireland	2000	Male	90 to 94	76% (66-85%)	24% (15-34%)	4.7 (3.1-6.7)	7.7% (2.6-13%)	8.4% (2.5-15%)	58 (17-99)	310 (100-530)
Ireland	2000	Male	95 plus	78% (63-88%)	22% (12-37%)	4.7 (2.6-7.1)	7.7% (1.3-14%)	8.4% (0.99-16%)	12 (1.4-23)	45 (7.4-82)
Ireland	2005	Female	All Ages	70% (66-74%)	30% (26-34%)	3.6 (3-4.1)	5.6% (4.3-7.2%)	7.7% (4.7-11%)	1100 (650-1600)	26000 (20000-33000)
Ireland	2005	Female	15 to 19	69% (57-80%)	31% (20-43%)	3.5 (2.4-4.8)	3.8% (2.6-5.3%)	13% (7.9-18%)	5.1 (3.2-7.3)	600 (410-800)
Ireland	2005	Female	20 to 24	83% (74-88%)	17% (12-26%)	5.1 (3.5-6.9)	5.7% (4.3-7.4%)	17% (12-23%)	8.5 (5.8-11)	1200 (880-1600)
Ireland	2005	Female	25 to 29	82% (75-88%)	18% (12-25%)	4.5 (3.2-5.9)	4.8% (3.6-6.1%)	15% (11-20%)	8.7 (6.1-11)	1100 (810-1500)
Ireland	2005	Female	30 to 34	80% (74-86%)	20% (14-26%)	4.2 (3-5.6)	4.2% (3.2-5.5%)	15% (11-19%)	10 (7.5-14)	1000 (760-1300)
Ireland	2005	Female	35 to 39	79% (73-85%)	21% (15-27%)	4.3 (3-5.9)	5% (3.7-6.5%)	16% (13-20%)	17 (13-21)	1300 (960-1600)
Ireland	2005	Female	40 to 44	78% (72-84%)	22% (16-28%)	4.6 (3.4-6.2)	6.1% (4.7-7.8%)	17% (14-21%)	27 (22-33)	1700 (1300-2100)
Ireland	2005	Female	45 to 49	79% (73-84%)	21% (16-27%)	5.3 (3.9-7.1)	7.7% (5.9-9.9%)	18% (14-23%)	43 (34-55)	2200 (1700-2800)
Ireland	2005	Female	50 to 54	77% (71-83%)	23% (17-29%)	5.4 (4-7.2)	8.5% (6.3-11%)	17% (13-22%)	57 (44-75)	2500 (1900-3300)
Ireland	2005	Female	55 to 59	75% (69-80%)	25% (20-31%)	5.3 (3.9-6.9)	8.4% (6.1-11%)	15% (11-19%)	71 (53-95)	2700 (2000-3500)
Ireland	2005	Female	60 to 64	72% (67-78%)	28% (22-33%)	4.9 (3.5-6.6)	7.8% (5.3-11%)	12% (8.4-17%)	75 (52-100)	2400 (1600-3300)
Ireland	2005	Female	65 to 69	66% (59-73%)	34% (27-41%)	4 (3-5.2)	6.3% (4-8.7%)	9.1% (6-13%)	74 (48-100)	1900 (1200-2800)
Ireland	2005	Female	70 to 74	61% (54-67%)	39% (33-46%)	3.2 (2.2-4.5)	5% (2.5-8.1%)	6.7% (3.3-11%)	82 (41-130)	1700 (850-2700)
Ireland	2005	Female	75 to 79	58% (48-67%)	42% (33-52%)	3 (2-4.4)	4.6% (1.7-8.2%)	5.7% (2.1-11%)	110 (40-200)	1700 (640-3100)
Ireland	2005	Female	80 to 84	53% (40-66%)	47% (34-60%)	2.9 (2-4.1)	4.9% (1.9-8.6%)	5.8% (2.2-10%)	160 (59-290)	1800 (740-3200)
Ireland	2005	Female	85 to 89	50% (38-62%)	50% (38-62%)	3 (1.9-4.3)	5.3% (2.2-9.6%)	6.2% (2.3-11%)	160 (59-290)	1300 (520-2300)
Ireland	2005	Female	90 to 94	49% (37-60%)	51% (40-63%)	3.1 (1.8-4.8)	5.6% (1.8-11%)	6.5% (1.9-13%)	110 (32-220)	620 (200-1200)
Ireland	2005	Female	95 plus	50% (34-66%)	50% (34-66%)	3.3 (1.8-5.5)	6.6% (1.4-13%)	7.6% (1.4-15%)	50 (9.2-100)	180 (38-340)
Ireland	2005	Male	All Ages	75% (70-80%)	25% (20-30%)	5.2 (4.5-6)	11% (9.6-13%)	13% (9.8-16%)	1900 (1400-2400)	58000 (49000-69000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	2005	Male	15 to 19	68% (55-80%)	32% (20-45%)	4 (2.6-5.6)	9.2% (6.3-13%)	18% (11-26%)	18 (11-26)	1700 (1200-2300)
Ireland	2005	Male	20 to 24	82% (72-88%)	18% (12-28%)	6.8 (4.8-9.2)	18% (14-23%)	29% (20-37%)	50 (34-65)	4600 (3400-5700)
Ireland	2005	Male	25 to 29	84% (77-90%)	16% (10-23%)	6.4 (4.7-8.2)	15% (12-19%)	26% (19-34%)	45 (32-59)	4100 (3200-5200)
Ireland	2005	Male	30 to 34	85% (79-90%)	15% (9.7-21%)	5.9 (4.3-7.9)	13% (10-16%)	24% (17-30%)	41 (30-53)	3400 (2600-4200)
Ireland	2005	Male	35 to 39	86% (80-91%)	14% (9-20%)	5.5 (3.9-7.5)	12% (9.1-15%)	23% (17-29%)	43 (31-56)	3100 (2300-3800)
Ireland	2005	Male	40 to 44	87% (80-91%)	13% (8.6-20%)	6.2 (4.5-8.1)	13% (10-16%)	24% (18-29%)	61 (45-76)	3600 (2700-4400)
Ireland	2005	Male	45 to 49	87% (81-92%)	13% (8.3-19%)	8.1 (5.9-11)	15% (12-18%)	25% (20-30%)	89 (70-110)	4500 (3600-5300)
Ireland	2005	Male	50 to 54	86% (81-91%)	14% (9.4-19%)	9.1 (6.7-12)	15% (12-19%)	24% (19-28%)	130 (100-150)	5300 (4300-6400)
Ireland	2005	Male	55 to 59	84% (79-88%)	16% (12-21%)	8.9 (6.8-12)	15% (11-18%)	20% (16-25%)	160 (120-200)	5800 (4600-7100)
Ireland	2005	Male	60 to 64	80% (74-85%)	20% (15-26%)	8.1 (6-11)	13% (10-16%)	17% (13-21%)	170 (130-220)	5200 (4000-6600)
Ireland	2005	Male	65 to 69	77% (70-83%)	23% (17-30%)	6.8 (5.2-8.7)	12% (8.9-15%)	15% (11-18%)	190 (140-250)	4900 (3700-6200)
Ireland	2005	Male	70 to 74	74% (66-80%)	26% (20-34%)	5.6 (4-7.5)	9.6% (6-13%)	11% (6.8-15%)	210 (120-290)	4200 (2500-5700)
Ireland	2005	Male	75 to 79	74% (66-81%)	26% (19-34%)	5 (3.4-7)	8.4% (4.1-12%)	9.5% (4.4-14%)	210 (100-320)	3300 (1600-5000)
Ireland	2005	Male	80 to 84	75% (63-84%)	25% (16-37%)	5 (3.5-6.7)	8.5% (4.5-13%)	9.4% (4.7-14%)	230 (110-340)	2600 (1400-3800)
Ireland	2005	Male	85 to 89	74% (62-83%)	26% (17-38%)	4.8 (3.3-6.7)	8.2% (4-13%)	9% (4.1-14%)	150 (71-240)	1200 (620-1900)
Ireland	2005	Male	90 to 94	76% (64-85%)	24% (15-36%)	4.9 (3.2-7)	7.9% (3.2-13%)	8.7% (3.1-14%)	68 (24-110)	370 (150-600)
Ireland	2005	Male	95 plus	77% (63-88%)	23% (12-37%)	4.9 (2.8-7.5)	8% (1.9-14%)	8.8% (1.7-16%)	16 (3-28)	57 (14-100)
Ireland	2010	Female	All Ages	69% (65-73%)	31% (27-35%)	3.3 (2.7-4)	5% (3.8-6.5%)	6.9% (4.1-10%)	970 (570-1500)	25000 (19000-32000)
Ireland	2010	Female	15 to 19	71% (59-82%)	29% (18-41%)	3.3 (2.2-4.8)	3.6% (2.4-5%)	12% (7.3-18%)	4.3 (2.6-6.3)	540 (370-740)
Ireland	2010	Female	20 to 24	82% (74-88%)	18% (12-26%)	4.8 (3.3-6.7)	5.4% (4-6.8%)	17% (11-23%)	6.1 (3.9-8.5)	1000 (740-1400)
Ireland	2010	Female	25 to 29	82% (75-88%)	18% (12-25%)	4.1 (3-5.6)	4.6% (3.5-5.9%)	14% (9.2-19%)	8.4 (5.4-11)	1200 (850-1500)
Ireland	2010	Female	30 to 34	80% (74-86%)	20% (14-26%)	3.9 (2.7-5.3)	3.9% (2.9-5.1%)	13% (9.7-17%)	10 (7.2-13)	1100 (790-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	2010	Female	35 to 39	79% (72-85%)	21% (15-28%)	4 (2.8-5.7)	4.5% (3.4-5.8%)	15% (12-19%)	16 (12-20)	1300 (980-1700)
Ireland	2010	Female	40 to 44	78% (71-84%)	22% (16-29%)	4.3 (3-5.8)	5.5% (4.2-7.2%)	17% (13-20%)	25 (20-31)	1600 (1300-2000)
Ireland	2010	Female	45 to 49	79% (73-84%)	21% (16-27%)	4.9 (3.5-6.6)	7.2% (5.5-9.2%)	18% (14-22%)	44 (35-55)	2300 (1800-2900)
Ireland	2010	Female	50 to 54	76% (70-81%)	24% (19-30%)	5 (3.6-6.7)	7.4% (5.5-9.7%)	16% (13-20%)	52 (40-67)	2300 (1800-3100)
Ireland	2010	Female	55 to 59	74% (68-79%)	26% (21-32%)	4.9 (3.6-6.5)	7.5% (5.5-9.9%)	14% (11-18%)	64 (50-84)	2500 (1800-3300)
Ireland	2010	Female	60 to 64	73% (67-79%)	27% (21-33%)	4.6 (3.3-6.3)	7.1% (5-9.8%)	12% (8.5-16%)	77 (56-100)	2500 (1700-3400)
Ireland	2010	Female	65 to 69	67% (61-74%)	33% (26-39%)	3.8 (2.8-5.1)	5.7% (3.8-8%)	8.7% (6-12%)	70 (48-95)	1900 (1200-2600)
Ireland	2010	Female	70 to 74	61% (53-68%)	39% (32-47%)	3 (2.1-4.2)	4.5% (2.3-7%)	6.4% (3.5-9.9%)	71 (39-110)	1500 (790-2400)
Ireland	2010	Female	75 to 79	56% (45-67%)	44% (33-55%)	2.6 (1.7-3.9)	3.9% (1.6-7%)	5% (2.1-9.2%)	85 (35-160)	1400 (550-2600)
Ireland	2010	Female	80 to 84	53% (40-65%)	47% (35-60%)	2.7 (1.7-3.8)	4.1% (1.7-7.2%)	5% (1.9-9%)	120 (46-220)	1400 (580-2500)
Ireland	2010	Female	85 to 89	51% (38-61%)	49% (39-62%)	2.8 (1.7-4.1)	4.5% (1.7-8.1%)	5.3% (1.8-9.8%)	150 (51-280)	1200 (450-2200)
Ireland	2010	Female	90 to 94	49% (36-62%)	51% (38-64%)	2.9 (1.6-4.5)	4.7% (1.3-9.4%)	5.5% (1.3-11%)	100 (25-210)	580 (160-1100)
Ireland	2010	Female	95 plus	50% (33-66%)	50% (34-67%)	3.1 (1.6-5.2)	5.7% (0.82-12%)	6.6% (0.7-14%)	61 (6.5-130)	220 (31-440)
Ireland	2010	Male	All Ages	74% (69-79%)	26% (21-31%)	4.9 (4-5.8)	11% (8.8-13%)	12% (9.1-16%)	1800 (1300-2300)	56000 (46000-67000)
Ireland	2010	Male	15 to 19	68% (54-80%)	32% (20-46%)	3.7 (2.4-5.3)	8.2% (5.6-11%)	17% (10-26%)	13 (7.8-20)	1400 (920-1900)
Ireland	2010	Male	20 to 24	81% (71-88%)	19% (12-29%)	6.4 (4.5-8.9)	17% (13-21%)	27% (18-35%)	37 (25-49)	3700 (2800-4700)
Ireland	2010	Male	25 to 29	84% (77-89%)	16% (11-23%)	6 (4.4-8)	14% (11-18%)	24% (17-32%)	39 (27-51)	3800 (2900-4800)
Ireland	2010	Male	30 to 34	86% (79-90%)	14% (9.7-21%)	5.6 (4-7.6)	12% (9.4-15%)	22% (16-28%)	40 (29-53)	3500 (2600-4300)
Ireland	2010	Male	35 to 39	86% (80-91%)	14% (9-20%)	5.2 (3.7-7.3)	11% (8.7-14%)	22% (16-28%)	46 (33-60)	3300 (2500-4200)
Ireland	2010	Male	40 to 44	86% (79-91%)	14% (8.7-21%)	5.8 (4.3-7.9)	12% (9.5-15%)	23% (18-29%)	61 (46-76)	3700 (2800-4500)
Ireland	2010	Male	45 to 49	87% (81-92%)	13% (8.4-19%)	7.6 (5.4-10)	14% (11-17%)	25% (20-30%)	92 (71-110)	4600 (3700-5600)
Ireland	2010	Male	50 to 54	86% (81-91%)	14% (9.3-19%)	8.5 (6.2-11)	15% (12-18%)	24% (19-28%)	120 (97-140)	5200 (4200-6300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	2010	Male	55 to 59	84% (79-88%)	16% (12-21%)	8.4 (6.3-11)	14% (11-17%)	21% (17-25%)	150 (120-180)	5600 (4600-6800)
Ireland	2010	Male	60 to 64	81% (75-86%)	19% (14-25%)	7.8 (5.7-11)	13% (9.8-16%)	17% (13-21%)	180 (140-230)	5700 (4500-7000)
Ireland	2010	Male	65 to 69	77% (70-83%)	23% (17-30%)	6.5 (4.9-8.5)	11% (8.3-14%)	14% (11-18%)	190 (140-240)	4800 (3600-6100)
Ireland	2010	Male	70 to 74	74% (66-80%)	26% (20-34%)	5.2 (3.8-7.1)	8.8% (5.6-12%)	11% (6.5-15%)	180 (110-250)	3700 (2300-5000)
Ireland	2010	Male	75 to 79	74% (66-82%)	26% (18-34%)	4.8 (3.2-6.7)	7.6% (3.8-11%)	8.7% (4.2-13%)	190 (89-280)	3000 (1500-4500)
Ireland	2010	Male	80 to 84	74% (62-83%)	26% (17-38%)	4.6 (3.2-6.3)	7.5% (3.9-11%)	8.4% (4.1-13%)	200 (96-300)	2300 (1200-3500)
Ireland	2010	Male	85 to 89	72% (60-82%)	28% (18-40%)	4.4 (2.9-6.1)	7% (3.3-11%)	7.8% (3.4-12%)	150 (67-250)	1300 (580-2000)
Ireland	2010	Male	90 to 94	74% (63-84%)	26% (16-37%)	4.4 (2.8-6.5)	6.7% (2.3-11%)	7.4% (2.2-13%)	67 (20-120)	370 (130-620)
Ireland	2010	Male	95 plus	76% (61-87%)	24% (13-39%)	4.4 (2.5-7)	6.9% (1.5-13%)	7.6% (1.3-14%)	18 (3.2-34)	67 (15-120)
Ireland	2016	Female	All Ages	68% (64-73%)	32% (27-36%)	3.1 (2.3-4.2)	4.7% (3.2-6.7%)	6.7% (3.3-11%)	990 (480-1600)	24000 (16000-34000)
Ireland	2016	Female	15 to 19	71% (57-82%)	29% (18-43%)	3.2 (1.9-4.9)	3.3% (2.2-4.7%)	12% (6.4-18%)	3.6 (1.9-5.6)	480 (320-710)
Ireland	2016	Female	20 to 24	80% (68-87%)	20% (13-32%)	4.4 (2.7-6.8)	5.1% (3.8-6.7%)	16% (9.5-23%)	3.4 (2-5.1)	640 (440-880)
Ireland	2016	Female	25 to 29	79% (69-87%)	21% (13-31%)	3.8 (2.4-5.6)	4.3% (3.2-5.7%)	13% (8.2-19%)	5 (2.9-7.4)	800 (550-1100)
Ireland	2016	Female	30 to 34	78% (67-86%)	22% (14-33%)	3.6 (2.3-5.4)	3.7% (2.7-4.8%)	13% (8.8-17%)	8.5 (5.5-12)	1000 (690-1400)
Ireland	2016	Female	35 to 39	77% (68-85%)	23% (15-32%)	3.7 (2.4-5.6)	4.1% (3-5.5%)	15% (11-19%)	14 (10-20)	1200 (910-1700)
Ireland	2016	Female	40 to 44	77% (68-85%)	23% (15-32%)	4.1 (2.6-5.9)	5.1% (3.6-6.7%)	16% (12-20%)	24 (17-33)	1600 (1200-2200)
Ireland	2016	Female	45 to 49	78% (68-85%)	22% (15-32%)	4.6 (3-6.7)	6.5% (4.6-8.7%)	17% (13-21%)	39 (28-54)	2100 (1500-2900)
Ireland	2016	Female	50 to 54	77% (69-84%)	23% (16-31%)	4.9 (3.1-7)	6.9% (4.8-9.3%)	15% (11-20%)	52 (37-71)	2400 (1700-3300)
Ireland	2016	Female	55 to 59	76% (67-83%)	24% (17-33%)	4.8 (3.2-7)	7% (4.9-9.8%)	13% (9.8-18%)	63 (43-91)	2400 (1700-3500)
Ireland	2016	Female	60 to 64	73% (64-80%)	27% (20-36%)	4.4 (2.9-6.5)	6.5% (4.1-9.7%)	11% (7.2-16%)	73 (46-110)	2400 (1500-3600)
Ireland	2016	Female	65 to 69	69% (60-77%)	31% (23-40%)	3.8 (2.5-5.4)	5.3% (3.1-8.1%)	8.3% (5.2-12%)	79 (49-120)	2200 (1200-3300)
Ireland	2016	Female	70 to 74	65% (55-72%)	35% (28-45%)	3.1 (1.9-4.7)	4.3% (2-7.4%)	6.3% (3.2-10%)	76 (37-130)	1600 (740-2800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	2016	Female	75 to 79	60% (47-72%)	40% (28-53%)	2.8 (1.6-4.4)	3.8% (1.2-7.6%)	5% (1.8-10%)	87 (30-170)	1400 (460-2800)
Ireland	2016	Female	80 to 84	56% (42-69%)	44% (31-58%)	2.8 (1.7-4.3)	4% (1.3-7.5%)	4.9% (1.5-9.4%)	120 (36-230)	1400 (470-2700)
Ireland	2016	Female	85 to 89	53% (40-65%)	47% (35-60%)	2.8 (1.6-4.4)	4.3% (1.1-8.3%)	5.1% (1.1-10%)	140 (32-280)	1200 (280-2300)
Ireland	2016	Female	90 to 94	52% (38-64%)	48% (36-62%)	3 (1.6-4.8)	4.6% (0.79-9.4%)	5.3% (0.7-11%)	130 (17-280)	720 (120-1500)
Ireland	2016	Female	95 plus	52% (35-68%)	48% (32-65%)	3.1 (1.5-5.4)	5.4% (0.48-11%)	6.3% (0.35-14%)	75 (4-160)	260 (22-550)
Ireland	2016	Male	All Ages	71% (66-77%)	29% (23-34%)	4.5 (3.3-6.1)	9.9% (7.3-12%)	11% (6.9-15%)	1800 (1100-2600)	53000 (39000-70000)
Ireland	2016	Male	15 to 19	66% (49-80%)	34% (20-51%)	3.5 (2-5.7)	7.4% (4.9-11%)	16% (9.1-26%)	11 (5.8-18)	1200 (790-1700)
Ireland	2016	Male	20 to 24	77% (63-87%)	23% (13-37%)	5.9 (3.7-8.9)	16% (12-20%)	25% (16-35%)	23 (14-34)	2400 (1700-3200)
Ireland	2016	Male	25 to 29	80% (69-87%)	20% (13-31%)	5.5 (3.6-8.1)	14% (10-17%)	23% (15-31%)	24 (15-34)	2500 (1800-3300)
Ireland	2016	Male	30 to 34	82% (71-89%)	18% (11-29%)	5.2 (3.3-7.4)	11% (8.3-14%)	21% (14-29%)	32 (20-45)	2800 (2000-3700)
Ireland	2016	Male	35 to 39	83% (74-90%)	17% (9.7-26%)	4.8 (3.1-7.2)	11% (7.8-14%)	21% (14-28%)	39 (25-56)	3000 (2100-4000)
Ireland	2016	Male	40 to 44	84% (75-91%)	16% (9.1-25%)	5.5 (3.6-7.7)	12% (8.4-15%)	23% (16-29%)	59 (40-82)	3600 (2700-4800)
Ireland	2016	Male	45 to 49	85% (75-91%)	15% (9.2-25%)	7.1 (4.6-10)	13% (10-17%)	24% (18-30%)	88 (62-120)	4500 (3400-5900)
Ireland	2016	Male	50 to 54	84% (75-90%)	16% (10-25%)	8 (5.3-11)	14% (11-17%)	23% (17-28%)	120 (87-150)	5200 (4000-6700)
Ireland	2016	Male	55 to 59	82% (74-88%)	18% (12-26%)	8 (5.4-11)	13% (10-16%)	20% (16-25%)	150 (110-190)	5500 (4000-7000)
Ireland	2016	Male	60 to 64	80% (71-86%)	20% (14-29%)	7.4 (4.9-11)	12% (8.7-15%)	17% (12-21%)	170 (120-230)	5500 (3900-7100)
Ireland	2016	Male	65 to 69	76% (67-84%)	24% (16-33%)	6.2 (4.1-8.7)	10% (7.1-14%)	14% (9.1-18%)	210 (140-290)	5600 (3700-7400)
Ireland	2016	Male	70 to 74	75% (65-82%)	25% (18-35%)	5.2 (3.2-7.6)	8.4% (4.5-12%)	10% (5.4-15%)	190 (100-280)	4100 (2100-5900)
Ireland	2016	Male	75 to 79	74% (63-83%)	26% (17-37%)	4.6 (2.8-7)	7.1% (3-11%)	8.2% (3.4-13%)	190 (77-320)	3100 (1300-5000)
Ireland	2016	Male	80 to 84	73% (60-83%)	27% (17-40%)	4.4 (2.7-6.4)	6.8% (3-11%)	7.7% (3.1-13%)	210 (86-350)	2500 (1100-4100)
Ireland	2016	Male	85 to 89	72% (59-82%)	28% (18-41%)	4.2 (2.6-6.4)	6.5% (2.3-11%)	7.2% (2.2-12%)	160 (47-280)	1300 (440-2200)
Ireland	2016	Male	90 to 94	74% (62-83%)	26% (17-38%)	4.3 (2.5-6.6)	6.2% (1.6-11%)	6.8% (1.4-13%)	94 (19-180)	530 (130-950)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ireland	2016	Male	95 plus	76% (60-87%)	24% (13-40%)	4.3 (2.3-7.3)	6.4% (0.77-12%)	7.1% (0.48-14%)	27 (1.8-54)	97 (12-190)
Israel	1990	Female	All Ages	39% (34-44%)	61% (56-66%)	0.45 (0.38-0.52)	0.15% (-0.38-0.65%)	-0.63% (-1.7-0.4%)	-92 (-260-59)	800 (-1900-3400)
Israel	1990	Female	15 to 19	41% (29-54%)	59% (46-71%)	0.46 (0.27-0.71)	0.86% (0.46-1.4%)	2.5% (0.84-4.6%)	1.5 (0.49-2.6)	190 (100-300)
Israel	1990	Female	20 to 24	46% (32-61%)	54% (39-68%)	0.62 (0.35-0.97)	1.2% (0.73-1.8%)	2.8% (1.1-5%)	1.8 (0.69-3.2)	260 (160-390)
Israel	1990	Female	25 to 29	48% (33-64%)	52% (36-67%)	0.62 (0.37-0.95)	1.1% (0.66-1.5%)	2.7% (1.2-4.4%)	1.7 (0.74-2.8)	220 (140-320)
Israel	1990	Female	30 to 34	49% (33-66%)	51% (34-67%)	0.64 (0.38-0.95)	1.1% (0.68-1.5%)	3.1% (1.7-4.7%)	2.6 (1.4-4.1)	240 (150-350)
Israel	1990	Female	35 to 39	50% (35-66%)	50% (34-65%)	0.69 (0.42-1)	1.2% (0.73-1.7%)	3.6% (2.2-5.2%)	4.5 (2.7-6.8)	310 (200-440)
Israel	1990	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.77 (0.48-1.1)	1.4% (0.85-2.1%)	3.8% (2.3-5.6%)	7 (4.3-11)	380 (240-550)
Israel	1990	Female	45 to 49	47% (34-58%)	53% (42-66%)	0.85 (0.56-1.2)	1.5% (0.84-2.3%)	3.7% (2.1-5.3%)	7.1 (4.1-11)	320 (180-470)
Israel	1990	Female	50 to 54	45% (34-56%)	55% (44-66%)	0.83 (0.54-1.2)	1.3% (0.57-2.2%)	2.8% (1.3-4.3%)	8.4 (3.8-14)	310 (130-510)
Israel	1990	Female	55 to 59	44% (33-53%)	56% (47-67%)	0.72 (0.46-1)	0.69% (-0.094-1.6%)	1.4% (0.057-2.8%)	6.9 (0.29-14)	200 (-28-440)
Israel	1990	Female	60 to 64	43% (32-53%)	57% (47-68%)	0.66 (0.42-0.96)	-0.048% (-1-0.87%)	0.099% (-1.3-1.4%)	0.83 (-12-13)	-18 (-370-320)
Israel	1990	Female	65 to 69	42% (32-52%)	58% (48-68%)	0.64 (0.42-0.94)	-0.44% (-1.5-0.56%)	-0.48% (-1.9-0.88%)	-6.3 (-26-12)	-190 (-630-230)
Israel	1990	Female	70 to 74	40% (31-50%)	60% (50-69%)	0.61 (0.38-0.89)	-0.74% (-2-0.3%)	-0.89% (-2.4-0.47%)	-15 (-41-8)	-300 (-790-120)
Israel	1990	Female	75 to 79	37% (27-49%)	63% (51-73%)	0.55 (0.33-0.82)	-1% (-2.2--0.025%)	-1.3% (-2.8-0.0021%)	-33 (-74-0.056)	-480 (-1100--10)
Israel	1990	Female	80 to 84	35% (24-45%)	65% (55-76%)	0.49 (0.3-0.71)	-1.1% (-2.2--0.21%)	-1.3% (-2.7--0.22%)	-36 (-73--5.9)	-380 (-770--72)
Israel	1990	Female	85 to 89	33% (23-43%)	67% (57-77%)	0.44 (0.26-0.69)	-1.1% (-2.2--0.28%)	-1.3% (-2.6--0.33%)	-27 (-53--6.6)	-200 (-390--48)
Israel	1990	Female	90 to 94	33% (23-43%)	67% (57-77%)	0.42 (0.23-0.69)	-1.1% (-2.1--0.3%)	-1.4% (-2.6--0.37%)	-13 (-25--3.5)	-66 (-130--17)
Israel	1990	Female	95 plus	34% (23-47%)	66% (53-77%)	0.42 (0.21-0.71)	-1.2% (-2.5--0.32%)	-1.5% (-3.1--0.41%)	-4.6 (-9.3--1.3)	-15 (-31--3.8)
Israel	1990	Male	All Ages	44% (38-50%)	56% (50-62%)	0.54 (0.47-0.62)	0.87% (0.043-1.7%)	-0.17% (-1.5-1.2%)	-27 (-240-200)	4800 (230-9400)
Israel	1990	Male	15 to 19	41% (30-54%)	59% (46-70%)	0.37 (0.21-0.57)	1.9% (0.99-3.1%)	3.2% (1.1-6.1%)	4.8 (1.7-9.3)	510 (260-840)
Israel	1990	Male	20 to 24	48% (33-63%)	52% (37-67%)	0.59 (0.34-0.89)	3.6% (2.1-5.4%)	4.6% (1.7-8.2%)	8 (2.9-14)	880 (500-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	1990	Male	25 to 29	52% (35-68%)	48% (32-65%)	0.67 (0.41-1)	3.4% (2.1-4.8%)	4.8% (2-7.8%)	7.1 (3-12)	770 (470-1100)
Israel	1990	Male	30 to 34	56% (38-74%)	44% (26-62%)	0.71 (0.42-1.1)	3.2% (2.1-4.3%)	4.9% (2.5-7.5%)	7.9 (4-12)	730 (470-1000)
Israel	1990	Male	35 to 39	59% (42-76%)	41% (24-58%)	0.73 (0.44-1.1)	2.8% (1.8-3.8%)	4.6% (2.3-6.8%)	8.9 (4.7-13)	720 (470-970)
Israel	1990	Male	40 to 44	61% (44-77%)	39% (23-56%)	0.87 (0.55-1.2)	2.6% (1.6-3.7%)	4.1% (1.8-6.2%)	10 (4.7-16)	670 (400-940)
Israel	1990	Male	45 to 49	61% (43-76%)	39% (24-57%)	1.2 (0.71-1.7)	2.4% (1.2-3.7%)	3.4% (1.3-5.5%)	9.7 (3.5-16)	540 (260-820)
Israel	1990	Male	50 to 54	61% (46-75%)	39% (25-54%)	1.3 (0.82-1.9)	2% (0.44-3.4%)	2.5% (-0.018-4.7%)	11 (-0.068-21)	520 (110-890)
Israel	1990	Male	55 to 59	60% (49-70%)	40% (30-51%)	1.3 (0.88-1.8)	1.1% (-0.46-2.6%)	1.2% (-0.99-3.3%)	8.7 (-7.6-25)	380 (-160-900)
Israel	1990	Male	60 to 64	60% (48-70%)	40% (30-52%)	1.2 (0.83-1.7)	0.53% (-1.1-2.2%)	0.46% (-1.7-2.6%)	5 (-19-29)	210 (-440-860)
Israel	1990	Male	65 to 69	61% (49-70%)	39% (30-51%)	1.1 (0.78-1.6)	-0.11% (-1.6-1.5%)	-0.27% (-2.2-1.7%)	-4.4 (-37-30)	-50 (-800-730)
Israel	1990	Male	70 to 74	60% (49-68%)	40% (32-51%)	0.97 (0.64-1.4)	-0.49% (-1.9-0.89%)	-0.69% (-2.5-1%)	-12 (-43-18)	-200 (-760-370)
Israel	1990	Male	75 to 79	57% (44-69%)	43% (31-56%)	0.84 (0.53-1.2)	-0.85% (-2.3-0.43%)	-1.1% (-2.8-0.45%)	-29 (-74-11)	-380 (-1000-180)
Israel	1990	Male	80 to 84	53% (41-65%)	47% (35-59%)	0.73 (0.46-1)	-0.91% (-2.2-0.26%)	-1.1% (-2.7-0.27%)	-30 (-71-7.2)	-290 (-730-84)
Israel	1990	Male	85 to 89	50% (38-62%)	50% (38-62%)	0.64 (0.38-0.96)	-0.87% (-2.1-0.17%)	-1.1% (-2.5-0.13%)	-19 (-45-2.4)	-140 (-320-26)
Israel	1990	Male	90 to 94	50% (38-62%)	50% (38-62%)	0.6 (0.35-0.94)	-1% (-2.3-0.021%)	-1.3% (-2.8-0.038%)	-10 (-22-0.3)	-49 (-110-0.99)
Israel	1990	Male	95 plus	52% (37-66%)	48% (34-63%)	0.58 (0.3-0.98)	-1.1% (-2.5--0.0044%)	-1.5% (-3.1-0.098%)	-3.7 (-7.8-0.25)	-11 (-25-0.043)
Israel	1995	Female	All Ages	37% (33-43%)	63% (57-67%)	0.42 (0.36-0.49)	0.22% (-0.23-0.64%)	-0.44% (-1.4-0.41%)	-78 (-240-69)	1300 (-1400-3800)
Israel	1995	Female	15 to 19	39% (28-52%)	61% (48-72%)	0.42 (0.24-0.65)	0.81% (0.43-1.3%)	2.4% (0.7-4.4%)	1.4 (0.42-2.7)	200 (100-310)
Israel	1995	Female	20 to 24	45% (31-60%)	55% (40-69%)	0.57 (0.33-0.9)	1.2% (0.7-1.7%)	2.8% (0.97-5%)	1.9 (0.63-3.5)	310 (180-460)
Israel	1995	Female	25 to 29	47% (31-63%)	53% (37-69%)	0.57 (0.33-0.88)	1% (0.65-1.4%)	2.6% (1.1-4.2%)	1.6 (0.68-2.8)	250 (150-360)
Israel	1995	Female	30 to 34	48% (31-65%)	52% (35-69%)	0.59 (0.35-0.88)	0.98% (0.64-1.4%)	2.9% (1.6-4.4%)	2.5 (1.4-3.9)	250 (160-360)
Israel	1995	Female	35 to 39	49% (33-65%)	51% (35-67%)	0.64 (0.38-0.96)	1.1% (0.66-1.5%)	3.4% (2-4.9%)	4.1 (2.4-6.1)	300 (190-420)
Israel	1995	Female	40 to 44	48% (34-64%)	52% (36-66%)	0.7 (0.43-1)	1.2% (0.74-1.8%)	3.6% (2.3-5.5%)	7.5 (4.5-11)	410 (250-610)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	1995	Female	45 to 49	45% (31-57%)	55% (43-69%)	0.76 (0.49-1.1)	1.4% (0.79-2%)	3.5% (2.2-5.2%)	10 (5.9-15)	450 (270-670)
Israel	1995	Female	50 to 54	43% (32-55%)	57% (45-68%)	0.74 (0.48-1.1)	1.2% (0.6-2%)	2.8% (1.5-4.3%)	8.4 (4.5-13)	310 (150-500)
Israel	1995	Female	55 to 59	42% (31-52%)	58% (48-69%)	0.64 (0.41-0.92)	0.76% (0.091-1.5%)	1.6% (0.51-2.9%)	8.4 (2.6-15)	240 (28-460)
Israel	1995	Female	60 to 64	41% (30-52%)	59% (48-70%)	0.59 (0.36-0.87)	0.11% (-0.66-0.88%)	0.44% (-0.78-1.6%)	3.8 (-6.9-14)	43 (-270-340)
Israel	1995	Female	65 to 69	40% (29-50%)	60% (50-71%)	0.57 (0.37-0.83)	-0.24% (-1.1-0.57%)	-0.13% (-1.3-1%)	-1.9 (-20-14)	-120 (-530-270)
Israel	1995	Female	70 to 74	38% (29-48%)	62% (52-71%)	0.54 (0.34-0.79)	-0.47% (-1.4-0.37%)	-0.49% (-1.7-0.63%)	-11 (-38-14)	-260 (-780-200)
Israel	1995	Female	75 to 79	36% (25-47%)	64% (53-75%)	0.48 (0.29-0.72)	-0.69% (-1.7-0.1%)	-0.82% (-2.1-0.2%)	-20 (-51-5.2)	-330 (-780-47)
Israel	1995	Female	80 to 84	33% (23-43%)	67% (57-77%)	0.43 (0.26-0.64)	-0.9% (-1.9--0.11%)	-1.1% (-2.3--0.12%)	-35 (-76--3.8)	-380 (-800--47)
Israel	1995	Female	85 to 89	31% (21-41%)	69% (59-79%)	0.39 (0.22-0.61)	-0.94% (-1.9--0.21%)	-1.1% (-2.3--0.25%)	-33 (-67--7.2)	-250 (-500--56)
Israel	1995	Female	90 to 94	31% (22-42%)	69% (58-78%)	0.37 (0.2-0.63)	-0.99% (-1.9--0.24%)	-1.2% (-2.3--0.29%)	-19 (-37--4.4)	-95 (-190--23)
Israel	1995	Female	95 plus	33% (21-45%)	67% (55-79%)	0.37 (0.18-0.64)	-1.1% (-2.2--0.24%)	-1.4% (-2.7--0.33%)	-7.6 (-15--1.9)	-25 (-50--5.7)
Israel	1995	Male	All Ages	43% (37-49%)	57% (51-63%)	0.52 (0.45-0.6)	1% (0.31-1.8%)	0.14% (-1-1.3%)	26 (-200-250)	6500 (1900-11000)
Israel	1995	Male	15 to 19	39% (28-52%)	61% (48-72%)	0.34 (0.19-0.53)	1.7% (0.87-2.9%)	2.9% (0.89-5.7%)	4.6 (1.4-9.3)	510 (250-850)
Israel	1995	Male	20 to 24	46% (31-62%)	54% (38-69%)	0.54 (0.31-0.82)	3.4% (1.9-5.4%)	4.4% (1.5-8.2%)	11 (3.4-20)	1100 (600-1800)
Israel	1995	Male	25 to 29	50% (33-67%)	50% (33-67%)	0.61 (0.35-0.93)	3.3% (1.9-4.8%)	4.6% (1.7-8%)	8.1 (2.9-14)	860 (500-1300)
Israel	1995	Male	30 to 34	54% (35-72%)	46% (28-65%)	0.65 (0.38-0.98)	3% (1.9-4.3%)	4.7% (2.2-7.4%)	8.6 (4-14)	800 (490-1100)
Israel	1995	Male	35 to 39	58% (40-74%)	42% (26-60%)	0.67 (0.41-1)	2.8% (1.8-4%)	4.5% (2.4-6.7%)	10 (5.3-15)	790 (510-1100)
Israel	1995	Male	40 to 44	60% (42-76%)	40% (24-58%)	0.81 (0.51-1.2)	2.5% (1.5-3.5%)	3.8% (1.8-5.7%)	12 (5.7-19)	820 (500-1200)
Israel	1995	Male	45 to 49	60% (41-76%)	40% (24-59%)	1.1 (0.64-1.6)	2.3% (1.2-3.4%)	3.3% (1.3-5.2%)	15 (6.2-24)	820 (410-1200)
Israel	1995	Male	50 to 54	60% (43-74%)	40% (26-57%)	1.2 (0.76-1.8)	1.9% (0.72-3.2%)	2.6% (0.61-4.5%)	13 (2.9-22)	580 (230-950)
Israel	1995	Male	55 to 59	59% (46-69%)	41% (31-54%)	1.2 (0.8-1.6)	1.3% (0.071-2.5%)	1.6% (-0.2-3.4%)	13 (-1.5-27)	490 (27-970)
Israel	1995	Male	60 to 64	58% (46-69%)	42% (31-54%)	1.2 (0.76-1.6)	0.85% (-0.5-2.1%)	0.95% (-0.89-2.7%)	12 (-11-33)	380 (-220-980)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	1995	Male	65 to 69	59% (48-68%)	41% (32-52%)	1 (0.71-1.4)	0.23% (-1.1-1.6%)	0.19% (-1.5-1.9%)	3.4 (-26-33)	120 (-560-810)
Israel	1995	Male	70 to 74	58% (47-67%)	42% (33-53%)	0.9 (0.6-1.3)	-0.22% (-1.4-1%)	-0.32% (-1.8-1.2%)	-7.9 (-44-29)	-120 (-790-580)
Israel	1995	Male	75 to 79	55% (43-67%)	45% (33-57%)	0.77 (0.48-1.1)	-0.49% (-1.7-0.6%)	-0.65% (-2.1-0.67%)	-16 (-52-17)	-210 (-730-270)
Israel	1995	Male	80 to 84	51% (38-63%)	49% (37-62%)	0.66 (0.41-0.95)	-0.64% (-1.8-0.36%)	-0.81% (-2.1-0.39%)	-23 (-62-11)	-230 (-640-130)
Israel	1995	Male	85 to 89	48% (35-60%)	52% (40-65%)	0.57 (0.34-0.86)	-0.64% (-1.7-0.25%)	-0.81% (-2.1-0.23%)	-19 (-49-5.7)	-130 (-360-51)
Israel	1995	Male	90 to 94	48% (36-61%)	52% (39-64%)	0.54 (0.31-0.84)	-0.8% (-2.0-0.99%)	-1% (-2.4-0.076%)	-12 (-28-0.93)	-58 (-140-7.6)
Israel	1995	Male	95 plus	50% (35-65%)	50% (35-65%)	0.53 (0.27-0.91)	-0.9% (-2.1-0.093%)	-1.2% (-2.7-0.018%)	-5 (-11-0.07)	-15 (-37-1.5)
Israel	2000	Female	All Ages	36% (32-41%)	64% (59-68%)	0.45 (0.39-0.51)	0.45% (0.11-0.82%)	0.082% (-0.6-0.74%)	15 (-110-130)	2900 (750-5100)
Israel	2000	Female	15 to 19	41% (30-53%)	59% (47-70%)	0.47 (0.29-0.72)	0.84% (0.47-1.3%)	2.6% (0.94-4.6%)	1.7 (0.6-3.1)	220 (120-340)
Israel	2000	Female	20 to 24	46% (33-60%)	54% (40-67%)	0.65 (0.39-1)	1.2% (0.76-1.8%)	3.2% (1.3-5.5%)	2.2 (0.87-3.8)	340 (210-510)
Israel	2000	Female	25 to 29	47% (33-62%)	53% (38-67%)	0.64 (0.38-0.96)	1% (0.65-1.4%)	3% (1.4-4.7%)	2.1 (0.96-3.3)	290 (190-420)
Israel	2000	Female	30 to 34	48% (33-64%)	52% (36-67%)	0.65 (0.4-0.97)	1% (0.66-1.5%)	3.3% (2-4.9%)	3.1 (1.9-4.8)	290 (190-420)
Israel	2000	Female	35 to 39	48% (35-64%)	52% (36-65%)	0.7 (0.42-1.1)	1.1% (0.7-1.6%)	3.8% (2.4-5.5%)	4.8 (2.9-7)	340 (220-470)
Israel	2000	Female	40 to 44	46% (33-61%)	54% (39-67%)	0.74 (0.46-1.1)	1.2% (0.78-1.8%)	4.1% (2.7-5.9%)	7.5 (4.8-11)	410 (260-600)
Israel	2000	Female	45 to 49	41% (28-52%)	59% (48-72%)	0.74 (0.49-1)	1.3% (0.81-2%)	3.8% (2.4-5.3%)	12 (7.5-17)	530 (330-780)
Israel	2000	Female	50 to 54	38% (28-49%)	62% (51-72%)	0.7 (0.45-1)	1.3% (0.7-2%)	3.1% (1.9-4.4%)	14 (8.3-20)	500 (280-770)
Israel	2000	Female	55 to 59	37% (28-48%)	63% (52-72%)	0.61 (0.4-0.91)	0.86% (0.32-1.5%)	2% (1.1-3.2%)	9.6 (5.1-16)	270 (100-490)
Israel	2000	Female	60 to 64	36% (26-46%)	64% (54-74%)	0.56 (0.35-0.85)	0.44% (-0.19-1.2%)	1.1% (0.11-2.1%)	8.4 (0.89-17)	170 (-74-440)
Israel	2000	Female	65 to 69	35% (26-45%)	65% (55-74%)	0.54 (0.34-0.8)	0.21% (-0.42-0.9%)	0.6% (-0.31-1.6%)	7.7 (-3.9-21)	96 (-210-420)
Israel	2000	Female	70 to 74	34% (25-43%)	66% (57-75%)	0.51 (0.33-0.77)	0.031% (-0.67-0.74%)	0.25% (-0.63-1.2%)	5.3 (-13-26)	17 (-370-420)
Israel	2000	Female	75 to 79	31% (23-42%)	69% (58-77%)	0.46 (0.28-0.7)	-0.25% (-1-0.4%)	-0.22% (-1.2-0.66%)	-6.4 (-35-19)	-140 (-580-220)
Israel	2000	Female	80 to 84	30% (21-39%)	70% (61-79%)	0.42 (0.26-0.63)	-0.44% (-1.1-0.14%)	-0.49% (-1.4-0.24%)	-16 (-44-7.7)	-190 (-500-61)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	2000	Female	85 to 89	29% (20-39%)	71% (61-80%)	0.39 (0.23-0.61)	-0.56% (-1.3-0.002%)	-0.68% (-1.6-0.017%)	-19 (-44-0.49)	-150 (-340-0.49)
Israel	2000	Female	90 to 94	29% (20-39%)	71% (61-80%)	0.38 (0.21-0.62)	-0.63% (-1.3--0.068%)	-0.78% (-1.6--0.084%)	-15 (-30--1.7)	-75 (-160--8.1)
Israel	2000	Female	95 plus	30% (20-41%)	70% (59-80%)	0.38 (0.18-0.63)	-0.69% (-1.5--0.023%)	-0.88% (-1.9--0.079%)	-7.4 (-16--0.69)	-24 (-52--0.79)
Israel	2000	Male	All Ages	47% (41-53%)	53% (47-59%)	0.62 (0.54-0.7)	1.7% (1-2.5%)	1.1% (0.13-2.1%)	210 (23-400)	12000 (6800-17000)
Israel	2000	Male	15 to 19	48% (39-59%)	52% (41-61%)	0.46 (0.28-0.67)	2.1% (1-3.5%)	3.7% (1.2-7%)	6.5 (2.1-12)	650 (320-1100)
Israel	2000	Male	20 to 24	55% (44-68%)	45% (32-56%)	0.72 (0.45-1)	3.9% (2.2-6%)	5.5% (2.1-9.5%)	15 (5.6-26)	1500 (800-2300)
Israel	2000	Male	25 to 29	59% (45-72%)	41% (28-55%)	0.8 (0.51-1.1)	3.8% (2.3-5.5%)	5.8% (2.4-9.3%)	13 (5.6-22)	1200 (710-1800)
Israel	2000	Male	30 to 34	60% (45-75%)	40% (25-55%)	0.81 (0.51-1.2)	3.6% (2.3-5%)	6.2% (3.3-9.2%)	13 (6.6-19)	1000 (650-1400)
Israel	2000	Male	35 to 39	62% (47-76%)	38% (24-53%)	0.8 (0.51-1.2)	3.7% (2.5-5.1%)	6.3% (3.8-8.9%)	17 (10-24)	1100 (770-1500)
Israel	2000	Male	40 to 44	63% (47-78%)	37% (22-53%)	0.97 (0.64-1.4)	3.6% (2.5-4.8%)	6% (3.8-8.2%)	21 (13-30)	1200 (820-1600)
Israel	2000	Male	45 to 49	59% (43-74%)	41% (26-57%)	1.2 (0.73-1.8)	3.3% (2.1-4.5%)	5.3% (3.3-7.2%)	27 (17-39)	1400 (870-1900)
Israel	2000	Male	50 to 54	56% (42-69%)	44% (31-58%)	1.2 (0.79-1.8)	2.7% (1.6-3.8%)	3.9% (2.2-5.7%)	28 (15-41)	1200 (680-1700)
Israel	2000	Male	55 to 59	57% (50-65%)	43% (35-50%)	1.3 (0.92-1.7)	2% (0.9-3.2%)	2.8% (1.1-4.4%)	21 (7.6-33)	770 (330-1200)
Israel	2000	Male	60 to 64	57% (49-65%)	43% (35-51%)	1.3 (0.87-1.8)	1.6% (0.36-2.8%)	2% (0.4-3.7%)	23 (4.8-43)	710 (160-1300)
Israel	2000	Male	65 to 69	56% (48-64%)	44% (36-52%)	1.1 (0.79-1.5)	0.97% (-0.08-2.1%)	1.2% (-0.16-2.7%)	21 (-2.8-48)	520 (-47-1200)
Israel	2000	Male	70 to 74	54% (46-62%)	46% (38-54%)	0.93 (0.65-1.3)	0.46% (-0.44-1.5%)	0.56% (-0.57-1.8%)	13 (-14-43)	260 (-240-830)
Israel	2000	Male	75 to 79	49% (40-60%)	51% (40-60%)	0.75 (0.49-1.1)	0.15% (-0.67-1.1%)	0.16% (-0.84-1.3%)	4.8 (-24-38)	81 (-360-560)
Israel	2000	Male	80 to 84	45% (34-57%)	55% (43-66%)	0.64 (0.4-0.92)	-0.0069% (-0.82-0.73%)	-0.04% (-1-0.86%)	-1.1 (-27-24)	-2.3 (-290-260)
Israel	2000	Male	85 to 89	43% (31-54%)	57% (46-69%)	0.56 (0.33-0.86)	-0.14% (-0.92-0.5%)	-0.22% (-1.1-0.54%)	-4.7 (-24-12)	-27 (-180-97)
Israel	2000	Male	90 to 94	43% (31-54%)	57% (46-69%)	0.52 (0.3-0.82)	-0.3% (-1.1-0.34%)	-0.43% (-1.4-0.34%)	-5.8 (-19-4.7)	-25 (-92-28)
Israel	2000	Male	95 plus	45% (31-61%)	55% (39-69%)	0.51 (0.26-0.89)	-0.37% (-1.3-0.36%)	-0.55% (-1.7-0.33%)	-3.1 (-9.2-1.9)	-8.3 (-29-8.2)
Israel	2005	Female	All Ages	35% (31-39%)	65% (61-69%)	0.5 (0.43-0.58)	0.64% (0.34-0.97%)	0.42% (-0.18-1%)	82 (-35-200)	4400 (2300-6400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	2005	Female	15 to 19	40% (30-53%)	60% (47-70%)	0.54 (0.32-0.83)	0.86% (0.5-1.3%)	2.7% (1.1-4.7%)	1.6 (0.68-2.8)	220 (130-340)
Israel	2005	Female	20 to 24	46% (33-60%)	54% (40-67%)	0.74 (0.44-1.2)	1.3% (0.84-1.9%)	3.4% (1.6-5.5%)	2.5 (1.2-4.2)	390 (240-570)
Israel	2005	Female	25 to 29	47% (33-62%)	53% (38-67%)	0.73 (0.44-1.1)	1.1% (0.71-1.5%)	3.3% (1.7-5%)	2.4 (1.2-3.7)	350 (220-500)
Israel	2005	Female	30 to 34	48% (32-63%)	52% (37-68%)	0.75 (0.45-1.1)	1.1% (0.71-1.5%)	3.7% (2.2-5.3%)	3.4 (2-5.2)	350 (240-480)
Israel	2005	Female	35 to 39	48% (34-64%)	52% (36-66%)	0.8 (0.46-1.2)	1.2% (0.77-1.7%)	4.2% (2.7-6%)	5.2 (3.2-7.8)	380 (240-530)
Israel	2005	Female	40 to 44	45% (33-61%)	55% (39-67%)	0.85 (0.52-1.2)	1.4% (0.87-2.1%)	4.6% (3-6.6%)	8.6 (5.5-12)	480 (310-690)
Israel	2005	Female	45 to 49	39% (28-51%)	61% (49-72%)	0.84 (0.55-1.2)	1.5% (0.93-2.1%)	4.2% (2.8-5.7%)	12 (8.1-17)	560 (370-800)
Israel	2005	Female	50 to 54	35% (26-46%)	65% (54-74%)	0.74 (0.48-1.1)	1.3% (0.77-2%)	3.4% (2.2-4.6%)	15 (9.8-22)	580 (350-850)
Israel	2005	Female	55 to 59	33% (25-42%)	67% (58-75%)	0.62 (0.4-0.89)	1.1% (0.55-1.7%)	2.5% (1.5-3.7%)	16 (9.8-24)	490 (250-780)
Israel	2005	Female	60 to 64	32% (24-42%)	68% (58-76%)	0.56 (0.35-0.87)	0.73% (0.19-1.4%)	1.7% (0.86-2.7%)	12 (5.7-19)	270 (77-500)
Israel	2005	Female	65 to 69	32% (24-40%)	68% (60-76%)	0.56 (0.36-0.83)	0.6% (0.046-1.3%)	1.3% (0.5-2.2%)	14 (5.5-25)	260 (21-550)
Israel	2005	Female	70 to 74	30% (22-39%)	70% (61-78%)	0.51 (0.32-0.78)	0.41% (-0.17-1%)	0.84% (0.058-1.7%)	15 (1.1-30)	210 (-82-520)
Israel	2005	Female	75 to 79	28% (20-38%)	72% (62-80%)	0.46 (0.28-0.71)	0.13% (-0.46-0.72%)	0.31% (-0.45-1.1%)	8.8 (-12-31)	74 (-260-410)
Israel	2005	Female	80 to 84	28% (19-37%)	72% (63-81%)	0.45 (0.27-0.68)	-0.15% (-0.8-0.41%)	-0.11% (-0.9-0.58%)	-4.3 (-35-22)	-79 (-430-210)
Israel	2005	Female	85 to 89	27% (18-37%)	73% (63-82%)	0.43 (0.26-0.67)	-0.33% (-0.95-0.2%)	-0.39% (-1.2-0.26%)	-14 (-41-9.1)	-110 (-330-67)
Israel	2005	Female	90 to 94	27% (19-36%)	73% (64-81%)	0.41 (0.23-0.67)	-0.41% (-1-0.14%)	-0.51% (-1.2-0.16%)	-11 (-27-3.3)	-56 (-140-20)
Israel	2005	Female	95 plus	29% (19-40%)	71% (60-81%)	0.42 (0.21-0.7)	-0.48% (-1.2-0.16%)	-0.62% (-1.5-0.14%)	-6.3 (-15-1.4)	-20 (-50-6.7)
Israel	2005	Male	All Ages	47% (42-52%)	53% (48-58%)	0.72 (0.62-0.82)	2% (1.3-2.6%)	1.5% (0.65-2.4%)	290 (120-470)	14000 (9200-19000)
Israel	2005	Male	15 to 19	49% (41-60%)	51% (40-59%)	0.54 (0.33-0.8)	2.1% (1.1-3.3%)	3.9% (1.4-6.9%)	6.1 (2.1-11)	630 (330-990)
Israel	2005	Male	20 to 24	56% (46-68%)	44% (32-54%)	0.84 (0.54-1.2)	4% (2.3-5.9%)	5.7% (2.3-9.3%)	15 (6.1-26)	1500 (860-2300)
Israel	2005	Male	25 to 29	60% (48-73%)	40% (27-52%)	0.94 (0.63-1.3)	3.8% (2.4-5.2%)	6% (2.7-9.4%)	14 (6-22)	1300 (830-1900)
Israel	2005	Male	30 to 34	61% (47-74%)	39% (26-53%)	0.95 (0.6-1.3)	3.6% (2.4-4.8%)	6.3% (3.4-9.4%)	14 (7.7-21)	1200 (790-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	2005	Male	35 to 39	61% (47-75%)	39% (25-53%)	0.92 (0.58-1.3)	3.6% (2.4-4.8%)	6.5% (3.9-9%)	16 (9.1-22)	1100 (750-1500)
Israel	2005	Male	40 to 44	64% (49-77%)	36% (23-51%)	1.1 (0.73-1.6)	3.7% (2.6-5%)	6.5% (4.2-8.7%)	22 (14-31)	1300 (880-1700)
Israel	2005	Male	45 to 49	58% (42-73%)	42% (27-58%)	1.3 (0.84-2)	3.6% (2.4-5%)	6% (3.9-8.2%)	29 (18-41)	1400 (960-2000)
Israel	2005	Male	50 to 54	54% (40-67%)	46% (33-60%)	1.4 (0.93-2)	3.1% (2-4.5%)	4.9% (3.1-6.8%)	37 (23-53)	1600 (1000-2200)
Israel	2005	Male	55 to 59	55% (49-62%)	45% (38-51%)	1.5 (1-1.9)	2.5% (1.4-3.6%)	3.6% (2.1-5.3%)	36 (20-53)	1300 (730-1900)
Israel	2005	Male	60 to 64	56% (49-62%)	44% (38-51%)	1.4 (0.98-2)	2% (0.87-3.2%)	2.7% (1.2-4.3%)	29 (12-47)	880 (390-1400)
Israel	2005	Male	65 to 69	55% (48-61%)	45% (39-52%)	1.2 (0.9-1.7)	1.4% (0.44-2.5%)	1.8% (0.53-3.2%)	29 (8.7-53)	710 (210-1300)
Israel	2005	Male	70 to 74	53% (46-59%)	47% (41-54%)	1 (0.73-1.4)	0.87% (0.045-1.9%)	1.1% (0.064-2.3%)	24 (1.3-51)	470 (24-990)
Israel	2005	Male	75 to 79	48% (39-56%)	52% (44-61%)	0.83 (0.54-1.2)	0.51% (-0.19-1.4%)	0.61% (-0.23-1.6%)	17 (-6.6-45)	260 (-95-680)
Israel	2005	Male	80 to 84	42% (31-53%)	58% (47-69%)	0.67 (0.42-0.96)	0.3% (-0.35-0.99%)	0.34% (-0.45-1.2%)	10 (-13-37)	120 (-140-410)
Israel	2005	Male	85 to 89	38% (28-49%)	62% (51-72%)	0.57 (0.35-0.87)	0.094% (-0.51-0.69%)	0.062% (-0.65-0.76%)	1.7 (-18-20)	23 (-130-170)
Israel	2005	Male	90 to 94	38% (28-49%)	62% (51-72%)	0.54 (0.31-0.83)	-0.072% (-0.69-0.54%)	-0.16% (-0.9-0.55%)	-2.3 (-13-8)	-6.4 (-63-48)
Israel	2005	Male	95 plus	40% (26-55%)	60% (45-74%)	0.53 (0.26-0.92)	-0.13% (-0.89-0.52%)	-0.27% (-1.2-0.51%)	-1.7 (-7.5-3.2)	-3.3 (-23-13)
Israel	2010	Female	All Ages	34% (29-38%)	66% (62-71%)	0.58 (0.48-0.69)	0.78% (0.47-1.2%)	0.77% (0.17-1.5%)	160 (32-290)	5700 (3300-8200)
Israel	2010	Female	15 to 19	38% (27-51%)	62% (49-73%)	0.59 (0.33-0.95)	0.78% (0.46-1.2%)	2.8% (1.3-4.9%)	1.2 (0.54-2.2)	200 (120-310)
Israel	2010	Female	20 to 24	43% (29-58%)	57% (42-71%)	0.82 (0.45-1.3)	1.2% (0.82-1.8%)	3.7% (1.8-6.1%)	2.1 (0.98-3.5)	370 (240-550)
Israel	2010	Female	25 to 29	45% (30-61%)	55% (39-70%)	0.83 (0.48-1.3)	1.1% (0.73-1.5%)	3.6% (1.8-5.5%)	2.3 (1.1-3.6)	360 (230-510)
Israel	2010	Female	30 to 34	47% (30-64%)	53% (36-70%)	0.86 (0.5-1.3)	1.1% (0.69-1.5%)	4.1% (2.5-5.9%)	3.7 (2.2-5.6)	380 (260-560)
Israel	2010	Female	35 to 39	47% (31-63%)	53% (37-69%)	0.91 (0.52-1.4)	1.2% (0.77-1.8%)	4.8% (3-6.9%)	6.3 (3.7-9.4)	460 (300-660)
Israel	2010	Female	40 to 44	44% (29-59%)	56% (41-71%)	0.95 (0.55-1.4)	1.4% (0.84-2.1%)	5.2% (3.4-7.2%)	8.8 (5.4-13)	510 (310-720)
Israel	2010	Female	45 to 49	39% (27-50%)	61% (50-73%)	0.97 (0.62-1.4)	1.6% (0.99-2.3%)	4.9% (3.4-6.6%)	13 (8.3-18)	610 (390-850)
Israel	2010	Female	50 to 54	35% (26-46%)	65% (54-74%)	0.89 (0.58-1.3)	1.5% (0.93-2.2%)	4% (2.7-5.4%)	16 (11-23)	640 (390-930)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	2010	Female	55 to 59	33% (25-42%)	67% (58-75%)	0.75 (0.48-1.1)	1.3% (0.67-1.9%)	3.1% (2-4.3%)	21 (13-30)	660 (370-990)
Israel	2010	Female	60 to 64	32% (23-42%)	68% (58-77%)	0.68 (0.41-1)	0.93% (0.37-1.6%)	2.2% (1.2-3.3%)	19 (11-30)	480 (200-840)
Israel	2010	Female	65 to 69	32% (24-40%)	68% (60-76%)	0.66 (0.42-0.98)	0.82% (0.27-1.5%)	1.8% (0.98-2.7%)	17 (9.6-27)	350 (120-610)
Israel	2010	Female	70 to 74	30% (22-38%)	70% (62-78%)	0.61 (0.38-0.93)	0.72% (0.17-1.4%)	1.4% (0.56-2.2%)	22 (8.9-37)	360 (79-690)
Israel	2010	Female	75 to 79	28% (20-38%)	72% (62-80%)	0.56 (0.33-0.88)	0.43% (-0.11-1.1%)	0.79% (0.072-1.6%)	20 (1.8-42)	240 (-60-590)
Israel	2010	Female	80 to 84	28% (19-36%)	72% (64-81%)	0.53 (0.32-0.8)	0.15% (-0.44-0.76%)	0.29% (-0.44-1.1%)	11 (-18-41)	85 (-240-430)
Israel	2010	Female	85 to 89	27% (18-36%)	73% (64-82%)	0.5 (0.3-0.79)	-0.041% (-0.64-0.55%)	-0.029% (-0.76-0.69%)	-1.2 (-32-29)	-17 (-270-230)
Israel	2010	Female	90 to 94	27% (18-37%)	73% (63-82%)	0.48 (0.25-0.8)	-0.12% (-0.63-0.47%)	-0.16% (-0.78-0.55%)	-4.2 (-20-14)	-21 (-110-82)
Israel	2010	Female	95 plus	28% (18-39%)	72% (61-82%)	0.49 (0.23-0.86)	-0.2% (-0.88-0.53%)	-0.3% (-1.1-0.56%)	-3.6 (-13-6.8)	-9.9 (-44-26)
Israel	2010	Male	All Ages	41% (35-47%)	59% (53-65%)	0.77 (0.66-0.93)	1.8% (1.4-2.7%)	1.8% (0.91-2.6%)	360 (180-550)	15000 (10000-20000)
Israel	2010	Male	15 to 19	38% (27-51%)	62% (49-73%)	0.49 (0.26-0.81)	1.8% (1-2.9%)	3.5% (1.4-6.6%)	4.7 (1.8-8.6)	550 (310-850)
Israel	2010	Male	20 to 24	45% (30-60%)	55% (40-70%)	0.79 (0.44-1.3)	3.7% (2.3-5.6%)	5.5% (2.3-9.5%)	12 (5-21)	1300 (760-2000)
Israel	2010	Male	25 to 29	49% (32-65%)	51% (35-68%)	0.89 (0.53-1.4)	3.5% (2.3-4.9%)	6% (2.7-9.7%)	11 (5.1-18)	1200 (740-1700)
Israel	2010	Male	30 to 34	53% (34-71%)	47% (29-66%)	0.96 (0.56-1.5)	3.3% (2.2-4.5%)	6.3% (3.5-9.6%)	13 (6.9-20)	1200 (770-1600)
Israel	2010	Male	35 to 39	56% (39-74%)	44% (26-61%)	0.99 (0.58-1.5)	3.2% (2.1-4.4%)	6.2% (3.7-9%)	16 (9.1-23)	1200 (820-1600)
Israel	2010	Male	40 to 44	58% (40-75%)	42% (25-60%)	1.2 (0.73-1.7)	3.4% (2.5-4.7%)	6.5% (4.3-9.1%)	21 (14-30)	1300 (900-1700)
Israel	2010	Male	45 to 49	55% (39-72%)	45% (28-61%)	1.5 (0.87-2.3)	3.6% (2.5-5.1%)	6.5% (4.4-9.1%)	28 (19-42)	1400 (980-2000)
Israel	2010	Male	50 to 54	53% (40-66%)	47% (34-60%)	1.6 (0.99-2.3)	3.3% (2.1-4.7%)	5.4% (3.5-7.6%)	34 (21-49)	1500 (940-2100)
Israel	2010	Male	55 to 59	54% (47-60%)	46% (40-53%)	1.7 (1.2-2.2)	2.9% (1.7-4.1%)	4.3% (2.6-6%)	44 (26-64)	1700 (970-2400)
Israel	2010	Male	60 to 64	54% (47-61%)	46% (39-53%)	1.6 (1.1-2.2)	2.4% (1.3-3.7%)	3.4% (1.8-5.1%)	48 (25-73)	1500 (790-2400)
Israel	2010	Male	65 to 69	53% (47-60%)	47% (40-53%)	1.4 (0.99-1.9)	1.8% (0.83-3%)	2.4% (1.2-3.9%)	39 (18-64)	980 (440-1700)
Israel	2010	Male	70 to 74	51% (44-58%)	49% (42-56%)	1.2 (0.82-1.6)	1.2% (0.36-2.3%)	1.6% (0.5-2.9%)	33 (11-60)	650 (200-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	2010	Male	75 to 79	47% (38-56%)	53% (44-62%)	0.95 (0.62-1.3)	0.82% (0.11-1.7%)	1% (0.14-2%)	28 (4.2-59)	440 (63-910)
Israel	2010	Male	80 to 84	41% (30-52%)	59% (48-70%)	0.78 (0.49-1.1)	0.54% (-0.071-1.3%)	0.63% (-0.12-1.5%)	20 (-3.7-47)	230 (-31-550)
Israel	2010	Male	85 to 89	38% (27-49%)	62% (51-73%)	0.66 (0.38-1.1)	0.3% (-0.26-0.94%)	0.31% (-0.39-1.1%)	9.2 (-12-33)	86 (-73-270)
Israel	2010	Male	90 to 94	38% (27-48%)	62% (52-73%)	0.62 (0.35-0.98)	0.13% (-0.46-0.78%)	0.066% (-0.66-0.8%)	1.3 (-13-16)	16 (-58-96)
Israel	2010	Male	95 plus	39% (26-55%)	61% (45-74%)	0.61 (0.3-1.1)	0.075% (-0.67-0.8%)	-0.046% (-0.93-0.79%)	-0.35 (-7.1-6)	2.3 (-20-24)
Israel	2016	Female	All Ages	36% (32-41%)	64% (59-68%)	0.64 (0.49-0.85)	0.82% (0.39-1.3%)	0.76% (0.018-1.6%)	170 (4.4-380)	6300 (2900-10000)
Israel	2016	Female	15 to 19	38% (27-51%)	62% (49-73%)	0.6 (0.33-1)	0.76% (0.47-1.1%)	2.8% (1.2-5.1%)	1.2 (0.48-2.2)	220 (130-340)
Israel	2016	Female	20 to 24	43% (29-59%)	57% (41-71%)	0.84 (0.45-1.4)	1.2% (0.83-1.8%)	3.7% (1.7-6.3%)	1.9 (0.84-3.3)	380 (230-560)
Israel	2016	Female	25 to 29	46% (30-62%)	54% (38-70%)	0.85 (0.46-1.4)	1.1% (0.73-1.6%)	3.6% (1.9-5.7%)	2.1 (1-3.5)	360 (240-520)
Israel	2016	Female	30 to 34	47% (30-65%)	53% (35-70%)	0.88 (0.47-1.4)	1.1% (0.7-1.6%)	4.2% (2.4-6.2%)	3.5 (1.9-5.6)	400 (250-580)
Israel	2016	Female	35 to 39	48% (32-64%)	52% (36-68%)	0.95 (0.53-1.5)	1.3% (0.78-1.9%)	5% (3-7.4%)	6.7 (3.9-11)	510 (320-780)
Israel	2016	Female	40 to 44	46% (32-62%)	54% (38-68%)	1 (0.59-1.6)	1.4% (0.84-2.2%)	5.6% (3.5-7.9%)	10 (6-17)	610 (380-920)
Israel	2016	Female	45 to 49	44% (30-56%)	56% (44-70%)	1.1 (0.69-1.7)	1.7% (1-2.5%)	5.4% (3.5-7.4%)	14 (8.6-22)	690 (420-1000)
Israel	2016	Female	50 to 54	42% (30-54%)	58% (46-70%)	1.1 (0.66-1.7)	1.7% (0.9-2.6%)	4.5% (2.8-6.3%)	18 (10-27)	710 (390-1100)
Israel	2016	Female	55 to 59	41% (30-52%)	59% (48-70%)	0.96 (0.58-1.5)	1.4% (0.69-2.3%)	3.6% (2.2-5.2%)	21 (12-32)	660 (320-1100)
Israel	2016	Female	60 to 64	40% (29-51%)	60% (49-71%)	0.88 (0.52-1.4)	1% (0.31-1.9%)	2.5% (1.2-4%)	23 (11-39)	580 (170-1100)
Israel	2016	Female	65 to 69	39% (28-49%)	61% (51-72%)	0.85 (0.53-1.3)	0.86% (0.13-1.7%)	2% (0.93-3.3%)	27 (12-45)	530 (82-1000)
Israel	2016	Female	70 to 74	37% (28-47%)	63% (53-72%)	0.79 (0.48-1.3)	0.73% (-0.023-1.7%)	1.5% (0.47-2.7%)	24 (7.3-46)	380 (-11-860)
Israel	2016	Female	75 to 79	34% (24-46%)	66% (54-76%)	0.71 (0.4-1.2)	0.45% (-0.25-1.3%)	0.85% (-0.066-2%)	19 (-1.4-44)	230 (-130-690)
Israel	2016	Female	80 to 84	32% (22-42%)	68% (58-78%)	0.63 (0.36-0.98)	0.17% (-0.51-0.96%)	0.32% (-0.53-1.3%)	12 (-19-49)	92 (-280-540)
Israel	2016	Female	85 to 89	30% (21-40%)	70% (60-79%)	0.57 (0.32-0.93)	-0.02% (-0.73-0.7%)	-0.008% (-0.88-0.86%)	-0.41 (-43-44)	-10 (-350-350)
Israel	2016	Female	90 to 94	30% (20-40%)	70% (60-80%)	0.55 (0.28-0.93)	-0.1% (-0.73-0.6%)	-0.15% (-0.93-0.69%)	-5.4 (-32-23)	-25 (-180-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Israel	2016	Female	95 plus	32% (20-44%)	68% (56-80%)	0.56 (0.26-1)	-0.18% (-1-0.72%)	-0.29% (-1.3-0.74%)	-6.2 (-27-15)	-16 (-87-61)
Israel	2016	Male	All Ages	43% (37-49%)	57% (51-63%)	0.84 (0.63-1.1)	2.1% (1.4-3%)	1.8% (0.72-3%)	410 (170-710)	17000 (11000-25000)
Israel	2016	Male	15 to 19	38% (26-51%)	62% (49-74%)	0.5 (0.25-0.85)	1.8% (1-2.9%)	3.6% (1.3-6.6%)	4.8 (1.7-9.3)	580 (330-940)
Israel	2016	Male	20 to 24	45% (30-61%)	55% (39-70%)	0.8 (0.42-1.3)	3.7% (2.3-5.6%)	5.6% (2.2-9.8%)	12 (4.7-22)	1300 (760-2000)
Israel	2016	Male	25 to 29	49% (32-66%)	51% (34-68%)	0.91 (0.52-1.4)	3.5% (2.2-5.1%)	6.1% (2.7-10%)	11 (4.7-20)	1200 (730-1800)
Israel	2016	Male	30 to 34	53% (34-70%)	47% (30-66%)	0.98 (0.55-1.5)	3.3% (2.1-4.6%)	6.5% (3.5-10%)	13 (6.8-21)	1200 (760-1700)
Israel	2016	Male	35 to 39	57% (38-74%)	43% (26-62%)	1 (0.57-1.6)	3.2% (2.2-4.6%)	6.5% (3.7-9.5%)	17 (9.2-27)	1300 (840-1900)
Israel	2016	Male	40 to 44	59% (41-76%)	41% (24-59%)	1.2 (0.72-1.9)	3.4% (2.3-4.9%)	6.8% (4.3-9.6%)	25 (14-39)	1500 (960-2200)
Israel	2016	Male	45 to 49	59% (40-75%)	41% (25-60%)	1.6 (0.88-2.6)	3.8% (2.4-5.6%)	7.1% (4.7-10%)	32 (18-51)	1600 (1000-2500)
Israel	2016	Male	50 to 54	59% (43-73%)	41% (27-57%)	1.8 (1.1-2.8)	3.5% (2.1-5.3%)	6% (3.7-8.8%)	37 (21-59)	1600 (950-2600)
Israel	2016	Male	55 to 59	58% (45-68%)	42% (32-55%)	1.8 (1.2-2.7)	3% (1.8-4.7%)	4.7% (2.8-7.2%)	41 (23-67)	1600 (870-2500)
Israel	2016	Male	60 to 64	57% (45-68%)	43% (32-55%)	1.7 (1.1-2.6)	2.6% (1.3-4.3%)	3.7% (1.9-6%)	55 (26-91)	1700 (830-2900)
Israel	2016	Male	65 to 69	58% (46-67%)	42% (33-54%)	1.6 (1-2.3)	2% (0.82-3.6%)	2.7% (1.2-4.7%)	62 (25-110)	1600 (570-2800)
Israel	2016	Male	70 to 74	56% (46-66%)	44% (34-54%)	1.4 (0.83-2)	1.4% (0.3-2.7%)	1.8% (0.47-3.4%)	41 (11-81)	830 (180-1700)
Israel	2016	Male	75 to 79	54% (42-66%)	46% (34-58%)	1.2 (0.69-1.8)	0.93% (0.0064-2.2%)	1.1% (0.0032-2.6%)	31 (0.083-70)	480 (2.8-1100)
Israel	2016	Male	80 to 84	50% (37-62%)	50% (38-63%)	1 (0.61-1.5)	0.59% (-0.23-1.7%)	0.68% (-0.3-1.8%)	22 (-10-62)	270 (-100-740)
Israel	2016	Male	85 to 89	47% (34-59%)	53% (41-66%)	0.87 (0.49-1.4)	0.32% (-0.42-1.2%)	0.3% (-0.62-1.3%)	11 (-21-46)	110 (-140-400)
Israel	2016	Male	90 to 94	47% (35-60%)	53% (40-65%)	0.82 (0.44-1.4)	0.11% (-0.64-1%)	0.013% (-0.9-1%)	0.28 (-23-26)	17 (-110-160)
Israel	2016	Male	95 plus	49% (34-64%)	51% (36-66%)	0.81 (0.38-1.5)	0.032% (-0.89-0.97%)	-0.13% (-1.2-0.95%)	-1.8 (-17-13)	1.7 (-50-54)
Italy	1990	Female	All Ages	62% (57-66%)	38% (34-43%)	3.5 (2.8-4.2)	6.6% (4.8-8.9%)	10% (6.8-15%)	27000 (18000-39000)	510000 (370000-690000)
Italy	1990	Female	15 to 19	64% (49-77%)	36% (23-51%)	2.6 (1.6-3.6)	2.4% (1.3-3.7%)	10% (5-15%)	55 (28-86)	5400 (3100-8100)
Italy	1990	Female	20 to 24	72% (60-82%)	28% (18-40%)	3.2 (2-4.6)	2.8% (1.7-4.1%)	11% (5.9-16%)	80 (43-120)	7900 (5000-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	1990	Female	25 to 29	75% (63-83%)	25% (17-37%)	3.5 (2.3-4.9)	2.8% (1.7-3.9%)	10% (6.1-14%)	93 (57-130)	8500 (5500-12000)
Italy	1990	Female	30 to 34	75% (62-83%)	25% (17-38%)	3.8 (2.6-5.3)	3.2% (2.1-4.3%)	11% (7.7-15%)	120 (81-160)	9400 (6300-13000)
Italy	1990	Female	35 to 39	73% (62-81%)	27% (19-38%)	4.2 (2.9-5.9)	3.9% (2.7-5.4%)	13% (9.6-17%)	180 (130-240)	12000 (8600-16000)
Italy	1990	Female	40 to 44	70% (60-78%)	30% (22-40%)	4.4 (3.2-6)	4.9% (3.5-6.6%)	14% (11-18%)	320 (240-410)	18000 (13000-24000)
Italy	1990	Female	45 to 49	65% (54-73%)	35% (27-46%)	4.5 (3.2-6.2)	6.1% (4.3-8.2%)	14% (11-19%)	460 (340-600)	23000 (16000-30000)
Italy	1990	Female	50 to 54	63% (54-71%)	37% (29-46%)	4.8 (3.5-6.5)	7.2% (5.2-9.9%)	14% (11-19%)	760 (570-1000)	33000 (24000-45000)
Italy	1990	Female	55 to 59	63% (53-71%)	37% (29-47%)	5.3 (3.9-7)	8.5% (6.1-12%)	15% (11-20%)	1200 (900-1600)	44000 (32000-61000)
Italy	1990	Female	60 to 64	63% (52-71%)	37% (29-48%)	5.3 (3.7-7.3)	9% (6.1-12%)	14% (9.9-19%)	1800 (1300-2400)	56000 (38000-78000)
Italy	1990	Female	65 to 69	61% (51-70%)	39% (30-49%)	4.8 (3.5-6.3)	9.1% (6.1-13%)	13% (9-18%)	2600 (1800-3600)	68000 (45000-92000)
Italy	1990	Female	70 to 74	59% (49-67%)	41% (33-51%)	4.2 (3-5.8)	8.7% (5.5-13%)	12% (7.4-17%)	2500 (1600-3600)	51000 (32000-74000)
Italy	1990	Female	75 to 79	55% (42-66%)	45% (34-58%)	3.8 (2.5-5.6)	8.1% (4.3-13%)	10% (5.5-16%)	4500 (2500-7100)	70000 (38000-110000)
Italy	1990	Female	80 to 84	51% (39-63%)	49% (37-61%)	3.7 (2.5-5.2)	7.9% (4.5-13%)	9.6% (5.4-15%)	5400 (3000-8600)	60000 (34000-96000)
Italy	1990	Female	85 to 89	48% (35-60%)	52% (40-65%)	3.5 (2.2-5.2)	7.4% (3.7-13%)	8.7% (4.1-15%)	4200 (2000-7200)	33000 (16000-56000)
Italy	1990	Female	90 to 94	47% (35-59%)	53% (41-65%)	3.5 (2-5.3)	7.2% (2.8-13%)	8.4% (3.2-15%)	2200 (820-4000)	12000 (4600-21000)
Italy	1990	Female	95 plus	48% (33-63%)	52% (37-67%)	3.6 (1.9-5.8)	7.6% (2.3-14%)	8.8% (2.5-17%)	740 (210-1400)	2600 (780-4800)
Italy	1990	Male	All Ages	78% (73-82%)	22% (18-27%)	6.1 (5.1-7.4)	12% (10-15%)	16% (12-20%)	46000 (35000-57000)	1100000 (890000-1300000)
Italy	1990	Male	15 to 19	69% (54-82%)	31% (18-46%)	2.8 (1.8-4)	6.4% (3.4-9.6%)	13% (6.2-20%)	230 (110-350)	18000 (10000-28000)
Italy	1990	Male	20 to 24	80% (69-88%)	20% (12-31%)	4.7 (3.3-6.4)	11% (7.1-15%)	20% (12-27%)	510 (320-720)	39000 (26000-54000)
Italy	1990	Male	25 to 29	84% (74-90%)	16% (9.9-26%)	6.5 (4.8-8.6)	11% (7.9-15%)	20% (14-27%)	610 (410-810)	43000 (31000-57000)
Italy	1990	Male	30 to 34	87% (78-93%)	13% (7.1-22%)	7 (5.2-9.2)	11% (8-14%)	20% (14-26%)	560 (400-720)	37000 (28000-47000)
Italy	1990	Male	35 to 39	89% (82-94%)	11% (6-18%)	6 (4.4-8.2)	9.8% (7-13%)	18% (13-24%)	520 (360-690)	32000 (23000-42000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	1990	Male	40 to 44	89% (83-94%)	11% (6.1-17%)	6.6 (4.8-8.7)	11% (7.8-14%)	19% (14-24%)	780 (560-1000)	42000 (31000-54000)
Italy	1990	Male	45 to 49	88% (81-93%)	12% (7.1-19%)	8.7 (6.4-11)	14% (11-17%)	21% (17-25%)	1300 (1000-1500)	59000 (48000-71000)
Italy	1990	Male	50 to 54	86% (79-91%)	14% (9.3-21%)	10 (7.6-13)	15% (12-18%)	21% (17-25%)	2100 (1800-2500)	87000 (71000-100000)
Italy	1990	Male	55 to 59	83% (77-88%)	17% (12-23%)	11 (8.2-13)	15% (12-18%)	19% (15-24%)	3300 (2500-4000)	110000 (88000-140000)
Italy	1990	Male	60 to 64	83% (76-88%)	17% (12-24%)	11 (8.1-14)	15% (11-18%)	18% (13-22%)	4700 (3500-5800)	140000 (100000-170000)
Italy	1990	Male	65 to 69	84% (78-89%)	16% (11-22%)	10 (8-13)	15% (11-19%)	17% (13-22%)	6100 (4500-7700)	150000 (110000-190000)
Italy	1990	Male	70 to 74	84% (78-88%)	16% (12-22%)	9.5 (7.4-12)	15% (11-18%)	17% (12-21%)	5100 (3800-6500)	100000 (77000-130000)
Italy	1990	Male	75 to 79	81% (73-88%)	19% (12-27%)	8.6 (6.5-11)	14% (10-18%)	15% (11-20%)	7700 (5700-10000)	120000 (89000-150000)
Italy	1990	Male	80 to 84	79% (69-86%)	21% (14-31%)	7.9 (6.1-10)	13% (9.6-17%)	15% (10-19%)	6700 (4800-8700)	75000 (54000-97000)
Italy	1990	Male	85 to 89	78% (67-86%)	22% (14-33%)	7.4 (5.4-9.9)	12% (8.8-17%)	14% (9.5-19%)	3700 (2500-5000)	29000 (20000-39000)
Italy	1990	Male	90 to 94	79% (69-86%)	21% (14-31%)	7.2 (4.8-9.9)	12% (7.4-16%)	13% (8.2-18%)	1400 (860-1900)	7400 (4800-10000)
Italy	1990	Male	95 plus	80% (67-89%)	20% (11-33%)	6.9 (4.1-10)	11% (5.8-16%)	13% (6.1-18%)	330 (160-480)	1200 (610-1700)
Italy	1995	Female	All Ages	64% (61-68%)	36% (32-39%)	3.1 (2.6-3.7)	5.3% (3.6-7.2%)	7.8% (4.7-12%)	21000 (13000-32000)	410000 (290000-560000)
Italy	1995	Female	15 to 19	66% (53-77%)	34% (23-47%)	2.2 (1.4-3.2)	2.1% (1.2-3.2%)	8.9% (4.3-14%)	39 (19-61)	4000 (2200-6100)
Italy	1995	Female	20 to 24	75% (65-83%)	25% (17-35%)	2.8 (1.8-4.1)	2.5% (1.5-3.7%)	9.6% (5.1-14%)	62 (32-92)	6500 (4000-9600)
Italy	1995	Female	25 to 29	77% (68-84%)	23% (16-32%)	3 (2.1-4.2)	2.3% (1.5-3.3%)	8% (4.6-11%)	75 (43-110)	7300 (4600-10000)
Italy	1995	Female	30 to 34	77% (68-84%)	23% (16-32%)	3.3 (2.4-4.5)	2.5% (1.7-3.4%)	7.6% (5.2-10%)	100 (70-140)	8500 (5800-12000)
Italy	1995	Female	35 to 39	73% (65-81%)	27% (19-35%)	3.6 (2.4-5)	3.1% (2.2-4.4%)	10% (7.2-13%)	150 (110-200)	10000 (7200-14000)
Italy	1995	Female	40 to 44	72% (66-78%)	28% (22-34%)	3.9 (2.8-5.2)	4.1% (2.8-5.6%)	12% (8.9-15%)	240 (180-310)	14000 (10000-19000)
Italy	1995	Female	45 to 49	70% (65-75%)	30% (25-35%)	4.1 (2.9-5.5)	5.3% (3.7-7.1%)	13% (9.7-16%)	440 (330-560)	22000 (16000-29000)
Italy	1995	Female	50 to 54	66% (61-72%)	34% (28-39%)	4.3 (3.1-5.7)	6.1% (4.3-8.4%)	13% (9.5-16%)	580 (440-760)	25000 (18000-34000)
Italy	1995	Female	55 to 59	65% (59-71%)	35% (29-41%)	4.6 (3.5-6.1)	7.1% (5-9.6%)	13% (9.7-17%)	960 (710-1300)	36000 (26000-48000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	1995	Female	60 to 64	67% (62-72%)	33% (28-38%)	4.8 (3.6-6.4)	7.6% (5.1-10%)	12% (8.7-16%)	1400 (970-1900)	44000 (29000-60000)
Italy	1995	Female	65 to 69	65% (57-71%)	35% (29-43%)	4.3 (3.3-5.6)	7.3% (4.8-10%)	11% (7.3-15%)	2000 (1300-2700)	51000 (34000-71000)
Italy	1995	Female	70 to 74	59% (48-67%)	41% (33-52%)	3.5 (2.5-4.9)	6.6% (3.6-10%)	9% (5-13%)	2600 (1500-4000)	55000 (30000-84000)
Italy	1995	Female	75 to 79	54% (42-66%)	46% (34-58%)	3.2 (2.1-4.5)	5.9% (2.7-9.9%)	7.5% (3.3-13%)	2300 (1000-3900)	37000 (16000-61000)
Italy	1995	Female	80 to 84	51% (38-62%)	49% (38-62%)	3 (2-4.3)	5.7% (2.7-9.5%)	7% (3.2-12%)	4100 (1900-6800)	47000 (22000-77000)
Italy	1995	Female	85 to 89	48% (36-60%)	52% (40-64%)	2.9 (1.8-4.3)	5.4% (2.3-9.5%)	6.3% (2.6-12%)	3600 (1500-6600)	29000 (12000-53000)
Italy	1995	Female	90 to 94	47% (35-60%)	53% (40-65%)	2.9 (1.7-4.4)	5.2% (1.8-10%)	6.1% (2-12%)	2000 (640-4000)	11000 (3800-21000)
Italy	1995	Female	95 plus	48% (32-63%)	52% (37-68%)	3 (1.6-4.9)	5.7% (1.3-11%)	6.6% (1.4-13%)	790 (160-1600)	2800 (670-5600)
Italy	1995	Male	All Ages	81% (77-85%)	19% (15-23%)	5.5 (4.6-6.6)	11% (9.2-13%)	14% (12-17%)	41000 (33000-50000)	970000 (810000-1200000)
Italy	1995	Male	15 to 19	74% (61-84%)	26% (16-39%)	2.4 (1.6-3.4)	5.7% (2.8-8.6%)	12% (5.4-18%)	160 (72-240)	13000 (6900-19000)
Italy	1995	Male	20 to 24	84% (76-90%)	16% (9.9-24%)	4.1 (3-5.6)	9.9% (6.1-14%)	18% (11-25%)	400 (240-570)	32000 (20000-43000)
Italy	1995	Male	25 to 29	87% (81-92%)	13% (8.5-19%)	5.7 (4.3-7.5)	9.9% (6.8-13%)	18% (12-24%)	490 (320-660)	36000 (25000-47000)
Italy	1995	Male	30 to 34	89% (83-94%)	11% (6.4-17%)	6.1 (4.5-7.9)	8.1% (5.8-11%)	13% (9.2-18%)	490 (330-650)	34000 (24000-44000)
Italy	1995	Male	35 to 39	91% (86-95%)	8.9% (5.4-14%)	5.1 (3.7-6.9)	7.3% (5-10%)	12% (8.2-17%)	430 (280-590)	27000 (19000-37000)
Italy	1995	Male	40 to 44	91% (86-95%)	8.8% (5.4-14%)	5.5 (4.2-7.3)	8.5% (5.9-11%)	15% (10-20%)	560 (390-760)	31000 (22000-42000)
Italy	1995	Male	45 to 49	90% (85-94%)	10% (6.3-15%)	7.4 (5.6-9.5)	12% (8.9-14%)	19% (14-23%)	1100 (870-1400)	55000 (42000-66000)
Italy	1995	Male	50 to 54	88% (83-92%)	12% (8.5-17%)	8.7 (6.6-11)	13% (11-16%)	19% (16-23%)	1700 (1400-2000)	69000 (57000-82000)
Italy	1995	Male	55 to 59	86% (82-89%)	14% (11-18%)	9.5 (7.4-12)	14% (11-17%)	18% (15-22%)	2800 (2300-3300)	98000 (80000-120000)
Italy	1995	Male	60 to 64	86% (82-89%)	14% (11-18%)	9.6 (7.4-13)	14% (11-17%)	17% (14-21%)	3900 (3100-4800)	120000 (95000-140000)
Italy	1995	Male	65 to 69	86% (81-90%)	14% (10-19%)	9 (7-11)	14% (11-17%)	16% (13-20%)	5500 (4300-6800)	140000 (110000-170000)
Italy	1995	Male	70 to 74	84% (77-88%)	16% (12-23%)	8 (6.2-10)	13% (10-16%)	15% (12-19%)	6500 (5100-8100)	130000 (100000-160000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	1995	Male	75 to 79	81% (72-88%)	19% (12-28%)	7.2 (5.5-9.4)	12% (9.2-15%)	14% (10-17%)	4800 (3600-6100)	74000 (56000-94000)
Italy	1995	Male	80 to 84	79% (68-87%)	21% (13-32%)	6.7 (5-8.5)	12% (8.4-15%)	13% (9.3-17%)	6500 (4700-8400)	73000 (54000-94000)
Italy	1995	Male	85 to 89	78% (67-85%)	22% (15-33%)	6.3 (4.5-8.2)	11% (7.2-14%)	12% (7.9-16%)	4200 (2800-5600)	33000 (23000-45000)
Italy	1995	Male	90 to 94	79% (69-86%)	21% (14-31%)	6.1 (4.1-8.2)	10% (5.5-14%)	11% (5.9-16%)	1500 (810-2200)	8300 (4600-12000)
Italy	1995	Male	95 plus	80% (67-89%)	20% (11-33%)	5.9 (3.6-8.7)	9.7% (3.8-15%)	11% (4-16%)	380 (140-590)	1400 (550-2100)
Italy	2000	Female	All Ages	60% (57-64%)	40% (36-43%)	2.6 (2.1-3.1)	4% (2.6-5.6%)	5.8% (3.2-9.1%)	16000 (9200-25000)	310000 (200000-430000)
Italy	2000	Female	15 to 19	61% (47-74%)	39% (26-53%)	1.8 (1.1-2.6)	1.7% (0.91-2.6%)	7.5% (3.5-12%)	25 (12-42)	2600 (1400-4000)
Italy	2000	Female	20 to 24	71% (58-80%)	29% (20-42%)	2.3 (1.4-3.4)	2.1% (1.3-3.2%)	8.6% (4.4-13%)	42 (21-66)	4400 (2600-6700)
Italy	2000	Female	25 to 29	75% (65-83%)	25% (17-35%)	2.6 (1.8-3.6)	2% (1.3-2.9%)	8.7% (4.9-13%)	58 (32-85)	5700 (3600-8000)
Italy	2000	Female	30 to 34	75% (66-83%)	25% (17-34%)	2.8 (2-3.9)	2.2% (1.4-3.1%)	9% (5.9-12%)	84 (55-120)	7000 (4700-9500)
Italy	2000	Female	35 to 39	72% (63-80%)	28% (20-37%)	3 (2.1-4.3)	2.7% (1.8-3.7%)	9.9% (7.1-13%)	140 (99-180)	9500 (6600-13000)
Italy	2000	Female	40 to 44	70% (62-77%)	30% (23-38%)	3.3 (2.4-4.4)	3.4% (2.4-4.7%)	11% (7.9-14%)	210 (150-270)	12000 (8600-16000)
Italy	2000	Female	45 to 49	66% (59-73%)	34% (27-41%)	3.4 (2.4-4.6)	4.2% (2.9-5.8%)	11% (8.2-14%)	310 (230-400)	16000 (11000-21000)
Italy	2000	Female	50 to 54	62% (55-69%)	38% (31-45%)	3.5 (2.5-4.7)	4.9% (3.4-6.7%)	11% (8.1-14%)	500 (380-660)	22000 (16000-29000)
Italy	2000	Female	55 to 59	62% (55-69%)	38% (31-45%)	3.8 (2.9-5)	5.6% (3.9-7.8%)	11% (8-14%)	660 (500-870)	25000 (18000-34000)
Italy	2000	Female	60 to 64	61% (53-67%)	39% (33-47%)	3.7 (2.7-5.1)	5.7% (3.7-8.2%)	9.7% (6.8-13%)	980 (690-1300)	31000 (21000-44000)
Italy	2000	Female	65 to 69	57% (48-65%)	43% (35-52%)	3.2 (2.4-4.3)	5.2% (3.2-7.5%)	8.1% (5.4-11%)	1300 (850-1800)	34000 (21000-48000)
Italy	2000	Female	70 to 74	54% (46-62%)	46% (38-54%)	2.8 (2-3.9)	4.8% (2.4-7.5%)	6.8% (3.6-11%)	1800 (940-2700)	36000 (19000-57000)
Italy	2000	Female	75 to 79	49% (39-60%)	51% (40-61%)	2.5 (1.6-3.6)	4.2% (1.8-7.3%)	5.5% (2.4-9.4%)	2300 (1000-4000)	37000 (16000-63000)
Italy	2000	Female	80 to 84	47% (35-58%)	53% (42-65%)	2.4 (1.6-3.4)	4.1% (1.7-7%)	5.1% (2.1-8.9%)	2100 (870-3700)	24000 (11000-42000)
Italy	2000	Female	85 to 89	46% (34-58%)	54% (42-66%)	2.4 (1.5-3.6)	4% (1.5-7.3%)	4.8% (1.6-8.9%)	3100 (1000-5700)	25000 (9500-46000)
Italy	2000	Female	90 to 94	45% (33-57%)	55% (43-67%)	2.4 (1.4-3.7)	3.9% (1-7.7%)	4.6% (1-9.2%)	2000 (460-4100)	11000 (2900-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	2000	Female	95 plus	46% (31-61%)	54% (39-69%)	2.5 (1.3-4.1)	4.4% (0.59-9.2%)	5.1% (0.42-11%)	850 (70-1800)	3000 (390-6300)
Italy	2000	Male	All Ages	81% (77-85%)	19% (15-23%)	4.8 (4-5.8)	9.9% (8-12%)	13% (9.6-16%)	35000 (27000-44000)	810000 (650000-970000)
Italy	2000	Male	15 to 19	71% (58-82%)	29% (18-42%)	2 (1.4-2.9)	4.7% (2.3-7.1%)	11% (4.7-17%)	97 (44-150)	8200 (4200-12000)
Italy	2000	Male	20 to 24	82% (74-89%)	18% (11-26%)	3.5 (2.5-4.7)	9% (5.5-13%)	17% (9.3-25%)	280 (160-410)	22000 (14000-31000)
Italy	2000	Male	25 to 29	87% (81-91%)	13% (8.8-19%)	4.9 (3.6-6.4)	9.6% (6.2-13%)	20% (12-27%)	410 (250-570)	30000 (20000-40000)
Italy	2000	Male	30 to 34	89% (83-93%)	11% (6.5-17%)	5.2 (3.9-6.9)	8.2% (5.5-11%)	17% (11-23%)	410 (270-560)	29000 (20000-38000)
Italy	2000	Male	35 to 39	91% (86-95%)	9.2% (5.4-14%)	4.5 (3.3-6.1)	6.8% (4.5-9.3%)	13% (8.7-18%)	390 (260-550)	25000 (17000-35000)
Italy	2000	Male	40 to 44	91% (86-95%)	9% (5.4-14%)	4.9 (3.7-6.4)	7.3% (5-10%)	13% (9.1-19%)	470 (320-650)	27000 (18000-37000)
Italy	2000	Male	45 to 49	90% (85-94%)	10% (6.3-15%)	6.5 (4.8-8.3)	10% (7.2-13%)	17% (12-21%)	850 (600-1100)	41000 (30000-52000)
Italy	2000	Male	50 to 54	88% (84-91%)	12% (8.6-16%)	7.5 (5.7-9.4)	12% (9.1-14%)	18% (14-21%)	1500 (1200-1800)	63000 (49000-76000)
Italy	2000	Male	55 to 59	85% (82-89%)	15% (11-18%)	7.9 (6.2-9.9)	12% (9.8-15%)	17% (14-20%)	2000 (1700-2400)	73000 (59000-87000)
Italy	2000	Male	60 to 64	85% (82-89%)	15% (11-18%)	7.9 (6.1-10)	12% (9.8-15%)	16% (12-19%)	3100 (2500-3700)	94000 (76000-110000)
Italy	2000	Male	65 to 69	86% (82-90%)	14% (10-18%)	7.6 (6.1-9.5)	12% (9.4-15%)	14% (11-18%)	4200 (3300-5100)	110000 (85000-130000)
Italy	2000	Male	70 to 74	85% (81-89%)	15% (11-19%)	7 (5.4-8.8)	11% (8.5-14%)	13% (10-17%)	5400 (4000-6800)	110000 (81000-140000)
Italy	2000	Male	75 to 79	83% (77-88%)	17% (12-23%)	6.4 (4.8-8.2)	10% (7.4-14%)	12% (8.3-15%)	5900 (4100-7700)	92000 (65000-120000)
Italy	2000	Male	80 to 84	80% (71-87%)	20% (13-29%)	5.8 (4.5-7.5)	9.7% (6.7-13%)	11% (7.4-15%)	3900 (2700-5300)	44000 (31000-59000)
Italy	2000	Male	85 to 89	78% (68-85%)	22% (15-32%)	5.4 (4-7.1)	8.9% (5.4-12%)	10% (5.9-14%)	4200 (2400-5800)	33000 (20000-47000)
Italy	2000	Male	90 to 94	79% (69-86%)	21% (14-31%)	5.2 (3.5-7.2)	8.2% (4-12%)	9.1% (4.2-14%)	1800 (820-2700)	9700 (4700-15000)
Italy	2000	Male	95 plus	80% (67-89%)	20% (11-33%)	5 (3-7.4)	7.9% (2.2-13%)	8.7% (2.1-14%)	400 (95-650)	1400 (410-2300)
Italy	2005	Female	All Ages	59% (55-62%)	41% (38-45%)	2.3 (1.9-2.8)	3.3% (2.1-4.6%)	4.8% (2.5-7.6%)	14000 (7100-22000)	250000 (170000-350000)
Italy	2005	Female	15 to 19	61% (47-74%)	39% (26-53%)	1.6 (0.99-2.4)	1.4% (0.79-2.1%)	6.8% (3.2-11%)	17 (8-27)	1900 (1100-3000)
Italy	2005	Female	20 to 24	71% (60-79%)	29% (21-40%)	2.1 (1.3-3.1)	1.8% (1.1-2.6%)	7.8% (4.1-12%)	26 (14-41)	3200 (1900-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	2005	Female	25 to 29	75% (66-82%)	25% (18-34%)	2.3 (1.6-3.2)	1.7% (1-2.4%)	7.7% (4.4-11%)	35 (20-51)	4000 (2500-5600)
Italy	2005	Female	30 to 34	75% (66-82%)	25% (18-34%)	2.5 (1.8-3.5)	1.8% (1.2-2.5%)	8.1% (5.3-11%)	58 (38-80)	5400 (3600-7700)
Italy	2005	Female	35 to 39	71% (62-79%)	29% (21-38%)	2.7 (1.9-3.8)	2.2% (1.4-3.1%)	9% (6.4-12%)	110 (76-140)	7800 (5300-11000)
Italy	2005	Female	40 to 44	68% (60-75%)	32% (25-40%)	2.9 (2.1-3.9)	2.9% (2-4%)	9.6% (7.1-12%)	190 (140-250)	11000 (7900-15000)
Italy	2005	Female	45 to 49	65% (57-71%)	35% (29-43%)	3 (2.1-4)	3.6% (2.5-4.9%)	9.8% (7.5-13%)	280 (210-350)	14000 (9900-19000)
Italy	2005	Female	50 to 54	61% (54-67%)	39% (33-46%)	3.1 (2.2-4.1)	4.2% (2.8-5.7%)	9.5% (7.2-12%)	390 (300-500)	17000 (12000-23000)
Italy	2005	Female	55 to 59	62% (56-67%)	38% (33-44%)	3.4 (2.6-4.6)	4.8% (3.3-6.6%)	9.6% (7.3-13%)	620 (470-810)	23000 (17000-32000)
Italy	2005	Female	60 to 64	59% (53-66%)	41% (34-47%)	3.3 (2.4-4.4)	4.8% (3-6.8%)	8.5% (6-12%)	730 (510-990)	23000 (15000-33000)
Italy	2005	Female	65 to 69	55% (48-62%)	45% (38-52%)	2.9 (2.1-3.7)	4.3% (2.7-6.1%)	7% (4.7-9.6%)	1000 (670-1400)	27000 (17000-37000)
Italy	2005	Female	70 to 74	53% (46-59%)	47% (41-54%)	2.5 (1.8-3.4)	3.9% (2.1-6%)	5.9% (3.3-8.8%)	1300 (730-2000)	28000 (15000-43000)
Italy	2005	Female	75 to 79	47% (38-57%)	53% (43-62%)	2.1 (1.4-3)	3.4% (1.4-5.7%)	4.6% (2-7.6%)	1700 (760-2900)	28000 (12000-47000)
Italy	2005	Female	80 to 84	45% (34-56%)	55% (44-66%)	2.1 (1.4-3)	3.3% (1.3-5.7%)	4.2% (1.6-7.5%)	2500 (950-4400)	29000 (12000-51000)
Italy	2005	Female	85 to 89	46% (34-57%)	54% (43-66%)	2.2 (1.3-3.2)	3.2% (1-5.9%)	3.9% (1.1-7.2%)	1900 (520-3500)	16000 (4700-29000)
Italy	2005	Female	90 to 94	45% (33-56%)	55% (44-67%)	2.2 (1.3-3.3)	3.1% (0.56-6.3%)	3.6% (0.49-7.6%)	1900 (260-4100)	11000 (2000-23000)
Italy	2005	Female	95 plus	46% (31-60%)	54% (40-69%)	2.2 (1.2-3.7)	3.4% (0.039-7.7%)	3.9% (-0.2-9.1%)	950 (-48-2200)	3500 (39-7600)
Italy	2005	Male	All Ages	81% (78-85%)	19% (15-22%)	4.4 (3.6-5.2)	8.7% (6.7-11%)	11% (7.9-14%)	31000 (22000-40000)	690000 (530000-840000)
Italy	2005	Male	15 to 19	71% (58-82%)	29% (18-42%)	1.8 (1.2-2.6)	4% (2.1-6.1%)	9.8% (4.4-16%)	70 (31-110)	6200 (3400-9400)
Italy	2005	Male	20 to 24	83% (74-89%)	17% (11-26%)	3.2 (2.2-4.4)	7.8% (4.6-11%)	16% (8.3-24%)	180 (96-270)	15000 (9100-22000)
Italy	2005	Male	25 to 29	87% (80-91%)	13% (8.5-20%)	4.4 (3.2-5.8)	8.4% (5.5-11%)	19% (11-26%)	270 (170-390)	21000 (14000-29000)
Italy	2005	Male	30 to 34	89% (83-94%)	11% (6.4-17%)	4.7 (3.4-6.2)	7.2% (4.8-9.8%)	17% (11-23%)	300 (190-430)	22000 (15000-30000)
Italy	2005	Male	35 to 39	91% (86-94%)	9.3% (5.6-14%)	4 (2.9-5.4)	5.9% (3.9-8.3%)	13% (8-18%)	310 (200-440)	21000 (14000-29000)
Italy	2005	Male	40 to 44	91% (86-94%)	9.1% (5.6-14%)	4.3 (3.3-5.7)	6.3% (4.3-8.8%)	12% (8-17%)	430 (280-610)	25000 (17000-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	2005	Male	45 to 49	90% (85-94%)	9.9% (6.5-15%)	5.8 (4.3-7.4)	8.8% (6.1-12%)	15% (11-20%)	740 (510-950)	37000 (26000-47000)
Italy	2005	Male	50 to 54	88% (84-91%)	12% (8.6-16%)	6.7 (5.1-8.3)	10% (7.7-13%)	16% (12-20%)	1200 (860-1400)	49000 (37000-60000)
Italy	2005	Male	55 to 59	85% (82-88%)	15% (12-18%)	7 (5.5-8.7)	11% (8.4-13%)	15% (12-19%)	1900 (1400-2300)	68000 (53000-82000)
Italy	2005	Male	60 to 64	85% (82-89%)	15% (11-18%)	7.1 (5.5-8.9)	11% (8.3-13%)	14% (11-18%)	2300 (1800-2900)	72000 (56000-90000)
Italy	2005	Male	65 to 69	86% (83-90%)	14% (10-17%)	6.8 (5.3-8.5)	11% (8.1-13%)	13% (10-16%)	3400 (2600-4200)	88000 (68000-110000)
Italy	2005	Male	70 to 74	85% (81-88%)	15% (12-19%)	6.2 (4.8-7.8)	9.9% (6.9-13%)	12% (8.3-15%)	4400 (3000-5600)	89000 (63000-110000)
Italy	2005	Male	75 to 79	83% (76-88%)	17% (12-24%)	5.6 (4.3-7.1)	9.1% (5.9-12%)	10% (6.8-14%)	5100 (3300-6800)	80000 (52000-110000)
Italy	2005	Male	80 to 84	81% (72-87%)	19% (13-28%)	5.2 (4-6.7)	8.5% (5.3-12%)	9.6% (5.9-14%)	5200 (3200-7200)	59000 (38000-81000)
Italy	2005	Male	85 to 89	79% (71-86%)	21% (14-29%)	4.9 (3.6-6.5)	7.8% (4.3-11%)	8.7% (4.6-13%)	2700 (1400-4000)	22000 (12000-32000)
Italy	2005	Male	90 to 94	80% (71-87%)	20% (13-29%)	4.8 (3.3-6.5)	7% (3-11%)	7.8% (3.1-12%)	2000 (770-3100)	11000 (4600-17000)
Italy	2005	Male	95 plus	81% (68-89%)	19% (11-32%)	4.6 (2.8-6.7)	6.8% (1.7-12%)	7.4% (1.3-13%)	520 (93-930)	1900 (470-3300)
Italy	2010	Female	All Ages	54% (51-58%)	46% (42-49%)	2.1 (1.6-2.6)	3% (1.9-4.2%)	4.3% (1.9-7.1%)	13000 (6000-22000)	230000 (140000-330000)
Italy	2010	Female	15 to 19	58% (44-71%)	42% (29-56%)	1.4 (0.87-2.2)	1.2% (0.68-1.8%)	5.9% (2.8-9.5%)	13 (6.4-21)	1700 (970-2500)
Italy	2010	Female	20 to 24	69% (57-78%)	31% (22-43%)	1.9 (1.2-3)	1.6% (1-2.4%)	7.2% (3.8-11%)	20 (10-31)	2700 (1700-4000)
Italy	2010	Female	25 to 29	71% (61-79%)	29% (21-39%)	2.1 (1.4-3)	1.5% (0.96-2.1%)	7.1% (4.1-11%)	23 (13-34)	3000 (1900-4400)
Italy	2010	Female	30 to 34	71% (63-79%)	29% (21-37%)	2.3 (1.6-3.3)	1.6% (1-2.3%)	7.7% (4.9-11%)	44 (28-60)	4300 (2700-6000)
Italy	2010	Female	35 to 39	66% (58-75%)	34% (25-42%)	2.4 (1.6-3.5)	1.9% (1.3-2.8%)	8.5% (5.9-11%)	88 (61-120)	6700 (4500-9400)
Italy	2010	Female	40 to 44	62% (53-69%)	38% (31-47%)	2.5 (1.7-3.4)	2.5% (1.7-3.4%)	8.8% (6.3-11%)	160 (120-210)	10000 (6900-14000)
Italy	2010	Female	45 to 49	59% (52-65%)	41% (35-48%)	2.6 (1.8-3.6)	3.1% (2.1-4.4%)	8.9% (6.7-11%)	270 (200-350)	14000 (9700-19000)
Italy	2010	Female	50 to 54	57% (51-63%)	43% (37-49%)	2.8 (1.9-3.7)	3.7% (2.5-5.1%)	8.6% (6.4-11%)	370 (270-480)	16000 (11000-22000)
Italy	2010	Female	55 to 59	58% (53-63%)	42% (37-47%)	3.1 (2.2-4.2)	4.3% (2.8-6%)	8.7% (6.2-12%)	510 (370-690)	20000 (13000-27000)
Italy	2010	Female	60 to 64	55% (49-60%)	45% (40-51%)	2.9 (2.1-4)	4.1% (2.5-5.9%)	7.6% (5.1-10%)	700 (480-960)	23000 (14000-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	2010	Female	65 to 69	54% (48-60%)	46% (40-52%)	2.7 (2-3.5)	3.8% (2.3-5.4%)	6.4% (4.2-8.8%)	790 (530-1100)	21000 (13000-30000)
Italy	2010	Female	70 to 74	51% (45-57%)	49% (43-55%)	2.3 (1.6-3.2)	3.4% (1.8-5.3%)	5.3% (2.9-7.8%)	1100 (620-1700)	24000 (12000-37000)
Italy	2010	Female	75 to 79	44% (36-52%)	56% (48-64%)	1.9 (1.2-2.8)	2.9% (1.2-5%)	4.1% (1.8-6.7%)	1400 (640-2400)	23000 (9800-40000)
Italy	2010	Female	80 to 84	44% (34-55%)	56% (45-66%)	2 (1.3-2.8)	3% (1.2-5.3%)	3.9% (1.5-6.9%)	2200 (850-4000)	26000 (10000-46000)
Italy	2010	Female	85 to 89	44% (34-54%)	56% (46-66%)	2 (1.3-3)	3% (0.91-5.5%)	3.6% (1-6.8%)	2800 (760-5200)	23000 (6700-41000)
Italy	2010	Female	90 to 94	43% (32-54%)	57% (46-68%)	2 (1.2-3.1)	2.9% (0.42-5.9%)	3.4% (0.3-7.1%)	1500 (130-3200)	8400 (1200-17000)
Italy	2010	Female	95 plus	44% (30-60%)	56% (40-70%)	2.1 (1.1-3.6)	3.3% (0.13-7.6%)	3.8% (-0.026-9.1%)	1300 (-8.7-3100)	4600 (170-11000)
Italy	2010	Male	All Ages	80% (77-84%)	20% (16-23%)	4.2 (3.3-5.3)	8.1% (5.9-10%)	10% (6.7-14%)	30000 (19000-40000)	640000 (470000-810000)
Italy	2010	Male	15 to 19	69% (55-81%)	31% (19-45%)	1.7 (1.1-2.5)	3.4% (1.9-5.1%)	9% (4.2-14%)	53 (25-85)	5100 (2900-7600)
Italy	2010	Male	20 to 24	82% (73-89%)	18% (11-27%)	3 (2.1-4.4)	6.9% (4.3-9.9%)	15% (8-23%)	130 (69-200)	12000 (7400-17000)
Italy	2010	Male	25 to 29	86% (79-91%)	14% (9.3-21%)	4.2 (2.9-5.8)	7.4% (4.8-10%)	18% (11-26%)	180 (110-260)	15000 (9500-20000)
Italy	2010	Male	30 to 34	88% (81-93%)	12% (7.1-19%)	4.5 (3.2-6.1)	6.6% (4.3-9.2%)	16% (10-23%)	220 (140-310)	17000 (11000-23000)
Italy	2010	Male	35 to 39	89% (84-94%)	11% (6.2-16%)	3.8 (2.7-5.2)	5.5% (3.6-7.7%)	13% (7.7-18%)	250 (160-360)	18000 (12000-24000)
Italy	2010	Male	40 to 44	90% (85-94%)	9.9% (5.9-15%)	4.1 (2.9-5.5)	5.9% (3.9-8.3%)	12% (8-17%)	390 (250-540)	23000 (15000-32000)
Italy	2010	Male	45 to 49	90% (85-93%)	10% (6.6-15%)	5.5 (4-7.4)	8.2% (5.5-11%)	15% (10-20%)	750 (500-990)	38000 (26000-49000)
Italy	2010	Male	50 to 54	88% (84-91%)	12% (8.7-16%)	6.4 (4.7-8.4)	9.8% (7-13%)	16% (11-20%)	1100 (800-1400)	48000 (34000-60000)
Italy	2010	Male	55 to 59	85% (82-88%)	15% (12-18%)	6.7 (5.1-8.7)	10% (7.7-13%)	15% (11-18%)	1600 (1200-1900)	57000 (43000-71000)
Italy	2010	Male	60 to 64	84% (80-88%)	16% (12-20%)	6.7 (5-8.6)	10% (7.4-13%)	13% (10-17%)	2300 (1700-2900)	72000 (53000-91000)
Italy	2010	Male	65 to 69	86% (83-90%)	14% (10-17%)	6.5 (5-8.4)	9.8% (7.1-12%)	12% (9.1-16%)	2800 (2000-3500)	72000 (53000-91000)
Italy	2010	Male	70 to 74	85% (82-88%)	15% (12-18%)	6 (4.5-7.9)	9.2% (6.2-12%)	11% (7.5-15%)	3900 (2600-5100)	81000 (55000-110000)
Italy	2010	Male	75 to 79	82% (76-87%)	18% (13-24%)	5.4 (3.9-7.1)	8.3% (5.1-11%)	9.6% (5.9-13%)	4500 (2700-6200)	72000 (45000-98000)
Italy	2010	Male	80 to 84	82% (74-88%)	18% (12-26%)	5.1 (3.8-6.7)	7.9% (4.6-11%)	8.9% (5-13%)	5100 (2900-7500)	60000 (35000-86000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	2010	Male	85 to 89	81% (74-87%)	19% (13-26%)	4.9 (3.5-6.5)	7.4% (3.6-11%)	8.2% (3.8-13%)	4200 (1900-6400)	34000 (17000-51000)
Italy	2010	Male	90 to 94	81% (73-88%)	19% (12-27%)	4.7 (3.1-6.4)	6.6% (2.4-11%)	7.3% (2.3-12%)	1500 (480-2500)	8500 (3100-14000)
Italy	2010	Male	95 plus	81% (69-90%)	19% (9.9-31%)	4.5 (2.6-6.6)	6.4% (0.92-12%)	7% (0.56-13%)	720 (58-1400)	2700 (380-4800)
Italy	2016	Female	All Ages	59% (55-63%)	41% (37-45%)	2.1 (1.5-2.9)	2.8% (1.4-4.5%)	3.9% (1.2-7.2%)	13000 (4300-26000)	230000 (110000-370000)
Italy	2016	Female	15 to 19	63% (48-76%)	37% (24-52%)	1.5 (0.82-2.3)	1.1% (0.65-1.8%)	6% (2.8-10%)	12 (5.7-21)	1500 (880-2400)
Italy	2016	Female	20 to 24	72% (59-82%)	28% (18-41%)	1.9 (1.1-3)	1.5% (0.93-2.3%)	7.1% (3.5-11%)	19 (9.5-31)	2700 (1600-4000)
Italy	2016	Female	25 to 29	74% (62-83%)	26% (17-38%)	2.1 (1.3-3.3)	1.4% (0.88-2.1%)	7% (3.8-11%)	21 (11-32)	2700 (1600-4000)
Italy	2016	Female	30 to 34	74% (63-83%)	26% (17-37%)	2.3 (1.4-3.6)	1.5% (0.89-2.3%)	7.5% (4.4-11%)	34 (19-51)	3400 (1900-5000)
Italy	2016	Female	35 to 39	72% (61-81%)	28% (19-39%)	2.5 (1.5-3.8)	1.9% (1.1-2.8%)	8.5% (5.5-12%)	79 (49-110)	5900 (3600-8700)
Italy	2016	Female	40 to 44	69% (58-77%)	31% (23-42%)	2.6 (1.6-3.9)	2.4% (1.5-3.6%)	9.1% (6.1-12%)	160 (100-220)	9600 (6000-14000)
Italy	2016	Female	45 to 49	63% (52-72%)	37% (28-48%)	2.7 (1.6-4)	3% (1.8-4.4%)	9% (6-12%)	290 (190-410)	15000 (9100-22000)
Italy	2016	Female	50 to 54	60% (51-68%)	40% (32-49%)	2.8 (1.7-4.1)	3.5% (2.1-5.2%)	8.5% (5.8-12%)	390 (270-560)	17000 (11000-26000)
Italy	2016	Female	55 to 59	58% (50-66%)	42% (34-50%)	2.9 (1.8-4.3)	3.9% (2.3-6%)	8.2% (5.3-12%)	500 (320-730)	19000 (11000-29000)
Italy	2016	Female	60 to 64	57% (48-66%)	43% (34-52%)	2.9 (1.8-4.3)	3.9% (2.1-6.1%)	7.3% (4.6-11%)	650 (390-990)	21000 (11000-33000)
Italy	2016	Female	65 to 69	58% (50-66%)	42% (34-50%)	2.7 (1.8-4)	3.6% (1.9-5.8%)	6.2% (3.6-9.5%)	810 (460-1200)	22000 (11000-34000)
Italy	2016	Female	70 to 74	55% (47-64%)	45% (36-53%)	2.4 (1.5-3.5)	3.2% (1.3-5.7%)	5% (2.3-8.3%)	1100 (470-1800)	22000 (8900-39000)
Italy	2016	Female	75 to 79	52% (41-62%)	48% (38-59%)	2.1 (1.3-3.2)	2.8% (0.8-5.5%)	3.9% (1.2-7.5%)	1300 (410-2500)	22000 (6200-42000)
Italy	2016	Female	80 to 84	49% (37-60%)	51% (40-63%)	2.1 (1.2-3.1)	2.8% (0.63-5.5%)	3.6% (0.87-7.2%)	2100 (520-4200)	25000 (5500-49000)
Italy	2016	Female	85 to 89	46% (34-58%)	54% (42-66%)	2 (1.1-3.1)	2.6% (0.35-5.6%)	3.2% (0.38-6.9%)	2700 (310-5800)	22000 (2900-46000)
Italy	2016	Female	90 to 94	45% (33-57%)	55% (43-67%)	2 (1.1-3.1)	2.5% (-0.0012-5.8%)	3% (-0.15-7%)	2400 (-120-5700)	13000 (-6.3-31000)
Italy	2016	Female	95 plus	46% (31-62%)	54% (38-69%)	2 (0.95-3.6)	2.8% (-0.39-7.2%)	3.2% (-0.75-8.7%)	1000 (-230-2700)	3600 (-470-9100)
Italy	2016	Male	All Ages	80% (76-83%)	20% (17-24%)	4 (2.7-5.4)	7.2% (4.4-9.9%)	8.7% (4.5-13%)	28000 (14000-41000)	580000 (360000-800000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Italy	2016	Male	15 to 19	69% (55-82%)	31% (18-45%)	1.6 (0.95-2.6)	3.1% (1.8-4.8%)	8.6% (4-14%)	44 (20-72)	4300 (2500-6400)
Italy	2016	Male	20 to 24	80% (68-88%)	20% (12-32%)	2.7 (1.7-4.1)	6.3% (3.8-9.2%)	14% (7-21%)	120 (59-180)	11000 (6700-16000)
Italy	2016	Male	25 to 29	84% (74-90%)	16% (9.8-26%)	3.8 (2.4-5.7)	6.9% (4.2-9.8%)	17% (9.2-25%)	150 (81-230)	13000 (7800-18000)
Italy	2016	Male	30 to 34	87% (78-93%)	13% (7.3-22%)	4.1 (2.6-6.1)	6.1% (3.8-8.7%)	15% (8.8-23%)	160 (91-240)	13000 (7900-18000)
Italy	2016	Male	35 to 39	89% (82-94%)	11% (6.2-18%)	3.5 (2.3-5.1)	5% (3-7.3%)	12% (6.6-18%)	200 (110-310)	15000 (9000-21000)
Italy	2016	Male	40 to 44	89% (82-94%)	11% (6.2-18%)	3.8 (2.5-5.5)	5.3% (3.3-7.9%)	11% (6.9-17%)	330 (200-500)	20000 (13000-30000)
Italy	2016	Male	45 to 49	88% (81-93%)	12% (7.1-19%)	5.1 (3.3-7.4)	7.4% (4.5-10%)	14% (8.7-19%)	720 (440-1000)	37000 (23000-51000)
Italy	2016	Male	50 to 54	87% (82-91%)	13% (9.1-18%)	5.9 (3.9-8.3)	8.7% (5.7-12%)	15% (9.7-20%)	1100 (680-1400)	46000 (30000-62000)
Italy	2016	Male	55 to 59	84% (80-88%)	16% (12-20%)	6.2 (4.1-8.8)	9.2% (6.1-12%)	14% (9.2-18%)	1400 (930-1900)	53000 (35000-70000)
Italy	2016	Male	60 to 64	84% (79-88%)	16% (12-21%)	6.2 (4.2-8.8)	9.1% (5.9-12%)	12% (8.1-16%)	1900 (1300-2600)	62000 (40000-82000)
Italy	2016	Male	65 to 69	86% (81-89%)	14% (11-19%)	6 (4.1-8.4)	8.8% (5.6-12%)	11% (7.3-15%)	2600 (1600-3500)	68000 (44000-93000)
Italy	2016	Male	70 to 74	85% (81-89%)	15% (11-19%)	5.6 (3.7-7.9)	8.2% (4.8-12%)	10% (5.8-14%)	3400 (2000-4800)	72000 (42000-100000)
Italy	2016	Male	75 to 79	83% (76-88%)	17% (12-24%)	5 (3.3-7.2)	7.3% (3.7-11%)	8.5% (4.3-13%)	3900 (2000-5900)	64000 (32000-96000)
Italy	2016	Male	80 to 84	81% (71-88%)	19% (12-29%)	4.7 (3.1-6.7)	6.7% (3-11%)	7.6% (3.3-12%)	4700 (2000-7500)	56000 (24000-87000)
Italy	2016	Male	85 to 89	80% (70-86%)	20% (14-30%)	4.4 (2.8-6.5)	6.1% (2.1-10%)	6.8% (2.1-12%)	4100 (1200-7000)	34000 (12000-57000)
Italy	2016	Male	90 to 94	80% (72-87%)	20% (13-28%)	4.3 (2.6-6.4)	5.5% (1.1-10%)	6.1% (0.83-11%)	2300 (310-4400)	13000 (2500-24000)
Italy	2016	Male	95 plus	81% (69-89%)	19% (11-31%)	4.1 (2.3-6.5)	5.4% (0.18-11%)	5.8% (-0.3-12%)	520 (-28-1100)	1900 (64-4000)
Luxembourg	1990	Female	All Ages	75% (67-83%)	25% (17-33%)	2.1 (2-2.3)	3.9% (2.4-5.5%)	4% (0.9-7.3%)	77 (17-140)	2300 (1400-3200)
Luxembourg	1990	Female	15 to 19	72% (53-86%)	28% (14-47%)	1.6 (1-2.3)	2.6% (1.4-4%)	8.5% (4-14%)	0.31 (0.15-0.51)	32 (18-48)
Luxembourg	1990	Female	20 to 24	77% (59-89%)	23% (11-41%)	2 (1.3-2.9)	3.5% (2.2-5%)	9.3% (4.9-14%)	0.61 (0.31-0.95)	66 (42-93)
Luxembourg	1990	Female	25 to 29	79% (64-90%)	21% (9.6-36%)	2.2 (1.5-2.9)	3.2% (2.1-4.4%)	9.7% (5.6-14%)	0.79 (0.45-1.1)	77 (51-110)
Luxembourg	1990	Female	30 to 34	81% (65-91%)	19% (8.6-35%)	2.3 (1.6-3.1)	3.4% (2.3-4.5%)	11% (7.7-15%)	1.1 (0.7-1.4)	85 (59-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	1990	Female	35 to 39	82% (67-92%)	18% (8.1-33%)	2.4 (1.6-3.5)	4.3% (3-5.8%)	13% (9.2-17%)	1.8 (1.2-2.3)	110 (80-150)
Luxembourg	1990	Female	40 to 44	82% (66-92%)	18% (7.7-34%)	2.6 (1.8-3.5)	5.6% (3.9-7.3%)	14% (10-17%)	3 (2.2-3.8)	160 (120-200)
Luxembourg	1990	Female	45 to 49	82% (64-92%)	18% (7.8-36%)	2.9 (2-3.9)	6.6% (4.8-8.7%)	15% (11-18%)	4.2 (3.1-5.3)	190 (140-240)
Luxembourg	1990	Female	50 to 54	82% (65-92%)	18% (8-35%)	3.2 (2.3-4.2)	7.3% (5.4-9.4%)	14% (11-17%)	6.1 (4.7-7.7)	240 (180-320)
Luxembourg	1990	Female	55 to 59	81% (65-92%)	19% (7.9-35%)	3.6 (2.6-4.8)	7.7% (5.4-10%)	13% (9.6-17%)	8.4 (6.1-11)	290 (210-390)
Luxembourg	1990	Female	60 to 64	80% (61-92%)	20% (8.2-39%)	3.5 (2.5-4.9)	6.8% (4.2-9.8%)	10% (6.7-14%)	11 (7-15)	320 (200-460)
Luxembourg	1990	Female	65 to 69	78% (59-91%)	22% (9.4-41%)	3 (2.2-4)	5% (2.5-7.9%)	7.2% (3.6-11%)	11 (5.6-17)	270 (130-420)
Luxembourg	1990	Female	70 to 74	75% (57-88%)	25% (12-43%)	2.5 (1.7-3.5)	3.4% (0.27-6.8%)	4.7% (0.49-9.1%)	8.3 (0.85-16)	160 (13-310)
Luxembourg	1990	Female	75 to 79	73% (53-87%)	27% (13-47%)	2.2 (1.4-3.2)	1.8% (-1.7-5.7%)	2.2% (-2.2-7%)	7.3 (-7.4-23)	110 (-110-350)
Luxembourg	1990	Female	80 to 84	70% (49-85%)	30% (15-51%)	2.1 (1.4-3.1)	1.5% (-1.8-5.1%)	1.7% (-2.5-6.2%)	7.1 (-10-25)	83 (-99-280)
Luxembourg	1990	Female	85 to 89	68% (49-82%)	32% (18-51%)	2.1 (1.3-3)	1.2% (-1.8-4.6%)	1.2% (-2.4-5.3%)	4.1 (-8-17)	36 (-55-140)
Luxembourg	1990	Female	90 to 94	67% (48-83%)	33% (17-52%)	2.1 (1.2-3.1)	1.1% (-2.1-4.6%)	1% (-2.9-5.3%)	1.6 (-4.5-8.6)	10 (-21-45)
Luxembourg	1990	Female	95 plus	67% (43-85%)	33% (15-57%)	2 (1.1-3.4)	1.1% (-2.6-5.5%)	0.98% (-3.5-6.3%)	0.34 (-1.2-2.2)	1.6 (-3.8-7.9)
Luxembourg	1990	Male	All Ages	82% (75-89%)	18% (11-25%)	4.6 (4.3-5)	13% (11-15%)	15% (11-18%)	280 (220-350)	8100 (6900-9500)
Luxembourg	1990	Male	15 to 19	75% (55-89%)	25% (11-45%)	2.9 (1.9-4.1)	8.4% (4.9-12%)	14% (7.6-22%)	1.8 (0.97-2.7)	150 (83-220)
Luxembourg	1990	Male	20 to 24	80% (63-91%)	20% (9.1-37%)	4.4 (3.1-6)	14% (9.8-19%)	21% (13-30%)	4.3 (2.7-6)	350 (240-480)
Luxembourg	1990	Male	25 to 29	83% (70-93%)	17% (7.4-30%)	5 (3.7-6.6)	14% (10-18%)	22% (15-30%)	5.3 (3.6-7.1)	410 (300-530)
Luxembourg	1990	Male	30 to 34	86% (72-94%)	14% (5.8-28%)	5.4 (3.9-7)	13% (10-17%)	23% (16-29%)	5.6 (4-7.2)	400 (300-500)
Luxembourg	1990	Male	35 to 39	88% (76-95%)	12% (4.5-24%)	5.4 (3.9-7.2)	14% (11-18%)	23% (17-29%)	7.5 (5.4-9.6)	460 (350-580)
Luxembourg	1990	Male	40 to 44	89% (77-96%)	11% (3.9-23%)	5.8 (4.3-7.4)	14% (11-18%)	23% (17-28%)	9.5 (7-12)	510 (390-630)
Luxembourg	1990	Male	45 to 49	90% (74-96%)	10% (3.7-26%)	6.6 (4.8-8.7)	17% (12-20%)	23% (18-28%)	14 (11-18)	660 (510-800)
Luxembourg	1990	Male	50 to 54	90% (77-97%)	9.9% (3.4-23%)	7.1 (5.2-9.2)	17% (13-20%)	22% (18-27%)	20 (16-24)	800 (630-950)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	1990	Male	55 to 59	90% (79-97%)	10% (3.4-21%)	7.4 (5.6-9.4)	16% (13-20%)	21% (16-25%)	27 (21-33)	940 (730-1100)
Luxembourg	1990	Male	60 to 64	89% (76-97%)	11% (3.5-24%)	7.3 (5.4-9.5)	15% (11-18%)	18% (13-22%)	35 (24-44)	1000 (730-1300)
Luxembourg	1990	Male	65 to 69	89% (76-96%)	11% (3.8-24%)	6.6 (5.1-8.3)	13% (9.5-17%)	15% (11-19%)	32 (22-40)	770 (550-960)
Luxembourg	1990	Male	70 to 74	88% (76-95%)	12% (4.7-24%)	5.9 (4.2-7.7)	11% (6.5-16%)	13% (7.3-18%)	30 (17-42)	590 (340-820)
Luxembourg	1990	Male	75 to 79	87% (74-95%)	13% (5.1-26%)	5.4 (3.6-7.4)	10% (4.8-15%)	11% (5.4-17%)	36 (17-53)	550 (260-800)
Luxembourg	1990	Male	80 to 84	86% (71-94%)	14% (5.9-29%)	5.2 (3.7-6.9)	10% (5.4-14%)	11% (5.7-16%)	32 (17-47)	350 (190-520)
Luxembourg	1990	Male	85 to 89	84% (71-93%)	16% (6.7-29%)	4.9 (3.5-6.7)	9.2% (4.7-14%)	10% (4.9-16%)	18 (8.5-28)	140 (70-220)
Luxembourg	1990	Male	90 to 94	84% (71-94%)	16% (6.2-29%)	4.8 (3.2-6.9)	8.3% (3.2-14%)	9.1% (3-15%)	5.9 (2-9.9)	33 (12-53)
Luxembourg	1990	Male	95 plus	84% (67-95%)	16% (4.9-33%)	4.6 (2.6-7.3)	7.7% (1.4-14%)	8.4% (1.2-16%)	0.8 (0.12-1.5)	3 (0.56-5.3)
Luxembourg	1995	Female	All Ages	75% (67-82%)	25% (18-33%)	2 (1.8-2.3)	3.6% (2.3-5%)	3.5% (0.76-6.6%)	65 (14-120)	2000 (1300-2800)
Luxembourg	1995	Female	15 to 19	72% (54-87%)	28% (13-46%)	1.5 (1-2.2)	2.8% (1.5-4.3%)	7.7% (3.7-12%)	0.43 (0.21-0.68)	39 (21-60)
Luxembourg	1995	Female	20 to 24	77% (60-89%)	23% (11-40%)	1.9 (1.3-2.8)	3.3% (2.2-4.7%)	8.7% (4.6-13%)	0.54 (0.28-0.85)	59 (37-83)
Luxembourg	1995	Female	25 to 29	79% (64-90%)	21% (9.7-36%)	2.1 (1.5-2.8)	2.8% (1.9-3.8%)	9.1% (5.5-13%)	0.58 (0.35-0.85)	64 (44-90)
Luxembourg	1995	Female	30 to 34	81% (66-91%)	19% (8.5-34%)	2.2 (1.5-3)	3.6% (2.5-4.8%)	11% (7.3-15%)	1.4 (0.94-1.9)	110 (73-140)
Luxembourg	1995	Female	35 to 39	82% (66-92%)	18% (8.2-34%)	2.3 (1.6-3.2)	3.8% (2.7-5.1%)	13% (9.1-16%)	1.7 (1.2-2.2)	110 (80-140)
Luxembourg	1995	Female	40 to 44	82% (66-92%)	18% (7.7-34%)	2.5 (1.7-3.4)	5% (3.5-6.6%)	14% (10-17%)	2.8 (2.1-3.6)	160 (120-200)
Luxembourg	1995	Female	45 to 49	82% (65-92%)	18% (7.9-35%)	2.7 (1.9-3.7)	6.1% (4.4-8%)	14% (11-17%)	4.3 (3.3-5.3)	200 (150-250)
Luxembourg	1995	Female	50 to 54	82% (65-92%)	18% (8-35%)	3.1 (2.2-4.1)	6.6% (4.6-8.4%)	13% (10-16%)	5.1 (3.9-6.3)	210 (150-260)
Luxembourg	1995	Female	55 to 59	81% (65-92%)	19% (8-35%)	3.5 (2.5-4.6)	6.8% (4.7-8.9%)	13% (9.5-16%)	6.5 (4.9-8.2)	230 (170-300)
Luxembourg	1995	Female	60 to 64	80% (61-92%)	20% (8.1-39%)	3.4 (2.4-4.7)	6.2% (3.7-8.8%)	10% (6.6-14%)	8.7 (5.7-12)	260 (160-370)
Luxembourg	1995	Female	65 to 69	78% (59-90%)	22% (9.6-41%)	2.9 (2.1-3.9)	4.6% (2.4-7.1%)	7% (3.9-10%)	9.2 (5-14)	230 (120-350)
Luxembourg	1995	Female	70 to 74	75% (58-88%)	25% (12-42%)	2.4 (1.6-3.4)	3.1% (0.4-6%)	4.4% (0.74-8.4%)	8.6 (1.4-16)	170 (21-320)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	1995	Female	75 to 79	73% (53-87%)	27% (13-47%)	2.1 (1.3-3.1)	1.7% (-1.2-4.9%)	2.1% (-1.5-6.2%)	4.8 (-3.5-14)	76 (-55-220)
Luxembourg	1995	Female	80 to 84	70% (49-85%)	30% (15-51%)	2.1 (1.3-3)	1.2% (-1.7-4.6%)	1.4% (-2.3-5.5%)	5.1 (-8.5-20)	62 (-87-230)
Luxembourg	1995	Female	85 to 89	68% (49-83%)	32% (17-51%)	2 (1.3-2.9)	0.95% (-1.7-3.9%)	0.93% (-2.2-4.4%)	3.5 (-8.4-17)	33 (-59-140)
Luxembourg	1995	Female	90 to 94	67% (48-83%)	33% (17-52%)	2 (1.2-3)	0.81% (-2.1-4%)	0.71% (-2.7-4.5%)	1.3 (-5.1-8.6)	9.6 (-25-48)
Luxembourg	1995	Female	95 plus	67% (44-85%)	33% (15-56%)	2 (1-3.3)	0.86% (-2.6-5%)	0.71% (-3.3-5.6%)	0.39 (-1.9-3.1)	2 (-5.8-11)
Luxembourg	1995	Male	All Ages	82% (74-89%)	18% (11-26%)	4.4 (4-4.9)	12% (9.8-14%)	14% (10-17%)	260 (190-320)	7400 (6200-8700)
Luxembourg	1995	Male	15 to 19	75% (55-89%)	25% (11-45%)	2.8 (1.8-3.9)	6.7% (4.2-9.6%)	14% (7.6-21%)	1 (0.56-1.6)	95 (58-140)
Luxembourg	1995	Male	20 to 24	80% (62-91%)	20% (8.9-38%)	4.2 (2.9-5.8)	13% (8.9-17%)	20% (12-28%)	2.9 (1.8-4.2)	260 (180-350)
Luxembourg	1995	Male	25 to 29	83% (69-93%)	17% (7-31%)	4.9 (3.5-6.4)	12% (9.2-16%)	20% (14-27%)	4.4 (3-5.9)	360 (260-470)
Luxembourg	1995	Male	30 to 34	86% (72-95%)	14% (5.1-28%)	5.2 (3.8-6.8)	12% (9.1-15%)	21% (15-27%)	5.2 (3.7-6.9)	390 (290-490)
Luxembourg	1995	Male	35 to 39	88% (74-95%)	12% (4.6-26%)	5.2 (3.7-6.9)	13% (9.7-17%)	22% (16-28%)	7.5 (5.4-9.5)	460 (350-580)
Luxembourg	1995	Male	40 to 44	89% (76-96%)	11% (3.8-24%)	5.6 (4.1-7.3)	14% (10-17%)	22% (17-28%)	9.2 (6.8-12)	500 (380-620)
Luxembourg	1995	Male	45 to 49	90% (75-96%)	10% (3.7-25%)	6.3 (4.6-8.4)	15% (11-19%)	23% (17-28%)	13 (9.6-16)	610 (460-740)
Luxembourg	1995	Male	50 to 54	90% (77-96%)	10% (3.7-23%)	6.8 (4.9-8.8)	16% (12-19%)	22% (17-26%)	18 (14-21)	720 (560-860)
Luxembourg	1995	Male	55 to 59	90% (78-97%)	10% (3.5-22%)	7.1 (5.4-9.1)	16% (12-19%)	20% (16-24%)	24 (19-29)	840 (660-1000)
Luxembourg	1995	Male	60 to 64	90% (76-96%)	10% (3.6-24%)	6.9 (5.1-9.1)	14% (10-17%)	17% (12-22%)	30 (21-37)	880 (640-1100)
Luxembourg	1995	Male	65 to 69	89% (76-96%)	11% (3.8-24%)	6.3 (4.9-8)	12% (8.7-15%)	15% (10-18%)	36 (25-46)	890 (640-1100)
Luxembourg	1995	Male	70 to 74	88% (77-95%)	12% (4.8-23%)	5.7 (4-7.4)	10% (6.1-14%)	12% (6.8-17%)	27 (16-38)	540 (320-750)
Luxembourg	1995	Male	75 to 79	87% (74-95%)	13% (4.9-26%)	5.2 (3.5-7.2)	9.3% (4.3-14%)	10% (4.5-15%)	25 (11-37)	380 (170-560)
Luxembourg	1995	Male	80 to 84	86% (71-94%)	14% (5.8-29%)	5 (3.5-6.6)	9% (4.4-13%)	9.9% (4.5-15%)	28 (13-42)	320 (160-470)
Luxembourg	1995	Male	85 to 89	84% (70-93%)	16% (6.6-30%)	4.7 (3.3-6.5)	8.3% (4.1-13%)	9.1% (4.4-15%)	18 (8.6-29)	150 (73-230)
Luxembourg	1995	Male	90 to 94	84% (69-94%)	16% (5.8-31%)	4.6 (3-6.6)	7.6% (2.7-13%)	8.3% (2.7-14%)	6.5 (2-11)	36 (13-60)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	1995	Male	95 plus	84% (66-95%)	16% (5-34%)	4.5 (2.5-7)	7.1% (1.4-13%)	7.8% (1-15%)	1.2 (0.16-2.3)	4.6 (0.86-8.5)
Luxembourg	2000	Female	All Ages	75% (67-82%)	25% (18-33%)	2.3 (2-2.7)	4.1% (2.8-5.7%)	4.8% (2-8%)	84 (34-140)	2300 (1600-3100)
Luxembourg	2000	Female	15 to 19	69% (50-85%)	31% (15-50%)	1.7 (1-2.4)	2% (1.2-2.9%)	8% (4-13%)	0.21 (0.1-0.33)	25 (15-37)
Luxembourg	2000	Female	20 to 24	74% (56-88%)	26% (12-44%)	2.1 (1.4-3.2)	2.7% (1.9-3.7%)	9.5% (5.3-15%)	0.28 (0.15-0.42)	42 (27-59)
Luxembourg	2000	Female	25 to 29	78% (64-90%)	22% (9.9-36%)	2.4 (1.6-3.3)	2.9% (2-3.9%)	10% (6.3-14%)	0.57 (0.36-0.81)	63 (44-85)
Luxembourg	2000	Female	30 to 34	81% (66-92%)	19% (8.5-34%)	2.5 (1.8-3.5)	3.3% (2.4-4.5%)	12% (8.5-16%)	1.2 (0.82-1.6)	96 (70-130)
Luxembourg	2000	Female	35 to 39	83% (69-92%)	17% (7.7-31%)	2.7 (1.9-3.8)	3.5% (2.5-4.6%)	14% (10-17%)	1.5 (1.1-1.8)	110 (78-140)
Luxembourg	2000	Female	40 to 44	83% (67-92%)	17% (7.5-33%)	2.9 (2-4)	5.3% (3.9-7%)	15% (12-18%)	3.3 (2.6-4.1)	180 (140-230)
Luxembourg	2000	Female	45 to 49	83% (66-92%)	17% (7.6-34%)	3.2 (2.2-4.5)	6.2% (4.5-8.1%)	15% (12-19%)	4.4 (3.4-5.5)	210 (160-270)
Luxembourg	2000	Female	50 to 54	82% (66-92%)	18% (7.7-34%)	3.6 (2.5-4.8)	7% (5-9%)	15% (11-18%)	5.7 (4.3-7.1)	240 (180-300)
Luxembourg	2000	Female	55 to 59	82% (66-92%)	18% (7.7-34%)	4 (2.8-5.6)	7.9% (5.5-10%)	14% (11-18%)	7.5 (5.7-9.5)	270 (190-350)
Luxembourg	2000	Female	60 to 64	81% (63-92%)	19% (7.9-37%)	4 (2.7-5.7)	6.8% (4.4-9.7%)	12% (8.1-16%)	8 (5.4-11)	250 (160-350)
Luxembourg	2000	Female	65 to 69	78% (60-91%)	22% (9.3-40%)	3.4 (2.3-4.7)	5.3% (3.2-8.1%)	8.5% (5.4-12%)	8.8 (5.6-13)	230 (130-340)
Luxembourg	2000	Female	70 to 74	75% (58-88%)	25% (12-42%)	2.7 (1.8-3.9)	4% (1.3-7.1%)	5.8% (2.1-10%)	10 (3.7-18)	210 (68-360)
Luxembourg	2000	Female	75 to 79	72% (53-87%)	28% (13-47%)	2.4 (1.4-3.6)	2.8% (-0.14-6.1%)	3.7% (-0.086-8.1%)	9.5 (-0.21-20)	150 (-7.7-320)
Luxembourg	2000	Female	80 to 84	70% (49-85%)	30% (15-51%)	2.3 (1.5-3.4)	2.4% (-0.54-5.8%)	2.9% (-0.83-7%)	8.2 (-2.3-20)	96 (-22-230)
Luxembourg	2000	Female	85 to 89	67% (48-82%)	33% (18-52%)	2.2 (1.4-3.3)	2.1% (-0.69-5.3%)	2.3% (-1-6.2%)	8.3 (-3.6-23)	71 (-23-180)
Luxembourg	2000	Female	90 to 94	66% (47-83%)	34% (17-53%)	2.2 (1.3-3.4)	1.8% (-1.1-5.5%)	1.9% (-1.6-6.2%)	4.6 (-3.8-15)	28 (-16-82)
Luxembourg	2000	Female	95 plus	66% (42-85%)	34% (15-58%)	2.2 (1.1-3.8)	1.9% (-1.7-6.5%)	1.9% (-2.4-7.5%)	1.5 (-1.8-5.6)	5.9 (-5.3-21)
Luxembourg	2000	Male	All Ages	83% (75-89%)	17% (11-25%)	5.1 (4.4-5.9)	13% (11-15%)	15% (12-19%)	280 (210-340)	7800 (6600-9100)
Luxembourg	2000	Male	15 to 19	74% (54-89%)	26% (11-46%)	3.1 (2-4.5)	6.4% (4.1-9.1%)	15% (8.5-22%)	0.94 (0.54-1.4)	90 (58-130)
Luxembourg	2000	Male	20 to 24	80% (63-91%)	20% (9.2-37%)	4.8 (3.3-6.8)	13% (9.5-18%)	22% (14-31%)	2.9 (1.9-4.1)	250 (180-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	2000	Male	25 to 29	84% (72-93%)	16% (7.2-28%)	5.6 (3.8-7.5)	14% (10-18%)	23% (16-31%)	4.5 (3-6.1)	360 (260-470)
Luxembourg	2000	Male	30 to 34	87% (75-95%)	13% (5.4-25%)	6 (4.2-8)	13% (9.5-16%)	24% (17-31%)	5.2 (3.8-6.8)	390 (290-490)
Luxembourg	2000	Male	35 to 39	89% (78-96%)	11% (4.3-22%)	6 (4.4-8.1)	13% (10-16%)	25% (19-31%)	7 (5.3-8.8)	450 (350-570)
Luxembourg	2000	Male	40 to 44	90% (78-96%)	9.9% (3.6-22%)	6.4 (4.7-8.6)	14% (11-18%)	25% (20-31%)	9.7 (7.7-12)	540 (430-650)
Luxembourg	2000	Male	45 to 49	91% (77-97%)	9.2% (3.4-23%)	7.3 (5.1-9.9)	16% (13-20%)	26% (20-30%)	14 (11-17)	690 (540-820)
Luxembourg	2000	Male	50 to 54	91% (79-97%)	9.1% (3.2-21%)	7.9 (5.6-10)	16% (13-20%)	24% (20-29%)	18 (14-21)	750 (600-890)
Luxembourg	2000	Male	55 to 59	91% (80-97%)	9.4% (3.3-20%)	8.2 (6-11)	16% (13-20%)	22% (18-26%)	23 (18-27)	800 (660-950)
Luxembourg	2000	Male	60 to 64	90% (78-97%)	9.9% (3.3-22%)	7.9 (5.8-11)	15% (12-19%)	19% (15-24%)	28 (22-34)	840 (650-1000)
Luxembourg	2000	Male	65 to 69	89% (78-96%)	11% (3.6-22%)	7.3 (5.5-9.4)	13% (10-17%)	16% (13-20%)	33 (26-41)	830 (650-1000)
Luxembourg	2000	Male	70 to 74	89% (77-95%)	11% (4.6-23%)	6.5 (4.5-8.8)	12% (7.6-16%)	14% (8.9-19%)	39 (25-51)	770 (500-1000)
Luxembourg	2000	Male	75 to 79	87% (75-95%)	13% (5-25%)	6 (3.9-8.5)	11% (5.6-15%)	12% (6.3-17%)	30 (16-42)	470 (250-650)
Luxembourg	2000	Male	80 to 84	86% (71-94%)	14% (6-29%)	5.6 (3.9-7.7)	11% (6.2-15%)	12% (6.8-17%)	26 (15-37)	290 (170-410)
Luxembourg	2000	Male	85 to 89	84% (71-93%)	16% (6.5-29%)	5.4 (3.7-7.6)	9.9% (5.3-15%)	11% (5.7-16%)	23 (12-34)	180 (100-270)
Luxembourg	2000	Male	90 to 94	84% (71-94%)	16% (6.2-29%)	5.2 (3.3-7.6)	9.1% (3.9-14%)	10% (4.1-16%)	9.5 (3.9-15)	52 (22-81)
Luxembourg	2000	Male	95 plus	84% (67-95%)	16% (5.1-33%)	5.1 (2.8-8.2)	8.5% (2.5-14%)	9.4% (2.3-16%)	1.9 (0.48-3.3)	7.1 (2.1-12)
Luxembourg	2005	Female	All Ages	76% (68-83%)	24% (17-32%)	3 (2.5-3.5)	5.1% (3.7-6.7%)	7% (4-11%)	120 (70-190)	2900 (2100-3800)
Luxembourg	2005	Female	15 to 19	69% (50-85%)	31% (15-50%)	2.1 (1.3-3.1)	2.2% (1.4-3.3%)	9.1% (4.8-14%)	0.27 (0.14-0.42)	31 (19-46)
Luxembourg	2005	Female	20 to 24	75% (56-88%)	25% (12-44%)	2.7 (1.7-4.1)	3.1% (2.1-4.3%)	11% (6.4-16%)	0.37 (0.22-0.56)	51 (34-71)
Luxembourg	2005	Female	25 to 29	79% (65-90%)	21% (9.8-35%)	3 (2-4.3)	3.2% (2.2-4.3%)	12% (7.6-16%)	0.63 (0.41-0.87)	67 (47-90)
Luxembourg	2005	Female	30 to 34	81% (68-92%)	19% (8.4-32%)	3.3 (2.2-4.6)	3.2% (2.3-4.3%)	14% (10-18%)	0.95 (0.7-1.2)	84 (62-110)
Luxembourg	2005	Female	35 to 39	83% (70-93%)	17% (7.2-30%)	3.4 (2.3-4.8)	4.1% (3.1-5.4%)	16% (12-19%)	1.9 (1.5-2.4)	130 (100-170)
Luxembourg	2005	Female	40 to 44	83% (69-93%)	17% (7.5-31%)	3.7 (2.6-5)	4.8% (3.6-6.3%)	17% (13-20%)	2.9 (2.3-3.5)	170 (130-210)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	2005	Female	45 to 49	83% (67-92%)	17% (7.9-33%)	4 (2.7-5.6)	6.7% (4.9-8.8%)	17% (14-21%)	5.1 (4-6.2)	250 (190-320)
Luxembourg	2005	Female	50 to 54	83% (68-92%)	17% (8-32%)	4.5 (3.2-6.1)	7.8% (5.8-10%)	17% (13-21%)	6.8 (5.2-8.4)	290 (220-360)
Luxembourg	2005	Female	55 to 59	82% (67-92%)	18% (7.7-33%)	5.1 (3.6-7.1)	8.4% (6.1-11%)	16% (13-20%)	8.2 (6.4-10)	310 (230-400)
Luxembourg	2005	Female	60 to 64	81% (65-92%)	19% (7.8-35%)	5 (3.4-7.1)	8.4% (5.8-11%)	14% (10-19%)	9.5 (6.9-13)	300 (210-410)
Luxembourg	2005	Female	65 to 69	79% (62-91%)	21% (8.9-38%)	4.3 (3-5.8)	6.9% (4.5-9.7%)	11% (7.4-15%)	10 (7-14)	280 (180-390)
Luxembourg	2005	Female	70 to 74	75% (57-88%)	25% (12-43%)	3.4 (2.2-4.8)	5.5% (2.8-8.8%)	8.3% (4.4-13%)	12 (6.4-19)	260 (130-410)
Luxembourg	2005	Female	75 to 79	72% (52-87%)	28% (13-48%)	3 (1.8-4.4)	4.5% (1.3-8.2%)	6% (1.9-11%)	15 (4.7-28)	240 (72-460)
Luxembourg	2005	Female	80 to 84	70% (49-85%)	30% (15-51%)	2.9 (1.8-4.2)	4.3% (1.4-7.7%)	5.3% (1.6-9.7%)	18 (5.6-33)	210 (67-390)
Luxembourg	2005	Female	85 to 89	68% (50-82%)	32% (18-50%)	2.9 (1.8-4.2)	4% (0.97-7.9%)	4.7% (0.98-9.4%)	14 (3-29)	120 (30-230)
Luxembourg	2005	Female	90 to 94	68% (49-84%)	32% (16-51%)	2.9 (1.7-4.3)	3.8% (0.39-8.2%)	4.2% (0.071-9.7%)	11 (0.19-25)	64 (6.5-140)
Luxembourg	2005	Female	95 plus	67% (45-86%)	33% (14-55%)	2.9 (1.5-4.8)	3.8% (-0.49-9.6%)	4.3% (-0.78-11%)	4.6 (-0.84-12)	17 (-2.3-43)
Luxembourg	2005	Male	All Ages	84% (77-90%)	16% (10-23%)	6.5 (5.5-7.6)	14% (12-16%)	18% (14-21%)	300 (240-360)	8300 (7100-9600)
Luxembourg	2005	Male	15 to 19	75% (55-89%)	25% (11-45%)	3.9 (2.5-5.8)	6.5% (4.5-9.2%)	17% (11-26%)	0.92 (0.57-1.3)	92 (63-130)
Luxembourg	2005	Male	20 to 24	81% (64-91%)	19% (8.7-36%)	6.1 (4.1-8.7)	14% (11-19%)	25% (17-35%)	3.2 (2.1-4.3)	280 (210-370)
Luxembourg	2005	Male	25 to 29	85% (74-94%)	15% (6.2-26%)	7.1 (5-9.7)	13% (10-17%)	27% (20-35%)	3.3 (2.4-4.4)	290 (220-360)
Luxembourg	2005	Male	30 to 34	88% (77-95%)	12% (4.6-23%)	7.6 (5.4-10)	13% (9.9-16%)	27% (21-34%)	4.4 (3.3-5.5)	340 (270-410)
Luxembourg	2005	Male	35 to 39	90% (79-96%)	10% (3.9-21%)	7.6 (5.5-10)	14% (11-17%)	28% (22-34%)	7.1 (5.5-8.6)	460 (370-560)
Luxembourg	2005	Male	40 to 44	91% (81-96%)	9.4% (3.6-19%)	8.1 (5.9-11)	15% (12-18%)	29% (24-34%)	10 (8.4-12)	590 (480-690)
Luxembourg	2005	Male	45 to 49	91% (80-97%)	8.8% (3.3-20%)	9.2 (6.4-12)	17% (14-20%)	29% (24-33%)	15 (13-18)	730 (610-850)
Luxembourg	2005	Male	50 to 54	92% (81-97%)	8.5% (3.1-19%)	10 (7.1-13)	18% (15-21%)	27% (23-31%)	21 (17-24)	860 (730-1000)
Luxembourg	2005	Male	55 to 59	91% (82-97%)	8.7% (3.1-18%)	10 (7.7-14)	17% (14-20%)	24% (20-28%)	24 (20-28)	870 (730-1000)
Luxembourg	2005	Male	60 to 64	91% (80-97%)	9.1% (3.2-20%)	10 (7.2-14)	16% (13-19%)	21% (17-25%)	27 (22-32)	830 (660-990)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	2005	Male	65 to 69	90% (80-97%)	9.6% (3.2-20%)	9.3 (6.8-12)	15% (12-18%)	19% (15-23%)	31 (25-38)	790 (640-970)
Luxembourg	2005	Male	70 to 74	90% (80-96%)	10% (3.9-20%)	8.4 (5.9-11)	14% (10-17%)	17% (12-20%)	38 (29-48)	780 (590-970)
Luxembourg	2005	Male	75 to 79	89% (78-96%)	11% (4.3-22%)	7.7 (5.1-11)	13% (8.4-16%)	15% (9.7-19%)	44 (28-57)	690 (460-900)
Luxembourg	2005	Male	80 to 84	87% (73-95%)	13% (5.5-27%)	7.2 (5-9.6)	13% (8.6-16%)	14% (9.7-19%)	35 (24-46)	400 (280-520)
Luxembourg	2005	Male	85 to 89	84% (70-94%)	16% (6.5-30%)	6.7 (4.5-9.3)	12% (7.7-16%)	13% (8.5-18%)	21 (13-29)	170 (110-230)
Luxembourg	2005	Male	90 to 94	84% (69-94%)	16% (6-31%)	6.5 (4-9.5)	11% (6.4-16%)	13% (6.9-18%)	13 (7-18)	70 (40-96)
Luxembourg	2005	Male	95 plus	84% (67-95%)	16% (4.9-33%)	6.3 (3.4-10)	11% (4.7-16%)	12% (4.9-18%)	3.4 (1.4-5.1)	12 (5.4-18)
Luxembourg	2010	Female	All Ages	76% (68-82%)	24% (18-32%)	3.4 (2.8-4.2)	5.7% (4.2-7.5%)	8.3% (5.2-12%)	150 (94-230)	3400 (2600-4400)
Luxembourg	2010	Female	15 to 19	72% (55-86%)	28% (14-45%)	2.5 (1.6-3.7)	2.4% (1.5-3.6%)	9.9% (5.6-15%)	0.32 (0.18-0.49)	37 (24-54)
Luxembourg	2010	Female	20 to 24	77% (61-89%)	23% (11-39%)	3.2 (2.1-4.9)	3.1% (2.2-4.2%)	12% (7.4-18%)	0.37 (0.23-0.55)	54 (37-76)
Luxembourg	2010	Female	25 to 29	80% (66-90%)	20% (9.6-34%)	3.5 (2.3-5)	3.2% (2.2-4.2%)	13% (8.1-17%)	0.68 (0.44-0.91)	76 (52-99)
Luxembourg	2010	Female	30 to 34	81% (67-92%)	19% (8-33%)	3.7 (2.4-5.2)	3.4% (2.5-4.5%)	15% (11-19%)	1 (0.75-1.3)	93 (68-120)
Luxembourg	2010	Female	35 to 39	83% (69-93%)	17% (7.2-31%)	3.9 (2.6-5.5)	3.9% (2.9-5.2%)	17% (13-21%)	1.6 (1.2-2)	120 (91-150)
Luxembourg	2010	Female	40 to 44	82% (66-92%)	18% (8.1-34%)	4.1 (2.7-5.7)	4.7% (3.5-6.3%)	17% (14-21%)	2.7 (2.2-3.4)	170 (130-220)
Luxembourg	2010	Female	45 to 49	82% (65-91%)	18% (8.6-35%)	4.5 (3-6.5)	7.1% (5.3-9.3%)	18% (15-23%)	6.1 (4.8-7.7)	300 (230-390)
Luxembourg	2010	Female	50 to 54	83% (68-92%)	17% (7.7-32%)	5.2 (3.7-7.3)	8.1% (6.1-10%)	18% (14-22%)	7.7 (6.1-9.7)	340 (260-440)
Luxembourg	2010	Female	55 to 59	81% (67-92%)	19% (8.4-33%)	5.8 (4.1-8.1)	8.9% (6.6-12%)	17% (14-22%)	9.3 (7.4-12)	360 (270-470)
Luxembourg	2010	Female	60 to 64	80% (63-91%)	20% (8.8-37%)	5.6 (3.8-8.2)	8.8% (6.2-12%)	15% (11-20%)	11 (8.5-15)	360 (260-490)
Luxembourg	2010	Female	65 to 69	79% (61-91%)	21% (8.9-39%)	4.9 (3.4-6.6)	7.6% (5.2-10%)	12% (8.8-16%)	11 (7.8-14)	290 (200-400)
Luxembourg	2010	Female	70 to 74	75% (59-88%)	25% (12-41%)	3.9 (2.5-5.7)	6.2% (3.5-9.6%)	9.5% (5.7-14%)	12 (7.4-18)	270 (150-410)
Luxembourg	2010	Female	75 to 79	72% (52-87%)	28% (13-48%)	3.4 (2-5.2)	5.3% (2.3-9.1%)	7.3% (3.3-12%)	16 (7.2-27)	260 (110-470)
Luxembourg	2010	Female	80 to 84	71% (50-86%)	29% (14-50%)	3.4 (2.2-4.9)	5.5% (2.5-9.3%)	6.9% (3.2-12%)	24 (11-40)	290 (130-490)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	2010	Female	85 to 89	70% (52-84%)	30% (16-48%)	3.4 (2.1-5.1)	5.3% (2.2-9.5%)	6.3% (2.4-12%)	26 (9.6-47)	220 (86-390)
Luxembourg	2010	Female	90 to 94	69% (51-84%)	31% (16-49%)	3.4 (2-5.3)	5.1% (1.2-10%)	5.9% (1.1-12%)	15 (2.6-30)	84 (20-160)
Luxembourg	2010	Female	95 plus	68% (47-86%)	32% (14-53%)	3.3 (1.7-5.9)	5.3% (0.24-11%)	6% (0.085-13%)	7.7 (0.11-17)	28 (1.3-60)
Luxembourg	2010	Male	All Ages	84% (76-90%)	16% (10-24%)	7.4 (6-9)	13% (11-16%)	18% (15-21%)	310 (250-370)	8100 (6900-9400)
Luxembourg	2010	Male	15 to 19	75% (55-89%)	25% (11-45%)	4.4 (2.8-6.6)	6% (4.1-8.4%)	19% (11-28%)	0.78 (0.48-1.2)	87 (60-120)
Luxembourg	2010	Male	20 to 24	81% (64-91%)	19% (8.6-36%)	6.9 (4.5-10)	14% (10-18%)	27% (19-37%)	2.7 (1.9-3.6)	260 (190-330)
Luxembourg	2010	Male	25 to 29	85% (74-93%)	15% (6.6-26%)	8 (5.5-11)	12% (9.2-15%)	28% (21-36%)	2.6 (1.9-3.4)	260 (200-320)
Luxembourg	2010	Male	30 to 34	88% (77-95%)	12% (4.9-23%)	8.5 (6-12)	12% (9-14%)	29% (22-35%)	3.4 (2.6-4.2)	290 (220-350)
Luxembourg	2010	Male	35 to 39	90% (79-96%)	10% (3.9-21%)	8.5 (5.9-12)	13% (10-16%)	30% (24-36%)	5.5 (4.3-6.6)	380 (310-450)
Luxembourg	2010	Male	40 to 44	90% (79-96%)	9.9% (3.6-21%)	9 (6.5-12)	13% (11-16%)	30% (25-35%)	8 (6.6-9.4)	480 (400-570)
Luxembourg	2010	Male	45 to 49	91% (77-97%)	9.2% (3.5-23%)	10 (7.2-14)	16% (14-19%)	30% (26-34%)	15 (13-17)	750 (640-870)
Luxembourg	2010	Male	50 to 54	91% (80-97%)	8.5% (3.1-20%)	11 (7.9-15)	17% (14-20%)	28% (24-32%)	20 (17-23)	860 (740-990)
Luxembourg	2010	Male	55 to 59	91% (81-97%)	9.1% (3.2-19%)	12 (8.4-16)	17% (14-20%)	25% (21-29%)	25 (21-29)	900 (770-1100)
Luxembourg	2010	Male	60 to 64	91% (79-97%)	9.4% (3.2-21%)	11 (7.9-16)	16% (13-18%)	21% (18-25%)	27 (22-32)	850 (690-1000)
Luxembourg	2010	Male	65 to 69	90% (80-96%)	9.8% (3.6-20%)	10 (7.6-14)	15% (12-18%)	19% (16-23%)	29 (23-34)	750 (610-900)
Luxembourg	2010	Male	70 to 74	90% (79-96%)	10% (4.1-21%)	9.3 (6.4-13)	14% (11-17%)	17% (13-21%)	34 (27-42)	720 (560-900)
Luxembourg	2010	Male	75 to 79	89% (77-95%)	11% (4.6-23%)	8.6 (5.7-12)	13% (9.3-16%)	15% (11-19%)	39 (28-50)	640 (470-800)
Luxembourg	2010	Male	80 to 84	86% (73-94%)	14% (5.9-27%)	8.1 (5.7-11)	13% (9.5-16%)	15% (11-19%)	48 (36-63)	570 (430-730)
Luxembourg	2010	Male	85 to 89	84% (70-93%)	16% (6.8-30%)	7.5 (4.8-11)	12% (8.7-16%)	14% (9.6-18%)	30 (21-40)	240 (170-320)
Luxembourg	2010	Male	90 to 94	84% (70-94%)	16% (6.3-30%)	7.4 (4.5-11)	12% (7.2-16%)	13% (8-18%)	11 (7-16)	63 (40-85)
Luxembourg	2010	Male	95 plus	84% (68-95%)	16% (5.2-32%)	7.2 (3.9-12)	11% (5.8-16%)	13% (6.4-18%)	4 (2-5.8)	15 (7.6-21)
Luxembourg	2016	Female	All Ages	75% (67-82%)	25% (18-33%)	3.4 (2.5-4.6)	5.5% (3.7-7.7%)	7.8% (4.4-13%)	170 (92-270)	3700 (2500-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	2016	Female	15 to 19	72% (53-86%)	28% (14-47%)	2.5 (1.4-3.9)	2.2% (1.4-3.4%)	9.6% (5.2-15%)	0.32 (0.17-0.5)	38 (23-56)
Luxembourg	2016	Female	20 to 24	76% (59-89%)	24% (11-41%)	3.2 (1.9-5.1)	3.1% (2.1-4.3%)	12% (7-18%)	0.46 (0.26-0.69)	65 (44-91)
Luxembourg	2016	Female	25 to 29	79% (64-90%)	21% (10-36%)	3.4 (2-5.2)	3.1% (2.2-4.2%)	12% (7.9-17%)	0.71 (0.46-1)	84 (59-110)
Luxembourg	2016	Female	30 to 34	80% (64-92%)	20% (8.5-36%)	3.6 (2.2-5.6)	3.3% (2.4-4.6%)	15% (11-20%)	1.2 (0.85-1.7)	110 (76-150)
Luxembourg	2016	Female	35 to 39	81% (66-92%)	19% (8.3-34%)	3.7 (2.2-5.8)	3.9% (2.8-5.4%)	17% (12-21%)	1.9 (1.3-2.4)	140 (99-180)
Luxembourg	2016	Female	40 to 44	82% (66-92%)	18% (7.8-34%)	4.1 (2.5-6)	4.7% (3.4-6.4%)	18% (14-22%)	2.9 (2.2-3.8)	180 (130-240)
Luxembourg	2016	Female	45 to 49	82% (64-92%)	18% (7.9-36%)	4.5 (2.8-6.8)	6.6% (4.6-8.8%)	18% (14-23%)	6.1 (4.5-7.9)	310 (220-410)
Luxembourg	2016	Female	50 to 54	81% (64-92%)	19% (8.1-36%)	5 (3.2-7.2)	7.6% (5.5-10%)	17% (14-22%)	8.7 (6.4-12)	390 (280-520)
Luxembourg	2016	Female	55 to 59	81% (64-92%)	19% (8.3-36%)	5.6 (3.7-8.4)	8.4% (6.1-11%)	17% (13-22%)	10 (7.6-14)	400 (290-560)
Luxembourg	2016	Female	60 to 64	79% (60-92%)	21% (8.2-40%)	5.5 (3.5-8.6)	8.3% (5.7-12%)	15% (11-19%)	12 (8.6-17)	390 (270-550)
Luxembourg	2016	Female	65 to 69	78% (59-90%)	22% (9.5-41%)	4.7 (3-7)	7.2% (4.6-10%)	12% (8-16%)	12 (8.4-18)	340 (220-490)
Luxembourg	2016	Female	70 to 74	75% (57-88%)	25% (12-43%)	3.9 (2.3-6.1)	6.1% (3.1-9.9%)	9.3% (5.3-15%)	13 (7.1-20)	270 (140-450)
Luxembourg	2016	Female	75 to 79	72% (52-87%)	28% (13-48%)	3.4 (1.9-5.7)	5.3% (2.1-9.9%)	7.2% (3.1-13%)	16 (6.6-30)	270 (100-500)
Luxembourg	2016	Female	80 to 84	70% (48-85%)	30% (15-52%)	3.3 (1.9-5.2)	5.2% (2-9.7%)	6.6% (2.5-12%)	23 (9.2-44)	280 (110-530)
Luxembourg	2016	Female	85 to 89	67% (48-82%)	33% (18-52%)	3.2 (1.8-5)	4.9% (1.7-9.6%)	5.9% (1.8-11%)	29 (8.9-58)	240 (80-470)
Luxembourg	2016	Female	90 to 94	66% (47-83%)	34% (17-53%)	3.1 (1.7-5.4)	4.6% (0.97-9.7%)	5.3% (0.83-11%)	21 (3.2-46)	120 (24-250)
Luxembourg	2016	Female	95 plus	66% (43-85%)	34% (15-57%)	3.1 (1.5-5.7)	4.8% (0.14-11%)	5.4% (-0.078-13%)	7.7 (-0.11-18)	28 (0.85-62)
Luxembourg	2016	Male	All Ages	83% (76-89%)	17% (11-24%)	7.3 (5.3-9.8)	13% (11-15%)	17% (13-20%)	330 (260-410)	8600 (7000-10000)
Luxembourg	2016	Male	15 to 19	75% (55-89%)	25% (11-45%)	4.4 (2.5-6.8)	5.9% (3.9-8.4%)	18% (10-28%)	0.84 (0.47-1.3)	94 (61-130)
Luxembourg	2016	Male	20 to 24	80% (63-91%)	20% (8.9-37%)	6.7 (4.1-10)	12% (8.8-16%)	26% (17-35%)	2.2 (1.4-3.1)	240 (170-320)
Luxembourg	2016	Male	25 to 29	83% (70-93%)	17% (7.3-30%)	7.7 (4.8-12)	11% (8.8-14%)	27% (19-35%)	2.8 (1.9-3.8)	280 (220-370)
Luxembourg	2016	Male	30 to 34	86% (71-94%)	14% (5.6-29%)	8.2 (5.1-12)	11% (8.6-14%)	27% (20-34%)	4 (2.8-5.2)	330 (250-430)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Luxembourg	2016	Male	35 to 39	88% (75-95%)	12% (4.7-25%)	8.1 (5.3-12)	12% (9.2-15%)	29% (22-35%)	5.4 (4-7)	380 (290-480)
Luxembourg	2016	Male	40 to 44	89% (77-96%)	11% (3.9-23%)	8.7 (5.8-13)	13% (10-16%)	30% (24-35%)	8.2 (6.3-10)	490 (390-600)
Luxembourg	2016	Male	45 to 49	90% (76-96%)	10% (3.8-24%)	10 (6.4-15)	15% (12-18%)	29% (25-34%)	14 (11-18)	720 (580-870)
Luxembourg	2016	Male	50 to 54	90% (79-96%)	9.9% (3.6-21%)	11 (6.9-16)	16% (13-19%)	28% (24-31%)	22 (18-27)	960 (780-1100)
Luxembourg	2016	Male	55 to 59	90% (79-97%)	10% (3.2-21%)	11 (7.5-16)	16% (13-19%)	24% (21-28%)	26 (21-32)	980 (800-1200)
Luxembourg	2016	Male	60 to 64	89% (76-97%)	11% (3.3-24%)	11 (7.2-16)	15% (12-18%)	21% (17-24%)	29 (23-35)	910 (720-1100)
Luxembourg	2016	Male	65 to 69	89% (75-96%)	11% (3.9-25%)	10 (6.7-14)	14% (11-17%)	19% (15-22%)	33 (26-40)	860 (680-1000)
Luxembourg	2016	Male	70 to 74	88% (76-95%)	12% (4.7-24%)	8.9 (5.8-13)	13% (9.8-16%)	16% (13-20%)	33 (25-42)	700 (520-880)
Luxembourg	2016	Male	75 to 79	87% (74-95%)	13% (5.4-26%)	8.2 (5.1-13)	12% (8.5-15%)	14% (10-19%)	37 (26-48)	610 (440-780)
Luxembourg	2016	Male	80 to 84	86% (71-94%)	14% (6-29%)	7.8 (5-12)	12% (8.3-15%)	14% (9.7-18%)	47 (32-63)	560 (390-740)
Luxembourg	2016	Male	85 to 89	84% (69-93%)	16% (6.7-31%)	7.5 (4.7-11)	11% (7.6-15%)	13% (8.6-18%)	45 (29-60)	370 (250-490)
Luxembourg	2016	Male	90 to 94	84% (70-94%)	16% (5.8-30%)	7.3 (4.3-12)	11% (6.6-15%)	12% (7.3-17%)	19 (11-26)	100 (62-140)
Luxembourg	2016	Male	95 plus	84% (67-95%)	16% (4.5-33%)	7.1 (3.5-12)	11% (4.7-15%)	13% (5-18%)	3.9 (1.6-5.7)	14 (6.3-20)
Malta	1990	Female	All Ages	73% (64-81%)	27% (19-36%)	0.98 (0.87-1.1)	-0.18% (-1.4-1.2%)	-1.8% (-4.5-0.89%)	-24 (-60-11)	-78 (-630-500)
Malta	1990	Female	15 to 19	70% (50-86%)	30% (14-50%)	0.97 (0.63-1.4)	1.2% (0.7-1.7%)	4.1% (1.5-7.1%)	0.068 (0.025-0.12)	14 (8.1-21)
Malta	1990	Female	20 to 24	75% (57-88%)	25% (12-43%)	1.3 (0.85-1.7)	1.9% (1.3-2.6%)	3.9% (1.7-6.3%)	0.11 (0.048-0.19)	26 (16-38)
Malta	1990	Female	25 to 29	78% (62-90%)	22% (10-38%)	1.2 (0.82-1.5)	1.5% (0.99-2%)	3.1% (1.2-5.3%)	0.11 (0.042-0.19)	27 (18-39)
Malta	1990	Female	30 to 34	79% (62-91%)	21% (9.4-38%)	1.1 (0.77-1.5)	1.2% (0.78-1.7%)	3.3% (1.5-5.2%)	0.19 (0.083-0.31)	27 (17-38)
Malta	1990	Female	35 to 39	81% (65-92%)	19% (8.1-35%)	1.2 (0.82-1.6)	1.4% (0.82-2.1%)	4.2% (2.1-6.3%)	0.43 (0.21-0.66)	33 (20-48)
Malta	1990	Female	40 to 44	81% (65-92%)	19% (8-35%)	1.2 (0.88-1.7)	1.6% (0.77-2.5%)	4.6% (2.1-7.1%)	0.84 (0.41-1.3)	46 (23-72)
Malta	1990	Female	45 to 49	81% (63-92%)	19% (8.4-37%)	1.3 (0.89-1.8)	1.7% (0.76-3%)	4.8% (2.4-7.4%)	0.74 (0.37-1.2)	32 (14-54)
Malta	1990	Female	50 to 54	81% (63-91%)	19% (8.7-37%)	1.4 (1-1.9)	1.6% (0.37-3%)	3.9% (1.4-6.7%)	1.3 (0.44-2.1)	43 (9.5-80)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	1990	Female	55 to 59	80% (63-91%)	20% (8.8-37%)	1.6 (1.1-2.2)	1.2% (-0.56-3%)	2.4% (-0.44-5.5%)	1.2 (-0.22-2.8)	34 (-18-86)
Malta	1990	Female	60 to 64	78% (59-91%)	22% (8.6-41%)	1.6 (1.1-2.3)	-0.19% (-2.5-2.2%)	0.027% (-3.4-3.5%)	0.02 (-2.8-2.8)	-6.3 (-85-73)
Malta	1990	Female	65 to 69	76% (56-90%)	24% (9.8-44%)	1.5 (1.1-2.1)	-1.2% (-3.9-1.4%)	-1.5% (-5-2%)	-1.9 (-6.4-2.5)	-51 (-150-55)
Malta	1990	Female	70 to 74	74% (54-87%)	26% (13-46%)	1.4 (0.92-2.1)	-1.8% (-4.5-0.91%)	-2.3% (-5.8-1.3%)	-3.7 (-9.6-2.2)	-72 (-180-36)
Malta	1990	Female	75 to 79	71% (51-86%)	29% (14-49%)	1.3 (0.83-2)	-2.3% (-4.8-0.58%)	-2.8% (-6.1-0.61%)	-6.2 (-14-1.4)	-89 (-200-21)
Malta	1990	Female	80 to 84	68% (46-84%)	32% (16-54%)	1.3 (0.79-1.8)	-2.7% (-5.5-0.047%)	-3.2% (-6.6-0.06%)	-8.7 (-18-0.15)	-90 (-190-1.6)
Malta	1990	Female	85 to 89	65% (45-81%)	35% (19-55%)	1.2 (0.7-1.8)	-2.3% (-4.8-0.016%)	-2.9% (-5.8-0.023%)	-5.4 (-11-0.043)	-39 (-80-0.3)
Malta	1990	Female	90 to 94	64% (45-81%)	36% (19-55%)	1.2 (0.63-1.8)	-2.1% (-4.3-0.23%)	-2.7% (-5.4-0.065%)	-2.2 (-4.4-0.055)	-11 (-22-1.2)
Malta	1990	Female	95 plus	64% (40-84%)	36% (16-60%)	1.1 (0.55-2)	-2% (-4.4-0.46%)	-2.7% (-5.7-0.3%)	-0.83 (-1.8-0.092)	-2.6 (-5.7-0.62)
Malta	1990	Male	All Ages	78% (68-86%)	22% (14-32%)	1.7 (1.6-1.9)	2.1% (0.43-3.8%)	0.64% (-2.3-3.5%)	9 (-32-50)	980 (200-1800)
Malta	1990	Male	15 to 19	72% (51-88%)	28% (12-49%)	1.4 (0.9-2)	3.9% (2.5-5.5%)	6.9% (3.3-11%)	0.49 (0.24-0.79)	57 (37-81)
Malta	1990	Male	20 to 24	78% (59-90%)	22% (10-41%)	2.1 (1.4-2.9)	7.4% (5.1-9.9%)	9.8% (5.1-15%)	0.9 (0.48-1.4)	110 (77-160)
Malta	1990	Male	25 to 29	81% (67-92%)	19% (8.3-33%)	2 (1.4-2.6)	5.9% (4.3-7.8%)	8.1% (4.4-13%)	0.87 (0.46-1.3)	120 (80-160)
Malta	1990	Male	30 to 34	84% (68-94%)	16% (6.2-32%)	2 (1.4-2.6)	4.6% (3.2-6.2%)	7% (3.8-11%)	0.85 (0.46-1.3)	97 (66-130)
Malta	1990	Male	35 to 39	87% (74-95%)	13% (5.2-26%)	2 (1.4-2.7)	3.9% (2.7-5.3%)	6.1% (3.1-9.4%)	0.82 (0.41-1.3)	80 (54-110)
Malta	1990	Male	40 to 44	88% (74-96%)	12% (4.2-26%)	2.4 (1.8-3.1)	3.7% (2.2-5.4%)	5.6% (2.1-9%)	1.4 (0.49-2.2)	99 (55-140)
Malta	1990	Male	45 to 49	89% (71-96%)	11% (4.1-29%)	3 (2.1-4.1)	4.4% (2-7.2%)	6% (2-11%)	1.5 (0.48-2.7)	80 (34-130)
Malta	1990	Male	50 to 54	89% (75-96%)	11% (4-25%)	3.3 (2.4-4.5)	4.3% (1.4-7.4%)	5.7% (1.2-10%)	2.7 (0.52-5)	120 (40-210)
Malta	1990	Male	55 to 59	89% (76-96%)	11% (3.7-24%)	3.3 (2.4-4.3)	3.6% (0.72-6.8%)	4.2% (0.2-8.7%)	3.1 (0.16-6.5)	120 (24-240)
Malta	1990	Male	60 to 64	88% (74-96%)	12% (3.9-26%)	3.1 (2.2-4.3)	2.4% (-0.98-6.1%)	2.5% (-1.9-7.2%)	3.1 (-2.2-9)	100 (-43-270)
Malta	1990	Male	65 to 69	88% (73-96%)	12% (4.2-27%)	2.8 (2.1-3.6)	1% (-2.3-4.3%)	0.87% (-3.1-4.9%)	1.5 (-5.3-8.2)	48 (-110-200)
Malta	1990	Male	70 to 74	87% (72-94%)	13% (5.6-28%)	2.4 (1.7-3.3)	0.13% (-2.9-3.2%)	-0.12% (-3.7-3.5%)	-0.23 (-7.5-7.2)	6 (-130-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	1990	Male	75 to 79	85% (70-94%)	15% (5.8-30%)	2.2 (1.4-3.1)	-0.53% (-3.4-2.5%)	-0.87% (-4.3-2.6%)	-2 (-9.9-6.1)	-20 (-130-95)
Malta	1990	Male	80 to 84	83% (68-93%)	17% (7-32%)	2 (1.4-2.8)	-0.99% (-3.7-1.7%)	-1.4% (-4.7-1.7%)	-3 (-10-3.9)	-26 (-99-45)
Malta	1990	Male	85 to 89	81% (66-92%)	19% (7.8-34%)	1.9 (1.3-2.7)	-1% (-3.4-1.3%)	-1.6% (-4.5-1.2%)	-1.8 (-5.2-1.4)	-11 (-36-13)
Malta	1990	Male	90 to 94	82% (66-93%)	18% (7.1-34%)	1.8 (1.1-2.7)	-1.2% (-3.5-1.2%)	-1.8% (-4.7-1.2%)	-0.87 (-2.3-0.56)	-3.5 (-11-3.7)
Malta	1990	Male	95 plus	83% (64-94%)	17% (5.6-36%)	1.8 (0.95-2.9)	-1.1% (-3.4-1.5%)	-1.8% (-4.8-1.3%)	-0.24 (-0.63-0.18)	-0.6 (-1.9-0.84)
Malta	1995	Female	All Ages	73% (65-80%)	27% (20-35%)	1.1 (0.94-1.2)	0.2% (-0.93-1.4%)	-1.1% (-3.5-1.5%)	-14 (-47-19)	90 (-400-600)
Malta	1995	Female	15 to 19	70% (51-85%)	30% (15-49%)	1 (0.67-1.4)	1.2% (0.77-1.7%)	3.6% (1.4-6%)	0.099 (0.038-0.17)	17 (10-26)
Malta	1995	Female	20 to 24	75% (57-88%)	25% (12-43%)	1.3 (0.9-1.8)	1.9% (1.3-2.6%)	4% (1.8-6.4%)	0.1 (0.045-0.17)	27 (17-40)
Malta	1995	Female	25 to 29	77% (61-90%)	23% (10-39%)	1.2 (0.84-1.7)	1.5% (1-2%)	3.3% (1.5-5.4%)	0.081 (0.036-0.13)	21 (14-30)
Malta	1995	Female	30 to 34	79% (62-91%)	21% (8.9-38%)	1.2 (0.81-1.6)	1.2% (0.77-1.7%)	3.3% (1.8-5.1%)	0.15 (0.077-0.24)	24 (15-34)
Malta	1995	Female	35 to 39	81% (65-91%)	19% (8.6-35%)	1.2 (0.86-1.7)	1.3% (0.81-1.8%)	4.3% (2.4-6.4%)	0.35 (0.19-0.54)	30 (19-44)
Malta	1995	Female	40 to 44	81% (64-92%)	19% (8.1-36%)	1.3 (0.91-1.8)	1.6% (0.87-2.5%)	5% (2.8-7.5%)	0.71 (0.4-1.1)	41 (24-63)
Malta	1995	Female	45 to 49	81% (62-92%)	19% (8.4-38%)	1.4 (0.94-1.9)	1.7% (0.83-2.8%)	5.3% (3-7.8%)	1.2 (0.69-1.8)	53 (26-84)
Malta	1995	Female	50 to 54	81% (63-91%)	19% (8.5-37%)	1.5 (1.1-2)	2% (0.77-3.3%)	4.9% (2.5-7.4%)	1.2 (0.6-1.9)	43 (17-70)
Malta	1995	Female	55 to 59	80% (63-91%)	20% (8.6-37%)	1.7 (1.2-2.3)	1.6% (-0.0049-3.2%)	3.4% (0.78-6.3%)	1.7 (0.37-3.1)	48 (-0.14-99)
Malta	1995	Female	60 to 64	78% (60-91%)	22% (8.6-40%)	1.7 (1.2-2.4)	0.66% (-1.4-2.6%)	1.4% (-1.7-4.5%)	0.98 (-1.2-3.1)	21 (-42-86)
Malta	1995	Female	65 to 69	76% (56-90%)	24% (9.7-44%)	1.6 (1.1-2.2)	-0.47% (-2.8-1.9%)	-0.46% (-3.7-2.9%)	-0.5 (-4.1-3.1)	-17 (-100-69)
Malta	1995	Female	70 to 74	74% (55-87%)	26% (13-45%)	1.5 (0.98-2.2)	-1.1% (-3.6-1.4%)	-1.4% (-4.8-2%)	-2.5 (-8.3-3.5)	-50 (-160-64)
Malta	1995	Female	75 to 79	70% (50-86%)	30% (14-50%)	1.4 (0.85-2.1)	-1.6% (-4.1-0.93%)	-2.1% (-5.3-1.2%)	-4.1 (-10-2.4)	-60 (-150-34)
Malta	1995	Female	80 to 84	67% (45-84%)	33% (16-55%)	1.3 (0.82-1.9)	-2% (-4.5-0.43%)	-2.4% (-5.5-0.47%)	-5.8 (-13-1.1)	-60 (-140-13)
Malta	1995	Female	85 to 89	65% (45-81%)	35% (19-55%)	1.3 (0.75-1.9)	-1.7% (-3.9-0.49%)	-2.2% (-4.8-0.44%)	-5 (-11-0.99)	-36 (-82-10)
Malta	1995	Female	90 to 94	64% (44-81%)	36% (19-56%)	1.2 (0.66-1.9)	-1.5% (-3.5-0.48%)	-2.1% (-4.5-0.34%)	-2.2 (-4.8-0.36)	-11 (-24-3.3)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	1995	Female	95 plus	64% (41-84%)	36% (16-59%)	1.2 (0.58-2.1)	-1.5% (-3.6-0.79%)	-2.2% (-4.9-0.55%)	-0.73 (-1.7-0.19)	-2.2 (-5.2-1.1)
Malta	1995	Male	All Ages	78% (69-86%)	22% (14-31%)	1.9 (1.7-2.1)	2.6% (0.99-4.2%)	1.3% (-1.4-4%)	19 (-20-58)	1200 (470-2000)
Malta	1995	Male	15 to 19	71% (51-87%)	29% (13-49%)	1.5 (0.95-2.1)	3.8% (2.5-5.4%)	6.8% (3.3-11%)	0.54 (0.26-0.84)	64 (41-92)
Malta	1995	Male	20 to 24	77% (58-90%)	23% (9.7-42%)	2.2 (1.5-3)	7.4% (5.1-9.9%)	9.8% (5.3-15%)	0.96 (0.52-1.4)	120 (81-170)
Malta	1995	Male	25 to 29	81% (65-92%)	19% (7.6-35%)	2.1 (1.5-2.7)	6.2% (4.3-8.1%)	8.3% (4.5-12%)	0.84 (0.45-1.3)	100 (69-140)
Malta	1995	Male	30 to 34	84% (67-94%)	16% (6.1-33%)	2.1 (1.5-2.7)	4.7% (3.3-6.3%)	7.1% (3.8-11%)	0.86 (0.47-1.3)	96 (66-130)
Malta	1995	Male	35 to 39	86% (72-95%)	14% (5.1-28%)	2.1 (1.5-3)	4.2% (2.8-5.6%)	6.5% (3.5-9.7%)	1.1 (0.57-1.7)	97 (64-130)
Malta	1995	Male	40 to 44	88% (74-96%)	12% (4.1-26%)	2.5 (1.8-3.3)	3.9% (2.5-5.5%)	6.4% (3-9.6%)	1.2 (0.57-1.9)	92 (57-130)
Malta	1995	Male	45 to 49	88% (72-96%)	12% (4.1-28%)	3.2 (2.3-4.4)	4.8% (2.7-7.3%)	7.2% (3.4-12%)	2.8 (1.3-4.7)	160 (86-240)
Malta	1995	Male	50 to 54	89% (74-96%)	11% (4-26%)	3.5 (2.5-4.7)	4.9% (2.2-8%)	6.7% (2.3-12%)	2.4 (0.81-4.1)	110 (47-170)
Malta	1995	Male	55 to 59	88% (76-96%)	12% (4-24%)	3.5 (2.6-4.6)	4.3% (1.8-7.5%)	5.4% (1.7-9.9%)	4 (1.3-7.3)	150 (63-260)
Malta	1995	Male	60 to 64	88% (73-96%)	12% (4.1-27%)	3.3 (2.4-4.5)	3.2% (0.28-6.9%)	3.7% (-0.2-8.5%)	3.9 (-0.2-9)	130 (11-270)
Malta	1995	Male	65 to 69	88% (73-96%)	12% (4.2-27%)	3 (2.2-3.9)	1.8% (-1.4-6%)	1.9% (-1.7-5.5%)	3.1 (-2.8-9.2)	88 (-46-230)
Malta	1995	Male	70 to 74	87% (73-95%)	13% (5.4-27%)	2.6 (1.8-3.5)	0.73% (-2.3-3.8%)	0.56% (-3-4.3%)	1.2 (-6.5-9.3)	35 (-110-180)
Malta	1995	Male	75 to 79	85% (70-94%)	15% (5.9-30%)	2.3 (1.5-3.2)	0.041% (-2.8-3%)	-0.23% (-3.6-3.2%)	-0.52 (-8-7.1)	1.6 (-110-110)
Malta	1995	Male	80 to 84	83% (67-93%)	17% (6.6-33%)	2.1 (1.4-2.9)	-0.33% (-2.9-2%)	-0.63% (-3.6-2.1%)	-1.4 (-8-4.6)	-8.6 (-77-55)
Malta	1995	Male	85 to 89	82% (65-92%)	18% (7.5-35%)	2 (1.3-2.8)	-0.53% (-2.9-1.7%)	-0.93% (-3.8-1.7%)	-1.4 (-5.6-2.5)	-7 (-38-22)
Malta	1995	Male	90 to 94	82% (64-93%)	18% (6.8-36%)	1.9 (1.2-2.9)	-0.65% (-2.8-1.6%)	-1.2% (-3.9-1.5%)	-0.73 (-2.4-0.93)	-2.6 (-11-5.8)
Malta	1995	Male	95 plus	82% (63-95%)	18% (5.3-37%)	1.9 (1-3)	-0.7% (-3-2%)	-1.3% (-4.3-1.9%)	-0.24 (-0.77-0.34)	-0.53 (-2.3-1.5)
Malta	2000	Female	All Ages	73% (65-81%)	27% (19-35%)	1 (0.91-1.2)	0.16% (-0.91-1.3%)	-1.2% (-3.5-1.1%)	-18 (-52-17)	74 (-410-580)
Malta	2000	Female	15 to 19	70% (51-86%)	30% (14-49%)	0.96 (0.64-1.3)	1.2% (0.73-1.8%)	3.3% (1.2-5.6%)	0.1 (0.036-0.18)	17 (9.9-26)
Malta	2000	Female	20 to 24	75% (57-88%)	25% (12-43%)	1.3 (0.84-1.7)	1.9% (1.3-2.5%)	3.6% (1.5-5.7%)	0.13 (0.056-0.21)	32 (20-45)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	2000	Female	25 to 29	78% (62-90%)	22% (10-38%)	1.2 (0.78-1.6)	1.4% (1-1.9%)	3% (1.2-4.9%)	0.096 (0.039-0.16)	23 (16-34)
Malta	2000	Female	30 to 34	80% (63-91%)	20% (9-37%)	1.1 (0.75-1.6)	1.2% (0.81-1.7%)	3.4% (1.7-5.2%)	0.15 (0.076-0.23)	20 (13-29)
Malta	2000	Female	35 to 39	81% (65-91%)	19% (8.7-35%)	1.2 (0.79-1.6)	1.3% (0.83-1.8%)	4.3% (2.5-6.3%)	0.33 (0.19-0.5)	29 (19-41)
Malta	2000	Female	40 to 44	81% (65-92%)	19% (8.1-35%)	1.2 (0.86-1.6)	1.4% (0.74-2.1%)	4.8% (2.8-7.2%)	0.59 (0.33-0.9)	36 (20-54)
Malta	2000	Female	45 to 49	81% (62-92%)	19% (8.3-38%)	1.3 (0.9-1.8)	1.7% (0.77-2.7%)	4.7% (2.5-7.2%)	1.1 (0.6-1.8)	50 (24-79)
Malta	2000	Female	50 to 54	81% (63-92%)	19% (8.5-37%)	1.4 (1-1.9)	1.8% (0.75-3%)	4.6% (2.5-7%)	1.9 (1-2.9)	67 (28-110)
Malta	2000	Female	55 to 59	80% (63-91%)	20% (8.6-37%)	1.6 (1.1-2.2)	1.6% (0.031-3.1%)	3.4% (0.81-6.1%)	1.4 (0.31-2.5)	39 (0.83-79)
Malta	2000	Female	60 to 64	78% (59-91%)	22% (8.6-41%)	1.6 (1.1-2.3)	0.71% (-1-2.4%)	1.6% (-1.1-4.4%)	1.2 (-0.85-3.4)	26 (-36-90)
Malta	2000	Female	65 to 69	77% (55-90%)	23% (9.8-45%)	1.5 (1.1-2.1)	-0.28% (-2.2-1.7%)	-0.13% (-2.9-2.9%)	-0.14 (-3.1-3)	-10 (-82-65)
Malta	2000	Female	70 to 74	74% (55-87%)	26% (13-45%)	1.4 (0.93-2)	-1% (-3.3-1.2%)	-1.3% (-4.3-1.7%)	-2.1 (-7.2-3)	-44 (-140-53)
Malta	2000	Female	75 to 79	71% (50-86%)	29% (14-50%)	1.3 (0.83-2)	-1.7% (-4-0.62%)	-2.1% (-5.3-0.75%)	-5.4 (-13-2)	-79 (-190-31)
Malta	2000	Female	80 to 84	68% (45-84%)	32% (16-55%)	1.3 (0.78-1.8)	-2% (-4.3-0.3%)	-2.5% (-5.3-0.29%)	-6.7 (-14-0.83)	-70 (-150-10)
Malta	2000	Female	85 to 89	65% (45-81%)	35% (19-55%)	1.2 (0.69-1.8)	-1.9% (-4.1-0.27%)	-2.4% (-5-0.29%)	-5.8 (-12-0.64)	-42 (-91-6.1)
Malta	2000	Female	90 to 94	64% (45-82%)	36% (18-55%)	1.2 (0.64-1.8)	-1.7% (-3.6-0.27%)	-2.3% (-4.7-0.2%)	-3.3 (-6.8-0.28)	-16 (-34-2.6)
Malta	2000	Female	95 plus	64% (40-84%)	36% (16-60%)	1.1 (0.54-2)	-1.7% (-3.7-0.56%)	-2.3% (-4.9-0.4%)	-1.3 (-2.9-0.22)	-4.2 (-9.4-1.4)
Malta	2000	Male	All Ages	80% (70-87%)	20% (13-30%)	1.9 (1.7-2.1)	2.4% (0.79-4%)	1% (-1.9-3.7%)	17 (-28-58)	1200 (400-2000)
Malta	2000	Male	15 to 19	72% (51-88%)	28% (12-49%)	1.4 (0.89-2)	3.8% (2.4-5.3%)	6.6% (3.2-11%)	0.51 (0.25-0.82)	59 (38-85)
Malta	2000	Male	20 to 24	78% (59-90%)	22% (10-41%)	2.1 (1.4-2.9)	7% (4.9-9.4%)	9.2% (4.9-14%)	0.99 (0.54-1.5)	130 (90-180)
Malta	2000	Male	25 to 29	82% (67-92%)	18% (8.1-33%)	2 (1.4-2.6)	5.9% (4.2-7.8%)	7.9% (4.1-12%)	0.95 (0.51-1.5)	110 (77-150)
Malta	2000	Male	30 to 34	85% (69-94%)	15% (6.1-31%)	2 (1.4-2.6)	4.5% (3.2-6%)	7.1% (3.9-11%)	0.62 (0.34-0.95)	74 (52-100)
Malta	2000	Male	35 to 39	87% (74-95%)	13% (5.1-26%)	2 (1.4-2.8)	4% (2.7-5.4%)	6.4% (3.5-9.8%)	0.99 (0.52-1.6)	92 (61-130)
Malta	2000	Male	40 to 44	88% (74-96%)	12% (4.2-26%)	2.4 (1.8-3.1)	3.8% (2.4-5.2%)	6.2% (2.9-9.5%)	1.3 (0.61-2)	99 (60-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	2000	Male	45 to 49	89% (71-96%)	11% (4.1-29%)	3 (2.2-4.1)	4.6% (2.5-6.9%)	6.9% (3.3-11%)	2.6 (1.2-4.3)	140 (77-220)
Malta	2000	Male	50 to 54	89% (75-96%)	11% (3.9-25%)	3.4 (2.4-4.4)	4.7% (2.2-7.3%)	6.6% (2.6-11%)	4.4 (1.6-7.3)	200 (91-310)
Malta	2000	Male	55 to 59	89% (77-96%)	11% (3.6-23%)	3.3 (2.5-4.3)	4% (1.5-6.8%)	5% (1.5-8.9%)	3.1 (0.9-5.6)	120 (44-200)
Malta	2000	Male	60 to 64	88% (73-96%)	12% (3.8-27%)	3.1 (2.2-4.2)	2.9% (0.084-5.9%)	3.3% (-0.5-7.1%)	3.8 (-0.62-8.4)	130 (3.6-260)
Malta	2000	Male	65 to 69	88% (73-96%)	12% (4.1-27%)	2.8 (2.1-3.6)	1.5% (-1.4-4.2%)	1.5% (-2.1-4.9%)	2.6 (-3.6-8.6)	76 (-68-220)
Malta	2000	Male	70 to 74	87% (73-95%)	13% (5.4-27%)	2.4 (1.7-3.3)	0.57% (-2.3-3.4%)	0.41% (-3.1-3.8%)	0.99 (-7.1-9)	31 (-120-180)
Malta	2000	Male	75 to 79	85% (70-94%)	15% (5.7-30%)	2.2 (1.4-3.1)	-0.16% (-3-2.6%)	-0.44% (-3.7-2.8%)	-1.2 (-10-7.8)	-7.5 (-130-120)
Malta	2000	Male	80 to 84	84% (68-93%)	16% (6.9-32%)	2 (1.4-2.8)	-0.5% (-3.1-1.9%)	-0.82% (-3.8-2%)	-2 (-9.2-4.7)	-15 (-88-57)
Malta	2000	Male	85 to 89	82% (66-92%)	18% (7.8-34%)	1.9 (1.2-2.7)	-0.69% (-2.9-1.6%)	-1.1% (-3.8-1.5%)	-1.8 (-6.5-2.6)	-10 (-44-23)
Malta	2000	Male	90 to 94	82% (67-93%)	18% (7.1-33%)	1.8 (1.1-2.7)	-0.84% (-3-1.3%)	-1.3% (-4-1.3%)	-1 (-3.2-1)	-4.1 (-15-6.5)
Malta	2000	Male	95 plus	82% (63-94%)	18% (5.6-37%)	1.8 (0.97-2.9)	-0.87% (-3.1-1.5%)	-1.5% (-4.4-1.4%)	-0.36 (-1.1-0.34)	-0.9 (-3.3-1.6)
Malta	2005	Female	All Ages	73% (64-80%)	27% (20-36%)	1 (0.88-1.2)	0.27% (-0.6-1.2%)	-0.89% (-2.8-0.99%)	-14 (-44-16)	130 (-280-570)
Malta	2005	Female	15 to 19	74% (56-87%)	26% (13-44%)	0.99 (0.65-1.3)	1.1% (0.69-1.6%)	3.6% (1.3-6%)	0.067 (0.024-0.11)	14 (8.3-21)
Malta	2005	Female	20 to 24	78% (62-90%)	22% (10-38%)	1.3 (0.87-1.7)	1.9% (1.3-2.6%)	3.5% (1.6-5.5%)	0.11 (0.047-0.18)	27 (17-39)
Malta	2005	Female	25 to 29	80% (66-91%)	20% (9.5-34%)	1.2 (0.8-1.6)	1.4% (0.98-2%)	2.9% (1.3-4.6%)	0.099 (0.041-0.16)	25 (16-35)
Malta	2005	Female	30 to 34	80% (65-91%)	20% (9.3-35%)	1.1 (0.74-1.5)	1.2% (0.82-1.7%)	3.4% (1.9-5.1%)	0.15 (0.081-0.24)	22 (14-32)
Malta	2005	Female	35 to 39	80% (65-91%)	20% (9.2-35%)	1.1 (0.76-1.6)	1.3% (0.81-1.8%)	4.3% (2.6-6.3%)	0.28 (0.16-0.42)	25 (16-36)
Malta	2005	Female	40 to 44	79% (62-91%)	21% (9.3-38%)	1.2 (0.81-1.6)	1.3% (0.74-2%)	4.8% (2.8-7.1%)	0.49 (0.28-0.74)	31 (18-47)
Malta	2005	Female	45 to 49	79% (60-90%)	21% (10-40%)	1.2 (0.84-1.6)	1.6% (0.8-2.5%)	4.6% (2.7-6.7%)	1 (0.59-1.6)	47 (24-72)
Malta	2005	Female	50 to 54	79% (61-90%)	21% (10-39%)	1.3 (0.94-1.8)	1.7% (0.74-2.7%)	4.5% (2.6-6.7%)	1.5 (0.8-2.2)	53 (25-84)
Malta	2005	Female	55 to 59	77% (59-89%)	23% (11-41%)	1.4 (1-2)	1.7% (0.48-3%)	3.7% (1.6-6%)	2.3 (0.98-3.8)	68 (20-120)
Malta	2005	Female	60 to 64	75% (55-89%)	25% (11-45%)	1.5 (1-2.1)	1% (-0.34-2.4%)	2.3% (0.0071-4.6%)	1.6 (0.0048-3.2)	36 (-11-88)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	2005	Female	65 to 69	74% (54-88%)	26% (12-46%)	1.4 (1-1.9)	0.031% (-1.4-1.5%)	0.44% (-1.7-2.7%)	0.5 (-2-3)	1.4 (-61-67)
Malta	2005	Female	70 to 74	69% (50-84%)	31% (16-50%)	1.2 (0.78-1.8)	-0.57% (-2.2-1.1%)	-0.65% (-3-1.6%)	-1.1 (-5-2.7)	-25 (-100-48)
Malta	2005	Female	75 to 79	62% (41-81%)	38% (19-59%)	1.1 (0.66-1.6)	-1.1% (-2.9-0.55%)	-1.4% (-3.8-0.73%)	-3.9 (-10-1.9)	-57 (-150-28)
Malta	2005	Female	80 to 84	63% (40-80%)	37% (20-60%)	1.1 (0.66-1.6)	-1.6% (-3.6-0.13%)	-2.1% (-4.5-0.1%)	-6.7 (-14-0.34)	-70 (-150-5.5)
Malta	2005	Female	85 to 89	63% (43-80%)	37% (20-57%)	1.1 (0.67-1.7)	-1.7% (-3.6-0.18%)	-2.2% (-4.7-0.13%)	-5.9 (-12-0.38)	-43 (-92-4.7)
Malta	2005	Female	90 to 94	62% (43-80%)	38% (20-57%)	1.1 (0.61-1.7)	-1.5% (-3.3-0.2%)	-2.1% (-4.2-0.15%)	-3.1 (-6.5-0.24)	-15 (-33-2.1)
Malta	2005	Female	95 plus	63% (39-83%)	37% (17-61%)	1.1 (0.52-1.9)	-1.5% (-3.5-0.38%)	-2.1% (-4.6-0.26%)	-1.6 (-3.5-0.2)	-5 (-11-1.4)
Malta	2005	Male	All Ages	82% (74-89%)	18% (11-26%)	2 (1.8-2.2)	2.5% (0.98-3.9%)	1.1% (-1.5-3.5%)	17 (-23-54)	1200 (490-1900)
Malta	2005	Male	15 to 19	74% (54-88%)	26% (12-46%)	1.4 (0.92-2)	3.6% (2.3-5%)	6.5% (3.2-10%)	0.44 (0.21-0.7)	56 (36-80)
Malta	2005	Male	20 to 24	81% (64-91%)	19% (8.6-36%)	2.1 (1.5-2.9)	7% (4.8-9.2%)	9.2% (4.8-14%)	0.91 (0.48-1.4)	120 (82-170)
Malta	2005	Male	25 to 29	85% (73-93%)	15% (6.6-27%)	2 (1.5-2.7)	5.7% (4.1-7.4%)	7.9% (4.3-12%)	0.84 (0.44-1.3)	110 (78-150)
Malta	2005	Male	30 to 34	88% (76-95%)	12% (4.7-24%)	2 (1.4-2.7)	4.5% (3.2-6.1%)	7.2% (4-11%)	0.72 (0.38-1.1)	83 (57-110)
Malta	2005	Male	35 to 39	89% (78-96%)	11% (4.2-22%)	2 (1.4-2.8)	4% (2.8-5.4%)	6.9% (3.8-10%)	0.73 (0.38-1.1)	71 (48-97)
Malta	2005	Male	40 to 44	90% (78-96%)	10% (4-22%)	2.4 (1.7-3.1)	3.8% (2.5-5.3%)	6.5% (3.3-9.8%)	1.3 (0.66-2)	97 (64-140)
Malta	2005	Male	45 to 49	89% (76-96%)	11% (3.9-24%)	3 (2.2-4.1)	4.3% (2.6-6.4%)	7.2% (3.7-11%)	2.1 (1.1-3.4)	130 (75-190)
Malta	2005	Male	50 to 54	90% (78-96%)	9.8% (3.6-22%)	3.4 (2.5-4.4)	4.6% (2.2-7.2%)	6.8% (2.7-11%)	3.6 (1.4-6)	170 (80-260)
Malta	2005	Male	55 to 59	90% (80-97%)	9.5% (3.3-20%)	3.4 (2.6-4.2)	3.9% (1.7-6.5%)	5.2% (1.9-9.1%)	4.8 (1.7-8.5)	190 (82-320)
Malta	2005	Male	60 to 64	88% (75-96%)	12% (3.9-25%)	3.1 (2.2-4.1)	3% (0.5-5.7%)	3.5% (0.17-7.1%)	3 (0.14-6.1)	98 (16-190)
Malta	2005	Male	65 to 69	88% (75-96%)	12% (4.2-25%)	2.7 (2.1-3.5)	1.7% (-0.79-4.2%)	1.8% (-1.5-5%)	2.9 (-2.3-7.8)	84 (-37-200)
Malta	2005	Male	70 to 74	88% (75-95%)	12% (5.1-25%)	2.4 (1.7-3.3)	0.7% (-1.7-3.1%)	0.56% (-2.4-3.5%)	1.2 (-5.2-7.5)	35 (-85-160)
Malta	2005	Male	75 to 79	85% (71-94%)	15% (5.8-29%)	2.1 (1.4-3)	0.08% (-2.3-2.6%)	-0.18% (-2.9-2.8%)	-0.47 (-8-7.2)	3.7 (-110-120)
Malta	2005	Male	80 to 84	84% (68-93%)	16% (6.6-32%)	2 (1.3-2.7)	-0.4% (-2.7-1.9%)	-0.71% (-3.5-2%)	-1.9 (-9.9-5.4)	-13 (-95-64)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	2005	Male	85 to 89	83% (67-93%)	17% (7-33%)	1.9 (1.2-2.6)	-0.55% (-2.7-1.6%)	-0.93% (-3.6-1.5%)	-1.6 (-6.3-2.7)	-8.7 (-43-24)
Malta	2005	Male	90 to 94	83% (68-93%)	17% (6.6-32%)	1.8 (1.2-2.7)	-0.69% (-2.8-1.4%)	-1.2% (-3.7-1.3%)	-0.98 (-3.2-1.1)	-3.7 (-15-7.2)
Malta	2005	Male	95 plus	83% (66-95%)	17% (5.1-34%)	1.8 (0.95-2.9)	-0.62% (-2.7-1.7%)	-1.2% (-3.8-1.6%)	-0.33 (-1.1-0.45)	-0.76 (-3.3-2)
Malta	2010	Female	All Ages	72% (64-80%)	28% (20-36%)	1.1 (0.92-1.3)	0.44% (-0.39-1.3%)	-0.53% (-2.4-1.4%)	-8.3 (-37-22)	220 (-190-640)
Malta	2010	Female	15 to 19	73% (57-87%)	27% (13-43%)	1 (0.68-1.4)	1.1% (0.71-1.6%)	3.6% (1.5-6%)	0.063 (0.025-0.11)	15 (9.3-23)
Malta	2010	Female	20 to 24	78% (62-89%)	22% (11-38%)	1.4 (0.91-1.9)	1.8% (1.2-2.5%)	3.7% (1.8-5.8%)	0.081 (0.039-0.13)	22 (14-32)
Malta	2010	Female	25 to 29	80% (66-91%)	20% (9.4-34%)	1.2 (0.86-1.6)	1.5% (1-2.1%)	2.9% (1.2-4.7%)	0.13 (0.054-0.21)	23 (15-32)
Malta	2010	Female	30 to 34	80% (65-91%)	20% (9.3-35%)	1.2 (0.79-1.6)	1.2% (0.77-1.6%)	3.6% (2.1-5.2%)	0.16 (0.091-0.24)	26 (17-37)
Malta	2010	Female	35 to 39	79% (63-91%)	21% (9.4-37%)	1.2 (0.8-1.7)	1.2% (0.76-1.7%)	4.6% (2.8-6.5%)	0.24 (0.15-0.36)	25 (16-35)
Malta	2010	Female	40 to 44	79% (61-91%)	21% (9.4-39%)	1.2 (0.84-1.7)	1.3% (0.76-2%)	5.1% (3.1-7.4%)	0.44 (0.27-0.66)	29 (17-43)
Malta	2010	Female	45 to 49	79% (59-90%)	21% (10-41%)	1.3 (0.86-1.8)	1.5% (0.77-2.3%)	4.9% (3-7.1%)	0.86 (0.51-1.3)	40 (21-63)
Malta	2010	Female	50 to 54	78% (60-90%)	22% (10-40%)	1.4 (0.96-1.8)	1.7% (0.74-2.8%)	4.8% (2.7-6.9%)	1.5 (0.83-2.2)	54 (25-87)
Malta	2010	Female	55 to 59	76% (60-89%)	24% (11-40%)	1.5 (1.1-2.1)	1.7% (0.49-2.9%)	4.2% (2-6.5%)	2 (0.91-3)	59 (17-100)
Malta	2010	Female	60 to 64	75% (55-89%)	25% (11-45%)	1.5 (1.1-2.2)	1.2% (-0.094-2.7%)	2.8% (0.61-5%)	2.5 (0.56-4.5)	60 (-4.5-120)
Malta	2010	Female	65 to 69	73% (53-88%)	27% (12-47%)	1.5 (1.1-2)	0.43% (-0.75-1.7%)	1.3% (-0.62-3.2%)	1.3 (-0.6-3.2)	19 (-33-75)
Malta	2010	Female	70 to 74	69% (50-84%)	31% (16-50%)	1.3 (0.85-1.9)	-0.15% (-1.5-1.4%)	0.018% (-2.1-2.3%)	0.023 (-3.2-3.4)	-7.1 (-70-63)
Malta	2010	Female	75 to 79	63% (41-81%)	37% (19-59%)	1.1 (0.69-1.7)	-0.85% (-2.5-0.89%)	-1.1% (-3.5-1.1%)	-2.2 (-6.6-2.2)	-33 (-98-33)
Malta	2010	Female	80 to 84	63% (41-81%)	37% (19-59%)	1.2 (0.72-1.7)	-1.2% (-3.2-0.62%)	-1.6% (-4.1-0.69%)	-5.1 (-13-2.3)	-54 (-140-28)
Malta	2010	Female	85 to 89	63% (44-80%)	37% (20-56%)	1.2 (0.71-1.8)	-1.4% (-3.2-0.43%)	-1.8% (-4.1-0.46%)	-5.9 (-14-1.5)	-43 (-100-14)
Malta	2010	Female	90 to 94	62% (43-80%)	38% (20-57%)	1.2 (0.65-1.9)	-1.2% (-2.9-0.51%)	-1.7% (-3.9-0.5%)	-3.1 (-7-0.84)	-15 (-36-6.1)
Malta	2010	Female	95 plus	63% (39-83%)	37% (17-61%)	1.1 (0.54-2)	-1.1% (-3-0.86%)	-1.7% (-4.1-0.86%)	-1.2 (-3.1-0.64)	-3.8 (-10-2.9)
Malta	2010	Male	All Ages	83% (75-89%)	17% (11-25%)	2.2 (1.9-2.5)	2.8% (1.4-4.2%)	1.7% (-0.76-4%)	28 (-13-68)	1500 (730-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	2010	Male	15 to 19	73% (53-88%)	27% (12-47%)	1.5 (0.92-2.1)	3.4% (2.3-4.8%)	6.4% (3.2-10%)	0.33 (0.16-0.53)	45 (30-64)
Malta	2010	Male	20 to 24	81% (65-91%)	19% (8.8-35%)	2.3 (1.5-3.1)	6.9% (4.9-9.2%)	9.3% (4.8-14%)	0.83 (0.42-1.3)	120 (82-170)
Malta	2010	Male	25 to 29	85% (73-93%)	15% (6.9-27%)	2.2 (1.5-2.8)	5.7% (4.1-7.5%)	8.2% (4.3-12%)	0.79 (0.41-1.2)	110 (77-150)
Malta	2010	Male	30 to 34	88% (76-95%)	12% (5-24%)	2.1 (1.5-2.9)	4.8% (3.3-6.4%)	7.5% (4.2-11%)	1 (0.56-1.5)	110 (73-150)
Malta	2010	Male	35 to 39	89% (78-96%)	11% (4.1-22%)	2.2 (1.5-3.1)	4.3% (2.9-5.7%)	7.3% (4.1-11%)	1 (0.54-1.5)	89 (59-120)
Malta	2010	Male	40 to 44	90% (78-96%)	10% (3.8-22%)	2.5 (1.8-3.4)	4.2% (2.8-5.8%)	7.3% (3.7-11%)	1.3 (0.68-2)	93 (60-130)
Malta	2010	Male	45 to 49	90% (75-96%)	10% (4-25%)	3.2 (2.3-4.3)	4.7% (2.9-6.9%)	8.1% (4.4-12%)	2.3 (1.3-3.6)	130 (81-190)
Malta	2010	Male	50 to 54	90% (77-96%)	9.7% (3.6-23%)	3.6 (2.6-4.7)	5% (2.8-7.6%)	7.6% (3.6-12%)	3.9 (1.9-6.3)	180 (98-270)
Malta	2010	Male	55 to 59	90% (80-97%)	9.5% (3.3-20%)	3.6 (2.7-4.5)	4.4% (2.1-6.9%)	6% (2.3-9.8%)	4.9 (1.9-8.2)	200 (92-310)
Malta	2010	Male	60 to 64	88% (75-96%)	12% (3.9-25%)	3.3 (2.4-4.3)	3.5% (1-6.2%)	4.3% (0.93-8%)	6.8 (1.4-12)	220 (64-400)
Malta	2010	Male	65 to 69	87% (74-96%)	13% (4.2-26%)	2.9 (2.2-3.7)	2.3% (-0.012-4.5%)	2.6% (-0.46-5.5%)	4.3 (-0.68-9.1)	120 (-0.65-240)
Malta	2010	Male	70 to 74	87% (75-95%)	13% (5.1-25%)	2.6 (1.8-3.5)	1.4% (-0.87-3.7%)	1.4% (-1.5-4.4%)	3.2 (-3.4-9.9)	79 (-49-210)
Malta	2010	Male	75 to 79	85% (70-94%)	15% (6-30%)	2.3 (1.5-3.2)	0.49% (-1.8-3.1%)	0.28% (-2.5-3.4%)	0.68 (-5.7-7.9)	21 (-73-130)
Malta	2010	Male	80 to 84	83% (67-93%)	17% (7.1-33%)	2.1 (1.4-2.9)	-0.016% (-2.4-2.3%)	-0.3% (-3.2-2.5%)	-0.78 (-8.2-6.7)	-0.44 (-80-78)
Malta	2010	Male	85 to 89	82% (67-93%)	18% (7.3-33%)	2 (1.3-2.8)	-0.29% (-2.5-1.9%)	-0.63% (-3.3-1.9%)	-1.5 (-7.7-4.5)	-5.9 (-51-39)
Malta	2010	Male	90 to 94	83% (68-93%)	17% (6.8-32%)	1.9 (1.2-2.8)	-0.32% (-2.4-1.8%)	-0.74% (-3.3-1.9%)	-0.75 (-3.3-1.9)	-2.1 (-15-12)
Malta	2010	Male	95 plus	83% (65-95%)	17% (5.3-35%)	1.9 (1-3)	-0.32% (-2.4-2.1%)	-0.82% (-3.3-2.1%)	-0.26 (-1.1-0.7)	-0.45 (-3.2-2.9)
Malta	2016	Female	All Ages	74% (66-81%)	26% (19-34%)	1.3 (0.92-1.7)	0.6% (-0.42-1.8%)	-0.21% (-2.3-2.1%)	-3.3 (-38-36)	300 (-220-920)
Malta	2016	Female	15 to 19	70% (51-85%)	30% (15-49%)	1.1 (0.64-1.7)	1.1% (0.71-1.6%)	3.7% (1.5-6.2%)	0.059 (0.023-0.1)	14 (8.7-21)
Malta	2016	Female	20 to 24	75% (57-88%)	25% (12-43%)	1.4 (0.86-2.1)	1.9% (1.3-2.6%)	3.9% (1.9-6.5%)	0.11 (0.047-0.19)	31 (20-47)
Malta	2016	Female	25 to 29	78% (62-90%)	22% (10-38%)	1.3 (0.83-1.9)	1.6% (1.1-2.2%)	3.4% (1.5-5.5%)	0.12 (0.053-0.2)	24 (15-34)
Malta	2016	Female	30 to 34	79% (62-91%)	21% (9.1-38%)	1.3 (0.76-1.9)	1.3% (0.81-1.8%)	4.2% (2.4-6.4%)	0.11 (0.058-0.18)	18 (11-27)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	2016	Female	35 to 39	81% (64-91%)	19% (8.8-36%)	1.3 (0.8-2)	1.2% (0.77-1.8%)	5.2% (3.1-7.7%)	0.27 (0.15-0.45)	28 (18-41)
Malta	2016	Female	40 to 44	81% (65-92%)	19% (8.1-35%)	1.4 (0.85-2.1)	1.4% (0.8-2.2%)	5.8% (3.5-8.9%)	0.53 (0.29-0.86)	35 (20-54)
Malta	2016	Female	45 to 49	81% (63-92%)	19% (8.3-37%)	1.5 (0.88-2.3)	1.6% (0.82-2.6%)	5.8% (3.5-8.7%)	0.81 (0.45-1.3)	39 (21-65)
Malta	2016	Female	50 to 54	80% (63-91%)	20% (8.6-37%)	1.6 (0.94-2.4)	1.8% (0.78-3.1%)	5.4% (3-8.4%)	1.4 (0.7-2.2)	53 (23-90)
Malta	2016	Female	55 to 59	80% (63-91%)	20% (8.7-37%)	1.8 (1.1-2.7)	1.8% (0.52-3.4%)	4.8% (2.2-7.6%)	2 (0.88-3.4)	63 (19-120)
Malta	2016	Female	60 to 64	78% (59-91%)	22% (8.6-41%)	1.8 (1.1-2.8)	1.5% (0.011-3.3%)	3.3% (0.78-6.2%)	2.2 (0.51-4.4)	57 (0.48-130)
Malta	2016	Female	65 to 69	76% (56-90%)	24% (9.8-44%)	1.7 (1.1-2.6)	0.75% (-0.65-2.4%)	1.8% (-0.52-4.3%)	1.9 (-0.55-4.6)	37 (-32-120)
Malta	2016	Female	70 to 74	74% (55-87%)	26% (13-45%)	1.6 (0.97-2.5)	0.21% (-1.4-2.3%)	0.58% (-2.1-3.5%)	0.91 (-3.1-5.4)	11 (-72-110)
Malta	2016	Female	75 to 79	70% (50-86%)	30% (14-50%)	1.5 (0.87-2.4)	-0.54% (-2.5-1.8%)	-0.78% (-3.5-2.2%)	-2.1 (-9.9-6.1)	-31 (-150-100)
Malta	2016	Female	80 to 84	68% (46-84%)	32% (16-54%)	1.4 (0.82-2.2)	-0.85% (-2.9-1.4%)	-1.2% (-3.8-1.7%)	-3.4 (-11-4.9)	-35 (-120-60)
Malta	2016	Female	85 to 89	65% (45-81%)	35% (19-55%)	1.3 (0.74-2.1)	-0.93% (-2.9-1%)	-1.3% (-3.7-1.2%)	-3.6 (-10-3.2)	-25 (-77-28)
Malta	2016	Female	90 to 94	64% (44-81%)	36% (19-56%)	1.3 (0.69-2.1)	-0.91% (-2.8-1.1%)	-1.4% (-3.8-1.1%)	-3.1 (-8.5-2.4)	-14 (-43-16)
Malta	2016	Female	95 plus	64% (40-84%)	36% (16-60%)	1.3 (0.56-2.4)	-0.88% (-2.8-1.2%)	-1.4% (-4-1.3%)	-1.5 (-4.1-1.3)	-4.1 (-13-5.8)
Malta	2016	Male	All Ages	82% (74-88%)	18% (12-26%)	2.4 (1.7-3.2)	3.3% (1.3-5.7%)	2.3% (-0.46-5.8%)	43 (-8.9-110)	1800 (780-3200)
Malta	2016	Male	15 to 19	72% (51-87%)	28% (13-49%)	1.6 (0.87-2.5)	3.4% (2.3-4.8%)	6.5% (3.1-11%)	0.27 (0.13-0.47)	40 (26-58)
Malta	2016	Male	20 to 24	78% (59-90%)	22% (9.7-41%)	2.4 (1.4-3.5)	7.2% (5.1-9.6%)	9.6% (5-15%)	0.69 (0.33-1.2)	110 (72-150)
Malta	2016	Male	25 to 29	81% (67-92%)	19% (7.8-33%)	2.3 (1.4-3.3)	6% (4.4-7.9%)	8.4% (4.4-14%)	0.71 (0.34-1.2)	110 (72-150)
Malta	2016	Male	30 to 34	84% (69-94%)	16% (5.9-31%)	2.2 (1.4-3.3)	5% (3.5-6.8%)	7.9% (4.2-12%)	0.91 (0.45-1.5)	100 (69-150)
Malta	2016	Male	35 to 39	86% (73-95%)	14% (5.2-27%)	2.3 (1.4-3.5)	4.5% (3-6.2%)	7.9% (4.2-12%)	1.1 (0.59-1.8)	100 (69-150)
Malta	2016	Male	40 to 44	88% (74-96%)	12% (4.4-26%)	2.7 (1.7-4)	4.7% (2.9-6.8%)	8.4% (4.7-13%)	1.6 (0.81-2.5)	110 (69-170)
Malta	2016	Male	45 to 49	89% (73-96%)	11% (4.1-27%)	3.5 (2.1-5.2)	5.5% (3.2-8.4%)	9.8% (5.3-16%)	2.3 (1.2-3.8)	130 (76-210)
Malta	2016	Male	50 to 54	89% (76-96%)	11% (4.1-24%)	3.8 (2.4-5.6)	5.9% (3.2-9.3%)	9.3% (4.7-15%)	4.4 (2.1-7.4)	200 (110-320)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malta	2016	Male	55 to 59	89% (77-96%)	11% (3.5-23%)	3.8 (2.4-5.6)	5.3% (2.4-9%)	7.6% (3.2-13%)	6 (2.3-11)	230 (110-400)
Malta	2016	Male	60 to 64	88% (72-96%)	12% (4-28%)	3.6 (2.2-5.3)	4.4% (1.3-8.3%)	5.7% (1.4-11%)	7 (1.7-14)	230 (69-440)
Malta	2016	Male	65 to 69	88% (72-96%)	12% (4.3-28%)	3.2 (2.1-4.6)	3.1% (0.55-6.7%)	3.7% (0.22-8.3%)	8 (0.53-18)	220 (41-470)
Malta	2016	Male	70 to 74	87% (73-95%)	13% (5.3-27%)	2.7 (1.8-4.2)	2% (-0.28-5.6%)	2.3% (-0.72-6.7%)	5.2 (-1.6-15)	120 (-16-330)
Malta	2016	Male	75 to 79	85% (71-94%)	15% (5.6-29%)	2.4 (1.5-3.8)	1.2% (-1.2-4.8%)	1.1% (-1.7-5.3%)	4 (-6-19)	79 (-72-300)
Malta	2016	Male	80 to 84	84% (66-93%)	16% (6.9-34%)	2.3 (1.4-3.5)	0.64% (-1.9-3.8%)	0.41% (-2.6-4%)	1.3 (-8-13)	27 (-75-160)
Malta	2016	Male	85 to 89	82% (65-92%)	18% (7.7-35%)	2.1 (1.3-3.2)	0.32% (-1.9-3%)	0.025% (-2.7-3.1%)	0.065 (-5.7-7.1)	6.2 (-38-61)
Malta	2016	Male	90 to 94	82% (66-93%)	18% (6.5-34%)	2.1 (1.2-3.2)	0.15% (-2.1-2.9%)	-0.2% (-2.9-3.1%)	-0.26 (-3.7-4.1)	1.3 (-17-23)
Malta	2016	Male	95 plus	82% (64-95%)	18% (5.1-36%)	2 (0.99-3.4)	0.15% (-2.1-3.3%)	-0.29% (-3.3-6%)	-0.12 (-1.3-1.5)	0.27 (-3.8-5.9)
Netherlands	1990	Female	All Ages	79% (76-83%)	21% (17-24%)	1.8 (1.7-2)	2.7% (1.6-3.8%)	2.9% (0.86-5.1%)	1800 (530-3200)	52000 (31000-73000)
Netherlands	1990	Female	15 to 19	77% (65-87%)	23% (13-35%)	1.6 (1.2-2.2)	2% (1.2-2.9%)	7.6% (3.9-12%)	11 (5.2-16)	1100 (660-1600)
Netherlands	1990	Female	20 to 24	81% (71-88%)	19% (12-29%)	1.7 (1.2-2.3)	2.3% (1.5-3.3%)	7.5% (3.6-12%)	14 (6.9-23)	1800 (1200-2500)
Netherlands	1990	Female	25 to 29	83% (73-89%)	17% (11-27%)	1.4 (0.98-1.9)	1.8% (1.1-2.6%)	5.8% (2.1-9.5%)	14 (5-23)	1600 (950-2200)
Netherlands	1990	Female	30 to 34	83% (73-90%)	17% (10-27%)	1.5 (1.1-2)	1.9% (1.1-2.8%)	6.6% (3.2-10%)	22 (11-34)	1800 (1100-2500)
Netherlands	1990	Female	35 to 39	84% (76-90%)	16% (10-24%)	2.2 (1.6-2.9)	2.9% (1.9-4%)	9.8% (6.3-13%)	45 (29-61)	2800 (1900-3700)
Netherlands	1990	Female	40 to 44	84% (76-90%)	16% (9.6-24%)	2.7 (2.1-3.3)	3.9% (2.7-5.2%)	11% (8.5-14%)	86 (64-110)	4500 (3400-5900)
Netherlands	1990	Female	45 to 49	84% (76-90%)	16% (10-24%)	3 (2.3-3.7)	4.7% (3.4-6.2%)	11% (8.6-14%)	110 (81-140)	4900 (3700-6200)
Netherlands	1990	Female	50 to 54	85% (78-90%)	15% (9.7-22%)	3.1 (2.4-3.9)	5% (3.6-6.7%)	11% (8-13%)	130 (100-170)	5300 (3900-6700)
Netherlands	1990	Female	55 to 59	87% (82-92%)	13% (8.3-18%)	3.2 (2.5-3.9)	4.9% (3.3-6.7%)	8.9% (6.4-12%)	170 (120-220)	5800 (4000-7800)
Netherlands	1990	Female	60 to 64	87% (81-92%)	13% (8.4-19%)	3.1 (2.4-3.9)	4.2% (2.4-6.3%)	6.8% (4.1-10%)	190 (120-280)	5800 (3300-8500)
Netherlands	1990	Female	65 to 69	84% (78-90%)	16% (10-22%)	2.8 (2.2-3.4)	3.4% (1.5-5.4%)	5.1% (2.3-8.1%)	220 (100-350)	5500 (2500-8700)
Netherlands	1990	Female	70 to 74	81% (73-87%)	19% (13-27%)	2.5 (1.8-3.2)	2.7% (0.51-5%)	3.8% (0.73-7%)	220 (43-410)	4500 (810-8300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	1990	Female	75 to 79	77% (65-85%)	23% (15-35%)	2.2 (1.5-2.9)	1.9% (-0.29-4.3%)	2.4% (-0.47-5.6%)	220 (-43-500)	3500 (-550-7800)
Netherlands	1990	Female	80 to 84	74% (61-83%)	26% (17-39%)	1.9 (1.4-2.6)	1.3% (-0.59-3.6%)	1.5% (-0.84-4.3%)	190 (-100-510)	2200 (-970-5800)
Netherlands	1990	Female	85 to 89	72% (60-81%)	28% (19-40%)	1.8 (1.2-2.5)	1.1% (-0.72-3%)	1.1% (-1-3.4%)	140 (-120-410)	1200 (-800-3300)
Netherlands	1990	Female	90 to 94	71% (58-82%)	29% (18-42%)	1.7 (1.1-2.4)	0.67% (-1.3-2.8%)	0.65% (-1.6-3.1%)	45 (-110-220)	290 (-530-1200)
Netherlands	1990	Female	95 plus	71% (54-84%)	29% (16-46%)	1.6 (0.9-2.5)	0.46% (-1.8-3%)	0.36% (-2.3-3.3%)	9.1 (-57-84)	48 (-180-300)
Netherlands	1990	Male	All Ages	87% (84-90%)	13% (10-16%)	3.5 (3.2-3.8)	6.4% (4.7-8.1%)	6.7% (4.1-9.2%)	4400 (2700-6200)	140000 (100000-170000)
Netherlands	1990	Male	15 to 19	80% (70-87%)	20% (13-30%)	2.7 (2-3.5)	5.5% (3.4-7.8%)	13% (7.1-18%)	37 (20-54)	3300 (2100-4600)
Netherlands	1990	Male	20 to 24	87% (79-92%)	13% (7.8-21%)	4.2 (3.3-5.2)	10% (7.3-14%)	20% (13-27%)	89 (57-120)	8000 (5700-10000)
Netherlands	1990	Male	25 to 29	89% (82-93%)	11% (6.6-18%)	3.7 (2.9-4.5)	8.5% (6-11%)	17% (11-23%)	85 (55-120)	7300 (5200-9500)
Netherlands	1990	Male	30 to 34	91% (85-95%)	9% (5.2-15%)	3.6 (2.8-4.5)	7.1% (5-9.3%)	15% (9.1-20%)	82 (52-110)	6400 (4300-8400)
Netherlands	1990	Male	35 to 39	93% (89-96%)	7.1% (4.1-11%)	4.1 (3.2-5)	7% (5-9.1%)	14% (8.9-18%)	99 (64-130)	6700 (4800-8700)
Netherlands	1990	Male	40 to 44	94% (90-97%)	5.8% (3.1-10%)	4.5 (3.6-5.4)	7.3% (5-9.7%)	13% (8.2-18%)	150 (97-210)	8800 (6000-12000)
Netherlands	1990	Male	45 to 49	95% (91-97%)	5.2% (3.1-8.7%)	4.9 (4-6)	7.8% (5.1-11%)	12% (7.9-17%)	180 (110-260)	9100 (6100-12000)
Netherlands	1990	Male	50 to 54	95% (92-97%)	4.9% (3-7.8%)	5.4 (4.4-6.3)	8.4% (5.5-11%)	12% (7.7-16%)	260 (170-360)	11000 (7400-15000)
Netherlands	1990	Male	55 to 59	95% (92-97%)	5.2% (3.3-7.9%)	5.9 (4.9-6.9)	9.3% (6.1-12%)	12% (7.6-16%)	430 (270-570)	15000 (10000-20000)
Netherlands	1990	Male	60 to 64	94% (91-96%)	6% (3.8-8.9%)	5.7 (4.7-6.8)	8.6% (5.3-12%)	10% (6.1-15%)	560 (330-770)	17000 (10000-23000)
Netherlands	1990	Male	65 to 69	93% (89-95%)	7.2% (4.5-11%)	4.9 (4.1-5.7)	7% (4-9.9%)	8.1% (4.3-12%)	650 (350-930)	16000 (9200-23000)
Netherlands	1990	Male	70 to 74	91% (87-94%)	8.7% (5.9-13%)	4.3 (3.4-5.1)	5.5% (2.6-8.4%)	6.1% (2.6-9.5%)	580 (250-910)	12000 (5400-18000)
Netherlands	1990	Male	75 to 79	89% (84-94%)	11% (6.4-16%)	3.8 (2.9-4.7)	4.4% (1.7-7.4%)	4.7% (1.6-8%)	520 (170-900)	8100 (3300-14000)
Netherlands	1990	Male	80 to 84	88% (81-93%)	12% (7-19%)	3.4 (2.6-4.3)	3.8% (1.4-6.4%)	4% (1.1-7%)	400 (110-690)	4600 (1700-7800)
Netherlands	1990	Male	85 to 89	87% (80-92%)	13% (7.8-20%)	3.2 (2.3-4.1)	3.2% (0.98-5.7%)	3.4% (0.76-6.1%)	220 (50-410)	1900 (570-3300)
Netherlands	1990	Male	90 to 94	87% (80-92%)	13% (7.6-20%)	3 (2.1-4)	2.6% (0.32-5.5%)	2.6% (-0.092-6%)	79 (-2.7-180)	470 (58-980)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	1990	Male	95 plus	88% (79-94%)	12% (5.7-21%)	2.9 (1.9-4.1)	2.3% (-0.48-5.8%)	2.2% (-1-6.3%)	20 (-9.1-58)	83 (-17-210)
Netherlands	1995	Female	All Ages	78% (75-82%)	22% (18-25%)	1.8 (1.6-1.9)	2.7% (1.7-3.8%)	3.1% (1.2-5.1%)	2100 (790-3400)	56000 (36000-76000)
Netherlands	1995	Female	15 to 19	79% (69-88%)	21% (12-31%)	1.7 (1.2-2.2)	1.9% (1.2-2.9%)	7.6% (4-12%)	8.4 (4.2-13)	910 (560-1300)
Netherlands	1995	Female	20 to 24	81% (72-87%)	19% (13-28%)	1.7 (1.2-2.3)	2.2% (1.4-3.1%)	7.3% (3.6-11%)	12 (5.7-19)	1500 (980-2100)
Netherlands	1995	Female	25 to 29	81% (72-88%)	19% (12-28%)	1.3 (0.95-1.7)	1.7% (1.1-2.4%)	5.4% (2.2-8.7%)	13 (5-21)	1500 (920-2200)
Netherlands	1995	Female	30 to 34	82% (73-89%)	18% (11-27%)	1.4 (1.1-1.9)	1.7% (1.1-2.5%)	6% (3-9%)	20 (10-30)	1700 (1100-2400)
Netherlands	1995	Female	35 to 39	83% (76-89%)	17% (11-24%)	2.1 (1.6-2.8)	2.7% (1.8-3.8%)	9.2% (5.8-12%)	47 (30-63)	3000 (2000-3900)
Netherlands	1995	Female	40 to 44	84% (77-90%)	16% (10-23%)	2.6 (2.1-3.2)	3.8% (2.7-5%)	11% (7.8-14%)	84 (62-110)	4500 (3300-5700)
Netherlands	1995	Female	45 to 49	83% (77-88%)	17% (12-23%)	2.9 (2.3-3.6)	4.6% (3.3-6.1%)	11% (8.7-14%)	140 (110-170)	6300 (4800-7900)
Netherlands	1995	Female	50 to 54	84% (78-89%)	16% (11-22%)	3 (2.4-3.7)	5.1% (3.8-6.6%)	11% (8.5-13%)	150 (120-190)	6100 (4600-7600)
Netherlands	1995	Female	55 to 59	85% (80-90%)	15% (10-20%)	3 (2.4-3.6)	4.8% (3.5-6.5%)	8.9% (6.9-11%)	170 (130-210)	5800 (4400-7600)
Netherlands	1995	Female	60 to 64	83% (77-89%)	17% (11-23%)	2.6 (2-3.3)	4% (2.6-5.7%)	6.6% (4.4-9%)	180 (120-260)	5400 (3500-7700)
Netherlands	1995	Female	65 to 69	78% (71-86%)	22% (14-29%)	2.3 (1.8-3)	3.2% (1.7-4.8%)	4.8% (2.6-7%)	210 (110-300)	5100 (2600-7600)
Netherlands	1995	Female	70 to 74	79% (71-85%)	21% (15-29%)	2.3 (1.7-3)	2.8% (0.89-4.8%)	4% (1.3-6.7%)	260 (87-450)	5200 (1500-8800)
Netherlands	1995	Female	75 to 79	77% (67-86%)	23% (14-33%)	2.2 (1.5-2.9)	2.2% (0.057-4.4%)	2.8% (0.078-5.7%)	250 (7-510)	3800 (110-7900)
Netherlands	1995	Female	80 to 84	74% (63-84%)	26% (16-37%)	1.9 (1.4-2.6)	1.6% (-0.27-3.7%)	1.9% (-0.43-4.5%)	240 (-54-570)	2700 (-480-6400)
Netherlands	1995	Female	85 to 89	72% (60-82%)	28% (18-40%)	1.8 (1.2-2.4)	1.3% (-0.52-3.2%)	1.4% (-0.75-3.8%)	190 (-100-510)	1600 (-660-4000)
Netherlands	1995	Female	90 to 94	72% (60-82%)	28% (18-40%)	1.7 (1.1-2.4)	0.88% (-1.1-3%)	0.92% (-1.3-3.4%)	79 (-120-300)	470 (-570-1600)
Netherlands	1995	Female	95 plus	72% (57-84%)	28% (16-43%)	1.6 (0.95-2.5)	0.69% (-1.5-3.3%)	0.64% (-1.9-3.6%)	20 (-62-120)	89 (-190-430)
Netherlands	1995	Male	All Ages	89% (86-91%)	11% (9.2-14%)	3.4 (3.2-3.7)	6.4% (4.7-7.9%)	6.6% (4.1-8.9%)	4500 (2800-6100)	140000 (100000-170000)
Netherlands	1995	Male	15 to 19	81% (74-87%)	19% (13-26%)	2.7 (2.1-3.3)	5.5% (3.4-7.7%)	13% (6.9-18%)	31 (17-44)	2800 (1700-3900)
Netherlands	1995	Male	20 to 24	89% (83-93%)	11% (7.4-17%)	4.1 (3.3-5.1)	10% (7.1-13%)	20% (13-27%)	76 (48-100)	6900 (4800-8900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	1995	Male	25 to 29	90% (86-93%)	9.5% (6.5-14%)	3.6 (2.9-4.4)	8% (5.6-10%)	17% (11-23%)	75 (47-100)	6700 (4700-8800)
Netherlands	1995	Male	30 to 34	92% (88-95%)	7.9% (5-12%)	3.5 (2.8-4.4)	6.8% (4.6-9%)	14% (8.5-19%)	85 (51-120)	6600 (4500-8700)
Netherlands	1995	Male	35 to 39	94% (91-96%)	5.9% (3.6-8.9%)	4 (3.2-4.8)	6.7% (4.7-8.7%)	13% (8.7-17%)	100 (65-130)	6900 (4900-8800)
Netherlands	1995	Male	40 to 44	95% (92-97%)	4.8% (2.7-7.8%)	4.5 (3.6-5.3)	7.4% (5.2-9.7%)	13% (9-18%)	150 (99-200)	8500 (6000-11000)
Netherlands	1995	Male	45 to 49	96% (93-97%)	4.3% (2.9-6.5%)	4.9 (4.1-5.7)	8% (5.6-11%)	13% (8.9-17%)	240 (160-320)	12000 (8400-15000)
Netherlands	1995	Male	50 to 54	96% (94-97%)	4.2% (2.8-6.1%)	5.3 (4.4-6.1)	8.6% (6.1-11%)	13% (8.8-16%)	290 (200-370)	12000 (8800-15000)
Netherlands	1995	Male	55 to 59	95% (94-97%)	4.6% (3.2-6.2%)	5.8 (4.9-6.7)	9.3% (6.6-12%)	13% (8.6-16%)	410 (280-530)	15000 (10000-19000)
Netherlands	1995	Male	60 to 64	94% (93-96%)	5.5% (3.9-7.4%)	5.5 (4.7-6.6)	8.6% (5.5-11%)	11% (6.4-14%)	550 (330-750)	17000 (10000-22000)
Netherlands	1995	Male	65 to 69	93% (90-95%)	7.2% (5-9.7%)	4.7 (4-5.4)	6.9% (4.4-9.3%)	8% (4.8-11%)	610 (360-840)	15000 (9400-20000)
Netherlands	1995	Male	70 to 74	91% (87-94%)	8.7% (6-13%)	4 (3.2-4.8)	5.3% (2.7-7.9%)	5.9% (2.8-9%)	610 (280-920)	12000 (6300-18000)
Netherlands	1995	Male	75 to 79	90% (84-94%)	10% (6.2-16%)	3.6 (2.8-4.5)	4.2% (1.6-6.7%)	4.5% (1.5-7.5%)	490 (170-820)	7800 (3000-12000)
Netherlands	1995	Male	80 to 84	88% (82-93%)	12% (6.8-18%)	3.3 (2.6-4.1)	3.7% (1.3-6%)	3.9% (1.1-6.5%)	420 (120-710)	4800 (1700-7900)
Netherlands	1995	Male	85 to 89	87% (80-92%)	13% (7.7-20%)	3.1 (2.3-3.9)	3.2% (0.98-5.6%)	3.3% (0.84-6%)	240 (61-440)	2000 (640-3500)
Netherlands	1995	Male	90 to 94	88% (80-93%)	12% (7.3-20%)	3 (2.1-3.9)	2.6% (0.3-5.3%)	2.7% (-0.037-5.7%)	86 (-1.2-180)	500 (57-1000)
Netherlands	1995	Male	95 plus	88% (79-94%)	12% (5.7-21%)	2.8 (1.8-4)	2.4% (-0.2-5.8%)	2.4% (-0.66-6.3%)	21 (-5.8-57)	85 (-7.1-210)
Netherlands	2000	Female	All Ages	80% (76-83%)	20% (17-24%)	1.8 (1.7-2)	2.7% (1.8-3.7%)	3% (1.4-4.7%)	2200 (970-3500)	58000 (39000-77000)
Netherlands	2000	Female	15 to 19	79% (68-87%)	21% (13-32%)	1.6 (1.2-2.1)	1.8% (1.1-2.6%)	7.2% (3.8-11%)	7.8 (4.1-12)	860 (520-1300)
Netherlands	2000	Female	20 to 24	81% (72-88%)	19% (12-28%)	1.6 (1.1-2.2)	2.1% (1.4-2.9%)	6.9% (3.4-11%)	8.8 (4.3-14)	1200 (780-1700)
Netherlands	2000	Female	25 to 29	82% (73-89%)	18% (11-27%)	1.3 (0.92-1.7)	1.6% (1-2.2%)	5.2% (2.3-8.4%)	9.9 (4.3-16)	1200 (800-1700)
Netherlands	2000	Female	30 to 34	83% (74-89%)	17% (11-26%)	1.4 (1-1.8)	1.6% (1-2.3%)	5.8% (3-8.6%)	18 (9-26)	1600 (1000-2200)
Netherlands	2000	Female	35 to 39	84% (77-90%)	16% (10-23%)	2.1 (1.5-2.7)	2.5% (1.6-3.5%)	8.9% (5.8-12%)	45 (29-61)	2900 (1900-3800)
Netherlands	2000	Female	40 to 44	85% (78-90%)	15% (9.8-22%)	2.6 (2.1-3.2)	3.6% (2.4-4.8%)	10% (7.6-13%)	84 (62-110)	4400 (3200-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	2000	Female	45 to 49	85% (79-89%)	15% (11-21%)	2.9 (2.3-3.6)	4.6% (3.3-6.1%)	11% (8.4-14%)	140 (110-180)	6300 (4700-8000)
Netherlands	2000	Female	50 to 54	86% (81-90%)	14% (10-19%)	3 (2.4-3.7)	5.1% (3.7-6.6%)	11% (8.8-13%)	200 (160-250)	8000 (6200-10000)
Netherlands	2000	Female	55 to 59	88% (84-91%)	12% (8.6-16%)	3.1 (2.5-3.7)	5% (3.7-6.6%)	9.5% (7.6-12%)	200 (160-250)	6800 (5100-8700)
Netherlands	2000	Female	60 to 64	87% (83-91%)	13% (9.1-17%)	2.9 (2.3-3.5)	4.3% (2.8-6%)	7.3% (5.1-9.5%)	200 (140-270)	5900 (3900-8100)
Netherlands	2000	Female	65 to 69	82% (76-88%)	18% (12-24%)	2.5 (2-3.1)	3.3% (1.9-4.9%)	5.2% (3.2-7.2%)	210 (130-300)	5100 (2900-7500)
Netherlands	2000	Female	70 to 74	80% (73-85%)	20% (15-27%)	2.3 (1.7-2.9)	2.8% (1.4-6%)	4% (1.7-6.6%)	250 (110-420)	4900 (1800-8200)
Netherlands	2000	Female	75 to 79	77% (67-84%)	23% (16-33%)	2 (1.5-2.7)	2.1% (0.19-4.1%)	2.7% (0.35-5.3%)	270 (35-550)	4200 (360-8500)
Netherlands	2000	Female	80 to 84	73% (62-83%)	27% (17-38%)	1.8 (1.3-2.5)	1.5% (-0.26-3.3%)	1.8% (-0.36-4%)	230 (-46-520)	2600 (-430-5800)
Netherlands	2000	Female	85 to 89	72% (60-81%)	28% (19-40%)	1.7 (1.2-2.3)	1.2% (-0.59-2.9%)	1.3% (-0.77-3.5%)	190 (-110-500)	1600 (-820-3900)
Netherlands	2000	Female	90 to 94	71% (60-81%)	29% (19-40%)	1.6 (1.1-2.3)	0.75% (-1-2.6%)	0.79% (-1.3-3%)	79 (-120-300)	460 (-620-1600)
Netherlands	2000	Female	95 plus	71% (55-84%)	29% (16-45%)	1.5 (0.87-2.3)	0.58% (-1.4-2.9%)	0.53% (-1.8-3.2%)	20 (-68-120)	90 (-210-440)
Netherlands	2000	Male	All Ages	89% (87-91%)	11% (8.5-13%)	3.3 (3.1-3.6)	6.3% (4.8-7.8%)	6.5% (4.3-8.6%)	4500 (3000-6000)	140000 (110000-170000)
Netherlands	2000	Male	15 to 19	82% (76-88%)	18% (12-24%)	2.6 (2-3.3)	5.4% (3.3-7.6%)	13% (6.8-18%)	29 (16-43)	2700 (1700-3700)
Netherlands	2000	Male	20 to 24	89% (84-93%)	11% (6.9-16%)	4 (3.1-4.9)	9.9% (6.8-13%)	19% (12-27%)	63 (40-86)	5700 (3900-7400)
Netherlands	2000	Male	25 to 29	91% (87-94%)	8.8% (5.7-13%)	3.5 (2.8-4.3)	7.7% (5.2-10%)	17% (10-23%)	62 (37-86)	5600 (3800-7200)
Netherlands	2000	Male	30 to 34	93% (89-95%)	7.3% (4.6-11%)	3.4 (2.7-4.2)	6.5% (4.4-8.7%)	15% (8.8-20%)	75 (44-110)	6000 (4000-8000)
Netherlands	2000	Male	35 to 39	94% (92-97%)	5.8% (3.5-8.5%)	3.8 (3.1-4.6)	6.6% (4.7-8.5%)	14% (9.5-19%)	100 (66-140)	6900 (4900-9000)
Netherlands	2000	Male	40 to 44	95% (93-97%)	4.7% (2.6-7.5%)	4.2 (3.5-5)	7.4% (5.4-9.7%)	14% (9.9-19%)	150 (110-210)	8800 (6300-12000)
Netherlands	2000	Male	45 to 49	96% (94-97%)	4.1% (2.7-6.1%)	4.6 (3.9-5.4)	8.2% (5.9-11%)	14% (9.9-18%)	230 (160-300)	11000 (8200-15000)
Netherlands	2000	Male	50 to 54	96% (94-97%)	3.9% (2.5-5.5%)	5 (4.3-5.9)	8.8% (6.4-11%)	13% (9.7-17%)	370 (270-470)	16000 (12000-20000)
Netherlands	2000	Male	55 to 59	96% (94-97%)	4.1% (2.9-5.7%)	5.5 (4.8-6.4)	9.5% (6.9-12%)	13% (9.5-17%)	460 (330-580)	16000 (12000-21000)
Netherlands	2000	Male	60 to 64	95% (93-97%)	5% (3.5-6.7%)	5.4 (4.6-6.3)	8.7% (6-11%)	11% (7.5-15%)	540 (350-710)	16000 (11000-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	2000	Male	65 to 69	94% (91-96%)	6.2% (4.3-8.5%)	4.6 (4-5.3)	6.9% (4.5-9.1%)	8.2% (5.4-11%)	580 (380-790)	15000 (9900-19000)
Netherlands	2000	Male	70 to 74	93% (90-95%)	7.2% (5.3-9.6%)	4.1 (3.4-4.8)	5.2% (2.7-7.6%)	5.9% (2.8-8.7%)	570 (280-850)	12000 (6200-17000)
Netherlands	2000	Male	75 to 79	91% (87-94%)	9.1% (5.8-13%)	3.6 (2.9-4.3)	4% (1.6-6.3%)	4.3% (1.5-7%)	520 (180-850)	8200 (3400-13000)
Netherlands	2000	Male	80 to 84	89% (83-93%)	11% (7-17%)	3.2 (2.5-3.9)	3.4% (1.3-5.6%)	3.6% (1.1-6.1%)	390 (120-660)	4500 (1700-7500)
Netherlands	2000	Male	85 to 89	88% (81-92%)	12% (7.5-19%)	3 (2.2-3.7)	2.9% (1.1-5.1%)	3.1% (0.89-5.5%)	240 (71-430)	2000 (750-3500)
Netherlands	2000	Male	90 to 94	88% (82-93%)	12% (7.1-18%)	2.9 (2-3.7)	2.4% (0.34-4.8%)	2.5% (0.12-5.2%)	85 (4.1-180)	500 (69-1000)
Netherlands	2000	Male	95 plus	88% (80-94%)	12% (5.6-20%)	2.7 (1.8-3.8)	2.2% (-0.2-5.1%)	2.2% (-0.66-5.5%)	21 (-6.4-54)	85 (-7.6-200)
Netherlands	2005	Female	All Ages	81% (78-84%)	19% (16-22%)	1.8 (1.7-2)	2.7% (1.9-3.6%)	3.1% (1.7-4.7%)	2200 (1100-3300)	57000 (40000-74000)
Netherlands	2005	Female	15 to 19	79% (69-87%)	21% (13-31%)	1.5 (1.1-2)	1.5% (0.95-2.2%)	6.8% (3.6-10%)	5.7 (3-8.8)	730 (470-1100)
Netherlands	2005	Female	20 to 24	82% (74-89%)	18% (11-26%)	1.6 (1.1-2.2)	2% (1.3-2.7%)	6.8% (3.1-11%)	7.6 (3.5-12)	1100 (720-1600)
Netherlands	2005	Female	25 to 29	84% (76-90%)	16% (10-24%)	1.3 (0.94-1.7)	1.5% (0.93-2%)	5.3% (2.1-8.5%)	7.1 (2.9-11)	960 (620-1300)
Netherlands	2005	Female	30 to 34	84% (77-90%)	16% (9.8-23%)	1.4 (1-1.9)	1.5% (0.94-2.1%)	6% (3.2-8.8%)	15 (7.5-22)	1300 (800-1800)
Netherlands	2005	Female	35 to 39	84% (77-90%)	16% (9.8-23%)	2 (1.5-2.7)	2.2% (1.4-3.1%)	8.8% (5.8-12%)	37 (25-51)	2400 (1600-3300)
Netherlands	2005	Female	40 to 44	85% (78-91%)	15% (9.2-22%)	2.6 (2.1-3.2)	3.2% (2.2-4.4%)	10% (7.5-13%)	77 (55-98)	4100 (3000-5300)
Netherlands	2005	Female	45 to 49	86% (81-90%)	14% (10-19%)	2.9 (2.3-3.4)	4.2% (3-5.5%)	11% (8.1-13%)	130 (97-160)	5800 (4400-7300)
Netherlands	2005	Female	50 to 54	87% (82-91%)	13% (9.3-18%)	3 (2.4-3.6)	4.8% (3.5-6.2%)	10% (8.3-13%)	180 (140-220)	7100 (5400-8800)
Netherlands	2005	Female	55 to 59	89% (86-92%)	11% (8.1-14%)	3.1 (2.5-3.7)	4.9% (3.6-6.4%)	9.5% (7.6-11%)	240 (190-300)	8300 (6200-10000)
Netherlands	2005	Female	60 to 64	89% (85-92%)	11% (8.2-15%)	3 (2.4-3.7)	4.3% (3-5.9%)	7.7% (5.7-9.7%)	220 (160-280)	6400 (4500-8500)
Netherlands	2005	Female	65 to 69	85% (80-89%)	15% (11-20%)	2.7 (2.2-3.2)	3.4% (2.2-4.8%)	5.7% (4-7.4%)	210 (150-280)	5100 (3200-7000)
Netherlands	2005	Female	70 to 74	82% (76-86%)	18% (14-24%)	2.4 (1.8-3)	2.9% (1.3-4.5%)	4.3% (2.2-6.5%)	250 (120-360)	4800 (2200-7400)
Netherlands	2005	Female	75 to 79	77% (68-84%)	23% (16-32%)	2 (1.5-2.6)	2.1% (0.43-3.8%)	2.9% (0.75-5.1%)	260 (65-450)	3900 (800-6900)
Netherlands	2005	Female	80 to 84	72% (60-81%)	28% (19-40%)	1.7 (1.2-2.3)	1.6% (0.092-3.1%)	2% (0.16-3.9%)	270 (21-520)	3000 (180-5900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	2005	Female	85 to 89	71% (60-80%)	29% (20-40%)	1.6 (1.1-2.2)	1.2% (-0.34-2.9%)	1.4% (-0.41-3.4%)	200 (-55-480)	1600 (-440-3700)
Netherlands	2005	Female	90 to 94	70% (58-80%)	30% (20-42%)	1.5 (0.98-2.1)	0.79% (-0.87-2.5%)	0.86% (-1.1-2.9%)	89 (-110-300)	510 (-550-1600)
Netherlands	2005	Female	95 plus	70% (54-84%)	30% (16-46%)	1.4 (0.82-2.2)	0.61% (-1.2-2.7%)	0.58% (-1.5-3%)	25 (-65-130)	110 (-210-480)
Netherlands	2005	Male	All Ages	90% (88-92%)	10% (8.1-12%)	3.2 (3-3.5)	6% (4.7-7.5%)	6.4% (4.4-8.4%)	4200 (2900-5600)	130000 (99000-160000)
Netherlands	2005	Male	15 to 19	84% (77-89%)	16% (11-23%)	2.5 (2-3.2)	4.4% (2.7-6.1%)	12% (6.4-18%)	21 (11-31)	2100 (1400-3000)
Netherlands	2005	Male	20 to 24	90% (84-93%)	10% (6.6-16%)	3.8 (3-4.7)	8.4% (5.9-11%)	19% (11-26%)	45 (27-62)	4400 (3100-5900)
Netherlands	2005	Male	25 to 29	91% (88-94%)	8.5% (5.8-12%)	3.3 (2.6-4.2)	6.8% (4.8-9%)	16% (9.7-23%)	43 (25-59)	3900 (2700-5100)
Netherlands	2005	Male	30 to 34	93% (90-96%)	6.9% (4.4-10%)	3.3 (2.6-4.1)	5.8% (3.8-8%)	14% (8.2-20%)	54 (31-77)	4400 (2900-5800)
Netherlands	2005	Male	35 to 39	94% (92-97%)	5.5% (3.4-8.2%)	3.6 (2.9-4.4)	6% (4.1-7.9%)	14% (8.9-19%)	85 (53-120)	5900 (4100-7800)
Netherlands	2005	Male	40 to 44	95% (93-97%)	4.6% (2.6-7.4%)	4 (3.3-4.8)	6.8% (4.9-9%)	15% (10-19%)	140 (94-180)	7900 (5700-10000)
Netherlands	2005	Male	45 to 49	96% (94-97%)	4% (2.7-5.9%)	4.4 (3.7-5.3)	7.8% (5.7-10%)	14% (10-18%)	210 (150-270)	10000 (7700-13000)
Netherlands	2005	Male	50 to 54	96% (95-98%)	3.7% (2.5-5.2%)	4.9 (4.1-5.7)	8.6% (6.5-11%)	14% (10-17%)	320 (240-400)	14000 (10000-17000)
Netherlands	2005	Male	55 to 59	96% (95-97%)	3.9% (2.8-5.3%)	5.3 (4.6-6.1)	9.2% (7.1-11%)	14% (10-17%)	510 (400-640)	19000 (15000-23000)
Netherlands	2005	Male	60 to 64	95% (94-97%)	4.7% (3.3-6.3%)	5.2 (4.4-6.1)	8.5% (6.2-11%)	11% (8.3-14%)	530 (390-680)	16000 (12000-20000)
Netherlands	2005	Male	65 to 69	94% (92-96%)	5.9% (4.1-8%)	4.5 (3.9-5.1)	6.7% (4.6-8.6%)	8.3% (5.7-11%)	520 (360-690)	13000 (9400-17000)
Netherlands	2005	Male	70 to 74	93% (91-95%)	7% (5.2-9.1%)	3.9 (3.3-4.6)	5.1% (3-7.1%)	5.9% (3.4-8.4%)	520 (290-730)	11000 (6300-15000)
Netherlands	2005	Male	75 to 79	92% (88-94%)	8.2% (5.5-12%)	3.5 (2.9-4.2)	3.9% (1.9-6%)	4.3% (2-6.8%)	480 (210-750)	7700 (3700-12000)
Netherlands	2005	Male	80 to 84	90% (85-94%)	9.9% (6.2-15%)	3.2 (2.6-3.8)	3.3% (1.3-5.3%)	3.5% (1.2-5.8%)	410 (150-700)	4900 (2000-8000)
Netherlands	2005	Male	85 to 89	89% (84-93%)	11% (7-16%)	3 (2.3-3.6)	2.8% (0.98-4.9%)	2.9% (0.88-5.3%)	230 (70-420)	2000 (700-3500)
Netherlands	2005	Male	90 to 94	89% (83-93%)	11% (6.7-17%)	2.8 (2.1-3.6)	2.3% (0.36-4.6%)	2.3% (0.11-5%)	91 (4.3-190)	540 (84-1100)
Netherlands	2005	Male	95 plus	89% (81-95%)	11% (5.2-19%)	2.7 (1.7-3.7)	2.1% (-0.23-4.9%)	2% (-0.57-5.2%)	21 (-5.8-52)	84 (-8.8-200)
Netherlands	2010	Female	All Ages	82% (79-85%)	18% (15-21%)	1.9 (1.7-2.1)	2.6% (1.8-3.4%)	3.1% (1.7-4.5%)	2200 (1200-3200)	55000 (39000-71000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	2010	Female	15 to 19	81% (71-88%)	19% (12-29%)	1.5 (1.1-2)	1.4% (0.87-2%)	6.6% (3.5-10%)	4.7 (2.5-7.4)	680 (430-970)
Netherlands	2010	Female	20 to 24	84% (76-90%)	16% (10-24%)	1.6 (1.1-2.2)	1.9% (1.3-2.5%)	6.8% (3-11%)	6.1 (2.7-9.8)	1100 (710-1600)
Netherlands	2010	Female	25 to 29	85% (77-90%)	15% (9.7-23%)	1.3 (0.93-1.7)	1.4% (0.93-2%)	5.4% (1.9-8.8%)	6.3 (2.3-10)	940 (590-1300)
Netherlands	2010	Female	30 to 34	85% (78-91%)	15% (9.4-22%)	1.4 (1.1-1.9)	1.4% (0.91-2%)	6.1% (3.1-9.1%)	11 (5.5-16)	1000 (670-1400)
Netherlands	2010	Female	35 to 39	85% (78-90%)	15% (9.7-22%)	2 (1.5-2.6)	2% (1.3-2.9%)	8.8% (5.8-12%)	28 (18-38)	1900 (1300-2600)
Netherlands	2010	Female	40 to 44	86% (79-91%)	14% (9.3-21%)	2.5 (2-3.1)	2.9% (2-3.9%)	10% (7.4-13%)	63 (45-80)	3500 (2500-4600)
Netherlands	2010	Female	45 to 49	86% (81-90%)	14% (10-19%)	2.8 (2.3-3.5)	3.7% (2.6-5%)	10% (7.9-13%)	110 (85-140)	5200 (3800-6700)
Netherlands	2010	Female	50 to 54	87% (82-91%)	13% (9.5-18%)	3 (2.4-3.6)	4.4% (3.2-5.7%)	9.8% (7.8-12%)	170 (130-200)	6700 (5000-8400)
Netherlands	2010	Female	55 to 59	89% (86-92%)	11% (7.8-14%)	3.1 (2.5-3.8)	4.6% (3.4-5.9%)	9% (7.3-11%)	210 (170-260)	7300 (5500-9300)
Netherlands	2010	Female	60 to 64	89% (86-92%)	11% (7.9-14%)	3 (2.4-3.7)	4.1% (2.8-5.5%)	7.5% (5.9-9.3%)	260 (200-320)	7600 (5300-10000)
Netherlands	2010	Female	65 to 69	86% (82-90%)	14% (9.8-18%)	2.7 (2.2-3.2)	3.4% (2.2-4.7%)	5.9% (4.3-7.4%)	230 (170-290)	5500 (3700-7400)
Netherlands	2010	Female	70 to 74	83% (78-87%)	17% (13-22%)	2.4 (1.9-3)	2.7% (1.4-4.2%)	4.4% (2.6-6.3%)	230 (140-330)	4500 (2300-6700)
Netherlands	2010	Female	75 to 79	78% (70-85%)	22% (15-30%)	2 (1.5-2.6)	2.1% (0.59-3.6%)	3% (1.2-4.9%)	240 (91-390)	3600 (1000-6200)
Netherlands	2010	Female	80 to 84	74% (63-83%)	26% (17-37%)	1.8 (1.3-2.4)	1.6% (0.19-3.1%)	2.1% (0.42-4%)	260 (54-490)	2800 (350-5500)
Netherlands	2010	Female	85 to 89	73% (62-82%)	27% (18-38%)	1.7 (1.2-2.3)	1.2% (-0.26-2.7%)	1.5% (-0.24-3.2%)	220 (-36-500)	1800 (-370-4000)
Netherlands	2010	Female	90 to 94	72% (61-82%)	28% (18-39%)	1.6 (1-2.2)	0.79% (-0.75-2.4%)	0.89% (-0.9-2.8%)	96 (-100-300)	550 (-510-1700)
Netherlands	2010	Female	95 plus	72% (57-84%)	28% (16-43%)	1.5 (0.87-2.2)	0.6% (-1.2-2.7%)	0.58% (-1.4-3%)	30 (-74-150)	130 (-250-570)
Netherlands	2010	Male	All Ages	90% (88-92%)	10% (8.1-12%)	3.2 (2.9-3.5)	5.8% (4.5-7.2%)	6.2% (4.4-8%)	4100 (2900-5400)	120000 (95000-150000)
Netherlands	2010	Male	15 to 19	84% (77-89%)	16% (11-23%)	2.5 (1.9-3.1)	3.8% (2.5-5.4%)	11% (6.1-17%)	17 (8.8-25)	1900 (1200-2600)
Netherlands	2010	Male	20 to 24	89% (84-93%)	11% (6.9-16%)	3.7 (2.8-4.6)	7.8% (5.5-10%)	18% (11-25%)	41 (24-58)	4200 (2900-5600)
Netherlands	2010	Male	25 to 29	91% (87-94%)	9% (5.9-13%)	3.2 (2.5-4)	6.3% (4.3-8.2%)	16% (8.6-22%)	35 (20-50)	3500 (2400-4600)
Netherlands	2010	Male	30 to 34	93% (89-95%)	7.1% (4.5-11%)	3.1 (2.4-4)	5.3% (3.6-7.2%)	14% (7.4-20%)	39 (21-56)	3300 (2200-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	2010	Male	35 to 39	94% (92-97%)	5.5% (3.4-8%)	3.5 (2.8-4.3)	5.6% (3.8-7.4%)	14% (8.5-19%)	63 (40-88)	4500 (3200-6000)
Netherlands	2010	Male	40 to 44	95% (93-97%)	4.6% (2.6-7.4%)	3.9 (3.2-4.7)	6.2% (4.4-8.3%)	14% (9.8-19%)	110 (77-150)	6700 (4900-8800)
Netherlands	2010	Male	45 to 49	96% (94-97%)	4.1% (2.7-6.1%)	4.3 (3.6-5.1)	7.2% (5.3-9.4%)	14% (10-18%)	190 (140-250)	9500 (7100-12000)
Netherlands	2010	Male	50 to 54	96% (95-98%)	3.7% (2.5-5.2%)	4.7 (4-5.5)	8.2% (6.2-10%)	14% (10-17%)	290 (220-370)	13000 (9700-16000)
Netherlands	2010	Male	55 to 59	96% (95-97%)	3.9% (2.8-5.3%)	5.1 (4.4-6)	9% (7-11%)	14% (11-17%)	450 (350-550)	16000 (13000-20000)
Netherlands	2010	Male	60 to 64	95% (94-97%)	4.7% (3.3-6.3%)	5 (4.2-5.9)	8.3% (6.3-10%)	12% (8.6-14%)	610 (460-760)	19000 (14000-24000)
Netherlands	2010	Male	65 to 69	94% (92-96%)	6% (4.2-8.1%)	4.3 (3.7-5)	6.6% (4.7-8.4%)	8.5% (6-11%)	530 (380-680)	14000 (9900-18000)
Netherlands	2010	Male	70 to 74	93% (91-95%)	7% (5.3-9.2%)	3.8 (3.1-4.5)	5% (3.2-6.8%)	6.1% (3.8-8.3%)	480 (310-650)	10000 (6300-14000)
Netherlands	2010	Male	75 to 79	92% (88-95%)	8% (5.3-12%)	3.4 (2.7-4.1)	3.9% (2.2-5.7%)	4.4% (2.4-6.6%)	470 (250-690)	7600 (4200-11000)
Netherlands	2010	Male	80 to 84	90% (86-94%)	9.5% (6-14%)	3.1 (2.5-3.8)	3.2% (1.5-5%)	3.4% (1.5-5.5%)	400 (170-640)	4800 (2300-7500)
Netherlands	2010	Male	85 to 89	90% (85-93%)	10% (6.8-15%)	2.9 (2.3-3.6)	2.6% (0.91-4.4%)	2.7% (0.8-4.8%)	260 (77-460)	2300 (820-3800)
Netherlands	2010	Male	90 to 94	90% (84-94%)	10% (6.4-16%)	2.8 (2-3.6)	2.1% (0.22-4.2%)	2.1% (-0.11-4.6%)	93 (-4.8-200)	560 (58-1100)
Netherlands	2010	Male	95 plus	89% (81-95%)	11% (5.3-19%)	2.6 (1.7-3.5)	1.9% (-0.27-4.4%)	1.8% (-0.71-4.7%)	23 (-8.9-59)	96 (-14-220)
Netherlands	2016	Female	All Ages	82% (78-84%)	18% (16-22%)	1.9 (1.5-2.5)	2.7% (1.6-4%)	3.3% (1.4-5.4%)	2400 (1100-4000)	58000 (36000-85000)
Netherlands	2016	Female	15 to 19	78% (67-87%)	22% (13-33%)	1.5 (0.96-2.1)	1.3% (0.83-1.9%)	6.5% (3.3-10%)	4.2 (2.1-6.8)	630 (390-940)
Netherlands	2016	Female	20 to 24	82% (72-89%)	18% (11-28%)	1.6 (1-2.3)	1.8% (1.2-2.6%)	6.8% (2.7-11%)	6.2 (2.5-10)	1100 (740-1700)
Netherlands	2016	Female	25 to 29	83% (74-90%)	17% (10-26%)	1.3 (0.84-1.8)	1.5% (0.95-2.1%)	5.4% (1.9-9.1%)	5.8 (2-9.9)	960 (610-1400)
Netherlands	2016	Female	30 to 34	84% (74-91%)	16% (9.5-26%)	1.4 (0.92-2)	1.4% (0.94-2.1%)	6.2% (3.2-9.4%)	9.9 (5.2-16)	1000 (670-1500)
Netherlands	2016	Female	35 to 39	84% (77-90%)	16% (9.7-23%)	2 (1.4-2.9)	2% (1.3-2.9%)	9.1% (5.8-13%)	20 (12-29)	1500 (950-2100)
Netherlands	2016	Female	40 to 44	85% (77-91%)	15% (9.5-23%)	2.5 (1.8-3.5)	2.8% (1.8-4%)	10% (7.3-14%)	56 (39-76)	3200 (2100-4500)
Netherlands	2016	Female	45 to 49	85% (78-90%)	15% (10-22%)	2.8 (2-3.8)	3.7% (2.4-5.1%)	11% (7.7-14%)	110 (75-140)	5000 (3400-6900)
Netherlands	2016	Female	50 to 54	86% (81-91%)	14% (9.5-19%)	3 (2.2-4.1)	4.3% (3-5.9%)	10% (7.8-13%)	170 (130-220)	6900 (4800-9400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	2016	Female	55 to 59	89% (86-92%)	11% (7.7-14%)	3.1 (2.3-4.2)	4.5% (3-6.1%)	9.1% (7.1-12%)	200 (150-260)	7200 (5000-9900)
Netherlands	2016	Female	60 to 64	89% (86-92%)	11% (7.6-14%)	3.1 (2.2-4.1)	4.1% (2.6-5.9%)	7.6% (5.6-10%)	240 (170-320)	7200 (4600-10000)
Netherlands	2016	Female	65 to 69	87% (82-91%)	13% (9.4-18%)	2.8 (2.1-3.7)	3.5% (2.1-5.2%)	6.2% (4.3-8.5%)	310 (210-430)	7600 (4600-12000)
Netherlands	2016	Female	70 to 74	84% (79-88%)	16% (12-21%)	2.5 (1.8-3.4)	3% (1.4-5%)	4.9% (2.7-7.4%)	250 (140-390)	5100 (2300-8300)
Netherlands	2016	Female	75 to 79	80% (72-87%)	20% (13-28%)	2.2 (1.5-3)	2.4% (0.62-4.5%)	3.5% (1.2-6%)	250 (90-440)	3900 (1100-7500)
Netherlands	2016	Female	80 to 84	77% (66-85%)	23% (15-34%)	2 (1.3-2.8)	1.9% (0.21-4%)	2.5% (0.47-5.1%)	300 (53-590)	3400 (390-7100)
Netherlands	2016	Female	85 to 89	75% (65-83%)	25% (17-35%)	1.8 (1.2-2.6)	1.4% (-0.22-3.4%)	1.8% (-0.24-4.2%)	270 (-36-640)	2200 (-350-5300)
Netherlands	2016	Female	90 to 94	74% (64-83%)	26% (17-36%)	1.7 (1.1-2.5)	1% (-0.65-3.2%)	1.2% (-0.8-3.7%)	160 (-110-510)	910 (-560-2900)
Netherlands	2016	Female	95 plus	74% (58-85%)	26% (15-42%)	1.6 (0.86-2.5)	0.78% (-1.1-3.4%)	0.78% (-1.4-3.8%)	48 (-90-230)	200 (-280-820)
Netherlands	2016	Male	All Ages	89% (87-91%)	11% (8.8-13%)	3.2 (2.5-4)	6% (4.1-8.1%)	6.4% (3.9-9.4%)	4700 (2800-6900)	130000 (90000-180000)
Netherlands	2016	Male	15 to 19	80% (72-88%)	20% (12-28%)	2.4 (1.6-3.3)	3.5% (2.2-4.9%)	11% (5.7-17%)	14 (7.4-22)	1600 (1000-2300)
Netherlands	2016	Male	20 to 24	87% (80-92%)	13% (7.7-20%)	3.6 (2.5-4.9)	7.5% (5.1-10%)	18% (10-25%)	40 (23-59)	4300 (2900-5900)
Netherlands	2016	Male	25 to 29	89% (84-93%)	11% (6.6-16%)	3.1 (2.2-4.3)	6.4% (4.3-8.4%)	16% (8.3-23%)	35 (18-51)	3700 (2500-5100)
Netherlands	2016	Male	30 to 34	91% (86-95%)	8.6% (5-14%)	3.1 (2.2-4.3)	5.6% (3.6-7.5%)	14% (7.5-21%)	39 (21-57)	3500 (2300-4800)
Netherlands	2016	Male	35 to 39	93% (89-96%)	6.8% (3.9-11%)	3.5 (2.5-4.6)	5.8% (3.9-7.9%)	15% (8.9-21%)	47 (27-68)	3500 (2400-4800)
Netherlands	2016	Male	40 to 44	94% (90-97%)	5.6% (3-9.7%)	3.9 (2.8-5.1)	6.5% (4.4-8.8%)	15% (10-21%)	100 (67-150)	6400 (4300-8700)
Netherlands	2016	Male	45 to 49	95% (92-97%)	4.8% (2.9-7.9%)	4.3 (3.2-5.5)	7.4% (5.1-10%)	15% (10-20%)	180 (120-250)	9400 (6400-13000)
Netherlands	2016	Male	50 to 54	96% (93-97%)	4.3% (2.7-6.5%)	4.7 (3.4-6.2)	8.4% (5.8-11%)	15% (10-19%)	310 (210-410)	13000 (9200-18000)
Netherlands	2016	Male	55 to 59	96% (94-97%)	4.3% (3-6%)	5.1 (3.8-6.7)	9.2% (6.7-12%)	14% (11-18%)	440 (310-600)	16000 (12000-22000)
Netherlands	2016	Male	60 to 64	95% (93-97%)	4.9% (3.5-6.9%)	5 (3.7-6.6)	8.6% (6-11%)	12% (8.6-16%)	580 (400-790)	18000 (12000-25000)
Netherlands	2016	Male	65 to 69	94% (91-96%)	6.1% (4.1-8.5%)	4.3 (3.2-5.6)	7.1% (4.7-9.6%)	9.3% (6.3-13%)	760 (490-1100)	20000 (13000-28000)
Netherlands	2016	Male	70 to 74	93% (90-95%)	7.3% (5.2-10%)	3.8 (2.8-4.9)	5.4% (3.2-8.1%)	6.7% (3.9-10%)	580 (340-880)	12000 (7000-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Netherlands	2016	Male	75 to 79	91% (87-94%)	8.9% (5.7-13%)	3.4 (2.5-4.5)	4.2% (2.1-6.9%)	4.9% (2.4-8.1%)	530 (260-880)	8700 (4200-14000)
Netherlands	2016	Male	80 to 84	89% (83-93%)	11% (6.5-17%)	3 (2.2-4.2)	3.5% (1.5-6.3%)	3.9% (1.6-7.1%)	520 (210-950)	6200 (2700-11000)
Netherlands	2016	Male	85 to 89	88% (81-93%)	12% (7.5-19%)	2.8 (1.9-3.8)	2.9% (0.98-5.4%)	3.1% (0.91-5.9%)	350 (100-690)	3000 (1000-5600)
Netherlands	2016	Male	90 to 94	88% (81-93%)	12% (7-19%)	2.6 (1.8-3.7)	2.4% (0.26-4.9%)	2.5% (0.031-5.6%)	160 (2.1-360)	960 (99-2000)
Netherlands	2016	Male	95 plus	88% (79-94%)	12% (5.6-21%)	2.5 (1.5-3.8)	2.1% (-0.18-5.1%)	2.1% (-0.55-5.7%)	34 (-8.5-89)	140 (-11-330)
Norway	1990	Female	All Ages	92% (90-93%)	8.4% (6.6-10%)	1.7 (1.5-2)	1.2% (-0.43-3%)	-0.24% (-3.5-3.3%)	-51 (-740-700)	7300 (-2500-17000)
Norway	1990	Female	15 to 19	90% (85-95%)	9.5% (5.4-15%)	1.4 (0.85-2.2)	2.7% (1.5-4.2%)	8.6% (3.4-14%)	3.9 (1.5-6.6)	440 (230-660)
Norway	1990	Female	20 to 24	94% (89-97%)	6% (3.2-11%)	1.8 (0.98-2.9)	3.6% (2.4-5%)	8.8% (3.6-14%)	4.2 (1.7-7)	710 (450-1000)
Norway	1990	Female	25 to 29	95% (92-97%)	4.7% (2.6-8.3%)	1.8 (1-2.9)	3.3% (2.1-4.7%)	8.3% (3.1-14%)	5.3 (2-8.8)	690 (420-980)
Norway	1990	Female	30 to 34	96% (93-98%)	4.1% (2.1-7.4%)	1.9 (1.2-3)	2.9% (1.7-4.2%)	8.4% (3.8-13%)	7 (3.2-11)	650 (390-940)
Norway	1990	Female	35 to 39	96% (92-98%)	4.3% (2.2-7.8%)	2 (1.1-3.2)	3.2% (1.8-4.7%)	9.2% (4.9-14%)	11 (5.7-16)	770 (460-1100)
Norway	1990	Female	40 to 44	95% (92-98%)	4.7% (2.4-8.4%)	2.3 (1.3-3.3)	3.8% (2.2-5.6%)	10% (5.9-14%)	19 (11-27)	1100 (660-1600)
Norway	1990	Female	45 to 49	95% (91-97%)	5% (2.5-9.4%)	2.6 (1.5-4.2)	4.6% (2.5-7.1%)	11% (6.1-15%)	26 (15-38)	1200 (710-1900)
Norway	1990	Female	50 to 54	94% (90-97%)	5.5% (3.1-9.6%)	2.8 (1.7-4.3)	4.5% (2.4-7.1%)	9.2% (5.2-14%)	27 (15-40)	1100 (600-1700)
Norway	1990	Female	55 to 59	94% (89-97%)	6.1% (3.3-11%)	2.9 (1.7-4.3)	4% (1.5-6.9%)	7.1% (2.8-12%)	33 (13-54)	1100 (430-1900)
Norway	1990	Female	60 to 64	93% (88-96%)	6.9% (3.7-12%)	2.7 (1.6-4.2)	2.8% (-0.019-6.4%)	4.5% (0.18-9.8%)	36 (1.5-79)	1100 (-6.8-2400)
Norway	1990	Female	65 to 69	92% (86-96%)	8.1% (4.1-14%)	2.4 (1.6-3.4)	1.2% (-1.6-4.4%)	1.9% (-2.3-6.4%)	27 (-32-88)	630 (-840-2200)
Norway	1990	Female	70 to 74	90% (84-94%)	9.7% (5.8-16%)	2.1 (1.2-3.2)	-0.00048% (-3.1-3.6%)	0.08% (-4.2-5%)	1.8 (-92-110)	0.66 (-1800-2100)
Norway	1990	Female	75 to 79	88% (81-94%)	12% (6.3-19%)	1.9 (1.1-3.1)	-1.1% (-4.3-2.9%)	-1.4% (-5.6-3.7%)	-47 (-190-120)	-680 (-2800-1900)
Norway	1990	Female	80 to 84	86% (78-92%)	14% (7.9-22%)	1.9 (1.2-2.9)	-1.4% (-4.6-2.3%)	-1.7% (-5.7-2.7%)	-75 (-240-120)	-810 (-2600-1400)
Norway	1990	Female	85 to 89	84% (75-91%)	16% (9.5-25%)	1.9 (1.1-2.8)	-1.4% (-4.3-2.2%)	-1.8% (-5.2-2.3%)	-75 (-220-99)	-550 (-1700-850)
Norway	1990	Female	90 to 94	83% (73-90%)	17% (9.8-27%)	1.8 (0.96-3)	-1.3% (-4.2-2.6%)	-1.7% (-5.2-2.9%)	-43 (-130-73)	-210 (-660-420)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	1990	Female	95 plus	82% (69-91%)	18% (8.5-31%)	1.8 (0.77-3.3)	-1.1% (-4.3-3.9%)	-1.4% (-5.2-4.3%)	-13 (-47-39)	-41 (-160-140)
Norway	1990	Male	All Ages	94% (93-96%)	5.9% (4.3-7.5%)	2 (1.7-2.3)	3.3% (1.2-5.3%)	0.82% (-2.5-4.1%)	190 (-580-950)	22000 (7900-36000)
Norway	1990	Male	15 to 19	90% (84-94%)	9.8% (5.7-16%)	1.4 (0.92-2.1)	6.3% (3.3-9.3%)	10% (4-16%)	14 (5.5-22)	1300 (710-2000)
Norway	1990	Male	20 to 24	94% (90-97%)	5.7% (3.1-10%)	2.1 (1.3-3.1)	11% (7.2-15%)	13% (6.4-22%)	24 (11-38)	2700 (1700-3700)
Norway	1990	Male	25 to 29	96% (93-98%)	3.9% (2.1-6.9%)	2.3 (1.4-3.3)	10% (7.2-14%)	14% (7-20%)	23 (12-35)	2600 (1700-3500)
Norway	1990	Male	30 to 34	97% (94-99%)	2.9% (1.4-5.5%)	2.3 (1.5-3.5)	9% (6.3-12%)	13% (7.2-19%)	26 (14-38)	2400 (1700-3200)
Norway	1990	Male	35 to 39	97% (95-99%)	2.6% (1.2-4.6%)	2.3 (1.4-3.4)	8.6% (6.1-11%)	13% (7.6-18%)	32 (19-46)	2500 (1800-3300)
Norway	1990	Male	40 to 44	98% (95-99%)	2.5% (1.2-4.6%)	2.5 (1.6-3.5)	8.3% (5.9-11%)	12% (7.5-16%)	44 (28-61)	2900 (2000-3800)
Norway	1990	Male	45 to 49	98% (95-99%)	2.4% (1.1-5.1%)	3 (1.8-4.3)	7.7% (4.7-11%)	10% (5.3-15%)	47 (24-72)	2600 (1600-3800)
Norway	1990	Male	50 to 54	97% (95-99%)	2.6% (1.1-4.9%)	3.2 (2.1-4.6)	6.5% (3.4-10%)	7.9% (3.4-13%)	44 (19-74)	2100 (1100-3300)
Norway	1990	Male	55 to 59	97% (95-99%)	2.8% (1.3-5.2%)	3.4 (2.3-4.8)	4.6% (1.1-8.6%)	5% (0.077-10%)	46 (0.71-97)	1900 (450-3600)
Norway	1990	Male	60 to 64	97% (94-98%)	3.3% (1.5-6.2%)	3.3 (2.1-4.8)	2.8% (-0.9-7.2%)	2.7% (-2.1-8.3%)	42 (-34-130)	1600 (-520-4200)
Norway	1990	Male	65 to 69	96% (93-98%)	3.9% (1.8-7.2%)	2.9 (2.1-4)	0.91% (-2.7-4.6%)	0.5% (-3.9-5.1%)	13 (-100-130)	690 (-2000-3500)
Norway	1990	Male	70 to 74	96% (92-98%)	4.4% (2.4-7.7%)	2.5 (1.7-3.7)	-0.19% (-3.5-3.4%)	-0.66% (-4.7-3.7%)	-23 (-160-130)	-150 (-2800-2700)
Norway	1990	Male	75 to 79	95% (91-98%)	4.9% (2.4-8.6%)	2.3 (1.4-3.4)	-0.64% (-3.8-2.9%)	-1.1% (-4.8-3%)	-44 (-200-120)	-440 (-2600-2000)
Norway	1990	Male	80 to 84	94% (90-97%)	5.5% (2.6-10%)	2.2 (1.5-3.1)	-0.82% (-3.7-2.3%)	-1.2% (-4.7-2.3%)	-47 (-180-88)	-380 (-1800-1100)
Norway	1990	Male	85 to 89	94% (89-97%)	6% (3.1-11%)	2.2 (1.4-3.1)	-0.76% (-3.4-2%)	-1.1% (-4.3-2.1%)	-30 (-110-55)	-170 (-790-460)
Norway	1990	Male	90 to 94	94% (89-97%)	5.9% (2.9-11%)	2.1 (1.3-3.2)	-0.86% (-3.4-2.2%)	-1.3% (-4.3-2.3%)	-15 (-51-27)	-63 (-250-170)
Norway	1990	Male	95 plus	94% (89-98%)	5.8% (2.3-11%)	2.1 (1.1-3.5)	-0.77% (-3.6-3%)	-1.2% (-4.5-3.2%)	-41 (-15-11)	-11 (-50-43)
Norway	1995	Female	All Ages	92% (90-94%)	7.8% (6.2-9.7%)	1.7 (1.5-2)	1.3% (-0.096-2.9%)	-0.1% (-3-3.3%)	-23 (-660-720)	7700 (-530-17000)
Norway	1995	Female	15 to 19	92% (87-95%)	8.5% (4.8-13%)	1.4 (0.84-2.2)	2.6% (1.4-3.9%)	8.1% (2.9-14%)	3.1 (1.1-5.2)	360 (190-530)
Norway	1995	Female	20 to 24	95% (91-97%)	5.2% (2.9-8.9%)	1.8 (0.96-2.9)	3.5% (2.3-4.8%)	8.1% (2.9-14%)	3.7 (1.3-6.1)	650 (400-920)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	1995	Female	25 to 29	96% (93-98%)	4% (2.2-6.9%)	1.8 (1.1-2.9)	2.8% (1.9-3.9%)	7.6% (2.8-13%)	3.9 (1.4-6.6)	610 (390-860)
Norway	1995	Female	30 to 34	96% (94-98%)	3.5% (1.7-6.1%)	1.9 (1.2-3)	2.6% (1.5-3.7%)	7.8% (3.7-12%)	6.1 (2.8-9.8)	590 (350-860)
Norway	1995	Female	35 to 39	96% (93-98%)	3.7% (1.8-6.5%)	2 (1.1-3.1)	2.8% (1.7-4.2%)	8.9% (4.8-13%)	9.7 (5.1-14)	700 (410-1000)
Norway	1995	Female	40 to 44	96% (93-98%)	4% (2.1-6.9%)	2.3 (1.3-3.3)	3.6% (2-5.3%)	10% (6.1-14%)	18 (11-25)	1000 (600-1400)
Norway	1995	Female	45 to 49	96% (92-98%)	4.4% (2.1-7.8%)	2.6 (1.6-4.2)	4.4% (2.4-6.8%)	11% (6.5-15%)	31 (19-44)	1500 (840-2200)
Norway	1995	Female	50 to 54	95% (92-97%)	4.9% (2.7-8.4%)	2.8 (1.7-4.3)	4.6% (2.5-7.2%)	9.5% (5.7-14%)	34 (21-49)	1400 (780-2100)
Norway	1995	Female	55 to 59	95% (91-97%)	5.4% (3.1-9.3%)	2.9 (1.7-4.4)	4.2% (1.8-6.9%)	7.8% (3.9-12%)	35 (18-52)	1200 (520-1900)
Norway	1995	Female	60 to 64	94% (89-96%)	6.4% (3.6-11%)	2.8 (1.6-4.3)	3% (0.3-6.4%)	5.1% (0.95-9.9%)	35 (6.7-68)	1000 (100-2000)
Norway	1995	Female	65 to 69	92% (87-96%)	8.1% (4.3-13%)	2.4 (1.6-3.4)	1.5% (-1-4.4%)	2.5% (-1.2-6.4%)	29 (-14-74)	660 (-440-1900)
Norway	1995	Female	70 to 74	90% (84-94%)	10% (6.1-16%)	2.1 (1.2-3.2)	0.27% (-2.3-3.4%)	0.54% (-3.1-4.7%)	11 (-67-100)	160 (-1400-2000)
Norway	1995	Female	75 to 79	87% (80-93%)	13% (7.1-20%)	1.9 (1-3)	-0.76% (-3.6-2.9%)	-0.98% (-4.6-3.7%)	-32 (-150-120)	-490 (-2300-1900)
Norway	1995	Female	80 to 84	85% (76-92%)	15% (8.5-24%)	1.8 (1.1-2.7)	-1.2% (-4-2.1%)	-1.5% (-4.9-2.5%)	-65 (-220-110)	-710 (-2400-1300)
Norway	1995	Female	85 to 89	83% (74-90%)	17% (10-26%)	1.8 (1-2.8)	-1.4% (-4-2.1%)	-1.7% (-5-2.3%)	-79 (-240-110)	-590 (-1800-900)
Norway	1995	Female	90 to 94	81% (71-89%)	19% (11-29%)	1.8 (0.92-2.9)	-1.3% (-4.1-2.5%)	-1.7% (-5-2.7%)	-51 (-150-82)	-250 (-770-470)
Norway	1995	Female	95 plus	80% (66-91%)	20% (9.2-34%)	1.7 (0.71-3.2)	-1.1% (-4.1-3.7%)	-1.4% (-5-4.2%)	-17 (-58-49)	-52 (-190-180)
Norway	1995	Male	All Ages	95% (94-96%)	4.8% (3.6-6%)	2 (1.8-2.3)	3.2% (1.3-5.1%)	0.85% (-2.3-4.2%)	190 (-510-920)	21000 (8300-34000)
Norway	1995	Male	15 to 19	92% (88-95%)	7.8% (4.6-12%)	1.5 (0.94-2.2)	5.7% (3.1-8.4%)	9.7% (3.7-16%)	9.2 (3.6-15)	940 (500-1400)
Norway	1995	Male	20 to 24	96% (93-97%)	4.4% (2.6-7.1%)	2.1 (1.3-3.2)	9.7% (6.4-13%)	13% (5.5-20%)	18 (7.4-28)	2100 (1400-3000)
Norway	1995	Male	25 to 29	97% (95-98%)	3% (1.8-4.6%)	2.3 (1.4-3.4)	8.8% (6.2-12%)	12% (5.5-19%)	19 (9-30)	2300 (1500-3000)
Norway	1995	Male	30 to 34	98% (96-99%)	2.2% (1.1-3.7%)	2.4 (1.5-3.5)	7.4% (5-9.8%)	11% (5.4-16%)	19 (10-29)	1900 (1300-2600)
Norway	1995	Male	35 to 39	98% (97-99%)	2% (0.99-3.4%)	2.3 (1.4-3.4)	7.1% (5.1-9.3%)	11% (6.3-15%)	24 (14-34)	2000 (1400-2600)
Norway	1995	Male	40 to 44	98% (97-99%)	1.9% (0.93-3.4%)	2.5 (1.6-3.5)	7.4% (5.4-9.6%)	11% (7.3-16%)	34 (21-47)	2300 (1600-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	1995	Male	45 to 49	98% (97-99%)	1.9% (0.84-3.4%)	3 (1.9-4.3)	7.6% (4.9-11%)	11% (5.9-15%)	55 (30-80)	3000 (1900-4200)
Norway	1995	Male	50 to 54	98% (96-99%)	2% (0.93-3.6%)	3.3 (2.1-4.6)	6.5% (3.7-9.8%)	8.3% (4-13%)	51 (24-81)	2500 (1400-3600)
Norway	1995	Male	55 to 59	98% (96-99%)	2.3% (1-4.2%)	3.4 (2.2-4.9)	5.2% (2-8.9%)	6.1% (1.4-11%)	48 (11-91)	2000 (770-3500)
Norway	1995	Male	60 to 64	97% (95-99%)	2.6% (1.2-4.7%)	3.4 (2-5)	3.4% (-0.19-7.5%)	3.6% (-1.2-9%)	43 (-15-110)	1600 (-87-3500)
Norway	1995	Male	65 to 69	97% (94-98%)	3.3% (1.6-5.8%)	3 (2.1-4.1)	1.4% (-1.9-5%)	1.1% (-3-5.6%)	24 (-63-120)	860 (-1200-3100)
Norway	1995	Male	70 to 74	96% (93-98%)	4% (2.2-6.6%)	2.6 (1.7-3.7)	0.23% (-2.8-3.8%)	-0.14% (-3.8-4%)	-4.6 (-130-130)	180 (-2100-2800)
Norway	1995	Male	75 to 79	95% (92-98%)	4.6% (2.2-7.8%)	2.3 (1.4-3.4)	-0.53% (-3.4-3%)	-0.98% (-4.5-3%)	-40 (-180-120)	-370 (-2400-2100)
Norway	1995	Male	80 to 84	95% (91-98%)	5.1% (2.5-9.4%)	2.3 (1.5-3.2)	-0.83% (-3.8-2.3%)	-1.3% (-4.7-2.3%)	-51 (-190-91)	-420 (-1900-1200)
Norway	1995	Male	85 to 89	94% (90-97%)	5.5% (3-9.6%)	2.2 (1.4-3.2)	-0.84% (-3.4-2.1%)	-1.3% (-4.3-2%)	-36 (-120-58)	-210 (-860-530)
Norway	1995	Male	90 to 94	94% (90-97%)	5.5% (2.8-9.9%)	2.2 (1.3-3.3)	-0.92% (-3.4-2.2%)	-1.3% (-4.3-2.2%)	-18 (-57-29)	-74 (-280-180)
Norway	1995	Male	95 plus	94% (89-98%)	5.6% (2.2-11%)	2.1 (1.1-3.5)	-0.71% (-3.6-3.2%)	-1.1% (-4.5-3.3%)	-4.2 (-16-12)	-11 (-53-49)
Norway	2000	Female	All Ages	92% (90-94%)	8% (6.3-9.8%)	1.9 (1.7-2.1)	1.7% (0.39-3.3%)	0.45% (-2.4-3.8%)	100 (-530-850)	9900 (2100-19000)
Norway	2000	Female	15 to 19	91% (87-95%)	8.7% (5-13%)	1.5 (0.88-2.4)	2.7% (1.4-4.1%)	7.9% (3.1-13%)	3.2 (1.2-5.4)	360 (190-550)
Norway	2000	Female	20 to 24	95% (91-97%)	5.3% (2.9-9%)	1.9 (1-3.2)	3.6% (2.3-5%)	7.9% (2.9-13%)	3.6 (1.3-6.1)	590 (380-850)
Norway	2000	Female	25 to 29	96% (93-98%)	4% (2.2-7%)	2 (1.1-3.1)	3% (1.9-4.2%)	7.6% (2.7-13%)	4.5 (1.6-7.5)	640 (400-900)
Norway	2000	Female	30 to 34	96% (94-98%)	3.5% (1.8-6.2%)	2.1 (1.3-3.2)	2.5% (1.6-3.8%)	7.5% (3.7-12%)	6.2 (3.1-9.8)	620 (380-900)
Norway	2000	Female	35 to 39	96% (93-98%)	3.8% (1.8-6.6%)	2.2 (1.2-3.4)	2.8% (1.7-4.2%)	8.5% (4.6-13%)	9.7 (5.3-14)	720 (420-1000)
Norway	2000	Female	40 to 44	96% (93-98%)	4.1% (2.1-7%)	2.5 (1.4-3.6)	3.4% (1.9-5.1%)	9.7% (5.8-14%)	16 (9.6-23)	940 (560-1400)
Norway	2000	Female	45 to 49	96% (92-98%)	4.4% (2.2-8.1%)	2.8 (1.6-4.4)	4.4% (2.5-6.6%)	11% (6.8-15%)	28 (18-40)	1400 (810-2100)
Norway	2000	Female	50 to 54	95% (91-97%)	5% (2.8-8.6%)	3 (1.8-4.6)	4.7% (2.5-7.2%)	9.7% (5.9-14%)	42 (26-61)	1700 (1000-2700)
Norway	2000	Female	55 to 59	94% (91-97%)	5.5% (3.2-9.3%)	3.1 (1.8-4.7)	4.6% (2.2-7.4%)	8.4% (4.8-12%)	45 (26-65)	1600 (800-2400)
Norway	2000	Female	60 to 64	93% (89-96%)	6.6% (3.6-11%)	3 (1.7-4.6)	3.6% (0.99-6.8%)	6.1% (2.3-10%)	41 (15-69)	1200 (330-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	2000	Female	65 to 69	92% (86-95%)	8.3% (4.5-14%)	2.6 (1.7-3.7)	2.2% (-0.057-4.8%)	3.7% (0.37-7.1%)	34 (3.4-67)	800 (-22-1700)
Norway	2000	Female	70 to 74	90% (84-94%)	10% (6.4-16%)	2.2 (1.3-3.5)	0.99% (-1.6-3.9%)	1.6% (-1.8-5.4%)	26 (-31-92)	470 (-740-1900)
Norway	2000	Female	75 to 79	87% (79-93%)	13% (7.5-21%)	2 (1.1-3.3)	-0.028% (-2.8-3.5%)	-0.012% (-3.5-4.5%)	-0.32 (-110-140)	-18 (-1700-2300)
Norway	2000	Female	80 to 84	85% (75-91%)	15% (8.6-25%)	2 (1.2-3)	-0.63% (-3.5-2.7%)	-0.82% (-4.2-3.3%)	-37 (-190-150)	-390 (-2200-1800)
Norway	2000	Female	85 to 89	82% (73-89%)	18% (11-27%)	1.9 (1.1-3)	-0.92% (-3.7-2.7%)	-1.2% (-4.5-3%)	-60 (-230-140)	-430 (-1700-1200)
Norway	2000	Female	90 to 94	81% (71-89%)	19% (11-29%)	1.9 (0.98-3.2)	-1% (-3.9-3.2%)	-1.4% (-4.8-3.5%)	-47 (-160-120)	-220 (-830-690)
Norway	2000	Female	95 plus	80% (65-91%)	20% (9.3-35%)	1.9 (0.77-3.5)	-0.82% (-3.9-4.5%)	-1.2% (-4.9-5.1%)	-16 (-65-68)	-46 (-220-240)
Norway	2000	Male	All Ages	95% (94-96%)	4.9% (3.7-6.2%)	2.2 (1.9-2.5)	4% (2.3-5.7%)	1.8% (-1-4.9%)	380 (-220-1000)	25000 (14000-36000)
Norway	2000	Male	15 to 19	92% (88-95%)	7.9% (4.8-12%)	1.6 (1-2.4)	5.8% (3.2-8.5%)	9.8% (3.8-16%)	9.5 (3.7-15)	960 (530-1400)
Norway	2000	Male	20 to 24	96% (93-97%)	4.4% (2.6-7.3%)	2.3 (1.4-3.4)	9.8% (6.3-13%)	12% (5.6-19%)	19 (8.6-30)	2100 (1300-2900)
Norway	2000	Male	25 to 29	97% (95-98%)	3% (1.8-4.8%)	2.5 (1.5-3.5)	8.6% (5.8-12%)	11% (5.4-17%)	21 (9.9-31)	2200 (1500-3100)
Norway	2000	Male	30 to 34	98% (96-99%)	2.2% (1.1-3.8%)	2.6 (1.7-3.7)	7% (4.8-9.5%)	10% (5.1-16%)	20 (10-32)	2000 (1300-2700)
Norway	2000	Male	35 to 39	98% (97-99%)	2.1% (1-3.3%)	2.5 (1.6-3.6)	6.7% (4.6-9%)	10% (5.3-15%)	25 (13-36)	2000 (1300-2700)
Norway	2000	Male	40 to 44	98% (97-99%)	1.9% (0.94-3.3%)	2.7 (1.8-3.8)	7.1% (4.8-9.5%)	11% (6.6-15%)	34 (20-47)	2300 (1600-3000)
Norway	2000	Male	45 to 49	98% (96-99%)	1.9% (0.89-3.7%)	3.2 (2.1-4.6)	7.9% (5.3-11%)	12% (7.5-17%)	51 (32-71)	2800 (2000-3800)
Norway	2000	Male	50 to 54	98% (96-99%)	2.1% (0.89-3.7%)	3.5 (2.3-4.8)	7.5% (4.9-11%)	10% (6.1-15%)	74 (44-110)	3500 (2200-4900)
Norway	2000	Male	55 to 59	98% (96-99%)	2.3% (1.1-4.2%)	3.7 (2.4-5.1)	6.5% (3.8-10%)	8.3% (4.4-13%)	71 (38-110)	2900 (1600-4400)
Norway	2000	Male	60 to 64	97% (95-99%)	2.7% (1.4-4.7%)	3.6 (2.3-5.3)	4.8% (1.6-8.7%)	5.7% (1.2-11%)	61 (13-120)	2100 (670-3700)
Norway	2000	Male	65 to 69	97% (94-98%)	3.4% (1.6-5.8%)	3.2 (2.2-4.3)	2.7% (-0.12-6%)	2.9% (-0.79-7%)	47 (-13-110)	1400 (-61-3000)
Norway	2000	Male	70 to 74	96% (93-98%)	4.1% (2.3-6.7%)	2.8 (1.8-4)	1.3% (-1.4-4.6%)	1.2% (-2.2-5.2%)	30 (-57-130)	800 (-880-2800)
Norway	2000	Male	75 to 79	95% (92-98%)	4.6% (2.3-7.9%)	2.5 (1.5-3.8)	0.33% (-2.4-3.7%)	0.013% (-3.3-4%)	0.54 (-130-150)	230 (-1700-2600)
Norway	2000	Male	80 to 84	95% (91-98%)	5.2% (2.5-9.4%)	2.4 (1.6-3.5)	-0.28% (-3.1-2.7%)	-0.68% (-3.9-2.7%)	-28 (-160-110)	-150 (-1600-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	2000	Male	85 to 89	94% (90-97%)	5.6% (3-9.5%)	2.4 (1.5-3.4)	-0.54% (-3.2-2.4%)	-0.97% (-4.1-2.4%)	-30 (-130-76)	-150 (-890-680)
Norway	2000	Male	90 to 94	94% (90-97%)	5.5% (2.8-9.9%)	2.3 (1.4-3.5)	-0.75% (-3.4-2.6%)	-1.2% (-4.4-2.6%)	-18 (-64-38)	-66 (-310-230)
Norway	2000	Male	95 plus	94% (89-98%)	5.6% (2.2-11%)	2.3 (1.2-3.8)	-0.55% (-3.4-3.5%)	-1% (-4.3-3.6%)	-3.9 (-17-14)	-8.6 (-53-54)
Norway	2005	Female	All Ages	91% (89-93%)	8.5% (6.6-11%)	2.1 (1.8-2.4)	2.3% (0.97-3.8%)	1.4% (-1.4-4.8%)	300 (-300-1000)	13000 (5400-21000)
Norway	2005	Female	15 to 19	90% (84-95%)	9.7% (5.4-16%)	1.6 (0.95-2.6)	2.4% (1.4-3.6%)	8% (3.3-13%)	2.8 (1.2-4.7)	350 (200-520)
Norway	2005	Female	20 to 24	94% (89-97%)	6.1% (3.3-11%)	2.1 (1-3.5)	3.5% (2.2-5.1%)	8.4% (3-15%)	3.9 (1.4-6.8)	580 (350-840)
Norway	2005	Female	25 to 29	95% (92-97%)	4.7% (2.6-8.3%)	2.1 (1.2-3.5)	3% (1.9-4.3%)	8.1% (3.2-13%)	4.2 (1.6-6.8)	550 (330-780)
Norway	2005	Female	30 to 34	96% (93-98%)	4.1% (2.1-7.5%)	2.2 (1.4-3.5)	2.5% (1.5-3.8%)	8.3% (4.1-13%)	6 (3-9.4)	580 (340-860)
Norway	2005	Female	35 to 39	96% (92-98%)	4.4% (2.2-7.9%)	2.4 (1.3-3.8)	2.8% (1.6-4.2%)	9.2% (5.1-13%)	10 (5.7-15)	750 (440-1100)
Norway	2005	Female	40 to 44	95% (91-98%)	4.7% (2.4-8.5%)	2.7 (1.6-4)	3.4% (1.8-5.1%)	10% (6-14%)	16 (9.7-23)	960 (540-1400)
Norway	2005	Female	45 to 49	95% (90-97%)	5.1% (2.5-9.6%)	3.1 (1.8-4.8)	4.5% (2.6-6.7%)	11% (7.5-16%)	29 (19-42)	1400 (860-2100)
Norway	2005	Female	50 to 54	94% (90-97%)	5.6% (3.1-9.7%)	3.3 (2-5)	5% (2.8-7.4%)	11% (7.1-15%)	41 (27-57)	1700 (1000-2600)
Norway	2005	Female	55 to 59	94% (89-97%)	6.1% (3.4-11%)	3.4 (2-5)	4.9% (2.7-7.5%)	9.1% (5.8-13%)	55 (35-77)	2000 (1100-3000)
Norway	2005	Female	60 to 64	93% (88-96%)	7% (3.7-12%)	3.2 (1.8-4.9)	4.2% (1.7-7%)	7.1% (3.6-11%)	53 (27-82)	1600 (650-2600)
Norway	2005	Female	65 to 69	92% (86-96%)	8.2% (4.2-14%)	2.9 (1.9-4.1)	3% (0.85-5.3%)	5% (2.1-8%)	44 (18-70)	1100 (290-1900)
Norway	2005	Female	70 to 74	90% (84-94%)	9.8% (6-16%)	2.5 (1.4-3.8)	1.9% (-0.59-5%)	2.9% (-0.32-6.9%)	39 (-4.2-91)	730 (-240-1900)
Norway	2005	Female	75 to 79	88% (81-94%)	12% (6.4-19%)	2.3 (1.3-3.8)	0.92% (-1.9-4.5%)	1.3% (-2.2-5.7%)	31 (-52-140)	460 (-920-2200)
Norway	2005	Female	80 to 84	86% (77-92%)	14% (7.8-23%)	2.2 (1.4-3.5)	0.23% (-2.7-3.7%)	0.26% (-3.3-4.4%)	11 (-130-180)	140 (-1500-2200)
Norway	2005	Female	85 to 89	84% (75-91%)	16% (9.4-25%)	2.2 (1.3-3.4)	-0.14% (-3-3.5%)	-0.29% (-3.8-4%)	-14 (-190-200)	-67 (-1500-1700)
Norway	2005	Female	90 to 94	83% (72-90%)	17% (10-28%)	2.2 (1.2-3.7)	-0.37% (-3.4-4%)	-0.63% (-4.3-4.5%)	-23 (-160-170)	-87 (-790-940)
Norway	2005	Female	95 plus	81% (69-91%)	19% (8.7-31%)	2.1 (0.92-4)	-0.2% (-3.6-5.7%)	-0.46% (-4.3-6.5%)	-7.3 (-69-100)	-13 (-240-370)
Norway	2005	Male	All Ages	94% (92-96%)	5.9% (4.3-7.6%)	2.4 (2.1-2.8)	4.6% (3.1-6.3%)	3% (0.47-5.9%)	600 (93-1200)	27000 (18000-37000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	2005	Male	15 to 19	90% (83-94%)	10% (5.6-17%)	1.7 (1-2.6)	5.2% (3.1-7.4%)	9.7% (4-16%)	7.5 (3.2-12)	840 (500-1200)
Norway	2005	Male	20 to 24	94% (90-97%)	5.9% (3.2-10%)	2.5 (1.5-3.8)	9.3% (6.1-13%)	12% (5.5-19%)	16 (7.5-26)	1800 (1100-2600)
Norway	2005	Male	25 to 29	96% (93-98%)	4% (2.2-7.1%)	2.7 (1.6-4)	8.2% (5.7-11%)	12% (5.7-18%)	16 (7.8-24)	1700 (1100-2400)
Norway	2005	Male	30 to 34	97% (94-99%)	3% (1.4-5.9%)	2.7 (1.8-4)	6.7% (4.6-9%)	11% (5.7-17%)	17 (8.7-26)	1600 (1100-2200)
Norway	2005	Male	35 to 39	97% (95-99%)	2.7% (1.3-4.9%)	2.7 (1.7-4)	6.3% (4.3-8.6%)	11% (5.9-15%)	23 (13-33)	1800 (1200-2500)
Norway	2005	Male	40 to 44	97% (95-99%)	2.5% (1.2-4.7%)	3 (1.9-4.2)	6.5% (4.4-8.9%)	11% (6.6-16%)	28 (17-40)	1900 (1300-2600)
Norway	2005	Male	45 to 49	98% (95-99%)	2.5% (1.1-5%)	3.5 (2.3-5)	7.8% (5.4-11%)	13% (8.1-17%)	48 (31-68)	2700 (1900-3600)
Norway	2005	Male	50 to 54	97% (95-99%)	2.6% (1.1-5.2%)	3.8 (2.5-5.2)	8.1% (5.5-11%)	12% (8-17%)	70 (46-100)	3300 (2200-4600)
Norway	2005	Male	55 to 59	97% (94-99%)	2.9% (1.3-5.6%)	4 (2.7-5.4)	7.5% (4.9-11%)	10% (6.3-15%)	96 (59-140)	3800 (2400-5500)
Norway	2005	Male	60 to 64	97% (93-98%)	3.4% (1.5-6.6%)	3.9 (2.4-5.6)	6.1% (3.1-9.5%)	7.7% (3.5-12%)	90 (43-150)	3000 (1500-4700)
Norway	2005	Male	65 to 69	96% (92-98%)	4% (1.8-7.7%)	3.5 (2.4-4.7)	4.1% (1.5-7.1%)	4.9% (1.7-8.5%)	69 (23-120)	1900 (710-3300)
Norway	2005	Male	70 to 74	95% (92-98%)	4.5% (2.4-8%)	3 (1.9-4.3)	2.6% (-0.1-5.7%)	2.8% (-0.43-6.7%)	57 (-8.2-130)	1300 (-52-2800)
Norway	2005	Male	75 to 79	95% (91-98%)	5% (2.4-9%)	2.8 (1.6-4.1)	1.5% (-1.2-4.8%)	1.5% (-1.8-5.3%)	44 (-52-160)	840 (-640-2700)
Norway	2005	Male	80 to 84	94% (90-97%)	5.6% (2.6-10%)	2.7 (1.7-3.9)	0.76% (-1.8-3.8%)	0.52% (-2.5-3.9%)	21 (-99-160)	390 (-940-2000)
Norway	2005	Male	85 to 89	94% (89-97%)	6.1% (3-11%)	2.6 (1.7-3.8)	0.33% (-2.1-3.4%)	0.011% (-2.8-3.5%)	0.33 (-94-120)	99 (-620-1000)
Norway	2005	Male	90 to 94	94% (89-97%)	6% (2.8-11%)	2.5 (1.6-3.9)	0.017% (-2.6-3.6%)	-0.38% (-3.4-3.6%)	-6.2 (-56-59)	1.8 (-260-350)
Norway	2005	Male	95 plus	94% (89-98%)	5.9% (2.3-11%)	2.5 (1.3-4.2)	0.16% (-2.8-4.9%)	-0.25% (-3.7-5.1%)	-1.1 (-17-22)	2.9 (-53-88)
Norway	2010	Female	All Ages	91% (89-93%)	8.6% (6.8-11%)	2.1 (1.8-2.5)	2.4% (1.1-3.9%)	1.7% (-1-5%)	380 (-210-1100)	14000 (6400-22000)
Norway	2010	Female	15 to 19	90% (84-95%)	9.8% (5.4-16%)	1.6 (0.92-2.6)	2.4% (1.4-3.6%)	7.9% (3.3-13%)	3.2 (1.3-5.4)	390 (230-580)
Norway	2010	Female	20 to 24	94% (89-97%)	6.2% (3.3-11%)	2.1 (1.1-3.5)	3.4% (2.2-4.9%)	8.5% (3.1-15%)	3.9 (1.4-6.7)	630 (380-900)
Norway	2010	Female	25 to 29	95% (91-97%)	4.8% (2.6-8.7%)	2.2 (1.2-3.6)	3% (1.8-4.2%)	8.3% (3.2-14%)	4.2 (1.6-6.9)	570 (340-840)
Norway	2010	Female	30 to 34	96% (92-98%)	4.2% (2.1-7.5%)	2.3 (1.3-3.7)	2.5% (1.5-3.7%)	8.8% (4.4-14%)	5.1 (2.6-8.2)	520 (320-770)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	2010	Female	35 to 39	96% (92-98%)	4.4% (2.1-8.3%)	2.4 (1.3-3.9)	2.6% (1.5-3.9%)	9.5% (5.3-14%)	8.6 (4.8-12)	670 (390-990)
Norway	2010	Female	40 to 44	95% (91-98%)	4.8% (2.4-8.6%)	2.7 (1.5-4.2)	3% (1.6-4.5%)	10% (5.9-14%)	14 (8.5-20)	890 (510-1300)
Norway	2010	Female	45 to 49	95% (90-98%)	5.2% (2.5-9.9%)	3.1 (1.8-4.9)	4.1% (2.4-6.4%)	11% (7.5-16%)	26 (17-37)	1300 (780-2000)
Norway	2010	Female	50 to 54	94% (90-97%)	5.7% (3-10%)	3.3 (2-5.1)	4.9% (2.9-7.4%)	11% (7.4-15%)	41 (28-57)	1800 (1100-2600)
Norway	2010	Female	55 to 59	94% (89-97%)	6.2% (3.4-11%)	3.5 (2-5.2)	4.8% (2.7-7.2%)	9.4% (6.2-13%)	51 (33-70)	1800 (1000-2800)
Norway	2010	Female	60 to 64	93% (87-96%)	7% (3.8-13%)	3.3 (1.8-5.1)	4.1% (1.8-7.1%)	7.2% (3.8-11%)	62 (33-96)	1900 (810-3100)
Norway	2010	Female	65 to 69	92% (86-96%)	8.2% (4.3-14%)	2.9 (1.9-4.2)	3.2% (1.1-5.6%)	5.3% (2.6-8.3%)	55 (27-85)	1300 (480-2300)
Norway	2010	Female	70 to 74	90% (84-94%)	9.8% (5.8-16%)	2.5 (1.4-3.9)	2.1% (-0.25-5%)	3.4% (0.38-7%)	43 (4.9-88)	820 (-97-1900)
Norway	2010	Female	75 to 79	88% (80-93%)	12% (6.5-20%)	2.3 (1.2-3.9)	1.3% (-1.3-4.5%)	1.9% (-1.4-5.8%)	37 (-27-120)	540 (-560-2000)
Norway	2010	Female	80 to 84	86% (77-92%)	14% (7.8-23%)	2.3 (1.4-3.6)	0.58% (-2.1-3.9%)	0.78% (-2.6-4.7%)	27 (-87-170)	300 (-1000-2000)
Norway	2010	Female	85 to 89	84% (75-90%)	16% (9.8-25%)	2.3 (1.3-3.6)	0.2% (-2.6-3.6%)	0.15% (-3.2-4.3%)	7.7 (-160-220)	99 (-1300-1900)
Norway	2010	Female	90 to 94	83% (73-90%)	17% (9.9-27%)	2.2 (1.2-3.8)	-0.067% (-3-4.4%)	-0.26% (-3.9-4.9%)	-11 (-160-210)	-17 (-830-1200)
Norway	2010	Female	95 plus	81% (68-91%)	19% (8.7-32%)	2.2 (0.91-4.1)	0.04% (-3.4-6.1%)	-0.18% (-4.2-7%)	-3.5 (-83-140)	3.5 (-270-490)
Norway	2010	Male	All Ages	94% (93-96%)	5.8% (4.3-7.4%)	2.5 (2.1-2.9)	4.8% (3.4-6.4%)	3.5% (1.2-6.2%)	700 (240-1200)	29000 (20000-39000)
Norway	2010	Male	15 to 19	90% (84-94%)	10% (5.8-16%)	1.7 (1-2.7)	4.8% (3-7%)	9.5% (3.9-16%)	6.8 (2.8-11)	820 (490-1200)
Norway	2010	Male	20 to 24	94% (90-97%)	5.9% (3.2-10%)	2.5 (1.5-3.9)	9.2% (6.2-13%)	12% (5.5-20%)	15 (6.7-25)	1900 (1200-2700)
Norway	2010	Male	25 to 29	96% (93-98%)	4% (2.2-7.1%)	2.7 (1.6-4.1)	8.3% (5.7-11%)	12% (5.9-18%)	16 (7.9-25)	1800 (1200-2500)
Norway	2010	Male	30 to 34	97% (94-99%)	2.9% (1.4-5.7%)	2.8 (1.8-4.1)	6.7% (4.6-9.1%)	11% (5.7-18%)	15 (7.8-24)	1500 (1000-2100)
Norway	2010	Male	35 to 39	97% (95-99%)	2.7% (1.2-4.7%)	2.7 (1.6-4.1)	6% (4.1-8.2%)	11% (5.7-16%)	19 (10-28)	1600 (1100-2200)
Norway	2010	Male	40 to 44	97% (95-99%)	2.5% (1.2-4.7%)	3 (1.9-4.3)	6.2% (4.3-8.5%)	11% (6.9-16%)	29 (18-42)	2000 (1300-2700)
Norway	2010	Male	45 to 49	98% (95-99%)	2.5% (1.1-5.1%)	3.5 (2.2-5)	7.4% (5-10%)	12% (7.9-17%)	46 (29-65)	2600 (1800-3600)
Norway	2010	Male	50 to 54	97% (95-99%)	2.6% (1.1-5.2%)	3.8 (2.6-5.3)	8% (5.5-11%)	12% (8.1-17%)	70 (46-97)	3300 (2300-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	2010	Male	55 to 59	97% (95-99%)	2.9% (1.3-5.4%)	4 (2.7-5.7)	7.9% (5.2-11%)	11% (7.3-16%)	96 (62-140)	3800 (2500-5300)
Norway	2010	Male	60 to 64	97% (94-98%)	3.3% (1.6-6.3%)	4 (2.5-5.8)	6.5% (3.7-10%)	8.5% (4.7-13%)	120 (65-180)	3900 (2200-6000)
Norway	2010	Male	65 to 69	96% (93-98%)	3.9% (1.8-7.4%)	3.5 (2.4-4.8)	4.7% (2.3-7.3%)	5.7% (2.7-9.2%)	96 (45-150)	2600 (1300-4100)
Norway	2010	Male	70 to 74	96% (92-98%)	4.5% (2.4-7.6%)	3.1 (2-4.5)	3% (0.5-5.8%)	3.5% (0.51-6.8%)	66 (9.7-130)	1500 (250-2900)
Norway	2010	Male	75 to 79	95% (91-98%)	5% (2.4-8.9%)	2.8 (1.7-4.3)	2% (-0.56-5.1%)	2.1% (-0.98-5.8%)	54 (-25-150)	970 (-270-2500)
Norway	2010	Male	80 to 84	94% (90-97%)	5.6% (2.6-10%)	2.7 (1.8-4)	1.2% (-1.2-3.9%)	1.1% (-1.8-4.2%)	37 (-64-150)	560 (-580-1900)
Norway	2010	Male	85 to 89	94% (89-97%)	6.1% (3.1-11%)	2.6 (1.7-3.9)	0.74% (-1.5-3.6%)	0.51% (-2.2-3.8%)	18 (-79-140)	250 (-510-1200)
Norway	2010	Male	90 to 94	94% (89-97%)	6% (3-11%)	2.6 (1.6-4)	0.37% (-2.1-3.9%)	0.049% (-2.9-4%)	0.97 (-58-80)	46 (-260-470)
Norway	2010	Male	95 plus	94% (88-98%)	5.9% (2.3-12%)	2.6 (1.3-4.4)	0.37% (-2.7-5.5%)	-0.0011% (-3.4-5.8%)	-0.0028 (-19-33)	8.5 (-62-130)
Norway	2016	Female	All Ages	91% (89-93%)	8.6% (6.7-11%)	2.1 (1.6-2.8)	2.4% (1.1-4%)	1.8% (-0.9-5.4%)	380 (-200-1200)	14000 (6200-23000)
Norway	2016	Female	15 to 19	90% (84-95%)	9.9% (5.5-16%)	1.6 (0.84-2.8)	2.1% (1.3-3.3%)	7.9% (3-14%)	2.3 (0.84-4.1)	330 (200-500)
Norway	2016	Female	20 to 24	94% (89-97%)	6.3% (3.3-11%)	2.1 (1-3.5)	3.3% (2.1-4.7%)	8.7% (3.2-15%)	3.3 (1.1-5.8)	630 (390-930)
Norway	2016	Female	25 to 29	95% (91-97%)	4.8% (2.6-8.7%)	2.1 (1.1-3.6)	2.7% (1.8-3.9%)	8.3% (3.2-14%)	3.5 (1.3-5.9)	600 (370-870)
Norway	2016	Female	30 to 34	96% (92-98%)	4.2% (2.1-7.7%)	2.3 (1.2-3.7)	2.4% (1.5-3.6%)	8.9% (4.4-14%)	4.9 (2.4-8)	550 (330-820)
Norway	2016	Female	35 to 39	96% (92-98%)	4.4% (2.1-8.3%)	2.4 (1.2-4.1)	2.5% (1.5-3.9%)	9.6% (5.3-14%)	7.2 (3.9-11)	600 (340-890)
Norway	2016	Female	40 to 44	95% (91-98%)	4.8% (2.4-8.8%)	2.7 (1.5-4.3)	2.9% (1.6-4.6%)	10% (6.2-15%)	13 (7.5-19)	860 (470-1300)
Norway	2016	Female	45 to 49	95% (90-97%)	5.2% (2.5-9.9%)	3.1 (1.7-5)	4% (2.2-6.2%)	11% (7.2-16%)	27 (16-40)	1400 (780-2100)
Norway	2016	Female	50 to 54	94% (90-97%)	5.7% (3.1-10%)	3.3 (2-5.3)	4.7% (2.8-7.3%)	11% (7.4-15%)	40 (26-56)	1700 (1000-2600)
Norway	2016	Female	55 to 59	94% (89-97%)	6.2% (3.4-11%)	3.4 (2-5.4)	4.7% (2.5-7.5%)	9.6% (6.2-13%)	50 (31-72)	1800 (970-2900)
Norway	2016	Female	60 to 64	93% (87-96%)	7.1% (3.8-13%)	3.3 (1.7-5.3)	4% (1.7-7%)	7.3% (3.9-11%)	56 (30-90)	1700 (740-3000)
Norway	2016	Female	65 to 69	92% (86-96%)	8.2% (4.3-14%)	2.9 (1.7-4.4)	3.1% (0.94-5.7%)	5.3% (2.4-8.6%)	68 (31-110)	1700 (490-3100)
Norway	2016	Female	70 to 74	90% (84-94%)	9.9% (5.9-16%)	2.5 (1.3-4.1)	2% (-0.27-5%)	3.4% (0.45-7.1%)	49 (6-110)	940 (-120-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	2016	Female	75 to 79	88% (80-93%)	12% (6.7-20%)	2.3 (1.2-3.9)	1.3% (-1.1-4.7%)	2% (-1.1-6.3%)	38 (-20-120)	560 (-480-2000)
Norway	2016	Female	80 to 84	86% (77-92%)	14% (8-23%)	2.2 (1.2-3.6)	0.66% (-1.8-3.9%)	0.89% (-2.2-4.8%)	26 (-64-140)	290 (-810-1700)
Norway	2016	Female	85 to 89	84% (74-90%)	16% (9.8-26%)	2.2 (1.2-3.6)	0.25% (-2.4-3.9%)	0.23% (-2.9-4.6%)	9.5 (-120-200)	110 (-1000-1600)
Norway	2016	Female	90 to 94	82% (73-90%)	18% (10-27%)	2.2 (1.1-3.8)	-0.038% (-2.8-4.4%)	-0.22% (-3.6-5%)	-11 (-170-230)	-11 (-880-1300)
Norway	2016	Female	95 plus	81% (67-91%)	19% (8.8-33%)	2.1 (0.84-4.1)	0.0068% (-3.2-5.5%)	-0.22% (-4-6.3%)	-5.4 (-95-150)	0.86 (-310-540)
Norway	2016	Male	All Ages	94% (93-96%)	5.7% (4.2-7.3%)	2.5 (1.9-3.3)	4.9% (3.2-6.9%)	3.7% (1.2-6.7%)	760 (250-1400)	31000 (20000-44000)
Norway	2016	Male	15 to 19	90% (83-94%)	10% (5.7-17%)	1.7 (0.99-2.8)	4.4% (2.8-6.3%)	9.2% (3.7-16%)	5.1 (2-8.8)	710 (430-1000)
Norway	2016	Male	20 to 24	94% (89-97%)	5.9% (3.2-11%)	2.5 (1.4-4)	9% (6-12%)	12% (5.3-19%)	13 (6-22)	1900 (1200-2800)
Norway	2016	Male	25 to 29	96% (93-98%)	4% (2.2-7.2%)	2.7 (1.6-4.1)	8.2% (5.7-11%)	12% (5.6-19%)	15 (7.5-25)	2000 (1300-2700)
Norway	2016	Male	30 to 34	97% (94-99%)	3% (1.4-5.8%)	2.8 (1.7-4.3)	6.7% (4.5-9.1%)	11% (5.5-18%)	15 (7.5-25)	1700 (1100-2300)
Norway	2016	Male	35 to 39	97% (95-99%)	2.7% (1.3-4.9%)	2.7 (1.5-4.3)	6.2% (4.1-8.5%)	11% (6-17%)	18 (9.5-28)	1600 (1000-2200)
Norway	2016	Male	40 to 44	97% (95-99%)	2.6% (1.2-4.8%)	3 (1.8-4.6)	6.2% (4.3-8.6%)	12% (7.2-18%)	27 (16-41)	1900 (1300-2700)
Norway	2016	Male	45 to 49	97% (95-99%)	2.5% (1.2-5.3%)	3.5 (2.1-5.2)	7.2% (4.9-10%)	13% (8.1-18%)	46 (29-69)	2700 (1800-3900)
Norway	2016	Male	50 to 54	97% (95-99%)	2.6% (1.1-5.2%)	3.8 (2.4-5.6)	7.9% (5.4-11%)	13% (8.2-18%)	68 (42-100)	3300 (2200-4800)
Norway	2016	Male	55 to 59	97% (94-99%)	2.9% (1.3-5.6%)	4 (2.5-6.1)	7.9% (5.2-11%)	12% (7.5-17%)	95 (60-150)	3800 (2500-5700)
Norway	2016	Male	60 to 64	97% (94-99%)	3.4% (1.5-6.4%)	3.9 (2.3-6.2)	6.6% (3.8-11%)	9% (5.1-14%)	110 (59-180)	3600 (2000-5700)
Norway	2016	Male	65 to 69	96% (93-98%)	3.9% (1.7-7.1%)	3.5 (2.2-5.2)	4.8% (2.4-8%)	6.1% (3.1-10%)	120 (61-210)	3400 (1700-5800)
Norway	2016	Male	70 to 74	95% (92-98%)	4.5% (2.4-7.8%)	3.1 (1.8-4.7)	3.2% (0.78-6.5%)	3.8% (0.85-7.9%)	86 (19-180)	1900 (450-4000)
Norway	2016	Male	75 to 79	95% (91-98%)	5.1% (2.4-8.8%)	2.8 (1.6-4.4)	2.2% (-0.26-5.8%)	2.4% (-0.5-6.5%)	62 (-13-170)	1100 (-130-2800)
Norway	2016	Male	80 to 84	94% (89-97%)	5.7% (2.6-11%)	2.7 (1.6-4.2)	1.4% (-0.82-4.6%)	1.3% (-1.4-5%)	43 (-46-160)	610 (-370-2000)
Norway	2016	Male	85 to 89	94% (89-97%)	6.2% (3.1-11%)	2.6 (1.5-4.1)	0.93% (-1.3-4.1%)	0.75% (-1.9-4.4%)	26 (-64-150)	300 (-430-1300)
Norway	2016	Male	90 to 94	94% (89-97%)	6.1% (2.9-11%)	2.6 (1.5-4.1)	0.53% (-2.4-2%)	0.26% (-2.7-4.5%)	6.5 (-67-110)	83 (-300-660)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Norway	2016	Male	95 plus	94% (88-98%)	5.9% (2.2-12%)	2.5 (1.3-4.3)	0.43% (-2.5-5.2%)	0.091% (-3.3-5.7%)	0.68 (-25-43)	13 (-75-160)
Portugal	1990	Female	All Ages	47% (41-53%)	53% (47-59%)	3 (2.4-3.6)	8.7% (6.7-11%)	14% (10-18%)	6700 (4900-8800)	130000 (100000-160000)
Portugal	1990	Female	15 to 19	44% (31-60%)	56% (40-69%)	2 (1.1-3)	2.9% (1.7-4.4%)	9% (4.9-14%)	15 (8.3-24)	1400 (810-2100)
Portugal	1990	Female	20 to 24	51% (35-67%)	49% (33-65%)	2.2 (1.2-3.4)	3.4% (2.1-5.1%)	9.8% (5.6-15%)	20 (12-31)	1800 (1200-2600)
Portugal	1990	Female	25 to 29	54% (35-70%)	46% (30-65%)	2.1 (1.3-3.2)	3.3% (2.1-4.6%)	9.9% (6.2-14%)	23 (14-33)	1900 (1200-2600)
Portugal	1990	Female	30 to 34	57% (37-73%)	43% (27-63%)	2.7 (1.7-4)	4.2% (3-5.7%)	13% (9.6-17%)	36 (27-47)	2500 (1800-3300)
Portugal	1990	Female	35 to 39	59% (42-74%)	41% (26-58%)	4.1 (2.6-5.9)	6.3% (4.7-8.4%)	18% (14-23%)	67 (52-85)	4000 (3100-5000)
Portugal	1990	Female	40 to 44	60% (45-75%)	40% (25-55%)	4.9 (3.4-6.8)	8.3% (6.3-11%)	21% (17-25%)	110 (88-140)	5800 (4600-7200)
Portugal	1990	Female	45 to 49	62% (48-72%)	38% (28-52%)	5.2 (3.6-6.9)	9.9% (7.5-13%)	21% (17-27%)	160 (130-200)	7200 (5600-9100)
Portugal	1990	Female	50 to 54	58% (47-68%)	42% (32-53%)	5.1 (3.7-6.9)	11% (8.2-14%)	21% (17-26%)	230 (180-280)	9100 (7200-11000)
Portugal	1990	Female	55 to 59	51% (39-60%)	49% (40-61%)	4.8 (3.4-6.5)	11% (8.4-15%)	20% (15-25%)	320 (240-400)	11000 (8500-14000)
Portugal	1990	Female	60 to 64	48% (35-59%)	52% (41-65%)	4.6 (3.1-6.6)	11% (8.1-15%)	18% (13-23%)	420 (310-550)	13000 (9100-16000)
Portugal	1990	Female	65 to 69	51% (39-61%)	49% (39-61%)	4.6 (3.2-6.2)	12% (8.8-16%)	17% (13-22%)	640 (470-830)	16000 (11000-20000)
Portugal	1990	Female	70 to 74	48% (38-58%)	52% (42-62%)	4.1 (2.8-5.7)	12% (8.4-17%)	16% (11-22%)	820 (570-1100)	16000 (11000-22000)
Portugal	1990	Female	75 to 79	41% (29-53%)	59% (47-71%)	3.4 (2.1-5)	11% (7.2-16%)	13% (8.8-19%)	1100 (730-1600)	16000 (11000-24000)
Portugal	1990	Female	80 to 84	37% (26-48%)	63% (52-74%)	3.1 (1.9-4.4)	10% (6.4-15%)	12% (7.7-18%)	1300 (780-1800)	13000 (8400-19000)
Portugal	1990	Female	85 to 89	38% (26-50%)	62% (50-74%)	3.1 (1.9-4.7)	9.7% (5.7-15%)	11% (6.6-17%)	950 (550-1400)	7200 (4200-11000)
Portugal	1990	Female	90 to 94	39% (28-51%)	61% (49-72%)	3.1 (1.7-4.9)	9.2% (4.9-14%)	11% (5.7-17%)	400 (210-630)	2100 (1100-3300)
Portugal	1990	Female	95 plus	43% (28-60%)	57% (40-72%)	3.3 (1.6-5.7)	9.6% (4.3-16%)	11% (5-18%)	130 (57-210)	440 (200-710)
Portugal	1990	Male	All Ages	71% (65-77%)	29% (23-35%)	8.6 (7.1-10)	19% (16-22%)	24% (18-29%)	13000 (9700-15000)	330000 (270000-380000)
Portugal	1990	Male	15 to 19	61% (45-76%)	39% (24-55%)	3.7 (2.3-5.5)	12% (7.4-17%)	18% (11-26%)	110 (66-160)	8600 (5500-12000)
Portugal	1990	Male	20 to 24	70% (55-83%)	30% (17-45%)	6.3 (4.1-8.8)	21% (14-28%)	29% (19-40%)	220 (150-300)	17000 (12000-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	1990	Male	25 to 29	75% (60-86%)	25% (14-40%)	8.9 (6.3-12)	22% (17-29%)	33% (24-43%)	250 (180-320)	17000 (13000-22000)
Portugal	1990	Male	30 to 34	79% (64-89%)	21% (11-36%)	11 (7.8-14)	22% (17-27%)	34% (27-41%)	260 (210-320)	17000 (13000-20000)
Portugal	1990	Male	35 to 39	82% (70-90%)	18% (9.6-30%)	12 (8.4-16)	22% (18-26%)	34% (28-40%)	290 (230-340)	16000 (13000-19000)
Portugal	1990	Male	40 to 44	83% (71-92%)	17% (8.3-29%)	13 (9.9-18)	22% (18-26%)	34% (29-39%)	360 (300-410)	18000 (15000-21000)
Portugal	1990	Male	45 to 49	84% (73-91%)	16% (9.5-27%)	15 (12-20)	23% (19-27%)	33% (28-38%)	450 (380-520)	20000 (17000-23000)
Portugal	1990	Male	50 to 54	85% (76-90%)	15% (9.5-24%)	16 (12-20)	24% (20-28%)	32% (27-37%)	670 (560-780)	26000 (22000-30000)
Portugal	1990	Male	55 to 59	87% (80-91%)	13% (8.7-20%)	15 (11-19)	23% (19-28%)	30% (24-35%)	960 (770-1100)	32000 (26000-38000)
Portugal	1990	Male	60 to 64	86% (78-90%)	14% (9.5-22%)	14 (10-18)	22% (17-26%)	27% (21-32%)	1200 (940-1500)	35000 (27000-42000)
Portugal	1990	Male	65 to 69	82% (75-87%)	18% (13-25%)	13 (9.7-16)	21% (16-26%)	25% (19-30%)	1500 (1100-1900)	36000 (27000-44000)
Portugal	1990	Male	70 to 74	80% (72-85%)	20% (15-28%)	12 (8.8-16)	20% (15-25%)	24% (17-29%)	1600 (1200-2000)	31000 (22000-38000)
Portugal	1990	Male	75 to 79	78% (69-86%)	22% (14-31%)	11 (8-15)	19% (13-25%)	22% (15-28%)	1900 (1300-2400)	28000 (19000-35000)
Portugal	1990	Male	80 to 84	77% (66-85%)	23% (15-34%)	10 (7.1-13)	19% (13-24%)	21% (14-27%)	1600 (1100-2100)	17000 (12000-23000)
Portugal	1990	Male	85 to 89	75% (64-84%)	25% (16-36%)	9 (6.2-12)	18% (12-23%)	20% (13-26%)	890 (610-1200)	6800 (4700-8900)
Portugal	1990	Male	90 to 94	77% (66-85%)	23% (15-34%)	8.3 (5.4-12)	17% (12-22%)	19% (13-24%)	280 (190-360)	1500 (1000-1900)
Portugal	1990	Male	95 plus	78% (65-88%)	22% (12-35%)	7.7 (4.4-12)	16% (10-22%)	18% (11-24%)	58 (36-79)	200 (130-280)
Portugal	1995	Female	All Ages	47% (42-53%)	53% (47-58%)	2.8 (2.3-3.4)	7.8% (6-10%)	12% (9-16%)	6200 (4500-8000)	110000 (91000-140000)
Portugal	1995	Female	15 to 19	44% (30-59%)	56% (41-70%)	1.8 (1-2.7)	2.5% (1.4-3.9%)	8.1% (4.3-13%)	12 (6.3-19)	1100 (650-1700)
Portugal	1995	Female	20 to 24	50% (34-67%)	50% (33-66%)	1.9 (1.1-3.1)	2.8% (1.7-4.2%)	7.9% (4.5-12%)	16 (9-25)	1500 (980-2300)
Portugal	1995	Female	25 to 29	54% (35-70%)	46% (30-65%)	1.9 (1.2-2.8)	2.7% (1.7-3.8%)	7.7% (4.7-11%)	18 (11-25)	1500 (1000-2100)
Portugal	1995	Female	30 to 34	56% (36-73%)	44% (27-64%)	2.5 (1.6-3.6)	3.7% (2.6-4.9%)	11% (8-14%)	33 (24-43)	2300 (1700-3000)
Portugal	1995	Female	35 to 39	58% (42-73%)	42% (27-58%)	3.7 (2.4-5.2)	5.4% (3.9-7.2%)	16% (12-19%)	58 (44-73)	3500 (2700-4400)
Portugal	1995	Female	40 to 44	60% (45-74%)	40% (26-55%)	4.5 (3.1-6.2)	7.3% (5.6-9.5%)	19% (15-23%)	99 (79-120)	5300 (4100-6500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	1995	Female	45 to 49	61% (48-72%)	39% (28-52%)	4.8 (3.3-6.4)	8.5% (6.4-11%)	19% (15-24%)	140 (110-170)	6500 (5100-8100)
Portugal	1995	Female	50 to 54	58% (46-68%)	42% (32-54%)	4.7 (3.3-6.3)	9.7% (7.2-12%)	19% (15-24%)	190 (150-240)	7600 (5900-9700)
Portugal	1995	Female	55 to 59	50% (39-60%)	50% (40-61%)	4.4 (3.1-6.1)	9.9% (7.3-13%)	18% (14-23%)	250 (190-330)	8900 (6800-12000)
Portugal	1995	Female	60 to 64	48% (35-59%)	52% (41-65%)	4.2 (2.8-6)	10% (7.1-13%)	16% (12-21%)	360 (270-470)	11000 (7800-14000)
Portugal	1995	Female	65 to 69	50% (38-60%)	50% (40-62%)	4.2 (2.9-5.6)	11% (7.6-14%)	16% (12-20%)	550 (400-710)	14000 (9800-18000)
Portugal	1995	Female	70 to 74	48% (37-58%)	52% (42-63%)	3.8 (2.6-5.2)	11% (7.4-15%)	15% (10-20%)	810 (560-1100)	16000 (11000-22000)
Portugal	1995	Female	75 to 79	40% (30-53%)	60% (47-70%)	3.1 (2-4.6)	9.6% (6.1-14%)	12% (7.8-18%)	890 (580-1300)	13000 (8600-19000)
Portugal	1995	Female	80 to 84	37% (26-49%)	63% (51-74%)	2.8 (1.8-4.1)	9% (5.5-13%)	11% (6.6-16%)	1100 (690-1700)	12000 (7500-18000)
Portugal	1995	Female	85 to 89	37% (26-49%)	63% (51-74%)	2.8 (1.7-4.2)	8.7% (5-13%)	10% (5.8-16%)	960 (550-1500)	7300 (4200-11000)
Portugal	1995	Female	90 to 94	39% (28-52%)	61% (48-72%)	2.8 (1.6-4.5)	8.2% (4.1-13%)	9.7% (4.7-15%)	470 (230-750)	2500 (1200-3900)
Portugal	1995	Female	95 plus	43% (27-60%)	57% (40-73%)	3.1 (1.5-5.3)	8.7% (3.6-14%)	10% (4.1-17%)	150 (61-250)	520 (210-870)
Portugal	1995	Male	All Ages	72% (65-78%)	28% (22-35%)	8.3 (7-10)	18% (15-21%)	23% (18-27%)	13000 (10000-15000)	320000 (260000-370000)
Portugal	1995	Male	15 to 19	61% (44-76%)	39% (24-56%)	3.5 (2.1-5.2)	10% (6.4-15%)	16% (9.5-25%)	80 (47-120)	6500 (4100-9400)
Portugal	1995	Male	20 to 24	70% (55-83%)	30% (17-45%)	5.9 (4-8.2)	18% (13-24%)	25% (16-35%)	190 (130-270)	15000 (11000-20000)
Portugal	1995	Male	25 to 29	76% (62-86%)	24% (14-38%)	8.2 (5.8-11)	18% (14-23%)	25% (18-33%)	210 (160-280)	15000 (12000-19000)
Portugal	1995	Male	30 to 34	79% (64-89%)	21% (11-36%)	10 (7.3-13)	18% (14-22%)	27% (21-32%)	240 (190-300)	16000 (12000-19000)
Portugal	1995	Male	35 to 39	82% (69-90%)	18% (9.7-31%)	11 (7.9-15)	20% (16-24%)	29% (24-34%)	290 (240-350)	17000 (14000-20000)
Portugal	1995	Male	40 to 44	83% (71-91%)	17% (8.6-29%)	12 (9.3-16)	21% (17-25%)	32% (27-36%)	370 (310-430)	19000 (16000-22000)
Portugal	1995	Male	45 to 49	84% (73-91%)	16% (9.3-27%)	14 (11-18)	22% (18-26%)	32% (27-36%)	480 (400-560)	22000 (18000-25000)
Portugal	1995	Male	50 to 54	85% (76-91%)	15% (9.3-24%)	15 (11-19)	22% (18-26%)	31% (26-35%)	580 (480-670)	23000 (19000-26000)
Portugal	1995	Male	55 to 59	86% (79-91%)	14% (8.7-21%)	14 (11-18)	22% (18-26%)	29% (24-34%)	840 (680-980)	28000 (23000-33000)
Portugal	1995	Male	60 to 64	85% (78-90%)	15% (9.7-22%)	13 (9.6-17)	21% (17-25%)	26% (20-31%)	1100 (900-1400)	33000 (26000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	1995	Male	65 to 69	82% (74-87%)	18% (13-26%)	12 (9.2-15)	20% (15-24%)	24% (18-29%)	1500 (1100-1800)	35000 (27000-42000)
Portugal	1995	Male	70 to 74	79% (71-85%)	21% (15-29%)	11 (8.2-14)	20% (14-24%)	23% (17-28%)	1800 (1300-2200)	34000 (26000-42000)
Portugal	1995	Male	75 to 79	78% (68-86%)	22% (14-32%)	10 (7.5-14)	19% (13-23%)	21% (15-26%)	1700 (1200-2100)	25000 (18000-31000)
Portugal	1995	Male	80 to 84	76% (65-85%)	24% (15-35%)	9.3 (6.5-12)	18% (13-23%)	20% (14-26%)	1700 (1200-2200)	19000 (13000-24000)
Portugal	1995	Male	85 to 89	75% (63-84%)	25% (16-37%)	8.3 (5.7-11)	17% (12-22%)	19% (14-25%)	1000 (750-1400)	8000 (5800-10000)
Portugal	1995	Male	90 to 94	76% (65-85%)	24% (15-35%)	7.7 (5.1-11)	16% (11-21%)	18% (13-23%)	370 (270-480)	1900 (1400-2500)
Portugal	1995	Male	95 plus	78% (63-89%)	22% (11-37%)	7.1 (4.1-11)	15% (9.7-21%)	17% (11-24%)	75 (47-100)	260 (170-360)
Portugal	2000	Female	All Ages	45% (40-51%)	55% (49-60%)	2.4 (2-2.9)	6.4% (4.8-8.3%)	9.9% (6.9-14%)	5100 (3500-7000)	93000 (72000-120000)
Portugal	2000	Female	15 to 19	38% (26-54%)	62% (46-74%)	1.4 (0.77-2.1)	1.8% (1.1-2.8%)	6.1% (3.2-9.9%)	6.7 (3.5-11)	680 (400-1000)
Portugal	2000	Female	20 to 24	45% (30-62%)	55% (38-70%)	1.5 (0.87-2.5)	2.2% (1.4-3.2%)	6% (3.4-9.5%)	9.8 (5.4-15)	1100 (700-1600)
Portugal	2000	Female	25 to 29	51% (34-68%)	49% (32-66%)	1.6 (0.95-2.4)	2.1% (1.3-2.8%)	6.1% (3.7-8.6%)	12 (6.8-16)	1100 (770-1600)
Portugal	2000	Female	30 to 34	56% (38-72%)	44% (28-62%)	2.2 (1.4-3.1)	2.9% (2.1-3.9%)	8.8% (6.4-11%)	24 (17-31)	1700 (1300-2300)
Portugal	2000	Female	35 to 39	59% (44-74%)	41% (26-56%)	3.3 (2.2-4.6)	4.6% (3.4-6.1%)	14% (10-17%)	52 (40-65)	3200 (2400-4000)
Portugal	2000	Female	40 to 44	59% (45-73%)	41% (27-55%)	3.9 (2.7-5.4)	6.2% (4.7-8.1%)	16% (13-19%)	86 (69-110)	4600 (3600-5600)
Portugal	2000	Female	45 to 49	59% (46-70%)	41% (30-54%)	4.1 (2.9-5.3)	7.2% (5.4-9.4%)	17% (13-21%)	120 (97-150)	5700 (4600-7300)
Portugal	2000	Female	50 to 54	55% (45-66%)	45% (34-55%)	4 (2.8-5.4)	7.9% (6-10%)	16% (13-21%)	160 (130-210)	6700 (5100-8600)
Portugal	2000	Female	55 to 59	48% (38-57%)	52% (43-62%)	3.7 (2.6-5.1)	8% (5.9-11%)	15% (12-20%)	190 (140-240)	6800 (5100-8900)
Portugal	2000	Female	60 to 64	45% (34-56%)	55% (44-66%)	3.5 (2.4-5)	8% (5.7-11%)	14% (9.7-18%)	270 (190-360)	8200 (5600-11000)
Portugal	2000	Female	65 to 69	47% (36-57%)	53% (43-64%)	3.5 (2.4-4.8)	8.3% (5.8-11%)	13% (9.2-17%)	410 (290-540)	10000 (7300-14000)
Portugal	2000	Female	70 to 74	44% (34-55%)	56% (45-66%)	3.1 (2.1-4.2)	8.5% (5.5-12%)	12% (7.9-16%)	630 (420-880)	12000 (8200-18000)
Portugal	2000	Female	75 to 79	39% (28-51%)	61% (49-72%)	2.6 (1.7-3.9)	7.6% (4.7-12%)	9.8% (6-15%)	790 (480-1200)	12000 (7300-18000)
Portugal	2000	Female	80 to 84	37% (26-47%)	63% (53-74%)	2.5 (1.6-3.6)	7.3% (4.3-11%)	9% (5.3-14%)	840 (490-1300)	9200 (5400-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	2000	Female	85 to 89	37% (26-48%)	63% (52-74%)	2.4 (1.5-3.7)	7.1% (3.8-11%)	8.4% (4.5-13%)	870 (470-1400)	6700 (3600-11000)
Portugal	2000	Female	90 to 94	39% (27-51%)	61% (49-73%)	2.5 (1.4-4)	6.7% (3-11%)	7.8% (3.4-13%)	460 (200-770)	2400 (1100-4000)
Portugal	2000	Female	95 plus	43% (27-60%)	57% (40-73%)	2.7 (1.3-4.6)	7% (2.3-13%)	8.1% (2.5-15%)	170 (50-300)	570 (180-1000)
Portugal	2000	Male	All Ages	74% (68-80%)	26% (20-32%)	7.8 (6.5-9.3)	18% (15-21%)	22% (18-26%)	12000 (10000-14000)	300000 (250000-340000)
Portugal	2000	Male	15 to 19	60% (44-75%)	40% (25-56%)	3 (1.8-4.5)	8.4% (5.3-12%)	15% (8.4-22%)	47 (27-70)	4000 (2600-5700)
Portugal	2000	Male	20 to 24	71% (56-83%)	29% (17-44%)	5.2 (3.6-7.3)	16% (11-22%)	23% (15-32%)	140 (91-190)	11000 (8100-15000)
Portugal	2000	Male	25 to 29	77% (65-87%)	23% (13-35%)	7.5 (5.3-9.8)	17% (13-21%)	24% (18-32%)	180 (130-240)	13000 (10000-17000)
Portugal	2000	Male	30 to 34	82% (69-90%)	18% (9.7-31%)	9.2 (6.8-12)	17% (14-21%)	26% (20-31%)	220 (170-270)	14000 (12000-17000)
Portugal	2000	Male	35 to 39	84% (74-91%)	16% (8.6-26%)	10 (7.2-13)	18% (15-22%)	28% (23-33%)	290 (240-340)	17000 (14000-19000)
Portugal	2000	Male	40 to 44	85% (75-92%)	15% (7.9-25%)	11 (8.5-15)	21% (17-24%)	31% (27-36%)	390 (330-440)	20000 (17000-23000)
Portugal	2000	Male	45 to 49	86% (78-91%)	14% (8.6-22%)	13 (9.9-17)	22% (18-26%)	32% (27-36%)	500 (430-570)	22000 (19000-26000)
Portugal	2000	Male	50 to 54	86% (79-91%)	14% (8.9-21%)	13 (10-17)	22% (18-25%)	30% (25-34%)	590 (490-680)	23000 (19000-27000)
Portugal	2000	Male	55 to 59	87% (82-92%)	13% (8.4-18%)	12 (9.3-16)	21% (17-25%)	28% (23-32%)	700 (590-820)	24000 (20000-28000)
Portugal	2000	Male	60 to 64	86% (81-91%)	14% (9.2-19%)	12 (8.6-15)	20% (16-23%)	25% (21-29%)	960 (780-1100)	28000 (23000-33000)
Portugal	2000	Male	65 to 69	83% (78-88%)	17% (12-22%)	11 (8.3-14)	19% (15-23%)	23% (19-28%)	1300 (1100-1600)	32000 (26000-39000)
Portugal	2000	Male	70 to 74	81% (75-86%)	19% (14-25%)	10 (7.6-13)	19% (15-23%)	22% (17-26%)	1700 (1300-2000)	32000 (25000-39000)
Portugal	2000	Male	75 to 79	79% (71-86%)	21% (14-29%)	9.3 (6.7-12)	18% (13-22%)	20% (15-25%)	1900 (1400-2300)	28000 (21000-34000)
Portugal	2000	Male	80 to 84	77% (67-85%)	23% (15-33%)	8.2 (5.8-11)	17% (12-21%)	19% (14-24%)	1500 (1200-2000)	17000 (13000-21000)
Portugal	2000	Male	85 to 89	76% (64-85%)	24% (15-36%)	7.4 (5-9.8)	16% (11-20%)	18% (13-23%)	1100 (830-1500)	8700 (6500-11000)
Portugal	2000	Male	90 to 94	77% (66-85%)	23% (15-34%)	6.8 (4.5-9.6)	15% (10-19%)	16% (11-22%)	440 (310-580)	2300 (1600-3000)
Portugal	2000	Male	95 plus	78% (64-88%)	22% (12-36%)	6.3 (3.6-9.7)	14% (7.5-19%)	15% (8.3-22%)	90 (48-130)	320 (170-450)
Portugal	2005	Female	All Ages	44% (40-49%)	56% (51-60%)	2.2 (1.8-2.7)	5.2% (3.7-6.8%)	7.8% (5.1-11%)	4000 (2600-5500)	74000 (54000-95000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	2005	Female	15 to 19	35% (25-49%)	65% (51-75%)	1.1 (0.66-1.8)	1.5% (0.91-2.2%)	5.4% (2.9-8.5%)	4 (2.1-6.4)	440 (270-670)
Portugal	2005	Female	20 to 24	45% (32-60%)	55% (40-68%)	1.4 (0.83-2.3)	1.9% (1.3-2.7%)	5.8% (3.3-8.8%)	6.4 (3.6-9.8)	800 (530-1100)
Portugal	2005	Female	25 to 29	53% (38-68%)	47% (32-62%)	1.6 (0.99-2.3)	1.8% (1.2-2.4%)	5.9% (3.7-8.4%)	8.9 (5.5-13)	960 (660-1300)
Portugal	2005	Female	30 to 34	58% (42-72%)	42% (28-58%)	2.1 (1.4-2.9)	2.4% (1.7-3.2%)	8.1% (5.9-11%)	19 (13-24)	1500 (1100-1900)
Portugal	2005	Female	35 to 39	59% (46-73%)	41% (27-54%)	3.1 (2.1-4.3)	3.9% (2.9-5.2%)	13% (9.8-16%)	41 (31-51)	2600 (2000-3300)
Portugal	2005	Female	40 to 44	58% (45-71%)	42% (29-55%)	3.6 (2.5-4.8)	5.1% (3.8-6.6%)	15% (12-18%)	71 (56-87)	3900 (3000-4800)
Portugal	2005	Female	45 to 49	57% (47-66%)	43% (34-53%)	3.6 (2.7-4.9)	6.1% (4.6-8.2%)	15% (12-19%)	100 (83-130)	5000 (4000-6300)
Portugal	2005	Female	50 to 54	53% (45-61%)	47% (39-55%)	3.5 (2.6-4.7)	6.6% (4.9-8.8%)	14% (11-18%)	130 (100-170)	5600 (4300-7100)
Portugal	2005	Female	55 to 59	46% (39-53%)	54% (47-61%)	3.3 (2.4-4.5)	6.7% (4.9-8.8%)	13% (10-17%)	160 (130-210)	6000 (4500-7800)
Portugal	2005	Female	60 to 64	43% (36-50%)	57% (50-64%)	3.1 (2.2-4.2)	6.6% (4.6-9%)	12% (8.7-16%)	200 (140-260)	6200 (4400-8300)
Portugal	2005	Female	65 to 69	43% (36-51%)	57% (49-64%)	3 (2.2-3.9)	6.6% (4.6-9.1%)	11% (7.8-14%)	300 (220-390)	7600 (5400-10000)
Portugal	2005	Female	70 to 74	40% (33-48%)	60% (52-67%)	2.6 (1.8-3.5)	6.5% (4.9-3%)	9.5% (6-13%)	450 (290-630)	9100 (5800-13000)
Portugal	2005	Female	75 to 79	35% (27-43%)	65% (57-73%)	2.2 (1.4-3)	5.7% (3.4-8.6%)	7.6% (4.5-11%)	580 (340-860)	8900 (5200-13000)
Portugal	2005	Female	80 to 84	32% (23-42%)	68% (58-77%)	2 (1.2-2.8)	5.5% (3-8.3%)	6.8% (3.8-10%)	730 (400-1100)	8100 (4500-12000)
Portugal	2005	Female	85 to 89	33% (23-44%)	67% (56-77%)	2 (1.2-3.1)	5.3% (2.8-8.6%)	6.4% (3.3-10%)	610 (310-990)	4800 (2500-7700)
Portugal	2005	Female	90 to 94	35% (24-47%)	65% (53-76%)	2.1 (1.2-3.4)	5% (2.2-8.8%)	5.9% (2.5-11%)	410 (170-720)	2200 (930-3800)
Portugal	2005	Female	95 plus	39% (25-55%)	61% (45-75%)	2.3 (1.1-4)	5.4% (1.6-10%)	6.3% (1.7-12%)	170 (45-320)	580 (170-1100)
Portugal	2005	Male	All Ages	76% (71-81%)	24% (19-29%)	7.7 (6.3-9.2)	17% (14-20%)	21% (17-24%)	11000 (9500-13000)	270000 (240000-310000)
Portugal	2005	Male	15 to 19	61% (48-75%)	39% (25-52%)	2.9 (1.8-4.3)	7.5% (4.9-10%)	15% (8.7-21%)	29 (17-42)	2600 (1700-3600)
Portugal	2005	Male	20 to 24	73% (60-84%)	27% (16-40%)	5.2 (3.6-7.1)	16% (11-21%)	25% (17-35%)	87 (58-120)	7600 (5600-10000)
Portugal	2005	Male	25 to 29	79% (67-88%)	21% (12-33%)	7.2 (5.2-9.6)	16% (12-21%)	28% (20-36%)	130 (90-160)	9900 (7500-12000)
Portugal	2005	Male	30 to 34	82% (70-91%)	18% (9.2-30%)	8.8 (6.4-11)	15% (12-19%)	26% (20-31%)	170 (130-210)	12000 (9200-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	2005	Male	35 to 39	85% (74-92%)	15% (8-26%)	9.6 (6.9-13)	17% (13-20%)	27% (22-32%)	230 (190-280)	14000 (11000-16000)
Portugal	2005	Male	40 to 44	86% (76-92%)	14% (7.6-24%)	11 (8-14)	19% (16-22%)	30% (26-33%)	370 (320-420)	19000 (16000-21000)
Portugal	2005	Male	45 to 49	87% (80-92%)	13% (8.1-20%)	13 (9.5-16)	21% (18-25%)	31% (27-35%)	510 (440-580)	23000 (20000-26000)
Portugal	2005	Male	50 to 54	87% (82-92%)	13% (8.2-18%)	13 (9.6-17)	22% (18-25%)	30% (26-34%)	630 (530-710)	25000 (21000-28000)
Portugal	2005	Male	55 to 59	89% (84-92%)	11% (7.8-16%)	12 (9-15)	20% (17-24%)	27% (23-31%)	730 (620-840)	25000 (21000-29000)
Portugal	2005	Male	60 to 64	87% (83-91%)	13% (9-17%)	11 (8.1-14)	19% (16-22%)	25% (21-28%)	810 (690-930)	24000 (20000-28000)
Portugal	2005	Male	65 to 69	84% (78-88%)	16% (12-22%)	10 (7.9-13)	18% (15-21%)	23% (19-26%)	1100 (930-1300)	27000 (23000-32000)
Portugal	2005	Male	70 to 74	82% (77-86%)	18% (14-23%)	9.8 (7.4-12)	18% (14-21%)	21% (17-25%)	1500 (1200-1800)	29000 (24000-35000)
Portugal	2005	Male	75 to 79	81% (74-87%)	19% (13-26%)	9.1 (6.8-12)	17% (13-20%)	19% (15-23%)	1700 (1400-2100)	26000 (21000-32000)
Portugal	2005	Male	80 to 84	79% (69-86%)	21% (14-31%)	8 (5.7-10)	16% (12-19%)	18% (14-22%)	1700 (1300-2100)	19000 (14000-23000)
Portugal	2005	Male	85 to 89	76% (65-85%)	24% (15-35%)	7 (4.8-9.4)	14% (11-18%)	16% (12-21%)	1000 (740-1300)	7800 (5900-10000)
Portugal	2005	Male	90 to 94	77% (67-86%)	23% (14-33%)	6.5 (4.3-9.1)	13% (9-17%)	15% (9.9-20%)	480 (320-640)	2600 (1800-3400)
Portugal	2005	Male	95 plus	79% (65-89%)	21% (11-35%)	6 (3.4-9.3)	12% (6.1-18%)	14% (6.6-20%)	110 (53-160)	400 (190-570)
Portugal	2010	Female	All Ages	46% (42-50%)	54% (50-58%)	2.2 (1.7-2.8)	4.5% (3.1-6.2%)	6.6% (4.1-9.5%)	3400 (2100-5000)	64000 (45000-85000)
Portugal	2010	Female	15 to 19	39% (28-54%)	61% (46-72%)	1.2 (0.7-2)	1.3% (0.8-1.9%)	5.3% (3-8.4%)	2.9 (1.7-4.6)	370 (230-540)
Portugal	2010	Female	20 to 24	49% (36-64%)	51% (36-64%)	1.5 (0.88-2.4)	1.9% (1.3-2.7%)	6.1% (3.6-9.5%)	5.4 (3.1-8.3)	680 (460-960)
Portugal	2010	Female	25 to 29	57% (42-71%)	43% (29-58%)	1.6 (1-2.4)	1.7% (1.2-2.3%)	6.2% (3.9-8.8%)	6.3 (4-8.9)	750 (510-1000)
Portugal	2010	Female	30 to 34	60% (45-74%)	40% (26-55%)	2.1 (1.4-3)	2.2% (1.5-3%)	8.5% (6.1-11%)	16 (11-20)	1300 (920-1700)
Portugal	2010	Female	35 to 39	60% (47-73%)	40% (27-53%)	3 (2-4.2)	3.3% (2.5-4.5%)	12% (9-15%)	35 (27-44)	2300 (1700-3000)
Portugal	2010	Female	40 to 44	59% (47-71%)	41% (29-53%)	3.5 (2.4-4.7)	4.5% (3.3-5.9%)	14% (11-17%)	60 (46-74)	3400 (2500-4200)
Portugal	2010	Female	45 to 49	60% (52-68%)	40% (32-48%)	3.7 (2.6-5)	5.6% (4.1-7.4%)	15% (11-19%)	97 (76-120)	4700 (3600-6000)
Portugal	2010	Female	50 to 54	55% (48-62%)	45% (38-52%)	3.6 (2.6-4.8)	6% (4.4-8%)	14% (11-18%)	120 (97-160)	5200 (4000-6700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	2010	Female	55 to 59	45% (39-51%)	55% (49-61%)	3.1 (2.2-4.2)	5.8% (4.2-7.7%)	12% (9.4-16%)	140 (110-180)	5200 (3900-7000)
Portugal	2010	Female	60 to 64	43% (37-49%)	57% (51-63%)	3 (2.1-4.1)	5.7% (3.9-7.9%)	11% (7.9-14%)	170 (130-230)	5600 (3800-7600)
Portugal	2010	Female	65 to 69	45% (38-51%)	55% (49-62%)	2.9 (2.1-4)	5.8% (3.9-8.1%)	9.9% (7-13%)	230 (160-300)	6000 (4100-8200)
Portugal	2010	Female	70 to 74	40% (34-47%)	60% (53-66%)	2.5 (1.7-3.5)	5.5% (3.4-8.2%)	8.4% (5.4-12%)	340 (220-490)	7000 (4400-10000)
Portugal	2010	Female	75 to 79	33% (26-41%)	67% (59-74%)	2 (1.3-2.8)	4.7% (2.7-7.3%)	6.5% (3.8-9.8%)	450 (260-690)	7100 (4000-11000)
Portugal	2010	Female	80 to 84	31% (22-40%)	69% (60-78%)	1.8 (1.1-2.6)	4.5% (2.3-7.1%)	5.7% (2.9-9%)	600 (310-950)	6800 (3400-11000)
Portugal	2010	Female	85 to 89	31% (22-42%)	69% (58-78%)	1.8 (1.1-2.8)	4.4% (2-7.4%)	5.3% (2.3-9.1%)	620 (280-1100)	4900 (2200-8300)
Portugal	2010	Female	90 to 94	33% (23-44%)	67% (56-77%)	1.9 (1-3)	4.2% (1.6-7.5%)	4.9% (1.8-9%)	350 (130-640)	1900 (710-3400)
Portugal	2010	Female	95 plus	37% (23-52%)	63% (48-77%)	2 (0.98-3.5)	4.5% (1.2-8.8%)	5.3% (1.3-10%)	190 (47-380)	670 (180-1300)
Portugal	2010	Male	All Ages	76% (72-81%)	24% (19-28%)	7.4 (5.8-9.2)	16% (14-18%)	19% (17-22%)	11000 (9000-12000)	250000 (210000-280000)
Portugal	2010	Male	15 to 19	62% (48-76%)	38% (24-52%)	2.8 (1.7-4.3)	5.9% (4.1-8.2%)	14% (8.4-20%)	16 (9.9-23)	1700 (1200-2300)
Portugal	2010	Male	20 to 24	75% (61-85%)	25% (15-39%)	5 (3.3-7.1)	14% (10-18%)	24% (16-33%)	49 (32-68)	4800 (3500-6200)
Portugal	2010	Male	25 to 29	79% (68-87%)	21% (13-32%)	6.8 (4.7-9.3)	14% (11-18%)	28% (20-36%)	74 (53-96)	6300 (4800-7900)
Portugal	2010	Male	30 to 34	82% (70-90%)	18% (9.8-30%)	8.1 (5.6-11)	14% (11-17%)	27% (21-33%)	120 (90-140)	8400 (6700-10000)
Portugal	2010	Male	35 to 39	84% (74-91%)	16% (8.8-26%)	8.9 (6.2-12)	15% (12-18%)	27% (23-32%)	180 (150-220)	11000 (9200-13000)
Portugal	2010	Male	40 to 44	85% (76-92%)	15% (8.1-24%)	10 (7.3-14)	18% (15-21%)	30% (26-34%)	310 (260-350)	16000 (14000-18000)
Portugal	2010	Male	45 to 49	86% (80-91%)	14% (8.6-20%)	12 (8.7-15)	20% (17-23%)	31% (27-34%)	470 (420-530)	21000 (19000-24000)
Portugal	2010	Male	50 to 54	88% (82-92%)	12% (8.3-18%)	12 (8.9-16)	21% (18-24%)	29% (26-33%)	630 (550-700)	25000 (22000-28000)
Portugal	2010	Male	55 to 59	89% (85-92%)	11% (7.7-15%)	11 (8.4-15)	19% (17-23%)	27% (23-30%)	720 (630-820)	25000 (22000-28000)
Portugal	2010	Male	60 to 64	88% (83-91%)	12% (9.1-17%)	10 (7.6-14)	18% (15-21%)	24% (20-27%)	810 (690-930)	24000 (20000-28000)
Portugal	2010	Male	65 to 69	83% (78-87%)	17% (13-22%)	9.6 (7.1-12)	17% (14-20%)	21% (18-25%)	920 (780-1100)	23000 (19000-27000)
Portugal	2010	Male	70 to 74	82% (77-86%)	18% (14-23%)	9.2 (6.8-12)	16% (13-19%)	20% (17-23%)	1200 (1000-1500)	25000 (20000-29000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	2010	Male	75 to 79	82% (76-87%)	18% (13-24%)	8.7 (6.4-12)	15% (12-19%)	18% (15-22%)	1600 (1300-1900)	24000 (19000-29000)
Portugal	2010	Male	80 to 84	79% (71-86%)	21% (14-29%)	7.7 (5.5-10)	14% (11-18%)	17% (13-21%)	1600 (1200-2000)	18000 (14000-22000)
Portugal	2010	Male	85 to 89	77% (67-85%)	23% (15-33%)	6.7 (4.5-9.3)	13% (9.4-17%)	15% (11-20%)	1200 (850-1600)	9500 (6700-12000)
Portugal	2010	Male	90 to 94	78% (68-86%)	22% (14-32%)	6.2 (4-9)	12% (7.5-16%)	14% (8.3-19%)	480 (290-660)	2600 (1600-3500)
Portugal	2010	Male	95 plus	79% (66-89%)	21% (11-34%)	5.7 (3.2-8.9)	12% (5.4-17%)	13% (5.8-19%)	150 (66-210)	520 (240-750)
Portugal	2016	Female	All Ages	47% (42-52%)	53% (48-58%)	2.3 (1.6-3.2)	4.2% (2.6-6.3%)	6.4% (3.4-10%)	3500 (1800-5600)	61000 (38000-90000)
Portugal	2016	Female	15 to 19	44% (31-59%)	56% (41-69%)	1.3 (0.71-2.3)	1.2% (0.76-1.8%)	5.5% (3-8.8%)	2.4 (1.3-3.8)	320 (190-490)
Portugal	2016	Female	20 to 24	51% (35-66%)	49% (34-65%)	1.5 (0.79-2.5)	1.7% (1.2-2.5%)	6% (3.4-9.7%)	3.8 (2.2-6.2)	550 (350-790)
Portugal	2016	Female	25 to 29	54% (36-70%)	46% (30-64%)	1.5 (0.82-2.4)	1.5% (1-2.1%)	6% (3.6-8.9%)	4 (2.4-6)	520 (340-710)
Portugal	2016	Female	30 to 34	57% (36-73%)	43% (27-64%)	2 (1.1-3)	1.9% (1.3-2.6%)	7.9% (5.2-11%)	9.9 (6.4-14)	880 (600-1200)
Portugal	2016	Female	35 to 39	59% (44-73%)	41% (27-56%)	2.9 (1.7-4.4)	2.9% (1.9-4.1%)	11% (8-15%)	29 (20-40)	2000 (1400-2800)
Portugal	2016	Female	40 to 44	61% (47-74%)	39% (26-53%)	3.5 (2.2-5.2)	4.1% (2.9-5.7%)	14% (10-17%)	52 (38-69)	3000 (2100-4000)
Portugal	2016	Female	45 to 49	62% (50-72%)	38% (28-50%)	3.7 (2.3-5.5)	5% (3.5-7%)	14% (11-19%)	84 (62-110)	4200 (2900-5600)
Portugal	2016	Female	50 to 54	57% (47-66%)	43% (34-53%)	3.6 (2.3-5.2)	5.4% (3.7-7.5%)	13% (9.8-18%)	110 (81-150)	4900 (3400-6700)
Portugal	2016	Female	55 to 59	45% (37-55%)	55% (45-63%)	3 (1.9-4.5)	5.3% (3.5-7.5%)	12% (8.3-16%)	130 (90-170)	4700 (3200-6800)
Portugal	2016	Female	60 to 64	42% (32-52%)	58% (48-68%)	2.9 (1.8-4.4)	5.1% (3.2-7.5%)	10% (6.7-14%)	160 (110-230)	5200 (3300-7700)
Portugal	2016	Female	65 to 69	46% (37-55%)	54% (45-63%)	3 (1.9-4.5)	5.4% (3.2-8.2%)	9.6% (6.2-14%)	220 (140-320)	6000 (3600-8800)
Portugal	2016	Female	70 to 74	43% (33-53%)	57% (47-67%)	2.6 (1.6-4)	5.2% (2.7-8%)	8.2% (4.5-12%)	270 (150-420)	5700 (3000-9100)
Portugal	2016	Female	75 to 79	35% (26-47%)	65% (53-74%)	2 (1.2-3.2)	4.4% (2.1-7.4%)	6.3% (3.1-10%)	420 (200-690)	6700 (3200-11000)
Portugal	2016	Female	80 to 84	34% (24-45%)	66% (55-76%)	2 (1.1-3.1)	4.4% (1.9-7.7%)	5.7% (2.6-9.9%)	650 (290-1100)	7300 (3300-13000)
Portugal	2016	Female	85 to 89	35% (24-46%)	65% (54-76%)	2 (1.1-3.2)	4.4% (1.7-7.9%)	5.4% (2-9.7%)	690 (270-1300)	5400 (2100-9900)
Portugal	2016	Female	90 to 94	37% (26-49%)	63% (51-74%)	2 (1.1-3.4)	4.2% (1.3-8.2%)	5% (1.5-9.8%)	460 (130-900)	2500 (760-4800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Portugal	2016	Female	95 plus	41% (26-57%)	59% (43-74%)	2.2 (1-4)	4.6% (0.95-9.2%)	5.4% (0.96-11%)	240 (43-490)	820 (170-1600)
Portugal	2016	Male	All Ages	75% (70-80%)	25% (20-30%)	7.3 (5.1-10)	15% (12-17%)	18% (14-21%)	10000 (8100-12000)	230000 (190000-260000)
Portugal	2016	Male	15 to 19	61% (45-76%)	39% (24-55%)	2.7 (1.4-4.3)	5.5% (3.8-7.9%)	13% (7.7-20%)	13 (7.5-21)	1500 (990-2100)
Portugal	2016	Male	20 to 24	70% (55-83%)	30% (17-45%)	4.5 (2.7-6.9)	13% (9.1-17%)	22% (14-33%)	39 (24-58)	4100 (2900-5400)
Portugal	2016	Male	25 to 29	76% (61-87%)	24% (13-39%)	6.4 (3.9-9.7)	14% (10-18%)	27% (18-36%)	52 (35-71)	4600 (3300-5900)
Portugal	2016	Male	30 to 34	79% (63-90%)	21% (10-37%)	7.7 (4.8-11)	14% (10-17%)	27% (20-34%)	86 (61-110)	6200 (4700-7800)
Portugal	2016	Male	35 to 39	82% (69-91%)	18% (9-31%)	8.5 (5.2-13)	14% (11-18%)	28% (21-33%)	170 (130-210)	11000 (8100-13000)
Portugal	2016	Male	40 to 44	83% (71-92%)	17% (8.3-29%)	9.6 (6.2-14)	16% (13-20%)	29% (24-33%)	270 (220-320)	14000 (11000-16000)
Portugal	2016	Male	45 to 49	84% (75-91%)	16% (8.9-25%)	11 (7.3-16)	18% (15-22%)	30% (26-34%)	410 (340-470)	18000 (16000-21000)
Portugal	2016	Male	50 to 54	86% (80-91%)	14% (8.7-20%)	12 (7.7-17)	19% (16-22%)	29% (25-32%)	580 (490-670)	23000 (20000-27000)
Portugal	2016	Male	55 to 59	89% (84-92%)	11% (7.8-16%)	11 (7.5-16)	18% (16-21%)	26% (23-30%)	640 (550-740)	22000 (19000-26000)
Portugal	2016	Male	60 to 64	87% (82-91%)	13% (9-18%)	10 (6.8-15)	17% (14-20%)	23% (20-26%)	800 (660-940)	24000 (20000-28000)
Portugal	2016	Male	65 to 69	84% (78-88%)	16% (12-22%)	9.5 (6.4-14)	16% (13-19%)	21% (17-24%)	900 (740-1100)	22000 (18000-27000)
Portugal	2016	Male	70 to 74	81% (75-86%)	19% (14-25%)	8.9 (5.9-13)	15% (12-18%)	19% (16-22%)	1000 (810-1200)	20000 (16000-24000)
Portugal	2016	Male	75 to 79	80% (71-86%)	20% (14-29%)	8.1 (5.2-12)	14% (11-17%)	17% (13-20%)	1400 (1100-1800)	22000 (17000-27000)
Portugal	2016	Male	80 to 84	77% (66-85%)	23% (15-34%)	7.2 (4.5-11)	13% (9.5-17%)	15% (11-20%)	1700 (1200-2200)	19000 (13000-24000)
Portugal	2016	Male	85 to 89	76% (65-85%)	24% (15-35%)	6.4 (3.8-9.7)	12% (7.7-16%)	14% (8.7-19%)	1300 (800-1700)	10000 (6300-13000)
Portugal	2016	Male	90 to 94	77% (66-86%)	23% (14-34%)	6 (3.4-9.1)	11% (6.1-15%)	13% (6.6-18%)	630 (320-880)	3300 (1800-4600)
Portugal	2016	Male	95 plus	79% (65-89%)	21% (11-35%)	5.5 (2.8-9)	11% (4-16%)	12% (4.3-18%)	160 (60-250)	580 (220-860)
Spain	1990	Female	All Ages	55% (51-60%)	45% (40-49%)	3.2 (2.7-3.8)	6.8% (5.1-8.8%)	10% (7.1-14%)	16000 (11000-22000)	330000 (250000-410000)
Spain	1990	Female	15 to 19	56% (43-68%)	44% (32-57%)	2.8 (1.8-3.9)	3.2% (1.9-4.7%)	11% (6.1-17%)	59 (33-91)	5600 (3400-8100)
Spain	1990	Female	20 to 24	67% (54-77%)	33% (23-46%)	3.6 (2.5-5)	4.2% (2.8-5.8%)	13% (8.4-18%)	86 (56-120)	8300 (5800-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	1990	Female	25 to 29	68% (55-77%)	32% (23-45%)	3.6 (2.5-4.9)	3.9% (2.8-5.3%)	12% (8-16%)	95 (66-130)	8300 (5800-11000)
Spain	1990	Female	30 to 34	67% (54-77%)	33% (23-46%)	3.6 (2.3-5)	4.1% (3-5.5%)	13% (9.7-17%)	110 (84-150)	8600 (6200-11000)
Spain	1990	Female	35 to 39	64% (53-73%)	36% (27-47%)	3.8 (2.6-5.3)	4.7% (3.5-6.3%)	15% (11-19%)	150 (110-190)	9600 (7200-12000)
Spain	1990	Female	40 to 44	64% (52-74%)	36% (26-48%)	4.2 (2.9-5.7)	5.9% (4.4-7.9%)	17% (13-21%)	230 (180-300)	13000 (9900-17000)
Spain	1990	Female	45 to 49	65% (53-74%)	35% (26-47%)	4.9 (3.4-6.5)	7.4% (5.5-10%)	18% (14-24%)	330 (260-450)	16000 (12000-22000)
Spain	1990	Female	50 to 54	65% (54-73%)	35% (27-46%)	5.2 (3.8-7)	8.8% (6.5-12%)	18% (14-24%)	490 (380-650)	21000 (16000-27000)
Spain	1990	Female	55 to 59	62% (50-71%)	38% (29-50%)	5.2 (3.7-6.8)	9.4% (6.9-12%)	17% (13-23%)	800 (600-1100)	30000 (22000-39000)
Spain	1990	Female	60 to 64	59% (48-69%)	41% (31-52%)	4.8 (3.4-6.5)	9.6% (7-13%)	16% (12-21%)	1100 (830-1400)	34000 (25000-46000)
Spain	1990	Female	65 to 69	56% (46-65%)	44% (35-54%)	4.3 (3.1-5.8)	9.5% (6.7-13%)	14% (10-19%)	1500 (1100-1900)	38000 (27000-50000)
Spain	1990	Female	70 to 74	51% (41-61%)	49% (39-59%)	3.7 (2.6-5.2)	8.9% (6.1-12%)	12% (8.6-17%)	1800 (1300-2500)	37000 (25000-51000)
Spain	1990	Female	75 to 79	46% (34-57%)	54% (43-66%)	3.2 (2.1-4.6)	7.9% (4.9-12%)	10% (6.2-15%)	2400 (1500-3600)	37000 (22000-55000)
Spain	1990	Female	80 to 84	42% (30-52%)	58% (48-70%)	2.9 (1.8-4.1)	7.3% (4.4-11%)	9% (5.3-14%)	2900 (1700-4500)	32000 (19000-50000)
Spain	1990	Female	85 to 89	39% (28-51%)	61% (49-72%)	2.7 (1.6-4.1)	6.8% (3.6-11%)	8.1% (4.1-13%)	2400 (1200-4000)	19000 (9800-30000)
Spain	1990	Female	90 to 94	39% (27-51%)	61% (49-73%)	2.7 (1.5-4.2)	6.4% (2.9-11%)	7.4% (3.2-13%)	1100 (480-2000)	6000 (2700-10000)
Spain	1990	Female	95 plus	40% (26-54%)	60% (46-74%)	2.8 (1.4-4.7)	6.5% (2.3-12%)	7.6% (2.5-14%)	410 (140-750)	1400 (500-2500)
Spain	1990	Male	All Ages	74% (69-79%)	26% (21-31%)	6.8 (5.8-8)	15% (12-17%)	17% (14-21%)	30000 (24000-36000)	830000 (680000-970000)
Spain	1990	Male	15 to 19	67% (54-79%)	33% (21-46%)	4.8 (3.3-6.7)	11% (6.7-15%)	20% (12-29%)	300 (180-440)	24000 (16000-34000)
Spain	1990	Male	20 to 24	75% (63-84%)	25% (16-37%)	7.5 (5.5-9.9)	18% (13-24%)	28% (19-38%)	690 (460-940)	52000 (36000-69000)
Spain	1990	Male	25 to 29	78% (67-86%)	22% (14-33%)	7.4 (5.6-9.6)	16% (11-20%)	24% (17-31%)	660 (470-860)	46000 (34000-59000)
Spain	1990	Male	30 to 34	80% (68-87%)	20% (13-32%)	7.5 (5.5-9.7)	14% (11-18%)	23% (17-29%)	610 (460-760)	39000 (31000-49000)
Spain	1990	Male	35 to 39	81% (72-88%)	19% (12-28%)	8 (5.8-10)	16% (12-19%)	26% (21-31%)	650 (510-780)	38000 (30000-45000)
Spain	1990	Male	40 to 44	83% (75-90%)	17% (9.9-25%)	9.1 (6.9-12)	17% (14-21%)	28% (23-32%)	890 (750-1000)	46000 (39000-54000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	1990	Male	45 to 49	86% (78-91%)	14% (8.8-22%)	11 (8.1-14)	19% (16-22%)	28% (24-33%)	1200 (1000-1400)	55000 (47000-64000)
Spain	1990	Male	50 to 54	87% (81-92%)	13% (8.3-19%)	11 (8.9-15)	19% (16-23%)	27% (22-31%)	1600 (1400-1900)	65000 (54000-75000)
Spain	1990	Male	55 to 59	88% (82-92%)	12% (8-18%)	12 (9.2-15)	19% (15-22%)	24% (19-29%)	2600 (2100-3100)	89000 (72000-110000)
Spain	1990	Male	60 to 64	87% (81-92%)	13% (8.3-19%)	11 (8.9-14)	17% (13-21%)	22% (17-26%)	3200 (2500-3900)	94000 (74000-110000)
Spain	1990	Male	65 to 69	86% (79-91%)	14% (9.3-21%)	10 (7.9-13)	16% (12-20%)	19% (14-24%)	3800 (2800-4700)	92000 (69000-110000)
Spain	1990	Male	70 to 74	84% (77-89%)	16% (11-23%)	8.8 (6.7-11)	14% (11-18%)	17% (12-21%)	3600 (2600-4400)	69000 (51000-85000)
Spain	1990	Male	75 to 79	81% (71-88%)	19% (12-29%)	7.6 (5.6-10)	13% (9.4-16%)	15% (11-19%)	3800 (2700-4800)	57000 (41000-71000)
Spain	1990	Male	80 to 84	77% (66-85%)	23% (15-34%)	6.6 (4.8-8.6)	12% (8.4-15%)	13% (9.4-17%)	3500 (2500-4600)	38000 (27000-50000)
Spain	1990	Male	85 to 89	73% (61-82%)	27% (18-39%)	5.8 (3.9-8.2)	11% (7.5-14%)	12% (8.4-16%)	2100 (1500-2900)	17000 (12000-22000)
Spain	1990	Male	90 to 94	72% (60-81%)	28% (19-40%)	5.4 (3.5-7.6)	9.7% (6.6-13%)	11% (7.3-15%)	760 (510-1000)	4000 (2800-5500)
Spain	1990	Male	95 plus	72% (57-83%)	28% (17-43%)	5 (2.9-7.6)	9.2% (5-13%)	10% (5.4-15%)	190 (97-270)	660 (360-940)
Spain	1995	Female	All Ages	56% (51-60%)	44% (40-49%)	3 (2.5-3.6)	5.8% (4.4-7.6%)	8.7% (5.9-12%)	14000 (9600-19000)	280000 (210000-360000)
Spain	1995	Female	15 to 19	55% (43-67%)	45% (33-57%)	2.5 (1.6-3.6)	2.5% (1.5-3.6%)	9.8% (5.3-15%)	38 (20-58)	4000 (2500-5700)
Spain	1995	Female	20 to 24	67% (54-77%)	33% (23-46%)	3.3 (2.3-4.5)	3.4% (2.3-4.6%)	11% (7-16%)	59 (37-84)	6600 (4500-9000)
Spain	1995	Female	25 to 29	68% (55-78%)	32% (22-45%)	3.3 (2.2-4.5)	3.1% (2.2-4.1%)	8.5% (5.7-12%)	69 (47-95)	6800 (4700-9100)
Spain	1995	Female	30 to 34	67% (53-76%)	33% (24-47%)	3.3 (2.2-4.6)	3.1% (2.2-4.1%)	8.3% (5.9-11%)	90 (64-120)	7400 (5300-9800)
Spain	1995	Female	35 to 39	64% (52-74%)	36% (26-48%)	3.4 (2.4-4.8)	4% (2.9-5.2%)	12% (8.7-15%)	140 (100-180)	9300 (6800-12000)
Spain	1995	Female	40 to 44	64% (53-74%)	36% (26-47%)	3.8 (2.7-5.3)	5.2% (3.9-6.9%)	14% (11-18%)	200 (160-260)	12000 (9000-15000)
Spain	1995	Female	45 to 49	65% (53-74%)	35% (26-47%)	4.5 (3.2-6)	6.5% (4.8-8.7%)	16% (12-21%)	310 (240-400)	16000 (12000-20000)
Spain	1995	Female	50 to 54	65% (54-73%)	35% (27-46%)	4.8 (3.4-6.4)	7.6% (5.5-10%)	16% (12-22%)	420 (320-560)	18000 (14000-25000)
Spain	1995	Female	55 to 59	62% (51-71%)	38% (29-49%)	4.7 (3.3-6.3)	8.2% (6-11%)	16% (12-21%)	560 (420-750)	21000 (16000-29000)
Spain	1995	Female	60 to 64	59% (47-69%)	41% (31-53%)	4.4 (3.1-6)	8.3% (6-11%)	14% (11-19%)	900 (670-1200)	29000 (21000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	1995	Female	65 to 69	56% (45-65%)	44% (35-55%)	4 (2.8-5.2)	8.2% (5.7-11%)	13% (9.2-17%)	1200 (900-1600)	33000 (23000-44000)
Spain	1995	Female	70 to 74	51% (41-61%)	49% (39-59%)	3.4 (2.4-4.7)	7.6% (5-11%)	11% (7.4-15%)	1700 (1100-2300)	34000 (23000-49000)
Spain	1995	Female	75 to 79	45% (34-57%)	55% (43-66%)	2.9 (1.8-4.2)	6.7% (4.1-10%)	8.8% (5.3-13%)	1900 (1200-2900)	30000 (18000-45000)
Spain	1995	Female	80 to 84	41% (30-52%)	59% (48-70%)	2.6 (1.6-3.7)	6.1% (3.6-9.5%)	7.6% (4.5-12%)	2500 (1500-3800)	28000 (17000-43000)
Spain	1995	Female	85 to 89	39% (28-50%)	61% (50-72%)	2.5 (1.5-3.7)	5.7% (3-9.2%)	6.8% (3.5-11%)	2300 (1200-3800)	18000 (9700-30000)
Spain	1995	Female	90 to 94	39% (28-50%)	61% (50-72%)	2.5 (1.4-3.8)	5.4% (2.4-9.2%)	6.3% (2.6-11%)	1300 (560-2300)	7200 (3100-12000)
Spain	1995	Female	95 plus	40% (26-53%)	60% (47-74%)	2.6 (1.3-4.3)	5.7% (1.8-10%)	6.6% (1.9-12%)	480 (130-890)	1700 (510-3000)
Spain	1995	Male	All Ages	75% (70-80%)	25% (20-30%)	6.4 (5.4-7.5)	13% (11-15%)	15% (13-18%)	29000 (24000-34000)	750000 (640000-890000)
Spain	1995	Male	15 to 19	67% (53-79%)	33% (21-47%)	4.3 (3-6)	8.2% (5.4-12%)	18% (11-26%)	180 (110-260)	16000 (11000-22000)
Spain	1995	Male	20 to 24	75% (63-84%)	25% (16-37%)	6.8 (4.9-9)	15% (10-20%)	25% (17-34%)	450 (300-610)	36000 (26000-48000)
Spain	1995	Male	25 to 29	77% (66-85%)	23% (15-34%)	6.7 (5-8.6)	12% (8.8-15%)	18% (13-23%)	470 (340-620)	35000 (26000-45000)
Spain	1995	Male	30 to 34	80% (68-87%)	20% (13-32%)	6.8 (5-8.9)	9.7% (7.5-12%)	14% (10-17%)	490 (360-620)	33000 (26000-41000)
Spain	1995	Male	35 to 39	81% (72-88%)	19% (12-28%)	7.2 (5.3-9.4)	12% (9.1-15%)	18% (14-22%)	590 (460-730)	36000 (28000-43000)
Spain	1995	Male	40 to 44	83% (74-90%)	17% (9.9-26%)	8.1 (6.1-10)	15% (12-18%)	24% (20-27%)	790 (660-920)	41000 (34000-49000)
Spain	1995	Male	45 to 49	86% (78-91%)	14% (8.7-22%)	9.6 (7.3-12)	17% (14-20%)	25% (22-29%)	1200 (1000-1400)	56000 (47000-65000)
Spain	1995	Male	50 to 54	87% (81-92%)	13% (7.9-19%)	10 (8.1-13)	18% (14-21%)	25% (21-29%)	1500 (1300-1800)	62000 (52000-72000)
Spain	1995	Male	55 to 59	88% (82-92%)	12% (8.2-18%)	11 (8.4-13)	17% (14-21%)	23% (19-27%)	2000 (1600-2400)	69000 (56000-82000)
Spain	1995	Male	60 to 64	87% (81-92%)	13% (8.3-19%)	10 (8-13)	16% (13-20%)	20% (16-24%)	3000 (2400-3600)	89000 (71000-110000)
Spain	1995	Male	65 to 69	86% (80-91%)	14% (9.2-20%)	9.2 (7.1-12)	15% (12-18%)	18% (14-22%)	3700 (2800-4500)	89000 (71000-110000)
Spain	1995	Male	70 to 74	84% (77-89%)	16% (11-23%)	8 (6.1-10)	13% (10-16%)	16% (12-19%)	4000 (3100-5000)	79000 (61000-98000)
Spain	1995	Male	75 to 79	81% (72-88%)	19% (12-28%)	6.9 (5.1-9.2)	12% (9.1-15%)	13% (10-17%)	3400 (2700-4400)	52000 (40000-65000)
Spain	1995	Male	80 to 84	77% (66-85%)	23% (15-34%)	6 (4.2-7.9)	11% (8.1-14%)	12% (9.1-16%)	3300 (2500-4200)	37000 (27000-47000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	1995	Male	85 to 89	73% (62-82%)	27% (18-38%)	5.2 (3.5-7.2)	9.5% (6.9-13%)	11% (7.6-14%)	2300 (1600-3000)	18000 (13000-24000)
Spain	1995	Male	90 to 94	72% (60-81%)	28% (19-40%)	4.8 (3.1-6.7)	8.4% (5.5-11%)	9.5% (6.1-13%)	910 (580-1300)	4900 (3200-6800)
Spain	1995	Male	95 plus	71% (56-83%)	29% (17-44%)	4.5 (2.6-6.8)	8% (3.9-12%)	8.9% (4-13%)	210 (96-320)	760 (370-1100)
Spain	2000	Female	All Ages	51% (48-55%)	49% (45-52%)	2.5 (2.1-3)	4.4% (3.4-5.7%)	6.2% (4.3-8.6%)	11000 (7400-15000)	220000 (170000-270000)
Spain	2000	Female	15 to 19	51% (41-62%)	49% (38-59%)	2 (1.4-2.9)	2.1% (1.3-3.1%)	8.4% (4.6-13%)	26 (14-41)	2900 (1800-4200)
Spain	2000	Female	20 to 24	65% (55-74%)	35% (26-45%)	2.9 (2.1-4)	3% (2.1-4.1%)	11% (6.8-15%)	46 (29-65)	5600 (3900-7600)
Spain	2000	Female	25 to 29	68% (59-75%)	32% (25-41%)	3 (2.1-3.9)	2.9% (2-3.9%)	10% (7-14%)	56 (37-77)	6000 (4200-8000)
Spain	2000	Female	30 to 34	65% (56-72%)	35% (28-44%)	2.9 (2-3.9)	3% (2.2-3.9%)	10% (7.4-13%)	80 (58-100)	6800 (4900-9000)
Spain	2000	Female	35 to 39	62% (54-69%)	38% (31-46%)	3 (2.1-4)	3.5% (2.6-4.6%)	11% (8.7-14%)	120 (97-160)	8600 (6600-11000)
Spain	2000	Female	40 to 44	63% (55-71%)	37% (29-45%)	3.4 (2.5-4.5)	4.6% (3.5-5.9%)	13% (10-16%)	200 (160-250)	12000 (9000-15000)
Spain	2000	Female	45 to 49	64% (58-71%)	36% (29-42%)	4 (3-5.2)	5.8% (4.4-7.6%)	15% (12-18%)	280 (220-350)	14000 (11000-18000)
Spain	2000	Female	50 to 54	60% (53-67%)	40% (33-47%)	4 (3-5.1)	6.3% (4.7-8.4%)	14% (11-18%)	370 (290-480)	16000 (13000-22000)
Spain	2000	Female	55 to 59	54% (47-60%)	46% (40-53%)	3.5 (2.6-4.7)	6.4% (4.8-8.5%)	13% (10-17%)	440 (340-570)	17000 (13000-22000)
Spain	2000	Female	60 to 64	47% (41-54%)	53% (46-59%)	3 (2.2-4.2)	6.1% (4.5-8.3%)	11% (8.4-14%)	540 (410-710)	17000 (13000-23000)
Spain	2000	Female	65 to 69	43% (36-50%)	57% (50-64%)	2.6 (1.9-3.4)	5.8% (4.2-7.7%)	9.4% (7-12%)	860 (640-1100)	23000 (17000-30000)
Spain	2000	Female	70 to 74	37% (31-44%)	63% (56-69%)	2.1 (1.5-2.9)	5.2% (3.6-7.1%)	7.6% (5.4-11%)	1100 (810-1600)	24000 (16000-33000)
Spain	2000	Female	75 to 79	30% (23-38%)	70% (62-77%)	1.6 (1.1-2.3)	4.3% (2.8-6.2%)	5.7% (3.7-8.4%)	1400 (880-2000)	22000 (14000-32000)
Spain	2000	Female	80 to 84	31% (22-39%)	69% (61-78%)	1.7 (1.1-2.4)	4.1% (2.5-6.3%)	5.2% (3.1-8.1%)	1700 (1000-2600)	19000 (11000-29000)
Spain	2000	Female	85 to 89	32% (22-42%)	68% (58-78%)	1.8 (1.1-2.6)	4.1% (2.2-6.8%)	4.9% (2.6-8.1%)	1800 (980-3000)	15000 (7900-24000)
Spain	2000	Female	90 to 94	32% (23-41%)	68% (59-77%)	1.8 (1-2.7)	3.8% (1.6-6.6%)	4.5% (1.8-8%)	1200 (460-2100)	6300 (2600-11000)
Spain	2000	Female	95 plus	33% (21-45%)	67% (55-79%)	1.8 (0.93-3.1)	4.1% (1.3-7.8%)	4.7% (1.4-9.1%)	500 (150-960)	1800 (570-3300)
Spain	2000	Male	All Ages	78% (75-82%)	22% (18-25%)	6.2 (5.3-7.3)	13% (11-15%)	15% (12-17%)	27000 (23000-33000)	720000 (620000-840000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	2000	Male	15 to 19	68% (56-79%)	32% (21-44%)	4 (2.8-5.4)	7.7% (5-11%)	17% (10-25%)	140 (80-200)	12000 (7700-16000)
Spain	2000	Male	20 to 24	77% (67-84%)	23% (16-33%)	6.4 (4.7-8.2)	15% (10-19%)	26% (17-36%)	380 (250-510)	32000 (23000-41000)
Spain	2000	Male	25 to 29	81% (74-86%)	19% (14-26%)	6.4 (4.9-8.1)	13% (9.8-17%)	24% (17-31%)	410 (280-530)	32000 (24000-41000)
Spain	2000	Male	30 to 34	83% (76-88%)	17% (12-24%)	6.6 (5-8.2)	12% (8.6-15%)	20% (15-26%)	460 (340-590)	32000 (24000-39000)
Spain	2000	Male	35 to 39	84% (78-89%)	16% (11-22%)	6.8 (5.2-8.7)	12% (9.4-15%)	21% (16-26%)	600 (460-740)	36000 (28000-44000)
Spain	2000	Male	40 to 44	86% (79-91%)	14% (9.1-21%)	7.8 (5.9-9.8)	14% (12-17%)	24% (20-28%)	840 (700-1000)	44000 (37000-53000)
Spain	2000	Male	45 to 49	88% (84-92%)	12% (7.8-16%)	9.2 (7.4-11)	17% (14-20%)	26% (22-29%)	1200 (1000-1300)	54000 (46000-62000)
Spain	2000	Male	50 to 54	90% (86-93%)	10% (7-14%)	10 (8-12)	17% (14-20%)	24% (21-28%)	1600 (1400-1900)	64000 (55000-75000)
Spain	2000	Male	55 to 59	90% (86-93%)	10% (7.3-14%)	10 (8.1-12)	17% (14-20%)	22% (19-26%)	1900 (1600-2300)	68000 (57000-80000)
Spain	2000	Male	60 to 64	89% (85-92%)	11% (8-15%)	9.7 (7.7-12)	16% (13-19%)	20% (16-24%)	2300 (1900-2800)	70000 (58000-83000)
Spain	2000	Male	65 to 69	87% (83-91%)	13% (9.4-17%)	8.5 (6.8-10)	14% (11-17%)	17% (14-21%)	3400 (2800-4100)	85000 (69000-100000)
Spain	2000	Male	70 to 74	84% (78-88%)	16% (12-22%)	7.1 (5.6-9)	12% (10-15%)	15% (12-18%)	3900 (3100-4700)	77000 (62000-93000)
Spain	2000	Male	75 to 79	79% (72-86%)	21% (14-28%)	6 (4.6-7.9)	11% (8.6-13%)	12% (9.8-15%)	3900 (3100-4900)	60000 (48000-75000)
Spain	2000	Male	80 to 84	75% (64-83%)	25% (17-36%)	5.1 (3.7-6.8)	9.5% (7.1-12%)	11% (8-14%)	2900 (2200-3800)	33000 (24000-42000)
Spain	2000	Male	85 to 89	71% (60-80%)	29% (20-40%)	4.5 (3.1-6.1)	8.4% (6-11%)	9.4% (6.5-13%)	2200 (1500-2900)	17000 (12000-23000)
Spain	2000	Male	90 to 94	70% (59-80%)	30% (20-41%)	4.1 (2.7-5.8)	7.3% (4.5-10%)	8.2% (5-11%)	990 (600-1400)	5300 (3300-7500)
Spain	2000	Male	95 plus	70% (55-82%)	30% (18-45%)	3.9 (2.2-5.9)	6.9% (2.9-11%)	7.6% (2.9-12%)	260 (100-410)	950 (400-1500)
Spain	2005	Female	All Ages	52% (49-55%)	48% (45-51%)	2.5 (2.1-2.9)	4% (3-5%)	5.3% (3.7-7.3%)	9800 (6800-13000)	200000 (160000-260000)
Spain	2005	Female	15 to 19	53% (44-63%)	47% (37-56%)	2 (1.4-2.9)	1.9% (1.2-2.7%)	8.1% (4.5-12%)	18 (10-28)	2200 (1400-3200)
Spain	2005	Female	20 to 24	68% (60-76%)	32% (24-40%)	3 (2.1-4)	2.9% (2-3.9%)	11% (6.6-15%)	34 (21-48)	4600 (3100-6300)
Spain	2005	Female	25 to 29	69% (61-76%)	31% (24-39%)	2.9 (2.1-3.9)	2.6% (1.8-3.5%)	10% (6.7-14%)	46 (30-62)	5600 (3900-7700)
Spain	2005	Female	30 to 34	66% (58-73%)	34% (27-42%)	2.8 (2-3.8)	2.6% (1.9-3.5%)	10% (7.2-13%)	68 (49-89)	6400 (4600-8400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	2005	Female	35 to 39	63% (56-70%)	37% (30-44%)	3 (2.2-3.9)	3.2% (2.4-4.2%)	11% (8.6-14%)	120 (92-150)	8600 (6400-11000)
Spain	2005	Female	40 to 44	65% (58-71%)	35% (29-42%)	3.4 (2.5-4.4)	4.2% (3.2-5.5%)	12% (10-15%)	200 (160-240)	12000 (9200-15000)
Spain	2005	Female	45 to 49	65% (59-70%)	35% (30-41%)	3.9 (3-5.1)	5.5% (4.1-7.3%)	14% (11-17%)	310 (250-390)	16000 (12000-20000)
Spain	2005	Female	50 to 54	60% (54-66%)	40% (34-46%)	3.8 (2.8-5)	6% (4.5-7.9%)	13% (10-17%)	360 (280-470)	16000 (13000-22000)
Spain	2005	Female	55 to 59	56% (50-61%)	44% (39-50%)	3.6 (2.7-4.6)	6% (4.5-8.1%)	12% (9.9-16%)	450 (360-580)	18000 (14000-23000)
Spain	2005	Female	60 to 64	49% (44-55%)	51% (45-56%)	3 (2.3-4)	5.6% (4.1-7.6%)	10% (8-14%)	500 (390-660)	17000 (12000-22000)
Spain	2005	Female	65 to 69	43% (38-49%)	57% (51-62%)	2.5 (1.9-3.2)	5.1% (3.8-6.8%)	8.7% (6.6-11%)	620 (470-790)	17000 (12000-22000)
Spain	2005	Female	70 to 74	37% (31-42%)	63% (58-69%)	2 (1.4-2.7)	4.6% (3.1-6.2%)	7% (4.9-9.3%)	980 (700-1300)	21000 (15000-29000)
Spain	2005	Female	75 to 79	29% (24-36%)	71% (64-76%)	1.5 (1.1-2.2)	3.8% (2.4-5.4%)	5.2% (3.4-7.4%)	1200 (800-1800)	20000 (13000-28000)
Spain	2005	Female	80 to 84	28% (20-36%)	72% (64-80%)	1.5 (0.95-2.1)	3.5% (2.2-5.3%)	4.5% (2.8-6.8%)	1600 (1000-2500)	19000 (12000-28000)
Spain	2005	Female	85 to 89	28% (20-36%)	72% (64-80%)	1.5 (0.93-2.2)	3.4% (1.9-5.5%)	4.1% (2.2-6.7%)	1600 (860-2600)	13000 (7000-20000)
Spain	2005	Female	90 to 94	28% (20-36%)	72% (64-80%)	1.5 (0.87-2.3)	3.1% (1.4-5.5%)	3.6% (1.5-6.5%)	1100 (470-2000)	6000 (2700-11000)
Spain	2005	Female	95 plus	29% (19-41%)	71% (59-81%)	1.5 (0.79-2.5)	3.3% (1-6.3%)	3.8% (1.1-7.5%)	530 (150-1000)	1900 (580-3600)
Spain	2005	Male	All Ages	79% (76-83%)	21% (17-24%)	6.1 (5.2-7.2)	12% (10-14%)	13% (11-16%)	27000 (22000-31000)	700000 (600000-810000)
Spain	2005	Male	15 to 19	70% (58-80%)	30% (20-42%)	3.9 (2.8-5.3)	7.2% (4.7-9.9%)	16% (9.8-24%)	100 (61-150)	9400 (6200-13000)
Spain	2005	Male	20 to 24	78% (69-85%)	22% (15-31%)	6.3 (4.7-8.2)	14% (9.8-18%)	26% (18-35%)	270 (180-360)	24000 (17000-31000)
Spain	2005	Male	25 to 29	82% (75-87%)	18% (13-25%)	6.3 (4.8-7.9)	12% (8.9-15%)	24% (17-32%)	340 (230-440)	29000 (21000-37000)
Spain	2005	Male	30 to 34	84% (77-89%)	16% (11-23%)	6.4 (4.9-8)	11% (8.1-14%)	22% (16-28%)	390 (290-510)	29000 (22000-37000)
Spain	2005	Male	35 to 39	84% (79-90%)	16% (10-21%)	6.7 (5-8.5)	11% (8.7-14%)	21% (16-26%)	540 (410-670)	34000 (26000-43000)
Spain	2005	Male	40 to 44	86% (80-91%)	14% (8.6-20%)	7.5 (5.9-9.5)	13% (10-16%)	23% (19-26%)	820 (680-960)	44000 (36000-53000)
Spain	2005	Male	45 to 49	89% (85-92%)	11% (7.6-15%)	8.9 (7-11)	16% (13-18%)	25% (21-28%)	1200 (1100-1400)	57000 (49000-66000)
Spain	2005	Male	50 to 54	90% (86-93%)	10% (6.7-14%)	9.6 (7.6-12)	16% (14-19%)	24% (20-27%)	1600 (1400-1800)	64000 (55000-74000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	2005	Male	55 to 59	90% (87-93%)	10% (7.4-13%)	9.7 (7.8-12)	16% (13-19%)	22% (19-25%)	2000 (1700-2300)	72000 (61000-84000)
Spain	2005	Male	60 to 64	89% (86-92%)	11% (8.2-14%)	9.3 (7.3-12)	15% (12-18%)	19% (16-23%)	2300 (1900-2700)	69000 (58000-82000)
Spain	2005	Male	65 to 69	87% (83-90%)	13% (9.6-17%)	8.2 (6.5-10)	14% (11-16%)	17% (14-20%)	2700 (2200-3200)	67000 (55000-80000)
Spain	2005	Male	70 to 74	84% (80-88%)	16% (12-20%)	6.9 (5.4-8.6)	12% (9.8-14%)	14% (12-17%)	3700 (3000-4400)	74000 (61000-89000)
Spain	2005	Male	75 to 79	79% (72-86%)	21% (14-28%)	5.7 (4.4-7.5)	10% (8.1-13%)	12% (9.4-15%)	3900 (3100-4800)	60000 (48000-74000)
Spain	2005	Male	80 to 84	73% (62-81%)	27% (19-38%)	4.6 (3.3-6.3)	8.7% (6.4-11%)	9.9% (7.3-13%)	3500 (2600-4500)	40000 (30000-51000)
Spain	2005	Male	85 to 89	69% (58-78%)	31% (22-42%)	4 (2.8-5.5)	7.4% (5.1-9.9%)	8.4% (5.7-11%)	2000 (1400-2700)	16000 (11000-21000)
Spain	2005	Male	90 to 94	68% (56-78%)	32% (22-44%)	3.7 (2.4-5.3)	6.4% (3.9-9.3%)	7.2% (4.2-10%)	990 (580-1400)	5400 (3200-7700)
Spain	2005	Male	95 plus	68% (53-81%)	32% (19-47%)	3.5 (2.5-5.4)	6% (2.5-9.4%)	6.6% (2.5-11%)	300 (110-480)	1100 (460-1700)
Spain	2010	Female	All Ages	54% (51-57%)	46% (43-49%)	2.6 (2-3.2)	3.8% (2.8-4.9%)	5.1% (3.5-7.2%)	9600 (6500-13000)	200000 (150000-270000)
Spain	2010	Female	15 to 19	57% (47-68%)	43% (32-53%)	2.2 (1.5-3.2)	1.7% (1.1-2.4%)	7.8% (4.4-12%)	13 (7.1-19)	1800 (1200-2700)
Spain	2010	Female	20 to 24	71% (62-78%)	29% (22-38%)	3.1 (2.1-4.3)	2.6% (1.8-3.5%)	10% (6.7-15%)	23 (15-32)	3600 (2400-4900)
Spain	2010	Female	25 to 29	70% (62-77%)	30% (23-38%)	3 (2-4.2)	2.3% (1.7-3.2%)	9.9% (6.5-14%)	32 (21-45)	4400 (3000-6100)
Spain	2010	Female	30 to 34	67% (59-74%)	33% (26-41%)	2.9 (2-4.1)	2.3% (1.6-3.1%)	9.9% (7-13%)	55 (38-74)	5700 (4000-7800)
Spain	2010	Female	35 to 39	66% (59-73%)	34% (27-41%)	3.1 (2.2-4.1)	2.8% (2.1-3.8%)	11% (8.5-14%)	100 (78-130)	8000 (5700-10000)
Spain	2010	Female	40 to 44	66% (60-73%)	34% (27-40%)	3.5 (2.5-4.6)	3.9% (2.9-5.1%)	13% (9.9-16%)	190 (150-240)	12000 (9000-15000)
Spain	2010	Female	45 to 49	66% (59-71%)	34% (29-41%)	3.9 (2.8-5.3)	5% (3.6-6.7%)	13% (10-17%)	310 (240-410)	16000 (12000-22000)
Spain	2010	Female	50 to 54	63% (57-68%)	37% (32-43%)	4 (2.8-5.4)	5.7% (4.1-7.8%)	13% (10-17%)	410 (320-540)	19000 (14000-25000)
Spain	2010	Female	55 to 59	60% (55-65%)	40% (35-45%)	3.9 (2.8-5.3)	6% (4.3-8.1%)	12% (9.7-16%)	470 (370-630)	19000 (14000-25000)
Spain	2010	Female	60 to 64	54% (48-60%)	46% (40-52%)	3.4 (2.4-4.6)	5.5% (3.8-7.6%)	10% (7.8-14%)	530 (400-710)	18000 (13000-25000)
Spain	2010	Female	65 to 69	47% (42-54%)	53% (46-58%)	2.8 (2-3.7)	4.9% (3.4-6.7%)	8.6% (6.4-11%)	590 (430-760)	17000 (12000-23000)
Spain	2010	Female	70 to 74	39% (33-46%)	61% (54-67%)	2.1 (1.5-2.9)	4.3% (2.9-6%)	6.8% (4.7-9.2%)	730 (510-990)	16000 (11000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	2010	Female	75 to 79	32% (26-39%)	68% (61-74%)	1.7 (1.1-2.4)	3.7% (2.3-5.4%)	5.2% (3.4-7.6%)	1100 (740-1600)	19000 (12000-28000)
Spain	2010	Female	80 to 84	30% (22-38%)	70% (62-78%)	1.5 (0.97-2.2)	3.4% (2-5.2%)	4.4% (2.7-6.8%)	1600 (970-2400)	18000 (11000-29000)
Spain	2010	Female	85 to 89	29% (21-38%)	71% (62-79%)	1.5 (0.9-2.2)	3.1% (1.7-5.1%)	3.8% (2-6.3%)	1700 (910-2800)	14000 (7500-23000)
Spain	2010	Female	90 to 94	28% (20-37%)	72% (63-80%)	1.5 (0.85-2.3)	2.9% (1.3-5.1%)	3.4% (1.4-6.2%)	1100 (460-2000)	6100 (2700-11000)
Spain	2010	Female	95 plus	30% (20-42%)	70% (58-80%)	1.6 (0.81-2.6)	3.2% (1-6%)	3.7% (1.1-7.1%)	650 (190-1300)	2300 (750-4300)
Spain	2010	Male	All Ages	79% (76-83%)	21% (17-24%)	6 (4.8-7.4)	11% (9.3-13%)	13% (10-15%)	25000 (21000-30000)	640000 (540000-740000)
Spain	2010	Male	15 to 19	71% (60-81%)	29% (19-40%)	3.9 (2.7-5.4)	5.6% (3.7-7.6%)	15% (9.4-22%)	55 (34-79)	6100 (4100-8200)
Spain	2010	Male	20 to 24	79% (71-86%)	21% (14-29%)	6.2 (4.4-8.3)	12% (8.7-15%)	25% (17-34%)	150 (100-200)	16000 (12000-20000)
Spain	2010	Male	25 to 29	82% (76-88%)	18% (12-24%)	6.2 (4.5-8.2)	11% (7.9-13%)	23% (16-31%)	200 (140-260)	19000 (14000-25000)
Spain	2010	Male	30 to 34	83% (76-89%)	17% (11-24%)	6.2 (4.5-8.1)	9.2% (6.7-12%)	21% (15-28%)	270 (190-360)	23000 (17000-29000)
Spain	2010	Male	35 to 39	84% (79-89%)	16% (11-21%)	6.4 (4.7-8.4)	9.7% (7.2-12%)	21% (15-26%)	400 (290-500)	28000 (20000-35000)
Spain	2010	Male	40 to 44	86% (80-91%)	14% (8.9-20%)	7.3 (5.4-9.4)	12% (9-14%)	22% (18-27%)	690 (540-820)	39000 (31000-47000)
Spain	2010	Male	45 to 49	89% (84-92%)	11% (7.9-16%)	8.6 (6.5-11)	14% (12-17%)	24% (20-27%)	1200 (980-1300)	55000 (47000-64000)
Spain	2010	Male	50 to 54	90% (86-93%)	9.9% (6.9-14%)	9.4 (7-12)	15% (13-18%)	23% (20-27%)	1600 (1400-1900)	67000 (57000-76000)
Spain	2010	Male	55 to 59	90% (86-92%)	10% (7.6-14%)	9.5 (7.3-12)	15% (13-18%)	21% (18-24%)	1900 (1600-2200)	68000 (58000-79000)
Spain	2010	Male	60 to 64	89% (85-92%)	11% (8.2-15%)	9 (6.8-12)	14% (12-17%)	18% (16-21%)	2300 (1900-2600)	70000 (58000-82000)
Spain	2010	Male	65 to 69	87% (84-91%)	13% (9.1-16%)	8 (6.1-10)	13% (11-15%)	16% (14-19%)	2500 (2100-3000)	65000 (54000-76000)
Spain	2010	Male	70 to 74	85% (81-89%)	15% (11-19%)	6.9 (5.1-8.9)	11% (9.2-14%)	14% (11-17%)	2800 (2300-3300)	57000 (47000-69000)
Spain	2010	Male	75 to 79	81% (74-87%)	19% (13-26%)	5.8 (4.3-7.8)	9.8% (7.5-12%)	12% (8.9-14%)	3600 (2800-4500)	57000 (44000-71000)
Spain	2010	Male	80 to 84	75% (64-83%)	25% (17-36%)	4.7 (3.3-6.5)	8.3% (5.9-11%)	9.6% (6.7-12%)	3500 (2500-4500)	40000 (28000-52000)
Spain	2010	Male	85 to 89	70% (59-79%)	30% (21-41%)	4 (2.7-5.8)	7% (4.7-9.4%)	7.9% (5.2-11%)	2500 (1700-3400)	20000 (14000-28000)
Spain	2010	Male	90 to 94	69% (57-78%)	31% (22-43%)	3.7 (2.3-5.4)	6% (3.4-8.5%)	6.7% (3.7-9.6%)	990 (540-1400)	5500 (3100-7900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	2010	Male	95 plus	69% (53-81%)	31% (19-47%)	3.5 (2-5.5)	5.7% (2.4-9.1%)	6.2% (2.4-10%)	340 (130-560)	1300 (530-2000)
Spain	2016	Female	All Ages	55% (51-59%)	45% (41-49%)	2.7 (1.9-3.7)	3.9% (2.6-5.6%)	5.5% (3.2-8.4%)	12000 (6700-18000)	210000 (140000-300000)
Spain	2016	Female	15 to 19	55% (42-67%)	45% (33-58%)	2.1 (1.2-3.4)	1.5% (0.98-2.2%)	7.3% (3.9-12%)	9.6 (5.1-15)	1500 (950-2200)
Spain	2016	Female	20 to 24	66% (53-77%)	34% (23-47%)	2.8 (1.7-4.4)	2.4% (1.6-3.2%)	9.9% (5.7-15%)	16 (9.3-25)	2800 (1900-4000)
Spain	2016	Female	25 to 29	67% (55-77%)	33% (23-45%)	2.8 (1.7-4.3)	2.1% (1.4-3%)	9.6% (5.9-14%)	21 (12-31)	3000 (2000-4300)
Spain	2016	Female	30 to 34	66% (54-76%)	34% (24-46%)	2.8 (1.7-4.3)	2.1% (1.4-3%)	10% (6.5-14%)	38 (24-55)	4100 (2800-5700)
Spain	2016	Female	35 to 39	64% (52-74%)	36% (26-48%)	2.9 (1.8-4.4)	2.6% (1.8-3.6%)	11% (7.8-15%)	90 (63-120)	7200 (4900-9700)
Spain	2016	Female	40 to 44	64% (53-73%)	36% (27-47%)	3.3 (2.2-4.9)	3.4% (2.4-4.9%)	12% (9.1-16%)	170 (130-220)	11000 (7600-14000)
Spain	2016	Female	45 to 49	67% (56-74%)	33% (26-44%)	4 (2.5-5.8)	4.7% (3.2-6.6%)	13% (10-18%)	310 (230-420)	16000 (11000-22000)
Spain	2016	Female	50 to 54	67% (59-74%)	33% (26-41%)	4.3 (2.8-6.3)	5.6% (3.9-7.8%)	13% (9.9-18%)	450 (320-610)	20000 (14000-28000)
Spain	2016	Female	55 to 59	64% (57-70%)	36% (30-43%)	4.2 (2.7-6.2)	5.9% (4-8.3%)	12% (9.2-17%)	540 (390-740)	21000 (15000-29000)
Spain	2016	Female	60 to 64	59% (51-67%)	41% (33-49%)	3.8 (2.5-5.5)	5.6% (3.6-7.8%)	11% (7.6-15%)	540 (380-740)	18000 (12000-26000)
Spain	2016	Female	65 to 69	54% (46-62%)	46% (38-54%)	3.2 (2-4.7)	5.1% (3.2-7.4%)	9.2% (6.2-13%)	690 (470-970)	20000 (12000-28000)
Spain	2016	Female	70 to 74	46% (37-56%)	54% (44-63%)	2.6 (1.5-4)	4.6% (2.6-6.9%)	7.5% (4.6-11%)	810 (510-1200)	18000 (11000-27000)
Spain	2016	Female	75 to 79	39% (29-49%)	61% (51-71%)	2.1 (1.3-3.3)	4% (2.2-6.4%)	5.7% (3.3-8.8%)	970 (560-1500)	16000 (9100-26000)
Spain	2016	Female	80 to 84	36% (26-46%)	64% (54-74%)	1.9 (1.1-3)	3.7% (2-6.3%)	4.9% (2.6-8.2%)	1800 (980-3100)	22000 (12000-37000)
Spain	2016	Female	85 to 89	35% (24-45%)	65% (55-76%)	1.8 (1-3.1)	3.6% (1.6-6.3%)	4.4% (2-7.9%)	2200 (1000-3900)	18000 (8200-32000)
Spain	2016	Female	90 to 94	35% (25-45%)	65% (55-75%)	1.8 (0.98-3.2)	3.4% (1.3-6.3%)	4% (1.5-7.7%)	1800 (670-3500)	10000 (3900-19000)
Spain	2016	Female	95 plus	36% (23-49%)	64% (51-77%)	1.9 (0.92-3.4)	3.7% (1-7.2%)	4.3% (1-8.4%)	980 (230-1900)	3400 (910-6600)
Spain	2016	Male	All Ages	78% (74-82%)	22% (18-26%)	5.8 (4.2-8)	10% (8.3-13%)	12% (9-15%)	25000 (19000-31000)	600000 (490000-710000)
Spain	2016	Male	15 to 19	67% (52-79%)	33% (21-48%)	3.6 (2.2-5.5)	4.8% (3.2-6.6%)	14% (8.2-22%)	37 (21-57)	4600 (3100-6400)
Spain	2016	Male	20 to 24	74% (63-84%)	26% (16-37%)	5.6 (3.6-8.3)	11% (7.5-14%)	23% (15-32%)	100 (62-140)	12000 (8300-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Spain	2016	Male	25 to 29	77% (67-85%)	23% (15-33%)	5.6 (3.6-8.1)	9.5% (6.8-12%)	22% (14-31%)	110 (71-150)	12000 (8600-16000)
Spain	2016	Male	30 to 34	79% (67-87%)	21% (13-33%)	5.7 (3.7-8.2)	8.3% (5.9-11%)	20% (13-27%)	160 (110-220)	14000 (10000-19000)
Spain	2016	Male	35 to 39	81% (72-88%)	19% (12-28%)	6 (4-8.7)	8.5% (6.1-11%)	20% (14-26%)	290 (200-380)	22000 (16000-28000)
Spain	2016	Male	40 to 44	83% (74-90%)	17% (10-26%)	6.8 (4.6-9.7)	10% (7.6-13%)	22% (17-26%)	560 (420-690)	32000 (25000-40000)
Spain	2016	Male	45 to 49	86% (80-91%)	14% (8.5-20%)	8.1 (5.6-12)	13% (10-16%)	23% (19-27%)	1000 (830-1200)	49000 (40000-58000)
Spain	2016	Male	50 to 54	89% (84-93%)	11% (7.2-16%)	9 (6.2-13)	14% (12-17%)	23% (19-26%)	1500 (1300-1800)	64000 (53000-74000)
Spain	2016	Male	55 to 59	90% (86-93%)	10% (7.2-14%)	9.3 (6.4-13)	14% (12-17%)	21% (18-24%)	2000 (1600-2300)	70000 (58000-83000)
Spain	2016	Male	60 to 64	89% (86-92%)	11% (7.5-14%)	9 (6.1-13)	14% (11-16%)	19% (16-22%)	2100 (1700-2500)	65000 (53000-77000)
Spain	2016	Male	65 to 69	88% (84-91%)	12% (8.7-16%)	8 (5.6-11)	12% (10-15%)	16% (13-19%)	2600 (2100-3200)	68000 (55000-81000)
Spain	2016	Male	70 to 74	86% (82-90%)	14% (10-18%)	7 (4.7-10)	11% (8.3-13%)	14% (11-17%)	2800 (2100-3500)	58000 (45000-71000)
Spain	2016	Male	75 to 79	83% (76-89%)	17% (11-24%)	6.1 (4.1-8.9)	9.6% (6.8-12%)	11% (8.3-15%)	2800 (2000-3600)	45000 (32000-57000)
Spain	2016	Male	80 to 84	79% (69-86%)	21% (14-31%)	5.2 (3.3-7.6)	8.3% (5.4-11%)	9.6% (6.3-13%)	3800 (2500-5200)	45000 (29000-59000)
Spain	2016	Male	85 to 89	74% (63-83%)	26% (17-37%)	4.4 (2.7-6.9)	7.1% (4-9.9%)	8.1% (4.5-11%)	3100 (1700-4300)	25000 (14000-35000)
Spain	2016	Male	90 to 94	73% (62-82%)	27% (18-38%)	4.1 (2.4-6.4)	6.2% (3-9.3%)	7% (3.3-11%)	1700 (800-2700)	9500 (4700-14000)
Spain	2016	Male	95 plus	73% (58-83%)	27% (17-42%)	3.8 (2.1-6.2)	5.9% (2-9.7%)	6.5% (2-11%)	450 (140-760)	1600 (550-2700)
Sweden	1990	Female	All Ages	86% (84-88%)	14% (12-16%)	3 (2.5-3.4)	4.5% (2.4-7%)	5.5% (1.2-11%)	2600 (540-5000)	56000 (31000-85000)
Sweden	1990	Female	15 to 19	81% (73-87%)	19% (13-27%)	2.5 (1.6-3.6)	2.9% (1.9-4.4%)	12% (6.2-18%)	8.6 (4.6-13)	880 (540-1300)
Sweden	1990	Female	20 to 24	87% (80-92%)	13% (8-20%)	3 (1.8-4.5)	3.9% (2.6-5.6%)	13% (6.6-21%)	12 (6-19)	1400 (940-2000)
Sweden	1990	Female	25 to 29	90% (84-93%)	10% (6.6-16%)	2.8 (1.8-4)	3.6% (2.2-5.1%)	12% (5.6-18%)	15 (6.9-23)	1500 (910-2100)
Sweden	1990	Female	30 to 34	91% (85-95%)	8.8% (5.2-15%)	2.8 (1.8-4)	3.2% (1.9-4.6%)	11% (5.6-18%)	17 (8.3-26)	1400 (800-2000)
Sweden	1990	Female	35 to 39	91% (87-95%)	8.5% (5.3-13%)	2.9 (1.8-4.1)	3.6% (2.1-5.3%)	12% (6.5-17%)	28 (15-40)	1800 (1000-2600)
Sweden	1990	Female	40 to 44	92% (87-95%)	8.3% (5-13%)	3.3 (2.2-4.6)	4.5% (2.7-6.4%)	13% (8.3-17%)	53 (34-72)	2900 (1800-4100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	1990	Female	45 to 49	92% (87-95%)	8% (4.8-13%)	4.1 (2.7-5.9)	5.9% (3.8-8.6%)	14% (9.6-19%)	85 (57-120)	4000 (2600-5700)
Sweden	1990	Female	50 to 54	92% (88-95%)	7.7% (4.6-12%)	4.7 (3.2-6.5)	6.6% (4.3-9.3%)	14% (9.9-18%)	95 (67-130)	4000 (2700-5500)
Sweden	1990	Female	55 to 59	93% (89-96%)	7.3% (4.3-11%)	5.1 (3.6-7.2)	7.5% (4.8-11%)	13% (8.9-19%)	140 (92-190)	5000 (3200-7100)
Sweden	1990	Female	60 to 64	92% (87-95%)	8.1% (4.8-13%)	5.1 (3.4-7.4)	7% (3.5-11%)	11% (6-17%)	190 (100-290)	5800 (2900-8900)
Sweden	1990	Female	65 to 69	90% (85-94%)	9.7% (5.7-15%)	4.6 (3.3-6)	6% (2.8-9.4%)	8.7% (4-14%)	250 (120-400)	6500 (3000-10000)
Sweden	1990	Female	70 to 74	88% (82-92%)	12% (8.3-18%)	3.9 (2.6-5.4)	4.9% (1.9-1%)	6.6% (1.5-12%)	290 (64-550)	6000 (1300-11000)
Sweden	1990	Female	75 to 79	84% (77-90%)	16% (10-23%)	3.4 (2.1-4.9)	4% (-0.57-9.2%)	5% (-0.81-11%)	350 (-56-810)	5600 (-760-13000)
Sweden	1990	Female	80 to 84	82% (73-88%)	18% (12-27%)	3.3 (2.2-4.5)	3.7% (-0.76-8.8%)	4.3% (-1.3-11%)	410 (-130-1000)	4900 (-950-12000)
Sweden	1990	Female	85 to 89	80% (73-86%)	20% (14-27%)	3.2 (2.1-4.6)	3.5% (-0.84-8.8%)	3.9% (-1.4-10%)	360 (-130-980)	3000 (-750-7600)
Sweden	1990	Female	90 to 94	80% (71-86%)	20% (14-29%)	3.2 (2-4.8)	3.1% (-1.5-9%)	3.4% (-2.2-11%)	190 (-120-580)	1100 (-480-3100)
Sweden	1990	Female	95 plus	78% (67-87%)	22% (13-33%)	3.1 (1.6-5)	3.1% (-2.4-10%)	3.4% (-3.2-12%)	65 (-62-230)	250 (-190-820)
Sweden	1990	Male	All Ages	86% (84-88%)	14% (12-16%)	2.3 (2-2.6)	4.4% (2.3-6.7%)	2.5% (-0.87-6%)	1200 (-420-2900)	60000 (31000-90000)
Sweden	1990	Male	15 to 19	77% (67-85%)	23% (15-33%)	1.5 (0.94-2.2)	5.3% (3.3-7.6%)	9.2% (4.4-15%)	16 (7.8-26)	1800 (1100-2500)
Sweden	1990	Male	20 to 24	86% (78-92%)	14% (8.3-22%)	2.2 (1.4-3.2)	10% (7.1-14%)	13% (6.9-21%)	34 (18-55)	4100 (2800-5800)
Sweden	1990	Male	25 to 29	90% (84-94%)	10% (6-16%)	2.4 (1.5-3.4)	9.9% (6.9-13%)	14% (6.8-20%)	40 (20-61)	4400 (3000-5900)
Sweden	1990	Male	30 to 34	92% (86-96%)	7.8% (4.2-14%)	2.4 (1.5-3.5)	8.8% (6-12%)	13% (7.1-20%)	44 (24-66)	4100 (2800-5600)
Sweden	1990	Male	35 to 39	93% (88-96%)	7.2% (4-12%)	2.3 (1.4-3.3)	8.4% (6.1-11%)	13% (8.2-19%)	58 (36-83)	4500 (3200-5900)
Sweden	1990	Male	40 to 44	93% (89-96%)	6.8% (3.7-11%)	2.5 (1.6-3.5)	8.7% (6.4-11%)	14% (9.3-18%)	98 (65-130)	6100 (4500-7800)
Sweden	1990	Male	45 to 49	94% (89-96%)	6.5% (3.7-11%)	3.1 (2-4.5)	8.6% (5.8-12%)	13% (8.1-18%)	130 (79-180)	6600 (4500-8900)
Sweden	1990	Male	50 to 54	94% (90-97%)	6.1% (3.4-10%)	3.7 (2.6-5.1)	8.1% (5.2-12%)	11% (6.6-16%)	130 (78-190)	5900 (3800-8400)
Sweden	1990	Male	55 to 59	94% (91-97%)	5.7% (3.2-9.5%)	4.3 (2.9-5.9)	7.7% (4.1-12%)	9.6% (4.5-16%)	180 (83-290)	6600 (3500-10000)
Sweden	1990	Male	60 to 64	94% (90-97%)	6.2% (3.3-10%)	4.3 (2.9-6.2)	6% (1.8-11%)	7% (1.6-14%)	210 (47-400)	6700 (2000-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	1990	Male	65 to 69	93% (88-96%)	7.3% (4-12%)	3.8 (2.8-5)	3.5% (0.032-7.3%)	3.7% (-0.83-8.6%)	180 (-42-420)	5100 (47-11000)
Sweden	1990	Male	70 to 74	91% (86-94%)	9.4% (5.9-14%)	3.2 (2.2-4.4)	1.7% (-2.5-7%)	1.5% (-2.9-6.3%)	100 (-200-440)	2700 (-3100-9000)
Sweden	1990	Male	75 to 79	88% (81-93%)	12% (7.3-19%)	2.8 (1.8-4)	0.81% (-2.8-4.7%)	0.5% (-3.7-5%)	45 (-330-450)	1200 (-4300-7300)
Sweden	1990	Male	80 to 84	85% (77-91%)	15% (8.9-23%)	2.7 (1.8-3.6)	0.3% (-2.9-3.8%)	-0.073% (-3.8-3.9%)	-6.7 (-350-360)	330 (-3200-4300)
Sweden	1990	Male	85 to 89	84% (76-90%)	16% (10-24%)	2.5 (1.7-3.5)	0.045% (-3-3.4%)	-0.34% (-3.8-3.5%)	-21 (-230-210)	24 (-1500-1800)
Sweden	1990	Male	90 to 94	84% (75-90%)	16% (10-25%)	2.4 (1.5-3.6)	-0.088% (-3.2-3.6%)	-0.5% (-4.2-3.8%)	-13 (-110-94)	-13 (-490-540)
Sweden	1990	Male	95 plus	83% (73-90%)	17% (9.7-27%)	2.3 (1.3-3.8)	0.05% (-3.1-4.8%)	-0.36% (-4.1-5.1%)	-2.3 (-26-33)	1.3 (-81-120)
Sweden	1995	Female	All Ages	86% (83-88%)	14% (12-17%)	3.2 (2.7-3.7)	5% (3-7.4%)	6.5% (2-12%)	3000 (940-5500)	62000 (36000-90000)
Sweden	1995	Female	15 to 19	80% (72-87%)	20% (13-28%)	2.6 (1.7-3.8)	2.8% (1.8-4.1%)	12% (6.2-18%)	6.5 (3.4-10)	730 (450-1000)
Sweden	1995	Female	20 to 24	87% (79-92%)	13% (8.3-21%)	3.2 (1.9-4.8)	4.2% (2.8-5.9%)	14% (6.9-21%)	11 (5.4-17)	1500 (920-2000)
Sweden	1995	Female	25 to 29	89% (84-93%)	11% (6.6-16%)	3 (2-4.3)	3.6% (2.4-4.9%)	12% (6-18%)	12 (6-18)	1500 (980-2100)
Sweden	1995	Female	30 to 34	91% (85-95%)	8.9% (5.2-15%)	3 (2-4.4)	3.2% (2.1-4.6%)	11% (5.9-17%)	16 (8.1-24)	1500 (930-2100)
Sweden	1995	Female	35 to 39	92% (87-95%)	8.5% (5.1-13%)	3.1 (1.9-4.6)	3.5% (2-5%)	12% (7.2-18%)	24 (14-34)	1700 (1000-2400)
Sweden	1995	Female	40 to 44	92% (87-95%)	8.2% (5-13%)	3.6 (2.4-5)	4.4% (2.8-6.2%)	14% (9.2-18%)	44 (29-58)	2500 (1600-3400)
Sweden	1995	Female	45 to 49	92% (87-95%)	7.8% (4.8-13%)	4.4 (2.9-6.2)	5.9% (3.9-8.4%)	15% (11-20%)	89 (63-120)	4400 (3000-6000)
Sweden	1995	Female	50 to 54	92% (87-95%)	7.6% (4.6-13%)	5.1 (3.6-6.8)	7% (4.8-9.7%)	15% (11-19%)	120 (92-170)	5300 (3700-7300)
Sweden	1995	Female	55 to 59	93% (89-96%)	7.3% (4.3-11%)	5.6 (3.9-7.8)	7.7% (5.1-11%)	14% (9.8-19%)	140 (97-190)	5200 (3500-7200)
Sweden	1995	Female	60 to 64	92% (87-95%)	8.1% (4.8-13%)	5.6 (3.7-8)	7.5% (4.2-11%)	12% (7.4-18%)	180 (110-260)	5700 (3300-8300)
Sweden	1995	Female	65 to 69	90% (85-94%)	9.8% (5.9-15%)	4.9 (3.6-6.5)	6.6% (3.4-9.9%)	9.8% (5.2-14%)	230 (130-350)	6200 (3200-9100)
Sweden	1995	Female	70 to 74	88% (82-92%)	12% (8.5-18%)	4.2 (2.8-5.8)	5.6% (2-9.8%)	7.8% (3-14%)	330 (120-570)	6800 (2400-12000)
Sweden	1995	Female	75 to 79	84% (77-90%)	16% (10-23%)	3.7 (2.3-5.3)	4.9% (0.54-10%)	6.2% (0.65-13%)	400 (42-830)	6400 (740-13000)
Sweden	1995	Female	80 to 84	82% (73-88%)	18% (12-27%)	3.6 (2.4-4.9)	4.7% (0.19-9.6%)	5.6% (0.016-12%)	520 (1.5-1100)	6100 (250-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	1995	Female	85 to 89	80% (72-86%)	20% (14-28%)	3.5 (2.3-5)	4.3% (0.068-9.8%)	5% (-0.27-12%)	510 (-28-1200)	4200 (71-9400)
Sweden	1995	Female	90 to 94	79% (71-86%)	21% (14-29%)	3.4 (2.2-5.2)	3.9% (-0.7-10%)	4.4% (-1.3-12%)	290 (-85-780)	1700 (-300-4100)
Sweden	1995	Female	95 plus	78% (67-87%)	22% (13-33%)	3.3 (1.8-5.4)	3.9% (-1.6-11%)	4.3% (-2.1-13%)	100 (-51-310)	390 (-160-1100)
Sweden	1995	Male	All Ages	86% (83-88%)	14% (12-17%)	2.5 (2.1-2.8)	5.2% (3.2-7.3%)	3.4% (0.2-6.7%)	1600 (96-3200)	67000 (40000-95000)
Sweden	1995	Male	15 to 19	77% (67-85%)	23% (15-33%)	1.6 (0.99-2.3)	4.7% (3-6.8%)	9.3% (4.5-15%)	10 (4.9-16)	1200 (790-1800)
Sweden	1995	Male	20 to 24	86% (78-92%)	14% (8.1-22%)	2.3 (1.5-3.5)	9.9% (7-13%)	13% (6.7-21%)	26 (13-41)	3500 (2300-4900)
Sweden	1995	Male	25 to 29	90% (83-94%)	10% (5.9-17%)	2.5 (1.6-3.7)	9.5% (6.8-12%)	13% (6.6-20%)	31 (15-47)	4000 (2600-5200)
Sweden	1995	Male	30 to 34	92% (86-96%)	7.9% (4.1-14%)	2.6 (1.7-3.7)	8.5% (6-11%)	13% (6.8-20%)	39 (20-59)	4000 (2700-5300)
Sweden	1995	Male	35 to 39	93% (88-96%)	7.3% (4-12%)	2.4 (1.6-3.6)	8.3% (6-11%)	13% (8.1-19%)	51 (31-74)	4200 (3000-5500)
Sweden	1995	Male	40 to 44	93% (89-96%)	6.9% (3.7-11%)	2.7 (1.8-3.8)	8.8% (6.5-11%)	14% (9.3-19%)	81 (53-110)	5400 (3900-6900)
Sweden	1995	Male	45 to 49	93% (89-96%)	6.5% (3.6-11%)	3.3 (2.1-4.7)	9.1% (6.4-12%)	14% (9.1-19%)	130 (85-180)	7300 (5100-9600)
Sweden	1995	Male	50 to 54	94% (90-97%)	6.1% (3.4-10%)	3.9 (2.8-5.4)	9.2% (6.4-12%)	13% (8.7-18%)	180 (120-250)	8200 (5700-11000)
Sweden	1995	Male	55 to 59	94% (90-97%)	5.7% (3.2-9.6%)	4.6 (3.2-6.4)	8.9% (5.3-13%)	12% (6.6-18%)	200 (110-300)	7600 (4700-11000)
Sweden	1995	Male	60 to 64	94% (90-96%)	6.2% (3.5-10%)	4.7 (3.1-6.6)	7.3% (3.3-12%)	8.9% (3.4-15%)	220 (86-370)	7100 (3300-12000)
Sweden	1995	Male	65 to 69	93% (88-96%)	7.5% (4.2-12%)	4.1 (3-5.4)	4.8% (1.4-8.4%)	5.4% (0.99-10%)	210 (40-410)	5800 (1700-11000)
Sweden	1995	Male	70 to 74	90% (85-94%)	9.6% (6-15%)	3.5 (2.3-4.7)	2.8% (-0.86-6.7%)	2.8% (-1.4-7.6%)	180 (-97-490)	4200 (-1300-10000)
Sweden	1995	Male	75 to 79	87% (80-93%)	13% (7.4-20%)	3 (1.9-4.3)	1.7% (-1.7-5.7%)	1.6% (-2.5-6.3%)	130 (-210-510)	2500 (-2600-8100)
Sweden	1995	Male	80 to 84	85% (76-91%)	15% (9.2-24%)	2.8 (1.9-3.9)	1.1% (-2.2-4.6%)	0.83% (-3-4.8%)	75 (-280-430)	1300 (-2500-5200)
Sweden	1995	Male	85 to 89	84% (75-89%)	16% (11-25%)	2.7 (1.8-3.8)	0.82% (-2.1-4.2%)	0.53% (-3-4.3%)	37 (-210-300)	510 (-1300-2600)
Sweden	1995	Male	90 to 94	83% (75-89%)	17% (11-25%)	2.6 (1.6-3.9)	0.65% (-2.4-4.6%)	0.34% (-3.3-4.8%)	11 (-100-150)	120 (-460-830)
Sweden	1995	Male	95 plus	83% (72-90%)	17% (9.8-28%)	2.5 (1.3-4.1)	0.76% (-2.7-5.8%)	0.45% (-3.5-6.2%)	3.3 (-26-47)	23 (-82-170)
Sweden	2000	Female	All Ages	86% (83-88%)	14% (12-17%)	3.3 (2.9-3.8)	5.2% (3.3-7.5%)	7% (2.7-12%)	3400 (1300-5800)	64000 (40000-90000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	2000	Female	15 to 19	80% (72-87%)	20% (13-28%)	2.7 (1.7-3.8)	2.8% (1.7-4.1%)	12% (6.1-18%)	6.5 (3.4-10)	710 (440-1000)
Sweden	2000	Female	20 to 24	86% (79-92%)	14% (8.4-21%)	3.3 (2-4.9)	3.9% (2.6-5.4%)	14% (6.9-21%)	8.7 (4.4-13)	1200 (750-1600)
Sweden	2000	Female	25 to 29	89% (84-93%)	11% (6.8-16%)	3.1 (2-4.6)	3.2% (2.2-4.4%)	12% (6.3-18%)	9.4 (4.8-14)	1200 (790-1700)
Sweden	2000	Female	30 to 34	91% (85-95%)	8.9% (5.1-15%)	3.1 (2-4.5)	2.8% (1.8-3.9%)	12% (6.4-17%)	12 (6.7-18)	1200 (810-1700)
Sweden	2000	Female	35 to 39	91% (87-95%)	8.6% (5.1-13%)	3.2 (2-4.7)	3.3% (2-4.8%)	13% (7.3-18%)	23 (13-33)	1700 (1000-2400)
Sweden	2000	Female	40 to 44	92% (87-95%)	8.2% (5-13%)	3.7 (2.5-5.2)	4.2% (2.7-6%)	14% (9.7-19%)	39 (26-52)	2300 (1500-3200)
Sweden	2000	Female	45 to 49	92% (87-95%)	7.8% (4.8-13%)	4.5 (3-6.5)	5.8% (3.8-8.1%)	15% (11-20%)	75 (54-99)	3700 (2500-5000)
Sweden	2000	Female	50 to 54	93% (88-95%)	7.5% (4.6-12%)	5.3 (3.7-7.1)	6.9% (4.8-9.5%)	15% (11-19%)	130 (99-170)	5600 (4000-7600)
Sweden	2000	Female	55 to 59	93% (89-96%)	7.3% (4.3-11%)	5.8 (4-8)	7.8% (5.4-11%)	14% (11-19%)	180 (130-240)	6700 (4600-9000)
Sweden	2000	Female	60 to 64	92% (87-95%)	8.2% (4.8-13%)	5.8 (3.9-8.1)	7.7% (4.7-11%)	13% (8.3-18%)	190 (120-270)	6000 (3700-8500)
Sweden	2000	Female	65 to 69	90% (84-94%)	10% (5.8-16%)	5.1 (3.7-6.7)	6.9% (4.2-9.9%)	10% (6.5-14%)	220 (140-310)	5800 (3400-8200)
Sweden	2000	Female	70 to 74	87% (81-91%)	13% (8.9-19%)	4.3 (2.9-6)	5.9% (2.5-9.8%)	8.3% (3.7-14%)	290 (130-480)	6100 (2700-10000)
Sweden	2000	Female	75 to 79	83% (75-89%)	17% (11-25%)	3.7 (2.3-5.5)	5.2% (1.1-9.9%)	6.7% (1.5-13%)	420 (93-790)	6800 (1500-13000)
Sweden	2000	Female	80 to 84	80% (71-87%)	20% (13-29%)	3.6 (2.4-5)	5.1% (0.85-9.9%)	6.1% (0.84-12%)	560 (75-1100)	6600 (1100-13000)
Sweden	2000	Female	85 to 89	79% (71-86%)	21% (14-29%)	3.5 (2.3-5)	4.9% (0.71-10%)	5.7% (0.54-12%)	620 (59-1300)	5100 (760-10000)
Sweden	2000	Female	90 to 94	79% (70-85%)	21% (15-30%)	3.5 (2.2-5.3)	4.5% (-0.034-10%)	5.1% (-0.42-12%)	410 (-34-960)	2300 (-17-5200)
Sweden	2000	Female	95 plus	78% (66-87%)	22% (13-34%)	3.4 (1.9-5.5)	4.5% (-1-11%)	5% (-1.4-13%)	160 (-47-430)	600 (-130-1500)
Sweden	2000	Male	All Ages	86% (84-89%)	14% (11-16%)	2.7 (2.3-3.1)	6% (4.1-8%)	4.8% (1.6-8.4%)	2200 (740-3800)	74000 (51000-100000)
Sweden	2000	Male	15 to 19	77% (67-85%)	23% (15-33%)	1.7 (1.1-2.6)	4.6% (2.9-6.6%)	9.8% (4.9-15%)	11 (5.5-17)	1200 (780-1800)
Sweden	2000	Male	20 to 24	86% (79-92%)	14% (8-21%)	2.5 (1.6-3.9)	9.4% (6.3-13%)	14% (7.2-21%)	27 (14-41)	3100 (2000-4200)
Sweden	2000	Male	25 to 29	90% (84-94%)	9.7% (5.8-16%)	2.8 (1.8-4)	8.7% (6.2-11%)	14% (7.2-21%)	30 (15-45)	3300 (2200-4400)
Sweden	2000	Male	30 to 34	93% (87-96%)	7.4% (4-13%)	2.8 (1.8-4.1)	7.7% (5.4-10%)	14% (8-21%)	35 (19-52)	3400 (2300-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	2000	Male	35 to 39	93% (89-96%)	7% (4-11%)	2.7 (1.7-4)	7.8% (5.6-10%)	15% (9-21%)	48 (30-68)	3800 (2700-5100)
Sweden	2000	Male	40 to 44	93% (89-96%)	6.6% (3.6-11%)	2.9 (1.9-4.2)	8.7% (6.4-11%)	16% (11-21%)	74 (50-98)	4700 (3500-6000)
Sweden	2000	Male	45 to 49	94% (90-96%)	6.3% (3.5-10%)	3.6 (2.3-5.2)	9.7% (6.8-13%)	16% (11-22%)	120 (84-160)	6400 (4600-8500)
Sweden	2000	Male	50 to 54	94% (90-97%)	5.9% (3.3-9.8%)	4.3 (3-5.7)	10% (7.5-13%)	16% (11-21%)	210 (150-280)	9300 (6700-12000)
Sweden	2000	Male	55 to 59	95% (91-97%)	5.5% (3.1-9.1%)	5 (3.5-6.8)	10% (7.1-14%)	15% (9.9-20%)	280 (190-380)	10000 (7200-14000)
Sweden	2000	Male	60 to 64	94% (90-97%)	5.9% (3.2-9.7%)	5.1 (3.5-7.1)	8.9% (5.1-13%)	12% (6.4-17%)	270 (150-410)	8700 (5000-13000)
Sweden	2000	Male	65 to 69	93% (88-96%)	7.1% (4-12%)	4.5 (3.3-5.8)	6.3% (3-9.7%)	7.6% (3.3-12%)	250 (110-410)	6800 (3200-11000)
Sweden	2000	Male	70 to 74	91% (86-94%)	9.4% (5.9-14%)	3.8 (2.5-5.2)	4.1% (0.53-8.1%)	4.6% (0.11-9.4%)	240 (6-490)	5200 (720-10000)
Sweden	2000	Male	75 to 79	87% (80-93%)	13% (7.4-20%)	3.3 (2.1-4.7)	2.9% (-0.57-7.1%)	3% (-1.1-8%)	240 (-89-640)	4100 (-800-10000)
Sweden	2000	Male	80 to 84	85% (76-91%)	15% (9.2-24%)	3.1 (2.1-4.2)	2.1% (-1.4-5.9%)	2% (-2.1-6.4%)	180 (-180-570)	2400 (-1500-6700)
Sweden	2000	Male	85 to 89	83% (75-89%)	17% (11-25%)	2.9 (1.9-4.1)	1.8% (-1.3-5.3%)	1.6% (-1.9-5.6%)	120 (-150-420)	1200 (-910-3600)
Sweden	2000	Male	90 to 94	83% (74-89%)	17% (11-26%)	2.8 (1.7-4.2)	1.5% (-1.6-5.7%)	1.4% (-2.2-6.1%)	52 (-86-230)	350 (-380-1300)
Sweden	2000	Male	95 plus	83% (72-90%)	17% (9.8-28%)	2.7 (1.4-4.4)	1.6% (-2-6.8%)	1.4% (-2.7-7.4%)	13 (-26-71)	61 (-76-260)
Sweden	2005	Female	All Ages	86% (84-88%)	14% (12-16%)	3.3 (2.9-3.8)	5% (3.3-6.9%)	6.6% (2.9-11%)	3200 (1400-5400)	60000 (39000-83000)
Sweden	2005	Female	15 to 19	80% (72-87%)	20% (13-28%)	2.6 (1.7-3.8)	2.5% (1.5-3.7%)	11% (5.6-18%)	6.5 (3.3-10)	740 (440-1100)
Sweden	2005	Female	20 to 24	87% (80-92%)	13% (8.1-20%)	3.1 (1.9-4.8)	3.8% (2.4-5.5%)	14% (6.1-22%)	9.2 (4.2-15)	1200 (740-1700)
Sweden	2005	Female	25 to 29	90% (84-94%)	10% (6.5-16%)	3 (2-4.4)	3.1% (2-4.2%)	12% (5.5-18%)	8.2 (3.9-13)	1100 (690-1500)
Sweden	2005	Female	30 to 34	91% (86-95%)	8.8% (5.3-14%)	3 (2-4.4)	2.8% (1.8-3.9%)	11% (6.2-17%)	12 (6.6-18)	1200 (780-1700)
Sweden	2005	Female	35 to 39	92% (87-95%)	8.5% (5.2-13%)	3.1 (1.9-4.6)	3% (1.8-4.3%)	12% (7-17%)	21 (12-30)	1500 (930-2200)
Sweden	2005	Female	40 to 44	92% (87-95%)	8.2% (4.9-13%)	3.6 (2.4-5.1)	3.8% (2.4-5.3%)	14% (9.2-18%)	35 (24-46)	2100 (1300-2900)
Sweden	2005	Female	45 to 49	92% (87-95%)	7.8% (4.7-13%)	4.5 (2.9-6.5)	5.2% (3.4-7.5%)	15% (11-20%)	63 (45-84)	3200 (2100-4400)
Sweden	2005	Female	50 to 54	92% (88-96%)	7.5% (4.4-12%)	5.2 (3.6-7.1)	6.7% (4.6-9.1%)	15% (11-19%)	110 (80-140)	4700 (3300-6300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	2005	Female	55 to 59	93% (89-96%)	7.2% (4.3-11%)	5.7 (4.1-7.7)	7.4% (5.1-10%)	14% (11-18%)	180 (140-230)	6800 (4800-9000)
Sweden	2005	Female	60 to 64	92% (87-95%)	8% (4.7-13%)	5.7 (3.9-8)	7.4% (4.7-10%)	12% (8.5-17%)	220 (150-300)	7000 (4600-9700)
Sweden	2005	Female	65 to 69	90% (85-94%)	9.8% (5.7-15%)	5 (3.7-6.6)	6.6% (4.9-2%)	10% (6.5-13%)	210 (140-290)	5700 (3500-8000)
Sweden	2005	Female	70 to 74	87% (81-91%)	13% (8.5-19%)	4.2 (2.8-5.8)	5.7% (2.7-9%)	8% (4.1-12%)	250 (120-390)	5300 (2500-8400)
Sweden	2005	Female	75 to 79	83% (75-89%)	17% (11-25%)	3.7 (2.3-5.4)	5% (1.2-9.2%)	6.5% (1.8-12%)	340 (94-610)	5500 (1400-10000)
Sweden	2005	Female	80 to 84	80% (70-87%)	20% (13-30%)	3.5 (2.3-4.9)	4.7% (0.93-9.2%)	5.8% (0.99-11%)	520 (87-1000)	6100 (1200-12000)
Sweden	2005	Female	85 to 89	78% (69-85%)	22% (15-31%)	3.4 (2.2-4.8)	4.5% (0.92-9.1%)	5.3% (0.83-11%)	570 (88-1200)	4800 (950-9500)
Sweden	2005	Female	90 to 94	77% (68-85%)	23% (15-32%)	3.3 (2.1-5)	4.2% (0.27-9.6%)	4.8% (-0.019-11%)	410 (-1.6-990)	2300 (140-5300)
Sweden	2005	Female	95 plus	77% (64-86%)	23% (14-36%)	3.2 (1.7-5.2)	4.2% (-0.89-11%)	4.7% (-1.2-13%)	190 (-49-510)	710 (-140-1800)
Sweden	2005	Male	All Ages	87% (85-89%)	13% (11-15%)	2.7 (2.4-3.2)	6.1% (4.4-8.1%)	4.9% (2.1-8.3%)	2200 (920-3700)	74000 (52000-97000)
Sweden	2005	Male	15 to 19	78% (68-85%)	22% (15-32%)	1.7 (1-2.6)	4.4% (2.9-6.3%)	9.4% (4.5-15%)	11 (5.1-17)	1300 (850-1900)
Sweden	2005	Male	20 to 24	87% (79-92%)	13% (7.8-21%)	2.5 (1.6-3.8)	9.3% (6.3-12%)	13% (6.6-20%)	26 (13-39)	3100 (2000-4200)
Sweden	2005	Male	25 to 29	91% (85-94%)	9.3% (5.5-15%)	2.7 (1.7-4)	8.6% (6.1-11%)	14% (6.9-20%)	26 (13-40)	3000 (2100-4100)
Sweden	2005	Male	30 to 34	93% (87-96%)	7.1% (3.9-13%)	2.8 (1.7-4.1)	7.2% (5.1-9.4%)	14% (7.4-20%)	27 (15-40)	2900 (2000-3800)
Sweden	2005	Male	35 to 39	93% (89-96%)	6.7% (3.7-11%)	2.6 (1.6-3.8)	7.1% (5.1-9.3%)	14% (8.3-20%)	40 (23-56)	3400 (2400-4500)
Sweden	2005	Male	40 to 44	94% (90-96%)	6.3% (3.5-10%)	2.9 (1.9-4.1)	8.1% (6-10%)	15% (10-20%)	68 (46-90)	4500 (3300-5800)
Sweden	2005	Male	45 to 49	94% (90-97%)	5.9% (3.5-10%)	3.5 (2.2-5.1)	9.3% (6.7-12%)	16% (11-22%)	110 (74-150)	5800 (4200-7700)
Sweden	2005	Male	50 to 54	94% (91-97%)	5.6% (3.2-9.2%)	4.2 (3-5.7)	10% (7.6-13%)	16% (12-21%)	180 (130-240)	8000 (5800-10000)
Sweden	2005	Male	55 to 59	95% (92-97%)	5.3% (3-8.5%)	4.9 (3.5-6.7)	10% (7.4-14%)	15% (11-20%)	290 (200-380)	11000 (7900-15000)
Sweden	2005	Male	60 to 64	94% (91-97%)	5.7% (3.2-9.3%)	5 (3.5-7)	9.2% (5.8-13%)	12% (7.6-17%)	330 (210-470)	11000 (6900-15000)
Sweden	2005	Male	65 to 69	93% (89-96%)	6.8% (3.8-11%)	4.4 (3.3-5.7)	6.7% (3.6-9.9%)	8.2% (4.3-12%)	280 (140-420)	7400 (4000-11000)
Sweden	2005	Male	70 to 74	91% (86-94%)	9% (5.9-14%)	3.8 (2.6-5.2)	4.5% (1.2-8.3%)	5.2% (1.1-9.7%)	230 (48-440)	5100 (1300-9200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	2005	Male	75 to 79	88% (81-93%)	12% (7.4-19%)	3.3 (2.1-4.8)	3% (-0.057-7.1%)	3.2% (-0.48-8.1%)	210 (-31-520)	3700 (-64-8500)
Sweden	2005	Male	80 to 84	85% (76-91%)	15% (9.2-24%)	3 (2.1-4.1)	2.1% (-0.99-5.8%)	2% (-1.6-6.3%)	180 (-150-560)	2500 (-1200-6800)
Sweden	2005	Male	85 to 89	83% (75-89%)	17% (11-25%)	2.8 (1.9-4)	1.8% (-0.99-5%)	1.7% (-1.5-5.4%)	130 (-120-420)	1300 (-690-3600)
Sweden	2005	Male	90 to 94	82% (73-89%)	18% (11-27%)	2.7 (1.7-4)	1.6% (-1.3-5.4%)	1.4% (-1.9-5.8%)	60 (-79-250)	410 (-340-1400)
Sweden	2005	Male	95 plus	82% (71-90%)	18% (10-29%)	2.6 (1.4-4.3)	1.6% (-1.8-6.4%)	1.4% (-2.6-7%)	17 (-30-84)	78 (-87-310)
Sweden	2010	Female	All Ages	86% (84-88%)	14% (12-16%)	3.2 (2.6-3.7)	4.4% (2.8-6.1%)	5.8% (2.4-9.8%)	2700 (1100-4500)	53000 (34000-74000)
Sweden	2010	Female	15 to 19	80% (72-87%)	20% (13-28%)	2.4 (1.5-3.5)	2.3% (1.4-3.5%)	10% (4.6-17%)	6.1 (2.7-9.8)	730 (440-1100)
Sweden	2010	Female	20 to 24	87% (79-92%)	13% (8.1-21%)	2.9 (1.8-4.5)	3.6% (2.4-5.3%)	13% (5.6-21%)	9.2 (3.9-15)	1300 (790-1800)
Sweden	2010	Female	25 to 29	90% (84-93%)	10% (6.5-16%)	2.8 (1.8-4.2)	3% (2-4.2%)	11% (5.2-17%)	8.1 (3.8-13)	1100 (720-1600)
Sweden	2010	Female	30 to 34	91% (86-95%)	8.6% (5.1-14%)	2.8 (1.8-4.2)	2.6% (1.7-3.8%)	11% (5.8-17%)	9.9 (5.2-15)	1100 (680-1600)
Sweden	2010	Female	35 to 39	92% (87-95%)	8.3% (5-13%)	3 (1.8-4.2)	2.7% (1.7-3.9%)	12% (6.7-16%)	17 (9.7-24)	1300 (810-1900)
Sweden	2010	Female	40 to 44	92% (88-95%)	8% (4.9-12%)	3.4 (2.2-4.8)	3.3% (2.1-4.8%)	13% (8.6-17%)	30 (20-40)	1900 (1200-2700)
Sweden	2010	Female	45 to 49	92% (88-95%)	7.6% (4.7-12%)	4.2 (2.8-6.2)	4.7% (3-6.8%)	15% (10-20%)	56 (40-76)	2900 (1900-4100)
Sweden	2010	Female	50 to 54	93% (88-95%)	7.3% (4.5-12%)	4.9 (3.4-6.8)	6% (3.9-8.4%)	14% (11-19%)	89 (66-120)	4000 (2700-5500)
Sweden	2010	Female	55 to 59	93% (89-96%)	7.1% (4.3-11%)	5.4 (3.7-7.3)	6.7% (4.4-9.3%)	13% (9.8-18%)	130 (95-170)	5100 (3400-7000)
Sweden	2010	Female	60 to 64	92% (87-95%)	7.9% (4.8-13%)	5.4 (3.6-7.6)	6.6% (4.2-9.6%)	11% (7.9-16%)	200 (140-280)	6700 (4100-9500)
Sweden	2010	Female	65 to 69	90% (85-94%)	9.7% (5.7-15%)	4.8 (3.4-6.3)	5.9% (3.6-8.3%)	9.2% (6.2-12%)	220 (150-310)	6100 (3800-8600)
Sweden	2010	Female	70 to 74	88% (82-91%)	12% (8.6-18%)	4 (2.6-5.7)	5% (2.4-8.2%)	7.3% (4.1-11%)	220 (120-350)	4800 (2300-7700)
Sweden	2010	Female	75 to 79	84% (76-89%)	16% (11-24%)	3.5 (2.2-5.2)	4.3% (0.94-8%)	5.8% (1.6-10%)	260 (72-470)	4300 (980-8100)
Sweden	2010	Female	80 to 84	81% (71-87%)	19% (13-29%)	3.3 (2.2-4.6)	4% (0.59-8%)	5% (0.58-10%)	370 (42-750)	4500 (660-9000)
Sweden	2010	Female	85 to 89	78% (69-85%)	22% (15-31%)	3.2 (2.1-4.5)	3.9% (0.59-7.7%)	4.6% (0.54-9.2%)	500 (59-1000)	4200 (620-8400)
Sweden	2010	Female	90 to 94	77% (68-85%)	23% (15-32%)	3.1 (1.9-4.6)	3.6% (-0.11-8.1%)	4.1% (-0.41-9.6%)	370 (-38-870)	2100 (-68-4700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	2010	Female	95 plus	77% (65-86%)	23% (14-35%)	3 (1.6-4.9)	3.7% (-0.96-9.4%)	4.1% (-1.3-11%)	190 (-59-510)	700 (-170-1800)
Sweden	2010	Male	All Ages	87% (85-89%)	13% (11-15%)	2.7 (2.2-3.1)	5.8% (4.2-7.5%)	4.6% (2-7.6%)	2000 (870-3300)	71000 (50000-92000)
Sweden	2010	Male	15 to 19	77% (67-85%)	23% (15-33%)	1.6 (0.97-2.4)	4.3% (2.9-6.1%)	8.9% (3.9-14%)	9.7 (4.3-16)	1300 (870-1900)
Sweden	2010	Male	20 to 24	86% (78-92%)	14% (8-22%)	2.4 (1.5-3.6)	9.3% (6.5-12%)	12% (5.6-20%)	25 (12-41)	3600 (2300-4900)
Sweden	2010	Male	25 to 29	90% (84-94%)	9.8% (5.8-16%)	2.6 (1.6-3.8)	8.7% (6.1-11%)	12% (6-19%)	26 (13-41)	3300 (2200-4500)
Sweden	2010	Male	30 to 34	93% (87-96%)	7.3% (4-13%)	2.6 (1.7-3.9)	7.2% (5.1-9.5%)	13% (6.5-19%)	25 (13-38)	2800 (2000-3800)
Sweden	2010	Male	35 to 39	93% (89-96%)	6.7% (3.8-11%)	2.5 (1.6-3.7)	6.7% (4.9-8.7%)	13% (7.5-18%)	32 (18-45)	3100 (2200-4000)
Sweden	2010	Male	40 to 44	94% (90-96%)	6.3% (3.6-10%)	2.7 (1.8-3.9)	7.4% (5.4-9.5%)	14% (9.3-19%)	55 (36-74)	4000 (2900-5100)
Sweden	2010	Male	45 to 49	94% (90-96%)	6% (3.6-9.8%)	3.4 (2.2-5)	8.5% (6.1-11%)	15% (10-21%)	95 (64-130)	5400 (3800-7200)
Sweden	2010	Male	50 to 54	94% (91-97%)	5.6% (3.2-9.2%)	4 (2.8-5.5)	9.5% (6.9-13%)	15% (11-21%)	150 (110-200)	7000 (5100-9100)
Sweden	2010	Male	55 to 59	95% (91-97%)	5.3% (3-8.6%)	4.7 (3.3-6.5)	9.9% (6.9-13%)	15% (10-20%)	230 (160-310)	8800 (6200-12000)
Sweden	2010	Male	60 to 64	94% (91-97%)	5.7% (3.2-9.2%)	4.8 (3.2-6.8)	8.8% (5.6-13%)	12% (7.6-17%)	330 (200-470)	11000 (6800-15000)
Sweden	2010	Male	65 to 69	93% (89-96%)	6.8% (3.8-11%)	4.2 (3.1-5.6)	6.6% (3.8-9.7%)	8.4% (4.8-12%)	320 (180-470)	8600 (5000-13000)
Sweden	2010	Male	70 to 74	91% (87-94%)	8.9% (5.7-13%)	3.6 (2.4-5)	4.4% (1.6-7.6%)	5.2% (1.7-9.2%)	230 (74-400)	5000 (1900-8700)
Sweden	2010	Male	75 to 79	88% (82-93%)	12% (7.3-18%)	3.1 (2-4.5)	2.9% (0.22-6.3%)	3.1% (-0.11-7.2%)	180 (-6.2-410)	3200 (230-7000)
Sweden	2010	Male	80 to 84	85% (78-91%)	15% (9-22%)	2.9 (1.9-4)	1.9% (-0.82-5%)	1.8% (-1.4-5.6%)	140 (-110-430)	2000 (-860-5300)
Sweden	2010	Male	85 to 89	83% (74-89%)	17% (11-26%)	2.7 (1.8-3.8)	1.6% (-0.92-4.6%)	1.5% (-1.5-4.9%)	130 (-120-420)	1300 (-710-3600)
Sweden	2010	Male	90 to 94	82% (73-89%)	18% (11-27%)	2.6 (1.6-3.9)	1.4% (-1.2-5.1%)	1.3% (-1.8-5.6%)	63 (-88-270)	430 (-360-1500)
Sweden	2010	Male	95 plus	82% (71-90%)	18% (10-29%)	2.5 (1.3-4.1)	1.5% (-1.6-6%)	1.4% (-2.2-6.6%)	19 (-32-94)	87 (-96-360)
Sweden	2016	Female	All Ages	86% (84-88%)	14% (12-16%)	3.2 (2.3-4.2)	4.5% (2.6-6.8%)	6.2% (2-11%)	2900 (930-5300)	56000 (31000-88000)
Sweden	2016	Female	15 to 19	80% (72-87%)	20% (13-28%)	2.4 (1.4-3.8)	2.2% (1.3-3.4%)	11% (4.5-18%)	4.4 (1.8-7.6)	560 (320-820)
Sweden	2016	Female	20 to 24	87% (80-92%)	13% (8.2-20%)	3 (1.7-4.9)	3.6% (2.3-5.3%)	13% (5.4-22%)	10 (4.1-18)	1400 (880-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	2016	Female	25 to 29	89% (84-93%)	11% (6.6-16%)	2.9 (1.7-4.5)	3.1% (2-4.3%)	11% (5-18%)	9.8 (4.1-16)	1300 (830-1900)
Sweden	2016	Female	30 to 34	91% (86-95%)	8.8% (5-14%)	2.9 (1.7-4.5)	2.7% (1.7-4%)	11% (5.5-18%)	11 (5.4-18)	1200 (710-1700)
Sweden	2016	Female	35 to 39	92% (87-95%)	8.4% (5.1-13%)	3 (1.8-4.7)	2.8% (1.6-4.1%)	12% (6.7-18%)	15 (8.3-23)	1200 (740-1800)
Sweden	2016	Female	40 to 44	92% (87-95%)	8.1% (4.9-13%)	3.5 (2.1-5.3)	3.5% (2.1-5.1%)	14% (8.7-19%)	30 (19-44)	1900 (1100-2800)
Sweden	2016	Female	45 to 49	92% (87-95%)	7.7% (4.7-13%)	4.3 (2.5-6.6)	4.7% (2.8-7.1%)	15% (9.9-21%)	60 (39-86)	3100 (1900-4700)
Sweden	2016	Female	50 to 54	93% (88-95%)	7.4% (4.5-12%)	5 (3.1-7.3)	5.9% (3.6-8.6%)	15% (11-20%)	94 (64-130)	4200 (2600-6100)
Sweden	2016	Female	55 to 59	93% (89-96%)	7.2% (4.3-11%)	5.5 (3.5-8.4)	6.7% (4-9.7%)	14% (9.7-19%)	130 (87-180)	5100 (3000-7600)
Sweden	2016	Female	60 to 64	92% (87-95%)	8% (4.8-13%)	5.5 (3.4-8.3)	6.7% (3.6-10%)	12% (7.6-17%)	170 (110-250)	5800 (3200-8800)
Sweden	2016	Female	65 to 69	90% (85-94%)	9.7% (5.7-15%)	4.9 (3.3-7)	6% (3.3-9.3%)	9.5% (5.9-14%)	260 (160-390)	7300 (4000-11000)
Sweden	2016	Female	70 to 74	88% (82-91%)	12% (8.6-18%)	4.1 (2.6-6.2)	5.1% (2.2-8.8%)	7.6% (3.8-12%)	280 (140-460)	6100 (2600-11000)
Sweden	2016	Female	75 to 79	84% (76-89%)	16% (11-24%)	3.6 (2.2-5.4)	4.5% (0.93-8.6%)	6.1% (1.6-11%)	280 (71-550)	4800 (1000-9200)
Sweden	2016	Female	80 to 84	81% (71-87%)	19% (13-29%)	3.4 (2-4.9)	4.3% (0.55-8.9%)	5.5% (0.67-11%)	370 (45-810)	4500 (580-9700)
Sweden	2016	Female	85 to 89	78% (69-85%)	22% (15-31%)	3.2 (2-5)	4.2% (0.32-9%)	5.1% (0.18-11%)	500 (17-1100)	4200 (310-9200)
Sweden	2016	Female	90 to 94	78% (68-85%)	22% (15-32%)	3.2 (1.9-5.1)	3.9% (-0.23-9.2%)	4.6% (-0.5-11%)	470 (-53-1200)	2700 (-150-6300)
Sweden	2016	Female	95 plus	77% (66-86%)	23% (14-34%)	3.1 (1.5-5.2)	4% (-0.95-11%)	4.5% (-1.3-13%)	230 (-62-630)	820 (-180-2200)
Sweden	2016	Male	All Ages	87% (85-89%)	13% (11-15%)	2.7 (2-3.6)	5.7% (3.7-8.2%)	5.1% (2-9%)	2400 (960-4300)	73000 (46000-110000)
Sweden	2016	Male	15 to 19	77% (67-85%)	23% (15-33%)	1.6 (0.93-2.7)	3.9% (2.5-5.7%)	8.9% (3.9-15%)	7.1 (3-12)	990 (620-1400)
Sweden	2016	Male	20 to 24	86% (78-92%)	14% (8.3-22%)	2.4 (1.4-3.9)	8.5% (5.7-12%)	12% (5-20%)	27 (11-46)	3600 (2300-5200)
Sweden	2016	Male	25 to 29	90% (84-94%)	9.8% (5.8-16%)	2.7 (1.6-4.2)	8% (5.4-11%)	12% (5.6-20%)	29 (13-50)	3600 (2300-5000)
Sweden	2016	Male	30 to 34	93% (87-96%)	7.3% (4-13%)	2.7 (1.6-4.2)	6.9% (4.6-9.5%)	13% (6.2-20%)	29 (14-47)	2900 (2000-4200)
Sweden	2016	Male	35 to 39	93% (89-96%)	6.8% (3.9-11%)	2.6 (1.5-4)	6.1% (4.2-8.3%)	13% (6.9-19%)	29 (15-45)	2600 (1700-3600)
Sweden	2016	Male	40 to 44	94% (90-96%)	6.4% (3.6-10%)	2.8 (1.7-4.3)	6.5% (4.5-9%)	14% (8.4-20%)	50 (29-73)	3500 (2400-4800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sweden	2016	Male	45 to 49	94% (90-97%)	6% (3.4-10%)	3.5 (2.1-5.5)	7.8% (5.3-11%)	15% (9.8-21%)	95 (59-140)	5300 (3500-7400)
Sweden	2016	Male	50 to 54	94% (91-97%)	5.6% (3.2-9.1%)	4.2 (2.7-6.1)	9% (6.1-13%)	15% (10-22%)	150 (96-220)	6900 (4600-9800)
Sweden	2016	Male	55 to 59	95% (92-97%)	5.2% (3-8.4%)	4.9 (3-7.3)	9.6% (6.3-14%)	15% (9.7-21%)	210 (140-310)	8400 (5400-12000)
Sweden	2016	Male	60 to 64	94% (91-97%)	5.7% (3.2-9.2%)	4.9 (3-7.3)	8.7% (5-13%)	12% (7-17%)	280 (160-410)	9100 (5000-13000)
Sweden	2016	Male	65 to 69	93% (89-96%)	6.9% (3.9-11%)	4.4 (2.9-6.2)	6.8% (3.7-10%)	8.9% (4.8-14%)	370 (200-590)	10000 (5500-16000)
Sweden	2016	Male	70 to 74	91% (86-94%)	8.8% (5.8-14%)	3.7 (2.3-5.4)	4.7% (1.6-8.7%)	5.8% (1.8-11%)	320 (100-610)	7200 (2500-13000)
Sweden	2016	Male	75 to 79	88% (82-93%)	12% (7-18%)	3.3 (1.9-4.9)	3.3% (0.43-7.2%)	3.7% (0.16-8.3%)	240 (11-550)	4200 (510-9300)
Sweden	2016	Male	80 to 84	86% (78-91%)	14% (8.7-22%)	3 (1.9-4.4)	2.5% (-0.38-6.5%)	2.6% (-0.85-7.4%)	190 (-67-560)	2600 (-380-6700)
Sweden	2016	Male	85 to 89	83% (75-89%)	17% (11-25%)	2.8 (1.8-4.2)	2.2% (-0.58-5.9%)	2.2% (-1-6.6%)	180 (-83-520)	1600 (-450-4500)
Sweden	2016	Male	90 to 94	82% (74-89%)	18% (11-26%)	2.7 (1.5-4.3)	1.9% (-1.6-6.1%)	1.9% (-1.6-6.7%)	110 (-92-400)	700 (-380-2200)
Sweden	2016	Male	95 plus	82% (72-90%)	18% (10-28%)	2.6 (1.3-4.3)	1.9% (-1.4-7.1%)	1.9% (-2-7.9%)	33 (-36-140)	140 (-99-510)
Switzerland	1990	Female	All Ages	88% (85-90%)	12% (9.8-15%)	3.5 (3.2-3.9)	4.9% (3.3-6.6%)	5.4% (1.8-8.9%)	1700 (550-2700)	45000 (31000-60000)
Switzerland	1990	Female	15 to 19	86% (75-92%)	14% (7.6-25%)	3.1 (2.1-4.4)	3.9% (2.3-5.6%)	14% (7.4-20%)	10 (5.6-15)	930 (570-1300)
Switzerland	1990	Female	20 to 24	89% (81-94%)	11% (6.1-19%)	3.9 (2.5-5.5)	5.1% (3.2-7.2%)	16% (9-23%)	20 (11-29)	1800 (1200-2500)
Switzerland	1990	Female	25 to 29	90% (82-95%)	10% (5-18%)	4.1 (2.8-5.5)	4.7% (3.1-6.5%)	16% (9.3-22%)	23 (13-33)	2000 (1300-2700)
Switzerland	1990	Female	30 to 34	90% (82-95%)	10% (5.5-18%)	4.3 (3.2-5.7)	4.7% (3.2-6.4%)	16% (10-22%)	27 (17-36)	2000 (1400-2600)
Switzerland	1990	Female	35 to 39	89% (83-94%)	11% (6.4-17%)	4.6 (3.5-6)	5.3% (3.9-7%)	18% (13-23%)	36 (26-46)	2300 (1700-2900)
Switzerland	1990	Female	40 to 44	89% (83-94%)	11% (6.5-17%)	4.8 (3.6-6.2)	6.5% (4.9-8.5%)	19% (15-24%)	58 (45-72)	3200 (2500-4100)
Switzerland	1990	Female	45 to 49	91% (87-94%)	9% (6.3-13%)	5 (3.9-6.2)	7.8% (6.1-10%)	19% (16-23%)	86 (69-110)	4100 (3300-5000)
Switzerland	1990	Female	50 to 54	92% (89-94%)	7.9% (5.8-11%)	4.9 (3.8-6.2)	8.6% (6.6-11%)	18% (15-22%)	100 (81-130)	4300 (3400-5300)
Switzerland	1990	Female	55 to 59	92% (90-94%)	7.6% (5.7-10%)	4.8 (3.8-6.2)	8.4% (6.5-11%)	16% (13-20%)	130 (100-150)	4500 (3500-5600)
Switzerland	1990	Female	60 to 64	92% (90-94%)	7.6% (5.8-10%)	4.6 (3.6-5.8)	7.3% (5-9.8%)	12% (8.8-16%)	140 (100-190)	4400 (3000-5900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	1990	Female	65 to 69	92% (89-94%)	7.9% (5.8-11%)	4.1 (3.3-5.2)	5.9% (3.5-8.3%)	9.1% (5.3-13%)	160 (96-230)	4200 (2400-5900)
Switzerland	1990	Female	70 to 74	90% (87-93%)	9.7% (7.4-13%)	3.7 (2.8-4.7)	4.5% (1.6-7.5%)	6.3% (2.3-11%)	150 (56-260)	3200 (1200-5300)
Switzerland	1990	Female	75 to 79	87% (82-91%)	13% (8.9-18%)	3.3 (2.5-4.3)	3.4% (0.38-6.8%)	4.2% (-0.07-8.6%)	180 (-3.2-380)	3000 (320-6000)
Switzerland	1990	Female	80 to 84	85% (78-89%)	15% (11-22%)	3.2 (2.4-4.2)	3% (-0.24-6.4%)	3.4% (-0.87-7.8%)	220 (-54-500)	2700 (-230-5800)
Switzerland	1990	Female	85 to 89	83% (77-88%)	17% (12-23%)	3.1 (2.3-4.1)	2.8% (-0.73-6.5%)	3% (-1.4-7.5%)	200 (-89-500)	1700 (-450-4000)
Switzerland	1990	Female	90 to 94	83% (76-88%)	17% (12-24%)	3.1 (2.1-4.2)	2.3% (-1.4-6.3%)	2.4% (-2.2-7.3%)	91 (-84-280)	570 (-350-1600)
Switzerland	1990	Female	95 plus	82% (73-89%)	18% (11-27%)	3 (1.8-4.4)	2.1% (-2-7.2%)	2.1% (-3.1-8.1%)	26 (-39-100)	110 (-110-370)
Switzerland	1990	Male	All Ages	90% (87-93%)	9.8% (7.1-13%)	3.6 (3.2-3.9)	9.2% (7.1-11%)	10% (6.7-13%)	3200 (2100-4300)	94000 (73000-120000)
Switzerland	1990	Male	15 to 19	84% (73-92%)	16% (8-27%)	2.3 (1.5-3.3)	6.9% (3.9-10%)	12% (6-19%)	25 (12-39)	2100 (1200-3200)
Switzerland	1990	Male	20 to 24	89% (81-95%)	11% (5.4-19%)	3.3 (2.2-4.7)	12% (7.1-17%)	17% (8.2-25%)	69 (33-100)	5700 (3400-8000)
Switzerland	1990	Male	25 to 29	91% (85-96%)	8.7% (4.4-15%)	3.4 (2.3-4.9)	11% (6.2-15%)	16% (7.7-25%)	74 (34-110)	5800 (3300-8300)
Switzerland	1990	Male	30 to 34	93% (86-97%)	7.4% (3.5-14%)	3.6 (2.5-5.2)	9.4% (5.8-13%)	15% (7.8-23%)	68 (35-100)	4800 (2900-6800)
Switzerland	1990	Male	35 to 39	93% (87-97%)	7% (3.2-13%)	3.9 (2.8-5.2)	9.2% (6.1-12%)	16% (9.3-22%)	66 (38-93)	4400 (2800-5800)
Switzerland	1990	Male	40 to 44	94% (88-97%)	6.3% (2.9-12%)	4.5 (3.4-5.6)	11% (7.6-14%)	18% (12-24%)	110 (72-140)	5900 (4200-7800)
Switzerland	1990	Male	45 to 49	95% (91-97%)	5.2% (2.9-8.8%)	5.4 (4.3-6.4)	13% (9.9-16%)	20% (16-25%)	160 (120-200)	7600 (6000-9600)
Switzerland	1990	Male	50 to 54	95% (92-97%)	5% (3.1-7.7%)	5.8 (4.6-7)	13% (10-17%)	20% (15-24%)	210 (160-260)	8500 (6500-10000)
Switzerland	1990	Male	55 to 59	94% (92-96%)	5.6% (3.9-7.9%)	5.8 (4.7-6.9)	13% (9.7-15%)	17% (13-21%)	270 (200-340)	9400 (7300-12000)
Switzerland	1990	Male	60 to 64	95% (92-96%)	5.4% (3.7-7.5%)	5.7 (4.7-6.9)	11% (8-14%)	14% (9.6-18%)	310 (220-410)	9400 (6600-12000)
Switzerland	1990	Male	65 to 69	96% (94-97%)	4.3% (2.8-6.1%)	5.5 (4.6-6.6)	9.8% (6.5-13%)	12% (7.4-16%)	380 (250-520)	9400 (6100-13000)
Switzerland	1990	Male	70 to 74	96% (94-97%)	4.1% (2.7-6%)	5.3 (4.3-6.3)	8.5% (5.2-12%)	9.7% (5.6-14%)	370 (220-540)	7400 (4500-11000)
Switzerland	1990	Male	75 to 79	95% (93-97%)	4.7% (2.9-7.3%)	5 (4.1-6)	7.5% (4-11%)	8.3% (4.1-12%)	430 (210-650)	6700 (3500-9900)
Switzerland	1990	Male	80 to 84	94% (91-97%)	5.7% (3.5-8.9%)	4.7 (3.7-5.6)	6.6% (3.1-10%)	7.2% (3-11%)	390 (160-610)	4500 (2100-6800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	1990	Male	85 to 89	93% (89-96%)	6.8% (4.3-11%)	4.4 (3.4-5.4)	5.7% (1.9-9.3%)	6.1% (1.6-10%)	230 (65-390)	1900 (630-3100)
Switzerland	1990	Male	90 to 94	93% (89-96%)	7.2% (4.4-11%)	4.2 (3-5.3)	4.6% (0.65-8.5%)	4.8% (0.21-9.4%)	75 (3.2-150)	430 (61-800)
Switzerland	1990	Male	95 plus	92% (86-96%)	7.8% (3.9-14%)	3.9 (2.5-5.3)	3.9% (-0.49-8.5%)	3.9% (-1.2-9.3%)	14 (-4.5-34)	57 (-7.1-130)
Switzerland	1995	Female	All Ages	88% (85-90%)	12% (9.8-15%)	3.2 (2.9-3.6)	4% (2.5-5.5%)	3.8% (0.29-7.1%)	1200 (94-2300)	37000 (24000-51000)
Switzerland	1995	Female	15 to 19	86% (75-92%)	14% (7.6-25%)	2.8 (1.9-3.9)	3.2% (1.9-4.6%)	12% (6.2-18%)	7.5 (3.9-11)	720 (440-1000)
Switzerland	1995	Female	20 to 24	89% (80-94%)	11% (6.1-20%)	3.5 (2.2-5)	4.2% (2.7-5.9%)	14% (7.4-20%)	13 (7-19)	1300 (880-1900)
Switzerland	1995	Female	25 to 29	90% (82-95%)	10% (5-18%)	3.7 (2.5-5)	3.7% (2.4-5.1%)	12% (6.8-17%)	17 (9.4-24)	1600 (1100-2200)
Switzerland	1995	Female	30 to 34	90% (83-95%)	10% (5.2-17%)	3.9 (2.9-5.2)	3.6% (2.5-4.9%)	11% (7.1-15%)	23 (14-32)	1800 (1300-2400)
Switzerland	1995	Female	35 to 39	89% (83-94%)	11% (6.3-17%)	4.2 (3.2-5.3)	4.4% (3.1-5.7%)	15% (11-19%)	31 (22-41)	2100 (1500-2700)
Switzerland	1995	Female	40 to 44	89% (83-94%)	11% (6.5-17%)	4.4 (3.3-5.6)	5.8% (4.3-7.6%)	17% (13-21%)	53 (40-66)	2900 (2200-3700)
Switzerland	1995	Female	45 to 49	91% (87-94%)	9% (6.4-13%)	4.5 (3.5-5.6)	6.7% (5.1-8.5%)	17% (14-21%)	76 (61-93)	3700 (2900-4500)
Switzerland	1995	Female	50 to 54	92% (89-94%)	7.9% (5.7-11%)	4.5 (3.4-5.7)	7.6% (5.8-9.6%)	16% (13-20%)	100 (84-130)	4400 (3400-5400)
Switzerland	1995	Female	55 to 59	92% (90-94%)	7.6% (5.8-10%)	4.4 (3.5-5.6)	7.5% (5.8-9.5%)	15% (12-18%)	110 (90-140)	4100 (3200-5000)
Switzerland	1995	Female	60 to 64	92% (90-94%)	7.7% (5.8-10%)	4.2 (3.3-5.5)	6.3% (4.3-8.6%)	11% (8.1-14%)	120 (87-160)	3700 (2600-5000)
Switzerland	1995	Female	65 to 69	92% (89-94%)	8% (5.9-11%)	3.8 (3-4.7)	5% (2.9-7.1%)	7.9% (4.7-11%)	130 (79-190)	3400 (2000-4800)
Switzerland	1995	Female	70 to 74	90% (87-93%)	9.8% (7.5-13%)	3.3 (2.6-4.2)	3.5% (1.1-6.1%)	5.1% (1.5-8.6%)	130 (39-230)	2800 (920-4800)
Switzerland	1995	Female	75 to 79	87% (82-91%)	13% (9.3-18%)	3 (2.3-3.9)	2.3% (-0.46-5.2%)	2.9% (-0.87-6.9%)	100 (-30-250)	1800 (-340-4000)
Switzerland	1995	Female	80 to 84	85% (78-89%)	15% (11-22%)	2.9 (2.2-3.8)	1.8% (-1.2-4.9%)	1.9% (-2-5.9%)	120 (-120-360)	1600 (-980-4300)
Switzerland	1995	Female	85 to 89	83% (77-88%)	17% (12-23%)	2.9 (2.1-3.8)	1.5% (-1.6-4.9%)	1.5% (-2.7-5.7%)	110 (-200-410)	1100 (-1200-3400)
Switzerland	1995	Female	90 to 94	83% (76-88%)	17% (12-24%)	2.8 (1.9-3.9)	1.2% (-2.3-5.2%)	0.97% (-3.4-5.9%)	48 (-170-290)	370 (-740-1600)
Switzerland	1995	Female	95 plus	82% (73-89%)	18% (11-27%)	2.7 (1.6-4)	1.1% (-3.1-6.2%)	0.82% (-4.1-7.1%)	15 (-76-130)	81 (-230-460)
Switzerland	1995	Male	All Ages	90% (87-93%)	9.7% (7.1-13%)	3.2 (2.9-3.6)	7.7% (5.9-9.7%)	7.9% (4.8-11%)	2500 (1500-3600)	78000 (59000-98000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	1995	Male	15 to 19	84% (73-92%)	16% (7.9-27%)	2.1 (1.3-3)	5.8% (3.4-8.7%)	11% (5.3-18%)	16 (7.7-26)	1500 (880-2300)
Switzerland	1995	Male	20 to 24	89% (80-94%)	11% (5.6-20%)	3 (1.9-4.3)	10% (6.1-14%)	15% (6.6-22%)	45 (20-68)	4100 (2500-5700)
Switzerland	1995	Male	25 to 29	91% (85-96%)	8.6% (4.3-15%)	3.1 (2.2-4.5)	8.6% (5.2-12%)	13% (5.6-19%)	47 (21-72)	4200 (2500-5900)
Switzerland	1995	Male	30 to 34	93% (86-97%)	7.4% (3.4-14%)	3.3 (2.3-4.7)	7% (4.5-9.7%)	11% (5.3-16%)	49 (24-75)	4000 (2500-5600)
Switzerland	1995	Male	35 to 39	93% (87-97%)	7% (3.1-13%)	3.5 (2.6-4.7)	7.4% (4.9-9.9%)	12% (7.2-17%)	57 (33-81)	4000 (2700-5400)
Switzerland	1995	Male	40 to 44	94% (88-97%)	6.4% (3-12%)	4 (3.1-5.1)	9.1% (6.5-12%)	15% (11-20%)	86 (57-110)	5000 (3600-6500)
Switzerland	1995	Male	45 to 49	95% (91-97%)	5.2% (2.9-8.7%)	4.8 (3.9-5.8)	11% (8.5-14%)	18% (14-23%)	150 (110-190)	7200 (5600-9000)
Switzerland	1995	Male	50 to 54	95% (92-97%)	5% (3.1-7.6%)	5.2 (4.1-6.2)	12% (8.9-15%)	18% (13-22%)	200 (150-250)	8400 (6400-10000)
Switzerland	1995	Male	55 to 59	94% (92-96%)	5.6% (4-8%)	5.2 (4.2-6.1)	11% (8.6-14%)	15% (12-19%)	220 (170-280)	8000 (6000-9900)
Switzerland	1995	Male	60 to 64	95% (92-96%)	5.4% (3.7-7.8%)	5.1 (4.2-6.1)	9.7% (7-13%)	13% (8.8-16%)	260 (180-350)	7900 (5600-10000)
Switzerland	1995	Male	65 to 69	96% (94-97%)	4.3% (2.9-6.2%)	5 (4.1-5.9)	8.2% (5.3-11%)	9.9% (6.1-14%)	290 (180-400)	7300 (4700-9900)
Switzerland	1995	Male	70 to 74	96% (94-97%)	4% (2.7-6.1%)	4.7 (3.9-5.7)	6.7% (3.7-9.8%)	7.7% (3.9-12%)	310 (160-470)	6400 (3500-9300)
Switzerland	1995	Male	75 to 79	95% (93-97%)	4.7% (2.9-7.5%)	4.5 (3.6-5.4)	5.7% (2.5-8.8%)	6.2% (2.4-10%)	280 (110-450)	4500 (2000-7000)
Switzerland	1995	Male	80 to 84	94% (91-97%)	5.6% (3.5-9%)	4.2 (3.4-5.1)	4.7% (1.4-7.9%)	4.9% (1.1-8.8%)	260 (62-460)	3200 (1000-5400)
Switzerland	1995	Male	85 to 89	93% (90-96%)	6.7% (4.3-10%)	3.9 (3-4.9)	3.8% (0.62-7.2%)	3.9% (0.16-7.9%)	170 (6.8-340)	1400 (240-2700)
Switzerland	1995	Male	90 to 94	93% (89-96%)	7.3% (4.4-11%)	3.7 (2.7-4.7)	2.9% (-0.56-6.6%)	2.8% (-1.2-7.2%)	58 (-24-150)	360 (-70-830)
Switzerland	1995	Male	95 plus	92% (87-96%)	7.8% (3.9-13%)	3.5 (2.3-4.8)	2.4% (-1.4-6.5%)	2.2% (-2.2-7%)	11 (-11-35)	49 (-27-130)
Switzerland	2000	Female	All Ages	88% (85-90%)	12% (9.8-15%)	3 (2.7-3.3)	3.3% (2.1-4.7%)	2.8% (-0.23-5.9%)	910 (-72-1900)	31000 (20000-43000)
Switzerland	2000	Female	15 to 19	86% (75-92%)	14% (7.5-25%)	2.5 (1.7-3.6)	2.5% (1.5-3.7%)	11% (5.7-16%)	5.5 (2.8-8.3)	580 (350-840)
Switzerland	2000	Female	20 to 24	89% (80-94%)	11% (5.9-20%)	3.2 (2-4.7)	3.5% (2.3-4.9%)	13% (6.6-20%)	8.4 (4.3-13)	990 (640-1400)
Switzerland	2000	Female	25 to 29	90% (82-95%)	10% (5.1-18%)	3.4 (2.3-4.6)	3.3% (2.1-4.5%)	12% (6.8-18%)	11 (6.1-17)	1200 (770-1600)
Switzerland	2000	Female	30 to 34	90% (82-95%)	10% (5.3-18%)	3.6 (2.6-4.8)	3.2% (2.2-4.3%)	13% (8.7-18%)	17 (11-24)	1500 (1000-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	2000	Female	35 to 39	89% (82-94%)	11% (6.4-18%)	3.9 (2.9-4.9)	3.9% (2.8-5.1%)	15% (11-19%)	30 (21-39)	2100 (1500-2600)
Switzerland	2000	Female	40 to 44	89% (84-94%)	11% (6.5-16%)	4.1 (3-5.2)	4.8% (3.5-6.2%)	16% (12-19%)	43 (33-55)	2500 (1900-3100)
Switzerland	2000	Female	45 to 49	91% (88-94%)	8.7% (6.4-12%)	4.2 (3.3-5.3)	5.6% (4.2-7.1%)	16% (13-19%)	60 (48-74)	3000 (2300-3700)
Switzerland	2000	Female	50 to 54	92% (90-94%)	7.6% (5.7-9.9%)	4.2 (3.2-5.3)	6.5% (4.9-8.3%)	15% (12-18%)	92 (72-110)	3900 (3000-4800)
Switzerland	2000	Female	55 to 59	93% (91-94%)	7.4% (5.7-9.3%)	4.1 (3.3-5.1)	6.8% (5.3-8.6%)	14% (11-16%)	120 (96-140)	4200 (3300-5100)
Switzerland	2000	Female	60 to 64	93% (91-94%)	7.4% (5.8-9.4%)	3.9 (3-5)	5.8% (4.1-7.8%)	11% (8.1-13%)	110 (84-140)	3400 (2500-4400)
Switzerland	2000	Female	65 to 69	92% (90-94%)	7.9% (6-10%)	3.5 (2.8-4.3)	4.3% (2.7-6.1%)	7.4% (4.9-10%)	110 (73-150)	2800 (1700-3900)
Switzerland	2000	Female	70 to 74	90% (88-92%)	9.6% (7.6-12%)	3.1 (2.4-3.9)	3% (1.1-5.1%)	4.7% (1.8-7.7%)	110 (44-180)	2200 (810-3800)
Switzerland	2000	Female	75 to 79	87% (84-91%)	13% (9.1-16%)	2.8 (2.1-3.6)	1.9% (-0.46-4.2%)	2.5% (-0.79-5.6%)	94 (-29-220)	1600 (-390-3700)
Switzerland	2000	Female	80 to 84	85% (79-89%)	15% (11-21%)	2.7 (2-3.4)	1.1% (-1.5-3.7%)	1.2% (-2.2-4.6%)	61 (-120-240)	860 (-1100-2800)
Switzerland	2000	Female	85 to 89	83% (77-88%)	17% (12-23%)	2.6 (1.9-3.4)	0.73% (-2.3-3.7%)	0.56% (-3.3-4.3%)	41 (-240-310)	520 (-1600-2600)
Switzerland	2000	Female	90 to 94	83% (77-88%)	17% (12-23%)	2.6 (1.7-3.5)	0.39% (-3.4%)	0.082% (-4.1-4.5%)	4.7 (-250-260)	150 (-1100-1400)
Switzerland	2000	Female	95 plus	82% (73-89%)	18% (11-27%)	2.5 (1.5-3.6)	0.31% (-3.5-4.7%)	-0.058% (-4.9-5.2%)	-1.4 (-110-120)	30 (-350-450)
Switzerland	2000	Male	All Ages	91% (88-93%)	9.3% (6.8-12%)	3 (2.7-3.3)	6.7% (5-8.4%)	6.5% (3.7-9.5%)	2000 (1100-2900)	64000 (48000-83000)
Switzerland	2000	Male	15 to 19	84% (73-92%)	16% (8.2-27%)	1.9 (1.2-2.7)	4.9% (2.9-7.3%)	10% (4.7-17%)	12 (5.5-21)	1300 (730-1900)
Switzerland	2000	Male	20 to 24	89% (81-95%)	11% (5.5-19%)	2.7 (1.7-3.8)	9.2% (5.7-13%)	14% (6.2-21%)	29 (13-44)	2900 (1800-4100)
Switzerland	2000	Male	25 to 29	91% (84-96%)	8.7% (4.3-16%)	2.9 (1.9-4.1)	8.2% (5.1-11%)	13% (5.7-21%)	31 (13-49)	3000 (1800-4200)
Switzerland	2000	Male	30 to 34	93% (86-97%)	7.4% (3.5-14%)	3 (2.1-4.3)	6.8% (4.3-9.3%)	12% (5.6-19%)	35 (15-54)	3100 (1900-4300)
Switzerland	2000	Male	35 to 39	93% (87-97%)	7% (3.1-13%)	3.2 (2.3-4.2)	7% (4.6-9.3%)	14% (7.1-19%)	50 (25-71)	3700 (2300-4800)
Switzerland	2000	Male	40 to 44	94% (88-97%)	6.4% (3-12%)	3.7 (2.8-4.6)	8.2% (5.8-11%)	15% (10-20%)	76 (50-100)	4500 (3200-5800)
Switzerland	2000	Male	45 to 49	95% (92-97%)	5% (2.9-8.4%)	4.4 (3.6-5.2)	9.9% (7.5-13%)	17% (13-22%)	120 (86-150)	5800 (4400-7200)
Switzerland	2000	Male	50 to 54	96% (93-97%)	4.5% (2.9-6.6%)	4.8 (3.8-5.6)	10% (7.9-13%)	17% (13-21%)	180 (130-220)	7400 (5700-9200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	2000	Male	55 to 59	95% (94-96%)	5% (3.7-6.5%)	4.8 (3.9-5.6)	9.8% (7.5-12%)	14% (11-18%)	210 (160-270)	7700 (5900-9700)
Switzerland	2000	Male	60 to 64	95% (94-96%)	4.9% (3.6-6.4%)	4.7 (3.9-5.6)	8.7% (6.4-11%)	12% (8.4-15%)	220 (150-290)	6700 (4900-8800)
Switzerland	2000	Male	65 to 69	96% (95-97%)	3.9% (2.8-5.4%)	4.5 (3.8-5.3)	7.2% (4.8-9.8%)	9.1% (5.7-12%)	240 (150-330)	6100 (4000-8300)
Switzerland	2000	Male	70 to 74	96% (95-97%)	3.6% (2.5-5.2%)	4.3 (3.6-5.2)	5.7% (3.3-8.4%)	6.8% (3.6-10%)	240 (130-360)	5000 (2900-7500)
Switzerland	2000	Male	75 to 79	96% (94-97%)	4.2% (2.6-6.4%)	4.1 (3.3-4.9)	4.4% (1.8-7.4%)	4.9% (1.6-8.5%)	230 (77-400)	3800 (1500-6200)
Switzerland	2000	Male	80 to 84	95% (92-97%)	5.3% (3.3-8%)	3.8 (3.1-4.6)	3.3% (0.57-6.4%)	3.4% (0.059-7.1%)	160 (2.9-340)	2100 (330-4000)
Switzerland	2000	Male	85 to 89	94% (90-96%)	6.4% (4.1-9.6%)	3.6 (2.8-4.4)	2.4% (-0.31-5.6%)	2.3% (-1-6.2%)	100 (-44-270)	970 (-130-2300)
Switzerland	2000	Male	90 to 94	93% (89-96%)	7% (4.2-11%)	3.4 (2.5-4.2)	1.7% (-1.3-5.1%)	1.4% (-2.2-5.5%)	35 (-52-130)	250 (-190-760)
Switzerland	2000	Male	95 plus	92% (87-96%)	7.7% (3.8-13%)	3.2 (2.1-4.3)	1.3% (-2-4.7%)	0.84% (-3.1-5%)	5 (-19-30)	31 (-48-120)
Switzerland	2005	Female	All Ages	88% (85-90%)	12% (9.7-15%)	2.9 (2.6-3.2)	3.1% (2-4.2%)	2.8% (-0.16-5.5%)	870 (-52-1800)	29000 (19000-39000)
Switzerland	2005	Female	15 to 19	85% (75-92%)	15% (7.6-25%)	2.4 (1.6-3.4)	2.3% (1.3-3.3%)	10% (4.8-16%)	5.1 (2.4-7.9)	560 (330-800)
Switzerland	2005	Female	20 to 24	89% (80-94%)	11% (6.1-20%)	3.1 (1.9-4.5)	3.1% (2-4.3%)	12% (5.9-19%)	7.1 (3.3-11)	910 (580-1300)
Switzerland	2005	Female	25 to 29	90% (82-95%)	10% (5.1-18%)	3.3 (2.2-4.5)	2.9% (1.9-3.8%)	12% (6.1-18%)	8.5 (4.4-12)	960 (620-1300)
Switzerland	2005	Female	30 to 34	90% (82-94%)	10% (5.6-18%)	3.5 (2.5-4.6)	2.9% (2-3.8%)	13% (8.2-18%)	13 (8.3-18)	1200 (800-1600)
Switzerland	2005	Female	35 to 39	89% (83-94%)	11% (6.4-17%)	3.7 (2.9-4.7)	3.3% (2.4-4.3%)	14% (10-18%)	23 (17-30)	1700 (1200-2100)
Switzerland	2005	Female	40 to 44	89% (84-94%)	11% (6.5-16%)	3.9 (3.1-4.9)	4.1% (3-5.3%)	15% (12-19%)	40 (30-50)	2400 (1800-3000)
Switzerland	2005	Female	45 to 49	92% (89-94%)	8.4% (6.3-11%)	4.1 (3.2-5)	5.2% (3.9-6.7%)	15% (12-18%)	61 (49-75)	3000 (2400-3700)
Switzerland	2005	Female	50 to 54	93% (91-94%)	7.3% (5.7-9.1%)	4.1 (3.2-5)	5.7% (4.3-7.2%)	15% (12-17%)	76 (60-91)	3200 (2500-3900)
Switzerland	2005	Female	55 to 59	93% (91-94%)	7.4% (5.9-9.1%)	3.9 (3.2-4.9)	5.9% (4.6-7.5%)	13% (11-15%)	100 (85-130)	3700 (3000-4600)
Switzerland	2005	Female	60 to 64	93% (91-94%)	7.2% (5.7-8.8%)	3.7 (3-4.8)	5.4% (4-7.1%)	10% (8.2-13%)	110 (88-140)	3400 (2500-4500)
Switzerland	2005	Female	65 to 69	93% (91-94%)	7.3% (5.7-9.3%)	3.4 (2.7-4.2)	4.3% (2.9-5.9%)	7.8% (5.5-10%)	110 (74-140)	2600 (1800-3600)
Switzerland	2005	Female	70 to 74	91% (88-92%)	9.4% (7.6-12%)	3 (2.3-3.7)	3% (1.3-4.7%)	5% (2.5-7.6%)	110 (52-160)	2100 (990-3300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	2005	Female	75 to 79	87% (83-91%)	13% (9.4-17%)	2.7 (2.1-3.5)	1.9% (-0.011-3.9%)	2.7% (0.024-5.4%)	93 (0.81-190)	1500 (-10-3100)
Switzerland	2005	Female	80 to 84	85% (79-89%)	15% (11-21%)	2.6 (2-3.3)	1.1% (-1.1-3.3%)	1.3% (-1.7-4.3%)	72 (-97-250)	940 (-930-2900)
Switzerland	2005	Female	85 to 89	83% (77-87%)	17% (13-23%)	2.5 (1.8-3.2)	0.71% (-1.9-3.3%)	0.62% (-3-3.9%)	40 (-200-250)	460 (-1300-2100)
Switzerland	2005	Female	90 to 94	83% (75-88%)	17% (12-25%)	2.5 (1.7-3.3)	0.38% (-2.7-3.6%)	0.14% (-3.9-4.2%)	8.5 (-230-250)	150 (-1100-1400)
Switzerland	2005	Female	95 plus	82% (73-89%)	18% (11-27%)	2.4 (1.5-3.5)	0.26% (-3.5-4.4%)	-0.049% (-4.7-5%)	-1.4 (-140-140)	31 (-420-510)
Switzerland	2005	Male	All Ages	91% (88-93%)	8.9% (6.6-12%)	3 (2.7-3.3)	6.3% (4.7-8%)	6.1% (3.4-8.8%)	1800 (990-2700)	59000 (44000-76000)
Switzerland	2005	Male	15 to 19	84% (73-92%)	16% (8.1-27%)	1.8 (1.1-2.6)	4.5% (2.7-6.6%)	9.8% (4.2-16%)	10 (4.4-17)	1100 (680-1700)
Switzerland	2005	Male	20 to 24	89% (80-94%)	11% (5.6-20%)	2.6 (1.7-3.8)	8.9% (5.8-12%)	14% (5.8-21%)	23 (9.5-36)	2700 (1700-3800)
Switzerland	2005	Male	25 to 29	91% (85-96%)	8.8% (4.3-15%)	2.8 (1.8-3.9)	7.6% (5-10%)	13% (5.3-20%)	21 (8.1-33)	2400 (1500-3300)
Switzerland	2005	Male	30 to 34	93% (86-97%)	7.5% (3.5-14%)	2.9 (2-4.1)	6.2% (3.9-8.6%)	12% (5.5-20%)	23 (10-37)	2300 (1400-3200)
Switzerland	2005	Male	35 to 39	93% (87-97%)	7.1% (3.1-13%)	3.1 (2.2-4.1)	6.4% (4.1-8.6%)	13% (6.5-19%)	41 (21-59)	3100 (2000-4200)
Switzerland	2005	Male	40 to 44	94% (88-97%)	6.4% (3.1-12%)	3.5 (2.7-4.4)	7.2% (5.1-9.4%)	14% (9.4-20%)	67 (42-94)	4100 (2900-5500)
Switzerland	2005	Male	45 to 49	95% (92-97%)	4.9% (2.9-7.8%)	4.2 (3.5-5)	8.9% (6.8-11%)	16% (12-21%)	110 (78-140)	5500 (4200-7000)
Switzerland	2005	Male	50 to 54	96% (94-97%)	4.2% (2.8-6.1%)	4.6 (3.8-5.5)	9.8% (7.4-12%)	16% (12-20%)	150 (110-190)	6500 (5000-8100)
Switzerland	2005	Male	55 to 59	95% (94-96%)	4.8% (3.6-6.2%)	4.6 (3.9-5.4)	9.4% (7.1-12%)	14% (10-18%)	210 (150-260)	7500 (5700-9300)
Switzerland	2005	Male	60 to 64	95% (94-97%)	4.7% (3.4-6.1%)	4.6 (3.8-5.4)	8.3% (6.2-11%)	12% (8.6-15%)	230 (170-290)	7100 (5200-8900)
Switzerland	2005	Male	65 to 69	96% (95-97%)	3.7% (2.6-5.1%)	4.4 (3.7-5.3)	6.8% (4.8-9.1%)	8.9% (6.1-12%)	210 (140-280)	5400 (3800-7300)
Switzerland	2005	Male	70 to 74	97% (95-98%)	3.4% (2.3-4.8%)	4.2 (3.5-5.1)	5.6% (3.4-7.8%)	6.9% (3.8-9.9%)	220 (130-320)	4700 (2800-6700)
Switzerland	2005	Male	75 to 79	96% (94-97%)	4.1% (2.5-6.3%)	4 (3.3-4.8)	4.3% (2-6.8%)	4.9% (1.9-7.9%)	210 (81-340)	3500 (1600-5500)
Switzerland	2005	Male	80 to 84	95% (92-97%)	5.2% (3.2-8.3%)	3.7 (3-4.4)	3.1% (0.53-5.7%)	3.2% (-0.052-6.4%)	160 (-2.7-340)	2100 (340-4000)
Switzerland	2005	Male	85 to 89	94% (90-96%)	6.3% (4.1-9.5%)	3.5 (2.7-4.2)	2.2% (-0.53-5.1%)	2% (-1.3-5.5%)	88 (-55-240)	870 (-210-2000)
Switzerland	2005	Male	90 to 94	93% (89-96%)	6.9% (4.2-11%)	3.3 (2.4-4.1)	1.5% (-1.4-4.7%)	1.2% (-2.3-5.1%)	33 (-61-140)	250 (-240-800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	2005	Male	95 plus	92% (87-96%)	7.6% (3.9-13%)	3.1 (2-4.1)	1% (-2.3-4.5%)	0.6% (-3.4-4.8%)	4.8 (-27-38)	33 (-76-150)
Switzerland	2010	Female	All Ages	88% (86-91%)	12% (9.4-14%)	2.9 (2.5-3.3)	2.9% (1.9-4%)	2.7% (0.019-5.4%)	900 (6.4-1800)	28000 (18000-38000)
Switzerland	2010	Female	15 to 19	85% (75-92%)	15% (7.6-25%)	2.3 (1.5-3.3)	1.9% (1.2-2.8%)	9.4% (4.6-14%)	3.6 (1.8-5.7)	450 (280-640)
Switzerland	2010	Female	20 to 24	89% (80-94%)	11% (6-20%)	3 (1.9-4.4)	2.8% (1.9-3.8%)	12% (5.7-18%)	5.2 (2.5-7.9)	800 (530-1100)
Switzerland	2010	Female	25 to 29	90% (82-95%)	10% (5.2-18%)	3.1 (2.1-4.3)	2.6% (1.7-3.4%)	11% (5.6-17%)	7.2 (3.4-11)	900 (570-1200)
Switzerland	2010	Female	30 to 34	90% (82-95%)	10% (5.3-18%)	3.3 (2.3-4.6)	2.5% (1.7-3.4%)	12% (7.7-17%)	10 (6.4-14)	980 (670-1300)
Switzerland	2010	Female	35 to 39	89% (83-94%)	11% (6.2-17%)	3.6 (2.7-4.6)	3% (2.2-3.9%)	14% (10-18%)	18 (13-22)	1300 (990-1700)
Switzerland	2010	Female	40 to 44	89% (84-94%)	11% (6.5-16%)	3.8 (2.9-4.9)	3.7% (2.7-4.8%)	15% (11-18%)	34 (26-42)	2100 (1500-2600)
Switzerland	2010	Female	45 to 49	92% (89-94%)	8.3% (6.2-11%)	3.9 (3.1-4.9)	4.6% (3.5-6%)	15% (12-18%)	59 (47-72)	3000 (2300-3700)
Switzerland	2010	Female	50 to 54	93% (91-94%)	7.2% (5.5-9%)	3.9 (3.1-4.9)	5.4% (4.1-6.9%)	14% (12-17%)	80 (64-96)	3400 (2600-4200)
Switzerland	2010	Female	55 to 59	93% (91-94%)	7.1% (5.6-8.6%)	3.9 (3.1-4.8)	5.6% (4.3-7.1%)	13% (10-15%)	98 (79-120)	3500 (2700-4300)
Switzerland	2010	Female	60 to 64	93% (92-95%)	6.9% (5.4-8.5%)	3.7 (2.9-4.8)	5.1% (3.7-6.7%)	10% (8.1-12%)	120 (91-150)	3500 (2600-4600)
Switzerland	2010	Female	65 to 69	93% (91-95%)	7% (5.5-8.8%)	3.3 (2.7-4.1)	4.2% (2.8-5.7%)	7.9% (5.8-9.9%)	120 (88-150)	3000 (2000-3900)
Switzerland	2010	Female	70 to 74	91% (89-93%)	8.9% (7.3-11%)	2.9 (2.2-3.7)	2.9% (1.4-4.5%)	5.2% (2.9-7.4%)	100 (57-150)	2000 (960-3200)
Switzerland	2010	Female	75 to 79	88% (84-91%)	12% (9.1-16%)	2.6 (2-3.4)	1.9% (0.12-3.7%)	2.9% (0.43-5.4%)	93 (14-170)	1500 (84-2900)
Switzerland	2010	Female	80 to 84	85% (80-89%)	15% (11-20%)	2.5 (1.9-3.2)	1.1% (-0.99-3.1%)	1.4% (-1.4-4.1%)	77 (-80-230)	950 (-830-2700)
Switzerland	2010	Female	85 to 89	84% (78-88%)	16% (12-22%)	2.4 (1.8-3.2)	0.74% (-1.7-3.1%)	0.73% (-2.5-3.7%)	55 (-180-290)	560 (-1300-2400)
Switzerland	2010	Female	90 to 94	83% (77-88%)	17% (12-23%)	2.4 (1.6-3.3)	0.43% (-2.5-3.5%)	0.26% (-3.5-4%)	15 (-210-240)	170 (-950-1300)
Switzerland	2010	Female	95 plus	82% (73-89%)	18% (11-27%)	2.3 (1.4-3.4)	0.36% (-3.3-4.2%)	0.12% (-4.2-4.9%)	4.3 (-150-170)	52 (-460-600)
Switzerland	2010	Male	All Ages	91% (88-94%)	8.7% (6.4-12%)	2.9 (2.5-3.3)	5.8% (4.4-7.4%)	5.7% (3.1-8.2%)	1700 (930-2600)	55000 (41000-71000)
Switzerland	2010	Male	15 to 19	84% (72-92%)	16% (8-28%)	1.7 (1.1-2.5)	4.1% (2.5-5.8%)	9.3% (3.7-16%)	8.1 (3.1-14)	1000 (600-1500)
Switzerland	2010	Male	20 to 24	89% (80-94%)	11% (5.6-20%)	2.5 (1.6-3.6)	8% (5.4-11%)	13% (5.2-21%)	16 (6.2-26)	2300 (1500-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	2010	Male	25 to 29	91% (84-96%)	8.8% (4.3-16%)	2.7 (1.8-3.8)	7% (4.7-9.5%)	13% (5-20%)	17 (6.7-28)	2300 (1400-3100)
Switzerland	2010	Male	30 to 34	92% (86-96%)	7.5% (3.6-14%)	2.8 (1.9-4.1)	5.6% (3.7-7.8%)	12% (5.3-19%)	19 (8.3-29)	2000 (1300-2700)
Switzerland	2010	Male	35 to 39	93% (87-97%)	7.2% (3.2-13%)	2.9 (2.1-3.9)	5.7% (3.8-7.7%)	13% (6.7-19%)	29 (15-42)	2300 (1500-3100)
Switzerland	2010	Male	40 to 44	94% (88-97%)	6.5% (3-12%)	3.3 (2.5-4.3)	6.5% (4.6-8.7%)	14% (9-20%)	57 (36-80)	3600 (2500-4800)
Switzerland	2010	Male	45 to 49	95% (92-97%)	4.9% (2.8-8.1%)	4 (3.4-4.9)	8% (5.9-10%)	16% (11-20%)	100 (73-130)	5300 (4000-6700)
Switzerland	2010	Male	50 to 54	96% (94-97%)	4.2% (2.7-6%)	4.4 (3.6-5.3)	8.9% (6.7-11%)	15% (12-20%)	140 (110-190)	6300 (4800-7900)
Switzerland	2010	Male	55 to 59	95% (94-96%)	4.7% (3.6-6%)	4.4 (3.7-5.2)	8.8% (6.8-11%)	14% (11-17%)	180 (140-230)	6700 (5200-8400)
Switzerland	2010	Male	60 to 64	95% (94-97%)	4.6% (3.4-5.9%)	4.3 (3.6-5.2)	7.9% (5.9-10%)	11% (8.5-14%)	220 (160-280)	6900 (5200-8700)
Switzerland	2010	Male	65 to 69	96% (95-97%)	3.6% (2.6-5%)	4.2 (3.5-5.1)	6.5% (4.6-8.5%)	8.7% (6.1-11%)	230 (150-300)	5900 (4100-7700)
Switzerland	2010	Male	70 to 74	97% (95-98%)	3.4% (2.4-4.8%)	4 (3.3-4.8)	5.3% (3.3-7.5%)	6.7% (4-9.5%)	200 (120-290)	4300 (2600-6000)
Switzerland	2010	Male	75 to 79	96% (94-97%)	4.1% (2.6-6%)	3.8 (3.1-4.6)	4.1% (2-6.3%)	4.8% (2.1-7.5%)	200 (88-320)	3400 (1600-5200)
Switzerland	2010	Male	80 to 84	95% (92-97%)	5.1% (3.2-7.7%)	3.5 (2.8-4.3)	2.8% (0.51-5.1%)	3% (0.0069-5.8%)	160 (0.36-300)	2000 (360-3700)
Switzerland	2010	Male	85 to 89	94% (91-96%)	6.1% (4-9.1%)	3.3 (2.5-4.1)	2% (-0.28-4.5%)	2.6% (-1-4.9%)	100 (-51-260)	980 (-130-2100)
Switzerland	2010	Male	90 to 94	93% (90-96%)	6.7% (4.2-10%)	3.1 (2.3-4)	1.4% (-1.3-4.2%)	1.2% (-2.1-4.5%)	34 (-61-130)	250 (-220-760)
Switzerland	2010	Male	95 plus	93% (88-96%)	7.5% (3.6-12%)	2.9 (1.9-4)	0.96% (-2.1-4.2%)	0.61% (-3-4.4%)	6.6 (-33-48)	42 (-94-180)
Switzerland	2016	Female	All Ages	88% (86-91%)	12% (9.5-14%)	2.8 (2.2-3.6)	2.8% (1.5-4.3%)	2.6% (-0.42-5.6%)	880 (-140-2000)	28000 (14000-43000)
Switzerland	2016	Female	15 to 19	85% (75-92%)	15% (7.5-25%)	2.2 (1.3-3.3)	1.8% (1.1-2.6%)	9.2% (4.8-15%)	2.7 (1.3-4.5)	380 (230-550)
Switzerland	2016	Female	20 to 24	89% (80-94%)	11% (6.2-20%)	2.9 (1.8-4.4)	2.8% (1.9-3.9%)	11% (5.3-17%)	4.7 (2-8)	790 (500-1200)
Switzerland	2016	Female	25 to 29	90% (82-95%)	10% (5.2-18%)	3.1 (2-4.6)	2.5% (1.6-3.4%)	11% (5.6-17%)	6.4 (2.9-11)	890 (570-1300)
Switzerland	2016	Female	30 to 34	90% (82-95%)	10% (5.2-18%)	3.3 (2.2-4.7)	2.4% (1.6-3.4%)	12% (7.6-17%)	9.8 (5.3-16)	990 (640-1400)
Switzerland	2016	Female	35 to 39	89% (83-94%)	11% (6.4-17%)	3.5 (2.4-4.9)	2.8% (1.9-4%)	14% (10-18%)	15 (8.8-25)	1200 (790-1700)
Switzerland	2016	Female	40 to 44	89% (83-93%)	11% (6.5-17%)	3.8 (2.6-5.2)	3.4% (2.3-4.9%)	15% (11-18%)	27 (17-43)	1700 (1100-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	2016	Female	45 to 49	91% (88-94%)	8.6% (6.3-12%)	3.9 (2.8-5.2)	4.3% (2.9-6%)	15% (12-18%)	52 (34-78)	2700 (1800-3800)
Switzerland	2016	Female	50 to 54	93% (90-94%)	7.3% (5.5-9.5%)	3.9 (2.7-5.3)	5.1% (3.5-7%)	14% (11-17%)	83 (55-120)	3600 (2300-5200)
Switzerland	2016	Female	55 to 59	93% (91-94%)	7% (5.5-8.8%)	3.8 (2.8-5.1)	5.3% (3.7-7.2%)	13% (10-15%)	100 (65-150)	3600 (2400-5200)
Switzerland	2016	Female	60 to 64	93% (91-94%)	7% (5.5-8.8%)	3.6 (2.6-5)	4.7% (3.2-6.5%)	10% (7.5-13%)	100 (66-150)	3200 (2000-4500)
Switzerland	2016	Female	65 to 69	93% (91-94%)	7.2% (5.5-9.3%)	3.3 (2.4-4.4)	4% (2.4-5.8%)	7.8% (5.4-10%)	120 (75-190)	3100 (1800-4700)
Switzerland	2016	Female	70 to 74	91% (89-93%)	8.9% (7.1-11%)	2.9 (2-3.9)	2.8% (1.1-4.9%)	5.1% (2.6-7.8%)	110 (54-190)	2300 (830-4000)
Switzerland	2016	Female	75 to 79	88% (84-91%)	12% (8.9-16%)	2.6 (1.8-3.6)	1.9% (-0.12-4%)	2.9% (0.25-5.8%)	88 (7.3-180)	1400 (-92-3200)
Switzerland	2016	Female	80 to 84	85% (80-89%)	15% (11-20%)	2.5 (1.7-3.5)	1.1% (-1.1-3.5%)	1.5% (-1.6-4.5%)	75 (-85-240)	950 (-920-2900)
Switzerland	2016	Female	85 to 89	84% (78-88%)	16% (12-22%)	2.4 (1.6-3.4)	0.73% (-1.8-3.4%)	0.73% (-2.6-4.2%)	55 (-190-320)	570 (-1400-2700)
Switzerland	2016	Female	90 to 94	83% (77-88%)	17% (12-23%)	2.4 (1.5-3.4)	0.41% (-2.4-3.6%)	0.23% (-3.5-4.2%)	17 (-260-310)	200 (-1200-1700)
Switzerland	2016	Female	95 plus	83% (74-90%)	17% (10-26%)	2.3 (1.3-3.5)	0.31% (-3.2-4.1%)	0.064% (-4.3-4.8%)	2 (-150-170)	45 (-460-590)
Switzerland	2016	Male	All Ages	91% (89-94%)	8.7% (6.5-11%)	2.9 (2.2-3.7)	5.7% (3.8-7.9%)	5.5% (2.4-8.9%)	1800 (730-3300)	57000 (35000-87000)
Switzerland	2016	Male	15 to 19	84% (73-92%)	16% (8.1-27%)	1.7 (1-2.6)	4.1% (2.5-5.9%)	9.2% (3.2-16%)	6.6 (2.2-12)	880 (530-1300)
Switzerland	2016	Male	20 to 24	89% (80-94%)	11% (5.8-20%)	2.5 (1.5-3.7)	8.2% (5.5-11%)	13% (4.9-21%)	14 (5.1-26)	2200 (1400-3300)
Switzerland	2016	Male	25 to 29	91% (84-96%)	8.9% (4.5-16%)	2.6 (1.6-4)	7.2% (4.7-9.9%)	12% (4.9-21%)	17 (6.2-32)	2400 (1500-3400)
Switzerland	2016	Male	30 to 34	92% (85-97%)	7.6% (3.4-15%)	2.8 (1.7-4.2)	5.7% (3.8-8.1%)	12% (5.2-19%)	19 (7.7-36)	2200 (1400-3100)
Switzerland	2016	Male	35 to 39	93% (87-97%)	7.3% (3.3-13%)	2.9 (1.9-4.2)	5.6% (3.7-7.9%)	13% (6.6-19%)	26 (12-43)	2200 (1400-3200)
Switzerland	2016	Male	40 to 44	93% (88-97%)	6.6% (3-12%)	3.3 (2.3-4.7)	6.4% (4.3-9.1%)	14% (9.2-21%)	47 (25-76)	3000 (1900-4400)
Switzerland	2016	Male	45 to 49	95% (91-97%)	5.2% (3-8.7%)	3.9 (2.9-5.2)	7.7% (5.3-11%)	16% (11-22%)	93 (54-150)	4900 (3200-7300)
Switzerland	2016	Male	50 to 54	95% (93-97%)	4.5% (2.9-6.7%)	4.3 (3.1-5.7)	8.5% (5.9-12%)	15% (11-20%)	150 (90-240)	6700 (4400-10000)
Switzerland	2016	Male	55 to 59	95% (93-96%)	5% (3.7-6.6%)	4.3 (3.2-5.7)	8.7% (6.1-12%)	14% (10-18%)	200 (120-300)	7300 (4600-11000)
Switzerland	2016	Male	60 to 64	95% (94-97%)	4.8% (3.5-6.3%)	4.2 (3.1-5.6)	7.8% (5.4-11%)	12% (8.3-15%)	210 (130-320)	6500 (4200-9700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Switzerland	2016	Male	65 to 69	96% (95-97%)	3.8% (2.7-5.3%)	4.1 (3-5.3)	6.5% (4.1-9.3%)	9% (5.9-12%)	240 (140-390)	6400 (3700-10000)
Switzerland	2016	Male	70 to 74	96% (95-97%)	3.6% (2.5-5.1%)	3.9 (2.9-5.1)	5.3% (2.9-8.1%)	6.9% (3.7-10%)	240 (120-410)	5100 (2600-8500)
Switzerland	2016	Male	75 to 79	96% (94-97%)	4.2% (2.7-6.3%)	3.7 (2.7-4.8)	4.1% (1.6-6.9%)	4.9% (1.8-8.2%)	200 (65-390)	3500 (1300-6200)
Switzerland	2016	Male	80 to 84	95% (92-97%)	5.3% (3.3-8.3%)	3.4 (2.5-4.5)	2.9% (0.29-5.9%)	3.1% (-0.14-6.7%)	180 (-9.7-410)	2300 (220-4700)
Switzerland	2016	Male	85 to 89	94% (90-96%)	6.4% (4.1-9.5%)	3.2 (2.3-4.3)	2.1% (-0.56-5.4%)	2.1% (-1.2-5.9%)	120 (-70-350)	1200 (-320-3000)
Switzerland	2016	Male	90 to 94	93% (89-96%)	6.9% (4.2-11%)	3.1 (2-4.2)	1.5% (-1.3-4.8%)	1.3% (-2.1-5.2%)	51 (-87-200)	360 (-340-1100)
Switzerland	2016	Male	95 plus	92% (87-96%)	7.6% (3.8-13%)	2.9 (1.7-4.1)	1% (-2.1-4.6%)	0.74% (-3.2-5%)	8.7 (-37-61)	50 (-100-220)
United Kingdom	1990	Female	All Ages	82% (80-85%)	18% (15-20%)	2.8 (1.9-4.1)	3.8% (1.4-6.9%)	4.5% (0.011-10%)	15000 (37-33000)	340000 (120000-610000)
United Kingdom	1990	Female	15 to 19	79% (70-85%)	21% (15-30%)	2.2 (1.3-3.6)	2.5% (1.5-3.6%)	8.4% (4.5-13%)	45 (24-73)	5200 (3400-7600)
United Kingdom	1990	Female	20 to 24	83% (74-89%)	17% (11-26%)	2.8 (1.6-4.6)	3.5% (2.3-4.9%)	9.9% (5.1-16%)	71 (36-110)	10000 (6600-14000)
United Kingdom	1990	Female	25 to 29	85% (77-90%)	15% (9.9-23%)	2.8 (1.6-4.6)	3.2% (2.1-4.4%)	10% (5.4-15%)	88 (47-140)	10000 (6700-15000)
United Kingdom	1990	Female	30 to 34	87% (78-92%)	13% (8.4-22%)	3 (1.8-4.7)	3.2% (2-4.6%)	11% (6.2-16%)	120 (69-170)	10000 (6600-14000)
United Kingdom	1990	Female	35 to 39	88% (82-93%)	12% (7.3-18%)	3.2 (1.9-5.1)	3.8% (2.4-5.6%)	12% (7.6-17%)	190 (120-270)	13000 (8400-18000)
United Kingdom	1990	Female	40 to 44	89% (83-93%)	11% (6.6-17%)	3.5 (2-5.6)	4.5% (2.8-6.6%)	12% (8.2-17%)	350 (230-480)	19000 (13000-27000)
United Kingdom	1990	Female	45 to 49	89% (82-93%)	11% (6.5-18%)	4 (2.2-6.3)	5.6% (3.3-8.2%)	13% (7.8-18%)	490 (300-690)	23000 (14000-33000)
United Kingdom	1990	Female	50 to 54	89% (83-93%)	11% (6.6-17%)	4.3 (2.5-6.8)	6.2% (3.4-9.1%)	12% (6.9-17%)	680 (400-980)	27000 (15000-40000)
United Kingdom	1990	Female	55 to 59	89% (83-93%)	11% (6.6-17%)	4.6 (2.7-7.4)	6.4% (3.1-9.8%)	10% (5.5-16%)	980 (520-1500)	34000 (16000-52000)
United Kingdom	1990	Female	60 to 64	88% (81-93%)	12% (7.4-19%)	4.6 (2.6-7.4)	5.7% (1.9-9.9%)	8.2% (2.9-14%)	1400 (470-2300)	39000 (13000-68000)
United Kingdom	1990	Female	65 to 69	85% (78-91%)	15% (8.7-22%)	4.1 (2.5-6.2)	4.6% (0.84-8.7%)	6.2% (1.2-12%)	1700 (310-3200)	41000 (7100-78000)
United Kingdom	1990	Female	70 to 74	83% (76-89%)	17% (11-24%)	3.7 (2.1-5.9)	3.7% (-0.38-8.7%)	4.7% (-0.59-11%)	1600 (-210-3800)	32000 (-3400-74000)
United Kingdom	1990	Female	75 to 79	80% (71-87%)	20% (13-29%)	3.4 (1.9-5.4)	2.9% (-1.4-8.3%)	3.5% (-1.9-10%)	1800 (-980-5400)	29000 (-14000-82000)
United Kingdom	1990	Female	80 to 84	78% (67-86%)	22% (14-33%)	3.2 (1.9-5.1)	2.8% (-1.4-8.1%)	3.3% (-1.8-9.6%)	2100 (-1100-6200)	25000 (-12000-70000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	1990	Female	85 to 89	76% (67-83%)	24% (17-33%)	3.1 (1.8-5)	2.9% (-1.2-8.2%)	3.3% (-1.6-9.6%)	1900 (-930-5600)	15000 (-6500-44000)
United Kingdom	1990	Female	90 to 94	76% (66-83%)	24% (17-34%)	3.1 (1.7-5.2)	2.6% (-1.4-8.3%)	3% (-1.9-9.6%)	980 (-630-3200)	5500 (-3000-17000)
United Kingdom	1990	Female	95 plus	75% (61-85%)	25% (15-39%)	3 (1.5-5.3)	2.7% (-1.8-8.7%)	3.1% (-2.4-10%)	380 (-300-1300)	1400 (-910-4500)
United Kingdom	1990	Male	All Ages	86% (83-89%)	14% (11-17%)	2.6 (1.7-3.7)	3.2% (0.7-6.1%)	2.2% (-1.4-6.4%)	6800 (-4500-20000)	300000 (63000-580000)
United Kingdom	1990	Male	15 to 19	78% (69-85%)	22% (15-31%)	1.7 (0.98-2.8)	4.8% (3-6.9%)	8.5% (4.3-13%)	120 (61-190)	12000 (7600-18000)
United Kingdom	1990	Male	20 to 24	85% (77-91%)	15% (9.1-23%)	2.6 (1.5-4.2)	8.9% (6.1-12%)	13% (6.6-19%)	270 (140-400)	30000 (20000-41000)
United Kingdom	1990	Male	25 to 29	88% (81-93%)	12% (7.5-19%)	2.8 (1.7-4.6)	8.3% (5.7-11%)	13% (6.8-19%)	270 (140-410)	28000 (19000-39000)
United Kingdom	1990	Male	30 to 34	90% (83-94%)	10% (5.8-17%)	3 (1.8-4.6)	7.1% (4.7-10%)	12% (6.7-18%)	250 (140-390)	23000 (15000-31000)
United Kingdom	1990	Male	35 to 39	92% (87-95%)	8.5% (5-13%)	3 (1.8-4.7)	6.1% (4-8.8%)	10% (5.3-15%)	270 (140-410)	20000 (13000-29000)
United Kingdom	1990	Male	40 to 44	93% (88-96%)	7.3% (4-12%)	3.2 (1.9-5.1)	5.3% (3.1-8%)	8.1% (3.9-13%)	350 (170-580)	22000 (13000-34000)
United Kingdom	1990	Male	45 to 49	94% (89-96%)	6.4% (3.7-11%)	3.7 (2.1-5.8)	4.9% (2-8.6%)	6.7% (1.8-13%)	410 (110-790)	22000 (9100-39000)
United Kingdom	1990	Male	50 to 54	94% (90-97%)	5.7% (3.1-9.9%)	4.1 (2.3-6.4)	4.4% (1-8.6%)	5.6% (0.48-12%)	530 (45-1100)	24000 (5200-46000)
United Kingdom	1990	Male	55 to 59	95% (91-97%)	5.4% (3-8.6%)	4.4 (2.6-7)	4.1% (0.27-8.8%)	4.8% (-0.35-11%)	750 (-55-1700)	28000 (1900-60000)
United Kingdom	1990	Male	60 to 64	95% (91-97%)	5.5% (3.1-9.1%)	4.3 (2.5-7)	3.2% (-0.8-8.4%)	3.5% (-1.6-9.7%)	920 (-420-2600)	30000 (-7400-76000)
United Kingdom	1990	Male	65 to 69	94% (90-97%)	6% (3.3-9.7%)	3.9 (2.4-5.9)	2.1% (-1.5-6.4%)	2.2% (-2.1-7.3%)	900 (-890-3000)	25000 (-17000-74000)
United Kingdom	1990	Male	70 to 74	93% (89-96%)	6.9% (4.2-11%)	3.5 (2-5.5)	1.4% (-2.2-5.6%)	1.2% (-2.9-6.1%)	580 (-1300-2800)	14000 (-22000-57000)
United Kingdom	1990	Male	75 to 79	92% (87-95%)	8% (4.7-13%)	3.2 (1.8-5.1)	0.91% (-2.5-5.3%)	0.73% (-3-5.7%)	420 (-1800-3300)	8700 (-23000-52000)
United Kingdom	1990	Male	80 to 84	91% (84-95%)	9.3% (5.5-16%)	3 (1.8-4.7)	0.94% (-2.2-5.1%)	0.78% (-2.7-5.5%)	400 (-1400-2800)	5800 (-13000-31000)
United Kingdom	1990	Male	85 to 89	90% (84-94%)	10% (6.4-16%)	2.9 (1.7-4.6)	1% (-1.8-5.1%)	0.9% (-2.4-5.4%)	260 (-690-1600)	2600 (-4700-13000)
United Kingdom	1990	Male	90 to 94	89% (83-93%)	11% (6.5-17%)	2.9 (1.6-4.8)	0.88% (-2-5%)	0.71% (-2.7-5.4%)	72 (-270-540)	530 (-1200-3000)
United Kingdom	1990	Male	95 plus	89% (82-94%)	11% (6-18%)	2.8 (1.4-5)	0.98% (-2.2-5.8%)	0.79% (-2.9-6.4%)	19 (-70-150)	95 (-220-560)
United Kingdom	1995	Female	All Ages	83% (80-86%)	17% (14-20%)	2.8 (1.9-4.1)	3.8% (1.4-6.8%)	4.6% (0.22-9.9%)	15000 (740-33000)	330000 (130000-580000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	1995	Female	15 to 19	80% (72-86%)	20% (14-28%)	2.2 (1.3-3.6)	2.3% (1.5-3.4%)	8.1% (4.4-12%)	37 (20-57)	4400 (2900-6400)
United Kingdom	1995	Female	20 to 24	84% (76-89%)	16% (11-24%)	2.8 (1.6-4.6)	3.4% (2.3-4.7%)	9.7% (5-15%)	59 (30-89)	8400 (5600-12000)
United Kingdom	1995	Female	25 to 29	85% (78-90%)	15% (9.7-22%)	2.8 (1.6-4.6)	3.1% (2.1-4.3%)	9.7% (5.3-15%)	83 (46-130)	9900 (6600-14000)
United Kingdom	1995	Female	30 to 34	87% (80-92%)	13% (8.2-20%)	3 (1.8-4.8)	3.1% (2.1-4.5%)	11% (6.5-15%)	130 (81-190)	11000 (7400-16000)
United Kingdom	1995	Female	35 to 39	89% (83-93%)	11% (7.3-17%)	3.2 (1.9-5)	3.8% (2.5-5.3%)	12% (8.3-17%)	200 (140-280)	14000 (9200-19000)
United Kingdom	1995	Female	40 to 44	90% (84-93%)	10% (6.5-16%)	3.5 (2.1-5.5)	4.7% (3-6.5%)	13% (9-17%)	330 (230-440)	18000 (12000-25000)
United Kingdom	1995	Female	45 to 49	90% (84-94%)	10% (6.3-16%)	4 (2.3-6.3)	5.6% (3.4-8%)	13% (8.6-18%)	570 (370-790)	27000 (17000-38000)
United Kingdom	1995	Female	50 to 54	90% (84-94%)	10% (6.2-16%)	4.3 (2.5-6.8)	6.2% (3.7-9.1%)	12% (7.7-17%)	720 (450-1000)	29000 (17000-42000)
United Kingdom	1995	Female	55 to 59	89% (84-93%)	11% (6.6-16%)	4.6 (2.7-7.4)	6.3% (3.2-9.5%)	11% (5.9-16%)	920 (500-1300)	32000 (16000-48000)
United Kingdom	1995	Female	60 to 64	88% (81-92%)	12% (7.5-19%)	4.5 (2.6-7.3)	5.6% (2.1-9.6%)	8.4% (3.3-14%)	1100 (450-1900)	33000 (12000-57000)
United Kingdom	1995	Female	65 to 69	86% (78-91%)	14% (9.1-22%)	4.1 (2.5-6.3)	4.5% (1-8.4%)	6.3% (1.5-12%)	1400 (350-2700)	35000 (8000-66000)
United Kingdom	1995	Female	70 to 74	83% (76-89%)	17% (11-24%)	3.7 (2.1-5.7)	3.6% (-0.11-8.3%)	4.7% (-0.18-11%)	1800 (-67-4000)	35000 (-960-79000)
United Kingdom	1995	Female	75 to 79	81% (72-87%)	19% (13-28%)	3.4 (1.9-5.6)	2.9% (-1.2-8%)	3.6% (-1.6-10%)	1600 (-720-4600)	25000 (-10000-70000)
United Kingdom	1995	Female	80 to 84	78% (68-86%)	22% (14-32%)	3.2 (1.9-5.2)	2.9% (-1.1-8%)	3.4% (-1.5-9.4%)	2200 (-950-6000)	25000 (-9500-68000)
United Kingdom	1995	Female	85 to 89	77% (68-84%)	23% (16-32%)	3.2 (1.8-5)	3% (-1-8.3%)	3.4% (-1.4-9.7%)	2200 (-860-6200)	18000 (-6000-49000)
United Kingdom	1995	Female	90 to 94	76% (67-84%)	24% (16-33%)	3.1 (1.8-5.2)	2.8% (-1.3-8.6%)	3.2% (-1.6-10%)	1300 (-670-4100)	7300 (-3300-22000)
United Kingdom	1995	Female	95 plus	75% (63-85%)	25% (15-37%)	3.1 (1.6-5.4)	2.9% (-1.6-9.3%)	3.3% (-2-11%)	530 (-320-1700)	1900 (-1000-6100)
United Kingdom	1995	Male	All Ages	87% (85-90%)	13% (10-15%)	2.6 (1.8-3.8)	3.8% (1.5-6.6%)	2.7% (-0.75-7.1%)	8300 (-2300-22000)	340000 (130000-590000)
United Kingdom	1995	Male	15 to 19	80% (72-87%)	20% (13-28%)	1.8 (1-2.9)	4.7% (3-6.6%)	8.6% (4.5-13%)	96 (50-150)	10000 (6600-14000)
United Kingdom	1995	Male	20 to 24	86% (79-92%)	14% (8.4-21%)	2.7 (1.5-4.3)	9.1% (6.2-12%)	12% (6.5-19%)	220 (120-330)	26000 (18000-35000)
United Kingdom	1995	Male	25 to 29	89% (82-93%)	11% (7.1-18%)	2.9 (1.7-4.6)	8.5% (6-11%)	12% (6.6-19%)	270 (140-410)	29000 (20000-40000)
United Kingdom	1995	Male	30 to 34	91% (84-94%)	9.2% (5.5-16%)	3 (1.8-4.8)	7.4% (5.2-10%)	12% (6.9-18%)	310 (180-460)	28000 (19000-37000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	1995	Male	35 to 39	92% (88-95%)	7.9% (4.8-12%)	3 (1.8-4.7)	6.8% (4.7-9.2%)	11% (6.8-17%)	320 (190-470)	24000 (17000-33000)
United Kingdom	1995	Male	40 to 44	93% (89-96%)	6.8% (3.8-11%)	3.3 (2-5.2)	6.3% (4.1-9.1%)	10% (6-15%)	400 (230-610)	25000 (16000-35000)
United Kingdom	1995	Male	45 to 49	94% (90-97%)	5.8% (3.4-9.9%)	3.7 (2.2-6)	6.1% (3.4-9.4%)	9% (4.4-15%)	590 (290-980)	31000 (17000-49000)
United Kingdom	1995	Male	50 to 54	95% (91-97%)	5.2% (2.9-8.7%)	4.2 (2.4-6.6)	5.6% (2.4-9.4%)	7.6% (2.7-13%)	690 (240-1200)	31000 (13000-52000)
United Kingdom	1995	Male	55 to 59	95% (92-97%)	4.9% (2.8-8%)	4.5 (2.6-7.1)	5.1% (1.4-9.4%)	6.3% (1.4-12%)	880 (190-1700)	33000 (9100-62000)
United Kingdom	1995	Male	60 to 64	95% (92-97%)	5.1% (2.8-8.3%)	4.4 (2.5-7.2)	4% (0.073-9%)	4.6% (-0.37-11%)	1000 (-81-2400)	32000 (600-70000)
United Kingdom	1995	Male	65 to 69	94% (91-97%)	5.6% (3.1-9%)	4 (2.4-6.1)	2.7% (-0.66-6.9%)	2.9% (-1.2-8%)	1000 (-430-2800)	27000 (-6700-69000)
United Kingdom	1995	Male	70 to 74	94% (90-96%)	6.4% (4-10%)	3.5 (2.1-5.5)	1.7% (-1.7-5.9%)	1.7% (-2.4-6.6%)	840 (-1200-3300)	19000 (-20000-67000)
United Kingdom	1995	Male	75 to 79	93% (88-96%)	7.4% (4.3-12%)	3.2 (1.8-5.3)	1.2% (-2.1-5.6%)	1.1% (-2.7-6.2%)	530 (-1400-3100)	10000 (-18000-48000)
United Kingdom	1995	Male	80 to 84	91% (86-95%)	8.5% (4.8-14%)	3.1 (1.9-4.9)	1.2% (-1.9-5.4%)	1% (-2.5-5.8%)	550 (-1300-3000)	7600 (-12000-34000)
United Kingdom	1995	Male	85 to 89	91% (86-94%)	9.4% (6-14%)	3 (1.8-4.7)	1.3% (-1.6-5.3%)	1.2% (-2.1-5.7%)	420 (-730-2000)	4000 (-4800-16000)
United Kingdom	1995	Male	90 to 94	90% (85-94%)	9.8% (6.3-15%)	3 (1.7-4.9)	1.2% (-1.8-5.6%)	1.1% (-2.5-6.1%)	150 (-340-850)	1000 (-1600-4700)
United Kingdom	1995	Male	95 plus	90% (83-94%)	10% (5.5-17%)	2.9 (1.5-5)	1.3% (-2.6-2%)	1.2% (-2.7-6.8%)	35 (-80-200)	160 (-240-740)
United Kingdom	2000	Female	All Ages	83% (81-86%)	17% (14-19%)	2.9 (1.9-4.2)	3.9% (1.8-6.6%)	4.8% (0.83-10%)	15000 (2600-32000)	330000 (150000-560000)
United Kingdom	2000	Female	15 to 19	80% (72-86%)	20% (14-28%)	2.2 (1.3-3.7)	2.2% (1.4-3.1%)	7.8% (4.3-12%)	35 (19-56)	4200 (2700-6100)
United Kingdom	2000	Female	20 to 24	83% (75-89%)	17% (11-25%)	2.8 (1.6-4.8)	3.2% (2.2-4.4%)	9.4% (4.9-14%)	50 (26-77)	7100 (4700-9800)
United Kingdom	2000	Female	25 to 29	85% (78-90%)	15% (9.8-22%)	2.8 (1.6-4.6)	3% (2.1-4.1%)	9.7% (5.6-15%)	70 (40-110)	8400 (5600-12000)
United Kingdom	2000	Female	30 to 34	87% (79-92%)	13% (8.2-21%)	3 (1.8-4.8)	3.1% (2.1-4.4%)	11% (7.3-15%)	130 (85-180)	11000 (7600-15000)
United Kingdom	2000	Female	35 to 39	89% (83-93%)	11% (7.3-17%)	3.2 (1.9-5.2)	3.9% (2.6-5.4%)	13% (8.9-17%)	230 (160-310)	15000 (10000-21000)
United Kingdom	2000	Female	40 to 44	89% (83-93%)	11% (6.6-17%)	3.6 (2.2-5.7)	4.8% (3.2-6.7%)	14% (10-18%)	360 (260-470)	20000 (14000-27000)
United Kingdom	2000	Female	45 to 49	90% (83-94%)	10% (6.3-17%)	4 (2.3-6.3)	5.8% (3.7-8.1%)	14% (9.7-19%)	530 (370-710)	25000 (17000-34000)
United Kingdom	2000	Female	50 to 54	90% (84-94%)	10% (6.2-16%)	4.4 (2.5-7)	6.3% (3.9-8.8%)	13% (8.6-18%)	830 (560-1100)	34000 (21000-47000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	2000	Female	55 to 59	89% (84-93%)	11% (6.6-16%)	4.7 (2.7-7.6)	6.3% (3.6-9.1%)	11% (6.9-16%)	940 (580-1300)	33000 (18000-47000)
United Kingdom	2000	Female	60 to 64	88% (81-93%)	12% (7.5-19%)	4.6 (2.7-7.6)	5.6% (2.5-8.9%)	8.9% (4.4-14%)	1100 (530-1700)	32000 (14000-51000)
United Kingdom	2000	Female	65 to 69	86% (78-91%)	14% (9-22%)	4.2 (2.6-6.5)	4.6% (1.6-8.1%)	6.7% (2.4-12%)	1200 (450-2200)	31000 (11000-54000)
United Kingdom	2000	Female	70 to 74	83% (75-89%)	17% (11-25%)	3.7 (2.1-6)	3.8% (0.23-8%)	5.1% (0.43-10%)	1500 (130-3200)	30000 (1800-63000)
United Kingdom	2000	Female	75 to 79	81% (72-87%)	19% (13-28%)	3.4 (1.9-5.8)	3.1% (-0.66-7.7%)	3.9% (-0.84-9.7%)	1900 (-400-4600)	29000 (-6100-72000)
United Kingdom	2000	Female	80 to 84	78% (68-86%)	22% (14-32%)	3.3 (2-5.4)	3.1% (-0.62-7.7%)	3.7% (-0.8-9.5%)	2000 (-430-5100)	23000 (-4500-59000)
United Kingdom	2000	Female	85 to 89	77% (68-84%)	23% (16-32%)	3.2 (1.9-5.2)	3.1% (-0.66-8%)	3.6% (-0.87-9.5%)	2300 (-550-6000)	19000 (-3900-49000)
United Kingdom	2000	Female	90 to 94	76% (67-84%)	24% (16-33%)	3.2 (1.8-5.3)	2.9% (-1-8.1%)	3.3% (-1.3-9.7%)	1500 (-610-4400)	8300 (-3000-24000)
United Kingdom	2000	Female	95 plus	75% (63-85%)	25% (15-37%)	3.1 (1.5-5.5)	2.9% (-1.3-9.1%)	3.3% (-1.8-11%)	650 (-350-2100)	2400 (-1100-7400)
United Kingdom	2000	Male	All Ages	87% (85-90%)	13% (10-15%)	2.7 (1.8-4)	4.5% (2.3-7.2%)	3.6% (0.4-7.6%)	11000 (1200-22000)	390000 (210000-630000)
United Kingdom	2000	Male	15 to 19	80% (72-86%)	20% (14-28%)	1.8 (1-3)	4.5% (2.9-6.3%)	8.7% (4.6-13%)	92 (48-140)	9900 (6300-14000)
United Kingdom	2000	Male	20 to 24	86% (79-91%)	14% (8.6-21%)	2.7 (1.5-4.5)	8.7% (6.1-11%)	12% (6.7-19%)	190 (100-280)	22000 (15000-29000)
United Kingdom	2000	Male	25 to 29	89% (82-93%)	11% (7-18%)	2.9 (1.7-4.6)	8.3% (5.9-11%)	13% (6.9-19%)	240 (130-360)	25000 (17000-34000)
United Kingdom	2000	Male	30 to 34	91% (85-95%)	9.2% (5.2-15%)	3 (1.8-4.8)	7.7% (5.5-10%)	13% (8-19%)	330 (200-470)	28000 (20000-38000)
United Kingdom	2000	Male	35 to 39	92% (88-95%)	7.9% (4.7-12%)	3.1 (1.8-5)	7.5% (5.3-10%)	13% (8.7-19%)	420 (270-580)	30000 (21000-40000)
United Kingdom	2000	Male	40 to 44	93% (89-96%)	6.8% (3.9-11%)	3.3 (2-5.4)	7.5% (5.3-10%)	13% (8.8-18%)	510 (350-730)	31000 (22000-41000)
United Kingdom	2000	Male	45 to 49	94% (90-97%)	5.9% (3.5-10%)	3.8 (2.2-6.1)	7.6% (4.9-11%)	12% (7.5-18%)	690 (440-1000)	35000 (23000-50000)
United Kingdom	2000	Male	50 to 54	95% (91-97%)	5.2% (2.9-8.9%)	4.2 (2.4-6.7)	7% (4-11%)	10% (5.7-16%)	1000 (550-1600)	43000 (25000-65000)
United Kingdom	2000	Male	55 to 59	95% (92-97%)	4.9% (2.8-7.9%)	4.5 (2.7-7.3)	6.3% (3-10%)	8.4% (3.8-14%)	1100 (500-1800)	40000 (20000-66000)
United Kingdom	2000	Male	60 to 64	95% (92-97%)	5.1% (2.9-8.3%)	4.5 (2.6-7.4)	5.1% (1.6-9.4%)	6.2% (1.7-12%)	1200 (330-2300)	37000 (12000-69000)
United Kingdom	2000	Male	65 to 69	94% (91-97%)	5.6% (3.1-9%)	4.1 (2.5-6.4)	3.6% (0.49-7.5%)	4.1% (0.28-8.8%)	1200 (80-2500)	31000 (4100-64000)
United Kingdom	2000	Male	70 to 74	94% (90-96%)	6.4% (4-10%)	3.6 (2.1-5.9)	2.4% (-0.84-6.5%)	2.5% (-1.3-7.6%)	1000 (-540-3100)	23000 (-7600-63000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	2000	Male	75 to 79	93% (88-96%)	7.3% (4.4-12%)	3.3 (1.9-5.5)	1.7% (-1.4-6.1%)	1.7% (-1.9-6.8%)	900 (-1000-3600)	16000 (-12000-56000)
United Kingdom	2000	Male	80 to 84	92% (86-95%)	8.3% (5-14%)	3.2 (1.9-5.1)	1.6% (-1.4-5.5%)	1.5% (-1.9-6%)	690 (-890-2700)	9100 (-8100-31000)
United Kingdom	2000	Male	85 to 89	91% (86-94%)	9.2% (5.9-14%)	3.1 (1.9-5)	1.7% (-1.2-5.6%)	1.6% (-1.7-6.1%)	610 (-630-2300)	5600 (-3900-18000)
United Kingdom	2000	Male	90 to 94	90% (86-94%)	9.6% (6.1-14%)	3 (1.7-5.1)	1.5% (-1.3-5.9%)	1.5% (-1.9-6.6%)	250 (-320-1100)	1600 (-1400-6200)
United Kingdom	2000	Male	95 plus	90% (83-94%)	10% (5.7-17%)	3 (1.5-5.4)	1.7% (-1.5-6.8%)	1.6% (-2.1-7.5%)	63 (-83-310)	270 (-240-1100)
United Kingdom	2005	Female	All Ages	83% (81-86%)	17% (14-19%)	3 (2-4.3)	4% (2-6.3%)	5% (1.3-9.6%)	15000 (4000-29000)	330000 (170000-520000)
United Kingdom	2005	Female	15 to 19	80% (72-86%)	20% (14-28%)	2.2 (1.3-3.7)	2.1% (1.4-3.1%)	7.6% (4.1-12%)	33 (18-52)	4400 (2800-6400)
United Kingdom	2005	Female	20 to 24	83% (75-89%)	17% (11-25%)	2.8 (1.6-4.7)	3.2% (2.3-4.4%)	9.1% (4.9-14%)	48 (26-76)	7800 (5100-11000)
United Kingdom	2005	Female	25 to 29	85% (78-90%)	15% (9.8-22%)	2.9 (1.7-4.8)	3.1% (2.2-4.2%)	9.5% (5.6-14%)	61 (36-92)	8000 (5300-11000)
United Kingdom	2005	Female	30 to 34	87% (79-92%)	13% (8.2-21%)	3 (1.8-4.9)	3.2% (2.2-4.3%)	11% (7.4-15%)	110 (74-150)	10000 (6900-14000)
United Kingdom	2005	Female	35 to 39	89% (83-93%)	11% (7.2-17%)	3.2 (1.9-5.1)	3.9% (2.7-5.3%)	13% (9.5-18%)	230 (160-300)	15000 (11000-21000)
United Kingdom	2005	Female	40 to 44	89% (83-93%)	11% (6.5-17%)	3.6 (2.1-5.7)	4.8% (3.3-6.5%)	14% (10-18%)	390 (280-500)	22000 (15000-29000)
United Kingdom	2005	Female	45 to 49	90% (83-94%)	10% (6.3-17%)	4.1 (2.4-6.6)	5.8% (3.9-8.1%)	15% (10-19%)	560 (400-730)	27000 (18000-36000)
United Kingdom	2005	Female	50 to 54	90% (84-94%)	10% (6.4-16%)	4.5 (2.6-7.2)	6.4% (4.1-8.7%)	13% (9.4-18%)	750 (520-990)	31000 (20000-42000)
United Kingdom	2005	Female	55 to 59	89% (84-93%)	11% (6.6-16%)	4.7 (2.8-7.6)	6.3% (3.8-8.9%)	12% (7.7-16%)	1100 (700-1400)	38000 (23000-52000)
United Kingdom	2005	Female	60 to 64	88% (82-93%)	12% (7.3-18%)	4.7 (2.7-7.6)	5.7% (3-8.5%)	9.3% (5.4-13%)	1100 (630-1600)	33000 (17000-49000)
United Kingdom	2005	Female	65 to 69	86% (78-91%)	14% (8.6-22%)	4.2 (2.6-6.6)	4.7% (2-7.6%)	7.1% (3.4-11%)	1200 (570-1900)	30000 (12000-49000)
United Kingdom	2005	Female	70 to 74	83% (76-89%)	17% (11-24%)	3.8 (2.1-6.1)	3.9% (0.91-7.5%)	5.5% (1.5-10%)	1400 (370-2500)	27000 (5900-52000)
United Kingdom	2005	Female	75 to 79	80% (72-87%)	20% (13-28%)	3.5 (1.9-5.7)	3.3% (-0.011-7.4%)	4.3% (0.062-9.3%)	1700 (25-3700)	26000 (-98-59000)
United Kingdom	2005	Female	80 to 84	78% (68-86%)	22% (14-32%)	3.3 (2-5.4)	3.2% (-0.11-7.4%)	3.9% (-0.13-9.1%)	2300 (-75-5200)	26000 (-940-60000)
United Kingdom	2005	Female	85 to 89	77% (67-83%)	23% (17-33%)	3.2 (1.9-5.3)	3.2% (-0.37-7.7%)	3.8% (-0.49-9.1%)	2100 (-280-5100)	17000 (-1900-42000)
United Kingdom	2005	Female	90 to 94	76% (67-83%)	24% (17-33%)	3.2 (1.8-5.4)	2.9% (-0.71-7.7%)	3.4% (-0.91-9.2%)	1600 (-430-4300)	8900 (-2200-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	2005	Female	95 plus	75% (62-85%)	25% (15-38%)	3.1 (1.5-5.6)	3% (-1.8-3%)	3.4% (-1.4-9.8%)	780 (-330-2200)	2800 (-960-8000)
United Kingdom	2005	Male	All Ages	88% (85-90%)	12% (10-15%)	2.8 (1.9-4)	5.1% (3.1-7.6%)	4.5% (1.5-8.4%)	13000 (4100-23000)	430000 (260000-640000)
United Kingdom	2005	Male	15 to 19	80% (72-86%)	20% (14-28%)	1.8 (1-3.1)	4.3% (2.8-5.9%)	8.8% (4.7-14%)	83 (44-130)	9600 (6200-14000)
United Kingdom	2005	Male	20 to 24	86% (79-91%)	14% (8.5-21%)	2.8 (1.5-4.5)	8.4% (6.1-11%)	12% (6.9-19%)	170 (96-270)	22000 (15000-30000)
United Kingdom	2005	Male	25 to 29	89% (82-93%)	11% (6.9-18%)	2.9 (1.7-4.9)	8% (5.8-10%)	13% (7.5-19%)	190 (110-280)	21000 (15000-29000)
United Kingdom	2005	Male	30 to 34	91% (85-95%)	9.2% (5.4-15%)	3.1 (1.8-4.9)	7.6% (5.6-10%)	13% (8.5-19%)	280 (180-390)	25000 (18000-33000)
United Kingdom	2005	Male	35 to 39	92% (88-95%)	7.9% (4.8-12%)	3.1 (1.8-4.9)	7.7% (5.6-10%)	14% (9.6-19%)	440 (300-590)	31000 (23000-40000)
United Kingdom	2005	Male	40 to 44	93% (89-96%)	6.8% (3.9-11%)	3.4 (2-5.3)	8% (5.8-11%)	14% (10-19%)	620 (440-830)	36000 (27000-48000)
United Kingdom	2005	Male	45 to 49	94% (90-97%)	5.9% (3.4-10%)	3.9 (2.3-6.2)	8.4% (5.7-11%)	14% (9.5-19%)	820 (560-1100)	40000 (28000-55000)
United Kingdom	2005	Male	50 to 54	95% (91-97%)	5.2% (3-8.9%)	4.3 (2.5-7)	8.1% (5.2-11%)	12% (7.8-18%)	1000 (650-1500)	44000 (28000-63000)
United Kingdom	2005	Male	55 to 59	95% (92-97%)	4.9% (2.9-7.7%)	4.7 (2.7-7.4)	7.3% (4.2-11%)	10% (5.8-15%)	1400 (800-2100)	51000 (30000-76000)
United Kingdom	2005	Male	60 to 64	95% (92-97%)	5% (2.9-8.1%)	4.6 (2.7-7.5)	6.1% (2.9-9.9%)	7.8% (3.6-13%)	1400 (670-2400)	44000 (21000-72000)
United Kingdom	2005	Male	65 to 69	94% (91-97%)	5.6% (3.1-9.2%)	4.2 (2.6-6.5)	4.5% (1.5-8%)	5.4% (1.7-9.8%)	1300 (430-2500)	35000 (12000-63000)
United Kingdom	2005	Male	70 to 74	94% (90-96%)	6.4% (4-10%)	3.7 (2.1-5.9)	3.2% (0.27-6.9%)	3.6% (0.095-8.1%)	1200 (33-2800)	26000 (2300-57000)
United Kingdom	2005	Male	75 to 79	93% (89-96%)	7.3% (4.4-11%)	3.4 (1.9-5.6)	2.4% (-0.55-6.2%)	2.5% (-0.92-7%)	1100 (-420-3200)	19000 (-4400-50000)
United Kingdom	2005	Male	80 to 84	92% (86-95%)	8.3% (4.9-14%)	3.3 (2-5.3)	2.1% (-0.61-5.8%)	2.1% (-0.97-6.3%)	1100 (-500-3200)	14000 (-3900-38000)
United Kingdom	2005	Male	85 to 89	91% (86-94%)	9.1% (5.8-14%)	3.2 (1.9-5.1)	2.1% (-0.61-6%)	2.1% (-0.99-6.5%)	750 (-350-2300)	6600 (-1900-19000)
United Kingdom	2005	Male	90 to 94	91% (86-94%)	9.5% (6.1-14%)	3.1 (1.8-5.3)	2% (-0.91-6.4%)	2% (-1.3-7.1%)	390 (-260-1400)	2400 (-1100-7800)
United Kingdom	2005	Male	95 plus	90% (83-94%)	10% (5.5-17%)	3 (1.5-5.5)	2% (-1.1-7.2%)	2% (-1.7-8.1%)	110 (-92-450)	460 (-260-1600)
United Kingdom	2010	Female	All Ages	83% (80-85%)	17% (15-20%)	3 (2-4.3)	4% (2.2-6%)	5.2% (1.8-9.2%)	15000 (5300-27000)	330000 (190000-490000)
United Kingdom	2010	Female	15 to 19	79% (71-86%)	21% (14-29%)	2.2 (1.2-3.7)	2% (1.3-2.8%)	7.3% (3.9-12%)	28 (15-44)	4000 (2600-5700)
United Kingdom	2010	Female	20 to 24	83% (75-89%)	17% (11-25%)	2.8 (1.6-4.7)	3.1% (2.2-4.2%)	9% (5-14%)	44 (24-70)	7800 (5200-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	2010	Female	25 to 29	85% (77-90%)	15% (10-23%)	2.9 (1.6-4.7)	3% (2.1-4%)	9.3% (5.5-14%)	62 (36-93)	8600 (5800-12000)
United Kingdom	2010	Female	30 to 34	87% (79-92%)	13% (8.3-21%)	3 (1.7-4.8)	3.2% (2.2-4.4%)	11% (7.6-15%)	100 (70-140)	9600 (6800-13000)
United Kingdom	2010	Female	35 to 39	88% (82-93%)	12% (7.4-18%)	3.2 (1.9-5.1)	3.9% (2.8-5.3%)	13% (9.6-17%)	200 (150-270)	14000 (10000-19000)
United Kingdom	2010	Female	40 to 44	89% (83-93%)	11% (6.6-17%)	3.6 (2.1-5.6)	4.8% (3.3-6.5%)	14% (11-18%)	380 (280-480)	21000 (16000-28000)
United Kingdom	2010	Female	45 to 49	90% (83-94%)	10% (6.4-17%)	4.1 (2.3-6.5)	5.6% (3.8-7.6%)	15% (11-19%)	580 (420-740)	28000 (19000-37000)
United Kingdom	2010	Female	50 to 54	90% (84-94%)	10% (6.2-16%)	4.5 (2.6-7)	6.2% (4.2-8.5%)	14% (9.7-18%)	760 (540-980)	31000 (21000-42000)
United Kingdom	2010	Female	55 to 59	89% (84-93%)	11% (6.5-16%)	4.8 (2.9-7.7)	6.2% (4-8.6%)	12% (8.3-16%)	920 (640-1200)	33000 (21000-45000)
United Kingdom	2010	Female	60 to 64	88% (82-93%)	12% (7.5-18%)	4.7 (2.7-7.6)	5.5% (3.2-8%)	9.4% (6-13%)	1200 (750-1700)	36000 (21000-52000)
United Kingdom	2010	Female	65 to 69	86% (78-91%)	14% (9.1-22%)	4.2 (2.5-6.5)	4.6% (2.3-7.2%)	7.3% (4-11%)	1100 (640-1700)	29000 (15000-45000)
United Kingdom	2010	Female	70 to 74	83% (75-88%)	17% (12-25%)	3.8 (2.2-6.1)	3.9% (1.3-7%)	5.6% (2.1-9.7%)	1300 (480-2200)	26000 (8500-47000)
United Kingdom	2010	Female	75 to 79	80% (72-87%)	20% (13-28%)	3.5 (1.9-5.8)	3.4% (0.47-7%)	4.5% (0.78-8.9%)	1500 (260-3000)	24000 (3400-49000)
United Kingdom	2010	Female	80 to 84	78% (67-86%)	22% (14-33%)	3.3 (2-5.4)	3.3% (0.2-7%)	4.1% (0.36-8.8%)	2000 (180-4300)	24000 (1400-51000)
United Kingdom	2010	Female	85 to 89	76% (66-83%)	24% (17-34%)	3.2 (1.9-5.3)	3.2% (-0.064-7.5%)	3.9% (-0.11-8.9%)	2400 (-69-5500)	20000 (-390-45000)
United Kingdom	2010	Female	90 to 94	75% (66-83%)	25% (17-34%)	3.1 (1.7-5.3)	3% (-0.34-7.4%)	3.5% (-0.44-8.7%)	1500 (-190-3900)	8700 (-1000-22000)
United Kingdom	2010	Female	95 plus	75% (62-84%)	25% (16-38%)	3.1 (1.5-5.5)	3.1% (-0.72-8.1%)	3.6% (-0.99-9.6%)	930 (-260-2500)	3400 (-790-8800)
United Kingdom	2010	Male	All Ages	87% (85-90%)	13% (10-15%)	2.9 (1.9-4.2)	5.4% (3.5-7.6%)	5.1% (2.3-8.7%)	14000 (6200-24000)	450000 (290000-640000)
United Kingdom	2010	Male	15 to 19	79% (70-86%)	21% (14-30%)	1.8 (1-3)	3.9% (2.6-5.5%)	8.5% (4.5-13%)	66 (35-100)	8300 (5400-12000)
United Kingdom	2010	Male	20 to 24	86% (78-91%)	14% (8.6-22%)	2.8 (1.5-4.6)	8.1% (5.8-11%)	12% (6.6-19%)	160 (84-250)	21000 (14000-29000)
United Kingdom	2010	Male	25 to 29	89% (82-93%)	11% (7.1-18%)	3 (1.7-4.9)	7.8% (5.8-10%)	13% (7.4-19%)	190 (110-290)	22000 (16000-30000)
United Kingdom	2010	Male	30 to 34	91% (84-95%)	9.4% (5.4-16%)	3.1 (1.8-4.9)	7.5% (5.5-10%)	14% (8.7-19%)	250 (160-360)	23000 (17000-31000)
United Kingdom	2010	Male	35 to 39	92% (88-95%)	8% (4.7-12%)	3.1 (1.8-4.8)	7.8% (5.7-10%)	15% (10-20%)	400 (280-540)	28000 (21000-37000)
United Kingdom	2010	Male	40 to 44	93% (89-96%)	6.9% (3.9-11%)	3.4 (2-5.3)	8.3% (6.1-11%)	15% (11-20%)	650 (470-850)	38000 (28000-49000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	2010	Male	45 to 49	94% (90-97%)	5.9% (3.5-10%)	4 (2.2-6.3)	8.6% (6-11%)	15% (11-20%)	900 (630-1200)	44000 (32000-59000)
United Kingdom	2010	Male	50 to 54	95% (91-97%)	5.3% (3-9%)	4.4 (2.6-7)	8.5% (5.7-12%)	14% (9-19%)	1100 (730-1500)	47000 (31000-64000)
United Kingdom	2010	Male	55 to 59	95% (92-97%)	4.9% (2.8-7.7%)	4.7 (2.9-7.6)	7.9% (4.9-11%)	12% (7.2-16%)	1300 (840-1900)	49000 (30000-69000)
United Kingdom	2010	Male	60 to 64	95% (92-97%)	5% (2.8-8.2%)	4.7 (2.7-7.6)	6.6% (3.7-10%)	9% (4.9-14%)	1700 (920-2600)	52000 (29000-79000)
United Kingdom	2010	Male	65 to 69	94% (91-97%)	5.6% (3.1-8.9%)	4.2 (2.6-6.5)	5.1% (2.5-8.3%)	6.5% (3.2-10%)	1500 (730-2500)	39000 (19000-64000)
United Kingdom	2010	Male	70 to 74	94% (90-96%)	6.4% (4-10%)	3.8 (2.2-6)	3.7% (1.1-7.2%)	4.4% (1.1-8.6%)	1400 (360-2700)	29000 (8500-56000)
United Kingdom	2010	Male	75 to 79	93% (88-96%)	7.3% (4.4-12%)	3.5 (1.9-5.8)	2.9% (0.25-6.3%)	3.2% (0.069-7.2%)	1300 (28-2900)	21000 (1800-48000)
United Kingdom	2010	Male	80 to 84	92% (86-95%)	8.3% (4.9-14%)	3.4 (2-5.4)	2.5% (0.068-6%)	2.7% (-0.16-6.6%)	1300 (-77-3100)	16000 (420-37000)
United Kingdom	2010	Male	85 to 89	91% (86-94%)	9.2% (5.9-14%)	3.3 (1.9-5.3)	2.4% (-0.096-6%)	2.6% (-0.34-6.7%)	1100 (-150-2900)	9700 (-380-24000)
United Kingdom	2010	Male	90 to 94	90% (85-94%)	9.6% (6.1-15%)	3.2 (1.8-5.3)	2.3% (-0.41-6.3%)	2.4% (-0.72-7%)	510 (-150-1500)	3000 (-580-8200)
United Kingdom	2010	Male	95 plus	90% (83-94%)	10% (5.7-17%)	3.1 (1.5-5.4)	2.4% (-0.75-7.1%)	2.5% (-1.2-8%)	180 (-85-570)	700 (-220-2100)
United Kingdom	2016	Female	All Ages	83% (80-86%)	17% (14-20%)	3 (2-4.5)	4% (2.3-6%)	5.3% (2-9.3%)	16000 (6300-28000)	340000 (200000-510000)
United Kingdom	2016	Female	15 to 19	79% (71-86%)	21% (14-29%)	2.2 (1.3-3.6)	1.8% (1.2-2.6%)	7.2% (3.8-11%)	23 (12-37)	350 (2300-5100)
United Kingdom	2016	Female	20 to 24	83% (75-89%)	17% (11-25%)	2.9 (1.6-4.8)	3% (2.1-4%)	9% (4.8-14%)	43 (23-68)	7900 (5300-11000)
United Kingdom	2016	Female	25 to 29	85% (77-90%)	15% (9.9-23%)	2.9 (1.6-5)	2.9% (2.1-4%)	9.3% (5.6-14%)	60 (36-90)	8600 (5700-12000)
United Kingdom	2016	Female	30 to 34	87% (79-92%)	13% (8.4-21%)	3 (1.7-5.1)	3% (2.2-4.2%)	11% (7.8-15%)	100 (73-140)	9900 (7100-13000)
United Kingdom	2016	Female	35 to 39	88% (82-93%)	12% (7.4-18%)	3.2 (1.8-5.2)	3.8% (2.7-5.1%)	14% (10-18%)	170 (120-220)	12000 (8600-15000)
United Kingdom	2016	Female	40 to 44	89% (83-93%)	11% (6.6-17%)	3.6 (2.1-5.6)	4.6% (3.2-6.3%)	15% (11-18%)	350 (260-440)	20000 (14000-26000)
United Kingdom	2016	Female	45 to 49	90% (83-94%)	10% (6.3-17%)	4.1 (2.3-6.4)	5.5% (3.7-7.5%)	15% (11-19%)	600 (440-760)	29000 (20000-39000)
United Kingdom	2016	Female	50 to 54	90% (83-94%)	10% (6.3-17%)	4.5 (2.6-7.1)	6.1% (4.1-8.2%)	14% (10-18%)	810 (590-1000)	34000 (24000-45000)
United Kingdom	2016	Female	55 to 59	89% (84-93%)	11% (6.6-16%)	4.8 (2.9-7.7)	6.2% (4.1-8.4%)	12% (8.6-16%)	940 (660-1200)	34000 (22000-46000)
United Kingdom	2016	Female	60 to 64	88% (81-93%)	12% (7.5-19%)	4.7 (2.7-7.4)	5.5% (3.3-7.9%)	9.6% (6.3-13%)	1100 (730-1500)	34000 (20000-49000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	2016	Female	65 to 69	86% (78-91%)	14% (9-22%)	4.2 (2.6-6.5)	4.6% (2.5-7%)	7.4% (4.3-11%)	1400 (820-2000)	36000 (20000-55000)
United Kingdom	2016	Female	70 to 74	83% (76-88%)	17% (12-24%)	3.8 (2.2-6.3)	4% (1.5-6.9%)	5.9% (2.6-9.7%)	1300 (580-2200)	27000 (10000-47000)
United Kingdom	2016	Female	75 to 79	81% (72-87%)	19% (13-28%)	3.5 (1.9-5.9)	3.5% (0.78-7%)	4.8% (1.2-9.2%)	1600 (390-3000)	25000 (5500-50000)
United Kingdom	2016	Female	80 to 84	78% (68-86%)	22% (14-32%)	3.4 (2-5.5)	3.4% (0.48-7.1%)	4.4% (0.72-8.8%)	2200 (360-4400)	25000 (3500-51000)
United Kingdom	2016	Female	85 to 89	76% (67-84%)	24% (16-33%)	3.2 (1.9-5.3)	3.4% (0.24-7.3%)	4.2% (0.32-9%)	2500 (190-5400)	20000 (1500-44000)
United Kingdom	2016	Female	90 to 94	76% (66-83%)	24% (17-34%)	3.2 (1.8-5.5)	3.1% (-0.061-7.4%)	3.7% (-0.14-8.8%)	2200 (-80-5200)	12000 (-230-28000)
United Kingdom	2016	Female	95 plus	75% (62-84%)	25% (16-38%)	3 (1.5-5.5)	3.3% (-0.5-8.1%)	3.8% (-0.7-9.8%)	1000 (-190-2700)	3700 (-570-9300)
United Kingdom	2016	Male	All Ages	88% (85-90%)	12% (10-15%)	3 (2-4.4)	5.5% (3.6-7.8%)	5.3% (2.4-8.9%)	16000 (7300-27000)	480000 (310000-690000)
United Kingdom	2016	Male	15 to 19	80% (72-86%)	20% (14-28%)	1.9 (1.1-3.1)	3.8% (2.6-5.3%)	8.4% (4.2-14%)	59 (30-95)	7800 (5200-11000)
United Kingdom	2016	Male	20 to 24	86% (79-91%)	14% (8.5-21%)	2.9 (1.6-4.7)	8% (5.8-11%)	12% (6.6-19%)	150 (82-240)	22000 (15000-30000)
United Kingdom	2016	Male	25 to 29	89% (82-93%)	11% (7.1-18%)	3.1 (1.7-5.2)	7.8% (5.7-10%)	13% (7.2-19%)	180 (110-280)	23000 (16000-31000)
United Kingdom	2016	Male	30 to 34	90% (84-94%)	9.5% (5.5-16%)	3.2 (1.8-5.2)	7.4% (5.4-9.8%)	14% (8.8-19%)	250 (160-360)	23000 (17000-31000)
United Kingdom	2016	Male	35 to 39	92% (87-95%)	8.2% (4.8-13%)	3.2 (1.8-5)	7.7% (5.7-10%)	15% (10-20%)	340 (230-450)	25000 (18000-32000)
United Kingdom	2016	Male	40 to 44	93% (88-96%)	7.1% (4-12%)	3.5 (2-5.3)	8.3% (6.1-11%)	16% (11-20%)	600 (430-780)	35000 (27000-45000)
United Kingdom	2016	Male	45 to 49	94% (90-96%)	6.1% (3.5-10%)	4 (2.2-6.3)	8.7% (6-12%)	15% (11-21%)	930 (650-1200)	46000 (33000-61000)
United Kingdom	2016	Male	50 to 54	95% (91-97%)	5.4% (3.1-9%)	4.5 (2.6-7.1)	8.6% (5.8-12%)	14% (9.6-19%)	1200 (820-1600)	51000 (35000-69000)
United Kingdom	2016	Male	55 to 59	95% (92-97%)	5.1% (2.9-8.2%)	4.9 (2.9-7.7)	8.1% (5.3-11%)	12% (7.9-17%)	1400 (910-2000)	51000 (33000-72000)
United Kingdom	2016	Male	60 to 64	95% (91-97%)	5.1% (2.8-8.5%)	4.8 (2.8-7.7)	6.9% (4-10%)	9.6% (5.5-14%)	1700 (950-2500)	52000 (30000-77000)
United Kingdom	2016	Male	65 to 69	94% (91-97%)	5.6% (3.1-9.1%)	4.3 (2.7-6.7)	5.4% (2.9-8.4%)	7% (3.7-11%)	2000 (1000-3100)	52000 (27000-80000)
United Kingdom	2016	Male	70 to 74	94% (90-96%)	6.4% (4-10%)	3.9 (2.3-6.4)	4.1% (1.6-7.3%)	4.9% (1.8-8.9%)	1600 (590-2900)	33000 (13000-61000)
United Kingdom	2016	Male	75 to 79	93% (88-96%)	7.5% (4.4-12%)	3.6 (2-5.9)	3.2% (0.6-6.6%)	3.6% (0.52-7.6%)	1500 (220-3200)	25000 (4600-54000)
United Kingdom	2016	Male	80 to 84	91% (86-95%)	8.6% (5.1-14%)	3.4 (2-5.5)	2.8% (0.25-6.2%)	3.1% (0.07-7%)	1600 (37-3700)	20000 (1800-44000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Kingdom	2016	Male	85 to 89	90% (85-94%)	9.6% (6-15%)	3.3 (1.9-5.3)	2.7% (0.18-6.3%)	2.9% (-0.044-7.1%)	1500 (-22-3600)	12000 (810-29000)
United Kingdom	2016	Male	90 to 94	90% (84-94%)	10% (6.3-16%)	3.2 (1.8-5.4)	2.5% (-0.17-6.6%)	2.7% (-0.49-7.4%)	910 (-160-2500)	5300 (-340-14000)
United Kingdom	2016	Male	95 plus	90% (83-94%)	10% (5.6-17%)	3.1 (1.6-5.4)	2.6% (-0.54-7.3%)	2.7% (-0.97-8.2%)	250 (-87-730)	950 (-200-2700)
Argentina	1990	Female	All Ages	90% (87-92%)	10% (7.8-13%)	3.5 (3-4.1)	5.1% (3.3-6.9%)	6.6% (2.5-11%)	7500 (2800-13000)	230000 (150000-320000)
Argentina	1990	Female	15 to 19	91% (85-96%)	8.6% (4.3-15%)	4 (2.4-5.8)	4.8% (3-6.8%)	13% (7.4-19%)	89 (51-130)	8300 (5200-12000)
Argentina	1990	Female	20 to 24	94% (90-97%)	5.7% (2.9-10%)	5.7 (3.7-8.3)	7% (4.9-9.2%)	16% (10-22%)	130 (81-170)	12000 (8500-16000)
Argentina	1990	Female	25 to 29	95% (91-97%)	5.2% (2.6-9.2%)	6 (4-8.6)	6.9% (5.1-8.9%)	16% (12-21%)	150 (110-190)	13000 (9300-16000)
Argentina	1990	Female	30 to 34	94% (90-97%)	5.9% (3.3-10%)	5.9 (4-8.3)	6.9% (5-9.1%)	16% (11-20%)	190 (130-250)	13000 (9600-17000)
Argentina	1990	Female	35 to 39	92% (87-96%)	7.6% (3.8-13%)	5.4 (3.6-7.8)	7.3% (5-9.8%)	16% (11-21%)	260 (180-350)	16000 (11000-21000)
Argentina	1990	Female	40 to 44	92% (87-96%)	8.1% (4.5-13%)	5.2 (3.3-7.7)	8% (5.2-11%)	16% (10-22%)	340 (220-470)	18000 (11000-25000)
Argentina	1990	Female	45 to 49	93% (87-96%)	7.5% (3.9-13%)	5.2 (3.2-7.7)	8.7% (5.3-12%)	16% (10-22%)	430 (270-600)	20000 (12000-27000)
Argentina	1990	Female	50 to 54	93% (88-96%)	7.2% (3.8-12%)	4.9 (3-7.3)	8.9% (5.1-13%)	15% (8.9-21%)	540 (320-760)	21000 (13000-31000)
Argentina	1990	Female	55 to 59	93% (88-96%)	7.4% (4.1-12%)	4.8 (3-7)	8.9% (4.7-13%)	14% (7.7-20%)	700 (390-1000)	24000 (13000-35000)
Argentina	1990	Female	60 to 64	92% (85-96%)	8.4% (4.5-15%)	4.4 (2.7-6.8)	7.7% (2.7-13%)	11% (4.1-18%)	790 (300-1300)	23000 (8200-38000)
Argentina	1990	Female	65 to 69	90% (82-95%)	10% (4.7-18%)	3.8 (2.5-5.5)	6.3% (2.5-11%)	8.5% (3.4-15%)	800 (310-1400)	19000 (7300-33000)
Argentina	1990	Female	70 to 74	88% (80-94%)	12% (6-20%)	3.4 (2-5.2)	5% (0.33-10%)	6.5% (0.5-13%)	780 (60-1600)	15000 (1000-31000)
Argentina	1990	Female	75 to 79	87% (78-94%)	13% (6.3-22%)	3.3 (1.8-5.1)	4% (-1.5-10%)	4.8% (-1.9-12%)	790 (-310-2000)	12000 (-4200-31000)
Argentina	1990	Female	80 to 84	86% (76-93%)	14% (7-24%)	3.2 (2-4.8)	3.7% (-1.2-9.3%)	4.2% (-1.8-11%)	700 (-290-1800)	7900 (-2700-20000)
Argentina	1990	Female	85 to 89	85% (75-92%)	15% (8.1-25%)	3.1 (1.9-4.8)	3.4% (-1.6-9.4%)	3.8% (-2.1-11%)	520 (-300-1500)	4100 (-1800-11000)
Argentina	1990	Female	90 to 94	85% (75-92%)	15% (8-25%)	3.1 (1.8-5)	3.2% (-2-10%)	3.5% (-2.6-12%)	250 (-190-830)	1400 (-800-4300)
Argentina	1990	Female	95 plus	85% (72-93%)	15% (6.9-28%)	3.2 (1.6-5.5)	3.3% (-2.8-11%)	3.5% (-3.6-13%)	81 (-83-300)	300 (-250-1000)
Argentina	1990	Male	All Ages	94% (92-96%)	5.7% (3.9-7.9%)	3.5 (3-4.1)	7.4% (5.3-9.5%)	7.3% (3.9-11%)	10000 (5700-16000)	410000 (300000-530000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	1990	Male	15 to 19	94% (89-97%)	5.7% (2.6-11%)	3.7 (2.3-5.5)	8.9% (6-12%)	15% (9-21%)	210 (130-300)	18000 (12000-25000)
Argentina	1990	Male	20 to 24	96% (92-98%)	4.3% (2.1-8.1%)	5.6 (3.6-8.1)	15% (11-20%)	22% (14-30%)	370 (240-500)	33000 (23000-43000)
Argentina	1990	Male	25 to 29	96% (92-98%)	4.3% (2-7.9%)	6.3 (4.2-8.7)	15% (12-19%)	23% (16-30%)	370 (250-500)	31000 (23000-39000)
Argentina	1990	Male	30 to 34	96% (92-98%)	4.3% (2-8.3%)	6.2 (4.3-8.8)	14% (11-18%)	21% (15-28%)	410 (290-540)	29000 (21000-37000)
Argentina	1990	Male	35 to 39	96% (92-98%)	4.1% (1.8-7.5%)	5.6 (3.7-8)	13% (9.5-18%)	20% (13-26%)	500 (340-670)	31000 (22000-41000)
Argentina	1990	Male	40 to 44	96% (93-98%)	3.7% (1.6-7.2%)	5.2 (3.3-7.6)	13% (8.8-18%)	18% (11-25%)	670 (410-930)	36000 (23000-48000)
Argentina	1990	Male	45 to 49	97% (94-99%)	3.1% (1.3-6.3%)	5.1 (3.2-7.5)	13% (8.2-18%)	17% (10-24%)	880 (520-1200)	40000 (25000-56000)
Argentina	1990	Male	50 to 54	97% (94-99%)	2.8% (1.2-5.9%)	5 (3.1-7.5)	12% (7.1-18%)	15% (8.3-22%)	1100 (620-1700)	45000 (25000-65000)
Argentina	1990	Male	55 to 59	97% (94-99%)	3% (1.2-6%)	4.8 (3-7)	11% (5.6-17%)	13% (6.4-20%)	1400 (680-2100)	47000 (24000-71000)
Argentina	1990	Male	60 to 64	97% (93-99%)	3.3% (1.3-7%)	4.5 (2.6-6.9)	8.7% (3.1-15%)	10% (3.4-17%)	1400 (490-2500)	41000 (15000-70000)
Argentina	1990	Male	65 to 69	96% (92-99%)	3.6% (1.4-7.8%)	4.1 (2.8-5.9)	6.4% (1.9-12%)	7.2% (2-13%)	1200 (330-2200)	28000 (8500-52000)
Argentina	1990	Male	70 to 74	96% (92-98%)	3.9% (1.6-8%)	3.7 (2.2-5.4)	4.5% (-0.027-9.9%)	5% (-0.33-11%)	850 (-56-1900)	17000 (-99-37000)
Argentina	1990	Male	75 to 79	96% (92-98%)	4.2% (1.8-8.2%)	3.4 (1.9-5.1)	3.1% (-1.3-8.7%)	3.3% (-1.8-9.8%)	590 (-320-1700)	9200 (-3800-25000)
Argentina	1990	Male	80 to 84	95% (90-98%)	4.8% (2.1-10%)	3.1 (1.9-4.5)	2.2% (-1.5-6.8%)	2.2% (-2-7.4%)	320 (-290-1100)	3900 (-2600-12000)
Argentina	1990	Male	85 to 89	94% (89-97%)	5.7% (2.5-11%)	2.9 (1.8-4.4)	1.6% (-1.8-5.9%)	1.5% (-2.5-6.4%)	140 (-230-600)	1300 (-1400-4700)
Argentina	1990	Male	90 to 94	94% (89-98%)	6% (2.4-11%)	2.8 (1.5-4.4)	1.2% (-2.3-6.1%)	0.98% (-2.9-6.6%)	38 (-110-260)	270 (-520-1400)
Argentina	1990	Male	95 plus	94% (87-98%)	6% (2-13%)	2.8 (1.3-4.8)	1.3% (-2.6-7.2%)	1% (-3.5-7.9%)	10 (-35-79)	50 (-100-280)
Argentina	1995	Female	All Ages	90% (87-92%)	9.9% (7.6-13%)	2.8 (2.4-3.2)	3.7% (2.3-5.4%)	4.1% (0.82-8%)	4900 (980-9300)	170000 (100000-250000)
Argentina	1995	Female	15 to 19	91% (85-96%)	8.5% (4.2-15%)	3.1 (2-4.6)	4.1% (2.6-6%)	11% (5.9-17%)	84 (46-130)	8400 (5200-12000)
Argentina	1995	Female	20 to 24	94% (90-97%)	5.6% (2.9-10%)	4.5 (2.8-6.6)	6.1% (4.2-8.2%)	13% (7.8-18%)	110 (67-160)	12000 (8500-16000)
Argentina	1995	Female	25 to 29	95% (91-97%)	5.2% (2.7-9.3%)	4.8 (3.2-6.7)	5.9% (4.2-7.7%)	13% (8.4-17%)	120 (81-170)	11000 (7900-15000)
Argentina	1995	Female	30 to 34	94% (90-97%)	5.8% (3.2-10%)	4.7 (3.1-6.5)	5.5% (3.8-7.5%)	13% (8.5-17%)	150 (98-200)	11000 (7700-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	1995	Female	35 to 39	92% (87-96%)	7.6% (3.7-13%)	4.3 (2.8-6.1)	5.6% (3.6-7.7%)	12% (8.1-17%)	190 (120-260)	12000 (8000-17000)
Argentina	1995	Female	40 to 44	92% (87-96%)	8.1% (4.3-13%)	4.1 (2.6-5.9)	6.2% (3.7-8.9%)	13% (7.8-18%)	280 (170-390)	15000 (9100-21000)
Argentina	1995	Female	45 to 49	93% (87-96%)	7.5% (3.9-13%)	4.1 (2.6-6)	6.8% (3.9-10%)	13% (7.6-18%)	370 (230-530)	17000 (10000-25000)
Argentina	1995	Female	50 to 54	93% (88-96%)	7.2% (4-12%)	3.9 (2.4-5.7)	6.9% (3.6-11%)	12% (6.4-17%)	440 (240-650)	17000 (9100-26000)
Argentina	1995	Female	55 to 59	93% (88-96%)	7.3% (4-12%)	3.8 (2.3-5.4)	6.6% (3-10%)	10% (5.2-16%)	500 (250-770)	17000 (8000-27000)
Argentina	1995	Female	60 to 64	92% (85-96%)	8.3% (4.2-15%)	3.5 (2-5.2)	5.3% (1.2-9.7%)	7.8% (2.1-14%)	530 (140-960)	15000 (3400-29000)
Argentina	1995	Female	65 to 69	90% (82-95%)	10% (4.8-18%)	3 (1.9-4.2)	4% (0.74-7.9%)	5.6% (1.3-11%)	540 (130-1000)	13000 (2500-26000)
Argentina	1995	Female	70 to 74	89% (80-94%)	11% (6.1-20%)	2.7 (1.6-4)	2.9% (-0.82-7.2%)	3.8% (-0.97-9.3%)	500 (-120-1200)	9500 (-2600-24000)
Argentina	1995	Female	75 to 79	87% (78-94%)	13% (6.4-22%)	2.6 (1.4-4)	1.9% (-2.1-6.6%)	2.3% (-2.7-8.3%)	370 (-430-1300)	5800 (-6300-19000)
Argentina	1995	Female	80 to 84	86% (76-93%)	14% (6.8-24%)	2.5 (1.6-3.7)	1.6% (-2.1-5.9%)	1.8% (-2.8-7%)	310 (-490-1200)	3700 (-5000-14000)
Argentina	1995	Female	85 to 89	85% (76-92%)	15% (8-24%)	2.5 (1.5-3.8)	1.4% (-2.2-5.8%)	1.5% (-2.8-6.8%)	240 (-460-1100)	2100 (-3300-8500)
Argentina	1995	Female	90 to 94	85% (76-92%)	15% (7.8-24%)	2.5 (1.4-4)	1.3% (-2.6-6.7%)	1.2% (-3.4-7.6%)	110 (-310-700)	710 (-1500-3700)
Argentina	1995	Female	95 plus	85% (73-93%)	15% (6.5-27%)	2.5 (1.3-4.3)	1.3% (-3.1-7.9%)	1.2% (-4.1-9.1%)	38 (-130-280)	160 (-390-980)
Argentina	1995	Male	All Ages	94% (92-96%)	5.7% (3.8-8%)	3.3 (2.9-3.8)	7.5% (5.6-9.6%)	6.9% (3.9-10%)	10000 (5600-16000)	430000 (310000-540000)
Argentina	1995	Male	15 to 19	94% (89-97%)	5.8% (2.6-11%)	3.5 (2.1-5.1)	9.6% (6.7-13%)	15% (9.1-21%)	260 (160-380)	25000 (17000-33000)
Argentina	1995	Male	20 to 24	96% (92-98%)	4.3% (2.1-7.8%)	5.3 (3.4-7.6)	16% (12-21%)	21% (14-29%)	460 (290-630)	43000 (31000-56000)
Argentina	1995	Male	25 to 29	96% (92-98%)	4.3% (1.9-8.1%)	5.8 (4-8.2)	15% (12-19%)	20% (14-27%)	440 (300-590)	37000 (28000-47000)
Argentina	1995	Male	30 to 34	96% (91-98%)	4.3% (2-8.6%)	5.8 (4-8.1)	13% (10-17%)	19% (13-25%)	410 (290-550)	30000 (22000-39000)
Argentina	1995	Male	35 to 39	96% (92-98%)	4.1% (1.8-7.8%)	5.2 (3.4-7.3)	13% (9.2-17%)	18% (12-24%)	490 (320-660)	31000 (22000-41000)
Argentina	1995	Male	40 to 44	96% (93-98%)	3.7% (1.5-7.5%)	4.9 (3.1-7)	13% (8.7-17%)	17% (11-24%)	640 (400-890)	35000 (24000-47000)
Argentina	1995	Male	45 to 49	97% (93-99%)	3.1% (1.3-6.7%)	4.8 (3-7)	13% (8.1-18%)	16% (9.9-23%)	880 (540-1300)	41000 (27000-57000)
Argentina	1995	Male	50 to 54	97% (94-99%)	2.9% (1.1-6.1%)	4.6 (2.9-7)	12% (6.9-17%)	14% (8.3-22%)	1100 (590-1600)	43000 (25000-63000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	1995	Male	55 to 59	97% (94-99%)	3% (1.2-6%)	4.5 (2.8-6.5)	10% (5.6-15%)	12% (6.5-19%)	1200 (640-1900)	42000 (23000-63000)
Argentina	1995	Male	60 to 64	97% (93-99%)	3.3% (1.3-6.7%)	4.2 (2.5-6.3)	8.1% (3.3-14%)	9.4% (3.7-16%)	1300 (510-2300)	37000 (15000-65000)
Argentina	1995	Male	65 to 69	96% (92-99%)	3.5% (1.3-7.5%)	3.8 (2.5-5.5)	5.9% (1.7-11%)	6.6% (1.7-12%)	1100 (300-2100)	28000 (8200-51000)
Argentina	1995	Male	70 to 74	96% (92-98%)	3.8% (1.7-7.6%)	3.5 (2.1-5)	4.2% (0.23-9%)	4.7% (-0.0042-10%)	880 (-0.81-2000)	17000 (970-37000)
Argentina	1995	Male	75 to 79	96% (92-98%)	4.1% (1.8-7.9%)	3.2 (1.8-4.7)	3% (-0.88-7.9%)	3.1% (-1.3-8.7%)	560 (-230-1500)	8800 (-2600-23000)
Argentina	1995	Male	80 to 84	95% (91-98%)	4.8% (2.1-9.4%)	2.9 (1.8-4.2)	2.1% (-1.3-6.2%)	2.1% (-1.8-6.7%)	320 (-280-1000)	3900 (-2400-11000)
Argentina	1995	Male	85 to 89	94% (89-97%)	5.7% (2.6-11%)	2.7 (1.6-4)	1.4% (-1.4-5.1%)	1.3% (-2.1-5.5%)	140 (-220-580)	1300 (-1300-4600)
Argentina	1995	Male	90 to 94	94% (89-97%)	6% (2.7-11%)	2.6 (1.5-4)	0.99% (-2-5.2%)	0.79% (-2.7-5.6%)	37 (-130-270)	270 (-550-1400)
Argentina	1995	Male	95 plus	94% (87-98%)	6% (2.2-13%)	2.5 (1.2-4.3)	1.1% (-2.4-6%)	0.83% (-3.2-6.5%)	10 (-40-83)	52 (-120-300)
Argentina	2000	Female	All Ages	90% (88-92%)	9.6% (7.6-12%)	2.6 (2.2-3)	3.2% (1.9-4.7%)	3.2% (0.38-6.5%)	4200 (490-8500)	150000 (88000-220000)
Argentina	2000	Female	15 to 19	93% (88-96%)	7.2% (3.7-12%)	2.9 (1.8-4.3)	4% (2.4-5.7%)	10% (5.6-16%)	78 (42-120)	7800 (4800-11000)
Argentina	2000	Female	20 to 24	95% (91-97%)	5.1% (2.7-9.3%)	4.1 (2.7-5.9)	5.5% (3.9-7.3%)	12% (7.2-17%)	110 (63-150)	12000 (8600-16000)
Argentina	2000	Female	25 to 29	95% (91-97%)	5% (2.6-8.9%)	4.3 (2.9-6)	5.1% (3.6-6.7%)	11% (7.5-16%)	100 (68-150)	10000 (7300-14000)
Argentina	2000	Female	30 to 34	94% (90-97%)	6% (3.3-10%)	4.2 (2.9-5.9)	4.6% (3.2-6.3%)	11% (7.2-15%)	120 (77-170)	9300 (6400-13000)
Argentina	2000	Female	35 to 39	92% (87-96%)	7.9% (3.8-13%)	3.8 (2.4-5.5)	4.7% (3-6.5%)	11% (6.7-15%)	150 (96-220)	10000 (6600-14000)
Argentina	2000	Female	40 to 44	92% (87-96%)	8.1% (4.3-13%)	3.7 (2.4-5.4)	5.3% (3.1-7.6%)	11% (6.5-16%)	230 (130-340)	13000 (7400-18000)
Argentina	2000	Female	45 to 49	93% (88-96%)	7.2% (4-12%)	3.7 (2.3-5.5)	5.8% (3.2-8.9%)	11% (6.4-17%)	340 (190-500)	15000 (8500-23000)
Argentina	2000	Female	50 to 54	93% (88-96%)	7.4% (4.2-12%)	3.5 (2.1-5.1)	6% (3-9.4%)	10% (5.5-16%)	420 (220-650)	17000 (8300-26000)
Argentina	2000	Female	55 to 59	93% (88-96%)	7.4% (4.1-12%)	3.4 (2.1-4.9)	5.7% (2.4-9.2%)	9.2% (4.5-14%)	470 (230-740)	16000 (6900-26000)
Argentina	2000	Female	60 to 64	92% (86-95%)	8.5% (4.5-14%)	3.1 (1.8-4.7)	4.5% (1.1-8.5%)	6.9% (2-13%)	460 (140-840)	13000 (3100-25000)
Argentina	2000	Female	65 to 69	89% (80-95%)	11% (5.1-20%)	2.6 (1.7-3.7)	3.3% (0.51-6.6%)	4.7% (0.92-9.2%)	440 (86-860)	10000 (1600-21000)
Argentina	2000	Female	70 to 74	88% (79-94%)	12% (6.2-21%)	2.4 (1.4-3.6)	2.3% (-0.87-6%)	3.1% (-0.93-7.7%)	430 (-120-1100)	8000 (-3200-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	2000	Female	75 to 79	87% (78-94%)	13% (6.3-22%)	2.3 (1.3-3.7)	1.4% (-2-5.6%)	1.7% (-2.4-6.9%)	290 (-430-1200)	4500 (-6600-19000)
Argentina	2000	Female	80 to 84	86% (76-93%)	14% (7-24%)	2.3 (1.4-3.3)	1.1% (-2.1-4.7%)	1.2% (-2.8-5.6%)	240 (-540-1100)	2800 (-5600-12000)
Argentina	2000	Female	85 to 89	85% (75-92%)	15% (8.3-25%)	2.2 (1.3-3.4)	0.9% (-2.2-4.8%)	0.89% (-2.8-5.5%)	180 (-540-1100)	1600 (-3900-8300)
Argentina	2000	Female	90 to 94	85% (75-92%)	15% (7.9-25%)	2.2 (1.2-3.5)	0.71% (-2.4-5.1%)	0.61% (-3.2-5.8%)	75 (-390-720)	530 (-1800-3900)
Argentina	2000	Female	95 plus	85% (73-93%)	15% (6.7-27%)	2.2 (1.1-3.9)	0.73% (-3.1-6.1%)	0.56% (-3.9-7%)	25 (-180-310)	130 (-550-1100)
Argentina	2000	Male	All Ages	95% (93-97%)	4.9% (3.4-6.7%)	2.7 (2.4-3.1)	6.1% (4.5-7.9%)	4.8% (2.3-7.7%)	7400 (3500-12000)	350000 (250000-460000)
Argentina	2000	Male	15 to 19	95% (91-98%)	4.7% (2.3-8.7%)	2.8 (1.7-4.2)	9% (6-12%)	13% (7.6-20%)	240 (140-360)	23000 (15000-31000)
Argentina	2000	Male	20 to 24	96% (94-98%)	3.6% (1.8-6.3%)	4.2 (2.7-6.2)	15% (11-19%)	19% (12-26%)	480 (300-670)	46000 (33000-60000)
Argentina	2000	Male	25 to 29	96% (94-98%)	3.6% (1.7-6.4%)	4.7 (3.2-6.5)	14% (11-18%)	18% (12-25%)	420 (270-580)	37000 (27000-48000)
Argentina	2000	Male	30 to 34	96% (94-98%)	3.5% (1.7-6.3%)	4.7 (3.2-6.6)	11% (8.6-15%)	16% (10-22%)	360 (240-500)	27000 (20000-36000)
Argentina	2000	Male	35 to 39	96% (94-98%)	3.5% (1.6-6.5%)	4.2 (2.7-6)	10% (7.5-14%)	14% (9.3-20%)	370 (240-530)	25000 (18000-34000)
Argentina	2000	Male	40 to 44	97% (94-99%)	3.1% (1.2-6.2%)	3.9 (2.4-5.7)	10% (6.9-14%)	14% (8.5-19%)	490 (300-700)	28000 (19000-39000)
Argentina	2000	Male	45 to 49	97% (95-99%)	2.6% (1.1-5.2%)	3.8 (2.5-5.5)	9.9% (6.4-14%)	13% (7.6-19%)	670 (410-1000)	33000 (21000-47000)
Argentina	2000	Male	50 to 54	98% (95-99%)	2.4% (0.98-4.8%)	3.8 (2.4-5.5)	9% (5.2-14%)	11% (6.2-17%)	850 (470-1300)	35000 (20000-53000)
Argentina	2000	Male	55 to 59	98% (95-99%)	2.5% (1.1-4.9%)	3.7 (2.3-5.4)	7.8% (4.2-12%)	9.2% (4.7-15%)	920 (460-1500)	32000 (17000-52000)
Argentina	2000	Male	60 to 64	97% (94-99%)	2.8% (1.2-5.7%)	3.4 (2-5.2)	5.9% (2.2-11%)	6.8% (2.4-13%)	880 (310-1700)	26000 (9800-49000)
Argentina	2000	Male	65 to 69	97% (93-99%)	3.4% (1.3-7.1%)	3 (2-4.3)	3.9% (0.63-7.6%)	4.3% (0.45-8.9%)	720 (75-1500)	18000 (2800-35000)
Argentina	2000	Male	70 to 74	96% (92-98%)	3.9% (1.7-8%)	2.7 (1.6-4)	2.5% (-0.53-6.3%)	2.7% (-0.84-7.2%)	550 (-170-1500)	11000 (-2400-29000)
Argentina	2000	Male	75 to 79	96% (92-98%)	4.3% (1.7-8.5%)	2.4 (1.4-3.8)	1.5% (-1.3-5.2%)	1.5% (-1.8-5.6%)	310 (-360-1200)	5100 (-4500-17000)
Argentina	2000	Male	80 to 84	95% (90-98%)	5% (2.2-10%)	2.2 (1.4-3.3)	0.86% (-1.6-3.9%)	0.75% (-2.2-4.2%)	130 (-380-730)	1800 (-3500-8100)
Argentina	2000	Male	85 to 89	94% (89-97%)	6% (2.6-11%)	2.1 (1.2-3.1)	0.5% (-1.7-3.1%)	0.32% (-2.2-3.2%)	39 (-270-400)	520 (-1800-3200)
Argentina	2000	Male	90 to 94	94% (88-97%)	6.4% (2.6-12%)	2 (1.1-3.2)	0.21% (-2.1-3%)	-0.037% (-2.7-3.2%)	-2.1 (-160-180)	69 (-720-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	2000	Male	95 plus	94% (87-98%)	6.3% (2.3-13%)	2 (0.92-3.5)	0.28% (-2.3-3.8%)	-0.021% (-3-4%)	-0.32 (-47-63)	17 (-140-240)
Argentina	2005	Female	All Ages	90% (87-92%)	9.9% (7.7-13%)	2.3 (2-2.7)	2.7% (1.4-4%)	2.4% (-0.25-5.2%)	3400 (-350-7500)	130000 (68000-190000)
Argentina	2005	Female	15 to 19	93% (88-96%)	7.4% (3.8-12%)	2.6 (1.7-3.9)	3.5% (2.2-5.1%)	9.5% (5-14%)	67 (35-100)	7100 (4400-10000)
Argentina	2005	Female	20 to 24	95% (90-97%)	5.3% (2.8-9.5%)	3.7 (2.4-5.3)	5.1% (3.6-6.7%)	11% (6.4-16%)	87 (50-130)	11000 (7600-15000)
Argentina	2005	Female	25 to 29	95% (91-97%)	5.2% (2.6-8.9%)	3.9 (2.7-5.4)	4.5% (3.2-5.9%)	10% (6.6-15%)	96 (60-140)	10000 (7300-14000)
Argentina	2005	Female	30 to 34	94% (90-97%)	6.2% (3.4-10%)	3.8 (2.5-5.3)	4% (2.7-5.4%)	9.5% (6.1-13%)	100 (64-140)	8600 (5800-12000)
Argentina	2005	Female	35 to 39	92% (87-96%)	8.2% (4-13%)	3.4 (2.2-4.9)	4% (2.5-5.6%)	9.3% (5.8-13%)	130 (78-180)	8700 (5600-12000)
Argentina	2005	Female	40 to 44	92% (87-95%)	8.3% (4.6-13%)	3.4 (2.1-5)	4.5% (2.6-6.6%)	9.9% (5.6-14%)	200 (110-290)	11000 (6200-16000)
Argentina	2005	Female	45 to 49	93% (88-96%)	7.4% (4.2-12%)	3.4 (2.2-5.1)	4.9% (2.6-7.7%)	9.7% (5.2-15%)	290 (150-440)	13000 (7200-20000)
Argentina	2005	Female	50 to 54	92% (87-96%)	7.7% (4.3-13%)	3.1 (1.9-4.6)	5% (2.3-8%)	8.8% (4.5-13%)	380 (190-580)	15000 (7100-24000)
Argentina	2005	Female	55 to 59	92% (87-96%)	7.7% (4.4-13%)	3 (1.9-4.4)	4.7% (1.9-7.7%)	7.7% (3.6-12%)	430 (200-680)	15000 (6200-24000)
Argentina	2005	Female	60 to 64	91% (85-95%)	8.8% (4.8-15%)	2.8 (1.6-4.2)	3.7% (0.71-7.2%)	5.8% (1.5-11%)	410 (110-770)	12000 (2200-22000)
Argentina	2005	Female	65 to 69	89% (80-95%)	11% (5.3-20%)	2.3 (1.5-3.3)	2.6% (0.14-5.3%)	3.8% (0.69-7.6%)	360 (64-710)	8200 (470-17000)
Argentina	2005	Female	70 to 74	87% (78-93%)	13% (6.6-22%)	2.1 (1.2-3.2)	1.7% (-1.1-4.9%)	2.4% (-1.1-6.4%)	320 (-150-850)	5800 (-3600-17000)
Argentina	2005	Female	75 to 79	87% (77-94%)	13% (6.4-23%)	2.1 (1.1-3.3)	0.86% (-2.1-4.6%)	1.1% (-2.5-5.6%)	210 (-470-1100)	3100 (-7200-16000)
Argentina	2005	Female	80 to 84	86% (74-93%)	14% (7.1-26%)	2 (1.2-3)	0.59% (-2.2-3.8%)	0.67% (-2.8-4.4%)	160 (-630-1000)	1800 (-6800-11000)
Argentina	2005	Female	85 to 89	85% (74-92%)	15% (8.4-26%)	1.9 (1.2-3)	0.47% (-2.1-3.6%)	0.44% (-2.6-4.2%)	100 (-610-940)	1000 (-4500-7600)
Argentina	2005	Female	90 to 94	85% (75-92%)	15% (8.1-25%)	2 (1.1-3.2)	0.34% (-2.5-3.9%)	0.22% (-3-4.3%)	34 (-460-650)	320 (-2300-3700)
Argentina	2005	Female	95 plus	85% (72-93%)	15% (6.7-28%)	2 (0.99-3.5)	0.37% (-2.9-5.2%)	0.18% (-3.6-5.8%)	11 (-220-350)	89 (-690-1200)
Argentina	2005	Male	All Ages	95% (93-97%)	4.8% (3.3-6.6%)	2.6 (2.2-2.9)	5.7% (4.2-7.4%)	4.2% (1.9-6.7%)	6700 (3100-11000)	320000 (240000-430000)
Argentina	2005	Male	15 to 19	95% (92-98%)	4.7% (2.3-8.4%)	2.6 (1.6-3.9)	8.4% (5.4-12%)	13% (6.9-19%)	220 (120-320)	21000 (14000-29000)
Argentina	2005	Male	20 to 24	96% (94-98%)	3.6% (1.9-6.4%)	3.9 (2.6-5.8)	15% (11-19%)	18% (11-26%)	410 (250-580)	42000 (30000-54000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	2005	Male	25 to 29	96% (94-98%)	3.5% (1.7-6.1%)	4.4 (3-6.1)	14% (10-17%)	18% (11-25%)	410 (260-560)	38000 (28000-50000)
Argentina	2005	Male	30 to 34	96% (94-98%)	3.5% (1.7-6.3%)	4.3 (2.9-6.1)	11% (8.3-15%)	16% (10-22%)	340 (220-470)	27000 (20000-36000)
Argentina	2005	Male	35 to 39	97% (94-98%)	3.5% (1.7-6.4%)	3.9 (2.5-5.6)	9.6% (6.9-13%)	13% (8.6-19%)	330 (210-460)	23000 (17000-32000)
Argentina	2005	Male	40 to 44	97% (94-99%)	3.1% (1.4-6.1%)	3.7 (2.2-5.5)	9.3% (6.5-13%)	13% (8.1-18%)	410 (270-600)	25000 (17000-34000)
Argentina	2005	Male	45 to 49	97% (95-99%)	2.6% (1.2-4.9%)	3.6 (2.3-5.4)	8.9% (5.9-13%)	11% (7-17%)	570 (350-870)	29000 (19000-41000)
Argentina	2005	Male	50 to 54	98% (95-99%)	2.4% (1.1-4.6%)	3.5 (2.1-5.2)	8.1% (4.9-13%)	10% (5.6-16%)	760 (420-1200)	32000 (19000-50000)
Argentina	2005	Male	55 to 59	97% (95-99%)	2.5% (1.1-5%)	3.4 (2.2-5)	6.9% (3.6-11%)	8.2% (4.1-14%)	860 (430-1400)	30000 (16000-49000)
Argentina	2005	Male	60 to 64	97% (94-99%)	2.8% (1.1-5.8%)	3.2 (1.9-4.7)	5.3% (2-9.7%)	6.1% (2.2-11%)	810 (290-1500)	24000 (9300-44000)
Argentina	2005	Male	65 to 69	97% (93-99%)	3.4% (1.4-7%)	2.8 (1.9-4)	3.4% (0.67-6.9%)	3.9% (0.53-8%)	620 (86-1300)	15000 (3000-31000)
Argentina	2005	Male	70 to 74	96% (92-98%)	4% (1.7-7.9%)	2.5 (1.5-3.7)	2.2% (-0.41-5.5%)	2.4% (-0.72-6.2%)	470 (-140-1200)	9700 (-1800-24000)
Argentina	2005	Male	75 to 79	96% (91-98%)	4.4% (1.8-8.7%)	2.2 (1.2-3.5)	1.3% (-1.2-4.5%)	1.3% (-1.6-4.9%)	290 (-360-1100)	4900 (-4600-16000)
Argentina	2005	Male	80 to 84	95% (90-98%)	5.2% (2.3-10%)	2.1 (1.3-3)	0.73% (-1.5-3.1%)	0.63% (-2-3.4%)	120 (-390-660)	1800 (-3700-7800)
Argentina	2005	Male	85 to 89	94% (88-97%)	6.2% (2.8-12%)	1.9 (1.1-2.8)	0.43% (-1.5-2.6%)	0.28% (-2-2.8%)	39 (-280-390)	510 (-1800-3100)
Argentina	2005	Male	90 to 94	93% (88-97%)	6.6% (2.9-12%)	1.8 (1-2.9)	0.13% (-1.9-2.5%)	-0.094% (-2.5-2.6%)	-6.1 (-160-170)	49 (-710-960)
Argentina	2005	Male	95 plus	94% (86-98%)	6.4% (2.3-14%)	1.8 (0.82-3.1)	0.15% (-2.1-3.3%)	-0.14% (-2.7-3.4%)	-2.6 (-50-64)	11 (-150-240)
Argentina	2010	Female	All Ages	90% (87-92%)	10% (7.8-13%)	2.1 (1.8-2.5)	2.2% (1-3.5%)	1.7% (-0.7-4.4%)	2600 (-1100-6800)	110000 (52000-180000)
Argentina	2010	Female	15 to 19	91% (84-96%)	8.9% (4.2-16%)	2.2 (1.3-3.4)	3.1% (1.8-4.6%)	8.2% (3.9-13%)	63 (29-99)	6400 (3800-9400)
Argentina	2010	Female	20 to 24	94% (89-97%)	5.7% (2.9-11%)	3.2 (2-4.7)	4.5% (3-6.1%)	10% (5.3-15%)	83 (44-120)	9900 (6500-14000)
Argentina	2010	Female	25 to 29	95% (91-97%)	5.2% (2.6-9.4%)	3.5 (2.3-5.1)	4.1% (2.9-5.6%)	9.5% (5.6-14%)	88 (52-130)	9400 (6200-13000)
Argentina	2010	Female	30 to 34	94% (90-97%)	5.9% (3.2-10%)	3.4 (2.3-4.9)	3.7% (2.5-5.1%)	8.8% (5.5-13%)	110 (69-160)	9400 (6400-13000)
Argentina	2010	Female	35 to 39	92% (88-96%)	7.7% (3.8-12%)	3.1 (2-4.6)	3.5% (2.2-4.9%)	8.3% (5.1-12%)	130 (78-180)	8700 (5500-13000)
Argentina	2010	Female	40 to 44	92% (86-96%)	8.3% (4.4-14%)	3 (1.9-4.5)	3.8% (2.1-5.7%)	8.5% (4.8-13%)	170 (95-260)	9500 (5300-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	2010	Female	45 to 49	92% (87-96%)	7.7% (4.3-13%)	3 (1.9-4.6)	4.1% (2-6.7%)	8.3% (4.4-13%)	240 (130-380)	11000 (5600-18000)
Argentina	2010	Female	50 to 54	93% (87-96%)	7.4% (4.1-13%)	2.8 (1.7-4.3)	4.1% (1.8-6.8%)	7.5% (3.7-12%)	320 (160-520)	13000 (5500-21000)
Argentina	2010	Female	55 to 59	92% (87-96%)	7.6% (4.1-13%)	2.7 (1.7-4)	3.8% (1.3-6.5%)	6.5% (2.8-10%)	390 (160-630)	13000 (4600-22000)
Argentina	2010	Female	60 to 64	91% (85-96%)	8.7% (4.4-15%)	2.5 (1.5-3.9)	3% (0.37-6.2%)	4.7% (1.1-9.3%)	380 (88-740)	10000 (1200-22000)
Argentina	2010	Female	65 to 69	90% (80-95%)	10% (4.8-20%)	2.1 (1.4-3.1)	2% (-0.29-4.6%)	3.1% (0.099-6.5%)	310 (10-650)	6800 (-980-16000)
Argentina	2010	Female	70 to 74	88% (79-94%)	12% (6.2-21%)	1.9 (1.1-3)	1.2% (-1.2-4%)	1.8% (-1.3-5.5%)	240 (-170-710)	4000 (-4200-14000)
Argentina	2010	Female	75 to 79	87% (77-93%)	13% (6.7-23%)	1.9 (1-3)	0.41% (-2.2-3.5%)	0.63% (-2.7-4.5%)	120 (-480-840)	1500 (-7700-13000)
Argentina	2010	Female	80 to 84	85% (74-93%)	15% (7.3-26%)	1.8 (1.1-2.8)	0.14% (-2.3-3%)	0.18% (-2.8-3.7%)	46 (-730-950)	480 (-7800-11000)
Argentina	2010	Female	85 to 89	84% (74-91%)	16% (8.6-26%)	1.7 (1-2.7)	0.017% (-2.4-2.9%)	-0.059% (-2.9-3.2%)	-16 (-810-880)	45 (-5900-7100)
Argentina	2010	Female	90 to 94	84% (75-92%)	16% (8.2-25%)	1.8 (0.96-2.9)	-0.11% (-2.6-3.1%)	-0.27% (-3.1-3.5%)	-50 (-580-670)	-120 (-2900-3600)
Argentina	2010	Female	95 plus	84% (72-94%)	16% (6.5-28%)	1.8 (0.87-3.1)	-0.097% (-3-4%)	-0.33% (-3.7-4.6%)	-25 (-280-350)	-30 (-900-1200)
Argentina	2010	Male	All Ages	94% (92-96%)	5.8% (4-7.9%)	2.8 (2.3-3.3)	6% (4.4-7.7%)	4.8% (2.5-7.5%)	7900 (4000-12000)	350000 (250000-450000)
Argentina	2010	Male	15 to 19	94% (89-97%)	6.1% (2.7-11%)	2.7 (1.7-4.1)	8.1% (5-11%)	13% (7.2-20%)	230 (120-350)	21000 (13000-29000)
Argentina	2010	Male	20 to 24	95% (92-98%)	4.5% (2.2-8.2%)	4.2 (2.7-6.3)	14% (9.8-19%)	20% (12-28%)	440 (280-630)	41000 (28000-55000)
Argentina	2010	Male	25 to 29	95% (91-98%)	4.6% (1.9-8.7%)	4.7 (3.2-6.6)	13% (9.6-17%)	20% (13-27%)	430 (280-610)	36000 (26000-48000)
Argentina	2010	Male	30 to 34	95% (91-98%)	4.5% (2-9.1%)	4.6 (3.1-6.6)	11% (8-15%)	17% (11-24%)	430 (270-600)	32000 (22000-42000)
Argentina	2010	Male	35 to 39	96% (92-98%)	4.3% (1.8-8.3%)	4.1 (2.7-5.9)	9.5% (6.6-13%)	15% (9.2-21%)	380 (240-530)	25000 (17000-34000)
Argentina	2010	Male	40 to 44	96% (92-98%)	3.9% (1.5-7.7%)	3.9 (2.4-5.7)	9% (6.2-12%)	13% (8.3-19%)	430 (270-620)	24000 (16000-34000)
Argentina	2010	Male	45 to 49	97% (93-99%)	3.3% (1.4-6.9%)	3.8 (2.4-5.6)	8.8% (5.6-13%)	12% (7.4-18%)	590 (350-880)	28000 (18000-41000)
Argentina	2010	Male	50 to 54	97% (94-99%)	3.1% (1.3-6.3%)	3.7 (2.2-5.5)	8.2% (4.8-13%)	11% (6.1-17%)	790 (450-1300)	33000 (19000-50000)
Argentina	2010	Male	55 to 59	97% (93-99%)	3.3% (1.3-6.5%)	3.6 (2.3-5.3)	7.3% (3.9-12%)	9% (4.8-15%)	960 (510-1600)	34000 (18000-54000)
Argentina	2010	Male	60 to 64	97% (92-99%)	3.5% (1.3-7.5%)	3.4 (2-5.2)	5.8% (2.5-11%)	6.9% (2.8-13%)	980 (400-1800)	29000 (12000-53000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	2010	Male	65 to 69	96% (92-99%)	3.7% (1.4-7.8%)	3.1 (2-4.4)	4.2% (1.3-7.8%)	4.8% (1.4-9.1%)	800 (240-1500)	20000 (6300-37000)
Argentina	2010	Male	70 to 74	96% (91-98%)	4.1% (1.7-8.7%)	2.7 (1.7-4.1)	2.9% (0.044-6.5%)	3.2% (-0.16-7.4%)	610 (-30-1400)	12000 (180-28000)
Argentina	2010	Male	75 to 79	95% (91-98%)	4.5% (1.9-8.9%)	2.5 (1.4-3.9)	1.9% (-0.8-5.4%)	2% (-1.1-6%)	420 (-240-1300)	6900 (-2800-20000)
Argentina	2010	Male	80 to 84	95% (89-98%)	5.3% (2.3-11%)	2.3 (1.4-3.3)	1.2% (-1.1-3.8%)	1.1% (-1.6-4.1%)	240 (-330-890)	3100 (-3000-10000)
Argentina	2010	Male	85 to 89	94% (88-97%)	6.3% (2.8-12%)	2 (1.2-3.1)	0.83% (-1.2-3.2%)	0.72% (-1.7-3.3%)	120 (-280-540)	1200 (-1700-4400)
Argentina	2010	Male	90 to 94	93% (88-97%)	6.6% (2.9-12%)	2 (1.1-3.1)	0.52% (-1.6-3.1%)	0.33% (-2.1-3.3%)	25 (-160-260)	230 (-710-1400)
Argentina	2010	Male	95 plus	94% (86-98%)	6.4% (2.4-14%)	2 (0.9-3.4)	0.52% (-1.8-3.9%)	0.27% (-2.4-4.1%)	5.9 (-53-90)	45 (-160-340)
Argentina	2016	Female	All Ages	90% (87-92%)	10% (7.9-13%)	2.5 (1.9-3.4)	2.9% (1.2-4.9%)	2.9% (-0.25-6.7%)	4700 (-400-11000)	160000 (66000-270000)
Argentina	2016	Female	15 to 19	91% (84-96%)	9% (4.4-16%)	2.6 (1.5-4.2)	3.4% (1.9-5.3%)	9.7% (4.5-16%)	69 (32-110)	7000 (4000-11000)
Argentina	2016	Female	20 to 24	94% (90-97%)	5.8% (2.9-10%)	3.8 (2.2-6)	4.9% (3.2-6.9%)	12% (6.3-18%)	95 (49-150)	11000 (7100-16000)
Argentina	2016	Female	25 to 29	95% (90-97%)	5.3% (2.6-9.6%)	4.1 (2.4-6.3)	4.6% (3.1-6.4%)	12% (6.6-17%)	100 (58-160)	11000 (6800-15000)
Argentina	2016	Female	30 to 34	94% (90-97%)	6% (3.2-10%)	4.1 (2.5-6.2)	4.1% (2.7-5.8%)	11% (6.5-16%)	120 (72-180)	10000 (6500-15000)
Argentina	2016	Female	35 to 39	92% (87-96%)	7.7% (3.7-13%)	3.7 (2.2-5.7)	4% (2.4-5.9%)	9.9% (5.8-15%)	160 (92-240)	11000 (6900-17000)
Argentina	2016	Female	40 to 44	92% (86-95%)	8.4% (4.5-14%)	3.6 (2-5.6)	4.3% (2.4-6.7%)	10% (5.7-15%)	220 (120-350)	12000 (7000-20000)
Argentina	2016	Female	45 to 49	92% (86-96%)	7.8% (4.2-14%)	3.5 (2-5.7)	4.7% (2.4-7.7%)	9.7% (5.3-16%)	280 (150-460)	13000 (6800-22000)
Argentina	2016	Female	50 to 54	93% (87-96%)	7.5% (4-13%)	3.3 (1.8-5.3)	4.9% (2.1-8.3%)	9% (4.5-15%)	380 (180-630)	15000 (6500-26000)
Argentina	2016	Female	55 to 59	92% (87-96%)	7.6% (4.1-13%)	3.2 (1.9-5.1)	4.8% (1.7-8.4%)	8.1% (3.4-14%)	490 (210-820)	17000 (5900-29000)
Argentina	2016	Female	60 to 64	91% (85-96%)	8.7% (4.3-15%)	3 (1.6-4.9)	4% (0.78-8.3%)	6.3% (1.8-12%)	530 (150-1000)	15000 (3000-31000)
Argentina	2016	Female	65 to 69	89% (80-95%)	11% (4.9-20%)	2.5 (1.5-4)	3% (0.19-6.6%)	4.5% (0.83-9.5%)	500 (89-1000)	12000 (750-25000)
Argentina	2016	Female	70 to 74	88% (79-94%)	12% (6.1-21%)	2.2 (1.2-3.7)	2.1% (-0.76-6%)	3.1% (-0.59-8.1%)	420 (-83-1100)	7900 (-3000-22000)
Argentina	2016	Female	75 to 79	87% (77-93%)	13% (6.7-23%)	2.2 (1.1-3.6)	1.4% (-1.6-5.7%)	1.9% (-1.9-7.2%)	330 (-340-1300)	5000 (-5800-20000)
Argentina	2016	Female	80 to 84	86% (74-93%)	14% (7.3-26%)	2.1 (1.2-3.4)	1.2% (-1.8-4.9%)	1.4% (-2.2-6%)	360 (-580-1500)	4200 (-6500-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Argentina	2016	Female	85 to 89	85% (73-92%)	15% (8.4-27%)	2 (1.2-3.4)	1.1% (-1.8-4.8%)	1.2% (-2.2-5.5%)	340 (-680-1600)	2900 (-4900-13000)
Argentina	2016	Female	90 to 94	84% (74-92%)	16% (8.3-26%)	2.1 (1.1-3.5)	0.96% (-2.5-3%)	0.96% (-2.6-6%)	220 (-590-1400)	1300 (-2800-7200)
Argentina	2016	Female	95 plus	84% (71-93%)	16% (6.9-29%)	2.1 (0.95-3.8)	1% (-2.3-6.5%)	0.96% (-3.1-7.6%)	93 (-300-730)	400 (-920-2500)
Argentina	2016	Male	All Ages	94% (92-96%)	5.7% (3.9-7.7%)	2.6 (1.9-3.5)	5.8% (3.8-8%)	4.5% (2-7.8%)	7700 (3400-13000)	350000 (230000-490000)
Argentina	2016	Male	15 to 19	94% (89-97%)	6.1% (2.7-11%)	2.6 (1.4-4.2)	8% (4.8-12%)	13% (6.6-21%)	240 (120-370)	21000 (13000-31000)
Argentina	2016	Male	20 to 24	95% (92-98%)	4.5% (2.3-8.3%)	3.9 (2.3-6)	14% (9.3-19%)	19% (11-28%)	460 (260-690)	43000 (28000-60000)
Argentina	2016	Male	25 to 29	95% (91-98%)	4.5% (2-8.6%)	4.3 (2.5-6.6)	13% (9-18%)	19% (11-28%)	440 (250-650)	38000 (25000-52000)
Argentina	2016	Male	30 to 34	95% (91-98%)	4.5% (2-9%)	4.3 (2.6-6.6)	11% (7.4-15%)	17% (10-24%)	400 (240-590)	30000 (20000-42000)
Argentina	2016	Male	35 to 39	96% (92-98%)	4.4% (1.9-8.2%)	3.9 (2.3-6)	9.1% (5.9-13%)	14% (8-22%)	410 (240-620)	28000 (18000-40000)
Argentina	2016	Male	40 to 44	96% (92-98%)	3.9% (1.5-7.8%)	3.6 (2-5.8)	8.6% (5.6-13%)	13% (7.8-20%)	460 (270-700)	26000 (17000-39000)
Argentina	2016	Male	45 to 49	97% (93-99%)	3.2% (1.4-6.9%)	3.5 (2.1-5.6)	8.2% (5.1-13%)	12% (6.9-18%)	540 (320-880)	27000 (17000-41000)
Argentina	2016	Male	50 to 54	97% (94-99%)	3% (1.2-6%)	3.4 (1.9-5.3)	7.7% (4.4-12%)	10% (5.5-17%)	720 (400-1200)	30000 (17000-49000)
Argentina	2016	Male	55 to 59	97% (94-99%)	3.2% (1.3-6.4%)	3.4 (2-5.3)	6.8% (3.5-12%)	8.5% (4.3-15%)	910 (450-1600)	32000 (17000-54000)
Argentina	2016	Male	60 to 64	97% (93-99%)	3.4% (1.4-7.3%)	3.1 (1.7-5.1)	5.4% (2.1-11%)	6.5% (2.5-13%)	960 (360-1900)	28000 (11000-56000)
Argentina	2016	Male	65 to 69	96% (92-99%)	3.7% (1.4-7.9%)	2.8 (1.7-4.5)	3.9% (1.1-7.8%)	4.5% (1.2-9.3%)	830 (220-1700)	21000 (5900-42000)
Argentina	2016	Male	70 to 74	96% (92-98%)	4.1% (1.8-8.2%)	2.5 (1.4-4)	2.6% (-0.094-6.4%)	2.9% (-0.4-7.1%)	610 (-80-1500)	12000 (-440-30000)
Argentina	2016	Male	75 to 79	95% (91-98%)	4.5% (1.8-8.8%)	2.3 (1.2-3.8)	1.7% (-0.83-5.3%)	1.8% (-1.2-5.9%)	380 (-260-1300)	6300 (-3100-20000)
Argentina	2016	Male	80 to 84	95% (89-98%)	5.3% (2.2-11%)	2.1 (1.2-3.3)	1.1% (-1.2-3.9%)	1% (-1.6-4.3%)	220 (-350-960)	2900 (-3400-11000)
Argentina	2016	Male	85 to 89	94% (88-97%)	6.4% (2.8-12%)	1.9 (1.1-3.1)	0.73% (-1.3-3.4%)	0.62% (-1.8-3.6%)	110 (-320-660)	1100 (-2000-5400)
Argentina	2016	Male	90 to 94	93% (87-97%)	6.8% (2.9-13%)	1.8 (0.96-3.1)	0.44% (-1.7-3.5%)	0.25% (-2.2-3.8%)	25 (-210-360)	250 (-970-2000)
Argentina	2016	Male	95 plus	93% (86-98%)	6.7% (2.4-14%)	1.8 (0.81-3.3)	0.45% (-1.8-4.1%)	0.19% (-2.6-4.3%)	5.5 (-74-130)	51 (-210-450)
Chile	1990	Female	All Ages	76% (70-81%)	24% (19-30%)	2.6 (2.3-2.9)	4.1% (3-5.5%)	5% (2.2-8.4%)	1700 (740-2900)	66000 (46000-86000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	1990	Female	15 to 19	77% (63-88%)	23% (12-37%)	2.7 (1.7-4)	4.1% (2.6-5.8%)	11% (5.8-16%)	26 (14-40)	3000 (2000-4300)
Chile	1990	Female	20 to 24	83% (72-91%)	17% (9-28%)	4 (2.6-5.7)	6.3% (4.7-8.2%)	14% (8.8-19%)	40 (25-56)	5500 (3900-7400)
Chile	1990	Female	25 to 29	85% (75-92%)	15% (8-25%)	4.4 (3-6.1)	6.2% (4.7-7.9%)	14% (10-19%)	50 (35-69)	5500 (4000-7200)
Chile	1990	Female	30 to 34	85% (75-92%)	15% (7.9-25%)	4.4 (2.9-6.2)	5.8% (4.3-7.5%)	14% (9.9-18%)	59 (42-79)	5100 (3700-6500)
Chile	1990	Female	35 to 39	84% (73-92%)	16% (8-27%)	4 (2.5-5.7)	5.6% (4-7.5%)	13% (9.4-18%)	71 (49-95)	4900 (3500-6500)
Chile	1990	Female	40 to 44	84% (72-92%)	16% (8.4-28%)	3.7 (2.2-5.5)	6.3% (4.3-8.3%)	13% (9.2-17%)	96 (67-130)	5400 (3700-7100)
Chile	1990	Female	45 to 49	83% (70-91%)	17% (8.5-30%)	3.7 (2.2-5.7)	6.7% (4.5-9.3%)	13% (8.6-17%)	130 (85-180)	5900 (4000-8200)
Chile	1990	Female	50 to 54	82% (69-91%)	18% (9.1-31%)	3.5 (2.1-5.4)	7.1% (4.3-10%)	12% (7.5-17%)	150 (94-220)	6100 (3700-8700)
Chile	1990	Female	55 to 59	81% (68-90%)	19% (9.5-32%)	3.4 (2-5)	6.9% (3.7-10%)	11% (6-15%)	180 (100-270)	6300 (3400-9200)
Chile	1990	Female	60 to 64	80% (65-90%)	20% (9.8-35%)	3.1 (1.7-4.9)	5.8% (2-9.6%)	8.2% (3-13%)	200 (72-330)	5600 (2000-9600)
Chile	1990	Female	65 to 69	77% (62-89%)	23% (11-38%)	2.7 (1.7-4.1)	4.7% (1.4-8.4%)	6.4% (2-11%)	180 (59-320)	4400 (1300-7700)
Chile	1990	Female	70 to 74	74% (60-85%)	26% (15-40%)	2.4 (1.4-3.9)	3.6% (-0.22-7.9%)	4.6% (-0.21-9.9%)	170 (-7.7-380)	3300 (-200-7300)
Chile	1990	Female	75 to 79	72% (57-84%)	28% (16-43%)	2.3 (1.2-3.8)	2.6% (-1.3-7.4%)	3.1% (-1.7-9%)	150 (-82-430)	2200 (-1100-6400)
Chile	1990	Female	80 to 84	70% (52-83%)	30% (17-48%)	2.2 (1.2-3.4)	2.2% (-1.3-6.4%)	2.5% (-1.7-7.6%)	120 (-85-370)	1400 (-820-4000)
Chile	1990	Female	85 to 89	68% (51-81%)	32% (19-49%)	2.1 (1.2-3.3)	1.9% (-1.4-6.2%)	2% (-2-7.3%)	80 (-77-280)	670 (-500-2200)
Chile	1990	Female	90 to 94	67% (53-80%)	33% (20-47%)	2.1 (1.1-3.5)	1.7% (-1.5-6.4%)	1.8% (-2.2-7.8%)	35 (-41-140)	210 (-190-780)
Chile	1990	Female	95 plus	68% (48-83%)	32% (17-52%)	2.1 (0.97-3.8)	1.9% (-2.2-8.7%)	1.9% (-2.9-9.9%)	15 (-24-79)	60 (-71-270)
Chile	1990	Male	All Ages	84% (79-89%)	16% (11-21%)	2.7 (2.4-3)	8.6% (7-10%)	8.6% (6-12%)	3700 (2600-5000)	170000 (130000-200000)
Chile	1990	Male	15 to 19	83% (71-91%)	17% (8.9-29%)	2.7 (1.7-3.9)	8.3% (5.2-12%)	13% (7.2-20%)	79 (42-120)	7500 (4800-11000)
Chile	1990	Male	20 to 24	88% (78-94%)	12% (6.5-22%)	4.1 (2.8-5.9)	15% (11-20%)	20% (12-29%)	210 (120-290)	19000 (13000-25000)
Chile	1990	Male	25 to 29	89% (82-95%)	11% (5.5-18%)	4.7 (3.1-6.4)	16% (12-21%)	22% (15-30%)	240 (160-330)	19000 (14000-25000)
Chile	1990	Male	30 to 34	91% (82-96%)	9.3% (4.4-18%)	4.7 (3.1-6.5)	15% (11-20%)	21% (15-29%)	250 (170-330)	17000 (12000-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	1990	Male	35 to 39	91% (83-96%)	8.8% (4.1-17%)	4.2 (2.7-6)	14% (10-18%)	20% (14-27%)	250 (170-320)	15000 (11000-19000)
Chile	1990	Male	40 to 44	92% (84-96%)	8.4% (4-16%)	3.9 (2.4-5.7)	14% (10-18%)	20% (14-26%)	290 (200-390)	15000 (11000-20000)
Chile	1990	Male	45 to 49	92% (83-96%)	8.2% (3.9-17%)	3.9 (2.4-5.9)	14% (11-19%)	19% (14-25%)	360 (260-480)	16000 (12000-22000)
Chile	1990	Male	50 to 54	92% (84-96%)	8.3% (3.9-16%)	3.7 (2.3-5.6)	13% (9.4-18%)	17% (12-23%)	380 (270-520)	15000 (11000-20000)
Chile	1990	Male	55 to 59	91% (84-96%)	8.7% (4-16%)	3.6 (2.2-5.3)	12% (7.8-16%)	14% (9.6-20%)	420 (280-590)	14000 (9500-20000)
Chile	1990	Male	60 to 64	91% (82-96%)	9.3% (4.2-18%)	3.4 (2-5.2)	9.1% (5.1-14%)	11% (6-17%)	420 (230-660)	12000 (6800-19000)
Chile	1990	Male	65 to 69	90% (81-96%)	10% (4.5-19%)	3.1 (2-4.4)	6.6% (3.3-11%)	7.8% (3.7-13%)	320 (150-530)	7700 (3800-13000)
Chile	1990	Male	70 to 74	89% (81-95%)	11% (5.4-19%)	2.7 (1.6-4.2)	4.6% (0.94-9.3%)	5.2% (0.91-11%)	250 (43-510)	4800 (960-9700)
Chile	1990	Male	75 to 79	88% (79-94%)	12% (5.8-21%)	2.5 (1.3-4)	2.9% (-0.48-7.7%)	3.1% (-0.85-8.6%)	160 (-44-440)	2400 (-400-6500)
Chile	1990	Male	80 to 84	86% (75-93%)	14% (6.9-25%)	2.3 (1.3-3.5)	1.8% (-0.98-5.2%)	1.8% (-1.5-5.7%)	74 (-61-240)	920 (-520-2700)
Chile	1990	Male	85 to 89	84% (72-92%)	16% (8.5-28%)	2 (1.2-3.1)	1.2% (-1.2-4.2%)	1.1% (-1.8-4.6%)	27 (-46-110)	270 (-270-910)
Chile	1990	Male	90 to 94	83% (71-92%)	17% (8.4-29%)	1.9 (1-3.2)	0.83% (-1.5-3.8%)	0.6% (-2.2-4.2%)	6 (-22-42)	52 (-93-240)
Chile	1990	Male	95 plus	83% (67-93%)	17% (7-33%)	1.9 (0.89-3.3)	0.55% (-2.2-4.8%)	0.21% (-3.1-5.4%)	0.71 (-11-19)	7.5 (-31-68)
Chile	1995	Female	All Ages	76% (70-82%)	24% (18-30%)	2 (1.8-2.3)	3% (2.1-4.1%)	3.1% (0.72-5.7%)	1100 (260-2100)	48000 (31000-65000)
Chile	1995	Female	15 to 19	77% (63-88%)	23% (12-37%)	2.1 (1.3-3.2)	3.2% (2.1-4.5%)	8.3% (4.4-13%)	16 (8.7-26)	2200 (1400-3100)
Chile	1995	Female	20 to 24	83% (72-91%)	17% (8.9-28%)	3.1 (1.9-4.5)	5.3% (3.8-7%)	11% (6.3-16%)	26 (14-38)	4300 (3000-6000)
Chile	1995	Female	25 to 29	85% (74-92%)	15% (7.9-26%)	3.4 (2.3-4.8)	5% (3.7-6.4%)	11% (7.4-16%)	33 (21-46)	4400 (3200-5900)
Chile	1995	Female	30 to 34	85% (74-92%)	15% (8.1-26%)	3.4 (2.3-4.9)	4.4% (3.2-5.7%)	11% (7.7-15%)	40 (28-55)	4000 (2900-5200)
Chile	1995	Female	35 to 39	84% (73-92%)	16% (8.3-27%)	3.1 (2-4.5)	4.4% (3.1-5.8%)	10% (7.1-14%)	56 (38-76)	4100 (3000-5500)
Chile	1995	Female	40 to 44	83% (72-91%)	17% (8.5-28%)	2.9 (1.7-4.2)	4.5% (3-6.1%)	10% (6.7-14%)	73 (47-100)	4300 (2900-5800)
Chile	1995	Female	45 to 49	83% (68-91%)	17% (8.6-32%)	2.8 (1.7-4.3)	4.8% (3.1-7%)	9.7% (6.2-14%)	93 (59-130)	4500 (2800-6400)
Chile	1995	Female	50 to 54	82% (69-91%)	18% (9.4-31%)	2.7 (1.6-4.2)	5% (2.8-7.6%)	8.9% (5.4-13%)	120 (70-170)	4800 (2800-7000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	1995	Female	55 to 59	81% (69-91%)	19% (9-31%)	2.6 (1.6-4)	4.8% (2.4-7.4%)	7.9% (4.3-12%)	130 (71-190)	4400 (2300-6800)
Chile	1995	Female	60 to 64	80% (65-90%)	20% (10-35%)	2.4 (1.4-3.8)	3.9% (1.1-7%)	5.8% (1.9-10%)	140 (45-240)	3900 (1100-7000)
Chile	1995	Female	65 to 69	77% (60-89%)	23% (11-40%)	2.1 (1.3-3.2)	2.9% (0.41-5.7%)	4.1% (0.87-7.9%)	130 (26-240)	3000 (450-5900)
Chile	1995	Female	70 to 74	74% (58-85%)	26% (15-42%)	1.9 (1.1-3.1)	2% (-0.72-5.3%)	2.6% (-0.82-6.9%)	100 (-33-270)	2000 (-720-5200)
Chile	1995	Female	75 to 79	71% (55-84%)	29% (16-45%)	1.8 (0.91-2.9)	1.2% (-1.6-4.7%)	1.4% (-2.5-7%)	67 (-95-270)	1000 (-1400-4100)
Chile	1995	Female	80 to 84	69% (52-83%)	31% (17-48%)	1.7 (0.95-2.6)	0.84% (-1.7-3.7%)	0.94% (-2.2-4.4%)	47 (-110-220)	570 (-1100-2500)
Chile	1995	Female	85 to 89	67% (50-80%)	33% (20-50%)	1.6 (0.89-2.6)	0.63% (-1.8-3.7%)	0.6% (-2.4-4.2%)	27 (-110-190)	260 (-760-1600)
Chile	1995	Female	90 to 94	67% (51-80%)	33% (20-49%)	1.6 (0.83-2.7)	0.54% (-1.9-4%)	0.45% (-2.4-4.5%)	12 (-66-120)	92 (-310-670)
Chile	1995	Female	95 plus	67% (48-83%)	33% (17-52%)	1.6 (0.75-3)	0.61% (-2.4-5.4%)	0.46% (-3.1-6.2%)	5 (-34-68)	27 (-100-230)
Chile	1995	Male	All Ages	84% (79-88%)	16% (12-21%)	2.5 (2.2-2.8)	8% (6.6-9.7%)	7.6% (5.2-10%)	3300 (2200-4500)	150000 (120000-180000)
Chile	1995	Male	15 to 19	82% (71-91%)	18% (9.2-29%)	2.4 (1.5-3.6)	7.5% (4.9-11%)	12% (6.8-18%)	64 (34-98)	6400 (4200-8900)
Chile	1995	Male	20 to 24	87% (78-93%)	13% (6.7-22%)	3.8 (2.5-5.5)	15% (10-19%)	19% (11-27%)	170 (100-250)	16000 (11000-22000)
Chile	1995	Male	25 to 29	89% (81-94%)	11% (5.6-19%)	4.3 (2.9-5.8)	15% (11-19%)	20% (13-27%)	210 (140-290)	18000 (13000-23000)
Chile	1995	Male	30 to 34	90% (82-95%)	9.6% (4.7-18%)	4.2 (2.9-5.9)	13% (9.8-17%)	19% (13-26%)	210 (150-300)	16000 (11000-20000)
Chile	1995	Male	35 to 39	91% (84-96%)	9.1% (4.2-16%)	3.8 (2.5-5.4)	12% (9.1-16%)	18% (13-24%)	220 (150-290)	14000 (10000-18000)
Chile	1995	Male	40 to 44	91% (84-96%)	8.7% (4-16%)	3.6 (2.2-5.1)	12% (8.9-16%)	17% (12-23%)	250 (170-340)	14000 (10000-18000)
Chile	1995	Male	45 to 49	92% (82-96%)	8.5% (3.8-18%)	3.5 (2.2-5.3)	12% (8.9-16%)	17% (12-22%)	300 (220-410)	14000 (10000-19000)
Chile	1995	Male	50 to 54	91% (83-96%)	8.6% (3.9-17%)	3.4 (2.1-5.1)	12% (8.2-16%)	15% (11-21%)	350 (240-480)	14000 (10000-19000)
Chile	1995	Male	55 to 59	91% (83-96%)	9.1% (4.1-17%)	3.3 (2-4.7)	10% (7.1-14%)	13% (9-18%)	350 (240-490)	12000 (8300-16000)
Chile	1995	Male	60 to 64	90% (81-96%)	9.7% (4.2-19%)	3.1 (1.8-4.6)	8% (4.7-12%)	9.8% (5.7-15%)	360 (210-550)	10000 (6100-16000)
Chile	1995	Male	65 to 69	90% (81-96%)	10% (4.4-19%)	2.8 (1.8-4)	6% (3.2-9.5%)	7.1% (3.8-11%)	310 (160-500)	7500 (4000-12000)
Chile	1995	Male	70 to 74	89% (80-95%)	11% (5.2-20%)	2.5 (1.5-3.8)	4% (1.1-7.8%)	4.6% (1.1-9.1%)	230 (56-460)	4500 (1200-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	1995	Male	75 to 79	88% (79-94%)	12% (6-21%)	2.2 (1.2-3.5)	2.6% (-0.25-6.2%)	2.8% (-0.43-7.1%)	140 (-22-360)	2200 (-210-5500)
Chile	1995	Male	80 to 84	86% (75-93%)	14% (7-25%)	2 (1.2-3.1)	1.6% (-0.74-4.3%)	1.7% (-1.1-4.9%)	75 (-51-220)	910 (-410-2500)
Chile	1995	Male	85 to 89	83% (71-91%)	17% (8.6-29%)	1.8 (1.1-2.8)	1.1% (-1-3.4%)	0.99% (-1.5-3.8%)	32 (-49-120)	300 (-300-980)
Chile	1995	Male	90 to 94	82% (69-92%)	18% (8.5-31%)	1.7 (0.93-2.8)	0.68% (-1.4-3.3%)	0.51% (-2-3.6%)	7.8 (-30-55)	63 (-130-300)
Chile	1995	Male	95 plus	82% (65-93%)	18% (6.7-35%)	1.7 (0.8-3)	0.62% (-1.6-4.3%)	0.37% (-2.4-4.6%)	1.7 (-11-22)	11 (-30-80)
Chile	2000	Female	All Ages	76% (70-82%)	24% (18-30%)	2 (1.7-2.2)	2.9% (2.1-3.9%)	3.1% (1-5.5%)	1200 (390-2100)	48000 (33000-64000)
Chile	2000	Female	15 to 19	76% (62-87%)	24% (13-38%)	2 (1.2-3)	3% (2-4.2%)	8.4% (4.4-13%)	15 (8.1-24)	2200 (1400-3200)
Chile	2000	Female	20 to 24	83% (71-91%)	17% (9.3-29%)	3 (1.8-4.3)	5.2% (3.8-6.7%)	11% (6.1-16%)	23 (13-33)	4100 (2900-5700)
Chile	2000	Female	25 to 29	85% (74-92%)	15% (8.1-26%)	3.3 (2.1-4.6)	4.8% (3.6-6.2%)	11% (7.3-16%)	28 (18-40)	4200 (3000-5600)
Chile	2000	Female	30 to 34	85% (73-92%)	15% (8.2-27%)	3.3 (2.2-4.6)	4% (2.9-5.2%)	11% (7.3-14%)	36 (25-50)	4000 (2800-5300)
Chile	2000	Female	35 to 39	84% (72-92%)	16% (8.4-28%)	3 (1.9-4.2)	3.8% (2.7-5%)	10% (6.9-14%)	47 (32-64)	3900 (2700-5200)
Chile	2000	Female	40 to 44	83% (72-91%)	17% (8.7-28%)	2.8 (1.7-4.1)	4% (2.6-5.5%)	9.9% (6.4-13%)	66 (42-90)	4200 (2800-5700)
Chile	2000	Female	45 to 49	83% (68-91%)	17% (8.7-32%)	2.7 (1.7-4)	4.2% (2.6-6%)	9.4% (6-13%)	90 (57-130)	4500 (2900-6300)
Chile	2000	Female	50 to 54	82% (69-91%)	18% (9.3-31%)	2.6 (1.5-4)	4.4% (2.6-6.6%)	8.5% (5.3-12%)	110 (67-160)	4500 (2700-6600)
Chile	2000	Female	55 to 59	81% (69-91%)	19% (9.2-31%)	2.5 (1.5-3.7)	4.4% (2.4-6.6%)	7.8% (4.6-11%)	130 (78-190)	4600 (2500-6800)
Chile	2000	Female	60 to 64	79% (65-90%)	21% (10-35%)	2.3 (1.3-3.7)	3.7% (1.4-6.3%)	6% (2.7-9.9%)	130 (57-210)	3600 (1400-6200)
Chile	2000	Female	65 to 69	76% (60-89%)	24% (11-40%)	2 (1.2-3.1)	3% (0.95-5.4%)	4.5% (1.7-7.7%)	130 (51-230)	3100 (990-5600)
Chile	2000	Female	70 to 74	74% (58-85%)	26% (15-42%)	1.8 (0.99-2.9)	2.1% (-0.23-4.8%)	3.1% (-0.003-6.6%)	120 (-0.12-260)	2200 (-250-5000)
Chile	2000	Female	75 to 79	71% (55-84%)	29% (16-45%)	1.7 (0.85-2.8)	1.3% (-1.1-4.3%)	1.7% (-1.2-5.5%)	85 (-60-270)	1200 (-1000-4100)
Chile	2000	Female	80 to 84	69% (51-82%)	31% (18-49%)	1.6 (0.88-2.5)	0.99% (-1.1-3.5%)	1.2% (-1.4-4.3%)	67 (-78-240)	750 (-840-2700)
Chile	2000	Female	85 to 89	67% (50-80%)	33% (20-50%)	1.5 (0.85-2.4)	0.8% (-1.2-3.6%)	0.89% (-1.6-4%)	45 (-79-210)	390 (-550-1700)
Chile	2000	Female	90 to 94	66% (51-80%)	34% (20-49%)	1.5 (0.8-2.5)	0.69% (-1.5-3.7%)	0.67% (-1.9-4.4%)	24 (-68-150)	150 (-330-840)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	2000	Female	95 plus	67% (47-83%)	33% (17-53%)	1.6 (0.7-2.9)	0.73% (-1.8-4.7%)	0.64% (-2.4-5.3%)	11 (-41-92)	49 (-120-320)
Chile	2000	Male	All Ages	84% (79-89%)	16% (11-21%)	2.1 (1.9-2.4)	7.2% (5.9-8.4%)	6.6% (4.7-8.5%)	2900 (2000-3800)	130000 (110000-160000)
Chile	2000	Male	15 to 19	82% (70-91%)	18% (9.3-30%)	2 (1.2-3.1)	6.7% (4.5-9.3%)	11% (5.7-17%)	51 (27-79)	5700 (3700-8000)
Chile	2000	Male	20 to 24	87% (77-93%)	13% (6.9-23%)	3.2 (2-4.7)	13% (9.5-17%)	17% (9.7-24%)	120 (73-180)	13000 (9400-18000)
Chile	2000	Male	25 to 29	89% (81-94%)	11% (5.6-19%)	3.6 (2.4-5)	13% (9.8-17%)	17% (11-24%)	160 (94-220)	15000 (10000-19000)
Chile	2000	Male	30 to 34	90% (81-95%)	9.8% (4.7-19%)	3.5 (2.4-5)	12% (8.9-15%)	16% (11-22%)	180 (120-240)	14000 (10000-18000)
Chile	2000	Male	35 to 39	91% (83-96%)	9.1% (4.2-17%)	3.2 (2.1-4.5)	11% (8.2-13%)	16% (11-21%)	190 (130-250)	13000 (9600-17000)
Chile	2000	Male	40 to 44	91% (84-96%)	8.7% (4-16%)	3 (1.9-4.3)	11% (7.9-13%)	16% (11-20%)	230 (160-300)	13000 (9900-17000)
Chile	2000	Male	45 to 49	92% (83-96%)	8.5% (3.8-17%)	2.9 (1.8-4.5)	11% (7.9-14%)	15% (11-20%)	270 (200-360)	13000 (9900-17000)
Chile	2000	Male	50 to 54	91% (83-96%)	8.7% (4-17%)	2.8 (1.7-4.2)	9.7% (7.2-13%)	13% (9.8-18%)	300 (220-400)	12000 (9000-16000)
Chile	2000	Male	55 to 59	91% (83-96%)	9.1% (4.1-17%)	2.7 (1.7-3.9)	8.7% (6.1-12%)	12% (8.3-15%)	330 (230-450)	12000 (8200-15000)
Chile	2000	Male	60 to 64	90% (82-96%)	9.7% (4.2-18%)	2.6 (1.5-3.9)	7% (4.5-10%)	8.9% (5.8-13%)	290 (190-430)	8600 (5500-13000)
Chile	2000	Male	65 to 69	90% (80-95%)	10% (4.7-20%)	2.3 (1.5-3.3)	5.2% (3-7.7%)	6.4% (3.8-9.5%)	280 (170-420)	6700 (4000-10000)
Chile	2000	Male	70 to 74	89% (79-95%)	11% (5.4-21%)	2.1 (1.2-3.2)	3.5% (1.3-6.3%)	4.2% (1.5-7.5%)	210 (76-390)	4200 (1600-7600)
Chile	2000	Male	75 to 79	87% (78-94%)	13% (6.3-22%)	1.9 (0.99-2.9)	2.2% (0.11-4.9%)	2.5% (0.044-5.6%)	130 (2.4-300)	2000 (100-4600)
Chile	2000	Male	80 to 84	85% (74-93%)	15% (7.1-26%)	1.7 (0.96-2.6)	1.4% (-0.33-3.5%)	1.6% (-0.55-4%)	75 (-27-190)	890 (-200-2200)
Chile	2000	Male	85 to 89	83% (70-91%)	17% (9-30%)	1.5 (0.87-2.3)	0.99% (-0.58-2.8%)	0.99% (-0.84-3.1%)	34 (-29-110)	310 (-180-890)
Chile	2000	Male	90 to 94	82% (68-92%)	18% (8.4-32%)	1.5 (0.75-2.4)	0.55% (-1.1-2.5%)	0.44% (-1.5-2.7%)	8.4 (-28-54)	65 (-120-300)
Chile	2000	Male	95 plus	82% (66-93%)	18% (6.8-34%)	1.5 (0.67-2.6)	0.53% (-1.3-3.2%)	0.34% (-1.8-3.5%)	2.1 (-12-22)	13 (-32-80)
Chile	2005	Female	All Ages	77% (71-82%)	23% (18-29%)	2.1 (1.8-2.4)	3% (2.2-4%)	3.3% (1.3-5.6%)	1400 (530-2300)	52000 (36000-69000)
Chile	2005	Female	15 to 19	76% (63-87%)	24% (13-37%)	2 (1.2-3.1)	3% (2-4.2%)	8.9% (4.6-14%)	17 (8.8-28)	2400 (1600-3400)
Chile	2005	Female	20 to 24	83% (72-91%)	17% (8.7-28%)	3 (1.8-4.4)	5.2% (3.8-6.8%)	12% (6.4-17%)	23 (12-33)	4300 (2900-5900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	2005	Female	25 to 29	85% (74-92%)	15% (8-26%)	3.3 (2.2-4.7)	4.9% (3.7-6.3%)	12% (7.5-16%)	27 (17-38)	4100 (2900-5500)
Chile	2005	Female	30 to 34	85% (73-92%)	15% (7.9-27%)	3.3 (2.2-4.8)	4.1% (3-5.4%)	11% (7.4-15%)	34 (23-47)	3800 (2700-5100)
Chile	2005	Female	35 to 39	84% (72-92%)	16% (8.2-28%)	3.1 (1.9-4.4)	3.9% (2.8-5.2%)	11% (7.2-14%)	50 (34-68)	4200 (3000-5500)
Chile	2005	Female	40 to 44	83% (71-91%)	17% (8.7-29%)	2.9 (1.7-4.2)	4% (2.7-5.5%)	10% (6.8-14%)	70 (46-97)	4500 (3100-6000)
Chile	2005	Female	45 to 49	83% (68-91%)	17% (8.8-32%)	2.8 (1.7-4.2)	4.3% (2.8-6.1%)	9.7% (6.5-13%)	98 (65-140)	5000 (3300-7000)
Chile	2005	Female	50 to 54	82% (68-91%)	18% (9.3-32%)	2.7 (1.6-4)	4.2% (2.5-6.3%)	8.6% (5.3-12%)	120 (71-170)	4900 (2900-7300)
Chile	2005	Female	55 to 59	81% (68-90%)	19% (9.8-32%)	2.6 (1.6-3.8)	4.2% (2.3-6.2%)	7.6% (4.6-11%)	130 (80-190)	4600 (2600-6900)
Chile	2005	Female	60 to 64	79% (65-90%)	21% (10-35%)	2.4 (1.4-3.7)	3.6% (1.4-6.2%)	6% (2.8-9.8%)	150 (68-240)	4200 (1700-7300)
Chile	2005	Female	65 to 69	77% (60-89%)	23% (11-40%)	2.1 (1.3-3.1)	3% (1.1-5.3%)	4.7% (2-7.8%)	140 (59-230)	3200 (1100-5700)
Chile	2005	Female	70 to 74	73% (57-85%)	27% (15-43%)	1.8 (1-2.9)	2.2% (0.023-4.9%)	3.3% (0.39-6.6%)	140 (16-280)	2500 (25-5400)
Chile	2005	Female	75 to 79	71% (55-84%)	29% (16-45%)	1.7 (0.89-2.8)	1.5% (-0.75-4.3%)	2.1% (-0.78-5.6%)	110 (-44-310)	1600 (-790-4800)
Chile	2005	Female	80 to 84	69% (51-82%)	31% (18-49%)	1.6 (0.9-2.5)	1.3% (-0.8-3.8%)	1.6% (-0.92-4.7%)	100 (-59-300)	1100 (-710-3500)
Chile	2005	Female	85 to 89	67% (50-80%)	33% (20-50%)	1.6 (0.89-2.5)	1.3% (-0.99-3.7%)	1.3% (-1.2-4.4%)	76 (-71-260)	620 (-550-2100)
Chile	2005	Female	90 to 94	66% (51-80%)	34% (20-49%)	1.6 (0.84-2.6)	1% (-1.1-4%)	1.1% (-1.4-4.7%)	47 (-57-200)	280 (-300-1100)
Chile	2005	Female	95 plus	67% (46-83%)	33% (17-54%)	1.6 (0.74-2.9)	1.1% (-1.4-5%)	1.1% (-1.8-5.7%)	25 (-40-120)	96 (-120-430)
Chile	2005	Male	All Ages	85% (80-89%)	15% (11-20%)	2.3 (2-2.6)	7.4% (6.2-8.7%)	6.8% (4.9-8.8%)	3200 (2200-4100)	140000 (120000-170000)
Chile	2005	Male	15 to 19	82% (70-91%)	18% (9-30%)	2.1 (1.3-3.3)	6.8% (4.5-9.4%)	11% (5.9-17%)	57 (30-88)	6300 (4100-8900)
Chile	2005	Male	20 to 24	87% (77-93%)	13% (6.7-23%)	3.3 (2.1-4.9)	13% (9.7-17%)	18% (11-25%)	130 (76-180)	14000 (9700-19000)
Chile	2005	Male	25 to 29	89% (81-94%)	11% (5.7-19%)	3.7 (2.4-5.2)	13% (10-17%)	18% (12-25%)	140 (89-200)	14000 (9900-18000)
Chile	2005	Male	30 to 34	90% (81-95%)	9.9% (4.9-19%)	3.7 (2.5-5.2)	12% (8.9-15%)	17% (12-24%)	160 (110-230)	13000 (9800-17000)
Chile	2005	Male	35 to 39	91% (83-96%)	9.3% (4.2-17%)	3.3 (2.2-4.7)	11% (8.4-14%)	17% (12-22%)	200 (140-260)	14000 (10000-18000)
Chile	2005	Male	40 to 44	91% (84-96%)	8.8% (4.1-16%)	3.1 (2-4.5)	11% (8.3-14%)	17% (12-21%)	240 (180-310)	14000 (11000-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	2005	Male	45 to 49	91% (83-96%)	8.6% (4-17%)	3.1 (1.9-4.7)	11% (8.1-14%)	16% (12-21%)	300 (220-390)	15000 (11000-19000)
Chile	2005	Male	50 to 54	91% (83-96%)	8.6% (4-17%)	3 (1.8-4.5)	9.6% (7-13%)	14% (10-18%)	320 (230-420)	13000 (9800-18000)
Chile	2005	Male	55 to 59	91% (83-96%)	8.9% (4.1-17%)	2.9 (1.8-4.2)	8.5% (6.1-11%)	12% (8.2-15%)	340 (240-460)	12000 (8600-16000)
Chile	2005	Male	60 to 64	90% (82-96%)	9.5% (4.1-18%)	2.7 (1.6-4.1)	7% (4.4-10%)	9.1% (5.8-13%)	350 (220-510)	10000 (6600-15000)
Chile	2005	Male	65 to 69	90% (81-96%)	10% (4.3-19%)	2.5 (1.6-3.5)	5.4% (3.2-8%)	6.8% (4.1-10%)	290 (180-430)	7100 (4200-11000)
Chile	2005	Male	70 to 74	89% (80-95%)	11% (5.2-20%)	2.2 (1.3-3.3)	3.9% (1.7-6.6%)	4.6% (2-7.9%)	260 (110-440)	5000 (2200-8600)
Chile	2005	Male	75 to 79	88% (78-94%)	12% (5.9-22%)	2 (1-3.1)	2.6% (0.55-5.3%)	3% (0.61-6%)	180 (36-370)	2800 (600-5600)
Chile	2005	Male	80 to 84	86% (74-93%)	14% (6.9-26%)	1.8 (1-2.7)	1.8% (0.051-4%)	2% (-0.026-4.5%)	110 (-1.5-250)	1300 (38-2900)
Chile	2005	Male	85 to 89	83% (71-91%)	17% (8.7-29%)	1.6 (0.92-2.4)	1.3% (-0.19-3.1%)	1.4% (-0.38-3.4%)	54 (-15-130)	460 (-69-1100)
Chile	2005	Male	90 to 94	82% (70-92%)	18% (8.4-30%)	1.5 (0.78-2.5)	0.91% (-0.5-2.7%)	0.89% (-0.82-3.1%)	18 (-17-63)	120 (-68-360)
Chile	2005	Male	95 plus	82% (67-93%)	18% (6.9-33%)	1.5 (0.68-2.6)	0.8% (-0.95-3.3%)	0.65% (-1.5-3.6%)	5.4 (-13-30)	26 (-32-110)
Chile	2010	Female	All Ages	77% (71-82%)	23% (18-29%)	2.1 (1.8-2.4)	3% (2.1-4%)	3.3% (1.4-5.6%)	1500 (630-2500)	55000 (38000-73000)
Chile	2010	Female	15 to 19	76% (63-87%)	24% (13-37%)	2 (1.2-3)	3% (1.9-4.3%)	8.4% (4.1-14%)	18 (8.7-29)	2400 (1500-3600)
Chile	2010	Female	20 to 24	83% (71-91%)	17% (8.9-29%)	3 (1.8-4.4)	5.1% (3.5-6.7%)	11% (5.9-16%)	28 (15-42)	4700 (3200-6600)
Chile	2010	Female	25 to 29	85% (75-92%)	15% (8.1-25%)	3.3 (2.2-4.8)	4.7% (3.5-6.2%)	11% (7.2-16%)	29 (19-42)	4200 (2900-5600)
Chile	2010	Female	30 to 34	85% (73-92%)	15% (7.8-27%)	3.4 (2.2-4.9)	4% (2.8-5.4%)	11% (7.2-15%)	35 (23-49)	3700 (2500-4900)
Chile	2010	Female	35 to 39	84% (72-92%)	16% (8.1-28%)	3.1 (1.9-4.4)	3.9% (2.7-5.2%)	11% (7.3-15%)	49 (33-67)	4000 (2800-5300)
Chile	2010	Female	40 to 44	83% (71-92%)	17% (8.5-29%)	2.9 (1.7-4.2)	3.9% (2.6-5.5%)	10% (6.7-14%)	75 (48-100)	4700 (3200-6400)
Chile	2010	Female	45 to 49	83% (69-91%)	17% (9.2-31%)	2.8 (1.7-4.2)	4.2% (2.7-6%)	9.7% (6.4-13%)	110 (70-150)	5400 (3600-7600)
Chile	2010	Female	50 to 54	82% (69-91%)	18% (9.4-31%)	2.7 (1.6-4)	4.2% (2.5-6.2%)	8.6% (5.5-12%)	130 (84-190)	5500 (3400-8000)
Chile	2010	Female	55 to 59	81% (68-91%)	19% (9.4-32%)	2.6 (1.5-3.8)	4% (2.2-5.9%)	7.6% (4.6-11%)	150 (86-210)	5200 (2900-7700)
Chile	2010	Female	60 to 64	79% (65-90%)	21% (10-35%)	2.4 (1.3-3.6)	3.5% (1.5-5.9%)	6% (2.9-9.4%)	150 (75-250)	4500 (1900-7500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	2010	Female	65 to 69	76% (60-88%)	24% (12-40%)	2.1 (1.3-3.1)	2.9% (1.1-5%)	4.8% (2.2-7.4%)	160 (75-250)	3700 (1500-6300)
Chile	2010	Female	70 to 74	73% (58-85%)	27% (15-42%)	1.8 (1-2.9)	2.3% (0.22-4.7%)	3.5% (0.73-6.7%)	140 (30-270)	2600 (250-5500)
Chile	2010	Female	75 to 79	71% (55-84%)	29% (16-45%)	1.7 (0.89-2.8)	1.6% (-0.46-4.3%)	2.3% (-0.37-5.6%)	130 (-20-310)	1800 (-540-4900)
Chile	2010	Female	80 to 84	68% (50-82%)	32% (18-50%)	1.6 (0.92-2.5)	1.3% (-0.7-3.6%)	1.7% (-0.74-4.5%)	120 (-52-310)	1300 (-710-3700)
Chile	2010	Female	85 to 89	66% (49-79%)	34% (21-51%)	1.5 (0.88-2.5)	1.1% (-0.82-3.5%)	1.4% (-1.4-3%)	90 (-65-280)	730 (-510-2400)
Chile	2010	Female	90 to 94	66% (50-80%)	34% (20-50%)	1.6 (0.82-2.6)	1% (-1-3.9%)	1.2% (-1.3-4.6%)	56 (-62-220)	330 (-320-1200)
Chile	2010	Female	95 plus	66% (46-83%)	34% (17-54%)	1.6 (0.72-2.9)	1.1% (-1.5-5%)	1.1% (-1.9-5.7%)	32 (-56-160)	120 (-170-570)
Chile	2010	Male	All Ages	85% (80-89%)	15% (11-20%)	2.8 (2.5-3.2)	8% (6.5-9.5%)	7.8% (5.8-10%)	4000 (2900-5200)	170000 (130000-200000)
Chile	2010	Male	15 to 19	82% (70-91%)	18% (9.4-30%)	2.6 (1.5-3.9)	7.2% (4.7-10%)	12% (6.6-19%)	65 (34-98)	6800 (4400-9400)
Chile	2010	Male	20 to 24	87% (78-93%)	13% (6.6-22%)	4.1 (2.6-5.9)	14% (10-18%)	20% (12-28%)	160 (93-220)	16000 (11000-22000)
Chile	2010	Male	25 to 29	89% (80-94%)	11% (5.7-20%)	4.6 (3-6.3)	14% (10-18%)	20% (14-28%)	170 (110-230)	15000 (11000-20000)
Chile	2010	Male	30 to 34	90% (81-95%)	9.8% (4.7-19%)	4.5 (3-6.5)	12% (9.2-16%)	20% (13-27%)	170 (110-230)	13000 (9300-17000)
Chile	2010	Male	35 to 39	91% (83-96%)	9.2% (4.3-17%)	4 (2.6-5.8)	11% (8.5-15%)	19% (13-24%)	200 (140-270)	14000 (10000-18000)
Chile	2010	Male	40 to 44	91% (84-96%)	8.7% (4-16%)	3.8 (2.4-5.5)	11% (8.4-14%)	18% (13-23%)	270 (190-360)	16000 (12000-20000)
Chile	2010	Male	45 to 49	91% (82-96%)	8.6% (4-18%)	3.8 (2.4-5.6)	11% (8.4-15%)	17% (13-22%)	360 (260-480)	18000 (13000-23000)
Chile	2010	Male	50 to 54	91% (83-96%)	8.7% (4.1-17%)	3.6 (2.2-5.3)	10% (7.4-14%)	15% (11-20%)	400 (280-540)	17000 (12000-22000)
Chile	2010	Male	55 to 59	91% (83-96%)	9.2% (4.1-17%)	3.5 (2.2-5.1)	9.2% (6.3-12%)	13% (8.9-17%)	410 (290-560)	15000 (10000-20000)
Chile	2010	Male	60 to 64	90% (81-96%)	9.8% (4.3-19%)	3.3 (2-5)	7.8% (4.9-11%)	10% (6.5-15%)	420 (260-620)	13000 (7900-18000)
Chile	2010	Male	65 to 69	89% (80-96%)	11% (4.5-20%)	3 (2-4.2)	6.3% (3.8-9.3%)	7.9% (4.8-12%)	400 (240-600)	9900 (6000-15000)
Chile	2010	Male	70 to 74	88% (80-95%)	12% (5.4-20%)	2.7 (1.6-4.1)	4.9% (2.3-8.1%)	5.9% (2.8-9.7%)	330 (160-550)	6500 (3100-11000)
Chile	2010	Male	75 to 79	87% (78-94%)	13% (6.4-22%)	2.4 (1.3-3.8)	3.6% (1.1-7%)	4.2% (1.2-8.2%)	280 (76-540)	4300 (1300-8300)
Chile	2010	Male	80 to 84	85% (74-93%)	15% (7.2-26%)	2.1 (1.3-3.2)	2.7% (0.69-5.3%)	3% (0.67-6.1%)	200 (44-390)	2300 (610-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	2010	Male	85 to 89	82% (70-91%)	18% (8.9-30%)	1.9 (1.1-3)	2% (0.29-4.3%)	2.2% (0.18-4.9%)	110 (8.4-230)	890 (130-1900)
Chile	2010	Male	90 to 94	82% (69-92%)	18% (8.3-31%)	1.8 (0.96-3)	1.6% (-0.081-3.9%)	1.7% (-0.26-4.4%)	40 (-6.2-110)	240 (-12-600)
Chile	2010	Male	95 plus	82% (65-93%)	18% (6.7-35%)	1.8 (0.83-3.2)	1.5% (-0.63-4.9%)	1.5% (-1-5.4%)	17 (-12-62)	70 (-28-220)
Chile	2016	Female	All Ages	77% (71-82%)	23% (18-29%)	2.7 (2-3.5)	3.7% (2.4-5.4%)	4.6% (2.2-8%)	2400 (1000-4400)	73000 (45000-110000)
Chile	2016	Female	15 to 19	76% (62-87%)	24% (13-38%)	2.5 (1.4-4.1)	3.3% (2.1-4.9%)	11% (5.2-17%)	17 (7.4-30)	2300 (1500-3600)
Chile	2016	Female	20 to 24	83% (71-91%)	17% (9.2-29%)	3.8 (2.2-6)	5.5% (4-7.4%)	14% (7.9-21%)	28 (14-48)	4800 (3100-6900)
Chile	2016	Female	25 to 29	84% (73-92%)	16% (8.3-27%)	4.2 (2.5-6.4)	5.1% (3.7-6.7%)	14% (8.9-20%)	34 (18-55)	4900 (3300-6700)
Chile	2016	Female	30 to 34	84% (73-92%)	16% (8.2-27%)	4.2 (2.5-6.5)	4.4% (3.1-5.9%)	13% (8.3-18%)	38 (21-63)	4200 (2800-6000)
Chile	2016	Female	35 to 39	84% (72-92%)	16% (8.5-28%)	3.8 (2.2-5.9)	4.4% (2.9-6.2%)	13% (8.6-18%)	52 (29-88)	4400 (2800-6400)
Chile	2016	Female	40 to 44	83% (71-91%)	17% (8.5-29%)	3.6 (2-5.6)	4.5% (2.9-6.7%)	12% (8.2-17%)	78 (42-130)	5100 (3100-7700)
Chile	2016	Female	45 to 49	82% (66-91%)	18% (9.1-34%)	3.5 (2-5.4)	4.7% (3-7%)	11% (7.5-16%)	120 (66-190)	6400 (3700-9800)
Chile	2016	Female	50 to 54	82% (67-90%)	18% (9.7-33%)	3.3 (2-5.1)	4.9% (2.9-7.6%)	10% (6.6-15%)	160 (90-260)	7200 (4000-11000)
Chile	2016	Female	55 to 59	81% (68-91%)	19% (9.4-32%)	3.2 (1.9-5)	4.9% (2.6-7.5%)	9.3% (5.7-13%)	200 (110-330)	7300 (3800-12000)
Chile	2016	Female	60 to 64	79% (64-90%)	21% (10-36%)	2.9 (1.6-4.7)	4.4% (1.9-7.4%)	7.6% (3.9-12%)	220 (110-380)	6800 (2800-12000)
Chile	2016	Female	65 to 69	76% (60-88%)	24% (12-40%)	2.6 (1.5-4.1)	4% (1.7-6.5%)	6.3% (3.3-9.8%)	230 (110-410)	5800 (2500-10000)
Chile	2016	Female	70 to 74	73% (56-85%)	27% (15-44%)	2.3 (1.3-3.8)	3.3% (0.9-6.7%)	4.9% (1.7-9.3%)	240 (75-480)	4700 (1200-9800)
Chile	2016	Female	75 to 79	70% (54-84%)	30% (16-46%)	2.1 (1.1-3.6)	2.7% (0.11-6.3%)	3.7% (0.34-8.4%)	220 (20-520)	3400 (130-8400)
Chile	2016	Female	80 to 84	68% (50-82%)	32% (18-50%)	2 (1.1-3.3)	2.4% (0.031-5.6%)	3.1% (0.17-7.4%)	240 (12-590)	2800 (45-7000)
Chile	2016	Female	85 to 89	66% (49-80%)	34% (20-51%)	1.9 (1-3.3)	2.2% (-0.1-5.6%)	2.8% (-0.13-7%)	230 (-13-590)	1900 (-90-4800)
Chile	2016	Female	90 to 94	66% (50-80%)	34% (20-50%)	1.9 (0.97-3.3)	2.2% (-0.4-6.2%)	2.6% (-0.61-7.4%)	160 (-38-470)	940 (-170-2600)
Chile	2016	Female	95 plus	67% (46-82%)	33% (18-54%)	2 (0.88-3.7)	2.5% (-1.1-7.8%)	2.8% (-1.4-9.1%)	100 (-54-340)	360 (-160-1100)
Chile	2016	Male	All Ages	85% (81-89%)	15% (11-19%)	2.9 (2.2-3.7)	7.8% (5.9-9.9%)	7.4% (5-10%)	4300 (2500-6600)	170000 (120000-240000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chile	2016	Male	15 to 19	82% (69-90%)	18% (9.5-31%)	2.5 (1.4-4.1)	7.1% (4.5-10%)	13% (6.7-20%)	53 (24-89)	5900 (3600-8900)
Chile	2016	Male	20 to 24	87% (77-93%)	13% (6.9-23%)	3.9 (2.3-6.2)	14% (9.8-19%)	20% (11-30%)	140 (74-230)	15000 (9900-22000)
Chile	2016	Male	25 to 29	89% (80-94%)	11% (5.7-20%)	4.4 (2.8-6.5)	14% (10-18%)	21% (13-29%)	160 (93-270)	16000 (11000-22000)
Chile	2016	Male	30 to 34	90% (81-95%)	10% (4.8-19%)	4.5 (2.8-6.7)	12% (8.6-16%)	20% (13-28%)	170 (98-280)	14000 (9200-20000)
Chile	2016	Male	35 to 39	91% (83-96%)	9.4% (4.3-17%)	4 (2.4-6.2)	11% (8.1-15%)	19% (12-26%)	180 (100-290)	13000 (8400-19000)
Chile	2016	Male	40 to 44	91% (83-96%)	9% (4-17%)	3.8 (2.2-6)	11% (7.9-15%)	18% (13-24%)	240 (140-380)	14000 (9300-21000)
Chile	2016	Male	45 to 49	91% (82-96%)	8.8% (4-18%)	3.7 (2.2-5.5)	11% (7.8-15%)	17% (12-23%)	340 (200-540)	17000 (11000-26000)
Chile	2016	Male	50 to 54	91% (83-96%)	8.9% (4.1-17%)	3.5 (2.1-5.4)	10% (7.1-14%)	15% (11-20%)	430 (250-680)	18000 (11000-28000)
Chile	2016	Male	55 to 59	91% (83-96%)	9.3% (4.3-17%)	3.5 (2.1-5.2)	9% (6.1-13%)	13% (8.8-18%)	460 (270-730)	17000 (10000-25000)
Chile	2016	Male	60 to 64	90% (81-96%)	10% (4.4-19%)	3.2 (1.8-5.1)	7.6% (4.6-11%)	10% (6.4-15%)	480 (260-770)	14000 (7900-23000)
Chile	2016	Male	65 to 69	89% (80-95%)	11% (4.8-20%)	2.9 (1.9-4.4)	6.2% (3.6-9.6%)	7.9% (4.6-12%)	460 (250-770)	11000 (6100-19000)
Chile	2016	Male	70 to 74	88% (79-94%)	12% (5.6-21%)	2.6 (1.4-4.1)	4.7% (2.3-8.3%)	5.8% (2.8-10%)	390 (170-730)	7800 (3400-14000)
Chile	2016	Male	75 to 79	87% (77-94%)	13% (6.4-23%)	2.3 (1.2-3.9)	3.5% (1.7-7%)	4.1% (1.2-8.3%)	300 (82-620)	4600 (1400-9600)
Chile	2016	Male	80 to 84	85% (73-92%)	15% (7.6-27%)	2.1 (1.2-3.3)	2.6% (0.65-5.4%)	3% (0.73-6.3%)	230 (56-500)	2700 (680-5700)
Chile	2016	Male	85 to 89	82% (68-91%)	18% (9.2-32%)	1.9 (1-3.1)	2.1% (0.35-4.6%)	2.3% (0.25-5.4%)	140 (15-330)	1200 (200-2600)
Chile	2016	Male	90 to 94	81% (67-91%)	19% (8.8-33%)	1.8 (0.91-3.1)	1.7% (-0.015-4.6%)	1.8% (-0.26-5.1%)	59 (-8.2-170)	360 (-3.4-970)
Chile	2016	Male	95 plus	81% (64-93%)	19% (7.2-36%)	1.8 (0.82-3.3)	1.7% (-0.58-5.8%)	1.7% (-0.96-6.5%)	23 (-13-90)	91 (-30-300)
Uruguay	1990	Female	All Ages	55% (49-61%)	45% (39-51%)	2.2 (2-2.5)	3.5% (2.3-4.8%)	3.9% (1.4-6.5%)	530 (200-890)	16000 (10000-22000)
Uruguay	1990	Female	15 to 19	61% (44-75%)	39% (25-56%)	2.2 (1.3-3.5)	3.3% (1.9-5%)	8.4% (4.5-14%)	5.6 (3-9.2)	510 (300-770)
Uruguay	1990	Female	20 to 24	72% (58-82%)	28% (18-42%)	3.6 (2.2-5.3)	4.7% (3.2-6.6%)	12% (7.5-18%)	7.2 (4.3-10)	690 (460-980)
Uruguay	1990	Female	25 to 29	72% (57-82%)	28% (18-43%)	3.9 (2.6-5.6)	4.8% (3.4-6.4%)	13% (8.8-18%)	8.8 (5.9-12)	750 (520-1000)
Uruguay	1990	Female	30 to 34	69% (56-80%)	31% (20-44%)	3.9 (2.5-5.5)	5.1% (3.5-6.8%)	13% (8.5-17%)	13 (8.3-17)	890 (620-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	1990	Female	35 to 39	66% (52-78%)	34% (22-48%)	3.4 (2.2-5)	5.2% (3.5-7.4%)	12% (7.7-16%)	16 (10-22)	960 (640-1300)
Uruguay	1990	Female	40 to 44	64% (50-76%)	36% (24-50%)	3.1 (1.8-4.6)	5.4% (3.3-7.9%)	11% (6.9-16%)	21 (13-30)	1100 (660-1500)
Uruguay	1990	Female	45 to 49	63% (48-75%)	37% (25-52%)	3 (1.7-4.5)	5.8% (3.5-8.6%)	11% (6.6-16%)	29 (17-42)	1300 (780-1900)
Uruguay	1990	Female	50 to 54	62% (48-74%)	38% (26-52%)	2.9 (1.8-4.4)	6.1% (3.4-9.3%)	10% (5.9-15%)	39 (23-58)	1500 (870-2300)
Uruguay	1990	Female	55 to 59	61% (48-74%)	39% (26-52%)	2.9 (1.8-4.3)	6.2% (3.5-9.5%)	9.5% (5.5-14%)	56 (32-83)	1900 (1000-2800)
Uruguay	1990	Female	60 to 64	58% (43-71%)	42% (29-57%)	2.6 (1.5-4.1)	5.2% (2.1-8.8%)	7.4% (3.2-13%)	61 (26-110)	1800 (730-3100)
Uruguay	1990	Female	65 to 69	52% (38-66%)	48% (34-62%)	2.1 (1.3-3.2)	4.1% (1.6-7.1%)	5.5% (2.3-9.4%)	61 (25-100)	1500 (580-2600)
Uruguay	1990	Female	70 to 74	47% (34-61%)	53% (39-66%)	1.8 (0.99-2.8)	3% (0.16-6.4%)	3.8% (0.2-8.2%)	56 (3.1-120)	1100 (58-2400)
Uruguay	1990	Female	75 to 79	42% (29-57%)	58% (43-71%)	1.6 (0.84-2.7)	2.2% (-0.79-5.8%)	2.6% (-1-7%)	53 (-21-140)	810 (-310-2200)
Uruguay	1990	Female	80 to 84	40% (27-53%)	60% (47-73%)	1.4 (0.78-2.3)	1.8% (-0.73-4.8%)	2.1% (-1-5.7%)	47 (-23-130)	530 (-210-1400)
Uruguay	1990	Female	85 to 89	39% (26-53%)	61% (47-74%)	1.3 (0.72-2.1)	1.6% (-0.84-4.6%)	1.7% (-1.1-5.2%)	36 (-23-110)	290 (-150-830)
Uruguay	1990	Female	90 to 94	39% (27-53%)	61% (47-73%)	1.3 (0.67-2.2)	1.4% (-1-4.8%)	1.5% (-1.3-5.5%)	15 (-14-57)	86 (-62-300)
Uruguay	1990	Female	95 plus	41% (26-59%)	59% (41-74%)	1.4 (0.6-2.6)	1.4% (-1.5-5.7%)	1.4% (-2-6.5%)	5.3 (-7.6-24)	20 (-22-85)
Uruguay	1990	Male	All Ages	65% (58-70%)	35% (30-42%)	2.3 (2-2.6)	5% (3.6-6.7%)	4.6% (2.6-7.2%)	760 (420-1200)	27000 (19000-37000)
Uruguay	1990	Male	15 to 19	68% (52-80%)	32% (20-48%)	2.2 (1.3-3.5)	6% (3.8-8.9%)	10% (6.1-16%)	13 (7.7-20)	1100 (710-1600)
Uruguay	1990	Male	20 to 24	76% (62-85%)	24% (15-38%)	3.6 (2.3-5.4)	11% (7.6-15%)	16% (10-24%)	24 (15-35)	2000 (1300-2700)
Uruguay	1990	Male	25 to 29	76% (63-85%)	24% (15-37%)	4 (2.7-5.6)	11% (7.6-15%)	17% (11-24%)	27 (18-37)	2000 (1400-2600)
Uruguay	1990	Male	30 to 34	76% (62-85%)	24% (15-38%)	3.9 (2.6-5.7)	9.7% (6.4-13%)	15% (9.3-21%)	27 (17-39)	1800 (1200-2500)
Uruguay	1990	Male	35 to 39	75% (62-85%)	25% (15-38%)	3.5 (2.2-5.1)	8.4% (5.4-12%)	13% (7.7-18%)	27 (16-39)	1600 (1000-2300)
Uruguay	1990	Male	40 to 44	75% (63-85%)	25% (15-37%)	3.3 (1.9-4.8)	8.2% (4.9-12%)	12% (6.6-17%)	36 (20-53)	1900 (1100-2600)
Uruguay	1990	Male	45 to 49	76% (62-85%)	24% (15-38%)	3.3 (1.9-5)	8.2% (5-12%)	11% (6.3-17%)	52 (30-80)	2300 (1400-3500)
Uruguay	1990	Male	50 to 54	75% (63-85%)	25% (15-37%)	3.2 (1.9-4.8)	8.1% (4.6-12%)	10% (5.6-16%)	75 (41-120)	2900 (1700-4400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	1990	Male	55 to 59	74% (62-84%)	26% (16-38%)	3 (1.9-4.4)	7.3% (3.9-11%)	8.7% (4.7-14%)	110 (58-170)	3600 (2000-5600)
Uruguay	1990	Male	60 to 64	72% (60-82%)	28% (18-40%)	2.8 (1.6-4.2)	6% (2.7-10%)	7% (3.1-12%)	120 (53-200)	3300 (1500-5600)
Uruguay	1990	Male	65 to 69	69% (56-80%)	31% (20-44%)	2.5 (1.6-3.6)	4.2% (1.6-7.8%)	4.8% (1.7-9.1%)	94 (34-180)	2200 (850-4200)
Uruguay	1990	Male	70 to 74	66% (54-77%)	34% (23-46%)	2.2 (1.2-3.5)	2.9% (0.23-6.7%)	3.2% (0.073-7.5%)	68 (1.6-160)	1300 (100-3100)
Uruguay	1990	Male	75 to 79	63% (49-76%)	37% (24-51%)	2 (1-3.2)	1.9% (-0.66-5.5%)	2% (-0.95-6.1%)	47 (-22-140)	730 (-260-2100)
Uruguay	1990	Male	80 to 84	60% (43-74%)	40% (26-57%)	1.7 (0.93-2.6)	1.3% (-0.77-3.9%)	1.3% (-1.1-4.3%)	25 (-20-81)	300 (-180-910)
Uruguay	1990	Male	85 to 89	55% (39-70%)	45% (30-61%)	1.4 (0.73-2.2)	0.96% (-0.71-3.1%)	0.93% (-0.99-3.4%)	12 (-13-43)	110 (-79-340)
Uruguay	1990	Male	90 to 94	54% (38-69%)	46% (31-62%)	1.3 (0.66-2.3)	0.66% (-1-2.9%)	0.58% (-1.3-3.1%)	3 (-6.9-16)	20 (-31-89)
Uruguay	1990	Male	95 plus	55% (35-74%)	45% (26-65%)	1.3 (0.53-2.5)	0.58% (-1.3-3.8%)	0.43% (-1.8-4.1%)	0.58 (-2.5-5.6)	3.1 (-7-20)
Uruguay	1995	Female	All Ages	56% (50-61%)	44% (39-50%)	2.2 (1.9-2.5)	3.4% (2.2-4.7%)	3.8% (1.5-6.4%)	530 (210-900)	15000 (10000-21000)
Uruguay	1995	Female	15 to 19	61% (46-75%)	39% (25-54%)	2.1 (1.2-3.3)	3.1% (1.8-4.8%)	8.2% (4.4-13%)	5.4 (2.9-8.6)	500 (290-750)
Uruguay	1995	Female	20 to 24	72% (59-82%)	28% (18-41%)	3.5 (2.2-5)	4.4% (2.9-6.2%)	12% (7.2-17%)	7.3 (4.4-11)	700 (460-980)
Uruguay	1995	Female	25 to 29	72% (58-82%)	28% (18-42%)	3.8 (2.5-5.3)	4.8% (3.3-6.5%)	13% (8.4-17%)	9.2 (6.2-13)	760 (520-1000)
Uruguay	1995	Female	30 to 34	70% (55-80%)	30% (20-45%)	3.8 (2.5-5.4)	4.8% (3.3-6.5%)	12% (8.4-17%)	12 (8.3-17)	870 (590-1200)
Uruguay	1995	Female	35 to 39	66% (52-78%)	34% (22-48%)	3.3 (2.1-4.8)	4.9% (3.2-6.9%)	12% (7.5-16%)	16 (10-22)	970 (630-1300)
Uruguay	1995	Female	40 to 44	64% (51-77%)	36% (23-49%)	3 (1.8-4.5)	5.3% (3.1-7.6%)	11% (6.6-16%)	21 (13-30)	1100 (660-1600)
Uruguay	1995	Female	45 to 49	63% (48-75%)	37% (25-52%)	2.9 (1.7-4.4)	5.6% (3.3-8.3%)	11% (6.3-16%)	28 (17-41)	1300 (760-1800)
Uruguay	1995	Female	50 to 54	63% (49-74%)	37% (26-51%)	2.9 (1.7-4.3)	5.8% (3.2-8.8%)	9.9% (5.7-15%)	37 (21-54)	1400 (820-2200)
Uruguay	1995	Female	55 to 59	62% (49-74%)	38% (26-51%)	2.8 (1.7-4.2)	5.9% (3.2-8.9%)	9.4% (5.4-14%)	47 (27-70)	1600 (900-2400)
Uruguay	1995	Female	60 to 64	59% (44-72%)	41% (28-56%)	2.6 (1.5-4.1)	5% (2-8.7%)	7.3% (3.2-12%)	57 (25-95)	1700 (700-2800)
Uruguay	1995	Female	65 to 69	53% (39-67%)	47% (33-61%)	2.1 (1.3-3.2)	4% (1.5-6.9%)	5.5% (2.3-9.4%)	62 (26-110)	1500 (590-2600)
Uruguay	1995	Female	70 to 74	48% (35-61%)	52% (39-65%)	1.8 (0.97-2.8)	2.9% (0.23-6.1%)	3.8% (0.35-7.9%)	58 (5.4-120)	1100 (92-2400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	1995	Female	75 to 79	43% (30-58%)	57% (42-70%)	1.6 (0.81-2.7)	2.2% (-0.69-5.7%)	2.6% (-0.86-7%)	51 (-17-140)	780 (-240-2000)
Uruguay	1995	Female	80 to 84	40% (27-55%)	60% (45-73%)	1.4 (0.78-2.2)	1.9% (-0.6-4.9%)	2.2% (-0.7-5.8%)	52 (-17-140)	590 (-190-1500)
Uruguay	1995	Female	85 to 89	39% (27-53%)	61% (47-73%)	1.3 (0.72-2.1)	1.6% (-0.6-4.3%)	1.8% (-0.79-5.1%)	42 (-18-120)	340 (-120-920)
Uruguay	1995	Female	90 to 94	40% (27-54%)	60% (46-73%)	1.3 (0.68-2.2)	1.5% (-0.75-4.7%)	1.6% (-1.1-5.4%)	19 (-14-64)	110 (-57-340)
Uruguay	1995	Female	95 plus	42% (26-60%)	58% (40-74%)	1.4 (0.61-2.6)	1.5% (-1.2-5.5%)	1.6% (-1.6-6.4%)	7.1 (-7.3-28)	27 (-22-99)
Uruguay	1995	Male	All Ages	65% (59-71%)	35% (29-41%)	2.6 (2.3-3)	6.1% (4.6-7.8%)	5.9% (3.7-8.6%)	1000 (630-1500)	34000 (26000-44000)
Uruguay	1995	Male	15 to 19	68% (53-81%)	32% (19-47%)	2.5 (1.4-3.9)	6.9% (4.4-10%)	11% (6.8-17%)	18 (10-27)	1500 (910-2100)
Uruguay	1995	Male	20 to 24	76% (62-85%)	24% (15-38%)	4.1 (2.5-5.9)	12% (8.2-17%)	18% (12-26%)	33 (21-48)	2600 (1800-3600)
Uruguay	1995	Male	25 to 29	77% (64-86%)	23% (14-36%)	4.5 (2.9-6.3)	12% (8.6-17%)	19% (13-26%)	34 (23-47)	2400 (1700-3200)
Uruguay	1995	Male	30 to 34	76% (62-85%)	24% (15-38%)	4.4 (2.9-6.3)	11% (7.4-15%)	17% (11-23%)	33 (21-45)	2100 (1500-2900)
Uruguay	1995	Male	35 to 39	75% (63-85%)	25% (15-37%)	3.9 (2.5-5.7)	9.4% (6.3-13%)	15% (9.2-20%)	34 (21-48)	2000 (1300-2800)
Uruguay	1995	Male	40 to 44	75% (63-85%)	25% (15-37%)	3.7 (2.3-5.5)	9.6% (5.9-13%)	14% (8.2-19%)	44 (26-62)	2300 (1400-3100)
Uruguay	1995	Male	45 to 49	76% (62-85%)	24% (15-38%)	3.7 (2.2-5.5)	9.7% (6-14%)	13% (7.9-19%)	64 (39-94)	2800 (1800-4100)
Uruguay	1995	Male	50 to 54	76% (64-85%)	24% (15-36%)	3.6 (2.2-5.5)	9.7% (5.8-14%)	12% (7.2-18%)	89 (52-130)	3400 (2100-5100)
Uruguay	1995	Male	55 to 59	74% (63-84%)	26% (16-37%)	3.5 (2.2-5.1)	9.1% (5.5-13%)	11% (6.5-16%)	120 (71-170)	3900 (2400-5700)
Uruguay	1995	Male	60 to 64	72% (60-83%)	28% (17-40%)	3.2 (1.9-4.9)	7.5% (4-12%)	8.8% (4.5-14%)	140 (75-230)	4100 (2100-6500)
Uruguay	1995	Male	65 to 69	69% (57-80%)	31% (20-43%)	2.8 (1.8-4.1)	5.6% (2.7-9.4%)	6.4% (3-11%)	140 (64-230)	3200 (1500-5400)
Uruguay	1995	Male	70 to 74	67% (54-77%)	33% (23-46%)	2.5 (1.4-3.8)	4% (1.1-8%)	4.5% (1.1-9.1%)	100 (25-210)	2000 (540-4000)
Uruguay	1995	Male	75 to 79	64% (50-78%)	36% (22-50%)	2.2 (1.2-3.6)	2.9% (0.065-6.8%)	3.1% (-0.011-7.5%)	70 (-0.26-170)	1100 (22-2500)
Uruguay	1995	Male	80 to 84	60% (43-75%)	40% (25-57%)	1.9 (1.1-2.9)	2.1% (-0.1-5%)	2.2% (-0.31-5.5%)	46 (-6.4-110)	530 (-26-1300)
Uruguay	1995	Male	85 to 89	56% (39-70%)	44% (30-61%)	1.6 (0.87-2.5)	1.7% (-0.11-4.3%)	1.8% (-0.29-4.7%)	26 (-4.3-68)	210 (-13-530)
Uruguay	1995	Male	90 to 94	54% (38-69%)	46% (31-62%)	1.5 (0.77-2.5)	1.3% (-0.33-4%)	1.4% (-0.56-4.4%)	8.2 (-3.4-27)	48 (-12-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	1995	Male	95 plus	55% (35-74%)	45% (26-65%)	1.5 (0.61-2.8)	1.3% (-0.72-4.7%)	1.2% (-1.1-5.2%)	2.1 (-1.7-8.7)	8.4 (-4.7-31)
Uruguay	2000	Female	All Ages	54% (49-59%)	46% (41-51%)	2.2 (1.9-2.4)	3.5% (2.4-4.8%)	4% (1.9-6.4%)	570 (280-920)	16000 (11000-21000)
Uruguay	2000	Female	15 to 19	65% (51-77%)	35% (23-49%)	2.3 (1.4-3.5)	3.1% (1.9-4.7%)	9.1% (4.9-14%)	4.7 (2.5-7.4)	440 (270-650)
Uruguay	2000	Female	20 to 24	73% (60-83%)	27% (17-40%)	3.6 (2.3-5.2)	4.4% (2.9-6.2%)	13% (7.6-18%)	7.6 (4.6-11)	730 (480-990)
Uruguay	2000	Female	25 to 29	72% (60-81%)	28% (19-40%)	3.9 (2.6-5.3)	4.6% (3.2-6%)	13% (8.8-17%)	8.9 (6-12)	760 (540-1000)
Uruguay	2000	Female	30 to 34	69% (57-79%)	31% (21-43%)	3.8 (2.6-5.3)	4.5% (3.2-6%)	12% (8.6-16%)	10 (7.3-14)	760 (540-990)
Uruguay	2000	Female	35 to 39	64% (52-76%)	36% (24-48%)	3.2 (2.1-4.7)	4.5% (3-6.3%)	12% (7.7-16%)	14 (9.4-19)	880 (590-1200)
Uruguay	2000	Female	40 to 44	61% (49-73%)	39% (27-51%)	2.9 (1.7-4.3)	5% (3.1-7.2%)	11% (6.7-15%)	22 (13-30)	1100 (700-1600)
Uruguay	2000	Female	45 to 49	60% (48-72%)	40% (28-52%)	2.8 (1.7-4.2)	5.5% (3.3-8.1%)	11% (6.7-16%)	29 (18-42)	1300 (790-1900)
Uruguay	2000	Female	50 to 54	60% (47-71%)	40% (29-53%)	2.8 (1.7-4.2)	5.9% (3.4-8.9%)	10% (6.1-15%)	38 (23-57)	1500 (890-2300)
Uruguay	2000	Female	55 to 59	60% (47-71%)	40% (29-53%)	2.8 (1.7-4.2)	6% (3.4-9.1%)	9.6% (5.7-14%)	48 (29-71)	1600 (950-2500)
Uruguay	2000	Female	60 to 64	55% (43-68%)	45% (32-57%)	2.4 (1.4-3.9)	5% (2.4-8.3%)	7.5% (3.8-12%)	52 (26-84)	1500 (710-2500)
Uruguay	2000	Female	65 to 69	49% (36-63%)	51% (37-64%)	1.9 (1.2-2.9)	4% (1.8-6.7%)	5.7% (2.7-9.2%)	60 (28-98)	1500 (650-2400)
Uruguay	2000	Female	70 to 74	43% (31-56%)	57% (44-69%)	1.6 (0.88-2.6)	3% (0.78-5.9%)	4% (1.2-7.7%)	61 (18-120)	1200 (300-2300)
Uruguay	2000	Female	75 to 79	39% (28-54%)	61% (46-72%)	1.5 (0.76-2.5)	2.3% (-0.069-5.3%)	2.9% (-0.054-6.7%)	60 (-1.1-130)	910 (-25-2000)
Uruguay	2000	Female	80 to 84	38% (26-52%)	62% (48-74%)	1.4 (0.74-2.2)	2.1% (0.03-4.7%)	2.5% (0.011-5.6%)	59 (0.25-130)	660 (9.3-1500)
Uruguay	2000	Female	85 to 89	38% (26-52%)	62% (48-74%)	1.3 (0.72-2.1)	1.9% (-0.18-4.5%)	2.2% (-0.24-5.2%)	53 (-5.7-130)	420 (-39-1000)
Uruguay	2000	Female	90 to 94	39% (27-53%)	61% (47-73%)	1.3 (0.69-2.3)	1.8% (-0.51-4.9%)	2% (-0.74-5.8%)	33 (-12-94)	180 (-53-500)
Uruguay	2000	Female	95 plus	42% (26-59%)	58% (41-74%)	1.4 (0.6-2.6)	1.8% (-0.83-5.8%)	2% (-1.1-6.7%)	12 (-6.4-40)	43 (-19-140)
Uruguay	2000	Male	All Ages	71% (65-75%)	29% (25-35%)	2.6 (2.3-2.9)	6.2% (4.7-7.9%)	5.8% (3.7-8.3%)	960 (620-1400)	33000 (25000-43000)
Uruguay	2000	Male	15 to 19	76% (64-85%)	24% (15-36%)	2.6 (1.5-3.9)	7.1% (4.4-10%)	12% (7.1-18%)	16 (9.4-24)	1300 (840-1900)
Uruguay	2000	Male	20 to 24	81% (71-88%)	19% (12-29%)	4.1 (2.6-5.8)	13% (8.4-17%)	19% (12-27%)	37 (23-52)	2900 (1900-3900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	2000	Male	25 to 29	82% (74-88%)	18% (12-26%)	4.5 (3-6.1)	12% (8.6-16%)	19% (12-26%)	36 (24-49)	2600 (1800-3400)
Uruguay	2000	Male	30 to 34	81% (73-88%)	19% (12-27%)	4.4 (3-6.1)	10% (7.3-14%)	17% (11-23%)	30 (20-41)	2000 (1400-2600)
Uruguay	2000	Male	35 to 39	79% (71-86%)	21% (14-29%)	3.7 (2.4-5.3)	9.3% (6.4-13%)	14% (9.4-20%)	34 (22-47)	2000 (1300-2700)
Uruguay	2000	Male	40 to 44	78% (70-87%)	22% (13-30%)	3.6 (2.2-5.2)	9.2% (5.9-13%)	14% (8.5-19%)	41 (25-59)	2200 (1400-3000)
Uruguay	2000	Male	45 to 49	80% (72-87%)	20% (13-28%)	3.6 (2.2-5.2)	9.5% (6-14%)	13% (8.2-18%)	61 (39-88)	2800 (1800-3900)
Uruguay	2000	Male	50 to 54	80% (72-87%)	20% (13-28%)	3.5 (2.2-5.2)	9.5% (5.8-14%)	12% (7.4-18%)	87 (53-130)	3300 (2100-4900)
Uruguay	2000	Male	55 to 59	78% (69-85%)	22% (15-31%)	3.3 (2.1-4.9)	8.7% (5.3-13%)	11% (6.6-16%)	110 (67-160)	3600 (2300-5300)
Uruguay	2000	Male	60 to 64	76% (67-83%)	24% (17-33%)	3.1 (1.8-4.6)	7.5% (4.2-12%)	8.9% (5-14%)	120 (70-200)	3500 (1900-5500)
Uruguay	2000	Male	65 to 69	74% (64-82%)	26% (18-36%)	2.8 (1.8-4)	5.5% (2.8-9.1%)	6.4% (3.1-11%)	130 (63-210)	3000 (1500-5000)
Uruguay	2000	Male	70 to 74	71% (61-80%)	29% (20-39%)	2.4 (1.4-3.7)	3.9% (1.2-7.4%)	4.4% (1.3-8.5%)	100 (31-200)	2000 (630-3800)
Uruguay	2000	Male	75 to 79	69% (57-80%)	31% (20-43%)	2.2 (1.2-3.5)	2.9% (0.32-6.4%)	3.2% (0.26-7.2%)	76 (6.1-170)	1200 (130-2600)
Uruguay	2000	Male	80 to 84	63% (47-77%)	37% (23-53%)	1.8 (1-2.7)	2% (0.04-4.4%)	2.1% (-0.08-4.9%)	43 (-1.6-100)	490 (10-1100)
Uruguay	2000	Male	85 to 89	56% (40-71%)	44% (29-60%)	1.4 (0.78-2.3)	1.6% (0.09-3.5%)	1.7% (-0.046-3.9%)	25 (-0.69-58)	200 (12-460)
Uruguay	2000	Male	90 to 94	54% (39-70%)	46% (30-61%)	1.3 (0.68-2.3)	1.3% (-0.17-3.4%)	1.3% (-0.31-3.7%)	9.7 (-2.3-27)	56 (-7.2-150)
Uruguay	2000	Male	95 plus	56% (37-74%)	44% (26-63%)	1.3 (0.55-2.5)	1.3% (-0.39-4.2%)	1.3% (-0.66-4.6%)	2.7 (-1.3-9.3)	11 (-3.2-33)
Uruguay	2005	Female	All Ages	53% (48-58%)	47% (42-52%)	2 (1.7-2.2)	3% (2-4.2%)	3.3% (1.5-5.3%)	490 (220-810)	13000 (9000-18000)
Uruguay	2005	Female	15 to 19	65% (51-76%)	35% (24-49%)	2.1 (1.3-3.2)	2.8% (1.7-4.4%)	8.5% (4.6-14%)	4.2 (2.3-6.7)	400 (250-600)
Uruguay	2005	Female	20 to 24	73% (60-82%)	27% (18-40%)	3.3 (2.1-4.8)	4.1% (2.7-5.9%)	12% (7.2-17%)	6.2 (3.6-9.1)	610 (400-860)
Uruguay	2005	Female	25 to 29	71% (60-81%)	29% (19-40%)	3.5 (2.3-4.9)	4.1% (2.8-5.5%)	12% (7.6-16%)	7.5 (4.9-10)	670 (460-890)
Uruguay	2005	Female	30 to 34	68% (56-78%)	32% (22-44%)	3.5 (2.3-4.9)	4.2% (3-5.7%)	11% (7.6-15%)	11 (7.2-14)	760 (530-1000)
Uruguay	2005	Female	35 to 39	63% (51-74%)	37% (26-49%)	2.9 (1.9-4.3)	3.8% (2.6-5.4%)	10% (6.7-14%)	11 (7.1-15)	690 (460-940)
Uruguay	2005	Female	40 to 44	60% (48-72%)	40% (28-52%)	2.6 (1.6-3.8)	4.4% (2.7-6.3%)	9.7% (5.9-14%)	19 (11-26)	980 (600-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	2005	Female	45 to 49	59% (47-71%)	41% (29-53%)	2.5 (1.5-3.8)	4.7% (2.8-6.9%)	9.4% (5.7-14%)	25 (16-37)	1200 (720-1700)
Uruguay	2005	Female	50 to 54	59% (46-71%)	41% (29-54%)	2.5 (1.5-3.8)	5.1% (2.9-7.9%)	8.9% (5.2-13%)	35 (21-52)	1400 (790-2100)
Uruguay	2005	Female	55 to 59	59% (47-70%)	41% (30-53%)	2.5 (1.6-3.7)	5.1% (2.8-7.7%)	8.3% (4.7-12%)	43 (24-64)	1500 (800-2200)
Uruguay	2005	Female	60 to 64	54% (42-67%)	46% (33-58%)	2.2 (1.3-3.4)	4.3% (1.9-7.3%)	6.5% (3.1-11%)	44 (21-73)	1300 (570-2200)
Uruguay	2005	Female	65 to 69	48% (36-62%)	52% (38-64%)	1.7 (1.1-2.6)	3.3% (1.5-5.6%)	4.8% (2.3-7.9%)	46 (22-76)	1100 (500-1800)
Uruguay	2005	Female	70 to 74	42% (30-56%)	58% (44-70%)	1.4 (0.79-2.3)	2.5% (0.59-4.8%)	3.4% (0.92-6.5%)	49 (13-94)	950 (220-1900)
Uruguay	2005	Female	75 to 79	38% (27-53%)	62% (47-73%)	1.3 (0.7-2.2)	1.9% (-0.1-4.5%)	2.4% (-0.077-5.6%)	53 (-1.7-120)	790 (-41-1900)
Uruguay	2005	Female	80 to 84	37% (25-51%)	63% (49-75%)	1.2 (0.68-1.9)	1.6% (-0.15-3.8%)	2% (-0.17-4.6%)	51 (-4.6-120)	580 (-53-1400)
Uruguay	2005	Female	85 to 89	37% (26-51%)	63% (49-74%)	1.2 (0.65-1.9)	1.5% (-0.23-3.7%)	1.8% (-0.29-4.4%)	46 (-7.8-120)	370 (-56-900)
Uruguay	2005	Female	90 to 94	38% (26-52%)	62% (48-74%)	1.2 (0.61-2.1)	1.5% (-0.46-4.2%)	1.7% (-0.64-4.9%)	31 (-12-91)	170 (-54-480)
Uruguay	2005	Female	95 plus	41% (26-58%)	59% (42-74%)	1.3 (0.56-2.4)	1.5% (-0.8-5%)	1.7% (-1-5.7%)	13 (-8-44)	47 (-25-150)
Uruguay	2005	Male	All Ages	70% (65-75%)	30% (25-35%)	2.4 (2.1-2.8)	5.6% (4.3-7.1%)	5% (3.2-7.3%)	840 (530-1200)	29000 (22000-37000)
Uruguay	2005	Male	15 to 19	76% (64-85%)	24% (15-36%)	2.4 (1.4-3.7)	6.6% (4-9.6%)	12% (6.6-18%)	14 (7.7-21)	1200 (730-1700)
Uruguay	2005	Male	20 to 24	81% (71-88%)	19% (12-29%)	3.8 (2.4-5.6)	12% (8-17%)	18% (11-26%)	28 (17-40)	2300 (1500-3100)
Uruguay	2005	Male	25 to 29	82% (73-88%)	18% (12-27%)	4.2 (2.8-5.8)	11% (8-15%)	18% (12-25%)	31 (20-43)	2300 (1600-3000)
Uruguay	2005	Male	30 to 34	80% (72-87%)	20% (13-28%)	4.1 (2.7-5.7)	9.6% (6.6-13%)	15% (10-21%)	28 (19-39)	1900 (1300-2600)
Uruguay	2005	Male	35 to 39	78% (71-86%)	22% (14-29%)	3.5 (2.2-5)	8% (5.5-11%)	13% (8.2-18%)	24 (16-35)	1500 (1000-2100)
Uruguay	2005	Male	40 to 44	78% (70-86%)	22% (14-30%)	3.3 (2.1-4.8)	8.2% (5.3-11%)	12% (7.7-17%)	36 (22-51)	1900 (1200-2700)
Uruguay	2005	Male	45 to 49	79% (71-87%)	21% (13-29%)	3.4 (2.1-4.9)	8.6% (5.6-12%)	12% (7.6-17%)	54 (34-79)	2500 (1600-3500)
Uruguay	2005	Male	50 to 54	79% (70-86%)	21% (14-30%)	3.3 (2-4.9)	8.3% (5.1-12%)	11% (6.5-16%)	74 (44-110)	2900 (1800-4300)
Uruguay	2005	Male	55 to 59	77% (68-84%)	23% (16-32%)	3.1 (1.9-4.4)	7.5% (4.5-11%)	9.3% (5.6-14%)	91 (55-140)	3000 (1800-4500)
Uruguay	2005	Male	60 to 64	75% (66-83%)	25% (17-34%)	2.9 (1.7-4.3)	6.5% (3.6-10%)	7.7% (4.2-12%)	100 (57-170)	2900 (1700-4700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	2005	Male	65 to 69	73% (64-82%)	27% (18-36%)	2.6 (1.7-3.7)	5% (2.5-8.1%)	5.8% (3-9.6%)	100 (51-170)	2400 (1200-3900)
Uruguay	2005	Male	70 to 74	71% (61-79%)	29% (21-39%)	2.3 (1.4-3.5)	3.6% (1.2-6.7%)	4.1% (1.3-7.7%)	91 (30-170)	1800 (580-3300)
Uruguay	2005	Male	75 to 79	69% (56-79%)	31% (21-44%)	2.1 (1.1-3.3)	2.7% (0.4-5.8%)	3% (0.35-6.5%)	76 (8.9-170)	1200 (170-2600)
Uruguay	2005	Male	80 to 84	63% (47-77%)	37% (23-53%)	1.7 (0.95-2.6)	1.9% (0.15-4.1%)	2.1% (0.099-4.6%)	47 (2.2-100)	530 (44-1200)
Uruguay	2005	Male	85 to 89	55% (39-70%)	45% (30-61%)	1.3 (0.72-2.1)	1.5% (0.14-3.2%)	1.6% (0.033-3.5%)	25 (0.54-57)	200 (19-450)
Uruguay	2005	Male	90 to 94	54% (38-69%)	46% (31-62%)	1.2 (0.63-2.1)	1.2% (-0.089-3.2%)	1.2% (-0.21-3.4%)	10 (-1.7-28)	58 (-4.4-150)
Uruguay	2005	Male	95 plus	55% (36-74%)	45% (26-64%)	1.2 (0.52-2.3)	1.2% (-0.38-3.9%)	1.2% (-0.56-4.2%)	3 (-1.4-11)	12 (-3.7-38)
Uruguay	2010	Female	All Ages	57% (51-62%)	43% (38-49%)	1.8 (1.5-2.1)	2.5% (1.6-3.6%)	2.5% (0.84-4.6%)	390 (130-710)	11000 (7000-16000)
Uruguay	2010	Female	15 to 19	62% (47-76%)	38% (24-53%)	1.7 (0.96-2.7)	2.5% (1.5-3.8%)	7.2% (3.7-12%)	3.6 (1.9-5.9)	360 (220-550)
Uruguay	2010	Female	20 to 24	73% (60-83%)	27% (17-40%)	2.8 (1.7-4.2)	4.1% (2.6-5.9%)	10% (5.7-16%)	6.5 (3.6-9.9)	640 (400-930)
Uruguay	2010	Female	25 to 29	73% (60-83%)	27% (17-40%)	3.1 (2-4.4)	3.9% (2.7-5.4%)	10% (6.6-15%)	6.7 (4.2-9.7)	610 (400-850)
Uruguay	2010	Female	30 to 34	71% (57-80%)	29% (20-43%)	3.1 (2-4.4)	3.7% (2.5-5.1%)	10% (6.6-14%)	8.8 (5.8-12)	680 (460-930)
Uruguay	2010	Female	35 to 39	67% (53-78%)	33% (22-47%)	2.7 (1.7-4.1)	3.6% (2.4-5.1%)	9.2% (6-13%)	11 (7.1-15)	710 (470-980)
Uruguay	2010	Female	40 to 44	65% (52-76%)	35% (24-48%)	2.5 (1.4-3.6)	3.7% (2.2-5.3%)	9% (5.3-13%)	14 (8.1-20)	750 (450-1100)
Uruguay	2010	Female	45 to 49	64% (50-76%)	36% (24-50%)	2.4 (1.4-3.6)	4.1% (2.4-6.2%)	8.4% (5-12%)	23 (13-33)	1000 (600-1500)
Uruguay	2010	Female	50 to 54	63% (50-74%)	37% (26-50%)	2.3 (1.4-3.6)	4.2% (2.3-6.7%)	7.6% (4.4-12%)	31 (17-47)	1200 (670-1900)
Uruguay	2010	Female	55 to 59	63% (50-74%)	37% (26-50%)	2.3 (1.4-3.5)	4.2% (2.1-6.5%)	6.9% (3.9-11%)	35 (19-53)	1200 (630-1900)
Uruguay	2010	Female	60 to 64	59% (46-72%)	41% (28-54%)	2.1 (1.2-3.4)	3.6% (1.3-6.3%)	5.5% (2.4-9.5%)	38 (17-65)	1100 (430-1900)
Uruguay	2010	Female	65 to 69	54% (39-68%)	46% (32-61%)	1.7 (1-2.7)	2.7% (0.94-4.9%)	4.1% (1.7-7.2%)	38 (15-67)	900 (310-1700)
Uruguay	2010	Female	70 to 74	49% (35-62%)	51% (38-65%)	1.5 (0.79-2.3)	2% (0.12-4.3%)	2.8% (0.37-5.9%)	37 (4.8-77)	710 (44-1500)
Uruguay	2010	Female	75 to 79	44% (31-59%)	56% (41-69%)	1.3 (0.69-2.2)	1.4% (-0.52-3.8%)	1.8% (-0.56-4.8%)	38 (-12-99)	560 (-210-1500)
Uruguay	2010	Female	80 to 84	42% (28-57%)	58% (43-72%)	1.2 (0.65-1.9)	1.1% (-0.5-3.2%)	1.4% (-0.6-4%)	39 (-17-110)	430 (-200-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	2010	Female	85 to 89	41% (28-55%)	59% (45-72%)	1.1 (0.61-1.8)	0.97% (-0.5-2.9%)	1.2% (-0.6-3.6%)	34 (-17-100)	270 (-130-850)
Uruguay	2010	Female	90 to 94	41% (28-55%)	59% (45-72%)	1.1 (0.57-1.9)	0.92% (-0.65-3.2%)	1% (-0.81-3.7%)	21 (-16-76)	120 (-84-410)
Uruguay	2010	Female	95 plus	44% (27-62%)	56% (38-73%)	1.2 (0.52-2.2)	0.97% (-0.99-4%)	1% (-1.3-4.6%)	9 (-11-40)	34 (-35-140)
Uruguay	2010	Male	All Ages	67% (61-72%)	33% (28-39%)	2.4 (2.1-2.8)	5.9% (4.5-7.4%)	5.4% (3.6-7.6%)	890 (590-1300)	30000 (23000-39000)
Uruguay	2010	Male	15 to 19	69% (54-81%)	31% (19-46%)	2.2 (1.3-3.4)	6.8% (4.1-10%)	11% (6.2-17%)	17 (9.6-27)	1400 (840-2100)
Uruguay	2010	Male	20 to 24	76% (64-85%)	24% (15-36%)	3.7 (2.3-5.4)	12% (7.9-17%)	18% (11-26%)	36 (21-51)	2800 (1800-3800)
Uruguay	2010	Male	25 to 29	77% (64-86%)	23% (14-36%)	4.1 (2.7-5.6)	12% (8.3-17%)	19% (12-26%)	35 (22-49)	2500 (1700-3300)
Uruguay	2010	Male	30 to 34	77% (64-86%)	23% (14-36%)	4 (2.6-5.7)	10% (6.8-14%)	16% (10-23%)	32 (21-46)	2200 (1500-3000)
Uruguay	2010	Male	35 to 39	76% (64-85%)	24% (15-36%)	3.5 (2.2-5)	8.4% (5.7-12%)	13% (8.6-19%)	31 (20-44)	1900 (1300-2600)
Uruguay	2010	Male	40 to 44	76% (64-85%)	24% (15-36%)	3.3 (2-4.8)	8.2% (5.4-11%)	13% (8-18%)	34 (21-48)	1800 (1200-2500)
Uruguay	2010	Male	45 to 49	76% (64-85%)	24% (15-36%)	3.3 (2-4.9)	8.4% (5.5-12%)	12% (7.5-18%)	52 (33-76)	2400 (1600-3500)
Uruguay	2010	Male	50 to 54	76% (65-85%)	24% (15-35%)	3.2 (2-4.8)	8.2% (5.2-12%)	11% (6.8-16%)	73 (46-110)	2900 (1900-4300)
Uruguay	2010	Male	55 to 59	75% (63-84%)	25% (16-37%)	3.1 (2-4.5)	7.7% (4.9-11%)	9.7% (6.1-14%)	92 (57-130)	3100 (1900-4400)
Uruguay	2010	Male	60 to 64	73% (61-83%)	27% (17-39%)	2.8 (1.7-4.4)	6.5% (3.7-10%)	7.9% (4.5-12%)	100 (58-160)	2900 (1700-4500)
Uruguay	2010	Male	65 to 69	70% (58-80%)	30% (20-42%)	2.5 (1.6-3.7)	5.2% (2.9-8.3%)	6.1% (3.4-9.7%)	100 (59-170)	2500 (1400-4000)
Uruguay	2010	Male	70 to 74	68% (55-78%)	32% (22-45%)	2.3 (1.3-3.5)	3.9% (1.7-7%)	4.4% (1.9-7.9%)	90 (37-160)	1700 (730-3100)
Uruguay	2010	Male	75 to 79	65% (49-78%)	35% (22-51%)	2 (1.1-3.3)	2.9% (0.8-5.9%)	3.3% (0.85-6.7%)	82 (21-170)	1200 (350-2500)
Uruguay	2010	Male	80 to 84	61% (45-76%)	39% (24-55%)	1.7 (0.97-2.6)	2.2% (0.57-4.4%)	2.4% (0.59-4.8%)	59 (14-120)	670 (180-1300)
Uruguay	2010	Male	85 to 89	57% (40-71%)	43% (29-60%)	1.4 (0.79-2.3)	1.7% (0.37-3.7%)	1.8% (0.31-4.1%)	34 (5.9-76)	280 (61-600)
Uruguay	2010	Male	90 to 94	55% (39-70%)	45% (30-61%)	1.3 (0.69-2.3)	1.4% (0.11-3.4%)	1.5% (0.04-3.9%)	13 (0.34-33)	71 (5.7-180)
Uruguay	2010	Male	95 plus	57% (36-75%)	43% (25-64%)	1.3 (0.58-2.5)	1.4% (-0.16-4%)	1.4% (-0.36-4.3%)	3.5 (-0.89-11)	14 (-1.5-38)
Uruguay	2016	Female	All Ages	57% (51-62%)	43% (38-49%)	2.1 (1.6-2.8)	3% (1.7-4.7%)	3.2% (1.1-6.1%)	530 (180-1000)	14000 (7800-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	2016	Female	15 to 19	62% (47-76%)	38% (24-53%)	2 (1.1-3.3)	2.7% (1.5-4.1%)	8.3% (4.1-14%)	3.6 (1.8-6.2)	370 (210-560)
Uruguay	2016	Female	20 to 24	73% (60-83%)	27% (17-40%)	3.3 (1.9-5.2)	4.1% (2.6-5.9%)	12% (6.6-19%)	6.2 (3.3-9.6)	650 (400-920)
Uruguay	2016	Female	25 to 29	73% (60-83%)	27% (17-40%)	3.6 (2.1-5.3)	4% (2.6-5.6%)	12% (7.3-18%)	6.9 (4-10)	640 (430-900)
Uruguay	2016	Female	30 to 34	71% (58-81%)	29% (19-42%)	3.6 (2.1-5.4)	4.2% (2.7-6%)	12% (7.1-16%)	9.7 (5.9-14)	730 (470-1000)
Uruguay	2016	Female	35 to 39	67% (54-79%)	33% (21-46%)	3.2 (1.8-4.9)	4% (2.5-5.9%)	11% (6.5-15%)	12 (7.3-18)	800 (500-1200)
Uruguay	2016	Female	40 to 44	65% (52-77%)	35% (23-48%)	2.8 (1.6-4.6)	4.3% (2.4-6.5%)	10% (5.8-15%)	18 (10-27)	980 (560-1500)
Uruguay	2016	Female	45 to 49	64% (50-77%)	36% (23-50%)	2.8 (1.5-4.4)	4.4% (2.6-6.8%)	9.5% (5.6-14%)	22 (13-33)	1000 (590-1600)
Uruguay	2016	Female	50 to 54	64% (51-75%)	36% (25-49%)	2.7 (1.5-4.3)	4.7% (2.5-7.3%)	8.6% (4.9-13%)	32 (18-50)	1300 (720-2100)
Uruguay	2016	Female	55 to 59	63% (50-75%)	37% (25-50%)	2.7 (1.5-4.2)	4.8% (2.5-7.7%)	8% (4.5-12%)	43 (24-68)	1500 (790-2400)
Uruguay	2016	Female	60 to 64	60% (46-72%)	40% (28-54%)	2.4 (1.3-4.1)	4.1% (1.6-7.3%)	6.4% (2.8-11%)	45 (20-80)	1300 (530-2400)
Uruguay	2016	Female	65 to 69	54% (40-68%)	46% (32-60%)	2 (1.1-3.2)	3.3% (1.2-6.1%)	4.9% (1.9-8.8%)	46 (18-84)	1100 (410-2100)
Uruguay	2016	Female	70 to 74	49% (35-62%)	51% (38-65%)	1.7 (0.88-2.8)	2.6% (0.47-5.5%)	3.7% (0.8-7.6%)	47 (11-97)	920 (170-1900)
Uruguay	2016	Female	75 to 79	44% (31-60%)	56% (40-69%)	1.5 (0.76-2.6)	2% (-0.2-5.1%)	2.7% (-0.16-6.6%)	48 (-2.8-120)	730 (-77-1900)
Uruguay	2016	Female	80 to 84	42% (29-56%)	58% (44-71%)	1.4 (0.74-2.2)	1.7% (-0.23-4.3%)	2.1% (-0.24-5.4%)	59 (-6.5-150)	670 (-94-1700)
Uruguay	2016	Female	85 to 89	41% (27-55%)	59% (45-73%)	1.3 (0.66-2.2)	1.6% (-0.18-4.1%)	1.9% (-0.22-5.1%)	61 (-7.1-160)	490 (-51-1300)
Uruguay	2016	Female	90 to 94	41% (28-55%)	59% (45-72%)	1.3 (0.62-2.2)	1.5% (-0.43-4.3%)	1.8% (-0.54-5.3%)	46 (-14-140)	250 (-71-730)
Uruguay	2016	Female	95 plus	44% (28-61%)	56% (39-72%)	1.4 (0.57-2.7)	1.6% (-0.7-5.2%)	1.8% (-0.93-6.2%)	20 (-10-68)	73 (-31-240)
Uruguay	2016	Male	All Ages	67% (61-73%)	33% (27-39%)	2.2 (1.6-2.9)	5.5% (3.9-7.4%)	4.8% (3-7.2%)	810 (490-1200)	28000 (20000-39000)
Uruguay	2016	Male	15 to 19	69% (54-81%)	31% (19-46%)	2 (1.1-3.3)	6.4% (3.6-9.9%)	11% (5.5-17%)	15 (7.6-24)	1200 (700-1900)
Uruguay	2016	Male	20 to 24	76% (63-85%)	24% (15-37%)	3.3 (1.9-5.3)	11% (6.9-16%)	17% (9.3-26%)	33 (18-49)	2600 (1600-3700)
Uruguay	2016	Male	25 to 29	77% (65-87%)	23% (13-35%)	3.7 (2.2-5.5)	11% (7.3-16%)	18% (10-26%)	33 (19-49)	2500 (1500-3400)
Uruguay	2016	Male	30 to 34	77% (64-86%)	23% (14-36%)	3.6 (2.2-5.4)	9.5% (6.1-14%)	15% (8.9-22%)	28 (17-42)	1900 (1200-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uruguay	2016	Male	35 to 39	76% (64-85%)	24% (15-36%)	3.2 (1.8-4.9)	8.1% (5.3-12%)	13% (7.7-19%)	30 (18-45)	1800 (1200-2700)
Uruguay	2016	Male	40 to 44	76% (64-86%)	24% (14-36%)	3 (1.7-4.8)	7.7% (4.8-11%)	12% (7.2-18%)	36 (21-54)	1900 (1200-2800)
Uruguay	2016	Male	45 to 49	76% (64-86%)	24% (14-36%)	3 (1.7-4.7)	7.8% (5-12%)	11% (7.1-17%)	45 (28-69)	2100 (1400-3100)
Uruguay	2016	Male	50 to 54	76% (65-85%)	24% (15-35%)	2.9 (1.6-4.6)	7.5% (4.6-11%)	10% (6.1-16%)	65 (39-99)	2600 (1600-3900)
Uruguay	2016	Male	55 to 59	75% (64-84%)	25% (16-36%)	2.8 (1.7-4.4)	7% (4.2-10%)	8.9% (5.3-13%)	88 (52-130)	3000 (1800-4500)
Uruguay	2016	Male	60 to 64	73% (61-83%)	27% (17-39%)	2.6 (1.4-4.2)	6% (3.2-9.8%)	7.3% (3.9-12%)	97 (53-160)	2800 (1500-4600)
Uruguay	2016	Male	65 to 69	70% (57-80%)	30% (20-43%)	2.3 (1.4-3.6)	4.7% (2.4-7.7%)	5.5% (2.8-9.1%)	95 (48-160)	2300 (1200-3800)
Uruguay	2016	Male	70 to 74	67% (55-77%)	33% (23-45%)	2 (1.1-3.3)	3.4% (1.3-6.3%)	4% (1.5-7.3%)	82 (31-150)	1600 (620-2900)
Uruguay	2016	Male	75 to 79	65% (51-79%)	35% (21-49%)	1.8 (0.93-3.1)	2.6% (0.68-5.6%)	3% (0.74-6.3%)	67 (16-140)	1000 (250-2200)
Uruguay	2016	Male	80 to 84	61% (45-76%)	39% (24-55%)	1.5 (0.82-2.5)	1.9% (0.36-4.1%)	2.1% (0.37-4.5%)	49 (8.7-110)	560 (110-1200)
Uruguay	2016	Male	85 to 89	56% (40-71%)	44% (29-60%)	1.3 (0.65-2.2)	1.4% (0.24-3.5%)	1.6% (0.17-3.8%)	32 (3.4-79)	260 (43-630)
Uruguay	2016	Male	90 to 94	55% (39-70%)	45% (30-61%)	1.2 (0.59-2.2)	1.2% (-0.029-3.4%)	1.2% (-0.14-3.7%)	14 (-1.6-42)	81 (-2-230)
Uruguay	2016	Male	95 plus	56% (37-75%)	44% (25-63%)	1.2 (0.49-2.3)	1.3% (-0.28-4.2%)	1.3% (-0.49-4.6%)	4.1 (-1.6-15)	16 (-3.4-53)
Canada	1990	Female	All Ages	80% (76-83%)	20% (17-24%)	2.5 (2.2-2.8)	2.6% (1.4-3.7%)	1.2% (-1.2-3.9%)	1100 (-1000-3400)	82000 (46000-120000)
Canada	1990	Female	15 to 19	68% (56-78%)	32% (22-44%)	2.1 (1.2-3.1)	3.4% (2.1-5%)	11% (5.8-18%)	36 (19-58)	3700 (2300-5400)
Canada	1990	Female	20 to 24	86% (77-92%)	14% (8-23%)	4.1 (2.7-5.6)	5.4% (3.9-7.2%)	17% (10-24%)	65 (41-92)	7300 (5200-9600)
Canada	1990	Female	25 to 29	89% (82-93%)	11% (6.6-18%)	4.1 (2.8-5.3)	5% (3.7-6.6%)	16% (11-22%)	84 (56-110)	8800 (6400-12000)
Canada	1990	Female	30 to 34	90% (84-95%)	9.7% (5.4-16%)	3.9 (2.7-5.2)	4.7% (3.4-6.3%)	15% (10-19%)	100 (71-140)	9100 (6600-12000)
Canada	1990	Female	35 to 39	91% (85-95%)	9.3% (5.3-15%)	3.6 (2.5-4.9)	4.8% (3.3-6.3%)	14% (9.8-18%)	130 (90-170)	9200 (6400-12000)
Canada	1990	Female	40 to 44	91% (85-95%)	9.4% (5.5-15%)	3.4 (2.3-4.8)	4.9% (3.3-6.7%)	13% (8.7-17%)	160 (110-210)	9600 (6400-13000)
Canada	1990	Female	45 to 49	90% (84-94%)	9.7% (5.5-16%)	3.3 (2.2-4.7)	5.2% (3.3-7.3%)	11% (7.8-15%)	190 (130-250)	9200 (6200-13000)
Canada	1990	Female	50 to 54	89% (83-93%)	11% (6.5-17%)	3.1 (2-4.4)	4.9% (2.9-7.3%)	9.2% (5.8-13%)	200 (120-280)	8400 (5100-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	1990	Female	55 to 59	87% (79-93%)	13% (7.4-21%)	2.7 (1.8-4)	4.1% (2-6.3%)	6.9% (3.6-10%)	230 (120-330)	8000 (4000-12000)
Canada	1990	Female	60 to 64	84% (74-91%)	16% (9.5-26%)	2.4 (1.4-3.7)	2.9% (0.63-5.5%)	4.4% (1.2-8%)	220 (61-410)	6600 (1500-13000)
Canada	1990	Female	65 to 69	79% (68-88%)	21% (12-32%)	2.1 (1.4-3)	1.6% (-0.62-3.9%)	2.4% (-0.58-5.6%)	190 (-43-420)	4400 (-1600-11000)
Canada	1990	Female	70 to 74	75% (64-84%)	25% (16-36%)	1.8 (1.1-2.7)	0.42% (-1.8-2.9%)	0.62% (-2.3-3.8%)	60 (-220-360)	1100 (-4700-7600)
Canada	1990	Female	75 to 79	71% (57-83%)	29% (17-43%)	1.6 (0.93-2.5)	-0.43% (-2.8-2.4%)	-0.65% (-3.7-2.7%)	-82 (-470-350)	-1100 (-7000-5900)
Canada	1990	Female	80 to 84	68% (52-81%)	32% (19-48%)	1.5 (0.94-2.3)	-0.76% (-3.2-1.8%)	-1.1% (-4.1-2.1%)	-160 (-580-290)	-1500 (-6100-3500)
Canada	1990	Female	85 to 89	66% (50-80%)	34% (20-50%)	1.5 (0.91-2.3)	-0.79% (-3.1-1.9%)	-1.2% (-4.1-1.9%)	-160 (-530-240)	-960 (-3800-2400)
Canada	1990	Female	90 to 94	68% (52-81%)	32% (19-48%)	1.6 (0.94-2.5)	-0.8% (-3.5-2.3%)	-1.3% (-4.4-2.5%)	-110 (-390-220)	-450 (-2000-1300)
Canada	1990	Female	95 plus	71% (52-86%)	29% (14-48%)	1.8 (0.94-2.9)	-0.72% (-4.1-3.7%)	-1.3% (-5.3-4%)	-58 (-240-190)	-130 (-750-690)
Canada	1990	Male	All Ages	85% (82-88%)	15% (12-18%)	2.8 (2.5-3.1)	4.7% (3.2-6.2%)	2.5% (0.37-4.8%)	2600 (400-5100)	170000 (120000-240000)
Canada	1990	Male	15 to 19	69% (57-79%)	31% (21-43%)	2.2 (1.3-3.3)	7.2% (4.2-11%)	13% (6.8-19%)	120 (63-180)	10000 (6000-15000)
Canada	1990	Male	20 to 24	88% (81-94%)	12% (6.5-19%)	4.6 (3.1-6.3)	15% (10-20%)	23% (15-32%)	290 (180-400)	25000 (17000-33000)
Canada	1990	Male	25 to 29	92% (87-95%)	7.8% (4.5-13%)	4.8 (3.4-6.3)	13% (9.4-17%)	21% (14-29%)	320 (210-430)	26000 (19000-34000)
Canada	1990	Male	30 to 34	94% (89-97%)	6.1% (3.2-11%)	4.6 (3.2-6.1)	11% (7.8-14%)	18% (12-25%)	300 (200-420)	23000 (16000-30000)
Canada	1990	Male	35 to 39	94% (90-97%)	5.6% (2.9-9.5%)	4 (2.8-5.5)	8.9% (6.2-12%)	15% (9-20%)	270 (170-380)	19000 (13000-25000)
Canada	1990	Male	40 to 44	94% (90-97%)	5.6% (3-9.7%)	3.7 (2.4-5.2)	7.6% (4.8-11%)	12% (6.5-17%)	260 (140-390)	16000 (10000-22000)
Canada	1990	Male	45 to 49	94% (89-97%)	6% (3-11%)	3.5 (2.3-5)	7.1% (4.3-10%)	9.8% (5.3-15%)	270 (150-420)	14000 (8800-21000)
Canada	1990	Male	50 to 54	94% (89-97%)	6.4% (3.3-11%)	3.2 (2-4.7)	6% (3.1-9.5%)	7.6% (3.5-13%)	280 (130-480)	13000 (6700-21000)
Canada	1990	Male	55 to 59	93% (88-96%)	7% (3.7-12%)	2.9 (1.9-4.3)	4.7% (1.9-7.9%)	5.5% (1.8-9.8%)	330 (110-590)	13000 (5300-21000)
Canada	1990	Male	60 to 64	92% (86-96%)	8.1% (4.1-14%)	2.7 (1.6-4.1)	3.2% (0.39-6.7%)	3.4% (-0.094-7.9%)	320 (-8.6-740)	11000 (1400-22000)
Canada	1990	Male	65 to 69	90% (84-95%)	9.7% (4.8-16%)	2.4 (1.6-3.3)	1.7% (-0.76-4.6%)	1.7% (-1.3-5.1%)	210 (-160-630)	6200 (-2700-17000)
Canada	1990	Male	70 to 74	89% (81-94%)	11% (6.3-19%)	2 (1.2-2.9)	0.55% (-1.7-3%)	0.32% (-2.4-3.2%)	45 (-330-450)	1700 (-5400-9700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	1990	Male	75 to 79	87% (78-93%)	13% (6.7-22%)	1.7 (1-2.6)	-0.18% (-2.2-2.2%)	-0.53% (-3-2.2%)	-84 (-480-350)	-480 (-6200-5900)
Canada	1990	Male	80 to 84	86% (75-93%)	14% (7.3-25%)	1.6 (1-2.3)	-0.51% (-2.5-1.6%)	-0.88% (-3.2-1.5%)	-120 (-430-210)	-860 (-4100-2700)
Canada	1990	Male	85 to 89	85% (74-92%)	15% (7.9-26%)	1.6 (0.98-2.4)	-0.68% (-2.5-1.4%)	-1.1% (-3.3-1.2%)	-100 (-310-110)	-550 (-2000-1100)
Canada	1990	Male	90 to 94	86% (76-93%)	14% (7.1-24%)	1.6 (0.98-2.5)	-0.83% (-2.8-1.3%)	-1.3% (-3.8-1.2%)	-56 (-150-48)	-210 (-710-340)
Canada	1990	Male	95 plus	87% (75-94%)	13% (5.5-25%)	1.7 (0.92-2.8)	-0.86% (-3.2-2%)	-1.5% (-4.3-1.7%)	-23 (-67-26)	-54 (-200-120)
Canada	1995	Female	All Ages	80% (77-84%)	20% (16-23%)	2.4 (2.1-2.6)	2.2% (1.3-3.3%)	0.91% (-1.3-3.3%)	910 (-1300-3300)	77000 (42000-110000)
Canada	1995	Female	15 to 19	69% (58-79%)	31% (21-42%)	2 (1.2-3)	3.3% (2-4.8%)	11% (5.3-17%)	34 (17-54)	3700 (2200-5400)
Canada	1995	Female	20 to 24	87% (78-92%)	13% (7.9-22%)	3.8 (2.5-5.3)	5% (3.5-6.6%)	15% (9-22%)	51 (30-73)	6300 (4400-8500)
Canada	1995	Female	25 to 29	89% (83-93%)	11% (6.5-17%)	3.8 (2.7-5.2)	4.6% (3.3-6%)	14% (9.2-20%)	61 (39-83)	6800 (4900-9000)
Canada	1995	Female	30 to 34	90% (84-95%)	9.5% (5.4-16%)	3.7 (2.5-5)	4.2% (2.9-5.6%)	13% (8.8-18%)	89 (60-120)	8200 (5800-11000)
Canada	1995	Female	35 to 39	91% (85-95%)	9.2% (5.2-15%)	3.4 (2.3-4.6)	4.3% (3-5.6%)	13% (8.8-17%)	130 (87-170)	9100 (6400-12000)
Canada	1995	Female	40 to 44	91% (85-95%)	9.3% (5.4-15%)	3.2 (2.1-4.4)	4.4% (2.9-6.1%)	12% (7.8-16%)	160 (110-220)	9700 (6500-13000)
Canada	1995	Female	45 to 49	90% (84-94%)	9.6% (5.6-16%)	3.1 (2.1-4.3)	4.6% (3-6.5%)	10% (7.4-14%)	210 (150-280)	10000 (7100-14000)
Canada	1995	Female	50 to 54	89% (83-94%)	11% (6.4-17%)	2.9 (1.9-4.1)	4.5% (2.6-6.6%)	8.8% (5.6-12%)	220 (140-300)	9100 (5400-13000)
Canada	1995	Female	55 to 59	87% (79-93%)	13% (7.2-21%)	2.6 (1.7-3.8)	3.7% (1.8-5.8%)	6.5% (3.6-9.6%)	210 (120-310)	7500 (3800-12000)
Canada	1995	Female	60 to 64	84% (75-91%)	16% (9.4-25%)	2.3 (1.4-3.5)	2.6% (0.56-5%)	4.2% (1.4-7.5%)	200 (66-370)	5800 (1300-11000)
Canada	1995	Female	65 to 69	80% (69-88%)	20% (12-31%)	2 (1.3-2.8)	1.5% (-0.4-3.6%)	2.4% (-0.27-5.1%)	180 (-20-380)	4000 (-980-9600)
Canada	1995	Female	70 to 74	76% (65-84%)	24% (16-35%)	1.7 (1.1-2.5)	0.39% (-1.6-2.6%)	0.71% (-1.9-3.6%)	77 (-210-390)	1100 (-4700-7700)
Canada	1995	Female	75 to 79	72% (58-83%)	28% (17-42%)	1.5 (0.89-2.3)	-0.44% (-2.6-2%)	-0.55% (-3.3-2.5%)	-74 (-440-330)	-1200 (-6800-5400)
Canada	1995	Female	80 to 84	69% (54-81%)	31% (19-46%)	1.5 (0.89-2.1)	-0.85% (-3-1.5%)	-1.1% (-3.8-1.6%)	-190 (-640-280)	-2000 (-6800-3400)
Canada	1995	Female	85 to 89	67% (52-80%)	33% (20-48%)	1.4 (0.84-2.2)	-0.95% (-3.1-1.5%)	-1.3% (-4-1.6%)	-210 (-640-260)	-1400 (-4700-2300)
Canada	1995	Female	90 to 94	68% (52-81%)	32% (19-48%)	1.5 (0.89-2.4)	-0.99% (-3.5-1.9%)	-1.4% (-4.3-2%)	-160 (-480-220)	-690 (-2400-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	1995	Female	95 plus	71% (52-86%)	29% (14-48%)	1.7 (0.89-2.8)	-0.96% (-4.2-3.2%)	-1.5% (-5.3-3.3%)	-85 (-300-180)	-220 (-940-720)
Canada	1995	Male	All Ages	85% (82-88%)	15% (12-18%)	2.6 (2.3-2.9)	4.3% (3-5.7%)	2.3% (0.37-4.4%)	2600 (410-4900)	170000 (110000-220000)
Canada	1995	Male	15 to 19	69% (57-79%)	31% (21-43%)	2 (1.2-3.1)	6.6% (4-9.7%)	12% (6.1-19%)	97 (50-150)	9000 (5300-13000)
Canada	1995	Male	20 to 24	88% (81-94%)	12% (6.3-19%)	4.3 (2.9-6)	13% (9.1-18%)	22% (13-30%)	230 (140-330)	21000 (14000-28000)
Canada	1995	Male	25 to 29	92% (87-96%)	7.8% (4.5-13%)	4.5 (3.2-6.1)	12% (8.4-15%)	20% (13-27%)	230 (150-320)	20000 (14000-26000)
Canada	1995	Male	30 to 34	94% (89-97%)	6.1% (3.1-11%)	4.3 (3-5.8)	9.3% (6.6-12%)	15% (9.6-21%)	260 (170-370)	20000 (15000-27000)
Canada	1995	Male	35 to 39	94% (90-97%)	5.7% (3-9.5%)	3.8 (2.6-5.1)	8% (5.4-11%)	13% (7.6-18%)	270 (160-390)	19000 (13000-26000)
Canada	1995	Male	40 to 44	94% (90-97%)	5.6% (3-9.8%)	3.4 (2.3-4.8)	7.1% (4.5-9.8%)	11% (6.3-16%)	280 (160-400)	17000 (11000-23000)
Canada	1995	Male	45 to 49	94% (88-97%)	6% (3-12%)	3.3 (2.2-4.5)	6.6% (4.1-9.4%)	9.5% (5.4-14%)	310 (180-460)	16000 (10000-23000)
Canada	1995	Male	50 to 54	94% (89-97%)	6.4% (3.3-11%)	3 (1.9-4.4)	5.6% (3.1-8.6%)	7.4% (3.7-12%)	300 (150-480)	14000 (7400-21000)
Canada	1995	Male	55 to 59	93% (88-96%)	7% (3.6-12%)	2.7 (1.8-4)	4.4% (2-7.2%)	5.4% (2.1-9.1%)	300 (120-510)	11000 (5200-19000)
Canada	1995	Male	60 to 64	92% (86-96%)	8.1% (4-14%)	2.5 (1.5-3.8)	3.2% (0.63-6.2%)	3.6% (0.29-7.4%)	300 (25-620)	10000 (2100-20000)
Canada	1995	Male	65 to 69	90% (83-95%)	9.7% (4.9-17%)	2.2 (1.5-3.1)	1.9% (-0.32-4.5%)	2% (-0.73-5%)	250 (-90-630)	6900 (-1200-16000)
Canada	1995	Male	70 to 74	89% (82-94%)	11% (6.4-18%)	1.9 (1.2-2.7)	0.78% (-1.2-3%)	0.66% (-1.7-3.3%)	100 (-270-510)	2800 (-4500-11000)
Canada	1995	Male	75 to 79	87% (78-93%)	13% (6.8-22%)	1.6 (0.98-2.5)	-0.034% (-2-2.1%)	-0.32% (-2.6-2.1%)	-52 (-430-330)	-94 (-5600-5700)
Canada	1995	Male	80 to 84	86% (74-93%)	14% (7.2-26%)	1.5 (0.97-2.2)	-0.45% (-2.2-1.5%)	-0.77% (-2.9-1.4%)	-130 (-470-230)	-900 (-4500-3100)
Canada	1995	Male	85 to 89	85% (73-92%)	15% (7.8-27%)	1.5 (0.9-2.2)	-0.65% (-2.4-1.2%)	-1% (-3.1-1.1%)	-110 (-340-120)	-630 (-2300-1100)
Canada	1995	Male	90 to 94	86% (76-93%)	14% (7.1-24%)	1.5 (0.91-2.3)	-0.83% (-2.7-1.1%)	-1.3% (-3.5-0.99%)	-69 (-180-54)	-270 (-840-380)
Canada	1995	Male	95 plus	87% (76-94%)	13% (5.6-24%)	1.6 (0.86-2.6)	-0.86% (-3-1.6%)	-1.4% (-3.9-1.4%)	-24 (-66-23)	-58 (-200-110)
Canada	2000	Female	All Ages	81% (78-84%)	19% (16-22%)	2.4 (2.2-2.7)	2.3% (1.4-3.3%)	1.1% (-0.97-3.4%)	1200 (-1000-3600)	82000 (49000-120000)
Canada	2000	Female	15 to 19	76% (69-83%)	24% (17-31%)	2.2 (1.4-3.2)	3.4% (2.1-5%)	12% (5.9-18%)	37 (19-57)	4000 (2500-5700)
Canada	2000	Female	20 to 24	88% (81-93%)	12% (7.2-19%)	4 (2.6-5.6)	5.1% (3.6-6.7%)	16% (9.4-22%)	52 (32-74)	6500 (4600-8700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	2000	Female	25 to 29	90% (85-93%)	10% (6.6-15%)	3.9 (2.7-5.3)	4.5% (3.2-5.8%)	14% (9.3-20%)	51 (33-70)	6100 (4300-8100)
Canada	2000	Female	30 to 34	90% (85-94%)	9.5% (5.7-15%)	3.7 (2.6-5.1)	4.1% (2.9-5.4%)	13% (8.9-18%)	69 (46-93)	6700 (4600-8900)
Canada	2000	Female	35 to 39	91% (87-95%)	9% (5.1-13%)	3.4 (2.4-4.6)	4.1% (2.8-5.5%)	13% (8.7-16%)	120 (82-160)	9000 (6200-12000)
Canada	2000	Female	40 to 44	91% (87-95%)	8.8% (5.3-13%)	3.3 (2.2-4.5)	4.3% (2.8-5.9%)	12% (8-15%)	170 (120-230)	10000 (6900-14000)
Canada	2000	Female	45 to 49	91% (85-94%)	9.2% (5.5-15%)	3.2 (2.2-4.4)	4.5% (2.9-6.2%)	11% (7.6-14%)	220 (160-290)	11000 (7300-15000)
Canada	2000	Female	50 to 54	90% (84-93%)	10% (6.6-16%)	2.9 (2-4.2)	4.5% (2.6-6.5%)	9.1% (6-12%)	270 (170-360)	11000 (6900-16000)
Canada	2000	Female	55 to 59	87% (80-92%)	13% (7.6-20%)	2.6 (1.8-3.7)	3.7% (1.9-5.5%)	6.8% (4-9.5%)	240 (140-340)	8500 (4500-13000)
Canada	2000	Female	60 to 64	84% (76-90%)	16% (9.9-24%)	2.3 (1.4-3.4)	2.7% (0.82-4.9%)	4.5% (1.9-7.6%)	210 (88-350)	6000 (1900-11000)
Canada	2000	Female	65 to 69	79% (71-87%)	21% (13-29%)	2 (1.4-2.8)	1.6% (-0.049-3.5%)	2.8% (0.49-5.1%)	190 (33-350)	4200 (-130-9100)
Canada	2000	Female	70 to 74	74% (65-82%)	26% (18-35%)	1.7 (1.1-2.5)	0.69% (-1-2.7%)	1.2% (-1.1-3.7%)	120 (-110-370)	2000 (-3000-7800)
Canada	2000	Female	75 to 79	70% (59-81%)	30% (19-41%)	1.5 (0.91-2.3)	-0.069% (-1.9-2.2%)	-0.016% (-2.3-2.7%)	-2.5 (-350-390)	-210 (-5600-6600)
Canada	2000	Female	80 to 84	69% (53-81%)	31% (19-47%)	1.5 (0.92-2.2)	-0.53% (-2.5-1.7%)	-0.7% (-3.1-2%)	-120 (-540-340)	-1300 (-6000-4200)
Canada	2000	Female	85 to 89	67% (51-81%)	33% (19-49%)	1.5 (0.85-2.2)	-0.71% (-2.7-1.6%)	-1% (-3.5-1.7%)	-200 (-660-330)	-1300 (-4800-2900)
Canada	2000	Female	90 to 94	68% (53-82%)	32% (18-47%)	1.6 (0.91-2.4)	-0.74% (-3.2-2%)	-1.1% (-4-2%)	-150 (-530-270)	-620 (-2700-1700)
Canada	2000	Female	95 plus	71% (52-86%)	29% (14-48%)	1.7 (0.91-2.9)	-0.68% (-3.8-3.4%)	-1.1% (-4.8-3.6%)	-77 (-330-250)	-190 (-1000-940)
Canada	2000	Male	All Ages	88% (85-90%)	12% (10-15%)	2.8 (2.5-3.1)	4.6% (3.3-5.9%)	2.7% (0.82-4.6%)	3000 (920-5300)	18000 (130000-230000)
Canada	2000	Male	15 to 19	79% (72-86%)	21% (14-28%)	2.5 (1.5-3.7)	7% (4.4-10%)	14% (7.4-21%)	99 (52-150)	9400 (5900-13000)
Canada	2000	Male	20 to 24	91% (86-95%)	8.7% (5.2-14%)	4.7 (3.2-6.4)	14% (9.5-18%)	23% (14-32%)	220 (130-300)	20000 (14000-26000)
Canada	2000	Male	25 to 29	93% (90-96%)	6.5% (4-9.7%)	4.7 (3.2-6.3)	12% (8.7-15%)	22% (14-30%)	190 (120-260)	17000 (12000-22000)
Canada	2000	Male	30 to 34	95% (92-97%)	5.1% (2.8-8%)	4.5 (3.1-6.1)	9.9% (7.2-13%)	18% (12-26%)	210 (130-290)	17000 (12000-23000)
Canada	2000	Male	35 to 39	95% (92-97%)	4.9% (2.7-7.9%)	3.9 (2.8-5.3)	8.5% (5.8-11%)	15% (9.1-21%)	270 (160-360)	19000 (13000-25000)
Canada	2000	Male	40 to 44	95% (92-97%)	4.8% (2.7-7.8%)	3.6 (2.4-5)	7.7% (5-10%)	13% (7.5-18%)	310 (180-440)	19000 (12000-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	2000	Male	45 to 49	95% (91-97%)	5.1% (2.8-8.7%)	3.4 (2.4-4.7)	7.2% (4.6-10%)	11% (6.7-16%)	360 (220-510)	19000 (12000-26000)
Canada	2000	Male	50 to 54	94% (91-97%)	5.6% (3.1-9.4%)	3.2 (2-4.5)	6.2% (3.6-9.1%)	8.6% (4.8-13%)	390 (210-590)	18000 (10000-26000)
Canada	2000	Male	55 to 59	94% (90-96%)	6.2% (3.5-9.9%)	2.9 (1.9-4.1)	4.8% (2.5-7.5%)	6.2% (2.9-9.9%)	360 (160-570)	14000 (7300-21000)
Canada	2000	Male	60 to 64	93% (89-96%)	7% (3.9-11%)	2.7 (1.7-4)	3.6% (1.2-6.6%)	4.4% (1.3-8.2%)	330 (94-620)	11000 (3500-20000)
Canada	2000	Male	65 to 69	91% (86-96%)	8.6% (4.4-14%)	2.3 (1.6-3.3)	2.4% (0.26-4.7%)	2.7% (0.03-5.7%)	300 (3.3-630)	8000 (840-16000)
Canada	2000	Male	70 to 74	89% (83-94%)	11% (6.3-17%)	1.9 (1.2-2.7)	1.2% (-0.57-3.3%)	1.3% (-0.89-3.8%)	190 (-130-560)	4400 (-2000-12000)
Canada	2000	Male	75 to 79	87% (80-93%)	13% (6.7-20%)	1.7 (1-2.5)	0.37% (-1.3-2.4%)	0.18% (-1.8-2.5%)	33 (-320-440)	1200 (-4200-7500)
Canada	2000	Male	80 to 84	86% (75-93%)	14% (7.4-25%)	1.5 (0.98-2.2)	-0.17% (-1.7-1.7%)	-0.45% (-2.4-1.7%)	-73 (-380-270)	-350 (-3600-3500)
Canada	2000	Male	85 to 89	84% (74-92%)	16% (7.9-26%)	1.5 (0.9-2.2)	-0.44% (-2-1.3%)	-0.79% (-2.7-1.2%)	-100 (-340-160)	-510 (-2300-1500)
Canada	2000	Male	90 to 94	85% (75-92%)	15% (7.6-25%)	1.5 (0.92-2.4)	-0.63% (-2.4-1.3%)	-1% (-3.1-1.2%)	-65 (-200-72)	-240 (-900-500)
Canada	2000	Male	95 plus	87% (75-94%)	13% (5.9-25%)	1.6 (0.87-2.7)	-0.65% (-2.7-1.9%)	-1.1% (-3.6-1.8%)	-23 (-72-35)	-51 (-220-150)
Canada	2005	Female	All Ages	82% (79-84%)	18% (16-21%)	2.6 (2.3-2.8)	2.6% (1.7-3.5%)	1.6% (-0.2-3.8%)	1900 (-220-4300)	95000 (61000-130000)
Canada	2005	Female	15 to 19	76% (69-83%)	24% (17-31%)	2.3 (1.4-3.4)	3.4% (2.2-4.9%)	12% (6.2-18%)	36 (18-55)	4000 (2600-5700)
Canada	2005	Female	20 to 24	88% (81-93%)	12% (7.2-19%)	4.2 (2.8-5.9)	5.2% (3.7-6.8%)	16% (9.9-23%)	57 (35-80)	7200 (5100-9500)
Canada	2005	Female	25 to 29	90% (85-93%)	10% (6.5-15%)	4.2 (2.9-5.6)	4.5% (3.4-5.9%)	15% (9.7-20%)	52 (34-71)	6400 (4600-8500)
Canada	2005	Female	30 to 34	90% (86-94%)	9.5% (5.8-14%)	3.9 (2.7-5.5)	4.2% (3-5.4%)	14% (9.4-19%)	66 (44-90)	6500 (4600-8700)
Canada	2005	Female	35 to 39	91% (87-95%)	9% (5.1-13%)	3.6 (2.5-4.9)	4.2% (2.9-5.5%)	13% (9.2-17%)	100 (71-130)	7800 (5500-10000)
Canada	2005	Female	40 to 44	91% (87-95%)	8.8% (5.3-13%)	3.5 (2.3-4.7)	4.4% (3-6%)	12% (8.3-16%)	170 (120-220)	11000 (7400-14000)
Canada	2005	Female	45 to 49	91% (85-95%)	9.3% (5.5-15%)	3.4 (2.3-4.6)	4.7% (3.2-6.4%)	11% (8-14%)	250 (180-330)	13000 (8900-17000)
Canada	2005	Female	50 to 54	89% (84-93%)	11% (6.6-16%)	3.1 (2.1-4.4)	4.6% (2.9-6.6%)	9.5% (6.5-13%)	300 (210-400)	13000 (8300-18000)
Canada	2005	Female	55 to 59	87% (80-92%)	13% (7.8-20%)	2.8 (1.9-3.9)	3.9% (2.3-5.7%)	7.3% (4.7-9.9%)	310 (200-420)	11000 (6600-16000)
Canada	2005	Female	60 to 64	84% (76-90%)	16% (10-24%)	2.4 (1.5-3.5)	2.9% (1.2-5%)	5% (2.5-7.7%)	260 (130-400)	7800 (3100-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	2005	Female	65 to 69	79% (70-87%)	21% (13-30%)	2.1 (1.5-2.9)	2% (0.46-3.7%)	3.3% (1.2-5.5%)	220 (77-360)	5000 (1200-9300)
Canada	2005	Female	70 to 74	74% (65-82%)	26% (18-35%)	1.8 (1.1-2.5)	1.1% (-0.57-3%)	1.8% (-0.38-4.2%)	170 (-36-400)	2900 (-1600-8100)
Canada	2005	Female	75 to 79	70% (58-80%)	30% (20-42%)	1.6 (0.95-2.4)	0.37% (-1.2-2.4%)	0.6% (-1.4-3.1%)	85 (-200-440)	1100 (-3600-7000)
Canada	2005	Female	80 to 84	68% (53-81%)	32% (19-47%)	1.5 (0.97-2.2)	-0.004% (-1.8-2%)	-0.017% (-2.2-2.5%)	-3.3 (-440-480)	-8.7 (-5000-5700)
Canada	2005	Female	85 to 89	67% (51-80%)	33% (20-49%)	1.5 (0.91-2.3)	-0.26% (-2.1-2%)	-0.48% (-2.8-2.1%)	-96 (-560-430)	-500 (-4200-3900)
Canada	2005	Female	90 to 94	68% (52-82%)	32% (18-48%)	1.7 (0.96-2.6)	-0.24% (-2.6-2.5%)	-0.53% (-3.2-2.7%)	-85 (-520-430)	-250 (-2600-2500)
Canada	2005	Female	95 plus	71% (52-86%)	29% (14-48%)	1.8 (0.95-3)	-0.082% (-3.3-4%)	-0.42% (-4-4.3%)	-35 (-340-350)	-27 (-1100-1300)
Canada	2005	Male	All Ages	88% (86-90%)	12% (9.8-14%)	2.9 (2.6-3.2)	5% (3.8-6.3%)	3.3% (1.6-5.2%)	3900 (1800-6100)	200000 (150000-260000)
Canada	2005	Male	15 to 19	80% (72-86%)	20% (14-28%)	2.6 (1.6-3.8)	7% (4.4-10%)	14% (7.8-22%)	96 (53-140)	9400 (6000-13000)
Canada	2005	Male	20 to 24	91% (86-95%)	8.7% (5.3-14%)	4.9 (3.2-6.8)	14% (9.7-18%)	23% (14-32%)	220 (140-310)	21000 (15000-27000)
Canada	2005	Male	25 to 29	93% (90-96%)	6.5% (4-9.9%)	5 (3.4-6.7)	12% (8.9-16%)	22% (15-30%)	200 (130-270)	18000 (13000-23000)
Canada	2005	Male	30 to 34	95% (92-97%)	5.1% (2.9-8.4%)	4.7 (3.2-6.5)	10% (7.3-14%)	19% (13-27%)	190 (130-270)	16000 (11000-21000)
Canada	2005	Male	35 to 39	95% (92-97%)	4.8% (2.7-7.8%)	4.2 (3-5.7)	8.8% (6.1-12%)	16% (10-22%)	230 (140-310)	16000 (11000-22000)
Canada	2005	Male	40 to 44	95% (92-97%)	4.8% (2.7-7.9%)	3.8 (2.6-5.4)	8.1% (5.4-11%)	14% (8.7-20%)	330 (210-470)	20000 (14000-28000)
Canada	2005	Male	45 to 49	95% (91-97%)	5.1% (2.7-8.7%)	3.6 (2.5-5.1)	7.9% (5.2-11%)	12% (8-17%)	440 (280-610)	23000 (15000-31000)
Canada	2005	Male	50 to 54	94% (91-97%)	5.5% (3-9%)	3.4 (2.2-4.8)	7.1% (4.6-10%)	10% (6.5-15%)	500 (310-720)	22000 (14000-31000)
Canada	2005	Male	55 to 59	94% (90-97%)	6.2% (3.5-10%)	3 (2-4.3)	5.6% (3.3-8.3%)	7.5% (4.4-11%)	500 (290-760)	19000 (11000-28000)
Canada	2005	Male	60 to 64	93% (89-96%)	7% (3.9-11%)	2.8 (1.8-4.2)	4.3% (1.9-7.3%)	5.5% (2.3-9.4%)	450 (190-760)	14000 (6400-24000)
Canada	2005	Male	65 to 69	91% (86-95%)	8.6% (4.6-14%)	2.5 (1.7-3.4)	3% (1.1-5.4%)	3.6% (1.2-6.5%)	360 (120-650)	9600 (3600-17000)
Canada	2005	Male	70 to 74	89% (84-94%)	11% (6.4-16%)	2 (1.3-2.9)	1.8% (0.11-3.8%)	2.1% (-0.026-4.5%)	290 (-3.5-620)	6200 (370-13000)
Canada	2005	Male	75 to 79	88% (80-93%)	12% (6.6-20%)	1.8 (1.1-2.7)	0.94% (-0.68-2.9%)	0.91% (-1-3.2%)	160 (-180-550)	3000 (-2100-9300)
Canada	2005	Male	80 to 84	86% (75-93%)	14% (7.3-25%)	1.6 (1-2.4)	0.3% (-1.2-2.1%)	0.1% (-1.7-2.1%)	20 (-330-410)	760 (-3000-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	2005	Male	85 to 89	85% (73-92%)	15% (8-27%)	1.6 (0.99-2.3)	-0.076% (-1.5-1.7%)	-0.37% (-2.1-1.6%)	-53 (-310-220)	-95 (-2000-2100)
Canada	2005	Male	90 to 94	86% (76-93%)	14% (7.2-24%)	1.6 (0.99-2.5)	-0.26% (-1.9-1.6%)	-0.62% (-2.6-1.5%)	-49 (-200-120)	-130 (-910-780)
Canada	2005	Male	95 plus	87% (76-94%)	13% (5.6-24%)	1.7 (0.92-2.9)	-0.26% (-2.4-2.3%)	-0.69% (-3.2-2.3%)	-17 (-79-57)	-26 (-240-230)
Canada	2010	Female	All Ages	80% (77-84%)	20% (16-23%)	2.7 (2.4-3)	2.8% (2-3.7%)	2.2% (0.51-4.2%)	2700 (610-5100)	110000 (76000-150000)
Canada	2010	Female	15 to 19	68% (55-78%)	32% (22-45%)	2.2 (1.3-3.2)	3% (2-4.3%)	11% (5.9-17%)	31 (16-49)	3700 (2300-5200)
Canada	2010	Female	20 to 24	86% (76-92%)	14% (8-24%)	4.3 (2.9-6)	5.1% (3.7-6.7%)	17% (10-23%)	57 (35-81)	7300 (5100-9700)
Canada	2010	Female	25 to 29	89% (82-93%)	11% (6.8-18%)	4.4 (3-6)	4.7% (3.5-6%)	16% (10-21%)	60 (41-83)	7300 (5200-9500)
Canada	2010	Female	30 to 34	90% (83-95%)	10% (5.5-17%)	4.1 (2.8-5.8)	4.3% (3.1-5.6%)	15% (10-19%)	73 (50-98)	7200 (5200-9500)
Canada	2010	Female	35 to 39	90% (85-95%)	9.5% (5.3-15%)	3.9 (2.6-5.3)	4.3% (3-5.6%)	14% (9.8-18%)	98 (69-130)	7800 (5500-10000)
Canada	2010	Female	40 to 44	90% (85-95%)	9.5% (5.5-15%)	3.6 (2.4-5.1)	4.6% (3.1-6.1%)	13% (9.3-17%)	160 (110-200)	9800 (6600-13000)
Canada	2010	Female	45 to 49	90% (83-94%)	10% (5.7-17%)	3.5 (2.4-4.8)	4.8% (3.3-6.5%)	12% (8.8-15%)	260 (190-330)	13000 (9600-18000)
Canada	2010	Female	50 to 54	89% (82-93%)	11% (6.7-18%)	3.3 (2.2-4.6)	4.8% (3.1-6.7%)	9.9% (7.3-13%)	340 (240-440)	15000 (10000-20000)
Canada	2010	Female	55 to 59	87% (79-93%)	13% (7.4-21%)	2.9 (2-4.1)	4.2% (2.4-6%)	7.9% (5.3-11%)	350 (230-470)	13000 (7700-19000)
Canada	2010	Female	60 to 64	83% (74-90%)	17% (9.8-26%)	2.6 (1.6-3.8)	3.2% (1.4-5.2%)	5.6% (3.2-8.4%)	340 (190-500)	10000 (4700-17000)
Canada	2010	Female	65 to 69	79% (68-87%)	21% (13-32%)	2.2 (1.5-3.1)	2.3% (0.83-4%)	3.9% (1.8-6%)	280 (130-430)	6800 (2400-12000)
Canada	2010	Female	70 to 74	75% (63-84%)	25% (16-37%)	1.9 (1.2-2.8)	1.5% (-0.088-3.3%)	2.5% (0.36-4.8%)	230 (32-440)	4100 (-230-9000)
Canada	2010	Female	75 to 79	70% (55-82%)	30% (18-45%)	1.7 (1-2.6)	0.89% (-0.64-2.9%)	1.4% (-0.64-3.7%)	180 (-85-490)	2500 (-1900-8300)
Canada	2010	Female	80 to 84	67% (51-81%)	33% (19-49%)	1.6 (0.97-2.3)	0.61% (-1.2-2.4%)	0.8% (-1.2-3%)	150 (-230-570)	1700 (-2900-6800)
Canada	2010	Female	85 to 89	65% (49-79%)	35% (21-51%)	1.6 (0.93-2.4)	0.32% (-1.4-2.4%)	0.25% (-1.9-2.7%)	57 (-440-620)	760 (-3400-5500)
Canada	2010	Female	90 to 94	67% (51-81%)	33% (19-49%)	1.7 (1-2.6)	0.33% (-1.9-3%)	0.16% (-2.4-3.3%)	28 (-440-590)	390 (-2200-3600)
Canada	2010	Female	95 plus	71% (52-86%)	29% (14-48%)	2 (1-3.2)	0.54% (-2.6-4.5%)	0.33% (-3.3-4.9%)	36 (-360-540)	240 (-1100-2000)
Canada	2010	Male	All Ages	86% (84-89%)	14% (11-16%)	3 (2.7-3.4)	5.2% (4-6.5%)	3.9% (2.2-5.6%)	4700 (2700-6900)	220000 (170000-270000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	2010	Male	15 to 19	69% (58-80%)	31% (20-42%)	2.4 (1.4-3.6)	6% (3.7-8.5%)	13% (7-20%)	78 (41-120)	7800 (4900-11000)
Canada	2010	Male	20 to 24	89% (81-94%)	11% (6.1-19%)	5 (3.4-7.1)	13% (9.1-17%)	23% (15-32%)	210 (130-280)	19000 (14000-25000)
Canada	2010	Male	25 to 29	92% (87-95%)	7.7% (4.5-13%)	5.2 (3.6-7)	12% (8.6-15%)	23% (15-30%)	200 (140-270)	18000 (13000-23000)
Canada	2010	Male	30 to 34	94% (89-97%)	6.1% (3-11%)	4.9 (3.3-6.8)	9.9% (7.3-13%)	20% (13-28%)	190 (130-260)	16000 (11000-20000)
Canada	2010	Male	35 to 39	94% (90-97%)	5.7% (2.9-9.8%)	4.3 (3-6)	8.9% (6.3-12%)	18% (11-24%)	210 (140-280)	15000 (11000-20000)
Canada	2010	Male	40 to 44	94% (90-97%)	5.6% (2.9-9.8%)	4 (2.6-5.6)	8.4% (5.8-11%)	15% (10-21%)	290 (190-390)	18000 (12000-24000)
Canada	2010	Male	45 to 49	94% (89-97%)	6% (3-11%)	3.8 (2.6-5.2)	8.2% (5.7-11%)	14% (9.2-19%)	450 (300-630)	24000 (17000-32000)
Canada	2010	Male	50 to 54	94% (89-97%)	6.5% (3.5-11%)	3.5 (2.3-4.9)	7.6% (5.1-11%)	12% (7.5-16%)	590 (380-840)	26000 (18000-36000)
Canada	2010	Male	55 to 59	93% (88-96%)	6.9% (3.7-12%)	3.2 (2.1-4.4)	6.4% (4.1-9%)	9% (5.8-13%)	630 (400-890)	24000 (15000-34000)
Canada	2010	Male	60 to 64	92% (86-96%)	8% (4.1-14%)	2.9 (1.8-4.3)	5% (2.6-7.8%)	6.6% (3.5-10%)	610 (320-960)	20000 (11000-31000)
Canada	2010	Male	65 to 69	90% (83-95%)	9.7% (4.8-17%)	2.6 (1.7-3.5)	3.6% (1.7-5.8%)	4.5% (2.1-7.3%)	480 (230-790)	13000 (6300-21000)
Canada	2010	Male	70 to 74	89% (82-94%)	11% (6.3-18%)	2.1 (1.4-3.1)	2.4% (0.69-4.3%)	2.9% (0.78-5.2%)	360 (98-660)	7800 (2200-14000)
Canada	2010	Male	75 to 79	87% (78-93%)	13% (6.7-22%)	1.9 (1.2-2.8)	1.5% (-0.14-3.4%)	1.6% (-0.35-3.7%)	260 (-58-620)	4600 (-440-11000)
Canada	2010	Male	80 to 84	86% (75-93%)	14% (7.4-25%)	1.8 (1.1-2.5)	0.8% (-0.68-2.4%)	0.71% (-1.2-2.6%)	140 (-230-500)	2100 (-1900-6500)
Canada	2010	Male	85 to 89	85% (74-92%)	15% (7.8-26%)	1.7 (1.1-2.5)	0.36% (-1.1-2%)	0.13% (-1.6-2.1%)	22 (-280-360)	580 (-1700-3400)
Canada	2010	Male	90 to 94	86% (76-93%)	14% (7.4-24%)	1.8 (1.1-2.7)	0.17% (-1.5-2.1%)	-0.13% (-2.1-2.1%)	-12 (-190-200)	98 (-870-1200)
Canada	2010	Male	95 plus	87% (75-94%)	13% (5.8-25%)	1.9 (1-3.1)	0.2% (-1.9-2.9%)	-0.16% (-2.7-2.8%)	-5.4 (-91-97)	28 (-260-390)
Canada	2016	Female	All Ages	81% (77-84%)	19% (16-23%)	2.7 (2.2-3.2)	2.8% (1.8-3.8%)	2.2% (0.21-4.4%)	2900 (290-5800)	120000 (74000-160000)
Canada	2016	Female	15 to 19	68% (56-79%)	32% (21-44%)	2.2 (1.3-3.4)	3% (1.9-4.3%)	11% (6-18%)	30 (16-47)	3600 (2300-5100)
Canada	2016	Female	20 to 24	86% (77-92%)	14% (8.2-23%)	4.3 (2.7-6.3)	5% (3.6-6.6%)	17% (10-24%)	59 (36-84)	7700 (5400-10000)
Canada	2016	Female	25 to 29	89% (82-93%)	11% (6.6-18%)	4.3 (2.9-6.1)	4.6% (3.4-6%)	16% (10-22%)	63 (41-89)	7700 (5500-10000)
Canada	2016	Female	30 to 34	90% (84-95%)	9.7% (5.4-16%)	4.1 (2.8-5.8)	4.4% (3.1-5.7%)	15% (10-20%)	81 (55-110)	8000 (5700-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	2016	Female	35 to 39	91% (86-95%)	9.2% (5.1-14%)	3.8 (2.5-5.5)	4.3% (3-5.8%)	14% (9.7-18%)	100 (69-130)	7900 (5500-11000)
Canada	2016	Female	40 to 44	91% (85-95%)	9.2% (5.3-15%)	3.6 (2.3-5.2)	4.6% (3.1-6.3%)	13% (9.2-18%)	150 (100-200)	9400 (6400-13000)
Canada	2016	Female	45 to 49	90% (83-95%)	9.7% (5.2-17%)	3.5 (2.3-5)	4.8% (3.2-6.6%)	12% (8.8-16%)	240 (170-320)	13000 (8700-17000)
Canada	2016	Female	50 to 54	89% (82-94%)	11% (6.4-18%)	3.3 (2.1-4.7)	4.7% (3-6.8%)	10% (7.1-14%)	350 (250-480)	16000 (10000-23000)
Canada	2016	Female	55 to 59	87% (79-93%)	13% (7.3-21%)	2.9 (1.9-4.2)	4.1% (2.4-6.1%)	8% (5.4-11%)	400 (260-540)	15000 (8600-22000)
Canada	2016	Female	60 to 64	84% (75-90%)	16% (9.7-25%)	2.6 (1.6-3.9)	3.1% (1.3-5.2%)	5.7% (3.2-8.7%)	380 (210-570)	12000 (5200-19000)
Canada	2016	Female	65 to 69	79% (68-88%)	21% (12-32%)	2.2 (1.5-3.1)	2.3% (0.68-4.1%)	3.9% (1.8-6.1%)	350 (160-560)	8500 (2600-15000)
Canada	2016	Female	70 to 74	75% (64-84%)	25% (16-36%)	1.9 (1.2-2.9)	1.5% (-0.14-3.4%)	2.5% (0.35-4.8%)	260 (36-520)	4700 (-480-11000)
Canada	2016	Female	75 to 79	71% (57-83%)	29% (17-43%)	1.7 (0.97-2.7)	0.85% (-0.78-2.9%)	1.3% (-0.75-4%)	180 (-100-540)	2500 (-2200-8700)
Canada	2016	Female	80 to 84	68% (52-82%)	32% (18-48%)	1.6 (0.98-2.4)	0.56% (-1.1-2.5%)	0.75% (-1.4-3.2%)	140 (-270-600)	1600 (-3400-7500)
Canada	2016	Female	85 to 89	66% (50-80%)	34% (20-50%)	1.6 (0.93-2.5)	0.35% (-1.5-2.4%)	0.28% (-1.9-2.8%)	70 (-470-700)	890 (-3700-6100)
Canada	2016	Female	90 to 94	68% (53-81%)	32% (19-47%)	1.8 (1-2.8)	0.35% (-1.9-3.1%)	0.18% (-2.4-3.4%)	40 (-530-760)	520 (-2800-4500)
Canada	2016	Female	95 plus	72% (53-86%)	28% (14-47%)	2 (1-3.2)	0.53% (-2.7-4.6%)	0.31% (-3.5-5.1%)	45 (-500-730)	310 (-1500-2700)
Canada	2016	Male	All Ages	87% (84-89%)	13% (11-16%)	3 (2.5-3.6)	5.1% (3.8-6.4%)	3.7% (1.9-5.6%)	5100 (2600-7700)	230000 (170000-300000)
Canada	2016	Male	15 to 19	69% (57-80%)	31% (20-43%)	2.3 (1.3-3.6)	5.9% (3.7-8.5%)	13% (6.9-20%)	73 (38-110)	7500 (4600-11000)
Canada	2016	Male	20 to 24	88% (81-94%)	12% (6.3-19%)	5 (3.2-7.1)	12% (8.8-17%)	23% (14-32%)	210 (130-290)	20000 (14000-27000)
Canada	2016	Male	25 to 29	92% (87-96%)	7.8% (4.4-13%)	5.2 (3.5-7.2)	11% (8.4-15%)	22% (15-31%)	210 (130-280)	19000 (14000-25000)
Canada	2016	Male	30 to 34	94% (89-97%)	6.2% (3-11%)	4.9 (3.3-6.8)	9.8% (7-13%)	20% (13-28%)	200 (130-280)	17000 (12000-22000)
Canada	2016	Male	35 to 39	94% (90-97%)	5.7% (2.9-9.6%)	4.3 (2.9-6.2)	8.8% (6.3-12%)	18% (11-24%)	210 (130-290)	15000 (10000-20000)
Canada	2016	Male	40 to 44	94% (90-97%)	5.6% (2.9-9.9%)	3.9 (2.5-5.7)	8.5% (5.7-12%)	16% (9.7-22%)	270 (170-390)	17000 (11000-23000)
Canada	2016	Male	45 to 49	94% (89-97%)	6% (2.9-11%)	3.7 (2.5-5.3)	8.1% (5.6-11%)	14% (9.1-20%)	410 (270-570)	22000 (15000-30000)
Canada	2016	Male	50 to 54	94% (89-97%)	6.5% (3.4-11%)	3.5 (2.2-5)	7.6% (4.9-11%)	12% (7.5-17%)	630 (400-900)	28000 (18000-40000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Canada	2016	Male	55 to 59	93% (88-96%)	7.1% (3.7-12%)	3.1 (2.1-4.5)	6.4% (4.1-9.3%)	9.1% (5.8-13%)	720 (460-1100)	28000 (18000-40000)
Canada	2016	Male	60 to 64	92% (86-96%)	8.2% (4.2-14%)	2.8 (1.8-4.2)	4.9% (2.7-7.8%)	6.7% (3.7-11%)	690 (380-1100)	23000 (12000-36000)
Canada	2016	Male	65 to 69	90% (83-95%)	9.8% (5-17%)	2.5 (1.7-3.5)	3.6% (1.7-5.8%)	4.6% (2.1-7.4%)	610 (280-990)	16000 (7600-27000)
Canada	2016	Male	70 to 74	89% (82-94%)	11% (6.4-18%)	2.1 (1.3-3.1)	2.4% (0.59-4.5%)	2.8% (0.7-5.3%)	420 (100-800)	9000 (2300-17000)
Canada	2016	Male	75 to 79	87% (78-93%)	13% (6.9-22%)	1.9 (1.1-2.9)	1.4% (-0.11-3.4%)	1.6% (-0.34-3.9%)	270 (-59-680)	4800 (-370-12000)
Canada	2016	Male	80 to 84	85% (75-92%)	15% (7.6-25%)	1.7 (1.1-2.5)	0.84% (-0.64-2.4%)	0.77% (-1-2.7%)	160 (-210-560)	2500 (-1900-7300)
Canada	2016	Male	85 to 89	85% (74-92%)	15% (8-26%)	1.7 (1-2.5)	0.46% (-0.94-2%)	0.24% (-1.5-2.1%)	48 (-290-410)	870 (-1800-3800)
Canada	2016	Male	90 to 94	86% (76-93%)	14% (7.3-24%)	1.8 (1-2.8)	0.23% (-1.4-2.1%)	-0.064% (-2-2.1%)	-7.8 (-250-260)	180 (-1100-1600)
Canada	2016	Male	95 plus	87% (76-94%)	13% (5.8-24%)	1.9 (1-3.1)	0.21% (-1.9-3%)	-0.15% (-2.6-3%)	-6.7 (-120-140)	40 (-360-560)
United States	1990	Female	All Ages	60% (57-63%)	40% (37-43%)	1.8 (1.6-2)	2% (1.3-2.6%)	0.46% (-0.71-1.6%)	4800 (-7400-17000)	690000 (450000-930000)
United States	1990	Female	15 to 19	53% (45-61%)	47% (39-55%)	1.8 (1.4-2.2)	3.7% (2.5-4.9%)	9.3% (5.7-14%)	370 (230-550)	41000 (29000-56000)
United States	1990	Female	20 to 24	75% (67-81%)	25% (19-33%)	3.6 (3.1-4.1)	6.1% (4.8-7.4%)	14% (11-18%)	690 (510-870)	86000 (66000-110000)
United States	1990	Female	25 to 29	79% (72-84%)	21% (16-28%)	3.2 (2.8-3.6)	5.3% (4.2-6.3%)	12% (9.1-15%)	790 (610-990)	91000 (71000-110000)
United States	1990	Female	30 to 34	80% (74-85%)	20% (15-26%)	2.8 (2.4-3.3)	4.6% (3.7-5.5%)	10% (8.2-13%)	960 (760-1200)	90000 (72000-110000)
United States	1990	Female	35 to 39	80% (74-85%)	20% (15-26%)	2.6 (2.3-3)	4.5% (3.7-5.5%)	10% (8.6-12%)	1200 (990-1400)	90000 (75000-110000)
United States	1990	Female	40 to 44	78% (72-84%)	22% (16-28%)	2.4 (2.1-2.8)	4.5% (3.6-5.4%)	9.8% (8-12%)	1500 (1200-1700)	90000 (74000-110000)
United States	1990	Female	45 to 49	76% (68-82%)	24% (18-32%)	2.2 (1.9-2.6)	4.1% (3.2-5%)	8% (6.4-9.6%)	1500 (1200-1800)	76000 (60000-91000)
United States	1990	Female	50 to 54	73% (65-79%)	27% (21-35%)	1.9 (1.6-2.3)	3.4% (2.5-4.4%)	5.8% (4.5-7.3%)	1500 (1100-1800)	62000 (47000-79000)
United States	1990	Female	55 to 59	69% (60-77%)	31% (23-40%)	1.7 (1.4-2)	2.4% (1.5-3.3%)	3.8% (2.5-5.2%)	1400 (930-1900)	50000 (32000-68000)
United States	1990	Female	60 to 64	63% (54-72%)	37% (28-46%)	1.4 (1.2-1.7)	1.4% (0.53-2.3%)	2.1% (0.8-3.4%)	1300 (490-2000)	37000 (14000-61000)
United States	1990	Female	65 to 69	57% (48-67%)	43% (33-52%)	1.3 (1-1.6)	0.71% (-0.21-1.6%)	1% (-0.29-2.4%)	910 (-260-2100)	22000 (-6100-50000)
United States	1990	Female	70 to 74	53% (41-63%)	47% (37-59%)	1.1 (0.82-1.4)	0.089% (-0.96-1.1%)	0.11% (-1.3-1.5%)	120 (-1500-1700)	2700 (-28000-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	1990	Female	75 to 79	47% (35-59%)	53% (41-65%)	0.93 (0.67-1.3)	-0.41% (-1.4-0.56%)	-0.61% (-2-0.7%)	-880 (-2900-1000)	-11000 (-40000-16000)
United States	1990	Female	80 to 84	42% (30-53%)	58% (47-70%)	0.81 (0.56-1.1)	-0.7% (-1.7-0.22%)	-0.99% (-2.3-0.28%)	-1600 (-3700-460)	-16000 (-38000-5000)
United States	1990	Female	85 to 89	40% (29-51%)	60% (49-71%)	0.76 (0.5-1.1)	-0.93% (-2-0.043%)	-1.3% (-2.7--0.041%)	-2000 (-4200--64)	-14000 (-30000-690)
United States	1990	Female	90 to 94	41% (30-52%)	59% (48-70%)	0.78 (0.52-1.1)	-1.2% (-2.3--0.072%)	-1.6% (-3.1--0.23%)	-1700 (-3300--250)	-8000 (-16000--520)
United States	1990	Female	95 plus	44% (30-58%)	56% (42-70%)	0.85 (0.53-1.2)	-1.5% (-3--0.21%)	-2.1% (-4--0.44%)	-1200 (-2200--250)	-3400 (-6800--490)
United States	1990	Male	All Ages	72% (69-74%)	28% (26-31%)	3.2 (2.9-3.6)	6.2% (4.9-7.5%)	3.8% (2.1-5.6%)	42000 (23000-63000)	2500000 (1900000-3000000)
United States	1990	Male	15 to 19	57% (50-64%)	43% (36-50%)	2.8 (2.3-3.3)	9.5% (7.1-12%)	15% (11-20%)	1700 (1200-2200)	140000 (110000-180000)
United States	1990	Male	20 to 24	82% (76-87%)	18% (13-24%)	6.4 (5.6-7.3)	19% (15-23%)	28% (22-33%)	4400 (3500-5300)	360000 (300000-430000)
United States	1990	Male	25 to 29	87% (83-90%)	13% (10-17%)	5.7 (5.1-6.5)	15% (13-18%)	22% (18-26%)	4200 (3400-5100)	340000 (280000-400000)
United States	1990	Male	30 to 34	90% (86-92%)	10% (7.7-14%)	5.1 (4.5-5.8)	12% (10-14%)	17% (14-20%)	4100 (3400-4900)	310000 (260000-360000)
United States	1990	Male	35 to 39	90% (87-93%)	10% (7.4-13%)	4.7 (4.1-5.3)	11% (9.5-13%)	15% (13-18%)	4200 (3500-4900)	280000 (230000-320000)
United States	1990	Male	40 to 44	90% (86-92%)	10% (7.6-14%)	4.3 (3.8-4.9)	10% (8.8-12%)	14% (12-16%)	4200 (3500-4900)	240000 (210000-280000)
United States	1990	Male	45 to 49	89% (85-92%)	11% (8-15%)	3.9 (3.4-4.5)	9.1% (7.6-11%)	12% (9.6-14%)	3900 (3200-4800)	200000 (160000-240000)
United States	1990	Male	50 to 54	88% (84-91%)	12% (8.7-16%)	3.6 (3.1-4.1)	7.3% (5.7-9%)	9% (6.7-11%)	3800 (2800-4800)	160000 (130000-200000)
United States	1990	Male	55 to 59	86% (82-90%)	14% (9.7-18%)	3.2 (2.8-3.7)	5.5% (4-7.3%)	6.5% (4.4-8.9%)	3900 (2600-5300)	150000 (110000-190000)
United States	1990	Male	60 to 64	84% (79-89%)	16% (11-21%)	2.9 (2.5-3.4)	3.8% (2.2-5.6%)	4.2% (2.1-6.4%)	4000 (2000-6000)	130000 (73000-180000)
United States	1990	Male	65 to 69	82% (76-87%)	18% (13-24%)	2.7 (2.2-3.1)	2.5% (0.87-4.2%)	2.5% (0.55-4.6%)	3200 (700-5900)	91000 (32000-160000)
United States	1990	Male	70 to 74	79% (72-85%)	21% (15-28%)	2.4 (1.9-2.9)	1.4% (-0.22-3.1%)	1.2% (-0.74-3.3%)	1800 (-1100-5000)	47000 (-7600-100000)
United States	1990	Male	75 to 79	76% (67-83%)	24% (17-33%)	2.1 (1.6-2.5)	0.61% (-0.91-2.2%)	0.31% (-1.5-2.2%)	490 (-2400-3600)	17000 (-25000-59000)
United States	1990	Male	80 to 84	73% (62-80%)	27% (20-38%)	1.8 (1.4-2.2)	0.11% (-1.3-1.6%)	-0.25% (-2-1.5%)	-340 (-2800-2100)	2000 (-23000-27000)
United States	1990	Male	85 to 89	71% (61-79%)	29% (21-39%)	1.7 (1.3-2.2)	-0.21% (-1.7-1.2%)	-0.66% (-2.4-1.1%)	-610 (-2200-1000)	-1800 (-14000-9900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	1990	Male	90 to 94	72% (63-80%)	28% (20-37%)	1.7 (1.3-2.2)	-0.59% (-2.3-1%)	-1.2% (-3.3-0.81%)	-510 (-1400-360)	-1600 (-6200-2800)
United States	1990	Male	95 plus	74% (62-83%)	26% (17-38%)	1.7 (1.2-2.3)	-0.79% (-2.7-1.1%)	-1.5% (-3.9-0.75%)	-250 (-640-120)	-520 (-1800-720)
United States	1995	Female	All Ages	60% (57-63%)	40% (37-43%)	1.7 (1.5-1.9)	1.9% (1.3-2.5%)	0.37% (-0.72-1.5%)	4200 (-8200-17000)	690000 (460000-940000)
United States	1995	Female	15 to 19	52% (44-61%)	48% (39-56%)	1.7 (1.4-2.1)	3.5% (2.5-4.7%)	8.9% (5.5-13%)	350 (220-510)	41000 (29000-56000)
United States	1995	Female	20 to 24	74% (65-80%)	26% (20-35%)	3.5 (3-4)	5.8% (4.6-7%)	13% (9.7-17%)	570 (420-730)	77000 (59000-98000)
United States	1995	Female	25 to 29	78% (71-83%)	22% (17-29%)	3.1 (2.7-3.5)	4.9% (4-5.8%)	10% (7.7-13%)	610 (460-780)	78000 (60000-97000)
United States	1995	Female	30 to 34	80% (73-85%)	20% (15-27%)	2.7 (2.4-3.2)	4.3% (3.5-5.1%)	8.8% (6.9-11%)	830 (650-1000)	84000 (67000-100000)
United States	1995	Female	35 to 39	80% (74-85%)	20% (15-26%)	2.6 (2.2-3)	4.3% (3.6-5.2%)	9.1% (7.4-11%)	1300 (1000-1500)	97000 (80000-110000)
United States	1995	Female	40 to 44	78% (73-84%)	22% (16-27%)	2.4 (2.1-2.8)	4.3% (3.5-5.2%)	8.9% (7.3-10%)	1600 (1300-1900)	99000 (82000-120000)
United States	1995	Female	45 to 49	76% (69-82%)	24% (18-31%)	2.2 (1.9-2.6)	3.9% (3.1-4.7%)	7.5% (6.1-9%)	1800 (1400-2100)	91000 (74000-110000)
United States	1995	Female	50 to 54	73% (66-79%)	27% (21-34%)	1.9 (1.6-2.3)	3.1% (2.3-4%)	5.5% (4.2-6.9%)	1600 (1200-2000)	69000 (52000-87000)
United States	1995	Female	55 to 59	69% (60-76%)	31% (24-40%)	1.6 (1.3-1.9)	2.2% (1.4-3.1%)	3.5% (2.3-4.8%)	1300 (860-1800)	48000 (32000-67000)
United States	1995	Female	60 to 64	63% (54-72%)	37% (28-46%)	1.4 (1.1-1.7)	1.3% (0.55-2.2%)	1.9% (0.74-3.1%)	1100 (410-1700)	32000 (13000-53000)
United States	1995	Female	65 to 69	58% (49-67%)	42% (33-51%)	1.3 (0.99-1.6)	0.72% (-0.14-1.6%)	0.98% (-0.22-2.2%)	830 (-190-1900)	21000 (-4200-46000)
United States	1995	Female	70 to 74	54% (42-64%)	46% (36-58%)	1.1 (0.84-1.4)	0.18% (-0.72-1.1%)	0.21% (-1.1-1.4%)	250 (-1300-1700)	5900 (-23000-34000)
United States	1995	Female	75 to 79	48% (37-60%)	52% (40-63%)	0.95 (0.69-1.3)	-0.29% (-1.2-0.6%)	-0.48% (-1.7-0.71%)	-730 (-2500-1100)	-8700 (-35000-18000)
United States	1995	Female	80 to 84	43% (31-53%)	57% (47-69%)	0.81 (0.56-1.1)	-0.59% (-1.5-0.25%)	-0.86% (-2-0.27%)	-1600 (-3600-480)	-15000 (-36000-6200)
United States	1995	Female	85 to 89	40% (29-51%)	60% (49-71%)	0.75 (0.5-1)	-0.82% (-1.8-0.064%)	-1.2% (-2.5--0.0043%)	-2100 (-4500--7.8)	-14000 (-32000-1100)
United States	1995	Female	90 to 94	41% (30-52%)	59% (48-70%)	0.78 (0.52-1.1)	-1.1% (-2.2--0.078%)	-1.5% (-2.9--0.22%)	-2000 (-3800--290)	-9200 (-18000--660)
United States	1995	Female	95 plus	45% (31-58%)	55% (42-69%)	0.85 (0.53-1.2)	-1.5% (-2.9--0.21%)	-2% (-3.9--0.44%)	-1400 (-2700--300)	-4100 (-8000--580)
United States	1995	Male	All Ages	72% (69-74%)	28% (26-31%)	3.1 (2.8-3.4)	5.9% (4.7-7%)	3.5% (1.9-5.2%)	42000 (22000-61000)	240000 (190000-300000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	1995	Male	15 to 19	56% (49-63%)	44% (37-51%)	2.6 (2.2-3.1)	8.9% (6.7-11%)	14% (10-19%)	1600 (1100-2100)	140000 (110000-180000)
United States	1995	Male	20 to 24	81% (76-86%)	19% (14-24%)	6.1 (5.3-6.9)	18% (15-21%)	27% (21-32%)	3900 (3100-4700)	330000 (270000-390000)
United States	1995	Male	25 to 29	87% (83-90%)	13% (10-17%)	5.5 (4.9-6.2)	14% (12-17%)	21% (17-25%)	3300 (2700-4000)	280000 (230000-330000)
United States	1995	Male	30 to 34	89% (86-92%)	11% (7.8-14%)	4.9 (4.3-5.6)	11% (9.1-12%)	14% (12-17%)	3500 (2800-4100)	270000 (220000-320000)
United States	1995	Male	35 to 39	90% (86-92%)	10% (7.6-14%)	4.5 (4-5.1)	9.9% (8.3-11%)	13% (11-15%)	4000 (3300-4700)	280000 (230000-320000)
United States	1995	Male	40 to 44	89% (86-92%)	11% (7.7-14%)	4.1 (3.6-4.7)	9.5% (8-11%)	12% (10-14%)	4600 (3800-5300)	270000 (230000-310000)
United States	1995	Male	45 to 49	89% (85-92%)	11% (7.9-15%)	3.8 (3.3-4.3)	8.7% (7.3-10%)	11% (9.4-13%)	4800 (4000-5700)	240000 (210000-290000)
United States	1995	Male	50 to 54	88% (84-91%)	12% (8.7-16%)	3.5 (3-4)	7.1% (5.7-8.6%)	8.9% (6.9-11%)	4300 (3300-5400)	190000 (150000-230000)
United States	1995	Male	55 to 59	86% (82-90%)	14% (9.9-18%)	3.1 (2.6-3.6)	5.4% (4-7%)	6.5% (4.5-8.6%)	3800 (2600-5100)	150000 (110000-190000)
United States	1995	Male	60 to 64	84% (79-88%)	16% (12-21%)	2.8 (2.4-3.3)	3.9% (2.4-5.5%)	4.3% (2.4-6.4%)	3600 (2000-5300)	120000 (72000-170000)
United States	1995	Male	65 to 69	82% (76-87%)	18% (13-24%)	2.6 (2.2-3)	2.5% (1.1-4.1%)	2.6% (0.73-4.6%)	3200 (880-5500)	89000 (37000-150000)
United States	1995	Male	70 to 74	79% (73-85%)	21% (15-27%)	2.3 (1.9-2.8)	1.5% (0.0083-3.1%)	1.4% (-0.47-3.3%)	2200 (-730-5200)	54000 (300-110000)
United States	1995	Male	75 to 79	77% (69-83%)	23% (17-31%)	2 (1.6-2.5)	0.7% (-0.68-2.2%)	0.42% (-1.3-2.3%)	700 (-2200-3800)	21000 (-20000-65000)
United States	1995	Male	80 to 84	73% (64-81%)	27% (19-36%)	1.8 (1.4-2.2)	0.17% (-1.2-1.5%)	-0.17% (-1.9-1.4%)	-270 (-2900-2300)	3400 (-23000-30000)
United States	1995	Male	85 to 89	71% (62-79%)	29% (21-38%)	1.6 (1.2-2.1)	-0.18% (-1.6-1.2%)	-0.6% (-2.3-1%)	-670 (-2600-1200)	-1800 (-16000-13000)
United States	1995	Male	90 to 94	72% (63-80%)	28% (20-37%)	1.7 (1.2-2.1)	-0.56% (-2.2-0.99%)	-1.1% (-3.1-0.78%)	-600 (-1700-420)	-1900 (-7300-3400)
United States	1995	Male	95 plus	74% (63-83%)	26% (17-37%)	1.7 (1.2-2.2)	-0.81% (-2.6-0.9%)	-1.5% (-3.8-0.61%)	-270 (-690-110)	-600 (-1900-680)
United States	2000	Female	All Ages	60% (58-63%)	40% (37-42%)	1.7 (1.5-1.9)	1.8% (1.2-2.5%)	0.37% (-0.76-1.5%)	4500 (-9200-19000)	710000 (460000-980000)
United States	2000	Female	15 to 19	52% (44-60%)	48% (40-56%)	1.7 (1.3-2.1)	3.2% (2.2-4.2%)	8.8% (5.3-13%)	340 (210-500)	41000 (29000-54000)
United States	2000	Female	20 to 24	73% (65-80%)	27% (20-35%)	3.5 (3-4)	5.1% (4.2-6.2%)	13% (9.6-17%)	560 (420-720)	71000 (56000-88000)
United States	2000	Female	25 to 29	77% (71-82%)	23% (18-29%)	3.1 (2.7-3.5)	4.5% (3.6-5.3%)	11% (7.9-13%)	540 (400-680)	67000 (53000-82000)
United States	2000	Female	30 to 34	79% (73-84%)	21% (16-27%)	2.8 (2.4-3.2)	4.1% (3.3-4.9%)	9.2% (7.2-11%)	700 (540-850)	71000 (57000-88000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	2000	Female	35 to 39	80% (75-85%)	20% (15-25%)	2.6 (2.2-3)	4.2% (3.4-5%)	9.3% (7.5-11%)	1200 (960-1400)	92000 (76000-110000)
United States	2000	Female	40 to 44	79% (74-84%)	21% (16-26%)	2.5 (2.1-2.8)	4.3% (3.5-5.2%)	9% (7.3-11%)	1800 (1400-2100)	110000 (89000-130000)
United States	2000	Female	45 to 49	77% (70-83%)	23% (17-30%)	2.3 (2-2.7)	4% (3.2-4.9%)	7.8% (6.3-9.3%)	2000 (1600-2400)	100000 (85000-130000)
United States	2000	Female	50 to 54	75% (68-80%)	25% (20-32%)	2 (1.7-2.4)	3.2% (2.4-4.2%)	5.7% (4.4-7.3%)	2000 (1500-2500)	87000 (66000-110000)
United States	2000	Female	55 to 59	71% (63-78%)	29% (22-37%)	1.7 (1.4-2)	2.3% (1.5-3.2%)	3.7% (2.4-5.1%)	1600 (1000-2200)	58000 (38000-80000)
United States	2000	Female	60 to 64	65% (57-72%)	35% (28-43%)	1.5 (1.2-1.8)	1.3% (0.54-2.2%)	2% (0.77-3.3%)	1100 (430-1800)	34000 (13000-55000)
United States	2000	Female	65 to 69	59% (48-69%)	41% (31-52%)	1.3 (1-1.6)	0.76% (-0.089-1.7%)	1.1% (-0.095-2.3%)	820 (-72-1700)	20000 (-2300-44000)
United States	2000	Female	70 to 74	55% (44-65%)	45% (35-56%)	1.2 (0.88-1.5)	0.37% (-0.65-1.2%)	0.37% (-0.87-1.6%)	420 (-1000-1800)	8400 (-19000-36000)
United States	2000	Female	75 to 79	51% (40-62%)	49% (38-60%)	1 (0.76-1.4)	-0.21% (-1.1-0.68%)	-0.34% (-1.5-0.85%)	-560 (-2400-1400)	-6700 (-33000-22000)
United States	2000	Female	80 to 84	46% (34-57%)	54% (43-66%)	0.91 (0.64-1.2)	-0.55% (-1.4-0.31%)	-0.79% (-1.9-0.37%)	-1500 (-3700-700)	-15000 (-38000-8300)
United States	2000	Female	85 to 89	43% (32-55%)	57% (45-68%)	0.83 (0.56-1.2)	-0.8% (-1.8-0.085%)	-1.1% (-2.5-0.033%)	-2300 (-5100-68)	-16000 (-35000-1700)
United States	2000	Female	90 to 94	45% (33-56%)	55% (44-67%)	0.87 (0.6-1.2)	-1.1% (-2.2--0.056%)	-1.5% (-3--0.2%)	-2400 (-4600--310)	-11000 (-22000--540)
United States	2000	Female	95 plus	50% (36-63%)	50% (37-64%)	0.98 (0.62-1.4)	-1.6% (-3.1--0.19%)	-2.1% (-4.2--0.39%)	-1700 (-3400--310)	-5200 (-10000--630)
United States	2000	Male	All Ages	72% (70-75%)	28% (25-30%)	3 (2.7-3.4)	6% (4.9-7.1%)	3.8% (2.2-5.4%)	44000 (25000-63000)	2500000 (2000000-3000000)
United States	2000	Male	15 to 19	54% (48-61%)	46% (39-52%)	2.5 (2.1-3)	7.9% (5.9-10%)	14% (9.8-19%)	1300 (950-1800)	120000 (94000-160000)
United States	2000	Male	20 to 24	81% (75-86%)	19% (14-25%)	6 (5.3-6.9)	16% (13-19%)	26% (21-32%)	3500 (2800-4200)	290000 (240000-350000)
United States	2000	Male	25 to 29	87% (83-90%)	13% (10-17%)	5.5 (4.8-6.2)	23% (11-16%)	23% (18-28%)	2900 (2300-3500)	240000 (200000-290000)
United States	2000	Male	30 to 34	89% (86-92%)	11% (7.9-14%)	4.9 (4.3-5.5)	11% (9.6-13%)	18% (15-22%)	2800 (2300-3400)	220000 (190000-270000)
United States	2000	Male	35 to 39	90% (87-92%)	10% (7.6-13%)	4.5 (3.9-5.1)	11% (8.8-12%)	16% (13-19%)	3700 (3100-4400)	260000 (210000-300000)
United States	2000	Male	40 to 44	90% (87-92%)	10% (7.6-13%)	4.1 (3.6-4.7)	10% (8.5-12%)	14% (12-17%)	4900 (4000-5700)	280000 (240000-330000)
United States	2000	Male	45 to 49	89% (86-92%)	11% (7.6-14%)	3.8 (3.3-4.4)	9.4% (7.9-11%)	13% (11-15%)	5700 (4700-6600)	290000 (240000-330000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	2000	Male	50 to 54	89% (85-92%)	11% (8.4-15%)	3.5 (3-4)	7.8% (6.4-9.3%)	10% (8-12%)	5600 (4400-6700)	250000 (200000-300000)
United States	2000	Male	55 to 59	87% (83-91%)	13% (9.2-17%)	3.2 (2.7-3.6)	6% (4.5-7.4%)	7.2% (5.1-9.3%)	4600 (3300-6000)	180000 (140000-230000)
United States	2000	Male	60 to 64	85% (80-89%)	15% (11-20%)	2.8 (2.4-3.3)	4.3% (2.9-5.8%)	4.9% (2.9-6.8%)	3800 (2300-5400)	130000 (86000-170000)
United States	2000	Male	65 to 69	82% (75-87%)	18% (13-25%)	2.6 (2.1-3)	2.9% (1.6-4.4%)	3.1% (1.3-5%)	3200 (1400-5200)	91000 (49000-140000)
United States	2000	Male	70 to 74	80% (73-85%)	20% (15-27%)	2.3 (1.9-2.8)	1.8% (0.45-3.3%)	1.8% (0.041-3.6%)	2600 (58-5200)	62000 (15000-110000)
United States	2000	Male	75 to 79	78% (71-84%)	22% (16-29%)	2.1 (1.7-2.5)	0.97% (-0.39-2.4%)	0.75% (-0.95-2.5%)	1300 (-1600-4300)	30000 (-12000-75000)
United States	2000	Male	80 to 84	75% (66-82%)	25% (18-34%)	1.8 (1.4-2.3)	0.35% (-0.95-1.7%)	0.03% (-1.6-1.7%)	50 (-2600-2800)	7400 (-20000-37000)
United States	2000	Male	85 to 89	73% (65-81%)	27% (19-35%)	1.7 (1.3-2.1)	-0.079% (-1.4-1.2%)	-0.48% (-2.2-1.1%)	-620 (-2800-1500)	-900 (-17000-15000)
United States	2000	Male	90 to 94	75% (66-81%)	25% (19-34%)	1.7 (1.3-2.2)	-0.52% (-2.1-0.93%)	-1% (-3-0.76%)	-670 (-1900-490)	-2100 (-8300-3800)
United States	2000	Male	95 plus	76% (66-83%)	24% (17-34%)	1.7 (1.2-2.3)	-0.78% (-2.6-0.85%)	-1.4% (-3.7-0.63%)	-300 (-780-130)	-670 (-2200-760)
United States	2005	Female	All Ages	61% (58-63%)	39% (37-42%)	1.8 (1.6-2)	2.1% (1.5-2.8%)	0.87% (-0.13-1.9%)	11000 (-1600-23000)	840000 (600000-1100000)
United States	2005	Female	15 to 19	50% (43-58%)	50% (42-57%)	1.7 (1.4-2.1)	3.1% (2.2-4%)	8.8% (5.3-13%)	340 (210-490)	41000 (30000-54000)
United States	2005	Female	20 to 24	72% (65-79%)	28% (21-35%)	3.7 (3.1-4.2)	5.2% (4.3-6.2%)	13% (9.6-16%)	630 (470-780)	78000 (62000-97000)
United States	2005	Female	25 to 29	77% (71-82%)	23% (18-29%)	3.2 (2.8-3.7)	4.6% (3.8-5.4%)	11% (8.3-13%)	580 (450-710)	70000 (56000-86000)
United States	2005	Female	30 to 34	79% (73-84%)	21% (16-27%)	2.9 (2.5-3.3)	4.1% (3.3-4.9%)	9.3% (7.3-11%)	680 (530-820)	69000 (56000-84000)
United States	2005	Female	35 to 39	79% (74-84%)	21% (16-26%)	2.7 (2.4-3.2)	4.2% (3.4-5%)	9.4% (7.7-11%)	1100 (880-1300)	84000 (68000-100000)
United States	2005	Female	40 to 44	79% (74-84%)	21% (16-26%)	2.6 (2.3-3)	4.4% (3.6-5.3%)	9.3% (7.7-11%)	1800 (1500-2200)	110000 (92000-130000)
United States	2005	Female	45 to 49	78% (71-83%)	22% (17-29%)	2.4 (2.1-2.8)	4.3% (3.5-5.2%)	8.3% (6.8-9.8%)	2500 (2000-2900)	130000 (100000-150000)
United States	2005	Female	50 to 54	75% (69-81%)	25% (19-31%)	2.2 (1.8-2.5)	3.7% (2.9-4.6%)	6.6% (5.3-8%)	2600 (2100-3100)	110000 (87000-140000)
United States	2005	Female	55 to 59	71% (64-78%)	29% (22-36%)	1.9 (1.6-2.2)	2.7% (1.9-3.5%)	4.6% (3.4-5.8%)	2300 (1700-2900)	82000 (58000-110000)
United States	2005	Female	60 to 64	66% (58-73%)	34% (27-42%)	1.6 (1.3-1.9)	1.7% (0.87-2.5%)	2.7% (1.6-3.8%)	1600 (960-2300)	47000 (25000-70000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	2005	Female	65 to 69	58% (49-66%)	42% (34-51%)	1.3 (1.1-1.6)	1% (0.26-1.8%)	1.6% (0.64-2.6%)	1200 (470-1900)	27000 (6700-49000)
United States	2005	Female	70 to 74	53% (45-61%)	47% (39-55%)	1.2 (0.94-1.5)	0.55% (-0.19-1.4%)	0.88% (-0.13-1.9%)	880 (-130-1900)	16000 (-5400-38000)
United States	2005	Female	75 to 79	50% (42-59%)	50% (41-58%)	1.1 (0.86-1.3)	0.11% (-0.63-0.9%)	0.17% (-0.8-1.2%)	240 (-1200-1700)	3400 (-19000-27000)
United States	2005	Female	80 to 84	47% (36-57%)	53% (43-64%)	0.98 (0.71-1.3)	-0.23% (-1-0.56%)	-0.33% (-1.4-0.69%)	-650 (-2600-1300)	-6400 (-29000-16000)
United States	2005	Female	85 to 89	45% (35-55%)	55% (45-65%)	0.93 (0.66-1.2)	-0.53% (-1.4-0.29%)	-0.78% (-1.9-0.3%)	-1600 (-4000-620)	-11000 (-30000-5900)
United States	2005	Female	90 to 94	47% (37-56%)	53% (44-63%)	0.97 (0.7-1.3)	-0.85% (-1.9-0.14%)	-1.2% (-2.5-0.086%)	-2000 (-4100-140)	-9000 (-21000-1500)
United States	2005	Female	95 plus	52% (39-63%)	48% (37-61%)	1.1 (0.71-1.5)	-1.3% (-2.7-0.014%)	-1.8% (-3.5--0.16%)	-1500 (-3100--140)	-4600 (-9600-52)
United States	2005	Male	All Ages	72% (70-75%)	28% (25-30%)	3.2 (2.8-3.5)	6.6% (5.5-7.7%)	4.7% (3.1-6.2%)	56000 (37000-74000)	280000 (230000-340000)
United States	2005	Male	15 to 19	53% (46-60%)	47% (40-54%)	2.6 (2.1-3)	7.7% (5.9-9.8%)	14% (9.8-18%)	1300 (970-1800)	13000 (96000-160000)
United States	2005	Male	20 to 24	80% (75-85%)	20% (15-25%)	6.3 (5.4-7.2)	16% (13-19%)	25% (20-30%)	4000 (3200-4800)	33000 (270000-390000)
United States	2005	Male	25 to 29	87% (83-90%)	13% (10-17%)	5.7 (5-6.5)	14% (12-17%)	22% (18-27%)	3200 (2600-3900)	26000 (22000-310000)
United States	2005	Male	30 to 34	89% (86-92%)	11% (8-14%)	5.1 (4.4-5.8)	12% (10-14%)	19% (15-23%)	2900 (2300-3400)	22000 (19000-260000)
United States	2005	Male	35 to 39	90% (86-92%)	10% (7.6-14%)	4.7 (4.1-5.3)	11% (9.2-13%)	17% (14-20%)	3400 (2800-4000)	24000 (20000-270000)
United States	2005	Male	40 to 44	90% (86-92%)	10% (7.7-14%)	4.3 (3.8-4.9)	10% (8.9-12%)	15% (13-17%)	5000 (4100-5800)	29000 (25000-340000)
United States	2005	Male	45 to 49	89% (86-92%)	11% (7.6-14%)	4 (3.5-4.6)	10% (8.6-12%)	14% (12-16%)	6800 (5700-7800)	34000 (29000-390000)
United States	2005	Male	50 to 54	89% (85-92%)	11% (8.3-15%)	3.7 (3.2-4.2)	8.9% (7.4-10%)	12% (9.6-14%)	7500 (6200-8800)	33000 (27000-380000)
United States	2005	Male	55 to 59	87% (83-91%)	13% (9.4-17%)	3.3 (2.8-3.8)	7% (5.6-8.5%)	8.9% (6.9-11%)	7000 (5400-8500)	26000 (21000-320000)
United States	2005	Male	60 to 64	85% (80-89%)	15% (11-20%)	3 (2.6-3.4)	5.1% (3.7-6.5%)	6% (4.2-8%)	5300 (3600-7000)	17000 (13000-220000)
United States	2005	Male	65 to 69	82% (76-86%)	18% (14-24%)	2.7 (2.3-3.1)	3.6% (2.3-4.9%)	4.1% (2.4-5.8%)	4000 (2400-5700)	11000 (7300-160000)
United States	2005	Male	70 to 74	80% (74-84%)	20% (16-26%)	2.4 (2-2.9)	2.4% (1.1-3.7%)	2.6% (0.94-4.2%)	3200 (1200-5100)	7300 (3400-110000)
United States	2005	Male	75 to 79	78% (72-83%)	22% (17-28%)	2.1 (1.8-2.6)	1.5% (0.29-2.7%)	1.4% (-0.13-2.9%)	2200 (-210-4600)	43000 (8600-81000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	2005	Male	80 to 84	76% (67-82%)	24% (18-33%)	1.9 (1.5-2.4)	0.83% (-0.31-2.1%)	0.59% (-0.89-2.1%)	1000 (-1500-3600)	19000 (-7200-48000)
United States	2005	Male	85 to 89	75% (67-80%)	25% (20-33%)	1.8 (1.5-2.3)	0.34% (-0.9-1.7%)	-0.027% (-1.6-1.5%)	-37 (-2200-2100)	4400 (-12000-21000)
United States	2005	Male	90 to 94	75% (68-81%)	25% (19-32%)	1.9 (1.4-2.3)	-0.13% (-1.6-1.3%)	-0.61% (-2.4-1.1%)	-460 (-1800-800)	-600 (-7300-6000)
United States	2005	Male	95 plus	76% (67-83%)	24% (17-33%)	1.8 (1.3-2.4)	-0.4% (-2.1-1.3%)	-0.99% (-3.1-1%)	-240 (-730-250)	-390 (-2000-1300)
United States	2010	Female	All Ages	59% (57-62%)	41% (38-43%)	1.9 (1.6-2.1)	2.3% (1.7-3%)	1.3% (0.33-2.4%)	17000 (4000-30000)	950000 (680000-1200000)
United States	2010	Female	15 to 19	37% (31-44%)	63% (56-69%)	1.3 (1-1.6)	2.4% (1.8-3.1%)	7.1% (4.4-10%)	220 (140-320)	32000 (24000-42000)
United States	2010	Female	20 to 24	72% (64-78%)	28% (22-36%)	3.7 (3.1-4.4)	5.1% (4.2-6.1%)	13% (9.9-16%)	620 (470-780)	79000 (62000-97000)
United States	2010	Female	25 to 29	78% (72-83%)	22% (17-28%)	3.4 (2.9-4)	4.7% (3.9-5.6%)	11% (8.5-14%)	660 (500-810)	78000 (63000-95000)
United States	2010	Female	30 to 34	79% (73-84%)	21% (16-27%)	3 (2.5-3.5)	4.2% (3.5-5.1%)	9.8% (7.6-12%)	720 (560-870)	72000 (58000-89000)
United States	2010	Female	35 to 39	79% (74-84%)	21% (16-26%)	2.8 (2.4-3.4)	4.2% (3.5-5.1%)	10% (8.1-12%)	1000 (840-1200)	81000 (66000-97000)
United States	2010	Female	40 to 44	79% (74-84%)	21% (16-26%)	2.7 (2.3-3.2)	4.4% (3.6-5.4%)	10% (8.2-12%)	1700 (1400-1900)	100000 (83000-120000)
United States	2010	Female	45 to 49	78% (71-83%)	22% (17-29%)	2.5 (2.1-3)	4.5% (3.6-5.6%)	9% (7.5-11%)	2600 (2200-3100)	130000 (110000-160000)
United States	2010	Female	50 to 54	75% (69-81%)	25% (19-31%)	2.3 (1.9-2.7)	4.1% (3.2-5.1%)	7.4% (5.9-8.9%)	3200 (2500-3800)	140000 (110000-170000)
United States	2010	Female	55 to 59	72% (65-79%)	28% (21-35%)	1.9 (1.6-2.3)	3.1% (2.2-4%)	5.4% (4.1-6.7%)	2900 (2200-3600)	100000 (75000-140000)
United States	2010	Female	60 to 64	66% (59-73%)	34% (27-41%)	1.7 (1.4-2)	2% (1.1-2.9%)	3.4% (2.3-4.6%)	2400 (1600-3200)	68000 (40000-99000)
United States	2010	Female	65 to 69	59% (51-67%)	41% (33-49%)	1.4 (1.1-1.8)	1.3% (0.53-2.1%)	2.2% (1.2-3.2%)	1800 (940-2600)	41000 (17000-65000)
United States	2010	Female	70 to 74	54% (45-61%)	46% (39-55%)	1.2 (0.97-1.5)	0.82% (0.075-1.6%)	1.4% (0.42-2.3%)	1300 (410-2300)	23000 (2200-45000)
United States	2010	Female	75 to 79	50% (42-59%)	50% (41-58%)	1.1 (0.88-1.4)	0.39% (-0.33-1.2%)	0.58% (-0.31-1.6%)	760 (-410-2000)	11000 (-9000-33000)
United States	2010	Female	80 to 84	47% (36-57%)	53% (43-64%)	1 (0.74-1.3)	0.072% (-0.67-0.87%)	0.084% (-0.87-1%)	150 (-1600-1900)	2000 (-18000-24000)
United States	2010	Female	85 to 89	46% (35-56%)	54% (44-65%)	0.98 (0.69-1.3)	-0.22% (-1-0.65%)	-0.37% (-1.4-0.66%)	-800 (-3000-1400)	-4700 (-22000-14000)
United States	2010	Female	90 to 94	47% (37-56%)	53% (44-63%)	1 (0.72-1.3)	-0.53% (-1.5-0.47%)	-0.81% (-2-0.4%)	-1400 (-3400-680)	-5900 (-17000-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	2010	Female	95 plus	51% (38-63%)	49% (37-62%)	1.1 (0.72-1.5)	-0.89% (-2.2-0.42%)	-1.3% (-2.9-0.28%)	-1300 (-2800-270)	-3600 (-8700-1800)
United States	2010	Male	All Ages	72% (70-75%)	28% (25-30%)	3.2 (2.8-3.7)	6.8% (5.6-8%)	5.2% (3.7-6.8%)	64000 (46000-84000)	300000 (250000-360000)
United States	2010	Male	15 to 19	48% (42-55%)	52% (45-58%)	2.4 (1.9-2.9)	6.7% (5.2-8.4%)	13% (9.2-17%)	1000 (720-1300)	100000 (78000-130000)
United States	2010	Male	20 to 24	81% (75-86%)	19% (14-25%)	6.4 (5.5-7.6)	15% (12-18%)	25% (20-30%)	3600 (2900-4400)	300000 (250000-360000)
United States	2010	Male	25 to 29	87% (83-90%)	13% (10-17%)	5.8 (5-6.8)	14% (11-16%)	22% (18-27%)	3300 (2600-3900)	270000 (220000-320000)
United States	2010	Male	30 to 34	89% (86-92%)	11% (8-14%)	5.2 (4.4-6)	12% (10-14%)	19% (16-23%)	2900 (2300-3500)	230000 (190000-270000)
United States	2010	Male	35 to 39	89% (86-92%)	11% (7.8-14%)	4.8 (4.1-5.6)	11% (9.3-13%)	18% (14-21%)	3200 (2600-3800)	220000 (180000-260000)
United States	2010	Male	40 to 44	89% (86-92%)	11% (7.8-14%)	4.4 (3.7-5.1)	11% (8.9-12%)	16% (13-19%)	4300 (3500-5000)	250000 (210000-300000)
United States	2010	Male	45 to 49	89% (86-92%)	11% (7.8-14%)	4.1 (3.5-4.8)	10% (8.8-12%)	15% (13-17%)	6700 (5600-7800)	330000 (280000-390000)
United States	2010	Male	50 to 54	89% (85-92%)	11% (8.4-15%)	3.7 (3.1-4.3)	9.6% (8-11%)	13% (11-15%)	8700 (7200-10000)	380000 (310000-440000)
United States	2010	Male	55 to 59	87% (83-91%)	13% (9.4-17%)	3.4 (2.8-4)	7.9% (6.4-9.5%)	10% (8.2-12%)	8900 (7100-11000)	330000 (270000-400000)
United States	2010	Male	60 to 64	85% (81-89%)	15% (11-19%)	3.1 (2.6-3.6)	5.8% (4.4-7.3%)	7.3% (5.4-9.2%)	7600 (5600-9600)	240000 (190000-310000)
United States	2010	Male	65 to 69	82% (77-86%)	18% (14-23%)	2.8 (2.3-3.3)	4.2% (2.9-5.6%)	4.9% (3.3-6.7%)	5400 (3600-7300)	150000 (100000-200000)
United States	2010	Male	70 to 74	80% (74-84%)	20% (16-26%)	2.5 (2-3)	2.8% (1.6-4.2%)	3.2% (1.7-4.8%)	3900 (2000-5900)	88000 (50000-130000)
United States	2010	Male	75 to 79	78% (72-83%)	22% (17-28%)	2.2 (1.8-2.7)	1.9% (0.73-3.1%)	1.9% (0.47-3.5%)	2700 (670-5000)	52000 (20000-88000)
United States	2010	Male	80 to 84	76% (67-82%)	24% (18-33%)	2 (1.6-2.5)	1.3% (0.18-2.4%)	1.1% (-0.26-2.5%)	1800 (-440-4300)	29000 (4300-58000)
United States	2010	Male	85 to 89	75% (67-81%)	25% (19-33%)	1.9 (1.5-2.4)	0.76% (-0.38-1.9%)	0.44% (-0.95-1.8%)	670 (-1400-2800)	11000 (-5500-29000)
United States	2010	Male	90 to 94	75% (68-81%)	25% (19-32%)	1.9 (1.4-2.4)	0.28% (-1-1.7%)	-0.16% (-1.8-1.5%)	-140 (-1600-1300)	1600 (-5900-9500)
United States	2010	Male	95 plus	76% (67-83%)	24% (17-33%)	1.9 (1.3-2.5)	0.036% (-1.5-1.7%)	-0.51% (-2.4-1.5%)	-150 (-710-430)	46 (-1800-2100)
United States	2016	Female	All Ages	60% (57-63%)	40% (37-43%)	1.9 (1.4-2.4)	2.3% (1.5-3.3%)	1.4% (0.2-2.7%)	19000 (2700-37000)	100000 (66000-140000)
United States	2016	Female	15 to 19	35% (29-43%)	65% (57-71%)	1.2 (0.85-1.7)	2.4% (1.8-3.1%)	7% (4.2-11%)	200 (120-310)	30000 (22000-40000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	2016	Female	20 to 24	72% (64-79%)	28% (21-36%)	3.7 (2.8-4.9)	5.1% (4-6.3%)	13% (9.3-17%)	670 (480-890)	84000 (64000-100000)
United States	2016	Female	25 to 29	79% (73-84%)	21% (16-27%)	3.4 (2.6-4.4)	4.8% (3.7-5.9%)	11% (8.3-15%)	740 (540-970)	84000 (65000-110000)
United States	2016	Female	30 to 34	79% (73-85%)	21% (15-27%)	3 (2.2-3.9)	4.4% (3.4-5.4%)	10% (7.5-13%)	870 (650-1100)	83000 (64000-110000)
United States	2016	Female	35 to 39	80% (74-85%)	20% (15-26%)	2.8 (2.1-3.7)	4.4% (3.4-5.5%)	10% (7.8-13%)	1100 (860-1400)	85000 (66000-110000)
United States	2016	Female	40 to 44	79% (73-84%)	21% (16-27%)	2.7 (2-3.4)	4.5% (3.4-5.8%)	10% (7.9-13%)	1600 (1300-2100)	100000 (77000-130000)
United States	2016	Female	45 to 49	78% (71-83%)	22% (17-29%)	2.5 (1.9-3.3)	4.6% (3.4-5.9%)	9.3% (7.3-11%)	2400 (1900-2900)	120000 (90000-150000)
United States	2016	Female	50 to 54	76% (70-81%)	24% (19-30%)	2.3 (1.7-2.9)	4.2% (3-5.6%)	7.6% (5.7-9.5%)	3300 (2500-4200)	140000 (100000-180000)
United States	2016	Female	55 to 59	73% (66-79%)	27% (21-34%)	2 (1.4-2.5)	3.2% (2.1-4.5%)	5.6% (3.9-7.4%)	3500 (2500-4600)	120000 (81000-170000)
United States	2016	Female	60 to 64	67% (60-74%)	33% (26-40%)	1.7 (1.2-2.2)	2.1% (1.1-3.2%)	3.6% (2.1-5.1%)	2900 (1600-4100)	82000 (42000-130000)
United States	2016	Female	65 to 69	60% (52-68%)	40% (32-48%)	1.5 (1.1-1.9)	1.4% (0.48-2.4%)	2.4% (1.1-3.7%)	2500 (1200-3900)	57000 (20000-98000)
United States	2016	Female	70 to 74	55% (44-64%)	45% (36-56%)	1.3 (0.9-1.8)	0.91% (0.03-2%)	1.5% (0.36-2.8%)	1800 (430-3400)	32000 (1100-69000)
United States	2016	Female	75 to 79	51% (40-63%)	49% (37-60%)	1.1 (0.77-1.6)	0.46% (-0.37-1.5%)	0.69% (-0.41-1.9%)	960 (-560-2700)	14000 (-11000-43000)
United States	2016	Female	80 to 84	48% (35-58%)	52% (42-65%)	1 (0.67-1.4)	0.11% (-0.72-1%)	0.14% (-0.94-1.3%)	250 (-1600-2200)	2900 (-19000-27000)
United States	2016	Female	85 to 89	46% (34-57%)	54% (43-66%)	0.96 (0.6-1.4)	-0.19% (-1.1-0.81%)	-0.32% (-1.5-0.85%)	-720 (-3300-1900)	-4200 (-24000-18000)
United States	2016	Female	90 to 94	47% (36-58%)	53% (42-64%)	0.99 (0.62-1.4)	-0.51% (-1.6-0.63%)	-0.78% (-2.1-0.56%)	-1600 (-4300-1200)	-7100 (-21000-8600)
United States	2016	Female	95 plus	51% (37-64%)	49% (36-63%)	1.1 (0.64-1.6)	-0.92% (-2.4-0.57%)	-1.3% (-3.1-0.36%)	-1600 (-3700-430)	-4600 (-12000-2800)
United States	2016	Male	All Ages	73% (71-75%)	27% (25-29%)	3.2 (2.5-4.1)	6.7% (5.1-8.5%)	5.1% (3.3-7.2%)	71000 (46000-100000)	320000 (240000-410000)
United States	2016	Male	15 to 19	49% (42-55%)	51% (45-58%)	2.4 (1.7-3.2)	6.8% (5-8.9%)	13% (8.8-18%)	940 (620-1300)	93000 (68000-120000)
United States	2016	Male	20 to 24	81% (75-86%)	19% (14-25%)	6.4 (4.9-8.4)	15% (12-19%)	25% (19-32%)	3800 (2900-4800)	320000 (250000-390000)
United States	2016	Male	25 to 29	87% (83-90%)	13% (10-17%)	5.8 (4.5-7.5)	14% (11-17%)	22% (17-28%)	3700 (2800-4700)	300000 (240000-360000)
United States	2016	Male	30 to 34	89% (86-92%)	11% (7.9-14%)	5.2 (3.9-6.7)	12% (9.6-15%)	19% (15-24%)	3500 (2600-4400)	260000 (210000-320000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United States	2016	Male	35 to 39	90% (86-92%)	10% (7.7-14%)	4.7 (3.6-6.1)	11% (8.9-14%)	18% (14-22%)	3500 (2700-4300)	240000 (190000-290000)
United States	2016	Male	40 to 44	89% (86-92%)	11% (7.8-14%)	4.3 (3.3-5.6)	11% (8.4-13%)	16% (13-20%)	4100 (3200-5100)	240000 (190000-300000)
United States	2016	Male	45 to 49	89% (86-92%)	11% (7.8-14%)	4 (3.1-5.2)	11% (8.3-13%)	15% (12-19%)	5900 (4600-7200)	300000 (240000-360000)
United States	2016	Male	50 to 54	88% (85-91%)	12% (8.5-15%)	3.7 (2.8-4.7)	9.7% (7.6-12%)	13% (10-16%)	8800 (6900-11000)	380000 (300000-470000)
United States	2016	Male	55 to 59	87% (83-91%)	13% (9.3-17%)	3.3 (2.5-4.2)	7.9% (6-10%)	10% (7.8-13%)	10000 (7700-13000)	380000 (280000-480000)
United States	2016	Male	60 to 64	85% (81-89%)	15% (11-19%)	3 (2.2-3.9)	5.9% (4.1-7.8%)	7.4% (5.1-9.9%)	9000 (6200-12000)	290000 (200000-390000)
United States	2016	Male	65 to 69	82% (77-87%)	18% (13-23%)	2.7 (2-3.5)	4.2% (2.7-6%)	5.1% (3.1-7.4%)	7300 (4500-11000)	200000 (130000-290000)
United States	2016	Male	70 to 74	80% (73-85%)	20% (15-27%)	2.4 (1.8-3.2)	2.8% (1.4-4.7%)	3.2% (1.5-5.4%)	4900 (2300-8200)	110000 (57000-180000)
United States	2016	Male	75 to 79	78% (71-84%)	22% (16-29%)	2.2 (1.6-3)	1.9% (0.61-3.5%)	1.9% (0.42-3.9%)	3100 (670-6100)	58000 (19000-110000)
United States	2016	Male	80 to 84	76% (67-83%)	24% (17-33%)	2 (1.4-2.7)	1.3% (0.076-2.7%)	1.1% (-0.33-2.8%)	1900 (-560-4800)	30000 (1800-65000)
United States	2016	Male	85 to 89	75% (66-82%)	25% (18-34%)	1.9 (1.3-2.5)	0.75% (-0.46-2.2%)	0.46% (-1.1-2.2%)	790 (-1800-3800)	12000 (-7600-37000)
United States	2016	Male	90 to 94	75% (67-82%)	25% (18-33%)	1.8 (1.3-2.6)	0.29% (-1.1-2%)	-0.12% (-1.8-1.9%)	-140 (-2200-2300)	2300 (-8500-15000)
United States	2016	Male	95 plus	76% (66-84%)	24% (16-34%)	1.8 (1.2-2.6)	-0.017% (-1.7-1.9%)	-0.54% (-2.6-1.7%)	-230 (-1100-740)	-26 (-3100-3400)
Antigua and Barbuda	1990	Female	All Ages	38% (32-45%)	62% (55-68%)	0.24 (0.15-0.33)	0.86% (0.52-1.2%)	0.13% (-0.43-0.64%)	0.26 (-0.82-1.2)	63 (37-89)
Antigua and Barbuda	1990	Female	15 to 19	44% (30-61%)	56% (39-70%)	0.28 (0.14-0.49)	1.5% (1.1-2%)	2.5% (1.7-3.5%)	0.038 (0.026-0.055)	5.1 (3.6-6.9)
Antigua and Barbuda	1990	Female	20 to 24	50% (34-66%)	50% (34-66%)	0.41 (0.2-0.67)	2.6% (1.9-3.5%)	2.7% (1.9-3.6%)	0.053 (0.036-0.074)	12 (7.8-16)
Antigua and Barbuda	1990	Female	25 to 29	50% (33-66%)	50% (34-67%)	0.42 (0.21-0.68)	2.4% (1.8-3.1%)	2.5% (1.8-3.4%)	0.058 (0.039-0.081)	11 (7.5-14)
Antigua and Barbuda	1990	Female	30 to 34	48% (31-65%)	52% (35-69%)	0.41 (0.21-0.66)	2.3% (1.7-3%)	3.4% (2.6-4.5%)	0.065 (0.047-0.088)	8.2 (5.8-11)
Antigua and Barbuda	1990	Female	35 to 39	47% (32-64%)	53% (36-68%)	0.38 (0.18-0.65)	2.3% (1.7-2.9%)	3.4% (2.5-4.4%)	0.084 (0.061-0.11)	8.2 (6-11)
Antigua and Barbuda	1990	Female	40 to 44	46% (31-64%)	54% (36-69%)	0.36 (0.17-0.63)	2.4% (1.7-3.1%)	3.3% (2.3-4.5%)	0.14 (0.093-0.2)	9.1 (6.5-12)
Antigua and Barbuda	1990	Female	45 to 49	44% (28-61%)	56% (39-72%)	0.35 (0.16-0.64)	1.8% (1.1-2.4%)	2.6% (1.6-3.7%)	0.11 (0.064-0.16)	6.2 (3.8-8.4)
Antigua and Barbuda	1990	Female	50 to 54	40% (25-57%)	60% (43-75%)	0.31 (0.14-0.57)	1.4% (0.66-2%)	2% (0.87-3%)	0.083 (0.037-0.13)	3.8 (1.8-5.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	1990	Female	55 to 59	36% (22-53%)	64% (47-78%)	0.27 (0.12-0.49)	0.85% (0.12-1.5%)	1.2% (0.11-2%)	0.065 (0.0067-0.12)	2.5 (0.36-4.4)
Antigua and Barbuda	1990	Female	60 to 64	32% (20-48%)	68% (52-80%)	0.23 (0.1-0.43)	0.36% (-0.41-0.94%)	0.44% (-0.58-1.3%)	0.054 (-0.073-0.15)	1.9 (-2.1-4.9)
Antigua and Barbuda	1990	Female	65 to 69	30% (19-44%)	70% (56-81%)	0.2 (0.09-0.36)	-0.018% (-0.81-0.59%)	-0.046% (-1-0.75%)	-0.0077 (-0.18-0.12)	-0.09 (-4.2-3.1)
Antigua and Barbuda	1990	Female	70 to 74	27% (17-42%)	73% (58-83%)	0.17 (0.07-0.31)	-0.13% (-0.89-0.44%)	-0.17% (-1.1-0.55%)	-0.042 (-0.27-0.13)	-0.77 (-5.2-2.6)
Antigua and Barbuda	1990	Female	75 to 79	24% (14-38%)	76% (62-86%)	0.15 (0.062-0.28)	-0.24% (-1.1-0.33%)	-0.31% (-1.3-0.38%)	-0.11 (-0.47-0.14)	-1.5 (-6.7-2.2)
Antigua and Barbuda	1990	Female	80 to 84	22% (13-35%)	78% (65-87%)	0.14 (0.062-0.25)	-0.27% (-0.91-0.23%)	-0.37% (-1.2-0.23%)	-0.11 (-0.35-0.07)	-1.1 (-3.6-0.91)
Antigua and Barbuda	1990	Female	85 to 89	22% (13-33%)	78% (67-87%)	0.14 (0.062-0.27)	-0.32% (-1-0.13%)	-0.46% (-1.3-0.09%)	-0.11 (-0.29-0.02)	-0.67 (-2.1-0.29)
Antigua and Barbuda	1990	Female	90 to 94	25% (15-37%)	75% (63-85%)	0.16 (0.064-0.33)	-0.38% (-1.1-0.12%)	-0.56% (-1.4-0.05%)	-0.071 (-0.18-0.0067)	-0.3 (-0.84-0.091)
Antigua and Barbuda	1990	Female	95 plus	29% (15-46%)	71% (54-85%)	0.19 (0.065-0.4)	-0.49% (-1.3-0.13%)	-0.76% (-1.8--0.012%)	-0.031 (-0.074--0.00047)	-0.08 (-0.22-0.022)
Antigua and Barbuda	1990	Male	All Ages	60% (52-68%)	40% (32-48%)	0.5 (0.31-0.67)	3.4% (2.8-4.1%)	2.5% (1.8-3.4%)	5.3 (3.7-7.2)	280 (220-340)
Antigua and Barbuda	1990	Male	15 to 19	61% (44-76%)	39% (24-56%)	0.39 (0.2-0.67)	3.2% (2.3-4.2%)	3.7% (2.3-5.6%)	0.087 (0.052-0.13)	11 (7.8-15)
Antigua and Barbuda	1990	Male	20 to 24	70% (53-83%)	30% (17-47%)	0.71 (0.37-1.1)	6.5% (5-8.3%)	5.9% (3.7-8.2%)	0.18 (0.11-0.27)	30 (21-40)
Antigua and Barbuda	1990	Male	25 to 29	73% (56-85%)	27% (15-44%)	0.85 (0.45-1.3)	6.6% (5.1-8.3%)	6.3% (4.3-8.6%)	0.26 (0.18-0.37)	34 (24-44)
Antigua and Barbuda	1990	Male	30 to 34	74% (57-85%)	26% (15-43%)	0.89 (0.46-1.4)	6.4% (5.2-7.9%)	7% (5.1-9%)	0.25 (0.18-0.33)	27 (20-34)
Antigua and Barbuda	1990	Male	35 to 39	74% (58-86%)	26% (14-42%)	0.84 (0.43-1.4)	6.8% (5.5-8.3%)	7.8% (5.8-10%)	0.34 (0.25-0.45)	28 (22-35)
Antigua and Barbuda	1990	Male	40 to 44	73% (58-85%)	27% (15-42%)	0.81 (0.39-1.4)	7.1% (5.6-8.7%)	8.2% (6.2-10%)	0.49 (0.36-0.64)	31 (23-39)
Antigua and Barbuda	1990	Male	45 to 49	71% (53-84%)	29% (16-47%)	0.76 (0.38-1.3)	6.3% (5.1-7.8%)	7.6% (5.8-9.6%)	0.41 (0.3-0.53)	24 (18-30)
Antigua and Barbuda	1990	Male	50 to 54	70% (53-83%)	30% (17-47%)	0.7 (0.34-1.2)	5.6% (4.3-7%)	6.5% (4.8-8.3%)	0.52 (0.37-0.69)	23 (17-29)
Antigua and Barbuda	1990	Male	55 to 59	68% (50-82%)	32% (18-50%)	0.65 (0.31-1.2)	4.5% (3.2-6%)	5% (3.4-6.9%)	0.52 (0.35-0.72)	19 (13-25)
Antigua and Barbuda	1990	Male	60 to 64	66% (47-83%)	34% (17-53%)	0.59 (0.28-1.1)	3% (2-4.4%)	3.3% (2-4.8%)	0.76 (0.46-1.1)	23 (15-33)
Antigua and Barbuda	1990	Male	65 to 69	65% (47-80%)	35% (20-53%)	0.53 (0.26-0.91)	2.5% (1.6-3.5%)	2.8% (1.7-4%)	0.56 (0.34-0.8)	14 (8.7-20)
Antigua and Barbuda	1990	Male	70 to 74	62% (43-77%)	38% (23-57%)	0.44 (0.21-0.81)	1.4% (0.6-2.3%)	1.5% (0.52-2.5%)	0.52 (0.18-0.87)	11 (4.5-17)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	1990	Male	75 to 79	57% (37-75%)	43% (25-63%)	0.37 (0.17-0.66)	0.81% (0.046-1.5%)	0.8% (-0.073-1.6%)	0.28 (-0.025-0.56)	4.7 (0.27-8.6)
Antigua and Barbuda	1990	Male	80 to 84	52% (33-70%)	48% (30-67%)	0.32 (0.15-0.55)	0.5% (-0.14-1.1%)	0.47% (-0.29-1.1%)	0.1 (-0.068-0.25)	1.4 (-0.42-2.9)
Antigua and Barbuda	1990	Male	85 to 89	47% (30-64%)	53% (36-70%)	0.28 (0.13-0.51)	0.4% (-0.21-0.88%)	0.33% (-0.4-0.91%)	0.039 (-0.047-0.11)	0.41 (-0.21-0.91)
Antigua and Barbuda	1990	Male	90 to 94	47% (31-64%)	53% (36-69%)	0.28 (0.13-0.54)	0.21% (-0.44-0.7%)	0.044% (-0.75-0.63%)	0.0018 (-0.031-0.026)	0.052 (-0.11-0.17)
Antigua and Barbuda	1990	Male	95 plus	50% (31-70%)	50% (30-69%)	0.31 (0.13-0.62)	0.13% (-0.63-0.73%)	-0.16% (-1.1-0.55%)	-0.00096 (-0.0063-0.0031)	0.003 (-0.015-0.017)
Antigua and Barbuda	1995	Female	All Ages	38% (31-45%)	62% (55-69%)	0.32 (0.22-0.43)	0.83% (0.5-1.2%)	0.15% (-0.41-0.7%)	0.32 (-0.85-1.4)	67 (39-96)
Antigua and Barbuda	1995	Female	15 to 19	44% (30-60%)	56% (40-70%)	0.37 (0.18-0.64)	1.5% (1.1-2.1%)	2.5% (1.7-3.5%)	0.038 (0.025-0.053)	4.9 (3.4-6.7)
Antigua and Barbuda	1995	Female	20 to 24	50% (34-66%)	50% (34-66%)	0.54 (0.27-0.91)	2.5% (1.8-3.3%)	2.7% (1.8-3.9%)	0.051 (0.034-0.073)	9.5 (6.5-13)
Antigua and Barbuda	1995	Female	25 to 29	49% (33-66%)	51% (34-67%)	0.56 (0.31-0.9)	2.3% (1.7-3%)	2.5% (1.7-3.5%)	0.067 (0.043-0.094)	11 (7.8-15)
Antigua and Barbuda	1995	Female	30 to 34	48% (31-64%)	52% (36-69%)	0.54 (0.29-0.84)	2% (1.5-2.7%)	3.1% (2.2-4%)	0.059 (0.041-0.083)	8.9 (6.2-12)
Antigua and Barbuda	1995	Female	35 to 39	47% (31-64%)	53% (36-69%)	0.5 (0.26-0.81)	2% (1.5-2.6%)	3.2% (2.4-4.3%)	0.085 (0.06-0.12)	8.6 (6-11)
Antigua and Barbuda	1995	Female	40 to 44	46% (31-64%)	54% (36-69%)	0.47 (0.23-0.78)	2.3% (1.6-3%)	3.3% (2.3-4.6%)	0.19 (0.13-0.26)	12 (8.4-16)
Antigua and Barbuda	1995	Female	45 to 49	43% (27-61%)	57% (39-73%)	0.46 (0.24-0.82)	1.8% (1.1-2.5%)	2.6% (1.5-4%)	0.13 (0.074-0.2)	7.2 (4.3-10)
Antigua and Barbuda	1995	Female	50 to 54	40% (25-57%)	60% (43-75%)	0.41 (0.21-0.73)	1.4% (0.63-2.2%)	2% (0.87-3.3%)	0.14 (0.059-0.23)	6.1 (2.7-9.8)
Antigua and Barbuda	1995	Female	55 to 59	35% (22-53%)	65% (47-78%)	0.35 (0.16-0.63)	0.87% (0.13-1.6%)	1.2% (0.14-2.3%)	0.081 (0.009-0.15)	3 (0.48-5.3)
Antigua and Barbuda	1995	Female	60 to 64	32% (19-49%)	68% (51-81%)	0.3 (0.14-0.57)	0.3% (-0.57-1%)	0.37% (-0.71-1.3%)	0.042 (-0.081-0.16)	1.4 (-2.5-4.7)
Antigua and Barbuda	1995	Female	65 to 69	29% (18-44%)	71% (56-82%)	0.27 (0.12-0.48)	-0.048% (-0.91-0.6%)	-0.057% (-1.2-0.78%)	-0.01 (-0.21-0.15)	-0.29 (-5.3-3.6)
Antigua and Barbuda	1995	Female	70 to 74	27% (17-41%)	73% (59-83%)	0.22 (0.1-0.4)	-0.19% (-1-0.41%)	-0.22% (-1.2-0.53%)	-0.051 (-0.29-0.12)	-1.1 (-5.7-2.3)
Antigua and Barbuda	1995	Female	75 to 79	24% (14-38%)	76% (62-86%)	0.19 (0.086-0.37)	-0.28% (-1-0.28%)	-0.36% (-1.3-0.35%)	-0.11 (-0.41-0.11)	-1.6 (-5.9-1.6)
Antigua and Barbuda	1995	Female	80 to 84	22% (13-34%)	78% (66-87%)	0.18 (0.085-0.34)	-0.32% (-1-0.18%)	-0.43% (-1.3-0.2%)	-0.14 (-0.44-0.07)	-1.4 (-4.6-0.84)
Antigua and Barbuda	1995	Female	85 to 89	22% (13-34%)	78% (66-87%)	0.19 (0.086-0.35)	-0.36% (-1-0.11%)	-0.51% (-1.3-0.061%)	-0.14 (-0.38-0.018)	-0.94 (-2.7-0.29)
Antigua and Barbuda	1995	Female	90 to 94	24% (14-37%)	76% (63-86%)	0.21 (0.089-0.42)	-0.37% (-1-0.15%)	-0.57% (-1.4-0.09%)	-0.074 (-0.18-0.011)	-0.31 (-0.83-0.12)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	1995	Female	95 plus	29% (16-46%)	71% (54-84%)	0.26 (0.087-0.54)	-0.44% (-1.2-0.21%)	-0.75% (-1.8-0.095%)	-0.034 (-0.081-0.0041)	-0.085 (-0.24-0.04)
Antigua and Barbuda	1995	Male	All Ages	60% (53-68%)	40% (32-47%)	0.67 (0.46-0.89)	3.5% (2.8-4.3%)	2.7% (1.8-3.6%)	6 (4-8.2)	320 (250-390)
Antigua and Barbuda	1995	Male	15 to 19	60% (44-76%)	40% (24-56%)	0.52 (0.27-0.88)	3.5% (2.5-4.7%)	4.1% (2.4-6.2%)	0.098 (0.058-0.15)	12 (8.2-17)
Antigua and Barbuda	1995	Male	20 to 24	70% (53-83%)	30% (17-47%)	0.95 (0.54-1.5)	6.9% (5.1-8.7%)	6.4% (4.2-9%)	0.19 (0.12-0.27)	28 (19-37)
Antigua and Barbuda	1995	Male	25 to 29	73% (56-84%)	27% (16-44%)	1.1 (0.65-1.7)	6.7% (5.3-8.4%)	6.4% (4.3-8.6%)	0.29 (0.19-0.4)	37 (27-49)
Antigua and Barbuda	1995	Male	30 to 34	74% (56-85%)	26% (15-44%)	1.2 (0.67-1.8)	6.2% (5-7.7%)	6.6% (4.9-8.6%)	0.35 (0.25-0.47)	36 (27-46)
Antigua and Barbuda	1995	Male	35 to 39	73% (58-86%)	27% (14-42%)	1.1 (0.62-1.8)	6.3% (5.1-7.6%)	7.2% (5.5-9.3%)	0.35 (0.26-0.46)	31 (24-40)
Antigua and Barbuda	1995	Male	40 to 44	72% (57-84%)	28% (16-43%)	1.1 (0.58-1.7)	6.6% (5.2-8.2%)	7.9% (6-10%)	0.54 (0.39-0.71)	35 (27-45)
Antigua and Barbuda	1995	Male	45 to 49	71% (53-84%)	29% (16-47%)	1 (0.55-1.7)	6% (4.7-7.6%)	7.3% (5.5-9.4%)	0.47 (0.34-0.62)	27 (21-35)
Antigua and Barbuda	1995	Male	50 to 54	69% (52-83%)	31% (17-48%)	0.93 (0.48-1.6)	5.6% (4.1-7.3%)	6.5% (4.6-8.6%)	0.75 (0.52-1)	33 (24-44)
Antigua and Barbuda	1995	Male	55 to 59	68% (50-82%)	32% (18-50%)	0.85 (0.44-1.4)	4.6% (3.1-6.4%)	5.3% (3.4-7.4%)	0.77 (0.49-1.1)	28 (18-38)
Antigua and Barbuda	1995	Male	60 to 64	66% (47-82%)	34% (18-53%)	0.78 (0.37-1.4)	3.2% (1.9-4.7%)	3.5% (2-5.3%)	0.6 (0.33-0.91)	18 (11-26)
Antigua and Barbuda	1995	Male	65 to 69	65% (46-80%)	35% (20-54%)	0.69 (0.35-1.2)	2.6% (1.5-3.8%)	2.9% (1.6-4.3%)	0.65 (0.36-0.98)	16 (9.4-24)
Antigua and Barbuda	1995	Male	70 to 74	62% (43-76%)	38% (24-57%)	0.59 (0.29-1)	1.5% (0.57-2.5%)	1.6% (0.52-2.8%)	0.39 (0.13-0.69)	8.1 (3-14)
Antigua and Barbuda	1995	Male	75 to 79	57% (37-74%)	43% (26-63%)	0.49 (0.23-0.86)	0.91% (0.15-1.7%)	0.94% (0.016-1.9%)	0.3 (0.0048-0.61)	5.1 (0.87-9.4)
Antigua and Barbuda	1995	Male	80 to 84	51% (32-69%)	49% (31-68%)	0.42 (0.2-0.7)	0.56% (-0.12-1.2%)	0.54% (-0.26-1.3%)	0.17 (-0.078-0.42)	2.2 (-0.46-4.8)
Antigua and Barbuda	1995	Male	85 to 89	46% (29-63%)	54% (37-71%)	0.37 (0.18-0.68)	0.44% (-0.14-1%)	0.39% (-0.32-1%)	0.071 (-0.059-0.19)	0.71 (-0.24-1.6)
Antigua and Barbuda	1995	Male	90 to 94	47% (31-64%)	53% (36-69%)	0.37 (0.17-0.68)	0.27% (-0.35-0.84%)	0.1% (-0.65-0.78%)	0.0058 (-0.036-0.043)	0.092 (-0.12-0.29)
Antigua and Barbuda	1995	Male	95 plus	50% (30-69%)	50% (31-70%)	0.4 (0.16-0.77)	0.21% (-0.55-0.88%)	-0.1% (-0.99-0.74%)	-0.0013 (-0.013-0.0098)	0.011 (-0.03-0.048)
Antigua and Barbuda	2000	Female	All Ages	37% (31-45%)	63% (55-69%)	0.46 (0.33-0.6)	0.95% (0.6-1.3%)	0.37% (-0.25-1%)	0.78 (-0.52-2.3)	85 (52-120)
Antigua and Barbuda	2000	Female	15 to 19	43% (29-59%)	57% (41-71%)	0.51 (0.26-0.87)	1.5% (1-2.1%)	2.4% (1.5-3.6%)	0.042 (0.027-0.063)	5.7 (3.9-8)
Antigua and Barbuda	2000	Female	20 to 24	49% (33-65%)	51% (35-67%)	0.76 (0.4-1.3)	2.5% (1.8-3.4%)	2.9% (1.8-4.3%)	0.062 (0.038-0.093)	9.5 (6.4-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	2000	Female	25 to 29	48% (32-65%)	52% (35-68%)	0.78 (0.42-1.2)	2.3% (1.7-3%)	2.4% (1.6-3.5%)	0.054 (0.034-0.079)	10 (7.2-15)
Antigua and Barbuda	2000	Female	30 to 34	47% (30-64%)	53% (36-70%)	0.75 (0.41-1.2)	2% (1.4-2.7%)	2.8% (2-3.8%)	0.093 (0.063-0.13)	13 (8.6-18)
Antigua and Barbuda	2000	Female	35 to 39	46% (31-64%)	54% (36-69%)	0.69 (0.38-1.1)	1.9% (1.4-2.6%)	3.2% (2.3-4.2%)	0.13 (0.09-0.18)	12 (8.8-17)
Antigua and Barbuda	2000	Female	40 to 44	45% (30-63%)	55% (37-70%)	0.66 (0.34-1.1)	2.2% (1.4-3%)	3.5% (2.2-4.9%)	0.23 (0.14-0.33)	14 (9.5-20)
Antigua and Barbuda	2000	Female	45 to 49	43% (27-60%)	57% (40-73%)	0.64 (0.33-1.1)	1.7% (0.99-2.6%)	2.9% (1.7-4.4%)	0.18 (0.11-0.28)	9.8 (5.6-15)
Antigua and Barbuda	2000	Female	50 to 54	39% (25-56%)	61% (44-75%)	0.58 (0.29-0.99)	1.6% (0.62-2.6%)	2.3% (0.97-3.9%)	0.18 (0.072-0.3)	7.4 (2.9-13)
Antigua and Barbuda	2000	Female	55 to 59	35% (22-52%)	65% (48-78%)	0.49 (0.23-0.88)	0.97% (0.094-2%)	1.4% (0.21-2.8%)	0.13 (0.018-0.25)	4.5 (0.47-9.1)
Antigua and Barbuda	2000	Female	60 to 64	31% (19-47%)	69% (53-81%)	0.42 (0.19-0.77)	0.43% (-0.51-1.4%)	0.58% (-0.63-1.9%)	0.063 (-0.068-0.21)	1.8 (-2.1-6.2)
Antigua and Barbuda	2000	Female	65 to 69	29% (18-43%)	71% (57-82%)	0.37 (0.18-0.65)	0.062% (-0.76-0.87%)	0.13% (-0.97-1.2%)	0.019 (-0.14-0.18)	0.31 (-3.8-4.2)
Antigua and Barbuda	2000	Female	70 to 74	26% (17-40%)	74% (60-83%)	0.31 (0.14-0.55)	-0.11% (-0.96-0.63%)	-0.094% (-1.2-0.85%)	-0.023 (-0.27-0.21)	-0.67 (-5.8-3.8)
Antigua and Barbuda	2000	Female	75 to 79	23% (14-37%)	77% (63-86%)	0.27 (0.12-0.51)	-0.19% (-0.96-0.46%)	-0.23% (-1.1-0.59%)	-0.056 (-0.28-0.15)	-0.88 (-4.2-2.1)
Antigua and Barbuda	2000	Female	80 to 84	22% (12-34%)	78% (66-88%)	0.26 (0.12-0.47)	-0.22% (-0.93-0.42%)	-0.29% (-1.2-0.51%)	-0.095 (-0.39-0.16)	-0.97 (-4.2-1.8)
Antigua and Barbuda	2000	Female	85 to 89	22% (13-33%)	78% (67-87%)	0.26 (0.12-0.47)	-0.26% (-0.92-0.34%)	-0.39% (-1.2-0.33%)	-0.12 (-0.37-0.1)	-0.76 (-2.7-0.96)
Antigua and Barbuda	2000	Female	90 to 94	24% (14-36%)	76% (64-86%)	0.29 (0.13-0.57)	-0.26% (-0.93-0.45%)	-0.43% (-1.3-0.42%)	-0.077 (-0.23-0.075)	-0.29 (-1.1-0.5)
Antigua and Barbuda	2000	Female	95 plus	28% (15-45%)	72% (55-85%)	0.35 (0.13-0.74)	-0.25% (-1.1-0.65%)	-0.52% (-1.6-0.59%)	-0.03 (-0.09-0.033)	-0.06 (-0.26-0.16)
Antigua and Barbuda	2000	Male	All Ages	60% (52-68%)	40% (32-48%)	0.95 (0.69-1.2)	3.9% (3-4.7%)	3.1% (2.1-4.2%)	7.5 (4.9-10)	380 (300-480)
Antigua and Barbuda	2000	Male	15 to 19	60% (42-75%)	40% (25-58%)	0.73 (0.36-1.2)	4% (2.8-5.6%)	4.8% (2.8-7.3%)	0.16 (0.091-0.25)	17 (11-24)
Antigua and Barbuda	2000	Male	20 to 24	69% (52-83%)	31% (17-48%)	1.3 (0.76-2.1)	7.6% (5.6-9.8%)	7.5% (4.9-11%)	0.31 (0.2-0.46)	34 (24-45)
Antigua and Barbuda	2000	Male	25 to 29	72% (56-84%)	28% (16-44%)	1.6 (0.91-2.4)	7.1% (5.5-8.7%)	6.5% (4.3-8.8%)	0.24 (0.16-0.33)	33 (24-44)
Antigua and Barbuda	2000	Male	30 to 34	73% (55-85%)	27% (15-45%)	1.7 (0.97-2.5)	6.3% (5-7.9%)	6.4% (4.6-8.5%)	0.43 (0.3-0.59)	44 (33-57)
Antigua and Barbuda	2000	Male	35 to 39	73% (57-85%)	27% (15-43%)	1.6 (0.9-2.3)	5.9% (4.7-7.3%)	6.3% (4.6-8.3%)	0.42 (0.29-0.58)	39 (29-50)
Antigua and Barbuda	2000	Male	40 to 44	72% (57-85%)	28% (15-43%)	1.5 (0.82-2.3)	6.4% (5-7.9%)	7.4% (5.6-9.4%)	0.62 (0.45-0.82)	42 (32-55)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	2000	Male	45 to 49	70% (51-84%)	30% (16-49%)	1.4 (0.8-2.3)	6.2% (4.6-8%)	7.4% (5.3-9.8%)	0.67 (0.47-0.91)	38 (28-51)
Antigua and Barbuda	2000	Male	50 to 54	69% (51-83%)	31% (17-49%)	1.3 (0.69-2.2)	6% (4.3-8.1%)	7.1% (4.8-9.7%)	0.89 (0.6-1.2)	40 (27-54)
Antigua and Barbuda	2000	Male	55 to 59	67% (49-82%)	33% (18-51%)	1.2 (0.61-2)	5.2% (3.5-7.4%)	6% (3.9-8.7%)	0.99 (0.63-1.4)	36 (24-51)
Antigua and Barbuda	2000	Male	60 to 64	65% (46-82%)	35% (18-54%)	1.1 (0.52-1.9)	3.8% (2.2-5.9%)	4.3% (2.5-6.9%)	0.68 (0.39-1.1)	21 (12-32)
Antigua and Barbuda	2000	Male	65 to 69	64% (45-80%)	36% (20-55%)	0.97 (0.5-1.6)	2.9% (1.6-4.8%)	3.3% (1.7-5.6%)	0.57 (0.29-0.95)	14 (7.7-23)
Antigua and Barbuda	2000	Male	70 to 74	61% (43-76%)	39% (24-57%)	0.82 (0.4-1.4)	1.9% (0.73-3.3%)	2.1% (0.72-3.7%)	0.59 (0.2-1)	12 (4.7-21)
Antigua and Barbuda	2000	Male	75 to 79	56% (37-74%)	44% (26-63%)	0.69 (0.33-1.2)	1.3% (0.4-2.5%)	1.4% (0.34-2.7%)	0.38 (0.092-0.74)	5.9 (1.8-11)
Antigua and Barbuda	2000	Male	80 to 84	50% (32-68%)	50% (32-68%)	0.58 (0.28-1)	0.9% (0.17-1.8%)	0.94% (0.094-2%)	0.27 (0.027-0.58)	3.3 (0.59-6.7)
Antigua and Barbuda	2000	Male	85 to 89	45% (28-63%)	55% (37-72%)	0.51 (0.25-0.9)	0.72% (0.1-1.4%)	0.7% (-0.039-1.6%)	0.18 (-0.0098-0.39)	1.6 (0.23-3.3)
Antigua and Barbuda	2000	Male	90 to 94	46% (30-64%)	54% (36-70%)	0.52 (0.24-0.95)	0.46% (-0.18-1.3%)	0.32% (-0.45-1.3%)	0.033 (-0.048-0.13)	0.29 (-0.11-0.82)
Antigua and Barbuda	2000	Male	95 plus	49% (29-69%)	51% (31-71%)	0.57 (0.23-1)	0.41% (-0.39-1.4%)	0.12% (-0.87-1.2%)	0.0026 (-0.019-0.027)	0.037 (-0.036-0.12)
Antigua and Barbuda	2005	Female	All Ages	37% (31-45%)	63% (55-69%)	0.54 (0.38-0.72)	1.1% (0.68-1.6%)	0.61% (-0.095-1.5%)	1.3 (-0.22-3.2)	100 (63-150)
Antigua and Barbuda	2005	Female	15 to 19	43% (29-59%)	57% (41-71%)	0.6 (0.3-1)	1.4% (0.95-2%)	2.8% (1.8-4.2%)	0.035 (0.022-0.053)	5.4 (3.6-7.8)
Antigua and Barbuda	2005	Female	20 to 24	49% (33-64%)	51% (36-67%)	0.89 (0.47-1.5)	2.6% (1.8-3.5%)	3.1% (2-4.8%)	0.072 (0.045-0.11)	11 (7.4-16)
Antigua and Barbuda	2005	Female	25 to 29	48% (32-65%)	52% (35-68%)	0.92 (0.47-1.5)	2.4% (1.8-3.2%)	3% (2-4.3%)	0.052 (0.034-0.077)	8.8 (6.1-13)
Antigua and Barbuda	2005	Female	30 to 34	47% (30-64%)	53% (36-70%)	0.9 (0.48-1.4)	2.4% (1.7-3.1%)	3.8% (2.6-5.1%)	0.12 (0.084-0.17)	13 (9.5-18)
Antigua and Barbuda	2005	Female	35 to 39	46% (30-63%)	54% (37-70%)	0.82 (0.45-1.3)	2.1% (1.5-2.8%)	3.9% (2.8-5.3%)	0.19 (0.13-0.27)	16 (11-21)
Antigua and Barbuda	2005	Female	40 to 44	44% (29-62%)	56% (38-71%)	0.78 (0.41-1.2)	2.2% (1.4-3.1%)	4% (2.6-5.7%)	0.24 (0.16-0.35)	15 (9.5-21)
Antigua and Barbuda	2005	Female	45 to 49	43% (27-60%)	57% (40-73%)	0.75 (0.39-1.3)	2% (1.2-3%)	3.5% (2.2-5.4%)	0.23 (0.14-0.35)	12 (7-18)
Antigua and Barbuda	2005	Female	50 to 54	39% (25-56%)	61% (44-75%)	0.68 (0.34-1.2)	1.9% (0.82-3.2%)	2.8% (1.3-4.7%)	0.28 (0.13-0.46)	11 (4.9-19)
Antigua and Barbuda	2005	Female	55 to 59	34% (22-51%)	66% (49-78%)	0.58 (0.27-1)	1.2% (0.21-2.3%)	1.9% (0.5-3.6%)	0.13 (0.034-0.25)	4.5 (0.77-9)
Antigua and Barbuda	2005	Female	60 to 64	31% (19-48%)	69% (52-81%)	0.5 (0.23-0.92)	0.66% (-0.33-1.8%)	1% (-0.31-2.6%)	0.13 (-0.04-0.32)	3.6 (-1.8-9.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	2005	Female	65 to 69	28% (18-43%)	72% (57-82%)	0.43 (0.21-0.79)	0.26% (-0.65-1.2%)	0.45% (-0.68-1.7%)	0.07 (-0.11-0.26)	1.3 (-3.4-6.3)
Antigua and Barbuda	2005	Female	70 to 74	26% (16-40%)	74% (60-84%)	0.36 (0.17-0.67)	0.036% (-0.8-0.85%)	0.13% (-0.95-1.2%)	0.024 (-0.18-0.23)	0.17 (-3.9-4.3)
Antigua and Barbuda	2005	Female	75 to 79	23% (14-37%)	77% (63-86%)	0.32 (0.14-0.6)	-0.089% (-0.92-0.68%)	-0.085% (-1.1-0.88%)	-0.021 (-0.28-0.21)	-0.43 (-4.4-3.1)
Antigua and Barbuda	2005	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.3 (0.14-0.53)	-0.11% (-0.83-0.56%)	-0.15% (-1-0.74%)	-0.048 (-0.33-0.23)	-0.51 (-3.7-2.5)
Antigua and Barbuda	2005	Female	85 to 89	21% (13-32%)	79% (68-87%)	0.3 (0.14-0.57)	-0.13% (-0.83-0.58%)	-0.22% (-1.1-0.66%)	-0.068 (-0.34-0.2)	-0.39 (-2.5-1.7)
Antigua and Barbuda	2005	Female	90 to 94	24% (14-37%)	76% (63-86%)	0.35 (0.15-0.68)	-0.16% (-0.95-0.71%)	-0.3% (-1.2-0.75%)	-0.074 (-0.32-0.19)	-0.25 (-1.5-1.1)
Antigua and Barbuda	2005	Female	95 plus	28% (15-45%)	72% (55-85%)	0.41 (0.15-0.86)	-0.12% (-1.1-1.1%)	-0.32% (-1.5-1%)	-0.03 (-0.14-0.095)	-0.044 (-0.4-0.39)
Antigua and Barbuda	2005	Male	All Ages	60% (52-68%)	40% (32-48%)	1.1 (0.79-1.5)	4.6% (3.6-5.6%)	3.8% (2.6-5.3%)	9.3 (6.3-13)	470 (360-580)
Antigua and Barbuda	2005	Male	15 to 19	59% (42-75%)	41% (25-58%)	0.85 (0.44-1.4)	4.3% (3-5.9%)	5.4% (3.2-8.3%)	0.19 (0.11-0.29)	20 (13-29)
Antigua and Barbuda	2005	Male	20 to 24	69% (52-83%)	31% (17-48%)	1.6 (0.87-2.5)	8.4% (6.2-11%)	8.7% (5.6-12%)	0.5 (0.32-0.74)	51 (36-69)
Antigua and Barbuda	2005	Male	25 to 29	72% (56-84%)	28% (16-44%)	1.9 (1-2.9)	8.6% (6.7-11%)	9.3% (6.2-13%)	0.32 (0.21-0.45)	36 (27-48)
Antigua and Barbuda	2005	Male	30 to 34	73% (55-85%)	27% (15-45%)	2 (1.2-3)	8.1% (6.3-10%)	9.3% (6.8-13%)	0.61 (0.43-0.84)	53 (40-70)
Antigua and Barbuda	2005	Male	35 to 39	73% (56-85%)	27% (15-44%)	1.9 (1.1-2.8)	7.2% (5.7-8.9%)	9% (6.8-12%)	0.6 (0.43-0.79)	52 (40-67)
Antigua and Barbuda	2005	Male	40 to 44	72% (57-85%)	28% (15-43%)	1.8 (0.98-2.7)	7.4% (5.8-9.4%)	9.7% (7.3-12%)	0.8 (0.58-1.1)	51 (38-66)
Antigua and Barbuda	2005	Male	45 to 49	70% (52-84%)	30% (16-48%)	1.7 (0.93-2.8)	7.3% (5.4-9.7%)	9.3% (6.7-13%)	0.95 (0.66-1.3)	51 (37-68)
Antigua and Barbuda	2005	Male	50 to 54	69% (51-83%)	31% (17-49%)	1.5 (0.8-2.5)	6.7% (4.7-9.3%)	8.3% (5.6-12%)	1 (0.67-1.5)	46 (31-63)
Antigua and Barbuda	2005	Male	55 to 59	67% (50-82%)	33% (18-50%)	1.4 (0.7-2.3)	5.8% (3.7-8.4%)	7% (4.5-10%)	0.94 (0.6-1.4)	34 (22-50)
Antigua and Barbuda	2005	Male	60 to 64	65% (46-81%)	35% (19-54%)	1.3 (0.62-2.3)	4.4% (2.7-6.8%)	5.3% (3.2-8.3%)	0.82 (0.5-1.3)	26 (16-40)
Antigua and Barbuda	2005	Male	65 to 69	64% (45-80%)	36% (20-55%)	1.2 (0.58-2)	3.4% (1.9-5.7%)	4% (2.2-6.8%)	0.86 (0.46-1.4)	22 (12-36)
Antigua and Barbuda	2005	Male	70 to 74	61% (42-76%)	39% (24-58%)	0.97 (0.48-1.7)	2.2% (0.85-4%)	2.5% (0.86-4.5%)	0.59 (0.21-1.1)	12 (4.6-22)
Antigua and Barbuda	2005	Male	75 to 79	56% (36-73%)	44% (27-64%)	0.81 (0.38-1.5)	1.5% (0.42-3%)	1.6% (0.37-3.3%)	0.49 (0.12-1)	7.9 (2.2-16)
Antigua and Barbuda	2005	Male	80 to 84	50% (31-68%)	50% (32-69%)	0.69 (0.34-1.2)	1% (0.23-2.2%)	1.1% (0.16-2.4%)	0.33 (0.045-0.72)	4 (0.83-8.3)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	2005	Male	85 to 89	45% (29-62%)	55% (38-71%)	0.61 (0.29-1.1)	0.85% (0.19-1.7%)	0.85% (0.035-1.8%)	0.18 (0.0072-0.39)	1.7 (0.37-3.3)
Antigua and Barbuda	2005	Male	90 to 94	45% (30-62%)	55% (38-70%)	0.61 (0.28-1.1)	0.63% (-0.052-1.6%)	0.51% (-0.33-1.7%)	0.074 (-0.048-0.24)	0.58 (-0.047-1.5)
Antigua and Barbuda	2005	Male	95 plus	48% (29-68%)	52% (32-71%)	0.65 (0.27-1.3)	0.59% (-0.29-1.9%)	0.33% (-0.75-1.9%)	0.012 (-0.028-0.068)	0.091 (-0.046-0.3)
Antigua and Barbuda	2010	Female	All Ages	37% (31-44%)	63% (56-69%)	0.61 (0.4-0.86)	1.3% (0.77-1.9%)	0.94% (0.16-2%)	2.1 (0.34-4.3)	120 (72-180)
Antigua and Barbuda	2010	Female	15 to 19	42% (29-59%)	58% (41-71%)	0.67 (0.33-1.2)	1.4% (0.93-2%)	2.8% (1.7-4.3%)	0.039 (0.023-0.061)	5.9 (3.9-8.7)
Antigua and Barbuda	2010	Female	20 to 24	48% (32-64%)	52% (36-68%)	1 (0.49-1.7)	2.5% (1.7-3.4%)	3.4% (2-5.2%)	0.061 (0.036-0.097)	11 (7.1-15)
Antigua and Barbuda	2010	Female	25 to 29	48% (31-64%)	52% (36-69%)	1 (0.51-1.7)	2.5% (1.8-3.4%)	3.4% (2-5.2%)	0.07 (0.04-0.11)	11 (7.2-15)
Antigua and Barbuda	2010	Female	30 to 34	46% (30-64%)	54% (36-70%)	0.99 (0.5-1.6)	2.6% (1.8-3.5%)	4.2% (2.8-6%)	0.11 (0.069-0.16)	11 (7.5-15)
Antigua and Barbuda	2010	Female	35 to 39	45% (30-63%)	55% (37-70%)	0.91 (0.47-1.5)	2.2% (1.6-3%)	4.3% (2.9-5.9%)	0.17 (0.11-0.25)	14 (9.9-20)
Antigua and Barbuda	2010	Female	40 to 44	44% (29-62%)	56% (38-71%)	0.86 (0.42-1.5)	2.2% (1.4-3.2%)	4.3% (2.8-6.4%)	0.31 (0.2-0.47)	19 (12-28)
Antigua and Barbuda	2010	Female	45 to 49	42% (26-60%)	58% (40-74%)	0.84 (0.43-1.5)	2.1% (1.1-3.3%)	3.9% (2.3-6.1%)	0.31 (0.19-0.5)	16 (8.9-25)
Antigua and Barbuda	2010	Female	50 to 54	38% (24-55%)	62% (45-76%)	0.76 (0.38-1.3)	2% (0.89-3.5%)	3.3% (1.6-5.5%)	0.38 (0.18-0.65)	15 (6.7-27)
Antigua and Barbuda	2010	Female	55 to 59	34% (21-51%)	66% (49-79%)	0.65 (0.3-1.1)	1.3% (0.34-2.6%)	2.3% (0.84-4.2%)	0.22 (0.075-0.42)	7.7 (2-15)
Antigua and Barbuda	2010	Female	60 to 64	30% (18-46%)	70% (54-82%)	0.55 (0.25-1)	0.87% (-0.17-2.2%)	1.4% (-0.055-3.3%)	0.14 (-0.0054-0.33)	3.9 (-0.78-9.8)
Antigua and Barbuda	2010	Female	65 to 69	28% (18-42%)	72% (58-82%)	0.48 (0.22-0.88)	0.46% (-0.44-1.6%)	0.79% (-0.43-2.3%)	0.12 (-0.06-0.34)	2.5 (-2.3-8.1)
Antigua and Barbuda	2010	Female	70 to 74	26% (16-40%)	74% (60-84%)	0.41 (0.19-0.77)	0.25% (-0.63-1.2%)	0.44% (-0.71-1.8%)	0.085 (-0.14-0.34)	1.3 (-3.3-6.4)
Antigua and Barbuda	2010	Female	75 to 79	23% (14-37%)	77% (63-86%)	0.36 (0.15-0.7)	0.12% (-0.74-1%)	0.19% (-0.89-1.3%)	0.049 (-0.22-0.34)	0.58 (-3.5-5)
Antigua and Barbuda	2010	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.34 (0.14-0.62)	0.048% (-0.69-0.8%)	0.059% (-0.85-1%)	0.017 (-0.24-0.29)	0.19 (-2.8-3.3)
Antigua and Barbuda	2010	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.34 (0.15-0.64)	0.044% (-0.69-0.92%)	-0.0058% (-0.92-1.1%)	-0.002 (-0.31-0.34)	0.14 (-2.2-2.8)
Antigua and Barbuda	2010	Female	90 to 94	23% (14-35%)	77% (65-86%)	0.39 (0.15-0.75)	0.061% (-0.75-1.1%)	-0.039% (-1-1.2%)	-0.0093 (-0.25-0.28)	0.09 (-1.1-1.6)
Antigua and Barbuda	2010	Female	95 plus	27% (15-45%)	73% (55-85%)	0.47 (0.15-1)	0.14% (-0.86-1.6%)	-0.032% (-1.3-1.6%)	-0.0033 (-0.13-0.18)	0.062 (-0.37-0.68)
Antigua and Barbuda	2010	Male	All Ages	60% (52-68%)	40% (32-48%)	1.3 (0.84-1.7)	5% (3.9-6.5%)	4.5% (3-6.3%)	11 (7.1-16)	530 (390-680)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	2010	Male	15 to 19	59% (42-75%)	41% (25-58%)	0.97 (0.47-1.7)	4.4% (3-6.3%)	5.8% (3.4-8.9%)	0.22 (0.13-0.35)	24 (15-34)
Antigua and Barbuda	2010	Male	20 to 24	69% (52-82%)	31% (18-48%)	1.8 (0.96-3)	8.7% (6.4-12%)	9.4% (6-14%)	0.49 (0.3-0.74)	51 (36-70)
Antigua and Barbuda	2010	Male	25 to 29	72% (55-84%)	28% (16-45%)	2.1 (1.1-3.5)	9% (6.8-12%)	10% (6.4-14%)	0.4 (0.26-0.59)	44 (32-60)
Antigua and Barbuda	2010	Male	30 to 34	73% (55-85%)	27% (15-45%)	2.2 (1.2-3.5)	8.8% (6.7-12%)	10% (7.1-15%)	0.54 (0.37-0.79)	46 (33-61)
Antigua and Barbuda	2010	Male	35 to 39	73% (57-85%)	27% (15-43%)	2.1 (1.1-3.4)	7.8% (6-9.9%)	10% (7.5-14%)	0.57 (0.39-0.79)	47 (34-61)
Antigua and Barbuda	2010	Male	40 to 44	71% (56-85%)	29% (15-44%)	2 (1-3.3)	7.6% (5.7-9.9%)	10% (7.6-14%)	0.98 (0.68-1.3)	64 (46-86)
Antigua and Barbuda	2010	Male	45 to 49	70% (51-84%)	30% (16-49%)	1.9 (1-3.2)	7.5% (5.5-10%)	9.9% (7-14%)	1.1 (0.79-1.6)	61 (44-84)
Antigua and Barbuda	2010	Male	50 to 54	68% (50-83%)	32% (17-50%)	1.7 (0.88-2.9)	7% (4.8-10%)	9% (6-13%)	1.2 (0.79-1.8)	56 (38-80)
Antigua and Barbuda	2010	Male	55 to 59	67% (49-82%)	33% (18-51%)	1.6 (0.78-2.6)	6.3% (4-9.4%)	7.7% (4.7-12%)	1.4 (0.88-2.2)	52 (33-79)
Antigua and Barbuda	2010	Male	60 to 64	65% (46-82%)	35% (18-54%)	1.5 (0.66-2.6)	4.9% (2.9-7.9%)	6% (3.5-9.6%)	0.96 (0.55-1.6)	29 (17-47)
Antigua and Barbuda	2010	Male	65 to 69	64% (44-79%)	36% (21-56%)	1.3 (0.63-2.2)	3.8% (2-6.5%)	4.5% (2.4-7.7%)	0.95 (0.5-1.6)	24 (13-41)
Antigua and Barbuda	2010	Male	70 to 74	60% (42-76%)	40% (24-58%)	1.1 (0.5-1.9)	2.5% (1.1-4.6%)	2.9% (1.1-5.3%)	0.76 (0.29-1.4)	16 (6.7-28)
Antigua and Barbuda	2010	Male	75 to 79	55% (35-73%)	45% (27-65%)	0.9 (0.41-1.6)	1.8% (0.63-3.7%)	2% (0.61-4.2%)	0.5 (0.15-1.1)	8 (2.7-16)
Antigua and Barbuda	2010	Male	80 to 84	49% (31-67%)	51% (33-69%)	0.76 (0.36-1.4)	1.3% (0.37-2.6%)	1.4% (0.34-2.9%)	0.39 (0.093-0.83)	4.7 (1.4-9.6)
Antigua and Barbuda	2010	Male	85 to 89	45% (27-61%)	55% (39-73%)	0.67 (0.3-1.2)	1.1% (0.34-2.2%)	1.1% (0.21-2.4%)	0.24 (0.046-0.53)	2.2 (0.7-4.5)
Antigua and Barbuda	2010	Male	90 to 94	45% (29-61%)	55% (39-71%)	0.67 (0.29-1.3)	0.89% (0.18-2.1%)	0.82% (-0.041-2.2%)	0.099 (-0.0051-0.27)	0.68 (0.13-1.6)
Antigua and Barbuda	2010	Male	95 plus	47% (28-67%)	53% (33-72%)	0.73 (0.27-1.5)	0.88% (-0.042-2.6%)	0.68% (-0.42-2.7%)	0.032 (-0.02-0.12)	0.18 (-0.0085-0.5)
Antigua and Barbuda	2016	Female	All Ages	37% (30-44%)	63% (56-70%)	0.66 (0.37-1)	1.4% (0.79-2.3%)	1.2% (0.34-2.6%)	2.9 (0.79-6.4)	150 (79-240)
Antigua and Barbuda	2016	Female	15 to 19	42% (29-58%)	58% (42-71%)	0.71 (0.32-1.3)	1.4% (0.91-2.1%)	2.7% (1.5-4.5%)	0.036 (0.019-0.06)	5.6 (3.5-8.5)
Antigua and Barbuda	2016	Female	20 to 24	48% (32-64%)	52% (36-68%)	1.1 (0.46-2)	2.5% (1.6-3.5%)	3.4% (1.9-5.6%)	0.06 (0.032-0.1)	12 (7.3-17)
Antigua and Barbuda	2016	Female	25 to 29	47% (31-64%)	53% (36-69%)	1.1 (0.48-2)	2.6% (1.8-3.6%)	3.6% (2-5.7%)	0.082 (0.044-0.14)	12 (8-18)
Antigua and Barbuda	2016	Female	30 to 34	46% (29-63%)	54% (37-71%)	1.1 (0.47-1.9)	2.7% (1.8-3.7%)	4.3% (2.7-6.4%)	0.14 (0.083-0.22)	14 (9.2-20)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	2016	Female	35 to 39	45% (29-62%)	55% (38-71%)	0.98 (0.42-1.8)	2.4% (1.6-3.5%)	4.5% (2.9-6.8%)	0.14 (0.085-0.23)	11 (7.2-17)
Antigua and Barbuda	2016	Female	40 to 44	44% (29-62%)	56% (38-71%)	0.92 (0.39-1.8)	2.3% (1.4-3.5%)	4.6% (2.9-7.1%)	0.25 (0.15-0.41)	16 (9.5-25)
Antigua and Barbuda	2016	Female	45 to 49	42% (26-59%)	58% (41-74%)	0.89 (0.41-1.7)	2.2% (1.2-3.6%)	4.1% (2.3-6.7%)	0.45 (0.24-0.77)	23 (12-38)
Antigua and Barbuda	2016	Female	50 to 54	38% (24-55%)	62% (45-76%)	0.79 (0.35-1.5)	2.2% (0.98-3.9%)	3.6% (1.8-6.3%)	0.54 (0.27-0.97)	22 (10-40)
Antigua and Barbuda	2016	Female	55 to 59	33% (20-50%)	67% (50-80%)	0.67 (0.28-1.2)	1.6% (0.55-3%)	2.7% (1.1-5.1%)	0.36 (0.15-0.68)	12 (4.4-24)
Antigua and Barbuda	2016	Female	60 to 64	30% (18-46%)	70% (54-82%)	0.58 (0.23-1.1)	1.1% (-0.0021-2.7%)	1.7% (0.17-4%)	0.29 (0.026-0.69)	8.2 (-0.016-21)
Antigua and Barbuda	2016	Female	65 to 69	28% (17-42%)	72% (58-83%)	0.51 (0.2-0.9)	0.64% (-0.29-1.9%)	1.1% (-0.25-2.8%)	0.14 (-0.035-0.38)	3.1 (-1.4-9.4)
Antigua and Barbuda	2016	Female	70 to 74	25% (16-38%)	75% (62-84%)	0.43 (0.17-0.84)	0.43% (-0.4-1.7%)	0.69% (-0.43-2.4%)	0.15 (-0.091-0.52)	2.5 (-2.3-10)
Antigua and Barbuda	2016	Female	75 to 79	22% (14-35%)	78% (65-86%)	0.38 (0.14-0.76)	0.29% (-0.53-1.4%)	0.41% (-0.62-1.9%)	0.12 (-0.2-0.57)	1.7 (-3.2-8.5)
Antigua and Barbuda	2016	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.35 (0.14-0.7)	0.2% (-0.52-1.2%)	0.24% (-0.67-1.5%)	0.076 (-0.2-0.5)	0.88 (-2.3-5.6)
Antigua and Barbuda	2016	Female	85 to 89	20% (12-31%)	80% (69-88%)	0.35 (0.14-0.7)	0.2% (-0.54-1.2%)	0.18% (-0.76-1.4%)	0.053 (-0.22-0.42)	0.55 (-1.6-3.4)
Antigua and Barbuda	2016	Female	90 to 94	23% (13-35%)	77% (65-87%)	0.4 (0.15-0.81)	0.19% (-0.61-1.4%)	0.12% (-0.87-1.6%)	0.029 (-0.21-0.39)	0.29 (-0.93-2.2)
Antigua and Barbuda	2016	Female	95 plus	27% (14-43%)	73% (57-86%)	0.48 (0.15-1.1)	0.29% (-0.76-2%)	0.14% (-1.1-2.2%)	0.013 (-0.11-0.21)	0.11 (-0.29-0.77)
Antigua and Barbuda	2016	Male	All Ages	60% (52-68%)	40% (32-48%)	1.4 (0.75-2.1)	5.3% (3.8-7.2%)	4.9% (3.1-7.5%)	13 (7.6-20)	600 (410-840)
Antigua and Barbuda	2016	Male	15 to 19	59% (42-75%)	41% (25-58%)	1 (0.43-1.9)	4.3% (2.9-6.2%)	5.8% (3.3-9.1%)	0.22 (0.12-0.35)	23 (15-34)
Antigua and Barbuda	2016	Male	20 to 24	68% (51-82%)	32% (18-49%)	1.9 (0.84-3.3)	8.6% (6-12%)	9.5% (5.7-15%)	0.5 (0.29-0.81)	54 (37-77)
Antigua and Barbuda	2016	Male	25 to 29	71% (54-84%)	29% (16-46%)	2.3 (1-4)	9.1% (6.7-12%)	10% (6.4-16%)	0.41 (0.25-0.65)	46 (32-63)
Antigua and Barbuda	2016	Male	30 to 34	72% (55-85%)	28% (15-45%)	2.3 (1.1-4)	8.9% (6.5-12%)	11% (6.8-16%)	0.62 (0.39-0.93)	54 (37-74)
Antigua and Barbuda	2016	Male	35 to 39	72% (56-85%)	28% (15-44%)	2.2 (1-3.9)	8.2% (6-11%)	11% (7.3-15%)	0.48 (0.29-0.73)	39 (26-54)
Antigua and Barbuda	2016	Male	40 to 44	71% (56-84%)	29% (16-44%)	2.1 (0.94-3.9)	7.8% (5.7-11%)	11% (7.7-16%)	0.74 (0.47-1.1)	49 (34-70)
Antigua and Barbuda	2016	Male	45 to 49	70% (50-84%)	30% (16-50%)	2 (0.99-3.6)	7.7% (5.5-11%)	10% (7-15%)	1.5 (0.91-2.2)	79 (52-110)
Antigua and Barbuda	2016	Male	50 to 54	68% (50-83%)	32% (17-50%)	1.8 (0.85-3.3)	7.1% (4.7-11%)	9.3% (5.9-14%)	1.5 (0.92-2.4)	70 (44-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Antigua and Barbuda	2016	Male	55 to 59	66% (48-82%)	34% (18-52%)	1.7 (0.73-3)	6.4% (3.9-10%)	8% (4.7-13%)	1.8 (1-2.9)	68 (39-100)
Antigua and Barbuda	2016	Male	60 to 64	64% (45-81%)	36% (19-55%)	1.5 (0.61-2.8)	5.2% (2.8-8.6%)	6.3% (3.3-11%)	1.7 (0.87-2.8)	51 (27-83)
Antigua and Barbuda	2016	Male	65 to 69	63% (44-79%)	37% (21-56%)	1.4 (0.57-2.4)	4% (1.9-7%)	4.8% (2.3-8.7%)	1.1 (0.53-2)	28 (14-49)
Antigua and Barbuda	2016	Male	70 to 74	60% (41-76%)	40% (24-59%)	1.1 (0.47-2.1)	2.7% (1.1-5.4%)	3.2% (1.2-6.4%)	0.78 (0.3-1.6)	16 (6.5-32)
Antigua and Barbuda	2016	Male	75 to 79	55% (35-73%)	45% (27-65%)	0.95 (0.37-1.8)	2% (0.65-4.2%)	2.2% (0.65-4.9%)	0.7 (0.2-1.6)	11 (3.6-24)
Antigua and Barbuda	2016	Male	80 to 84	49% (30-67%)	51% (33-70%)	0.8 (0.33-1.5)	1.4% (0.36-3.3%)	1.6% (0.32-3.6%)	0.43 (0.087-1)	5.1 (1.3-12)
Antigua and Barbuda	2016	Male	85 to 89	44% (27-61%)	56% (39-73%)	0.7 (0.29-1.4)	1.2% (0.31-2.7%)	1.2% (0.2-3%)	0.24 (0.038-0.57)	2.1 (0.54-4.7)
Antigua and Barbuda	2016	Male	90 to 94	44% (29-61%)	56% (39-71%)	0.71 (0.28-1.4)	1% (0.21-2.6%)	0.96% (0.0016-2.9%)	0.12 (0.0002-0.35)	0.77 (0.16-2)
Antigua and Barbuda	2016	Male	95 plus	47% (28-67%)	53% (33-72%)	0.76 (0.26-1.6)	1.1% (0.078-3%)	0.87% (-0.34-3.2%)	0.031 (-0.013-0.11)	0.16 (0.012-0.45)
The Bahamas	1990	Female	All Ages	36% (30-44%)	64% (56-70%)	1.8 (1.5-2.1)	4.6% (3.6-5.8%)	6.3% (4.5-8.5%)	39 (27-51)	1500 (1200-1900)
The Bahamas	1990	Female	15 to 19	42% (29-58%)	58% (42-71%)	2 (1.1-3.3)	3.2% (2-4.9%)	7.2% (4.3-11%)	0.51 (0.29-0.79)	51 (32-76)
The Bahamas	1990	Female	20 to 24	48% (32-65%)	52% (35-68%)	3.1 (1.8-4.5)	5.7% (4-7.6%)	9.6% (6.3-13%)	1.2 (0.77-1.6)	110 (76-140)
The Bahamas	1990	Female	25 to 29	48% (31-65%)	52% (35-69%)	3.2 (1.9-4.6)	6.7% (4.8-8.6%)	9.8% (6.8-13%)	2.2 (1.5-2.9)	170 (120-230)
The Bahamas	1990	Female	30 to 34	46% (29-63%)	54% (37-71%)	3 (1.8-4.6)	7.2% (5.2-9.5%)	10% (7-13%)	2.8 (2-3.8)	190 (140-250)
The Bahamas	1990	Female	35 to 39	45% (30-63%)	55% (37-70%)	2.8 (1.6-4.3)	7.3% (5.2-9.7%)	10% (7.3-14%)	3 (2.1-4.1)	180 (120-240)
The Bahamas	1990	Female	40 to 44	44% (29-62%)	56% (38-71%)	2.7 (1.4-4.2)	7.3% (4.9-10%)	11% (7.7-16%)	2.7 (1.8-3.8)	140 (96-200)
The Bahamas	1990	Female	45 to 49	42% (26-59%)	58% (41-74%)	2.6 (1.4-4.3)	7.1% (4.6-10%)	11% (7.3-16%)	2.4 (1.6-3.6)	110 (75-170)
The Bahamas	1990	Female	50 to 54	38% (24-56%)	62% (44-76%)	2.3 (1.2-3.9)	7.4% (4.4-11%)	11% (6.7-16%)	3.2 (2-4.9)	130 (80-200)
The Bahamas	1990	Female	55 to 59	34% (21-51%)	66% (49-79%)	2 (0.99-3.4)	6.8% (3.8-10%)	9.5% (5.4-14%)	3.4 (1.9-5.1)	120 (66-180)
The Bahamas	1990	Female	60 to 64	30% (18-46%)	70% (54-82%)	1.7 (0.83-3.1)	5.5% (2.8-9.1%)	7.7% (4-13%)	2.5 (1.3-4.1)	75 (38-120)
The Bahamas	1990	Female	65 to 69	28% (17-42%)	72% (58-83%)	1.5 (0.77-2.5)	5% (2.4-8.4%)	6.6% (3.2-11%)	2.7 (1.4-4.5)	65 (33-110)
The Bahamas	1990	Female	70 to 74	25% (16-38%)	75% (62-84%)	1.2 (0.62-2.3)	4.3% (1.8-7.9%)	5.4% (2.3-9.8%)	3.1 (1.3-5.8)	59 (25-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	1990	Female	75 to 79	23% (13-36%)	77% (64-87%)	1.1 (0.47-2)	3.6% (1.3-7%)	4.4% (1.6-8.4%)	3.2 (1.2-6)	47 (17-89)
The Bahamas	1990	Female	80 to 84	21% (12-33%)	79% (67-88%)	1 (0.48-1.7)	3.4% (1.4-6.4%)	4.1% (1.7-7.6%)	2.6 (1.1-4.9)	29 (12-52)
The Bahamas	1990	Female	85 to 89	21% (12-32%)	79% (68-88%)	1 (0.5-1.8)	3.5% (1.4-6.6%)	4.1% (1.6-7.8%)	1.8 (0.69-3.4)	14 (5.3-26)
The Bahamas	1990	Female	90 to 94	23% (14-35%)	77% (65-86%)	1.2 (0.53-2.2)	3.9% (1.4-7.8%)	4.6% (1.6-9.2%)	0.88 (0.31-1.8)	4.6 (1.7-9.1)
The Bahamas	1990	Female	95 plus	27% (15-44%)	73% (56-85%)	1.4 (0.52-2.9)	4.6% (1-9.6%)	5.3% (1.1-11%)	0.27 (0.056-0.59)	0.95 (0.22-2)
The Bahamas	1990	Male	All Ages	57% (49-66%)	43% (34-51%)	3.7 (3.2-4.2)	12% (9.9-14%)	15% (12-17%)	120 (97-140)	4800 (4100-5600)
The Bahamas	1990	Male	15 to 19	59% (42-75%)	41% (25-58%)	3 (1.6-4.6)	8.9% (5.7-13%)	13% (7.8-19%)	2.3 (1.4-3.5)	200 (130-280)
The Bahamas	1990	Male	20 to 24	69% (52-83%)	31% (17-48%)	5.5 (3.5-7.8)	17% (12-22%)	21% (15-29%)	5.3 (3.6-7.4)	430 (300-560)
The Bahamas	1990	Male	25 to 29	72% (55-84%)	28% (16-45%)	6.6 (4.4-9.3)	18% (13-22%)	21% (15-28%)	8.8 (6.3-11)	630 (470-800)
The Bahamas	1990	Male	30 to 34	73% (56-85%)	27% (15-44%)	6.9 (4.6-9.6)	16% (13-20%)	20% (15-24%)	8.3 (6.3-10)	540 (420-670)
The Bahamas	1990	Male	35 to 39	73% (57-85%)	27% (15-43%)	6.6 (4.1-9.7)	16% (13-20%)	20% (16-24%)	8 (6.1-9.7)	460 (370-560)
The Bahamas	1990	Male	40 to 44	72% (56-84%)	28% (16-44%)	6.3 (3.7-9.7)	17% (14-21%)	21% (17-26%)	8.5 (6.8-10)	440 (350-530)
The Bahamas	1990	Male	45 to 49	70% (51-84%)	30% (16-49%)	6 (3.4-9.7)	18% (14-22%)	21% (17-26%)	11 (8.1-13)	470 (370-580)
The Bahamas	1990	Male	50 to 54	69% (50-83%)	31% (17-50%)	5.6 (3.2-8.9)	17% (13-21%)	21% (16-25%)	11 (8.6-13)	430 (340-520)
The Bahamas	1990	Male	55 to 59	67% (49-82%)	33% (18-51%)	5.1 (2.8-8)	17% (12-21%)	20% (15-25%)	11 (8.1-14)	360 (270-450)
The Bahamas	1990	Male	60 to 64	65% (46-82%)	35% (18-54%)	4.6 (2.5-7.7)	15% (10-19%)	17% (12-23%)	10 (7.1-13)	290 (200-380)
The Bahamas	1990	Male	65 to 69	64% (45-79%)	36% (21-55%)	4.1 (2.3-6.5)	14% (9.6-18%)	16% (11-21%)	11 (7.4-14)	250 (170-330)
The Bahamas	1990	Male	70 to 74	60% (42-76%)	40% (24-58%)	3.4 (1.8-6)	11% (7-15%)	13% (8-18%)	7.3 (4.5-10)	140 (87-190)
The Bahamas	1990	Male	75 to 79	55% (36-74%)	45% (26-64%)	2.9 (1.5-5)	9.2% (5.1-14%)	10% (5.9-15%)	7.6 (4.2-11)	110 (61-160)
The Bahamas	1990	Male	80 to 84	49% (31-67%)	51% (33-69%)	2.4 (1.2-4)	7.9% (4.4-12%)	9% (4.8-14%)	4.5 (2.5-7)	49 (27-74)
The Bahamas	1990	Male	85 to 89	45% (28-62%)	55% (38-72%)	2.1 (1.1-3.7)	7% (3.6-11%)	7.8% (3.9-13%)	2.2 (1.1-3.6)	17 (8.7-28)
The Bahamas	1990	Male	90 to 94	45% (30-62%)	55% (38-70%)	2.2 (1-3.9)	6.5% (3-11%)	7.3% (3.2-12%)	0.76 (0.33-1.3)	4 (1.8-6.7)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	1990	Male	95 plus	48% (28-68%)	52% (32-72%)	2.3 (0.96-4.3)	6.7% (2.6-11%)	7.5% (2.6-13%)	0.17 (0.058-0.3)	0.61 (0.24-1.1)
The Bahamas	1995	Female	All Ages	36% (30-43%)	64% (57-70%)	1.5 (1.3-1.8)	3.9% (3.1-4.9%)	5.2% (3.6-7.1%)	37 (26-51)	1400 (1100-1800)
The Bahamas	1995	Female	15 to 19	42% (28-58%)	58% (42-72%)	1.8 (0.92-2.9)	2.6% (1.7-3.9%)	5.8% (3.5-9%)	0.36 (0.21-0.55)	39 (24-58)
The Bahamas	1995	Female	20 to 24	48% (32-64%)	52% (36-68%)	2.6 (1.5-4)	4.3% (3-5.8%)	6.8% (4.2-9.6%)	0.82 (0.51-1.2)	85 (58-120)
The Bahamas	1995	Female	25 to 29	47% (31-64%)	53% (36-69%)	2.7 (1.6-4)	5.2% (3.7-6.9%)	6.8% (4.6-9.4%)	1.7 (1.1-2.5)	140 (99-190)
The Bahamas	1995	Female	30 to 34	45% (29-63%)	55% (37-71%)	2.6 (1.5-4.1)	5.3% (3.9-7.2%)	6.8% (4.7-9.5%)	2.7 (1.8-3.8)	190 (130-250)
The Bahamas	1995	Female	35 to 39	45% (29-63%)	55% (37-71%)	2.4 (1.4-3.8)	5.6% (4-7.6%)	7.5% (5.2-10%)	3.1 (2.1-4.3)	190 (130-250)
The Bahamas	1995	Female	40 to 44	43% (28-62%)	57% (38-72%)	2.3 (1.2-3.7)	6% (4.1-8.5%)	9% (6-13%)	2.9 (1.9-4.1)	150 (100-210)
The Bahamas	1995	Female	45 to 49	42% (25-59%)	58% (41-75%)	2.2 (1.2-3.7)	5.7% (3.7-8.9%)	9% (5.8-14%)	2.4 (1.6-3.7)	120 (74-180)
The Bahamas	1995	Female	50 to 54	38% (24-55%)	62% (45-76%)	2 (1.1-3.4)	6.3% (3.6-10%)	9.2% (5.4-14%)	3.2 (1.9-5.1)	130 (73-200)
The Bahamas	1995	Female	55 to 59	34% (20-50%)	66% (50-80%)	1.7 (0.84-2.9)	5.6% (3-9%)	7.9% (4.4-12%)	3.4 (1.8-5.4)	120 (63-190)
The Bahamas	1995	Female	60 to 64	30% (18-47%)	70% (53-82%)	1.5 (0.71-2.6)	4.5% (2.1-7.4%)	6.4% (3.2-11%)	2.4 (1.2-3.9)	71 (34-120)
The Bahamas	1995	Female	65 to 69	27% (17-41%)	73% (59-83%)	1.3 (0.66-2.1)	4.1% (1.8-7.1%)	5.4% (2.5-9%)	2.9 (1.3-4.9)	70 (32-120)
The Bahamas	1995	Female	70 to 74	25% (16-39%)	75% (61-84%)	1.1 (0.54-1.9)	3.5% (1.3-6.6%)	4.3% (1.6-8.1%)	2.8 (1.1-5.3)	53 (19-99)
The Bahamas	1995	Female	75 to 79	22% (13-36%)	78% (64-87%)	0.94 (0.43-1.7)	2.9% (0.91-5.7%)	3.6% (1.1-7.1%)	2.5 (0.79-4.9)	36 (11-74)
The Bahamas	1995	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.88 (0.42-1.5)	2.8% (1-5.3%)	3.3% (1.2-6.3%)	2.5 (0.94-4.7)	27 (10-51)
The Bahamas	1995	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.9 (0.41-1.5)	2.8% (0.97-5.6%)	3.3% (1.1-6.6%)	2.1 (0.68-4.1)	16 (5.4-31)
The Bahamas	1995	Female	90 to 94	23% (13-35%)	77% (65-87%)	1 (0.46-1.9)	3.2% (0.99-6.6%)	3.7% (1.1-7.8%)	1 (0.3-2.2)	5.4 (1.6-11)
The Bahamas	1995	Female	95 plus	27% (14-44%)	73% (56-86%)	1.2 (0.43-2.5)	3.7% (0.77-8.5%)	4.3% (0.72-10%)	0.34 (0.057-0.79)	1.2 (0.25-2.7)
The Bahamas	1995	Male	All Ages	58% (49-66%)	42% (34-51%)	3.2 (2.7-3.7)	10% (8.5-12%)	12% (9.7-14%)	110 (91-140)	4600 (3900-5500)
The Bahamas	1995	Male	15 to 19	59% (41-74%)	41% (26-59%)	2.5 (1.3-4.1)	6.5% (4.4-9.1%)	11% (6.7-16%)	0.94 (0.56-1.4)	95 (65-130)
The Bahamas	1995	Male	20 to 24	69% (51-82%)	31% (18-49%)	4.7 (3-6.9)	14% (10-19%)	18% (12-24%)	4.9 (3.3-6.8)	390 (280-530)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	1995	Male	25 to 29	72% (54-84%)	28% (16-46%)	5.7 (3.8-8)	14% (10-17%)	16% (11-21%)	6.5 (4.5-8.5)	480 (350-610)
The Bahamas	1995	Male	30 to 34	72% (54-85%)	28% (15-46%)	5.9 (3.9-8.3)	12% (9-15%)	13% (9.5-17%)	9.7 (7.1-13)	630 (470-800)
The Bahamas	1995	Male	35 to 39	72% (56-85%)	28% (15-44%)	5.6 (3.5-8.3)	12% (9-15%)	13% (9.8-17%)	8.4 (6.2-11)	500 (380-620)
The Bahamas	1995	Male	40 to 44	71% (55-84%)	29% (16-45%)	5.4 (3.1-8.1)	13% (9.9-16%)	15% (11-19%)	9.8 (7.2-12)	500 (380-630)
The Bahamas	1995	Male	45 to 49	70% (51-84%)	30% (16-49%)	5.1 (2.9-8.1)	14% (11-17%)	17% (13-20%)	8.5 (6.4-11)	390 (300-490)
The Bahamas	1995	Male	50 to 54	68% (50-83%)	32% (17-50%)	4.7 (2.6-7.6)	14% (11-18%)	17% (13-21%)	9.9 (7.2-12)	390 (290-490)
The Bahamas	1995	Male	55 to 59	66% (49-82%)	34% (18-51%)	4.3 (2.3-6.8)	14% (9.9-18%)	17% (12-22%)	12 (8.1-15)	390 (280-510)
The Bahamas	1995	Male	60 to 64	64% (45-81%)	36% (19-55%)	3.9 (2.1-6.5)	13% (8.3-17%)	15% (9.9-20%)	9.7 (6.4-13)	280 (180-370)
The Bahamas	1995	Male	65 to 69	63% (44-79%)	37% (21-56%)	3.5 (1.9-5.5)	12% (7.7-16%)	14% (8.7-19%)	12 (7.5-16)	270 (180-370)
The Bahamas	1995	Male	70 to 74	60% (41-75%)	40% (25-59%)	2.9 (1.6-5)	9.4% (5.3-14%)	11% (6-16%)	7.5 (4.3-11)	140 (82-220)
The Bahamas	1995	Male	75 to 79	54% (35-73%)	46% (27-65%)	2.4 (1.2-4.3)	7.9% (4.1-12%)	8.8% (4.5-14%)	7.1 (3.7-11)	100 (53-160)
The Bahamas	1995	Male	80 to 84	49% (30-67%)	51% (33-70%)	2 (1-3.4)	6.6% (3.4-10%)	7.5% (3.8-12%)	4 (2-6.3)	43 (23-69)
The Bahamas	1995	Male	85 to 89	44% (27-61%)	56% (39-73%)	1.8 (0.87-3.1)	5.6% (2.6-9.4%)	6.3% (2.8-11%)	2.5 (1.1-4.1)	19 (8.9-32)
The Bahamas	1995	Male	90 to 94	45% (29-62%)	55% (38-71%)	1.8 (0.88-3.2)	5.2% (2.3-9.2%)	5.8% (2.4-10%)	0.87 (0.35-1.6)	4.7 (2-8.2)
The Bahamas	1995	Male	95 plus	48% (29-68%)	52% (32-71%)	2 (0.79-3.7)	5.4% (1.9-10%)	6% (1.7-12%)	0.21 (0.061-0.4)	0.76 (0.26-1.4)
The Bahamas	2000	Female	All Ages	35% (29-43%)	65% (57-71%)	1.3 (1.1-1.6)	3.1% (2.4-4%)	3.9% (2.5-5.5%)	33 (21-47)	1200 (920-1600)
The Bahamas	2000	Female	15 to 19	41% (28-58%)	59% (42-72%)	1.5 (0.8-2.5)	2% (1.3-3%)	4.4% (2.5-7.2%)	0.26 (0.14-0.44)	32 (20-49)
The Bahamas	2000	Female	20 to 24	47% (31-63%)	53% (37-69%)	2.2 (1.2-3.5)	3.2% (2.2-4.3%)	4.3% (2.5-6.4%)	0.45 (0.26-0.69)	55 (37-79)
The Bahamas	2000	Female	25 to 29	47% (30-63%)	53% (37-70%)	2.3 (1.4-3.5)	3.4% (2.4-4.6%)	4% (2.5-5.7%)	0.9 (0.56-1.3)	88 (60-120)
The Bahamas	2000	Female	30 to 34	45% (28-62%)	55% (38-72%)	2.2 (1.3-3.5)	4.6% (3.3-6.3%)	5.9% (4.2-8.3%)	2.1 (1.5-3)	150 (110-210)
The Bahamas	2000	Female	35 to 39	44% (29-62%)	56% (38-71%)	2 (1.2-3.1)	4.1% (2.9-5.6%)	5.5% (3.8-7.7%)	2.5 (1.7-3.6)	160 (110-220)
The Bahamas	2000	Female	40 to 44	43% (28-61%)	57% (39-72%)	1.9 (1.1-3)	4.3% (2.9-6.1%)	6.1% (4.1-8.7%)	2.6 (1.7-3.8)	140 (94-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	2000	Female	45 to 49	41% (25-58%)	59% (42-75%)	1.9 (1-3.1)	3.9% (2.5-6%)	5.8% (3.8-8.9%)	2.1 (1.3-3.2)	100 (66-160)
The Bahamas	2000	Female	50 to 54	37% (23-54%)	63% (46-77%)	1.7 (0.89-2.9)	4.8% (2.8-7.7%)	6.9% (4-11%)	2.9 (1.7-4.6)	120 (68-190)
The Bahamas	2000	Female	55 to 59	33% (20-50%)	67% (50-80%)	1.4 (0.69-2.5)	4.4% (2.3-7.3%)	6.2% (3.3-10%)	2.8 (1.5-4.6)	97 (51-160)
The Bahamas	2000	Female	60 to 64	29% (17-45%)	71% (55-83%)	1.2 (0.57-2.2)	3.5% (1.6-6.1%)	4.9% (2.4-8.4%)	2.4 (1.1-4.1)	70 (32-120)
The Bahamas	2000	Female	65 to 69	27% (17-41%)	73% (59-83%)	1.1 (0.54-1.8)	3.1% (1.3-5.5%)	4.1% (1.8-7.2%)	2.7 (1.2-4.7)	64 (26-110)
The Bahamas	2000	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.89 (0.43-1.6)	2.7% (0.86-5.2%)	3.4% (1.1-6.5%)	2.8 (0.91-5.2)	53 (17-100)
The Bahamas	2000	Female	75 to 79	22% (13-35%)	78% (65-87%)	0.79 (0.35-1.4)	2.3% (0.52-4.7%)	2.8% (0.69-5.8%)	2.4 (0.57-4.9)	35 (7.9-71)
The Bahamas	2000	Female	80 to 84	20% (11-32%)	80% (68-89%)	0.74 (0.34-1.3)	2.1% (0.7-4.2%)	2.6% (0.83-5%)	2.3 (0.74-4.5)	25 (8-49)
The Bahamas	2000	Female	85 to 89	20% (12-31%)	80% (69-88%)	0.75 (0.36-1.3)	2.1% (0.69-4.4%)	2.5% (0.75-5.2%)	1.9 (0.59-4.1)	15 (4.7-31)
The Bahamas	2000	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.85 (0.37-1.7)	2.3% (0.63-5.3%)	2.7% (0.62-6.2%)	1.2 (0.27-2.7)	6.2 (1.6-14)
The Bahamas	2000	Female	95 plus	26% (14-43%)	74% (57-86%)	1 (0.35-2.1)	2.8% (0.46-6.8%)	3.1% (0.37-7.8%)	0.39 (0.047-0.99)	1.4 (0.23-3.4)
The Bahamas	2000	Male	All Ages	58% (49-66%)	42% (34-51%)	2.8 (2.3-3.2)	8.9% (7.4-10%)	10% (7.7-12%)	100 (78-130)	4200 (3500-5000)
The Bahamas	2000	Male	15 to 19	58% (41-74%)	42% (26-59%)	2.2 (1.1-3.5)	6.6% (4.2-9.5%)	9.8% (5.7-15%)	1.5 (0.87-2.3)	130 (85-190)
The Bahamas	2000	Male	20 to 24	68% (51-82%)	32% (18-49%)	4 (2.4-5.8)	14% (10-19%)	18% (11-24%)	4.3 (2.8-6.1)	340 (240-460)
The Bahamas	2000	Male	25 to 29	71% (53-84%)	29% (16-47%)	4.8 (3.1-6.9)	14% (10-19%)	17% (12-23%)	6.7 (4.6-9.1)	490 (350-640)
The Bahamas	2000	Male	30 to 34	72% (54-85%)	28% (15-46%)	5 (3.3-7.2)	10% (7.7-13%)	11% (7.9-15%)	6.1 (4.3-8.1)	420 (310-540)
The Bahamas	2000	Male	35 to 39	72% (56-84%)	28% (16-44%)	4.8 (3-6.9)	8.7% (6.3-11%)	9.3% (6.4-12%)	6.4 (4.3-8.8)	410 (290-530)
The Bahamas	2000	Male	40 to 44	71% (55-84%)	29% (16-45%)	4.5 (2.7-6.8)	9.2% (6.8-12%)	10% (7.2-13%)	7.1 (5-9.4)	390 (280-500)
The Bahamas	2000	Male	45 to 49	69% (50-83%)	31% (17-50%)	4.3 (2.4-6.9)	11% (8.1-14%)	13% (9.1-17%)	8.9 (6.3-12)	420 (300-540)
The Bahamas	2000	Male	50 to 54	68% (50-83%)	32% (17-50%)	3.9 (2.2-6.3)	12% (8.5-15%)	14% (9.8-18%)	9.8 (6.6-13)	390 (280-510)
The Bahamas	2000	Male	55 to 59	66% (48-81%)	34% (19-52%)	3.6 (1.9-5.8)	12% (7.9-16%)	14% (9.3-19%)	11 (6.8-14)	360 (240-480)
The Bahamas	2000	Male	60 to 64	64% (44-81%)	36% (19-56%)	3.3 (1.7-5.5)	11% (6.5-15%)	13% (7.6-18%)	10 (6-14)	290 (180-400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	2000	Male	65 to 69	62% (43-78%)	38% (22-57%)	2.9 (1.5-4.6)	10% (6.2-14%)	12% (6.9-16%)	11 (6.3-15)	250 (150-350)
The Bahamas	2000	Male	70 to 74	59% (41-75%)	41% (25-59%)	2.4 (1.3-4.1)	7.6% (4-12%)	8.7% (4.4-14%)	7 (3.5-11)	130 (69-210)
The Bahamas	2000	Male	75 to 79	54% (34-72%)	46% (28-66%)	2 (0.98-3.7)	6.3% (2.9-10%)	7.1% (3.2-12%)	6.4 (3-11)	93 (43-150)
The Bahamas	2000	Male	80 to 84	48% (30-66%)	52% (34-70%)	1.7 (0.85-2.8)	5.2% (2.5-8.6%)	5.8% (2.8-9.6%)	4 (1.9-6.7)	44 (22-72)
The Bahamas	2000	Male	85 to 89	43% (27-60%)	57% (40-73%)	1.5 (0.75-2.5)	4.4% (2.1-7.7%)	4.8% (2.3-8.6%)	2.3 (1.1-4.2)	18 (8.8-32)
The Bahamas	2000	Male	90 to 94	44% (28-61%)	56% (39-72%)	1.5 (0.75-2.7)	4.1% (1.8-7.7%)	4.5% (1.8-8.7%)	1.1 (0.44-2.1)	5.8 (2.5-11)
The Bahamas	2000	Male	95 plus	47% (27-67%)	53% (33-73%)	1.6 (0.65-3.1)	4.3% (1.4-8.4%)	4.7% (1.3-9.6%)	0.26 (0.072-0.52)	0.96 (0.31-1.9)
The Bahamas	2005	Female	All Ages	35% (29-43%)	65% (57-71%)	1.2 (1-1.5)	2.9% (2.2-3.8%)	3.7% (2.4-5.3%)	35 (22-50)	1200 (940-1600)
The Bahamas	2005	Female	15 to 19	41% (28-57%)	59% (43-72%)	1.3 (0.71-2.3)	2% (1.3-2.9%)	4.4% (2.5-7%)	0.27 (0.15-0.44)	33 (20-49)
The Bahamas	2005	Female	20 to 24	47% (31-63%)	53% (37-69%)	2 (1.1-3.1)	3.2% (2.2-4.3%)	4.7% (2.8-7.1%)	0.49 (0.29-0.76)	61 (40-86)
The Bahamas	2005	Female	25 to 29	46% (30-63%)	54% (37-70%)	2.1 (1.2-3.2)	3.7% (2.6-5%)	4.9% (3.1-7.2%)	0.99 (0.62-1.5)	93 (64-130)
The Bahamas	2005	Female	30 to 34	44% (28-62%)	56% (38-72%)	2 (1.1-3.1)	3.8% (2.7-5.3%)	5% (3.4-7.2%)	1.5 (0.99-2.1)	110 (77-150)
The Bahamas	2005	Female	35 to 39	43% (28-62%)	57% (38-72%)	1.8 (1-2.9)	4.1% (2.9-5.8%)	5.9% (4.1-8.4%)	2.3 (1.6-3.3)	140 (100-200)
The Bahamas	2005	Female	40 to 44	42% (28-61%)	58% (39-72%)	1.7 (0.93-2.8)	4.2% (2.9-6%)	6.5% (4.5-9.3%)	2.5 (1.7-3.6)	140 (94-200)
The Bahamas	2005	Female	45 to 49	40% (24-58%)	60% (42-76%)	1.7 (0.91-2.8)	4.2% (2.7-6.2%)	6.8% (4.4-10%)	2.7 (1.8-4.1)	130 (83-190)
The Bahamas	2005	Female	50 to 54	37% (22-54%)	63% (46-78%)	1.5 (0.82-2.5)	4.4% (2.6-6.8%)	6.6% (4.1-10%)	3.3 (2-5.1)	140 (81-210)
The Bahamas	2005	Female	55 to 59	33% (20-49%)	67% (51-80%)	1.3 (0.65-2.2)	4% (2.2-6.6%)	5.8% (3.3-9.3%)	3.3 (1.9-5.3)	110 (62-190)
The Bahamas	2005	Female	60 to 64	29% (17-45%)	71% (55-83%)	1.1 (0.54-1.9)	3.2% (1.5-5.4%)	4.6% (2.3-7.8%)	2.6 (1.3-4.4)	77 (36-130)
The Bahamas	2005	Female	65 to 69	27% (17-40%)	73% (60-83%)	0.95 (0.5-1.6)	2.8% (1.2-4.8%)	3.8% (1.7-6.3%)	2.7 (1.2-4.6)	64 (28-110)
The Bahamas	2005	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.8 (0.39-1.4)	2.3% (0.77-4.6%)	3.1% (1.1-5.8%)	3 (1.1-5.7)	56 (18-110)
The Bahamas	2005	Female	75 to 79	22% (12-35%)	78% (65-88%)	0.7 (0.32-1.3)	2% (0.57-4.1%)	2.5% (0.77-5.2%)	2.6 (0.78-5.3)	38 (11-78)
The Bahamas	2005	Female	80 to 84	20% (11-32%)	80% (68-89%)	0.65 (0.31-1.1)	1.9% (0.66-3.6%)	2.3% (0.83-4.5%)	2.6 (0.91-5)	28 (10-54)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	2005	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.66 (0.32-1.2)	1.8% (0.56-3.7%)	2.1% (0.65-4.4%)	1.9 (0.58-4)	15 (4.7-31)
The Bahamas	2005	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.76 (0.34-1.6)	2% (0.5-4.6%)	2.3% (0.53-5.4%)	1.2 (0.27-2.9)	6.5 (1.6-15)
The Bahamas	2005	Female	95 plus	26% (13-43%)	74% (57-87%)	0.91 (0.37-1.9)	2.4% (0.37-5.6%)	2.6% (0.28-6.7%)	0.5 (0.052-1.3)	1.8 (0.27-4.3)
The Bahamas	2005	Male	All Ages	59% (51-67%)	41% (33-49%)	2.6 (2.2-3)	9.3% (7.6-11%)	10% (8.1-13%)	110 (87-140)	4700 (3900-5500)
The Bahamas	2005	Male	15 to 19	58% (40-74%)	42% (26-60%)	1.9 (1-3.1)	6.5% (4.2-9.5%)	9.4% (5.6-14%)	1.7 (0.97-2.6)	150 (95-210)
The Bahamas	2005	Male	20 to 24	68% (50-82%)	32% (18-50%)	3.6 (2.1-5.3)	13% (9.3-18%)	16% (10-23%)	4.9 (3.1-7.1)	390 (270-540)
The Bahamas	2005	Male	25 to 29	71% (53-84%)	29% (16-47%)	4.3 (2.7-6.2)	14% (10-19%)	17% (11-23%)	6.2 (4.2-8.7)	460 (330-620)
The Bahamas	2005	Male	30 to 34	72% (54-84%)	28% (16-46%)	4.5 (2.9-6.5)	13% (9.6-16%)	15% (11-20%)	6.6 (4.8-8.8)	440 (330-570)
The Bahamas	2005	Male	35 to 39	71% (56-84%)	29% (16-44%)	4.3 (2.6-6.3)	11% (8.3-14%)	13% (9.2-17%)	8.6 (6-11)	520 (370-660)
The Bahamas	2005	Male	40 to 44	70% (54-84%)	30% (16-46%)	4.1 (2.4-6.2)	11% (8.2-14%)	13% (9.4-17%)	9 (6.3-12)	480 (350-620)
The Bahamas	2005	Male	45 to 49	69% (49-83%)	31% (17-51%)	3.9 (2.2-6.1)	12% (8.6-16%)	15% (10-19%)	11 (7.8-15)	530 (380-680)
The Bahamas	2005	Male	50 to 54	67% (49-82%)	33% (18-51%)	3.5 (2-5.7)	12% (8.1-16%)	15% (9.7-19%)	11 (7.3-15)	450 (310-600)
The Bahamas	2005	Male	55 to 59	65% (47-80%)	35% (20-53%)	3.2 (1.8-5.2)	11% (7.3-15%)	14% (8.9-19%)	11 (7-15)	390 (250-530)
The Bahamas	2005	Male	60 to 64	64% (43-80%)	36% (20-57%)	2.9 (1.6-4.9)	9.8% (5.7-14%)	12% (7.1-17%)	10 (6-14)	290 (180-420)
The Bahamas	2005	Male	65 to 69	62% (43-78%)	38% (22-57%)	2.6 (1.4-4.2)	9% (5.5-13%)	11% (6.3-16%)	10 (6.1-15)	240 (150-350)
The Bahamas	2005	Male	70 to 74	59% (40-75%)	41% (25-60%)	2.2 (1.1-3.7)	6.8% (3.6-11%)	8% (4-13%)	7.3 (3.7-12)	140 (72-230)
The Bahamas	2005	Male	75 to 79	54% (34-72%)	46% (28-66%)	1.8 (0.89-3.2)	5.5% (2.6-9.4%)	6.4% (2.9-11%)	5.8 (2.6-9.7)	86 (40-140)
The Bahamas	2005	Male	80 to 84	48% (30-66%)	52% (34-70%)	1.5 (0.78-2.5)	4.6% (2.3-7.7%)	5.3% (2.5-8.9%)	4.2 (2-7.1)	46 (23-78)
The Bahamas	2005	Male	85 to 89	43% (27-60%)	57% (40-73%)	1.3 (0.66-2.3)	3.9% (1.9-6.9%)	4.4% (2.1-7.9%)	2.3 (1.1-4.2)	18 (8.9-33)
The Bahamas	2005	Male	90 to 94	43% (28-61%)	57% (39-72%)	1.3 (0.66-2.5)	3.6% (1.6-6.8%)	3.9% (1.6-7.8%)	0.97 (0.4-1.9)	5.3 (2.3-10)
The Bahamas	2005	Male	95 plus	46% (27-67%)	54% (33-73%)	1.5 (0.58-2.8)	3.7% (1.2-7.5%)	4% (1.1-8.5%)	0.31 (0.082-0.65)	1.1 (0.36-2.3)
The Bahamas	2010	Female	All Ages	35% (29-42%)	65% (58-71%)	1.2 (0.95-1.5)	2.9% (2.1-3.9%)	3.6% (2.3-5.3%)	39 (25-59)	1400 (1000-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	2010	Female	15 to 19	41% (27-57%)	59% (43-73%)	1.3 (0.66-2.1)	1.9% (1.3-3%)	4.3% (2.4-6.8%)	0.29 (0.16-0.49)	35 (21-54)
The Bahamas	2010	Female	20 to 24	47% (30-63%)	53% (37-70%)	1.9 (1-3)	3.5% (2.4-4.9%)	5.7% (3.3-8.8%)	0.64 (0.36-1)	74 (48-110)
The Bahamas	2010	Female	25 to 29	46% (30-63%)	54% (37-70%)	2 (1.1-3)	3.5% (2.5-4.7%)	4.7% (2.9-7%)	0.91 (0.55-1.4)	90 (61-130)
The Bahamas	2010	Female	30 to 34	44% (28-62%)	56% (38-72%)	1.9 (1.1-3)	4% (2.9-5.6%)	5.8% (3.9-8.3%)	1.5 (1-2.2)	110 (79-160)
The Bahamas	2010	Female	35 to 39	43% (28-61%)	57% (39-72%)	1.8 (0.99-2.9)	4.1% (2.9-5.7%)	6.1% (4.3-8.6%)	2.3 (1.5-3.2)	140 (99-200)
The Bahamas	2010	Female	40 to 44	42% (27-61%)	58% (39-73%)	1.7 (0.89-2.7)	4.1% (2.8-6%)	6.7% (4.5-9.6%)	2.6 (1.7-3.7)	140 (94-210)
The Bahamas	2010	Female	45 to 49	40% (25-58%)	60% (42-75%)	1.6 (0.85-2.7)	3.9% (2.5-6%)	6.4% (4.1-9.7%)	3.1 (1.9-4.7)	150 (94-230)
The Bahamas	2010	Female	50 to 54	37% (23-54%)	63% (46-77%)	1.4 (0.77-2.4)	4.1% (2.4-6.5%)	6.4% (3.9-9.9%)	3.9 (2.4-6.2)	160 (93-250)
The Bahamas	2010	Female	55 to 59	33% (20-49%)	67% (51-80%)	1.2 (0.62-2.1)	3.8% (2.1-6.2%)	5.6% (3.3-9.1%)	3.9 (2.3-6.3)	140 (77-220)
The Bahamas	2010	Female	60 to 64	29% (17-45%)	71% (55-83%)	1 (0.51-1.8)	3% (1.4-5.4%)	4.4% (2.2-7.6%)	3 (1.5-5.2)	88 (43-160)
The Bahamas	2010	Female	65 to 69	27% (17-40%)	73% (60-83%)	0.91 (0.49-1.5)	2.6% (1.2-4.5%)	3.6% (1.8-6.2%)	2.9 (1.4-5.1)	70 (33-130)
The Bahamas	2010	Female	70 to 74	24% (15-37%)	76% (63-85%)	0.77 (0.38-1.5)	2.2% (0.81-4.5%)	3% (1.2-5.8%)	3.4 (1.3-6.7)	63 (22-130)
The Bahamas	2010	Female	75 to 79	21% (13-35%)	79% (65-87%)	0.67 (0.31-1.2)	1.9% (0.55-3.9%)	2.4% (0.76-5%)	3 (0.93-6.2)	43 (13-91)
The Bahamas	2010	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.63 (0.29-1.1)	1.8% (0.58-3.5%)	2.2% (0.72-4.3%)	3.1 (1-6.3)	34 (11-68)
The Bahamas	2010	Female	85 to 89	20% (11-30%)	80% (70-89%)	0.64 (0.3-1.2)	1.7% (0.54-3.8%)	2% (0.61-4.5%)	2.3 (0.7-4.9)	18 (5.6-38)
The Bahamas	2010	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.73 (0.31-1.5)	1.9% (0.46-4.5%)	2.2% (0.47-5.3%)	1.3 (0.28-3.2)	7.1 (1.7-16)
The Bahamas	2010	Female	95 plus	26% (13-43%)	74% (57-87%)	0.89 (0.31-1.9)	2.3% (0.3-5.8%)	2.5% (0.18-6.7%)	0.57 (0.039-1.5)	2.1 (0.26-5.2)
The Bahamas	2010	Male	All Ages	59% (51-67%)	41% (33-49%)	2.6 (2.1-3.2)	9.4% (7.7-11%)	10% (7.9-13%)	120 (94-160)	5200 (4200-6200)
The Bahamas	2010	Male	15 to 19	58% (40-74%)	42% (26-60%)	1.9 (0.98-3)	6.4% (4-9.3%)	9.2% (5.2-14%)	1.8 (1-2.8)	160 (100-230)
The Bahamas	2010	Male	20 to 24	68% (50-82%)	32% (18-50%)	3.5 (2-5.1)	13% (9.1-19%)	16% (10-24%)	6.3 (4-9.3)	500 (330-690)
The Bahamas	2010	Male	25 to 29	71% (54-83%)	29% (17-46%)	4.2 (2.5-6.1)	14% (9.6-18%)	17% (11-24%)	6.2 (4-8.7)	470 (330-630)
The Bahamas	2010	Male	30 to 34	72% (54-85%)	28% (15-46%)	4.3 (2.7-6.4)	13% (9.5-17%)	16% (11-21%)	6.5 (4.5-8.9)	440 (320-580)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	2010	Male	35 to 39	71% (55-84%)	29% (16-45%)	4.1 (2.5-6.3)	13% (9.2-17%)	15% (11-20%)	11 (7.6-15)	640 (460-830)
The Bahamas	2010	Male	40 to 44	70% (55-83%)	30% (17-45%)	3.9 (2.3-6.1)	12% (8.7-16%)	15% (10-20%)	11 (7.4-14)	580 (400-740)
The Bahamas	2010	Male	45 to 49	69% (49-83%)	31% (17-51%)	3.7 (2.1-5.9)	11% (7.9-15%)	14% (9.7-18%)	12 (8.1-16)	570 (400-740)
The Bahamas	2010	Male	50 to 54	67% (49-83%)	33% (17-51%)	3.4 (1.9-5.4)	11% (7.6-15%)	14% (9.6-19%)	12 (7.9-16)	490 (330-650)
The Bahamas	2010	Male	55 to 59	65% (47-81%)	35% (19-53%)	3.1 (1.7-5)	11% (7.1-15%)	14% (8.8-19%)	12 (8-17)	430 (280-600)
The Bahamas	2010	Male	60 to 64	63% (44-80%)	37% (20-56%)	2.8 (1.5-4.7)	9.4% (5.6-14%)	12% (7-17%)	11 (6.6-16)	320 (200-460)
The Bahamas	2010	Male	65 to 69	62% (43-78%)	38% (22-57%)	2.5 (1.4-4)	8.4% (5-12%)	10% (6-15%)	10 (6.1-15)	250 (150-360)
The Bahamas	2010	Male	70 to 74	58% (41-74%)	42% (26-59%)	2.1 (1.1-3.8)	6.5% (3.4-11%)	7.7% (3.9-13%)	8.4 (4.2-14)	160 (84-270)
The Bahamas	2010	Male	75 to 79	53% (34-72%)	47% (28-66%)	1.7 (0.85-3.1)	5.2% (2.5-9%)	6% (2.8-10%)	6.1 (2.8-10)	91 (43-150)
The Bahamas	2010	Male	80 to 84	48% (29-66%)	52% (34-71%)	1.5 (0.72-2.4)	4.3% (2.1-7.4%)	4.9% (2.4-8.4%)	4.7 (2.3-8.1)	51 (26-88)
The Bahamas	2010	Male	85 to 89	43% (27-60%)	57% (40-73%)	1.3 (0.63-2.3)	3.7% (1.7-6.6%)	4.1% (1.9-7.4%)	2.7 (1.2-4.9)	21 (10-38)
The Bahamas	2010	Male	90 to 94	43% (27-60%)	57% (40-73%)	1.3 (0.61-2.4)	3.4% (1.5-6.6%)	3.7% (1.5-7.5%)	1.1 (0.43-2.2)	6 (2.6-12)
The Bahamas	2010	Male	95 plus	46% (26-66%)	54% (34-74%)	1.4 (0.53-2.7)	3.5% (1.2-7%)	3.8% (1-8.2%)	0.31 (0.086-0.67)	1.2 (0.38-2.4)
The Bahamas	2016	Female	All Ages	35% (29-42%)	65% (58-71%)	1.1 (0.65-1.6)	2.7% (1.6-4%)	3.1% (1.5-5.4%)	44 (21-75)	1500 (870-2300)
The Bahamas	2016	Female	15 to 19	41% (27-57%)	59% (43-73%)	1.2 (0.52-2.1)	1.9% (1.1-2.9%)	4% (2.1-7%)	0.24 (0.12-0.44)	30 (18-48)
The Bahamas	2016	Female	20 to 24	46% (30-62%)	54% (38-70%)	1.7 (0.81-3)	3.2% (2.1-4.7%)	5% (2.6-7.9%)	0.58 (0.29-1)	73 (45-110)
The Bahamas	2016	Female	25 to 29	46% (29-63%)	54% (37-71%)	1.8 (0.84-3)	3.5% (2.3-5%)	4.9% (2.8-7.6%)	0.99 (0.52-1.6)	97 (61-140)
The Bahamas	2016	Female	30 to 34	44% (28-62%)	56% (38-72%)	1.7 (0.83-2.9)	3.8% (2.5-5.5%)	5.5% (3.4-8.3%)	1.7 (0.97-2.6)	130 (80-190)
The Bahamas	2016	Female	35 to 39	43% (28-61%)	57% (39-72%)	1.6 (0.78-2.8)	3.9% (2.5-5.7%)	5.9% (3.8-8.7%)	2 (1.2-3.1)	130 (80-190)
The Bahamas	2016	Female	40 to 44	42% (27-61%)	58% (39-73%)	1.5 (0.71-2.6)	3.8% (2.3-5.6%)	6.1% (3.8-9.4%)	2.5 (1.5-3.9)	140 (85-220)
The Bahamas	2016	Female	45 to 49	40% (24-57%)	60% (43-76%)	1.4 (0.68-2.6)	3.7% (2.2-5.9%)	6.2% (3.8-9.9%)	3 (1.8-4.7)	140 (83-230)
The Bahamas	2016	Female	50 to 54	37% (22-54%)	63% (46-78%)	1.3 (0.59-2.3)	3.9% (2.1-6.4%)	6% (3.4-9.7%)	4.5 (2.5-7.4)	180 (96-300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	2016	Female	55 to 59	32% (20-49%)	68% (51-80%)	1.1 (0.47-2)	3.5% (1.7-5.9%)	5.2% (2.7-8.5%)	4.9 (2.5-8.4)	170 (81-290)
The Bahamas	2016	Female	60 to 64	29% (17-44%)	71% (56-83%)	0.94 (0.42-1.8)	2.7% (1.1-5.1%)	3.9% (1.7-7.1%)	3.9 (1.7-7.2)	110 (46-220)
The Bahamas	2016	Female	65 to 69	26% (16-40%)	74% (60-84%)	0.82 (0.36-1.5)	2.3% (0.89-4.5%)	3.2% (1.3-6%)	3.8 (1.5-7.5)	90 (33-180)
The Bahamas	2016	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.7 (0.3-1.4)	2% (0.5-4.5%)	2.6% (0.74-5.7%)	3.6 (1-8.1)	67 (17-150)
The Bahamas	2016	Female	75 to 79	21% (13-34%)	79% (66-87%)	0.61 (0.24-1.2)	1.7% (0.29-4%)	2.1% (0.43-4.9%)	3.6 (0.69-8.7)	52 (8.9-130)
The Bahamas	2016	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.57 (0.24-1.1)	1.6% (0.28-3.6%)	1.9% (0.38-4.3%)	3.9 (0.75-9.1)	42 (7.5-98)
The Bahamas	2016	Female	85 to 89	20% (11-31%)	80% (69-89%)	0.58 (0.23-1.1)	1.5% (0.26-3.6%)	1.8% (0.27-4.2%)	2.9 (0.45-7.1)	22 (3.8-54)
The Bahamas	2016	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.65 (0.25-1.3)	1.6% (0.18-4.1%)	1.8% (0.16-4.7%)	1.6 (0.14-4.2)	8.4 (0.94-22)
The Bahamas	2016	Female	95 plus	26% (14-43%)	74% (57-86%)	0.78 (0.25-1.7)	1.9% (0.16-5.1%)	2.1% (0.061-6%)	0.59 (0.016-1.7)	2.2 (0.17-5.9)
The Bahamas	2016	Male	All Ages	59% (52-67%)	41% (33-48%)	2.4 (1.5-3.4)	8.7% (6.1-11%)	9% (5.6-13%)	130 (82-190)	5400 (3700-7200)
The Bahamas	2016	Male	15 to 19	57% (40-74%)	43% (26-60%)	1.7 (0.79-3)	5.9% (3.6-8.9%)	8.5% (4.6-14%)	1.4 (0.73-2.3)	130 (78-200)
The Bahamas	2016	Male	20 to 24	68% (51-82%)	32% (18-49%)	3.1 (1.6-5.2)	12% (7.9-17%)	15% (8.3-22%)	5.5 (3.1-8.4)	450 (280-660)
The Bahamas	2016	Male	25 to 29	71% (53-83%)	29% (17-47%)	3.7 (1.9-5.9)	13% (8.6-18%)	16% (9.6-24%)	6.3 (3.7-9.6)	480 (310-680)
The Bahamas	2016	Male	30 to 34	71% (54-85%)	29% (15-46%)	3.9 (2-6.4)	12% (8.3-17%)	15% (9.6-21%)	7.6 (4.8-11)	520 (340-730)
The Bahamas	2016	Male	35 to 39	71% (55-85%)	29% (15-45%)	3.7 (1.9-6.1)	12% (8.1-16%)	14% (9.1-20%)	9.1 (5.7-13)	540 (360-740)
The Bahamas	2016	Male	40 to 44	70% (54-83%)	30% (17-46%)	3.5 (1.8-6)	11% (7.4-15%)	14% (8.6-18%)	10 (6.3-14)	540 (360-750)
The Bahamas	2016	Male	45 to 49	69% (49-83%)	31% (17-51%)	3.3 (1.6-5.7)	11% (6.9-15%)	13% (8.1-19%)	11 (6.8-16)	540 (340-760)
The Bahamas	2016	Male	50 to 54	67% (48-82%)	33% (18-52%)	3 (1.5-5.2)	11% (6.4-15%)	13% (7.8-19%)	14 (7.9-20)	570 (340-800)
The Bahamas	2016	Male	55 to 59	65% (47-81%)	35% (19-53%)	2.8 (1.3-4.9)	9.8% (5.7-14%)	12% (7-18%)	15 (8.2-22)	510 (290-750)
The Bahamas	2016	Male	60 to 64	63% (43-80%)	37% (20-57%)	2.5 (1.2-4.5)	8.3% (4.4-13%)	10% (5.3-16%)	13 (6.6-21)	380 (200-600)
The Bahamas	2016	Male	65 to 69	62% (43-78%)	38% (22-57%)	2.2 (1.1-4)	7.4% (3.7-12%)	8.9% (4.3-15%)	13 (6.5-21)	310 (160-510)
The Bahamas	2016	Male	70 to 74	58% (39-74%)	42% (26-61%)	1.9 (0.86-3.4)	5.6% (2.5-9.9%)	6.7% (2.9-12%)	9.5 (4.2-17)	180 (83-330)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Bahamas	2016	Male	75 to 79	53% (34-72%)	47% (28-66%)	1.6 (0.7-3)	4.5% (1.9-8.5%)	5.2% (2.1-9.9%)	8.4 (3.3-16)	130 (50-240)
The Bahamas	2016	Male	80 to 84	47% (29-66%)	53% (34-71%)	1.3 (0.58-2.4)	3.7% (1.5-7%)	4.2% (1.7-8.1%)	5.6 (2.2-11)	62 (25-120)
The Bahamas	2016	Male	85 to 89	43% (26-59%)	57% (41-74%)	1.1 (0.48-2.2)	3.2% (1.2-6.3%)	3.5% (1.2-7.1%)	3 (1.1-6.1)	24 (8.9-47)
The Bahamas	2016	Male	90 to 94	43% (27-60%)	57% (40-73%)	1.1 (0.46-2.2)	2.8% (0.92-6.2%)	3.1% (0.89-7%)	1.2 (0.37-2.8)	6.9 (2.3-15)
The Bahamas	2016	Male	95 plus	46% (27-66%)	54% (34-73%)	1.2 (0.43-2.5)	3% (0.78-6.6%)	3.1% (0.57-7.5%)	0.35 (0.065-0.83)	1.3 (0.34-3)
Barbados	1990	Female	All Ages	32% (26-39%)	68% (61-74%)	1.2 (1-1.4)	2.2% (1.4-3.1%)	2.3% (0.97-4.1%)	27 (11-48)	790 (510-1100)
Barbados	1990	Female	15 to 19	37% (24-53%)	63% (47-76%)	1.3 (0.7-2.2)	1.9% (1.2-2.9%)	3.8% (2-6.3%)	0.24 (0.12-0.4)	28 (17-43)
Barbados	1990	Female	20 to 24	43% (28-59%)	57% (41-72%)	2 (1.1-3.1)	3.4% (2.3-4.7%)	5.6% (3.2-8.4%)	0.44 (0.25-0.67)	54 (36-76)
Barbados	1990	Female	25 to 29	43% (26-60%)	57% (40-74%)	2 (1.2-3)	3.4% (2.4-4.5%)	5.5% (3.5-7.7%)	0.5 (0.32-0.72)	59 (40-80)
Barbados	1990	Female	30 to 34	41% (25-59%)	59% (41-75%)	2 (1.1-2.9)	3.2% (2.3-4.3%)	5.6% (3.7-7.9%)	0.62 (0.4-0.89)	60 (41-81)
Barbados	1990	Female	35 to 39	40% (25-58%)	60% (42-75%)	1.8 (1-2.9)	3.5% (2.4-4.9%)	6.1% (4-8.7%)	0.91 (0.6-1.4)	65 (44-94)
Barbados	1990	Female	40 to 44	39% (25-57%)	61% (43-75%)	1.7 (0.91-2.8)	3.7% (2.3-5.5%)	6.5% (4-9.8%)	1.1 (0.7-1.7)	66 (41-99)
Barbados	1990	Female	45 to 49	37% (22-53%)	63% (47-78%)	1.7 (0.87-2.9)	3.8% (2.2-6.4%)	6.3% (3.6-11%)	1.1 (0.64-1.9)	55 (31-93)
Barbados	1990	Female	50 to 54	33% (20-50%)	67% (50-80%)	1.5 (0.77-2.6)	3.8% (1.9-6.7%)	5.8% (3-10%)	1.5 (0.77-2.7)	63 (31-110)
Barbados	1990	Female	55 to 59	29% (17-45%)	71% (55-83%)	1.2 (0.61-2.1)	3.1% (1.4-5.5%)	4.7% (2.3-8.1%)	1.5 (0.72-2.5)	54 (25-94)
Barbados	1990	Female	60 to 64	26% (15-41%)	74% (59-85%)	1.1 (0.52-1.8)	2.6% (0.86-5.1%)	3.6% (1.3-6.9%)	2.2 (0.79-4.2)	65 (22-130)
Barbados	1990	Female	65 to 69	24% (14-37%)	76% (63-86%)	0.92 (0.47-1.5)	2.1% (0.64-4.1%)	2.8% (0.87-5.4%)	2.2 (0.68-4.2)	53 (16-100)
Barbados	1990	Female	70 to 74	21% (13-34%)	79% (66-87%)	0.77 (0.37-1.4)	1.7% (0.23-3.9%)	2.2% (0.33-4.8%)	2.8 (0.42-6.3)	54 (7.2-120)
Barbados	1990	Female	75 to 79	19% (11-31%)	81% (69-89%)	0.67 (0.31-1.2)	1.3% (-0.058-3.3%)	1.6% (-0.086-4.1%)	2.8 (-0.15-6.9)	42 (-1.8-100)
Barbados	1990	Female	80 to 84	17% (9.7-28%)	83% (72-90%)	0.62 (0.3-1.1)	1.3% (0.11-3.1%)	1.5% (0.11-3.6%)	3 (0.2-7.2)	33 (2.9-76)
Barbados	1990	Female	85 to 89	17% (10-28%)	83% (72-90%)	0.64 (0.3-1.2)	1.3% (0.12-3.2%)	1.5% (0.086-3.6%)	2.8 (0.17-6.8)	22 (1.9-53)
Barbados	1990	Female	90 to 94	19% (11-31%)	81% (69-89%)	0.73 (0.32-1.4)	1.5% (0.036-3.9%)	1.7% (-0.045-4.5%)	1.9 (-0.05-5.1)	10 (0.24-26)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	1990	Female	95 plus	23% (12-39%)	77% (61-88%)	0.87 (0.32-1.8)	1.8% (-0.12-4.9%)	1.9% (-0.29-5.7%)	0.71 (-0.1-2.1)	2.6 (-0.17-7.1)
Barbados	1990	Male	All Ages	56% (47-65%)	44% (35-53%)	2.7 (2.4-3.1)	7.5% (6-9.1%)	7.6% (5.4-10%)	82 (57-110)	2800 (2200-3400)
Barbados	1990	Male	15 to 19	54% (37-71%)	46% (29-63%)	2 (1.1-3.2)	5.3% (3.5-7.7%)	8.2% (4.8-13%)	0.83 (0.48-1.3)	84 (55-120)
Barbados	1990	Male	20 to 24	65% (47-80%)	35% (20-53%)	3.8 (2.4-5.6)	11% (8.3-15%)	15% (9.9-22%)	2.4 (1.6-3.5)	220 (160-300)
Barbados	1990	Male	25 to 29	68% (49-82%)	32% (18-51%)	4.5 (2.9-6.4)	12% (9.1-16%)	17% (11-22%)	3.4 (2.3-4.7)	280 (200-360)
Barbados	1990	Male	30 to 34	69% (50-84%)	31% (16-50%)	4.7 (3.1-6.7)	12% (8.9-15%)	16% (12-21%)	3.9 (2.8-5.1)	280 (200-350)
Barbados	1990	Male	35 to 39	69% (52-83%)	31% (17-48%)	4.5 (2.8-6.7)	11% (8.6-15%)	16% (11-20%)	3.9 (2.8-5.1)	250 (180-320)
Barbados	1990	Male	40 to 44	68% (52-82%)	32% (18-48%)	4.3 (2.4-6.6)	11% (7.7-14%)	16% (11-20%)	3.4 (2.3-4.5)	190 (140-250)
Barbados	1990	Male	45 to 49	66% (45-81%)	34% (19-55%)	4 (2.1-6.6)	12% (8.1-15%)	15% (10-20%)	4.2 (2.7-5.5)	190 (130-250)
Barbados	1990	Male	50 to 54	64% (45-80%)	36% (20-55%)	3.6 (2-6.1)	11% (7.2-15%)	14% (9.2-19%)	4.5 (2.9-6.2)	180 (120-250)
Barbados	1990	Male	55 to 59	62% (44-79%)	38% (21-56%)	3.3 (1.8-5.5)	11% (6.6-15%)	13% (8.2-18%)	5.9 (3.6-8.2)	200 (120-280)
Barbados	1990	Male	60 to 64	60% (39-78%)	40% (22-61%)	3 (1.6-5.2)	9.1% (5.2-13%)	11% (6.2-15%)	7.3 (4.1-10)	210 (120-300)
Barbados	1990	Male	65 to 69	59% (39-76%)	41% (24-61%)	2.6 (1.5-4.2)	8.4% (4.8-12%)	9.8% (5.6-14%)	9.1 (5.3-13)	220 (130-320)
Barbados	1990	Male	70 to 74	55% (37-72%)	45% (28-63%)	2.2 (1.1-3.8)	6.3% (3.1-10%)	7.2% (3.5-12%)	9.8 (4.7-16)	190 (92-300)
Barbados	1990	Male	75 to 79	50% (31-69%)	50% (31-69%)	1.8 (0.89-3.1)	4.9% (2.1-8.1%)	5.5% (2.2-9.2%)	9.2 (3.8-16)	140 (58-230)
Barbados	1990	Male	80 to 84	44% (27-63%)	56% (37-73%)	1.5 (0.78-2.5)	4.1% (1.8-7.1%)	4.5% (1.9-7.9%)	7.3 (3-13)	79 (34-140)
Barbados	1990	Male	85 to 89	40% (24-57%)	60% (43-76%)	1.3 (0.66-2.4)	3.4% (1.3-6.4%)	3.7% (1.4-7.2%)	4.4 (1.7-8.5)	34 (14-65)
Barbados	1990	Male	90 to 94	40% (25-57%)	60% (43-75%)	1.3 (0.64-2.5)	3% (0.94-6.2%)	3.2% (0.9-7%)	1.5 (0.42-3.2)	8.2 (2.6-17)
Barbados	1990	Male	95 plus	43% (24-64%)	57% (36-76%)	1.5 (0.59-2.9)	3.1% (0.64-6.8%)	3.3% (0.4-7.6%)	0.31 (0.038-0.71)	1.2 (0.23-2.5)
Barbados	1995	Female	All Ages	31% (26-38%)	69% (62-74%)	1.1 (0.94-1.3)	1.9% (1.3-2.8%)	1.9% (0.78-3.5%)	23 (9.2-44)	720 (470-1000)
Barbados	1995	Female	15 to 19	36% (24-53%)	64% (47-76%)	1.2 (0.62-1.9)	1.8% (1.1-2.8%)	3.5% (1.7-6%)	0.18 (0.087-0.31)	21 (12-34)
Barbados	1995	Female	20 to 24	42% (27-59%)	58% (41-73%)	1.8 (0.96-2.9)	3.2% (2-4.4%)	5.1% (2.8-7.8%)	0.42 (0.23-0.66)	51 (32-74)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	1995	Female	25 to 29	42% (26-59%)	58% (41-74%)	1.8 (1.1-2.7)	3.2% (2.2-4.3%)	4.9% (3-7.1%)	0.43 (0.26-0.64)	51 (34-70)
Barbados	1995	Female	30 to 34	41% (25-57%)	59% (43-75%)	1.8 (1-2.6)	3% (2.1-4.1%)	5% (3.3-7%)	0.6 (0.38-0.86)	58 (39-79)
Barbados	1995	Female	35 to 39	40% (25-57%)	60% (43-75%)	1.6 (0.93-2.5)	3.2% (2.2-4.5%)	5.4% (3.6-7.8%)	1.1 (0.68-1.6)	74 (50-110)
Barbados	1995	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.5 (0.8-2.4)	3.2% (2-4.9%)	5.7% (3.5-8.4%)	1.2 (0.72-1.8)	71 (43-110)
Barbados	1995	Female	45 to 49	36% (22-53%)	64% (47-78%)	1.5 (0.78-2.5)	3.2% (1.8-5.3%)	5.4% (3.1-8.8%)	1.2 (0.71-2)	62 (35-100)
Barbados	1995	Female	50 to 54	33% (20-50%)	67% (50-80%)	1.3 (0.68-2.2)	3.2% (1.5-5.7%)	5% (2.5-8.5%)	1.4 (0.67-2.5)	57 (27-100)
Barbados	1995	Female	55 to 59	29% (17-45%)	71% (55-83%)	1.1 (0.55-1.9)	2.6% (1.1-4.7%)	4% (1.8-7%)	1.4 (0.62-2.4)	49 (21-86)
Barbados	1995	Female	60 to 64	26% (15-41%)	74% (59-85%)	0.95 (0.47-1.6)	2.2% (0.63-4.4%)	3% (0.97-5.9%)	1.7 (0.53-3.3)	50 (14-100)
Barbados	1995	Female	65 to 69	24% (14-37%)	76% (63-86%)	0.82 (0.42-1.4)	1.7% (0.41-3.6%)	2.3% (0.63-4.6%)	1.9 (0.52-3.8)	47 (11-95)
Barbados	1995	Female	70 to 74	21% (13-34%)	79% (66-87%)	0.69 (0.33-1.2)	1.4% (0.041-3.3%)	1.8% (0.11-4.3%)	2.1 (0.12-4.8)	39 (1.1-93)
Barbados	1995	Female	75 to 79	19% (11-31%)	81% (69-89%)	0.6 (0.28-1.1)	1% (-0.24-2.7%)	1.3% (-0.28-3.4%)	2.1 (-0.46-5.5)	31 (-7.3-83)
Barbados	1995	Female	80 to 84	17% (9.6-28%)	83% (72-90%)	0.56 (0.27-0.98)	1% (-0.053-2.6%)	1.2% (-0.072-3.1%)	2.6 (-0.15-6.5)	29 (-1.4-72)
Barbados	1995	Female	85 to 89	17% (9.9-27%)	83% (73-90%)	0.57 (0.27-1)	1.1% (-0.017-2.6%)	1.2% (-0.073-3%)	2.5 (-0.15-6.3)	19 (-0.31-48)
Barbados	1995	Female	90 to 94	19% (11-30%)	81% (70-89%)	0.66 (0.29-1.3)	1.2% (-0.055-3.4%)	1.3% (-0.14-4%)	1.8 (-0.19-5.2)	9.6 (-0.43-27)
Barbados	1995	Female	95 plus	23% (12-38%)	77% (62-88%)	0.79 (0.27-1.6)	1.5% (-0.24-4.4%)	1.6% (-0.42-5%)	0.82 (-0.22-2.7)	3 (-0.49-9)
Barbados	1995	Male	All Ages	56% (48-64%)	44% (36-52%)	2.5 (2.2-2.9)	6.8% (5.4-8.3%)	6.5% (4.5-8.9%)	78 (52-110)	2600 (2100-3300)
Barbados	1995	Male	15 to 19	54% (37-71%)	46% (29-63%)	1.8 (0.93-2.9)	5.1% (3.4-7.5%)	7.5% (4.4-12%)	0.73 (0.41-1.1)	71 (46-100)
Barbados	1995	Male	20 to 24	65% (46-80%)	35% (20-54%)	3.3 (2-5)	10% (7.2-14%)	13% (8.5-19%)	2.1 (1.3-3)	190 (130-260)
Barbados	1995	Male	25 to 29	68% (49-82%)	32% (18-51%)	4 (2.5-5.7)	11% (7.7-14%)	14% (8.9-19%)	2.8 (1.8-3.9)	230 (160-300)
Barbados	1995	Male	30 to 34	69% (49-83%)	31% (17-51%)	4.2 (2.7-5.8)	9.6% (7.2-12%)	12% (8.3-16%)	3.5 (2.4-4.7)	250 (180-330)
Barbados	1995	Male	35 to 39	69% (52-83%)	31% (17-48%)	4 (2.5-5.9)	9.2% (6.7-12%)	12% (8.4-16%)	3.6 (2.4-4.8)	230 (170-300)
Barbados	1995	Male	40 to 44	68% (52-82%)	32% (18-48%)	3.8 (2.2-5.7)	9.3% (6.5-12%)	13% (8.4-17%)	3.8 (2.4-5.1)	220 (150-280)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	1995	Male	45 to 49	66% (45-81%)	34% (19-55%)	3.5 (2-5.7)	10% (6.7-13%)	13% (8.4-17%)	5.3 (3.4-7.3)	250 (170-340)
Barbados	1995	Male	50 to 54	64% (45-80%)	36% (20-55%)	3.2 (1.7-5.1)	9.8% (6.3-13%)	12% (7.6-17%)	4.5 (2.8-6.3)	190 (120-250)
Barbados	1995	Male	55 to 59	62% (43-78%)	38% (22-57%)	2.9 (1.6-4.7)	9.5% (5.7-13%)	12% (7-16%)	6.2 (3.6-8.8)	210 (130-300)
Barbados	1995	Male	60 to 64	60% (39-78%)	40% (22-61%)	2.7 (1.4-4.5)	8% (4.4-12%)	9.7% (5.4-14%)	6.6 (3.6-9.8)	190 (110-290)
Barbados	1995	Male	65 to 69	58% (39-76%)	42% (24-61%)	2.3 (1.3-3.8)	7.2% (4-11%)	8.5% (4.7-13%)	8.7 (4.8-13)	210 (120-310)
Barbados	1995	Male	70 to 74	55% (36-71%)	45% (29-64%)	1.9 (1-3.4)	5.4% (2.5-9%)	6.2% (2.8-10%)	8.2 (3.7-14)	160 (70-260)
Barbados	1995	Male	75 to 79	49% (30-69%)	51% (31-70%)	1.6 (0.8-2.7)	4.3% (1.7-7.4%)	4.8% (1.9-8.4%)	8.1 (3-14)	120 (47-210)
Barbados	1995	Male	80 to 84	43% (26-62%)	57% (38-74%)	1.3 (0.69-2.3)	3.4% (1.4-6.1%)	3.8% (1.5-6.9%)	7 (2.8-13)	77 (32-140)
Barbados	1995	Male	85 to 89	39% (23-56%)	61% (44-77%)	1.2 (0.58-2.1)	2.8% (1-5.5%)	3% (1.1-6.1%)	4.3 (1.5-8.6)	34 (13-66)
Barbados	1995	Male	90 to 94	39% (24-57%)	61% (43-76%)	1.2 (0.56-2.2)	2.4% (0.68-5.4%)	2.6% (0.61-6%)	1.8 (0.44-4.2)	10 (2.8-22)
Barbados	1995	Male	95 plus	43% (24-63%)	57% (37-76%)	1.3 (0.52-2.5)	2.5% (0.4-5.9%)	2.7% (0.21-6.4%)	0.48 (0.038-1.2)	1.8 (0.28-4.2)
Barbados	2000	Female	All Ages	30% (25-37%)	70% (63-75%)	1 (0.88-1.2)	1.8% (1.2-2.5%)	1.8% (0.82-3.2%)	22 (9.3-38)	690 (460-960)
Barbados	2000	Female	15 to 19	36% (23-52%)	64% (48-77%)	1.1 (0.58-1.9)	1.6% (0.96-2.4%)	1.6% (1.6-5.4%)	0.12 (0.061-0.22)	17 (10-27)
Barbados	2000	Female	20 to 24	42% (26-58%)	58% (42-74%)	1.7 (0.9-2.7)	3.1% (2-4.4%)	4.6% (2.5-7.4%)	0.35 (0.19-0.56)	42 (27-61)
Barbados	2000	Female	25 to 29	41% (25-58%)	59% (42-75%)	1.7 (0.95-2.7)	3.2% (2.2-4.4%)	4.6% (2.7-6.7%)	0.49 (0.28-0.77)	53 (35-75)
Barbados	2000	Female	30 to 34	39% (24-56%)	61% (44-76%)	1.7 (0.95-2.6)	2.9% (2-3.9%)	4.3% (2.8-6.2%)	0.55 (0.34-0.82)	52 (35-73)
Barbados	2000	Female	35 to 39	39% (24-56%)	61% (44-76%)	1.5 (0.89-2.4)	2.9% (2-4%)	4.7% (3.1-6.8%)	0.92 (0.58-1.4)	67 (45-97)
Barbados	2000	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.5 (0.78-2.3)	2.9% (1.8-4.4%)	5% (3.1-7.4%)	1.2 (0.77-1.9)	74 (46-110)
Barbados	2000	Female	45 to 49	36% (21-53%)	64% (47-79%)	1.4 (0.75-2.3)	2.9% (1.7-4.8%)	4.9% (3-7.8%)	1.4 (0.82-2.3)	71 (41-110)
Barbados	2000	Female	50 to 54	32% (19-48%)	68% (52-81%)	1.2 (0.68-2)	2.8% (1.4-4.8%)	4.5% (2.4-7.5%)	1.5 (0.77-2.5)	63 (32-110)
Barbados	2000	Female	55 to 59	28% (16-44%)	72% (56-84%)	1 (0.54-1.8)	2.5% (1.1-4.5%)	3.8% (1.9-6.5%)	1.4 (0.69-2.4)	49 (23-87)
Barbados	2000	Female	60 to 64	25% (14-40%)	75% (60-86%)	0.89 (0.44-1.6)	2% (0.64-4%)	2.9% (1.1-5.4%)	1.5 (0.56-3)	46 (14-94)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	2000	Female	65 to 69	23% (14-36%)	77% (64-86%)	0.77 (0.4-1.3)	1.5% (0.37-3.1%)	2.2% (0.66-4.2%)	1.6 (0.46-3)	38 (9.7-76)
Barbados	2000	Female	70 to 74	20% (12-33%)	80% (67-88%)	0.65 (0.32-1.2)	1.2% (0.041-3%)	1.7% (0.17-3.9%)	2 (0.2-4.8)	38 (1.3-96)
Barbados	2000	Female	75 to 79	18% (10-30%)	82% (70-90%)	0.56 (0.26-1)	0.94% (-0.16-2.5%)	1.2% (-0.15-3.1%)	1.8 (-0.22-4.6)	26 (-4.7-70)
Barbados	2000	Female	80 to 84	17% (9.3-28%)	83% (72-91%)	0.53 (0.25-0.92)	0.94% (0.025-2.3%)	1.2% (0.047-2.9%)	2 (0.078-5)	22 (0.6-55)
Barbados	2000	Female	85 to 89	17% (9.5-26%)	83% (74-90%)	0.54 (0.26-0.99)	0.95% (0.084-2.4%)	1.1% (0.067-2.7%)	2.3 (0.13-5.7)	18 (1.5-45)
Barbados	2000	Female	90 to 94	19% (11-29%)	81% (71-89%)	0.62 (0.27-1.2)	1.1% (-0.0093-3.1%)	1.2% (-0.086-3.6%)	1.7 (-0.12-5)	9.3 (-0.078-27)
Barbados	2000	Female	95 plus	22% (11-37%)	78% (63-89%)	0.75 (0.26-1.5)	1.3% (-0.16-4%)	1.4% (-0.33-4.7%)	0.83 (-0.19-2.7)	3.1 (-0.37-9.4)
Barbados	2000	Male	All Ages	55% (47-63%)	45% (37-53%)	2.4 (2.1-2.7)	6.3% (5.1-7.7%)	6.1% (4.2-8.1%)	71 (50-97)	2500 (2000-3100)
Barbados	2000	Male	15 to 19	53% (35-70%)	47% (30-65%)	1.7 (0.86-2.7)	5% (3.2-7.3%)	7.1% (4-11%)	0.74 (0.42-1.2)	72 (46-110)
Barbados	2000	Male	20 to 24	64% (46-79%)	36% (21-54%)	3.2 (1.8-4.8)	10% (7.3-14%)	13% (8.4-19%)	1.9 (1.1-2.8)	170 (120-230)
Barbados	2000	Male	25 to 29	67% (49-82%)	33% (18-51%)	3.9 (2.3-5.7)	11% (7.6-15%)	14% (8.7-19%)	2.9 (1.8-4.2)	230 (160-310)
Barbados	2000	Male	30 to 34	68% (49-83%)	32% (17-51%)	4 (2.5-5.8)	8.4% (6.3-11%)	10% (7-14%)	2.3 (1.6-3.3)	180 (130-240)
Barbados	2000	Male	35 to 39	68% (51-82%)	32% (18-49%)	3.8 (2.3-5.6)	7.9% (5.9-10%)	9.8% (6.7-13%)	2.9 (1.9-3.9)	200 (140-260)
Barbados	2000	Male	40 to 44	67% (50-82%)	33% (18-50%)	3.6 (2.1-5.3)	8.3% (5.9-11%)	11% (7.4-15%)	3.7 (2.4-5)	220 (150-290)
Barbados	2000	Male	45 to 49	65% (44-80%)	35% (20-56%)	3.4 (1.9-5.2)	9.1% (6.3-12%)	12% (7.9-15%)	5.4 (3.6-7.5)	260 (180-350)
Barbados	2000	Male	50 to 54	63% (43-79%)	37% (21-57%)	3 (1.7-4.8)	9.1% (5.9-12%)	12% (7.2-16%)	6.2 (3.8-8.7)	260 (160-350)
Barbados	2000	Male	55 to 59	61% (43-77%)	39% (23-57%)	2.8 (1.5-4.4)	8.6% (5.2-12%)	11% (6.5-15%)	5.5 (3.4-7.8)	200 (120-270)
Barbados	2000	Male	60 to 64	59% (39-77%)	41% (23-61%)	2.5 (1.3-4.2)	7.2% (4.1-11%)	8.8% (4.9-13%)	6.2 (3.4-9.3)	180 (100-270)
Barbados	2000	Male	65 to 69	57% (38-75%)	43% (25-62%)	2.2 (1.2-3.6)	6.4% (3.5-9.8%)	7.7% (4.2-12%)	7.3 (4-11)	180 (100-270)
Barbados	2000	Male	70 to 74	53% (35-71%)	47% (29-65%)	1.8 (0.97-3.1)	4.8% (2.3-8.2%)	5.6% (2.7-9.6%)	7.5 (3.6-13)	150 (72-250)
Barbados	2000	Male	75 to 79	48% (29-67%)	52% (33-71%)	1.5 (0.74-2.6)	3.8% (1.6-6.9%)	4.4% (1.8-7.7%)	6.5 (2.7-12)	99 (42-170)
Barbados	2000	Male	80 to 84	42% (25-61%)	58% (39-75%)	1.2 (0.64-2.1)	3.1% (1.4-5.3%)	3.4% (1.5-6%)	5.8 (2.5-9.8)	64 (28-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	2000	Male	85 to 89	38% (22-55%)	62% (45-78%)	1.1 (0.54-1.9)	2.6% (1.1-5%)	2.8% (1.1-5.6%)	4 (1.6-7.9)	32 (13-61)
Barbados	2000	Male	90 to 94	38% (24-55%)	62% (45-76%)	1.1 (0.52-2.1)	2.3% (0.77-4.9%)	2.4% (0.74-5.5%)	1.9 (0.58-4.2)	10 (3.6-23)
Barbados	2000	Male	95 plus	42% (23-62%)	58% (38-77%)	1.2 (0.48-2.4)	2.4% (0.57-5.3%)	2.5% (0.42-6%)	0.62 (0.1-1.5)	2.4 (0.57-5.3)
Barbados	2005	Female	All Ages	30% (25-37%)	70% (63-75%)	1.1 (0.92-1.2)	2% (1.4-2.7%)	2.1% (1-3.3%)	24 (12-40)	760 (540-1000)
Barbados	2005	Female	15 to 19	36% (23-52%)	64% (48-77%)	1.2 (0.59-2)	1.7% (1-2.6%)	3.7% (1.8-6.3%)	0.14 (0.066-0.24)	18 (11-28)
Barbados	2005	Female	20 to 24	42% (26-58%)	58% (42-74%)	1.7 (0.92-2.9)	3.3% (2.1-4.7%)	5.2% (2.8-8.1%)	0.4 (0.21-0.65)	46 (29-66)
Barbados	2005	Female	25 to 29	41% (25-58%)	59% (42-75%)	1.8 (1-2.8)	3.3% (2.2-4.6%)	4.9% (2.9-7.2%)	0.45 (0.27-0.68)	48 (32-68)
Barbados	2005	Female	30 to 34	39% (24-56%)	61% (44-76%)	1.7 (0.95-2.6)	3.2% (2.2-4.4%)	5.2% (3.3-7.5%)	0.66 (0.41-0.98)	58 (40-81)
Barbados	2005	Female	35 to 39	39% (24-56%)	61% (44-76%)	1.6 (0.92-2.5)	3.1% (2.1-4.3%)	5.3% (3.6-7.5%)	0.87 (0.56-1.3)	63 (42-89)
Barbados	2005	Female	40 to 44	37% (23-56%)	63% (44-77%)	1.5 (0.82-2.4)	3.2% (2-4.7%)	5.6% (3.7-8.1%)	1.4 (0.87-2)	80 (52-120)
Barbados	2005	Female	45 to 49	35% (21-52%)	65% (48-79%)	1.4 (0.78-2.3)	3.1% (1.9-4.9%)	5.4% (3.3-8.5%)	1.8 (1.1-2.8)	89 (55-140)
Barbados	2005	Female	50 to 54	32% (19-48%)	68% (52-81%)	1.3 (0.7-2.1)	2.9% (1.5-4.8%)	4.8% (2.7-7.8%)	1.7 (0.98-2.8)	75 (41-130)
Barbados	2005	Female	55 to 59	28% (17-44%)	72% (56-83%)	1.1 (0.56-1.8)	2.6% (1.4-4.5%)	4.1% (2.3-6.9%)	1.9 (1-3.1)	67 (35-110)
Barbados	2005	Female	60 to 64	25% (15-39%)	75% (61-85%)	0.92 (0.45-1.6)	2.1% (0.8-3.9%)	3.2% (1.4-5.7%)	1.7 (0.72-2.9)	50 (20-92)
Barbados	2005	Female	65 to 69	23% (14-36%)	77% (64-86%)	0.8 (0.41-1.3)	1.7% (0.56-3.1%)	2.5% (0.94-4.3%)	1.6 (0.63-2.9)	40 (13-75)
Barbados	2005	Female	70 to 74	20% (13-32%)	80% (68-87%)	0.67 (0.34-1.2)	1.3% (0.22-3%)	1.9% (0.45-4.1%)	1.9 (0.43-4)	36 (5.9-81)
Barbados	2005	Female	75 to 79	18% (11-29%)	82% (71-89%)	0.58 (0.28-1)	1.1% (0.006-2.6%)	1.4% (0.079-3.4%)	2.1 (0.11-5)	31 (0.17-77)
Barbados	2005	Female	80 to 84	17% (9.4-27%)	83% (73-91%)	0.55 (0.26-0.94)	1% (0.13-2.4%)	1.3% (0.18-3%)	2.3 (0.32-5.2)	25 (3.4-60)
Barbados	2005	Female	85 to 89	17% (9.5-26%)	83% (74-91%)	0.56 (0.26-0.99)	1% (0.11-2.4%)	1.2% (0.11-2.8%)	2.3 (0.22-5.4)	18 (1.9-43)
Barbados	2005	Female	90 to 94	19% (11-30%)	81% (70-89%)	0.63 (0.27-1.2)	1.2% (0.017-3.2%)	1.3% (-0.027-3.6%)	2.1 (-0.041-5.6)	11 (0.16-30)
Barbados	2005	Female	95 plus	22% (11-38%)	78% (62-89%)	0.76 (0.27-1.6)	1.5% (-0.073-4.2%)	1.6% (-0.2-4.7%)	1 (-0.13-3)	3.7 (-0.18-11)
Barbados	2005	Male	All Ages	56% (47-64%)	44% (36-53%)	2.5 (2.2-2.8)	6.9% (5.7-8.2%)	6.7% (4.9-8.8%)	77 (56-100)	2700 (2200-3300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	2005	Male	15 to 19	53% (35-70%)	47% (30-65%)	1.7 (0.89-2.9)	5.3% (3.4-7.7%)	7.8% (4.5-12%)	0.7 (0.39-1.1)	68 (44-100)
Barbados	2005	Male	20 to 24	64% (45-80%)	36% (20-55%)	3.3 (1.9-5.1)	11% (7.6-15%)	14% (8.9-21%)	1.8 (1.1-2.7)	170 (110-230)
Barbados	2005	Male	25 to 29	67% (48-82%)	33% (18-52%)	4 (2.4-5.9)	12% (8.3-16%)	15% (9.8-22%)	2.7 (1.7-3.8)	210 (150-290)
Barbados	2005	Male	30 to 34	68% (49-83%)	32% (17-51%)	4.2 (2.6-6.1)	11% (7.8-14%)	14% (9.9-19%)	2.8 (1.9-3.7)	210 (150-270)
Barbados	2005	Male	35 to 39	68% (51-82%)	32% (18-49%)	3.9 (2.4-5.9)	9.7% (7.2-13%)	13% (9.5-18%)	3.2 (2.2-4.3)	210 (150-270)
Barbados	2005	Male	40 to 44	67% (50-81%)	33% (19-50%)	3.7 (2.2-5.5)	9.4% (6.8-12%)	13% (9.4-17%)	4 (2.7-5.3)	230 (170-300)
Barbados	2005	Male	45 to 49	65% (44-80%)	35% (20-56%)	3.4 (2-5.3)	10% (7.2-13%)	14% (9.6-18%)	6.5 (4.4-8.7)	310 (220-410)
Barbados	2005	Male	50 to 54	63% (43-79%)	37% (21-57%)	3.1 (1.8-4.8)	9.8% (6.5-13%)	13% (8.6-17%)	7.2 (4.7-9.7)	300 (200-400)
Barbados	2005	Male	55 to 59	61% (42-78%)	39% (22-58%)	2.9 (1.6-4.5)	8.9% (5.6-12%)	12% (7.2-16%)	7.5 (4.6-10)	270 (170-370)
Barbados	2005	Male	60 to 64	59% (39-77%)	41% (23-61%)	2.6 (1.4-4.3)	7.7% (4.3-11%)	9.6% (5.4-14%)	6.8 (3.8-9.8)	200 (110-290)
Barbados	2005	Male	65 to 69	57% (37-75%)	43% (25-63%)	2.3 (1.2-3.6)	6.7% (3.9-10%)	8.2% (4.8-12%)	7.2 (4.2-11)	180 (100-270)
Barbados	2005	Male	70 to 74	53% (35-70%)	47% (30-65%)	1.9 (0.98-3.1)	4.9% (2.5-8.2%)	5.9% (3-9.8%)	6.9 (3.4-12)	140 (70-230)
Barbados	2005	Male	75 to 79	48% (30-67%)	52% (33-70%)	1.5 (0.77-2.7)	3.9% (1.7-6.9%)	4.5% (2-7.9%)	7 (3-12)	110 (47-190)
Barbados	2005	Male	80 to 84	42% (25-61%)	58% (39-75%)	1.3 (0.67-2.1)	3.2% (1.5-5.5%)	3.6% (1.6-6.3%)	5.9 (2.7-10)	66 (30-120)
Barbados	2005	Male	85 to 89	38% (22-55%)	62% (45-78%)	1.1 (0.56-2)	2.7% (1.2-4.9%)	3% (1.3-5.6%)	4 (1.7-7.5)	32 (14-58)
Barbados	2005	Male	90 to 94	38% (23-55%)	62% (45-77%)	1.1 (0.54-2.1)	2.4% (0.86-5%)	2.6% (0.81-5.6%)	2.1 (0.63-4.4)	11 (4-24)
Barbados	2005	Male	95 plus	41% (23-62%)	59% (38-77%)	1.2 (0.49-2.5)	2.6% (0.71-5.6%)	2.7% (0.59-6.3%)	0.71 (0.15-1.6)	2.7 (0.73-5.9)
Barbados	2010	Female	All Ages	30% (25-36%)	70% (64-75%)	1.1 (0.97-1.3)	2.2% (1.6-3%)	2.4% (1.4-3.9%)	28 (16-45)	860 (620-1200)
Barbados	2010	Female	15 to 19	36% (23-52%)	64% (48-77%)	1.3 (0.64-2.2)	1.9% (1.1-2.9%)	3.9% (1.9-7%)	0.17 (0.082-0.3)	20 (12-32)
Barbados	2010	Female	20 to 24	41% (26-58%)	59% (42-74%)	1.9 (1-3.1)	3.5% (2.3-4.9%)	5.6% (3-8.9%)	0.38 (0.2-0.62)	43 (28-63)
Barbados	2010	Female	25 to 29	41% (25-58%)	59% (42-75%)	1.9 (1-3)	3.4% (2.3-4.8%)	5.3% (3.1-8%)	0.46 (0.26-0.71)	49 (32-70)
Barbados	2010	Female	30 to 34	39% (24-57%)	61% (43-76%)	1.8 (1-2.9)	3.4% (2.2-4.7%)	5.5% (3.4-8%)	0.61 (0.38-0.91)	54 (36-77)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	2010	Female	35 to 39	39% (24-56%)	61% (44-76%)	1.7 (0.95-2.8)	3.4% (2.3-4.8%)	5.8% (3.8-8.4%)	1 (0.67-1.6)	73 (50-100)
Barbados	2010	Female	40 to 44	37% (24-55%)	63% (45-76%)	1.6 (0.87-2.6)	3.3% (2.1-4.9%)	5.9% (3.7-8.7%)	1.3 (0.84-2)	78 (50-120)
Barbados	2010	Female	45 to 49	35% (21-53%)	65% (47-79%)	1.5 (0.84-2.5)	3.3% (2-5.1%)	5.6% (3.4-8.9%)	1.9 (1.2-3)	97 (60-150)
Barbados	2010	Female	50 to 54	32% (19-49%)	68% (51-81%)	1.4 (0.77-2.2)	3.3% (1.8-5.3%)	5.4% (3.2-8.4%)	2.5 (1.5-4)	100 (59-170)
Barbados	2010	Female	55 to 59	28% (16-44%)	72% (56-84%)	1.2 (0.6-1.9)	2.8% (1.5-4.8%)	4.5% (2.6-7.3%)	2.3 (1.3-3.8)	84 (46-140)
Barbados	2010	Female	60 to 64	25% (14-39%)	75% (61-86%)	0.99 (0.5-1.7)	2.2% (0.92-4%)	3.4% (1.6-5.9%)	1.9 (0.89-3.4)	60 (25-110)
Barbados	2010	Female	65 to 69	23% (14-36%)	77% (64-86%)	0.86 (0.44-1.4)	1.9% (0.77-3.5%)	2.8% (1.3-4.8%)	2 (0.91-3.5)	49 (20-91)
Barbados	2010	Female	70 to 74	20% (13-33%)	80% (67-87%)	0.72 (0.36-1.3)	1.5% (0.37-3.2%)	2.2% (0.69-4.4%)	1.9 (0.59-3.8)	37 (9.6-80)
Barbados	2010	Female	75 to 79	18% (11-30%)	82% (70-89%)	0.63 (0.3-1.1)	1.3% (0.19-3%)	1.7% (0.32-3.9%)	2.4 (0.45-5.4)	36 (5.6-84)
Barbados	2010	Female	80 to 84	17% (9.3-27%)	83% (73-91%)	0.59 (0.29-1)	1.2% (0.21-2.6%)	1.5% (0.3-3.3%)	2.7 (0.52-5.8)	30 (5.8-67)
Barbados	2010	Female	85 to 89	17% (9.5-26%)	83% (74-91%)	0.59 (0.27-1.1)	1.2% (0.23-2.8%)	1.5% (0.25-3.4%)	2.9 (0.51-6.7)	23 (4.3-52)
Barbados	2010	Female	90 to 94	18% (11-29%)	82% (71-89%)	0.67 (0.3-1.2)	1.5% (0.18-3.6%)	1.7% (0.14-4.2%)	2.6 (0.23-6.6)	14 (1.7-34)
Barbados	2010	Female	95 plus	22% (11-38%)	78% (62-89%)	0.81 (0.29-1.7)	1.8% (0.074-4.6%)	1.9% (-0.058-5.4%)	1.4 (-0.042-3.9)	5 (0.21-13)
Barbados	2010	Male	All Ages	55% (47-63%)	45% (37-53%)	2.7 (2.3-3)	7.3% (6-8.6%)	7.3% (5.4-9.5%)	86 (63-110)	2900 (2400-3600)
Barbados	2010	Male	15 to 19	53% (35-70%)	47% (30-65%)	1.9 (0.93-3)	5.1% (3.2-7.5%)	8% (4.4-13%)	0.6 (0.33-0.95)	61 (39-89)
Barbados	2010	Male	20 to 24	64% (46-79%)	36% (21-54%)	3.5 (2-5.5)	11% (7.7-15%)	15% (9.1-21%)	1.6 (0.94-2.3)	150 (100-200)
Barbados	2010	Male	25 to 29	67% (48-81%)	33% (19-52%)	4.3 (2.5-6.3)	12% (8.5-16%)	16% (10-22%)	2.5 (1.6-3.5)	200 (140-270)
Barbados	2010	Male	30 to 34	68% (48-83%)	32% (17-52%)	4.5 (2.7-6.6)	11% (8.3-15%)	16% (11-21%)	2.8 (1.9-3.8)	200 (150-260)
Barbados	2010	Male	35 to 39	68% (51-82%)	32% (18-49%)	4.3 (2.5-6.2)	10% (7.5-13%)	15% (10-19%)	3.3 (2.3-4.4)	220 (160-280)
Barbados	2010	Male	40 to 44	67% (50-81%)	33% (19-50%)	4 (2.4-5.9)	10% (7.2-13%)	15% (10-19%)	3.9 (2.7-5)	220 (160-290)
Barbados	2010	Male	45 to 49	65% (44-80%)	35% (20-56%)	3.7 (2.1-5.6)	11% (7.6-14%)	14% (10-18%)	6.8 (4.7-8.9)	330 (230-420)
Barbados	2010	Male	50 to 54	63% (43-80%)	37% (20-57%)	3.4 (2-5.2)	10% (6.9-13%)	13% (9.1-17%)	8.2 (5.4-11)	340 (240-450)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	2010	Male	55 to 59	61% (42-78%)	39% (22-58%)	3.1 (1.7-4.8)	9.5% (6.2-13%)	12% (8.2-17%)	9 (5.9-12)	320 (220-440)
Barbados	2010	Male	60 to 64	59% (38-77%)	41% (23-62%)	2.8 (1.5-4.5)	8.2% (4.8-11%)	10% (6-14%)	9.5 (5.6-14)	290 (170-400)
Barbados	2010	Male	65 to 69	57% (37-75%)	43% (25-63%)	2.5 (1.4-3.9)	7.3% (4.3-10%)	9% (5.4-13%)	8.5 (5-12)	210 (130-310)
Barbados	2010	Male	70 to 74	53% (35-70%)	47% (30-65%)	2 (1.1-3.5)	5.5% (2.8-8.7%)	6.6% (3.3-11%)	8.1 (4.1-13)	160 (83-260)
Barbados	2010	Male	75 to 79	48% (29-67%)	52% (33-71%)	1.7 (0.81-2.9)	4.2% (1.9-7.2%)	5% (2.2-8.5%)	7.4 (3.2-13)	110 (49-190)
Barbados	2010	Male	80 to 84	42% (25-61%)	58% (39-75%)	1.4 (0.7-2.3)	3.5% (1.6-5.9%)	4% (1.8-6.7%)	6.5 (2.9-11)	73 (34-120)
Barbados	2010	Male	85 to 89	38% (22-55%)	62% (45-78%)	1.2 (0.62-2.1)	2.9% (1.3-5.2%)	3.3% (1.4-5.9%)	4.5 (1.9-8.1)	36 (16-64)
Barbados	2010	Male	90 to 94	38% (23-56%)	62% (44-77%)	1.2 (0.59-2.2)	2.7% (1-5.3%)	3% (1-6%)	2.4 (0.83-4.9)	13 (5-26)
Barbados	2010	Male	95 plus	41% (23-61%)	59% (39-77%)	1.3 (0.53-2.6)	2.9% (0.86-6.2%)	3.1% (0.7-7.1%)	0.84 (0.19-1.9)	3.2 (0.92-6.6)
Barbados	2016	Female	All Ages	30% (24-36%)	70% (64-76%)	1.2 (0.88-1.6)	2.3% (1.5-3.5%)	2.6% (1.2-4.3%)	35 (17-61)	980 (600-1500)
Barbados	2016	Female	15 to 19	36% (23-52%)	64% (48-77%)	1.3 (0.64-2.3)	1.9% (1.1-3.1%)	4.1% (1.9-7.2%)	0.16 (0.072-0.29)	20 (11-32)
Barbados	2016	Female	20 to 24	41% (26-58%)	59% (42-74%)	2 (0.96-3.5)	3.5% (2.2-5.1%)	5.7% (3-9.4%)	0.36 (0.18-0.62)	43 (26-63)
Barbados	2016	Female	25 to 29	41% (25-58%)	59% (42-75%)	2 (1-3.5)	3.4% (2.3-4.9%)	5.5% (3-8.5%)	0.41 (0.22-0.67)	45 (29-65)
Barbados	2016	Female	30 to 34	39% (24-57%)	61% (43-76%)	2 (1-3.3)	3.4% (2.2-4.9%)	5.7% (3.4-8.8%)	0.61 (0.36-0.98)	55 (35-80)
Barbados	2016	Female	35 to 39	39% (24-56%)	61% (44-76%)	1.8 (0.94-3.1)	3.5% (2.3-5.1%)	6.1% (3.7-9.2%)	0.93 (0.56-1.4)	66 (42-97)
Barbados	2016	Female	40 to 44	37% (24-56%)	63% (44-76%)	1.7 (0.89-2.8)	3.5% (2.1-5.3%)	6.2% (3.8-9.4%)	1.3 (0.77-2)	79 (49-120)
Barbados	2016	Female	45 to 49	35% (21-52%)	65% (48-79%)	1.6 (0.86-2.8)	3.5% (2-5.7%)	5.9% (3.6-9.6%)	1.8 (1.1-3)	90 (53-150)
Barbados	2016	Female	50 to 54	32% (19-47%)	68% (53-81%)	1.4 (0.79-2.4)	3.4% (1.9-5.9%)	5.6% (3.2-9.2%)	2.6 (1.5-4.4)	110 (61-190)
Barbados	2016	Female	55 to 59	28% (17-44%)	72% (56-83%)	1.2 (0.64-2.1)	3% (1.6-5.3%)	4.8% (2.7-8%)	3.1 (1.7-5.1)	110 (59-190)
Barbados	2016	Female	60 to 64	25% (15-39%)	75% (61-85%)	1 (0.51-1.8)	2.5% (1-4.8%)	3.7% (1.7-6.7%)	2.9 (1.4-5.4)	88 (37-170)
Barbados	2016	Female	65 to 69	23% (14-35%)	77% (65-86%)	0.91 (0.46-1.6)	2.1% (0.85-4.2%)	3.1% (1.4-5.7%)	3.1 (1.4-5.8)	78 (31-150)
Barbados	2016	Female	70 to 74	21% (13-33%)	79% (67-87%)	0.77 (0.38-1.4)	1.7% (0.42-3.7%)	2.5% (0.72-5.1%)	2.8 (0.82-5.8)	55 (13-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	2016	Female	75 to 79	18% (11-30%)	82% (70-89%)	0.67 (0.31-1.2)	1.4% (0.21-3.4%)	1.9% (0.32-4.3%)	3.1 (0.56-7.1)	47 (7-110)
Barbados	2016	Female	80 to 84	17% (9.3-27%)	83% (73-91%)	0.62 (0.3-1.1)	1.4% (0.26-3.1%)	1.7% (0.34-3.9%)	3.6 (0.71-8.1)	40 (7.5-93)
Barbados	2016	Female	85 to 89	17% (9.4-26%)	83% (74-91%)	0.63 (0.29-1.2)	1.4% (0.27-3.2%)	1.7% (0.29-3.9%)	3.7 (0.66-8.7)	29 (5.5-67)
Barbados	2016	Female	90 to 94	18% (11-29%)	82% (71-89%)	0.71 (0.3-1.4)	1.6% (0.22-3.9%)	1.9% (0.2-4.5%)	3.3 (0.37-8.1)	18 (2.4-43)
Barbados	2016	Female	95 plus	22% (11-37%)	78% (63-89%)	0.86 (0.29-1.8)	2% (0.11-5.4%)	2.2% (-0.00055-6.2%)	1.6 (-0.0004-4.4)	5.6 (0.29-15)
Barbados	2016	Male	All Ages	55% (47-63%)	45% (37-53%)	2.8 (2.1-3.6)	7.5% (5.7-9.4%)	7.5% (5.2-9.9%)	100 (68-130)	3200 (2400-4100)
Barbados	2016	Male	15 to 19	53% (35-71%)	47% (29-65%)	2 (0.93-3.4)	5.3% (3.3-8%)	8.4% (4.5-14%)	0.61 (0.31-1)	61 (38-94)
Barbados	2016	Male	20 to 24	64% (45-79%)	36% (21-55%)	3.8 (2-6.2)	11% (7.6-15%)	16% (9.2-23%)	1.6 (0.92-2.4)	150 (97-210)
Barbados	2016	Male	25 to 29	67% (47-82%)	33% (18-53%)	4.6 (2.5-7.1)	12% (8.7-16%)	17% (11-23%)	2.3 (1.4-3.3)	190 (130-250)
Barbados	2016	Male	30 to 34	68% (48-83%)	32% (17-52%)	4.8 (2.7-7.4)	12% (8.4-15%)	17% (11-22%)	2.8 (1.8-3.8)	200 (140-260)
Barbados	2016	Male	35 to 39	68% (51-82%)	32% (18-49%)	4.5 (2.5-7.2)	11% (7.7-14%)	16% (11-21%)	3.1 (2-4.3)	200 (140-270)
Barbados	2016	Male	40 to 44	67% (49-81%)	33% (19-51%)	4.3 (2.4-6.7)	10% (7.3-13%)	15% (10-20%)	3.8 (2.6-5.2)	220 (160-290)
Barbados	2016	Male	45 to 49	65% (44-80%)	35% (20-56%)	3.9 (2.3-6.3)	11% (7.8-14%)	15% (11-19%)	6.4 (4.3-8.5)	310 (210-400)
Barbados	2016	Male	50 to 54	63% (44-80%)	37% (20-56%)	3.6 (2-5.8)	10% (7-13%)	14% (9.4-18%)	8.3 (5.5-11)	350 (230-460)
Barbados	2016	Male	55 to 59	61% (43-78%)	39% (22-57%)	3.2 (1.8-5.3)	9.7% (6.3-13%)	13% (8.2-17%)	11 (6.7-15)	380 (240-530)
Barbados	2016	Male	60 to 64	59% (38-77%)	41% (23-62%)	2.9 (1.5-5)	8.4% (5.1-12%)	11% (6.4-15%)	12 (6.9-17)	350 (210-500)
Barbados	2016	Male	65 to 69	57% (37-75%)	43% (25-63%)	2.6 (1.4-4.3)	7.4% (4.3-11%)	9.2% (5.3-13%)	12 (7-19)	310 (180-460)
Barbados	2016	Male	70 to 74	53% (35-71%)	47% (29-65%)	2.1 (1.1-3.7)	5.7% (2.8-9%)	6.9% (3.4-11%)	10 (5-16)	200 (100-320)
Barbados	2016	Male	75 to 79	48% (29-68%)	52% (32-71%)	1.8 (0.85-3.2)	4.5% (2-7.7%)	5.3% (2.4-9%)	8.8 (3.9-15)	140 (61-230)
Barbados	2016	Male	80 to 84	42% (25-61%)	58% (39-75%)	1.5 (0.74-2.5)	3.7% (1.7-6.4%)	4.2% (1.9-7.3%)	7.7 (3.5-13)	86 (40-150)
Barbados	2016	Male	85 to 89	38% (22-54%)	62% (46-78%)	1.3 (0.61-2.2)	3.1% (1.4-5.6%)	3.5% (1.5-6.4%)	5.2 (2.2-9.5)	42 (18-75)
Barbados	2016	Male	90 to 94	38% (24-54%)	62% (46-76%)	1.3 (0.59-2.3)	2.9% (1-5.5%)	3.1% (1-6.2%)	3 (0.99-6)	17 (6-32)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Barbados	2016	Male	95 plus	41% (22-61%)	59% (39-78%)	1.4 (0.51-2.7)	3.1% (0.89-6.2%)	3.3% (0.75-7%)	0.99 (0.22-2.2)	3.8 (1-7.7)
Belize	1990	Female	All Ages	22% (18-26%)	78% (74-82%)	0.84 (0.69-1.1)	1.5% (1.2-2%)	2.4% (1.6-3.4%)	11 (7.1-16)	430 (330-560)
Belize	1990	Female	15 to 19	28% (20-38%)	72% (62-80%)	1.1 (0.63-1.8)	2.1% (1.3-3.3%)	4.2% (2.4-6.8%)	0.27 (0.15-0.45)	29 (18-43)
Belize	1990	Female	20 to 24	38% (27-49%)	62% (51-73%)	1.9 (1.1-2.8)	3.9% (2.7-5.5%)	6.9% (4.3-10%)	0.5 (0.31-0.74)	51 (35-72)
Belize	1990	Female	25 to 29	38% (27-49%)	62% (51-73%)	2 (1.2-2.9)	4.1% (3-5.5%)	7% (4.6-9.9%)	0.54 (0.35-0.79)	51 (35-69)
Belize	1990	Female	30 to 34	37% (25-49%)	63% (51-75%)	2 (1.2-3.1)	4.1% (2.9-5.6%)	7.4% (4.9-10%)	0.5 (0.32-0.71)	39 (27-54)
Belize	1990	Female	35 to 39	36% (24-47%)	64% (53-76%)	1.8 (1-3)	4.2% (2.9-6%)	7.6% (5-11%)	0.56 (0.36-0.81)	36 (24-51)
Belize	1990	Female	40 to 44	34% (23-44%)	66% (56-77%)	1.6 (0.89-2.6)	4.3% (2.8-6.2%)	7.6% (4.9-11%)	0.6 (0.37-0.87)	33 (21-47)
Belize	1990	Female	45 to 49	31% (20-41%)	69% (59-80%)	1.6 (0.82-2.6)	4.3% (2.6-6.6%)	7.1% (4.3-11%)	0.63 (0.37-0.96)	29 (17-45)
Belize	1990	Female	50 to 54	27% (19-37%)	73% (63-81%)	1.4 (0.75-2.4)	4.4% (2.5-7%)	6.8% (3.8-11%)	0.86 (0.48-1.3)	34 (19-54)
Belize	1990	Female	55 to 59	23% (15-32%)	77% (68-85%)	1.1 (0.58-1.9)	3.9% (2.2-6.4%)	5.8% (3.3-9.3%)	0.95 (0.52-1.6)	32 (18-54)
Belize	1990	Female	60 to 64	19% (12-28%)	81% (72-88%)	0.9 (0.45-1.6)	3.1% (1.5-5.4%)	4.3% (2.2-7.4%)	0.95 (0.47-1.7)	27 (13-49)
Belize	1990	Female	65 to 69	18% (12-25%)	82% (75-88%)	0.77 (0.4-1.3)	2.6% (1.3-4.5%)	3.5% (1.8-5.9%)	0.9 (0.44-1.5)	21 (10-36)
Belize	1990	Female	70 to 74	16% (10-22%)	84% (78-90%)	0.62 (0.32-1.1)	2.1% (0.91-4%)	2.8% (1.2-5.2%)	0.88 (0.38-1.7)	16 (6.9-32)
Belize	1990	Female	75 to 79	13% (8.4-20%)	87% (80-92%)	0.53 (0.25-0.95)	1.6% (0.59-3.3%)	2.1% (0.74-4.1%)	0.79 (0.28-1.6)	11 (4-24)
Belize	1990	Female	80 to 84	12% (7.6-17%)	88% (83-92%)	0.5 (0.26-0.85)	1.5% (0.57-2.8%)	1.8% (0.69-3.4%)	0.82 (0.3-1.6)	8.8 (3.3-17)
Belize	1990	Female	85 to 89	13% (7.5-19%)	87% (81-92%)	0.52 (0.26-0.92)	1.4% (0.5-2.9%)	1.7% (0.58-3.5%)	0.61 (0.21-1.2)	4.7 (1.6-9.4)
Belize	1990	Female	90 to 94	15% (9.1-22%)	85% (78-91%)	0.61 (0.28-1.2)	1.6% (0.4-3.7%)	1.8% (0.4-4.4%)	0.42 (0.096-1)	2.2 (0.57-5.3)
Belize	1990	Female	95 plus	18% (9.5-30%)	82% (70-90%)	0.75 (0.28-1.5)	1.8% (0.15-4.7%)	2% (0.044-5.4%)	0.17 (0.0039-0.47)	0.62 (0.052-1.6)
Belize	1990	Male	All Ages	46% (39-54%)	54% (46-61%)	2.1 (1.8-2.5)	5.2% (4.2-6.3%)	7.3% (5.5-9.4%)	42 (32-54)	1700 (1400-2100)
Belize	1990	Male	15 to 19	50% (38-61%)	50% (39-62%)	1.9 (1.1-2.9)	5.9% (3.9-8.3%)	8.4% (5.1-13%)	1 (0.63-1.6)	93 (61-130)
Belize	1990	Male	20 to 24	63% (51-73%)	37% (27-49%)	4 (2.7-5.8)	13% (9.2-17%)	16% (11-23%)	2.6 (1.7-3.7)	220 (150-290)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	1990	Male	25 to 29	70% (58-79%)	30% (21-42%)	5.1 (3.4-7.1)	15% (11-19%)	19% (14-25%)	3.3 (2.2-4.5)	240 (170-320)
Belize	1990	Male	30 to 34	74% (62-83%)	26% (17-38%)	5.2 (3.4-7.3)	15% (11-18%)	20% (14-25%)	2.9 (1.9-3.8)	190 (140-240)
Belize	1990	Male	35 to 39	75% (64-84%)	25% (16-36%)	4.9 (3-7.4)	14% (11-18%)	20% (14-25%)	2.7 (1.8-3.6)	160 (110-210)
Belize	1990	Male	40 to 44	73% (62-82%)	27% (18-38%)	4.9 (2.8-7.5)	14% (10-18%)	19% (14-25%)	2.6 (1.7-3.4)	140 (94-170)
Belize	1990	Male	45 to 49	70% (55-80%)	30% (20-45%)	4.5 (2.5-7.3)	14% (9.7-17%)	18% (13-23%)	2.5 (1.7-3.3)	110 (79-150)
Belize	1990	Male	50 to 54	68% (56-78%)	32% (22-44%)	4 (2.3-6.4)	13% (8.5-16%)	16% (11-21%)	3 (1.9-4.1)	120 (80-160)
Belize	1990	Male	55 to 59	67% (54-78%)	33% (22-46%)	3.7 (2.1-6)	12% (7.8-16%)	15% (9.6-20%)	3.5 (2.2-4.9)	120 (78-170)
Belize	1990	Male	60 to 64	65% (50-77%)	35% (23-50%)	3.4 (1.8-5.7)	10% (6.2-15%)	13% (7.6-18%)	4.1 (2.5-6)	120 (72-170)
Belize	1990	Male	65 to 69	62% (49-74%)	38% (26-51%)	2.9 (1.6-4.6)	9% (5.3-13%)	11% (6.4-16%)	3.6 (2.1-5.4)	86 (51-130)
Belize	1990	Male	70 to 74	58% (45-70%)	42% (30-55%)	2.3 (1.2-3.9)	6.9% (3.5-11%)	8.2% (4.1-13%)	3.2 (1.6-5.2)	62 (31-100)
Belize	1990	Male	75 to 79	54% (39-69%)	46% (31-61%)	1.9 (0.95-3.2)	5.4% (2.4-9.3%)	6.3% (2.7-11%)	2.6 (1.1-4.4)	39 (17-66)
Belize	1990	Male	80 to 84	47% (32-61%)	53% (39-68%)	1.6 (0.83-2.6)	4.5% (1.9-8%)	5.2% (2.1-9.3%)	2.3 (0.93-4)	25 (11-43)
Belize	1990	Male	85 to 89	39% (26-54%)	61% (46-74%)	1.4 (0.7-2.4)	3.8% (1.7-7.1%)	4.3% (1.8-8.1%)	1.4 (0.56-2.6)	11 (4.6-20)
Belize	1990	Male	90 to 94	38% (24-51%)	62% (49-76%)	1.3 (0.65-2.3)	3.3% (1.3-6.6%)	3.7% (1.3-7.5%)	0.71 (0.25-1.4)	3.8 (1.4-7.4)
Belize	1990	Male	95 plus	39% (23-56%)	61% (44-77%)	1.4 (0.57-2.7)	3.4% (0.91-7.3%)	3.6% (0.75-8.2%)	0.21 (0.041-0.46)	0.76 (0.2-1.6)
Belize	1995	Female	All Ages	23% (19-27%)	77% (73-81%)	0.84 (0.67-1)	1.8% (1.4-2.3%)	2.5% (1.7-3.6%)	12 (7.7-17)	480 (360-620)
Belize	1995	Female	15 to 19	29% (20-39%)	71% (61-80%)	1.1 (0.58-1.7)	2% (1.2-3%)	3.9% (2.1-6.2%)	0.26 (0.14-0.42)	28 (17-43)
Belize	1995	Female	20 to 24	38% (27-50%)	62% (50-73%)	1.8 (1.1-2.7)	3.8% (2.5-5.4%)	6.1% (3.8-9.2%)	0.5 (0.3-0.76)	51 (34-73)
Belize	1995	Female	25 to 29	38% (27-50%)	62% (50-73%)	1.9 (1.2-2.8)	4% (2.8-5.4%)	6.1% (4-8.6%)	0.55 (0.36-0.8)	50 (35-68)
Belize	1995	Female	30 to 34	37% (26-50%)	63% (50-74%)	1.9 (1.1-2.9)	3.8% (2.7-5.1%)	6.3% (4.3-8.7%)	0.66 (0.44-0.94)	53 (37-72)
Belize	1995	Female	35 to 39	36% (24-47%)	64% (53-76%)	1.7 (1-2.7)	3.6% (2.5-5.1%)	6.5% (4.5-9.2%)	0.57 (0.38-0.82)	39 (26-54)
Belize	1995	Female	40 to 44	34% (23-46%)	66% (54-77%)	1.5 (0.85-2.5)	4.2% (2.7-6.1%)	6.8% (4.3-9.8%)	0.9 (0.55-1.3)	48 (31-69)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	1995	Female	45 to 49	31% (20-42%)	69% (58-80%)	1.5 (0.79-2.5)	4.1% (2.4-6.4%)	6.4% (3.8-10%)	0.86 (0.5-1.4)	40 (23-64)
Belize	1995	Female	50 to 54	27% (19-37%)	73% (63-81%)	1.3 (0.71-2.2)	4.2% (2.2-6.8%)	6.2% (3.3-9.8%)	0.92 (0.49-1.5)	36 (19-58)
Belize	1995	Female	55 to 59	23% (15-32%)	77% (68-85%)	1.1 (0.55-1.8)	3.7% (1.9-6%)	5.2% (2.8-8.4%)	1.1 (0.56-1.7)	35 (19-59)
Belize	1995	Female	60 to 64	20% (13-29%)	80% (71-87%)	0.86 (0.43-1.5)	3% (1.4-5.3%)	4% (1.9-7%)	1.1 (0.5-1.9)	30 (14-54)
Belize	1995	Female	65 to 69	18% (11-25%)	82% (75-89%)	0.72 (0.38-1.2)	2.3% (1-4.1%)	3% (1.4-5.3%)	1.1 (0.48-1.8)	25 (11-44)
Belize	1995	Female	70 to 74	16% (10-22%)	84% (78-90%)	0.58 (0.3-1)	1.9% (0.7-3.7%)	2.4% (0.97-4.6%)	0.92 (0.37-1.7)	17 (6.7-33)
Belize	1995	Female	75 to 79	13% (8.5-20%)	87% (80-91%)	0.5 (0.24-0.88)	1.4% (0.44-2.9%)	1.8% (0.55-3.6%)	0.67 (0.21-1.4)	9.8 (3-20)
Belize	1995	Female	80 to 84	12% (7.7-18%)	88% (82-92%)	0.47 (0.24-0.79)	1.3% (0.46-2.5%)	1.6% (0.57-3.1%)	0.78 (0.29-1.5)	8.4 (3.1-16)
Belize	1995	Female	85 to 89	13% (7.6-19%)	87% (81-92%)	0.49 (0.24-0.88)	1.3% (0.41-2.6%)	1.5% (0.49-3.1%)	0.67 (0.22-1.4)	5.2 (1.7-11)
Belize	1995	Female	90 to 94	15% (9.2-23%)	85% (77-91%)	0.59 (0.26-1.1)	1.4% (0.31-3.3%)	1.6% (0.3-4%)	0.37 (0.068-0.92)	2 (0.43-4.7)
Belize	1995	Female	95 plus	19% (9.7-30%)	81% (70-90%)	0.73 (0.26-1.4)	1.6% (0.072-4.3%)	1.8% (-0.055-5%)	0.17 (-0.0053-0.49)	0.62 (0.028-1.7)
Belize	1995	Male	All Ages	48% (41-55%)	52% (45-59%)	2.1 (1.8-2.5)	6.6% (5.2-8.1%)	8.2% (6-11%)	55 (41-72)	2200 (1800-2800)
Belize	1995	Male	15 to 19	50% (38-61%)	50% (39-62%)	1.8 (0.99-2.8)	6.1% (4-8.8%)	8.2% (5-13%)	1.4 (0.77-2.1)	120 (75-170)
Belize	1995	Male	20 to 24	63% (51-74%)	37% (26-49%)	3.8 (2.4-5.4)	13% (9.2-18%)	16% (10-23%)	3.3 (2.1-4.9)	270 (180-370)
Belize	1995	Male	25 to 29	70% (58-79%)	30% (21-42%)	4.8 (3.2-6.8)	15% (11-19%)	19% (13-25%)	3.5 (2.3-4.9)	260 (180-340)
Belize	1995	Male	30 to 34	74% (62-83%)	26% (17-38%)	4.9 (3.2-6.8)	14% (10-17%)	18% (13-23%)	4 (2.7-5.5)	270 (190-350)
Belize	1995	Male	35 to 39	75% (64-84%)	25% (16-36%)	4.6 (2.9-6.7)	14% (9.9-17%)	18% (12-23%)	3.7 (2.4-5)	220 (150-290)
Belize	1995	Male	40 to 44	74% (63-82%)	26% (18-37%)	4.5 (2.6-6.8)	14% (9.6-17%)	18% (12-23%)	3.5 (2.4-4.8)	190 (130-250)
Belize	1995	Male	45 to 49	70% (57-79%)	30% (21-43%)	4.2 (2.4-6.6)	13% (9.3-17%)	17% (11-22%)	3.7 (2.4-4.8)	170 (120-220)
Belize	1995	Male	50 to 54	68% (56-78%)	32% (22-44%)	3.8 (2.1-6.1)	12% (8-16%)	15% (9.8-20%)	3.6 (2.3-4.8)	140 (92-190)
Belize	1995	Male	55 to 59	67% (55-78%)	33% (22-45%)	3.5 (1.9-5.6)	11% (7.3-16%)	14% (8.6-19%)	4.5 (2.8-6.4)	150 (95-210)
Belize	1995	Male	60 to 64	65% (51-76%)	35% (24-49%)	3.1 (1.7-5.4)	9.9% (5.7-15%)	12% (6.6-17%)	4.7 (2.7-7)	130 (78-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	1995	Male	65 to 69	62% (49-74%)	38% (26-51%)	2.7 (1.6-4.3)	8.6% (4.8-13%)	9.9% (5.5-15%)	6 (3.4-9.2)	140 (81-210)
Belize	1995	Male	70 to 74	58% (46-71%)	42% (29-54%)	2.2 (1.2-3.6)	6.7% (3.2-11%)	7.6% (3.5-12%)	4.6 (2.1-7.5)	86 (40-140)
Belize	1995	Male	75 to 79	54% (39-69%)	46% (31-61%)	1.8 (0.88-3)	5.3% (2.1-9.3%)	5.9% (2.3-10%)	3.4 (1.4-6.1)	50 (20-87)
Belize	1995	Male	80 to 84	47% (32-61%)	53% (39-68%)	1.5 (0.75-2.5)	4.2% (1.6-7.6%)	4.6% (1.7-8.7%)	2.9 (1.1-5.2)	31 (12-57)
Belize	1995	Male	85 to 89	39% (27-54%)	61% (46-73%)	1.3 (0.64-2.2)	3.4% (1.4-6.6%)	3.8% (1.4-7.5%)	1.7 (0.65-3.3)	13 (5.3-25)
Belize	1995	Male	90 to 94	38% (25-51%)	62% (49-75%)	1.2 (0.59-2.2)	2.9% (1-6.1%)	3.2% (0.97-6.8%)	0.62 (0.19-1.3)	3.4 (1.2-7)
Belize	1995	Male	95 plus	39% (23-57%)	61% (43-77%)	1.3 (0.53-2.5)	3% (0.79-6.7%)	3.2% (0.59-7.5%)	0.22 (0.042-0.53)	0.83 (0.22-1.9)
Belize	2000	Female	All Ages	21% (18-25%)	79% (75-82%)	0.74 (0.6-0.9)	1.7% (1.3-2.2%)	2.2% (1.5-3.2%)	13 (8.6-19)	540 (420-700)
Belize	2000	Female	15 to 19	24% (17-33%)	76% (67-83%)	0.82 (0.46-1.3)	1.7% (1.1-2.6%)	3% (1.7-4.8%)	0.28 (0.15-0.46)	31 (20-47)
Belize	2000	Female	20 to 24	33% (25-42%)	67% (58-75%)	1.4 (0.91-2.2)	3.1% (2.2-4.3%)	4.4% (2.7-6.6%)	0.51 (0.31-0.75)	56 (39-79)
Belize	2000	Female	25 to 29	36% (28-44%)	64% (56-72%)	1.7 (1.1-2.3)	3.3% (2.4-4.5%)	4.5% (3.1-6.4%)	0.65 (0.43-0.95)	62 (44-85)
Belize	2000	Female	30 to 34	34% (26-43%)	66% (57-74%)	1.7 (1.1-2.4)	3.3% (2.4-4.5%)	4.7% (3.3-6.6%)	0.71 (0.47-1.1)	57 (40-80)
Belize	2000	Female	35 to 39	32% (24-41%)	68% (59-76%)	1.4 (0.86-2.2)	3.4% (2.4-4.7%)	5.2% (3.7-7.1%)	0.87 (0.59-1.3)	57 (39-80)
Belize	2000	Female	40 to 44	30% (23-39%)	70% (61-77%)	1.3 (0.75-2)	3.5% (2.3-5.1%)	5.4% (3.6-7.6%)	0.95 (0.62-1.4)	51 (34-73)
Belize	2000	Female	45 to 49	28% (20-36%)	72% (64-80%)	1.2 (0.69-2)	3.5% (2.1-5.3%)	5.1% (3.2-7.9%)	1 (0.63-1.6)	48 (29-72)
Belize	2000	Female	50 to 54	23% (16-31%)	77% (69-84%)	1 (0.55-1.7)	3.5% (2-5.6%)	5% (2.8-7.8%)	1.1 (0.61-1.8)	44 (23-69)
Belize	2000	Female	55 to 59	19% (13-26%)	81% (74-87%)	0.8 (0.43-1.4)	3.1% (1.7-5.1%)	4.2% (2.4-6.7%)	1 (0.59-1.7)	35 (19-57)
Belize	2000	Female	60 to 64	16% (11-23%)	84% (77-89%)	0.66 (0.34-1.2)	2.5% (1.1-4.3%)	3.1% (1.5-5.5%)	1.1 (0.52-1.9)	31 (14-55)
Belize	2000	Female	65 to 69	15% (9.6-21%)	85% (79-90%)	0.56 (0.3-0.93)	2% (0.9-3.5%)	2.5% (1.2-4.4%)	1.1 (0.51-1.9)	25 (12-45)
Belize	2000	Female	70 to 74	13% (8.7-19%)	87% (81-91%)	0.47 (0.24-0.82)	1.5% (0.58-3%)	1.9% (0.75-3.8%)	1 (0.39-2)	19 (6.9-37)
Belize	2000	Female	75 to 79	12% (7.7-18%)	88% (82-92%)	0.42 (0.2-0.75)	1.2% (0.33-2.6%)	1.5% (0.41-3.1%)	0.76 (0.21-1.6)	11 (3-23)
Belize	2000	Female	80 to 84	12% (7.5-17%)	88% (83-93%)	0.42 (0.22-0.71)	1.2% (0.37-2.3%)	1.4% (0.43-2.7%)	0.74 (0.23-1.5)	7.9 (2.5-15)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	2000	Female	85 to 89	13% (7.5-18%)	87% (82-93%)	0.45 (0.22-0.8)	1.1% (0.32-2.4%)	1.3% (0.35-2.7%)	0.64 (0.17-1.4)	4.9 (1.3-10)
Belize	2000	Female	90 to 94	15% (8.9-22%)	85% (78-91%)	0.53 (0.25-1)	1.2% (0.24-3.1%)	1.4% (0.22-3.6%)	0.39 (0.063-1)	2.1 (0.41-5.1)
Belize	2000	Female	95 plus	18% (9.6-30%)	82% (70-90%)	0.65 (0.25-1.4)	1.4% (0.027-4%)	1.5% (-0.11-4.6%)	0.14 (-0.011-0.42)	0.5 (0.0099-1.5)
Belize	2000	Male	All Ages	52% (46-58%)	48% (42-54%)	2.3 (1.9-2.7)	7.5% (6-9.1%)	8.6% (6.5-11%)	74 (55-94)	3200 (2500-3900)
Belize	2000	Male	15 to 19	52% (44-59%)	48% (41-56%)	1.8 (1.1-2.6)	6.5% (4.4-9.2%)	8.4% (5.4-12%)	2.2 (1.3-3.4)	180 (120-270)
Belize	2000	Male	20 to 24	68% (61-74%)	32% (26-39%)	3.9 (2.7-5.4)	14% (10-19%)	17% (12-23%)	5.8 (3.8-8.4)	440 (300-600)
Belize	2000	Male	25 to 29	76% (70-80%)	24% (20-30%)	5 (3.5-6.8)	16% (12-20%)	19% (14-25%)	6.6 (4.6-9)	460 (330-620)
Belize	2000	Male	30 to 34	77% (71-82%)	23% (18-29%)	4.9 (3.5-6.6)	13% (10-17%)	16% (12-21%)	5.4 (3.8-7.2)	350 (250-460)
Belize	2000	Male	35 to 39	78% (72-83%)	22% (17-28%)	4.6 (3.1-6.6)	15% (11-19%)	18% (13-24%)	6.1 (4.2-8.2)	350 (250-450)
Belize	2000	Male	40 to 44	77% (72-83%)	23% (17-28%)	4.7 (2.9-6.9)	14% (10-18%)	17% (12-22%)	5.8 (3.9-7.7)	300 (200-390)
Belize	2000	Male	45 to 49	73% (67-79%)	27% (21-33%)	4.2 (2.6-6.3)	13% (9.1-17%)	16% (11-21%)	5.3 (3.4-7)	240 (160-310)
Belize	2000	Male	50 to 54	71% (64-77%)	29% (23-36%)	3.8 (2.1-5.8)	12% (8.2-16%)	15% (9.4-19%)	5.3 (3.4-7.2)	210 (130-280)
Belize	2000	Male	55 to 59	70% (62-77%)	30% (23-38%)	3.5 (2.1-5.6)	11% (7.1-16%)	13% (8.2-19%)	5.1 (3.1-7.2)	170 (110-240)
Belize	2000	Male	60 to 64	68% (59-77%)	32% (23-41%)	3.2 (1.9-5.3)	9.9% (5.7-14%)	11% (6.4-17%)	5.9 (3.4-9)	170 (97-250)
Belize	2000	Male	65 to 69	64% (54-72%)	36% (28-46%)	2.6 (1.7-4)	8.2% (4.6-12%)	9.3% (5.1-14%)	5.9 (3.3-9.1)	140 (78-210)
Belize	2000	Male	70 to 74	57% (47-67%)	43% (33-53%)	1.9 (1.1-3.1)	5.9% (2.8-9.9%)	6.6% (3.1-11%)	5.4 (2.4-9.2)	100 (47-170)
Belize	2000	Male	75 to 79	54% (41-66%)	46% (34-59%)	1.6 (0.83-2.6)	4.7% (2-8.1%)	5.2% (2.1-9%)	3.8 (1.6-6.6)	56 (23-95)
Belize	2000	Male	80 to 84	46% (32-59%)	54% (41-68%)	1.3 (0.73-2.2)	3.7% (1.5-6.8%)	4.1% (1.6-7.5%)	2.7 (1.1-4.9)	29 (12-53)
Belize	2000	Male	85 to 89	38% (25-52%)	62% (48-75%)	1.1 (0.56-1.9)	3% (1.2-5.5%)	3.2% (1.3-6.1%)	1.5 (0.59-2.9)	11 (4.8-22)
Belize	2000	Male	90 to 94	36% (23-50%)	64% (50-77%)	1.1 (0.52-2)	2.6% (0.98-5.2%)	2.7% (0.95-5.8%)	0.57 (0.2-1.2)	3.2 (1.2-6.5)
Belize	2000	Male	95 plus	37% (21-54%)	63% (46-79%)	1.1 (0.47-2.2)	2.7% (0.79-5.8%)	2.8% (0.59-6.3%)	0.16 (0.034-0.36)	0.6 (0.18-1.3)
Belize	2005	Female	All Ages	21% (18-25%)	79% (75-82%)	0.81 (0.66-1)	2% (1.6-2.5%)	2.7% (1.9-3.7%)	16 (11-22)	670 (520-860)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	2005	Female	15 to 19	23% (17-32%)	77% (68-83%)	0.88 (0.49-1.4)	1.9% (1.2-2.9%)	3.5% (1.9-5.6%)	0.32 (0.18-0.54)	35 (22-53)
Belize	2005	Female	20 to 24	33% (25-42%)	67% (58-75%)	1.6 (0.93-2.4)	3.4% (2.4-4.7%)	5.1% (3.2-7.6%)	0.65 (0.39-0.99)	70 (47-96)
Belize	2005	Female	25 to 29	35% (28-44%)	65% (56-72%)	1.8 (1.2-2.5)	3.6% (2.6-4.8%)	5.3% (3.6-7.5%)	0.78 (0.51-1.1)	72 (51-99)
Belize	2005	Female	30 to 34	34% (26-44%)	66% (56-74%)	1.7 (1.1-2.6)	3.4% (2.5-4.6%)	5% (3.5-7%)	0.97 (0.64-1.4)	77 (55-110)
Belize	2005	Female	35 to 39	31% (23-41%)	69% (59-77%)	1.5 (0.91-2.3)	3.4% (2.4-4.7%)	5.2% (3.7-7.2%)	1.2 (0.79-1.7)	77 (53-110)
Belize	2005	Female	40 to 44	30% (23-38%)	70% (62-77%)	1.4 (0.8-2.1)	3.5% (2.4-5%)	5.6% (3.8-7.7%)	1.3 (0.88-1.9)	71 (48-100)
Belize	2005	Female	45 to 49	28% (19-36%)	72% (64-81%)	1.3 (0.74-2.2)	3.9% (2.5-5.9%)	5.8% (3.8-8.8%)	1.3 (0.86-2)	61 (39-93)
Belize	2005	Female	50 to 54	23% (16-30%)	77% (70-84%)	1.1 (0.59-1.9)	3.9% (2.3-6.1%)	5.6% (3.3-8.6%)	1.4 (0.82-2.1)	54 (32-86)
Belize	2005	Female	55 to 59	18% (13-26%)	82% (74-87%)	0.86 (0.45-1.5)	3.4% (1.9-5.5%)	4.7% (2.8-7.4%)	1.2 (0.73-1.9)	41 (24-67)
Belize	2005	Female	60 to 64	16% (11-23%)	84% (77-89%)	0.71 (0.36-1.3)	2.7% (1.3-4.8%)	3.6% (1.8-6.1%)	1.1 (0.57-2)	32 (15-57)
Belize	2005	Female	65 to 69	15% (9.9-20%)	85% (80-90%)	0.6 (0.32-1)	2.3% (1.2-3.9%)	2.9% (1.5-4.9%)	1.2 (0.62-2.1)	29 (14-50)
Belize	2005	Female	70 to 74	13% (8.7-19%)	87% (81-91%)	0.5 (0.26-0.88)	1.8% (0.77-3.3%)	2.3% (1-4.1%)	1.1 (0.51-2.1)	21 (9-39)
Belize	2005	Female	75 to 79	12% (7.5-18%)	88% (82-92%)	0.45 (0.22-0.8)	1.5% (0.52-2.8%)	1.8% (0.66-3.4%)	1 (0.37-2)	15 (5.3-29)
Belize	2005	Female	80 to 84	11% (7.3-16%)	89% (84-93%)	0.44 (0.23-0.75)	1.4% (0.53-2.6%)	1.6% (0.64-3%)	0.86 (0.33-1.6)	9.1 (3.4-17)
Belize	2005	Female	85 to 89	12% (7.4-18%)	88% (82-93%)	0.48 (0.23-0.83)	1.3% (0.45-2.6%)	1.5% (0.5-3.1%)	0.64 (0.21-1.3)	4.9 (1.7-9.8)
Belize	2005	Female	90 to 94	14% (8.8-22%)	86% (78-91%)	0.56 (0.26-1.1)	1.4% (0.4-3.2%)	1.6% (0.4-3.8%)	0.42 (0.1-0.98)	2.2 (0.6-5)
Belize	2005	Female	95 plus	18% (9.5-29%)	82% (71-90%)	0.69 (0.26-1.4)	1.6% (0.19-4.3%)	1.8% (0.088-4.8%)	0.17 (0.0087-0.47)	0.62 (0.07-1.6)
Belize	2005	Male	All Ages	53% (47-58%)	47% (42-53%)	2.5 (2.1-2.9)	8.4% (6.8-10%)	9.8% (7.7-12%)	82 (64-100)	3700 (3000-4500)
Belize	2005	Male	15 to 19	51% (44-59%)	49% (41-56%)	1.9 (1.1-2.8)	7% (4.6-10%)	9.5% (5.9-14%)	2.4 (1.4-3.7)	200 (130-290)
Belize	2005	Male	20 to 24	68% (61-73%)	32% (27-39%)	4.2 (2.8-5.8)	15% (11-20%)	18% (12-25%)	6.8 (4.3-9.8)	510 (350-710)
Belize	2005	Male	25 to 29	76% (70-80%)	24% (20-30%)	5.4 (3.8-7.4)	16% (12-20%)	20% (14-26%)	6.6 (4.5-9.5)	470 (330-640)
Belize	2005	Male	30 to 34	77% (72-82%)	23% (18-28%)	5.3 (3.8-7.2)	14% (11-18%)	18% (13-23%)	6.9 (4.9-9.6)	450 (330-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	2005	Male	35 to 39	77% (72-83%)	23% (17-28%)	4.9 (3.3-7)	14% (11-18%)	18% (13-23%)	7.3 (5.2-9.8)	430 (310-560)
Belize	2005	Male	40 to 44	77% (72-83%)	23% (17-28%)	5 (3.2-7.4)	14% (10-17%)	17% (12-22%)	7 (4.9-9.3)	360 (270-480)
Belize	2005	Male	45 to 49	73% (67-79%)	27% (21-33%)	4.6 (2.8-6.9)	14% (10-18%)	17% (12-22%)	6.7 (4.6-8.7)	300 (210-390)
Belize	2005	Male	50 to 54	71% (64-77%)	29% (23-36%)	4 (2.3-6.2)	13% (9.1-17%)	16% (11-20%)	6.2 (4.2-8.2)	240 (170-320)
Belize	2005	Male	55 to 59	69% (62-77%)	31% (23-38%)	3.8 (2.3-5.7)	12% (8-16%)	15% (9.4-19%)	6.1 (3.8-8.2)	210 (130-280)
Belize	2005	Male	60 to 64	68% (59-76%)	32% (24-41%)	3.4 (2-5.5)	11% (6.3-15%)	12% (7.3-17%)	5.7 (3.4-8)	160 (98-230)
Belize	2005	Male	65 to 69	63% (54-72%)	37% (28-46%)	2.8 (1.7-4.3)	8.7% (5.3-13%)	10% (6.1-15%)	6.2 (3.6-9.1)	150 (85-210)
Belize	2005	Male	70 to 74	57% (47-67%)	43% (33-53%)	2.1 (1.2-3.5)	6.5% (3.4-10%)	7.3% (3.7-12%)	5 (2.5-8.2)	95 (49-150)
Belize	2005	Male	75 to 79	53% (40-66%)	47% (34-60%)	1.7 (0.89-2.8)	5.1% (2.3-8.6%)	5.7% (2.5-9.4%)	4.5 (1.8-7.7)	66 (28-110)
Belize	2005	Male	80 to 84	46% (31-59%)	54% (41-69%)	1.4 (0.76-2.3)	4.2% (1.9-7.1%)	4.6% (2-8%)	3 (1.3-5.1)	32 (15-55)
Belize	2005	Male	85 to 89	37% (25-52%)	63% (48-75%)	1.2 (0.6-2)	3.3% (1.5-5.8%)	3.6% (1.5-6.5%)	1.4 (0.58-2.5)	11 (4.8-19)
Belize	2005	Male	90 to 94	36% (23-49%)	64% (51-77%)	1.2 (0.55-2.1)	2.9% (1.1-5.7%)	3.1% (1.1-6.3%)	0.56 (0.2-1.1)	3.1 (1.2-5.9)
Belize	2005	Male	95 plus	37% (21-54%)	63% (46-79%)	1.2 (0.5-2.3)	3% (0.89-6.4%)	3.1% (0.74-7.1%)	0.17 (0.04-0.39)	0.65 (0.19-1.4)
Belize	2010	Female	All Ages	21% (18-25%)	79% (75-82%)	0.95 (0.75-1.2)	2.4% (1.8-3.1%)	3.3% (2.4-4.5%)	22 (15-30)	900 (690-1200)
Belize	2010	Female	15 to 19	23% (16-32%)	77% (68-84%)	1 (0.55-1.7)	2.1% (1.3-3.1%)	4% (2.1-6.5%)	0.45 (0.24-0.74)	47 (28-70)
Belize	2010	Female	20 to 24	33% (24-42%)	67% (58-76%)	1.8 (1-2.8)	3.8% (2.6-5.2%)	5.9% (3.5-8.9%)	0.9 (0.53-1.4)	90 (60-130)
Belize	2010	Female	25 to 29	35% (27-44%)	65% (56-73%)	2 (1.3-2.9)	3.8% (2.7-5.2%)	5.6% (3.7-8%)	1 (0.67-1.5)	93 (64-130)
Belize	2010	Female	30 to 34	34% (25-43%)	66% (57-75%)	2 (1.2-3.1)	3.7% (2.6-5%)	5.7% (3.9-7.8%)	1.1 (0.72-1.6)	86 (58-120)
Belize	2010	Female	35 to 39	31% (23-41%)	69% (59-77%)	1.7 (1-2.7)	3.8% (2.7-5.3%)	5.9% (4.1-8%)	1.6 (1-2.2)	99 (68-140)
Belize	2010	Female	40 to 44	30% (22-38%)	70% (62-78%)	1.5 (0.91-2.4)	3.7% (2.5-5.2%)	6% (4.2-8.3%)	1.5 (0.97-2.1)	82 (55-120)
Belize	2010	Female	45 to 49	27% (19-35%)	73% (65-81%)	1.5 (0.83-2.5)	4.3% (2.8-6.4%)	6.4% (4.1-9.5%)	2.3 (1.5-3.5)	110 (67-160)
Belize	2010	Female	50 to 54	23% (16-31%)	77% (69-84%)	1.2 (0.66-2.1)	4% (2.2-6.4%)	6% (3.6-9.3%)	1.9 (1.2-3.1)	79 (46-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	2010	Female	55 to 59	18% (13-26%)	82% (74-87%)	0.97 (0.52-1.7)	3.6% (2-5.8%)	5.2% (3.1-8.1%)	1.8 (1-2.8)	61 (35-99)
Belize	2010	Female	60 to 64	16% (11-23%)	84% (77-89%)	0.8 (0.4-1.4)	3.1% (1.5-5.1%)	4.1% (2.1-6.6%)	1.8 (0.93-3)	50 (25-87)
Belize	2010	Female	65 to 69	14% (9.5-20%)	86% (80-90%)	0.67 (0.35-1.1)	2.6% (1.4-4.3%)	3.4% (1.9-5.5%)	1.6 (0.87-2.7)	38 (20-63)
Belize	2010	Female	70 to 74	13% (8.4-19%)	87% (81-92%)	0.56 (0.29-1)	2.1% (0.99-3.7%)	2.7% (1.4-4.8%)	1.2 (0.62-2.2)	24 (11-43)
Belize	2010	Female	75 to 79	12% (7.5-17%)	88% (83-93%)	0.51 (0.24-0.92)	1.8% (0.69-3.5%)	2.2% (0.89-4.2%)	1.4 (0.59-2.7)	21 (8.3-41)
Belize	2010	Female	80 to 84	11% (7.3-17%)	89% (83-93%)	0.51 (0.26-0.86)	1.7% (0.69-3.2%)	2% (0.84-3.8%)	1.3 (0.52-2.3)	13 (5.4-25)
Belize	2010	Female	85 to 89	12% (7.3-18%)	88% (82-93%)	0.55 (0.25-0.99)	1.7% (0.65-3.5%)	2% (0.75-4.1%)	0.93 (0.35-1.9)	7 (2.7-14)
Belize	2010	Female	90 to 94	14% (8.8-21%)	86% (79-91%)	0.64 (0.29-1.2)	1.9% (0.56-4.2%)	2.1% (0.59-4.9%)	0.54 (0.15-1.2)	2.8 (0.86-6.4)
Belize	2010	Female	95 plus	18% (9.3-29%)	82% (71-91%)	0.79 (0.29-1.6)	2.1% (0.36-5.1%)	2.4% (0.28-6%)	0.23 (0.027-0.59)	0.83 (0.14-2)
Belize	2010	Male	All Ages	54% (48-59%)	46% (41-52%)	2.9 (2.4-3.6)	9.5% (7.7-11%)	11% (8.8-14%)	110 (85-140)	4600 (3700-5600)
Belize	2010	Male	15 to 19	51% (43-59%)	49% (41-57%)	2.2 (1.2-3.4)	7.6% (4.9-11%)	10% (6.3-16%)	2.9 (1.7-4.6)	240 (150-370)
Belize	2010	Male	20 to 24	67% (61-73%)	33% (27-39%)	4.8 (3.1-6.9)	16% (12-21%)	20% (14-28%)	7.4 (4.8-11)	560 (390-780)
Belize	2010	Male	25 to 29	75% (70-80%)	25% (20-30%)	6.2 (4.1-8.6)	16% (12-21%)	21% (15-28%)	7.1 (4.7-9.6)	500 (350-670)
Belize	2010	Male	30 to 34	77% (71-82%)	23% (18-29%)	6 (4.2-8.4)	15% (12-19%)	20% (15-25%)	6.5 (4.5-8.8)	420 (300-560)
Belize	2010	Male	35 to 39	77% (72-83%)	23% (17-28%)	5.6 (3.8-8.1)	14% (11-18%)	18% (13-23%)	6.9 (4.8-9.4)	410 (300-540)
Belize	2010	Male	40 to 44	77% (72-83%)	23% (17-28%)	5.7 (3.6-8.4)	15% (11-18%)	19% (14-24%)	8.3 (5.9-11)	430 (310-560)
Belize	2010	Male	45 to 49	73% (67-79%)	27% (21-33%)	5.2 (3.1-8)	14% (11-18%)	18% (13-22%)	9.7 (7-13)	450 (330-570)
Belize	2010	Male	50 to 54	70% (64-77%)	30% (23-36%)	4.6 (2.7-7.2)	14% (9.9-17%)	17% (12-21%)	10 (6.9-13)	400 (280-520)
Belize	2010	Male	55 to 59	69% (61-77%)	31% (23-39%)	4.3 (2.6-6.7)	13% (9.1-17%)	16% (11-20%)	11 (7.2-14)	360 (250-470)
Belize	2010	Male	60 to 64	68% (59-76%)	32% (24-41%)	3.9 (2.3-6.4)	12% (7.7-16%)	14% (8.9-18%)	9.9 (6.4-14)	280 (180-390)
Belize	2010	Male	65 to 69	63% (54-72%)	37% (28-46%)	3.2 (1.9-4.9)	10% (6.2-14%)	12% (7.1-16%)	8.7 (5.4-12)	210 (130-290)
Belize	2010	Male	70 to 74	57% (46-66%)	43% (34-54%)	2.4 (1.3-3.8)	7.5% (4-11%)	8.5% (4.5-13%)	7.7 (4.2-12)	150 (81-220)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	2010	Male	75 to 79	53% (40-66%)	47% (34-60%)	2 (1-3.2)	5.9% (2.8-9.2%)	6.6% (3-10%)	6.2 (2.8-10)	91 (43-150)
Belize	2010	Male	80 to 84	46% (31-58%)	54% (42-69%)	1.6 (0.87-2.6)	5% (2.3-8%)	5.6% (2.5-9%)	4.1 (1.9-7)	45 (21-75)
Belize	2010	Male	85 to 89	37% (25-51%)	63% (49-75%)	1.4 (0.67-2.4)	4% (1.7-7%)	4.4% (1.8-7.8%)	2 (0.84-3.6)	16 (6.9-27)
Belize	2010	Male	90 to 94	35% (23-49%)	65% (51-77%)	1.3 (0.61-2.5)	3.5% (1.4-6.7%)	3.8% (1.4-7.5%)	0.7 (0.25-1.4)	3.8 (1.5-7.2)
Belize	2010	Male	95 plus	36% (20-53%)	64% (47-80%)	1.4 (0.55-2.7)	3.5% (1.2-7.1%)	3.8% (0.98-8%)	0.21 (0.054-0.45)	0.77 (0.25-1.6)
Belize	2016	Female	All Ages	25% (21-29%)	75% (71-79%)	1.2 (0.88-1.7)	2.9% (2-4.1%)	4.1% (2.7-5.9%)	30 (19-44)	1200 (850-1700)
Belize	2016	Female	15 to 19	28% (20-38%)	72% (62-80%)	1.3 (0.7-2.3)	2.4% (1.5-3.8%)	5.1% (2.6-8.6%)	0.58 (0.29-0.99)	59 (35-94)
Belize	2016	Female	20 to 24	37% (26-49%)	63% (51-74%)	2.2 (1.2-3.6)	4.2% (2.8-6%)	7.3% (4.1-11%)	1.1 (0.59-1.7)	110 (70-160)
Belize	2016	Female	25 to 29	37% (26-49%)	63% (51-74%)	2.3 (1.3-3.7)	4.2% (2.9-5.7%)	6.6% (4.2-9.6%)	1.3 (0.78-2.1)	120 (76-170)
Belize	2016	Female	30 to 34	36% (25-49%)	64% (51-75%)	2.3 (1.3-3.8)	4.3% (2.9-6%)	6.8% (4.5-9.6%)	1.7 (1-2.6)	130 (80-180)
Belize	2016	Female	35 to 39	35% (24-46%)	65% (54-76%)	2.1 (1.2-3.5)	4.1% (2.6-5.9%)	6.7% (4.3-9.5%)	1.7 (1-2.7)	110 (71-170)
Belize	2016	Female	40 to 44	33% (23-45%)	67% (55-77%)	1.9 (1-3.1)	4% (2.5-6%)	6.9% (4.5-10%)	1.8 (1.1-3)	100 (63-160)
Belize	2016	Female	45 to 49	31% (19-41%)	69% (59-81%)	1.8 (0.89-3.1)	4.6% (2.8-7%)	7.2% (4.5-11%)	2.8 (1.6-4.3)	130 (75-200)
Belize	2016	Female	50 to 54	27% (18-36%)	73% (64-82%)	1.6 (0.81-2.8)	4.7% (2.6-7.7%)	7.1% (4-11%)	3.2 (1.8-5.2)	130 (70-210)
Belize	2016	Female	55 to 59	22% (15-31%)	78% (69-85%)	1.3 (0.63-2.3)	4.2% (2.3-7.1%)	6.1% (3.5-9.9%)	3 (1.6-4.9)	100 (54-170)
Belize	2016	Female	60 to 64	19% (12-28%)	81% (72-88%)	1 (0.5-1.9)	3.6% (1.7-6.2%)	4.8% (2.4-8%)	2.6 (1.3-4.5)	76 (35-130)
Belize	2016	Female	65 to 69	18% (11-25%)	82% (75-89%)	0.89 (0.43-1.5)	3.1% (1.5-5.4%)	4.1% (2-6.9%)	2.3 (1.1-3.9)	53 (24-94)
Belize	2016	Female	70 to 74	15% (10-22%)	85% (78-90%)	0.73 (0.36-1.3)	2.6% (1-4.8%)	3.3% (1.4-6.2%)	1.9 (0.81-3.5)	36 (15-69)
Belize	2016	Female	75 to 79	13% (8.5-20%)	87% (80-92%)	0.62 (0.28-1.1)	2.2% (0.77-4.3%)	2.7% (0.99-5.2%)	1.8 (0.66-3.5)	26 (9.3-52)
Belize	2016	Female	80 to 84	12% (7.4-17%)	88% (83-93%)	0.58 (0.28-1)	2% (0.82-4%)	2.4% (1-4.8%)	1.6 (0.62-3.1)	17 (6.6-34)
Belize	2016	Female	85 to 89	13% (7.4-19%)	87% (81-93%)	0.6 (0.28-1.2)	2% (0.76-4.3%)	2.3% (0.85-5.1%)	1.4 (0.5-3)	10 (3.9-23)
Belize	2016	Female	90 to 94	14% (8.8-22%)	86% (78-91%)	0.7 (0.3-1.4)	2.2% (0.66-4.8%)	2.5% (0.69-5.6%)	0.81 (0.22-1.8)	4.2 (1.2-9.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Belize	2016	Female	95 plus	18% (9.5-30%)	82% (70-90%)	0.87 (0.31-1.8)	2.5% (0.49-5.8%)	2.8% (0.45-6.7%)	0.31 (0.05-0.75)	1.1 (0.21-2.6)
Belize	2016	Male	All Ages	52% (45-58%)	48% (42-55%)	3.1 (2.3-4.1)	9.8% (7.6-12%)	12% (8.8-14%)	130 (90-170)	5300 (3900-6900)
Belize	2016	Male	15 to 19	49% (38-61%)	51% (39-62%)	2.2 (1.2-3.6)	7.5% (4.6-11%)	11% (6.1-16%)	3 (1.5-5)	250 (140-400)
Belize	2016	Male	20 to 24	63% (50-74%)	37% (26-50%)	4.7 (2.6-7.3)	15% (11-21%)	20% (13-27%)	7.8 (4.2-13)	600 (350-920)
Belize	2016	Male	25 to 29	70% (57-79%)	30% (21-43%)	5.9 (3.6-9.1)	16% (11-21%)	21% (14-28%)	7.8 (4.2-12)	570 (340-800)
Belize	2016	Male	30 to 34	73% (61-82%)	27% (18-39%)	6 (3.6-9.2)	15% (11-20%)	20% (14-26%)	7.6 (4.4-12)	500 (310-720)
Belize	2016	Male	35 to 39	74% (63-83%)	26% (17-37%)	5.7 (3.3-9)	14% (10-18%)	19% (13-24%)	7.5 (4.4-11)	450 (290-630)
Belize	2016	Male	40 to 44	73% (62-82%)	27% (18-38%)	5.6 (3.1-8.8)	14% (11-18%)	19% (13-24%)	9.4 (5.8-13)	490 (330-680)
Belize	2016	Male	45 to 49	69% (55-79%)	31% (21-45%)	5.1 (2.7-8.4)	14% (10-18%)	18% (13-23%)	11 (7.5-16)	520 (350-720)
Belize	2016	Male	50 to 54	67% (55-77%)	33% (23-45%)	4.6 (2.3-7.8)	14% (9.7-17%)	17% (12-21%)	13 (8.7-18)	530 (350-710)
Belize	2016	Male	55 to 59	66% (54-77%)	34% (23-46%)	4.3 (2.2-7.3)	13% (8.8-17%)	16% (11-20%)	14 (8.9-19)	470 (300-640)
Belize	2016	Male	60 to 64	64% (50-76%)	36% (24-50%)	3.8 (2-6.7)	11% (7.1-15%)	13% (8.2-18%)	12 (7.6-18)	350 (220-510)
Belize	2016	Male	65 to 69	61% (48-73%)	39% (27-52%)	3.3 (1.8-5.5)	9.9% (6-14%)	11% (6.8-16%)	11 (6.4-16)	260 (150-370)
Belize	2016	Male	70 to 74	58% (44-69%)	42% (31-56%)	2.6 (1.3-4.5)	7.8% (4-12%)	8.9% (4.6-13%)	8.1 (4.2-12)	150 (82-230)
Belize	2016	Male	75 to 79	54% (38-68%)	46% (32-62%)	2.2 (1-3.8)	6.2% (2.8-9.8%)	7% (3-11%)	6.4 (2.8-10)	94 (42-150)
Belize	2016	Male	80 to 84	46% (32-60%)	54% (40-68%)	1.8 (0.88-3.2)	5.3% (2.4-8.6%)	6% (2.6-9.7%)	4.7 (2.1-7.7)	51 (24-83)
Belize	2016	Male	85 to 89	39% (25-53%)	61% (47-75%)	1.6 (0.71-2.9)	4.4% (1.9-7.9%)	4.9% (2-8.9%)	2.7 (1.1-4.9)	21 (9.1-38)
Belize	2016	Male	90 to 94	36% (24-51%)	64% (49-76%)	1.5 (0.67-2.8)	3.9% (1.5-7.1%)	4.3% (1.5-8.1%)	0.95 (0.33-1.8)	5.2 (2-9.3)
Belize	2016	Male	95 plus	37% (21-55%)	63% (45-79%)	1.5 (0.61-3.1)	3.9% (1.3-8%)	4.2% (1.1-9%)	0.24 (0.062-0.51)	0.88 (0.28-1.8)
Cuba	1990	Female	All Ages	31% (25-38%)	69% (62-75%)	0.88 (0.73-1)	1.8% (1.3-2.5%)	1.6% (0.55-2.6%)	510 (180-860)	24000 (16000-32000)
Cuba	1990	Female	15 to 19	36% (23-52%)	64% (48-77%)	0.94 (0.48-1.5)	2.5% (1.2-4.2%)	5% (2-9.3%)	20 (8-38)	1800 (940-3100)
Cuba	1990	Female	20 to 24	42% (27-59%)	58% (41-73%)	1.4 (0.79-2.2)	3.8% (2.3-5.9%)	7.1% (3.3-12%)	35 (16-58)	3300 (2000-5000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	1990	Female	25 to 29	41% (26-59%)	59% (41-74%)	1.5 (0.84-2.2)	3.6% (2.3-5.2%)	6.6% (3.5-10%)	33 (18-53)	3100 (1900-4400)
Cuba	1990	Female	30 to 34	40% (24-57%)	60% (43-76%)	1.4 (0.79-2.2)	3.2% (2.1-4.7%)	5.8% (3.2-9.5%)	24 (13-39)	2000 (1300-2900)
Cuba	1990	Female	35 to 39	39% (25-57%)	61% (43-75%)	1.3 (0.74-2.1)	2.9% (1.8-4.4%)	5.3% (2.8-8.4%)	29 (15-46)	2000 (1200-3000)
Cuba	1990	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.2 (0.64-2)	2.8% (1.5-4.5%)	4.8% (2.4-7.9%)	34 (17-56)	2000 (1100-3000)
Cuba	1990	Female	45 to 49	36% (21-52%)	64% (48-79%)	1.2 (0.6-2.1)	2.8% (1.4-4.8%)	4.6% (2-8%)	43 (19-76)	2100 (1000-3600)
Cuba	1990	Female	50 to 54	32% (19-49%)	68% (51-81%)	1.1 (0.55-1.8)	2.5% (1.1-4.4%)	3.9% (1.6-7%)	47 (20-87)	1900 (870-3500)
Cuba	1990	Female	55 to 59	28% (17-44%)	72% (56-83%)	0.89 (0.46-1.6)	2.1% (0.86-3.8%)	3% (1.2-5.6%)	45 (18-84)	1600 (660-2900)
Cuba	1990	Female	60 to 64	25% (15-40%)	75% (60-85%)	0.76 (0.36-1.4)	1.6% (0.41-3.3%)	2.3% (0.52-4.6%)	45 (11-93)	1300 (340-2700)
Cuba	1990	Female	65 to 69	23% (14-36%)	77% (64-86%)	0.66 (0.33-1.1)	1.2% (0.2-2.6%)	1.7% (0.25-3.5%)	43 (6.1-87)	1000 (160-2100)
Cuba	1990	Female	70 to 74	21% (13-33%)	79% (67-87%)	0.55 (0.27-1)	0.87% (-0.17-2.2%)	1.1% (-0.24-2.9%)	38 (-8.1-97)	740 (-140-1900)
Cuba	1990	Female	75 to 79	18% (11-30%)	82% (70-89%)	0.48 (0.22-0.91)	0.57% (-0.36-1.7%)	0.69% (-0.51-2.2%)	30 (-23-96)	470 (-300-1400)
Cuba	1990	Female	80 to 84	17% (9.4-27%)	83% (73-91%)	0.44 (0.21-0.77)	0.43% (-0.38-1.4%)	0.49% (-0.54-1.7%)	23 (-25-80)	270 (-230-890)
Cuba	1990	Female	85 to 89	17% (9.6-26%)	83% (74-90%)	0.45 (0.21-0.8)	0.3% (-0.5-1.3%)	0.27% (-0.7-1.5%)	12 (-32-66)	120 (-200-520)
Cuba	1990	Female	90 to 94	19% (11-30%)	81% (70-89%)	0.52 (0.22-1)	0.29% (-0.74-1.7%)	0.22% (-1-2%)	5.4 (-26-49)	44 (-110-270)
Cuba	1990	Female	95 plus	22% (11-38%)	78% (62-89%)	0.62 (0.21-1.3)	0.39% (-0.95-2.5%)	0.26% (-1.4-2.8%)	2.1 (-11-22)	13 (-31-80)
Cuba	1990	Male	All Ages	55% (47-63%)	45% (37-53%)	2 (1.7-2.3)	6.2% (4.9-7.7%)	5.4% (3.4-7.9%)	2200 (1400-3200)	94000 (72000-120000)
Cuba	1990	Male	15 to 19	53% (36-71%)	47% (29-64%)	1.4 (0.79-2.3)	5% (3.1-7.5%)	7.3% (4-12%)	44 (24-73)	4100 (2500-6200)
Cuba	1990	Male	20 to 24	64% (45-80%)	36% (20-55%)	2.7 (1.7-4)	11% (7.3-15%)	14% (8.6-20%)	120 (74-180)	11000 (7100-15000)
Cuba	1990	Male	25 to 29	67% (48-82%)	33% (18-52%)	3.3 (2.1-4.7)	12% (8.3-16%)	16% (11-23%)	150 (96-210)	12000 (8200-16000)
Cuba	1990	Male	30 to 34	69% (49-83%)	31% (17-51%)	3.4 (2.1-5)	12% (8.3-16%)	17% (11-24%)	120 (81-170)	8600 (5900-11000)
Cuba	1990	Male	35 to 39	68% (51-82%)	32% (18-49%)	3.3 (2-5)	11% (7.5-15%)	16% (10-22%)	140 (84-190)	8600 (5800-12000)
Cuba	1990	Male	40 to 44	67% (51-81%)	33% (19-49%)	3.1 (1.8-4.7)	11% (6.8-15%)	15% (9-21%)	150 (90-210)	8000 (5300-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	1990	Male	45 to 49	65% (44-80%)	35% (20-56%)	2.9 (1.6-4.7)	9.8% (6.1-14%)	13% (7.6-19%)	170 (100-250)	8300 (5200-12000)
Cuba	1990	Male	50 to 54	63% (44-80%)	37% (20-56%)	2.6 (1.5-4.2)	8.7% (5.2-13%)	11% (6.4-17%)	190 (100-280)	7900 (4700-12000)
Cuba	1990	Male	55 to 59	61% (44-78%)	39% (22-56%)	2.4 (1.3-3.9)	7.7% (4.2-12%)	9.7% (5.2-15%)	200 (100-310)	6900 (3800-11000)
Cuba	1990	Male	60 to 64	59% (40-78%)	41% (22-60%)	2.2 (1.1-3.7)	6.1% (2.8-10%)	7.4% (3.2-13%)	200 (90-340)	6000 (2700-10000)
Cuba	1990	Male	65 to 69	58% (39-75%)	42% (25-61%)	1.9 (1-3.1)	4.9% (2.1-8.6%)	5.8% (2.4-10%)	200 (83-360)	5000 (2200-8700)
Cuba	1990	Male	70 to 74	54% (35-71%)	46% (29-65%)	1.6 (0.79-2.8)	3.6% (1.2-7.3%)	4.1% (1.2-8.6%)	190 (54-390)	3700 (1200-7400)
Cuba	1990	Male	75 to 79	49% (30-68%)	51% (32-70%)	1.3 (0.65-2.3)	2.6% (0.57-5.5%)	2.9% (0.49-6.3%)	160 (27-350)	2400 (530-5200)
Cuba	1990	Male	80 to 84	43% (26-62%)	57% (38-74%)	1.1 (0.56-1.8)	1.8% (0.24-4.1%)	2% (0.15-4.7%)	110 (8.4-250)	1200 (160-2800)
Cuba	1990	Male	85 to 89	38% (23-55%)	62% (45-77%)	0.94 (0.47-1.6)	1.3% (0.0043-3.5%)	1.3% (-0.2-3.9%)	61 (-9.4-180)	520 (1.7-1400)
Cuba	1990	Male	90 to 94	39% (24-57%)	61% (43-76%)	0.96 (0.44-1.7)	0.97% (-0.37-3.3%)	0.91% (-0.7-3.6%)	21 (-16-86)	140 (-52-460)
Cuba	1990	Male	95 plus	42% (23-62%)	58% (38-77%)	1 (0.42-2)	0.96% (-0.76-3.8%)	0.78% (-1.3-4.2%)	4.8 (-8-26)	24 (-18-95)
Cuba	1995	Female	All Ages	32% (26-39%)	68% (61-74%)	0.85 (0.72-1)	1.7% (1.1-2.3%)	1.4% (0.4-2.4%)	480 (140-870)	22000 (15000-31000)
Cuba	1995	Female	15 to 19	37% (24-53%)	63% (47-76%)	0.9 (0.45-1.5)	2.2% (1.1-3.7%)	4.7% (1.9-8.7%)	11 (4.4-20)	1000 (540-1700)
Cuba	1995	Female	20 to 24	43% (27-60%)	57% (40-73%)	1.4 (0.75-2.1)	3.5% (2.1-5.3%)	6.4% (2.9-11%)	25 (11-44)	2600 (1500-4000)
Cuba	1995	Female	25 to 29	42% (26-59%)	58% (41-74%)	1.4 (0.8-2.1)	3.3% (2.2-4.6%)	5.9% (3.1-9.3%)	27 (14-42)	2700 (1700-3800)
Cuba	1995	Female	30 to 34	41% (25-58%)	59% (42-75%)	1.3 (0.8-2.1)	2.9% (2-4.2%)	5.3% (2.9-8.4%)	28 (16-44)	2500 (1600-3600)
Cuba	1995	Female	35 to 39	40% (26-58%)	60% (42-74%)	1.2 (0.71-2)	2.7% (1.7-4.1%)	5% (2.8-7.8%)	26 (15-41)	1900 (1100-2700)
Cuba	1995	Female	40 to 44	39% (24-57%)	61% (43-76%)	1.2 (0.6-1.9)	2.6% (1.4-4%)	4.6% (2.3-7.4%)	36 (17-59)	2100 (1100-3200)
Cuba	1995	Female	45 to 49	37% (22-54%)	63% (46-78%)	1.1 (0.58-1.9)	2.6% (1.3-4.5%)	4.3% (1.9-7.5%)	43 (20-77)	2100 (1000-3500)
Cuba	1995	Female	50 to 54	33% (20-49%)	67% (51-80%)	1 (0.53-1.7)	2.4% (1-4.2%)	3.7% (1.6-6.6%)	49 (21-89)	2000 (890-3600)
Cuba	1995	Female	55 to 59	29% (17-45%)	71% (55-83%)	0.85 (0.43-1.5)	1.9% (0.79-3.5%)	2.9% (1.1-5.3%)	49 (19-90)	1700 (690-3100)
Cuba	1995	Female	60 to 64	26% (15-41%)	74% (59-85%)	0.73 (0.35-1.3)	1.5% (0.34-3.1%)	2.1% (0.43-4.3%)	45 (9.5-94)	1300 (300-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	1995	Female	65 to 69	24% (15-37%)	76% (63-85%)	0.63 (0.33-1.1)	1.1% (0.11-2.3%)	1.5% (0.12-3.2%)	42 (3.4-89)	1000 (110-2200)
Cuba	1995	Female	70 to 74	21% (13-34%)	79% (66-87%)	0.53 (0.26-0.95)	0.76% (-0.23-2%)	0.98% (-0.33-2.6%)	36 (-12-94)	690 (-210-1800)
Cuba	1995	Female	75 to 79	19% (11-31%)	81% (69-89%)	0.46 (0.21-0.85)	0.49% (-0.43-1.6%)	0.59% (-0.6-2%)	27 (-28-92)	420 (-370-1400)
Cuba	1995	Female	80 to 84	17% (9.9-28%)	83% (72-90%)	0.43 (0.2-0.74)	0.37% (-0.4-1.3%)	0.41% (-0.55-1.6%)	22 (-29-83)	270 (-280-930)
Cuba	1995	Female	85 to 89	17% (10-27%)	83% (73-90%)	0.44 (0.21-0.79)	0.25% (-0.5-1.2%)	0.21% (-0.73-1.4%)	11 (-37-71)	120 (-240-570)
Cuba	1995	Female	90 to 94	20% (11-31%)	80% (69-89%)	0.51 (0.22-1)	0.24% (-0.65-1.6%)	0.15% (-0.97-1.9%)	4.3 (-28-54)	44 (-120-290)
Cuba	1995	Female	95 plus	23% (12-39%)	77% (61-88%)	0.61 (0.21-1.3)	0.33% (-0.88-2.3%)	0.17% (-1.4-2.6%)	1.7 (-14-27)	14 (-37-97)
Cuba	1995	Male	All Ages	56% (48-65%)	44% (35-52%)	1.9 (1.7-2.2)	6.3% (4.9-7.9%)	5.2% (3.3-7.6%)	2300 (1400-3300)	96000 (75000-120000)
Cuba	1995	Male	15 to 19	54% (36-71%)	46% (29-64%)	1.3 (0.73-2.2)	4.9% (3-7.5%)	7.1% (3.9-12%)	29 (15-49)	2700 (1700-4100)
Cuba	1995	Male	20 to 24	65% (46-80%)	35% (20-54%)	2.6 (1.6-3.8)	10% (6.9-14%)	13% (8-20%)	100 (60-150)	9000 (6100-12000)
Cuba	1995	Male	25 to 29	68% (49-82%)	32% (18-51%)	3.1 (2-4.4)	11% (7.9-15%)	15% (10-22%)	140 (90-200)	11000 (7900-15000)
Cuba	1995	Male	30 to 34	69% (49-83%)	31% (17-51%)	3.2 (2-4.5)	11% (7.9-15%)	16% (10-21%)	160 (100-220)	11000 (7900-15000)
Cuba	1995	Male	35 to 39	69% (52-83%)	31% (17-48%)	3 (1.8-4.5)	11% (7.5-14%)	15% (9.8-21%)	130 (82-180)	8300 (5800-11000)
Cuba	1995	Male	40 to 44	68% (51-82%)	32% (18-49%)	2.9 (1.7-4.4)	10% (6.8-14%)	14% (8.9-20%)	160 (99-230)	9000 (6000-12000)
Cuba	1995	Male	45 to 49	66% (45-81%)	34% (19-55%)	2.7 (1.5-4.5)	9.8% (6.3-14%)	13% (7.7-19%)	180 (110-270)	8900 (5700-13000)
Cuba	1995	Male	50 to 54	64% (45-80%)	36% (20-55%)	2.5 (1.4-4.1)	8.6% (5.2-13%)	11% (6.3-16%)	210 (120-310)	8700 (5300-13000)
Cuba	1995	Male	55 to 59	62% (44-78%)	38% (22-56%)	2.3 (1.2-3.6)	7.5% (4.4-11%)	9.4% (5.3-14%)	230 (130-350)	8000 (4700-12000)
Cuba	1995	Male	60 to 64	60% (40-78%)	40% (22-60%)	2 (1.1-3.5)	6% (3-10%)	7.2% (3.4-12%)	210 (99-350)	6200 (3100-10000)
Cuba	1995	Male	65 to 69	58% (39-76%)	42% (24-61%)	1.8 (0.98-3)	4.6% (2.1-8.2%)	5.4% (2.3-10%)	210 (89-380)	5100 (2300-9000)
Cuba	1995	Male	70 to 74	55% (36-71%)	45% (29-64%)	1.5 (0.75-2.6)	3.3% (1.1-7%)	3.8% (1.2-8.2%)	180 (56-390)	3600 (1200-7600)
Cuba	1995	Male	75 to 79	49% (30-69%)	51% (31-70%)	1.2 (0.59-2.2)	2.3% (0.48-5.4%)	2.6% (0.42-6.2%)	150 (24-360)	2300 (480-5300)
Cuba	1995	Male	80 to 84	44% (26-62%)	56% (38-74%)	1 (0.52-1.7)	1.6% (0.21-3.8%)	1.7% (0.1-4.3%)	100 (5.8-260)	1200 (160-2800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	1995	Male	85 to 89	39% (23-56%)	61% (44-77%)	0.89 (0.44-1.6)	1.1% (-0.071-3.2%)	1.1% (-0.29-3.6%)	52 (-14-170)	460 (-32-1400)
Cuba	1995	Male	90 to 94	39% (25-56%)	61% (44-75%)	0.9 (0.43-1.6)	0.77% (-0.39-2.7%)	0.67% (-0.72-3%)	17 (-18-77)	120 (-60-430)
Cuba	1995	Male	95 plus	43% (24-63%)	57% (37-76%)	0.98 (0.41-1.9)	0.77% (-0.68-3.3%)	0.53% (-1.3-3.5%)	4.1 (-10-27)	25 (-22-100)
Cuba	2000	Female	All Ages	33% (27-40%)	67% (60-73%)	0.86 (0.73-1)	1.6% (1.1-2.2%)	1.3% (0.43-2.4%)	470 (150-850)	21000 (14000-28000)
Cuba	2000	Female	15 to 19	38% (25-55%)	62% (45-75%)	0.92 (0.47-1.6)	1.8% (1.1-2.9%)	4.3% (2-7.8%)	6.7 (3.1-12)	750 (450-1200)
Cuba	2000	Female	20 to 24	44% (29-60%)	56% (40-71%)	1.4 (0.75-2.2)	3.1% (2-4.5%)	5.8% (2.8-9.7%)	11 (5.6-19)	1400 (850-2000)
Cuba	2000	Female	25 to 29	43% (27-60%)	57% (40-73%)	1.4 (0.8-2.1)	3.1% (2.1-4.3%)	5.4% (2.9-8.4%)	18 (10-29)	2100 (1400-2900)
Cuba	2000	Female	30 to 34	42% (26-59%)	58% (41-74%)	1.4 (0.81-2.1)	2.8% (1.9-3.8%)	4.9% (3-7.5%)	22 (13-34)	2200 (1500-3100)
Cuba	2000	Female	35 to 39	41% (26-58%)	59% (42-74%)	1.3 (0.74-1.9)	2.5% (1.7-3.7%)	4.7% (2.7-7.1%)	29 (17-45)	2300 (1500-3200)
Cuba	2000	Female	40 to 44	40% (25-58%)	60% (42-75%)	1.2 (0.63-1.9)	2.5% (1.5-3.7%)	4.5% (2.5-6.9%)	30 (17-46)	1800 (1100-2700)
Cuba	2000	Female	45 to 49	38% (23-55%)	62% (45-77%)	1.1 (0.6-2)	2.5% (1.4-4%)	4.2% (2.2-7.1%)	44 (23-74)	2200 (1200-3500)
Cuba	2000	Female	50 to 54	34% (21-50%)	66% (50-79%)	1 (0.53-1.7)	2.4% (1.1-4%)	3.7% (1.7-6.2%)	50 (23-84)	2000 (1000-3400)
Cuba	2000	Female	55 to 59	30% (18-46%)	70% (54-82%)	0.86 (0.44-1.5)	1.9% (0.83-3.4%)	2.8% (1.2-5.1%)	51 (21-93)	1800 (750-3100)
Cuba	2000	Female	60 to 64	27% (16-42%)	73% (58-84%)	0.73 (0.36-1.3)	1.5% (0.43-3%)	2.1% (0.62-4.2%)	49 (14-98)	1500 (430-2900)
Cuba	2000	Female	65 to 69	24% (15-38%)	76% (62-85%)	0.64 (0.33-1.1)	1.1% (0.16-2.3%)	1.5% (0.18-3.1%)	43 (5.3-90)	1000 (150-2200)
Cuba	2000	Female	70 to 74	22% (14-35%)	78% (65-86%)	0.54 (0.27-0.98)	0.74% (-0.23-1.9%)	0.96% (-0.33-2.5%)	36 (-12-94)	700 (-210-1800)
Cuba	2000	Female	75 to 79	20% (12-32%)	80% (68-88%)	0.47 (0.21-0.87)	0.54% (-0.33-1.6%)	0.66% (-0.49-2.1%)	30 (-22-93)	470 (-290-1400)
Cuba	2000	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.44 (0.21-0.76)	0.39% (-0.31-1.3%)	0.44% (-0.49-1.6%)	23 (-25-83)	280 (-220-960)
Cuba	2000	Female	85 to 89	18% (10-29%)	82% (71-90%)	0.45 (0.21-0.81)	0.29% (-0.4-1.2%)	0.27% (-0.65-1.5%)	13 (-32-72)	140 (-190-590)
Cuba	2000	Female	90 to 94	20% (12-32%)	80% (68-88%)	0.51 (0.22-0.99)	0.28% (-0.51-1.5%)	0.19% (-0.84-1.8%)	6.1 (-26-57)	60 (-100-320)
Cuba	2000	Female	95 plus	24% (12-40%)	76% (60-88%)	0.61 (0.22-1.2)	0.36% (-0.72-2%)	0.16% (-1.3-2.4%)	2 (-16-29)	20 (-39-110)
Cuba	2000	Male	All Ages	58% (50-66%)	42% (34-50%)	1.9 (1.7-2.2)	6.4% (5-8%)	5.4% (3.6-7.7%)	2300 (1500-3300)	96000 (75000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	2000	Male	15 to 19	55% (38-72%)	45% (28-62%)	1.4 (0.74-2.2)	4.4% (2.9-6.6%)	6.9% (3.8-11%)	20 (11-33)	2000 (1300-3000)
Cuba	2000	Male	20 to 24	66% (48-80%)	34% (20-52%)	2.6 (1.6-3.9)	9.8% (6.8-13%)	13% (8.2-20%)	55 (33-81)	5200 (3600-7200)
Cuba	2000	Male	25 to 29	69% (51-83%)	31% (17-49%)	3.1 (1.9-4.4)	11% (7.7-14%)	15% (9.7-22%)	110 (68-150)	9000 (6300-12000)
Cuba	2000	Male	30 to 34	70% (51-84%)	30% (16-49%)	3.2 (2.1-4.4)	11% (7.8-14%)	16% (11-21%)	130 (89-180)	10000 (7300-13000)
Cuba	2000	Male	35 to 39	70% (53-84%)	30% (16-47%)	3 (1.9-4.3)	11% (7.7-14%)	16% (11-21%)	160 (110-220)	11000 (7800-14000)
Cuba	2000	Male	40 to 44	69% (52-83%)	31% (17-48%)	2.9 (1.7-4.3)	11% (7.6-15%)	15% (10-21%)	150 (100-210)	8800 (6200-12000)
Cuba	2000	Male	45 to 49	67% (47-82%)	33% (18-53%)	2.7 (1.5-4.3)	11% (7.2-14%)	14% (9.2-20%)	210 (140-300)	10000 (7200-14000)
Cuba	2000	Male	50 to 54	65% (46-81%)	35% (19-54%)	2.5 (1.4-4)	9.3% (6-13%)	12% (7.7-17%)	230 (150-330)	9700 (6400-14000)
Cuba	2000	Male	55 to 59	63% (45-79%)	37% (21-55%)	2.2 (1.2-3.5)	7.9% (4.9-12%)	9.9% (5.9-15%)	260 (160-390)	9100 (5700-13000)
Cuba	2000	Male	60 to 64	61% (41-78%)	39% (22-59%)	2 (1.1-3.3)	6.2% (3.4-10%)	7.5% (3.9-13%)	240 (130-410)	7300 (3900-12000)
Cuba	2000	Male	65 to 69	60% (40-77%)	40% (23-60%)	1.8 (0.98-2.9)	4.7% (2.4-8.1%)	5.7% (2.7-9.9%)	220 (100-380)	5400 (2700-9100)
Cuba	2000	Male	70 to 74	56% (37-73%)	44% (27-63%)	1.5 (0.77-2.6)	3.4% (1.3-6.8%)	3.9% (1.4-8%)	190 (68-390)	3800 (1500-7500)
Cuba	2000	Male	75 to 79	51% (32-70%)	49% (30-68%)	1.2 (0.6-2.2)	2.4% (0.66-5.1%)	2.7% (0.63-5.9%)	150 (36-330)	2300 (640-5000)
Cuba	2000	Male	80 to 84	45% (27-63%)	55% (37-73%)	1 (0.52-1.7)	1.6% (0.34-3.7%)	1.8% (0.23-4.3%)	99 (13-240)	1200 (240-2700)
Cuba	2000	Male	85 to 89	40% (24-57%)	60% (43-76%)	0.9 (0.44-1.6)	1.1% (0.066-3%)	1.2% (-0.1-3.5%)	54 (-4.7-160)	480 (28-1300)
Cuba	2000	Male	90 to 94	41% (26-58%)	59% (42-74%)	0.92 (0.43-1.7)	0.83% (-0.26-2.8%)	0.75% (-0.58-3.1%)	19 (-15-81)	140 (-42-450)
Cuba	2000	Male	95 plus	44% (25-64%)	56% (36-75%)	1 (0.41-1.9)	0.84% (-0.57-3.3%)	0.6% (-1.2-3.6%)	5.3 (-11-32)	31 (-21-120)
Cuba	2005	Female	All Ages	34% (28-42%)	66% (58-72%)	0.96 (0.82-1.1)	1.7% (1.2-2.3%)	1.5% (0.63-2.6%)	580 (240-970)	22000 (15000-30000)
Cuba	2005	Female	15 to 19	40% (26-56%)	60% (44-74%)	1 (0.53-1.8)	1.7% (1.1-2.6%)	4.5% (2.3-7.8%)	5.9 (3.1-10)	750 (450-1100)
Cuba	2005	Female	20 to 24	46% (30-62%)	54% (38-70%)	1.5 (0.81-2.5)	3% (2-4.3%)	5.8% (3-9.2%)	8.6 (4.5-14)	1200 (780-1800)
Cuba	2005	Female	25 to 29	45% (28-62%)	55% (38-72%)	1.6 (0.87-2.5)	3% (2.1-4.1%)	5.4% (3.2-8.4%)	10 (5.8-15)	1300 (850-1800)
Cuba	2005	Female	30 to 34	43% (27-61%)	57% (39-73%)	1.5 (0.88-2.3)	2.7% (1.9-3.8%)	5.1% (3.2-7.5%)	17 (11-26)	1900 (1300-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	2005	Female	35 to 39	42% (27-60%)	58% (40-73%)	1.4 (0.83-2.1)	2.5% (1.8-3.4%)	4.9% (3.2-7.1%)	26 (16-37)	2100 (1500-2900)
Cuba	2005	Female	40 to 44	41% (27-59%)	59% (41-73%)	1.3 (0.73-2)	2.5% (1.6-3.6%)	4.9% (3-7.2%)	39 (24-57)	2500 (1600-3400)
Cuba	2005	Female	45 to 49	39% (24-57%)	61% (43-76%)	1.3 (0.67-2.1)	2.6% (1.6-4%)	4.7% (2.6-7.4%)	42 (24-67)	2100 (1300-3200)
Cuba	2005	Female	50 to 54	36% (22-53%)	64% (47-78%)	1.1 (0.61-1.8)	2.5% (1.3-3.9%)	4.1% (2.1-6.4%)	57 (29-91)	2400 (1300-3700)
Cuba	2005	Female	55 to 59	31% (19-48%)	69% (52-81%)	0.96 (0.5-1.6)	2.2% (1.1-3.7%)	3.3% (1.6-5.6%)	60 (30-100)	2100 (1100-3500)
Cuba	2005	Female	60 to 64	28% (17-43%)	72% (57-83%)	0.82 (0.4-1.4)	1.8% (0.67-3.2%)	2.5% (0.97-4.6%)	62 (24-120)	1800 (710-3400)
Cuba	2005	Female	65 to 69	26% (16-39%)	74% (61-84%)	0.71 (0.37-1.2)	1.3% (0.41-2.5%)	1.8% (0.57-3.4%)	57 (18-110)	1400 (440-2600)
Cuba	2005	Female	70 to 74	23% (14-37%)	77% (63-86%)	0.6 (0.3-1.1)	0.93% (-0.024-2.1%)	1.2% (-0.062-2.8%)	48 (-2.4-110)	930 (-22-2100)
Cuba	2005	Female	75 to 79	21% (13-33%)	79% (67-87%)	0.52 (0.24-0.96)	0.71% (-0.15-1.9%)	0.9% (-0.24-2.3%)	43 (-11-110)	660 (-140-1700)
Cuba	2005	Female	80 to 84	19% (11-30%)	81% (70-89%)	0.49 (0.24-0.84)	0.6% (-0.096-1.5%)	0.72% (-0.18-1.9%)	39 (-9.8-110)	460 (-69-1200)
Cuba	2005	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.5 (0.24-0.88)	0.54% (-0.19-1.5%)	0.58% (-0.33-1.8%)	31 (-18-94)	280 (-91-750)
Cuba	2005	Female	90 to 94	21% (13-33%)	79% (67-87%)	0.57 (0.25-1.1)	0.56% (-0.33-2%)	0.54% (-0.58-2.4%)	21 (-22-89)	140 (-81-480)
Cuba	2005	Female	95 plus	25% (13-42%)	75% (58-87%)	0.68 (0.25-1.4)	0.67% (-0.52-2.7%)	0.57% (-0.97-3.1%)	9.6 (-16-52)	49 (-38-190)
Cuba	2005	Male	All Ages	60% (51-67%)	40% (33-49%)	2.1 (1.9-2.4)	6.9% (5.5-8.4%)	6.1% (4.2-8.4%)	2700 (1900-3700)	100000 (82000-120000)
Cuba	2005	Male	15 to 19	57% (39-73%)	43% (27-61%)	1.5 (0.8-2.4)	4.4% (2.9-6.3%)	7.2% (4.1-12%)	19 (11-31)	2100 (1400-3100)
Cuba	2005	Male	20 to 24	67% (50-82%)	33% (18-50%)	2.8 (1.7-4.2)	9.6% (6.7-13%)	14% (8.4-20%)	43 (26-63)	4500 (3100-6100)
Cuba	2005	Male	25 to 29	70% (52-83%)	30% (17-48%)	3.3 (2.1-4.9)	11% (7.9-15%)	16% (10-23%)	63 (40-89)	5600 (4000-7400)
Cuba	2005	Male	30 to 34	71% (52-84%)	29% (16-48%)	3.5 (2.2-4.9)	11% (8-14%)	17% (12-23%)	110 (76-150)	8600 (6300-11000)
Cuba	2005	Male	35 to 39	71% (54-84%)	29% (16-46%)	3.3 (2.1-4.7)	11% (8.2-14%)	17% (12-22%)	150 (110-200)	10000 (7800-13000)
Cuba	2005	Male	40 to 44	70% (53-83%)	30% (17-47%)	3.1 (1.9-4.5)	11% (8.2-14%)	17% (12-22%)	210 (150-280)	12000 (9100-16000)
Cuba	2005	Male	45 to 49	68% (47-82%)	32% (18-53%)	2.9 (1.6-4.6)	11% (8.2-15%)	16% (11-22%)	220 (150-290)	11000 (7600-14000)
Cuba	2005	Male	50 to 54	66% (48-82%)	34% (18-52%)	2.7 (1.5-4.2)	10% (7.1-14%)	14% (9.5-19%)	280 (190-390)	12000 (8100-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	2005	Male	55 to 59	64% (47-80%)	36% (20-53%)	2.5 (1.4-3.9)	8.9% (5.8-13%)	11% (7.3-16%)	300 (190-440)	11000 (7000-15000)
Cuba	2005	Male	60 to 64	62% (42-79%)	38% (21-58%)	2.2 (1.2-3.7)	7% (4-11%)	8.6% (4.8-14%)	310 (170-490)	9100 (5200-14000)
Cuba	2005	Male	65 to 69	61% (41-77%)	39% (23-59%)	2 (1.1-3.1)	5.5% (3.1-8.9%)	6.7% (3.7-11%)	290 (160-470)	7000 (3900-11000)
Cuba	2005	Male	70 to 74	57% (39-73%)	43% (27-61%)	1.6 (0.84-2.8)	4.1% (1.8-7.4%)	4.8% (2-8.9%)	240 (100-450)	4700 (2100-8700)
Cuba	2005	Male	75 to 79	52% (33-71%)	48% (29-67%)	1.4 (0.68-2.4)	2.9% (1-5.6%)	3.3% (1.1-6.7%)	190 (61-390)	3000 (1000-5800)
Cuba	2005	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.1 (0.58-1.9)	2% (0.57-4.1%)	2.3% (0.55-4.8%)	130 (32-280)	1500 (440-3100)
Cuba	2005	Male	85 to 89	42% (25-59%)	58% (41-75%)	1 (0.49-1.7)	1.5% (0.3-3.5%)	1.6% (0.18-4%)	77 (8.5-190)	650 (140-1500)
Cuba	2005	Male	90 to 94	42% (27-59%)	58% (41-73%)	1 (0.48-1.8)	1.2% (-0.049-3.3%)	1.2% (-0.32-3.8%)	35 (-9.3-110)	220 (-9.3-620)
Cuba	2005	Male	95 plus	45% (26-65%)	55% (35-74%)	1.1 (0.46-2.1)	1.2% (-0.37-3.9%)	1.1% (-0.86-4.4%)	11 (-8.9-46)	52 (-17-170)
Cuba	2010	Female	All Ages	35% (29-42%)	65% (58-71%)	1 (0.87-1.2)	1.8% (1.3-2.4%)	1.8% (0.86-3%)	720 (340-1200)	24000 (17000-33000)
Cuba	2010	Female	15 to 19	40% (27-57%)	60% (43-73%)	1.1 (0.54-1.9)	1.8% (1.1-2.7%)	4.8% (2.5-8.2%)	4.7 (2.5-8.1)	630 (380-960)
Cuba	2010	Female	20 to 24	46% (30-63%)	54% (37-70%)	1.6 (0.86-2.7)	3% (2-4.2%)	5.9% (3.2-9.4%)	8.8 (4.8-14)	1300 (850-1900)
Cuba	2010	Female	25 to 29	46% (30-63%)	54% (37-70%)	1.7 (0.93-2.7)	3% (2.1-4.1%)	5.7% (3.3-8.7%)	8.5 (5-13)	1200 (800-1700)
Cuba	2010	Female	30 to 34	44% (28-61%)	56% (39-72%)	1.6 (0.92-2.6)	2.7% (1.9-3.7%)	5.5% (3.5-8%)	11 (6.7-16)	1200 (840-1700)
Cuba	2010	Female	35 to 39	43% (28-61%)	57% (39-72%)	1.5 (0.88-2.4)	2.6% (1.8-3.5%)	5.2% (3.4-7.4%)	21 (14-31)	1900 (1300-2600)
Cuba	2010	Female	40 to 44	42% (27-60%)	58% (40-73%)	1.4 (0.78-2.2)	2.5% (1.7-3.7%)	5.2% (3.3-7.7%)	37 (23-53)	2400 (1600-3400)
Cuba	2010	Female	45 to 49	40% (24-57%)	60% (43-76%)	1.4 (0.75-2.2)	2.7% (1.7-4.1%)	5.1% (3.1-7.8%)	59 (36-91)	3000 (1900-4500)
Cuba	2010	Female	50 to 54	36% (23-54%)	64% (46-77%)	1.2 (0.68-1.9)	2.7% (1.5-4.1%)	4.6% (2.6-6.9%)	57 (33-89)	2400 (1400-3700)
Cuba	2010	Female	55 to 59	32% (20-48%)	68% (52-80%)	1 (0.56-1.7)	2.3% (1.3-3.8%)	3.7% (2.1-6%)	72 (42-120)	2600 (1500-4200)
Cuba	2010	Female	60 to 64	28% (17-44%)	72% (56-83%)	0.88 (0.44-1.5)	1.9% (0.85-3.4%)	2.8% (1.3-4.9%)	71 (32-130)	2100 (940-3800)
Cuba	2010	Female	65 to 69	26% (16-40%)	74% (60-84%)	0.77 (0.41-1.3)	1.5% (0.63-2.8%)	2.2% (0.84-3.8%)	73 (29-130)	1800 (710-3200)
Cuba	2010	Female	70 to 74	24% (15-37%)	76% (63-85%)	0.65 (0.33-1.1)	1.1% (0.12-2.5%)	1.5% (0.17-3.3%)	65 (7.4-140)	1300 (130-2800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	2010	Female	75 to 79	21% (12-34%)	79% (66-88%)	0.57 (0.26-1.1)	0.89% (-0.011-2.2%)	1.2% (-0.051-2.8%)	58 (-2.6-140)	890 (-11-2100)
Cuba	2010	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.54 (0.26-0.93)	0.8% (0.059-1.8%)	1% (0.0099-2.3%)	59 (0.59-140)	680 (49-1500)
Cuba	2010	Female	85 to 89	20% (11-30%)	80% (70-89%)	0.55 (0.26-0.99)	0.78% (0.031-1.9%)	0.89% (-0.083-2.3%)	52 (-4.8-140)	450 (17-1100)
Cuba	2010	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.62 (0.28-1.2)	0.87% (-0.082-2.5%)	0.93% (-0.31-3%)	40 (-13-130)	250 (-22-730)
Cuba	2010	Female	95 plus	26% (13-42%)	74% (58-87%)	0.75 (0.28-1.6)	1.1% (-0.29-3.7%)	1% (-0.66-4.3%)	20 (-13-82)	88 (-24-300)
Cuba	2010	Male	All Ages	60% (53-68%)	40% (32-47%)	2.3 (2-2.7)	7.4% (6-9%)	6.8% (4.9-9.2%)	3200 (2300-4400)	110000 (91000-140000)
Cuba	2010	Male	15 to 19	57% (40-73%)	43% (27-60%)	1.6 (0.83-2.6)	4.4% (2.9-6.2%)	7.8% (4.4-12%)	15 (8.9-25)	1800 (1200-2600)
Cuba	2010	Male	20 to 24	67% (51-82%)	33% (18-49%)	3 (1.7-4.5)	9.6% (6.8-13%)	14% (8.6-21%)	44 (27-66)	4800 (3300-6500)
Cuba	2010	Male	25 to 29	70% (53-83%)	30% (17-47%)	3.6 (2.2-5.4)	11% (7.7-14%)	17% (11-24%)	55 (35-78)	5000 (3500-6700)
Cuba	2010	Male	30 to 34	71% (53-85%)	29% (15-47%)	3.7 (2.3-5.4)	11% (8-14%)	18% (12-24%)	73 (50-98)	5700 (4100-7400)
Cuba	2010	Male	35 to 39	71% (54-84%)	29% (16-46%)	3.5 (2.2-5.2)	11% (8.3-14%)	18% (13-23%)	140 (97-180)	9300 (6900-12000)
Cuba	2010	Male	40 to 44	70% (54-84%)	30% (16-46%)	3.3 (2-4.9)	12% (8.6-15%)	19% (14-24%)	210 (160-280)	12000 (9400-16000)
Cuba	2010	Male	45 to 49	68% (49-83%)	32% (17-51%)	3.2 (1.9-4.8)	12% (8.8-15%)	18% (13-23%)	320 (230-410)	16000 (12000-20000)
Cuba	2010	Male	50 to 54	67% (49-82%)	33% (18-51%)	2.9 (1.7-4.5)	11% (8.1-15%)	16% (11-21%)	300 (210-410)	13000 (9000-17000)
Cuba	2010	Male	55 to 59	65% (47-80%)	35% (20-53%)	2.6 (1.5-4.2)	9.9% (6.7-13%)	13% (8.7-18%)	380 (250-530)	13000 (9000-18000)
Cuba	2010	Male	60 to 64	63% (43-80%)	37% (20-57%)	2.4 (1.3-4)	7.9% (4.8-12%)	9.9% (5.9-15%)	370 (220-560)	11000 (6600-16000)
Cuba	2010	Male	65 to 69	61% (42-78%)	39% (22-58%)	2.1 (1.2-3.3)	6.3% (3.8-9.7%)	7.7% (4.5-12%)	370 (220-580)	9000 (5400-14000)
Cuba	2010	Male	70 to 74	58% (39-74%)	42% (26-61%)	1.8 (0.93-3)	4.7% (2.3-8.1%)	5.7% (2.7-10%)	320 (150-560)	6300 (3100-11000)
Cuba	2010	Male	75 to 79	53% (34-71%)	47% (29-66%)	1.5 (0.74-2.6)	3.4% (1.4-6.3%)	4% (1.5-7.6%)	240 (94-460)	3700 (1500-6900)
Cuba	2010	Male	80 to 84	47% (29-65%)	53% (35-71%)	1.2 (0.64-2.1)	2.4% (0.83-4.6%)	2.8% (0.87-5.5%)	170 (54-350)	2000 (700-3900)
Cuba	2010	Male	85 to 89	42% (26-60%)	58% (40-74%)	1.1 (0.54-1.9)	1.9% (0.57-4%)	2% (0.49-4.8%)	110 (27-250)	910 (280-2000)
Cuba	2010	Male	90 to 94	43% (27-60%)	57% (40-73%)	1.1 (0.53-2)	1.6% (0.21-3.9%)	1.6% (0.00089-4.4%)	54 (0.03-150)	320 (42-800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	2010	Male	95 plus	45% (26-66%)	55% (34-74%)	1.2 (0.49-2.3)	1.6% (-0.16-4.6%)	1.5% (-0.62-5.2%)	17 (-7-59)	75 (-7.7-220)
Cuba	2016	Female	All Ages	34% (29-41%)	66% (59-71%)	1.1 (0.83-1.5)	2% (1.3-2.9%)	2.1% (0.96-3.7%)	910 (420-1600)	28000 (18000-41000)
Cuba	2016	Female	15 to 19	40% (27-57%)	60% (43-73%)	1.2 (0.56-2.1)	1.8% (1.1-2.9%)	5.3% (2.7-9%)	4.3 (2.2-7.5)	580 (350-910)
Cuba	2016	Female	20 to 24	46% (30-63%)	54% (37-70%)	1.8 (0.9-3.1)	3.2% (2.1-4.5%)	6.7% (3.5-11%)	7.8 (4.2-13)	1200 (740-1700)
Cuba	2016	Female	25 to 29	46% (30-63%)	54% (37-70%)	1.8 (0.96-3)	3.1% (2.2-4.3%)	6.1% (3.4-9.5%)	9.7 (5.3-15)	1400 (900-2000)
Cuba	2016	Female	30 to 34	44% (28-61%)	56% (39-72%)	1.8 (0.93-2.9)	2.9% (2-4%)	6% (3.7-9%)	11 (6.5-16)	1300 (840-1800)
Cuba	2016	Female	35 to 39	43% (28-61%)	57% (39-72%)	1.6 (0.92-2.7)	2.7% (1.8-3.9%)	5.9% (3.6-8.8%)	14 (8.7-22)	1300 (850-1800)
Cuba	2016	Female	40 to 44	42% (27-60%)	58% (40-73%)	1.5 (0.79-2.6)	2.7% (1.8-4.1%)	5.9% (3.5-9.2%)	33 (20-52)	2200 (1400-3200)
Cuba	2016	Female	45 to 49	40% (25-57%)	60% (43-75%)	1.5 (0.77-2.5)	2.9% (1.8-4.6%)	5.8% (3.5-8.9%)	62 (37-96)	3200 (2000-5000)
Cuba	2016	Female	50 to 54	36% (23-53%)	64% (47-77%)	1.3 (0.71-2.2)	2.9% (1.6-4.5%)	5.1% (2.9-7.9%)	90 (50-140)	3900 (2200-6200)
Cuba	2016	Female	55 to 59	32% (19-49%)	68% (51-81%)	1.1 (0.58-1.9)	2.5% (1.4-4.3%)	4.1% (2.3-6.9%)	82 (45-140)	2900 (1600-4800)
Cuba	2016	Female	60 to 64	29% (17-44%)	71% (56-83%)	0.96 (0.49-1.7)	2.2% (1-4.1%)	3.2% (1.5-6%)	94 (44-180)	2800 (1300-5300)
Cuba	2016	Female	65 to 69	26% (16-40%)	74% (60-84%)	0.84 (0.44-1.4)	1.8% (0.7-3.4%)	2.5% (0.99-4.8%)	90 (36-170)	2200 (860-4200)
Cuba	2016	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.71 (0.34-1.3)	1.4% (0.27-3%)	1.9% (0.41-4.2%)	83 (17-180)	1600 (320-3600)
Cuba	2016	Female	75 to 79	21% (12-35%)	79% (65-88%)	0.62 (0.28-1.2)	1.1% (0.099-2.6%)	1.4% (0.1-3.4%)	81 (5.7-200)	1200 (120-3000)
Cuba	2016	Female	80 to 84	20% (11-32%)	80% (68-89%)	0.58 (0.28-1)	0.97% (0.14-2.2%)	1.2% (0.09-2.9%)	84 (6.6-200)	980 (130-2300)
Cuba	2016	Female	85 to 89	20% (11-30%)	80% (70-89%)	0.59 (0.27-1.1)	0.95% (0.1-2.4%)	1.1% (0.016-3%)	73 (1.1-190)	630 (72-1600)
Cuba	2016	Female	90 to 94	22% (13-33%)	78% (67-87%)	0.66 (0.3-1.3)	1.1% (-0.03-3%)	1.2% (-0.23-3.7%)	58 (-12-180)	360 (-11-1000)
Cuba	2016	Female	95 plus	26% (13-43%)	74% (57-87%)	0.79 (0.29-1.7)	1.3% (-0.17-4.1%)	1.3% (-0.51-5%)	29 (-11-110)	120 (-16-390)
Cuba	2016	Male	All Ages	61% (53-68%)	39% (32-47%)	2.5 (1.9-3.3)	8% (6.1-10%)	7.4% (5.1-11%)	3900 (2600-5600)	130000 (99000-170000)
Cuba	2016	Male	15 to 19	57% (40-74%)	43% (26-60%)	1.7 (0.84-3)	4.6% (2.9-6.8%)	8.4% (4.6-14%)	14 (7.8-23)	1700 (1000-2400)
Cuba	2016	Male	20 to 24	68% (50-82%)	32% (18-50%)	3.2 (1.8-5.3)	10% (6.9-14%)	16% (9.1-23%)	41 (24-63)	4400 (3000-6100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cuba	2016	Male	25 to 29	71% (53-83%)	29% (17-47%)	3.9 (2.3-6.1)	11% (7.7-14%)	17% (11-24%)	64 (39-91)	6000 (4100-7900)
Cuba	2016	Male	30 to 34	71% (53-85%)	29% (15-47%)	4.1 (2.3-6.2)	11% (8-15%)	18% (12-25%)	75 (48-100)	6000 (4200-7800)
Cuba	2016	Male	35 to 39	71% (54-84%)	29% (16-46%)	3.8 (2.2-5.9)	11% (8.4-15%)	19% (13-26%)	91 (61-130)	6300 (4600-8300)
Cuba	2016	Male	40 to 44	70% (54-84%)	30% (16-46%)	3.6 (2-5.8)	12% (9-16%)	20% (14-26%)	200 (130-260)	11000 (8100-15000)
Cuba	2016	Male	45 to 49	68% (49-83%)	32% (17-51%)	3.4 (2-5.3)	13% (9.2-16%)	19% (14-25%)	330 (230-440)	16000 (12000-21000)
Cuba	2016	Male	50 to 54	67% (49-82%)	33% (18-51%)	3.1 (1.8-5)	12% (8.8-16%)	17% (12-22%)	490 (340-650)	20000 (15000-27000)
Cuba	2016	Male	55 to 59	65% (47-80%)	35% (20-53%)	2.9 (1.6-4.7)	11% (7.4-15%)	14% (9.9-20%)	450 (300-630)	16000 (11000-22000)
Cuba	2016	Male	60 to 64	63% (43-80%)	37% (20-57%)	2.6 (1.4-4.4)	8.7% (5.3-13%)	11% (6.7-16%)	480 (280-720)	14000 (8400-21000)
Cuba	2016	Male	65 to 69	62% (42-78%)	38% (22-58%)	2.3 (1.3-3.7)	7% (4.1-11%)	8.6% (5-13%)	460 (260-700)	11000 (6500-17000)
Cuba	2016	Male	70 to 74	58% (40-74%)	42% (26-60%)	1.9 (0.97-3.4)	5.3% (2.5-9.1%)	6.4% (3.1-11%)	400 (190-690)	7800 (3800-13000)
Cuba	2016	Male	75 to 79	53% (34-72%)	47% (28-66%)	1.6 (0.77-2.9)	3.9% (1.6-7.1%)	4.6% (1.8-8.5%)	330 (120-630)	5100 (2000-9400)
Cuba	2016	Male	80 to 84	47% (29-66%)	53% (34-71%)	1.3 (0.68-2.4)	2.7% (0.96-5.5%)	3.2% (1-6.6%)	240 (79-510)	2800 (970-5700)
Cuba	2016	Male	85 to 89	43% (26-60%)	57% (40-74%)	1.2 (0.59-2.1)	2.2% (0.69-4.8%)	2.5% (0.67-5.7%)	150 (42-360)	1300 (420-2900)
Cuba	2016	Male	90 to 94	43% (27-60%)	57% (40-73%)	1.2 (0.57-2.2)	1.9% (0.35-4.8%)	2% (0.14-5.5%)	79 (5.5-210)	460 (83-1100)
Cuba	2016	Male	95 plus	45% (27-66%)	55% (34-73%)	1.3 (0.52-2.5)	2% (0.022-5.4%)	2% (-0.37-6.2%)	25 (-4.7-82)	110 (1.1-300)
Dominica	1990	Female	All Ages	35% (29-42%)	65% (58-71%)	1.4 (1.2-1.7)	3.4% (2.5-4.5%)	4.2% (2.5-6.2%)	9.5 (5.7-14)	290 (210-380)
Dominica	1990	Female	15 to 19	42% (28-58%)	58% (42-72%)	1.7 (0.97-2.7)	2.5% (1.6-3.5%)	6.4% (4-9.7%)	0.074 (0.044-0.11)	8.9 (5.7-13)
Dominica	1990	Female	20 to 24	48% (32-64%)	52% (36-68%)	2.6 (1.5-3.8)	4.7% (3.4-6.3%)	9.1% (6.1-13%)	0.18 (0.11-0.25)	19 (13-27)
Dominica	1990	Female	25 to 29	48% (31-64%)	52% (36-69%)	2.7 (1.6-3.8)	5.2% (3.8-6.8%)	9.3% (6.4-12%)	0.23 (0.16-0.32)	22 (16-30)
Dominica	1990	Female	30 to 34	46% (29-63%)	54% (37-71%)	2.6 (1.6-3.9)	5.2% (3.8-6.8%)	9.4% (6.7-13%)	0.25 (0.17-0.34)	20 (14-26)
Dominica	1990	Female	35 to 39	45% (30-62%)	55% (38-70%)	2.4 (1.4-3.8)	5.3% (3.7-7.3%)	9.4% (6.6-13%)	0.31 (0.21-0.44)	21 (15-28)
Dominica	1990	Female	40 to 44	44% (29-62%)	56% (38-71%)	2.3 (1.2-3.7)	5.9% (3.8-8.4%)	9.8% (6.5-14%)	0.44 (0.28-0.63)	24 (16-34)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	1990	Female	45 to 49	42% (26-59%)	58% (41-74%)	2.2 (1.2-3.7)	5.4% (3.4-8.1%)	9.2% (5.9-14%)	0.39 (0.25-0.59)	19 (12-29)
Dominica	1990	Female	50 to 54	38% (24-55%)	62% (45-76%)	2 (1.1-3.4)	6% (3.5-9.3%)	8.7% (5.2-13%)	0.7 (0.41-1.1)	28 (17-44)
Dominica	1990	Female	55 to 59	34% (20-51%)	66% (49-80%)	1.7 (0.87-2.8)	5.4% (3-8.7%)	7.4% (4.1-12%)	0.74 (0.42-1.2)	26 (14-42)
Dominica	1990	Female	60 to 64	30% (18-46%)	70% (54-82%)	1.4 (0.73-2.5)	4.6% (2.1-7.9%)	5.8% (2.7-10%)	0.86 (0.4-1.5)	25 (11-43)
Dominica	1990	Female	65 to 69	28% (18-42%)	72% (58-82%)	1.3 (0.68-2)	3.9% (1.8-6.7%)	4.9% (2.3-8.2%)	0.88 (0.42-1.5)	21 (9.8-36)
Dominica	1990	Female	70 to 74	25% (16-39%)	75% (61-84%)	1.1 (0.54-1.9)	3.2% (1.1-6.2%)	3.9% (1.4-7.3%)	1 (0.36-1.9)	19 (6.6-36)
Dominica	1990	Female	75 to 79	22% (13-36%)	78% (64-87%)	0.92 (0.43-1.7)	2.7% (0.83-5.3%)	3.2% (0.99-6.4%)	0.94 (0.29-1.8)	14 (4.1-27)
Dominica	1990	Female	80 to 84	21% (12-32%)	79% (68-88%)	0.86 (0.42-1.4)	2.6% (0.89-5%)	2.9% (0.98-5.6%)	1.1 (0.37-2.1)	12 (3.9-23)
Dominica	1990	Female	85 to 89	21% (12-31%)	79% (69-88%)	0.87 (0.43-1.5)	2.5% (0.82-5.1%)	2.8% (0.88-5.7%)	0.82 (0.26-1.7)	6.2 (2-13)
Dominica	1990	Female	90 to 94	23% (14-35%)	77% (65-86%)	1 (0.45-1.9)	2.8% (0.71-6.4%)	3.1% (0.73-7.2%)	0.43 (0.1-1)	2.2 (0.58-5.1)
Dominica	1990	Female	95 plus	27% (15-44%)	73% (56-85%)	1.2 (0.44-2.4)	3.3% (0.41-8%)	3.6% (0.32-9%)	0.15 (0.013-0.37)	0.52 (0.068-1.3)
Dominica	1990	Male	All Ages	56% (48-65%)	44% (35-52%)	3 (2.7-3.3)	8.6% (7.2-10%)	9.8% (7.4-12%)	24 (18-31)	840 (700-990)
Dominica	1990	Male	15 to 19	59% (42-75%)	41% (25-58%)	2.5 (1.4-3.9)	6.8% (4.5-9.7%)	10% (6.5-15%)	0.35 (0.22-0.53)	33 (22-47)
Dominica	1990	Male	20 to 24	69% (52-83%)	31% (17-48%)	4.7 (3-6.5)	13% (10-17%)	19% (13-25%)	0.77 (0.53-1)	69 (51-90)
Dominica	1990	Male	25 to 29	72% (55-84%)	28% (16-45%)	5.6 (3.7-7.6)	15% (12-19%)	20% (15-25%)	1.3 (0.97-1.7)	99 (75-120)
Dominica	1990	Male	30 to 34	73% (55-85%)	27% (15-45%)	5.9 (3.8-8.1)	15% (12-18%)	20% (15-25%)	1.1 (0.88-1.4)	79 (61-96)
Dominica	1990	Male	35 to 39	72% (56-85%)	28% (15-44%)	5.6 (3.6-8.2)	13% (11-16%)	20% (15-24%)	0.81 (0.61-1)	54 (42-67)
Dominica	1990	Male	40 to 44	71% (55-85%)	29% (15-45%)	5.4 (3.1-8.1)	15% (11-18%)	19% (15-24%)	1.3 (0.97-1.7)	71 (53-88)
Dominica	1990	Male	45 to 49	70% (51-83%)	30% (17-49%)	5 (2.9-7.9)	13% (10-17%)	19% (14-23%)	1.2 (0.86-1.5)	57 (44-71)
Dominica	1990	Male	50 to 54	68% (51-83%)	32% (17-49%)	4.7 (2.6-7.3)	13% (9.5-17%)	17% (12-22%)	1.6 (1.1-2.1)	66 (47-84)
Dominica	1990	Male	55 to 59	66% (48-82%)	34% (18-52%)	4.3 (2.4-6.9)	12% (8.6-16%)	16% (11-20%)	1.8 (1.2-2.4)	63 (44-83)
Dominica	1990	Male	60 to 64	65% (45-81%)	35% (19-55%)	3.9 (2.2-6.6)	11% (7.1-15%)	13% (8.5-18%)	2.2 (1.4-3)	64 (41-86)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	1990	Male	65 to 69	63% (44-79%)	37% (21-56%)	3.5 (2-5.5)	10% (6.6-14%)	12% (7.7-16%)	2.5 (1.6-3.5)	59 (38-81)
Dominica	1990	Male	70 to 74	60% (42-76%)	40% (24-58%)	2.9 (1.5-4.9)	8.3% (4.4-12%)	9.3% (4.9-14%)	3 (1.6-4.5)	56 (30-84)
Dominica	1990	Male	75 to 79	55% (35-74%)	45% (26-65%)	2.4 (1.2-4.2)	6.5% (3.2-10%)	7.3% (3.5-11%)	2.3 (1.1-3.5)	33 (16-52)
Dominica	1990	Male	80 to 84	49% (30-68%)	51% (32-70%)	2.1 (1.1-3.3)	5.6% (2.7-9%)	6.2% (2.9-9.9%)	1.9 (0.89-3)	20 (9.7-33)
Dominica	1990	Male	85 to 89	45% (28-62%)	55% (38-72%)	1.8 (0.94-3)	4.9% (2.2-8.1%)	5.4% (2.3-9.2%)	1.3 (0.58-2.3)	10 (4.5-17)
Dominica	1990	Male	90 to 94	45% (29-62%)	55% (38-71%)	1.8 (0.89-3.3)	4.5% (1.8-8.2%)	4.9% (1.9-9.3%)	0.58 (0.21-1.1)	3.1 (1.2-5.7)
Dominica	1990	Male	95 plus	48% (28-68%)	52% (32-72%)	2 (0.81-3.7)	4.6% (1.4-9%)	5% (1.3-10%)	0.15 (0.039-0.31)	0.56 (0.16-1.1)
Dominica	1995	Female	All Ages	35% (29-42%)	65% (58-71%)	1.4 (1.2-1.6)	3.2% (2.3-4.3%)	3.8% (2.2-5.8%)	9.4 (5.2-15)	280 (200-380)
Dominica	1995	Female	15 to 19	42% (28-58%)	58% (42-72%)	1.7 (0.91-2.7)	2.4% (1.5-3.5%)	5.8% (3.5-9%)	0.065 (0.039-0.099)	7.9 (4.9-12)
Dominica	1995	Female	20 to 24	48% (32-64%)	52% (36-68%)	2.5 (1.4-3.7)	4.4% (3.1-5.9%)	8.2% (5.3-12%)	0.15 (0.098-0.22)	17 (12-24)
Dominica	1995	Female	25 to 29	47% (30-64%)	53% (36-70%)	2.6 (1.5-3.7)	5.1% (3.7-6.8%)	8.4% (5.8-11%)	0.26 (0.17-0.35)	24 (17-31)
Dominica	1995	Female	30 to 34	45% (29-63%)	55% (37-71%)	2.5 (1.5-3.7)	5.2% (3.8-6.9%)	8.6% (6.2-11%)	0.34 (0.24-0.47)	26 (19-35)
Dominica	1995	Female	35 to 39	45% (30-62%)	55% (38-70%)	2.3 (1.4-3.5)	5% (3.5-6.9%)	8.6% (6-12%)	0.33 (0.22-0.46)	22 (15-30)
Dominica	1995	Female	40 to 44	44% (29-61%)	56% (39-71%)	2.2 (1.2-3.5)	5.3% (3.4-7.6%)	9% (6-13%)	0.42 (0.27-0.61)	24 (15-34)
Dominica	1995	Female	45 to 49	41% (25-59%)	59% (41-75%)	2.1 (1.1-3.5)	5% (3.1-7.8%)	8.5% (5.4-13%)	0.39 (0.25-0.61)	19 (12-30)
Dominica	1995	Female	50 to 54	38% (24-56%)	62% (44-76%)	1.9 (1-3.3)	5.3% (3-8.4%)	8.1% (4.7-12%)	0.56 (0.32-0.87)	23 (13-36)
Dominica	1995	Female	55 to 59	34% (21-50%)	66% (50-79%)	1.6 (0.86-2.8)	4.9% (2.6-8.1%)	7% (3.8-11%)	0.69 (0.38-1.1)	24 (13-40)
Dominica	1995	Female	60 to 64	30% (18-47%)	70% (53-82%)	1.4 (0.69-2.4)	4.2% (1.8-7.5%)	5.4% (2.4-9.4%)	0.81 (0.36-1.4)	23 (10-42)
Dominica	1995	Female	65 to 69	28% (17-41%)	72% (59-83%)	1.2 (0.63-2)	3.5% (1.5-6.3%)	4.4% (1.9-7.8%)	0.81 (0.36-1.4)	19 (8.2-35)
Dominica	1995	Female	70 to 74	25% (16-38%)	75% (62-84%)	1 (0.52-1.8)	2.9% (0.79-5.8%)	3.5% (0.97-6.8%)	0.97 (0.28-2)	18 (4.8-37)
Dominica	1995	Female	75 to 79	22% (13-36%)	78% (64-87%)	0.88 (0.41-1.6)	2.4% (0.58-5.1%)	2.9% (0.71-5.9%)	0.94 (0.23-1.9)	14 (3.3-28)
Dominica	1995	Female	80 to 84	21% (12-32%)	79% (68-88%)	0.83 (0.39-1.4)	2.3% (0.67-4.7%)	2.7% (0.74-5.2%)	1.1 (0.3-2.1)	11 (3.2-22)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	1995	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.84 (0.41-1.5)	2.3% (0.66-4.9%)	2.6% (0.69-5.6%)	0.87 (0.23-1.9)	6.6 (1.8-14)
Dominica	1995	Female	90 to 94	23% (13-35%)	77% (65-87%)	0.97 (0.42-1.8)	2.5% (0.61-6%)	2.8% (0.58-6.9%)	0.5 (0.11-1.2)	2.6 (0.63-6.2)
Dominica	1995	Female	95 plus	27% (14-44%)	73% (56-86%)	1.2 (0.42-2.3)	3% (0.3-7.3%)	3.3% (0.2-8.3%)	0.2 (0.012-0.49)	0.69 (0.068-1.7)
Dominica	1995	Male	All Ages	57% (49-65%)	43% (35-51%)	2.9 (2.6-3.3)	8.6% (7.2-10%)	9.1% (6.8-12%)	26 (19-33)	900 (750-1100)
Dominica	1995	Male	15 to 19	59% (41-74%)	41% (26-59%)	2.4 (1.3-3.7)	7% (4.7-10%)	9.9% (6.1-15%)	0.4 (0.24-0.61)	35 (23-52)
Dominica	1995	Male	20 to 24	69% (51-82%)	31% (18-49%)	4.5 (2.8-6.4)	13% (10-17%)	18% (12-24%)	0.78 (0.53-1)	69 (51-90)
Dominica	1995	Male	25 to 29	72% (55-84%)	28% (16-45%)	5.4 (3.6-7.3)	15% (12-19%)	19% (14-24%)	1.6 (1.1-2)	110 (85-140)
Dominica	1995	Male	30 to 34	73% (54-85%)	27% (15-46%)	5.6 (3.6-7.8)	14% (11-17%)	18% (14-23%)	1.2 (0.88-1.5)	83 (64-100)
Dominica	1995	Male	35 to 39	72% (56-85%)	28% (15-44%)	5.3 (3.4-7.9)	13% (10-16%)	18% (13-22%)	1.1 (0.82-1.4)	70 (54-87)
Dominica	1995	Male	40 to 44	71% (56-84%)	29% (16-44%)	5.1 (3-7.6)	14% (10-17%)	18% (13-22%)	1.5 (1-1.9)	78 (57-99)
Dominica	1995	Male	45 to 49	70% (50-83%)	30% (17-50%)	4.8 (2.8-7.6)	13% (9.9-16%)	17% (13-21%)	1.4 (1-1.8)	67 (50-86)
Dominica	1995	Male	50 to 54	68% (50-83%)	32% (17-50%)	4.4 (2.5-6.9)	13% (9-16%)	16% (11-20%)	1.8 (1.2-2.3)	72 (50-92)
Dominica	1995	Male	55 to 59	66% (49-82%)	34% (18-51%)	4.1 (2.2-6.6)	12% (7.8-15%)	15% (10-19%)	1.7 (1.1-2.2)	60 (41-80)
Dominica	1995	Male	60 to 64	65% (45-81%)	35% (19-55%)	3.7 (2-6.2)	10% (6.4-14%)	12% (7.5-17%)	2.4 (1.5-3.3)	68 (42-94)
Dominica	1995	Male	65 to 69	63% (44-79%)	37% (21-56%)	3.3 (1.9-5.3)	9.4% (6-13%)	11% (6.9-15%)	2.5 (1.5-3.5)	59 (37-82)
Dominica	1995	Male	70 to 74	60% (41-75%)	40% (25-59%)	2.8 (1.5-4.7)	7.7% (4-12%)	8.5% (4.5-13%)	3.2 (1.7-5)	59 (32-92)
Dominica	1995	Male	75 to 79	54% (35-73%)	46% (27-65%)	2.3 (1.2-3.9)	6% (2.9-9.4%)	6.7% (3.1-11%)	2.2 (1-3.5)	32 (16-51)
Dominica	1995	Male	80 to 84	49% (31-67%)	51% (33-69%)	1.9 (1-3.1)	5.2% (2.3-8.3%)	5.6% (2.5-9.1%)	2 (0.85-3.2)	21 (9.4-34)
Dominica	1995	Male	85 to 89	44% (28-60%)	56% (40-72%)	1.7 (0.87-2.8)	4.3% (1.9-7.4%)	4.7% (2-8.2%)	1.3 (0.57-2.3)	10 (4.5-17)
Dominica	1995	Male	90 to 94	45% (29-62%)	55% (38-71%)	1.7 (0.82-3)	3.9% (1.5-7.3%)	4.2% (1.6-8.3%)	0.62 (0.23-1.2)	3.3 (1.3-6.3)
Dominica	1995	Male	95 plus	48% (28-68%)	52% (32-72%)	1.8 (0.76-3.4)	4.1% (1.2-8%)	4.4% (1-9%)	0.2 (0.047-0.42)	0.74 (0.21-1.5)
Dominica	2000	Female	All Ages	35% (29-41%)	65% (59-71%)	1.3 (1.1-1.5)	2.8% (2-3.9%)	3.3% (1.9-5.2%)	7.4 (4.1-12)	230 (160-320)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	2000	Female	15 to 19	41% (27-57%)	59% (43-73%)	1.5 (0.81-2.5)	2.3% (1.4-3.5%)	5.3% (3.1-8.4%)	0.068 (0.039-0.11)	8 (5-12)
Dominica	2000	Female	20 to 24	47% (30-64%)	53% (36-70%)	2.3 (1.3-3.5)	4.1% (2.8-5.7%)	7.2% (4.5-10%)	0.12 (0.075-0.18)	14 (9.7-20)
Dominica	2000	Female	25 to 29	47% (30-63%)	53% (37-70%)	2.4 (1.4-3.6)	4.2% (3.1-5.5%)	7% (4.7-9.6%)	0.15 (0.099-0.21)	16 (11-22)
Dominica	2000	Female	30 to 34	45% (29-62%)	55% (38-71%)	2.3 (1.4-3.5)	4.4% (3.2-5.8%)	7.4% (5.2-10%)	0.24 (0.16-0.33)	20 (14-27)
Dominica	2000	Female	35 to 39	44% (29-62%)	56% (38-71%)	2.1 (1.3-3.3)	4.4% (3-6.1%)	7.6% (5.5-10%)	0.3 (0.2-0.42)	21 (15-29)
Dominica	2000	Female	40 to 44	43% (28-61%)	57% (39-72%)	2 (1.1-3.2)	4.6% (3-6.5%)	8% (5.4-11%)	0.34 (0.22-0.5)	20 (13-29)
Dominica	2000	Female	45 to 49	41% (25-58%)	59% (42-75%)	1.9 (1-3.3)	4.5% (2.8-7.2%)	7.7% (4.9-12%)	0.39 (0.25-0.63)	20 (12-32)
Dominica	2000	Female	50 to 54	37% (23-54%)	63% (46-77%)	1.7 (0.92-2.9)	4.5% (2.4-7.2%)	7.2% (4.1-11%)	0.44 (0.25-0.7)	18 (10-30)
Dominica	2000	Female	55 to 59	33% (20-50%)	67% (50-80%)	1.5 (0.73-2.5)	4.1% (2.1-6.9%)	6.1% (3.3-10%)	0.52 (0.27-0.85)	18 (9.5-31)
Dominica	2000	Female	60 to 64	29% (17-45%)	71% (55-83%)	1.3 (0.61-2.2)	3.5% (1.5-6.5%)	4.8% (2.1-8.6%)	0.62 (0.28-1.1)	18 (7.7-34)
Dominica	2000	Female	65 to 69	27% (17-40%)	73% (60-83%)	1.1 (0.58-1.8)	3% (1.1-5.5%)	3.9% (1.6-6.9%)	0.64 (0.26-1.2)	15 (6-29)
Dominica	2000	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.93 (0.46-1.7)	2.4% (0.58-5.2%)	3% (0.8-6.2%)	0.7 (0.19-1.4)	13 (3.2-28)
Dominica	2000	Female	75 to 79	22% (13-35%)	78% (65-87%)	0.81 (0.37-1.5)	2% (0.41-4.4%)	2.4% (0.55-5.2%)	0.69 (0.16-1.5)	10 (2.2-22)
Dominica	2000	Female	80 to 84	20% (12-32%)	80% (68-88%)	0.77 (0.36-1.3)	1.9% (0.49-4%)	2.2% (0.57-4.6%)	0.81 (0.19-1.7)	8.7 (2.1-18)
Dominica	2000	Female	85 to 89	20% (12-31%)	80% (69-88%)	0.78 (0.38-1.4)	1.9% (0.53-4.2%)	2.2% (0.57-4.8%)	0.7 (0.17-1.5)	5.3 (1.4-11)
Dominica	2000	Female	90 to 94	22% (13-35%)	78% (65-87%)	0.88 (0.38-1.7)	2.2% (0.51-5%)	2.4% (0.48-5.8%)	0.45 (0.087-1.1)	2.4 (0.51-5.5)
Dominica	2000	Female	95 plus	26% (14-43%)	74% (57-86%)	1 (0.38-2.1)	2.5% (0.26-6.2%)	2.8% (0.16-7%)	0.19 (0.011-0.47)	0.68 (0.071-1.6)
Dominica	2000	Male	All Ages	57% (49-65%)	43% (35-51%)	2.8 (2.4-3.2)	7.7% (6.4-9.1%)	8% (6-10%)	21 (15-28)	770 (630-920)
Dominica	2000	Male	15 to 19	58% (41-74%)	42% (26-59%)	2.2 (1.2-3.5)	6.5% (4.2-9.4%)	9.3% (5.5-14%)	0.34 (0.2-0.52)	31 (20-45)
Dominica	2000	Male	20 to 24	68% (50-82%)	32% (18-50%)	4.1 (2.6-5.9)	12% (9.3-16%)	16% (11-22%)	0.59 (0.4-0.82)	54 (40-71)
Dominica	2000	Male	25 to 29	71% (54-84%)	29% (16-46%)	5 (3.2-7)	14% (10-17%)	17% (12-22%)	1.1 (0.75-1.4)	81 (61-100)
Dominica	2000	Male	30 to 34	72% (54-85%)	28% (15-46%)	5.2 (3.4-7.4)	12% (9.4-15%)	15% (12-19%)	0.94 (0.69-1.2)	70 (53-87)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	2000	Male	35 to 39	72% (56-85%)	28% (15-44%)	4.9 (3.1-7.2)	12% (9.2-15%)	15% (11-19%)	1.1 (0.82-1.5)	73 (55-93)
Dominica	2000	Male	40 to 44	71% (55-84%)	29% (16-45%)	4.7 (2.7-7)	12% (8.8-15%)	15% (11-19%)	1.1 (0.82-1.5)	66 (48-82)
Dominica	2000	Male	45 to 49	69% (49-83%)	31% (17-51%)	4.4 (2.5-6.9)	12% (8.7-15%)	15% (11-19%)	1.2 (0.88-1.6)	62 (45-80)
Dominica	2000	Male	50 to 54	67% (49-82%)	33% (18-51%)	4.1 (2.3-6.2)	12% (8.2-15%)	15% (10-18%)	1.6 (1.1-2)	65 (45-84)
Dominica	2000	Male	55 to 59	66% (47-81%)	34% (19-53%)	3.8 (2-5.9)	11% (7.1-14%)	13% (9.1-17%)	1.6 (1.1-2.1)	57 (39-75)
Dominica	2000	Male	60 to 64	64% (44-81%)	36% (19-56%)	3.4 (1.8-5.8)	9.2% (5.5-13%)	11% (6.8-15%)	1.9 (1.2-2.7)	57 (34-79)
Dominica	2000	Male	65 to 69	62% (43-78%)	38% (22-57%)	3 (1.7-4.9)	8.3% (5.2-12%)	9.8% (6.1-14%)	2.1 (1.3-3)	51 (31-73)
Dominica	2000	Male	70 to 74	59% (40-75%)	41% (25-60%)	2.5 (1.3-4.4)	6.6% (3.5-10%)	7.4% (3.8-12%)	2.4 (1.3-3.9)	47 (25-74)
Dominica	2000	Male	75 to 79	54% (35-73%)	46% (27-65%)	2.1 (1.1-3.7)	5.2% (2.4-8.5%)	5.8% (2.6-9.5%)	1.9 (0.84-3.1)	28 (13-46)
Dominica	2000	Male	80 to 84	48% (30-66%)	52% (34-70%)	1.8 (0.92-2.9)	4.4% (2-7.1%)	4.8% (2.1-7.8%)	1.7 (0.72-2.8)	18 (8-29)
Dominica	2000	Male	85 to 89	43% (27-60%)	57% (40-73%)	1.5 (0.78-2.6)	3.6% (1.6-6.3%)	3.9% (1.7-7%)	1 (0.44-1.8)	8 (3.7-14)
Dominica	2000	Male	90 to 94	44% (29-61%)	56% (39-71%)	1.6 (0.75-2.8)	3.3% (1.4-6.2%)	3.6% (1.4-7%)	0.5 (0.19-1)	2.7 (1.1-5.3)
Dominica	2000	Male	95 plus	47% (27-67%)	53% (33-73%)	1.7 (0.71-3.2)	3.6% (1.1-7.1%)	3.8% (0.96-7.9%)	0.19 (0.048-0.4)	0.72 (0.21-1.4)
Dominica	2005	Female	All Ages	35% (29-42%)	65% (58-71%)	1.3 (1.1-1.5)	2.6% (1.9-3.5%)	3.1% (1.8-4.9%)	7.1 (4-11)	220 (160-300)
Dominica	2005	Female	15 to 19	41% (27-58%)	59% (42-73%)	1.5 (0.77-2.3)	2.1% (1.4-3.1%)	5.3% (3.1-8.5%)	0.055 (0.032-0.088)	7 (4.4-10)
Dominica	2005	Female	20 to 24	47% (31-63%)	53% (37-69%)	2.2 (1.2-3.3)	4.1% (2.8-5.6%)	7.3% (4.5-11%)	0.14 (0.083-0.2)	15 (10-21)
Dominica	2005	Female	25 to 29	46% (30-64%)	54% (36-70%)	2.2 (1.3-3.4)	3.8% (2.7-5.1%)	6.9% (4.5-9.7%)	0.1 (0.065-0.15)	12 (8.5-17)
Dominica	2005	Female	30 to 34	44% (28-62%)	56% (38-72%)	2.1 (1.3-3.3)	4.1% (3-5.6%)	7.4% (5.1-10%)	0.21 (0.14-0.29)	17 (12-24)
Dominica	2005	Female	35 to 39	43% (28-61%)	57% (39-72%)	2 (1.2-3.1)	4% (2.9-5.6%)	7.4% (5.3-10%)	0.26 (0.18-0.37)	19 (13-26)
Dominica	2005	Female	40 to 44	42% (27-60%)	58% (40-73%)	1.9 (1-3)	4.2% (2.8-6.1%)	7.7% (5.3-11%)	0.36 (0.24-0.52)	21 (14-30)
Dominica	2005	Female	45 to 49	40% (25-57%)	60% (43-75%)	1.8 (0.98-3)	4.4% (2.8-6.9%)	7.4% (4.7-11%)	0.44 (0.27-0.7)	21 (13-34)
Dominica	2005	Female	50 to 54	37% (23-53%)	63% (47-77%)	1.6 (0.89-2.7)	4.1% (2.3-6.7%)	6.8% (4.1-10%)	0.46 (0.26-0.72)	19 (11-30)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	2005	Female	55 to 59	32% (20-49%)	68% (51-80%)	1.4 (0.72-2.3)	3.7% (1.8-6.3%)	5.7% (3.1-9.3%)	0.44 (0.24-0.74)	16 (8.2-27)
Dominica	2005	Female	60 to 64	29% (17-45%)	71% (55-83%)	1.2 (0.6-2.1)	3.1% (1.2-5.7%)	4.4% (2-7.8%)	0.48 (0.22-0.88)	14 (5.7-27)
Dominica	2005	Female	65 to 69	27% (17-41%)	73% (59-83%)	1 (0.55-1.7)	2.7% (1.1-4.9%)	3.7% (1.7-6.4%)	0.6 (0.26-1.1)	14 (5.7-26)
Dominica	2005	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.87 (0.43-1.6)	2.2% (0.59-4.8%)	2.9% (0.82-5.8%)	0.68 (0.19-1.4)	13 (3.3-26)
Dominica	2005	Female	75 to 79	22% (13-35%)	78% (65-87%)	0.76 (0.36-1.4)	1.9% (0.29-4.1%)	2.3% (0.41-4.9%)	0.68 (0.13-1.5)	9.8 (1.6-21)
Dominica	2005	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.71 (0.34-1.2)	1.8% (0.47-3.8%)	2.1% (0.58-4.3%)	0.77 (0.21-1.5)	8.2 (2.2-17)
Dominica	2005	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.72 (0.36-1.3)	1.8% (0.52-3.7%)	2.1% (0.58-4.3%)	0.72 (0.2-1.5)	5.5 (1.5-11)
Dominica	2005	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.83 (0.37-1.5)	2.1% (0.49-4.9%)	2.3% (0.49-5.6%)	0.5 (0.1-1.2)	2.6 (0.61-6.1)
Dominica	2005	Female	95 plus	26% (14-43%)	74% (57-86%)	0.98 (0.35-1.9)	2.4% (0.19-5.9%)	2.6% (0.12-6.7%)	0.22 (0.01-0.56)	0.79 (0.062-1.9)
Dominica	2005	Male	All Ages	57% (50-65%)	43% (35-50%)	2.7 (2.4-3)	7.8% (6.5-9.2%)	7.9% (6-10%)	22 (16-29)	800 (660-960)
Dominica	2005	Male	15 to 19	58% (40-74%)	42% (26-60%)	2.1 (1.1-3.2)	6.2% (4.1-8.9%)	9.3% (5.6-14%)	0.31 (0.19-0.47)	29 (19-42)
Dominica	2005	Male	20 to 24	68% (50-82%)	32% (18-50%)	3.9 (2.4-5.5)	12% (9-16%)	17% (11-23%)	0.64 (0.43-0.91)	58 (41-78)
Dominica	2005	Male	25 to 29	71% (53-84%)	29% (16-47%)	4.7 (3-6.7)	14% (10-18%)	18% (13-24%)	0.88 (0.6-1.2)	68 (49-89)
Dominica	2005	Male	30 to 34	72% (54-85%)	28% (15-46%)	4.9 (3.1-7)	13% (9.9-16%)	17% (13-22%)	1 (0.74-1.4)	73 (56-94)
Dominica	2005	Male	35 to 39	71% (55-85%)	29% (15-45%)	4.6 (2.9-6.8)	12% (9.1-15%)	16% (12-21%)	0.97 (0.69-1.3)	65 (49-81)
Dominica	2005	Male	40 to 44	70% (54-84%)	30% (16-46%)	4.4 (2.7-6.4)	13% (9.5-16%)	16% (12-21%)	1.6 (1.1-2)	86 (64-110)
Dominica	2005	Male	45 to 49	69% (49-83%)	31% (17-51%)	4.2 (2.4-6.5)	12% (9-15%)	16% (12-20%)	1.5 (1.1-2)	74 (54-95)
Dominica	2005	Male	50 to 54	67% (49-82%)	33% (18-51%)	3.8 (2.1-6)	11% (8-15%)	15% (10-19%)	1.7 (1.2-2.3)	73 (50-94)
Dominica	2005	Male	55 to 59	65% (48-81%)	35% (19-52%)	3.5 (1.9-5.6)	11% (7.1-14%)	13% (8.7-18%)	1.8 (1.2-2.5)	65 (43-87)
Dominica	2005	Male	60 to 64	63% (44-80%)	37% (20-56%)	3.2 (1.7-5.4)	9% (5.4-13%)	11% (6.5-15%)	2 (1.2-2.8)	58 (35-82)
Dominica	2005	Male	65 to 69	62% (43-78%)	38% (22-57%)	2.8 (1.6-4.5)	7.9% (4.9-11%)	9.5% (5.9-13%)	2 (1.3-2.9)	50 (31-72)
Dominica	2005	Male	70 to 74	59% (40-75%)	41% (25-60%)	2.4 (1.3-4.1)	6.3% (3.2-9.9%)	7.1% (3.6-11%)	2.3 (1.2-3.6)	44 (23-69)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	2005	Male	75 to 79	53% (35-72%)	47% (28-65%)	2 (0.98-3.5)	4.9% (2.3-8.3%)	5.5% (2.5-9.2%)	1.9 (0.87-3.3)	29 (13-48)
Dominica	2005	Male	80 to 84	48% (30-65%)	52% (35-70%)	1.6 (0.86-2.6)	4% (1.9-6.6%)	4.4% (2-7.3%)	1.6 (0.74-2.6)	17 (8.2-29)
Dominica	2005	Male	85 to 89	43% (26-60%)	57% (40-74%)	1.4 (0.72-2.4)	3.3% (1.5-6%)	3.6% (1.6-6.7%)	1.1 (0.48-2)	8.4 (3.8-15)
Dominica	2005	Male	90 to 94	43% (28-61%)	57% (39-72%)	1.4 (0.7-2.6)	3.1% (1.3-5.9%)	3.3% (1.2-6.5%)	0.51 (0.19-1)	2.8 (1.1-5.4)
Dominica	2005	Male	95 plus	46% (27-67%)	54% (33-73%)	1.6 (0.66-3)	3.3% (0.97-6.7%)	3.5% (0.84-7.4%)	0.2 (0.047-0.42)	0.72 (0.21-1.5)
Dominica	2010	Female	All Ages	35% (29-42%)	65% (58-71%)	1.3 (1.1-1.6)	2.5% (1.8-3.5%)	3.1% (1.7-4.9%)	7.8 (4.3-13)	230 (160-310)
Dominica	2010	Female	15 to 19	41% (27-57%)	59% (43-73%)	1.5 (0.79-2.4)	1.9% (1.2-2.7%)	4.6% (2.6-7.3%)	0.04 (0.022-0.064)	5.8 (3.6-8.9)
Dominica	2010	Female	20 to 24	46% (31-64%)	54% (36-69%)	2.2 (1.2-3.4)	4% (2.7-5.6%)	6.7% (4.1-10%)	0.14 (0.084-0.22)	16 (10-22)
Dominica	2010	Female	25 to 29	46% (29-63%)	54% (37-71%)	2.2 (1.3-3.5)	3.9% (2.8-5.4%)	6.5% (4-9.3%)	0.14 (0.088-0.2)	15 (10-21)
Dominica	2010	Female	30 to 34	44% (28-62%)	56% (38-72%)	2.2 (1.3-3.4)	3.5% (2.5-4.8%)	6.5% (4.4-9.2%)	0.12 (0.081-0.18)	12 (8.5-17)
Dominica	2010	Female	35 to 39	44% (29-61%)	56% (39-71%)	2 (1.1-3.1)	3.9% (2.7-5.4%)	7.1% (5-9.9%)	0.25 (0.17-0.36)	18 (12-25)
Dominica	2010	Female	40 to 44	42% (27-60%)	58% (40-73%)	1.9 (1.1-3)	4.2% (2.8-6.2%)	7.5% (5.1-11%)	0.38 (0.26-0.56)	22 (15-32)
Dominica	2010	Female	45 to 49	40% (25-57%)	60% (43-75%)	1.8 (0.98-3)	4.2% (2.6-6.7%)	7.2% (4.6-11%)	0.48 (0.31-0.76)	24 (15-38)
Dominica	2010	Female	50 to 54	37% (23-54%)	63% (46-77%)	1.6 (0.87-2.7)	4.3% (2.4-6.8%)	6.7% (4-10%)	0.55 (0.32-0.86)	23 (13-36)
Dominica	2010	Female	55 to 59	32% (20-49%)	68% (51-80%)	1.4 (0.71-2.4)	3.6% (1.8-6.1%)	5.6% (3.1-9.2%)	0.5 (0.27-0.81)	18 (9.2-31)
Dominica	2010	Female	60 to 64	29% (17-45%)	71% (55-83%)	1.2 (0.59-2.1)	3.1% (1.3-5.7%)	4.4% (2.1-7.8%)	0.55 (0.25-0.99)	16 (6.8-31)
Dominica	2010	Female	65 to 69	26% (16-41%)	74% (59-84%)	1 (0.55-1.7)	2.6% (1.1-4.8%)	3.7% (1.6-6.4%)	0.56 (0.25-0.97)	13 (5.4-24)
Dominica	2010	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.87 (0.43-1.6)	2.3% (0.62-4.7%)	2.9% (0.9-5.8%)	0.74 (0.23-1.5)	14 (4-28)
Dominica	2010	Female	75 to 79	22% (13-34%)	78% (66-87%)	0.77 (0.35-1.4)	1.9% (0.36-4.2%)	2.4% (0.5-5.1%)	0.77 (0.16-1.6)	11 (2.1-25)
Dominica	2010	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.72 (0.34-1.2)	1.9% (0.54-3.9%)	2.2% (0.62-4.6%)	0.87 (0.25-1.8)	9.3 (2.7-19)
Dominica	2010	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.73 (0.34-1.3)	1.9% (0.52-4%)	2.1% (0.56-4.6%)	0.82 (0.21-1.8)	6.3 (1.7-13)
Dominica	2010	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.83 (0.37-1.6)	2.1% (0.47-5%)	2.3% (0.45-5.6%)	0.62 (0.12-1.5)	3.3 (0.73-7.7)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	2010	Female	95 plus	26% (13-43%)	74% (57-87%)	1 (0.36-2.1)	2.4% (0.25-6.3%)	2.6% (0.15-7%)	0.29 (0.016-0.77)	1 (0.1-2.6)
Dominica	2010	Male	All Ages	58% (50-66%)	42% (34-50%)	2.8 (2.3-3.3)	8% (6.6-9.5%)	8.2% (6.1-10%)	25 (18-31)	890 (710-1100)
Dominica	2010	Male	15 to 19	58% (41-74%)	42% (26-59%)	2.1 (1.2-3.3)	5.9% (3.8-8.4%)	8.9% (5.3-13%)	0.26 (0.15-0.39)	25 (16-36)
Dominica	2010	Male	20 to 24	68% (50-82%)	32% (18-50%)	4 (2.5-5.8)	13% (9.1-17%)	16% (11-23%)	0.84 (0.55-1.2)	71 (50-97)
Dominica	2010	Male	25 to 29	71% (53-83%)	29% (17-47%)	4.8 (3-7)	13% (9.8-17%)	18% (12-24%)	0.89 (0.6-1.2)	70 (52-92)
Dominica	2010	Male	30 to 34	72% (54-85%)	28% (15-46%)	5 (3-7.3)	13% (10-17%)	17% (13-23%)	1.1 (0.76-1.4)	73 (54-91)
Dominica	2010	Male	35 to 39	71% (55-84%)	29% (16-45%)	4.7 (3-7.1)	11% (8.7-14%)	16% (11-20%)	0.82 (0.58-1.1)	57 (43-73)
Dominica	2010	Male	40 to 44	70% (55-84%)	30% (16-45%)	4.5 (2.7-6.8)	13% (9.5-16%)	16% (12-21%)	1.7 (1.2-2.2)	93 (68-120)
Dominica	2010	Male	45 to 49	69% (49-82%)	31% (18-51%)	4.2 (2.5-6.5)	12% (9.1-16%)	16% (11-20%)	2.1 (1.5-2.8)	100 (74-130)
Dominica	2010	Male	50 to 54	67% (49-82%)	33% (18-51%)	3.9 (2.1-6)	12% (8.3-15%)	15% (10-19%)	2.1 (1.5-2.7)	86 (61-110)
Dominica	2010	Male	55 to 59	65% (47-81%)	35% (19-53%)	3.5 (1.9-5.5)	11% (7-14%)	13% (8.5-17%)	2.3 (1.4-3.1)	80 (52-110)
Dominica	2010	Male	60 to 64	63% (44-80%)	37% (20-56%)	3.2 (1.7-5.4)	9.1% (5.5-13%)	11% (6.5-15%)	2.3 (1.3-3.3)	67 (40-96)
Dominica	2010	Male	65 to 69	62% (42-78%)	38% (22-58%)	2.9 (1.6-4.5)	8.1% (5.1-12%)	9.6% (6-14%)	2.3 (1.4-3.3)	57 (35-81)
Dominica	2010	Male	70 to 74	58% (40-74%)	42% (26-60%)	2.4 (1.3-4.1)	6.3% (3.4-9.8%)	7.2% (3.8-11%)	2.3 (1.2-3.6)	45 (24-71)
Dominica	2010	Male	75 to 79	53% (34-72%)	47% (28-66%)	2 (1-3.5)	5% (2.4-8.2%)	5.6% (2.6-9.1%)	2.1 (0.99-3.5)	32 (15-53)
Dominica	2010	Male	80 to 84	48% (29-66%)	52% (34-71%)	1.7 (0.86-2.7)	4% (2-6.6%)	4.4% (2.1-7.4%)	1.7 (0.79-2.7)	18 (8.9-30)
Dominica	2010	Male	85 to 89	43% (27-60%)	57% (40-73%)	1.5 (0.74-2.6)	3.4% (1.6-6.1%)	3.7% (1.7-6.8%)	1.1 (0.51-2)	8.9 (4.1-16)
Dominica	2010	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.5 (0.71-2.7)	3.1% (1.3-5.9%)	3.4% (1.3-6.6%)	0.51 (0.2-1)	2.8 (1.2-5.3)
Dominica	2010	Male	95 plus	46% (26-66%)	54% (34-74%)	1.6 (0.64-3.1)	3.3% (1.1-6.9%)	3.6% (0.97-7.8%)	0.18 (0.05-0.4)	0.69 (0.22-1.4)
Dominica	2016	Female	All Ages	35% (29-42%)	65% (58-71%)	1.4 (1-1.9)	2.7% (1.7-4.1%)	3.4% (1.6-5.8%)	9.5 (4.6-17)	280 (160-420)
Dominica	2016	Female	15 to 19	41% (27-57%)	59% (43-73%)	1.5 (0.76-2.6)	2.2% (1.3-3.4%)	5.3% (2.9-8.9%)	0.058 (0.031-0.1)	7.2 (4.4-11)
Dominica	2016	Female	20 to 24	47% (31-63%)	53% (37-69%)	2.3 (1.2-3.8)	4.1% (2.7-5.8%)	7.3% (4.2-11%)	0.14 (0.079-0.23)	16 (10-24)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	2016	Female	25 to 29	46% (30-63%)	54% (37-70%)	2.3 (1.3-3.9)	4.2% (2.9-5.9%)	7.2% (4.4-11%)	0.17 (0.098-0.26)	17 (11-25)
Dominica	2016	Female	30 to 34	44% (28-61%)	56% (39-72%)	2.2 (1.2-3.6)	3.9% (2.8-5.5%)	7.4% (4.8-11%)	0.18 (0.11-0.28)	16 (11-24)
Dominica	2016	Female	35 to 39	43% (29-61%)	57% (39-71%)	2.1 (1.1-3.3)	4% (2.8-5.8%)	7.6% (5.2-11%)	0.25 (0.16-0.38)	18 (12-26)
Dominica	2016	Female	40 to 44	42% (27-60%)	58% (40-73%)	1.9 (1-3.3)	4.3% (2.7-6.5%)	7.9% (5.2-12%)	0.39 (0.24-0.63)	23 (14-35)
Dominica	2016	Female	45 to 49	40% (25-58%)	60% (42-75%)	1.9 (0.96-3.3)	4.4% (2.6-7%)	7.5% (4.7-12%)	0.56 (0.32-0.9)	27 (15-45)
Dominica	2016	Female	50 to 54	37% (23-54%)	63% (46-77%)	1.7 (0.86-2.9)	4.5% (2.4-7.3%)	7% (4-11%)	0.77 (0.42-1.3)	32 (17-54)
Dominica	2016	Female	55 to 59	32% (20-49%)	68% (51-80%)	1.4 (0.7-2.5)	3.9% (1.9-6.6%)	5.9% (3-9.9%)	0.69 (0.35-1.2)	24 (12-43)
Dominica	2016	Female	60 to 64	29% (17-45%)	71% (55-83%)	1.2 (0.58-2.2)	3.3% (1.3-6.2%)	4.7% (2.1-8.4%)	0.77 (0.33-1.4)	23 (9-44)
Dominica	2016	Female	65 to 69	27% (16-41%)	73% (59-84%)	1.1 (0.55-1.9)	2.9% (1.1-5.5%)	3.9% (1.7-7.1%)	0.77 (0.3-1.4)	19 (6.7-36)
Dominica	2016	Female	70 to 74	24% (15-37%)	76% (63-85%)	0.91 (0.44-1.7)	2.5% (0.66-5.1%)	3.2% (0.99-6.4%)	0.88 (0.25-1.8)	17 (4.4-36)
Dominica	2016	Female	75 to 79	22% (13-35%)	78% (65-87%)	0.79 (0.34-1.5)	2.1% (0.46-4.7%)	2.6% (0.62-5.7%)	0.93 (0.21-2.2)	14 (2.8-33)
Dominica	2016	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.74 (0.34-1.3)	2% (0.53-4.5%)	2.4% (0.62-5.3%)	1 (0.26-2.2)	11 (2.7-25)
Dominica	2016	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.75 (0.34-1.4)	2% (0.53-4.5%)	2.3% (0.58-5.3%)	0.9 (0.23-2.1)	6.9 (1.8-16)
Dominica	2016	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.86 (0.37-1.7)	2.2% (0.48-5.3%)	2.5% (0.46-6.1%)	0.69 (0.13-1.7)	3.7 (0.78-8.9)
Dominica	2016	Female	95 plus	26% (13-43%)	74% (57-87%)	1 (0.36-2.2)	2.5% (0.35-6.5%)	2.8% (0.23-7.3%)	0.33 (0.029-0.86)	1.2 (0.16-3)
Dominica	2016	Male	All Ages	59% (50-66%)	41% (34-50%)	2.9 (2.2-3.9)	8.4% (6.5-10%)	8.5% (6-11%)	29 (19-40)	1000 (770-1300)
Dominica	2016	Male	15 to 19	58% (40-74%)	42% (26-60%)	2.2 (1.1-3.7)	6.5% (4.1-9.6%)	9.8% (5.6-15%)	0.33 (0.18-0.55)	30 (18-47)
Dominica	2016	Male	20 to 24	68% (50-81%)	32% (19-50%)	4.1 (2.4-6.5)	13% (9.4-18%)	17% (11-24%)	0.95 (0.57-1.4)	79 (52-110)
Dominica	2016	Male	25 to 29	71% (53-84%)	29% (16-47%)	4.9 (2.8-7.6)	14% (10-18%)	19% (12-25%)	1.1 (0.71-1.6)	86 (59-120)
Dominica	2016	Male	30 to 34	72% (53-85%)	28% (15-47%)	5.1 (3-7.8)	14% (10-17%)	18% (12-24%)	1.2 (0.82-1.7)	86 (61-120)
Dominica	2016	Male	35 to 39	71% (55-84%)	29% (16-45%)	4.8 (2.8-7.6)	12% (9.1-16%)	17% (12-22%)	1 (0.69-1.4)	67 (47-90)
Dominica	2016	Male	40 to 44	70% (54-84%)	30% (16-46%)	4.6 (2.6-7.4)	13% (9.3-16%)	17% (12-21%)	1.7 (1.1-2.4)	94 (66-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominica	2016	Male	45 to 49	69% (49-83%)	31% (17-51%)	4.3 (2.4-7.1)	12% (9-16%)	16% (11-20%)	2.2 (1.5-2.9)	100 (74-140)
Dominica	2016	Male	50 to 54	67% (49-82%)	33% (18-51%)	4 (2.1-6.5)	12% (8.3-15%)	15% (10-19%)	2.7 (1.8-3.8)	110 (78-150)
Dominica	2016	Male	55 to 59	65% (48-81%)	35% (19-52%)	3.6 (1.9-6.1)	11% (7-14%)	13% (8.4-18%)	2.7 (1.7-3.8)	97 (61-140)
Dominica	2016	Male	60 to 64	63% (43-80%)	37% (20-57%)	3.3 (1.7-5.7)	9.2% (5.3-13%)	11% (6.4-15%)	2.8 (1.6-4.1)	84 (48-120)
Dominica	2016	Male	65 to 69	62% (43-78%)	38% (22-57%)	2.9 (1.5-4.8)	8.2% (4.8-12%)	9.6% (5.6-14%)	2.9 (1.6-4.2)	70 (40-100)
Dominica	2016	Male	70 to 74	58% (40-74%)	42% (26-60%)	2.5 (1.2-4.3)	6.4% (3.3-10%)	7.3% (3.7-12%)	2.7 (1.3-4.5)	53 (26-85)
Dominica	2016	Male	75 to 79	53% (34-72%)	47% (28-66%)	2.1 (0.98-3.8)	5.1% (2.3-8.3%)	5.7% (2.5-9.5%)	2.5 (1.1-4.3)	37 (16-64)
Dominica	2016	Male	80 to 84	48% (30-66%)	52% (34-70%)	1.7 (0.84-3)	4.1% (1.9-7%)	4.6% (2-7.9%)	1.9 (0.83-3.4)	21 (9.4-37)
Dominica	2016	Male	85 to 89	43% (26-60%)	57% (40-74%)	1.5 (0.73-2.8)	3.5% (1.5-6.3%)	3.8% (1.6-7.2%)	1.2 (0.5-2.3)	9.7 (4.1-18)
Dominica	2016	Male	90 to 94	43% (28-61%)	57% (39-72%)	1.5 (0.71-2.9)	3.2% (1.3-6.2%)	3.5% (1.3-6.8%)	0.58 (0.21-1.1)	3.2 (1.2-6.2)
Dominica	2016	Male	95 plus	46% (26-66%)	54% (34-74%)	1.6 (0.63-3.2)	3.4% (1.1-7.2%)	3.7% (0.98-8.2%)	0.2 (0.05-0.44)	0.75 (0.24-1.6)
Dominican Republic	1990	Female	All Ages	47% (41-53%)	53% (47-59%)	0.84 (0.72-0.96)	1.4% (1.1-1.8%)	2.1% (1.3-2.9%)	340 (220-480)	15000 (12000-19000)
Dominican Republic	1990	Female	15 to 19	57% (45-68%)	43% (32-55%)	1.1 (0.71-1.6)	2.1% (1.3-3.1%)	4.3% (2.2-6.6%)	9.7 (4.7-17)	1000 (600-1600)
Dominican Republic	1990	Female	20 to 24	64% (52-74%)	36% (26-48%)	1.7 (1.2-2.4)	4.1% (2.8-5.6%)	7.1% (4.3-10%)	21 (12-32)	2100 (1400-2900)
Dominican Republic	1990	Female	25 to 29	63% (50-73%)	37% (27-50%)	1.6 (1.1-2.3)	4.1% (2.8-5.7%)	7% (4.1-10%)	22 (13-34)	1900 (1300-2700)
Dominican Republic	1990	Female	30 to 34	60% (47-70%)	40% (30-53%)	1.6 (1-2.2)	4% (2.6-5.5%)	7% (4.2-10%)	24 (14-35)	1800 (1100-2500)
Dominican Republic	1990	Female	35 to 39	57% (44-68%)	43% (32-56%)	1.5 (0.91-2.1)	3.8% (2.5-5.5%)	6.7% (4.1-9.9%)	25 (15-38)	1500 (970-2200)
Dominican Republic	1990	Female	40 to 44	54% (43-65%)	46% (35-57%)	1.3 (0.81-1.9)	3.6% (2.1-5.2%)	6.3% (3.8-9.5%)	24 (14-37)	1300 (780-1900)
Dominican Republic	1990	Female	45 to 49	51% (36-63%)	49% (37-64%)	1.3 (0.77-1.9)	3.5% (1.9-5.4%)	5.9% (3.3-9.1%)	26 (14-40)	1200 (670-1800)
Dominican Republic	1990	Female	50 to 54	47% (34-60%)	53% (40-66%)	1.1 (0.62-1.7)	3.2% (1.6-5.1%)	5.2% (2.7-8.2%)	30 (15-48)	1200 (600-1800)
Dominican Republic	1990	Female	55 to 59	43% (29-59%)	57% (41-71%)	0.87 (0.49-1.4)	2.7% (1.4-4.5%)	4.2% (2.2-6.9%)	29 (15-48)	950 (500-1600)
Dominican Republic	1990	Female	60 to 64	38% (24-52%)	62% (48-76%)	0.77 (0.41-1.3)	2.1% (0.86-3.8%)	3.1% (1.3-5.5%)	26 (11-48)	720 (280-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	1990	Female	65 to 69	32% (21-45%)	68% (55-79%)	0.71 (0.41-1.1)	1.9% (0.73-3.3%)	2.7% (1.1-4.6%)	26 (10-45)	580 (210-1000)
Dominican Republic	1990	Female	70 to 74	28% (18-41%)	72% (59-82%)	0.64 (0.35-1.1)	1.8% (0.6-3.3%)	2.4% (0.81-4.2%)	27 (8.8-48)	490 (150-870)
Dominican Republic	1990	Female	75 to 79	26% (16-38%)	74% (62-84%)	0.59 (0.28-1)	1.4% (0.21-2.7%)	1.7% (0.29-3.4%)	21 (3.4-42)	290 (44-600)
Dominican Republic	1990	Female	80 to 84	24% (14-37%)	76% (63-86%)	0.54 (0.28-0.91)	1.1% (0.098-2.3%)	1.4% (0.084-2.8%)	15 (0.94-30)	160 (14-310)
Dominican Republic	1990	Female	85 to 89	24% (15-37%)	76% (63-85%)	0.53 (0.29-0.91)	0.83% (-0.12-1.9%)	0.93% (-0.22-2.2%)	8.7 (-2.1-21)	69 (-9.4-160)
Dominican Republic	1990	Female	90 to 94	27% (17-40%)	73% (60-83%)	0.59 (0.29-1)	0.68% (-0.33-2.1%)	0.68% (-0.56-2.3%)	3.6 (-2.9-12)	22 (-11-66)
Dominican Republic	1990	Female	95 plus	32% (17-50%)	68% (50-83%)	0.69 (0.3-1.3)	0.62% (-0.7-2.5%)	0.54% (-1-2.7%)	1 (-2-5.3)	4.8 (-5.2-19)
Dominican Republic	1990	Male	All Ages	70% (64-76%)	30% (24-36%)	1.4 (1.2-1.5)	3.6% (2.9-4.3%)	4.6% (3.4-5.8%)	980 (730-1300)	47000 (38000-57000)
Dominican Republic	1990	Male	15 to 19	75% (65-83%)	25% (17-35%)	1.2 (0.82-1.8)	5.2% (3.4-7.2%)	7.5% (4.3-11%)	31 (18-45)	2900 (1900-4000)
Dominican Republic	1990	Male	20 to 24	85% (77-90%)	15% (10-23%)	2.2 (1.6-2.9)	10% (7.9-13%)	13% (8.9-18%)	73 (50-98)	6400 (4700-8300)
Dominican Republic	1990	Male	25 to 29	86% (78-90%)	14% (9.7-22%)	2.5 (1.9-3.4)	11% (8.3-14%)	14% (10-18%)	80 (57-110)	6400 (4800-8100)
Dominican Republic	1990	Male	30 to 34	86% (79-90%)	14% (9.8-21%)	2.9 (2-3.8)	11% (8.6-14%)	15% (11-20%)	85 (60-110)	5900 (4400-7600)
Dominican Republic	1990	Male	35 to 39	85% (77-90%)	15% (10-23%)	3 (2-4.1)	11% (8.5-15%)	15% (11-20%)	87 (61-110)	5300 (3900-6900)
Dominican Republic	1990	Male	40 to 44	83% (76-89%)	17% (11-24%)	2.8 (1.8-3.9)	11% (7.5-14%)	14% (9.9-19%)	82 (56-110)	4400 (3100-5900)
Dominican Republic	1990	Male	45 to 49	81% (71-88%)	19% (12-29%)	2.3 (1.5-3.3)	8.8% (6.2-12%)	12% (8-16%)	73 (48-100)	3400 (2400-4700)
Dominican Republic	1990	Male	50 to 54	79% (70-86%)	21% (14-30%)	2 (1.2-2.9)	7.2% (4.7-10%)	9.5% (6-13%)	74 (46-100)	3000 (2000-4300)
Dominican Republic	1990	Male	55 to 59	77% (67-85%)	23% (15-33%)	1.9 (1.2-2.9)	6.3% (4-9.2%)	8.3% (5.2-12%)	77 (48-110)	2700 (1700-4000)
Dominican Republic	1990	Male	60 to 64	76% (64-85%)	24% (15-36%)	1.8 (1.1-2.9)	5.5% (3.1-8.7%)	7% (3.7-11%)	81 (43-130)	2400 (1300-3800)
Dominican Republic	1990	Male	65 to 69	76% (64-86%)	24% (14-36%)	1.7 (1.1-2.4)	4.6% (2.6-6.9%)	5.7% (3.1-8.7%)	74 (40-120)	1800 (1000-2800)
Dominican Republic	1990	Male	70 to 74	73% (59-84%)	27% (16-41%)	1.4 (0.83-2.1)	3.7% (1.7-6%)	4.5% (2-7.3%)	64 (27-100)	1200 (560-2000)
Dominican Republic	1990	Male	75 to 79	68% (51-81%)	32% (19-49%)	1.2 (0.71-2)	2.9% (1-5.1%)	3.3% (1.1-6%)	48 (15-88)	710 (240-1300)
Dominican Republic	1990	Male	80 to 84	62% (44-76%)	38% (24-56%)	1 (0.61-1.6)	2.1% (0.63-3.7%)	2.4% (0.54-4.3%)	29 (6.5-53)	320 (95-570)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	1990	Male	85 to 89	58% (40-73%)	42% (27-60%)	0.87 (0.48-1.4)	1.5% (0.33-2.8%)	1.6% (0.16-3.2%)	15 (1.6-29)	130 (26-230)
Dominican Republic	1990	Male	90 to 94	58% (42-73%)	42% (27-58%)	0.85 (0.45-1.4)	0.96% (-0.26-2.4%)	0.88% (-0.63-2.5%)	4.2 (-3-12)	27 (-7.4-67)
Dominican Republic	1990	Male	95 plus	61% (40-79%)	39% (21-60%)	0.88 (0.41-1.6)	0.73% (-0.7-2.4%)	0.46% (-1.3-2.5%)	0.68 (-1.9-3.7)	4.3 (-4.1-14)
Dominican Republic	1995	Female	All Ages	47% (41-53%)	53% (47-59%)	0.99 (0.85-1.1)	1.7% (1.3-2.2%)	2.7% (1.8-3.7%)	460 (300-630)	19000 (14000-23000)
Dominican Republic	1995	Female	15 to 19	57% (46-68%)	43% (32-54%)	1.3 (0.84-1.9)	2.4% (1.5-3.4%)	5.3% (3-8.1%)	11 (6.2-18)	1200 (740-1700)
Dominican Republic	1995	Female	20 to 24	64% (53-74%)	36% (26-47%)	2 (1.3-2.8)	4.1% (2.9-5.5%)	7.1% (4.4-10%)	22 (13-32)	2200 (1500-3100)
Dominican Republic	1995	Female	25 to 29	63% (50-73%)	37% (27-50%)	1.9 (1.3-2.7)	3.9% (2.7-5.3%)	6% (3.8-8.9%)	24 (15-34)	2100 (1500-2900)
Dominican Republic	1995	Female	30 to 34	60% (46-71%)	40% (29-54%)	1.8 (1.2-2.6)	3.8% (2.6-5.2%)	6.2% (3.9-9%)	27 (17-38)	2000 (1400-2800)
Dominican Republic	1995	Female	35 to 39	57% (43-69%)	43% (31-57%)	1.7 (1-2.5)	3.7% (2.5-5.3%)	6.4% (4-9.4%)	30 (19-43)	1900 (1300-2600)
Dominican Republic	1995	Female	40 to 44	54% (42-65%)	46% (35-58%)	1.6 (0.95-2.2)	3.8% (2.5-5.4%)	6.6% (4.2-9.6%)	32 (20-46)	1700 (1100-2400)
Dominican Republic	1995	Female	45 to 49	50% (36-63%)	50% (37-64%)	1.5 (0.87-2.2)	3.8% (2.3-5.6%)	6.4% (3.8-9.6%)	33 (20-49)	1500 (920-2300)
Dominican Republic	1995	Female	50 to 54	46% (34-60%)	54% (40-66%)	1.3 (0.72-2)	3.7% (2-5.7%)	5.9% (3.2-9.1%)	34 (19-55)	1300 (730-2100)
Dominican Republic	1995	Female	55 to 59	43% (29-59%)	57% (41-71%)	1 (0.58-1.7)	3.2% (1.8-5%)	5% (2.9-7.8%)	37 (21-58)	1200 (700-1900)
Dominican Republic	1995	Female	60 to 64	38% (24-53%)	62% (47-76%)	0.9 (0.49-1.5)	2.6% (1.2-4.5%)	3.9% (2-6.6%)	36 (17-62)	1000 (470-1700)
Dominican Republic	1995	Female	65 to 69	32% (20-44%)	68% (56-80%)	0.82 (0.47-1.3)	2.4% (1.1-3.9%)	3.4% (1.7-5.5%)	37 (18-60)	850 (410-1400)
Dominican Republic	1995	Female	70 to 74	28% (18-40%)	72% (60-82%)	0.75 (0.4-1.2)	2.4% (0.93-4%)	3.2% (1.3-5.4%)	40 (16-68)	740 (290-1300)
Dominican Republic	1995	Female	75 to 79	26% (16-39%)	74% (61-84%)	0.7 (0.34-1.2)	1.9% (0.62-3.4%)	2.5% (0.83-4.5%)	34 (11-61)	480 (160-890)
Dominican Republic	1995	Female	80 to 84	24% (15-37%)	76% (63-85%)	0.63 (0.33-1)	1.7% (0.56-2.9%)	2.1% (0.66-3.7%)	29 (9.4-53)	310 (100-550)
Dominican Republic	1995	Female	85 to 89	25% (15-37%)	75% (63-85%)	0.62 (0.33-1)	1.4% (0.28-2.7%)	1.6% (0.24-3.2%)	19 (3-40)	150 (30-300)
Dominican Republic	1995	Female	90 to 94	27% (17-40%)	73% (60-83%)	0.7 (0.35-1.2)	1.2% (0.056-3%)	1.4% (-0.08-3.4%)	9.9 (-0.58-24)	55 (2.5-130)
Dominican Republic	1995	Female	95 plus	32% (18-51%)	68% (49-82%)	0.81 (0.36-1.5)	1.2% (-0.34-3.5%)	1.2% (-0.59-3.9%)	3.4 (-1.6-11)	13 (-3.7-38)
Dominican Republic	1995	Male	All Ages	70% (64-75%)	30% (25-36%)	1.6 (1.4-1.8)	4.2% (3.4-5.1%)	5.3% (4-6.7%)	1200 (910-1500)	56000 (45000-68000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	1995	Male	15 to 19	75% (65-83%)	25% (17-35%)	1.4 (0.93-2)	5.6% (3.7-7.6%)	8.6% (5-13%)	33 (19-48)	3000 (2000-4200)
Dominican Republic	1995	Male	20 to 24	85% (77-90%)	15% (10-23%)	2.6 (1.8-3.3)	10% (7.6-14%)	13% (9-18%)	76 (52-100)	6700 (4900-8600)
Dominican Republic	1995	Male	25 to 29	86% (78-90%)	14% (9.6-22%)	2.9 (2.1-3.9)	10% (7.4-13%)	12% (8-17%)	85 (60-110)	6800 (5100-8800)
Dominican Republic	1995	Male	30 to 34	86% (78-90%)	14% (9.6-22%)	3.3 (2.3-4.4)	10% (7.6-13%)	13% (8.9-17%)	98 (70-130)	6900 (5000-8900)
Dominican Republic	1995	Male	35 to 39	85% (77-90%)	15% (10-23%)	3.4 (2.3-4.7)	10% (7.6-13%)	13% (9.2-18%)	100 (74-140)	6400 (4800-8300)
Dominican Republic	1995	Male	40 to 44	83% (76-89%)	17% (11-24%)	3.2 (2.1-4.4)	10% (7.1-13%)	13% (8.9-18%)	100 (71-140)	5600 (3900-7400)
Dominican Republic	1995	Male	45 to 49	81% (71-87%)	19% (13-29%)	2.7 (1.7-3.7)	9% (6.3-12%)	12% (7.9-16%)	94 (62-130)	4400 (3100-6100)
Dominican Republic	1995	Male	50 to 54	79% (71-85%)	21% (15-29%)	2.3 (1.4-3.4)	7.7% (5.2-11%)	10% (6.6-14%)	86 (55-120)	3500 (2300-4900)
Dominican Republic	1995	Male	55 to 59	77% (67-85%)	23% (15-33%)	2.2 (1.4-3.3)	7.1% (4.5-10%)	9.1% (5.8-14%)	97 (61-150)	3400 (2200-5000)
Dominican Republic	1995	Male	60 to 64	76% (64-85%)	24% (15-36%)	2.1 (1.3-3.2)	6.3% (3.7-9.9%)	8% (4.4-13%)	110 (61-170)	3200 (1800-5000)
Dominican Republic	1995	Male	65 to 69	76% (63-86%)	24% (14-37%)	1.9 (1.2-2.7)	5.4% (3.2-8.1%)	6.7% (3.8-10%)	100 (60-160)	2500 (1500-3800)
Dominican Republic	1995	Male	70 to 74	73% (59-84%)	27% (16-41%)	1.6 (0.98-2.4)	4.3% (2.1-6.9%)	5.3% (2.6-8.3%)	91 (43-150)	1700 (850-2700)
Dominican Republic	1995	Male	75 to 79	68% (51-81%)	32% (19-49%)	1.4 (0.8-2.2)	3.4% (1.4-5.9%)	4% (1.6-6.9%)	72 (29-130)	1100 (430-1800)
Dominican Republic	1995	Male	80 to 84	62% (44-76%)	38% (24-56%)	1.2 (0.68-1.8)	2.6% (1.1-4.4%)	3% (1.2-5.1%)	52 (20-89)	570 (230-960)
Dominican Republic	1995	Male	85 to 89	57% (40-73%)	43% (27-60%)	0.98 (0.53-1.6)	1.9% (0.72-3.4%)	2.1% (0.64-3.8%)	28 (8.4-50)	220 (81-390)
Dominican Republic	1995	Male	90 to 94	58% (41-73%)	42% (27-59%)	0.96 (0.5-1.6)	1.3% (0.11-2.8%)	1.3% (-0.12-3.1%)	8.9 (-0.82-21)	54 (4.6-120)
Dominican Republic	1995	Male	95 plus	61% (41-79%)	39% (21-59%)	1 (0.45-1.8)	1.1% (-0.32-3.1%)	0.95% (-0.83-3.2%)	2 (-1.7-6.7)	9.5 (-2.7-25)
Dominican Republic	2000	Female	All Ages	44% (39-49%)	56% (51-61%)	1 (0.88-1.2)	1.7% (1.4-2.1%)	2.6% (1.8-3.6%)	480 (320-660)	19000 (14000-23000)
Dominican Republic	2000	Female	15 to 19	58% (48-67%)	42% (33-52%)	1.4 (0.94-2)	2.5% (1.6-3.6%)	6% (3.5-8.8%)	13 (7.4-19)	1300 (840-1900)
Dominican Republic	2000	Female	20 to 24	63% (53-72%)	37% (28-47%)	2.1 (1.5-2.9)	3.9% (2.8-5.1%)	6.5% (4-9.4%)	20 (13-28)	2100 (1500-2800)
Dominican Republic	2000	Female	25 to 29	60% (50-69%)	40% (31-50%)	2 (1.4-2.7)	3.3% (2.4-4.3%)	4.5% (2.7-6.8%)	21 (14-28)	2000 (1400-2700)
Dominican Republic	2000	Female	30 to 34	56% (46-65%)	44% (35-54%)	1.8 (1.3-2.6)	2.9% (2.1-3.9%)	4% (2.6-5.9%)	23 (15-32)	1900 (1300-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	2000	Female	35 to 39	53% (43-63%)	47% (37-57%)	1.6 (1.1-2.4)	2.9% (2.1-4.1%)	4.5% (2.7-6.9%)	26 (17-37)	1800 (1200-2400)
Dominican Republic	2000	Female	40 to 44	49% (40-59%)	51% (41-60%)	1.5 (0.95-2.1)	3% (2-4.3%)	4.8% (3.1-7%)	29 (19-42)	1600 (1100-2300)
Dominican Republic	2000	Female	45 to 49	45% (34-56%)	55% (44-66%)	1.4 (0.9-2.1)	3.1% (2-4.6%)	5.2% (3.3-7.6%)	32 (20-48)	1500 (980-2300)
Dominican Republic	2000	Female	50 to 54	41% (30-53%)	59% (47-70%)	1.2 (0.68-1.9)	3.1% (1.9-4.8%)	5.2% (3.1-7.8%)	32 (19-51)	1300 (780-2000)
Dominican Republic	2000	Female	55 to 59	36% (23-51%)	64% (49-77%)	0.89 (0.51-1.5)	2.8% (1.7-4.3%)	4.5% (2.8-6.8%)	31 (19-49)	1100 (650-1700)
Dominican Republic	2000	Female	60 to 64	31% (19-45%)	69% (55-81%)	0.77 (0.41-1.3)	2.5% (1.3-4%)	3.8% (2.1-6.1%)	35 (19-56)	990 (530-1600)
Dominican Republic	2000	Female	65 to 69	27% (17-38%)	73% (62-83%)	0.74 (0.42-1.2)	2.4% (1.3-3.7%)	3.5% (1.9-5.5%)	40 (22-63)	920 (490-1500)
Dominican Republic	2000	Female	70 to 74	25% (16-36%)	75% (64-84%)	0.72 (0.39-1.2)	2.4% (1.1-3.9%)	3.4% (1.7-5.4%)	45 (22-74)	830 (400-1400)
Dominican Republic	2000	Female	75 to 79	24% (15-36%)	76% (64-85%)	0.72 (0.37-1.2)	2% (0.73-3.7%)	2.7% (0.99-4.9%)	40 (14-72)	580 (210-1000)
Dominican Republic	2000	Female	80 to 84	24% (15-36%)	76% (64-85%)	0.69 (0.37-1.1)	1.9% (0.68-3.3%)	2.3% (0.82-4.2%)	40 (14-72)	430 (150-770)
Dominican Republic	2000	Female	85 to 89	24% (15-37%)	76% (63-85%)	0.68 (0.35-1.1)	1.6% (0.5-3.1%)	1.9% (0.49-3.7%)	28 (7.3-57)	220 (66-430)
Dominican Republic	2000	Female	90 to 94	27% (17-40%)	73% (60-83%)	0.75 (0.38-1.3)	1.5% (0.27-3.4%)	1.7% (0.19-4%)	16 (1.7-36)	86 (15-190)
Dominican Republic	2000	Female	95 plus	32% (17-50%)	68% (50-83%)	0.88 (0.39-1.6)	1.5% (-0.12-4%)	1.6% (-0.42-4.7%)	5.6 (-1.5-16)	21 (-1.7-56)
Dominican Republic	2000	Male	All Ages	73% (69-77%)	27% (23-31%)	1.9 (1.7-2.1)	4.5% (3.7-5.5%)	5.4% (4.1-6.9%)	1400 (1000-1800)	62000 (50000-75000)
Dominican Republic	2000	Male	15 to 19	79% (73-85%)	21% (15-27%)	1.7 (1.2-2.4)	6% (4.1-8.3%)	10% (6.2-14%)	36 (22-53)	3400 (2300-4700)
Dominican Republic	2000	Male	20 to 24	87% (83-91%)	13% (9.4-17%)	3 (2.2-3.9)	11% (7.9-14%)	14% (9.7-19%)	79 (55-110)	7000 (5000-9000)
Dominican Republic	2000	Male	25 to 29	88% (85-91%)	12% (8.8-15%)	3.4 (2.6-4.4)	9.4% (6.7-12%)	11% (6.9-16%)	87 (61-110)	7100 (5200-9000)
Dominican Republic	2000	Male	30 to 34	88% (84-91%)	12% (9.2-16%)	3.8 (2.9-4.9)	8.6% (6.3-11%)	10% (6.8-14%)	100 (73-130)	7200 (5300-9200)
Dominican Republic	2000	Male	35 to 39	87% (82-90%)	13% (9.7-18%)	3.9 (2.7-5.2)	8.9% (6.5-12%)	11% (7.2-15%)	110 (79-150)	7100 (5200-9300)
Dominican Republic	2000	Male	40 to 44	85% (80-89%)	15% (11-20%)	3.6 (2.4-4.9)	8.6% (6.1-11%)	11% (7.1-14%)	120 (80-160)	6500 (4600-8400)
Dominican Republic	2000	Male	45 to 49	82% (76-87%)	18% (13-24%)	3 (2-4.1)	8% (5.6-11%)	10% (6.9-14%)	110 (74-150)	5300 (3700-7200)
Dominican Republic	2000	Male	50 to 54	79% (73-84%)	21% (16-27%)	2.5 (1.6-3.5)	7.2% (4.7-10%)	9.2% (5.9-13%)	97 (62-140)	4100 (2700-5800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	2000	Male	55 to 59	75% (68-82%)	25% (18-32%)	2.3 (1.5-3.3)	6.8% (4.5-10%)	8.8% (5.6-13%)	97 (61-150)	3400 (2200-5100)
Dominican Republic	2000	Male	60 to 64	74% (63-83%)	26% (17-37%)	2.2 (1.4-3.3)	6.2% (3.7-9.7%)	7.9% (4.5-12%)	110 (66-180)	3400 (2000-5300)
Dominican Republic	2000	Male	65 to 69	72% (59-83%)	28% (17-41%)	1.9 (1.2-2.6)	5.2% (3.2-7.8%)	6.5% (3.9-9.7%)	120 (69-180)	2800 (1700-4200)
Dominican Republic	2000	Male	70 to 74	68% (53-80%)	32% (20-47%)	1.5 (0.91-2.3)	4.2% (2.3-6.7%)	5.2% (2.8-8.2%)	100 (56-160)	2000 (1100-3100)
Dominican Republic	2000	Male	75 to 79	63% (48-78%)	37% (22-52%)	1.4 (0.79-2.2)	3.5% (1.6-5.9%)	4.1% (1.9-7.1%)	86 (40-150)	1300 (600-2200)
Dominican Republic	2000	Male	80 to 84	60% (42-75%)	40% (25-58%)	1.2 (0.71-1.8)	2.8% (1.3-4.7%)	3.2% (1.4-5.5%)	70 (30-120)	770 (350-1300)
Dominican Republic	2000	Male	85 to 89	56% (39-72%)	44% (28-61%)	1 (0.56-1.7)	2.1% (0.91-3.6%)	2.3% (0.87-4.1%)	39 (15-67)	310 (130-540)
Dominican Republic	2000	Male	90 to 94	56% (40-72%)	44% (28-60%)	1 (0.54-1.7)	1.5% (0.39-3.2%)	1.6% (0.12-3.5%)	14 (1-30)	81 (20-170)
Dominican Republic	2000	Male	95 plus	59% (38-77%)	41% (23-62%)	1.1 (0.48-1.8)	1.4% (0.029-3.5%)	1.3% (-0.42-3.7%)	3.3 (-1.1-9.7)	15 (0.31-36)
Dominican Republic	2005	Female	All Ages	44% (40-48%)	56% (52-60%)	1.1 (0.91-1.2)	1.6% (1.3-2.1%)	2.4% (1.6-3.4%)	490 (310-710)	19000 (14000-24000)
Dominican Republic	2005	Female	15 to 19	59% (52-67%)	41% (33-48%)	1.5 (1-2.1)	2.5% (1.7-3.4%)	5.8% (3.6-8.4%)	13 (8.2-20)	1400 (880-1900)
Dominican Republic	2005	Female	20 to 24	63% (57-70%)	37% (30-43%)	2.2 (1.6-3)	3.8% (2.7-5%)	6.5% (4-9.4%)	21 (14-30)	2200 (1600-3000)
Dominican Republic	2005	Female	25 to 29	61% (54-67%)	39% (33-46%)	2 (1.5-2.7)	3.1% (2.2-4.1%)	4% (2.4-6.3%)	21 (14-29)	2000 (1500-2700)
Dominican Republic	2005	Female	30 to 34	57% (49-63%)	43% (37-51%)	1.9 (1.3-2.5)	2.6% (1.8-3.6%)	3.4% (2.1-5.1%)	22 (14-32)	1900 (1300-2600)
Dominican Republic	2005	Female	35 to 39	52% (45-59%)	48% (41-55%)	1.7 (1.2-2.4)	2.5% (1.7-3.5%)	3.4% (2.1-5.3%)	24 (16-35)	1700 (1200-2400)
Dominican Republic	2005	Female	40 to 44	48% (40-55%)	52% (45-60%)	1.5 (1-2)	2.5% (1.7-3.5%)	3.8% (2.5-5.5%)	28 (19-41)	1700 (1100-2400)
Dominican Republic	2005	Female	45 to 49	43% (34-51%)	57% (49-66%)	1.3 (0.88-1.9)	2.6% (1.7-3.9%)	4.2% (2.7-6.3%)	32 (19-48)	1600 (990-2300)
Dominican Republic	2005	Female	50 to 54	38% (28-49%)	62% (51-72%)	1.1 (0.66-1.7)	2.6% (1.5-4.1%)	4.3% (2.5-6.5%)	32 (18-51)	1300 (750-2000)
Dominican Republic	2005	Female	55 to 59	32% (21-47%)	68% (53-79%)	0.81 (0.46-1.4)	2.3% (1.4-3.7%)	3.8% (2.4-5.8%)	29 (17-47)	1000 (610-1600)
Dominican Republic	2005	Female	60 to 64	27% (17-40%)	73% (60-83%)	0.7 (0.38-1.2)	2.1% (1.1-3.5%)	3.3% (1.8-5.4%)	30 (15-49)	860 (460-1400)
Dominican Republic	2005	Female	65 to 69	25% (16-36%)	75% (64-84%)	0.7 (0.4-1.1)	2.1% (1.1-3.4%)	3.2% (1.8-5%)	38 (20-61)	900 (470-1400)
Dominican Republic	2005	Female	70 to 74	24% (15-35%)	76% (65-85%)	0.7 (0.38-1.2)	2.1% (0.95-3.6%)	3.1% (1.5-5.1%)	46 (21-80)	870 (400-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	2005	Female	75 to 79	24% (15-35%)	76% (65-85%)	0.72 (0.37-1.2)	1.8% (0.56-3.5%)	2.5% (0.75-4.7%)	43 (13-83)	630 (180-1200)
Dominican Republic	2005	Female	80 to 84	24% (14-36%)	76% (64-86%)	0.69 (0.38-1.1)	1.7% (0.58-3.2%)	2.2% (0.75-4.1%)	46 (15-84)	490 (160-910)
Dominican Republic	2005	Female	85 to 89	24% (14-37%)	76% (63-86%)	0.68 (0.36-1.1)	1.5% (0.46-3%)	1.8% (0.53-3.6%)	35 (9.3-68)	270 (81-530)
Dominican Republic	2005	Female	90 to 94	27% (16-39%)	73% (61-84%)	0.76 (0.38-1.3)	1.5% (0.25-3.4%)	1.7% (0.16-4.1%)	20 (1.7-47)	110 (17-240)
Dominican Republic	2005	Female	95 plus	31% (17-50%)	69% (50-83%)	0.88 (0.38-1.6)	1.5% (-0.14-4.1%)	1.6% (-0.45-4.7%)	6.9 (-2-20)	27 (-2.6-71)
Dominican Republic	2005	Male	All Ages	74% (70-78%)	26% (22-30%)	2 (1.8-2.2)	4.4% (3.6-5.4%)	5% (3.8-6.5%)	1500 (1100-1900)	65000 (52000-80000)
Dominican Republic	2005	Male	15 to 19	79% (73-85%)	21% (15-27%)	1.8 (1.2-2.4)	5.9% (4.1-7.9%)	9.6% (6.1-14%)	37 (22-54)	3500 (2400-4800)
Dominican Republic	2005	Male	20 to 24	87% (83-91%)	13% (9.2-17%)	3.1 (2.3-3.9)	11% (7.8-14%)	14% (9.4-19%)	83 (56-110)	7400 (5300-9600)
Dominican Republic	2005	Male	25 to 29	88% (84-91%)	12% (8.7-16%)	3.5 (2.6-4.5)	9.2% (6.5-12%)	11% (6.7-16%)	89 (62-120)	7300 (5400-9300)
Dominican Republic	2005	Male	30 to 34	88% (84-91%)	12% (9-16%)	3.9 (2.9-5)	8.1% (5.9-11%)	9.3% (6.1-13%)	100 (72-140)	7300 (5400-9500)
Dominican Republic	2005	Male	35 to 39	87% (83-90%)	13% (9.5-17%)	4 (2.8-5.3)	7.7% (5.5-10%)	9% (5.8-13%)	110 (76-160)	7200 (5100-9700)
Dominican Republic	2005	Male	40 to 44	86% (81-89%)	14% (11-19%)	3.7 (2.5-5)	7.7% (5.4-10%)	9.1% (6.1-13%)	120 (83-170)	6900 (4800-9200)
Dominican Republic	2005	Male	45 to 49	83% (78-88%)	17% (12-22%)	3.1 (2.2-4.3)	7.2% (4.9-9.9%)	8.7% (5.7-12%)	120 (80-170)	5900 (4000-8100)
Dominican Republic	2005	Male	50 to 54	82% (76-86%)	18% (14-24%)	2.7 (1.8-3.9)	6.8% (4.3-9.8%)	8.5% (5.2-13%)	110 (66-170)	4800 (2900-6900)
Dominican Republic	2005	Male	55 to 59	79% (72-85%)	21% (15-28%)	2.5 (1.7-3.6)	6.7% (4.2-10%)	8.6% (5.2-13%)	110 (66-170)	3900 (2400-5900)
Dominican Republic	2005	Male	60 to 64	75% (64-84%)	25% (16-36%)	2.3 (1.4-3.5)	5.9% (3.3-9.2%)	7.4% (4.1-12%)	110 (61-180)	3200 (1800-5200)
Dominican Republic	2005	Male	65 to 69	72% (59-83%)	28% (17-41%)	1.9 (1.2-2.7)	4.8% (2.8-7.3%)	6% (3.4-9.2%)	110 (63-180)	2800 (1600-4300)
Dominican Republic	2005	Male	70 to 74	68% (53-81%)	32% (19-47%)	1.5 (0.92-2.3)	4% (2.2-6.4%)	4.9% (2.7-8%)	110 (57-180)	2100 (1100-3400)
Dominican Republic	2005	Male	75 to 79	63% (48-77%)	37% (23-52%)	1.4 (0.81-2.2)	3.3% (1.4-5.7%)	3.9% (1.6-6.8%)	94 (40-170)	1400 (620-2500)
Dominican Republic	2005	Male	80 to 84	59% (41-74%)	41% (26-59%)	1.2 (0.7-1.8)	2.5% (1.1-4.4%)	2.9% (1.2-5.2%)	80 (33-140)	890 (380-1600)
Dominican Republic	2005	Male	85 to 89	55% (39-71%)	45% (29-61%)	1 (0.56-1.7)	1.9% (0.73-3.3%)	2.1% (0.68-3.7%)	43 (14-79)	350 (130-630)
Dominican Republic	2005	Male	90 to 94	56% (40-71%)	44% (29-60%)	1 (0.53-1.7)	1.4% (0.25-3%)	1.4% (-0.03-3.3%)	15 (-0.32-35)	90 (17-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	2005	Male	95 plus	58% (37-77%)	42% (23-63%)	1 (0.46-1.9)	1.3% (-0.087-3.3%)	1.1% (-0.53-3.5%)	3.6 (-1.7-11)	17 (-1.2-42)
Dominican Republic	2010	Female	All Ages	44% (41-48%)	56% (52-59%)	1.1 (0.93-1.3)	1.7% (1.2-2.2%)	2.4% (1.3-3.5%)	520 (280-800)	20000 (14000-26000)
Dominican Republic	2010	Female	15 to 19	58% (51-65%)	42% (35-49%)	1.5 (1-2.1)	2.4% (1.6-3.3%)	5.5% (3.3-8%)	12 (7.3-19)	1300 (860-1900)
Dominican Republic	2010	Female	20 to 24	63% (57-69%)	37% (31-43%)	2.2 (1.6-3)	3.8% (2.7-5.1%)	6.9% (4.3-10%)	19 (12-28)	2200 (1500-3000)
Dominican Republic	2010	Female	25 to 29	61% (56-67%)	39% (33-44%)	2.1 (1.5-2.8)	3.3% (2.3-4.3%)	4.8% (2.8-7.4%)	20 (13-29)	2100 (1500-2800)
Dominican Republic	2010	Female	30 to 34	57% (51-62%)	43% (38-49%)	1.9 (1.3-2.6)	2.7% (1.9-3.8%)	3.8% (2.3-5.7%)	21 (13-30)	1900 (1300-2600)
Dominican Republic	2010	Female	35 to 39	51% (45-57%)	49% (43-55%)	1.7 (1.1-2.3)	2.5% (1.7-3.5%)	3.6% (2.2-5.5%)	22 (14-33)	1700 (1100-2400)
Dominican Republic	2010	Female	40 to 44	46% (41-52%)	54% (48-59%)	1.4 (0.99-2)	2.4% (1.6-3.4%)	3.8% (2.4-5.5%)	26 (17-39)	1600 (1100-2300)
Dominican Republic	2010	Female	45 to 49	41% (34-48%)	59% (52-66%)	1.3 (0.86-1.8)	2.4% (1.5-3.7%)	4.1% (2.6-6.2%)	30 (18-47)	1600 (950-2300)
Dominican Republic	2010	Female	50 to 54	41% (31-54%)	59% (46-69%)	1.2 (0.75-1.9)	2.6% (1.4-4.1%)	4.4% (2.4-6.9%)	35 (18-58)	1500 (820-2400)
Dominican Republic	2010	Female	55 to 59	42% (28-58%)	58% (42-72%)	1.1 (0.64-1.8)	2.4% (1.2-4.1%)	4.1% (2.2-6.8%)	36 (18-61)	1300 (640-2100)
Dominican Republic	2010	Female	60 to 64	37% (23-51%)	63% (49-77%)	1 (0.55-1.7)	2.1% (0.82-3.8%)	3.3% (1.4-6%)	34 (14-65)	1000 (390-1900)
Dominican Republic	2010	Female	65 to 69	31% (20-43%)	69% (57-80%)	0.92 (0.53-1.5)	2.1% (0.89-3.6%)	3.1% (1.4-5.4%)	39 (17-70)	920 (390-1600)
Dominican Republic	2010	Female	70 to 74	27% (18-40%)	73% (60-82%)	0.83 (0.45-1.4)	2% (0.67-3.8%)	2.9% (1-5.2%)	49 (17-93)	940 (320-1800)
Dominican Republic	2010	Female	75 to 79	25% (15-37%)	75% (63-85%)	0.77 (0.38-1.3)	1.8% (0.4-3.5%)	2.4% (0.52-4.7%)	49 (11-100)	730 (160-1500)
Dominican Republic	2010	Female	80 to 84	23% (14-35%)	77% (65-86%)	0.7 (0.37-1.2)	1.6% (0.43-2.9%)	2% (0.51-3.7%)	52 (13-100)	570 (150-1100)
Dominican Republic	2010	Female	85 to 89	24% (14-36%)	76% (64-86%)	0.68 (0.36-1.2)	1.4% (0.42-2.9%)	1.7% (0.41-3.4%)	41 (9.2-85)	320 (89-650)
Dominican Republic	2010	Female	90 to 94	26% (16-40%)	74% (60-84%)	0.76 (0.37-1.4)	1.4% (0.16-3.2%)	1.5% (-0.0021-3.8%)	23 (-0.031-58)	130 (14-320)
Dominican Republic	2010	Female	95 plus	31% (17-49%)	69% (51-83%)	0.88 (0.37-1.7)	1.4% (-0.21-3.9%)	1.5% (-0.55-4.6%)	8.4 (-3.2-26)	34 (-5-93)
Dominican Republic	2010	Male	All Ages	73% (69-78%)	27% (22-31%)	2 (1.8-2.3)	4.7% (3.7-5.8%)	5.1% (3.6-6.9%)	1600 (1100-2100)	69000 (52000-87000)
Dominican Republic	2010	Male	15 to 19	76% (69-83%)	24% (17-31%)	1.7 (1.1-2.4)	5.7% (3.8-7.8%)	9.5% (5.5-14%)	36 (21-54)	3500 (2300-4900)
Dominican Republic	2010	Male	20 to 24	86% (81-90%)	14% (9.7-19%)	3 (2.2-4)	11% (7.9-14%)	15% (9.8-21%)	82 (54-110)	7500 (5200-9800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	2010	Male	25 to 29	87% (82-91%)	13% (9.2-18%)	3.5 (2.5-4.5)	9.8% (6.8-13%)	12% (7.4-18%)	92 (60-130)	7600 (5300-9900)
Dominican Republic	2010	Male	30 to 34	87% (82-90%)	13% (9.8-18%)	3.9 (2.8-5.1)	8.6% (6.1-11%)	10% (6.7-15%)	100 (67-140)	7400 (5200-9700)
Dominican Republic	2010	Male	35 to 39	86% (81-90%)	14% (10-19%)	4 (2.8-5.4)	8.1% (5.6-11%)	9.7% (6-14%)	110 (74-160)	7300 (5000-10000)
Dominican Republic	2010	Male	40 to 44	85% (79-89%)	15% (11-21%)	3.8 (2.5-5)	7.7% (5.4-10%)	9.3% (6.1-13%)	130 (80-180)	7000 (4700-9600)
Dominican Republic	2010	Male	45 to 49	83% (76-88%)	17% (12-24%)	3.2 (2.2-4.4)	7.2% (4.8-10%)	8.9% (5.6-13%)	130 (79-190)	6300 (4100-9100)
Dominican Republic	2010	Male	50 to 54	80% (74-86%)	20% (14-26%)	2.7 (1.8-3.9)	6.6% (3.9-9.8%)	8.2% (4.6-13%)	130 (70-190)	5400 (3100-8000)
Dominican Republic	2010	Male	55 to 59	78% (71-84%)	22% (16-29%)	2.6 (1.7-3.8)	6.7% (3.9-10%)	8.6% (5-13%)	130 (74-210)	4700 (2800-7300)
Dominican Republic	2010	Male	60 to 64	76% (65-85%)	24% (15-35%)	2.4 (1.5-3.7)	6% (3.1-9.7%)	7.5% (3.7-13%)	130 (62-210)	3800 (2000-6300)
Dominican Republic	2010	Male	65 to 69	75% (63-86%)	25% (14-37%)	2.1 (1.4-3.1)	5% (2.8-7.7%)	6.2% (3.4-10%)	120 (63-190)	2900 (1600-4600)
Dominican Republic	2010	Male	70 to 74	72% (58-84%)	28% (16-42%)	1.8 (1.1-2.7)	4.1% (2-6.8%)	5% (2.3-8.3%)	120 (55-200)	2300 (1100-3900)
Dominican Republic	2010	Male	75 to 79	66% (50-80%)	34% (20-50%)	1.6 (0.89-2.4)	3.2% (1.2-5.9%)	3.8% (1.3-7%)	110 (36-200)	1600 (620-3000)
Dominican Republic	2010	Male	80 to 84	61% (43-75%)	39% (25-57%)	1.3 (0.76-2)	2.5% (0.97-4.4%)	2.8% (1-5.2%)	93 (34-170)	1100 (420-1900)
Dominican Republic	2010	Male	85 to 89	56% (39-72%)	44% (28-61%)	1.1 (0.6-1.8)	1.8% (0.57-3.5%)	2% (0.47-3.9%)	51 (12-100)	420 (130-790)
Dominican Republic	2010	Male	90 to 94	56% (40-72%)	44% (28-60%)	1 (0.55-1.7)	1.3% (0.016-3%)	1.3% (-0.34-3.2%)	17 (-4.6-43)	110 (1.3-240)
Dominican Republic	2010	Male	95 plus	59% (38-77%)	41% (23-62%)	1.1 (0.49-2)	1.1% (-0.4-3.3%)	0.93% (-0.97-3.6%)	3.8 (-4-14)	19 (-6.7-55)
Dominican Republic	2016	Female	All Ages	43% (39-47%)	57% (53-61%)	1.2 (0.9-1.5)	2% (1.3-2.8%)	2.8% (1.5-4.4%)	680 (330-1100)	23000 (15000-33000)
Dominican Republic	2016	Female	15 to 19	55% (47-62%)	45% (38-53%)	1.5 (0.95-2.3)	2.3% (1.5-3.5%)	5.6% (3.1-8.8%)	12 (6.4-19)	1300 (780-1900)
Dominican Republic	2016	Female	20 to 24	62% (55-68%)	38% (32-45%)	2.3 (1.5-3.4)	3.9% (2.7-5.4%)	7.4% (4.4-11%)	19 (11-31)	2300 (1500-3200)
Dominican Republic	2016	Female	25 to 29	61% (55-67%)	39% (33-45%)	2.3 (1.5-3.3)	3.6% (2.5-5%)	6% (3.5-9.1%)	21 (12-32)	2200 (1500-3100)
Dominican Republic	2016	Female	30 to 34	56% (49-62%)	44% (38-51%)	2 (1.3-3)	3% (2.1-4.3%)	4.7% (2.8-7.1%)	22 (13-33)	2000 (1300-2900)
Dominican Republic	2016	Female	35 to 39	50% (43-58%)	50% (42-57%)	1.7 (1.1-2.6)	2.7% (1.8-3.9%)	4.4% (2.5-6.7%)	23 (14-36)	1800 (1200-2600)
Dominican Republic	2016	Female	40 to 44	46% (39-53%)	54% (47-61%)	1.5 (0.97-2.2)	2.6% (1.7-3.9%)	4.5% (2.7-6.8%)	28 (16-44)	1800 (1100-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	2016	Female	45 to 49	41% (33-50%)	59% (50-67%)	1.4 (0.85-2.1)	2.7% (1.6-4.1%)	4.7% (2.8-7.3%)	34 (19-55)	1800 (1100-2800)
Dominican Republic	2016	Female	50 to 54	42% (31-54%)	58% (46-69%)	1.4 (0.75-2.2)	2.9% (1.5-4.7%)	5.1% (2.7-8.2%)	44 (22-76)	1900 (990-3200)
Dominican Republic	2016	Female	55 to 59	42% (28-57%)	58% (43-72%)	1.2 (0.64-2)	2.7% (1.3-4.7%)	4.7% (2.4-7.7%)	49 (24-83)	1700 (870-3100)
Dominican Republic	2016	Female	60 to 64	36% (23-52%)	64% (48-77%)	1.1 (0.56-1.8)	2.3% (0.88-4.3%)	3.7% (1.5-6.6%)	48 (18-94)	1400 (540-2800)
Dominican Republic	2016	Female	65 to 69	31% (20-43%)	69% (57-80%)	0.98 (0.52-1.6)	2.4% (0.92-4.2%)	3.6% (1.4-6.1%)	55 (21-100)	1300 (520-2500)
Dominican Republic	2016	Female	70 to 74	27% (18-40%)	73% (60-82%)	0.9 (0.46-1.6)	2.3% (0.68-4.4%)	3.3% (1-6.2%)	61 (18-120)	1200 (340-2300)
Dominican Republic	2016	Female	75 to 79	25% (15-37%)	75% (63-85%)	0.84 (0.4-1.5)	2% (0.48-4.2%)	2.7% (0.68-5.7%)	68 (15-140)	1000 (230-2100)
Dominican Republic	2016	Female	80 to 84	23% (14-36%)	77% (64-86%)	0.76 (0.38-1.3)	1.9% (0.56-3.6%)	2.3% (0.73-4.6%)	78 (23-160)	850 (260-1700)
Dominican Republic	2016	Female	85 to 89	24% (14-36%)	76% (64-86%)	0.74 (0.38-1.3)	1.7% (0.42-3.4%)	2% (0.4-4.1%)	65 (13-130)	510 (130-1000)
Dominican Republic	2016	Female	90 to 94	26% (16-39%)	74% (61-84%)	0.83 (0.41-1.5)	1.7% (0.14-4%)	1.9% (-0.022-4.7%)	38 (-0.5-97)	210 (18-520)
Dominican Republic	2016	Female	95 plus	31% (17-49%)	69% (51-83%)	0.96 (0.38-1.8)	1.7% (-0.12-4.9%)	1.8% (-0.45-5.6%)	15 (-3.7-46)	58 (-3.7-160)
Dominican Republic	2016	Male	All Ages	72% (67-75%)	28% (25-33%)	2.1 (1.6-2.7)	5.4% (3.8-7.1%)	5.7% (3.5-8.2%)	1900 (1100-2700)	79000 (54000-110000)
Dominican Republic	2016	Male	15 to 19	73% (66-80%)	27% (20-34%)	1.6 (0.97-2.5)	5.6% (3.6-7.9%)	9.3% (5.2-14%)	35 (19-58)	3400 (2100-5000)
Dominican Republic	2016	Male	20 to 24	85% (80-89%)	15% (11-20%)	3.1 (2.1-4.4)	11% (8-15%)	16% (9.8-22%)	87 (52-130)	8000 (5400-11000)
Dominican Republic	2016	Male	25 to 29	86% (82-90%)	14% (10-18%)	3.6 (2.4-5)	11% (7.7-14%)	14% (8.7-20%)	100 (61-150)	8400 (5700-12000)
Dominican Republic	2016	Male	30 to 34	85% (81-89%)	15% (11-19%)	4 (2.7-5.6)	9.8% (6.7-13%)	12% (7.7-18%)	110 (68-160)	8200 (5600-11000)
Dominican Republic	2016	Male	35 to 39	84% (79-88%)	16% (12-21%)	4 (2.5-5.7)	9.1% (6-12%)	12% (6.9-17%)	120 (71-180)	7900 (5000-11000)
Dominican Republic	2016	Male	40 to 44	83% (78-87%)	17% (13-22%)	3.8 (2.4-5.5)	8.6% (5.7-12%)	11% (6.8-15%)	140 (80-200)	7700 (4800-11000)
Dominican Republic	2016	Male	45 to 49	81% (75-86%)	19% (14-25%)	3.2 (2.1-4.7)	7.8% (5.1-11%)	9.8% (6.1-14%)	140 (82-220)	7000 (4400-10000)
Dominican Republic	2016	Male	50 to 54	77% (70-82%)	23% (18-30%)	2.6 (1.6-3.9)	7% (4.2-10%)	8.9% (5.1-14%)	140 (77-230)	6200 (3600-9400)
Dominican Republic	2016	Male	55 to 59	75% (68-82%)	25% (18-32%)	2.5 (1.6-3.7)	6.9% (3.9-11%)	8.9% (4.9-14%)	160 (87-270)	5900 (3300-9500)
Dominican Republic	2016	Male	60 to 64	75% (64-84%)	25% (16-36%)	2.5 (1.4-4)	6.4% (3.2-11%)	8.2% (4-14%)	170 (79-290)	5200 (2600-8700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Dominican Republic	2016	Male	65 to 69	75% (62-86%)	25% (14-38%)	2.3 (1.4-3.4)	5.6% (2.9-9.1%)	7% (3.6-12%)	160 (80-270)	4000 (2100-6600)
Dominican Republic	2016	Male	70 to 74	72% (58-84%)	28% (16-42%)	1.9 (1.1-3)	4.5% (2-7.8%)	5.5% (2.4-9.9%)	140 (57-250)	2700 (1200-4800)
Dominican Republic	2016	Male	75 to 79	67% (50-81%)	33% (19-50%)	1.7 (0.92-2.7)	3.6% (1.2-6.8%)	4.3% (1.3-8.2%)	130 (39-250)	2000 (640-3800)
Dominican Republic	2016	Male	80 to 84	61% (43-75%)	39% (25-57%)	1.4 (0.78-2.2)	2.8% (1-5.2%)	3.2% (1-6%)	120 (40-240)	1400 (500-2600)
Dominican Republic	2016	Male	85 to 89	56% (39-72%)	44% (28-61%)	1.1 (0.63-1.9)	2.1% (0.59-4.1%)	2.2% (0.48-4.6%)	73 (16-150)	600 (180-1200)
Dominican Republic	2016	Male	90 to 94	56% (39-72%)	44% (28-61%)	1.1 (0.58-1.9)	1.5% (0.064-3.6%)	1.5% (-0.31-4%)	26 (-5.6-70)	160 (7-390)
Dominican Republic	2016	Male	95 plus	59% (38-77%)	41% (23-62%)	1.1 (0.51-2.1)	1.3% (-0.35-3.9%)	1.1% (-0.88-4.2%)	5.9 (-4.6-22)	29 (-7.3-87)
Grenada	1990	Female	All Ages	31% (25-37%)	69% (63-75%)	1.2 (1.1-1.4)	2.9% (1.9-3.9%)	3.8% (2-6%)	14 (7.4-22)	400 (270-550)
Grenada	1990	Female	15 to 19	40% (26-56%)	60% (44-74%)	1.6 (0.91-2.6)	2.3% (1.3-3.4%)	4.8% (2.6-7.7%)	0.14 (0.075-0.23)	15 (9.1-23)
Grenada	1990	Female	20 to 24	46% (30-62%)	54% (38-70%)	2.5 (1.4-3.8)	4% (2.7-5.5%)	7.1% (4.3-10%)	0.21 (0.13-0.32)	22 (14-31)
Grenada	1990	Female	25 to 29	45% (29-63%)	55% (37-71%)	2.6 (1.5-3.8)	4.4% (3.1-6%)	7.6% (5-11%)	0.3 (0.19-0.44)	27 (18-38)
Grenada	1990	Female	30 to 34	43% (27-61%)	57% (39-73%)	2.5 (1.5-3.7)	4.5% (3-6.2%)	7.6% (5-11%)	0.36 (0.22-0.55)	27 (18-40)
Grenada	1990	Female	35 to 39	43% (28-60%)	57% (40-72%)	2.3 (1.3-3.7)	4.8% (3.1-7%)	8.1% (5.2-12%)	0.44 (0.26-0.67)	27 (17-40)
Grenada	1990	Female	40 to 44	42% (26-60%)	58% (40-74%)	2.2 (1.1-3.5)	5.1% (3-7.6%)	8.5% (4.9-12%)	0.51 (0.28-0.77)	27 (15-40)
Grenada	1990	Female	45 to 49	39% (24-57%)	61% (43-76%)	2.1 (1.1-3.6)	5.4% (3.1-8.8%)	8.5% (4.8-14%)	0.56 (0.31-0.93)	26 (14-43)
Grenada	1990	Female	50 to 54	36% (22-53%)	64% (47-78%)	1.9 (1-3.2)	5.4% (2.8-9.1%)	7.9% (4.2-13%)	0.97 (0.5-1.6)	38 (20-65)
Grenada	1990	Female	55 to 59	32% (19-48%)	68% (52-81%)	1.6 (0.83-2.6)	5% (2.4-8.5%)	7% (3.4-11%)	1 (0.45-1.7)	33 (15-56)
Grenada	1990	Female	60 to 64	28% (17-44%)	72% (56-83%)	1.3 (0.67-2.3)	4.1% (1.5-7.2%)	5.4% (2.1-9.4%)	1.4 (0.53-2.5)	40 (15-71)
Grenada	1990	Female	65 to 69	26% (16-39%)	74% (61-84%)	1.2 (0.63-1.9)	3.6% (1.4-6.3%)	4.6% (1.9-7.8%)	1.8 (0.7-3)	41 (16-71)
Grenada	1990	Female	70 to 74	23% (14-37%)	77% (63-86%)	0.99 (0.51-1.7)	2.9% (0.74-5.7%)	3.6% (0.95-7.1%)	2.1 (0.53-4.2)	39 (9.6-78)
Grenada	1990	Female	75 to 79	21% (12-34%)	79% (66-88%)	0.86 (0.39-1.5)	2.3% (0.34-4.7%)	2.7% (0.39-5.7%)	1.3 (0.18-2.7)	18 (2.8-38)
Grenada	1990	Female	80 to 84	19% (11-30%)	81% (70-89%)	0.8 (0.38-1.3)	2.1% (0.47-4.3%)	2.5% (0.53-5.1%)	1.1 (0.23-2.3)	12 (2.6-25)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	1990	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.82 (0.41-1.5)	2.2% (0.53-4.7%)	2.5% (0.58-5.6%)	0.98 (0.23-2.2)	7.4 (1.8-16)
Grenada	1990	Female	90 to 94	21% (13-34%)	79% (66-87%)	0.94 (0.42-1.8)	2.4% (0.41-5.8%)	2.8% (0.41-6.8%)	0.54 (0.083-1.3)	2.8 (0.5-6.7)
Grenada	1990	Female	95 plus	25% (13-42%)	75% (58-87%)	1.1 (0.42-2.3)	2.9% (0.17-7.6%)	3.2% (0.053-8.7%)	0.17 (0.0028-0.47)	0.61 (0.035-1.6)
Grenada	1990	Male	All Ages	52% (43-61%)	48% (39-57%)	2.6 (2.4-2.9)	8.1% (6.5-9.9%)	10% (7.5-13%)	36 (26-47)	1200 (980-1500)
Grenada	1990	Male	15 to 19	57% (40-73%)	43% (27-60%)	2.4 (1.4-3.7)	6.1% (3.9-8.9%)	8.5% (5-13%)	0.59 (0.34-0.93)	53 (34-79)
Grenada	1990	Male	20 to 24	67% (50-82%)	33% (18-50%)	4.6 (3-6.4)	13% (9.1-17%)	16% (11-22%)	1.3 (0.8-1.9)	100 (72-140)
Grenada	1990	Male	25 to 29	70% (52-83%)	30% (17-48%)	5.5 (3.6-7.5)	14% (10-17%)	18% (13-23%)	1.5 (0.97-2)	110 (78-150)
Grenada	1990	Male	30 to 34	71% (53-84%)	29% (16-47%)	5.8 (3.8-8.1)	14% (11-17%)	18% (14-23%)	1.5 (1-2.1)	100 (75-130)
Grenada	1990	Male	35 to 39	71% (55-84%)	29% (16-45%)	5.5 (3.4-8.3)	13% (10-17%)	18% (13-23%)	1.6 (1.1-2.2)	96 (69-130)
Grenada	1990	Male	40 to 44	70% (54-84%)	30% (16-46%)	5.3 (3-8.1)	14% (9.9-17%)	18% (13-23%)	1.5 (1-2.1)	81 (57-110)
Grenada	1990	Male	45 to 49	68% (49-82%)	32% (18-51%)	5 (2.8-7.7)	14% (10-17%)	18% (13-22%)	1.5 (1.1-2)	68 (49-90)
Grenada	1990	Male	50 to 54	67% (49-82%)	33% (18-51%)	4.6 (2.6-7.2)	13% (9.1-17%)	16% (11-21%)	2.7 (1.8-3.6)	110 (72-140)
Grenada	1990	Male	55 to 59	65% (46-81%)	35% (19-54%)	4.2 (2.3-6.6)	13% (8.4-17%)	16% (10-20%)	2.7 (1.7-3.7)	92 (60-120)
Grenada	1990	Male	60 to 64	63% (43-80%)	37% (20-57%)	3.8 (2.1-6.2)	11% (6.3-15%)	13% (7.5-18%)	4.5 (2.5-6.3)	130 (73-180)
Grenada	1990	Male	65 to 69	61% (42-78%)	39% (22-58%)	3.3 (1.9-5.2)	10% (6.3-14%)	12% (7.3-17%)	4.6 (2.7-6.5)	110 (64-150)
Grenada	1990	Male	70 to 74	58% (39-74%)	42% (26-61%)	2.8 (1.4-4.6)	8.4% (4.3-13%)	9.7% (4.9-15%)	4.8 (2.4-7.4)	90 (46-140)
Grenada	1990	Male	75 to 79	53% (34-71%)	47% (29-66%)	2.3 (1.2-4)	6.6% (2.9-11%)	7.5% (3.2-12%)	3.1 (1.3-5.1)	45 (20-73)
Grenada	1990	Male	80 to 84	47% (29-65%)	53% (35-71%)	2 (1-3.2)	5.5% (2.5-9.1%)	6.3% (2.8-10%)	2.1 (0.91-3.5)	23 (9.9-37)
Grenada	1990	Male	85 to 89	42% (26-59%)	58% (41-74%)	1.7 (0.88-2.9)	4.7% (2-8.3%)	5.3% (2.2-9.5%)	1.2 (0.48-2.2)	9.2 (3.8-16)
Grenada	1990	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.7 (0.84-3.1)	4.3% (1.5-8.3%)	4.8% (1.5-9.5%)	0.38 (0.12-0.75)	2 (0.69-3.9)
Grenada	1990	Male	95 plus	46% (26-66%)	54% (34-74%)	1.9 (0.78-3.5)	4.4% (1.1-8.8%)	4.8% (0.99-10%)	0.077 (0.015-0.16)	0.28 (0.07-0.57)
Grenada	1995	Female	All Ages	31% (25-38%)	69% (62-75%)	1.2 (1.1-1.4)	2.9% (2-4%)	3.7% (2-5.9%)	14 (7.5-22)	390 (270-550)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	1995	Female	15 to 19	39% (26-56%)	61% (44-74%)	1.6 (0.91-2.6)	2.3% (1.4-3.4%)	4.7% (2.5-7.6%)	0.13 (0.07-0.22)	15 (8.6-22)
Grenada	1995	Female	20 to 24	46% (30-62%)	54% (38-70%)	2.5 (1.4-3.7)	4% (2.7-5.5%)	6.9% (4.1-10%)	0.23 (0.13-0.35)	24 (16-35)
Grenada	1995	Female	25 to 29	45% (29-62%)	55% (38-71%)	2.6 (1.5-3.9)	4.4% (3-6%)	7.3% (4.6-10%)	0.23 (0.14-0.35)	21 (14-30)
Grenada	1995	Female	30 to 34	43% (27-61%)	57% (39-73%)	2.5 (1.4-3.7)	4.5% (3-6.2%)	7.4% (4.8-11%)	0.42 (0.25-0.64)	32 (21-46)
Grenada	1995	Female	35 to 39	42% (27-60%)	58% (40-73%)	2.3 (1.3-3.6)	4.7% (3.1-6.8%)	7.8% (4.9-11%)	0.53 (0.31-0.8)	33 (21-50)
Grenada	1995	Female	40 to 44	41% (27-59%)	59% (41-73%)	2.2 (1.2-3.4)	5% (2.9-7.5%)	8.2% (4.8-12%)	0.57 (0.33-0.87)	30 (18-45)
Grenada	1995	Female	45 to 49	39% (24-55%)	61% (45-76%)	2.1 (1.1-3.6)	5.3% (3-8.6%)	8.1% (4.7-13%)	0.6 (0.34-0.98)	28 (16-45)
Grenada	1995	Female	50 to 54	36% (22-53%)	64% (47-78%)	1.9 (1-3.3)	5.3% (2.7-9%)	7.7% (4-13%)	0.66 (0.34-1.1)	26 (13-45)
Grenada	1995	Female	55 to 59	31% (19-49%)	69% (51-81%)	1.6 (0.82-2.7)	4.9% (2.2-8.3%)	6.8% (3.3-11%)	1 (0.47-1.7)	34 (16-57)
Grenada	1995	Female	60 to 64	28% (16-44%)	72% (56-84%)	1.4 (0.68-2.3)	4% (1.6-7.1%)	5.4% (2.3-9.4%)	1 (0.41-1.8)	29 (11-52)
Grenada	1995	Female	65 to 69	26% (16-39%)	74% (61-84%)	1.2 (0.63-2)	3.5% (1.4-6.2%)	4.5% (1.8-7.8%)	1.6 (0.66-2.8)	37 (15-65)
Grenada	1995	Female	70 to 74	23% (15-35%)	77% (65-85%)	0.98 (0.51-1.7)	2.8% (0.79-5.6%)	3.6% (1.1-6.9%)	1.7 (0.5-3.3)	32 (9-62)
Grenada	1995	Female	75 to 79	21% (12-33%)	79% (67-88%)	0.85 (0.39-1.5)	2.3% (0.41-4.8%)	2.8% (0.5-5.8%)	1.9 (0.34-3.9)	27 (4.7-57)
Grenada	1995	Female	80 to 84	19% (11-30%)	81% (70-89%)	0.8 (0.38-1.4)	2.2% (0.55-4.4%)	2.5% (0.64-5.3%)	1.4 (0.35-2.9)	15 (3.8-30)
Grenada	1995	Female	85 to 89	19% (11-30%)	81% (70-89%)	0.82 (0.41-1.5)	2.2% (0.6-4.7%)	2.5% (0.66-5.6%)	0.95 (0.24-2.1)	7.1 (2-16)
Grenada	1995	Female	90 to 94	21% (13-33%)	79% (67-87%)	0.95 (0.41-1.8)	2.5% (0.46-6%)	2.8% (0.47-7%)	0.62 (0.1-1.5)	3.2 (0.62-7.7)
Grenada	1995	Female	95 plus	25% (13-42%)	75% (58-87%)	1.1 (0.41-2.4)	2.9% (0.15-7.7%)	3.3% (0.061-8.8%)	0.24 (0.0042-0.63)	0.82 (0.042-2.1)
Grenada	1995	Male	All Ages	52% (44-61%)	48% (39-56%)	2.6 (2.4-2.9)	8.5% (7-10%)	10% (7.4-13%)	35 (26-46)	1200 (1000-1500)
Grenada	1995	Male	15 to 19	56% (39-73%)	44% (27-61%)	2.4 (1.3-3.7)	6.2% (4-9.2%)	8.5% (5-13%)	0.62 (0.34-0.99)	55 (34-82)
Grenada	1995	Male	20 to 24	67% (49-81%)	33% (19-51%)	4.6 (2.9-6.5)	13% (9.2-17%)	16% (11-22%)	1.6 (0.98-2.2)	130 (86-170)
Grenada	1995	Male	25 to 29	70% (52-83%)	30% (17-48%)	5.5 (3.6-7.7)	14% (10-17%)	17% (12-23%)	1.3 (0.87-1.8)	96 (69-130)
Grenada	1995	Male	30 to 34	71% (53-84%)	29% (16-47%)	5.8 (3.7-8.1)	13% (10-16%)	17% (13-22%)	1.8 (1.2-2.4)	120 (87-160)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	1995	Male	35 to 39	71% (54-84%)	29% (16-46%)	5.4 (3.4-8.1)	13% (9.9-16%)	17% (12-21%)	1.9 (1.3-2.5)	110 (81-150)
Grenada	1995	Male	40 to 44	70% (54-83%)	30% (17-46%)	5.2 (3-7.9)	13% (9.7-17%)	17% (12-21%)	1.9 (1.3-2.6)	100 (73-130)
Grenada	1995	Male	45 to 49	68% (48-83%)	32% (17-52%)	4.9 (2.8-7.8)	14% (10-17%)	17% (13-22%)	1.8 (1.3-2.4)	83 (60-110)
Grenada	1995	Male	50 to 54	66% (48-82%)	34% (18-52%)	4.6 (2.6-7.2)	13% (9.1-17%)	16% (11-21%)	1.9 (1.3-2.6)	75 (51-100)
Grenada	1995	Male	55 to 59	65% (47-80%)	35% (20-53%)	4.2 (2.3-6.7)	13% (8.3-17%)	15% (10-20%)	3.3 (2.1-4.5)	110 (74-150)
Grenada	1995	Male	60 to 64	62% (42-80%)	38% (20-58%)	3.8 (2-6.3)	11% (6.6-15%)	13% (7.6-18%)	2.8 (1.7-4)	81 (48-110)
Grenada	1995	Male	65 to 69	61% (42-77%)	39% (23-58%)	3.3 (1.9-5.3)	10% (6.2-14%)	12% (7.2-17%)	5 (2.9-7.2)	120 (70-170)
Grenada	1995	Male	70 to 74	57% (38-74%)	43% (26-62%)	2.8 (1.4-4.7)	8.2% (4.3-13%)	9.5% (4.9-15%)	3.8 (1.9-6)	72 (37-110)
Grenada	1995	Male	75 to 79	52% (34-70%)	48% (30-66%)	2.3 (1.2-3.9)	6.4% (2.9-10%)	7.2% (3.3-12%)	3.7 (1.6-6.1)	54 (24-88)
Grenada	1995	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.9 (0.99-3.1)	5.3% (2.4-8.7%)	6% (2.6-9.8%)	2.2 (0.95-3.7)	24 (11-40)
Grenada	1995	Male	85 to 89	42% (25-59%)	58% (41-75%)	1.7 (0.86-2.8)	4.6% (1.9-8%)	5.1% (2-9.2%)	1.2 (0.49-2.2)	9.5 (3.9-17)
Grenada	1995	Male	90 to 94	42% (27-59%)	58% (41-73%)	1.7 (0.81-3.1)	4.1% (1.5-7.8%)	4.6% (1.5-9%)	0.48 (0.16-0.94)	2.6 (0.92-4.9)
Grenada	1995	Male	95 plus	46% (26-66%)	54% (34-74%)	1.9 (0.77-3.5)	4.2% (1-8.5%)	4.7% (0.85-9.8%)	0.12 (0.021-0.25)	0.42 (0.098-0.87)
Grenada	2000	Female	All Ages	32% (26-38%)	68% (62-74%)	1.2 (1.1-1.4)	2.7% (1.9-3.7%)	3.3% (1.7-5.4%)	13 (6.6-21)	370 (250-520)
Grenada	2000	Female	15 to 19	39% (26-56%)	61% (44-74%)	1.6 (0.89-2.5)	2.2% (1.3-3.3%)	4.5% (2.4-7.3%)	0.14 (0.073-0.23)	16 (9.5-24)
Grenada	2000	Female	20 to 24	45% (29-62%)	55% (38-71%)	2.4 (1.4-3.6)	3.8% (2.6-5.2%)	6.3% (3.8-9.4%)	0.22 (0.12-0.33)	24 (16-34)
Grenada	2000	Female	25 to 29	44% (28-62%)	56% (38-72%)	2.4 (1.4-3.7)	4% (2.7-5.6%)	6.4% (3.9-9.2%)	0.24 (0.13-0.36)	22 (14-32)
Grenada	2000	Female	30 to 34	43% (27-60%)	57% (40-73%)	2.4 (1.3-3.6)	4.1% (2.7-5.7%)	6.6% (4.2-9.4%)	0.28 (0.16-0.44)	22 (14-32)
Grenada	2000	Female	35 to 39	42% (27-59%)	58% (41-73%)	2.2 (1.3-3.4)	4.3% (2.7-6.2%)	6.9% (4.4-9.9%)	0.52 (0.3-0.79)	33 (20-49)
Grenada	2000	Female	40 to 44	41% (26-59%)	59% (41-74%)	2.1 (1.1-3.3)	4.6% (2.7-6.9%)	7.4% (4.5-11%)	0.59 (0.34-0.89)	32 (19-48)
Grenada	2000	Female	45 to 49	39% (23-56%)	61% (44-77%)	2 (1-3.4)	4.8% (2.7-8%)	7.4% (4.3-12%)	0.63 (0.35-1)	29 (16-48)
Grenada	2000	Female	50 to 54	35% (21-52%)	65% (48-79%)	1.8 (0.93-3.1)	4.8% (2.3-8.3%)	6.9% (3.5-11%)	0.68 (0.34-1.1)	27 (13-46)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	2000	Female	55 to 59	31% (19-48%)	69% (52-81%)	1.5 (0.77-2.5)	4.3% (2-7.5%)	6% (2.8-10%)	0.82 (0.39-1.4)	28 (13-48)
Grenada	2000	Female	60 to 64	28% (16-43%)	72% (57-84%)	1.3 (0.64-2.2)	3.5% (1.3-6.6%)	4.8% (1.8-8.7%)	0.98 (0.37-1.8)	28 (9.7-53)
Grenada	2000	Female	65 to 69	25% (15-39%)	75% (61-85%)	1.1 (0.59-1.9)	3.1% (1.1-5.6%)	4% (1.5-7%)	1 (0.38-1.8)	24 (8.5-45)
Grenada	2000	Female	70 to 74	23% (14-36%)	77% (64-86%)	0.94 (0.48-1.7)	2.5% (0.56-5.3%)	3.2% (0.84-6.4%)	1.6 (0.42-3.4)	30 (7.1-65)
Grenada	2000	Female	75 to 79	20% (12-34%)	80% (66-88%)	0.82 (0.38-1.5)	2% (0.23-4.4%)	2.5% (0.29-5.3%)	1.4 (0.16-3.1)	20 (2.2-45)
Grenada	2000	Female	80 to 84	19% (11-30%)	81% (70-89%)	0.77 (0.36-1.3)	2% (0.42-4.1%)	2.3% (0.51-4.8%)	1.5 (0.34-3.2)	16 (3.7-35)
Grenada	2000	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.78 (0.38-1.4)	2% (0.46-4.5%)	2.3% (0.49-5.2%)	1.2 (0.26-2.7)	9.1 (2.1-20)
Grenada	2000	Female	90 to 94	21% (12-32%)	79% (68-88%)	0.89 (0.39-1.7)	2.2% (0.35-5.4%)	2.6% (0.36-6.4%)	0.59 (0.082-1.5)	3.1 (0.49-7.5)
Grenada	2000	Female	95 plus	24% (13-41%)	76% (59-87%)	1.1 (0.39-2.2)	2.7% (0.19-6.9%)	3% (0.086-8%)	0.26 (0.0075-0.71)	0.92 (0.069-2.4)
Grenada	2000	Male	All Ages	53% (44-62%)	47% (38-56%)	2.6 (2.4-2.9)	8.6% (7.1-10%)	9.5% (7.2-12%)	36 (27-47)	1300 (1100-1600)
Grenada	2000	Male	15 to 19	56% (38-73%)	44% (27-62%)	2.3 (1.3-3.5)	6.2% (4.1-9%)	8.4% (5-13%)	0.75 (0.42-1.2)	66 (41-99)
Grenada	2000	Male	20 to 24	66% (49-81%)	34% (19-51%)	4.4 (2.8-6.1)	13% (9.3-17%)	16% (11-22%)	1.7 (1.1-2.4)	140 (92-180)
Grenada	2000	Male	25 to 29	70% (51-83%)	30% (17-49%)	5.3 (3.4-7.4)	14% (10-17%)	17% (12-22%)	1.7 (1.1-2.4)	130 (88-170)
Grenada	2000	Male	30 to 34	71% (51-84%)	29% (16-49%)	5.5 (3.4-7.8)	13% (10-17%)	17% (13-21%)	1.6 (1-2.2)	110 (75-140)
Grenada	2000	Male	35 to 39	70% (54-84%)	30% (16-46%)	5.2 (3.3-7.8)	13% (9.6-16%)	16% (12-20%)	2.1 (1.4-2.8)	130 (89-160)
Grenada	2000	Male	40 to 44	69% (53-83%)	31% (17-47%)	5 (2.9-7.4)	13% (9.4-16%)	16% (12-20%)	2.1 (1.4-2.9)	110 (80-150)
Grenada	2000	Male	45 to 49	67% (47-82%)	33% (18-53%)	4.7 (2.7-7.4)	13% (9.5-17%)	16% (12-20%)	2.5 (1.7-3.3)	110 (80-150)
Grenada	2000	Male	50 to 54	66% (47-81%)	34% (19-53%)	4.3 (2.4-6.8)	13% (8.6-16%)	15% (10-20%)	2.4 (1.5-3.3)	96 (64-130)
Grenada	2000	Male	55 to 59	64% (46-80%)	36% (20-54%)	4 (2.2-6.3)	12% (8-16%)	15% (9.3-19%)	2.6 (1.7-3.6)	88 (57-120)
Grenada	2000	Male	60 to 64	62% (41-79%)	38% (21-59%)	3.6 (1.9-6)	10% (6-14%)	12% (7-17%)	3.4 (1.9-4.9)	97 (57-140)
Grenada	2000	Male	65 to 69	60% (41-77%)	40% (23-59%)	3.2 (1.8-5)	9.5% (5.6-13%)	11% (6.5-16%)	3.6 (2.1-5.2)	85 (51-120)
Grenada	2000	Male	70 to 74	57% (38-73%)	43% (27-62%)	2.6 (1.4-4.5)	7.5% (3.7-12%)	8.6% (4.2-13%)	4.6 (2.2-7.4)	87 (43-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	2000	Male	75 to 79	52% (32-71%)	48% (29-68%)	2.2 (1.1-3.7)	5.8% (2.6-9.5%)	6.6% (2.9-11%)	3 (1.3-5.1)	44 (20-73)
Grenada	2000	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.8 (0.93-2.9)	4.8% (2.1-8%)	5.4% (2.4-9.1%)	2.4 (1-4.1)	26 (12-43)
Grenada	2000	Male	85 to 89	41% (25-58%)	59% (42-75%)	1.6 (0.82-2.7)	4.1% (1.8-7.4%)	4.6% (1.9-8.3%)	1.1 (0.47-2)	8.8 (3.8-15)
Grenada	2000	Male	90 to 94	42% (26-59%)	58% (41-74%)	1.6 (0.77-2.9)	3.8% (1.4-7.4%)	4.2% (1.4-8.3%)	0.43 (0.15-0.87)	2.3 (0.86-4.6)
Grenada	2000	Male	95 plus	45% (26-65%)	55% (35-74%)	1.7 (0.72-3.4)	4% (1.1-8.1%)	4.3% (0.93-9.3%)	0.13 (0.027-0.27)	0.46 (0.12-0.95)
Grenada	2005	Female	All Ages	33% (27-40%)	67% (60-73%)	1.3 (1.1-1.4)	2.7% (1.8-3.7%)	3% (1.5-5.1%)	13 (6.4-22)	380 (260-540)
Grenada	2005	Female	15 to 19	39% (26-56%)	61% (44-74%)	1.5 (0.86-2.4)	2.1% (1.3-3.2%)	4.3% (2.4-7%)	0.15 (0.08-0.25)	17 (10-26)
Grenada	2005	Female	20 to 24	45% (29-62%)	55% (38-71%)	2.3 (1.4-3.3)	3.7% (2.5-5.1%)	6.1% (3.7-9%)	0.26 (0.15-0.39)	28 (19-40)
Grenada	2005	Female	25 to 29	45% (28-62%)	55% (38-72%)	2.3 (1.4-3.6)	4% (2.7-5.5%)	6.2% (3.7-8.9%)	0.29 (0.16-0.43)	27 (17-37)
Grenada	2005	Female	30 to 34	43% (27-60%)	57% (40-73%)	2.3 (1.3-3.5)	4.1% (2.8-5.7%)	6.4% (4.1-9.2%)	0.32 (0.18-0.48)	24 (15-35)
Grenada	2005	Female	35 to 39	42% (27-59%)	58% (41-73%)	2.1 (1.2-3.3)	4.3% (2.8-6.1%)	6.7% (4.3-9.8%)	0.41 (0.24-0.63)	26 (16-38)
Grenada	2005	Female	40 to 44	41% (26-59%)	59% (41-74%)	2 (1.1-3.2)	4.4% (2.6-6.7%)	7% (4.2-11%)	0.73 (0.43-1.1)	39 (23-60)
Grenada	2005	Female	45 to 49	39% (24-56%)	61% (44-76%)	1.9 (1-3.2)	4.6% (2.5-7.8%)	7% (3.9-11%)	0.8 (0.43-1.3)	37 (20-61)
Grenada	2005	Female	50 to 54	35% (22-51%)	65% (49-78%)	1.7 (0.91-2.9)	4.5% (2.1-7.9%)	6.4% (3.3-11%)	0.79 (0.39-1.3)	31 (15-53)
Grenada	2005	Female	55 to 59	31% (19-47%)	69% (53-81%)	1.5 (0.75-2.4)	4.1% (1.8-7.1%)	5.6% (2.7-9.7%)	0.78 (0.37-1.3)	27 (12-46)
Grenada	2005	Female	60 to 64	28% (16-44%)	72% (56-84%)	1.2 (0.62-2.2)	3.3% (1-6.3%)	4.4% (1.6-8.2%)	0.84 (0.31-1.6)	24 (7.6-47)
Grenada	2005	Female	65 to 69	26% (16-39%)	74% (61-84%)	1.1 (0.58-1.8)	2.8% (0.9-5.4%)	3.7% (1.3-6.8%)	1.1 (0.39-2)	25 (8.3-49)
Grenada	2005	Female	70 to 74	23% (14-36%)	77% (64-86%)	0.9 (0.45-1.6)	2.2% (0.36-4.7%)	2.7% (0.57-5.9%)	1 (0.21-2.2)	19 (3.1-42)
Grenada	2005	Female	75 to 79	20% (12-33%)	80% (67-88%)	0.78 (0.35-1.4)	1.8% (0.12-4.1%)	2.1% (0.2-5%)	1.5 (0.14-3.4)	21 (1.5-50)
Grenada	2005	Female	80 to 84	19% (11-30%)	81% (70-89%)	0.73 (0.35-1.2)	1.8% (0.31-3.8%)	2.1% (0.37-4.5%)	1.5 (0.26-3.2)	16 (2.6-35)
Grenada	2005	Female	85 to 89	19% (11-30%)	81% (70-89%)	0.74 (0.36-1.3)	1.8% (0.35-4.1%)	2.1% (0.39-4.7%)	1.4 (0.25-3.2)	10 (1.9-24)
Grenada	2005	Female	90 to 94	21% (12-33%)	79% (67-88%)	0.85 (0.39-1.6)	2% (0.32-5%)	2.3% (0.29-5.7%)	0.78 (0.096-1.9)	4.1 (0.62-10)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	2005	Female	95 plus	25% (13-41%)	75% (59-87%)	1 (0.38-2.1)	2.4% (0.093-6.4%)	2.7% (-0.021-7.3%)	0.26 (-0.002-0.69)	0.89 (0.033-2.3)
Grenada	2005	Male	All Ages	55% (47-64%)	45% (36-53%)	2.7 (2.5-3)	8.9% (7.3-10%)	9.3% (7-12%)	38 (28-49)	1400 (1200-1700)
Grenada	2005	Male	15 to 19	56% (39-73%)	44% (27-61%)	2.2 (1.3-3.3)	6.1% (3.9-8.8%)	8.2% (4.9-12%)	0.81 (0.42-1.3)	71 (43-110)
Grenada	2005	Male	20 to 24	67% (49-82%)	33% (18-51%)	4.2 (2.7-5.9)	13% (9-17%)	15% (10-21%)	2 (1.2-3)	160 (110-230)
Grenada	2005	Male	25 to 29	70% (52-83%)	30% (17-48%)	5 (3.2-7)	14% (10-18%)	17% (12-22%)	1.9 (1.2-2.7)	140 (94-190)
Grenada	2005	Male	30 to 34	71% (52-84%)	29% (16-48%)	5.3 (3.3-7.5)	13% (10-17%)	17% (12-22%)	1.9 (1.2-2.6)	120 (87-160)
Grenada	2005	Male	35 to 39	71% (54-84%)	29% (16-46%)	5 (3.1-7.5)	13% (9.8-16%)	17% (12-21%)	1.8 (1.2-2.4)	110 (76-140)
Grenada	2005	Male	40 to 44	69% (54-83%)	31% (17-46%)	4.8 (2.7-7.2)	13% (9.2-16%)	16% (11-21%)	2.7 (1.8-3.7)	140 (100-190)
Grenada	2005	Male	45 to 49	68% (47-82%)	32% (18-53%)	4.5 (2.6-7)	13% (9.3-17%)	16% (11-20%)	2.9 (1.9-4)	130 (91-180)
Grenada	2005	Male	50 to 54	66% (47-81%)	34% (19-53%)	4.1 (2.3-6.5)	12% (8.5-16%)	15% (10-20%)	3.2 (2.1-4.3)	130 (86-170)
Grenada	2005	Male	55 to 59	64% (46-80%)	36% (20-54%)	3.8 (2.1-6.1)	12% (7.5-16%)	14% (8.8-19%)	3 (1.9-4.2)	100 (67-140)
Grenada	2005	Male	60 to 64	62% (42-79%)	38% (21-58%)	3.4 (1.8-5.8)	10% (5.6-14%)	12% (6.5-16%)	2.9 (1.6-4.1)	82 (47-120)
Grenada	2005	Male	65 to 69	61% (41-77%)	39% (23-59%)	3 (1.7-4.8)	8.9% (5.1-13%)	10% (5.9-15%)	3.8 (2.2-5.5)	89 (51-130)
Grenada	2005	Male	70 to 74	57% (39-74%)	43% (26-61%)	2.5 (1.3-4.3)	6.8% (3.3-11%)	7.7% (3.7-12%)	3.1 (1.5-5)	60 (29-95)
Grenada	2005	Male	75 to 79	52% (33-71%)	48% (29-67%)	2.1 (1-3.6)	5.4% (2.3-9%)	6.1% (2.6-10%)	3.6 (1.5-6.1)	53 (23-89)
Grenada	2005	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.7 (0.89-2.8)	4.5% (2-7.4%)	5% (2.1-8.5%)	2.4 (1-4.1)	26 (12-44)
Grenada	2005	Male	85 to 89	41% (25-59%)	59% (41-75%)	1.5 (0.77-2.6)	3.8% (1.6-6.9%)	4.3% (1.8-7.8%)	1.3 (0.52-2.3)	9.8 (4.2-18)
Grenada	2005	Male	90 to 94	42% (26-59%)	58% (41-74%)	1.6 (0.75-2.9)	3.5% (1.3-7%)	3.8% (1.3-7.8%)	0.43 (0.15-0.87)	2.3 (0.87-4.6)
Grenada	2005	Male	95 plus	45% (26-65%)	55% (35-74%)	1.7 (0.69-3.2)	3.6% (0.97-7.6%)	3.9% (0.87-8.5%)	0.12 (0.026-0.25)	0.42 (0.11-0.89)
Grenada	2010	Female	All Ages	33% (27-40%)	67% (60-73%)	1.2 (1.1-1.4)	2.5% (1.7-3.5%)	2.7% (1.4-4.5%)	11 (5.7-19)	360 (250-500)
Grenada	2010	Female	15 to 19	39% (26-55%)	61% (45-74%)	1.4 (0.78-2.3)	2% (1.3-3.1%)	4.1% (2.2-6.8%)	0.12 (0.065-0.21)	14 (8.4-21)
Grenada	2010	Female	20 to 24	45% (29-61%)	55% (39-71%)	2.1 (1.2-3.2)	3.5% (2.4-4.9%)	5.7% (3.4-8.5%)	0.26 (0.16-0.41)	30 (20-43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	2010	Female	25 to 29	44% (28-61%)	56% (39-72%)	2.1 (1.3-3.2)	3.7% (2.5-5.1%)	5.7% (3.5-8.3%)	0.31 (0.18-0.48)	30 (20-43)
Grenada	2010	Female	30 to 34	42% (26-60%)	58% (40-74%)	2.1 (1.2-3.2)	3.7% (2.5-5.3%)	5.8% (3.7-8.6%)	0.34 (0.2-0.52)	27 (17-38)
Grenada	2010	Female	35 to 39	42% (26-60%)	58% (40-74%)	1.9 (1.1-3.1)	3.9% (2.5-5.8%)	6.2% (3.9-9.5%)	0.39 (0.22-0.61)	25 (15-37)
Grenada	2010	Female	40 to 44	40% (26-59%)	60% (41-74%)	1.8 (0.96-3)	4.1% (2.4-6.2%)	6.5% (3.9-9.8%)	0.49 (0.28-0.75)	26 (15-41)
Grenada	2010	Female	45 to 49	39% (23-56%)	61% (44-77%)	1.8 (0.91-3)	4.2% (2.2-7.1%)	6.4% (3.5-11%)	0.91 (0.5-1.5)	42 (23-70)
Grenada	2010	Female	50 to 54	35% (21-51%)	65% (49-79%)	1.6 (0.84-2.7)	4.1% (1.9-7.1%)	5.9% (2.9-9.8%)	0.93 (0.46-1.6)	37 (18-64)
Grenada	2010	Female	55 to 59	31% (18-47%)	69% (53-82%)	1.3 (0.67-2.2)	3.6% (1.4-6.5%)	5% (2.3-8.7%)	0.83 (0.37-1.5)	28 (12-51)
Grenada	2010	Female	60 to 64	27% (16-43%)	73% (57-84%)	1.1 (0.55-2)	2.9% (0.86-5.7%)	3.9% (1.3-7.3%)	0.72 (0.24-1.4)	21 (6.2-41)
Grenada	2010	Female	65 to 69	25% (16-39%)	75% (61-84%)	0.99 (0.53-1.6)	2.4% (0.69-4.7%)	3.1% (1-5.9%)	0.82 (0.27-1.6)	19 (5.6-39)
Grenada	2010	Female	70 to 74	23% (14-35%)	77% (65-86%)	0.82 (0.41-1.5)	1.8% (0.16-4.1%)	2.3% (0.32-4.9%)	0.96 (0.14-2.1)	18 (1.7-41)
Grenada	2010	Female	75 to 79	20% (12-33%)	80% (67-88%)	0.72 (0.32-1.3)	1.5% (-0.025-3.7%)	1.8% (0.019-4.5%)	0.86 (0.0093-2.1)	12 (-0.23-31)
Grenada	2010	Female	80 to 84	19% (10-30%)	81% (70-90%)	0.67 (0.32-1.2)	1.5% (0.18-3.4%)	1.7% (0.23-4%)	1.2 (0.16-2.8)	13 (1.6-30)
Grenada	2010	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.69 (0.33-1.2)	1.5% (0.24-3.7%)	1.8% (0.25-4.2%)	1.2 (0.17-2.9)	9.3 (1.4-22)
Grenada	2010	Female	90 to 94	21% (12-33%)	79% (67-88%)	0.78 (0.33-1.5)	1.7% (0.19-4.4%)	2% (0.15-5%)	0.79 (0.062-2)	4.2 (0.46-10)
Grenada	2010	Female	95 plus	24% (13-41%)	76% (59-87%)	0.95 (0.34-2)	2.1% (0.024-6%)	2.3% (-0.088-6.8%)	0.32 (-0.012-0.95)	1.1 (0.013-3.2)
Grenada	2010	Male	All Ages	56% (48-65%)	44% (35-52%)	2.7 (2.4-3)	8.6% (7.1-10%)	8.9% (6.7-11%)	36 (27-47)	1400 (1100-1800)
Grenada	2010	Male	15 to 19	56% (38-73%)	44% (27-62%)	2 (1.1-3.1)	5.8% (3.8-8.5%)	7.8% (4.7-12%)	0.65 (0.37-1)	58 (35-86)
Grenada	2010	Male	20 to 24	66% (48-81%)	34% (19-52%)	3.8 (2.5-5.4)	12% (8.5-16%)	15% (9.6-20%)	2.1 (1.2-3.1)	170 (110-240)
Grenada	2010	Male	25 to 29	69% (52-83%)	31% (17-48%)	4.6 (3.1-6.4)	13% (9.6-17%)	16% (11-22%)	2.1 (1.3-3)	160 (110-220)
Grenada	2010	Male	30 to 34	70% (51-84%)	30% (16-49%)	4.9 (3.1-7.1)	13% (9.6-16%)	16% (12-21%)	1.9 (1.2-2.6)	130 (87-170)
Grenada	2010	Male	35 to 39	70% (54-84%)	30% (16-46%)	4.6 (2.8-6.9)	12% (9-16%)	16% (11-21%)	1.9 (1.3-2.7)	110 (81-160)
Grenada	2010	Male	40 to 44	69% (53-83%)	31% (17-47%)	4.4 (2.5-6.7)	12% (8.6-16%)	16% (11-20%)	2 (1.3-2.8)	110 (72-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	2010	Male	45 to 49	67% (47-82%)	33% (18-53%)	4.1 (2.3-6.5)	12% (8.5-16%)	15% (10-19%)	3.2 (2.1-4.5)	150 (99-200)
Grenada	2010	Male	50 to 54	66% (47-81%)	34% (19-53%)	3.8 (2.1-6.1)	12% (7.7-15%)	14% (9.2-19%)	3.3 (2.1-4.6)	130 (87-180)
Grenada	2010	Male	55 to 59	64% (46-80%)	36% (20-54%)	3.5 (1.9-5.3)	11% (6.6-15%)	13% (7.7-17%)	3.6 (2.1-5)	120 (74-170)
Grenada	2010	Male	60 to 64	62% (41-79%)	38% (21-59%)	3.1 (1.7-5.3)	9.2% (5-13%)	11% (5.8-15%)	3 (1.6-4.5)	87 (47-130)
Grenada	2010	Male	65 to 69	60% (41-77%)	40% (23-59%)	2.8 (1.6-4.4)	8.3% (4.7-12%)	9.6% (5.5-14%)	2.9 (1.7-4.3)	70 (40-100)
Grenada	2010	Male	70 to 74	57% (38-73%)	43% (27-62%)	2.3 (1.2-3.9)	6.2% (3.1-10%)	7.1% (3.5-12%)	3.1 (1.5-5.1)	59 (29-98)
Grenada	2010	Male	75 to 79	52% (32-71%)	48% (29-68%)	1.9 (0.94-3.4)	5% (2.1-8.7%)	5.6% (2.3-9.9%)	2.4 (1-4.3)	36 (16-64)
Grenada	2010	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.6 (0.83-2.6)	4.1% (1.8-7%)	4.6% (1.9-8%)	2.2 (0.94-3.9)	24 (11-42)
Grenada	2010	Male	85 to 89	41% (25-58%)	59% (42-75%)	1.4 (0.71-2.5)	3.4% (1.5-6.4%)	3.8% (1.5-7.2%)	1.4 (0.56-2.6)	11 (4.5-20)
Grenada	2010	Male	90 to 94	41% (26-59%)	59% (41-74%)	1.4 (0.66-2.6)	3.1% (1.1-6.3%)	3.4% (1.1-7.1%)	0.49 (0.16-1)	2.7 (0.95-5.4)
Grenada	2010	Male	95 plus	44% (25-65%)	56% (35-75%)	1.5 (0.61-3)	3.2% (0.88-7%)	3.5% (0.72-7.9%)	0.12 (0.026-0.28)	0.46 (0.12-1)
Grenada	2016	Female	All Ages	33% (27-40%)	67% (60-73%)	1.2 (0.95-1.6)	2.5% (1.6-3.7%)	2.8% (1.3-4.6%)	12 (5.2-21)	360 (220-550)
Grenada	2016	Female	15 to 19	38% (25-54%)	62% (46-75%)	1.4 (0.7-2.3)	2% (1.2-3.2%)	4% (2-7.1%)	0.099 (0.048-0.18)	11 (6.6-19)
Grenada	2016	Female	20 to 24	44% (29-61%)	56% (39-71%)	2 (1.1-3.3)	3.5% (2.3-5%)	5.5% (2.9-8.8%)	0.2 (0.1-0.34)	23 (15-35)
Grenada	2016	Female	25 to 29	44% (27-61%)	56% (39-73%)	2.1 (1.2-3.4)	3.6% (2.4-5.3%)	5.6% (3.3-8.6%)	0.31 (0.17-0.54)	31 (20-47)
Grenada	2016	Female	30 to 34	42% (26-60%)	58% (40-74%)	2 (1.1-3.3)	3.6% (2.3-5.2%)	5.7% (3.5-8.6%)	0.39 (0.21-0.65)	32 (19-49)
Grenada	2016	Female	35 to 39	41% (26-59%)	59% (41-74%)	1.9 (1-3.1)	3.8% (2.3-5.6%)	6.1% (3.8-9.2%)	0.45 (0.24-0.76)	29 (17-47)
Grenada	2016	Female	40 to 44	40% (25-58%)	60% (42-75%)	1.8 (0.92-3)	3.9% (2.2-6.3%)	6.4% (3.8-9.9%)	0.5 (0.27-0.83)	27 (15-45)
Grenada	2016	Female	45 to 49	38% (22-55%)	62% (45-78%)	1.7 (0.89-3.1)	4.1% (2.2-7.1%)	6.3% (3.5-11%)	0.6 (0.31-1)	28 (15-49)
Grenada	2016	Female	50 to 54	34% (21-51%)	66% (49-79%)	1.5 (0.79-2.7)	3.9% (1.8-7%)	5.7% (2.9-9.9%)	1 (0.52-1.9)	42 (20-77)
Grenada	2016	Female	55 to 59	30% (18-46%)	70% (54-82%)	1.3 (0.66-2.3)	3.4% (1.5-6.3%)	4.8% (2.2-8.6%)	1.1 (0.5-2)	37 (16-70)
Grenada	2016	Female	60 to 64	27% (16-42%)	73% (58-84%)	1.1 (0.52-2)	2.8% (0.84-5.6%)	3.8% (1.3-7.4%)	0.9 (0.3-1.8)	26 (7.7-53)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	2016	Female	65 to 69	25% (15-38%)	75% (62-85%)	0.95 (0.49-1.6)	2.3% (0.67-4.6%)	3.1% (1-5.8%)	0.82 (0.27-1.6)	19 (5.6-38)
Grenada	2016	Female	70 to 74	22% (14-35%)	78% (65-86%)	0.81 (0.39-1.4)	1.9% (0.28-4.3%)	2.4% (0.45-5.3%)	0.89 (0.16-2)	17 (2.2-39)
Grenada	2016	Female	75 to 79	20% (12-32%)	80% (68-88%)	0.71 (0.31-1.3)	1.5% (0.059-3.7%)	1.9% (0.15-4.5%)	1 (0.076-2.5)	15 (0.55-36)
Grenada	2016	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.65 (0.31-1.2)	1.5% (0.21-3.4%)	1.8% (0.26-4%)	1.2 (0.17-2.7)	12 (1.7-29)
Grenada	2016	Female	85 to 89	18% (11-28%)	82% (72-89%)	0.66 (0.31-1.2)	1.5% (0.21-3.7%)	1.8% (0.22-4.2%)	1.1 (0.13-2.6)	8.1 (1.2-20)
Grenada	2016	Female	90 to 94	20% (12-32%)	80% (68-88%)	0.76 (0.32-1.5)	1.7% (0.2-4.4%)	1.9% (0.18-5.2%)	0.83 (0.076-2.2)	4.4 (0.53-12)
Grenada	2016	Female	95 plus	24% (12-40%)	76% (60-88%)	0.91 (0.32-2)	2% (0.049-5.5%)	2.3% (-0.062-6.4%)	0.34 (-0.0094-0.96)	1.2 (0.028-3.3)
Grenada	2016	Male	All Ages	56% (48-65%)	44% (35-52%)	2.7 (2.1-3.5)	8.5% (6.6-11%)	8.7% (6.1-11%)	39 (26-52)	1500 (1100-1900)
Grenada	2016	Male	15 to 19	55% (38-72%)	45% (28-62%)	2 (1-3.2)	5.8% (3.7-8.7%)	7.8% (4.4-12%)	0.54 (0.28-0.93)	48 (28-76)
Grenada	2016	Male	20 to 24	66% (48-81%)	34% (19-52%)	3.8 (2.2-5.6)	12% (8.1-16%)	15% (8.9-21%)	1.6 (0.86-2.6)	130 (80-190)
Grenada	2016	Male	25 to 29	69% (51-83%)	31% (17-49%)	4.5 (2.8-6.8)	13% (9-17%)	16% (11-22%)	2.2 (1.2-3.4)	170 (110-240)
Grenada	2016	Male	30 to 34	70% (51-84%)	30% (16-49%)	4.7 (2.8-7.2)	12% (8.8-16%)	16% (11-21%)	2.3 (1.3-3.4)	160 (98-220)
Grenada	2016	Male	35 to 39	70% (53-84%)	30% (16-47%)	4.5 (2.7-6.9)	12% (8.4-16%)	16% (11-20%)	2.1 (1.3-3.1)	130 (83-180)
Grenada	2016	Male	40 to 44	69% (52-83%)	31% (17-48%)	4.3 (2.3-6.8)	12% (8.2-16%)	15% (9.9-20%)	2.3 (1.4-3.3)	120 (79-170)
Grenada	2016	Male	45 to 49	67% (46-82%)	33% (18-54%)	4 (2.2-6.6)	12% (8.2-16%)	15% (10-19%)	2.5 (1.6-3.6)	120 (75-160)
Grenada	2016	Male	50 to 54	65% (46-80%)	35% (20-54%)	3.7 (2-6)	11% (7.4-15%)	14% (9-18%)	4 (2.4-5.6)	160 (99-220)
Grenada	2016	Male	55 to 59	63% (45-79%)	37% (21-55%)	3.4 (1.8-5.6)	10% (6.2-14%)	12% (7.3-17%)	4.2 (2.4-6.2)	140 (84-210)
Grenada	2016	Male	60 to 64	61% (41-79%)	39% (21-59%)	3 (1.6-5.4)	8.9% (4.8-13%)	10% (5.6-15%)	4 (2-6.2)	110 (60-180)
Grenada	2016	Male	65 to 69	60% (40-76%)	40% (24-60%)	2.7 (1.4-4.4)	8.1% (4.4-12%)	9.4% (5-14%)	3.6 (1.9-5.5)	85 (46-130)
Grenada	2016	Male	70 to 74	56% (38-73%)	44% (27-62%)	2.2 (1.1-3.9)	6.1% (2.9-10%)	6.9% (3.3-12%)	2.7 (1.2-4.5)	51 (24-86)
Grenada	2016	Male	75 to 79	51% (32-70%)	49% (30-68%)	1.9 (0.85-3.3)	4.8% (2-8.5%)	5.4% (2.1-9.6%)	2.6 (1-4.7)	39 (16-69)
Grenada	2016	Male	80 to 84	45% (27-63%)	55% (37-73%)	1.5 (0.77-2.6)	3.9% (1.6-7.1%)	4.4% (1.7-7.9%)	2.1 (0.77-3.8)	23 (8.8-41)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Grenada	2016	Male	85 to 89	40% (24-57%)	60% (43-76%)	1.4 (0.66-2.4)	3.3% (1.3-6.4%)	3.7% (1.4-7.2%)	1.3 (0.48-2.5)	10 (3.9-20)
Grenada	2016	Male	90 to 94	40% (26-58%)	60% (42-74%)	1.4 (0.61-2.6)	2.9% (0.98-6.3%)	3.2% (0.92-7%)	0.62 (0.18-1.4)	3.4 (1.1-7.3)
Grenada	2016	Male	95 plus	44% (25-65%)	56% (35-75%)	1.5 (0.58-3)	3.1% (0.81-7%)	3.3% (0.66-7.9%)	0.15 (0.03-0.36)	0.56 (0.15-1.3)
Guyana	1990	Female	All Ages	29% (23-36%)	71% (64-77%)	0.46 (0.37-0.59)	1.2% (0.83-1.7%)	1.3% (0.66-2.1%)	31 (15-50)	1600 (1100-2200)
Guyana	1990	Female	15 to 19	35% (22-51%)	65% (49-78%)	0.53 (0.32-0.83)	1.7% (1-2.5%)	2.7% (1.4-4.5%)	1.1 (0.58-1.9)	110 (67-170)
Guyana	1990	Female	20 to 24	41% (26-58%)	59% (42-74%)	0.82 (0.5-1.2)	2.9% (2-4%)	3.7% (2.2-5.8%)	2.2 (1.3-3.4)	220 (150-310)
Guyana	1990	Female	25 to 29	41% (25-58%)	59% (42-75%)	0.87 (0.55-1.3)	3% (2.1-4.1%)	3.8% (2.5-5.5%)	2.5 (1.6-3.6)	220 (160-300)
Guyana	1990	Female	30 to 34	39% (23-56%)	61% (44-77%)	0.87 (0.51-1.3)	3% (2.2-4%)	4.2% (2.9-5.7%)	2.6 (1.9-3.7)	200 (150-270)
Guyana	1990	Female	35 to 39	38% (24-55%)	62% (45-76%)	0.84 (0.5-1.3)	3.1% (2.3-4.1%)	4.4% (3.2-6%)	3 (2.1-4.2)	190 (140-260)
Guyana	1990	Female	40 to 44	37% (23-55%)	63% (45-77%)	0.8 (0.47-1.2)	2.9% (1.9-4%)	4.1% (2.7-5.9%)	3.3 (2.1-4.7)	170 (120-240)
Guyana	1990	Female	45 to 49	35% (20-52%)	65% (48-80%)	0.74 (0.39-1.2)	2.5% (1.5-3.9%)	3.5% (2-5.5%)	3.1 (1.8-4.8)	140 (82-220)
Guyana	1990	Female	50 to 54	32% (19-48%)	68% (52-81%)	0.67 (0.36-1.1)	2.3% (0.95-3.9%)	3% (1.2-5.2%)	3.1 (1.3-5.4)	120 (51-210)
Guyana	1990	Female	55 to 59	28% (16-43%)	72% (57-84%)	0.59 (0.31-1)	1.9% (0.58-3.7%)	2.4% (0.7-4.7%)	3.3 (0.95-6.4)	110 (33-210)
Guyana	1990	Female	60 to 64	24% (14-39%)	76% (61-86%)	0.52 (0.27-0.92)	1.2% (-0.037-2.9%)	1.5% (-0.046-3.5%)	2.4 (-0.077-5.8)	67 (-2-160)
Guyana	1990	Female	65 to 69	22% (13-35%)	78% (65-87%)	0.44 (0.24-0.73)	0.76% (-0.31-2%)	0.94% (-0.38-2.5%)	2 (-0.81-5.3)	46 (-18-120)
Guyana	1990	Female	70 to 74	20% (12-32%)	80% (68-88%)	0.37 (0.2-0.65)	0.5% (-0.44-1.6%)	0.61% (-0.52-1.9%)	1.1 (-0.94-3.6)	21 (-18-64)
Guyana	1990	Female	75 to 79	18% (10-29%)	82% (71-90%)	0.32 (0.16-0.57)	0.32% (-0.55-1.2%)	0.37% (-0.67-1.4%)	0.49 (-0.88-1.9)	6.9 (-12-26)
Guyana	1990	Female	80 to 84	16% (9.1-26%)	84% (74-91%)	0.29 (0.15-0.49)	0.32% (-0.33-1%)	0.38% (-0.41-1.3%)	0.63 (-0.71-2.1)	6.9 (-7.2-22)
Guyana	1990	Female	85 to 89	16% (9.2-25%)	84% (75-91%)	0.29 (0.15-0.5)	0.25% (-0.33-0.88%)	0.26% (-0.44-1%)	0.38 (-0.67-1.5)	3.3 (-4.5-12)
Guyana	1990	Female	90 to 94	17% (10-28%)	83% (72-90%)	0.33 (0.16-0.58)	0.23% (-0.45-1%)	0.2% (-0.6-1.2%)	0.12 (-0.36-0.69)	0.83 (-1.6-3.7)
Guyana	1990	Female	95 plus	21% (11-35%)	79% (65-89%)	0.39 (0.16-0.74)	0.22% (-0.71-1.3%)	0.14% (-0.98-1.5%)	0.022 (-0.16-0.24)	0.14 (-0.46-0.86)
Guyana	1990	Male	All Ages	56% (47-64%)	44% (36-53%)	3.9 (3.4-4.4)	10% (8.7-12%)	14% (12-17%)	460 (380-540)	18000 (15000-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	1990	Male	15 to 19	59% (42-74%)	41% (26-58%)	3.6 (2.5-4.8)	11% (7.6-14%)	16% (11-21%)	10 (7.1-14)	800 (580-1100)
Guyana	1990	Male	20 to 24	69% (52-83%)	31% (17-48%)	6.9 (5.2-8.8)	21% (16-26%)	27% (20-33%)	30 (22-38)	2100 (1600-2700)
Guyana	1990	Male	25 to 29	72% (55-84%)	28% (16-45%)	7.7 (5.9-9.6)	21% (17-26%)	27% (20-32%)	33 (25-41)	2200 (1700-2700)
Guyana	1990	Male	30 to 34	73% (55-85%)	27% (15-45%)	7.7 (5.6-9.9)	20% (16-24%)	26% (20-31%)	33 (27-41)	2000 (1600-2400)
Guyana	1990	Male	35 to 39	72% (56-85%)	28% (15-44%)	7.2 (5.2-9.3)	20% (16-24%)	26% (21-30%)	29 (23-35)	1600 (1300-1900)
Guyana	1990	Male	40 to 44	71% (56-84%)	29% (16-44%)	6.8 (4.9-9.1)	20% (16-24%)	25% (21-30%)	30 (25-37)	1500 (1200-1800)
Guyana	1990	Male	45 to 49	70% (51-84%)	30% (16-49%)	6.4 (4.3-8.8)	21% (17-25%)	25% (20-30%)	39 (30-47)	1600 (1300-2000)
Guyana	1990	Male	50 to 54	68% (50-83%)	32% (17-50%)	5.7 (3.8-8.1)	19% (15-24%)	23% (18-28%)	37 (29-47)	1400 (1100-1700)
Guyana	1990	Male	55 to 59	66% (49-81%)	34% (19-51%)	5 (3-7.7)	18% (13-23%)	21% (15-27%)	41 (30-52)	1300 (960-1700)
Guyana	1990	Male	60 to 64	65% (45-82%)	35% (18-55%)	4.5 (2.6-7.3)	16% (11-21%)	18% (12-24%)	46 (31-62)	1300 (840-1700)
Guyana	1990	Male	65 to 69	63% (44-79%)	37% (21-56%)	4.1 (2.4-6.3)	15% (11-20%)	17% (12-23%)	44 (30-60)	1000 (700-1400)
Guyana	1990	Male	70 to 74	60% (42-76%)	40% (24-58%)	3.4 (2-5.5)	12% (7.7-17%)	14% (8.7-20%)	36 (23-51)	660 (420-930)
Guyana	1990	Male	75 to 79	55% (35-74%)	45% (26-65%)	2.9 (1.5-4.8)	10% (5.7-15%)	11% (6.2-17%)	22 (12-33)	310 (170-460)
Guyana	1990	Male	80 to 84	49% (31-67%)	51% (33-69%)	2.4 (1.4-4)	8% (4.6-12%)	9.3% (5.3-14%)	15 (8.2-23)	160 (90-240)
Guyana	1990	Male	85 to 89	45% (28-62%)	55% (38-72%)	2.1 (1.2-3.6)	6.7% (3.7-10%)	7.7% (4.2-12%)	9.9 (5.4-16)	76 (42-120)
Guyana	1990	Male	90 to 94	45% (29-63%)	55% (37-71%)	2.1 (1.1-3.6)	6% (2.9-9.9%)	6.9% (3.2-11%)	3 (1.4-5)	16 (7.4-26)
Guyana	1990	Male	95 plus	48% (28-68%)	52% (32-72%)	2.2 (0.98-3.9)	6.2% (2.4-11%)	7% (2.4-13%)	0.71 (0.24-1.3)	2.5 (0.95-4.4)
Guyana	1995	Female	All Ages	29% (23-36%)	71% (64-77%)	0.53 (0.41-0.66)	1.3% (0.93-1.8%)	1.4% (0.67-2.3%)	37 (17-60)	1800 (1200-2500)
Guyana	1995	Female	15 to 19	35% (23-51%)	65% (49-77%)	0.62 (0.37-0.96)	1.7% (1.1-2.6%)	2.8% (1.5-4.8%)	1.1 (0.59-1.9)	110 (70-170)
Guyana	1995	Female	20 to 24	41% (26-58%)	59% (42-74%)	0.94 (0.57-1.4)	2.9% (2-3.9%)	3.6% (2.2-5.6%)	2.3 (1.4-3.6)	230 (160-320)
Guyana	1995	Female	25 to 29	41% (25-58%)	59% (42-75%)	1 (0.63-1.5)	3.1% (2.2-4.2%)	3.8% (2.4-5.4%)	2.8 (1.8-4.2)	240 (170-330)
Guyana	1995	Female	30 to 34	39% (24-56%)	61% (44-76%)	1 (0.58-1.5)	3% (2.2-4%)	4% (2.7-5.4%)	2.8 (1.9-4)	210 (150-280)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	1995	Female	35 to 39	38% (24-55%)	62% (45-76%)	0.97 (0.57-1.5)	3.2% (2.3-4.3%)	4.5% (3.2-6.2%)	3.3 (2.3-4.5)	200 (150-270)
Guyana	1995	Female	40 to 44	37% (23-55%)	63% (45-77%)	0.92 (0.53-1.4)	3% (2.1-4.3%)	4.4% (3-6.3%)	3.4 (2.2-4.8)	180 (120-250)
Guyana	1995	Female	45 to 49	35% (21-52%)	65% (48-79%)	0.85 (0.46-1.3)	2.7% (1.5-4.1%)	3.7% (2.1-5.8%)	3.3 (1.9-5.2)	150 (86-230)
Guyana	1995	Female	50 to 54	32% (19-48%)	68% (52-81%)	0.76 (0.42-1.2)	2.5% (1.1-4.3%)	3.3% (1.5-5.7%)	3.5 (1.6-6.1)	130 (62-230)
Guyana	1995	Female	55 to 59	28% (16-43%)	72% (57-84%)	0.68 (0.36-1.2)	2.2% (0.76-4.2%)	2.9% (1-5.4%)	4 (1.4-7.5)	130 (45-240)
Guyana	1995	Female	60 to 64	25% (14-40%)	75% (60-86%)	0.59 (0.31-1.1)	1.5% (0.098-3.3%)	1.9% (0.13-4.1%)	2.7 (0.19-6.2)	76 (5.2-170)
Guyana	1995	Female	65 to 69	22% (14-35%)	78% (65-86%)	0.51 (0.28-0.84)	0.96% (-0.15-2.4%)	1.2% (-0.17-2.9%)	2.4 (-0.36-5.8)	55 (-8.3-130)
Guyana	1995	Female	70 to 74	20% (12-32%)	80% (68-88%)	0.42 (0.23-0.75)	0.62% (-0.42-1.9%)	0.77% (-0.46-2.3%)	2 (-1.2-6)	35 (-25-110)
Guyana	1995	Female	75 to 79	18% (10-29%)	82% (71-90%)	0.36 (0.18-0.64)	0.39% (-0.57-1.3%)	0.47% (-0.66-1.6%)	1.8 (-2.6-6.3)	25 (-38-89)
Guyana	1995	Female	80 to 84	16% (9.1-26%)	84% (74-91%)	0.34 (0.18-0.56)	0.42% (-0.3-1.2%)	0.5% (-0.37-1.4%)	0.83 (-0.6-2.4)	8.9 (-6.3-25)
Guyana	1995	Female	85 to 89	16% (9.3-25%)	84% (75-91%)	0.34 (0.17-0.59)	0.39% (-0.24-1.2%)	0.43% (-0.33-1.4%)	0.43 (-0.33-1.4)	3.5 (-2.1-11)
Guyana	1995	Female	90 to 94	18% (11-28%)	82% (72-89%)	0.38 (0.19-0.68)	0.39% (-0.34-1.4%)	0.38% (-0.49-1.5%)	0.33 (-0.42-1.3)	2 (-1.8-7)
Guyana	1995	Female	95 plus	21% (11-35%)	79% (65-89%)	0.45 (0.19-0.86)	0.41% (-0.58-1.8%)	0.35% (-0.82-2%)	0.056 (-0.13-0.31)	0.25 (-0.36-1.1)
Guyana	1995	Male	All Ages	55% (47-64%)	45% (36-53%)	4.3 (3.7-5)	11% (9.6-13%)	15% (13-18%)	510 (420-610)	20000 (17000-23000)
Guyana	1995	Male	15 to 19	59% (41-74%)	41% (26-59%)	4.1 (2.8-5.5)	12% (8.3-15%)	17% (12-23%)	10 (7.1-14)	810 (590-1100)
Guyana	1995	Male	20 to 24	69% (51-82%)	31% (18-49%)	7.8 (5.8-10)	21% (16-26%)	26% (19-33%)	27 (19-34)	1900 (1400-2400)
Guyana	1995	Male	25 to 29	72% (55-84%)	28% (16-45%)	8.6 (6.6-11)	21% (16-25%)	25% (19-31%)	35 (26-43)	2300 (1700-2800)
Guyana	1995	Male	30 to 34	72% (55-85%)	28% (15-45%)	8.7 (6.3-11)	19% (16-23%)	24% (19-29%)	33 (26-40)	2000 (1600-2400)
Guyana	1995	Male	35 to 39	72% (56-85%)	28% (15-44%)	8 (5.8-11)	19% (15-23%)	24% (19-28%)	32 (25-39)	1800 (1400-2100)
Guyana	1995	Male	40 to 44	71% (56-84%)	29% (16-44%)	7.6 (5.4-10)	20% (16-24%)	24% (20-29%)	40 (33-49)	2000 (1600-2400)
Guyana	1995	Male	45 to 49	70% (50-83%)	30% (17-50%)	7.1 (4.8-9.7)	20% (16-24%)	24% (20-29%)	43 (34-53)	1900 (1500-2300)
Guyana	1995	Male	50 to 54	68% (50-83%)	32% (17-50%)	6.5 (4.4-9.1)	19% (15-24%)	23% (18-28%)	43 (33-53)	1600 (1300-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	1995	Male	55 to 59	66% (49-82%)	34% (18-51%)	5.8 (3.5-9)	19% (14-23%)	22% (16-27%)	44 (33-56)	1400 (1100-1800)
Guyana	1995	Male	60 to 64	65% (46-81%)	35% (19-54%)	5.2 (3-8.6)	16% (12-21%)	19% (14-24%)	46 (33-60)	1300 (890-1700)
Guyana	1995	Male	65 to 69	63% (45-79%)	37% (21-55%)	4.6 (2.7-7)	16% (11-20%)	18% (13-23%)	45 (32-60)	1000 (740-1400)
Guyana	1995	Male	70 to 74	60% (41-75%)	40% (25-59%)	3.9 (2.2-6.2)	13% (8.9-18%)	15% (10-21%)	39 (27-54)	720 (490-1000)
Guyana	1995	Male	75 to 79	55% (35-73%)	45% (27-65%)	3.2 (1.8-5.3)	9.9% (6.1-14%)	12% (7.3-17%)	28 (17-41)	410 (250-590)
Guyana	1995	Male	80 to 84	49% (30-67%)	51% (33-70%)	2.7 (1.5-4.4)	9.3% (5.7-14%)	10% (6.3-15%)	22 (13-32)	230 (140-340)
Guyana	1995	Male	85 to 89	44% (27-62%)	56% (38-73%)	2.4 (1.3-3.9)	7.8% (4.4-12%)	8.8% (5-13%)	9.7 (5.4-15)	73 (41-110)
Guyana	1995	Male	90 to 94	45% (29-62%)	55% (38-71%)	2.4 (1.3-4)	7% (3.5-11%)	7.9% (3.8-13%)	5.6 (2.7-9)	29 (14-47)
Guyana	1995	Male	95 plus	48% (28-68%)	52% (32-72%)	2.5 (1.1-4.5)	7.2% (2.9-12%)	8% (3-14%)	0.92 (0.35-1.6)	3.2 (1.3-5.5)
Guyana	2000	Female	All Ages	28% (23-35%)	72% (65-77%)	0.45 (0.36-0.57)	1.2% (0.77-1.6%)	1.1% (0.5-1.9%)	26 (11-42)	1500 (980-2000)
Guyana	2000	Female	15 to 19	35% (22-51%)	65% (49-78%)	0.52 (0.31-0.82)	1.5% (0.92-2.3%)	2.5% (1.3-4.4%)	0.71 (0.35-1.2)	76 (46-120)
Guyana	2000	Female	20 to 24	41% (26-57%)	59% (43-74%)	0.79 (0.48-1.2)	2.7% (1.8-3.9%)	3.3% (1.9-5.3%)	2.1 (1.2-3.4)	200 (140-300)
Guyana	2000	Female	25 to 29	40% (24-57%)	60% (43-76%)	0.84 (0.52-1.3)	2.8% (2-3.8%)	3.3% (2-4.8%)	2.8 (1.8-4.2)	240 (170-330)
Guyana	2000	Female	30 to 34	39% (23-56%)	61% (44-77%)	0.84 (0.49-1.3)	2.7% (2-3.6%)	3.5% (2.5-4.9%)	3.1 (2.1-4.3)	220 (160-300)
Guyana	2000	Female	35 to 39	38% (24-55%)	62% (45-76%)	0.8 (0.47-1.2)	2.8% (2.1-3.7%)	3.9% (2.8-5.3%)	3.6 (2.6-5)	220 (160-300)
Guyana	2000	Female	40 to 44	36% (23-55%)	64% (45-77%)	0.76 (0.45-1.2)	2.3% (1.5-3.3%)	3.4% (2.2-4.8%)	3.2 (2.1-4.6)	170 (120-240)
Guyana	2000	Female	45 to 49	35% (20-52%)	65% (48-80%)	0.71 (0.38-1.1)	1.8% (0.88-3.1%)	2.5% (1.2-4.2%)	2.8 (1.3-4.6)	130 (60-210)
Guyana	2000	Female	50 to 54	31% (19-48%)	69% (52-81%)	0.65 (0.35-1.1)	1.4% (0.26-2.8%)	1.8% (0.33-3.7%)	2.3 (0.42-4.7)	89 (17-180)
Guyana	2000	Female	55 to 59	27% (16-43%)	73% (57-84%)	0.58 (0.32-0.98)	1.2% (-0.12-2.7%)	1.5% (-0.11-3.4%)	1.8 (-0.13-4.1)	59 (-6.5-140)
Guyana	2000	Female	60 to 64	24% (14-39%)	76% (61-86%)	0.51 (0.26-0.89)	0.69% (-0.52-2.1%)	0.89% (-0.62-2.6%)	1.4 (-0.91-4.1)	37 (-28-110)
Guyana	2000	Female	65 to 69	22% (13-34%)	78% (66-87%)	0.44 (0.24-0.72)	0.31% (-0.74-1.4%)	0.41% (-0.84-1.7%)	0.64 (-1.3-2.6)	14 (-31-59)
Guyana	2000	Female	70 to 74	20% (12-31%)	80% (69-88%)	0.36 (0.19-0.63)	0.17% (-0.75-1.2%)	0.24% (-0.84-1.4%)	0.5 (-1.8-3)	8 (-36-55)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	2000	Female	75 to 79	17% (10-28%)	83% (72-90%)	0.31 (0.15-0.54)	0.083% (-0.73-0.86%)	0.12% (-0.85-1.1%)	0.23 (-1.6-2.1)	2.8 (-24-29)
Guyana	2000	Female	80 to 84	16% (9.2-26%)	84% (74-91%)	0.29 (0.15-0.49)	0.19% (-0.46-0.83%)	0.24% (-0.55-1.1%)	0.46 (-1.1-2)	4.8 (-12-21)
Guyana	2000	Female	85 to 89	16% (9.3-25%)	84% (75-91%)	0.29 (0.15-0.51)	0.21% (-0.35-0.79%)	0.23% (-0.46-0.98%)	0.17 (-0.33-0.73)	1.4 (-2.3-5.5)
Guyana	2000	Female	90 to 94	17% (10-27%)	83% (73-90%)	0.32 (0.16-0.57)	0.18% (-0.44-0.98%)	0.15% (-0.6-1.1%)	0.061 (-0.25-0.46)	0.44 (-1.1-2.4)
Guyana	2000	Female	95 plus	20% (11-34%)	80% (66-89%)	0.37 (0.15-0.7)	0.16% (-0.66-1.3%)	0.072% (-0.87-1.4%)	0.017 (-0.21-0.33)	0.15 (-0.62-1.2)
Guyana	2000	Male	All Ages	56% (47-64%)	44% (36-53%)	3.9 (3.3-4.5)	12% (9.8-14%)	15% (12-17%)	460 (380-550)	19000 (16000-22000)
Guyana	2000	Male	15 to 19	58% (41-74%)	42% (26-59%)	3.5 (2.4-4.8)	11% (7.5-14%)	15% (11-21%)	8.3 (5.6-11)	660 (470-870)
Guyana	2000	Male	20 to 24	68% (51-82%)	32% (18-49%)	6.8 (5-8.7)	20% (15-25%)	26% (19-32%)	26 (19-33)	1900 (1400-2400)
Guyana	2000	Male	25 to 29	71% (54-84%)	29% (16-46%)	7.5 (5.8-9.5)	21% (16-25%)	25% (19-31%)	39 (29-49)	2600 (2000-3200)
Guyana	2000	Male	30 to 34	72% (54-85%)	28% (15-46%)	7.5 (5.5-9.6)	20% (16-24%)	24% (19-28%)	45 (35-55)	2700 (2100-3300)
Guyana	2000	Male	35 to 39	72% (56-84%)	28% (16-44%)	7 (5-9.2)	19% (15-22%)	23% (18-28%)	37 (29-46)	2000 (1600-2500)
Guyana	2000	Male	40 to 44	71% (55-84%)	29% (16-45%)	6.6 (4.8-8.9)	19% (15-22%)	23% (19-28%)	39 (31-48)	1900 (1600-2400)
Guyana	2000	Male	45 to 49	69% (50-83%)	31% (17-50%)	6.2 (4.2-8.3)	18% (15-22%)	22% (18-26%)	35 (27-43)	1500 (1200-1900)
Guyana	2000	Male	50 to 54	68% (50-83%)	32% (17-50%)	5.6 (3.7-7.8)	18% (14-22%)	21% (16-26%)	41 (31-52)	1600 (1200-2000)
Guyana	2000	Male	55 to 59	66% (48-81%)	34% (19-52%)	4.9 (2.8-7.7)	16% (12-21%)	19% (13-25%)	36 (25-47)	1200 (810-1500)
Guyana	2000	Male	60 to 64	64% (44-81%)	36% (19-56%)	4.3 (2.5-7)	14% (9.7-19%)	17% (11-22%)	38 (25-51)	1000 (700-1400)
Guyana	2000	Male	65 to 69	63% (44-79%)	37% (21-56%)	3.9 (2.3-6)	14% (9.3-18%)	16% (11-21%)	35 (24-48)	810 (550-1100)
Guyana	2000	Male	70 to 74	59% (41-75%)	41% (25-59%)	3.3 (1.8-5.2)	11% (6.6-16%)	13% (7.6-19%)	35 (20-51)	640 (380-940)
Guyana	2000	Male	75 to 79	54% (35-73%)	46% (27-65%)	2.7 (1.5-4.5)	8.8% (4.9-13%)	10% (5.6-15%)	21 (12-32)	310 (170-460)
Guyana	2000	Male	80 to 84	48% (30-66%)	52% (34-70%)	2.3 (1.3-3.7)	7.5% (4.2-11%)	8.7% (4.8-13%)	16 (9.1-25)	170 (97-260)
Guyana	2000	Male	85 to 89	44% (27-61%)	56% (39-73%)	2 (1.1-3.3)	6.2% (3.4-9.5%)	7.1% (3.8-11%)	3.8 (2.1-5.9)	30 (16-46)
Guyana	2000	Male	90 to 94	44% (28-61%)	56% (39-72%)	2 (1-3.4)	5.8% (2.8-9.7%)	6.5% (3-11%)	2.4 (1.1-4)	13 (5.9-21)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	2000	Male	95 plus	47% (28-68%)	53% (32-72%)	2.1 (0.94-3.9)	6.1% (2.2-11%)	6.7% (2.2-12%)	1 (0.33-1.8)	3.5 (1.3-6.2)
Guyana	2005	Female	All Ages	23% (19-29%)	77% (71-81%)	0.3 (0.25-0.37)	0.93% (0.61-1.3%)	0.81% (0.25-1.4%)	20 (6.3-35)	1200 (770-1600)
Guyana	2005	Female	15 to 19	27% (18-41%)	73% (59-82%)	0.33 (0.2-0.52)	1.2% (0.71-2%)	1.7% (0.67-3.3%)	0.57 (0.22-1.1)	62 (36-99)
Guyana	2005	Female	20 to 24	32% (21-46%)	68% (54-79%)	0.48 (0.31-0.72)	2% (1.3-2.9%)	2.2% (1.2-3.5%)	1.3 (0.66-2.1)	130 (85-190)
Guyana	2005	Female	25 to 29	33% (23-49%)	67% (51-77%)	0.55 (0.36-0.8)	2% (1.4-2.6%)	2% (1.3-3%)	1.6 (1-2.4)	150 (110-210)
Guyana	2005	Female	30 to 34	32% (21-47%)	68% (53-79%)	0.55 (0.35-0.81)	2% (1.5-2.6%)	2.3% (1.6-3.1%)	2 (1.4-2.9)	160 (120-220)
Guyana	2005	Female	35 to 39	31% (22-46%)	69% (54-78%)	0.53 (0.35-0.78)	2.1% (1.6-2.7%)	2.7% (2-3.6%)	2.7 (1.9-3.7)	170 (130-230)
Guyana	2005	Female	40 to 44	32% (22-47%)	68% (53-78%)	0.54 (0.36-0.8)	2% (1.4-2.7%)	2.8% (2-3.8%)	3.2 (2.2-4.4)	170 (120-230)
Guyana	2005	Female	45 to 49	31% (20-45%)	69% (55-80%)	0.51 (0.32-0.77)	1.6% (0.93-2.5%)	2.2% (1.3-3.4%)	3.2 (1.8-4.8)	140 (85-220)
Guyana	2005	Female	50 to 54	29% (17-44%)	71% (56-83%)	0.48 (0.27-0.76)	1.3% (0.39-2.3%)	1.8% (0.66-3.1%)	2.7 (0.99-4.7)	100 (29-180)
Guyana	2005	Female	55 to 59	27% (15-42%)	73% (58-85%)	0.46 (0.25-0.78)	0.98% (-0.1-2%)	1.3% (-0.0035-2.6%)	2.3 (-0.0061-4.6)	72 (-7.5-150)
Guyana	2005	Female	60 to 64	24% (14-38%)	76% (62-86%)	0.4 (0.21-0.71)	0.42% (-0.61-1.5%)	0.58% (-0.68-1.8%)	1 (-1.2-3.2)	26 (-36-87)
Guyana	2005	Female	65 to 69	21% (13-34%)	79% (66-87%)	0.35 (0.19-0.57)	0.043% (-0.84-0.81%)	0.11% (-0.92-1.1%)	0.23 (-1.9-2.2)	2.6 (-48-50)
Guyana	2005	Female	70 to 74	19% (12-31%)	81% (69-88%)	0.29 (0.15-0.51)	-0.047% (-0.91-0.75%)	-0.017% (-1-0.89%)	-0.032 (-1.9-1.8)	-1.9 (-38-31)
Guyana	2005	Female	75 to 79	17% (9.8-28%)	83% (72-90%)	0.25 (0.12-0.43)	-0.11% (-0.91-0.53%)	-0.11% (-1.1-0.64%)	-0.27 (-2.5-1.6)	-4.5 (-37-22)
Guyana	2005	Female	80 to 84	15% (8.7-25%)	85% (75-91%)	0.23 (0.12-0.38)	-0.048% (-0.73-0.46%)	-0.055% (-0.84-0.56%)	-0.11 (-1.7-1.1)	-1.2 (-18-12)
Guyana	2005	Female	85 to 89	15% (9-24%)	85% (76-91%)	0.23 (0.12-0.38)	0.032% (-0.46-0.53%)	0.014% (-0.62-0.65%)	0.021 (-0.92-0.99)	0.46 (-6.6-7.6)
Guyana	2005	Female	90 to 94	17% (9.9-27%)	83% (73-90%)	0.25 (0.13-0.45)	-0.042% (-0.63-0.58%)	-0.1% (-0.78-0.61%)	-0.071 (-0.54-0.42)	-0.17 (-2.6-2.3)
Guyana	2005	Female	95 plus	20% (10-33%)	80% (67-90%)	0.3 (0.13-0.56)	-0.065% (-0.79-0.72%)	-0.18% (-1-0.77%)	-0.028 (-0.17-0.12)	-0.04 (-0.49-0.45)
Guyana	2005	Male	All Ages	57% (49-65%)	43% (35-51%)	3.2 (2.9-3.6)	12% (10-14%)	14% (12-16%)	480 (390-570)	20000 (16000-23000)
Guyana	2005	Male	15 to 19	57% (42-72%)	43% (28-58%)	2.8 (2-3.7)	9.9% (7-13%)	13% (9.2-18%)	8.2 (5.6-11)	640 (450-850)
Guyana	2005	Male	20 to 24	71% (56-83%)	29% (17-44%)	5.9 (4.6-7.2)	21% (16-26%)	25% (19-32%)	23 (17-30)	1600 (1200-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	2005	Male	25 to 29	74% (61-85%)	26% (15-39%)	6.6 (5.4-7.9)	21% (16-26%)	25% (19-31%)	32 (24-40)	2100 (1600-2600)
Guyana	2005	Male	30 to 34	74% (61-85%)	26% (15-39%)	6.5 (5.1-7.9)	19% (15-22%)	22% (18-27%)	36 (28-44)	2200 (1700-2600)
Guyana	2005	Male	35 to 39	74% (61-85%)	26% (15-39%)	6 (4.8-7.3)	17% (14-20%)	21% (17-25%)	40 (32-49)	2200 (1800-2700)
Guyana	2005	Male	40 to 44	73% (61-84%)	27% (16-39%)	5.8 (4.6-7.2)	18% (15-21%)	22% (19-25%)	47 (38-57)	2300 (1900-2800)
Guyana	2005	Male	45 to 49	73% (57-84%)	27% (16-43%)	5.5 (4.2-6.9)	18% (15-21%)	22% (19-26%)	53 (43-64)	2300 (1900-2800)
Guyana	2005	Male	50 to 54	69% (52-82%)	31% (18-48%)	4.6 (3.2-6.1)	17% (13-21%)	20% (16-24%)	53 (42-66)	2000 (1600-2500)
Guyana	2005	Male	55 to 59	65% (47-81%)	35% (19-53%)	3.9 (2.3-6)	15% (10-20%)	17% (12-23%)	46 (31-61)	1500 (1000-2000)
Guyana	2005	Male	60 to 64	64% (44-80%)	36% (20-56%)	3.5 (2-5.5)	13% (8.1-18%)	15% (9.2-21%)	35 (22-49)	970 (600-1300)
Guyana	2005	Male	65 to 69	62% (43-78%)	38% (22-57%)	3.1 (1.9-4.7)	12% (7.4-16%)	14% (8.5-19%)	37 (23-52)	860 (540-1200)
Guyana	2005	Male	70 to 74	59% (40-74%)	41% (26-60%)	2.6 (1.5-4.2)	9.4% (5.1-14%)	10% (5.6-16%)	24 (13-37)	450 (240-680)
Guyana	2005	Male	75 to 79	54% (34-72%)	46% (28-66%)	2.2 (1.2-3.6)	7.3% (3.4-12%)	8.2% (3.8-13%)	21 (9.7-34)	300 (140-480)
Guyana	2005	Male	80 to 84	48% (30-66%)	52% (34-70%)	1.8 (1-2.9)	5.9% (3-9.6%)	6.7% (3.4-11%)	13 (6.4-21)	140 (69-220)
Guyana	2005	Male	85 to 89	43% (27-60%)	57% (40-73%)	1.6 (0.89-2.6)	4.6% (2.5-7.6%)	5.3% (2.8-8.7%)	5.6 (2.9-9.2)	44 (23-72)
Guyana	2005	Male	90 to 94	43% (28-61%)	57% (39-72%)	1.6 (0.87-2.7)	4.4% (1.9-7.8%)	4.7% (2-8.7%)	1.9 (0.8-3.4)	9.9 (4.4-18)
Guyana	2005	Male	95 plus	46% (27-66%)	54% (34-73%)	1.7 (0.75-3.1)	4.6% (1.5-8.9%)	5% (1.4-9.9%)	0.55 (0.16-1.1)	1.9 (0.61-3.7)
Guyana	2010	Female	All Ages	23% (19-29%)	77% (71-81%)	0.27 (0.22-0.32)	0.88% (0.55-1.2%)	0.71% (0.2-1.3%)	18 (5-33)	1100 (690-1600)
Guyana	2010	Female	15 to 19	27% (17-41%)	73% (59-83%)	0.28 (0.17-0.45)	1.1% (0.59-1.9%)	1.6% (0.49-3.1%)	0.69 (0.21-1.4)	77 (41-130)
Guyana	2010	Female	20 to 24	31% (20-45%)	69% (55-80%)	0.42 (0.26-0.63)	1.9% (1.2-2.8%)	2.1% (1.1-3.4%)	0.93 (0.47-1.6)	100 (64-150)
Guyana	2010	Female	25 to 29	32% (21-47%)	68% (53-79%)	0.47 (0.31-0.7)	1.9% (1.4-2.6%)	2% (1.3-3%)	1.1 (0.69-1.7)	110 (75-150)
Guyana	2010	Female	30 to 34	31% (20-46%)	69% (54-80%)	0.47 (0.3-0.72)	2% (1.5-2.7%)	2.4% (1.6-3.3%)	1.9 (1.3-2.7)	150 (110-200)
Guyana	2010	Female	35 to 39	31% (21-46%)	69% (54-79%)	0.46 (0.3-0.7)	2.2% (1.6-2.8%)	2.9% (2.1-3.8%)	2.8 (2-3.8)	180 (130-230)
Guyana	2010	Female	40 to 44	32% (22-47%)	68% (53-78%)	0.47 (0.31-0.69)	1.9% (1.3-2.5%)	2.7% (1.9-3.7%)	3.2 (2.2-4.5)	170 (120-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	2010	Female	45 to 49	30% (20-44%)	70% (56-80%)	0.44 (0.27-0.67)	1.5% (0.79-2.2%)	2.1% (1.2-3.2%)	3.2 (1.7-4.8)	140 (78-220)
Guyana	2010	Female	50 to 54	29% (17-44%)	71% (56-83%)	0.42 (0.24-0.66)	1.2% (0.32-2%)	1.7% (0.6-2.9%)	3.1 (1.1-5.4)	120 (32-200)
Guyana	2010	Female	55 to 59	26% (16-42%)	74% (58-84%)	0.4 (0.22-0.66)	0.82% (-0.14-1.7%)	1.1% (-0.032-2.2%)	2.3 (-0.064-4.6)	72 (-14-150)
Guyana	2010	Female	60 to 64	23% (13-37%)	77% (63-87%)	0.35 (0.18-0.61)	0.32% (-0.62-1.2%)	0.46% (-0.65-1.5%)	0.97 (-1.4-3.3)	24 (-47-87)
Guyana	2010	Female	65 to 69	21% (12-33%)	79% (67-88%)	0.3 (0.16-0.5)	-0.054% (-0.9-0.65%)	-0.02% (-1-0.83%)	-0.042 (-2.3-1.9)	-3.3 (-55-41)
Guyana	2010	Female	70 to 74	19% (12-31%)	81% (69-88%)	0.25 (0.13-0.44)	-0.14% (-0.91-0.55%)	-0.14% (-1-0.65%)	-0.34 (-2.6-1.6)	-7.7 (-50-29)
Guyana	2010	Female	75 to 79	17% (9.8-27%)	83% (73-90%)	0.22 (0.11-0.39)	-0.25% (-1.1-0.37%)	-0.28% (-1.2-0.42%)	-0.63 (-2.7-0.92)	-8.9 (-38-14)
Guyana	2010	Female	80 to 84	15% (8.5-25%)	85% (75-91%)	0.2 (0.1-0.34)	-0.17% (-0.87-0.32%)	-0.21% (-1-0.36%)	-0.43 (-2.1-0.75)	-4.2 (-22-8)
Guyana	2010	Female	85 to 89	15% (8.5-24%)	85% (76-92%)	0.2 (0.1-0.33)	-0.12% (-0.68-0.34%)	-0.17% (-0.83-0.38%)	-0.26 (-1.3-0.58)	-1.6 (-8.9-4.5)
Guyana	2010	Female	90 to 94	17% (9.9-26%)	83% (74-90%)	0.22 (0.11-0.38)	-0.066% (-0.61-0.46%)	-0.16% (-0.8-0.48%)	-0.15 (-0.77-0.46)	-0.38 (-3.5-2.5)
Guyana	2010	Female	95 plus	19% (10-33%)	81% (67-90%)	0.26 (0.11-0.49)	-0.096% (-0.77-0.55%)	-0.26% (-1.1-0.56%)	-0.047 (-0.19-0.1)	-0.071 (-0.56-0.41)
Guyana	2010	Male	All Ages	57% (49-65%)	43% (35-51%)	2.9 (2.6-3.3)	12% (9.8-13%)	13% (11-16%)	460 (370-560)	19000 (16000-22000)
Guyana	2010	Male	15 to 19	57% (41-71%)	43% (29-59%)	2.5 (1.8-3.3)	9% (6.3-12%)	12% (8.3-17%)	9.5 (6.3-13)	750 (520-1000)
Guyana	2010	Male	20 to 24	71% (56-82%)	29% (18-44%)	5.3 (4.1-6.6)	19% (15-24%)	24% (18-31%)	21 (16-28)	1500 (1200-1900)
Guyana	2010	Male	25 to 29	74% (61-84%)	26% (16-39%)	5.9 (4.6-7.2)	20% (16-24%)	24% (19-30%)	25 (19-32)	1700 (1300-2100)
Guyana	2010	Male	30 to 34	74% (60-85%)	26% (15-40%)	5.8 (4.5-7.2)	20% (16-24%)	24% (19-28%)	32 (25-39)	1900 (1500-2300)
Guyana	2010	Male	35 to 39	74% (61-85%)	26% (15-39%)	5.3 (4.1-6.6)	19% (15-22%)	23% (19-27%)	39 (31-47)	2100 (1700-2500)
Guyana	2010	Male	40 to 44	73% (60-84%)	27% (16-40%)	5.2 (4.1-6.3)	17% (15-20%)	22% (18-25%)	47 (38-57)	2300 (1900-2800)
Guyana	2010	Male	45 to 49	72% (57-84%)	28% (16-43%)	4.9 (3.7-6.1)	17% (14-20%)	21% (17-25%)	56 (45-69)	2400 (2000-3000)
Guyana	2010	Male	50 to 54	68% (51-82%)	32% (18-49%)	4.1 (2.8-5.4)	15% (12-18%)	18% (14-22%)	53 (41-66)	2000 (1600-2500)
Guyana	2010	Male	55 to 59	65% (47-81%)	35% (19-53%)	3.5 (2.1-5.2)	13% (8.9-18%)	16% (10-21%)	48 (31-66)	1600 (1000-2200)
Guyana	2010	Male	60 to 64	63% (43-80%)	37% (20-57%)	3.1 (1.7-4.9)	11% (6.5-16%)	13% (7.6-19%)	41 (24-59)	1100 (670-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	2010	Male	65 to 69	61% (42-78%)	39% (22-58%)	2.7 (1.6-4.1)	10% (6.2-15%)	12% (6.8-17%)	30 (18-44)	680 (410-1000)
Guyana	2010	Male	70 to 74	58% (40-74%)	42% (26-60%)	2.3 (1.3-3.6)	8% (4.1-13%)	9% (4.6-14%)	28 (14-45)	520 (270-830)
Guyana	2010	Male	75 to 79	53% (33-72%)	47% (28-67%)	1.9 (1-3.2)	6.4% (2.8-11%)	7.1% (3-12%)	16 (6.8-28)	230 (100-390)
Guyana	2010	Male	80 to 84	47% (29-65%)	53% (35-71%)	1.6 (0.88-2.6)	5% (2.5-8.3%)	5.6% (2.7-9.4%)	10 (4.9-17)	110 (54-180)
Guyana	2010	Male	85 to 89	43% (26-60%)	57% (40-74%)	1.4 (0.77-2.3)	3.9% (1.9-6.6%)	4.3% (2.1-7.5%)	4.8 (2.3-8.2)	37 (19-63)
Guyana	2010	Male	90 to 94	43% (27-60%)	57% (40-73%)	1.4 (0.74-2.3)	3.2% (1.5-6%)	3.6% (1.5-6.8%)	1.8 (0.73-3.4)	9.8 (4.5-18)
Guyana	2010	Male	95 plus	45% (26-66%)	55% (34-74%)	1.5 (0.64-2.8)	3.5% (1.2-7%)	3.8% (0.99-8.1%)	0.3 (0.079-0.64)	1.1 (0.37-2.4)
Guyana	2016	Female	All Ages	28% (23-35%)	72% (65-77%)	0.35 (0.26-0.46)	0.95% (0.51-1.4%)	0.81% (0.18-1.5%)	20 (4.3-37)	1100 (590-1700)
Guyana	2016	Female	15 to 19	34% (22-51%)	66% (49-78%)	0.38 (0.22-0.61)	1.3% (0.68-2%)	2% (0.6-3.8%)	0.78 (0.24-1.5)	89 (45-140)
Guyana	2016	Female	20 to 24	40% (25-56%)	60% (44-75%)	0.57 (0.32-0.88)	2.2% (1.4-3.2%)	2.6% (1.2-4.4%)	1.5 (0.7-2.5)	160 (100-240)
Guyana	2016	Female	25 to 29	39% (23-56%)	61% (44-77%)	0.61 (0.35-0.99)	2.1% (1.5-3%)	2.5% (1.4-3.9%)	1 (0.56-1.7)	100 (67-140)
Guyana	2016	Female	30 to 34	37% (22-54%)	63% (46-78%)	0.6 (0.33-0.99)	2.1% (1.5-3%)	2.7% (1.8-4%)	1.2 (0.74-1.8)	96 (66-130)
Guyana	2016	Female	35 to 39	36% (23-54%)	64% (46-77%)	0.58 (0.31-0.94)	2.2% (1.6-3%)	3.1% (2.1-4.3%)	1.9 (1.2-2.8)	130 (88-170)
Guyana	2016	Female	40 to 44	35% (22-53%)	65% (47-78%)	0.55 (0.31-0.87)	1.9% (1.2-2.7%)	2.9% (1.9-4.1%)	2.5 (1.6-3.7)	140 (90-190)
Guyana	2016	Female	45 to 49	33% (19-51%)	67% (49-81%)	0.51 (0.28-0.84)	1.5% (0.69-2.4%)	2.2% (1-3.5%)	3.1 (1.4-5.1)	140 (63-220)
Guyana	2016	Female	50 to 54	30% (18-46%)	70% (54-82%)	0.46 (0.25-0.78)	1.2% (0.31-2.2%)	1.8% (0.61-3.2%)	3.4 (1.1-6)	130 (34-230)
Guyana	2016	Female	55 to 59	26% (15-42%)	74% (58-85%)	0.42 (0.21-0.71)	0.93% (0.033-2%)	1.3% (0.16-2.7%)	3.3 (0.4-6.9)	110 (3.9-230)
Guyana	2016	Female	60 to 64	23% (13-38%)	77% (62-87%)	0.37 (0.18-0.63)	0.45% (-0.47-1.4%)	0.64% (-0.54-1.8%)	1.6 (-1.4-4.7)	41 (-44-130)
Guyana	2016	Female	65 to 69	21% (13-33%)	79% (67-87%)	0.32 (0.16-0.53)	0.076% (-0.76-0.86%)	0.15% (-0.87-1.1%)	0.36 (-2.2-2.7)	5.5 (-55-60)
Guyana	2016	Female	70 to 74	19% (11-30%)	81% (70-89%)	0.26 (0.13-0.46)	-0.049% (-0.8-0.71%)	-0.025% (-0.96-0.9%)	-0.06 (-2.4-2.3)	-2.7 (-46-40)
Guyana	2016	Female	75 to 79	17% (9.7-28%)	83% (72-90%)	0.23 (0.11-0.41)	-0.12% (-0.87-0.5%)	-0.14% (-1-0.58%)	-0.38 (-2.6-1.6)	-5.5 (-38-22)
Guyana	2016	Female	80 to 84	15% (8.8-25%)	85% (75-91%)	0.21 (0.1-0.36)	-0.072% (-0.72-0.46%)	-0.1% (-0.85-0.53%)	-0.17 (-1.5-0.88)	-1.5 (-15-9.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guyana	2016	Female	85 to 89	15% (8.8-24%)	85% (76-91%)	0.21 (0.1-0.36)	-0.044% (-0.56-0.45%)	-0.095% (-0.72-0.5%)	-0.088 (-0.68-0.46)	-0.36 (-4.6-3.7)
Guyana	2016	Female	90 to 94	17% (9.8-26%)	83% (74-90%)	0.23 (0.11-0.41)	-0.046% (-0.63-0.53%)	-0.13% (-0.84-0.57%)	-0.08 (-0.51-0.34)	-0.17 (-2.3-2)
Guyana	2016	Female	95 plus	20% (10-33%)	80% (67-90%)	0.27 (0.11-0.56)	-0.084% (-0.8-0.66%)	-0.24% (-1.1-0.62%)	-0.061 (-0.29-0.16)	-0.087 (-0.82-0.69)
Guyana	2016	Male	All Ages	57% (48-66%)	43% (34-52%)	3.1 (2.4-3.9)	12% (9.1-14%)	13% (9.9-17%)	460 (340-590)	19000 (14000-23000)
Guyana	2016	Male	15 to 19	57% (40-74%)	43% (26-60%)	2.6 (1.7-3.7)	9% (5.9-13%)	13% (8-19%)	10 (6.3-15)	810 (540-1100)
Guyana	2016	Male	20 to 24	67% (50-82%)	33% (18-50%)	5.1 (3.4-7.1)	18% (14-24%)	23% (17-31%)	27 (19-36)	1900 (1400-2500)
Guyana	2016	Male	25 to 29	71% (53-83%)	29% (17-47%)	5.7 (3.8-7.8)	19% (14-24%)	24% (17-30%)	24 (17-31)	1600 (1200-2000)
Guyana	2016	Male	30 to 34	71% (53-85%)	29% (15-47%)	5.7 (3.7-8.1)	18% (14-23%)	22% (17-28%)	25 (18-32)	1500 (1100-1900)
Guyana	2016	Male	35 to 39	71% (55-84%)	29% (16-45%)	5.3 (3.4-7.5)	17% (13-22%)	22% (16-27%)	25 (18-32)	1400 (1000-1800)
Guyana	2016	Male	40 to 44	70% (54-84%)	30% (16-46%)	5 (3.4-7)	17% (13-21%)	21% (16-26%)	32 (23-41)	1600 (1200-2000)
Guyana	2016	Male	45 to 49	68% (49-83%)	32% (17-51%)	4.7 (3.1-6.6)	16% (13-20%)	20% (16-25%)	50 (36-64)	2200 (1600-2800)
Guyana	2016	Male	50 to 54	67% (48-82%)	33% (18-52%)	4.2 (2.7-6.1)	15% (11-19%)	18% (13-23%)	60 (43-80)	2300 (1700-3000)
Guyana	2016	Male	55 to 59	65% (47-80%)	35% (20-53%)	3.6 (2.1-5.6)	13% (8.8-18%)	16% (10-21%)	63 (40-89)	2100 (1300-2900)
Guyana	2016	Male	60 to 64	63% (43-80%)	37% (20-57%)	3.2 (1.7-5.1)	11% (6.3-17%)	13% (7.3-19%)	48 (27-72)	1300 (760-2000)
Guyana	2016	Male	65 to 69	62% (42-78%)	38% (22-58%)	2.8 (1.6-4.5)	10% (5.8-15%)	12% (6.7-18%)	40 (22-61)	930 (520-1400)
Guyana	2016	Male	70 to 74	58% (40-74%)	42% (26-60%)	2.4 (1.3-4)	8.1% (4-13%)	9.2% (4.4-15%)	25 (12-39)	460 (230-730)
Guyana	2016	Male	75 to 79	53% (34-72%)	47% (28-66%)	2 (1-3.5)	6.4% (2.8-11%)	7.2% (3-12%)	21 (8.8-36)	300 (130-520)
Guyana	2016	Male	80 to 84	47% (29-66%)	53% (34-71%)	1.7 (0.88-2.8)	5.1% (2.2-9%)	5.7% (2.5-10%)	10 (4.5-18)	110 (49-190)
Guyana	2016	Male	85 to 89	42% (26-59%)	58% (41-74%)	1.4 (0.74-2.4)	4% (1.8-7.2%)	4.5% (1.9-8.3%)	3.2 (1.4-5.8)	25 (11-45)
Guyana	2016	Male	90 to 94	42% (28-60%)	58% (40-72%)	1.4 (0.72-2.4)	3.4% (1.4-6.6%)	3.7% (1.3-7.5%)	1.6 (0.58-3.1)	8.7 (3.5-17)
Guyana	2016	Male	95 plus	45% (26-65%)	55% (35-74%)	1.5 (0.64-2.9)	3.6% (1-7.8%)	3.9% (0.9-8.9%)	0.47 (0.11-1.1)	1.8 (0.52-3.9)
Haiti	1990	Female	All Ages	25% (21-30%)	75% (70-79%)	0.74 (0.63-0.85)	1.4% (0.96-2%)	2.1% (1.2-3.2%)	1000 (590-1600)	42000 (29000-58000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	1990	Female	15 to 19	33% (22-46%)	67% (54-78%)	1 (0.56-1.6)	2.8% (1.4-4.5%)	4% (1.9-6.8%)	28 (13-47)	2300 (1200-3700)
Haiti	1990	Female	20 to 24	38% (26-52%)	62% (48-74%)	1.5 (0.89-2.3)	4.5% (2.6-6.8%)	5.7% (3-9.2%)	57 (30-93)	4400 (2600-6800)
Haiti	1990	Female	25 to 29	37% (24-50%)	63% (50-76%)	1.5 (0.92-2.3)	4.7% (2.9-7.1%)	5.8% (3.3-9.1%)	67 (38-110)	4700 (2900-7100)
Haiti	1990	Female	30 to 34	36% (22-49%)	64% (51-78%)	1.5 (0.87-2.3)	4.5% (2.7-6.9%)	5.6% (3.1-8.9%)	66 (37-110)	4200 (2500-6500)
Haiti	1990	Female	35 to 39	37% (24-51%)	63% (49-76%)	1.4 (0.82-2.3)	4.8% (2.8-7.4%)	6.1% (3.5-9.5%)	82 (45-130)	4400 (2600-6900)
Haiti	1990	Female	40 to 44	36% (24-50%)	64% (50-76%)	1.4 (0.74-2.2)	4.9% (2.6-8.1%)	6.3% (3.3-10%)	82 (41-140)	4000 (2100-6500)
Haiti	1990	Female	45 to 49	35% (21-49%)	65% (51-79%)	1.3 (0.69-2.2)	4.7% (2.3-7.8%)	5.9% (2.9-9.9%)	95 (44-170)	4000 (1900-6900)
Haiti	1990	Female	50 to 54	31% (19-45%)	69% (55-81%)	1.2 (0.62-2)	4.6% (2-8.2%)	5.6% (2.4-10%)	120 (50-210)	4300 (1900-7900)
Haiti	1990	Female	55 to 59	26% (16-40%)	74% (60-84%)	0.97 (0.5-1.7)	3.8% (1.4-7.1%)	4.5% (1.7-8.5%)	110 (40-200)	3500 (1300-6400)
Haiti	1990	Female	60 to 64	23% (14-36%)	77% (64-86%)	0.82 (0.4-1.5)	2.8% (0.72-5.8%)	3.3% (0.8-6.8%)	91 (22-180)	2500 (610-5000)
Haiti	1990	Female	65 to 69	21% (13-32%)	79% (68-87%)	0.71 (0.37-1.2)	2.1% (0.38-4.4%)	2.4% (0.44-5%)	80 (14-170)	1800 (320-3800)
Haiti	1990	Female	70 to 74	19% (12-29%)	81% (71-88%)	0.59 (0.29-1.1)	1.6% (-0.019-4%)	1.8% (-0.043-4.4%)	66 (-1.4-160)	1200 (-14-3000)
Haiti	1990	Female	75 to 79	17% (9.9-26%)	83% (74-90%)	0.5 (0.23-0.92)	1.2% (-0.19-3.2%)	1.3% (-0.23-3.6%)	45 (-7.6-120)	630 (-100-1700)
Haiti	1990	Female	80 to 84	15% (9-23%)	85% (77-91%)	0.47 (0.24-0.79)	1.1% (-0.0089-2.7%)	1.3% (-0.032-3.1%)	32 (-0.88-78)	340 (-2.6-800)
Haiti	1990	Female	85 to 89	15% (8.9-23%)	85% (77-91%)	0.48 (0.23-0.87)	1.1% (0.0096-2.8%)	1.2% (-0.012-3.1%)	14 (-0.14-37)	100 (0.9-270)
Haiti	1990	Female	90 to 94	17% (10-26%)	83% (74-90%)	0.54 (0.24-1.1)	1.2% (-0.0014-3.2%)	1.3% (-0.069-3.6%)	4.3 (-0.23-12)	23 (-0.027-61)
Haiti	1990	Female	95 plus	20% (10-33%)	80% (67-90%)	0.64 (0.23-1.3)	1.3% (-0.2-4%)	1.4% (-0.3-4.6%)	1.1 (-0.23-3.4)	3.8 (-0.58-12)
Haiti	1990	Male	All Ages	44% (37-51%)	56% (49-63%)	1.7 (1.5-1.9)	3.3% (2.6-4.2%)	5.1% (3.7-6.5%)	2400 (1700-3100)	100000 (80000-130000)
Haiti	1990	Male	15 to 19	55% (41-68%)	45% (32-59%)	1.7 (1-2.7)	6.1% (3.8-9%)	7.8% (4.7-12%)	60 (34-97)	4900 (3000-7400)
Haiti	1990	Male	20 to 24	66% (52-77%)	34% (23-48%)	3.3 (2.2-4.5)	12% (8.1-16%)	14% (9-19%)	160 (100-240)	12000 (8100-17000)
Haiti	1990	Male	25 to 29	66% (52-77%)	34% (23-48%)	3.9 (2.6-5.3)	13% (9-17%)	15% (10-21%)	190 (130-270)	13000 (9100-18000)
Haiti	1990	Male	30 to 34	65% (51-76%)	35% (24-49%)	4 (2.6-5.6)	13% (9.6-18%)	16% (11-21%)	210 (140-290)	13000 (8700-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	1990	Male	35 to 39	63% (49-75%)	37% (25-51%)	3.7 (2.3-5.5)	13% (8.7-17%)	15% (10-21%)	210 (140-290)	11000 (7800-16000)
Haiti	1990	Male	40 to 44	61% (49-73%)	39% (27-51%)	3.5 (2.1-5.3)	12% (8.1-16%)	15% (9.4-20%)	180 (120-260)	9100 (5900-13000)
Haiti	1990	Male	45 to 49	60% (44-72%)	40% (28-56%)	3.3 (1.9-5.1)	12% (7.4-16%)	14% (8.5-20%)	210 (130-300)	9200 (5700-13000)
Haiti	1990	Male	50 to 54	57% (43-70%)	43% (30-57%)	3 (1.7-4.7)	11% (6.3-15%)	13% (7.3-18%)	210 (120-310)	8100 (4600-12000)
Haiti	1990	Male	55 to 59	54% (39-69%)	46% (31-61%)	2.7 (1.5-4.3)	9.7% (5.5-14%)	12% (6.4-17%)	210 (110-310)	6800 (3900-10000)
Haiti	1990	Male	60 to 64	52% (36-69%)	48% (31-64%)	2.4 (1.3-4.2)	8.3% (4.1-13%)	9.6% (4.7-15%)	190 (93-310)	5400 (2700-8600)
Haiti	1990	Male	65 to 69	51% (34-69%)	49% (31-66%)	2.2 (1.2-3.4)	7.1% (3.5-11%)	8.2% (3.9-13%)	180 (88-300)	4300 (2100-6900)
Haiti	1990	Male	70 to 74	47% (31-65%)	53% (35-69%)	1.8 (0.9-3)	5.4% (2.1-9.4%)	6.1% (2.4-11%)	150 (58-280)	2900 (1100-5000)
Haiti	1990	Male	75 to 79	42% (27-60%)	58% (40-73%)	1.5 (0.7-2.5)	4.2% (1.4-7.6%)	4.7% (1.5-8.5%)	110 (38-210)	1600 (550-2900)
Haiti	1990	Male	80 to 84	37% (23-54%)	63% (46-77%)	1.2 (0.63-2.1)	3.4% (1.2-6.3%)	3.7% (1.2-7.1%)	60 (20-110)	640 (230-1200)
Haiti	1990	Male	85 to 89	34% (21-48%)	66% (52-79%)	1.1 (0.56-1.8)	2.7% (0.99-5.3%)	3% (1-6%)	27 (9.1-55)	210 (74-410)
Haiti	1990	Male	90 to 94	35% (22-49%)	65% (51-78%)	1.1 (0.53-2)	2.3% (0.64-5%)	2.5% (0.57-5.7%)	8.3 (1.9-19)	45 (13-97)
Haiti	1990	Male	95 plus	38% (23-57%)	62% (43-77%)	1.2 (0.5-2.2)	2.4% (0.36-5.7%)	2.5% (0.17-6.5%)	1.9 (0.13-4.7)	7 (1-17)
Haiti	1995	Female	All Ages	29% (24-34%)	71% (66-76%)	0.74 (0.64-0.84)	1.3% (0.89-1.8%)	1.7% (0.96-2.7%)	870 (480-1300)	37000 (25000-51000)
Haiti	1995	Female	15 to 19	37% (26-50%)	63% (50-74%)	0.99 (0.56-1.5)	2.5% (1.3-4%)	3.7% (1.7-6.1%)	28 (13-47)	2300 (1200-3600)
Haiti	1995	Female	20 to 24	44% (32-57%)	56% (43-68%)	1.6 (0.95-2.3)	3.6% (2.2-5.6%)	4.5% (2.4-7.3%)	51 (27-81)	4000 (2400-6000)
Haiti	1995	Female	25 to 29	41% (29-54%)	59% (46-71%)	1.5 (0.97-2.2)	3.4% (2.1-5.3%)	3.9% (2.2-6.5%)	59 (34-95)	4200 (2600-6400)
Haiti	1995	Female	30 to 34	39% (26-53%)	61% (47-74%)	1.5 (0.84-2.2)	3.2% (1.9-4.9%)	3.7% (2.1-6.1%)	62 (35-98)	3900 (2400-6000)
Haiti	1995	Female	35 to 39	42% (29-55%)	58% (45-71%)	1.5 (0.88-2.3)	3.7% (2.1-5.7%)	4.5% (2.5-7.2%)	74 (42-120)	4100 (2400-6400)
Haiti	1995	Female	40 to 44	44% (32-57%)	56% (43-68%)	1.5 (0.85-2.4)	4% (2.1-6.5%)	5% (2.6-8.3%)	85 (43-140)	4100 (2100-6800)
Haiti	1995	Female	45 to 49	41% (27-54%)	59% (46-73%)	1.4 (0.76-2.3)	3.9% (1.8-6.7%)	4.8% (2.2-8.4%)	81 (36-140)	3400 (1500-5900)
Haiti	1995	Female	50 to 54	33% (20-47%)	67% (53-80%)	1.1 (0.59-1.8)	3.7% (1.4-6.7%)	4.5% (1.7-8.2%)	96 (37-180)	3500 (1400-6700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	1995	Female	55 to 59	26% (17-40%)	74% (60-83%)	0.85 (0.43-1.5)	2.9% (1-5.4%)	3.5% (1.2-6.5%)	90 (30-170)	2900 (960-5300)
Haiti	1995	Female	60 to 64	23% (14-36%)	77% (64-86%)	0.71 (0.35-1.3)	2.1% (0.31-4.4%)	2.5% (0.38-5.2%)	73 (11-160)	2000 (290-4200)
Haiti	1995	Female	65 to 69	21% (13-33%)	79% (67-87%)	0.62 (0.33-1)	1.5% (0.056-3.4%)	1.7% (0.067-3.8%)	59 (2.3-130)	1300 (48-2900)
Haiti	1995	Female	70 to 74	19% (12-29%)	81% (71-88%)	0.51 (0.25-0.9)	1.1% (-0.25-2.9%)	1.2% (-0.25-3.3%)	45 (-8.7-120)	810 (-180-2100)
Haiti	1995	Female	75 to 79	17% (9.9-26%)	83% (74-90%)	0.44 (0.2-0.81)	0.79% (-0.41-2.4%)	0.88% (-0.47-2.7%)	31 (-17-97)	440 (-220-1400)
Haiti	1995	Female	80 to 84	15% (9-24%)	85% (76-91%)	0.41 (0.2-0.7)	0.76% (-0.2-2.1%)	0.85% (-0.26-2.3%)	24 (-7.2-66)	250 (-64-690)
Haiti	1995	Female	85 to 89	15% (9-23%)	85% (77-91%)	0.42 (0.2-0.76)	0.74% (-0.21-2.2%)	0.81% (-0.28-2.5%)	11 (-3.7-33)	82 (-24-240)
Haiti	1995	Female	90 to 94	17% (10-26%)	83% (74-90%)	0.48 (0.22-0.93)	0.81% (-0.22-2.6%)	0.88% (-0.33-2.9%)	3.6 (-1.3-12)	19 (-5.4-60)
Haiti	1995	Female	95 plus	20% (11-33%)	80% (67-89%)	0.57 (0.21-1.2)	0.93% (-0.45-3.2%)	0.97% (-0.61-3.5%)	0.91 (-0.57-3.3)	3.4 (-1.6-11)
Haiti	1995	Male	All Ages	48% (40-55%)	52% (45-60%)	1.6 (1.4-1.8)	3.2% (2.5-4%)	4.6% (3.4-5.9%)	2200 (1600-2800)	97000 (77000-120000)
Haiti	1995	Male	15 to 19	60% (47-72%)	40% (28-53%)	1.7 (1-2.5)	5.9% (3.8-8.5%)	7.8% (4.8-12%)	65 (37-100)	5300 (3400-7900)
Haiti	1995	Male	20 to 24	70% (59-79%)	30% (21-41%)	3.2 (2.2-4.2)	11% (7.4-15%)	12% (7.9-17%)	150 (94-220)	11000 (7700-16000)
Haiti	1995	Male	25 to 29	69% (57-78%)	31% (22-43%)	3.6 (2.4-4.9)	10% (6.7-14%)	11% (7.1-16%)	180 (120-250)	12000 (8500-17000)
Haiti	1995	Male	30 to 34	69% (57-78%)	31% (22-43%)	3.7 (2.5-5.1)	9.8% (6.8-13%)	11% (7.2-15%)	200 (130-280)	12000 (8500-17000)
Haiti	1995	Male	35 to 39	67% (56-77%)	33% (23-44%)	3.5 (2.3-5.1)	9.7% (6.5-14%)	11% (7-16%)	200 (130-290)	11000 (7400-16000)
Haiti	1995	Male	40 to 44	63% (51-74%)	37% (26-49%)	3.2 (1.9-4.8)	9.2% (5.9-13%)	11% (6.7-15%)	190 (120-280)	9600 (6200-14000)
Haiti	1995	Male	45 to 49	62% (46-74%)	38% (26-54%)	3 (1.7-4.8)	9.2% (5.6-13%)	11% (6.2-16%)	170 (100-260)	7600 (4600-11000)
Haiti	1995	Male	50 to 54	61% (48-73%)	39% (27-52%)	2.8 (1.6-4.4)	9% (5.2-13%)	11% (5.9-16%)	200 (110-290)	7600 (4300-11000)
Haiti	1995	Male	55 to 59	56% (41-70%)	44% (30-59%)	2.5 (1.4-3.8)	8% (4.4-12%)	9.4% (5.1-14%)	190 (100-300)	6300 (3500-9700)
Haiti	1995	Male	60 to 64	52% (35-69%)	48% (31-65%)	2.1 (1.1-3.6)	6.6% (3.2-11%)	7.7% (3.6-13%)	170 (81-290)	4800 (2400-8100)
Haiti	1995	Male	65 to 69	51% (34-68%)	49% (32-66%)	1.9 (1-2.9)	5.6% (2.7-9.3%)	6.5% (3-11%)	150 (72-260)	3600 (1700-5900)
Haiti	1995	Male	70 to 74	47% (30-65%)	53% (35-70%)	1.6 (0.81-2.6)	4.3% (1.6-7.9%)	4.8% (1.7-9%)	130 (44-240)	2400 (860-4300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	1995	Male	75 to 79	42% (27-60%)	58% (40-73%)	1.3 (0.63-2.2)	3.3% (1.1-6.3%)	3.7% (1.1-7%)	91 (28-180)	1300 (440-2500)
Haiti	1995	Male	80 to 84	37% (23-54%)	63% (46-77%)	1.1 (0.56-1.8)	2.6% (0.89-5%)	2.8% (0.92-5.5%)	50 (16-97)	540 (180-1000)
Haiti	1995	Male	85 to 89	34% (21-48%)	66% (52-79%)	0.95 (0.49-1.6)	2% (0.66-4.2%)	2.2% (0.63-4.7%)	23 (6.6-49)	180 (60-370)
Haiti	1995	Male	90 to 94	35% (22-50%)	65% (50-78%)	0.96 (0.46-1.7)	1.7% (0.41-3.8%)	1.8% (0.28-4.2%)	7.1 (1.1-17)	40 (9.5-88)
Haiti	1995	Male	95 plus	38% (22-57%)	62% (43-78%)	1.1 (0.42-1.9)	1.7% (0.15-4.4%)	1.8% (-0.086-4.9%)	1.7 (-0.082-4.8)	6.6 (0.54-17)
Haiti	2000	Female	All Ages	24% (20-29%)	76% (71-80%)	0.67 (0.58-0.77)	1.2% (0.83-1.7%)	1.7% (0.91-2.6%)	850 (460-1300)	35000 (24000-49000)
Haiti	2000	Female	15 to 19	31% (22-43%)	69% (57-78%)	0.89 (0.51-1.4)	2.2% (1.2-3.4%)	3.3% (1.7-5.4%)	26 (13-44)	2200 (1200-3600)
Haiti	2000	Female	20 to 24	36% (26-48%)	64% (52-74%)	1.4 (0.83-2)	2.9% (1.8-4.5%)	3.5% (1.8-5.8%)	43 (23-70)	3600 (2200-5300)
Haiti	2000	Female	25 to 29	34% (24-46%)	66% (54-76%)	1.3 (0.81-2)	2.3% (1.3-3.5%)	2.4% (1.3-4.1%)	44 (24-71)	3300 (2100-4900)
Haiti	2000	Female	30 to 34	31% (20-44%)	69% (56-80%)	1.2 (0.7-1.9)	2% (1.2-3.2%)	2.1% (1.1-3.6%)	46 (26-75)	3100 (1900-4800)
Haiti	2000	Female	35 to 39	33% (22-45%)	67% (55-78%)	1.2 (0.69-1.9)	2.3% (1.4-3.7%)	2.7% (1.5-4.4%)	60 (34-98)	3400 (2000-5300)
Haiti	2000	Female	40 to 44	35% (25-47%)	65% (53-75%)	1.2 (0.71-2)	2.8% (1.6-4.5%)	3.3% (1.8-5.4%)	70 (37-110)	3500 (1900-5500)
Haiti	2000	Female	45 to 49	35% (22-47%)	65% (53-78%)	1.3 (0.7-2)	3.1% (1.6-5.3%)	3.8% (1.9-6.6%)	81 (41-140)	3500 (1800-6000)
Haiti	2000	Female	50 to 54	31% (20-45%)	69% (55-80%)	1.1 (0.59-1.9)	3.4% (1.5-6.2%)	4.2% (1.8-7.5%)	90 (39-170)	3300 (1500-6200)
Haiti	2000	Female	55 to 59	26% (16-39%)	74% (61-84%)	0.91 (0.46-1.5)	3% (1.2-5.4%)	3.6% (1.4-6.5%)	94 (35-170)	3000 (1100-5600)
Haiti	2000	Female	60 to 64	23% (14-36%)	77% (64-86%)	0.76 (0.37-1.4)	2.3% (0.54-4.7%)	2.7% (0.66-5.6%)	83 (20-170)	2300 (540-4600)
Haiti	2000	Female	65 to 69	21% (13-32%)	79% (68-87%)	0.66 (0.35-1.1)	1.7% (0.2-3.6%)	2% (0.26-4.2%)	71 (9.1-150)	1600 (180-3400)
Haiti	2000	Female	70 to 74	19% (12-29%)	81% (71-88%)	0.55 (0.28-0.94)	1.3% (-0.067-3.2%)	1.5% (-0.078-3.7%)	54 (-3-140)	980 (-51-2400)
Haiti	2000	Female	75 to 79	16% (9.9-26%)	84% (74-90%)	0.47 (0.22-0.85)	0.97% (-0.25-2.6%)	1.1% (-0.29-3%)	38 (-10-100)	540 (-130-1500)
Haiti	2000	Female	80 to 84	15% (9-23%)	85% (77-91%)	0.44 (0.22-0.75)	0.92% (-0.048-2.3%)	1% (-0.084-2.5%)	31 (-2.6-76)	320 (-17-790)
Haiti	2000	Female	85 to 89	15% (8.9-23%)	85% (77-91%)	0.45 (0.22-0.82)	0.88% (-0.12-2.4%)	0.98% (-0.18-2.6%)	15 (-2.8-42)	120 (-15-310)
Haiti	2000	Female	90 to 94	17% (10-26%)	83% (74-90%)	0.51 (0.23-0.98)	0.95% (-0.098-2.8%)	1% (-0.18-3.2%)	5.2 (-0.91-16)	28 (-2.8-79)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	2000	Female	95 plus	20% (10-33%)	80% (67-90%)	0.61 (0.22-1.2)	1.1% (-0.29-3.5%)	1.2% (-0.45-3.9%)	1.3 (-0.51-4.5)	4.8 (-1.3-15)
Haiti	2000	Male	All Ages	46% (40-53%)	54% (47-60%)	1.7 (1.5-1.9)	3.5% (2.7-4.4%)	4.8% (3.5-6.2%)	2300 (1600-2900)	100000 (78000-130000)
Haiti	2000	Male	15 to 19	58% (46-70%)	42% (30-54%)	1.8 (1.1-2.6)	5.8% (3.8-8.3%)	7.9% (4.8-11%)	71 (42-110)	5900 (3900-8700)
Haiti	2000	Male	20 to 24	69% (57-78%)	31% (22-43%)	3.3 (2.3-4.5)	11% (7.5-15%)	13% (8.3-18%)	160 (100-240)	12000 (8400-17000)
Haiti	2000	Male	25 to 29	68% (56-77%)	32% (23-44%)	3.8 (2.6-5.2)	9.2% (6.1-13%)	10% (6.2-15%)	170 (110-240)	12000 (8000-16000)
Haiti	2000	Male	30 to 34	67% (56-76%)	33% (24-44%)	3.9 (2.6-5.4)	8.2% (5.7-11%)	9% (5.8-13%)	190 (120-270)	12000 (8100-16000)
Haiti	2000	Male	35 to 39	65% (54-75%)	35% (25-46%)	3.6 (2.3-5.3)	7.9% (5.1-11%)	8.7% (5.3-13%)	200 (130-280)	11000 (7400-16000)
Haiti	2000	Male	40 to 44	61% (50-72%)	39% (28-50%)	3.3 (2-4.9)	7.7% (5-11%)	8.6% (5.4-12%)	190 (120-280)	9700 (6300-14000)
Haiti	2000	Male	45 to 49	60% (47-72%)	40% (28-53%)	3.1 (1.8-4.8)	8% (5-11%)	9.1% (5.6-13%)	200 (120-290)	8800 (5500-12000)
Haiti	2000	Male	50 to 54	59% (46-70%)	41% (30-54%)	2.9 (1.7-4.5)	8.3% (4.9-12%)	9.7% (5.5-14%)	180 (99-260)	6900 (4000-10000)
Haiti	2000	Male	55 to 59	54% (40-68%)	46% (32-60%)	2.5 (1.4-3.9)	7.9% (4.4-12%)	9.3% (5.1-14%)	200 (100-310)	6600 (3600-10000)
Haiti	2000	Male	60 to 64	51% (35-68%)	49% (32-65%)	2.2 (1.2-3.9)	6.8% (3.4-11%)	7.9% (3.9-13%)	190 (91-320)	5300 (2700-8800)
Haiti	2000	Male	65 to 69	51% (33-68%)	49% (32-67%)	2 (1.1-3.1)	6% (3-9.5%)	6.9% (3.3-11%)	180 (84-290)	4100 (2000-6600)
Haiti	2000	Male	70 to 74	47% (30-64%)	53% (36-70%)	1.7 (0.86-2.8)	4.6% (1.7-8.2%)	5.2% (1.9-9.4%)	140 (50-250)	2600 (970-4600)
Haiti	2000	Male	75 to 79	42% (27-60%)	58% (40-73%)	1.4 (0.68-2.4)	3.5% (1.3-6.6%)	4% (1.4-7.4%)	99 (35-190)	1400 (520-2600)
Haiti	2000	Male	80 to 84	37% (22-53%)	63% (47-78%)	1.1 (0.6-1.9)	2.8% (1-5.3%)	3.1% (1.1-6.1%)	58 (20-110)	630 (230-1200)
Haiti	2000	Male	85 to 89	33% (21-47%)	67% (53-79%)	1 (0.52-1.7)	2.2% (0.8-4.5%)	2.5% (0.78-5.1%)	28 (8.9-56)	210 (76-430)
Haiti	2000	Male	90 to 94	34% (22-49%)	66% (51-78%)	1 (0.5-1.8)	1.9% (0.57-4.1%)	2.1% (0.46-4.6%)	9.2 (2-21)	51 (15-110)
Haiti	2000	Male	95 plus	38% (23-57%)	62% (43-77%)	1.1 (0.46-2.1)	2% (0.33-4.7%)	2.1% (0.11-5.2%)	2.3 (0.12-5.8)	9 (1.5-21)
Haiti	2005	Female	All Ages	24% (20-29%)	76% (71-80%)	0.7 (0.6-0.79)	1.2% (0.83-1.7%)	1.6% (0.88-2.5%)	880 (460-1400)	36000 (24000-51000)
Haiti	2005	Female	15 to 19	31% (22-43%)	69% (57-78%)	0.9 (0.5-1.4)	2% (1.1-3.2%)	3% (1.5-5%)	24 (12-40)	2100 (1200-3300)
Haiti	2005	Female	20 to 24	36% (25-48%)	64% (52-75%)	1.4 (0.84-2)	2.9% (1.7-4.4%)	3.6% (1.8-5.8%)	45 (24-73)	3800 (2300-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	2005	Female	25 to 29	34% (24-46%)	66% (54-76%)	1.3 (0.83-2)	2.2% (1.3-3.4%)	2.3% (1.2-3.9%)	45 (25-73)	3500 (2200-5200)
Haiti	2005	Female	30 to 34	31% (20-44%)	69% (56-80%)	1.2 (0.69-1.9)	1.7% (1-2.7%)	1.7% (0.92-2.9%)	41 (23-71)	2900 (1700-4600)
Haiti	2005	Female	35 to 39	33% (22-45%)	67% (55-78%)	1.2 (0.71-1.9)	2% (1.2-3.2%)	2.2% (1.2-3.6%)	56 (30-95)	3300 (1900-5300)
Haiti	2005	Female	40 to 44	35% (24-47%)	65% (53-76%)	1.3 (0.7-2)	2.3% (1.3-3.7%)	2.7% (1.4-4.4%)	73 (35-120)	3600 (1900-6000)
Haiti	2005	Female	45 to 49	34% (22-47%)	66% (53-78%)	1.3 (0.7-2.1)	2.8% (1.4-4.7%)	3.3% (1.7-5.8%)	81 (40-150)	3500 (1700-6200)
Haiti	2005	Female	50 to 54	31% (19-44%)	69% (56-81%)	1.1 (0.6-1.9)	3.2% (1.3-5.7%)	3.9% (1.6-6.9%)	100 (41-190)	3700 (1500-7000)
Haiti	2005	Female	55 to 59	26% (16-39%)	74% (61-84%)	0.92 (0.46-1.6)	2.9% (1.2-5.1%)	3.5% (1.4-6.4%)	89 (35-170)	2900 (1100-5300)
Haiti	2005	Female	60 to 64	23% (14-36%)	77% (64-86%)	0.78 (0.38-1.4)	2.2% (0.55-4.6%)	2.7% (0.66-5.5%)	84 (20-180)	2300 (510-4800)
Haiti	2005	Female	65 to 69	21% (13-32%)	79% (68-87%)	0.68 (0.35-1.1)	1.7% (0.22-3.7%)	2% (0.25-4.3%)	77 (9.9-160)	1700 (220-3700)
Haiti	2005	Female	70 to 74	19% (12-30%)	81% (70-88%)	0.56 (0.28-0.98)	1.3% (-0.056-3.2%)	1.5% (-0.063-3.7%)	61 (-2.7-150)	1100 (-48-2800)
Haiti	2005	Female	75 to 79	16% (9.8-26%)	84% (74-90%)	0.48 (0.22-0.89)	0.98% (-0.25-2.7%)	1.1% (-0.29-3.1%)	41 (-11-110)	580 (-150-1500)
Haiti	2005	Female	80 to 84	15% (9.1-24%)	85% (76-91%)	0.45 (0.22-0.75)	0.93% (-0.064-2.2%)	1% (-0.098-2.6%)	33 (-3.4-80)	340 (-26-830)
Haiti	2005	Female	85 to 89	15% (9.1-23%)	85% (77-91%)	0.45 (0.22-0.81)	0.89% (-0.1-2.3%)	0.99% (-0.17-2.6%)	18 (-3.1-47)	140 (-16-350)
Haiti	2005	Female	90 to 94	17% (10-26%)	83% (74-90%)	0.51 (0.24-0.98)	0.96% (-0.08-2.7%)	1.1% (-0.17-3.1%)	6.7 (-1.1-20)	36 (-3.1-100)
Haiti	2005	Female	95 plus	19% (10-32%)	81% (68-90%)	0.6 (0.22-1.2)	1.1% (-0.24-3.5%)	1.2% (-0.4-3.9%)	1.7 (-0.58-5.7)	6.2 (-1.4-20)
Haiti	2005	Male	All Ages	47% (41-54%)	53% (46-59%)	1.8 (1.6-2)	3.5% (2.6-4.5%)	4.7% (3.4-6.2%)	2300 (1600-3100)	100000 (75000-130000)
Haiti	2005	Male	15 to 19	58% (45-71%)	42% (29-55%)	1.8 (1.1-2.6)	5.4% (3.6-7.8%)	7.2% (4.5-11%)	65 (38-110)	5600 (3600-8400)
Haiti	2005	Male	20 to 24	69% (57-78%)	31% (22-43%)	3.3 (2.3-4.4)	11% (7.6-15%)	13% (8.8-18%)	170 (110-260)	14000 (9100-20000)
Haiti	2005	Male	25 to 29	68% (57-77%)	32% (23-43%)	3.8 (2.6-5.1)	9.6% (6.4-14%)	11% (6.5-16%)	180 (110-260)	13000 (8300-18000)
Haiti	2005	Male	30 to 34	67% (55-76%)	33% (24-45%)	3.9 (2.6-5.4)	7.9% (5.3-11%)	8.6% (5.4-13%)	170 (110-250)	11000 (7400-16000)
Haiti	2005	Male	35 to 39	65% (55-75%)	35% (25-45%)	3.6 (2.3-5.3)	7.3% (4.7-10%)	8% (4.8-12%)	190 (110-290)	11000 (6900-16000)
Haiti	2005	Male	40 to 44	61% (50-72%)	39% (28-50%)	3.3 (2-4.9)	6.9% (4.4-9.9%)	7.7% (4.7-11%)	190 (110-300)	9900 (6100-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	2005	Male	45 to 49	60% (47-71%)	40% (29-53%)	3.1 (1.9-4.9)	7.4% (4.7-11%)	8.4% (5-12%)	200 (120-300)	9000 (5500-13000)
Haiti	2005	Male	50 to 54	59% (47-70%)	41% (30-53%)	2.9 (1.7-4.6)	7.9% (4.7-11%)	9.2% (5.4-13%)	210 (110-310)	8100 (4600-12000)
Haiti	2005	Male	55 to 59	54% (40-68%)	46% (32-60%)	2.5 (1.4-4)	7.8% (4.4-12%)	9.2% (5.2-14%)	180 (97-280)	6100 (3400-9400)
Haiti	2005	Male	60 to 64	51% (35-68%)	49% (32-65%)	2.2 (1.2-3.9)	6.7% (3.3-11%)	7.8% (3.7-13%)	190 (90-320)	5500 (2700-9000)
Haiti	2005	Male	65 to 69	51% (33-68%)	49% (32-67%)	2 (1.1-3.2)	5.9% (2.9-9.5%)	6.8% (3.3-11%)	190 (86-310)	4400 (2100-7300)
Haiti	2005	Male	70 to 74	47% (30-65%)	53% (35-70%)	1.7 (0.87-2.8)	4.5% (1.8-8%)	5.2% (2-9.2%)	150 (55-260)	2800 (1100-4800)
Haiti	2005	Male	75 to 79	41% (27-60%)	59% (40-73%)	1.4 (0.68-2.4)	3.5% (1.3-6.4%)	3.9% (1.3-7.3%)	100 (34-190)	1500 (520-2700)
Haiti	2005	Male	80 to 84	36% (22-54%)	64% (46-78%)	1.1 (0.59-1.9)	2.8% (1-5.2%)	3.1% (1-5.9%)	62 (21-120)	670 (240-1300)
Haiti	2005	Male	85 to 89	33% (21-47%)	67% (53-79%)	1 (0.52-1.7)	2.2% (0.77-4.4%)	2.4% (0.75-4.9%)	29 (9.2-60)	230 (80-460)
Haiti	2005	Male	90 to 94	34% (22-49%)	66% (51-78%)	1 (0.49-1.8)	1.9% (0.58-4.2%)	2.1% (0.5-4.6%)	9.9 (2.4-22)	55 (16-120)
Haiti	2005	Male	95 plus	37% (22-56%)	63% (44-78%)	1.1 (0.45-2.1)	2% (0.34-4.7%)	2.1% (0.17-5.3%)	2.6 (0.21-6.7)	9.9 (1.8-24)
Haiti	2010	Female	All Ages	22% (18-26%)	78% (74-82%)	0.68 (0.57-0.81)	0.54% (0.26-1.1%)	0.77% (0.33-1.6%)	1000 (550-1600)	40000 (27000-57000)
Haiti	2010	Female	15 to 19	28% (19-39%)	72% (61-81%)	0.89 (0.49-1.4)	0.4% (0.14-1.1%)	0.37% (0.11-1.2%)	24 (12-41)	2100 (1200-3300)
Haiti	2010	Female	20 to 24	31% (22-42%)	69% (58-78%)	1.3 (0.78-1.9)	0.8% (0.31-2.1%)	0.69% (0.22-2%)	44 (24-72)	3800 (2400-5700)
Haiti	2010	Female	25 to 29	29% (20-40%)	71% (60-80%)	1.2 (0.75-1.8)	0.83% (0.35-1.9%)	0.72% (0.26-2%)	50 (27-82)	3900 (2500-6000)
Haiti	2010	Female	30 to 34	27% (17-39%)	73% (61-83%)	1.2 (0.67-1.8)	0.81% (0.37-1.7%)	0.72% (0.29-1.7%)	46 (26-79)	3200 (2000-5000)
Haiti	2010	Female	35 to 39	27% (18-39%)	73% (61-82%)	1.1 (0.61-1.7)	0.95% (0.43-1.9%)	0.92% (0.37-2.1%)	54 (28-94)	3200 (1800-5200)
Haiti	2010	Female	40 to 44	27% (18-38%)	73% (62-82%)	1 (0.55-1.7)	1.1% (0.5-2.1%)	1.2% (0.47-2.4%)	69 (35-120)	3500 (1900-5700)
Haiti	2010	Female	45 to 49	28% (17-39%)	72% (61-83%)	1.1 (0.59-1.9)	1.4% (0.6-2.8%)	1.5% (0.58-3.2%)	86 (41-160)	3800 (1900-6700)
Haiti	2010	Female	50 to 54	28% (18-42%)	72% (58-82%)	1.1 (0.58-2)	1.8% (0.7-3.8%)	2% (0.74-4.4%)	110 (49-210)	4200 (1800-7700)
Haiti	2010	Female	55 to 59	26% (16-40%)	74% (60-84%)	1 (0.51-1.8)	1.8% (0.65-3.7%)	2% (0.67-4.3%)	120 (49-220)	3800 (1600-7000)
Haiti	2010	Female	60 to 64	23% (13-36%)	77% (64-87%)	0.85 (0.41-1.6)	1.5% (0.38-3.2%)	1.6% (0.42-3.5%)	95 (28-200)	2600 (780-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	2010	Female	65 to 69	21% (13-32%)	79% (68-87%)	0.74 (0.39-1.2)	1.3% (0.27-2.8%)	1.4% (0.29-3.2%)	95 (21-200)	2200 (490-4400)
Haiti	2010	Female	70 to 74	19% (12-29%)	81% (71-88%)	0.62 (0.31-1.1)	1.1% (0.1-2.7%)	1.2% (0.11-3%)	84 (8.3-200)	1500 (150-3700)
Haiti	2010	Female	75 to 79	16% (9.8-26%)	84% (74-90%)	0.53 (0.24-0.98)	0.89% (-0.048-2.4%)	0.97% (-0.068-2.6%)	61 (-4.4-160)	870 (-55-2200)
Haiti	2010	Female	80 to 84	15% (9.1-24%)	85% (76-91%)	0.5 (0.24-0.85)	0.88% (0.053-2.2%)	0.96% (0.049-2.4%)	48 (2.8-110)	510 (36-1200)
Haiti	2010	Female	85 to 89	15% (9-24%)	85% (76-91%)	0.51 (0.24-0.91)	0.9% (0.048-2.3%)	0.97% (0.023-2.5%)	27 (0.46-70)	210 (8.6-510)
Haiti	2010	Female	90 to 94	17% (10-26%)	83% (74-90%)	0.57 (0.27-1.1)	1% (0.035-2.6%)	1.1% (-0.016-2.9%)	11 (-0.16-29)	57 (1.9-150)
Haiti	2010	Female	95 plus	20% (11-32%)	80% (68-89%)	0.68 (0.26-1.4)	1.3% (-0.12-3.5%)	1.3% (-0.23-3.9%)	3.2 (-0.56-10)	12 (-1.1-35)
Haiti	2010	Male	All Ages	47% (41-53%)	53% (47-59%)	2 (1.8-2.3)	1.2% (0.6-2.9%)	1.7% (0.77-4.1%)	2800 (1900-3900)	120000 (88000-160000)
Haiti	2010	Male	15 to 19	55% (44-68%)	45% (32-56%)	1.8 (1.1-2.7)	0.61% (0.21-1.9%)	0.53% (0.16-1.7%)	67 (39-110)	5800 (3600-8900)
Haiti	2010	Male	20 to 24	68% (58-77%)	32% (23-42%)	3.6 (2.5-4.8)	1.6% (0.61-5.2%)	1.5% (0.49-5.2%)	190 (120-300)	15000 (10000-22000)
Haiti	2010	Male	25 to 29	68% (59-77%)	32% (23-41%)	4.3 (2.9-5.8)	2% (0.79-6%)	1.8% (0.65-5.8%)	230 (140-350)	16000 (11000-24000)
Haiti	2010	Male	30 to 34	66% (57-74%)	34% (26-43%)	4.3 (2.9-5.9)	2.3% (0.96-6%)	2.1% (0.81-6%)	210 (130-320)	14000 (9100-20000)
Haiti	2010	Male	35 to 39	63% (54-73%)	37% (27-46%)	3.9 (2.4-5.7)	2.5% (1.1-6.2%)	2.4% (0.97-6.3%)	200 (120-300)	11000 (7200-16000)
Haiti	2010	Male	40 to 44	61% (52-71%)	39% (29-48%)	3.6 (2.2-5.5)	2.7% (1.3-6.1%)	2.6% (1.1-6.5%)	220 (130-330)	11000 (7000-16000)
Haiti	2010	Male	45 to 49	61% (50-71%)	39% (29-50%)	3.5 (2.2-5.4)	3.1% (1.5-6.9%)	3.1% (1.4-7.4%)	250 (140-370)	11000 (6800-16000)
Haiti	2010	Male	50 to 54	58% (47-69%)	42% (31-53%)	3.2 (1.9-4.9)	3.4% (1.5-7.6%)	3.5% (1.4-8.3%)	250 (140-380)	9800 (5600-14000)
Haiti	2010	Male	55 to 59	54% (42-66%)	46% (34-58%)	2.8 (1.6-4.4)	3.8% (1.7-7.8%)	3.9% (1.6-8.7%)	250 (140-390)	8500 (5000-13000)
Haiti	2010	Male	60 to 64	51% (35-68%)	49% (32-65%)	2.5 (1.3-4.3)	3.6% (1.6-7.1%)	3.7% (1.5-8.1%)	210 (110-340)	6000 (3100-9600)
Haiti	2010	Male	65 to 69	50% (33-67%)	50% (33-67%)	2.2 (1.2-3.6)	3.7% (1.7-6.9%)	3.9% (1.7-7.5%)	230 (120-380)	5500 (2900-8800)
Haiti	2010	Male	70 to 74	47% (30-64%)	53% (36-70%)	1.9 (0.96-3.1)	3.1% (1.2-6%)	3.3% (1.2-6.5%)	190 (76-330)	3600 (1500-6100)
Haiti	2010	Male	75 to 79	41% (27-59%)	59% (41-73%)	1.5 (0.74-2.6)	2.6% (0.89-5.1%)	2.7% (0.87-5.5%)	130 (46-250)	1900 (710-3500)
Haiti	2010	Male	80 to 84	37% (23-54%)	63% (46-77%)	1.3 (0.66-2.1)	2.2% (0.76-4.4%)	2.3% (0.74-4.8%)	81 (30-150)	880 (330-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	2010	Male	85 to 89	33% (21-48%)	67% (52-79%)	1.1 (0.57-1.9)	1.9% (0.72-3.8%)	2% (0.68-4.2%)	40 (14-78)	310 (120-600)
Haiti	2010	Male	90 to 94	34% (22-48%)	66% (52-78%)	1.1 (0.55-2)	1.8% (0.54-3.7%)	1.9% (0.45-4%)	13 (3.4-28)	73 (22-150)
Haiti	2010	Male	95 plus	37% (22-55%)	63% (45-78%)	1.2 (0.51-2.3)	1.9% (0.34-4.3%)	1.9% (0.22-4.7%)	3.6 (0.46-8.8)	14 (2.8-32)
Haiti	2016	Female	All Ages	23% (19-27%)	77% (73-81%)	0.74 (0.56-0.97)	1.7% (1-2.6%)	2.3% (1.1-3.9%)	1100 (530-2000)	44000 (26000-67000)
Haiti	2016	Female	15 to 19	30% (22-41%)	70% (59-78%)	0.98 (0.52-1.6)	1.8% (1-2.9%)	2.7% (1.3-4.5%)	23 (11-39)	2100 (1100-3300)
Haiti	2016	Female	20 to 24	33% (25-44%)	67% (56-75%)	1.4 (0.8-2.2)	3.4% (2-5.1%)	4.6% (2.4-7.4%)	42 (21-74)	3800 (2200-6000)
Haiti	2016	Female	25 to 29	30% (22-41%)	70% (59-78%)	1.3 (0.79-2.1)	3.2% (2.1-4.8%)	4.1% (2.2-6.6%)	49 (25-83)	4000 (2400-6100)
Haiti	2016	Female	30 to 34	28% (19-39%)	72% (61-81%)	1.2 (0.7-1.9)	2.8% (1.7-4.1%)	3.5% (2-5.5%)	56 (30-93)	4000 (2400-6100)
Haiti	2016	Female	35 to 39	28% (20-41%)	72% (59-80%)	1.2 (0.63-1.9)	2.9% (1.8-4.5%)	3.7% (2.1-5.8%)	68 (33-120)	4000 (2200-6500)
Haiti	2016	Female	40 to 44	28% (20-39%)	72% (61-80%)	1.1 (0.58-1.9)	3.1% (1.7-5.1%)	4.1% (2.2-6.7%)	76 (37-130)	3900 (2000-6500)
Haiti	2016	Female	45 to 49	26% (17-39%)	74% (61-83%)	1.1 (0.53-1.9)	3.3% (1.7-5.7%)	4.4% (2.3-7.5%)	91 (43-170)	4000 (2000-7200)
Haiti	2016	Female	50 to 54	27% (17-40%)	73% (60-83%)	1.1 (0.54-1.9)	3.8% (1.7-6.7%)	4.8% (2.2-8.6%)	120 (53-230)	4700 (2100-8700)
Haiti	2016	Female	55 to 59	26% (16-39%)	74% (61-84%)	1 (0.49-1.8)	3.5% (1.4-6.6%)	4.4% (1.7-8.3%)	130 (50-250)	4100 (1600-8100)
Haiti	2016	Female	60 to 64	23% (13-36%)	77% (64-87%)	0.86 (0.41-1.5)	2.7% (0.74-5.6%)	3.3% (0.89-6.8%)	110 (30-240)	3100 (830-6700)
Haiti	2016	Female	65 to 69	21% (13-32%)	79% (68-87%)	0.75 (0.39-1.3)	2.2% (0.43-4.8%)	2.6% (0.48-5.4%)	98 (18-210)	2200 (420-4800)
Haiti	2016	Female	70 to 74	19% (12-29%)	81% (71-88%)	0.63 (0.29-1.2)	1.7% (0.12-4.3%)	2% (0.14-5%)	92 (6.4-230)	1700 (120-4100)
Haiti	2016	Female	75 to 79	16% (9.7-26%)	84% (74-90%)	0.54 (0.24-1)	1.3% (-0.033-3.6%)	1.5% (-0.03-4.1%)	75 (-1.7-200)	1100 (-26-2900)
Haiti	2016	Female	80 to 84	15% (8.8-24%)	85% (76-91%)	0.5 (0.24-0.89)	1.2% (0.012-3.1%)	1.4% (-0.011-3.6%)	64 (-0.5-170)	670 (6.9-1700)
Haiti	2016	Female	85 to 89	15% (8.9-23%)	85% (77-91%)	0.51 (0.23-0.98)	1.2% (0.072-3.3%)	1.3% (0.017-3.7%)	31 (0.38-89)	240 (14-660)
Haiti	2016	Female	90 to 94	17% (9.9-26%)	83% (74-90%)	0.57 (0.25-1.1)	1.3% (0.027-3.5%)	1.4% (-0.056-4%)	14 (-0.54-39)	71 (1.5-200)
Haiti	2016	Female	95 plus	20% (10-32%)	80% (68-90%)	0.68 (0.25-1.4)	1.4% (-0.19-4.2%)	1.5% (-0.33-4.8%)	3.9 (-0.86-12)	14 (-1.9-43)
Haiti	2016	Male	All Ages	50% (44-55%)	50% (45-56%)	2.2 (1.7-2.9)	5.5% (3.8-7.4%)	7% (4.7-9.5%)	3200 (2000-4600)	140000 (96000-190000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Haiti	2016	Male	15 to 19	57% (47-68%)	43% (32-53%)	1.9 (1.1-2.9)	4.8% (3.1-7.2%)	6.1% (3.6-9.6%)	72 (39-120)	6100 (3700-9600)
Haiti	2016	Male	20 to 24	70% (61-78%)	30% (22-39%)	3.8 (2.4-5.6)	12% (8.3-17%)	15% (9.6-21%)	210 (120-330)	16000 (10000-24000)
Haiti	2016	Male	25 to 29	70% (61-77%)	30% (23-39%)	4.5 (2.9-6.5)	13% (9.1-18%)	16% (11-22%)	250 (150-380)	18000 (12000-26000)
Haiti	2016	Male	30 to 34	68% (60-76%)	32% (24-40%)	4.5 (2.9-6.6)	12% (8.8-16%)	15% (10-21%)	280 (160-430)	18000 (11000-26000)
Haiti	2016	Male	35 to 39	65% (56-74%)	35% (26-44%)	4.1 (2.5-6.2)	11% (7.6-15%)	13% (8.6-19%)	260 (150-390)	15000 (9000-22000)
Haiti	2016	Male	40 to 44	63% (55-72%)	37% (28-45%)	3.9 (2.2-5.9)	11% (7.2-14%)	13% (8.3-17%)	240 (140-370)	12000 (7400-18000)
Haiti	2016	Male	45 to 49	63% (53-72%)	37% (28-47%)	3.7 (2.2-5.8)	11% (6.9-15%)	13% (8.1-18%)	270 (150-410)	12000 (7300-18000)
Haiti	2016	Male	50 to 54	60% (50-70%)	40% (30-50%)	3.4 (1.9-5.4)	10% (6.2-14%)	12% (7.2-17%)	300 (160-450)	12000 (6600-17000)
Haiti	2016	Male	55 to 59	55% (44-67%)	45% (33-56%)	2.9 (1.6-4.7)	9.5% (5.6-14%)	12% (6.6-17%)	290 (160-450)	9900 (5500-15000)
Haiti	2016	Male	60 to 64	51% (36-68%)	49% (32-64%)	2.5 (1.3-4.4)	7.9% (4.1-12%)	9.4% (4.8-15%)	280 (130-450)	7800 (3900-13000)
Haiti	2016	Male	65 to 69	50% (33-68%)	50% (32-67%)	2.3 (1.2-3.7)	6.9% (3.6-11%)	8.1% (4.1-13%)	240 (110-400)	5500 (2600-9100)
Haiti	2016	Male	70 to 74	47% (30-64%)	53% (36-70%)	1.9 (0.9-3.3)	5.3% (2.1-9.2%)	6.1% (2.3-11%)	210 (77-370)	3900 (1500-6900)
Haiti	2016	Male	75 to 79	41% (27-60%)	59% (40-73%)	1.5 (0.75-2.8)	4.1% (1.4-7.7%)	4.6% (1.5-8.7%)	160 (52-300)	2300 (790-4300)
Haiti	2016	Male	80 to 84	37% (22-53%)	63% (47-78%)	1.3 (0.67-2.2)	3.3% (1.1-6.2%)	3.7% (1.2-7%)	100 (33-200)	1100 (390-2100)
Haiti	2016	Male	85 to 89	33% (21-47%)	67% (53-79%)	1.1 (0.56-2)	2.7% (0.94-5.5%)	3% (0.94-6.3%)	44 (14-92)	340 (120-700)
Haiti	2016	Male	90 to 94	34% (21-49%)	66% (51-79%)	1.1 (0.52-2.1)	2.3% (0.66-5%)	2.5% (0.58-5.7%)	16 (3.7-36)	88 (25-190)
Haiti	2016	Male	95 plus	37% (22-55%)	63% (45-78%)	1.2 (0.5-2.3)	2.4% (0.44-5.6%)	2.5% (0.23-6.3%)	3.9 (0.37-9.8)	15 (2.8-35)
Jamaica	1990	Female	All Ages	25% (20-31%)	75% (69-80%)	0.82 (0.67-0.99)	1.5% (0.92-2.1%)	2% (0.78-3.6%)	150 (57-270)	4900 (3100-7300)
Jamaica	1990	Female	15 to 19	31% (20-47%)	69% (53-80%)	1 (0.55-1.8)	1.6% (1-2.4%)	3.3% (1.6-5.6%)	1.8 (0.89-3.2)	250 (140-380)
Jamaica	1990	Female	20 to 24	37% (23-53%)	63% (47-77%)	1.6 (0.88-2.5)	3% (2-4.1%)	5.1% (2.8-7.8%)	3.3 (1.8-5.3)	460 (300-670)
Jamaica	1990	Female	25 to 29	37% (22-54%)	63% (46-78%)	1.6 (0.95-2.5)	3.1% (2.2-4.3%)	5.2% (3-7.8%)	3.7 (2.2-5.6)	450 (310-640)
Jamaica	1990	Female	30 to 34	35% (21-52%)	65% (48-79%)	1.6 (0.89-2.5)	3.1% (2-4.4%)	5.2% (3.1-7.9%)	4.6 (2.7-7.1)	430 (280-620)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	1990	Female	35 to 39	34% (21-51%)	66% (49-79%)	1.4 (0.76-2.4)	2.9% (1.8-4.3%)	5% (2.8-7.8%)	4.7 (2.6-7.4)	350 (220-530)
Jamaica	1990	Female	40 to 44	33% (20-50%)	67% (50-80%)	1.4 (0.68-2.3)	3.1% (1.7-4.9%)	5.4% (2.7-8.9%)	6 (3-9.9)	350 (190-570)
Jamaica	1990	Female	45 to 49	31% (18-47%)	69% (53-82%)	1.3 (0.64-2.3)	3.2% (1.6-5.5%)	5.3% (2.6-9.2%)	7.5 (3.6-13)	370 (180-630)
Jamaica	1990	Female	50 to 54	28% (16-44%)	72% (56-84%)	1.2 (0.58-2.1)	3.3% (1.4-5.9%)	5% (2-8.8%)	10 (4.1-18)	410 (170-740)
Jamaica	1990	Female	55 to 59	24% (14-39%)	76% (61-86%)	0.97 (0.46-1.7)	2.7% (0.93-5.2%)	4.1% (1.4-7.5%)	11 (3.5-20)	370 (130-700)
Jamaica	1990	Female	60 to 64	21% (12-36%)	79% (64-88%)	0.82 (0.38-1.5)	2.3% (0.47-4.6%)	3.1% (0.69-6.3%)	13 (2.7-26)	370 (76-760)
Jamaica	1990	Female	65 to 69	20% (12-31%)	80% (69-88%)	0.71 (0.36-1.2)	1.9% (0.38-4%)	2.5% (0.51-5.4%)	14 (2.8-29)	330 (65-710)
Jamaica	1990	Female	70 to 74	17% (10-28%)	83% (72-90%)	0.6 (0.28-1.1)	1.5% (0.074-3.7%)	2% (0.096-4.9%)	14 (0.69-35)	260 (14-660)
Jamaica	1990	Female	75 to 79	15% (8.7-26%)	85% (74-91%)	0.51 (0.22-0.98)	1.2% (-0.018-3.1%)	1.6% (-0.038-3.9%)	14 (-0.32-35)	200 (-2.6-520)
Jamaica	1990	Female	80 to 84	14% (7.6-23%)	86% (77-92%)	0.48 (0.21-0.85)	1.2% (0.11-2.7%)	1.5% (0.13-3.5%)	16 (1.4-38)	180 (18-420)
Jamaica	1990	Female	85 to 89	14% (7.7-23%)	86% (77-92%)	0.49 (0.23-0.91)	1.2% (0.15-2.9%)	1.4% (0.15-3.5%)	12 (1.3-30)	96 (12-230)
Jamaica	1990	Female	90 to 94	16% (9-26%)	84% (74-91%)	0.56 (0.24-1.1)	1.4% (0.12-3.5%)	1.6% (0.066-4.2%)	8.9 (0.35-23)	47 (3.8-120)
Jamaica	1990	Female	95 plus	19% (9.5-33%)	81% (67-91%)	0.68 (0.24-1.5)	1.7% (-0.018-4.7%)	1.9% (-0.14-5.4%)	3.8 (-0.28-11)	13 (-0.15-36)
Jamaica	1990	Male	All Ages	45% (36-54%)	55% (46-64%)	1.9 (1.5-2.2)	4.2% (3.2-5.3%)	5.4% (3.4-7.5%)	390 (250-530)	14000 (11000-18000)
Jamaica	1990	Male	15 to 19	48% (31-66%)	52% (34-69%)	1.6 (0.86-2.6)	4.6% (2.9-6.6%)	7.3% (3.8-11%)	6.4 (3.3-10)	700 (450-1000)
Jamaica	1990	Male	20 to 24	60% (41-77%)	40% (23-59%)	3.1 (1.9-4.5)	10% (7.1-13%)	14% (8.6-20%)	17 (10-26)	1700 (1200-2300)
Jamaica	1990	Male	25 to 29	63% (44-79%)	37% (21-56%)	3.8 (2.4-5.6)	11% (7.7-14%)	15% (9.6-21%)	19 (12-27)	1600 (1200-2200)
Jamaica	1990	Male	30 to 34	64% (43-80%)	36% (20-57%)	4 (2.4-6.1)	10% (7.3-13%)	15% (10-21%)	18 (11-25)	1400 (960-1800)
Jamaica	1990	Male	35 to 39	64% (46-80%)	36% (20-54%)	3.8 (2.2-6)	9.4% (6.5-13%)	14% (9.1-20%)	16 (10-24)	1100 (750-1500)
Jamaica	1990	Male	40 to 44	63% (45-79%)	37% (21-55%)	3.6 (1.9-5.9)	8.8% (5.9-12%)	13% (8.1-18%)	17 (10-24)	970 (630-1300)
Jamaica	1990	Male	45 to 49	61% (40-77%)	39% (23-60%)	3.3 (1.7-5.6)	8.4% (5.3-12%)	12% (7.2-17%)	18 (11-26)	920 (570-1300)
Jamaica	1990	Male	50 to 54	59% (39-77%)	41% (23-61%)	3 (1.5-5)	7.9% (4.7-11%)	11% (6.3-16%)	22 (12-31)	920 (550-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	1990	Male	55 to 59	57% (38-75%)	43% (25-62%)	2.7 (1.4-4.5)	7.6% (4.3-11%)	10% (5.7-15%)	28 (15-40)	990 (570-1400)
Jamaica	1990	Male	60 to 64	54% (34-74%)	46% (26-66%)	2.4 (1.2-4.2)	6.7% (3.3-10%)	8.6% (4.3-13%)	34 (17-52)	1000 (510-1500)
Jamaica	1990	Male	65 to 69	53% (33-71%)	47% (29-67%)	2.1 (1.1-3.6)	6.2% (3.1-9.7%)	7.8% (4-12%)	42 (21-65)	1000 (510-1600)
Jamaica	1990	Male	70 to 74	49% (30-67%)	51% (33-70%)	1.7 (0.82-3.1)	4.9% (2.1-8.5%)	6.1% (2.6-11%)	42 (17-73)	810 (340-1400)
Jamaica	1990	Male	75 to 79	44% (26-63%)	56% (37-74%)	1.4 (0.68-2.6)	3.9% (1.5-6.9%)	4.7% (1.8-8.5%)	38 (15-69)	580 (230-1000)
Jamaica	1990	Male	80 to 84	38% (22-56%)	62% (44-78%)	1.2 (0.59-2.1)	3.3% (1.3-5.9%)	3.8% (1.5-7%)	35 (13-64)	380 (150-690)
Jamaica	1990	Male	85 to 89	34% (19-50%)	66% (50-81%)	1 (0.49-1.9)	2.7% (0.98-5.3%)	3.1% (1.1-6.3%)	23 (7.9-45)	180 (64-340)
Jamaica	1990	Male	90 to 94	34% (21-51%)	66% (49-79%)	1 (0.46-1.9)	2.5% (0.73-5%)	2.8% (0.73-5.8%)	9.5 (2.4-19)	51 (15-100)
Jamaica	1990	Male	95 plus	37% (20-57%)	63% (43-80%)	1.1 (0.44-2.3)	2.5% (0.49-5.6%)	2.7% (0.35-6.4%)	2.4 (0.31-5.6)	8.9 (1.8-20)
Jamaica	1995	Female	All Ages	25% (20-31%)	75% (69-80%)	0.79 (0.65-0.95)	1.4% (0.85-2.1%)	1.7% (0.55-3.1%)	140 (46-250)	4700 (2900-7100)
Jamaica	1995	Female	15 to 19	32% (20-47%)	68% (53-80%)	0.98 (0.51-1.7)	1.6% (0.94-2.4%)	3% (1.4-5.2%)	1.7 (0.82-3)	230 (130-360)
Jamaica	1995	Female	20 to 24	37% (23-54%)	63% (46-77%)	1.5 (0.83-2.4)	2.7% (1.8-3.9%)	4.2% (2.2-6.5%)	3.2 (1.6-5.1)	440 (280-650)
Jamaica	1995	Female	25 to 29	37% (22-54%)	63% (46-78%)	1.5 (0.89-2.4)	2.8% (2-3.8%)	4.1% (2.4-6.1%)	3.9 (2.1-6)	460 (310-670)
Jamaica	1995	Female	30 to 34	36% (21-53%)	64% (47-79%)	1.5 (0.84-2.4)	2.6% (1.8-3.7%)	4.1% (2.4-6.2%)	4.5 (2.5-7)	440 (290-630)
Jamaica	1995	Female	35 to 39	35% (21-52%)	65% (48-79%)	1.4 (0.75-2.2)	2.6% (1.6-3.9%)	4.2% (2.5-6.6%)	5.2 (2.9-8.6)	400 (240-620)
Jamaica	1995	Female	40 to 44	33% (21-51%)	67% (49-79%)	1.3 (0.65-2.1)	2.7% (1.4-4.4%)	4.6% (2.4-7.6%)	6.2 (3.2-11)	370 (200-610)
Jamaica	1995	Female	45 to 49	31% (18-48%)	69% (52-82%)	1.2 (0.59-2.2)	2.8% (1.4-4.9%)	4.5% (2.2-8.2%)	7.3 (3.4-13)	360 (170-650)
Jamaica	1995	Female	50 to 54	28% (16-45%)	72% (55-84%)	1.1 (0.54-2)	2.8% (1.1-5.1%)	4.3% (1.8-7.7%)	9 (3.7-17)	370 (150-700)
Jamaica	1995	Female	55 to 59	25% (14-40%)	75% (60-86%)	0.92 (0.44-1.6)	2.4% (0.77-4.7%)	3.5% (1.2-6.6%)	9.4 (3.1-19)	330 (110-650)
Jamaica	1995	Female	60 to 64	22% (12-36%)	78% (64-88%)	0.78 (0.36-1.4)	1.9% (0.25-4.1%)	2.6% (0.38-5.6%)	10 (1.5-22)	290 (40-670)
Jamaica	1995	Female	65 to 69	20% (12-31%)	80% (69-88%)	0.68 (0.35-1.2)	1.6% (0.16-3.7%)	2.1% (0.29-4.8%)	11 (1.5-25)	260 (29-610)
Jamaica	1995	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.56 (0.27-1)	1.2% (-0.12-3.3%)	1.6% (-0.1-4.2%)	11 (-0.68-31)	220 (-23-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	1995	Female	75 to 79	16% (8.9-26%)	84% (74-91%)	0.49 (0.21-0.93)	0.98% (-0.2-2.9%)	1.2% (-0.23-3.5%)	11 (-2.1-34)	170 (-30-500)
Jamaica	1995	Female	80 to 84	14% (7.6-24%)	86% (76-92%)	0.46 (0.21-0.82)	0.97% (-0.068-2.5%)	1.2% (-0.078-3%)	15 (-1-39)	160 (-10-410)
Jamaica	1995	Female	85 to 89	14% (8-23%)	86% (77-92%)	0.46 (0.22-0.85)	1% (-0.015-2.6%)	1.2% (-0.069-3.2%)	15 (-0.93-39)	120 (-1.7-300)
Jamaica	1995	Female	90 to 94	16% (9.1-26%)	84% (74-91%)	0.53 (0.22-1)	1.2% (-0.071-3.2%)	1.3% (-0.13-3.8%)	8.6 (-0.86-24)	46 (-2.8-130)
Jamaica	1995	Female	95 plus	19% (9.5-34%)	81% (66-90%)	0.65 (0.22-1.3)	1.4% (-0.17-4.2%)	1.6% (-0.33-4.9%)	4 (-0.85-13)	14 (-1.7-43)
Jamaica	1995	Male	All Ages	46% (37-55%)	54% (45-63%)	1.8 (1.5-2.1)	4.3% (3.3-5.4%)	5% (3.2-7%)	400 (250-570)	15000 (11000-19000)
Jamaica	1995	Male	15 to 19	49% (31-67%)	51% (33-69%)	1.5 (0.8-2.4)	4.5% (2.8-6.6%)	7% (3.5-12%)	6.6 (3.4-11)	700 (430-1000)
Jamaica	1995	Male	20 to 24	60% (41-77%)	40% (23-59%)	2.9 (1.8-4.3)	9.5% (6.5-13%)	12% (7.5-19%)	18 (10-27)	1700 (1100-2400)
Jamaica	1995	Male	25 to 29	64% (44-79%)	36% (21-56%)	3.6 (2.2-5.1)	9.6% (6.8-13%)	13% (7.9-19%)	20 (12-29)	1700 (1200-2400)
Jamaica	1995	Male	30 to 34	65% (44-80%)	35% (20-56%)	3.7 (2.2-5.5)	8.9% (6.2-12%)	12% (7.7-17%)	19 (12-29)	1500 (1000-2100)
Jamaica	1995	Male	35 to 39	64% (46-80%)	36% (20-54%)	3.5 (2-5.5)	8.4% (5.9-12%)	12% (7.5-17%)	20 (12-29)	1300 (890-1900)
Jamaica	1995	Male	40 to 44	63% (45-79%)	37% (21-55%)	3.3 (1.8-5.3)	8% (5.1-11%)	11% (6.8-16%)	19 (11-28)	1100 (710-1600)
Jamaica	1995	Male	45 to 49	61% (39-77%)	39% (23-61%)	3.1 (1.6-5.3)	7.5% (4.7-11%)	10% (6.1-15%)	20 (12-31)	1000 (620-1500)
Jamaica	1995	Male	50 to 54	59% (39-77%)	41% (23-61%)	2.8 (1.4-4.7)	7.2% (4.1-10%)	9.7% (5.4-14%)	24 (13-35)	1000 (580-1500)
Jamaica	1995	Male	55 to 59	57% (39-75%)	43% (25-61%)	2.5 (1.3-4.2)	6.9% (3.8-10%)	9.1% (5-13%)	29 (15-43)	1000 (570-1500)
Jamaica	1995	Male	60 to 64	55% (35-74%)	45% (26-65%)	2.3 (1.1-4)	6.1% (3.1-9.4%)	7.8% (3.8-12%)	34 (17-53)	1000 (510-1600)
Jamaica	1995	Male	65 to 69	53% (34-71%)	47% (29-66%)	2 (1.1-3.3)	5.6% (2.8-9.2%)	7% (3.5-11%)	43 (21-71)	1000 (530-1700)
Jamaica	1995	Male	70 to 74	49% (31-67%)	51% (33-69%)	1.6 (0.78-2.9)	4.4% (1.8-7.7%)	5.3% (2.2-9.4%)	41 (16-72)	800 (330-1400)
Jamaica	1995	Male	75 to 79	44% (26-64%)	56% (36-74%)	1.3 (0.62-2.5)	3.4% (1.2-6.3%)	4.1% (1.5-7.5%)	34 (12-63)	510 (190-940)
Jamaica	1995	Male	80 to 84	38% (22-56%)	62% (44-78%)	1.1 (0.55-1.9)	2.8% (1.1-5.2%)	3.2% (1.2-6.1%)	31 (11-58)	340 (130-640)
Jamaica	1995	Male	85 to 89	34% (19-50%)	66% (50-81%)	0.96 (0.47-1.8)	2.3% (0.82-4.6%)	2.7% (0.87-5.4%)	24 (8.1-49)	190 (68-390)
Jamaica	1995	Male	90 to 94	34% (21-51%)	66% (49-79%)	0.98 (0.44-1.8)	2.1% (0.58-4.5%)	2.3% (0.57-5.1%)	12 (2.9-26)	64 (18-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	1995	Male	95 plus	38% (20-58%)	62% (42-80%)	1.1 (0.41-2.1)	2.2% (0.4-5.1%)	2.3% (0.24-5.7%)	3.3 (0.35-8.2)	12 (2.3-29)
Jamaica	2000	Female	All Ages	26% (21-33%)	74% (67-79%)	0.75 (0.62-0.89)	1.3% (0.74-1.9%)	1.3% (0.38-2.6%)	120 (34-230)	4400 (2600-6600)
Jamaica	2000	Female	15 to 19	33% (21-48%)	67% (52-79%)	0.93 (0.49-1.6)	1.4% (0.87-2.1%)	2.6% (1.2-4.4%)	1.4 (0.63-2.4)	200 (120-310)
Jamaica	2000	Female	20 to 24	38% (24-55%)	62% (45-76%)	1.4 (0.75-2.2)	2.5% (1.6-3.4%)	3.3% (1.7-5.2%)	2.4 (1.2-3.9)	370 (240-530)
Jamaica	2000	Female	25 to 29	38% (23-55%)	62% (45-77%)	1.4 (0.81-2.2)	2.5% (1.7-3.5%)	3% (1.7-4.6%)	3.8 (2.1-5.8)	450 (300-640)
Jamaica	2000	Female	30 to 34	36% (22-53%)	64% (47-78%)	1.4 (0.78-2.1)	2.3% (1.6-3.3%)	3.1% (1.9-4.8%)	4.7 (2.8-7.4)	460 (300-660)
Jamaica	2000	Female	35 to 39	35% (22-52%)	65% (48-78%)	1.3 (0.7-2)	2.3% (1.5-3.4%)	3.3% (2-5.2%)	5.9 (3.3-9.6)	450 (290-690)
Jamaica	2000	Female	40 to 44	34% (21-52%)	66% (48-79%)	1.2 (0.62-2)	2.3% (1.3-3.8%)	3.7% (2-6%)	7 (3.8-11)	420 (240-660)
Jamaica	2000	Female	45 to 49	32% (18-49%)	68% (51-82%)	1.2 (0.57-2)	2.4% (1.3-4.3%)	3.8% (1.9-6.8%)	7.6 (3.7-13)	370 (190-670)
Jamaica	2000	Female	50 to 54	29% (17-45%)	71% (55-83%)	1 (0.52-1.8)	2.3% (0.88-4.4%)	3.5% (1.4-6.5%)	8.4 (3.4-16)	350 (130-650)
Jamaica	2000	Female	55 to 59	25% (15-40%)	75% (60-85%)	0.85 (0.4-1.5)	1.9% (0.53-3.8%)	2.8% (0.87-5.4%)	8.2 (2.6-16)	290 (80-590)
Jamaica	2000	Female	60 to 64	22% (13-36%)	78% (64-87%)	0.72 (0.33-1.3)	1.5% (0.011-3.3%)	2% (0.12-4.5%)	8.5 (0.49-19)	250 (1.9-570)
Jamaica	2000	Female	65 to 69	20% (12-32%)	80% (68-88%)	0.63 (0.32-1.1)	1.1% (-0.071-2.9%)	1.5% (0.008-3.9%)	9 (0.047-23)	210 (-13-560)
Jamaica	2000	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.53 (0.26-1)	0.88% (-0.3-2.8%)	1.2% (-0.31-3.6%)	9.2 (-2.2-27)	170 (-57-530)
Jamaica	2000	Female	75 to 79	16% (9.2-27%)	84% (73-91%)	0.46 (0.2-0.88)	0.72% (-0.4-2.3%)	0.93% (-0.45-3%)	8.5 (-4.1-27)	120 (-66-410)
Jamaica	2000	Female	80 to 84	15% (8.1-25%)	85% (75-92%)	0.43 (0.19-0.76)	0.69% (-0.26-2.1%)	0.84% (-0.31-2.5%)	10 (-3.8-31)	110 (-40-340)
Jamaica	2000	Female	85 to 89	15% (8.2-24%)	85% (76-92%)	0.44 (0.21-0.8)	0.74% (-0.19-2.2%)	0.85% (-0.23-2.6%)	11 (-2.9-34)	86 (-21-260)
Jamaica	2000	Female	90 to 94	17% (9.5-26%)	83% (74-91%)	0.49 (0.21-0.96)	0.84% (-0.21-2.7%)	0.93% (-0.33-3.2%)	8.2 (-2.9-28)	45 (-11-140)
Jamaica	2000	Female	95 plus	20% (9.7-34%)	80% (66-90%)	0.6 (0.21-1.2)	1% (-0.36-3.5%)	1.1% (-0.57-4.1%)	3.4 (-1.8-13)	13 (-4.4-44)
Jamaica	2000	Male	All Ages	47% (39-56%)	53% (44-61%)	1.7 (1.4-2)	4.2% (3.2-5.2%)	4.4% (2.9-6.2%)	400 (260-570)	16000 (12000-20000)
Jamaica	2000	Male	15 to 19	49% (32-67%)	51% (33-68%)	1.4 (0.75-2.3)	4.2% (2.6-6.1%)	6.3% (3.3-10%)	6 (3.1-10)	650 (410-960)
Jamaica	2000	Male	20 to 24	61% (42-77%)	39% (23-58%)	2.7 (1.6-4)	9% (6.3-12%)	11% (6.8-17%)	17 (10-26)	1600 (1100-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	2000	Male	25 to 29	64% (45-80%)	36% (20-55%)	3.3 (2-4.8)	9% (6.2-12%)	11% (6.8-16%)	21 (13-31)	1800 (1200-2500)
Jamaica	2000	Male	30 to 34	65% (45-81%)	35% (19-55%)	3.5 (2.1-5)	8.1% (5.7-11%)	10% (6.5-14%)	23 (14-33)	1700 (1200-2400)
Jamaica	2000	Male	35 to 39	65% (47-80%)	35% (20-53%)	3.2 (1.9-5)	7.3% (5-9.9%)	9.1% (5.7-13%)	22 (14-33)	1500 (1000-2100)
Jamaica	2000	Male	40 to 44	64% (47-79%)	36% (21-53%)	3.1 (1.7-4.8)	6.9% (4.6-9.6%)	8.9% (5.4-13%)	23 (14-33)	1400 (880-1900)
Jamaica	2000	Male	45 to 49	62% (41-78%)	38% (22-59%)	2.9 (1.5-4.9)	6.7% (4.2-9.4%)	8.6% (5-13%)	22 (13-34)	1100 (700-1700)
Jamaica	2000	Male	50 to 54	60% (40-78%)	40% (22-60%)	2.6 (1.3-4.5)	6.3% (3.7-9.3%)	8.1% (4.4-12%)	25 (13-37)	1100 (610-1600)
Jamaica	2000	Male	55 to 59	58% (40-76%)	42% (24-60%)	2.3 (1.2-4)	6.1% (3.2-9.3%)	7.8% (4.1-12%)	30 (16-47)	1100 (580-1700)
Jamaica	2000	Male	60 to 64	55% (35-74%)	45% (26-65%)	2.1 (1-3.6)	5.4% (2.6-8.6%)	6.7% (3.1-11%)	34 (16-53)	1000 (480-1600)
Jamaica	2000	Male	65 to 69	54% (35-72%)	46% (28-65%)	1.8 (0.97-3)	4.9% (2.4-8.2%)	5.9% (2.9-9.9%)	42 (20-71)	1000 (500-1700)
Jamaica	2000	Male	70 to 74	50% (32-67%)	50% (33-68%)	1.5 (0.75-2.7)	3.8% (1.5-6.7%)	4.5% (1.8-8%)	41 (17-75)	820 (330-1400)
Jamaica	2000	Male	75 to 79	45% (27-65%)	55% (35-73%)	1.2 (0.58-2.2)	2.9% (1.1-5.5%)	3.4% (1.2-6.4%)	33 (12-65)	510 (190-990)
Jamaica	2000	Male	80 to 84	39% (22-57%)	61% (43-78%)	1 (0.51-1.8)	2.3% (0.86-4.4%)	2.6% (0.96-5%)	27 (9.9-52)	310 (110-570)
Jamaica	2000	Male	85 to 89	35% (20-50%)	65% (50-80%)	0.9 (0.44-1.6)	1.9% (0.69-3.9%)	2.1% (0.7-4.4%)	21 (6.8-43)	170 (61-340)
Jamaica	2000	Male	90 to 94	35% (21-51%)	65% (49-79%)	0.9 (0.43-1.7)	1.7% (0.46-3.7%)	1.8% (0.42-4.2%)	12 (2.7-27)	67 (18-140)
Jamaica	2000	Male	95 plus	38% (20-59%)	62% (41-80%)	1 (0.4-1.9)	1.8% (0.3-4.3%)	1.8% (0.12-4.8%)	4.3 (0.28-11)	17 (2.8-40)
Jamaica	2005	Female	All Ages	26% (21-32%)	74% (68-79%)	0.81 (0.67-0.98)	1.5% (0.91-2.2%)	1.6% (0.62-3.1%)	150 (56-280)	5200 (3200-7700)
Jamaica	2005	Female	15 to 19	32% (20-48%)	68% (52-80%)	0.97 (0.51-1.7)	1.5% (0.91-2.3%)	3% (1.5-5.2%)	1.5 (0.74-2.6)	210 (130-330)
Jamaica	2005	Female	20 to 24	38% (23-54%)	62% (46-77%)	1.5 (0.78-2.4)	2.7% (1.8-3.8%)	3.8% (2-6.1%)	3.3 (1.7-5.4)	430 (280-630)
Jamaica	2005	Female	25 to 29	37% (22-54%)	63% (46-78%)	1.5 (0.84-2.4)	2.6% (1.8-3.7%)	3.4% (1.9-5.1%)	3.9 (2.1-6.2)	450 (300-640)
Jamaica	2005	Female	30 to 34	36% (21-53%)	64% (47-79%)	1.5 (0.81-2.3)	2.4% (1.6-3.5%)	3.2% (1.9-5.1%)	4.5 (2.6-7.1)	440 (280-640)
Jamaica	2005	Female	35 to 39	35% (22-52%)	65% (48-78%)	1.3 (0.74-2.2)	2.4% (1.6-3.6%)	3.7% (2.2-5.6%)	6.9 (4.1-11)	520 (340-780)
Jamaica	2005	Female	40 to 44	34% (21-52%)	66% (48-79%)	1.3 (0.66-2.1)	2.5% (1.4-3.8%)	3.9% (2.2-6.2%)	8.8 (4.8-14)	540 (300-840)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	2005	Female	45 to 49	32% (18-49%)	68% (51-82%)	1.2 (0.6-2.1)	2.6% (1.4-4.5%)	4.1% (2.1-7%)	11 (5.5-18)	530 (280-920)
Jamaica	2005	Female	50 to 54	29% (17-45%)	71% (55-83%)	1.1 (0.55-1.9)	2.5% (1.1-4.6%)	3.9% (1.7-6.8%)	11 (5.1-20)	470 (210-850)
Jamaica	2005	Female	55 to 59	25% (15-40%)	75% (60-85%)	0.91 (0.43-1.6)	2.1% (0.66-4.2%)	3.2% (1.1-6%)	9.6 (3.3-18)	340 (100-660)
Jamaica	2005	Female	60 to 64	22% (13-36%)	78% (64-87%)	0.77 (0.36-1.4)	1.7% (0.2-3.7%)	2.4% (0.4-4.9%)	10 (1.8-21)	300 (37-650)
Jamaica	2005	Female	65 to 69	20% (12-32%)	80% (68-88%)	0.67 (0.34-1.1)	1.4% (0.076-3.2%)	1.9% (0.25-4.3%)	10 (1.3-24)	250 (14-590)
Jamaica	2005	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.56 (0.26-1.1)	1.1% (-0.15-3.1%)	1.5% (-0.093-4%)	11 (-0.75-31)	220 (-28-630)
Jamaica	2005	Female	75 to 79	16% (9.2-26%)	84% (74-91%)	0.49 (0.22-0.92)	0.87% (-0.3-2.6%)	1.1% (-0.3-3.4%)	11 (-3.1-33)	170 (-60-510)
Jamaica	2005	Female	80 to 84	15% (8.1-24%)	85% (76-92%)	0.45 (0.21-0.8)	0.84% (-0.098-2.4%)	1% (-0.12-2.9%)	13 (-1.6-38)	150 (-18-420)
Jamaica	2005	Female	85 to 89	15% (8.1-23%)	85% (77-92%)	0.46 (0.22-0.84)	0.9% (-0.085-2.5%)	1.1% (-0.11-3%)	14 (-1.4-39)	110 (-9.8-300)
Jamaica	2005	Female	90 to 94	16% (9.3-27%)	84% (73-91%)	0.52 (0.22-1)	1% (-0.12-3%)	1.2% (-0.2-3.6%)	11 (-1.9-33)	59 (-6.5-170)
Jamaica	2005	Female	95 plus	20% (9.6-35%)	80% (65-90%)	0.63 (0.22-1.3)	1.2% (-0.29-3.8%)	1.4% (-0.45-4.5%)	5.9 (-1.9-19)	22 (-4.9-66)
Jamaica	2005	Male	All Ages	47% (39-56%)	53% (44-61%)	1.8 (1.5-2.2)	4.9% (3.8-6.1%)	5% (3.5-6.9%)	460 (310-650)	18000 (14000-24000)
Jamaica	2005	Male	15 to 19	49% (31-67%)	51% (33-69%)	1.5 (0.79-2.4)	4.8% (2.9-7.1%)	7.3% (3.7-12%)	8.6 (4.3-14)	850 (510-1300)
Jamaica	2005	Male	20 to 24	61% (42-77%)	39% (23-58%)	2.9 (1.7-4.4)	11% (7.1-15%)	13% (7.7-20%)	28 (16-41)	2300 (1500-3300)
Jamaica	2005	Male	25 to 29	64% (44-79%)	36% (21-56%)	3.5 (2.1-5.2)	11% (7.3-15%)	14% (8.3-20%)	29 (17-43)	2200 (1500-3100)
Jamaica	2005	Male	30 to 34	65% (44-80%)	35% (20-56%)	3.7 (2.2-5.5)	9.5% (6.5-13%)	12% (7.7-17%)	28 (18-41)	2000 (1400-2800)
Jamaica	2005	Male	35 to 39	65% (47-80%)	35% (20-53%)	3.5 (2-5.3)	8.1% (5.7-11%)	10% (6.6-15%)	26 (17-39)	1700 (1200-2400)
Jamaica	2005	Male	40 to 44	63% (46-79%)	37% (21-54%)	3.3 (1.8-5.1)	7.2% (4.9-10%)	9.4% (6-14%)	26 (16-38)	1500 (1000-2200)
Jamaica	2005	Male	45 to 49	61% (41-78%)	39% (22-59%)	3 (1.6-4.9)	7% (4.5-9.7%)	9.2% (5.6-13%)	25 (15-36)	1300 (800-1800)
Jamaica	2005	Male	50 to 54	59% (40-77%)	41% (23-60%)	2.7 (1.4-4.6)	6.5% (3.9-9.4%)	8.6% (5.1-13%)	29 (17-43)	1300 (770-1900)
Jamaica	2005	Male	55 to 59	57% (39-76%)	43% (24-61%)	2.5 (1.3-4.1)	6.3% (3.5-9.4%)	8.3% (4.5-12%)	31 (17-47)	1100 (630-1700)
Jamaica	2005	Male	60 to 64	55% (35-74%)	45% (26-65%)	2.2 (1.1-3.9)	5.7% (2.9-8.8%)	7.1% (3.6-11%)	36 (18-58)	1100 (550-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	2005	Male	65 to 69	54% (34-72%)	46% (28-66%)	1.9 (1-3.3)	5.1% (2.6-8.4%)	6.4% (3.2-10%)	39 (20-64)	970 (490-1600)
Jamaica	2005	Male	70 to 74	50% (31-67%)	50% (33-69%)	1.6 (0.77-2.9)	4% (1.7-7.1%)	4.8% (2-8.7%)	40 (17-73)	790 (330-1400)
Jamaica	2005	Male	75 to 79	44% (27-64%)	56% (36-73%)	1.3 (0.62-2.3)	3.1% (1.2-5.7%)	3.7% (1.4-6.8%)	36 (14-67)	560 (220-1100)
Jamaica	2005	Male	80 to 84	39% (22-57%)	61% (43-78%)	1.1 (0.52-1.9)	2.5% (1-4.7%)	2.9% (1.2-5.4%)	33 (13-61)	370 (150-700)
Jamaica	2005	Male	85 to 89	34% (19-51%)	66% (49-81%)	0.94 (0.46-1.7)	2.1% (0.81-4.2%)	2.4% (0.86-4.8%)	24 (8.7-48)	190 (74-370)
Jamaica	2005	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.95 (0.44-1.8)	1.9% (0.61-4.1%)	2.1% (0.59-4.7%)	14 (3.8-31)	77 (24-160)
Jamaica	2005	Male	95 plus	38% (20-58%)	62% (42-80%)	1 (0.43-2)	2% (0.45-4.6%)	2.1% (0.3-5.2%)	5.7 (0.8-14)	22 (4.7-50)
Jamaica	2010	Female	All Ages	27% (22-33%)	73% (67-78%)	0.93 (0.75-1.2)	1.7% (1-2.6%)	2% (0.8-3.8%)	200 (77-360)	6300 (3800-9600)
Jamaica	2010	Female	15 to 19	32% (20-48%)	68% (52-80%)	1.1 (0.55-1.9)	1.7% (1-2.7%)	3.5% (1.7-6.1%)	2.1 (1-3.7)	270 (160-420)
Jamaica	2010	Female	20 to 24	38% (23-55%)	62% (45-77%)	1.6 (0.86-2.6)	2.9% (1.9-4.1%)	4.6% (2.4-7.4%)	3.5 (1.8-5.7)	480 (300-690)
Jamaica	2010	Female	25 to 29	38% (22-54%)	62% (46-78%)	1.7 (0.95-2.6)	3% (2-4.2%)	4.5% (2.5-6.9%)	4.1 (2.3-6.5)	490 (310-720)
Jamaica	2010	Female	30 to 34	36% (21-53%)	64% (47-79%)	1.6 (0.89-2.6)	2.8% (1.9-4%)	4.5% (2.5-7.1%)	4.6 (2.5-7.4)	450 (290-660)
Jamaica	2010	Female	35 to 39	35% (22-52%)	65% (48-78%)	1.5 (0.81-2.4)	2.8% (1.8-4.1%)	4.6% (2.8-7.3%)	6.5 (3.7-10)	500 (320-740)
Jamaica	2010	Female	40 to 44	34% (21-52%)	66% (48-79%)	1.4 (0.75-2.3)	2.8% (1.6-4.3%)	4.9% (2.8-7.6%)	9.5 (5.3-15)	580 (330-910)
Jamaica	2010	Female	45 to 49	32% (18-49%)	68% (51-82%)	1.4 (0.67-2.3)	2.9% (1.5-4.9%)	4.9% (2.6-8.1%)	12 (6.4-21)	610 (320-1100)
Jamaica	2010	Female	50 to 54	29% (17-45%)	71% (55-83%)	1.2 (0.6-2.1)	2.9% (1.4-5.1%)	4.6% (2.1-7.8%)	16 (7.1-27)	650 (290-1200)
Jamaica	2010	Female	55 to 59	25% (14-41%)	75% (59-86%)	1 (0.47-1.8)	2.6% (0.95-4.8%)	3.8% (1.5-7%)	14 (5.7-26)	500 (190-940)
Jamaica	2010	Female	60 to 64	22% (12-37%)	78% (63-88%)	0.86 (0.4-1.6)	2.1% (0.47-4.5%)	3% (0.79-6.1%)	14 (3.7-29)	420 (93-860)
Jamaica	2010	Female	65 to 69	20% (12-32%)	80% (68-88%)	0.74 (0.38-1.3)	1.7% (0.3-4%)	2.4% (0.5-5.1%)	14 (3.2-30)	340 (62-750)
Jamaica	2010	Female	70 to 74	18% (11-29%)	82% (71-89%)	0.63 (0.29-1.2)	1.4% (0.015-3.7%)	1.9% (0.12-4.8%)	15 (0.95-39)	290 (2.8-760)
Jamaica	2010	Female	75 to 79	16% (9.3-27%)	84% (73-91%)	0.54 (0.23-1)	1.2% (-0.14-3.3%)	1.5% (-0.12-4.2%)	16 (-1.4-44)	240 (-30-670)
Jamaica	2010	Female	80 to 84	15% (7.9-24%)	85% (76-92%)	0.51 (0.23-0.92)	1.1% (-0.033-2.9%)	1.4% (-0.026-3.6%)	21 (-0.42-54)	230 (-6.9-580)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	2010	Female	85 to 89	15% (8.3-23%)	85% (77-92%)	0.51 (0.24-0.96)	1.2% (0.005-3%)	1.4% (-0.0075-3.6%)	20 (-0.11-52)	160 (0.63-410)
Jamaica	2010	Female	90 to 94	16% (9.3-26%)	84% (74-91%)	0.58 (0.25-1.1)	1.4% (0.018-3.6%)	1.6% (-0.043-4.4%)	15 (-0.42-41)	80 (1.1-210)
Jamaica	2010	Female	95 plus	20% (9.8-34%)	80% (66-90%)	0.7 (0.24-1.5)	1.6% (-0.13-4.6%)	1.8% (-0.28-5.4%)	7.9 (-1.2-24)	29 (-2.3-82)
Jamaica	2010	Male	All Ages	49% (40-58%)	51% (42-60%)	2.1 (1.7-2.6)	5.6% (4.2-7.3%)	5.9% (3.9-8%)	600 (390-850)	22000 (17000-29000)
Jamaica	2010	Male	15 to 19	49% (32-67%)	51% (33-68%)	1.6 (0.86-2.7)	5.1% (3.1-7.7%)	7.8% (4-13%)	9.8 (4.9-16)	960 (590-1400)
Jamaica	2010	Male	20 to 24	60% (42-77%)	40% (23-58%)	3.2 (1.8-4.9)	11% (7.2-15%)	14% (8.2-21%)	26 (15-41)	2300 (1500-3300)
Jamaica	2010	Male	25 to 29	64% (44-79%)	36% (21-56%)	3.9 (2.3-5.8)	11% (7.6-16%)	16% (9.4-23%)	28 (17-43)	2300 (1500-3200)
Jamaica	2010	Male	30 to 34	65% (45-80%)	35% (20-55%)	4.1 (2.4-6.2)	11% (7.2-15%)	15% (9.3-22%)	26 (16-39)	1900 (1200-2600)
Jamaica	2010	Male	35 to 39	65% (48-80%)	35% (20-52%)	3.8 (2.2-6)	9.7% (6.6-14%)	14% (8.6-20%)	30 (18-45)	1900 (1200-2700)
Jamaica	2010	Male	40 to 44	64% (47-79%)	36% (21-53%)	3.6 (2-5.6)	9.2% (6-13%)	13% (8-19%)	36 (21-54)	2000 (1300-2900)
Jamaica	2010	Male	45 to 49	62% (41-78%)	38% (22-59%)	3.3 (1.8-5.4)	8.6% (5.5-12%)	12% (7.3-17%)	42 (24-61)	2000 (1300-2800)
Jamaica	2010	Male	50 to 54	59% (40-77%)	41% (23-60%)	3 (1.6-5)	7.8% (4.7-11%)	10% (5.9-15%)	43 (25-63)	1800 (1100-2600)
Jamaica	2010	Male	55 to 59	57% (39-75%)	43% (25-61%)	2.8 (1.4-4.7)	7.5% (4.2-11%)	9.7% (5.4-14%)	50 (27-72)	1800 (1000-2600)
Jamaica	2010	Male	60 to 64	55% (35-74%)	45% (26-65%)	2.5 (1.3-4.4)	6.6% (3.4-9.8%)	8.2% (4.2-12%)	52 (26-80)	1600 (810-2400)
Jamaica	2010	Male	65 to 69	54% (35-72%)	46% (28-65%)	2.2 (1.1-3.7)	6% (3-9.4%)	7.3% (3.7-12%)	54 (28-87)	1300 (680-2100)
Jamaica	2010	Male	70 to 74	50% (31-67%)	50% (33-69%)	1.8 (0.83-3.2)	4.6% (1.9-8%)	5.5% (2.3-9.4%)	51 (22-91)	1000 (440-1800)
Jamaica	2010	Male	75 to 79	45% (27-64%)	55% (36-73%)	1.5 (0.67-2.6)	3.7% (1.4-6.7%)	4.3% (1.6-7.9%)	46 (18-85)	710 (280-1300)
Jamaica	2010	Male	80 to 84	39% (23-57%)	61% (43-77%)	1.2 (0.59-2.2)	3.1% (1.3-5.6%)	3.5% (1.4-6.4%)	46 (19-85)	520 (210-950)
Jamaica	2010	Male	85 to 89	35% (20-51%)	65% (49-80%)	1.1 (0.51-1.9)	2.6% (0.98-4.9%)	2.9% (1.1-5.6%)	37 (14-72)	300 (110-560)
Jamaica	2010	Male	90 to 94	35% (21-51%)	65% (49-79%)	1.1 (0.48-2)	2.4% (0.8-4.8%)	2.6% (0.82-5.5%)	20 (6.2-42)	110 (36-220)
Jamaica	2010	Male	95 plus	37% (20-58%)	63% (42-80%)	1.2 (0.46-2.3)	2.4% (0.53-5.4%)	2.6% (0.42-6%)	7.5 (1.2-18)	28 (6.2-62)
Jamaica	2016	Female	All Ages	27% (22-33%)	73% (67-78%)	1 (0.75-1.4)	1.9% (1.1-3.1%)	2.3% (0.79-4.4%)	240 (80-470)	7300 (4000-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	2016	Female	15 to 19	33% (20-49%)	67% (51-80%)	1.2 (0.58-2.1)	1.8% (1-2.9%)	3.8% (1.8-7%)	2.2 (0.95-4)	270 (150-440)
Jamaica	2016	Female	20 to 24	38% (24-55%)	62% (45-76%)	1.8 (0.87-3)	3.1% (1.9-4.4%)	5.1% (2.5-8.3%)	3.8 (1.7-6.5)	520 (320-770)
Jamaica	2016	Female	25 to 29	38% (23-55%)	62% (45-77%)	1.8 (0.96-3)	3.1% (2.1-4.5%)	4.9% (2.6-7.7%)	4.3 (2.1-7.4)	530 (330-790)
Jamaica	2016	Female	30 to 34	36% (22-53%)	64% (47-78%)	1.7 (0.95-2.9)	3% (1.9-4.3%)	4.9% (2.8-7.8%)	5.1 (2.5-9)	520 (320-800)
Jamaica	2016	Female	35 to 39	35% (21-52%)	65% (48-79%)	1.6 (0.81-2.7)	2.9% (1.8-4.4%)	5% (2.9-8%)	6.4 (3.3-12)	510 (300-800)
Jamaica	2016	Female	40 to 44	34% (21-51%)	66% (49-79%)	1.5 (0.74-2.5)	3% (1.6-4.9%)	5.2% (2.8-8.6%)	9.1 (4.4-17)	570 (300-980)
Jamaica	2016	Female	45 to 49	32% (18-48%)	68% (52-82%)	1.4 (0.68-2.5)	3.1% (1.6-5.5%)	5.2% (2.7-8.8%)	13 (6.1-24)	660 (320-1200)
Jamaica	2016	Female	50 to 54	29% (17-45%)	71% (55-83%)	1.3 (0.59-2.3)	3.1% (1.4-5.6%)	4.9% (2.2-8.5%)	17 (7.4-32)	740 (320-1400)
Jamaica	2016	Female	55 to 59	25% (15-40%)	75% (60-85%)	1.1 (0.5-1.9)	2.8% (0.99-5.5%)	4.1% (1.7-7.7%)	20 (7.8-39)	720 (250-1500)
Jamaica	2016	Female	60 to 64	22% (13-36%)	78% (64-87%)	0.92 (0.42-1.7)	2.3% (0.46-4.9%)	3.2% (0.87-6.6%)	19 (4.9-40)	580 (130-1300)
Jamaica	2016	Female	65 to 69	20% (12-32%)	80% (68-88%)	0.8 (0.38-1.4)	1.9% (0.28-4.4%)	2.7% (0.61-5.6%)	19 (3.9-39)	450 (77-1000)
Jamaica	2016	Female	70 to 74	18% (11-30%)	82% (70-89%)	0.67 (0.3-1.3)	1.6% (0.03-4.2%)	2.1% (0.13-5.3%)	18 (1.1-46)	350 (6.1-910)
Jamaica	2016	Female	75 to 79	16% (9.2-27%)	84% (73-91%)	0.58 (0.24-1.1)	1.3% (-0.16-3.6%)	1.7% (-0.18-4.7%)	19 (-2.2-49)	280 (-39-760)
Jamaica	2016	Female	80 to 84	15% (8.3-24%)	85% (76-92%)	0.54 (0.24-1)	1.3% (-0.021-3.5%)	1.6% (-0.041-4.3%)	26 (-0.71-69)	280 (-4.9-760)
Jamaica	2016	Female	85 to 89	15% (8.1-23%)	85% (77-92%)	0.55 (0.23-1.1)	1.3% (0.067-3.4%)	1.6% (0.055-4.1%)	27 (0.97-70)	210 (11-540)
Jamaica	2016	Female	90 to 94	16% (9.3-26%)	84% (74-91%)	0.62 (0.26-1.3)	1.5% (0.072-4%)	1.7% (-0.0062-4.7%)	20 (-0.074-54)	110 (4.8-280)
Jamaica	2016	Female	95 plus	20% (9.6-34%)	80% (66-90%)	0.76 (0.27-1.6)	1.8% (-0.044-5%)	2% (-0.2-5.8%)	9.5 (-0.95-27)	34 (-0.89-94)
Jamaica	2016	Male	All Ages	50% (42-58%)	50% (42-58%)	2.4 (1.7-3.3)	6.1% (4.4-8.1%)	6.3% (3.9-8.7%)	720 (420-1100)	26000 (18000-36000)
Jamaica	2016	Male	15 to 19	50% (32-68%)	50% (32-68%)	1.8 (0.87-3)	5.4% (3.2-8.4%)	8.5% (4.1-14%)	10 (4.7-18)	990 (560-1600)
Jamaica	2016	Male	20 to 24	61% (42-77%)	39% (23-58%)	3.4 (1.8-5.6)	11% (7.7-17%)	15% (8.8-24%)	28 (15-48)	2500 (1600-3800)
Jamaica	2016	Male	25 to 29	64% (45-79%)	36% (21-55%)	4.2 (2.4-6.9)	12% (8.2-17%)	17% (10-25%)	33 (19-53)	2700 (1700-3900)
Jamaica	2016	Male	30 to 34	65% (45-81%)	35% (19-55%)	4.4 (2.5-7.1)	11% (7.7-15%)	16% (9.9-23%)	31 (18-48)	2200 (1500-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jamaica	2016	Male	35 to 39	65% (48-80%)	35% (20-52%)	4.1 (2.3-6.7)	10% (7.1-14%)	15% (9.3-21%)	29 (17-45)	1900 (1200-2800)
Jamaica	2016	Male	40 to 44	64% (47-79%)	36% (21-53%)	3.9 (2-6.3)	9.7% (6.4-13%)	14% (8.5-19%)	35 (20-53)	2000 (1200-2900)
Jamaica	2016	Male	45 to 49	62% (40-78%)	38% (22-60%)	3.6 (1.8-6.2)	9.1% (5.8-13%)	12% (7.8-17%)	46 (26-71)	2300 (1400-3300)
Jamaica	2016	Male	50 to 54	60% (41-77%)	40% (23-59%)	3.2 (1.6-5.5)	8.2% (5-12%)	11% (6.4-15%)	55 (29-85)	2300 (1300-3500)
Jamaica	2016	Male	55 to 59	57% (39-75%)	43% (25-61%)	2.9 (1.4-5.1)	7.7% (4.2-11%)	10% (5.4-14%)	61 (30-95)	2200 (1100-3300)
Jamaica	2016	Male	60 to 64	55% (35-74%)	45% (26-65%)	2.6 (1.3-4.8)	6.8% (3.5-10%)	8.4% (4.4-13%)	68 (32-110)	2100 (1000-3300)
Jamaica	2016	Male	65 to 69	54% (34-72%)	46% (28-66%)	2.3 (1.1-4)	6.2% (3.2-9.8%)	7.6% (3.8-12%)	72 (34-120)	1800 (870-2900)
Jamaica	2016	Male	70 to 74	50% (32-67%)	50% (33-68%)	1.9 (0.88-3.5)	4.9% (2-8.3%)	5.8% (2.4-10%)	63 (25-110)	1200 (510-2200)
Jamaica	2016	Male	75 to 79	45% (27-64%)	55% (36-73%)	1.6 (0.68-2.9)	3.9% (1.5-7%)	4.6% (1.7-8.3%)	56 (19-100)	850 (310-1500)
Jamaica	2016	Male	80 to 84	39% (23-57%)	61% (43-77%)	1.3 (0.6-2.3)	3.3% (1.2-5.9%)	3.8% (1.4-6.7%)	55 (20-100)	620 (230-1100)
Jamaica	2016	Male	85 to 89	35% (20-51%)	65% (49-80%)	1.1 (0.51-2.1)	2.8% (1-5.4%)	3.2% (1.1-6.1%)	43 (14-84)	340 (120-650)
Jamaica	2016	Male	90 to 94	35% (21-51%)	65% (49-79%)	1.1 (0.49-2.2)	2.6% (0.81-5.1%)	2.8% (0.83-5.8%)	26 (7.5-54)	140 (44-280)
Jamaica	2016	Male	95 plus	37% (20-58%)	63% (42-80%)	1.2 (0.46-2.5)	2.6% (0.63-5.8%)	2.8% (0.5-6.6%)	9.2 (1.6-21)	35 (8.7-77)
Saint Lucia	1990	Female	All Ages	34% (28-41%)	66% (59-72%)	1.3 (1.1-1.5)	3.4% (2.5-4.5%)	4.6% (2.8-6.8%)	19 (12-28)	650 (490-850)
Saint Lucia	1990	Female	15 to 19	43% (29-59%)	57% (41-71%)	1.7 (0.95-2.7)	3% (1.9-4.4%)	7% (4.2-10%)	0.3 (0.18-0.46)	30 (20-44)
Saint Lucia	1990	Female	20 to 24	49% (33-65%)	51% (35-67%)	2.5 (1.5-3.8)	4.9% (3.5-6.6%)	9.6% (6.5-13%)	0.45 (0.3-0.62)	46 (33-62)
Saint Lucia	1990	Female	25 to 29	48% (32-65%)	52% (35-68%)	2.6 (1.6-3.9)	5.8% (4.2-7.7%)	9.9% (7-13%)	0.69 (0.48-0.93)	59 (43-78)
Saint Lucia	1990	Female	30 to 34	46% (30-64%)	54% (36-70%)	2.5 (1.5-3.9)	5.6% (4-7.3%)	9.7% (7-13%)	0.62 (0.44-0.83)	48 (35-63)
Saint Lucia	1990	Female	35 to 39	45% (30-63%)	55% (37-70%)	2.4 (1.4-3.8)	6.1% (4.4-8.3%)	10% (7.4-14%)	0.75 (0.54-1)	47 (34-64)
Saint Lucia	1990	Female	40 to 44	44% (30-63%)	56% (37-70%)	2.3 (1.2-3.7)	6.6% (4.4-9.2%)	11% (7.1-15%)	1 (0.7-1.4)	55 (37-77)
Saint Lucia	1990	Female	45 to 49	42% (27-59%)	58% (41-73%)	2.2 (1.2-3.7)	6.5% (4.1-9.9%)	10% (6.4-15%)	1.1 (0.72-1.7)	52 (33-78)
Saint Lucia	1990	Female	50 to 54	39% (24-56%)	61% (44-76%)	2 (1-3.3)	7% (4.2-11%)	9.8% (5.9-15%)	1.7 (1-2.6)	67 (40-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	1990	Female	55 to 59	34% (21-51%)	66% (49-79%)	1.7 (0.86-2.8)	5.6% (3-9%)	8.2% (4.6-13%)	1.5 (0.81-2.3)	51 (28-81)
Saint Lucia	1990	Female	60 to 64	31% (18-47%)	69% (53-82%)	1.4 (0.71-2.5)	4.9% (2.2-8.5%)	6.3% (2.8-11%)	1.9 (0.86-3.3)	55 (24-95)
Saint Lucia	1990	Female	65 to 69	28% (18-42%)	72% (58-82%)	1.2 (0.67-2)	3.9% (1.7-6.9%)	5.2% (2.3-8.8%)	1.9 (0.86-3.2)	45 (19-77)
Saint Lucia	1990	Female	70 to 74	26% (16-39%)	74% (61-84%)	1 (0.53-1.9)	3.1% (0.89-6.3%)	4% (1.3-7.9%)	2.1 (0.65-4.1)	40 (12-80)
Saint Lucia	1990	Female	75 to 79	23% (13-37%)	77% (63-87%)	0.9 (0.41-1.6)	2.4% (0.52-5.1%)	3.2% (0.69-6.6%)	1.6 (0.34-3.2)	23 (5-49)
Saint Lucia	1990	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.84 (0.4-1.4)	2.4% (0.68-4.9%)	2.9% (0.81-5.9%)	1.7 (0.47-3.4)	18 (5-36)
Saint Lucia	1990	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.86 (0.42-1.6)	2.4% (0.67-5%)	2.8% (0.74-5.9%)	1 (0.27-2.1)	7.6 (2.1-16)
Saint Lucia	1990	Female	90 to 94	23% (14-36%)	77% (64-86%)	0.99 (0.44-1.9)	2.6% (0.62-5.9%)	3.1% (0.62-6.9%)	0.56 (0.11-1.3)	2.9 (0.67-6.5)
Saint Lucia	1990	Female	95 plus	28% (15-45%)	72% (55-85%)	1.2 (0.45-2.3)	3.1% (0.36-7.3%)	3.5% (0.25-8.5%)	0.21 (0.016-0.51)	0.74 (0.086-1.7)
Saint Lucia	1990	Male	All Ages	55% (47-63%)	45% (37-53%)	2.7 (2.4-3)	8.8% (7.1-10%)	11% (8.3-14%)	53 (40-66)	1900 (1600-2300)
Saint Lucia	1990	Male	15 to 19	60% (42-75%)	40% (25-58%)	2.5 (1.4-3.8)	6.6% (4.3-9.4%)	11% (6.6-16%)	0.8 (0.49-1.2)	73 (48-100)
Saint Lucia	1990	Male	20 to 24	69% (52-83%)	31% (17-48%)	4.6 (3.1-6.4)	14% (11-19%)	19% (14-26%)	2.4 (1.7-3.3)	190 (140-250)
Saint Lucia	1990	Male	25 to 29	72% (56-84%)	28% (16-44%)	5.5 (3.7-7.5)	15% (12-19%)	22% (16-28%)	2.4 (1.7-3.1)	180 (140-230)
Saint Lucia	1990	Male	30 to 34	73% (56-85%)	27% (15-44%)	5.8 (3.8-8)	16% (12-20%)	22% (17-27%)	2.7 (2.1-3.4)	180 (140-220)
Saint Lucia	1990	Male	35 to 39	73% (57-85%)	27% (15-43%)	5.5 (3.5-8.2)	16% (12-20%)	21% (16-27%)	3 (2.2-3.7)	170 (130-210)
Saint Lucia	1990	Male	40 to 44	72% (57-85%)	28% (15-43%)	5.3 (3.1-7.8)	16% (12-20%)	21% (16-26%)	3.1 (2.3-3.8)	160 (120-200)
Saint Lucia	1990	Male	45 to 49	70% (52-84%)	30% (16-48%)	5 (2.9-8)	16% (12-20%)	20% (15-25%)	3.4 (2.5-4.3)	150 (110-190)
Saint Lucia	1990	Male	50 to 54	69% (51-83%)	31% (17-49%)	4.6 (2.6-7.1)	14% (10-18%)	19% (13-23%)	3.5 (2.4-4.4)	140 (99-180)
Saint Lucia	1990	Male	55 to 59	67% (49-82%)	33% (18-51%)	4.2 (2.3-6.6)	13% (9.3-17%)	17% (12-22%)	4.3 (3-5.6)	150 (100-190)
Saint Lucia	1990	Male	60 to 64	65% (46-82%)	35% (18-54%)	3.8 (2.1-6.4)	12% (7.4-16%)	14% (8.7-19%)	5.6 (3.5-7.7)	160 (100-220)
Saint Lucia	1990	Male	65 to 69	64% (45-79%)	36% (21-55%)	3.4 (2-5.4)	11% (6.5-15%)	13% (7.9-18%)	5.5 (3.4-7.6)	130 (82-180)
Saint Lucia	1990	Male	70 to 74	60% (42-76%)	40% (24-58%)	2.9 (1.6-4.8)	8.4% (4.5-13%)	10% (5.3-15%)	6.4 (3.5-9.9)	120 (66-190)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	1990	Male	75 to 79	56% (36-74%)	44% (26-64%)	2.4 (1.2-4.1)	6.5% (3.1-10%)	7.7% (3.6-12%)	4.3 (2-6.9)	64 (31-100)
Saint Lucia	1990	Male	80 to 84	50% (32-68%)	50% (32-68%)	2.1 (1.1-3.4)	5.6% (2.7-9.2%)	6.4% (2.9-10%)	3.3 (1.5-5.4)	36 (17-58)
Saint Lucia	1990	Male	85 to 89	45% (29-62%)	55% (38-71%)	1.8 (0.93-3.1)	4.8% (2.1-8.2%)	5.4% (2.2-9.5%)	1.4 (0.59-2.6)	11 (4.9-19)
Saint Lucia	1990	Male	90 to 94	46% (30-63%)	54% (37-70%)	1.8 (0.89-3.2)	4.3% (1.6-7.9%)	4.8% (1.7-9%)	0.51 (0.18-0.97)	2.7 (1-5.1)
Saint Lucia	1990	Male	95 plus	49% (29-69%)	51% (31-71%)	1.9 (0.79-3.8)	4.4% (1.1-8.9%)	4.8% (0.98-10%)	0.13 (0.027-0.27)	0.47 (0.12-0.95)
Saint Lucia	1995	Female	All Ages	35% (29-42%)	65% (58-71%)	1.5 (1.3-1.8)	3.8% (2.8-4.9%)	5.1% (3.3-7.3%)	21 (14-31)	700 (520-930)
Saint Lucia	1995	Female	15 to 19	42% (29-59%)	58% (41-71%)	1.9 (1-3)	3.1% (2-4.6%)	7.4% (4.5-11%)	0.26 (0.16-0.4)	27 (17-38)
Saint Lucia	1995	Female	20 to 24	48% (32-64%)	52% (36-68%)	2.8 (1.7-4.4)	4.9% (3.5-6.6%)	10% (6.5-14%)	0.41 (0.27-0.56)	44 (31-60)
Saint Lucia	1995	Female	25 to 29	48% (31-65%)	52% (35-69%)	2.9 (1.8-4.2)	5.9% (4.3-7.7%)	10% (7.2-13%)	0.68 (0.47-0.91)	59 (43-77)
Saint Lucia	1995	Female	30 to 34	46% (29-64%)	54% (36-71%)	2.8 (1.7-4.2)	5.4% (3.9-7.1%)	9.8% (7-13%)	0.67 (0.48-0.87)	54 (39-70)
Saint Lucia	1995	Female	35 to 39	45% (30-63%)	55% (37-70%)	2.7 (1.6-4.1)	6.1% (4.4-8.3%)	11% (7.7-14%)	0.9 (0.64-1.2)	58 (42-79)
Saint Lucia	1995	Female	40 to 44	44% (29-62%)	56% (38-71%)	2.5 (1.4-4)	6.8% (4.6-9.4%)	11% (7.5-15%)	1.1 (0.76-1.5)	60 (41-84)
Saint Lucia	1995	Female	45 to 49	42% (27-59%)	58% (41-73%)	2.4 (1.3-4)	6.8% (4.3-10%)	10% (6.6-15%)	1.3 (0.82-1.8)	60 (38-88)
Saint Lucia	1995	Female	50 to 54	38% (24-55%)	62% (45-76%)	2.2 (1.2-3.7)	7.1% (4.3-11%)	10% (6.4-15%)	1.5 (0.96-2.3)	62 (38-96)
Saint Lucia	1995	Female	55 to 59	34% (21-51%)	66% (49-79%)	1.9 (0.95-3.1)	6.1% (3.3-9.9%)	8.7% (4.9-14%)	1.6 (0.89-2.5)	56 (31-91)
Saint Lucia	1995	Female	60 to 64	30% (18-47%)	70% (53-82%)	1.6 (0.79-2.8)	5.4% (2.4-9.4%)	7% (3.2-12%)	2.3 (1-3.9)	65 (29-110)
Saint Lucia	1995	Female	65 to 69	28% (18-42%)	72% (58-82%)	1.4 (0.75-2.3)	4.4% (1.9-7.7%)	5.7% (2.6-9.8%)	1.9 (0.86-3.2)	45 (20-78)
Saint Lucia	1995	Female	70 to 74	25% (16-39%)	75% (61-84%)	1.2 (0.59-2.1)	3.6% (1.2-7.1%)	4.5% (1.6-8.7%)	2.4 (0.84-4.6)	45 (15-89)
Saint Lucia	1995	Female	75 to 79	22% (13-36%)	78% (64-87%)	1 (0.45-1.8)	2.7% (0.77-5.5%)	3.6% (1.1-7.2%)	1.9 (0.56-3.8)	30 (8.4-58)
Saint Lucia	1995	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.95 (0.45-1.6)	2.6% (0.91-5.1%)	3.3% (1.2-6.5%)	2 (0.69-3.9)	22 (7.7-43)
Saint Lucia	1995	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.97 (0.49-1.7)	2.6% (0.92-5.1%)	3.2% (1-6.3%)	1.3 (0.43-2.6)	10 (3.6-20)
Saint Lucia	1995	Female	90 to 94	23% (14-35%)	77% (65-86%)	1.1 (0.5-2.1)	3.1% (0.96-6.7%)	3.6% (1-7.9%)	0.78 (0.22-1.7)	4.1 (1.2-8.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	1995	Female	95 plus	27% (15-44%)	73% (56-85%)	1.3 (0.49-2.6)	3.6% (0.63-8.1%)	4.2% (0.54-9.5%)	0.3 (0.038-0.67)	1 (0.18-2.3)
Saint Lucia	1995	Male	All Ages	56% (48-64%)	44% (36-52%)	3.2 (2.8-3.6)	10% (8.3-12%)	12% (9.4-14%)	59 (47-72)	2200 (1900-2500)
Saint Lucia	1995	Male	15 to 19	59% (42-75%)	41% (25-58%)	2.8 (1.5-4.3)	7.5% (4.8-11%)	12% (7.2-18%)	0.85 (0.51-1.3)	75 (49-110)
Saint Lucia	1995	Male	20 to 24	69% (52-83%)	31% (17-48%)	5.1 (3.3-7.1)	16% (12-20%)	21% (15-28%)	2.6 (1.8-3.5)	210 (150-270)
Saint Lucia	1995	Male	25 to 29	72% (55-84%)	28% (16-45%)	6.1 (4.1-8.4)	17% (13-22%)	23% (17-29%)	3.3 (2.4-4.2)	230 (180-300)
Saint Lucia	1995	Male	30 to 34	73% (55-85%)	27% (15-45%)	6.4 (4.2-8.7)	17% (13-21%)	22% (17-28%)	3.7 (2.8-4.5)	240 (190-290)
Saint Lucia	1995	Male	35 to 39	72% (57-85%)	28% (15-43%)	6.1 (3.9-8.8)	16% (13-20%)	22% (17-27%)	3.5 (2.7-4.3)	210 (160-250)
Saint Lucia	1995	Male	40 to 44	72% (56-85%)	28% (15-44%)	5.8 (3.4-8.7)	16% (13-20%)	21% (17-26%)	3.5 (2.7-4.3)	180 (140-220)
Saint Lucia	1995	Male	45 to 49	70% (51-84%)	30% (16-49%)	5.5 (3.2-9)	16% (12-20%)	21% (16-25%)	3.8 (2.8-4.6)	170 (130-210)
Saint Lucia	1995	Male	50 to 54	69% (51-83%)	31% (17-49%)	5.1 (2.9-8.2)	15% (11-19%)	19% (14-24%)	4.2 (3-5.3)	170 (120-210)
Saint Lucia	1995	Male	55 to 59	67% (50-81%)	33% (19-50%)	4.7 (2.6-7.5)	14% (10-17%)	18% (13-22%)	4.3 (3.1-5.4)	150 (110-190)
Saint Lucia	1995	Male	60 to 64	65% (46-81%)	35% (19-54%)	4.3 (2.3-7.1)	12% (7.8-16%)	15% (9.8-20%)	4.9 (3.2-6.5)	140 (95-190)
Saint Lucia	1995	Male	65 to 69	63% (45-79%)	37% (21-55%)	3.8 (2.2-5.9)	11% (7.3-15%)	14% (8.8-18%)	5.7 (3.6-7.6)	140 (88-180)
Saint Lucia	1995	Male	70 to 74	60% (42-76%)	40% (24-58%)	3.2 (1.7-5.3)	9.1% (5.1-13%)	11% (6.1-16%)	6.6 (3.8-9.6)	130 (73-180)
Saint Lucia	1995	Male	75 to 79	55% (36-74%)	45% (26-64%)	2.7 (1.4-4.6)	7% (3.7-11%)	8.3% (4.4-12%)	5.3 (2.8-7.9)	80 (42-120)
Saint Lucia	1995	Male	80 to 84	49% (31-67%)	51% (33-69%)	2.3 (1.2-3.7)	6.1% (3.1-9.2%)	7% (3.5-11%)	4.4 (2.1-6.7)	48 (24-73)
Saint Lucia	1995	Male	85 to 89	45% (28-62%)	55% (38-72%)	2 (1-3.4)	5.1% (2.6-8.2%)	5.8% (2.8-9.6%)	1.9 (0.94-3.1)	15 (7.5-24)
Saint Lucia	1995	Male	90 to 94	45% (29-62%)	55% (38-71%)	2 (0.98-3.5)	4.7% (2.1-8.1%)	5.3% (2.3-9.3%)	0.66 (0.28-1.2)	3.6 (1.6-6.2)
Saint Lucia	1995	Male	95 plus	48% (29-68%)	52% (32-71%)	2.1 (0.88-4.1)	4.9% (1.6-9.2%)	5.4% (1.5-10%)	0.17 (0.046-0.33)	0.6 (0.19-1.1)
Saint Lucia	2000	Female	All Ages	35% (29-42%)	65% (58-71%)	1.6 (1.4-1.8)	3.6% (2.7-4.7%)	4.7% (3-6.9%)	20 (13-30)	660 (490-880)
Saint Lucia	2000	Female	15 to 19	41% (28-58%)	59% (42-72%)	1.9 (1-3)	2.6% (1.6-3.8%)	6.8% (4.1-10%)	0.2 (0.11-0.3)	23 (15-34)
Saint Lucia	2000	Female	20 to 24	47% (31-64%)	53% (36-69%)	2.8 (1.6-4.3)	4.6% (3.3-6.3%)	8.8% (5.7-12%)	0.4 (0.25-0.56)	43 (30-58)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	2000	Female	25 to 29	47% (30-64%)	53% (36-70%)	2.9 (1.7-4.2)	4.7% (3.4-6.1%)	8.6% (5.9-12%)	0.43 (0.29-0.59)	43 (31-58)
Saint Lucia	2000	Female	30 to 34	45% (29-63%)	55% (37-71%)	2.8 (1.7-4.2)	4.5% (3.3-5.8%)	7.8% (5.6-10%)	0.51 (0.36-0.67)	45 (32-59)
Saint Lucia	2000	Female	35 to 39	44% (29-62%)	56% (38-71%)	2.6 (1.5-4.1)	5% (3.6-6.8%)	9.2% (6.6-12%)	0.79 (0.56-1.1)	55 (39-74)
Saint Lucia	2000	Female	40 to 44	43% (28-61%)	57% (39-72%)	2.5 (1.3-3.9)	6.1% (4-8.4%)	9.9% (6.8-14%)	1.3 (0.91-1.8)	71 (48-99)
Saint Lucia	2000	Female	45 to 49	41% (26-58%)	59% (42-74%)	2.4 (1.3-3.9)	6.2% (4-9.3%)	9.5% (6.2-14%)	1.3 (0.84-1.9)	61 (40-91)
Saint Lucia	2000	Female	50 to 54	37% (23-55%)	63% (45-77%)	2.1 (1.2-3.6)	6.4% (3.9-10%)	9.5% (6-14%)	1.5 (0.97-2.3)	62 (38-95)
Saint Lucia	2000	Female	55 to 59	33% (20-51%)	67% (49-80%)	1.8 (0.94-3)	5.5% (2.9-9.2%)	8% (4.5-13%)	1.4 (0.78-2.2)	49 (26-82)
Saint Lucia	2000	Female	60 to 64	30% (18-46%)	70% (54-82%)	1.6 (0.77-2.8)	5% (2.2-9%)	6.5% (3-11%)	1.9 (0.88-3.3)	56 (25-100)
Saint Lucia	2000	Female	65 to 69	27% (17-41%)	73% (59-83%)	1.4 (0.72-2.3)	4% (1.7-7%)	5.4% (2.4-9.2%)	1.8 (0.79-3)	43 (19-77)
Saint Lucia	2000	Female	70 to 74	25% (16-38%)	75% (62-84%)	1.1 (0.57-2)	3.4% (1.1-6.9%)	4.3% (1.4-8.7%)	1.9 (0.61-3.7)	35 (11-74)
Saint Lucia	2000	Female	75 to 79	22% (13-36%)	78% (64-87%)	0.99 (0.46-1.8)	2.8% (0.69-5.7%)	3.6% (0.96-7.3%)	1.9 (0.5-3.8)	28 (7-59)
Saint Lucia	2000	Female	80 to 84	20% (12-33%)	80% (67-88%)	0.93 (0.46-1.6)	2.5% (0.83-4.9%)	3.2% (1.1-6.4%)	2 (0.67-3.8)	22 (7.6-43)
Saint Lucia	2000	Female	85 to 89	20% (12-32%)	80% (68-88%)	0.95 (0.48-1.7)	2.5% (0.92-4.8%)	3.1% (1.1-6.1%)	1.7 (0.59-3.3)	13 (4.9-26)
Saint Lucia	2000	Female	90 to 94	23% (13-35%)	77% (65-87%)	1.1 (0.48-2.1)	2.9% (0.95-6.2%)	3.5% (1-7.5%)	1 (0.3-2.2)	5.5 (1.7-12)
Saint Lucia	2000	Female	95 plus	27% (14-44%)	73% (56-86%)	1.3 (0.47-2.6)	3.4% (0.68-7.4%)	4% (0.68-8.9%)	0.32 (0.054-0.71)	1.1 (0.22-2.4)
Saint Lucia	2000	Male	All Ages	56% (48-65%)	44% (35-52%)	3.3 (3-3.7)	10% (8.4-12%)	12% (9.4-14%)	58 (47-70)	2200 (1900-2600)
Saint Lucia	2000	Male	15 to 19	58% (41-74%)	42% (26-59%)	2.7 (1.5-4.3)	7.8% (5.1-11%)	12% (7.1-18%)	1.1 (0.66-1.7)	95 (62-140)
Saint Lucia	2000	Male	20 to 24	68% (51-82%)	32% (18-49%)	5.1 (3.3-7.2)	16% (11-21%)	21% (14-28%)	2.7 (1.9-3.7)	210 (150-280)
Saint Lucia	2000	Male	25 to 29	71% (54-84%)	29% (16-46%)	6.1 (4-8.5)	17% (13-21%)	22% (16-29%)	3.1 (2.3-4.1)	230 (170-290)
Saint Lucia	2000	Male	30 to 34	72% (54-85%)	28% (15-46%)	6.4 (4.3-8.8)	16% (12-20%)	22% (16-27%)	3.2 (2.4-4)	220 (170-270)
Saint Lucia	2000	Male	35 to 39	72% (56-85%)	28% (15-44%)	6.1 (3.9-8.7)	15% (12-19%)	21% (16-26%)	3.5 (2.7-4.3)	210 (170-260)
Saint Lucia	2000	Male	40 to 44	71% (55-84%)	29% (16-45%)	5.8 (3.5-8.5)	15% (12-19%)	21% (16-25%)	3.8 (2.9-4.7)	200 (160-250)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	2000	Male	45 to 49	69% (50-83%)	31% (17-50%)	5.5 (3.1-8.7)	15% (12-19%)	20% (15-24%)	4.2 (3.1-5.1)	190 (150-240)
Saint Lucia	2000	Male	50 to 54	68% (49-83%)	32% (17-51%)	5.1 (2.9-8.1)	15% (11-18%)	18% (14-23%)	4.4 (3.3-5.5)	180 (130-220)
Saint Lucia	2000	Male	55 to 59	66% (48-81%)	34% (19-52%)	4.7 (2.6-7.4)	14% (10-18%)	17% (12-22%)	4.7 (3.4-6)	160 (120-200)
Saint Lucia	2000	Male	60 to 64	64% (45-81%)	36% (19-55%)	4.2 (2.2-7)	12% (7.6-15%)	14% (9.5-19%)	4.5 (3-5.9)	130 (89-170)
Saint Lucia	2000	Male	65 to 69	63% (44-79%)	37% (21-56%)	3.8 (2.1-6)	11% (7-14%)	13% (8.8-18%)	5.1 (3.4-6.7)	120 (82-170)
Saint Lucia	2000	Male	70 to 74	59% (41-75%)	41% (25-59%)	3.2 (1.7-5.3)	8.8% (5.1-13%)	10% (6-15%)	5.8 (3.4-8.3)	110 (65-160)
Saint Lucia	2000	Male	75 to 79	54% (35-72%)	46% (28-65%)	2.6 (1.4-4.5)	6.7% (3.6-10%)	8% (4.3-12%)	4.6 (2.5-7)	70 (38-110)
Saint Lucia	2000	Male	80 to 84	48% (30-66%)	52% (34-70%)	2.2 (1.1-3.6)	5.6% (3-8.4%)	6.6% (3.5-10%)	4 (2.1-6.1)	45 (24-68)
Saint Lucia	2000	Male	85 to 89	44% (27-61%)	56% (39-73%)	1.9 (0.97-3.2)	4.7% (2.4-7.4%)	5.5% (2.7-8.8%)	2.3 (1.1-3.7)	19 (9.7-30)
Saint Lucia	2000	Male	90 to 94	44% (28-61%)	56% (39-72%)	1.9 (0.95-3.4)	4.3% (2-7.2%)	4.9% (2.1-8.4%)	0.8 (0.35-1.4)	4.4 (2.1-7.5)
Saint Lucia	2000	Male	95 plus	47% (28-68%)	53% (32-72%)	2.1 (0.87-4.1)	4.4% (1.5-8.1%)	4.9% (1.4-9.4%)	0.17 (0.049-0.32)	0.63 (0.22-1.2)
Saint Lucia	2005	Female	All Ages	35% (29-43%)	65% (57-71%)	1.6 (1.4-1.8)	3.5% (2.6-4.5%)	4.4% (2.9-6.4%)	20 (13-29)	660 (500-870)
Saint Lucia	2005	Female	15 to 19	41% (28-57%)	59% (43-72%)	1.8 (0.98-2.8)	2.3% (1.5-3.4%)	6.8% (4-11%)	0.17 (0.1-0.26)	21 (13-31)
Saint Lucia	2005	Female	20 to 24	47% (31-64%)	53% (36-69%)	2.6 (1.5-4.1)	4.8% (3.3-6.6%)	9.1% (5.8-13%)	0.53 (0.33-0.75)	53 (36-72)
Saint Lucia	2005	Female	25 to 29	46% (30-64%)	54% (36-70%)	2.7 (1.6-4.1)	4.8% (3.4-6.4%)	9% (6.1-12%)	0.46 (0.31-0.63)	45 (32-60)
Saint Lucia	2005	Female	30 to 34	45% (28-62%)	55% (38-72%)	2.6 (1.6-3.9)	5% (3.6-6.6%)	8.8% (6.1-12%)	0.73 (0.51-0.99)	57 (41-75)
Saint Lucia	2005	Female	35 to 39	44% (29-61%)	56% (39-71%)	2.4 (1.4-3.8)	4.8% (3.4-6.5%)	9.3% (6.7-12%)	0.78 (0.56-1.1)	54 (38-73)
Saint Lucia	2005	Female	40 to 44	42% (27-61%)	58% (39-73%)	2.3 (1.3-3.6)	5.2% (3.6-7.4%)	9.5% (6.6-13%)	1.2 (0.82-1.6)	67 (46-94)
Saint Lucia	2005	Female	45 to 49	40% (25-58%)	60% (42-75%)	2.2 (1.2-3.7)	5.2% (3.3-8.1%)	9.1% (6-13%)	1.2 (0.78-1.8)	59 (38-88)
Saint Lucia	2005	Female	50 to 54	37% (23-54%)	63% (46-77%)	2 (1.1-3.3)	5.9% (3.6-9.4%)	9.1% (5.7-14%)	1.6 (0.98-2.4)	63 (39-97)
Saint Lucia	2005	Female	55 to 59	33% (20-49%)	67% (51-80%)	1.7 (0.88-2.8)	5.1% (2.8-8.3%)	7.6% (4.3-12%)	1.4 (0.82-2.3)	51 (28-83)
Saint Lucia	2005	Female	60 to 64	29% (17-45%)	71% (55-83%)	1.5 (0.73-2.5)	4.6% (2.1-8%)	6.2% (2.9-11%)	1.7 (0.82-2.9)	50 (23-87)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	2005	Female	65 to 69	27% (17-41%)	73% (59-83%)	1.3 (0.69-2.1)	3.8% (1.7-6.8%)	5% (2.3-8.7%)	1.8 (0.79-3.1)	42 (19-75)
Saint Lucia	2005	Female	70 to 74	24% (15-38%)	76% (62-85%)	1.1 (0.54-1.9)	3.1% (1.1-6.1%)	4% (1.5-7.9%)	1.8 (0.64-3.5)	34 (12-68)
Saint Lucia	2005	Female	75 to 79	22% (13-34%)	78% (66-87%)	0.92 (0.42-1.7)	2.6% (0.65-5.5%)	3.3% (0.88-7%)	1.4 (0.35-2.9)	21 (5.1-44)
Saint Lucia	2005	Female	80 to 84	20% (11-32%)	80% (68-89%)	0.86 (0.42-1.5)	2.3% (0.79-4.7%)	2.9% (0.98-5.8%)	1.9 (0.64-3.8)	21 (7.4-43)
Saint Lucia	2005	Female	85 to 89	20% (11-31%)	80% (69-89%)	0.88 (0.43-1.6)	2.2% (0.78-4.3%)	2.7% (0.93-5.4%)	1.7 (0.6-3.5)	14 (5.1-28)
Saint Lucia	2005	Female	90 to 94	22% (13-34%)	78% (66-87%)	1 (0.46-1.9)	2.5% (0.78-5.5%)	3% (0.83-6.7%)	1.3 (0.38-2.9)	7.2 (2.2-15)
Saint Lucia	2005	Female	95 plus	26% (14-43%)	74% (57-86%)	1.2 (0.44-2.5)	3% (0.6-6.6%)	3.5% (0.57-8%)	0.43 (0.071-0.99)	1.5 (0.31-3.5)
Saint Lucia	2005	Male	All Ages	58% (49-66%)	42% (34-51%)	3.3 (3-3.6)	10% (8.7-12%)	11% (9.1-14%)	63 (51-76)	2400 (2100-2800)
Saint Lucia	2005	Male	15 to 19	58% (41-74%)	42% (26-59%)	2.5 (1.4-3.9)	7.5% (4.9-11%)	12% (7-17%)	1.1 (0.68-1.7)	97 (63-140)
Saint Lucia	2005	Male	20 to 24	68% (50-82%)	32% (18-50%)	4.8 (3.1-6.7)	15% (11-20%)	20% (14-28%)	2.9 (2-4)	230 (170-300)
Saint Lucia	2005	Male	25 to 29	71% (54-83%)	29% (17-46%)	5.7 (3.8-7.9)	17% (13-22%)	22% (16-30%)	3.8 (2.7-5)	270 (200-340)
Saint Lucia	2005	Male	30 to 34	72% (54-85%)	28% (15-46%)	6 (3.9-8.4)	16% (13-21%)	22% (17-28%)	3.6 (2.7-4.6)	240 (180-300)
Saint Lucia	2005	Male	35 to 39	72% (55-85%)	28% (15-45%)	5.7 (3.7-8.3)	16% (12-20%)	21% (16-27%)	4.2 (3.2-5.3)	250 (190-310)
Saint Lucia	2005	Male	40 to 44	70% (55-84%)	30% (16-45%)	5.5 (3.3-8)	15% (12-19%)	20% (16-25%)	4.6 (3.5-5.7)	240 (190-300)
Saint Lucia	2005	Male	45 to 49	69% (49-83%)	31% (17-51%)	5.1 (3-8.3)	15% (11-18%)	20% (14-24%)	4.8 (3.6-5.9)	220 (170-280)
Saint Lucia	2005	Male	50 to 54	67% (49-82%)	33% (18-51%)	4.7 (2.7-7.4)	14% (10-17%)	18% (13-22%)	4.3 (3.1-5.4)	180 (130-220)
Saint Lucia	2005	Male	55 to 59	65% (47-81%)	35% (19-53%)	4.3 (2.4-6.9)	13% (9.4-17%)	17% (12-21%)	5.1 (3.6-6.6)	170 (130-230)
Saint Lucia	2005	Male	60 to 64	64% (44-80%)	36% (20-56%)	3.9 (2.1-6.5)	12% (7.5-16%)	14% (8.8-18%)	5.4 (3.5-7.3)	150 (100-210)
Saint Lucia	2005	Male	65 to 69	62% (43-78%)	38% (22-57%)	3.5 (2-5.6)	10% (6.5-14%)	13% (7.9-17%)	5.7 (3.6-7.7)	140 (86-190)
Saint Lucia	2005	Male	70 to 74	59% (41-75%)	41% (25-59%)	2.9 (1.5-4.9)	8.2% (4.5-12%)	9.6% (5.3-14%)	5.4 (3-8)	100 (59-150)
Saint Lucia	2005	Male	75 to 79	54% (35-72%)	46% (28-65%)	2.4 (1.2-4.2)	6.5% (3.2-10%)	7.5% (3.6-11%)	4.3 (2.1-6.5)	63 (31-97)
Saint Lucia	2005	Male	80 to 84	48% (29-66%)	52% (34-71%)	2 (1.1-3.3)	5.2% (2.6-8%)	6% (3-9.3%)	3.9 (2-6.1)	44 (23-67)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	2005	Male	85 to 89	43% (26-60%)	57% (40-74%)	1.8 (0.9-3)	4.3% (2.1-6.9%)	5% (2.4-8.2%)	2.6 (1.2-4.3)	21 (11-34)
Saint Lucia	2005	Male	90 to 94	43% (28-61%)	57% (39-72%)	1.8 (0.89-3.2)	3.9% (1.8-6.7%)	4.4% (1.8-7.8%)	1.2 (0.51-2.1)	6.8 (3-12)
Saint Lucia	2005	Male	95 plus	46% (27-66%)	54% (34-73%)	2 (0.79-3.9)	4.1% (1.4-7.6%)	4.5% (1.3-8.8%)	0.26 (0.076-0.52)	1 (0.36-1.9)
Saint Lucia	2010	Female	All Ages	35% (29-42%)	65% (58-71%)	1.4 (1.2-1.6)	3% (2.3-3.9%)	3.8% (2.4-5.6%)	18 (12-27)	630 (470-800)
Saint Lucia	2010	Female	15 to 19	40% (27-57%)	60% (43-73%)	1.5 (0.83-2.5)	2.2% (1.4-3.2%)	5.3% (3-8.4%)	0.17 (0.095-0.26)	20 (13-30)
Saint Lucia	2010	Female	20 to 24	46% (31-63%)	54% (37-69%)	2.3 (1.3-3.6)	4.1% (2.8-5.7%)	7.5% (4.6-11%)	0.39 (0.24-0.56)	42 (28-58)
Saint Lucia	2010	Female	25 to 29	46% (29-63%)	54% (37-71%)	2.4 (1.4-3.6)	4.1% (2.9-5.5%)	7.4% (4.7-10%)	0.4 (0.26-0.57)	42 (29-57)
Saint Lucia	2010	Female	30 to 34	44% (28-61%)	56% (39-72%)	2.3 (1.3-3.5)	4.1% (2.9-5.5%)	7.3% (4.9-9.9%)	0.53 (0.36-0.73)	45 (31-60)
Saint Lucia	2010	Female	35 to 39	43% (28-61%)	57% (39-72%)	2.1 (1.2-3.3)	4.2% (2.9-5.7%)	8.1% (5.7-11%)	0.72 (0.5-0.97)	51 (35-69)
Saint Lucia	2010	Female	40 to 44	42% (27-60%)	58% (40-73%)	2 (1.1-3.2)	4.8% (3.2-6.8%)	8.5% (5.8-12%)	1.3 (0.9-1.8)	73 (49-100)
Saint Lucia	2010	Female	45 to 49	40% (25-58%)	60% (42-75%)	1.9 (1.1-3.1)	4.7% (3.1-7.1%)	8% (5.4-12%)	1.5 (1-2.2)	73 (48-110)
Saint Lucia	2010	Female	50 to 54	37% (23-53%)	63% (47-77%)	1.7 (0.98-2.8)	4.7% (2.8-7.3%)	7.7% (4.9-11%)	1.6 (1-2.4)	66 (40-100)
Saint Lucia	2010	Female	55 to 59	32% (20-48%)	68% (52-80%)	1.5 (0.77-2.4)	4.2% (2.3-6.8%)	6.5% (3.7-10%)	1.5 (0.83-2.3)	51 (29-85)
Saint Lucia	2010	Female	60 to 64	29% (17-44%)	71% (56-83%)	1.3 (0.62-2.2)	3.4% (1.6-5.9%)	5.1% (2.5-8.5%)	1.4 (0.68-2.3)	41 (19-71)
Saint Lucia	2010	Female	65 to 69	26% (16-40%)	74% (60-84%)	1.1 (0.6-1.8)	3% (1.4-5.3%)	4.2% (2-7.2%)	1.6 (0.75-2.8)	38 (17-67)
Saint Lucia	2010	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.92 (0.46-1.6)	2.4% (0.83-4.9%)	3.4% (1.3-6.6%)	1.5 (0.54-2.9)	28 (9.3-57)
Saint Lucia	2010	Female	75 to 79	21% (13-35%)	79% (65-87%)	0.8 (0.37-1.4)	2% (0.42-4.3%)	2.7% (0.58-5.6%)	1.4 (0.29-2.9)	20 (4.4-44)
Saint Lucia	2010	Female	80 to 84	20% (11-32%)	80% (68-89%)	0.76 (0.36-1.3)	1.8% (0.53-3.8%)	2.3% (0.72-4.9%)	1.5 (0.44-3.1)	17 (5.2-36)
Saint Lucia	2010	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.77 (0.38-1.3)	1.8% (0.59-3.7%)	2.2% (0.69-4.7%)	1.5 (0.44-3.1)	12 (3.7-24)
Saint Lucia	2010	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.88 (0.38-1.6)	2% (0.57-4.3%)	2.4% (0.61-5.4%)	1.1 (0.29-2.5)	6.3 (1.7-14)
Saint Lucia	2010	Female	95 plus	26% (13-43%)	74% (57-87%)	1.1 (0.38-2.1)	2.4% (0.43-5.5%)	2.8% (0.31-6.7%)	0.48 (0.055-1.2)	1.8 (0.32-4.2)
Saint Lucia	2010	Male	All Ages	59% (50-67%)	41% (33-50%)	3.1 (2.7-3.5)	9.5% (8-11%)	10% (8.1-12%)	64 (51-79)	2500 (2100-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	2010	Male	15 to 19	57% (40-74%)	43% (26-60%)	2.2 (1.2-3.5)	6.3% (4.1-9.1%)	9.6% (5.6-15%)	0.88 (0.51-1.3)	80 (51-110)
Saint Lucia	2010	Male	20 to 24	68% (50-82%)	32% (18-50%)	4.2 (2.7-6.2)	14% (9.6-18%)	18% (12-25%)	2.5 (1.6-3.5)	200 (140-270)
Saint Lucia	2010	Male	25 to 29	71% (54-84%)	29% (16-46%)	5.1 (3.2-7.2)	16% (11-21%)	20% (14-27%)	3.7 (2.6-5.1)	260 (190-350)
Saint Lucia	2010	Male	30 to 34	71% (53-85%)	29% (15-47%)	5.3 (3.3-7.5)	15% (11-19%)	20% (14-26%)	3.6 (2.6-4.6)	240 (180-300)
Saint Lucia	2010	Male	35 to 39	71% (55-85%)	29% (15-45%)	5 (3.2-7.4)	14% (10-17%)	19% (14-24%)	3.4 (2.5-4.4)	210 (160-260)
Saint Lucia	2010	Male	40 to 44	70% (54-83%)	30% (17-46%)	4.8 (2.9-7.1)	13% (9.9-17%)	18% (13-23%)	4.5 (3.3-5.7)	240 (180-300)
Saint Lucia	2010	Male	45 to 49	68% (48-83%)	32% (17-52%)	4.5 (2.7-6.9)	13% (10-17%)	18% (13-22%)	5.7 (4.1-7.2)	270 (200-340)
Saint Lucia	2010	Male	50 to 54	67% (48-82%)	33% (18-52%)	4.2 (2.4-6.4)	12% (8.9-16%)	16% (12-21%)	5.2 (3.6-6.6)	210 (150-270)
Saint Lucia	2010	Male	55 to 59	65% (47-81%)	35% (19-53%)	3.8 (2.2-5.9)	12% (8.3-15%)	15% (10-20%)	6 (4.1-7.9)	210 (140-270)
Saint Lucia	2010	Male	60 to 64	63% (43-80%)	37% (20-57%)	3.4 (1.9-5.6)	10% (6.2-14%)	12% (7.7-17%)	5.9 (3.6-8.1)	170 (110-240)
Saint Lucia	2010	Male	65 to 69	62% (43-78%)	38% (22-57%)	3.1 (1.7-4.8)	9% (5.6-13%)	11% (6.9-15%)	5.4 (3.3-7.5)	130 (82-180)
Saint Lucia	2010	Male	70 to 74	58% (40-74%)	42% (26-60%)	2.6 (1.3-4.4)	7% (3.8-11%)	8.3% (4.4-13%)	5.4 (2.8-8.6)	110 (57-170)
Saint Lucia	2010	Male	75 to 79	53% (34-72%)	47% (28-66%)	2.1 (1.1-3.7)	5.6% (2.7-8.9%)	6.5% (3.1-10%)	4.4 (2.1-7)	67 (32-110)
Saint Lucia	2010	Male	80 to 84	47% (29-65%)	53% (35-71%)	1.8 (0.9-2.9)	4.3% (2.1-7.1%)	5.1% (2.4-8.4%)	3.4 (1.6-5.6)	38 (19-62)
Saint Lucia	2010	Male	85 to 89	42% (26-60%)	58% (40-74%)	1.6 (0.78-2.7)	3.7% (1.7-6.3%)	4.2% (2-7.4%)	2.2 (1-4)	18 (8.8-32)
Saint Lucia	2010	Male	90 to 94	43% (27-60%)	57% (40-73%)	1.6 (0.76-2.8)	3.2% (1.4-5.9%)	3.7% (1.5-6.9%)	1.2 (0.48-2.3)	7 (3-13)
Saint Lucia	2010	Male	95 plus	46% (27-66%)	54% (34-73%)	1.7 (0.68-3.3)	3.4% (1.2-6.8%)	3.7% (1-7.8%)	0.37 (0.099-0.78)	1.5 (0.49-2.9)
Saint Lucia	2016	Female	All Ages	35% (29-42%)	65% (58-71%)	1.4 (1-1.8)	3% (2.1-4.2%)	3.7% (2.1-5.7%)	19 (11-30)	650 (450-900)
Saint Lucia	2016	Female	15 to 19	40% (27-56%)	60% (44-73%)	1.5 (0.73-2.5)	2.2% (1.4-3.3%)	5.8% (3.1-9.2%)	0.17 (0.088-0.27)	19 (12-30)
Saint Lucia	2016	Female	20 to 24	46% (30-63%)	54% (37-70%)	2.2 (1.2-3.6)	3.9% (2.6-5.5%)	7.7% (4.5-11%)	0.35 (0.2-0.52)	40 (26-57)
Saint Lucia	2016	Female	25 to 29	45% (29-63%)	55% (37-71%)	2.2 (1.2-3.6)	4% (2.8-5.5%)	7.6% (4.7-11%)	0.37 (0.23-0.56)	39 (26-55)
Saint Lucia	2016	Female	30 to 34	44% (28-61%)	56% (39-72%)	2.2 (1.1-3.6)	3.9% (2.7-5.5%)	7.5% (5-11%)	0.52 (0.34-0.74)	45 (31-62)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	2016	Female	35 to 39	43% (28-61%)	57% (39-72%)	2 (1.1-3.3)	3.9% (2.7-5.5%)	8.1% (5.6-11%)	0.64 (0.43-0.91)	46 (31-64)
Saint Lucia	2016	Female	40 to 44	42% (27-61%)	58% (39-73%)	1.9 (0.99-3.2)	4.5% (2.9-6.6%)	8.4% (5.6-12%)	1.2 (0.77-1.7)	67 (44-98)
Saint Lucia	2016	Female	45 to 49	40% (24-57%)	60% (43-76%)	1.8 (0.99-3.1)	4.4% (2.8-6.7%)	7.8% (5.2-12%)	1.5 (0.96-2.1)	72 (47-110)
Saint Lucia	2016	Female	50 to 54	36% (22-54%)	64% (46-78%)	1.6 (0.87-2.8)	4.4% (2.6-6.9%)	7.5% (4.7-11%)	1.9 (1.1-2.8)	77 (46-120)
Saint Lucia	2016	Female	55 to 59	32% (19-48%)	68% (52-81%)	1.4 (0.71-2.4)	3.9% (2-6.6%)	6.2% (3.4-10%)	1.8 (1-3)	65 (35-110)
Saint Lucia	2016	Female	60 to 64	28% (17-44%)	72% (56-83%)	1.2 (0.56-2.1)	3.2% (1.4-5.9%)	4.9% (2.1-8.5%)	1.5 (0.69-2.7)	46 (19-82)
Saint Lucia	2016	Female	65 to 69	26% (16-40%)	74% (60-84%)	1 (0.52-1.8)	2.8% (1.1-5.1%)	3.9% (1.6-7%)	1.7 (0.69-3)	41 (15-75)
Saint Lucia	2016	Female	70 to 74	24% (14-37%)	76% (63-86%)	0.87 (0.41-1.6)	2.2% (0.68-4.6%)	3.2% (1.1-6.4%)	1.5 (0.49-3)	29 (8.5-59)
Saint Lucia	2016	Female	75 to 79	21% (13-34%)	79% (66-87%)	0.76 (0.33-1.4)	1.8% (0.38-4%)	2.5% (0.55-5.4%)	1.4 (0.3-3.1)	21 (4.3-47)
Saint Lucia	2016	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.71 (0.34-1.3)	1.6% (0.41-3.7%)	2.2% (0.51-4.8%)	1.7 (0.39-3.8)	19 (4.6-42)
Saint Lucia	2016	Female	85 to 89	19% (11-30%)	81% (70-89%)	0.72 (0.34-1.3)	1.7% (0.48-3.7%)	2.1% (0.53-4.6%)	1.6 (0.41-3.6)	13 (3.5-28)
Saint Lucia	2016	Female	90 to 94	21% (13-33%)	79% (67-87%)	0.81 (0.34-1.6)	1.9% (0.44-4.2%)	2.2% (0.46-5.3%)	0.93 (0.19-2.2)	5.1 (1.2-11)
Saint Lucia	2016	Female	95 plus	25% (13-43%)	75% (57-87%)	0.98 (0.34-2)	2.2% (0.35-5.6%)	2.5% (0.21-6.8%)	0.55 (0.045-1.5)	2 (0.33-5.1)
Saint Lucia	2016	Male	All Ages	59% (51-67%)	41% (33-49%)	3 (2.3-3.9)	9.4% (7.4-11%)	9.8% (7.2-13%)	65 (47-85)	2500 (2000-3100)
Saint Lucia	2016	Male	15 to 19	57% (39-73%)	43% (27-61%)	2.1 (1.1-3.6)	6.4% (4-9.3%)	10% (5.9-15%)	0.78 (0.44-1.2)	71 (45-100)
Saint Lucia	2016	Male	20 to 24	67% (51-82%)	33% (18-49%)	3.9 (2.4-6.2)	13% (9.3-18%)	18% (11-25%)	2.4 (1.5-3.4)	200 (140-270)
Saint Lucia	2016	Male	25 to 29	70% (53-83%)	30% (17-47%)	4.8 (2.8-7.3)	15% (11-20%)	20% (13-28%)	3.3 (2.1-4.6)	240 (170-320)
Saint Lucia	2016	Male	30 to 34	71% (53-85%)	29% (15-47%)	5 (2.9-7.5)	14% (11-19%)	20% (14-26%)	3.3 (2.3-4.5)	220 (160-290)
Saint Lucia	2016	Male	35 to 39	71% (55-84%)	29% (16-45%)	4.7 (2.7-7.2)	13% (9.6-17%)	19% (13-24%)	3.4 (2.3-4.5)	210 (150-270)
Saint Lucia	2016	Male	40 to 44	70% (54-83%)	30% (17-46%)	4.5 (2.4-6.9)	13% (9.2-16%)	18% (12-23%)	4 (2.7-5.2)	220 (160-280)
Saint Lucia	2016	Male	45 to 49	68% (48-83%)	32% (17-52%)	4.2 (2.4-6.8)	13% (9.3-17%)	17% (12-22%)	5.7 (4-7.4)	270 (190-350)
Saint Lucia	2016	Male	50 to 54	67% (49-82%)	33% (18-51%)	3.9 (2.1-6.3)	12% (8.3-15%)	16% (11-20%)	6.4 (4.3-8.4)	270 (180-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Lucia	2016	Male	55 to 59	65% (46-80%)	35% (20-54%)	3.6 (2-5.9)	11% (7.4-15%)	14% (9.3-19%)	7 (4.4-9.4)	240 (160-330)
Saint Lucia	2016	Male	60 to 64	63% (43-80%)	37% (20-57%)	3.2 (1.6-5.5)	9.6% (5.3-14%)	12% (6.6-17%)	6.5 (3.7-9.2)	190 (110-270)
Saint Lucia	2016	Male	65 to 69	62% (42-78%)	38% (22-58%)	2.9 (1.5-4.8)	8.5% (4.9-12%)	10% (6-15%)	6.1 (3.4-8.9)	150 (84-210)
Saint Lucia	2016	Male	70 to 74	58% (39-74%)	42% (26-61%)	2.4 (1.2-4.2)	6.5% (3.2-10%)	7.7% (3.8-12%)	5 (2.5-8)	98 (49-160)
Saint Lucia	2016	Male	75 to 79	53% (34-71%)	47% (29-66%)	2 (0.98-3.6)	5.1% (2.3-8.6%)	6% (2.7-10%)	4.6 (2-7.6)	69 (32-120)
Saint Lucia	2016	Male	80 to 84	47% (29-65%)	53% (35-71%)	1.7 (0.83-2.9)	4% (1.8-6.9%)	4.7% (2.1-8.1%)	3.5 (1.6-6.2)	40 (18-69)
Saint Lucia	2016	Male	85 to 89	42% (26-59%)	58% (41-74%)	1.5 (0.69-2.6)	3.4% (1.6-6%)	3.9% (1.8-7.1%)	2.1 (0.95-3.9)	17 (8.1-31)
Saint Lucia	2016	Male	90 to 94	42% (27-59%)	58% (41-73%)	1.5 (0.67-2.6)	3% (1.2-5.8%)	3.4% (1.2-6.7%)	1 (0.36-2)	5.7 (2.3-11)
Saint Lucia	2016	Male	95 plus	45% (26-66%)	55% (34-74%)	1.6 (0.61-3.2)	3.2% (0.94-6.6%)	3.4% (0.76-7.5%)	0.4 (0.089-0.87)	1.6 (0.44-3.2)
Saint Vincent and the Grenadines	1990	Female	All Ages	35% (28-42%)	65% (58-72%)	0.77 (0.64-0.9)	1.4% (0.83-2.1%)	1.2% (0.071-2.6%)	4.4 (0.26-9.3)	210 (120-310)
Saint Vincent and the Grenadines	1990	Female	15 to 19	43% (29-59%)	57% (41-71%)	1 (0.57-1.6)	2% (1.3-2.9%)	3.9% (2.3-6.1%)	0.14 (0.081-0.23)	16 (10-23)
Saint Vincent and the Grenadines	1990	Female	20 to 24	49% (33-65%)	51% (35-67%)	1.5 (0.92-2.2)	3.4% (2.3-4.8%)	5.3% (3.3-7.7%)	0.23 (0.14-0.34)	26 (17-36)
Saint Vincent and the Grenadines	1990	Female	25 to 29	49% (32-66%)	51% (34-68%)	1.6 (0.97-2.3)	3.7% (2.6-4.8%)	5.5% (3.5-7.9%)	0.29 (0.18-0.43)	28 (20-39)
Saint Vincent and the Grenadines	1990	Female	30 to 34	47% (30-64%)	53% (36-70%)	1.5 (0.88-2.4)	3.6% (2.5-4.8%)	5.5% (3.6-7.8%)	0.3 (0.19-0.43)	24 (17-34)
Saint Vincent and the Grenadines	1990	Female	35 to 39	46% (31-63%)	54% (37-69%)	1.4 (0.8-2.3)	3.4% (2.1-4.9%)	5.2% (3.1-7.8%)	0.3 (0.18-0.46)	20 (13-29)
Saint Vincent and the Grenadines	1990	Female	40 to 44	45% (30-63%)	55% (37-70%)	1.4 (0.7-2.3)	3.3% (1.9-5.3%)	5.2% (2.9-8.3%)	0.31 (0.17-0.5)	17 (9.8-27)
Saint Vincent and the Grenadines	1990	Female	45 to 49	43% (27-60%)	57% (40-73%)	1.3 (0.69-2.2)	3.1% (1.4-5.2%)	4.6% (2.1-7.7%)	0.3 (0.14-0.51)	14 (6.7-23)
Saint Vincent and the Grenadines	1990	Female	50 to 54	39% (25-57%)	61% (43-75%)	1.2 (0.63-2)	2.8% (0.89-5.2%)	3.9% (1.3-7.2%)	0.48 (0.16-0.88)	19 (6.4-36)
Saint Vincent and the Grenadines	1990	Female	55 to 59	35% (22-52%)	65% (48-78%)	1 (0.51-1.7)	2.2% (0.47-4.4%)	3% (0.69-5.9%)	0.45 (0.1-0.9)	15 (3.2-31)
Saint Vincent and the Grenadines	1990	Female	60 to 64	31% (19-47%)	69% (53-81%)	0.86 (0.44-1.5)	1.5% (-0.19-3.6%)	2% (-0.18-4.7%)	0.44 (-0.038-1)	12 (-1.5-30)
Saint Vincent and the Grenadines	1990	Female	65 to 69	29% (18-43%)	71% (57-82%)	0.74 (0.39-1.2)	0.97% (-0.52-2.8%)	1.3% (-0.54-3.6%)	0.39 (-0.17-1.1)	8.9 (-4.8-26)
Saint Vincent and the Grenadines	1990	Female	70 to 74	26% (17-40%)	74% (60-83%)	0.63 (0.31-1.1)	0.47% (-1.2-3%)	0.61% (-1.2-2.9%)	0.24 (-0.48-1.1)	4.3 (-9.3-21)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	1990	Female	75 to 79	23% (14-36%)	77% (64-86%)	0.54 (0.24-1)	0.26% (-1.1-1.9%)	0.31% (-1.3-2.4%)	0.14 (-0.57-1)	2 (-8.3-15)
Saint Vincent and the Grenadines	1990	Female	80 to 84	21% (12-34%)	79% (66-88%)	0.51 (0.24-0.89)	0.28% (-0.89-1.7%)	0.31% (-1.1-2%)	0.14 (-0.49-0.92)	1.6 (-5.1-9.9)
Saint Vincent and the Grenadines	1990	Female	85 to 89	21% (13-33%)	79% (67-87%)	0.52 (0.25-0.92)	0.28% (-0.76-1.8%)	0.28% (-0.94-2%)	0.11 (-0.39-0.83)	1 (-2.7-6.4)
Saint Vincent and the Grenadines	1990	Female	90 to 94	24% (14-37%)	76% (63-86%)	0.59 (0.27-1.1)	0.45% (-0.73-2.3%)	0.45% (-0.96-2.6%)	0.093 (-0.2-0.55)	0.56 (-0.9-2.8)
Saint Vincent and the Grenadines	1990	Female	95 plus	28% (15-46%)	72% (54-85%)	0.71 (0.26-1.4)	0.62% (-1.1-3.3%)	0.58% (-1.4-3.7%)	0.037 (-0.09-0.23)	0.15 (-0.26-0.81)
Saint Vincent and the Grenadines	1990	Male	All Ages	55% (47-64%)	45% (36-53%)	1.6 (1.3-1.8)	5.8% (4.5-7.2%)	6.4% (4.6-8.7%)	24 (16-32)	980 (780-1200)
Saint Vincent and the Grenadines	1990	Male	15 to 19	60% (43-75%)	40% (25-57%)	1.5 (0.86-2.2)	5.7% (3.9-7.7%)	7.5% (4.8-11%)	0.61 (0.39-0.93)	57 (39-81)
Saint Vincent and the Grenadines	1990	Male	20 to 24	69% (52-83%)	31% (17-48%)	2.7 (1.8-3.8)	11% (8.2-14%)	12% (8.7-17%)	1.3 (0.87-1.8)	120 (87-160)
Saint Vincent and the Grenadines	1990	Male	25 to 29	72% (56-84%)	28% (16-44%)	3.3 (2.2-4.5)	12% (9.1-15%)	14% (9.8-18%)	1.4 (0.97-2)	120 (90-160)
Saint Vincent and the Grenadines	1990	Male	30 to 34	73% (56-85%)	27% (15-44%)	3.4 (2.2-4.9)	12% (9.2-15%)	15% (11-19%)	1.3 (0.89-1.8)	96 (71-130)
Saint Vincent and the Grenadines	1990	Male	35 to 39	73% (57-85%)	27% (15-43%)	3.2 (2-5)	12% (9.1-15%)	15% (11-20%)	1.2 (0.83-1.6)	77 (57-100)
Saint Vincent and the Grenadines	1990	Male	40 to 44	72% (56-85%)	28% (15-44%)	3.1 (1.7-4.9)	12% (8.8-16%)	15% (11-21%)	1.2 (0.84-1.7)	69 (49-91)
Saint Vincent and the Grenadines	1990	Male	45 to 49	71% (52-84%)	29% (16-48%)	3 (1.6-4.8)	12% (8.3-16%)	15% (10-20%)	1.4 (0.94-1.9)	67 (47-91)
Saint Vincent and the Grenadines	1990	Male	50 to 54	69% (51-83%)	31% (17-49%)	2.7 (1.5-4.3)	11% (7.5-15%)	14% (9-19%)	2 (1.3-2.8)	82 (55-110)
Saint Vincent and the Grenadines	1990	Male	55 to 59	67% (50-82%)	33% (18-50%)	2.5 (1.3-4)	10% (6.6-14%)	12% (7.9-17%)	2.2 (1.4-3.2)	77 (50-110)
Saint Vincent and the Grenadines	1990	Male	60 to 64	66% (46-82%)	34% (18-54%)	2.3 (1.2-3.8)	8.1% (4.6-12%)	9.4% (5.3-15%)	2.4 (1.4-3.7)	69 (40-110)
Saint Vincent and the Grenadines	1990	Male	65 to 69	64% (46-80%)	36% (20-54%)	2 (1.1-3.2)	6.7% (3.8-10%)	7.7% (4.3-12%)	2.5 (1.4-4)	60 (33-95)
Saint Vincent and the Grenadines	1990	Male	70 to 74	61% (43-76%)	39% (24-57%)	1.7 (0.86-2.9)	4.7% (1.9-8.4%)	5.3% (2.1-9.5%)	2.3 (0.85-4.2)	44 (18-79)
Saint Vincent and the Grenadines	1990	Male	75 to 79	56% (37-74%)	44% (26-63%)	1.4 (0.7-2.4)	3.6% (1.3-6.9%)	4.1% (1.4-7.8%)	1.8 (0.6-3.7)	27 (9.4-53)
Saint Vincent and the Grenadines	1990	Male	80 to 84	51% (32-69%)	49% (31-68%)	1.2 (0.62-2)	2.8% (0.96-5.5%)	3.1% (1-6.2%)	1.1 (0.36-2.2)	12 (4.1-23)
Saint Vincent and the Grenadines	1990	Male	85 to 89	46% (29-63%)	54% (37-71%)	1.1 (0.53-1.9)	2.2% (0.7-4.6%)	2.4% (0.65-5.1%)	0.6 (0.16-1.3)	4.7 (1.5-9.9)
Saint Vincent and the Grenadines	1990	Male	90 to 94	46% (30-63%)	54% (37-70%)	1.1 (0.53-1.9)	1.9% (0.36-4.3%)	2% (0.21-4.8%)	0.2 (0.021-0.48)	1.1 (0.22-2.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	1990	Male	95 plus	49% (30-69%)	51% (31-70%)	1.2 (0.47-2.2)	1.9% (0.074-5.1%)	1.9% (-0.27-5.6%)	0.039 (-0.0051-0.11)	0.15 (0.0061-0.41)
Saint Vincent and the Grenadines	1995	Female	All Ages	35% (29-42%)	65% (58-71%)	0.78 (0.66-0.9)	1.3% (0.74-2%)	1% (-0.11-2.4%)	4 (-0.4-9.1)	200 (110-310)
Saint Vincent and the Grenadines	1995	Female	15 to 19	43% (29-59%)	57% (41-71%)	0.98 (0.53-1.6)	1.8% (1.1-2.7%)	3.4% (1.9-5.5%)	0.12 (0.067-0.19)	14 (8.4-20)
Saint Vincent and the Grenadines	1995	Female	20 to 24	49% (33-64%)	51% (36-67%)	1.5 (0.85-2.2)	3% (2-4.2%)	4.3% (2.5-6.4%)	0.2 (0.12-0.3)	24 (15-33)
Saint Vincent and the Grenadines	1995	Female	25 to 29	48% (32-65%)	52% (35-68%)	1.5 (0.92-2.2)	3.3% (2.3-4.4%)	4.5% (2.8-6.6%)	0.25 (0.15-0.38)	25 (17-34)
Saint Vincent and the Grenadines	1995	Female	30 to 34	46% (30-64%)	54% (36-70%)	1.5 (0.88-2.2)	3.2% (2.2-4.4%)	4.5% (2.9-6.5%)	0.35 (0.21-0.52)	28 (19-39)
Saint Vincent and the Grenadines	1995	Female	35 to 39	46% (31-63%)	54% (37-69%)	1.4 (0.77-2.2)	3% (1.9-4.5%)	4.4% (2.6-6.7%)	0.36 (0.21-0.57)	24 (15-36)
Saint Vincent and the Grenadines	1995	Female	40 to 44	44% (29-63%)	56% (37-71%)	1.3 (0.69-2.1)	3% (1.7-4.8%)	4.6% (2.5-7.3%)	0.39 (0.2-0.63)	21 (12-34)
Saint Vincent and the Grenadines	1995	Female	45 to 49	42% (27-60%)	58% (40-73%)	1.2 (0.66-2.1)	2.8% (1.2-4.9%)	4% (1.7-7%)	0.35 (0.15-0.61)	17 (7-28)
Saint Vincent and the Grenadines	1995	Female	50 to 54	39% (24-56%)	61% (44-76%)	1.1 (0.6-1.9)	2.5% (0.67-4.9%)	3.5% (1-6.6%)	0.35 (0.1-0.66)	14 (3.7-26)
Saint Vincent and the Grenadines	1995	Female	55 to 59	34% (21-52%)	66% (48-79%)	0.95 (0.48-1.6)	1.8% (0.19-4%)	2.5% (0.34-5.4%)	0.46 (0.062-1)	15 (1.6-34)
Saint Vincent and the Grenadines	1995	Female	60 to 64	31% (19-48%)	69% (52-81%)	0.81 (0.41-1.5)	1.2% (-0.41-3.4%)	1.6% (-0.46-4.4%)	0.36 (-0.1-0.96)	10 (-3.7-28)
Saint Vincent and the Grenadines	1995	Female	65 to 69	28% (18-43%)	72% (57-82%)	0.71 (0.37-1.2)	0.64% (-0.78-2.4%)	0.86% (-0.85-2.9%)	0.28 (-0.27-0.94)	6.1 (-7.3-22)
Saint Vincent and the Grenadines	1995	Female	70 to 74	26% (16-39%)	74% (61-84%)	0.6 (0.29-1)	0.23% (-1.2-1.9%)	0.33% (-1.4-2.3%)	0.14 (-0.59-1)	2.3 (-12-19)
Saint Vincent and the Grenadines	1995	Female	75 to 79	23% (13-37%)	77% (63-87%)	0.52 (0.23-0.98)	0.1% (-1.2-1.8%)	0.14% (-1.4-2.1%)	0.071 (-0.72-1.1)	0.93 (-11-15)
Saint Vincent and the Grenadines	1995	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.49 (0.23-0.85)	0.21% (-0.93-1.7%)	0.24% (-1.1-2%)	0.12 (-0.55-1)	1.3 (-6-11)
Saint Vincent and the Grenadines	1995	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.5 (0.24-0.89)	0.18% (-0.85-1.6%)	0.17% (-1-1.9%)	0.069 (-0.42-0.78)	0.67 (-3-5.9)
Saint Vincent and the Grenadines	1995	Female	90 to 94	24% (14-37%)	76% (63-86%)	0.58 (0.25-1.1)	0.3% (-0.89-2.1%)	0.26% (-1.1-2.4%)	0.063 (-0.27-0.58)	0.43 (-1.3-3)
Saint Vincent and the Grenadines	1995	Female	95 plus	28% (14-46%)	72% (54-86%)	0.69 (0.25-1.4)	0.47% (-1.1-3.2%)	0.4% (-1.5-3.5%)	0.032 (-0.12-0.28)	0.15 (-0.36-0.98)
Saint Vincent and the Grenadines	1995	Male	All Ages	57% (48-65%)	43% (35-52%)	1.6 (1.4-1.9)	5.8% (4.6-7.2%)	5.9% (4.1-8.1%)	24 (17-33)	1000 (810-1200)
Saint Vincent and the Grenadines	1995	Male	15 to 19	60% (42-75%)	40% (25-58%)	1.4 (0.8-2.2)	5.5% (3.8-7.7%)	7.2% (4.5-11%)	0.63 (0.37-0.95)	58 (38-84)
Saint Vincent and the Grenadines	1995	Male	20 to 24	69% (52-83%)	31% (17-48%)	2.6 (1.7-3.7)	11% (7.9-14%)	12% (8.1-16%)	1.4 (0.87-2)	120 (87-160)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	1995	Male	25 to 29	72% (56-84%)	28% (16-44%)	3.1 (2.1-4.3)	11% (8.4-14%)	13% (8.9-17%)	1.3 (0.91-1.8)	110 (82-140)
Saint Vincent and the Grenadines	1995	Male	30 to 34	73% (55-85%)	27% (15-45%)	3.3 (2.1-4.5)	11% (8.1-13%)	13% (9.2-17%)	1.5 (1-2)	110 (80-140)
Saint Vincent and the Grenadines	1995	Male	35 to 39	73% (57-85%)	27% (15-43%)	3.1 (1.9-4.6)	11% (8.1-14%)	13% (9.6-18%)	1.4 (0.98-2)	94 (69-120)
Saint Vincent and the Grenadines	1995	Male	40 to 44	72% (57-85%)	28% (15-43%)	3 (1.7-4.6)	11% (7.9-14%)	13% (9.5-18%)	1.4 (0.96-2)	80 (57-110)
Saint Vincent and the Grenadines	1995	Male	45 to 49	70% (51-84%)	30% (16-49%)	2.8 (1.5-4.6)	11% (7.4-14%)	13% (9-18%)	1.5 (1-2.2)	75 (51-100)
Saint Vincent and the Grenadines	1995	Male	50 to 54	69% (51-83%)	31% (17-49%)	2.6 (1.5-4.1)	9.8% (6.6-14%)	12% (7.7-17%)	1.7 (1.1-2.4)	69 (44-97)
Saint Vincent and the Grenadines	1995	Male	55 to 59	67% (49-82%)	33% (18-51%)	2.4 (1.3-3.8)	9.2% (5.8-13%)	11% (7-16%)	2.4 (1.5-3.5)	83 (53-120)
Saint Vincent and the Grenadines	1995	Male	60 to 64	65% (46-82%)	35% (18-54%)	2.2 (1.2-3.6)	7.4% (4.1-12%)	8.7% (4.7-14%)	2.3 (1.3-3.7)	68 (37-110)
Saint Vincent and the Grenadines	1995	Male	65 to 69	64% (45-80%)	36% (20-55%)	1.9 (1-3.1)	6.2% (3.3-9.8%)	7.1% (3.8-11%)	2.6 (1.4-4.1)	61 (33-95)
Saint Vincent and the Grenadines	1995	Male	70 to 74	60% (42-76%)	40% (24-58%)	1.6 (0.82-2.7)	4.3% (1.7-7.7%)	4.8% (1.9-8.9%)	2.1 (0.85-4)	41 (17-75)
Saint Vincent and the Grenadines	1995	Male	75 to 79	55% (36-73%)	45% (27-64%)	1.3 (0.66-2.3)	3.2% (1-6.4%)	3.6% (1.1-7.2%)	1.9 (0.56-3.8)	28 (9.1-56)
Saint Vincent and the Grenadines	1995	Male	80 to 84	50% (31-67%)	50% (33-69%)	1.1 (0.58-1.9)	2.3% (0.66-4.7%)	2.5% (0.6-5.2%)	1.1 (0.26-2.2)	12 (3.4-24)
Saint Vincent and the Grenadines	1995	Male	85 to 89	45% (28-62%)	55% (38-72%)	1 (0.51-1.7)	1.8% (0.4-3.9%)	1.8% (0.28-4.2%)	0.56 (0.089-1.3)	4.6 (1-10)
Saint Vincent and the Grenadines	1995	Male	90 to 94	46% (30-63%)	54% (37-70%)	1 (0.49-1.9)	1.5% (0.15-3.7%)	1.5% (-0.06-4.1%)	0.21 (-0.0085-0.57)	1.2 (0.12-3)
Saint Vincent and the Grenadines	1995	Male	95 plus	49% (30-69%)	51% (31-70%)	1.1 (0.45-2.1)	1.6% (-0.073-4.4%)	1.5% (-0.41-4.8%)	0.053 (-0.014-0.17)	0.22 (-0.01-0.61)
Saint Vincent and the Grenadines	2000	Female	All Ages	35% (29-42%)	65% (58-71%)	0.8 (0.66-0.93)	1.2% (0.64-1.9%)	0.92% (-0.13-2.2%)	3.5 (-0.52-8.6)	190 (94-290)
Saint Vincent and the Grenadines	2000	Female	15 to 19	42% (29-59%)	58% (41-71%)	0.97 (0.52-1.6)	1.7% (1.1-2.6%)	3.2% (1.7-5.4%)	0.11 (0.059-0.18)	12 (7.7-19)
Saint Vincent and the Grenadines	2000	Female	20 to 24	48% (32-64%)	52% (36-68%)	1.4 (0.79-2.2)	2.7% (1.8-3.8%)	3.7% (2.1-5.7%)	0.17 (0.093-0.26)	21 (13-29)
Saint Vincent and the Grenadines	2000	Female	25 to 29	48% (31-65%)	52% (35-69%)	1.5 (0.88-2.2)	2.8% (1.9-3.7%)	3.5% (2.1-5.1%)	0.19 (0.11-0.29)	20 (13-28)
Saint Vincent and the Grenadines	2000	Female	30 to 34	46% (29-63%)	54% (37-71%)	1.4 (0.82-2.2)	2.8% (1.9-3.9%)	3.9% (2.4-5.5%)	0.26 (0.15-0.39)	22 (14-31)
Saint Vincent and the Grenadines	2000	Female	35 to 39	45% (30-62%)	55% (38-70%)	1.3 (0.77-2.1)	2.6% (1.7-3.9%)	3.8% (2.4-5.7%)	0.36 (0.21-0.58)	25 (16-37)
Saint Vincent and the Grenadines	2000	Female	40 to 44	44% (29-62%)	56% (38-71%)	1.3 (0.68-2)	2.7% (1.5-4.2%)	4% (2.2-6.2%)	0.41 (0.22-0.67)	23 (12-37)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	2000	Female	45 to 49	42% (26-58%)	58% (42-74%)	1.2 (0.64-2)	2.4% (1-4.2%)	3.5% (1.6-5.9%)	0.39 (0.18-0.68)	18 (8.1-32)
Saint Vincent and the Grenadines	2000	Female	50 to 54	38% (24-55%)	62% (45-76%)	1.1 (0.58-1.8)	2.2% (0.54-4.4%)	3.1% (0.86-5.9%)	0.39 (0.11-0.76)	15 (3.8-30)
Saint Vincent and the Grenadines	2000	Female	55 to 59	34% (21-51%)	66% (49-79%)	0.93 (0.47-1.6)	1.6% (0.078-3.8%)	2.2% (0.25-4.9%)	0.31 (0.033-0.7)	10 (0.46-24)
Saint Vincent and the Grenadines	2000	Female	60 to 64	30% (18-47%)	70% (53-82%)	0.79 (0.39-1.4)	1% (-0.55-3.2%)	1.4% (-0.6-4.2%)	0.36 (-0.15-1)	9.8 (-5.1-30)
Saint Vincent and the Grenadines	2000	Female	65 to 69	28% (18-42%)	72% (58-82%)	0.69 (0.36-1.1)	0.53% (-0.86-2.2%)	0.75% (-0.93-2.7%)	0.23 (-0.28-0.81)	4.8 (-7.8-19)
Saint Vincent and the Grenadines	2000	Female	70 to 74	25% (16-40%)	75% (60-84%)	0.59 (0.29-1)	0.2% (-1.2-1.9%)	0.31% (-1.4-2.4%)	0.13 (-0.57-1)	1.9 (-12-19)
Saint Vincent and the Grenadines	2000	Female	75 to 79	23% (13-36%)	77% (64-87%)	0.51 (0.23-0.92)	0.087% (-1.2-1.7%)	0.13% (-1.4-2%)	0.068 (-0.7-1)	0.79 (-11-15)
Saint Vincent and the Grenadines	2000	Female	80 to 84	21% (12-33%)	79% (67-88%)	0.48 (0.23-0.81)	0.1% (-0.97-1.4%)	0.12% (-1.1-1.7%)	0.068 (-0.67-0.99)	0.77 (-7.3-11)
Saint Vincent and the Grenadines	2000	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.49 (0.23-0.86)	0.12% (-0.9-1.5%)	0.097% (-1.1-1.6%)	0.043 (-0.48-0.72)	0.48 (-3.5-5.7)
Saint Vincent and the Grenadines	2000	Female	90 to 94	23% (14-36%)	77% (64-86%)	0.55 (0.24-1.1)	0.22% (-0.88-1.9%)	0.18% (-1.1-2.1%)	0.041 (-0.27-0.48)	0.31 (-1.2-2.5)
Saint Vincent and the Grenadines	2000	Female	95 plus	27% (14-45%)	73% (55-86%)	0.66 (0.24-1.3)	0.38% (-1.1-2.8%)	0.29% (-1.5-3.1%)	0.025 (-0.13-0.27)	0.13 (-0.39-0.93)
Saint Vincent and the Grenadines	2000	Male	All Ages	57% (49-66%)	43% (34-51%)	1.7 (1.5-1.9)	5.8% (4.6-7.1%)	5.7% (4-7.6%)	25 (18-34)	1100 (850-1300)
Saint Vincent and the Grenadines	2000	Male	15 to 19	59% (42-75%)	41% (25-58%)	1.4 (0.74-2.2)	5.4% (3.6-7.7%)	7% (4.2-10%)	0.63 (0.37-1)	57 (38-85)
Saint Vincent and the Grenadines	2000	Male	20 to 24	69% (51-83%)	31% (17-49%)	2.6 (1.6-3.7)	10% (7.4-13%)	11% (7.2-15%)	1.4 (0.88-2)	120 (85-170)
Saint Vincent and the Grenadines	2000	Male	25 to 29	72% (55-84%)	28% (16-45%)	3.1 (2-4.4)	9.3% (6.9-12%)	9.9% (6.8-14%)	1.2 (0.78-1.7)	100 (75-140)
Saint Vincent and the Grenadines	2000	Male	30 to 34	73% (55-85%)	27% (15-45%)	3.2 (2.1-4.6)	7.9% (6-10%)	8.3% (5.7-11%)	1 (0.7-1.5)	84 (61-110)
Saint Vincent and the Grenadines	2000	Male	35 to 39	72% (56-85%)	28% (15-44%)	3 (1.9-4.5)	9.7% (7.3-12%)	11% (8.1-15%)	1.8 (1.2-2.4)	120 (85-150)
Saint Vincent and the Grenadines	2000	Male	40 to 44	71% (56-84%)	29% (16-44%)	2.9 (1.7-4.4)	10% (7.5-13%)	12% (8.6-17%)	2 (1.4-2.7)	110 (80-150)
Saint Vincent and the Grenadines	2000	Male	45 to 49	70% (51-83%)	30% (17-49%)	2.8 (1.5-4.4)	9.4% (6.6-13%)	11% (7.6-15%)	1.8 (1.2-2.5)	90 (63-120)
Saint Vincent and the Grenadines	2000	Male	50 to 54	68% (50-83%)	32% (17-50%)	2.5 (1.4-4.1)	9.2% (6.1-13%)	11% (7-15%)	2 (1.3-2.8)	83 (53-120)
Saint Vincent and the Grenadines	2000	Male	55 to 59	66% (49-82%)	34% (18-51%)	2.3 (1.3-3.8)	8.6% (5.3-12%)	10% (6.2-15%)	2.1 (1.3-3.1)	73 (47-110)
Saint Vincent and the Grenadines	2000	Male	60 to 64	65% (45-81%)	35% (19-55%)	2.1 (1.1-3.6)	7% (3.9-11%)	8.1% (4.5-13%)	2.6 (1.4-4.1)	76 (42-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	2000	Male	65 to 69	63% (44-79%)	37% (21-56%)	1.9 (1-3)	5.9% (3.1-9.4%)	6.7% (3.5-11%)	2.6 (1.4-4.2)	62 (33-100)
Saint Vincent and the Grenadines	2000	Male	70 to 74	60% (41-75%)	40% (25-59%)	1.6 (0.81-2.7)	4.2% (1.8-7.6%)	4.7% (2-8.4%)	2.3 (0.98-4.1)	44 (19-79)
Saint Vincent and the Grenadines	2000	Male	75 to 79	55% (35-73%)	45% (27-65%)	1.3 (0.65-2.3)	3% (1-5.9%)	3.3% (1.1-6.6%)	1.7 (0.55-3.4)	26 (8.6-51)
Saint Vincent and the Grenadines	2000	Male	80 to 84	49% (31-67%)	51% (33-69%)	1.1 (0.57-1.8)	2.2% (0.72-4.3%)	2.3% (0.71-4.7%)	1.1 (0.33-2.3)	12 (4.1-25)
Saint Vincent and the Grenadines	2000	Male	85 to 89	44% (28-62%)	56% (38-72%)	0.96 (0.49-1.6)	1.7% (0.48-3.5%)	1.8% (0.43-3.8%)	0.6 (0.15-1.3)	4.9 (1.4-10)
Saint Vincent and the Grenadines	2000	Male	90 to 94	45% (29-62%)	55% (38-71%)	0.98 (0.49-1.8)	1.4% (0.26-3.5%)	1.5% (0.098-3.9%)	0.23 (0.016-0.6)	1.3 (0.24-3.3)
Saint Vincent and the Grenadines	2000	Male	95 plus	48% (28-68%)	52% (32-72%)	1.1 (0.44-2)	1.5% (0.035-4.2%)	1.4% (-0.29-4.5%)	0.063 (-0.014-0.2)	0.27 (0.0058-0.73)
Saint Vincent and the Grenadines	2005	Female	All Ages	36% (29-43%)	64% (57-71%)	0.9 (0.75-1.1)	1.5% (0.82-2.2%)	1.3% (0.09-2.7%)	4.8 (0.35-10)	210 (120-320)
Saint Vincent and the Grenadines	2005	Female	15 to 19	42% (28-58%)	58% (42-72%)	1 (0.55-1.7)	1.9% (1.2-2.9%)	3.7% (2-6%)	0.11 (0.059-0.19)	12 (7.5-19)
Saint Vincent and the Grenadines	2005	Female	20 to 24	48% (32-64%)	52% (36-68%)	1.6 (0.85-2.4)	3% (2-4.2%)	4.5% (2.5-6.8%)	0.18 (0.095-0.27)	21 (13-30)
Saint Vincent and the Grenadines	2005	Female	25 to 29	47% (30-64%)	53% (36-70%)	1.6 (0.91-2.4)	3% (2-4.1%)	4.1% (2.4-6.2%)	0.19 (0.1-0.29)	21 (13-29)
Saint Vincent and the Grenadines	2005	Female	30 to 34	45% (29-62%)	55% (38-71%)	1.5 (0.89-2.4)	2.8% (1.9-3.9%)	4.2% (2.7-6%)	0.22 (0.13-0.34)	20 (13-28)
Saint Vincent and the Grenadines	2005	Female	35 to 39	45% (30-62%)	55% (38-70%)	1.4 (0.85-2.2)	2.8% (1.8-4%)	4.4% (2.8-6.5%)	0.3 (0.19-0.47)	21 (14-31)
Saint Vincent and the Grenadines	2005	Female	40 to 44	43% (29-61%)	57% (39-71%)	1.3 (0.74-2.2)	2.8% (1.5-4.4%)	4.5% (2.5-7%)	0.48 (0.27-0.76)	27 (15-43)
Saint Vincent and the Grenadines	2005	Female	45 to 49	42% (26-59%)	58% (41-74%)	1.3 (0.7-2.1)	2.7% (1.2-4.5%)	4% (2-6.6%)	0.52 (0.25-0.86)	25 (11-41)
Saint Vincent and the Grenadines	2005	Female	50 to 54	38% (24-55%)	62% (45-76%)	1.2 (0.63-2)	2.5% (0.79-4.8%)	3.7% (1.3-6.7%)	0.53 (0.18-0.96)	21 (6.5-39)
Saint Vincent and the Grenadines	2005	Female	55 to 59	34% (20-51%)	66% (49-80%)	1 (0.51-1.7)	2% (0.37-4.3%)	2.8% (0.7-5.8%)	0.45 (0.11-0.93)	15 (2.8-33)
Saint Vincent and the Grenadines	2005	Female	60 to 64	30% (18-46%)	70% (54-82%)	0.86 (0.42-1.5)	1.4% (-0.26-3.7%)	1.9% (-0.2-5%)	0.34 (-0.037-0.87)	9.4 (-1.8-25)
Saint Vincent and the Grenadines	2005	Female	65 to 69	28% (17-42%)	72% (58-83%)	0.75 (0.4-1.2)	0.85% (-0.67-2.7%)	1.2% (-0.69-3.4%)	0.39 (-0.23-1.1)	8.5 (-6.9-26)
Saint Vincent and the Grenadines	2005	Female	70 to 74	25% (16-39%)	75% (61-84%)	0.63 (0.31-1.1)	0.54% (-0.9-2.5%)	0.75% (-1-3.1%)	0.29 (-0.37-1.2)	4.9 (-8.1-23)
Saint Vincent and the Grenadines	2005	Female	75 to 79	22% (13-35%)	78% (65-87%)	0.54 (0.24-0.97)	0.3% (-1-2%)	0.39% (-1.2-2.4%)	0.2 (-0.61-1.3)	2.7 (-9.2-18)
Saint Vincent and the Grenadines	2005	Female	80 to 84	21% (12-32%)	79% (68-88%)	0.51 (0.25-0.88)	0.34% (-0.81-1.8%)	0.4% (-0.95-2.1%)	0.23 (-0.55-1.2)	2.5 (-5.8-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	2005	Female	85 to 89	21% (12-32%)	79% (68-88%)	0.52 (0.25-0.9)	0.36% (-0.7-1.8%)	0.38% (-0.83-2%)	0.2 (-0.43-1)	1.6 (-3.1-8)
Saint Vincent and the Grenadines	2005	Female	90 to 94	23% (14-36%)	77% (64-86%)	0.59 (0.27-1.1)	0.46% (-0.76-2.2%)	0.45% (-0.98-2.6%)	0.11 (-0.25-0.64)	0.69 (-1.2-3.4)
Saint Vincent and the Grenadines	2005	Female	95 plus	27% (14-44%)	73% (56-86%)	0.71 (0.25-1.4)	0.64% (-0.93-3.1%)	0.59% (-1.3-3.5%)	0.051 (-0.11-0.3)	0.22 (-0.32-1)
Saint Vincent and the Grenadines	2005	Male	All Ages	59% (50-66%)	41% (34-50%)	1.9 (1.7-2.2)	6.9% (5.7-8.3%)	6.8% (5-8.9%)	30 (22-39)	1300 (1000-1500)
Saint Vincent and the Grenadines	2005	Male	15 to 19	59% (41-75%)	41% (25-59%)	1.5 (0.81-2.4)	5.9% (3.9-8.2%)	7.6% (4.6-12%)	0.64 (0.38-0.99)	57 (37-83)
Saint Vincent and the Grenadines	2005	Male	20 to 24	69% (51-82%)	31% (18-49%)	2.7 (1.7-4)	11% (8.1-15%)	13% (8.4-18%)	1.5 (0.99-2.2)	130 (92-180)
Saint Vincent and the Grenadines	2005	Male	25 to 29	71% (54-84%)	29% (16-46%)	3.3 (2.1-4.8)	11% (8.5-15%)	13% (9-18%)	1.5 (0.98-2.2)	120 (85-160)
Saint Vincent and the Grenadines	2005	Male	30 to 34	72% (55-85%)	28% (15-45%)	3.4 (2.2-4.8)	11% (8.3-14%)	13% (9.5-18%)	1.6 (1.1-2.2)	110 (81-150)
Saint Vincent and the Grenadines	2005	Male	35 to 39	72% (57-85%)	28% (15-43%)	3.3 (2.1-4.8)	11% (8-13%)	13% (9.4-17%)	1.7 (1.2-2.4)	110 (82-150)
Saint Vincent and the Grenadines	2005	Male	40 to 44	71% (56-84%)	29% (16-44%)	3.1 (1.8-4.7)	11% (8.1-14%)	14% (9.8-18%)	2.5 (1.8-3.4)	140 (100-190)
Saint Vincent and the Grenadines	2005	Male	45 to 49	70% (51-84%)	30% (16-49%)	2.9 (1.6-4.6)	11% (7.7-14%)	13% (9.2-18%)	2.7 (1.9-3.7)	130 (94-180)
Saint Vincent and the Grenadines	2005	Male	50 to 54	68% (50-83%)	32% (17-50%)	2.7 (1.5-4.2)	10% (6.8-14%)	12% (8.1-17%)	2.7 (1.7-3.7)	110 (73-150)
Saint Vincent and the Grenadines	2005	Male	55 to 59	66% (49-81%)	34% (19-51%)	2.5 (1.3-4)	9.3% (5.9-13%)	11% (7.1-16%)	2.7 (1.7-3.8)	93 (58-130)
Saint Vincent and the Grenadines	2005	Male	60 to 64	64% (45-81%)	36% (19-55%)	2.2 (1.2-3.8)	7.6% (4.3-12%)	8.9% (5.1-14%)	2.4 (1.4-3.7)	71 (41-110)
Saint Vincent and the Grenadines	2005	Male	65 to 69	63% (44-79%)	37% (21-56%)	2 (1.1-3.2)	6.4% (3.5-10%)	7.4% (4.1-12%)	3 (1.7-4.8)	73 (40-110)
Saint Vincent and the Grenadines	2005	Male	70 to 74	60% (42-76%)	40% (24-58%)	1.7 (0.89-2.8)	4.6% (2-8.2%)	5.3% (2.3-9.2%)	2.5 (1.1-4.4)	48 (21-84)
Saint Vincent and the Grenadines	2005	Male	75 to 79	55% (35-73%)	45% (27-65%)	1.4 (0.68-2.3)	3.4% (1.2-6.3%)	3.8% (1.3-7.1%)	1.9 (0.65-3.6)	29 (10-53)
Saint Vincent and the Grenadines	2005	Male	80 to 84	49% (30-67%)	51% (33-70%)	1.2 (0.6-1.9)	2.5% (0.93-4.7%)	2.8% (0.93-5.2%)	1.2 (0.43-2.4)	14 (5.1-26)
Saint Vincent and the Grenadines	2005	Male	85 to 89	44% (28-61%)	56% (39-72%)	1 (0.54-1.7)	2% (0.72-4%)	2.2% (0.68-4.4%)	0.74 (0.23-1.5)	5.9 (2.1-12)
Saint Vincent and the Grenadines	2005	Male	90 to 94	44% (29-62%)	56% (38-71%)	1 (0.51-1.9)	1.7% (0.46-4%)	1.8% (0.29-4.4%)	0.29 (0.047-0.69)	1.7 (0.42-3.8)
Saint Vincent and the Grenadines	2005	Male	95 plus	47% (28-67%)	53% (33-72%)	1.1 (0.45-2.2)	1.8% (0.21-4.7%)	1.8% (-0.091-5.3%)	0.082 (-0.0042-0.24)	0.33 (0.04-0.87)
Saint Vincent and the Grenadines	2010	Female	All Ages	36% (29-43%)	64% (57-71%)	0.99 (0.79-1.2)	1.7% (0.94-2.6%)	1.6% (0.3-3.3%)	6.1 (1.1-12)	250 (130-380)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	2010	Female	15 to 19	41% (28-58%)	59% (42-72%)	1.1 (0.58-1.9)	2% (1.2-3%)	4% (2-6.8%)	0.11 (0.057-0.19)	12 (7.4-20)
Saint Vincent and the Grenadines	2010	Female	20 to 24	47% (31-64%)	53% (36-69%)	1.7 (0.89-2.6)	3.1% (2-4.4%)	4.9% (2.7-7.6%)	0.17 (0.09-0.27)	21 (13-30)
Saint Vincent and the Grenadines	2010	Female	25 to 29	47% (30-64%)	53% (36-70%)	1.7 (0.98-2.7)	3.2% (2.1-4.3%)	4.8% (2.8-7.4%)	0.2 (0.11-0.32)	22 (14-31)
Saint Vincent and the Grenadines	2010	Female	30 to 34	45% (29-63%)	55% (37-71%)	1.7 (0.95-2.6)	2.9% (2-4.1%)	4.6% (2.8-6.7%)	0.23 (0.14-0.36)	21 (13-29)
Saint Vincent and the Grenadines	2010	Female	35 to 39	44% (29-62%)	56% (38-71%)	1.5 (0.88-2.4)	2.8% (1.9-4.2%)	4.6% (3-6.9%)	0.3 (0.19-0.47)	21 (14-31)
Saint Vincent and the Grenadines	2010	Female	40 to 44	43% (28-61%)	57% (39-72%)	1.4 (0.76-2.3)	2.9% (1.6-4.5%)	4.8% (2.7-7.4%)	0.43 (0.24-0.68)	24 (13-38)
Saint Vincent and the Grenadines	2010	Female	45 to 49	41% (25-58%)	59% (42-75%)	1.4 (0.73-2.3)	2.9% (1.4-4.9%)	4.4% (2.2-7.5%)	0.66 (0.32-1.1)	31 (15-54)
Saint Vincent and the Grenadines	2010	Female	50 to 54	38% (23-55%)	62% (45-77%)	1.3 (0.68-2.1)	2.8% (0.95-5.3%)	4.1% (1.5-7.6%)	0.72 (0.27-1.3)	29 (9.7-53)
Saint Vincent and the Grenadines	2010	Female	55 to 59	33% (20-50%)	67% (50-80%)	1.1 (0.54-1.9)	2.2% (0.52-4.7%)	3.2% (0.95-6.3%)	0.63 (0.18-1.2)	21 (4.9-43)
Saint Vincent and the Grenadines	2010	Female	60 to 64	30% (18-45%)	70% (55-82%)	0.91 (0.45-1.6)	1.6% (-0.057-4.1%)	2.3% (0.11-5.4%)	0.49 (0.024-1.2)	14 (-0.46-34)
Saint Vincent and the Grenadines	2010	Female	65 to 69	27% (17-41%)	73% (59-83%)	0.8 (0.43-1.3)	1.1% (-0.47-3.2%)	1.5% (-0.46-4%)	0.36 (-0.11-0.99)	8.2 (-3.4-24)
Saint Vincent and the Grenadines	2010	Female	70 to 74	25% (15-38%)	75% (62-85%)	0.68 (0.33-1.2)	0.81% (-0.75-3%)	1.1% (-0.86-3.8%)	0.46 (-0.35-1.7)	8.3 (-7.7-31)
Saint Vincent and the Grenadines	2010	Female	75 to 79	22% (13-35%)	78% (65-87%)	0.6 (0.26-1.1)	0.57% (-0.82-2.7%)	0.71% (-0.95-3.2%)	0.33 (-0.46-1.5)	4.8 (-7.1-22)
Saint Vincent and the Grenadines	2010	Female	80 to 84	20% (12-32%)	80% (68-88%)	0.56 (0.27-0.98)	0.62% (-0.63-2.3%)	0.73% (-0.77-2.6%)	0.39 (-0.41-1.4)	4.2 (-4.3-15)
Saint Vincent and the Grenadines	2010	Female	85 to 89	20% (12-31%)	80% (69-88%)	0.56 (0.27-1)	0.59% (-0.61-2.3%)	0.64% (-0.76-2.6%)	0.32 (-0.37-1.3)	2.6 (-2.6-9.9)
Saint Vincent and the Grenadines	2010	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.64 (0.27-1.2)	0.74% (-0.63-2.9%)	0.78% (-0.82-3.4%)	0.22 (-0.23-0.98)	1.3 (-1.1-5.2)
Saint Vincent and the Grenadines	2010	Female	95 plus	26% (14-44%)	74% (56-86%)	0.76 (0.27-1.6)	1% (-0.75-3.9%)	1% (-1.1-4.5%)	0.095 (-0.1-0.43)	0.37 (-0.28-1.5)
Saint Vincent and the Grenadines	2010	Male	All Ages	59% (51-67%)	41% (33-49%)	2.1 (1.8-2.6)	7.7% (6.2-9.3%)	7.7% (5.6-10%)	34 (25-45)	1400 (1100-1700)
Saint Vincent and the Grenadines	2010	Male	15 to 19	58% (41-74%)	42% (26-59%)	1.6 (0.87-2.6)	6.2% (4-8.9%)	8.2% (4.9-12%)	0.69 (0.4-1.1)	60 (38-90)
Saint Vincent and the Grenadines	2010	Male	20 to 24	68% (51-82%)	32% (18-49%)	3 (1.8-4.5)	12% (8.4-16%)	14% (9-20%)	1.6 (1-2.4)	140 (93-190)
Saint Vincent and the Grenadines	2010	Male	25 to 29	71% (54-84%)	29% (16-46%)	3.6 (2.2-5.3)	12% (9.1-16%)	15% (9.9-21%)	1.7 (1.1-2.4)	130 (94-180)
Saint Vincent and the Grenadines	2010	Male	30 to 34	72% (54-85%)	28% (15-46%)	3.8 (2.4-5.5)	12% (8.9-15%)	15% (10-20%)	1.7 (1.2-2.4)	120 (90-160)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	2010	Male	35 to 39	72% (56-85%)	28% (15-44%)	3.5 (2.2-5.2)	11% (8.6-15%)	14% (10-19%)	1.9 (1.3-2.6)	120 (88-160)
Saint Vincent and the Grenadines	2010	Male	40 to 44	71% (54-84%)	29% (16-46%)	3.4 (1.9-5.1)	11% (8.3-15%)	14% (10-19%)	2.4 (1.6-3.2)	130 (95-170)
Saint Vincent and the Grenadines	2010	Male	45 to 49	69% (50-83%)	31% (17-50%)	3.2 (1.8-4.9)	11% (7.8-15%)	14% (9.3-18%)	3.4 (2.3-4.6)	160 (120-220)
Saint Vincent and the Grenadines	2010	Male	50 to 54	68% (49-82%)	32% (18-51%)	2.9 (1.6-4.6)	11% (7.1-15%)	13% (8.3-18%)	3.7 (2.4-5.1)	150 (100-210)
Saint Vincent and the Grenadines	2010	Male	55 to 59	66% (48-81%)	34% (19-52%)	2.7 (1.5-4.3)	10% (6.3-14%)	12% (7.6-17%)	3.6 (2.2-5.1)	120 (78-180)
Saint Vincent and the Grenadines	2010	Male	60 to 64	64% (44-81%)	36% (19-56%)	2.4 (1.3-4.2)	8.2% (4.6-13%)	9.7% (5.4-15%)	3.1 (1.7-4.9)	92 (51-140)
Saint Vincent and the Grenadines	2010	Male	65 to 69	63% (44-79%)	37% (21-56%)	2.2 (1.2-3.6)	7% (3.8-11%)	8.1% (4.3-13%)	2.8 (1.5-4.5)	67 (36-110)
Saint Vincent and the Grenadines	2010	Male	70 to 74	59% (40-75%)	41% (25-60%)	1.8 (0.96-3)	5.2% (2.4-9.2%)	5.9% (2.7-10%)	2.9 (1.3-5.4)	57 (26-100)
Saint Vincent and the Grenadines	2010	Male	75 to 79	54% (35-72%)	46% (28-65%)	1.5 (0.75-2.6)	3.8% (1.5-7.1%)	4.3% (1.6-7.9%)	2.1 (0.79-4)	32 (12-60)
Saint Vincent and the Grenadines	2010	Male	80 to 84	48% (30-66%)	52% (34-70%)	1.3 (0.65-2.1)	2.9% (1.1-5.6%)	3.2% (1.2-6.2%)	1.4 (0.49-2.7)	16 (5.8-30)
Saint Vincent and the Grenadines	2010	Male	85 to 89	44% (27-61%)	56% (39-73%)	1.1 (0.56-1.9)	2.4% (0.91-4.9%)	2.6% (0.88-5.4%)	0.84 (0.29-1.8)	6.7 (2.5-14)
Saint Vincent and the Grenadines	2010	Male	90 to 94	44% (28-61%)	56% (39-72%)	1.1 (0.53-2)	2% (0.57-4.6%)	2.1% (0.43-5%)	0.33 (0.067-0.79)	1.9 (0.53-4.2)
Saint Vincent and the Grenadines	2010	Male	95 plus	47% (27-67%)	53% (33-73%)	1.2 (0.49-2.3)	2.1% (0.31-5.3%)	2.1% (0.015-5.8%)	0.098 (0.00067-0.27)	0.39 (0.057-0.97)
Saint Vincent and the Grenadines	2016	Female	All Ages	35% (29-42%)	65% (58-71%)	1 (0.74-1.4)	1.9% (0.97-3.1%)	1.9% (0.44-3.8%)	7.6 (1.7-16)	280 (140-470)
Saint Vincent and the Grenadines	2016	Female	15 to 19	41% (28-58%)	59% (42-72%)	1.2 (0.55-2)	2% (1.1-3.2%)	4.1% (1.9-7.1%)	0.1 (0.047-0.18)	11 (6.4-18)
Saint Vincent and the Grenadines	2016	Female	20 to 24	47% (31-63%)	53% (37-69%)	1.7 (0.89-2.9)	3.2% (1.9-4.8%)	5.2% (2.6-8.7%)	0.17 (0.083-0.29)	20 (12-31)
Saint Vincent and the Grenadines	2016	Female	25 to 29	46% (30-64%)	54% (36-70%)	1.8 (0.96-2.8)	3.3% (2.1-4.7%)	5% (2.8-7.9%)	0.2 (0.1-0.35)	22 (13-33)
Saint Vincent and the Grenadines	2016	Female	30 to 34	44% (28-62%)	56% (38-72%)	1.7 (0.91-2.8)	3.1% (2-4.6%)	4.9% (2.9-7.5%)	0.27 (0.15-0.45)	23 (14-35)
Saint Vincent and the Grenadines	2016	Female	35 to 39	44% (28-61%)	56% (39-72%)	1.6 (0.86-2.7)	3% (1.9-4.5%)	4.9% (2.9-7.4%)	0.35 (0.2-0.58)	24 (15-38)
Saint Vincent and the Grenadines	2016	Female	40 to 44	42% (27-61%)	58% (39-73%)	1.5 (0.79-2.5)	3.1% (1.6-5.1%)	5.1% (2.7-8.3%)	0.48 (0.25-0.8)	27 (14-43)
Saint Vincent and the Grenadines	2016	Female	45 to 49	40% (25-58%)	60% (42-75%)	1.4 (0.74-2.5)	3.1% (1.4-5.5%)	4.7% (2.4-8.2%)	0.62 (0.31-1.1)	29 (14-51)
Saint Vincent and the Grenadines	2016	Female	50 to 54	37% (23-55%)	63% (45-77%)	1.3 (0.68-2.3)	3% (1.1-5.8%)	4.4% (1.8-8.1%)	0.92 (0.36-1.7)	37 (13-70)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	2016	Female	55 to 59	32% (20-49%)	68% (51-80%)	1.1 (0.54-1.9)	2.4% (0.73-4.9%)	3.5% (1.2-6.6%)	0.89 (0.3-1.7)	30 (9.1-60)
Saint Vincent and the Grenadines	2016	Female	60 to 64	29% (18-45%)	71% (55-82%)	0.94 (0.46-1.7)	1.8% (0.11-4.3%)	2.5% (0.33-5.6%)	0.71 (0.092-1.6)	20 (1.3-50)
Saint Vincent and the Grenadines	2016	Female	65 to 69	27% (17-41%)	73% (59-83%)	0.82 (0.44-1.4)	1.3% (-0.28-3.7%)	1.8% (-0.22-4.6%)	0.55 (-0.072-1.5)	13 (-2.6-35)
Saint Vincent and the Grenadines	2016	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.69 (0.33-1.3)	1% (-0.65-3.3%)	1.3% (-0.67-4.1%)	0.43 (-0.21-1.4)	7.8 (-4.7-26)
Saint Vincent and the Grenadines	2016	Female	75 to 79	22% (13-35%)	78% (65-87%)	0.61 (0.26-1.2)	0.74% (-0.73-2.9%)	0.91% (-0.87-3.5%)	0.48 (-0.45-1.9)	6.9 (-6.9-28)
Saint Vincent and the Grenadines	2016	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.57 (0.26-1.1)	0.78% (-0.56-2.8%)	0.92% (-0.66-3.3%)	0.57 (-0.4-2.1)	6.1 (-4.2-22)
Saint Vincent and the Grenadines	2016	Female	85 to 89	20% (11-31%)	80% (69-89%)	0.57 (0.26-1.1)	0.79% (-0.47-2.7%)	0.88% (-0.59-3.1%)	0.42 (-0.27-1.5)	3.3 (-1.9-11)
Saint Vincent and the Grenadines	2016	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.65 (0.28-1.3)	0.91% (-0.5-3.3%)	0.99% (-0.7-3.7%)	0.29 (-0.2-1.1)	1.6 (-0.89-5.7)
Saint Vincent and the Grenadines	2016	Female	95 plus	26% (14-43%)	74% (57-86%)	0.79 (0.28-1.6)	1.2% (-0.68-4.5%)	1.2% (-0.96-5.1%)	0.13 (-0.1-0.58)	0.51 (-0.29-2)
Saint Vincent and the Grenadines	2016	Male	All Ages	59% (51-67%)	41% (33-49%)	2.3 (1.7-3)	8.2% (6.3-10%)	8.2% (5.6-11%)	42 (28-57)	1600 (1200-2100)
Saint Vincent and the Grenadines	2016	Male	15 to 19	58% (40-74%)	42% (26-60%)	1.7 (0.82-2.8)	6.3% (3.9-9.3%)	8.3% (4.8-13%)	0.63 (0.33-1)	55 (33-86)
Saint Vincent and the Grenadines	2016	Male	20 to 24	68% (50-82%)	32% (18-50%)	3.1 (1.7-5)	12% (8.3-17%)	14% (8.9-21%)	1.6 (0.92-2.5)	130 (86-200)
Saint Vincent and the Grenadines	2016	Male	25 to 29	71% (53-84%)	29% (16-47%)	3.7 (2.2-5.8)	13% (8.9-17%)	16% (9.8-22%)	1.7 (1-2.6)	130 (87-190)
Saint Vincent and the Grenadines	2016	Male	30 to 34	72% (54-85%)	28% (15-46%)	3.9 (2.3-6.1)	12% (8.9-16%)	15% (10-21%)	1.9 (1.2-2.7)	130 (89-180)
Saint Vincent and the Grenadines	2016	Male	35 to 39	71% (55-85%)	29% (15-45%)	3.7 (2.1-5.7)	12% (8.4-15%)	15% (10-20%)	2.1 (1.3-2.9)	130 (91-180)
Saint Vincent and the Grenadines	2016	Male	40 to 44	70% (54-84%)	30% (16-46%)	3.5 (1.9-5.5)	12% (8.4-16%)	15% (10-21%)	2.6 (1.7-3.6)	140 (96-190)
Saint Vincent and the Grenadines	2016	Male	45 to 49	69% (48-83%)	31% (17-52%)	3.3 (1.8-5.3)	12% (8-16%)	15% (9.4-20%)	3.4 (2.2-4.7)	160 (110-220)
Saint Vincent and the Grenadines	2016	Male	50 to 54	67% (49-83%)	33% (17-51%)	3 (1.7-4.9)	11% (7.4-15%)	14% (8.9-19%)	5 (3.2-7.1)	200 (130-290)
Saint Vincent and the Grenadines	2016	Male	55 to 59	65% (48-81%)	35% (19-52%)	2.8 (1.5-4.6)	10% (6.3-14%)	12% (7.6-17%)	5.5 (3.3-7.9)	190 (120-270)
Saint Vincent and the Grenadines	2016	Male	60 to 64	63% (44-81%)	37% (19-56%)	2.5 (1.3-4.2)	8.4% (4.6-13%)	9.9% (5.3-15%)	4.6 (2.4-7.1)	130 (70-210)
Saint Vincent and the Grenadines	2016	Male	65 to 69	62% (43-78%)	38% (22-57%)	2.2 (1.2-3.6)	7.1% (3.8-11%)	8.2% (4.4-13%)	3.8 (2-6.1)	92 (48-140)
Saint Vincent and the Grenadines	2016	Male	70 to 74	58% (41-74%)	42% (26-59%)	1.9 (0.94-3.3)	5.4% (2.4-9.5%)	6.1% (2.6-11%)	2.8 (1.2-5.1)	54 (24-95)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saint Vincent and the Grenadines	2016	Male	75 to 79	53% (35-71%)	47% (29-65%)	1.6 (0.74-2.7)	4% (1.5-7.7%)	4.5% (1.6-8.7%)	2.5 (0.9-5)	38 (14-73)
Saint Vincent and the Grenadines	2016	Male	80 to 84	48% (29-66%)	52% (34-71%)	1.3 (0.64-2.3)	3.1% (1.1-6.1%)	3.4% (1.2-6.8%)	1.9 (0.67-3.8)	21 (7.7-41)
Saint Vincent and the Grenadines	2016	Male	85 to 89	43% (27-60%)	57% (40-73%)	1.2 (0.56-2.1)	2.6% (0.87-5.3%)	2.8% (0.86-6%)	1 (0.32-2.2)	8.1 (2.8-17)
Saint Vincent and the Grenadines	2016	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.2 (0.52-2.2)	2.2% (0.57-5%)	2.3% (0.44-5.5%)	0.41 (0.079-1)	2.4 (0.62-5.4)
Saint Vincent and the Grenadines	2016	Male	95 plus	46% (27-66%)	54% (34-73%)	1.2 (0.49-2.5)	2.3% (0.3-5.6%)	2.3% (0.011-6.4%)	0.12 (0.00057-0.33)	0.47 (0.063-1.2)
Suriname	1990	Female	All Ages	33% (27-40%)	67% (60-73%)	0.91 (0.75-1.1)	1.7% (1.2-2.3%)	2.2% (1.2-3.4%)	25 (14-39)	1100 (770-1400)
Suriname	1990	Female	15 to 19	40% (26-56%)	60% (44-74%)	1.1 (0.58-1.8)	2.5% (1.4-4.1%)	5% (2.2-8.5%)	0.76 (0.34-1.3)	72 (39-120)
Suriname	1990	Female	20 to 24	46% (30-62%)	54% (38-70%)	1.7 (0.95-2.5)	4% (2.5-5.8%)	6.9% (3.7-11%)	1.3 (0.67-2)	120 (76-180)
Suriname	1990	Female	25 to 29	45% (29-62%)	55% (38-71%)	1.7 (1.1-2.5)	4% (2.7-5.4%)	6.6% (4-9.7%)	1.3 (0.82-2)	120 (84-170)
Suriname	1990	Female	30 to 34	43% (27-61%)	57% (39-73%)	1.6 (0.95-2.5)	3.8% (2.6-5.3%)	6.5% (4.1-9.7%)	1.3 (0.85-1.9)	100 (71-150)
Suriname	1990	Female	35 to 39	43% (28-60%)	57% (40-72%)	1.5 (0.85-2.5)	3.7% (2.4-5.3%)	6.5% (4-9.4%)	1.4 (0.86-2)	91 (59-130)
Suriname	1990	Female	40 to 44	41% (27-60%)	59% (40-73%)	1.5 (0.73-2.4)	3.7% (2.2-5.6%)	6.4% (3.7-9.8%)	1.5 (0.87-2.4)	85 (51-130)
Suriname	1990	Female	45 to 49	39% (24-57%)	61% (43-76%)	1.4 (0.75-2.3)	3.7% (2-5.9%)	6% (3.3-9.7%)	1.9 (1-3.1)	89 (49-150)
Suriname	1990	Female	50 to 54	36% (22-52%)	64% (48-78%)	1.3 (0.66-2.1)	3.6% (1.8-6.2%)	5.5% (2.7-9.2%)	2.5 (1.2-4.2)	100 (51-170)
Suriname	1990	Female	55 to 59	32% (19-48%)	68% (52-81%)	1.1 (0.53-1.8)	3.2% (1.4-5.6%)	4.5% (2-7.9%)	2.9 (1.3-5.2)	98 (43-180)
Suriname	1990	Female	60 to 64	28% (17-44%)	72% (56-83%)	0.91 (0.44-1.6)	2.4% (0.68-4.7%)	3.2% (0.92-6.3%)	2.3 (0.67-4.5)	67 (20-130)
Suriname	1990	Female	65 to 69	26% (16-40%)	74% (60-84%)	0.79 (0.41-1.4)	1.9% (0.47-3.8%)	2.5% (0.61-5%)	2 (0.49-4.1)	47 (12-95)
Suriname	1990	Female	70 to 74	23% (14-36%)	77% (64-86%)	0.67 (0.34-1.2)	1.4% (0.022-3.4%)	1.8% (0.038-4.3%)	1.5 (0.034-3.6)	28 (0.41-67)
Suriname	1990	Female	75 to 79	21% (12-33%)	79% (67-88%)	0.57 (0.26-1.1)	1.1% (-0.18-2.8%)	1.3% (-0.27-3.4%)	1.2 (-0.24-3.1)	17 (-3.1-47)
Suriname	1990	Female	80 to 84	19% (11-30%)	81% (70-89%)	0.54 (0.25-0.9)	0.94% (-0.078-2.3%)	1.1% (-0.14-2.8%)	1.1 (-0.14-2.9)	12 (-1.1-31)
Suriname	1990	Female	85 to 89	19% (11-30%)	81% (70-89%)	0.55 (0.27-0.96)	0.88% (-0.084-2.4%)	0.99% (-0.17-2.7%)	0.94 (-0.16-2.6)	7.5 (-0.72-20)
Suriname	1990	Female	90 to 94	21% (12-33%)	79% (67-88%)	0.63 (0.27-1.2)	0.97% (-0.14-2.8%)	1.1% (-0.24-3.3%)	0.56 (-0.13-1.7)	3.1 (-0.42-9.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	1990	Female	95 plus	25% (13-42%)	75% (58-87%)	0.75 (0.27-1.5)	1.2% (-0.39-3.9%)	1.3% (-0.63-4.5%)	0.17 (-0.088-0.62)	0.66 (-0.22-2.1)
Suriname	1990	Male	All Ages	54% (46-63%)	46% (37-54%)	2 (1.6-2.3)	5.2% (4-6.5%)	6.6% (4.5-8.9%)	92 (63-120)	4100 (3200-5100)
Suriname	1990	Male	15 to 19	57% (40-73%)	43% (27-60%)	1.7 (0.92-2.6)	5.6% (3.5-8.3%)	8% (4.5-12%)	2 (1.1-3.2)	190 (110-270)
Suriname	1990	Male	20 to 24	67% (50-81%)	33% (19-50%)	3.1 (1.9-4.5)	12% (8-16%)	15% (9.3-21%)	6.3 (3.9-9)	520 (360-710)
Suriname	1990	Male	25 to 29	70% (53-83%)	30% (17-47%)	3.7 (2.4-5.2)	13% (9.4-17%)	17% (12-23%)	7.4 (5-10)	560 (400-750)
Suriname	1990	Male	30 to 34	71% (53-84%)	29% (16-47%)	3.9 (2.5-5.6)	13% (9.3-17%)	17% (12-23%)	6.3 (4.3-8.5)	430 (310-570)
Suriname	1990	Male	35 to 39	71% (55-84%)	29% (16-45%)	3.7 (2.2-5.6)	13% (8.4-17%)	16% (10-22%)	6.9 (4.3-9.4)	410 (270-550)
Suriname	1990	Male	40 to 44	70% (54-83%)	30% (17-46%)	3.5 (1.9-5.4)	12% (7.7-16%)	15% (9.5-22%)	6.3 (3.9-8.9)	340 (220-460)
Suriname	1990	Male	45 to 49	68% (48-83%)	32% (17-52%)	3.3 (1.9-5.4)	11% (7.1-15%)	14% (8.7-20%)	7 (4.2-9.9)	330 (210-450)
Suriname	1990	Male	50 to 54	67% (48-82%)	33% (18-52%)	3 (1.7-5)	9.8% (5.5-14%)	12% (6.6-18%)	7.9 (4.2-12)	320 (180-470)
Suriname	1990	Male	55 to 59	65% (46-80%)	35% (20-54%)	2.8 (1.5-4.5)	9% (4.9-13%)	11% (5.8-17%)	9.6 (4.9-15)	330 (170-500)
Suriname	1990	Male	60 to 64	63% (43-79%)	37% (21-57%)	2.5 (1.3-4.4)	7.6% (3.8-12%)	9% (4.5-15%)	8.8 (4.3-15)	260 (130-410)
Suriname	1990	Male	65 to 69	61% (42-78%)	39% (22-58%)	2.2 (1.2-3.6)	6.5% (3.1-11%)	7.7% (3.6-13%)	7.7 (3.7-13)	190 (92-310)
Suriname	1990	Male	70 to 74	58% (39-74%)	42% (26-61%)	1.8 (0.95-3.1)	4.9% (2-9.4%)	5.7% (2.2-11%)	5.4 (2.1-10)	100 (42-200)
Suriname	1990	Male	75 to 79	53% (34-71%)	47% (29-66%)	1.5 (0.74-2.7)	3.8% (1.3-7.3%)	4.3% (1.4-8.4%)	4.3 (1.4-8.4)	64 (22-120)
Suriname	1990	Male	80 to 84	47% (29-65%)	53% (35-71%)	1.3 (0.66-2.2)	2.9% (0.95-5.6%)	3.2% (0.95-6.4%)	2.9 (0.85-5.7)	32 (10-63)
Suriname	1990	Male	85 to 89	42% (26-59%)	58% (41-74%)	1.1 (0.56-2)	2.5% (0.81-4.9%)	2.7% (0.77-5.6%)	1.8 (0.52-3.8)	15 (4.8-30)
Suriname	1990	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.2 (0.54-2.1)	2.2% (0.53-5.1%)	2.3% (0.41-5.7%)	0.85 (0.15-2.1)	4.8 (1.2-11)
Suriname	1990	Male	95 plus	46% (26-66%)	54% (34-74%)	1.3 (0.52-2.4)	2.4% (0.31-5.9%)	2.4% (0.032-6.6%)	0.22 (0.0029-0.59)	0.84 (0.11-2.1)
Suriname	1995	Female	All Ages	32% (26-40%)	68% (60-74%)	0.85 (0.71-1)	1.6% (1.1-2.1%)	2% (1.1-3.3%)	25 (13-41)	1100 (750-1500)
Suriname	1995	Female	15 to 19	39% (26-56%)	61% (44-74%)	1 (0.53-1.7)	2.3% (1.2-3.7%)	4.6% (2-8.4%)	0.6 (0.25-1.1)	59 (31-95)
Suriname	1995	Female	20 to 24	45% (30-62%)	55% (38-70%)	1.6 (0.86-2.4)	3.6% (2.2-5.3%)	6.2% (3.1-9.9%)	1.1 (0.55-1.8)	110 (68-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	1995	Female	25 to 29	45% (28-62%)	55% (38-72%)	1.6 (0.97-2.4)	3.5% (2.4-4.9%)	5.8% (3.4-8.7%)	1.2 (0.68-1.8)	110 (72-150)
Suriname	1995	Female	30 to 34	43% (27-61%)	57% (39-73%)	1.6 (0.92-2.3)	3.4% (2.3-4.6%)	5.7% (3.6-8.2%)	1.4 (0.91-2.1)	120 (79-160)
Suriname	1995	Female	35 to 39	42% (27-60%)	58% (40-73%)	1.4 (0.81-2.3)	3.4% (2.2-4.8%)	5.7% (3.6-8.2%)	1.6 (0.99-2.3)	110 (69-150)
Suriname	1995	Female	40 to 44	41% (27-60%)	59% (40-73%)	1.4 (0.72-2.2)	3.4% (2-5.1%)	5.8% (3.4-8.8%)	1.8 (1-2.7)	97 (59-140)
Suriname	1995	Female	45 to 49	39% (24-56%)	61% (44-76%)	1.3 (0.69-2.2)	3.3% (1.8-5.5%)	5.4% (2.9-8.8%)	1.9 (1-3.1)	89 (48-150)
Suriname	1995	Female	50 to 54	36% (22-53%)	64% (47-78%)	1.2 (0.61-2)	3.3% (1.6-5.7%)	5% (2.4-8.6%)	2.3 (1.1-4.1)	93 (44-160)
Suriname	1995	Female	55 to 59	31% (19-48%)	69% (52-81%)	1 (0.5-1.7)	2.8% (1.1-5.1%)	4.1% (1.7-7.4%)	2.6 (1-4.7)	89 (35-160)
Suriname	1995	Female	60 to 64	28% (16-44%)	72% (56-84%)	0.85 (0.41-1.5)	2.2% (0.58-4.2%)	3% (0.78-5.7%)	2.5 (0.66-4.8)	72 (19-140)
Suriname	1995	Female	65 to 69	26% (16-39%)	74% (61-84%)	0.74 (0.38-1.3)	1.7% (0.33-3.6%)	2.3% (0.43-4.7%)	2.2 (0.43-4.6)	52 (10-110)
Suriname	1995	Female	70 to 74	23% (15-36%)	77% (64-85%)	0.62 (0.31-1.1)	1.3% (-0.034-3.1%)	1.7% (-0.015-4%)	1.9 (-0.015-4.4)	35 (-0.86-83)
Suriname	1995	Female	75 to 79	21% (12-33%)	79% (67-88%)	0.54 (0.24-1)	0.96% (-0.23-2.7%)	1.2% (-0.28-3.3%)	1.2 (-0.29-3.4)	18 (-4.3-49)
Suriname	1995	Female	80 to 84	19% (11-31%)	81% (69-89%)	0.51 (0.24-0.88)	0.9% (-0.11-2.4%)	1.1% (-0.18-2.9%)	1.2 (-0.19-3.1)	13 (-1.7-33)
Suriname	1995	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.52 (0.25-0.93)	0.86% (-0.099-2.3%)	0.98% (-0.14-2.7%)	0.96 (-0.14-2.6)	7.6 (-0.89-20)
Suriname	1995	Female	90 to 94	21% (12-33%)	79% (67-88%)	0.59 (0.26-1.2)	0.94% (-0.14-2.9%)	1% (-0.24-3.3%)	0.62 (-0.15-2)	3.5 (-0.48-10)
Suriname	1995	Female	95 plus	25% (13-41%)	75% (59-87%)	0.71 (0.26-1.4)	1.1% (-0.37-3.7%)	1.2% (-0.61-4.3%)	0.26 (-0.13-0.94)	0.97 (-0.32-3.2)
Suriname	1995	Male	All Ages	54% (46-63%)	46% (37-54%)	1.8 (1.5-2.2)	5.1% (3.9-6.4%)	6.4% (4.3-8.6%)	97 (67-130)	4300 (3400-5300)
Suriname	1995	Male	15 to 19	57% (39-73%)	43% (27-61%)	1.5 (0.8-2.4)	5.4% (3.3-8%)	7.5% (4-12%)	1.8 (0.95-2.9)	160 (99-250)
Suriname	1995	Male	20 to 24	67% (50-81%)	33% (19-50%)	2.9 (1.7-4.2)	11% (7.4-15%)	14% (8.3-20%)	5.7 (3.4-8.3)	480 (320-650)
Suriname	1995	Male	25 to 29	70% (53-83%)	30% (17-47%)	3.5 (2.2-4.9)	12% (8.7-16%)	16% (10-22%)	7.3 (4.8-10)	560 (390-740)
Suriname	1995	Male	30 to 34	71% (53-84%)	29% (16-47%)	3.6 (2.3-5.1)	12% (8.3-15%)	15% (10-21%)	8.3 (5.6-11)	560 (400-750)
Suriname	1995	Male	35 to 39	71% (54-84%)	29% (16-46%)	3.4 (2-5)	11% (7.4-15%)	14% (9.1-20%)	7.6 (4.8-11)	460 (310-610)
Suriname	1995	Male	40 to 44	70% (54-83%)	30% (17-46%)	3.2 (1.8-4.9)	11% (6.8-15%)	14% (8.4-19%)	7 (4.3-10)	380 (250-520)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	1995	Male	45 to 49	68% (48-82%)	32% (18-52%)	3.1 (1.7-4.9)	10% (6.6-15%)	13% (7.8-19%)	7.1 (4.3-10)	330 (210-480)
Suriname	1995	Male	50 to 54	66% (48-82%)	34% (18-52%)	2.8 (1.5-4.7)	9.3% (5.2-14%)	12% (6.2-17%)	7.9 (4.2-12)	320 (180-470)
Suriname	1995	Male	55 to 59	65% (46-80%)	35% (20-54%)	2.6 (1.4-4.1)	8.6% (4.7-13%)	11% (5.7-16%)	9.2 (4.9-14)	320 (170-490)
Suriname	1995	Male	60 to 64	62% (42-79%)	38% (21-58%)	2.3 (1.2-4)	7.3% (3.6-12%)	8.7% (4.2-14%)	9.7 (4.6-16)	280 (140-450)
Suriname	1995	Male	65 to 69	61% (41-77%)	39% (23-59%)	2 (1.1-3.4)	6.1% (3-10%)	7.3% (3.5-12%)	8.7 (4.2-15)	210 (100-350)
Suriname	1995	Male	70 to 74	57% (40-74%)	43% (26-60%)	1.7 (0.91-3)	4.7% (1.9-8.8%)	5.4% (2.1-10%)	6.7 (2.6-13)	130 (52-240)
Suriname	1995	Male	75 to 79	52% (33-71%)	48% (29-67%)	1.4 (0.66-2.6)	3.6% (1.2-7.1%)	4.1% (1.3-8.2%)	4 (1.3-8)	60 (21-120)
Suriname	1995	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.2 (0.59-2)	2.7% (0.95-5.3%)	3% (0.98-6%)	2.7 (0.89-5.3)	30 (11-59)
Suriname	1995	Male	85 to 89	42% (25-59%)	58% (41-75%)	1.1 (0.52-1.9)	2.3% (0.75-4.5%)	2.5% (0.76-5.1%)	1.8 (0.56-3.8)	15 (5-30)
Suriname	1995	Male	90 to 94	42% (27-59%)	58% (41-73%)	1.1 (0.48-2)	2% (0.5-4.7%)	2.1% (0.4-5.3%)	0.79 (0.15-2)	4.5 (1.1-11)
Suriname	1995	Male	95 plus	46% (26-66%)	54% (34-74%)	1.2 (0.46-2.3)	2.1% (0.3-5.4%)	2.2% (0.013-6%)	0.26 (0.0016-0.73)	1 (0.15-2.6)
Suriname	2000	Female	All Ages	32% (27-39%)	68% (61-73%)	0.95 (0.79-1.1)	1.8% (1.3-2.5%)	2.5% (1.3-4%)	36 (19-58)	1300 (920-1900)
Suriname	2000	Female	15 to 19	39% (26-55%)	61% (45-74%)	1.2 (0.59-2)	2.5% (1.3-4.1%)	5.1% (2.2-8.8%)	0.77 (0.33-1.4)	73 (39-120)
Suriname	2000	Female	20 to 24	45% (29-62%)	55% (38-71%)	1.7 (0.95-2.7)	3.7% (2.2-5.4%)	6.3% (3.2-10%)	1.1 (0.51-1.7)	110 (62-160)
Suriname	2000	Female	25 to 29	44% (28-62%)	56% (38-72%)	1.8 (1.1-2.7)	3.4% (2.3-4.7%)	5.3% (3-7.9%)	1.1 (0.62-1.7)	110 (71-150)
Suriname	2000	Female	30 to 34	43% (27-61%)	57% (39-73%)	1.7 (1-2.6)	3.3% (2.3-4.6%)	5.4% (3.5-7.9%)	1.4 (0.87-2)	110 (77-160)
Suriname	2000	Female	35 to 39	42% (27-60%)	58% (40-73%)	1.6 (0.93-2.5)	3.3% (2.1-4.7%)	5.5% (3.5-7.9%)	2 (1.2-2.8)	130 (85-180)
Suriname	2000	Female	40 to 44	41% (26-59%)	59% (41-74%)	1.5 (0.8-2.4)	3.5% (2.1-5.2%)	5.8% (3.4-8.7%)	2.3 (1.4-3.5)	130 (78-190)
Suriname	2000	Female	45 to 49	39% (24-56%)	61% (44-76%)	1.5 (0.78-2.4)	3.6% (1.9-5.8%)	5.7% (3.1-9.2%)	2.6 (1.4-4.2)	120 (66-200)
Suriname	2000	Female	50 to 54	35% (21-51%)	65% (49-79%)	1.3 (0.7-2.3)	3.7% (1.7-6.3%)	5.5% (2.6-9.4%)	3 (1.4-5.1)	120 (55-200)
Suriname	2000	Female	55 to 59	31% (18-47%)	69% (53-82%)	1.1 (0.54-1.9)	3.3% (1.4-5.9%)	4.8% (2.1-8.6%)	3.4 (1.5-6)	110 (50-200)
Suriname	2000	Female	60 to 64	28% (16-43%)	72% (57-84%)	0.94 (0.46-1.7)	2.7% (0.84-5.2%)	3.7% (1.2-6.8%)	3.6 (1.1-6.8)	100 (31-190)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	2000	Female	65 to 69	25% (16-39%)	75% (61-84%)	0.83 (0.43-1.4)	2.3% (0.57-4.5%)	3% (0.77-5.8%)	3.8 (0.97-7.4)	87 (22-170)
Suriname	2000	Female	70 to 74	23% (14-36%)	77% (64-86%)	0.7 (0.35-1.2)	1.8% (0.25-4.2%)	2.3% (0.32-5.1%)	3.4 (0.46-7.3)	62 (8-140)
Suriname	2000	Female	75 to 79	20% (12-33%)	80% (67-88%)	0.61 (0.27-1.1)	1.4% (0.025-3.4%)	1.7% (0.024-4.3%)	2.6 (0.035-6.3)	37 (0.66-91)
Suriname	2000	Female	80 to 84	19% (11-30%)	81% (70-89%)	0.57 (0.26-1)	1.3% (0.14-3.1%)	1.6% (0.17-3.8%)	2.2 (0.24-5.2)	24 (2.7-55)
Suriname	2000	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.58 (0.27-1.1)	1.3% (0.17-3%)	1.5% (0.17-3.5%)	1.5 (0.17-3.7)	12 (1.6-28)
Suriname	2000	Female	90 to 94	21% (12-32%)	79% (68-88%)	0.66 (0.3-1.3)	1.4% (0.089-3.6%)	1.5% (0.04-4.2%)	0.95 (0.025-2.6)	5.1 (0.35-13)
Suriname	2000	Female	95 plus	25% (13-41%)	75% (59-87%)	0.79 (0.28-1.6)	1.7% (-0.14-4.6%)	1.7% (-0.32-5.2%)	0.42 (-0.08-1.3)	1.5 (-0.13-4.5)
Suriname	2000	Male	All Ages	54% (46-63%)	46% (37-54%)	2.1 (1.7-2.5)	5.9% (4.6-7.3%)	7.4% (5.2-9.7%)	130 (89-170)	5400 (4200-6500)
Suriname	2000	Male	15 to 19	56% (38-73%)	44% (27-62%)	1.7 (0.9-2.8)	5.9% (3.5-9%)	8.3% (4.4-13%)	2.5 (1.3-4)	220 (130-340)
Suriname	2000	Male	20 to 24	67% (49-81%)	33% (19-51%)	3.2 (1.9-4.7)	12% (8.1-17%)	16% (9.5-22%)	6.2 (3.7-8.8)	500 (330-680)
Suriname	2000	Male	25 to 29	70% (52-83%)	30% (17-48%)	3.9 (2.4-5.5)	13% (8.8-17%)	16% (11-22%)	8 (5.2-11)	600 (420-800)
Suriname	2000	Male	30 to 34	71% (52-84%)	29% (16-48%)	4 (2.6-5.7)	12% (8.6-15%)	15% (11-21%)	8.7 (6-12)	590 (430-770)
Suriname	2000	Male	35 to 39	70% (54-84%)	30% (16-46%)	3.8 (2.4-5.6)	11% (7.7-15%)	14% (9.6-19%)	10 (6.7-14)	620 (440-800)
Suriname	2000	Male	40 to 44	69% (53-83%)	31% (17-47%)	3.7 (2.1-5.6)	11% (7.3-15%)	14% (8.9-19%)	9.7 (6.1-13)	520 (350-690)
Suriname	2000	Male	45 to 49	68% (47-82%)	32% (18-53%)	3.4 (1.9-5.5)	11% (7-15%)	14% (8.3-19%)	9.8 (6-14)	460 (300-640)
Suriname	2000	Male	50 to 54	66% (47-81%)	34% (19-53%)	3.2 (1.7-5.1)	10% (5.8-15%)	13% (6.9-18%)	10 (5.5-15)	410 (230-580)
Suriname	2000	Male	55 to 59	64% (46-80%)	36% (20-54%)	2.9 (1.6-4.7)	9.7% (5.4-14%)	12% (6.4-18%)	12 (6.3-17)	400 (220-580)
Suriname	2000	Male	60 to 64	62% (42-79%)	38% (21-58%)	2.6 (1.4-4.4)	8.5% (4.4-13%)	10% (5.2-16%)	13 (6.4-20)	360 (190-560)
Suriname	2000	Male	65 to 69	61% (41-77%)	39% (23-59%)	2.3 (1.3-3.7)	7.4% (3.8-12%)	8.8% (4.5-14%)	13 (6.6-21)	310 (160-480)
Suriname	2000	Male	70 to 74	57% (38-73%)	43% (27-62%)	1.9 (0.98-3.3)	5.7% (2.3-10%)	6.6% (2.7-12%)	10 (4.1-19)	200 (83-350)
Suriname	2000	Male	75 to 79	52% (32-70%)	48% (30-68%)	1.6 (0.74-2.8)	4.4% (1.6-8.2%)	5% (1.8-9.5%)	7.1 (2.5-13)	100 (37-190)
Suriname	2000	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.3 (0.64-2.2)	3.5% (1.3-6.5%)	3.9% (1.4-7.4%)	4.1 (1.4-7.8)	45 (16-84)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	2000	Male	85 to 89	41% (25-59%)	59% (41-75%)	1.2 (0.58-2)	2.8% (1.1-5.5%)	3.1% (1.1-6.2%)	2.3 (0.81-4.7)	18 (6.9-36)
Suriname	2000	Male	90 to 94	42% (27-59%)	58% (41-73%)	1.2 (0.54-2.2)	2.5% (0.83-5.5%)	2.7% (0.75-6.1%)	1.1 (0.31-2.5)	6.2 (2.1-13)
Suriname	2000	Male	95 plus	45% (26-66%)	55% (34-74%)	1.3 (0.51-2.5)	2.7% (0.67-6.3%)	2.8% (0.48-7%)	0.35 (0.059-0.88)	1.3 (0.33-3.1)
Suriname	2005	Female	All Ages	32% (27-40%)	68% (60-73%)	1.2 (0.95-1.4)	2.3% (1.6-3.2%)	3.3% (1.8-5.1%)	52 (29-81)	1800 (1200-2400)
Suriname	2005	Female	15 to 19	39% (26-56%)	61% (44-74%)	1.4 (0.7-2.3)	2.9% (1.5-4.8%)	6.2% (2.8-11%)	0.99 (0.45-1.7)	92 (48-150)
Suriname	2005	Female	20 to 24	45% (29-61%)	55% (39-71%)	2 (1.1-3.3)	4% (2.4-5.8%)	7.1% (3.6-11%)	1.3 (0.68-2.1)	130 (81-190)
Suriname	2005	Female	25 to 29	44% (28-61%)	56% (39-72%)	2.1 (1.2-3.2)	3.5% (2.3-4.8%)	5.4% (3.1-8.2%)	1 (0.58-1.6)	100 (67-150)
Suriname	2005	Female	30 to 34	42% (26-60%)	58% (40-74%)	2 (1.2-3.2)	3.2% (2.2-4.5%)	5.1% (3.2-7.6%)	1.3 (0.83-2)	110 (79-160)
Suriname	2005	Female	35 to 39	42% (27-59%)	58% (41-73%)	1.9 (1.1-2.9)	3.6% (2.4-5%)	6% (3.8-8.5%)	2 (1.3-3)	140 (90-190)
Suriname	2005	Female	40 to 44	41% (26-59%)	59% (41-74%)	1.8 (0.94-2.9)	3.8% (2.3-5.6%)	6.4% (3.8-9.4%)	3.2 (1.9-4.7)	170 (110-250)
Suriname	2005	Female	45 to 49	39% (23-56%)	61% (44-77%)	1.7 (0.9-2.9)	4.1% (2.3-6.6%)	6.6% (3.7-10%)	3.7 (2.1-6)	170 (99-280)
Suriname	2005	Female	50 to 54	35% (21-52%)	65% (48-79%)	1.5 (0.79-2.7)	4.4% (2.2-7.4%)	6.6% (3.4-11%)	4.4 (2.2-7.3)	170 (87-290)
Suriname	2005	Female	55 to 59	31% (18-47%)	69% (53-82%)	1.3 (0.65-2.3)	4.1% (1.9-7.1%)	5.9% (2.8-10%)	4.6 (2.2-7.7)	150 (71-270)
Suriname	2005	Female	60 to 64	27% (16-43%)	73% (57-84%)	1.1 (0.54-2)	3.5% (1.4-6.5%)	4.7% (1.8-8.5%)	4.9 (1.9-8.9)	140 (53-260)
Suriname	2005	Female	65 to 69	25% (16-39%)	75% (61-84%)	0.97 (0.52-1.6)	3.1% (1.1-5.6%)	4% (1.5-7.2%)	5.6 (2.1-10)	130 (47-240)
Suriname	2005	Female	70 to 74	23% (14-36%)	77% (64-86%)	0.81 (0.41-1.5)	2.5% (0.68-5.2%)	3.2% (0.89-6.5%)	5.7 (1.5-12)	110 (28-220)
Suriname	2005	Female	75 to 79	20% (12-33%)	80% (67-88%)	0.71 (0.32-1.3)	2% (0.39-4.5%)	2.5% (0.51-5.5%)	4.6 (0.92-10)	66 (13-150)
Suriname	2005	Female	80 to 84	19% (11-29%)	81% (71-89%)	0.66 (0.32-1.1)	1.9% (0.49-4.1%)	2.3% (0.58-4.9%)	4 (1-8.5)	42 (11-90)
Suriname	2005	Female	85 to 89	19% (11-29%)	81% (71-89%)	0.67 (0.32-1.2)	1.8% (0.43-3.9%)	2.1% (0.47-4.7%)	2.7 (0.6-5.9)	21 (4.9-45)
Suriname	2005	Female	90 to 94	21% (12-33%)	79% (67-88%)	0.76 (0.34-1.5)	1.9% (0.35-4.6%)	2.2% (0.32-5.3%)	1.4 (0.21-3.4)	7.3 (1.3-17)
Suriname	2005	Female	95 plus	24% (13-41%)	76% (59-87%)	0.91 (0.32-1.9)	2.2% (0.077-5.7%)	2.4% (-0.012-6.6%)	0.6 (-0.0028-1.6)	2.1 (0.075-5.6)
Suriname	2005	Male	All Ages	55% (47-64%)	45% (36-53%)	2.6 (2.1-3.1)	7.4% (5.8-9%)	9.1% (6.6-12%)	180 (130-230)	7100 (5600-8500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	2005	Male	15 to 19	56% (38-73%)	44% (27-62%)	2 (1.1-3.2)	6.9% (4.1-11%)	9.8% (5.4-16%)	3.3 (1.8-5.3)	280 (170-430)
Suriname	2005	Male	20 to 24	67% (48-81%)	33% (19-52%)	3.8 (2.2-5.6)	14% (9.4-19%)	18% (11-26%)	9.2 (5.6-13)	710 (470-980)
Suriname	2005	Male	25 to 29	70% (51-83%)	30% (17-49%)	4.6 (2.8-6.6)	15% (10-19%)	19% (13-25%)	9.1 (6.1-12)	660 (470-860)
Suriname	2005	Male	30 to 34	71% (51-84%)	29% (16-49%)	4.8 (3-6.9)	13% (9.3-16%)	16% (12-21%)	9.8 (7-13)	660 (480-840)
Suriname	2005	Male	35 to 39	70% (54-84%)	30% (16-46%)	4.5 (2.7-6.6)	13% (8.9-16%)	16% (11-21%)	12 (8.5-16)	720 (520-910)
Suriname	2005	Male	40 to 44	69% (53-83%)	31% (17-47%)	4.3 (2.4-6.5)	12% (8.2-16%)	15% (10-20%)	14 (9.3-19)	760 (510-980)
Suriname	2005	Male	45 to 49	68% (47-82%)	32% (18-53%)	4 (2.3-6.4)	12% (8.2-16%)	15% (10-20%)	15 (9.8-20)	680 (460-900)
Suriname	2005	Male	50 to 54	66% (47-81%)	34% (19-53%)	3.7 (2-5.9)	12% (7.2-16%)	15% (8.7-20%)	15 (8.9-21)	600 (370-830)
Suriname	2005	Male	55 to 59	64% (46-80%)	36% (20-54%)	3.4 (1.8-5.4)	11% (6.5-16%)	14% (7.9-19%)	16 (9-22)	530 (310-730)
Suriname	2005	Male	60 to 64	62% (42-79%)	38% (21-58%)	3 (1.6-5.2)	9.8% (5.2-14%)	12% (6.1-17%)	16 (8.4-24)	470 (240-680)
Suriname	2005	Male	65 to 69	60% (41-77%)	40% (23-59%)	2.7 (1.5-4.4)	8.9% (4.8-13%)	10% (5.7-16%)	18 (9.6-27)	420 (230-620)
Suriname	2005	Male	70 to 74	57% (38-73%)	43% (27-62%)	2.2 (1.1-3.9)	7.1% (3.3-12%)	8.2% (3.7-14%)	16 (7.3-27)	300 (140-510)
Suriname	2005	Male	75 to 79	52% (32-71%)	48% (29-68%)	1.9 (0.88-3.3)	5.6% (2.3-9.6%)	6.4% (2.6-11%)	11 (4.4-20)	170 (67-290)
Suriname	2005	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.5 (0.77-2.6)	4.5% (1.8-7.9%)	5.1% (2-9%)	6.8 (2.7-12)	74 (30-130)
Suriname	2005	Male	85 to 89	41% (25-57%)	59% (43-75%)	1.4 (0.68-2.3)	3.6% (1.4-6.7%)	4% (1.5-7.6%)	3.5 (1.3-6.7)	28 (11-51)
Suriname	2005	Male	90 to 94	41% (26-58%)	59% (42-74%)	1.4 (0.62-2.5)	3.2% (1.2-6.3%)	3.5% (1.2-7.2%)	1.4 (0.47-2.9)	7.7 (2.8-15)
Suriname	2005	Male	95 plus	44% (25-64%)	56% (36-75%)	1.5 (0.58-2.9)	3.4% (1-7.1%)	3.7% (0.9-8.1%)	0.5 (0.12-1.1)	1.9 (0.56-3.9)
Suriname	2010	Female	All Ages	32% (27-39%)	68% (61-73%)	1.3 (1-1.6)	2.7% (1.8-3.7%)	3.8% (2.1-5.8%)	63 (34-96)	2100 (1400-2900)
Suriname	2010	Female	15 to 19	39% (26-55%)	61% (45-74%)	1.5 (0.75-2.6)	3.2% (1.6-5.4%)	7.5% (3.2-13%)	1 (0.43-1.8)	91 (47-150)
Suriname	2010	Female	20 to 24	45% (29-61%)	55% (39-71%)	2.3 (1.2-3.7)	4.6% (2.8-6.8%)	9% (4.8-14%)	1.6 (0.83-2.6)	160 (94-230)
Suriname	2010	Female	25 to 29	44% (28-61%)	56% (39-72%)	2.3 (1.3-3.6)	4% (2.6-5.7%)	7% (4-11%)	1.4 (0.76-2.2)	130 (86-190)
Suriname	2010	Female	30 to 34	42% (26-60%)	58% (40-74%)	2.2 (1.3-3.6)	3.9% (2.6-5.5%)	6.7% (3.9-10%)	1.5 (0.84-2.2)	120 (77-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	2010	Female	35 to 39	41% (27-59%)	59% (41-73%)	2.1 (1.2-3.3)	3.9% (2.5-5.7%)	7% (4.3-10%)	2.3 (1.4-3.4)	160 (100-220)
Suriname	2010	Female	40 to 44	40% (26-59%)	60% (41-74%)	2 (1.1-3.2)	4.3% (2.6-6.4%)	7.6% (4.6-11%)	3.4 (2-4.9)	190 (110-270)
Suriname	2010	Female	45 to 49	38% (23-56%)	62% (44-77%)	1.9 (1-3.2)	4.6% (2.5-7.2%)	7.5% (4.3-12%)	4.9 (2.8-7.8)	230 (130-370)
Suriname	2010	Female	50 to 54	35% (21-52%)	65% (48-79%)	1.7 (0.88-2.9)	4.8% (2.5-7.9%)	7.4% (3.9-12%)	5.7 (3-9.2)	230 (120-380)
Suriname	2010	Female	55 to 59	31% (18-47%)	69% (53-82%)	1.4 (0.72-2.6)	4.5% (2.2-7.6%)	6.5% (3.2-11%)	5.7 (2.8-9.6)	190 (93-330)
Suriname	2010	Female	60 to 64	27% (16-42%)	73% (58-84%)	1.2 (0.57-2.2)	3.8% (1.4-7%)	5.2% (2-9.3%)	5.4 (2.1-9.5)	150 (58-290)
Suriname	2010	Female	65 to 69	25% (16-38%)	75% (62-84%)	1.1 (0.55-1.8)	3.4% (1.4-6%)	4.5% (1.9-7.8%)	6 (2.4-10)	140 (56-250)
Suriname	2010	Female	70 to 74	23% (14-35%)	77% (65-86%)	0.89 (0.44-1.6)	2.8% (0.81-5.6%)	3.6% (1.1-7.5%)	6.2 (1.8-13)	120 (33-240)
Suriname	2010	Female	75 to 79	20% (12-33%)	80% (67-88%)	0.77 (0.36-1.5)	2.3% (0.56-5.1%)	2.9% (0.7-6.2%)	5.8 (1.4-13)	84 (20-180)
Suriname	2010	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.73 (0.33-1.3)	2.2% (0.58-4.6%)	2.6% (0.68-5.5%)	5.2 (1.3-11)	56 (15-120)
Suriname	2010	Female	85 to 89	18% (11-28%)	82% (72-89%)	0.74 (0.34-1.4)	2.1% (0.57-4.5%)	2.5% (0.63-5.2%)	3.9 (0.99-8.3)	30 (7.8-63)
Suriname	2010	Female	90 to 94	20% (12-32%)	80% (68-88%)	0.84 (0.36-1.6)	2.2% (0.52-5.1%)	2.6% (0.54-6%)	2.1 (0.44-4.7)	11 (2.5-25)
Suriname	2010	Female	95 plus	24% (12-41%)	76% (59-88%)	1 (0.36-2.1)	2.7% (0.32-6.5%)	3% (0.24-7.7%)	0.77 (0.061-2)	2.7 (0.32-6.8)
Suriname	2010	Male	All Ages	56% (48-64%)	44% (36-52%)	2.9 (2.3-3.6)	8.7% (6.8-11%)	10% (7.6-13%)	220 (160-270)	8400 (6700-10000)
Suriname	2010	Male	15 to 19	56% (38-72%)	44% (28-62%)	2.2 (1.2-3.6)	7.6% (4.4-12%)	11% (5.8-18%)	3.4 (1.8-5.4)	280 (160-430)
Suriname	2010	Male	20 to 24	66% (48-81%)	34% (19-52%)	4.2 (2.4-6.4)	16% (10-21%)	21% (13-28%)	10 (6.3-15)	790 (520-1100)
Suriname	2010	Male	25 to 29	69% (51-82%)	31% (18-49%)	5.1 (3.1-7.5)	16% (12-21%)	21% (15-28%)	12 (8-15)	820 (590-1100)
Suriname	2010	Male	30 to 34	71% (51-84%)	29% (16-49%)	5.3 (3.2-7.8)	15% (11-20%)	20% (14-27%)	11 (7.4-14)	690 (510-870)
Suriname	2010	Male	35 to 39	70% (54-84%)	30% (16-46%)	5 (3-7.6)	14% (9.6-18%)	18% (12-24%)	13 (8.9-17)	770 (550-1000)
Suriname	2010	Male	40 to 44	69% (53-83%)	31% (17-47%)	4.7 (2.7-7.3)	14% (9.6-18%)	18% (12-23%)	16 (11-21)	840 (580-1100)
Suriname	2010	Male	45 to 49	67% (46-82%)	33% (18-54%)	4.5 (2.5-7.1)	13% (9-18%)	17% (11-22%)	21 (13-27)	950 (640-1200)
Suriname	2010	Male	50 to 54	66% (47-81%)	34% (19-53%)	4.1 (2.2-6.6)	13% (7.9-17%)	16% (9.6-21%)	20 (12-27)	810 (500-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	2010	Male	55 to 59	64% (45-79%)	36% (21-55%)	3.7 (2-6.2)	12% (7.5-17%)	15% (9.2-20%)	20 (13-28)	690 (430-950)
Suriname	2010	Male	60 to 64	62% (42-79%)	38% (21-58%)	3.4 (1.8-5.9)	11% (6.2-15%)	13% (7.4-18%)	19 (11-27)	550 (320-770)
Suriname	2010	Male	65 to 69	60% (40-77%)	40% (23-60%)	3 (1.7-5)	9.7% (5.7-14%)	11% (6.6-17%)	20 (11-29)	470 (270-690)
Suriname	2010	Male	70 to 74	56% (38-73%)	44% (27-62%)	2.5 (1.2-4.3)	7.9% (4-12%)	9.1% (4.5-14%)	19 (9.3-30)	350 (180-560)
Suriname	2010	Male	75 to 79	51% (32-70%)	49% (30-68%)	2 (1-3.8)	6.2% (2.8-10%)	7.1% (3.2-12%)	15 (6.5-25)	220 (97-360)
Suriname	2010	Male	80 to 84	46% (28-64%)	54% (36-72%)	1.7 (0.85-2.9)	5.1% (2.3-8.5%)	5.7% (2.5-9.6%)	9.2 (4-16)	100 (45-170)
Suriname	2010	Male	85 to 89	41% (24-58%)	59% (42-76%)	1.5 (0.71-2.6)	4.2% (1.7-7.4%)	4.6% (1.8-8.2%)	5.1 (2-9.1)	40 (16-70)
Suriname	2010	Male	90 to 94	41% (26-58%)	59% (42-74%)	1.5 (0.68-2.8)	3.7% (1.4-6.7%)	4% (1.4-7.7%)	1.9 (0.67-3.7)	11 (4-20)
Suriname	2010	Male	95 plus	44% (25-64%)	56% (36-75%)	1.6 (0.64-3.2)	3.9% (1.2-7.8%)	4.2% (1-8.7%)	0.56 (0.14-1.2)	2.1 (0.63-4.2)
Suriname	2016	Female	All Ages	32% (27-39%)	68% (61-73%)	1.4 (1-1.9)	3% (1.9-4.3%)	4.1% (2.2-6.3%)	74 (39-120)	2400 (1400-3400)
Suriname	2016	Female	15 to 19	38% (25-55%)	62% (45-75%)	1.6 (0.78-2.8)	3.2% (1.6-5.5%)	7.6% (3.2-14%)	1 (0.42-1.9)	95 (47-160)
Suriname	2016	Female	20 to 24	44% (28-61%)	56% (39-72%)	2.4 (1.2-4.1)	4.6% (2.8-7%)	9.5% (4.8-15%)	1.4 (0.72-2.4)	140 (82-210)
Suriname	2016	Female	25 to 29	44% (27-61%)	56% (39-73%)	2.5 (1.3-4.1)	4.4% (2.8-6.5%)	8.3% (4.7-13%)	1.6 (0.87-2.6)	150 (95-220)
Suriname	2016	Female	30 to 34	42% (26-60%)	58% (40-74%)	2.4 (1.3-4)	4.1% (2.6-5.9%)	7.5% (4.1-11%)	1.8 (0.97-2.8)	150 (89-210)
Suriname	2016	Female	35 to 39	41% (26-58%)	59% (42-74%)	2.2 (1.2-3.7)	4.1% (2.5-6.2%)	7.7% (4.5-11%)	2.2 (1.2-3.3)	150 (89-220)
Suriname	2016	Female	40 to 44	40% (25-58%)	60% (42-75%)	2.1 (1.1-3.4)	4.4% (2.5-6.6%)	7.8% (4.4-12%)	3.4 (1.9-5.1)	190 (110-290)
Suriname	2016	Female	45 to 49	38% (23-55%)	62% (45-77%)	2 (0.99-3.6)	4.6% (2.5-7.4%)	7.8% (4.3-12%)	4.7 (2.6-7.4)	230 (120-360)
Suriname	2016	Female	50 to 54	34% (21-50%)	66% (50-79%)	1.8 (0.89-3.2)	4.9% (2.5-8.1%)	7.7% (4.1-12%)	7.1 (3.7-11)	290 (150-470)
Suriname	2016	Female	55 to 59	30% (18-46%)	70% (54-82%)	1.5 (0.74-2.8)	4.6% (2.1-7.9%)	6.7% (3.3-11%)	7.4 (3.6-12)	260 (120-440)
Suriname	2016	Female	60 to 64	27% (16-42%)	73% (58-84%)	1.3 (0.58-2.4)	3.9% (1.5-7.1%)	5.4% (2.2-9.8%)	6.7 (2.7-12)	200 (73-370)
Suriname	2016	Female	65 to 69	25% (15-38%)	75% (62-85%)	1.1 (0.57-2)	3.6% (1.3-6.3%)	4.8% (1.9-8.2%)	6.8 (2.6-12)	160 (61-290)
Suriname	2016	Female	70 to 74	22% (14-36%)	78% (64-86%)	0.95 (0.46-1.8)	3% (0.95-6.4%)	3.9% (1.3-8%)	6.8 (2.2-14)	130 (41-260)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	2016	Female	75 to 79	20% (12-32%)	80% (68-88%)	0.83 (0.38-1.6)	2.5% (0.61-5.4%)	3.2% (0.79-6.7%)	6.7 (1.7-15)	98 (24-210)
Suriname	2016	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.77 (0.37-1.4)	2.4% (0.69-5%)	2.9% (0.82-6%)	6.6 (1.9-14)	71 (21-150)
Suriname	2016	Female	85 to 89	18% (10-28%)	82% (72-90%)	0.78 (0.35-1.4)	2.3% (0.66-4.8%)	2.7% (0.74-5.7%)	5.6 (1.5-12)	43 (12-89)
Suriname	2016	Female	90 to 94	20% (12-32%)	80% (68-88%)	0.89 (0.36-1.7)	2.4% (0.59-5.4%)	2.8% (0.61-6.5%)	3.1 (0.69-7.2)	16 (4-37)
Suriname	2016	Female	95 plus	24% (12-40%)	76% (60-88%)	1.1 (0.37-2.2)	2.8% (0.42-6.5%)	3.2% (0.35-7.8%)	1.2 (0.13-3)	4.2 (0.61-10)
Suriname	2016	Male	All Ages	56% (48-64%)	44% (36-52%)	3.1 (2.3-4.2)	9.3% (7-12%)	11% (7.7-14%)	240 (160-310)	9000 (6800-11000)
Suriname	2016	Male	15 to 19	56% (37-73%)	44% (27-63%)	2.3 (1.2-3.9)	7.6% (4.3-12%)	12% (5.8-19%)	3.5 (1.7-5.9)	300 (170-480)
Suriname	2016	Male	20 to 24	66% (48-81%)	34% (19-52%)	4.4 (2.4-7.1)	16% (10-21%)	21% (13-30%)	9 (5.2-13)	700 (450-960)
Suriname	2016	Male	25 to 29	69% (51-82%)	31% (18-49%)	5.3 (3-8.5)	17% (12-22%)	23% (15-31%)	11 (7.4-15)	810 (560-1100)
Suriname	2016	Male	30 to 34	70% (51-84%)	30% (16-49%)	5.5 (3.2-8.8)	16% (11-21%)	22% (16-29%)	12 (8.6-17)	810 (580-1000)
Suriname	2016	Male	35 to 39	70% (54-84%)	30% (16-46%)	5.3 (3.1-8.4)	15% (11-20%)	21% (14-27%)	12 (8.3-17)	730 (520-960)
Suriname	2016	Male	40 to 44	69% (52-82%)	31% (18-48%)	5 (2.7-8.1)	14% (10-19%)	19% (13-25%)	16 (11-21)	830 (570-1100)
Suriname	2016	Male	45 to 49	67% (46-82%)	33% (18-54%)	4.7 (2.5-8)	14% (9.2-18%)	18% (12-24%)	20 (13-27)	930 (610-1200)
Suriname	2016	Male	50 to 54	65% (47-81%)	35% (19-53%)	4.3 (2.2-7.2)	13% (8-17%)	16% (9.9-21%)	25 (16-34)	1000 (640-1400)
Suriname	2016	Male	55 to 59	63% (46-79%)	37% (21-54%)	3.9 (2-6.7)	12% (7.9-17%)	15% (9.8-20%)	26 (16-36)	900 (580-1200)
Suriname	2016	Male	60 to 64	61% (41-78%)	39% (22-59%)	3.5 (1.7-6.2)	11% (6.2-15%)	13% (7.3-18%)	23 (13-34)	680 (380-970)
Suriname	2016	Male	65 to 69	60% (40-77%)	40% (23-60%)	3.1 (1.6-5.4)	9.8% (5.6-14%)	12% (6.5-17%)	22 (12-32)	520 (290-770)
Suriname	2016	Male	70 to 74	56% (38-73%)	44% (27-62%)	2.6 (1.3-4.7)	7.9% (4-12%)	9.1% (4.6-14%)	19 (9.3-30)	360 (180-560)
Suriname	2016	Male	75 to 79	51% (32-70%)	49% (30-68%)	2.1 (1-4.1)	6.2% (2.8-10%)	7.1% (3.2-12%)	15 (6.7-26)	230 (100-380)
Suriname	2016	Male	80 to 84	45% (27-63%)	55% (37-73%)	1.8 (0.87-3.2)	5.2% (2.4-8.7%)	5.8% (2.6-9.8%)	11 (4.9-19)	120 (55-200)
Suriname	2016	Male	85 to 89	40% (24-57%)	60% (43-76%)	1.6 (0.75-2.8)	4.3% (1.8-7.5%)	4.8% (1.9-8.5%)	6.6 (2.6-12)	51 (21-91)
Suriname	2016	Male	90 to 94	40% (26-57%)	60% (43-74%)	1.6 (0.69-2.9)	3.8% (1.4-7%)	4.2% (1.4-7.8%)	2.6 (0.87-4.9)	14 (5.2-26)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Suriname	2016	Male	95 plus	43% (25-63%)	57% (37-75%)	1.7 (0.63-3.2)	4.1% (1.2-8%)	4.4% (1.1-9%)	0.77 (0.2-1.5)	2.9 (0.89-5.6)
Trinidad and Tobago	1990	Female	All Ages	31% (26-38%)	69% (62-74%)	0.69 (0.59-0.8)	1.1% (0.53-1.8%)	0.85% (-0.16-2.1%)	33 (-6-83)	1900 (900-3200)
Trinidad and Tobago	1990	Female	15 to 19	38% (25-55%)	62% (45-75%)	0.85 (0.45-1.4)	1.8% (1-2.9%)	3.4% (1.6-6%)	1.2 (0.56-2.1)	130 (79-220)
Trinidad and Tobago	1990	Female	20 to 24	44% (29-61%)	56% (39-71%)	1.3 (0.72-2)	3.2% (2-4.7%)	4.9% (2.6-8%)	2.6 (1.3-4.2)	270 (160-400)
Trinidad and Tobago	1990	Female	25 to 29	44% (28-60%)	56% (40-72%)	1.3 (0.8-1.9)	2.8% (1.9-3.8%)	3.9% (2.2-5.9%)	2.4 (1.3-3.7)	260 (170-370)
Trinidad and Tobago	1990	Female	30 to 34	42% (26-59%)	58% (41-74%)	1.3 (0.74-1.9)	2.8% (1.9-3.9%)	4.2% (2.6-6.1%)	3.1 (2-4.5)	260 (170-360)
Trinidad and Tobago	1990	Female	35 to 39	41% (27-59%)	59% (41-73%)	1.2 (0.66-1.8)	2.6% (1.7-3.8%)	4.2% (2.6-6.3%)	3.4 (2.1-5.1)	230 (150-330)
Trinidad and Tobago	1990	Female	40 to 44	40% (26-58%)	60% (42-74%)	1.1 (0.6-1.8)	2.3% (1.1-3.8%)	3.7% (1.8-6.1%)	3.4 (1.7-5.8)	190 (94-320)
Trinidad and Tobago	1990	Female	45 to 49	38% (23-55%)	62% (45-77%)	1.1 (0.55-1.8)	2.1% (0.58-4.2%)	3.1% (0.91-6%)	4.5 (1.3-8.6)	210 (57-400)
Trinidad and Tobago	1990	Female	50 to 54	35% (21-51%)	65% (49-79%)	0.95 (0.5-1.6)	1.6% (-0.19-4%)	2.2% (-0.12-5.3%)	4.2 (-0.23-10)	160 (-18-390)
Trinidad and Tobago	1990	Female	55 to 59	30% (18-47%)	70% (53-82%)	0.8 (0.4-1.4)	1.1% (-0.63-3.3%)	1.4% (-0.75-4.1%)	3.2 (-1.7-9.6)	110 (-61-340)
Trinidad and Tobago	1990	Female	60 to 64	27% (16-42%)	73% (58-84%)	0.68 (0.34-1.2)	0.49% (-1.1-2.5%)	0.62% (-1.4-3.1%)	1.8 (-4.1-9.1)	52 (-120-260)
Trinidad and Tobago	1990	Female	65 to 69	25% (15-38%)	75% (62-85%)	0.6 (0.31-0.99)	0.22% (-1.1-1.9%)	0.29% (-1.3-2.3%)	1.1 (-5.3-8.9)	25 (-130-210)
Trinidad and Tobago	1990	Female	70 to 74	22% (14-35%)	78% (65-86%)	0.5 (0.25-0.91)	0.098% (-1.2-1.7%)	0.13% (-1.4-2%)	0.58 (-6.4-8.9)	9.8 (-130-170)
Trinidad and Tobago	1990	Female	75 to 79	20% (11-32%)	80% (68-89%)	0.43 (0.19-0.79)	0.0039% (-1.1-1.4%)	-0.0009% (-1.3-1.6%)	-0.0095 (-6.4-7.7)	0.29 (-93-120)
Trinidad and Tobago	1990	Female	80 to 84	18% (10-29%)	82% (71-90%)	0.4 (0.18-0.68)	0.1% (-0.77-1.2%)	0.11% (-0.96-1.4%)	0.46 (-4.1-6.4)	5.8 (-44-69)
Trinidad and Tobago	1990	Female	85 to 89	18% (11-28%)	82% (72-89%)	0.41 (0.19-0.72)	0.19% (-0.67-1.4%)	0.18% (-0.82-1.6%)	0.6 (-2.7-5.2)	5.5 (-19-41)
Trinidad and Tobago	1990	Female	90 to 94	20% (12-32%)	80% (68-88%)	0.46 (0.21-0.85)	0.26% (-0.7-1.7%)	0.24% (-0.92-1.9%)	0.36 (-1.4-2.8)	2.4 (-6.3-15)
Trinidad and Tobago	1990	Female	95 plus	24% (12-40%)	76% (60-88%)	0.55 (0.2-1.1)	0.35% (-0.96-2.5%)	0.27% (-1.2-2.7%)	0.11 (-0.51-1.1)	0.56 (-1.6-3.9)
Trinidad and Tobago	1990	Male	All Ages	53% (45-62%)	47% (38-55%)	1.5 (1.3-1.7)	4.9% (3.5-6.2%)	4.8% (2.9-7.1%)	230 (140-350)	10000 (7400-13000)
Trinidad and Tobago	1990	Male	15 to 19	56% (38-72%)	44% (28-62%)	1.3 (0.68-2)	4.7% (2.9-6.9%)	6.3% (3.3-10%)	4.5 (2.4-7.2)	420 (270-620)
Trinidad and Tobago	1990	Male	20 to 24	66% (48-81%)	34% (19-52%)	2.4 (1.5-3.4)	9.6% (6.5-13%)	11% (6.9-17%)	12 (7.5-18)	1000 (690-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	1990	Male	25 to 29	69% (51-83%)	31% (17-49%)	2.9 (1.9-4)	10% (7.2-14%)	12% (7.7-17%)	16 (10-23)	1300 (880-1700)
Trinidad and Tobago	1990	Male	30 to 34	70% (51-84%)	30% (16-49%)	3 (1.9-4.2)	9.9% (7.1-13%)	12% (8.1-17%)	18 (12-25)	1200 (870-1700)
Trinidad and Tobago	1990	Male	35 to 39	70% (53-84%)	30% (16-47%)	2.8 (1.7-4.3)	9.1% (6-12%)	11% (7.3-16%)	15 (9.8-22)	980 (660-1300)
Trinidad and Tobago	1990	Male	40 to 44	69% (52-83%)	31% (17-48%)	2.7 (1.5-4.1)	8.6% (5.1-12%)	11% (6-16%)	16 (8.8-23)	890 (540-1300)
Trinidad and Tobago	1990	Male	45 to 49	67% (47-82%)	33% (18-53%)	2.5 (1.4-4.1)	7.8% (4.3-12%)	9.4% (4.9-15%)	18 (9.5-30)	880 (480-1400)
Trinidad and Tobago	1990	Male	50 to 54	65% (47-81%)	35% (19-53%)	2.3 (1.3-3.8)	6.7% (3-11%)	7.8% (3.3-13%)	20 (8.1-34)	820 (360-1400)
Trinidad and Tobago	1990	Male	55 to 59	63% (45-79%)	37% (21-55%)	2.1 (1.2-3.4)	6% (2.6-10%)	6.8% (2.8-12%)	21 (8.5-36)	720 (310-1300)
Trinidad and Tobago	1990	Male	60 to 64	61% (41-78%)	39% (22-59%)	1.9 (1-3.3)	4.8% (1.5-9.5%)	5.4% (1.6-11%)	21 (6.3-43)	620 (190-1200)
Trinidad and Tobago	1990	Male	65 to 69	60% (41-77%)	40% (23-59%)	1.7 (0.92-2.7)	4.2% (1.4-8%)	4.6% (1.5-9.1%)	23 (7.7-46)	560 (180-1100)
Trinidad and Tobago	1990	Male	70 to 74	56% (37-72%)	44% (28-63%)	1.4 (0.69-2.4)	3.1% (0.64-6.6%)	3.3% (0.58-7.3%)	19 (3.3-41)	360 (73-770)
Trinidad and Tobago	1990	Male	75 to 79	51% (32-71%)	49% (29-68%)	1.2 (0.56-2)	2.3% (0.26-5.3%)	2.5% (0.21-5.9%)	14 (1.2-33)	210 (25-480)
Trinidad and Tobago	1990	Male	80 to 84	45% (27-64%)	55% (36-73%)	0.97 (0.48-1.7)	1.8% (0.31-4.1%)	1.9% (0.25-4.4%)	7.8 (1-19)	88 (15-200)
Trinidad and Tobago	1990	Male	85 to 89	41% (25-58%)	59% (42-75%)	0.85 (0.42-1.5)	1.3% (0.12-3.3%)	1.4% (0.012-3.6%)	3.4 (0.03-8.9)	28 (2.5-70)
Trinidad and Tobago	1990	Male	90 to 94	41% (26-58%)	59% (42-74%)	0.86 (0.42-1.5)	1.1% (-0.11-3.1%)	1.1% (-0.3-3.4%)	1 (-0.28-3.3)	6.2 (-0.61-18)
Trinidad and Tobago	1990	Male	95 plus	44% (25-65%)	56% (35-75%)	0.93 (0.39-1.8)	1.1% (-0.39-3.6%)	0.98% (-0.71-3.7%)	0.2 (-0.15-0.79)	0.9 (-0.31-2.9)
Trinidad and Tobago	1995	Female	All Ages	31% (25-38%)	69% (62-75%)	0.62 (0.53-0.72)	0.9% (0.35-1.5%)	0.51% (-0.34-1.6%)	21 (-14-66)	1600 (620-2700)
Trinidad and Tobago	1995	Female	15 to 19	37% (24-53%)	63% (47-76%)	0.73 (0.38-1.2)	1.6% (0.91-2.6%)	2.9% (1.2-5.3%)	1.1 (0.47-2.1)	130 (71-210)
Trinidad and Tobago	1995	Female	20 to 24	43% (27-59%)	57% (41-73%)	1.1 (0.61-1.7)	2.7% (1.7-4%)	4% (1.9-6.6%)	2.1 (1-3.6)	230 (140-350)
Trinidad and Tobago	1995	Female	25 to 29	42% (26-60%)	58% (40-74%)	1.1 (0.67-1.7)	2.5% (1.7-3.6%)	3.2% (1.7-5.2%)	2 (1-3.3)	210 (140-300)
Trinidad and Tobago	1995	Female	30 to 34	40% (25-58%)	60% (42-75%)	1.1 (0.64-1.7)	2.4% (1.6-3.3%)	3.4% (2.1-5%)	2.9 (1.8-4.4)	260 (170-360)
Trinidad and Tobago	1995	Female	35 to 39	39% (25-57%)	61% (43-75%)	1 (0.57-1.6)	2.3% (1.4-3.3%)	3.4% (2-5.2%)	3.4 (2-5.2)	230 (150-340)
Trinidad and Tobago	1995	Female	40 to 44	38% (24-57%)	62% (43-76%)	0.95 (0.52-1.5)	1.9% (0.89-3.2%)	2.9% (1.3-4.9%)	3.6 (1.6-6.1)	200 (96-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	1995	Female	45 to 49	36% (22-54%)	64% (46-78%)	0.91 (0.47-1.5)	1.6% (0.24-3.4%)	2.4% (0.45-4.9%)	3.6 (0.72-7.6)	170 (26-360)
Trinidad and Tobago	1995	Female	50 to 54	33% (20-49%)	67% (51-80%)	0.82 (0.44-1.4)	1.1% (-0.43-3.1%)	1.5% (-0.54-4.1%)	3.1 (-1.2-8.4)	120 (-52-340)
Trinidad and Tobago	1995	Female	55 to 59	29% (18-45%)	71% (55-82%)	0.69 (0.35-1.2)	0.66% (-0.8-2.4%)	0.89% (-0.99-3.1%)	2.2 (-2.4-7.9)	74 (-90-270)
Trinidad and Tobago	1995	Female	60 to 64	26% (15-40%)	74% (60-85%)	0.59 (0.29-1)	0.15% (-1.3-1.8%)	0.2% (-1.6-2.2%)	0.62 (-5-6.9)	17 (-150-200)
Trinidad and Tobago	1995	Female	65 to 69	23% (14-36%)	77% (64-86%)	0.51 (0.26-0.86)	-0.12% (-1.3-1.3%)	-0.13% (-1.6-1.7%)	-0.49 (-6.2-6.5)	-13 (-150-160)
Trinidad and Tobago	1995	Female	70 to 74	21% (13-34%)	79% (66-87%)	0.43 (0.21-0.81)	-0.18% (-1.3-1.1%)	-0.2% (-1.6-1.4%)	-0.93 (-7.3-6.1)	-19 (-140-110)
Trinidad and Tobago	1995	Female	75 to 79	19% (11-31%)	81% (69-89%)	0.38 (0.17-0.71)	-0.22% (-1.2-0.89%)	-0.27% (-1.5-1%)	-1.3 (-7-4.9)	-19 (-100-75)
Trinidad and Tobago	1995	Female	80 to 84	17% (9.5-28%)	83% (72-90%)	0.35 (0.16-0.6)	-0.12% (-0.94-0.81%)	-0.16% (-1.1-0.92%)	-0.79 (-5.5-4.6)	-7.6 (-58-52)
Trinidad and Tobago	1995	Female	85 to 89	17% (9.8-27%)	83% (73-90%)	0.36 (0.16-0.63)	-0.012% (-0.8-1%)	-0.054% (-0.98-1.1%)	-0.22 (-4-4.5)	-0.42 (-29-36)
Trinidad and Tobago	1995	Female	90 to 94	19% (11-30%)	81% (70-89%)	0.41 (0.17-0.77)	0.069% (-0.84-1.4%)	0.0093% (-1.1-1.5%)	0.018 (-2.1-3)	0.84 (-10-17)
Trinidad and Tobago	1995	Female	95 plus	23% (12-39%)	77% (61-88%)	0.49 (0.18-1)	0.14% (-1.1-2.1%)	0.035% (-1.4-2.3%)	0.021 (-0.83-1.4)	0.35 (-2.5-4.9)
Trinidad and Tobago	1995	Male	All Ages	53% (45-62%)	47% (38-55%)	1.4 (1.2-1.6)	4.3% (3.1-5.6%)	3.9% (2.2-5.8%)	200 (110-310)	9300 (6600-12000)
Trinidad and Tobago	1995	Male	15 to 19	54% (36-71%)	46% (29-64%)	1.1 (0.57-1.8)	4.2% (2.6-6.3%)	5.8% (3-9.3%)	3.8 (2-6.2)	380 (230-570)
Trinidad and Tobago	1995	Male	20 to 24	65% (46-80%)	35% (20-54%)	2.1 (1.3-3.1)	8.8% (6-12%)	10% (6.2-16%)	11 (6.2-16)	940 (630-1300)
Trinidad and Tobago	1995	Male	25 to 29	68% (50-82%)	32% (18-50%)	2.5 (1.6-3.6)	9.3% (6.4-13%)	11% (6.8-16%)	13 (8.3-20)	1000 (710-1400)
Trinidad and Tobago	1995	Male	30 to 34	69% (49-83%)	31% (17-51%)	2.6 (1.7-3.7)	8.5% (5.9-12%)	10% (6.5-15%)	18 (11-26)	1200 (860-1700)
Trinidad and Tobago	1995	Male	35 to 39	69% (52-83%)	31% (17-48%)	2.5 (1.5-3.6)	7.5% (4.9-10%)	9% (5.5-13%)	16 (9.5-23)	1000 (660-1400)
Trinidad and Tobago	1995	Male	40 to 44	68% (51-82%)	32% (18-49%)	2.4 (1.4-3.5)	6.8% (4-9.8%)	8.2% (4.5-12%)	16 (8.7-25)	910 (540-1300)
Trinidad and Tobago	1995	Male	45 to 49	66% (44-81%)	34% (19-56%)	2.2 (1.2-3.5)	6.4% (3.4-10%)	7.6% (3.8-12%)	18 (9.1-30)	890 (460-1400)
Trinidad and Tobago	1995	Male	50 to 54	64% (44-80%)	36% (20-56%)	2 (1.1-3.2)	5.3% (2.1-9.1%)	6.2% (2.2-11%)	18 (6.3-31)	750 (300-1300)
Trinidad and Tobago	1995	Male	55 to 59	62% (43-78%)	38% (22-57%)	1.8 (0.98-2.9)	4.6% (1.6-8.2%)	5.2% (1.7-9.5%)	18 (6.1-33)	640 (230-1200)
Trinidad and Tobago	1995	Male	60 to 64	60% (40-77%)	40% (23-60%)	1.6 (0.86-2.8)	3.7% (0.74-7.7%)	4.1% (0.77-8.8%)	17 (3-36)	490 (98-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	1995	Male	65 to 69	58% (39-75%)	42% (25-61%)	1.4 (0.78-2.4)	3.1% (0.69-6.4%)	3.4% (0.74-7.2%)	17 (3.6-35)	410 (94-840)
Trinidad and Tobago	1995	Male	70 to 74	54% (36-71%)	46% (29-64%)	1.2 (0.6-2.1)	2.3% (0.15-5.4%)	2.4% (0.1-5.9%)	15 (0.61-36)	290 (19-690)
Trinidad and Tobago	1995	Male	75 to 79	49% (31-68%)	51% (32-69%)	1 (0.48-1.8)	1.7% (0.028-4.2%)	1.8% (-0.098-4.5%)	10 (-0.77-26)	160 (2.6-390)
Trinidad and Tobago	1995	Male	80 to 84	43% (26-61%)	57% (39-74%)	0.83 (0.41-1.4)	1.3% (0.047-3.1%)	1.3% (-0.015-3.3%)	6.9 (-0.08-17)	78 (3-190)
Trinidad and Tobago	1995	Male	85 to 89	39% (23-56%)	61% (44-77%)	0.73 (0.35-1.2)	0.93% (-0.12-2.5%)	0.93% (-0.26-2.7%)	2.8 (-0.77-8.2)	24 (-3.1-64)
Trinidad and Tobago	1995	Male	90 to 94	39% (25-56%)	61% (44-75%)	0.73 (0.35-1.4)	0.72% (-0.32-2.3%)	0.66% (-0.52-2.5%)	0.77 (-0.6-2.9)	4.9 (-2.2-16)
Trinidad and Tobago	1995	Male	95 plus	42% (24-63%)	58% (37-76%)	0.8 (0.32-1.5)	0.69% (-0.56-2.8%)	0.54% (-0.92-2.9%)	0.16 (-0.28-0.89)	0.82 (-0.66-3.3)
Trinidad and Tobago	2000	Female	All Ages	30% (24-37%)	70% (63-76%)	0.6 (0.52-0.69)	0.75% (0.27-1.3%)	0.33% (-0.42-1.2%)	15 (-19-56)	1400 (510-2400)
Trinidad and Tobago	2000	Female	15 to 19	35% (22-52%)	65% (48-78%)	0.67 (0.36-1.1)	1.5% (0.82-2.4%)	2.5% (1.1-4.8%)	1.2 (0.5-2.3)	140 (77-220)
Trinidad and Tobago	2000	Female	20 to 24	41% (26-58%)	59% (42-74%)	1 (0.56-1.6)	2.3% (1.5-3.3%)	2.8% (1.4-4.6%)	1.8 (0.92-3)	220 (140-320)
Trinidad and Tobago	2000	Female	25 to 29	40% (25-58%)	60% (42-75%)	1 (0.59-1.6)	2.1% (1.4-2.9%)	2.3% (1.2-3.7%)	1.7 (0.88-2.7)	190 (120-270)
Trinidad and Tobago	2000	Female	30 to 34	39% (23-56%)	61% (44-77%)	1 (0.57-1.5)	2.1% (1.4-2.9%)	2.6% (1.6-3.9%)	2.2 (1.3-3.2)	200 (130-270)
Trinidad and Tobago	2000	Female	35 to 39	38% (24-55%)	62% (45-76%)	0.93 (0.52-1.4)	1.9% (1.2-2.6%)	2.6% (1.6-3.9%)	3.2 (1.9-4.7)	230 (140-320)
Trinidad and Tobago	2000	Female	40 to 44	37% (23-55%)	63% (45-77%)	0.87 (0.47-1.4)	1.7% (0.84-2.8%)	2.5% (1.2-4.1%)	3.8 (1.8-6.1)	210 (110-340)
Trinidad and Tobago	2000	Female	45 to 49	35% (20-52%)	65% (48-80%)	0.84 (0.43-1.4)	1.3% (0.21-2.8%)	1.9% (0.39-4%)	3.4 (0.69-6.9)	160 (27-340)
Trinidad and Tobago	2000	Female	50 to 54	32% (19-47%)	68% (53-81%)	0.75 (0.4-1.3)	0.88% (-0.44-2.6%)	1.3% (-0.56-3.5%)	2.8 (-1.3-7.7)	110 (-56-320)
Trinidad and Tobago	2000	Female	55 to 59	28% (17-43%)	72% (57-83%)	0.63 (0.32-1.1)	0.44% (-0.84-2%)	0.63% (-1-2.6%)	1.8 (-2.8-7.4)	57 (-110-260)
Trinidad and Tobago	2000	Female	60 to 64	24% (14-39%)	76% (61-86%)	0.54 (0.26-0.98)	-0.038% (-1.3-1.5%)	-0.025% (-1.6-1.8%)	-0.084 (-5.5-6.4)	-4.7 (-170-190)
Trinidad and Tobago	2000	Female	65 to 69	22% (14-35%)	78% (65-86%)	0.46 (0.23-0.78)	-0.26% (-1.4-0.99%)	-0.28% (-1.6-1.2%)	-1.1 (-6.3-4.9)	-30 (-160-110)
Trinidad and Tobago	2000	Female	70 to 74	20% (12-32%)	80% (68-88%)	0.39 (0.19-0.71)	-0.3% (-1.3-0.83%)	-0.34% (-1.6-1%)	-1.5 (-7.1-4.7)	-32 (-140-90)
Trinidad and Tobago	2000	Female	75 to 79	18% (10-29%)	82% (71-90%)	0.34 (0.16-0.62)	-0.31% (-1.2-0.69%)	-0.38% (-1.5-0.81%)	-1.9 (-7.6-4.1)	-29 (-110-60)
Trinidad and Tobago	2000	Female	80 to 84	16% (9.3-27%)	84% (73-91%)	0.32 (0.15-0.54)	-0.21% (-0.97-0.65%)	-0.26% (-1.2-0.75%)	-1.3 (-6-3.8)	-14 (-63-42)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	2000	Female	85 to 89	16% (9.3-25%)	84% (75-91%)	0.32 (0.15-0.58)	-0.098% (-0.81-0.81%)	-0.15% (-1-0.89%)	-0.68 (-4.5-4)	-3.9 (-33-32)
Trinidad and Tobago	2000	Female	90 to 94	18% (11-29%)	82% (71-89%)	0.37 (0.16-0.69)	-0.00017% (-0.79-1.2%)	-0.069% (-1-1.3%)	-0.17 (-2.6-3.4)	-0.0012 (-12-18)
Trinidad and Tobago	2000	Female	95 plus	22% (11-37%)	78% (63-89%)	0.45 (0.16-0.93)	0.074% (-0.99-1.8%)	-0.043% (-1.3-1.9%)	-0.035 (-1-1.5)	0.24 (-3.2-5.6)
Trinidad and Tobago	2000	Male	All Ages	53% (44-62%)	47% (38-56%)	1.4 (1.2-1.5)	3.9% (2.8-5.1%)	3.4% (2-5.2%)	190 (110-280)	9100 (6500-12000)
Trinidad and Tobago	2000	Male	15 to 19	52% (34-70%)	48% (30-66%)	1 (0.53-1.6)	4% (2.5-5.9%)	5.4% (2.8-8.9%)	4.1 (2.1-6.7)	420 (260-620)
Trinidad and Tobago	2000	Male	20 to 24	63% (44-79%)	37% (21-56%)	1.9 (1.1-2.9)	8.3% (5.6-12%)	9.6% (5.6-15%)	11 (6.4-17)	970 (630-1400)
Trinidad and Tobago	2000	Male	25 to 29	67% (47-81%)	33% (19-53%)	2.3 (1.4-3.4)	8.1% (5.4-11%)	8.9% (5.3-13%)	12 (7.1-18)	980 (650-1400)
Trinidad and Tobago	2000	Male	30 to 34	68% (47-82%)	32% (18-53%)	2.4 (1.5-3.6)	7.4% (5.1-10%)	8.4% (5.2-12%)	13 (8.1-19)	930 (630-1300)
Trinidad and Tobago	2000	Male	35 to 39	67% (50-82%)	33% (18-50%)	2.3 (1.4-3.4)	6.4% (4.3-8.9%)	7.2% (4.4-10%)	16 (9.8-24)	1100 (690-1500)
Trinidad and Tobago	2000	Male	40 to 44	66% (50-81%)	34% (19-50%)	2.2 (1.3-3.3)	6.1% (3.8-8.9%)	7% (4-11%)	17 (10-26)	990 (620-1400)
Trinidad and Tobago	2000	Male	45 to 49	64% (43-80%)	36% (20-57%)	2 (1.1-3.3)	5.5% (3.1-8.7%)	6.3% (3.3-10%)	18 (9.4-29)	900 (500-1400)
Trinidad and Tobago	2000	Male	50 to 54	62% (43-79%)	38% (21-57%)	1.8 (1-2.9)	4.7% (2-8%)	5.3% (2.1-9.3%)	19 (7.7-34)	820 (350-1400)
Trinidad and Tobago	2000	Male	55 to 59	60% (42-77%)	40% (23-58%)	1.7 (0.91-2.7)	4% (1.5-7.3%)	4.6% (1.5-8.3%)	19 (6.6-36)	690 (250-1200)
Trinidad and Tobago	2000	Male	60 to 64	58% (37-76%)	42% (24-63%)	1.5 (0.79-2.6)	3.2% (0.59-6.8%)	3.6% (0.64-7.7%)	16 (2.9-35)	490 (88-1000)
Trinidad and Tobago	2000	Male	65 to 69	57% (37-74%)	43% (26-63%)	1.3 (0.71-2.2)	2.7% (0.5-5.6%)	2.9% (0.46-6.1%)	15 (2.4-32)	370 (70-770)
Trinidad and Tobago	2000	Male	70 to 74	53% (35-70%)	47% (30-65%)	1.1 (0.55-1.9)	1.9% (0.069-4.6%)	2% (-0.061-5.1%)	12 (-0.35-30)	230 (8.7-580)
Trinidad and Tobago	2000	Male	75 to 79	48% (29-67%)	52% (33-71%)	0.9 (0.43-1.6)	1.4% (-0.097-3.6%)	1.4% (-0.17-3.9%)	8.6 (-1-23)	130 (-9.4-350)
Trinidad and Tobago	2000	Male	80 to 84	42% (25-60%)	58% (40-75%)	0.74 (0.37-1.2)	1% (0.0041-2.5%)	1.1% (-0.092-2.7%)	5.4 (-0.47-14)	63 (0.27-150)
Trinidad and Tobago	2000	Male	85 to 89	37% (22-54%)	63% (46-78%)	0.65 (0.32-1.1)	0.74% (-0.12-2%)	0.72% (-0.25-2.2%)	2.4 (-0.83-7)	21 (-3.3-57)
Trinidad and Tobago	2000	Male	90 to 94	38% (23-55%)	62% (45-77%)	0.66 (0.31-1.2)	0.55% (-0.32-1.9%)	0.48% (-0.53-2%)	0.68 (-0.74-2.9)	4.6 (-2.6-16)
Trinidad and Tobago	2000	Male	95 plus	41% (22-61%)	59% (39-78%)	0.72 (0.29-1.3)	0.52% (-0.59-2.3%)	0.35% (-0.93-2.4%)	0.13 (-0.34-0.87)	0.74 (-0.81-3.2)
Trinidad and Tobago	2005	Female	All Ages	32% (26-39%)	68% (61-74%)	0.69 (0.59-0.79)	0.91% (0.41-1.4%)	0.49% (-0.27-1.4%)	21 (-12-60)	1700 (740-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	2005	Female	15 to 19	37% (24-53%)	63% (47-76%)	0.73 (0.4-1.2)	1.6% (0.99-2.6%)	3% (1.4-5.3%)	1.2 (0.53-2.1)	130 (76-200)
Trinidad and Tobago	2005	Female	20 to 24	42% (27-60%)	58% (40-73%)	1.1 (0.61-1.7)	2.5% (1.7-3.5%)	3.5% (1.9-5.6%)	2 (1.1-3.2)	250 (160-360)
Trinidad and Tobago	2005	Female	25 to 29	42% (26-59%)	58% (41-74%)	1.1 (0.64-1.8)	2.4% (1.6-3.4%)	3% (1.7-4.8%)	2.1 (1.1-3.3)	230 (150-330)
Trinidad and Tobago	2005	Female	30 to 34	40% (24-58%)	60% (42-76%)	1.1 (0.62-1.7)	2.3% (1.6-3.3%)	3.1% (1.9-4.6%)	2.4 (1.4-3.5)	210 (140-300)
Trinidad and Tobago	2005	Female	35 to 39	39% (25-57%)	61% (43-75%)	1 (0.57-1.6)	2.2% (1.4-3.1%)	3.4% (2.1-4.9%)	3.1 (1.9-4.5)	220 (140-310)
Trinidad and Tobago	2005	Female	40 to 44	38% (24-56%)	62% (44-76%)	0.94 (0.5-1.5)	1.8% (0.97-2.9%)	2.9% (1.5-4.6%)	4.3 (2.2-6.8)	250 (130-390)
Trinidad and Tobago	2005	Female	45 to 49	36% (22-53%)	64% (47-78%)	0.91 (0.48-1.5)	1.6% (0.51-3.1%)	2.6% (1-4.7%)	4.7 (1.8-8.6)	220 (70-410)
Trinidad and Tobago	2005	Female	50 to 54	33% (20-50%)	67% (50-80%)	0.81 (0.42-1.4)	1.1% (-0.12-2.7%)	1.8% (-0.015-4%)	3.8 (-0.032-8.5)	150 (-16-350)
Trinidad and Tobago	2005	Female	55 to 59	29% (17-45%)	71% (55-83%)	0.68 (0.35-1.2)	0.66% (-0.59-2.2%)	1% (-0.62-3%)	2.7 (-1.7-8.1)	87 (-76-290)
Trinidad and Tobago	2005	Female	60 to 64	26% (15-41%)	74% (59-85%)	0.59 (0.29-1)	0.11% (-1.2-1.6%)	0.21% (-1.4-2.1%)	0.7 (-4.6-6.8)	14 (-150-200)
Trinidad and Tobago	2005	Female	65 to 69	23% (14-36%)	77% (64-86%)	0.51 (0.27-0.84)	-0.16% (-1.3-1%)	-0.13% (-1.5-1.4%)	-0.5 (-5.9-5.2)	-19 (-150-120)
Trinidad and Tobago	2005	Female	70 to 74	21% (13-33%)	79% (67-87%)	0.42 (0.21-0.77)	-0.26% (-1.3-0.86%)	-0.27% (-1.6-1.1%)	-1.1 (-6.5-4.5)	-27 (-130-87)
Trinidad and Tobago	2005	Female	75 to 79	19% (11-31%)	81% (69-89%)	0.37 (0.17-0.69)	-0.3% (-1.3-0.69%)	-0.35% (-1.5-0.82%)	-1.7 (-7.5-4)	-27 (-120-61)
Trinidad and Tobago	2005	Female	80 to 84	17% (9.5-28%)	83% (72-90%)	0.34 (0.16-0.59)	-0.19% (-0.93-0.64%)	-0.24% (-1.2-0.77%)	-1.3 (-6.1-4)	-14 (-68-44)
Trinidad and Tobago	2005	Female	85 to 89	17% (9.6-27%)	83% (73-90%)	0.35 (0.16-0.63)	-0.097% (-0.88-0.82%)	-0.15% (-1.1-0.95%)	-0.68 (-4.9-4.2)	-4 (-36-33)
Trinidad and Tobago	2005	Female	90 to 94	19% (11-31%)	81% (69-89%)	0.4 (0.18-0.76)	0.00018% (-0.84-1.2%)	-0.069% (-1.1-1.3%)	-0.19 (-3-3.8)	0.0048 (-14-20)
Trinidad and Tobago	2005	Female	95 plus	23% (12-39%)	77% (61-88%)	0.48 (0.17-1)	0.092% (-0.99-1.7%)	-0.027% (-1.3-1.9%)	-0.028 (-1.4-1.9)	0.38 (-4-7)
Trinidad and Tobago	2005	Male	All Ages	56% (48-64%)	44% (36-52%)	1.5 (1.4-1.7)	4.9% (3.7-6.1%)	4.3% (2.7-6%)	230 (150-330)	11000 (8500-14000)
Trinidad and Tobago	2005	Male	15 to 19	54% (36-71%)	46% (29-64%)	1.1 (0.57-1.7)	4.7% (2.9-7.1%)	6.3% (3.2-10%)	5.7 (2.9-9.4)	520 (310-800)
Trinidad and Tobago	2005	Male	20 to 24	65% (45-80%)	35% (20-55%)	2 (1.3-3)	9.4% (6.3-13%)	11% (6.7-17%)	18 (11-28)	1500 (990-2200)
Trinidad and Tobago	2005	Male	25 to 29	68% (48-82%)	32% (18-52%)	2.5 (1.5-3.6)	10% (6.7-14%)	12% (7.1-17%)	19 (11-27)	1400 (900-1900)
Trinidad and Tobago	2005	Male	30 to 34	69% (49-83%)	31% (17-51%)	2.6 (1.6-3.7)	9.2% (6.3-13%)	11% (7-16%)	18 (11-26)	1200 (830-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	2005	Male	35 to 39	69% (52-83%)	31% (17-48%)	2.5 (1.5-3.6)	7.9% (5.5-11%)	9.6% (6.3-14%)	15 (10-22)	990 (670-1300)
Trinidad and Tobago	2005	Male	40 to 44	67% (51-82%)	33% (18-49%)	2.3 (1.4-3.4)	7.4% (4.9-10%)	9% (5.7-13%)	22 (14-31)	1200 (790-1700)
Trinidad and Tobago	2005	Male	45 to 49	65% (45-81%)	35% (19-55%)	2.2 (1.2-3.5)	6.5% (3.9-9.7%)	7.8% (4.5-12%)	23 (13-36)	1100 (680-1700)
Trinidad and Tobago	2005	Male	50 to 54	64% (45-80%)	36% (20-55%)	2 (1.1-3.1)	5.5% (2.7-8.9%)	6.5% (3.2-11%)	23 (11-37)	980 (470-1600)
Trinidad and Tobago	2005	Male	55 to 59	62% (43-78%)	38% (22-57%)	1.8 (0.97-2.8)	4.5% (1.9-7.7%)	5.3% (2.2-9.1%)	23 (9.3-40)	830 (350-1400)
Trinidad and Tobago	2005	Male	60 to 64	59% (39-77%)	41% (23-61%)	1.6 (0.86-2.8)	3.5% (0.88-6.9%)	4% (1-8.1%)	19 (4.7-38)	590 (140-1200)
Trinidad and Tobago	2005	Male	65 to 69	58% (38-75%)	42% (25-62%)	1.4 (0.79-2.3)	2.9% (0.75-5.7%)	3.3% (0.79-6.5%)	17 (4.1-34)	430 (110-840)
Trinidad and Tobago	2005	Male	70 to 74	54% (35-71%)	46% (29-65%)	1.2 (0.61-2)	2% (0.12-4.9%)	2.2% (0.1-5.4%)	12 (0.55-29)	240 (14-560)
Trinidad and Tobago	2005	Male	75 to 79	49% (30-69%)	51% (31-70%)	0.98 (0.48-1.7)	1.5% (0.025-3.8%)	1.6% (-0.038-4.2%)	9.2 (-0.21-23)	140 (2.3-360)
Trinidad and Tobago	2005	Male	80 to 84	43% (26-62%)	57% (38-74%)	0.82 (0.42-1.3)	1.2% (0.059-2.7%)	1.2% (-0.017-2.9%)	6.2 (-0.089-15)	73 (3.6-160)
Trinidad and Tobago	2005	Male	85 to 89	39% (23-55%)	61% (45-77%)	0.71 (0.36-1.2)	0.9% (-0.023-2.3%)	0.9% (-0.15-2.5%)	3.1 (-0.5-8.6)	27 (-0.68-69)
Trinidad and Tobago	2005	Male	90 to 94	39% (24-56%)	61% (44-76%)	0.72 (0.35-1.3)	0.69% (-0.25-2.2%)	0.63% (-0.43-2.3%)	0.97 (-0.68-3.6)	6.3 (-2.2-20)
Trinidad and Tobago	2005	Male	95 plus	42% (23-62%)	58% (38-77%)	0.78 (0.32-1.5)	0.61% (-0.65-2.5%)	0.44% (-0.98-2.6%)	0.18 (-0.4-1.1)	0.99 (-1-4.4)
Trinidad and Tobago	2010	Female	All Ages	32% (26-39%)	68% (61-74%)	0.74 (0.63-0.88)	1% (0.48-1.6%)	0.66% (-0.15-1.7%)	28 (-6.5-72)	1900 (840-3000)
Trinidad and Tobago	2010	Female	15 to 19	37% (24-53%)	63% (47-76%)	0.78 (0.41-1.3)	1.7% (1-2.7%)	3.3% (1.6-5.9%)	0.94 (0.45-1.7)	110 (63-170)
Trinidad and Tobago	2010	Female	20 to 24	43% (27-59%)	57% (41-73%)	1.2 (0.64-1.9)	2.8% (1.8-4%)	4.2% (2.3-6.9%)	2 (1.1-3.3)	240 (150-360)
Trinidad and Tobago	2010	Female	25 to 29	42% (26-59%)	58% (41-74%)	1.2 (0.68-1.8)	2.5% (1.7-3.5%)	3.3% (1.8-5.2%)	2.3 (1.2-3.6)	260 (170-370)
Trinidad and Tobago	2010	Female	30 to 34	41% (25-58%)	59% (42-75%)	1.1 (0.66-1.8)	2.5% (1.8-3.5%)	3.6% (2.3-5.4%)	2.7 (1.7-4.1)	240 (160-340)
Trinidad and Tobago	2010	Female	35 to 39	40% (25-58%)	60% (42-75%)	1.1 (0.63-1.7)	2.3% (1.5-3.3%)	3.7% (2.4-5.5%)	3.1 (2-4.6)	220 (150-320)
Trinidad and Tobago	2010	Female	40 to 44	38% (24-57%)	62% (43-76%)	1 (0.53-1.6)	2% (1.1-3.2%)	3.4% (1.9-5.5%)	3.8 (2.1-6)	220 (120-350)
Trinidad and Tobago	2010	Female	45 to 49	37% (22-53%)	63% (47-78%)	0.96 (0.5-1.6)	1.8% (0.69-3.2%)	3% (1.4-5.3%)	5.4 (2.6-9.7)	260 (100-470)
Trinidad and Tobago	2010	Female	50 to 54	33% (20-50%)	67% (50-80%)	0.86 (0.46-1.5)	1.3% (0.085-3%)	2.2% (0.48-4.4%)	5 (1.1-10)	200 (13-430)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	2010	Female	55 to 59	29% (17-44%)	71% (56-83%)	0.73 (0.37-1.2)	0.84% (-0.37-2.3%)	1.4% (-0.22-3.3%)	3.7 (-0.59-8.8)	120 (-52-320)
Trinidad and Tobago	2010	Female	60 to 64	26% (15-41%)	74% (59-85%)	0.62 (0.32-1.1)	0.26% (-0.95-1.7%)	0.43% (-1.1-2.3%)	1.5 (-3.9-7.8)	36 (-140-240)
Trinidad and Tobago	2010	Female	65 to 69	24% (15-37%)	76% (63-85%)	0.54 (0.28-0.9)	0.021% (-1.1-1.2%)	0.12% (-1.3-1.6%)	0.47 (-5.2-6.8)	2.7 (-140-160)
Trinidad and Tobago	2010	Female	70 to 74	21% (13-34%)	79% (66-87%)	0.45 (0.23-0.85)	-0.15% (-1.2-1%)	-0.14% (-1.5-1.3%)	-0.58 (-6.3-5.5)	-17 (-130-110)
Trinidad and Tobago	2010	Female	75 to 79	19% (11-31%)	81% (69-89%)	0.4 (0.18-0.73)	-0.21% (-1.2-0.86%)	-0.25% (-1.5-1.1%)	-1.2 (-7.3-5)	-19 (-110-74)
Trinidad and Tobago	2010	Female	80 to 84	17% (9.9-28%)	83% (72-90%)	0.37 (0.18-0.63)	-0.11% (-0.87-0.81%)	-0.14% (-1.1-0.99%)	-0.73 (-5.7-5.1)	-7.6 (-63-59)
Trinidad and Tobago	2010	Female	85 to 89	17% (10-27%)	83% (73-90%)	0.37 (0.17-0.68)	0.004% (-0.74-1%)	-0.039% (-0.95-1.2%)	-0.18 (-4.2-5.4)	0.18 (-32-44)
Trinidad and Tobago	2010	Female	90 to 94	19% (11-31%)	81% (69-89%)	0.43 (0.18-0.81)	0.085% (-0.8-1.3%)	0.018% (-1.1-1.5%)	0.051 (-3.1-4.4)	1.6 (-14-24)
Trinidad and Tobago	2010	Female	95 plus	23% (12-39%)	77% (61-88%)	0.52 (0.19-1.1)	0.18% (-0.95-2%)	0.046% (-1.3-2.2%)	0.05 (-1.5-2.4)	0.82 (-4.4-9)
Trinidad and Tobago	2010	Male	All Ages	57% (49-65%)	43% (35-51%)	1.7 (1.5-2)	5.5% (4.2-7%)	4.8% (3.2-6.7%)	280 (190-390)	13000 (9900-17000)
Trinidad and Tobago	2010	Male	15 to 19	54% (36-71%)	46% (29-64%)	1.1 (0.6-1.9)	5.1% (3.1-8%)	6.8% (3.4-11%)	5.3 (2.6-8.8)	460 (260-720)
Trinidad and Tobago	2010	Male	20 to 24	65% (46-80%)	35% (20-54%)	2.2 (1.3-3.3)	10% (6.6-15%)	12% (6.9-18%)	20 (12-31)	1600 (990-2300)
Trinidad and Tobago	2010	Male	25 to 29	68% (49-82%)	32% (18-51%)	2.6 (1.6-3.8)	11% (7.2-15%)	13% (7.7-18%)	25 (15-35)	1800 (1200-2500)
Trinidad and Tobago	2010	Male	30 to 34	69% (50-83%)	31% (17-50%)	2.8 (1.8-4)	10% (6.9-14%)	12% (7.6-18%)	22 (14-32)	1500 (1000-2100)
Trinidad and Tobago	2010	Male	35 to 39	69% (52-83%)	31% (17-48%)	2.6 (1.6-3.9)	8.8% (5.9-12%)	11% (7.1-16%)	18 (11-26)	1100 (770-1600)
Trinidad and Tobago	2010	Male	40 to 44	68% (51-82%)	32% (18-49%)	2.5 (1.5-3.8)	8% (5.1-11%)	10% (6.2-14%)	19 (12-27)	1100 (670-1500)
Trinidad and Tobago	2010	Male	45 to 49	66% (45-81%)	34% (19-55%)	2.3 (1.3-3.5)	7.1% (4.2-10%)	8.7% (5.1-13%)	26 (15-40)	1300 (790-1900)
Trinidad and Tobago	2010	Male	50 to 54	64% (44-80%)	36% (20-56%)	2.1 (1.2-3.3)	6.1% (3.2-9.5%)	7.3% (3.8-12%)	27 (14-44)	1200 (620-1800)
Trinidad and Tobago	2010	Male	55 to 59	62% (43-78%)	38% (22-57%)	1.9 (1.1-3)	5.2% (2.6-8.3%)	6.3% (3.1-10%)	27 (13-45)	1000 (490-1600)
Trinidad and Tobago	2010	Male	60 to 64	60% (40-78%)	40% (22-60%)	1.7 (0.95-2.9)	4.1% (1.4-7.5%)	4.8% (1.6-8.8%)	25 (8.5-48)	790 (260-1400)
Trinidad and Tobago	2010	Male	65 to 69	58% (39-75%)	42% (25-61%)	1.5 (0.87-2.5)	3.4% (1.2-6.3%)	3.9% (1.4-7.4%)	22 (8-41)	560 (210-1000)
Trinidad and Tobago	2010	Male	70 to 74	55% (36-72%)	45% (28-64%)	1.3 (0.67-2.2)	2.4% (0.47-5.3%)	2.6% (0.48-6%)	15 (2.8-34)	310 (65-700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	2010	Male	75 to 79	50% (31-69%)	50% (31-69%)	1.1 (0.51-1.9)	1.8% (0.26-4.3%)	2% (0.25-4.7%)	12 (1.5-28)	180 (26-420)
Trinidad and Tobago	2010	Male	80 to 84	44% (26-62%)	56% (38-74%)	0.88 (0.46-1.5)	1.4% (0.24-3.2%)	1.5% (0.19-3.4%)	7.5 (0.97-18)	89 (16-200)
Trinidad and Tobago	2010	Male	85 to 89	39% (24-56%)	61% (44-76%)	0.76 (0.38-1.3)	1.1% (0.12-2.7%)	1.1% (0.037-2.9%)	4.5 (0.15-12)	38 (4.3-97)
Trinidad and Tobago	2010	Male	90 to 94	39% (24-56%)	61% (44-76%)	0.77 (0.37-1.4)	0.87% (-0.16-2.7%)	0.82% (-0.33-2.8%)	1.6 (-0.64-5.4)	9.9 (-1.9-30)
Trinidad and Tobago	2010	Male	95 plus	42% (23-63%)	58% (37-77%)	0.84 (0.34-1.6)	0.79% (-0.56-2.9%)	0.64% (-0.85-3.1%)	0.33 (-0.44-1.6)	1.6 (-1.1-5.8)
Trinidad and Tobago	2016	Female	All Ages	32% (26-39%)	68% (61-74%)	0.8 (0.6-1)	1.1% (0.44-2%)	0.86% (-0.18-2.3%)	40 (-7.9-110)	2200 (820-3800)
Trinidad and Tobago	2016	Female	15 to 19	37% (24-53%)	63% (47-76%)	0.85 (0.41-1.6)	1.8% (1-2.8%)	3.6% (1.6-6.7%)	0.79 (0.35-1.5)	90 (50-140)
Trinidad and Tobago	2016	Female	20 to 24	42% (27-59%)	58% (41-73%)	1.3 (0.65-2.1)	2.9% (1.9-4.3%)	4.6% (2.4-7.7%)	1.6 (0.77-2.6)	190 (120-280)
Trinidad and Tobago	2016	Female	25 to 29	42% (26-59%)	58% (41-74%)	1.3 (0.69-2.1)	2.7% (1.8-3.8%)	3.7% (2-5.9%)	2.1 (1.1-3.4)	240 (150-350)
Trinidad and Tobago	2016	Female	30 to 34	40% (25-57%)	60% (43-75%)	1.3 (0.7-2)	2.6% (1.7-3.8%)	3.9% (2.4-5.9%)	3.3 (1.9-4.9)	300 (200-430)
Trinidad and Tobago	2016	Female	35 to 39	40% (25-57%)	60% (43-75%)	1.2 (0.65-1.9)	2.4% (1.5-3.8%)	4.1% (2.5-6.2%)	3.6 (2.1-5.5)	260 (160-400)
Trinidad and Tobago	2016	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.1 (0.56-1.8)	2.2% (1.2-3.6%)	3.8% (2-6.1%)	4.1 (2.2-6.9)	240 (130-410)
Trinidad and Tobago	2016	Female	45 to 49	36% (22-53%)	64% (47-78%)	1 (0.53-1.8)	2% (0.86-3.7%)	3.3% (1.6-6.1%)	4.8 (2.3-8.9)	240 (100-440)
Trinidad and Tobago	2016	Female	50 to 54	33% (19-50%)	67% (50-81%)	0.93 (0.49-1.6)	1.6% (0.27-3.5%)	2.5% (0.57-5.4%)	6.1 (1.4-13)	250 (43-580)
Trinidad and Tobago	2016	Female	55 to 59	29% (17-44%)	71% (56-83%)	0.79 (0.39-1.3)	1.1% (-0.26-2.9%)	1.7% (-0.15-4.1%)	5.2 (-0.46-13)	180 (-42-460)
Trinidad and Tobago	2016	Female	60 to 64	26% (15-41%)	74% (59-85%)	0.68 (0.33-1.2)	0.54% (-0.78-2.2%)	0.78% (-0.93-2.9%)	3.1 (-3.8-11)	89 (-130-360)
Trinidad and Tobago	2016	Female	65 to 69	23% (14-37%)	77% (63-86%)	0.59 (0.3-0.99)	0.26% (-0.96-1.7%)	0.41% (-1.2-2.3%)	1.8 (-5.4-10)	39 (-140-260)
Trinidad and Tobago	2016	Female	70 to 74	21% (13-33%)	79% (67-87%)	0.5 (0.25-0.96)	0.076% (-1.1-1.6%)	0.14% (-1.3-2%)	0.71 (-6.7-10)	9.9 (-140-200)
Trinidad and Tobago	2016	Female	75 to 79	19% (11-31%)	81% (69-89%)	0.43 (0.2-0.85)	-0.004% (-1-1.3%)	-0.0019% (-1.3-1.6%)	-0.017 (-7.3-8.8)	-0.49 (-110-140)
Trinidad and Tobago	2016	Female	80 to 84	17% (9.7-27%)	83% (73-90%)	0.4 (0.19-0.71)	0.087% (-0.75-1.2%)	0.091% (-0.93-1.5%)	0.49 (-5.3-8.2)	6.7 (-58-95)
Trinidad and Tobago	2016	Female	85 to 89	17% (9.8-27%)	83% (73-90%)	0.41 (0.18-0.74)	0.19% (-0.6-1.4%)	0.17% (-0.8-1.6%)	0.86 (-4-7.8)	8.8 (-29-64)
Trinidad and Tobago	2016	Female	90 to 94	19% (11-30%)	81% (70-89%)	0.46 (0.2-0.9)	0.28% (-0.7-1.8%)	0.25% (-0.93-2%)	0.88 (-3.3-7)	6.3 (-15-40)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Trinidad and Tobago	2016	Female	95 plus	23% (12-38%)	77% (62-88%)	0.56 (0.2-1.2)	0.41% (-0.84-2.6%)	0.32% (-1.2-2.9%)	0.43 (-1.6-3.9)	2.3 (-4.6-14)
Trinidad and Tobago	2016	Male	All Ages	57% (50-65%)	43% (35-50%)	1.9 (1.4-2.4)	5.8% (4-7.7%)	5.1% (3.1-7.7%)	320 (200-490)	14000 (9900-19000)
Trinidad and Tobago	2016	Male	15 to 19	54% (36-71%)	46% (29-64%)	1.3 (0.61-2.2)	5.2% (3.1-7.9%)	7.1% (3.5-12%)	4.1 (2.1-6.8)	370 (210-580)
Trinidad and Tobago	2016	Male	20 to 24	65% (47-80%)	35% (20-53%)	2.4 (1.3-3.7)	11% (6.8-15%)	13% (7.3-19%)	14 (8.1-22)	1200 (730-1700)
Trinidad and Tobago	2016	Male	25 to 29	68% (49-82%)	32% (18-51%)	2.9 (1.6-4.3)	11% (7.4-16%)	14% (7.9-20%)	21 (12-31)	1600 (1000-2200)
Trinidad and Tobago	2016	Male	30 to 34	69% (49-83%)	31% (17-51%)	3 (1.8-4.6)	11% (7.1-15%)	13% (8-19%)	26 (16-39)	1800 (1200-2600)
Trinidad and Tobago	2016	Male	35 to 39	69% (52-82%)	31% (18-48%)	2.8 (1.7-4.5)	9.4% (6.2-13%)	12% (7.2-17%)	22 (13-32)	1400 (890-1900)
Trinidad and Tobago	2016	Male	40 to 44	67% (51-82%)	33% (18-49%)	2.7 (1.5-4.3)	8.7% (5.4-12%)	11% (6.5-16%)	22 (13-32)	1200 (750-1700)
Trinidad and Tobago	2016	Male	45 to 49	65% (44-81%)	35% (19-56%)	2.5 (1.4-4.1)	7.7% (4.4-12%)	9.6% (5.3-15%)	24 (13-38)	1200 (680-1800)
Trinidad and Tobago	2016	Male	50 to 54	64% (44-80%)	36% (20-56%)	2.3 (1.3-3.7)	6.6% (3.3-10%)	8.1% (4.1-13%)	32 (16-53)	1400 (710-2300)
Trinidad and Tobago	2016	Male	55 to 59	61% (43-78%)	39% (22-57%)	2 (1.1-3.2)	5.7% (2.6-9.2%)	6.8% (3.1-11%)	36 (17-59)	1300 (610-2200)
Trinidad and Tobago	2016	Male	60 to 64	59% (39-77%)	41% (23-61%)	1.8 (0.98-3.1)	4.6% (1.5-8.1%)	5.3% (1.7-9.5%)	34 (11-61)	1100 (340-1900)
Trinidad and Tobago	2016	Male	65 to 69	58% (39-75%)	42% (25-61%)	1.6 (0.9-2.7)	3.9% (1.4-7.4%)	4.5% (1.5-8.5%)	31 (11-58)	780 (280-1500)
Trinidad and Tobago	2016	Male	70 to 74	54% (36-71%)	46% (29-64%)	1.4 (0.71-2.5)	2.8% (0.61-6.2%)	3.1% (0.58-7%)	23 (4.3-50)	460 (95-1000)
Trinidad and Tobago	2016	Male	75 to 79	49% (30-68%)	51% (32-70%)	1.1 (0.54-2)	2.2% (0.32-5.3%)	2.4% (0.31-5.8%)	17 (2.2-41)	260 (40-620)
Trinidad and Tobago	2016	Male	80 to 84	43% (26-62%)	57% (38-74%)	0.94 (0.45-1.6)	1.7% (0.31-4%)	1.8% (0.24-4.4%)	10 (1.4-26)	120 (23-300)
Trinidad and Tobago	2016	Male	85 to 89	39% (23-55%)	61% (45-77%)	0.82 (0.38-1.5)	1.3% (0.16-3.3%)	1.4% (0.072-3.6%)	5.8 (0.32-15)	49 (5.9-120)
Trinidad and Tobago	2016	Male	90 to 94	39% (24-56%)	61% (44-76%)	0.82 (0.38-1.6)	1.1% (-0.07-3.2%)	1.1% (-0.25-3.4%)	2.3 (-0.53-7.1)	14 (-0.87-39)
Trinidad and Tobago	2016	Male	95 plus	42% (23-62%)	58% (38-77%)	0.89 (0.35-1.8)	1.1% (-0.37-3.6%)	0.91% (-0.75-3.8%)	0.5 (-0.4-2.1)	2.2 (-0.77-8)
Bolivia	1990	Female	All Ages	42% (38-46%)	58% (54-62%)	0.68 (0.55-0.83)	1.2% (0.67-1.7%)	1.8% (0.79-2.9%)	580 (260-960)	24000 (14000-36000)
Bolivia	1990	Female	15 to 19	37% (27-45%)	63% (55-73%)	0.51 (0.29-0.8)	1.7% (0.48-3.2%)	2.4% (0.21-5.1%)	13 (1-27)	1100 (310-2200)
Bolivia	1990	Female	20 to 24	57% (48-65%)	43% (35-52%)	1.1 (0.65-1.6)	3.8% (1.5-6.7%)	5.3% (1.4-9.7%)	33 (8.6-63)	2600 (1000-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	1990	Female	25 to 29	63% (54-70%)	37% (30-46%)	1.3 (0.86-1.9)	4.4% (2.1-7.2%)	6.3% (2.6-11%)	39 (16-66)	2800 (1300-4500)
Bolivia	1990	Female	30 to 34	66% (56-72%)	34% (28-44%)	1.4 (0.93-2.1)	4.5% (2.2-7.5%)	6.6% (3-11%)	43 (19-74)	2700 (1300-4400)
Bolivia	1990	Female	35 to 39	66% (56-72%)	34% (28-44%)	1.4 (0.87-2.1)	4.2% (2-7%)	6.2% (2.7-10%)	46 (20-77)	2600 (1200-4100)
Bolivia	1990	Female	40 to 44	65% (57-71%)	35% (29-43%)	1.4 (0.87-2.1)	4.2% (2-6.9%)	6.3% (2.9-10%)	45 (20-75)	2200 (990-3600)
Bolivia	1990	Female	45 to 49	64% (54-71%)	36% (29-46%)	1.4 (0.88-2.2)	4.1% (1.7-7.2%)	6.1% (2.7-11%)	51 (22-90)	2100 (890-3800)
Bolivia	1990	Female	50 to 54	62% (51-71%)	38% (29-49%)	1.4 (0.83-2.2)	4.1% (1.7-7.3%)	5.9% (2.6-10%)	55 (24-95)	2000 (830-3500)
Bolivia	1990	Female	55 to 59	60% (44-76%)	40% (24-56%)	1.4 (0.78-2.3)	3.9% (1.4-7%)	5.4% (2-9.4%)	65 (24-120)	2000 (730-3700)
Bolivia	1990	Female	60 to 64	57% (40-76%)	43% (24-60%)	1.3 (0.74-2.2)	3.1% (0.7-6.4%)	4.1% (1.1-8.1%)	70 (17-140)	1900 (430-3800)
Bolivia	1990	Female	65 to 69	53% (37-70%)	47% (30-63%)	1.2 (0.66-1.8)	2.1% (0.017-4.6%)	2.8% (0.21-5.7%)	58 (4.2-120)	1300 (9.3-2700)
Bolivia	1990	Female	70 to 74	48% (33-64%)	52% (36-67%)	0.96 (0.52-1.6)	1.3% (-0.51-3.4%)	1.7% (-0.55-4.3%)	42 (-14-110)	720 (-290-1900)
Bolivia	1990	Female	75 to 79	42% (28-59%)	58% (41-72%)	0.75 (0.37-1.3)	0.61% (-0.99-2.5%)	0.77% (-1.1-3%)	19 (-27-74)	250 (-410-1000)
Bolivia	1990	Female	80 to 84	36% (22-55%)	64% (45-78%)	0.62 (0.32-1)	0.24% (-0.98-1.6%)	0.29% (-1.2-1.9%)	4.8 (-19-31)	48 (-200-320)
Bolivia	1990	Female	85 to 89	34% (21-49%)	66% (51-79%)	0.56 (0.3-0.96)	-0.057% (-1.1-1.1%)	-0.087% (-1.3-1.3%)	-0.84 (-13-12)	-4.8 (-92-89)
Bolivia	1990	Female	90 to 94	35% (23-51%)	65% (49-77%)	0.56 (0.29-0.99)	-0.35% (-1.3-0.71%)	-0.46% (-1.7-0.8%)	-1.7 (-6.4-3.1)	-7.8 (-31-16)
Bolivia	1990	Female	95 plus	39% (23-58%)	61% (42-77%)	0.58 (0.24-1.1)	-0.56% (-1.8-0.79%)	-0.76% (-2.2-0.85%)	-0.86 (-2.6-0.95)	-2.5 (-7.9-3.5)
Bolivia	1990	Male	All Ages	46% (41-51%)	54% (49-59%)	1.1 (0.94-1.4)	3.1% (2.4-3.9%)	4.6% (3.3-6%)	1600 (1100-2100)	69000 (53000-87000)
Bolivia	1990	Male	15 to 19	38% (28-47%)	62% (53-72%)	0.6 (0.33-0.97)	3.5% (2-5.2%)	4.3% (2-7.1%)	24 (11-40)	2300 (1300-3500)
Bolivia	1990	Male	20 to 24	73% (63-79%)	27% (21-37%)	1.7 (1.1-2.4)	9.5% (6.2-13%)	12% (6.8-17%)	86 (48-130)	7000 (4500-9900)
Bolivia	1990	Male	25 to 29	80% (73-85%)	20% (15-27%)	2.2 (1.4-3.2)	12% (8.2-16%)	15% (9.4-20%)	110 (68-160)	8100 (5400-11000)
Bolivia	1990	Male	30 to 34	82% (74-86%)	18% (14-26%)	2.5 (1.7-3.6)	13% (8.7-17%)	16% (11-22%)	130 (80-180)	8100 (5600-11000)
Bolivia	1990	Male	35 to 39	78% (70-83%)	22% (17-30%)	2.6 (1.6-3.7)	13% (8.8-18%)	16% (11-23%)	140 (90-200)	8100 (5400-11000)
Bolivia	1990	Male	40 to 44	77% (70-83%)	23% (17-30%)	2.6 (1.6-3.9)	13% (8.6-17%)	16% (11-22%)	140 (90-200)	7000 (4700-9700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	1990	Male	45 to 49	79% (70-84%)	21% (16-30%)	2.7 (1.7-4.1)	12% (8-17%)	16% (10-22%)	160 (98-230)	6900 (4500-9900)
Bolivia	1990	Male	50 to 54	76% (69-82%)	24% (18-31%)	2.7 (1.6-3.9)	11% (7.1-16%)	14% (8.8-20%)	150 (90-220)	5700 (3500-8200)
Bolivia	1990	Male	55 to 59	70% (61-77%)	30% (23-39%)	2.5 (1.5-3.8)	9.8% (6-14%)	12% (7.4-18%)	150 (92-230)	5100 (3100-7600)
Bolivia	1990	Male	60 to 64	70% (59-77%)	30% (23-41%)	2.4 (1.4-3.8)	8.1% (4.4-13%)	9.8% (5.3-16%)	160 (85-260)	4500 (2500-7200)
Bolivia	1990	Male	65 to 69	73% (61-83%)	27% (17-39%)	2.1 (1.3-3.2)	6.1% (3.1-9.9%)	7.4% (3.8-12%)	140 (66-220)	3200 (1600-5100)
Bolivia	1990	Male	70 to 74	71% (56-83%)	29% (17-44%)	1.7 (0.97-2.6)	4.2% (1.6-7.3%)	4.9% (1.7-8.5%)	100 (36-180)	1900 (710-3300)
Bolivia	1990	Male	75 to 79	61% (43-78%)	39% (22-57%)	1.2 (0.61-2)	2.7% (0.71-5.2%)	3.2% (0.72-6.2%)	65 (14-130)	940 (230-1800)
Bolivia	1990	Male	80 to 84	53% (34-70%)	47% (30-66%)	0.92 (0.5-1.5)	1.8% (0.43-3.4%)	2% (0.4-3.9%)	28 (5.3-54)	310 (73-570)
Bolivia	1990	Male	85 to 89	49% (32-65%)	51% (35-68%)	0.8 (0.44-1.4)	1.1% (0.099-2.3%)	1.2% (0.007-2.6%)	9.3 (0.055-20)	74 (6.6-160)
Bolivia	1990	Male	90 to 94	49% (33-66%)	51% (34-67%)	0.78 (0.4-1.4)	0.64% (-0.36-1.8%)	0.59% (-0.62-1.9%)	1.7 (-1.7-5.5)	11 (-6-30)
Bolivia	1990	Male	95 plus	51% (30-71%)	49% (29-70%)	0.8 (0.35-1.5)	0.29% (-0.89-1.7%)	0.083% (-1.3-1.8%)	0.062 (-0.96-1.3)	0.85 (-2.6-5)
Bolivia	1995	Female	All Ages	42% (38-47%)	58% (53-62%)	0.63 (0.51-0.76)	1.1% (0.59-1.6%)	1.5% (0.56-2.6%)	470 (170-820)	20000 (11000-31000)
Bolivia	1995	Female	15 to 19	36% (27-45%)	64% (55-73%)	0.46 (0.26-0.73)	1.4% (0.45-2.6%)	2% (0.27-4.1%)	10 (1.3-22)	1000 (320-1800)
Bolivia	1995	Female	20 to 24	57% (47-65%)	43% (35-53%)	0.97 (0.59-1.4)	3.1% (1.2-5.3%)	4.3% (1-7.9%)	26 (6.3-49)	2200 (910-3800)
Bolivia	1995	Female	25 to 29	63% (54-70%)	37% (30-46%)	1.2 (0.77-1.8)	3.6% (1.7-5.8%)	5.2% (2-9.1%)	30 (11-54)	2300 (1100-3700)
Bolivia	1995	Female	30 to 34	66% (56-72%)	34% (28-44%)	1.3 (0.85-1.9)	3.6% (1.8-5.9%)	5.4% (2.4-9.3%)	33 (15-57)	2100 (1000-3500)
Bolivia	1995	Female	35 to 39	66% (57-71%)	34% (29-43%)	1.3 (0.79-1.9)	3.4% (1.6-5.8%)	5.2% (2.2-8.9%)	36 (16-62)	2000 (920-3400)
Bolivia	1995	Female	40 to 44	65% (56-71%)	35% (29-44%)	1.3 (0.78-1.9)	3.4% (1.6-5.7%)	5.3% (2.3-8.9%)	41 (18-69)	2000 (920-3300)
Bolivia	1995	Female	45 to 49	64% (54-71%)	36% (29-46%)	1.3 (0.78-2)	3.4% (1.3-5.8%)	5.2% (2.1-8.8%)	41 (17-71)	1700 (730-3000)
Bolivia	1995	Female	50 to 54	62% (52-71%)	38% (29-48%)	1.3 (0.76-2)	3.4% (1.4-6.1%)	5% (2.2-8.9%)	50 (22-87)	1800 (750-3200)
Bolivia	1995	Female	55 to 59	60% (44-76%)	40% (24-56%)	1.3 (0.72-2.1)	3.3% (1.1-6%)	4.6% (1.6-8.3%)	54 (19-99)	1700 (550-3200)
Bolivia	1995	Female	60 to 64	57% (40-75%)	43% (25-60%)	1.2 (0.66-2)	2.5% (0.41-5.3%)	3.4% (0.65-7%)	55 (10-110)	1500 (230-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	1995	Female	65 to 69	53% (36-71%)	47% (29-64%)	1.1 (0.61-1.7)	1.7% (-0.17-3.8%)	2.2% (-0.084-4.8%)	50 (-1.7-110)	1100 (-110-2400)
Bolivia	1995	Female	70 to 74	48% (32-64%)	52% (36-68%)	0.86 (0.47-1.4)	0.97% (-0.7-2.9%)	1.3% (-0.74-3.5%)	34 (-20-98)	570 (-400-1700)
Bolivia	1995	Female	75 to 79	41% (27-59%)	59% (41-73%)	0.67 (0.34-1.2)	0.4% (-0.98-1.9%)	0.53% (-1.1-2.4%)	14 (-28-66)	180 (-420-900)
Bolivia	1995	Female	80 to 84	36% (22-53%)	64% (47-78%)	0.55 (0.29-0.94)	0.086% (-1-1.2%)	0.12% (-1.2-1.5%)	2.5 (-25-31)	23 (-270-320)
Bolivia	1995	Female	85 to 89	34% (21-48%)	66% (52-79%)	0.51 (0.27-0.87)	-0.16% (-1.2-0.85%)	-0.21% (-1.4-0.98%)	-2.8 (-19-13)	-19 (-140-98)
Bolivia	1995	Female	90 to 94	35% (23-50%)	65% (50-77%)	0.51 (0.26-0.91)	-0.44% (-1.4-0.49%)	-0.57% (-1.8-0.55%)	-2.9 (-9.1-2.8)	-13 (-43-15)
Bolivia	1995	Female	95 plus	38% (23-57%)	62% (43-77%)	0.53 (0.23-1)	-0.62% (-1.8-0.55%)	-0.84% (-2.3-0.56%)	-1.2 (-3.2-0.77)	-3.4 (-9.9-3)
Bolivia	1995	Male	All Ages	47% (42-51%)	53% (49-58%)	1.1 (0.89-1.3)	3% (2.3-3.8%)	4.3% (3-5.7%)	1400 (960-1900)	62000 (46000-78000)
Bolivia	1995	Male	15 to 19	38% (28-47%)	62% (53-72%)	0.56 (0.31-0.88)	3.2% (1.9-4.7%)	3.9% (1.9-6.4%)	22 (11-37)	2200 (1300-3300)
Bolivia	1995	Male	20 to 24	72% (63-79%)	28% (21-37%)	1.5 (0.99-2.2)	8.6% (5.7-12%)	10% (6.1-15%)	74 (44-110)	6400 (4100-8800)
Bolivia	1995	Male	25 to 29	80% (73-85%)	20% (15-27%)	2.1 (1.3-3)	11% (7.2-14%)	13% (8.4-18%)	93 (58-130)	7100 (4800-9400)
Bolivia	1995	Male	30 to 34	82% (75-86%)	18% (14-25%)	2.4 (1.6-3.3)	11% (7.4-14%)	14% (8.8-19%)	100 (63-140)	6800 (4600-9100)
Bolivia	1995	Male	35 to 39	78% (71-83%)	22% (17-29%)	2.4 (1.5-3.5)	11% (7.3-15%)	14% (9.2-20%)	110 (71-170)	6600 (4400-9300)
Bolivia	1995	Male	40 to 44	77% (69-82%)	23% (18-31%)	2.4 (1.5-3.6)	11% (7.3-15%)	14% (9.3-20%)	130 (81-180)	6500 (4300-9100)
Bolivia	1995	Male	45 to 49	78% (70-84%)	22% (16-30%)	2.5 (1.5-3.9)	11% (6.9-15%)	14% (8.9-20%)	130 (79-190)	5800 (3700-8200)
Bolivia	1995	Male	50 to 54	76% (69-82%)	24% (18-31%)	2.5 (1.5-3.7)	9.7% (6.2-14%)	13% (8-18%)	140 (87-210)	5500 (3500-7900)
Bolivia	1995	Male	55 to 59	70% (61-77%)	30% (23-39%)	2.3 (1.4-3.6)	8.7% (5.3-13%)	11% (6.7-16%)	130 (78-200)	4500 (2700-6600)
Bolivia	1995	Male	60 to 64	69% (59-77%)	31% (23-41%)	2.2 (1.3-3.6)	7.1% (3.9-11%)	8.8% (4.8-14%)	140 (74-220)	3900 (2100-6300)
Bolivia	1995	Male	65 to 69	73% (60-83%)	27% (17-40%)	2 (1.2-3)	5.3% (2.6-8.8%)	6.5% (3.2-11%)	130 (63-210)	3000 (1500-4900)
Bolivia	1995	Male	70 to 74	71% (56-83%)	29% (17-44%)	1.5 (0.88-2.4)	3.6% (1.5-6.1%)	4.3% (1.7-7.3%)	98 (38-170)	1800 (760-3100)
Bolivia	1995	Male	75 to 79	61% (43-78%)	39% (22-57%)	1.1 (0.56-1.8)	2.3% (0.74-4.4%)	2.7% (0.78-5.2%)	60 (17-120)	870 (270-1700)
Bolivia	1995	Male	80 to 84	53% (34-69%)	47% (31-66%)	0.84 (0.46-1.4)	1.5% (0.37-2.8%)	1.7% (0.34-3.3%)	30 (5.7-56)	320 (75-600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	1995	Male	85 to 89	48% (31-65%)	52% (35-69%)	0.74 (0.4-1.3)	0.97% (0.025-2%)	1% (-0.067-2.2%)	11 (-0.67-23)	87 (2.3-180)
Bolivia	1995	Male	90 to 94	49% (33-66%)	51% (34-67%)	0.73 (0.38-1.3)	0.54% (-0.31-1.5%)	0.48% (-0.6-1.7%)	1.8 (-2.3-6.2)	12 (-7.1-35)
Bolivia	1995	Male	95 plus	52% (31-71%)	48% (29-69%)	0.75 (0.33-1.4)	0.26% (-0.85-1.5%)	0.051% (-1.3-1.5%)	0.046 (-1.1-1.3)	0.93 (-2.9-5.2)
Bolivia	2000	Female	All Ages	42% (38-47%)	58% (53-62%)	0.62 (0.5-0.76)	0.96% (0.56-1.4%)	1.3% (0.51-2.3%)	380 (140-660)	16000 (9200-25000)
Bolivia	2000	Female	15 to 19	36% (27-44%)	64% (56-73%)	0.44 (0.25-0.69)	1.2% (0.5-2%)	1.8% (0.34-3.5%)	7.2 (1.4-14)	780 (320-1300)
Bolivia	2000	Female	20 to 24	56% (46-64%)	44% (36-54%)	0.93 (0.57-1.4)	2.6% (1.2-4.2%)	3.7% (1.1-6.7%)	18 (5.2-33)	1700 (830-2800)
Bolivia	2000	Female	25 to 29	63% (53-70%)	37% (30-47%)	1.2 (0.73-1.7)	2.9% (1.6-4.5%)	4.5% (1.8-7.6%)	21 (8.4-36)	1800 (970-2800)
Bolivia	2000	Female	30 to 34	65% (56-72%)	35% (28-44%)	1.3 (0.81-1.8)	2.9% (1.6-4.6%)	4.7% (2.1-8%)	23 (10-39)	1600 (880-2600)
Bolivia	2000	Female	35 to 39	65% (55-71%)	35% (29-45%)	1.2 (0.77-1.8)	2.8% (1.5-4.5%)	4.6% (2.3-7.5%)	27 (13-44)	1600 (850-2400)
Bolivia	2000	Female	40 to 44	65% (55-71%)	35% (29-45%)	1.2 (0.75-1.9)	2.8% (1.3-4.6%)	4.7% (2.2-7.7%)	31 (15-51)	1500 (760-2500)
Bolivia	2000	Female	45 to 49	64% (54-71%)	36% (29-46%)	1.3 (0.75-1.9)	2.8% (1.3-4.9%)	4.6% (2.1-7.9%)	37 (17-65)	1600 (700-2700)
Bolivia	2000	Female	50 to 54	62% (51-71%)	38% (29-49%)	1.2 (0.7-1.9)	2.8% (1.1-5.1%)	4.5% (2-7.8%)	39 (17-67)	1400 (570-2500)
Bolivia	2000	Female	55 to 59	59% (42-75%)	41% (25-58%)	1.2 (0.66-2)	2.8% (0.85-5.1%)	4.2% (1.4-7.6%)	46 (16-84)	1500 (450-2700)
Bolivia	2000	Female	60 to 64	56% (39-74%)	44% (26-61%)	1.2 (0.63-2)	2.2% (0.43-4.5%)	3.2% (0.83-6.3%)	43 (11-87)	1100 (230-2400)
Bolivia	2000	Female	65 to 69	52% (35-69%)	48% (31-65%)	1 (0.56-1.6)	1.5% (-0.033-3.4%)	2.2% (0.16-4.5%)	40 (3-85)	830 (-18-1900)
Bolivia	2000	Female	70 to 74	47% (31-64%)	53% (36-69%)	0.82 (0.45-1.3)	0.93% (-0.51-2.5%)	1.3% (-0.47-3.3%)	33 (-11-85)	540 (-300-1500)
Bolivia	2000	Female	75 to 79	40% (27-58%)	60% (42-73%)	0.64 (0.32-1.1)	0.42% (-0.77-1.8%)	0.6% (-0.87-2.3%)	16 (-22-59)	190 (-360-800)
Bolivia	2000	Female	80 to 84	35% (22-53%)	65% (47-78%)	0.53 (0.28-0.9)	0.13% (-0.89-1.1%)	0.18% (-1.1-1.4%)	4.4 (-25-34)	40 (-270-330)
Bolivia	2000	Female	85 to 89	33% (21-47%)	67% (53-79%)	0.48 (0.26-0.83)	-0.12% (-1-0.75%)	-0.15% (-1.3-0.89%)	-2.5 (-21-15)	-17 (-150-110)
Bolivia	2000	Female	90 to 94	34% (22-49%)	66% (51-78%)	0.48 (0.24-0.85)	-0.4% (-1.4-0.46%)	-0.53% (-1.7-0.51%)	-3.7 (-12-3.6)	-17 (-58-20)
Bolivia	2000	Female	95 plus	37% (22-56%)	63% (44-78%)	0.5 (0.22-0.97)	-0.56% (-1.7-0.41%)	-0.78% (-2.2-0.39%)	-1.4 (-4.1-0.72)	-4.1 (-13-3.2)
Bolivia	2000	Male	All Ages	47% (42-52%)	53% (48-58%)	1.1 (0.87-1.3)	3% (2.3-3.8%)	4% (2.8-5.4%)	1300 (870-1700)	57000 (43000-71000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	2000	Male	15 to 19	37% (28-47%)	63% (53-72%)	0.54 (0.3-0.87)	3% (1.9-4.3%)	3.6% (1.9-5.9%)	20 (9.9-32)	2100 (1300-3100)
Bolivia	2000	Male	20 to 24	72% (62-79%)	28% (21-38%)	1.5 (0.96-2.2)	7.8% (5.3-10%)	9.3% (5.5-14%)	66 (39-99)	6100 (4000-8200)
Bolivia	2000	Male	25 to 29	80% (72-85%)	20% (15-28%)	2 (1.3-2.9)	9.3% (6.4-12%)	12% (7.1-16%)	80 (49-110)	6500 (4400-8600)
Bolivia	2000	Male	30 to 34	81% (74-86%)	19% (14-26%)	2.3 (1.5-3.3)	9.5% (6.6-13%)	12% (7.9-17%)	86 (53-120)	6000 (4100-8100)
Bolivia	2000	Male	35 to 39	78% (70-83%)	22% (17-30%)	2.3 (1.4-3.4)	9.6% (6.4-13%)	13% (7.9-18%)	95 (58-140)	5800 (3800-8000)
Bolivia	2000	Male	40 to 44	77% (70-82%)	23% (18-30%)	2.4 (1.5-3.6)	9.5% (6.4-13%)	13% (8.3-18%)	110 (69-150)	5600 (3800-7800)
Bolivia	2000	Male	45 to 49	78% (69-84%)	22% (16-31%)	2.4 (1.4-3.7)	9.3% (6-13%)	12% (7.8-18%)	120 (76-170)	5700 (3700-7900)
Bolivia	2000	Male	50 to 54	76% (68-81%)	24% (19-32%)	2.4 (1.4-3.5)	8.7% (5.5-13%)	11% (7.2-17%)	120 (76-180)	4800 (3100-7000)
Bolivia	2000	Male	55 to 59	70% (60-77%)	30% (23-40%)	2.3 (1.3-3.5)	7.7% (4.9-11%)	10% (6.1-15%)	130 (80-200)	4500 (2800-6600)
Bolivia	2000	Male	60 to 64	69% (58-77%)	31% (23-42%)	2.1 (1.2-3.4)	6.4% (3.5-10%)	7.9% (4.3-13%)	120 (66-200)	3600 (1900-5800)
Bolivia	2000	Male	65 to 69	73% (60-82%)	27% (18-40%)	1.9 (1.1-2.9)	4.8% (2.5-7.8%)	5.8% (2.9-9.5%)	120 (57-190)	2700 (1400-4400)
Bolivia	2000	Male	70 to 74	70% (55-82%)	30% (18-45%)	1.5 (0.87-2.3)	3.2% (1.4-5.4%)	3.9% (1.6-6.5%)	98 (40-170)	1900 (810-3100)
Bolivia	2000	Male	75 to 79	60% (42-77%)	40% (23-58%)	1 (0.54-1.7)	2.1% (0.65-3.9%)	2.4% (0.73-4.6%)	61 (18-120)	900 (270-1700)
Bolivia	2000	Male	80 to 84	52% (33-69%)	48% (31-67%)	0.8 (0.43-1.3)	1.4% (0.4-2.6%)	1.6% (0.4-3%)	34 (8.5-64)	370 (100-670)
Bolivia	2000	Male	85 to 89	47% (31-63%)	53% (37-69%)	0.7 (0.37-1.2)	0.9% (0.049-1.8%)	0.97% (-0.078-2%)	13 (-1-28)	110 (5.8-210)
Bolivia	2000	Male	90 to 94	48% (31-65%)	52% (35-69%)	0.69 (0.34-1.2)	0.51% (-0.3-1.4%)	0.46% (-0.53-1.5%)	2.5 (-2.9-8.3)	17 (-9.7-46)
Bolivia	2000	Male	95 plus	51% (30-70%)	49% (30-70%)	0.71 (0.3-1.3)	0.26% (-0.65-1.3%)	0.073% (-1.1-1.3%)	0.091 (-1.4-1.6)	1.3 (-3.3-6.7)
Bolivia	2005	Female	All Ages	40% (37-44%)	60% (56-63%)	0.6 (0.49-0.73)	0.96% (0.57-1.4%)	1.3% (0.49-2.2%)	370 (140-620)	15000 (9100-22000)
Bolivia	2005	Female	15 to 19	27% (22-35%)	73% (65-78%)	0.32 (0.19-0.51)	0.98% (0.53-1.5%)	1.4% (0.38-2.6%)	4.8 (1.3-9.3)	630 (320-990)
Bolivia	2005	Female	20 to 24	52% (46-58%)	48% (42-54%)	0.84 (0.52-1.3)	2.2% (1.2-3.5%)	3.1% (1-5.5%)	13 (4.4-24)	1500 (820-2300)
Bolivia	2005	Female	25 to 29	62% (56-67%)	38% (33-44%)	1.1 (0.74-1.6)	2.5% (1.5-3.8%)	3.9% (1.7-6.4%)	17 (7.6-30)	1600 (930-2400)
Bolivia	2005	Female	30 to 34	64% (58-69%)	36% (31-42%)	1.2 (0.8-1.7)	2.5% (1.4-3.8%)	4.1% (2-6.6%)	19 (9.1-31)	1400 (830-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	2005	Female	35 to 39	64% (58-69%)	36% (31-42%)	1.2 (0.75-1.7)	2.4% (1.3-3.8%)	4.2% (2.1-6.7%)	23 (12-38)	1400 (750-2200)
Bolivia	2005	Female	40 to 44	63% (58-68%)	37% (32-42%)	1.2 (0.75-1.7)	2.4% (1.2-4%)	4.3% (2.2-7%)	28 (14-46)	1400 (700-2300)
Bolivia	2005	Female	45 to 49	63% (57-68%)	37% (32-43%)	1.2 (0.76-1.8)	2.5% (1.1-4.4%)	4.3% (2.1-7.2%)	34 (16-57)	1400 (650-2500)
Bolivia	2005	Female	50 to 54	60% (50-69%)	40% (31-50%)	1.2 (0.69-1.8)	2.6% (1.1-4.5%)	4.3% (2-7.2%)	42 (20-71)	1500 (670-2700)
Bolivia	2005	Female	55 to 59	59% (41-75%)	41% (25-59%)	1.2 (0.66-2)	2.6% (0.91-4.6%)	4.1% (1.6-7.1%)	44 (18-77)	1400 (470-2500)
Bolivia	2005	Female	60 to 64	56% (38-74%)	44% (26-62%)	1.1 (0.61-1.9)	2.1% (0.46-4.2%)	3.1% (0.92-6%)	46 (14-90)	1200 (260-2500)
Bolivia	2005	Female	65 to 69	52% (35-69%)	48% (31-65%)	1 (0.55-1.6)	1.5% (0.033-3.1%)	2.2% (0.27-4.3%)	39 (4.7-77)	820 (17-1700)
Bolivia	2005	Female	70 to 74	46% (31-63%)	54% (37-69%)	0.8 (0.44-1.3)	0.95% (-0.37-2.4%)	1.4% (-0.27-3.2%)	34 (-6.7-79)	550 (-210-1400)
Bolivia	2005	Female	75 to 79	39% (26-56%)	61% (44-74%)	0.62 (0.31-1.1)	0.48% (-0.69-1.7%)	0.7% (-0.77-2.2%)	21 (-22-63)	250 (-340-880)
Bolivia	2005	Female	80 to 84	34% (20-51%)	66% (49-80%)	0.51 (0.26-0.87)	0.2% (-0.8-1.1%)	0.29% (-0.95-1.4%)	8.6 (-29-42)	78 (-310-430)
Bolivia	2005	Female	85 to 89	32% (20-46%)	68% (54-80%)	0.46 (0.24-0.77)	-0.058% (-0.96-0.7%)	-0.076% (-1.2-0.84%)	-1.6 (-26-19)	-11 (-190-140)
Bolivia	2005	Female	90 to 94	33% (21-48%)	67% (52-79%)	0.46 (0.23-0.8)	-0.33% (-1.2-0.48%)	-0.45% (-1.5-0.54%)	-4.5 (-16-5.5)	-20 (-74-29)
Bolivia	2005	Female	95 plus	36% (21-55%)	64% (45-79%)	0.48 (0.2-0.93)	-0.51% (-1.6-0.45%)	-0.72% (-2-0.46%)	-2.1 (-5.9-1.3)	-5.9 (-18-5.1)
Bolivia	2005	Male	All Ages	51% (47-54%)	49% (46-53%)	1.2 (0.96-1.4)	3.1% (2.4-3.9%)	3.9% (2.7-5.2%)	1200 (850-1600)	55000 (42000-70000)
Bolivia	2005	Male	15 to 19	36% (32-42%)	64% (58-68%)	0.51 (0.3-0.78)	2.8% (1.8-4%)	3.4% (1.8-5.5%)	17 (8.7-27)	1900 (1200-2800)
Bolivia	2005	Male	20 to 24	75% (71-79%)	25% (21-29%)	1.6 (1-2.3)	7.7% (5.1-10%)	9.4% (5.7-13%)	62 (37-91)	5900 (3900-8100)
Bolivia	2005	Male	25 to 29	83% (80-86%)	17% (14-20%)	2.2 (1.4-3.1)	9% (6.3-12%)	12% (6.8-16%)	77 (46-110)	6500 (4300-8800)
Bolivia	2005	Male	30 to 34	84% (81-86%)	16% (14-19%)	2.4 (1.6-3.4)	8.9% (6.3-12%)	12% (7.7-16%)	80 (49-110)	5800 (4000-7800)
Bolivia	2005	Male	35 to 39	81% (77-84%)	19% (16-23%)	2.4 (1.6-3.6)	8.9% (6.1-12%)	12% (7.7-17%)	87 (53-130)	5500 (3600-7500)
Bolivia	2005	Male	40 to 44	80% (76-83%)	20% (17-24%)	2.5 (1.6-3.7)	8.8% (6.1-13%)	12% (8-18%)	96 (61-140)	5200 (3600-7300)
Bolivia	2005	Male	45 to 49	81% (78-84%)	19% (16-22%)	2.6 (1.6-3.9)	8.8% (5.9-12%)	12% (7.8-17%)	110 (70-160)	5200 (3500-7400)
Bolivia	2005	Male	50 to 54	79% (75-82%)	21% (18-25%)	2.5 (1.5-3.8)	8.1% (5.2-12%)	11% (6.8-16%)	120 (76-190)	5000 (3200-7300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	2005	Male	55 to 59	72% (67-76%)	28% (24-33%)	2.4 (1.4-3.5)	7.3% (4.5-11%)	9.6% (5.7-14%)	120 (72-180)	4200 (2500-6300)
Bolivia	2005	Male	60 to 64	72% (66-76%)	28% (24-34%)	2.2 (1.3-3.4)	6% (3.3-9.4%)	7.6% (4.1-12%)	130 (67-210)	3800 (2000-6100)
Bolivia	2005	Male	65 to 69	74% (62-83%)	26% (17-38%)	1.9 (1.2-2.9)	4.4% (2.3-7%)	5.4% (2.8-8.7%)	110 (54-170)	2600 (1300-4100)
Bolivia	2005	Male	70 to 74	69% (54-82%)	31% (18-46%)	1.4 (0.82-2.2)	2.9% (1.2-4.9%)	3.5% (1.4-6%)	88 (35-150)	1700 (700-2900)
Bolivia	2005	Male	75 to 79	59% (41-77%)	41% (23-59%)	1 (0.53-1.7)	1.9% (0.66-3.5%)	2.3% (0.74-4.2%)	64 (20-120)	940 (320-1700)
Bolivia	2005	Male	80 to 84	51% (32-68%)	49% (32-68%)	0.77 (0.41-1.3)	1.3% (0.41-2.4%)	1.5% (0.39-2.9%)	40 (10-75)	440 (130-800)
Bolivia	2005	Male	85 to 89	47% (31-63%)	53% (37-69%)	0.68 (0.37-1.2)	0.86% (0.052-1.7%)	0.94% (-0.066-1.9%)	16 (-1.1-33)	130 (8.1-260)
Bolivia	2005	Male	90 to 94	47% (31-64%)	53% (36-69%)	0.67 (0.34-1.2)	0.5% (-0.27-1.3%)	0.45% (-0.51-1.4%)	3.3 (-3.7-10)	23 (-12-58)
Bolivia	2005	Male	95 plus	49% (28-69%)	51% (31-72%)	0.69 (0.29-1.3)	0.28% (-0.58-1.3%)	0.095% (-1-1.3%)	0.18 (-1.9-2.4)	2.1 (-4.5-9.7)
Bolivia	2010	Female	All Ages	41% (38-45%)	59% (55-62%)	0.65 (0.53-0.79)	1.1% (0.65-1.7%)	1.4% (0.52-2.4%)	430 (160-770)	17000 (9900-26000)
Bolivia	2010	Female	15 to 19	27% (22-37%)	73% (63-78%)	0.34 (0.19-0.54)	0.99% (0.55-1.5%)	1.4% (0.46-2.7%)	4.8 (1.6-9.4)	640 (360-1000)
Bolivia	2010	Female	20 to 24	52% (47-59%)	48% (41-53%)	0.87 (0.54-1.3)	2.2% (1.3-3.4%)	3.2% (1.2-5.5%)	13 (4.3-23)	1500 (830-2300)
Bolivia	2010	Female	25 to 29	62% (56-67%)	38% (33-44%)	1.2 (0.77-1.7)	2.5% (1.5-3.7%)	3.9% (1.9-6.5%)	17 (7.5-29)	1600 (940-2500)
Bolivia	2010	Female	30 to 34	63% (58-69%)	37% (31-42%)	1.2 (0.82-1.8)	2.4% (1.5-3.6%)	4% (2.1-6.5%)	19 (9.9-32)	1500 (890-2300)
Bolivia	2010	Female	35 to 39	63% (58-69%)	37% (31-42%)	1.2 (0.76-1.8)	2.4% (1.3-3.7%)	4.2% (2.1-6.7%)	24 (12-40)	1500 (820-2400)
Bolivia	2010	Female	40 to 44	63% (57-68%)	37% (32-43%)	1.2 (0.77-1.8)	2.5% (1.2-4%)	4.5% (2.2-7.2%)	30 (15-50)	1500 (760-2500)
Bolivia	2010	Female	45 to 49	62% (56-68%)	38% (32-44%)	1.3 (0.79-1.9)	2.6% (1.2-4.4%)	4.5% (2.3-7.5%)	38 (19-66)	1600 (760-2800)
Bolivia	2010	Female	50 to 54	60% (50-69%)	40% (31-50%)	1.2 (0.73-1.9)	2.7% (1.1-4.6%)	4.5% (2.1-7.5%)	47 (22-83)	1700 (740-3100)
Bolivia	2010	Female	55 to 59	59% (41-75%)	41% (25-59%)	1.3 (0.7-2)	2.7% (1-4.8%)	4.3% (1.9-7.3%)	58 (26-100)	1800 (710-3300)
Bolivia	2010	Female	60 to 64	56% (38-74%)	44% (26-62%)	1.2 (0.65-2.1)	2.2% (0.47-4.4%)	3.3% (0.99-6.4%)	52 (15-100)	1400 (290-2800)
Bolivia	2010	Female	65 to 69	52% (35-69%)	48% (31-65%)	1 (0.59-1.6)	1.6% (0.076-3.2%)	2.4% (0.35-4.5%)	51 (7.4-100)	1100 (53-2300)
Bolivia	2010	Female	70 to 74	46% (31-63%)	54% (37-69%)	0.84 (0.45-1.4)	1.1% (-0.32-2.6%)	1.6% (-0.23-3.6%)	42 (-6-96)	690 (-200-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	2010	Female	75 to 79	39% (26-56%)	61% (44-74%)	0.65 (0.32-1.1)	0.61% (-0.6-1.8%)	0.86% (-0.64-2.4%)	28 (-20-78)	360 (-360-1100)
Bolivia	2010	Female	80 to 84	34% (21-52%)	66% (48-79%)	0.53 (0.28-0.9)	0.3% (-0.67-1.2%)	0.4% (-0.77-1.6%)	16 (-30-60)	150 (-330-610)
Bolivia	2010	Female	85 to 89	32% (20-46%)	68% (54-80%)	0.48 (0.25-0.79)	0.0051% (-0.87-0.79%)	-0.0038% (-1-0.93%)	-0.14 (-34-29)	1.5 (-240-220)
Bolivia	2010	Female	90 to 94	33% (21-48%)	67% (52-79%)	0.48 (0.24-0.83)	-0.29% (-1.2-0.52%)	-0.4% (-1.5-0.61%)	-5.8 (-23-9)	-26 (-110-48)
Bolivia	2010	Female	95 plus	36% (21-54%)	64% (46-79%)	0.49 (0.21-0.91)	-0.47% (-1.6-0.53%)	-0.67% (-2-0.53%)	-3 (-9.1-2.3)	-8.4 (-28-9.1)
Bolivia	2010	Male	All Ages	52% (48-55%)	48% (45-52%)	1.3 (1.1-1.6)	3.5% (2.7-4.5%)	4.2% (2.9-5.7%)	1400 (870-1900)	60000 (43000-78000)
Bolivia	2010	Male	15 to 19	36% (31-40%)	64% (60-69%)	0.54 (0.32-0.81)	2.8% (1.8-3.9%)	3.5% (1.8-5.7%)	17 (8.5-28)	2000 (1300-2900)
Bolivia	2010	Male	20 to 24	75% (72-78%)	25% (22-28%)	1.7 (1.1-2.4)	7.7% (5.3-10%)	9.6% (5.7-14%)	62 (34-94)	6000 (4000-8400)
Bolivia	2010	Male	25 to 29	83% (80-85%)	17% (15-20%)	2.3 (1.5-3.3)	8.9% (6.1-12%)	12% (6.8-17%)	79 (43-120)	6800 (4300-9500)
Bolivia	2010	Male	30 to 34	84% (81-86%)	16% (14-19%)	2.6 (1.7-3.6)	8.8% (6.1-12%)	12% (7.3-17%)	85 (49-130)	6300 (4200-8800)
Bolivia	2010	Male	35 to 39	81% (77-83%)	19% (17-23%)	2.6 (1.7-3.8)	8.9% (6-12%)	13% (7.9-18%)	94 (56-140)	6000 (3800-8400)
Bolivia	2010	Male	40 to 44	79% (76-82%)	21% (18-24%)	2.6 (1.7-3.8)	8.9% (6.2-12%)	13% (8.3-18%)	100 (62-150)	5700 (3600-8000)
Bolivia	2010	Male	45 to 49	81% (78-84%)	19% (16-22%)	2.8 (1.8-4.2)	8.8% (6-13%)	12% (8.3-18%)	120 (74-190)	5600 (3700-8500)
Bolivia	2010	Male	50 to 54	78% (75-81%)	22% (19-25%)	2.7 (1.7-4)	8.2% (5.4-12%)	11% (7.1-16%)	130 (78-210)	5400 (3300-8300)
Bolivia	2010	Male	55 to 59	72% (67-76%)	28% (24-33%)	2.5 (1.6-3.7)	7.4% (4.7-11%)	9.9% (6.1-14%)	150 (82-220)	5100 (3000-7800)
Bolivia	2010	Male	60 to 64	71% (66-76%)	29% (24-34%)	2.4 (1.4-3.6)	6.2% (3.5-9.8%)	7.9% (4.3-13%)	140 (71-230)	4100 (2100-6600)
Bolivia	2010	Male	65 to 69	74% (62-83%)	26% (17-38%)	2.1 (1.3-3.1)	4.6% (2.4-7.3%)	5.7% (3-9.1%)	130 (66-210)	3100 (1600-5200)
Bolivia	2010	Male	70 to 74	69% (53-82%)	31% (18-47%)	1.5 (0.88-2.4)	3.1% (1.3-5.1%)	3.7% (1.5-6.3%)	98 (39-170)	1900 (800-3200)
Bolivia	2010	Male	75 to 79	59% (40-77%)	41% (23-60%)	1.1 (0.56-1.8)	2% (0.77-3.7%)	2.4% (0.87-4.4%)	72 (26-130)	1100 (400-2000)
Bolivia	2010	Male	80 to 84	51% (33-68%)	49% (32-67%)	0.83 (0.45-1.4)	1.4% (0.46-2.4%)	1.6% (0.43-2.9%)	52 (14-93)	580 (180-1000)
Bolivia	2010	Male	85 to 89	47% (30-63%)	53% (37-70%)	0.72 (0.39-1.2)	0.94% (0.12-1.8%)	1% (0.027-2.1%)	24 (0.57-48)	200 (25-370)
Bolivia	2010	Male	90 to 94	47% (30-64%)	53% (36-70%)	0.7 (0.36-1.2)	0.56% (-0.26-1.4%)	0.52% (-0.5-1.5%)	5.4 (-5-16)	36 (-17-92)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	2010	Male	95 plus	49% (28-69%)	51% (31-72%)	0.71 (0.29-1.3)	0.33% (-0.58-1.4%)	0.15% (-0.96-1.4%)	0.42 (-2.6-3.9)	3.8 (-6.2-15)
Bolivia	2016	Female	All Ages	44% (40-48%)	56% (52-60%)	0.8 (0.58-1.1)	1.4% (0.79-2.3%)	1.7% (0.62-3%)	590 (200-1100)	22000 (12000-36000)
Bolivia	2016	Female	15 to 19	35% (25-44%)	65% (56-75%)	0.5 (0.26-0.81)	1.2% (0.66-1.9%)	2% (0.71-3.8%)	6.2 (2.1-12)	780 (400-1300)
Bolivia	2016	Female	20 to 24	55% (45-63%)	45% (37-55%)	1.1 (0.61-1.7)	2.5% (1.5-3.7%)	3.9% (1.5-6.8%)	14 (5.2-26)	1600 (930-2500)
Bolivia	2016	Female	25 to 29	62% (52-69%)	38% (31-48%)	1.4 (0.81-2.1)	2.7% (1.7-4.1%)	4.6% (2-7.7%)	18 (7.6-32)	1800 (1000-2800)
Bolivia	2016	Female	30 to 34	65% (54-71%)	35% (29-46%)	1.5 (0.87-2.3)	2.6% (1.5-4.1%)	4.8% (2.4-7.9%)	22 (10-40)	1800 (960-2900)
Bolivia	2016	Female	35 to 39	65% (55-71%)	35% (29-45%)	1.4 (0.83-2.3)	2.6% (1.4-4.3%)	5% (2.6-8.1%)	29 (13-53)	1900 (910-3100)
Bolivia	2016	Female	40 to 44	64% (54-71%)	36% (29-46%)	1.5 (0.83-2.3)	2.8% (1.3-4.6%)	5.3% (2.8-8.7%)	38 (17-67)	2000 (860-3400)
Bolivia	2016	Female	45 to 49	63% (52-70%)	37% (30-48%)	1.5 (0.82-2.4)	2.9% (1.3-5.2%)	5.3% (2.7-9%)	48 (22-88)	2100 (910-3800)
Bolivia	2016	Female	50 to 54	61% (50-70%)	39% (30-50%)	1.4 (0.79-2.3)	3.1% (1.3-5.5%)	5.2% (2.4-8.7%)	61 (27-110)	2200 (900-4100)
Bolivia	2016	Female	55 to 59	58% (41-74%)	42% (26-59%)	1.4 (0.75-2.3)	3.2% (1.3-5.6%)	5% (2.3-8.6%)	74 (32-140)	2300 (920-4500)
Bolivia	2016	Female	60 to 64	55% (38-74%)	45% (26-62%)	1.3 (0.71-2.4)	2.6% (0.71-5.3%)	3.9% (1.3-7.5%)	78 (25-160)	2100 (540-4500)
Bolivia	2016	Female	65 to 69	51% (35-69%)	49% (31-65%)	1.2 (0.64-2)	2.9% (0.31-4.1%)	2.9% (0.72-5.6%)	68 (15-140)	1500 (220-3200)
Bolivia	2016	Female	70 to 74	46% (31-63%)	54% (37-69%)	0.94 (0.48-1.6)	1.4% (-0.099-3.3%)	2% (0.074-4.4%)	61 (2.2-140)	1000 (-76-2600)
Bolivia	2016	Female	75 to 79	39% (25-57%)	61% (43-75%)	0.73 (0.34-1.3)	0.85% (-0.41-2.4%)	1.2% (-0.41-3.1%)	43 (-13-120)	560 (-240-1600)
Bolivia	2016	Female	80 to 84	34% (21-52%)	66% (48-79%)	0.6 (0.3-1)	0.48% (-0.6-1.6%)	0.63% (-0.71-1.9%)	31 (-34-100)	300 (-380-1000)
Bolivia	2016	Female	85 to 89	32% (20-46%)	68% (54-80%)	0.55 (0.28-0.97)	0.15% (-0.74-1.2%)	0.17% (-0.92-1.4%)	7 (-40-60)	56 (-290-440)
Bolivia	2016	Female	90 to 94	33% (21-48%)	67% (52-79%)	0.55 (0.27-0.99)	-0.17% (-1.2-0.94%)	-0.26% (-1.5-1.1%)	-5.8 (-34-25)	-23 (-160-130)
Bolivia	2016	Female	95 plus	36% (21-55%)	64% (45-79%)	0.58 (0.24-1.1)	-0.36% (-1.6-0.96%)	-0.55% (-2-0.99%)	-3.9 (-15-7.1)	-10 (-46-27)
Bolivia	2016	Male	All Ages	50% (45-54%)	50% (46-55%)	1.4 (1-1.9)	4.2% (3-5.6%)	4.7% (3.1-6.7%)	1600 (960-2500)	70000 (47000-100000)
Bolivia	2016	Male	15 to 19	36% (26-46%)	64% (54-74%)	0.62 (0.32-1)	3% (2-4.4%)	3.9% (2-6.6%)	19 (9.2-34)	2300 (1400-3400)
Bolivia	2016	Male	20 to 24	71% (61-78%)	29% (22-39%)	1.7 (1-2.7)	7.9% (5.4-11%)	9.8% (5.5-15%)	65 (34-100)	6500 (4100-9300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bolivia	2016	Male	25 to 29	79% (71-85%)	21% (15-29%)	2.3 (1.4-3.6)	9.1% (5.8-12%)	12% (6.3-18%)	83 (39-140)	7200 (4300-11000)
Bolivia	2016	Male	30 to 34	81% (73-86%)	19% (14-27%)	2.7 (1.7-4.1)	9.1% (5.8-13%)	12% (6.8-18%)	94 (44-150)	7100 (4200-11000)
Bolivia	2016	Male	35 to 39	77% (69-83%)	23% (17-31%)	2.7 (1.6-4.2)	9.2% (5.9-13%)	13% (7.5-20%)	110 (57-180)	7100 (4200-11000)
Bolivia	2016	Male	40 to 44	76% (68-82%)	24% (18-32%)	2.8 (1.6-4.5)	9.2% (6.1-13%)	13% (8.4-20%)	120 (69-200)	6800 (4100-11000)
Bolivia	2016	Male	45 to 49	77% (68-83%)	23% (17-32%)	2.8 (1.6-4.6)	9.2% (6.2-13%)	13% (8.5-19%)	140 (79-220)	6600 (4100-10000)
Bolivia	2016	Male	50 to 54	75% (67-81%)	25% (19-33%)	2.8 (1.6-4.3)	8.6% (5.4-13%)	12% (7.4-18%)	150 (84-250)	6200 (3600-9900)
Bolivia	2016	Male	55 to 59	69% (60-77%)	31% (23-40%)	2.6 (1.5-4.1)	7.9% (4.7-12%)	11% (6.2-16%)	170 (89-270)	5900 (3400-9200)
Bolivia	2016	Male	60 to 64	68% (58-77%)	32% (23-42%)	2.5 (1.4-4.3)	6.6% (3.7-11%)	8.5% (4.5-14%)	180 (88-310)	5200 (2700-8900)
Bolivia	2016	Male	65 to 69	72% (59-82%)	28% (18-41%)	2.2 (1.3-3.5)	5.1% (2.6-8.5%)	6.4% (3.3-11%)	150 (72-280)	3700 (1800-6600)
Bolivia	2016	Male	70 to 74	69% (54-82%)	31% (18-46%)	1.7 (0.91-2.8)	3.4% (1.5-6.1%)	4.2% (1.8-7.6%)	130 (55-240)	2500 (1100-4600)
Bolivia	2016	Male	75 to 79	59% (40-77%)	41% (23-60%)	1.2 (0.58-2.1)	2.3% (0.9-4.3%)	2.8% (1-5.4%)	91 (34-180)	1400 (530-2700)
Bolivia	2016	Male	80 to 84	51% (32-68%)	49% (32-68%)	0.91 (0.47-1.6)	1.6% (0.5-2.9%)	1.9% (0.52-3.5%)	71 (19-130)	790 (240-1500)
Bolivia	2016	Male	85 to 89	46% (30-63%)	54% (37-70%)	0.8 (0.41-1.4)	1.1% (0.24-2.2%)	1.2% (0.17-2.5%)	38 (5.2-80)	310 (64-630)
Bolivia	2016	Male	90 to 94	46% (30-64%)	54% (36-70%)	0.79 (0.38-1.4)	0.72% (-0.16-1.7%)	0.69% (-0.38-1.9%)	11 (-5.8-30)	71 (-16-170)
Bolivia	2016	Male	95 plus	49% (28-69%)	51% (31-72%)	0.8 (0.32-1.6)	0.5% (-0.47-1.8%)	0.32% (-0.9-1.9%)	1.4 (-4-8.2)	8.8 (-8.5-33)
Ecuador	1990	Female	All Ages	10% (7.9-13%)	90% (87-92%)	0.61 (0.47-0.77)	2% (1.6-2.5%)	3.2% (2.4-4.1%)	790 (580-1000)	32000 (25000-39000)
Ecuador	1990	Female	15 to 19	9.4% (5.4-15%)	91% (85-95%)	0.4 (0.19-0.72)	1.8% (0.99-3%)	2.9% (1.3-5.3%)	15 (6.8-27)	1400 (790-2400)
Ecuador	1990	Female	20 to 24	14% (8.2-20%)	86% (80-92%)	0.89 (0.47-1.5)	4% (2.5-6%)	6.3% (3.4-10%)	35 (19-55)	3200 (2000-4700)
Ecuador	1990	Female	25 to 29	16% (9.6-23%)	84% (77-90%)	1.2 (0.67-2)	5% (3.3-6.9%)	8.3% (5-12%)	42 (25-63)	3400 (2200-4800)
Ecuador	1990	Female	30 to 34	17% (11-26%)	83% (74-89%)	1.3 (0.73-2.2)	5.1% (3.4-7.1%)	8.7% (5.6-12%)	46 (29-66)	3300 (2300-4600)
Ecuador	1990	Female	35 to 39	19% (11-28%)	81% (72-89%)	1.4 (0.76-2.2)	5.2% (3.6-7.3%)	8.8% (6-12%)	53 (35-74)	3300 (2200-4500)
Ecuador	1990	Female	40 to 44	19% (12-27%)	81% (73-88%)	1.3 (0.7-2.1)	5.4% (3.6-7.5%)	9% (5.8-13%)	55 (35-77)	3000 (1900-4100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	1990	Female	45 to 49	17% (10-25%)	83% (75-90%)	1.1 (0.54-1.9)	4.9% (3.2-6.9%)	8.1% (5.1-11%)	54 (35-78)	2600 (1700-3700)
Ecuador	1990	Female	50 to 54	15% (9.7-23%)	85% (77-90%)	0.99 (0.48-1.7)	4.7% (3-6.8%)	7.3% (4.6-11%)	58 (36-85)	2400 (1500-3400)
Ecuador	1990	Female	55 to 59	15% (9.6-23%)	85% (77-90%)	1.1 (0.54-1.9)	4.9% (3.1-7.1%)	7.3% (4.8-11%)	67 (44-99)	2300 (1500-3400)
Ecuador	1990	Female	60 to 64	14% (8.3-21%)	86% (79-92%)	1 (0.49-1.8)	4.6% (3-6.7%)	6.5% (4.2-9.4%)	74 (47-110)	2200 (1400-3200)
Ecuador	1990	Female	65 to 69	11% (7-17%)	89% (83-93%)	0.82 (0.39-1.4)	3.9% (2.4-5.8%)	5.5% (3.4-8.1%)	70 (43-100)	1700 (1000-2500)
Ecuador	1990	Female	70 to 74	9.5% (6.1-14%)	91% (86-94%)	0.64 (0.31-1.2)	3.2% (1.8-5%)	4.3% (2.5-6.7%)	66 (39-100)	1300 (740-2000)
Ecuador	1990	Female	75 to 79	8.3% (4.6-13%)	92% (87-95%)	0.5 (0.22-0.95)	2.4% (1.3-3.9%)	3.1% (1.7-5.2%)	56 (30-93)	830 (440-1400)
Ecuador	1990	Female	80 to 84	7.2% (3.8-12%)	93% (88-96%)	0.4 (0.18-0.73)	1.9% (1-3.2%)	2.3% (1.2-3.9%)	45 (24-77)	490 (270-830)
Ecuador	1990	Female	85 to 89	6.7% (3.7-11%)	93% (89-96%)	0.36 (0.16-0.66)	1.5% (0.76-2.7%)	1.8% (0.88-3.3%)	32 (15-58)	240 (120-430)
Ecuador	1990	Female	90 to 94	7.3% (4.1-12%)	93% (88-96%)	0.37 (0.16-0.75)	1.3% (0.54-2.6%)	1.5% (0.59-3.1%)	16 (6.2-33)	83 (34-170)
Ecuador	1990	Female	95 plus	8.5% (4.2-15%)	91% (85-96%)	0.41 (0.15-0.86)	1.3% (0.42-2.9%)	1.4% (0.4-3.3%)	5.4 (1.5-13)	19 (6.5-44)
Ecuador	1990	Male	All Ages	21% (17-25%)	79% (75-83%)	1.9 (1.6-2.4)	6.6% (5.6-7.8%)	9.2% (7.8-11%)	2900 (2400-3400)	120000 (100000-150000)
Ecuador	1990	Male	15 to 19	17% (10-26%)	83% (74-90%)	0.88 (0.43-1.5)	4.9% (3-7.5%)	6.3% (3.4-10%)	51 (27-83)	4700 (2800-7200)
Ecuador	1990	Male	20 to 24	35% (22-48%)	65% (52-78%)	2.5 (1.4-4.1)	13% (9-17%)	16% (10-22%)	170 (110-240)	14000 (9600-19000)
Ecuador	1990	Male	25 to 29	42% (28-56%)	58% (44-72%)	3.5 (2-5.3)	16% (12-21%)	20% (14-27%)	210 (140-290)	16000 (11000-21000)
Ecuador	1990	Male	30 to 34	43% (28-59%)	57% (41-72%)	4.3 (2.6-6.5)	17% (13-21%)	22% (16-28%)	220 (160-290)	15000 (11000-19000)
Ecuador	1990	Male	35 to 39	42% (28-56%)	58% (44-72%)	4.5 (2.6-7.1)	17% (14-22%)	23% (18-29%)	240 (180-300)	14000 (11000-17000)
Ecuador	1990	Male	40 to 44	41% (28-55%)	59% (45-72%)	4.4 (2.4-7.1)	17% (14-21%)	23% (18-28%)	220 (170-280)	12000 (9100-14000)
Ecuador	1990	Male	45 to 49	42% (27-56%)	58% (44-73%)	4.6 (2.4-7.5)	17% (14-21%)	23% (18-28%)	250 (190-300)	11000 (8900-14000)
Ecuador	1990	Male	50 to 54	41% (28-54%)	59% (46-72%)	4.4 (2.3-7.2)	16% (12-19%)	21% (17-25%)	250 (190-300)	9700 (7800-12000)
Ecuador	1990	Male	55 to 59	36% (24-51%)	64% (49-76%)	4.1 (2.2-6.9)	14% (11-17%)	18% (15-23%)	250 (190-300)	8400 (6600-10000)
Ecuador	1990	Male	60 to 64	34% (22-49%)	66% (51-78%)	3.9 (2-6.8)	12% (9.4-15%)	15% (12-20%)	240 (190-310)	7000 (5400-8900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	1990	Male	65 to 69	34% (22-46%)	66% (54-78%)	3.3 (1.7-5.4)	11% (8-14%)	14% (10-17%)	230 (170-290)	5500 (4100-7000)
Ecuador	1990	Male	70 to 74	30% (19-42%)	70% (58-81%)	2.3 (1.2-4)	8.2% (5.7-11%)	10% (7.1-14%)	200 (140-270)	3800 (2600-5000)
Ecuador	1990	Male	75 to 79	24% (14-36%)	76% (64-86%)	1.5 (0.71-2.7)	5.6% (3.3-8.2%)	6.8% (4-9.9%)	150 (86-220)	2200 (1300-3100)
Ecuador	1990	Male	80 to 84	20% (11-30%)	80% (70-89%)	1.1 (0.5-1.9)	4.1% (2.4-6.1%)	4.8% (2.8-7.3%)	96 (56-140)	1000 (610-1500)
Ecuador	1990	Male	85 to 89	18% (9.9-27%)	82% (73-90%)	0.91 (0.42-1.6)	3.2% (1.7-5%)	3.7% (2-5.9%)	56 (29-88)	430 (230-670)
Ecuador	1990	Male	90 to 94	18% (11-28%)	82% (72-89%)	0.89 (0.4-1.7)	2.7% (1.3-4.7%)	3.1% (1.4-5.4%)	24 (11-42)	130 (63-230)
Ecuador	1990	Male	95 plus	20% (11-34%)	80% (66-89%)	0.94 (0.36-1.9)	2.6% (0.98-5%)	2.8% (0.92-5.6%)	6.7 (2.2-13)	25 (9.5-48)
Ecuador	1995	Female	All Ages	10% (8-13%)	90% (87-92%)	0.7 (0.53-0.9)	2.3% (1.9-2.9%)	3.7% (2.7-4.7%)	940 (700-1200)	36000 (29000-45000)
Ecuador	1995	Female	15 to 19	9.2% (5.3-14%)	91% (86-95%)	0.44 (0.21-0.79)	2% (1.1-3.3%)	3.1% (1.4-5.5%)	17 (8-31)	1700 (920-2800)
Ecuador	1995	Female	20 to 24	13% (8.1-20%)	87% (80-92%)	0.99 (0.53-1.7)	4.1% (2.6-6%)	6.6% (3.6-10%)	37 (21-57)	3400 (2200-4900)
Ecuador	1995	Female	25 to 29	15% (9.4-23%)	85% (77-91%)	1.4 (0.74-2.2)	5% (3.3-6.9%)	8.5% (5.1-12%)	46 (28-67)	3800 (2500-5200)
Ecuador	1995	Female	30 to 34	17% (10-26%)	83% (74-90%)	1.5 (0.81-2.4)	5% (3.5-7%)	8.9% (5.9-12%)	50 (33-70)	3600 (2500-4900)
Ecuador	1995	Female	35 to 39	19% (11-28%)	81% (72-89%)	1.5 (0.81-2.4)	5.1% (3.6-7%)	9.1% (6.2-12%)	56 (38-76)	3500 (2500-4700)
Ecuador	1995	Female	40 to 44	18% (11-27%)	82% (73-89%)	1.4 (0.76-2.3)	5.2% (3.6-7.1%)	9% (5.9-12%)	63 (42-87)	3500 (2400-4800)
Ecuador	1995	Female	45 to 49	16% (9.7-24%)	84% (76-90%)	1.2 (0.59-2.1)	5.1% (3.4-7%)	8.5% (5.6-12%)	64 (41-89)	3000 (2000-4100)
Ecuador	1995	Female	50 to 54	15% (9.6-22%)	85% (78-90%)	1.1 (0.56-1.9)	4.8% (3.1-7%)	7.8% (5-11%)	68 (43-97)	2800 (1800-4000)
Ecuador	1995	Female	55 to 59	15% (9.3-22%)	85% (78-91%)	1.2 (0.58-2.2)	5% (3.3-7.1%)	7.6% (5.1-11%)	77 (51-110)	2700 (1800-3800)
Ecuador	1995	Female	60 to 64	14% (7.9-21%)	86% (79-92%)	1.1 (0.53-2.1)	4.7% (3-6.8%)	6.8% (4.4-9.7%)	86 (56-120)	2500 (1600-3600)
Ecuador	1995	Female	65 to 69	11% (6.9-17%)	89% (83-93%)	0.92 (0.43-1.6)	4.2% (2.6-6.1%)	6% (3.8-8.5%)	87 (55-120)	2100 (1300-3000)
Ecuador	1995	Female	70 to 74	9.4% (6.1-14%)	91% (86-94%)	0.72 (0.35-1.3)	3.5% (2.1-5.4%)	4.8% (3-7.3%)	83 (51-130)	1600 (950-2500)
Ecuador	1995	Female	75 to 79	8.2% (4.8-13%)	92% (87-95%)	0.56 (0.24-1.1)	2.7% (1.5-4.2%)	3.5% (2-5.6%)	68 (38-110)	1000 (570-1600)
Ecuador	1995	Female	80 to 84	7.1% (3.8-12%)	93% (88-96%)	0.45 (0.2-0.83)	2.2% (1.2-3.5%)	2.7% (1.5-4.5%)	61 (33-100)	660 (370-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	1995	Female	85 to 89	6.6% (3.5-11%)	93% (89-97%)	0.4 (0.18-0.73)	1.8% (0.91-3%)	2.2% (1.1-3.7%)	46 (23-80)	350 (180-600)
Ecuador	1995	Female	90 to 94	7% (4-11%)	93% (89-96%)	0.41 (0.18-0.82)	1.5% (0.69-2.8%)	1.8% (0.79-3.5%)	24 (10-45)	120 (54-230)
Ecuador	1995	Female	95 plus	8.3% (4.2-14%)	92% (86-96%)	0.45 (0.16-0.93)	1.5% (0.57-3.2%)	1.7% (0.59-3.7%)	8.8 (3-19)	31 (12-65)
Ecuador	1995	Male	All Ages	21% (17-25%)	79% (75-83%)	2.3 (1.8-2.8)	7.3% (6.1-8.6%)	9.8% (8.3-11%)	3200 (2700-3800)	140000 (120000-170000)
Ecuador	1995	Male	15 to 19	17% (10-25%)	83% (75-90%)	1 (0.49-1.8)	5.3% (3.2-8%)	6.8% (3.6-11%)	63 (33-100)	5600 (3300-8600)
Ecuador	1995	Male	20 to 24	35% (21-48%)	65% (52-79%)	2.9 (1.6-4.8)	13% (9.3-18%)	16% (11-22%)	210 (130-290)	17000 (11000-22000)
Ecuador	1995	Male	25 to 29	41% (28-55%)	59% (45-72%)	4 (2.3-6.1)	16% (12-21%)	19% (13-26%)	250 (170-340)	18000 (13000-24000)
Ecuador	1995	Male	30 to 34	43% (28-58%)	57% (42-72%)	4.9 (2.9-7.3)	17% (13-21%)	22% (15-28%)	260 (180-340)	17000 (13000-22000)
Ecuador	1995	Male	35 to 39	41% (28-55%)	59% (45-72%)	5.1 (3-8.1)	17% (13-21%)	22% (17-28%)	260 (200-340)	16000 (12000-19000)
Ecuador	1995	Male	40 to 44	41% (28-55%)	59% (45-72%)	5.1 (2.8-8)	16% (13-20%)	22% (17-27%)	260 (200-320)	13000 (11000-17000)
Ecuador	1995	Male	45 to 49	42% (27-56%)	58% (44-73%)	5.2 (2.8-8.5)	16% (13-20%)	22% (17-27%)	260 (200-310)	12000 (9300-14000)
Ecuador	1995	Male	50 to 54	40% (28-53%)	60% (47-72%)	5 (2.7-8.3)	15% (12-18%)	20% (16-25%)	260 (210-320)	10000 (8400-13000)
Ecuador	1995	Male	55 to 59	36% (23-50%)	64% (50-77%)	4.7 (2.4-7.9)	13% (10-16%)	17% (14-22%)	250 (200-310)	8600 (6700-11000)
Ecuador	1995	Male	60 to 64	34% (21-48%)	66% (52-79%)	4.3 (2.3-7.7)	12% (9-15%)	15% (12-19%)	260 (200-330)	7400 (5800-9400)
Ecuador	1995	Male	65 to 69	33% (21-46%)	67% (54-79%)	3.7 (1.9-6.3)	11% (7.9-13%)	13% (10-17%)	260 (200-330)	6200 (4700-7900)
Ecuador	1995	Male	70 to 74	29% (18-41%)	71% (59-82%)	2.6 (1.3-4.5)	8.3% (6-11%)	10% (7.5-14%)	220 (160-290)	4300 (3100-5700)
Ecuador	1995	Male	75 to 79	24% (14-35%)	76% (65-86%)	1.7 (0.79-3.1)	5.9% (3.7-8.3%)	7.2% (4.6-10%)	160 (100-230)	2400 (1500-3400)
Ecuador	1995	Male	80 to 84	19% (11-29%)	81% (71-89%)	1.2 (0.57-2.1)	4.3% (2.7-6.3%)	5.2% (3.3-7.5%)	120 (74-170)	1300 (820-1900)
Ecuador	1995	Male	85 to 89	17% (9.8-27%)	83% (73-90%)	1 (0.47-1.8)	3.5% (2.1-5.3%)	4.2% (2.3-6.3%)	77 (43-120)	590 (340-890)
Ecuador	1995	Male	90 to 94	18% (10-27%)	82% (73-90%)	1 (0.45-1.9)	3.1% (1.6-4.9%)	3.6% (1.7-5.7%)	34 (16-54)	180 (92-290)
Ecuador	1995	Male	95 plus	20% (10-33%)	80% (67-90%)	1.1 (0.4-2.2)	3% (1.3-5.4%)	3.3% (1.3-6.2%)	9.8 (3.6-18)	36 (15-64)
Ecuador	2000	Female	All Ages	9.6% (7.7-12%)	90% (88-92%)	0.82 (0.62-1)	2.6% (2.2-3.2%)	4.2% (3.3-5.2%)	1100 (860-1400)	40000 (33000-49000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	2000	Female	15 to 19	9% (5.3-14%)	91% (86-95%)	0.54 (0.27-0.93)	2.2% (1.2-3.4%)	3.6% (1.8-6.2%)	19 (9.5-33)	1900 (1100-2900)
Ecuador	2000	Female	20 to 24	12% (7.3-17%)	88% (83-93%)	1.1 (0.59-1.8)	4% (2.7-5.6%)	6.3% (3.8-9.5%)	35 (21-53)	3300 (2200-4700)
Ecuador	2000	Female	25 to 29	13% (8.2-19%)	87% (81-92%)	1.4 (0.78-2.3)	4.5% (3.1-6.2%)	7.7% (5-11%)	42 (27-59)	3600 (2500-4800)
Ecuador	2000	Female	30 to 34	16% (10-23%)	84% (77-90%)	1.7 (0.96-2.7)	4.8% (3.4-6.4%)	8.7% (6.1-12%)	49 (34-67)	3700 (2700-4800)
Ecuador	2000	Female	35 to 39	19% (12-27%)	81% (73-88%)	1.9 (1.1-2.9)	5.3% (3.9-7.1%)	9.7% (7.1-13%)	62 (45-80)	4000 (2900-5100)
Ecuador	2000	Female	40 to 44	18% (12-25%)	82% (75-88%)	1.7 (0.96-2.7)	5.3% (3.8-7.1%)	9.5% (6.9-12%)	69 (49-90)	3800 (2800-5000)
Ecuador	2000	Female	45 to 49	14% (8.7-21%)	86% (79-91%)	1.3 (0.66-2.2)	4.8% (3.4-6.5%)	8.4% (5.9-11%)	72 (51-96)	3500 (2500-4600)
Ecuador	2000	Female	50 to 54	12% (8.1-18%)	88% (82-92%)	1.1 (0.58-1.9)	4.7% (3.2-6.4%)	7.6% (5.3-10%)	72 (50-98)	2900 (2100-4000)
Ecuador	2000	Female	55 to 59	13% (8-19%)	87% (81-92%)	1.2 (0.62-2.2)	4.9% (3.4-6.8%)	7.6% (5.5-10%)	85 (61-120)	3000 (2100-4200)
Ecuador	2000	Female	60 to 64	12% (7.2-18%)	88% (82-93%)	1.2 (0.58-2.2)	4.8% (3.4-6.7%)	7% (5-9.7%)	96 (68-130)	2800 (2000-3900)
Ecuador	2000	Female	65 to 69	10% (6.6-15%)	90% (85-93%)	1 (0.5-1.7)	4.6% (3.1-6.2%)	6.5% (4.4-8.7%)	110 (73-140)	2600 (1800-3500)
Ecuador	2000	Female	70 to 74	9% (5.7-14%)	91% (86-94%)	0.85 (0.41-1.6)	4.1% (2.6-5.8%)	5.6% (3.7-8%)	110 (72-150)	2100 (1400-3000)
Ecuador	2000	Female	75 to 79	8.1% (4.6-13%)	92% (87-95%)	0.68 (0.29-1.3)	3.2% (1.9-4.7%)	4.2% (2.6-6.3%)	92 (56-130)	1400 (820-2000)
Ecuador	2000	Female	80 to 84	7% (3.7-11%)	93% (89-96%)	0.55 (0.24-0.99)	2.5% (1.5-4%)	3.3% (1.9-5.1%)	80 (47-120)	870 (520-1300)
Ecuador	2000	Female	85 to 89	6.5% (3.5-10%)	94% (90-97%)	0.49 (0.22-0.9)	2.2% (1.2-3.5%)	2.7% (1.5-4.3%)	64 (35-100)	490 (270-790)
Ecuador	2000	Female	90 to 94	6.9% (3.9-11%)	93% (89-96%)	0.5 (0.22-0.97)	1.9% (0.92-3.3%)	2.3% (1.1-3.9%)	38 (18-65)	200 (95-340)
Ecuador	2000	Female	95 plus	8.1% (4-14%)	92% (86-96%)	0.55 (0.2-1.1)	1.9% (0.79-3.7%)	2.2% (0.84-4.5%)	14 (5.3-28)	49 (20-95)
Ecuador	2000	Male	All Ages	23% (20-27%)	77% (73-80%)	3.1 (2.5-3.8)	8.4% (7-9.9%)	11% (9.1-13%)	3700 (3100-4400)	160000 (130000-190000)
Ecuador	2000	Male	15 to 19	21% (14-30%)	79% (70-86%)	1.6 (0.8-2.7)	7.2% (4.3-11%)	9.8% (5.5-15%)	96 (53-150)	8100 (4900-12000)
Ecuador	2000	Male	20 to 24	39% (26-51%)	61% (49-74%)	4.2 (2.5-6.5)	16% (11-21%)	20% (13-26%)	280 (190-370)	21000 (15000-28000)
Ecuador	2000	Male	25 to 29	42% (30-54%)	58% (46-70%)	5.1 (3-7.6)	16% (12-21%)	20% (14-27%)	280 (190-370)	20000 (15000-26000)
Ecuador	2000	Male	30 to 34	44% (32-57%)	56% (43-68%)	6.3 (3.9-9.3)	17% (13-22%)	22% (16-29%)	290 (210-380)	19000 (14000-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	2000	Male	35 to 39	42% (30-55%)	58% (45-70%)	6.5 (3.9-9.9)	17% (13-21%)	23% (18-28%)	290 (220-370)	17000 (13000-21000)
Ecuador	2000	Male	40 to 44	40% (29-52%)	60% (48-71%)	6.2 (3.7-9.4)	16% (13-19%)	22% (17-27%)	280 (220-350)	15000 (12000-18000)
Ecuador	2000	Male	45 to 49	44% (31-55%)	56% (45-69%)	6.8 (3.9-11)	16% (13-19%)	22% (18-27%)	310 (250-380)	14000 (12000-17000)
Ecuador	2000	Male	50 to 54	44% (32-56%)	56% (44-68%)	6.9 (3.9-11)	15% (13-19%)	21% (17-25%)	300 (250-370)	12000 (9800-15000)
Ecuador	2000	Male	55 to 59	37% (25-49%)	63% (51-75%)	6 (3.3-9.9)	13% (11-16%)	18% (14-22%)	290 (230-350)	9900 (8000-12000)
Ecuador	2000	Male	60 to 64	30% (19-43%)	70% (57-81%)	4.8 (2.5-8.4)	11% (8.6-14%)	14% (11-18%)	270 (210-340)	7700 (6100-9700)
Ecuador	2000	Male	65 to 69	30% (19-42%)	70% (58-81%)	4.1 (2.1-6.8)	10% (7.6-13%)	13% (9.8-16%)	280 (210-350)	6700 (5100-8400)
Ecuador	2000	Male	70 to 74	29% (19-40%)	71% (60-81%)	3.2 (1.6-5.5)	8.5% (6.4-11%)	11% (8.1-14%)	270 (200-340)	5100 (3900-6600)
Ecuador	2000	Male	75 to 79	23% (14-35%)	77% (65-86%)	2 (0.97-3.8)	6.2% (4.3-8.3%)	7.6% (5.3-10%)	200 (140-270)	2900 (2100-4000)
Ecuador	2000	Male	80 to 84	19% (11-29%)	81% (71-89%)	1.5 (0.71-2.5)	4.5% (3.1-6.1%)	5.5% (3.8-7.6%)	140 (97-190)	1600 (1100-2200)
Ecuador	2000	Male	85 to 89	17% (9.7-27%)	83% (73-90%)	1.3 (0.59-2.2)	3.8% (2.4-5.5%)	4.5% (2.9-6.7%)	96 (61-140)	750 (470-1100)
Ecuador	2000	Male	90 to 94	18% (11-27%)	82% (73-89%)	1.3 (0.56-2.3)	3.5% (2-5.2%)	4% (2.3-6%)	48 (27-73)	260 (150-390)
Ecuador	2000	Male	95 plus	20% (10-34%)	80% (66-90%)	1.3 (0.5-2.7)	3.4% (1.7-5.7%)	3.8% (1.7-6.6%)	14 (6.1-23)	50 (24-83)
Ecuador	2005	Female	All Ages	9.5% (7.7-12%)	90% (88-92%)	0.95 (0.72-1.2)	2.8% (2.3-3.4%)	4.5% (3.7-5.5%)	1200 (990-1500)	43000 (36000-52000)
Ecuador	2005	Female	15 to 19	8.7% (5.2-13%)	91% (87-95%)	0.61 (0.3-1.1)	2.3% (1.3-3.6%)	4% (2-6.7%)	21 (10-35)	2000 (1200-3100)
Ecuador	2005	Female	20 to 24	11% (6.9-17%)	89% (83-93%)	1.2 (0.65-2)	4% (2.7-5.6%)	6.6% (4.1-9.7%)	36 (22-53)	3400 (2300-4900)
Ecuador	2005	Female	25 to 29	13% (7.9-19%)	87% (81-92%)	1.6 (0.88-2.6)	4.4% (3.1-5.9%)	7.7% (5-11%)	41 (27-57)	3600 (2500-4900)
Ecuador	2005	Female	30 to 34	16% (10-23%)	84% (77-90%)	2 (1.1-3.2)	4.4% (3.2-5.9%)	8.3% (5.8-11%)	45 (31-62)	3500 (2600-4600)
Ecuador	2005	Female	35 to 39	18% (12-26%)	82% (74-88%)	2.1 (1.2-3.3)	4.9% (3.7-6.5%)	9.5% (7.1-12%)	58 (43-77)	3900 (2900-5000)
Ecuador	2005	Female	40 to 44	17% (11-25%)	83% (75-89%)	1.9 (1.1-3)	5% (3.7-6.6%)	9.3% (7-12%)	69 (52-90)	3900 (2900-5100)
Ecuador	2005	Female	45 to 49	14% (8.4-20%)	86% (80-92%)	1.5 (0.75-2.5)	4.6% (3.3-6.1%)	8.2% (6-11%)	74 (54-98)	3600 (2700-4700)
Ecuador	2005	Female	50 to 54	12% (7.9-18%)	88% (82-92%)	1.3 (0.65-2.2)	4.4% (3.1-5.9%)	7.4% (5.4-9.8%)	81 (59-110)	3400 (2500-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	2005	Female	55 to 59	13% (7.7-19%)	87% (81-92%)	1.4 (0.71-2.4)	4.8% (3.5-6.6%)	7.6% (5.6-10%)	93 (68-130)	3300 (2400-4500)
Ecuador	2005	Female	60 to 64	12% (7-18%)	88% (82-93%)	1.4 (0.67-2.4)	4.8% (3.4-6.6%)	7.2% (5.4-9.7%)	110 (77-140)	3100 (2300-4300)
Ecuador	2005	Female	65 to 69	10% (6.4-15%)	90% (85-94%)	1.2 (0.56-2)	4.6% (3.3-6.2%)	6.8% (5-8.9%)	120 (87-150)	2800 (2100-3800)
Ecuador	2005	Female	70 to 74	8.7% (5.5-13%)	91% (87-94%)	0.95 (0.46-1.8)	4.2% (2.9-5.9%)	6% (4.3-8.2%)	130 (89-180)	2400 (1700-3400)
Ecuador	2005	Female	75 to 79	7.8% (4.3-13%)	92% (87-96%)	0.76 (0.33-1.4)	3.4% (2.2-4.9%)	4.7% (3.1-6.6%)	120 (76-170)	1700 (1100-2500)
Ecuador	2005	Female	80 to 84	6.7% (3.6-11%)	93% (89-96%)	0.61 (0.28-1.1)	2.8% (1.8-4.1%)	3.7% (2.4-5.4%)	100 (65-150)	1100 (720-1600)
Ecuador	2005	Female	85 to 89	6.3% (3.4-10%)	94% (90-97%)	0.55 (0.24-1)	2.4% (1.4-3.7%)	3.1% (1.8-4.7%)	81 (47-120)	620 (370-940)
Ecuador	2005	Female	90 to 94	6.7% (3.8-11%)	93% (89-96%)	0.57 (0.25-1.1)	2.2% (1.1-3.5%)	2.7% (1.3-4.4%)	53 (26-86)	280 (140-450)
Ecuador	2005	Female	95 plus	8% (3.9-14%)	92% (86-96%)	0.63 (0.22-1.3)	2.2% (1-4.1%)	2.6% (1.1-4.8%)	21 (9-39)	73 (33-130)
Ecuador	2005	Male	All Ages	23% (20-27%)	77% (73-80%)	3.7 (2.9-4.5)	8.6% (7.1-10%)	11% (9.1-13%)	4000 (3300-4700)	170000 (140000-200000)
Ecuador	2005	Male	15 to 19	21% (14-30%)	79% (70-86%)	1.8 (0.9-3.1)	7.3% (4.5-11%)	10% (5.8-16%)	99 (55-150)	8400 (5200-12000)
Ecuador	2005	Male	20 to 24	39% (26-50%)	61% (50-74%)	4.8 (2.8-7.4)	16% (11-21%)	19% (13-27%)	290 (200-410)	23000 (16000-30000)
Ecuador	2005	Male	25 to 29	42% (29-53%)	58% (47-71%)	5.8 (3.4-8.8)	16% (11-20%)	20% (13-26%)	280 (190-390)	21000 (15000-27000)
Ecuador	2005	Male	30 to 34	44% (31-56%)	56% (44-69%)	7.2 (4.5-11)	16% (12-20%)	21% (15-28%)	290 (200-380)	19000 (14000-24000)
Ecuador	2005	Male	35 to 39	41% (30-55%)	59% (45-70%)	7.4 (4.5-11)	16% (12-20%)	21% (16-27%)	280 (210-360)	17000 (13000-21000)
Ecuador	2005	Male	40 to 44	39% (29-52%)	61% (48-71%)	7 (4-11)	15% (12-18%)	21% (16-26%)	290 (220-360)	15000 (12000-19000)
Ecuador	2005	Male	45 to 49	43% (31-56%)	57% (44-69%)	7.8 (4.6-12)	15% (12-19%)	21% (17-26%)	320 (260-400)	15000 (12000-18000)
Ecuador	2005	Male	50 to 54	43% (31-54%)	57% (46-69%)	8 (4.5-12)	14% (12-18%)	20% (16-24%)	350 (280-430)	14000 (11000-17000)
Ecuador	2005	Male	55 to 59	36% (25-49%)	64% (51-75%)	6.9 (3.8-11)	13% (10-15%)	17% (14-21%)	300 (240-370)	10000 (8400-13000)
Ecuador	2005	Male	60 to 64	30% (19-42%)	70% (58-81%)	5.4 (2.9-9.6)	10% (8.1-13%)	14% (11-17%)	280 (220-360)	8300 (6500-10000)
Ecuador	2005	Male	65 to 69	29% (19-41%)	71% (59-81%)	4.6 (2.5-7.6)	9.4% (7.2-12%)	12% (9.6-15%)	290 (220-360)	6900 (5300-8800)
Ecuador	2005	Male	70 to 74	28% (18-39%)	72% (61-82%)	3.6 (1.9-6.2)	8.3% (6.3-11%)	11% (8.1-13%)	290 (220-370)	5700 (4300-7200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	2005	Male	75 to 79	23% (14-35%)	77% (65-86%)	2.3 (1.1-4.3)	6.1% (4.5-8.2%)	7.7% (5.6-10%)	230 (170-310)	3500 (2500-4600)
Ecuador	2005	Male	80 to 84	18% (11-28%)	82% (72-89%)	1.7 (0.81-2.8)	4.5% (3.2-6.1%)	5.6% (4-7.7%)	170 (120-230)	1800 (1300-2500)
Ecuador	2005	Male	85 to 89	17% (9.4-26%)	83% (74-91%)	1.5 (0.68-2.5)	4% (2.6-5.6%)	4.8% (3.2-6.7%)	110 (75-160)	890 (590-1300)
Ecuador	2005	Male	90 to 94	17% (10-26%)	83% (74-90%)	1.4 (0.66-2.7)	3.7% (2.3-5.4%)	4.3% (2.6-6.5%)	63 (38-93)	340 (210-500)
Ecuador	2005	Male	95 plus	19% (10-32%)	81% (68-90%)	1.5 (0.58-3.1)	3.7% (1.9-6%)	4.2% (2.1-7%)	19 (9.5-32)	70 (36-110)
Ecuador	2010	Female	All Ages	10% (8.3-13%)	90% (87-92%)	1.1 (0.79-1.4)	2.9% (2.4-3.5%)	4.7% (3.8-5.7%)	1400 (1100-1700)	47000 (38000-57000)
Ecuador	2010	Female	15 to 19	8.6% (5-14%)	91% (86-95%)	0.61 (0.29-1.1)	2.1% (1.2-3.4%)	3.9% (1.8-6.9%)	19 (8.9-33)	1900 (1100-3000)
Ecuador	2010	Female	20 to 24	13% (7.4-19%)	87% (81-93%)	1.4 (0.7-2.4)	3.8% (2.5-5.3%)	6.4% (3.9-9.6%)	33 (20-51)	3400 (2300-4700)
Ecuador	2010	Female	25 to 29	15% (9-22%)	85% (78-91%)	1.9 (0.98-3.2)	4.3% (3.1-5.8%)	7.7% (4.9-11%)	42 (27-60)	3800 (2600-5100)
Ecuador	2010	Female	30 to 34	16% (9.6-24%)	84% (76-90%)	2 (1.1-3.4)	4.1% (2.9-5.5%)	7.6% (5.2-10%)	43 (30-60)	3500 (2500-4700)
Ecuador	2010	Female	35 to 39	18% (10-27%)	82% (73-90%)	2.1 (1.1-3.3)	4.3% (3.2-5.9%)	8.4% (5.9-11%)	53 (38-70)	3700 (2700-4900)
Ecuador	2010	Female	40 to 44	17% (11-26%)	83% (74-89%)	2 (1.1-3.2)	4.5% (3.3-6.1%)	8.7% (6.4-12%)	67 (48-89)	3900 (2800-5200)
Ecuador	2010	Female	45 to 49	15% (9-23%)	85% (77-91%)	1.7 (0.84-2.9)	4.5% (3.2-6.1%)	8.3% (5.9-11%)	79 (56-110)	3900 (2800-5300)
Ecuador	2010	Female	50 to 54	14% (9-21%)	86% (79-91%)	1.5 (0.77-2.6)	4.5% (3.1-6.3%)	7.8% (5.6-10%)	94 (67-130)	4000 (2800-5400)
Ecuador	2010	Female	55 to 59	14% (8.9-21%)	86% (79-91%)	1.6 (0.81-2.9)	4.8% (3.4-6.7%)	7.9% (5.8-11%)	110 (83-160)	4100 (3000-5800)
Ecuador	2010	Female	60 to 64	13% (7.7-20%)	87% (80-92%)	1.6 (0.74-2.8)	5% (3.5-6.9%)	7.5% (5.5-10%)	120 (90-170)	3700 (2700-5100)
Ecuador	2010	Female	65 to 69	11% (6.5-16%)	89% (84-94%)	1.3 (0.62-2.2)	4.6% (3.4-6.3%)	6.9% (5.3-9.2%)	140 (100-180)	3300 (2500-4600)
Ecuador	2010	Female	70 to 74	8.8% (5.7-13%)	91% (87-94%)	0.99 (0.47-1.8)	4.2% (3-5.8%)	6.1% (4.5-8.3%)	140 (100-190)	2800 (2000-3900)
Ecuador	2010	Female	75 to 79	7.7% (4.3-12%)	92% (88-96%)	0.77 (0.32-1.5)	3.5% (2.3-4.9%)	4.8% (3.3-6.6%)	140 (95-200)	2100 (1400-2900)
Ecuador	2010	Female	80 to 84	6.6% (3.4-11%)	93% (89-97%)	0.61 (0.27-1.1)	2.8% (1.9-4.1%)	3.8% (2.6-5.5%)	130 (84-180)	1400 (920-2000)
Ecuador	2010	Female	85 to 89	6.1% (3.2-10%)	94% (90-97%)	0.55 (0.24-1)	2.4% (1.5-3.7%)	3.1% (1.9-4.8%)	99 (61-150)	760 (480-1200)
Ecuador	2010	Female	90 to 94	6.5% (3.7-11%)	94% (89-96%)	0.56 (0.24-1.1)	2.3% (1.2-3.6%)	2.7% (1.5-4.5%)	66 (35-110)	340 (190-550)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	2010	Female	95 plus	7.7% (3.8-13%)	92% (87-96%)	0.62 (0.23-1.3)	2.3% (1-4.1%)	2.7% (1.2-4.9%)	27 (12-49)	92 (43-170)
Ecuador	2010	Male	All Ages	22% (18-26%)	78% (74-82%)	3.7 (2.9-4.7)	8.4% (6.7-10%)	10% (8.7-12%)	4000 (3300-4800)	170000 (140000-200000)
Ecuador	2010	Male	15 to 19	16% (9.4-24%)	84% (76-91%)	1.4 (0.67-2.6)	5.9% (3.6-9%)	8.4% (4.4-14%)	76 (39-120)	6800 (4100-10000)
Ecuador	2010	Male	20 to 24	33% (20-46%)	67% (54-80%)	4.1 (2.2-6.8)	14% (9.1-19%)	17% (11-25%)	250 (160-370)	20000 (13000-27000)
Ecuador	2010	Male	25 to 29	40% (26-53%)	60% (47-74%)	5.7 (3.2-8.9)	15% (10-20%)	19% (12-26%)	280 (180-380)	20000 (14000-27000)
Ecuador	2010	Male	30 to 34	41% (27-56%)	59% (44-73%)	7.1 (4-11)	15% (10-19%)	19% (13-27%)	260 (170-370)	18000 (13000-23000)
Ecuador	2010	Male	35 to 39	40% (26-54%)	60% (46-74%)	7.4 (4.2-12)	14% (11-19%)	20% (14-26%)	270 (190-370)	17000 (12000-21000)
Ecuador	2010	Male	40 to 44	39% (26-53%)	61% (47-74%)	7.3 (3.9-12)	14% (11-18%)	20% (15-25%)	280 (210-370)	15000 (11000-19000)
Ecuador	2010	Male	45 to 49	40% (25-54%)	60% (46-75%)	7.5 (4-12)	14% (10-17%)	19% (15-24%)	310 (230-380)	14000 (11000-18000)
Ecuador	2010	Male	50 to 54	38% (26-52%)	62% (48-74%)	7.2 (3.9-12)	13% (9.9-16%)	18% (14-22%)	330 (260-410)	13000 (11000-17000)
Ecuador	2010	Male	55 to 59	34% (22-49%)	66% (51-78%)	6.7 (3.6-11)	11% (8.8-14%)	15% (12-19%)	330 (260-420)	12000 (9100-15000)
Ecuador	2010	Male	60 to 64	32% (20-46%)	68% (54-80%)	6.2 (3.2-11)	10% (7.9-13%)	14% (11-17%)	300 (230-390)	9000 (6800-12000)
Ecuador	2010	Male	65 to 69	32% (20-44%)	68% (56-80%)	5.3 (2.7-8.9)	9.5% (7.1-12%)	12% (9.6-16%)	330 (250-430)	8100 (6000-11000)
Ecuador	2010	Male	70 to 74	28% (17-40%)	72% (60-83%)	3.7 (1.8-6.5)	7.9% (6-10%)	10% (7.7-13%)	310 (240-400)	6100 (4600-8000)
Ecuador	2010	Male	75 to 79	23% (13-34%)	77% (66-87%)	2.4 (1.1-4.5)	6% (4.4-8.1%)	7.6% (5.5-10%)	260 (190-340)	3900 (2800-5200)
Ecuador	2010	Male	80 to 84	18% (10-29%)	82% (71-90%)	1.7 (0.82-3)	4.6% (3.3-6.2%)	5.7% (4.1-8%)	190 (140-270)	2200 (1500-3000)
Ecuador	2010	Male	85 to 89	16% (9.1-26%)	84% (74-91%)	1.5 (0.67-2.6)	4% (2.7-5.7%)	4.8% (3.2-6.9%)	140 (90-190)	1100 (720-1500)
Ecuador	2010	Male	90 to 94	17% (9.8-26%)	83% (74-90%)	1.4 (0.64-2.8)	3.7% (2.3-5.4%)	4.3% (2.7-6.4%)	72 (44-110)	390 (240-570)
Ecuador	2010	Male	95 plus	19% (9.7-31%)	81% (69-90%)	1.5 (0.56-3.1)	3.7% (2-6%)	4.2% (2.2-7%)	22 (12-38)	82 (44-130)
Ecuador	2016	Female	All Ages	11% (8.5-13%)	89% (87-92%)	1.1 (0.76-1.6)	2.9% (2.4-3.6%)	4.6% (3.7-5.7%)	1500 (1200-1900)	50000 (40000-62000)
Ecuador	2016	Female	15 to 19	8.7% (5.1-14%)	91% (86-95%)	0.62 (0.27-1.2)	1.9% (1.1-3.2%)	3.5% (1.5-6.3%)	17 (7.2-30)	1800 (970-2800)
Ecuador	2016	Female	20 to 24	13% (7.6-19%)	87% (81-92%)	1.4 (0.66-2.5)	3.5% (2.4-4.9%)	5.8% (3.4-8.7%)	29 (17-45)	3200 (2100-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	2016	Female	25 to 29	15% (8.9-21%)	85% (79-91%)	1.9 (0.96-3.3)	3.8% (2.8-5.1%)	6.7% (4.2-9.4%)	36 (23-53)	3600 (2500-4900)
Ecuador	2016	Female	30 to 34	16% (9.7-24%)	84% (76-90%)	2.1 (1.1-3.5)	3.7% (2.7-5%)	6.8% (4.6-9.5%)	40 (27-57)	3400 (2500-4700)
Ecuador	2016	Female	35 to 39	18% (11-26%)	82% (74-89%)	2.1 (1.1-3.5)	4% (2.9-5.5%)	7.8% (5.5-10%)	51 (35-70)	3700 (2700-5000)
Ecuador	2016	Female	40 to 44	18% (11-26%)	82% (74-89%)	2 (1-3.4)	4.1% (3-5.6%)	8.1% (5.8-11%)	65 (46-88)	4000 (2900-5400)
Ecuador	2016	Female	45 to 49	16% (8.9-23%)	84% (77-91%)	1.7 (0.81-3)	4.2% (2.9-5.8%)	7.9% (5.5-11%)	82 (57-110)	4200 (3000-5800)
Ecuador	2016	Female	50 to 54	14% (9-21%)	86% (79-91%)	1.5 (0.73-2.7)	4.3% (3-5.9%)	7.5% (5.3-10%)	99 (69-130)	4300 (3000-5900)
Ecuador	2016	Female	55 to 59	14% (8.8-21%)	86% (79-91%)	1.6 (0.78-3)	4.6% (3.2-6.3%)	7.6% (5.6-10%)	120 (89-170)	4500 (3200-6300)
Ecuador	2016	Female	60 to 64	13% (7.5-20%)	87% (80-93%)	1.5 (0.71-2.9)	4.6% (3.3-6.6%)	7.2% (5.2-9.7%)	150 (110-210)	4600 (3300-6600)
Ecuador	2016	Female	65 to 69	11% (6.4-16%)	89% (84-94%)	1.3 (0.6-2.2)	4.4% (3.2-6.1%)	6.7% (5.1-8.9%)	150 (110-210)	3700 (2700-5200)
Ecuador	2016	Female	70 to 74	8.8% (5.6-13%)	91% (87-94%)	0.98 (0.43-1.8)	4% (2.8-5.6%)	6% (4.3-8%)	160 (110-220)	3200 (2200-4300)
Ecuador	2016	Female	75 to 79	7.7% (4.3-13%)	92% (87-96%)	0.77 (0.31-1.5)	3.3% (2.2-4.6%)	4.7% (3.2-6.5%)	160 (110-210)	2300 (1600-3400)
Ecuador	2016	Female	80 to 84	6.7% (3.5-11%)	93% (89-97%)	0.62 (0.26-1.2)	2.8% (1.8-4.1%)	3.8% (2.6-5.4%)	140 (96-210)	1600 (1100-2300)
Ecuador	2016	Female	85 to 89	6.2% (3.3-10%)	94% (90-97%)	0.56 (0.25-1.1)	2.4% (1.4-3.7%)	3.1% (1.9-4.7%)	120 (72-180)	930 (560-1400)
Ecuador	2016	Female	90 to 94	6.7% (3.8-11%)	93% (89-96%)	0.57 (0.25-1.1)	2.2% (1.2-3.7%)	2.7% (1.5-4.4%)	83 (44-140)	430 (230-720)
Ecuador	2016	Female	95 plus	7.9% (3.9-14%)	92% (86-96%)	0.63 (0.23-1.4)	2.3% (1.1-4.1%)	2.7% (1.2-5%)	33 (15-61)	110 (53-210)
Ecuador	2016	Male	All Ages	22% (18-27%)	78% (73-82%)	3.7 (2.7-5.1)	8.1% (6.6-9.7%)	9.9% (8.3-12%)	4200 (3400-5100)	170000 (140000-210000)
Ecuador	2016	Male	15 to 19	15% (9.3-23%)	85% (77-91%)	1.4 (0.63-2.6)	5.3% (3.2-8%)	7.4% (3.7-12%)	65 (33-110)	6000 (3600-9000)
Ecuador	2016	Male	20 to 24	33% (20-45%)	67% (55-80%)	4.1 (2-6.8)	13% (8.6-17%)	16% (9.9-23%)	220 (140-330)	18000 (12000-26000)
Ecuador	2016	Male	25 to 29	39% (26-53%)	61% (47-74%)	5.6 (2.9-9.2)	14% (9.6-18%)	17% (11-24%)	260 (160-360)	19000 (13000-26000)
Ecuador	2016	Male	30 to 34	41% (26-56%)	59% (44-74%)	6.9 (3.8-11)	13% (9.5-17%)	18% (12-24%)	250 (160-350)	17000 (12000-23000)
Ecuador	2016	Male	35 to 39	39% (26-53%)	61% (47-74%)	7.2 (4.1-12)	13% (9.8-17%)	18% (13-24%)	260 (180-350)	16000 (12000-21000)
Ecuador	2016	Male	40 to 44	38% (26-53%)	62% (47-74%)	7.1 (3.7-12)	12% (9.6-16%)	18% (13-23%)	270 (200-360)	15000 (11000-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ecuador	2016	Male	45 to 49	40% (25-54%)	60% (46-75%)	7.3 (3.7-12)	13% (9.7-16%)	18% (14-23%)	310 (240-400)	15000 (12000-19000)
Ecuador	2016	Male	50 to 54	38% (26-51%)	62% (49-74%)	7 (3.5-12)	12% (9.3-15%)	17% (13-21%)	340 (270-440)	14000 (11000-18000)
Ecuador	2016	Male	55 to 59	34% (22-48%)	66% (52-78%)	6.5 (3.2-11)	10% (8-13%)	15% (12-18%)	350 (270-440)	12000 (9500-16000)
Ecuador	2016	Male	60 to 64	32% (20-46%)	68% (54-80%)	6.1 (2.9-11)	9.5% (7.3-12%)	13% (10-16%)	370 (280-480)	11000 (8400-14000)
Ecuador	2016	Male	65 to 69	31% (20-44%)	69% (56-80%)	5.2 (2.6-8.9)	9% (6.7-12%)	12% (9.2-15%)	360 (270-460)	8900 (6600-11000)
Ecuador	2016	Male	70 to 74	27% (17-39%)	73% (61-83%)	3.6 (1.7-6.5)	7.6% (5.5-9.8%)	9.8% (7.4-12%)	350 (260-460)	6900 (5100-9000)
Ecuador	2016	Male	75 to 79	22% (13-34%)	78% (66-87%)	2.3 (1.1-4.5)	5.8% (4.1-7.8%)	7.3% (5.3-9.9%)	290 (200-390)	4400 (3100-5900)
Ecuador	2016	Male	80 to 84	18% (10-28%)	82% (72-90%)	1.7 (0.76-3)	4.4% (3.1-5.9%)	5.5% (3.9-7.6%)	220 (150-300)	2400 (1700-3400)
Ecuador	2016	Male	85 to 89	16% (8.9-25%)	84% (75-91%)	1.4 (0.63-2.6)	3.8% (2.6-5.4%)	4.6% (3.1-6.5%)	160 (100-220)	1200 (820-1800)
Ecuador	2016	Male	90 to 94	16% (9.6-25%)	84% (75-90%)	1.4 (0.58-2.7)	3.5% (2.2-5.1%)	4.1% (2.5-6%)	85 (52-120)	460 (290-670)
Ecuador	2016	Male	95 plus	18% (9.5-31%)	82% (69-91%)	1.4 (0.55-3)	3.5% (1.9-5.7%)	4% (2.1-6.7%)	25 (13-42)	94 (51-150)
Peru	1990	Female	All Ages	68% (64-71%)	32% (29-36%)	1.6 (1.4-1.9)	2.6% (1.9-3.3%)	4.4% (2.9-6.1%)	2600 (1700-3600)	110000 (77000-130000)
Peru	1990	Female	15 to 19	58% (48-65%)	42% (35-52%)	1.3 (0.75-1.9)	3.6% (1.8-5.8%)	6.6% (2.9-11%)	68 (31-120)	5900 (3000-9400)
Peru	1990	Female	20 to 24	84% (78-88%)	16% (12-22%)	2.5 (1.6-3.6)	7% (4.2-10%)	12% (6.7-17%)	150 (83-230)	12000 (7500-17000)
Peru	1990	Female	25 to 29	87% (83-90%)	13% (10-17%)	3 (2-4.2)	7.6% (4.9-11%)	14% (8.6-19%)	160 (99-230)	12000 (7400-16000)
Peru	1990	Female	30 to 34	89% (85-91%)	11% (8.8-15%)	3.2 (2.1-4.5)	7.3% (4.9-10%)	13% (8.6-19%)	160 (100-240)	11000 (6900-15000)
Peru	1990	Female	35 to 39	89% (85-91%)	11% (8.8-15%)	3.1 (2-4.3)	7.1% (4.5-10%)	13% (8-17%)	170 (100-250)	9800 (6100-14000)
Peru	1990	Female	40 to 44	89% (86-92%)	11% (8.3-14%)	3.1 (1.9-4.5)	7.1% (4.3-10%)	13% (7.6-18%)	170 (100-240)	8600 (5100-12000)
Peru	1990	Female	45 to 49	90% (86-92%)	9.8% (7.5-14%)	3.2 (2-4.9)	7.4% (4.5-11%)	13% (8-19%)	190 (120-280)	8500 (5100-12000)
Peru	1990	Female	50 to 54	90% (86-92%)	10% (7.9-14%)	3.2 (1.9-4.9)	7.7% (4.6-12%)	13% (7.9-19%)	220 (130-320)	8500 (5000-13000)
Peru	1990	Female	55 to 59	88% (83-91%)	12% (9.1-17%)	3.1 (1.9-4.8)	7.7% (4.5-12%)	12% (7.5-18%)	250 (150-370)	8200 (4800-12000)
Peru	1990	Female	60 to 64	85% (80-89%)	15% (11-20%)	3 (1.8-4.6)	7.1% (3.5-11%)	10% (5.4-16%)	260 (130-410)	7300 (3700-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	1990	Female	65 to 69	82% (75-87%)	18% (13-25%)	2.8 (1.7-4.1)	6.1% (3-9.7%)	8.8% (4.4-14%)	250 (120-390)	5800 (2700-9300)
Peru	1990	Female	70 to 74	79% (71-84%)	21% (16-29%)	2.4 (1.3-3.7)	4.6% (1.5-8.3%)	6.5% (2.4-12%)	210 (76-370)	3900 (1300-6900)
Peru	1990	Female	75 to 79	76% (66-84%)	24% (16-34%)	2 (1.1-3.2)	3.2% (0.35-6.8%)	4.3% (0.58-9%)	160 (23-330)	2300 (260-4800)
Peru	1990	Female	80 to 84	70% (58-79%)	30% (21-42%)	1.7 (0.96-2.6)	2.2% (-0.29-5.3%)	2.8% (-0.37-6.7%)	99 (-13-240)	1100 (-140-2500)
Peru	1990	Female	85 to 89	64% (51-75%)	36% (25-49%)	1.5 (0.86-2.4)	1.4% (-0.72-4.3%)	1.7% (-0.96-5.2%)	48 (-27-150)	370 (-190-1100)
Peru	1990	Female	90 to 94	62% (50-73%)	38% (27-50%)	1.5 (0.77-2.4)	0.78% (-1.3-3.6%)	0.83% (-1.8-4.4%)	12 (-27-67)	72 (-120-350)
Peru	1990	Female	95 plus	62% (46-74%)	38% (26-54%)	1.4 (0.67-2.4)	0.46% (-1.8-3.9%)	0.36% (-2.4-4.6%)	1.8 (-12-23)	9.3 (-37-79)
Peru	1990	Male	All Ages	75% (71-79%)	25% (21-29%)	2.8 (2.4-3.3)	5.9% (5-7%)	9.3% (7.3-11%)	6500 (5000-8000)	270000 (230000-320000)
Peru	1990	Male	15 to 19	61% (50-70%)	39% (30-50%)	1.7 (0.97-2.5)	6.6% (4.1-9.4%)	9.1% (5.1-13%)	150 (85-220)	13000 (7800-18000)
Peru	1990	Male	20 to 24	93% (90-95%)	7.1% (4.9-10%)	3.9 (2.5-5.4)	15% (11-19%)	19% (13-25%)	400 (280-530)	32000 (23000-42000)
Peru	1990	Male	25 to 29	96% (94-97%)	4.4% (3.2-6.3%)	5 (3.4-7)	18% (13-22%)	24% (17-31%)	450 (320-610)	33000 (24000-43000)
Peru	1990	Male	30 to 34	97% (96-98%)	2.9% (2-4.3%)	5.6 (4-7.6)	18% (14-22%)	25% (19-32%)	430 (320-550)	28000 (22000-35000)
Peru	1990	Male	35 to 39	98% (97-99%)	2.2% (1.5-3.4%)	5.9 (3.9-8.3)	18% (14-23%)	26% (19-33%)	460 (330-600)	27000 (20000-34000)
Peru	1990	Male	40 to 44	98% (97-99%)	2% (1.2-3%)	6 (3.8-8.6)	18% (14-23%)	26% (19-33%)	470 (340-610)	24000 (18000-31000)
Peru	1990	Male	45 to 49	98% (97-99%)	2.1% (1.3-3.3%)	6 (3.7-9.2)	18% (13-23%)	26% (19-32%)	530 (370-680)	24000 (17000-30000)
Peru	1990	Male	50 to 54	98% (97-99%)	1.9% (1.2-3%)	6.1 (3.7-9.2)	18% (13-22%)	24% (17-30%)	580 (400-730)	23000 (16000-29000)
Peru	1990	Male	55 to 59	98% (97-99%)	1.7% (1-2.6%)	6.1 (3.9-9)	17% (12-21%)	23% (17-29%)	630 (450-800)	21000 (15000-27000)
Peru	1990	Male	60 to 64	98% (97-99%)	1.6% (0.93-2.6%)	5.6 (3.4-8.4)	15% (9.8-20%)	19% (13-26%)	630 (410-820)	18000 (12000-23000)
Peru	1990	Male	65 to 69	98% (97-99%)	1.8% (1.1-2.9%)	4.8 (3-7.2)	13% (7.8-17%)	16% (9.9-22%)	570 (350-780)	13000 (8500-19000)
Peru	1990	Male	70 to 74	98% (96-98%)	2.5% (1.5-3.9%)	4 (2.3-6)	9.6% (5.1-15%)	12% (6.5-19%)	450 (240-690)	8500 (4600-13000)
Peru	1990	Male	75 to 79	96% (93-98%)	3.8% (2.1-6.6%)	3.2 (1.7-5.1)	7.1% (2.9-12%)	8.6% (3.4-15%)	340 (130-600)	5000 (2000-8600)
Peru	1990	Male	80 to 84	95% (91-97%)	5.5% (3.2-9.3%)	2.6 (1.5-4.1)	5.4% (2.1-9.7%)	6.3% (2.3-12%)	210 (77-390)	2300 (890-4100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	1990	Male	85 to 89	93% (89-96%)	6.6% (3.6-11%)	2.4 (1.4-3.8)	3.9% (1-7.8%)	4.5% (1-9.1%)	110 (24-220)	830 (220-1700)
Peru	1990	Male	90 to 94	93% (89-97%)	6.7% (3.4-11%)	2.4 (1.3-3.9)	2.9% (-0.015-6.9%)	3.1% (-0.36-7.9%)	37 (-4.4-97)	210 (-1.1-500)
Peru	1990	Male	95 plus	94% (88-97%)	6.4% (2.7-12%)	2.3 (1.2-4)	2.2% (-0.85-7.2%)	2.2% (-1.5-8.1%)	8.1 (-5.6-30)	33 (-13-100)
Peru	1995	Female	All Ages	69% (65-72%)	31% (28-35%)	1.6 (1.3-1.9)	2.9% (2.1-3.7%)	4.5% (2.8-6.4%)	2700 (1600-3800)	110000 (78000-140000)
Peru	1995	Female	15 to 19	58% (48-66%)	42% (34-52%)	1.2 (0.7-1.8)	3.2% (1.6-5.4%)	5.9% (2.6-9.7%)	65 (29-110)	5700 (2800-8900)
Peru	1995	Female	20 to 24	84% (79-88%)	16% (12-21%)	2.4 (1.5-3.3)	6.3% (3.7-9%)	11% (6-16%)	140 (77-210)	11000 (6800-16000)
Peru	1995	Female	25 to 29	87% (83-90%)	13% (9.9-17%)	2.7 (1.8-3.9)	6.7% (4.2-9.1%)	12% (7.2-17%)	150 (93-220)	11000 (7200-16000)
Peru	1995	Female	30 to 34	89% (85-91%)	11% (8.9-15%)	2.9 (2-4)	6.3% (4.1-8.8%)	12% (7.4-16%)	160 (99-220)	10000 (6800-15000)
Peru	1995	Female	35 to 39	89% (85-91%)	11% (8.7-15%)	2.8 (1.8-4)	6.2% (4-8.7%)	11% (6.9-15%)	170 (110-240)	9800 (6400-14000)
Peru	1995	Female	40 to 44	89% (86-92%)	11% (8.3-14%)	2.8 (1.8-4.1)	6.2% (3.9-8.9%)	11% (6.7-16%)	180 (110-260)	9100 (5700-13000)
Peru	1995	Female	45 to 49	90% (86-93%)	9.8% (7.4-14%)	2.9 (1.8-4.5)	6.6% (4-9.9%)	11% (7.1-16%)	200 (120-300)	8900 (5400-13000)
Peru	1995	Female	50 to 54	90% (86-92%)	10% (7.9-14%)	2.9 (1.8-4.5)	7.1% (4-11%)	12% (7.1-17%)	240 (140-350)	9000 (5100-14000)
Peru	1995	Female	55 to 59	88% (83-91%)	12% (9.1-17%)	2.9 (1.8-4.3)	7.1% (4.1-11%)	11% (6.7-16%)	260 (160-390)	8800 (5200-13000)
Peru	1995	Female	60 to 64	85% (80-89%)	15% (11-20%)	2.8 (1.7-4.2)	6.4% (3.2-10%)	9.3% (4.9-15%)	280 (150-450)	8000 (4100-13000)
Peru	1995	Female	65 to 69	82% (76-87%)	18% (13-24%)	2.6 (1.6-3.8)	5.5% (2.5-9%)	7.7% (3.7-13%)	280 (130-450)	6400 (3000-10000)
Peru	1995	Female	70 to 74	79% (72-85%)	21% (15-28%)	2.2 (1.2-3.4)	4.1% (1.1-7.6%)	5.6% (1.7-10%)	230 (70-410)	4100 (1200-7700)
Peru	1995	Female	75 to 79	76% (67-84%)	24% (16-33%)	1.8 (0.99-3)	2.7% (-0.017-6%)	3.6% (0.099-7.8%)	160 (4.4-350)	2300 (-14-5000)
Peru	1995	Female	80 to 84	71% (60-80%)	29% (20-40%)	1.6 (0.91-2.4)	1.7% (-0.62-4.5%)	2.2% (-0.81-5.7%)	98 (-35-260)	1000 (-360-2700)
Peru	1995	Female	85 to 89	65% (52-75%)	35% (25-48%)	1.4 (0.8-2.2)	1% (-0.95-3.5%)	1.2% (-1.2-4.3%)	47 (-49-170)	370 (-340-1300)
Peru	1995	Female	90 to 94	63% (51-74%)	37% (26-49%)	1.4 (0.73-2.2)	0.44% (-1.7-3.1%)	0.42% (-2.1-3.7%)	8.9 (-46-81)	58 (-220-420)
Peru	1995	Female	95 plus	63% (48-75%)	37% (25-52%)	1.3 (0.65-2.3)	0.14% (-2.1-3.3%)	-0.02% (-2.7-3.8%)	-0.15 (-20-27)	4 (-60-93)
Peru	1995	Male	All Ages	76% (73-80%)	24% (20-27%)	2.8 (2.4-3.3)	6.9% (5.7-8.1%)	10% (7.8-12%)	6800 (5200-8600)	290000 (240000-340000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	1995	Male	15 to 19	61% (50-70%)	39% (30-50%)	1.6 (0.92-2.4)	6.2% (3.9-8.6%)	8.8% (5-13%)	140 (75-200)	12000 (7600-17000)
Peru	1995	Male	20 to 24	93% (90-95%)	7.1% (4.9-10%)	3.7 (2.4-5.1)	14% (11-18%)	19% (13-25%)	380 (260-500)	31000 (22000-40000)
Peru	1995	Male	25 to 29	96% (94-97%)	4.4% (3.2-6.1%)	4.7 (3.2-6.7)	17% (13-21%)	23% (17-29%)	460 (330-600)	34000 (25000-43000)
Peru	1995	Male	30 to 34	97% (96-98%)	2.8% (1.9-4.2%)	5.4 (3.7-7.3)	17% (13-21%)	24% (18-30%)	470 (350-600)	31000 (24000-39000)
Peru	1995	Male	35 to 39	98% (97-99%)	2.2% (1.5-3.3%)	5.6 (3.7-8)	17% (13-22%)	25% (18-31%)	490 (350-640)	29000 (21000-37000)
Peru	1995	Male	40 to 44	98% (97-99%)	1.9% (1.2-3%)	5.8 (3.7-8.2)	17% (13-22%)	25% (18-31%)	510 (360-650)	26000 (19000-33000)
Peru	1995	Male	45 to 49	98% (97-99%)	2% (1.3-3.2%)	5.7 (3.6-8.7)	17% (13-22%)	25% (18-31%)	570 (400-740)	26000 (18000-33000)
Peru	1995	Male	50 to 54	98% (97-99%)	1.9% (1.2-2.9%)	5.8 (3.5-8.7)	17% (12-21%)	23% (16-29%)	600 (410-770)	23000 (16000-30000)
Peru	1995	Male	55 to 59	98% (97-99%)	1.6% (1-2.5%)	5.8 (3.7-8.5)	16% (11-20%)	22% (15-27%)	660 (460-870)	22000 (16000-29000)
Peru	1995	Male	60 to 64	98% (97-99%)	1.6% (0.92-2.6%)	5.3 (3.2-8)	14% (9.2-19%)	18% (12-24%)	680 (440-920)	19000 (13000-26000)
Peru	1995	Male	65 to 69	98% (97-99%)	1.8% (1.1-2.9%)	4.6 (2.9-6.8)	12% (7.3-16%)	15% (9.2-21%)	620 (390-890)	15000 (9100-21000)
Peru	1995	Male	70 to 74	98% (96-98%)	2.4% (1.5-3.8%)	3.8 (2.3-5.8)	8.7% (4.6-14%)	11% (5.7-17%)	500 (260-790)	9400 (5000-15000)
Peru	1995	Male	75 to 79	96% (94-98%)	3.8% (2.1-6.4%)	3 (1.6-4.8)	6.3% (2.6-11%)	7.7% (3.1-14%)	360 (150-640)	5200 (2200-9200)
Peru	1995	Male	80 to 84	95% (91-97%)	5.3% (3-9.1%)	2.5 (1.5-3.8)	4.7% (1.7-8.6%)	5.5% (1.9-10%)	230 (75-430)	2400 (860-4600)
Peru	1995	Male	85 to 89	94% (89-96%)	6.4% (3.5-11%)	2.3 (1.3-3.6)	3.4% (0.68-7.1%)	3.8% (0.63-8.1%)	120 (20-260)	960 (200-2000)
Peru	1995	Male	90 to 94	94% (89-97%)	6.5% (3.3-11%)	2.3 (1.2-3.7)	2.4% (-0.26-6.2%)	2.6% (-0.66-7.2%)	41 (-11-110)	230 (-24-580)
Peru	1995	Male	95 plus	94% (88-97%)	6.2% (2.6-12%)	2.2 (1.1-3.8)	1.9% (-0.95-6.4%)	1.8% (-1.7-7%)	8.7 (-7.6-33)	36 (-18-120)
Peru	2000	Female	All Ages	69% (66-72%)	31% (28-34%)	1.5 (1.3-1.8)	2.6% (1.9-3.3%)	4% (2.5-5.7%)	2000 (1200-2900)	79000 (58000-100000)
Peru	2000	Female	15 to 19	60% (54-66%)	40% (34-46%)	1.2 (0.72-1.7)	2.3% (1.3-3.6%)	4.6% (2.1-7.5%)	37 (17-62)	3700 (2100-5700)
Peru	2000	Female	20 to 24	84% (81-87%)	16% (13-19%)	2.2 (1.4-3.2)	4.5% (2.8-6.4%)	8.2% (4.6-12%)	81 (44-120)	7600 (4600-11000)
Peru	2000	Female	25 to 29	86% (84-89%)	14% (11-16%)	2.5 (1.7-3.5)	4.7% (3-6.6%)	9% (5.4-13%)	88 (53-130)	7300 (4700-10000)
Peru	2000	Female	30 to 34	88% (85-90%)	12% (9.9-15%)	2.7 (1.8-3.7)	4.5% (2.9-6.2%)	9% (5.8-12%)	95 (61-130)	6900 (4500-9500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	2000	Female	35 to 39	88% (85-90%)	12% (9.9-15%)	2.5 (1.7-3.6)	4.4% (2.9-6.2%)	9% (5.8-12%)	110 (71-160)	6800 (4500-9600)
Peru	2000	Female	40 to 44	88% (85-91%)	12% (9.2-15%)	2.5 (1.6-3.7)	4.6% (3-6.7%)	9.4% (6.1-13%)	130 (83-180)	6800 (4400-9600)
Peru	2000	Female	45 to 49	90% (87-92%)	10% (8.2-13%)	2.7 (1.7-4.2)	5.1% (3.1-7.6%)	9.8% (6.5-14%)	160 (100-230)	7200 (4500-11000)
Peru	2000	Female	50 to 54	89% (86-91%)	11% (8.7-14%)	2.7 (1.7-4.1)	5.7% (3.2-8.7%)	10% (6.3-15%)	190 (120-280)	7400 (4400-11000)
Peru	2000	Female	55 to 59	86% (81-89%)	14% (11-19%)	2.5 (1.5-3.7)	5.5% (3.3-8.3%)	9.2% (5.8-13%)	200 (120-280)	6600 (4100-9800)
Peru	2000	Female	60 to 64	84% (79-87%)	16% (13-21%)	2.4 (1.5-3.7)	5.1% (2.7-8.3%)	7.9% (4.6-12%)	220 (120-340)	6200 (3300-9900)
Peru	2000	Female	65 to 69	82% (75-87%)	18% (13-25%)	2.4 (1.5-3.6)	4.5% (2.2-7.4%)	6.8% (3.7-11%)	230 (120-360)	5300 (2600-8700)
Peru	2000	Female	70 to 74	79% (72-84%)	21% (16-28%)	2.1 (1.2-3.2)	3.4% (1.1-6.3%)	5% (1.8-9%)	200 (73-350)	3600 (1200-6700)
Peru	2000	Female	75 to 79	76% (67-84%)	24% (16-33%)	1.7 (0.93-2.8)	2.4% (0.27-5.1%)	3.4% (0.53-6.9%)	150 (23-310)	2100 (220-4500)
Peru	2000	Female	80 to 84	70% (59-80%)	30% (20-41%)	1.5 (0.84-2.3)	1.6% (-0.29-3.9%)	2% (-0.34-5%)	94 (-15-230)	1000 (-180-2500)
Peru	2000	Female	85 to 89	64% (52-75%)	36% (25-48%)	1.3 (0.73-2.1)	0.98% (-0.61-3.2%)	1.2% (-0.87-3.9%)	49 (-36-160)	380 (-240-1200)
Peru	2000	Female	90 to 94	62% (49-73%)	38% (27-51%)	1.3 (0.68-2.1)	0.42% (-1.3-2.5%)	0.42% (-1.7-3.2%)	11 (-47-85)	71 (-210-430)
Peru	2000	Female	95 plus	62% (47-75%)	38% (25-53%)	1.2 (0.61-2.1)	0.15% (-1.8-2.9%)	-0.0018% (-2.4-3.3%)	-0.033 (-22-29)	5.7 (-66-100)
Peru	2000	Male	All Ages	80% (77-83%)	20% (17-23%)	2.8 (2.4-3.3)	6.9% (5.8-8.2%)	9.5% (7.3-12%)	5900 (4400-7600)	250000 (200000-300000)
Peru	2000	Male	15 to 19	68% (62-74%)	32% (26-38%)	1.7 (1-2.6)	5.7% (3.8-7.8%)	8.5% (5.1-12%)	100 (60-150)	10000 (6500-14000)
Peru	2000	Male	20 to 24	94% (93-96%)	5.7% (4.4-7.4%)	3.7 (2.4-5.1)	13% (9.3-16%)	17% (12-22%)	280 (190-380)	25000 (19000-32000)
Peru	2000	Male	25 to 29	96% (95-97%)	3.7% (2.9-4.8%)	4.6 (3.1-6.4)	14% (11-18%)	20% (14-26%)	320 (230-430)	26000 (19000-33000)
Peru	2000	Male	30 to 34	98% (97-98%)	2.5% (1.8-3.3%)	5.2 (3.6-7)	14% (11-18%)	21% (16-27%)	350 (260-460)	25000 (19000-31000)
Peru	2000	Male	35 to 39	98% (97-99%)	1.9% (1.3-2.6%)	5.4 (3.6-7.6)	14% (11-18%)	22% (16-27%)	400 (280-520)	24000 (18000-31000)
Peru	2000	Male	40 to 44	98% (98-99%)	1.7% (1.1-2.3%)	5.6 (3.6-7.9)	15% (11-18%)	22% (16-28%)	420 (290-550)	22000 (16000-29000)
Peru	2000	Male	45 to 49	98% (97-99%)	1.8% (1.3-2.6%)	5.5 (3.5-8.4)	15% (11-19%)	22% (16-28%)	490 (340-650)	23000 (16000-30000)
Peru	2000	Male	50 to 54	98% (97-99%)	1.8% (1.2-2.6%)	5.5 (3.3-8.3)	15% (10-19%)	21% (14-26%)	530 (360-700)	21000 (14000-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	2000	Male	55 to 59	99% (98-99%)	1.5% (0.96-2.2%)	5.6 (3.6-8.2)	14% (9.6-18%)	20% (14-25%)	570 (390-750)	20000 (14000-26000)
Peru	2000	Male	60 to 64	98% (98-99%)	1.6% (0.97-2.4%)	4.9 (3-7.4)	12% (8-17%)	16% (11-22%)	610 (390-840)	18000 (12000-24000)
Peru	2000	Male	65 to 69	98% (97-99%)	1.9% (1.1-3%)	4.3 (2.7-6.3)	10% (6.4-14%)	13% (8.2-19%)	590 (360-840)	14000 (8700-20000)
Peru	2000	Male	70 to 74	98% (96-98%)	2.4% (1.5-3.9%)	3.6 (2.1-5.5)	7.5% (4-12%)	9.5% (5.1-15%)	470 (250-750)	8900 (4900-14000)
Peru	2000	Male	75 to 79	96% (93-98%)	3.8% (2.1-6.6%)	2.8 (1.6-4.5)	5.4% (2.3-9.6%)	6.6% (2.8-12%)	330 (140-600)	4900 (2100-8900)
Peru	2000	Male	80 to 84	95% (91-97%)	5.4% (3.1-9.5%)	2.3 (1.4-3.6)	4% (1.5-7.3%)	4.7% (1.7-8.9%)	230 (80-420)	2500 (950-4600)
Peru	2000	Male	85 to 89	94% (89-96%)	6.5% (3.5-11%)	2.1 (1.2-3.3)	3% (0.68-5.9%)	3.3% (0.6-6.9%)	130 (23-260)	990 (230-2000)
Peru	2000	Male	90 to 94	93% (89-97%)	6.6% (3.3-11%)	2.1 (1.1-3.4)	2.1% (-0.17-5.1%)	2.2% (-0.55-5.9%)	44 (-11-120)	250 (-21-630)
Peru	2000	Male	95 plus	94% (88-97%)	6.3% (2.7-12%)	2.1 (1-3.6)	1.7% (-0.8-5.5%)	1.6% (-1.5-6.1%)	9.4 (-8.8-36)	40 (-19-130)
Peru	2005	Female	All Ages	69% (65-71%)	31% (29-35%)	1.5 (1.3-1.8)	2.5% (1.8-3.2%)	3.8% (2.3-5.6%)	1900 (1200-2900)	72000 (53000-95000)
Peru	2005	Female	15 to 19	58% (51-64%)	42% (36-49%)	1.1 (0.68-1.6)	1.9% (1.1-2.8%)	3.7% (1.7-6.1%)	26 (12-43)	3000 (1700-4500)
Peru	2005	Female	20 to 24	83% (80-87%)	17% (13-20%)	2.1 (1.3-3)	3.7% (2.4-5.2%)	6.9% (3.6-10%)	57 (29-85)	5900 (3800-8400)
Peru	2005	Female	25 to 29	86% (82-88%)	14% (12-18%)	2.4 (1.6-3.3)	3.8% (2.5-5.2%)	7.5% (4.4-11%)	65 (37-95)	6000 (3900-8300)
Peru	2005	Female	30 to 34	87% (84-90%)	13% (10-16%)	2.6 (1.7-3.5)	3.7% (2.5-5.1%)	7.7% (4.8-11%)	71 (44-100)	5700 (3800-7900)
Peru	2005	Female	35 to 39	87% (84-90%)	13% (10-16%)	2.4 (1.6-3.4)	3.7% (2.4-5.3%)	7.9% (5.2-11%)	89 (56-130)	5800 (3800-8300)
Peru	2005	Female	40 to 44	87% (84-90%)	13% (9.8-16%)	2.4 (1.6-3.5)	3.9% (2.5-5.7%)	8.5% (5.5-12%)	110 (73-160)	6200 (3900-8800)
Peru	2005	Female	45 to 49	89% (86-91%)	11% (8.7-14%)	2.6 (1.7-4)	4.4% (2.5-6.7%)	8.9% (5.8-13%)	140 (91-220)	6700 (4000-10000)
Peru	2005	Female	50 to 54	88% (85-91%)	12% (9.3-15%)	2.6 (1.6-4)	4.9% (2.7-7.6%)	9.1% (5.7-14%)	180 (110-270)	7200 (4200-11000)
Peru	2005	Female	55 to 59	85% (80-89%)	15% (11-20%)	2.4 (1.5-3.7)	4.8% (2.8-7.4%)	8.3% (5.2-12%)	190 (120-280)	6300 (3800-9600)
Peru	2005	Female	60 to 64	83% (78-87%)	17% (13-22%)	2.3 (1.4-3.7)	4.6% (2.4-7.5%)	7.4% (4.1-12%)	200 (110-320)	5800 (3100-9400)
Peru	2005	Female	65 to 69	81% (74-86%)	19% (14-26%)	2.3 (1.4-3.4)	4.1% (2.1-6.7%)	6.5% (3.5-9.9%)	220 (120-360)	5200 (2600-8500)
Peru	2005	Female	70 to 74	78% (69-84%)	22% (16-31%)	2 (1.1-3.1)	3.2% (1.1-5.8%)	4.8% (1.9-8.3%)	210 (83-370)	3900 (1300-7100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	2005	Female	75 to 79	74% (64-83%)	26% (17-36%)	1.6 (0.87-2.7)	2.3% (0.41-4.7%)	3.4% (0.84-6.6%)	190 (45-380)	2700 (480-5700)
Peru	2005	Female	80 to 84	68% (56-77%)	32% (23-44%)	1.4 (0.77-2.2)	1.5% (-0.073-3.5%)	2.1% (-0.07-4.6%)	110 (-3.7-250)	1200 (-56-2700)
Peru	2005	Female	85 to 89	62% (49-73%)	38% (27-51%)	1.2 (0.67-2)	0.97% (-0.43-2.8%)	1.2% (-0.62-3.6%)	57 (-29-170)	440 (-200-1300)
Peru	2005	Female	90 to 94	60% (47-71%)	40% (29-53%)	1.2 (0.62-1.9)	0.49% (-1-2.3%)	0.52% (-1.4-2.9%)	16 (-43-89)	98 (-190-470)
Peru	2005	Female	95 plus	60% (44-72%)	40% (28-56%)	1.2 (0.55-2)	0.24% (-1.5-2.6%)	0.13% (-2-3%)	1.5 (-24-37)	12 (-73-130)
Peru	2005	Male	All Ages	80% (78-83%)	20% (17-22%)	2.9 (2.5-3.3)	6.7% (5.4-7.9%)	8.8% (6.6-11%)	5200 (3800-6800)	220000 (180000-270000)
Peru	2005	Male	15 to 19	67% (61-73%)	33% (27-39%)	1.7 (1-2.5)	4.9% (3.4-6.8%)	7.6% (4.5-11%)	79 (46-120)	8400 (5600-12000)
Peru	2005	Male	20 to 24	94% (92-96%)	5.8% (4.5-7.8%)	3.6 (2.3-5)	11% (8.2-14%)	15% (10-21%)	220 (140-310)	21000 (15000-28000)
Peru	2005	Male	25 to 29	96% (95-97%)	3.9% (3.1-5%)	4.5 (3-6.2)	13% (9.4-16%)	18% (12-24%)	280 (190-380)	23000 (17000-30000)
Peru	2005	Male	30 to 34	97% (97-98%)	2.6% (1.9-3.5%)	5.1 (3.6-6.7)	12% (9.3-16%)	18% (13-24%)	290 (200-390)	21000 (16000-27000)
Peru	2005	Male	35 to 39	98% (97-99%)	2% (1.4-2.7%)	5.3 (3.6-7.4)	13% (9.2-16%)	20% (14-25%)	330 (230-450)	21000 (15000-28000)
Peru	2005	Male	40 to 44	98% (98-99%)	1.7% (1.2-2.4%)	5.5 (3.6-7.7)	13% (9.2-16%)	20% (14-26%)	380 (260-500)	21000 (15000-27000)
Peru	2005	Male	45 to 49	98% (97-99%)	1.9% (1.3-2.7%)	5.4 (3.4-8)	13% (9.1-17%)	20% (14-26%)	430 (290-580)	20000 (14000-27000)
Peru	2005	Male	50 to 54	98% (97-99%)	1.9% (1.2-2.7%)	5.4 (3.2-7.9)	13% (8.8-17%)	19% (13-25%)	480 (320-650)	20000 (13000-26000)
Peru	2005	Male	55 to 59	98% (98-99%)	1.6% (1-2.3%)	5.4 (3.5-8)	13% (8.6-17%)	18% (12-24%)	520 (340-700)	18000 (12000-24000)
Peru	2005	Male	60 to 64	98% (97-99%)	1.7% (1.1-2.7%)	4.7 (2.9-7.2)	11% (7-15%)	15% (9.7-21%)	530 (330-740)	16000 (10000-22000)
Peru	2005	Male	65 to 69	98% (97-99%)	2% (1.2-3.3%)	4.1 (2.6-6.1)	8.9% (5.4-13%)	12% (7.3-17%)	520 (320-760)	13000 (7900-19000)
Peru	2005	Male	70 to 74	97% (96-98%)	2.6% (1.6-4.1%)	3.4 (2-5.2)	6.6% (3.5-11%)	8.6% (4.6-14%)	440 (240-710)	8700 (4700-14000)
Peru	2005	Male	75 to 79	96% (93-98%)	4.1% (2.2-7.2%)	2.7 (1.5-4.3)	4.7% (1.9-8.6%)	5.8% (2.3-11%)	320 (130-600)	4900 (2000-8900)
Peru	2005	Male	80 to 84	94% (90-97%)	5.9% (3.3-10%)	2.2 (1.3-3.4)	3.5% (1.3-6.4%)	4.2% (1.4-7.8%)	220 (77-430)	2500 (900-4700)
Peru	2005	Male	85 to 89	93% (88-96%)	7% (3.8-12%)	2 (1.1-3.2)	2.7% (0.69-5.4%)	3.1% (0.64-6.3%)	140 (27-280)	1100 (280-2200)
Peru	2005	Male	90 to 94	93% (88-96%)	7.1% (3.6-12%)	2 (1.1-3.2)	2% (-0.2-4.8%)	2.1% (-0.49-5.5%)	51 (-12-130)	300 (-30-710)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	2005	Male	95 plus	93% (87-97%)	6.8% (2.8-13%)	2 (0.97-3.4)	1.7% (-0.73-5.4%)	1.6% (-1.4-6.1%)	12 (-11-46)	52 (-23-170)
Peru	2010	Female	All Ages	69% (66-71%)	31% (29-34%)	1.4 (1.2-1.7)	2.3% (1.7-3.1%)	3.4% (2-4.9%)	1900 (1100-2800)	71000 (50000-94000)
Peru	2010	Female	15 to 19	56% (52-60%)	44% (40-48%)	0.99 (0.62-1.5)	1.6% (0.91-2.4%)	3.1% (1.3-5.2%)	21 (8.9-35)	2500 (1500-3800)
Peru	2010	Female	20 to 24	83% (81-86%)	17% (14-19%)	2 (1.3-2.9)	3.3% (2.1-4.7%)	5.9% (2.9-9.2%)	46 (22-72)	5300 (3300-7700)
Peru	2010	Female	25 to 29	86% (84-88%)	14% (12-16%)	2.3 (1.5-3.2)	3.4% (2.2-4.8%)	6.7% (3.7-10%)	54 (29-82)	5400 (3300-7700)
Peru	2010	Female	30 to 34	87% (85-89%)	13% (11-15%)	2.4 (1.7-3.4)	3.4% (2.2-4.8%)	7.1% (4.3-10%)	66 (39-96)	5500 (3500-7900)
Peru	2010	Female	35 to 39	87% (85-89%)	13% (11-15%)	2.3 (1.5-3.2)	3.3% (2.1-4.8%)	7.1% (4.4-10%)	80 (49-120)	5500 (3500-7800)
Peru	2010	Female	40 to 44	88% (86-90%)	12% (10-14%)	2.3 (1.5-3.3)	3.6% (2.3-5.3%)	7.7% (4.9-11%)	110 (69-160)	6100 (3800-9000)
Peru	2010	Female	45 to 49	89% (87-90%)	11% (9.5-13%)	2.4 (1.5-3.7)	3.8% (2.2-5.9%)	7.9% (5-11%)	140 (86-210)	6500 (3800-10000)
Peru	2010	Female	50 to 54	87% (85-89%)	13% (11-15%)	2.4 (1.5-3.6)	4.2% (2.4-6.7%)	8% (5-12%)	180 (110-280)	7200 (4200-11000)
Peru	2010	Female	55 to 59	85% (83-88%)	15% (12-17%)	2.3 (1.4-3.3)	4.4% (2.4-6.9%)	7.7% (4.6-11%)	210 (120-310)	7100 (4000-11000)
Peru	2010	Female	60 to 64	81% (79-84%)	19% (16-21%)	2.1 (1.3-3.3)	4.1% (2-6.9%)	6.6% (3.6-11%)	210 (120-350)	6200 (3100-10000)
Peru	2010	Female	65 to 69	77% (73-81%)	23% (19-27%)	2 (1.3-2.9)	3.6% (1.9-5.9%)	5.6% (3.1-8.9%)	220 (120-340)	5000 (2600-8200)
Peru	2010	Female	70 to 74	71% (67-76%)	29% (24-33%)	1.6 (0.87-2.4)	2.7% (1.1-4.8%)	4.2% (1.8-7%)	200 (86-350)	3700 (1400-6700)
Peru	2010	Female	75 to 79	68% (61-74%)	32% (26-39%)	1.3 (0.7-2)	2% (0.5-3.8%)	2.9% (0.88-5.3%)	170 (51-310)	2400 (620-4600)
Peru	2010	Female	80 to 84	61% (51-70%)	39% (30-49%)	1.1 (0.6-1.6)	1.3% (0.12-2.8%)	1.8% (0.22-3.7%)	120 (14-260)	1300 (120-2900)
Peru	2010	Female	85 to 89	53% (41-64%)	47% (36-59%)	0.9 (0.5-1.5)	0.82% (-0.21-2.1%)	1% (-0.31-2.6%)	63 (-19-160)	490 (-130-1200)
Peru	2010	Female	90 to 94	51% (39-61%)	49% (39-61%)	0.87 (0.46-1.4)	0.43% (-0.63-1.8%)	0.46% (-0.93-2.1%)	18 (-37-84)	110 (-160-450)
Peru	2010	Female	95 plus	51% (37-64%)	49% (36-63%)	0.86 (0.4-1.5)	0.21% (-1-1.8%)	0.11% (-1.5-2.1%)	1.7 (-23-33)	14 (-67-120)
Peru	2010	Male	All Ages	82% (80-84%)	18% (16-20%)	2.9 (2.4-3.4)	6.7% (5.4-8.2%)	8.4% (6.2-11%)	5500 (4000-7500)	230000 (180000-290000)
Peru	2010	Male	15 to 19	68% (64-72%)	32% (28-36%)	1.6 (0.96-2.4)	4.5% (3.1-6.3%)	6.9% (4.1-10%)	71 (42-110)	7700 (5200-11000)
Peru	2010	Male	20 to 24	94% (93-95%)	5.8% (4.6-7.4%)	3.4 (2.2-4.9)	10% (7.4-14%)	14% (9.2-20%)	210 (130-310)	21000 (14000-27000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	2010	Male	25 to 29	96% (95-97%)	3.8% (3.1-4.6%)	4.3 (2.8-5.9)	12% (8.8-16%)	18% (12-24%)	290 (190-400)	24000 (17000-31000)
Peru	2010	Male	30 to 34	98% (97-98%)	2.4% (1.8-3.2%)	4.9 (3.5-6.6)	12% (8.6-15%)	18% (13-25%)	310 (210-420)	23000 (16000-30000)
Peru	2010	Male	35 to 39	98% (97-99%)	1.9% (1.4-2.6%)	5.1 (3.4-7.1)	12% (8.4-16%)	19% (13-26%)	340 (230-480)	22000 (15000-30000)
Peru	2010	Male	40 to 44	98% (98-99%)	1.8% (1.2-2.4%)	5.2 (3.5-7.5)	12% (8.7-16%)	19% (13-25%)	390 (270-540)	22000 (15000-29000)
Peru	2010	Male	45 to 49	98% (98-99%)	1.8% (1.3-2.5%)	5.2 (3.3-7.6)	12% (8.5-16%)	19% (13-26%)	460 (310-630)	22000 (15000-30000)
Peru	2010	Male	50 to 54	98% (98-99%)	1.7% (1.2-2.3%)	5.3 (3.3-7.9)	12% (8.2-16%)	18% (12-24%)	510 (340-700)	21000 (14000-29000)
Peru	2010	Male	55 to 59	98% (98-99%)	1.6% (1-2.2%)	5.2 (3.4-7.6)	12% (7.5-16%)	17% (11-23%)	560 (350-780)	20000 (13000-27000)
Peru	2010	Male	60 to 64	99% (98-99%)	1.5% (0.96-2.2%)	4.8 (2.9-7.2)	10% (6.3-15%)	14% (8.7-20%)	580 (350-840)	17000 (10000-24000)
Peru	2010	Male	65 to 69	98% (97-99%)	1.8% (1.2-2.6%)	4.1 (2.6-5.9)	8.3% (4.9-13%)	11% (6.6-17%)	530 (300-790)	13000 (7600-19000)
Peru	2010	Male	70 to 74	97% (96-98%)	2.6% (1.8-3.6%)	3.3 (2-4.9)	6.1% (3.1-9.9%)	7.8% (4.1-13%)	460 (230-750)	9000 (4700-15000)
Peru	2010	Male	75 to 79	96% (94-97%)	4.1% (2.5-6.2%)	2.6 (1.5-4)	4.3% (1.7-7.8%)	5.3% (2-9.8%)	360 (130-660)	5400 (2100-9900)
Peru	2010	Male	80 to 84	94% (90-96%)	6.3% (3.8-10%)	2 (1.2-3.1)	3.1% (1-5.8%)	3.7% (1.1-7%)	240 (74-460)	2700 (920-5000)
Peru	2010	Male	85 to 89	92% (87-96%)	7.6% (4.4-13%)	1.8 (1.1-2.9)	2.4% (0.56-4.8%)	2.7% (0.45-5.6%)	150 (26-330)	1200 (290-2500)
Peru	2010	Male	90 to 94	92% (87-96%)	7.7% (4.3-13%)	1.8 (0.98-2.9)	1.7% (-0.2-4.3%)	1.8% (-0.61-5%)	57 (-20-160)	330 (-40-850)
Peru	2010	Male	95 plus	93% (86-97%)	7.3% (3.2-14%)	1.8 (0.88-3.2)	1.4% (-0.85-4.6%)	1.2% (-1.5-5.1%)	13 (-14-51)	56 (-35-190)
Peru	2016	Female	All Ages	69% (67-72%)	31% (28-33%)	1.4 (1.1-1.9)	2.1% (1.3-3.1%)	3% (1.5-4.7%)	1800 (890-3000)	66000 (40000-99000)
Peru	2016	Female	15 to 19	56% (52-59%)	44% (41-48%)	0.96 (0.56-1.5)	1.5% (0.85-2.3%)	3% (1.3-5.3%)	15 (6.2-28)	2000 (1200-3200)
Peru	2016	Female	20 to 24	84% (81-86%)	16% (14-19%)	2 (1.2-3)	3% (1.9-4.3%)	5.5% (2.6-9.1%)	33 (15-58)	4500 (2700-6700)
Peru	2016	Female	25 to 29	86% (84-88%)	14% (12-16%)	2.2 (1.4-3.3)	3.1% (2-4.4%)	6.3% (3.3-9.8%)	41 (21-69)	4900 (3100-7200)
Peru	2016	Female	30 to 34	87% (85-89%)	13% (11-15%)	2.4 (1.5-3.4)	3% (1.9-4.3%)	6.7% (3.8-10%)	50 (29-80)	4800 (3000-7100)
Peru	2016	Female	35 to 39	87% (85-89%)	13% (11-15%)	2.3 (1.4-3.4)	2.9% (1.8-4.4%)	6.9% (4.2-10%)	65 (36-110)	4900 (3000-7700)
Peru	2016	Female	40 to 44	88% (86-90%)	12% (10-14%)	2.3 (1.4-3.5)	3.1% (1.8-4.8%)	7.4% (4.7-11%)	90 (51-150)	5300 (3000-8700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	2016	Female	45 to 49	89% (87-90%)	11% (9.6-13%)	2.4 (1.5-3.6)	3.3% (1.7-5.5%)	7.5% (4.6-11%)	120 (67-200)	6000 (3200-10000)
Peru	2016	Female	50 to 54	87% (85-89%)	13% (11-15%)	2.3 (1.4-3.4)	3.6% (1.8-6%)	7.5% (4.5-11%)	170 (90-280)	6800 (3300-11000)
Peru	2016	Female	55 to 59	85% (83-88%)	15% (12-17%)	2.2 (1.4-3.4)	3.8% (2-6.2%)	7.2% (4.2-11%)	200 (100-330)	6900 (3300-12000)
Peru	2016	Female	60 to 64	81% (78-84%)	19% (16-22%)	2 (1.2-3.3)	3.5% (1.6-6.4%)	6.1% (3.1-10%)	220 (100-390)	6400 (2700-11000)
Peru	2016	Female	65 to 69	76% (72-80%)	24% (20-28%)	1.9 (1.1-2.9)	3.1% (1.3-5.3%)	5.2% (2.6-8.3%)	220 (100-380)	5100 (2100-9000)
Peru	2016	Female	70 to 74	71% (66-75%)	29% (25-34%)	1.5 (0.81-2.3)	2.4% (0.85-4.5%)	3.9% (1.7-6.8%)	200 (76-370)	3600 (1200-7000)
Peru	2016	Female	75 to 79	67% (60-74%)	33% (26-40%)	1.2 (0.63-2)	1.8% (0.35-3.6%)	2.7% (0.73-5.2%)	170 (44-330)	2300 (460-5000)
Peru	2016	Female	80 to 84	60% (50-70%)	40% (30-50%)	1 (0.57-1.6)	1.2% (0.0016-2.7%)	1.7% (0.11-3.6%)	120 (8-270)	1300 (1.7-2900)
Peru	2016	Female	85 to 89	52% (41-62%)	48% (38-59%)	0.85 (0.46-1.4)	0.73% (-0.32-2%)	0.95% (-0.41-2.6%)	79 (-34-220)	610 (-240-1700)
Peru	2016	Female	90 to 94	50% (38-61%)	50% (39-62%)	0.82 (0.42-1.4)	0.36% (-0.68-1.8%)	0.38% (-0.95-2.1%)	24 (-61-140)	150 (-300-750)
Peru	2016	Female	95 plus	50% (35-63%)	50% (37-65%)	0.82 (0.36-1.5)	0.15% (-1-1.9%)	0.035% (-1.5-2.2%)	0.87 (-34-51)	15 (-100-190)
Peru	2016	Male	All Ages	83% (81-85%)	17% (15-19%)	2.8 (2.2-3.7)	6.4% (4.7-8.3%)	7.6% (5-11%)	5200 (3100-7900)	220000 (150000-300000)
Peru	2016	Male	15 to 19	67% (64-71%)	33% (29-36%)	1.5 (0.87-2.5)	4.2% (2.8-5.9%)	6.6% (3.8-11%)	53 (28-88)	6400 (4100-9700)
Peru	2016	Male	20 to 24	94% (93-95%)	5.8% (4.6-7.3%)	3.3 (2-5)	9.8% (7-13%)	14% (8.6-20%)	170 (97-260)	18000 (12000-25000)
Peru	2016	Male	25 to 29	96% (95-97%)	3.8% (3.1-4.6%)	4.2 (2.6-6.2)	11% (8-15%)	17% (10-24%)	230 (140-360)	21000 (14000-30000)
Peru	2016	Male	30 to 34	98% (97-98%)	2.4% (1.9-3.2%)	4.7 (3.2-6.8)	11% (7.9-15%)	18% (12-25%)	260 (150-400)	20000 (14000-29000)
Peru	2016	Male	35 to 39	98% (97-99%)	1.9% (1.4-2.5%)	4.9 (3.1-7.4)	11% (7.4-15%)	18% (12-25%)	300 (170-460)	20000 (13000-29000)
Peru	2016	Male	40 to 44	98% (98-99%)	1.8% (1.2-2.4%)	5 (3.2-7.6)	11% (7.3-15%)	18% (12-25%)	340 (200-530)	20000 (13000-29000)
Peru	2016	Male	45 to 49	98% (98-99%)	1.8% (1.3-2.5%)	4.9 (3.1-7.5)	11% (7.1-15%)	18% (12-25%)	410 (240-630)	21000 (13000-30000)
Peru	2016	Male	50 to 54	98% (98-99%)	1.7% (1.2-2.3%)	5 (3-7.6)	11% (6.9-15%)	17% (11-23%)	500 (290-780)	21000 (13000-32000)
Peru	2016	Male	55 to 59	98% (98-99%)	1.6% (1-2.2%)	5 (3.1-7.6)	11% (6.7-15%)	16% (9.9-22%)	550 (310-850)	20000 (12000-30000)
Peru	2016	Male	60 to 64	99% (98-99%)	1.5% (0.95-2.2%)	4.6 (2.8-7.3)	9.6% (5.4-14%)	13% (7.6-20%)	610 (320-970)	18000 (10000-29000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Peru	2016	Male	65 to 69	98% (97-99%)	1.8% (1.2-2.7%)	3.9 (2.3-5.9)	7.6% (4.2-12%)	10% (5.6-16%)	550 (280-920)	14000 (7100-23000)
Peru	2016	Male	70 to 74	97% (96-98%)	2.6% (1.8-3.7%)	3.1 (1.8-4.8)	5.4% (2.5-9.3%)	7.1% (3.3-12%)	430 (180-800)	8700 (3800-16000)
Peru	2016	Male	75 to 79	96% (94-97%)	4.1% (2.6-6.2%)	2.4 (1.3-4)	3.8% (1.4-7.3%)	4.8% (1.7-9.2%)	340 (110-680)	5200 (1800-10000)
Peru	2016	Male	80 to 84	93% (89-96%)	6.6% (4-11%)	1.9 (1.1-3.1)	2.8% (0.81-5.5%)	3.3% (0.86-6.6%)	240 (57-500)	2800 (800-5500)
Peru	2016	Male	85 to 89	92% (87-96%)	7.8% (4.5-13%)	1.7 (0.98-2.9)	2.1% (0.36-4.6%)	2.4% (0.25-5.4%)	170 (17-380)	1400 (240-3000)
Peru	2016	Male	90 to 94	92% (87-96%)	7.8% (4.3-13%)	1.7 (0.91-2.9)	1.6% (-0.37-4.3%)	1.6% (-0.76-5.1%)	70 (-33-210)	420 (-100-1200)
Peru	2016	Male	95 plus	93% (87-97%)	7.4% (3.2-13%)	1.7 (0.81-3.1)	1.3% (-0.83-4.5%)	1.1% (-1.5-5.1%)	16 (-22-74)	76 (-48-280)
Colombia	1990	Female	All Ages	46% (40-52%)	54% (48-60%)	1 (0.81-1.2)	1.9% (1.1-2.7%)	2.5% (0.74-4.3%)	1800 (530-3100)	77000 (47000-110000)
Colombia	1990	Female	15 to 19	44% (31-57%)	56% (43-69%)	1.1 (0.63-1.7)	2.6% (1.5-4%)	4.7% (2.2-7.6%)	54 (26-88)	5600 (3200-8600)
Colombia	1990	Female	20 to 24	59% (46-71%)	41% (29-54%)	1.6 (0.94-2.6)	4.1% (2.5-5.9%)	6.4% (3.2-10%)	92 (45-150)	9600 (5600-14000)
Colombia	1990	Female	25 to 29	60% (46-71%)	40% (29-54%)	1.5 (0.93-2.2)	3.7% (2.3-5.2%)	5.8% (2.9-9%)	84 (41-130)	8100 (5000-12000)
Colombia	1990	Female	30 to 34	60% (47-71%)	40% (29-53%)	1.5 (0.91-2.1)	3.2% (1.9-4.6%)	5.3% (2.8-8%)	75 (40-120)	6400 (3800-9200)
Colombia	1990	Female	35 to 39	60% (48-71%)	40% (29-52%)	1.5 (0.98-2.2)	3% (1.8-4.4%)	5.3% (2.9-8%)	87 (46-130)	5900 (3400-8700)
Colombia	1990	Female	40 to 44	61% (49-71%)	39% (29-51%)	1.7 (1.1-2.4)	3.1% (1.7-4.7%)	5.6% (2.9-8.5%)	95 (50-140)	5400 (3000-8200)
Colombia	1990	Female	45 to 49	61% (48-72%)	39% (28-52%)	1.9 (1.1-2.9)	3.7% (1.8-6.2%)	6.2% (3-10%)	130 (60-210)	5900 (2900-9900)
Colombia	1990	Female	50 to 54	61% (47-73%)	39% (27-53%)	2 (1.2-3.1)	4.1% (1.7-7%)	6.3% (2.7-11%)	170 (75-290)	6700 (2800-11000)
Colombia	1990	Female	55 to 59	60% (45-74%)	40% (26-55%)	2.1 (1.2-3.3)	4.2% (1.5-7.6%)	6.1% (2.3-11%)	210 (77-370)	7000 (2500-13000)
Colombia	1990	Female	60 to 64	57% (41-72%)	43% (28-59%)	1.9 (1-3.1)	3.5% (0.66-7.2%)	4.8% (1-9.7%)	220 (44-440)	6200 (1100-13000)
Colombia	1990	Female	65 to 69	53% (37-68%)	47% (32-63%)	1.6 (0.78-2.5)	2.7% (0.1-5.5%)	3.6% (0.27-7.4%)	190 (14-380)	4400 (180-9200)
Colombia	1990	Female	70 to 74	49% (34-64%)	51% (36-66%)	1.3 (0.58-2.3)	1.9% (-0.54-4.7%)	2.4% (-0.65-6.1%)	150 (-41-370)	2800 (-860-7100)
Colombia	1990	Female	75 to 79	43% (28-60%)	57% (40-72%)	1.1 (0.48-2)	1.1% (-0.96-3.6%)	1.4% (-1.3-4.6%)	92 (-82-290)	1400 (-1200-4500)
Colombia	1990	Female	80 to 84	39% (24-55%)	61% (45-76%)	1 (0.48-1.7)	0.88% (-0.98-3%)	1.1% (-1.2-3.7%)	66 (-77-230)	740 (-800-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	1990	Female	85 to 89	39% (25-54%)	61% (46-75%)	1 (0.49-1.7)	0.73% (-1.1-2.8%)	0.81% (-1.5-3.3%)	44 (-81-180)	360 (-570-1400)
Colombia	1990	Female	90 to 94	42% (28-57%)	58% (43-72%)	1.1 (0.56-2)	0.47% (-1.4-2.8%)	0.42% (-1.9-3.3%)	13 (-57-100)	87 (-270-520)
Colombia	1990	Female	95 plus	46% (29-65%)	54% (35-71%)	1.3 (0.54-2.3)	0.36% (-1.9-3.6%)	0.2% (-2.7-4.1%)	2.1 (-29-44)	16 (-86-160)
Colombia	1990	Male	All Ages	86% (82-90%)	14% (10-18%)	2.8 (2.3-3.2)	9.7% (7.5-12%)	11% (8.5-14%)	12000 (8800-15000)	570000 (440000-710000)
Colombia	1990	Male	15 to 19	84% (73-91%)	16% (8.5-27%)	2.5 (1.5-3.7)	11% (6.5-16%)	14% (7.8-21%)	600 (320-900)	46000 (26000-68000)
Colombia	1990	Male	20 to 24	93% (88-96%)	6.9% (4-12%)	4.3 (2.9-6.1)	19% (12-26%)	22% (13-30%)	1600 (940-2200)	110000 (71000-150000)
Colombia	1990	Male	25 to 29	95% (91-97%)	5.1% (3-8.5%)	4.7 (3.3-6.5)	20% (13-27%)	23% (14-32%)	1600 (980-2200)	100000 (68000-140000)
Colombia	1990	Male	30 to 34	95% (92-97%)	4.7% (2.7-8%)	4.8 (3.3-6.6)	18% (13-25%)	22% (15-31%)	1200 (800-1700)	75000 (51000-100000)
Colombia	1990	Male	35 to 39	95% (92-97%)	5.3% (3.1-8.2%)	4.9 (3.2-7.2)	17% (12-24%)	21% (14-30%)	990 (650-1400)	56000 (37000-77000)
Colombia	1990	Male	40 to 44	95% (91-97%)	5.4% (3.3-8.5%)	5.2 (3.4-7.3)	16% (11-21%)	21% (14-29%)	750 (500-1100)	38000 (26000-52000)
Colombia	1990	Male	45 to 49	95% (92-97%)	4.8% (2.6-8.4%)	5.6 (3.5-8.2)	15% (10-21%)	20% (13-27%)	670 (450-940)	30000 (20000-41000)
Colombia	1990	Male	50 to 54	94% (91-97%)	5.7% (3.3-9.3%)	5.8 (3.7-8.5)	14% (9.3-19%)	18% (12-24%)	710 (460-940)	28000 (18000-37000)
Colombia	1990	Male	55 to 59	92% (87-95%)	8% (4.6-13%)	6.1 (3.8-9)	14% (9.4-18%)	17% (12-23%)	790 (530-1000)	27000 (18000-35000)
Colombia	1990	Male	60 to 64	90% (84-94%)	10% (5.8-16%)	5.4 (3.2-8.2)	12% (7.2-16%)	14% (8.5-19%)	840 (500-1200)	24000 (15000-33000)
Colombia	1990	Male	65 to 69	88% (82-93%)	12% (6.7-18%)	4.1 (2.5-6.1)	9% (4.5-13%)	11% (5.4-16%)	690 (340-1000)	17000 (8700-24000)
Colombia	1990	Male	70 to 74	86% (79-92%)	14% (8.3-21%)	2.9 (1.7-4.7)	5.7% (2-10%)	6.6% (2.2-12%)	470 (160-840)	9100 (3200-16000)
Colombia	1990	Male	75 to 79	84% (74-92%)	16% (8.5-26%)	2.3 (1.2-3.8)	3.8% (0.87-8%)	4.3% (0.8-9.3%)	300 (55-660)	4600 (1100-9800)
Colombia	1990	Male	80 to 84	81% (69-90%)	19% (10-31%)	2 (1.1-3.2)	3.1% (0.67-6.5%)	3.4% (0.56-7.4%)	200 (34-430)	2300 (520-4800)
Colombia	1990	Male	85 to 89	79% (67-88%)	21% (12-33%)	1.9 (1-3)	2.6% (0.47-5.7%)	2.8% (0.24-6.4%)	120 (10-270)	960 (170-2200)
Colombia	1990	Male	90 to 94	80% (68-89%)	20% (11-32%)	2 (1-3.3)	2.3% (-0.11-5.8%)	2.3% (-0.45-6.4%)	44 (-8.5-120)	260 (-12-660)
Colombia	1990	Male	95 plus	81% (67-91%)	19% (9-33%)	2.1 (1-3.8)	2.2% (-0.73-6.6%)	2% (-1.5-7.4%)	10 (-7.5-37)	44 (-15-130)
Colombia	1995	Female	All Ages	52% (47-57%)	48% (43-53%)	1.1 (0.88-1.3)	1.9% (1.1-2.8%)	2.4% (0.48-4.4%)	1800 (370-3400)	82000 (46000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	1995	Female	15 to 19	54% (43-64%)	46% (36-57%)	1.3 (0.79-1.9)	3.1% (1.7-4.7%)	5.8% (2.6-9.2%)	74 (33-120)	7200 (3800-11000)
Colombia	1995	Female	20 to 24	67% (57-75%)	33% (25-43%)	1.8 (1.1-2.7)	4.4% (2.6-6.3%)	7.3% (3.5-11%)	97 (46-150)	9800 (5600-14000)
Colombia	1995	Female	25 to 29	65% (56-74%)	35% (26-44%)	1.6 (1-2.2)	3.7% (2.2-5.3%)	6.3% (3.1-9.7%)	92 (45-140)	8700 (5200-13000)
Colombia	1995	Female	30 to 34	65% (55-73%)	35% (27-45%)	1.5 (0.99-2.2)	3.1% (1.8-4.5%)	5.6% (2.8-8.7%)	87 (45-140)	7100 (4300-11000)
Colombia	1995	Female	35 to 39	64% (55-73%)	36% (27-45%)	1.5 (1-2.2)	2.9% (1.7-4.3%)	5.4% (2.9-8.2%)	98 (52-150)	6500 (3600-9700)
Colombia	1995	Female	40 to 44	63% (54-72%)	37% (28-46%)	1.7 (1.1-2.3)	2.8% (1.5-4.3%)	5.5% (2.9-8.3%)	110 (55-160)	6000 (3300-9100)
Colombia	1995	Female	45 to 49	63% (52-73%)	37% (27-48%)	1.9 (1.2-2.8)	3.2% (1.5-5.5%)	6% (2.9-9.7%)	140 (65-220)	6400 (2900-11000)
Colombia	1995	Female	50 to 54	63% (51-74%)	37% (26-49%)	2 (1.2-3)	3.6% (1.4-6.4%)	5.9% (2.5-10%)	170 (73-290)	6600 (2600-12000)
Colombia	1995	Female	55 to 59	64% (51-75%)	36% (25-49%)	2.1 (1.2-3.2)	3.9% (1.2-7.1%)	5.8% (2.1-10%)	210 (74-370)	7000 (2300-13000)
Colombia	1995	Female	60 to 64	62% (48-74%)	38% (26-52%)	1.9 (1.1-3.1)	3.3% (0.49-7%)	4.5% (0.75-9.4%)	230 (38-460)	6400 (950-14000)
Colombia	1995	Female	65 to 69	57% (43-70%)	43% (30-57%)	1.6 (0.85-2.5)	2.4% (-0.14-5.4%)	3.2% (-0.092-7.1%)	190 (-5.6-430)	4500 (-250-10000)
Colombia	1995	Female	70 to 74	57% (44-69%)	43% (31-56%)	1.5 (0.7-2.5)	1.6% (-0.95-4.8%)	2.2% (-1.1-6.1%)	150 (-74-420)	2800 (-1700-8100)
Colombia	1995	Female	75 to 79	52% (37-67%)	48% (33-63%)	1.3 (0.58-2.3)	0.92% (-1.4-3.8%)	1.2% (-1.8-4.9%)	89 (-130-360)	1300 (-2000-5300)
Colombia	1995	Female	80 to 84	43% (28-60%)	57% (40-72%)	1.1 (0.51-1.8)	0.7% (-1.2-2.9%)	0.86% (-1.4-3.5%)	60 (-100-250)	660 (-1100-2700)
Colombia	1995	Female	85 to 89	41% (27-56%)	59% (44-73%)	1 (0.51-1.7)	0.54% (-1.2-2.6%)	0.6% (-1.6-3%)	40 (-110-200)	340 (-780-1600)
Colombia	1995	Female	90 to 94	45% (30-60%)	55% (40-70%)	1.1 (0.57-1.9)	0.3% (-1.6-2.6%)	0.23% (-2.1-3%)	8.8 (-81-120)	73 (-380-630)
Colombia	1995	Female	95 plus	49% (32-67%)	51% (33-68%)	1.2 (0.55-2.2)	0.11% (-2.1-3.2%)	-0.11% (-2.9-3.5%)	-1.6 (-40-49)	6.2 (-120-170)
Colombia	1995	Male	All Ages	88% (84-91%)	12% (9.2-16%)	2.8 (2.3-3.2)	10% (7.6-12%)	11% (8.2-14%)	12000 (9200-16000)	62000 (47000-78000)
Colombia	1995	Male	15 to 19	87% (80-93%)	13% (7.2-20%)	2.5 (1.6-3.7)	12% (6.7-18%)	15% (7.7-22%)	780 (400-1200)	60000 (32000-88000)
Colombia	1995	Male	20 to 24	94% (91-96%)	5.8% (3.7-9.4%)	4.2 (2.8-5.9)	19% (12-26%)	22% (13-30%)	1600 (950-2300)	120000 (73000-160000)
Colombia	1995	Male	25 to 29	95% (92-97%)	4.7% (2.8-7.9%)	4.4 (3.1-6.2)	19% (12-26%)	22% (14-30%)	1600 (970-2200)	100000 (69000-140000)
Colombia	1995	Male	30 to 34	96% (93-97%)	4.4% (2.6-7.1%)	4.6 (3.2-6.3)	17% (12-24%)	21% (14-29%)	1300 (830-1800)	80000 (54000-110000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	1995	Male	35 to 39	95% (92-97%)	5% (3.1-7.7%)	4.6 (3.1-6.6)	16% (11-22%)	20% (13-28%)	1100 (710-1500)	61000 (41000-84000)
Colombia	1995	Male	40 to 44	95% (92-97%)	5% (3.1-7.6%)	4.9 (3.3-6.9)	15% (10-20%)	19% (13-27%)	850 (540-1200)	44000 (29000-59000)
Colombia	1995	Male	45 to 49	96% (92-98%)	4.4% (2.4-7.8%)	5.3 (3.4-7.7)	14% (9.1-19%)	19% (12-26%)	760 (470-1000)	35000 (23000-47000)
Colombia	1995	Male	50 to 54	95% (92-97%)	5.2% (3.1-8.2%)	5.6 (3.6-8)	14% (8.8-19%)	18% (11-24%)	750 (470-1000)	30000 (20000-40000)
Colombia	1995	Male	55 to 59	93% (89-96%)	7.2% (4.3-11%)	5.9 (3.9-8.5)	14% (9.2-18%)	17% (11-22%)	830 (550-1100)	28000 (19000-38000)
Colombia	1995	Male	60 to 64	91% (87-94%)	9% (5.6-13%)	5.3 (3.2-8)	12% (6.9-16%)	14% (8.2-19%)	870 (520-1200)	25000 (15000-35000)
Colombia	1995	Male	65 to 69	90% (85-94%)	10% (6.1-15%)	4 (2.5-6)	8.7% (4.5-13%)	10% (5.2-15%)	760 (380-1100)	18000 (9500-27000)
Colombia	1995	Male	70 to 74	89% (83-93%)	11% (7.3-17%)	3 (1.7-4.6)	5.4% (1.9-9.7%)	6.2% (2-11%)	510 (160-930)	9900 (3500-18000)
Colombia	1995	Male	75 to 79	87% (80-93%)	13% (7.4-20%)	2.4 (1.3-3.9)	3.5% (0.54-7.4%)	3.9% (0.37-8.4%)	310 (31-670)	4800 (720-10000)
Colombia	1995	Male	80 to 84	83% (72-91%)	17% (9.2-28%)	2 (1.1-3.2)	2.8% (0.58-6%)	3.1% (0.39-6.8%)	210 (26-460)	2400 (490-5000)
Colombia	1995	Male	85 to 89	80% (67-88%)	20% (12-33%)	1.8 (1-3)	2.3% (0.36-5.1%)	2.5% (0.17-5.8%)	130 (8.7-300)	1100 (160-2300)
Colombia	1995	Male	90 to 94	81% (69-89%)	19% (11-31%)	2 (1-3.2)	2% (-0.25-5.1%)	2% (-0.7-5.8%)	48 (-17-140)	290 (-36-780)
Colombia	1995	Male	95 plus	82% (69-91%)	18% (8.8-31%)	2.1 (1-3.7)	1.9% (-0.76-6.2%)	1.8% (-1.6-6.9%)	12 (-11-46)	53 (-21-170)
Colombia	2000	Female	All Ages	53% (48-58%)	47% (42-52%)	1 (0.85-1.3)	1.9% (1.1-2.7%)	2.1% (0.34-4%)	1700 (280-3300)	81000 (47000-120000)
Colombia	2000	Female	15 to 19	55% (44-65%)	45% (35-56%)	1.2 (0.73-1.8)	3% (1.6-4.7%)	5.6% (2.4-9.3%)	80 (35-130)	7700 (4200-12000)
Colombia	2000	Female	20 to 24	67% (57-75%)	33% (25-43%)	1.6 (0.97-2.5)	4.2% (2.4-6.1%)	6.8% (3.1-11%)	98 (45-160)	10000 (5900-15000)
Colombia	2000	Female	25 to 29	66% (56-74%)	34% (26-44%)	1.5 (0.93-2.1)	3.5% (2.1-5%)	5.9% (2.7-9.1%)	80 (37-130)	8000 (4800-11000)
Colombia	2000	Female	30 to 34	65% (55-73%)	35% (27-45%)	1.4 (0.92-2)	2.9% (1.7-4.2%)	5.2% (2.7-8.1%)	83 (43-130)	7200 (4300-11000)
Colombia	2000	Female	35 to 39	64% (55-73%)	36% (27-45%)	1.4 (0.98-2)	2.6% (1.5-3.8%)	5.1% (2.7-7.7%)	93 (49-140)	6600 (3800-9500)
Colombia	2000	Female	40 to 44	64% (54-72%)	36% (28-46%)	1.5 (1-2.1)	2.6% (1.4-4%)	5.2% (2.7-7.8%)	110 (59-170)	6400 (3500-9600)
Colombia	2000	Female	45 to 49	64% (54-74%)	36% (26-46%)	1.7 (1.1-2.5)	2.9% (1.3-4.7%)	5.5% (2.8-8.9%)	140 (70-230)	6700 (3100-11000)
Colombia	2000	Female	50 to 54	63% (52-73%)	37% (27-48%)	1.8 (1.1-2.8)	3.1% (1.2-5.4%)	5.4% (2.4-8.9%)	160 (73-270)	6500 (2800-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	2000	Female	55 to 59	64% (52-75%)	36% (25-48%)	1.9 (1.1-3)	3.3% (1.1-6.2%)	5.4% (2-9.4%)	190 (70-340)	6500 (2100-12000)
Colombia	2000	Female	60 to 64	62% (48-74%)	38% (26-52%)	1.8 (1-2.9)	2.9% (0.32-5.9%)	4.1% (0.66-8.3%)	200 (33-410)	5800 (700-12000)
Colombia	2000	Female	65 to 69	57% (44-71%)	43% (29-56%)	1.5 (0.77-2.3)	2.1% (-0.15-4.8%)	2.9% (-0.025-6.3%)	180 (-1.6-400)	4300 (-320-9600)
Colombia	2000	Female	70 to 74	57% (44-69%)	43% (31-56%)	1.4 (0.63-2.3)	1.3% (-0.92-4.4%)	1.9% (-1.1-5.7%)	140 (-84-440)	2600 (-1700-8300)
Colombia	2000	Female	75 to 79	53% (38-67%)	47% (33-62%)	1.2 (0.56-2.1)	0.7% (-1.4-3.3%)	0.94% (-1.7-4.3%)	77 (-140-350)	1100 (-2100-5100)
Colombia	2000	Female	80 to 84	44% (28-60%)	56% (40-72%)	1 (0.49-1.7)	0.51% (-1.2-2.5%)	0.62% (-1.5-3.1%)	51 (-120-260)	570 (-1300-2800)
Colombia	2000	Female	85 to 89	41% (27-57%)	59% (43-73%)	0.97 (0.47-1.6)	0.36% (-1.2-2.2%)	0.37% (-1.7-2.6%)	29 (-130-200)	260 (-910-1500)
Colombia	2000	Female	90 to 94	45% (31-61%)	55% (39-69%)	1.1 (0.53-1.8)	0.11% (-1.6-2.2%)	-0.0062% (-2.1-2.4%)	-0.3 (-110-120)	35 (-500-660)
Colombia	2000	Female	95 plus	49% (32-67%)	51% (33-68%)	1.2 (0.53-2.1)	-0.066% (-2.1-2.7%)	-0.34% (-2.9-2.9%)	-6.2 (-54-54)	-5 (-160-200)
Colombia	2000	Male	All Ages	88% (85-91%)	12% (8.8-15%)	2.7 (2.2-3.1)	9.8% (7.4-12%)	10% (7.6-13%)	12000 (9000-16000)	620000 (470000-780000)
Colombia	2000	Male	15 to 19	87% (80-93%)	13% (7.2-20%)	2.3 (1.4-3.4)	11% (6-17%)	14% (7-21%)	760 (380-1100)	59000 (31000-86000)
Colombia	2000	Male	20 to 24	94% (91-96%)	5.8% (3.7-9.4%)	3.9 (2.5-5.6)	18% (11-25%)	20% (12-29%)	1600 (940-2400)	120000 (73000-170000)
Colombia	2000	Male	25 to 29	95% (92-97%)	4.7% (2.8-7.9%)	4.1 (2.8-5.8)	18% (11-25%)	20% (12-29%)	1500 (870-2100)	98000 (63000-140000)
Colombia	2000	Male	30 to 34	96% (93-98%)	4.4% (2.5-7.1%)	4.2 (3-5.9)	16% (11-22%)	19% (12-28%)	1200 (790-1700)	78000 (52000-110000)
Colombia	2000	Male	35 to 39	95% (92-97%)	5% (3.1-7.6%)	4.3 (2.8-6.1)	15% (10-20%)	19% (12-26%)	1100 (710-1500)	62000 (42000-84000)
Colombia	2000	Male	40 to 44	95% (92-97%)	5% (3.2-7.7%)	4.5 (3-6.3)	14% (9.1-18%)	18% (11-24%)	890 (570-1200)	46000 (31000-63000)
Colombia	2000	Male	45 to 49	96% (92-98%)	4.4% (2.4-7.7%)	4.9 (3.1-7.1)	13% (8.2-18%)	17% (11-24%)	830 (520-1200)	39000 (25000-54000)
Colombia	2000	Male	50 to 54	95% (92-97%)	5.2% (3-8.2%)	5.2 (3.3-7.3)	12% (7.8-17%)	16% (10-22%)	770 (470-1100)	32000 (20000-43000)
Colombia	2000	Male	55 to 59	93% (89-96%)	7.2% (4.3-11%)	5.4 (3.5-7.8)	12% (8-17%)	16% (9.9-21%)	780 (500-1100)	27000 (18000-37000)
Colombia	2000	Male	60 to 64	91% (87-95%)	9% (5.4-13%)	4.9 (3-7.3)	10% (5.9-15%)	13% (7-18%)	810 (450-1200)	24000 (14000-34000)
Colombia	2000	Male	65 to 69	90% (85-94%)	10% (6-15%)	3.7 (2.3-5.4)	7.7% (3.9-12%)	9.1% (4.4-14%)	710 (340-1100)	17000 (8700-26000)
Colombia	2000	Male	70 to 74	89% (83-93%)	11% (7.1-17%)	2.8 (1.7-4.3)	4.8% (1.6-8.8%)	5.4% (1.7-10%)	510 (160-940)	10000 (3400-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	2000	Male	75 to 79	87% (80-93%)	13% (7.2-20%)	2.2 (1.2-3.6)	3.1% (0.47-6.7%)	3.5% (0.29-7.6%)	310 (26-690)	4900 (760-11000)
Colombia	2000	Male	80 to 84	83% (72-91%)	17% (9-28%)	1.9 (1.1-3)	2.5% (0.48-5.3%)	2.7% (0.37-6%)	210 (30-480)	2500 (480-5300)
Colombia	2000	Male	85 to 89	80% (69-88%)	20% (12-31%)	1.7 (0.94-2.7)	2% (0.25-4.4%)	2.1% (0.067-4.9%)	120 (4-300)	1100 (130-2400)
Colombia	2000	Male	90 to 94	81% (70-89%)	19% (11-30%)	1.8 (0.95-2.9)	1.6% (-0.29-4.4%)	1.5% (-0.7-4.9%)	47 (-21-150)	300 (-54-830)
Colombia	2000	Male	95 plus	81% (69-91%)	19% (9-31%)	1.9 (0.91-3.3)	1.5% (-0.81-4.9%)	1.2% (-1.6-5.2%)	11 (-14-46)	52 (-29-180)
Colombia	2005	Female	All Ages	46% (41-51%)	54% (49-59%)	0.94 (0.75-1.1)	1.7% (1-2.5%)	2% (0.42-3.7%)	1700 (370-3200)	76000 (47000-110000)
Colombia	2005	Female	15 to 19	43% (32-56%)	57% (44-68%)	0.95 (0.55-1.5)	2.3% (1.3-3.6%)	4.4% (1.9-7.3%)	52 (23-89)	5700 (3200-8600)
Colombia	2005	Female	20 to 24	57% (44-69%)	43% (31-56%)	1.4 (0.78-2.2)	3.5% (2.2-5%)	5.7% (2.6-9.1%)	76 (35-120)	8900 (5300-13000)
Colombia	2005	Female	25 to 29	56% (43-67%)	44% (33-57%)	1.2 (0.72-1.8)	2.9% (1.9-4.1%)	4.8% (2.3-7.6%)	61 (29-96)	7000 (4400-10000)
Colombia	2005	Female	30 to 34	55% (42-67%)	45% (33-58%)	1.2 (0.68-1.7)	2.4% (1.6-3.5%)	4.3% (2.3-6.6%)	58 (30-90)	5600 (3600-8100)
Colombia	2005	Female	35 to 39	55% (43-67%)	45% (33-57%)	1.2 (0.76-1.7)	2.3% (1.4-3.2%)	4.4% (2.5-6.4%)	75 (43-110)	5900 (3800-8400)
Colombia	2005	Female	40 to 44	54% (42-65%)	46% (35-58%)	1.3 (0.8-1.8)	2.3% (1.4-3.3%)	4.5% (2.6-6.5%)	98 (56-140)	6100 (3700-8900)
Colombia	2005	Female	45 to 49	56% (41-68%)	44% (32-59%)	1.5 (0.88-2.3)	2.5% (1.3-4.1%)	5% (2.6-7.8%)	130 (71-210)	6600 (3600-10000)
Colombia	2005	Female	50 to 54	60% (46-72%)	40% (28-54%)	1.7 (1.1-2.7)	2.8% (1.2-4.8%)	5.2% (2.5-8.2%)	170 (83-270)	7200 (3400-12000)
Colombia	2005	Female	55 to 59	61% (47-74%)	39% (26-53%)	1.8 (1.1-2.9)	3.1% (1.3-5.5%)	5.3% (2.4-8.9%)	200 (91-340)	7100 (3000-13000)
Colombia	2005	Female	60 to 64	58% (43-73%)	42% (27-57%)	1.7 (0.92-2.8)	2.7% (0.6-5.3%)	4.1% (1.1-7.7%)	190 (51-370)	5700 (1300-11000)
Colombia	2005	Female	65 to 69	55% (39-70%)	45% (30-61%)	1.5 (0.74-2.2)	2.1% (0.17-4.4%)	3% (0.34-6%)	190 (22-380)	4500 (390-9200)
Colombia	2005	Female	70 to 74	52% (37-65%)	48% (35-63%)	1.2 (0.55-2.1)	1.4% (-0.46-3.9%)	2% (-0.48-5.2%)	160 (-40-430)	3100 (-950-8400)
Colombia	2005	Female	75 to 79	46% (32-63%)	54% (37-68%)	1 (0.46-1.8)	0.86% (-0.84-3%)	1.1% (-1-3.8%)	110 (-100-370)	1600 (-1500-5400)
Colombia	2005	Female	80 to 84	40% (25-57%)	60% (43-75%)	0.92 (0.45-1.5)	0.62% (-0.85-2.4%)	0.76% (-1.1-2.9%)	77 (-110-300)	850 (-1100-3200)
Colombia	2005	Female	85 to 89	38% (24-54%)	62% (46-76%)	0.9 (0.43-1.5)	0.44% (-1-2%)	0.46% (-1.3-2.4%)	44 (-130-230)	390 (-920-1800)
Colombia	2005	Female	90 to 94	42% (28-57%)	58% (43-72%)	1 (0.5-1.7)	0.19% (-1.3-2.1%)	0.085% (-1.8-2.3%)	5.2 (-110-140)	72 (-520-780)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	2005	Female	95 plus	46% (30-64%)	54% (36-70%)	1.1 (0.49-1.9)	0.038% (-1.8-2.4%)	-0.21% (-2.6-2.7%)	-5.5 (-68-71)	4.2 (-200-260)
Colombia	2005	Male	All Ages	89% (86-92%)	11% (8.5-14%)	2.9 (2.4-3.3)	8.9% (6.9-11%)	9.1% (6.6-12%)	10000 (7600-13000)	520000 (400000-650000)
Colombia	2005	Male	15 to 19	87% (80-93%)	13% (7.1-20%)	2.4 (1.5-3.4)	10% (5.8-15%)	14% (7-20%)	490 (250-740)	40000 (23000-58000)
Colombia	2005	Male	20 to 24	94% (91-96%)	5.8% (3.7-9.5%)	3.9 (2.6-5.7)	17% (11-23%)	20% (11-28%)	1200 (700-1700)	92000 (59000-130000)
Colombia	2005	Male	25 to 29	95% (92-97%)	4.7% (2.8-8%)	4.2 (2.8-5.9)	17% (11-23%)	20% (12-28%)	1100 (700-1600)	81000 (53000-110000)
Colombia	2005	Male	30 to 34	96% (93-97%)	4.4% (2.6-7.1%)	4.3 (3-6)	15% (11-20%)	19% (12-26%)	910 (590-1300)	60000 (41000-80000)
Colombia	2005	Male	35 to 39	95% (92-97%)	5% (3-7.6%)	4.3 (2.8-6.2)	13% (9.3-18%)	18% (12-24%)	810 (540-1100)	49000 (34000-66000)
Colombia	2005	Male	40 to 44	95% (92-97%)	5% (3.1-7.5%)	4.6 (3-6.4)	12% (8.6-16%)	17% (11-23%)	740 (490-1000)	41000 (28000-54000)
Colombia	2005	Male	45 to 49	96% (92-98%)	4.4% (2.4-7.8%)	4.9 (3.2-7.1)	12% (8-16%)	16% (11-23%)	740 (470-1000)	36000 (24000-48000)
Colombia	2005	Male	50 to 54	95% (92-97%)	5.2% (3.1-8.3%)	5.2 (3.4-7.4)	12% (7.5-16%)	16% (9.9-21%)	780 (500-1100)	33000 (22000-44000)
Colombia	2005	Male	55 to 59	93% (89-96%)	7.2% (4.3-11%)	5.5 (3.5-7.9)	11% (7.5-15%)	15% (9.7-20%)	810 (520-1100)	29000 (19000-39000)
Colombia	2005	Male	60 to 64	91% (87-95%)	9% (5.5-13%)	4.9 (3.1-7.3)	9.9% (5.6-14%)	12% (6.7-17%)	780 (430-1100)	24000 (13000-33000)
Colombia	2005	Male	65 to 69	90% (85-94%)	10% (5.9-15%)	3.8 (2.4-5.4)	7.5% (3.7-11%)	8.9% (4.2-14%)	720 (340-1100)	18000 (8700-27000)
Colombia	2005	Male	70 to 74	89% (84-93%)	11% (7-16%)	2.8 (1.7-4.3)	4.6% (1.6-8.5%)	5.2% (1.7-10%)	510 (160-970)	10000 (3600-19000)
Colombia	2005	Male	75 to 79	88% (80-93%)	12% (7.2-20%)	2.2 (1.2-3.7)	2.9% (0.37-6.6%)	3.2% (0.16-7.5%)	340 (17-800)	5400 (680-12000)
Colombia	2005	Male	80 to 84	84% (73-91%)	16% (8.9-27%)	1.9 (1.1-3)	2.3% (0.38-5%)	2.5% (0.22-5.6%)	250 (22-550)	2900 (490-6200)
Colombia	2005	Male	85 to 89	80% (69-89%)	20% (11-31%)	1.7 (0.96-2.8)	1.9% (0.25-4.2%)	2% (0.011-4.6%)	150 (0.81-350)	1300 (170-2900)
Colombia	2005	Male	90 to 94	80% (69-89%)	20% (11-31%)	1.8 (0.95-2.9)	1.6% (-0.27-4.4%)	1.5% (-0.75-4.9%)	58 (-29-190)	360 (-64-1100)
Colombia	2005	Male	95 plus	81% (68-91%)	19% (8.9-32%)	1.9 (0.93-3.3)	1.5% (-0.68-5.1%)	1.3% (-1.4-5.7%)	15 (-17-67)	71 (-33-250)
Colombia	2010	Female	All Ages	44% (38-49%)	56% (51-62%)	0.97 (0.76-1.2)	1.9% (1.2-2.6%)	2.1% (0.55-3.9%)	2000 (500-3600)	86000 (54000-120000)
Colombia	2010	Female	15 to 19	39% (28-52%)	61% (48-72%)	0.9 (0.52-1.5)	2.2% (1.4-3.4%)	4.2% (1.9-7.2%)	42 (19-72)	5100 (3000-7700)
Colombia	2010	Female	20 to 24	52% (38-65%)	48% (35-62%)	1.3 (0.71-2.2)	3.5% (2.3-5%)	5.4% (2.5-8.7%)	62 (29-100)	8500 (5400-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	2010	Female	25 to 29	52% (38-64%)	48% (36-62%)	1.2 (0.68-1.8)	3.2% (2.2-4.4%)	4.6% (2.3-7.2%)	57 (29-90)	7800 (5100-11000)
Colombia	2010	Female	30 to 34	51% (37-64%)	49% (36-63%)	1.1 (0.64-1.7)	2.7% (1.8-3.7%)	4.1% (2.2-6.3%)	55 (30-85)	6500 (4200-9100)
Colombia	2010	Female	35 to 39	51% (38-64%)	49% (36-62%)	1.2 (0.7-1.7)	2.4% (1.6-3.3%)	4.2% (2.5-6.2%)	65 (38-97)	6000 (4000-8300)
Colombia	2010	Female	40 to 44	50% (37-62%)	50% (38-63%)	1.2 (0.76-1.8)	2.3% (1.5-3.3%)	4.4% (2.7-6.3%)	93 (56-130)	6600 (4300-9300)
Colombia	2010	Female	45 to 49	53% (37-65%)	47% (35-63%)	1.5 (0.86-2.3)	2.6% (1.5-4%)	5.1% (2.9-7.8%)	140 (81-220)	7500 (4400-12000)
Colombia	2010	Female	50 to 54	58% (44-72%)	42% (28-56%)	1.8 (1.1-2.7)	3% (1.5-4.8%)	5.5% (2.9-8.6%)	200 (110-310)	8500 (4400-14000)
Colombia	2010	Female	55 to 59	60% (45-73%)	40% (27-55%)	1.9 (1.1-3)	3.1% (1.4-5.4%)	5.5% (2.7-9.2%)	240 (110-390)	8600 (3800-15000)
Colombia	2010	Female	60 to 64	57% (40-72%)	43% (28-60%)	1.7 (0.94-2.9)	2.7% (0.8-5.2%)	4.4% (1.4-8.1%)	230 (74-410)	6900 (2000-13000)
Colombia	2010	Female	65 to 69	53% (37-69%)	47% (31-63%)	1.5 (0.74-2.3)	2.3% (0.45-4.5%)	3.4% (0.71-6.5%)	220 (47-420)	5300 (1000-10000)
Colombia	2010	Female	70 to 74	49% (34-64%)	51% (36-66%)	1.2 (0.55-2.1)	1.7% (-0.13-4%)	2.3% (-0.094-5.4%)	200 (-7.8-460)	3700 (-310-9100)
Colombia	2010	Female	75 to 79	43% (28-60%)	57% (40-72%)	1 (0.47-1.9)	1.1% (-0.51-3.1%)	1.4% (-0.64-4%)	150 (-67-420)	2200 (-1100-6200)
Colombia	2010	Female	80 to 84	38% (24-55%)	62% (45-76%)	0.94 (0.44-1.6)	0.8% (-0.57-2.5%)	0.97% (-0.73-3.2%)	110 (-85-370)	1300 (-880-4100)
Colombia	2010	Female	85 to 89	38% (24-53%)	62% (47-76%)	0.95 (0.46-1.6)	0.62% (-0.81-2.2%)	0.67% (-1.1-2.6%)	78 (-130-300)	670 (-920-2400)
Colombia	2010	Female	90 to 94	41% (28-56%)	59% (44-72%)	1 (0.53-1.8)	0.38% (-1.1-2.3%)	0.31% (-1.5-2.6%)	24 (-120-200)	190 (-550-1100)
Colombia	2010	Female	95 plus	45% (28-63%)	55% (37-72%)	1.1 (0.49-2.1)	0.26% (-1.6-2.9%)	0.038% (-2.3-3.3%)	1.3 (-79-110)	35 (-220-390)
Colombia	2010	Male	All Ages	88% (85-91%)	12% (9.1-15%)	3.1 (2.5-3.6)	9.1% (7.2-11%)	9.1% (6.7-12%)	11000 (8000-14000)	530000 (420000-650000)
Colombia	2010	Male	15 to 19	84% (72-91%)	16% (8.6-28%)	2.3 (1.4-3.4)	9.7% (5.8-14%)	13% (7-20%)	450 (240-690)	37000 (22000-54000)
Colombia	2010	Male	20 to 24	93% (88-96%)	7% (4.1-12%)	4 (2.5-5.8)	17% (11-23%)	20% (12-29%)	1100 (660-1600)	83000 (55000-120000)
Colombia	2010	Male	25 to 29	95% (91-97%)	5.2% (3-9%)	4.3 (2.9-6.2)	17% (12-23%)	21% (13-29%)	1100 (740-1600)	81000 (55000-110000)
Colombia	2010	Male	30 to 34	95% (92-97%)	4.8% (2.7-8.1%)	4.5 (3.1-6.3)	15% (11-20%)	20% (13-28%)	920 (620-1300)	62000 (44000-82000)
Colombia	2010	Male	35 to 39	95% (91-97%)	5.4% (3.2-8.5%)	4.5 (2.9-6.4)	14% (9.7-19%)	19% (12-26%)	740 (480-1000)	46000 (31000-62000)
Colombia	2010	Male	40 to 44	95% (91-97%)	5.5% (3.3-8.5%)	4.7 (3.1-6.6)	12% (8.7-16%)	18% (12-24%)	720 (470-990)	41000 (29000-55000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	2010	Male	45 to 49	95% (91-97%)	4.9% (2.6-8.9%)	5.1 (3.2-7.3)	12% (8-16%)	17% (11-23%)	760 (500-1100)	38000 (26000-52000)
Colombia	2010	Male	50 to 54	94% (91-97%)	5.7% (3.3-9.4%)	5.4 (3.5-7.7)	11% (7.7-15%)	16% (10-21%)	830 (540-1100)	36000 (24000-48000)
Colombia	2010	Male	55 to 59	92% (87-95%)	8% (4.6-13%)	5.6 (3.6-8.2)	11% (7.6-15%)	15% (10-20%)	930 (620-1200)	34000 (24000-46000)
Colombia	2010	Male	60 to 64	90% (84-94%)	10% (5.9-16%)	5 (3.1-7.7)	9.6% (5.7-13%)	12% (7-17%)	880 (510-1200)	27000 (16000-38000)
Colombia	2010	Male	65 to 69	88% (82-93%)	12% (6.6-18%)	3.8 (2.3-5.6)	7.4% (3.9-11%)	9.1% (4.6-14%)	760 (390-1200)	19000 (10000-29000)
Colombia	2010	Male	70 to 74	86% (79-92%)	14% (8.3-21%)	2.7 (1.6-4.4)	4.6% (1.7-8.2%)	5.3% (1.9-9.8%)	550 (190-990)	11000 (4200-20000)
Colombia	2010	Male	75 to 79	84% (74-91%)	16% (8.6-26%)	2.1 (1.1-3.5)	3% (0.65-6.2%)	3.3% (0.58-7.2%)	380 (64-820)	6000 (1300-13000)
Colombia	2010	Male	80 to 84	81% (68-90%)	19% (10-32%)	1.8 (1-2.9)	2.4% (0.54-5.1%)	2.6% (0.38-5.8%)	290 (43-670)	3500 (760-7500)
Colombia	2010	Male	85 to 89	78% (66-87%)	22% (13-34%)	1.7 (0.91-2.8)	2% (0.36-4.5%)	2.1% (0.14-5%)	190 (13-460)	1600 (310-3700)
Colombia	2010	Male	90 to 94	78% (66-88%)	22% (12-34%)	1.8 (0.93-2.9)	1.7% (-0.13-4.7%)	1.6% (-0.53-5.1%)	80 (-26-250)	500 (-37-1400)
Colombia	2010	Male	95 plus	80% (65-91%)	20% (9.4-35%)	1.9 (0.92-3.4)	1.6% (-0.55-5.3%)	1.4% (-1.2-5.8%)	21 (-18-86)	97 (-34-320)
Colombia	2016	Female	All Ages	50% (44-55%)	50% (45-56%)	1.1 (0.84-1.5)	2% (1.2-3.1%)	2.3% (0.6-4.2%)	2200 (570-4100)	93000 (53000-140000)
Colombia	2016	Female	15 to 19	43% (30-55%)	57% (45-70%)	1 (0.54-1.7)	2.3% (1.4-3.6%)	4.6% (2-8.2%)	38 (17-69)	4800 (2700-7500)
Colombia	2016	Female	20 to 24	58% (45-70%)	42% (30-55%)	1.5 (0.79-2.6)	3.8% (2.5-5.4%)	6.1% (2.8-10%)	59 (26-100)	8400 (5200-13000)
Colombia	2016	Female	25 to 29	60% (45-71%)	40% (29-55%)	1.4 (0.8-2.2)	3.4% (2.3-4.8%)	5.4% (2.5-8.8%)	58 (27-96)	7900 (5100-11000)
Colombia	2016	Female	30 to 34	59% (45-71%)	41% (29-55%)	1.4 (0.79-2.1)	2.8% (1.8-4%)	4.8% (2.6-7.4%)	60 (32-95)	7100 (4400-10000)
Colombia	2016	Female	35 to 39	59% (47-71%)	41% (29-53%)	1.4 (0.85-2.2)	2.6% (1.6-3.7%)	4.9% (2.8-7.6%)	73 (40-120)	6700 (4200-9700)
Colombia	2016	Female	40 to 44	60% (47-71%)	40% (29-53%)	1.5 (0.95-2.3)	2.4% (1.4-3.7%)	5.2% (2.9-7.7%)	93 (52-140)	6500 (3800-10000)
Colombia	2016	Female	45 to 49	60% (46-72%)	40% (28-54%)	1.8 (1-2.7)	2.6% (1.4-4.2%)	5.5% (2.9-8.7%)	150 (78-240)	8000 (4200-13000)
Colombia	2016	Female	50 to 54	60% (47-73%)	40% (27-53%)	1.9 (1.1-2.9)	2.8% (1.3-4.7%)	5.6% (2.9-8.8%)	210 (110-340)	9300 (4300-16000)
Colombia	2016	Female	55 to 59	60% (45-73%)	40% (27-55%)	1.9 (1.1-3)	3% (1.2-5.2%)	5.4% (2.6-9.1%)	260 (120-430)	9400 (3900-16000)
Colombia	2016	Female	60 to 64	56% (41-72%)	44% (28-59%)	1.7 (0.93-2.9)	2.6% (0.72-5.2%)	4.4% (1.4-8.1%)	270 (88-500)	8300 (2300-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	2016	Female	65 to 69	53% (37-68%)	47% (32-63%)	1.5 (0.74-2.4)	2.2% (0.46-4.6%)	3.4% (0.86-6.7%)	270 (66-540)	6700 (1300-14000)
Colombia	2016	Female	70 to 74	49% (34-63%)	51% (37-66%)	1.2 (0.55-2.2)	1.7% (-0.074-4.1%)	2.4% (-0.013-5.6%)	230 (-1.1-540)	4500 (-200-11000)
Colombia	2016	Female	75 to 79	43% (28-60%)	57% (40-72%)	1 (0.45-1.9)	1.1% (-0.44-3.3%)	1.5% (-0.61-4.3%)	170 (-69-490)	2500 (-1100-7300)
Colombia	2016	Female	80 to 84	38% (24-54%)	62% (46-76%)	0.94 (0.44-1.6)	0.84% (-0.55-2.6%)	1% (-0.72-3.2%)	130 (-93-420)	1500 (-1000-4800)
Colombia	2016	Female	85 to 89	37% (23-52%)	63% (48-77%)	0.94 (0.44-1.6)	0.64% (-0.72-2.4%)	0.71% (-1.1-2.9%)	90 (-140-360)	780 (-910-2900)
Colombia	2016	Female	90 to 94	40% (27-55%)	60% (45-73%)	1 (0.49-1.9)	0.42% (-1-2.3%)	0.35% (-1.5-2.7%)	31 (-130-240)	240 (-600-1300)
Colombia	2016	Female	95 plus	44% (28-63%)	56% (37-72%)	1.1 (0.46-2.1)	0.31% (-1.5-2.9%)	0.073% (-2.3-3.2%)	3 (-89-130)	52 (-260-490)
Colombia	2016	Male	All Ages	89% (86-91%)	11% (8.7-14%)	3.2 (2.4-4.2)	9.2% (7-12%)	8.8% (6.1-12%)	11000 (7400-15000)	520000 (380000-660000)
Colombia	2016	Male	15 to 19	83% (72-91%)	17% (8.8-28%)	2.3 (1.3-3.6)	9.5% (5.5-14%)	13% (6.7-21%)	360 (180-560)	31000 (17000-45000)
Colombia	2016	Male	20 to 24	93% (88-96%)	7.1% (4.1-12%)	4 (2.4-6.1)	17% (11-23%)	21% (12-29%)	940 (530-1400)	74000 (46000-100000)
Colombia	2016	Male	25 to 29	95% (91-97%)	5.2% (3-9.1%)	4.3 (2.7-6.5)	17% (12-23%)	21% (13-30%)	990 (610-1500)	73000 (48000-100000)
Colombia	2016	Male	30 to 34	95% (92-97%)	4.8% (2.7-8.3%)	4.5 (2.9-6.7)	15% (11-21%)	20% (13-28%)	870 (560-1300)	61000 (42000-84000)
Colombia	2016	Male	35 to 39	95% (91-97%)	5.4% (3.1-8.6%)	4.5 (2.8-6.9)	14% (9.3-19%)	19% (12-27%)	710 (440-1100)	46000 (31000-64000)
Colombia	2016	Male	40 to 44	95% (91-97%)	5.5% (3.3-8.6%)	4.8 (2.9-7.1)	12% (8.3-17%)	18% (11-25%)	620 (380-900)	37000 (25000-51000)
Colombia	2016	Male	45 to 49	95% (91-97%)	4.9% (2.6-8.9%)	5.1 (3.1-7.7)	12% (7.5-16%)	17% (10-24%)	730 (430-1100)	39000 (25000-54000)
Colombia	2016	Male	50 to 54	94% (90-97%)	5.8% (3.4-9.5%)	5.4 (3.4-8)	11% (7.2-15%)	16% (10-22%)	870 (530-1200)	40000 (26000-54000)
Colombia	2016	Male	55 to 59	92% (87-95%)	8% (4.6-13%)	5.6 (3.5-8.4)	11% (7.2-14%)	15% (9.6-20%)	1000 (640-1400)	38000 (25000-52000)
Colombia	2016	Male	60 to 64	90% (84-94%)	10% (5.9-16%)	5 (3-7.9)	9.3% (5.3-13%)	12% (6.7-17%)	1000 (570-1500)	33000 (19000-46000)
Colombia	2016	Male	65 to 69	88% (82-93%)	12% (6.7-18%)	3.8 (2.2-5.7)	7.2% (3.6-11%)	8.9% (4.2-14%)	940 (460-1500)	24000 (12000-37000)
Colombia	2016	Male	70 to 74	86% (79-92%)	14% (8.3-21%)	2.7 (1.5-4.5)	4.5% (1.8-8.5%)	5.3% (1.9-10%)	630 (230-1200)	13000 (5000-24000)
Colombia	2016	Male	75 to 79	84% (74-91%)	16% (8.6-26%)	2.1 (1.1-3.6)	2.9% (0.71-6.6%)	3.3% (0.59-7.7%)	410 (73-950)	6700 (1500-15000)
Colombia	2016	Male	80 to 84	81% (68-89%)	19% (11-32%)	1.8 (1-3)	2.4% (0.51-5.2%)	2.6% (0.32-6%)	330 (43-750)	4000 (850-8700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Colombia	2016	Male	85 to 89	78% (65-87%)	22% (13-35%)	1.7 (0.89-2.9)	2% (0.35-4.7%)	2.1% (0.11-5.2%)	220 (11-550)	1900 (330-4400)
Colombia	2016	Male	90 to 94	78% (67-87%)	22% (13-33%)	1.8 (0.93-3.1)	1.7% (-0.11-4.7%)	1.6% (-0.51-5.1%)	96 (-30-300)	620 (-43-1700)
Colombia	2016	Male	95 plus	79% (65-90%)	21% (9.6-35%)	1.9 (0.89-3.5)	1.7% (-0.6-5.6%)	1.4% (-1.3-6%)	27 (-26-120)	130 (-50-450)
Costa Rica	1990	Female	All Ages	65% (58-70%)	35% (30-42%)	1 (0.88-1.2)	2.2% (1.5-3%)	3.8% (2-5.8%)	190 (100-290)	6300 (4300-8400)
Costa Rica	1990	Female	15 to 19	65% (49-77%)	35% (23-51%)	1.1 (0.62-1.7)	1.8% (1-2.7%)	3.8% (1.9-6.2%)	2 (0.94-3.2)	240 (130-360)
Costa Rica	1990	Female	20 to 24	77% (65-86%)	23% (14-35%)	1.3 (0.75-2)	2.5% (1.6-3.5%)	4.3% (2.3-6.8%)	2.7 (1.5-4.4)	380 (240-560)
Costa Rica	1990	Female	25 to 29	77% (65-85%)	23% (15-35%)	1.3 (0.79-1.9)	2.3% (1.5-3.3%)	4.1% (2.2-6.1%)	3 (1.6-4.5)	370 (230-540)
Costa Rica	1990	Female	30 to 34	76% (65-85%)	24% (15-35%)	1.3 (0.84-1.9)	2.2% (1.4-3.1%)	4.3% (2.7-6.3%)	4 (2.5-5.9)	360 (220-520)
Costa Rica	1990	Female	35 to 39	77% (66-85%)	23% (15-34%)	1.5 (0.9-2.2)	2.5% (1.5-3.6%)	5.4% (3.2-7.8%)	5.4 (3.2-7.8)	360 (220-530)
Costa Rica	1990	Female	40 to 44	78% (68-86%)	22% (14-32%)	1.8 (1.1-2.6)	2.8% (1.6-4.2%)	6.5% (4.2-9.1%)	7.5 (4.8-11)	420 (250-610)
Costa Rica	1990	Female	45 to 49	79% (67-87%)	21% (13-33%)	2.2 (1.3-3.3)	3.5% (1.8-5.6%)	7.3% (4.4-11%)	9.9 (5.8-15)	470 (250-750)
Costa Rica	1990	Female	50 to 54	76% (65-84%)	24% (16-35%)	2.6 (1.5-3.8)	4.3% (2.2-6.7%)	7.9% (4.6-12%)	14 (7.6-21)	560 (290-880)
Costa Rica	1990	Female	55 to 59	70% (57-80%)	30% (20-43%)	2.7 (1.6-4.2)	5.2% (2.6-8.4%)	8.6% (4.8-14%)	20 (11-31)	700 (360-1100)
Costa Rica	1990	Female	60 to 64	63% (49-76%)	37% (24-51%)	2.4 (1.3-4)	5% (2.3-8.6%)	7.5% (3.8-13%)	23 (12-40)	700 (330-1200)
Costa Rica	1990	Female	65 to 69	60% (46-74%)	40% (26-54%)	2.1 (1.2-3.2)	4.4% (2.1-7.3%)	6.4% (3.1-10%)	24 (12-39)	580 (270-950)
Costa Rica	1990	Female	70 to 74	56% (43-70%)	44% (30-57%)	1.9 (1.1-3.1)	3.9% (1.5-7.1%)	5.3% (2.2-9.7%)	24 (9.9-45)	470 (180-870)
Costa Rica	1990	Female	75 to 79	49% (34-65%)	51% (35-66%)	1.6 (0.82-2.6)	2.9% (0.71-6%)	3.7% (0.88-7.4%)	22 (5.4-44)	330 (78-670)
Costa Rica	1990	Female	80 to 84	45% (29-60%)	55% (40-71%)	1.4 (0.73-2.2)	2.1% (0.34-4.5%)	2.6% (0.28-5.6%)	16 (1.7-35)	180 (30-380)
Costa Rica	1990	Female	85 to 89	44% (29-60%)	56% (40-71%)	1.4 (0.71-2.3)	1.6% (-0.22-4%)	1.9% (-0.36-5%)	10 (-2-26)	81 (-11-200)
Costa Rica	1990	Female	90 to 94	48% (33-63%)	52% (37-67%)	1.4 (0.72-2.4)	1.1% (-0.78-4%)	1.2% (-1.2-4.7%)	3.6 (-3.6-14)	21 (-15-74)
Costa Rica	1990	Female	95 plus	52% (34-71%)	48% (29-66%)	1.4 (0.6-2.7)	0.7% (-1.6-4.3%)	0.6% (-2.3-5%)	0.59 (-2.2-5)	2.9 (-6.6-17)
Costa Rica	1990	Male	All Ages	69% (62-76%)	31% (24-38%)	2.4 (2.1-2.7)	6.9% (5.8-8.1%)	8.6% (6.4-11%)	570 (420-730)	24000 (20000-29000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	1990	Male	15 to 19	60% (42-77%)	40% (23-58%)	1.5 (0.75-2.4)	5.4% (3.5-7.8%)	7.7% (4.3-12%)	8.6 (4.8-14)	900 (580-1300)
Costa Rica	1990	Male	20 to 24	81% (69-89%)	19% (11-31%)	2.9 (1.7-4.5)	11% (7.7-15%)	14% (8.5-21%)	24 (14-36)	2300 (1600-3200)
Costa Rica	1990	Male	25 to 29	84% (74-91%)	16% (9.1-26%)	3.7 (2.3-5.5)	12% (8.8-17%)	17% (11-24%)	33 (20-47)	2700 (1900-3700)
Costa Rica	1990	Male	30 to 34	85% (75-92%)	15% (8.2-25%)	4.2 (2.7-5.9)	13% (9.1-17%)	18% (12-25%)	36 (24-49)	2600 (1800-3500)
Costa Rica	1990	Male	35 to 39	86% (77-92%)	14% (7.9-23%)	4.4 (2.5-6.6)	13% (9.1-17%)	19% (13-26%)	37 (25-51)	2300 (1600-3100)
Costa Rica	1990	Male	40 to 44	86% (78-93%)	14% (7.4-22%)	4.5 (2.6-6.6)	13% (9-17%)	19% (13-25%)	40 (27-53)	2200 (1600-2900)
Costa Rica	1990	Male	45 to 49	87% (77-93%)	13% (7.3-23%)	4.8 (2.7-8.1)	12% (8.3-16%)	19% (12-24%)	40 (26-52)	1900 (1300-2500)
Costa Rica	1990	Male	50 to 54	87% (78-93%)	13% (7.4-22%)	5.4 (3.1-8.6)	13% (8.5-16%)	18% (12-22%)	49 (33-63)	2000 (1400-2600)
Costa Rica	1990	Male	55 to 59	86% (76-93%)	14% (7.1-24%)	5.7 (3-8.9)	12% (7.7-16%)	16% (10-21%)	55 (35-72)	2000 (1300-2600)
Costa Rica	1990	Male	60 to 64	83% (71-92%)	17% (7.9-29%)	5 (2.6-8.4)	10% (6-14%)	13% (7.6-18%)	58 (34-79)	1700 (1000-2400)
Costa Rica	1990	Male	65 to 69	80% (67-90%)	20% (9.6-33%)	3.8 (2.2-6)	8.5% (4.7-12%)	10% (5.8-15%)	54 (30-77)	1300 (780-1900)
Costa Rica	1990	Male	70 to 74	76% (62-86%)	24% (14-38%)	2.9 (1.6-4.9)	6.2% (2.7-10%)	7.4% (3.1-12%)	45 (18-76)	890 (380-1500)
Costa Rica	1990	Male	75 to 79	70% (53-83%)	30% (17-47%)	2.3 (1.1-4)	4.5% (1.3-8.3%)	5.2% (1.4-9.6%)	36 (9.7-68)	550 (170-1000)
Costa Rica	1990	Male	80 to 84	64% (46-79%)	36% (21-54%)	1.9 (0.93-3.1)	3.7% (1.1-7.2%)	4.2% (1.1-8.2%)	28 (7.4-54)	310 (94-590)
Costa Rica	1990	Male	85 to 89	61% (44-76%)	39% (24-56%)	1.8 (0.85-3.1)	3.2% (0.75-6.8%)	3.6% (0.65-7.7%)	17 (3.1-37)	140 (32-290)
Costa Rica	1990	Male	90 to 94	63% (45-78%)	37% (22-55%)	1.9 (0.86-3.7)	3% (0.32-6.8%)	3.3% (0.06-7.8%)	7.7 (0.14-19)	44 (4.7-99)
Costa Rica	1990	Male	95 plus	66% (46-83%)	34% (17-54%)	2.2 (0.78-4.3)	3.1% (-0.22-7.4%)	3.3% (-0.64-8.4%)	2.1 (-0.42-5.4)	8.4 (-0.57-20)
Costa Rica	1995	Female	All Ages	65% (58-71%)	35% (29-42%)	1.1 (0.94-1.3)	2.5% (1.7-3.3%)	4.2% (2.5-6.3%)	250 (140-370)	7700 (5400-10000)
Costa Rica	1995	Female	15 to 19	65% (49-77%)	35% (23-51%)	1.1 (0.64-1.7)	1.8% (1.1-2.6%)	3.9% (1.9-6.4%)	2.4 (1.2-4)	290 (170-440)
Costa Rica	1995	Female	20 to 24	77% (64-85%)	23% (15-36%)	1.3 (0.75-2.1)	2.5% (1.6-3.6%)	4.5% (2.3-7.1%)	2.8 (1.4-4.6)	390 (240-570)
Costa Rica	1995	Female	25 to 29	76% (64-85%)	24% (15-36%)	1.3 (0.78-1.9)	2.3% (1.5-3.3%)	4.1% (2.2-6.1%)	3.2 (1.8-4.8)	390 (250-570)
Costa Rica	1995	Female	30 to 34	76% (65-85%)	24% (15-35%)	1.4 (0.86-2)	2.2% (1.4-3.2%)	4.5% (2.9-6.5%)	4.7 (3-6.8)	430 (280-600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	1995	Female	35 to 39	77% (67-85%)	23% (15-33%)	1.5 (0.9-2.2)	2.5% (1.5-3.7%)	5.7% (3.6-7.9%)	6.9 (4.3-9.8)	470 (290-670)
Costa Rica	1995	Female	40 to 44	78% (68-86%)	22% (14-32%)	1.8 (1.2-2.6)	3% (1.8-4.4%)	6.6% (4.5-9.1%)	9.5 (6.3-13)	520 (330-740)
Costa Rica	1995	Female	45 to 49	78% (67-86%)	22% (14-33%)	2.2 (1.3-3.4)	3.6% (2-5.6%)	7.7% (4.8-11%)	13 (8.4-20)	630 (360-990)
Costa Rica	1995	Female	50 to 54	76% (65-84%)	24% (16-35%)	2.6 (1.6-3.9)	4.4% (2.4-6.7%)	8.3% (5.2-12%)	16 (10-24)	680 (380-1000)
Costa Rica	1995	Female	55 to 59	69% (57-80%)	31% (20-43%)	2.7 (1.6-4.2)	5.2% (2.9-8.4%)	8.7% (5-14%)	23 (13-36)	820 (450-1300)
Costa Rica	1995	Female	60 to 64	63% (49-76%)	37% (24-51%)	2.4 (1.4-4)	5.1% (2.5-8.6%)	7.9% (4.2-13%)	27 (14-44)	820 (410-1400)
Costa Rica	1995	Female	65 to 69	60% (46-73%)	40% (27-54%)	2.1 (1.3-3.4)	4.7% (2.4-7.7%)	6.7% (3.6-11%)	31 (16-51)	760 (380-1200)
Costa Rica	1995	Female	70 to 74	56% (42-70%)	44% (30-58%)	1.9 (1.1-3.1)	4.2% (1.9-7.5%)	5.8% (2.8-10%)	32 (15-56)	630 (290-1100)
Costa Rica	1995	Female	75 to 79	49% (33-65%)	51% (35-67%)	1.6 (0.85-2.7)	3.3% (1-6.2%)	4.3% (1.4-8%)	28 (9-53)	420 (130-800)
Costa Rica	1995	Female	80 to 84	44% (29-60%)	56% (40-71%)	1.4 (0.75-2.2)	2.5% (0.68-4.9%)	3.1% (0.83-6.1%)	23 (6-45)	250 (70-500)
Costa Rica	1995	Female	85 to 89	44% (29-59%)	56% (41-71%)	1.4 (0.74-2.3)	1.9% (0.18-4.4%)	2.3% (0.096-5.3%)	15 (0.65-36)	120 (12-280)
Costa Rica	1995	Female	90 to 94	47% (33-62%)	53% (38-67%)	1.4 (0.72-2.4)	1.5% (-0.45-4.4%)	1.6% (-0.7-5.2%)	6.8 (-2.8-22)	38 (-11-120)
Costa Rica	1995	Female	95 plus	51% (33-70%)	49% (30-67%)	1.4 (0.6-2.7)	1% (-1.2-4.7%)	1% (-1.8-5.4%)	1.5 (-2.5-7.7)	6.2 (-6.8-27)
Costa Rica	1995	Male	All Ages	70% (63-77%)	30% (23-37%)	2.5 (2.2-2.9)	7.6% (6.4-9%)	9.1% (6.9-11%)	710 (540-890)	29000 (25000-34000)
Costa Rica	1995	Male	15 to 19	60% (42-76%)	40% (24-58%)	1.5 (0.78-2.5)	5.5% (3.6-8.2%)	7.9% (4.3-13%)	12 (6.3-18)	1200 (730-1700)
Costa Rica	1995	Male	20 to 24	81% (69-89%)	19% (11-31%)	3 (1.7-4.7)	12% (7.8-16%)	15% (8.6-22%)	28 (16-41)	2500 (1700-3500)
Costa Rica	1995	Male	25 to 29	84% (75-91%)	16% (9.1-25%)	3.8 (2.2-5.7)	12% (8.6-17%)	16% (10-24%)	37 (23-54)	3000 (2000-4100)
Costa Rica	1995	Male	30 to 34	85% (75-92%)	15% (8.2-25%)	4.3 (2.7-6.3)	12% (8.9-16%)	17% (12-23%)	43 (30-60)	3100 (2200-4200)
Costa Rica	1995	Male	35 to 39	86% (77-92%)	14% (8-23%)	4.4 (2.5-6.6)	12% (8.7-16%)	18% (12-24%)	47 (31-63)	2900 (2000-3800)
Costa Rica	1995	Male	40 to 44	86% (78-93%)	14% (7.4-22%)	4.5 (2.7-6.7)	13% (8.9-17%)	19% (13-24%)	51 (35-67)	2800 (1900-3600)
Costa Rica	1995	Male	45 to 49	87% (77-93%)	13% (7.3-23%)	4.9 (2.8-8.1)	13% (8.6-16%)	19% (12-24%)	55 (36-72)	2600 (1800-3400)
Costa Rica	1995	Male	50 to 54	87% (78-92%)	13% (7.6-22%)	5.5 (3.1-8.6)	13% (8.7-16%)	18% (12-23%)	59 (40-75)	2400 (1700-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	1995	Male	55 to 59	86% (76-93%)	14% (7.1-24%)	5.8 (3.2-9.3)	12% (8.1-16%)	16% (11-21%)	66 (44-87)	2400 (1600-3100)
Costa Rica	1995	Male	60 to 64	83% (71-92%)	17% (7.9-29%)	5.1 (2.7-8.5)	10% (6.4-14%)	13% (8.1-18%)	69 (42-93)	2100 (1300-2800)
Costa Rica	1995	Male	65 to 69	81% (68-90%)	19% (9.9-32%)	3.9 (2.3-6.2)	8.7% (5-12%)	11% (6.1-15%)	68 (39-96)	1700 (980-2300)
Costa Rica	1995	Male	70 to 74	76% (62-86%)	24% (14-38%)	3 (1.6-5.1)	6.5% (2.9-11%)	7.7% (3.3-13%)	58 (25-95)	1100 (510-1900)
Costa Rica	1995	Male	75 to 79	69% (53-83%)	31% (17-47%)	2.3 (1.1-4.1)	4.6% (1.5-8.4%)	5.4% (1.6-9.7%)	42 (13-75)	640 (210-1100)
Costa Rica	1995	Male	80 to 84	63% (46-78%)	37% (22-54%)	1.9 (0.95-3.2)	3.9% (1.4-7.1%)	4.4% (1.4-8.2%)	35 (11-64)	390 (140-710)
Costa Rica	1995	Male	85 to 89	60% (44-76%)	40% (24-56%)	1.8 (0.85-3.2)	3.4% (1.1-6.8%)	3.8% (0.99-7.8%)	24 (6.1-48)	190 (57-370)
Costa Rica	1995	Male	90 to 94	63% (46-78%)	37% (22-54%)	2 (0.89-3.8)	3.3% (0.53-7.2%)	3.6% (0.36-8.2%)	12 (1.2-27)	66 (10-140)
Costa Rica	1995	Male	95 plus	66% (47-83%)	34% (17-53%)	2.2 (0.82-4.4)	3.4% (0.017-7.8%)	3.6% (-0.39-8.7%)	3.5 (-0.38-8.5)	13 (0.068-31)
Costa Rica	2000	Female	All Ages	63% (57-68%)	37% (32-43%)	1 (0.84-1.2)	2.3% (1.7-3%)	3.8% (2.4-5.6%)	250 (160-370)	7900 (5700-10000)
Costa Rica	2000	Female	15 to 19	68% (54-79%)	32% (21-46%)	1.1 (0.68-1.7)	1.9% (1.1-2.9%)	4.1% (2-6.6%)	3.1 (1.5-4.9)	360 (200-540)
Costa Rica	2000	Female	20 to 24	77% (66-85%)	23% (15-34%)	1.2 (0.74-1.9)	2.4% (1.6-3.4%)	4.2% (2.2-6.8%)	3.1 (1.6-4.9)	430 (270-640)
Costa Rica	2000	Female	25 to 29	74% (64-83%)	26% (17-36%)	1.1 (0.68-1.7)	2.2% (1.5-3.1%)	3.9% (2.2-5.8%)	3 (1.7-4.5)	380 (240-540)
Costa Rica	2000	Female	30 to 34	73% (62-82%)	27% (18-38%)	1.1 (0.74-1.7)	2% (1.3-2.8%)	4.1% (2.6-5.7%)	4.1 (2.7-5.8)	400 (260-580)
Costa Rica	2000	Female	35 to 39	74% (65-83%)	26% (17-35%)	1.3 (0.8-1.9)	2.2% (1.4-3.1%)	5.2% (3.5-7.2%)	6.6 (4.4-9.2)	470 (310-670)
Costa Rica	2000	Female	40 to 44	74% (65-83%)	26% (17-35%)	1.5 (0.99-2.2)	2.7% (1.7-4%)	6.3% (4.4-8.3%)	11 (7.5-14)	600 (390-840)
Costa Rica	2000	Female	45 to 49	75% (64-83%)	25% (17-36%)	1.9 (1.1-2.8)	3.3% (1.9-4.9%)	6.9% (4.4-9.9%)	14 (9-20)	660 (380-1000)
Costa Rica	2000	Female	50 to 54	73% (62-81%)	27% (19-38%)	2.2 (1.4-3.3)	3.9% (2.3-6%)	7.5% (5-11%)	19 (12-27)	770 (460-1200)
Costa Rica	2000	Female	55 to 59	65% (52-76%)	35% (24-48%)	2.2 (1.3-3.5)	4.4% (2.6-6.9%)	7.6% (4.8-11%)	22 (14-33)	800 (470-1200)
Costa Rica	2000	Female	60 to 64	56% (43-70%)	44% (30-57%)	1.9 (1.1-3.1)	4.3% (2.3-6.9%)	6.9% (4-11%)	26 (15-40)	770 (430-1200)
Costa Rica	2000	Female	65 to 69	54% (41-67%)	46% (33-59%)	1.7 (0.96-2.6)	3.9% (2.1-6.1%)	5.8% (3.4-8.9%)	28 (17-44)	700 (400-1100)
Costa Rica	2000	Female	70 to 74	50% (38-64%)	50% (36-62%)	1.5 (0.85-2.5)	3.6% (1.8-6.2%)	5% (2.7-8.3%)	32 (17-54)	630 (320-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	2000	Female	75 to 79	45% (30-61%)	55% (39-70%)	1.3 (0.7-2.3)	2.9% (1.1-5.2%)	3.8% (1.5-6.9%)	29 (11-52)	430 (170-780)
Costa Rica	2000	Female	80 to 84	42% (27-58%)	58% (42-73%)	1.2 (0.64-2)	2.2% (0.72-4.1%)	2.8% (0.98-5.3%)	22 (7.8-42)	250 (85-460)
Costa Rica	2000	Female	85 to 89	42% (28-57%)	58% (43-72%)	1.2 (0.66-2)	1.7% (0.3-3.7%)	2.1% (0.35-4.6%)	17 (2.7-36)	130 (23-280)
Costa Rica	2000	Female	90 to 94	45% (31-60%)	55% (40-69%)	1.3 (0.65-2.2)	1.3% (-0.28-3.5%)	1.5% (-0.49-4.2%)	7.9 (-2.7-23)	45 (-9.8-120)
Costa Rica	2000	Female	95 plus	50% (32-68%)	50% (32-68%)	1.3 (0.54-2.4)	0.88% (-0.92-3.7%)	0.87% (-1.4-4.3%)	1.7 (-2.8-8.6)	7.4 (-7.6-31)
Costa Rica	2000	Male	All Ages	73% (66-79%)	27% (21-34%)	2.6 (2.2-3)	7.7% (6.5-9.1%)	8.8% (6.7-11%)	770 (590-960)	33000 (27000-39000)
Costa Rica	2000	Male	15 to 19	69% (53-81%)	31% (19-47%)	1.8 (0.92-2.8)	6.2% (4-9%)	9.2% (5.2-14%)	16 (8.8-25)	1500 (970-2200)
Costa Rica	2000	Male	20 to 24	85% (76-91%)	15% (9.5-24%)	3.1 (1.8-4.8)	12% (8.1-17%)	15% (9-23%)	35 (21-52)	3100 (2100-4300)
Costa Rica	2000	Male	25 to 29	85% (77-91%)	15% (9.1-23%)	3.6 (2.1-5.5)	13% (8.8-18%)	17% (11-25%)	38 (23-55)	3100 (2100-4300)
Costa Rica	2000	Male	30 to 34	86% (78-92%)	14% (8-22%)	4.1 (2.7-6.1)	12% (8.8-17%)	17% (12-24%)	44 (30-62)	3200 (2300-4300)
Costa Rica	2000	Male	35 to 39	86% (79-92%)	14% (7.8-21%)	4.2 (2.5-6.3)	12% (8.3-16%)	17% (12-24%)	50 (33-68)	3200 (2200-4300)
Costa Rica	2000	Male	40 to 44	86% (77-92%)	14% (7.9-23%)	4.2 (2.5-6.3)	12% (8.3-16%)	18% (12-23%)	61 (40-81)	3300 (2300-4400)
Costa Rica	2000	Male	45 to 49	86% (77-92%)	14% (7.9-23%)	4.6 (2.6-7.6)	12% (8.2-16%)	17% (12-23%)	62 (42-82)	3000 (2000-3900)
Costa Rica	2000	Male	50 to 54	87% (78-92%)	13% (7.8-22%)	5.2 (3-8.1)	12% (8.1-15%)	17% (11-21%)	68 (46-87)	2900 (2000-3700)
Costa Rica	2000	Male	55 to 59	86% (77-93%)	14% (7.5-23%)	5.5 (3.1-8.6)	11% (7.4-15%)	16% (10-20%)	70 (45-90)	2500 (1700-3300)
Costa Rica	2000	Male	60 to 64	83% (72-92%)	17% (8.3-28%)	4.8 (2.6-8)	9.7% (5.8-13%)	13% (7.4-17%)	69 (41-94)	2100 (1300-2900)
Costa Rica	2000	Male	65 to 69	81% (68-90%)	19% (9.9-32%)	3.7 (2.2-5.9)	8% (4.5-11%)	10% (5.6-14%)	70 (39-98)	1800 (1000-2400)
Costa Rica	2000	Male	70 to 74	77% (64-87%)	23% (13-36%)	3 (1.6-5.1)	6.1% (2.7-10%)	7.3% (3.2-12%)	64 (27-110)	1300 (580-2100)
Costa Rica	2000	Male	75 to 79	71% (56-84%)	29% (16-44%)	2.3 (1.1-4.1)	4.3% (1.5-7.8%)	5% (1.6-9.3%)	44 (15-82)	690 (240-1300)
Costa Rica	2000	Male	80 to 84	63% (46-78%)	37% (22-54%)	1.8 (0.91-3.1)	3.4% (1.3-6.3%)	3.9% (1.4-7.3%)	33 (12-62)	380 (140-700)
Costa Rica	2000	Male	85 to 89	60% (42-75%)	40% (25-58%)	1.7 (0.78-3)	3% (1-6.1%)	3.4% (0.99-6.9%)	25 (7.2-51)	200 (66-400)
Costa Rica	2000	Male	90 to 94	62% (45-77%)	38% (23-55%)	1.8 (0.8-3.6)	2.9% (0.49-6.5%)	3.1% (0.33-7.3%)	14 (1.5-33)	79 (13-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	2000	Male	95 plus	65% (45-83%)	35% (17-55%)	2.1 (0.74-4.1)	3% (0.049-7%)	3.1% (-0.34-7.8%)	4 (-0.44-10)	16 (0.26-37)
Costa Rica	2005	Female	All Ages	63% (57-68%)	37% (32-43%)	0.98 (0.81-1.2)	2.3% (1.7-2.9%)	3.7% (2.4-5.4%)	260 (170-380)	8200 (6100-11000)
Costa Rica	2005	Female	15 to 19	67% (55-78%)	33% (22-45%)	1 (0.62-1.6)	1.7% (1-2.6%)	4% (1.9-6.6%)	2.9 (1.4-4.8)	350 (200-510)
Costa Rica	2005	Female	20 to 24	76% (65-84%)	24% (16-35%)	1.1 (0.67-1.8)	2.4% (1.5-3.4%)	4.2% (2.2-6.6%)	3.4 (1.8-5.4)	470 (300-690)
Costa Rica	2005	Female	25 to 29	74% (63-82%)	26% (18-37%)	1.1 (0.62-1.6)	2.1% (1.4-2.9%)	3.8% (2.2-5.8%)	3 (1.7-4.5)	400 (250-570)
Costa Rica	2005	Female	30 to 34	72% (61-82%)	28% (18-39%)	1.1 (0.68-1.6)	2% (1.4-2.8%)	4.3% (2.9-5.9%)	4 (2.7-5.6)	390 (260-550)
Costa Rica	2005	Female	35 to 39	73% (63-82%)	27% (18-37%)	1.2 (0.7-1.8)	2% (1.3-2.9%)	5.2% (3.5-7%)	6 (4.1-8.3)	450 (300-620)
Costa Rica	2005	Female	40 to 44	74% (63-82%)	26% (18-37%)	1.4 (0.91-2)	2.5% (1.6-3.4%)	6.1% (4.4-8%)	11 (7.7-14)	610 (400-850)
Costa Rica	2005	Female	45 to 49	74% (63-82%)	26% (18-37%)	1.7 (1-2.6)	3% (1.8-4.5%)	6.7% (4.4-9.5%)	16 (11-23)	760 (450-1100)
Costa Rica	2005	Female	50 to 54	72% (61-81%)	28% (19-39%)	2 (1.2-3)	3.7% (2.2-5.4%)	7.3% (5-10%)	20 (13-27)	820 (510-1200)
Costa Rica	2005	Female	55 to 59	64% (51-76%)	36% (24-49%)	2.1 (1.2-3.2)	4.1% (2.4-6.2%)	7.4% (4.9-11%)	25 (17-37)	920 (560-1400)
Costa Rica	2005	Female	60 to 64	56% (41-69%)	44% (31-59%)	1.7 (0.97-2.8)	3.9% (2.3-6.2%)	6.6% (4.2-9.9%)	26 (17-39)	800 (470-1300)
Costa Rica	2005	Female	65 to 69	53% (40-67%)	47% (33-60%)	1.5 (0.91-2.4)	3.7% (2.2-5.5%)	5.8% (3.7-8.4%)	30 (19-43)	730 (450-1100)
Costa Rica	2005	Female	70 to 74	50% (37-63%)	50% (37-63%)	1.4 (0.8-2.3)	3.3% (1.8-5.4%)	4.9% (2.9-7.7%)	31 (19-50)	620 (350-1000)
Costa Rica	2005	Female	75 to 79	45% (30-60%)	55% (40-70%)	1.2 (0.66-2.1)	2.6% (1.2-4.7%)	3.6% (1.7-6.2%)	30 (14-52)	450 (210-810)
Costa Rica	2005	Female	80 to 84	41% (26-57%)	59% (43-74%)	1.1 (0.6-1.9)	2% (0.74-3.6%)	2.7% (1-4.9%)	23 (9.4-43)	260 (97-480)
Costa Rica	2005	Female	85 to 89	41% (28-56%)	59% (44-72%)	1.2 (0.62-1.9)	1.6% (0.33-3.3%)	2.1% (0.4-4.2%)	17 (3.3-36)	140 (28-280)
Costa Rica	2005	Female	90 to 94	45% (31-60%)	55% (40-69%)	1.2 (0.61-2.1)	1.3% (-0.19-3.4%)	1.5% (-0.36-4.1%)	9.4 (-2.3-26)	52 (-8-140)
Costa Rica	2005	Female	95 plus	49% (31-69%)	51% (31-69%)	1.2 (0.5-2.3)	0.94% (-0.8-3.6%)	0.98% (-1.2-4.3%)	2.6 (-3.1-11)	11 (-8.8-41)
Costa Rica	2005	Male	All Ages	74% (67-79%)	26% (21-33%)	2.5 (2.2-3)	7.6% (6.2-9%)	8.4% (6.4-11%)	780 (600-990)	34000 (28000-40000)
Costa Rica	2005	Male	15 to 19	69% (53-81%)	31% (19-47%)	1.6 (0.86-2.6)	6% (3.7-8.6%)	8.8% (4.8-14%)	16 (8.5-24)	1500 (960-2200)
Costa Rica	2005	Male	20 to 24	85% (75-90%)	15% (9.6-25%)	2.9 (1.6-4.4)	12% (7.7-16%)	15% (8.5-22%)	36 (21-53)	3300 (2200-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	2005	Male	25 to 29	85% (76-91%)	15% (9.2-24%)	3.4 (2-5.4)	12% (8.3-17%)	16% (9.7-24%)	38 (23-57)	3200 (2200-4500)
Costa Rica	2005	Male	30 to 34	86% (77-92%)	14% (8-23%)	3.9 (2.4-5.9)	12% (8.4-16%)	17% (11-24%)	39 (26-55)	2900 (2000-3900)
Costa Rica	2005	Male	35 to 39	86% (78-92%)	14% (8.1-22%)	4 (2.2-6.2)	11% (7.6-15%)	17% (11-23%)	45 (30-63)	3000 (2000-4000)
Costa Rica	2005	Male	40 to 44	85% (77-92%)	15% (8-23%)	4 (2.3-5.9)	11% (7.6-14%)	17% (11-22%)	59 (39-79)	3300 (2300-4400)
Costa Rica	2005	Male	45 to 49	86% (76-92%)	14% (8-24%)	4.3 (2.4-7.1)	11% (7.6-15%)	16% (11-22%)	71 (46-95)	3400 (2300-4600)
Costa Rica	2005	Male	50 to 54	86% (78-92%)	14% (8-22%)	4.9 (2.8-7.8)	11% (7.6-15%)	16% (11-21%)	76 (50-98)	3200 (2100-4200)
Costa Rica	2005	Male	55 to 59	86% (76-92%)	14% (7.5-24%)	5.2 (2.9-8.2)	11% (6.9-14%)	15% (9.7-20%)	81 (52-110)	3000 (1900-3900)
Costa Rica	2005	Male	60 to 64	83% (72-91%)	17% (8.5-28%)	4.5 (2.4-7.5)	9% (5.3-12%)	12% (7-16%)	70 (40-95)	2200 (1300-3000)
Costa Rica	2005	Male	65 to 69	80% (67-90%)	20% (10-33%)	3.5 (2.1-5.5)	7.3% (4.2-10%)	9.3% (5.3-13%)	70 (40-100)	1800 (1000-2600)
Costa Rica	2005	Male	70 to 74	77% (64-86%)	23% (14-36%)	2.8 (1.6-4.8)	5.5% (2.6-9%)	6.8% (3.1-11%)	59 (27-97)	1200 (550-2000)
Costa Rica	2005	Male	75 to 79	70% (54-83%)	30% (17-46%)	2.2 (1-3.8)	3.9% (1.3-7.2%)	4.6% (1.5-8.5%)	47 (15-87)	750 (260-1400)
Costa Rica	2005	Male	80 to 84	63% (45-78%)	37% (22-55%)	1.7 (0.84-2.9)	3% (1.1-5.5%)	3.5% (1.3-6.5%)	34 (12-64)	400 (150-730)
Costa Rica	2005	Male	85 to 89	59% (42-75%)	41% (25-58%)	1.6 (0.71-2.9)	2.7% (0.93-5.2%)	3% (0.92-6.1%)	23 (7.1-47)	190 (68-380)
Costa Rica	2005	Male	90 to 94	60% (43-76%)	40% (24-57%)	1.7 (0.71-3.3)	2.6% (0.55-5.8%)	2.8% (0.39-6.6%)	15 (2.1-35)	84 (18-190)
Costa Rica	2005	Male	95 plus	63% (43-81%)	37% (19-57%)	1.8 (0.64-3.8)	2.7% (0.16-6.4%)	2.7% (-0.22-7.2%)	4.4 (-0.36-12)	18 (1.1-42)
Costa Rica	2010	Female	All Ages	66% (60-71%)	34% (29-40%)	1.1 (0.89-1.3)	2.4% (1.7-3.1%)	4% (2.5-5.7%)	300 (190-440)	9400 (6700-12000)
Costa Rica	2010	Female	15 to 19	63% (47-76%)	37% (24-53%)	0.93 (0.51-1.5)	1.6% (0.9-2.3%)	3.5% (1.6-6.1%)	2.3 (1-4)	300 (180-450)
Costa Rica	2010	Female	20 to 24	76% (62-85%)	24% (15-38%)	1.1 (0.59-1.9)	2.3% (1.5-3.3%)	4.1% (2-6.7%)	3.5 (1.7-5.8)	490 (290-730)
Costa Rica	2010	Female	25 to 29	75% (63-85%)	25% (15-37%)	1.1 (0.62-1.7)	2.1% (1.4-3%)	4% (2.1-5.9%)	3.4 (1.8-5.1)	450 (280-650)
Costa Rica	2010	Female	30 to 34	75% (63-84%)	25% (16-37%)	1.1 (0.68-1.7)	2% (1.3-2.8%)	4.4% (2.9-6.3%)	4.4 (2.9-6.4)	430 (280-620)
Costa Rica	2010	Female	35 to 39	76% (65-85%)	24% (15-35%)	1.3 (0.7-2)	2.1% (1.3-3%)	5.2% (3.4-7.3%)	6.3 (4.2-9)	450 (290-650)
Costa Rica	2010	Female	40 to 44	77% (67-85%)	23% (15-33%)	1.5 (0.91-2.2)	2.4% (1.4-3.5%)	6.3% (4.4-8.3%)	11 (7.6-15)	620 (390-890)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	2010	Female	45 to 49	78% (65-86%)	22% (14-35%)	1.9 (1.1-2.8)	2.9% (1.6-4.5%)	7.1% (4.7-9.7%)	18 (11-25)	830 (490-1300)
Costa Rica	2010	Female	50 to 54	75% (64-84%)	25% (16-36%)	2.2 (1.3-3.2)	3.6% (2.1-5.4%)	7.5% (5.1-10%)	24 (16-33)	1000 (580-1500)
Costa Rica	2010	Female	55 to 59	69% (55-80%)	31% (20-45%)	2.3 (1.4-3.5)	4.3% (2.5-6.6%)	7.8% (5.1-11%)	31 (21-45)	1100 (680-1700)
Costa Rica	2010	Female	60 to 64	63% (48-75%)	37% (25-52%)	2 (1.1-3.3)	4.1% (2.3-6.3%)	7.2% (4.5-11%)	34 (21-50)	1000 (600-1600)
Costa Rica	2010	Female	65 to 69	60% (46-73%)	40% (27-54%)	1.8 (1.1-2.9)	3.8% (2.2-5.7%)	6.2% (4.1-9%)	34 (22-50)	850 (520-1300)
Costa Rica	2010	Female	70 to 74	55% (42-69%)	45% (31-58%)	1.6 (0.9-2.7)	3.4% (1.8-5.6%)	5.3% (3-8.1%)	36 (21-56)	720 (390-1200)
Costa Rica	2010	Female	75 to 79	48% (32-65%)	52% (35-68%)	1.4 (0.7-2.3)	2.6% (1.1-4.7%)	3.8% (1.8-6.7%)	33 (16-58)	510 (220-940)
Costa Rica	2010	Female	80 to 84	43% (28-59%)	57% (41-72%)	1.2 (0.61-2)	2% (0.74-3.8%)	2.8% (1.1-5.1%)	29 (12-53)	320 (120-610)
Costa Rica	2010	Female	85 to 89	42% (28-58%)	58% (42-72%)	1.2 (0.6-2)	1.6% (0.35-3.2%)	2.1% (0.46-4.2%)	21 (4.5-42)	170 (38-340)
Costa Rica	2010	Female	90 to 94	45% (31-60%)	55% (40-69%)	1.2 (0.6-2)	1.2% (-0.086-3.1%)	1.5% (-0.21-3.9%)	10 (-1.4-27)	60 (-4.1-150)
Costa Rica	2010	Female	95 plus	49% (32-69%)	51% (31-68%)	1.2 (0.48-2.3)	0.95% (-0.61-3.4%)	1% (-1.1-4.2%)	2.9 (-3-12)	12 (-8.1-45)
Costa Rica	2010	Male	All Ages	73% (66-79%)	27% (21-34%)	2.5 (2.1-3)	7.5% (6.1-9%)	8.3% (6.2-10%)	840 (630-1100)	36000 (29000-43000)
Costa Rica	2010	Male	15 to 19	58% (40-75%)	42% (25-60%)	1.3 (0.63-2.2)	5.1% (3.1-7.5%)	7.1% (3.5-12%)	12 (6.1-21)	1300 (780-1900)
Costa Rica	2010	Male	20 to 24	80% (67-88%)	20% (12-33%)	2.5 (1.4-4.1)	11% (7.2-15%)	13% (7.5-20%)	35 (20-54)	3300 (2100-4700)
Costa Rica	2010	Male	25 to 29	84% (73-91%)	16% (9.5-27%)	3.2 (1.8-5.1)	12% (8-16%)	16% (9.4-23%)	43 (26-63)	3600 (2400-5000)
Costa Rica	2010	Male	30 to 34	85% (74-92%)	15% (8.4-26%)	3.6 (2.2-5.5)	11% (8-16%)	16% (10-23%)	42 (27-61)	3200 (2200-4300)
Costa Rica	2010	Male	35 to 39	85% (76-92%)	15% (8.1-24%)	3.8 (2-5.9)	11% (7.2-15%)	16% (10-23%)	43 (27-61)	2800 (1900-3900)
Costa Rica	2010	Male	40 to 44	86% (77-92%)	14% (7.5-23%)	3.9 (2.2-5.9)	11% (7.2-14%)	16% (11-22%)	57 (37-78)	3200 (2200-4300)
Costa Rica	2010	Male	45 to 49	86% (76-93%)	14% (7.5-24%)	4.2 (2.3-7.1)	11% (7.1-14%)	16% (11-22%)	74 (48-99)	3600 (2400-4900)
Costa Rica	2010	Male	50 to 54	86% (77-92%)	14% (7.6-23%)	4.7 (2.8-7.4)	11% (7.3-14%)	16% (11-20%)	90 (59-120)	3800 (2500-5000)
Costa Rica	2010	Male	55 to 59	85% (75-93%)	15% (7.4-25%)	5 (2.8-7.7)	11% (7-14%)	15% (9.6-19%)	94 (60-120)	3400 (2200-4500)
Costa Rica	2010	Male	60 to 64	83% (70-92%)	17% (7.9-30%)	4.4 (2.3-7.2)	8.8% (5.1-12%)	12% (7-16%)	86 (50-120)	2700 (1600-3700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	2010	Male	65 to 69	80% (67-90%)	20% (9.9-33%)	3.4 (1.9-5.4)	7% (4.1-9.9%)	9.1% (5.3-13%)	74 (43-110)	1900 (1100-2800)
Costa Rica	2010	Male	70 to 74	75% (61-86%)	25% (14-39%)	2.6 (1.4-4.5)	5.1% (2.5-8.4%)	6.4% (3.1-11%)	60 (29-100)	1300 (600-2100)
Costa Rica	2010	Male	75 to 79	69% (53-83%)	31% (17-47%)	2 (0.92-3.6)	3.7% (1.3-6.7%)	4.4% (1.6-8%)	49 (18-89)	780 (300-1400)
Costa Rica	2010	Male	80 to 84	63% (45-78%)	37% (22-55%)	1.6 (0.78-2.8)	2.9% (1.1-5.2%)	3.4% (1.3-6.3%)	39 (15-71)	460 (180-860)
Costa Rica	2010	Male	85 to 89	59% (42-75%)	41% (25-58%)	1.5 (0.69-2.7)	2.5% (0.97-4.7%)	2.8% (1-5.6%)	26 (9.1-51)	220 (86-430)
Costa Rica	2010	Male	90 to 94	60% (43-75%)	40% (25-57%)	1.6 (0.66-3.1)	2.5% (0.59-5.2%)	2.7% (0.46-5.9%)	15 (2.6-33)	88 (21-190)
Costa Rica	2010	Male	95 plus	63% (43-81%)	37% (19-57%)	1.8 (0.62-3.6)	2.6% (0.28-6%)	2.7% (-0.079-6.8%)	4.6 (-0.14-12)	19 (2-44)
Costa Rica	2016	Female	All Ages	66% (60-71%)	34% (29-40%)	1.2 (0.87-1.6)	2.6% (1.8-3.6%)	4.1% (2.4-6.2%)	370 (220-570)	11000 (7400-16000)
Costa Rica	2016	Female	15 to 19	63% (46-76%)	37% (24-54%)	0.95 (0.49-1.5)	1.6% (0.9-2.4%)	3.6% (1.6-6.3%)	2.1 (0.91-3.6)	280 (160-440)
Costa Rica	2016	Female	20 to 24	75% (62-85%)	25% (15-38%)	1.1 (0.58-1.9)	2.3% (1.5-3.4%)	4.1% (2-7%)	3.2 (1.5-5.6)	480 (290-730)
Costa Rica	2016	Female	25 to 29	75% (63-84%)	25% (16-37%)	1.1 (0.6-1.9)	2.1% (1.4-3.1%)	4% (2.2-6.3%)	3.5 (1.9-5.5)	470 (290-690)
Costa Rica	2016	Female	30 to 34	75% (63-84%)	25% (16-37%)	1.1 (0.65-1.8)	2% (1.3-2.9%)	4.5% (2.9-6.6%)	4.9 (3.1-7.3)	490 (320-730)
Costa Rica	2016	Female	35 to 39	76% (64-85%)	24% (15-36%)	1.3 (0.68-2.1)	2.1% (1.4-3.2%)	5.3% (3.5-7.6%)	7 (4.4-10)	520 (330-780)
Costa Rica	2016	Female	40 to 44	76% (66-85%)	24% (15-34%)	1.6 (0.88-2.4)	2.5% (1.5-3.7%)	6.5% (4.3-8.9%)	11 (7-15)	610 (380-910)
Costa Rica	2016	Female	45 to 49	77% (65-86%)	23% (14-35%)	1.9 (1.1-3.1)	3% (1.6-4.8%)	7.2% (4.6-11%)	18 (11-26)	860 (500-1400)
Costa Rica	2016	Female	50 to 54	75% (63-83%)	25% (17-37%)	2.2 (1.3-3.5)	3.6% (2.1-5.7%)	7.6% (4.9-11%)	28 (18-41)	1200 (680-1900)
Costa Rica	2016	Female	55 to 59	68% (55-79%)	32% (21-45%)	2.3 (1.3-3.7)	4.3% (2.5-6.7%)	8% (5-12%)	40 (25-60)	1500 (860-2300)
Costa Rica	2016	Female	60 to 64	62% (47-75%)	38% (25-53%)	2.1 (1.1-3.4)	4.2% (2.4-6.7%)	7.3% (4.5-11%)	40 (25-61)	1200 (720-2000)
Costa Rica	2016	Female	65 to 69	59% (45-73%)	41% (27-55%)	1.8 (1-3)	3.9% (2.3-5.9%)	6.5% (4.2-9.7%)	45 (28-67)	1100 (670-1800)
Costa Rica	2016	Female	70 to 74	54% (41-68%)	46% (32-59%)	1.7 (0.9-2.8)	3.5% (1.8-5.9%)	5.5% (3.2-8.7%)	46 (26-77)	930 (480-1600)
Costa Rica	2016	Female	75 to 79	48% (32-64%)	52% (36-68%)	1.4 (0.68-2.4)	2.7% (1.2-5%)	4% (1.8-7%)	41 (19-74)	630 (270-1200)
Costa Rica	2016	Female	80 to 84	43% (27-59%)	57% (41-73%)	1.2 (0.58-2)	2% (0.69-4%)	2.8% (1-5.4%)	36 (13-68)	400 (140-800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Costa Rica	2016	Female	85 to 89	41% (28-56%)	59% (44-72%)	1.2 (0.58-2.1)	1.6% (0.33-3.5%)	2.1% (0.43-4.6%)	27 (5.3-59)	220 (47-470)
Costa Rica	2016	Female	90 to 94	44% (30-59%)	56% (41-70%)	1.2 (0.56-2.1)	1.3% (-0.084-3.4%)	1.5% (-0.23-4.2%)	15 (-2.3-42)	88 (-5.8-240)
Costa Rica	2016	Female	95 plus	49% (30-68%)	51% (32-70%)	1.2 (0.47-2.3)	0.98% (-0.59-3.5%)	1% (-1-4.2%)	4.4 (-4-18)	19 (-11-68)
Costa Rica	2016	Male	All Ages	74% (68-80%)	26% (20-32%)	2.7 (2-3.5)	7.9% (6.2-9.7%)	8.3% (5.9-11%)	1000 (700-1300)	41000 (31000-51000)
Costa Rica	2016	Male	15 to 19	58% (39-75%)	42% (25-61%)	1.3 (0.6-2.3)	5.1% (3.1-7.6%)	7.3% (3.4-12%)	11 (5.4-19)	1200 (710-1800)
Costa Rica	2016	Male	20 to 24	80% (67-88%)	20% (12-33%)	2.6 (1.3-4.3)	11% (7.2-16%)	14% (7.5-22%)	34 (19-54)	3300 (2100-4800)
Costa Rica	2016	Male	25 to 29	84% (73-90%)	16% (9.6-27%)	3.3 (1.8-5.4)	12% (8.1-17%)	16% (9.4-24%)	45 (25-69)	3900 (2600-5600)
Costa Rica	2016	Male	30 to 34	85% (74-91%)	15% (8.6-26%)	3.7 (2.1-5.9)	12% (8.1-16%)	17% (11-24%)	49 (30-72)	3800 (2500-5200)
Costa Rica	2016	Male	35 to 39	85% (76-92%)	15% (8.4-24%)	3.9 (1.9-6.3)	11% (7.3-15%)	17% (10-24%)	50 (29-73)	3400 (2100-4700)
Costa Rica	2016	Male	40 to 44	86% (76-92%)	14% (7.6-24%)	4 (2.1-6.4)	11% (7.3-15%)	17% (11-23%)	56 (35-78)	3200 (2100-4400)
Costa Rica	2016	Male	45 to 49	86% (76-92%)	14% (7.5-24%)	4.3 (2.2-7.5)	11% (7.1-15%)	17% (11-22%)	73 (47-100)	3600 (2400-5000)
Costa Rica	2016	Male	50 to 54	86% (77-92%)	14% (7.8-23%)	4.8 (2.6-7.8)	11% (7.3-15%)	16% (10-21%)	100 (66-140)	4400 (2900-5900)
Costa Rica	2016	Male	55 to 59	85% (75-93%)	15% (7.4-25%)	5 (2.8-8)	11% (7.2-14%)	15% (10-20%)	120 (80-160)	4500 (3000-6000)
Costa Rica	2016	Male	60 to 64	83% (70-92%)	17% (8-30%)	4.4 (2.3-7.3)	9% (5.3-12%)	12% (7.4-17%)	100 (60-140)	3200 (1900-4400)
Costa Rica	2016	Male	65 to 69	80% (67-90%)	20% (10-33%)	3.4 (1.9-5.4)	7.1% (4.1-10%)	9.3% (5.3-13%)	96 (55-140)	2500 (1500-3600)
Costa Rica	2016	Male	70 to 74	75% (60-86%)	25% (14-40%)	2.6 (1.4-4.5)	5.1% (2.4-8.5%)	6.5% (3.1-11%)	77 (36-130)	1600 (750-2600)
Costa Rica	2016	Male	75 to 79	69% (53-82%)	31% (18-47%)	2 (0.87-3.6)	3.7% (1.4-6.7%)	4.4% (1.6-8.1%)	60 (22-110)	970 (370-1800)
Costa Rica	2016	Male	80 to 84	62% (45-78%)	38% (22-55%)	1.7 (0.77-3)	2.9% (1.1-5.7%)	3.5% (1.3-6.8%)	49 (19-95)	590 (230-1100)
Costa Rica	2016	Male	85 to 89	59% (41-74%)	41% (26-59%)	1.5 (0.68-2.9)	2.6% (0.91-5%)	2.9% (0.96-5.8%)	36 (12-72)	310 (110-600)
Costa Rica	2016	Male	90 to 94	60% (43-75%)	40% (25-57%)	1.6 (0.66-3.2)	2.5% (0.66-5.4%)	2.7% (0.47-6.3%)	22 (4-50)	130 (34-280)
Costa Rica	2016	Male	95 plus	62% (42-81%)	38% (19-58%)	1.8 (0.62-3.6)	2.6% (0.32-5.9%)	2.7% (-0.052-6.8%)	6.8 (-0.13-17)	28 (3.4-63)
El Salvador	1990	Female	All Ages	32% (25-40%)	68% (60-75%)	0.48 (0.38-0.59)	1.3% (0.83-1.7%)	1.9% (0.88-3.2%)	270 (120-440)	11000 (6900-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	1990	Female	15 to 19	29% (16-46%)	71% (54-84%)	0.44 (0.21-0.79)	1.4% (0.64-2.5%)	1.9% (0.54-4%)	6.3 (1.8-13)	630 (290-1100)
El Salvador	1990	Female	20 to 24	38% (21-58%)	62% (42-79%)	0.56 (0.27-0.99)	2% (1.1-3.1%)	2.3% (0.83-4.3%)	7.1 (2.6-14)	810 (450-1300)
El Salvador	1990	Female	25 to 29	42% (23-62%)	58% (38-77%)	0.59 (0.28-0.99)	2.1% (1.4-3%)	2.5% (1.2-4%)	6.8 (3.2-11)	750 (460-1100)
El Salvador	1990	Female	30 to 34	43% (24-63%)	57% (37-76%)	0.63 (0.31-1)	2.3% (1.6-3.1%)	3% (1.7-4.5%)	8.1 (4.8-12)	730 (490-1000)
El Salvador	1990	Female	35 to 39	45% (26-65%)	55% (35-74%)	0.73 (0.36-1.2)	2.6% (1.8-3.7%)	3.9% (2.5-5.6%)	12 (7.6-17)	820 (560-1100)
El Salvador	1990	Female	40 to 44	46% (29-65%)	54% (35-71%)	0.88 (0.48-1.4)	2.9% (2-4.1%)	4.5% (2.9-6.6%)	16 (9.9-23)	880 (590-1200)
El Salvador	1990	Female	45 to 49	46% (26-66%)	54% (34-74%)	1.1 (0.53-1.8)	3.3% (2-5%)	5.2% (3.1-8.1%)	22 (13-33)	1000 (590-1500)
El Salvador	1990	Female	50 to 54	46% (27-66%)	54% (34-73%)	1.3 (0.66-2.1)	3.6% (2-5.3%)	5.5% (3.2-8.3%)	26 (15-39)	1100 (590-1600)
El Salvador	1990	Female	55 to 59	43% (25-63%)	57% (37-75%)	1.4 (0.68-2.4)	3.9% (2-6.4%)	5.8% (3.1-9.4%)	33 (18-55)	1100 (580-1900)
El Salvador	1990	Female	60 to 64	41% (22-62%)	59% (38-78%)	1.4 (0.65-2.4)	3.6% (1.4-6.5%)	5% (2.1-8.7%)	37 (15-65)	1000 (430-1900)
El Salvador	1990	Female	65 to 69	40% (22-60%)	60% (40-78%)	1.3 (0.61-2.2)	2.9% (0.91-5.4%)	3.9% (1.3-7.1%)	34 (12-62)	810 (260-1500)
El Salvador	1990	Female	70 to 74	36% (20-55%)	64% (45-80%)	1.1 (0.51-2.1)	2.2% (0.29-4.8%)	2.9% (0.43-6.3%)	28 (4.1-60)	530 (74-1200)
El Salvador	1990	Female	75 to 79	32% (16-53%)	68% (47-84%)	0.94 (0.38-1.8)	1.4% (-0.35-3.9%)	1.8% (-0.5-4.8%)	19 (-5.5-52)	280 (-72-770)
El Salvador	1990	Female	80 to 84	29% (14-49%)	71% (51-86%)	0.82 (0.31-1.6)	0.86% (-0.51-2.8%)	1% (-0.72-3.3%)	9.8 (-7-33)	110 (-70-350)
El Salvador	1990	Female	85 to 89	30% (15-48%)	70% (52-85%)	0.79 (0.34-1.5)	0.4% (-0.92-1.9%)	0.4% (-1.2-2.2%)	3.3 (-10-19)	30 (-70-140)
El Salvador	1990	Female	90 to 94	34% (18-52%)	66% (48-82%)	0.83 (0.34-1.6)	-0.041% (-1.4-1.8%)	-0.17% (-1.9-1.9%)	-0.78 (-8.8-8.6)	-1.1 (-41-50)
El Salvador	1990	Female	95 plus	38% (20-59%)	62% (41-80%)	0.85 (0.3-1.7)	-0.34% (-2-1.7%)	-0.59% (-2.7-1.9%)	-0.98 (-4.5-3.1)	-2.3 (-14-12)
El Salvador	1990	Male	All Ages	47% (37-57%)	53% (43-63%)	1.2 (0.99-1.4)	8.1% (6.4-9.5%)	10% (7.9-12%)	2000 (1600-2400)	96000 (76000-110000)
El Salvador	1990	Male	15 to 19	38% (20-59%)	62% (41-80%)	0.69 (0.31-1.3)	4.9% (2.6-8.2%)	5.3% (2.3-9.6%)	45 (19-82)	3900 (2000-6500)
El Salvador	1990	Male	20 to 24	58% (35-76%)	42% (24-65%)	1.5 (0.71-2.4)	11% (7-15%)	11% (6.2-16%)	140 (79-210)	11000 (6800-15000)
El Salvador	1990	Male	25 to 29	66% (46-82%)	34% (18-54%)	2 (1-3.2)	16% (11-21%)	16% (10-22%)	170 (110-230)	12000 (8600-16000)
El Salvador	1990	Male	30 to 34	71% (50-85%)	29% (15-50%)	2.3 (1.3-3.6)	19% (14-25%)	21% (14-27%)	180 (120-240)	12000 (8600-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	1990	Male	35 to 39	73% (55-86%)	27% (14-45%)	2.5 (1.3-4.1)	22% (16-27%)	25% (17-31%)	200 (140-260)	11000 (8300-14000)
El Salvador	1990	Male	40 to 44	74% (58-87%)	26% (13-42%)	2.8 (1.5-4.4)	22% (16-27%)	26% (18-32%)	200 (140-250)	10000 (7300-13000)
El Salvador	1990	Male	45 to 49	75% (54-87%)	25% (13-46%)	3 (1.5-5)	22% (16-27%)	27% (18-33%)	210 (140-270)	9300 (6700-12000)
El Salvador	1990	Male	50 to 54	74% (55-87%)	26% (13-45%)	3.2 (1.7-5)	20% (14-24%)	24% (17-31%)	200 (140-260)	8000 (5700-10000)
El Salvador	1990	Male	55 to 59	73% (54-87%)	27% (13-46%)	3.4 (1.8-5.4)	17% (12-23%)	22% (14-28%)	190 (120-240)	6400 (4300-8300)
El Salvador	1990	Male	60 to 64	71% (50-86%)	29% (14-50%)	3.1 (1.5-5.2)	14% (8.7-19%)	17% (11-23%)	160 (100-220)	4800 (3100-6400)
El Salvador	1990	Male	65 to 69	68% (46-84%)	32% (16-54%)	2.4 (1.2-4.1)	10% (6.6-15%)	13% (7.9-18%)	130 (83-190)	3200 (2000-4500)
El Salvador	1990	Male	70 to 74	63% (42-80%)	37% (20-58%)	1.8 (0.83-3.3)	7% (4.1-11%)	8.4% (4.8-13%)	93 (52-150)	1800 (1000-2900)
El Salvador	1990	Male	75 to 79	57% (34-78%)	43% (22-66%)	1.5 (0.56-3)	4.6% (2.1-8.5%)	5.2% (2.4-10%)	60 (27-120)	900 (420-1700)
El Salvador	1990	Male	80 to 84	53% (31-75%)	47% (25-69%)	1.3 (0.49-2.5)	3.1% (1.2-6.4%)	3.4% (1.2-7.1%)	32 (11-69)	360 (130-750)
El Salvador	1990	Male	85 to 89	52% (29-72%)	48% (28-71%)	1.2 (0.5-2.5)	2.4% (0.66-5.4%)	2.6% (0.59-6%)	17 (3.9-39)	130 (36-310)
El Salvador	1990	Male	90 to 94	54% (33-74%)	46% (26-67%)	1.3 (0.49-2.6)	1.9% (0.057-5.1%)	1.9% (-0.21-5.8%)	5.7 (-0.65-18)	34 (0.99-92)
El Salvador	1990	Male	95 plus	57% (32-78%)	43% (22-68%)	1.5 (0.47-3)	1.7% (-0.54-5.6%)	1.6% (-1-6.1%)	1.3 (-0.82-5.1)	5.8 (-1.8-18)
El Salvador	1995	Female	All Ages	32% (25-40%)	68% (60-75%)	0.52 (0.41-0.63)	1.6% (1.1-2.1%)	2.3% (1.1-3.7%)	310 (150-500)	12000 (8200-17000)
El Salvador	1995	Female	15 to 19	29% (16-47%)	71% (53-84%)	0.45 (0.21-0.82)	1.7% (0.77-3.1%)	2.5% (0.7-5.5%)	7.9 (2.1-17)	750 (320-1400)
El Salvador	1995	Female	20 to 24	38% (21-58%)	62% (42-79%)	0.58 (0.27-1)	2.3% (1.3-3.6%)	3% (1.2-5.6%)	8.3 (3.3-16)	910 (510-1500)
El Salvador	1995	Female	25 to 29	41% (22-61%)	59% (39-78%)	0.6 (0.29-1)	2.4% (1.6-3.4%)	3.2% (1.5-5.1%)	8.2 (4-14)	870 (560-1200)
El Salvador	1995	Female	30 to 34	43% (23-62%)	57% (38-77%)	0.65 (0.32-1.1)	2.6% (1.8-3.6%)	3.6% (2.2-5.6%)	10 (6.3-16)	890 (600-1200)
El Salvador	1995	Female	35 to 39	45% (26-64%)	55% (36-74%)	0.75 (0.36-1.3)	2.9% (2-4%)	4.4% (2.8-6.3%)	15 (9.5-21)	1000 (690-1400)
El Salvador	1995	Female	40 to 44	45% (28-65%)	55% (35-72%)	0.9 (0.47-1.5)	3% (2.1-4.2%)	4.8% (3.2-6.7%)	18 (12-25)	1000 (690-1400)
El Salvador	1995	Female	45 to 49	46% (27-65%)	54% (35-73%)	1.1 (0.53-1.8)	3.3% (2-5%)	5.3% (3.3-8%)	23 (14-35)	1100 (660-1600)
El Salvador	1995	Female	50 to 54	45% (27-66%)	55% (34-73%)	1.3 (0.67-2)	3.5% (2-5.2%)	5.6% (3.5-8.2%)	28 (17-41)	1100 (670-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	1995	Female	55 to 59	43% (25-64%)	57% (36-75%)	1.4 (0.69-2.5)	3.9% (2.1-6.3%)	5.8% (3.1-9.5%)	37 (20-59)	1300 (650-2100)
El Salvador	1995	Female	60 to 64	41% (22-62%)	59% (38-78%)	1.4 (0.68-2.4)	3.6% (1.5-6.4%)	5.1% (2.4-8.9%)	39 (18-69)	1100 (520-2000)
El Salvador	1995	Female	65 to 69	40% (22-60%)	60% (40-78%)	1.3 (0.63-2.3)	3% (1.1-5.5%)	4.1% (1.6-7.3%)	39 (15-70)	930 (340-1700)
El Salvador	1995	Female	70 to 74	36% (19-57%)	64% (43-81%)	1.2 (0.5-2.2)	2.4% (0.53-5.1%)	3.1% (0.72-6.6%)	36 (7.9-75)	670 (150-1500)
El Salvador	1995	Female	75 to 79	31% (15-52%)	69% (48-85%)	0.96 (0.37-1.9)	1.6% (-0.082-3.9%)	2% (-0.16-5%)	25 (-2-62)	360 (-20-910)
El Salvador	1995	Female	80 to 84	29% (13-48%)	71% (52-87%)	0.83 (0.32-1.6)	1% (-0.34-2.8%)	1.2% (-0.49-3.3%)	14 (-5.5-39)	150 (-51-420)
El Salvador	1995	Female	85 to 89	29% (15-47%)	71% (53-85%)	0.81 (0.34-1.5)	0.54% (-0.75-2.2%)	0.57% (-1-2.5%)	5.8 (-10-25)	49 (-68-190)
El Salvador	1995	Female	90 to 94	33% (17-51%)	67% (49-83%)	0.85 (0.33-1.6)	0.077% (-1.3-1.7%)	-0.025% (-1.7-2%)	-0.15 (-9.8-12)	2.8 (-45-62)
El Salvador	1995	Female	95 plus	38% (19-59%)	62% (41-81%)	0.87 (0.32-1.7)	-0.22% (-1.9-1.9%)	-0.45% (-2.5-2.1%)	-0.96 (-5.4-4.5)	-1.9 (-16-17)
El Salvador	1995	Male	All Ages	48% (38-58%)	52% (42-62%)	1.3 (1.1-1.5)	9.5% (7.7-11%)	11% (9-13%)	2200 (1700-2600)	100000 (82000-120000)
El Salvador	1995	Male	15 to 19	38% (20-60%)	62% (40-80%)	0.72 (0.32-1.3)	5.5% (2.7-9.2%)	6% (2.5-11%)	46 (19-84)	3900 (1900-6700)
El Salvador	1995	Male	20 to 24	58% (36-76%)	42% (24-64%)	1.5 (0.74-2.5)	12% (7.8-17%)	12% (7-19%)	130 (74-200)	10000 (6500-15000)
El Salvador	1995	Male	25 to 29	66% (45-82%)	34% (18-55%)	2 (1.1-3.3)	17% (12-22%)	17% (11-24%)	170 (110-240)	13000 (8700-17000)
El Salvador	1995	Male	30 to 34	71% (50-85%)	29% (15-50%)	2.4 (1.3-3.7)	20% (14-25%)	22% (15-28%)	210 (140-270)	13000 (9600-17000)
El Salvador	1995	Male	35 to 39	73% (55-86%)	27% (14-45%)	2.6 (1.4-4.1)	22% (16-28%)	25% (18-32%)	230 (160-300)	13000 (9600-16000)
El Salvador	1995	Male	40 to 44	74% (58-87%)	26% (13-42%)	2.8 (1.5-4.5)	22% (16-27%)	26% (19-32%)	220 (160-280)	11000 (8400-14000)
El Salvador	1995	Male	45 to 49	75% (54-87%)	25% (13-46%)	3.1 (1.6-5.2)	22% (16-27%)	26% (19-33%)	230 (160-290)	10000 (7500-13000)
El Salvador	1995	Male	50 to 54	74% (55-87%)	26% (13-45%)	3.3 (1.8-5.2)	20% (14-24%)	24% (17-31%)	210 (150-260)	8200 (6000-10000)
El Salvador	1995	Male	55 to 59	73% (54-87%)	27% (13-46%)	3.5 (1.8-5.7)	17% (12-22%)	22% (15-28%)	200 (140-260)	7000 (4800-9100)
El Salvador	1995	Male	60 to 64	71% (50-86%)	29% (14-50%)	3.2 (1.6-5.4)	14% (9.1-18%)	17% (11-23%)	180 (110-240)	5200 (3400-6900)
El Salvador	1995	Male	65 to 69	68% (46-84%)	32% (16-54%)	2.5 (1.2-4.1)	10% (6.7-15%)	13% (8.1-18%)	150 (95-210)	3600 (2300-5000)
El Salvador	1995	Male	70 to 74	63% (42-80%)	37% (20-58%)	1.9 (0.83-3.4)	7% (4.3-11%)	8.5% (5.1-13%)	100 (62-160)	2000 (1200-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	1995	Male	75 to 79	57% (34-78%)	43% (22-66%)	1.5 (0.57-3)	4.5% (2.3-8.1%)	5.3% (2.6-9.6%)	65 (31-120)	980 (490-1800)
El Salvador	1995	Male	80 to 84	53% (30-74%)	47% (26-70%)	1.3 (0.51-2.5)	3.1% (1.3-6.1%)	3.5% (1.3-7%)	38 (14-77)	430 (180-850)
El Salvador	1995	Male	85 to 89	51% (29-72%)	49% (28-71%)	1.3 (0.49-2.5)	2.4% (0.74-5.4%)	2.6% (0.65-6%)	19 (4.7-44)	150 (46-340)
El Salvador	1995	Male	90 to 94	54% (32-73%)	46% (27-68%)	1.4 (0.5-2.6)	1.9% (0.12-5.1%)	2% (-0.17-5.7%)	6.6 (-0.59-19)	39 (2.4-100)
El Salvador	1995	Male	95 plus	57% (32-78%)	43% (22-68%)	1.5 (0.48-3)	1.8% (-0.39-5.4%)	1.7% (-0.92-6%)	1.7 (-0.9-5.9)	7.3 (-1.5-22)
El Salvador	2000	Female	All Ages	32% (26-41%)	68% (59-74%)	0.59 (0.47-0.73)	1.9% (1.3-2.6%)	2.9% (1.5-4.5%)	370 (200-580)	13000 (9200-18000)
El Salvador	2000	Female	15 to 19	29% (16-46%)	71% (54-84%)	0.5 (0.23-0.9)	1.7% (0.85-3%)	2.8% (0.88-5.7%)	6.4 (2-13)	640 (320-1100)
El Salvador	2000	Female	20 to 24	38% (21-58%)	62% (42-79%)	0.63 (0.29-1.1)	2.4% (1.4-3.7%)	3.3% (1.3-6.2%)	8.2 (3.3-15)	920 (520-1500)
El Salvador	2000	Female	25 to 29	41% (22-62%)	59% (38-78%)	0.66 (0.3-1.1)	2.5% (1.6-3.5%)	3.4% (1.7-5.5%)	8.5 (4.4-14)	890 (580-1300)
El Salvador	2000	Female	30 to 34	43% (22-63%)	57% (37-78%)	0.72 (0.35-1.2)	2.6% (1.8-3.5%)	3.9% (2.4-5.9%)	9.8 (6.1-15)	870 (600-1200)
El Salvador	2000	Female	35 to 39	45% (26-64%)	55% (36-74%)	0.83 (0.41-1.4)	2.8% (1.9-3.8%)	4.6% (3-6.6%)	15 (9.4-21)	1000 (700-1400)
El Salvador	2000	Female	40 to 44	45% (28-65%)	55% (35-72%)	1 (0.52-1.6)	3% (2.1-4.2%)	5.1% (3.5-7%)	18 (13-26)	1000 (710-1400)
El Salvador	2000	Female	45 to 49	45% (26-66%)	55% (34-74%)	1.2 (0.59-2)	3.3% (2-4.8%)	5.7% (3.7-8.2%)	24 (15-34)	1100 (710-1700)
El Salvador	2000	Female	50 to 54	44% (26-65%)	56% (35-74%)	1.4 (0.72-2.3)	3.7% (2.1-5.5%)	5.9% (3.8-8.7%)	30 (19-43)	1200 (710-1800)
El Salvador	2000	Female	55 to 59	42% (24-63%)	58% (37-76%)	1.6 (0.76-2.7)	4.2% (2.3-6.8%)	6.3% (3.7-10%)	41 (24-66)	1400 (790-2300)
El Salvador	2000	Female	60 to 64	40% (21-62%)	60% (38-79%)	1.5 (0.74-2.8)	3.9% (1.9-6.7%)	5.7% (2.9-9.5%)	44 (23-74)	1300 (630-2200)
El Salvador	2000	Female	65 to 69	39% (21-59%)	61% (41-79%)	1.4 (0.69-2.4)	3.4% (1.6-6%)	4.9% (2.4-8.4%)	47 (23-81)	1100 (520-2000)
El Salvador	2000	Female	70 to 74	35% (19-56%)	65% (44-81%)	1.3 (0.56-2.4)	2.9% (1-5.9%)	3.9% (1.5-7.8%)	46 (17-91)	880 (310-1800)
El Salvador	2000	Female	75 to 79	31% (16-52%)	69% (48-84%)	1.1 (0.42-2.2)	2.1% (0.46-4.7%)	2.7% (0.63-6.1%)	36 (8.2-78)	530 (120-1200)
El Salvador	2000	Female	80 to 84	28% (13-49%)	72% (51-87%)	0.93 (0.35-1.9)	1.5% (0.092-3.6%)	1.8% (0.062-4.4%)	22 (0.76-55)	250 (15-600)
El Salvador	2000	Female	85 to 89	29% (14-47%)	71% (53-86%)	0.9 (0.37-1.7)	0.97% (-0.28-2.8%)	1.1% (-0.43-3.5%)	13 (-5-38)	100 (-30-300)
El Salvador	2000	Female	90 to 94	33% (17-52%)	67% (48-83%)	0.94 (0.39-1.7)	0.48% (-0.91-2.5%)	0.45% (-1.2-2.9%)	3.1 (-8.4-20)	20 (-37-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	2000	Female	95 plus	38% (19-59%)	62% (41-81%)	0.97 (0.35-1.9)	0.19% (-1.6-2.7%)	0.016% (-2.2-3%)	0.04 (-5.5-7.9)	2 (-17-28)
El Salvador	2000	Male	All Ages	48% (39-58%)	52% (42-61%)	1.5 (1.2-1.8)	10% (8.7-12%)	12% (9.9-14%)	2300 (1800-2700)	100000 (86000-120000)
El Salvador	2000	Male	15 to 19	38% (20-59%)	62% (41-80%)	0.81 (0.35-1.5)	5.6% (2.9-9.4%)	6.4% (2.6-12%)	41 (17-73)	3500 (1800-5900)
El Salvador	2000	Male	20 to 24	57% (35-75%)	43% (25-65%)	1.7 (0.83-2.8)	12% (8.1-18%)	13% (7.6-19%)	140 (83-210)	11000 (7100-16000)
El Salvador	2000	Male	25 to 29	66% (45-82%)	34% (18-55%)	2.3 (1.2-3.8)	17% (12-22%)	17% (11-24%)	170 (110-250)	13000 (9100-17000)
El Salvador	2000	Male	30 to 34	71% (50-85%)	29% (15-50%)	2.6 (1.5-4.1)	19% (14-25%)	21% (15-27%)	200 (140-260)	13000 (9900-17000)
El Salvador	2000	Male	35 to 39	73% (54-86%)	27% (14-46%)	2.9 (1.6-4.8)	21% (16-26%)	24% (18-30%)	220 (170-280)	13000 (9900-16000)
El Salvador	2000	Male	40 to 44	74% (56-87%)	26% (13-44%)	3.2 (1.7-5)	21% (17-26%)	26% (19-31%)	220 (160-270)	11000 (8600-14000)
El Salvador	2000	Male	45 to 49	74% (54-87%)	26% (13-46%)	3.5 (1.8-6)	21% (16-26%)	26% (19-32%)	230 (170-290)	10000 (8000-13000)
El Salvador	2000	Male	50 to 54	74% (54-87%)	26% (13-46%)	3.7 (2-5.9)	19% (14-24%)	23% (18-29%)	200 (150-260)	8100 (6200-10000)
El Salvador	2000	Male	55 to 59	73% (54-87%)	27% (13-46%)	4 (2-6.6)	17% (12-21%)	21% (15-27%)	210 (150-270)	7200 (5200-9200)
El Salvador	2000	Male	60 to 64	71% (50-86%)	29% (14-50%)	3.6 (1.8-6.1)	13% (9.3-17%)	17% (11-22%)	180 (120-240)	5400 (3700-7100)
El Salvador	2000	Male	65 to 69	67% (45-84%)	33% (16-55%)	2.8 (1.4-4.7)	10% (7-14%)	13% (8.6-18%)	160 (110-220)	3900 (2600-5400)
El Salvador	2000	Male	70 to 74	62% (41-80%)	38% (20-59%)	2.1 (0.94-3.9)	7.2% (4.3-11%)	8.7% (5.3-14%)	110 (68-180)	2200 (1300-3500)
El Salvador	2000	Male	75 to 79	56% (33-78%)	44% (22-67%)	1.7 (0.64-3.3)	4.9% (2.4-8.7%)	5.8% (2.8-10%)	82 (40-140)	1200 (620-2200)
El Salvador	2000	Male	80 to 84	52% (29-74%)	48% (26-71%)	1.5 (0.56-2.8)	3.6% (1.4-6.9%)	4% (1.6-7.9%)	48 (19-93)	540 (220-1000)
El Salvador	2000	Male	85 to 89	51% (29-72%)	49% (28-71%)	1.4 (0.55-2.8)	2.9% (0.92-6.2%)	3.2% (0.87-7.2%)	27 (7.4-62)	220 (71-490)
El Salvador	2000	Male	90 to 94	53% (32-73%)	47% (27-68%)	1.5 (0.55-3)	2.4% (0.28-5.9%)	2.5% (0.02-6.8%)	10 (0.079-27)	58 (6.9-150)
El Salvador	2000	Male	95 plus	56% (31-78%)	44% (22-69%)	1.7 (0.52-3.4)	2.4% (-0.18-6.6%)	2.3% (-0.73-7.4%)	2.7 (-0.84-8.6)	11 (-0.88-31)
El Salvador	2005	Female	All Ages	32% (26-40%)	68% (60-74%)	0.69 (0.54-0.83)	2.3% (1.6-3.2%)	3.4% (1.9-5.4%)	480 (270-770)	16000 (11000-22000)
El Salvador	2005	Female	15 to 19	28% (15-45%)	72% (55-85%)	0.54 (0.24-0.99)	1.8% (0.91-3%)	3% (1.1-6%)	7 (2.4-14)	690 (340-1200)
El Salvador	2005	Female	20 to 24	37% (20-57%)	63% (43-80%)	0.69 (0.32-1.2)	2.5% (1.4-4%)	3.6% (1.5-6.6%)	9.1 (3.8-17)	970 (540-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	2005	Female	25 to 29	40% (22-60%)	60% (40-78%)	0.73 (0.33-1.3)	2.5% (1.6-3.7%)	3.6% (1.8-6%)	9.8 (4.8-16)	960 (600-1400)
El Salvador	2005	Female	30 to 34	42% (22-62%)	58% (38-78%)	0.8 (0.39-1.3)	2.6% (1.7-3.6%)	4% (2.5-6.3%)	11 (7.1-18)	950 (630-1300)
El Salvador	2005	Female	35 to 39	44% (26-64%)	56% (36-74%)	0.92 (0.44-1.5)	2.8% (1.9-4%)	4.8% (3.2-6.9%)	16 (11-23)	1100 (720-1500)
El Salvador	2005	Female	40 to 44	45% (28-65%)	55% (35-72%)	1.1 (0.59-1.8)	3.1% (2.1-4.4%)	5.4% (3.7-7.4%)	21 (14-29)	1200 (800-1600)
El Salvador	2005	Female	45 to 49	45% (25-64%)	55% (36-75%)	1.4 (0.66-2.3)	3.5% (2.1-5.2%)	6% (3.9-8.6%)	28 (18-42)	1300 (820-2000)
El Salvador	2005	Female	50 to 54	44% (26-64%)	56% (36-74%)	1.6 (0.82-2.6)	3.9% (2.2-6%)	6.4% (4-9.3%)	36 (23-53)	1500 (860-2200)
El Salvador	2005	Female	55 to 59	42% (25-63%)	58% (37-75%)	1.8 (0.87-3.1)	4.6% (2.5-7.6%)	7% (4.1-11%)	47 (27-75)	1600 (900-2700)
El Salvador	2005	Female	60 to 64	40% (21-61%)	60% (39-79%)	1.8 (0.84-3.1)	4.5% (2.3-7.7%)	6.6% (3.5-11%)	57 (30-95)	1700 (850-2800)
El Salvador	2005	Female	65 to 69	39% (21-59%)	61% (41-79%)	1.6 (0.78-2.8)	4% (2-6.8%)	5.7% (3-9.4%)	61 (32-100)	1500 (730-2500)
El Salvador	2005	Female	70 to 74	35% (19-56%)	65% (44-81%)	1.4 (0.62-2.7)	3.4% (1.4-6.7%)	4.7% (2-9%)	61 (26-120)	1200 (480-2300)
El Salvador	2005	Female	75 to 79	31% (15-51%)	69% (49-85%)	1.2 (0.47-2.4)	2.5% (0.69-5.5%)	3.3% (0.93-7%)	51 (14-110)	760 (210-1600)
El Salvador	2005	Female	80 to 84	28% (13-47%)	72% (53-87%)	1 (0.4-2)	1.8% (0.41-4.1%)	2.3% (0.47-5.2%)	35 (7.1-80)	380 (82-870)
El Salvador	2005	Female	85 to 89	29% (15-47%)	71% (53-85%)	1 (0.42-1.9)	1.3% (-0.074-3.4%)	1.5% (-0.17-4.2%)	23 (-2.6-64)	190 (-10-490)
El Salvador	2005	Female	90 to 94	33% (17-50%)	67% (50-83%)	1.1 (0.43-2)	0.85% (-0.69-3.2%)	0.91% (-0.99-3.8%)	8.5 (-9.4-35)	50 (-40-190)
El Salvador	2005	Female	95 plus	38% (18-58%)	62% (42-82%)	1.1 (0.4-2.3)	0.59% (-1.3-3.6%)	0.51% (-1.8-4.2%)	1.8 (-6.6-15)	8.7 (-19-53)
El Salvador	2005	Male	All Ages	48% (39-58%)	52% (42-61%)	1.7 (1.4-2)	12% (9.6-13%)	13% (10-15%)	2400 (2000-2900)	110000 (94000-130000)
El Salvador	2005	Male	15 to 19	37% (19-58%)	63% (42-81%)	0.89 (0.38-1.7)	5.9% (3-10%)	6.8% (2.8-12%)	48 (20-89)	4100 (2100-6900)
El Salvador	2005	Male	20 to 24	57% (34-75%)	43% (25-66%)	1.9 (0.92-3.1)	13% (8.2-19%)	14% (7.9-20%)	150 (88-230)	12000 (7300-17000)
El Salvador	2005	Male	25 to 29	66% (45-81%)	34% (19-55%)	2.5 (1.3-4.2)	17% (12-23%)	18% (11-25%)	220 (140-320)	16000 (11000-21000)
El Salvador	2005	Male	30 to 34	70% (50-84%)	30% (16-50%)	3 (1.6-4.7)	20% (15-25%)	21% (15-28%)	220 (160-290)	14000 (11000-18000)
El Salvador	2005	Male	35 to 39	73% (55-86%)	27% (14-45%)	3.3 (1.8-5.4)	21% (17-27%)	24% (18-31%)	220 (170-290)	13000 (10000-17000)
El Salvador	2005	Male	40 to 44	74% (57-87%)	26% (13-43%)	3.6 (1.9-5.6)	21% (17-26%)	25% (20-32%)	230 (170-290)	12000 (9200-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	2005	Male	45 to 49	74% (53-87%)	26% (13-47%)	3.9 (2.1-6.8)	21% (16-26%)	26% (20-32%)	220 (170-280)	10000 (7800-13000)
El Salvador	2005	Male	50 to 54	74% (55-87%)	26% (13-45%)	4.2 (2.3-6.8)	19% (14-23%)	23% (18-29%)	210 (160-270)	8600 (6600-11000)
El Salvador	2005	Male	55 to 59	73% (54-87%)	27% (13-46%)	4.5 (2.4-7.3)	17% (12-21%)	20% (15-25%)	200 (150-260)	7100 (5300-9000)
El Salvador	2005	Male	60 to 64	71% (49-86%)	29% (14-51%)	4.1 (2-7)	13% (9.4-17%)	17% (12-21%)	180 (130-230)	5400 (3900-7000)
El Salvador	2005	Male	65 to 69	67% (46-84%)	33% (16-54%)	3.2 (1.6-5.3)	10% (7-14%)	13% (8.9-18%)	170 (110-230)	4100 (2800-5600)
El Salvador	2005	Male	70 to 74	63% (41-80%)	37% (20-59%)	2.4 (1.1-4.4)	7.2% (4.2-11%)	9% (5.2-14%)	120 (70-190)	2500 (1400-3900)
El Salvador	2005	Male	75 to 79	57% (33-77%)	43% (23-67%)	1.9 (0.74-3.8)	5.1% (2.5-8.7%)	6.1% (2.9-11%)	90 (43-160)	1400 (670-2400)
El Salvador	2005	Male	80 to 84	52% (29-74%)	48% (26-71%)	1.7 (0.64-3.2)	3.8% (1.5-7.3%)	4.4% (1.7-8.5%)	59 (22-110)	670 (270-1200)
El Salvador	2005	Male	85 to 89	50% (28-71%)	50% (29-72%)	1.6 (0.59-3.2)	3.2% (0.99-6.7%)	3.5% (0.94-7.8%)	40 (11-88)	320 (100-700)
El Salvador	2005	Male	90 to 94	52% (31-72%)	48% (28-69%)	1.7 (0.61-3.4)	2.7% (0.42-6.4%)	2.9% (0.21-7.3%)	17 (1.2-42)	97 (15-230)
El Salvador	2005	Male	95 plus	56% (30-77%)	44% (23-70%)	1.8 (0.57-3.8)	2.8% (0.022-7.1%)	2.8% (-0.5-8.1%)	4.4 (-0.79-13)	18 (0.14-47)
El Salvador	2010	Female	All Ages	33% (26-41%)	67% (59-74%)	0.78 (0.6-0.97)	2.7% (1.8-3.7%)	3.9% (2.1-6.1%)	590 (320-930)	19000 (12000-26000)
El Salvador	2010	Female	15 to 19	28% (15-45%)	72% (55-85%)	0.59 (0.26-1.1)	1.9% (0.91-3.3%)	3.3% (1.2-6.5%)	8.3 (3-16)	800 (380-1400)
El Salvador	2010	Female	20 to 24	37% (20-57%)	63% (43-80%)	0.75 (0.34-1.4)	2.6% (1.5-4.1%)	3.9% (1.6-7.3%)	8.8 (3.6-16)	970 (530-1500)
El Salvador	2010	Female	25 to 29	40% (21-60%)	60% (40-79%)	0.78 (0.35-1.4)	2.6% (1.6-3.9%)	3.9% (1.8-6.6%)	9.6 (4.5-16)	950 (580-1400)
El Salvador	2010	Female	30 to 34	41% (22-61%)	59% (39-78%)	0.84 (0.4-1.4)	2.8% (1.8-4%)	4.4% (2.6-6.9%)	13 (7.9-21)	1100 (690-1600)
El Salvador	2010	Female	35 to 39	43% (25-63%)	57% (37-75%)	0.97 (0.48-1.7)	2.9% (1.9-4.3%)	5.1% (3.3-7.5%)	18 (11-26)	1200 (770-1700)
El Salvador	2010	Female	40 to 44	44% (27-64%)	56% (36-73%)	1.2 (0.61-1.9)	3.1% (2.1-4.5%)	5.7% (3.9-8.1%)	22 (15-31)	1200 (810-1700)
El Salvador	2010	Female	45 to 49	44% (25-64%)	56% (36-75%)	1.5 (0.68-2.4)	3.6% (2.2-5.7%)	6.2% (3.9-9.4%)	31 (20-48)	1500 (880-2300)
El Salvador	2010	Female	50 to 54	44% (25-64%)	56% (36-75%)	1.7 (0.87-2.8)	4.2% (2.4-6.6%)	6.8% (4.1-10%)	42 (26-64)	1700 (1000-2700)
El Salvador	2010	Female	55 to 59	42% (24-62%)	58% (38-76%)	1.9 (0.94-3.3)	5% (2.8-8.2%)	7.6% (4.5-12%)	59 (34-97)	2100 (1100-3500)
El Salvador	2010	Female	60 to 64	40% (22-61%)	60% (39-78%)	1.9 (0.91-3.4)	5% (2.7-8.5%)	7.2% (4-12%)	68 (37-110)	2000 (1100-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	2010	Female	65 to 69	39% (21-59%)	61% (41-79%)	1.8 (0.86-3)	4.5% (2.3-7.6%)	6.4% (3.4-10%)	72 (38-120)	1800 (880-3000)
El Salvador	2010	Female	70 to 74	35% (19-55%)	65% (45-81%)	1.6 (0.7-3)	3.8% (1.7-7.5%)	5.3% (2.5-10%)	73 (34-140)	1400 (650-2800)
El Salvador	2010	Female	75 to 79	30% (15-51%)	70% (49-85%)	1.3 (0.52-2.6)	2.8% (0.93-6.2%)	3.8% (1.3-8%)	63 (22-130)	950 (320-2000)
El Salvador	2010	Female	80 to 84	28% (13-48%)	72% (52-87%)	1.1 (0.44-2.2)	2.1% (0.51-4.8%)	2.7% (0.63-5.9%)	47 (11-110)	520 (120-1200)
El Salvador	2010	Female	85 to 89	28% (14-46%)	72% (54-86%)	1.1 (0.45-2)	1.6% (0.074-3.9%)	1.9% (0.025-4.7%)	33 (0.43-83)	260 (12-650)
El Salvador	2010	Female	90 to 94	32% (16-50%)	68% (50-84%)	1.1 (0.44-2.2)	1.1% (-0.54-3.6%)	1.2% (-0.87-4.3%)	15 (-10-53)	84 (-43-280)
El Salvador	2010	Female	95 plus	37% (18-57%)	63% (43-82%)	1.2 (0.42-2.4)	0.85% (-1.2-4.1%)	0.81% (-1.7-4.8%)	3.9 (-8-23)	17 (-23-79)
El Salvador	2010	Male	All Ages	49% (40-59%)	51% (41-60%)	1.9 (1.5-2.3)	12% (10-15%)	13% (10-15%)	2500 (2100-3100)	110000 (94000-140000)
El Salvador	2010	Male	15 to 19	36% (19-58%)	64% (42-81%)	0.94 (0.39-1.8)	6.1% (3-11%)	7% (2.8-13%)	57 (24-100)	4700 (2300-8200)
El Salvador	2010	Male	20 to 24	56% (34-75%)	44% (25-66%)	2 (0.92-3.5)	13% (8.2-19%)	14% (7.7-22%)	140 (78-220)	11000 (6700-16000)
El Salvador	2010	Male	25 to 29	65% (44-82%)	35% (18-56%)	2.7 (1.4-4.7)	18% (12-25%)	19% (12-28%)	190 (120-280)	14000 (9200-19000)
El Salvador	2010	Male	30 to 34	70% (49-85%)	30% (15-51%)	3.2 (1.7-5.1)	20% (15-27%)	22% (16-31%)	250 (170-340)	16000 (11000-21000)
El Salvador	2010	Male	35 to 39	72% (54-86%)	28% (14-46%)	3.5 (1.9-5.8)	22% (17-28%)	25% (19-33%)	230 (170-310)	14000 (10000-17000)
El Salvador	2010	Male	40 to 44	74% (56-87%)	26% (13-44%)	3.9 (2-6.1)	22% (17-27%)	26% (20-33%)	230 (170-290)	12000 (9100-15000)
El Salvador	2010	Male	45 to 49	74% (53-87%)	26% (13-47%)	4.2 (2.2-7.2)	21% (16-26%)	25% (20-32%)	230 (180-300)	11000 (8400-13000)
El Salvador	2010	Male	50 to 54	74% (54-87%)	26% (13-46%)	4.5 (2.4-7.4)	19% (14-23%)	23% (18-29%)	220 (170-280)	9000 (6900-11000)
El Salvador	2010	Male	55 to 59	72% (54-87%)	28% (13-46%)	4.7 (2.5-7.8)	16% (12-20%)	20% (15-25%)	210 (160-270)	7400 (5500-9500)
El Salvador	2010	Male	60 to 64	70% (48-86%)	30% (14-52%)	4.3 (2.1-7.6)	13% (9.5-17%)	16% (11-21%)	190 (130-250)	5600 (4000-7400)
El Salvador	2010	Male	65 to 69	67% (45-84%)	33% (16-55%)	3.4 (1.7-5.7)	10% (7-14%)	13% (8.6-17%)	170 (110-220)	4100 (2800-5600)
El Salvador	2010	Male	70 to 74	62% (41-79%)	38% (21-59%)	2.6 (1.2-4.8)	7.2% (4-11%)	8.9% (5-14%)	140 (75-210)	2700 (1600-4200)
El Salvador	2010	Male	75 to 79	56% (34-77%)	44% (23-66%)	2 (0.81-4.1)	5.1% (2.4-8.7%)	6.2% (2.9-11%)	110 (49-180)	1600 (790-2800)
El Salvador	2010	Male	80 to 84	52% (29-73%)	48% (27-71%)	1.8 (0.67-3.6)	4% (1.7-7.4%)	4.6% (1.7-8.5%)	74 (28-140)	860 (330-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	2010	Male	85 to 89	50% (28-71%)	50% (29-72%)	1.7 (0.67-3.5)	3.4% (1.1-7%)	3.8% (1.1-8.3%)	53 (15-120)	440 (140-910)
El Salvador	2010	Male	90 to 94	52% (31-72%)	48% (28-69%)	1.8 (0.65-3.6)	3% (0.5-6.7%)	3.2% (0.3-7.7%)	27 (2.5-63)	150 (26-350)
El Salvador	2010	Male	95 plus	55% (29-77%)	45% (23-71%)	2 (0.57-4.2)	3% (0.16-7.5%)	3.1% (-0.32-8.8%)	7.6 (-0.78-21)	31 (1.5-77)
El Salvador	2016	Female	All Ages	33% (26-41%)	67% (59-74%)	0.87 (0.61-1.2)	2.9% (1.8-4.3%)	4% (2-6.7%)	680 (330-1200)	20000 (12000-30000)
El Salvador	2016	Female	15 to 19	27% (15-43%)	73% (57-85%)	0.6 (0.26-1.2)	1.8% (0.9-3.3%)	3.3% (1.1-6.8%)	6.7 (2.3-14)	680 (330-1200)
El Salvador	2016	Female	20 to 24	36% (19-56%)	64% (44-81%)	0.77 (0.34-1.5)	2.6% (1.5-4.1%)	4% (1.6-7.7%)	8.5 (3.4-16)	990 (560-1700)
El Salvador	2016	Female	25 to 29	39% (21-60%)	61% (40-79%)	0.81 (0.35-1.5)	2.6% (1.6-3.9%)	4.1% (1.9-7%)	9.3 (4.2-16)	960 (590-1500)
El Salvador	2016	Female	30 to 34	41% (21-61%)	59% (39-79%)	0.89 (0.4-1.6)	2.7% (1.7-4.1%)	4.7% (2.8-7.7%)	12 (7-20)	1000 (630-1500)
El Salvador	2016	Female	35 to 39	43% (25-63%)	57% (37-75%)	1 (0.48-1.8)	2.9% (1.9-4.5%)	5.3% (3.4-8.3%)	18 (11-28)	1200 (720-1800)
El Salvador	2016	Female	40 to 44	44% (27-64%)	56% (36-73%)	1.2 (0.62-2.1)	3.1% (2-4.7%)	5.9% (4-8.6%)	23 (15-34)	1300 (820-1900)
El Salvador	2016	Female	45 to 49	44% (23-64%)	56% (36-77%)	1.5 (0.67-2.6)	3.7% (2-6%)	6.5% (3.9-10%)	34 (19-54)	1600 (880-2600)
El Salvador	2016	Female	50 to 54	43% (24-64%)	57% (36-76%)	1.7 (0.87-3.1)	4.3% (2.2-7.1%)	7% (4-11%)	47 (26-79)	1900 (1000-3300)
El Salvador	2016	Female	55 to 59	41% (23-62%)	59% (38-77%)	1.9 (0.93-3.5)	5% (2.6-8.6%)	7.7% (4.3-13%)	65 (34-110)	2300 (1200-4000)
El Salvador	2016	Female	60 to 64	39% (21-60%)	61% (40-79%)	1.9 (0.87-3.6)	5% (2.5-8.9%)	7.3% (3.8-12%)	78 (40-140)	2300 (1100-4100)
El Salvador	2016	Female	65 to 69	37% (20-58%)	63% (42-80%)	1.8 (0.81-3.2)	4.6% (2.2-8.4%)	6.6% (3.3-12%)	80 (39-140)	2000 (940-3600)
El Salvador	2016	Female	70 to 74	34% (18-53%)	66% (47-82%)	1.6 (0.7-3.1)	3.9% (1.6-7.5%)	5.4% (2.4-10%)	82 (35-160)	1600 (660-3100)
El Salvador	2016	Female	75 to 79	29% (15-49%)	71% (51-85%)	1.3 (0.53-2.7)	2.9% (0.94-6.5%)	3.9% (1.4-8.5%)	76 (25-170)	1200 (360-2600)
El Salvador	2016	Female	80 to 84	26% (12-46%)	74% (54-88%)	1.1 (0.41-2.3)	2.2% (0.52-5.3%)	2.8% (0.61-6.6%)	65 (14-160)	720 (160-1800)
El Salvador	2016	Female	85 to 89	27% (14-44%)	73% (56-86%)	1.1 (0.44-2.1)	1.7% (0.11-4.2%)	2% (0.021-5.2%)	47 (0.51-120)	370 (24-930)
El Salvador	2016	Female	90 to 94	31% (16-49%)	69% (51-84%)	1.1 (0.44-2.3)	1.3% (-0.5-4%)	1.4% (-0.79-4.8%)	22 (-12-77)	130 (-49-400)
El Salvador	2016	Female	95 plus	36% (17-57%)	64% (43-83%)	1.2 (0.41-2.5)	1% (-1.2-4.4%)	1% (-1.7-5.3%)	7.2 (-12-37)	30 (-35-130)
El Salvador	2016	Male	All Ages	51% (42-61%)	49% (39-58%)	2.2 (1.6-2.9)	13% (11-16%)	13% (9.9-16%)	2600 (2000-3400)	120000 (90000-140000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
El Salvador	2016	Male	15 to 19	36% (19-57%)	64% (43-81%)	0.98 (0.39-1.9)	6.1% (3-11%)	7.3% (2.9-14%)	47 (19-88)	4000 (1900-7000)
El Salvador	2016	Male	20 to 24	56% (34-75%)	44% (25-66%)	2.1 (0.96-3.8)	13% (8-20%)	15% (7.6-23%)	140 (73-230)	11000 (6600-17000)
El Salvador	2016	Male	25 to 29	65% (43-82%)	35% (18-57%)	2.9 (1.4-4.9)	18% (12-25%)	20% (12-29%)	200 (120-300)	14000 (9500-21000)
El Salvador	2016	Male	30 to 34	70% (49-84%)	30% (16-51%)	3.4 (1.8-5.6)	21% (16-27%)	23% (16-32%)	230 (160-310)	15000 (11000-20000)
El Salvador	2016	Male	35 to 39	72% (54-86%)	28% (14-46%)	3.7 (1.9-6.4)	22% (17-29%)	26% (19-35%)	230 (170-320)	14000 (10000-18000)
El Salvador	2016	Male	40 to 44	73% (56-86%)	27% (14-44%)	4.1 (2.1-6.7)	22% (17-27%)	27% (21-35%)	230 (170-310)	12000 (9000-16000)
El Salvador	2016	Male	45 to 49	74% (52-87%)	26% (13-48%)	4.4 (2.1-7.8)	21% (16-26%)	26% (20-33%)	240 (170-310)	11000 (8100-14000)
El Salvador	2016	Male	50 to 54	73% (53-87%)	27% (13-47%)	4.7 (2.4-7.8)	19% (14-23%)	23% (18-29%)	230 (170-300)	9500 (6900-12000)
El Salvador	2016	Male	55 to 59	72% (53-86%)	28% (14-47%)	5 (2.5-8.4)	16% (12-20%)	20% (15-26%)	230 (160-300)	7900 (5800-10000)
El Salvador	2016	Male	60 to 64	70% (48-86%)	30% (14-52%)	4.5 (2.1-8)	16% (9.4-17%)	16% (12-21%)	200 (140-280)	6100 (4300-8100)
El Salvador	2016	Male	65 to 69	67% (44-84%)	33% (16-56%)	3.5 (1.7-6.1)	10% (7-14%)	13% (8.8-17%)	170 (110-230)	4200 (2800-5700)
El Salvador	2016	Male	70 to 74	62% (41-80%)	38% (20-59%)	2.6 (1.1-4.9)	7.2% (4-11%)	9.1% (5-14%)	140 (74-220)	2800 (1500-4400)
El Salvador	2016	Male	75 to 79	56% (32-77%)	44% (23-68%)	2.1 (0.75-4.2)	5.2% (2.4-8.7%)	6.4% (2.9-11%)	120 (52-200)	1800 (850-3200)
El Salvador	2016	Male	80 to 84	51% (28-73%)	49% (27-72%)	1.8 (0.64-3.7)	4.1% (1.6-7.6%)	4.8% (1.8-9%)	99 (36-180)	1100 (440-2000)
El Salvador	2016	Male	85 to 89	49% (27-70%)	51% (30-73%)	1.7 (0.62-3.5)	3.5% (1-7%)	4% (1-8.3%)	79 (20-160)	640 (190-1300)
El Salvador	2016	Male	90 to 94	51% (29-72%)	49% (28-71%)	1.9 (0.63-3.8)	3.2% (0.55-6.8%)	3.4% (0.37-7.9%)	41 (4.5-93)	230 (41-500)
El Salvador	2016	Male	95 plus	54% (29-77%)	46% (23-71%)	2 (0.57-4.4)	3.2% (0.19-7.5%)	3.3% (-0.29-8.7%)	13 (-1.2-35)	54 (3.2-130)
Guatemala	1990	Female	All Ages	10% (8.4-12%)	90% (88-92%)	1.2 (0.9-1.4)	2.5% (2.1-3.1%)	4.3% (3.5-5.3%)	1400 (1100-1800)	55000 (44000-67000)
Guatemala	1990	Female	15 to 19	9.4% (6.8-13%)	91% (87-93%)	0.99 (0.55-1.6)	3.1% (1.9-4.4%)	4.7% (2.7-7%)	33 (19-48)	2700 (1700-3900)
Guatemala	1990	Female	20 to 24	15% (11-20%)	85% (80-89%)	1.7 (0.94-2.7)	5.2% (3.6-7%)	7.2% (4.7-9.8%)	56 (34-79)	4500 (3000-6100)
Guatemala	1990	Female	25 to 29	18% (12-23%)	82% (77-88%)	1.9 (1.1-3)	6.8% (4.9-8.9%)	9.1% (6.3-12%)	72 (47-100)	5300 (3700-7000)
Guatemala	1990	Female	30 to 34	19% (12-25%)	81% (75-88%)	2.1 (1.2-3.2)	8% (6-10%)	11% (8.1-14%)	89 (61-120)	5800 (4300-7600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	1990	Female	35 to 39	21% (14-26%)	79% (74-86%)	2.5 (1.5-4)	9.1% (7.1-11%)	13% (9.9-15%)	110 (82-150)	6600 (4900-8500)
Guatemala	1990	Female	40 to 44	20% (14-25%)	80% (75-86%)	2.9 (1.7-4.4)	9.3% (7.4-12%)	13% (11-16%)	120 (88-160)	6100 (4600-7800)
Guatemala	1990	Female	45 to 49	17% (10-24%)	83% (76-90%)	3.1 (1.6-5.1)	8.8% (6.9-11%)	13% (10-16%)	110 (79-140)	4800 (3600-6100)
Guatemala	1990	Female	50 to 54	15% (9.9-22%)	85% (78-90%)	3.1 (1.7-5.1)	8.1% (6.2-10%)	12% (8.9-14%)	110 (81-150)	4300 (3200-5700)
Guatemala	1990	Female	55 to 59	14% (8.9-20%)	86% (80-91%)	3.2 (1.7-5.3)	7.4% (5.5-9.4%)	10% (7.9-13%)	120 (85-150)	3900 (2900-5100)
Guatemala	1990	Female	60 to 64	12% (7.4-18%)	88% (82-93%)	2.8 (1.4-4.9)	6.4% (4.7-8.5%)	8.3% (6.2-11%)	120 (86-170)	3400 (2500-4600)
Guatemala	1990	Female	65 to 69	11% (6.8-16%)	89% (84-93%)	2.4 (1.3-4)	5.9% (4.3-8%)	7.5% (5.5-10%)	120 (83-160)	2700 (1900-3700)
Guatemala	1990	Female	70 to 74	9.7% (6.2-14%)	90% (86-94%)	2.1 (1-3.7)	5.1% (3.6-7.2%)	6.3% (4.5-8.8%)	110 (72-150)	1900 (1400-2800)
Guatemala	1990	Female	75 to 79	8.5% (5.1-13%)	91% (87-95%)	1.7 (0.78-3.1)	4.3% (2.9-6.3%)	5.1% (3.4-7.4%)	88 (57-130)	1300 (830-1900)
Guatemala	1990	Female	80 to 84	7.7% (4.4-12%)	92% (88-96%)	1.4 (0.69-2.5)	3.7% (2.3-5.4%)	4.4% (2.7-6.4%)	66 (41-100)	700 (440-1000)
Guatemala	1990	Female	85 to 89	8% (4.7-13%)	92% (87-95%)	1.4 (0.69-2.6)	3.6% (2.1-5.7%)	4.2% (2.5-6.7%)	50 (29-79)	370 (220-580)
Guatemala	1990	Female	90 to 94	9.5% (5.7-14%)	90% (86-94%)	1.5 (0.71-2.8)	3.4% (2-5.5%)	4.1% (2.3-6.6%)	25 (15-41)	130 (76-210)
Guatemala	1990	Female	95 plus	12% (6.5-18%)	88% (82-93%)	1.7 (0.62-3.3)	3.4% (1.7-5.5%)	3.9% (1.8-6.5%)	7.3 (3.5-12)	25 (13-42)
Guatemala	1990	Male	All Ages	10% (8.5-13%)	90% (87-92%)	1.4 (1.1-1.7)	5.3% (4.4-6.1%)	7.5% (6.3-8.6%)	2800 (2200-3400)	130000 (100000-160000)
Guatemala	1990	Male	15 to 19	9% (5.5-14%)	91% (86-95%)	0.66 (0.32-1.2)	4.4% (3.1-6.2%)	4.8% (3-7.4%)	45 (26-71)	4400 (2900-6400)
Guatemala	1990	Male	20 to 24	19% (12-28%)	81% (72-88%)	2.1 (1.1-3.4)	12% (8.7-16%)	12% (8.1-17%)	160 (100-230)	13000 (9500-18000)
Guatemala	1990	Male	25 to 29	25% (16-34%)	75% (66-84%)	3.1 (1.7-5)	16% (13-20%)	17% (13-23%)	230 (160-300)	17000 (12000-22000)
Guatemala	1990	Male	30 to 34	26% (16-37%)	74% (63-84%)	3.5 (1.9-5.6)	19% (15-23%)	21% (16-27%)	280 (210-370)	18000 (14000-23000)
Guatemala	1990	Male	35 to 39	26% (16-35%)	74% (65-84%)	3.5 (1.9-5.8)	20% (17-24%)	23% (19-28%)	310 (230-400)	18000 (14000-22000)
Guatemala	1990	Male	40 to 44	25% (17-35%)	75% (65-83%)	3.5 (1.9-5.7)	19% (16-23%)	23% (20-28%)	290 (220-370)	15000 (11000-18000)
Guatemala	1990	Male	45 to 49	26% (16-36%)	74% (64-84%)	3.6 (1.9-6)	18% (16-22%)	23% (19-27%)	270 (200-340)	12000 (9300-15000)
Guatemala	1990	Male	50 to 54	25% (17-35%)	75% (65-83%)	3.8 (2.1-6.2)	16% (13-19%)	20% (17-24%)	250 (190-310)	9800 (7700-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	1990	Male	55 to 59	23% (15-32%)	77% (68-85%)	3.9 (2-6.4)	14% (11-16%)	17% (14-21%)	240 (180-300)	8000 (6300-10000)
Guatemala	1990	Male	60 to 64	20% (12-29%)	80% (71-88%)	3.2 (1.6-5.9)	11% (8.7-13%)	13% (11-16%)	220 (170-280)	6200 (4800-7900)
Guatemala	1990	Male	65 to 69	18% (11-25%)	82% (75-89%)	2.4 (1.3-4)	8.7% (7-11%)	11% (8.5-13%)	190 (140-240)	4400 (3400-5600)
Guatemala	1990	Male	70 to 74	15% (9.7-22%)	85% (78-90%)	1.7 (0.88-2.9)	6.6% (5.1-8.6%)	7.8% (6.1-10%)	140 (110-190)	2700 (2000-3600)
Guatemala	1990	Male	75 to 79	12% (7.2-18%)	88% (82-93%)	1.3 (0.58-2.3)	4.7% (3.4-6.3%)	5.5% (3.9-7.3%)	94 (66-130)	1400 (960-1900)
Guatemala	1990	Male	80 to 84	10% (6-15%)	90% (85-94%)	0.97 (0.46-1.7)	3.4% (2.3-4.6%)	3.9% (2.7-5.4%)	53 (35-75)	580 (390-800)
Guatemala	1990	Male	85 to 89	10% (5.9-16%)	90% (84-94%)	0.88 (0.42-1.6)	2.9% (1.9-4.2%)	3.3% (2.1-4.9%)	29 (18-43)	230 (140-330)
Guatemala	1990	Male	90 to 94	12% (7.2-19%)	88% (81-93%)	1 (0.44-2)	2.7% (1.6-4.1%)	3% (1.7-4.7%)	12 (7-20)	67 (40-110)
Guatemala	1990	Male	95 plus	15% (7.8-26%)	85% (74-92%)	1.2 (0.46-2.5)	2.7% (1.3-4.4%)	2.9% (1.3-5.1%)	3 (1.4-5.3)	11 (5.9-19)
Guatemala	1995	Female	All Ages	8.8% (7.3-10%)	91% (90-93%)	0.94 (0.73-1.2)	2.5% (2.1-3%)	4.5% (3.7-5.4%)	1400 (1100-1700)	51000 (41000-63000)
Guatemala	1995	Female	15 to 19	8.5% (5.4-12%)	92% (88-95%)	0.84 (0.44-1.4)	2.6% (1.5-3.9%)	4.2% (2.3-6.5%)	28 (15-43)	2400 (1400-3500)
Guatemala	1995	Female	20 to 24	13% (8.1-18%)	87% (82-92%)	1.3 (0.68-2.1)	4.2% (2.8-5.9%)	5.9% (3.6-8.5%)	43 (25-64)	3700 (2400-5200)
Guatemala	1995	Female	25 to 29	14% (8.7-20%)	86% (80-91%)	1.4 (0.77-2.3)	5.3% (3.7-7.1%)	7.4% (4.9-10%)	51 (31-74)	4100 (2700-5700)
Guatemala	1995	Female	30 to 34	15% (9.2-22%)	85% (78-91%)	1.5 (0.85-2.4)	6.4% (4.9-8.2%)	8.9% (6.5-12%)	69 (46-95)	4800 (3400-6300)
Guatemala	1995	Female	35 to 39	15% (9.2-21%)	85% (79-91%)	1.7 (0.93-2.8)	7.3% (5.7-9.3%)	10% (8-13%)	89 (63-120)	5300 (3900-6900)
Guatemala	1995	Female	40 to 44	15% (10-22%)	85% (78-90%)	2.1 (1.2-3.3)	8.1% (6.5-10%)	12% (9.6-15%)	110 (79-140)	5600 (4200-7400)
Guatemala	1995	Female	45 to 49	16% (9.5-22%)	84% (78-90%)	2.6 (1.3-4.4)	8.2% (6.4-10%)	12% (9.8-15%)	110 (82-150)	5100 (3800-6700)
Guatemala	1995	Female	50 to 54	16% (10-22%)	84% (78-90%)	3 (1.6-5)	8.1% (6.2-10%)	12% (9.3-15%)	110 (84-150)	4400 (3300-5800)
Guatemala	1995	Female	55 to 59	14% (9.3-20%)	86% (80-91%)	3 (1.6-5.2)	7.5% (5.6-9.6%)	11% (8.3-14%)	120 (85-150)	3900 (2900-5200)
Guatemala	1995	Female	60 to 64	12% (7.6-18%)	88% (82-92%)	2.6 (1.4-4.7)	6.4% (4.7-8.5%)	8.7% (6.5-11%)	120 (89-170)	3600 (2500-4900)
Guatemala	1995	Female	65 to 69	11% (6.8-15%)	89% (85-93%)	2.3 (1.2-3.8)	5.9% (4.3-7.9%)	7.8% (5.8-10%)	130 (92-180)	3100 (2100-4200)
Guatemala	1995	Female	70 to 74	9.8% (6.3-14%)	90% (86-94%)	2 (0.99-3.4)	5.2% (3.7-7.3%)	6.7% (4.8-9.3%)	120 (86-180)	2300 (1600-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	1995	Female	75 to 79	8.7% (5.1-13%)	91% (87-95%)	1.6 (0.74-2.9)	4.3% (2.9-6.3%)	5.3% (3.6-7.7%)	94 (61-140)	1400 (900-2000)
Guatemala	1995	Female	80 to 84	7.8% (4.5-12%)	92% (88-96%)	1.4 (0.66-2.4)	3.8% (2.4-5.6%)	4.5% (2.9-6.7%)	80 (49-120)	840 (520-1300)
Guatemala	1995	Female	85 to 89	8.1% (4.6-12%)	92% (88-95%)	1.3 (0.65-2.4)	3.7% (2.2-5.7%)	4.4% (2.6-6.9%)	56 (32-89)	410 (250-650)
Guatemala	1995	Female	90 to 94	9.6% (5.9-15%)	90% (85-94%)	1.4 (0.68-2.7)	3.6% (2.1-5.7%)	4.2% (2.5-6.8%)	28 (16-45)	140 (83-230)
Guatemala	1995	Female	95 plus	12% (6.7-18%)	88% (82-93%)	1.6 (0.6-3.1)	3.6% (1.7-5.8%)	4.1% (1.9-6.8%)	9.8 (4.4-16)	34 (16-56)
Guatemala	1995	Male	All Ages	10% (8.4-13%)	90% (87-92%)	1.3 (1-1.6)	6.5% (5.4-7.6%)	8.8% (7.6-10%)	3500 (2700-4300)	160000 (130000-200000)
Guatemala	1995	Male	15 to 19	9.1% (5.6-14%)	91% (86-94%)	0.64 (0.31-1.1)	4.4% (3-6.2%)	4.9% (2.9-7.6%)	54 (31-87)	5200 (3400-7700)
Guatemala	1995	Male	20 to 24	19% (12-27%)	81% (73-88%)	2 (1.1-3.3)	12% (8.8-16%)	13% (8.4-18%)	200 (130-300)	17000 (12000-23000)
Guatemala	1995	Male	25 to 29	25% (16-34%)	75% (66-84%)	2.9 (1.6-4.8)	16% (13-21%)	18% (13-23%)	280 (190-380)	20000 (15000-27000)
Guatemala	1995	Male	30 to 34	26% (16-37%)	74% (63-84%)	3.3 (1.8-5.2)	19% (16-24%)	22% (17-27%)	350 (260-470)	23000 (18000-29000)
Guatemala	1995	Male	35 to 39	26% (17-35%)	74% (65-83%)	3.4 (1.8-5.6)	21% (18-25%)	24% (20-29%)	400 (310-520)	23000 (18000-28000)
Guatemala	1995	Male	40 to 44	26% (17-35%)	74% (65-83%)	3.4 (1.8-5.5)	21% (18-24%)	25% (21-29%)	380 (290-480)	19000 (15000-23000)
Guatemala	1995	Male	45 to 49	26% (16-36%)	74% (64-84%)	3.5 (1.8-5.8)	19% (17-23%)	24% (21-28%)	360 (280-460)	16000 (13000-20000)
Guatemala	1995	Male	50 to 54	25% (17-35%)	75% (65-83%)	3.6 (1.9-5.8)	17% (14-20%)	21% (18-25%)	300 (230-380)	12000 (9100-15000)
Guatemala	1995	Male	55 to 59	23% (15-32%)	77% (68-85%)	3.7 (1.9-6.2)	15% (12-17%)	18% (15-22%)	270 (210-350)	9100 (7100-12000)
Guatemala	1995	Male	60 to 64	20% (13-29%)	80% (71-87%)	3.1 (1.6-5.6)	11% (9.3-14%)	14% (12-17%)	250 (190-320)	7100 (5500-9000)
Guatemala	1995	Male	65 to 69	18% (11-25%)	82% (75-89%)	2.3 (1.2-3.8)	9.1% (7.3-11%)	11% (9-14%)	220 (170-290)	5200 (4000-6600)
Guatemala	1995	Male	70 to 74	15% (9.8-22%)	85% (78-90%)	1.7 (0.85-2.8)	6.7% (5.2-8.5%)	8% (6.2-10%)	170 (130-230)	3200 (2400-4200)
Guatemala	1995	Male	75 to 79	12% (7-19%)	88% (81-93%)	1.2 (0.54-2.2)	4.8% (3.5-6.4%)	5.6% (4-7.5%)	110 (78-160)	1700 (1200-2300)
Guatemala	1995	Male	80 to 84	10% (5.9-15%)	90% (85-94%)	0.92 (0.43-1.6)	3.5% (2.4-4.8%)	4% (2.7-5.6%)	67 (45-96)	730 (490-1000)
Guatemala	1995	Male	85 to 89	10% (5.8-16%)	90% (84-94%)	0.84 (0.39-1.5)	3% (1.9-4.3%)	3.4% (2.1-5%)	36 (23-53)	270 (180-400)
Guatemala	1995	Male	90 to 94	12% (7.2-19%)	88% (81-93%)	0.97 (0.42-1.9)	2.7% (1.6-4.2%)	3.1% (1.7-4.8%)	14 (7.6-22)	75 (43-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	1995	Male	95 plus	16% (7.8-26%)	84% (74-92%)	1.2 (0.44-2.4)	2.8% (1.4-4.6%)	3% (1.4-5.1%)	3.9 (1.8-6.7)	15 (7.3-25)
Guatemala	2000	Female	All Ages	7.2% (6-8.4%)	93% (92-94%)	0.7 (0.53-0.88)	2.5% (2.1-3%)	4.6% (3.8-5.5%)	1400 (1100-1800)	49000 (39000-61000)
Guatemala	2000	Female	15 to 19	6.9% (4.4-10%)	93% (90-96%)	0.61 (0.32-1)	2.1% (1.2-3.3%)	3.4% (1.8-5.5%)	23 (11-39)	2100 (1200-3300)
Guatemala	2000	Female	20 to 24	9.5% (6.2-14%)	91% (86-94%)	0.87 (0.47-1.5)	3.3% (2.2-4.7%)	4.6% (2.7-6.9%)	35 (19-54)	3200 (2000-4600)
Guatemala	2000	Female	25 to 29	10% (6.6-15%)	90% (85-93%)	0.94 (0.52-1.5)	4% (2.9-5.5%)	5.5% (3.5-7.8%)	39 (24-58)	3400 (2400-4700)
Guatemala	2000	Female	30 to 34	11% (7.1-17%)	89% (83-93%)	1 (0.58-1.7)	5% (3.8-6.4%)	7.1% (5.1-9.4%)	53 (35-75)	4000 (2800-5300)
Guatemala	2000	Female	35 to 39	12% (7.3-17%)	88% (83-93%)	1.2 (0.66-2)	5.8% (4.5-7.4%)	8.6% (6.6-11%)	71 (49-99)	4400 (3200-6000)
Guatemala	2000	Female	40 to 44	12% (7.7-17%)	88% (83-92%)	1.5 (0.78-2.3)	6.7% (5.3-8.4%)	10% (8.1-12%)	92 (66-130)	4900 (3600-6600)
Guatemala	2000	Female	45 to 49	12% (7.7-18%)	88% (82-92%)	1.8 (0.94-3.2)	7% (5.6-8.9%)	11% (8.7-13%)	100 (76-140)	4800 (3600-6400)
Guatemala	2000	Female	50 to 54	12% (8.3-18%)	88% (82-92%)	2.1 (1.2-3.6)	7.2% (5.6-9.2%)	11% (8.7-13%)	120 (89-160)	4700 (3600-6400)
Guatemala	2000	Female	55 to 59	12% (7.8-17%)	88% (83-92%)	2.3 (1.2-3.9)	6.9% (5.3-8.9%)	10% (7.9-13%)	120 (84-150)	4000 (2900-5300)
Guatemala	2000	Female	60 to 64	11% (6.8-16%)	89% (84-93%)	2.1 (1.1-3.8)	6.3% (4.6-8.3%)	8.8% (6.8-11%)	120 (90-170)	3500 (2500-4800)
Guatemala	2000	Female	65 to 69	10% (6.5-15%)	90% (85-93%)	1.9 (1-3.2)	5.9% (4.4-8%)	8.2% (6.2-11%)	140 (100-200)	3400 (2400-4700)
Guatemala	2000	Female	70 to 74	9.3% (6.1-14%)	91% (86-94%)	1.7 (0.85-2.9)	5.3% (3.9-7.2%)	7% (5.2-9.5%)	150 (100-210)	2700 (2000-3900)
Guatemala	2000	Female	75 to 79	8.1% (4.7-12%)	92% (88-95%)	1.4 (0.62-2.5)	4.2% (3-6%)	5.5% (3.9-7.7%)	120 (77-170)	1700 (1100-2400)
Guatemala	2000	Female	80 to 84	7.4% (4.3-11%)	93% (89-96%)	1.2 (0.56-2)	3.7% (2.4-5.3%)	4.6% (3.1-6.6%)	89 (58-130)	940 (620-1400)
Guatemala	2000	Female	85 to 89	7.7% (4.4-12%)	92% (88-96%)	1.1 (0.56-2.1)	3.6% (2.2-5.4%)	4.4% (2.7-6.6%)	63 (39-100)	470 (290-740)
Guatemala	2000	Female	90 to 94	9% (5.6-14%)	91% (86-94%)	1.2 (0.58-2.3)	3.5% (2-5.5%)	4.2% (2.3-6.7%)	32 (18-51)	160 (93-260)
Guatemala	2000	Female	95 plus	11% (6-17%)	89% (83-94%)	1.4 (0.5-2.7)	3.5% (1.6-5.9%)	4.1% (1.8-7.1%)	11 (4.5-19)	38 (17-63)
Guatemala	2000	Male	All Ages	12% (9.8-14%)	88% (86-90%)	1.3 (1.1-1.7)	6.9% (5.8-8.2%)	9.1% (7.9-10%)	3600 (2800-4500)	170000 (130000-210000)
Guatemala	2000	Male	15 to 19	9.7% (6.5-14%)	90% (86-94%)	0.64 (0.33-1.1)	4.3% (2.9-6.3%)	4.7% (2.8-7.5%)	60 (33-100)	5800 (3800-8800)
Guatemala	2000	Male	20 to 24	25% (19-32%)	75% (68-81%)	2.5 (1.5-3.8)	13% (9.7-18%)	14% (9.7-20%)	260 (160-390)	21000 (15000-30000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	2000	Male	25 to 29	31% (23-37%)	69% (63-77%)	3.5 (2.2-5.2)	17% (13-21%)	18% (14-24%)	330 (230-450)	24000 (18000-32000)
Guatemala	2000	Male	30 to 34	25% (19-33%)	75% (67-81%)	3 (1.8-4.5)	17% (14-21%)	19% (15-24%)	330 (240-430)	22000 (16000-28000)
Guatemala	2000	Male	35 to 39	24% (18-32%)	76% (68-82%)	2.9 (1.7-4.7)	19% (16-22%)	22% (18-26%)	380 (280-500)	21000 (17000-27000)
Guatemala	2000	Male	40 to 44	26% (20-33%)	74% (67-80%)	3.2 (1.8-4.9)	19% (17-22%)	23% (19-26%)	380 (290-490)	19000 (15000-24000)
Guatemala	2000	Male	45 to 49	26% (19-34%)	74% (66-81%)	3.3 (1.9-5.3)	18% (16-21%)	22% (19-25%)	380 (280-480)	17000 (13000-21000)
Guatemala	2000	Male	50 to 54	26% (19-33%)	74% (67-81%)	3.4 (2-5.4)	16% (14-19%)	20% (17-23%)	330 (250-430)	13000 (9900-16000)
Guatemala	2000	Male	55 to 59	23% (17-30%)	77% (70-83%)	3.5 (2-5.7)	14% (11-16%)	17% (15-20%)	260 (200-340)	9000 (7100-11000)
Guatemala	2000	Male	60 to 64	19% (13-25%)	81% (75-87%)	2.7 (1.4-4.6)	11% (8.9-13%)	13% (11-15%)	230 (180-300)	6800 (5400-8500)
Guatemala	2000	Male	65 to 69	16% (11-22%)	84% (78-89%)	1.9 (1.1-3.1)	8.4% (7-10%)	10% (8.7-13%)	210 (170-270)	5100 (4000-6400)
Guatemala	2000	Male	70 to 74	16% (11-21%)	84% (79-89%)	1.6 (0.87-2.6)	6.7% (5.3-8.2%)	8.2% (6.5-10%)	200 (150-260)	3700 (2800-4900)
Guatemala	2000	Male	75 to 79	13% (8.3-20%)	87% (80-92%)	1.2 (0.6-2.2)	5.1% (3.7-6.6%)	6.1% (4.5-7.9%)	140 (100-200)	2100 (1500-2900)
Guatemala	2000	Male	80 to 84	10% (6-15%)	90% (85-94%)	0.87 (0.43-1.5)	3.5% (2.5-4.8%)	4.1% (2.8-5.6%)	83 (56-110)	900 (630-1200)
Guatemala	2000	Male	85 to 89	9.7% (5.5-15%)	90% (85-94%)	0.74 (0.34-1.4)	2.8% (1.8-4.2%)	3.2% (2-4.8%)	42 (26-63)	320 (200-480)
Guatemala	2000	Male	90 to 94	11% (6.6-18%)	89% (82-93%)	0.83 (0.36-1.6)	2.7% (1.5-4.1%)	3% (1.7-4.8%)	17 (9.1-27)	90 (51-140)
Guatemala	2000	Male	95 plus	15% (7.2-25%)	85% (75-93%)	1 (0.38-2)	2.7% (1.3-4.5%)	2.9% (1.3-5.1%)	4.3 (1.9-7.4)	16 (7.9-27)
Guatemala	2005	Female	All Ages	7.2% (6-8.5%)	93% (92-94%)	0.7 (0.54-0.87)	2.7% (2.3-3.3%)	5% (4.2-6%)	1600 (1200-2000)	54000 (43000-67000)
Guatemala	2005	Female	15 to 19	6.9% (4.5-10%)	93% (90-96%)	0.59 (0.31-0.99)	2% (1.1-3.1%)	3.2% (1.7-5.2%)	22 (12-37)	2100 (1300-3300)
Guatemala	2005	Female	20 to 24	9.4% (6-14%)	91% (86-94%)	0.84 (0.45-1.4)	3.1% (2.1-4.4%)	4.3% (2.5-6.5%)	35 (20-56)	3400 (2100-4900)
Guatemala	2005	Female	25 to 29	10% (6.5-15%)	90% (85-93%)	0.89 (0.49-1.5)	3.7% (2.6-4.9%)	5.2% (3.4-7.2%)	41 (25-60)	3600 (2500-5000)
Guatemala	2005	Female	30 to 34	11% (6.8-17%)	89% (83-93%)	0.99 (0.54-1.6)	4.5% (3.4-5.8%)	6.9% (5.2-8.8%)	54 (37-76)	4000 (2900-5300)
Guatemala	2005	Female	35 to 39	11% (7.2-16%)	89% (84-93%)	1.2 (0.61-2)	5.4% (4.2-6.9%)	8.5% (6.6-11%)	72 (51-100)	4500 (3300-6000)
Guatemala	2005	Female	40 to 44	12% (7.5-16%)	88% (84-92%)	1.4 (0.74-2.2)	6.3% (5-8%)	9.9% (8-12%)	95 (70-130)	5100 (3800-6600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	2005	Female	45 to 49	12% (7.4-17%)	88% (83-93%)	1.7 (0.9-3)	6.9% (5.4-8.8%)	11% (8.7-13%)	110 (83-150)	5200 (3900-6800)
Guatemala	2005	Female	50 to 54	12% (8.2-18%)	88% (82-92%)	2.1 (1.1-3.4)	7.2% (5.5-9.3%)	11% (8.6-13%)	140 (100-180)	5500 (4100-7400)
Guatemala	2005	Female	55 to 59	12% (7.7-17%)	88% (83-92%)	2.2 (1.2-3.8)	6.9% (5.2-9.1%)	10% (8-13%)	150 (110-200)	5000 (3600-6900)
Guatemala	2005	Female	60 to 64	11% (6.7-16%)	89% (84-93%)	2.1 (1.1-3.7)	6.5% (4.8-8.7%)	9.2% (7.1-12%)	140 (100-190)	4100 (2900-5600)
Guatemala	2005	Female	65 to 69	10% (6.4-15%)	90% (85-94%)	1.9 (1-3.2)	6.2% (4.6-8.3%)	8.6% (6.7-11%)	160 (120-220)	3800 (2700-5200)
Guatemala	2005	Female	70 to 74	9.2% (6.1-13%)	91% (87-94%)	1.6 (0.84-2.8)	5.5% (4-7.5%)	7.5% (5.7-9.9%)	170 (120-230)	3200 (2200-4400)
Guatemala	2005	Female	75 to 79	8% (4.8-12%)	92% (88-95%)	1.3 (0.61-2.4)	4.4% (3.1-6.1%)	5.8% (4.2-7.9%)	150 (99-210)	2100 (1400-3100)
Guatemala	2005	Female	80 to 84	7.3% (4.3-11%)	93% (89-96%)	1.1 (0.55-1.9)	3.7% (2.5-5.2%)	4.7% (3.2-6.7%)	120 (75-170)	1200 (810-1800)
Guatemala	2005	Female	85 to 89	7.6% (4.3-12%)	92% (88-96%)	1.1 (0.54-2.1)	3.4% (2.2-5.2%)	4.3% (2.7-6.5%)	80 (50-120)	600 (380-920)
Guatemala	2005	Female	90 to 94	8.9% (5.5-14%)	91% (86-95%)	1.2 (0.56-2.2)	3.5% (2-5.5%)	4.2% (2.4-6.7%)	43 (25-70)	220 (130-360)
Guatemala	2005	Female	95 plus	11% (6-17%)	89% (83-94%)	1.3 (0.48-2.6)	3.6% (1.6-5.9%)	4.2% (1.8-7.1%)	14 (5.9-24)	49 (22-82)
Guatemala	2005	Male	All Ages	12% (10-14%)	88% (86-90%)	1.4 (1.1-1.7)	7.9% (6.6-9.4%)	9.9% (8.6-11%)	4300 (3400-5500)	200000 (160000-250000)
Guatemala	2005	Male	15 to 19	9.9% (6.5-14%)	90% (86-93%)	0.62 (0.31-1)	4.3% (2.9-6.3%)	4.6% (2.7-7.4%)	77 (44-130)	7100 (4500-11000)
Guatemala	2005	Male	20 to 24	26% (19-33%)	74% (67-81%)	2.4 (1.4-3.7)	13% (9.5-18%)	14% (9.5-21%)	340 (210-510)	27000 (18000-38000)
Guatemala	2005	Male	25 to 29	31% (24-38%)	69% (62-76%)	3.4 (2.1-5)	17% (13-21%)	19% (14-25%)	420 (290-590)	30000 (22000-40000)
Guatemala	2005	Male	30 to 34	26% (19-34%)	74% (66-81%)	2.9 (1.7-4.3)	17% (14-21%)	19% (15-24%)	380 (270-520)	25000 (19000-33000)
Guatemala	2005	Male	35 to 39	24% (18-31%)	76% (69-82%)	2.8 (1.6-4.5)	19% (16-22%)	22% (18-26%)	410 (300-560)	24000 (18000-31000)
Guatemala	2005	Male	40 to 44	26% (20-33%)	74% (67-80%)	3.1 (1.8-4.9)	20% (17-23%)	24% (21-27%)	440 (340-570)	22000 (17000-28000)
Guatemala	2005	Male	45 to 49	27% (19-34%)	73% (66-81%)	3.2 (1.8-5.1)	19% (16-22%)	23% (20-27%)	430 (320-550)	19000 (15000-24000)
Guatemala	2005	Male	50 to 54	26% (19-33%)	74% (67-81%)	3.3 (1.9-5.3)	17% (14-19%)	21% (18-24%)	380 (290-500)	15000 (12000-19000)
Guatemala	2005	Male	55 to 59	23% (17-30%)	77% (70-83%)	3.4 (2-5.4)	14% (12-16%)	18% (15-20%)	330 (260-430)	11000 (9100-14000)
Guatemala	2005	Male	60 to 64	19% (13-26%)	81% (74-87%)	2.6 (1.4-4.5)	11% (9.1-13%)	14% (12-16%)	260 (210-330)	7500 (6000-9400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	2005	Male	65 to 69	16% (11-22%)	84% (78-89%)	1.9 (1.1-3.1)	8.6% (7.2-10%)	11% (9.2-13%)	240 (190-310)	5700 (4500-7200)
Guatemala	2005	Male	70 to 74	16% (11-22%)	84% (78-89%)	1.6 (0.86-2.7)	6.9% (5.5-8.4%)	8.6% (6.9-10%)	220 (160-290)	4200 (3200-5400)
Guatemala	2005	Male	75 to 79	14% (8.5-20%)	86% (80-91%)	1.2 (0.58-2.2)	5.1% (3.8-6.6%)	6.2% (4.6-7.9%)	180 (120-240)	2600 (1800-3500)
Guatemala	2005	Male	80 to 84	11% (6.3-16%)	89% (84-94%)	0.86 (0.42-1.5)	3.6% (2.5-4.9%)	4.2% (3-5.7%)	110 (74-150)	1200 (810-1600)
Guatemala	2005	Male	85 to 89	9.9% (5.7-15%)	90% (85-94%)	0.72 (0.35-1.3)	2.8% (1.9-4.1%)	3.2% (2.1-4.7%)	57 (37-83)	440 (290-640)
Guatemala	2005	Male	90 to 94	11% (6.6-18%)	89% (82-93%)	0.79 (0.35-1.6)	2.6% (1.5-4.1%)	2.9% (1.6-4.7%)	21 (12-35)	120 (67-190)
Guatemala	2005	Male	95 plus	14% (7-24%)	86% (76-93%)	0.93 (0.35-1.9)	2.6% (1.3-4.4%)	2.8% (1.2-4.9%)	5.5 (2.3-10)	21 (10-35)
Guatemala	2010	Female	All Ages	8.7% (7.3-10%)	91% (90-93%)	0.91 (0.68-1.1)	3.2% (2.6-4%)	5.6% (4.6-6.8%)	1900 (1400-2400)	63000 (48000-79000)
Guatemala	2010	Female	15 to 19	7.8% (5-11%)	92% (89-95%)	0.72 (0.37-1.2)	2.2% (1.3-3.3%)	3.6% (1.8-5.8%)	26 (14-44)	2500 (1400-3800)
Guatemala	2010	Female	20 to 24	12% (7.3-17%)	88% (83-93%)	1.1 (0.57-1.8)	3.5% (2.3-4.9%)	5.2% (3-7.7%)	39 (21-62)	3700 (2300-5400)
Guatemala	2010	Female	25 to 29	13% (8.1-19%)	87% (81-92%)	1.2 (0.66-2)	4% (2.9-5.4%)	6% (3.9-8.5%)	49 (29-74)	4300 (2900-5900)
Guatemala	2010	Female	30 to 34	13% (8.4-20%)	87% (80-92%)	1.3 (0.71-2)	4.7% (3.6-6.2%)	7.5% (5.4-9.8%)	63 (42-91)	4800 (3400-6400)
Guatemala	2010	Female	35 to 39	14% (8.4-20%)	86% (80-92%)	1.5 (0.76-2.5)	5.5% (4.1-7.2%)	9% (6.8-11%)	79 (53-110)	5100 (3700-6900)
Guatemala	2010	Female	40 to 44	14% (9.2-20%)	86% (80-91%)	1.8 (1-2.9)	6.3% (4.8-8.3%)	11% (8.3-13%)	99 (71-130)	5500 (4000-7400)
Guatemala	2010	Female	45 to 49	15% (8.7-21%)	85% (79-91%)	2.2 (1.1-3.8)	7% (5.3-9.1%)	11% (8.7-14%)	130 (93-170)	6000 (4300-8100)
Guatemala	2010	Female	50 to 54	14% (9.5-21%)	86% (79-90%)	2.5 (1.4-4.2)	7.4% (5.5-9.8%)	11% (8.5-14%)	160 (110-220)	6400 (4500-8800)
Guatemala	2010	Female	55 to 59	13% (8.5-19%)	87% (81-92%)	2.6 (1.4-4.5)	7.3% (5.3-9.8%)	10% (8-14%)	180 (130-250)	6300 (4400-8800)
Guatemala	2010	Female	60 to 64	12% (7-17%)	88% (83-93%)	2.3 (1.1-4.3)	6.6% (4.8-9%)	9.2% (6.9-12%)	180 (130-250)	5300 (3700-7200)
Guatemala	2010	Female	65 to 69	10% (6.5-15%)	90% (85-93%)	2 (1-3.4)	6.3% (4.7-8.6%)	8.6% (6.6-11%)	180 (130-260)	4300 (3100-6100)
Guatemala	2010	Female	70 to 74	9.3% (6-14%)	91% (86-94%)	1.8 (0.88-3.1)	5.5% (4.1-7.5%)	7.5% (5.6-10%)	180 (130-260)	3500 (2400-5000)
Guatemala	2010	Female	75 to 79	8.2% (4.7-13%)	92% (87-95%)	1.4 (0.65-2.6)	4.5% (3.1-6.3%)	5.9% (4.3-8.2%)	170 (120-240)	2500 (1700-3600)
Guatemala	2010	Female	80 to 84	7.3% (4.3-11%)	93% (89-96%)	1.2 (0.57-2.1)	3.7% (2.5-5.4%)	4.8% (3.2-6.8%)	150 (98-220)	1600 (1100-2400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	2010	Female	85 to 89	7.4% (4.1-12%)	93% (88-96%)	1.1 (0.55-2.1)	3.4% (2.2-5.1%)	4.2% (2.7-6.4%)	110 (66-160)	800 (500-1200)
Guatemala	2010	Female	90 to 94	8.7% (5.5-14%)	91% (86-95%)	1.2 (0.56-2.3)	3.4% (2-5.3%)	4.1% (2.4-6.5%)	55 (32-90)	290 (170-460)
Guatemala	2010	Female	95 plus	11% (6.1-17%)	89% (83-94%)	1.4 (0.51-2.7)	3.5% (1.6-5.7%)	4.1% (1.8-6.9%)	17 (7.2-28)	58 (26-95)
Guatemala	2010	Male	All Ages	11% (8.8-13%)	89% (87-91%)	1.3 (1-1.6)	8% (6.8-9.4%)	9.7% (8.3-11%)	4300 (3300-5400)	200000 (160000-250000)
Guatemala	2010	Male	15 to 19	8.6% (5.2-13%)	91% (87-95%)	0.56 (0.27-1)	4.1% (2.8-6%)	4.3% (2.5-7%)	75 (41-120)	7200 (4700-11000)
Guatemala	2010	Male	20 to 24	18% (11-26%)	82% (74-89%)	1.7 (0.93-2.8)	11% (7.6-15%)	11% (6.9-17%)	260 (150-400)	22000 (14000-31000)
Guatemala	2010	Male	25 to 29	23% (15-33%)	77% (67-85%)	2.6 (1.4-4.2)	14% (11-19%)	15% (11-21%)	370 (240-540)	28000 (20000-39000)
Guatemala	2010	Male	30 to 34	25% (16-35%)	75% (65-84%)	2.9 (1.6-4.6)	16% (13-20%)	18% (14-24%)	410 (290-560)	27000 (20000-36000)
Guatemala	2010	Male	35 to 39	24% (15-34%)	76% (66-85%)	3 (1.6-4.9)	18% (15-22%)	21% (17-26%)	420 (300-560)	25000 (18000-32000)
Guatemala	2010	Male	40 to 44	24% (16-33%)	76% (67-84%)	3 (1.6-4.9)	19% (16-22%)	23% (19-27%)	430 (310-550)	22000 (17000-27000)
Guatemala	2010	Male	45 to 49	25% (15-34%)	75% (66-85%)	3.1 (1.6-5.2)	18% (15-21%)	22% (19-25%)	420 (310-550)	19000 (14000-24000)
Guatemala	2010	Male	50 to 54	24% (16-33%)	76% (67-84%)	3.2 (1.7-5.5)	16% (13-19%)	20% (17-23%)	400 (300-510)	16000 (12000-20000)
Guatemala	2010	Male	55 to 59	22% (14-31%)	78% (69-86%)	3.3 (1.7-5.6)	13% (11-16%)	16% (14-19%)	350 (260-460)	12000 (9200-15000)
Guatemala	2010	Male	60 to 64	19% (12-28%)	81% (72-88%)	2.8 (1.4-5.1)	11% (8.6-13%)	13% (11-16%)	300 (230-390)	8800 (6600-11000)
Guatemala	2010	Male	65 to 69	17% (11-24%)	83% (76-89%)	2.1 (1.1-3.6)	8.7% (7.2-11%)	11% (9.2-13%)	250 (190-320)	6100 (4600-7700)
Guatemala	2010	Male	70 to 74	15% (9.5-21%)	85% (79-91%)	1.5 (0.78-2.6)	6.7% (5.2-8.3%)	8.3% (6.7-10%)	220 (170-290)	4300 (3200-5600)
Guatemala	2010	Male	75 to 79	12% (6.7-18%)	88% (82-93%)	1.1 (0.5-2)	4.8% (3.6-6.2%)	5.9% (4.4-7.5%)	180 (130-240)	2700 (1900-3600)
Guatemala	2010	Male	80 to 84	9.8% (5.5-15%)	90% (85-94%)	0.83 (0.39-1.5)	3.5% (2.6-4.7%)	4.2% (3.1-5.6%)	130 (91-180)	1400 (1000-1900)
Guatemala	2010	Male	85 to 89	9.4% (5.3-15%)	91% (85-95%)	0.73 (0.35-1.3)	2.8% (1.9-4%)	3.2% (2.1-4.7%)	75 (49-110)	590 (390-840)
Guatemala	2010	Male	90 to 94	11% (6.4-17%)	89% (83-94%)	0.8 (0.36-1.5)	2.6% (1.5-4%)	2.9% (1.7-4.5%)	32 (18-50)	170 (100-270)
Guatemala	2010	Male	95 plus	14% (6.7-23%)	86% (77-93%)	0.96 (0.35-1.9)	2.7% (1.4-4.4%)	2.9% (1.4-5%)	8.1 (3.7-14)	30 (15-51)
Guatemala	2016	Female	All Ages	8.9% (7.4-10%)	91% (90-93%)	1 (0.73-1.4)	3.6% (2.7-4.6%)	5.9% (4.8-7.3%)	2200 (1400-3100)	70000 (49000-98000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	2016	Female	15 to 19	7.8% (4.9-11%)	92% (89-95%)	0.79 (0.39-1.4)	2.2% (1.3-3.4%)	3.7% (1.9-6.1%)	28 (12-50)	2600 (1400-4200)
Guatemala	2016	Female	20 to 24	11% (7.3-16%)	89% (84-93%)	1.2 (0.58-2.1)	3.4% (2.3-4.8%)	5.3% (3-7.8%)	40 (21-68)	4000 (2500-5900)
Guatemala	2016	Female	25 to 29	13% (7.9-19%)	87% (81-92%)	1.3 (0.65-2.2)	3.9% (2.8-5.3%)	6.1% (4-8.7%)	48 (26-80)	4300 (2800-6300)
Guatemala	2016	Female	30 to 34	13% (8-20%)	87% (80-92%)	1.4 (0.73-2.3)	4.6% (3.4-6.2%)	7.7% (5.6-10%)	68 (38-110)	5100 (3400-7400)
Guatemala	2016	Female	35 to 39	14% (8.4-20%)	86% (80-92%)	1.6 (0.81-2.7)	5.4% (3.9-7.3%)	9.4% (7.1-12%)	91 (51-150)	5900 (3700-8900)
Guatemala	2016	Female	40 to 44	14% (8.9-20%)	86% (80-91%)	1.9 (1-3.2)	6.2% (4.6-8.3%)	11% (8.5-13%)	120 (69-180)	6400 (4100-9600)
Guatemala	2016	Female	45 to 49	14% (8.4-20%)	86% (80-92%)	2.4 (1.2-4.2)	6.7% (4.9-9.1%)	11% (8.5-14%)	140 (86-220)	6800 (4300-10000)
Guatemala	2016	Female	50 to 54	14% (9.1-20%)	86% (80-91%)	2.7 (1.4-4.6)	7.2% (5.2-9.8%)	11% (8.3-14%)	180 (110-270)	7200 (4500-11000)
Guatemala	2016	Female	55 to 59	13% (8.5-19%)	87% (81-91%)	2.8 (1.4-5)	7.1% (5.1-9.8%)	10% (7.8-14%)	200 (120-310)	6800 (4200-10000)
Guatemala	2016	Female	60 to 64	11% (6.9-17%)	89% (83-93%)	2.5 (1.2-4.7)	6.4% (4.6-8.9%)	9.2% (6.9-12%)	210 (130-320)	6000 (3800-9000)
Guatemala	2016	Female	65 to 69	10% (6.4-14%)	90% (86-94%)	2.2 (1.1-3.8)	6.1% (4.5-8.3%)	8.6% (6.5-11%)	230 (140-340)	5400 (3400-7900)
Guatemala	2016	Female	70 to 74	9.2% (6-13%)	91% (87-94%)	1.9 (0.95-3.5)	5.4% (3.9-7.4%)	7.4% (5.5-10%)	200 (130-300)	3900 (2500-5800)
Guatemala	2016	Female	75 to 79	8% (4.6-12%)	92% (88-95%)	1.6 (0.66-2.9)	4.3% (3-6.2%)	5.8% (4.2-8.1%)	180 (100-270)	2600 (1600-4000)
Guatemala	2016	Female	80 to 84	7.1% (4.1-10%)	93% (90-96%)	1.3 (0.59-2.3)	3.6% (2.5-5.2%)	4.7% (3.2-6.7%)	180 (110-270)	1900 (1200-2900)
Guatemala	2016	Female	85 to 89	7.2% (4.1-11%)	93% (89-96%)	1.2 (0.55-2.2)	3.3% (2.1-5%)	4.2% (2.7-6.3%)	140 (83-220)	1100 (650-1700)
Guatemala	2016	Female	90 to 94	8.4% (5.2-13%)	92% (87-95%)	1.3 (0.57-2.5)	3.3% (2-5.2%)	4.1% (2.4-6.5%)	80 (45-130)	420 (240-680)
Guatemala	2016	Female	95 plus	10% (5.7-16%)	90% (84-94%)	1.5 (0.54-3)	3.5% (1.7-5.6%)	4.2% (2-6.9%)	27 (12-47)	93 (45-160)
Guatemala	2016	Male	All Ages	11% (9.2-14%)	89% (86-91%)	1.5 (1-2)	8.8% (7.3-10%)	10% (8.7-12%)	4900 (3400-6800)	23000 (16000-31000)
Guatemala	2016	Male	15 to 19	8.3% (5.1-13%)	92% (87-95%)	0.59 (0.27-1.1)	4.2% (2.8-6.2%)	4.5% (2.6-7.3%)	78 (38-140)	7600 (4500-12000)
Guatemala	2016	Male	20 to 24	18% (11-25%)	82% (75-89%)	1.9 (0.95-3.2)	11% (7.6-15%)	11% (7-17%)	270 (140-450)	23000 (15000-36000)
Guatemala	2016	Male	25 to 29	23% (15-32%)	77% (68-85%)	2.7 (1.5-4.6)	14% (11-19%)	15% (11-21%)	360 (220-560)	28000 (19000-40000)
Guatemala	2016	Male	30 to 34	24% (15-35%)	76% (65-85%)	3.1 (1.6-5.1)	16% (13-20%)	19% (14-24%)	450 (300-670)	31000 (22000-43000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guatemala	2016	Male	35 to 39	24% (15-33%)	76% (67-85%)	3.1 (1.6-5.2)	18% (15-22%)	22% (17-26%)	500 (330-750)	29000 (21000-41000)
Guatemala	2016	Male	40 to 44	24% (15-33%)	76% (67-85%)	3.2 (1.6-5.3)	18% (15-22%)	23% (19-27%)	510 (320-730)	26000 (18000-36000)
Guatemala	2016	Male	45 to 49	24% (15-34%)	76% (66-85%)	3.3 (1.7-5.8)	18% (15-21%)	22% (19-26%)	480 (310-710)	22000 (15000-30000)
Guatemala	2016	Male	50 to 54	23% (16-33%)	77% (67-84%)	3.4 (1.7-5.9)	16% (13-19%)	20% (17-23%)	450 (290-650)	18000 (12000-25000)
Guatemala	2016	Male	55 to 59	21% (14-30%)	79% (70-86%)	3.5 (1.7-6.3)	13% (11-16%)	17% (14-20%)	380 (250-530)	13000 (8900-18000)
Guatemala	2016	Male	60 to 64	19% (12-28%)	81% (72-88%)	3 (1.4-5.7)	10% (8.5-13%)	13% (11-16%)	340 (220-470)	9700 (6700-13000)
Guatemala	2016	Male	65 to 69	17% (10-24%)	83% (76-90%)	2.2 (1.1-3.9)	8.6% (7-11%)	11% (9.2-13%)	320 (210-440)	7500 (5300-10000)
Guatemala	2016	Male	70 to 74	14% (9.2-21%)	86% (79-91%)	1.6 (0.77-2.8)	6.7% (5.3-8.4%)	8.4% (6.9-10%)	260 (170-360)	5000 (3400-6800)
Guatemala	2016	Male	75 to 79	11% (6.6-18%)	89% (82-93%)	1.2 (0.49-2.2)	4.8% (3.6-6.1%)	6% (4.6-7.5%)	200 (130-280)	2900 (1900-4100)
Guatemala	2016	Male	80 to 84	9.6% (5.5-14%)	90% (86-95%)	0.88 (0.41-1.6)	3.6% (2.7-4.7%)	4.3% (3.2-5.7%)	160 (100-220)	1700 (1200-2400)
Guatemala	2016	Male	85 to 89	9.2% (5.1-15%)	91% (85-95%)	0.77 (0.34-1.4)	2.9% (2-4.1%)	3.3% (2.2-4.8%)	110 (67-160)	840 (550-1200)
Guatemala	2016	Male	90 to 94	11% (6.2-17%)	89% (83-94%)	0.85 (0.37-1.7)	2.6% (1.6-3.9%)	2.9% (1.8-4.5%)	53 (31-83)	290 (180-440)
Guatemala	2016	Male	95 plus	13% (6.4-22%)	87% (78-94%)	1 (0.37-2.1)	2.7% (1.5-4.4%)	2.9% (1.4-5%)	15 (7.1-25)	55 (29-90)
Honduras	1990	Female	All Ages	5.4% (4.5-6.6%)	95% (93-96%)	0.19 (0.16-0.23)	1.2% (1-1.5%)	2.3% (1.8-2.9%)	370 (280-480)	11000 (9000-14000)
Honduras	1990	Female	15 to 19	5.5% (3.4-8.1%)	95% (92-97%)	0.19 (0.096-0.33)	0.87% (0.53-1.3%)	0.94% (0.44-1.7%)	2.6 (1.2-4.6)	350 (210-540)
Honduras	1990	Female	20 to 24	8.3% (5.1-12%)	92% (88-95%)	0.28 (0.14-0.47)	1.5% (0.97-2.2%)	1.4% (0.71-2.4%)	4.5 (2.3-7.6)	600 (380-900)
Honduras	1990	Female	25 to 29	8.7% (5.4-13%)	91% (87-95%)	0.28 (0.15-0.47)	1.6% (1.1-2.2%)	1.6% (0.92-2.6%)	5.2 (2.9-8.4)	590 (400-850)
Honduras	1990	Female	30 to 34	9% (5.4-14%)	91% (86-95%)	0.3 (0.16-0.49)	1.9% (1.4-2.5%)	2.3% (1.6-3.2%)	8 (5.3-11)	650 (460-880)
Honduras	1990	Female	35 to 39	9.5% (5.6-14%)	91% (86-94%)	0.35 (0.18-0.59)	2.3% (1.7-3.1%)	3% (2.1-4.2%)	12 (8-18)	770 (540-1100)
Honduras	1990	Female	40 to 44	9.6% (6-14%)	90% (86-94%)	0.43 (0.23-0.69)	2.7% (1.9-3.7%)	3.7% (2.5-5.2%)	15 (9.6-21)	780 (540-1100)
Honduras	1990	Female	45 to 49	9.6% (5.5-14%)	90% (86-94%)	0.52 (0.25-0.91)	3.4% (2.3-4.9%)	4.7% (3.1-6.8%)	20 (13-30)	900 (590-1300)
Honduras	1990	Female	50 to 54	9.7% (6.1-14%)	90% (86-94%)	0.62 (0.32-1)	4% (2.8-5.6%)	5.3% (3.6-7.3%)	29 (19-42)	1100 (740-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	1990	Female	55 to 59	9.7% (5.6-16%)	90% (84-94%)	0.69 (0.32-1.2)	4.5% (3.1-6.2%)	5.8% (4-8.1%)	41 (28-58)	1300 (900-1900)
Honduras	1990	Female	60 to 64	8.8% (4.8-15%)	91% (85-95%)	0.63 (0.3-1.2)	4% (2.7-5.7%)	5% (3.4-7%)	44 (29-62)	1200 (810-1700)
Honduras	1990	Female	65 to 69	8.1% (4.6-13%)	92% (87-95%)	0.56 (0.27-1)	3.9% (2.6-5.4%)	4.7% (3.2-6.6%)	52 (35-73)	1200 (790-1700)
Honduras	1990	Female	70 to 74	7.1% (4.3-12%)	93% (88-96%)	0.48 (0.24-0.86)	3.3% (2.2-4.7%)	3.9% (2.7-5.5%)	45 (29-64)	810 (540-1200)
Honduras	1990	Female	75 to 79	5.8% (3.3-9.7%)	94% (90-97%)	0.38 (0.16-0.71)	2.5% (1.6-3.7%)	2.9% (1.9-4.4%)	39 (25-58)	550 (360-820)
Honduras	1990	Female	80 to 84	5.1% (2.8-8.2%)	95% (92-97%)	0.32 (0.14-0.57)	2% (1.3-3%)	2.3% (1.5-3.5%)	25 (16-39)	260 (170-400)
Honduras	1990	Female	85 to 89	5.3% (2.9-8.7%)	95% (91-97%)	0.32 (0.14-0.61)	1.7% (1.1-2.7%)	2% (1.2-3.2%)	17 (10-27)	130 (77-200)
Honduras	1990	Female	90 to 94	6.4% (3.6-10%)	94% (90-96%)	0.35 (0.15-0.7)	1.4% (0.8-2.5%)	1.6% (0.88-2.9%)	8.1 (4.5-15)	43 (24-75)
Honduras	1990	Female	95 plus	8% (4.3-13%)	92% (87-96%)	0.38 (0.14-0.78)	1.5% (0.78-2.9%)	1.7% (0.82-3.2%)	3.3 (1.6-6.2)	12 (5.9-21)
Honduras	1990	Male	All Ages	16% (13-19%)	84% (81-87%)	1.1 (0.98-1.3)	3.7% (2.9-4.7%)	5.5% (4.3-6.7%)	830 (580-1200)	37000 (27000-50000)
Honduras	1990	Male	15 to 19	14% (9-20%)	86% (80-91%)	0.61 (0.3-1.1)	3.4% (2.1-5.2%)	3.8% (2-6.6%)	13 (6.1-24)	1400 (850-2200)
Honduras	1990	Male	20 to 24	29% (20-39%)	71% (61-80%)	1.7 (0.93-2.8)	8.9% (5.8-13%)	10% (5.8-16%)	48 (25-82)	4200 (2600-6500)
Honduras	1990	Male	25 to 29	34% (24-45%)	66% (55-76%)	2.4 (1.3-3.7)	11% (7.5-15%)	13% (8-19%)	60 (32-100)	4600 (2900-7000)
Honduras	1990	Male	30 to 34	35% (25-47%)	65% (53-75%)	2.7 (1.5-4.1)	12% (8.5-16%)	15% (9.9-20%)	66 (39-100)	4400 (2800-6400)
Honduras	1990	Male	35 to 39	34% (24-44%)	66% (56-76%)	2.8 (1.5-4.6)	12% (8.9-17%)	16% (11-21%)	70 (40-110)	4100 (2600-6100)
Honduras	1990	Male	40 to 44	33% (23-43%)	67% (57-77%)	3 (1.5-4.8)	13% (9.4-17%)	16% (12-22%)	70 (43-110)	3600 (2400-5200)
Honduras	1990	Male	45 to 49	33% (22-43%)	67% (57-78%)	3.2 (1.5-5.6)	13% (9.4-16%)	16% (12-21%)	72 (45-110)	3300 (2100-4800)
Honduras	1990	Male	50 to 54	31% (22-41%)	69% (59-78%)	3.3 (1.7-5.5)	12% (8.7-15%)	15% (11-19%)	79 (50-110)	3100 (2100-4400)
Honduras	1990	Male	55 to 59	27% (19-36%)	73% (64-81%)	3.2 (1.6-5.5)	9.9% (7.3-13%)	12% (9-16%)	78 (52-120)	2700 (1800-3900)
Honduras	1990	Male	60 to 64	25% (17-35%)	75% (65-83%)	2.7 (1.3-4.9)	8% (5.9-11%)	9.7% (7.1-13%)	70 (46-100)	2000 (1400-2900)
Honduras	1990	Male	65 to 69	24% (14-36%)	76% (64-86%)	2.1 (1-3.6)	6.9% (4.8-9.4%)	8.4% (5.9-12%)	67 (43-100)	1600 (1100-2300)
Honduras	1990	Male	70 to 74	22% (13-34%)	78% (66-87%)	1.5 (0.71-2.9)	5.3% (3.4-7.7%)	6.3% (4.1-9.1%)	51 (30-77)	960 (590-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	1990	Male	75 to 79	18% (11-28%)	82% (72-89%)	1.1 (0.45-2.2)	3.9% (2.2-5.8%)	4.6% (2.6-6.9%)	39 (21-62)	580 (320-900)
Honduras	1990	Male	80 to 84	16% (8.9-25%)	84% (75-91%)	0.95 (0.4-1.7)	3.3% (1.9-5%)	3.8% (2.1-5.9%)	27 (15-42)	290 (170-450)
Honduras	1990	Male	85 to 89	15% (8.9-24%)	85% (76-91%)	0.92 (0.37-1.8)	2.8% (1.4-4.4%)	3.2% (1.6-5.1%)	16 (7.7-26)	120 (63-200)
Honduras	1990	Male	90 to 94	18% (11-27%)	82% (73-89%)	1.1 (0.42-2.2)	2.7% (1.1-4.5%)	3% (1.1-5.2%)	7.1 (2.8-13)	39 (17-66)
Honduras	1990	Male	95 plus	22% (12-36%)	78% (64-88%)	1.3 (0.41-2.8)	3% (1.1-5.3%)	3.3% (1-6.1%)	2.1 (0.64-3.9)	7.8 (2.8-14)
Honduras	1995	Female	All Ages	5.6% (4.6-6.9%)	94% (93-95%)	0.21 (0.17-0.26)	1.4% (1.2-1.7%)	2.6% (2-3.3%)	420 (310-550)	13000 (9700-16000)
Honduras	1995	Female	15 to 19	5.5% (3.5-8.2%)	94% (92-96%)	0.2 (0.1-0.36)	0.91% (0.57-1.4%)	1% (0.5-1.8%)	2.8 (1.4-5)	400 (240-620)
Honduras	1995	Female	20 to 24	8.4% (5.2-12%)	92% (88-95%)	0.3 (0.15-0.52)	1.6% (1-2.3%)	1.5% (0.78-2.6%)	4.7 (2.4-8.1)	650 (410-960)
Honduras	1995	Female	25 to 29	8.8% (5.5-13%)	91% (87-94%)	0.31 (0.16-0.51)	1.7% (1.2-2.2%)	1.7% (1-2.7%)	5.6 (3.2-9.3)	660 (440-940)
Honduras	1995	Female	30 to 34	9% (5.4-14%)	91% (86-95%)	0.32 (0.17-0.53)	2% (1.4-2.6%)	2.4% (1.7-3.3%)	8.7 (5.7-13)	730 (520-1000)
Honduras	1995	Female	35 to 39	9.6% (5.7-14%)	90% (86-94%)	0.38 (0.19-0.64)	2.3% (1.7-3.2%)	3.2% (2.3-4.4%)	13 (8.4-19)	840 (580-1200)
Honduras	1995	Female	40 to 44	9.8% (6.1-14%)	90% (86-94%)	0.46 (0.24-0.74)	2.7% (1.9-3.7%)	3.8% (2.6-5.2%)	17 (11-26)	940 (640-1400)
Honduras	1995	Female	45 to 49	9.8% (5.7-14%)	90% (86-94%)	0.56 (0.26-0.98)	3.4% (2.3-4.9%)	4.7% (3.2-6.8%)	23 (15-35)	1000 (670-1500)
Honduras	1995	Female	50 to 54	9.8% (6.2-14%)	90% (86-94%)	0.66 (0.35-1.1)	4% (2.8-5.6%)	5.3% (3.6-7.4%)	30 (20-45)	1200 (750-1700)
Honduras	1995	Female	55 to 59	9.8% (5.6-16%)	90% (84-94%)	0.74 (0.34-1.3)	4.5% (3.1-6.2%)	5.9% (4.1-8%)	42 (28-62)	1400 (930-2000)
Honduras	1995	Female	60 to 64	8.9% (4.8-15%)	91% (85-95%)	0.68 (0.32-1.3)	4% (2.8-5.7%)	5.1% (3.5-7.1%)	47 (31-68)	1300 (870-1900)
Honduras	1995	Female	65 to 69	8.1% (4.6-13%)	92% (87-95%)	0.6 (0.3-1.1)	3.9% (2.7-5.5%)	4.8% (3.4-6.7%)	56 (37-80)	1300 (860-1800)
Honduras	1995	Female	70 to 74	7.3% (4.3-12%)	93% (88-96%)	0.52 (0.25-0.92)	3.4% (2.3-4.9%)	4.1% (2.7-5.8%)	58 (38-84)	1100 (700-1500)
Honduras	1995	Female	75 to 79	5.9% (3.3-10%)	94% (90-97%)	0.42 (0.18-0.77)	2.6% (1.7-3.8%)	3.1% (2-4.6%)	43 (28-64)	610 (400-910)
Honduras	1995	Female	80 to 84	5.1% (2.8-8.4%)	95% (92-97%)	0.34 (0.15-0.63)	2.1% (1.4-3.2%)	2.5% (1.6-3.9%)	32 (20-51)	340 (210-520)
Honduras	1995	Female	85 to 89	5.3% (3-8.9%)	95% (91-97%)	0.34 (0.15-0.63)	1.8% (1.1-2.8%)	2.1% (1.3-3.3%)	23 (14-36)	170 (110-270)
Honduras	1995	Female	90 to 94	6.4% (3.7-10%)	94% (90-96%)	0.37 (0.16-0.73)	1.5% (0.85-2.6%)	1.8% (0.95-3.1%)	10 (5.5-18)	54 (30-94)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	1995	Female	95 plus	8% (4.3-14%)	92% (86-96%)	0.41 (0.15-0.84)	1.7% (0.8-3%)	1.8% (0.85-3.4%)	4.5 (2.1-8.3)	16 (7.8-29)
Honduras	1995	Male	All Ages	17% (14-20%)	83% (80-86%)	1.3 (1.1-1.4)	4.4% (3.4-5.7%)	6.2% (4.9-7.7%)	990 (660-1400)	43000 (30000-60000)
Honduras	1995	Male	15 to 19	14% (9.3-21%)	86% (79-91%)	0.66 (0.32-1.2)	3.5% (2.2-5.6%)	4.1% (2-7.1%)	16 (7.1-30)	1700 (980-2800)
Honduras	1995	Male	20 to 24	29% (20-39%)	71% (61-80%)	1.8 (0.97-2.9)	9.1% (6-13%)	10% (6-16%)	52 (25-95)	4600 (2700-7400)
Honduras	1995	Male	25 to 29	34% (24-45%)	66% (55-76%)	2.5 (1.4-4.1)	11% (7.4-15%)	13% (7.9-19%)	64 (33-120)	5000 (3000-8100)
Honduras	1995	Male	30 to 34	35% (24-48%)	65% (52-76%)	2.8 (1.6-4.3)	11% (8-15%)	14% (9.5-20%)	68 (36-110)	4700 (2900-7200)
Honduras	1995	Male	35 to 39	34% (24-45%)	66% (55-76%)	3 (1.6-5)	12% (8.6-16%)	15% (10-21%)	76 (41-120)	4500 (2700-7000)
Honduras	1995	Male	40 to 44	34% (24-44%)	66% (56-76%)	3.2 (1.7-5.1)	13% (9.2-16%)	16% (12-21%)	84 (47-130)	4400 (2700-6500)
Honduras	1995	Male	45 to 49	33% (22-44%)	67% (56-78%)	3.4 (1.7-6)	13% (9.4-16%)	16% (12-20%)	86 (50-130)	3900 (2400-5600)
Honduras	1995	Male	50 to 54	31% (22-41%)	69% (59-78%)	3.5 (1.9-5.9)	12% (8.8-15%)	15% (11-19%)	87 (53-130)	3500 (2200-5000)
Honduras	1995	Male	55 to 59	28% (19-37%)	72% (63-81%)	3.4 (1.7-6)	10% (7.4-13%)	12% (9.1-16%)	88 (56-130)	3000 (2000-4400)
Honduras	1995	Male	60 to 64	25% (17-36%)	75% (64-83%)	2.9 (1.3-5.2)	8.2% (6.1-11%)	9.8% (7.3-13%)	87 (56-130)	2500 (1700-3800)
Honduras	1995	Male	65 to 69	25% (15-37%)	75% (63-85%)	2.2 (1.1-3.9)	7.1% (5.1-9.7%)	8.5% (6-12%)	85 (53-130)	2000 (1300-3000)
Honduras	1995	Male	70 to 74	22% (14-34%)	78% (66-86%)	1.6 (0.77-3.1)	5.5% (3.5-7.8%)	6.5% (4.1-9.3%)	73 (42-120)	1400 (820-2100)
Honduras	1995	Male	75 to 79	18% (11-29%)	82% (71-89%)	1.2 (0.49-2.4)	4.1% (2.3-6.1%)	4.8% (2.7-7%)	48 (25-78)	710 (380-1100)
Honduras	1995	Male	80 to 84	16% (9.2-25%)	84% (75-91%)	1 (0.44-1.9)	3.4% (2-5.2%)	3.9% (2.2-6.1%)	35 (19-57)	390 (220-610)
Honduras	1995	Male	85 to 89	15% (9-24%)	85% (76-91%)	0.97 (0.4-1.9)	2.9% (1.5-4.7%)	3.3% (1.7-5.4%)	24 (12-41)	190 (94-310)
Honduras	1995	Male	90 to 94	18% (11-28%)	82% (72-89%)	1.1 (0.45-2.3)	2.8% (1.2-4.7%)	3.2% (1.3-5.4%)	10 (4.2-18)	57 (25-96)
Honduras	1995	Male	95 plus	22% (13-37%)	78% (63-87%)	1.4 (0.43-3)	3.2% (1.2-5.6%)	3.5% (1.1-6.4%)	3.3 (0.97-6)	12 (4.3-22)
Honduras	2000	Female	All Ages	5.7% (4.7-6.9%)	94% (93-95%)	0.24 (0.2-0.29)	1.7% (1.3-2%)	2.9% (2.3-3.6%)	500 (350-670)	15000 (11000-19000)
Honduras	2000	Female	15 to 19	5.5% (3.5-8.2%)	94% (92-97%)	0.22 (0.11-0.4)	0.96% (0.61-1.5%)	1.1% (0.56-2.1%)	3.2 (1.4-5.8)	450 (270-700)
Honduras	2000	Female	20 to 24	8.3% (5-12%)	92% (88-95%)	0.33 (0.16-0.57)	1.6% (1.1-2.4%)	1.6% (0.83-2.7%)	5 (2.5-9)	740 (450-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	2000	Female	25 to 29	8.7% (5.4-13%)	91% (87-95%)	0.33 (0.18-0.57)	1.7% (1.2-2.3%)	1.7% (1-2.7%)	5.7 (3.1-10)	700 (450-1000)
Honduras	2000	Female	30 to 34	9% (5.5-14%)	91% (86-94%)	0.36 (0.19-0.58)	2% (1.5-2.6%)	2.5% (1.7-3.5%)	9.3 (5.5-15)	810 (550-1200)
Honduras	2000	Female	35 to 39	9.5% (5.5-14%)	91% (86-94%)	0.41 (0.21-0.68)	2.4% (1.7-3.2%)	3.3% (2.3-4.5%)	14 (8.2-23)	940 (620-1400)
Honduras	2000	Female	40 to 44	9.6% (5.7-14%)	90% (86-94%)	0.5 (0.26-0.81)	2.7% (1.9-3.8%)	3.8% (2.6-5.3%)	19 (11-30)	1000 (650-1600)
Honduras	2000	Female	45 to 49	9.6% (5.7-14%)	90% (86-94%)	0.61 (0.29-1.1)	3.4% (2.3-4.9%)	4.8% (3.3-6.9%)	28 (17-44)	1300 (810-2000)
Honduras	2000	Female	50 to 54	9.7% (6.3-14%)	90% (86-94%)	0.73 (0.38-1.2)	4% (2.7-5.7%)	5.4% (3.7-7.5%)	36 (22-55)	1400 (870-2100)
Honduras	2000	Female	55 to 59	9.8% (5.7-16%)	90% (84-94%)	0.82 (0.38-1.5)	4.5% (3.1-6.2%)	6% (4.1-8.1%)	45 (29-68)	1500 (960-2200)
Honduras	2000	Female	60 to 64	8.9% (4.7-15%)	91% (85-95%)	0.76 (0.35-1.5)	4.1% (2.9-5.7%)	5.2% (3.6-7.3%)	51 (33-75)	1400 (920-2100)
Honduras	2000	Female	65 to 69	8.1% (4.7-13%)	92% (87-95%)	0.67 (0.34-1.2)	4% (2.8-5.6%)	5% (3.5-7%)	64 (42-94)	1500 (960-2100)
Honduras	2000	Female	70 to 74	7.2% (4.3-12%)	93% (88-96%)	0.58 (0.28-1)	3.6% (2.4-5%)	4.3% (2.9-6.1%)	66 (42-96)	1200 (760-1700)
Honduras	2000	Female	75 to 79	5.9% (3.3-9.5%)	94% (90-97%)	0.46 (0.2-0.85)	2.8% (1.8-4%)	3.3% (2.2-4.8%)	59 (37-88)	840 (520-1200)
Honduras	2000	Female	80 to 84	5.1% (2.7-8.3%)	95% (92-97%)	0.38 (0.17-0.69)	2.3% (1.5-3.4%)	2.7% (1.7-4%)	46 (29-70)	490 (300-740)
Honduras	2000	Female	85 to 89	5.3% (2.9-8.7%)	95% (91-97%)	0.37 (0.17-0.7)	2% (1.3-3.1%)	2.3% (1.5-3.6%)	29 (18-46)	210 (140-340)
Honduras	2000	Female	90 to 94	6.3% (3.7-10%)	94% (90-96%)	0.41 (0.17-0.81)	1.7% (0.95-2.8%)	2% (1.1-3.4%)	14 (7.2-24)	71 (38-120)
Honduras	2000	Female	95 plus	8% (4.2-13%)	92% (87-96%)	0.45 (0.16-0.91)	1.8% (0.88-3.3%)	2% (0.94-3.8%)	5.1 (2.4-9.6)	18 (8.9-33)
Honduras	2000	Male	All Ages	17% (14-20%)	83% (80-86%)	1.5 (1.3-1.7)	5.2% (4-6.6%)	6.9% (5.5-8.5%)	1200 (760-1700)	50000 (34000-71000)
Honduras	2000	Male	15 to 19	14% (9.2-20%)	86% (80-91%)	0.75 (0.36-1.3)	3.9% (2.4-6.2%)	4.6% (2.4-8.2%)	20 (8.7-40)	2100 (1200-3500)
Honduras	2000	Male	20 to 24	29% (20-38%)	71% (62-80%)	2.1 (1.1-3.4)	9.8% (6.4-14%)	11% (6.7-18%)	64 (30-130)	5500 (3200-9700)
Honduras	2000	Male	25 to 29	34% (24-45%)	66% (55-76%)	2.8 (1.6-4.6)	11% (7.8-16%)	13% (8.5-20%)	71 (35-130)	5500 (3200-9000)
Honduras	2000	Male	30 to 34	35% (25-47%)	65% (53-75%)	3.2 (1.9-4.9)	12% (8.2-16%)	14% (9.7-20%)	76 (38-130)	5200 (3200-8300)
Honduras	2000	Male	35 to 39	34% (24-45%)	66% (55-76%)	3.4 (1.8-5.5)	12% (8.7-17%)	16% (11-21%)	85 (45-140)	5100 (3000-7900)
Honduras	2000	Male	40 to 44	34% (24-44%)	66% (56-76%)	3.6 (1.9-5.8)	13% (9.1-17%)	16% (12-21%)	95 (52-150)	5000 (3000-7500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	2000	Male	45 to 49	33% (22-43%)	67% (57-78%)	3.8 (1.9-6.6)	13% (9.5-16%)	16% (12-21%)	110 (61-160)	4900 (3100-7200)
Honduras	2000	Male	50 to 54	31% (22-41%)	69% (59-78%)	3.9 (2.1-6.5)	12% (8.9-15%)	15% (11-19%)	110 (63-160)	4200 (2700-6000)
Honduras	2000	Male	55 to 59	28% (19-37%)	72% (63-81%)	3.9 (1.9-6.8)	10% (7.5-13%)	12% (9.1-16%)	98 (60-150)	3400 (2100-4900)
Honduras	2000	Male	60 to 64	25% (17-35%)	75% (65-83%)	3.3 (1.5-5.9)	8.3% (6.1-11%)	10% (7.3-14%)	98 (60-150)	2800 (1800-4300)
Honduras	2000	Male	65 to 69	25% (15-37%)	75% (63-85%)	2.6 (1.2-4.5)	7.3% (5.2-9.8%)	8.8% (6.2-12%)	100 (63-160)	2500 (1500-3700)
Honduras	2000	Male	70 to 74	22% (14-34%)	78% (66-86%)	1.9 (0.87-3.5)	5.8% (3.8-8.1%)	6.9% (4.5-9.8%)	91 (52-150)	1700 (1000-2800)
Honduras	2000	Male	75 to 79	18% (11-30%)	82% (70-89%)	1.4 (0.56-2.7)	4.3% (2.7-6.4%)	5.1% (3-7.4%)	70 (37-120)	1000 (560-1600)
Honduras	2000	Male	80 to 84	16% (9.2-25%)	84% (75-91%)	1.1 (0.49-2.1)	3.7% (2.2-5.4%)	4.2% (2.5-6.3%)	49 (27-79)	530 (300-850)
Honduras	2000	Male	85 to 89	15% (8.9-24%)	85% (76-91%)	1.1 (0.44-2.1)	3.1% (1.6-4.9%)	3.5% (1.8-5.6%)	30 (16-49)	230 (120-370)
Honduras	2000	Male	90 to 94	18% (11-27%)	82% (73-89%)	1.3 (0.49-2.6)	3% (1.5-5%)	3.4% (1.5-5.6%)	15 (6.7-26)	83 (39-140)
Honduras	2000	Male	95 plus	22% (12-36%)	78% (64-88%)	1.6 (0.47-3.3)	3.4% (1.3-5.7%)	3.7% (1.2-6.5%)	4.6 (1.5-8.2)	17 (6.7-29)
Honduras	2005	Female	All Ages	5.8% (4.8-7%)	94% (93-95%)	0.27 (0.22-0.33)	1.9% (1.5-2.3%)	3.2% (2.6-4%)	590 (400-800)	17000 (12000-23000)
Honduras	2005	Female	15 to 19	5.4% (3.4-7.9%)	95% (92-97%)	0.24 (0.12-0.42)	0.99% (0.64-1.5%)	1.2% (0.61-2.3%)	3.4 (1.6-6.3)	490 (300-770)
Honduras	2005	Female	20 to 24	8.1% (5-12%)	92% (88-95%)	0.35 (0.18-0.61)	1.7% (1.1-2.5%)	1.8% (0.94-3.1%)	5.7 (2.7-10)	840 (520-1300)
Honduras	2005	Female	25 to 29	8.5% (5.3-13%)	91% (87-95%)	0.36 (0.18-0.59)	1.8% (1.3-2.5%)	1.9% (1.2-3%)	6.6 (3.3-12)	820 (510-1200)
Honduras	2005	Female	30 to 34	8.8% (5.3-14%)	91% (86-95%)	0.38 (0.2-0.62)	2.1% (1.5-2.8%)	2.7% (1.9-3.9%)	10 (5.6-18)	900 (580-1300)
Honduras	2005	Female	35 to 39	9.2% (5.4-14%)	91% (86-95%)	0.44 (0.23-0.72)	2.5% (1.8-3.4%)	3.6% (2.6-5%)	17 (9.1-28)	1100 (700-1700)
Honduras	2005	Female	40 to 44	9.4% (5.6-14%)	91% (86-94%)	0.53 (0.28-0.85)	2.9% (2-4%)	4.2% (2.9-5.7%)	23 (12-36)	1200 (730-1900)
Honduras	2005	Female	45 to 49	9.4% (5.4-14%)	91% (86-95%)	0.66 (0.32-1.2)	3.6% (2.5-5.2%)	5.3% (3.6-7.5%)	34 (20-54)	1500 (920-2400)
Honduras	2005	Female	50 to 54	9.5% (6.1-14%)	90% (86-94%)	0.78 (0.41-1.3)	4.1% (2.9-5.8%)	5.7% (4-7.9%)	45 (27-72)	1800 (1100-2700)
Honduras	2005	Female	55 to 59	9.6% (5.5-16%)	90% (84-94%)	0.87 (0.41-1.6)	4.6% (3.1-6.2%)	6.1% (4.3-8.2%)	53 (32-79)	1800 (1100-2600)
Honduras	2005	Female	60 to 64	8.8% (4.7-15%)	91% (85-95%)	0.82 (0.38-1.6)	4.2% (2.9-5.9%)	5.4% (3.8-7.4%)	55 (34-82)	1500 (970-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	2005	Female	65 to 69	8.1% (4.7-13%)	92% (87-95%)	0.74 (0.37-1.3)	4.2% (3-5.7%)	5.2% (3.7-7.1%)	68 (44-100)	1600 (1000-2300)
Honduras	2005	Female	70 to 74	7.2% (4.4-12%)	93% (88-96%)	0.64 (0.31-1.1)	3.7% (2.5-5.1%)	4.5% (3.1-6.3%)	76 (46-110)	1400 (850-2100)
Honduras	2005	Female	75 to 79	5.9% (3.3-9.8%)	94% (90-97%)	0.51 (0.22-0.93)	2.9% (1.9-4.1%)	3.5% (2.3-4.9%)	69 (41-100)	970 (600-1400)
Honduras	2005	Female	80 to 84	5.1% (2.7-8.2%)	95% (92-97%)	0.42 (0.18-0.75)	2.4% (1.6-3.6%)	2.9% (1.9-4.3%)	65 (41-98)	680 (430-1000)
Honduras	2005	Female	85 to 89	5.3% (2.8-8.5%)	95% (91-97%)	0.41 (0.18-0.77)	2.1% (1.3-3.3%)	2.5% (1.5-3.9%)	38 (23-60)	280 (170-450)
Honduras	2005	Female	90 to 94	6.3% (3.6-10%)	94% (90-96%)	0.44 (0.19-0.88)	1.9% (1-3.1%)	2.2% (1.2-3.7%)	15 (8.1-27)	80 (44-140)
Honduras	2005	Female	95 plus	7.8% (4.2-13%)	92% (87-96%)	0.49 (0.18-0.99)	2% (1-3.5%)	2.2% (1.1-4.1%)	6 (2.9-11)	21 (11-38)
Honduras	2005	Male	All Ages	18% (15-21%)	82% (79-85%)	1.7 (1.5-1.9)	5.9% (4.6-7.6%)	7.6% (6.1-9.3%)	1400 (890-2100)	58000 (39000-84000)
Honduras	2005	Male	15 to 19	14% (9-20%)	86% (80-91%)	0.8 (0.38-1.4)	4% (2.4-6.3%)	4.9% (2.5-8.5%)	23 (9.5-49)	2400 (1300-4300)
Honduras	2005	Male	20 to 24	29% (20-38%)	71% (62-80%)	2.2 (1.2-3.6)	10% (6.7-15%)	12% (7.2-18%)	74 (36-140)	6400 (3600-11000)
Honduras	2005	Male	25 to 29	34% (24-45%)	66% (55-76%)	3.1 (1.7-4.9)	12% (8.1-16%)	14% (8.8-20%)	83 (40-150)	6500 (3800-11000)
Honduras	2005	Male	30 to 34	35% (24-47%)	65% (53-76%)	3.5 (2-5.3)	12% (8.6-16%)	15% (10-21%)	84 (45-150)	5800 (3600-9300)
Honduras	2005	Male	35 to 39	34% (23-45%)	66% (55-77%)	3.6 (1.9-5.9)	13% (9.1-17%)	16% (11-22%)	96 (53-160)	5800 (3500-9000)
Honduras	2005	Male	40 to 44	33% (23-44%)	67% (56-77%)	3.9 (2-6.2)	13% (9.7-17%)	17% (12-22%)	110 (60-170)	5800 (3600-8800)
Honduras	2005	Male	45 to 49	33% (22-43%)	67% (57-78%)	4.2 (2.1-7.2)	13% (9.9-17%)	17% (13-21%)	130 (74-190)	5800 (3700-8400)
Honduras	2005	Male	50 to 54	31% (22-41%)	69% (59-78%)	4.3 (2.3-7.2)	12% (9.3-16%)	15% (12-20%)	130 (80-210)	5400 (3400-8000)
Honduras	2005	Male	55 to 59	28% (19-37%)	72% (63-81%)	4.3 (2.1-7.5)	10% (7.6-14%)	13% (9.5-17%)	120 (70-180)	4100 (2600-6200)
Honduras	2005	Male	60 to 64	25% (17-36%)	75% (64-83%)	3.7 (1.7-6.6)	8.4% (6.2-11%)	10% (7.5-14%)	110 (67-170)	3200 (2000-4800)
Honduras	2005	Male	65 to 69	25% (15-37%)	75% (63-85%)	2.8 (1.4-5)	7.5% (5.3-10%)	8.9% (6.4-12%)	120 (69-180)	2800 (1700-4300)
Honduras	2005	Male	70 to 74	22% (14-34%)	78% (66-86%)	2.1 (0.98-4)	6% (4-8.4%)	7.1% (4.8-10%)	110 (64-190)	2100 (1300-3400)
Honduras	2005	Male	75 to 79	19% (11-29%)	81% (71-89%)	1.5 (0.62-3)	4.5% (2.8-6.5%)	5.3% (3.3-7.7%)	85 (46-140)	1300 (690-2100)
Honduras	2005	Male	80 to 84	16% (9.4-26%)	84% (74-91%)	1.3 (0.55-2.3)	3.8% (2.3-5.6%)	4.4% (2.7-6.5%)	67 (38-110)	730 (410-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	2005	Male	85 to 89	15% (8.8-24%)	85% (76-91%)	1.2 (0.49-2.3)	3.3% (1.9-5%)	3.7% (2.1-5.8%)	40 (22-65)	310 (170-490)
Honduras	2005	Male	90 to 94	17% (11-26%)	83% (74-89%)	1.3 (0.52-2.7)	3.1% (1.6-5.1%)	3.5% (1.7-5.8%)	18 (9-30)	98 (51-160)
Honduras	2005	Male	95 plus	21% (12-34%)	79% (66-88%)	1.6 (0.49-3.5)	3.5% (1.5-5.8%)	3.8% (1.4-6.6%)	6.4 (2.3-11)	24 (9.9-40)
Honduras	2010	Female	All Ages	5.2% (4.4-6.2%)	95% (94-96%)	0.22 (0.17-0.28)	1.8% (1.4-2.3%)	3% (2.4-3.8%)	600 (380-850)	17000 (12000-23000)
Honduras	2010	Female	15 to 19	5.5% (3.9-7.4%)	95% (93-96%)	0.22 (0.12-0.36)	0.93% (0.6-1.4%)	1.1% (0.6-1.9%)	3.1 (1.5-5.5)	490 (310-730)
Honduras	2010	Female	20 to 24	7.9% (5.6-10%)	92% (90-94%)	0.3 (0.16-0.51)	1.6% (1.1-2.3%)	1.6% (0.88-2.6%)	5.2 (2.5-9.6)	850 (530-1300)
Honduras	2010	Female	25 to 29	7.3% (5.1-10%)	93% (90-95%)	0.27 (0.15-0.44)	1.6% (1.2-2.2%)	1.6% (1-2.5%)	6 (3-11)	830 (530-1200)
Honduras	2010	Female	30 to 34	6.6% (4.6-10%)	93% (90-95%)	0.25 (0.14-0.42)	1.8% (1.4-2.4%)	2.3% (1.7-3.3%)	9.5 (5-17)	900 (590-1400)
Honduras	2010	Female	35 to 39	6.6% (4.4-10%)	93% (90-96%)	0.28 (0.15-0.48)	2.2% (1.6-2.9%)	3.1% (2.3-4.2%)	15 (7.7-25)	1000 (640-1600)
Honduras	2010	Female	40 to 44	6.4% (4.2-9.5%)	94% (90-96%)	0.32 (0.17-0.54)	2.4% (1.7-3.2%)	3.5% (2.5-4.8%)	20 (11-34)	1200 (690-1800)
Honduras	2010	Female	45 to 49	6.7% (4.2-9.6%)	93% (90-96%)	0.41 (0.2-0.74)	2.9% (2-4.1%)	4.3% (3-6%)	30 (17-49)	1400 (830-2200)
Honduras	2010	Female	50 to 54	8.2% (5.3-12%)	92% (88-95%)	0.59 (0.32-1)	3.6% (2.4-4.9%)	5% (3.5-6.8%)	44 (26-69)	1700 (1000-2600)
Honduras	2010	Female	55 to 59	9.5% (5.4-16%)	91% (84-95%)	0.77 (0.37-1.4)	4.2% (2.9-5.7%)	5.8% (4.1-7.7%)	61 (36-92)	2000 (1300-3000)
Honduras	2010	Female	60 to 64	8.8% (4.6-15%)	91% (85-95%)	0.73 (0.34-1.4)	3.9% (2.8-5.6%)	5.1% (3.6-7.1%)	60 (36-90)	1700 (1000-2500)
Honduras	2010	Female	65 to 69	8.1% (4.6-13%)	92% (87-95%)	0.65 (0.33-1.2)	3.9% (2.7-5.3%)	4.9% (3.5-6.8%)	68 (41-100)	1600 (990-2300)
Honduras	2010	Female	70 to 74	7.2% (4.2-12%)	93% (88-96%)	0.56 (0.27-1)	3.4% (2.3-4.8%)	4.3% (2.9-6%)	74 (43-110)	1400 (810-2000)
Honduras	2010	Female	75 to 79	5.8% (3.2-9.8%)	94% (90-97%)	0.44 (0.19-0.81)	2.7% (1.8-3.8%)	3.3% (2.2-4.6%)	70 (40-110)	1000 (580-1500)
Honduras	2010	Female	80 to 84	5% (2.7-8.1%)	95% (92-97%)	0.36 (0.16-0.66)	2.2% (1.5-3.3%)	2.7% (1.8-3.9%)	71 (43-110)	750 (460-1100)
Honduras	2010	Female	85 to 89	5.1% (2.8-8.4%)	95% (92-97%)	0.35 (0.16-0.67)	1.9% (1.2-3%)	2.3% (1.5-3.6%)	45 (26-73)	340 (200-540)
Honduras	2010	Female	90 to 94	6% (3.5-9.7%)	94% (90-97%)	0.38 (0.16-0.75)	1.7% (0.97-2.7%)	1.9% (1.1-3.3%)	17 (9.3-29)	88 (50-150)
Honduras	2010	Female	95 plus	7.5% (4-13%)	92% (87-96%)	0.41 (0.15-0.83)	1.8% (0.93-3.2%)	2% (0.99-3.7%)	5.5 (2.7-10)	19 (10-34)
Honduras	2010	Male	All Ages	22% (20-24%)	78% (76-80%)	1.8 (1.6-2.1)	6.6% (5.2-8.4%)	7.9% (6.4-9.7%)	1600 (1000-2500)	68000 (46000-100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	2010	Male	15 to 19	16% (14-19%)	84% (81-86%)	0.83 (0.46-1.3)	4.1% (2.6-5.9%)	5% (2.7-8.1%)	25 (12-49)	2600 (1500-4500)
Honduras	2010	Male	20 to 24	36% (32-40%)	64% (60-68%)	2.5 (1.5-3.9)	11% (7.3-16%)	13% (7.7-20%)	89 (43-170)	7700 (4500-13000)
Honduras	2010	Male	25 to 29	42% (37-46%)	58% (54-63%)	3.4 (2.1-5.2)	12% (8.8-17%)	16% (10-22%)	100 (53-180)	8000 (4900-13000)
Honduras	2010	Male	30 to 34	40% (35-44%)	60% (56-65%)	3.5 (2.3-5)	13% (9-17%)	16% (11-22%)	100 (54-170)	7000 (4300-11000)
Honduras	2010	Male	35 to 39	38% (34-42%)	62% (58-66%)	3.6 (2-5.5)	13% (9.5-17%)	17% (12-23%)	110 (59-180)	6500 (4000-10000)
Honduras	2010	Male	40 to 44	37% (33-42%)	63% (58-67%)	3.9 (2.1-5.8)	14% (10-17%)	18% (14-23%)	130 (72-210)	6800 (4200-11000)
Honduras	2010	Male	45 to 49	37% (32-41%)	63% (59-68%)	4.2 (2.3-7)	14% (11-17%)	18% (14-22%)	150 (83-230)	6700 (4200-10000)
Honduras	2010	Male	50 to 54	33% (29-37%)	67% (63-71%)	4.1 (2.3-6.6)	12% (9.8-15%)	16% (13-19%)	160 (94-240)	6200 (4000-9300)
Honduras	2010	Male	55 to 59	28% (24-33%)	72% (67-76%)	3.9 (2.1-6.3)	10% (8-13%)	13% (10-16%)	150 (88-220)	5100 (3200-7500)
Honduras	2010	Male	60 to 64	26% (18-35%)	74% (65-82%)	3.3 (1.6-5.9)	8.4% (6.2-11%)	10% (7.6-13%)	130 (76-200)	3700 (2300-5500)
Honduras	2010	Male	65 to 69	25% (15-37%)	75% (63-85%)	2.5 (1.2-4.4)	7.3% (5.2-9.8%)	8.8% (6.3-12%)	120 (72-200)	3000 (1800-4600)
Honduras	2010	Male	70 to 74	22% (14-34%)	78% (66-86%)	1.8 (0.86-3.5)	5.7% (3.8-8.1%)	6.8% (4.4-9.5%)	110 (65-190)	2200 (1300-3500)
Honduras	2010	Male	75 to 79	18% (11-29%)	82% (71-89%)	1.3 (0.56-2.6)	4.3% (2.5-6.2%)	5% (2.9-7.4%)	94 (48-160)	1400 (730-2300)
Honduras	2010	Male	80 to 84	16% (9.2-25%)	84% (75-91%)	1.1 (0.47-2)	3.6% (2.2-5.4%)	4.2% (2.5-6.3%)	78 (44-130)	850 (490-1400)
Honduras	2010	Male	85 to 89	15% (8.7-24%)	85% (76-91%)	1 (0.43-2)	3.1% (1.7-4.8%)	3.5% (1.9-5.4%)	49 (25-82)	380 (200-620)
Honduras	2010	Male	90 to 94	17% (10-26%)	83% (74-90%)	1.2 (0.46-2.4)	2.9% (1.4-4.8%)	3.2% (1.5-5.4%)	22 (10-37)	120 (58-190)
Honduras	2010	Male	95 plus	21% (11-34%)	79% (66-89%)	1.4 (0.43-3)	3.2% (1.3-5.7%)	3.5% (1.2-6.4%)	6.8 (2.2-12)	25 (9.8-44)
Honduras	2016	Female	All Ages	5.5% (4.7-6.5%)	95% (94-95%)	0.25 (0.18-0.33)	2% (1.5-2.5%)	3.1% (2.4-3.9%)	690 (460-970)	20000 (14000-27000)
Honduras	2016	Female	15 to 19	5.6% (3.9-7.5%)	94% (93-96%)	0.23 (0.11-0.38)	0.97% (0.62-1.4%)	1.2% (0.6-2%)	3.1 (1.5-5.6)	500 (320-790)
Honduras	2016	Female	20 to 24	8.1% (5.6-11%)	92% (89-94%)	0.31 (0.16-0.55)	1.7% (1.2-2.4%)	1.7% (0.92-2.9%)	5.6 (2.7-10)	950 (590-1400)
Honduras	2016	Female	25 to 29	7.4% (5.2-10%)	93% (90-95%)	0.28 (0.15-0.47)	1.7% (1.2-2.2%)	1.7% (1.1-2.7%)	6.6 (3.2-12)	940 (610-1400)
Honduras	2016	Female	30 to 34	6.7% (4.6-10%)	93% (90-95%)	0.26 (0.15-0.45)	1.8% (1.4-2.5%)	2.4% (1.7-3.5%)	11 (5.4-19)	1100 (680-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	2016	Female	35 to 39	6.7% (4.4-10%)	93% (90-96%)	0.28 (0.14-0.51)	2.1% (1.5-2.9%)	3.2% (2.3-4.4%)	17 (8.6-30)	1200 (740-1900)
Honduras	2016	Female	40 to 44	6.4% (4.3-9.6%)	94% (90-96%)	0.32 (0.16-0.54)	2.3% (1.6-3.1%)	3.5% (2.5-4.8%)	23 (12-38)	1300 (750-2000)
Honduras	2016	Female	45 to 49	6.7% (4.2-9.7%)	93% (90-96%)	0.41 (0.2-0.75)	2.8% (2-3.9%)	4.2% (3-5.8%)	34 (19-56)	1600 (950-2500)
Honduras	2016	Female	50 to 54	8.2% (5.4-12%)	92% (88-95%)	0.6 (0.31-1)	3.5% (2.4-4.7%)	5% (3.4-6.8%)	49 (27-78)	2000 (1100-3000)
Honduras	2016	Female	55 to 59	9.5% (5.5-16%)	90% (84-94%)	0.78 (0.37-1.4)	4.1% (2.8-5.7%)	5.6% (4-7.5%)	67 (41-100)	2300 (1400-3500)
Honduras	2016	Female	60 to 64	8.8% (4.7-15%)	91% (85-95%)	0.73 (0.34-1.4)	3.8% (2.7-5.3%)	5% (3.6-6.9%)	74 (46-110)	2100 (1300-3200)
Honduras	2016	Female	65 to 69	8.1% (4.7-13%)	92% (87-95%)	0.66 (0.33-1.2)	3.8% (2.6-5.2%)	4.8% (3.4-6.6%)	80 (51-120)	1900 (1200-2700)
Honduras	2016	Female	70 to 74	7.2% (4.3-12%)	93% (88-96%)	0.57 (0.26-1.1)	3.4% (2.2-4.8%)	4.2% (2.8-6.1%)	80 (48-120)	1500 (910-2300)
Honduras	2016	Female	75 to 79	5.8% (3.3-9.6%)	94% (90-97%)	0.45 (0.18-0.85)	2.6% (1.7-3.9%)	3.2% (2.2-4.7%)	74 (43-110)	1100 (630-1600)
Honduras	2016	Female	80 to 84	5% (2.7-8.1%)	95% (92-97%)	0.37 (0.16-0.68)	2.2% (1.4-3.3%)	2.6% (1.7-4%)	83 (50-130)	870 (520-1300)
Honduras	2016	Female	85 to 89	5.1% (2.8-8.5%)	95% (91-97%)	0.35 (0.15-0.7)	1.9% (1.2-3%)	2.3% (1.4-3.5%)	56 (33-90)	420 (250-660)
Honduras	2016	Female	90 to 94	6.1% (3.4-9.8%)	94% (90-97%)	0.38 (0.16-0.76)	1.7% (0.95-2.8%)	1.9% (1.1-3.2%)	23 (13-39)	120 (68-200)
Honduras	2016	Female	95 plus	7.6% (4-13%)	92% (87-96%)	0.41 (0.15-0.83)	1.8% (0.94-3.2%)	2% (1-3.7%)	7.4 (3.6-14)	26 (14-47)
Honduras	2016	Male	All Ages	23% (21-25%)	77% (75-79%)	2 (1.5-2.5)	7.2% (5.7-9%)	8.2% (6.6-10%)	1900 (1200-2900)	79000 (53000-110000)
Honduras	2016	Male	15 to 19	16% (14-19%)	84% (81-86%)	0.84 (0.42-1.4)	4.2% (2.7-6.2%)	5.2% (2.7-8.7%)	26 (12-53)	2700 (1600-4700)
Honduras	2016	Male	20 to 24	36% (32-40%)	64% (60-68%)	2.6 (1.4-4.1)	11% (7.5-16%)	14% (8-22%)	98 (48-190)	8600 (5100-14000)
Honduras	2016	Male	25 to 29	42% (37-46%)	58% (54-63%)	3.5 (2-5.6)	13% (9.1-18%)	16% (10-24%)	120 (61-200)	9100 (5700-14000)
Honduras	2016	Male	30 to 34	40% (35-44%)	60% (56-65%)	3.6 (2.2-5.4)	12% (9.2-17%)	16% (12-22%)	120 (65-200)	8200 (5300-13000)
Honduras	2016	Male	35 to 39	38% (33-42%)	62% (58-67%)	3.6 (1.9-5.7)	13% (9.4-17%)	17% (12-23%)	120 (68-210)	7700 (4800-12000)
Honduras	2016	Male	40 to 44	37% (33-42%)	63% (58-67%)	3.9 (2.1-6)	14% (10-17%)	18% (14-23%)	140 (80-230)	7500 (4800-12000)
Honduras	2016	Male	45 to 49	37% (32-41%)	63% (59-68%)	4.2 (2.4-7.2)	14% (11-17%)	18% (14-22%)	170 (97-260)	7700 (4900-12000)
Honduras	2016	Male	50 to 54	33% (29-37%)	67% (63-71%)	4.1 (2.3-6.7)	12% (9.6-15%)	16% (13-19%)	180 (110-280)	7200 (4700-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Honduras	2016	Male	55 to 59	28% (24-33%)	72% (67-76%)	3.9 (2.1-6.5)	10% (7.9-13%)	13% (9.9-16%)	170 (110-270)	6000 (3800-8800)
Honduras	2016	Male	60 to 64	26% (18-35%)	74% (65-82%)	3.3 (1.5-6.1)	8.3% (6.2-11%)	10% (7.5-13%)	170 (100-260)	4800 (3100-7400)
Honduras	2016	Male	65 to 69	25% (15-38%)	75% (62-85%)	2.5 (1.2-4.5)	7.2% (5.2-9.6%)	8.7% (6.2-12%)	150 (92-240)	3700 (2300-5600)
Honduras	2016	Male	70 to 74	22% (14-35%)	78% (65-86%)	1.8 (0.84-3.5)	5.7% (3.7-7.9%)	6.7% (4.4-9.5%)	130 (73-210)	2400 (1500-3900)
Honduras	2016	Male	75 to 79	19% (12-29%)	81% (71-88%)	1.3 (0.55-2.6)	4.3% (2.5-6.3%)	5% (2.9-7.4%)	100 (53-170)	1500 (830-2500)
Honduras	2016	Male	80 to 84	16% (9.4-26%)	84% (74-91%)	1.1 (0.48-2)	3.6% (2.2-5.3%)	4.2% (2.5-6.2%)	95 (52-150)	1000 (580-1700)
Honduras	2016	Male	85 to 89	15% (8.8-24%)	85% (76-91%)	1.1 (0.43-2.1)	3.1% (1.7-4.8%)	3.5% (1.8-5.5%)	64 (32-100)	500 (270-800)
Honduras	2016	Male	90 to 94	17% (11-26%)	83% (74-89%)	1.2 (0.46-2.4)	2.9% (1.4-4.9%)	3.2% (1.5-5.5%)	30 (14-52)	170 (80-280)
Honduras	2016	Male	95 plus	21% (12-34%)	79% (66-88%)	1.4 (0.44-3)	3.2% (1.3-5.6%)	3.5% (1.2-6.3%)	9.7 (3.4-18)	37 (15-64)
Mexico	1990	Female	All Ages	29% (23-35%)	71% (65-77%)	0.34 (0.27-0.42)	1% (0.74-1.4%)	2.1% (1.3-3%)	4000 (2600-5800)	120000 (86000-160000)
Mexico	1990	Female	15 to 19	26% (15-39%)	74% (61-85%)	0.31 (0.15-0.52)	0.78% (0.47-1.2%)	1.3% (0.57-2.2%)	36 (16-63)	4500 (2600-6800)
Mexico	1990	Female	20 to 24	37% (21-52%)	63% (48-79%)	0.37 (0.18-0.63)	1.1% (0.7-1.7%)	1.4% (0.64-2.4%)	43 (19-75)	5900 (3500-8800)
Mexico	1990	Female	25 to 29	41% (23-57%)	59% (43-77%)	0.38 (0.19-0.64)	1.2% (0.83-1.7%)	1.8% (1-2.7%)	56 (32-86)	6000 (4000-8400)
Mexico	1990	Female	30 to 34	43% (25-59%)	57% (41-75%)	0.42 (0.21-0.68)	1.6% (1.1-2.1%)	2.9% (2.2-3.7%)	99 (75-130)	7200 (5200-9300)
Mexico	1990	Female	35 to 39	45% (28-59%)	55% (41-72%)	0.5 (0.26-0.83)	2.2% (1.7-2.9%)	4.6% (3.7-5.6%)	190 (150-230)	10000 (8100-13000)
Mexico	1990	Female	40 to 44	45% (30-60%)	55% (40-70%)	0.6 (0.33-0.95)	2.6% (1.9-3.4%)	5.3% (4.4-6.3%)	230 (190-280)	11000 (8100-13000)
Mexico	1990	Female	45 to 49	43% (26-58%)	57% (42-74%)	0.73 (0.36-1.2)	2.9% (2-4%)	5.6% (4.4-7%)	310 (250-400)	12000 (8600-16000)
Mexico	1990	Female	50 to 54	43% (26-58%)	57% (42-74%)	0.89 (0.47-1.4)	2.9% (1.7-4.2%)	5.1% (3.8-6.8%)	350 (250-460)	12000 (7300-17000)
Mexico	1990	Female	55 to 59	43% (27-59%)	57% (41-73%)	1.1 (0.55-1.8)	2.8% (1.5-4.7%)	4.8% (3-7.2%)	420 (270-640)	13000 (6900-21000)
Mexico	1990	Female	60 to 64	43% (26-61%)	57% (39-74%)	1.2 (0.62-2)	2.5% (1-4.6%)	4.1% (2.1-6.6%)	440 (230-710)	11000 (4700-20000)
Mexico	1990	Female	65 to 69	41% (24-57%)	59% (43-76%)	1.3 (0.67-2)	2.5% (1-4.4%)	3.7% (1.9-6.1%)	470 (240-770)	10000 (4300-18000)
Mexico	1990	Female	70 to 74	35% (21-51%)	65% (49-79%)	1.2 (0.59-2.1)	2.4% (0.84-4.6%)	3.5% (1.4-6.1%)	440 (180-790)	8100 (2700-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	1990	Female	75 to 79	28% (15-43%)	72% (57-85%)	0.96 (0.43-1.7)	1.9% (0.63-3.7%)	2.6% (1-4.8%)	400 (160-750)	5800 (1900-11000)
Mexico	1990	Female	80 to 84	24% (13-38%)	76% (62-87%)	0.78 (0.35-1.4)	1.3% (0.36-2.6%)	1.8% (0.54-3.3%)	280 (86-540)	3000 (790-5900)
Mexico	1990	Female	85 to 89	25% (13-39%)	75% (61-87%)	0.77 (0.36-1.4)	1% (0.065-2.2%)	1.2% (0.08-2.7%)	190 (12-410)	1400 (88-3300)
Mexico	1990	Female	90 to 94	28% (16-43%)	72% (57-84%)	0.83 (0.36-1.5)	0.64% (-0.41-2.1%)	0.71% (-0.55-2.4%)	65 (-51-220)	370 (-230-1200)
Mexico	1990	Female	95 plus	33% (17-49%)	67% (51-83%)	0.9 (0.33-1.8)	0.44% (-0.89-2.4%)	0.39% (-1.2-2.7%)	12 (-38-87)	60 (-120-310)
Mexico	1990	Male	All Ages	46% (40-51%)	54% (49-60%)	2.9 (2.5-3.3)	8.6% (7.6-9.6%)	12% (11-14%)	31000 (27000-35000)	1300000 (1100000-1500000)
Mexico	1990	Male	15 to 19	34% (24-45%)	66% (55-76%)	1.3 (0.83-1.9)	5.7% (4-8%)	7.6% (5-11%)	520 (340-760)	46000 (32000-63000)
Mexico	1990	Male	20 to 24	63% (51-73%)	37% (27-49%)	3.5 (2.5-4.7)	14% (11-18%)	18% (13-23%)	1600 (1100-2000)	120000 (93000-160000)
Mexico	1990	Male	25 to 29	73% (63-81%)	27% (19-37%)	4.8 (3.4-6.3)	18% (14-22%)	22% (17-27%)	2000 (1500-2500)	140000 (110000-180000)
Mexico	1990	Male	30 to 34	76% (65-83%)	24% (17-35%)	5.6 (4-7.3)	20% (16-23%)	24% (20-29%)	2100 (1700-2600)	140000 (110000-170000)
Mexico	1990	Male	35 to 39	77% (68-84%)	23% (16-32%)	6.2 (4.4-8.3)	21% (18-25%)	27% (23-31%)	2500 (2100-2900)	140000 (120000-170000)
Mexico	1990	Male	40 to 44	78% (69-85%)	22% (15-31%)	6.7 (5-8.9)	22% (18-25%)	28% (24-32%)	2400 (2100-2800)	130000 (110000-140000)
Mexico	1990	Male	45 to 49	78% (67-85%)	22% (15-33%)	7.2 (4.9-10)	21% (18-24%)	27% (24-30%)	2800 (2400-3100)	130000 (110000-140000)
Mexico	1990	Male	50 to 54	78% (69-85%)	22% (15-31%)	8.2 (5.8-11)	20% (17-23%)	26% (22-29%)	2800 (2400-3100)	110000 (95000-130000)
Mexico	1990	Male	55 to 59	78% (69-86%)	22% (14-31%)	10 (7.1-13)	19% (15-22%)	23% (20-27%)	3000 (2500-3500)	100000 (85000-120000)
Mexico	1990	Male	60 to 64	76% (65-85%)	24% (15-35%)	9.4 (6.5-13)	16% (13-20%)	20% (16-23%)	2800 (2300-3300)	82000 (66000-100000)
Mexico	1990	Male	65 to 69	70% (58-81%)	30% (19-42%)	6.8 (4.6-9.3)	14% (11-17%)	17% (13-20%)	2600 (2100-3100)	63000 (50000-77000)
Mexico	1990	Male	70 to 74	61% (49-73%)	39% (27-51%)	4.5 (2.9-6.5)	10% (8.1-13%)	13% (9.9-15%)	1900 (1500-2300)	37000 (29000-46000)
Mexico	1990	Male	75 to 79	51% (37-65%)	49% (35-63%)	3.1 (1.9-4.8)	7.7% (5.7-10%)	9.1% (6.7-12%)	1500 (1100-2000)	23000 (17000-30000)
Mexico	1990	Male	80 to 84	45% (32-58%)	55% (42-68%)	2.6 (1.6-3.9)	6.1% (4.4-8.1%)	7.2% (5.2-9.5%)	1100 (800-1500)	13000 (9100-17000)
Mexico	1990	Male	85 to 89	43% (31-56%)	57% (44-69%)	2.5 (1.5-3.8)	5.4% (3.8-7.4%)	6.2% (4.3-8.5%)	780 (540-1100)	6300 (4400-8500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	1990	Male	90 to 94	46% (34-58%)	54% (42-66%)	2.6 (1.6-4)	5% (3.1-6.9%)	5.6% (3.3-8%)	370 (220-540)	2100 (1300-3000)
Mexico	1990	Male	95 plus	51% (35-65%)	49% (35-65%)	2.8 (1.6-4.6)	5% (2.5-7.7%)	5.4% (2.5-8.7%)	110 (49-170)	410 (200-620)
Mexico	1995	Female	All Ages	30% (25-36%)	70% (64-75%)	0.37 (0.3-0.46)	1.2% (0.82-1.6%)	2.3% (1.4-3.4%)	4800 (2900-7000)	140000 (94000-190000)
Mexico	1995	Female	15 to 19	28% (17-41%)	72% (59-83%)	0.34 (0.17-0.56)	0.8% (0.48-1.3%)	1.4% (0.61-2.4%)	35 (16-62)	4400 (2600-6800)
Mexico	1995	Female	20 to 24	38% (23-53%)	62% (47-77%)	0.39 (0.19-0.66)	1.1% (0.69-1.6%)	1.4% (0.67-2.5%)	44 (20-75)	6500 (3900-9800)
Mexico	1995	Female	25 to 29	41% (25-56%)	59% (44-75%)	0.4 (0.2-0.66)	1.1% (0.74-1.6%)	1.7% (1-2.6%)	52 (31-81)	5900 (3900-8300)
Mexico	1995	Female	30 to 34	43% (26-58%)	57% (42-74%)	0.43 (0.23-0.68)	1.4% (0.94-1.9%)	2.8% (2.2-3.6%)	98 (75-130)	7100 (5000-9200)
Mexico	1995	Female	35 to 39	45% (28-59%)	55% (41-72%)	0.51 (0.27-0.83)	1.9% (1.3-2.5%)	4.4% (3.6-5.4%)	190 (150-230)	10000 (7500-13000)
Mexico	1995	Female	40 to 44	44% (30-58%)	56% (42-70%)	0.61 (0.33-0.97)	2.2% (1.5-2.9%)	5.1% (4.2-6.2%)	250 (210-310)	11000 (8000-14000)
Mexico	1995	Female	45 to 49	42% (25-56%)	58% (44-75%)	0.74 (0.36-1.3)	2.6% (1.7-3.8%)	5.4% (4.2-6.8%)	320 (260-410)	12000 (8200-17000)
Mexico	1995	Female	50 to 54	42% (26-56%)	58% (44-74%)	0.9 (0.48-1.5)	2.7% (1.6-4.1%)	5.1% (3.7-6.7%)	390 (290-520)	13000 (7900-19000)
Mexico	1995	Female	55 to 59	42% (27-57%)	58% (43-73%)	1.1 (0.56-1.8)	2.9% (1.6-4.8%)	4.9% (3.2-7.3%)	470 (310-710)	14000 (7900-23000)
Mexico	1995	Female	60 to 64	42% (27-59%)	58% (41-73%)	1.2 (0.64-2)	2.7% (1.1-4.9%)	4.2% (2.3-6.8%)	530 (280-870)	14000 (5900-25000)
Mexico	1995	Female	65 to 69	42% (26-58%)	58% (42-74%)	1.4 (0.75-2.2)	2.6% (1.1-4.8%)	4% (2-6.7%)	590 (290-980)	13000 (5200-23000)
Mexico	1995	Female	70 to 74	38% (24-54%)	62% (46-76%)	1.4 (0.7-2.3)	2.6% (0.82-5.1%)	3.8% (1.5-6.7%)	620 (240-1100)	11000 (3500-22000)
Mexico	1995	Female	75 to 79	31% (18-46%)	69% (54-82%)	1.1 (0.52-2)	2.1% (0.6-4.1%)	2.9% (0.98-5.4%)	450 (150-870)	6600 (1900-13000)
Mexico	1995	Female	80 to 84	26% (15-40%)	74% (60-85%)	0.91 (0.42-1.6)	1.5% (0.37-3%)	2% (0.54-3.8%)	360 (100-700)	3800 (920-7700)
Mexico	1995	Female	85 to 89	26% (15-42%)	74% (58-85%)	0.86 (0.42-1.6)	1.1% (0.081-2.4%)	1.3% (0.11-3%)	240 (20-550)	1900 (140-4400)
Mexico	1995	Female	90 to 94	30% (17-44%)	70% (56-83%)	0.92 (0.43-1.7)	0.72% (-0.42-2.3%)	0.8% (-0.56-2.7%)	93 (-65-310)	530 (-320-1700)
Mexico	1995	Female	95 plus	34% (19-51%)	66% (49-81%)	0.99 (0.37-1.9)	0.5% (-0.93-2.7%)	0.45% (-1.3-3.1%)	20 (-57-140)	92 (-170-500)
Mexico	1995	Male	All Ages	49% (44-54%)	51% (46-56%)	3.2 (2.8-3.7)	9.3% (8.1-10%)	12% (11-14%)	33000 (29000-37000)	1400000 (1200000-1600000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	1995	Male	15 to 19	40% (30-51%)	60% (49-70%)	1.6 (1-2.3)	6.4% (4.5-8.8%)	8.9% (6-13%)	550 (370-790)	48000 (34000-66000)
Mexico	1995	Male	20 to 24	66% (55-76%)	34% (24-45%)	3.9 (2.8-5.1)	15% (11-18%)	18% (14-24%)	1700 (1300-2200)	130000 (100000-170000)
Mexico	1995	Male	25 to 29	73% (63-81%)	27% (19-37%)	5 (3.7-6.6)	17% (14-21%)	21% (17-27%)	2000 (1500-2500)	140000 (110000-170000)
Mexico	1995	Male	30 to 34	76% (66-83%)	24% (17-34%)	5.8 (4.2-7.5)	18% (15-22%)	23% (19-28%)	2200 (1800-2700)	140000 (120000-170000)
Mexico	1995	Male	35 to 39	77% (68-84%)	23% (16-32%)	6.5 (4.7-8.7)	20% (17-23%)	26% (22-30%)	2600 (2200-3100)	150000 (130000-180000)
Mexico	1995	Male	40 to 44	78% (69-85%)	22% (15-31%)	7 (5.1-9.3)	20% (17-23%)	27% (24-31%)	2700 (2400-3100)	140000 (120000-160000)
Mexico	1995	Male	45 to 49	78% (67-85%)	22% (15-33%)	7.4 (5.1-10)	20% (17-23%)	26% (23-29%)	2800 (2500-3200)	130000 (110000-150000)
Mexico	1995	Male	50 to 54	78% (68-85%)	22% (15-32%)	8.5 (6.1-11)	19% (16-23%)	25% (22-28%)	3000 (2600-3400)	120000 (100000-140000)
Mexico	1995	Male	55 to 59	78% (69-85%)	22% (15-31%)	10 (7.4-14)	18% (15-22%)	23% (19-27%)	3100 (2600-3600)	110000 (88000-130000)
Mexico	1995	Male	60 to 64	75% (64-84%)	25% (16-36%)	9.6 (6.7-14)	16% (12-19%)	19% (15-23%)	3000 (2400-3600)	90000 (70000-110000)
Mexico	1995	Male	65 to 69	68% (55-79%)	32% (21-45%)	6.7 (4.5-9.3)	13% (10-16%)	16% (12-19%)	2700 (2200-3300)	67000 (52000-83000)
Mexico	1995	Male	70 to 74	57% (44-69%)	43% (31-56%)	4.2 (2.6-6.1)	9.7% (7.4-12%)	12% (9-14%)	2300 (1800-2800)	44000 (34000-56000)
Mexico	1995	Male	75 to 79	46% (33-61%)	54% (39-67%)	2.9 (1.7-4.5)	7% (5.2-9.3%)	8.3% (6.2-11%)	1400 (1100-1900)	22000 (16000-29000)
Mexico	1995	Male	80 to 84	41% (29-55%)	59% (45-71%)	2.5 (1.5-3.7)	5.7% (4.1-7.7%)	6.7% (4.8-8.9%)	1200 (840-1600)	13000 (9500-18000)
Mexico	1995	Male	85 to 89	41% (29-54%)	59% (46-71%)	2.4 (1.4-3.7)	5.1% (3.6-6.9%)	5.8% (4.1-7.9%)	860 (600-1200)	6900 (4900-9400)
Mexico	1995	Male	90 to 94	44% (32-56%)	56% (44-68%)	2.5 (1.5-3.9)	4.7% (3-6.5%)	5.2% (3.2-7.4%)	420 (260-590)	2300 (1500-3300)
Mexico	1995	Male	95 plus	48% (32-63%)	52% (37-68%)	2.7 (1.5-4.4)	4.7% (2.5-7.1%)	5.1% (2.4-7.9%)	130 (61-200)	490 (260-750)
Mexico	2000	Female	All Ages	28% (25-32%)	72% (68-75%)	0.33 (0.27-0.41)	1.3% (0.99-1.7%)	2.5% (1.8-3.5%)	5200 (3700-7100)	150000 (110000-190000)
Mexico	2000	Female	15 to 19	32% (24-41%)	68% (59-76%)	0.41 (0.24-0.62)	0.88% (0.52-1.4%)	1.6% (0.71-2.7%)	37 (17-63)	4700 (2800-7200)
Mexico	2000	Female	20 to 24	37% (27-48%)	63% (52-73%)	0.39 (0.22-0.62)	1.1% (0.71-1.6%)	1.4% (0.69-2.4%)	39 (18-63)	6200 (3800-9300)
Mexico	2000	Female	25 to 29	35% (25-46%)	65% (54-75%)	0.34 (0.19-0.54)	1.1% (0.74-1.5%)	1.6% (1-2.4%)	47 (30-69)	6100 (4100-8300)
Mexico	2000	Female	30 to 34	35% (24-46%)	65% (54-76%)	0.35 (0.21-0.54)	1.2% (0.87-1.6%)	2.6% (2.1-3.3%)	87 (69-110)	6900 (5100-9000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	2000	Female	35 to 39	35% (25-47%)	65% (53-75%)	0.4 (0.22-0.64)	1.7% (1.3-2.2%)	4.1% (3.5-4.9%)	170 (140-200)	9500 (7400-12000)
Mexico	2000	Female	40 to 44	35% (26-47%)	65% (53-74%)	0.48 (0.28-0.75)	2% (1.5-2.7%)	4.9% (4.1-5.7%)	250 (210-290)	11000 (8900-14000)
Mexico	2000	Female	45 to 49	35% (24-47%)	65% (53-76%)	0.61 (0.34-1)	2.4% (1.6-3.3%)	5.1% (4.2-6.4%)	340 (280-430)	14000 (9800-18000)
Mexico	2000	Female	50 to 54	35% (25-47%)	65% (53-75%)	0.75 (0.45-1.2)	2.8% (1.8-3.8%)	5% (3.9-6.3%)	420 (330-530)	15000 (10000-19000)
Mexico	2000	Female	55 to 59	36% (25-48%)	64% (52-75%)	0.92 (0.51-1.5)	2.9% (1.9-4.5%)	4.9% (3.5-6.7%)	530 (380-730)	16000 (10000-24000)
Mexico	2000	Female	60 to 64	36% (24-50%)	64% (50-76%)	1 (0.58-1.7)	2.9% (1.7-4.5%)	4.3% (2.8-6.4%)	590 (390-870)	16000 (9100-24000)
Mexico	2000	Female	65 to 69	33% (22-46%)	67% (54-78%)	1.1 (0.61-1.7)	2.8% (1.7-4.4%)	4.1% (2.7-6%)	690 (450-1000)	15000 (9000-24000)
Mexico	2000	Female	70 to 74	28% (18-39%)	72% (61-82%)	1 (0.54-1.6)	2.7% (1.4-4.5%)	3.8% (2.2-6%)	710 (410-1100)	13000 (7100-22000)
Mexico	2000	Female	75 to 79	22% (15-33%)	78% (67-85%)	0.83 (0.43-1.4)	2.1% (1.1-3.7%)	2.9% (1.6-4.8%)	580 (310-950)	8400 (4200-15000)
Mexico	2000	Female	80 to 84	20% (13-28%)	80% (72-87%)	0.7 (0.39-1.1)	1.5% (0.72-2.6%)	2% (1-3.4%)	380 (190-640)	4100 (1900-7100)
Mexico	2000	Female	85 to 89	20% (13-29%)	80% (71-87%)	0.67 (0.36-1.1)	1.1% (0.33-2.1%)	1.3% (0.43-2.6%)	240 (76-460)	1900 (580-3700)
Mexico	2000	Female	90 to 94	23% (15-32%)	77% (68-85%)	0.72 (0.36-1.3)	0.71% (-0.11-1.9%)	0.81% (-0.22-2.2%)	95 (-27-260)	560 (-89-1500)
Mexico	2000	Female	95 plus	27% (16-39%)	73% (61-84%)	0.78 (0.31-1.5)	0.54% (-0.5-2.2%)	0.49% (-0.83-2.5%)	24 (-40-120)	110 (-100-450)
Mexico	2000	Male	All Ages	50% (46-55%)	50% (45-54%)	3.5 (3-3.9)	9.8% (8.5-11%)	13% (11-14%)	33000 (29000-37000)	1400000 (1200000-1600000)
Mexico	2000	Male	15 to 19	44% (35-53%)	56% (47-65%)	1.8 (1.2-2.5)	6.5% (4.7-8.9%)	9.4% (6.4-13%)	500 (340-700)	45000 (32000-61000)
Mexico	2000	Male	20 to 24	67% (58-75%)	33% (25-42%)	4 (2.9-5.3)	14% (11-18%)	18% (13-24%)	1400 (1000-1800)	120000 (89000-150000)
Mexico	2000	Male	25 to 29	72% (63-79%)	28% (21-37%)	5 (3.6-6.5)	16% (13-20%)	21% (16-26%)	1800 (1400-2200)	140000 (110000-170000)
Mexico	2000	Male	30 to 34	75% (66-82%)	25% (18-34%)	5.7 (4.3-7.3)	17% (14-20%)	22% (19-27%)	2000 (1700-2400)	140000 (120000-160000)
Mexico	2000	Male	35 to 39	76% (69-83%)	24% (17-31%)	6.4 (4.7-8.6)	19% (16-22%)	25% (21-29%)	2400 (2100-2800)	150000 (120000-170000)
Mexico	2000	Male	40 to 44	77% (69-84%)	23% (16-31%)	6.9 (5.2-9.1)	19% (17-22%)	26% (23-29%)	2700 (2400-3000)	140000 (120000-160000)
Mexico	2000	Male	45 to 49	77% (67-84%)	23% (16-33%)	7.3 (5.2-10)	19% (16-22%)	25% (22-28%)	2900 (2600-3300)	140000 (120000-160000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	2000	Male	50 to 54	77% (68-84%)	23% (16-32%)	8.4 (6.1-11)	18% (15-21%)	23% (20-27%)	3000 (2600-3400)	120000 (100000-140000)
Mexico	2000	Male	55 to 59	78% (70-85%)	22% (15-30%)	11 (7.6-14)	17% (14-21%)	22% (18-25%)	3200 (2700-3800)	110000 (93000-140000)
Mexico	2000	Male	60 to 64	76% (66-84%)	24% (16-34%)	10 (7.2-14)	15% (12-19%)	18% (15-22%)	3100 (2500-3800)	94000 (74000-120000)
Mexico	2000	Male	65 to 69	69% (57-79%)	31% (21-43%)	7 (4.9-9.7)	13% (9.8-16%)	15% (12-18%)	3000 (2400-3700)	75000 (58000-92000)
Mexico	2000	Male	70 to 74	59% (46-70%)	41% (30-54%)	4.5 (2.9-6.5)	9.6% (7.5-12%)	12% (9.1-14%)	2500 (1900-3000)	49000 (38000-62000)
Mexico	2000	Male	75 to 79	50% (37-64%)	50% (36-63%)	3.3 (2-4.9)	7.4% (5.4-9.6%)	8.7% (6.5-11%)	1900 (1400-2500)	30000 (22000-39000)
Mexico	2000	Male	80 to 84	45% (32-58%)	55% (42-68%)	2.9 (1.8-4.2)	6.1% (4.5-8%)	7.1% (5.3-9.4%)	1300 (930-1700)	15000 (11000-20000)
Mexico	2000	Male	85 to 89	44% (32-55%)	56% (45-68%)	2.6 (1.6-4)	5.2% (3.7-7%)	6% (4.2-8.1%)	890 (620-1200)	7300 (5200-9900)
Mexico	2000	Male	90 to 94	46% (35-57%)	54% (43-65%)	2.7 (1.7-4.1)	4.7% (3.1-6.5%)	5.3% (3.3-7.5%)	440 (280-620)	2600 (1700-3600)
Mexico	2000	Male	95 plus	51% (35-65%)	49% (35-65%)	3 (1.6-4.7)	4.8% (2.6-7.2%)	5.2% (2.6-8%)	150 (74-230)	570 (310-870)
Mexico	2005	Female	All Ages	26% (23-29%)	74% (71-77%)	0.33 (0.26-0.39)	1.4% (1.1-1.8%)	2.6% (1.9-3.5%)	6000 (4300-7900)	160000 (120000-200000)
Mexico	2005	Female	15 to 19	26% (20-34%)	74% (66-80%)	0.33 (0.2-0.51)	0.85% (0.53-1.3%)	1.4% (0.62-2.4%)	33 (15-56)	4600 (2700-6800)
Mexico	2005	Female	20 to 24	36% (27-45%)	64% (55-73%)	0.38 (0.21-0.59)	1.1% (0.74-1.6%)	1.4% (0.62-2.3%)	37 (16-60)	6200 (3800-9200)
Mexico	2005	Female	25 to 29	36% (27-45%)	64% (55-73%)	0.34 (0.21-0.54)	1% (0.68-1.4%)	1.5% (0.91-2.3%)	45 (27-67)	5900 (3900-8200)
Mexico	2005	Female	30 to 34	34% (26-44%)	66% (56-74%)	0.34 (0.21-0.52)	1.1% (0.73-1.5%)	2.3% (1.8-3%)	80 (62-100)	6600 (4600-8700)
Mexico	2005	Female	35 to 39	34% (26-44%)	66% (56-74%)	0.38 (0.22-0.6)	1.4% (1-1.9%)	3.7% (3.1-4.5%)	160 (130-190)	8800 (6700-11000)
Mexico	2005	Female	40 to 44	34% (25-44%)	66% (56-75%)	0.46 (0.28-0.71)	1.8% (1.3-2.4%)	4.4% (3.8-5.2%)	250 (210-290)	11000 (8800-14000)
Mexico	2005	Female	45 to 49	33% (22-43%)	67% (57-78%)	0.56 (0.31-0.91)	2.2% (1.5-3%)	4.7% (3.9-5.7%)	360 (300-440)	14000 (11000-18000)
Mexico	2005	Female	50 to 54	32% (22-42%)	68% (58-78%)	0.67 (0.4-1)	2.5% (1.7-3.4%)	4.7% (3.8-5.8%)	470 (380-580)	16000 (12000-22000)
Mexico	2005	Female	55 to 59	32% (23-44%)	68% (56-77%)	0.82 (0.46-1.3)	2.8% (1.9-4.1%)	4.6% (3.5-6.2%)	570 (430-770)	17000 (12000-25000)
Mexico	2005	Female	60 to 64	32% (21-45%)	68% (55-79%)	0.93 (0.52-1.5)	2.8% (1.8-4.3%)	4.3% (3-6%)	680 (470-950)	18000 (11000-27000)
Mexico	2005	Female	65 to 69	30% (20-42%)	70% (58-80%)	0.99 (0.56-1.5)	2.9% (1.8-4.3%)	4.2% (2.9-5.8%)	770 (530-1100)	17000 (11000-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	2005	Female	70 to 74	26% (17-36%)	74% (64-83%)	0.96 (0.54-1.6)	2.8% (1.6-4.5%)	3.9% (2.4-5.9%)	860 (530-1300)	16000 (9100-26000)
Mexico	2005	Female	75 to 79	21% (14-30%)	79% (70-86%)	0.8 (0.42-1.3)	2.2% (1.2-3.7%)	3% (1.8-4.8%)	720 (420-1200)	11000 (5600-18000)
Mexico	2005	Female	80 to 84	18% (12-26%)	82% (74-88%)	0.67 (0.38-1.1)	1.6% (0.83-2.7%)	2.1% (1.1-3.5%)	510 (270-840)	5500 (2800-9400)
Mexico	2005	Female	85 to 89	19% (12-27%)	81% (73-88%)	0.66 (0.37-1.1)	1.1% (0.39-2%)	1.4% (0.5-2.6%)	270 (98-510)	2100 (730-4100)
Mexico	2005	Female	90 to 94	22% (14-30%)	78% (70-86%)	0.7 (0.36-1.2)	0.69% (-0.092-1.8%)	0.78% (-0.16-2.2%)	110 (-23-310)	670 (-90-1800)
Mexico	2005	Female	95 plus	25% (16-37%)	75% (63-84%)	0.75 (0.3-1.4)	0.48% (-0.52-2.1%)	0.41% (-0.83-2.3%)	26 (-52-150)	130 (-140-570)
Mexico	2005	Male	All Ages	50% (46-54%)	50% (46-54%)	3.6 (3.1-4.1)	10% (8.7-11%)	13% (11-14%)	35000 (30000-40000)	1400000 (1200000-1600000)
Mexico	2005	Male	15 to 19	40% (33-49%)	60% (51-67%)	1.7 (1.1-2.3)	6.2% (4.5-8.4%)	8.5% (5.8-12%)	440 (300-630)	43000 (31000-58000)
Mexico	2005	Male	20 to 24	66% (57-74%)	34% (26-43%)	3.9 (2.8-5.3)	14% (11-18%)	18% (13-23%)	1300 (930-1700)	110000 (87000-140000)
Mexico	2005	Male	25 to 29	72% (64-78%)	28% (22-36%)	4.9 (3.6-6.4)	16% (13-19%)	20% (16-26%)	1600 (1300-2100)	130000 (110000-160000)
Mexico	2005	Male	30 to 34	74% (67-81%)	26% (19-33%)	5.7 (4.3-7.3)	17% (14-19%)	22% (18-26%)	2000 (1700-2400)	140000 (120000-170000)
Mexico	2005	Male	35 to 39	75% (68-82%)	25% (18-32%)	6.2 (4.6-8.3)	17% (15-20%)	23% (20-28%)	2400 (2000-2800)	150000 (120000-170000)
Mexico	2005	Male	40 to 44	75% (68-82%)	25% (18-32%)	6.8 (5.1-8.8)	18% (15-21%)	24% (21-27%)	2700 (2400-3100)	150000 (130000-170000)
Mexico	2005	Male	45 to 49	76% (66-83%)	24% (17-34%)	7.2 (5.1-9.7)	18% (15-20%)	23% (21-26%)	3100 (2700-3500)	140000 (120000-170000)
Mexico	2005	Male	50 to 54	76% (67-83%)	24% (17-33%)	8.2 (5.9-11)	17% (14-20%)	22% (19-25%)	3300 (2800-3800)	140000 (110000-160000)
Mexico	2005	Male	55 to 59	76% (68-84%)	24% (16-32%)	10 (7.3-13)	16% (13-19%)	20% (17-24%)	3400 (2800-4000)	120000 (96000-150000)
Mexico	2005	Male	60 to 64	74% (64-83%)	26% (17-36%)	9.7 (6.9-13)	14% (11-17%)	17% (14-21%)	3400 (2700-4100)	100000 (79000-130000)
Mexico	2005	Male	65 to 69	68% (56-79%)	32% (21-44%)	6.9 (4.7-9.5)	12% (9.2-15%)	14% (11-18%)	3200 (2500-3900)	79000 (61000-98000)
Mexico	2005	Male	70 to 74	60% (48-71%)	40% (29-52%)	4.7 (3.1-6.6)	9.6% (7.3-12%)	11% (8.9-14%)	2800 (2200-3600)	56000 (43000-71000)
Mexico	2005	Male	75 to 79	52% (40-65%)	48% (35-60%)	3.5 (2.2-5.2)	7.4% (5.4-9.6%)	8.8% (6.5-11%)	2200 (1600-2900)	34000 (25000-45000)
Mexico	2005	Male	80 to 84	47% (34-58%)	53% (42-66%)	3 (2-4.4)	6.3% (4.6-8.2%)	7.3% (5.4-9.7%)	1700 (1300-2300)	20000 (15000-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	2005	Male	85 to 89	44% (32-55%)	56% (45-68%)	2.7 (1.8-4.1)	5.1% (3.7-6.9%)	5.9% (4.2-8%)	990 (700-1300)	8200 (5900-11000)
Mexico	2005	Male	90 to 94	46% (35-58%)	54% (42-65%)	2.8 (1.7-4.2)	4.6% (3-6.4%)	5.2% (3.3-7.3%)	540 (340-770)	3100 (2000-4400)
Mexico	2005	Male	95 plus	51% (36-65%)	49% (35-64%)	3 (1.7-4.7)	4.7% (2.5-7%)	5.1% (2.4-7.9%)	170 (80-260)	660 (350-990)
Mexico	2010	Female	All Ages	29% (25-32%)	71% (68-75%)	0.33 (0.27-0.41)	1.3% (0.88-1.7%)	2.4% (1.7-3.2%)	6000 (4200-8200)	150000 (110000-200000)
Mexico	2010	Female	15 to 19	24% (16-32%)	76% (68-84%)	0.26 (0.15-0.41)	0.77% (0.48-1.2%)	1.1% (0.46-2.1%)	30 (12-56)	4600 (2700-6800)
Mexico	2010	Female	20 to 24	35% (25-46%)	65% (54-75%)	0.32 (0.17-0.52)	1.1% (0.69-1.6%)	1.2% (0.49-2.2%)	36 (14-64)	6500 (4000-9700)
Mexico	2010	Female	25 to 29	38% (28-48%)	62% (52-72%)	0.32 (0.19-0.51)	0.97% (0.62-1.4%)	1.4% (0.78-2.3%)	45 (24-70)	5900 (3800-8400)
Mexico	2010	Female	30 to 34	39% (29-50%)	61% (50-71%)	0.34 (0.2-0.53)	0.92% (0.51-1.3%)	2.1% (1.6-2.8%)	77 (57-100)	6000 (3500-8400)
Mexico	2010	Female	35 to 39	39% (29-50%)	61% (50-71%)	0.39 (0.22-0.62)	1.1% (0.68-1.5%)	3.2% (2.6-3.9%)	150 (120-190)	7900 (5100-11000)
Mexico	2010	Female	40 to 44	40% (30-51%)	60% (49-70%)	0.47 (0.29-0.71)	1.3% (0.76-1.9%)	3.8% (3.2-4.5%)	230 (190-280)	9500 (6100-13000)
Mexico	2010	Female	45 to 49	40% (28-52%)	60% (48-72%)	0.61 (0.34-1)	1.6% (0.93-2.4%)	4% (3.2-5%)	350 (270-430)	12000 (7500-17000)
Mexico	2010	Female	50 to 54	40% (29-51%)	60% (49-71%)	0.75 (0.46-1.1)	1.9% (1.1-2.9%)	4.1% (3.2-5.2%)	480 (370-600)	15000 (8800-21000)
Mexico	2010	Female	55 to 59	38% (27-50%)	62% (50-73%)	0.83 (0.48-1.4)	2.3% (1.4-3.4%)	4.2% (3.1-5.6%)	610 (450-830)	18000 (11000-26000)
Mexico	2010	Female	60 to 64	34% (21-48%)	66% (52-79%)	0.85 (0.46-1.4)	2.4% (1.5-3.7%)	3.8% (2.7-5.3%)	670 (470-930)	17000 (11000-26000)
Mexico	2010	Female	65 to 69	32% (20-47%)	68% (53-80%)	0.91 (0.49-1.5)	2.6% (1.6-3.8%)	3.8% (2.7-5.3%)	800 (560-1100)	17000 (11000-26000)
Mexico	2010	Female	70 to 74	29% (18-40%)	71% (60-82%)	0.92 (0.51-1.6)	2.5% (1.3-4.1%)	3.5% (2.1-5.4%)	860 (520-1300)	15000 (8100-25000)
Mexico	2010	Female	75 to 79	23% (14-35%)	77% (65-86%)	0.78 (0.39-1.3)	2% (1-3.4%)	2.8% (1.5-4.3%)	780 (440-1200)	11000 (5600-18000)
Mexico	2010	Female	80 to 84	21% (13-31%)	79% (69-87%)	0.67 (0.34-1.1)	1.3% (0.55-2.3%)	1.8% (0.82-3%)	520 (230-860)	5500 (2300-9500)
Mexico	2010	Female	85 to 89	22% (14-34%)	78% (66-86%)	0.68 (0.35-1.2)	0.82% (0.088-1.8%)	1.1% (0.13-2.2%)	290 (36-600)	2200 (220-4800)
Mexico	2010	Female	90 to 94	27% (17-38%)	73% (62-83%)	0.76 (0.38-1.4)	0.39% (-0.46-1.5%)	0.42% (-0.69-1.8%)	72 (-120-310)	480 (-550-1800)
Mexico	2010	Female	95 plus	32% (19-46%)	68% (54-81%)	0.84 (0.34-1.6)	0.14% (-1-1.7%)	-0.05% (-1.5-2%)	-4.2 (-130-170)	55 (-380-640)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	2010	Male	All Ages	52% (49-57%)	48% (43-51%)	3.4 (2.9-3.8)	10% (8.9-12%)	12% (11-14%)	40000 (35000-46000)	1600000 (1400000-1900000)
Mexico	2010	Male	15 to 19	38% (30-47%)	62% (53-70%)	1.4 (0.9-1.9)	5.6% (4-7.7%)	7.2% (4.7-11%)	490 (330-720)	47000 (33000-64000)
Mexico	2010	Male	20 to 24	66% (57-74%)	34% (26-43%)	3.4 (2.4-4.6)	13% (10-17%)	16% (11-21%)	1600 (1100-2100)	130000 (99000-170000)
Mexico	2010	Male	25 to 29	73% (66-80%)	27% (20-34%)	4.4 (3.2-5.8)	15% (12-19%)	19% (14-24%)	2000 (1500-2600)	160000 (120000-190000)
Mexico	2010	Male	30 to 34	76% (68-82%)	24% (18-32%)	5.1 (3.8-6.5)	16% (13-19%)	21% (16-25%)	2400 (1900-2900)	160000 (130000-200000)
Mexico	2010	Male	35 to 39	76% (69-83%)	24% (17-31%)	5.6 (4.1-7.3)	17% (14-20%)	22% (18-26%)	2900 (2300-3400)	170000 (140000-200000)
Mexico	2010	Male	40 to 44	77% (69-84%)	23% (16-31%)	6 (4.5-7.8)	17% (14-20%)	22% (19-26%)	3100 (2600-3500)	160000 (140000-190000)
Mexico	2010	Male	45 to 49	78% (69-84%)	22% (16-31%)	6.5 (4.6-8.7)	17% (14-19%)	22% (19-25%)	3400 (2900-3900)	160000 (140000-190000)
Mexico	2010	Male	50 to 54	77% (69-84%)	23% (16-31%)	7.4 (5.4-9.7)	16% (13-19%)	21% (18-24%)	3800 (3200-4300)	160000 (130000-180000)
Mexico	2010	Male	55 to 59	76% (68-83%)	24% (17-32%)	8.8 (6.4-12)	15% (12-18%)	19% (16-22%)	3900 (3200-4600)	140000 (110000-170000)
Mexico	2010	Male	60 to 64	72% (61-82%)	28% (18-39%)	8 (5.6-11)	13% (10-16%)	16% (13-19%)	3600 (2900-4400)	110000 (85000-140000)
Mexico	2010	Male	65 to 69	68% (56-79%)	32% (21-44%)	6 (4.1-8.3)	11% (8.6-14%)	14% (11-17%)	3500 (2800-4300)	87000 (67000-110000)
Mexico	2010	Male	70 to 74	63% (51-74%)	37% (26-49%)	4.4 (2.9-6.2)	9.1% (6.9-12%)	11% (8.3-14%)	3100 (2400-4000)	62000 (47000-80000)
Mexico	2010	Male	75 to 79	55% (42-68%)	45% (32-58%)	3.3 (2-4.9)	7% (4.9-9.2%)	8.2% (5.7-11%)	2500 (1800-3300)	39000 (27000-52000)
Mexico	2010	Male	80 to 84	49% (36-61%)	51% (39-64%)	2.8 (1.8-4.1)	5.7% (4-7.7%)	6.6% (4.6-9%)	1900 (1300-2500)	21000 (15000-29000)
Mexico	2010	Male	85 to 89	47% (35-59%)	53% (41-65%)	2.6 (1.6-3.8)	4.8% (3.3-6.6%)	5.5% (3.7-7.7%)	1300 (880-1800)	11000 (7400-15000)
Mexico	2010	Male	90 to 94	50% (39-62%)	50% (38-61%)	2.7 (1.7-4.1)	4.2% (2.5-6.1%)	4.7% (2.7-6.9%)	590 (330-860)	3500 (2100-5000)
Mexico	2010	Male	95 plus	56% (41-70%)	44% (30-59%)	2.9 (1.7-4.6)	4.2% (1.9-6.6%)	4.5% (1.7-7.5%)	220 (83-370)	910 (420-1500)
Mexico	2016	Female	All Ages	39% (35-43%)	61% (57-65%)	0.43 (0.32-0.56)	1.2% (0.67-1.7%)	2.1% (1.2-3.1%)	6100 (3400-9200)	150000 (91000-230000)
Mexico	2016	Female	15 to 19	30% (19-42%)	70% (58-81%)	0.3 (0.15-0.51)	0.77% (0.44-1.2%)	1.3% (0.46-2.5%)	31 (11-61)	4500 (2500-7000)
Mexico	2016	Female	20 to 24	48% (35-58%)	52% (42-65%)	0.41 (0.22-0.67)	1.1% (0.69-1.7%)	1.5% (0.53-2.7%)	43 (16-79)	7100 (4000-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	2016	Female	25 to 29	54% (41-62%)	46% (38-59%)	0.43 (0.25-0.67)	0.98% (0.57-1.5%)	1.6% (0.72-2.7%)	50 (22-84)	6200 (3500-9200)
Mexico	2016	Female	30 to 34	53% (41-61%)	47% (39-59%)	0.44 (0.26-0.68)	0.87% (0.41-1.3%)	2.1% (1.4-3%)	78 (52-110)	5800 (3000-8700)
Mexico	2016	Female	35 to 39	52% (43-61%)	48% (39-57%)	0.48 (0.28-0.78)	0.98% (0.47-1.5%)	3.1% (2.4-4%)	150 (120-200)	7200 (3700-11000)
Mexico	2016	Female	40 to 44	51% (42-61%)	49% (39-58%)	0.57 (0.35-0.86)	1% (0.44-1.7%)	3.5% (2.8-4.4%)	240 (190-310)	8700 (3900-13000)
Mexico	2016	Female	45 to 49	49% (38-59%)	51% (41-62%)	0.69 (0.4-1.1)	1.3% (0.53-2.1%)	3.7% (2.8-4.8%)	360 (270-480)	11000 (5100-18000)
Mexico	2016	Female	50 to 54	48% (38-57%)	52% (43-62%)	0.82 (0.49-1.2)	1.6% (0.68-2.6%)	3.7% (2.6-5%)	500 (360-680)	14000 (6500-23000)
Mexico	2016	Female	55 to 59	47% (35-57%)	53% (43-65%)	0.95 (0.55-1.5)	1.8% (0.8-3.2%)	3.7% (2.5-5.4%)	650 (440-930)	17000 (8000-29000)
Mexico	2016	Female	60 to 64	45% (29-61%)	55% (39-71%)	1 (0.55-1.7)	1.9% (0.73-3.5%)	3.4% (2-5.3%)	770 (440-1200)	18000 (6800-32000)
Mexico	2016	Female	65 to 69	41% (25-57%)	59% (43-75%)	1.1 (0.53-1.7)	2.1% (0.92-3.6%)	3.4% (1.9-5.1%)	850 (480-1300)	18000 (7800-30000)
Mexico	2016	Female	70 to 74	35% (20-49%)	65% (51-80%)	1 (0.49-1.8)	2.2% (0.86-3.8%)	3.2% (1.6-5.2%)	890 (440-1500)	16000 (6100-28000)
Mexico	2016	Female	75 to 79	28% (15-43%)	72% (57-85%)	0.83 (0.36-1.5)	1.8% (0.57-3.2%)	2.5% (1-4.2%)	780 (330-1300)	11000 (3600-20000)
Mexico	2016	Female	80 to 84	25% (14-39%)	75% (61-86%)	0.74 (0.33-1.3)	1.1% (0.052-2.2%)	1.5% (0.22-2.8%)	520 (76-980)	5400 (240-11000)
Mexico	2016	Female	85 to 89	28% (15-43%)	72% (57-85%)	0.77 (0.34-1.4)	0.54% (-0.41-1.6%)	0.73% (-0.48-2%)	240 (-160-640)	1800 (-1300-5200)
Mexico	2016	Female	90 to 94	33% (19-47%)	67% (53-81%)	0.85 (0.37-1.6)	0.092% (-0.94-1.3%)	0.045% (-1.3-1.6%)	11 (-300-380)	160 (-1600-2200)
Mexico	2016	Female	95 plus	38% (22-53%)	62% (47-78%)	0.91 (0.34-1.8)	-0.17% (-1.5-1.5%)	-0.44% (-2.1-1.7%)	-53 (-250-210)	-93 (-780-820)
Mexico	2016	Male	All Ages	57% (53-61%)	43% (39-47%)	3.4 (2.6-4.3)	11% (9-12%)	12% (9.9-14%)	43000 (36000-51000)	1700000 (1500000-2000000)
Mexico	2016	Male	15 to 19	40% (29-51%)	60% (49-71%)	1.3 (0.79-2)	5.4% (3.8-7.6%)	7.3% (4.3-11%)	430 (260-650)	42000 (29000-60000)
Mexico	2016	Male	20 to 24	69% (60-77%)	31% (23-40%)	3.3 (2.2-4.6)	13% (9.8-17%)	16% (11-22%)	1500 (1000-2100)	130000 (96000-170000)
Mexico	2016	Male	25 to 29	77% (69-83%)	23% (17-31%)	4.3 (2.9-5.9)	15% (12-19%)	19% (14-24%)	1900 (1400-2400)	150000 (110000-190000)
Mexico	2016	Male	30 to 34	78% (71-84%)	22% (16-29%)	4.9 (3.5-6.7)	16% (12-19%)	20% (16-25%)	2200 (1700-2700)	160000 (120000-190000)
Mexico	2016	Male	35 to 39	78% (72-84%)	22% (16-28%)	5.3 (3.6-7.6)	16% (13-20%)	22% (17-26%)	2700 (2200-3300)	170000 (140000-200000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mexico	2016	Male	40 to 44	79% (72-85%)	21% (15-28%)	5.7 (4-7.8)	16% (13-19%)	22% (18-26%)	3200 (2700-3800)	180000 (150000-210000)
Mexico	2016	Male	45 to 49	79% (71-85%)	21% (15-29%)	6 (4.1-8.3)	16% (13-19%)	21% (18-24%)	3800 (3100-4400)	180000 (150000-210000)
Mexico	2016	Male	50 to 54	78% (70-84%)	22% (16-30%)	6.8 (4.8-9.4)	15% (13-18%)	20% (17-23%)	4200 (3600-4900)	180000 (150000-210000)
Mexico	2016	Male	55 to 59	78% (70-84%)	22% (16-30%)	8.2 (5.7-11)	15% (12-18%)	18% (15-22%)	4500 (3700-5400)	160000 (130000-200000)
Mexico	2016	Male	60 to 64	75% (64-84%)	25% (16-36%)	7.7 (5.2-11)	13% (9.8-16%)	16% (13-19%)	4400 (3600-5400)	140000 (110000-170000)
Mexico	2016	Male	65 to 69	69% (56-80%)	31% (20-44%)	5.5 (3.6-8.1)	11% (8.1-13%)	13% (10-16%)	4000 (3100-4900)	100000 (77000-130000)
Mexico	2016	Male	70 to 74	60% (47-71%)	40% (29-53%)	3.7 (2.2-5.5)	8% (5.8-11%)	9.6% (7.1-13%)	3200 (2400-4300)	65000 (48000-87000)
Mexico	2016	Male	75 to 79	50% (36-64%)	50% (36-64%)	2.6 (1.5-4.1)	5.9% (3.9-8.3%)	7% (4.6-9.8%)	2500 (1700-3600)	39000 (26000-55000)
Mexico	2016	Male	80 to 84	46% (32-59%)	54% (41-68%)	2.3 (1.3-3.7)	4.9% (3.1-7.1%)	5.7% (3.7-8.3%)	2000 (1300-2900)	23000 (15000-33000)
Mexico	2016	Male	85 to 89	47% (34-60%)	53% (40-66%)	2.3 (1.3-3.8)	4.2% (2.5-6.2%)	4.9% (2.9-7.2%)	1300 (790-2000)	11000 (6700-17000)
Mexico	2016	Male	90 to 94	52% (39-64%)	48% (36-61%)	2.6 (1.5-4)	3.8% (1.9-6%)	4.2% (1.9-6.9%)	740 (340-1200)	4400 (2200-6900)
Mexico	2016	Male	95 plus	57% (41-71%)	43% (29-59%)	2.8 (1.5-4.5)	3.7% (1.4-6.5%)	4% (1.1-7.3%)	270 (73-490)	1100 (400-1900)
Nicaragua	1990	Female	All Ages	14% (11-17%)	86% (83-89%)	0.38 (0.3-0.48)	1% (0.81-1.3%)	2% (1.5-2.6%)	150 (110-200)	6200 (4900-7700)
Nicaragua	1990	Female	15 to 19	15% (9.2-22%)	85% (78-91%)	0.37 (0.19-0.63)	1.3% (0.76-2%)	1.9% (0.9-3.5%)	2.7 (1.3-5)	350 (210-560)
Nicaragua	1990	Female	20 to 24	26% (16-36%)	74% (64-84%)	0.56 (0.3-0.93)	2.2% (1.3-3.1%)	2.8% (1.4-4.7%)	4.3 (2.1-7.3)	540 (320-790)
Nicaragua	1990	Female	25 to 29	25% (16-35%)	75% (65-84%)	0.57 (0.31-0.9)	2.2% (1.5-3.1%)	3.1% (1.7-4.9%)	4.2 (2.3-6.6)	480 (310-680)
Nicaragua	1990	Female	30 to 34	25% (16-36%)	75% (64-84%)	0.62 (0.35-1)	2.3% (1.6-3.2%)	3.6% (2.3-5.3%)	5.2 (3.3-7.8)	480 (340-680)
Nicaragua	1990	Female	35 to 39	25% (16-34%)	75% (66-84%)	0.72 (0.37-1.2)	2.6% (1.7-3.7%)	4.2% (2.7-6.2%)	7.3 (4.6-11)	520 (340-740)
Nicaragua	1990	Female	40 to 44	25% (17-35%)	75% (65-83%)	0.84 (0.42-1.3)	3.1% (2-4.4%)	5.4% (3.4-7.8%)	7.9 (4.9-11)	450 (290-640)
Nicaragua	1990	Female	45 to 49	28% (17-37%)	72% (63-83%)	1.1 (0.52-1.8)	3.7% (2.2-5.7%)	6.5% (3.9-9.8%)	11 (6.6-17)	540 (330-840)
Nicaragua	1990	Female	50 to 54	26% (17-36%)	74% (64-83%)	1.2 (0.62-2.1)	4.2% (2.6-6.3%)	7.2% (4.6-11%)	14 (8.9-20)	580 (370-870)
Nicaragua	1990	Female	55 to 59	20% (13-28%)	80% (72-87%)	1.2 (0.58-2)	4.4% (2.7-6.4%)	7.1% (4.6-10%)	18 (11-26)	620 (400-920)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	1990	Female	60 to 64	15% (9.2-22%)	85% (78-91%)	1 (0.49-1.8)	3.8% (2.4-5.7%)	6.1% (4-8.8%)	17 (11-25)	520 (340-770)
Nicaragua	1990	Female	65 to 69	13% (8-18%)	87% (82-92%)	0.83 (0.41-1.5)	3.4% (2.1-4.9%)	5.3% (3.4-7.7%)	16 (11-24)	410 (260-610)
Nicaragua	1990	Female	70 to 74	12% (7.5-17%)	88% (83-92%)	0.69 (0.34-1.2)	2.9% (1.8-4.5%)	4.5% (2.8-6.8%)	15 (9.3-23)	300 (180-460)
Nicaragua	1990	Female	75 to 79	11% (6.4-18%)	89% (82-94%)	0.58 (0.27-1.1)	2.3% (1.3-3.8%)	3.3% (1.9-5.3%)	12 (7-20)	190 (110-310)
Nicaragua	1990	Female	80 to 84	10% (5.5-18%)	90% (82-94%)	0.51 (0.24-0.9)	1.9% (1-3.3%)	2.5% (1.4-4.3%)	7.9 (4.3-14)	89 (48-150)
Nicaragua	1990	Female	85 to 89	11% (5.6-17%)	89% (83-94%)	0.5 (0.23-0.94)	1.6% (0.79-3.1%)	2.1% (0.97-4%)	6.6 (3.1-13)	52 (25-99)
Nicaragua	1990	Female	90 to 94	12% (6.9-20%)	88% (80-93%)	0.54 (0.24-1.1)	1.4% (0.51-3.1%)	1.7% (0.53-3.9%)	3.6 (1.1-8.1)	19 (6.8-42)
Nicaragua	1990	Female	95 plus	15% (8-27%)	85% (73-92%)	0.6 (0.22-1.2)	1.4% (0.19-3.4%)	1.6% (0.051-4.3%)	1.2 (0.039-3.3)	4.6 (0.62-11)
Nicaragua	1990	Male	All Ages	44% (35-53%)	56% (47-65%)	1.6 (1.4-1.9)	4.4% (3.8-5%)	6.6% (5.7-7.7%)	650 (560-760)	32000 (27000-37000)
Nicaragua	1990	Male	15 to 19	39% (25-55%)	61% (45-75%)	1 (0.49-1.7)	4.8% (2.9-7.2%)	6% (3.1-10%)	17 (8.6-27)	1600 (980-2500)
Nicaragua	1990	Male	20 to 24	67% (51-79%)	33% (21-49%)	2.5 (1.4-3.9)	12% (7.9-16%)	14% (8.4-21%)	46 (27-68)	4000 (2600-5500)
Nicaragua	1990	Male	25 to 29	75% (60-86%)	25% (14-40%)	3.3 (1.9-5.1)	15% (11-20%)	19% (12-26%)	59 (38-83)	4600 (3200-6100)
Nicaragua	1990	Male	30 to 34	75% (60-86%)	25% (14-40%)	3.6 (2.2-5.5)	16% (12-20%)	21% (15-27%)	57 (41-73)	4100 (3100-5100)
Nicaragua	1990	Male	35 to 39	71% (57-82%)	29% (18-43%)	3.8 (2.1-6.1)	17% (13-21%)	23% (17-29%)	59 (43-76)	3700 (2800-4600)
Nicaragua	1990	Male	40 to 44	70% (56-82%)	30% (18-44%)	4 (2.1-6.4)	17% (14-21%)	24% (18-29%)	54 (41-67)	2900 (2300-3500)
Nicaragua	1990	Male	45 to 49	72% (57-84%)	28% (16-43%)	4.3 (2.2-7.6)	17% (13-20%)	23% (18-28%)	58 (45-71)	2700 (2100-3300)
Nicaragua	1990	Male	50 to 54	73% (59-84%)	27% (16-41%)	4.5 (2.3-7.6)	15% (11-18%)	21% (17-26%)	53 (42-65)	2200 (1800-2700)
Nicaragua	1990	Male	55 to 59	71% (56-83%)	29% (17-44%)	4.5 (2.3-7.8)	13% (10-16%)	18% (14-22%)	56 (43-69)	2000 (1500-2500)
Nicaragua	1990	Male	60 to 64	66% (49-80%)	34% (20-51%)	3.7 (1.8-6.7)	11% (7.9-13%)	14% (11-18%)	53 (40-67)	1600 (1200-2000)
Nicaragua	1990	Male	65 to 69	60% (42-77%)	40% (23-58%)	2.7 (1.4-4.7)	8.6% (6.3-11%)	12% (8.4-15%)	42 (31-55)	1100 (790-1400)
Nicaragua	1990	Male	70 to 74	54% (35-70%)	46% (30-65%)	2 (0.94-3.5)	6.9% (4.5-9.4%)	8.9% (5.8-12%)	36 (24-49)	710 (470-990)
Nicaragua	1990	Male	75 to 79	47% (29-65%)	53% (35-71%)	1.5 (0.67-2.9)	5.3% (3.1-7.7%)	6.6% (3.8-9.4%)	27 (16-39)	410 (240-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	1990	Male	80 to 84	42% (25-59%)	58% (41-75%)	1.3 (0.6-2.2)	4.4% (2.6-6.4%)	5.3% (3.1-7.6%)	16 (9.5-24)	180 (110-270)
Nicaragua	1990	Male	85 to 89	40% (24-58%)	60% (42-76%)	1.2 (0.55-2.3)	3.9% (2-6.1%)	4.5% (2.3-7.2%)	12 (6-19)	93 (50-140)
Nicaragua	1990	Male	90 to 94	43% (27-61%)	57% (39-73%)	1.4 (0.56-2.7)	3.6% (1.5-6.1%)	4.1% (1.6-7.1%)	6.1 (2.4-11)	34 (14-57)
Nicaragua	1990	Male	95 plus	49% (28-70%)	51% (30-72%)	1.7 (0.56-3.7)	3.9% (1.2-6.9%)	4.3% (1.1-7.9%)	2 (0.48-3.6)	7.4 (2.2-13)
Nicaragua	1995	Female	All Ages	14% (11-17%)	86% (83-89%)	0.37 (0.29-0.47)	1.2% (0.95-1.5%)	2.3% (1.7-3.1%)	200 (140-260)	7300 (5800-9100)
Nicaragua	1995	Female	15 to 19	15% (9.2-21%)	85% (79-91%)	0.36 (0.18-0.6)	1.2% (0.76-2%)	1.8% (0.84-3.3%)	3.2 (1.5-6)	400 (240-650)
Nicaragua	1995	Female	20 to 24	26% (17-36%)	74% (64-83%)	0.54 (0.29-0.91)	2.1% (1.3-3%)	2.6% (1.3-4.4%)	4.6 (2.2-7.9)	600 (370-900)
Nicaragua	1995	Female	25 to 29	25% (17-35%)	75% (65-83%)	0.55 (0.3-0.9)	2.1% (1.5-2.9%)	2.9% (1.6-4.5%)	4.4 (2.5-7)	520 (340-740)
Nicaragua	1995	Female	30 to 34	25% (16-36%)	75% (64-84%)	0.6 (0.33-0.95)	2.2% (1.6-3.1%)	3.4% (2.2-5%)	5.7 (3.7-8.3)	520 (360-720)
Nicaragua	1995	Female	35 to 39	24% (15-34%)	76% (66-85%)	0.68 (0.36-1.1)	2.5% (1.7-3.5%)	4.1% (2.7-6.1%)	8.2 (5.4-12)	580 (400-850)
Nicaragua	1995	Female	40 to 44	25% (17-35%)	75% (65-83%)	0.79 (0.41-1.3)	2.9% (2-4.1%)	5.3% (3.5-7.4%)	11 (7.1-15)	630 (420-870)
Nicaragua	1995	Female	45 to 49	28% (17-38%)	72% (62-83%)	0.99 (0.49-1.7)	3.8% (2.4-5.7%)	6.3% (4-9.6%)	13 (8.3-21)	630 (390-950)
Nicaragua	1995	Female	50 to 54	26% (17-36%)	74% (64-83%)	1.1 (0.59-2)	4.2% (2.6-6.4%)	7% (4.5-10%)	17 (11-24)	680 (440-1000)
Nicaragua	1995	Female	55 to 59	20% (13-28%)	80% (72-87%)	1.1 (0.54-1.9)	4.3% (2.7-6.3%)	6.9% (4.4-9.9%)	19 (12-28)	680 (430-990)
Nicaragua	1995	Female	60 to 64	15% (9.2-23%)	85% (77-91%)	0.96 (0.46-1.8)	4% (2.5-6%)	6.1% (4-8.9%)	22 (14-32)	640 (420-950)
Nicaragua	1995	Female	65 to 69	13% (8.2-18%)	87% (82-92%)	0.8 (0.4-1.4)	3.5% (2.3-5.3%)	5.3% (3.5-7.8%)	22 (14-32)	540 (350-800)
Nicaragua	1995	Female	70 to 74	12% (7.7-17%)	88% (83-92%)	0.67 (0.33-1.2)	3% (1.8-4.7%)	4.4% (2.8-6.8%)	21 (13-32)	400 (250-630)
Nicaragua	1995	Female	75 to 79	11% (6.5-18%)	89% (82-94%)	0.57 (0.25-1)	2.4% (1.3-3.8%)	3.3% (1.9-5.4%)	16 (9.1-26)	240 (140-390)
Nicaragua	1995	Female	80 to 84	11% (5.7-18%)	89% (82-94%)	0.5 (0.24-0.88)	2% (1.1-3.4%)	2.6% (1.4-4.4%)	11 (6-19)	120 (67-210)
Nicaragua	1995	Female	85 to 89	11% (5.8-18%)	89% (82-94%)	0.49 (0.23-0.94)	1.7% (0.77-3.1%)	2.1% (0.95-3.9%)	9.7 (4.4-19)	75 (36-140)
Nicaragua	1995	Female	90 to 94	13% (7.2-20%)	87% (80-93%)	0.53 (0.24-1)	1.4% (0.46-3.2%)	1.7% (0.48-3.9%)	5.6 (1.6-13)	30 (9.6-67)
Nicaragua	1995	Female	95 plus	16% (8-28%)	84% (72-92%)	0.58 (0.21-1.2)	1.4% (0.13-3.6%)	1.5% (-0.006-4.3%)	2 (-0.0078-5.5)	7.2 (0.7-19)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	1995	Male	All Ages	45% (37-54%)	55% (46-63%)	1.6 (1.3-1.8)	5.4% (4.8-6.1%)	8% (6.9-9.1%)	870 (750-1000)	41000 (35000-47000)
Nicaragua	1995	Male	15 to 19	40% (25-56%)	60% (44-75%)	0.98 (0.48-1.7)	4.8% (2.9-7.2%)	6.1% (3.2-10%)	18 (9.6-30)	1800 (1100-2700)
Nicaragua	1995	Male	20 to 24	68% (52-80%)	32% (20-48%)	2.4 (1.4-3.8)	12% (8.1-16%)	15% (8.8-21%)	53 (32-78)	4600 (3100-6300)
Nicaragua	1995	Male	25 to 29	76% (61-86%)	24% (14-39%)	3.2 (1.9-4.9)	15% (11-20%)	19% (13-26%)	61 (42-84)	4800 (3500-6400)
Nicaragua	1995	Male	30 to 34	76% (61-86%)	24% (14-39%)	3.5 (2.1-5.3)	17% (13-21%)	22% (16-27%)	74 (56-95)	5100 (3900-6400)
Nicaragua	1995	Male	35 to 39	71% (57-83%)	29% (17-43%)	3.7 (2-5.9)	18% (14-22%)	24% (18-29%)	86 (65-110)	5200 (4100-6300)
Nicaragua	1995	Male	40 to 44	70% (56-82%)	30% (18-44%)	3.9 (2-6.1)	18% (14-22%)	24% (19-29%)	85 (66-100)	4500 (3500-5500)
Nicaragua	1995	Male	45 to 49	73% (57-84%)	27% (16-43%)	4.1 (2-7.2)	18% (14-21%)	23% (18-28%)	75 (58-90)	3400 (2700-4100)
Nicaragua	1995	Male	50 to 54	73% (59-84%)	27% (16-41%)	4.3 (2.3-7.4)	16% (12-19%)	21% (17-25%)	75 (58-91)	3000 (2400-3600)
Nicaragua	1995	Male	55 to 59	71% (57-84%)	29% (16-43%)	4.4 (2.2-7.6)	14% (11-17%)	18% (15-22%)	73 (57-90)	2500 (2000-3100)
Nicaragua	1995	Male	60 to 64	67% (49-81%)	33% (19-51%)	3.6 (1.8-6.5)	11% (8.4-14%)	14% (11-18%)	69 (52-88)	2000 (1500-2600)
Nicaragua	1995	Male	65 to 69	61% (42-77%)	39% (23-58%)	2.6 (1.4-4.5)	9% (6.7-12%)	12% (8.6-15%)	61 (44-78)	1500 (1100-1900)
Nicaragua	1995	Male	70 to 74	54% (37-71%)	46% (29-63%)	1.9 (0.93-3.4)	7.1% (4.6-9.6%)	9% (5.9-12%)	50 (32-68)	980 (640-1300)
Nicaragua	1995	Male	75 to 79	48% (30-66%)	52% (34-70%)	1.5 (0.63-2.8)	5.3% (3.1-7.8%)	6.6% (3.7-9.6%)	36 (20-52)	540 (310-780)
Nicaragua	1995	Male	80 to 84	42% (25-60%)	58% (40-75%)	1.2 (0.57-2.2)	4.4% (2.6-6.5%)	5.2% (2.9-7.8%)	24 (14-36)	270 (150-400)
Nicaragua	1995	Male	85 to 89	40% (24-58%)	60% (42-76%)	1.2 (0.52-2.2)	3.9% (2-6.2%)	4.5% (2.3-7.3%)	17 (8.4-27)	130 (69-210)
Nicaragua	1995	Male	90 to 94	44% (28-62%)	56% (38-72%)	1.3 (0.54-2.7)	3.7% (1.5-6.2%)	4.1% (1.6-7.2%)	8.5 (3.3-15)	46 (19-78)
Nicaragua	1995	Male	95 plus	49% (29-70%)	51% (30-71%)	1.7 (0.52-3.5)	3.9% (1.1-7%)	4.3% (0.97-8.1%)	2.8 (0.64-5.3)	10 (3-19)
Nicaragua	2000	Female	All Ages	13% (11-16%)	87% (84-89%)	0.33 (0.26-0.41)	1.3% (1.1-1.6%)	2.6% (1.9-3.4%)	210 (150-280)	7600 (6000-9300)
Nicaragua	2000	Female	15 to 19	15% (10-20%)	85% (80-90%)	0.34 (0.18-0.54)	1.2% (0.7-1.8%)	1.7% (0.75-3.2%)	3.2 (1.4-6)	420 (250-650)
Nicaragua	2000	Female	20 to 24	25% (18-33%)	75% (67-82%)	0.5 (0.29-0.79)	1.9% (1.2-2.8%)	2.3% (1.1-4%)	3.9 (1.9-6.7)	600 (370-870)
Nicaragua	2000	Female	25 to 29	24% (17-31%)	76% (69-83%)	0.48 (0.29-0.74)	1.9% (1.4-2.6%)	2.5% (1.5-3.8%)	3.8 (2.4-5.8)	520 (360-740)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	2000	Female	30 to 34	21% (15-29%)	79% (71-85%)	0.48 (0.28-0.75)	2% (1.4-2.6%)	3% (2.1-4.1%)	4.7 (3.3-6.6)	480 (350-650)
Nicaragua	2000	Female	35 to 39	20% (14-28%)	80% (72-86%)	0.51 (0.28-0.86)	2.1% (1.5-2.9%)	3.7% (2.6-5.2%)	6.7 (4.7-9.4)	510 (370-690)
Nicaragua	2000	Female	40 to 44	20% (14-28%)	80% (72-86%)	0.58 (0.31-0.95)	2.5% (1.8-3.3%)	4.6% (3.3-6.2%)	10 (7.3-14)	610 (440-810)
Nicaragua	2000	Female	45 to 49	22% (15-30%)	78% (70-85%)	0.74 (0.38-1.3)	3.1% (2.1-4.5%)	5.5% (3.8-7.7%)	14 (10-21)	710 (480-1000)
Nicaragua	2000	Female	50 to 54	21% (14-29%)	79% (71-86%)	0.85 (0.44-1.5)	3.7% (2.5-5.5%)	6.2% (4.3-8.9%)	15 (10-22)	610 (420-880)
Nicaragua	2000	Female	55 to 59	16% (11-23%)	84% (77-89%)	0.87 (0.44-1.5)	3.9% (2.6-5.6%)	6.2% (4.2-8.7%)	20 (14-29)	710 (470-1000)
Nicaragua	2000	Female	60 to 64	14% (8.5-20%)	86% (80-92%)	0.84 (0.42-1.5)	3.8% (2.5-5.6%)	5.9% (4-8.3%)	23 (16-33)	680 (460-980)
Nicaragua	2000	Female	65 to 69	13% (8-18%)	87% (82-92%)	0.76 (0.39-1.3)	3.6% (2.4-5.3%)	5.4% (3.7-7.8%)	25 (17-36)	610 (410-890)
Nicaragua	2000	Female	70 to 74	12% (7.6-17%)	88% (83-92%)	0.64 (0.31-1.2)	3% (1.9-4.7%)	4.5% (2.9-6.8%)	24 (15-36)	460 (290-710)
Nicaragua	2000	Female	75 to 79	11% (6.2-19%)	89% (81-94%)	0.54 (0.24-1)	2.4% (1.4-3.8%)	3.4% (2-5.3%)	20 (12-31)	300 (180-480)
Nicaragua	2000	Female	80 to 84	10% (5.4-18%)	90% (82-95%)	0.47 (0.22-0.82)	2% (1.1-3.3%)	2.7% (1.5-4.4%)	15 (8.5-24)	160 (93-260)
Nicaragua	2000	Female	85 to 89	11% (5.8-17%)	89% (83-94%)	0.46 (0.21-0.88)	1.6% (0.78-3%)	2% (0.94-3.8%)	12 (5.6-22)	92 (45-170)
Nicaragua	2000	Female	90 to 94	12% (7-20%)	88% (80-93%)	0.49 (0.22-0.96)	1.4% (0.48-3%)	1.6% (0.49-3.6%)	7.2 (2.2-17)	39 (14-85)
Nicaragua	2000	Female	95 plus	15% (7.6-27%)	85% (73-92%)	0.54 (0.2-1.1)	1.2% (0.15-3.3%)	1.4% (0.024-4%)	2.7 (0.045-7.9)	10 (1.3-27)
Nicaragua	2000	Male	All Ages	52% (44-61%)	48% (39-56%)	1.8 (1.5-2.1)	6.6% (5.8-7.4%)	9% (7.8-10%)	950 (820-1100)	45000 (39000-52000)
Nicaragua	2000	Male	15 to 19	53% (37-67%)	47% (33-63%)	1.4 (0.73-2.2)	5.9% (3.6-8.6%)	8.2% (4.5-13%)	27 (15-43)	2500 (1500-3700)
Nicaragua	2000	Male	20 to 24	77% (64-85%)	23% (15-36%)	3 (1.8-4.6)	13% (9.2-18%)	17% (11-24%)	65 (40-92)	5600 (3700-7600)
Nicaragua	2000	Male	25 to 29	81% (71-88%)	19% (12-29%)	3.7 (2.3-5.6)	16% (12-20%)	21% (15-28%)	69 (48-93)	5600 (4200-7200)
Nicaragua	2000	Male	30 to 34	79% (68-87%)	21% (13-32%)	3.8 (2.5-5.6)	17% (13-21%)	22% (17-28%)	74 (56-93)	5300 (4100-6500)
Nicaragua	2000	Male	35 to 39	72% (62-82%)	28% (18-38%)	3.7 (2.1-5.8)	18% (14-21%)	23% (18-29%)	85 (64-100)	5200 (4100-6300)
Nicaragua	2000	Male	40 to 44	71% (60-81%)	29% (19-40%)	3.9 (2.1-6)	17% (14-21%)	24% (19-28%)	94 (73-110)	5000 (4000-6000)
Nicaragua	2000	Male	45 to 49	74% (62-83%)	26% (17-38%)	4.2 (2.2-7.1)	17% (14-20%)	23% (18-27%)	99 (78-120)	4600 (3700-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	2000	Male	50 to 54	73% (61-83%)	27% (17-39%)	4.3 (2.3-7.2)	15% (12-18%)	20% (16-24%)	77 (62-93)	3100 (2500-3700)
Nicaragua	2000	Male	55 to 59	70% (56-82%)	30% (18-44%)	4.2 (2.1-7.3)	13% (10-16%)	17% (14-21%)	75 (59-91)	2600 (2100-3200)
Nicaragua	2000	Male	60 to 64	65% (48-79%)	35% (21-52%)	3.4 (1.6-6.1)	10% (7.7-13%)	13% (10-16%)	66 (51-84)	2000 (1500-2500)
Nicaragua	2000	Male	65 to 69	60% (41-76%)	40% (24-59%)	2.5 (1.3-4.3)	8.4% (6.3-11%)	11% (8.1-14%)	65 (48-84)	1600 (1200-2000)
Nicaragua	2000	Male	70 to 74	54% (36-70%)	46% (30-64%)	1.9 (0.9-3.4)	6.7% (4.4-9.2%)	8.6% (5.6-12%)	54 (35-75)	1100 (710-1500)
Nicaragua	2000	Male	75 to 79	48% (30-66%)	52% (34-70%)	1.4 (0.62-2.8)	5% (2.9-7.3%)	6.3% (3.7-9.1%)	38 (23-56)	590 (350-860)
Nicaragua	2000	Male	80 to 84	43% (26-60%)	57% (40-74%)	1.2 (0.57-2.1)	4.2% (2.4-6.1%)	5% (2.9-7.4%)	27 (15-40)	310 (180-450)
Nicaragua	2000	Male	85 to 89	41% (25-59%)	59% (41-75%)	1.2 (0.53-2.2)	3.7% (2-5.9%)	4.3% (2.2-6.8%)	20 (10-32)	160 (83-250)
Nicaragua	2000	Male	90 to 94	44% (28-62%)	56% (38-72%)	1.3 (0.54-2.6)	3.5% (1.5-5.9%)	4% (1.5-6.9%)	11 (4.1-18)	58 (25-97)
Nicaragua	2000	Male	95 plus	49% (29-71%)	51% (29-71%)	1.6 (0.53-3.5)	3.7% (1.1-6.8%)	4.1% (0.97-8%)	3.5 (0.82-6.8)	13 (4-24)
Nicaragua	2005	Female	All Ages	13% (11-16%)	87% (84-89%)	0.36 (0.28-0.44)	1.6% (1.3-2%)	3% (2.2-4%)	260 (190-340)	8600 (6800-11000)
Nicaragua	2005	Female	15 to 19	14% (9.8-20%)	86% (80-90%)	0.34 (0.19-0.56)	1.1% (0.69-1.7%)	1.7% (0.77-3.2%)	2.6 (1.1-4.8)	380 (230-570)
Nicaragua	2005	Female	20 to 24	25% (18-33%)	75% (67-82%)	0.51 (0.3-0.82)	1.9% (1.2-2.6%)	2.3% (1.1-3.9%)	3.7 (1.9-6.3)	620 (390-910)
Nicaragua	2005	Female	25 to 29	23% (17-30%)	77% (70-83%)	0.49 (0.29-0.75)	1.9% (1.3-2.5%)	2.4% (1.5-3.5%)	3.4 (2-5)	540 (360-760)
Nicaragua	2005	Female	30 to 34	21% (15-29%)	79% (71-85%)	0.49 (0.28-0.77)	1.9% (1.4-2.5%)	3% (2.1-4.1%)	4.5 (3.2-6.1)	510 (360-690)
Nicaragua	2005	Female	35 to 39	19% (13-27%)	81% (73-87%)	0.52 (0.27-0.87)	2.1% (1.5-2.8%)	3.8% (2.7-5.2%)	6.9 (4.9-9.5)	540 (390-740)
Nicaragua	2005	Female	40 to 44	20% (13-27%)	80% (73-87%)	0.59 (0.31-0.97)	2.5% (1.8-3.3%)	4.8% (3.4-6.3%)	10 (7.2-14)	620 (450-820)
Nicaragua	2005	Female	45 to 49	22% (14-30%)	78% (70-86%)	0.75 (0.39-1.3)	3% (2.1-4.2%)	5.5% (3.9-7.8%)	16 (11-22)	780 (540-1100)
Nicaragua	2005	Female	50 to 54	20% (14-29%)	80% (71-86%)	0.86 (0.44-1.5)	3.6% (2.5-5.2%)	6.3% (4.5-8.8%)	21 (15-30)	890 (620-1300)
Nicaragua	2005	Female	55 to 59	16% (10-23%)	84% (77-90%)	0.88 (0.45-1.5)	4.1% (2.8-5.8%)	6.4% (4.4-8.8%)	24 (17-33)	830 (570-1200)
Nicaragua	2005	Female	60 to 64	13% (8.5-20%)	87% (80-92%)	0.85 (0.42-1.5)	3.9% (2.6-5.5%)	5.9% (4.2-8.3%)	26 (18-36)	770 (540-1100)
Nicaragua	2005	Female	65 to 69	12% (7.9-18%)	88% (82-92%)	0.77 (0.39-1.3)	3.7% (2.5-5.3%)	5.5% (3.9-7.7%)	29 (20-41)	690 (480-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	2005	Female	70 to 74	12% (7.6-17%)	88% (83-92%)	0.66 (0.33-1.2)	3.2% (2.1-4.8%)	4.8% (3.2-7%)	28 (19-42)	550 (360-830)
Nicaragua	2005	Female	75 to 79	11% (6.5-18%)	89% (82-94%)	0.57 (0.26-1)	2.6% (1.5-4%)	3.7% (2.3-5.5%)	25 (16-39)	380 (230-590)
Nicaragua	2005	Female	80 to 84	10% (5.6-18%)	90% (82-94%)	0.49 (0.23-0.87)	2.2% (1.3-3.5%)	2.9% (1.7-4.7%)	22 (13-36)	240 (140-390)
Nicaragua	2005	Female	85 to 89	10% (5.7-17%)	90% (83-94%)	0.47 (0.22-0.91)	1.7% (0.88-3.1%)	2.2% (1.1-4.1%)	18 (9.1-33)	140 (72-260)
Nicaragua	2005	Female	90 to 94	12% (7-20%)	88% (80-93%)	0.51 (0.23-0.98)	1.4% (0.52-3.1%)	1.7% (0.56-3.8%)	10 (3.4-23)	57 (21-120)
Nicaragua	2005	Female	95 plus	15% (7.6-26%)	85% (74-92%)	0.56 (0.21-1.1)	1.3% (0.2-3.4%)	1.5% (0.064-4.1%)	4.1 (0.19-12)	15 (2.4-41)
Nicaragua	2005	Male	All Ages	54% (46-62%)	46% (38-54%)	2 (1.7-2.3)	7.8% (6.9-8.9%)	10% (8.6-11%)	1100 (940-1300)	52000 (45000-59000)
Nicaragua	2005	Male	15 to 19	52% (37-67%)	48% (33-63%)	1.4 (0.75-2.4)	5.9% (3.7-8.7%)	8.4% (4.6-13%)	25 (13-39)	2400 (1500-3600)
Nicaragua	2005	Male	20 to 24	77% (64-85%)	23% (15-36%)	3.2 (1.9-4.8)	14% (9.4-18%)	17% (11-25%)	76 (47-110)	6400 (4400-8700)
Nicaragua	2005	Male	25 to 29	81% (70-89%)	19% (11-30%)	3.8 (2.3-5.7)	16% (12-20%)	20% (15-28%)	80 (56-110)	6400 (4700-8200)
Nicaragua	2005	Male	30 to 34	79% (68-87%)	21% (13-32%)	3.9 (2.5-5.8)	17% (13-20%)	22% (17-27%)	85 (64-110)	6100 (4700-7400)
Nicaragua	2005	Male	35 to 39	72% (63-82%)	28% (18-37%)	3.8 (2.1-6)	17% (14-21%)	23% (18-28%)	90 (70-110)	5500 (4400-6700)
Nicaragua	2005	Male	40 to 44	71% (60-81%)	29% (19-40%)	4 (2.2-6.2)	17% (14-21%)	23% (19-28%)	100 (82-130)	5500 (4400-6600)
Nicaragua	2005	Male	45 to 49	74% (62-83%)	26% (17-38%)	4.4 (2.3-7.4)	17% (14-20%)	22% (18-26%)	120 (93-140)	5400 (4400-6400)
Nicaragua	2005	Male	50 to 54	73% (61-83%)	27% (17-39%)	4.4 (2.4-7.4)	15% (12-18%)	20% (16-23%)	100 (82-120)	4200 (3400-5000)
Nicaragua	2005	Male	55 to 59	70% (57-82%)	30% (18-43%)	4.4 (2.3-7.5)	13% (11-16%)	17% (13-20%)	83 (65-100)	2900 (2300-3500)
Nicaragua	2005	Male	60 to 64	65% (49-79%)	35% (21-51%)	3.5 (1.8-6.5)	10% (7.9-13%)	13% (10-16%)	75 (59-94)	2200 (1800-2800)
Nicaragua	2005	Male	65 to 69	60% (42-77%)	40% (23-58%)	2.7 (1.4-4.6)	8.4% (6.3-11%)	11% (8.2-14%)	71 (54-92)	1800 (1300-2300)
Nicaragua	2005	Male	70 to 74	55% (37-71%)	45% (29-63%)	2 (0.95-3.6)	6.8% (4.5-9.2%)	8.6% (5.7-12%)	64 (42-87)	1300 (850-1700)
Nicaragua	2005	Male	75 to 79	48% (31-66%)	52% (34-69%)	1.5 (0.66-2.9)	5% (2.9-7.2%)	6.2% (3.6-9%)	46 (27-68)	720 (420-1000)
Nicaragua	2005	Male	80 to 84	43% (27-61%)	57% (39-73%)	1.3 (0.61-2.3)	4.2% (2.5-6.2%)	5% (2.9-7.5%)	37 (21-55)	420 (240-620)
Nicaragua	2005	Male	85 to 89	41% (25-59%)	59% (41-75%)	1.2 (0.55-2.3)	3.7% (1.9-5.9%)	4.3% (2.2-6.9%)	29 (15-46)	230 (120-370)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	2005	Male	90 to 94	43% (27-61%)	57% (39-73%)	1.4 (0.54-2.7)	3.5% (1.5-5.9%)	3.9% (1.5-6.8%)	15 (5.9-26)	84 (36-140)
Nicaragua	2005	Male	95 plus	48% (28-69%)	52% (31-72%)	1.6 (0.5-3.5)	3.7% (1.1-6.8%)	4% (0.88-7.7%)	4.9 (1.1-9.6)	19 (5.4-34)
Nicaragua	2010	Female	All Ages	14% (12-17%)	86% (83-88%)	0.37 (0.29-0.47)	1.8% (1.4-2.2%)	3.1% (2.3-4.2%)	290 (220-400)	9500 (7600-12000)
Nicaragua	2010	Female	15 to 19	15% (10-20%)	85% (80-90%)	0.34 (0.18-0.56)	1.1% (0.66-1.7%)	1.6% (0.67-3.1%)	2.2 (0.93-4.1)	350 (210-540)
Nicaragua	2010	Female	20 to 24	25% (18-33%)	75% (67-82%)	0.5 (0.29-0.82)	1.8% (1.2-2.6%)	2.1% (1.1-3.7%)	3 (1.5-5.3)	590 (370-870)
Nicaragua	2010	Female	25 to 29	24% (17-31%)	76% (69-83%)	0.48 (0.29-0.74)	1.8% (1.3-2.4%)	2.2% (1.4-3.2%)	3.3 (2-4.8)	570 (390-810)
Nicaragua	2010	Female	30 to 34	21% (15-29%)	79% (71-85%)	0.47 (0.27-0.75)	1.8% (1.3-2.4%)	2.8% (2.1-3.8%)	4.4 (3.2-6)	540 (390-750)
Nicaragua	2010	Female	35 to 39	20% (13-28%)	80% (72-87%)	0.5 (0.27-0.84)	2% (1.5-2.6%)	3.7% (2.7-4.9%)	7 (5-9.6)	580 (430-780)
Nicaragua	2010	Female	40 to 44	20% (13-28%)	80% (72-87%)	0.57 (0.31-0.95)	2.3% (1.7-3.1%)	4.6% (3.3-6%)	11 (7.7-14)	660 (480-870)
Nicaragua	2010	Female	45 to 49	22% (14-30%)	78% (70-86%)	0.73 (0.38-1.2)	2.8% (1.9-4.1%)	5.3% (3.7-7.4%)	16 (11-23)	810 (560-1100)
Nicaragua	2010	Female	50 to 54	21% (14-29%)	79% (71-86%)	0.85 (0.45-1.5)	3.5% (2.4-5%)	6.1% (4.4-8.4%)	25 (18-35)	1000 (730-1500)
Nicaragua	2010	Female	55 to 59	16% (11-23%)	84% (77-89%)	0.85 (0.43-1.5)	3.8% (2.6-5.3%)	6.2% (4.3-8.4%)	30 (21-42)	1100 (740-1500)
Nicaragua	2010	Female	60 to 64	14% (8.4-20%)	86% (80-92%)	0.83 (0.42-1.5)	3.9% (2.7-5.6%)	5.8% (4-8.1%)	30 (21-43)	890 (620-1300)
Nicaragua	2010	Female	65 to 69	13% (8.2-18%)	87% (82-92%)	0.75 (0.38-1.3)	3.5% (2.4-5.2%)	5.3% (3.7-7.5%)	32 (22-44)	770 (530-1100)
Nicaragua	2010	Female	70 to 74	12% (7.8-18%)	88% (82-92%)	0.65 (0.32-1.2)	3.2% (2.1-4.8%)	4.7% (3.2-7%)	33 (22-48)	640 (420-970)
Nicaragua	2010	Female	75 to 79	11% (6.5-19%)	89% (81-93%)	0.55 (0.25-1)	2.5% (1.5-3.9%)	3.5% (2.2-5.4%)	28 (17-43)	420 (260-650)
Nicaragua	2010	Female	80 to 84	10% (5.5-18%)	90% (82-94%)	0.48 (0.23-0.85)	2.1% (1.2-3.5%)	2.8% (1.7-4.6%)	27 (17-44)	300 (180-480)
Nicaragua	2010	Female	85 to 89	10% (5.7-17%)	90% (83-94%)	0.46 (0.21-0.87)	1.6% (0.89-3%)	2.1% (1.1-3.8%)	24 (12-44)	180 (99-340)
Nicaragua	2010	Female	90 to 94	12% (6.8-20%)	88% (80-93%)	0.49 (0.22-0.98)	1.3% (0.48-2.8%)	1.6% (0.49-3.4%)	14 (4.3-30)	74 (26-160)
Nicaragua	2010	Female	95 plus	15% (7.6-27%)	85% (73-92%)	0.54 (0.2-1.1)	1.2% (0.15-3.2%)	1.3% (-0.024-3.7%)	4.9 (-0.088-14)	19 (2.3-50)
Nicaragua	2010	Male	All Ages	56% (48-64%)	44% (36-52%)	2.1 (1.7-2.4)	8.5% (7.5-9.7%)	10% (8.9-12%)	1200 (1000-1400)	55000 (47000-62000)
Nicaragua	2010	Male	15 to 19	52% (36-67%)	48% (33-64%)	1.4 (0.7-2.3)	5.5% (3.5-8.2%)	7.8% (4.2-12%)	21 (11-33)	2100 (1300-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	2010	Male	20 to 24	77% (64-85%)	23% (15-36%)	3 (1.8-4.7)	13% (8.8-17%)	16% (10-23%)	62 (38-88)	5600 (3900-7700)
Nicaragua	2010	Male	25 to 29	81% (70-89%)	19% (11-30%)	3.7 (2.2-5.5)	15% (11-19%)	19% (14-27%)	85 (59-120)	6900 (5200-8900)
Nicaragua	2010	Male	30 to 34	79% (68-87%)	21% (13-32%)	3.7 (2.4-5.6)	16% (13-20%)	21% (16-27%)	89 (68-110)	6500 (5000-7900)
Nicaragua	2010	Male	35 to 39	72% (62-82%)	28% (18-38%)	3.6 (2-5.8)	17% (13-20%)	23% (18-28%)	100 (78-130)	6300 (4900-7700)
Nicaragua	2010	Male	40 to 44	71% (60-81%)	29% (19-40%)	3.8 (2.2-6)	17% (14-20%)	23% (18-27%)	110 (85-130)	5700 (4600-6900)
Nicaragua	2010	Male	45 to 49	74% (62-83%)	26% (17-38%)	4.2 (2.1-7)	16% (13-19%)	22% (18-26%)	110 (90-130)	5300 (4300-6300)
Nicaragua	2010	Male	50 to 54	73% (61-83%)	27% (17-39%)	4.2 (2.2-7.1)	14% (11-17%)	19% (16-22%)	120 (95-140)	4900 (4000-5800)
Nicaragua	2010	Male	55 to 59	71% (57-82%)	29% (18-43%)	4.1 (2.1-7.1)	13% (10-15%)	16% (13-20%)	110 (91-140)	4000 (3200-4900)
Nicaragua	2010	Male	60 to 64	66% (48-80%)	34% (20-52%)	3.4 (1.7-6.1)	10% (7.8-13%)	13% (9.7-16%)	84 (64-110)	2500 (1900-3100)
Nicaragua	2010	Male	65 to 69	60% (42-77%)	40% (23-58%)	2.5 (1.3-4.4)	8.2% (6.2-11%)	11% (8-13%)	76 (58-98)	1900 (1400-2400)
Nicaragua	2010	Male	70 to 74	55% (37-71%)	45% (29-63%)	1.9 (0.93-3.5)	6.5% (4.4-8.9%)	8.2% (5.6-11%)	68 (46-95)	1400 (930-1900)
Nicaragua	2010	Male	75 to 79	48% (30-66%)	52% (34-70%)	1.5 (0.63-2.8)	4.8% (2.9-7%)	6% (3.6-8.8%)	53 (31-77)	820 (490-1200)
Nicaragua	2010	Male	80 to 84	44% (27-61%)	56% (39-73%)	1.2 (0.58-2.2)	4% (2.3-5.9%)	4.8% (2.7-7.1%)	45 (25-67)	510 (290-750)
Nicaragua	2010	Male	85 to 89	41% (26-58%)	59% (42-74%)	1.2 (0.52-2.2)	3.5% (1.8-5.5%)	4.1% (2.1-6.5%)	36 (19-58)	290 (150-460)
Nicaragua	2010	Male	90 to 94	43% (27-61%)	57% (39-73%)	1.3 (0.52-2.5)	3.3% (1.4-5.5%)	3.7% (1.4-6.4%)	20 (7.8-35)	110 (48-190)
Nicaragua	2010	Male	95 plus	48% (28-70%)	52% (30-72%)	1.6 (0.49-3.4)	3.4% (1-6.4%)	3.7% (0.85-7.3%)	6.5 (1.5-13)	25 (7.5-47)
Nicaragua	2016	Female	All Ages	16% (13-19%)	84% (81-87%)	0.48 (0.34-0.66)	2.1% (1.6-2.7%)	3.4% (2.4-4.8%)	360 (240-520)	12000 (8400-16000)
Nicaragua	2016	Female	15 to 19	14% (9-21%)	86% (79-91%)	0.35 (0.17-0.64)	1.1% (0.64-1.8%)	1.7% (0.67-3.5%)	2 (0.79-4.2)	330 (190-530)
Nicaragua	2016	Female	20 to 24	25% (16-35%)	75% (65-84%)	0.53 (0.26-0.96)	1.9% (1.2-2.7%)	2.3% (1.1-4.4%)	3 (1.3-5.9)	590 (350-920)
Nicaragua	2016	Female	25 to 29	25% (16-34%)	75% (66-84%)	0.53 (0.28-0.9)	1.9% (1.3-2.6%)	2.5% (1.4-3.8%)	3.4 (1.8-5.6)	610 (390-880)
Nicaragua	2016	Female	30 to 34	24% (16-34%)	76% (66-84%)	0.58 (0.31-0.98)	1.9% (1.4-2.6%)	3.2% (2.2-4.7%)	5.4 (3.4-8.2)	650 (440-920)
Nicaragua	2016	Female	35 to 39	24% (15-33%)	76% (67-85%)	0.66 (0.34-1.1)	2.1% (1.5-2.9%)	4.2% (2.9-5.9%)	9.1 (5.7-14)	740 (500-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	2016	Female	40 to 44	24% (16-34%)	76% (66-84%)	0.77 (0.4-1.3)	2.5% (1.7-3.5%)	5.1% (3.5-7.1%)	14 (8.6-20)	850 (560-1200)
Nicaragua	2016	Female	45 to 49	27% (16-37%)	73% (63-84%)	0.98 (0.48-1.7)	3.1% (2-4.8%)	6% (3.9-8.9%)	20 (12-31)	1000 (660-1600)
Nicaragua	2016	Female	50 to 54	25% (17-35%)	75% (65-83%)	1.1 (0.58-2)	3.8% (2.5-5.7%)	6.7% (4.5-9.8%)	28 (18-43)	1200 (760-1900)
Nicaragua	2016	Female	55 to 59	19% (12-27%)	81% (73-88%)	1.1 (0.53-1.9)	4.1% (2.7-6%)	6.7% (4.4-9.4%)	40 (26-61)	1400 (920-2200)
Nicaragua	2016	Female	60 to 64	15% (9-23%)	85% (77-91%)	0.97 (0.46-1.8)	3.9% (2.5-5.8%)	6% (4-8.6%)	48 (30-72)	1400 (880-2200)
Nicaragua	2016	Female	65 to 69	13% (8.2-19%)	87% (81-92%)	0.82 (0.4-1.5)	3.5% (2.3-5.2%)	5.4% (3.6-7.7%)	34 (21-52)	840 (520-1300)
Nicaragua	2016	Female	70 to 74	12% (7.7-18%)	88% (82-92%)	0.68 (0.32-1.3)	3.1% (2-4.8%)	4.7% (3.1-7%)	35 (21-54)	690 (430-1100)
Nicaragua	2016	Female	75 to 79	11% (6.4-19%)	89% (81-94%)	0.58 (0.26-1.1)	2.4% (1.4-3.9%)	3.5% (2.1-5.6%)	31 (18-49)	470 (270-770)
Nicaragua	2016	Female	80 to 84	10% (5.4-18%)	90% (82-95%)	0.5 (0.23-0.94)	2.1% (1.2-3.4%)	2.8% (1.6-4.6%)	32 (18-53)	360 (200-590)
Nicaragua	2016	Female	85 to 89	10% (5.4-17%)	90% (83-95%)	0.48 (0.21-0.9)	1.7% (0.85-3.1%)	2.1% (1-4%)	29 (14-53)	230 (110-410)
Nicaragua	2016	Female	90 to 94	12% (6.7-19%)	88% (81-93%)	0.51 (0.21-1)	1.4% (0.47-3.1%)	1.7% (0.49-3.8%)	19 (5.5-42)	100 (34-220)
Nicaragua	2016	Female	95 plus	15% (7.4-26%)	85% (74-93%)	0.56 (0.2-1.2)	1.3% (0.16-3.5%)	1.4% (-0.00015-4.1%)	7.4 (-0.00076-22)	28 (3.6-76)
Nicaragua	2016	Male	All Ages	53% (44-60%)	47% (40-56%)	2.1 (1.6-2.7)	9.5% (8.1-11%)	11% (9.1-13%)	1400 (1100-1900)	63000 (50000-79000)
Nicaragua	2016	Male	15 to 19	39% (25-55%)	61% (45-75%)	0.98 (0.44-1.9)	4.6% (2.9-6.9%)	6.1% (3.1-11%)	15 (7.4-27)	1700 (1000-2600)
Nicaragua	2016	Male	20 to 24	67% (51-79%)	33% (21-49%)	2.4 (1.2-4.1)	11% (7.7-16%)	14% (8-21%)	56 (30-88)	5200 (3300-7500)
Nicaragua	2016	Male	25 to 29	75% (60-86%)	25% (14-40%)	3.2 (1.8-5.2)	14% (10-19%)	18% (12-25%)	84 (52-120)	7000 (4800-9500)
Nicaragua	2016	Male	30 to 34	75% (61-86%)	25% (14-39%)	3.5 (2-5.7)	15% (12-19%)	20% (15-26%)	100 (70-140)	7500 (5500-9800)
Nicaragua	2016	Male	35 to 39	71% (57-82%)	29% (18-43%)	3.7 (1.9-6)	17% (13-20%)	23% (17-28%)	120 (85-170)	7700 (5600-10000)
Nicaragua	2016	Male	40 to 44	70% (56-82%)	30% (18-44%)	3.9 (2.1-6.2)	17% (13-21%)	23% (18-28%)	130 (95-190)	7200 (5300-9500)
Nicaragua	2016	Male	45 to 49	72% (57-84%)	28% (16-43%)	4.2 (2-7.5)	16% (13-20%)	22% (17-26%)	130 (95-180)	6300 (4700-8200)
Nicaragua	2016	Male	50 to 54	73% (60-84%)	27% (16-40%)	4.5 (2.3-7.7)	15% (12-18%)	20% (16-24%)	130 (95-180)	5500 (4000-7200)
Nicaragua	2016	Male	55 to 59	71% (57-84%)	29% (16-43%)	4.6 (2.2-8.1)	13% (10-16%)	17% (13-20%)	150 (110-210)	5300 (3900-7100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nicaragua	2016	Male	60 to 64	67% (49-81%)	33% (19-51%)	3.8 (1.8-6.9)	10% (7.8-13%)	13% (9.9-16%)	140 (96-180)	4100 (2900-5400)
Nicaragua	2016	Male	65 to 69	61% (43-77%)	39% (23-57%)	2.8 (1.4-4.8)	8.4% (6.3-11%)	11% (8.1-14%)	84 (58-120)	2100 (1500-2900)
Nicaragua	2016	Male	70 to 74	55% (37-71%)	45% (29-63%)	2 (0.93-3.7)	6.6% (4.4-8.9%)	8.4% (5.7-11%)	78 (50-110)	1600 (1000-2200)
Nicaragua	2016	Male	75 to 79	48% (31-66%)	52% (34-69%)	1.5 (0.64-3)	4.9% (3-7%)	6.1% (3.7-8.8%)	65 (37-96)	1000 (580-1500)
Nicaragua	2016	Male	80 to 84	43% (26-61%)	57% (39-74%)	1.3 (0.57-2.4)	4% (2.3-5.9%)	4.8% (2.8-7.2%)	59 (32-89)	670 (380-1000)
Nicaragua	2016	Male	85 to 89	40% (24-58%)	60% (42-76%)	1.2 (0.51-2.3)	3.6% (1.9-5.7%)	4.1% (2.1-6.6%)	49 (25-80)	390 (210-630)
Nicaragua	2016	Male	90 to 94	42% (26-60%)	58% (40-74%)	1.3 (0.52-2.6)	3.3% (1.5-5.6%)	3.7% (1.5-6.4%)	28 (11-49)	160 (67-260)
Nicaragua	2016	Male	95 plus	47% (27-68%)	53% (32-73%)	1.5 (0.49-3.3)	3.4% (1.1-6.2%)	3.7% (0.96-6.9%)	9.6 (2.5-18)	37 (12-67)
Panama	1990	Female	All Ages	23% (17-31%)	77% (69-83%)	0.62 (0.52-0.74)	2% (1.5-2.7%)	3.8% (2.2-5.8%)	170 (98-260)	5200 (3800-7000)
Panama	1990	Female	15 to 19	20% (10-35%)	80% (65-90%)	0.51 (0.22-0.97)	1.4% (0.82-2.4%)	2.4% (0.99-4.9%)	1.8 (0.69-3.4)	210 (120-350)
Panama	1990	Female	20 to 24	28% (14-45%)	72% (55-86%)	0.68 (0.3-1.2)	2.3% (1.4-3.4%)	3.2% (1.4-5.8%)	2.5 (1.1-4.7)	330 (190-500)
Panama	1990	Female	25 to 29	30% (15-50%)	70% (50-85%)	0.72 (0.33-1.3)	2.3% (1.4-3.4%)	3.4% (1.6-5.8%)	3 (1.4-5.1)	330 (190-490)
Panama	1990	Female	30 to 34	33% (16-53%)	67% (47-84%)	0.81 (0.4-1.4)	2.2% (1.4-3.3%)	3.7% (2-5.9%)	3.3 (1.8-5.4)	300 (190-430)
Panama	1990	Female	35 to 39	35% (18-55%)	65% (45-82%)	0.96 (0.42-1.6)	2.3% (1.4-3.6%)	4.1% (2.3-6.8%)	3.8 (2.1-6.3)	280 (170-430)
Panama	1990	Female	40 to 44	35% (19-56%)	65% (44-81%)	1.2 (0.6-2)	2.9% (1.7-4.5%)	5.4% (3.2-8.4%)	6 (3.4-9.4)	340 (200-510)
Panama	1990	Female	45 to 49	35% (17-55%)	65% (45-83%)	1.4 (0.61-2.5)	3.7% (2-6%)	6.8% (3.6-11%)	8.7 (4.8-14)	410 (220-670)
Panama	1990	Female	50 to 54	35% (18-55%)	65% (45-82%)	1.6 (0.74-2.8)	4.4% (2.5-7%)	7.8% (4.4-12%)	11 (6.3-17)	460 (260-730)
Panama	1990	Female	55 to 59	33% (17-54%)	67% (46-83%)	1.8 (0.81-3.3)	5.3% (2.8-8.5%)	8.5% (4.8-13%)	16 (8.8-25)	550 (300-890)
Panama	1990	Female	60 to 64	30% (15-51%)	70% (49-85%)	1.7 (0.72-3.4)	5% (2.5-8.2%)	7.5% (3.8-12%)	17 (8.8-29)	510 (250-850)
Panama	1990	Female	65 to 69	28% (13-48%)	72% (52-87%)	1.5 (0.66-2.8)	4.9% (2.5-8.2%)	6.9% (3.5-11%)	21 (11-35)	510 (250-860)
Panama	1990	Female	70 to 74	25% (12-43%)	75% (57-88%)	1.3 (0.53-2.7)	4.3% (1.9-8.2%)	5.8% (2.5-11%)	24 (10-44)	450 (190-850)
Panama	1990	Female	75 to 79	22% (9.7-39%)	78% (61-90%)	1.1 (0.38-2.2)	3.3% (1.2-6.6%)	4.3% (1.5-8.4%)	20 (6.9-40)	290 (100-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	1990	Female	80 to 84	20% (8.7-36%)	80% (64-91%)	0.95 (0.33-1.9)	2.5% (0.78-5.4%)	3.2% (0.98-6.9%)	14 (4.3-30)	150 (48-320)
Panama	1990	Female	85 to 89	20% (9.5-36%)	80% (64-91%)	0.94 (0.38-1.9)	2.1% (0.35-4.9%)	2.5% (0.39-6.1%)	11 (1.7-26)	84 (14-200)
Panama	1990	Female	90 to 94	24% (11-42%)	76% (58-89%)	1 (0.4-2.1)	1.7% (-0.098-4.7%)	2% (-0.27-5.8%)	5.4 (-0.76-16)	29 (-1.8-83)
Panama	1990	Female	95 plus	30% (12-52%)	70% (48-88%)	1.1 (0.35-2.5)	1.4% (-0.91-5.4%)	1.5% (-1.4-6.4%)	1.5 (-1.4-6.3)	5.6 (-3.7-22)
Panama	1990	Male	All Ages	40% (31-51%)	60% (49-69%)	1.9 (1.7-2.1)	6.5% (5.5-7.6%)	8.4% (6.5-11%)	520 (400-660)	21000 (17000-25000)
Panama	1990	Male	15 to 19	28% (13-46%)	72% (54-87%)	0.87 (0.33-1.6)	4.2% (2.5-6.7%)	5.2% (2.3-9.3%)	7.9 (3.5-14)	820 (470-1300)
Panama	1990	Male	20 to 24	47% (26-68%)	53% (32-74%)	2 (0.89-3.5)	9.9% (6-14%)	11% (5.8-19%)	24 (12-39)	2200 (1300-3200)
Panama	1990	Male	25 to 29	57% (34-77%)	43% (23-66%)	2.8 (1.4-4.6)	12% (7.8-17%)	15% (8.6-23%)	33 (18-50)	2500 (1600-3600)
Panama	1990	Male	30 to 34	64% (38-82%)	36% (18-62%)	3.4 (1.8-5.3)	13% (8.3-17%)	17% (10-23%)	32 (20-47)	2200 (1500-3100)
Panama	1990	Male	35 to 39	67% (45-83%)	33% (17-55%)	3.9 (2-6.2)	13% (8.3-18%)	18% (11-25%)	32 (19-46)	1900 (1200-2700)
Panama	1990	Male	40 to 44	68% (47-84%)	32% (16-53%)	4.3 (2.2-6.7)	12% (8-17%)	18% (11-24%)	32 (20-45)	1700 (1100-2300)
Panama	1990	Male	45 to 49	68% (44-84%)	32% (16-56%)	4.6 (2.2-7.9)	12% (7.9-16%)	18% (11-24%)	35 (21-47)	1600 (1000-2100)
Panama	1990	Male	50 to 54	68% (46-85%)	32% (15-54%)	4.8 (2.4-8.2)	12% (8.3-16%)	17% (12-22%)	38 (25-50)	1500 (1100-2000)
Panama	1990	Male	55 to 59	66% (45-85%)	34% (15-55%)	5.1 (2.5-8.7)	12% (8.1-16%)	16% (11-21%)	44 (29-59)	1500 (1000-2100)
Panama	1990	Male	60 to 64	63% (40-82%)	37% (18-60%)	4.5 (2.1-8.1)	10% (6.7-14%)	13% (8.6-18%)	48 (31-67)	1400 (910-2000)
Panama	1990	Male	65 to 69	59% (35-79%)	41% (21-65%)	3.5 (1.6-6)	9.2% (5.6-13%)	11% (6.9-16%)	52 (31-74)	1300 (750-1800)
Panama	1990	Male	70 to 74	54% (33-74%)	46% (26-67%)	2.5 (1.1-4.9)	6.9% (3.4-11%)	8.3% (4-14%)	49 (24-82)	940 (470-1600)
Panama	1990	Male	75 to 79	47% (25-69%)	53% (31-75%)	1.9 (0.66-4.2)	5.1% (1.8-9.2%)	5.9% (2-11%)	37 (12-67)	550 (190-990)
Panama	1990	Male	80 to 84	42% (20-64%)	58% (36-80%)	1.7 (0.6-3.5)	4.3% (1.6-7.9%)	5.1% (1.8-9.4%)	25 (8.7-47)	280 (100-510)
Panama	1990	Male	85 to 89	40% (20-64%)	60% (36-80%)	1.6 (0.59-3.4)	3.8% (1.2-7.3%)	4.4% (1.3-8.6%)	18 (5.1-36)	140 (44-280)
Panama	1990	Male	90 to 94	44% (23-67%)	56% (33-77%)	1.8 (0.62-3.9)	3.6% (0.8-7.4%)	4.1% (0.72-8.7%)	9.3 (1.6-20)	51 (11-110)
Panama	1990	Male	95 plus	50% (23-74%)	50% (26-77%)	2.1 (0.64-4.8)	3.6% (0.19-8.4%)	4% (-0.19-9.6%)	2.9 (-0.14-7.1)	11 (0.55-25)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	1995	Female	All Ages	23% (17-31%)	77% (69-83%)	0.66 (0.55-0.78)	2.1% (1.5-2.7%)	3.8% (2.3-5.9%)	190 (110-290)	5700 (4100-7600)
Panama	1995	Female	15 to 19	20% (10-34%)	80% (66-90%)	0.52 (0.22-1)	1.4% (0.83-2.3%)	2.5% (1-4.9%)	1.8 (0.74-3.5)	220 (120-360)
Panama	1995	Female	20 to 24	27% (14-46%)	73% (54-86%)	0.69 (0.3-1.3)	2.2% (1.4-3.4%)	3.1% (1.3-5.5%)	2.7 (1.1-4.8)	360 (210-550)
Panama	1995	Female	25 to 29	30% (15-49%)	70% (51-85%)	0.73 (0.34-1.3)	2.2% (1.4-3.2%)	3.2% (1.5-5.4%)	3 (1.4-5.1)	340 (210-500)
Panama	1995	Female	30 to 34	32% (15-53%)	68% (47-85%)	0.81 (0.4-1.4)	2.1% (1.3-3%)	3.2% (1.8-5.1%)	3.3 (1.8-5.3)	310 (200-450)
Panama	1995	Female	35 to 39	34% (18-53%)	66% (47-82%)	0.95 (0.45-1.6)	2.1% (1.3-3.2%)	3.7% (2.1-5.8%)	4.1 (2.3-6.6)	310 (200-460)
Panama	1995	Female	40 to 44	35% (19-55%)	65% (45-81%)	1.2 (0.62-1.9)	2.7% (1.6-4.2%)	5.1% (3-7.8%)	6.4 (3.8-9.9)	370 (230-570)
Panama	1995	Female	45 to 49	35% (17-55%)	65% (45-83%)	1.4 (0.65-2.5)	3.5% (1.9-5.6%)	6.5% (3.7-10%)	9 (4.9-14)	440 (250-690)
Panama	1995	Female	50 to 54	34% (17-55%)	66% (45-83%)	1.7 (0.78-2.8)	4.2% (2.4-6.6%)	7.4% (4.3-11%)	12 (6.9-19)	520 (300-810)
Panama	1995	Female	55 to 59	32% (17-53%)	68% (47-83%)	1.8 (0.83-3.3)	4.8% (2.6-7.8%)	8% (4.6-13%)	15 (8.6-24)	540 (300-870)
Panama	1995	Female	60 to 64	30% (15-51%)	70% (49-85%)	1.7 (0.77-3.3)	4.9% (2.5-8%)	7.5% (4-12%)	18 (9.7-30)	550 (290-920)
Panama	1995	Female	65 to 69	28% (14-48%)	72% (52-86%)	1.6 (0.67-2.9)	4.9% (2.5-8.1%)	7% (3.7-11%)	23 (12-37)	560 (290-920)
Panama	1995	Female	70 to 74	26% (12-43%)	74% (57-88%)	1.4 (0.57-2.8)	4.3% (1.9-8.1%)	6% (2.7-11%)	24 (11-46)	470 (210-890)
Panama	1995	Female	75 to 79	22% (10-39%)	78% (61-90%)	1.2 (0.41-2.3)	3.4% (1.3-6.7%)	4.5% (1.7-8.7%)	23 (8.8-45)	340 (130-670)
Panama	1995	Female	80 to 84	20% (8.6-37%)	80% (63-91%)	1 (0.37-2.1)	2.7% (0.89-5.9%)	3.4% (1.1-7.5%)	19 (6-41)	210 (68-450)
Panama	1995	Female	85 to 89	21% (9.3-36%)	79% (64-91%)	0.98 (0.38-2)	2.2% (0.49-5.1%)	2.7% (0.54-6.3%)	14 (2.8-33)	110 (23-250)
Panama	1995	Female	90 to 94	25% (11-43%)	75% (57-89%)	1.1 (0.41-2.2)	1.8% (0.032-4.9%)	2.2% (-0.1-6.1%)	6.9 (-0.32-19)	38 (0.62-100)
Panama	1995	Female	95 plus	30% (13-52%)	70% (48-87%)	1.2 (0.37-2.6)	1.5% (-0.72-5.7%)	1.7% (-1.1-6.9%)	2.1 (-1.4-8.8)	8.1 (-3.7-30)
Panama	1995	Male	All Ages	42% (32-53%)	58% (47-68%)	2.1 (1.9-2.3)	6.8% (5.8-8.1%)	8.6% (6.8-11%)	620 (490-780)	25000 (21000-29000)
Panama	1995	Male	15 to 19	28% (13-47%)	72% (53-87%)	0.89 (0.34-1.7)	4.3% (2.6-6.9%)	5.3% (2.4-9.5%)	8.4 (3.8-15)	870 (490-1400)
Panama	1995	Male	20 to 24	47% (25-69%)	53% (31-75%)	2.1 (0.92-3.5)	10% (6.1-15%)	12% (5.8-19%)	30 (15-49)	2600 (1500-3900)
Panama	1995	Male	25 to 29	57% (34-77%)	43% (23-66%)	2.9 (1.5-4.8)	12% (7.4-17%)	14% (8-22%)	37 (21-57)	2900 (1800-4100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	1995	Male	30 to 34	64% (40-82%)	36% (18-60%)	3.5 (1.9-5.3)	12% (8-16%)	15% (9.4-21%)	41 (25-59)	2800 (1800-3800)
Panama	1995	Male	35 to 39	67% (45-83%)	33% (17-55%)	3.9 (2.1-6.3)	12% (8-17%)	17% (10-23%)	39 (24-56)	2400 (1600-3300)
Panama	1995	Male	40 to 44	68% (48-84%)	32% (16-52%)	4.3 (2.3-6.8)	12% (7.7-16%)	17% (11-23%)	39 (24-53)	2000 (1300-2700)
Panama	1995	Male	45 to 49	68% (43-84%)	32% (16-57%)	4.7 (2.3-8.2)	12% (7.8-16%)	17% (11-23%)	43 (27-58)	2000 (1300-2700)
Panama	1995	Male	50 to 54	68% (44-85%)	32% (15-56%)	5 (2.5-8.3)	12% (8.3-16%)	17% (11-22%)	46 (31-61)	1900 (1300-2500)
Panama	1995	Male	55 to 59	66% (44-84%)	34% (16-56%)	5.2 (2.5-8.7)	12% (8.2-16%)	16% (11-21%)	52 (35-70)	1800 (1200-2400)
Panama	1995	Male	60 to 64	63% (40-82%)	37% (18-60%)	4.7 (2.2-8.3)	10% (6.8-14%)	13% (8.8-18%)	54 (36-76)	1600 (1000-2200)
Panama	1995	Male	65 to 69	59% (36-79%)	41% (21-64%)	3.6 (1.6-6.2)	9.1% (5.4-13%)	11% (6.8-16%)	57 (35-81)	1400 (850-2000)
Panama	1995	Male	70 to 74	53% (32-74%)	47% (26-68%)	2.6 (1.1-5.2)	6.9% (3.4-11%)	8.3% (4.2-14%)	53 (26-87)	1000 (510-1700)
Panama	1995	Male	75 to 79	47% (24-70%)	53% (30-76%)	2 (0.72-4.2)	5.1% (1.8-9.1%)	6% (2.1-11%)	44 (16-81)	660 (240-1200)
Panama	1995	Male	80 to 84	42% (21-65%)	58% (35-79%)	1.7 (0.61-3.7)	4.5% (1.8-8.2%)	5.3% (2-9.5%)	36 (13-65)	390 (150-720)
Panama	1995	Male	85 to 89	41% (20-63%)	59% (37-80%)	1.7 (0.61-3.5)	3.9% (1.2-7.3%)	4.5% (1.4-8.5%)	23 (7.1-44)	180 (58-340)
Panama	1995	Male	90 to 94	44% (22-67%)	56% (33-78%)	1.9 (0.64-4)	3.6% (0.92-7.2%)	4.1% (0.87-8.4%)	11 (2.3-22)	60 (15-120)
Panama	1995	Male	95 plus	49% (24-74%)	51% (26-76%)	2.2 (0.63-4.9)	3.7% (0.38-7.9%)	4% (0.061-9.3%)	3.4 (0.052-7.7)	13 (1.3-28)
Panama	2000	Female	All Ages	23% (18-31%)	77% (69-82%)	0.7 (0.59-0.84)	2.1% (1.6-2.8%)	4% (2.5-5.9%)	210 (130-320)	6300 (4600-8300)
Panama	2000	Female	15 to 19	20% (10-34%)	80% (66-90%)	0.53 (0.23-1)	1.4% (0.8-2.3%)	2.4% (0.99-4.8%)	1.7 (0.69-3.4)	210 (120-360)
Panama	2000	Female	20 to 24	27% (13-45%)	73% (55-87%)	0.71 (0.3-1.3)	2.1% (1.3-3.2%)	2.8% (1.2-5%)	2.5 (1.1-4.5)	350 (210-540)
Panama	2000	Female	25 to 29	30% (14-50%)	70% (50-86%)	0.75 (0.34-1.3)	2.2% (1.4-3.1%)	3% (1.4-5%)	3 (1.4-5.1)	360 (220-540)
Panama	2000	Female	30 to 34	32% (15-54%)	68% (46-85%)	0.83 (0.4-1.4)	1.8% (1.2-2.6%)	2.6% (1.4-4.1%)	2.9 (1.6-4.7)	310 (200-460)
Panama	2000	Female	35 to 39	33% (17-53%)	67% (47-83%)	0.97 (0.45-1.6)	2% (1.3-2.9%)	3.4% (2-5.2%)	4.4 (2.5-6.9)	340 (220-500)
Panama	2000	Female	40 to 44	34% (19-55%)	66% (45-81%)	1.2 (0.6-1.9)	2.5% (1.6-3.9%)	4.8% (2.9-7.2%)	7 (4.1-11)	420 (250-620)
Panama	2000	Female	45 to 49	34% (17-54%)	66% (46-83%)	1.4 (0.64-2.5)	3.2% (1.9-5.1%)	6.2% (3.6-9.8%)	9.5 (5.6-15)	480 (280-750)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	2000	Female	50 to 54	34% (18-55%)	66% (45-82%)	1.7 (0.79-2.8)	4% (2.4-6.3%)	7% (4.2-11%)	13 (8-21)	570 (340-890)
Panama	2000	Female	55 to 59	32% (16-52%)	68% (48-84%)	1.8 (0.84-3.3)	4.6% (2.7-7.6%)	7.8% (4.5-12%)	17 (9.8-27)	620 (350-1000)
Panama	2000	Female	60 to 64	30% (14-50%)	70% (50-86%)	1.8 (0.77-3.4)	4.9% (2.6-7.9%)	7.5% (4.2-12%)	21 (12-34)	640 (350-1100)
Panama	2000	Female	65 to 69	28% (14-48%)	72% (52-86%)	1.7 (0.71-3)	4.9% (2.7-8.2%)	7.2% (4.1-12%)	26 (14-41)	630 (340-1000)
Panama	2000	Female	70 to 74	25% (12-43%)	75% (57-88%)	1.4 (0.59-2.8)	4.4% (2.1-8%)	6.2% (3-11%)	27 (13-48)	540 (260-970)
Panama	2000	Female	75 to 79	22% (9.5-38%)	78% (62-90%)	1.2 (0.42-2.4)	3.5% (1.4-6.6%)	4.7% (1.9-8.7%)	26 (11-48)	390 (150-730)
Panama	2000	Female	80 to 84	20% (8.2-37%)	80% (63-92%)	1 (0.36-2.2)	2.9% (1.1-6%)	3.7% (1.4-7.8%)	23 (8.7-49)	250 (95-530)
Panama	2000	Female	85 to 89	20% (9.1-36%)	80% (64-91%)	1 (0.4-2)	2.3% (0.7-5%)	2.9% (0.84-6.5%)	16 (4.6-37)	130 (39-290)
Panama	2000	Female	90 to 94	24% (11-42%)	76% (58-89%)	1.1 (0.42-2.3)	1.9% (0.25-4.8%)	2.3% (0.16-6.2%)	7.9 (0.54-21)	44 (5.5-110)
Panama	2000	Female	95 plus	29% (13-51%)	71% (49-87%)	1.2 (0.37-2.6)	1.6% (-0.4-5.3%)	1.8% (-0.84-6.7%)	2.6 (-1.2-9.4)	10 (-2.6-33)
Panama	2000	Male	All Ages	42% (32-53%)	58% (47-68%)	2.2 (2-2.4)	6.6% (5.6-7.7%)	8.4% (6.7-10%)	620 (490-780)	25000 (21000-30000)
Panama	2000	Male	15 to 19	27% (13-47%)	73% (53-87%)	0.93 (0.35-1.8)	4.3% (2.5-7%)	5.4% (2.4-10%)	8.5 (3.7-16)	890 (500-1500)
Panama	2000	Male	20 to 24	46% (25-67%)	54% (33-75%)	2.1 (0.92-3.8)	9.8% (6-15%)	11% (5.6-18%)	26 (13-44)	2400 (1400-3600)
Panama	2000	Male	25 to 29	57% (33-76%)	43% (24-67%)	3 (1.5-5)	11% (7.1-15%)	13% (7.3-20%)	33 (18-50)	2700 (1800-3900)
Panama	2000	Male	30 to 34	63% (39-82%)	37% (18-61%)	3.6 (1.9-5.5)	11% (7.1-14%)	13% (8-18%)	36 (21-51)	2600 (1700-3500)
Panama	2000	Male	35 to 39	66% (45-83%)	34% (17-55%)	4 (2.1-6.5)	10% (6.9-14%)	14% (8.6-19%)	35 (22-50)	2200 (1500-3100)
Panama	2000	Male	40 to 44	67% (46-83%)	33% (17-54%)	4.4 (2.4-7)	11% (7.1-14%)	15% (9.4-20%)	39 (24-54)	2100 (1400-2900)
Panama	2000	Male	45 to 49	67% (43-83%)	33% (17-57%)	4.8 (2.3-8.3)	11% (7.3-14%)	16% (10-20%)	41 (27-55)	2000 (1300-2600)
Panama	2000	Male	50 to 54	67% (45-83%)	33% (17-55%)	5.1 (2.6-8.5)	11% (8-15%)	16% (11-20%)	50 (34-66)	2000 (1400-2700)
Panama	2000	Male	55 to 59	65% (44-84%)	35% (16-56%)	5.4 (2.6-9)	11% (7.8-15%)	15% (10-20%)	54 (37-73)	1900 (1300-2600)
Panama	2000	Male	60 to 64	63% (39-82%)	37% (18-61%)	4.8 (2.2-8.7)	10% (6.9-14%)	13% (9-18%)	61 (40-85)	1800 (1200-2500)
Panama	2000	Male	65 to 69	59% (35-78%)	41% (22-65%)	3.7 (1.7-6.4)	8.8% (5.4-12%)	11% (7.1-16%)	60 (37-85)	1500 (910-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	2000	Male	70 to 74	52% (31-72%)	48% (28-69%)	2.7 (1.2-5.3)	6.8% (3.5-11%)	8.3% (4.3-13%)	54 (28-87)	1100 (560-1700)
Panama	2000	Male	75 to 79	46% (24-69%)	54% (31-76%)	2 (0.73-4.4)	5% (2-8.7%)	6% (2.3-11%)	45 (17-78)	680 (270-1200)
Panama	2000	Male	80 to 84	41% (20-63%)	59% (37-80%)	1.7 (0.62-3.8)	4.4% (1.8-7.8%)	5.2% (2.1-9.3%)	39 (15-70)	430 (170-770)
Panama	2000	Male	85 to 89	39% (19-62%)	61% (38-81%)	1.7 (0.6-3.7)	3.9% (1.4-7%)	4.5% (1.5-8.2%)	25 (8.6-47)	200 (73-370)
Panama	2000	Male	90 to 94	43% (22-66%)	57% (34-78%)	1.9 (0.64-4.1)	3.5% (1.1-6.7%)	4% (1.1-7.9%)	11 (2.9-22)	62 (19-120)
Panama	2000	Male	95 plus	48% (23-73%)	52% (27-77%)	2.2 (0.63-5)	3.6% (0.6-7.4%)	4% (0.37-8.6%)	3.5 (0.34-7.8)	14 (2.3-28)
Panama	2005	Female	All Ages	23% (17-30%)	77% (70-83%)	0.73 (0.61-0.87)	2.3% (1.7-3%)	4.1% (2.6-6.2%)	250 (160-380)	7200 (5300-9500)
Panama	2005	Female	15 to 19	19% (9.6-32%)	81% (68-90%)	0.53 (0.23-1)	1.4% (0.78-2.2%)	2.4% (0.98-4.7%)	1.8 (0.73-3.5)	220 (120-380)
Panama	2005	Female	20 to 24	26% (13-44%)	74% (56-87%)	0.7 (0.29-1.3)	2.1% (1.3-3.1%)	2.8% (1.2-5.1%)	2.6 (1.1-4.7)	360 (220-540)
Panama	2005	Female	25 to 29	29% (13-47%)	71% (53-87%)	0.74 (0.32-1.3)	2.1% (1.4-3.1%)	2.9% (1.3-4.9%)	2.9 (1.3-5)	370 (230-540)
Panama	2005	Female	30 to 34	31% (14-51%)	69% (49-86%)	0.82 (0.38-1.4)	1.9% (1.3-2.7%)	2.7% (1.5-4.4%)	3.2 (1.7-5.3)	350 (220-510)
Panama	2005	Female	35 to 39	32% (16-52%)	68% (48-84%)	0.97 (0.43-1.7)	2.1% (1.3-3%)	3.5% (2-5.5%)	4.9 (2.8-7.8)	390 (250-570)
Panama	2005	Female	40 to 44	33% (17-54%)	67% (46-83%)	1.2 (0.6-1.9)	2.4% (1.5-3.7%)	4.8% (2.9-7.2%)	7.3 (4.3-11)	460 (280-670)
Panama	2005	Female	45 to 49	33% (17-53%)	67% (47-83%)	1.4 (0.64-2.5)	3.1% (1.9-4.9%)	6% (3.4-9.5%)	11 (6.2-17)	560 (330-880)
Panama	2005	Female	50 to 54	32% (16-54%)	68% (46-84%)	1.6 (0.78-2.8)	3.9% (2.3-6.2%)	7.1% (4.2-11%)	15 (9.2-23)	660 (390-1000)
Panama	2005	Female	55 to 59	31% (16-51%)	69% (49-84%)	1.8 (0.85-3.3)	4.7% (2.7-7.6%)	7.8% (4.6-12%)	20 (12-32)	740 (430-1200)
Panama	2005	Female	60 to 64	29% (14-49%)	71% (51-86%)	1.8 (0.78-3.4)	4.8% (2.6-7.9%)	7.5% (4.3-12%)	25 (14-38)	760 (420-1200)
Panama	2005	Female	65 to 69	27% (13-46%)	73% (54-87%)	1.6 (0.71-2.9)	4.8% (2.7-7.7%)	7.2% (4.1-11%)	28 (16-45)	710 (390-1100)
Panama	2005	Female	70 to 74	24% (12-42%)	76% (58-88%)	1.4 (0.57-2.8)	4.4% (2.2-7.7%)	6.3% (3.3-11%)	32 (17-55)	630 (320-1100)
Panama	2005	Female	75 to 79	21% (9.2-38%)	79% (62-91%)	1.2 (0.43-2.4)	3.5% (1.5-6.5%)	4.7% (2-8.7%)	29 (13-54)	440 (180-830)
Panama	2005	Female	80 to 84	19% (8.1-35%)	81% (65-92%)	1 (0.35-2.1)	2.8% (1.1-5.7%)	3.7% (1.5-7.5%)	26 (10-53)	290 (120-590)
Panama	2005	Female	85 to 89	20% (8.9-35%)	80% (65-91%)	1 (0.41-2.1)	2.5% (0.84-5.3%)	3.1% (1-6.7%)	22 (7.2-47)	170 (58-360)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	2005	Female	90 to 94	24% (11-42%)	76% (58-89%)	1.1 (0.43-2.4)	2.1% (0.36-5.3%)	2.6% (0.36-6.6%)	13 (1.7-33)	70 (11-170)
Panama	2005	Female	95 plus	29% (13-51%)	71% (49-87%)	1.2 (0.39-2.7)	1.9% (-0.29-5.9%)	2.2% (-0.59-7.4%)	4.7 (-1.3-16)	17 (-2.5-55)
Panama	2005	Male	All Ages	43% (33-54%)	57% (46-67%)	2.4 (2.2-2.6)	7% (5.9-8.2%)	8.6% (6.9-11%)	720 (560-890)	28000 (23000-33000)
Panama	2005	Male	15 to 19	27% (13-46%)	73% (54-87%)	0.92 (0.35-1.8)	4.4% (2.5-7.2%)	5.4% (2.3-10%)	9.3 (4-18)	960 (530-1600)
Panama	2005	Male	20 to 24	46% (24-68%)	54% (32-76%)	2.1 (0.91-3.9)	10% (6-15%)	11% (5.7-19%)	29 (14-49)	2600 (1500-4100)
Panama	2005	Male	25 to 29	56% (33-76%)	44% (24-67%)	3 (1.5-5.1)	11% (7.2-16%)	14% (7.5-20%)	36 (19-55)	2900 (1900-4200)
Panama	2005	Male	30 to 34	62% (38-81%)	38% (19-62%)	3.7 (2-5.7)	11% (7.5-15%)	14% (8.8-20%)	39 (24-57)	2900 (1900-3900)
Panama	2005	Male	35 to 39	65% (44-82%)	35% (18-56%)	4.1 (2.2-6.6)	11% (7.3-15%)	15% (9.1-20%)	41 (26-58)	2600 (1700-3600)
Panama	2005	Male	40 to 44	67% (47-83%)	33% (17-53%)	4.5 (2.4-7)	11% (7.1-14%)	15% (9.7-20%)	45 (28-62)	2500 (1700-3300)
Panama	2005	Male	45 to 49	67% (42-83%)	33% (17-58%)	4.9 (2.4-8.4)	11% (7.4-15%)	16% (11-21%)	49 (32-67)	2400 (1600-3200)
Panama	2005	Male	50 to 54	66% (43-83%)	34% (17-57%)	5.2 (2.7-8.7)	11% (7.7-14%)	15% (11-20%)	55 (38-73)	2300 (1600-3000)
Panama	2005	Male	55 to 59	65% (43-83%)	35% (17-57%)	5.4 (2.7-9.1)	11% (7.9-15%)	15% (10-20%)	65 (44-86)	2300 (1600-3100)
Panama	2005	Male	60 to 64	62% (39-82%)	38% (18-61%)	4.9 (2.2-8.7)	10% (6.7-14%)	13% (8.6-18%)	69 (45-96)	2100 (1300-2900)
Panama	2005	Male	65 to 69	58% (34-78%)	42% (22-66%)	3.7 (1.7-6.5)	8.7% (5.4-12%)	11% (7.1-15%)	67 (42-96)	1700 (1100-2400)
Panama	2005	Male	70 to 74	52% (31-73%)	48% (27-69%)	2.7 (1.2-5.5)	6.8% (3.6-11%)	8.4% (4.4-13%)	62 (33-99)	1200 (640-1900)
Panama	2005	Male	75 to 79	45% (24-68%)	55% (32-76%)	2.1 (0.73-4.5)	5.1% (2.1-8.7%)	6.1% (2.5-11%)	50 (19-86)	760 (300-1300)
Panama	2005	Male	80 to 84	41% (20-63%)	59% (37-80%)	1.8 (0.63-3.9)	4.4% (1.8-7.8%)	5.3% (2.1-9.5%)	44 (18-78)	500 (210-870)
Panama	2005	Male	85 to 89	39% (19-62%)	61% (38-81%)	1.7 (0.61-3.7)	3.9% (1.4-7.2%)	4.6% (1.6-8.4%)	32 (11-61)	260 (96-480)
Panama	2005	Male	90 to 94	42% (22-65%)	58% (35-78%)	1.9 (0.65-4.1)	3.7% (1.2-7.1%)	4.3% (1.2-8.5%)	17 (4.9-34)	96 (31-190)
Panama	2005	Male	95 plus	47% (22-72%)	53% (28-78%)	2.2 (0.63-5)	3.8% (0.64-7.8%)	4.2% (0.46-9.1%)	5.6 (0.62-12)	21 (3.5-44)
Panama	2010	Female	All Ages	23% (18-30%)	77% (70-82%)	0.82 (0.68-0.97)	2.5% (1.8-3.3%)	4.5% (2.9-6.7%)	310 (190-460)	8700 (6400-11000)
Panama	2010	Female	15 to 19	19% (9.9-32%)	81% (68-90%)	0.56 (0.24-1.1)	1.4% (0.8-2.3%)	2.5% (1-4.8%)	1.9 (0.8-3.8)	240 (130-400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	2010	Female	20 to 24	26% (13-43%)	74% (57-87%)	0.75 (0.32-1.4)	2.2% (1.3-3.2%)	2.9% (1.3-5.3%)	2.8 (1.2-5.1)	390 (220-590)
Panama	2010	Female	25 to 29	29% (14-48%)	71% (52-86%)	0.8 (0.35-1.4)	2.2% (1.4-3.1%)	3% (1.3-5.1%)	3.2 (1.4-5.6)	400 (240-590)
Panama	2010	Female	30 to 34	31% (14-51%)	69% (49-86%)	0.87 (0.4-1.6)	1.9% (1.3-2.8%)	2.9% (1.6-4.6%)	3.6 (1.9-5.8)	380 (250-560)
Panama	2010	Female	35 to 39	32% (17-51%)	68% (49-83%)	1 (0.46-1.7)	2.1% (1.4-3.2%)	3.8% (2.3-5.9%)	5.6 (3.2-8.7)	440 (290-660)
Panama	2010	Female	40 to 44	33% (18-54%)	67% (46-82%)	1.2 (0.64-2)	2.5% (1.6-3.7%)	4.8% (3-7.1%)	8.2 (5-12)	520 (320-770)
Panama	2010	Female	45 to 49	33% (16-52%)	67% (48-84%)	1.5 (0.69-2.6)	3.3% (1.9-5.2%)	6.2% (3.6-9.7%)	14 (7.9-21)	700 (410-1100)
Panama	2010	Female	50 to 54	33% (17-54%)	67% (46-83%)	1.8 (0.85-3)	4.2% (2.5-6.5%)	7.6% (4.6-11%)	20 (13-30)	870 (540-1300)
Panama	2010	Female	55 to 59	31% (16-53%)	69% (47-84%)	2 (0.92-3.5)	4.9% (2.9-7.9%)	8.3% (5-13%)	25 (15-38)	930 (550-1500)
Panama	2010	Female	60 to 64	29% (14-50%)	71% (50-86%)	1.9 (0.82-3.7)	5% (2.8-8.2%)	7.9% (4.6-12%)	30 (17-47)	920 (520-1500)
Panama	2010	Female	65 to 69	28% (13-47%)	72% (53-87%)	1.8 (0.75-3.1)	5% (2.8-8%)	7.6% (4.4-12%)	35 (20-53)	870 (490-1400)
Panama	2010	Female	70 to 74	25% (12-42%)	75% (58-88%)	1.5 (0.64-3)	4.6% (2.4-8.1%)	6.7% (3.6-11%)	39 (21-66)	770 (410-1300)
Panama	2010	Female	75 to 79	21% (9.2-38%)	79% (62-91%)	1.3 (0.46-2.6)	3.7% (1.6-6.7%)	5.1% (2.3-9.2%)	35 (15-64)	540 (230-990)
Panama	2010	Female	80 to 84	19% (8.1-36%)	81% (64-92%)	1.1 (0.37-2.3)	3% (1.2-5.8%)	4% (1.7-7.9%)	31 (13-64)	350 (150-710)
Panama	2010	Female	85 to 89	20% (8.5-35%)	80% (65-92%)	1.1 (0.41-2.2)	2.6% (0.96-5.5%)	3.4% (1.2-7.1%)	27 (9.6-59)	210 (78-460)
Panama	2010	Female	90 to 94	23% (11-41%)	77% (59-89%)	1.2 (0.45-2.5)	2.4% (0.56-5.6%)	2.9% (0.56-7.2%)	19 (3.4-46)	100 (23-240)
Panama	2010	Female	95 plus	29% (12-51%)	71% (49-88%)	1.3 (0.4-2.9)	2.1% (-0.14-6.3%)	2.5% (-0.44-8%)	7.1 (-1.3-23)	26 (-1.7-78)
Panama	2010	Male	All Ages	43% (34-54%)	57% (46-66%)	2.6 (2.3-2.8)	7.6% (6.3-8.8%)	9.1% (7.4-11%)	890 (700-1100)	35000 (28000-41000)
Panama	2010	Male	15 to 19	26% (12-45%)	74% (55-88%)	0.96 (0.35-1.8)	4.7% (2.5-7.9%)	5.7% (2.5-11%)	13 (5.6-25)	1200 (650-2100)
Panama	2010	Male	20 to 24	45% (24-67%)	55% (33-76%)	2.2 (0.95-4)	11% (6-16%)	12% (5.9-20%)	40 (19-67)	3400 (1900-5200)
Panama	2010	Male	25 to 29	56% (33-76%)	44% (24-67%)	3.2 (1.5-5.4)	12% (7.3-17%)	14% (7.7-22%)	46 (25-71)	3600 (2200-5300)
Panama	2010	Male	30 to 34	62% (39-81%)	38% (19-61%)	3.9 (2-6.2)	12% (7.9-16%)	15% (9.3-22%)	48 (29-70)	3400 (2200-4700)
Panama	2010	Male	35 to 39	65% (44-82%)	35% (18-56%)	4.3 (2.3-7)	11% (7.6-15%)	15% (9.5-21%)	48 (30-69)	3100 (2000-4200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	2010	Male	40 to 44	66% (46-83%)	34% (17-54%)	4.7 (2.6-7.3)	11% (7.6-15%)	16% (10-21%)	54 (35-74)	3000 (2000-3900)
Panama	2010	Male	45 to 49	67% (43-83%)	33% (17-57%)	5.1 (2.5-8.7)	11% (7.7-15%)	16% (11-21%)	62 (41-82)	3000 (2000-3900)
Panama	2010	Male	50 to 54	67% (45-83%)	33% (17-55%)	5.5 (2.9-9)	11% (8.1-15%)	16% (12-21%)	70 (49-92)	2900 (2100-3800)
Panama	2010	Male	55 to 59	65% (43-83%)	35% (17-57%)	5.8 (2.9-9.8)	11% (8-15%)	15% (11-20%)	79 (54-100)	2800 (1900-3700)
Panama	2010	Male	60 to 64	62% (39-82%)	38% (18-61%)	5.2 (2.4-9.2)	10% (7-14%)	13% (9.1-18%)	86 (57-120)	2600 (1700-3600)
Panama	2010	Male	65 to 69	59% (35-78%)	41% (22-65%)	4 (1.8-6.9)	9.1% (5.8-12%)	12% (7.6-16%)	84 (54-120)	2100 (1300-2900)
Panama	2010	Male	70 to 74	52% (31-74%)	48% (26-69%)	2.9 (1.3-5.7)	7% (3.9-11%)	8.7% (4.9-13%)	74 (41-110)	1500 (810-2300)
Panama	2010	Male	75 to 79	46% (24-69%)	54% (31-76%)	2.2 (0.78-4.8)	5.3% (2.3-9.1%)	6.5% (2.8-11%)	61 (26-110)	940 (400-1600)
Panama	2010	Male	80 to 84	41% (19-63%)	59% (37-81%)	1.9 (0.69-4.1)	4.5% (1.9-8%)	5.5% (2.3-9.7%)	53 (22-94)	600 (250-1100)
Panama	2010	Male	85 to 89	39% (19-61%)	61% (39-81%)	1.8 (0.66-3.9)	4.1% (1.7-7.4%)	4.9% (1.9-8.9%)	41 (16-74)	330 (130-590)
Panama	2010	Male	90 to 94	42% (21-65%)	58% (35-79%)	2 (0.68-4.3)	4% (1.3-7.5%)	4.6% (1.4-8.9%)	25 (7.5-48)	140 (46-260)
Panama	2010	Male	95 plus	47% (21-72%)	53% (28-79%)	2.3 (0.66-5.3)	4% (0.84-8.2%)	4.5% (0.55-9.5%)	7.9 (1-17)	30 (5.9-61)
Panama	2016	Female	All Ages	23% (18-31%)	77% (69-82%)	0.92 (0.7-1.2)	2.7% (1.9-3.7%)	4.7% (2.8-7.1%)	370 (220-550)	10000 (6900-14000)
Panama	2016	Female	15 to 19	19% (9.3-32%)	81% (68-91%)	0.6 (0.23-1.2)	1.4% (0.81-2.4%)	2.5% (0.98-5.1%)	1.9 (0.76-3.9)	250 (140-430)
Panama	2016	Female	20 to 24	26% (12-44%)	74% (56-88%)	0.8 (0.31-1.5)	2.2% (1.4-3.4%)	3.1% (1.3-5.6%)	2.7 (1.1-5.3)	400 (240-640)
Panama	2016	Female	25 to 29	28% (14-47%)	72% (53-86%)	0.85 (0.35-1.6)	2.1% (1.4-3.1%)	3% (1.3-5.1%)	3 (1.3-5.5)	410 (250-620)
Panama	2016	Female	30 to 34	30% (14-50%)	70% (50-86%)	0.95 (0.41-1.7)	2.1% (1.3-3.1%)	3.4% (1.8-5.5%)	4 (2-6.8)	420 (260-640)
Panama	2016	Female	35 to 39	32% (16-52%)	68% (48-84%)	1.1 (0.48-1.9)	2.2% (1.4-3.3%)	4% (2.3-6.2%)	5.5 (3.1-9.3)	470 (290-730)
Panama	2016	Female	40 to 44	33% (18-53%)	67% (47-82%)	1.3 (0.67-2.3)	2.6% (1.5-3.9%)	5.2% (3.2-7.8%)	8.9 (5.2-14)	590 (360-900)
Panama	2016	Female	45 to 49	33% (16-52%)	67% (48-84%)	1.6 (0.75-2.9)	3.3% (1.9-5.2%)	6.5% (3.7-10%)	15 (8.2-24)	800 (450-1300)
Panama	2016	Female	50 to 54	32% (16-53%)	68% (47-84%)	1.9 (0.86-3.3)	4.1% (2.3-6.4%)	7.5% (4.4-11%)	22 (12-35)	1000 (570-1600)
Panama	2016	Female	55 to 59	30% (16-50%)	70% (50-84%)	2 (0.95-3.7)	4.8% (2.8-7.9%)	8.3% (4.9-13%)	29 (17-48)	1100 (630-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	2016	Female	60 to 64	28% (13-49%)	72% (51-87%)	2 (0.87-3.8)	5% (2.8-8.2%)	8% (4.6-13%)	35 (20-57)	1100 (610-1800)
Panama	2016	Female	65 to 69	27% (13-46%)	73% (54-87%)	1.9 (0.81-3.3)	5.1% (2.9-8.3%)	7.9% (4.7-12%)	42 (24-67)	1100 (600-1800)
Panama	2016	Female	70 to 74	24% (12-41%)	76% (59-88%)	1.6 (0.64-3.2)	4.6% (2.4-7.8%)	6.9% (3.8-11%)	45 (23-75)	910 (460-1500)
Panama	2016	Female	75 to 79	20% (9.3-36%)	80% (64-91%)	1.3 (0.49-2.7)	3.7% (1.6-6.7%)	5.3% (2.3-9.4%)	42 (19-77)	660 (280-1200)
Panama	2016	Female	80 to 84	18% (7.9-34%)	82% (66-92%)	1.1 (0.4-2.3)	3% (1.2-6%)	4.1% (1.7-8.1%)	39 (16-77)	450 (180-860)
Panama	2016	Female	85 to 89	19% (8.2-33%)	81% (67-92%)	1.1 (0.42-2.3)	2.7% (1.3-5.5%)	3.5% (1.3-7.1%)	35 (12-72)	280 (100-580)
Panama	2016	Female	90 to 94	23% (10-40%)	77% (60-90%)	1.2 (0.47-2.6)	2.6% (0.62-5.9%)	3.2% (0.66-7.5%)	26 (5.4-62)	140 (33-330)
Panama	2016	Female	95 plus	28% (12-50%)	72% (50-88%)	1.3 (0.42-3)	2.3% (-0.077-6.5%)	2.7% (-0.4-8.1%)	10 (-1.6-31)	38 (-1.1-110)
Panama	2016	Male	All Ages	44% (35-54%)	56% (46-65%)	2.9 (2.3-3.6)	8.1% (6.6-9.8%)	9.5% (7.4-12%)	1000 (760-1400)	39000 (31000-50000)
Panama	2016	Male	15 to 19	26% (13-45%)	74% (55-87%)	1 (0.39-2.1)	4.9% (2.7-8.4%)	6.1% (2.6-12%)	14 (5.8-28)	1300 (700-2300)
Panama	2016	Male	20 to 24	45% (24-67%)	55% (33-76%)	2.4 (1-4.5)	11% (6.3-17%)	13% (6.1-21%)	39 (19-67)	3400 (2000-5400)
Panama	2016	Male	25 to 29	56% (33-75%)	44% (25-67%)	3.5 (1.6-6.2)	13% (8-18%)	15% (8.6-24%)	49 (26-77)	3900 (2400-5800)
Panama	2016	Male	30 to 34	62% (38-81%)	38% (19-62%)	4.2 (2.1-7)	13% (8.5-17%)	16% (10-24%)	51 (30-78)	3700 (2400-5300)
Panama	2016	Male	35 to 39	65% (43-82%)	35% (18-57%)	4.7 (2.3-7.7)	12% (8.2-17%)	17% (11-23%)	52 (31-79)	3400 (2200-4800)
Panama	2016	Male	40 to 44	66% (46-83%)	34% (17-54%)	5.2 (2.6-8.3)	12% (8.1-16%)	17% (11-22%)	60 (37-87)	3400 (2300-4700)
Panama	2016	Male	45 to 49	67% (42-83%)	33% (17-58%)	5.6 (2.7-9.6)	12% (8.3-15%)	17% (12-22%)	70 (44-97)	3500 (2300-4700)
Panama	2016	Male	50 to 54	66% (44-83%)	34% (17-56%)	6 (3.1-9.9)	12% (8.4-15%)	17% (12-21%)	81 (55-110)	3500 (2400-4600)
Panama	2016	Male	55 to 59	65% (43-83%)	35% (17-57%)	6.2 (3-10)	11% (8-15%)	16% (11-20%)	93 (62-130)	3400 (2300-4700)
Panama	2016	Male	60 to 64	62% (38-81%)	38% (19-62%)	5.6 (2.6-9.9)	10% (7-14%)	13% (9.3-18%)	100 (64-140)	3000 (2000-4300)
Panama	2016	Male	65 to 69	58% (35-79%)	42% (21-65%)	4.3 (2-7.4)	9.2% (5.9-12%)	12% (7.8-16%)	98 (63-140)	2500 (1600-3500)
Panama	2016	Male	70 to 74	52% (31-73%)	48% (27-69%)	3.1 (1.4-6.1)	7.1% (4-11%)	8.9% (5.1-13%)	86 (48-130)	1700 (980-2700)
Panama	2016	Male	75 to 79	45% (23-69%)	55% (31-77%)	2.3 (0.8-4.9)	5.4% (2.4-8.9%)	6.6% (3-11%)	73 (31-120)	1100 (480-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Panama	2016	Male	80 to 84	40% (19-63%)	60% (37-81%)	2 (0.7-4.4)	4.6% (2.1-7.9%)	5.7% (2.5-9.6%)	65 (28-110)	750 (320-1300)
Panama	2016	Male	85 to 89	39% (19-62%)	61% (38-81%)	1.9 (0.66-4.3)	4.3% (1.8-7.4%)	5.1% (2.1-9%)	51 (21-91)	420 (180-740)
Panama	2016	Male	90 to 94	41% (20-65%)	59% (35-80%)	2.1 (0.73-4.6)	4.1% (1.5-7.4%)	4.8% (1.6-8.9%)	33 (11-62)	180 (67-340)
Panama	2016	Male	95 plus	46% (21-72%)	54% (28-79%)	2.4 (0.68-5.9)	4.2% (1-8.2%)	4.7% (0.82-9.7%)	11 (2-24)	43 (10-84)
Venezuela	1990	Female	All Ages	24% (18-31%)	76% (69-82%)	1 (0.82-1.2)	3.2% (2.4-4.1%)	6.4% (4-8.9%)	2300 (1500-3300)	69000 (51000-91000)
Venezuela	1990	Female	15 to 19	21% (11-36%)	79% (64-89%)	0.85 (0.39-1.6)	2.1% (1.1-3.5%)	3.7% (1.6-6.9%)	21 (9.1-39)	2300 (1200-4000)
Venezuela	1990	Female	20 to 24	29% (14-48%)	71% (52-86%)	1.1 (0.51-2)	3% (1.8-4.6%)	4.6% (2.2-8.1%)	30 (14-52)	3500 (2000-5500)
Venezuela	1990	Female	25 to 29	32% (16-52%)	68% (48-84%)	1.2 (0.56-2.1)	3% (1.8-4.3%)	4.6% (2.3-7.6%)	33 (16-55)	3500 (2100-5300)
Venezuela	1990	Female	30 to 34	34% (16-55%)	66% (45-84%)	1.3 (0.66-2.3)	3.1% (1.9-4.7%)	5.2% (2.9-8.2%)	41 (23-67)	3500 (2100-5300)
Venezuela	1990	Female	35 to 39	36% (18-56%)	64% (44-82%)	1.6 (0.72-2.6)	3.7% (2.1-5.5%)	6.3% (3.4-9.8%)	58 (31-92)	3800 (2200-5800)
Venezuela	1990	Female	40 to 44	37% (21-58%)	63% (42-79%)	1.9 (0.98-3.2)	4.6% (2.7-7.1%)	8.1% (4.7-12%)	85 (49-130)	4700 (2700-7100)
Venezuela	1990	Female	45 to 49	37% (18-57%)	63% (43-82%)	2.4 (1-4.1)	5.8% (3.2-9.2%)	9.7% (5.3-15%)	110 (60-180)	5400 (2900-8500)
Venezuela	1990	Female	50 to 54	36% (19-58%)	64% (42-81%)	2.7 (1.3-4.7)	7.2% (4.1-11%)	11% (6.6-17%)	170 (99-250)	6800 (4000-10000)
Venezuela	1990	Female	55 to 59	34% (18-56%)	66% (44-82%)	2.9 (1.4-5.2)	7.8% (4.5-12%)	11% (6.7-17%)	220 (130-330)	7500 (4300-12000)
Venezuela	1990	Female	60 to 64	32% (16-54%)	68% (46-84%)	2.9 (1.2-5.5)	7.9% (4.4-13%)	11% (6.1-17%)	260 (150-420)	7700 (4300-12000)
Venezuela	1990	Female	65 to 69	30% (15-51%)	70% (49-85%)	2.7 (1.1-4.9)	7.9% (4.1-13%)	10% (5.5-17%)	290 (160-470)	7000 (3700-11000)
Venezuela	1990	Female	70 to 74	27% (14-45%)	73% (55-86%)	2.3 (0.93-4.5)	7.2% (3.5-12%)	9.2% (4.6-15%)	300 (140-500)	5600 (2700-9300)
Venezuela	1990	Female	75 to 79	23% (11-41%)	77% (59-89%)	1.9 (0.68-3.9)	6% (2.3-11%)	7.4% (2.9-14%)	260 (100-490)	3800 (1500-7000)
Venezuela	1990	Female	80 to 84	21% (9.1-39%)	79% (61-91%)	1.6 (0.54-3.4)	5.2% (1.9-10%)	6.4% (2.3-12%)	220 (77-420)	2300 (850-4500)
Venezuela	1990	Female	85 to 89	22% (11-38%)	78% (62-89%)	1.6 (0.62-3.2)	4.9% (1.7-9.7%)	5.9% (2.1-12%)	150 (54-310)	1200 (420-2300)
Venezuela	1990	Female	90 to 94	26% (12-44%)	74% (56-88%)	1.7 (0.65-3.5)	4.7% (1.4-10%)	5.6% (1.5-12%)	69 (19-150)	360 (110-770)
Venezuela	1990	Female	95 plus	32% (14-54%)	68% (46-86%)	1.9 (0.61-4.1)	4.6% (0.64-11%)	5.3% (0.58-13%)	20 (2.2-50)	70 (9.8-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	1990	Male	All Ages	41% (32-52%)	59% (48-68%)	3.1 (2.7-3.5)	8.7% (7.2-10%)	12% (9.8-14%)	5900 (4800-7300)	250000 (200000-290000)
Venezuela	1990	Male	15 to 19	29% (14-49%)	71% (51-86%)	1.4 (0.57-2.7)	6.4% (3.5-11%)	8.2% (3.7-15%)	120 (56-230)	11000 (5900-19000)
Venezuela	1990	Male	20 to 24	49% (27-70%)	51% (30-73%)	3.3 (1.6-5.7)	14% (8.6-21%)	17% (9.2-27%)	360 (190-570)	29000 (17000-43000)
Venezuela	1990	Male	25 to 29	59% (36-78%)	41% (22-64%)	4.7 (2.4-7.6)	17% (11-23%)	21% (13-31%)	420 (250-610)	30000 (20000-43000)
Venezuela	1990	Male	30 to 34	65% (40-82%)	35% (18-60%)	5.6 (3.1-8.5)	17% (12-22%)	23% (15-30%)	400 (270-540)	27000 (18000-35000)
Venezuela	1990	Male	35 to 39	68% (48-84%)	32% (16-52%)	6.3 (3.3-10)	17% (12-22%)	23% (16-31%)	380 (260-510)	22000 (16000-29000)
Venezuela	1990	Male	40 to 44	69% (50-85%)	31% (15-50%)	7 (3.7-11)	17% (13-22%)	23% (17-29%)	400 (290-520)	21000 (16000-27000)
Venezuela	1990	Male	45 to 49	70% (46-85%)	30% (15-54%)	7.5 (3.6-13)	16% (12-21%)	22% (16-28%)	410 (290-540)	19000 (14000-24000)
Venezuela	1990	Male	50 to 54	69% (47-85%)	31% (15-53%)	8 (4.1-13)	16% (11-20%)	20% (15-26%)	490 (350-630)	19000 (14000-25000)
Venezuela	1990	Male	55 to 59	68% (46-85%)	32% (15-54%)	8.5 (4.1-14)	15% (11-20%)	19% (13-24%)	550 (380-730)	19000 (13000-25000)
Venezuela	1990	Male	60 to 64	65% (41-83%)	35% (17-59%)	7.6 (3.5-13)	13% (9.2-18%)	16% (11-22%)	580 (400-800)	17000 (12000-23000)
Venezuela	1990	Male	65 to 69	61% (37-80%)	39% (20-63%)	5.8 (2.7-10)	12% (7.8-17%)	14% (9.4-20%)	550 (360-770)	13000 (8600-18000)
Venezuela	1990	Male	70 to 74	55% (33-75%)	45% (25-67%)	4.2 (1.8-8.3)	9.7% (5.9-14%)	11% (6.8-17%)	460 (270-680)	8700 (5300-13000)
Venezuela	1990	Male	75 to 79	48% (25-71%)	52% (29-75%)	3.2 (1.2-6.9)	7.7% (3.8-12%)	8.9% (4.3-14%)	340 (170-540)	5000 (2500-8000)
Venezuela	1990	Male	80 to 84	43% (22-66%)	57% (34-78%)	2.8 (1-5.6)	7% (3.5-11%)	8% (3.9-13%)	240 (120-400)	2600 (1300-4300)
Venezuela	1990	Male	85 to 89	42% (21-65%)	58% (35-79%)	2.7 (1-5.7)	6.6% (3.1-11%)	7.6% (3.4-13%)	150 (71-260)	1200 (560-2000)
Venezuela	1990	Male	90 to 94	46% (25-68%)	54% (32-75%)	3 (1.1-6.4)	6.4% (2.6-10%)	7.3% (2.8-12%)	65 (25-110)	350 (140-570)
Venezuela	1990	Male	95 plus	50% (25-74%)	50% (26-75%)	3.4 (1-7.7)	6.6% (2.2-12%)	7.4% (2.1-14%)	17 (4.9-32)	64 (20-110)
Venezuela	1995	Female	All Ages	23% (18-31%)	77% (69-82%)	0.97 (0.79-1.2)	3.1% (2.3-4.1%)	6.1% (3.9-8.7%)	2500 (1600-3600)	73000 (54000-97000)
Venezuela	1995	Female	15 to 19	21% (10-36%)	79% (64-90%)	0.8 (0.35-1.5)	2% (1.1-3.3%)	3.5% (1.5-6.6%)	22 (9.3-43)	2500 (1300-4200)
Venezuela	1995	Female	20 to 24	29% (14-47%)	71% (53-86%)	1.1 (0.47-2)	2.9% (1.7-4.5%)	4.4% (2-8%)	30 (14-53)	3500 (2000-5600)
Venezuela	1995	Female	25 to 29	31% (16-51%)	69% (49-84%)	1.1 (0.52-2)	2.9% (1.8-4.2%)	4.4% (2.2-7.4%)	33 (16-55)	3600 (2200-5500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	1995	Female	30 to 34	33% (16-53%)	67% (47-84%)	1.2 (0.61-2.1)	2.9% (1.8-4.2%)	4.8% (2.7-7.7%)	43 (24-69)	3800 (2300-5600)
Venezuela	1995	Female	35 to 39	35% (18-55%)	65% (45-82%)	1.5 (0.71-2.5)	3.3% (1.9-5%)	5.8% (3.1-8.9%)	61 (32-96)	4200 (2400-6300)
Venezuela	1995	Female	40 to 44	35% (19-56%)	65% (44-81%)	1.8 (0.91-3)	4.2% (2.5-6.6%)	7.3% (4.2-11%)	91 (50-140)	5100 (3000-7900)
Venezuela	1995	Female	45 to 49	36% (18-55%)	64% (45-82%)	2.2 (0.96-3.7)	5.2% (3-8.4%)	8.8% (4.9-14%)	120 (69-200)	6000 (3300-9700)
Venezuela	1995	Female	50 to 54	35% (18-56%)	65% (44-82%)	2.5 (1.2-4.5)	6.5% (3.7-10%)	10% (5.9-16%)	170 (95-270)	7100 (4000-11000)
Venezuela	1995	Female	55 to 59	33% (17-55%)	67% (45-83%)	2.8 (1.3-5.1)	7.2% (4-11%)	11% (6-17%)	220 (120-340)	7600 (4300-12000)
Venezuela	1995	Female	60 to 64	31% (16-52%)	69% (48-84%)	2.8 (1.2-5.3)	7.3% (4-12%)	10% (5.6-16%)	270 (150-420)	7800 (4200-13000)
Venezuela	1995	Female	65 to 69	30% (15-49%)	70% (51-85%)	2.5 (1-4.5)	7.4% (3.8-12%)	9.9% (5.1-16%)	310 (160-500)	7600 (3800-12000)
Venezuela	1995	Female	70 to 74	26% (12-44%)	74% (56-88%)	2.2 (0.87-4.3)	6.7% (3.3-11%)	8.7% (4.3-15%)	310 (150-530)	6000 (2900-10000)
Venezuela	1995	Female	75 to 79	22% (9.8-39%)	78% (61-90%)	1.7 (0.63-3.6)	5.6% (2.1-10%)	6.9% (2.6-13%)	270 (100-500)	4000 (1500-7200)
Venezuela	1995	Female	80 to 84	20% (8.5-38%)	80% (62-91%)	1.5 (0.5-3.2)	4.8% (1.7-9.1%)	6% (2.1-11%)	240 (86-450)	2600 (940-4900)
Venezuela	1995	Female	85 to 89	21% (9.4-37%)	79% (63-91%)	1.5 (0.57-3)	4.4% (1.6-8.9%)	5.4% (1.9-11%)	170 (58-340)	1300 (460-2600)
Venezuela	1995	Female	90 to 94	25% (12-43%)	75% (57-88%)	1.6 (0.59-3.3)	4.3% (1.3-9.2%)	5.1% (1.5-11%)	83 (24-180)	440 (130-920)
Venezuela	1995	Female	95 plus	31% (13-52%)	69% (48-87%)	1.8 (0.54-4)	4.1% (0.54-10%)	4.9% (0.45-12%)	24 (2.2-60)	84 (11-200)
Venezuela	1995	Male	All Ages	42% (32-53%)	58% (47-68%)	3 (2.7-3.4)	9.1% (7.5-11%)	12% (9.9-14%)	6900 (5500-8500)	290000 (230000-350000)
Venezuela	1995	Male	15 to 19	29% (14-48%)	71% (52-86%)	1.4 (0.53-2.6)	6.4% (3.4-11%)	8% (3.6-15%)	160 (70-300)	14000 (7100-24000)
Venezuela	1995	Male	20 to 24	48% (26-70%)	52% (30-74%)	3.1 (1.5-5.5)	14% (8.1-21%)	17% (8.7-26%)	450 (230-720)	35000 (20000-53000)
Venezuela	1995	Male	25 to 29	58% (36-78%)	42% (22-64%)	4.4 (2.3-7.3)	16% (11-23%)	20% (12-30%)	510 (300-750)	36000 (23000-52000)
Venezuela	1995	Male	30 to 34	64% (41-82%)	36% (18-59%)	5.4 (3-8.1)	16% (12-21%)	22% (14-29%)	500 (330-680)	33000 (23000-43000)
Venezuela	1995	Male	35 to 39	68% (46-84%)	32% (16-54%)	6 (3.2-9.7)	16% (11-22%)	22% (15-30%)	470 (310-640)	27000 (19000-36000)
Venezuela	1995	Male	40 to 44	69% (50-84%)	31% (16-50%)	6.7 (3.6-10)	16% (12-21%)	22% (16-29%)	500 (350-660)	26000 (19000-33000)
Venezuela	1995	Male	45 to 49	69% (45-84%)	31% (16-55%)	7.2 (3.5-12)	16% (11-20%)	21% (15-27%)	520 (370-670)	23000 (17000-30000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	1995	Male	50 to 54	68% (46-84%)	32% (16-54%)	7.6 (3.9-13)	15% (11-20%)	20% (14-25%)	540 (390-720)	22000 (16000-28000)
Venezuela	1995	Male	55 to 59	67% (45-85%)	33% (15-55%)	8 (3.9-13)	14% (10-19%)	18% (13-23%)	580 (410-770)	20000 (14000-26000)
Venezuela	1995	Male	60 to 64	64% (40-83%)	36% (17-60%)	7.1 (3.4-13)	13% (8.8-17%)	15% (11-21%)	610 (420-830)	18000 (12000-24000)
Venezuela	1995	Male	65 to 69	60% (36-79%)	40% (21-64%)	5.5 (2.5-9.6)	11% (7.5-16%)	14% (8.9-19%)	610 (400-850)	15000 (9500-20000)
Venezuela	1995	Male	70 to 74	54% (32-74%)	46% (26-68%)	4 (1.7-7.7)	9.1% (5.6-13%)	11% (6.3-16%)	500 (300-740)	9600 (5800-14000)
Venezuela	1995	Male	75 to 79	47% (25-70%)	53% (30-75%)	3 (1.1-6.6)	7.2% (3.5-11%)	8.3% (3.9-13%)	370 (180-600)	5400 (2600-8700)
Venezuela	1995	Male	80 to 84	42% (21-65%)	58% (35-79%)	2.6 (1-5.2)	6.5% (3.2-10%)	7.5% (3.7-12%)	270 (130-450)	3000 (1500-4800)
Venezuela	1995	Male	85 to 89	41% (20-64%)	59% (36-80%)	2.6 (0.94-5.3)	6.1% (2.8-10%)	7% (3.1-12%)	170 (78-290)	1400 (610-2300)
Venezuela	1995	Male	90 to 94	45% (23-67%)	55% (33-77%)	2.9 (0.99-6.2)	5.8% (2.4-9.7%)	6.7% (2.6-11%)	72 (27-120)	390 (160-650)
Venezuela	1995	Male	95 plus	50% (24-74%)	50% (26-76%)	3.3 (1-7.5)	6.1% (1.9-11%)	6.8% (1.8-13%)	19 (5-35)	71 (22-130)
Venezuela	2000	Female	All Ages	23% (17-30%)	77% (70-83%)	0.92 (0.75-1.1)	3% (2.2-3.9%)	5.7% (3.6-8.2%)	2400 (1500-3500)	72000 (52000-94000)
Venezuela	2000	Female	15 to 19	20% (10-34%)	80% (66-90%)	0.72 (0.32-1.3)	1.8% (1-3%)	3.3% (1.4-6.3%)	21 (8.8-40)	2400 (1300-4100)
Venezuela	2000	Female	20 to 24	27% (13-44%)	73% (56-87%)	0.96 (0.41-1.7)	2.7% (1.6-4.1%)	4% (1.8-7%)	29 (13-52)	3600 (2100-5600)
Venezuela	2000	Female	25 to 29	30% (15-50%)	70% (50-85%)	1 (0.46-1.8)	2.6% (1.6-3.9%)	4% (1.9-6.8%)	30 (14-52)	3400 (2000-5200)
Venezuela	2000	Female	30 to 34	32% (15-52%)	68% (48-85%)	1.1 (0.56-2)	2.6% (1.6-3.8%)	4.3% (2.3-7%)	37 (20-62)	3500 (2100-5200)
Venezuela	2000	Female	35 to 39	34% (17-54%)	66% (46-83%)	1.3 (0.61-2.3)	2.8% (1.7-4.4%)	5.1% (2.7-8.1%)	56 (30-91)	4100 (2400-6300)
Venezuela	2000	Female	40 to 44	34% (19-55%)	66% (45-81%)	1.6 (0.8-2.8)	3.6% (2.1-5.7%)	6.4% (3.7-10%)	87 (48-140)	5100 (2900-8100)
Venezuela	2000	Female	45 to 49	34% (17-54%)	66% (46-83%)	2 (0.89-3.5)	4.7% (2.6-7.7%)	7.9% (4.3-13%)	130 (71-210)	6400 (3500-10000)
Venezuela	2000	Female	50 to 54	34% (17-54%)	66% (46-83%)	2.3 (1.1-4.1)	5.7% (3.2-8.9%)	9.1% (5.1-14%)	170 (98-270)	7300 (4200-12000)
Venezuela	2000	Female	55 to 59	32% (16-51%)	68% (49-84%)	2.5 (1.2-4.5)	6.1% (3.4-10%)	9.4% (5.4-15%)	200 (110-320)	7200 (4000-12000)
Venezuela	2000	Female	60 to 64	30% (14-50%)	70% (50-86%)	2.5 (1-4.8)	6.2% (3.3-10%)	8.9% (4.8-14%)	240 (130-390)	7200 (3800-12000)
Venezuela	2000	Female	65 to 69	28% (14-48%)	72% (52-86%)	2.3 (0.96-4.2)	6.4% (3.3-11%)	8.7% (4.5-14%)	290 (150-480)	7100 (3700-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	2000	Female	70 to 74	25% (12-42%)	75% (58-88%)	2 (0.8-4)	5.9% (2.9-10%)	7.8% (3.7-13%)	310 (150-520)	5900 (2800-10000)
Venezuela	2000	Female	75 to 79	22% (9.8-39%)	78% (61-90%)	1.6 (0.59-3.3)	4.9% (1.9-9.1%)	6.2% (2.4-11%)	270 (100-490)	4000 (1500-7300)
Venezuela	2000	Female	80 to 84	20% (8.5-37%)	80% (63-91%)	1.4 (0.47-3)	4.1% (1.5-7.9%)	5.3% (1.9-10%)	220 (78-430)	2400 (880-4800)
Venezuela	2000	Female	85 to 89	20% (9.1-36%)	80% (64-91%)	1.4 (0.53-2.8)	3.8% (1.3-7.8%)	4.7% (1.6-9.7%)	160 (54-350)	1300 (450-2700)
Venezuela	2000	Female	90 to 94	24% (11-42%)	76% (58-89%)	1.5 (0.54-3)	3.6% (1-8%)	4.4% (1.2-9.9%)	90 (24-200)	480 (140-1000)
Venezuela	2000	Female	95 plus	29% (13-52%)	71% (48-87%)	1.6 (0.49-3.5)	3.6% (0.45-9%)	4.2% (0.32-11%)	28 (2.2-75)	100 (13-250)
Venezuela	2000	Male	All Ages	41% (31-53%)	59% (47-69%)	2.9 (2.5-3.4)	9.5% (7.9-11%)	12% (9.7-14%)	7600 (6100-9200)	320000 (260000-400000)
Venezuela	2000	Male	15 to 19	27% (13-46%)	73% (54-87%)	1.3 (0.5-2.4)	6.1% (3.1-11%)	7.4% (3.2-14%)	190 (80-360)	16000 (8000-28000)
Venezuela	2000	Male	20 to 24	46% (25-68%)	54% (32-75%)	2.9 (1.3-5.1)	14% (7.5-21%)	15% (7.8-25%)	580 (290-960)	43000 (24000-69000)
Venezuela	2000	Male	25 to 29	57% (33-77%)	43% (23-67%)	4.1 (2-6.9)	16% (9.9-23%)	19% (11-28%)	590 (340-880)	41000 (26000-59000)
Venezuela	2000	Male	30 to 34	63% (39-82%)	37% (18-61%)	5 (2.8-7.8)	16% (11-21%)	20% (13-28%)	550 (350-760)	36000 (25000-48000)
Venezuela	2000	Male	35 to 39	66% (44-83%)	34% (17-56%)	5.6 (3-9)	15% (11-21%)	21% (14-28%)	540 (360-740)	32000 (22000-43000)
Venezuela	2000	Male	40 to 44	67% (47-84%)	33% (16-53%)	6.2 (3.3-9.6)	15% (11-20%)	21% (15-27%)	560 (400-760)	29000 (21000-39000)
Venezuela	2000	Male	45 to 49	67% (44-83%)	33% (17-56%)	6.6 (3.2-11)	15% (11-20%)	20% (14-26%)	600 (420-800)	28000 (20000-36000)
Venezuela	2000	Male	50 to 54	67% (45-83%)	33% (17-55%)	7 (3.6-12)	14% (10-19%)	19% (13-24%)	620 (440-830)	25000 (18000-33000)
Venezuela	2000	Male	55 to 59	65% (43-84%)	35% (16-57%)	7.4 (3.5-13)	13% (9.6-18%)	17% (12-22%)	610 (430-820)	21000 (15000-28000)
Venezuela	2000	Male	60 to 64	63% (39-82%)	37% (18-61%)	6.7 (3.1-12)	12% (8.1-16%)	14% (10-20%)	600 (420-840)	18000 (12000-24000)
Venezuela	2000	Male	65 to 69	59% (35-78%)	41% (22-65%)	5.1 (2.3-8.9)	11% (7-15%)	13% (8.4-18%)	630 (420-880)	15000 (10000-21000)
Venezuela	2000	Male	70 to 74	52% (31-73%)	48% (27-69%)	3.7 (1.5-7.5)	8.4% (5-12%)	9.8% (5.9-15%)	530 (310-790)	10000 (6100-15000)
Venezuela	2000	Male	75 to 79	46% (24-69%)	54% (31-76%)	2.8 (1-6.2)	6.6% (3.2-11%)	7.7% (3.7-13%)	380 (180-630)	5700 (2800-9300)
Venezuela	2000	Male	80 to 84	41% (20-64%)	59% (36-80%)	2.4 (0.88-4.8)	5.9% (2.8-9.7%)	6.8% (3.2-11%)	290 (140-470)	3200 (1500-5200)
Venezuela	2000	Male	85 to 89	40% (20-63%)	60% (37-80%)	2.4 (0.86-5)	5.5% (2.4-9.3%)	6.3% (2.7-11%)	180 (77-310)	1400 (620-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	2000	Male	90 to 94	43% (23-66%)	57% (34-77%)	2.6 (0.92-5.7)	5.3% (2-9%)	6.1% (2.2-11%)	80 (28-140)	440 (170-750)
Venezuela	2000	Male	95 plus	49% (23-73%)	51% (27-77%)	3 (0.88-7.1)	5.5% (1.7-10%)	6.2% (1.5-12%)	22 (5.5-42)	83 (25-150)
Venezuela	2005	Female	All Ages	22% (17-30%)	78% (70-83%)	0.93 (0.76-1.1)	3% (2.2-4%)	5.6% (3.5-8.1%)	2600 (1600-3800)	78000 (57000-100000)
Venezuela	2005	Female	15 to 19	19% (9.4-32%)	81% (68-91%)	0.69 (0.3-1.3)	1.8% (0.99-2.9%)	3.2% (1.4-6.1%)	23 (9.8-43)	2600 (1400-4400)
Venezuela	2005	Female	20 to 24	26% (12-44%)	74% (56-88%)	0.92 (0.4-1.7)	2.6% (1.5-4.1%)	3.9% (1.7-7%)	31 (14-57)	3800 (2100-6000)
Venezuela	2005	Female	25 to 29	29% (13-47%)	71% (53-87%)	0.99 (0.41-1.7)	2.6% (1.6-3.8%)	3.8% (1.7-6.6%)	33 (14-59)	3700 (2200-5700)
Venezuela	2005	Female	30 to 34	31% (14-51%)	69% (49-86%)	1.1 (0.5-1.9)	2.5% (1.5-3.8%)	4.1% (2.2-6.9%)	38 (20-65)	3500 (2200-5400)
Venezuela	2005	Female	35 to 39	32% (17-52%)	68% (48-83%)	1.3 (0.59-2.2)	2.7% (1.6-4.2%)	4.9% (2.6-7.7%)	55 (29-88)	4100 (2400-6300)
Venezuela	2005	Female	40 to 44	33% (17-53%)	67% (47-83%)	1.5 (0.78-2.6)	3.3% (1.9-5.1%)	6.1% (3.5-9.6%)	90 (52-140)	5500 (3300-8600)
Venezuela	2005	Female	45 to 49	33% (16-52%)	67% (48-84%)	1.9 (0.86-3.2)	4.4% (2.5-7.1%)	7.5% (4.2-12%)	140 (79-230)	7100 (4100-11000)
Venezuela	2005	Female	50 to 54	32% (16-53%)	68% (47-84%)	2.2 (1.1-3.8)	5.5% (3.1-8.5%)	8.7% (5-14%)	200 (110-320)	8400 (4800-13000)
Venezuela	2005	Female	55 to 59	31% (15-51%)	69% (49-85%)	2.4 (1.1-4.3)	5.9% (3.3-9.5%)	9% (5.1-14%)	230 (130-370)	8500 (4700-13000)
Venezuela	2005	Female	60 to 64	29% (14-49%)	71% (51-86%)	2.4 (1-4.6)	5.8% (3-9.5%)	8.5% (4.6-14%)	250 (130-400)	7600 (3900-12000)
Venezuela	2005	Female	65 to 69	27% (13-47%)	73% (53-87%)	2.2 (0.94-4.1)	5.9% (3-10%)	8.3% (4.3-14%)	300 (150-490)	7400 (3800-12000)
Venezuela	2005	Female	70 to 74	25% (12-42%)	75% (58-88%)	1.9 (0.77-3.8)	5.5% (2.7-9.4%)	7.4% (3.6-12%)	320 (160-540)	6200 (3000-10000)
Venezuela	2005	Female	75 to 79	21% (9.1-37%)	79% (63-91%)	1.6 (0.56-3.3)	4.7% (1.9-8.8%)	6% (2.4-11%)	300 (120-560)	4500 (1800-8200)
Venezuela	2005	Female	80 to 84	19% (8.2-35%)	81% (65-92%)	1.3 (0.46-2.8)	3.9% (1.5-7.5%)	5.1% (1.9-9.9%)	250 (90-490)	2800 (1000-5500)
Venezuela	2005	Female	85 to 89	19% (8.9-34%)	81% (66-91%)	1.3 (0.51-2.7)	3.6% (1.3-7.2%)	4.5% (1.6-9.3%)	190 (66-390)	1500 (550-3100)
Venezuela	2005	Female	90 to 94	23% (11-40%)	77% (60-89%)	1.4 (0.53-3.1)	3.6% (1-7.9%)	4.4% (1.2-9.8%)	120 (32-270)	610 (180-1400)
Venezuela	2005	Female	95 plus	28% (12-50%)	72% (50-88%)	1.6 (0.49-3.6)	3.5% (0.45-8.8%)	4.2% (0.35-11%)	39 (3.2-100)	140 (17-350)
Venezuela	2005	Male	All Ages	41% (31-53%)	59% (47-69%)	3 (2.6-3.5)	9.9% (7.9-12%)	12% (9.7-14%)	8400 (6800-11000)	370000 (290000-470000)
Venezuela	2005	Male	15 to 19	26% (12-45%)	74% (55-88%)	1.2 (0.46-2.3)	6.1% (3-11%)	7.3% (3-15%)	230 (94-440)	19000 (9100-34000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	2005	Male	20 to 24	45% (24-67%)	55% (33-76%)	2.8 (1.2-5)	14% (7.3-22%)	15% (7.3-26%)	720 (340-1200)	53000 (28000-87000)
Venezuela	2005	Male	25 to 29	55% (33-75%)	45% (25-67%)	4 (1.9-6.6)	16% (9.7-24%)	19% (10-29%)	760 (410-1200)	52000 (31000-78000)
Venezuela	2005	Male	30 to 34	62% (38-80%)	38% (20-62%)	4.8 (2.6-7.6)	16% (11-22%)	20% (13-29%)	630 (380-910)	40000 (27000-56000)
Venezuela	2005	Male	35 to 39	65% (43-83%)	35% (17-57%)	5.5 (2.9-8.9)	15% (10-21%)	21% (14-29%)	580 (370-820)	34000 (23000-47000)
Venezuela	2005	Male	40 to 44	66% (46-83%)	34% (17-54%)	6.1 (3.2-9.4)	15% (10-20%)	20% (14-27%)	610 (420-830)	32000 (23000-43000)
Venezuela	2005	Male	45 to 49	67% (42-83%)	33% (17-58%)	6.5 (3.2-11)	14% (10-19%)	19% (14-25%)	650 (440-860)	30000 (21000-39000)
Venezuela	2005	Male	50 to 54	66% (43-84%)	34% (16-57%)	7 (3.5-12)	14% (9.8-18%)	18% (13-24%)	710 (490-940)	28000 (20000-37000)
Venezuela	2005	Male	55 to 59	64% (43-83%)	36% (17-57%)	7.3 (3.5-12)	13% (9-17%)	16% (11-22%)	700 (480-960)	24000 (17000-33000)
Venezuela	2005	Male	60 to 64	62% (38-81%)	38% (19-62%)	6.5 (3-12)	11% (7.5-15%)	14% (9.5-19%)	630 (430-890)	19000 (13000-26000)
Venezuela	2005	Male	65 to 69	58% (34-77%)	42% (23-66%)	5 (2.3-8.9)	9.9% (6.5-14%)	12% (8-17%)	620 (420-900)	15000 (10000-22000)
Venezuela	2005	Male	70 to 74	52% (31-73%)	48% (27-69%)	3.6 (1.6-7.3)	7.9% (4.8-12%)	9.5% (5.6-14%)	530 (320-810)	10000 (6300-16000)
Venezuela	2005	Male	75 to 79	45% (23-68%)	55% (32-77%)	2.8 (0.98-6.2)	6.3% (3-10%)	7.4% (3.5-12%)	420 (200-690)	6300 (3000-10000)
Venezuela	2005	Male	80 to 84	40% (19-63%)	60% (37-81%)	2.4 (0.82-4.7)	5.6% (2.6-9.4%)	6.5% (2.9-11%)	310 (140-530)	3500 (1600-5900)
Venezuela	2005	Male	85 to 89	39% (19-62%)	61% (38-81%)	2.3 (0.81-4.9)	5.2% (2.2-8.9%)	6% (2.5-10%)	200 (84-350)	1600 (690-2700)
Venezuela	2005	Male	90 to 94	42% (21-65%)	58% (35-79%)	2.5 (0.84-5.6)	5.1% (1.9-8.8%)	5.8% (2.1-10%)	98 (35-170)	540 (210-930)
Venezuela	2005	Male	95 plus	47% (22-72%)	53% (28-78%)	2.9 (0.81-7)	5.4% (1.6-10%)	6% (1.5-12%)	28 (7-53)	100 (32-200)
Venezuela	2010	Female	All Ages	22% (16-28%)	78% (72-84%)	0.99 (0.78-1.2)	3.1% (2.2-4.2%)	5.6% (3.5-8.2%)	3000 (1800-4300)	88000 (62000-120000)
Venezuela	2010	Female	15 to 19	18% (8.9-31%)	82% (69-91%)	0.7 (0.31-1.3)	1.8% (0.95-2.9%)	3.2% (1.3-6.1%)	23 (9.6-44)	2600 (1400-4500)
Venezuela	2010	Female	20 to 24	25% (12-42%)	75% (58-88%)	0.92 (0.39-1.7)	2.6% (1.5-4%)	3.8% (1.7-7.1%)	32 (14-59)	4100 (2400-6400)
Venezuela	2010	Female	25 to 29	27% (13-46%)	73% (54-87%)	0.97 (0.4-1.8)	2.5% (1.6-3.8%)	3.7% (1.6-6.5%)	35 (15-62)	4000 (2400-6200)
Venezuela	2010	Female	30 to 34	29% (13-49%)	71% (51-87%)	1.1 (0.48-1.9)	2.4% (1.5-3.7%)	3.9% (2-6.5%)	41 (20-67)	3900 (2400-5900)
Venezuela	2010	Female	35 to 39	31% (15-49%)	69% (51-85%)	1.3 (0.56-2.2)	2.7% (1.6-4.1%)	4.7% (2.4-7.4%)	55 (28-89)	4200 (2500-6300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	2010	Female	40 to 44	32% (17-50%)	68% (50-83%)	1.5 (0.78-2.6)	3.2% (1.9-5%)	5.9% (3.3-9.1%)	91 (50-150)	5600 (3300-8800)
Venezuela	2010	Female	45 to 49	32% (15-51%)	68% (49-85%)	1.9 (0.83-3.4)	4.2% (2.3-6.7%)	7.4% (4-12%)	150 (83-250)	7900 (4300-13000)
Venezuela	2010	Female	50 to 54	31% (16-52%)	69% (48-84%)	2.2 (1-3.9)	5.2% (2.9-8.3%)	8.4% (4.8-13%)	220 (120-350)	9600 (5600-15000)
Venezuela	2010	Female	55 to 59	30% (15-50%)	70% (50-85%)	2.4 (1.1-4.3)	5.8% (3.2-9.5%)	8.8% (4.8-14%)	280 (160-460)	10000 (5700-17000)
Venezuela	2010	Female	60 to 64	28% (13-48%)	72% (52-87%)	2.4 (1-4.5)	5.8% (3-9.5%)	8.3% (4.5-14%)	300 (160-500)	9200 (4900-15000)
Venezuela	2010	Female	65 to 69	27% (13-45%)	73% (55-87%)	2.2 (0.92-4.1)	5.7% (2.8-9.6%)	8.1% (4.1-13%)	320 (160-520)	8100 (4100-13000)
Venezuela	2010	Female	70 to 74	24% (12-40%)	76% (60-88%)	1.9 (0.77-4)	5.3% (2.6-9%)	7.2% (3.5-12%)	340 (160-590)	6800 (3200-12000)
Venezuela	2010	Female	75 to 79	20% (9.2-36%)	80% (64-91%)	1.6 (0.57-3.3)	4.5% (1.8-8.3%)	5.8% (2.3-11%)	320 (130-600)	4900 (2000-9100)
Venezuela	2010	Female	80 to 84	18% (7.7-34%)	82% (66-92%)	1.4 (0.45-2.8)	3.9% (1.4-7.5%)	5% (1.8-9.8%)	310 (110-590)	3400 (1300-6600)
Venezuela	2010	Female	85 to 89	19% (8.6-34%)	81% (66-91%)	1.3 (0.51-2.8)	3.6% (1.3-7.4%)	4.6% (1.6-9.5%)	240 (83-500)	1900 (670-4000)
Venezuela	2010	Female	90 to 94	22% (10-40%)	78% (60-90%)	1.5 (0.55-3.2)	3.6% (1.1-8.1%)	4.5% (1.2-10%)	150 (41-330)	800 (240-1800)
Venezuela	2010	Female	95 plus	27% (11-50%)	73% (50-89%)	1.6 (0.48-3.7)	3.6% (0.56-8.9%)	4.4% (0.41-11%)	53 (5.2-140)	190 (28-480)
Venezuela	2010	Male	All Ages	41% (32-52%)	59% (48-68%)	3.2 (2.6-3.8)	10% (8.1-13%)	12% (9.5-15%)	9900 (7800-12000)	430000 (340000-550000)
Venezuela	2010	Male	15 to 19	25% (12-44%)	75% (56-88%)	1.2 (0.47-2.4)	6.4% (3-12%)	7.5% (3-15%)	280 (120-560)	22000 (10000-43000)
Venezuela	2010	Male	20 to 24	44% (23-66%)	56% (34-77%)	2.8 (1.2-5)	14% (7.4-22%)	15% (7.4-26%)	870 (420-1500)	63000 (33000-110000)
Venezuela	2010	Male	25 to 29	54% (32-74%)	46% (26-68%)	4 (1.8-7)	16% (9.6-24%)	19% (10-29%)	900 (490-1400)	61000 (36000-92000)
Venezuela	2010	Male	30 to 34	61% (37-80%)	39% (20-63%)	4.9 (2.5-7.9)	16% (10-23%)	20% (12-29%)	760 (460-1100)	48000 (31000-68000)
Venezuela	2010	Male	35 to 39	64% (43-81%)	36% (19-57%)	5.5 (2.8-9.2)	16% (10-22%)	21% (13-29%)	630 (400-910)	37000 (25000-52000)
Venezuela	2010	Male	40 to 44	65% (44-82%)	35% (18-56%)	6.1 (3.1-9.7)	15% (10-19%)	20% (14-27%)	640 (430-860)	34000 (24000-45000)
Venezuela	2010	Male	45 to 49	66% (41-82%)	34% (18-59%)	6.6 (3.2-11)	14% (9.8-18%)	19% (13-25%)	720 (500-950)	34000 (24000-44000)
Venezuela	2010	Male	50 to 54	65% (42-82%)	35% (18-58%)	6.9 (3.6-12)	14% (9.5-18%)	18% (13-23%)	830 (570-1100)	33000 (23000-44000)
Venezuela	2010	Male	55 to 59	63% (41-83%)	37% (17-59%)	7.3 (3.5-12)	13% (8.9-17%)	16% (11-21%)	870 (590-1200)	30000 (20000-41000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	2010	Male	60 to 64	61% (38-81%)	39% (19-62%)	6.6 (3-12)	11% (7.4-15%)	13% (9-19%)	780 (520-1100)	23000 (16000-33000)
Venezuela	2010	Male	65 to 69	57% (33-77%)	43% (23-67%)	5.1 (2.3-9.1)	9.8% (6.3-14%)	12% (7.7-17%)	710 (460-1000)	18000 (11000-25000)
Venezuela	2010	Male	70 to 74	51% (30-72%)	49% (28-70%)	3.7 (1.5-7.8)	7.8% (4.6-12%)	9.4% (5.5-14%)	600 (350-910)	12000 (6900-18000)
Venezuela	2010	Male	75 to 79	44% (23-68%)	56% (32-77%)	2.8 (0.95-6.3)	6.1% (3-10%)	7.1% (3.4-12%)	480 (230-780)	7300 (3500-12000)
Venezuela	2010	Male	80 to 84	39% (19-62%)	61% (38-81%)	2.4 (0.82-4.7)	5.4% (2.6-9.1%)	6.3% (2.8-11%)	380 (180-640)	4200 (2000-7100)
Venezuela	2010	Male	85 to 89	37% (18-60%)	63% (40-82%)	2.3 (0.84-4.8)	5% (2.4-8.6%)	5.8% (2.6-10%)	240 (110-420)	1900 (890-3300)
Venezuela	2010	Male	90 to 94	40% (21-64%)	60% (36-79%)	2.5 (0.82-5.6)	5.1% (2-8.7%)	5.7% (2.1-10%)	120 (46-210)	670 (270-1200)
Venezuela	2010	Male	95 plus	45% (20-71%)	55% (29-80%)	2.9 (0.78-7.2)	5.4% (1.7-9.9%)	6% (1.6-11%)	36 (9.8-69)	130 (41-250)
Venezuela	2016	Female	All Ages	22% (17-29%)	78% (71-83%)	1.1 (0.77-1.5)	3.3% (2.2-4.5%)	5.6% (3.3-8.4%)	3600 (2000-5700)	100000 (67000-150000)
Venezuela	2016	Female	15 to 19	18% (8.6-31%)	82% (69-91%)	0.7 (0.28-1.4)	1.7% (0.91-3.1%)	3.2% (1.2-6.6%)	22 (8.6-47)	2600 (1300-4700)
Venezuela	2016	Female	20 to 24	25% (12-42%)	75% (58-88%)	0.94 (0.36-1.9)	2.6% (1.5-4.2%)	3.9% (1.6-7.4%)	32 (12-63)	4100 (2200-6900)
Venezuela	2016	Female	25 to 29	27% (13-46%)	73% (54-87%)	1 (0.38-2)	2.6% (1.5-4.1%)	3.9% (1.6-7%)	39 (15-75)	4400 (2400-7300)
Venezuela	2016	Female	30 to 34	29% (13-51%)	71% (49-87%)	1.1 (0.48-2.1)	2.5% (1.5-4%)	4% (1.9-7.1%)	46 (20-86)	4400 (2600-7300)
Venezuela	2016	Female	35 to 39	31% (15-51%)	69% (49-85%)	1.3 (0.55-2.5)	2.8% (1.5-4.4%)	4.9% (2.4-8.3%)	63 (30-110)	4900 (2700-7800)
Venezuela	2016	Female	40 to 44	32% (17-51%)	68% (49-83%)	1.6 (0.75-2.8)	3.2% (1.8-5.2%)	6% (3.1-9.7%)	92 (48-160)	5800 (3100-9700)
Venezuela	2016	Female	45 to 49	32% (16-51%)	68% (49-84%)	1.9 (0.8-3.5)	4.1% (2.2-6.7%)	7.3% (3.7-12%)	160 (81-280)	8300 (4400-14000)
Venezuela	2016	Female	50 to 54	31% (15-51%)	69% (49-85%)	2.2 (1-4)	5.1% (2.8-8.5%)	8.2% (4.6-13%)	270 (150-470)	12000 (6500-20000)
Venezuela	2016	Female	55 to 59	29% (15-50%)	71% (50-85%)	2.4 (1-4.6)	5.6% (3-9.2%)	8.5% (4.7-14%)	350 (180-570)	13000 (6700-21000)
Venezuela	2016	Female	60 to 64	28% (13-47%)	72% (53-87%)	2.4 (0.96-4.6)	5.5% (2.8-9.1%)	8.1% (4.2-13%)	360 (170-610)	11000 (5400-19000)
Venezuela	2016	Female	65 to 69	26% (12-45%)	74% (55-88%)	2.2 (0.84-4.2)	5.5% (2.7-9.2%)	7.9% (3.9-13%)	380 (180-660)	9800 (4500-17000)
Venezuela	2016	Female	70 to 74	24% (11-41%)	76% (59-89%)	2 (0.77-4.1)	5.2% (2.5-9%)	7.1% (3.5-12%)	410 (190-760)	8300 (3900-15000)
Venezuela	2016	Female	75 to 79	20% (8.9-37%)	80% (63-91%)	1.6 (0.56-3.6)	4.4% (1.7-8.1%)	5.7% (2.3-10%)	400 (150-760)	6100 (2400-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Venezuela	2016	Female	80 to 84	18% (7.6-33%)	82% (67-92%)	1.3 (0.44-2.8)	3.8% (1.3-7.2%)	4.9% (1.7-9.3%)	370 (130-720)	4200 (1500-8200)
Venezuela	2016	Female	85 to 89	18% (8.5-33%)	82% (67-91%)	1.3 (0.48-2.8)	3.5% (1.2-7.3%)	4.5% (1.5-9.3%)	310 (98-640)	2400 (810-5000)
Venezuela	2016	Female	90 to 94	22% (9.7-39%)	78% (61-90%)	1.4 (0.53-3.1)	3.5% (0.98-7.6%)	4.4% (1.1-9.6%)	200 (50-450)	1100 (290-2300)
Venezuela	2016	Female	95 plus	27% (11-48%)	73% (52-89%)	1.6 (0.5-3.8)	3.6% (0.52-8.9%)	4.3% (0.39-11%)	74 (6.5-190)	260 (39-640)
Venezuela	2016	Male	All Ages	42% (33-53%)	58% (47-67%)	3.4 (2.5-4.4)	10% (8-13%)	12% (9.3-15%)	11000 (8300-15000)	490000 (350000-660000)
Venezuela	2016	Male	15 to 19	25% (12-43%)	75% (57-88%)	1.2 (0.45-2.5)	6.5% (3-12%)	7.6% (3-15%)	280 (110-580)	23000 (10000-44000)
Venezuela	2016	Male	20 to 24	44% (23-66%)	56% (34-77%)	2.9 (1.2-5.4)	14% (7.1-23%)	15% (7.1-27%)	880 (400-1600)	65000 (32000-110000)
Venezuela	2016	Male	25 to 29	54% (32-74%)	46% (26-68%)	4.1 (1.8-7.4)	16% (9.5-25%)	19% (9.8-30%)	1000 (510-1700)	68000 (38000-110000)
Venezuela	2016	Male	30 to 34	61% (37-80%)	39% (20-63%)	5 (2.3-8.7)	17% (11-24%)	21% (12-30%)	850 (490-1300)	55000 (34000-82000)
Venezuela	2016	Male	35 to 39	64% (42-82%)	36% (18-58%)	5.6 (2.6-10)	16% (10-22%)	21% (13-30%)	720 (430-1100)	43000 (27000-63000)
Venezuela	2016	Male	40 to 44	65% (45-82%)	35% (18-55%)	6.2 (3-10)	15% (10-19%)	20% (13-27%)	650 (400-960)	35000 (23000-50000)
Venezuela	2016	Male	45 to 49	66% (41-82%)	34% (18-59%)	6.7 (3.1-12)	14% (9.8-18%)	19% (13-25%)	770 (490-1100)	37000 (24000-51000)
Venezuela	2016	Male	50 to 54	65% (42-82%)	35% (18-58%)	7 (3.6-13)	14% (9.5-18%)	18% (13-23%)	1000 (670-1500)	43000 (28000-59000)
Venezuela	2016	Male	55 to 59	63% (41-82%)	37% (18-59%)	7.3 (3.4-13)	13% (8.7-17%)	16% (11-21%)	1100 (720-1600)	39000 (25000-55000)
Venezuela	2016	Male	60 to 64	60% (37-80%)	40% (20-63%)	6.5 (2.8-12)	11% (7.2-15%)	13% (8.9-18%)	970 (610-1400)	29000 (18000-42000)
Venezuela	2016	Male	65 to 69	56% (33-77%)	44% (23-67%)	5 (2.2-9)	9.4% (6.1-13%)	12% (7.5-16%)	870 (530-1300)	21000 (14000-32000)
Venezuela	2016	Male	70 to 74	50% (30-71%)	50% (29-70%)	3.6 (1.5-7.6)	7.5% (4.5-11%)	9% (5.4-13%)	730 (410-1100)	14000 (8300-22000)
Venezuela	2016	Male	75 to 79	44% (22-66%)	56% (34-78%)	2.8 (0.93-6.2)	6% (2.8-9.6%)	7% (3.3-11%)	600 (280-1000)	9100 (4300-15000)
Venezuela	2016	Male	80 to 84	39% (18-61%)	61% (39-82%)	2.4 (0.78-4.9)	5.3% (2.4-8.9%)	6.1% (2.7-10%)	470 (200-810)	5200 (2300-8900)
Venezuela	2016	Male	85 to 89	37% (18-60%)	63% (40-82%)	2.3 (0.77-5.2)	4.9% (2.2-8.7%)	5.7% (2.5-10%)	310 (130-540)	2400 (1100-4200)
Venezuela	2016	Male	90 to 94	40% (19-62%)	60% (38-81%)	2.5 (0.77-5.6)	4.9% (2-8.4%)	5.5% (2.1-9.8%)	160 (57-280)	850 (340-1500)
Venezuela	2016	Male	95 plus	44% (20-70%)	56% (30-80%)	2.8 (0.76-7)	5.2% (1.5-9.6%)	5.8% (1.4-11%)	46 (11-90)	170 (48-320)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	1990	Female	All Ages	45% (37-52%)	55% (48-63%)	0.95 (0.83-1.1)	1.9% (1.5-2.4%)	2.2% (1.4-3.1%)	8200 (5000-12000)	430000 (330000-540000)
Brazil	1990	Female	15 to 19	47% (32-63%)	53% (37-68%)	0.94 (0.5-1.5)	2% (1.3-3.1%)	4% (2.1-6.6%)	180 (93-300)	20000 (13000-30000)
Brazil	1990	Female	20 to 24	57% (39-73%)	43% (27-61%)	1.5 (0.86-2.4)	3.8% (2.6-5.2%)	6.1% (3.5-9.2%)	340 (190-510)	40000 (27000-55000)
Brazil	1990	Female	25 to 29	59% (40-74%)	41% (26-60%)	1.7 (1-2.6)	4.4% (3.2-5.7%)	6.9% (4.3-9.8%)	450 (280-640)	48000 (34000-64000)
Brazil	1990	Female	30 to 34	59% (40-75%)	41% (25-60%)	1.7 (0.95-2.7)	4.5% (3.3-6%)	7.4% (4.8-11%)	560 (360-810)	49000 (35000-65000)
Brazil	1990	Female	35 to 39	58% (41-73%)	42% (27-59%)	1.7 (0.92-2.7)	4.6% (3.2-6.4%)	7.8% (4.9-11%)	680 (430-980)	48000 (33000-67000)
Brazil	1990	Female	40 to 44	55% (39-70%)	45% (30-61%)	1.7 (0.89-2.6)	4.8% (3-6.9%)	8% (4.6-12%)	830 (480-1200)	48000 (30000-68000)
Brazil	1990	Female	45 to 49	50% (31-65%)	50% (35-69%)	1.6 (0.79-2.7)	4.5% (2.6-6.9%)	7.1% (4-11%)	870 (490-1400)	43000 (25000-64000)
Brazil	1990	Female	50 to 54	45% (28-62%)	55% (38-72%)	1.5 (0.75-2.6)	4.3% (2.3-6.9%)	6.5% (3.4-11%)	1000 (540-1600)	42000 (24000-66000)
Brazil	1990	Female	55 to 59	42% (27-59%)	58% (41-73%)	1.4 (0.72-2.4)	3.9% (2-6.3%)	5.7% (2.9-9.4%)	1100 (540-1700)	37000 (20000-60000)
Brazil	1990	Female	60 to 64	41% (24-58%)	59% (42-76%)	1.2 (0.64-2.3)	2.9% (1.1-5.4%)	3.9% (1.3-7.5%)	930 (310-1800)	27000 (10000-51000)
Brazil	1990	Female	65 to 69	42% (26-57%)	58% (43-74%)	0.96 (0.5-1.6)	1.7% (0.37-3.5%)	2.2% (0.41-4.7%)	610 (110-1300)	15000 (3300-30000)
Brazil	1990	Female	70 to 74	39% (25-54%)	61% (46-75%)	0.67 (0.32-1.2)	0.63% (-0.48-1.9%)	0.74% (-0.73-2.4%)	240 (-230-760)	4800 (-3800-15000)
Brazil	1990	Female	75 to 79	33% (20-49%)	67% (51-80%)	0.52 (0.23-0.95)	0.19% (-0.83-1.3%)	0.16% (-1.1-1.5%)	58 (-400-550)	1200 (-5300-8500)
Brazil	1990	Female	80 to 84	27% (15-41%)	73% (59-85%)	0.5 (0.24-0.84)	0.34% (-0.47-1.3%)	0.34% (-0.64-1.5%)	110 (-220-500)	1500 (-2000-5600)
Brazil	1990	Female	85 to 89	24% (13-37%)	76% (63-87%)	0.51 (0.24-0.93)	0.52% (-0.28-1.6%)	0.53% (-0.43-1.8%)	110 (-94-390)	980 (-510-3100)
Brazil	1990	Female	90 to 94	25% (14-38%)	75% (62-86%)	0.6 (0.27-1.1)	0.68% (-0.18-2.1%)	0.7% (-0.36-2.4%)	69 (-35-240)	420 (-110-1300)
Brazil	1990	Female	95 plus	28% (14-46%)	72% (54-86%)	0.73 (0.29-1.5)	0.97% (-0.2-2.7%)	0.99% (-0.43-3.1%)	33 (-14-100)	130 (-27-370)
Brazil	1990	Male	All Ages	68% (63-73%)	32% (27-37%)	1.8 (1.6-2)	6.6% (5.6-7.7%)	7.5% (5.9-9.2%)	39000 (31000-48000)	1900000 (1600000-2300000)
Brazil	1990	Male	15 to 19	70% (59-81%)	30% (19-41%)	1.3 (0.75-2)	6.2% (3.7-9.3%)	8.5% (4.6-13%)	1100 (590-1700)	90000 (53000-130000)
Brazil	1990	Male	20 to 24	81% (72-88%)	19% (12-28%)	2.8 (1.8-3.8)	13% (9.2-17%)	16% (10-21%)	2800 (1900-3900)	220000 (150000-290000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	1990	Male	25 to 29	83% (73-89%)	17% (11-27%)	3.5 (2.4-4.7)	15% (12-19%)	18% (13-24%)	3500 (2500-4600)	270000 (200000-340000)
Brazil	1990	Male	30 to 34	83% (73-89%)	17% (11-27%)	3.6 (2.4-4.9)	16% (12-19%)	19% (14-24%)	3700 (2800-4700)	260000 (200000-320000)
Brazil	1990	Male	35 to 39	83% (75-89%)	17% (11-25%)	3.3 (2.2-4.6)	15% (12-19%)	18% (14-23%)	3700 (2800-4700)	230000 (180000-280000)
Brazil	1990	Male	40 to 44	82% (73-88%)	18% (12-27%)	3.2 (2-4.6)	14% (11-18%)	17% (13-22%)	3800 (2800-4800)	200000 (150000-250000)
Brazil	1990	Male	45 to 49	80% (69-87%)	20% (13-31%)	3.4 (2.1-5)	14% (10-18%)	17% (12-22%)	3900 (2700-5100)	180000 (130000-230000)
Brazil	1990	Male	50 to 54	78% (68-86%)	22% (14-32%)	3.3 (1.9-5)	12% (8.2-16%)	15% (9.6-20%)	4200 (2700-5600)	170000 (110000-220000)
Brazil	1990	Male	55 to 59	77% (67-85%)	23% (15-33%)	3.1 (1.9-4.7)	10% (6.9-14%)	12% (8-17%)	3900 (2500-5500)	130000 (87000-180000)
Brazil	1990	Male	60 to 64	76% (66-85%)	24% (15-34%)	2.8 (1.7-4.4)	7.9% (4.5-12%)	9.2% (5-14%)	3500 (1900-5300)	100000 (56000-150000)
Brazil	1990	Male	65 to 69	75% (65-84%)	25% (16-35%)	2.3 (1.4-3.5)	5.5% (2.8-8.9%)	6.3% (3.1-10%)	2500 (1200-4100)	59000 (30000-96000)
Brazil	1990	Male	70 to 74	71% (59-80%)	29% (20-41%)	1.7 (0.96-2.6)	3.3% (1.1-6.1%)	3.6% (1-6.9%)	1400 (400-2800)	28000 (9100-52000)
Brazil	1990	Male	75 to 79	64% (51-76%)	36% (24-49%)	1.2 (0.64-1.9)	1.8% (0.23-3.8%)	1.9% (0.097-4.2%)	750 (38-1600)	12000 (1500-24000)
Brazil	1990	Male	80 to 84	55% (42-67%)	45% (33-58%)	0.93 (0.54-1.4)	1.2% (0.037-2.5%)	1.2% (-0.12-2.7%)	360 (-39-830)	4300 (130-9200)
Brazil	1990	Male	85 to 89	47% (35-59%)	53% (41-65%)	0.8 (0.46-1.3)	0.75% (-0.14-1.8%)	0.67% (-0.33-1.9%)	92 (-45-260)	870 (-160-2200)
Brazil	1990	Male	90 to 94	44% (32-56%)	56% (44-68%)	0.73 (0.4-1.2)	0.48% (-0.29-1.4%)	0.32% (-0.58-1.4%)	16 (-29-70)	140 (-87-420)
Brazil	1990	Male	95 plus	43% (29-58%)	57% (42-71%)	0.68 (0.32-1.2)	0.48% (-0.35-1.5%)	0.19% (-0.86-1.3%)	2.5 (-11-18)	25 (-19-79)
Brazil	1995	Female	All Ages	45% (38-53%)	55% (47-62%)	1.2 (1-1.3)	2.5% (2-3.1%)	3.1% (2.1-4.3%)	12000 (8000-17000)	560000 (440000-690000)
Brazil	1995	Female	15 to 19	46% (31-63%)	54% (37-69%)	1.1 (0.58-1.8)	2.3% (1.4-3.5%)	4.7% (2.5-7.7%)	230 (120-380)	25000 (15000-37000)
Brazil	1995	Female	20 to 24	56% (38-72%)	44% (28-62%)	1.8 (0.98-2.9)	4.1% (2.8-5.6%)	6.9% (4-10%)	390 (230-570)	44000 (30000-61000)
Brazil	1995	Female	25 to 29	58% (39-74%)	42% (26-61%)	2 (1.2-3.1)	4.6% (3.4-6%)	7.5% (4.7-10%)	500 (320-700)	52000 (36000-69000)
Brazil	1995	Female	30 to 34	58% (39-74%)	42% (26-61%)	2 (1.1-3.1)	4.8% (3.5-6.2%)	8.1% (5.3-11%)	680 (450-930)	59000 (42000-77000)
Brazil	1995	Female	35 to 39	58% (40-73%)	42% (27-60%)	2 (1.1-3.2)	5% (3.6-6.8%)	8.8% (5.8-12%)	860 (570-1200)	60000 (42000-82000)
Brazil	1995	Female	40 to 44	55% (38-70%)	45% (30-62%)	2 (1-3)	5.3% (3.5-7.4%)	9.1% (5.6-13%)	1100 (670-1600)	63000 (42000-88000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	1995	Female	45 to 49	50% (31-65%)	50% (35-69%)	1.9 (0.95-3.2)	5.1% (3.1-7.7%)	8.4% (4.8-13%)	1200 (690-1900)	58000 (35000-87000)
Brazil	1995	Female	50 to 54	46% (28-63%)	54% (37-72%)	1.9 (0.94-3.1)	5.1% (2.9-8%)	7.9% (4.5-12%)	1300 (760-2000)	54000 (32000-82000)
Brazil	1995	Female	55 to 59	44% (29-61%)	56% (39-71%)	1.8 (0.95-3.1)	4.9% (2.7-7.6%)	7.2% (4-11%)	1500 (840-2400)	54000 (30000-84000)
Brazil	1995	Female	60 to 64	45% (29-61%)	55% (39-71%)	1.7 (0.89-2.9)	3.9% (1.7-6.8%)	5.4% (2.3-9.4%)	1400 (610-2400)	41000 (18000-72000)
Brazil	1995	Female	65 to 69	45% (31-60%)	55% (40-69%)	1.3 (0.72-2.1)	2.6% (0.92-4.7%)	3.4% (1.2-6.3%)	1100 (380-2100)	27000 (9600-49000)
Brazil	1995	Female	70 to 74	43% (29-57%)	57% (43-71%)	0.9 (0.46-1.5)	1.2% (-0.21-2.9%)	1.5% (-0.3-3.7%)	530 (-110-1300)	10000 (-1800-25000)
Brazil	1995	Female	75 to 79	36% (22-52%)	64% (48-78%)	0.69 (0.32-1.2)	0.61% (-0.58-2.1%)	0.71% (-0.77-2.5%)	280 (-300-970)	4300 (-4100-14000)
Brazil	1995	Female	80 to 84	29% (16-44%)	71% (56-84%)	0.65 (0.32-1.1)	0.82% (-0.18-2%)	0.92% (-0.28-2.4%)	350 (-110-910)	4000 (-880-9900)
Brazil	1995	Female	85 to 89	26% (15-40%)	74% (60-85%)	0.67 (0.31-1.2)	1% (0.069-2.5%)	1.2% (-0.0038-2.9%)	320 (-1.1-820)	2600 (170-6200)
Brazil	1995	Female	90 to 94	27% (15-40%)	73% (60-85%)	0.78 (0.36-1.4)	1.3% (0.16-3.2%)	1.4% (0.069-3.7%)	180 (9.2-470)	1000 (120-2400)
Brazil	1995	Female	95 plus	30% (15-48%)	70% (52-85%)	0.96 (0.38-1.9)	1.7% (0.13-4%)	1.9% (0.0018-4.6%)	75 (0.07-190)	270 (21-650)
Brazil	1995	Male	All Ages	69% (64-74%)	31% (26-36%)	2.2 (2-2.4)	8.2% (7.1-9.4%)	9.2% (7.4-11%)	50000 (40000-61000)	2400000 (2100000-2800000)
Brazil	1995	Male	15 to 19	70% (58-81%)	30% (19-42%)	1.5 (0.85-2.3)	6.8% (4.2-10%)	9.4% (5.2-15%)	1400 (750-2100)	110000 (66000-160000)
Brazil	1995	Male	20 to 24	81% (71-88%)	19% (12-29%)	3.2 (2.1-4.4)	14% (10-18%)	17% (12-24%)	3400 (2200-4600)	260000 (180000-340000)
Brazil	1995	Male	25 to 29	82% (72-89%)	18% (11-28%)	4 (2.7-5.5)	16% (12-20%)	19% (14-25%)	4000 (2900-5200)	300000 (220000-380000)
Brazil	1995	Male	30 to 34	82% (73-89%)	18% (11-27%)	4.2 (2.8-5.7)	16% (13-20%)	19% (15-24%)	4300 (3300-5500)	300000 (240000-370000)
Brazil	1995	Male	35 to 39	83% (74-89%)	17% (11-26%)	3.8 (2.6-5.4)	16% (13-19%)	19% (15-24%)	4600 (3500-5700)	280000 (220000-340000)
Brazil	1995	Male	40 to 44	81% (73-88%)	19% (12-27%)	3.8 (2.4-5.3)	16% (12-19%)	19% (14-24%)	4900 (3700-6100)	260000 (200000-320000)
Brazil	1995	Male	45 to 49	79% (69-87%)	21% (13-31%)	3.9 (2.4-5.9)	15% (11-19%)	19% (14-24%)	5100 (3700-6400)	230000 (170000-290000)
Brazil	1995	Male	50 to 54	78% (68-86%)	22% (14-32%)	3.8 (2.3-5.6)	14% (9.8-17%)	17% (12-22%)	5000 (3400-6400)	200000 (140000-250000)
Brazil	1995	Male	55 to 59	77% (68-86%)	23% (14-32%)	3.7 (2.2-5.5)	12% (8.4-16%)	15% (10-19%)	5100 (3500-6800)	170000 (120000-230000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	1995	Male	60 to 64	78% (69-86%)	22% (14-31%)	3.5 (2.1-5.3)	9.7% (5.9-14%)	11% (6.8-16%)	4500 (2700-6400)	130000 (78000-180000)
Brazil	1995	Male	65 to 69	77% (68-85%)	23% (15-32%)	2.9 (1.9-4.3)	7.1% (4-11%)	8.1% (4.5-12%)	3800 (2100-5700)	89000 (50000-130000)
Brazil	1995	Male	70 to 74	73% (63-81%)	27% (19-37%)	2.1 (1.2-3.2)	4.4% (1.8-7.5%)	4.9% (2-8.7%)	2300 (900-3900)	43000 (18000-74000)
Brazil	1995	Male	75 to 79	66% (53-78%)	34% (22-47%)	1.5 (0.81-2.4)	2.6% (0.68-5%)	2.8% (0.61-5.6%)	1200 (250-2400)	18000 (4800-35000)
Brazil	1995	Male	80 to 84	57% (44-69%)	43% (31-56%)	1.2 (0.68-1.8)	1.8% (0.4-3.5%)	1.9% (0.32-3.7%)	670 (110-1300)	7500 (1700-15000)
Brazil	1995	Male	85 to 89	49% (37-62%)	51% (38-63%)	1 (0.57-1.6)	1.3% (0.21-2.7%)	1.3% (0.056-2.9%)	240 (11-540)	2000 (330-4400)
Brazil	1995	Male	90 to 94	47% (35-59%)	53% (41-65%)	0.92 (0.51-1.5)	0.9% (-0.0071-2.1%)	0.81% (-0.25-2.2%)	49 (-15-140)	320 (-2.6-770)
Brazil	1995	Male	95 plus	47% (32-62%)	53% (38-68%)	0.88 (0.42-1.6)	0.81% (-0.19-2.4%)	0.59% (-0.57-2.4%)	8.8 (-8.7-36)	48 (-11-140)
Brazil	2000	Female	All Ages	47% (40-54%)	53% (46-60%)	1.5 (1.3-1.7)	3.1% (2.5-3.8%)	4% (2.8-5.4%)	16000 (11000-22000)	690000 (550000-840000)
Brazil	2000	Female	15 to 19	48% (33-64%)	52% (36-67%)	1.4 (0.74-2.2)	2.5% (1.6-3.7%)	5.8% (3.1-9.1%)	270 (150-430)	28000 (18000-42000)
Brazil	2000	Female	20 to 24	58% (41-73%)	42% (27-59%)	2.2 (1.2-3.4)	4.3% (3.1-5.9%)	8.3% (5-12%)	450 (270-660)	51000 (35000-69000)
Brazil	2000	Female	25 to 29	59% (40-74%)	41% (26-60%)	2.4 (1.4-3.7)	4.7% (3.5-6.2%)	8.5% (5.6-12%)	520 (340-720)	54000 (39000-71000)
Brazil	2000	Female	30 to 34	58% (39-74%)	42% (26-61%)	2.4 (1.3-3.6)	4.9% (3.7-6.3%)	9.1% (6.1-12%)	700 (470-950)	61000 (45000-78000)
Brazil	2000	Female	35 to 39	57% (41-72%)	43% (28-59%)	2.3 (1.3-3.7)	5.2% (3.8-6.9%)	9.7% (6.6-13%)	980 (660-1300)	70000 (49000-91000)
Brazil	2000	Female	40 to 44	54% (38-70%)	46% (30-62%)	2.3 (1.2-3.5)	5.5% (3.7-7.7%)	10% (6.4-14%)	1300 (830-1800)	75000 (52000-100000)
Brazil	2000	Female	45 to 49	49% (31-65%)	51% (35-69%)	2.2 (1.1-3.7)	5.5% (3.5-8%)	9.5% (5.7-14%)	1500 (910-2300)	74000 (47000-110000)
Brazil	2000	Female	50 to 54	47% (30-64%)	53% (36-70%)	2.3 (1.2-3.8)	5.7% (3.5-8.6%)	9.2% (5.5-14%)	1800 (1100-2600)	72000 (45000-110000)
Brazil	2000	Female	55 to 59	46% (31-63%)	54% (37-69%)	2.3 (1.2-3.8)	5.7% (3.2-8.4%)	8.6% (4.9-13%)	1900 (1100-2800)	66000 (38000-98000)
Brazil	2000	Female	60 to 64	47% (31-63%)	53% (37-69%)	2.1 (1.1-3.6)	4.7% (2.3-7.7%)	6.7% (3.2-11%)	1900 (920-3200)	57000 (26000-93000)
Brazil	2000	Female	65 to 69	48% (32-63%)	52% (37-68%)	1.6 (0.91-2.6)	3.4% (1.5-5.8%)	4.6% (2-7.9%)	1600 (690-2700)	38000 (17000-64000)
Brazil	2000	Female	70 to 74	47% (33-61%)	53% (39-67%)	1.2 (0.61-2)	1.8% (0.2-3.8%)	2.4% (0.22-5%)	1000 (92-2100)	19000 (2000-42000)
Brazil	2000	Female	75 to 79	41% (26-57%)	59% (43-74%)	0.96 (0.46-1.7)	1.1% (-0.32-3%)	1.4% (-0.46-3.7%)	600 (-200-1600)	9200 (-2600-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	2000	Female	80 to 84	33% (19-49%)	67% (51-81%)	0.9 (0.43-1.5)	1.4% (0.16-2.9%)	1.6% (0.13-3.6%)	700 (53-1500)	7800 (890-16000)
Brazil	2000	Female	85 to 89	29% (16-43%)	71% (57-84%)	0.89 (0.41-1.6)	1.6% (0.43-3.5%)	1.9% (0.43-4.2%)	660 (150-1500)	5200 (1400-11000)
Brazil	2000	Female	90 to 94	29% (17-44%)	71% (56-83%)	1 (0.47-1.9)	2% (0.53-4.4%)	2.3% (0.53-5.1%)	450 (99-1000)	2400 (610-5200)
Brazil	2000	Female	95 plus	32% (16-52%)	68% (48-84%)	1.2 (0.51-2.5)	2.6% (0.56-5.4%)	2.9% (0.49-6.2%)	160 (28-360)	580 (120-1200)
Brazil	2000	Male	All Ages	71% (66-75%)	29% (25-34%)	2.7 (2.4-3)	9.9% (8.5-11%)	11% (8.7-12%)	61000 (50000-72000)	2900000 (2500000-3300000)
Brazil	2000	Male	15 to 19	73% (63-83%)	27% (17-37%)	1.9 (1.1-2.9)	8.2% (4.9-12%)	11% (6.4-17%)	1800 (1000-2700)	150000 (87000-210000)
Brazil	2000	Male	20 to 24	82% (74-89%)	18% (11-26%)	3.9 (2.6-5.2)	16% (12-21%)	20% (14-27%)	4400 (2900-5900)	340000 (240000-440000)
Brazil	2000	Male	25 to 29	82% (73-89%)	18% (11-27%)	4.6 (3.2-6.4)	18% (14-23%)	22% (16-29%)	4500 (3200-5800)	340000 (260000-420000)
Brazil	2000	Male	30 to 34	82% (72-89%)	18% (11-28%)	4.7 (3.2-6.6)	18% (14-22%)	22% (17-27%)	4600 (3600-5900)	320000 (260000-400000)
Brazil	2000	Male	35 to 39	82% (74-89%)	18% (11-26%)	4.3 (2.9-6.2)	17% (14-21%)	21% (16-27%)	5100 (3900-6400)	320000 (260000-390000)
Brazil	2000	Male	40 to 44	82% (74-89%)	18% (11-26%)	4.4 (2.8-6.1)	17% (13-20%)	21% (17-26%)	5800 (4400-7000)	310000 (250000-370000)
Brazil	2000	Male	45 to 49	80% (69-87%)	20% (13-31%)	4.5 (2.9-6.6)	16% (13-20%)	21% (16-25%)	6300 (4800-7700)	290000 (230000-350000)
Brazil	2000	Male	50 to 54	78% (67-85%)	22% (15-33%)	4.3 (2.6-6.4)	15% (11-18%)	19% (14-23%)	6200 (4500-7800)	250000 (180000-310000)
Brazil	2000	Male	55 to 59	77% (67-86%)	23% (14-33%)	4.2 (2.5-6.3)	13% (9.6-17%)	16% (12-21%)	5900 (4200-7500)	200000 (150000-250000)
Brazil	2000	Male	60 to 64	76% (66-85%)	24% (15-34%)	3.8 (2.3-5.9)	10% (6.8-14%)	13% (8-17%)	5400 (3500-7400)	160000 (100000-210000)
Brazil	2000	Male	65 to 69	75% (64-83%)	25% (17-36%)	3.1 (2-4.5)	7.8% (4.8-11%)	9.2% (5.6-13%)	4400 (2600-6200)	100000 (65000-150000)
Brazil	2000	Male	70 to 74	70% (59-79%)	30% (21-41%)	2.2 (1.3-3.5)	5.1% (2.5-8.2%)	5.8% (2.8-9.5%)	3100 (1500-5000)	60000 (30000-95000)
Brazil	2000	Male	75 to 79	64% (50-77%)	36% (23-50%)	1.6 (0.86-2.6)	3.2% (1.3-5.6%)	3.6% (1.3-6.4%)	1700 (630-3100)	26000 (10000-46000)
Brazil	2000	Male	80 to 84	55% (41-68%)	45% (32-59%)	1.3 (0.73-2)	2.3% (0.87-4.1%)	2.5% (0.86-4.5%)	980 (340-1800)	11000 (4200-20000)
Brazil	2000	Male	85 to 89	47% (34-61%)	53% (39-66%)	1.1 (0.59-1.8)	1.7% (0.56-3.4%)	1.8% (0.48-3.7%)	490 (130-1000)	3900 (1300-7900)
Brazil	2000	Male	90 to 94	46% (33-59%)	54% (41-67%)	1 (0.56-1.8)	1.3% (0.21-2.9%)	1.3% (0.057-3%)	150 (6.4-340)	870 (140-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	2000	Male	95 plus	47% (31-63%)	53% (37-69%)	1 (0.46-1.8)	1.1% (-0.058-3.1%)	1% (-0.31-3.2%)	25 (-7.7-78)	110 (-5.4-300)
Brazil	2005	Female	All Ages	44% (40-49%)	56% (51-60%)	1.5 (1.4-1.7)	3.1% (2.6-3.7%)	3.8% (2.8-5%)	17000 (12000-22000)	710000 (580000-860000)
Brazil	2005	Female	15 to 19	51% (40-63%)	49% (37-60%)	1.6 (1-2.5)	2.8% (1.9-3.8%)	6.7% (4.1-10%)	290 (180-440)	30000 (20000-42000)
Brazil	2005	Female	20 to 24	58% (47-69%)	42% (31-53%)	2.5 (1.6-3.5)	4.5% (3.4-5.9%)	9.1% (5.9-13%)	480 (310-670)	56000 (41000-74000)
Brazil	2005	Female	25 to 29	59% (48-69%)	41% (31-52%)	2.6 (1.8-3.6)	4.8% (3.8-6%)	9.1% (6.5-12%)	560 (410-750)	60000 (46000-77000)
Brazil	2005	Female	30 to 34	55% (44-66%)	45% (34-56%)	2.4 (1.6-3.3)	4.7% (3.7-5.8%)	8.9% (6.5-11%)	650 (470-820)	59000 (45000-74000)
Brazil	2005	Female	35 to 39	53% (43-64%)	47% (36-57%)	2.2 (1.4-3.2)	4.8% (3.6-6.1%)	9.2% (6.5-12%)	870 (620-1100)	65000 (48000-82000)
Brazil	2005	Female	40 to 44	51% (41-61%)	49% (39-59%)	2.2 (1.4-3.2)	5.2% (3.7-6.9%)	9.8% (6.6-13%)	1300 (880-1800)	78000 (56000-100000)
Brazil	2005	Female	45 to 49	46% (36-58%)	54% (42-64%)	2.2 (1.4-3.4)	5.4% (3.7-7.4%)	9.5% (6.4-13%)	1700 (1100-2300)	82000 (58000-110000)
Brazil	2005	Female	50 to 54	40% (30-51%)	60% (49-70%)	2 (1.2-3.1)	5.2% (3.5-7.2%)	8.5% (5.7-12%)	1800 (1200-2500)	76000 (52000-100000)
Brazil	2005	Female	55 to 59	36% (28-47%)	64% (53-72%)	1.8 (1.1-2.8)	4.8% (3-6.7%)	7.5% (4.9-11%)	1900 (1200-2600)	66000 (43000-93000)
Brazil	2005	Female	60 to 64	35% (24-47%)	65% (53-76%)	1.6 (0.89-2.6)	3.9% (2.2-6%)	5.7% (3.2-8.7%)	1700 (930-2500)	49000 (28000-76000)
Brazil	2005	Female	65 to 69	35% (25-47%)	65% (53-75%)	1.2 (0.7-1.9)	2.9% (1.6-4.5%)	4% (2.2-6.3%)	1500 (830-2400)	37000 (21000-58000)
Brazil	2005	Female	70 to 74	37% (26-49%)	63% (51-74%)	0.96 (0.52-1.6)	1.9% (0.66-3.5%)	2.5% (0.87-4.7%)	1100 (380-2000)	21000 (7600-40000)
Brazil	2005	Female	75 to 79	33% (21-46%)	67% (54-79%)	0.8 (0.42-1.3)	1.3% (0.17-2.7%)	1.7% (0.18-3.5%)	860 (96-1800)	13000 (1700-27000)
Brazil	2005	Female	80 to 84	25% (15-37%)	75% (63-85%)	0.7 (0.37-1.1)	1.4% (0.5-2.6%)	1.7% (0.59-3.1%)	840 (300-1600)	9400 (3500-17000)
Brazil	2005	Female	85 to 89	21% (12-30%)	79% (70-88%)	0.67 (0.34-1.2)	1.5% (0.66-2.9%)	1.8% (0.74-3.4%)	760 (310-1400)	6000 (2500-11000)
Brazil	2005	Female	90 to 94	21% (13-31%)	79% (69-87%)	0.76 (0.37-1.4)	1.8% (0.71-3.5%)	2.1% (0.75-4.2%)	540 (190-1100)	2900 (1100-5600)
Brazil	2005	Female	95 plus	23% (12-37%)	77% (63-88%)	0.92 (0.39-1.7)	2.3% (0.73-4.5%)	2.6% (0.74-5.2%)	220 (64-440)	790 (260-1500)
Brazil	2005	Male	All Ages	72% (68-75%)	28% (25-32%)	3 (2.7-3.3)	11% (9.3-12%)	11% (9.4-13%)	68000 (57000-79000)	320000 (280000-360000)
Brazil	2005	Male	15 to 19	76% (68-84%)	24% (16-32%)	2.2 (1.3-3.3)	8.9% (5.4-13%)	12% (7.1-18%)	2000 (1100-2900)	160000 (98000-220000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	2005	Male	20 to 24	83% (76-88%)	17% (12-24%)	4.2 (2.8-5.7)	17% (13-22%)	21% (15-28%)	4800 (3300-6400)	370000 (270000-480000)
Brazil	2005	Male	25 to 29	83% (76-89%)	17% (11-24%)	5 (3.5-6.7)	19% (14-23%)	23% (17-30%)	5000 (3700-6500)	370000 (290000-470000)
Brazil	2005	Male	30 to 34	83% (76-88%)	17% (12-24%)	5.1 (3.6-6.9)	18% (15-22%)	23% (18-29%)	4700 (3700-5900)	330000 (270000-400000)
Brazil	2005	Male	35 to 39	82% (76-88%)	18% (12-24%)	4.5 (3.1-6.3)	17% (14-21%)	22% (17-27%)	5100 (3900-6200)	320000 (250000-380000)
Brazil	2005	Male	40 to 44	81% (75-87%)	19% (13-25%)	4.5 (2.9-6.2)	17% (13-20%)	22% (17-27%)	6100 (4700-7300)	330000 (270000-390000)
Brazil	2005	Male	45 to 49	80% (72-86%)	20% (14-28%)	4.8 (3.1-6.9)	17% (13-20%)	22% (17-26%)	7100 (5600-8500)	330000 (270000-400000)
Brazil	2005	Male	50 to 54	78% (70-84%)	22% (16-30%)	4.6 (2.9-6.8)	15% (12-19%)	20% (15-24%)	7400 (5700-9000)	300000 (230000-360000)
Brazil	2005	Male	55 to 59	76% (68-83%)	24% (17-32%)	4.2 (2.8-6.2)	13% (10-17%)	17% (13-21%)	6900 (5200-8500)	240000 (180000-290000)
Brazil	2005	Male	60 to 64	75% (67-83%)	25% (17-33%)	3.9 (2.4-6)	11% (7.3-14%)	13% (8.9-17%)	5900 (4000-7700)	170000 (120000-220000)
Brazil	2005	Male	65 to 69	74% (65-82%)	26% (18-35%)	3.2 (2-4.7)	8.3% (5.3-11%)	9.9% (6.3-14%)	5200 (3300-7100)	130000 (81000-170000)
Brazil	2005	Male	70 to 74	70% (59-79%)	30% (21-41%)	2.3 (1.4-3.6)	5.4% (2.9-8.3%)	6.3% (3.3-9.9%)	3500 (1800-5500)	68000 (36000-110000)
Brazil	2005	Male	75 to 79	63% (50-75%)	37% (25-50%)	1.6 (0.91-2.6)	3.4% (1.5-5.6%)	3.8% (1.6-6.5%)	2200 (890-3800)	33000 (14000-57000)
Brazil	2005	Male	80 to 84	53% (40-66%)	47% (34-60%)	1.3 (0.72-2)	2.4% (1-4%)	2.6% (1.1-4.6%)	1200 (510-2200)	14000 (6200-24000)
Brazil	2005	Male	85 to 89	45% (33-57%)	55% (43-67%)	1.1 (0.61-1.7)	1.9% (0.77-3.4%)	2% (0.71-3.7%)	630 (230-1200)	5100 (2100-9300)
Brazil	2005	Male	90 to 94	43% (31-55%)	57% (45-69%)	1 (0.55-1.6)	1.5% (0.47-3%)	1.5% (0.34-3.1%)	230 (52-490)	1300 (410-2600)
Brazil	2005	Male	95 plus	44% (29-59%)	56% (41-71%)	0.97 (0.47-1.8)	1.3% (0.2-3.2%)	1.2% (-0.017-3.3%)	44 (-0.59-120)	180 (27-450)
Brazil	2010	Female	All Ages	43% (39-47%)	57% (53-61%)	1.4 (1.3-1.6)	3% (2.5-3.6%)	3.5% (2.6-4.6%)	17000 (12000-22000)	720000 (590000-860000)
Brazil	2010	Female	15 to 19	50% (39-62%)	50% (38-61%)	1.5 (0.95-2.3)	2.7% (1.9-3.8%)	6.3% (3.8-9.5%)	260 (160-400)	28000 (19000-40000)
Brazil	2010	Female	20 to 24	58% (47-68%)	42% (32-53%)	2.3 (1.4-3.4)	4.6% (3.4-5.9%)	8.7% (5.5-12%)	450 (290-640)	53000 (38000-70000)
Brazil	2010	Female	25 to 29	58% (48-68%)	42% (32-52%)	2.4 (1.6-3.4)	4.8% (3.7-5.9%)	8.7% (6.2-12%)	560 (400-760)	62000 (46000-79000)
Brazil	2010	Female	30 to 34	54% (44-64%)	46% (36-56%)	2.2 (1.5-3)	4.5% (3.5-5.6%)	8.2% (6.1-11%)	650 (480-840)	61000 (46000-76000)
Brazil	2010	Female	35 to 39	52% (42-63%)	48% (37-58%)	2.1 (1.3-3)	4.5% (3.4-5.7%)	8.5% (6-11%)	810 (570-1100)	62000 (46000-78000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	2010	Female	40 to 44	50% (40-60%)	50% (40-60%)	2.1 (1.3-2.9)	4.9% (3.5-6.5%)	9.1% (6.3-12%)	1200 (830-1600)	74000 (53000-96000)
Brazil	2010	Female	45 to 49	45% (35-56%)	55% (44-65%)	2.1 (1.3-3.1)	5% (3.5-6.8%)	8.8% (6.1-12%)	1600 (1100-2300)	83000 (59000-110000)
Brazil	2010	Female	50 to 54	40% (30-49%)	60% (51-70%)	1.9 (1.2-2.8)	4.8% (3.2-6.7%)	8% (5.2-11%)	1900 (1200-2600)	79000 (54000-110000)
Brazil	2010	Female	55 to 59	35% (26-44%)	65% (56-74%)	1.7 (1.1-2.6)	4.4% (2.9-6.2%)	6.9% (4.6-9.8%)	2000 (1300-2800)	71000 (48000-100000)
Brazil	2010	Female	60 to 64	32% (21-44%)	68% (56-79%)	1.3 (0.77-2.2)	3.5% (2-5.2%)	5.1% (2.9-7.8%)	1700 (970-2600)	51000 (30000-78000)
Brazil	2010	Female	65 to 69	31% (22-43%)	69% (57-78%)	1 (0.6-1.6)	2.5% (1.5-3.9%)	3.5% (2.1-5.5%)	1400 (820-2100)	34000 (20000-54000)
Brazil	2010	Female	70 to 74	31% (21-43%)	69% (57-79%)	0.76 (0.4-1.3)	1.6% (0.65-2.9%)	2.2% (0.88-3.8%)	1100 (430-1900)	21000 (8500-37000)
Brazil	2010	Female	75 to 79	28% (18-40%)	72% (60-82%)	0.63 (0.32-1.1)	1.1% (0.26-2.3%)	1.4% (0.29-2.9%)	800 (160-1600)	12000 (2900-25000)
Brazil	2010	Female	80 to 84	24% (14-35%)	76% (65-86%)	0.63 (0.34-1)	1.2% (0.45-2.3%)	1.5% (0.52-2.9%)	930 (320-1800)	10000 (3800-20000)
Brazil	2010	Female	85 to 89	21% (13-31%)	79% (69-87%)	0.66 (0.34-1.1)	1.4% (0.58-2.6%)	1.7% (0.65-3.2%)	860 (340-1700)	7000 (3000-13000)
Brazil	2010	Female	90 to 94	22% (13-32%)	78% (68-87%)	0.78 (0.38-1.3)	1.7% (0.66-3.3%)	2% (0.68-3.9%)	660 (230-1300)	3600 (1400-6800)
Brazil	2010	Female	95 plus	25% (13-38%)	75% (62-87%)	0.96 (0.41-1.8)	2.2% (0.68-4.2%)	2.5% (0.68-5%)	280 (77-570)	1000 (320-2000)
Brazil	2010	Male	All Ages	72% (68-75%)	28% (25-32%)	2.9 (2.6-3.2)	11% (9.2-12%)	11% (9-13%)	70000 (59000-82000)	330000 (280000-370000)
Brazil	2010	Male	15 to 19	75% (66-83%)	25% (17-34%)	2 (1.2-3.1)	8.5% (5.1-12%)	12% (6.5-17%)	1900 (1000-2800)	150000 (90000-210000)
Brazil	2010	Male	20 to 24	83% (76-88%)	17% (12-24%)	4 (2.6-5.6)	16% (12-21%)	20% (14-27%)	4600 (3100-6300)	350000 (250000-470000)
Brazil	2010	Male	25 to 29	83% (76-88%)	17% (12-24%)	4.8 (3.3-6.5)	18% (14-22%)	23% (16-29%)	5100 (3700-6700)	380000 (290000-480000)
Brazil	2010	Male	30 to 34	82% (75-88%)	18% (12-25%)	4.8 (3.4-6.5)	17% (14-21%)	22% (17-28%)	4900 (3800-6300)	350000 (280000-420000)
Brazil	2010	Male	35 to 39	82% (76-88%)	18% (12-24%)	4.4 (2.9-6)	17% (13-20%)	22% (17-27%)	4900 (3800-6000)	310000 (240000-370000)
Brazil	2010	Male	40 to 44	81% (74-87%)	19% (13-26%)	4.3 (2.8-6.1)	16% (13-19%)	21% (16-26%)	5800 (4500-7100)	320000 (260000-380000)
Brazil	2010	Male	45 to 49	79% (71-86%)	21% (14-29%)	4.5 (3-6.5)	16% (13-19%)	21% (17-25%)	7300 (5800-8700)	350000 (280000-410000)
Brazil	2010	Male	50 to 54	78% (70-84%)	22% (16-30%)	4.4 (2.7-6.3)	15% (12-18%)	19% (15-23%)	7900 (6000-9600)	320000 (250000-390000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	2010	Male	55 to 59	75% (67-83%)	25% (17-33%)	4 (2.6-5.9)	13% (9.8-16%)	16% (12-20%)	7700 (5800-9600)	270000 (200000-330000)
Brazil	2010	Male	60 to 64	75% (65-83%)	25% (17-35%)	3.7 (2.3-5.7)	10% (7-14%)	13% (8.5-17%)	6600 (4400-8700)	190000 (130000-260000)
Brazil	2010	Male	65 to 69	74% (65-82%)	26% (18-35%)	3.1 (2-4.5)	7.9% (5.1-11%)	9.5% (6-13%)	5300 (3300-7300)	130000 (83000-180000)
Brazil	2010	Male	70 to 74	69% (58-79%)	31% (21-42%)	2.2 (1.3-3.4)	5.1% (2.7-7.8%)	6% (3.1-9.4%)	3700 (2000-5800)	74000 (40000-110000)
Brazil	2010	Male	75 to 79	62% (50-75%)	38% (25-50%)	1.6 (0.86-2.5)	3.1% (1.4-5.3%)	3.6% (1.5-6.2%)	2200 (920-3800)	34000 (15000-58000)
Brazil	2010	Male	80 to 84	53% (40-65%)	47% (35-60%)	1.2 (0.72-1.9)	2.2% (1-3.8%)	2.5% (1-4.3%)	1400 (580-2400)	16000 (7200-27000)
Brazil	2010	Male	85 to 89	45% (33-57%)	55% (43-67%)	1 (0.6-1.7)	1.7% (0.74-3.2%)	1.9% (0.7-3.5%)	740 (280-1400)	6200 (2700-11000)
Brazil	2010	Male	90 to 94	43% (30-55%)	57% (45-70%)	0.97 (0.53-1.6)	1.4% (0.47-2.8%)	1.4% (0.35-3%)	290 (69-610)	1700 (550-3400)
Brazil	2010	Male	95 plus	43% (28-58%)	57% (42-72%)	0.92 (0.42-1.7)	1.3% (0.27-3.1%)	1.2% (0.039-3.2%)	55 (1.8-140)	230 (48-540)
Brazil	2016	Female	All Ages	42% (38-47%)	58% (53-62%)	1.5 (1.2-1.9)	3.2% (2.5-4%)	3.7% (2.5-5.2%)	21000 (14000-29000)	800000 (620000-1000000)
Brazil	2016	Female	15 to 19	49% (37-61%)	51% (39-63%)	1.6 (0.91-2.5)	2.8% (1.8-3.9%)	6.5% (3.7-10%)	260 (150-400)	29000 (19000-41000)
Brazil	2016	Female	20 to 24	57% (45-68%)	43% (32-55%)	2.4 (1.4-3.7)	4.8% (3.5-6.3%)	8.9% (5.5-13%)	400 (250-590)	51000 (36000-69000)
Brazil	2016	Female	25 to 29	56% (45-67%)	44% (33-55%)	2.5 (1.6-3.7)	5% (3.8-6.2%)	8.8% (6.1-12%)	490 (330-680)	59000 (43000-76000)
Brazil	2016	Female	30 to 34	53% (41-65%)	47% (35-59%)	2.3 (1.4-3.3)	4.7% (3.6-5.9%)	8.5% (6-11%)	660 (470-880)	66000 (50000-85000)
Brazil	2016	Female	35 to 39	52% (41-63%)	48% (37-59%)	2.2 (1.3-3.4)	4.7% (3.4-6%)	8.7% (6.1-12%)	880 (610-1200)	70000 (52000-90000)
Brazil	2016	Female	40 to 44	48% (38-59%)	52% (41-62%)	2.1 (1.3-3.2)	4.9% (3.5-6.5%)	9.1% (6.3-12%)	1200 (820-1600)	76000 (55000-99000)
Brazil	2016	Female	45 to 49	43% (32-56%)	57% (44-68%)	2.1 (1.3-3.3)	4.9% (3.5-6.7%)	8.8% (6-12%)	1600 (1100-2200)	82000 (58000-110000)
Brazil	2016	Female	50 to 54	40% (29-51%)	60% (49-71%)	2 (1.2-3)	4.9% (3.4-6.7%)	8.2% (5.5-11%)	2100 (1400-2900)	90000 (62000-120000)
Brazil	2016	Female	55 to 59	36% (27-47%)	64% (53-73%)	1.9 (1.1-2.9)	4.6% (3-6.3%)	7.3% (4.7-10%)	2400 (1500-3300)	86000 (56000-120000)
Brazil	2016	Female	60 to 64	34% (22-48%)	66% (52-78%)	1.6 (0.85-2.7)	3.8% (2.2-5.8%)	5.6% (3.2-8.8%)	2200 (1300-3500)	68000 (39000-100000)
Brazil	2016	Female	65 to 69	36% (24-49%)	64% (51-76%)	1.3 (0.69-2.1)	2.9% (1.6-4.7%)	4.2% (2.3-6.7%)	2000 (1100-3200)	50000 (28000-78000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	2016	Female	70 to 74	34% (22-47%)	66% (53-78%)	0.89 (0.45-1.5)	1.8% (0.71-3.4%)	2.5% (1-4.6%)	1300 (530-2500)	27000 (11000-50000)
Brazil	2016	Female	75 to 79	29% (18-42%)	71% (58-82%)	0.72 (0.35-1.3)	1.3% (0.32-2.6%)	1.6% (0.4-3.5%)	1100 (260-2300)	17000 (4200-35000)
Brazil	2016	Female	80 to 84	25% (15-36%)	75% (64-85%)	0.71 (0.36-1.2)	1.4% (0.49-2.6%)	1.7% (0.6-3.3%)	1300 (440-2500)	15000 (5300-28000)
Brazil	2016	Female	85 to 89	23% (14-32%)	77% (68-86%)	0.75 (0.37-1.3)	1.6% (0.64-3%)	1.9% (0.72-3.6%)	1300 (470-2400)	10000 (4100-19000)
Brazil	2016	Female	90 to 94	23% (14-33%)	77% (67-86%)	0.88 (0.41-1.6)	1.9% (0.68-3.6%)	2.2% (0.73-4.3%)	1000 (340-2100)	5700 (2100-11000)
Brazil	2016	Female	95 plus	25% (13-40%)	75% (60-87%)	1.1 (0.44-2.1)	2.3% (0.74-4.6%)	2.7% (0.72-5.5%)	460 (120-940)	1700 (540-3200)
Brazil	2016	Male	All Ages	71% (68-75%)	29% (25-32%)	3 (2.4-3.7)	11% (8.9-13%)	10% (8.1-13%)	76000 (60000-93000)	3400000 (2900000-4000000)
Brazil	2016	Male	15 to 19	71% (62-81%)	29% (19-38%)	1.9 (1-3)	8.1% (4.8-12%)	11% (5.8-17%)	1800 (1000-2800)	150000 (86000-220000)
Brazil	2016	Male	20 to 24	81% (74-87%)	19% (13-26%)	3.9 (2.5-5.8)	16% (11-22%)	20% (13-28%)	4300 (2800-6000)	330000 (230000-450000)
Brazil	2016	Male	25 to 29	82% (74-88%)	18% (12-26%)	4.8 (3.2-7)	18% (14-23%)	22% (16-30%)	4700 (3300-6300)	350000 (260000-460000)
Brazil	2016	Male	30 to 34	82% (74-88%)	18% (12-26%)	4.8 (3.2-7)	17% (14-21%)	22% (17-28%)	5000 (3800-6500)	360000 (280000-450000)
Brazil	2016	Male	35 to 39	81% (75-88%)	19% (12-25%)	4.4 (2.8-6.3)	16% (13-20%)	21% (16-27%)	5200 (3900-6600)	340000 (270000-410000)
Brazil	2016	Male	40 to 44	80% (73-86%)	20% (14-27%)	4.3 (2.7-6.3)	16% (13-20%)	21% (16-26%)	5900 (4500-7200)	330000 (260000-400000)
Brazil	2016	Male	45 to 49	78% (70-85%)	22% (15-30%)	4.5 (2.9-6.8)	16% (13-19%)	21% (16-25%)	7100 (5500-8600)	340000 (270000-410000)
Brazil	2016	Male	50 to 54	77% (69-84%)	23% (16-31%)	4.4 (2.7-6.4)	15% (11-18%)	19% (15-23%)	8600 (6600-11000)	360000 (270000-430000)
Brazil	2016	Male	55 to 59	75% (67-83%)	25% (17-33%)	4.1 (2.6-6)	13% (9.7-16%)	16% (12-20%)	8900 (6600-11000)	310000 (230000-380000)
Brazil	2016	Male	60 to 64	74% (63-83%)	26% (17-37%)	3.7 (2.2-5.8)	10% (6.7-14%)	13% (8.3-17%)	7900 (5200-11000)	230000 (160000-310000)
Brazil	2016	Male	65 to 69	73% (62-82%)	27% (18-38%)	3 (1.9-4.6)	7.9% (4.9-11%)	9.5% (5.9-13%)	6500 (4000-9100)	160000 (100000-220000)
Brazil	2016	Male	70 to 74	69% (58-79%)	31% (21-42%)	2.3 (1.3-3.6)	5.2% (2.8-8.2%)	6.1% (3.2-9.7%)	4300 (2200-6900)	84000 (44000-130000)
Brazil	2016	Male	75 to 79	64% (52-76%)	36% (24-48%)	1.7 (0.9-2.7)	3.3% (1.4-5.8%)	3.8% (1.5-6.8%)	2800 (1100-5100)	43000 (19000-77000)
Brazil	2016	Male	80 to 84	55% (42-67%)	45% (33-58%)	1.3 (0.74-2.2)	2.4% (1-4.3%)	2.6% (1-4.9%)	1800 (720-3400)	21000 (9100-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Brazil	2016	Male	85 to 89	47% (35-59%)	53% (41-65%)	1.1 (0.64-1.9)	1.8% (0.75-3.5%)	1.9% (0.7-3.9%)	1000 (370-2000)	8500 (3600-16000)
Brazil	2016	Male	90 to 94	45% (33-58%)	55% (42-67%)	1.1 (0.56-1.8)	1.5% (0.45-3.2%)	1.5% (0.31-3.5%)	480 (98-1100)	2800 (850-6100)
Brazil	2016	Male	95 plus	46% (31-60%)	54% (40-69%)	1 (0.46-1.9)	1.4% (0.25-3.3%)	1.3% (-0.0038-3.5%)	96 (-0.29-260)	410 (74-990)
Paraguay	1990	Female	All Ages	49% (42-55%)	51% (45-58%)	0.94 (0.74-1.2)	1.5% (1-2%)	1.7% (0.74-2.8%)	130 (60-220)	7800 (5400-10000)
Paraguay	1990	Female	15 to 19	62% (49-74%)	38% (26-51%)	1.2 (0.74-1.8)	2.1% (1.4-3%)	4.5% (2.5-7%)	4.4 (2.4-7)	540 (340-770)
Paraguay	1990	Female	20 to 24	71% (59-80%)	29% (20-41%)	1.9 (1.2-2.8)	3.8% (2.6-5.1%)	6% (3.6-8.9%)	7.5 (4.6-11)	1000 (670-1400)
Paraguay	1990	Female	25 to 29	71% (59-80%)	29% (20-41%)	1.9 (1.3-2.8)	3.8% (2.7-5%)	5.7% (3.5-8.4%)	7.6 (4.7-11)	910 (630-1200)
Paraguay	1990	Female	30 to 34	67% (54-77%)	33% (23-46%)	1.8 (1.2-2.7)	3.4% (2.3-4.7%)	5.3% (3.1-7.8%)	7.8 (4.7-12)	780 (530-1100)
Paraguay	1990	Female	35 to 39	60% (46-71%)	40% (29-54%)	1.7 (1-2.6)	3.3% (2-4.8%)	5.3% (2.9-8.2%)	9.4 (5.1-15)	740 (440-1100)
Paraguay	1990	Female	40 to 44	57% (44-68%)	43% (32-56%)	1.8 (1.1-2.6)	3.7% (2.2-5.7%)	6.6% (3.7-10%)	13 (7-20)	770 (460-1200)
Paraguay	1990	Female	45 to 49	55% (40-66%)	45% (34-60%)	1.9 (1.1-2.9)	4.3% (2.2-7%)	7.4% (3.6-12%)	17 (8.1-28)	820 (430-1300)
Paraguay	1990	Female	50 to 54	48% (36-60%)	52% (40-64%)	1.6 (0.96-2.5)	4.2% (2-7%)	6.9% (3.2-12%)	19 (9-33)	800 (390-1300)
Paraguay	1990	Female	55 to 59	39% (28-51%)	61% (49-72%)	1.3 (0.75-2.1)	3.5% (1.6-6.1%)	5.4% (2.3-9.4%)	18 (7.6-31)	620 (270-1100)
Paraguay	1990	Female	60 to 64	33% (22-46%)	67% (54-78%)	1.1 (0.57-1.8)	2.4% (0.65-4.9%)	3.5% (0.92-7.1%)	16 (4.2-33)	470 (130-970)
Paraguay	1990	Female	65 to 69	32% (22-43%)	68% (57-78%)	0.83 (0.47-1.3)	1.3% (0.0067-3.1%)	1.8% (-0.0025-4.4%)	9.8 (-0.014-23)	240 (1.2-560)
Paraguay	1990	Female	70 to 74	31% (22-42%)	69% (58-78%)	0.59 (0.33-0.99)	0.25% (-0.79-1.6%)	0.36% (-1.1-2.3%)	2.2 (-7.1-14)	42 (-130-270)
Paraguay	1990	Female	75 to 79	27% (18-39%)	73% (61-82%)	0.46 (0.24-0.8)	-0.15% (-1.1-0.88%)	-0.22% (-1.4-1.1%)	-1.6 (-10-7.9)	-20 (-150-120)
Paraguay	1990	Female	80 to 84	22% (13-32%)	78% (68-87%)	0.43 (0.23-0.71)	0.091% (-0.67-1%)	0.064% (-0.92-1.2%)	0.43 (-6.1-8.2)	8.1 (-60-92)
Paraguay	1990	Female	85 to 89	20% (12-29%)	80% (71-88%)	0.44 (0.23-0.76)	0.3% (-0.44-1.2%)	0.3% (-0.62-1.4%)	1.8 (-3.6-8.4)	17 (-24-66)
Paraguay	1990	Female	90 to 94	21% (13-30%)	79% (70-87%)	0.53 (0.25-0.96)	0.48% (-0.41-1.7%)	0.49% (-0.6-1.9%)	1.7 (-2-6.7)	10 (-8.5-36)
Paraguay	1990	Female	95 plus	23% (12-38%)	77% (62-88%)	0.66 (0.28-1.3)	0.78% (-0.41-2.6%)	0.81% (-0.66-3.1%)	0.99 (-0.8-3.8)	3.9 (-2.1-13)
Paraguay	1990	Male	All Ages	76% (70-81%)	24% (19-30%)	3.3 (2.9-3.9)	7.4% (6.3-8.6%)	11% (8.6-13%)	1000 (810-1200)	44000 (37000-52000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	1990	Male	15 to 19	77% (66-86%)	23% (14-34%)	3.1 (2.1-4.4)	8.2% (5.9-11%)	14% (9.4-20%)	22 (15-31)	2100 (1500-2900)
Paraguay	1990	Male	20 to 24	88% (81-93%)	12% (7.3-19%)	5.8 (4.2-7.6)	17% (13-22%)	26% (19-35%)	53 (39-71)	4800 (3700-6000)
Paraguay	1990	Male	25 to 29	89% (82-93%)	11% (6.6-18%)	5.9 (4.5-7.7)	17% (14-22%)	27% (20-34%)	54 (40-70)	4500 (3500-5500)
Paraguay	1990	Male	30 to 34	90% (84-94%)	9.7% (5.7-16%)	6.4 (4.8-8.3)	17% (14-21%)	27% (21-34%)	56 (43-71)	4100 (3300-5100)
Paraguay	1990	Male	35 to 39	91% (86-95%)	9% (5.3-14%)	7 (4.9-9.2)	18% (14-22%)	28% (21-35%)	64 (48-80)	4100 (3200-5000)
Paraguay	1990	Male	40 to 44	91% (85-94%)	9.1% (5.5-15%)	7.1 (4.9-9.3)	18% (14-22%)	28% (22-34%)	69 (53-85)	3800 (3000-4600)
Paraguay	1990	Male	45 to 49	90% (83-94%)	9.9% (5.6-17%)	7.2 (5-9.9)	18% (15-22%)	27% (22-32%)	80 (64-97)	3800 (3100-4600)
Paraguay	1990	Male	50 to 54	89% (82-93%)	11% (6.9-18%)	7 (4.8-9.5)	18% (14-22%)	25% (20-30%)	94 (74-110)	3800 (3100-4700)
Paraguay	1990	Male	55 to 59	87% (78-92%)	13% (8-22%)	6.4 (4.4-9)	17% (13-21%)	23% (17-28%)	95 (73-120)	3300 (2600-4100)
Paraguay	1990	Male	60 to 64	84% (75-91%)	16% (9.2-25%)	5.6 (3.7-8.1)	15% (10-18%)	19% (14-24%)	110 (77-140)	3200 (2300-4000)
Paraguay	1990	Male	65 to 69	81% (71-89%)	19% (11-29%)	4.9 (3.3-6.9)	13% (9-17%)	17% (11-22%)	110 (75-140)	2600 (1800-3500)
Paraguay	1990	Male	70 to 74	77% (65-85%)	23% (15-35%)	4.2 (2.6-6.1)	11% (5.9-15%)	13% (7.3-19%)	95 (53-130)	1800 (1100-2600)
Paraguay	1990	Male	75 to 79	72% (57-83%)	28% (17-43%)	2.9 (1.7-4.5)	7.1% (3.1-11%)	8.4% (3.7-14%)	64 (27-100)	950 (430-1500)
Paraguay	1990	Male	80 to 84	66% (51-79%)	34% (21-49%)	2.2 (1.3-3.4)	4.8% (2.1-8.5%)	5.6% (2.2-10%)	33 (13-61)	370 (160-660)
Paraguay	1990	Male	85 to 89	63% (46-76%)	37% (24-54%)	1.9 (1.1-3)	3.3% (1.1-6.8%)	3.7% (1.1-7.9%)	17 (4.8-35)	130 (44-270)
Paraguay	1990	Male	90 to 94	63% (48-76%)	37% (24-52%)	1.6 (0.92-2.7)	1.8% (-0.055-4.8%)	1.9% (-0.4-5.4%)	4.3 (-0.92-13)	26 (-0.77-69)
Paraguay	1990	Male	95 plus	64% (45-80%)	36% (20-55%)	1.3 (0.62-2.4)	1% (-0.68-4.2%)	0.78% (-1.3-4.5%)	0.53 (-0.88-3.1)	2.9 (-1.9-12)
Paraguay	1995	Female	All Ages	50% (43-56%)	50% (44-57%)	1.1 (0.83-1.3)	1.8% (1.2-2.5%)	2.1% (1-3.4%)	190 (92-310)	10000 (6900-14000)
Paraguay	1995	Female	15 to 19	64% (50-75%)	36% (25-50%)	1.4 (0.84-2)	2.4% (1.6-3.5%)	5.6% (3.1-8.8%)	6.4 (3.5-10)	720 (470-1000)
Paraguay	1995	Female	20 to 24	72% (60-81%)	28% (19-40%)	2.1 (1.3-3.1)	4.2% (2.9-5.6%)	7.5% (4.6-11%)	9 (5.5-13)	1100 (770-1500)
Paraguay	1995	Female	25 to 29	72% (60-81%)	28% (19-40%)	2.1 (1.4-3.1)	4.1% (2.9-5.4%)	6.8% (4.4-9.7%)	9 (5.8-13)	1100 (740-1400)
Paraguay	1995	Female	30 to 34	68% (54-78%)	32% (22-46%)	2 (1.3-3)	3.7% (2.5-5.1%)	6.3% (3.8-9.1%)	9.4 (5.7-14)	930 (630-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	1995	Female	35 to 39	61% (48-72%)	39% (28-52%)	1.9 (1.2-2.9)	3.5% (2.2-5.1%)	6.1% (3.5-9.4%)	12 (6.6-18)	900 (570-1300)
Paraguay	1995	Female	40 to 44	58% (45-68%)	42% (32-55%)	2 (1.2-2.8)	4.1% (2.4-6%)	7.4% (4.2-11%)	17 (9.7-26)	1000 (630-1500)
Paraguay	1995	Female	45 to 49	56% (42-67%)	44% (33-58%)	2 (1.2-3.2)	4.7% (2.3-7.5%)	8.2% (4-13%)	23 (11-36)	1100 (580-1700)
Paraguay	1995	Female	50 to 54	49% (37-61%)	51% (39-63%)	1.8 (1.1-2.8)	4.8% (2.3-7.7%)	7.7% (3.5-12%)	26 (12-43)	1100 (530-1700)
Paraguay	1995	Female	55 to 59	40% (28-51%)	60% (49-72%)	1.5 (0.85-2.4)	4% (1.9-6.9%)	6.1% (2.8-11%)	26 (12-45)	900 (430-1600)
Paraguay	1995	Female	60 to 64	34% (23-46%)	66% (54-77%)	1.2 (0.64-2)	2.9% (0.85-5.7%)	4.1% (1.2-8%)	20 (6.1-40)	600 (190-1200)
Paraguay	1995	Female	65 to 69	33% (23-44%)	67% (56-77%)	0.93 (0.52-1.5)	1.6% (0.22-3.7%)	2.3% (0.29-5.1%)	15 (2-36)	370 (45-850)
Paraguay	1995	Female	70 to 74	32% (23-43%)	68% (57-77%)	0.66 (0.37-1.1)	0.43% (-0.78-2%)	0.59% (-1.1-2.7%)	4.7 (-8.3-21)	87 (-160-410)
Paraguay	1995	Female	75 to 79	28% (19-40%)	72% (60-81%)	0.52 (0.28-0.9)	-0.012% (-1-1.2%)	-0.035% (-1.3-1.6%)	-0.29 (-12-14)	-2 (-170-210)
Paraguay	1995	Female	80 to 84	23% (14-33%)	77% (67-86%)	0.49 (0.27-0.82)	0.26% (-0.55-1.3%)	0.28% (-0.76-1.6%)	2.4 (-6.2-13)	29 (-61-150)
Paraguay	1995	Female	85 to 89	21% (12-31%)	79% (69-88%)	0.51 (0.26-0.88)	0.51% (-0.33-1.7%)	0.55% (-0.47-2%)	4.1 (-3.6-15)	35 (-22-120)
Paraguay	1995	Female	90 to 94	22% (13-32%)	78% (68-87%)	0.61 (0.29-1.1)	0.73% (-0.29-2.3%)	0.79% (-0.46-2.7%)	3.5 (-2-12)	20 (-8-63)
Paraguay	1995	Female	95 plus	24% (13-39%)	76% (61-87%)	0.77 (0.32-1.5)	1.1% (-0.25-3.4%)	1.2% (-0.48-4.1%)	2 (-0.8-6.7)	7.4 (-1.7-23)
Paraguay	1995	Male	All Ages	76% (70-82%)	24% (18-30%)	3.7 (3.1-4.3)	9% (7.5-11%)	13% (10-15%)	1400 (1100-1700)	58000 (49000-69000)
Paraguay	1995	Male	15 to 19	78% (66-86%)	22% (14-34%)	3.4 (2.3-4.8)	9.3% (6.7-13%)	16% (11-22%)	33 (22-47)	3000 (2100-4000)
Paraguay	1995	Male	20 to 24	88% (81-93%)	12% (7.2-19%)	6.3 (4.5-8.3)	19% (14-24%)	28% (20-37%)	72 (51-94)	6000 (4500-7700)
Paraguay	1995	Male	25 to 29	90% (83-93%)	10% (6.6-17%)	6.4 (4.8-8.5)	19% (15-24%)	28% (21-36%)	75 (55-98)	5900 (4600-7500)
Paraguay	1995	Male	30 to 34	90% (84-94%)	9.6% (5.7-16%)	6.9 (5.1-9)	19% (15-23%)	28% (22-35%)	78 (59-98)	5500 (4400-6900)
Paraguay	1995	Male	35 to 39	91% (86-95%)	8.9% (5.4-14%)	7.5 (5.2-9.9)	19% (15-23%)	29% (22-36%)	89 (68-110)	5500 (4300-6800)
Paraguay	1995	Male	40 to 44	91% (86-95%)	9% (5.4-14%)	7.6 (5.3-10)	19% (16-23%)	29% (23-34%)	110 (83-130)	5700 (4600-6800)
Paraguay	1995	Male	45 to 49	90% (83-94%)	9.8% (5.7-17%)	7.8 (5.4-11)	20% (16-23%)	28% (23-33%)	110 (92-140)	5300 (4400-6400)
Paraguay	1995	Male	50 to 54	89% (82-93%)	11% (6.8-18%)	7.6 (5.3-10)	19% (15-23%)	26% (21-31%)	120 (98-150)	5000 (4000-6000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	1995	Male	55 to 59	87% (78-92%)	13% (7.7-22%)	7 (4.7-9.8)	18% (14-22%)	24% (19-29%)	140 (110-170)	4700 (3700-5700)
Paraguay	1995	Male	60 to 64	84% (75-91%)	16% (9.2-25%)	6 (3.9-8.7)	16% (11-20%)	20% (15-25%)	130 (95-160)	3700 (2800-4600)
Paraguay	1995	Male	65 to 69	81% (71-89%)	19% (11-29%)	5.3 (3.6-7.5)	14% (10-18%)	18% (13-23%)	140 (100-180)	3500 (2500-4400)
Paraguay	1995	Male	70 to 74	77% (66-85%)	23% (15-34%)	4.5 (2.9-6.5)	12% (7-16%)	14% (8.4-20%)	130 (77-180)	2500 (1500-3400)
Paraguay	1995	Male	75 to 79	72% (58-83%)	28% (17-42%)	3.1 (1.9-4.9)	7.9% (3.9-12%)	9.4% (4.5-15%)	87 (41-130)	1300 (640-2000)
Paraguay	1995	Male	80 to 84	67% (51-80%)	33% (20-49%)	2.4 (1.4-3.6)	5.5% (2.5-9.3%)	6.4% (2.7-11%)	49 (21-84)	540 (250-920)
Paraguay	1995	Male	85 to 89	63% (48-76%)	37% (24-52%)	2.1 (1.2-3.2)	3.9% (1.5-7.7%)	4.4% (1.5-8.9%)	26 (8.8-53)	210 (76-400)
Paraguay	1995	Male	90 to 94	63% (48-77%)	37% (23-52%)	1.8 (1-2.9)	2.3% (0.37-5.7%)	2.5% (0.09-6.4%)	7.4 (0.28-19)	43 (6.7-100)
Paraguay	1995	Male	95 plus	65% (46-81%)	35% (19-54%)	1.5 (0.71-2.8)	1.5% (-0.44-5%)	1.2% (-1.1-5.6%)	1.1 (-0.99-5.1)	5.3 (-1.6-19)
Paraguay	2000	Female	All Ages	46% (40-52%)	54% (48-60%)	0.85 (0.67-1.1)	1.5% (1-2%)	1.5% (0.67-2.5%)	160 (69-260)	9000 (6100-12000)
Paraguay	2000	Female	15 to 19	63% (52-73%)	37% (27-48%)	1.2 (0.77-1.7)	2.2% (1.4-3.2%)	5% (2.7-7.8%)	6.6 (3.6-10)	770 (510-1100)
Paraguay	2000	Female	20 to 24	68% (59-76%)	32% (24-41%)	1.7 (1.2-2.4)	3.7% (2.5-4.9%)	6% (3.8-8.9%)	8.1 (4.9-12)	1100 (760-1600)
Paraguay	2000	Female	25 to 29	68% (60-76%)	32% (24-40%)	1.7 (1.2-2.3)	3.5% (2.6-4.6%)	5.4% (3.5-7.7%)	7.2 (4.6-10)	960 (670-1300)
Paraguay	2000	Female	30 to 34	64% (55-73%)	36% (27-45%)	1.6 (1.1-2.4)	3.2% (2.2-4.3%)	5% (3.1-7.3%)	7.9 (4.8-11)	870 (600-1200)
Paraguay	2000	Female	35 to 39	55% (46-65%)	45% (35-54%)	1.4 (0.93-2.2)	2.9% (1.9-4.1%)	4.7% (2.8-7.4%)	9.4 (5.5-15)	810 (550-1200)
Paraguay	2000	Female	40 to 44	51% (42-60%)	49% (40-58%)	1.5 (0.94-2.1)	3.1% (1.9-4.7%)	5.5% (3.2-8.5%)	14 (7.9-22)	900 (550-1300)
Paraguay	2000	Female	45 to 49	49% (38-59%)	51% (41-62%)	1.5 (0.95-2.3)	3.5% (1.9-5.7%)	6.1% (3.2-9.9%)	21 (11-34)	1000 (550-1600)
Paraguay	2000	Female	50 to 54	42% (32-52%)	58% (48-68%)	1.3 (0.81-2)	3.5% (1.7-5.8%)	5.5% (2.5-9.2%)	23 (11-38)	940 (460-1500)
Paraguay	2000	Female	55 to 59	31% (23-42%)	69% (58-77%)	1 (0.61-1.6)	2.7% (1.3-4.6%)	4% (1.8-7.1%)	20 (8.9-34)	700 (330-1200)
Paraguay	2000	Female	60 to 64	29% (20-39%)	71% (61-80%)	0.88 (0.49-1.4)	2% (0.61-4.2%)	2.8% (0.77-5.9%)	18 (4.8-37)	520 (160-1100)
Paraguay	2000	Female	65 to 69	31% (23-40%)	69% (60-77%)	0.77 (0.47-1.2)	1.2% (-0.026-2.8%)	1.6% (-0.11-3.9%)	11 (-0.78-27)	260 (-6.3-640)
Paraguay	2000	Female	70 to 74	29% (21-39%)	71% (61-79%)	0.52 (0.29-0.86)	0.27% (-0.65-1.4%)	0.38% (-0.86-1.9%)	3.5 (-8.1-18)	63 (-160-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	2000	Female	75 to 79	24% (16-35%)	76% (65-84%)	0.4 (0.21-0.68)	-0.021% (-0.78-0.86%)	-0.035% (-1-1.1%)	-0.35 (-11-11)	-4.1 (-160-170)
Paraguay	2000	Female	80 to 84	21% (13-31%)	79% (69-87%)	0.41 (0.23-0.68)	0.13% (-0.54-0.93%)	0.13% (-0.73-1.2%)	1.3 (-7.2-12)	18 (-74-130)
Paraguay	2000	Female	85 to 89	20% (12-29%)	80% (71-88%)	0.44 (0.23-0.75)	0.35% (-0.38-1.3%)	0.37% (-0.53-1.5%)	3.4 (-4.9-14)	30 (-32-110)
Paraguay	2000	Female	90 to 94	21% (13-31%)	79% (69-87%)	0.53 (0.25-0.93)	0.51% (-0.31-1.7%)	0.53% (-0.46-1.9%)	3 (-2.6-11)	18 (-11-57)
Paraguay	2000	Female	95 plus	24% (13-38%)	76% (62-87%)	0.66 (0.28-1.3)	0.8% (-0.31-2.6%)	0.83% (-0.53-3%)	1.8 (-1.1-6.3)	7.1 (-2.6-22)
Paraguay	2000	Male	All Ages	80% (75-84%)	20% (16-25%)	3.6 (3.1-4.2)	9.2% (7.7-11%)	12% (9.8-15%)	1600 (1200-1900)	67000 (57000-79000)
Paraguay	2000	Male	15 to 19	83% (76-89%)	17% (11-24%)	3.5 (2.4-4.7)	9.6% (6.8-13%)	17% (11-22%)	43 (28-59)	3800 (2700-5000)
Paraguay	2000	Male	20 to 24	90% (85-93%)	9.9% (6.6-15%)	6 (4.5-7.7)	18% (14-23%)	27% (20-35%)	87 (63-110)	7200 (5500-9200)
Paraguay	2000	Male	25 to 29	90% (86-93%)	9.7% (6.7-14%)	5.9 (4.4-7.8)	18% (14-23%)	27% (20-34%)	80 (59-100)	6300 (5000-8000)
Paraguay	2000	Male	30 to 34	92% (88-95%)	8.1% (5.3-12%)	6.6 (4.9-8.5)	18% (14-22%)	27% (20-34%)	87 (65-110)	6100 (4900-7700)
Paraguay	2000	Male	35 to 39	93% (90-95%)	7.2% (4.7-10%)	7.1 (5-9.4)	19% (15-23%)	28% (21-34%)	100 (77-130)	6400 (4900-7900)
Paraguay	2000	Male	40 to 44	92% (89-95%)	7.6% (5-11%)	7.2 (5.1-9.3)	19% (15-23%)	27% (22-33%)	120 (93-150)	6600 (5200-7900)
Paraguay	2000	Male	45 to 49	92% (88-95%)	8% (5.1-12%)	7.4 (5.3-9.9)	19% (15-23%)	27% (22-32%)	140 (120-170)	6800 (5500-8100)
Paraguay	2000	Male	50 to 54	91% (86-94%)	9.1% (5.9-14%)	7.3 (5.1-10)	18% (15-22%)	25% (20-30%)	150 (120-180)	6100 (4900-7400)
Paraguay	2000	Male	55 to 59	88% (81-92%)	12% (7.6-19%)	6.5 (4.5-8.9)	17% (13-21%)	22% (17-27%)	150 (120-190)	5300 (4100-6400)
Paraguay	2000	Male	60 to 64	84% (76-90%)	16% (10-24%)	5.4 (3.6-7.8)	14% (10-18%)	18% (13-23%)	150 (110-190)	4400 (3100-5500)
Paraguay	2000	Male	65 to 69	81% (73-88%)	19% (12-27%)	4.8 (3.4-6.5)	13% (8.7-17%)	16% (11-20%)	140 (95-180)	3400 (2300-4300)
Paraguay	2000	Male	70 to 74	78% (69-85%)	22% (15-31%)	4.1 (2.7-5.8)	10% (6-14%)	13% (7.3-18%)	140 (80-200)	2700 (1600-3700)
Paraguay	2000	Male	75 to 79	71% (57-82%)	29% (18-43%)	2.7 (1.6-4.2)	6.6% (3.2-10%)	7.8% (3.8-12%)	87 (41-140)	1300 (650-2100)
Paraguay	2000	Male	80 to 84	66% (50-79%)	34% (21-50%)	2 (1.2-3.2)	4.5% (2-7.8%)	5.1% (2.1-9.1%)	50 (21-89)	560 (250-960)
Paraguay	2000	Male	85 to 89	64% (49-77%)	36% (23-51%)	1.9 (1.1-2.9)	3.2% (1.2-6.1%)	3.6% (1.2-7%)	27 (8.9-53)	220 (80-420)
Paraguay	2000	Male	90 to 94	64% (50-78%)	36% (22-50%)	1.6 (0.93-2.6)	1.8% (0.16-4.3%)	1.9% (-0.17-4.8%)	7.4 (-0.68-19)	44 (4-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	2000	Male	95 plus	66% (47-81%)	34% (19-53%)	1.4 (0.65-2.5)	1.1% (-0.44-3.8%)	0.88% (-1.1-4%)	1 (-1.2-4.8)	5.4 (-2.1-19)
Paraguay	2005	Female	All Ages	47% (41-52%)	53% (48-59%)	0.76 (0.6-0.96)	1.3% (0.88-1.8%)	1.1% (0.33-2%)	130 (39-250)	8800 (5800-12000)
Paraguay	2005	Female	15 to 19	63% (52-73%)	37% (27-48%)	1 (0.66-1.5)	2% (1.3-2.9%)	4.4% (2.3-7.3%)	6.6 (3.4-11)	790 (500-1100)
Paraguay	2005	Female	20 to 24	68% (59-76%)	32% (24-41%)	1.5 (0.97-2.1)	3.4% (2.4-4.6%)	5.2% (3-7.8%)	8.4 (4.9-13)	1200 (850-1800)
Paraguay	2005	Female	25 to 29	68% (59-76%)	32% (24-41%)	1.4 (0.99-2)	3.2% (2.3-4.3%)	4.5% (2.8-6.8%)	7.2 (4.2-11)	1000 (720-1400)
Paraguay	2005	Female	30 to 34	64% (55-73%)	36% (27-45%)	1.4 (0.92-2)	2.8% (1.9-4%)	4.1% (2.5-6.4%)	7 (4.2-11)	830 (560-1200)
Paraguay	2005	Female	35 to 39	55% (46-64%)	45% (36-54%)	1.2 (0.77-1.9)	2.5% (1.7-3.7%)	3.9% (2.2-6.1%)	8.9 (5-14)	790 (510-1100)
Paraguay	2005	Female	40 to 44	51% (42-60%)	49% (40-58%)	1.2 (0.8-1.9)	2.7% (1.6-3.9%)	4.5% (2.5-7.1%)	14 (7.5-22)	880 (540-1300)
Paraguay	2005	Female	45 to 49	49% (38-59%)	51% (41-62%)	1.3 (0.82-2)	2.9% (1.5-4.7%)	4.8% (2.3-8.1%)	19 (9.2-33)	980 (510-1600)
Paraguay	2005	Female	50 to 54	42% (32-52%)	58% (48-68%)	1.1 (0.71-1.7)	2.8% (1.3-4.6%)	4.3% (1.8-7.3%)	23 (9.9-40)	960 (450-1600)
Paraguay	2005	Female	55 to 59	31% (23-42%)	69% (58-77%)	0.88 (0.52-1.4)	2.1% (0.89-3.8%)	3% (1.2-5.6%)	19 (7.8-35)	680 (290-1300)
Paraguay	2005	Female	60 to 64	29% (20-39%)	71% (61-80%)	0.76 (0.42-1.2)	1.4% (0.2-3.3%)	2% (0.28-4.6%)	15 (2.2-34)	440 (62-1000)
Paraguay	2005	Female	65 to 69	31% (23-40%)	69% (60-77%)	0.67 (0.4-1)	0.64% (-0.42-2%)	0.92% (-0.52-2.8%)	8.2 (-4.6-25)	190 (-130-600)
Paraguay	2005	Female	70 to 74	29% (21-39%)	71% (61-79%)	0.45 (0.26-0.75)	-0.068% (-0.9-0.85%)	-0.047% (-1.2-1.2%)	-0.48 (-12-12)	-18 (-240-220)
Paraguay	2005	Female	75 to 79	25% (16-35%)	75% (65-84%)	0.35 (0.18-0.6)	-0.24% (-0.93-0.46%)	-0.3% (-1.2-0.6%)	-4.1 (-16-7.8)	-62 (-240-120)
Paraguay	2005	Female	80 to 84	22% (14-31%)	78% (69-86%)	0.36 (0.2-0.6)	-0.084% (-0.76-0.58%)	-0.12% (-0.97-0.69%)	-1.7 (-13-9.9)	-15 (-140-110)
Paraguay	2005	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.39 (0.2-0.65)	0.12% (-0.54-0.91%)	0.097% (-0.75-1.1%)	1.2 (-8.8-13)	14 (-60-100)
Paraguay	2005	Female	90 to 94	21% (13-31%)	79% (69-87%)	0.47 (0.23-0.83)	0.26% (-0.49-1.2%)	0.23% (-0.72-1.3%)	1.7 (-5.2-10)	12 (-23-55)
Paraguay	2005	Female	95 plus	24% (13-39%)	76% (61-87%)	0.58 (0.25-1.1)	0.44% (-0.5-1.9%)	0.41% (-0.78-2.1%)	1.2 (-2.2-6)	5.2 (-5.8-21)
Paraguay	2005	Male	All Ages	81% (76-85%)	19% (15-24%)	3.2 (2.8-3.8)	9.2% (7.5-11%)	11% (8.5-14%)	1700 (1300-2100)	73000 (60000-87000)
Paraguay	2005	Male	15 to 19	84% (76-90%)	16% (10-24%)	3.1 (2.1-4.1)	9% (6.3-12%)	15% (9.9-21%)	45 (29-63)	4000 (2800-5300)
Paraguay	2005	Male	20 to 24	90% (85-93%)	9.8% (6.5-15%)	5.2 (3.9-6.7)	17% (13-22%)	24% (17-32%)	97 (69-130)	8200 (6100-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	2005	Male	25 to 29	90% (86-93%)	9.6% (6.6-14%)	5.1 (3.8-6.6)	17% (13-21%)	23% (17-31%)	90 (65-120)	7200 (5500-9100)
Paraguay	2005	Male	30 to 34	92% (88-95%)	8.1% (5.3-12%)	5.6 (4.2-7.4)	17% (13-21%)	24% (17-30%)	88 (64-110)	6300 (4800-8000)
Paraguay	2005	Male	35 to 39	93% (90-95%)	7.2% (4.6-10%)	6.2 (4.3-8.1)	17% (13-22%)	25% (18-31%)	110 (78-140)	6700 (5100-8500)
Paraguay	2005	Male	40 to 44	93% (89-95%)	7.5% (4.9-11%)	6.2 (4.3-8.2)	17% (13-21%)	24% (18-30%)	130 (95-160)	7000 (5300-8800)
Paraguay	2005	Male	45 to 49	92% (88-95%)	8% (5-12%)	6.4 (4.6-8.6)	17% (14-21%)	24% (19-29%)	160 (120-190)	7400 (5800-9000)
Paraguay	2005	Male	50 to 54	91% (87-94%)	9% (5.9-13%)	6.3 (4.4-8.4)	17% (13-20%)	22% (17-27%)	180 (130-220)	7300 (5600-9000)
Paraguay	2005	Male	55 to 59	88% (81-92%)	12% (7.5-19%)	5.5 (3.9-7.5)	15% (11-19%)	20% (14-25%)	180 (130-220)	6200 (4500-7700)
Paraguay	2005	Male	60 to 64	84% (76-90%)	16% (9.8-24%)	4.6 (3.1-6.6)	12% (8.3-16%)	15% (10-20%)	160 (100-210)	4600 (3100-6100)
Paraguay	2005	Male	65 to 69	82% (73-89%)	18% (11-27%)	4.1 (2.9-5.5)	11% (6.6-15%)	13% (8.1-18%)	150 (94-210)	3700 (2300-5000)
Paraguay	2005	Male	70 to 74	78% (69-85%)	22% (15-31%)	3.6 (2.4-4.9)	8.4% (4.4-13%)	10% (5.2-15%)	120 (62-180)	2300 (1200-3500)
Paraguay	2005	Male	75 to 79	71% (58-82%)	29% (18-42%)	2.3 (1.4-3.5)	5.2% (2.3-8.7%)	6.1% (2.6-10%)	83 (35-140)	1300 (560-2100)
Paraguay	2005	Male	80 to 84	66% (49-79%)	34% (21-51%)	1.8 (1.1-2.7)	3.4% (1.4-6%)	3.8% (1.4-6.9%)	48 (18-88)	550 (220-980)
Paraguay	2005	Male	85 to 89	64% (49-78%)	36% (22-51%)	1.6 (0.95-2.5)	2.3% (0.65-4.4%)	2.5% (0.55-5%)	24 (5.4-49)	200 (56-390)
Paraguay	2005	Male	90 to 94	65% (51-77%)	35% (23-49%)	1.4 (0.83-2.2)	1.2% (-0.21-3.1%)	1.2% (-0.62-3.4%)	6 (-3.2-17)	38 (-6.7-97)
Paraguay	2005	Male	95 plus	66% (47-81%)	34% (19-53%)	1.2 (0.57-2.1)	0.71% (-0.65-2.7%)	0.4% (-1.3-2.7%)	0.62 (-2.4-2)	4.5 (-4.1-17)
Paraguay	2010	Female	All Ages	52% (45-58%)	48% (42-55%)	0.87 (0.67-1.1)	1.4% (0.88-2%)	1.1% (0.22-2.1%)	150 (31-300)	10000 (6200-14000)
Paraguay	2010	Female	15 to 19	64% (51-75%)	36% (25-49%)	1 (0.61-1.5)	2.1% (1.3-3%)	4.5% (2.2-7.5%)	7.1 (3.6-12)	830 (520-1200)
Paraguay	2010	Female	20 to 24	72% (60-81%)	28% (19-40%)	1.5 (0.98-2.3)	3.5% (2.4-4.8%)	5.5% (3.2-8.5%)	9.9 (5.7-16)	1400 (930-2000)
Paraguay	2010	Female	25 to 29	72% (60-81%)	28% (19-40%)	1.5 (1-2.2)	3.3% (2.4-4.4%)	4.9% (2.9-7.4%)	9.2 (5.4-14)	1300 (860-1700)
Paraguay	2010	Female	30 to 34	68% (55-78%)	32% (22-45%)	1.5 (0.9-2.2)	2.9% (1.9-4%)	4.4% (2.5-6.7%)	8.8 (5-14)	990 (650-1400)
Paraguay	2010	Female	35 to 39	61% (48-73%)	39% (27-52%)	1.4 (0.81-2.1)	2.7% (1.7-3.8%)	4.4% (2.4-7.1%)	10 (5.4-16)	850 (540-1200)
Paraguay	2010	Female	40 to 44	58% (45-68%)	42% (32-55%)	1.4 (0.87-2.2)	2.8% (1.6-4.3%)	5% (2.5-7.9%)	16 (8.1-25)	980 (560-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	2010	Female	45 to 49	56% (41-67%)	44% (33-59%)	1.5 (0.88-2.3)	2.9% (1.3-5%)	5% (2.1-8.5%)	22 (9-37)	1100 (510-1800)
Paraguay	2010	Female	50 to 54	49% (37-61%)	51% (39-63%)	1.3 (0.81-2.1)	2.9% (1.2-5%)	4.6% (1.9-8%)	27 (11-47)	1100 (460-1900)
Paraguay	2010	Female	55 to 59	39% (28-52%)	61% (48-72%)	1.1 (0.64-1.8)	2.2% (0.74-4.3%)	3.3% (1.1-6.4%)	25 (8.8-49)	890 (300-1700)
Paraguay	2010	Female	60 to 64	34% (23-46%)	66% (54-77%)	0.89 (0.47-1.5)	1.4% (-0.00052-3.3%)	2% (0.025-4.6%)	18 (0.23-42)	520 (-0.21-1200)
Paraguay	2010	Female	65 to 69	33% (23-45%)	67% (55-77%)	0.7 (0.39-1.1)	0.52% (-0.57-2%)	0.75% (-0.7-2.7%)	7.9 (-7.1-29)	180 (-200-690)
Paraguay	2010	Female	70 to 74	32% (23-43%)	68% (57-77%)	0.5 (0.27-0.84)	-0.21% (-1.2-0.85%)	-0.22% (-1.4-1.1%)	-2.9 (-18-15)	-68 (-360-280)
Paraguay	2010	Female	75 to 79	28% (18-41%)	72% (59-82%)	0.39 (0.2-0.7)	-0.39% (-1.2-0.43%)	-0.48% (-1.5-0.57%)	-6.9 (-22-8)	-100 (-320-120)
Paraguay	2010	Female	80 to 84	23% (14-33%)	77% (67-86%)	0.36 (0.2-0.59)	-0.13% (-0.84-0.57%)	-0.18% (-1.1-0.72%)	-3 (-18-12)	-29 (-190-130)
Paraguay	2010	Female	85 to 89	20% (13-30%)	80% (70-87%)	0.38 (0.2-0.65)	0.083% (-0.56-0.83%)	0.048% (-0.75-1%)	0.81 (-12-16)	12 (-84-130)
Paraguay	2010	Female	90 to 94	22% (13-32%)	78% (68-87%)	0.45 (0.22-0.81)	0.21% (-0.5-1.1%)	0.17% (-0.71-1.3%)	1.7 (-6.8-13)	13 (-30-68)
Paraguay	2010	Female	95 plus	24% (13-40%)	76% (60-87%)	0.57 (0.24-1.1)	0.39% (-0.57-1.8%)	0.34% (-0.83-2.1%)	1.3 (-3-7.7)	5.9 (-8.3-28)
Paraguay	2010	Male	All Ages	79% (74-83%)	21% (17-26%)	3.1 (2.5-3.7)	9.7% (7.7-12%)	11% (8-14%)	1900 (1400-2400)	83000 (67000-100000)
Paraguay	2010	Male	15 to 19	78% (67-87%)	22% (13-33%)	2.5 (1.7-3.6)	8.2% (5.6-12%)	13% (8.1-19%)	44 (27-65)	3900 (2700-5500)
Paraguay	2010	Male	20 to 24	88% (81-93%)	12% (7.1-19%)	4.7 (3.3-6.3)	17% (12-21%)	23% (16-31%)	110 (74-150)	9100 (6600-12000)
Paraguay	2010	Male	25 to 29	89% (83-93%)	11% (6.6-17%)	4.8 (3.5-6.4)	17% (13-21%)	23% (16-31%)	110 (79-150)	8800 (6600-11000)
Paraguay	2010	Male	30 to 34	90% (84-94%)	9.8% (5.8-16%)	5.1 (3.7-6.8)	17% (13-21%)	23% (16-30%)	110 (77-150)	7700 (5800-9900)
Paraguay	2010	Male	35 to 39	91% (86-95%)	9.1% (5.5-14%)	5.5 (3.7-7.6)	17% (13-22%)	24% (17-31%)	120 (81-150)	7200 (5400-9100)
Paraguay	2010	Male	40 to 44	91% (85-95%)	9.2% (5.5-15%)	5.6 (3.7-7.7)	17% (13-21%)	24% (17-30%)	140 (100-180)	7500 (5600-9400)
Paraguay	2010	Male	45 to 49	90% (83-94%)	10% (5.6-17%)	5.7 (3.9-7.9)	17% (13-21%)	23% (17-28%)	170 (130-210)	8100 (6100-9900)
Paraguay	2010	Male	50 to 54	89% (82-93%)	11% (6.8-18%)	5.6 (3.8-7.7)	16% (12-20%)	21% (15-26%)	200 (140-250)	8100 (6000-10000)
Paraguay	2010	Male	55 to 59	86% (78-92%)	14% (8-22%)	5.1 (3.4-7.1)	15% (10-19%)	19% (13-24%)	220 (150-290)	7700 (5400-9900)
Paraguay	2010	Male	60 to 64	84% (75-91%)	16% (9.5-25%)	4.4 (2.9-6.4)	12% (7.7-16%)	15% (9.2-20%)	190 (120-270)	5700 (3700-7700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	2010	Male	65 to 69	81% (71-89%)	19% (11-29%)	3.9 (2.5-5.4)	10% (5.8-14%)	12% (6.9-17%)	180 (100-250)	4300 (2500-5900)
Paraguay	2010	Male	70 to 74	77% (65-85%)	23% (15-35%)	3.3 (2-4.7)	7.8% (4-12%)	9.3% (4.7-14%)	150 (74-230)	2900 (1500-4400)
Paraguay	2010	Male	75 to 79	72% (57-83%)	28% (17-43%)	2.3 (1.3-3.5)	4.9% (2-8.5%)	5.7% (2.3-10%)	87 (34-160)	1300 (540-2300)
Paraguay	2010	Male	80 to 84	67% (51-80%)	33% (20-49%)	1.7 (1-2.7)	3.2% (1.2-5.8%)	3.6% (1.2-6.6%)	56 (19-100)	630 (240-1200)
Paraguay	2010	Male	85 to 89	63% (47-77%)	37% (23-53%)	1.5 (0.87-2.4)	2.1% (0.55-4.2%)	2.3% (0.4-4.8%)	30 (5.4-62)	250 (63-490)
Paraguay	2010	Male	90 to 94	63% (48-77%)	37% (23-52%)	1.3 (0.73-2.2)	1.1% (-0.3-3.1%)	1.1% (-0.66-3.4%)	7 (-4.3-22)	46 (-12-130)
Paraguay	2010	Male	95 plus	64% (45-80%)	36% (20-55%)	1.1 (0.5-2)	0.66% (-0.7-2.7%)	0.35% (-1.3-2.7%)	0.71 (-2.7-5.5)	5.5 (-5.8-23)
Paraguay	2016	Female	All Ages	53% (46-59%)	47% (41-54%)	0.94 (0.65-1.3)	1.6% (0.86-2.4%)	1.2% (0.085-2.4%)	180 (13-380)	11000 (6200-17000)
Paraguay	2016	Female	15 to 19	64% (51-75%)	36% (25-49%)	1 (0.6-1.6)	2.1% (1.3-3.1%)	4.8% (2.4-8.2%)	7.4 (3.5-13)	860 (540-1300)
Paraguay	2016	Female	20 to 24	72% (60-81%)	28% (19-40%)	1.6 (0.95-2.5)	3.6% (2.4-5%)	6% (3.1-9.8%)	11 (5.6-18)	1500 (980-2200)
Paraguay	2016	Female	25 to 29	73% (60-81%)	27% (19-40%)	1.6 (1-2.5)	3.4% (2.4-4.7%)	5.3% (2.9-8.2%)	10 (5.6-17)	1400 (950-2000)
Paraguay	2016	Female	30 to 34	68% (55-78%)	32% (22-45%)	1.5 (0.88-2.3)	3% (2-4.3%)	4.7% (2.5-7.4%)	11 (5.6-18)	1300 (820-1800)
Paraguay	2016	Female	35 to 39	61% (48-73%)	39% (27-52%)	1.4 (0.8-2.3)	2.8% (1.7-4.2%)	4.7% (2.5-7.6%)	12 (6.2-21)	1100 (650-1600)
Paraguay	2016	Female	40 to 44	58% (45-69%)	42% (31-55%)	1.5 (0.83-2.3)	2.9% (1.6-4.6%)	5.2% (2.6-8.5%)	16 (7.7-28)	1000 (560-1700)
Paraguay	2016	Female	45 to 49	56% (42-67%)	44% (33-58%)	1.6 (0.87-2.6)	3% (1.2-5.2%)	5.2% (2.1-9.1%)	23 (9.6-41)	1200 (500-2100)
Paraguay	2016	Female	50 to 54	49% (37-61%)	51% (39-63%)	1.4 (0.78-2.3)	2.9% (1.1-5.5%)	4.8% (1.7-8.6%)	30 (11-55)	1200 (470-2300)
Paraguay	2016	Female	55 to 59	40% (28-52%)	60% (48-72%)	1.2 (0.62-2)	2.3% (0.65-4.7%)	3.5% (1.1-7.1%)	29 (8.9-58)	1000 (290-2100)
Paraguay	2016	Female	60 to 64	34% (23-46%)	66% (54-77%)	0.93 (0.47-1.6)	1.5% (-0.085-3.6%)	2.1% (-0.02-5%)	23 (-0.23-56)	690 (-39-1700)
Paraguay	2016	Female	65 to 69	33% (23-44%)	67% (56-77%)	0.72 (0.39-1.2)	0.61% (-0.56-2.2%)	0.89% (-0.69-3.1%)	12 (-9-40)	260 (-250-950)
Paraguay	2016	Female	70 to 74	32% (23-43%)	68% (57-77%)	0.52 (0.27-0.91)	-0.14% (-1.1-0.98%)	-0.12% (-1.4-1.4%)	-1.8 (-21-19)	-54 (-430-360)
Paraguay	2016	Female	75 to 79	28% (18-40%)	72% (60-82%)	0.41 (0.21-0.73)	-0.35% (-1.2-0.51%)	-0.43% (-1.5-0.66%)	-7.7 (-27-11)	-120 (-410-170)
Paraguay	2016	Female	80 to 84	23% (14-33%)	77% (67-86%)	0.39 (0.2-0.66)	-0.098% (-0.87-0.7%)	-0.14% (-1.1-0.86%)	-2.8 (-22-16)	-26 (-230-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Paraguay	2016	Female	85 to 89	21% (13-31%)	79% (69-87%)	0.4 (0.2-0.72)	0.13% (-0.62-1.1%)	0.1% (-0.83-1.3%)	1.8 (-14-22)	21 (-100-170)
Paraguay	2016	Female	90 to 94	22% (13-32%)	78% (68-87%)	0.47 (0.23-0.89)	0.28% (-0.49-1.4%)	0.24% (-0.73-1.6%)	2.9 (-8.8-20)	21 (-39-110)
Paraguay	2016	Female	95 plus	24% (13-39%)	76% (61-87%)	0.59 (0.24-1.2)	0.46% (-0.59-2.1%)	0.42% (-0.88-2.4%)	1.9 (-4.1-11)	8.8 (-11-40)
Paraguay	2016	Male	All Ages	80% (75-84%)	20% (16-25%)	3.4 (2.5-4.4)	11% (8-13%)	11% (7.6-15%)	2300 (1500-3100)	99000 (71000-130000)
Paraguay	2016	Male	15 to 19	78% (68-87%)	22% (13-32%)	2.6 (1.6-3.9)	8.7% (5.8-12%)	14% (8.3-20%)	49 (27-79)	4300 (2700-6400)
Paraguay	2016	Male	20 to 24	88% (81-93%)	12% (7.1-19%)	4.8 (3.1-7)	17% (12-23%)	24% (16-33%)	120 (74-180)	10000 (6900-14000)
Paraguay	2016	Male	25 to 29	90% (83-93%)	10% (6.5-17%)	5 (3.4-7.1)	18% (13-23%)	24% (16-33%)	130 (83-180)	10000 (7200-13000)
Paraguay	2016	Male	30 to 34	90% (85-94%)	9.6% (5.7-15%)	5.3 (3.6-7.5)	17% (13-22%)	24% (17-32%)	130 (90-180)	9600 (6900-13000)
Paraguay	2016	Male	35 to 39	91% (86-95%)	8.9% (5.3-14%)	5.7 (3.6-8.3)	18% (13-23%)	25% (17-32%)	140 (94-190)	8900 (6300-12000)
Paraguay	2016	Male	40 to 44	91% (86-95%)	9% (5.4-14%)	5.8 (3.6-8.5)	17% (13-22%)	24% (17-31%)	150 (98-200)	8100 (5700-11000)
Paraguay	2016	Male	45 to 49	90% (83-94%)	9.8% (5.5-17%)	6 (3.8-8.7)	17% (13-22%)	23% (17-29%)	180 (130-240)	8700 (6300-11000)
Paraguay	2016	Male	50 to 54	89% (83-93%)	11% (6.7-17%)	5.9 (3.7-8.7)	17% (12-21%)	22% (15-28%)	220 (150-290)	9200 (6300-12000)
Paraguay	2016	Male	55 to 59	87% (78-92%)	13% (7.8-22%)	5.3 (3.4-8.1)	15% (10-20%)	19% (13-25%)	250 (170-340)	8900 (6000-12000)
Paraguay	2016	Male	60 to 64	84% (75-91%)	16% (9.3-25%)	4.6 (2.8-7.1)	12% (7.8-17%)	15% (9.4-21%)	260 (160-370)	7700 (4800-11000)
Paraguay	2016	Male	65 to 69	81% (71-89%)	19% (11-29%)	4.1 (2.5-6)	11% (5.8-15%)	13% (7-18%)	240 (130-350)	5900 (3300-8400)
Paraguay	2016	Male	70 to 74	77% (66-85%)	23% (15-34%)	3.4 (2-5.2)	8.1% (3.7-13%)	9.6% (4.2-15%)	190 (85-310)	3800 (1700-6000)
Paraguay	2016	Male	75 to 79	72% (58-84%)	28% (16-42%)	2.4 (1.3-3.9)	5.2% (1.8-9.4%)	6% (2-11%)	130 (42-230)	1900 (670-3500)
Paraguay	2016	Male	80 to 84	67% (51-80%)	33% (20-49%)	1.8 (1-2.9)	3.4% (1.1-6.7%)	3.8% (1-7.6%)	77 (21-160)	870 (280-1700)
Paraguay	2016	Male	85 to 89	64% (48-77%)	36% (23-52%)	1.6 (0.88-2.6)	2.3% (0.45-5.1%)	2.5% (0.33-5.7%)	38 (4.8-87)	320 (60-690)
Paraguay	2016	Male	90 to 94	63% (49-77%)	37% (23-51%)	1.4 (0.77-2.4)	1.3% (-0.33-3.6%)	1.2% (-0.7-4%)	10 (-6.4-34)	67 (-18-190)
Paraguay	2016	Male	95 plus	65% (46-81%)	35% (19-54%)	1.2 (0.51-2.2)	0.77% (-0.7-3.2%)	0.45% (-1.3-3.3%)	1.2 (-3.5-8.6)	8.2 (-7.8-35)
Algeria	1990	Female	All Ages	9.9% (6.1-16%)	90% (84-94%)	0.16 (0.12-0.2)	0.23% (0.03-0.47%)	0.069% (-0.48-0.7%)	41 (-270-420)	9500 (1200-20000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	1990	Female	15 to 19	8.7% (2.8-20%)	91% (80-97%)	0.13 (0.037-0.31)	0.45% (0.24-0.74%)	0.5% (0.16-1.1%)	5.3 (1.7-13)	920 (480-1600)
Algeria	1990	Female	20 to 24	11% (3.3-26%)	89% (74-97%)	0.21 (0.065-0.43)	0.79% (0.47-1.2%)	0.72% (0.22-1.6%)	8.6 (2.6-20)	1600 (960-2600)
Algeria	1990	Female	25 to 29	13% (3.5-28%)	87% (72-96%)	0.25 (0.076-0.51)	0.9% (0.57-1.3%)	0.96% (0.31-2%)	11 (3.6-24)	1700 (1000-2600)
Algeria	1990	Female	30 to 34	14% (3.7-31%)	86% (69-96%)	0.3 (0.1-0.58)	0.9% (0.51-1.4%)	1.2% (0.39-2.3%)	14 (4.7-29)	1600 (870-2500)
Algeria	1990	Female	35 to 39	16% (4.2-32%)	84% (68-96%)	0.36 (0.12-0.69)	0.86% (0.42-1.6%)	1.3% (0.38-2.8%)	18 (5.3-39)	1400 (660-2600)
Algeria	1990	Female	40 to 44	17% (5.8-32%)	83% (68-94%)	0.4 (0.14-0.8)	0.87% (0.28-1.8%)	1.5% (0.32-3.4%)	18 (3.9-41)	1100 (330-2200)
Algeria	1990	Female	45 to 49	17% (5.9-33%)	83% (67-94%)	0.44 (0.14-0.92)	0.77% (0.023-2.1%)	1.3% (0.039-3.7%)	19 (0.55-52)	920 (26-2500)
Algeria	1990	Female	50 to 54	18% (7.4-35%)	82% (65-93%)	0.45 (0.14-0.95)	0.62% (-0.29-2%)	1% (-0.47-3.4%)	20 (-8.6-65)	800 (-370-2600)
Algeria	1990	Female	55 to 59	19% (8-37%)	81% (63-92%)	0.45 (0.13-0.98)	0.41% (-0.62-1.7%)	0.69% (-0.86-2.8%)	17 (-22-66)	550 (-860-2300)
Algeria	1990	Female	60 to 64	21% (8-40%)	79% (60-92%)	0.44 (0.11-1.1)	0.13% (-1-1.6%)	0.23% (-1.4-2.4%)	7.1 (-43-75)	170 (-1300-2200)
Algeria	1990	Female	65 to 69	21% (8.7-39%)	79% (61-91%)	0.39 (0.097-0.86)	-0.17% (-1.3-1%)	-0.2% (-1.8-1.5%)	-6.7 (-60-50)	-200 (-1500-1200)
Algeria	1990	Female	70 to 74	20% (7.4-37%)	80% (63-93%)	0.31 (0.068-0.79)	-0.36% (-1.5-0.78%)	-0.46% (-1.9-1%)	-21 (-90-47)	-420 (-1700-870)
Algeria	1990	Female	75 to 79	18% (6.6-38%)	82% (62-93%)	0.25 (0.054-0.75)	-0.42% (-1.5-0.54%)	-0.53% (-1.9-0.71%)	-25 (-91-34)	-370 (-1400-480)
Algeria	1990	Female	80 to 84	16% (5.6-37%)	84% (63-94%)	0.23 (0.048-0.69)	-0.37% (-1.4-0.45%)	-0.46% (-1.8-0.59%)	-18 (-67-23)	-190 (-720-250)
Algeria	1990	Female	85 to 89	17% (5.5-37%)	83% (63-94%)	0.23 (0.048-0.7)	-0.43% (-1.5-0.33%)	-0.54% (-1.9-0.39%)	-16 (-55-11)	-110 (-400-86)
Algeria	1990	Female	90 to 94	19% (6.2-41%)	81% (59-94%)	0.26 (0.052-0.77)	-0.52% (-1.6-0.32%)	-0.67% (-2-0.37%)	-8.4 (-25-4.7)	-41 (-120-26)
Algeria	1990	Female	95 plus	24% (7.2-47%)	76% (53-93%)	0.31 (0.056-0.94)	-0.59% (-2-0.53%)	-0.79% (-2.5-0.59%)	-3.1 (-9.7-2.3)	-9.5 (-32-8.3)
Algeria	1990	Male	All Ages	7.5% (5-11%)	92% (89-95%)	0.18 (0.14-0.24)	0.61% (0.41-0.88%)	0.67% (0.29-1.3%)	450 (190-860)	27000 (18000-40000)
Algeria	1990	Male	15 to 19	6.4% (2.7-13%)	94% (87-97%)	0.093 (0.029-0.2)	0.73% (0.42-1.2%)	0.71% (0.24-1.5%)	11 (3.3-23)	1600 (890-2600)
Algeria	1990	Male	20 to 24	9.8% (4-19%)	90% (81-96%)	0.21 (0.076-0.42)	1.6% (0.99-2.5%)	1.5% (0.58-3%)	27 (10-55)	3500 (2100-5500)
Algeria	1990	Male	25 to 29	12% (4.8-23%)	88% (77-95%)	0.32 (0.13-0.62)	1.9% (1.2-2.9%)	2.2% (0.89-4.1%)	38 (15-72)	3900 (2300-6200)
Algeria	1990	Male	30 to 34	14% (5-28%)	86% (72-95%)	0.41 (0.17-0.73)	2% (1.2-3.1%)	2.4% (1.1-4.4%)	42 (19-78)	3600 (2100-5900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	1990	Male	35 to 39	15% (5.7-28%)	85% (72-94%)	0.47 (0.18-0.9)	2% (1.1-3.3%)	2.6% (1.2-5%)	48 (21-91)	3400 (1900-5900)
Algeria	1990	Male	40 to 44	15% (6.8-27%)	85% (73-93%)	0.5 (0.18-1)	1.9% (0.96-3.4%)	2.5% (1-4.9%)	38 (16-76)	2400 (1200-4300)
Algeria	1990	Male	45 to 49	15% (5.8-28%)	85% (72-94%)	0.52 (0.16-1.1)	1.8% (0.81-3.5%)	2.4% (0.88-5.1%)	40 (15-84)	2100 (950-4000)
Algeria	1990	Male	50 to 54	14% (6.2-28%)	86% (72-94%)	0.52 (0.17-1.1)	1.5% (0.54-3.1%)	2% (0.53-4.3%)	43 (11-98)	1900 (680-4100)
Algeria	1990	Male	55 to 59	14% (5.9-27%)	86% (73-94%)	0.5 (0.15-1.1)	1.3% (0.39-2.9%)	1.7% (0.38-4%)	52 (12-130)	1900 (570-4400)
Algeria	1990	Male	60 to 64	13% (5.5-26%)	87% (74-94%)	0.45 (0.13-1)	1% (0.15-2.7%)	1.3% (0.11-3.5%)	47 (3.8-130)	1400 (220-3900)
Algeria	1990	Male	65 to 69	13% (5.9-24%)	87% (76-94%)	0.37 (0.11-0.81)	0.66% (-0.045-2%)	0.79% (-0.16-2.5%)	32 (-6.3-100)	810 (-56-2500)
Algeria	1990	Male	70 to 74	12% (5.5-21%)	88% (79-95%)	0.27 (0.08-0.66)	0.37% (-0.2-1.4%)	0.42% (-0.31-1.7%)	21 (-15-82)	430 (-220-1600)
Algeria	1990	Male	75 to 79	10% (4.5-20%)	90% (80-96%)	0.21 (0.053-0.57)	0.18% (-0.29-0.95%)	0.19% (-0.41-1.1%)	8.6 (-20-52)	150 (-250-800)
Algeria	1990	Male	80 to 84	9.1% (3.5-18%)	91% (82-97%)	0.18 (0.045-0.44)	0.14% (-0.25-0.7%)	0.14% (-0.34-0.82%)	4.7 (-12-27)	63 (-110-300)
Algeria	1990	Male	85 to 89	8.8% (3.4-18%)	91% (82-97%)	0.16 (0.041-0.41)	0.075% (-0.29-0.5%)	0.047% (-0.4-0.54%)	1.1 (-9.4-12)	16 (-60-100)
Algeria	1990	Male	90 to 94	9.7% (4-19%)	90% (81-96%)	0.17 (0.042-0.47)	0.01% (-0.38-0.5%)	-0.045% (-0.52-0.54%)	-0.45 (-5.2-5.5)	0.64 (-23-30)
Algeria	1990	Male	95 plus	12% (4.4-26%)	88% (74-96%)	0.2 (0.036-0.56)	0.0074% (-0.49-0.73%)	-0.085% (-0.68-0.77%)	0.23 (-1.9-2.1)	0.084 (-5.3-8.3)
Algeria	1995	Female	All Ages	10% (6-16%)	90% (84-94%)	0.16 (0.12-0.21)	0.23% (0.024-0.46%)	-0.00015% (-0.56-0.63%)	0.42 (-340-390)	9600 (970-20000)
Algeria	1995	Female	15 to 19	9% (2.9-20%)	91% (80-97%)	0.13 (0.036-0.3)	0.42% (0.23-0.68%)	0.43% (0.15-1%)	4.9 (1.6-12)	970 (530-1600)
Algeria	1995	Female	20 to 24	12% (3.4-26%)	88% (74-97%)	0.2 (0.067-0.43)	0.78% (0.48-1.2%)	0.65% (0.21-1.5%)	7.8 (2.6-18)	1800 (1000-2800)
Algeria	1995	Female	25 to 29	13% (3.7-28%)	87% (72-96%)	0.25 (0.08-0.5)	0.87% (0.55-1.3%)	0.87% (0.3-1.7%)	10 (3.5-21)	1800 (1100-2700)
Algeria	1995	Female	30 to 34	15% (3.9-31%)	85% (69-96%)	0.3 (0.096-0.56)	0.85% (0.5-1.3%)	1.1% (0.37-2.1%)	13 (4.7-27)	1700 (950-2600)
Algeria	1995	Female	35 to 39	16% (4.5-31%)	84% (69-95%)	0.35 (0.11-0.65)	0.8% (0.41-1.4%)	1.2% (0.39-2.6%)	17 (5.6-36)	1500 (740-2600)
Algeria	1995	Female	40 to 44	17% (6.4-32%)	83% (68-94%)	0.39 (0.14-0.74)	0.79% (0.28-1.6%)	1.4% (0.32-3.1%)	22 (5.1-50)	1400 (490-2800)
Algeria	1995	Female	45 to 49	18% (6.1-32%)	82% (68-94%)	0.42 (0.14-0.88)	0.69% (0.04-1.8%)	1.2% (0.02-3.4%)	18 (0.32-52)	920 (51-2500)
Algeria	1995	Female	50 to 54	18% (7.1-36%)	82% (64-93%)	0.44 (0.13-0.95)	0.53% (-0.27-1.8%)	0.91% (-0.42-3%)	17 (-7.5-56)	690 (-340-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	1995	Female	55 to 59	19% (7.7-39%)	81% (61-92%)	0.44 (0.12-1)	0.33% (-0.61-1.6%)	0.57% (-0.85-2.7%)	14 (-21-65)	460 (-870-2300)
Algeria	1995	Female	60 to 64	20% (8.1-39%)	80% (61-92%)	0.42 (0.11-1)	0.046% (-1-1.4%)	0.12% (-1.4-2.1%)	4 (-46-73)	72 (-1500-2200)
Algeria	1995	Female	65 to 69	21% (8.5-38%)	79% (62-91%)	0.36 (0.088-0.79)	-0.24% (-1.3-0.84%)	-0.29% (-1.8-1.2%)	-12 (-72-51)	-330 (-1800-1200)
Algeria	1995	Female	70 to 74	20% (7.9-38%)	80% (62-92%)	0.29 (0.072-0.71)	-0.4% (-1.4-0.58%)	-0.52% (-1.9-0.77%)	-23 (-84-35)	-460 (-1700-670)
Algeria	1995	Female	75 to 79	17% (6.1-38%)	83% (62-94%)	0.24 (0.048-0.71)	-0.46% (-1.7-0.42%)	-0.59% (-2.1-0.56%)	-31 (-110-29)	-470 (-1700-430)
Algeria	1995	Female	80 to 84	16% (5.1-37%)	84% (63-95%)	0.21 (0.043-0.63)	-0.42% (-1.6-0.33%)	-0.53% (-2-0.43%)	-23 (-83-19)	-240 (-900-200)
Algeria	1995	Female	85 to 89	17% (5.2-38%)	83% (62-95%)	0.22 (0.043-0.68)	-0.48% (-1.6-0.22%)	-0.61% (-2-0.28%)	-22 (-76-10)	-160 (-560-78)
Algeria	1995	Female	90 to 94	19% (6.1-42%)	81% (58-94%)	0.25 (0.047-0.74)	-0.58% (-1.8-0.17%)	-0.74% (-2.3-0.2%)	-14 (-43-3.7)	-68 (-210-21)
Algeria	1995	Female	95 plus	24% (6.7-48%)	76% (52-93%)	0.3 (0.051-0.89)	-0.66% (-2-0.32%)	-0.87% (-2.5-0.34%)	-4.6 (-13-1.9)	-14 (-44-7.2)
Algeria	1995	Male	All Ages	7.7% (5.2-11%)	92% (89-95%)	0.18 (0.13-0.23)	0.59% (0.4-0.85%)	0.59% (0.23-1.1%)	420 (170-800)	28000 (19000-41000)
Algeria	1995	Male	15 to 19	6.3% (2.6-13%)	94% (87-97%)	0.085 (0.026-0.18)	0.67% (0.39-1%)	0.56% (0.17-1.2%)	11 (3.4-23)	1800 (1000-2800)
Algeria	1995	Male	20 to 24	9.6% (3.8-19%)	90% (81-96%)	0.19 (0.068-0.38)	1.4% (0.83-2.1%)	1.1% (0.42-2.2%)	26 (10-54)	3800 (2300-5900)
Algeria	1995	Male	25 to 29	12% (4.4-23%)	88% (77-96%)	0.29 (0.11-0.55)	1.7% (1-2.6%)	1.7% (0.67-3.2%)	37 (15-68)	4100 (2400-6300)
Algeria	1995	Male	30 to 34	14% (5-27%)	86% (73-95%)	0.38 (0.15-0.67)	1.7% (1.1-2.6%)	2% (0.92-3.6%)	41 (19-76)	3800 (2200-5900)
Algeria	1995	Male	35 to 39	15% (5.8-27%)	85% (73-94%)	0.43 (0.17-0.81)	1.8% (1-2.8%)	2.2% (1.1-4%)	46 (20-84)	3500 (2000-5600)
Algeria	1995	Male	40 to 44	15% (6.8-28%)	85% (72-93%)	0.47 (0.18-0.88)	1.6% (0.88-2.7%)	2.1% (0.91-4%)	46 (19-89)	3000 (1600-5100)
Algeria	1995	Male	45 to 49	15% (5.7-27%)	85% (73-94%)	0.48 (0.15-1.1)	1.5% (0.74-3%)	2% (0.75-4.4%)	40 (14-86)	2100 (1000-4200)
Algeria	1995	Male	50 to 54	14% (6-28%)	86% (72-94%)	0.48 (0.15-1)	1.3% (0.47-2.8%)	1.7% (0.44-3.9%)	36 (9-88)	1700 (610-3700)
Algeria	1995	Male	55 to 59	14% (5.9-27%)	86% (73-94%)	0.46 (0.14-1)	1.1% (0.32-2.6%)	1.4% (0.29-3.5%)	42 (8.1-110)	1600 (450-3800)
Algeria	1995	Male	60 to 64	13% (5.7-26%)	87% (74-94%)	0.42 (0.12-1)	0.84% (0.076-2.3%)	1% (-0.0059-3%)	45 (-0.27-140)	1400 (120-3900)
Algeria	1995	Male	65 to 69	13% (5.8-24%)	87% (76-94%)	0.34 (0.1-0.74)	0.51% (-0.12-1.7%)	0.6% (-0.25-2.1%)	29 (-13-100)	770 (-180-2500)
Algeria	1995	Male	70 to 74	11% (5.1-21%)	89% (79-95%)	0.25 (0.076-0.6)	0.27% (-0.23-1.1%)	0.3% (-0.38-1.4%)	15 (-19-70)	340 (-280-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	1995	Male	75 to 79	9.9% (4.4-19%)	90% (81-96%)	0.19 (0.049-0.5)	0.12% (-0.3-0.75%)	0.11% (-0.45-0.87%)	6.3 (-26-49)	120 (-310-730)
Algeria	1995	Male	80 to 84	8.9% (3.5-17%)	91% (83-96%)	0.16 (0.04-0.39)	0.089% (-0.29-0.53%)	0.073% (-0.4-0.61%)	3.1 (-17-26)	49 (-160-290)
Algeria	1995	Male	85 to 89	8.7% (3.5-18%)	91% (82-97%)	0.15 (0.036-0.39)	0.031% (-0.33-0.38%)	-0.0039% (-0.45-0.41%)	-0.11 (-13-12)	8 (-82-97)
Algeria	1995	Male	90 to 94	9.8% (4-20%)	90% (80-96%)	0.16 (0.036-0.43)	-0.032% (-0.43-0.37%)	-0.094% (-0.6-0.39%)	-1.2 (-7.8-5.1)	-2.6 (-35-30)
Algeria	1995	Male	95 plus	12% (4.5-26%)	88% (74-96%)	0.18 (0.032-0.54)	-0.042% (-0.5-0.48%)	-0.14% (-0.74-0.49%)	-0.5 (-2.5-1.7)	-0.6 (-7.2-6.8)
Algeria	2000	Female	All Ages	11% (6.4-16%)	89% (84-94%)	0.19 (0.15-0.24)	0.28% (0.047-0.54%)	0.041% (-0.58-0.76%)	27 (-360-480)	12000 (1900-24000)
Algeria	2000	Female	15 to 19	9.1% (3-20%)	91% (80-97%)	0.14 (0.042-0.32)	0.46% (0.27-0.74%)	0.52% (0.18-1.1%)	5.4 (1.9-12)	1100 (610-1800)
Algeria	2000	Female	20 to 24	12% (3.5-24%)	88% (76-96%)	0.22 (0.072-0.47)	0.84% (0.53-1.3%)	0.76% (0.27-1.7%)	8.7 (3-19)	2000 (1200-3100)
Algeria	2000	Female	25 to 29	13% (3.8-28%)	87% (72-96%)	0.27 (0.085-0.54)	0.92% (0.58-1.3%)	0.99% (0.37-1.9%)	11 (4.1-22)	2000 (1200-3100)
Algeria	2000	Female	30 to 34	15% (3.7-31%)	85% (69-96%)	0.32 (0.1-0.62)	0.9% (0.54-1.4%)	1.2% (0.49-2.3%)	15 (5.9-28)	1900 (1100-3000)
Algeria	2000	Female	35 to 39	16% (4.4-31%)	84% (69-96%)	0.38 (0.12-0.72)	0.85% (0.44-1.5%)	1.4% (0.47-2.8%)	20 (6.8-42)	1800 (910-3000)
Algeria	2000	Female	40 to 44	17% (6-32%)	83% (68-94%)	0.43 (0.15-0.82)	0.85% (0.31-1.6%)	1.6% (0.43-3.4%)	26 (7-56)	1600 (570-3200)
Algeria	2000	Female	45 to 49	18% (6.1-33%)	82% (67-94%)	0.46 (0.15-0.95)	0.77% (0.079-1.9%)	1.4% (0.19-3.6%)	28 (3.8-74)	1400 (160-3500)
Algeria	2000	Female	50 to 54	19% (7.1-36%)	81% (64-93%)	0.48 (0.15-1)	0.63% (-0.2-1.9%)	1.1% (-0.26-3.6%)	22 (-5-69)	920 (-310-3000)
Algeria	2000	Female	55 to 59	19% (7.4-38%)	81% (62-93%)	0.48 (0.12-1.1)	0.44% (-0.52-1.9%)	0.77% (-0.76-3.1%)	19 (-17-77)	640 (-730-2800)
Algeria	2000	Female	60 to 64	20% (7.7-40%)	80% (60-92%)	0.46 (0.12-1.1)	0.14% (-0.99-1.7%)	0.26% (-1.4-2.5%)	9.8 (-49-95)	240 (-1600-2800)
Algeria	2000	Female	65 to 69	21% (8.9-38%)	79% (62-91%)	0.4 (0.1-0.88)	-0.17% (-1.2-1%)	-0.19% (-1.8-1.5%)	-9 (-81-70)	-280 (-2100-1700)
Algeria	2000	Female	70 to 74	19% (8.4-37%)	81% (63-92%)	0.31 (0.083-0.77)	-0.35% (-1.4-0.61%)	-0.46% (-1.9-0.89%)	-25 (-100-47)	-520 (-2100-890)
Algeria	2000	Female	75 to 79	17% (6.5-35%)	83% (65-93%)	0.25 (0.053-0.71)	-0.42% (-1.5-0.41%)	-0.55% (-1.9-0.54%)	-30 (-110-31)	-460 (-1600-470)
Algeria	2000	Female	80 to 84	16% (5.5-36%)	84% (64-95%)	0.23 (0.049-0.66)	-0.38% (-1.4-0.4%)	-0.49% (-1.8-0.49%)	-25 (-95-25)	-270 (-1000-280)
Algeria	2000	Female	85 to 89	16% (5.3-37%)	84% (63-95%)	0.23 (0.048-0.73)	-0.44% (-1.6-0.28%)	-0.56% (-1.9-0.33%)	-25 (-90-14)	-180 (-650-120)
Algeria	2000	Female	90 to 94	19% (6.3-42%)	81% (58-94%)	0.26 (0.051-0.78)	-0.52% (-1.7-0.26%)	-0.68% (-2.2-0.28%)	-17 (-55-7.3)	-84 (-270-42)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	2000	Female	95 plus	23% (6.9-48%)	77% (52-93%)	0.32 (0.054-0.96)	-0.6% (-2-0.56%)	-0.8% (-2.5-0.59%)	-6.9 (-22-5)	-21 (-69-20)
Algeria	2000	Male	All Ages	8.1% (5.6-12%)	92% (88-94%)	0.21 (0.16-0.26)	0.72% (0.48-1%)	0.71% (0.3-1.3%)	540 (230-1000)	34000 (23000-50000)
Algeria	2000	Male	15 to 19	6.3% (2.6-12%)	94% (88-97%)	0.093 (0.028-0.2)	0.73% (0.44-1.2%)	0.68% (0.23-1.5%)	12 (3.9-26)	2000 (1100-3200)
Algeria	2000	Male	20 to 24	9.8% (3.8-19%)	90% (81-96%)	0.21 (0.075-0.41)	1.6% (0.99-2.4%)	1.4% (0.53-2.7%)	31 (12-65)	4500 (2700-6900)
Algeria	2000	Male	25 to 29	12% (4.8-24%)	88% (76-95%)	0.33 (0.12-0.63)	1.9% (1.2-2.9%)	2.1% (0.88-3.9%)	44 (19-82)	4800 (2900-7400)
Algeria	2000	Male	30 to 34	14% (5-27%)	86% (73-95%)	0.42 (0.17-0.76)	1.9% (1.2-2.9%)	2.4% (1.1-4.2%)	47 (21-85)	4300 (2600-6800)
Algeria	2000	Male	35 to 39	15% (5.8-28%)	85% (72-94%)	0.48 (0.18-0.9)	1.9% (1.1-3.1%)	2.6% (1.2-4.7%)	54 (25-98)	4100 (2400-6600)
Algeria	2000	Male	40 to 44	15% (6.8-27%)	85% (73-93%)	0.51 (0.2-0.98)	1.8% (0.99-3%)	2.5% (1.1-4.7%)	55 (24-100)	3600 (1900-6000)
Algeria	2000	Male	45 to 49	15% (5.8-28%)	85% (72-94%)	0.53 (0.17-1.1)	1.7% (0.83-3.2%)	2.3% (0.9-4.7%)	61 (23-130)	3300 (1600-6200)
Algeria	2000	Male	50 to 54	14% (6-28%)	86% (72-94%)	0.52 (0.17-1.1)	1.5% (0.56-3%)	2% (0.54-4.3%)	49 (14-110)	2200 (850-4700)
Algeria	2000	Male	55 to 59	14% (5.7-26%)	86% (74-94%)	0.5 (0.15-1.1)	1.3% (0.39-2.8%)	1.7% (0.36-4%)	50 (11-120)	1900 (580-4400)
Algeria	2000	Male	60 to 64	13% (5.4-25%)	87% (75-95%)	0.44 (0.13-1.1)	0.96% (0.14-2.6%)	1.2% (0.074-3.4%)	53 (3.3-150)	1700 (230-4400)
Algeria	2000	Male	65 to 69	12% (5.5-23%)	88% (77-94%)	0.36 (0.11-0.79)	0.6% (-0.041-1.8%)	0.73% (-0.14-2.4%)	42 (-7.9-140)	1100 (-74-3500)
Algeria	2000	Male	70 to 74	11% (5-20%)	89% (80-95%)	0.27 (0.078-0.65)	0.33% (-0.2-1.3%)	0.38% (-0.31-1.5%)	24 (-21-100)	530 (-310-2000)
Algeria	2000	Male	75 to 79	9.9% (4.1-20%)	90% (80-96%)	0.21 (0.05-0.55)	0.17% (-0.28-0.87%)	0.17% (-0.4-1%)	10 (-26-65)	190 (-310-1000)
Algeria	2000	Male	80 to 84	8.9% (3.4-18%)	91% (82-97%)	0.18 (0.041-0.44)	0.12% (-0.26-0.64%)	0.11% (-0.37-0.73%)	5.8 (-19-37)	82 (-170-420)
Algeria	2000	Male	85 to 89	8.8% (3.2-18%)	91% (82-97%)	0.16 (0.038-0.44)	0.059% (-0.31-0.5%)	0.028% (-0.43-0.57%)	1 (-16-21)	20 (-100-170)
Algeria	2000	Male	90 to 94	9.9% (3.9-20%)	90% (80-96%)	0.18 (0.042-0.49)	-0.005% (-0.41-0.5%)	-0.064% (-0.57-0.52%)	-1.1 (-9.7-8.7)	-0.54 (-42-51)
Algeria	2000	Male	95 plus	12% (4.4-25%)	88% (75-96%)	0.2 (0.035-0.56)	-0.011% (-0.5-0.67%)	-0.11% (-0.72-0.7%)	-0.54 (-3.6-3.3)	-0.22 (-9.9-13)
Algeria	2005	Female	All Ages	11% (6.8-16%)	89% (84-93%)	0.24 (0.18-0.31)	0.39% (0.12-0.72%)	0.21% (-0.52-1.1%)	140 (-350-720)	17000 (5000-32000)
Algeria	2005	Female	15 to 19	8.7% (2.9-19%)	91% (81-97%)	0.16 (0.049-0.37)	0.49% (0.28-0.78%)	0.63% (0.23-1.4%)	5.6 (2-12)	1100 (630-1800)
Algeria	2005	Female	20 to 24	11% (3.6-24%)	89% (76-96%)	0.26 (0.083-0.52)	0.9% (0.57-1.4%)	0.94% (0.37-2%)	9.8 (3.8-21)	2300 (1400-3600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	2005	Female	25 to 29	13% (3.8-27%)	87% (73-96%)	0.32 (0.1-0.63)	0.99% (0.64-1.4%)	1.2% (0.46-2.3%)	13 (5-25)	2400 (1500-3600)
Algeria	2005	Female	30 to 34	15% (4.1-30%)	85% (70-96%)	0.38 (0.13-0.73)	0.98% (0.61-1.5%)	1.5% (0.63-2.8%)	18 (7.5-33)	2300 (1400-3400)
Algeria	2005	Female	35 to 39	17% (4.5-32%)	83% (68-95%)	0.45 (0.15-0.87)	0.96% (0.52-1.6%)	1.8% (0.66-3.5%)	25 (9-49)	2200 (1100-3600)
Algeria	2005	Female	40 to 44	17% (6.2-33%)	83% (67-94%)	0.5 (0.19-0.97)	1% (0.42-1.9%)	2.1% (0.68-4%)	36 (12-70)	2200 (910-4100)
Algeria	2005	Female	45 to 49	18% (6.3-33%)	82% (67-94%)	0.54 (0.19-1.1)	0.99% (0.27-2.3%)	1.9% (0.51-4.6%)	41 (9.8-97)	2000 (520-4800)
Algeria	2005	Female	50 to 54	18% (6.7-36%)	82% (64-93%)	0.55 (0.17-1.2)	0.88% (-0.013-2.3%)	1.6% (0.14-4.2%)	44 (3.5-120)	1800 (-29-4800)
Algeria	2005	Female	55 to 59	19% (7-37%)	81% (63-93%)	0.55 (0.14-1.2)	0.72% (-0.27-2.3%)	1.3% (-0.3-3.7%)	36 (-8.2-110)	1200 (-440-3800)
Algeria	2005	Female	60 to 64	20% (7.5-38%)	80% (62-92%)	0.52 (0.13-1.2)	0.4% (-0.71-2.1%)	0.66% (-0.92-3.2%)	25 (-36-120)	690 (-1300-3600)
Algeria	2005	Female	65 to 69	21% (8.7-37%)	79% (63-91%)	0.46 (0.12-1)	0.055% (-1-1.4%)	0.14% (-1.5-2.1%)	7.4 (-7.5-110)	110 (-2000-2700)
Algeria	2005	Female	70 to 74	19% (8.4-35%)	81% (65-92%)	0.36 (0.095-0.88)	-0.19% (-1.3-0.98%)	-0.22% (-1.7-1.3%)	-15 (-110-91)	-340 (-2300-1800)
Algeria	2005	Female	75 to 79	17% (6.7-34%)	83% (66-93%)	0.3 (0.064-0.82)	-0.3% (-1.3-0.73%)	-0.39% (-1.7-0.95%)	-28 (-120-67)	-430 (-1800-1000)
Algeria	2005	Female	80 to 84	16% (5.4-35%)	84% (65-95%)	0.27 (0.056-0.76)	-0.28% (-1.3-0.64%)	-0.37% (-1.6-0.8%)	-23 (-100-52)	-250 (-1100-610)
Algeria	2005	Female	85 to 89	16% (5.5-37%)	84% (63-94%)	0.28 (0.056-0.84)	-0.35% (-1.4-0.55%)	-0.46% (-1.7-0.66%)	-24 (-89-34)	-170 (-690-270)
Algeria	2005	Female	90 to 94	19% (6.4-41%)	81% (59-94%)	0.32 (0.064-0.96)	-0.43% (-1.6-0.64%)	-0.58% (-2.1-0.75%)	-18 (-65-23)	-87 (-320-130)
Algeria	2005	Female	95 plus	23% (7.3-49%)	77% (51-93%)	0.37 (0.066-1.2)	-0.48% (-1.8-1.1%)	-0.67% (-2.3-1.2%)	-7.7 (-27-14)	-23 (-88-52)
Algeria	2005	Male	All Ages	8.4% (5.7-12%)	92% (88-94%)	0.26 (0.2-0.33)	0.89% (0.6-1.2%)	0.93% (0.45-1.7%)	720 (340-1300)	43000 (29000-60000)
Algeria	2005	Male	15 to 19	6.2% (2.6-13%)	94% (87-97%)	0.11 (0.033-0.24)	0.79% (0.48-1.3%)	0.81% (0.27-1.8%)	12 (3.9-26)	2000 (1200-3300)
Algeria	2005	Male	20 to 24	9.7% (3.7-19%)	90% (81-96%)	0.24 (0.086-0.49)	1.7% (1.1-2.6%)	1.7% (0.65-3.3%)	34 (13-69)	5000 (3000-7700)
Algeria	2005	Male	25 to 29	12% (4.9-23%)	88% (77-95%)	0.38 (0.15-0.73)	2% (1.3-3.1%)	2.5% (1.1-4.7%)	49 (21-91)	5500 (3400-8600)
Algeria	2005	Male	30 to 34	14% (5-26%)	86% (74-95%)	0.48 (0.2-0.87)	2.1% (1.3-3.2%)	2.8% (1.3-5%)	53 (24-96)	4900 (2900-7500)
Algeria	2005	Male	35 to 39	15% (5.5-27%)	85% (73-94%)	0.54 (0.2-1)	2.1% (1.2-3.3%)	3.1% (1.4-5.4%)	60 (27-110)	4600 (2700-7500)
Algeria	2005	Male	40 to 44	15% (6.5-27%)	85% (73-94%)	0.58 (0.23-1.1)	2% (1.1-3.4%)	2.9% (1.3-5.4%)	67 (29-120)	4400 (2400-7300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	2005	Male	45 to 49	15% (6-28%)	85% (72-94%)	0.61 (0.2-1.2)	2% (0.96-3.6%)	2.9% (1.2-5.7%)	77 (32-160)	4200 (2100-7800)
Algeria	2005	Male	50 to 54	14% (6.1-28%)	86% (72-94%)	0.61 (0.2-1.2)	1.8% (0.69-3.4%)	2.5% (0.85-5%)	85 (27-170)	3800 (1500-7600)
Algeria	2005	Male	55 to 59	14% (5.8-27%)	86% (73-94%)	0.58 (0.18-1.3)	1.6% (0.55-3.2%)	2.1% (0.65-4.7%)	71 (22-160)	2700 (930-5700)
Algeria	2005	Male	60 to 64	13% (5.5-26%)	87% (74-94%)	0.52 (0.16-1.2)	1.2% (0.27-3%)	1.6% (0.27-4.2%)	69 (12-180)	2200 (480-5400)
Algeria	2005	Male	65 to 69	13% (5.6-24%)	87% (76-94%)	0.43 (0.13-0.94)	0.83% (0.086-2.3%)	1% (0.036-3%)	60 (2-170)	1600 (170-4400)
Algeria	2005	Male	70 to 74	11% (5-21%)	89% (79-95%)	0.32 (0.089-0.73)	0.5% (-0.068-1.6%)	0.6% (-0.16-2.1%)	44 (-11-150)	920 (-120-2900)
Algeria	2005	Male	75 to 79	9.8% (4.1-19%)	90% (81-96%)	0.24 (0.061-0.65)	0.29% (-0.15-1.2%)	0.32% (-0.25-1.5%)	23 (-18-110)	390 (-210-1700)
Algeria	2005	Male	80 to 84	8.7% (3.5-17%)	91% (83-96%)	0.2 (0.05-0.51)	0.2% (-0.18-0.86%)	0.21% (-0.28-1%)	13 (-18-65)	170 (-160-740)
Algeria	2005	Male	85 to 89	8.4% (3.4-17%)	92% (83-97%)	0.18 (0.044-0.46)	0.13% (-0.24-0.72%)	0.11% (-0.33-0.81%)	4.9 (-16-37)	54 (-96-300)
Algeria	2005	Male	90 to 94	9.5% (4-20%)	90% (80-96%)	0.19 (0.047-0.51)	0.059% (-0.33-0.71%)	0.01% (-0.49-0.79%)	0.23 (-11-18)	8.4 (-48-100)
Algeria	2005	Male	95 plus	12% (4.3-25%)	88% (75-96%)	0.23 (0.04-0.68)	0.061% (-0.45-0.87%)	-0.03% (-0.66-0.92%)	-0.19 (-4.1-5.7)	1.6 (-11-22)
Algeria	2010	Female	All Ages	12% (7.2-17%)	88% (83-93%)	0.26 (0.19-0.35)	0.41% (0.11-0.8%)	0.18% (-0.59-1.2%)	140 (-420-880)	18000 (4700-36000)
Algeria	2010	Female	15 to 19	8.8% (2.8-20%)	91% (80-97%)	0.17 (0.047-0.39)	0.5% (0.29-0.8%)	0.67% (0.24-1.5%)	5.1 (1.8-11)	1100 (620-1800)
Algeria	2010	Female	20 to 24	11% (3.3-25%)	89% (75-97%)	0.26 (0.074-0.59)	0.92% (0.56-1.4%)	0.98% (0.37-2.1%)	9.5 (3.6-20)	2300 (1300-3700)
Algeria	2010	Female	25 to 29	13% (3.6-27%)	87% (73-96%)	0.32 (0.092-0.66)	1% (0.65-1.5%)	1.2% (0.46-2.5%)	14 (5.1-27)	2700 (1600-4000)
Algeria	2010	Female	30 to 34	14% (3.8-30%)	86% (70-96%)	0.37 (0.12-0.72)	0.98% (0.62-1.5%)	1.5% (0.64-2.9%)	19 (7.9-36)	2600 (1500-3900)
Algeria	2010	Female	35 to 39	16% (4.6-31%)	84% (69-95%)	0.44 (0.15-0.83)	0.93% (0.51-1.5%)	1.8% (0.68-3.4%)	25 (9.4-49)	2300 (1300-3700)
Algeria	2010	Female	40 to 44	17% (6.2-32%)	83% (68-94%)	0.48 (0.18-0.93)	0.95% (0.42-1.8%)	2% (0.74-4.1%)	36 (13-74)	2300 (990-4400)
Algeria	2010	Female	45 to 49	17% (6-33%)	83% (67-94%)	0.52 (0.18-1.1)	0.92% (0.25-2.1%)	1.9% (0.5-4.3%)	43 (11-100)	2200 (600-5100)
Algeria	2010	Female	50 to 54	18% (6.7-36%)	82% (64-93%)	0.54 (0.17-1.1)	0.82% (-0.039-2.2%)	1.6% (0.067-4%)	45 (1.9-120)	1900 (-97-5200)
Algeria	2010	Female	55 to 59	19% (7.2-37%)	81% (63-93%)	0.54 (0.15-1.2)	0.67% (-0.35-2.1%)	1.2% (-0.35-3.6%)	46 (-14-140)	1600 (-820-5100)
Algeria	2010	Female	60 to 64	20% (7.6-40%)	80% (60-92%)	0.52 (0.14-1.3)	0.36% (-0.77-2%)	0.63% (-0.98-3.1%)	27 (-42-140)	720 (-1500-4100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	2010	Female	65 to 69	20% (8.4-39%)	80% (61-92%)	0.46 (0.12-1)	0.048% (-1-1.5%)	0.15% (-1.4-2.2%)	8 (-74-120)	100 (-2000-2900)
Algeria	2010	Female	70 to 74	19% (7.8-36%)	81% (64-92%)	0.36 (0.09-0.97)	-0.18% (-1.3-1.1%)	-0.2% (-1.7-1.5%)	-15 (-130-110)	-360 (-2600-2300)
Algeria	2010	Female	75 to 79	17% (6.5-34%)	83% (66-94%)	0.29 (0.065-0.85)	-0.28% (-1.2-0.79%)	-0.37% (-1.6-1%)	-32 (-140-89)	-500 (-2200-1400)
Algeria	2010	Female	80 to 84	16% (5.3-33%)	84% (67-95%)	0.26 (0.056-0.75)	-0.28% (-1.2-0.62%)	-0.36% (-1.5-0.75%)	-31 (-130-65)	-340 (-1400-720)
Algeria	2010	Female	85 to 89	16% (5.2-35%)	84% (65-95%)	0.26 (0.054-0.77)	-0.34% (-1.2-0.55%)	-0.45% (-1.6-0.63%)	-31 (-110-43)	-220 (-810-330)
Algeria	2010	Female	90 to 94	19% (6.2-40%)	81% (60-94%)	0.3 (0.06-0.9)	-0.43% (-1.5-0.59%)	-0.57% (-1.9-0.67%)	-21 (-71-25)	-100 (-350-140)
Algeria	2010	Female	95 plus	23% (7-48%)	77% (52-93%)	0.37 (0.063-1.1)	-0.5% (-1.8-0.88%)	-0.69% (-2.3-1%)	-10 (-34-15)	-31 (-110-55)
Algeria	2010	Male	All Ages	9% (6.1-13%)	91% (87-94%)	0.28 (0.21-0.37)	0.97% (0.64-1.4%)	0.94% (0.41-1.8%)	770 (340-1500)	45000 (30000-66000)
Algeria	2010	Male	15 to 19	6.4% (2.6-13%)	94% (87-97%)	0.11 (0.031-0.25)	0.81% (0.48-1.3%)	0.84% (0.28-1.9%)	10 (3.5-22)	1800 (1000-2900)
Algeria	2010	Male	20 to 24	9.7% (3.7-19%)	90% (81-96%)	0.24 (0.082-0.5)	1.7% (1.1-2.7%)	1.7% (0.65-3.5%)	31 (12-65)	4900 (3000-7800)
Algeria	2010	Male	25 to 29	12% (4.7-23%)	88% (77-95%)	0.38 (0.14-0.74)	2% (1.3-3%)	2.5% (1-4.7%)	49 (20-94)	5900 (3600-9100)
Algeria	2010	Male	30 to 34	14% (5-27%)	86% (73-95%)	0.48 (0.2-0.9)	2% (1.3-3.1%)	2.8% (1.3-5.1%)	55 (25-99)	5400 (3400-8400)
Algeria	2010	Male	35 to 39	15% (5.7-28%)	85% (72-94%)	0.55 (0.2-1)	2.1% (1.3-3.3%)	3.1% (1.5-5.6%)	60 (28-110)	4900 (2900-7600)
Algeria	2010	Male	40 to 44	15% (6.7-27%)	85% (73-93%)	0.58 (0.23-1.1)	2% (1.1-3.3%)	3% (1.3-5.5%)	66 (30-120)	4500 (2500-7600)
Algeria	2010	Male	45 to 49	15% (5.9-28%)	85% (72-94%)	0.6 (0.2-1.2)	1.9% (0.97-3.5%)	2.9% (1.2-5.7%)	80 (35-160)	4400 (2300-8100)
Algeria	2010	Male	50 to 54	14% (5.9-28%)	86% (72-94%)	0.59 (0.19-1.2)	1.7% (0.71-3.3%)	2.5% (0.89-5%)	86 (30-170)	4000 (1700-7700)
Algeria	2010	Male	55 to 59	13% (6-26%)	87% (74-94%)	0.57 (0.18-1.2)	1.5% (0.54-3.2%)	2.1% (0.68-4.7%)	94 (30-210)	3600 (1300-7700)
Algeria	2010	Male	60 to 64	13% (5.4-25%)	87% (75-95%)	0.51 (0.16-1.2)	1.2% (0.3-2.9%)	1.6% (0.33-4%)	78 (15-190)	2500 (630-5900)
Algeria	2010	Male	65 to 69	13% (5.7-23%)	87% (77-94%)	0.42 (0.13-0.93)	0.82% (0.11-2.2%)	1.1% (0.067-3%)	61 (3.8-180)	1600 (200-4400)
Algeria	2010	Male	70 to 74	11% (5.1-21%)	89% (79-95%)	0.31 (0.093-0.74)	0.5% (-0.037-1.6%)	0.61% (-0.13-2.1%)	46 (-9.7-160)	970 (-68-3200)
Algeria	2010	Male	75 to 79	9.8% (4.1-19%)	90% (81-96%)	0.24 (0.059-0.68)	0.29% (-0.14-1.2%)	0.33% (-0.25-1.5%)	29 (-22-130)	500 (-250-2100)
Algeria	2010	Male	80 to 84	8.7% (3.5-17%)	91% (83-97%)	0.2 (0.049-0.51)	0.2% (-0.18-0.83%)	0.21% (-0.28-0.99%)	18 (-25-83)	230 (-210-960)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	2010	Male	85 to 89	8.5% (3.5-17%)	91% (83-97%)	0.18 (0.049-0.47)	0.13% (-0.21-0.66%)	0.11% (-0.33-0.72%)	6.5 (-20-44)	72 (-120-370)
Algeria	2010	Male	90 to 94	9.7% (4-20%)	90% (80-96%)	0.2 (0.049-0.52)	0.06% (-0.34-0.66%)	0.0096% (-0.5-0.7%)	0.28 (-15-21)	11 (-65-130)
Algeria	2010	Male	95 plus	12% (4.5-25%)	88% (75-96%)	0.23 (0.042-0.65)	0.059% (-0.42-0.78%)	-0.037% (-0.66-0.85%)	-0.34 (-5.9-7.7)	2.2 (-16-30)
Algeria	2016	Female	All Ages	12% (7.1-17%)	88% (83-93%)	0.26 (0.15-0.4)	0.41% (0.043-0.93%)	0.13% (-0.74-1.3%)	110 (-630-1100)	19000 (2000-44000)
Algeria	2016	Female	15 to 19	8.6% (2.5-20%)	91% (80-98%)	0.16 (0.038-0.41)	0.5% (0.29-0.84%)	0.68% (0.23-1.7%)	3.8 (1.3-9.4)	820 (450-1400)
Algeria	2016	Female	20 to 24	11% (3.1-24%)	89% (76-97%)	0.25 (0.058-0.58)	0.92% (0.56-1.4%)	1% (0.33-2.3%)	7.8 (2.6-19)	2100 (1200-3300)
Algeria	2016	Female	25 to 29	12% (3.2-27%)	88% (73-97%)	0.29 (0.074-0.68)	0.99% (0.63-1.4%)	1.2% (0.43-2.7%)	12 (4.4-27)	2700 (1600-4200)
Algeria	2016	Female	30 to 34	14% (3.5-30%)	86% (70-97%)	0.34 (0.1-0.73)	0.96% (0.6-1.5%)	1.5% (0.58-3%)	19 (7.3-38)	2900 (1700-4600)
Algeria	2016	Female	35 to 39	16% (4.1-31%)	84% (69-96%)	0.41 (0.12-0.86)	0.9% (0.49-1.5%)	1.8% (0.67-3.6%)	28 (10-57)	2700 (1500-4600)
Algeria	2016	Female	40 to 44	16% (5.8-32%)	84% (68-94%)	0.46 (0.15-0.95)	0.9% (0.37-1.7%)	2% (0.64-4.1%)	39 (12-82)	2600 (1000-4900)
Algeria	2016	Female	45 to 49	17% (5.8-33%)	83% (67-94%)	0.51 (0.15-1.1)	0.87% (0.16-2.1%)	1.8% (0.38-4.5%)	47 (10-120)	2400 (460-5900)
Algeria	2016	Female	50 to 54	18% (6.8-34%)	82% (66-93%)	0.52 (0.15-1.2)	0.77% (-0.1-2.2%)	1.5% (0.016-4.2%)	51 (0.47-140)	2200 (-290-6300)
Algeria	2016	Female	55 to 59	19% (7.3-37%)	81% (63-93%)	0.53 (0.14-1.2)	0.62% (-0.43-2.2%)	1.2% (-0.44-3.9%)	49 (-19-160)	1700 (-1100-6000)
Algeria	2016	Female	60 to 64	20% (7.9-38%)	80% (62-92%)	0.51 (0.13-1.3)	0.32% (-0.77-2.2%)	0.6% (-0.99-3.5%)	34 (-58-190)	910 (-2200-6100)
Algeria	2016	Female	65 to 69	21% (8.8-37%)	79% (63-91%)	0.45 (0.12-1)	0.036% (-0.98-1.6%)	0.14% (-1.4-2.5%)	8.5 (-84-150)	87 (-2300-3800)
Algeria	2016	Female	70 to 74	19% (8.2-34%)	81% (66-92%)	0.36 (0.091-0.93)	-0.17% (-1.2-1.2%)	-0.2% (-1.6-1.6%)	-15 (-120-120)	-380 (-2600-2600)
Algeria	2016	Female	75 to 79	17% (6.5-34%)	83% (66-94%)	0.29 (0.067-0.83)	-0.28% (-1.2-0.85%)	-0.37% (-1.6-1.1%)	-37 (-160-110)	-580 (-2500-1800)
Algeria	2016	Female	80 to 84	16% (5.6-35%)	84% (65-94%)	0.27 (0.056-0.8)	-0.3% (-1.3-0.72%)	-0.38% (-1.7-0.91%)	-45 (-190-110)	-480 (-2100-1200)
Algeria	2016	Female	85 to 89	17% (5.6-37%)	83% (63-94%)	0.27 (0.055-0.87)	-0.36% (-1.4-0.56%)	-0.47% (-1.8-0.71%)	-49 (-190-75)	-360 (-1500-600)
Algeria	2016	Female	90 to 94	19% (7.1-40%)	81% (60-93%)	0.31 (0.062-0.93)	-0.43% (-1.6-0.72%)	-0.57% (-2-0.87%)	-33 (-110-49)	-160 (-570-280)
Algeria	2016	Female	95 plus	23% (7.9-48%)	77% (52-92%)	0.37 (0.063-1.2)	-0.48% (-1.8-1.1%)	-0.66% (-2.4-1.3%)	-13 (-46-25)	-38 (-150-90)
Algeria	2016	Male	All Ages	9.3% (6.3-13%)	91% (87-94%)	0.29 (0.17-0.44)	1% (0.65-1.6%)	0.93% (0.33-1.9%)	870 (300-1800)	50000 (30000-79000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Algeria	2016	Male	15 to 19	6.2% (2.6-12%)	94% (88-97%)	0.1 (0.029-0.25)	0.81% (0.49-1.3%)	0.84% (0.28-1.9%)	7.9 (2.7-18)	1400 (830-2400)
Algeria	2016	Male	20 to 24	9.6% (3.6-19%)	90% (81-96%)	0.24 (0.065-0.54)	1.8% (1.1-2.8%)	1.7% (0.62-3.8%)	27 (9.5-62)	4400 (2600-7000)
Algeria	2016	Male	25 to 29	12% (4.6-23%)	88% (77-95%)	0.37 (0.11-0.79)	2.1% (1.3-3.1%)	2.5% (0.97-4.9%)	48 (18-96)	5900 (3500-9600)
Algeria	2016	Male	30 to 34	14% (4.7-26%)	86% (74-95%)	0.47 (0.17-0.9)	2% (1.2-3.1%)	2.8% (1.2-5.2%)	60 (26-110)	6100 (3700-10000)
Algeria	2016	Male	35 to 39	15% (5.5-27%)	85% (73-94%)	0.53 (0.19-1)	2% (1.2-3.3%)	3% (1.4-5.6%)	69 (31-130)	5800 (3400-9500)
Algeria	2016	Male	40 to 44	15% (6.6-27%)	85% (73-93%)	0.56 (0.2-1.1)	1.9% (1.1-3.2%)	2.9% (1.3-5.6%)	72 (31-140)	5100 (2900-8700)
Algeria	2016	Male	45 to 49	15% (5.9-27%)	85% (73-94%)	0.58 (0.18-1.2)	1.8% (0.89-3.3%)	2.8% (1.1-5.5%)	85 (33-170)	4800 (2300-8900)
Algeria	2016	Male	50 to 54	14% (6-27%)	86% (73-94%)	0.57 (0.18-1.2)	1.6% (0.64-3.2%)	2.4% (0.79-5%)	96 (32-210)	4500 (1800-9200)
Algeria	2016	Male	55 to 59	14% (5.9-26%)	86% (74-94%)	0.55 (0.17-1.3)	1.4% (0.5-3.1%)	2% (0.59-4.6%)	100 (29-230)	4000 (1400-8600)
Algeria	2016	Male	60 to 64	13% (5.5-25%)	87% (75-94%)	0.49 (0.14-1.2)	1.1% (0.28-2.8%)	1.5% (0.3-4%)	110 (21-280)	3500 (860-8800)
Algeria	2016	Male	65 to 69	12% (5.4-23%)	88% (77-95%)	0.4 (0.11-0.94)	0.79% (0.09-2.2%)	1% (0.065-3%)	75 (4.6-220)	2000 (230-5500)
Algeria	2016	Male	70 to 74	11% (4.8-20%)	89% (80-95%)	0.3 (0.082-0.74)	0.49% (-0.04-1.6%)	0.61% (-0.11-2.1%)	50 (-9.1-170)	1100 (-81-3600)
Algeria	2016	Male	75 to 79	9.6% (3.9-19%)	90% (81-96%)	0.23 (0.054-0.66)	0.3% (-0.15-1.2%)	0.34% (-0.24-1.5%)	33 (-23-150)	570 (-280-2400)
Algeria	2016	Male	80 to 84	8.6% (3.3-17%)	91% (83-97%)	0.19 (0.046-0.52)	0.21% (-0.17-0.87%)	0.22% (-0.27-1%)	24 (-30-110)	310 (-260-1300)
Algeria	2016	Male	85 to 89	8.4% (3.3-17%)	92% (83-97%)	0.18 (0.045-0.5)	0.14% (-0.22-0.64%)	0.12% (-0.33-0.72%)	12 (-31-70)	120 (-200-590)
Algeria	2016	Male	90 to 94	9.6% (3.8-20%)	90% (80-96%)	0.19 (0.044-0.54)	0.07% (-0.31-0.68%)	0.024% (-0.45-0.73%)	1.2 (-22-37)	21 (-97-210)
Algeria	2016	Male	95 plus	12% (4.2-26%)	88% (74-96%)	0.22 (0.039-0.68)	0.07% (-0.41-0.85%)	-0.02% (-0.6-0.92%)	-0.28 (-8.7-14)	4.2 (-24-53)
Bahrain	1990	Female	All Ages	13% (7.2-20%)	87% (80-93%)	0.75 (0.56-0.95)	1.7% (0.85-2.8%)	3% (0.72-6.1%)	25 (6.1-50)	920 (450-1500)
Bahrain	1990	Female	15 to 19	11% (3.1-24%)	89% (76-97%)	0.55 (0.15-1.3)	1.1% (0.51-2.1%)	2.3% (0.89-5.1%)	0.21 (0.08-0.47)	25 (12-48)
Bahrain	1990	Female	20 to 24	14% (4-30%)	86% (70-96%)	0.89 (0.25-2)	1.7% (0.91-2.8%)	3.5% (1.3-7.1%)	0.38 (0.14-0.74)	48 (26-81)
Bahrain	1990	Female	25 to 29	16% (4.3-33%)	84% (67-96%)	1.1 (0.32-2.1)	2% (1.1-3.3%)	4.2% (1.7-7.7%)	0.79 (0.3-1.5)	80 (42-130)
Bahrain	1990	Female	30 to 34	18% (4.5-36%)	82% (64-96%)	1.2 (0.45-2.2)	2% (1.2-3.1%)	4.8% (2.1-7.9%)	0.83 (0.35-1.4)	79 (45-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	1990	Female	35 to 39	19% (5.2-37%)	81% (63-95%)	1.5 (0.52-2.6)	2.3% (1.1-3.7%)	5.3% (2.2-8.7%)	0.94 (0.39-1.6)	72 (36-110)
Bahrain	1990	Female	40 to 44	20% (7.2-36%)	80% (64-93%)	1.6 (0.62-3.1)	2.7% (1.2-4.8%)	5.9% (2.5-10%)	0.88 (0.38-1.6)	55 (25-96)
Bahrain	1990	Female	45 to 49	21% (7.4-38%)	79% (62-93%)	1.8 (0.59-3.7)	3.2% (1.1-6.4%)	5.9% (2.1-12%)	1.2 (0.44-2.5)	63 (22-130)
Bahrain	1990	Female	50 to 54	22% (8.9-39%)	78% (61-91%)	1.9 (0.61-4)	4% (0.95-8%)	5.9% (1.5-12%)	2.6 (0.64-5.1)	100 (26-210)
Bahrain	1990	Female	55 to 59	22% (9.6-42%)	78% (58-90%)	1.9 (0.57-4.2)	4.3% (0.78-9.3%)	5.8% (1.1-12%)	3.1 (0.57-6.8)	110 (19-240)
Bahrain	1990	Female	60 to 64	24% (9.8-45%)	76% (55-90%)	1.8 (0.5-4.4)	3.9% (0.22-9.4%)	4.9% (0.33-12%)	4.1 (0.27-9.8)	120 (6.6-280)
Bahrain	1990	Female	65 to 69	25% (11-43%)	75% (57-89%)	1.6 (0.45-3.8)	3.5% (-0.034-8.6%)	4.2% (-0.061-11%)	3.1 (-0.044-7.9)	73 (-0.7-190)
Bahrain	1990	Female	70 to 74	23% (10-43%)	77% (57-90%)	1.3 (0.34-3.6)	2.5% (-0.56-7.8%)	3% (-0.68-9.2%)	2.7 (-0.61-8.5)	50 (-11-160)
Bahrain	1990	Female	75 to 79	21% (8.3-43%)	79% (57-92%)	1.1 (0.24-3.3)	1.8% (-0.79-6.6%)	2.1% (-0.96-7.8%)	1.3 (-0.58-4.8)	19 (-8.2-70)
Bahrain	1990	Female	80 to 84	20% (6.2-41%)	80% (59-94%)	1 (0.19-2.9)	1.7% (-0.45-6.3%)	1.9% (-0.57-7.4%)	1.3 (-0.4-5)	14 (-3.8-52)
Bahrain	1990	Female	85 to 89	20% (6.5-42%)	80% (58-93%)	0.98 (0.19-2.9)	1.6% (-0.52-6.6%)	1.7% (-0.72-7.7%)	0.86 (-0.32-3.8)	6.8 (-2.2-29)
Bahrain	1990	Female	90 to 94	23% (7.5-45%)	77% (55-92%)	1.1 (0.22-3)	1.5% (-0.77-6%)	1.7% (-1-7.1%)	0.27 (-0.16-1.2)	1.5 (-0.74-6.2)
Bahrain	1990	Female	95 plus	27% (8.9-51%)	73% (49-91%)	1.3 (0.24-3.6)	1.6% (-1.2-7%)	1.7% (-1.5-8.1%)	0.095 (-0.086-0.44)	0.35 (-0.27-1.5)
Bahrain	1990	Male	All Ages	12% (7.6-18%)	88% (82-92%)	1.1 (0.92-1.3)	2.9% (2.2-3.7%)	3.8% (2.5-5.2%)	46 (29-67)	2200 (1600-2800)
Bahrain	1990	Male	15 to 19	7.9% (3.4-15%)	92% (85-97%)	0.4 (0.12-0.9)	1.7% (0.87-3.1%)	2.4% (0.98-5.2%)	0.48 (0.19-1)	48 (25-89)
Bahrain	1990	Male	20 to 24	12% (4.8-23%)	88% (77-95%)	0.89 (0.31-1.8)	3.6% (2.1-5.8%)	5.3% (2.3-9.4%)	1.5 (0.64-2.7)	140 (78-220)
Bahrain	1990	Male	25 to 29	15% (6.2-28%)	85% (72-94%)	1.4 (0.58-2.4)	4.5% (2.7-6.9%)	7.2% (3.7-12%)	3.5 (1.8-5.7)	280 (170-440)
Bahrain	1990	Male	30 to 34	17% (6.8-32%)	83% (68-93%)	1.7 (0.9-2.8)	4.6% (3-6.6%)	8% (4.7-12%)	4.5 (2.5-6.8)	340 (210-490)
Bahrain	1990	Male	35 to 39	19% (7.7-33%)	81% (67-92%)	2 (0.88-3.4)	4.6% (2.7-6.9%)	8% (4.3-12%)	4.5 (2.4-7.1)	310 (180-450)
Bahrain	1990	Male	40 to 44	19% (8.9-33%)	81% (67-91%)	2.2 (0.89-4)	4.6% (2.7-7.2%)	7.4% (4-12%)	3.8 (1.9-6.1)	220 (130-350)
Bahrain	1990	Male	45 to 49	19% (7.8-33%)	81% (67-92%)	2.3 (0.76-4.7)	4.8% (2.5-7.6%)	6.9% (3.5-11%)	3.3 (1.6-5.5)	160 (85-260)
Bahrain	1990	Male	50 to 54	18% (8-33%)	82% (67-92%)	2.2 (0.73-4.7)	4.5% (2.2-7.3%)	5.9% (2.7-9.9%)	3.8 (1.7-6.4)	160 (78-260)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	1990	Male	55 to 59	17% (7.7-31%)	83% (69-92%)	2.1 (0.67-4.6)	4.3% (2-7.4%)	5.3% (2.3-9.4%)	3.8 (1.6-6.8)	140 (58-240)
Bahrain	1990	Male	60 to 64	16% (6.9-31%)	84% (69-93%)	1.9 (0.56-4.6)	3.8% (1.4-6.9%)	4.4% (1.6-8.2%)	5 (1.8-9.4)	140 (54-270)
Bahrain	1990	Male	65 to 69	16% (7.1-28%)	84% (72-93%)	1.6 (0.48-3.5)	3.3% (1.1-6.2%)	3.8% (1.2-7.1%)	4.5 (1.5-8.6)	110 (38-200)
Bahrain	1990	Male	70 to 74	14% (6.5-25%)	86% (75-93%)	1.2 (0.36-2.8)	2.6% (0.54-5.5%)	2.9% (0.55-6%)	2.9 (0.58-6.2)	55 (11-110)
Bahrain	1990	Male	75 to 79	12% (5.3-24%)	88% (76-95%)	0.9 (0.22-2.4)	1.9% (0.26-4.5%)	2.1% (0.25-5%)	2 (0.25-4.9)	30 (4.2-71)
Bahrain	1990	Male	80 to 84	11% (4.5-22%)	89% (78-96%)	0.76 (0.19-2)	1.6% (0.29-3.9%)	1.8% (0.28-4.3%)	1.6 (0.28-4)	18 (3.4-43)
Bahrain	1990	Male	85 to 89	11% (4.5-22%)	89% (78-95%)	0.7 (0.18-1.8)	1.4% (0.15-3.3%)	1.5% (0.12-3.7%)	0.89 (0.068-2.3)	7.1 (0.79-18)
Bahrain	1990	Male	90 to 94	12% (5.2-24%)	88% (76-95%)	0.74 (0.19-1.9)	1.2% (0.022-3.2%)	1.3% (-0.069-3.6%)	0.22 (-0.011-0.61)	1.2 (0.022-3.3)
Bahrain	1990	Male	95 plus	15% (5.6-32%)	85% (68-94%)	0.84 (0.17-2.4)	1.1% (-0.21-3.6%)	1.2% (-0.4-3.9%)	0.047 (-0.016-0.16)	0.19 (-0.033-0.6)
Bahrain	1995	Female	All Ages	13% (7.7-20%)	87% (80-92%)	0.71 (0.53-0.92)	1.7% (0.76-2.8%)	2.6% (0.49-5.6%)	25 (4.8-54)	960 (440-1700)
Bahrain	1995	Female	15 to 19	11% (3.3-23%)	89% (77-97%)	0.5 (0.13-1.2)	0.97% (0.49-1.9%)	2.1% (0.79-4.7%)	0.2 (0.076-0.45)	25 (12-49)
Bahrain	1995	Female	20 to 24	14% (4.2-30%)	86% (70-96%)	0.8 (0.23-1.8)	1.6% (0.89-2.7%)	3.2% (1.2-6.6%)	0.43 (0.15-0.9)	55 (30-93)
Bahrain	1995	Female	25 to 29	16% (4.5-34%)	84% (66-96%)	0.97 (0.29-1.9)	1.9% (1-3.1%)	3.9% (1.5-7.2%)	0.75 (0.27-1.4)	79 (43-130)
Bahrain	1995	Female	30 to 34	18% (5.2-36%)	82% (64-95%)	1.1 (0.41-2.1)	1.9% (1.1-3.1%)	4.4% (2-7.3%)	0.95 (0.42-1.7)	88 (48-140)
Bahrain	1995	Female	35 to 39	19% (5.6-37%)	81% (63-94%)	1.3 (0.52-2.3)	2% (1-3.3%)	4.8% (2.2-7.7%)	1 (0.45-1.8)	83 (41-130)
Bahrain	1995	Female	40 to 44	20% (7.6-37%)	80% (63-92%)	1.5 (0.58-2.7)	2.6% (1.1-4.6%)	5.4% (2.3-9.3%)	1.3 (0.56-2.5)	81 (35-140)
Bahrain	1995	Female	45 to 49	21% (7.6-38%)	79% (62-92%)	1.6 (0.56-3.3)	2.7% (0.93-5.3%)	5.3% (1.9-10%)	1.3 (0.48-2.6)	72 (25-140)
Bahrain	1995	Female	50 to 54	22% (9-40%)	78% (60-91%)	1.7 (0.54-3.7)	3.4% (0.83-7.2%)	5.3% (1.4-11%)	2.1 (0.51-4.3)	89 (22-190)
Bahrain	1995	Female	55 to 59	23% (10-42%)	77% (58-90%)	1.7 (0.54-3.9)	3.8% (0.61-8.6%)	5.1% (0.89-12%)	3 (0.51-6.6)	100 (16-240)
Bahrain	1995	Female	60 to 64	24% (10-44%)	76% (56-90%)	1.7 (0.45-3.9)	3.3% (-0.11-8.7%)	4.2% (-0.07-11%)	3.9 (-0.061-10)	110 (-3.4-300)
Bahrain	1995	Female	65 to 69	25% (11-42%)	75% (58-89%)	1.5 (0.42-3.3)	2.9% (-0.41-8%)	3.4% (-0.45-9.5%)	3.5 (-0.49-9.8)	83 (-12-230)
Bahrain	1995	Female	70 to 74	23% (10-41%)	77% (59-90%)	1.2 (0.3-2.9)	1.9% (-0.87-6.7%)	2.2% (-1-7.7%)	2.7 (-1.1-9.5)	51 (-23-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	1995	Female	75 to 79	20% (8.2-40%)	80% (60-92%)	0.94 (0.21-2.7)	1.3% (-1.5-5.5%)	1.5% (-1.2-6.4%)	1.3 (-0.96-5.7)	19 (-14-87)
Bahrain	1995	Female	80 to 84	19% (6.8-40%)	81% (60-93%)	0.85 (0.18-2.5)	1.2% (-0.67-5.4%)	1.4% (-0.8-6%)	1 (-0.57-4.8)	11 (-5.7-52)
Bahrain	1995	Female	85 to 89	19% (6.6-41%)	81% (59-93%)	0.84 (0.18-2.5)	1.1% (-0.66-5.3%)	1.2% (-0.79-6%)	0.88 (-0.57-4.5)	7 (-4.1-34)
Bahrain	1995	Female	90 to 94	22% (7.6-44%)	78% (56-92%)	0.94 (0.19-2.7)	1.1% (-0.86-5.4%)	1.2% (-1.1-6.2%)	0.44 (-0.39-2.2)	2.4 (-1.9-12)
Bahrain	1995	Female	95 plus	27% (8.8-51%)	73% (49-91%)	1.1 (0.21-3.1)	1.2% (-1.5-5.7%)	1.2% (-1.8-6.5%)	0.099 (-0.15-0.51)	0.37 (-0.46-1.8)
Bahrain	1995	Male	All Ages	12% (8-18%)	88% (82-92%)	1 (0.83-1.2)	2.9% (2.2-3.7%)	3.5% (2.3-5%)	49 (30-71)	2400 (1800-3100)
Bahrain	1995	Male	15 to 19	8% (3.3-15%)	92% (85-97%)	0.35 (0.11-0.79)	1.5% (0.81-2.7%)	2.1% (0.91-4.5%)	0.46 (0.19-0.98)	48 (25-89)
Bahrain	1995	Male	20 to 24	12% (5-23%)	88% (77-95%)	0.78 (0.27-1.6)	3.2% (1.9-5.2%)	4.4% (2-8.2%)	1.5 (0.66-2.9)	150 (84-250)
Bahrain	1995	Male	25 to 29	15% (6-28%)	85% (72-94%)	1.2 (0.51-2.1)	4.2% (2.5-6.2%)	6.4% (3.2-10%)	3.5 (1.7-5.9)	290 (170-450)
Bahrain	1995	Male	30 to 34	17% (6.5-32%)	83% (68-94%)	1.5 (0.77-2.4)	4.3% (2.7-6.4%)	7.3% (4.1-11%)	4.6 (2.5-7.4)	350 (220-520)
Bahrain	1995	Male	35 to 39	19% (7.4-34%)	81% (66-93%)	1.8 (0.78-3)	4.4% (2.6-6.4%)	7.5% (4.1-11%)	4.9 (2.5-8)	340 (200-500)
Bahrain	1995	Male	40 to 44	19% (8.7-33%)	81% (67-91%)	1.9 (0.79-3.3)	4.3% (2.5-6.9%)	7% (3.6-11%)	4.6 (2.3-7.6)	280 (150-430)
Bahrain	1995	Male	45 to 49	18% (7.3-34%)	82% (66-93%)	1.9 (0.67-3.9)	4.4% (2.2-7.1%)	6.5% (3.2-10%)	4.5 (2.1-7.6)	230 (120-370)
Bahrain	1995	Male	50 to 54	18% (7.7-34%)	82% (66-92%)	1.9 (0.65-4.1)	4.1% (1.9-6.8%)	5.6% (2.5-9.4%)	3.9 (1.7-6.7)	170 (82-290)
Bahrain	1995	Male	55 to 59	17% (7.6-32%)	83% (68-92%)	1.9 (0.61-4)	4% (1.7-7%)	5% (2.1-9%)	3.8 (1.6-7)	140 (61-250)
Bahrain	1995	Male	60 to 64	16% (7.4-30%)	84% (70-93%)	1.7 (0.51-4)	3.5% (1.2-6.6%)	4.1% (1.4-7.8%)	4.4 (1.5-8.2)	130 (44-240)
Bahrain	1995	Male	65 to 69	16% (7.1-29%)	84% (71-93%)	1.4 (0.43-3)	3% (0.91-5.6%)	3.4% (0.99-6.5%)	4.6 (1.3-8.8)	110 (34-210)
Bahrain	1995	Male	70 to 74	14% (6.6-25%)	86% (75-93%)	1 (0.31-2.4)	2.2% (0.41-5%)	2.5% (0.42-5.5%)	3.3 (0.56-7.7)	63 (11-140)
Bahrain	1995	Male	75 to 79	12% (5.3-23%)	88% (77-95%)	0.75 (0.19-2.1)	1.6% (0.23-4.1%)	1.8% (0.25-4.6%)	2.3 (0.31-5.9)	34 (5-87)
Bahrain	1995	Male	80 to 84	11% (4.4-21%)	89% (79-96%)	0.64 (0.16-1.6)	1.4% (0.21-3.5%)	1.5% (0.2-3.8%)	1.3 (0.19-3.3)	15 (2.3-36)
Bahrain	1995	Male	85 to 89	11% (4.4-22%)	89% (78-96%)	0.59 (0.16-1.5)	1.2% (0.15-3%)	1.2% (0.12-3.2%)	0.99 (0.092-2.6)	7.9 (1-21)
Bahrain	1995	Male	90 to 94	12% (5-24%)	88% (76-95%)	0.62 (0.16-1.6)	1% (0.02-2.8%)	1% (-0.056-3.1%)	0.38 (-0.02-1.1)	2.2 (0.043-5.9)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	1995	Male	95 plus	15% (5.6-31%)	85% (69-94%)	0.71 (0.15-2)	0.95% (-0.21-3%)	0.94% (-0.39-3.3%)	0.062 (-0.025-0.22)	0.25 (-0.053-0.8)
Bahrain	2000	Female	All Ages	13% (7.6-21%)	87% (79-92%)	0.58 (0.42-0.73)	1.3% (0.56-2.3%)	1.8% (0.16-4.4%)	17 (1.5-43)	800 (360-1400)
Bahrain	2000	Female	15 to 19	11% (3.1-25%)	89% (75-97%)	0.4 (0.1-0.98)	0.9% (0.45-1.7%)	1.9% (0.74-4.4%)	0.23 (0.084-0.52)	27 (13-54)
Bahrain	2000	Female	20 to 24	14% (3.9-30%)	86% (70-96%)	0.64 (0.17-1.4)	1.5% (0.82-2.6%)	2.8% (0.98-5.9%)	0.46 (0.15-0.98)	58 (30-100)
Bahrain	2000	Female	25 to 29	16% (4.3-33%)	84% (67-96%)	0.77 (0.21-1.6)	1.6% (0.88-2.7%)	3.4% (1.1-6.3%)	0.6 (0.21-1.1)	69 (37-110)
Bahrain	2000	Female	30 to 34	17% (4.6-37%)	83% (63-95%)	0.89 (0.3-1.7)	1.7% (0.91-2.7%)	3.6% (1.5-6.4%)	0.86 (0.35-1.6)	82 (45-130)
Bahrain	2000	Female	35 to 39	19% (5.2-37%)	81% (63-95%)	1 (0.38-1.8)	1.7% (0.88-2.8%)	3.9% (1.7-6.5%)	1.2 (0.52-2.2)	94 (46-150)
Bahrain	2000	Female	40 to 44	20% (7-38%)	80% (62-93%)	1.1 (0.48-2)	2% (0.85-3.5%)	4.3% (1.8-7.4%)	1.6 (0.66-2.9)	100 (41-170)
Bahrain	2000	Female	45 to 49	20% (7.1-37%)	80% (63-93%)	1.2 (0.44-2.5)	2.1% (0.66-4.3%)	4.2% (1.4-8.6%)	1.3 (0.46-2.8)	72 (23-160)
Bahrain	2000	Female	50 to 54	21% (8.4-40%)	79% (60-92%)	1.3 (0.43-2.8)	2.4% (0.39-5.4%)	4% (0.9-9%)	1.3 (0.28-2.9)	57 (9.4-130)
Bahrain	2000	Female	55 to 59	22% (9.3-42%)	78% (58-91%)	1.3 (0.39-2.9)	2.5% (0.11-6.4%)	3.7% (0.39-9%)	2 (0.21-5)	72 (2.9-190)
Bahrain	2000	Female	60 to 64	24% (9.8-44%)	76% (56-90%)	1.3 (0.34-3)	2.1% (-0.6-6.7%)	2.8% (-0.58-8.4%)	2.5 (-0.56-7.3)	73 (-20-230)
Bahrain	2000	Female	65 to 69	24% (10-42%)	76% (58-90%)	1.1 (0.3-2.6)	1.6% (-1-5.7%)	2% (-1.1-6.9%)	2.1 (-1.2-7.6)	50 (-32-180)
Bahrain	2000	Female	70 to 74	23% (9.1-40%)	77% (60-91%)	0.89 (0.23-2.3)	0.88% (-1.3-4.9%)	1.1% (-1.5-5.7%)	1.6 (-2.2-8.4)	29 (-43-160)
Bahrain	2000	Female	75 to 79	20% (7.6-40%)	80% (60-92%)	0.73 (0.16-2.1)	0.48% (-1.2-3.8%)	0.58% (-1.4-4.5%)	0.4 (-0.98-3.2)	6.2 (-15-50)
Bahrain	2000	Female	80 to 84	19% (6.6-41%)	81% (59-93%)	0.66 (0.13-1.9)	0.48% (-0.93-3.4%)	0.54% (-1.1-4%)	0.46 (-0.95-3.3)	5.3 (-10-38)
Bahrain	2000	Female	85 to 89	19% (6.5-40%)	81% (60-93%)	0.66 (0.13-1.9)	0.45% (-1.1-3.4%)	0.48% (-1.3-3.8%)	0.3 (-0.79-2.5)	2.5 (-5.9-19)
Bahrain	2000	Female	90 to 94	22% (7.4-43%)	78% (57-93%)	0.74 (0.15-2.1)	0.45% (-1.3-3.6%)	0.45% (-1.6-4.1%)	0.18 (-0.63-1.7)	1.1 (-3.3-8.8)
Bahrain	2000	Female	95 plus	27% (9.1-51%)	73% (49-91%)	0.87 (0.17-2.4)	0.49% (-1.8-4.3%)	0.47% (-2.1-4.9%)	0.072 (-0.32-0.74)	0.29 (-1-2.6)
Bahrain	2000	Male	All Ages	12% (8-18%)	88% (82-92%)	0.81 (0.66-1)	3% (2.2-3.9%)	3.5% (2.2-5%)	55 (33-80)	2800 (2000-3800)
Bahrain	2000	Male	15 to 19	7.8% (3.3-15%)	92% (85-97%)	0.27 (0.079-0.63)	1.5% (0.81-2.7%)	2% (0.85-4.3%)	0.57 (0.23-1.2)	57 (30-110)
Bahrain	2000	Male	20 to 24	12% (4.9-23%)	88% (77-95%)	0.62 (0.22-1.3)	3.3% (1.8-5.5%)	4.3% (1.8-8.2%)	2.1 (0.86-4.2)	190 (100-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	2000	Male	25 to 29	15% (6.1-28%)	85% (72-94%)	0.97 (0.38-1.8)	4.2% (2.3-6.7%)	6.2% (2.9-11%)	4.3 (1.9-7.7)	340 (180-550)
Bahrain	2000	Male	30 to 34	17% (6.6-32%)	83% (68-93%)	1.2 (0.59-2)	4.4% (2.6-6.9%)	6.9% (3.7-11%)	5.9 (3-10)	420 (240-670)
Bahrain	2000	Male	35 to 39	19% (7.4-33%)	81% (67-93%)	1.4 (0.62-2.3)	4.4% (2.5-6.7%)	7.1% (3.7-11%)	6 (2.9-10)	390 (210-630)
Bahrain	2000	Male	40 to 44	19% (8.5-32%)	81% (68-92%)	1.5 (0.63-2.5)	4.3% (2.3-6.9%)	6.6% (3.2-11%)	6.6 (3.1-11)	380 (200-620)
Bahrain	2000	Male	45 to 49	18% (7.4-33%)	82% (67-93%)	1.5 (0.55-3)	4.1% (2-6.8%)	6% (2.8-10%)	6.7 (2.8-12)	330 (160-560)
Bahrain	2000	Male	50 to 54	18% (7.7-33%)	82% (67-92%)	1.5 (0.52-3.2)	3.8% (1.6-6.6%)	5.2% (2.1-9.1%)	4.9 (2-8.7)	210 (90-370)
Bahrain	2000	Male	55 to 59	17% (7.6-32%)	83% (68-92%)	1.5 (0.48-3.2)	3.6% (1.4-6.6%)	4.5% (1.7-8.4%)	3.8 (1.4-7)	140 (55-260)
Bahrain	2000	Male	60 to 64	16% (6.9-31%)	84% (69-93%)	1.3 (0.37-3.1)	3% (0.88-5.9%)	3.5% (1-7.1%)	3.7 (1-7.4)	110 (32-220)
Bahrain	2000	Male	65 to 69	15% (6.8-28%)	85% (72-93%)	1.1 (0.33-2.4)	2.4% (0.62-4.9%)	2.7% (0.65-5.7%)	3.7 (0.89-7.8)	90 (23-190)
Bahrain	2000	Male	70 to 74	14% (6.5-25%)	86% (75-94%)	0.79 (0.25-1.9)	1.7% (0.26-4.1%)	1.9% (0.28-4.6%)	2.9 (0.41-7.2)	57 (8.5-140)
Bahrain	2000	Male	75 to 79	12% (5.4-23%)	88% (77-95%)	0.61 (0.16-1.6)	1.2% (0.13-3.3%)	1.4% (0.12-3.8%)	1.6 (0.15-4.5)	25 (2.6-68)
Bahrain	2000	Male	80 to 84	11% (4.5-22%)	89% (78-95%)	0.51 (0.14-1.3)	1% (0.14-2.9%)	1.1% (0.13-3.2%)	1.1 (0.13-3.1)	13 (1.8-35)
Bahrain	2000	Male	85 to 89	11% (4.4-22%)	89% (78-96%)	0.47 (0.13-1.2)	0.85% (0.043-2.5%)	0.89% (-0.0072-2.7%)	0.58 (-0.0048-1.7)	4.8 (0.23-14)
Bahrain	2000	Male	90 to 94	12% (5-24%)	88% (76-95%)	0.49 (0.13-1.2)	0.71% (-0.09-2.2%)	0.71% (-0.2-2.4%)	0.3 (-0.084-1)	1.8 (-0.23-5.5)
Bahrain	2000	Male	95 plus	15% (5.4-30%)	85% (70-95%)	0.56 (0.12-1.6)	0.67% (-0.31-2.6%)	0.62% (-0.5-2.7%)	0.083 (-0.067-0.37)	0.35 (-0.16-1.3)
Bahrain	2005	Female	All Ages	13% (7.8-19%)	87% (81-92%)	0.52 (0.39-0.67)	1% (0.42-1.9%)	1.3% (-0.079-3.6%)	13 (-0.84-38)	720 (280-1300)
Bahrain	2005	Female	15 to 19	10% (2.9-21%)	90% (79-97%)	0.34 (0.086-0.8)	0.69% (0.36-1.3%)	1.4% (0.53-3.4%)	0.15 (0.051-0.34)	23 (11-41)
Bahrain	2005	Female	20 to 24	13% (3.8-28%)	87% (72-96%)	0.54 (0.15-1.2)	1.3% (0.7-2.1%)	2.2% (0.76-5%)	0.35 (0.12-0.78)	55 (30-92)
Bahrain	2005	Female	25 to 29	14% (3.9-30%)	86% (70-96%)	0.64 (0.19-1.3)	1.4% (0.8-2.2%)	2.6% (0.87-5.3%)	0.52 (0.16-1)	71 (40-110)
Bahrain	2005	Female	30 to 34	16% (4.2-33%)	84% (67-96%)	0.74 (0.25-1.3)	1.3% (0.8-2.1%)	2.9% (1.2-5.1%)	0.68 (0.27-1.2)	77 (44-120)
Bahrain	2005	Female	35 to 39	18% (4.8-33%)	82% (67-95%)	0.86 (0.32-1.5)	1.4% (0.73-2.3%)	3.3% (1.4-5.5%)	0.93 (0.39-1.6)	79 (41-130)
Bahrain	2005	Female	40 to 44	19% (7.3-33%)	81% (67-93%)	0.96 (0.41-1.7)	1.5% (0.7-2.7%)	3.6% (1.6-6.3%)	1.3 (0.54-2.2)	83 (37-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	2005	Female	45 to 49	19% (6.7-36%)	81% (64-93%)	1.1 (0.39-2.1)	1.6% (0.46-3.5%)	3.5% (1.2-7.4%)	1.5 (0.48-3.3)	86 (23-190)
Bahrain	2005	Female	50 to 54	20% (7.7-38%)	80% (62-92%)	1.1 (0.35-2.3)	1.7% (0.14-4.3%)	3.3% (0.66-7.5%)	1.5 (0.31-3.5)	70 (5.5-180)
Bahrain	2005	Female	55 to 59	21% (8.4-40%)	79% (60-92%)	1.1 (0.32-2.5)	1.9% (-0.11-5.4%)	2.9% (0.16-7.8%)	1.5 (0.085-4.1)	56 (-3.2-170)
Bahrain	2005	Female	60 to 64	22% (8.6-43%)	78% (57-91%)	1.1 (0.29-2.7)	1.5% (-0.77-5.4%)	2.1% (-0.8-7.1%)	1.6 (-0.57-5.5)	49 (-25-180)
Bahrain	2005	Female	65 to 69	23% (9.8-41%)	77% (59-90%)	0.97 (0.26-2.2)	1% (-1.2-4.7%)	1.4% (-1.3-5.9%)	1.7 (-1.7-7.5)	41 (-44-180)
Bahrain	2005	Female	70 to 74	21% (9.3-39%)	79% (61-91%)	0.77 (0.2-1.9)	0.5% (-1.4-4.1%)	0.62% (-1.6-4.8%)	0.81 (-2.2-6.4)	15 (-44-130)
Bahrain	2005	Female	75 to 79	19% (7.4-38%)	81% (62-93%)	0.63 (0.13-1.8)	0.21% (-1.3-3.2%)	0.27% (-1.5-3.7%)	0.32 (-1.9-4.6)	4.8 (-31-74)
Bahrain	2005	Female	80 to 84	18% (6.1-38%)	82% (62-94%)	0.57 (0.12-1.7)	0.23% (-1.1-2.6%)	0.26% (-1.2-3.1%)	0.25 (-1.2-3.1)	2.9 (-14-35)
Bahrain	2005	Female	85 to 89	18% (6.5-40%)	82% (60-93%)	0.57 (0.12-1.6)	0.2% (-1.1-2.7%)	0.2% (-1.3-3%)	0.14 (-0.9-2.2)	1.2 (-6.9-17)
Bahrain	2005	Female	90 to 94	21% (7.5-43%)	79% (57-93%)	0.64 (0.13-1.7)	0.18% (-1.4-2.8%)	0.16% (-1.6-3.2%)	0.052 (-0.54-1.1)	0.36 (-2.7-5.6)
Bahrain	2005	Female	95 plus	25% (8.3-50%)	75% (50-92%)	0.75 (0.14-2.2)	0.2% (-1.7-3.5%)	0.14% (-2.1-4%)	0.02 (-0.29-0.53)	0.11 (-0.93-1.9)
Bahrain	2005	Male	All Ages	12% (7.9-18%)	88% (82-92%)	0.74 (0.62-0.9)	2.4% (1.8-3%)	2.9% (1.8-4.1%)	45 (27-66)	2500 (1900-3300)
Bahrain	2005	Male	15 to 19	7.3% (3-14%)	93% (86-97%)	0.23 (0.067-0.53)	1.2% (0.71-2%)	1.6% (0.7-3.4%)	0.41 (0.18-0.88)	51 (30-90)
Bahrain	2005	Male	20 to 24	11% (4.5-22%)	89% (78-96%)	0.52 (0.17-1.1)	2.6% (1.6-4.1%)	3.4% (1.5-6.7%)	1.7 (0.71-3.3)	200 (120-330)
Bahrain	2005	Male	25 to 29	14% (5.8-27%)	86% (73-94%)	0.81 (0.33-1.5)	3.3% (2-4.9%)	5.2% (2.5-8.9%)	3.4 (1.6-5.9)	320 (190-500)
Bahrain	2005	Male	30 to 34	16% (6.3-31%)	84% (69-94%)	1 (0.49-1.7)	3.4% (2.2-4.9%)	5.9% (3.3-9.2%)	4.1 (2.3-6.6)	350 (220-510)
Bahrain	2005	Male	35 to 39	18% (7.3-32%)	82% (68-93%)	1.2 (0.52-2)	3.3% (2-4.9%)	6% (3.2-9.4%)	4.6 (2.4-7.6)	350 (210-530)
Bahrain	2005	Male	40 to 44	18% (8.2-31%)	82% (69-92%)	1.3 (0.53-2.2)	3.3% (1.9-5%)	5.6% (2.8-9%)	5.3 (2.6-8.9)	350 (190-530)
Bahrain	2005	Male	45 to 49	17% (6.8-32%)	83% (68-93%)	1.3 (0.46-2.5)	3.3% (1.7-5.4%)	5.1% (2.3-8.5%)	6 (2.5-10)	320 (160-530)
Bahrain	2005	Male	50 to 54	17% (7.1-32%)	83% (68-93%)	1.3 (0.42-2.6)	3.1% (1.2-5.2%)	4.3% (1.6-7.5%)	4.6 (1.8-8.3)	210 (84-380)
Bahrain	2005	Male	55 to 59	16% (7.2-30%)	84% (70-93%)	1.2 (0.4-2.8)	2.8% (1-5.2%)	3.7% (1.2-6.9%)	3.4 (1.1-6.5)	130 (45-250)
Bahrain	2005	Male	60 to 64	15% (6.7-30%)	85% (70-93%)	1.1 (0.31-2.9)	2.4% (0.62-5%)	2.9% (0.71-6%)	2.8 (0.67-5.7)	88 (22-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	2005	Male	65 to 69	15% (6.3-26%)	85% (74-94%)	0.91 (0.28-2.1)	1.9% (0.42-4.2%)	2.3% (0.45-4.8%)	3 (0.6-6.7)	75 (16-170)
Bahrain	2005	Male	70 to 74	13% (5.9-23%)	87% (77-94%)	0.67 (0.21-1.6)	1.4% (0.17-3.6%)	1.5% (0.18-4.1%)	2.4 (0.28-6.3)	47 (6.3-130)
Bahrain	2005	Male	75 to 79	11% (4.8-22%)	89% (78-95%)	0.51 (0.13-1.3)	0.95% (0.039-2.7%)	1% (0.0087-3.1%)	1.4 (0.013-4.3)	22 (0.92-66)
Bahrain	2005	Male	80 to 84	10% (4.1-20%)	90% (80-96%)	0.43 (0.11-1.1)	0.77% (0.064-2.2%)	0.83% (0.034-2.4%)	0.95 (0.035-2.8)	11 (0.92-32)
Bahrain	2005	Male	85 to 89	10% (4-20%)	90% (80-96%)	0.39 (0.1-0.99)	0.63% (0.001-1.9%)	0.66% (-0.046-2.1%)	0.55 (-0.038-1.8)	4.6 (0.007-14)
Bahrain	2005	Male	90 to 94	11% (4.4-23%)	89% (77-96%)	0.41 (0.1-1)	0.52% (-0.14-1.8%)	0.52% (-0.23-1.9%)	0.18 (-0.081-0.67)	1.1 (-0.28-3.8)
Bahrain	2005	Male	95 plus	14% (5.1-30%)	86% (70-95%)	0.48 (0.098-1.4)	0.5% (-0.34-2.2%)	0.44% (-0.52-2.3%)	0.058 (-0.068-0.3)	0.25 (-0.18-1.1)
Bahrain	2010	Female	All Ages	14% (7.8-21%)	86% (79-92%)	0.5 (0.36-0.67)	0.87% (0.3-1.7%)	0.98% (-0.31-3.1%)	12 (-3.8-37)	790 (260-1600)
Bahrain	2010	Female	15 to 19	9.7% (3.2-22%)	90% (78-97%)	0.29 (0.077-0.73)	0.64% (0.35-1.1%)	1.3% (0.51-3%)	0.15 (0.054-0.36)	26 (14-47)
Bahrain	2010	Female	20 to 24	12% (3.9-26%)	88% (74-96%)	0.46 (0.12-1.1)	1.1% (0.68-1.9%)	1.9% (0.67-4.3%)	0.37 (0.13-0.85)	69 (40-120)
Bahrain	2010	Female	25 to 29	14% (4.3-30%)	86% (70-96%)	0.55 (0.17-1.2)	1.2% (0.73-1.9%)	2.2% (0.76-4.4%)	0.57 (0.2-1.2)	98 (56-150)
Bahrain	2010	Female	30 to 34	16% (4.4-33%)	84% (67-96%)	0.64 (0.23-1.2)	1.2% (0.7-1.8%)	2.5% (1.1-4.5%)	0.68 (0.29-1.2)	93 (54-150)
Bahrain	2010	Female	35 to 39	18% (4.8-34%)	82% (66-95%)	0.76 (0.27-1.4)	1.1% (0.58-1.9%)	2.8% (1.2-5%)	0.97 (0.39-1.8)	95 (48-160)
Bahrain	2010	Female	40 to 44	19% (6.6-34%)	81% (66-93%)	0.85 (0.34-1.6)	1.2% (0.47-2.2%)	3.1% (1.3-5.8%)	1.3 (0.56-2.5)	95 (37-170)
Bahrain	2010	Female	45 to 49	19% (6.5-35%)	81% (65-93%)	0.92 (0.32-1.8)	1.3% (0.21-3%)	3% (0.92-6.3%)	1.6 (0.47-3.4)	90 (15-200)
Bahrain	2010	Female	50 to 54	20% (7.8-37%)	80% (63-92%)	0.96 (0.31-1.9)	1.3% (-0.15-3.5%)	2.7% (0.34-6.4%)	1.9 (0.23-4.6)	91 (-8.9-250)
Bahrain	2010	Female	55 to 59	21% (8.6-39%)	79% (61-91%)	0.98 (0.29-2.3)	1.4% (-0.37-4.3%)	2.3% (-0.2-6.4%)	1.9 (-0.16-5.3)	69 (-18-220)
Bahrain	2010	Female	60 to 64	22% (8.9-43%)	78% (57-91%)	0.95 (0.25-2.4)	1% (-0.96-4.6%)	1.6% (-1-6.1%)	1.2 (-0.78-4.6)	35 (-35-160)
Bahrain	2010	Female	65 to 69	23% (9.2-41%)	77% (59-91%)	0.84 (0.21-2.1)	0.56% (-1.4-3.7%)	0.82% (-1.6-4.8%)	0.94 (-2-5.6)	22 (-56-140)
Bahrain	2010	Female	70 to 74	21% (8.9-39%)	79% (61-91%)	0.67 (0.16-1.8)	0.12% (-1.7-3.3%)	0.19% (-1.9-4%)	0.32 (-3.4-7)	5.1 (-70-140)
Bahrain	2010	Female	75 to 79	19% (7.4-37%)	81% (63-93%)	0.54 (0.11-1.5)	-0.059% (-1.5-2.5%)	-0.067% (-1.8-3%)	-0.084 (-2.3-3.8)	-1.4 (-36-58)
Bahrain	2010	Female	80 to 84	17% (5.9-36%)	83% (64-94%)	0.48 (0.097-1.4)	-0.014% (-1.3-2.3%)	-0.028% (-1.6-2.6%)	-0.034 (-1.9-3)	-0.22 (-21-34)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	2010	Female	85 to 89	17% (6.3-38%)	83% (62-94%)	0.48 (0.098-1.5)	-0.026% (-1.2-1.9%)	-0.058% (-1.5-2.2%)	-0.051 (-1.2-1.8)	-0.21 (-9.3-15)
Bahrain	2010	Female	90 to 94	20% (7-41%)	80% (59-93%)	0.54 (0.11-1.6)	-0.037% (-1.4-2%)	-0.094% (-1.6-2.3%)	-0.036 (-0.62-0.88)	-0.087 (-3.1-5)
Bahrain	2010	Female	95 plus	25% (8.1-48%)	75% (52-92%)	0.64 (0.12-2)	-0.025% (-1.7-2.9%)	-0.12% (-2.1-3.1%)	-0.015 (-0.26-0.37)	-0.013 (-0.87-1.4)
Bahrain	2010	Male	All Ages	13% (8.2-19%)	87% (81-92%)	0.7 (0.53-0.92)	2.2% (1.6-2.8%)	2.6% (1.5-3.8%)	47 (27-73)	3100 (2200-4200)
Bahrain	2010	Male	15 to 19	7.3% (3-15%)	93% (85-97%)	0.2 (0.056-0.48)	1.1% (0.68-1.8%)	1.5% (0.65-3.1%)	0.38 (0.16-0.82)	52 (31-87)
Bahrain	2010	Male	20 to 24	11% (4.6-22%)	89% (78-95%)	0.45 (0.15-1)	2.4% (1.5-3.8%)	3.1% (1.4-6.1%)	1.7 (0.74-3.5)	220 (140-370)
Bahrain	2010	Male	25 to 29	14% (5.6-26%)	86% (74-94%)	0.7 (0.27-1.3)	2.8% (1.8-4.1%)	4.5% (2.2-7.8%)	4 (2-7.4)	460 (290-710)
Bahrain	2010	Male	30 to 34	16% (5.9-30%)	84% (70-94%)	0.89 (0.41-1.5)	2.7% (1.8-4%)	5.1% (2.8-8.2%)	4.7 (2.5-7.9)	480 (300-700)
Bahrain	2010	Male	35 to 39	17% (6.6-31%)	83% (69-93%)	1 (0.41-1.8)	2.7% (1.7-3.9%)	5.3% (2.8-8.6%)	4.4 (2.3-7.4)	400 (250-590)
Bahrain	2010	Male	40 to 44	17% (7.8-31%)	83% (69-92%)	1.1 (0.45-2)	2.6% (1.5-4.2%)	4.9% (2.4-8.1%)	5.2 (2.5-9)	390 (210-600)
Bahrain	2010	Male	45 to 49	17% (6.8-30%)	83% (70-93%)	1.1 (0.39-2.2)	2.7% (1.3-4.5%)	4.5% (2.1-7.6%)	5.6 (2.5-9.8)	330 (170-560)
Bahrain	2010	Male	50 to 54	16% (7-32%)	84% (68-93%)	1.1 (0.36-2.3)	2.5% (1-4.3%)	3.8% (1.4-6.6%)	5.9 (2.1-11)	300 (120-530)
Bahrain	2010	Male	55 to 59	15% (6.6-29%)	85% (71-93%)	1 (0.32-2.3)	2.3% (0.74-4.4%)	3.2% (0.94-6.2%)	5.2 (1.5-10)	210 (68-410)
Bahrain	2010	Male	60 to 64	15% (6.1-29%)	85% (71-94%)	0.94 (0.28-2.4)	1.9% (0.5-4%)	2.5% (0.62-5.3%)	3.4 (0.89-7.5)	120 (31-250)
Bahrain	2010	Male	65 to 69	14% (6.3-26%)	86% (74-94%)	0.77 (0.22-1.8)	1.5% (0.29-3.6%)	1.9% (0.32-4.4%)	2.4 (0.42-5.8)	63 (12-150)
Bahrain	2010	Male	70 to 74	13% (5.8-23%)	87% (77-94%)	0.58 (0.16-1.4)	1.1% (0.064-3.1%)	1.2% (0.05-3.6%)	1.9 (0.082-6.1)	40 (2.3-120)
Bahrain	2010	Male	75 to 79	11% (4.8-22%)	89% (78-95%)	0.44 (0.1-1.2)	0.73% (-0.059-2.4%)	0.81% (-0.09-2.7%)	1.1 (-0.12-3.8)	18 (-1.4-59)
Bahrain	2010	Male	80 to 84	10% (3.8-20%)	90% (80-96%)	0.37 (0.092-1)	0.58% (-0.036-2%)	0.62% (-0.083-2.2%)	0.8 (-0.11-2.7)	9.3 (-0.58-31)
Bahrain	2010	Male	85 to 89	9.9% (3.9-20%)	90% (80-96%)	0.34 (0.093-0.89)	0.48% (-0.057-1.6%)	0.5% (-0.11-1.7%)	0.43 (-0.099-1.5)	3.7 (-0.45-12)
Bahrain	2010	Male	90 to 94	11% (4.5-23%)	89% (77-95%)	0.36 (0.092-0.94)	0.4% (-0.17-1.5%)	0.38% (-0.26-1.6%)	0.15 (-0.1-0.66)	0.95 (-0.39-3.6)
Bahrain	2010	Male	95 plus	14% (5.2-29%)	86% (71-95%)	0.41 (0.082-1.3)	0.39% (-0.32-1.8%)	0.32% (-0.49-1.9%)	0.034 (-0.053-0.2)	0.16 (-0.13-0.78)
Bahrain	2016	Female	All Ages	14% (8.5-21%)	86% (79-91%)	0.52 (0.28-0.83)	0.87% (0.12-2%)	1% (-0.51-3.7%)	14 (-7.5-53)	880 (130-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	2016	Female	15 to 19	10% (3.1-22%)	90% (78-97%)	0.3 (0.067-0.8)	0.64% (0.35-1.2%)	1.4% (0.52-3.4%)	0.16 (0.058-0.38)	28 (15-54)
Bahrain	2016	Female	20 to 24	13% (3.8-28%)	87% (72-96%)	0.49 (0.11-1.2)	1.2% (0.67-1.9%)	2% (0.65-4.8%)	0.31 (0.098-0.78)	61 (34-110)
Bahrain	2016	Female	25 to 29	15% (4.5-31%)	85% (69-96%)	0.58 (0.15-1.4)	1.2% (0.71-2%)	2.3% (0.75-5%)	0.49 (0.15-1.2)	88 (49-140)
Bahrain	2016	Female	30 to 34	17% (4.5-34%)	83% (66-95%)	0.67 (0.21-1.4)	1.2% (0.65-2%)	2.6% (1-5.3%)	0.75 (0.26-1.6)	100 (55-180)
Bahrain	2016	Female	35 to 39	18% (4.9-35%)	82% (65-95%)	0.78 (0.24-1.5)	1.1% (0.53-2%)	2.9% (1-5.7%)	0.97 (0.31-2)	96 (46-170)
Bahrain	2016	Female	40 to 44	19% (7.2-35%)	81% (65-93%)	0.87 (0.31-1.8)	1.2% (0.38-2.4%)	3.1% (1.1-6.3%)	1.4 (0.47-2.9)	99 (32-200)
Bahrain	2016	Female	45 to 49	20% (7-37%)	80% (63-93%)	0.96 (0.31-2)	1.3% (0.15-3%)	3.1% (0.78-7%)	1.9 (0.48-4.5)	110 (14-270)
Bahrain	2016	Female	50 to 54	21% (8.3-39%)	79% (61-92%)	1 (0.31-2.2)	1.3% (-0.19-3.8%)	2.7% (0.22-6.9%)	2.4 (0.21-6.2)	110 (-16-320)
Bahrain	2016	Female	55 to 59	22% (9.2-41%)	78% (59-91%)	1 (0.3-2.4)	1.3% (-0.6-4.5%)	2.3% (-0.36-6.9%)	2.9 (-0.46-8.9)	110 (-47-390)
Bahrain	2016	Female	60 to 64	23% (9.7-41%)	77% (59-90%)	0.98 (0.24-2.5)	0.95% (-1.2-4.5%)	1.5% (-1.3-6.3%)	1.6 (-1.4-6.6)	49 (-62-230)
Bahrain	2016	Female	65 to 69	23% (9.5-41%)	77% (59-90%)	0.86 (0.23-2.1)	0.53% (-1.5-4.1%)	0.8% (-1.7-5.2%)	0.92 (-2.1-6.1)	21 (-60-160)
Bahrain	2016	Female	70 to 74	22% (9.2-39%)	78% (61-91%)	0.69 (0.15-1.8)	0.11% (-1.7-3.7%)	0.19% (-2-4.5%)	0.3 (-3.4-7.1)	4.6 (-73-140)
Bahrain	2016	Female	75 to 79	19% (7.1-37%)	81% (63-93%)	0.56 (0.1-1.7)	-0.05% (-1.6-2.8%)	-0.054% (-1.9-3.4%)	-0.098 (-3.4-6.2)	-1.7 (-55-98)
Bahrain	2016	Female	80 to 84	18% (6.2-39%)	82% (61-94%)	0.51 (0.1-1.6)	0.0014% (-1.4-2.4%)	-0.0075% (-1.6-2.9%)	-0.013 (-2.3-4)	0.022 (-27-46)
Bahrain	2016	Female	85 to 89	19% (6.3-40%)	81% (60-94%)	0.51 (0.096-1.7)	-0.037% (-1.4-2.4%)	-0.073% (-1.6-2.8%)	-0.071 (-1.6-2.5)	-0.33 (-12-21)
Bahrain	2016	Female	90 to 94	22% (7.1-43%)	78% (57-93%)	0.57 (0.11-1.8)	-0.051% (-1.5-2.2%)	-0.12% (-1.9-2.6%)	-0.069 (-1.1-1.4)	-0.19 (-5.8-8.1)
Bahrain	2016	Female	95 plus	26% (8.2-51%)	74% (49-92%)	0.67 (0.12-2.1)	-0.04% (-1.8-2.9%)	-0.15% (-2.3-3.3%)	-0.024 (-0.38-0.53)	-0.028 (-1.3-1.9)
Bahrain	2016	Male	All Ages	13% (8.3-19%)	87% (81-92%)	0.68 (0.38-1.1)	2.1% (1.3-3%)	2.5% (1.1-4%)	54 (24-94)	3300 (2000-5000)
Bahrain	2016	Male	15 to 19	7.2% (3-14%)	93% (86-97%)	0.2 (0.047-0.52)	1.1% (0.64-1.8%)	1.5% (0.63-3.3%)	0.44 (0.18-1)	63 (36-110)
Bahrain	2016	Male	20 to 24	11% (4.4-22%)	89% (78-96%)	0.45 (0.12-1.1)	2.4% (1.4-3.8%)	3.1% (1.2-6.4%)	1.8 (0.71-3.9)	250 (140-410)
Bahrain	2016	Male	25 to 29	14% (5.6-26%)	86% (74-94%)	0.69 (0.23-1.4)	2.8% (1.8-4.2%)	4.5% (2.1-8.3%)	3.9 (1.7-7.3)	440 (270-700)
Bahrain	2016	Male	30 to 34	16% (5.9-31%)	84% (69-94%)	0.88 (0.34-1.7)	2.7% (1.7-4.2%)	5% (2.4-8.5%)	4.9 (2.2-8.8)	490 (290-750)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bahrain	2016	Male	35 to 39	17% (6.8-32%)	83% (68-93%)	1 (0.34-2)	2.7% (1.6-4.1%)	5.3% (2.5-8.9%)	4.4 (1.9-8)	390 (220-610)
Bahrain	2016	Male	40 to 44	18% (8.1-31%)	82% (69-92%)	1.1 (0.4-2.1)	2.6% (1.5-4.2%)	4.8% (2.2-8.3%)	5.4 (2.3-10)	400 (210-670)
Bahrain	2016	Male	45 to 49	17% (6.7-31%)	83% (69-93%)	1.1 (0.37-2.3)	2.6% (1.2-4.5%)	4.4% (1.9-7.7%)	6.3 (2.5-12)	380 (180-670)
Bahrain	2016	Male	50 to 54	16% (7.3-32%)	84% (68-93%)	1.1 (0.36-2.4)	2.4% (0.96-4.3%)	3.7% (1.3-6.6%)	6.4 (2.1-13)	320 (130-580)
Bahrain	2016	Male	55 to 59	16% (6.7-30%)	84% (70-93%)	1 (0.31-2.4)	2.2% (0.68-4.4%)	3.1% (0.91-6.1%)	7.4 (2-16)	300 (90-610)
Bahrain	2016	Male	60 to 64	15% (6.4-29%)	85% (71-94%)	0.93 (0.25-2.4)	1.8% (0.42-4%)	2.4% (0.53-5.2%)	4 (0.84-8.8)	130 (30-290)
Bahrain	2016	Male	65 to 69	14% (6.2-26%)	86% (74-94%)	0.76 (0.21-1.8)	1.5% (0.19-3.4%)	1.8% (0.22-4.2%)	3.3 (0.41-8.5)	89 (11-210)
Bahrain	2016	Male	70 to 74	13% (6-24%)	87% (76-94%)	0.57 (0.14-1.5)	1% (0.062-2.9%)	1.2% (0.044-3.4%)	2.4 (0.098-7.2)	50 (3-150)
Bahrain	2016	Male	75 to 79	11% (5-22%)	89% (78-95%)	0.43 (0.099-1.3)	0.7% (-0.027-2.4%)	0.78% (-0.074-2.7%)	1.4 (-0.12-4.6)	22 (-0.97-72)
Bahrain	2016	Male	80 to 84	10% (4-20%)	90% (80-96%)	0.37 (0.093-1.1)	0.57% (-0.076-1.9%)	0.62% (-0.14-2.2%)	0.9 (-0.19-3.3)	11 (-1.3-37)
Bahrain	2016	Male	85 to 89	10% (4.2-20%)	90% (80-96%)	0.34 (0.088-0.93)	0.49% (-0.068-1.8%)	0.51% (-0.13-1.9%)	0.56 (-0.14-2.1)	4.8 (-0.67-17)
Bahrain	2016	Male	90 to 94	12% (4.7-23%)	88% (77-95%)	0.36 (0.091-0.99)	0.42% (-0.16-1.7%)	0.41% (-0.25-1.8%)	0.22 (-0.14-0.95)	1.4 (-0.51-5.2)
Bahrain	2016	Male	95 plus	14% (5.4-29%)	86% (71-95%)	0.42 (0.08-1.3)	0.42% (-0.28-1.8%)	0.36% (-0.46-1.9%)	0.056 (-0.073-0.3)	0.27 (-0.18-1.2)
Egypt	1990	Female	All Ages	1.4% (1-1.8%)	99% (98-99%)	0.21 (0.15-0.28)	0.5% (0.41-0.62%)	1.1% (0.85-1.4%)	2300 (1800-3000)	66000 (53000-81000)
Egypt	1990	Female	15 to 19	1.5% (0.72-2.6%)	99% (97-99%)	0.21 (0.083-0.44)	0.58% (0.37-0.88%)	0.78% (0.42-1.3%)	21 (11-35)	2500 (1600-3800)
Egypt	1990	Female	20 to 24	1.9% (0.92-3.4%)	98% (97-99%)	0.35 (0.15-0.65)	0.94% (0.64-1.3%)	1% (0.56-1.7%)	27 (15-45)	3900 (2600-5700)
Egypt	1990	Female	25 to 29	2% (0.98-3.4%)	98% (97-99%)	0.36 (0.16-0.65)	1% (0.72-1.4%)	1.2% (0.69-1.8%)	35 (21-56)	4300 (3000-6000)
Egypt	1990	Female	30 to 34	1.9% (0.92-3.4%)	98% (97-99%)	0.37 (0.17-0.67)	1% (0.73-1.4%)	1.4% (0.86-2.2%)	39 (24-61)	4100 (2900-5800)
Egypt	1990	Female	35 to 39	2% (0.92-3.5%)	98% (97-99%)	0.4 (0.18-0.76)	1% (0.74-1.5%)	1.5% (0.94-2.4%)	54 (34-84)	4100 (2900-5800)
Egypt	1990	Female	40 to 44	2% (0.97-3.7%)	98% (96-99%)	0.43 (0.18-0.81)	1.1% (0.73-1.5%)	1.8% (1.2-2.7%)	65 (44-100)	4100 (2800-6100)
Egypt	1990	Female	45 to 49	2% (0.89-3.6%)	98% (96-99%)	0.45 (0.18-0.84)	1.2% (0.79-1.8%)	1.9% (1.2-2.9%)	88 (56-140)	4300 (3000-6400)
Egypt	1990	Female	50 to 54	2% (0.93-3.9%)	98% (96-99%)	0.43 (0.17-0.85)	1.3% (0.88-2%)	2% (1.3-2.9%)	150 (94-220)	5800 (3900-8500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	1990	Female	55 to 59	2% (0.94-3.7%)	98% (96-99%)	0.39 (0.15-0.82)	1.3% (0.83-1.9%)	1.9% (1.2-2.8%)	160 (100-250)	5500 (3600-8300)
Egypt	1990	Female	60 to 64	1.9% (0.9-3.7%)	98% (96-99%)	0.35 (0.13-0.73)	1.3% (0.82-1.9%)	1.7% (1.1-2.6%)	220 (150-350)	6300 (4200-9800)
Egypt	1990	Female	65 to 69	1.9% (0.88-3.4%)	98% (97-99%)	0.3 (0.11-0.6)	1.3% (0.87-2%)	1.7% (1.2-2.7%)	280 (190-420)	6400 (4300-9700)
Egypt	1990	Female	70 to 74	1.9% (0.95-3.3%)	98% (97-99%)	0.24 (0.088-0.52)	1.4% (0.94-2%)	1.8% (1.2-2.6%)	330 (220-490)	6000 (4100-8800)
Egypt	1990	Female	75 to 79	1.7% (0.81-3.2%)	98% (97-99%)	0.2 (0.065-0.47)	1.2% (0.81-1.8%)	1.7% (1.1-2.4%)	220 (150-320)	3100 (2100-4600)
Egypt	1990	Female	80 to 84	1.5% (0.73-2.8%)	98% (97-99%)	0.18 (0.057-0.4)	1.4% (0.95-1.9%)	1.7% (1.2-2.4%)	260 (180-360)	2600 (1900-3600)
Egypt	1990	Female	85 to 89	1.5% (0.72-2.7%)	99% (97-99%)	0.16 (0.058-0.35)	1.4% (1-2%)	1.7% (1.2-2.4%)	220 (160-310)	1600 (1100-2200)
Egypt	1990	Female	90 to 94	1.6% (0.82-3%)	98% (97-99%)	0.17 (0.056-0.39)	1.4% (1-2%)	1.7% (1.2-2.5%)	100 (72-140)	500 (360-710)
Egypt	1990	Female	95 plus	2% (0.88-4.1%)	98% (96-99%)	0.2 (0.054-0.51)	1.5% (0.98-2.3%)	1.8% (1.2-2.7%)	34 (22-50)	110 (72-170)
Egypt	1990	Male	All Ages	1.1% (0.81-1.4%)	99% (99-99%)	0.17 (0.12-0.23)	0.69% (0.58-0.81%)	1.2% (1-1.4%)	2900 (2300-3400)	94000 (79000-110000)
Egypt	1990	Male	15 to 19	0.93% (0.44-1.6%)	99% (98-100%)	0.09 (0.033-0.19)	0.65% (0.41-0.98%)	0.59% (0.29-1.1%)	21 (10-40)	3000 (1900-4500)
Egypt	1990	Male	20 to 24	1.4% (0.62-2.4%)	99% (98-99%)	0.2 (0.085-0.38)	1.4% (0.95-1.9%)	1.2% (0.61-2.1%)	42 (21-72)	6100 (3900-8700)
Egypt	1990	Male	25 to 29	1.6% (0.81-2.8%)	98% (97-99%)	0.32 (0.15-0.6)	1.5% (1.1-2.1%)	1.5% (0.83-2.5%)	58 (31-96)	6500 (4300-9000)
Egypt	1990	Male	30 to 34	1.7% (0.87-3%)	98% (97-99%)	0.37 (0.17-0.64)	1.5% (1.1-2.1%)	1.7% (1.1-2.8%)	69 (42-110)	6300 (4300-8800)
Egypt	1990	Male	35 to 39	1.9% (0.91-3.2%)	98% (97-99%)	0.38 (0.17-0.69)	1.6% (1.2-2.2%)	2% (1.3-2.9%)	98 (65-140)	6900 (4900-9400)
Egypt	1990	Male	40 to 44	1.9% (1-3.3%)	98% (97-99%)	0.4 (0.17-0.72)	1.7% (1.3-2.2%)	2.2% (1.6-3%)	130 (94-180)	7600 (5700-10000)
Egypt	1990	Male	45 to 49	1.8% (0.82-3%)	98% (97-99%)	0.43 (0.18-0.8)	1.8% (1.3-2.3%)	2.3% (1.7-3.1%)	160 (120-220)	7600 (5700-9800)
Egypt	1990	Male	50 to 54	1.7% (0.86-2.9%)	98% (97-99%)	0.42 (0.17-0.8)	1.8% (1.3-2.4%)	2.3% (1.6-3%)	250 (180-330)	9600 (7100-13000)
Egypt	1990	Male	55 to 59	1.6% (0.79-2.8%)	98% (97-99%)	0.36 (0.14-0.7)	1.7% (1.2-2.2%)	2.1% (1.5-2.8%)	290 (210-390)	9600 (6900-13000)
Egypt	1990	Male	60 to 64	1.5% (0.7-2.8%)	98% (97-99%)	0.3 (0.11-0.62)	1.5% (1.1-2.1%)	1.9% (1.3-2.6%)	340 (240-480)	9400 (6700-13000)
Egypt	1990	Male	65 to 69	1.5% (0.71-2.6%)	99% (97-99%)	0.25 (0.096-0.52)	1.5% (1.1-2%)	1.8% (1.3-2.3%)	370 (270-500)	8400 (6100-11000)
Egypt	1990	Male	70 to 74	1.4% (0.69-2.4%)	99% (98-99%)	0.19 (0.074-0.42)	1.6% (1.1-2.1%)	1.9% (1.3-2.5%)	360 (260-490)	6500 (4600-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	1990	Male	75 to 79	1.2% (0.57-2.2%)	99% (98-99%)	0.15 (0.049-0.33)	1.4% (0.95-2%)	1.8% (1.2-2.5%)	230 (150-330)	3300 (2100-4500)
Egypt	1990	Male	80 to 84	1.1% (0.51-1.9%)	99% (98-99%)	0.12 (0.042-0.26)	1.6% (0.98-2.2%)	1.9% (1.1-2.6%)	210 (130-290)	2100 (1300-3000)
Egypt	1990	Male	85 to 89	1% (0.49-1.8%)	99% (98-100%)	0.11 (0.038-0.23)	1.6% (0.87-2.1%)	1.8% (0.99-2.5%)	150 (80-200)	1100 (590-1400)
Egypt	1990	Male	90 to 94	1.1% (0.53-2%)	99% (98-99%)	0.11 (0.037-0.24)	1.5% (0.73-2%)	1.7% (0.84-2.4%)	55 (26-75)	270 (130-370)
Egypt	1990	Male	95 plus	1.4% (0.55-2.7%)	99% (97-99%)	0.12 (0.03-0.27)	1.5% (0.71-2.2%)	1.8% (0.78-2.6%)	14 (6.2-20)	46 (21-67)
Egypt	1995	Female	All Ages	1.3% (1-1.7%)	99% (98-99%)	0.21 (0.15-0.29)	0.6% (0.49-0.75%)	1.3% (0.98-1.6%)	2500 (1900-3200)	69000 (55000-86000)
Egypt	1995	Female	15 to 19	1.5% (0.71-2.6%)	99% (97-99%)	0.22 (0.085-0.47)	0.57% (0.36-0.85%)	0.8% (0.42-1.4%)	21 (11-35)	2700 (1700-4000)
Egypt	1995	Female	20 to 24	1.8% (0.85-3.2%)	98% (97-99%)	0.34 (0.14-0.65)	0.92% (0.64-1.3%)	1% (0.55-1.7%)	23 (12-37)	3800 (2500-5400)
Egypt	1995	Female	25 to 29	1.7% (0.82-2.9%)	98% (97-99%)	0.33 (0.14-0.61)	0.97% (0.7-1.3%)	1.1% (0.67-1.7%)	28 (17-43)	4000 (2800-5600)
Egypt	1995	Female	30 to 34	1.6% (0.78-3%)	98% (97-99%)	0.33 (0.15-0.63)	0.96% (0.68-1.3%)	1.3% (0.82-2%)	34 (21-53)	4000 (2800-5600)
Egypt	1995	Female	35 to 39	1.8% (0.83-3.3%)	98% (97-99%)	0.4 (0.17-0.76)	0.99% (0.71-1.4%)	1.5% (0.97-2.3%)	51 (32-77)	4200 (2900-5800)
Egypt	1995	Female	40 to 44	2% (0.94-3.6%)	98% (96-99%)	0.45 (0.19-0.85)	1.1% (0.75-1.5%)	1.8% (1.2-2.8%)	64 (43-97)	4100 (2800-5800)
Egypt	1995	Female	45 to 49	2% (0.86-3.6%)	98% (96-99%)	0.47 (0.19-0.91)	1.1% (0.76-1.7%)	1.9% (1.2-2.9%)	100 (65-160)	5200 (3600-7600)
Egypt	1995	Female	50 to 54	2% (0.93-3.8%)	98% (96-99%)	0.46 (0.18-0.92)	1.3% (0.88-1.9%)	2% (1.3-3%)	140 (92-210)	5700 (3800-8300)
Egypt	1995	Female	55 to 59	2% (0.94-3.5%)	98% (96-99%)	0.42 (0.16-0.84)	1.3% (0.84-1.9%)	1.9% (1.2-2.8%)	160 (110-250)	5600 (3800-8400)
Egypt	1995	Female	60 to 64	1.9% (0.89-3.5%)	98% (97-99%)	0.37 (0.13-0.79)	1.3% (0.85-1.9%)	1.7% (1.2-2.6%)	240 (160-370)	6800 (4500-10000)
Egypt	1995	Female	65 to 69	1.9% (0.92-3.4%)	98% (97-99%)	0.32 (0.12-0.66)	1.3% (0.87-2%)	1.8% (1.2-2.7%)	300 (200-450)	6900 (4600-11000)
Egypt	1995	Female	70 to 74	1.9% (0.94-3.4%)	98% (97-99%)	0.26 (0.09-0.58)	1.4% (0.94-2.1%)	1.8% (1.2-2.6%)	360 (240-530)	6600 (4500-9700)
Egypt	1995	Female	75 to 79	1.7% (0.79-3.3%)	98% (97-99%)	0.21 (0.066-0.5)	1.3% (0.87-1.9%)	1.7% (1.2-2.5%)	300 (200-430)	4200 (2800-6100)
Egypt	1995	Female	80 to 84	1.5% (0.72-2.8%)	98% (97-99%)	0.18 (0.061-0.4)	1.4% (0.94-1.9%)	1.8% (1.2-2.5%)	290 (190-400)	3000 (2000-4100)
Egypt	1995	Female	85 to 89	1.5% (0.72-2.7%)	99% (97-99%)	0.17 (0.061-0.38)	1.4% (1-2%)	1.8% (1.2-2.4%)	220 (160-310)	1600 (1200-2200)
Egypt	1995	Female	90 to 94	1.7% (0.83-3%)	98% (97-99%)	0.19 (0.061-0.41)	1.4% (1-2%)	1.8% (1.2-2.5%)	100 (72-140)	500 (360-710)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	1995	Female	95 plus	2.1% (0.87-4.3%)	98% (96-99%)	0.22 (0.056-0.55)	1.5% (0.98-2.2%)	1.8% (1.2-2.7%)	34 (22-50)	110 (71-160)
Egypt	1995	Male	All Ages	1.1% (0.82-1.3%)	99% (99-99%)	0.18 (0.12-0.24)	0.83% (0.7-0.98%)	1.4% (1.2-1.7%)	3200 (2600-3800)	100000 (88000-120000)
Egypt	1995	Male	15 to 19	0.91% (0.44-1.6%)	99% (98-100%)	0.092 (0.034-0.19)	0.65% (0.42-0.94%)	0.6% (0.3-1.1%)	23 (11-42)	3500 (2200-5100)
Egypt	1995	Male	20 to 24	1.4% (0.65-2.3%)	99% (98-99%)	0.21 (0.088-0.4)	1.4% (0.93-1.9%)	1.2% (0.63-2.1%)	41 (21-71)	6100 (4000-8800)
Egypt	1995	Male	25 to 29	1.7% (0.86-2.8%)	98% (97-99%)	0.34 (0.15-0.63)	1.5% (1.1-2.1%)	1.6% (0.88-2.5%)	55 (30-89)	6400 (4400-8900)
Egypt	1995	Male	30 to 34	1.7% (0.88-3%)	98% (97-99%)	0.38 (0.18-0.67)	1.5% (1.1-2%)	1.8% (1.1-2.7%)	71 (45-110)	6700 (4700-9200)
Egypt	1995	Male	35 to 39	1.7% (0.85-2.9%)	98% (97-99%)	0.35 (0.16-0.65)	1.5% (1.1-2%)	2% (1.3-2.8%)	99 (68-140)	7200 (5300-9500)
Egypt	1995	Male	40 to 44	1.7% (0.93-3%)	98% (97-99%)	0.38 (0.17-0.68)	1.7% (1.3-2.2%)	2.2% (1.6-2.9%)	140 (100-190)	8300 (6200-11000)
Egypt	1995	Male	45 to 49	1.8% (0.85-3%)	98% (97-99%)	0.47 (0.2-0.85)	1.8% (1.4-2.4%)	2.4% (1.8-3.2%)	230 (160-300)	11000 (8000-14000)
Egypt	1995	Male	50 to 54	1.8% (0.92-3%)	98% (97-99%)	0.46 (0.19-0.88)	1.9% (1.4-2.5%)	2.4% (1.7-3.2%)	260 (190-350)	10000 (7500-13000)
Egypt	1995	Male	55 to 59	1.6% (0.8-2.8%)	98% (97-99%)	0.38 (0.14-0.74)	1.7% (1.2-2.3%)	2.1% (1.5-2.9%)	300 (220-410)	10000 (7200-13000)
Egypt	1995	Male	60 to 64	1.5% (0.69-2.7%)	99% (97-99%)	0.31 (0.11-0.64)	1.6% (1.1-2.2%)	1.9% (1.4-2.6%)	370 (270-510)	10000 (7300-14000)
Egypt	1995	Male	65 to 69	1.4% (0.66-2.5%)	99% (98-99%)	0.26 (0.094-0.52)	1.5% (1.1-2%)	1.9% (1.4-2.5%)	410 (300-550)	9300 (6800-12000)
Egypt	1995	Male	70 to 74	1.3% (0.67-2.3%)	99% (98-99%)	0.2 (0.075-0.42)	1.6% (1.1-2.1%)	1.9% (1.4-2.5%)	430 (310-570)	7700 (5600-10000)
Egypt	1995	Male	75 to 79	1.2% (0.55-2.1%)	99% (98-99%)	0.15 (0.05-0.34)	1.5% (1-2.1%)	1.8% (1.2-2.5%)	310 (200-420)	4200 (2900-5900)
Egypt	1995	Male	80 to 84	1% (0.46-1.8%)	99% (98-100%)	0.12 (0.043-0.27)	1.6% (0.99-2.2%)	1.9% (1.2-2.7%)	230 (140-330)	2400 (1500-3300)
Egypt	1995	Male	85 to 89	0.99% (0.47-1.7%)	99% (98-100%)	0.11 (0.038-0.23)	1.6% (0.89-2.1%)	1.9% (1-2.5%)	150 (81-200)	1100 (590-1400)
Egypt	1995	Male	90 to 94	1.1% (0.52-1.9%)	99% (98-99%)	0.11 (0.037-0.23)	1.5% (0.73-2.1%)	1.8% (0.85-2.5%)	52 (25-74)	260 (130-360)
Egypt	1995	Male	95 plus	1.3% (0.53-2.6%)	99% (97-99%)	0.12 (0.031-0.27)	1.5% (0.71-2.3%)	1.8% (0.8-2.7%)	14 (6.1-21)	46 (21-69)
Egypt	2000	Female	All Ages	1.4% (1.1-1.8%)	99% (98-99%)	0.22 (0.16-0.31)	0.69% (0.56-0.85%)	1.3% (1.1-1.7%)	2500 (1900-3200)	74000 (59000-91000)
Egypt	2000	Female	15 to 19	1.5% (0.75-2.6%)	99% (97-99%)	0.22 (0.087-0.44)	0.57% (0.37-0.83%)	0.84% (0.46-1.4%)	19 (11-33)	2900 (1800-4400)
Egypt	2000	Female	20 to 24	1.8% (0.85-3.3%)	98% (97-99%)	0.34 (0.14-0.66)	0.94% (0.64-1.3%)	1.1% (0.59-1.7%)	23 (13-38)	4400 (2800-6300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	2000	Female	25 to 29	1.7% (0.86-3.1%)	98% (97-99%)	0.33 (0.14-0.63)	0.98% (0.7-1.3%)	1.2% (0.71-1.8%)	24 (15-39)	4100 (2800-5700)
Egypt	2000	Female	30 to 34	1.7% (0.8-3.1%)	98% (97-99%)	0.33 (0.14-0.65)	0.96% (0.7-1.3%)	1.4% (0.87-2.1%)	31 (19-48)	4000 (2800-5700)
Egypt	2000	Female	35 to 39	1.9% (0.87-3.3%)	98% (97-99%)	0.4 (0.18-0.76)	0.97% (0.71-1.4%)	1.6% (1-2.4%)	49 (31-75)	4300 (3100-6100)
Egypt	2000	Female	40 to 44	2% (1-3.8%)	98% (96-99%)	0.45 (0.19-0.85)	1% (0.73-1.5%)	1.8% (1.2-2.8%)	69 (46-100)	4600 (3200-6600)
Egypt	2000	Female	45 to 49	2.1% (0.9-3.6%)	98% (96-99%)	0.47 (0.19-0.88)	1.1% (0.78-1.7%)	1.9% (1.2-2.9%)	100 (63-150)	5100 (3500-7600)
Egypt	2000	Female	50 to 54	2.1% (0.97-3.9%)	98% (96-99%)	0.46 (0.18-0.89)	1.3% (0.86-1.9%)	2% (1.3-3%)	180 (120-270)	7500 (5000-11000)
Egypt	2000	Female	55 to 59	2% (0.94-3.9%)	98% (96-99%)	0.41 (0.15-0.86)	1.3% (0.86-2%)	1.9% (1.3-2.9%)	180 (120-270)	6100 (4100-9200)
Egypt	2000	Female	60 to 64	2% (0.89-3.7%)	98% (96-99%)	0.36 (0.13-0.79)	1.3% (0.84-1.9%)	1.7% (1.2-2.6%)	250 (160-380)	7000 (4700-11000)
Egypt	2000	Female	65 to 69	2% (0.93-3.5%)	98% (97-99%)	0.31 (0.11-0.68)	1.3% (0.84-1.9%)	1.8% (1.2-2.7%)	310 (200-470)	7200 (4800-11000)
Egypt	2000	Female	70 to 74	1.9% (0.95-3.4%)	98% (97-99%)	0.25 (0.088-0.55)	1.3% (0.89-1.9%)	1.8% (1.2-2.6%)	390 (260-580)	7200 (4800-11000)
Egypt	2000	Female	75 to 79	1.7% (0.83-3.2%)	98% (97-99%)	0.21 (0.067-0.5)	1.2% (0.85-1.8%)	1.7% (1.1-2.4%)	310 (210-450)	4400 (3000-6300)
Egypt	2000	Female	80 to 84	1.6% (0.75-3%)	98% (97-99%)	0.18 (0.059-0.41)	1.3% (0.88-1.8%)	1.7% (1.1-2.4%)	280 (190-390)	2900 (1900-4100)
Egypt	2000	Female	85 to 89	1.5% (0.72-2.7%)	98% (97-99%)	0.17 (0.062-0.37)	1.3% (0.89-1.7%)	1.7% (1.2-2.3%)	180 (130-250)	1300 (950-1800)
Egypt	2000	Female	90 to 94	1.7% (0.86-3%)	98% (97-99%)	0.18 (0.061-0.4)	1.3% (0.9-1.8%)	1.7% (1.2-2.3%)	89 (63-120)	440 (310-620)
Egypt	2000	Female	95 plus	2% (0.88-4.1%)	98% (96-99%)	0.21 (0.057-0.53)	1.3% (0.86-2%)	1.7% (1.1-2.6%)	29 (18-44)	94 (61-140)
Egypt	2000	Male	All Ages	1.1% (0.84-1.4%)	99% (99-99%)	0.17 (0.12-0.24)	0.94% (0.79-1.1%)	1.5% (1.2-1.8%)	3400 (2800-4000)	11000 (9500-13000)
Egypt	2000	Male	15 to 19	0.91% (0.45-1.5%)	99% (98-100%)	0.086 (0.033-0.17)	0.67% (0.45-0.97%)	0.58% (0.29-1%)	23 (12-41)	3900 (2500-5900)
Egypt	2000	Male	20 to 24	1.4% (0.66-2.2%)	99% (98-99%)	0.2 (0.086-0.37)	1.4% (0.96-2%)	1.2% (0.61-2%)	44 (23-75)	7300 (4900-11000)
Egypt	2000	Male	25 to 29	1.7% (0.86-2.7%)	98% (97-99%)	0.32 (0.14-0.58)	1.5% (1.1-2.1%)	1.6% (0.88-2.5%)	49 (28-80)	6500 (4500-8900)
Egypt	2000	Male	30 to 34	1.7% (0.86-3%)	98% (97-99%)	0.36 (0.17-0.64)	1.5% (1.1-2%)	1.7% (1.1-2.7%)	61 (39-95)	6200 (4400-8500)
Egypt	2000	Male	35 to 39	1.7% (0.85-2.8%)	98% (97-99%)	0.34 (0.15-0.64)	1.5% (1.1-2%)	1.9% (1.4-2.8%)	96 (66-140)	7300 (5500-9800)
Egypt	2000	Male	40 to 44	1.7% (0.92-3%)	98% (97-99%)	0.37 (0.16-0.64)	1.6% (1.3-2.2%)	2.2% (1.6-2.9%)	160 (120-210)	9200 (6900-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	2000	Male	45 to 49	1.8% (0.85-3%)	98% (97-99%)	0.44 (0.18-0.82)	1.8% (1.3-2.3%)	2.4% (1.7-3.1%)	250 (180-330)	12000 (8700-15000)
Egypt	2000	Male	50 to 54	1.8% (0.94-2.9%)	98% (97-99%)	0.44 (0.19-0.84)	1.9% (1.4-2.4%)	2.4% (1.7-3.1%)	350 (250-470)	14000 (10000-18000)
Egypt	2000	Male	55 to 59	1.6% (0.77-2.8%)	98% (97-99%)	0.36 (0.14-0.71)	1.7% (1.2-2.3%)	2.1% (1.5-2.9%)	310 (230-420)	10000 (7600-14000)
Egypt	2000	Male	60 to 64	1.4% (0.68-2.7%)	99% (97-99%)	0.29 (0.1-0.61)	1.5% (1.1-2.1%)	1.9% (1.4-2.6%)	380 (270-520)	10000 (7500-14000)
Egypt	2000	Male	65 to 69	1.4% (0.65-2.5%)	99% (98-99%)	0.24 (0.09-0.48)	1.5% (1.1-2%)	1.9% (1.4-2.4%)	420 (300-560)	9500 (6900-13000)
Egypt	2000	Male	70 to 74	1.3% (0.66-2.2%)	99% (98-99%)	0.18 (0.071-0.41)	1.5% (1.1-2.1%)	1.9% (1.3-2.5%)	460 (320-620)	8200 (5800-11000)
Egypt	2000	Male	75 to 79	1.1% (0.54-2%)	99% (98-99%)	0.14 (0.047-0.33)	1.5% (1-2%)	1.8% (1.2-2.4%)	340 (230-480)	4800 (3300-6600)
Egypt	2000	Male	80 to 84	1% (0.48-1.8%)	99% (98-100%)	0.12 (0.038-0.27)	1.5% (0.96-2.2%)	1.9% (1.2-2.7%)	250 (150-360)	2600 (1600-3600)
Egypt	2000	Male	85 to 89	1% (0.46-1.7%)	99% (98-100%)	0.11 (0.036-0.23)	1.4% (0.82-2%)	1.8% (0.99-2.5%)	120 (69-170)	880 (500-1200)
Egypt	2000	Male	90 to 94	1.1% (0.52-1.9%)	99% (98-99%)	0.11 (0.036-0.23)	1.4% (0.7-1.9%)	1.7% (0.83-2.4%)	48 (24-68)	240 (120-340)
Egypt	2000	Male	95 plus	1.3% (0.56-2.7%)	99% (97-99%)	0.12 (0.029-0.28)	1.4% (0.67-2.2%)	1.8% (0.77-2.7%)	12 (5.4-19)	40 (19-62)
Egypt	2005	Female	All Ages	1.4% (1.1-1.9%)	99% (98-99%)	0.23 (0.17-0.32)	0.78% (0.63-0.97%)	1.5% (1.1-1.9%)	3000 (2300-3900)	87000 (69000-110000)
Egypt	2005	Female	15 to 19	1.5% (0.72-2.7%)	98% (97-99%)	0.22 (0.083-0.44)	0.56% (0.36-0.84%)	0.84% (0.45-1.5%)	19 (9.9-32)	3000 (1900-4600)
Egypt	2005	Female	20 to 24	1.8% (0.83-3.4%)	98% (97-99%)	0.33 (0.14-0.64)	0.95% (0.66-1.3%)	1.1% (0.59-1.8%)	25 (14-42)	5100 (3400-7500)
Egypt	2005	Female	25 to 29	1.7% (0.82-3%)	98% (97-99%)	0.33 (0.14-0.61)	0.98% (0.73-1.3%)	1.2% (0.71-1.9%)	27 (16-43)	4800 (3300-6900)
Egypt	2005	Female	30 to 34	1.7% (0.79-3%)	98% (97-99%)	0.33 (0.15-0.63)	0.96% (0.68-1.3%)	1.4% (0.86-2.1%)	31 (20-49)	4300 (3000-6100)
Egypt	2005	Female	35 to 39	1.9% (0.89-3.5%)	98% (97-99%)	0.4 (0.18-0.76)	0.98% (0.71-1.3%)	1.6% (1-2.4%)	49 (31-74)	4500 (3200-6400)
Egypt	2005	Female	40 to 44	2.1% (1-3.8%)	98% (96-99%)	0.45 (0.19-0.87)	1.1% (0.74-1.5%)	1.9% (1.2-2.9%)	79 (53-120)	5300 (3600-7400)
Egypt	2005	Female	45 to 49	2.1% (0.97-3.7%)	98% (96-99%)	0.47 (0.2-0.94)	1.1% (0.76-1.7%)	1.9% (1.3-2.9%)	120 (75-180)	6100 (4200-9100)
Egypt	2005	Female	50 to 54	2.2% (1-4.2%)	98% (96-99%)	0.47 (0.19-0.92)	1.4% (0.91-2%)	2% (1.3-3%)	210 (140-320)	8600 (5700-13000)
Egypt	2005	Female	55 to 59	2.1% (0.99-4.1%)	98% (96-99%)	0.43 (0.16-0.86)	1.4% (0.87-2.1%)	1.9% (1.2-3%)	270 (170-420)	9200 (6000-14000)
Egypt	2005	Female	60 to 64	2.1% (0.92-3.9%)	98% (96-99%)	0.38 (0.13-0.8)	1.3% (0.86-2%)	1.8% (1.2-2.7%)	280 (180-430)	8000 (5300-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	2005	Female	65 to 69	2.1% (0.97-3.6%)	98% (96-99%)	0.33 (0.12-0.67)	1.3% (0.86-2%)	1.8% (1.2-2.7%)	350 (230-530)	8100 (5400-12000)
Egypt	2005	Female	70 to 74	2% (1-3.6%)	98% (96-99%)	0.27 (0.097-0.58)	1.4% (0.91-2.1%)	1.8% (1.2-2.7%)	450 (300-660)	8300 (5600-12000)
Egypt	2005	Female	75 to 79	1.8% (0.88-3.4%)	98% (97-99%)	0.22 (0.07-0.51)	1.3% (0.85-1.9%)	1.7% (1.1-2.5%)	390 (260-570)	5500 (3700-8100)
Egypt	2005	Female	80 to 84	1.6% (0.75-3.1%)	98% (97-99%)	0.19 (0.062-0.41)	1.3% (0.9-1.9%)	1.7% (1.2-2.4%)	350 (230-500)	3600 (2400-5100)
Egypt	2005	Female	85 to 89	1.6% (0.76-2.8%)	98% (97-99%)	0.18 (0.062-0.41)	1.2% (0.88-1.7%)	1.7% (1.2-2.3%)	210 (150-280)	1500 (1100-2100)
Egypt	2005	Female	90 to 94	1.8% (0.87-3.2%)	98% (97-99%)	0.19 (0.06-0.42)	1.3% (0.88-1.8%)	1.7% (1.2-2.3%)	100 (72-140)	510 (360-720)
Egypt	2005	Female	95 plus	2.2% (0.91-4.4%)	98% (96-99%)	0.22 (0.058-0.54)	1.3% (0.87-2%)	1.7% (1.1-2.6%)	35 (22-53)	110 (73-170)
Egypt	2005	Male	All Ages	1.1% (0.88-1.5%)	99% (99-99%)	0.19 (0.13-0.26)	1% (0.88-1.2%)	1.6% (1.3-1.9%)	4000 (3300-4800)	130000 (110000-160000)
Egypt	2005	Male	15 to 19	0.92% (0.45-1.6%)	99% (98-100%)	0.087 (0.033-0.18)	0.67% (0.43-0.98%)	0.59% (0.29-1%)	25 (12-45)	4300 (2700-6500)
Egypt	2005	Male	20 to 24	1.4% (0.67-2.3%)	99% (98-99%)	0.2 (0.089-0.37)	1.4% (0.97-2%)	1.2% (0.62-2%)	50 (26-85)	8600 (5600-12000)
Egypt	2005	Male	25 to 29	1.7% (0.87-2.8%)	98% (97-99%)	0.33 (0.15-0.59)	1.5% (1.1-2.1%)	1.6% (0.87-2.5%)	57 (31-92)	7900 (5500-11000)
Egypt	2005	Male	30 to 34	1.7% (0.89-3%)	98% (97-99%)	0.37 (0.17-0.64)	1.5% (1.1-2%)	1.8% (1.1-2.7%)	61 (40-93)	6500 (4600-9100)
Egypt	2005	Male	35 to 39	1.7% (0.88-2.9%)	98% (97-99%)	0.34 (0.15-0.64)	1.5% (1.2-2%)	2% (1.4-2.9%)	93 (65-130)	7300 (5500-9500)
Egypt	2005	Male	40 to 44	1.7% (0.93-3%)	98% (97-99%)	0.37 (0.17-0.66)	1.7% (1.3-2.2%)	2.2% (1.6-3%)	170 (130-240)	10000 (7600-13000)
Egypt	2005	Male	45 to 49	1.8% (0.87-3.1%)	98% (97-99%)	0.44 (0.19-0.83)	1.8% (1.4-2.4%)	2.4% (1.8-3.1%)	300 (220-400)	14000 (11000-18000)
Egypt	2005	Male	50 to 54	1.8% (0.94-3%)	98% (97-99%)	0.44 (0.19-0.84)	1.9% (1.4-2.5%)	2.4% (1.7-3.2%)	430 (310-590)	17000 (12000-22000)
Egypt	2005	Male	55 to 59	1.6% (0.79-2.9%)	98% (97-99%)	0.37 (0.14-0.71)	1.7% (1.3-2.4%)	2.2% (1.6-2.9%)	480 (340-670)	16000 (11000-22000)
Egypt	2005	Male	60 to 64	1.5% (0.7-2.7%)	99% (97-99%)	0.3 (0.11-0.62)	1.6% (1.2-2.1%)	1.9% (1.4-2.6%)	420 (300-560)	11000 (8400-15000)
Egypt	2005	Male	65 to 69	1.4% (0.66-2.4%)	99% (98-99%)	0.25 (0.093-0.49)	1.5% (1.1-2%)	1.9% (1.4-2.5%)	460 (340-610)	10000 (7600-14000)
Egypt	2005	Male	70 to 74	1.3% (0.68-2.3%)	99% (98-99%)	0.19 (0.07-0.39)	1.5% (1.1-2%)	1.9% (1.4-2.5%)	500 (350-670)	9000 (6400-12000)
Egypt	2005	Male	75 to 79	1.2% (0.54-2.1%)	99% (98-99%)	0.15 (0.047-0.33)	1.5% (1-2%)	1.8% (1.2-2.5%)	410 (280-570)	5700 (4000-7900)
Egypt	2005	Male	80 to 84	1% (0.48-1.8%)	99% (98-100%)	0.12 (0.038-0.27)	1.5% (0.99-2.2%)	1.9% (1.2-2.6%)	310 (200-440)	3200 (2000-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	2005	Male	85 to 89	1% (0.46-1.8%)	99% (98-100%)	0.11 (0.038-0.23)	1.4% (0.82-2%)	1.8% (1-2.5%)	140 (81-200)	1000 (600-1400)
Egypt	2005	Male	90 to 94	1.1% (0.53-2%)	99% (98-99%)	0.11 (0.036-0.23)	1.3% (0.7-1.9%)	1.7% (0.84-2.4%)	56 (27-78)	280 (140-380)
Egypt	2005	Male	95 plus	1.3% (0.55-2.7%)	99% (97-99%)	0.11 (0.028-0.26)	1.4% (0.67-2.1%)	1.7% (0.77-2.6%)	15 (6.6-23)	49 (22-73)
Egypt	2010	Female	All Ages	1.5% (1.2-2%)	98% (98-99%)	0.24 (0.16-0.32)	0.84% (0.67-1.1%)	1.5% (1.1-1.9%)	3400 (2500-4500)	99000 (77000-120000)
Egypt	2010	Female	15 to 19	1.5% (0.76-2.7%)	98% (97-99%)	0.21 (0.081-0.41)	0.56% (0.36-0.81%)	0.84% (0.46-1.4%)	16 (8.9-28)	2700 (1700-4200)
Egypt	2010	Female	20 to 24	2% (0.96-3.5%)	98% (96-99%)	0.34 (0.14-0.63)	0.97% (0.67-1.4%)	1.1% (0.64-1.8%)	28 (16-45)	5700 (3800-8400)
Egypt	2010	Female	25 to 29	2% (1-3.5%)	98% (97-99%)	0.35 (0.16-0.65)	1% (0.74-1.4%)	1.3% (0.8-2%)	35 (21-54)	6000 (4100-8400)
Egypt	2010	Female	30 to 34	2% (0.97-3.5%)	98% (96-99%)	0.36 (0.16-0.64)	1% (0.73-1.4%)	1.5% (0.93-2.3%)	42 (25-63)	5500 (3900-7600)
Egypt	2010	Female	35 to 39	2.1% (0.99-3.7%)	98% (96-99%)	0.39 (0.17-0.76)	0.99% (0.73-1.3%)	1.6% (1.1-2.4%)	52 (34-79)	5000 (3500-6800)
Egypt	2010	Female	40 to 44	2.1% (1.1-4%)	98% (96-99%)	0.43 (0.18-0.81)	1.1% (0.75-1.5%)	1.9% (1.3-2.8%)	85 (56-130)	5700 (3900-8100)
Egypt	2010	Female	45 to 49	2.2% (0.98-4%)	98% (96-99%)	0.45 (0.18-0.87)	1.2% (0.78-1.7%)	1.9% (1.2-3%)	140 (88-210)	7100 (4800-11000)
Egypt	2010	Female	50 to 54	2.2% (1-4.2%)	98% (96-99%)	0.44 (0.17-0.85)	1.3% (0.89-2%)	2% (1.3-3%)	240 (150-360)	9800 (6500-15000)
Egypt	2010	Female	55 to 59	2.1% (1-4%)	98% (96-99%)	0.4 (0.15-0.8)	1.4% (0.89-2.1%)	1.9% (1.2-2.9%)	310 (190-470)	10000 (6700-16000)
Egypt	2010	Female	60 to 64	2.1% (0.95-4%)	98% (96-99%)	0.35 (0.12-0.74)	1.3% (0.82-2%)	1.7% (1.1-2.6%)	380 (250-590)	11000 (7100-17000)
Egypt	2010	Female	65 to 69	2.1% (0.99-3.6%)	98% (96-99%)	0.3 (0.11-0.62)	1.3% (0.84-2%)	1.8% (1.1-2.7%)	380 (240-580)	8700 (5600-13000)
Egypt	2010	Female	70 to 74	2% (1-3.5%)	98% (96-99%)	0.24 (0.082-0.53)	1.3% (0.88-2%)	1.7% (1.1-2.6%)	460 (300-710)	8500 (5600-13000)
Egypt	2010	Female	75 to 79	1.8% (0.85-3.5%)	98% (97-99%)	0.2 (0.06-0.48)	1.2% (0.81-1.8%)	1.6% (1.1-2.4%)	420 (270-620)	5900 (3900-8700)
Egypt	2010	Female	80 to 84	1.7% (0.74-3.1%)	98% (97-99%)	0.17 (0.053-0.37)	1.3% (0.85-1.9%)	1.7% (1.1-2.4%)	390 (260-570)	4100 (2600-5800)
Egypt	2010	Female	85 to 89	1.6% (0.75-2.8%)	98% (97-99%)	0.16 (0.057-0.36)	1.2% (0.85-1.7%)	1.6% (1.1-2.2%)	250 (180-350)	1800 (1300-2600)
Egypt	2010	Female	90 to 94	1.8% (0.88-3.1%)	98% (97-99%)	0.17 (0.055-0.39)	1.2% (0.84-1.7%)	1.6% (1.1-2.3%)	120 (84-170)	610 (420-880)
Egypt	2010	Female	95 plus	2.2% (0.93-4.4%)	98% (96-99%)	0.2 (0.054-0.51)	1.3% (0.82-2%)	1.7% (1.1-2.6%)	42 (27-65)	140 (90-210)
Egypt	2010	Male	All Ages	1.2% (0.92-1.6%)	99% (98-99%)	0.18 (0.12-0.25)	1.1% (0.92-1.3%)	1.6% (1.3-1.9%)	4500 (3600-5500)	150000 (130000-180000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	2010	Male	15 to 19	0.95% (0.44-1.7%)	99% (98-100%)	0.082 (0.028-0.17)	0.67% (0.43-1%)	0.57% (0.29-1.1%)	26 (13-47)	4200 (2600-6400)
Egypt	2010	Male	20 to 24	1.4% (0.65-2.5%)	99% (98-99%)	0.19 (0.078-0.35)	1.4% (0.98-2%)	1.1% (0.59-2%)	61 (31-110)	10000 (6600-15000)
Egypt	2010	Male	25 to 29	1.7% (0.86-2.8%)	98% (97-99%)	0.29 (0.13-0.54)	1.6% (1.1-2.1%)	1.5% (0.84-2.4%)	73 (40-120)	10000 (6800-14000)
Egypt	2010	Male	30 to 34	1.8% (0.87-3.2%)	98% (97-99%)	0.35 (0.16-0.63)	1.5% (1.1-2.1%)	1.7% (1.1-2.7%)	80 (50-120)	8900 (6300-12000)
Egypt	2010	Male	35 to 39	1.9% (0.96-3.4%)	98% (97-99%)	0.36 (0.16-0.67)	1.6% (1.2-2.1%)	2.1% (1.5-3%)	100 (70-150)	8300 (6000-11000)
Egypt	2010	Male	40 to 44	1.9% (1.1-3.5%)	98% (97-99%)	0.38 (0.17-0.7)	1.7% (1.3-2.2%)	2.3% (1.7-3.1%)	170 (130-240)	11000 (7900-14000)
Egypt	2010	Male	45 to 49	1.8% (0.85-3.1%)	98% (97-99%)	0.41 (0.17-0.75)	1.8% (1.3-2.3%)	2.4% (1.7-3.1%)	320 (230-430)	15000 (11000-20000)
Egypt	2010	Male	50 to 54	1.7% (0.87-3%)	98% (97-99%)	0.39 (0.15-0.74)	1.9% (1.3-2.5%)	2.3% (1.7-3.1%)	490 (340-670)	19000 (14000-25000)
Egypt	2010	Male	55 to 59	1.6% (0.78-2.9%)	98% (97-99%)	0.34 (0.13-0.65)	1.7% (1.2-2.4%)	2.1% (1.4-2.9%)	560 (380-780)	18000 (13000-25000)
Egypt	2010	Male	60 to 64	1.5% (0.74-2.8%)	98% (97-99%)	0.29 (0.1-0.6)	1.6% (1.1-2.1%)	1.9% (1.4-2.6%)	590 (410-810)	16000 (12000-22000)
Egypt	2010	Male	65 to 69	1.5% (0.72-2.6%)	98% (97-99%)	0.24 (0.088-0.46)	1.5% (1.1-2%)	1.8% (1.3-2.4%)	480 (350-640)	11000 (8000-15000)
Egypt	2010	Male	70 to 74	1.4% (0.72-2.4%)	99% (98-99%)	0.18 (0.063-0.4)	1.5% (1.1-2%)	1.8% (1.3-2.4%)	510 (360-690)	9300 (6500-12000)
Egypt	2010	Male	75 to 79	1.2% (0.58-2.1%)	99% (98-99%)	0.14 (0.044-0.32)	1.4% (0.94-2%)	1.7% (1.1-2.4%)	420 (280-590)	5800 (3900-8200)
Egypt	2010	Male	80 to 84	1.1% (0.53-1.9%)	99% (98-99%)	0.11 (0.038-0.25)	1.5% (0.92-2.1%)	1.8% (1.1-2.5%)	340 (210-490)	3500 (2200-4900)
Egypt	2010	Male	85 to 89	1% (0.48-1.9%)	99% (98-100%)	0.1 (0.035-0.22)	1.4% (0.75-1.9%)	1.7% (0.89-2.4%)	190 (97-260)	1300 (720-1800)
Egypt	2010	Male	90 to 94	1.1% (0.54-2%)	99% (98-99%)	0.099 (0.033-0.22)	1.3% (0.64-1.8%)	1.6% (0.75-2.3%)	69 (32-98)	340 (170-480)
Egypt	2010	Male	95 plus	1.4% (0.57-2.8%)	99% (97-99%)	0.11 (0.027-0.25)	1.3% (0.58-2%)	1.7% (0.67-2.6%)	18 (7.2-28)	59 (26-90)
Egypt	2016	Female	All Ages	1.5% (1.2-2%)	98% (98-99%)	0.24 (0.13-0.39)	0.87% (0.7-1.1%)	1.5% (1.1-1.9%)	3400 (2500-4600)	100000 (76000-130000)
Egypt	2016	Female	15 to 19	1.5% (0.76-2.9%)	98% (97-99%)	0.21 (0.073-0.45)	0.53% (0.34-0.78%)	0.78% (0.41-1.3%)	13 (6.8-23)	2500 (1500-3800)
Egypt	2016	Female	20 to 24	2% (0.93-3.6%)	98% (96-99%)	0.34 (0.13-0.73)	0.96% (0.65-1.3%)	1.1% (0.61-1.7%)	21 (11-34)	4900 (3200-7200)
Egypt	2016	Female	25 to 29	2% (1-3.6%)	98% (96-99%)	0.36 (0.14-0.7)	1% (0.73-1.4%)	1.2% (0.73-1.9%)	31 (17-49)	6300 (4300-8800)
Egypt	2016	Female	30 to 34	2% (0.94-3.6%)	98% (96-99%)	0.36 (0.15-0.74)	0.99% (0.72-1.3%)	1.4% (0.91-2.1%)	42 (26-64)	6300 (4500-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	2016	Female	35 to 39	2% (0.98-3.6%)	98% (96-99%)	0.39 (0.15-0.83)	0.95% (0.69-1.3%)	1.6% (1-2.3%)	55 (35-85)	5800 (4000-8000)
Egypt	2016	Female	40 to 44	2.1% (1-3.9%)	98% (96-99%)	0.43 (0.17-0.89)	1% (0.71-1.4%)	1.8% (1.2-2.6%)	80 (51-130)	5800 (4000-8400)
Egypt	2016	Female	45 to 49	2.1% (0.91-3.8%)	98% (96-99%)	0.45 (0.15-0.94)	1.1% (0.73-1.5%)	1.8% (1.2-2.7%)	130 (78-200)	6900 (4600-10000)
Egypt	2016	Female	50 to 54	2.2% (1-4.1%)	98% (96-99%)	0.44 (0.15-0.92)	1.2% (0.81-1.8%)	1.9% (1.2-2.8%)	230 (140-350)	9700 (6200-14000)
Egypt	2016	Female	55 to 59	2.1% (0.96-4%)	98% (96-99%)	0.4 (0.13-0.91)	1.3% (0.84-2%)	1.8% (1.2-2.8%)	330 (210-510)	11000 (7100-17000)
Egypt	2016	Female	60 to 64	2.1% (0.95-4%)	98% (96-99%)	0.35 (0.11-0.82)	1.2% (0.79-1.8%)	1.7% (1.1-2.5%)	350 (220-570)	10000 (6500-16000)
Egypt	2016	Female	65 to 69	2.1% (0.92-3.5%)	98% (97-99%)	0.3 (0.093-0.68)	1.2% (0.81-1.9%)	1.7% (1.1-2.5%)	440 (280-690)	10000 (6700-16000)
Egypt	2016	Female	70 to 74	2% (1-3.6%)	98% (96-99%)	0.25 (0.073-0.59)	1.3% (0.84-1.9%)	1.7% (1.1-2.5%)	460 (290-700)	8500 (5400-13000)
Egypt	2016	Female	75 to 79	1.8% (0.85-3.5%)	98% (97-99%)	0.21 (0.053-0.5)	1.2% (0.76-1.7%)	1.6% (1-2.3%)	370 (230-550)	5300 (3400-7800)
Egypt	2016	Female	80 to 84	1.6% (0.71-3%)	98% (97-99%)	0.18 (0.05-0.42)	1.3% (0.82-1.8%)	1.7% (1.1-2.3%)	390 (250-550)	4000 (2500-5700)
Egypt	2016	Female	85 to 89	1.6% (0.73-2.8%)	98% (97-99%)	0.16 (0.048-0.39)	1.2% (0.82-1.7%)	1.6% (1.1-2.2%)	270 (180-380)	2000 (1300-2800)
Egypt	2016	Female	90 to 94	1.7% (0.87-3.1%)	98% (97-99%)	0.17 (0.05-0.41)	1.2% (0.81-1.7%)	1.6% (1.1-2.2%)	150 (97-210)	730 (490-1000)
Egypt	2016	Female	95 plus	2.2% (0.92-4.4%)	98% (96-99%)	0.2 (0.049-0.53)	1.3% (0.77-1.9%)	1.7% (1-2.5%)	53 (32-80)	170 (110-260)
Egypt	2016	Male	All Ages	1.2% (0.91-1.5%)	99% (98-99%)	0.19 (0.1-0.3)	1.2% (0.98-1.4%)	1.7% (1.4-2%)	5000 (3700-6200)	170000 (130000-200000)
Egypt	2016	Male	15 to 19	0.95% (0.45-1.7%)	99% (98-100%)	0.083 (0.026-0.19)	0.67% (0.43-0.97%)	0.55% (0.27-1%)	23 (11-46)	4000 (2500-6000)
Egypt	2016	Male	20 to 24	1.4% (0.63-2.3%)	99% (98-99%)	0.19 (0.069-0.39)	1.4% (0.97-2%)	1.1% (0.57-1.8%)	51 (26-90)	9000 (5700-13000)
Egypt	2016	Male	25 to 29	1.6% (0.81-2.8%)	98% (97-99%)	0.29 (0.12-0.58)	1.5% (1.1-2%)	1.4% (0.78-2.1%)	71 (39-120)	11000 (7300-15000)
Egypt	2016	Male	30 to 34	1.8% (0.85-3.1%)	98% (97-99%)	0.34 (0.14-0.71)	1.5% (1.1-2%)	1.6% (1-2.5%)	88 (55-140)	10000 (7300-14000)
Egypt	2016	Male	35 to 39	1.9% (0.97-3.2%)	98% (97-99%)	0.36 (0.14-0.72)	1.6% (1.2-2%)	2% (1.4-2.7%)	120 (80-180)	11000 (7800-14000)
Egypt	2016	Male	40 to 44	1.9% (1-3.4%)	98% (97-99%)	0.38 (0.14-0.75)	1.7% (1.3-2.2%)	2.2% (1.7-3%)	180 (130-260)	11000 (8400-15000)
Egypt	2016	Male	45 to 49	1.8% (0.81-3.1%)	98% (97-99%)	0.4 (0.14-0.79)	1.8% (1.3-2.3%)	2.3% (1.7-3.1%)	310 (220-440)	15000 (11000-20000)
Egypt	2016	Male	50 to 54	1.7% (0.86-2.9%)	98% (97-99%)	0.38 (0.14-0.77)	1.8% (1.3-2.4%)	2.3% (1.6-3.1%)	510 (340-720)	20000 (14000-27000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Egypt	2016	Male	55 to 59	1.6% (0.79-2.8%)	98% (97-99%)	0.33 (0.12-0.7)	1.7% (1.2-2.3%)	2.1% (1.4-2.8%)	630 (410-900)	21000 (14000-29000)
Egypt	2016	Male	60 to 64	1.5% (0.71-2.7%)	98% (97-99%)	0.28 (0.087-0.64)	1.6% (1.1-2.1%)	1.9% (1.4-2.6%)	620 (420-880)	17000 (12000-24000)
Egypt	2016	Male	65 to 69	1.5% (0.7-2.5%)	99% (97-99%)	0.23 (0.075-0.49)	1.5% (1-2%)	1.8% (1.3-2.4%)	650 (440-900)	15000 (10000-20000)
Egypt	2016	Male	70 to 74	1.4% (0.68-2.4%)	99% (98-99%)	0.18 (0.053-0.41)	1.5% (1-2%)	1.8% (1.3-2.5%)	550 (370-760)	10000 (6700-14000)
Egypt	2016	Male	75 to 79	1.2% (0.56-2.1%)	99% (98-99%)	0.13 (0.035-0.32)	1.4% (0.9-2%)	1.8% (1.2-2.4%)	420 (250-610)	5800 (3600-8300)
Egypt	2016	Male	80 to 84	1.1% (0.48-1.9%)	99% (98-100%)	0.11 (0.033-0.25)	1.5% (0.92-2.1%)	1.8% (1.1-2.6%)	380 (230-550)	3800 (2400-5500)
Egypt	2016	Male	85 to 89	1% (0.48-1.9%)	99% (98-100%)	0.1 (0.029-0.25)	1.4% (0.74-1.9%)	1.8% (0.89-2.4%)	220 (110-300)	1600 (800-2200)
Egypt	2016	Male	90 to 94	1.1% (0.54-2%)	99% (98-99%)	0.1 (0.028-0.23)	1.3% (0.62-1.8%)	1.6% (0.77-2.3%)	89 (42-130)	440 (210-620)
Egypt	2016	Male	95 plus	1.4% (0.56-2.8%)	99% (97-99%)	0.11 (0.025-0.28)	1.4% (0.6-2.1%)	1.7% (0.7-2.6%)	24 (9.8-37)	79 (34-120)
Iran	1990	Female	All Ages	9.3% (5.6-14%)	91% (86-94%)	0.0015 (0.00058-0.003)	0.092% (0.067-0.12%)	0.062% (0.038-0.085%)	86 (53-120)	10000 (7100-14000)
Iran	1990	Female	20 to 24	11% (3.1-24%)	89% (76-97%)	0.002 (0.00043-0.0053)	0.35% (0.2-0.56%)	0.043% (0.024-0.072%)	1.2 (0.66-2)	1500 (790-2500)
Iran	1990	Female	25 to 29	12% (3.4-27%)	88% (73-97%)	0.0024 (0.0005-0.0065)	0.39% (0.24-0.58%)	0.052% (0.032-0.084%)	1.4 (0.85-2.3)	1600 (950-2500)
Iran	1990	Female	30 to 34	14% (3.5-30%)	86% (70-97%)	0.0029 (0.00063-0.0073)	0.37% (0.22-0.55%)	0.09% (0.053-0.15%)	2.5 (1.4-4.5)	1400 (830-2300)
Iran	1990	Female	35 to 39	16% (4.4-30%)	84% (70-96%)	0.0034 (0.00079-0.0084)	0.3% (0.19-0.44%)	0.11% (0.067-0.17%)	3.4 (1.9-5.4)	1100 (660-1600)
Iran	1990	Female	40 to 44	16% (5.5-30%)	84% (70-94%)	0.0038 (0.00095-0.0095)	0.26% (0.17-0.37%)	0.17% (0.091-0.28%)	4.8 (2.4-8)	750 (460-1100)
Iran	1990	Female	45 to 49	17% (5.5-34%)	83% (66-94%)	0.0042 (0.00086-0.011)	0.22% (0.15-0.3%)	0.18% (0.094-0.28%)	5.1 (2.6-8.6)	550 (360-790)
Iran	1990	Female	50 to 54	18% (6.5-36%)	82% (64-93%)	0.0043 (0.00092-0.012)	0.18% (0.12-0.25%)	0.17% (0.1-0.25%)	6.6 (3.5-10)	510 (330-730)
Iran	1990	Female	55 to 59	19% (6.9-38%)	81% (62-93%)	0.0043 (0.00089-0.012)	0.15% (0.1-0.21%)	0.16% (0.09-0.24%)	8 (4.3-13)	450 (290-630)
Iran	1990	Female	60 to 64	19% (7.3-38%)	81% (62-93%)	0.0041 (0.0008-0.012)	0.14% (0.093-0.2%)	0.16% (0.09-0.26%)	10 (5.5-17)	410 (250-610)
Iran	1990	Female	65 to 69	20% (8.5-36%)	80% (64-91%)	0.0035 (0.00066-0.01)	0.13% (0.086-0.2%)	0.16% (0.093-0.26%)	13 (7-22)	370 (240-560)
Iran	1990	Female	70 to 74	18% (7.7-34%)	82% (66-92%)	0.0028 (0.00054-0.0083)	0.12% (0.075-0.18%)	0.14% (0.081-0.23%)	11 (6-18)	230 (140-350)
Iran	1990	Female	75 to 79	17% (6.2-34%)	83% (66-94%)	0.0023 (0.00035-0.0078)	0.1% (0.063-0.15%)	0.12% (0.07-0.2%)	8.1 (4.5-13)	130 (78-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	1990	Female	80 to 84	15% (4.9-34%)	85% (66-95%)	0.0021 (0.00029-0.0068)	0.082% (0.054-0.12%)	0.096% (0.058-0.15%)	4.7 (2.8-7.6)	55 (35-83)
Iran	1990	Female	85 to 89	16% (4.6-37%)	84% (63-95%)	0.0021 (0.0003-0.0077)	0.071% (0.046-0.11%)	0.08% (0.048-0.13%)	3.5 (2.1-6)	29 (19-47)
Iran	1990	Female	90 to 94	18% (5.4-40%)	82% (60-95%)	0.0024 (0.00035-0.0085)	0.061% (0.038-0.098%)	0.065% (0.037-0.11%)	1.5 (0.88-2.8)	9.1 (5.7-15)
Iran	1990	Female	95 plus	22% (6.3-48%)	78% (52-94%)	0.0028 (0.00036-0.01)	0.064% (0.04-0.11%)	0.06% (0.032-0.11%)	0.5 (0.26-0.92)	2.2 (1.4-3.6)
Iran	1990	Male	All Ages	7% (4.8-9.7%)	93% (90-95%)	0.0017 (0.00062-0.0033)	0.14% (0.11-0.19%)	0.099% (0.074-0.13%)	200 (150-270)	20000 (15000-26000)
Iran	1990	Male	20 to 24	9.3% (3.5-18%)	91% (82-96%)	0.0019 (0.00047-0.0047)	0.49% (0.29-0.79%)	0.033% (0.016-0.059%)	2 (1-3.4)	3000 (1800-4700)
Iran	1990	Male	25 to 29	11% (4.3-22%)	89% (78-96%)	0.003 (0.00072-0.0074)	0.49% (0.31-0.73%)	0.057% (0.029-0.11%)	3.2 (1.7-6)	2800 (1700-4200)
Iran	1990	Male	30 to 34	13% (4.6-25%)	87% (75-95%)	0.0038 (0.00098-0.0092)	0.46% (0.31-0.67%)	0.098% (0.057-0.17%)	5.5 (3.3-9.5)	2400 (1500-3500)
Iran	1990	Male	35 to 39	14% (5.2-27%)	86% (73-95%)	0.0043 (0.0011-0.011)	0.45% (0.32-0.62%)	0.18% (0.1-0.3%)	10 (6.2-17)	2200 (1400-3100)
Iran	1990	Male	40 to 44	14% (6.3-26%)	86% (74-94%)	0.0046 (0.0012-0.012)	0.41% (0.28-0.57%)	0.22% (0.15-0.33%)	11 (7.3-17)	1500 (1000-2100)
Iran	1990	Male	45 to 49	14% (5.7-27%)	86% (73-94%)	0.0049 (0.0011-0.013)	0.39% (0.28-0.5%)	0.29% (0.2-0.43%)	15 (9.5-23)	1300 (880-1700)
Iran	1990	Male	50 to 54	14% (5.9-27%)	86% (73-94%)	0.0049 (0.0012-0.013)	0.33% (0.24-0.43%)	0.28% (0.2-0.4%)	19 (12-28)	1200 (860-1600)
Iran	1990	Male	55 to 59	13% (5.4-26%)	87% (74-95%)	0.0046 (0.001-0.012)	0.27% (0.2-0.35%)	0.26% (0.18-0.38%)	22 (14-34)	1000 (740-1500)
Iran	1990	Male	60 to 64	13% (5.3-25%)	87% (75-95%)	0.0042 (0.00086-0.012)	0.24% (0.17-0.32%)	0.25% (0.17-0.37%)	27 (17-43)	940 (660-1400)
Iran	1990	Male	65 to 69	12% (5.4-23%)	88% (77-95%)	0.0034 (0.00073-0.0093)	0.21% (0.15-0.28%)	0.23% (0.16-0.33%)	29 (18-46)	790 (530-1100)
Iran	1990	Male	70 to 74	11% (4.9-20%)	89% (80-95%)	0.0026 (0.00049-0.0073)	0.17% (0.12-0.23%)	0.19% (0.13-0.28%)	22 (14-34)	450 (310-670)
Iran	1990	Male	75 to 79	9.5% (4.1-18%)	91% (82-96%)	0.0019 (0.00035-0.0059)	0.14% (0.1-0.19%)	0.15% (0.11-0.22%)	15 (10-22)	240 (170-340)
Iran	1990	Male	80 to 84	8.4% (3.3-17%)	92% (83-97%)	0.0016 (0.0003-0.0048)	0.11% (0.078-0.16%)	0.12% (0.082-0.18%)	9.3 (5.9-14)	110 (72-150)
Iran	1990	Male	85 to 89	8.3% (3.3-17%)	92% (83-97%)	0.0015 (0.00029-0.0045)	0.094% (0.068-0.13%)	0.098% (0.068-0.15%)	4.6 (3.1-7)	40 (28-56)
Iran	1990	Male	90 to 94	9.4% (3.8-20%)	91% (80-96%)	0.0016 (0.00028-0.0048)	0.08% (0.058-0.11%)	0.078% (0.052-0.12%)	1.5 (0.99-2.4)	9.5 (6.8-14)
Iran	1990	Male	95 plus	12% (4.2-25%)	88% (75-96%)	0.0018 (0.00026-0.0064)	0.087% (0.063-0.12%)	0.068% (0.044-0.1%)	0.36 (0.23-0.55)	1.9 (1.3-2.6)
Iran	1995	Female	All Ages	9.7% (6.2-15%)	90% (85-94%)	0.002 (0.0009-0.004)	0.14% (0.11-0.18%)	0.096% (0.066-0.12%)	110 (71-140)	12000 (8300-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	1995	Female	20 to 24	11% (3.3-24%)	89% (76-97%)	0.0026 (0.00065-0.0068)	0.42% (0.24-0.66%)	0.074% (0.048-0.11%)	1.4 (0.86-2.2)	1700 (910-2800)
Iran	1995	Female	25 to 29	12% (3.4-26%)	88% (74-97%)	0.0031 (0.00077-0.0078)	0.45% (0.29-0.67%)	0.085% (0.058-0.12%)	1.7 (1-2.6)	1700 (1000-2700)
Iran	1995	Female	30 to 34	14% (3.5-30%)	86% (70-96%)	0.0037 (0.00093-0.0092)	0.42% (0.26-0.62%)	0.14% (0.087-0.21%)	2.9 (1.8-4.8)	1600 (930-2500)
Iran	1995	Female	35 to 39	15% (4.3-30%)	85% (70-96%)	0.0044 (0.0011-0.011)	0.34% (0.22-0.47%)	0.16% (0.1-0.23%)	4.1 (2.4-6.2)	1200 (770-1800)
Iran	1995	Female	40 to 44	16% (5.8-30%)	84% (70-94%)	0.0049 (0.0014-0.011)	0.29% (0.18-0.42%)	0.22% (0.13-0.33%)	6.6 (3.7-11)	1000 (620-1600)
Iran	1995	Female	45 to 49	17% (6-32%)	83% (68-94%)	0.0054 (0.0014-0.014)	0.24% (0.16-0.32%)	0.22% (0.12-0.33%)	6.5 (3.6-11)	690 (440-970)
Iran	1995	Female	50 to 54	18% (6.6-35%)	82% (65-93%)	0.0057 (0.0014-0.015)	0.2% (0.14-0.27%)	0.2% (0.12-0.29%)	6.7 (3.8-10)	500 (330-710)
Iran	1995	Female	55 to 59	19% (7.2-37%)	81% (63-93%)	0.0057 (0.0013-0.015)	0.17% (0.11-0.23%)	0.18% (0.11-0.28%)	8.5 (5-13)	460 (320-650)
Iran	1995	Female	60 to 64	20% (7.6-39%)	80% (61-92%)	0.0054 (0.0012-0.016)	0.15% (0.1-0.21%)	0.18% (0.11-0.27%)	12 (7.2-20)	470 (300-680)
Iran	1995	Female	65 to 69	20% (8.2-37%)	80% (63-92%)	0.0047 (0.00099-0.013)	0.14% (0.095-0.21%)	0.18% (0.11-0.27%)	15 (9-24)	420 (270-610)
Iran	1995	Female	70 to 74	19% (7.8-34%)	81% (66-92%)	0.0037 (0.00077-0.0098)	0.12% (0.084-0.19%)	0.16% (0.096-0.25%)	16 (9.6-26)	340 (220-510)
Iran	1995	Female	75 to 79	17% (6-35%)	83% (65-94%)	0.003 (0.00048-0.0095)	0.11% (0.072-0.16%)	0.13% (0.081-0.21%)	11 (6.8-17)	180 (120-260)
Iran	1995	Female	80 to 84	15% (5.2-34%)	85% (66-95%)	0.0027 (0.00041-0.0084)	0.088% (0.06-0.13%)	0.1% (0.067-0.16%)	6.1 (3.9-9.3)	70 (47-100)
Iran	1995	Female	85 to 89	16% (5.2-35%)	84% (65-95%)	0.0027 (0.00043-0.0092)	0.076% (0.051-0.12%)	0.086% (0.054-0.14%)	4.2 (2.6-6.9)	34 (22-53)
Iran	1995	Female	90 to 94	19% (6.1-41%)	81% (59-94%)	0.0031 (0.0005-0.01)	0.065% (0.044-0.1%)	0.071% (0.043-0.12%)	2 (1.2-3.4)	12 (7.7-18)
Iran	1995	Female	95 plus	23% (7.2-47%)	77% (53-93%)	0.0037 (0.00049-0.013)	0.068% (0.044-0.1%)	0.064% (0.036-0.11%)	0.65 (0.36-1.1)	2.8 (1.8-4.3)
Iran	1995	Male	All Ages	7.3% (5.1-10%)	93% (90-95%)	0.0022 (0.00099-0.0042)	0.22% (0.17-0.28%)	0.15% (0.12-0.18%)	250 (190-320)	23000 (17000-30000)
Iran	1995	Male	20 to 24	9.2% (3.5-18%)	91% (82-97%)	0.0024 (0.00067-0.0058)	0.62% (0.39-0.95%)	0.057% (0.034-0.092%)	2.5 (1.5-4.2)	3300 (2000-5400)
Iran	1995	Male	25 to 29	12% (4.6-22%)	88% (78-95%)	0.0039 (0.0011-0.0094)	0.61% (0.4-0.87%)	0.094% (0.059-0.16%)	4.1 (2.4-7.1)	3000 (2000-4500)
Iran	1995	Male	30 to 34	13% (4.9-26%)	87% (74-95%)	0.0049 (0.0015-0.011)	0.55% (0.37-0.8%)	0.15% (0.1-0.26%)	6.8 (4.2-12)	2600 (1700-3900)
Iran	1995	Male	35 to 39	15% (5.5-27%)	85% (73-95%)	0.0055 (0.0016-0.013)	0.53% (0.37-0.72%)	0.26% (0.18-0.41%)	13 (8.2-21)	2500 (1600-3500)
Iran	1995	Male	40 to 44	15% (6.4-27%)	85% (73-94%)	0.0059 (0.0018-0.014)	0.48% (0.34-0.64%)	0.3% (0.21-0.43%)	17 (11-26)	2200 (1500-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	1995	Male	45 to 49	14% (5.9-27%)	86% (73-94%)	0.0062 (0.0016-0.016)	0.45% (0.34-0.57%)	0.38% (0.27-0.52%)	20 (14-31)	1600 (1200-2200)
Iran	1995	Male	50 to 54	14% (5.8-27%)	86% (73-94%)	0.0061 (0.0015-0.016)	0.37% (0.28-0.46%)	0.34% (0.25-0.47%)	20 (13-30)	1200 (870-1600)
Iran	1995	Male	55 to 59	13% (5.4-26%)	87% (74-95%)	0.0058 (0.0015-0.014)	0.3% (0.22-0.38%)	0.3% (0.21-0.41%)	25 (16-35)	1100 (800-1500)
Iran	1995	Male	60 to 64	13% (5.2-25%)	87% (75-95%)	0.0052 (0.0013-0.014)	0.26% (0.19-0.34%)	0.28% (0.2-0.39%)	33 (22-48)	1100 (790-1500)
Iran	1995	Male	65 to 69	12% (5.2-23%)	88% (77-95%)	0.0043 (0.00099-0.011)	0.22% (0.17-0.29%)	0.25% (0.18-0.34%)	35 (23-49)	930 (680-1200)
Iran	1995	Male	70 to 74	11% (5-20%)	89% (80-95%)	0.0032 (0.00072-0.0082)	0.18% (0.14-0.24%)	0.21% (0.15-0.29%)	33 (22-47)	680 (480-930)
Iran	1995	Male	75 to 79	9.5% (4-18%)	90% (82-96%)	0.0024 (0.00047-0.0073)	0.15% (0.11-0.2%)	0.17% (0.12-0.23%)	20 (15-28)	320 (240-420)
Iran	1995	Male	80 to 84	8.4% (3.2-17%)	92% (83-97%)	0.002 (0.00037-0.0059)	0.12% (0.089-0.16%)	0.13% (0.094-0.18%)	12 (8.4-16)	140 (100-180)
Iran	1995	Male	85 to 89	8.2% (3.2-16%)	92% (84-97%)	0.0019 (0.00038-0.0056)	0.1% (0.077-0.14%)	0.11% (0.077-0.15%)	6.1 (4.3-8.5)	51 (37-68)
Iran	1995	Male	90 to 94	9.3% (3.8-19%)	91% (81-96%)	0.002 (0.00039-0.0057)	0.087% (0.065-0.12%)	0.086% (0.06-0.12%)	2.1 (1.4-2.9)	13 (9.3-17)
Iran	1995	Male	95 plus	12% (4.1-25%)	88% (75-96%)	0.0023 (0.00034-0.0071)	0.094% (0.071-0.12%)	0.076% (0.052-0.11%)	0.46 (0.32-0.65)	2.3 (1.7-3.1)
Iran	2000	Female	All Ages	10% (6.3-16%)	90% (84-94%)	0.0026 (0.0013-0.0048)	0.17% (0.13-0.22%)	0.11% (0.079-0.13%)	130 (94-160)	14000 (10000-19000)
Iran	2000	Female	20 to 24	11% (3.2-25%)	89% (75-97%)	0.0029 (0.00078-0.0077)	0.42% (0.24-0.65%)	0.079% (0.055-0.11%)	1.9 (1.3-3)	2200 (1200-3700)
Iran	2000	Female	25 to 29	12% (3.5-26%)	88% (74-97%)	0.0036 (0.00095-0.0087)	0.45% (0.28-0.65%)	0.093% (0.066-0.13%)	2.1 (1.4-2.9)	2000 (1200-3100)
Iran	2000	Female	30 to 34	14% (3.7-29%)	86% (71-96%)	0.0043 (0.0011-0.01)	0.42% (0.26-0.62%)	0.14% (0.096-0.2%)	3.3 (2.2-4.8)	1800 (1100-2800)
Iran	2000	Female	35 to 39	15% (4.2-31%)	85% (69-96%)	0.005 (0.0014-0.012)	0.33% (0.23-0.47%)	0.17% (0.12-0.23%)	4.7 (3-7.1)	1400 (890-2100)
Iran	2000	Female	40 to 44	16% (5.4-31%)	84% (69-95%)	0.0055 (0.0017-0.013)	0.28% (0.19-0.39%)	0.22% (0.14-0.3%)	7.3 (4.4-11)	1100 (720-1700)
Iran	2000	Female	45 to 49	16% (5.9-31%)	84% (69-94%)	0.006 (0.0016-0.014)	0.24% (0.16-0.32%)	0.22% (0.13-0.31%)	9.1 (5.2-14)	960 (630-1400)
Iran	2000	Female	50 to 54	17% (7-33%)	83% (67-93%)	0.0062 (0.0016-0.016)	0.2% (0.14-0.27%)	0.2% (0.13-0.27%)	8.7 (5.2-13)	650 (430-930)
Iran	2000	Female	55 to 59	18% (7.2-37%)	82% (63-93%)	0.0063 (0.0016-0.016)	0.16% (0.11-0.22%)	0.17% (0.11-0.25%)	8.6 (5.3-13)	470 (320-660)
Iran	2000	Female	60 to 64	19% (7.3-39%)	81% (61-93%)	0.006 (0.0013-0.017)	0.14% (0.1-0.2%)	0.17% (0.11-0.24%)	13 (7.9-19)	480 (330-680)
Iran	2000	Female	65 to 69	20% (8.3-37%)	80% (63-92%)	0.0053 (0.0011-0.013)	0.14% (0.097-0.18%)	0.17% (0.11-0.24%)	17 (11-25)	480 (330-670)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	2000	Female	70 to 74	19% (7.7-34%)	81% (66-92%)	0.0042 (0.00093-0.011)	0.12% (0.085-0.17%)	0.15% (0.099-0.22%)	19 (12-28)	390 (270-570)
Iran	2000	Female	75 to 79	17% (6.4-34%)	83% (66-94%)	0.0034 (0.00058-0.011)	0.1% (0.074-0.14%)	0.12% (0.084-0.18%)	17 (11-24)	260 (180-360)
Iran	2000	Female	80 to 84	15% (5.3-35%)	85% (65-95%)	0.0031 (0.00047-0.0097)	0.085% (0.061-0.12%)	0.098% (0.07-0.14%)	9.2 (6.2-13)	100 (75-150)
Iran	2000	Female	85 to 89	16% (5.1-36%)	84% (64-95%)	0.0031 (0.00055-0.011)	0.072% (0.052-0.1%)	0.081% (0.055-0.12%)	5 (3.3-7.7)	40 (29-59)
Iran	2000	Female	90 to 94	18% (5.8-40%)	82% (60-94%)	0.0036 (0.00061-0.012)	0.06% (0.043-0.087%)	0.064% (0.043-0.097%)	2.1 (1.4-3.2)	12 (8.5-18)
Iran	2000	Female	95 plus	23% (7.1-48%)	77% (52-93%)	0.0043 (0.00063-0.014)	0.062% (0.043-0.089%)	0.056% (0.035-0.088%)	0.71 (0.44-1.1)	3.2 (2.2-4.6)
Iran	2000	Male	All Ages	7.9% (5.4-11%)	92% (89-95%)	0.0028 (0.0014-0.0051)	0.27% (0.21-0.34%)	0.17% (0.14-0.21%)	310 (250-400)	28000 (21000-37000)
Iran	2000	Male	20 to 24	9.2% (3.6-18%)	91% (82-96%)	0.0028 (0.00082-0.0064)	0.6% (0.38-0.9%)	0.059% (0.039-0.09%)	3.8 (2.3-5.8)	4500 (2700-7100)
Iran	2000	Male	25 to 29	11% (4.4-22%)	89% (78-96%)	0.0043 (0.0014-0.0099)	0.6% (0.41-0.84%)	0.098% (0.064-0.18%)	5.3 (3.2-9.7)	3700 (2300-5400)
Iran	2000	Male	30 to 34	13% (4.7-26%)	87% (74-95%)	0.0055 (0.0017-0.013)	0.55% (0.37-0.79%)	0.16% (0.11-0.26%)	8.1 (5.1-14)	3000 (1900-4500)
Iran	2000	Male	35 to 39	14% (5.6-27%)	86% (73-94%)	0.0063 (0.0018-0.015)	0.54% (0.39-0.73%)	0.27% (0.19-0.42%)	15 (9.8-24)	2800 (1900-4000)
Iran	2000	Male	40 to 44	15% (6.5-26%)	85% (74-94%)	0.0067 (0.0021-0.016)	0.49% (0.36-0.65%)	0.31% (0.22-0.44%)	20 (14-30)	2600 (1800-3600)
Iran	2000	Male	45 to 49	14% (5.4-27%)	86% (73-95%)	0.007 (0.0019-0.016)	0.46% (0.35-0.58%)	0.39% (0.28-0.53%)	31 (21-45)	2400 (1700-3200)
Iran	2000	Male	50 to 54	14% (5.9-28%)	86% (72-94%)	0.007 (0.0019-0.018)	0.38% (0.29-0.48%)	0.35% (0.26-0.48%)	27 (19-38)	1600 (1200-2100)
Iran	2000	Male	55 to 59	13% (5.5-26%)	87% (74-94%)	0.0067 (0.0018-0.017)	0.31% (0.24-0.4%)	0.31% (0.23-0.42%)	26 (18-36)	1200 (860-1500)
Iran	2000	Male	60 to 64	13% (5.3-25%)	87% (75-95%)	0.006 (0.0015-0.016)	0.27% (0.2-0.35%)	0.29% (0.21-0.41%)	35 (24-49)	1200 (830-1600)
Iran	2000	Male	65 to 69	12% (5.3-23%)	88% (77-95%)	0.0049 (0.0012-0.012)	0.23% (0.18-0.3%)	0.26% (0.19-0.35%)	41 (29-58)	1100 (810-1500)
Iran	2000	Male	70 to 74	11% (4.6-20%)	89% (80-95%)	0.0036 (0.00086-0.0096)	0.19% (0.14-0.25%)	0.21% (0.15-0.29%)	38 (27-54)	780 (570-1100)
Iran	2000	Male	75 to 79	9.4% (3.8-18%)	91% (82-96%)	0.0027 (0.00053-0.0083)	0.15% (0.12-0.2%)	0.17% (0.12-0.23%)	31 (22-43)	470 (350-640)
Iran	2000	Male	80 to 84	8.4% (3.1-17%)	92% (83-97%)	0.0023 (0.00045-0.0066)	0.12% (0.091-0.16%)	0.13% (0.095-0.18%)	17 (12-24)	190 (140-270)
Iran	2000	Male	85 to 89	8.2% (3.3-16%)	92% (84-97%)	0.0022 (0.00047-0.0057)	0.1% (0.079-0.13%)	0.11% (0.079-0.15%)	7.7 (5.6-11)	65 (49-84)
Iran	2000	Male	90 to 94	9.2% (3.8-19%)	91% (81-96%)	0.0023 (0.00045-0.0064)	0.09% (0.068-0.12%)	0.088% (0.063-0.12%)	2.6 (1.9-3.7)	16 (12-21)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	2000	Male	95 plus	12% (4.1-24%)	88% (76-96%)	0.0026 (0.00038-0.008)	0.098% (0.074-0.13%)	0.079% (0.055-0.11%)	0.61 (0.43-0.83)	3 (2.3-3.9)
Iran	2005	Female	All Ages	11% (6.3-17%)	89% (83-94%)	0.0031 (0.0015-0.0053)	0.18% (0.14-0.23%)	0.11% (0.086-0.13%)	160 (110-210)	16000 (12000-21000)
Iran	2005	Female	20 to 24	11% (3.1-25%)	89% (75-97%)	0.0031 (0.00084-0.0077)	0.4% (0.24-0.63%)	0.085% (0.061-0.12%)	2.7 (1.7-3.9)	2800 (1500-4600)
Iran	2005	Female	25 to 29	12% (3.3-27%)	88% (73-97%)	0.0038 (0.00096-0.0089)	0.43% (0.28-0.63%)	0.1% (0.074-0.14%)	2.7 (1.8-3.8)	2500 (1400-3900)
Iran	2005	Female	30 to 34	14% (3.7-29%)	86% (71-96%)	0.0044 (0.0012-0.01)	0.4% (0.25-0.59%)	0.14% (0.1-0.19%)	3.3 (2.2-5)	1900 (1100-2900)
Iran	2005	Female	35 to 39	15% (4.2-30%)	85% (70-96%)	0.0052 (0.0015-0.013)	0.33% (0.22-0.46%)	0.17% (0.13-0.23%)	4.6 (3-6.8)	1500 (940-2200)
Iran	2005	Female	40 to 44	16% (5.7-31%)	84% (69-94%)	0.0058 (0.0018-0.013)	0.27% (0.18-0.38%)	0.22% (0.15-0.3%)	7.1 (4.4-11)	1200 (720-1900)
Iran	2005	Female	45 to 49	17% (6-31%)	83% (69-94%)	0.0064 (0.0017-0.015)	0.23% (0.16-0.32%)	0.22% (0.14-0.32%)	9.4 (5.7-15)	1000 (670-1500)
Iran	2005	Female	50 to 54	17% (6.8-34%)	83% (66-93%)	0.0067 (0.0017-0.016)	0.19% (0.14-0.26%)	0.2% (0.13-0.27%)	11 (7-17)	850 (570-1200)
Iran	2005	Female	55 to 59	18% (7.5-35%)	82% (65-92%)	0.0066 (0.0017-0.017)	0.16% (0.11-0.21%)	0.17% (0.11-0.24%)	10 (6.2-15)	560 (380-770)
Iran	2005	Female	60 to 64	19% (7.7-36%)	81% (64-92%)	0.0063 (0.0014-0.016)	0.14% (0.1-0.19%)	0.16% (0.11-0.23%)	13 (8-18)	480 (320-670)
Iran	2005	Female	65 to 69	20% (8.2-37%)	80% (63-92%)	0.0056 (0.0012-0.014)	0.13% (0.098-0.18%)	0.16% (0.11-0.23%)	18 (11-26)	510 (350-720)
Iran	2005	Female	70 to 74	19% (7.8-36%)	81% (64-92%)	0.0045 (0.00098-0.013)	0.14% (0.086-0.16%)	0.14% (0.098-0.2%)	24 (15-35)	500 (330-710)
Iran	2005	Female	75 to 79	17% (6.3-34%)	83% (66-94%)	0.0037 (0.00066-0.012)	0.1% (0.074-0.14%)	0.12% (0.083-0.16%)	22 (15-33)	340 (240-490)
Iran	2005	Female	80 to 84	15% (5.2-34%)	85% (66-95%)	0.0033 (0.00055-0.011)	0.083% (0.063-0.11%)	0.095% (0.069-0.13%)	15 (10-21)	170 (120-230)
Iran	2005	Female	85 to 89	16% (5.1-36%)	84% (64-95%)	0.0033 (0.00061-0.011)	0.069% (0.051-0.096%)	0.076% (0.054-0.11%)	8.3 (5.8-12)	67 (49-96)
Iran	2005	Female	90 to 94	18% (5.8-41%)	82% (59-94%)	0.0037 (0.00067-0.012)	0.057% (0.043-0.078%)	0.059% (0.042-0.086%)	2.7 (1.8-3.9)	16 (11-22)
Iran	2005	Female	95 plus	23% (6.4-47%)	77% (53-94%)	0.0044 (0.00064-0.013)	0.058% (0.043-0.081%)	0.051% (0.034-0.08%)	0.78 (0.51-1.2)	3.6 (2.5-5)
Iran	2005	Male	All Ages	8.7% (6-13%)	91% (87-94%)	0.0034 (0.0017-0.006)	0.3% (0.23-0.37%)	0.18% (0.15-0.22%)	390 (300-510)	34000 (26000-44000)
Iran	2005	Male	20 to 24	9.2% (3.6-18%)	91% (82-96%)	0.0029 (0.0009-0.0067)	0.56% (0.35-0.81%)	0.061% (0.041-0.092%)	5.9 (3.6-9.3)	5900 (3500-8900)
Iran	2005	Male	25 to 29	12% (4.4-23%)	88% (77-96%)	0.0046 (0.0015-0.01)	0.55% (0.38-0.77%)	0.1% (0.067-0.18%)	7.9 (4.9-14)	4500 (2900-6700)
Iran	2005	Male	30 to 34	13% (4.7-27%)	87% (73-95%)	0.0059 (0.002-0.013)	0.51% (0.34-0.72%)	0.16% (0.12-0.26%)	11 (7-17)	3400 (2200-4900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	2005	Male	35 to 39	15% (5.4-27%)	85% (73-95%)	0.0068 (0.0021-0.015)	0.54% (0.39-0.71%)	0.29% (0.2-0.43%)	19 (12-30)	3300 (2300-4700)
Iran	2005	Male	40 to 44	15% (6.5-26%)	85% (74-93%)	0.0072 (0.0023-0.016)	0.49% (0.36-0.64%)	0.33% (0.25-0.44%)	24 (16-36)	3000 (2100-4100)
Iran	2005	Male	45 to 49	14% (5.6-27%)	86% (73-94%)	0.0074 (0.002-0.017)	0.46% (0.37-0.58%)	0.41% (0.3-0.54%)	36 (25-52)	2800 (2100-3700)
Iran	2005	Male	50 to 54	14% (5.8-27%)	86% (73-94%)	0.0073 (0.0021-0.019)	0.39% (0.3-0.48%)	0.37% (0.28-0.48%)	37 (26-53)	2200 (1600-3000)
Iran	2005	Male	55 to 59	13% (5.6-25%)	87% (75-94%)	0.007 (0.0019-0.017)	0.32% (0.24-0.4%)	0.32% (0.24-0.44%)	30 (20-45)	1400 (980-1900)
Iran	2005	Male	60 to 64	13% (5.3-25%)	87% (75-95%)	0.0063 (0.0016-0.016)	0.28% (0.21-0.36%)	0.31% (0.22-0.41%)	34 (23-49)	1200 (830-1600)
Iran	2005	Male	65 to 69	12% (5.4-22%)	88% (78-95%)	0.0052 (0.0013-0.012)	0.25% (0.19-0.32%)	0.28% (0.21-0.38%)	45 (31-65)	1200 (870-1600)
Iran	2005	Male	70 to 74	11% (4.9-20%)	89% (80-95%)	0.0039 (0.00093-0.01)	0.19% (0.15-0.25%)	0.22% (0.16-0.3%)	51 (36-73)	1000 (750-1400)
Iran	2005	Male	75 to 79	9.6% (4-19%)	90% (81-96%)	0.0029 (0.00063-0.0084)	0.16% (0.12-0.2%)	0.18% (0.13-0.23%)	43 (31-60)	660 (490-880)
Iran	2005	Male	80 to 84	8.4% (3.4-17%)	92% (83-97%)	0.0024 (0.00051-0.0068)	0.12% (0.094-0.16%)	0.13% (0.097-0.18%)	26 (18-37)	300 (220-400)
Iran	2005	Male	85 to 89	8.2% (3.3-16%)	92% (84-97%)	0.0022 (0.0005-0.0061)	0.1% (0.08-0.14%)	0.11% (0.081-0.15%)	13 (9.1-18)	110 (78-140)
Iran	2005	Male	90 to 94	9.3% (3.6-19%)	91% (81-96%)	0.0024 (0.00049-0.0066)	0.091% (0.071-0.12%)	0.089% (0.067-0.12%)	3.7 (2.7-5.2)	23 (17-30)
Iran	2005	Male	95 plus	12% (4.2-24%)	88% (76-96%)	0.0028 (0.00043-0.0082)	0.1% (0.077-0.13%)	0.081% (0.059-0.11%)	0.85 (0.6-1.2)	4.2 (3.2-5.5)
Iran	2010	Female	All Ages	11% (6.7-17%)	89% (83-93%)	0.0034 (0.0017-0.0059)	0.21% (0.16-0.26%)	0.13% (0.1-0.15%)	180 (130-250)	18000 (13000-24000)
Iran	2010	Female	20 to 24	11% (3-24%)	89% (76-97%)	0.0032 (0.00074-0.0083)	0.43% (0.25-0.65%)	0.11% (0.081-0.16%)	3.1 (2-4.6)	2800 (1600-4600)
Iran	2010	Female	25 to 29	12% (3.4-27%)	88% (73-97%)	0.0039 (0.001-0.0092)	0.46% (0.3-0.67%)	0.13% (0.099-0.17%)	3.8 (2.4-5.3)	3300 (2000-5200)
Iran	2010	Female	30 to 34	14% (3.9-29%)	86% (71-96%)	0.0046 (0.0014-0.01)	0.42% (0.27-0.63%)	0.16% (0.13-0.21%)	4.4 (2.9-6.1)	2600 (1500-4100)
Iran	2010	Female	35 to 39	15% (4.3-31%)	85% (69-96%)	0.0054 (0.0015-0.013)	0.33% (0.22-0.47%)	0.19% (0.14-0.26%)	5.1 (3.3-7.7)	1700 (1000-2500)
Iran	2010	Female	40 to 44	16% (5.7-31%)	84% (69-94%)	0.006 (0.0018-0.013)	0.27% (0.17-0.39%)	0.24% (0.17-0.32%)	7.5 (4.7-11)	1300 (780-2000)
Iran	2010	Female	45 to 49	17% (6-32%)	83% (68-94%)	0.0065 (0.0017-0.015)	0.24% (0.17-0.33%)	0.24% (0.17-0.33%)	10 (6.2-15)	1200 (760-1700)
Iran	2010	Female	50 to 54	17% (6.6-34%)	83% (66-93%)	0.0068 (0.0017-0.015)	0.2% (0.14-0.27%)	0.23% (0.16-0.31%)	13 (8.2-19)	1000 (680-1500)
Iran	2010	Female	55 to 59	18% (6.6-36%)	82% (64-93%)	0.0069 (0.0017-0.017)	0.17% (0.12-0.23%)	0.21% (0.14-0.29%)	15 (9.2-23)	850 (570-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	2010	Female	60 to 64	19% (7.1-39%)	81% (61-93%)	0.0066 (0.0015-0.017)	0.16% (0.11-0.21%)	0.2% (0.13-0.29%)	17 (11-26)	640 (440-900)
Iran	2010	Female	65 to 69	20% (7.7-36%)	80% (64-92%)	0.0057 (0.0013-0.014)	0.15% (0.11-0.2%)	0.19% (0.13-0.27%)	19 (12-29)	530 (360-750)
Iran	2010	Female	70 to 74	18% (7.4-34%)	82% (66-93%)	0.0046 (0.00096-0.013)	0.13% (0.095-0.18%)	0.16% (0.11-0.24%)	24 (15-37)	500 (340-720)
Iran	2010	Female	75 to 79	16% (6.1-33%)	84% (67-94%)	0.0037 (0.00065-0.011)	0.11% (0.079-0.14%)	0.13% (0.091-0.17%)	25 (16-35)	390 (270-530)
Iran	2010	Female	80 to 84	15% (4.9-33%)	85% (67-95%)	0.0033 (0.00057-0.011)	0.089% (0.067-0.12%)	0.1% (0.075-0.14%)	19 (13-27)	220 (150-300)
Iran	2010	Female	85 to 89	15% (4.9-35%)	85% (65-95%)	0.0034 (0.0006-0.011)	0.073% (0.056-0.095%)	0.08% (0.061-0.11%)	11 (8-15)	91 (67-120)
Iran	2010	Female	90 to 94	18% (5.9-41%)	82% (59-94%)	0.0038 (0.00065-0.013)	0.058% (0.045-0.076%)	0.06% (0.045-0.081%)	3.8 (2.8-5.3)	23 (17-31)
Iran	2010	Female	95 plus	22% (6.8-48%)	78% (52-93%)	0.0046 (0.00067-0.015)	0.057% (0.043-0.077%)	0.051% (0.037-0.071%)	0.88 (0.62-1.2)	4 (2.9-5.4)
Iran	2010	Male	All Ages	9.2% (6.2-13%)	91% (87-94%)	0.0038 (0.0019-0.0064)	0.34% (0.27-0.42%)	0.22% (0.18-0.25%)	460 (350-610)	38000 (29000-49000)
Iran	2010	Male	20 to 24	9.3% (3.7-18%)	91% (82-96%)	0.0031 (0.00092-0.0072)	0.63% (0.41-0.92%)	0.084% (0.052-0.12%)	6.8 (3.9-10)	600 (3600-9000)
Iran	2010	Male	25 to 29	11% (4.7-22%)	89% (78-95%)	0.0047 (0.0016-0.0099)	0.63% (0.45-0.88%)	0.14% (0.092-0.2%)	11 (6.9-16)	5900 (3800-8600)
Iran	2010	Male	30 to 34	13% (4.9-25%)	87% (75-95%)	0.006 (0.0021-0.013)	0.58% (0.4-0.79%)	0.21% (0.15-0.3%)	14 (8.7-21)	4300 (2800-6000)
Iran	2010	Male	35 to 39	15% (5.6-26%)	85% (74-94%)	0.0069 (0.0022-0.015)	0.58% (0.43-0.75%)	0.34% (0.25-0.47%)	21 (14-32)	3700 (2500-5000)
Iran	2010	Male	40 to 44	15% (6.7-27%)	85% (73-93%)	0.0074 (0.0024-0.016)	0.53% (0.39-0.7%)	0.38% (0.29-0.5%)	27 (18-40)	3400 (2300-4800)
Iran	2010	Male	45 to 49	14% (5.7-27%)	86% (73-94%)	0.0076 (0.0022-0.017)	0.5% (0.39-0.62%)	0.46% (0.35-0.59%)	39 (26-56)	3100 (2300-4100)
Iran	2010	Male	50 to 54	14% (5.8-28%)	86% (72-94%)	0.0075 (0.0021-0.017)	0.41% (0.32-0.51%)	0.41% (0.31-0.52%)	43 (29-63)	2600 (1900-3500)
Iran	2010	Male	55 to 59	13% (5.5-26%)	87% (74-95%)	0.0072 (0.0019-0.017)	0.34% (0.26-0.42%)	0.36% (0.26-0.47%)	43 (29-64)	2000 (1400-2700)
Iran	2010	Male	60 to 64	13% (5.2-25%)	87% (75-95%)	0.0064 (0.0016-0.016)	0.3% (0.22-0.39%)	0.34% (0.25-0.45%)	40 (27-57)	1400 (960-1800)
Iran	2010	Male	65 to 69	12% (5.4-23%)	88% (77-95%)	0.0053 (0.0014-0.013)	0.27% (0.2-0.35%)	0.32% (0.24-0.43%)	43 (29-62)	1100 (800-1600)
Iran	2010	Male	70 to 74	11% (5-20%)	89% (80-95%)	0.004 (0.00096-0.011)	0.22% (0.17-0.29%)	0.26% (0.19-0.36%)	53 (36-76)	1100 (770-1500)
Iran	2010	Male	75 to 79	9.6% (4.1-18%)	90% (82-96%)	0.003 (0.00068-0.0085)	0.18% (0.14-0.23%)	0.21% (0.16-0.27%)	53 (37-74)	820 (600-1100)
Iran	2010	Male	80 to 84	8.6% (3.4-17%)	91% (83-97%)	0.0025 (0.00059-0.007)	0.14% (0.11-0.18%)	0.15% (0.11-0.2%)	34 (24-48)	400 (290-540)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	2010	Male	85 to 89	8.4% (3.3-17%)	92% (83-97%)	0.0024 (0.00053-0.0065)	0.11% (0.09-0.14%)	0.12% (0.092-0.16%)	17 (13-23)	140 (110-190)
Iran	2010	Male	90 to 94	9.4% (3.8-19%)	91% (81-96%)	0.0025 (0.00055-0.0071)	0.097% (0.077-0.12%)	0.095% (0.073-0.13%)	5.4 (4-7.2)	33 (26-43)
Iran	2010	Male	95 plus	12% (4.2-25%)	88% (75-96%)	0.0029 (0.0005-0.0091)	0.11% (0.083-0.14%)	0.086% (0.064-0.11%)	1 (0.75-1.4)	5.1 (3.9-6.7)
Iran	2016	Female	All Ages	12% (7.1-19%)	88% (81-93%)	0.0034 (0.0014-0.0064)	0.23% (0.18-0.29%)	0.14% (0.11-0.17%)	200 (140-270)	20000 (15000-27000)
Iran	2016	Female	20 to 24	11% (3.2-25%)	89% (75-97%)	0.003 (0.00063-0.0081)	0.47% (0.28-0.69%)	0.16% (0.11-0.24%)	2.4 (1.5-3.6)	2000 (1200-3200)
Iran	2016	Female	25 to 29	12% (3.3-27%)	88% (73-97%)	0.0036 (0.00074-0.0088)	0.5% (0.32-0.74%)	0.17% (0.12-0.23%)	3.9 (2.6-5.5)	3400 (1900-5200)
Iran	2016	Female	30 to 34	14% (3.7-29%)	86% (71-96%)	0.0041 (0.0011-0.0099)	0.46% (0.28-0.7%)	0.19% (0.15-0.25%)	5.9 (4.1-8.3)	3700 (2100-6000)
Iran	2016	Female	35 to 39	15% (4.5-30%)	85% (70-96%)	0.0049 (0.0014-0.012)	0.36% (0.23-0.51%)	0.21% (0.16-0.28%)	6.9 (4.5-10)	2500 (1500-3900)
Iran	2016	Female	40 to 44	16% (5.8-31%)	84% (69-94%)	0.0056 (0.0016-0.013)	0.28% (0.19-0.4%)	0.24% (0.17-0.33%)	8.1 (5.2-12)	1600 (960-2400)
Iran	2016	Female	45 to 49	17% (5.7-32%)	83% (68-94%)	0.0061 (0.0016-0.015)	0.24% (0.17-0.33%)	0.24% (0.17-0.34%)	10 (6.3-15)	1300 (840-1900)
Iran	2016	Female	50 to 54	18% (6.9-35%)	82% (65-93%)	0.0063 (0.0017-0.015)	0.21% (0.15-0.29%)	0.24% (0.16-0.35%)	15 (8.9-23)	1200 (770-1700)
Iran	2016	Female	55 to 59	19% (7.5-37%)	81% (63-93%)	0.0064 (0.0015-0.016)	0.18% (0.12-0.25%)	0.22% (0.14-0.32%)	18 (10-29)	1000 (620-1500)
Iran	2016	Female	60 to 64	20% (7.3-39%)	80% (61-93%)	0.0061 (0.0013-0.018)	0.17% (0.12-0.23%)	0.22% (0.14-0.31%)	25 (15-40)	970 (650-1400)
Iran	2016	Female	65 to 69	20% (8.4-37%)	80% (63-92%)	0.0054 (0.0012-0.014)	0.16% (0.11-0.22%)	0.21% (0.14-0.31%)	25 (15-40)	720 (490-1100)
Iran	2016	Female	70 to 74	19% (8.1-35%)	81% (65-92%)	0.0044 (0.00085-0.012)	0.14% (0.098-0.19%)	0.18% (0.12-0.26%)	22 (13-35)	470 (310-700)
Iran	2016	Female	75 to 79	17% (6.5-35%)	83% (65-94%)	0.0037 (0.00059-0.012)	0.11% (0.079-0.14%)	0.13% (0.092-0.18%)	21 (13-30)	330 (230-460)
Iran	2016	Female	80 to 84	16% (5.4-35%)	84% (65-95%)	0.0034 (0.00056-0.011)	0.09% (0.068-0.12%)	0.11% (0.076-0.14%)	18 (12-25)	210 (150-280)
Iran	2016	Female	85 to 89	17% (5.6-38%)	83% (62-94%)	0.0034 (0.00058-0.012)	0.072% (0.056-0.091%)	0.079% (0.06-0.1%)	13 (9.3-18)	110 (81-140)
Iran	2016	Female	90 to 94	20% (6.7-42%)	80% (58-93%)	0.0039 (0.00066-0.014)	0.058% (0.046-0.073%)	0.058% (0.045-0.077%)	4.6 (3.5-6.3)	29 (22-38)
Iran	2016	Female	95 plus	24% (7.7-48%)	76% (52-92%)	0.0046 (0.00061-0.016)	0.057% (0.043-0.076%)	0.048% (0.035-0.066%)	1.2 (0.88-1.7)	6 (4.4-8.2)
Iran	2016	Male	All Ages	9.6% (6.4-13%)	90% (87-94%)	0.0037 (0.0016-0.0071)	0.38% (0.31-0.46%)	0.25% (0.21-0.29%)	540 (410-710)	43000 (32000-54000)
Iran	2016	Male	20 to 24	9.3% (3.7-18%)	91% (82-96%)	0.0028 (0.00069-0.0071)	0.73% (0.47-1.1%)	0.13% (0.059-0.19%)	5.8 (2.7-9.4)	4500 (2800-6800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iran	2016	Male	25 to 29	12% (4.4-23%)	88% (77-96%)	0.0043 (0.0012-0.01)	0.73% (0.51-1%)	0.19% (0.11-0.29%)	12 (6.4-19)	6000 (3900-8800)
Iran	2016	Male	30 to 34	13% (4.6-25%)	87% (75-95%)	0.0055 (0.0016-0.012)	0.66% (0.46-0.91%)	0.27% (0.17-0.38%)	19 (11-30)	5900 (3800-8700)
Iran	2016	Male	35 to 39	15% (5.8-26%)	85% (74-94%)	0.0063 (0.0018-0.014)	0.63% (0.47-0.82%)	0.4% (0.28-0.55%)	28 (17-41)	4900 (3400-6700)
Iran	2016	Male	40 to 44	15% (6.8-27%)	85% (73-93%)	0.0068 (0.0018-0.016)	0.57% (0.42-0.73%)	0.42% (0.31-0.56%)	31 (21-43)	3900 (2700-5300)
Iran	2016	Male	45 to 49	15% (5.4-27%)	85% (73-95%)	0.007 (0.0017-0.017)	0.52% (0.41-0.66%)	0.49% (0.37-0.65%)	45 (30-66)	3700 (2700-5000)
Iran	2016	Male	50 to 54	14% (5.9-27%)	86% (73-94%)	0.0068 (0.0018-0.017)	0.42% (0.33-0.53%)	0.43% (0.32-0.55%)	51 (35-74)	3100 (2300-4200)
Iran	2016	Male	55 to 59	13% (5.5-26%)	87% (74-94%)	0.0065 (0.0017-0.017)	0.34% (0.27-0.44%)	0.37% (0.27-0.49%)	52 (35-75)	2400 (1800-3200)
Iran	2016	Male	60 to 64	13% (5.4-25%)	87% (75-95%)	0.0058 (0.0014-0.016)	0.31% (0.23-0.41%)	0.36% (0.25-0.49%)	64 (42-91)	2200 (1600-3000)
Iran	2016	Male	65 to 69	12% (5.4-23%)	88% (77-95%)	0.0048 (0.0012-0.012)	0.27% (0.2-0.35%)	0.32% (0.23-0.44%)	55 (36-80)	1500 (1000-2000)
Iran	2016	Male	70 to 74	11% (5-20%)	89% (80-95%)	0.0035 (0.0008-0.0095)	0.23% (0.17-0.3%)	0.28% (0.2-0.38%)	49 (33-70)	1000 (690-1400)
Iran	2016	Male	75 to 79	9.5% (4-19%)	90% (81-96%)	0.0027 (0.00054-0.0083)	0.18% (0.14-0.24%)	0.22% (0.16-0.29%)	50 (35-71)	780 (560-1100)
Iran	2016	Male	80 to 84	8.6% (3.3-17%)	91% (83-97%)	0.0023 (0.00047-0.0067)	0.14% (0.11-0.19%)	0.16% (0.12-0.22%)	38 (27-55)	440 (320-610)
Iran	2016	Male	85 to 89	8.4% (3.4-17%)	92% (83-97%)	0.0021 (0.00047-0.0062)	0.12% (0.091-0.15%)	0.12% (0.093-0.16%)	25 (18-34)	210 (160-280)
Iran	2016	Male	90 to 94	9.5% (3.8-19%)	91% (81-96%)	0.0023 (0.00047-0.0071)	0.098% (0.077-0.13%)	0.096% (0.073-0.13%)	7.8 (5.7-11)	49 (37-63)
Iran	2016	Male	95 plus	12% (4.3-25%)	88% (75-96%)	0.0027 (0.00038-0.0092)	0.11% (0.085-0.14%)	0.087% (0.064-0.12%)	1.7 (1.2-2.3)	8.5 (6.4-11)
Iraq	1990	Female	All Ages	11% (7-17%)	89% (83-93%)	0.45 (0.33-0.6)	1.1% (0.45-2%)	1.7% (0.22-4%)	990 (130-2300)	38000 (16000-65000)
Iraq	1990	Female	15 to 19	10% (3.3-23%)	90% (77-97%)	0.38 (0.11-0.88)	0.95% (0.47-1.8%)	1.6% (0.61-3.5%)	16 (5.9-35)	1600 (770-3100)
Iraq	1990	Female	20 to 24	13% (4.3-29%)	87% (71-96%)	0.62 (0.2-1.3)	1.7% (0.89-3.1%)	2.7% (0.96-5.3%)	30 (10-61)	2900 (1400-5000)
Iraq	1990	Female	25 to 29	15% (4.2-31%)	85% (69-96%)	0.75 (0.22-1.6)	2% (1-3.6%)	3.3% (1.3-6.4%)	32 (11-62)	2700 (1300-4800)
Iraq	1990	Female	30 to 34	17% (4.7-35%)	83% (65-95%)	0.9 (0.28-1.8)	2.3% (1.1-3.8%)	3.8% (1.5-6.7%)	36 (14-68)	2600 (1200-4400)
Iraq	1990	Female	35 to 39	19% (5.4-36%)	81% (64-95%)	1.1 (0.33-2.1)	2.4% (0.99-4.6%)	4% (1.5-7.9%)	54 (19-110)	3300 (1300-6300)
Iraq	1990	Female	40 to 44	20% (7.7-37%)	80% (63-92%)	1.2 (0.43-2.3)	2.9% (1.1-5.6%)	4.8% (1.8-9.3%)	55 (19-110)	2900 (1100-5600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	1990	Female	45 to 49	21% (7.7-38%)	79% (62-92%)	1.3 (0.44-2.7)	3.3% (1-6.7%)	5.1% (1.6-11%)	82 (25-180)	3700 (1200-7900)
Iraq	1990	Female	50 to 54	22% (8.6-40%)	78% (60-91%)	1.3 (0.42-2.9)	3.5% (0.83-7.7%)	5% (1.3-11%)	110 (27-250)	4200 (970-9800)
Iraq	1990	Female	55 to 59	23% (9-43%)	77% (57-91%)	1.3 (0.4-2.9)	3.5% (0.44-8%)	4.8% (0.71-11%)	130 (18-300)	4300 (520-10000)
Iraq	1990	Female	60 to 64	24% (9.5-45%)	76% (55-90%)	1.3 (0.33-2.9)	2.8% (-0.32-7.3%)	3.7% (-0.34-9.5%)	120 (-11-320)	3400 (-360-9200)
Iraq	1990	Female	65 to 69	24% (10-43%)	76% (57-90%)	1.1 (0.29-2.4)	2.3% (-0.56-6.5%)	2.9% (-0.65-8.2%)	120 (-27-350)	2900 (-710-8100)
Iraq	1990	Female	70 to 74	23% (9.2-41%)	77% (59-91%)	0.88 (0.21-2.1)	1.3% (-1.1-5.6%)	1.7% (-1.3-6.7%)	89 (-66-380)	1700 (-1300-7300)
Iraq	1990	Female	75 to 79	21% (7.5-41%)	79% (59-92%)	0.72 (0.15-2)	0.77% (-1.4-4.9%)	0.91% (-1.7-5.8%)	52 (-97-330)	760 (-1400-4900)
Iraq	1990	Female	80 to 84	19% (6.6-40%)	81% (60-93%)	0.65 (0.13-1.9)	0.72% (-1.1-4.2%)	0.83% (-1.4-4.8%)	37 (-61-220)	410 (-630-2400)
Iraq	1990	Female	85 to 89	19% (6.7-40%)	81% (60-93%)	0.64 (0.13-1.8)	0.59% (-1.1-3.9%)	0.66% (-1.3-4.6%)	20 (-40-140)	160 (-290-1000)
Iraq	1990	Female	90 to 94	22% (7.4-43%)	78% (57-93%)	0.72 (0.14-1.9)	0.55% (-1.3-4%)	0.6% (-1.6-4.6%)	7.7 (-21-61)	43 (-100-310)
Iraq	1990	Female	95 plus	27% (8.6-50%)	73% (50-91%)	0.85 (0.16-2.3)	0.63% (-1.8-4.9%)	0.66% (-2.3-5.7%)	2.4 (-8.3-21)	9.1 (-27-71)
Iraq	1990	Male	All Ages	8.7% (6.1-12%)	91% (88-94%)	0.5 (0.37-0.65)	1.4% (0.85-2.1%)	2% (1-3.2%)	1300 (620-2100)	55000 (34000-78000)
Iraq	1990	Male	15 to 19	7.7% (3.3-15%)	92% (85-97%)	0.27 (0.089-0.58)	1% (0.53-1.9%)	1.3% (0.47-2.7%)	20 (7.4-43)	2100 (1000-3900)
Iraq	1990	Male	20 to 24	12% (4.9-22%)	88% (78-95%)	0.62 (0.22-1.2)	2.4% (1.2-4.1%)	3% (1.2-5.8%)	60 (24-120)	5200 (2600-9500)
Iraq	1990	Male	25 to 29	15% (6.2-27%)	85% (73-94%)	0.97 (0.38-1.9)	3.3% (1.7-5.4%)	4.7% (2.1-8.2%)	72 (31-130)	5300 (2700-9400)
Iraq	1990	Male	30 to 34	17% (6.4-32%)	83% (68-94%)	1.2 (0.52-2.3)	3.7% (2-6%)	5.5% (2.6-9%)	75 (35-130)	5000 (2600-8200)
Iraq	1990	Male	35 to 39	18% (7.4-33%)	82% (67-93%)	1.4 (0.53-2.6)	4% (1.9-6.7%)	5.8% (2.5-9.8%)	110 (45-190)	6200 (2900-11000)
Iraq	1990	Male	40 to 44	19% (8.3-32%)	81% (68-92%)	1.5 (0.6-2.8)	4.1% (1.9-7%)	5.7% (2.5-9.9%)	92 (39-170)	4800 (2200-8400)
Iraq	1990	Male	45 to 49	18% (7.4-32%)	82% (68-93%)	1.6 (0.54-3.3)	4.2% (1.9-7.1%)	5.6% (2.3-9.5%)	120 (51-210)	5600 (2500-9700)
Iraq	1990	Male	50 to 54	18% (7.7-33%)	82% (67-92%)	1.6 (0.51-3.2)	3.9% (1.6-6.9%)	5% (1.9-8.9%)	130 (46-240)	5100 (2000-9300)
Iraq	1990	Male	55 to 59	17% (7.5-31%)	83% (69-92%)	1.5 (0.48-3.2)	3.7% (1.2-6.9%)	4.6% (1.4-8.8%)	140 (42-280)	4800 (1600-9400)
Iraq	1990	Male	60 to 64	16% (7.2-30%)	84% (70-93%)	1.3 (0.4-3)	3.1% (0.74-6.3%)	3.8% (0.82-7.7%)	130 (28-280)	3900 (940-7900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	1990	Male	65 to 69	15% (7.1-28%)	85% (72-93%)	1.1 (0.35-2.3)	2.6% (0.44-5.5%)	3.1% (0.47-6.7%)	130 (20-290)	3100 (520-6700)
Iraq	1990	Male	70 to 74	14% (6.4-25%)	86% (75-94%)	0.81 (0.24-1.9)	1.7% (0.057-4.4%)	2% (-0.0088-5.3%)	100 (-0.38-280)	2000 (66-5200)
Iraq	1990	Male	75 to 79	12% (5.5-24%)	88% (76-95%)	0.61 (0.16-1.6)	1.1% (-0.18-3.5%)	1.3% (-0.29-4.1%)	62 (-14-200)	930 (-150-3000)
Iraq	1990	Male	80 to 84	11% (4.4-21%)	89% (79-96%)	0.51 (0.13-1.2)	0.88% (-0.17-2.8%)	0.97% (-0.26-3.2%)	33 (-9.1-110)	380 (-73-1200)
Iraq	1990	Male	85 to 89	10% (4.4-20%)	90% (80-96%)	0.46 (0.13-1.1)	0.68% (-0.22-2.3%)	0.73% (-0.33-2.6%)	16 (-7.2-58)	130 (-44-440)
Iraq	1990	Male	90 to 94	12% (4.9-24%)	88% (76-95%)	0.49 (0.13-1.3)	0.6% (-0.33-2.3%)	0.63% (-0.48-2.6%)	5.5 (-4.2-23)	32 (-17-120)
Iraq	1990	Male	95 plus	15% (5.5-30%)	85% (70-95%)	0.56 (0.12-1.6)	0.54% (-0.56-2.8%)	0.51% (-0.77-3.1%)	1 (-1.5-6.2)	4.4 (-4.3-23)
Iraq	1995	Female	All Ages	12% (7.5-18%)	88% (82-92%)	0.35 (0.25-0.46)	0.77% (0.23-1.5%)	0.94% (-0.27-2.7%)	640 (-180-1900)	30000 (9000-58000)
Iraq	1995	Female	15 to 19	11% (3.5-23%)	89% (77-96%)	0.3 (0.091-0.65)	0.77% (0.4-1.4%)	1.2% (0.5-2.6%)	14 (5.7-31)	1500 (790-2800)
Iraq	1995	Female	20 to 24	14% (4.2-28%)	86% (72-96%)	0.47 (0.16-0.97)	1.4% (0.79-2.4%)	2% (0.77-3.9%)	28 (11-56)	2900 (1500-4800)
Iraq	1995	Female	25 to 29	15% (4.5-31%)	85% (69-95%)	0.57 (0.17-1.2)	1.6% (0.89-2.8%)	2.5% (1-4.7%)	36 (14-70)	3100 (1600-5300)
Iraq	1995	Female	30 to 34	17% (4.9-34%)	83% (66-95%)	0.67 (0.21-1.3)	1.8% (0.89-3%)	2.7% (1.1-4.9%)	37 (14-69)	2700 (1400-4800)
Iraq	1995	Female	35 to 39	19% (5.9-36%)	81% (64-94%)	0.79 (0.27-1.5)	1.9% (0.73-3.7%)	3% (0.99-5.9%)	43 (15-90)	2600 (990-5100)
Iraq	1995	Female	40 to 44	20% (7.9-37%)	80% (63-92%)	0.89 (0.34-1.7)	2.2% (0.74-4.5%)	3.6% (1.1-7.1%)	73 (22-150)	3700 (1200-7700)
Iraq	1995	Female	45 to 49	21% (7.6-39%)	79% (61-92%)	0.97 (0.31-2.1)	2.4% (0.56-5.6%)	3.6% (0.87-8.2%)	65 (15-150)	2900 (660-6700)
Iraq	1995	Female	50 to 54	22% (8.7-41%)	78% (59-91%)	1 (0.31-2.2)	2.5% (0.17-6.1%)	3.5% (0.37-8.6%)	92 (9.9-230)	3500 (250-9100)
Iraq	1995	Female	55 to 59	23% (10-43%)	77% (57-90%)	0.99 (0.31-2.2)	2.3% (-0.19-6%)	3.1% (-0.098-8.2%)	100 (-3-270)	3400 (-270-9000)
Iraq	1995	Female	60 to 64	24% (10-44%)	76% (56-90%)	0.94 (0.27-2.2)	1.6% (-0.94-5.5%)	2.1% (-1.1-7%)	85 (-45-290)	2400 (-1400-8200)
Iraq	1995	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.83 (0.23-1.9)	0.99% (-1.3-4.5%)	1.3% (-1.6-5.7%)	63 (-77-290)	1400 (-1900-6800)
Iraq	1995	Female	70 to 74	23% (10-40%)	77% (60-90%)	0.65 (0.17-1.6)	0.3% (-1.6-3.5%)	0.38% (-1.9-4.2%)	22 (-110-250)	400 (-2100-4600)
Iraq	1995	Female	75 to 79	21% (8.1-40%)	79% (60-92%)	0.52 (0.12-1.4)	-0.071% (-1.9-2.9%)	-0.091% (-2.2-3.3%)	-5.9 (-150-230)	-77 (-2100-3400)
Iraq	1995	Female	80 to 84	19% (6.5-40%)	81% (60-93%)	0.47 (0.1-1.4)	-0.059% (-1.6-2.3%)	-0.083% (-1.8-2.6%)	-4.6 (-100-150)	-39 (-1100-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	1995	Female	85 to 89	19% (6.6-41%)	81% (59-93%)	0.47 (0.094-1.3)	-0.15% (-1.6-2%)	-0.2% (-1.9-2.3%)	-7.2 (-69-83)	-46 (-510-630)
Iraq	1995	Female	90 to 94	22% (7.7-43%)	78% (57-92%)	0.53 (0.11-1.4)	-0.19% (-1.8-2.1%)	-0.27% (-2.1-2.5%)	-4.1 (-32-37)	-17 (-160-200)
Iraq	1995	Female	95 plus	27% (8.2-51%)	73% (49-92%)	0.63 (0.11-1.7)	-0.19% (-2.3-2.8%)	-0.31% (-2.9-3.3%)	-1.4 (-13-15)	-3.5 (-41-51)
Iraq	1995	Male	All Ages	9% (6.2-13%)	91% (87-94%)	0.37 (0.28-0.49)	1.1% (0.65-1.8%)	1.5% (0.62-2.6%)	1100 (470-2000)	53000 (31000-78000)
Iraq	1995	Male	15 to 19	7.9% (3.3-15%)	92% (85-97%)	0.2 (0.069-0.43)	0.85% (0.46-1.5%)	0.95% (0.37-2%)	20 (7.4-43)	2100 (1100-3800)
Iraq	1995	Male	20 to 24	12% (4.8-23%)	88% (77-95%)	0.45 (0.17-0.87)	1.9% (1-3.3%)	2.2% (0.93-4.3%)	61 (25-120)	5500 (2900-10000)
Iraq	1995	Male	25 to 29	15% (6.1-28%)	85% (72-94%)	0.71 (0.28-1.3)	2.6% (1.4-4.5%)	3.4% (1.5-6.2%)	88 (37-160)	6600 (3200-11000)
Iraq	1995	Male	30 to 34	17% (6.6-31%)	83% (69-93%)	0.91 (0.38-1.6)	3% (1.5-4.8%)	4.1% (1.8-7.1%)	81 (35-140)	5400 (2700-8900)
Iraq	1995	Male	35 to 39	19% (7.5-34%)	81% (66-92%)	1 (0.39-2)	3.2% (1.4-5.6%)	4.4% (1.6-8%)	86 (32-160)	5000 (2200-8800)
Iraq	1995	Male	40 to 44	19% (8.7-33%)	81% (67-91%)	1.1 (0.45-2.1)	3.2% (1.3-5.9%)	4.3% (1.6-8%)	110 (39-220)	5900 (2300-11000)
Iraq	1995	Male	45 to 49	19% (7.6-33%)	81% (67-92%)	1.2 (0.4-2.5)	3.3% (1.3-6.2%)	4.2% (1.5-8.1%)	100 (36-200)	4700 (1800-8900)
Iraq	1995	Male	50 to 54	18% (7.8-34%)	82% (66-92%)	1.1 (0.36-2.4)	3% (0.89-5.8%)	3.7% (0.99-7.4%)	120 (31-260)	4900 (1400-10000)
Iraq	1995	Male	55 to 59	17% (7.5-32%)	83% (68-93%)	1.1 (0.33-2.4)	2.8% (0.63-5.8%)	3.4% (0.66-7.2%)	130 (23-290)	4400 (960-9600)
Iraq	1995	Male	60 to 64	16% (6.9-31%)	84% (69-93%)	0.97 (0.27-2.2)	2.2% (0.25-5.3%)	2.7% (0.24-6.3%)	120 (11-290)	3500 (390-8200)
Iraq	1995	Male	65 to 69	16% (7.4-28%)	84% (72-93%)	0.79 (0.25-1.7)	1.7% (0.028-4.4%)	1.9% (-0.046-5.2%)	95 (-2.3-260)	2300 (38-6100)
Iraq	1995	Male	70 to 74	14% (7-25%)	86% (75-93%)	0.59 (0.18-1.4)	0.97% (-0.27-3.4%)	1.1% (-0.41-3.9%)	61 (-22-220)	1200 (-330-4300)
Iraq	1995	Male	75 to 79	12% (5.6-23%)	88% (77-94%)	0.44 (0.11-1.1)	0.56% (-0.4-2.5%)	0.6% (-0.53-2.8%)	35 (-30-170)	550 (-380-2500)
Iraq	1995	Male	80 to 84	11% (4.5-21%)	89% (79-96%)	0.37 (0.096-0.92)	0.38% (-0.42-1.8%)	0.39% (-0.54-2%)	17 (-24-88)	200 (-210-950)
Iraq	1995	Male	85 to 89	11% (4.2-20%)	89% (80-96%)	0.34 (0.09-0.87)	0.24% (-0.45-1.5%)	0.22% (-0.59-1.6%)	5.9 (-16-45)	56 (-110-350)
Iraq	1995	Male	90 to 94	12% (4.8-23%)	88% (77-95%)	0.36 (0.09-0.93)	0.2% (-0.49-1.4%)	0.16% (-0.68-1.6%)	1.7 (-7.7-18)	13 (-35-97)
Iraq	1995	Male	95 plus	15% (5.2-31%)	85% (69-95%)	0.41 (0.078-1.2)	0.13% (-0.77-1.6%)	0.041% (-1.1-1.7%)	0.11 (-3.1-4.9)	1.5 (-8.6-18)
Iraq	2000	Female	All Ages	12% (7.4-18%)	88% (82-93%)	0.23 (0.17-0.31)	0.34% (-0.062-0.85%)	0.053% (-0.87-1.2%)	39 (-650-860)	14000 (-2600-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	2000	Female	15 to 19	11% (3.6-24%)	89% (76-96%)	0.2 (0.058-0.44)	0.6% (0.35-1%)	0.88% (0.39-1.8%)	10 (4.4-22)	1200 (670-2200)
Iraq	2000	Female	20 to 24	14% (4.4-30%)	86% (70-96%)	0.31 (0.1-0.64)	1% (0.62-1.7%)	1.4% (0.54-2.7%)	20 (7.5-40)	2400 (1400-3900)
Iraq	2000	Female	25 to 29	16% (4.5-32%)	84% (68-95%)	0.37 (0.12-0.75)	1.2% (0.72-1.8%)	1.6% (0.73-3%)	26 (11-48)	2600 (1500-4200)
Iraq	2000	Female	30 to 34	17% (4.5-35%)	83% (65-95%)	0.43 (0.13-0.81)	1.1% (0.6-1.9%)	1.6% (0.62-3%)	30 (11-57)	2400 (1200-4100)
Iraq	2000	Female	35 to 39	19% (5.1-36%)	81% (64-95%)	0.5 (0.17-0.96)	1.1% (0.37-2.2%)	1.6% (0.41-3.5%)	30 (7.9-68)	1900 (700-3900)
Iraq	2000	Female	40 to 44	20% (7.4-36%)	80% (64-93%)	0.56 (0.2-1.1)	1.2% (0.23-2.6%)	1.9% (0.32-4.2%)	37 (6.2-83)	1900 (400-4100)
Iraq	2000	Female	45 to 49	21% (7.2-38%)	79% (62-93%)	0.61 (0.2-1.3)	1.2% (-0.13-3.2%)	1.8% (-0.036-4.8%)	51 (-0.94-140)	2200 (-250-6400)
Iraq	2000	Female	50 to 54	21% (8.6-40%)	79% (60-91%)	0.63 (0.19-1.4)	1% (-0.55-3.6%)	1.5% (-0.61-5%)	39 (-16-140)	1500 (-780-5300)
Iraq	2000	Female	55 to 59	22% (9.4-42%)	78% (58-91%)	0.62 (0.18-1.4)	0.71% (-1.3-1.1%)	1% (-1.2-4.2%)	38 (-43-150)	1200 (-1600-5200)
Iraq	2000	Female	60 to 64	24% (9.8-44%)	76% (56-90%)	0.59 (0.16-1.4)	0.12% (-1.9-2.8%)	0.21% (-2.2-3.6%)	9.6 (-100-160)	200 (-3000-4500)
Iraq	2000	Female	65 to 69	24% (11-42%)	76% (58-89%)	0.53 (0.14-1.2)	-0.34% (-2.1-1.8%)	-0.37% (-2.5-2.3%)	-21 (-140-130)	-550 (-3400-3000)
Iraq	2000	Female	70 to 74	23% (11-40%)	77% (60-89%)	0.42 (0.11-1)	-0.67% (-2.3-1.1%)	-0.78% (-2.7-1.4%)	-50 (-180-85)	-970 (-3400-1600)
Iraq	2000	Female	75 to 79	21% (7.8-39%)	79% (61-92%)	0.34 (0.074-0.93)	-0.79% (-2.4-0.56%)	-0.93% (-2.7-0.71%)	-64 (-190-45)	-930 (-2800-640)
Iraq	2000	Female	80 to 84	19% (6.7-40%)	81% (60-93%)	0.3 (0.063-0.86)	-0.71% (-2.3-0.42%)	-0.84% (-2.6-0.48%)	-54 (-170-30)	-570 (-1800-330)
Iraq	2000	Female	85 to 89	19% (6.8-41%)	81% (59-93%)	0.3 (0.062-0.9)	-0.77% (-2.2-0.33%)	-0.93% (-2.6-0.35%)	-39 (-110-14)	-280 (-810-120)
Iraq	2000	Female	90 to 94	22% (7.4-45%)	78% (55-93%)	0.35 (0.068-0.93)	-0.86% (-2.4-0.33%)	-1.1% (-2.9-0.33%)	-18 (-48-5.7)	-88 (-240-33)
Iraq	2000	Female	95 plus	27% (8.8-52%)	73% (48-91%)	0.41 (0.069-1.1)	-0.98% (-2.9-0.77%)	-1.2% (-3.5-0.85%)	-6.1 (-17-4.2)	-19 (-56-15)
Iraq	2000	Male	All Ages	9.4% (6.5-13%)	91% (87-94%)	0.25 (0.18-0.34)	0.81% (0.44-1.3%)	0.85% (0.23-1.7%)	710 (180-1500)	40000 (22000-64000)
Iraq	2000	Male	15 to 19	7.9% (3.3-15%)	92% (85-97%)	0.13 (0.04-0.29)	0.71% (0.4-1.2%)	0.72% (0.25-1.5%)	16 (5.6-37)	2000 (1000-3500)
Iraq	2000	Male	20 to 24	12% (4.9-24%)	88% (76-95%)	0.29 (0.11-0.59)	1.6% (0.88-2.7%)	1.6% (0.67-3.3%)	52 (21-110)	5100 (2800-9000)
Iraq	2000	Male	25 to 29	15% (6.6-28%)	85% (72-93%)	0.46 (0.19-0.84)	2% (1.1-3.3%)	2.5% (1.1-4.4%)	76 (33-140)	6100 (3300-10000)
Iraq	2000	Male	30 to 34	17% (6.6-32%)	83% (68-93%)	0.59 (0.25-1)	2.1% (1.2-3.5%)	2.7% (1.2-4.8%)	79 (35-140)	5600 (3000-9100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	2000	Male	35 to 39	19% (7.4-34%)	81% (66-93%)	0.68 (0.25-1.3)	2.2% (0.98-3.9%)	2.8% (1.1-5.5%)	69 (25-140)	4300 (1900-7900)
Iraq	2000	Male	40 to 44	19% (8.7-34%)	81% (66-91%)	0.72 (0.28-1.4)	2.1% (0.83-4%)	2.7% (0.85-5.4%)	67 (19-140)	3600 (1300-7300)
Iraq	2000	Male	45 to 49	18% (7.7-33%)	82% (67-92%)	0.75 (0.24-1.6)	2% (0.64-4.4%)	2.5% (0.65-5.7%)	91 (24-210)	4300 (1400-9400)
Iraq	2000	Male	50 to 54	18% (7.7-33%)	82% (67-92%)	0.73 (0.23-1.6)	1.7% (0.35-4.1%)	2.1% (0.23-5.2%)	71 (7.7-180)	2900 (560-7100)
Iraq	2000	Male	55 to 59	17% (7.3-32%)	83% (68-93%)	0.7 (0.21-1.6)	1.5% (0.1-3.9%)	1.8% (-0.028-4.8%)	81 (-1.2-220)	2900 (190-7600)
Iraq	2000	Male	60 to 64	16% (6.9-31%)	84% (69-93%)	0.63 (0.18-1.5)	1.1% (-0.2-3.5%)	1.3% (-0.35-4.2%)	68 (-18-230)	2000 (-340-6600)
Iraq	2000	Male	65 to 69	16% (7.1-29%)	84% (71-93%)	0.51 (0.15-1.1)	0.61% (-0.46-2.4%)	0.65% (-0.63-2.9%)	39 (-40-180)	1000 (-810-4100)
Iraq	2000	Male	70 to 74	14% (6.5-25%)	86% (75-94%)	0.38 (0.11-0.88)	0.23% (-0.58-1.7%)	0.21% (-0.76-1.9%)	14 (-49-130)	320 (-830-2400)
Iraq	2000	Male	75 to 79	12% (5.4-23%)	88% (77-95%)	0.29 (0.074-0.72)	0.053% (-0.66-1.2%)	0.016% (-0.85-1.4%)	1.1 (-55-86)	57 (-730-1300)
Iraq	2000	Male	80 to 84	11% (4.5-22%)	89% (78-96%)	0.24 (0.064-0.59)	-0.037% (-0.61-0.78%)	-0.087% (-0.76-0.83%)	-4.2 (-38-41)	-21 (-370-460)
Iraq	2000	Male	85 to 89	11% (4.3-21%)	89% (79-96%)	0.22 (0.058-0.56)	-0.1% (-0.62-0.53%)	-0.17% (-0.78-0.56%)	-5.2 (-24-17)	-26 (-170-140)
Iraq	2000	Male	90 to 94	12% (4.8-24%)	88% (76-95%)	0.23 (0.058-0.61)	-0.12% (-0.69-0.48%)	-0.21% (-0.9-0.49%)	-2.6 (-11-6.1)	-9.1 (-52-37)
Iraq	2000	Male	95 plus	15% (5.7-31%)	85% (69-94%)	0.26 (0.053-0.75)	-0.2% (-0.95-0.59%)	-0.34% (-1.3-0.56%)	-1.2 (-4.3-2)	-2.7 (-13-8.5)
Iraq	2005	Female	All Ages	12% (7.1-18%)	88% (82-93%)	0.18 (0.13-0.25)	0.16% (-0.15-0.52%)	-0.22% (-0.93-0.54%)	-180 (-800-470)	8100 (-8200-27000)
Iraq	2005	Female	15 to 19	11% (3.2-23%)	89% (77-97%)	0.15 (0.038-0.34)	0.44% (0.26-0.71%)	0.47% (0.21-0.96%)	8.7 (3.8-17)	1200 (690-1900)
Iraq	2005	Female	20 to 24	14% (4.1-29%)	86% (71-96%)	0.24 (0.073-0.49)	0.79% (0.49-1.2%)	0.75% (0.3-1.5%)	15 (6-30)	2200 (1300-3300)
Iraq	2005	Female	25 to 29	16% (4.3-31%)	84% (69-96%)	0.29 (0.081-0.58)	0.91% (0.58-1.4%)	0.99% (0.44-1.8%)	21 (8.8-39)	2400 (1400-3700)
Iraq	2005	Female	30 to 34	17% (4.3-35%)	83% (65-96%)	0.34 (0.11-0.64)	0.84% (0.43-1.4%)	1% (0.32-1.9%)	23 (7-47)	2200 (1100-3700)
Iraq	2005	Female	35 to 39	19% (4.9-36%)	81% (64-95%)	0.4 (0.13-0.74)	0.74% (0.23-1.5%)	1% (0.16-2.3%)	27 (4.9-61)	1900 (580-3900)
Iraq	2005	Female	40 to 44	20% (7.4-36%)	80% (64-93%)	0.44 (0.16-0.83)	0.78% (0.092-1.8%)	1.2% (0.16-2.8%)	30 (3.1-75)	1600 (180-3700)
Iraq	2005	Female	45 to 49	20% (7.4-37%)	80% (63-93%)	0.47 (0.16-0.96)	0.67% (-0.37-2.1%)	1% (-0.35-3.2%)	29 (-9.9-92)	1300 (-640-4100)
Iraq	2005	Female	50 to 54	21% (8.7-39%)	79% (61-91%)	0.48 (0.15-1.1)	0.46% (-0.79-2.2%)	0.73% (-0.93-3.2%)	30 (-38-140)	1100 (-1800-5300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	2005	Female	55 to 59	22% (9-41%)	78% (59-91%)	0.48 (0.13-1.1)	0.16% (-1.3-1.9%)	0.31% (-1.5-2.8%)	11 (-53-98)	260 (-2000-3200)
Iraq	2005	Female	60 to 64	23% (9.6-43%)	77% (57-90%)	0.46 (0.12-1.1)	-0.36% (-2.1-4%)	-0.39% (-2.4-1.9%)	-19 (-120-91)	-660 (-3700-2500)
Iraq	2005	Female	65 to 69	24% (11-42%)	76% (58-89%)	0.41 (0.1-0.91)	-0.73% (-2.2-0.72%)	-0.85% (-2.7-0.96%)	-53 (-170-62)	-1300 (-4100-1400)
Iraq	2005	Female	70 to 74	23% (9.9-41%)	77% (59-90%)	0.32 (0.081-0.81)	-0.92% (-2.4-0.3%)	-1.1% (-2.9-0.38%)	-77 (-210-25)	-1500 (-3900-450)
Iraq	2005	Female	75 to 79	20% (8-40%)	80% (60-92%)	0.26 (0.057-0.77)	-0.95% (-2.6-0.056%)	-1.1% (-3-0.069%)	-82 (-220-5.1)	-1200 (-3200-74)
Iraq	2005	Female	80 to 84	19% (6.1-40%)	81% (60-94%)	0.23 (0.048-0.7)	-0.86% (-2.3--0.005%)	-1% (-2.7--0.02%)	-69 (-190--1.3)	-740 (-2000--4.2)
Iraq	2005	Female	85 to 89	19% (6.3-41%)	81% (59-94%)	0.24 (0.041-0.73)	-0.91% (-2.4--0.076%)	-1.1% (-2.9--0.11%)	-49 (-130--4.6)	-360 (-950--29)
Iraq	2005	Female	90 to 94	22% (7.8-44%)	78% (56-92%)	0.27 (0.051-0.76)	-1% (-2.6--0.1%)	-1.2% (-3.1--0.15%)	-23 (-58--2.7)	-110 (-290--11)
Iraq	2005	Female	95 plus	27% (8.6-51%)	73% (49-91%)	0.32 (0.052-0.93)	-1.1% (-2.9-0.00071%)	-1.4% (-3.6--0.037%)	-7.3 (-19--0.18)	-24 (-61-0.016)
Iraq	2005	Male	All Ages	9.4% (6.4-13%)	91% (87-94%)	0.2 (0.14-0.27)	0.54% (0.29-0.91%)	0.47% (0.06-1.1%)	500 (64-1200)	34000 (18000-56000)
Iraq	2005	Male	15 to 19	7.9% (3.2-15%)	92% (85-97%)	0.1 (0.031-0.22)	0.46% (0.24-0.8%)	0.34% (0.12-0.79%)	15 (5.1-33)	2000 (1100-3500)
Iraq	2005	Male	20 to 24	12% (4.9-23%)	88% (77-95%)	0.23 (0.079-0.46)	0.86% (0.46-1.5%)	0.69% (0.25-1.5%)	46 (18-96)	4900 (2700-8300)
Iraq	2005	Male	25 to 29	15% (6.3-28%)	85% (72-94%)	0.35 (0.14-0.67)	1.2% (0.66-2%)	1.2% (0.47-2.3%)	67 (27-120)	5800 (3100-9600)
Iraq	2005	Male	30 to 34	17% (6.4-33%)	83% (67-94%)	0.45 (0.19-0.78)	1.3% (0.72-2.2%)	1.4% (0.61-2.6%)	71 (31-130)	5300 (2800-8800)
Iraq	2005	Male	35 to 39	19% (7.1-34%)	81% (66-93%)	0.52 (0.19-0.98)	1.4% (0.67-2.6%)	1.6% (0.57-3.2%)	70 (24-140)	4600 (2100-8400)
Iraq	2005	Male	40 to 44	19% (8.4-33%)	81% (67-92%)	0.55 (0.21-1.1)	1.4% (0.58-2.6%)	1.6% (0.5-3.3%)	56 (17-120)	3300 (1300-6400)
Iraq	2005	Male	45 to 49	18% (7.6-33%)	82% (67-92%)	0.58 (0.19-1.2)	1.3% (0.41-3%)	1.5% (0.31-3.9%)	56 (12-140)	2800 (860-6500)
Iraq	2005	Male	50 to 54	18% (7.7-33%)	82% (67-92%)	0.57 (0.18-1.3)	1.1% (0.056-2.9%)	1.2% (-0.11-3.6%)	63 (-5.7-190)	2800 (140-7500)
Iraq	2005	Male	55 to 59	17% (7.4-32%)	83% (68-93%)	0.55 (0.16-1.3)	0.88% (-0.16-2.8%)	0.96% (-0.33-3.4%)	46 (-16-170)	1700 (-310-5600)
Iraq	2005	Male	60 to 64	16% (7-31%)	84% (69-93%)	0.49 (0.14-1.1)	0.57% (-0.44-2.5%)	0.6% (-0.64-2.9%)	38 (-38-190)	1200 (-870-5400)
Iraq	2005	Male	65 to 69	15% (6.9-28%)	85% (72-93%)	0.39 (0.12-0.84)	0.2% (-0.64-1.5%)	0.17% (-0.84-1.7%)	12 (-56-120)	390 (-1200-2900)
Iraq	2005	Male	70 to 74	14% (6.6-25%)	86% (75-93%)	0.29 (0.083-0.69)	-0.021% (-0.69-0.96%)	-0.076% (-0.86-1%)	-5.5 (-65-77)	-32 (-1100-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	2005	Male	75 to 79	12% (5.2-23%)	88% (77-95%)	0.22 (0.057-0.58)	-0.11% (-0.7-0.64%)	-0.16% (-0.87-0.68%)	-11 (-58-46)	-120 (-780-690)
Iraq	2005	Male	80 to 84	11% (4.4-21%)	89% (79-96%)	0.19 (0.048-0.47)	-0.16% (-0.71-0.33%)	-0.22% (-0.89-0.35%)	-11 (-46-17)	-99 (-450-210)
Iraq	2005	Male	85 to 89	11% (4.2-22%)	89% (78-96%)	0.17 (0.045-0.44)	-0.19% (-0.68-0.2%)	-0.28% (-0.86-0.19%)	-8.9 (-28-6.4)	-54 (-190-57)
Iraq	2005	Male	90 to 94	12% (4.9-23%)	88% (77-95%)	0.18 (0.044-0.48)	-0.2% (-0.73-0.22%)	-0.3% (-0.96-0.19%)	-4 (-13-2.6)	-16 (-59-17)
Iraq	2005	Male	95 plus	15% (5.4-31%)	85% (69-95%)	0.2 (0.042-0.61)	-0.28% (-0.92-0.25%)	-0.42% (-1.2-0.18%)	-1.5 (-4.5-0.66)	-4 (-13-3.6)
Iraq	2010	Female	All Ages	12% (7.4-18%)	88% (82-93%)	0.18 (0.12-0.24)	0.15% (-0.16-0.53%)	-0.22% (-0.95-0.58%)	-200 (-870-490)	8200 (-8600-28000)
Iraq	2010	Female	15 to 19	10% (3.8-22%)	90% (78-96%)	0.14 (0.044-0.33)	0.51% (0.32-0.82%)	0.63% (0.3-1.2%)	8.8 (4.1-18)	1200 (740-2000)
Iraq	2010	Female	20 to 24	14% (4.5-28%)	86% (72-96%)	0.23 (0.07-0.47)	0.87% (0.55-1.3%)	0.92% (0.39-1.8%)	15 (6-30)	2200 (1300-3500)
Iraq	2010	Female	25 to 29	15% (4.4-31%)	85% (69-96%)	0.27 (0.081-0.56)	0.94% (0.61-1.4%)	1.1% (0.45-2.1%)	19 (7.2-40)	2400 (1400-3800)
Iraq	2010	Female	30 to 34	17% (4.5-35%)	83% (65-95%)	0.32 (0.098-0.61)	0.83% (0.43-1.3%)	1% (0.32-2%)	22 (6.6-49)	2200 (1100-3900)
Iraq	2010	Female	35 to 39	19% (5-37%)	81% (63-95%)	0.38 (0.12-0.7)	0.7% (0.22-1.4%)	0.99% (0.15-2.2%)	26 (4.1-63)	1900 (580-4000)
Iraq	2010	Female	40 to 44	20% (7.1-38%)	80% (62-93%)	0.42 (0.15-0.81)	0.71% (0.025-1.7%)	1.1% (0.099-2.8%)	36 (3.1-91)	1900 (70-4600)
Iraq	2010	Female	45 to 49	21% (7.2-38%)	79% (62-93%)	0.47 (0.15-1)	0.61% (-0.4-2%)	1% (-0.37-3.1%)	32 (-12-100)	1400 (-810-4500)
Iraq	2010	Female	50 to 54	21% (8.6-40%)	79% (60-91%)	0.48 (0.14-1)	0.4% (-0.86-2.1%)	0.67% (-1.1-3.2%)	25 (-37-120)	830 (-1700-4600)
Iraq	2010	Female	55 to 59	22% (9-41%)	78% (59-91%)	0.48 (0.14-1.1)	0.093% (-1.3-1.8%)	0.24% (-1.6-2.7%)	13 (-86-140)	240 (-3200-4700)
Iraq	2010	Female	60 to 64	23% (9.8-43%)	77% (57-90%)	0.45 (0.12-1.1)	-0.4% (-2-1.4%)	-0.44% (-2.5-1.9%)	-20 (-110-86)	-690 (-3400-2500)
Iraq	2010	Female	65 to 69	24% (11-41%)	76% (59-89%)	0.39 (0.11-0.91)	-0.74% (-2.3-0.71%)	-0.87% (-2.8-0.94%)	-56 (-180-58)	-1400 (-4400-1300)
Iraq	2010	Female	70 to 74	22% (9.7-39%)	78% (61-90%)	0.31 (0.079-0.78)	-0.88% (-2.3-0.26%)	-1% (-2.8-0.37%)	-79 (-210-25)	-1500 (-4000-440)
Iraq	2010	Female	75 to 79	20% (7.7-41%)	80% (59-92%)	0.25 (0.052-0.75)	-0.9% (-2.5-0.043%)	-1.1% (-2.9-0.018%)	-86 (-230-1.5)	-1300 (-3400-60)
Iraq	2010	Female	80 to 84	18% (6.3-40%)	82% (60-94%)	0.23 (0.047-0.7)	-0.83% (-2.3--0.013%)	-0.99% (-2.8--0.023%)	-71 (-200--1.6)	-750 (-2100--12)
Iraq	2010	Female	85 to 89	19% (6.5-41%)	81% (59-93%)	0.23 (0.044-0.67)	-0.9% (-2.5--0.097%)	-1.1% (-3--0.13%)	-50 (-140--6)	-370 (-1000--40)
Iraq	2010	Female	90 to 94	21% (7.5-43%)	79% (57-93%)	0.25 (0.05-0.68)	-0.98% (-2.5--0.11%)	-1.2% (-3--0.19%)	-23 (-58--3.5)	-120 (-290--13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	2010	Female	95 plus	26% (9-49%)	74% (51-91%)	0.3 (0.053-0.84)	-1.1% (-2.8--0.022%)	-1.4% (-3.4--0.084%)	-7.8 (-19--0.45)	-25 (-63--0.46)
Iraq	2010	Male	All Ages	9.2% (6.3-13%)	91% (87-94%)	0.19 (0.13-0.26)	0.56% (0.29-0.95%)	0.47% (0.0099-1.2%)	490 (11-1300)	36000 (18000-62000)
Iraq	2010	Male	15 to 19	7.7% (3.3-15%)	92% (85-97%)	0.097 (0.03-0.21)	0.58% (0.33-0.97%)	0.48% (0.16-1.1%)	15 (5-35)	2100 (1100-3500)
Iraq	2010	Male	20 to 24	12% (4.8-23%)	88% (77-95%)	0.22 (0.074-0.43)	1.2% (0.68-1.9%)	1% (0.39-2.1%)	46 (17-99)	5100 (2800-8700)
Iraq	2010	Male	25 to 29	15% (6.1-28%)	85% (72-94%)	0.34 (0.13-0.66)	1.5% (0.86-2.4%)	1.6% (0.66-2.9%)	65 (26-130)	5800 (3200-9900)
Iraq	2010	Male	30 to 34	17% (6.3-32%)	83% (68-94%)	0.43 (0.18-0.76)	1.5% (0.87-2.4%)	1.8% (0.81-3.1%)	73 (31-140)	5700 (3100-9400)
Iraq	2010	Male	35 to 39	18% (7.3-33%)	82% (67-93%)	0.49 (0.18-0.91)	1.5% (0.75-2.7%)	1.8% (0.67-3.5%)	73 (25-150)	5000 (2400-9100)
Iraq	2010	Male	40 to 44	19% (8.6-32%)	81% (68-91%)	0.52 (0.2-0.98)	1.4% (0.6-2.7%)	1.6% (0.52-3.5%)	71 (21-150)	4200 (1700-8300)
Iraq	2010	Male	45 to 49	18% (7.3-33%)	82% (67-93%)	0.55 (0.18-1.1)	1.3% (0.4-3%)	1.5% (0.28-3.9%)	60 (10-160)	3100 (910-7200)
Iraq	2010	Male	50 to 54	17% (7.5-33%)	83% (67-93%)	0.55 (0.17-1.2)	1% (0.038-2.7%)	1.2% (-0.14-3.4%)	50 (-6.1-150)	2200 (88-6200)
Iraq	2010	Male	55 to 59	17% (7.4-31%)	83% (69-93%)	0.52 (0.16-1.2)	0.8% (-0.24-2.6%)	0.87% (-0.45-3.2%)	55 (-27-210)	2100 (-610-7300)
Iraq	2010	Male	60 to 64	16% (6.8-30%)	84% (70-93%)	0.46 (0.13-1.1)	0.49% (-0.48-2.2%)	0.5% (-0.68-2.7%)	30 (-40-160)	990 (-930-4600)
Iraq	2010	Male	65 to 69	15% (7-28%)	85% (72-93%)	0.38 (0.12-0.85)	0.14% (-0.66-1.4%)	0.099% (-0.87-1.6%)	7.5 (-65-120)	300 (-1400-3000)
Iraq	2010	Male	70 to 74	14% (6.5-25%)	86% (75-93%)	0.28 (0.085-0.69)	-0.061% (-0.76-0.88%)	-0.13% (-0.96-0.96%)	-9.7 (-78-76)	-100 (-1300-1500)
Iraq	2010	Male	75 to 79	12% (5.4-23%)	88% (77-95%)	0.22 (0.053-0.58)	-0.13% (-0.76-0.5%)	-0.2% (-0.93-0.53%)	-14 (-67-39)	-160 (-890-610)
Iraq	2010	Male	80 to 84	11% (4.2-21%)	89% (79-96%)	0.18 (0.046-0.46)	-0.18% (-0.72-0.32%)	-0.25% (-0.9-0.31%)	-13 (-47-16)	-120 (-470-200)
Iraq	2010	Male	85 to 89	11% (4.4-20%)	89% (80-96%)	0.17 (0.044-0.42)	-0.21% (-0.7-0.16%)	-0.3% (-0.89-0.14%)	-9.3 (-28-4.3)	-57 (-190-45)
Iraq	2010	Male	90 to 94	12% (5.1-24%)	88% (76-95%)	0.18 (0.044-0.47)	-0.22% (-0.77-0.19%)	-0.32% (-1-0.16%)	-4.2 (-13-2.1)	-17 (-61-15)
Iraq	2010	Male	95 plus	15% (5.6-31%)	85% (69-94%)	0.2 (0.042-0.59)	-0.3% (-0.99-0.2%)	-0.46% (-1.3-0.15%)	-1.7 (-4.7-0.55)	-4.4 (-14-2.9)
Iraq	2016	Female	All Ages	12% (7.2-18%)	88% (82-93%)	0.21 (0.12-0.34)	0.24% (-0.15-0.84%)	-0.047% (-0.9-1.2%)	-43 (-890-1200)	15000 (-8800-54000)
Iraq	2016	Female	15 to 19	11% (3.4-24%)	89% (76-97%)	0.17 (0.041-0.43)	0.47% (0.27-0.79%)	0.52% (0.2-1.2%)	11 (4.5-23)	1500 (820-2500)
Iraq	2016	Female	20 to 24	14% (4-29%)	86% (71-96%)	0.27 (0.066-0.6)	0.84% (0.5-1.4%)	0.82% (0.32-1.8%)	18 (6.6-40)	2700 (1600-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	2016	Female	25 to 29	15% (4-33%)	85% (67-96%)	0.32 (0.078-0.7)	0.95% (0.58-1.5%)	1.1% (0.41-2.3%)	23 (8.7-54)	2900 (1700-4800)
Iraq	2016	Female	30 to 34	17% (5-36%)	83% (64-95%)	0.37 (0.11-0.78)	0.88% (0.43-1.6%)	1.1% (0.35-2.4%)	28 (8-68)	2700 (1300-5400)
Iraq	2016	Female	35 to 39	19% (5.9-36%)	81% (64-94%)	0.44 (0.14-0.89)	0.8% (0.23-1.7%)	1.2% (0.2-2.8%)	36 (5.6-97)	2600 (720-6100)
Iraq	2016	Female	40 to 44	20% (7.9-37%)	80% (63-92%)	0.49 (0.17-1)	0.84% (0.074-2.2%)	1.4% (0.16-3.6%)	51 (6.3-140)	2700 (240-7300)
Iraq	2016	Female	45 to 49	21% (7.2-39%)	79% (61-93%)	0.53 (0.16-1.2)	0.78% (-0.3-2.5%)	1.3% (-0.25-4%)	57 (-11-180)	2500 (-930-8200)
Iraq	2016	Female	50 to 54	22% (8.1-41%)	78% (59-92%)	0.55 (0.15-1.3)	0.61% (-0.81-2.9%)	1% (-0.95-4.3%)	47 (-42-210)	1700 (-2100-8000)
Iraq	2016	Female	55 to 59	23% (8.6-42%)	77% (58-91%)	0.55 (0.14-1.3)	0.35% (-1.2-2.7%)	0.6% (-1.4-3.7%)	30 (-67-190)	830 (-2700-6300)
Iraq	2016	Female	60 to 64	24% (9.7-42%)	76% (58-90%)	0.52 (0.12-1.3)	-0.13% (-1.8-2.2%)	-0.09% (-2.1-3%)	-5.7 (-160-210)	-340 (-4900-6100)
Iraq	2016	Female	65 to 69	24% (10-42%)	76% (58-90%)	0.46 (0.11-1.1)	-0.49% (-2.2-1.7%)	-0.57% (-2.6-2.2%)	-36 (-170-140)	-970 (-4200-3500)
Iraq	2016	Female	70 to 74	23% (9.9-41%)	77% (59-90%)	0.38 (0.086-0.99)	-0.71% (-2.2-1.1%)	-0.86% (-2.7-1.4%)	-65 (-200-97)	-1300 (-3900-1900)
Iraq	2016	Female	75 to 79	21% (8.4-40%)	79% (60-92%)	0.31 (0.062-0.95)	-0.8% (-2.3-0.66%)	-0.97% (-2.8-0.73%)	-82 (-240-57)	-1200 (-3500-900)
Iraq	2016	Female	80 to 84	19% (7-40%)	81% (60-93%)	0.28 (0.056-0.82)	-0.75% (-2.2-0.37%)	-0.91% (-2.7-0.41%)	-70 (-210-29)	-740 (-2200-360)
Iraq	2016	Female	85 to 89	20% (7-41%)	80% (59-93%)	0.28 (0.054-0.86)	-0.84% (-2.4-0.17%)	-1% (-2.9-0.13%)	-52 (-150-7.8)	-380 (-1100-77)
Iraq	2016	Female	90 to 94	23% (8.3-45%)	77% (55-92%)	0.31 (0.055-0.87)	-0.92% (-2.5-0.21%)	-1.1% (-3-0.2%)	-24 (-65-4.2)	-120 (-330-28)
Iraq	2016	Female	95 plus	27% (9.2-52%)	73% (48-91%)	0.36 (0.061-1)	-1% (-2.9-0.33%)	-1.3% (-3.5-0.32%)	-8.1 (-21-2)	-26 (-70-8.2)
Iraq	2016	Male	All Ages	9.2% (6.3-13%)	91% (87-94%)	0.22 (0.13-0.36)	0.64% (0.29-1.2%)	0.59% (0.032-1.5%)	760 (36-1900)	50000 (23000-93000)
Iraq	2016	Male	15 to 19	7.7% (3.2-15%)	92% (85-97%)	0.11 (0.031-0.26)	0.49% (0.26-0.89%)	0.37% (0.11-0.88%)	19 (5.8-44)	2500 (1300-4400)
Iraq	2016	Male	20 to 24	12% (4.9-23%)	88% (77-95%)	0.25 (0.077-0.52)	0.94% (0.48-1.7%)	0.74% (0.23-1.7%)	56 (19-120)	6200 (3300-11000)
Iraq	2016	Male	25 to 29	15% (6.2-27%)	85% (73-94%)	0.39 (0.13-0.82)	1.3% (0.68-2.2%)	1.2% (0.43-2.7%)	81 (26-170)	7300 (3800-13000)
Iraq	2016	Male	30 to 34	17% (6.6-32%)	83% (68-93%)	0.5 (0.18-1)	1.4% (0.76-2.6%)	1.6% (0.58-3.1%)	90 (33-190)	7000 (3500-13000)
Iraq	2016	Male	35 to 39	18% (7.4-33%)	82% (67-93%)	0.56 (0.19-1.1)	1.5% (0.71-3%)	1.7% (0.57-3.8%)	97 (29-220)	6600 (2900-13000)
Iraq	2016	Male	40 to 44	19% (8.4-33%)	81% (67-92%)	0.59 (0.22-1.2)	1.5% (0.62-3%)	1.7% (0.53-3.9%)	100 (27-230)	5900 (2300-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Iraq	2016	Male	45 to 49	18% (7.3-33%)	82% (67-93%)	0.61 (0.19-1.4)	1.4% (0.43-3.4%)	1.7% (0.33-4.3%)	110 (19-280)	5400 (1500-13000)
Iraq	2016	Male	50 to 54	17% (7.4-33%)	83% (67-93%)	0.61 (0.19-1.4)	1.2% (0.11-3.3%)	1.4% (-0.055-4.2%)	82 (-2.7-250)	3600 (330-10000)
Iraq	2016	Male	55 to 59	16% (7.2-31%)	84% (69-93%)	0.58 (0.15-1.4)	1% (-0.11-3.2%)	1.1% (-0.29-4%)	66 (-16-230)	2500 (-240-7800)
Iraq	2016	Male	60 to 64	16% (6.9-30%)	84% (70-93%)	0.52 (0.13-1.3)	0.69% (-0.37-2.8%)	0.75% (-0.56-3.3%)	62 (-45-280)	2000 (-1000-8200)
Iraq	2016	Male	65 to 69	15% (6.8-27%)	85% (73-93%)	0.43 (0.11-1)	0.33% (-0.57-2%)	0.32% (-0.79-2.3%)	25 (-61-190)	730 (-1300-4600)
Iraq	2016	Male	70 to 74	14% (6.2-25%)	86% (75-94%)	0.32 (0.086-0.73)	0.076% (-0.65-1.4%)	0.032% (-0.86-1.6%)	2.8 (-76-140)	150 (-1300-2700)
Iraq	2016	Male	75 to 79	12% (5.1-23%)	88% (77-95%)	0.24 (0.056-0.67)	-0.026% (-0.64-1%)	-0.079% (-0.81-1.1%)	-6.6 (-69-83)	-37 (-900-1300)
Iraq	2016	Male	80 to 84	11% (4.3-21%)	89% (79-96%)	0.2 (0.05-0.55)	-0.093% (-0.63-0.6%)	-0.16% (-0.79-0.64%)	-9.3 (-48-39)	-68 (-460-440)
Iraq	2016	Male	85 to 89	11% (4.2-20%)	89% (80-96%)	0.19 (0.048-0.49)	-0.14% (-0.62-0.42%)	-0.22% (-0.8-0.43%)	-7.5 (-28-16)	-42 (-190-130)
Iraq	2016	Male	90 to 94	12% (4.8-23%)	88% (77-95%)	0.19 (0.046-0.51)	-0.15% (-0.69-0.43%)	-0.25% (-0.92-0.4%)	-3.4 (-13-5.5)	-12 (-58-35)
Iraq	2016	Male	95 plus	14% (5.5-30%)	86% (70-94%)	0.22 (0.042-0.64)	-0.22% (-0.86-0.41%)	-0.36% (-1.2-0.35%)	-1.4 (-4.4-1.3)	-3.3 (-13-6.4)
Jordan	1990	Female	All Ages	10% (6.4-16%)	90% (84-94%)	0.057 (0.04-0.076)	-0.015% (-0.19-0.12%)	-0.42% (-0.94--0.056%)	-29 (-64--3.6)	-63 (-840-520)
Jordan	1990	Female	15 to 19	9.5% (3.3-21%)	90% (79-97%)	0.05 (0.016-0.11)	0.37% (0.23-0.55%)	0.34% (0.18-0.63%)	0.4 (0.21-0.75)	96 (57-150)
Jordan	1990	Female	20 to 24	12% (3.8-27%)	88% (73-96%)	0.079 (0.026-0.16)	0.67% (0.43-0.95%)	0.42% (0.19-0.75%)	0.48 (0.22-0.91)	150 (94-240)
Jordan	1990	Female	25 to 29	14% (3.8-29%)	86% (71-96%)	0.097 (0.029-0.2)	0.66% (0.45-0.92%)	0.39% (0.18-0.72%)	0.41 (0.19-0.8)	130 (82-190)
Jordan	1990	Female	30 to 34	16% (3.7-33%)	84% (67-96%)	0.12 (0.035-0.25)	0.56% (0.37-0.8%)	0.41% (0.18-0.74%)	0.4 (0.19-0.75)	90 (55-130)
Jordan	1990	Female	35 to 39	18% (4.5-35%)	82% (65-95%)	0.14 (0.041-0.28)	0.43% (0.24-0.63%)	0.43% (0.17-0.83%)	0.47 (0.18-0.93)	61 (34-93)
Jordan	1990	Female	40 to 44	19% (6.5-36%)	81% (64-94%)	0.16 (0.054-0.32)	0.29% (0.017-0.51%)	0.4% (0.031-0.84%)	0.56 (0.042-1.2)	43 (2.3-78)
Jordan	1990	Female	45 to 49	19% (6.6-36%)	81% (64-93%)	0.17 (0.052-0.35)	0.029% (-0.42-0.36%)	0.053% (-0.63-0.6%)	0.1 (-1.2-1.2)	4.8 (-70-57)
Jordan	1990	Female	50 to 54	20% (7.7-39%)	80% (61-92%)	0.17 (0.055-0.36)	-0.24% (-0.92-0.19%)	-0.28% (-1.2-0.31%)	-0.77 (-3.3-0.85)	-43 (-170-34)
Jordan	1990	Female	55 to 59	21% (7.9-41%)	79% (59-92%)	0.17 (0.052-0.37)	-0.49% (-1.5-0.028%)	-0.61% (-1.9-0.1%)	-2 (-6-0.32)	-85 (-260-5.6)
Jordan	1990	Female	60 to 64	22% (8.4-41%)	78% (59-92%)	0.16 (0.044-0.37)	-0.72% (-1.9--0.054%)	-0.9% (-2.4--0.059%)	-3.6 (-9.9--0.23)	-120 (-320--9.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	1990	Female	65 to 69	22% (9.6-40%)	78% (60-90%)	0.14 (0.036-0.31)	-0.86% (-2--0.14%)	-1.1% (-2.5--0.16%)	-5.4 (-13--0.72)	-140 (-330--22)
Jordan	1990	Female	70 to 74	21% (8.8-38%)	79% (62-91%)	0.11 (0.028-0.27)	-0.75% (-1.8--0.1%)	-0.92% (-2.2--0.11%)	-4.9 (-12--0.57)	-99 (-240--13)
Jordan	1990	Female	75 to 79	19% (7.1-38%)	81% (62-93%)	0.093 (0.021-0.25)	-0.67% (-1.9--0.081%)	-0.81% (-2.2--0.096%)	-5.1 (-14--0.6)	-78 (-210--9.1)
Jordan	1990	Female	80 to 84	17% (6.3-37%)	83% (63-94%)	0.084 (0.018-0.23)	-0.62% (-1.9--0.069%)	-0.75% (-2.2--0.083%)	-4.6 (-13--0.48)	-50 (-140--5.1)
Jordan	1990	Female	85 to 89	18% (6.2-38%)	82% (62-94%)	0.085 (0.018-0.25)	-0.62% (-1.9--0.066%)	-0.75% (-2.3--0.087%)	-3.2 (-9.8--0.35)	-24 (-74--2.5)
Jordan	1990	Female	90 to 94	21% (7.5-43%)	79% (57-92%)	0.097 (0.02-0.28)	-0.66% (-1.9--0.07%)	-0.82% (-2.3--0.097%)	-1.4 (-4--0.17)	-7.1 (-20--0.74)
Jordan	1990	Female	95 plus	25% (7.9-50%)	75% (50-92%)	0.12 (0.021-0.33)	-0.74% (-2.1--0.05%)	-0.93% (-2.6--0.091%)	-0.48 (-1.4--0.045)	-1.5 (-4.5--0.11)
Jordan	1990	Male	All Ages	8% (5.6-11%)	92% (89-94%)	0.065 (0.048-0.087)	0.3% (0.16-0.43%)	0.095% (-0.13-0.3%)	7.6 (-10-24)	1500 (780-2200)
Jordan	1990	Male	15 to 19	7.1% (3-14%)	93% (86-97%)	0.036 (0.012-0.079)	0.55% (0.33-0.83%)	0.34% (0.11-0.78%)	0.69 (0.21-1.6)	170 (100-270)
Jordan	1990	Male	20 to 24	11% (4.4-21%)	89% (79-96%)	0.081 (0.029-0.16)	1.2% (0.73-1.7%)	0.71% (0.23-1.5%)	1.7 (0.54-3.8)	370 (220-550)
Jordan	1990	Male	25 to 29	14% (5.3-26%)	86% (74-95%)	0.13 (0.05-0.25)	1.3% (0.86-1.8%)	1.1% (0.43-2%)	2.2 (0.88-4.3)	330 (220-490)
Jordan	1990	Male	30 to 34	15% (5.8-29%)	85% (71-94%)	0.16 (0.066-0.3)	1.1% (0.72-1.6%)	0.99% (0.39-1.9%)	1.8 (0.69-3.4)	220 (140-340)
Jordan	1990	Male	35 to 39	17% (6.5-30%)	83% (70-93%)	0.18 (0.068-0.36)	0.96% (0.57-1.4%)	0.88% (0.28-1.6%)	1.4 (0.45-2.7)	160 (89-230)
Jordan	1990	Male	40 to 44	17% (7.6-30%)	83% (70-92%)	0.2 (0.072-0.38)	0.68% (0.29-1.1%)	0.53% (-0.09-1.1%)	0.97 (-0.15-2.2)	100 (43-170)
Jordan	1990	Male	45 to 49	16% (6.6-30%)	84% (70-93%)	0.2 (0.068-0.43)	0.41% (-0.029-0.77%)	0.26% (-0.47-0.83%)	0.61 (-1.1-2)	69 (-4.8-130)
Jordan	1990	Male	50 to 54	16% (6.9-30%)	84% (70-93%)	0.2 (0.068-0.41)	0.22% (-0.27-0.59%)	0.1% (-0.68-0.65%)	0.32 (-2.1-2.1)	42 (-53-110)
Jordan	1990	Male	55 to 59	15% (6.6-29%)	85% (71-93%)	0.19 (0.062-0.38)	0.1% (-0.38-0.49%)	0.0077% (-0.71-0.56%)	0.029 (-2.8-2.2)	19 (-70-93)
Jordan	1990	Male	60 to 64	15% (6.2-28%)	85% (72-94%)	0.17 (0.05-0.38)	0.027% (-0.46-0.39%)	-0.042% (-0.7-0.45%)	-0.2 (-3.3-2.1)	4.7 (-81-68)
Jordan	1990	Male	65 to 69	14% (6-26%)	86% (74-94%)	0.14 (0.043-0.3)	-0.043% (-0.5-0.26%)	-0.1% (-0.74-0.31%)	-0.55 (-3.8-1.6)	-6.9 (-81-42)
Jordan	1990	Male	70 to 74	13% (5.8-23%)	87% (77-94%)	0.11 (0.032-0.25)	-0.035% (-0.41-0.21%)	-0.074% (-0.56-0.24%)	-0.39 (-2.9-1.3)	-4.4 (-51-27)
Jordan	1990	Male	75 to 79	11% (4.9-21%)	89% (79-95%)	0.08 (0.02-0.21)	-0.022% (-0.36-0.18%)	-0.051% (-0.49-0.22%)	-0.34 (-3.3-1.4)	-2.6 (-44-23)
Jordan	1990	Male	80 to 84	10% (3.7-20%)	90% (80-96%)	0.067 (0.017-0.16)	-0.016% (-0.33-0.16%)	-0.041% (-0.46-0.19%)	-0.3 (-3.3-1.4)	-1.5 (-33-16)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	1990	Male	85 to 89	9.7% (3.9-19%)	90% (81-96%)	0.061 (0.016-0.16)	-0.034% (-0.33-0.14%)	-0.07% (-0.44-0.14%)	-0.31 (-1.9-0.64)	-1.3 (-13-5.5)
Jordan	1990	Male	90 to 94	11% (4.5-22%)	89% (78-96%)	0.064 (0.016-0.16)	-0.054% (-0.35-0.12%)	-0.11% (-0.48-0.12%)	-0.15 (-0.69-0.17)	-0.48 (-3.1-1.1)
Jordan	1990	Male	95 plus	13% (5-28%)	87% (72-95%)	0.074 (0.015-0.21)	-0.045% (-0.41-0.15%)	-0.13% (-0.59-0.11%)	-0.046 (-0.21-0.04)	-0.065 (-0.59-0.22)
Jordan	1995	Female	All Ages	11% (6.2-17%)	89% (83-94%)	0.062 (0.044-0.084)	0.029% (-0.15-0.16%)	-0.37% (-0.88--0.025%)	-30 (-71--1.9)	160 (-790-920)
Jordan	1995	Female	15 to 19	9.5% (2.9-22%)	91% (78-97%)	0.05 (0.014-0.12)	0.38% (0.23-0.57%)	0.35% (0.19-0.64%)	0.46 (0.24-0.85)	120 (68-180)
Jordan	1995	Female	20 to 24	12% (3.5-27%)	88% (73-97%)	0.08 (0.025-0.17)	0.7% (0.43-0.99%)	0.43% (0.21-0.77%)	0.66 (0.32-1.3)	230 (130-340)
Jordan	1995	Female	25 to 29	14% (3.5-29%)	86% (71-96%)	0.097 (0.03-0.2)	0.68% (0.47-0.94%)	0.39% (0.19-0.72%)	0.59 (0.27-1.1)	200 (130-300)
Jordan	1995	Female	30 to 34	16% (4-32%)	84% (68-96%)	0.12 (0.036-0.23)	0.59% (0.37-0.84%)	0.43% (0.2-0.77%)	0.66 (0.31-1.3)	150 (94-230)
Jordan	1995	Female	35 to 39	17% (4.7-34%)	83% (66-95%)	0.14 (0.042-0.28)	0.45% (0.27-0.65%)	0.47% (0.21-0.89%)	0.75 (0.31-1.4)	99 (59-150)
Jordan	1995	Female	40 to 44	18% (6.4-34%)	82% (66-94%)	0.15 (0.049-0.31)	0.32% (0.079-0.56%)	0.47% (0.12-0.9%)	0.78 (0.19-1.6)	60 (14-110)
Jordan	1995	Female	45 to 49	19% (6.6-36%)	81% (64-93%)	0.17 (0.052-0.36)	0.081% (-0.33-0.41%)	0.15% (-0.47-0.68%)	0.32 (-1.1.6)	16 (-64-80)
Jordan	1995	Female	50 to 54	20% (7.9-38%)	80% (62-92%)	0.17 (0.052-0.37)	-0.17% (-0.8-0.25%)	-0.18% (-1-0.44%)	-0.6 (-3.3-1.5)	-38 (-180-56)
Jordan	1995	Female	55 to 59	21% (8.2-40%)	79% (60-92%)	0.17 (0.048-0.38)	-0.44% (-1.4-0.064%)	-0.53% (-1.8-0.14%)	-2.2 (-7.1-0.61)	-98 (-310-15)
Jordan	1995	Female	60 to 64	22% (8.4-43%)	78% (57-92%)	0.16 (0.04-0.38)	-0.68% (-1.9--0.019%)	-0.85% (-2.4--0.013%)	-4.3 (-13--0.065)	-150 (-420--4.3)
Jordan	1995	Female	65 to 69	23% (9.4-41%)	77% (59-91%)	0.14 (0.034-0.32)	-0.83% (-2--0.12%)	-1% (-2.5--0.14%)	-6.2 (-15--0.79)	-160 (-390--22)
Jordan	1995	Female	70 to 74	21% (8.6-38%)	79% (62-91%)	0.11 (0.027-0.28)	-0.71% (-1.7--0.084%)	-0.87% (-2.1--0.095%)	-5.9 (-15--0.6)	-120 (-290--14)
Jordan	1995	Female	75 to 79	18% (7.1-37%)	82% (63-93%)	0.09 (0.019-0.25)	-0.63% (-1.8--0.071%)	-0.76% (-2.2--0.082%)	-4.8 (-13--0.52)	-73 (-210--8.2)
Jordan	1995	Female	80 to 84	17% (5.9-37%)	83% (63-94%)	0.081 (0.016-0.23)	-0.58% (-1.7--0.034%)	-0.7% (-2.1--0.045%)	-4.1 (-12--0.25)	-45 (-130--2.9)
Jordan	1995	Female	85 to 89	17% (5.9-38%)	83% (62-94%)	0.082 (0.017-0.25)	-0.58% (-1.9--0.047%)	-0.7% (-2.3--0.062%)	-3.5 (-11--0.33)	-27 (-87--2.2)
Jordan	1995	Female	90 to 94	20% (6.8-42%)	80% (58-93%)	0.093 (0.018-0.28)	-0.61% (-1.8--0.058%)	-0.75% (-2.2--0.083%)	-1.8 (-5.3--0.2)	-9.1 (-27--0.87)
Jordan	1995	Female	95 plus	24% (8.3-48%)	76% (52-92%)	0.11 (0.02-0.31)	-0.68% (-2--0.05%)	-0.86% (-2.5--0.093%)	-0.56 (-1.6--0.059)	-1.8 (-5.2--0.13)
Jordan	1995	Male	All Ages	8.7% (5.7-12%)	91% (88-94%)	0.072 (0.052-0.095)	0.35% (0.21-0.49%)	0.13% (-0.096-0.34%)	12 (-9.1-33)	2200 (1300-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	1995	Male	15 to 19	6.9% (2.9-14%)	93% (86-97%)	0.035 (0.011-0.077)	0.56% (0.34-0.85%)	0.33% (0.11-0.74%)	0.74 (0.25-1.7)	200 (120-310)
Jordan	1995	Male	20 to 24	11% (4.2-20%)	89% (80-96%)	0.076 (0.029-0.14)	1.2% (0.74-1.7%)	0.68% (0.24-1.4%)	2.1 (0.73-4.5)	480 (300-740)
Jordan	1995	Male	25 to 29	13% (5.3-25%)	87% (75-95%)	0.12 (0.048-0.22)	1.3% (0.86-1.8%)	1% (0.47-1.9%)	3.2 (1.4-6.2)	510 (330-730)
Jordan	1995	Male	30 to 34	15% (5.7-29%)	85% (71-94%)	0.15 (0.064-0.28)	1.1% (0.76-1.6%)	0.96% (0.39-1.8%)	2.8 (1.1-5.4)	380 (240-570)
Jordan	1995	Male	35 to 39	17% (6.5-31%)	83% (69-93%)	0.18 (0.069-0.35)	0.97% (0.62-1.4%)	0.88% (0.28-1.6%)	2.3 (0.67-4.2)	260 (150-380)
Jordan	1995	Male	40 to 44	17% (7.6-30%)	83% (70-92%)	0.19 (0.066-0.38)	0.7% (0.32-1.1%)	0.57% (-0.036-1.2%)	1.4 (-0.093-3.1)	150 (69-240)
Jordan	1995	Male	45 to 49	16% (6.6-31%)	84% (69-93%)	0.2 (0.064-0.43)	0.44% (0.02-0.83%)	0.3% (-0.37-0.89%)	0.88 (-1.1-2.7)	93 (4.4-180)
Jordan	1995	Male	50 to 54	16% (6.7-30%)	84% (70-93%)	0.19 (0.062-0.42)	0.25% (-0.25-0.62%)	0.14% (-0.61-0.69%)	0.55 (-2.5-2.8)	59 (-56-150)
Jordan	1995	Male	55 to 59	15% (6.4-28%)	85% (72-94%)	0.18 (0.057-0.39)	0.12% (-0.35-0.48%)	0.037% (-0.68-0.54%)	0.2 (-3.7-3.1)	32 (-94-130)
Jordan	1995	Male	60 to 64	14% (6.2-28%)	86% (72-94%)	0.16 (0.046-0.37)	0.044% (-0.43-0.39%)	-0.018% (-0.65-0.46%)	-0.11 (-4.2-2.9)	11 (-100-98)
Jordan	1995	Male	65 to 69	14% (5.9-25%)	86% (75-94%)	0.14 (0.041-0.29)	-0.023% (-0.44-0.28%)	-0.076% (-0.66-0.32%)	-0.5 (-4.2-2)	-4.7 (-90-57)
Jordan	1995	Male	70 to 74	13% (5.4-23%)	87% (77-95%)	0.1 (0.029-0.23)	-0.015% (-0.36-0.22%)	-0.047% (-0.51-0.25%)	-0.3 (-3.3-1.6)	-2.3 (-57-34)
Jordan	1995	Male	75 to 79	11% (4.7-21%)	89% (79-95%)	0.078 (0.018-0.19)	-0.0044% (-0.34-0.2%)	-0.028% (-0.46-0.23%)	-0.16 (-2.6-1.3)	-0.49 (-35-21)
Jordan	1995	Male	80 to 84	9.8% (3.9-19%)	90% (81-96%)	0.065 (0.016-0.16)	0.00019% (-0.31-0.18%)	-0.021% (-0.37-0.2%)	-0.12 (-2.2-1.2)	0.008 (-22-14)
Jordan	1995	Male	85 to 89	9.5% (3.8-19%)	90% (81-96%)	0.059 (0.015-0.15)	-0.019% (-0.33-0.15%)	-0.052% (-0.43-0.16%)	-0.3 (-2.5-0.93)	-1 (-17-8)
Jordan	1995	Male	90 to 94	11% (4.4-22%)	89% (78-96%)	0.063 (0.015-0.17)	-0.043% (-0.36-0.13%)	-0.096% (-0.48-0.12%)	-0.23 (-1.2-0.3)	-0.64 (-5.3-2)
Jordan	1995	Male	95 plus	13% (5-29%)	87% (71-95%)	0.072 (0.014-0.2)	-0.035% (-0.4-0.17%)	-0.12% (-0.57-0.12%)	-0.055 (-0.27-0.058)	-0.066 (-0.76-0.32)
Jordan	2000	Female	All Ages	11% (6.3-17%)	89% (83-94%)	0.09 (0.065-0.12)	0.053% (-0.13-0.22%)	-0.39% (-0.96--0.0034%)	-34 (-85--0.25)	300 (-750-1200)
Jordan	2000	Female	15 to 19	9.5% (2.8-21%)	90% (79-97%)	0.071 (0.019-0.16)	0.4% (0.25-0.62%)	0.42% (0.22-0.83%)	0.61 (0.29-1.2)	140 (80-220)
Jordan	2000	Female	20 to 24	12% (3.6-26%)	88% (74-96%)	0.11 (0.037-0.24)	0.75% (0.49-1.1%)	0.55% (0.26-1%)	0.85 (0.4-1.6)	250 (160-380)
Jordan	2000	Female	25 to 29	14% (4-30%)	86% (70-96%)	0.14 (0.044-0.28)	0.75% (0.52-1%)	0.55% (0.25-1%)	0.91 (0.4-1.8)	250 (160-370)
Jordan	2000	Female	30 to 34	15% (4-33%)	85% (67-96%)	0.16 (0.051-0.32)	0.65% (0.42-0.94%)	0.61% (0.3-1.1%)	1.1 (0.49-2.1)	200 (120-300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	2000	Female	35 to 39	17% (4.5-34%)	83% (66-95%)	0.2 (0.061-0.39)	0.54% (0.31-0.82%)	0.71% (0.32-1.4%)	1.4 (0.59-2.8)	150 (87-230)
Jordan	2000	Female	40 to 44	18% (6.7-36%)	82% (64-93%)	0.22 (0.076-0.45)	0.42% (0.12-0.77%)	0.73% (0.24-1.5%)	1.5 (0.49-3.2)	97 (29-190)
Jordan	2000	Female	45 to 49	19% (6.5-35%)	81% (65-93%)	0.23 (0.073-0.53)	0.18% (-0.3-0.68%)	0.39% (-0.29-1.3%)	0.89 (-0.64-3.1)	37 (-62-140)
Jordan	2000	Female	50 to 54	19% (7-37%)	81% (63-93%)	0.24 (0.07-0.52)	-0.069% (-0.72-0.54%)	0.03% (-0.9-1%)	0.1 (-2.9-3.4)	-15 (-160-120)
Jordan	2000	Female	55 to 59	20% (7.4-38%)	80% (62-93%)	0.23 (0.065-0.52)	-0.35% (-1.3-0.33%)	-0.37% (-1.7-0.64%)	-1.8 (-7.9-3)	-91 (-330-86)
Jordan	2000	Female	60 to 64	21% (8.1-40%)	79% (60-92%)	0.22 (0.058-0.52)	-0.64% (-1.9-0.099%)	-0.76% (-2.3-0.23%)	-4.6 (-14-1.4)	-160 (-480-25)
Jordan	2000	Female	65 to 69	22% (9.4-38%)	78% (62-91%)	0.2 (0.055-0.42)	-0.87% (-2.1--0.14%)	-1.1% (-2.6--0.13%)	-7.7 (-18--0.93)	-200 (-480--31)
Jordan	2000	Female	70 to 74	20% (9-37%)	80% (63-91%)	0.16 (0.039-0.4)	-0.8% (-1.9--0.12%)	-0.96% (-2.3--0.13%)	-7.4 (-18--0.98)	-150 (-370--24)
Jordan	2000	Female	75 to 79	18% (7.1-36%)	82% (64-93%)	0.13 (0.027-0.36)	-0.73% (-2--0.11%)	-0.87% (-2.4--0.13%)	-6.6 (-18--0.93)	-100 (-280--15)
Jordan	2000	Female	80 to 84	17% (6-36%)	83% (64-94%)	0.12 (0.025-0.33)	-0.69% (-2--0.096%)	-0.82% (-2.4--0.12%)	-5.2 (-15--0.69)	-57 (-170--7.7)
Jordan	2000	Female	85 to 89	18% (6-39%)	82% (61-94%)	0.12 (0.023-0.36)	-0.7% (-2.1--0.084%)	-0.84% (-2.6--0.11%)	-4.2 (-13--0.54)	-32 (-98--3.7)
Jordan	2000	Female	90 to 94	20% (6.9-43%)	80% (57-93%)	0.14 (0.026-0.39)	-0.74% (-2.2--0.1%)	-0.91% (-2.6--0.13%)	-2.6 (-7.3--0.36)	-13 (-37--1.7)
Jordan	2000	Female	95 plus	25% (7.9-49%)	75% (51-92%)	0.16 (0.028-0.47)	-0.81% (-2.3--0.063%)	-1% (-2.8--0.11%)	-0.94 (-2.6--0.11)	-3.1 (-8.4--0.24)
Jordan	2000	Male	All Ages	8.6% (5.7-12%)	91% (88-94%)	0.1 (0.074-0.14)	0.43% (0.27-0.59%)	0.21% (-0.059-0.47%)	20 (-5.3-47)	2700 (1600-3900)
Jordan	2000	Male	15 to 19	6.7% (2.8-13%)	93% (87-97%)	0.048 (0.015-0.11)	0.6% (0.35-0.91%)	0.41% (0.14-0.9%)	0.97 (0.32-2.2)	230 (140-360)
Jordan	2000	Male	20 to 24	10% (4-20%)	90% (80-96%)	0.11 (0.038-0.2)	1.3% (0.82-1.8%)	0.86% (0.32-1.8%)	2.5 (0.87-5.3)	520 (320-790)
Jordan	2000	Male	25 to 29	13% (5.1-24%)	87% (76-95%)	0.17 (0.066-0.31)	1.4% (0.94-2%)	1.3% (0.6-2.2%)	3.9 (1.7-7.1)	570 (370-830)
Jordan	2000	Male	30 to 34	15% (5.5-29%)	85% (71-95%)	0.22 (0.092-0.39)	1.2% (0.85-1.8%)	1.2% (0.56-2.2%)	3.6 (1.6-6.5)	460 (290-680)
Jordan	2000	Male	35 to 39	16% (6.5-29%)	84% (71-93%)	0.25 (0.096-0.48)	1.1% (0.72-1.6%)	1.1% (0.47-2%)	3.2 (1.2-6.1)	340 (210-500)
Jordan	2000	Male	40 to 44	17% (7.3-30%)	83% (70-93%)	0.27 (0.1-0.52)	0.83% (0.44-1.3%)	0.79% (0.14-1.5%)	2.2 (0.39-4.4)	200 (110-330)
Jordan	2000	Male	45 to 49	16% (6.4-30%)	84% (70-94%)	0.28 (0.092-0.62)	0.57% (0.074-1.1%)	0.51% (-0.3-1.4%)	1.5 (-0.89-4.2)	120 (16-250)
Jordan	2000	Male	50 to 54	16% (6.7-31%)	84% (69-93%)	0.27 (0.086-0.59)	0.39% (-0.15-1%)	0.34% (-0.5-1.2%)	1.3 (-1.8-4.8)	90 (-33-230)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	2000	Male	55 to 59	15% (6.4-29%)	85% (71-94%)	0.26 (0.082-0.56)	0.26% (-0.27-0.82%)	0.23% (-0.53-1.1%)	1.2 (-2.8-5.9)	68 (-72-220)
Jordan	2000	Male	60 to 64	14% (6.1-28%)	86% (72-94%)	0.23 (0.064-0.53)	0.15% (-0.36-0.73%)	0.12% (-0.56-0.85%)	0.92 (-4.2-6.8)	43 (-100-220)
Jordan	2000	Male	65 to 69	14% (5.9-25%)	86% (75-94%)	0.19 (0.056-0.4)	0.043% (-0.39-0.43%)	0.011% (-0.62-0.53%)	0.1 (-4.7-4.2)	11 (-100-110)
Jordan	2000	Male	70 to 74	12% (5.4-22%)	88% (78-95%)	0.14 (0.041-0.33)	0.01% (-0.37-0.29%)	-0.013% (-0.49-0.34%)	-0.1 (-3.8-2.6)	1.9 (-68-53)
Jordan	2000	Male	75 to 79	11% (4.4-20%)	89% (80-96%)	0.11 (0.026-0.27)	0.00097% (-0.35-0.22%)	-0.021% (-0.46-0.26%)	-0.14 (-3.1-1.8)	0.069 (-45-28)
Jordan	2000	Male	80 to 84	9.5% (3.8-18%)	90% (82-96%)	0.09 (0.024-0.23)	-0.0026% (-0.31-0.19%)	-0.025% (-0.4-0.21%)	-0.13 (-2.1-1.1)	-0.19 (-21-13)
Jordan	2000	Male	85 to 89	9.4% (3.7-18%)	91% (82-96%)	0.083 (0.021-0.22)	-0.03% (-0.36-0.16%)	-0.068% (-0.46-0.16%)	-0.31 (-2.1-0.74)	-1.2 (-15-6.8)
Jordan	2000	Male	90 to 94	11% (4.1-21%)	89% (79-96%)	0.088 (0.021-0.24)	-0.056% (-0.38-0.14%)	-0.11% (-0.51-0.13%)	-0.36 (-1.6-0.41)	-1.1 (-7.2-2.8)
Jordan	2000	Male	95 plus	13% (4.7-27%)	87% (73-95%)	0.1 (0.02-0.29)	-0.047% (-0.44-0.18%)	-0.14% (-0.62-0.14%)	-0.11 (-0.5-0.11)	-0.15 (-1.4-0.61)
Jordan	2005	Female	All Ages	11% (6.5-17%)	89% (83-93%)	0.14 (0.1-0.18)	-0.11% (-0.13-0.35%)	-0.32% (-0.95-0.26%)	-32 (-97-25)	690 (-810-2200)
Jordan	2005	Female	15 to 19	9.3% (2.9-20%)	91% (80-97%)	0.1 (0.029-0.23)	0.42% (0.26-0.64%)	0.47% (0.23-0.93%)	0.66 (0.31-1.3)	150 (86-230)
Jordan	2005	Female	20 to 24	12% (3.5-26%)	88% (74-96%)	0.16 (0.052-0.35)	0.78% (0.51-1.1%)	0.66% (0.31-1.2%)	1.1 (0.51-2.2)	300 (180-460)
Jordan	2005	Female	25 to 29	14% (4-29%)	86% (71-96%)	0.19 (0.065-0.4)	0.81% (0.54-1.1%)	0.72% (0.33-1.4%)	1.2 (0.5-2.4)	290 (180-430)
Jordan	2005	Female	30 to 34	15% (4.3-33%)	85% (67-96%)	0.23 (0.079-0.46)	0.72% (0.47-1%)	0.83% (0.42-1.5%)	1.5 (0.68-2.9)	260 (160-390)
Jordan	2005	Female	35 to 39	17% (4.9-33%)	83% (67-95%)	0.28 (0.093-0.53)	0.65% (0.37-1%)	1% (0.45-1.9%)	2.3 (0.9-4.5)	220 (120-350)
Jordan	2005	Female	40 to 44	18% (6.7-34%)	82% (66-93%)	0.31 (0.11-0.6)	0.59% (0.23-1.1%)	1.2% (0.5-2.2%)	3 (1.2-6)	180 (71-330)
Jordan	2005	Female	45 to 49	18% (6.5-34%)	82% (66-94%)	0.33 (0.11-0.71)	0.41% (-0.12-1.1%)	0.9% (0.093-2.3%)	2.3 (0.23-6.3)	100 (-31-290)
Jordan	2005	Female	50 to 54	19% (7.3-36%)	81% (64-93%)	0.34 (0.11-0.74)	0.2% (-0.56-1.1%)	0.51% (-0.58-2%)	1.6 (-1.7-6.4)	45 (-130-240)
Jordan	2005	Female	55 to 59	20% (7.9-38%)	80% (62-92%)	0.34 (0.1-0.75)	-0.087% (-1.1-0.98%)	0.047% (-1.3-1.6%)	0.23 (-6.7-8.1)	-24 (-310-260)
Jordan	2005	Female	60 to 64	21% (8.4-42%)	79% (58-92%)	0.33 (0.087-0.78)	-0.45% (-1.8-0.65%)	-0.47% (-2.2-1%)	-3.8 (-17-8.3)	-140 (-600-220)
Jordan	2005	Female	65 to 69	21% (8.9-38%)	79% (62-91%)	0.29 (0.074-0.63)	-0.8% (-2.1-0.24%)	-0.95% (-2.6-0.34%)	-9.5 (-27-3.3)	-250 (-670-72)
Jordan	2005	Female	70 to 74	20% (8.2-36%)	80% (64-92%)	0.22 (0.054-0.56)	-0.77% (-2-0.025%)	-0.9% (-2.3-0.069%)	-10 (-26-0.78)	-200 (-510-6.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	2005	Female	75 to 79	17% (6.6-34%)	83% (66-93%)	0.18 (0.039-0.5)	-0.7% (-1.9--0.04%)	-0.83% (-2.2--0.041%)	-7.8 (-20--0.34)	-120 (-320--7)
Jordan	2005	Female	80 to 84	16% (5.5-35%)	84% (65-95%)	0.16 (0.035-0.46)	-0.67% (-1.9--0.065%)	-0.79% (-2.2--0.079%)	-5.8 (-16--0.53)	-64 (-180--5.8)
Jordan	2005	Female	85 to 89	17% (5.5-37%)	83% (63-94%)	0.16 (0.034-0.51)	-0.68% (-2--0.056%)	-0.81% (-2.3--0.076%)	-5 (-14--0.48)	-38 (-110--3.2)
Jordan	2005	Female	90 to 94	20% (6.6-43%)	80% (57-93%)	0.19 (0.037-0.55)	-0.72% (-2--0.06%)	-0.88% (-2.4--0.088%)	-2.9 (-8.3--0.29)	-15 (-41--1.2)
Jordan	2005	Female	95 plus	24% (7.3-49%)	76% (51-93%)	0.23 (0.039-0.67)	-0.8% (-2.2--0.00057%)	-1% (-2.7--0.031%)	-1.3 (-3.6--0.042)	-4.3 (-12--0.0028)
Jordan	2005	Male	All Ages	8.7% (5.8-12%)	91% (88-94%)	0.15 (0.11-0.19)	0.52% (0.32-0.75%)	0.33% (0.022-0.78%)	36 (2-85)	3500 (2100-5100)
Jordan	2005	Male	15 to 19	6.5% (2.8-13%)	93% (87-97%)	0.067 (0.022-0.15)	0.62% (0.38-0.94%)	0.43% (0.16-0.93%)	1 (0.35-2.3)	240 (140-370)
Jordan	2005	Male	20 to 24	10% (3.9-20%)	90% (80-96%)	0.15 (0.051-0.29)	1.3% (0.83-1.9%)	0.85% (0.34-1.7%)	2.8 (1-5.7)	590 (360-900)
Jordan	2005	Male	25 to 29	13% (4.8-24%)	87% (76-95%)	0.24 (0.095-0.44)	1.5% (1-2%)	1.4% (0.65-2.4%)	4 (1.8-7.5)	610 (400-890)
Jordan	2005	Male	30 to 34	15% (5.4-28%)	85% (72-95%)	0.3 (0.13-0.54)	1.4% (0.91-1.9%)	1.4% (0.69-2.5%)	3.9 (1.8-7)	530 (340-770)
Jordan	2005	Male	35 to 39	16% (6.3-29%)	84% (71-94%)	0.35 (0.14-0.65)	1.2% (0.82-1.8%)	1.4% (0.66-2.5%)	4.2 (1.9-7.8)	440 (280-660)
Jordan	2005	Male	40 to 44	16% (7.3-29%)	84% (71-93%)	0.37 (0.15-0.74)	1% (0.56-1.6%)	1.1% (0.38-2.2%)	3.6 (1.2-7.3)	320 (170-530)
Jordan	2005	Male	45 to 49	16% (6.4-29%)	84% (71-94%)	0.39 (0.14-0.81)	0.8% (0.24-1.6%)	0.86% (-0.0075-2.2%)	3.1 (-0.033-7.9)	210 (61-440)
Jordan	2005	Male	50 to 54	15% (6.5-29%)	85% (71-94%)	0.39 (0.13-0.86)	0.65% (0.013-1.7%)	0.73% (-0.23-2.2%)	2.9 (-0.94-9.3)	160 (2.9-410)
Jordan	2005	Male	55 to 59	14% (6.2-27%)	86% (73-94%)	0.37 (0.12-0.8)	0.52% (-0.088-1.6%)	0.59% (-0.26-2%)	3.2 (-1.4-11)	140 (-22-430)
Jordan	2005	Male	60 to 64	14% (5.8-27%)	86% (73-94%)	0.33 (0.096-0.74)	0.36% (-0.22-1.4%)	0.4% (-0.37-1.7%)	3.5 (-3.1-15)	120 (-68-480)
Jordan	2005	Male	65 to 69	13% (5.8-25%)	87% (75-94%)	0.27 (0.082-0.58)	0.21% (-0.29-0.97%)	0.23% (-0.4-1.2%)	2.4 (-4.3-13)	67 (-93-320)
Jordan	2005	Male	70 to 74	12% (5.4-22%)	88% (78-95%)	0.2 (0.057-0.48)	0.1% (-0.3-0.61%)	0.1% (-0.41-0.74%)	1 (-4-7.6)	25 (-71-150)
Jordan	2005	Male	75 to 79	10% (4.4-20%)	90% (80-96%)	0.15 (0.039-0.39)	0.049% (-0.33-0.43%)	0.039% (-0.43-0.49%)	0.34 (-3.9-4.5)	7.8 (-53-70)
Jordan	2005	Male	80 to 84	9.3% (3.7-19%)	91% (81-96%)	0.13 (0.033-0.33)	0.028% (-0.3-0.3%)	0.013% (-0.39-0.33%)	0.079 (-2.6-2.2)	2.3 (-26-25)
Jordan	2005	Male	85 to 89	9.1% (3.6-18%)	91% (82-96%)	0.12 (0.031-0.31)	-0.0074% (-0.32-0.22%)	-0.039% (-0.41-0.24%)	-0.19 (-2.1-1.2)	-0.34 (-14-9.6)
Jordan	2005	Male	90 to 94	10% (4.2-20%)	90% (80-96%)	0.12 (0.031-0.32)	-0.036% (-0.35-0.22%)	-0.089% (-0.47-0.21%)	-0.25 (-1.3-0.58)	-0.61 (-6.1-3.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	2005	Male	95 plus	13% (4.6-26%)	87% (74-95%)	0.14 (0.028-0.42)	-0.025% (-0.42-0.32%)	-0.11% (-0.59-0.27%)	-0.13 (-0.69-0.32)	-0.12 (-2-1.5)
Jordan	2010	Female	All Ages	11% (6.8-16%)	89% (84-93%)	0.14 (0.097-0.19)	0.17% (-0.029-0.4%)	-0.18% (-0.79-0.41%)	-17 (-75-39)	1200 (-180-2900)
Jordan	2010	Female	15 to 19	9% (3-19%)	91% (81-97%)	0.099 (0.028-0.23)	0.43% (0.27-0.64%)	0.51% (0.25-1%)	0.65 (0.3-1.4)	160 (94-250)
Jordan	2010	Female	20 to 24	12% (3.5-25%)	88% (75-96%)	0.16 (0.047-0.34)	0.82% (0.52-1.2%)	0.7% (0.33-1.4%)	0.92 (0.41-1.9)	320 (190-480)
Jordan	2010	Female	25 to 29	13% (3.8-27%)	87% (73-96%)	0.19 (0.056-0.39)	0.83% (0.57-1.1%)	0.77% (0.35-1.4%)	1.1 (0.48-2.2)	350 (220-510)
Jordan	2010	Female	30 to 34	15% (4-31%)	85% (69-96%)	0.23 (0.073-0.44)	0.74% (0.49-1.1%)	0.88% (0.41-1.6%)	1.5 (0.63-3)	300 (180-450)
Jordan	2010	Female	35 to 39	17% (4.8-32%)	83% (68-95%)	0.27 (0.088-0.52)	0.63% (0.38-0.97%)	1.1% (0.47-2%)	2.5 (1-4.7)	260 (150-410)
Jordan	2010	Female	40 to 44	18% (6.2-33%)	82% (67-94%)	0.3 (0.11-0.58)	0.55% (0.21-0.97%)	1.2% (0.56-2.3%)	3.1 (1.3-6)	200 (78-370)
Jordan	2010	Female	45 to 49	18% (6-34%)	82% (66-94%)	0.33 (0.11-0.65)	0.42% (-0.039-1%)	1.1% (0.31-2.4%)	3.2 (0.84-7.6)	140 (-13-350)
Jordan	2010	Female	50 to 54	19% (7.3-36%)	81% (64-93%)	0.34 (0.11-0.73)	0.23% (-0.42-1.1%)	0.68% (-0.24-2.2%)	2.5 (-0.91-8.3)	70 (-130-350)
Jordan	2010	Female	55 to 59	20% (7.8-38%)	80% (62-92%)	0.34 (0.1-0.78)	0.0028% (-0.87-0.96%)	0.26% (-0.91-1.8%)	1.2 (-4.1-8.4)	1.1 (-240-280)
Jordan	2010	Female	60 to 64	21% (8.5-40%)	79% (60-92%)	0.32 (0.084-0.78)	-0.27% (-1.4-0.75%)	-0.19% (-1.5-1.3%)	-1.3 (-10-8.6)	-81 (-410-240)
Jordan	2010	Female	65 to 69	21% (9.3-38%)	79% (62-91%)	0.28 (0.076-0.64)	-0.54% (-1.6-0.38%)	-0.61% (-2-0.63%)	-5.7 (-18-5.9)	-180 (-510-120)
Jordan	2010	Female	70 to 74	20% (8-36%)	80% (64-92%)	0.22 (0.054-0.57)	-0.57% (-1.6-0.17%)	-0.66% (-2-0.34%)	-7 (-22-3.2)	-160 (-470-49)
Jordan	2010	Female	75 to 79	17% (6.4-35%)	83% (65-94%)	0.18 (0.036-0.49)	-0.54% (-1.6-0.077%)	-0.65% (-1.9-0.13%)	-6.3 (-19-1.2)	-110 (-310-17)
Jordan	2010	Female	80 to 84	16% (5.5-36%)	84% (64-95%)	0.16 (0.033-0.47)	-0.53% (-1.7-0.0076%)	-0.64% (-2-0.033%)	-4.9 (-15-0.23)	-57 (-180-0.91)
Jordan	2010	Female	85 to 89	16% (5.5-37%)	84% (63-95%)	0.16 (0.034-0.5)	-0.56% (-1.7-0.0052%)	-0.68% (-2.1-0.015%)	-4.3 (-13-0.092)	-34 (-100-0.31)
Jordan	2010	Female	90 to 94	19% (6.5-41%)	81% (59-94%)	0.18 (0.037-0.55)	-0.6% (-1.7-0.025%)	-0.74% (-2.2-0.024%)	-3 (-9.1-0.11)	-15 (-46-0.65)
Jordan	2010	Female	95 plus	23% (7-48%)	77% (52-93%)	0.22 (0.038-0.65)	-0.64% (-1.9-0.1%)	-0.82% (-2.4-0.097%)	-1.3 (-3.7-0.15)	-4.1 (-12-0.71)
Jordan	2010	Male	All Ages	8.7% (5.9-12%)	91% (88-94%)	0.15 (0.11-0.21)	0.57% (0.37-0.82%)	0.38% (0.049-0.84%)	44 (5.4-100)	4200 (2600-6400)
Jordan	2010	Male	15 to 19	6.4% (2.7-13%)	94% (87-97%)	0.066 (0.019-0.15)	0.66% (0.4-1%)	0.5% (0.17-1.1%)	1.1 (0.35-2.5)	270 (160-420)
Jordan	2010	Male	20 to 24	9.9% (3.8-19%)	90% (81-96%)	0.15 (0.049-0.31)	1.4% (0.93-2%)	1.1% (0.41-2.3%)	3 (1.1-6.3)	640 (390-960)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	2010	Male	25 to 29	12% (4.8-23%)	88% (77-95%)	0.23 (0.087-0.45)	1.6% (1.1-2.2%)	1.6% (0.74-2.8%)	4.5 (1.9-8.3)	730 (470-1100)
Jordan	2010	Male	30 to 34	14% (5.2-27%)	86% (73-95%)	0.29 (0.12-0.52)	1.4% (0.97-2%)	1.5% (0.73-2.8%)	4.1 (1.9-7.6)	600 (390-900)
Jordan	2010	Male	35 to 39	16% (6.1-29%)	84% (71-94%)	0.33 (0.13-0.62)	1.3% (0.86-1.8%)	1.5% (0.67-2.6%)	4.8 (2.1-9)	550 (360-810)
Jordan	2010	Male	40 to 44	16% (7.1-28%)	84% (72-93%)	0.36 (0.15-0.69)	1% (0.6-1.6%)	1.2% (0.4-2.2%)	4.5 (1.4-9.3)	430 (240-710)
Jordan	2010	Male	45 to 49	16% (6.1-29%)	84% (71-94%)	0.38 (0.13-0.77)	0.82% (0.27-1.6%)	0.92% (0.039-2.2%)	4.3 (0.23-11)	310 (110-650)
Jordan	2010	Male	50 to 54	15% (6.2-30%)	85% (70-94%)	0.38 (0.12-0.81)	0.66% (0.03-1.6%)	0.76% (-0.16-2.2%)	4.2 (-0.9-13)	230 (11-590)
Jordan	2010	Male	55 to 59	14% (6-27%)	86% (73-94%)	0.36 (0.11-0.8)	0.54% (-0.05-1.5%)	0.62% (-0.21-2.1%)	3.7 (-1.2-13)	160 (-16-490)
Jordan	2010	Male	60 to 64	14% (5.7-26%)	86% (74-94%)	0.32 (0.092-0.76)	0.41% (-0.13-1.4%)	0.47% (-0.25-1.8%)	3.8 (-1.9-15)	130 (-39-480)
Jordan	2010	Male	65 to 69	13% (5.9-24%)	87% (76-94%)	0.26 (0.078-0.57)	0.24% (-0.24-1%)	0.27% (-0.38-1.3%)	3.1 (-3.8-15)	85 (-83-390)
Jordan	2010	Male	70 to 74	12% (5.3-21%)	88% (79-95%)	0.19 (0.054-0.45)	0.13% (-0.25-0.69%)	0.14% (-0.35-0.86%)	1.7 (-4.3-10)	40 (-78-220)
Jordan	2010	Male	75 to 79	10% (4.2-19%)	90% (81-96%)	0.14 (0.036-0.37)	0.08% (-0.26-0.48%)	0.08% (-0.32-0.59%)	0.85 (-3.5-6.2)	16 (-49-98)
Jordan	2010	Male	80 to 84	9.1% (3.5-18%)	91% (82-96%)	0.12 (0.029-0.32)	0.049% (-0.24-0.35%)	0.038% (-0.35-0.4%)	0.29 (-2.6-3.1)	5 (-26-35)
Jordan	2010	Male	85 to 89	9.1% (3.6-19%)	91% (81-96%)	0.12 (0.029-0.3)	0.015% (-0.29-0.25%)	-0.013% (-0.38-0.27%)	-0.073 (-2.2-1.6)	0.79 (-15-13)
Jordan	2010	Male	90 to 94	10% (4-21%)	90% (79-96%)	0.12 (0.03-0.35)	-0.013% (-0.33-0.28%)	-0.064% (-0.45-0.27%)	-0.19 (-1.4-0.81)	-0.24 (-6.2-5.1)
Jordan	2010	Male	95 plus	13% (4.6-27%)	87% (73-95%)	0.14 (0.027-0.43)	0.0056% (-0.39-0.38%)	-0.082% (-0.55-0.35%)	-0.083 (-0.54-0.35)	0.024 (-1.5-1.6)
Jordan	2016	Female	All Ages	12% (6.8-17%)	88% (83-93%)	0.15 (0.085-0.25)	0.23% (0.0088-0.51%)	-0.08% (-0.69-0.63%)	-8.7 (-82-69)	1800 (69-4300)
Jordan	2016	Female	15 to 19	9.3% (3-20%)	91% (80-97%)	0.1 (0.026-0.26)	0.41% (0.25-0.64%)	0.41% (0.18-0.85%)	0.74 (0.31-1.6)	190 (110-310)
Jordan	2016	Female	20 to 24	12% (3.3-26%)	88% (74-97%)	0.16 (0.045-0.39)	0.8% (0.5-1.2%)	0.6% (0.25-1.3%)	1.1 (0.44-2.5)	370 (220-570)
Jordan	2016	Female	25 to 29	13% (3.7-29%)	87% (71-96%)	0.2 (0.05-0.47)	0.82% (0.55-1.1%)	0.71% (0.3-1.4%)	1.3 (0.51-2.7)	390 (240-570)
Jordan	2016	Female	30 to 34	15% (3.8-31%)	85% (69-96%)	0.23 (0.061-0.53)	0.75% (0.49-1.1%)	0.84% (0.36-1.7%)	1.9 (0.75-4.1)	380 (230-590)
Jordan	2016	Female	35 to 39	17% (4.3-32%)	83% (68-96%)	0.27 (0.077-0.61)	0.65% (0.38-1%)	1.1% (0.43-2.2%)	2.9 (1-6)	320 (180-530)
Jordan	2016	Female	40 to 44	18% (6.3-34%)	82% (66-94%)	0.3 (0.1-0.64)	0.58% (0.23-1.1%)	1.3% (0.54-2.6%)	4.2 (1.6-9.3)	290 (110-550)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	2016	Female	45 to 49	18% (6.6-35%)	82% (65-93%)	0.33 (0.11-0.74)	0.48% (0.0077-1.2%)	1.1% (0.32-2.6%)	4.7 (1.2-12)	220 (4.1-560)
Jordan	2016	Female	50 to 54	19% (7.5-38%)	81% (62-93%)	0.35 (0.1-0.78)	0.32% (-0.33-1.2%)	0.81% (-0.13-2.5%)	4.2 (-0.69-13)	140 (-140-550)
Jordan	2016	Female	55 to 59	20% (8-40%)	80% (60-92%)	0.35 (0.1-0.85)	0.12% (-0.66-1.2%)	0.47% (-0.61-2.3%)	2.8 (-3.7-14)	48 (-260-470)
Jordan	2016	Female	60 to 64	22% (8.7-41%)	78% (59-91%)	0.34 (0.085-0.86)	-0.14% (-1.1-1.1%)	-0.0037% (-1.3-1.8%)	-0.0026 (-8.7-12)	-44 (-340-340)
Jordan	2016	Female	65 to 69	22% (9.7-40%)	78% (60-90%)	0.3 (0.08-0.78)	-0.4% (-1.4-0.62%)	-0.42% (-1.7-1.1%)	-3.5 (-15-8.9)	-130 (-430-200)
Jordan	2016	Female	70 to 74	20% (8.7-38%)	80% (62-91%)	0.24 (0.052-0.65)	-0.45% (-1.4-0.37%)	-0.52% (-1.8-0.6%)	-6 (-21-6.6)	-140 (-460-110)
Jordan	2016	Female	75 to 79	18% (6.7-37%)	82% (63-93%)	0.19 (0.034-0.56)	-0.47% (-1.4-0.22%)	-0.56% (-1.8-0.31%)	-7 (-23-3.8)	-120 (-370-54)
Jordan	2016	Female	80 to 84	17% (5.7-37%)	83% (63-94%)	0.17 (0.033-0.5)	-0.46% (-1.5-0.15%)	-0.56% (-1.8-0.19%)	-5.9 (-20-2)	-69 (-230-22)
Jordan	2016	Female	85 to 89	17% (5.6-38%)	83% (62-94%)	0.17 (0.033-0.53)	-0.5% (-1.6-0.096%)	-0.62% (-2-0.12%)	-5.4 (-17-0.99)	-42 (-130-8.3)
Jordan	2016	Female	90 to 94	20% (6.8-42%)	80% (58-93%)	0.2 (0.035-0.59)	-0.54% (-1.6-0.11%)	-0.68% (-2-0.13%)	-3.2 (-9.9-0.62)	-17 (-50-3.5)
Jordan	2016	Female	95 plus	24% (7.4-50%)	76% (50-93%)	0.23 (0.037-0.73)	-0.58% (-1.8-0.29%)	-0.76% (-2.3-0.28%)	-1.5 (-4.6-0.49)	-4.9 (-15-2.2)
Jordan	2016	Male	All Ages	9% (6-13%)	91% (87-94%)	0.16 (0.092-0.26)	0.6% (0.38-0.93%)	0.41% (0.054-1%)	60 (7.2-160)	5400 (3200-8500)
Jordan	2016	Male	15 to 19	6.5% (2.7-13%)	94% (87-97%)	0.067 (0.017-0.16)	0.58% (0.35-0.86%)	0.37% (0.11-0.86%)	1.4 (0.4-3.4)	320 (190-510)
Jordan	2016	Male	20 to 24	10% (3.9-19%)	90% (81-96%)	0.15 (0.045-0.35)	1.2% (0.74-1.8%)	0.71% (0.24-1.5%)	3.7 (1.2-8.4)	750 (440-1200)
Jordan	2016	Male	25 to 29	12% (5.1-24%)	88% (76-95%)	0.23 (0.076-0.51)	1.4% (0.93-2%)	1.2% (0.47-2.2%)	5.6 (2.2-11)	830 (520-1300)
Jordan	2016	Male	30 to 34	14% (5.2-27%)	86% (73-95%)	0.29 (0.11-0.61)	1.3% (0.85-1.9%)	1.2% (0.53-2.4%)	6 (2.4-13)	780 (490-1200)
Jordan	2016	Male	35 to 39	16% (5.9-29%)	84% (71-94%)	0.33 (0.12-0.69)	1.2% (0.8-1.9%)	1.3% (0.54-2.6%)	6.4 (2.4-14)	690 (420-1100)
Jordan	2016	Male	40 to 44	16% (7.2-28%)	84% (72-93%)	0.35 (0.13-0.73)	1% (0.56-1.8%)	1.1% (0.33-2.3%)	6.2 (1.7-15)	580 (300-1000)
Jordan	2016	Male	45 to 49	16% (6-28%)	84% (72-94%)	0.37 (0.12-0.77)	0.83% (0.32-1.7%)	0.91% (0.066-2.2%)	6.4 (0.47-18)	470 (170-1000)
Jordan	2016	Male	50 to 54	15% (6.5-29%)	85% (71-94%)	0.37 (0.12-0.85)	0.69% (0.1-1.7%)	0.79% (-0.1-2.5%)	6.4 (-0.86-20)	360 (52-960)
Jordan	2016	Male	55 to 59	14% (6.3-27%)	86% (73-94%)	0.35 (0.11-0.83)	0.56% (-0.0044-1.7%)	0.65% (-0.17-2.3%)	5.3 (-1.3-20)	230 (-2-760)
Jordan	2016	Male	60 to 64	14% (5.8-26%)	86% (74-94%)	0.32 (0.089-0.79)	0.44% (-0.092-1.6%)	0.52% (-0.24-2%)	4.5 (-2-19)	160 (-33-600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Jordan	2016	Male	65 to 69	13% (5.9-25%)	87% (75-94%)	0.26 (0.071-0.63)	0.28% (-0.19-1.2%)	0.33% (-0.29-1.5%)	3.4 (-2.9-16)	95 (-60-390)
Jordan	2016	Male	70 to 74	12% (5.2-21%)	88% (79-95%)	0.19 (0.05-0.49)	0.16% (-0.22-0.76%)	0.18% (-0.34-0.93%)	2.3 (-4.1-13)	53 (-72-250)
Jordan	2016	Male	75 to 79	10% (4.3-20%)	90% (80-96%)	0.15 (0.032-0.39)	0.11% (-0.23-0.57%)	0.11% (-0.32-0.71%)	1.6 (-4.5-9.7)	29 (-60-150)
Jordan	2016	Male	80 to 84	9.2% (3.5-19%)	91% (81-97%)	0.12 (0.025-0.32)	0.073% (-0.22-0.41%)	0.066% (-0.3-0.48%)	0.73 (-3.3-5.3)	11 (-32-62)
Jordan	2016	Male	85 to 89	8.9% (3.5-18%)	91% (82-96%)	0.11 (0.027-0.31)	0.04% (-0.23-0.33%)	0.016% (-0.32-0.35%)	0.13 (-2.5-3)	3 (-17-26)
Jordan	2016	Male	90 to 94	10% (4.2-20%)	90% (80-96%)	0.12 (0.028-0.32)	0.017% (-0.28-0.31%)	-0.029% (-0.39-0.33%)	-0.11 (-1.5-1.3)	0.43 (-6.7-7.8)
Jordan	2016	Male	95 plus	13% (4.6-26%)	87% (74-95%)	0.14 (0.025-0.43)	0.036% (-0.34-0.46%)	-0.048% (-0.51-0.43%)	-0.057 (-0.62-0.55)	0.18 (-1.7-2.3)
Kuwait	1990	Female	All Ages	10% (6-16%)	90% (84-94%)	0.0076 (0.0013-0.017)	0.28% (0.2-0.37%)	0.073% (-0.048-0.12%)	1.3 (-0.87-2.3)	480 (320-660)
Kuwait	1990	Female	15 to 19	8.8% (2.9-19%)	91% (81-97%)	0.0057 (0.00074-0.019)	0.3% (0.16-0.46%)	0.08% (0.047-0.14%)	0.044 (0.03-0.067)	41 (21-68)
Kuwait	1990	Female	20 to 24	11% (3.4-24%)	89% (76-97%)	0.009 (0.0012-0.026)	0.65% (0.4-0.99%)	0.091% (0.054-0.16%)	0.051 (0.034-0.079)	93 (53-150)
Kuwait	1990	Female	25 to 29	13% (3.8-27%)	87% (73-96%)	0.011 (0.0016-0.03)	0.7% (0.47-0.98%)	0.17% (0.11-0.26%)	0.091 (0.059-0.14)	96 (58-140)
Kuwait	1990	Female	30 to 34	15% (4-30%)	85% (70-96%)	0.013 (0.002-0.036)	0.62% (0.41-0.88%)	0.16% (0.11-0.25%)	0.083 (0.056-0.12)	80 (48-120)
Kuwait	1990	Female	35 to 39	16% (4.2-32%)	84% (68-96%)	0.015 (0.0024-0.04)	0.49% (0.33-0.69%)	0.2% (0.14-0.29%)	0.13 (0.081-0.18)	64 (39-94)
Kuwait	1990	Female	40 to 44	17% (6.2-32%)	83% (68-94%)	0.017 (0.0025-0.045)	0.37% (0.22-0.55%)	0.22% (0.14-0.32%)	0.14 (0.091-0.21)	41 (23-66)
Kuwait	1990	Female	45 to 49	17% (6.1-33%)	83% (67-94%)	0.018 (0.0023-0.05)	0.28% (0.14-0.41%)	0.19% (0.067-0.28%)	0.14 (0.049-0.22)	26 (13-40)
Kuwait	1990	Female	50 to 54	18% (7.3-35%)	82% (65-93%)	0.019 (0.0023-0.054)	0.2% (0.036-0.33%)	0.16% (-0.063-0.28%)	0.13 (-0.045-0.24)	15 (2.4-25)
Kuwait	1990	Female	55 to 59	19% (7.6-38%)	81% (62-92%)	0.018 (0.0018-0.057)	0.13% (-0.086-0.25%)	0.11% (-0.18-0.25%)	0.086 (-0.14-0.21)	6.4 (-4.3-12)
Kuwait	1990	Female	60 to 64	21% (7.8-41%)	79% (59-92%)	0.018 (0.0017-0.052)	0.082% (-0.19-0.21%)	0.073% (-0.28-0.23%)	0.073 (-0.27-0.24)	3.5 (-8.1-9.3)
Kuwait	1990	Female	65 to 69	21% (8.3-39%)	79% (61-92%)	0.016 (0.0014-0.047)	0.071% (-0.2-0.18%)	0.074% (-0.26-0.22%)	0.095 (-0.35-0.3)	3.1 (-8.7-8.3)
Kuwait	1990	Female	70 to 74	20% (8-36%)	80% (64-92%)	0.013 (0.0014-0.04)	0.066% (-0.2-0.16%)	0.072% (-0.27-0.19%)	0.072 (-0.26-0.2)	1.7 (-4.8-4.2)
Kuwait	1990	Female	75 to 79	17% (6.4-36%)	83% (64-94%)	0.01 (0.00098-0.034)	0.057% (-0.21-0.13%)	0.061% (-0.26-0.16%)	0.06 (-0.24-0.16)	1 (-3.6-2.6)
Kuwait	1990	Female	80 to 84	16% (5.1-36%)	84% (64-95%)	0.0092 (0.00077-0.032)	0.036% (-0.22-0.1%)	0.034% (-0.26-0.11%)	0.044 (-0.33-0.15)	0.6 (-3.5-1.7)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	1990	Female	85 to 89	16% (5.1-38%)	84% (62-95%)	0.0092 (0.00081-0.033)	0.02% (-0.2-0.072%)	0.013% (-0.25-0.076%)	0.011 (-0.21-0.068)	0.17 (-1.6-0.63)
Kuwait	1990	Female	90 to 94	19% (6.1-42%)	81% (58-94%)	0.01 (0.00078-0.036)	0.02% (-0.2-0.072%)	0.0085% (-0.27-0.073%)	0.0039 (-0.12-0.033)	0.059 (-0.58-0.22)
Kuwait	1990	Female	95 plus	23% (7.3-49%)	77% (51-93%)	0.013 (0.00094-0.045)	0.011% (-0.26-0.074%)	-0.015% (-0.37-0.057%)	-0.0026 (-0.064-0.01)	0.0087 (-0.21-0.06)
Kuwait	1990	Male	All Ages	9.4% (6.2-14%)	91% (86-94%)	0.01 (0.0018-0.023)	0.45% (0.34-0.57%)	0.25% (0.16-0.34%)	7.6 (4.8-10)	1100 (770-1400)
Kuwait	1990	Male	15 to 19	6.4% (2.6-13%)	94% (87-97%)	0.004 (0.00051-0.012)	0.36% (0.2-0.57%)	0.049% (0.023-0.12%)	0.066 (0.035-0.15)	62 (35-100)
Kuwait	1990	Male	20 to 24	9.8% (3.8-19%)	90% (81-96%)	0.0088 (0.0012-0.026)	0.72% (0.41-1.2%)	0.14% (0.067-0.3%)	0.22 (0.13-0.39)	120 (72-180)
Kuwait	1990	Male	25 to 29	12% (4.8-24%)	88% (76-95%)	0.014 (0.0022-0.038)	0.8% (0.49-1.2%)	0.17% (0.082-0.34%)	0.26 (0.15-0.45)	130 (86-190)
Kuwait	1990	Male	30 to 34	14% (5.3-27%)	86% (73-95%)	0.017 (0.0029-0.045)	0.8% (0.52-1.1%)	0.28% (0.16-0.51%)	0.55 (0.35-0.83)	180 (120-260)
Kuwait	1990	Male	35 to 39	16% (6.2-28%)	84% (72-94%)	0.02 (0.0032-0.05)	0.81% (0.57-1.1%)	0.45% (0.28-0.7%)	0.77 (0.51-1.1)	160 (110-220)
Kuwait	1990	Male	40 to 44	16% (6.8-28%)	84% (72-93%)	0.021 (0.0032-0.055)	0.75% (0.54-1%)	0.54% (0.35-0.75%)	0.86 (0.57-1.2)	130 (85-180)
Kuwait	1990	Male	45 to 49	15% (6-29%)	85% (71-94%)	0.022 (0.0028-0.063)	0.64% (0.48-0.81%)	0.59% (0.35-0.8%)	1.3 (0.75-1.8)	120 (84-160)
Kuwait	1990	Male	50 to 54	15% (6.3-29%)	85% (71-94%)	0.022 (0.003-0.064)	0.47% (0.29-0.62%)	0.45% (0.19-0.64%)	0.92 (0.4-1.4)	66 (41-90)
Kuwait	1990	Male	55 to 59	14% (6-27%)	86% (73-94%)	0.021 (0.0023-0.06)	0.35% (0.19-0.48%)	0.34% (0.073-0.54%)	0.73 (0.17-1.2)	40 (21-57)
Kuwait	1990	Male	60 to 64	13% (5.5-26%)	87% (74-94%)	0.019 (0.0021-0.053)	0.3% (0.12-0.44%)	0.32% (0.08-0.52%)	0.77 (0.19-1.3)	29 (12-43)
Kuwait	1990	Male	65 to 69	13% (5.8-24%)	87% (76-94%)	0.015 (0.0017-0.044)	0.24% (0.096-0.35%)	0.28% (0.074-0.45%)	0.41 (0.11-0.67)	12 (4.7-18)
Kuwait	1990	Male	70 to 74	12% (5.2-21%)	88% (79-95%)	0.012 (0.0015-0.035)	0.18% (0.054-0.28%)	0.21% (0.032-0.33%)	0.29 (0.047-0.49)	6.2 (1.9-9.7)
Kuwait	1990	Male	75 to 79	10% (4.3-19%)	90% (81-96%)	0.0088 (0.001-0.028)	0.17% (0.071-0.24%)	0.19% (0.061-0.29%)	0.17 (0.055-0.25)	2.7 (1.1-3.9)
Kuwait	1990	Male	80 to 84	9.1% (3.6-18%)	91% (82-96%)	0.0073 (0.00089-0.022)	0.13% (0.043-0.2%)	0.15% (0.036-0.22%)	0.16 (0.039-0.25)	1.9 (0.59-2.8)
Kuwait	1990	Male	85 to 89	8.8% (3.6-17%)	91% (83-96%)	0.0066 (0.0008-0.02)	0.12% (0.053-0.17%)	0.13% (0.04-0.19%)	0.12 (0.037-0.18)	1 (0.46-1.5)
Kuwait	1990	Male	90 to 94	9.9% (4-21%)	90% (79-96%)	0.007 (0.00076-0.022)	0.086% (0.017-0.12%)	0.079% (-0.0054-0.12%)	0.038 (-0.0026-0.058)	0.27 (0.055-0.39)
Kuwait	1990	Male	95 plus	12% (4.5-25%)	88% (75-96%)	0.008 (0.00071-0.028)	0.12% (0.039-0.17%)	0.086% (-0.025-0.13%)	0.012 (-0.0034-0.018)	0.07 (0.022-0.099)
Kuwait	1995	Female	All Ages	11% (6.1-17%)	89% (83-94%)	0.0074 (0.0028-0.014)	0.29% (0.21-0.38%)	0.079% (-0.056-0.13%)	1.1 (-0.74-1.9)	360 (240-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	1995	Female	15 to 19	8.9% (3-20%)	91% (80-97%)	0.0054 (0.001-0.016)	0.34% (0.18-0.51%)	0.12% (0.086-0.17%)	0.019 (0.013-0.028)	20 (10-32)
Kuwait	1995	Female	20 to 24	12% (3.6-25%)	88% (75-96%)	0.0085 (0.0018-0.023)	0.7% (0.42-1.1%)	0.12% (0.09-0.18%)	0.028 (0.02-0.042)	55 (31-89)
Kuwait	1995	Female	25 to 29	13% (3.8-28%)	87% (72-96%)	0.01 (0.0023-0.025)	0.74% (0.51-1%)	0.19% (0.14-0.28%)	0.07 (0.048-0.1)	85 (53-130)
Kuwait	1995	Female	30 to 34	15% (4.1-31%)	85% (69-96%)	0.012 (0.0031-0.027)	0.63% (0.41-0.92%)	0.18% (0.13-0.25%)	0.069 (0.046-0.098)	71 (44-110)
Kuwait	1995	Female	35 to 39	17% (4.5-32%)	83% (68-95%)	0.014 (0.0041-0.032)	0.49% (0.33-0.71%)	0.21% (0.15-0.29%)	0.096 (0.064-0.14)	51 (30-76)
Kuwait	1995	Female	40 to 44	17% (6.7-33%)	83% (67-93%)	0.016 (0.0046-0.037)	0.37% (0.24-0.55%)	0.22% (0.14-0.31%)	0.11 (0.063-0.15)	32 (19-50)
Kuwait	1995	Female	45 to 49	18% (6.1-34%)	82% (66-94%)	0.018 (0.0039-0.044)	0.28% (0.15-0.41%)	0.19% (0.028-0.28%)	0.085 (0.013-0.13)	16 (8.2-24)
Kuwait	1995	Female	50 to 54	19% (6.8-37%)	81% (63-93%)	0.018 (0.0036-0.049)	0.21% (0.031-0.33%)	0.16% (-0.066-0.3%)	0.096 (-0.037-0.18)	11 (1.5-18)
Kuwait	1995	Female	55 to 59	20% (7.7-38%)	80% (62-92%)	0.018 (0.0033-0.048)	0.13% (-0.082-0.24%)	0.098% (-0.18-0.24%)	0.06 (-0.11-0.16)	4.7 (-3-9.3)
Kuwait	1995	Female	60 to 64	21% (8.1-39%)	79% (61-92%)	0.017 (0.0028-0.045)	0.081% (-0.19-0.2%)	0.068% (-0.27-0.22%)	0.079 (-0.32-0.26)	4 (-9.8-10)
Kuwait	1995	Female	65 to 69	21% (9-38%)	79% (62-91%)	0.015 (0.0027-0.04)	0.08% (-0.16-0.19%)	0.084% (-0.23-0.22%)	0.06 (-0.17-0.16)	1.9 (-3.8-4.4)
Kuwait	1995	Female	70 to 74	20% (8.2-37%)	80% (63-92%)	0.012 (0.002-0.034)	0.082% (-0.17-0.16%)	0.091% (-0.22-0.19%)	0.15 (-0.35-0.32)	3.3 (-6.9-6.7)
Kuwait	1995	Female	75 to 79	18% (7-36%)	82% (64-93%)	0.0095 (0.0015-0.03)	0.072% (-0.15-0.14%)	0.08% (-0.2-0.17%)	0.077 (-0.19-0.16)	1.3 (-2.7-2.6)
Kuwait	1995	Female	80 to 84	17% (5.7-37%)	83% (63-94%)	0.0085 (0.0012-0.027)	0.048% (-0.17-0.1%)	0.048% (-0.2-0.11%)	0.037 (-0.16-0.088)	0.47 (-1.7-1)
Kuwait	1995	Female	85 to 89	17% (5.6-38%)	83% (62-94%)	0.0086 (0.0012-0.026)	0.033% (-0.14-0.071%)	0.029% (-0.18-0.076%)	0.026 (-0.16-0.072)	0.29 (-1.2-0.65)
Kuwait	1995	Female	90 to 94	20% (6.8-43%)	80% (57-93%)	0.01 (0.0012-0.032)	0.031% (-0.16-0.073%)	0.022% (-0.21-0.074%)	0.014 (-0.12-0.046)	0.12 (-0.62-0.3)
Kuwait	1995	Female	95 plus	24% (7.1-49%)	76% (51-93%)	0.012 (0.0014-0.04)	0.021% (-0.22-0.076%)	-0.00079% (-0.3-0.06%)	-0.00014 (-0.066-0.013)	0.02 (-0.21-0.072)
Kuwait	1995	Male	All Ages	10% (6.4-15%)	90% (85-94%)	0.01 (0.0041-0.019)	0.58% (0.47-0.72%)	0.32% (0.22-0.39%)	7.4 (5.1-9.6)	1100 (800-1400)
Kuwait	1995	Male	15 to 19	6.4% (2.6-13%)	94% (87-97%)	0.0037 (0.00071-0.011)	0.46% (0.27-0.69%)	0.073% (0.044-0.17%)	0.031 (0.018-0.076)	33 (19-53)
Kuwait	1995	Male	20 to 24	9.8% (3.9-19%)	90% (81-96%)	0.008 (0.0018-0.022)	1.1% (0.72-1.6%)	0.3% (0.2-0.5%)	0.22 (0.14-0.36)	120 (75-190)
Kuwait	1995	Male	25 to 29	12% (4.8-23%)	88% (77-95%)	0.012 (0.0032-0.028)	1.1% (0.79-1.6%)	0.32% (0.21-0.49%)	0.37 (0.24-0.58)	210 (140-320)
Kuwait	1995	Male	30 to 34	14% (5.3-28%)	86% (72-95%)	0.015 (0.0048-0.033)	1% (0.72-1.4%)	0.47% (0.34-0.65%)	0.63 (0.44-0.89)	210 (140-310)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	1995	Male	35 to 39	16% (5.9-29%)	84% (71-94%)	0.017 (0.0053-0.038)	0.94% (0.68-1.2%)	0.66% (0.49-0.84%)	0.84 (0.59-1.2)	170 (110-230)
Kuwait	1995	Male	40 to 44	16% (7-28%)	84% (72-93%)	0.019 (0.0055-0.043)	0.79% (0.59-1%)	0.63% (0.43-0.81%)	0.96 (0.63-1.3)	140 (95-190)
Kuwait	1995	Male	45 to 49	16% (6.2-29%)	84% (71-94%)	0.02 (0.0044-0.049)	0.66% (0.5-0.83%)	0.64% (0.38-0.83%)	0.88 (0.5-1.2)	76 (54-100)
Kuwait	1995	Male	50 to 54	15% (6.4-29%)	85% (71-94%)	0.02 (0.0041-0.052)	0.48% (0.32-0.64%)	0.48% (0.23-0.68%)	0.77 (0.36-1.1)	53 (33-72)
Kuwait	1995	Male	55 to 59	14% (6.3-27%)	86% (73-94%)	0.019 (0.0036-0.048)	0.35% (0.18-0.5%)	0.35% (0.098-0.55%)	0.51 (0.13-0.84)	26 (14-38)
Kuwait	1995	Male	60 to 64	14% (6.1-26%)	86% (74-94%)	0.017 (0.003-0.044)	0.31% (0.14-0.45%)	0.33% (0.1-0.54%)	0.59 (0.19-0.98)	21 (9.2-32)
Kuwait	1995	Male	65 to 69	13% (5.8-24%)	87% (76-94%)	0.014 (0.0027-0.036)	0.26% (0.13-0.38%)	0.31% (0.12-0.49%)	0.54 (0.2-0.87)	15 (7.5-23)
Kuwait	1995	Male	70 to 74	12% (5.3-21%)	88% (79-95%)	0.01 (0.0021-0.028)	0.2% (0.094-0.31%)	0.24% (0.09-0.38%)	0.38 (0.14-0.62)	8 (3.7-12)
Kuwait	1995	Male	75 to 79	10% (4.3-20%)	90% (80-96%)	0.0078 (0.0014-0.023)	0.19% (0.1-0.27%)	0.22% (0.1-0.33%)	0.29 (0.13-0.44)	4.5 (2.4-6.7)
Kuwait	1995	Male	80 to 84	9% (3.5-18%)	91% (82-97%)	0.0064 (0.0011-0.018)	0.15% (0.07-0.22%)	0.17% (0.068-0.26%)	0.15 (0.059-0.23)	1.7 (0.78-2.5)
Kuwait	1995	Male	85 to 89	8.8% (3.4-18%)	91% (82-97%)	0.0059 (0.0011-0.016)	0.14% (0.072-0.19%)	0.15% (0.071-0.22%)	0.12 (0.056-0.17)	1 (0.51-1.4)
Kuwait	1995	Male	90 to 94	10% (4-21%)	90% (79-96%)	0.0062 (0.001-0.019)	0.095% (0.034-0.13%)	0.091% (0.014-0.13%)	0.049 (0.0072-0.071)	0.33 (0.11-0.45)
Kuwait	1995	Male	95 plus	12% (4.6-26%)	88% (74-95%)	0.0071 (0.001-0.022)	0.13% (0.064-0.17%)	0.1% (0.012-0.14%)	0.019 (0.0021-0.027)	0.11 (0.051-0.15)
Kuwait	2000	Female	All Ages	11% (6.2-17%)	89% (83-94%)	0.0061 (0.0033-0.0093)	0.33% (0.25-0.42%)	0.15% (0.062-0.19%)	2.4 (0.97-3.2)	460 (320-630)
Kuwait	2000	Female	15 to 19	8.9% (2.8-19%)	91% (81-97%)	0.0042 (0.001-0.011)	0.34% (0.19-0.53%)	0.087% (0.067-0.12%)	0.018 (0.013-0.024)	27 (14-44)
Kuwait	2000	Female	20 to 24	11% (3.5-24%)	89% (76-97%)	0.0066 (0.0016-0.016)	0.71% (0.42-1%)	0.11% (0.083-0.15%)	0.028 (0.02-0.04)	65 (36-100)
Kuwait	2000	Female	25 to 29	13% (3.6-27%)	87% (73-96%)	0.0079 (0.0021-0.017)	0.75% (0.49-1.1%)	0.18% (0.14-0.24%)	0.067 (0.048-0.092)	92 (57-140)
Kuwait	2000	Female	30 to 34	15% (3.7-31%)	85% (69-96%)	0.0092 (0.0028-0.019)	0.65% (0.43-0.93%)	0.19% (0.14-0.24%)	0.088 (0.064-0.12)	88 (53-140)
Kuwait	2000	Female	35 to 39	16% (4.4-31%)	84% (69-96%)	0.011 (0.0037-0.022)	0.51% (0.35-0.71%)	0.24% (0.18-0.3%)	0.13 (0.088-0.17)	62 (39-92)
Kuwait	2000	Female	40 to 44	17% (5.9-32%)	83% (68-94%)	0.012 (0.0041-0.025)	0.4% (0.26-0.56%)	0.28% (0.21-0.36%)	0.18 (0.13-0.24)	46 (27-68)
Kuwait	2000	Female	45 to 49	17% (6.2-33%)	83% (67-94%)	0.013 (0.0042-0.029)	0.32% (0.21-0.44%)	0.28% (0.17-0.38%)	0.17 (0.099-0.24)	24 (15-36)
Kuwait	2000	Female	50 to 54	18% (6.8-36%)	82% (64-93%)	0.013 (0.0036-0.032)	0.27% (0.13-0.38%)	0.29% (0.1-0.42%)	0.23 (0.078-0.35)	19 (9.3-28)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	2000	Female	55 to 59	19% (7.2-37%)	81% (63-93%)	0.013 (0.0033-0.032)	0.2% (0.052-0.29%)	0.22% (0.02-0.34%)	0.17 (0.015-0.27)	9.4 (2.5-14)
Kuwait	2000	Female	60 to 64	20% (7.7-38%)	80% (62-92%)	0.013 (0.0028-0.033)	0.18% (-0.022-0.29%)	0.21% (-0.044-0.35%)	0.22 (-0.047-0.39)	7.8 (-0.97-13)
Kuwait	2000	Female	65 to 69	20% (8.6-37%)	80% (63-91%)	0.011 (0.0025-0.028)	0.18% (-0.0063-0.27%)	0.23% (-0.012-0.36%)	0.35 (-0.016-0.57)	9.3 (-0.35-15)
Kuwait	2000	Female	70 to 74	19% (8.2-36%)	81% (64-92%)	0.0088 (0.002-0.024)	0.16% (-0.032-0.24%)	0.19% (-0.036-0.29%)	0.22 (-0.041-0.34)	4.4 (-0.86-6.6)
Kuwait	2000	Female	75 to 79	17% (6.4-35%)	83% (65-94%)	0.0072 (0.0013-0.022)	0.13% (-0.032-0.19%)	0.16% (-0.036-0.24%)	0.27 (-0.067-0.41)	4.1 (-1.1-6.2)
Kuwait	2000	Female	80 to 84	16% (5.4-35%)	84% (65-95%)	0.0065 (0.0011-0.021)	0.098% (-0.024-0.14%)	0.11% (-0.035-0.16%)	0.11 (-0.037-0.17)	1.3 (-0.34-1.8)
Kuwait	2000	Female	85 to 89	16% (5.2-37%)	84% (63-95%)	0.0065 (0.0011-0.02)	0.071% (-0.037-0.098%)	0.077% (-0.048-0.11%)	0.042 (-0.025-0.061)	0.36 (-0.18-0.51)
Kuwait	2000	Female	90 to 94	19% (5.7-40%)	81% (60-94%)	0.0074 (0.0013-0.023)	0.071% (-0.05-0.1%)	0.074% (-0.065-0.11%)	0.047 (-0.04-0.069)	0.28 (-0.19-0.4)
Kuwait	2000	Female	95 plus	23% (7.1-48%)	77% (52-93%)	0.0092 (0.0013-0.029)	0.056% (-0.11-0.091%)	0.046% (-0.18-0.085%)	0.014 (-0.051-0.026)	0.071 (-0.15-0.12)
Kuwait	2000	Male	All Ages	10% (6.6-15%)	90% (85-93%)	0.0082 (0.0047-0.012)	0.62% (0.5-0.75%)	0.37% (0.3-0.44%)	10 (7.7-13)	1300 (980-1700)
Kuwait	2000	Male	15 to 19	6.4% (2.7-13%)	94% (87-97%)	0.0028 (0.00072-0.0075)	0.48% (0.29-0.74%)	0.076% (0.053-0.12%)	0.039 (0.026-0.063)	44 (25-71)
Kuwait	2000	Male	20 to 24	9.6% (3.7-18%)	90% (82-96%)	0.0061 (0.0018-0.014)	1.1% (0.68-1.6%)	0.27% (0.19-0.41%)	0.21 (0.14-0.33)	130 (78-210)
Kuwait	2000	Male	25 to 29	12% (4.6-23%)	88% (77-95%)	0.0092 (0.003-0.019)	1.1% (0.81-1.6%)	0.32% (0.22-0.46%)	0.38 (0.25-0.56)	240 (150-350)
Kuwait	2000	Male	30 to 34	14% (5.2-27%)	86% (73-95%)	0.012 (0.0046-0.022)	1% (0.72-1.4%)	0.47% (0.36-0.6%)	0.62 (0.45-0.84)	230 (150-330)
Kuwait	2000	Male	35 to 39	15% (5.9-28%)	85% (72-94%)	0.013 (0.0051-0.025)	0.95% (0.72-1.2%)	0.69% (0.54-0.87%)	1 (0.72-1.3)	200 (140-280)
Kuwait	2000	Male	40 to 44	15% (6.9-27%)	85% (73-93%)	0.014 (0.0053-0.027)	0.81% (0.6-1.1%)	0.68% (0.52-0.85%)	1.3 (0.93-1.7)	170 (120-230)
Kuwait	2000	Male	45 to 49	15% (5.9-28%)	85% (72-94%)	0.015 (0.0048-0.032)	0.71% (0.56-0.89%)	0.74% (0.54-0.93%)	1.4 (0.94-1.9)	110 (83-150)
Kuwait	2000	Male	50 to 54	14% (6.1-29%)	86% (71-94%)	0.015 (0.0043-0.034)	0.53% (0.38-0.68%)	0.56% (0.36-0.76%)	1.2 (0.73-1.7)	76 (52-100)
Kuwait	2000	Male	55 to 59	14% (5.8-27%)	86% (73-94%)	0.014 (0.0037-0.034)	0.42% (0.29-0.55%)	0.46% (0.27-0.65%)	0.95 (0.54-1.4)	46 (31-63)
Kuwait	2000	Male	60 to 64	13% (5.6-25%)	87% (75-94%)	0.013 (0.0031-0.032)	0.37% (0.23-0.5%)	0.42% (0.22-0.61%)	0.83 (0.46-1.3)	29 (17-40)
Kuwait	2000	Male	65 to 69	13% (5.5-23%)	87% (77-94%)	0.01 (0.0029-0.026)	0.29% (0.19-0.39%)	0.35% (0.19-0.51%)	0.73 (0.4-1.1)	20 (13-28)
Kuwait	2000	Male	70 to 74	11% (5.1-21%)	89% (79-95%)	0.0078 (0.002-0.02)	0.23% (0.15-0.33%)	0.29% (0.16-0.43%)	0.66 (0.38-0.99)	14 (8.4-20)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	2000	Male	75 to 79	9.8% (4.2-19%)	90% (81-96%)	0.0059 (0.0012-0.016)	0.21% (0.14-0.27%)	0.25% (0.16-0.34%)	0.39 (0.24-0.55)	6.1 (3.9-8.4)
Kuwait	2000	Male	80 to 84	8.8% (3.5-18%)	91% (82-96%)	0.005 (0.0011-0.013)	0.17% (0.11-0.24%)	0.2% (0.12-0.28%)	0.21 (0.13-0.3)	2.3 (1.5-3.2)
Kuwait	2000	Male	85 to 89	8.6% (3.4-17%)	91% (83-97%)	0.0045 (0.001-0.011)	0.15% (0.11-0.19%)	0.17% (0.12-0.23%)	0.11 (0.074-0.15)	0.92 (0.65-1.2)
Kuwait	2000	Male	90 to 94	9.7% (3.8-19%)	90% (81-96%)	0.0048 (0.001-0.013)	0.11% (0.069-0.14%)	0.11% (0.06-0.15%)	0.051 (0.028-0.068)	0.32 (0.2-0.42)
Kuwait	2000	Male	95 plus	12% (4.5-26%)	88% (74-95%)	0.0056 (0.001-0.017)	0.15% (0.097-0.19%)	0.13% (0.058-0.17%)	0.026 (0.011-0.035)	0.13 (0.086-0.17)
Kuwait	2005	Female	All Ages	11% (6.5-18%)	89% (82-94%)	0.0051 (0.0028-0.0078)	0.35% (0.27-0.44%)	0.18% (0.12-0.21%)	3 (1.9-3.8)	560 (390-750)
Kuwait	2005	Female	15 to 19	8.5% (2.8-19%)	91% (81-97%)	0.0033 (0.00075-0.0086)	0.34% (0.19-0.56%)	0.085% (0.068-0.11%)	0.016 (0.012-0.021)	27 (14-46)
Kuwait	2005	Female	20 to 24	11% (3.2-24%)	89% (76-97%)	0.0052 (0.0013-0.012)	0.73% (0.44-1.1%)	0.1% (0.078-0.13%)	0.028 (0.021-0.039)	83 (45-140)
Kuwait	2005	Female	25 to 29	12% (3.5-27%)	88% (73-96%)	0.0062 (0.0016-0.014)	0.76% (0.51-1.1%)	0.17% (0.13-0.22%)	0.064 (0.046-0.089)	110 (63-160)
Kuwait	2005	Female	30 to 34	14% (3.7-30%)	86% (70-96%)	0.0074 (0.0022-0.015)	0.66% (0.43-0.97%)	0.17% (0.14-0.22%)	0.079 (0.058-0.11)	97 (57-150)
Kuwait	2005	Female	35 to 39	16% (4.3-32%)	84% (68-96%)	0.0087 (0.0027-0.017)	0.52% (0.35-0.71%)	0.22% (0.17-0.28%)	0.12 (0.085-0.17)	74 (46-110)
Kuwait	2005	Female	40 to 44	17% (5.6-33%)	83% (67-94%)	0.0098 (0.0032-0.02)	0.4% (0.27-0.58%)	0.27% (0.2-0.34%)	0.16 (0.12-0.23)	49 (30-75)
Kuwait	2005	Female	45 to 49	17% (5.9-33%)	83% (67-94%)	0.011 (0.0029-0.023)	0.33% (0.22-0.46%)	0.29% (0.2-0.39%)	0.23 (0.15-0.32)	38 (25-54)
Kuwait	2005	Female	50 to 54	18% (6.8-35%)	82% (65-93%)	0.011 (0.0027-0.025)	0.29% (0.16-0.39%)	0.32% (0.2-0.45%)	0.25 (0.14-0.36)	22 (12-31)
Kuwait	2005	Female	55 to 59	18% (7.2-37%)	82% (63-93%)	0.011 (0.0026-0.027)	0.23% (0.11-0.31%)	0.27% (0.12-0.38%)	0.3 (0.14-0.44)	17 (7.5-24)
Kuwait	2005	Female	60 to 64	20% (7.4-38%)	80% (62-93%)	0.01 (0.0022-0.026)	0.21% (0.043-0.31%)	0.27% (0.076-0.4%)	0.35 (0.1-0.54)	12 (2.4-18)
Kuwait	2005	Female	65 to 69	20% (8.5-37%)	80% (63-91%)	0.0091 (0.0019-0.023)	0.2% (0.058-0.27%)	0.25% (0.078-0.35%)	0.42 (0.14-0.6)	11 (3.3-16)
Kuwait	2005	Female	70 to 74	19% (8.2-36%)	81% (64-92%)	0.0073 (0.0016-0.021)	0.17% (0.054-0.24%)	0.22% (0.081-0.31%)	0.52 (0.19-0.73)	11 (3.2-14)
Kuwait	2005	Female	75 to 79	17% (6.4-35%)	83% (65-94%)	0.006 (0.0011-0.019)	0.15% (0.051-0.2%)	0.18% (0.069-0.24%)	0.22 (0.087-0.31)	3.3 (1.2-4.6)
Kuwait	2005	Female	80 to 84	16% (5.3-35%)	84% (65-95%)	0.0054 (0.00095-0.017)	0.1% (0.058-0.14%)	0.11% (0.064-0.15%)	0.13 (0.073-0.18)	1.5 (0.84-2)
Kuwait	2005	Female	85 to 89	16% (5.4-36%)	84% (64-95%)	0.0055 (0.00092-0.017)	0.079% (0.047-0.1%)	0.088% (0.046-0.11%)	0.072 (0.038-0.096)	0.62 (0.35-0.81)
Kuwait	2005	Female	90 to 94	19% (6.3-40%)	81% (60-94%)	0.0063 (0.0011-0.02)	0.082% (0.018-0.11%)	0.089% (0.0091-0.12%)	0.032 (0.0031-0.044)	0.19 (0.04-0.25)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	2005	Female	95 plus	23% (7.7-48%)	77% (52-92%)	0.0076 (0.0011-0.024)	0.069% (-0.031-0.098%)	0.063% (-0.055-0.095%)	0.018 (-0.016-0.027)	0.084 (-0.039-0.12)
Kuwait	2005	Male	All Ages	10% (6.3-15%)	90% (85-94%)	0.0066 (0.0038-0.0099)	0.62% (0.5-0.74%)	0.39% (0.33-0.45%)	13 (9.7-16)	1600 (1200-2100)
Kuwait	2005	Male	15 to 19	6.1% (2.6-12%)	94% (88-97%)	0.0022 (0.00058-0.0055)	0.46% (0.28-0.7%)	0.066% (0.046-0.096%)	0.046 (0.03-0.067)	51 (29-81)
Kuwait	2005	Male	20 to 24	9.4% (3.7-18%)	91% (82-96%)	0.0048 (0.0014-0.01)	1% (0.66-1.5%)	0.18% (0.13-0.27%)	0.2 (0.14-0.31)	170 (100-260)
Kuwait	2005	Male	25 to 29	12% (4.6-23%)	88% (77-95%)	0.0072 (0.0024-0.015)	1% (0.71-1.4%)	0.23% (0.16-0.33%)	0.36 (0.24-0.55)	260 (160-380)
Kuwait	2005	Male	30 to 34	13% (4.9-26%)	87% (74-95%)	0.0092 (0.0035-0.017)	0.99% (0.68-1.3%)	0.38% (0.3-0.49%)	0.66 (0.49-0.9)	280 (180-410)
Kuwait	2005	Male	35 to 39	15% (5.5-27%)	85% (73-94%)	0.01 (0.0037-0.02)	0.91% (0.68-1.2%)	0.58% (0.46-0.74%)	1.1 (0.79-1.4)	230 (160-320)
Kuwait	2005	Male	40 to 44	15% (6.5-27%)	85% (73-94%)	0.011 (0.004-0.023)	0.79% (0.58-1%)	0.63% (0.49-0.79%)	1.3 (0.97-1.8)	190 (130-260)
Kuwait	2005	Male	45 to 49	15% (5.7-27%)	85% (73-94%)	0.012 (0.0034-0.026)	0.71% (0.55-0.89%)	0.76% (0.57-0.95%)	2 (1.4-2.6)	160 (120-200)
Kuwait	2005	Male	50 to 54	14% (5.9-28%)	86% (72-94%)	0.012 (0.0033-0.027)	0.56% (0.42-0.72%)	0.62% (0.42-0.83%)	1.5 (0.95-2.1)	86 (61-110)
Kuwait	2005	Male	55 to 59	13% (5.7-26%)	87% (74-94%)	0.012 (0.0029-0.027)	0.45% (0.32-0.6%)	0.52% (0.32-0.72%)	1.2 (0.71-1.7)	53 (37-72)
Kuwait	2005	Male	60 to 64	13% (5.4-25%)	87% (75-95%)	0.01 (0.0025-0.027)	0.41% (0.29-0.57%)	0.5% (0.31-0.72%)	1.1 (0.7-1.6)	38 (25-52)
Kuwait	2005	Male	65 to 69	12% (5.3-22%)	88% (78-95%)	0.0084 (0.0021-0.021)	0.32% (0.22-0.42%)	0.39% (0.25-0.54%)	1.1 (0.68-1.6)	29 (19-41)
Kuwait	2005	Male	70 to 74	11% (4.9-20%)	89% (80-95%)	0.0062 (0.0015-0.016)	0.25% (0.18-0.36%)	0.33% (0.21-0.47%)	0.84 (0.53-1.3)	18 (12-25)
Kuwait	2005	Male	75 to 79	9.5% (3.9-19%)	91% (81-96%)	0.0047 (0.001-0.014)	0.23% (0.17-0.3%)	0.29% (0.21-0.4%)	0.63 (0.42-0.88)	9.9 (6.8-13)
Kuwait	2005	Male	80 to 84	8.5% (3.4-17%)	91% (83-97%)	0.004 (0.00088-0.012)	0.19% (0.14-0.26%)	0.23% (0.15-0.32%)	0.31 (0.21-0.43)	3.5 (2.4-4.8)
Kuwait	2005	Male	85 to 89	8.4% (3.4-17%)	92% (83-97%)	0.0036 (0.00078-0.0094)	0.17% (0.12-0.21%)	0.19% (0.14-0.24%)	0.16 (0.11-0.21)	1.3 (0.96-1.7)
Kuwait	2005	Male	90 to 94	9.3% (3.7-19%)	91% (81-96%)	0.0038 (0.00073-0.01)	0.12% (0.091-0.15%)	0.13% (0.092-0.16%)	0.05 (0.036-0.066)	0.31 (0.23-0.4)
Kuwait	2005	Male	95 plus	12% (4.1-25%)	88% (75-96%)	0.0044 (0.00071-0.014)	0.15% (0.12-0.2%)	0.14% (0.093-0.18%)	0.022 (0.015-0.03)	0.11 (0.08-0.14)
Kuwait	2010	Female	All Ages	11% (6.5-18%)	89% (82-94%)	0.0057 (0.0019-0.0098)	0.39% (0.3-0.49%)	0.18% (0.11-0.22%)	3.9 (2.3-5.1)	850 (590-1200)
Kuwait	2010	Female	15 to 19	8.6% (2.8-18%)	91% (82-97%)	0.0035 (0.00054-0.0097)	0.36% (0.2-0.55%)	0.1% (0.083-0.14%)	0.024 (0.018-0.032)	41 (21-67)
Kuwait	2010	Female	20 to 24	11% (3.4-25%)	89% (75-97%)	0.0055 (0.00098-0.014)	0.77% (0.47-1.1%)	0.11% (0.087-0.15%)	0.038 (0.028-0.053)	130 (72-210)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	2010	Female	25 to 29	13% (3.6-27%)	87% (73-96%)	0.0065 (0.0014-0.015)	0.79% (0.53-1.1%)	0.18% (0.14-0.23%)	0.095 (0.068-0.13)	180 (110-270)
Kuwait	2010	Female	30 to 34	14% (3.8-29%)	86% (71-96%)	0.0077 (0.0017-0.016)	0.69% (0.44-1%)	0.18% (0.15-0.24%)	0.1 (0.077-0.14)	160 (91-240)
Kuwait	2010	Female	35 to 39	16% (4.3-30%)	84% (70-96%)	0.009 (0.0022-0.019)	0.53% (0.35-0.74%)	0.22% (0.17-0.28%)	0.15 (0.11-0.21)	110 (66-160)
Kuwait	2010	Female	40 to 44	16% (6-31%)	84% (69-94%)	0.01 (0.0026-0.022)	0.41% (0.27-0.6%)	0.27% (0.2-0.35%)	0.22 (0.15-0.3)	77 (44-120)
Kuwait	2010	Female	45 to 49	17% (6-31%)	83% (69-94%)	0.011 (0.0025-0.027)	0.34% (0.22-0.46%)	0.29% (0.21-0.4%)	0.22 (0.15-0.31)	42 (25-62)
Kuwait	2010	Female	50 to 54	17% (6.6-34%)	83% (66-93%)	0.011 (0.0025-0.027)	0.28% (0.16-0.4%)	0.33% (0.2-0.46%)	0.35 (0.2-0.51)	35 (19-52)
Kuwait	2010	Female	55 to 59	18% (7.1-36%)	82% (64-93%)	0.011 (0.0022-0.031)	0.23% (0.1-0.33%)	0.3% (0.15-0.42%)	0.34 (0.16-0.5)	20 (8.2-28)
Kuwait	2010	Female	60 to 64	19% (7.3-38%)	81% (62-93%)	0.011 (0.0019-0.031)	0.22% (0.055-0.32%)	0.3% (0.085-0.45%)	0.5 (0.15-0.78)	18 (4.5-28)
Kuwait	2010	Female	65 to 69	20% (8-37%)	80% (63-92%)	0.0094 (0.0016-0.024)	0.21% (0.059-0.3%)	0.28% (0.09-0.39%)	0.52 (0.16-0.78)	14 (3.5-20)
Kuwait	2010	Female	70 to 74	19% (7.4-35%)	81% (65-93%)	0.0075 (0.0012-0.023)	0.18% (0.042-0.25%)	0.23% (0.062-0.31%)	0.53 (0.14-0.75)	11 (2.4-15)
Kuwait	2010	Female	75 to 79	17% (5.9-35%)	83% (65-94%)	0.0061 (0.00092-0.02)	0.15% (0.02-0.2%)	0.19% (0.032-0.26%)	0.49 (0.086-0.71)	7.6 (1.1-11)
Kuwait	2010	Female	80 to 84	15% (4.8-33%)	85% (67-95%)	0.0054 (0.00079-0.018)	0.11% (0.0013-0.14%)	0.12% (0.0023-0.16%)	0.21 (0.0042-0.3)	2.4 (0.03-3.2)
Kuwait	2010	Female	85 to 89	16% (5-36%)	84% (64-95%)	0.0054 (0.00078-0.017)	0.081% (0.0083-0.11%)	0.089% (0.0032-0.12%)	0.082 (0.0032-0.11)	0.68 (0.065-0.93)
Kuwait	2010	Female	90 to 94	18% (5.9-40%)	82% (60-94%)	0.0062 (0.00086-0.02)	0.084% (-0.02-0.11%)	0.091% (-0.031-0.12%)	0.052 (-0.019-0.073)	0.31 (-0.078-0.42)
Kuwait	2010	Female	95 plus	22% (6.5-47%)	78% (53-94%)	0.0076 (0.00085-0.027)	0.068% (-0.075-0.1%)	0.062% (-0.12-0.1%)	0.011 (-0.02-0.017)	0.049 (-0.056-0.075)
Kuwait	2010	Male	All Ages	10% (6.5-15%)	90% (85-93%)	0.0072 (0.0027-0.012)	0.66% (0.54-0.79%)	0.4% (0.32-0.46%)	14 (11-18)	2000 (1500-2700)
Kuwait	2010	Male	15 to 19	6.1% (2.6-12%)	94% (88-97%)	0.0023 (0.00042-0.0061)	0.47% (0.29-0.72%)	0.075% (0.048-0.12%)	0.058 (0.035-0.097)	60 (34-98)
Kuwait	2010	Male	20 to 24	9.4% (3.6-18%)	91% (82-96%)	0.005 (0.0011-0.012)	1.1% (0.71-1.7%)	0.19% (0.14-0.31%)	0.2 (0.13-0.34)	210 (120-320)
Kuwait	2010	Male	25 to 29	12% (4.5-23%)	88% (77-95%)	0.0076 (0.0019-0.017)	1.2% (0.8-1.6%)	0.25% (0.18-0.37%)	0.41 (0.28-0.62)	350 (220-520)
Kuwait	2010	Male	30 to 34	13% (4.7-26%)	87% (74-95%)	0.0096 (0.0026-0.02)	1.1% (0.74-1.5%)	0.42% (0.32-0.55%)	0.69 (0.52-0.94)	360 (230-530)
Kuwait	2010	Male	35 to 39	15% (5.4-27%)	85% (73-95%)	0.011 (0.0028-0.023)	0.95% (0.7-1.3%)	0.63% (0.49-0.8%)	1.3 (0.94-1.7)	320 (220-460)
Kuwait	2010	Male	40 to 44	15% (6.7-26%)	85% (74-93%)	0.012 (0.0029-0.024)	0.8% (0.59-1.1%)	0.64% (0.49-0.8%)	1.5 (1-2)	230 (160-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	2010	Male	45 to 49	14% (5.6-26%)	86% (74-94%)	0.012 (0.0027-0.028)	0.71% (0.55-0.88%)	0.76% (0.57-0.95%)	2 (1.4-2.7)	180 (130-230)
Kuwait	2010	Male	50 to 54	14% (5.6-28%)	86% (72-94%)	0.012 (0.0026-0.03)	0.57% (0.43-0.71%)	0.65% (0.44-0.86%)	2.1 (1.4-2.9)	130 (95-170)
Kuwait	2010	Male	55 to 59	13% (5.5-25%)	87% (75-95%)	0.012 (0.0021-0.03)	0.44% (0.3-0.58%)	0.51% (0.31-0.72%)	1.3 (0.79-2)	63 (41-84)
Kuwait	2010	Male	60 to 64	13% (5.2-24%)	87% (76-95%)	0.011 (0.0019-0.028)	0.41% (0.28-0.57%)	0.53% (0.34-0.76%)	1.4 (0.85-2)	47 (31-66)
Kuwait	2010	Male	65 to 69	12% (5.2-22%)	88% (78-95%)	0.0086 (0.0015-0.022)	0.32% (0.22-0.43%)	0.4% (0.25-0.58%)	0.93 (0.54-1.4)	25 (16-36)
Kuwait	2010	Male	70 to 74	11% (4.9-20%)	89% (80-95%)	0.0064 (0.0013-0.017)	0.26% (0.18-0.36%)	0.33% (0.21-0.47%)	0.89 (0.55-1.3)	19 (12-27)
Kuwait	2010	Male	75 to 79	9.5% (3.9-19%)	91% (81-96%)	0.0049 (0.00081-0.015)	0.22% (0.16-0.28%)	0.29% (0.2-0.39%)	0.66 (0.45-0.93)	11 (7.5-14)
Kuwait	2010	Male	80 to 84	8.4% (3.2-17%)	92% (83-97%)	0.0041 (0.00072-0.012)	0.19% (0.13-0.25%)	0.23% (0.16-0.31%)	0.37 (0.25-0.52)	4.3 (3-5.8)
Kuwait	2010	Male	85 to 89	8.3% (3.3-16%)	92% (84-97%)	0.0037 (0.00068-0.01)	0.17% (0.13-0.21%)	0.2% (0.14-0.26%)	0.23 (0.16-0.3)	1.9 (1.4-2.5)
Kuwait	2010	Male	90 to 94	9.4% (4-19%)	91% (81-96%)	0.004 (0.0007-0.011)	0.12% (0.091-0.16%)	0.13% (0.09-0.17%)	0.058 (0.041-0.077)	0.38 (0.28-0.49)
Kuwait	2010	Male	95 plus	12% (4.3-25%)	88% (75-96%)	0.0046 (0.00063-0.015)	0.16% (0.12-0.21%)	0.14% (0.097-0.19%)	0.019 (0.013-0.025)	0.097 (0.071-0.13)
Kuwait	2016	Female	All Ages	12% (7-19%)	88% (81-93%)	0.013 (0.0027-0.028)	0.39% (0.28-0.5%)	0.15% (-0.0068-0.23%)	3.8 (-0.16-7.1)	1100 (710-1500)
Kuwait	2016	Female	15 to 19	8.7% (2.8-18%)	91% (82-97%)	0.0074 (0.00093-0.024)	0.36% (0.2-0.57%)	0.14% (0.093-0.22%)	0.024 (0.014-0.043)	38 (20-64)
Kuwait	2016	Female	20 to 24	11% (3.2-24%)	89% (76-97%)	0.012 (0.0015-0.037)	0.79% (0.49-1.2%)	0.14% (0.1-0.24%)	0.04 (0.022-0.08)	130 (74-220)
Kuwait	2016	Female	25 to 29	13% (3.5-26%)	87% (74-96%)	0.014 (0.0017-0.038)	0.82% (0.55-1.2%)	0.21% (0.14-0.33%)	0.098 (0.051-0.18)	210 (120-320)
Kuwait	2016	Female	30 to 34	14% (3.5-30%)	86% (70-96%)	0.016 (0.0024-0.042)	0.72% (0.46-1.1%)	0.21% (0.15-0.32%)	0.13 (0.072-0.23)	220 (130-350)
Kuwait	2016	Female	35 to 39	16% (4.2-31%)	84% (69-96%)	0.019 (0.0027-0.049)	0.55% (0.36-0.77%)	0.24% (0.17-0.36%)	0.19 (0.094-0.33)	160 (96-240)
Kuwait	2016	Female	40 to 44	17% (5.8-32%)	83% (68-94%)	0.021 (0.0032-0.054)	0.41% (0.23-0.61%)	0.28% (0.19-0.41%)	0.26 (0.13-0.45)	110 (56-160)
Kuwait	2016	Female	45 to 49	17% (6-32%)	83% (68-94%)	0.023 (0.0033-0.06)	0.33% (0.18-0.48%)	0.3% (0.16-0.44%)	0.35 (0.16-0.62)	78 (40-120)
Kuwait	2016	Female	50 to 54	18% (6.9-35%)	82% (65-93%)	0.024 (0.003-0.069)	0.25% (0.074-0.39%)	0.29% (0.1-0.45%)	0.39 (0.13-0.71)	44 (12-72)
Kuwait	2016	Female	55 to 59	18% (7.1-36%)	82% (64-93%)	0.023 (0.0027-0.074)	0.19% (-0.0049-0.31%)	0.26% (0.051-0.42%)	0.47 (0.08-0.86)	29 (-0.98-52)
Kuwait	2016	Female	60 to 64	19% (7.1-36%)	81% (64-93%)	0.022 (0.0025-0.073)	0.15% (-0.099-0.29%)	0.22% (-0.092-0.41%)	0.5 (-0.22-1)	18 (-12-38)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	2016	Female	65 to 69	19% (8.1-35%)	81% (65-92%)	0.019 (0.0022-0.058)	0.14% (-0.11-0.28%)	0.2% (-0.11-0.39%)	0.59 (-0.32-1.3)	15 (-11-33)
Kuwait	2016	Female	70 to 74	18% (7.6-34%)	82% (66-92%)	0.015 (0.002-0.047)	0.11% (-0.13-0.23%)	0.15% (-0.17-0.3%)	0.39 (-0.41-0.92)	7.7 (-8.9-18)
Kuwait	2016	Female	75 to 79	16% (6-34%)	84% (66-94%)	0.012 (0.0012-0.045)	0.088% (-0.17-0.19%)	0.11% (-0.2-0.24%)	0.24 (-0.4-0.58)	3.7 (-7.1-8.8)
Kuwait	2016	Female	80 to 84	15% (4.8-33%)	85% (67-95%)	0.011 (0.00099-0.041)	0.048% (-0.22-0.13%)	0.052% (-0.25-0.15%)	0.13 (-0.56-0.41)	1.5 (-6.4-4.6)
Kuwait	2016	Female	85 to 89	16% (4.9-35%)	84% (65-95%)	0.011 (0.00083-0.041)	0.026% (-0.23-0.1%)	0.024% (-0.27-0.11%)	0.042 (-0.46-0.21)	0.42 (-3.5-1.8)
Kuwait	2016	Female	90 to 94	18% (5.6-40%)	82% (60-94%)	0.013 (0.0011-0.047)	0.029% (-0.22-0.1%)	0.024% (-0.26-0.11%)	0.011 (-0.13-0.06)	0.091 (-0.68-0.35)
Kuwait	2016	Female	95 plus	23% (6.5-47%)	77% (53-94%)	0.016 (0.0011-0.065)	0.013% (-0.28-0.1%)	-0.0094% (-0.39-0.093%)	-0.0028 (-0.12-0.03)	0.018 (-0.39-0.14)
Kuwait	2016	Male	All Ages	10% (6.4-15%)	90% (85-94%)	0.015 (0.0029-0.032)	0.7% (0.55-0.84%)	0.4% (0.26-0.49%)	17 (9.6-26)	2600 (1900-3500)
Kuwait	2016	Male	15 to 19	5.9% (2.5-12%)	94% (88-97%)	0.0046 (0.00058-0.014)	0.49% (0.29-0.73%)	0.1% (0.052-0.23%)	0.087 (0.038-0.22)	76 (43-120)
Kuwait	2016	Male	20 to 24	9.1% (3.5-18%)	91% (82-96%)	0.01 (0.0014-0.029)	1.2% (0.77-1.8%)	0.28% (0.16-0.52%)	0.28 (0.13-0.59)	230 (140-370)
Kuwait	2016	Male	25 to 29	11% (4.4-22%)	89% (78-96%)	0.015 (0.0024-0.042)	1.2% (0.84-1.7%)	0.35% (0.21-0.62%)	0.53 (0.29-1)	390 (240-580)
Kuwait	2016	Male	30 to 34	13% (4.8-25%)	87% (75-95%)	0.019 (0.0033-0.047)	1.1% (0.79-1.6%)	0.5% (0.35-0.72%)	0.97 (0.56-1.6)	500 (310-730)
Kuwait	2016	Male	35 to 39	14% (5.5-26%)	86% (74-95%)	0.022 (0.0034-0.057)	1% (0.75-1.3%)	0.69% (0.5-0.92%)	1.6 (0.94-2.6)	450 (300-630)
Kuwait	2016	Male	40 to 44	14% (6.4-26%)	86% (74-94%)	0.024 (0.0039-0.064)	0.84% (0.59-1.1%)	0.64% (0.44-0.84%)	1.8 (0.99-2.9)	340 (210-480)
Kuwait	2016	Male	45 to 49	14% (5.3-26%)	86% (74-95%)	0.025 (0.0036-0.068)	0.7% (0.53-0.9%)	0.7% (0.41-0.93%)	2.4 (1.2-4)	250 (160-340)
Kuwait	2016	Male	50 to 54	13% (5.4-26%)	87% (74-95%)	0.025 (0.0036-0.068)	0.54% (0.34-0.7%)	0.58% (0.26-0.8%)	2.2 (0.86-3.8)	160 (94-220)
Kuwait	2016	Male	55 to 59	13% (5.4-24%)	87% (76-95%)	0.024 (0.0029-0.071)	0.4% (0.2-0.55%)	0.43% (0.089-0.67%)	1.9 (0.43-3.5)	100 (52-150)
Kuwait	2016	Male	60 to 64	12% (5.1-23%)	88% (77-95%)	0.021 (0.0026-0.068)	0.37% (0.18-0.53%)	0.45% (0.15-0.72%)	1.6 (0.46-2.8)	58 (27-90)
Kuwait	2016	Male	65 to 69	12% (4.8-21%)	88% (79-95%)	0.017 (0.0023-0.052)	0.29% (0.15-0.42%)	0.36% (0.13-0.56%)	1.2 (0.44-2.1)	34 (17-54)
Kuwait	2016	Male	70 to 74	10% (4.7-19%)	90% (81-95%)	0.013 (0.0018-0.038)	0.24% (0.13-0.36%)	0.3% (0.14-0.47%)	0.84 (0.35-1.5)	18 (8.8-29)
Kuwait	2016	Male	75 to 79	9.1% (3.8-18%)	91% (82-96%)	0.0097 (0.0011-0.029)	0.2% (0.12-0.28%)	0.26% (0.14-0.37%)	0.59 (0.29-0.99)	9.9 (5.4-15)
Kuwait	2016	Male	80 to 84	8.1% (3.2-17%)	92% (83-97%)	0.0081 (0.00082-0.026)	0.16% (0.085-0.23%)	0.2% (0.096-0.3%)	0.37 (0.16-0.61)	4.4 (2.3-6.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kuwait	2016	Male	85 to 89	8% (3.3-16%)	92% (84-97%)	0.0074 (0.00084-0.023)	0.15% (0.086-0.2%)	0.17% (0.091-0.24%)	0.28 (0.14-0.42)	2.4 (1.4-3.5)
Kuwait	2016	Male	90 to 94	9.1% (3.7-19%)	91% (81-96%)	0.0079 (0.0009-0.027)	0.11% (0.047-0.15%)	0.11% (0.025-0.16%)	0.092 (0.021-0.14)	0.65 (0.26-0.94)
Kuwait	2016	Male	95 plus	11% (4.1-24%)	89% (76-96%)	0.0092 (0.00075-0.034)	0.15% (0.069-0.2%)	0.12% (0.0065-0.18%)	0.025 (0.0011-0.039)	0.14 (0.064-0.2)
Lebanon	1990	Female	All Ages	15% (12-18%)	85% (82-88%)	0.35 (0.27-0.44)	0.52% (0.24-0.85%)	0.38% (-0.23-1%)	32 (-19-93)	2400 (1000-3800)
Lebanon	1990	Female	15 to 19	15% (9.5-22%)	85% (78-90%)	0.34 (0.18-0.58)	0.55% (0.33-0.86%)	0.59% (0.27-1.1%)	0.83 (0.37-1.6)	130 (75-200)
Lebanon	1990	Female	20 to 24	20% (13-28%)	80% (72-87%)	0.55 (0.32-0.88)	1.1% (0.73-1.6%)	1.1% (0.52-1.9%)	1.7 (0.77-3)	270 (170-410)
Lebanon	1990	Female	25 to 29	20% (13-29%)	80% (71-87%)	0.61 (0.35-0.94)	1.3% (0.88-1.9%)	1.5% (0.77-2.6%)	2.4 (1.2-4.2)	320 (200-490)
Lebanon	1990	Female	30 to 34	19% (12-28%)	81% (72-88%)	0.58 (0.34-0.92)	1.3% (0.83-1.9%)	1.7% (0.8-2.8%)	2.6 (1.2-4.7)	280 (180-440)
Lebanon	1990	Female	35 to 39	18% (11-27%)	82% (73-89%)	0.55 (0.32-0.88)	1.2% (0.76-1.9%)	1.8% (0.82-3%)	2.8 (1.3-4.9)	230 (140-350)
Lebanon	1990	Female	40 to 44	18% (11-27%)	82% (73-89%)	0.57 (0.33-0.88)	1.2% (0.63-2%)	2% (0.84-3.4%)	3.5 (1.4-6)	220 (110-350)
Lebanon	1990	Female	45 to 49	18% (10-26%)	82% (74-90%)	0.61 (0.33-0.99)	1.2% (0.54-2.2%)	1.9% (0.74-3.5%)	5.3 (1.9-10)	270 (120-470)
Lebanon	1990	Female	50 to 54	18% (11-26%)	82% (74-89%)	0.62 (0.35-1)	1.2% (0.36-2.2%)	1.7% (0.48-3.4%)	6.2 (1.7-12)	260 (81-500)
Lebanon	1990	Female	55 to 59	18% (11-28%)	82% (72-89%)	0.58 (0.32-0.93)	0.93% (0.085-1.9%)	1.4% (0.095-2.9%)	6.6 (0.43-14)	230 (19-470)
Lebanon	1990	Female	60 to 64	18% (11-28%)	82% (72-89%)	0.53 (0.29-0.89)	0.55% (-0.32-1.7%)	0.77% (-0.45-2.3%)	4.9 (-2.8-14)	140 (-80-420)
Lebanon	1990	Female	65 to 69	18% (12-27%)	82% (73-88%)	0.49 (0.27-0.78)	0.26% (-0.54-1.2%)	0.36% (-0.75-1.6%)	2.7 (-5.4-11)	62 (-130-270)
Lebanon	1990	Female	70 to 74	17% (11-27%)	83% (73-89%)	0.43 (0.2-0.75)	0.012% (-0.84-0.95%)	0.02% (-1.1-1.2%)	0.1 (-7.6-8.3)	1.7 (-150-160)
Lebanon	1990	Female	75 to 79	16% (9.9-25%)	84% (75-90%)	0.37 (0.17-0.68)	-0.14% (-0.95-0.7%)	-0.19% (-1.2-0.84%)	-1.8 (-12-8.3)	-25 (-170-120)
Lebanon	1990	Female	80 to 84	15% (9.1-23%)	85% (77-91%)	0.33 (0.15-0.57)	-0.17% (-0.89-0.52%)	-0.22% (-1.1-0.63%)	-2 (-9.5-5.5)	-20 (-100-58)
Lebanon	1990	Female	85 to 89	14% (8.6-21%)	86% (79-91%)	0.3 (0.14-0.55)	-0.24% (-0.88-0.46%)	-0.32% (-1.1-0.52%)	-2.1 (-7.4-3.6)	-15 (-54-28)
Lebanon	1990	Female	90 to 94	15% (9.4-23%)	85% (77-91%)	0.3 (0.14-0.58)	-0.31% (-0.97-0.41%)	-0.41% (-1.2-0.46%)	-1.3 (-3.8-1.4)	-5.9 (-19-8)
Lebanon	1990	Female	95 plus	17% (9.6-29%)	83% (71-90%)	0.33 (0.12-0.69)	-0.38% (-1.3-0.65%)	-0.51% (-1.6-0.69%)	-0.49 (-1.5-0.68)	-1.5 (-4.9-2.5)
Lebanon	1990	Male	All Ages	23% (19-27%)	77% (73-81%)	1.3 (1.1-1.5)	2.9% (2-3.9%)	3.7% (2.2-5.7%)	410 (230-640)	16000 (11000-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	1990	Male	15 to 19	18% (11-27%)	82% (73-89%)	0.67 (0.33-1.2)	1.4% (0.76-2.3%)	1.6% (0.79-2.9%)	5.2 (2.5-9.3)	470 (260-800)
Lebanon	1990	Male	20 to 24	28% (18-40%)	72% (60-82%)	1.5 (0.83-2.5)	2.9% (1.7-4.6%)	3.2% (1.8-5.4%)	15 (8-26)	1200 (710-2000)
Lebanon	1990	Male	25 to 29	34% (22-46%)	66% (54-78%)	2.2 (1.3-3.4)	4% (2.4-6.1%)	5% (2.7-8%)	19 (10-31)	1400 (810-2100)
Lebanon	1990	Male	30 to 34	35% (24-49%)	65% (51-76%)	2.5 (1.5-3.8)	4.4% (2.6-6.6%)	5.8% (3.2-9.2%)	19 (10-30)	1200 (730-1900)
Lebanon	1990	Male	35 to 39	37% (24-50%)	63% (50-76%)	2.6 (1.5-4)	4.7% (2.6-7.2%)	6.5% (3.3-10%)	18 (8.9-29)	1100 (580-1700)
Lebanon	1990	Male	40 to 44	36% (25-49%)	64% (51-75%)	2.6 (1.5-4)	4.9% (2.6-7.5%)	6.7% (3.3-10%)	18 (8.7-30)	980 (520-1500)
Lebanon	1990	Male	45 to 49	34% (22-46%)	66% (54-78%)	2.6 (1.4-4.1)	5.1% (2.5-8.1%)	6.8% (3.1-11%)	27 (12-46)	1300 (630-2100)
Lebanon	1990	Male	50 to 54	32% (21-44%)	68% (56-79%)	2.7 (1.6-4.2)	5.5% (3-8.5%)	7.1% (3.8-11%)	38 (19-61)	1500 (810-2400)
Lebanon	1990	Male	55 to 59	31% (19-45%)	69% (55-81%)	3 (1.7-4.8)	6.2% (3.4-9.6%)	7.7% (4-12%)	55 (27-87)	1900 (1000-3000)
Lebanon	1990	Male	60 to 64	29% (18-44%)	71% (56-82%)	2.8 (1.5-4.8)	5.7% (2.7-9.5%)	6.8% (3.1-11%)	69 (31-120)	2000 (950-3400)
Lebanon	1990	Male	65 to 69	28% (17-42%)	72% (58-83%)	2.3 (1.2-3.7)	4.6% (1.8-8%)	5.3% (1.9-9.4%)	63 (22-110)	1500 (570-2700)
Lebanon	1990	Male	70 to 74	27% (17-40%)	73% (60-83%)	1.7 (0.91-2.9)	3% (0.48-6.1%)	3.5% (0.39-7.2%)	35 (3.8-75)	680 (110-1400)
Lebanon	1990	Male	75 to 79	25% (16-38%)	75% (62-84%)	1.3 (0.64-2.4)	1.8% (-0.44-4.7%)	1.9% (-0.62-5.4%)	21 (-7.1-60)	320 (-79-880)
Lebanon	1990	Male	80 to 84	23% (14-35%)	77% (65-86%)	1.1 (0.55-1.9)	1.1% (-0.52-3.4%)	1.2% (-0.75-3.8%)	9.9 (-6.6-33)	120 (-58-360)
Lebanon	1990	Male	85 to 89	21% (13-32%)	79% (68-87%)	0.91 (0.46-1.5)	0.52% (-0.68-2.4%)	0.47% (-0.98-2.7%)	2.6 (-5.7-15)	26 (-35-110)
Lebanon	1990	Male	90 to 94	22% (14-33%)	78% (67-86%)	0.82 (0.41-1.5)	0.069% (-1.1-1.7%)	-0.077% (-1.5-1.9%)	-0.17 (-3.3-4.1)	0.91 (-15-23)
Lebanon	1990	Male	95 plus	25% (14-42%)	75% (58-86%)	0.79 (0.31-1.7)	-0.21% (-1.6-1.5%)	-0.46% (-2.1-1.6%)	-0.26 (-1.2-0.91)	-0.48 (-3.6-3.4)
Lebanon	1995	Female	All Ages	14% (12-16%)	86% (84-88%)	0.31 (0.24-0.38)	0.5% (0.23-0.79%)	0.23% (-0.3-0.85%)	19 (-25-70)	2100 (920-3400)
Lebanon	1995	Female	15 to 19	14% (9.6-20%)	86% (80-90%)	0.3 (0.16-0.49)	0.6% (0.37-0.9%)	0.84% (0.37-1.5%)	0.68 (0.29-1.2)	120 (73-190)
Lebanon	1995	Female	20 to 24	18% (13-25%)	82% (75-87%)	0.46 (0.29-0.7)	1.2% (0.78-1.7%)	1.4% (0.67-2.3%)	1.3 (0.62-2.4)	260 (160-390)
Lebanon	1995	Female	25 to 29	18% (12-25%)	82% (75-88%)	0.49 (0.3-0.74)	1.3% (0.87-1.8%)	1.6% (0.8-2.6%)	1.8 (0.83-3.1)	290 (190-430)
Lebanon	1995	Female	30 to 34	17% (11-24%)	83% (76-89%)	0.46 (0.29-0.71)	1.2% (0.81-1.7%)	1.6% (0.84-2.8%)	2.3 (1.1-4.1)	300 (190-450)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	1995	Female	35 to 39	17% (12-24%)	83% (76-88%)	0.47 (0.29-0.72)	1.1% (0.72-1.6%)	1.7% (0.86-2.9%)	2.8 (1.3-4.9)	260 (160-380)
Lebanon	1995	Female	40 to 44	17% (11-23%)	83% (77-89%)	0.48 (0.3-0.73)	1.1% (0.62-1.7%)	1.8% (0.77-3.1%)	3.3 (1.4-5.8)	220 (120-350)
Lebanon	1995	Female	45 to 49	16% (9.9-22%)	84% (78-90%)	0.5 (0.28-0.77)	1% (0.48-1.8%)	1.6% (0.62-3%)	3.7 (1.4-6.8)	200 (95-340)
Lebanon	1995	Female	50 to 54	16% (11-23%)	84% (77-89%)	0.5 (0.31-0.8)	0.94% (0.27-1.8%)	1.4% (0.33-2.8%)	5.1 (1.2-10)	220 (68-430)
Lebanon	1995	Female	55 to 59	17% (10-25%)	83% (75-90%)	0.49 (0.26-0.78)	0.73% (0.034-1.5%)	1.1% (0.023-2.3%)	5 (0.11-11)	180 (8.5-380)
Lebanon	1995	Female	60 to 64	17% (10-26%)	83% (74-90%)	0.45 (0.24-0.74)	0.39% (-0.33-1.2%)	0.55% (-0.49-1.8%)	3.6 (-3.2-11)	100 (-94-330)
Lebanon	1995	Female	65 to 69	17% (11-25%)	83% (75-89%)	0.4 (0.23-0.65)	0.12% (-0.58-0.83%)	0.18% (-0.76-1.1%)	1.5 (-6.6-9.5)	34 (-160-230)
Lebanon	1995	Female	70 to 74	16% (11-25%)	84% (75-89%)	0.35 (0.17-0.61)	-0.11% (-0.84-0.65%)	-0.13% (-1.1-0.83%)	-1.3 (-10-7.9)	-26 (-200-150)
Lebanon	1995	Female	75 to 79	15% (9.4-23%)	85% (77-91%)	0.3 (0.14-0.54)	-0.24% (-0.98-0.45%)	-0.3% (-1.2-0.56%)	-2.6 (-10-4.8)	-37 (-150-70)
Lebanon	1995	Female	80 to 84	14% (8.5-21%)	86% (79-91%)	0.27 (0.13-0.46)	-0.26% (-0.86-0.33%)	-0.32% (-1.1-0.41%)	-2.6 (-8.9-3.3)	-28 (-94-35)
Lebanon	1995	Female	85 to 89	13% (7.9-20%)	87% (80-92%)	0.25 (0.12-0.45)	-0.31% (-0.85-0.21%)	-0.39% (-1.1-0.23%)	-3 (-8.1-1.8)	-21 (-59-15)
Lebanon	1995	Female	90 to 94	14% (8.7-21%)	86% (79-91%)	0.25 (0.11-0.48)	-0.36% (-0.92-0.2%)	-0.47% (-1.2-0.22%)	-1.8 (-4.6-0.84)	-8.7 (-23-5)
Lebanon	1995	Female	95 plus	16% (9.3-26%)	84% (74-91%)	0.27 (0.097-0.56)	-0.43% (-1.2-0.33%)	-0.57% (-1.5-0.35%)	-0.69 (-1.8-0.41)	-2.1 (-5.8-1.6)
Lebanon	1995	Male	All Ages	24% (21-28%)	76% (72-79%)	1.3 (1.1-1.5)	2.9% (2-3.8%)	3.5% (1.9-5.3%)	290 (150-440)	12000 (8300-16000)
Lebanon	1995	Male	15 to 19	19% (12-27%)	81% (73-88%)	0.62 (0.32-1)	1.7% (1-2.6%)	2.7% (1.4-4.7%)	3.9 (2-6.8)	390 (230-620)
Lebanon	1995	Male	20 to 24	30% (21-40%)	70% (60-79%)	1.5 (0.83-2.3)	4.1% (2.6-6.2%)	6.3% (3.6-10%)	12 (6.9-20)	1100 (680-1700)
Lebanon	1995	Male	25 to 29	35% (25-44%)	65% (56-75%)	2 (1.2-3)	4.9% (3.1-7.1%)	8.3% (4.8-13%)	16 (8.6-24)	1200 (740-1800)
Lebanon	1995	Male	30 to 34	36% (26-47%)	64% (53-74%)	2.2 (1.4-3.2)	4.6% (2.9-6.7%)	8.1% (4.6-12%)	16 (8.7-24)	1100 (680-1600)
Lebanon	1995	Male	35 to 39	37% (27-48%)	63% (52-73%)	2.3 (1.5-3.5)	4.5% (2.6-6.7%)	7.6% (3.9-12%)	15 (7.6-25)	990 (540-1500)
Lebanon	1995	Male	40 to 44	36% (27-47%)	64% (53-73%)	2.4 (1.4-3.6)	4.3% (2.2-6.7%)	6.9% (3.2-11%)	15 (6.7-24)	850 (430-1300)
Lebanon	1995	Male	45 to 49	33% (22-43%)	67% (57-78%)	2.3 (1.3-3.5)	4.3% (2-6.7%)	6.3% (2.7-10%)	16 (6.7-26)	780 (380-1200)
Lebanon	1995	Male	50 to 54	31% (22-42%)	69% (58-78%)	2.4 (1.4-3.7)	4.6% (2.2-7.1%)	6.5% (2.9-10%)	25 (12-41)	1100 (540-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	1995	Male	55 to 59	31% (20-46%)	69% (54-80%)	2.8 (1.5-4.5)	5.5% (2.9-8.6%)	7.3% (3.7-12%)	38 (19-63)	1400 (700-2200)
Lebanon	1995	Male	60 to 64	30% (18-46%)	70% (54-82%)	2.7 (1.4-4.6)	5.1% (2.2-8.4%)	6.5% (2.6-11%)	45 (18-77)	1400 (570-2300)
Lebanon	1995	Male	65 to 69	29% (18-43%)	71% (57-82%)	2.2 (1.2-3.5)	3.8% (1.3-6.8%)	4.7% (1.4-8.6%)	44 (13-80)	1100 (360-2000)
Lebanon	1995	Male	70 to 74	28% (18-41%)	72% (59-82%)	1.6 (0.87-2.7)	2.4% (0.15-5.3%)	2.7% (-0.064-6.4%)	28 (-0.63-65)	580 (39-1300)
Lebanon	1995	Male	75 to 79	26% (16-39%)	74% (61-84%)	1.3 (0.6-2.2)	1.2% (-0.48-3.8%)	1.3% (-0.83-4.5%)	10 (-6.8-35)	180 (-70-540)
Lebanon	1995	Male	80 to 84	23% (14-35%)	77% (65-86%)	1 (0.51-1.7)	0.67% (-0.68-2.6%)	0.64% (-0.98-3%)	4.4 (-6.6-20)	60 (-59-230)
Lebanon	1995	Male	85 to 89	22% (13-33%)	78% (67-87%)	0.83 (0.42-1.4)	0.21% (-0.92-1.6%)	0.089% (-1.3-1.8%)	0.48 (-6.9-9.5)	10 (-43-76)
Lebanon	1995	Male	90 to 94	23% (14-35%)	77% (65-86%)	0.75 (0.37-1.4)	-0.17% (-1.3-1.1%)	-0.37% (-1.7-1.2%)	-0.91 (-4.2-2.9)	-2.5 (-19-17)
Lebanon	1995	Male	95 plus	26% (13-43%)	74% (57-87%)	0.73 (0.28-1.5)	-0.41% (-1.7-1.1%)	-0.71% (-2.3-1.1%)	-0.44 (-1.4-0.67)	-1 (-4.2-2.6)
Lebanon	2000	Female	All Ages	14% (12-16%)	86% (84-88%)	0.27 (0.21-0.33)	0.42% (0.2-0.66%)	0.085% (-0.38-0.6%)	6.4 (-28-47)	1600 (770-2700)
Lebanon	2000	Female	15 to 19	14% (9.4-20%)	86% (80-91%)	0.25 (0.14-0.41)	0.56% (0.35-0.83%)	0.68% (0.3-1.2%)	0.45 (0.2-0.8)	110 (64-170)
Lebanon	2000	Female	20 to 24	18% (12-24%)	82% (76-88%)	0.39 (0.24-0.6)	1.1% (0.71-1.6%)	1.2% (0.57-1.9%)	0.89 (0.42-1.5)	230 (140-360)
Lebanon	2000	Female	25 to 29	17% (12-24%)	83% (76-88%)	0.41 (0.26-0.63)	1.2% (0.81-1.6%)	1.3% (0.68-2.1%)	1.1 (0.56-1.9)	260 (160-370)
Lebanon	2000	Female	30 to 34	16% (11-24%)	84% (76-89%)	0.39 (0.24-0.6)	1.1% (0.72-1.5%)	1.4% (0.73-2.3%)	1.5 (0.74-2.5)	250 (160-370)
Lebanon	2000	Female	35 to 39	16% (11-23%)	84% (77-89%)	0.4 (0.24-0.61)	0.97% (0.64-1.4%)	1.5% (0.71-2.5%)	2.1 (1-3.8)	240 (150-350)
Lebanon	2000	Female	40 to 44	16% (11-23%)	84% (77-89%)	0.41 (0.25-0.62)	0.89% (0.52-1.4%)	1.5% (0.67-2.6%)	2.7 (1.1-4.7)	210 (120-330)
Lebanon	2000	Female	45 to 49	15% (9.7-21%)	85% (79-90%)	0.42 (0.24-0.65)	0.8% (0.4-1.3%)	1.4% (0.5-2.4%)	2.9 (1.1-5.3)	170 (85-280)
Lebanon	2000	Female	50 to 54	15% (11-22%)	85% (78-89%)	0.42 (0.25-0.66)	0.7% (0.22-1.3%)	1.1% (0.3-2.2%)	3 (0.79-5.9)	140 (43-250)
Lebanon	2000	Female	55 to 59	16% (10-25%)	84% (75-90%)	0.41 (0.23-0.65)	0.51% (-0.033-1.1%)	0.82% (-0.069-1.8%)	3.4 (-0.3-7.3)	120 (-8.1-260)
Lebanon	2000	Female	60 to 64	16% (9.7-25%)	84% (75-90%)	0.38 (0.21-0.64)	0.22% (-0.34-0.82%)	0.34% (-0.47-1.3%)	2 (-2.7-7.5)	58 (-88-210)
Lebanon	2000	Female	65 to 69	16% (10-24%)	84% (76-90%)	0.34 (0.19-0.55)	0.01% (-0.54-0.57%)	0.046% (-0.73-0.84%)	0.36 (-5.8-6.8)	2.9 (-150-160)
Lebanon	2000	Female	70 to 74	16% (10-24%)	84% (76-90%)	0.3 (0.15-0.53)	-0.17% (-0.75-0.39%)	-0.21% (-1-0.56%)	-2.1 (-9.9-5.5)	-45 (-200-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	2000	Female	75 to 79	15% (9.2-23%)	85% (77-91%)	0.26 (0.12-0.46)	-0.27% (-0.81-0.28%)	-0.34% (-1.1-0.39%)	-3.3 (-11-3.7)	-51 (-160-56)
Lebanon	2000	Female	80 to 84	13% (8.6-21%)	87% (79-91%)	0.23 (0.11-0.41)	-0.29% (-0.79-0.18%)	-0.36% (-1-0.23%)	-2.9 (-8.3-1.9)	-32 (-91-20)
Lebanon	2000	Female	85 to 89	13% (8-19%)	87% (81-92%)	0.21 (0.1-0.39)	-0.34% (-0.82-0.095%)	-0.44% (-1-0.1%)	-2.8 (-7-0.67)	-21 (-52-5.9)
Lebanon	2000	Female	90 to 94	14% (8.7-21%)	86% (79-91%)	0.22 (0.094-0.43)	-0.39% (-0.93-0.063%)	-0.51% (-1.2-0.049%)	-2.1 (-5-0.22)	-10 (-25-1.6)
Lebanon	2000	Female	95 plus	16% (9.1-26%)	84% (74-91%)	0.24 (0.079-0.51)	-0.46% (-1.2-0.13%)	-0.61% (-1.5-0.11%)	-0.88 (-2.2-0.17)	-2.7 (-7.3-0.73)
Lebanon	2000	Male	All Ages	24% (21-28%)	76% (72-79%)	1.1 (0.94-1.3)	2.2% (1.5-3%)	2.4% (1-4%)	180 (74-300)	8500 (5600-12000)
Lebanon	2000	Male	15 to 19	18% (12-26%)	82% (74-88%)	0.52 (0.27-0.88)	1.4% (0.91-2.2%)	2.3% (1.2-4%)	2.5 (1.3-4.4)	290 (180-450)
Lebanon	2000	Male	20 to 24	29% (20-39%)	71% (61-80%)	1.2 (0.7-1.9)	3.4% (2.2-5%)	5.4% (3.1-8.5%)	8.2 (4.6-14)	800 (510-1200)
Lebanon	2000	Male	25 to 29	34% (24-43%)	66% (57-76%)	1.7 (1.1-2.5)	4% (2.6-5.7%)	6.9% (4-10%)	11 (6.1-17)	920 (590-1300)
Lebanon	2000	Male	30 to 34	35% (25-47%)	65% (53-75%)	1.9 (1.1-2.7)	3.5% (2.2-5.3%)	6.5% (3.8-10%)	9.8 (5.4-16)	790 (480-1200)
Lebanon	2000	Male	35 to 39	36% (26-48%)	64% (52-74%)	2 (1.2-2.9)	3.2% (1.9-4.9%)	6% (3-9.5%)	9.7 (4.8-16)	710 (400-1100)
Lebanon	2000	Male	40 to 44	35% (26-46%)	65% (54-74%)	2 (1.2-3)	3% (1.6-4.8%)	5.2% (2.2-8.9%)	9.6 (4-16)	620 (320-970)
Lebanon	2000	Male	45 to 49	32% (21-42%)	68% (58-79%)	1.9 (1.1-3)	3% (1.4-5%)	4.8% (1.9-8.7%)	11 (3.9-19)	590 (280-990)
Lebanon	2000	Male	50 to 54	30% (21-40%)	70% (60-79%)	2 (1.2-3.2)	3.4% (1.5-5.4%)	5% (1.9-8.5%)	13 (5-23)	620 (260-1000)
Lebanon	2000	Male	55 to 59	31% (19-44%)	69% (56-81%)	2.4 (1.3-3.8)	4.1% (1.8-6.7%)	5.9% (2.3-9.8%)	24 (9.5-41)	920 (420-1600)
Lebanon	2000	Male	60 to 64	30% (18-45%)	70% (55-82%)	2.3 (1.2-3.8)	3.8% (1.3-6.7%)	5.1% (1.5-8.8%)	31 (9.9-57)	990 (360-1700)
Lebanon	2000	Male	65 to 69	28% (18-42%)	72% (58-82%)	1.8 (0.98-2.9)	2.7% (0.62-5.1%)	3.3% (0.5-6.7%)	27 (4.1-54)	710 (170-1400)
Lebanon	2000	Male	70 to 74	27% (17-40%)	73% (60-83%)	1.4 (0.69-2.2)	1.5% (-0.16-3.6%)	1.7% (-0.46-4.5%)	17 (-4.7-45)	370 (-42-920)
Lebanon	2000	Male	75 to 79	25% (15-38%)	75% (62-85%)	1 (0.5-1.8)	0.64% (-0.71-2.5%)	0.62% (-1.1-3%)	6.1 (-11-29)	120 (-130-480)
Lebanon	2000	Male	80 to 84	22% (13-35%)	78% (65-87%)	0.84 (0.42-1.5)	0.23% (-0.82-1.7%)	0.12% (-1.2-1.9%)	0.86 (-8.9-14)	22 (-82-170)
Lebanon	2000	Male	85 to 89	21% (13-32%)	79% (68-87%)	0.7 (0.34-1.2)	-0.09% (-1-1.1%)	-0.26% (-1.4-1.1%)	-1.4 (-7.3-5.9)	-4.4 (-49-52)
Lebanon	2000	Male	90 to 94	22% (14-35%)	78% (65-86%)	0.64 (0.31-1.2)	-0.36% (-1.4-0.72%)	-0.59% (-1.9-0.7%)	-1.7 (-5.4-2.1)	-6.5 (-25-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	2000	Male	95 plus	26% (13-43%)	74% (57-87%)	0.62 (0.24-1.3)	-0.55% (-1.8-0.62%)	-0.88% (-2.5-0.51%)	-0.7 (-2-0.42)	-1.8 (-5.9-2)
Lebanon	2005	Female	All Ages	13% (11-16%)	87% (84-89%)	0.23 (0.18-0.29)	0.37% (0.17-0.59%)	-0.0017% (-0.43-0.45%)	-0.11 (-33-35)	1500 (710-2500)
Lebanon	2005	Female	15 to 19	13% (9.3-19%)	87% (81-91%)	0.22 (0.12-0.35)	0.54% (0.33-0.82%)	0.58% (0.26-1%)	0.44 (0.2-0.76)	130 (76-210)
Lebanon	2005	Female	20 to 24	17% (12-24%)	83% (76-88%)	0.34 (0.21-0.51)	1.1% (0.7-1.6%)	0.99% (0.47-1.7%)	0.84 (0.39-1.5)	290 (170-440)
Lebanon	2005	Female	25 to 29	17% (12-24%)	83% (76-88%)	0.36 (0.22-0.55)	1.1% (0.75-1.5%)	1.1% (0.57-1.8%)	0.91 (0.47-1.5)	270 (170-400)
Lebanon	2005	Female	30 to 34	16% (11-23%)	84% (77-89%)	0.34 (0.21-0.53)	1% (0.68-1.4%)	1.2% (0.63-2%)	1 (0.52-1.7)	220 (140-320)
Lebanon	2005	Female	35 to 39	16% (11-23%)	84% (77-89%)	0.34 (0.21-0.52)	0.87% (0.59-1.2%)	1.3% (0.68-2.2%)	1.4 (0.72-2.5)	200 (130-290)
Lebanon	2005	Female	40 to 44	16% (11-22%)	84% (78-89%)	0.35 (0.22-0.51)	0.75% (0.47-1.1%)	1.3% (0.62-2.2%)	1.9 (0.94-3.3)	180 (110-270)
Lebanon	2005	Female	45 to 49	15% (9.3-21%)	85% (79-91%)	0.36 (0.21-0.56)	0.68% (0.37-1.1%)	1.2% (0.46-2.1%)	2.4 (1-4.4)	160 (88-260)
Lebanon	2005	Female	50 to 54	15% (10-21%)	85% (79-90%)	0.36 (0.22-0.58)	0.58% (0.22-1%)	1% (0.29-1.9%)	3 (0.86-5.7)	150 (59-270)
Lebanon	2005	Female	55 to 59	16% (9.8-24%)	84% (76-90%)	0.36 (0.2-0.56)	0.42% (-0.012-0.86%)	0.72% (0.03-1.5%)	2.8 (0.12-5.7)	100 (-3.4-220)
Lebanon	2005	Female	60 to 64	16% (9.4-25%)	84% (75-91%)	0.33 (0.18-0.57)	0.17% (-0.3-0.66%)	0.28% (-0.38-1%)	1.5 (-2.1-5.8)	43 (-75-170)
Lebanon	2005	Female	65 to 69	16% (10-24%)	84% (76-90%)	0.3 (0.16-0.49)	-0.022% (-0.49-0.41%)	0.01% (-0.69-0.69%)	0.086 (-5.6-5.5)	-6.8 (-150-130)
Lebanon	2005	Female	70 to 74	15% (10-23%)	85% (77-90%)	0.26 (0.13-0.45)	-0.18% (-0.68-0.26%)	-0.22% (-0.92-0.4%)	-2.6 (-10-4.5)	-59 (-230-83)
Lebanon	2005	Female	75 to 79	14% (9-22%)	86% (78-91%)	0.22 (0.1-0.41)	-0.27% (-0.75-0.15%)	-0.34% (-1-0.23%)	-3.4 (-9.8-2.3)	-54 (-150-29)
Lebanon	2005	Female	80 to 84	13% (8.4-21%)	87% (79-92%)	0.2 (0.07-0.35)	-0.3% (-0.76-0.068%)	-0.38% (-0.96-0.099%)	-3.7 (-9.4-1)	-41 (-100-9.7)
Lebanon	2005	Female	85 to 89	13% (7.8-19%)	87% (81-92%)	0.18 (0.088-0.34)	-0.36% (-0.8--0.0066%)	-0.45% (-1--0.01%)	-3.5 (-7.8--0.079)	-26 (-58--0.48)
Lebanon	2005	Female	90 to 94	13% (8.5-20%)	87% (80-91%)	0.18 (0.082-0.36)	-0.41% (-0.94--0.036%)	-0.53% (-1.2--0.059%)	-2.2 (-4.7--0.25)	-11 (-24--0.95)
Lebanon	2005	Female	95 plus	15% (8.7-25%)	85% (75-91%)	0.2 (0.07-0.43)	-0.48% (-1.1--0.0035%)	-0.64% (-1.5--0.049%)	-1.1 (-2.6--0.08)	-3.4 (-8.3--0.024)
Lebanon	2005	Male	All Ages	25% (21-28%)	75% (72-79%)	1 (0.87-1.2)	1.8% (1.2-2.6%)	1.7% (0.54-3.2%)	150 (47-280)	8000 (5100-11000)
Lebanon	2005	Male	15 to 19	18% (12-26%)	82% (74-88%)	0.45 (0.23-0.76)	1.3% (0.83-1.9%)	2.1% (1.1-3.7%)	2.2 (1.1-3.9)	280 (180-440)
Lebanon	2005	Male	20 to 24	28% (19-39%)	72% (61-81%)	1.1 (0.6-1.7)	3% (2-4.5%)	4.8% (2.6-7.9%)	6.7 (3.6-11)	740 (470-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	2005	Male	25 to 29	33% (24-43%)	67% (57-76%)	1.5 (0.91-2.2)	3.4% (2.3-4.8%)	6.1% (3.5-9.5%)	8.1 (4.5-13)	790 (500-1200)
Lebanon	2005	Male	30 to 34	34% (24-46%)	66% (54-76%)	1.6 (1-2.4)	3% (2-4.3%)	5.6% (3.2-9%)	7.3 (4-12)	660 (420-980)
Lebanon	2005	Male	35 to 39	36% (26-47%)	64% (53-74%)	1.7 (1.1-2.5)	2.7% (1.6-4.1%)	5% (2.6-8.1%)	8.6 (4.3-15)	710 (410-1000)
Lebanon	2005	Male	40 to 44	34% (25-45%)	66% (55-75%)	1.7 (1.1-2.5)	2.4% (1.3-3.7%)	4.2% (1.8-7.1%)	11 (4.3-18)	770 (420-1200)
Lebanon	2005	Male	45 to 49	31% (21-42%)	69% (58-79%)	1.6 (0.93-2.5)	2.3% (1.2-3.9%)	3.8% (1.5-6.9%)	11 (3.9-20)	660 (330-1100)
Lebanon	2005	Male	50 to 54	29% (20-40%)	71% (60-80%)	1.7 (1-2.7)	2.5% (1.1-4.4%)	3.9% (1.3-7.1%)	13 (4.1-24)	620 (250-1100)
Lebanon	2005	Male	55 to 59	30% (19-44%)	70% (56-81%)	2 (1.1-3.3)	3.1% (1.2-5.4%)	4.6% (1.6-8.2%)	22 (7.8-40)	870 (360-1500)
Lebanon	2005	Male	60 to 64	29% (17-44%)	71% (56-83%)	1.9 (1-3.3)	2.9% (0.84-5.5%)	3.9% (0.84-7.6%)	29 (6.3-57)	950 (270-1800)
Lebanon	2005	Male	65 to 69	28% (17-41%)	72% (59-83%)	1.6 (0.84-2.6)	1.9% (0.29-4%)	2.3% (0.089-5.2%)	22 (0.91-50)	620 (93-1300)
Lebanon	2005	Male	70 to 74	26% (17-39%)	74% (61-83%)	1.2 (0.62-2)	0.95% (-0.3-2.8%)	1% (-0.6-3.5%)	11 (-6.4-39)	280 (-86-810)
Lebanon	2005	Male	75 to 79	24% (15-37%)	76% (63-85%)	0.89 (0.43-1.6)	0.34% (-0.75-1.8%)	0.26% (-1.1-2.2%)	2.8 (-13-24)	72 (-160-400)
Lebanon	2005	Male	80 to 84	22% (13-34%)	78% (66-87%)	0.72 (0.35-1.2)	0.0089% (-0.92-1%)	-0.13% (-1.3-1.2%)	-1.3 (-13-12)	1.2 (-120-140)
Lebanon	2005	Male	85 to 89	20% (12-31%)	80% (69-88%)	0.59 (0.3-1)	-0.23% (-1-0.62%)	-0.41% (-1.4-0.62%)	-2.9 (-9.8-4.5)	-15 (-66-41)
Lebanon	2005	Male	90 to 94	21% (13-33%)	79% (67-87%)	0.53 (0.26-0.98)	-0.45% (-1.3-0.33%)	-0.68% (-1.8-0.3%)	-2.4 (-6.1-1.1)	-9.7 (-28-7.1)
Lebanon	2005	Male	95 plus	25% (13-41%)	75% (59-87%)	0.51 (0.19-1.1)	-0.61% (-1.7-0.27%)	-0.93% (-2.3-0.13%)	-1 (-2.7-0.15)	-2.8 (-7.8-1.3)
Lebanon	2010	Female	All Ages	14% (12-17%)	86% (83-88%)	0.27 (0.21-0.34)	0.4% (0.21-0.64%)	0.0083% (-0.44-0.49%)	0.7 (-37-40)	1800 (860-2900)
Lebanon	2010	Female	15 to 19	13% (8.6-20%)	87% (80-91%)	0.23 (0.12-0.39)	0.56% (0.34-0.86%)	0.67% (0.31-1.2%)	0.46 (0.22-0.81)	140 (80-230)
Lebanon	2010	Female	20 to 24	18% (11-26%)	82% (74-89%)	0.37 (0.21-0.6)	1.1% (0.72-1.6%)	1.1% (0.52-2%)	0.93 (0.42-1.7)	320 (200-500)
Lebanon	2010	Female	25 to 29	19% (12-27%)	81% (73-88%)	0.41 (0.24-0.63)	1.2% (0.8-1.6%)	1.3% (0.67-2.1%)	1.1 (0.55-1.9)	340 (210-500)
Lebanon	2010	Female	30 to 34	17% (11-26%)	83% (74-89%)	0.39 (0.22-0.62)	1% (0.68-1.5%)	1.4% (0.7-2.4%)	1.2 (0.6-2)	260 (170-400)
Lebanon	2010	Female	35 to 39	17% (10-25%)	83% (75-90%)	0.37 (0.21-0.58)	0.88% (0.6-1.2%)	1.5% (0.75-2.5%)	1.3 (0.68-2.4)	200 (130-290)
Lebanon	2010	Female	40 to 44	16% (10-25%)	84% (75-90%)	0.38 (0.22-0.58)	0.75% (0.45-1.1%)	1.4% (0.71-2.4%)	1.8 (0.86-3.1)	180 (100-270)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	2010	Female	45 to 49	16% (9.6-24%)	84% (76-90%)	0.41 (0.22-0.66)	0.7% (0.37-1.1%)	1.4% (0.56-2.5%)	2.4 (1-4.5)	170 (90-280)
Lebanon	2010	Female	50 to 54	16% (10-24%)	84% (76-90%)	0.42 (0.24-0.7)	0.62% (0.26-1.1%)	1.2% (0.46-2.3%)	3.1 (1.2-5.9)	160 (67-290)
Lebanon	2010	Female	55 to 59	17% (10-25%)	83% (75-90%)	0.4 (0.22-0.63)	0.46% (0.067-0.91%)	0.9% (0.14-1.7%)	3.3 (0.51-6.5)	130 (20-250)
Lebanon	2010	Female	60 to 64	17% (10-26%)	83% (74-90%)	0.36 (0.2-0.63)	0.22% (-0.2-0.72%)	0.43% (-0.29-1.3%)	2 (-1.3-6.2)	59 (-52-190)
Lebanon	2010	Female	65 to 69	17% (11-25%)	83% (75-89%)	0.33 (0.18-0.55)	0.037% (-0.39-0.48%)	0.12% (-0.56-0.83%)	0.79 (-3.6-5.4)	10 (-110-130)
Lebanon	2010	Female	70 to 74	16% (10-25%)	84% (75-90%)	0.28 (0.13-0.5)	-0.11% (-0.6-0.34%)	-0.12% (-0.83-0.51%)	-1.4 (-9.2-5.7)	-38 (-200-110)
Lebanon	2010	Female	75 to 79	15% (8.9-23%)	85% (77-91%)	0.24 (0.11-0.44)	-0.22% (-0.7-0.21%)	-0.28% (-0.95-0.3%)	-3.9 (-13-4.2)	-64 (-200-64)
Lebanon	2010	Female	80 to 84	13% (8.3-21%)	87% (79-92%)	0.21 (0.1-0.37)	-0.25% (-0.7-0.12%)	-0.32% (-0.9-0.17%)	-4 (-11-2.3)	-46 (-120-22)
Lebanon	2010	Female	85 to 89	13% (7.6-20%)	87% (80-92%)	0.2 (0.091-0.37)	-0.32% (-0.76-0.07%)	-0.42% (-0.97-0.052%)	-4.4 (-11-0.57)	-32 (-78-7)
Lebanon	2010	Female	90 to 94	14% (8.6-22%)	86% (78-91%)	0.2 (0.088-0.39)	-0.38% (-0.89-0.037%)	-0.5% (-1.2-0.0094%)	-2.9 (-6.6-0.053)	-14 (-32-1.4)
Lebanon	2010	Female	95 plus	16% (8.6-27%)	84% (73-91%)	0.22 (0.072-0.48)	-0.45% (-1.2-0.064%)	-0.62% (-1.5-0.025%)	-1.2 (-3-0.05)	-3.7 (-9.7-0.53)
Lebanon	2010	Male	All Ages	24% (20-28%)	76% (72-80%)	1.1 (0.91-1.3)	2% (1.3-2.7%)	1.8% (0.68-3.2%)	170 (65-310)	9300 (5900-13000)
Lebanon	2010	Male	15 to 19	16% (10-25%)	84% (75-90%)	0.44 (0.21-0.76)	1.3% (0.83-2%)	2.1% (1-3.7%)	2.2 (1.1-4)	310 (190-480)
Lebanon	2010	Male	20 to 24	26% (16-37%)	74% (63-84%)	1 (0.53-1.7)	3.1% (2.1-4.5%)	4.9% (2.6-8.1%)	6.7 (3.6-12)	790 (510-1200)
Lebanon	2010	Male	25 to 29	31% (21-42%)	69% (58-79%)	1.4 (0.85-2.3)	3.5% (2.4-5.1%)	6.4% (3.4-10%)	8.8 (4.6-14)	890 (570-1300)
Lebanon	2010	Male	30 to 34	33% (21-45%)	67% (55-79%)	1.6 (0.96-2.6)	3.2% (2.1-4.8%)	6.2% (3.4-10%)	8.3 (4.4-14)	770 (480-1100)
Lebanon	2010	Male	35 to 39	34% (22-47%)	66% (53-78%)	1.7 (1-2.7)	2.9% (1.8-4.3%)	5.7% (2.9-9%)	8.5 (4.2-14)	710 (420-1000)
Lebanon	2010	Male	40 to 44	34% (23-46%)	66% (54-77%)	1.8 (1-2.7)	2.6% (1.5-4.1%)	4.7% (2.2-7.9%)	11 (4.9-19)	830 (460-1300)
Lebanon	2010	Male	45 to 49	31% (19-43%)	69% (57-81%)	1.7 (0.95-2.7)	2.5% (1.3-4%)	4.3% (1.7-7.5%)	15 (6.1-28)	950 (460-1600)
Lebanon	2010	Male	50 to 54	29% (19-41%)	71% (59-81%)	1.8 (1.1-2.8)	2.7% (1.3-4.4%)	4.4% (1.8-7.5%)	18 (7-32)	900 (430-1500)
Lebanon	2010	Male	55 to 59	28% (18-42%)	72% (58-82%)	2 (1.1-3.3)	3.1% (1.4-5.2%)	4.8% (1.9-8.1%)	23 (8.8-39)	930 (430-1500)
Lebanon	2010	Male	60 to 64	26% (15-41%)	74% (59-85%)	1.9 (0.94-3.3)	2.9% (0.97-5.2%)	4% (1.1-7.4%)	29 (8.1-55)	990 (330-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	2010	Male	65 to 69	25% (16-38%)	75% (62-84%)	1.5 (0.8-2.5)	2% (0.46-4%)	2.6% (0.36-5.4%)	26 (3.5-55)	730 (170-1500)
Lebanon	2010	Male	70 to 74	24% (15-36%)	76% (64-85%)	1.1 (0.58-1.9)	1.1% (-0.073-2.8%)	1.3% (-0.3-3.6%)	16 (-3.8-44)	380 (-24-960)
Lebanon	2010	Male	75 to 79	22% (14-34%)	78% (66-86%)	0.87 (0.43-1.5)	0.51% (-0.52-1.9%)	0.49% (-0.9-2.3%)	6.3 (-12-30)	130 (-140-510)
Lebanon	2010	Male	80 to 84	20% (12-31%)	80% (69-88%)	0.71 (0.36-1.2)	0.17% (-0.64-1.3%)	0.063% (-1-1.4%)	0.74 (-12-16)	28 (-100-210)
Lebanon	2010	Male	85 to 89	19% (11-29%)	81% (71-89%)	0.59 (0.29-1)	-0.087% (-0.8-0.76%)	-0.25% (-1.2-0.79%)	-2.5 (-12-8.5)	-8.2 (-77-75)
Lebanon	2010	Male	90 to 94	20% (12-30%)	80% (70-88%)	0.53 (0.26-0.98)	-0.31% (-1.1-0.49%)	-0.52% (-1.5-0.45%)	-2.7 (-7.9-2.4)	-10 (-35-16)
Lebanon	2010	Male	95 plus	23% (12-39%)	77% (61-88%)	0.52 (0.19-1.1)	-0.46% (-1.4-0.42%)	-0.77% (-2-0.28%)	-1.2 (-3.1-0.41)	-2.9 (-9.2-2.6)
Lebanon	2016	Female	All Ages	14% (12-17%)	86% (83-88%)	0.3 (0.19-0.44)	0.48% (0.24-0.8%)	0.047% (-0.43-0.68%)	5.6 (-50-81)	2800 (1400-4900)
Lebanon	2016	Female	15 to 19	13% (8.5-20%)	87% (80-91%)	0.25 (0.11-0.45)	0.56% (0.34-0.86%)	0.57% (0.24-1.1%)	0.51 (0.22-0.97)	180 (100-280)
Lebanon	2016	Female	20 to 24	18% (11-26%)	82% (74-89%)	0.4 (0.19-0.71)	1.2% (0.74-1.7%)	1% (0.43-1.9%)	1.2 (0.5-2.3)	450 (260-690)
Lebanon	2016	Female	25 to 29	18% (12-26%)	82% (74-88%)	0.44 (0.22-0.77)	1.2% (0.83-1.7%)	1.2% (0.53-2.2%)	1.6 (0.69-2.8)	500 (320-750)
Lebanon	2016	Female	30 to 34	17% (11-26%)	83% (74-89%)	0.42 (0.22-0.72)	1.1% (0.72-1.6%)	1.4% (0.66-2.5%)	2 (0.89-3.6)	460 (280-700)
Lebanon	2016	Female	35 to 39	17% (9.8-25%)	83% (75-90%)	0.4 (0.21-0.7)	0.93% (0.62-1.4%)	1.5% (0.67-2.7%)	2.2 (1-4.1)	350 (220-520)
Lebanon	2016	Female	40 to 44	16% (10-24%)	84% (76-90%)	0.41 (0.21-0.69)	0.81% (0.47-1.3%)	1.5% (0.65-2.6%)	2.7 (1.1-4.9)	270 (160-430)
Lebanon	2016	Female	45 to 49	16% (9.4-23%)	84% (77-91%)	0.44 (0.21-0.78)	0.76% (0.37-1.3%)	1.5% (0.52-2.8%)	3.5 (1.2-6.9)	260 (120-440)
Lebanon	2016	Female	50 to 54	16% (10-23%)	84% (77-90%)	0.45 (0.22-0.81)	0.69% (0.27-1.3%)	1.3% (0.41-2.7%)	4.3 (1.4-8.8)	230 (89-460)
Lebanon	2016	Female	55 to 59	16% (10-25%)	84% (75-90%)	0.42 (0.21-0.73)	0.53% (0.11-1.1%)	1% (0.21-2.2%)	4.4 (0.96-9.7)	180 (36-400)
Lebanon	2016	Female	60 to 64	16% (9.9-26%)	84% (74-90%)	0.39 (0.19-0.71)	0.3% (-0.16-0.95%)	0.56% (-0.23-1.6%)	3.5 (-1.3-10)	110 (-62-350)
Lebanon	2016	Female	65 to 69	17% (11-24%)	83% (76-89%)	0.35 (0.18-0.63)	0.11% (-0.33-0.7%)	0.22% (-0.49-1.1%)	1.8 (-4.1-9.3)	39 (-130-260)
Lebanon	2016	Female	70 to 74	16% (10-24%)	84% (76-90%)	0.31 (0.14-0.59)	-0.041% (-0.51-0.5%)	-0.029% (-0.76-0.78%)	-0.33 (-8.5-8.5)	-14 (-180-180)
Lebanon	2016	Female	75 to 79	15% (9-23%)	85% (77-91%)	0.26 (0.11-0.5)	-0.15% (-0.63-0.37%)	-0.2% (-0.86-0.49%)	-3.5 (-15-8.5)	-58 (-230-140)
Lebanon	2016	Female	80 to 84	13% (8.1-21%)	87% (79-92%)	0.23 (0.099-0.44)	-0.19% (-0.62-0.25%)	-0.25% (-0.82-0.31%)	-5.3 (-18-6.8)	-59 (-200-84)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lebanon	2016	Female	85 to 89	13% (7.6-20%)	87% (80-92%)	0.22 (0.09-0.43)	-0.27% (-0.74-0.17%)	-0.36% (-0.97-0.2%)	-6.3 (-17-3.5)	-46 (-130-29)
Lebanon	2016	Female	90 to 94	14% (8.8-21%)	86% (79-91%)	0.22 (0.09-0.46)	-0.34% (-0.83-0.17%)	-0.46% (-1.1-0.15%)	-4.4 (-10-1.5)	-21 (-51-10)
Lebanon	2016	Female	95 plus	16% (8.7-27%)	84% (73-91%)	0.24 (0.08-0.54)	-0.41% (-1.1-0.22%)	-0.57% (-1.4-0.19%)	-2.2 (-5.5-0.7)	-6.6 (-18-3.5)
Lebanon	2016	Male	All Ages	24% (20-28%)	76% (72-80%)	1.2 (0.79-1.7)	2% (1.1-3.2%)	1.8% (0.39-3.7%)	260 (53-510)	13000 (7000-20000)
Lebanon	2016	Male	15 to 19	16% (10-25%)	84% (75-90%)	0.47 (0.2-0.92)	1.2% (0.72-1.9%)	1.7% (0.74-3.3%)	2.6 (1.2-5.2)	380 (220-620)
Lebanon	2016	Male	20 to 24	26% (16-37%)	74% (63-84%)	1.1 (0.5-1.9)	2.7% (1.7-4.2%)	3.6% (1.7-6.5%)	9.1 (4.3-17)	1100 (660-1700)
Lebanon	2016	Male	25 to 29	31% (21-42%)	69% (58-79%)	1.5 (0.8-2.6)	3.2% (2.1-4.8%)	5% (2.4-8.8%)	12 (5.7-21)	1200 (770-1900)
Lebanon	2016	Male	30 to 34	33% (22-46%)	67% (54-78%)	1.7 (0.93-3)	3.1% (1.9-4.6%)	5.2% (2.5-9.2%)	12 (5.6-20)	1100 (680-1800)
Lebanon	2016	Male	35 to 39	34% (22-47%)	66% (53-78%)	1.9 (0.95-3.2)	2.9% (1.7-4.6%)	5.2% (2.2-9.2%)	12 (4.9-21)	1000 (550-1600)
Lebanon	2016	Male	40 to 44	34% (23-46%)	66% (54-77%)	1.9 (0.94-3.3)	2.8% (1.3-4.5%)	4.8% (1.7-8.6%)	13 (4.3-23)	940 (450-1500)
Lebanon	2016	Male	45 to 49	31% (19-43%)	69% (57-81%)	1.9 (0.85-3.4)	2.7% (1.1-4.6%)	4.5% (1.3-8.2%)	18 (5.6-35)	1100 (460-1900)
Lebanon	2016	Male	50 to 54	29% (19-40%)	71% (60-81%)	1.9 (0.98-3.4)	2.8% (1.1-4.9%)	4.5% (1.3-8%)	30 (9.1-55)	1500 (580-2600)
Lebanon	2016	Male	55 to 59	28% (17-41%)	72% (59-83%)	2.2 (1.1-3.9)	3.2% (1.4-5.4%)	4.8% (1.8-8.4%)	37 (14-67)	1500 (660-2700)
Lebanon	2016	Male	60 to 64	26% (15-41%)	74% (59-85%)	2 (0.94-3.8)	3% (1-5.4%)	4.1% (1.2-7.7%)	38 (10-74)	1300 (420-2400)
Lebanon	2016	Male	65 to 69	25% (15-38%)	75% (62-85%)	1.6 (0.78-2.9)	2.2% (0.4-4.4%)	2.8% (0.29-6.1%)	35 (3.3-75)	970 (170-2000)
Lebanon	2016	Male	70 to 74	24% (15-36%)	76% (64-85%)	1.2 (0.56-2.2)	1.3% (-0.096-3.4%)	1.6% (-0.41-4.3%)	26 (-6.3-72)	610 (-44-1500)
Lebanon	2016	Male	75 to 79	22% (14-34%)	78% (66-86%)	0.93 (0.41-1.7)	0.71% (-0.51-2.5%)	0.73% (-0.86-2.9%)	15 (-17-61)	280 (-210-1000)
Lebanon	2016	Male	80 to 84	20% (12-31%)	80% (69-88%)	0.75 (0.34-1.4)	0.34% (-0.64-1.8%)	0.27% (-1-2%)	4.9 (-18-38)	85 (-160-470)
Lebanon	2016	Male	85 to 89	19% (11-29%)	81% (71-89%)	0.62 (0.27-1.2)	0.051% (-0.77-1.2%)	-0.078% (-1.1-1.3%)	-1.2 (-17-21)	7.2 (-110-180)
Lebanon	2016	Male	90 to 94	20% (12-31%)	80% (69-88%)	0.57 (0.25-1.1)	-0.21% (-1-0.82%)	-0.4% (-1.4-0.82%)	-3.6 (-13-7.6)	-12 (-60-45)
Lebanon	2016	Male	95 plus	23% (12-38%)	77% (62-88%)	0.57 (0.2-1.2)	-0.39% (-1.4-0.73%)	-0.68% (-1.9-0.67%)	-2 (-5.7-2)	-4.7 (-17-8.8)
Libya	1990	Female	All Ages	11% (6.2-16%)	89% (84-94%)	0.0087 (0.0047-0.014)	0.19% (0.13-0.25%)	0.13% (0.0085-0.22%)	9.4 (0.59-16)	1000 (660-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	1990	Female	15 to 19	9.4% (3-20%)	91% (80-97%)	0.0071 (0.0018-0.017)	0.32% (0.18-0.49%)	0.15% (0.083-0.27%)	0.2 (0.1-0.36)	100 (55-170)
Libya	1990	Female	20 to 24	12% (3.6-26%)	88% (74-96%)	0.011 (0.0032-0.027)	0.64% (0.41-0.92%)	0.19% (0.11-0.32%)	0.27 (0.15-0.46)	190 (110-300)
Libya	1990	Female	25 to 29	13% (4-29%)	87% (71-96%)	0.014 (0.0038-0.032)	0.69% (0.48-0.95%)	0.31% (0.17-0.55%)	0.39 (0.2-0.67)	170 (110-250)
Libya	1990	Female	30 to 34	15% (4.1-32%)	85% (68-96%)	0.016 (0.0046-0.035)	0.61% (0.42-0.88%)	0.33% (0.18-0.54%)	0.42 (0.23-0.7)	140 (85-200)
Libya	1990	Female	35 to 39	17% (4.4-33%)	83% (67-96%)	0.019 (0.0052-0.04)	0.46% (0.31-0.63%)	0.27% (0.17-0.39%)	0.39 (0.23-0.59)	96 (62-140)
Libya	1990	Female	40 to 44	18% (6-33%)	82% (67-94%)	0.021 (0.0067-0.046)	0.35% (0.22-0.5%)	0.29% (0.15-0.42%)	0.45 (0.24-0.7)	66 (40-97)
Libya	1990	Female	45 to 49	18% (6.3-34%)	82% (66-94%)	0.023 (0.0066-0.052)	0.27% (0.14-0.4%)	0.24% (0.054-0.38%)	0.49 (0.11-0.81)	52 (26-80)
Libya	1990	Female	50 to 54	19% (7.3-37%)	81% (63-93%)	0.023 (0.0066-0.053)	0.23% (0.06-0.36%)	0.26% (0.032-0.43%)	0.69 (0.081-1.2)	46 (12-76)
Libya	1990	Female	55 to 59	20% (7.7-39%)	80% (61-92%)	0.024 (0.0059-0.058)	0.19% (-0.025-0.35%)	0.25% (-0.048-0.47%)	0.82 (-0.16-1.6)	37 (-5-70)
Libya	1990	Female	60 to 64	21% (8.1-40%)	79% (60-92%)	0.023 (0.0055-0.056)	0.16% (-0.12-0.35%)	0.22% (-0.13-0.49%)	0.96 (-0.55-2.1)	32 (-23-68)
Libya	1990	Female	65 to 69	22% (9.2-39%)	78% (61-91%)	0.02 (0.0054-0.048)	0.15% (-0.1-0.33%)	0.21% (-0.13-0.48%)	1.1 (-0.66-2.5)	28 (-19-63)
Libya	1990	Female	70 to 74	21% (9.2-38%)	79% (62-91%)	0.016 (0.0041-0.043)	0.16% (-0.11-0.33%)	0.21% (-0.12-0.45%)	1.2 (-0.67-2.7)	24 (-17-51)
Libya	1990	Female	75 to 79	19% (7.3-38%)	81% (62-93%)	0.013 (0.0026-0.04)	0.13% (-0.15-0.28%)	0.17% (-0.2-0.38%)	0.99 (-1.1-2.2)	15 (-17-31)
Libya	1990	Female	80 to 84	17% (5.7-38%)	83% (62-94%)	0.012 (0.0024-0.034)	0.11% (-0.14-0.24%)	0.14% (-0.16-0.3%)	0.56 (-0.68-1.2)	6.3 (-7.2-13)
Libya	1990	Female	85 to 89	17% (5.8-38%)	83% (62-94%)	0.012 (0.0024-0.038)	0.084% (-0.17-0.19%)	0.097% (-0.21-0.24%)	0.32 (-0.66-0.78)	2.6 (-5.1-6)
Libya	1990	Female	90 to 94	20% (6.4-42%)	80% (58-94%)	0.013 (0.0025-0.04)	0.064% (-0.2-0.17%)	0.068% (-0.25-0.19%)	0.12 (-0.45-0.35)	0.72 (-2.2-1.9)
Libya	1990	Female	95 plus	24% (7.1-49%)	76% (51-93%)	0.016 (0.0026-0.05)	0.048% (-0.3-0.17%)	0.04% (-0.37-0.19%)	0.025 (-0.24-0.12)	0.12 (-0.77-0.45)
Libya	1990	Male	All Ages	8.5% (5.8-12%)	91% (88-94%)	0.011 (0.0057-0.018)	0.37% (0.29-0.46%)	0.29% (0.2-0.37%)	27 (19-35)	2300 (1700-3000)
Libya	1990	Male	15 to 19	6.6% (2.7-13%)	93% (87-97%)	0.0048 (0.0013-0.012)	0.47% (0.29-0.7%)	0.13% (0.072-0.24%)	0.28 (0.15-0.53)	170 (95-270)
Libya	1990	Male	20 to 24	10% (4.2-20%)	90% (80-96%)	0.011 (0.0035-0.025)	0.96% (0.6-1.4%)	0.22% (0.11-0.43%)	0.57 (0.28-1.1)	330 (200-520)
Libya	1990	Male	25 to 29	13% (5-24%)	87% (76-95%)	0.017 (0.0057-0.038)	0.99% (0.69-1.4%)	0.37% (0.19-0.62%)	0.9 (0.47-1.5)	310 (200-460)
Libya	1990	Male	30 to 34	14% (5.3-28%)	86% (72-95%)	0.022 (0.008-0.046)	0.91% (0.65-1.2%)	0.49% (0.28-0.77%)	1.3 (0.74-2)	290 (190-410)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	1990	Male	35 to 39	16% (6.1-29%)	84% (71-94%)	0.025 (0.0079-0.052)	0.9% (0.65-1.2%)	0.69% (0.44-1%)	2 (1.2-3)	270 (190-370)
Libya	1990	Male	40 to 44	16% (7.1-29%)	84% (71-93%)	0.027 (0.0091-0.058)	0.81% (0.59-1.1%)	0.72% (0.45-1.1%)	2.1 (1.3-3.1)	210 (150-300)
Libya	1990	Male	45 to 49	16% (6.4-29%)	84% (71-94%)	0.028 (0.0087-0.063)	0.75% (0.55-0.98%)	0.79% (0.51-1.1%)	2.7 (1.7-4.1)	190 (140-260)
Libya	1990	Male	50 to 54	15% (6.5-29%)	85% (71-93%)	0.027 (0.0081-0.062)	0.6% (0.42-0.78%)	0.67% (0.42-0.95%)	2.7 (1.7-4.1)	150 (100-200)
Libya	1990	Male	55 to 59	15% (6.3-29%)	85% (71-94%)	0.027 (0.007-0.062)	0.47% (0.31-0.63%)	0.54% (0.31-0.77%)	2.7 (1.5-4)	120 (76-160)
Libya	1990	Male	60 to 64	14% (5.9-27%)	86% (73-94%)	0.024 (0.0057-0.059)	0.36% (0.21-0.52%)	0.42% (0.19-0.65%)	2.6 (1.2-4.2)	89 (50-130)
Libya	1990	Male	65 to 69	14% (6.2-25%)	86% (75-94%)	0.02 (0.0059-0.045)	0.32% (0.17-0.48%)	0.39% (0.19-0.6%)	2.8 (1.3-4.3)	72 (37-110)
Libya	1990	Male	70 to 74	12% (5.7-22%)	88% (78-94%)	0.015 (0.0042-0.036)	0.28% (0.15-0.41%)	0.34% (0.18-0.52%)	2.3 (1.2-3.6)	47 (25-70)
Libya	1990	Male	75 to 79	11% (4.7-20%)	89% (80-95%)	0.011 (0.0026-0.03)	0.22% (0.1-0.32%)	0.26% (0.11-0.39%)	1.6 (0.69-2.4)	25 (11-36)
Libya	1990	Male	80 to 84	9.5% (3.8-19%)	90% (81-96%)	0.0093 (0.0023-0.025)	0.2% (0.1-0.29%)	0.23% (0.1-0.35%)	0.99 (0.44-1.5)	11 (5.3-17)
Libya	1990	Male	85 to 89	9.3% (3.7-18%)	91% (82-96%)	0.0085 (0.0023-0.022)	0.17% (0.089-0.23%)	0.18% (0.086-0.27%)	0.56 (0.25-0.84)	4.6 (2.4-6.5)
Libya	1990	Male	90 to 94	10% (4.2-21%)	90% (79-96%)	0.0089 (0.0022-0.024)	0.14% (0.052-0.2%)	0.14% (0.037-0.22%)	0.21 (0.052-0.33)	1.3 (0.47-1.9)
Libya	1990	Male	95 plus	13% (4.7-27%)	87% (73-95%)	0.01 (0.0019-0.032)	0.15% (0.048-0.21%)	0.12% (-0.0061-0.21%)	0.066 (-0.0034-0.11)	0.32 (0.11-0.48)
Libya	1995	Female	All Ages	11% (6.5-17%)	89% (83-93%)	0.0091 (0.0064-0.012)	0.22% (0.16-0.29%)	0.15% (0.029-0.24%)	11 (2.3-18)	1200 (820-1700)
Libya	1995	Female	15 to 19	9.3% (3-20%)	91% (80-97%)	0.0069 (0.0019-0.016)	0.33% (0.19-0.5%)	0.14% (0.084-0.25%)	0.2 (0.11-0.35)	110 (64-180)
Libya	1995	Female	20 to 24	12% (3.4-26%)	88% (74-97%)	0.011 (0.0033-0.025)	0.65% (0.4-0.96%)	0.19% (0.11-0.3%)	0.29 (0.17-0.48)	230 (140-370)
Libya	1995	Female	25 to 29	14% (3.8-28%)	86% (72-96%)	0.014 (0.004-0.028)	0.7% (0.48-0.96%)	0.3% (0.16-0.51%)	0.45 (0.25-0.78)	230 (140-340)
Libya	1995	Female	30 to 34	15% (4-32%)	85% (68-96%)	0.016 (0.0051-0.031)	0.62% (0.42-0.85%)	0.32% (0.19-0.53%)	0.48 (0.27-0.8)	170 (110-260)
Libya	1995	Female	35 to 39	17% (4.9-32%)	83% (68-95%)	0.019 (0.0059-0.037)	0.46% (0.31-0.62%)	0.27% (0.17-0.39%)	0.44 (0.26-0.64)	110 (72-160)
Libya	1995	Female	40 to 44	18% (6.3-32%)	82% (68-94%)	0.021 (0.0075-0.041)	0.36% (0.23-0.5%)	0.29% (0.17-0.42%)	0.55 (0.31-0.84)	84 (52-120)
Libya	1995	Female	45 to 49	18% (6.2-34%)	82% (66-94%)	0.022 (0.0073-0.046)	0.27% (0.15-0.4%)	0.24% (0.064-0.38%)	0.52 (0.14-0.88)	57 (29-85)
Libya	1995	Female	50 to 54	19% (7.3-36%)	81% (64-93%)	0.023 (0.0069-0.048)	0.23% (0.062-0.37%)	0.26% (0.059-0.43%)	0.79 (0.18-1.3)	52 (13-85)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	1995	Female	55 to 59	20% (8.2-40%)	80% (60-92%)	0.024 (0.0069-0.052)	0.19% (-0.0098-0.36%)	0.25% (-0.011-0.46%)	0.96 (-0.044-1.8)	43 (-2.4-81)
Libya	1995	Female	60 to 64	21% (8.7-40%)	79% (60-91%)	0.023 (0.0063-0.05)	0.16% (-0.092-0.34%)	0.22% (-0.11-0.47%)	1.1 (-0.57-2.4)	36 (-20-79)
Libya	1995	Female	65 to 69	22% (9.2-39%)	78% (61-91%)	0.02 (0.0055-0.043)	0.15% (-0.078-0.33%)	0.22% (-0.11-0.47%)	1.3 (-0.65-2.8)	34 (-17-73)
Libya	1995	Female	70 to 74	21% (9-38%)	79% (62-91%)	0.016 (0.0042-0.037)	0.17% (-0.08-0.32%)	0.22% (-0.088-0.44%)	1.6 (-0.69-3.2)	32 (-16-62)
Libya	1995	Female	75 to 79	18% (7.1-36%)	82% (64-93%)	0.013 (0.0027-0.035)	0.14% (-0.11-0.27%)	0.18% (-0.14-0.36%)	1.3 (-0.93-2.5)	20 (-16-37)
Libya	1995	Female	80 to 84	17% (5.8-36%)	83% (64-94%)	0.011 (0.0025-0.032)	0.12% (-0.1-0.23%)	0.15% (-0.12-0.3%)	0.78 (-0.63-1.5)	8.8 (-7.2-17)
Libya	1995	Female	85 to 89	17% (6.1-39%)	83% (61-94%)	0.012 (0.0027-0.035)	0.091% (-0.14-0.19%)	0.11% (-0.17-0.24%)	0.44 (-0.67-0.98)	3.6 (-5.8-7.5)
Libya	1995	Female	90 to 94	20% (7.1-43%)	80% (57-93%)	0.013 (0.0028-0.038)	0.067% (-0.19-0.16%)	0.072% (-0.24-0.19%)	0.16 (-0.53-0.44)	0.96 (-2.6-2.3)
Libya	1995	Female	95 plus	24% (7.7-50%)	76% (50-92%)	0.016 (0.003-0.047)	0.049% (-0.24-0.16%)	0.041% (-0.32-0.18%)	0.033 (-0.25-0.15)	0.16 (-0.79-0.53)
Libya	1995	Male	All Ages	8.7% (6-12%)	91% (88-94%)	0.011 (0.0077-0.014)	0.41% (0.33-0.5%)	0.32% (0.23-0.4%)	31 (23-41)	2700 (2000-3500)
Libya	1995	Male	15 to 19	6.6% (2.7-13%)	93% (87-97%)	0.0047 (0.0014-0.01)	0.48% (0.29-0.72%)	0.13% (0.072-0.24%)	0.31 (0.17-0.56)	190 (110-300)
Libya	1995	Male	20 to 24	10% (4-20%)	90% (80-96%)	0.01 (0.0037-0.021)	0.97% (0.63-1.4%)	0.21% (0.11-0.41%)	0.65 (0.32-1.3)	400 (250-620)
Libya	1995	Male	25 to 29	13% (5-24%)	87% (76-95%)	0.016 (0.0063-0.031)	1% (0.7-1.4%)	0.36% (0.19-0.58%)	1 (0.56-1.7)	370 (240-540)
Libya	1995	Male	30 to 34	14% (5.3-28%)	86% (72-95%)	0.021 (0.0084-0.038)	0.92% (0.64-1.2%)	0.48% (0.27-0.74%)	1.3 (0.78-2.1)	310 (210-440)
Libya	1995	Male	35 to 39	16% (6.2-29%)	84% (71-94%)	0.024 (0.0089-0.047)	0.9% (0.66-1.2%)	0.7% (0.44-1.1%)	2.2 (1.4-3.4)	310 (210-420)
Libya	1995	Male	40 to 44	16% (7.4-28%)	84% (72-93%)	0.025 (0.01-0.05)	0.81% (0.59-1.1%)	0.72% (0.46-1.1%)	2.6 (1.6-3.9)	270 (190-380)
Libya	1995	Male	45 to 49	16% (6.1-29%)	84% (71-94%)	0.026 (0.0091-0.053)	0.76% (0.57-0.99%)	0.81% (0.53-1.2%)	3.1 (1.9-4.8)	220 (160-300)
Libya	1995	Male	50 to 54	15% (6.4-30%)	85% (70-94%)	0.026 (0.0085-0.056)	0.61% (0.44-0.81%)	0.69% (0.44-0.98%)	3.2 (2-4.7)	180 (120-240)
Libya	1995	Male	55 to 59	14% (6.4-28%)	86% (72-94%)	0.025 (0.0078-0.054)	0.48% (0.33-0.66%)	0.56% (0.35-0.81%)	3.1 (1.8-4.6)	130 (90-190)
Libya	1995	Male	60 to 64	14% (5.8-27%)	86% (73-94%)	0.022 (0.0066-0.05)	0.38% (0.23-0.55%)	0.44% (0.23-0.68%)	3.1 (1.5-4.7)	100 (62-150)
Libya	1995	Male	65 to 69	13% (6-24%)	87% (76-94%)	0.018 (0.0058-0.039)	0.34% (0.2-0.51%)	0.42% (0.22-0.65%)	3.3 (1.8-5.1)	85 (49-130)
Libya	1995	Male	70 to 74	12% (5.5-22%)	88% (78-95%)	0.014 (0.0041-0.033)	0.29% (0.17-0.43%)	0.36% (0.2-0.56%)	3.1 (1.7-4.9)	62 (36-93)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	1995	Male	75 to 79	11% (4.5-20%)	89% (80-95%)	0.01 (0.0027-0.027)	0.23% (0.12-0.32%)	0.27% (0.13-0.4%)	2 (0.98-2.9)	31 (16-44)
Libya	1995	Male	80 to 84	9.4% (3.8-19%)	91% (81-96%)	0.0087 (0.0024-0.022)	0.21% (0.12-0.3%)	0.24% (0.13-0.37%)	1.2 (0.62-1.8)	13 (7.6-20)
Libya	1995	Male	85 to 89	9.2% (3.7-18%)	91% (82-96%)	0.008 (0.0023-0.02)	0.17% (0.095-0.24%)	0.19% (0.095-0.28%)	0.72 (0.35-1.1)	5.9 (3.2-8.2)
Libya	1995	Male	90 to 94	10% (4.1-21%)	90% (79-96%)	0.0084 (0.0023-0.023)	0.14% (0.066-0.2%)	0.15% (0.057-0.22%)	0.27 (0.11-0.42)	1.6 (0.76-2.3)
Libya	1995	Male	95 plus	13% (4.6-27%)	87% (73-95%)	0.0096 (0.002-0.028)	0.15% (0.052-0.22%)	0.13% (0.012-0.21%)	0.074 (0.0071-0.12)	0.35 (0.12-0.51)
Libya	2000	Female	All Ages	11% (6.6-17%)	89% (83-93%)	0.012 (0.0095-0.014)	0.24% (0.16-0.31%)	0.13% (-0.014-0.24%)	12 (-1.2-22)	1400 (950-2000)
Libya	2000	Female	15 to 19	9.3% (2.9-20%)	91% (80-97%)	0.0084 (0.0023-0.019)	0.33% (0.19-0.49%)	0.14% (0.082-0.26%)	0.21 (0.12-0.38)	120 (65-190)
Libya	2000	Female	20 to 24	12% (3.5-27%)	88% (73-96%)	0.014 (0.0043-0.029)	0.66% (0.41-0.96%)	0.19% (0.1-0.31%)	0.32 (0.17-0.54)	260 (150-410)
Libya	2000	Female	25 to 29	14% (3.9-29%)	86% (71-96%)	0.017 (0.0056-0.033)	0.71% (0.49-0.98%)	0.31% (0.16-0.55%)	0.55 (0.28-0.98)	280 (180-420)
Libya	2000	Female	30 to 34	15% (4.4-33%)	85% (67-96%)	0.02 (0.0068-0.036)	0.62% (0.4-0.88%)	0.33% (0.19-0.55%)	0.64 (0.35-1)	230 (140-340)
Libya	2000	Female	35 to 39	17% (4.8-32%)	83% (68-95%)	0.023 (0.0083-0.042)	0.46% (0.32-0.64%)	0.27% (0.16-0.4%)	0.56 (0.32-0.84)	150 (94-210)
Libya	2000	Female	40 to 44	18% (6.7-33%)	82% (67-93%)	0.026 (0.01-0.047)	0.35% (0.21-0.5%)	0.29% (0.15-0.43%)	0.67 (0.34-1)	99 (57-150)
Libya	2000	Female	45 to 49	18% (6.4-34%)	82% (66-94%)	0.027 (0.0091-0.055)	0.25% (0.1-0.39%)	0.22% (0.037-0.38%)	0.64 (0.096-1.1)	68 (27-110)
Libya	2000	Female	50 to 54	19% (7.2-36%)	81% (64-93%)	0.028 (0.0089-0.058)	0.21% (0.031-0.35%)	0.23% (0.0083-0.42%)	0.81 (0.028-1.5)	51 (6.3-90)
Libya	2000	Female	55 to 59	20% (7.8-39%)	80% (61-92%)	0.029 (0.0086-0.062)	0.15% (-0.086-0.33%)	0.2% (-0.089-0.44%)	0.94 (-0.41-2.1)	40 (-22-87)
Libya	2000	Female	60 to 64	21% (8.3-40%)	79% (60-92%)	0.028 (0.0076-0.063)	0.11% (-0.18-0.33%)	0.17% (-0.22-0.44%)	1 (-1.4-2.7)	31 (-49-88)
Libya	2000	Female	65 to 69	21% (9.2-39%)	79% (61-91%)	0.024 (0.0067-0.052)	0.11% (-0.15-0.31%)	0.16% (-0.17-0.43%)	1.2 (-1.3-3.4)	28 (-39-83)
Libya	2000	Female	70 to 74	20% (8.2-37%)	80% (63-92%)	0.019 (0.005-0.048)	0.12% (-0.16-0.31%)	0.17% (-0.18-0.43%)	1.5 (-1.7-3.9)	29 (-37-75)
Libya	2000	Female	75 to 79	18% (6.4-37%)	82% (63-94%)	0.015 (0.0033-0.043)	0.11% (-0.21-0.27%)	0.14% (-0.24-0.35%)	1.3 (-2.4-3.3)	20 (-37-49)
Libya	2000	Female	80 to 84	16% (5.5-36%)	84% (64-94%)	0.014 (0.0029-0.039)	0.091% (-0.18-0.23%)	0.11% (-0.23-0.29%)	0.78 (-1.6-2.1)	8.6 (-17-22)
Libya	2000	Female	85 to 89	17% (5.7-38%)	83% (62-94%)	0.014 (0.003-0.041)	0.058% (-0.21-0.18%)	0.067% (-0.25-0.22%)	0.36 (-1.3-1.2)	2.9 (-10-9.4)
Libya	2000	Female	90 to 94	20% (6.5-42%)	80% (58-94%)	0.016 (0.0035-0.046)	0.035% (-0.23-0.16%)	0.033% (-0.29-0.18%)	0.095 (-0.85-0.54)	0.64 (-4.3-2.9)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	2000	Female	95 plus	24% (7.8-48%)	76% (52-92%)	0.019 (0.0036-0.054)	0.02% (-0.31-0.16%)	0.005% (-0.39-0.18%)	0.005 (-0.41-0.18)	0.085 (-1.3-0.67)
Libya	2000	Male	All Ages	9% (6-13%)	91% (87-94%)	0.014 (0.012-0.016)	0.44% (0.35-0.54%)	0.32% (0.23-0.42%)	36 (26-48)	3200 (2400-4100)
Libya	2000	Male	15 to 19	6.5% (2.7-13%)	93% (87-97%)	0.0056 (0.0018-0.012)	0.47% (0.28-0.71%)	0.14% (0.072-0.27%)	0.37 (0.19-0.72)	200 (110-320)
Libya	2000	Male	20 to 24	10% (3.9-20%)	90% (80-96%)	0.013 (0.0045-0.024)	0.94% (0.6-1.4%)	0.23% (0.11-0.43%)	0.86 (0.4-1.6)	470 (280-730)
Libya	2000	Male	25 to 29	13% (4.9-24%)	87% (76-95%)	0.02 (0.0077-0.034)	0.98% (0.69-1.4%)	0.37% (0.18-0.6%)	1.4 (0.7-2.4)	470 (320-710)
Libya	2000	Male	30 to 34	14% (5.2-28%)	86% (72-95%)	0.025 (0.011-0.043)	0.9% (0.62-1.2%)	0.47% (0.26-0.74%)	1.7 (0.95-2.8)	390 (260-560)
Libya	2000	Male	35 to 39	16% (6-29%)	84% (71-94%)	0.029 (0.012-0.053)	0.89% (0.66-1.2%)	0.67% (0.4-1%)	2.5 (1.4-3.8)	340 (240-470)
Libya	2000	Male	40 to 44	16% (7.2-28%)	84% (72-93%)	0.031 (0.013-0.059)	0.8% (0.57-1.1%)	0.71% (0.45-1.1%)	3 (1.8-4.5)	310 (210-420)
Libya	2000	Male	45 to 49	16% (6.2-29%)	84% (71-94%)	0.032 (0.011-0.063)	0.75% (0.56-1%)	0.79% (0.5-1.2%)	3.9 (2.5-6.1)	280 (200-380)
Libya	2000	Male	50 to 54	15% (6.4-29%)	85% (71-94%)	0.032 (0.011-0.066)	0.6% (0.42-0.81%)	0.67% (0.41-0.98%)	3.6 (2.2-5.2)	200 (140-270)
Libya	2000	Male	55 to 59	14% (6.2-27%)	86% (73-94%)	0.03 (0.01-0.065)	0.47% (0.3-0.65%)	0.53% (0.29-0.8%)	3.4 (1.9-5.3)	150 (96-210)
Libya	2000	Male	60 to 64	14% (5.9-26%)	86% (74-94%)	0.027 (0.008-0.062)	0.36% (0.2-0.51%)	0.41% (0.19-0.64%)	3.2 (1.5-5.1)	110 (59-160)
Libya	2000	Male	65 to 69	13% (5.6-24%)	87% (76-94%)	0.022 (0.0072-0.047)	0.33% (0.18-0.48%)	0.4% (0.2-0.62%)	3.6 (1.7-5.7)	93 (52-140)
Libya	2000	Male	70 to 74	12% (5.3-21%)	88% (79-95%)	0.016 (0.0049-0.038)	0.29% (0.16-0.43%)	0.35% (0.18-0.55%)	3.5 (1.7-5.5)	70 (38-110)
Libya	2000	Male	75 to 79	10% (4.3-20%)	90% (80-96%)	0.012 (0.0031-0.032)	0.22% (0.11-0.32%)	0.26% (0.13-0.39%)	2.5 (1.1-3.7)	39 (20-56)
Libya	2000	Male	80 to 84	9.1% (3.6-18%)	91% (82-96%)	0.01 (0.0028-0.025)	0.2% (0.11-0.3%)	0.23% (0.12-0.37%)	1.5 (0.74-2.3)	17 (9-25)
Libya	2000	Male	85 to 89	9% (3.6-18%)	91% (82-96%)	0.0096 (0.0027-0.023)	0.16% (0.076-0.24%)	0.18% (0.073-0.27%)	0.83 (0.33-1.3)	6.8 (3.2-9.9)
Libya	2000	Male	90 to 94	10% (4-21%)	90% (79-96%)	0.01 (0.0028-0.026)	0.13% (0.047-0.2%)	0.13% (0.027-0.22%)	0.32 (0.065-0.53)	1.9 (0.68-2.9)
Libya	2000	Male	95 plus	13% (4.7-27%)	87% (73-95%)	0.012 (0.0024-0.033)	0.14% (0.03-0.22%)	0.12% (-0.017-0.21%)	0.085 (-0.012-0.15)	0.42 (0.092-0.66)
Libya	2005	Female	All Ages	12% (6.7-18%)	88% (82-93%)	0.017 (0.013-0.02)	0.24% (0.15-0.33%)	0.086% (-0.11-0.24%)	8.1 (-11-22)	1500 (910-2200)
Libya	2005	Female	15 to 19	9.2% (3.1-20%)	91% (80-97%)	0.011 (0.0031-0.026)	0.33% (0.19-0.52%)	0.17% (0.091-0.3%)	0.23 (0.12-0.42)	120 (64-200)
Libya	2005	Female	20 to 24	12% (3.6-26%)	88% (74-96%)	0.018 (0.0056-0.037)	0.68% (0.42-1%)	0.21% (0.11-0.38%)	0.34 (0.17-0.6)	280 (170-440)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	2005	Female	25 to 29	13% (3.7-29%)	87% (71-96%)	0.021 (0.007-0.041)	0.73% (0.5-1%)	0.34% (0.17-0.6%)	0.62 (0.29-1.1)	320 (200-470)
Libya	2005	Female	30 to 34	15% (3.9-31%)	85% (69-96%)	0.025 (0.0088-0.045)	0.64% (0.43-0.9%)	0.36% (0.2-0.6%)	0.77 (0.42-1.3)	280 (180-420)
Libya	2005	Female	35 to 39	17% (4.6-32%)	83% (68-95%)	0.029 (0.011-0.052)	0.47% (0.32-0.64%)	0.29% (0.16-0.44%)	0.71 (0.36-1.1)	190 (120-270)
Libya	2005	Female	40 to 44	17% (6.2-32%)	83% (68-94%)	0.033 (0.014-0.061)	0.35% (0.19-0.52%)	0.29% (0.12-0.44%)	0.78 (0.32-1.2)	120 (66-190)
Libya	2005	Female	45 to 49	18% (6.3-33%)	82% (67-94%)	0.036 (0.012-0.072)	0.23% (0.059-0.36%)	0.2% (-0.013-0.37%)	0.64 (-0.045-1.2)	73 (19-120)
Libya	2005	Female	50 to 54	19% (7.4-36%)	81% (64-93%)	0.037 (0.012-0.076)	0.17% (-0.055-0.33%)	0.2% (-0.079-0.42%)	0.81 (-0.27-1.7)	52 (-16-100)
Libya	2005	Female	55 to 59	20% (7.8-38%)	80% (62-92%)	0.038 (0.011-0.081)	0.097% (-0.18-0.29%)	0.14% (-0.24-0.41%)	0.64 (-1.1-1.9)	26 (-52-79)
Libya	2005	Female	60 to 64	21% (8-40%)	79% (60-92%)	0.037 (0.0097-0.086)	0.045% (-0.33-0.3%)	0.081% (-0.38-0.4%)	0.51 (-2.6-2.6)	13 (-99-87)
Libya	2005	Female	65 to 69	22% (8.5-39%)	78% (61-91%)	0.032 (0.0086-0.068)	0.031% (-0.28-0.28%)	0.069% (-0.33-0.39%)	0.56 (-2.8-3.2)	8.9 (-84-80)
Libya	2005	Female	70 to 74	20% (7.8-37%)	80% (63-92%)	0.026 (0.0065-0.066)	0.048% (-0.28-0.28%)	0.084% (-0.33-0.38%)	0.82 (-3.2-3.7)	13 (-70-73)
Libya	2005	Female	75 to 79	18% (6.8-37%)	82% (63-93%)	0.021 (0.0046-0.058)	0.041% (-0.34-0.25%)	0.062% (-0.42-0.32%)	0.65 (-4.3-3.6)	8.5 (-70-53)
Libya	2005	Female	80 to 84	17% (5.6-36%)	83% (64-94%)	0.019 (0.0041-0.052)	0.028% (-0.32-0.2%)	0.039% (-0.37-0.25%)	0.33 (-3-2.2)	3.3 (-35-24)
Libya	2005	Female	85 to 89	17% (5.9-38%)	83% (62-94%)	0.019 (0.004-0.057)	-0.0026% (-0.34-0.16%)	-0.0051% (-0.41-0.2%)	-0.035 (-2.8-1.4)	-0.17 (-22-11)
Libya	2005	Female	90 to 94	20% (6.9-42%)	80% (58-93%)	0.021 (0.0045-0.062)	-0.028% (-0.38-0.14%)	-0.043% (-0.46-0.17%)	-0.16 (-1.7-0.64)	-0.65 (-8.9-3.5)
Libya	2005	Female	95 plus	24% (7.6-48%)	76% (52-92%)	0.025 (0.0047-0.071)	-0.049% (-0.47-0.15%)	-0.08% (-0.6-0.16%)	-0.11 (-0.79-0.22)	-0.27 (-2.6-0.84)
Libya	2005	Male	All Ages	9.4% (6.2-13%)	91% (87-94%)	0.019 (0.016-0.022)	0.49% (0.38-0.6%)	0.33% (0.22-0.45%)	40 (26-55)	3600 (2700-4700)
Libya	2005	Male	15 to 19	6.5% (2.7-13%)	93% (87-97%)	0.0073 (0.0023-0.017)	0.5% (0.31-0.74%)	0.18% (0.083-0.35%)	0.46 (0.21-0.92)	210 (120-340)
Libya	2005	Male	20 to 24	10% (3.9-20%)	90% (80-96%)	0.016 (0.0059-0.033)	1% (0.65-1.5%)	0.29% (0.13-0.58%)	1.1 (0.49-2.1)	510 (310-780)
Libya	2005	Male	25 to 29	12% (4.8-24%)	88% (76-95%)	0.026 (0.01-0.046)	1% (0.72-1.4%)	0.43% (0.21-0.75%)	1.8 (0.87-3.2)	560 (370-810)
Libya	2005	Male	30 to 34	14% (5-28%)	86% (72-95%)	0.033 (0.015-0.057)	0.94% (0.66-1.3%)	0.53% (0.27-0.86%)	2.3 (1.2-3.8)	500 (330-730)
Libya	2005	Male	35 to 39	16% (5.9-29%)	84% (71-94%)	0.039 (0.016-0.068)	0.9% (0.64-1.2%)	0.71% (0.41-1.1%)	3.1 (1.8-5)	430 (300-590)
Libya	2005	Male	40 to 44	16% (7.2-28%)	84% (72-93%)	0.041 (0.017-0.078)	0.81% (0.59-1.1%)	0.73% (0.44-1.1%)	3.3 (1.9-4.9)	340 (230-460)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	2005	Male	45 to 49	16% (6.3-29%)	84% (71-94%)	0.042 (0.015-0.083)	0.75% (0.54-1%)	0.79% (0.48-1.2%)	4 (2.3-6)	290 (200-390)
Libya	2005	Male	50 to 54	15% (6.4-29%)	85% (71-94%)	0.042 (0.015-0.088)	0.59% (0.39-0.81%)	0.65% (0.36-0.98%)	4 (2.1-6)	220 (140-310)
Libya	2005	Male	55 to 59	14% (6.1-28%)	86% (72-94%)	0.04 (0.013-0.086)	0.45% (0.25-0.64%)	0.5% (0.21-0.78%)	3.3 (1.4-5.3)	150 (82-210)
Libya	2005	Male	60 to 64	14% (5.7-27%)	86% (73-94%)	0.036 (0.01-0.082)	0.33% (0.13-0.5%)	0.37% (0.11-0.62%)	3.2 (0.91-5.3)	110 (43-170)
Libya	2005	Male	65 to 69	13% (5.8-24%)	87% (76-94%)	0.029 (0.0094-0.063)	0.31% (0.15-0.47%)	0.37% (0.15-0.58%)	3.5 (1.4-5.8)	94 (45-150)
Libya	2005	Male	70 to 74	12% (5.4-21%)	88% (79-95%)	0.022 (0.0066-0.053)	0.27% (0.13-0.42%)	0.32% (0.12-0.52%)	3.6 (1.4-5.8)	73 (34-110)
Libya	2005	Male	75 to 79	10% (4.3-20%)	90% (80-96%)	0.017 (0.0043-0.042)	0.21% (0.088-0.31%)	0.24% (0.091-0.39%)	2.7 (0.93-4.3)	42 (19-64)
Libya	2005	Male	80 to 84	9.1% (3.5-18%)	91% (82-96%)	0.014 (0.0035-0.033)	0.19% (0.084-0.3%)	0.22% (0.085-0.36%)	1.7 (0.67-2.8)	20 (8.7-31)
Libya	2005	Male	85 to 89	8.9% (3.7-17%)	91% (83-96%)	0.013 (0.0036-0.03)	0.15% (0.063-0.23%)	0.17% (0.056-0.27%)	1 (0.34-1.6)	8.4 (3.5-13)
Libya	2005	Male	90 to 94	10% (4.2-20%)	90% (80-96%)	0.013 (0.0037-0.034)	0.12% (0.02-0.2%)	0.11% (-0.0064-0.21%)	0.34 (-0.019-0.64)	2.2 (0.37-3.6)
Libya	2005	Male	95 plus	13% (4.7-27%)	87% (73-95%)	0.016 (0.0032-0.044)	0.12% (-0.0041-0.22%)	0.094% (-0.069-0.21%)	0.086 (-0.065-0.19)	0.47 (-0.015-0.83)
Libya	2010	Female	All Ages	12% (7.5-18%)	88% (82-93%)	0.019 (0.015-0.023)	0.26% (0.16-0.35%)	0.088% (-0.14-0.25%)	8.7 (-13-24)	1700 (980-2400)
Libya	2010	Female	15 to 19	9.2% (3-20%)	91% (80-97%)	0.012 (0.0033-0.029)	0.34% (0.19-0.53%)	0.18% (0.099-0.32%)	0.2 (0.11-0.35)	110 (57-180)
Libya	2010	Female	20 to 24	12% (3.3-26%)	88% (74-97%)	0.019 (0.0056-0.041)	0.7% (0.44-1%)	0.22% (0.12-0.4%)	0.33 (0.17-0.61)	280 (170-460)
Libya	2010	Female	25 to 29	13% (3.7-28%)	87% (72-96%)	0.023 (0.0072-0.046)	0.74% (0.5-1%)	0.34% (0.17-0.6%)	0.58 (0.28-1.1)	330 (200-500)
Libya	2010	Female	30 to 34	15% (4.1-31%)	85% (69-96%)	0.026 (0.0092-0.047)	0.65% (0.43-0.92%)	0.37% (0.2-0.62%)	0.8 (0.44-1.4)	320 (200-470)
Libya	2010	Female	35 to 39	16% (4.4-32%)	84% (68-96%)	0.031 (0.011-0.055)	0.48% (0.33-0.66%)	0.31% (0.17-0.47%)	0.82 (0.46-1.3)	240 (150-340)
Libya	2010	Female	40 to 44	17% (6.4-33%)	83% (67-94%)	0.034 (0.014-0.063)	0.35% (0.2-0.52%)	0.3% (0.15-0.47%)	0.97 (0.47-1.5)	160 (87-250)
Libya	2010	Female	45 to 49	18% (6.2-33%)	82% (67-94%)	0.037 (0.013-0.074)	0.24% (0.083-0.38%)	0.21% (0.0076-0.4%)	0.78 (0.023-1.5)	92 (31-150)
Libya	2010	Female	50 to 54	19% (7.1-36%)	81% (64-93%)	0.039 (0.012-0.08)	0.17% (-0.045-0.32%)	0.21% (-0.057-0.43%)	0.91 (-0.26-1.9)	59 (-15-120)
Libya	2010	Female	55 to 59	19% (7.4-38%)	81% (62-93%)	0.039 (0.011-0.085)	0.096% (-0.2-0.29%)	0.14% (-0.22-0.43%)	0.76 (-1.3-2.3)	31 (-66-97)
Libya	2010	Female	60 to 64	21% (7.9-41%)	79% (59-92%)	0.038 (0.0098-0.091)	0.043% (-0.37-0.29%)	0.084% (-0.43-0.41%)	0.53 (-2.8-2.7)	13 (-110-86)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	2010	Female	65 to 69	21% (8.6-39%)	79% (61-91%)	0.033 (0.0088-0.075)	0.027% (-0.3-0.27%)	0.067% (-0.34-0.4%)	0.56 (-3-3.4)	8.3 (-94-85)
Libya	2010	Female	70 to 74	20% (8-37%)	80% (63-92%)	0.027 (0.0069-0.07)	0.045% (-0.29-0.26%)	0.083% (-0.34-0.37%)	0.88 (-3.9-3.9)	13 (-85-76)
Libya	2010	Female	75 to 79	18% (6.8-34%)	82% (66-93%)	0.021 (0.0047-0.058)	0.036% (-0.33-0.23%)	0.058% (-0.41-0.31%)	0.67 (-4.7-3.6)	8.5 (-78-55)
Libya	2010	Female	80 to 84	16% (5.5-35%)	84% (65-95%)	0.019 (0.0042-0.055)	0.026% (-0.3-0.2%)	0.036% (-0.36-0.27%)	0.35 (-3.5-2.6)	3.5 (-40-28)
Libya	2010	Female	85 to 89	17% (5.3-37%)	83% (63-95%)	0.019 (0.004-0.061)	-0.007% (-0.35-0.16%)	-0.011% (-0.41-0.2%)	-0.092 (-3.6-1.8)	-0.58 (-28-14)
Libya	2010	Female	90 to 94	19% (6.4-43%)	81% (57-94%)	0.022 (0.0042-0.067)	-0.036% (-0.39-0.14%)	-0.054% (-0.49-0.16%)	-0.27 (-2.4-0.81)	-1.1 (-12-4.5)
Libya	2010	Female	95 plus	23% (7-49%)	77% (51-93%)	0.026 (0.0045-0.077)	-0.058% (-0.52-0.15%)	-0.092% (-0.67-0.16%)	-0.16 (-1.2-0.28)	-0.42 (-3.8-1.1)
Libya	2010	Male	All Ages	9.8% (6.4-14%)	90% (86-94%)	0.021 (0.017-0.026)	0.54% (0.42-0.67%)	0.36% (0.22-0.49%)	44 (28-62)	3900 (2900-5200)
Libya	2010	Male	15 to 19	6.5% (2.7-13%)	94% (87-97%)	0.0077 (0.0024-0.017)	0.51% (0.31-0.77%)	0.2% (0.086-0.39%)	0.45 (0.2-0.89)	190 (110-300)
Libya	2010	Male	20 to 24	10% (3.9-20%)	90% (80-96%)	0.018 (0.0056-0.035)	1% (0.68-1.5%)	0.32% (0.14-0.62%)	1.1 (0.5-2.2)	520 (320-810)
Libya	2010	Male	25 to 29	13% (4.8-24%)	87% (76-95%)	0.028 (0.01-0.052)	1.1% (0.75-1.4%)	0.46% (0.21-0.85%)	1.8 (0.81-3.3)	570 (380-820)
Libya	2010	Male	30 to 34	14% (5.2-28%)	86% (72-95%)	0.035 (0.016-0.061)	0.97% (0.67-1.3%)	0.55% (0.29-0.93%)	2.4 (1.3-4.2)	560 (370-790)
Libya	2010	Male	35 to 39	16% (5.9-29%)	84% (71-94%)	0.04 (0.016-0.07)	0.93% (0.68-1.2%)	0.73% (0.42-1.1%)	3.6 (2-5.9)	530 (370-720)
Libya	2010	Male	40 to 44	16% (7.2-29%)	84% (71-93%)	0.042 (0.018-0.077)	0.83% (0.58-1.1%)	0.74% (0.45-1.1%)	4 (2.3-6.1)	430 (290-580)
Libya	2010	Male	45 to 49	16% (6.3-29%)	84% (71-94%)	0.044 (0.016-0.089)	0.76% (0.54-1%)	0.81% (0.5-1.2%)	4.6 (2.7-6.8)	330 (230-440)
Libya	2010	Male	50 to 54	15% (6.2-29%)	85% (71-94%)	0.044 (0.015-0.09)	0.6% (0.39-0.83%)	0.67% (0.37-1%)	4.4 (2.2-6.6)	250 (160-340)
Libya	2010	Male	55 to 59	14% (6.1-27%)	86% (73-94%)	0.042 (0.014-0.087)	0.45% (0.26-0.65%)	0.5% (0.22-0.79%)	3.9 (1.7-6.2)	170 (100-250)
Libya	2010	Male	60 to 64	14% (5.7-26%)	86% (74-94%)	0.037 (0.011-0.084)	0.33% (0.12-0.51%)	0.37% (0.094-0.61%)	3.2 (0.87-5.4)	110 (41-180)
Libya	2010	Male	65 to 69	13% (6-25%)	87% (75-94%)	0.031 (0.01-0.064)	0.3% (0.14-0.47%)	0.36% (0.14-0.61%)	3.9 (1.5-6.5)	100 (48-170)
Libya	2010	Male	70 to 74	12% (5.4-22%)	88% (78-95%)	0.023 (0.007-0.056)	0.26% (0.12-0.4%)	0.31% (0.12-0.51%)	3.9 (1.5-6.4)	79 (34-130)
Libya	2010	Male	75 to 79	10% (4.5-20%)	90% (80-96%)	0.017 (0.0045-0.045)	0.21% (0.08-0.33%)	0.25% (0.078-0.4%)	3.2 (0.93-5.3)	49 (20-78)
Libya	2010	Male	80 to 84	9.3% (3.6-19%)	91% (81-96%)	0.015 (0.0038-0.036)	0.19% (0.082-0.3%)	0.22% (0.085-0.37%)	2.1 (0.8-3.6)	24 (9.9-39)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	2010	Male	85 to 89	9.2% (3.8-18%)	91% (82-96%)	0.014 (0.0039-0.034)	0.16% (0.057-0.24%)	0.17% (0.046-0.29%)	1.3 (0.35-2.3)	11 (4-18)
Libya	2010	Male	90 to 94	10% (4.3-21%)	90% (79-96%)	0.015 (0.004-0.036)	0.12% (0.016-0.21%)	0.12% (-0.01-0.23%)	0.47 (-0.041-0.94)	3 (0.4-5.3)
Libya	2010	Male	95 plus	13% (4.6-27%)	87% (73-95%)	0.017 (0.0034-0.047)	0.13% (-0.016-0.23%)	0.097% (-0.093-0.21%)	0.11 (-0.11-0.25)	0.61 (-0.074-1.1)
Libya	2016	Female	All Ages	13% (7.6-19%)	87% (81-92%)	0.022 (0.013-0.033)	0.24% (0.13-0.35%)	0.076% (-0.19-0.25%)	8.8 (-21-30)	1700 (840-2500)
Libya	2016	Female	15 to 19	9.3% (3.1-20%)	91% (80-97%)	0.013 (0.0035-0.033)	0.31% (0.19-0.5%)	0.14% (0.066-0.27%)	0.2 (0.1-0.39)	100 (57-170)
Libya	2016	Female	20 to 24	12% (3.7-25%)	88% (75-96%)	0.021 (0.0055-0.051)	0.64% (0.4-0.92%)	0.18% (0.083-0.33%)	0.28 (0.14-0.55)	220 (130-350)
Libya	2016	Female	25 to 29	14% (3.9-29%)	86% (71-96%)	0.025 (0.0067-0.059)	0.71% (0.49-0.99%)	0.29% (0.13-0.57%)	0.54 (0.26-1.1)	300 (180-440)
Libya	2016	Female	30 to 34	15% (3.7-31%)	85% (69-96%)	0.029 (0.0078-0.063)	0.64% (0.42-0.9%)	0.33% (0.17-0.57%)	0.78 (0.4-1.4)	300 (190-460)
Libya	2016	Female	35 to 39	17% (4.4-32%)	83% (68-96%)	0.033 (0.01-0.065)	0.47% (0.31-0.65%)	0.29% (0.15-0.45%)	0.92 (0.48-1.5)	250 (160-370)
Libya	2016	Female	40 to 44	17% (6.1-33%)	83% (67-94%)	0.037 (0.014-0.073)	0.35% (0.21-0.5%)	0.29% (0.12-0.47%)	1.2 (0.48-2)	190 (110-290)
Libya	2016	Female	45 to 49	18% (6.2-34%)	82% (66-94%)	0.04 (0.014-0.08)	0.24% (0.059-0.38%)	0.21% (-0.014-0.39%)	1.1 (-0.073-2.2)	130 (33-210)
Libya	2016	Female	50 to 54	19% (7.3-36%)	81% (64-93%)	0.041 (0.013-0.088)	0.17% (-0.051-0.34%)	0.2% (-0.094-0.42%)	1.3 (-0.58-2.8)	84 (-26-170)
Libya	2016	Female	55 to 59	20% (7.8-38%)	80% (62-92%)	0.042 (0.012-0.098)	0.089% (-0.26-0.29%)	0.13% (-0.29-0.4%)	0.92 (-2.2-2.9)	38 (-110-120)
Libya	2016	Female	60 to 64	21% (8.4-40%)	79% (60-92%)	0.041 (0.01-0.098)	0.031% (-0.39-0.29%)	0.07% (-0.45-0.41%)	0.59 (-3.9-3.5)	12 (-160-110)
Libya	2016	Female	65 to 69	21% (9.1-39%)	79% (61-91%)	0.036 (0.0099-0.088)	0.015% (-0.36-0.26%)	0.053% (-0.42-0.37%)	0.49 (-4-3.4)	5 (-120-87)
Libya	2016	Female	70 to 74	20% (8.4-37%)	80% (63-92%)	0.029 (0.0074-0.076)	0.035% (-0.35-0.27%)	0.071% (-0.4-0.37%)	0.81 (-4.5-4.3)	11 (-110-84)
Libya	2016	Female	75 to 79	18% (6.7-37%)	82% (63-93%)	0.024 (0.0047-0.068)	0.023% (-0.41-0.25%)	0.042% (-0.47-0.33%)	0.56 (-6.2-4.4)	6.2 (-100-66)
Libya	2016	Female	80 to 84	17% (5.9-37%)	83% (63-94%)	0.022 (0.0047-0.066)	0.011% (-0.39-0.2%)	0.018% (-0.46-0.27%)	0.21 (-5.2-3.2)	1.8 (-62-34)
Libya	2016	Female	85 to 89	17% (5.9-39%)	83% (61-94%)	0.022 (0.0049-0.07)	-0.022% (-0.42-0.17%)	-0.029% (-0.51-0.2%)	-0.33 (-5.7-2.3)	-2.3 (-44-17)
Libya	2016	Female	90 to 94	20% (6.9-42%)	80% (58-93%)	0.024 (0.0054-0.074)	-0.052% (-0.42-0.15%)	-0.073% (-0.54-0.17%)	-0.49 (-3.5-1.1)	-2.2 (-19-6.3)
Libya	2016	Female	95 plus	24% (8.4-49%)	76% (51-92%)	0.029 (0.0051-0.088)	-0.076% (-0.58-0.15%)	-0.12% (-0.72-0.17%)	-0.29 (-1.8-0.42)	-0.79 (-6.2-1.6)
Libya	2016	Male	All Ages	10% (7-14%)	90% (86-93%)	0.025 (0.015-0.036)	0.5% (0.38-0.62%)	0.34% (0.21-0.48%)	52 (32-76)	4000 (2900-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Libya	2016	Male	15 to 19	6.6% (2.7-13%)	93% (87-97%)	0.0085 (0.0022-0.02)	0.41% (0.24-0.63%)	0.14% (0.055-0.31%)	0.48 (0.19-1.1)	180 (100-290)
Libya	2016	Male	20 to 24	10% (3.9-20%)	90% (80-96%)	0.019 (0.0056-0.045)	0.76% (0.47-1.2%)	0.2% (0.075-0.43%)	0.94 (0.39-2)	390 (250-620)
Libya	2016	Male	25 to 29	13% (4.8-24%)	87% (76-95%)	0.03 (0.0098-0.064)	0.87% (0.58-1.2%)	0.31% (0.13-0.62%)	1.5 (0.64-2.9)	460 (300-670)
Libya	2016	Male	30 to 34	14% (5.2-29%)	86% (71-95%)	0.038 (0.014-0.077)	0.85% (0.57-1.2%)	0.41% (0.21-0.73%)	2 (1.1-3.5)	460 (310-680)
Libya	2016	Male	35 to 39	16% (6.2-30%)	84% (70-94%)	0.043 (0.016-0.085)	0.86% (0.62-1.1%)	0.61% (0.33-0.97%)	3.6 (2-5.9)	520 (360-710)
Libya	2016	Male	40 to 44	16% (7.4-29%)	84% (71-93%)	0.046 (0.018-0.091)	0.78% (0.55-1.1%)	0.65% (0.37-1%)	4.7 (2.6-7.6)	500 (330-690)
Libya	2016	Male	45 to 49	16% (6.5-29%)	84% (71-94%)	0.048 (0.017-0.095)	0.72% (0.5-0.97%)	0.74% (0.42-1.1%)	6.4 (3.7-10)	460 (310-630)
Libya	2016	Male	50 to 54	15% (6.5-29%)	85% (71-94%)	0.047 (0.016-0.099)	0.58% (0.35-0.8%)	0.63% (0.31-0.96%)	5.9 (2.8-9.2)	320 (200-460)
Libya	2016	Male	55 to 59	14% (6.3-28%)	86% (72-94%)	0.045 (0.014-0.097)	0.44% (0.23-0.65%)	0.48% (0.19-0.79%)	5.1 (2-8.5)	220 (110-330)
Libya	2016	Male	60 to 64	14% (5.8-27%)	86% (73-94%)	0.04 (0.012-0.094)	0.32% (0.086-0.51%)	0.35% (0.056-0.62%)	4.5 (0.66-7.8)	150 (42-250)
Libya	2016	Male	65 to 69	13% (5.8-24%)	87% (76-94%)	0.033 (0.0096-0.073)	0.29% (0.095-0.46%)	0.33% (0.096-0.56%)	4.4 (1.2-7.8)	120 (39-190)
Libya	2016	Male	70 to 74	12% (5.3-22%)	88% (78-95%)	0.025 (0.0068-0.061)	0.25% (0.087-0.4%)	0.3% (0.087-0.49%)	4.5 (1.3-7.6)	91 (30-150)
Libya	2016	Male	75 to 79	10% (4.2-20%)	90% (80-96%)	0.019 (0.0045-0.05)	0.2% (0.046-0.34%)	0.12% (0.036-0.42%)	3.7 (0.56-6.6)	57 (13-96)
Libya	2016	Male	80 to 84	9.3% (3.5-19%)	91% (81-97%)	0.016 (0.004-0.04)	0.19% (0.058-0.31%)	0.21% (0.054-0.37%)	2.5 (0.6-4.3)	28 (8-46)
Libya	2016	Male	85 to 89	9% (3.6-18%)	91% (82-96%)	0.014 (0.0039-0.038)	0.15% (0.052-0.26%)	0.16% (0.034-0.3%)	1.6 (0.33-2.9)	13 (4.3-23)
Libya	2016	Male	90 to 94	10% (4.1-21%)	90% (79-96%)	0.015 (0.0041-0.04)	0.12% (-0.00044-0.22%)	0.12% (-0.027-0.24%)	0.61 (-0.15-1.3)	3.9 (-0.014-7)
Libya	2016	Male	95 plus	13% (4.5-27%)	87% (73-95%)	0.018 (0.0035-0.052)	0.13% (-0.036-0.24%)	0.095% (-0.11-0.23%)	0.16 (-0.18-0.39)	0.85 (-0.25-1.6)
Morocco	1990	Female	All Ages	1% (0.73-1.3%)	99% (99-99%)	0.14 (0.1-0.18)	0.57% (0.37-1%)	1.1% (0.62-2.1%)	830 (470-1700)	29000 (18000-50000)
Morocco	1990	Female	15 to 19	1.1% (0.49-1.8%)	99% (98-100%)	0.13 (0.051-0.26)	0.71% (0.36-1.3%)	1.3% (0.54-2.8%)	16 (6.7-35)	1500 (770-3000)
Morocco	1990	Female	20 to 24	1.4% (0.61-2.3%)	99% (98-99%)	0.22 (0.089-0.4)	1.2% (0.61-2.1%)	1.9% (0.79-4.1%)	31 (12-64)	2800 (1400-5000)
Morocco	1990	Female	25 to 29	1.4% (0.69-2.4%)	99% (98-99%)	0.23 (0.11-0.41)	1.3% (0.67-2.4%)	2.1% (0.85-4.8%)	35 (14-77)	2900 (1400-5600)
Morocco	1990	Female	30 to 34	1.4% (0.63-2.5%)	99% (98-99%)	0.24 (0.11-0.42)	1.2% (0.63-2.4%)	2.1% (0.82-4.8%)	35 (14-80)	2600 (1300-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	1990	Female	35 to 39	1.4% (0.68-2.4%)	99% (98-99%)	0.25 (0.11-0.47)	1.1% (0.59-2.1%)	1.9% (0.85-4.1%)	33 (14-70)	2100 (1100-3900)
Morocco	1990	Female	40 to 44	1.4% (0.7-2.5%)	99% (98-99%)	0.27 (0.11-0.51)	1.2% (0.6-2.2%)	2% (0.91-4.2%)	28 (13-58)	1500 (770-2900)
Morocco	1990	Female	45 to 49	1.3% (0.58-2.4%)	99% (98-99%)	0.27 (0.11-0.52)	1.2% (0.59-2.5%)	2% (0.88-4.3%)	41 (17-85)	1900 (880-3800)
Morocco	1990	Female	50 to 54	1.4% (0.64-2.6%)	99% (97-99%)	0.28 (0.11-0.55)	1.2% (0.59-2.6%)	2% (0.86-4.3%)	57 (25-120)	2200 (1100-4700)
Morocco	1990	Female	55 to 59	1.4% (0.65-2.6%)	99% (97-99%)	0.27 (0.11-0.53)	1.3% (0.58-2.4%)	1.9% (0.82-3.8%)	75 (33-150)	2500 (1200-4900)
Morocco	1990	Female	60 to 64	1.4% (0.63-2.7%)	99% (97-99%)	0.26 (0.095-0.52)	1.3% (0.59-2.7%)	1.9% (0.79-4%)	100 (44-220)	2900 (1300-5900)
Morocco	1990	Female	65 to 69	1.4% (0.67-2.5%)	99% (98-99%)	0.23 (0.089-0.45)	1.3% (0.55-2.9%)	1.7% (0.74-4%)	110 (44-250)	2500 (1000-5600)
Morocco	1990	Female	70 to 74	1.4% (0.66-2.4%)	99% (98-99%)	0.19 (0.071-0.38)	1.2% (0.5-2.7%)	1.5% (0.65-3.6%)	89 (36-220)	1600 (690-3800)
Morocco	1990	Female	75 to 79	1.2% (0.55-2.2%)	99% (98-99%)	0.16 (0.051-0.35)	1% (0.38-2.4%)	1.3% (0.48-3.2%)	65 (23-160)	910 (340-2200)
Morocco	1990	Female	80 to 84	1.1% (0.5-2.1%)	99% (98-100%)	0.14 (0.046-0.29)	1% (0.36-2.3%)	1.3% (0.44-3%)	50 (17-120)	510 (190-1200)
Morocco	1990	Female	85 to 89	1.1% (0.53-2%)	99% (98-99%)	0.13 (0.049-0.28)	0.96% (0.37-2.2%)	1.2% (0.45-2.9%)	39 (15-93)	290 (110-670)
Morocco	1990	Female	90 to 94	1.2% (0.58-2.3%)	99% (98-99%)	0.15 (0.052-0.33)	0.88% (0.34-2.1%)	1.1% (0.4-2.6%)	17 (6.3-42)	86 (32-200)
Morocco	1990	Female	95 plus	1.6% (0.62-3.3%)	98% (97-99%)	0.18 (0.049-0.42)	0.95% (0.26-2.5%)	1.2% (0.29-3.1%)	5.1 (1.3-14)	17 (4.6-44)
Morocco	1990	Male	All Ages	2.5% (2-3.2%)	97% (97-98%)	0.66 (0.56-0.79)	1.3% (0.85-2%)	2.2% (1.4-3.6%)	1800 (1100-3000)	71000 (47000-110000)
Morocco	1990	Male	15 to 19	2.1% (1.1-3.4%)	98% (97-99%)	0.32 (0.14-0.62)	1.3% (0.74-2.1%)	1.8% (0.82-3.4%)	30 (14-56)	2900 (1600-4900)
Morocco	1990	Male	20 to 24	3.4% (1.8-5.7%)	97% (94-98%)	0.81 (0.39-1.4)	2.8% (1.5-4.8%)	3.8% (1.8-7.3%)	84 (38-160)	7200 (3900-12000)
Morocco	1990	Male	25 to 29	4.4% (2.4-6.9%)	96% (93-98%)	1.3 (0.66-2.2)	3.6% (1.9-6.3%)	5.3% (2.5-10%)	120 (56-230)	8900 (4800-15000)
Morocco	1990	Male	30 to 34	4.7% (2.5-7.9%)	95% (92-97%)	1.5 (0.81-2.4)	3.8% (2-6.8%)	5.7% (2.7-11%)	130 (60-250)	8200 (4400-15000)
Morocco	1990	Male	35 to 39	5% (2.7-8.2%)	95% (92-97%)	1.7 (0.78-3)	3.9% (2.2-6.9%)	6% (3-11%)	130 (63-230)	7200 (4000-13000)
Morocco	1990	Male	40 to 44	5% (2.9-8.3%)	95% (92-97%)	1.8 (0.8-3.3)	3.7% (2.1-6.3%)	5.6% (3-9.8%)	91 (48-160)	4700 (2700-7900)
Morocco	1990	Male	45 to 49	4.8% (2.5-7.7%)	95% (92-98%)	1.7 (0.72-3.3)	3.7% (2-6.4%)	5.4% (2.8-9.6%)	110 (55-200)	4800 (2600-8500)
Morocco	1990	Male	50 to 54	4.3% (2.4-7%)	96% (93-98%)	1.6 (0.67-2.9)	3.1% (1.7-5.4%)	4.3% (2.4-7.8%)	130 (70-240)	5100 (2900-9000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	1990	Male	55 to 59	3.9% (2-7%)	96% (93-98%)	1.4 (0.57-2.7)	2.9% (1.5-5.2%)	3.9% (2-7.1%)	170 (85-310)	5500 (2900-9800)
Morocco	1990	Male	60 to 64	3.6% (1.8-6.6%)	96% (93-98%)	1.2 (0.46-2.3)	2.6% (1.4-5.2%)	3.5% (1.7-6.8%)	210 (100-410)	5800 (3000-11000)
Morocco	1990	Male	65 to 69	3.4% (1.7-5.7%)	97% (94-98%)	0.93 (0.41-1.7)	2.5% (1.3-4.8%)	3.2% (1.6-6.3%)	200 (100-410)	4700 (2400-9000)
Morocco	1990	Male	70 to 74	3.1% (1.7-5.3%)	97% (95-98%)	0.71 (0.3-1.4)	2.4% (1.3-4.6%)	3% (1.6-5.8%)	180 (90-340)	3200 (1700-6100)
Morocco	1990	Male	75 to 79	2.8% (1.5-4.8%)	97% (95-99%)	0.55 (0.21-1.1)	2.1% (1-3.8%)	2.5% (1.3-4.7%)	120 (61-230)	1700 (870-3200)
Morocco	1990	Male	80 to 84	2.5% (1.3-4.4%)	98% (96-99%)	0.45 (0.17-0.88)	1.9% (0.95-3.6%)	2.3% (1.1-4.4%)	79 (38-150)	810 (400-1500)
Morocco	1990	Male	85 to 89	2.4% (1.2-4.1%)	98% (96-99%)	0.4 (0.16-0.82)	1.7% (0.85-3.3%)	2% (1-4.1%)	50 (25-100)	370 (180-730)
Morocco	1990	Male	90 to 94	2.5% (1.3-4.3%)	97% (96-99%)	0.4 (0.15-0.8)	1.5% (0.79-3%)	1.8% (0.91-3.6%)	20 (9.7-39)	99 (51-190)
Morocco	1990	Male	95 plus	3.1% (1.4-5.9%)	97% (94-99%)	0.43 (0.13-1)	1.6% (0.66-3.2%)	1.8% (0.74-3.8%)	4.7 (1.9-9.9)	16 (7-33)
Morocco	1995	Female	All Ages	1% (0.77-1.3%)	99% (99-99%)	0.14 (0.11-0.19)	0.64% (0.43-1.1%)	1.1% (0.68-2.1%)	880 (510-1700)	30000 (20000-51000)
Morocco	1995	Female	15 to 19	1% (0.51-1.7%)	99% (98-99%)	0.13 (0.052-0.26)	0.66% (0.35-1.2%)	1.2% (0.53-2.6%)	15 (6.3-31)	1500 (750-2700)
Morocco	1995	Female	20 to 24	1.3% (0.6-2.3%)	99% (98-99%)	0.22 (0.088-0.41)	1.1% (0.59-2%)	1.8% (0.75-3.9%)	27 (10-54)	2500 (1300-4500)
Morocco	1995	Female	25 to 29	1.4% (0.65-2.5%)	99% (98-99%)	0.24 (0.11-0.43)	1.2% (0.66-2.2%)	2.1% (0.88-4.7%)	32 (13-70)	2700 (1400-5000)
Morocco	1995	Female	30 to 34	1.4% (0.61-2.5%)	99% (98-99%)	0.24 (0.11-0.42)	1.2% (0.65-2.2%)	2.1% (0.87-4.5%)	36 (15-79)	2700 (1400-5100)
Morocco	1995	Female	35 to 39	1.4% (0.64-2.4%)	99% (98-99%)	0.25 (0.11-0.45)	1.1% (0.59-2%)	1.9% (0.87-3.8%)	38 (17-76)	2500 (1300-4500)
Morocco	1995	Female	40 to 44	1.4% (0.69-2.5%)	99% (98-99%)	0.27 (0.12-0.5)	1.1% (0.6-2.1%)	2% (0.93-4%)	43 (20-86)	2300 (1200-4300)
Morocco	1995	Female	45 to 49	1.4% (0.61-2.4%)	99% (98-99%)	0.28 (0.11-0.55)	1.2% (0.58-2.2%)	2% (0.89-4%)	37 (17-75)	1700 (880-3300)
Morocco	1995	Female	50 to 54	1.4% (0.66-2.6%)	99% (97-99%)	0.29 (0.11-0.59)	1.2% (0.59-2.3%)	1.9% (0.88-3.9%)	52 (25-110)	2100 (1100-4200)
Morocco	1995	Female	55 to 59	1.4% (0.66-2.5%)	99% (97-99%)	0.28 (0.11-0.55)	1.2% (0.59-2.2%)	1.8% (0.83-3.4%)	72 (33-140)	2400 (1200-4500)
Morocco	1995	Female	60 to 64	1.4% (0.64-2.7%)	99% (97-99%)	0.26 (0.099-0.53)	1.2% (0.58-2.4%)	1.7% (0.78-3.6%)	99 (44-200)	2800 (1300-5500)
Morocco	1995	Female	65 to 69	1.4% (0.68-2.6%)	99% (97-99%)	0.23 (0.093-0.45)	1.2% (0.53-2.4%)	1.6% (0.72-3.5%)	120 (53-260)	2800 (1300-6000)
Morocco	1995	Female	70 to 74	1.3% (0.67-2.5%)	99% (98-99%)	0.19 (0.073-0.38)	1.1% (0.49-2.3%)	1.4% (0.64-3.2%)	110 (51-270)	2100 (950-4800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	1995	Female	75 to 79	1.2% (0.54-2.2%)	99% (98-99%)	0.15 (0.05-0.34)	0.93% (0.36-2.1%)	1.2% (0.47-2.7%)	77 (29-180)	1100 (430-2500)
Morocco	1995	Female	80 to 84	1.1% (0.49-2.1%)	99% (98-100%)	0.14 (0.043-0.29)	0.93% (0.35-2.1%)	1.2% (0.43-2.7%)	53 (19-120)	550 (200-1200)
Morocco	1995	Female	85 to 89	1.1% (0.52-2.1%)	99% (98-99%)	0.13 (0.048-0.28)	0.89% (0.35-2%)	1.1% (0.42-2.5%)	37 (14-86)	270 (110-610)
Morocco	1995	Female	90 to 94	1.2% (0.57-2.3%)	99% (98-99%)	0.15 (0.051-0.33)	0.81% (0.32-1.9%)	1% (0.37-2.3%)	19 (6.8-44)	93 (35-220)
Morocco	1995	Female	95 plus	1.6% (0.62-3.3%)	98% (97-99%)	0.18 (0.048-0.43)	0.89% (0.24-2.2%)	1.1% (0.27-2.7%)	6.5 (1.6-16)	22 (6-52)
Morocco	1995	Male	All Ages	2.6% (2-3.3%)	97% (97-98%)	0.7 (0.59-0.84)	1.4% (0.97-2.1%)	2.3% (1.5-3.5%)	1900 (1200-2900)	72000 (49000-110000)
Morocco	1995	Male	15 to 19	2% (1.1-3.4%)	98% (97-99%)	0.32 (0.14-0.63)	1.2% (0.7-2%)	1.7% (0.77-3.1%)	28 (13-54)	2900 (1600-4800)
Morocco	1995	Male	20 to 24	3.4% (1.7-5.6%)	97% (94-98%)	0.81 (0.39-1.5)	2.7% (1.5-4.4%)	3.5% (1.6-6.5%)	74 (33-140)	6600 (3600-11000)
Morocco	1995	Male	25 to 29	4.3% (2.4-6.9%)	96% (93-98%)	1.3 (0.64-2.2)	3.4% (1.9-5.7%)	5% (2.3-9.2%)	110 (49-200)	8100 (4400-14000)
Morocco	1995	Male	30 to 34	4.7% (2.5-8.2%)	95% (92-98%)	1.5 (0.83-2.5)	3.5% (1.9-6.1%)	5.3% (2.6-10%)	120 (60-240)	8300 (4500-15000)
Morocco	1995	Male	35 to 39	5% (2.7-8%)	95% (92-97%)	1.7 (0.8-2.8)	3.6% (2-6.3%)	5.5% (2.9-10%)	140 (71-260)	8100 (4600-14000)
Morocco	1995	Male	40 to 44	5.1% (2.8-8.2%)	95% (92-97%)	1.8 (0.82-3.2)	3.4% (2-5.6%)	5.2% (2.8-9%)	130 (71-230)	6800 (4000-11000)
Morocco	1995	Male	45 to 49	4.8% (2.4-7.5%)	95% (92-98%)	1.7 (0.75-3.3)	3.4% (1.9-5.9%)	5.1% (2.7-8.9%)	110 (57-190)	4800 (2700-8200)
Morocco	1995	Male	50 to 54	4.3% (2.4-7%)	96% (93-98%)	1.6 (0.67-2.9)	2.9% (1.6-4.8%)	4.1% (2.3-7.2%)	110 (59-180)	4200 (2400-7000)
Morocco	1995	Male	55 to 59	3.9% (2-6.8%)	96% (93-98%)	1.4 (0.56-2.7)	2.6% (1.4-4.6%)	3.6% (1.9-6.4%)	140 (78-260)	4800 (2700-8400)
Morocco	1995	Male	60 to 64	3.5% (1.8-6.5%)	96% (94-98%)	1.2 (0.47-2.3)	2.4% (1.3-4.5%)	3.2% (1.7-6.1%)	180 (95-350)	5100 (2700-9500)
Morocco	1995	Male	65 to 69	3.3% (1.8-5.6%)	97% (94-98%)	0.93 (0.41-1.7)	2.3% (1.2-4.4%)	3% (1.6-5.8%)	220 (120-440)	5200 (2800-9900)
Morocco	1995	Male	70 to 74	3.1% (1.7-5.3%)	97% (95-98%)	0.71 (0.3-1.4)	2.2% (1.2-4.1%)	2.8% (1.5-5.2%)	220 (110-400)	3900 (2100-7200)
Morocco	1995	Male	75 to 79	2.7% (1.4-4.7%)	97% (95-99%)	0.54 (0.21-1.1)	1.9% (0.95-3.3%)	2.3% (1.2-4.1%)	140 (69-250)	1900 (1000-3500)
Morocco	1995	Male	80 to 84	2.4% (1.2-4.2%)	98% (96-99%)	0.45 (0.16-0.86)	1.8% (0.85-3.2%)	2.1% (1-3.9%)	82 (39-150)	850 (420-1600)
Morocco	1995	Male	85 to 89	2.3% (1.2-3.9%)	98% (96-99%)	0.39 (0.16-0.79)	1.5% (0.79-2.9%)	1.8% (0.92-3.5%)	46 (23-88)	340 (170-630)
Morocco	1995	Male	90 to 94	2.4% (1.3-4.2%)	98% (96-99%)	0.39 (0.15-0.79)	1.4% (0.73-2.7%)	1.6% (0.83-3.2%)	20 (10-40)	100 (53-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	1995	Male	95 plus	2.9% (1.4-5.8%)	97% (94-99%)	0.42 (0.12-0.97)	1.4% (0.64-2.8%)	1.6% (0.7-3.3%)	5.4 (2.4-11)	19 (8.4-37)
Morocco	2000	Female	All Ages	0.93% (0.69-1.2%)	99% (99-99%)	0.13 (0.099-0.17)	0.61% (0.43-0.94%)	1.1% (0.67-1.9%)	810 (500-1400)	27000 (19000-43000)
Morocco	2000	Female	15 to 19	0.94% (0.45-1.6%)	99% (98-100%)	0.12 (0.048-0.25)	0.56% (0.32-0.99%)	1.1% (0.49-2.2%)	12 (5.3-24)	1300 (710-2200)
Morocco	2000	Female	20 to 24	1.1% (0.52-2%)	99% (98-99%)	0.19 (0.073-0.35)	0.88% (0.51-1.5%)	1.5% (0.66-3.1%)	19 (8-38)	2100 (1100-3600)
Morocco	2000	Female	25 to 29	1.2% (0.57-2%)	99% (98-99%)	0.2 (0.087-0.37)	0.99% (0.58-1.7%)	1.7% (0.76-3.6%)	22 (10-47)	2200 (1300-3800)
Morocco	2000	Female	30 to 34	1.1% (0.51-2.1%)	99% (98-99%)	0.2 (0.086-0.36)	0.98% (0.56-1.8%)	1.8% (0.76-3.8%)	26 (11-57)	2200 (1200-4000)
Morocco	2000	Female	35 to 39	1.1% (0.54-2%)	99% (98-99%)	0.21 (0.094-0.39)	0.89% (0.52-1.5%)	1.6% (0.74-2.9%)	30 (14-58)	2100 (1200-3700)
Morocco	2000	Female	40 to 44	1.1% (0.58-2%)	99% (98-99%)	0.23 (0.1-0.41)	0.92% (0.53-1.6%)	1.6% (0.82-3.2%)	39 (19-75)	2200 (1300-3900)
Morocco	2000	Female	45 to 49	1.1% (0.51-2%)	99% (98-99%)	0.24 (0.099-0.46)	0.93% (0.52-1.7%)	1.6% (0.79-3.1%)	42 (21-79)	2100 (1100-3700)
Morocco	2000	Female	50 to 54	1.2% (0.59-2.2%)	99% (98-99%)	0.25 (0.1-0.5)	0.97% (0.53-1.8%)	1.6% (0.81-2.9%)	39 (21-75)	1600 (890-2900)
Morocco	2000	Female	55 to 59	1.3% (0.63-2.4%)	99% (98-99%)	0.26 (0.1-0.51)	1% (0.54-1.8%)	1.6% (0.8-2.9%)	57 (29-110)	2000 (1000-3600)
Morocco	2000	Female	60 to 64	1.4% (0.62-2.6%)	99% (97-99%)	0.25 (0.095-0.51)	1.1% (0.52-2.1%)	1.6% (0.72-3.1%)	84 (38-170)	2400 (1200-4700)
Morocco	2000	Female	65 to 69	1.4% (0.65-2.4%)	99% (98-99%)	0.23 (0.09-0.44)	1.1% (0.5-2.1%)	1.5% (0.69-3%)	110 (50-220)	2500 (1200-5100)
Morocco	2000	Female	70 to 74	1.3% (0.67-2.4%)	99% (98-99%)	0.18 (0.073-0.37)	0.96% (0.47-2%)	1.3% (0.62-2.6%)	120 (56-250)	2200 (1100-4700)
Morocco	2000	Female	75 to 79	1.2% (0.56-2.2%)	99% (98-99%)	0.15 (0.051-0.34)	0.83% (0.36-1.7%)	1.1% (0.46-2.3%)	94 (39-200)	1300 (570-2800)
Morocco	2000	Female	80 to 84	1.1% (0.47-2.1%)	99% (98-100%)	0.14 (0.044-0.29)	0.83% (0.33-1.8%)	1% (0.41-2.3%)	62 (24-140)	650 (260-1400)
Morocco	2000	Female	85 to 89	1.1% (0.5-2%)	99% (98-99%)	0.13 (0.047-0.28)	0.78% (0.32-1.7%)	0.97% (0.39-2%)	36 (14-78)	270 (110-560)
Morocco	2000	Female	90 to 94	1.2% (0.57-2.2%)	99% (98-99%)	0.14 (0.051-0.32)	0.71% (0.29-1.5%)	0.87% (0.34-1.9%)	16 (6.3-35)	81 (33-170)
Morocco	2000	Female	95 plus	1.5% (0.6-3.1%)	98% (97-99%)	0.17 (0.049-0.41)	0.77% (0.24-1.7%)	0.92% (0.27-2.2%)	6.3 (1.9-15)	21 (6.8-48)
Morocco	2000	Male	All Ages	2.8% (2.3-3.5%)	97% (96-98%)	0.78 (0.65-0.92)	1.5% (1.1-2.2%)	2.3% (1.6-3.3%)	1900 (1200-2700)	73000 (51000-100000)
Morocco	2000	Male	15 to 19	2.2% (1.2-3.5%)	98% (96-99%)	0.35 (0.15-0.66)	1.2% (0.74-2.1%)	1.8% (0.83-3.4%)	29 (14-54)	3100 (1800-5000)
Morocco	2000	Male	20 to 24	3.7% (2-6.2%)	96% (94-98%)	0.9 (0.43-1.6)	2.7% (1.6-4.4%)	3.8% (1.7-6.6%)	75 (35-130)	6900 (4000-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	2000	Male	25 to 29	4.8% (2.7-7.4%)	95% (93-97%)	1.4 (0.72-2.4)	3.4% (2-5.5%)	5.2% (2.6-9.1%)	100 (48-180)	7800 (4400-13000)
Morocco	2000	Male	30 to 34	5.1% (2.7-8.5%)	95% (92-97%)	1.7 (0.9-2.7)	3.4% (2-5.9%)	5.3% (2.7-9.9%)	110 (57-200)	7700 (4400-13000)
Morocco	2000	Male	35 to 39	5.1% (2.9-8.4%)	95% (92-97%)	1.7 (0.85-2.8)	3.3% (2-5.5%)	5.4% (3-9.4%)	140 (75-240)	8200 (4900-13000)
Morocco	2000	Male	40 to 44	5% (3-8.3%)	95% (92-97%)	1.7 (0.84-3)	3.2% (2-5%)	4.9% (2.9-8.3%)	140 (81-240)	7700 (4700-12000)
Morocco	2000	Male	45 to 49	4.7% (2.5-7.5%)	95% (92-97%)	1.7 (0.78-3.2)	3.1% (1.8-5.1%)	4.7% (2.7-8.2%)	150 (84-260)	6800 (4000-11000)
Morocco	2000	Male	50 to 54	4.2% (2.4-6.9%)	96% (93-98%)	1.5 (0.67-2.9)	2.6% (1.6-4.2%)	3.8% (2.3-6.3%)	110 (62-180)	4200 (2600-6800)
Morocco	2000	Male	55 to 59	3.7% (2-6.5%)	96% (93-98%)	1.3 (0.53-2.6)	2.4% (1.3-4%)	3.3% (1.9-5.6%)	120 (67-200)	4000 (2300-6700)
Morocco	2000	Male	60 to 64	3.4% (1.7-6%)	97% (94-98%)	1.1 (0.43-2.2)	2.1% (1.2-3.8%)	2.8% (1.5-5.1%)	160 (84-280)	4400 (2400-7700)
Morocco	2000	Male	65 to 69	3.1% (1.7-5.3%)	97% (95-98%)	0.87 (0.39-1.6)	2% (1.1-3.6%)	2.6% (1.4-4.8%)	190 (100-350)	4300 (2400-7900)
Morocco	2000	Male	70 to 74	2.9% (1.6-5%)	97% (95-98%)	0.66 (0.28-1.3)	1.9% (1-3.4%)	2.4% (1.3-4.3%)	220 (120-390)	4100 (2200-7000)
Morocco	2000	Male	75 to 79	2.6% (1.4-4.5%)	97% (95-99%)	0.5 (0.19-1)	1.6% (0.87-2.9%)	2% (1.1-3.5%)	160 (83-280)	2200 (1200-3900)
Morocco	2000	Male	80 to 84	2.3% (1.2-4%)	98% (96-99%)	0.42 (0.15-0.82)	1.5% (0.77-2.8%)	1.8% (0.91-3.3%)	92 (46-170)	960 (480-1700)
Morocco	2000	Male	85 to 89	2.2% (1.1-3.6%)	98% (96-99%)	0.37 (0.15-0.74)	1.3% (0.69-2.4%)	1.6% (0.8-2.9%)	46 (23-85)	340 (170-620)
Morocco	2000	Male	90 to 94	2.3% (1.2-4%)	98% (96-99%)	0.37 (0.14-0.74)	1.2% (0.64-2.2%)	1.4% (0.75-2.7%)	18 (9.4-34)	92 (49-170)
Morocco	2000	Male	95 plus	2.8% (1.3-5.5%)	97% (95-99%)	0.4 (0.12-0.93)	1.2% (0.55-2.4%)	1.4% (0.61-2.8%)	5.4 (2.3-11)	19 (8.4-36)
Morocco	2005	Female	All Ages	0.95% (0.73-1.3%)	99% (99-99%)	0.13 (0.098-0.17)	0.63% (0.46-0.95%)	1% (0.66-1.7%)	830 (520-1400)	28000 (20000-42000)
Morocco	2005	Female	15 to 19	0.94% (0.45-1.6%)	99% (98-100%)	0.12 (0.044-0.23)	0.52% (0.29-0.9%)	1% (0.46-2.1%)	9.5 (4.3-20)	1100 (600-1900)
Morocco	2005	Female	20 to 24	1.1% (0.52-2%)	99% (98-99%)	0.18 (0.071-0.34)	0.83% (0.5-1.4%)	1.4% (0.66-2.9%)	17 (7.4-34)	2000 (1200-3400)
Morocco	2005	Female	25 to 29	1.1% (0.54-2%)	99% (98-99%)	0.18 (0.079-0.34)	0.94% (0.59-1.5%)	1.7% (0.76-3.4%)	20 (9.1-41)	2200 (1300-3500)
Morocco	2005	Female	30 to 34	1.1% (0.49-2%)	99% (98-100%)	0.18 (0.08-0.33)	0.92% (0.56-1.5%)	1.7% (0.77-3.4%)	23 (10-47)	2100 (1300-3500)
Morocco	2005	Female	35 to 39	1.1% (0.53-1.9%)	99% (98-99%)	0.2 (0.086-0.35)	0.84% (0.5-1.3%)	1.5% (0.75-2.8%)	28 (14-52)	2100 (1200-3400)
Morocco	2005	Female	40 to 44	1.1% (0.59-1.9%)	99% (98-99%)	0.21 (0.096-0.38)	0.87% (0.51-1.5%)	1.6% (0.84-2.9%)	38 (20-70)	2300 (1300-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	2005	Female	45 to 49	1.2% (0.53-2%)	99% (98-99%)	0.23 (0.095-0.41)	0.88% (0.51-1.6%)	1.5% (0.77-2.8%)	48 (24-88)	2400 (1400-4100)
Morocco	2005	Female	50 to 54	1.2% (0.59-2.2%)	99% (98-99%)	0.24 (0.097-0.46)	0.92% (0.51-1.6%)	1.5% (0.77-2.7%)	55 (29-99)	2300 (1300-4000)
Morocco	2005	Female	55 to 59	1.3% (0.62-2.4%)	99% (98-99%)	0.25 (0.098-0.48)	0.96% (0.51-1.7%)	1.5% (0.76-2.6%)	51 (26-95)	1800 (980-3200)
Morocco	2005	Female	60 to 64	1.4% (0.62-2.6%)	99% (97-99%)	0.24 (0.09-0.49)	1% (0.5-1.8%)	1.5% (0.7-2.7%)	75 (37-140)	2200 (1100-4000)
Morocco	2005	Female	65 to 69	1.4% (0.65-2.5%)	99% (98-99%)	0.21 (0.086-0.42)	0.97% (0.49-1.8%)	1.4% (0.67-2.6%)	100 (50-200)	2400 (1200-4600)
Morocco	2005	Female	70 to 74	1.3% (0.66-2.4%)	99% (98-99%)	0.17 (0.07-0.35)	0.87% (0.43-1.7%)	1.2% (0.56-2.3%)	110 (53-220)	2100 (1000-4200)
Morocco	2005	Female	75 to 79	1.2% (0.55-2.2%)	99% (98-99%)	0.14 (0.049-0.32)	0.75% (0.33-1.5%)	0.97% (0.41-2%)	110 (45-220)	1500 (650-3100)
Morocco	2005	Female	80 to 84	1.1% (0.5-2%)	99% (98-99%)	0.13 (0.043-0.28)	0.75% (0.31-1.6%)	0.95% (0.37-2%)	78 (31-160)	820 (340-1700)
Morocco	2005	Female	85 to 89	1.1% (0.51-2%)	99% (98-99%)	0.13 (0.047-0.27)	0.71% (0.31-1.5%)	0.88% (0.36-1.8%)	48 (20-100)	350 (150-750)
Morocco	2005	Female	90 to 94	1.2% (0.58-2.2%)	99% (98-99%)	0.14 (0.049-0.29)	0.64% (0.25-1.4%)	0.79% (0.3-1.7%)	18 (6.9-38)	91 (36-190)
Morocco	2005	Female	95 plus	1.5% (0.64-3.2%)	98% (97-99%)	0.17 (0.046-0.41)	0.7% (0.22-1.5%)	0.84% (0.24-1.9%)	6.2 (1.8-14)	21 (6.6-46)
Morocco	2005	Male	All Ages	2.9% (2.4-3.6%)	97% (96-98%)	0.77 (0.63-0.92)	1.6% (1.2-2.3%)	2.3% (1.6-3.2%)	1900 (1300-2700)	75000 (54000-100000)
Morocco	2005	Male	15 to 19	2.2% (1.2-3.5%)	98% (96-99%)	0.32 (0.13-0.6)	1.2% (0.71-1.9%)	1.7% (0.8-3.1%)	25 (12-46)	2700 (1700-4400)
Morocco	2005	Male	20 to 24	3.7% (2-6.1%)	96% (94-98%)	0.83 (0.38-1.5)	2.6% (1.5-4.1%)	3.6% (1.7-6.3%)	71 (34-130)	6800 (4000-11000)
Morocco	2005	Male	25 to 29	4.8% (2.6-7.3%)	95% (93-97%)	1.3 (0.64-2.3)	3.2% (1.9-5%)	4.9% (2.5-8.3%)	92 (45-160)	7400 (4400-11000)
Morocco	2005	Male	30 to 34	5% (2.8-8.3%)	95% (92-97%)	1.5 (0.81-2.5)	3.2% (2-5.3%)	5.1% (2.8-9.2%)	97 (52-170)	6900 (4100-11000)
Morocco	2005	Male	35 to 39	5.2% (3-8.3%)	95% (92-97%)	1.6 (0.81-2.7)	3.2% (2-5.1%)	5.2% (2.9-8.6%)	120 (68-200)	7300 (4600-12000)
Morocco	2005	Male	40 to 44	5.1% (3-8.2%)	95% (92-97%)	1.6 (0.78-2.8)	3% (1.9-4.7%)	4.7% (2.7-7.7%)	150 (83-240)	8000 (4800-12000)
Morocco	2005	Male	45 to 49	4.7% (2.4-7.5%)	95% (93-98%)	1.6 (0.72-2.8)	2.9% (1.8-4.7%)	4.5% (2.6-7.4%)	170 (97-280)	8000 (4800-13000)
Morocco	2005	Male	50 to 54	4.2% (2.4-6.7%)	96% (93-98%)	1.4 (0.64-2.5)	2.5% (1.5-3.9%)	3.6% (2.2-5.8%)	160 (93-260)	6400 (3900-10000)
Morocco	2005	Male	55 to 59	3.7% (2-6.5%)	96% (94-98%)	1.2 (0.5-2.4)	2.2% (1.3-3.6%)	3.1% (1.8-5.1%)	130 (75-220)	4600 (2700-7400)
Morocco	2005	Male	60 to 64	3.3% (1.7-6%)	97% (94-98%)	1 (0.4-2)	2% (1.1-3.5%)	2.7% (1.5-4.9%)	150 (86-270)	4400 (2500-7700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	2005	Male	65 to 69	3.1% (1.6-5.3%)	97% (95-98%)	0.8 (0.36-1.5)	1.9% (1.1-3.2%)	2.4% (1.4-4.3%)	180 (100-320)	4200 (2500-7300)
Morocco	2005	Male	70 to 74	2.9% (1.6-4.9%)	97% (95-98%)	0.61 (0.26-1.2)	1.8% (1-3%)	2.3% (1.3-4%)	200 (110-340)	3700 (2100-6200)
Morocco	2005	Male	75 to 79	2.5% (1.3-4.4%)	97% (96-99%)	0.47 (0.18-0.94)	1.5% (0.78-2.6%)	1.8% (0.98-3.2%)	170 (91-300)	2500 (1300-4200)
Morocco	2005	Male	80 to 84	2.2% (1.1-4%)	98% (96-99%)	0.38 (0.14-0.75)	1.4% (0.71-2.5%)	1.7% (0.86-3.1%)	110 (56-210)	1200 (600-2100)
Morocco	2005	Male	85 to 89	2.1% (1.1-3.7%)	98% (96-99%)	0.34 (0.13-0.67)	1.2% (0.63-2.2%)	1.4% (0.73-2.7%)	60 (30-110)	440 (230-820)
Morocco	2005	Male	90 to 94	2.3% (1.2-3.9%)	98% (96-99%)	0.34 (0.13-0.67)	1.1% (0.6-2%)	1.3% (0.69-2.4%)	21 (11-39)	110 (59-200)
Morocco	2005	Male	95 plus	2.8% (1.3-5.5%)	97% (95-99%)	0.37 (0.11-0.89)	1.1% (0.51-2.2%)	1.3% (0.55-2.6%)	5.7 (2.4-11)	20 (8.9-38)
Morocco	2010	Female	All Ages	1.1% (0.81-1.4%)	99% (99-99%)	0.12 (0.092-0.16)	0.62% (0.46-0.88%)	0.93% (0.59-1.5%)	780 (480-1300)	27000 (20000-39000)
Morocco	2010	Female	15 to 19	1% (0.48-1.7%)	99% (98-100%)	0.097 (0.038-0.2)	0.46% (0.26-0.76%)	0.89% (0.41-1.8%)	7.1 (3.2-14)	930 (510-1500)
Morocco	2010	Female	20 to 24	1.3% (0.58-2.3%)	99% (98-99%)	0.16 (0.062-0.3)	0.77% (0.47-1.2%)	1.4% (0.63-2.7%)	13 (6-26)	1800 (1000-2900)
Morocco	2010	Female	25 to 29	1.4% (0.61-2.4%)	99% (98-99%)	0.17 (0.073-0.31)	0.89% (0.57-1.4%)	1.7% (0.77-3.3%)	19 (8.6-38)	2200 (1300-3400)
Morocco	2010	Female	30 to 34	1.3% (0.59-2.4%)	99% (98-99%)	0.17 (0.074-0.31)	0.9% (0.54-1.4%)	1.7% (0.82-3.3%)	22 (10-43)	2200 (1300-3400)
Morocco	2010	Female	35 to 39	1.3% (0.64-2.4%)	99% (98-99%)	0.18 (0.08-0.33)	0.8% (0.51-1.3%)	1.5% (0.78-2.7%)	26 (13-47)	2000 (1200-3100)
Morocco	2010	Female	40 to 44	1.3% (0.67-2.4%)	99% (98-99%)	0.19 (0.085-0.35)	0.83% (0.49-1.4%)	1.6% (0.83-2.8%)	36 (18-63)	2200 (1300-3600)
Morocco	2010	Female	45 to 49	1.3% (0.56-2.3%)	99% (98-99%)	0.2 (0.083-0.37)	0.84% (0.47-1.5%)	1.5% (0.72-2.8%)	48 (23-91)	2400 (1300-4300)
Morocco	2010	Female	50 to 54	1.3% (0.62-2.5%)	99% (97-99%)	0.21 (0.083-0.4)	0.86% (0.47-1.5%)	1.4% (0.73-2.5%)	61 (31-110)	2600 (1400-4600)
Morocco	2010	Female	55 to 59	1.4% (0.64-2.5%)	99% (97-99%)	0.2 (0.082-0.37)	0.86% (0.45-1.5%)	1.3% (0.7-2.3%)	67 (34-120)	2400 (1300-4300)
Morocco	2010	Female	60 to 64	1.4% (0.63-2.7%)	99% (97-99%)	0.19 (0.074-0.39)	0.86% (0.45-1.5%)	1.3% (0.63-2.3%)	61 (30-110)	1800 (940-3200)
Morocco	2010	Female	65 to 69	1.4% (0.66-2.5%)	99% (97-99%)	0.17 (0.068-0.34)	0.82% (0.4-1.5%)	1.2% (0.56-2.2%)	81 (40-150)	2000 (1000-3800)
Morocco	2010	Female	70 to 74	1.4% (0.66-2.4%)	99% (98-99%)	0.14 (0.054-0.28)	0.71% (0.31-1.4%)	0.96% (0.42-2%)	92 (40-190)	1800 (790-3600)
Morocco	2010	Female	75 to 79	1.2% (0.54-2.3%)	99% (98-99%)	0.11 (0.038-0.25)	0.59% (0.23-1.3%)	0.78% (0.3-1.7%)	87 (33-190)	1300 (500-2800)
Morocco	2010	Female	80 to 84	1.1% (0.47-2.1%)	99% (98-100%)	0.1 (0.033-0.21)	0.58% (0.23-1.2%)	0.74% (0.28-1.6%)	72 (27-150)	770 (300-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	2010	Female	85 to 89	1.1% (0.5-2%)	99% (98-100%)	0.098 (0.036-0.21)	0.54% (0.21-1.1%)	0.67% (0.25-1.4%)	54 (20-120)	400 (160-850)
Morocco	2010	Female	90 to 94	1.2% (0.57-2.3%)	99% (98-99%)	0.11 (0.038-0.24)	0.5% (0.17-1.1%)	0.6% (0.2-1.4%)	22 (7.2-49)	110 (39-250)
Morocco	2010	Female	95 plus	1.6% (0.62-3.4%)	98% (97-99%)	0.14 (0.036-0.33)	0.55% (0.13-1.3%)	0.65% (0.14-1.6%)	6.4 (1.4-15)	22 (5.6-52)
Morocco	2010	Male	All Ages	2.9% (2.3-3.5%)	97% (96-98%)	0.58 (0.47-0.7)	1.6% (1.2-2.2%)	2.2% (1.6-3%)	1900 (1300-2600)	72000 (53000-96000)
Morocco	2010	Male	15 to 19	2% (1-3.3%)	98% (97-99%)	0.22 (0.09-0.44)	1% (0.61-1.6%)	1.4% (0.64-2.6%)	19 (8.7-36)	2300 (1300-3600)
Morocco	2010	Male	20 to 24	3.2% (1.7-5.3%)	97% (95-98%)	0.55 (0.24-1)	2.3% (1.4-3.4%)	2.9% (1.5-5.1%)	55 (27-97)	5700 (3500-8800)
Morocco	2010	Male	25 to 29	4.2% (2.3-6.6%)	96% (93-98%)	0.87 (0.4-1.5)	2.8% (1.7-4.3%)	4% (2.1-6.9%)	81 (41-140)	7000 (4200-11000)
Morocco	2010	Male	30 to 34	4.5% (2.4-7.7%)	95% (92-98%)	1 (0.52-1.7)	2.8% (1.7-4.6%)	4.4% (2.4-7.6%)	86 (46-160)	6400 (3900-10000)
Morocco	2010	Male	35 to 39	4.8% (2.6-7.9%)	95% (92-97%)	1.1 (0.54-2)	2.9% (1.8-4.6%)	4.6% (2.6-7.7%)	100 (57-170)	6400 (4100-10000)
Morocco	2010	Male	40 to 44	4.9% (2.8-8%)	95% (92-97%)	1.2 (0.54-2)	2.8% (1.8-4.3%)	4.3% (2.6-6.9%)	120 (71-200)	6800 (4300-10000)
Morocco	2010	Male	45 to 49	4.6% (2.3-7.4%)	95% (93-98%)	1.2 (0.53-2.1)	2.7% (1.7-4.3%)	4.1% (2.4-6.6%)	170 (97-280)	8000 (4800-12000)
Morocco	2010	Male	50 to 54	4.1% (2.3-6.7%)	96% (93-98%)	1.1 (0.48-1.9)	2.3% (1.4-3.5%)	3.3% (2-5.3%)	180 (100-280)	7200 (4400-11000)
Morocco	2010	Male	55 to 59	3.7% (1.9-6.5%)	96% (93-98%)	0.93 (0.4-1.8)	2.1% (1.2-3.3%)	2.9% (1.7-4.7%)	180 (100-290)	6200 (3700-9800)
Morocco	2010	Male	60 to 64	3.4% (1.8-6.2%)	97% (94-98%)	0.78 (0.32-1.6)	1.9% (1.1-3.2%)	2.5% (1.4-4.4%)	150 (89-260)	4400 (2600-7300)
Morocco	2010	Male	65 to 69	3.2% (1.7-5.5%)	97% (95-98%)	0.63 (0.28-1.2)	1.7% (1-2.9%)	2.3% (1.3-3.9%)	170 (98-290)	4000 (2300-6600)
Morocco	2010	Male	70 to 74	3% (1.6-5.1%)	97% (95-98%)	0.47 (0.21-0.94)	1.6% (0.95-2.7%)	2.1% (1.2-3.5%)	190 (110-320)	3600 (2100-6000)
Morocco	2010	Male	75 to 79	2.6% (1.4-4.7%)	97% (95-99%)	0.36 (0.14-0.74)	1.3% (0.7-2.3%)	1.6% (0.85-2.9%)	150 (78-260)	2200 (1200-3800)
Morocco	2010	Male	80 to 84	2.3% (1.2-4.1%)	98% (96-99%)	0.3 (0.11-0.6)	1.2% (0.61-2.1%)	1.5% (0.73-2.7%)	120 (56-210)	1200 (610-2200)
Morocco	2010	Male	85 to 89	2.2% (1.1-3.7%)	98% (96-99%)	0.27 (0.11-0.55)	1.1% (0.56-1.9%)	1.3% (0.66-2.2%)	78 (40-140)	590 (310-1000)
Morocco	2010	Male	90 to 94	2.4% (1.2-4.1%)	98% (96-99%)	0.27 (0.1-0.53)	0.95% (0.48-1.7%)	1.1% (0.56-2%)	30 (15-53)	150 (80-270)
Morocco	2010	Male	95 plus	2.9% (1.3-5.5%)	97% (94-99%)	0.29 (0.083-0.68)	0.95% (0.39-1.9%)	1.1% (0.4-2.2%)	6.9 (2.6-14)	24 (9.7-49)
Morocco	2016	Female	All Ages	1.1% (0.84-1.4%)	99% (99-99%)	0.11 (0.069-0.17)	0.57% (0.4-0.81%)	0.79% (0.45-1.3%)	700 (380-1200)	25000 (17000-36000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	2016	Female	15 to 19	1% (0.48-1.7%)	99% (98-100%)	0.087 (0.031-0.19)	0.41% (0.24-0.68%)	0.76% (0.31-1.6%)	4.9 (2.1-10)	720 (410-1200)
Morocco	2016	Female	20 to 24	1.3% (0.59-2.2%)	99% (98-99%)	0.14 (0.051-0.31)	0.72% (0.45-1.1%)	1.2% (0.53-2.2%)	9.7 (4.2-19)	1500 (910-2400)
Morocco	2016	Female	25 to 29	1.4% (0.63-2.4%)	99% (98-99%)	0.15 (0.057-0.3)	0.81% (0.52-1.2%)	1.4% (0.64-2.9%)	14 (6.2-28)	1900 (1200-2900)
Morocco	2016	Female	30 to 34	1.3% (0.63-2.4%)	99% (98-99%)	0.15 (0.059-0.3)	0.8% (0.52-1.2%)	1.5% (0.72-2.9%)	19 (8.9-37)	2100 (1300-3300)
Morocco	2016	Female	35 to 39	1.3% (0.65-2.4%)	99% (98-99%)	0.16 (0.067-0.33)	0.72% (0.45-1.1%)	1.3% (0.68-2.4%)	22 (11-40)	1900 (1200-2900)
Morocco	2016	Female	40 to 44	1.3% (0.69-2.4%)	99% (98-99%)	0.17 (0.066-0.34)	0.71% (0.43-1.1%)	1.4% (0.7-2.4%)	30 (15-54)	2000 (1200-3200)
Morocco	2016	Female	45 to 49	1.3% (0.56-2.3%)	99% (98-99%)	0.18 (0.065-0.35)	0.71% (0.41-1.2%)	1.3% (0.61-2.4%)	40 (19-74)	2200 (1200-3700)
Morocco	2016	Female	50 to 54	1.3% (0.64-2.5%)	99% (98-99%)	0.18 (0.066-0.36)	0.73% (0.41-1.3%)	1.2% (0.63-2.2%)	54 (28-98)	2400 (1400-4200)
Morocco	2016	Female	55 to 59	1.4% (0.64-2.5%)	99% (97-99%)	0.17 (0.067-0.35)	0.74% (0.4-1.3%)	1.2% (0.59-2.1%)	70 (36-130)	2600 (1400-4700)
Morocco	2016	Female	60 to 64	1.4% (0.61-2.7%)	99% (97-99%)	0.17 (0.058-0.37)	0.73% (0.37-1.3%)	1.1% (0.52-2%)	80 (39-140)	2400 (1200-4400)
Morocco	2016	Female	65 to 69	1.4% (0.62-2.5%)	99% (97-99%)	0.15 (0.052-0.32)	0.68% (0.32-1.3%)	0.97% (0.46-1.9%)	67 (31-130)	1700 (770-3300)
Morocco	2016	Female	70 to 74	1.3% (0.66-2.4%)	99% (98-99%)	0.12 (0.043-0.26)	0.58% (0.22-1.1%)	0.79% (0.29-1.6%)	70 (26-150)	1400 (530-2800)
Morocco	2016	Female	75 to 79	1.2% (0.56-2.2%)	99% (98-99%)	0.1 (0.032-0.23)	0.48% (0.17-0.99%)	0.63% (0.21-1.3%)	71 (23-150)	1100 (350-2200)
Morocco	2016	Female	80 to 84	1.1% (0.48-2%)	99% (98-100%)	0.089 (0.028-0.2)	0.47% (0.17-0.96%)	0.59% (0.2-1.3%)	64 (21-140)	690 (240-1400)
Morocco	2016	Female	85 to 89	1.1% (0.5-2%)	99% (98-99%)	0.086 (0.03-0.2)	0.43% (0.14-0.92%)	0.53% (0.16-1.2%)	53 (16-120)	400 (130-870)
Morocco	2016	Female	90 to 94	1.2% (0.57-2.2%)	99% (98-99%)	0.095 (0.029-0.23)	0.39% (0.11-0.87%)	0.48% (0.13-1.1%)	27 (7.5-63)	140 (42-320)
Morocco	2016	Female	95 plus	1.6% (0.65-3.3%)	98% (97-99%)	0.12 (0.029-0.3)	0.43% (0.082-1%)	0.5% (0.078-1.2%)	9 (1.4-22)	31 (6.3-76)
Morocco	2016	Male	All Ages	3% (2.4-3.7%)	97% (96-98%)	0.53 (0.34-0.78)	1.6% (1.2-2.1%)	2% (1.4-2.6%)	1900 (1400-2600)	72000 (55000-94000)
Morocco	2016	Male	15 to 19	2% (1-3.1%)	98% (97-99%)	0.19 (0.068-0.41)	0.93% (0.56-1.5%)	1.2% (0.52-2.3%)	15 (6.4-28)	1900 (1100-3000)
Morocco	2016	Male	20 to 24	3.2% (1.7-5.4%)	97% (95-98%)	0.48 (0.19-0.97)	2.1% (1.3-3.1%)	2.6% (1.3-4.5%)	49 (24-85)	5300 (3300-8000)
Morocco	2016	Male	25 to 29	4.1% (2.2-6.6%)	96% (93-98%)	0.76 (0.32-1.5)	2.5% (1.6-3.8%)	3.6% (1.9-6.1%)	74 (38-120)	6700 (4200-10000)
Morocco	2016	Male	30 to 34	4.5% (2.4-7.8%)	95% (92-98%)	0.91 (0.42-1.7)	2.6% (1.6-4%)	4% (2.2-6.8%)	89 (49-150)	6900 (4300-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Morocco	2016	Male	35 to 39	4.8% (2.6-7.9%)	95% (92-97%)	0.99 (0.42-1.9)	2.6% (1.7-4%)	4.2% (2.5-6.9%)	100 (60-170)	6700 (4300-10000)
Morocco	2016	Male	40 to 44	4.9% (2.8-8.3%)	95% (92-97%)	1 (0.44-2)	2.5% (1.7-3.8%)	4% (2.5-6.2%)	110 (66-180)	6300 (4100-9500)
Morocco	2016	Male	45 to 49	4.6% (2.3-7.4%)	95% (93-98%)	1 (0.42-1.9)	2.5% (1.6-3.8%)	3.9% (2.3-6.1%)	140 (83-220)	6800 (4300-10000)
Morocco	2016	Male	50 to 54	4.1% (2.3-7%)	96% (93-98%)	0.92 (0.37-1.7)	2.1% (1.4-3.2%)	3.1% (1.9-4.7%)	170 (100-270)	7200 (4500-11000)
Morocco	2016	Male	55 to 59	3.7% (2-6.6%)	96% (93-98%)	0.8 (0.3-1.6)	1.9% (1.2-3%)	2.7% (1.6-4.2%)	200 (120-330)	7000 (4200-11000)
Morocco	2016	Male	60 to 64	3.4% (1.7-6.1%)	97% (94-98%)	0.66 (0.25-1.4)	1.7% (0.98-2.8%)	2.3% (1.3-3.8%)	210 (120-340)	6000 (3600-9900)
Morocco	2016	Male	65 to 69	3.1% (1.7-5.3%)	97% (95-98%)	0.53 (0.21-1.1)	1.5% (0.92-2.4%)	2% (1.2-3.3%)	170 (100-280)	4100 (2500-6600)
Morocco	2016	Male	70 to 74	2.9% (1.6-5%)	97% (95-98%)	0.41 (0.16-0.89)	1.4% (0.81-2.3%)	1.8% (1-3.1%)	170 (96-290)	3200 (1900-5400)
Morocco	2016	Male	75 to 79	2.6% (1.4-4.7%)	97% (95-99%)	0.31 (0.11-0.7)	1.1% (0.57-2%)	1.4% (0.7-2.5%)	140 (72-260)	2100 (1100-3700)
Morocco	2016	Male	80 to 84	2.3% (1.2-4.1%)	98% (96-99%)	0.26 (0.089-0.55)	1% (0.51-1.8%)	1.3% (0.62-2.1%)	110 (55-200)	1200 (610-2100)
Morocco	2016	Male	85 to 89	2.2% (1.1-3.8%)	98% (96-99%)	0.23 (0.086-0.49)	0.89% (0.43-1.5%)	1.1% (0.49-1.8%)	80 (37-140)	610 (300-1100)
Morocco	2016	Male	90 to 94	2.4% (1.3-4.2%)	98% (96-99%)	0.23 (0.082-0.51)	0.8% (0.34-1.5%)	0.93% (0.37-1.7%)	40 (16-76)	210 (89-390)
Morocco	2016	Male	95 plus	2.9% (1.3-5.8%)	97% (94-99%)	0.25 (0.067-0.63)	0.8% (0.27-1.6%)	0.9% (0.26-1.9%)	10 (3-22)	38 (13-75)
Palestine	1990	Female	All Ages	9.8% (5.9-15%)	90% (85-94%)	0.017 (0.01-0.028)	0.1% (0.026-0.18%)	-0.05% (-0.26-0.087%)	-2.1 (-11-3.5)	320 (84-540)
Palestine	1990	Female	15 to 19	9.3% (2.8-21%)	91% (79-97%)	0.016 (0.004-0.038)	0.32% (0.19-0.48%)	0.2% (0.12-0.32%)	0.14 (0.072-0.23)	49 (26-77)
Palestine	1990	Female	20 to 24	12% (3.4-26%)	88% (74-97%)	0.025 (0.007-0.057)	0.63% (0.4-0.93%)	0.23% (0.12-0.37%)	0.17 (0.08-0.28)	92 (55-140)
Palestine	1990	Female	25 to 29	14% (3.8-30%)	86% (70-96%)	0.031 (0.0081-0.069)	0.64% (0.44-0.89%)	0.23% (0.096-0.37%)	0.16 (0.062-0.28)	85 (54-130)
Palestine	1990	Female	30 to 34	15% (3.9-32%)	85% (68-96%)	0.037 (0.011-0.076)	0.57% (0.37-0.81%)	0.26% (0.053-0.44%)	0.16 (0.033-0.28)	61 (37-94)
Palestine	1990	Female	35 to 39	17% (4.5-33%)	83% (67-95%)	0.043 (0.012-0.092)	0.43% (0.27-0.59%)	0.25% (0.034-0.45%)	0.18 (0.023-0.33)	43 (25-62)
Palestine	1990	Female	40 to 44	18% (6.1-34%)	82% (66-94%)	0.048 (0.015-0.1)	0.29% (0.12-0.45%)	0.19% (-0.072-0.39%)	0.16 (-0.058-0.35)	28 (11-45)
Palestine	1990	Female	45 to 49	18% (6.6-35%)	82% (65-93%)	0.051 (0.015-0.11)	0.14% (-0.12-0.31%)	0.03% (-0.39-0.27%)	0.032 (-0.38-0.3)	13 (-12-30)
Palestine	1990	Female	50 to 54	19% (7.6-37%)	81% (63-92%)	0.053 (0.015-0.12)	0.033% (-0.33-0.24%)	-0.057% (-0.57-0.25%)	-0.077 (-0.81-0.33)	3.1 (-31-23)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	1990	Female	55 to 59	20% (8-40%)	80% (60-92%)	0.053 (0.013-0.12)	-0.086% (-0.58-0.18%)	-0.17% (-0.86-0.21%)	-0.28 (-1.4-0.34)	-7.4 (-50-16)
Palestine	1990	Female	60 to 64	21% (7.9-41%)	79% (59-92%)	0.052 (0.011-0.13)	-0.17% (-0.74-0.16%)	-0.25% (-1.1-0.19%)	-0.51 (-2.2-0.41)	-14 (-64-14)
Palestine	1990	Female	65 to 69	22% (8.8-40%)	78% (60-91%)	0.046 (0.011-0.12)	-0.18% (-0.75-0.15%)	-0.23% (-0.96-0.18%)	-0.63 (-2.6-0.5)	-15 (-66-13)
Palestine	1990	Female	70 to 74	21% (8.4-38%)	79% (62-92%)	0.037 (0.0085-0.1)	-0.15% (-0.7-0.17%)	-0.18% (-0.89-0.2%)	-0.53 (-2.5-0.59)	-10 (-48-12)
Palestine	1990	Female	75 to 79	18% (6.5-38%)	82% (62-93%)	0.03 (0.0054-0.096)	-0.12% (-0.7-0.15%)	-0.14% (-0.85-0.17%)	-0.4 (-2.4-0.49)	-5.7 (-35-7.3)
Palestine	1990	Female	80 to 84	17% (5.9-37%)	83% (63-94%)	0.027 (0.0048-0.087)	-0.086% (-0.58-0.12%)	-0.1% (-0.67-0.14%)	-0.23 (-1.5-0.3)	-2.4 (-16-3.3)
Palestine	1990	Female	85 to 89	17% (6.1-38%)	83% (62-94%)	0.027 (0.0048-0.084)	-0.1% (-0.51-0.096%)	-0.13% (-0.61-0.1%)	-0.21 (-1-0.17)	-1.5 (-7.5-1.4)
Palestine	1990	Female	90 to 94	20% (6.7-43%)	80% (57-93%)	0.03 (0.0052-0.087)	-0.14% (-0.6-0.067%)	-0.18% (-0.72-0.066%)	-0.15 (-0.61-0.056)	-0.73 (-3.1-0.35)
Palestine	1990	Female	95 plus	24% (7.2-49%)	76% (51-93%)	0.035 (0.0055-0.11)	-0.19% (-0.79-0.066%)	-0.25% (-0.98-0.056%)	-0.081 (-0.32-0.018)	-0.25 (-1-0.09)
Palestine	1990	Male	All Ages	7.4% (4.9-10%)	93% (90-95%)	0.019 (0.011-0.03)	0.22% (0.14-0.33%)	0.098% (-0.024-0.2%)	4.4 (-1.2-9)	730 (470-1000)
Palestine	1990	Male	15 to 19	6.7% (2.8-14%)	93% (86-97%)	0.011 (0.0031-0.025)	0.42% (0.26-0.64%)	0.16% (0.071-0.28%)	0.21 (0.093-0.4)	80 (49-130)
Palestine	1990	Male	20 to 24	10% (4.2-20%)	90% (80-96%)	0.024 (0.0082-0.052)	0.81% (0.52-1.2%)	0.23% (0.11-0.45%)	0.36 (0.17-0.73)	160 (99-240)
Palestine	1990	Male	25 to 29	13% (5-24%)	87% (76-95%)	0.038 (0.013-0.079)	0.85% (0.6-1.2%)	0.31% (0.15-0.61%)	0.4 (0.19-0.75)	130 (89-190)
Palestine	1990	Male	30 to 34	15% (5.1-28%)	85% (72-95%)	0.049 (0.017-0.097)	0.78% (0.53-1.1%)	0.35% (0.16-0.61%)	0.34 (0.16-0.61)	92 (58-130)
Palestine	1990	Male	35 to 39	16% (6.2-29%)	84% (71-94%)	0.055 (0.019-0.11)	0.73% (0.5-1%)	0.43% (0.16-0.74%)	0.39 (0.14-0.69)	72 (47-100)
Palestine	1990	Male	40 to 44	16% (7-29%)	84% (71-93%)	0.059 (0.02-0.12)	0.56% (0.31-0.82%)	0.3% (-0.086-0.6%)	0.35 (-0.11-0.73)	61 (33-89)
Palestine	1990	Male	45 to 49	16% (6.3-29%)	84% (71-94%)	0.061 (0.019-0.13)	0.43% (0.097-0.66%)	0.27% (-0.24-0.61%)	0.37 (-0.34-0.91)	44 (10-70)
Palestine	1990	Male	50 to 54	15% (6.5-30%)	85% (70-94%)	0.06 (0.019-0.13)	0.29% (-0.041-0.52%)	0.18% (-0.35-0.53%)	0.3 (-0.57-0.9)	29 (-4.3-54)
Palestine	1990	Male	55 to 59	14% (6.4-27%)	86% (73-94%)	0.058 (0.016-0.13)	0.18% (-0.17-0.4%)	0.11% (-0.44-0.43%)	0.2 (-0.8-0.83)	16 (-16-38)
Palestine	1990	Male	60 to 64	14% (5.7-27%)	86% (73-94%)	0.053 (0.013-0.14)	0.14% (-0.23-0.37%)	0.1% (-0.41-0.43%)	0.19 (-0.83-0.82)	10 (-18-27)
Palestine	1990	Male	65 to 69	13% (5.8-25%)	87% (75-94%)	0.043 (0.012-0.1)	0.12% (-0.19-0.31%)	0.1% (-0.32-0.35%)	0.22 (-0.68-0.75)	7.6 (-13-19)
Palestine	1990	Male	70 to 74	12% (5.5-23%)	88% (77-95%)	0.033 (0.0084-0.083)	0.13% (-0.11-0.28%)	0.13% (-0.17-0.33%)	0.31 (-0.41-0.79)	7 (-6.2-16)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	1990	Male	75 to 79	11% (4.6-21%)	89% (79-95%)	0.025 (0.0056-0.071)	0.11% (-0.11-0.24%)	0.12% (-0.16-0.27%)	0.3 (-0.41-0.7)	4.9 (-4.8-10)
Palestine	1990	Male	80 to 84	9.5% (3.6-19%)	90% (81-96%)	0.021 (0.0046-0.055)	0.12% (-0.052-0.23%)	0.13% (-0.078-0.26%)	0.27 (-0.16-0.56)	3.1 (-1.3-6.1)
Palestine	1990	Male	85 to 89	9.3% (3.8-18%)	91% (82-96%)	0.019 (0.0043-0.049)	0.099% (-0.046-0.2%)	0.095% (-0.077-0.22%)	0.15 (-0.12-0.35)	1.3 (-0.63-2.7)
Palestine	1990	Male	90 to 94	10% (4.4-21%)	90% (79-96%)	0.02 (0.0045-0.056)	0.072% (-0.082-0.18%)	0.055% (-0.13-0.18%)	0.044 (-0.11-0.15)	0.34 (-0.38-0.85)
Palestine	1990	Male	95 plus	13% (4.8-27%)	87% (73-95%)	0.023 (0.0038-0.071)	0.068% (-0.14-0.18%)	0.024% (-0.23-0.15%)	0.0065 (-0.063-0.042)	0.071 (-0.15-0.19)
Palestine	1995	Female	All Ages	9.9% (6-16%)	90% (84-94%)	0.021 (0.015-0.027)	0.1% (0.011-0.19%)	-0.11% (-0.38-0.074%)	-4.7 (-17-3.1)	340 (34-600)
Palestine	1995	Female	15 to 19	9.6% (3.1-22%)	90% (78-97%)	0.019 (0.0054-0.043)	0.34% (0.2-0.53%)	0.24% (0.15-0.36%)	0.13 (0.083-0.21)	58 (33-93)
Palestine	1995	Female	20 to 24	12% (3.7-27%)	88% (73-96%)	0.03 (0.0094-0.062)	0.68% (0.44-1%)	0.26% (0.15-0.4%)	0.15 (0.085-0.23)	110 (62-170)
Palestine	1995	Female	25 to 29	14% (3.9-30%)	86% (70-96%)	0.036 (0.011-0.073)	0.69% (0.45-0.96%)	0.24% (0.12-0.37%)	0.15 (0.07-0.23)	100 (62-150)
Palestine	1995	Female	30 to 34	15% (4.2-33%)	85% (67-96%)	0.042 (0.015-0.08)	0.59% (0.37-0.84%)	0.27% (0.077-0.44%)	0.17 (0.048-0.29)	81 (46-120)
Palestine	1995	Female	35 to 39	17% (4.9-33%)	83% (67-95%)	0.05 (0.017-0.096)	0.45% (0.29-0.63%)	0.27% (0.056-0.48%)	0.18 (0.035-0.33)	51 (30-72)
Palestine	1995	Female	40 to 44	18% (6-33%)	82% (67-94%)	0.056 (0.019-0.11)	0.3% (0.12-0.47%)	0.2% (-0.079-0.42%)	0.17 (-0.068-0.36)	32 (13-52)
Palestine	1995	Female	45 to 49	18% (6.5-34%)	82% (66-93%)	0.06 (0.019-0.12)	0.13% (-0.12-0.31%)	0.01% (-0.41-0.28%)	0.011 (-0.44-0.3)	14 (-13-33)
Palestine	1995	Female	50 to 54	19% (7.5-37%)	81% (63-92%)	0.062 (0.019-0.13)	-0.003% (-0.4-0.24%)	-0.11% (-0.68-0.23%)	-0.16 (-1.1-0.34)	-0.29 (-41-26)
Palestine	1995	Female	55 to 59	20% (8.4-39%)	80% (61-92%)	0.062 (0.017-0.13)	-0.15% (-0.7-0.16%)	-0.26% (-1-0.17%)	-0.55 (-2.2-0.35)	-17 (-79-17)
Palestine	1995	Female	60 to 64	21% (8.6-41%)	79% (59-91%)	0.06 (0.014-0.14)	-0.25% (-0.94-0.12%)	-0.34% (-1.2-0.14%)	-0.92 (-3.3-0.37)	-27 (-97-13)
Palestine	1995	Female	65 to 69	22% (9.6-39%)	78% (61-90%)	0.053 (0.013-0.12)	-0.26% (-0.87-0.1%)	-0.32% (-1.1-0.12%)	-1.1 (-3.7-0.41)	-26 (-91-11)
Palestine	1995	Female	70 to 74	21% (8.7-37%)	79% (63-91%)	0.042 (0.01-0.11)	-0.21% (-0.85-0.11%)	-0.25% (-1-0.14%)	-1.1 (-4.3-0.6)	-21 (-84-11)
Palestine	1995	Female	75 to 79	19% (7.2-37%)	81% (63-93%)	0.034 (0.0068-0.095)	-0.17% (-0.8-0.13%)	-0.2% (-0.94-0.15%)	-0.76 (-3.5-0.56)	-11 (-51-8.9)
Palestine	1995	Female	80 to 84	17% (5.8-36%)	83% (64-94%)	0.031 (0.0063-0.088)	-0.13% (-0.61-0.1%)	-0.15% (-0.68-0.11%)	-0.41 (-2-0.3)	-4.4 (-22-3.3)
Palestine	1995	Female	85 to 89	17% (5.6-37%)	83% (63-94%)	0.03 (0.0062-0.089)	-0.14% (-0.6-0.077%)	-0.17% (-0.68-0.083%)	-0.34 (-1.5-0.16)	-2.5 (-11-1.4)
Palestine	1995	Female	90 to 94	20% (6.5-42%)	80% (58-94%)	0.034 (0.0068-0.097)	-0.19% (-0.68-0.06%)	-0.23% (-0.8-0.058%)	-0.24 (-0.84-0.062)	-1.2 (-4.2-0.37)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	1995	Female	95 plus	24% (7.4-48%)	76% (52-93%)	0.041 (0.007-0.12)	-0.24% (-0.94-0.064%)	-0.3% (-1.1-0.051%)	-0.11 (-0.43-0.019)	-0.36 (-1.4-0.098)
Palestine	1995	Male	All Ages	7.4% (5-11%)	93% (89-95%)	0.022 (0.016-0.03)	0.26% (0.16-0.36%)	0.1% (-0.042-0.23%)	5 (-2-11)	890 (540-1200)
Palestine	1995	Male	15 to 19	6.7% (2.8-13%)	93% (87-97%)	0.012 (0.0039-0.026)	0.45% (0.29-0.64%)	0.19% (0.085-0.32%)	0.3 (0.14-0.54)	100 (61-150)
Palestine	1995	Male	20 to 24	10% (4.1-20%)	90% (80-96%)	0.028 (0.01-0.054)	0.91% (0.6-1.3%)	0.3% (0.15-0.55%)	0.47 (0.23-0.87)	190 (120-280)
Palestine	1995	Male	25 to 29	13% (5.1-25%)	87% (75-95%)	0.043 (0.017-0.084)	0.95% (0.67-1.3%)	0.38% (0.18-0.7%)	0.49 (0.23-0.9)	170 (110-240)
Palestine	1995	Male	30 to 34	15% (5.3-28%)	85% (72-95%)	0.055 (0.022-0.1)	0.87% (0.61-1.2%)	0.41% (0.19-0.71%)	0.44 (0.2-0.76)	130 (84-180)
Palestine	1995	Male	35 to 39	16% (5.9-30%)	84% (70-94%)	0.063 (0.023-0.12)	0.78% (0.54-1.1%)	0.48% (0.17-0.86%)	0.46 (0.16-0.81)	90 (59-130)
Palestine	1995	Male	40 to 44	16% (7.1-28%)	84% (72-93%)	0.067 (0.025-0.13)	0.58% (0.32-0.84%)	0.31% (-0.15-0.68%)	0.33 (-0.16-0.72)	62 (33-93)
Palestine	1995	Male	45 to 49	16% (6.1-29%)	84% (71-94%)	0.07 (0.023-0.15)	0.43% (0.11-0.68%)	0.26% (-0.28-0.65%)	0.42 (-0.46-1.1)	54 (12-88)
Palestine	1995	Male	50 to 54	15% (6.4-30%)	85% (70-94%)	0.069 (0.023-0.15)	0.28% (-0.098-0.52%)	0.16% (-0.44-0.54%)	0.3 (-0.86-1)	32 (-12-63)
Palestine	1995	Male	55 to 59	14% (6.2-28%)	86% (72-94%)	0.066 (0.02-0.14)	0.16% (-0.25-0.41%)	0.076% (-0.52-0.45%)	0.18 (-1.2-1.1)	19 (-29-49)
Palestine	1995	Male	60 to 64	14% (5.9-27%)	86% (73-94%)	0.059 (0.016-0.14)	0.12% (-0.3-0.36%)	0.071% (-0.51-0.41%)	0.19 (-1.4-1.1)	12 (-31-36)
Palestine	1995	Male	65 to 69	13% (5.7-25%)	87% (75-94%)	0.049 (0.014-0.11)	0.098% (-0.22-0.3%)	0.076% (-0.36-0.34%)	0.21 (-1-0.95)	8.3 (-20-25)
Palestine	1995	Male	70 to 74	12% (5.3-22%)	88% (78-95%)	0.036 (0.011-0.086)	0.11% (-0.15-0.27%)	0.11% (-0.22-0.31%)	0.34 (-0.71-1)	7.9 (-12-20)
Palestine	1995	Male	75 to 79	11% (4.5-20%)	89% (80-96%)	0.027 (0.0068-0.073)	0.1% (-0.14-0.23%)	0.1% (-0.18-0.25%)	0.32 (-0.58-0.8)	5.3 (-7.2-12)
Palestine	1995	Male	80 to 84	9.5% (3.7-19%)	91% (81-96%)	0.023 (0.0056-0.06)	0.11% (-0.079-0.22%)	0.11% (-0.11-0.24%)	0.3 (-0.28-0.63)	3.5 (-2.4-6.9)
Palestine	1995	Male	85 to 89	9.4% (3.7-18%)	91% (82-96%)	0.022 (0.0053-0.055)	0.09% (-0.079-0.19%)	0.083% (-0.11-0.2%)	0.16 (-0.21-0.4)	1.5 (-1.3-3.1)
Palestine	1995	Male	90 to 94	11% (4.2-22%)	89% (78-96%)	0.023 (0.0053-0.063)	0.06% (-0.12-0.17%)	0.042% (-0.16-0.17%)	0.041 (-0.16-0.17)	0.34 (-0.66-0.96)
Palestine	1995	Male	95 plus	13% (4.7-26%)	87% (74-95%)	0.026 (0.0047-0.077)	0.051% (-0.18-0.17%)	0.0064% (-0.27-0.15%)	0.002 (-0.084-0.047)	0.061 (-0.22-0.21)
Palestine	2000	Female	All Ages	9.8% (5.6-16%)	90% (84-94%)	0.035 (0.028-0.042)	0.061% (-0.08-0.17%)	-0.26% (-0.73-0.033%)	-14 (-40-1.8)	230 (-300-640)
Palestine	2000	Female	15 to 19	9.3% (2.8-21%)	91% (79-97%)	0.031 (0.0091-0.071)	0.35% (0.21-0.54%)	0.21% (0.13-0.33%)	0.15 (0.093-0.22)	69 (39-110)
Palestine	2000	Female	20 to 24	12% (3.4-26%)	88% (74-97%)	0.049 (0.015-0.1)	0.68% (0.42-1%)	0.24% (0.14-0.38%)	0.16 (0.096-0.25)	130 (75-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	2000	Female	25 to 29	14% (3.6-29%)	86% (71-96%)	0.059 (0.017-0.12)	0.69% (0.46-0.95%)	0.23% (0.099-0.4%)	0.16 (0.068-0.27)	120 (77-180)
Palestine	2000	Female	30 to 34	15% (4-33%)	85% (67-96%)	0.069 (0.023-0.13)	0.6% (0.39-0.87%)	0.27% (0.071-0.47%)	0.19 (0.05-0.32)	99 (59-150)
Palestine	2000	Female	35 to 39	17% (4.5-33%)	83% (67-95%)	0.083 (0.028-0.15)	0.45% (0.27-0.65%)	0.3% (0.047-0.56%)	0.23 (0.035-0.43)	68 (37-100)
Palestine	2000	Female	40 to 44	18% (6.3-33%)	82% (67-94%)	0.092 (0.034-0.18)	0.29% (0.087-0.48%)	0.22% (-0.11-0.54%)	0.18 (-0.088-0.43)	36 (10-62)
Palestine	2000	Female	45 to 49	18% (6.3-34%)	82% (66-94%)	0.1 (0.032-0.2)	0.083% (-0.21-0.3%)	-0.044% (-0.56-0.32%)	-0.048 (-0.62-0.34)	10 (-24-37)
Palestine	2000	Female	50 to 54	19% (7.4-37%)	81% (63-93%)	0.1 (0.033-0.22)	-0.1% (-0.63-0.2%)	-0.24% (-1.1-0.22%)	-0.39 (-1.7-0.36)	-13 (-77-24)
Palestine	2000	Female	55 to 59	20% (7.7-39%)	80% (61-92%)	0.11 (0.03-0.22)	-0.33% (-1.1-0.12%)	-0.49% (-1.6-0.13%)	-1.2 (-3.8-0.29)	-42 (-140-15)
Palestine	2000	Female	60 to 64	21% (8.1-41%)	79% (59-92%)	0.1 (0.025-0.25)	-0.5% (-1.6-0.062%)	-0.65% (-2-0.063%)	-2.4 (-7.4-0.23)	-71 (-220-8.9)
Palestine	2000	Female	65 to 69	22% (9.1-39%)	78% (61-91%)	0.089 (0.023-0.2)	-0.52% (-1.5-0.037%)	-0.64% (-1.8-0.044%)	-2.9 (-8.3-0.2)	-72 (-200-5)
Palestine	2000	Female	70 to 74	20% (8.6-38%)	80% (62-91%)	0.07 (0.017-0.18)	-0.44% (-1.4-0.039%)	-0.52% (-1.7-0.047%)	-3 (-9.7-0.27)	-57 (-180-4.9)
Palestine	2000	Female	75 to 79	18% (6.4-39%)	82% (61-94%)	0.057 (0.011-0.16)	-0.35% (-1.4-0.063%)	-0.42% (-1.7-0.076%)	-2.2 (-9.1-0.42)	-33 (-130-6)
Palestine	2000	Female	80 to 84	17% (5.5-37%)	83% (63-94%)	0.05 (0.01-0.14)	-0.28% (-1.1-0.058%)	-0.32% (-1.3-0.067%)	-1.3 (-5-0.26)	-14 (-54-2.8)
Palestine	2000	Female	85 to 89	17% (5.5-37%)	83% (63-94%)	0.05 (0.011-0.15)	-0.29% (-1.1-0.032%)	-0.34% (-1.2-0.032%)	-0.82 (-3-0.079)	-6.1 (-23-0.68)
Palestine	2000	Female	90 to 94	19% (6-40%)	81% (60-94%)	0.057 (0.011-0.16)	-0.36% (-1.1--0.0053%)	-0.43% (-1.4--0.016%)	-0.54 (-1.7--0.02)	-2.7 (-8.5--0.04)
Palestine	2000	Female	95 plus	24% (7.1-47%)	76% (53-93%)	0.068 (0.012-0.2)	-0.45% (-1.6--0.0028%)	-0.55% (-1.9--0.024%)	-0.26 (-0.87--0.011)	-0.83 (-2.9--0.0053)
Palestine	2000	Male	All Ages	7.5% (5.1-11%)	92% (89-95%)	0.038 (0.032-0.046)	0.25% (0.13-0.36%)	0.054% (-0.15-0.23%)	3.6 (-9.7-15)	1100 (580-1600)
Palestine	2000	Male	15 to 19	6.7% (2.8-13%)	93% (87-97%)	0.02 (0.0065-0.042)	0.43% (0.27-0.6%)	0.22% (0.093-0.37%)	0.59 (0.25-1)	140 (88-210)
Palestine	2000	Male	20 to 24	10% (4-20%)	90% (80-96%)	0.046 (0.017-0.089)	0.82% (0.53-1.2%)	0.31% (0.14-0.59%)	0.86 (0.42-1.6)	260 (170-390)
Palestine	2000	Male	25 to 29	13% (5.1-25%)	87% (75-95%)	0.073 (0.03-0.13)	0.88% (0.61-1.2%)	0.38% (0.16-0.72%)	0.78 (0.33-1.4)	220 (140-310)
Palestine	2000	Male	30 to 34	15% (5.3-28%)	85% (72-95%)	0.094 (0.04-0.17)	0.83% (0.57-1.1%)	0.4% (0.16-0.74%)	0.64 (0.26-1.1)	160 (110-230)
Palestine	2000	Male	35 to 39	16% (6.1-29%)	84% (71-94%)	0.11 (0.042-0.19)	0.75% (0.5-1%)	0.45% (0.091-0.78%)	0.65 (0.13-1.1)	130 (81-180)
Palestine	2000	Male	40 to 44	16% (7.1-28%)	84% (72-93%)	0.11 (0.044-0.21)	0.54% (0.21-0.81%)	0.23% (-0.36-0.65%)	0.33 (-0.49-0.93)	75 (29-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	2000	Male	45 to 49	16% (6.3-29%)	84% (71-94%)	0.12 (0.041-0.24)	0.35% (-0.042-0.66%)	0.14% (-0.54-0.61%)	0.26 (-0.99-1.1)	49 (-5.8-93)
Palestine	2000	Male	50 to 54	15% (6.5-29%)	85% (71-94%)	0.12 (0.04-0.24)	0.18% (-0.32-0.48%)	0.011% (-0.82-0.48%)	0.029 (-2.2-1.3)	29 (-52-78)
Palestine	2000	Male	55 to 59	14% (6.3-28%)	86% (72-94%)	0.11 (0.036-0.23)	0.059% (-0.55-0.4%)	-0.073% (-0.93-0.42%)	-0.22 (-2.9-1.3)	8.7 (-80-59)
Palestine	2000	Male	60 to 64	14% (6-27%)	86% (73-94%)	0.1 (0.03-0.23)	0.0079% (-0.53-0.34%)	-0.076% (-0.84-0.38%)	-0.29 (-3.2-1.4)	1.2 (-77-49)
Palestine	2000	Male	65 to 69	13% (6.1-25%)	87% (75-94%)	0.082 (0.026-0.17)	-0.02% (-0.51-0.26%)	-0.074% (-0.7-0.29%)	-0.33 (-3.1-1.3)	-2.6 (-66-34)
Palestine	2000	Male	70 to 74	12% (5.7-22%)	88% (78-94%)	0.061 (0.019-0.14)	0.0093% (-0.4-0.25%)	-0.014% (-0.5-0.27%)	-0.073 (-2.6-1.4)	1.1 (-45-28)
Palestine	2000	Male	75 to 79	11% (4.6-20%)	89% (80-95%)	0.047 (0.012-0.12)	0.027% (-0.32-0.22%)	0.012% (-0.39-0.24%)	0.061 (-1.9-1.2)	2.1 (-25-17)
Palestine	2000	Male	80 to 84	9.5% (4-19%)	90% (81-96%)	0.04 (0.01-0.1)	0.064% (-0.21-0.23%)	0.058% (-0.25-0.25%)	0.23 (-0.97-0.97)	2.9 (-9.6-10)
Palestine	2000	Male	85 to 89	9.3% (3.7-18%)	91% (82-96%)	0.036 (0.0098-0.093)	0.048% (-0.21-0.2%)	0.036% (-0.25-0.21%)	0.093 (-0.65-0.53)	1 (-4.3-4.2)
Palestine	2000	Male	90 to 94	10% (4.3-21%)	90% (79-96%)	0.038 (0.0097-0.1)	0.011% (-0.27-0.17%)	-0.013% (-0.34-0.16%)	-0.017 (-0.42-0.2)	0.077 (-1.9-1.2)
Palestine	2000	Male	95 plus	13% (4.7-28%)	87% (72-95%)	0.044 (0.0085-0.13)	-0.017% (-0.41-0.16%)	-0.069% (-0.52-0.14%)	-0.027 (-0.21-0.055)	-0.025 (-0.6-0.24)
Palestine	2005	Female	All Ages	10% (6.1-15%)	90% (85-94%)	0.049 (0.04-0.058)	0.05% (-0.11-0.18%)	-0.35% (-0.91-0.011%)	-20 (-51-0.54)	200 (-420-690)
Palestine	2005	Female	15 to 19	9.2% (3-20%)	91% (80-97%)	0.041 (0.012-0.094)	0.37% (0.22-0.59%)	0.27% (0.17-0.43%)	0.17 (0.11-0.27)	81 (44-130)
Palestine	2005	Female	20 to 24	12% (3.7-26%)	88% (74-96%)	0.066 (0.023-0.13)	0.73% (0.46-1.1%)	0.31% (0.16-0.5%)	0.18 (0.093-0.29)	140 (84-230)
Palestine	2005	Female	25 to 29	13% (3.9-28%)	87% (72-96%)	0.08 (0.026-0.15)	0.71% (0.47-1%)	0.26% (0.093-0.45%)	0.16 (0.055-0.27)	140 (82-210)
Palestine	2005	Female	30 to 34	15% (4-31%)	85% (69-96%)	0.095 (0.032-0.18)	0.61% (0.39-0.89%)	0.29% (0.043-0.53%)	0.19 (0.027-0.34)	110 (62-170)
Palestine	2005	Female	35 to 39	17% (4.6-33%)	83% (67-95%)	0.11 (0.037-0.2)	0.46% (0.27-0.67%)	0.38% (0.092-0.68%)	0.29 (0.068-0.52)	78 (43-120)
Palestine	2005	Female	40 to 44	18% (6.6-34%)	82% (66-93%)	0.13 (0.049-0.24)	0.28% (0.058-0.49%)	0.27% (-0.14-0.63%)	0.27 (-0.14-0.65)	45 (8.4-83)
Palestine	2005	Female	45 to 49	18% (6.1-35%)	82% (65-94%)	0.14 (0.045-0.28)	0.07% (-0.29-0.31%)	-0.02% (-0.64-0.41%)	-0.024 (-0.77-0.49)	9.6 (-37-43)
Palestine	2005	Female	50 to 54	19% (7.3-37%)	81% (63-93%)	0.14 (0.046-0.3)	-0.11% (-0.65-0.21%)	-0.21% (-1-0.29%)	-0.34 (-1.6-0.46)	-15 (-84-28)
Palestine	2005	Female	55 to 59	20% (8.4-38%)	80% (62-92%)	0.14 (0.042-0.3)	-0.35% (-1.1-0.097%)	-0.49% (-1.6-0.15%)	-1.2 (-3.9-0.38)	-48 (-160-14)
Palestine	2005	Female	60 to 64	21% (8.1-39%)	79% (61-92%)	0.14 (0.034-0.31)	-0.53% (-1.5-0.037%)	-0.69% (-2-0.066%)	-2.5 (-7.2-0.24)	-80 (-230-5.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	2005	Female	65 to 69	21% (8.8-39%)	79% (61-91%)	0.12 (0.03-0.26)	-0.61% (-1.6--0.021%)	-0.75% (-2-0.021%)	-3.8 (-10-0.1)	-96 (-250-3.2)
Palestine	2005	Female	70 to 74	20% (8.8-37%)	80% (63-91%)	0.094 (0.024-0.24)	-0.56% (-1.6--0.015%)	-0.66% (-1.9--0.017%)	-4.1 (-12--0.1)	-79 (-230--2.2)
Palestine	2005	Female	75 to 79	18% (7-36%)	82% (64-93%)	0.076 (0.017-0.22)	-0.49% (-1.7-0.0083%)	-0.57% (-2-0.014%)	-3.5 (-12-0.09)	-51 (-170-0.87)
Palestine	2005	Female	80 to 84	17% (5.8-36%)	83% (64-94%)	0.069 (0.015-0.2)	-0.44% (-1.5-0.016%)	-0.51% (-1.7-0.017%)	-2.4 (-8.2-0.079)	-26 (-89-0.9)
Palestine	2005	Female	85 to 89	17% (5.8-37%)	83% (63-94%)	0.07 (0.015-0.21)	-0.45% (-1.5--0.017%)	-0.53% (-1.8--0.023%)	-1.7 (-5.8--0.077)	-13 (-43--0.49)
Palestine	2005	Female	90 to 94	20% (7.1-42%)	80% (58-93%)	0.08 (0.018-0.23)	-0.52% (-1.6--0.054%)	-0.64% (-1.9--0.08%)	-0.89 (-2.6--0.11)	-4.5 (-13--0.48)
Palestine	2005	Female	95 plus	25% (7.8-49%)	75% (51-92%)	0.095 (0.018-0.28)	-0.64% (-2--0.057%)	-0.79% (-2.4--0.089%)	-0.41 (-1.3--0.046)	-1.3 (-4.2--0.12)
Palestine	2005	Male	All Ages	7.8% (5.1-11%)	92% (89-95%)	0.054 (0.046-0.063)	0.27% (0.14-0.41%)	0.066% (-0.19-0.28%)	5.1 (-14-22)	1300 (680-2000)
Palestine	2005	Male	15 to 19	6.7% (2.8-14%)	93% (86-97%)	0.028 (0.0094-0.06)	0.48% (0.31-0.7%)	0.31% (0.12-0.56%)	0.99 (0.39-1.8)	190 (120-280)
Palestine	2005	Male	20 to 24	10% (4.1-21%)	90% (79-96%)	0.064 (0.024-0.12)	0.97% (0.65-1.4%)	0.52% (0.22-1%)	1.5 (0.64-3)	320 (210-480)
Palestine	2005	Male	25 to 29	13% (5-25%)	87% (75-95%)	0.1 (0.041-0.18)	1% (0.68-1.4%)	0.58% (0.22-1.1%)	1.3 (0.51-2.5)	270 (180-380)
Palestine	2005	Male	30 to 34	15% (5.3-29%)	85% (71-95%)	0.13 (0.056-0.22)	0.9% (0.6-1.3%)	0.54% (0.18-1%)	0.98 (0.31-1.9)	200 (130-290)
Palestine	2005	Male	35 to 39	16% (5.9-30%)	84% (70-94%)	0.15 (0.059-0.26)	0.79% (0.5-1.1%)	0.52% (0.024-1%)	0.88 (0.04-1.7)	160 (95-220)
Palestine	2005	Male	40 to 44	16% (7.1-29%)	84% (71-93%)	0.16 (0.063-0.29)	0.55% (0.2-0.86%)	0.27% (-0.39-0.73%)	0.51 (-0.74-1.4)	100 (35-160)
Palestine	2005	Male	45 to 49	16% (6.5-29%)	84% (71-94%)	0.16 (0.055-0.33)	0.34% (-0.14-0.68%)	0.15% (-0.65-0.68%)	0.36 (-1.6-1.7)	59 (-23-120)
Palestine	2005	Male	50 to 54	15% (6.5-30%)	85% (70-93%)	0.16 (0.054-0.33)	0.16% (-0.43-0.51%)	0.012% (-0.89-0.53%)	0.035 (-2.6-1.5)	28 (-70-86)
Palestine	2005	Male	55 to 59	14% (6.2-29%)	86% (71-94%)	0.15 (0.05-0.32)	0.056% (-0.55-0.43%)	-0.06% (-0.95-0.48%)	-0.24 (-3.8-1.9)	11 (-100-80)
Palestine	2005	Male	60 to 64	14% (6-27%)	86% (73-94%)	0.14 (0.04-0.32)	0.015% (-0.57-0.38%)	-0.056% (-0.84-0.45%)	-0.25 (-3.8-2)	2.5 (-99-63)
Palestine	2005	Male	65 to 69	13% (5.8-25%)	87% (75-94%)	0.11 (0.035-0.24)	-0.045% (-0.57-0.28%)	-0.097% (-0.78-0.32%)	-0.56 (-4.5-1.8)	-7.5 (-95-47)
Palestine	2005	Male	70 to 74	12% (5.3-21%)	88% (79-95%)	0.082 (0.025-0.2)	-0.0031% (-0.46-0.27%)	-0.026% (-0.58-0.3%)	-0.17 (-3.9-2.1)	-0.45 (-67-40)
Palestine	2005	Male	75 to 79	10% (4.3-20%)	90% (80-96%)	0.062 (0.017-0.16)	-0.0045% (-0.38-0.22%)	-0.022% (-0.46-0.24%)	-0.14 (-3-1.5)	-0.47 (-39-23)
Palestine	2005	Male	80 to 84	9.2% (3.7-18%)	91% (82-96%)	0.052 (0.014-0.14)	0.03% (-0.3-0.24%)	0.02% (-0.34-0.25%)	0.094 (-1.6-1.2)	1.6 (-16-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	2005	Male	85 to 89	9% (3.6-18%)	91% (82-96%)	0.048 (0.013-0.12)	0.0013% (-0.32-0.18%)	-0.016% (-0.38-0.19%)	-0.054 (-1.2-0.62)	0.031 (-8.4-4.9)
Palestine	2005	Male	90 to 94	10% (4.2-21%)	90% (79-96%)	0.051 (0.013-0.13)	-0.043% (-0.38-0.15%)	-0.074% (-0.46-0.14%)	-0.11 (-0.69-0.21)	-0.37 (-3.2-1.2)
Palestine	2005	Male	95 plus	13% (4.7-27%)	87% (73-95%)	0.059 (0.011-0.17)	-0.089% (-0.56-0.14%)	-0.15% (-0.7-0.11%)	-0.069 (-0.32-0.051)	-0.15 (-0.97-0.24)
Palestine	2010	Female	All Ages	10% (6-16%)	90% (84-94%)	0.042 (0.033-0.051)	0.048% (-0.11-0.17%)	-0.35% (-0.91--0.011%)	-26 (-65--0.76)	230 (-510-840)
Palestine	2010	Female	15 to 19	9.2% (2.9-21%)	91% (79-97%)	0.035 (0.01-0.082)	0.36% (0.21-0.56%)	0.25% (0.15-0.4%)	0.18 (0.11-0.28)	96 (52-160)
Palestine	2010	Female	20 to 24	12% (3.5-26%)	88% (74-96%)	0.057 (0.018-0.12)	0.71% (0.45-1%)	0.28% (0.15-0.45%)	0.19 (0.1-0.31)	170 (100-270)
Palestine	2010	Female	25 to 29	14% (3.8-29%)	86% (71-96%)	0.069 (0.022-0.14)	0.72% (0.48-1%)	0.24% (0.11-0.41%)	0.15 (0.068-0.26)	160 (98-240)
Palestine	2010	Female	30 to 34	15% (4.1-32%)	85% (68-96%)	0.082 (0.027-0.15)	0.61% (0.38-0.88%)	0.26% (0.058-0.47%)	0.19 (0.042-0.33)	130 (74-200)
Palestine	2010	Female	35 to 39	17% (4.7-33%)	83% (67-95%)	0.097 (0.032-0.18)	0.46% (0.26-0.65%)	0.36% (0.13-0.61%)	0.33 (0.11-0.56)	90 (51-140)
Palestine	2010	Female	40 to 44	18% (6.7-32%)	82% (68-93%)	0.11 (0.043-0.2)	0.28% (0.062-0.48%)	0.25% (-0.11-0.55%)	0.31 (-0.14-0.69)	55 (12-98)
Palestine	2010	Female	45 to 49	18% (6.3-34%)	82% (66-94%)	0.12 (0.04-0.23)	0.081% (-0.2-0.31%)	-0.013% (-0.5-0.36%)	-0.021 (-0.84-0.6)	15 (-37-59)
Palestine	2010	Female	50 to 54	19% (7.5-36%)	81% (64-92%)	0.12 (0.038-0.25)	-0.09% (-0.54-0.2%)	-0.19% (-0.88-0.26%)	-0.39 (-1.8-0.51)	-15 (-91-34)
Palestine	2010	Female	55 to 59	20% (8.2-39%)	80% (61-92%)	0.12 (0.034-0.26)	-0.31% (-1.1-0.11%)	-0.45% (-1.5-0.14%)	-1.4 (-4.5-0.42)	-53 (-180-18)
Palestine	2010	Female	60 to 64	21% (8.4-41%)	79% (59-92%)	0.11 (0.028-0.28)	-0.49% (-1.5-0.056%)	-0.64% (-2-0.058%)	-2.6 (-8.1-0.24)	-84 (-250-9.5)
Palestine	2010	Female	65 to 69	21% (8.9-39%)	79% (61-91%)	0.1 (0.027-0.22)	-0.58% (-1.6-0.011%)	-0.72% (-1.9-0.024%)	-4.5 (-12-0.15)	-110 (-290-2)
Palestine	2010	Female	70 to 74	20% (8.5-37%)	80% (63-92%)	0.079 (0.02-0.2)	-0.52% (-1.5-0.0044%)	-0.63% (-1.8-0.0081%)	-4.9 (-14-0.063)	-95 (-270-0.78)
Palestine	2010	Female	75 to 79	18% (7-36%)	82% (64-93%)	0.064 (0.014-0.18)	-0.49% (-1.6-0.0067%)	-0.57% (-1.9-0.0038%)	-4.7 (-15-0.031)	-67 (-220-0.89)
Palestine	2010	Female	80 to 84	16% (5.6-36%)	84% (64-94%)	0.058 (0.012-0.16)	-0.46% (-1.7-0.008%)	-0.53% (-1.9-0.0043%)	-3.7 (-14-0.03)	-39 (-140-0.65)
Palestine	2010	Female	85 to 89	17% (5.5-38%)	83% (62-95%)	0.058 (0.012-0.18)	-0.47% (-1.6--0.027%)	-0.55% (-1.8--0.041%)	-2.7 (-9--0.2)	-20 (-67--1.1)
Palestine	2010	Female	90 to 94	19% (6.7-41%)	81% (59-93%)	0.066 (0.014-0.19)	-0.51% (-1.5--0.05%)	-0.62% (-1.8--0.076%)	-1.5 (-4.3--0.18)	-7.3 (-22--0.72)
Palestine	2010	Female	95 plus	24% (7.7-48%)	76% (52-92%)	0.078 (0.015-0.22)	-0.61% (-2--0.045%)	-0.75% (-2.3--0.072%)	-0.57 (-1.8--0.055)	-1.9 (-5.9--0.14)
Palestine	2010	Male	All Ages	7.9% (5.3-11%)	92% (89-95%)	0.046 (0.038-0.056)	0.26% (0.14-0.39%)	0.048% (-0.18-0.25%)	4.4 (-17-22)	1500 (730-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	2010	Male	15 to 19	6.8% (2.9-14%)	93% (86-97%)	0.024 (0.008-0.052)	0.49% (0.31-0.72%)	0.25% (0.12-0.46%)	0.67 (0.31-1.2)	190 (120-290)
Palestine	2010	Male	20 to 24	11% (4.1-20%)	89% (80-96%)	0.054 (0.021-0.1)	1% (0.7-1.5%)	0.46% (0.2-0.91%)	1.1 (0.48-2.2)	350 (220-540)
Palestine	2010	Male	25 to 29	13% (5.1-25%)	87% (75-95%)	0.085 (0.034-0.16)	1.1% (0.75-1.5%)	0.56% (0.2-1.1%)	1 (0.37-1.9)	290 (180-420)
Palestine	2010	Male	30 to 34	15% (5.3-29%)	85% (71-95%)	0.11 (0.047-0.19)	0.93% (0.62-1.3%)	0.53% (0.16-1%)	0.88 (0.26-1.7)	220 (140-330)
Palestine	2010	Male	35 to 39	16% (6.2-30%)	84% (70-94%)	0.12 (0.047-0.22)	0.81% (0.53-1.1%)	0.54% (0.085-1%)	0.9 (0.14-1.7)	180 (110-250)
Palestine	2010	Male	40 to 44	16% (7.4-29%)	84% (71-93%)	0.13 (0.054-0.25)	0.59% (0.28-0.89%)	0.32% (-0.25-0.77%)	0.66 (-0.52-1.6)	130 (58-200)
Palestine	2010	Male	45 to 49	16% (6.6-29%)	84% (71-93%)	0.14 (0.047-0.28)	0.38% (-0.032-0.67%)	0.19% (-0.48-0.67%)	0.6 (-1.5-2.1)	86 (-6.7-160)
Palestine	2010	Male	50 to 54	15% (6.5-30%)	85% (70-93%)	0.14 (0.047-0.28)	0.17% (-0.35-0.5%)	0.027% (-0.75-0.52%)	0.099 (-2.7-1.9)	37 (-71-100)
Palestine	2010	Male	55 to 59	15% (6.4-28%)	85% (72-94%)	0.13 (0.042-0.27)	0.046% (-0.51-0.39%)	-0.07% (-0.86-0.43%)	-0.31 (-3.8-1.9)	9.3 (-100-80)
Palestine	2010	Male	60 to 64	14% (6-27%)	86% (73-94%)	0.12 (0.034-0.26)	0.0082% (-0.57-0.36%)	-0.065% (-0.84-0.42%)	-0.39 (-5.1-2.5)	1.9 (-130-80)
Palestine	2010	Male	65 to 69	14% (6.3-26%)	86% (74-94%)	0.096 (0.031-0.21)	-0.024% (-0.54-0.28%)	-0.074% (-0.74-0.33%)	-0.48 (-4.9-2.2)	-4.6 (-100-55)
Palestine	2010	Male	70 to 74	12% (5.8-22%)	88% (78-94%)	0.071 (0.023-0.17)	0.028% (-0.39-0.28%)	0.01% (-0.5-0.32%)	0.078 (-3.8-2.4)	4.8 (-65-49)
Palestine	2010	Male	75 to 79	11% (4.6-20%)	89% (80-95%)	0.054 (0.015-0.14)	0.0058% (-0.37-0.22%)	-0.012% (-0.45-0.24%)	-0.095 (-3.6-1.9)	0.75 (-48-29)
Palestine	2010	Male	80 to 84	9.5% (3.7-19%)	91% (81-96%)	0.045 (0.012-0.12)	0.022% (-0.3-0.22%)	0.011% (-0.34-0.24%)	0.071 (-2.2-1.5)	1.6 (-22-17)
Palestine	2010	Male	85 to 89	9.3% (3.7-18%)	91% (82-96%)	0.042 (0.011-0.11)	-0.016% (-0.33-0.16%)	-0.035% (-0.38-0.16%)	-0.16 (-1.7-0.74)	-0.59 (-12-5.8)
Palestine	2010	Male	90 to 94	10% (4.2-21%)	90% (79-96%)	0.044 (0.011-0.11)	-0.055% (-0.39-0.13%)	-0.085% (-0.47-0.12%)	-0.18 (-0.97-0.25)	-0.64 (-4.6-1.5)
Palestine	2010	Male	95 plus	13% (4.7-26%)	87% (74-95%)	0.05 (0.0097-0.13)	-0.09% (-0.55-0.13%)	-0.15% (-0.69-0.1%)	-0.087 (-0.41-0.061)	-0.2 (-1.2-0.28)
Palestine	2016	Female	All Ages	10% (6-16%)	90% (84-94%)	0.04 (0.026-0.061)	0.031% (-0.16-0.18%)	-0.41% (-1--0.021%)	-39 (-97--2)	180 (-880-1100)
Palestine	2016	Female	15 to 19	9.4% (3.2-21%)	91% (79-97%)	0.033 (0.0093-0.081)	0.36% (0.21-0.54%)	0.26% (0.15-0.4%)	0.24 (0.14-0.39)	110 (59-170)
Palestine	2016	Female	20 to 24	12% (3.8-26%)	88% (74-96%)	0.053 (0.015-0.12)	0.71% (0.44-1%)	0.28% (0.15-0.46%)	0.28 (0.15-0.45)	220 (130-360)
Palestine	2016	Female	25 to 29	14% (3.7-29%)	86% (71-96%)	0.064 (0.018-0.13)	0.73% (0.49-1%)	0.24% (0.11-0.4%)	0.23 (0.1-0.39)	210 (130-320)
Palestine	2016	Female	30 to 34	15% (3.8-33%)	85% (67-96%)	0.075 (0.021-0.15)	0.62% (0.38-0.87%)	0.27% (0.05-0.48%)	0.28 (0.053-0.49)	160 (92-250)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	2016	Female	35 to 39	17% (4.6-34%)	83% (66-95%)	0.089 (0.028-0.18)	0.46% (0.27-0.66%)	0.36% (0.11-0.64%)	0.48 (0.15-0.83)	120 (64-180)
Palestine	2016	Female	40 to 44	18% (6.5-34%)	82% (66-93%)	0.1 (0.034-0.21)	0.28% (0.039-0.48%)	0.23% (-0.12-0.55%)	0.41 (-0.22-0.96)	68 (10-120)
Palestine	2016	Female	45 to 49	19% (6.9-35%)	81% (65-93%)	0.11 (0.034-0.22)	0.066% (-0.3-0.3%)	-0.034% (-0.57-0.34%)	-0.082 (-1.4-0.82)	16 (-71-77)
Palestine	2016	Female	50 to 54	19% (7.7-37%)	81% (63-92%)	0.11 (0.032-0.25)	-0.12% (-0.65-0.21%)	-0.23% (-1-0.22%)	-0.79 (-3.5-0.78)	-30 (-170-53)
Palestine	2016	Female	55 to 59	20% (8.2-39%)	80% (61-92%)	0.11 (0.028-0.24)	-0.34% (-1.1-0.11%)	-0.47% (-1.6-0.14%)	-2.1 (-7-0.66)	-81 (-270-26)
Palestine	2016	Female	60 to 64	21% (8.4-40%)	79% (60-92%)	0.11 (0.025-0.26)	-0.49% (-1.5-0.06%)	-0.63% (-1.8-0.076%)	-3.5 (-10-0.43)	-110 (-320-13)
Palestine	2016	Female	65 to 69	22% (9.3-39%)	78% (61-91%)	0.093 (0.025-0.21)	-0.62% (-1.7--0.0035%)	-0.75% (-2--0.0081%)	-6.2 (-17--0.064)	-150 (-420--0.81)
Palestine	2016	Female	70 to 74	20% (8.3-37%)	80% (63-92%)	0.075 (0.016-0.19)	-0.56% (-1.6-0.014%)	-0.66% (-1.9-0.014%)	-6.9 (-19-0.14)	-130 (-370-3.3)
Palestine	2016	Female	75 to 79	18% (6.7-37%)	82% (63-93%)	0.062 (0.012-0.18)	-0.54% (-1.7-0.011%)	-0.62% (-2-0.011%)	-7.5 (-24-0.13)	-110 (-350-2.3)
Palestine	2016	Female	80 to 84	17% (5.9-37%)	83% (63-94%)	0.056 (0.01-0.17)	-0.51% (-1.8-0.0016%)	-0.58% (-2--0.00035%)	-6.4 (-22--0.0039)	-66 (-230-0.21)
Palestine	2016	Female	85 to 89	18% (5.8-39%)	82% (61-94%)	0.056 (0.011-0.18)	-0.52% (-1.7--0.033%)	-0.6% (-2--0.043%)	-4.1 (-13--0.28)	-30 (-99--1.7)
Palestine	2016	Female	90 to 94	20% (6.6-42%)	80% (58-93%)	0.064 (0.012-0.19)	-0.56% (-1.7--0.051%)	-0.66% (-1.9--0.07%)	-2.1 (-6.3--0.23)	-11 (-31--0.95)
Palestine	2016	Female	95 plus	25% (7.5-50%)	75% (50-93%)	0.075 (0.013-0.21)	-0.65% (-2--0.039%)	-0.78% (-2.4--0.058%)	-0.98 (-3--0.076)	-3.2 (-10--0.19)
Palestine	2016	Male	All Ages	7.9% (5.5-11%)	92% (89-95%)	0.043 (0.028-0.064)	0.29% (0.15-0.43%)	0.058% (-0.19-0.26%)	6.6 (-22-29)	1900 (920-2900)
Palestine	2016	Male	15 to 19	6.6% (2.8-13%)	93% (87-97%)	0.022 (0.0068-0.048)	0.47% (0.29-0.67%)	0.23% (0.1-0.45%)	0.76 (0.34-1.5)	210 (120-320)
Palestine	2016	Male	20 to 24	10% (4-20%)	90% (80-96%)	0.048 (0.017-0.098)	0.99% (0.65-1.4%)	0.42% (0.17-0.9%)	1.5 (0.6-3.1)	450 (280-680)
Palestine	2016	Male	25 to 29	13% (5.2-24%)	87% (76-95%)	0.076 (0.028-0.14)	1% (0.74-1.5%)	0.52% (0.19-1.1%)	1.4 (0.51-2.9)	380 (250-560)
Palestine	2016	Male	30 to 34	15% (5.5-28%)	85% (72-95%)	0.099 (0.039-0.18)	0.93% (0.63-1.3%)	0.5% (0.15-1%)	1.2 (0.37-2.3)	280 (180-420)
Palestine	2016	Male	35 to 39	16% (6.3-29%)	84% (71-94%)	0.11 (0.041-0.22)	0.81% (0.54-1.1%)	0.52% (0.091-1%)	1.3 (0.23-2.5)	230 (140-330)
Palestine	2016	Male	40 to 44	16% (7.5-29%)	84% (71-92%)	0.12 (0.045-0.24)	0.6% (0.24-0.88%)	0.33% (-0.25-0.74%)	0.94 (-0.7-2.1)	160 (69-250)
Palestine	2016	Male	45 to 49	16% (6.5-30%)	84% (70-93%)	0.12 (0.04-0.25)	0.39% (0.0046-0.68%)	0.22% (-0.41-0.67%)	0.93 (-1.7-2.9)	120 (1.3-210)
Palestine	2016	Male	50 to 54	15% (6.6-30%)	85% (70-93%)	0.12 (0.037-0.27)	0.18% (-0.32-0.5%)	0.036% (-0.68-0.5%)	0.2 (-3.9-2.9)	56 (-100-160)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Palestine	2016	Male	55 to 59	15% (6.3-28%)	85% (72-94%)	0.12 (0.036-0.26)	0.048% (-0.5-0.39%)	-0.059% (-0.81-0.42%)	-0.39 (-5.2-2.7)	14 (-150-110)
Palestine	2016	Male	60 to 64	14% (6-28%)	86% (72-94%)	0.11 (0.03-0.26)	0.011% (-0.59-0.34%)	-0.057% (-0.78-0.37%)	-0.39 (-5.4-2.6)	2.7 (-140-85)
Palestine	2016	Male	65 to 69	14% (6-25%)	86% (75-94%)	0.087 (0.025-0.21)	-0.013% (-0.52-0.29%)	-0.059% (-0.72-0.32%)	-0.53 (-6.5-2.8)	-3.5 (-140-74)
Palestine	2016	Male	70 to 74	12% (5.5-23%)	88% (77-95%)	0.065 (0.018-0.16)	0.029% (-0.41-0.28%)	0.011% (-0.52-0.32%)	0.11 (-5.6-3.2)	6.5 (-96-64)
Palestine	2016	Male	75 to 79	11% (4.5-21%)	89% (79-95%)	0.05 (0.012-0.14)	0.013% (-0.38-0.23%)	-0.0029% (-0.46-0.25%)	-0.034 (-5.1-2.8)	2.2 (-68-41)
Palestine	2016	Male	80 to 84	9.7% (3.6-19%)	90% (81-96%)	0.042 (0.01-0.12)	0.026% (-0.31-0.22%)	0.015% (-0.37-0.23%)	0.15 (-3.5-2.2)	2.8 (-34-24)
Palestine	2016	Male	85 to 89	9.5% (3.7-19%)	91% (81-96%)	0.039 (0.0089-0.11)	-0.01% (-0.36-0.17%)	-0.029% (-0.41-0.17%)	-0.18 (-2.5-1)	-0.53 (-18-8.3)
Palestine	2016	Male	90 to 94	11% (4.2-22%)	89% (78-96%)	0.041 (0.0094-0.11)	-0.054% (-0.42-0.13%)	-0.085% (-0.5-0.12%)	-0.22 (-1.3-0.31)	-0.78 (-6.2-1.9)
Palestine	2016	Male	95 plus	13% (4.5-28%)	87% (72-96%)	0.047 (0.0086-0.13)	-0.087% (-0.57-0.13%)	-0.14% (-0.68-0.19%)	-0.11 (-0.51-0.078)	-0.24 (-1.5-0.36)
Oman	1990	Female	All Ages	9.1% (5.9-14%)	91% (86-94%)	0.11 (0.073-0.15)	-0.11% (-0.046-0.27%)	-0.11% (-0.54-0.31%)	-4.3 (-20-11)	310 (-130-780)
Oman	1990	Female	15 to 19	8.8% (2.9-19%)	91% (81-97%)	0.1 (0.024-0.25)	0.46% (0.24-0.8%)	0.56% (0.11-1.5%)	0.28 (0.054-0.75)	47 (24-85)
Oman	1990	Female	20 to 24	11% (3.6-24%)	89% (76-96%)	0.16 (0.047-0.36)	0.81% (0.47-1.3%)	0.75% (0.14-1.7%)	0.42 (0.081-1)	81 (45-130)
Oman	1990	Female	25 to 29	13% (3.6-27%)	87% (73-96%)	0.19 (0.055-0.41)	0.87% (0.53-1.3%)	0.9% (0.19-2%)	0.53 (0.11-1.2)	85 (49-140)
Oman	1990	Female	30 to 34	14% (3.7-30%)	86% (70-96%)	0.23 (0.07-0.45)	0.78% (0.43-1.2%)	0.89% (0.2-2%)	0.58 (0.13-1.3)	76 (40-120)
Oman	1990	Female	35 to 39	16% (4.5-31%)	84% (69-95%)	0.27 (0.089-0.49)	0.68% (0.29-1.2%)	0.93% (0.16-2%)	0.69 (0.12-1.5)	62 (27-110)
Oman	1990	Female	40 to 44	17% (6.1-31%)	83% (69-94%)	0.3 (0.11-0.56)	0.54% (0.089-1.2%)	0.83% (0.023-2%)	0.72 (0.019-1.8)	47 (8.3-100)
Oman	1990	Female	45 to 49	17% (6.2-32%)	83% (68-94%)	0.32 (0.1-0.65)	0.32% (-0.33-1.1%)	0.48% (-0.6-1.8%)	0.52 (-0.62-1.9)	29 (-29-100)
Oman	1990	Female	50 to 54	18% (7-36%)	82% (64-93%)	0.34 (0.1-0.7)	0.16% (-0.65-1.1%)	0.27% (-0.91-1.7%)	0.34 (-1.2-2.2)	14 (-56-90)
Oman	1990	Female	55 to 59	19% (7.5-37%)	81% (63-92%)	0.34 (0.095-0.77)	-0.075% (-1.1-1%)	-0.083% (-1.7-1.5%)	-0.11 (-2.3-2.3)	-5.5 (-85-80)
Oman	1990	Female	60 to 64	20% (8-39%)	80% (61-92%)	0.33 (0.082-0.85)	-0.28% (-1.3-0.98%)	-0.36% (-1.8-1.4%)	-0.63 (-3.3-2.6)	-20 (-98-76)
Oman	1990	Female	65 to 69	21% (8.5-38%)	79% (62-92%)	0.29 (0.072-0.69)	-0.47% (-1.6-0.51%)	-0.61% (-2.1-0.66%)	-1.3 (-4.5-1.4)	-33 (-110-35)
Oman	1990	Female	70 to 74	20% (8-36%)	80% (64-92%)	0.23 (0.054-0.62)	-0.53% (-1.6-0.25%)	-0.69% (-2-0.33%)	-1.6 (-4.9-0.86)	-32 (-93-14)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	1990	Female	75 to 79	17% (6.4-36%)	83% (64-94%)	0.19 (0.034-0.56)	-0.56% (-1.6-0.064%)	-0.72% (-2-0.067%)	-1.7 (-5-0.15)	-25 (-72-3)
Oman	1990	Female	80 to 84	16% (5.4-35%)	84% (65-95%)	0.17 (0.03-0.54)	-0.53% (-1.7--0.013%)	-0.67% (-2.1--0.031%)	-1.2 (-3.9--0.059)	-13 (-41--0.33)
Oman	1990	Female	85 to 89	16% (5.4-37%)	84% (63-95%)	0.17 (0.032-0.53)	-0.61% (-1.9--0.054%)	-0.77% (-2.4--0.093%)	-1 (-3.2--0.13)	-7.5 (-24--0.67)
Oman	1990	Female	90 to 94	19% (6.5-42%)	81% (58-93%)	0.19 (0.036-0.59)	-0.69% (-2--0.06%)	-0.88% (-2.5--0.12%)	-0.54 (-1.5--0.071)	-2.6 (-7.5--0.24)
Oman	1990	Female	95 plus	23% (7.8-48%)	77% (52-92%)	0.23 (0.039-0.71)	-0.76% (-2.2--0.001%)	-0.99% (-2.9--0.061%)	-0.17 (-0.49--0.01)	-0.54 (-1.6--0.00077)
Oman	1990	Male	All Ages	8% (5.2-12%)	92% (88-95%)	0.15 (0.11-0.21)	0.5% (0.3-0.77%)	0.48% (0.14-0.92%)	22 (6.4-43)	1800 (1100-2700)
Oman	1990	Male	15 to 19	6.2% (2.5-12%)	94% (88-98%)	0.069 (0.019-0.17)	0.71% (0.4-1.2%)	0.67% (0.18-1.5%)	0.51 (0.14-1.2)	81 (44-130)
Oman	1990	Male	20 to 24	9.6% (3.7-19%)	90% (81-96%)	0.15 (0.047-0.32)	1.5% (0.87-2.4%)	1.3% (0.39-2.9%)	1.5 (0.44-3.3)	210 (120-340)
Oman	1990	Male	25 to 29	12% (4.5-23%)	88% (77-95%)	0.24 (0.084-0.46)	1.7% (0.97-2.7%)	1.8% (0.66-3.6%)	2.6 (0.89-5.1)	280 (160-450)
Oman	1990	Male	30 to 34	14% (4.8-26%)	86% (74-95%)	0.3 (0.12-0.56)	1.6% (0.89-2.6%)	1.8% (0.63-3.6%)	3.2 (1.1-6.4)	300 (170-500)
Oman	1990	Male	35 to 39	15% (5.6-27%)	85% (73-94%)	0.34 (0.14-0.61)	1.5% (0.8-2.5%)	1.8% (0.65-3.6%)	3.8 (1.3-7.6)	300 (160-510)
Oman	1990	Male	40 to 44	15% (6.8-27%)	85% (73-93%)	0.37 (0.14-0.68)	1.3% (0.64-2.2%)	1.5% (0.44-3.1%)	3.2 (0.94-6.8)	230 (110-410)
Oman	1990	Male	45 to 49	15% (6.1-28%)	85% (72-94%)	0.39 (0.13-0.79)	1.1% (0.46-2.3%)	1.3% (0.32-3.2%)	2.8 (0.7-7)	160 (64-350)
Oman	1990	Male	50 to 54	14% (6.2-28%)	86% (72-94%)	0.39 (0.13-0.84)	0.85% (0.15-2%)	0.98% (-0.037-2.7%)	1.9 (-0.064-5.4)	94 (17-230)
Oman	1990	Male	55 to 59	14% (6-26%)	86% (74-94%)	0.37 (0.11-0.82)	0.69% (-0.017-1.9%)	0.78% (-0.22-2.5%)	1.4 (-0.37-4.4)	56 (-1.4-160)
Oman	1990	Male	60 to 64	13% (5.6-26%)	87% (74-94%)	0.34 (0.09-0.82)	0.56% (-0.11-1.9%)	0.64% (-0.29-2.4%)	1 (-0.46-3.9)	35 (-6.9-120)
Oman	1990	Male	65 to 69	13% (5.6-24%)	87% (76-94%)	0.27 (0.082-0.64)	0.29% (-0.28-1.3%)	0.29% (-0.44-1.5%)	0.52 (-0.78-2.7)	16 (-14-69)
Oman	1990	Male	70 to 74	11% (5.2-21%)	89% (79-95%)	0.21 (0.056-0.52)	0.11% (-0.37-0.73%)	0.072% (-0.55-0.85%)	0.16 (-1.2-1.8)	5.5 (-19-38)
Oman	1990	Male	75 to 79	9.9% (4.2-19%)	90% (81-96%)	0.16 (0.036-0.43)	-0.0062% (-0.43-0.45%)	-0.06% (-0.6-0.49%)	-0.13 (-1.3-1.1)	-0.22 (-17-19)
Oman	1990	Male	80 to 84	8.8% (3.5-17%)	91% (83-97%)	0.13 (0.03-0.33)	0.02% (-0.35-0.35%)	-0.02% (-0.49-0.37%)	-0.03 (-0.74-0.57)	0.38 (-6.8-7)
Oman	1990	Male	85 to 89	8.6% (3.5-17%)	91% (83-97%)	0.12 (0.03-0.29)	-0.0095% (-0.34-0.26%)	-0.059% (-0.48-0.26%)	-0.062 (-0.49-0.28)	-0.085 (-3.2-2.4)
Oman	1990	Male	90 to 94	9.7% (3.9-20%)	90% (80-96%)	0.13 (0.03-0.34)	-0.073% (-0.44-0.19%)	-0.15% (-0.63-0.16%)	-0.07 (-0.29-0.076)	-0.21 (-1.3-0.56)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	1990	Male	95 plus	12% (4.4-25%)	88% (75-96%)	0.15 (0.029-0.44)	-0.083% (-0.56-0.22%)	-0.19% (-0.79-0.17%)	-0.025 (-0.1-0.022)	-0.043 (-0.29-0.11)
Oman	1995	Female	All Ages	9.2% (5.8-14%)	91% (86-94%)	0.1 (0.068-0.14)	0.11% (-0.065-0.28%)	-0.23% (-0.73-0.19%)	-7.6 (-24-6.3)	270 (-150-680)
Oman	1995	Female	15 to 19	9% (2.9-19%)	91% (81-97%)	0.097 (0.024-0.23)	0.46% (0.26-0.77%)	0.55% (0.12-1.4%)	0.29 (0.059-0.77)	57 (31-98)
Oman	1995	Female	20 to 24	11% (3.5-24%)	89% (76-96%)	0.15 (0.045-0.34)	0.83% (0.5-1.3%)	0.77% (0.18-1.9%)	0.32 (0.074-0.77)	75 (44-120)
Oman	1995	Female	25 to 29	13% (3.8-27%)	87% (73-96%)	0.18 (0.053-0.38)	0.87% (0.55-1.3%)	0.89% (0.2-2%)	0.46 (0.1-1.1)	90 (53-140)
Oman	1995	Female	30 to 34	15% (3.5-30%)	85% (70-96%)	0.21 (0.067-0.4)	0.77% (0.46-1.2%)	0.87% (0.22-1.9%)	0.57 (0.14-1.3)	88 (50-140)
Oman	1995	Female	35 to 39	16% (4.3-31%)	84% (69-96%)	0.25 (0.085-0.45)	0.65% (0.31-1%)	0.89% (0.16-1.8%)	0.67 (0.13-1.4)	68 (32-110)
Oman	1995	Female	40 to 44	17% (5.8-31%)	83% (69-94%)	0.28 (0.1-0.53)	0.5% (0.07-1%)	0.79% (0.038-1.9%)	0.63 (0.029-1.6)	44 (6.8-92)
Oman	1995	Female	45 to 49	17% (6.2-32%)	83% (68-94%)	0.3 (0.093-0.64)	0.27% (-0.33-1%)	0.42% (-0.53-1.8%)	0.36 (-0.43-1.6)	20 (-24-74)
Oman	1995	Female	50 to 54	18% (7.2-35%)	82% (65-93%)	0.31 (0.093-0.68)	0.094% (-0.71-0.98%)	0.18% (-1-1.6%)	0.22 (-1.3-1.9)	7.7 (-60-82)
Oman	1995	Female	55 to 59	19% (7.7-37%)	81% (63-92%)	0.31 (0.084-0.72)	-0.16% (-1.2-0.75%)	-0.2% (-1.7-1.1%)	-0.28 (-2.4-1.7)	-13 (-93-61)
Oman	1995	Female	60 to 64	20% (7.7-37%)	80% (63-92%)	0.3 (0.072-0.75)	-0.37% (-1.4-0.63%)	-0.46% (-1.8-1%)	-0.81 (-3.3-1.7)	-27 (-100-47)
Oman	1995	Female	65 to 69	20% (8.6-37%)	80% (63-91%)	0.26 (0.062-0.62)	-0.55% (-1.6-0.28%)	-0.71% (-2.1-0.38%)	-1.5 (-4.5-0.81)	-37 (-110-18)
Oman	1995	Female	70 to 74	19% (7.9-37%)	81% (63-92%)	0.21 (0.047-0.53)	-0.59% (-1.6-0.089%)	-0.75% (-2-0.1%)	-1.8 (-5.1-0.25)	-36 (-99-5.5)
Oman	1995	Female	75 to 79	17% (6.3-37%)	83% (63-94%)	0.17 (0.032-0.49)	-0.61% (-1.8-0.048%)	-0.77% (-2.3-0.053%)	-2.3 (-6.5-0.16)	-33 (-95-2.6)
Oman	1995	Female	80 to 84	16% (5.2-36%)	84% (64-95%)	0.16 (0.027-0.47)	-0.59% (-1.8-0.032%)	-0.73% (-2.2-0.056%)	-1.8 (-5.5-0.14)	-19 (-59-1.1)
Oman	1995	Female	85 to 89	17% (5.5-37%)	83% (63-94%)	0.16 (0.03-0.48)	-0.67% (-2--0.085%)	-0.84% (-2.5-0.12%)	-1.5 (-4.6-0.22)	-11 (-34-1.4)
Oman	1995	Female	90 to 94	19% (6-43%)	81% (57-94%)	0.18 (0.031-0.57)	-0.76% (-2.1--0.08%)	-0.96% (-2.6--0.12%)	-0.84 (-2.3--0.099)	-4.2 (-11--0.42)
Oman	1995	Female	95 plus	24% (6.8-49%)	76% (51-93%)	0.22 (0.034-0.68)	-0.84% (-2.4--0.021%)	-1.1% (-3--0.081%)	-0.29 (-0.81--0.021)	-0.92 (-2.6--0.023)
Oman	1995	Male	All Ages	8.7% (5.5-13%)	91% (87-95%)	0.16 (0.12-0.21)	0.66% (0.42-0.96%)	0.58% (0.18-1.1%)	27 (8.2-52)	2400 (1500-3500)
Oman	1995	Male	15 to 19	6.2% (2.6-12%)	94% (88-97%)	0.063 (0.018-0.15)	0.72% (0.41-1.2%)	0.66% (0.17-1.5%)	0.62 (0.16-1.4)	100 (58-170)
Oman	1995	Male	20 to 24	9.5% (3.9-18%)	90% (82-96%)	0.14 (0.046-0.3)	1.5% (0.93-2.4%)	1.3% (0.41-3%)	1.6 (0.5-3.6)	230 (140-370)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	1995	Male	25 to 29	12% (4.5-23%)	88% (77-95%)	0.21 (0.08-0.4)	1.6% (1-2.5%)	1.7% (0.58-3.3%)	3.7 (1.3-7.5)	440 (260-690)
Oman	1995	Male	30 to 34	14% (5-26%)	86% (74-95%)	0.27 (0.12-0.49)	1.5% (0.89-2.4%)	1.7% (0.62-3.3%)	4.6 (1.6-9.3)	460 (270-760)
Oman	1995	Male	35 to 39	15% (5.6-28%)	85% (72-94%)	0.31 (0.13-0.55)	1.4% (0.8-2.3%)	1.7% (0.63-3.2%)	4.5 (1.6-8.8)	380 (210-610)
Oman	1995	Male	40 to 44	15% (6.6-27%)	85% (73-93%)	0.33 (0.13-0.62)	1.2% (0.6-2%)	1.4% (0.44-2.7%)	3.9 (1.3-8.1)	290 (150-500)
Oman	1995	Male	45 to 49	15% (6.1-28%)	85% (72-94%)	0.35 (0.12-0.71)	1% (0.45-2%)	1.2% (0.31-2.8%)	3.1 (0.84-7.5)	190 (79-390)
Oman	1995	Male	50 to 54	14% (6.2-28%)	86% (72-94%)	0.35 (0.11-0.75)	0.73% (0.096-1.8%)	0.82% (-0.12-2.3%)	2.2 (-0.32-6.2)	110 (16-280)
Oman	1995	Male	55 to 59	14% (5.7-27%)	86% (73-94%)	0.34 (0.099-0.76)	0.57% (-0.068-1.6%)	0.62% (-0.29-2%)	1.6 (-0.75-5.3)	71 (-8.2-200)
Oman	1995	Male	60 to 64	13% (5.5-26%)	87% (74-94%)	0.3 (0.083-0.74)	0.46% (-0.15-1.6%)	0.51% (-0.33-2%)	1.2 (-0.77-4.8)	42 (-14-150)
Oman	1995	Male	65 to 69	13% (5.6-24%)	87% (76-94%)	0.25 (0.071-0.57)	0.23% (-0.27-1%)	0.22% (-0.46-1.2%)	0.45 (-0.94-2.6)	15 (-17-65)
Oman	1995	Male	70 to 74	11% (5.2-21%)	89% (79-95%)	0.18 (0.051-0.45)	0.083% (-0.37-0.62%)	0.045% (-0.56-0.7%)	0.096 (-1.2-1.5)	4.2 (-20-31)
Oman	1995	Male	75 to 79	9.9% (4.3-19%)	90% (81-96%)	0.14 (0.033-0.38)	-0.018% (-0.41-0.31%)	-0.077% (-0.6-0.31%)	-0.17 (-1.3-0.68)	-0.69 (-16-12)
Oman	1995	Male	80 to 84	8.8% (3.5-18%)	91% (82-96%)	0.12 (0.029-0.3)	0.012% (-0.35-0.27%)	-0.032% (-0.5-0.28%)	-0.054 (-0.84-0.47)	0.28 (-7.8-5.9)
Oman	1995	Male	85 to 89	8.6% (3.5-17%)	91% (83-97%)	0.11 (0.027-0.27)	-0.015% (-0.36-0.22%)	-0.066% (-0.5-0.22%)	-0.082 (-0.61-0.27)	-0.16 (-4-2.5)
Oman	1995	Male	90 to 94	9.7% (3.9-19%)	90% (81-96%)	0.11 (0.027-0.32)	-0.076% (-0.47-0.18%)	-0.15% (-0.64-0.15%)	-0.091 (-0.38-0.087)	-0.28 (-1.7-0.63)
Oman	1995	Male	95 plus	12% (4.3-26%)	88% (74-96%)	0.13 (0.026-0.4)	-0.081% (-0.57-0.23%)	-0.19% (-0.8-0.17%)	-0.032 (-0.13-0.028)	-0.054 (-0.38-0.15)
Oman	2000	Female	All Ages	9.3% (5.2-14%)	91% (86-95%)	0.13 (0.088-0.17)	0.19% (-0.042-0.44%)	-0.24% (-0.88-0.4%)	-7.3 (-26-12)	400 (-92-950)
Oman	2000	Female	15 to 19	8.8% (2.8-20%)	91% (80-97%)	0.11 (0.029-0.26)	0.48% (0.27-0.79%)	0.63% (0.15-1.6%)	0.36 (0.086-0.89)	69 (38-120)
Oman	2000	Female	20 to 24	11% (3.5-25%)	89% (75-97%)	0.18 (0.053-0.39)	0.88% (0.53-1.3%)	0.86% (0.22-2.1%)	0.53 (0.14-1.3)	130 (76-200)
Oman	2000	Female	25 to 29	13% (3.5-28%)	87% (72-96%)	0.22 (0.066-0.43)	0.93% (0.6-1.3%)	1% (0.27-2.3%)	0.55 (0.15-1.2)	110 (66-170)
Oman	2000	Female	30 to 34	14% (3.5-30%)	86% (70-97%)	0.25 (0.079-0.49)	0.83% (0.51-1.2%)	1% (0.31-2.1%)	0.63 (0.18-1.3)	98 (58-150)
Oman	2000	Female	35 to 39	16% (4.1-31%)	84% (69-96%)	0.29 (0.096-0.54)	0.71% (0.36-1.1%)	1.1% (0.3-2.2%)	0.77 (0.21-1.5)	77 (40-130)
Oman	2000	Female	40 to 44	16% (6-31%)	84% (69-94%)	0.33 (0.11-0.63)	0.58% (0.14-1.2%)	1% (0.19-2.3%)	0.76 (0.13-1.7)	51 (13-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	2000	Female	45 to 49	17% (6.2-33%)	83% (67-94%)	0.36 (0.11-0.75)	0.37% (-0.29-1.2%)	0.65% (-0.36-2.3%)	0.52 (-0.29-1.8)	27 (-21-93)
Oman	2000	Female	50 to 54	18% (6.8-35%)	82% (65-93%)	0.37 (0.11-0.84)	0.21% (-0.7-1.4%)	0.42% (-0.9-2.2%)	0.42 (-0.91-2.4)	15 (-50-100)
Oman	2000	Female	55 to 59	19% (7.3-38%)	81% (62-93%)	0.38 (0.1-0.87)	-0.067% (-1.2-1.2%)	-0.026% (-1.6-1.8%)	-0.041 (-2.7-2.9)	-5.8 (-110-100)
Oman	2000	Female	60 to 64	20% (7.5-40%)	80% (60-92%)	0.36 (0.086-0.87)	-0.3% (-1.5-1%)	-0.35% (-1.9-1.5%)	-0.69 (-3.8-2.9)	-25 (-120-83)
Oman	2000	Female	65 to 69	21% (8.4-38%)	79% (62-92%)	0.32 (0.077-0.72)	-0.54% (-1.7-0.49%)	-0.67% (-2.2-0.69%)	-1.5 (-5-1.6)	-40 (-130-36)
Oman	2000	Female	70 to 74	20% (8.1-37%)	80% (63-92%)	0.26 (0.059-0.66)	-0.61% (-1.7-0.22%)	-0.78% (-2.2-0.31%)	-1.9 (-5.5-0.78)	-39 (-110-14)
Oman	2000	Female	75 to 79	18% (6.6-36%)	82% (64-93%)	0.21 (0.042-0.63)	-0.63% (-1.8-0.081%)	-0.81% (-2.3-0.081%)	-2.2 (-6.3-0.22)	-32 (-94-4.2)
Oman	2000	Female	80 to 84	16% (5.4-36%)	84% (64-95%)	0.19 (0.035-0.55)	-0.6% (-1.8--0.008%)	-0.75% (-2.2--0.023%)	-2.1 (-6--0.065)	-22 (-65--0.28)
Oman	2000	Female	85 to 89	17% (5.2-37%)	83% (63-95%)	0.19 (0.035-0.58)	-0.67% (-2--0.06%)	-0.85% (-2.5--0.099%)	-2 (-5.8--0.23)	-14 (-43--1.3)
Oman	2000	Female	90 to 94	19% (6.1-41%)	81% (59-94%)	0.21 (0.038-0.65)	-0.76% (-2.1--0.051%)	-0.97% (-2.5--0.092%)	-1.1 (-3--0.11)	-5.6 (-15--0.39)
Oman	2000	Female	95 plus	24% (7.3-48%)	76% (52-93%)	0.26 (0.041-0.78)	-0.82% (-2.3-0.05%)	-1.1% (-3--0.02%)	-0.41 (-1.2--0.0076)	-1.3 (-3.7-0.079)
Oman	2000	Male	All Ages	8.2% (5.4-12%)	92% (88-95%)	0.17 (0.13-0.22)	0.78% (0.51-1.1%)	0.67% (0.23-1.3%)	28 (9.4-54)	2300 (1500-3400)
Oman	2000	Male	15 to 19	6.2% (2.5-12%)	94% (88-97%)	0.074 (0.022-0.16)	0.75% (0.42-1.2%)	0.7% (0.18-1.6%)	0.78 (0.21-1.8)	130 (69-200)
Oman	2000	Male	20 to 24	9.4% (3.6-18%)	91% (82-96%)	0.16 (0.055-0.32)	1.6% (0.95-2.4%)	1.4% (0.44-2.9%)	2.2 (0.7-4.6)	330 (190-510)
Oman	2000	Male	25 to 29	12% (4.5-22%)	88% (78-95%)	0.25 (0.098-0.46)	1.8% (1.1-2.7%)	1.9% (0.7-3.6%)	3.3 (1.2-6.3)	390 (230-610)
Oman	2000	Male	30 to 34	14% (4.6-26%)	86% (74-95%)	0.32 (0.13-0.57)	1.6% (0.98-2.6%)	1.9% (0.75-3.6%)	3.8 (1.4-7.2)	390 (220-620)
Oman	2000	Male	35 to 39	15% (5.6-27%)	85% (73-94%)	0.37 (0.15-0.68)	1.6% (0.88-2.5%)	2% (0.75-3.8%)	3.8 (1.4-7.3)	320 (180-520)
Oman	2000	Male	40 to 44	15% (6.6-27%)	85% (73-93%)	0.4 (0.15-0.75)	1.4% (0.73-2.4%)	1.7% (0.61-3.5%)	3.3 (1.2-6.8)	240 (130-430)
Oman	2000	Male	45 to 49	15% (5.8-27%)	85% (73-94%)	0.42 (0.14-0.91)	1.2% (0.54-2.6%)	1.5% (0.47-3.6%)	3.4 (1-8)	200 (85-420)
Oman	2000	Male	50 to 54	14% (5.9-28%)	86% (72-94%)	0.42 (0.13-0.9)	0.95% (0.26-2.3%)	1.1% (0.1-3.2%)	2.1 (0.19-5.8)	100 (28-250)
Oman	2000	Male	55 to 59	14% (5.7-26%)	86% (74-94%)	0.4 (0.12-0.88)	0.76% (0.044-2.1%)	0.89% (-0.13-2.8%)	2.3 (-0.33-7.1)	95 (5.2-260)
Oman	2000	Male	60 to 64	13% (5.5-25%)	87% (75-95%)	0.36 (0.1-0.85)	0.62% (-0.087-2.1%)	0.72% (-0.21-2.7%)	2.3 (-0.68-8.2)	76 (-12-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	2000	Male	65 to 69	13% (5.6-23%)	87% (77-94%)	0.29 (0.085-0.66)	0.31% (-0.25-1.4%)	0.33% (-0.41-1.6%)	0.97 (-1.2-4.9)	28 (-23-130)
Oman	2000	Male	70 to 74	11% (5.2-20%)	89% (80-95%)	0.22 (0.059-0.55)	0.13% (-0.36-0.78%)	0.099% (-0.54-0.89%)	0.28 (-1.5-2.4)	8.2 (-24-50)
Oman	2000	Male	75 to 79	9.6% (4-19%)	90% (81-96%)	0.16 (0.037-0.44)	-0.00013% (-0.41-0.42%)	-0.053% (-0.57-0.45%)	-0.13 (-1.5-1.1)	0.009 (-18-19)
Oman	2000	Male	80 to 84	8.6% (3.4-17%)	91% (83-97%)	0.14 (0.034-0.34)	0.018% (-0.34-0.35%)	-0.021% (-0.46-0.36%)	-0.042 (-0.95-0.77)	0.48 (-8.8-9.3)
Oman	2000	Male	85 to 89	8.4% (3.4-18%)	92% (82-97%)	0.13 (0.033-0.32)	-0.013% (-0.37-0.28%)	-0.061% (-0.49-0.27%)	-0.098 (-0.78-0.45)	-0.19 (-5.2-4)
Oman	2000	Male	90 to 94	9.5% (3.9-19%)	90% (81-96%)	0.14 (0.032-0.37)	-0.082% (-0.47-0.21%)	-0.16% (-0.65-0.18%)	-0.12 (-0.5-0.14)	-0.39 (-2.2-1)
Oman	2000	Male	95 plus	12% (4.3-26%)	88% (74-96%)	0.16 (0.031-0.46)	-0.094% (-0.6-0.27%)	-0.21% (-0.87-0.22%)	-0.046 (-0.19-0.05)	-0.084 (-0.53-0.25)
Oman	2005	Female	All Ages	9.5% (5.6-15%)	90% (85-94%)	0.16 (0.12-0.22)	0.33% (0.06-0.68%)	-0.095% (-0.82-0.82%)	-2.5 (-22-22)	650 (120-1400)
Oman	2005	Female	15 to 19	9% (2.9-20%)	91% (80-97%)	0.14 (0.039-0.33)	0.52% (0.28-0.86%)	0.76% (0.22-1.8%)	0.37 (0.11-0.88)	77 (41-130)
Oman	2005	Female	20 to 24	12% (3.3-27%)	88% (73-97%)	0.22 (0.065-0.51)	0.94% (0.57-1.5%)	1% (0.3-2.4%)	0.65 (0.18-1.5)	150 (88-250)
Oman	2005	Female	25 to 29	13% (3.4-29%)	87% (71-97%)	0.27 (0.078-0.56)	0.99% (0.64-1.4%)	1.2% (0.37-2.6%)	0.79 (0.24-1.7)	160 (98-230)
Oman	2005	Female	30 to 34	14% (3.7-30%)	86% (70-96%)	0.31 (0.095-0.58)	0.91% (0.59-1.3%)	1.3% (0.44-2.6%)	0.71 (0.24-1.4)	110 (65-160)
Oman	2005	Female	35 to 39	16% (4.7-30%)	84% (70-95%)	0.36 (0.12-0.66)	0.8% (0.43-1.3%)	1.4% (0.44-2.8%)	1 (0.3-2)	96 (51-160)
Oman	2005	Female	40 to 44	16% (6-31%)	84% (69-94%)	0.4 (0.15-0.77)	0.69% (0.23-1.4%)	1.4% (0.35-2.8%)	1.2 (0.29-2.5)	78 (25-160)
Oman	2005	Female	45 to 49	17% (6-32%)	83% (68-94%)	0.44 (0.14-0.9)	0.52% (-0.25-1.5%)	0.98% (-0.25-3.1%)	0.97 (-0.23-3)	52 (-23-160)
Oman	2005	Female	50 to 54	18% (7.1-34%)	82% (66-93%)	0.45 (0.14-0.96)	0.41% (-0.59-1.9%)	0.79% (-0.69-3.1%)	0.98 (-0.93-4)	39 (-52-180)
Oman	2005	Female	55 to 59	18% (7.5-36%)	82% (64-93%)	0.45 (0.13-1)	0.16% (-0.94-1.6%)	0.34% (-1.3-2.6%)	0.45 (-1.7-3.5)	13 (-74-130)
Oman	2005	Female	60 to 64	19% (7.6-38%)	81% (62-92%)	0.44 (0.11-1.1)	-0.083% (-1.4-1.6%)	-0.048% (-1.8-2.3%)	-0.089 (-3.3-4.3)	-6.7 (-120-130)
Oman	2005	Female	65 to 69	20% (8.1-37%)	80% (63-92%)	0.38 (0.096-0.85)	-0.38% (-1.6-0.97%)	-0.47% (-2.1-1.4%)	-0.98 (-4.3-2.9)	-27 (-110-69)
Oman	2005	Female	70 to 74	19% (7.8-35%)	81% (65-92%)	0.3 (0.072-0.78)	-0.51% (-1.6-0.53%)	-0.66% (-2.1-0.7%)	-1.4 (-4.6-1.6)	-30 (-94-31)
Oman	2005	Female	75 to 79	17% (6.5-34%)	83% (66-94%)	0.25 (0.05-0.68)	-0.53% (-1.6-0.29%)	-0.7% (-2.1-0.32%)	-1.8 (-5.6-0.88)	-28 (-86-15)
Oman	2005	Female	80 to 84	15% (5.3-35%)	85% (65-95%)	0.22 (0.044-0.65)	-0.49% (-1.5-0.14%)	-0.64% (-2-0.15%)	-1.7 (-5.2-0.41)	-19 (-58-5.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	2005	Female	85 to 89	16% (5.3-36%)	84% (64-95%)	0.23 (0.042-0.68)	-0.58% (-1.8-0.061%)	-0.76% (-2.3-0.043%)	-1.8 (-5.5-0.1)	-13 (-42-1.5)
Oman	2005	Female	90 to 94	19% (6.1-40%)	81% (60-94%)	0.26 (0.048-0.76)	-0.67% (-1.9-0.045%)	-0.88% (-2.4-0.0031%)	-1.3 (-3.5-0.0046)	-6.3 (-18-0.44)
Oman	2005	Female	95 plus	23% (6.8-48%)	77% (52-93%)	0.31 (0.051-0.91)	-0.71% (-2.2-0.31%)	-0.97% (-2.8-0.29%)	-0.48 (-1.4-0.14)	-1.5 (-4.6-0.65)
Oman	2005	Male	All Ages	8.6% (5.6-12%)	91% (88-94%)	0.23 (0.17-0.28)	1% (0.68-1.5%)	0.96% (0.38-1.8%)	42 (16-81)	3100 (1900-4500)
Oman	2005	Male	15 to 19	6.2% (2.6-12%)	94% (88-97%)	0.092 (0.028-0.21)	0.81% (0.47-1.3%)	0.82% (0.25-1.8%)	0.95 (0.29-2.1)	150 (84-240)
Oman	2005	Male	20 to 24	9.6% (3.6-19%)	90% (81-96%)	0.2 (0.071-0.42)	1.7% (1-2.6%)	1.6% (0.56-3.4%)	2.8 (0.97-5.9)	370 (210-590)
Oman	2005	Male	25 to 29	12% (4.7-23%)	88% (77-95%)	0.32 (0.12-0.59)	2% (1.2-3.1%)	2.2% (0.89-4.3%)	4.3 (1.7-8.3)	480 (270-760)
Oman	2005	Male	30 to 34	14% (5.1-27%)	86% (73-95%)	0.41 (0.18-0.72)	1.9% (1.1-2.9%)	2.4% (1-4.5%)	4.2 (1.7-8.1)	420 (250-670)
Oman	2005	Male	35 to 39	15% (5.7-29%)	85% (71-94%)	0.47 (0.19-0.86)	1.8% (1-3%)	2.5% (0.99-4.7%)	4.6 (1.8-9.1)	390 (230-640)
Oman	2005	Male	40 to 44	15% (6.8-27%)	85% (73-93%)	0.5 (0.19-0.94)	1.6% (0.87-2.8%)	2.2% (0.82-4.4%)	4.7 (1.7-9.5)	340 (180-590)
Oman	2005	Male	45 to 49	15% (6-28%)	85% (72-94%)	0.52 (0.18-1)	1.5% (0.65-2.9%)	2% (0.61-4.4%)	5.5 (1.7-13)	310 (130-630)
Oman	2005	Male	50 to 54	14% (6.3-29%)	86% (71-94%)	0.51 (0.16-1.1)	1.2% (0.35-2.8%)	1.5% (0.27-3.9%)	4.5 (0.78-12)	220 (65-510)
Oman	2005	Male	55 to 59	14% (5.9-27%)	86% (73-94%)	0.49 (0.15-1.1)	1% (0.13-2.7%)	1.3% (0.056-3.5%)	3.6 (0.15-10)	140 (19-370)
Oman	2005	Male	60 to 64	13% (5.3-26%)	87% (74-95%)	0.44 (0.12-1.1)	0.86% (0.0063-2.6%)	1% (-0.08-3.2%)	3.8 (-0.29-12)	120 (0.84-360)
Oman	2005	Male	65 to 69	12% (5.4-24%)	88% (76-95%)	0.36 (0.11-0.79)	0.49% (-0.16-1.8%)	0.55% (-0.3-2.1%)	2.3 (-1.2-8.9)	62 (-20-220)
Oman	2005	Male	70 to 74	11% (5.1-20%)	89% (80-95%)	0.26 (0.076-0.64)	0.23% (-0.31-1.2%)	0.23% (-0.46-1.3%)	0.87 (-1.7-5.3)	21 (-28-110)
Oman	2005	Male	75 to 79	9.6% (4.1-18%)	90% (82-96%)	0.2 (0.046-0.53)	0.067% (-0.39-0.7%)	0.023% (-0.55-0.77%)	0.075 (-1.8-2.5)	3.8 (-22-39)
Oman	2005	Male	80 to 84	8.6% (3.5-17%)	91% (83-96%)	0.17 (0.043-0.42)	0.075% (-0.32-0.58%)	0.045% (-0.43-0.61%)	0.11 (-1-1.5)	2.3 (-9.8-17)
Oman	2005	Male	85 to 89	8.5% (3.4-17%)	92% (83-97%)	0.16 (0.038-0.39)	0.035% (-0.33-0.43%)	-0.0082% (-0.46-0.45%)	-0.014 (-0.81-0.81)	0.56 (-5.3-6.9)
Oman	2005	Male	90 to 94	9.5% (3.8-20%)	90% (80-96%)	0.17 (0.039-0.44)	-0.05% (-0.48-0.34%)	-0.12% (-0.65-0.32%)	-0.12 (-0.62-0.3)	-0.29 (-2.7-2)
Oman	2005	Male	95 plus	12% (4.3-25%)	88% (75-96%)	0.19 (0.037-0.57)	-0.063% (-0.56-0.46%)	-0.18% (-0.83-0.4%)	-0.049 (-0.23-0.11)	-0.07 (-0.61-0.52)
Oman	2010	Female	All Ages	10% (6.1-17%)	90% (83-94%)	0.2 (0.14-0.27)	0.37% (0.048-0.77%)	-0.052% (-0.86-1%)	-1.6 (-27-31)	800 (100-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	2010	Female	15 to 19	8.9% (2.8-20%)	91% (80-97%)	0.15 (0.042-0.36)	0.52% (0.3-0.84%)	0.78% (0.24-1.8%)	0.34 (0.11-0.79)	75 (41-130)
Oman	2010	Female	20 to 24	11% (3.5-24%)	89% (76-97%)	0.23 (0.07-0.53)	0.97% (0.6-1.5%)	1.1% (0.36-2.4%)	0.62 (0.21-1.4)	160 (94-250)
Oman	2010	Female	25 to 29	13% (3.7-27%)	87% (73-96%)	0.28 (0.089-0.54)	1% (0.66-1.4%)	1.2% (0.43-2.4%)	0.81 (0.28-1.7)	180 (110-280)
Oman	2010	Female	30 to 34	14% (3.9-30%)	86% (70-96%)	0.32 (0.11-0.59)	0.91% (0.58-1.3%)	1.4% (0.54-2.6%)	0.92 (0.35-1.8)	160 (100-250)
Oman	2010	Female	35 to 39	16% (4.5-31%)	84% (69-95%)	0.39 (0.14-0.72)	0.82% (0.44-1.3%)	1.5% (0.53-3%)	1 (0.34-2)	110 (57-170)
Oman	2010	Female	40 to 44	17% (6.2-31%)	83% (69-94%)	0.44 (0.16-0.83)	0.72% (0.25-1.4%)	1.6% (0.47-3.3%)	1.2 (0.35-2.5)	86 (29-170)
Oman	2010	Female	45 to 49	17% (5.9-32%)	83% (68-94%)	0.48 (0.15-0.97)	0.57% (-0.16-1.6%)	1.2% (-0.032-3.4%)	1.1 (-0.028-3.2)	65 (-17-190)
Oman	2010	Female	50 to 54	18% (6.8-34%)	82% (66-93%)	0.49 (0.14-1.1)	0.51% (-0.48-2%)	1% (-0.49-3.6%)	1.4 (-0.66-4.9)	56 (-55-230)
Oman	2010	Female	55 to 59	19% (6.9-37%)	81% (63-93%)	0.49 (0.13-1.1)	0.27% (-0.89-1.9%)	0.53% (-1.3-3%)	0.94 (-2.2-5.6)	30 (-100-200)
Oman	2010	Female	60 to 64	20% (7.5-38%)	80% (62-93%)	0.47 (0.11-1.1)	0.019% (-1.5-1.8%)	0.096% (-1.8-2.7%)	0.21 (-3.9-5.9)	2 (-140-180)
Oman	2010	Female	65 to 69	20% (8.3-36%)	80% (64-92%)	0.41 (0.1-0.95)	-0.28% (-1.6-1.3%)	-0.32% (-2.1-7%)	-1 (-6.3-5.5)	-28 (-160-130)
Oman	2010	Female	70 to 74	19% (7.3-35%)	81% (65-93%)	0.33 (0.076-0.84)	-0.44% (-1.6-0.92%)	-0.54% (-2.1-3%)	-1.7 (-6.3-6)	-34 (-120-73)
Oman	2010	Female	75 to 79	17% (6.1-35%)	83% (65-94%)	0.27 (0.051-0.81)	-0.47% (-1.6-0.71%)	-0.6% (-2.1-0.88%)	-1.7 (-5.8-2.5)	-25 (-89-38)
Oman	2010	Female	80 to 84	16% (5.2-35%)	84% (65-95%)	0.25 (0.044-0.69)	-0.42% (-1.5-0.32%)	-0.56% (-1.9-0.37%)	-1.6 (-5.3-1)	-17 (-61-14)
Oman	2010	Female	85 to 89	16% (5.3-35%)	84% (65-95%)	0.25 (0.046-0.77)	-0.5% (-1.6-0.23%)	-0.67% (-2.1-0.22%)	-2 (-6-0.64)	-15 (-46-6.5)
Oman	2010	Female	90 to 94	19% (6.2-40%)	81% (60-94%)	0.28 (0.051-0.88)	-0.59% (-1.8-0.32%)	-0.8% (-2.3-0.28%)	-1.6 (-4.7-0.57)	-7.8 (-23-4.3)
Oman	2010	Female	95 plus	23% (6.8-48%)	77% (52-93%)	0.33 (0.055-1)	-0.61% (-2.1-0.69%)	-0.86% (-2.7-0.74%)	-0.71 (-2.2-0.6)	-2.2 (-7.1-2.4)
Oman	2010	Male	All Ages	9.5% (5.9-14%)	90% (86-94%)	0.28 (0.21-0.35)	1.2% (0.84-1.7%)	1.1% (0.51-2%)	64 (28-120)	4600 (3100-6700)
Oman	2010	Male	15 to 19	6.2% (2.6-13%)	94% (87-97%)	0.098 (0.029-0.23)	0.84% (0.49-1.4%)	0.89% (0.29-2%)	1.1 (0.37-2.6)	160 (89-270)
Oman	2010	Male	20 to 24	9.5% (3.8-19%)	91% (81-96%)	0.22 (0.072-0.46)	1.7% (1.1-2.9%)	1.7% (0.59-3.5%)	4.4 (1.6-9.3)	600 (350-980)
Oman	2010	Male	25 to 29	12% (4.5-23%)	88% (77-95%)	0.33 (0.13-0.62)	2% (1.3-3%)	2.3% (0.91-4.3%)	7.2 (2.7-14)	850 (510-1300)
Oman	2010	Male	30 to 34	14% (4.7-27%)	86% (73-95%)	0.43 (0.19-0.75)	1.9% (1.2-2.9%)	2.5% (1.1-4.5%)	6.4 (2.6-12)	650 (390-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	2010	Male	35 to 39	15% (5.5-28%)	85% (72-95%)	0.49 (0.19-0.92)	1.9% (1.1-3.1%)	2.7% (1.1-5.1%)	6.2 (2.7-12)	540 (310-880)
Oman	2010	Male	40 to 44	15% (6.6-27%)	85% (73-93%)	0.52 (0.2-1)	1.7% (0.94-3%)	2.5% (1-4.9%)	6.5 (2.6-13)	480 (250-820)
Oman	2010	Male	45 to 49	15% (5.8-28%)	85% (72-94%)	0.55 (0.18-1.1)	1.7% (0.75-3.2%)	2.3% (0.78-5%)	6.8 (2.4-14)	390 (170-760)
Oman	2010	Male	50 to 54	14% (5.9-28%)	86% (72-94%)	0.54 (0.16-1.2)	1.4% (0.45-3.1%)	1.8% (0.42-4.2%)	6.9 (1.7-16)	330 (110-740)
Oman	2010	Male	55 to 59	13% (5.6-26%)	87% (74-94%)	0.52 (0.16-1.2)	1.2% (0.33-2.8%)	1.6% (0.31-3.8%)	6.2 (1.2-16)	250 (65-580)
Oman	2010	Male	60 to 64	13% (5.4-25%)	87% (75-95%)	0.46 (0.13-1.1)	1% (0.092-2.7%)	1.2% (0.05-3.4%)	4.9 (0.2-14)	150 (14-420)
Oman	2010	Male	65 to 69	12% (5.2-23%)	88% (77-95%)	0.38 (0.11-0.84)	0.66% (-0.056-2.1%)	0.76% (-0.15-2.5%)	3.7 (-0.72-12)	98 (-9.1-300)
Oman	2010	Male	70 to 74	11% (5-21%)	89% (79-95%)	0.28 (0.081-0.71)	0.34% (-0.21-1.5%)	0.37% (-0.32-1.8%)	2 (-1.9-10)	45 (-28-210)
Oman	2010	Male	75 to 79	9.6% (4.1-19%)	90% (81-96%)	0.22 (0.049-0.59)	0.16% (-0.35-0.99%)	0.14% (-0.5-1.1%)	0.63 (-2.3-5.1)	13 (-28-80)
Oman	2010	Male	80 to 84	8.6% (3.3-18%)	91% (82-97%)	0.18 (0.043-0.48)	0.16% (-0.23-0.77%)	0.14% (-0.33-0.86%)	0.47 (-1.1-2.7)	6.5 (-9.7-33)
Oman	2010	Male	85 to 89	8.6% (3.4-17%)	91% (83-97%)	0.17 (0.041-0.43)	0.12% (-0.25-0.6%)	0.089% (-0.36-0.65%)	0.19 (-0.77-1.4)	2.2 (-4.9-12)
Oman	2010	Male	90 to 94	9.7% (3.9-20%)	90% (80-96%)	0.18 (0.041-0.46)	0.024% (-0.38-0.52%)	-0.035% (-0.55-0.5%)	-0.038 (-0.59-0.57)	0.16 (-2.6-3.5)
Oman	2010	Male	95 plus	12% (4.2-27%)	88% (73-96%)	0.21 (0.039-0.6)	0.012% (-0.48-0.65%)	-0.089% (-0.69-0.62%)	-0.031 (-0.24-0.21)	0.017 (-0.65-0.89)
Oman	2016	Female	All Ages	11% (6.2-17%)	89% (83-94%)	0.22 (0.12-0.36)	0.42% (0.096-0.94%)	0.023% (-0.74-1.3%)	0.89 (-28-48)	1100 (240-2500)
Oman	2016	Female	15 to 19	8.4% (2.7-20%)	92% (80-97%)	0.14 (0.032-0.39)	0.52% (0.29-0.88%)	0.81% (0.25-2%)	0.27 (0.084-0.67)	60 (33-110)
Oman	2016	Female	20 to 24	11% (3.4-24%)	89% (76-97%)	0.23 (0.056-0.57)	0.97% (0.6-1.5%)	1.1% (0.33-2.7%)	0.71 (0.21-1.8)	200 (120-320)
Oman	2016	Female	25 to 29	13% (3.3-27%)	87% (73-97%)	0.27 (0.078-0.6)	1% (0.64-1.5%)	1.2% (0.38-2.6%)	1.1 (0.33-2.3)	260 (160-400)
Oman	2016	Female	30 to 34	14% (3.9-31%)	86% (69-96%)	0.32 (0.1-0.63)	0.92% (0.57-1.4%)	1.4% (0.5-2.7%)	1.3 (0.49-2.7)	240 (150-380)
Oman	2016	Female	35 to 39	16% (4.6-31%)	84% (69-95%)	0.38 (0.13-0.76)	0.82% (0.43-1.4%)	1.5% (0.49-3.1%)	1.8 (0.59-3.7)	200 (100-340)
Oman	2016	Female	40 to 44	16% (5.8-30%)	84% (70-94%)	0.43 (0.15-0.88)	0.73% (0.24-1.5%)	1.6% (0.44-3.6%)	1.6 (0.45-3.7)	120 (38-260)
Oman	2016	Female	45 to 49	17% (6.3-32%)	83% (68-94%)	0.47 (0.13-1.1)	0.59% (-0.11-1.8%)	1.2% (-0.092-3.8%)	1.4 (-0.1-4.4)	82 (-17-250)
Oman	2016	Female	50 to 54	18% (6.9-35%)	82% (65-93%)	0.49 (0.12-1.2)	0.5% (-0.46-2.1%)	1.1% (-0.42-3.9%)	1.6 (-0.63-5.9)	68 (-64-280)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	2016	Female	55 to 59	19% (7.3-38%)	81% (62-93%)	0.49 (0.12-1.2)	0.3% (-0.86-2%)	0.63% (-1.2-3.4%)	1.2 (-2.3-6.5)	38 (-110-260)
Oman	2016	Female	60 to 64	20% (7.2-39%)	80% (61-93%)	0.48 (0.11-1.2)	0.077% (-1.3-2%)	0.23% (-1.6-3.1%)	0.62 (-4.3-8.3)	10 (-160-260)
Oman	2016	Female	65 to 69	20% (8.4-36%)	80% (64-92%)	0.42 (0.09-1)	-0.19% (-1.4-1.5%)	-0.18% (-1.8-2.2%)	-0.57 (-5.8-7.2)	-22 (-160-180)
Oman	2016	Female	70 to 74	19% (8.1-34%)	81% (66-92%)	0.34 (0.072-0.94)	-0.33% (-1.4-1.1%)	-0.42% (-1.7-1.7%)	-1.6 (-6.7-5.9)	-36 (-150-120)
Oman	2016	Female	75 to 79	17% (6.6-35%)	83% (65-93%)	0.28 (0.051-0.87)	-0.4% (-1.5-0.92%)	-0.52% (-1.9-1.2%)	-1.7 (-6.5-3.8)	-27 (-100-63)
Oman	2016	Female	80 to 84	16% (5.2-36%)	84% (64-95%)	0.25 (0.042-0.8)	-0.39% (-1.4-0.66%)	-0.51% (-1.7-0.83%)	-1.7 (-5.9-2.9)	-19 (-66-34)
Oman	2016	Female	85 to 89	16% (5.4-37%)	84% (63-95%)	0.25 (0.044-0.8)	-0.48% (-1.6-0.42%)	-0.63% (-2-0.43%)	-2.1 (-6.6-1.4)	-15 (-50-14)
Oman	2016	Female	90 to 94	19% (6.2-41%)	81% (59-94%)	0.29 (0.044-0.87)	-0.58% (-1.8-0.46%)	-0.77% (-2.4-0.5%)	-2.1 (-6.5-1.3)	-10 (-32-8.1)
Oman	2016	Female	95 plus	24% (7.1-48%)	76% (52-93%)	0.34 (0.048-1.1)	-0.59% (-2.1-0.87%)	-0.83% (-2.7-0.93%)	-1 (-3.3-1.1)	-3.1 (-11-4.6)
Oman	2016	Male	All Ages	11% (6.4-16%)	89% (84-94%)	0.32 (0.19-0.5)	1.4% (0.87-2%)	1.3% (0.54-2.4%)	110 (46-220)	8500 (5300-13000)
Oman	2016	Male	15 to 19	6% (2.5-12%)	94% (88-97%)	0.095 (0.026-0.24)	0.89% (0.53-1.5%)	1% (0.36-2.2%)	1 (0.39-2.3)	150 (88-260)
Oman	2016	Male	20 to 24	9.3% (3.6-19%)	91% (81-96%)	0.21 (0.062-0.49)	1.8% (1.1-2.9%)	1.7% (0.63-3.8%)	5.2 (1.9-12)	760 (450-1200)
Oman	2016	Male	25 to 29	12% (4.4-22%)	88% (78-96%)	0.32 (0.12-0.64)	2% (1.2-3%)	2.3% (0.87-4.5%)	12 (4.5-24)	1600 (940-2400)
Oman	2016	Male	30 to 34	13% (4.8-26%)	87% (74-95%)	0.4 (0.17-0.72)	1.9% (1.2-2.8%)	2.4% (0.98-4.4%)	14 (5.7-27)	1600 (1000-2500)
Oman	2016	Male	35 to 39	14% (5.4-27%)	86% (73-95%)	0.46 (0.17-0.91)	1.8% (1.1-3%)	2.6% (1.1-5%)	13 (5.3-25)	1100 (660-1900)
Oman	2016	Male	40 to 44	15% (6.4-27%)	85% (73-94%)	0.5 (0.17-1)	1.7% (0.89-3.1%)	2.4% (0.91-5.1%)	12 (4.4-25)	880 (460-1600)
Oman	2016	Male	45 to 49	14% (5.8-27%)	86% (73-94%)	0.53 (0.15-1.2)	1.6% (0.71-3.1%)	2.2% (0.75-4.9%)	14 (4.5-31)	810 (370-1600)
Oman	2016	Male	50 to 54	14% (5.9-27%)	86% (73-94%)	0.53 (0.14-1.3)	1.4% (0.46-3%)	1.8% (0.45-4.4%)	11 (2.7-26)	530 (170-1200)
Oman	2016	Male	55 to 59	13% (5.5-26%)	87% (74-95%)	0.5 (0.14-1.2)	1.2% (0.25-2.9%)	1.5% (0.19-3.9%)	10 (1.3-27)	410 (88-1000)
Oman	2016	Male	60 to 64	13% (5.2-25%)	87% (75-95%)	0.45 (0.12-1.1)	0.98% (0.099-2.7%)	1.2% (0.045-3.6%)	9.6 (0.36-28)	310 (34-860)
Oman	2016	Male	65 to 69	12% (5.3-22%)	88% (78-95%)	0.37 (0.1-0.85)	0.65% (-0.046-2.1%)	0.77% (-0.12-2.6%)	5.1 (-0.83-17)	140 (-9.9-430)
Oman	2016	Male	70 to 74	11% (5-19%)	89% (81-95%)	0.27 (0.067-0.7)	0.37% (-0.17-1.5%)	0.41% (-0.27-1.8%)	3.1 (-2-14)	68 (-31-280)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Oman	2016	Male	75 to 79	9.5% (4-19%)	90% (81-96%)	0.21 (0.042-0.6)	0.19% (-0.25-1.1%)	0.18% (-0.37-1.2%)	1.3 (-2.6-8.7)	24 (-32-140)
Oman	2016	Male	80 to 84	8.5% (3.2-17%)	91% (83-97%)	0.18 (0.035-0.51)	0.18% (-0.22-0.9%)	0.17% (-0.32-1%)	0.89 (-1.7-5.4)	12 (-15-61)
Oman	2016	Male	85 to 89	8.3% (3.3-17%)	92% (83-97%)	0.16 (0.037-0.45)	0.14% (-0.21-0.75%)	0.11% (-0.32-0.79%)	0.34 (-0.95-2.3)	3.7 (-5.6-19)
Oman	2016	Male	90 to 94	9.3% (3.7-19%)	91% (81-96%)	0.17 (0.037-0.47)	0.061% (-0.33-0.65%)	0.0067% (-0.47-0.67%)	0.0096 (-0.66-0.96)	0.53 (-2.9-5.7)
Oman	2016	Male	95 plus	12% (4.2-25%)	88% (75-96%)	0.2 (0.033-0.63)	0.058% (-0.45-0.79%)	-0.039% (-0.66-0.77%)	-0.015 (-0.25-0.3)	0.09 (-0.69-1.2)
Qatar	1990	Female	All Ages	12% (6.8-18%)	88% (82-93%)	0.36 (0.24-0.49)	0.9% (0.48-1.5%)	1.3% (0.31-2.8%)	5.1 (1.2-11)	300 (150-510)
Qatar	1990	Female	15 to 19	10% (3.6-22%)	90% (78-96%)	0.28 (0.071-0.72)	0.77% (0.38-1.5%)	1.4% (0.42-3.6%)	0.087 (0.025-0.22)	12 (5.4-23)
Qatar	1990	Female	20 to 24	13% (4.4-29%)	87% (71-96%)	0.45 (0.12-1.1)	1.3% (0.71-2.4%)	2.2% (0.67-5.5%)	0.18 (0.052-0.42)	25 (13-45)
Qatar	1990	Female	25 to 29	15% (4.6-31%)	85% (69-95%)	0.54 (0.16-1.1)	1.5% (0.84-2.5%)	2.9% (1.5-8.8%)	0.26 (0.088-0.54)	31 (17-52)
Qatar	1990	Female	30 to 34	17% (4.5-34%)	83% (66-96%)	0.62 (0.21-1.2)	1.6% (0.87-2.6%)	3.3% (1.4-6.2%)	0.44 (0.17-0.85)	40 (21-65)
Qatar	1990	Female	35 to 39	18% (5-35%)	82% (65-95%)	0.71 (0.27-1.3)	1.7% (0.85-3%)	3.8% (1.6-6.7%)	0.78 (0.29-1.5)	54 (26-96)
Qatar	1990	Female	40 to 44	19% (7.3-36%)	81% (64-93%)	0.8 (0.33-1.5)	1.9% (0.77-3.5%)	4% (1.5-7.4%)	0.85 (0.32-1.6)	47 (18-87)
Qatar	1990	Female	45 to 49	20% (7.3-37%)	80% (63-93%)	0.89 (0.29-1.9)	1.9% (0.44-4.3%)	3.7% (1.1-7.9%)	0.66 (0.17-1.5)	31 (7.2-74)
Qatar	1990	Female	50 to 54	21% (8-40%)	79% (60-92%)	0.93 (0.26-2.1)	1.9% (0.11-4.8%)	3.3% (0.6-7.8%)	0.63 (0.11-1.6)	26 (1.6-68)
Qatar	1990	Female	55 to 59	22% (8.6-43%)	78% (57-91%)	0.95 (0.24-2.3)	1.9% (-0.07-5.4%)	3.1% (0.24-8.1%)	0.54 (0.043-1.4)	19 (-0.74-53)
Qatar	1990	Female	60 to 64	23% (9.4-43%)	77% (57-91%)	0.92 (0.21-2.3)	1.4% (-0.69-4.9%)	2.1% (-0.69-7.1%)	0.36 (-0.12-1.2)	10 (-4.9-37)
Qatar	1990	Female	65 to 69	24% (10-42%)	76% (58-90%)	0.81 (0.2-2)	0.88% (-0.93-4%)	1.4% (-1.5-4%)	0.23 (-0.17-0.92)	5.1 (-5.2-23)
Qatar	1990	Female	70 to 74	23% (9.4-41%)	77% (59-91%)	0.66 (0.16-1.8)	0.34% (-1.3-3.6%)	0.53% (-1.6-4.7%)	0.1 (-0.33-0.92)	1.7 (-7-18)
Qatar	1990	Female	75 to 79	20% (7.8-42%)	80% (58-92%)	0.54 (0.11-1.7)	0.011% (-1.6-3%)	0.037% (-2-3.7%)	0.0063 (-0.41-0.74)	0.032 (-6.5-12)
Qatar	1990	Female	80 to 84	19% (6.5-40%)	81% (60-93%)	0.48 (0.092-1.5)	-0.029% (-1.5-2.5%)	-0.039% (-1.8-3.1%)	-0.012 (-0.49-0.8)	-0.11 (-5.5-8.9)
Qatar	1990	Female	85 to 89	19% (6.5-40%)	81% (60-94%)	0.47 (0.089-1.5)	-0.057% (-1.5-2.2%)	-0.087% (-1.8-2.7%)	-0.021 (-0.43-0.61)	-0.12 (-3.2-4.7)
Qatar	1990	Female	90 to 94	21% (7-44%)	79% (56-93%)	0.52 (0.092-1.6)	-0.099% (-1.8-2.4%)	-0.16% (-2.1-2.7%)	-0.02 (-0.25-0.32)	-0.076 (-1.3-1.7)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	1990	Female	95 plus	26% (8.2-51%)	74% (49-92%)	0.6 (0.11-1.8)	-0.11% (-2-2.8%)	-0.22% (-2.5-3.3%)	-0.0088 (-0.099-0.13)	-0.018 (-0.33-0.47)
Qatar	1990	Male	All Ages	13% (8.1-20%)	87% (80-92%)	0.65 (0.49-0.85)	2.2% (1.6-3%)	3% (1.8-4.3%)	25 (14-38)	1600 (1000-2200)
Qatar	1990	Male	15 to 19	7.6% (3.1-14%)	92% (86-97%)	0.2 (0.054-0.47)	1.2% (0.65-2%)	1.6% (0.61-3.6%)	0.19 (0.07-0.44)	25 (13-43)
Qatar	1990	Male	20 to 24	12% (4.8-23%)	88% (77-95%)	0.46 (0.14-0.98)	2.5% (1.5-4.3%)	3.3% (1.3-6.8%)	0.76 (0.27-1.6)	86 (48-150)
Qatar	1990	Male	25 to 29	15% (6.1-27%)	85% (73-94%)	0.7 (0.26-1.3)	3.1% (1.8-4.9%)	4.7% (2.1-8.5%)	2 (0.88-3.7)	190 (110-300)
Qatar	1990	Male	30 to 34	17% (6.4-32%)	83% (68-94%)	0.88 (0.39-1.5)	3.2% (2-4.9%)	5.2% (2.7-8.5%)	3.5 (1.7-6.4)	290 (170-460)
Qatar	1990	Male	35 to 39	18% (7.2-33%)	82% (67-93%)	0.98 (0.45-1.6)	3.3% (1.9-5.1%)	5.5% (2.7-9.1%)	4.2 (1.9-7.5)	300 (160-480)
Qatar	1990	Male	40 to 44	18% (8.5-32%)	82% (68-91%)	1 (0.46-1.8)	3.1% (1.5-5.2%)	5.1% (2.2-8.7%)	3.7 (1.5-6.8)	230 (110-390)
Qatar	1990	Male	45 to 49	18% (7.7-33%)	82% (67-92%)	1.1 (0.37-2.4)	3.1% (1.3-5.4%)	4.7% (1.7-9.1%)	3.1 (1.1-5.9)	170 (74-300)
Qatar	1990	Male	50 to 54	17% (7.6-32%)	83% (68-92%)	1.1 (0.36-2.4)	2.8% (1-5.4%)	4.1% (1.3-7.9%)	2.5 (0.76-5)	120 (44-230)
Qatar	1990	Male	55 to 59	16% (7.2-31%)	84% (69-93%)	1.1 (0.32-2.5)	2.6% (0.87-5.1%)	3.6% (1.1-7.3%)	1.8 (0.55-3.7)	72 (24-140)
Qatar	1990	Male	60 to 64	16% (7.1-30%)	84% (70-93%)	0.96 (0.26-2.3)	2.2% (0.58-4.6%)	2.8% (0.71-5.9%)	1.2 (0.29-2.7)	40 (10-86)
Qatar	1990	Male	65 to 69	15% (7-28%)	85% (72-93%)	0.78 (0.22-1.8)	1.7% (0.31-3.9%)	2.1% (0.35-4.8%)	0.72 (0.12-1.7)	19 (3.7-46)
Qatar	1990	Male	70 to 74	14% (6.5-25%)	86% (75-93%)	0.58 (0.17-1.5)	1.2% (0.098-3.1%)	1.4% (0.036-3.7%)	0.39 (0.01-1.1)	8.4 (0.74-23)
Qatar	1990	Male	75 to 79	12% (5.3-23%)	88% (77-95%)	0.45 (0.12-1.2)	0.87% (-0.013-2.6%)	0.94% (-0.13-2.9%)	0.21 (-0.026-0.67)	3.7 (-0.055-11)
Qatar	1990	Male	80 to 84	11% (4.2-21%)	89% (79-96%)	0.38 (0.089-1)	0.63% (-0.08-2%)	0.64% (-0.23-2.2%)	0.2 (-0.074-0.72)	2.7 (-0.35-8.7)
Qatar	1990	Male	85 to 89	11% (4.3-21%)	89% (79-96%)	0.34 (0.088-0.88)	0.47% (-0.18-1.6%)	0.43% (-0.35-1.7%)	0.11 (-0.089-0.45)	1.1 (-0.44-3.9)
Qatar	1990	Male	90 to 94	12% (4.7-23%)	88% (77-95%)	0.36 (0.084-0.92)	0.35% (-0.36-1.7%)	0.27% (-0.58-1.8%)	0.028 (-0.059-0.18)	0.23 (-0.23-1.1)
Qatar	1990	Male	95 plus	14% (5.5-29%)	86% (71-94%)	0.41 (0.079-1.3)	0.36% (-0.5-2%)	0.21% (-0.81-2.1%)	0.0052 (-0.021-0.052)	0.038 (-0.051-0.21)
Qatar	1995	Female	All Ages	13% (7.3-20%)	87% (80-93%)	0.29 (0.2-0.41)	0.71% (0.34-1.2%)	0.82% (-0.011-2%)	3.5 (-0.046-8.8)	260 (120-450)
Qatar	1995	Female	15 to 19	11% (3.4-23%)	89% (77-97%)	0.22 (0.048-0.58)	0.66% (0.34-1.2%)	1.1% (0.34-2.8%)	0.079 (0.024-0.2)	11 (5.7-21)
Qatar	1995	Female	20 to 24	14% (4.3-30%)	86% (70-96%)	0.35 (0.087-0.84)	1.2% (0.66-2.2%)	1.7% (0.57-4.4%)	0.11 (0.035-0.3)	17 (9.1-32)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	1995	Female	25 to 29	16% (4.6-34%)	84% (66-95%)	0.42 (0.12-0.89)	1.3% (0.79-2.3%)	2.3% (0.82-4.6%)	0.26 (0.091-0.55)	32 (18-54)
Qatar	1995	Female	30 to 34	18% (4.8-36%)	82% (64-95%)	0.48 (0.16-0.9)	1.3% (0.78-2.3%)	2.6% (1.1-4.9%)	0.55 (0.21-1.1)	52 (28-87)
Qatar	1995	Female	35 to 39	19% (5.1-37%)	81% (63-95%)	0.55 (0.2-1)	1.4% (0.66-2.4%)	2.9% (1.1-5.4%)	0.74 (0.27-1.5)	52 (24-95)
Qatar	1995	Female	40 to 44	20% (7.6-37%)	80% (63-92%)	0.61 (0.26-1.1)	1.4% (0.46-2.8%)	3% (1-5.7%)	0.76 (0.28-1.6)	41 (13-84)
Qatar	1995	Female	45 to 49	21% (7.4-38%)	79% (62-93%)	0.67 (0.21-1.4)	1.3% (0.1-3.2%)	2.5% (0.6-6%)	0.58 (0.13-1.4)	26 (2.1-66)
Qatar	1995	Female	50 to 54	22% (8.9-41%)	78% (59-91%)	0.71 (0.2-1.6)	1.2% (-0.23-3.5%)	2.2% (0.2-5.9%)	0.45 (0.038-1.2)	17 (-3.9-50)
Qatar	1995	Female	55 to 59	23% (9.5-43%)	77% (57-91%)	0.71 (0.19-1.7)	1% (-0.52-3.8%)	1.9% (-0.27-5.9%)	0.39 (-0.061-1.2)	12 (-6-44)
Qatar	1995	Female	60 to 64	24% (10-43%)	76% (57-90%)	0.69 (0.16-1.7)	0.52% (-1.2-3.3%)	1% (-1.2-4.8%)	0.21 (-0.27-1.1)	4.9 (-11-30)
Qatar	1995	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.6 (0.15-1.4)	0.057% (-1.5-2.4%)	0.32% (-1.7-3.3%)	0.072 (-0.39-0.74)	0.44 (-12-18)
Qatar	1995	Female	70 to 74	23% (10-41%)	77% (59-90%)	0.48 (0.13-1.3)	-0.33% (-1.9-1.9%)	-0.28% (-2.2-2.4%)	-0.06 (-0.48-0.55)	-1.8 (-10-11)
Qatar	1995	Female	75 to 79	21% (8.3-41%)	79% (59-92%)	0.39 (0.082-1.3)	-0.53% (-2-1.3%)	-0.59% (-2.3-1.7%)	-0.15 (-0.61-0.39)	-2.5 (-9.5-6.1)
Qatar	1995	Female	80 to 84	20% (6.7-40%)	80% (60-93%)	0.36 (0.066-1.1)	-0.54% (-2-0.95%)	-0.63% (-2.3-1.2%)	-0.19 (-0.74-0.34)	-2.2 (-8.7-3.7)
Qatar	1995	Female	85 to 89	20% (6.8-42%)	80% (58-93%)	0.35 (0.071-1.1)	-0.56% (-2-0.78%)	-0.66% (-2.4-0.98%)	-0.15 (-0.54-0.2)	-1.1 (-4.3-1.6)
Qatar	1995	Female	90 to 94	23% (7.7-45%)	77% (55-92%)	0.39 (0.073-1.1)	-0.62% (-2.2-0.94%)	-0.76% (-2.7-1.1%)	-0.095 (-0.33-0.13)	-0.48 (-1.7-0.75)
Qatar	1995	Female	95 plus	27% (8.9-52%)	73% (48-91%)	0.46 (0.072-1.3)	-0.68% (-2.5-1.4%)	-0.88% (-3.1-1.6%)	-0.038 (-0.13-0.069)	-0.12 (-0.45-0.26)
Qatar	1995	Male	All Ages	14% (8.5-20%)	86% (80-92%)	0.48 (0.36-0.64)	1.8% (1.2-2.4%)	2.2% (1.2-3.3%)	20 (10-32)	1300 (860-1900)
Qatar	1995	Male	15 to 19	7.8% (3.3-15%)	92% (85-97%)	0.15 (0.039-0.37)	1% (0.59-1.8%)	1.4% (0.52-3%)	0.13 (0.048-0.3)	17 (9.3-29)
Qatar	1995	Male	20 to 24	12% (5-23%)	88% (77-95%)	0.33 (0.1-0.71)	2.1% (1.3-3.4%)	2.4% (0.95-4.8%)	0.77 (0.28-1.7)	95 (53-160)
Qatar	1995	Male	25 to 29	15% (6.2-27%)	85% (73-94%)	0.51 (0.19-0.97)	2.6% (1.6-4%)	3.6% (1.7-6.5%)	1.9 (0.81-3.5)	190 (110-300)
Qatar	1995	Male	30 to 34	17% (6.6-32%)	83% (68-93%)	0.64 (0.27-1.1)	2.6% (1.7-3.9%)	3.9% (2.1-6.5%)	2.5 (1.3-4.4)	220 (130-330)
Qatar	1995	Male	35 to 39	19% (7.5-33%)	81% (67-93%)	0.72 (0.33-1.2)	2.6% (1.6-4%)	4.1% (2.1-6.8%)	3.5 (1.7-6)	260 (150-400)
Qatar	1995	Male	40 to 44	19% (8.6-33%)	81% (67-91%)	0.77 (0.36-1.3)	2.3% (1.2-3.9%)	3.5% (1.5-6.4%)	3.1 (1.2-5.8)	200 (98-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	1995	Male	45 to 49	19% (7.7-34%)	81% (66-92%)	0.82 (0.28-1.7)	2.3% (0.89-4.5%)	3.3% (0.99-6.9%)	2.6 (0.76-5.5)	140 (56-290)
Qatar	1995	Male	50 to 54	18% (7.9-35%)	82% (65-92%)	0.82 (0.27-1.9)	2.1% (0.67-4.4%)	2.9% (0.8-6.5%)	2 (0.5-4.5)	93 (30-200)
Qatar	1995	Male	55 to 59	17% (7.3-33%)	83% (67-93%)	0.78 (0.23-1.8)	1.9% (0.52-4.2%)	2.5% (0.57-5.7%)	1.5 (0.35-3.6)	60 (16-130)
Qatar	1995	Male	60 to 64	16% (7.1-31%)	84% (69-93%)	0.7 (0.2-1.7)	1.5% (0.25-3.7%)	1.9% (0.23-4.8%)	1 (0.13-2.6)	33 (5.6-80)
Qatar	1995	Male	65 to 69	16% (7.3-28%)	84% (72-93%)	0.57 (0.16-1.3)	1% (-0.019-2.9%)	1.2% (-0.11-3.6%)	0.5 (-0.042-1.5)	14 (-0.28-39)
Qatar	1995	Male	70 to 74	14% (6.8-25%)	86% (75-93%)	0.42 (0.12-1.1)	0.64% (-0.15-2.1%)	0.74% (-0.27-2.5%)	0.27 (-0.1-0.99)	6 (-1.4-21)
Qatar	1995	Male	75 to 79	12% (5.4-23%)	88% (77-95%)	0.32 (0.082-0.94)	0.47% (-0.22-1.8%)	0.5% (-0.37-2%)	0.19 (-0.13-0.78)	3.3 (-1.4-13)
Qatar	1995	Male	80 to 84	11% (4.4-22%)	89% (78-96%)	0.28 (0.065-0.77)	0.29% (-0.31-1.3%)	0.26% (-0.45-1.4%)	0.1 (-0.18-0.53)	1.5 (-1.4-6.7)
Qatar	1995	Male	85 to 89	11% (4.4-22%)	89% (78-96%)	0.25 (0.063-0.66)	0.19% (-0.34-0.95%)	0.13% (-0.52-0.99%)	0.034 (-0.14-0.28)	0.47 (-0.85-2.4)
Qatar	1995	Male	90 to 94	12% (5-25%)	88% (75-95%)	0.26 (0.06-0.65)	0.09% (-0.51-0.92%)	-0.0066% (-0.74-0.94%)	-0.0011 (-0.11-0.14)	0.084 (-0.48-0.85)
Qatar	1995	Male	95 plus	15% (5.5-32%)	85% (68-95%)	0.3 (0.057-0.82)	0.083% (-0.64-1.2%)	-0.077% (-1-1.2%)	-0.0027 (-0.034-0.042)	0.012 (-0.093-0.17)
Qatar	2000	Female	All Ages	12% (7-19%)	88% (81-93%)	0.28 (0.19-0.39)	0.59% (0.22-1.1%)	0.52% (-0.26-1.6%)	2.5 (-1.3-7.7)	240 (83-440)
Qatar	2000	Female	15 to 19	10% (3.1-23%)	90% (77-97%)	0.2 (0.044-0.51)	0.61% (0.32-1.1%)	0.98% (0.29-2.5%)	0.096 (0.027-0.26)	15 (7.7-29)
Qatar	2000	Female	20 to 24	13% (3.7-28%)	87% (72-96%)	0.32 (0.075-0.78)	1.1% (0.65-1.9%)	1.5% (0.44-3.8%)	0.15 (0.042-0.4)	25 (14-46)
Qatar	2000	Female	25 to 29	14% (4.1-31%)	86% (69-96%)	0.38 (0.096-0.87)	1.2% (0.71-2%)	2% (0.62-4.1%)	0.19 (0.062-0.43)	27 (15-45)
Qatar	2000	Female	30 to 34	16% (4.3-33%)	84% (67-96%)	0.43 (0.13-0.85)	1.2% (0.66-2%)	2.1% (0.79-4%)	0.34 (0.12-0.77)	37 (21-64)
Qatar	2000	Female	35 to 39	17% (4.6-33%)	83% (67-95%)	0.49 (0.16-0.88)	1.1% (0.56-2.1%)	2.3% (0.85-4.5%)	0.68 (0.23-1.4)	51 (25-92)
Qatar	2000	Female	40 to 44	18% (6.2-34%)	82% (66-94%)	0.54 (0.22-0.97)	1.1% (0.43-2.1%)	2.3% (0.84-4.4%)	0.8 (0.28-1.6)	46 (16-92)
Qatar	2000	Female	45 to 49	19% (6.7-35%)	81% (65-93%)	0.59 (0.2-1.2)	0.92% (-0.074-2.4%)	1.9% (0.3-4.5%)	0.61 (0.095-1.5)	29 (-2.1-75)
Qatar	2000	Female	50 to 54	20% (7.6-38%)	80% (62-92%)	0.63 (0.18-1.4)	0.77% (-0.51-2.6%)	1.5% (-0.25-4.3%)	0.42 (-0.069-1.2)	16 (-10-55)
Qatar	2000	Female	55 to 59	21% (8-40%)	79% (60-92%)	0.64 (0.17-1.5)	0.57% (-0.88-2.8%)	1.1% (-0.84-4.2%)	0.28 (-0.19-1.1)	8.3 (-14-42)
Qatar	2000	Female	60 to 64	22% (8.8-42%)	78% (58-91%)	0.62 (0.15-1.6)	0.095% (-1.5-2.6%)	0.32% (-1.7-3.4%)	0.088 (-0.47-1)	1.2 (-19-33)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	2000	Female	65 to 69	22% (9.8-40%)	78% (60-90%)	0.54 (0.14-1.3)	-0.28% (-1.8-1.7%)	-0.21% (-2.2-2.4%)	-0.059 (-0.6-0.7)	-2.8 (-19-16)
Qatar	2000	Female	70 to 74	21% (8.7-38%)	79% (62-91%)	0.43 (0.11-1.1)	-0.59% (-2.1-1.2%)	-0.68% (-2.5-1.5%)	-0.2 (-0.74-0.45)	-4.7 (-17-9.4)
Qatar	2000	Female	75 to 79	19% (7.1-38%)	81% (62-93%)	0.35 (0.072-1)	-0.69% (-2.3-0.82%)	-0.85% (-2.7-0.98%)	-0.22 (-0.75-0.25)	-3.6 (-12-4.1)
Qatar	2000	Female	80 to 84	17% (5.9-37%)	83% (63-94%)	0.31 (0.061-0.96)	-0.66% (-2.1-0.52%)	-0.81% (-2.6-0.59%)	-0.26 (-0.78-0.18)	-2.9 (-9.1-2.3)
Qatar	2000	Female	85 to 89	18% (6-39%)	82% (61-94%)	0.32 (0.062-0.98)	-0.73% (-2.2-0.39%)	-0.9% (-2.7-0.41%)	-0.24 (-0.73-0.11)	-1.8 (-5.7-1)
Qatar	2000	Female	90 to 94	21% (6.7-42%)	79% (58-93%)	0.36 (0.065-1)	-0.81% (-2.5-0.64%)	-1% (-3-0.66%)	-0.13 (-0.38-0.088)	-0.66 (-2-0.54)
Qatar	2000	Female	95 plus	25% (7.5-50%)	75% (50-92%)	0.43 (0.065-1.3)	-0.91% (-2.8-0.83%)	-1.2% (-3.5-0.94%)	-0.058 (-0.17-0.044)	-0.18 (-0.56-0.17)
Qatar	2000	Male	All Ages	12% (7.8-18%)	88% (82-92%)	0.44 (0.33-0.58)	1.7% (1.1-2.4%)	2.1% (1.1-3.2%)	23 (12-39)	1500 (970-2200)
Qatar	2000	Male	15 to 19	7% (3-13%)	93% (87-97%)	0.13 (0.036-0.31)	0.99% (0.56-1.6%)	1.2% (0.43-2.8%)	0.24 (0.076-0.53)	32 (17-54)
Qatar	2000	Male	20 to 24	11% (4.3-21%)	89% (79-96%)	0.3 (0.09-0.66)	2.2% (1.3-3.5%)	2.6% (1-5.3%)	0.64 (0.24-1.4)	76 (44-130)
Qatar	2000	Male	25 to 29	14% (5.3-25%)	86% (75-95%)	0.46 (0.15-0.93)	2.5% (1.5-4%)	3.4% (1.4-6.4%)	1.6 (0.62-3.1)	160 (95-270)
Qatar	2000	Male	30 to 34	15% (5.7-29%)	85% (71-94%)	0.58 (0.23-1)	2.5% (1.5-3.9%)	3.7% (1.8-6.4%)	2.5 (1.1-4.7)	220 (130-360)
Qatar	2000	Male	35 to 39	17% (6.6-30%)	83% (70-93%)	0.64 (0.29-1.1)	2.4% (1.4-3.8%)	3.8% (1.9-6.5%)	3.2 (1.5-5.8)	240 (140-390)
Qatar	2000	Male	40 to 44	17% (7.8-30%)	83% (70-92%)	0.68 (0.32-1.2)	2.3% (1.3-3.6%)	3.5% (1.6-6%)	4 (1.8-7.3)	260 (140-420)
Qatar	2000	Male	45 to 49	17% (6.5-31%)	83% (69-94%)	0.72 (0.26-1.5)	2.1% (0.96-4%)	3.1% (1.2-6.2%)	3.8 (1.4-7.9)	210 (95-400)
Qatar	2000	Male	50 to 54	16% (6.9-32%)	84% (68-93%)	0.71 (0.24-1.5)	1.9% (0.68-3.9%)	2.5% (0.75-5.6%)	2.7 (0.77-5.9)	130 (46-270)
Qatar	2000	Male	55 to 59	15% (6.4-29%)	85% (71-94%)	0.69 (0.22-1.5)	1.7% (0.49-3.7%)	2.2% (0.59-5%)	1.9 (0.5-4.3)	75 (22-170)
Qatar	2000	Male	60 to 64	15% (6.3-28%)	85% (72-94%)	0.62 (0.17-1.5)	1.4% (0.24-3.4%)	1.7% (0.25-4.3%)	1.2 (0.18-3.2)	41 (7.2-100)
Qatar	2000	Male	65 to 69	14% (6.2-26%)	86% (74-94%)	0.51 (0.15-1.2)	0.92% (0.0034-2.6%)	1.1% (-0.029-2.3%)	0.74 (-0.029-2.3)	20 (0.076-60)
Qatar	2000	Male	70 to 74	13% (5.8-23%)	87% (77-94%)	0.38 (0.11-0.94)	0.56% (-0.16-1.9%)	0.62% (-0.25-2.2%)	0.36 (-0.15-1.3)	7.9 (-2.3-27)
Qatar	2000	Male	75 to 79	11% (4.7-21%)	89% (79-95%)	0.29 (0.071-0.83)	0.39% (-0.21-1.5%)	0.39% (-0.37-1.6%)	0.18 (-0.17-0.77)	3.4 (-1.8-13)
Qatar	2000	Male	80 to 84	9.9% (4-19%)	90% (81-96%)	0.24 (0.059-0.64)	0.28% (-0.23-1.1%)	0.24% (-0.37-1.1%)	0.1 (-0.15-0.49)	1.6 (-1.3-6.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	2000	Male	85 to 89	9.6% (3.9-19%)	90% (81-96%)	0.22 (0.057-0.57)	0.18% (-0.28-0.78%)	0.12% (-0.42-0.78%)	0.036 (-0.13-0.24)	0.51 (-0.79-2.2)
Qatar	2000	Male	90 to 94	11% (4.5-21%)	89% (79-95%)	0.23 (0.056-0.57)	0.076% (-0.46-0.72%)	-0.025% (-0.64-0.71%)	-0.0035 (-0.094-0.1)	0.069 (-0.4-0.65)
Qatar	2000	Male	95 plus	13% (4.8-28%)	87% (72-95%)	0.27 (0.053-0.74)	0.058% (-0.6-0.95%)	-0.1% (-0.91-0.83%)	-0.0049 (-0.044-0.043)	0.012 (-0.12-0.19)
Qatar	2005	Female	All Ages	12% (6.9-18%)	88% (82-93%)	0.38 (0.25-0.53)	0.86% (0.41-1.4%)	1.1% (0.16-2.5%)	5.5 (0.78-13)	420 (210-710)
Qatar	2005	Female	15 to 19	9.4% (3-21%)	91% (79-97%)	0.26 (0.06-0.69)	0.69% (0.35-1.3%)	1.4% (0.43-3.5%)	0.1 (0.031-0.27)	15 (7.7-29)
Qatar	2005	Female	20 to 24	12% (3.6-26%)	88% (74-96%)	0.41 (0.097-0.98)	1.2% (0.7-2.1%)	2% (0.59-4.8%)	0.2 (0.06-0.51)	34 (19-59)
Qatar	2005	Female	25 to 29	13% (4-28%)	87% (72-96%)	0.48 (0.14-1)	1.3% (0.77-2.1%)	2.4% (0.73-4.9%)	0.38 (0.12-0.82)	56 (31-91)
Qatar	2005	Female	30 to 34	15% (4.2-32%)	85% (68-96%)	0.56 (0.17-1.1)	1.3% (0.76-2.1%)	2.7% (0.96-5.2%)	0.6 (0.21-1.2)	68 (39-110)
Qatar	2005	Female	35 to 39	17% (4.8-32%)	83% (68-95%)	0.65 (0.23-1.2)	1.4% (0.69-2.3%)	3.3% (1.3-6%)	0.79 (0.29-1.5)	63 (31-110)
Qatar	2005	Female	40 to 44	17% (6.6-33%)	83% (67-93%)	0.71 (0.3-1.3)	1.4% (0.58-2.6%)	3.3% (1.4-6%)	0.98 (0.37-1.9)	61 (25-110)
Qatar	2005	Female	45 to 49	18% (6.7-34%)	82% (66-93%)	0.78 (0.26-1.6)	1.3% (0.23-3.1%)	2.9% (0.87-6%)	1 (0.29-2.3)	54 (10-120)
Qatar	2005	Female	50 to 54	19% (7.5-37%)	81% (63-93%)	0.81 (0.23-1.7)	1.3% (-0.15-3.5%)	2.5% (0.33-6.1%)	0.8 (0.1-2)	35 (-4.2-98)
Qatar	2005	Female	55 to 59	20% (7.5-38%)	80% (62-93%)	0.83 (0.22-1.9)	1.2% (-0.41-4.2%)	2.2% (-0.12-6.3%)	0.63 (-0.038-1.9)	23 (-7.8-81)
Qatar	2005	Female	60 to 64	21% (7.9-40%)	79% (60-92%)	0.81 (0.19-2.1)	0.76% (-1-3.8%)	1.3% (-1.1-5.3%)	0.31 (-0.27-1.4)	9.2 (-13-48)
Qatar	2005	Female	65 to 69	21% (8.9-39%)	79% (61-91%)	0.71 (0.18-1.7)	0.34% (-1.3-3%)	0.62% (-1.5-4.2%)	0.17 (-0.43-1.1)	3.6 (-14-32)
Qatar	2005	Female	70 to 74	20% (8.3-37%)	80% (63-92%)	0.56 (0.15-1.5)	-0.087% (-1.7-2.6%)	-0.053% (-2.1-3.3%)	-0.015 (-0.56-0.96)	-0.69 (-13-21)
Qatar	2005	Female	75 to 79	18% (6.6-36%)	82% (64-93%)	0.46 (0.093-1.4)	-0.28% (-1.8-2%)	-0.36% (-2.2-2.2%)	-0.1 (-0.64-0.65)	-1.6 (-11-11)
Qatar	2005	Female	80 to 84	16% (5.3-36%)	84% (64-95%)	0.41 (0.076-1.2)	-0.27% (-1.6-1.5%)	-0.36% (-2-1.7%)	-0.12 (-0.67-0.6)	-1.3 (-7.8-7.5)
Qatar	2005	Female	85 to 89	17% (5.2-39%)	83% (61-95%)	0.41 (0.08-1.4)	-0.34% (-1.8-1.4%)	-0.46% (-2.2-1.6%)	-0.13 (-0.61-0.44)	-0.91 (-4.8-3.6)
Qatar	2005	Female	90 to 94	20% (5.9-43%)	80% (57-94%)	0.47 (0.086-1.4)	-0.39% (-2.1-1.7%)	-0.54% (-2.6-1.8%)	-0.087 (-0.41-0.29)	-0.4 (-2.1-1.7)
Qatar	2005	Female	95 plus	24% (7.2-48%)	76% (52-93%)	0.56 (0.09-1.7)	-0.44% (-2.5-2.2%)	-0.66% (-3.1-2.5%)	-0.034 (-0.16-0.13)	-0.095 (-0.53-0.47)
Qatar	2005	Male	All Ages	12% (7.6-17%)	88% (83-92%)	0.58 (0.43-0.75)	2.2% (1.5-3%)	2.9% (1.6-4.3%)	41 (22-66)	2500 (1700-3600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	2005	Male	15 to 19	6.5% (2.8-13%)	93% (87-97%)	0.16 (0.047-0.4)	1.2% (0.68-2%)	1.7% (0.7-3.6%)	0.27 (0.11-0.59)	33 (19-57)
Qatar	2005	Male	20 to 24	10% (4-20%)	90% (80-96%)	0.37 (0.11-0.83)	2.3% (1.4-3.9%)	2.9% (1.2-6%)	1.3 (0.5-2.8)	150 (90-260)
Qatar	2005	Male	25 to 29	12% (5.2-23%)	88% (77-95%)	0.57 (0.2-1.1)	2.7% (1.6-4.3%)	3.9% (1.6-7.3%)	2.8 (1.1-5.3)	280 (160-460)
Qatar	2005	Male	30 to 34	14% (5.6-28%)	86% (72-94%)	0.72 (0.3-1.3)	2.7% (1.6-4.3%)	4.3% (2-7.5%)	3.7 (1.6-6.8)	330 (190-530)
Qatar	2005	Male	35 to 39	16% (6.2-29%)	84% (71-94%)	0.81 (0.34-1.4)	2.8% (1.6-4.5%)	4.7% (2.2-8.1%)	4.6 (2.1-8)	340 (190-540)
Qatar	2005	Male	40 to 44	16% (7.2-29%)	84% (71-93%)	0.87 (0.38-1.6)	2.7% (1.5-4.6%)	4.5% (2.1-8%)	5.5 (2.4-10)	350 (180-580)
Qatar	2005	Male	45 to 49	16% (6-29%)	84% (71-94%)	0.9 (0.31-1.8)	2.7% (1.3-4.7%)	4.1% (1.7-7.5%)	7 (2.7-13)	380 (170-670)
Qatar	2005	Male	50 to 54	15% (6.2-29%)	85% (71-94%)	0.89 (0.3-1.9)	2.4% (0.93-4.5%)	3.5% (1.2-6.6%)	6.1 (2.1-12)	280 (110-540)
Qatar	2005	Male	55 to 59	14% (6.1-27%)	86% (73-94%)	0.86 (0.27-1.9)	2.2% (0.76-4.5%)	3.1% (1-6.4%)	4.8 (1.5-9.8)	190 (64-380)
Qatar	2005	Male	60 to 64	14% (5.9-26%)	86% (74-94%)	0.78 (0.22-1.9)	2.4% (0.52-4.1%)	2.4% (0.71-5.3%)	2.5 (0.67-5.5)	80 (23-180)
Qatar	2005	Male	65 to 69	13% (5.9-25%)	87% (75-94%)	0.64 (0.19-1.5)	1.5% (0.35-3.4%)	1.9% (0.43-4.3%)	1.1 (0.25-2.6)	30 (7.2-69)
Qatar	2005	Male	70 to 74	12% (5.3-22%)	88% (78-95%)	0.48 (0.13-1.2)	0.98% (0.12-2.7%)	1.2% (0.12-3.2%)	0.75 (0.075-2.2)	16 (2.1-45)
Qatar	2005	Male	75 to 79	10% (4.4-20%)	90% (80-96%)	0.36 (0.084-1)	0.77% (0.02-2.1%)	0.77% (-0.095-2.4%)	0.34 (-0.037-1)	6 (0.15-18)
Qatar	2005	Male	80 to 84	9.1% (3.5-18%)	91% (82-96%)	0.3 (0.072-0.83)	0.59% (-0.0049-1.7%)	0.58% (-0.13-1.9%)	0.24 (-0.054-0.81)	3.3 (-0.024-9.6)
Qatar	2005	Male	85 to 89	8.9% (3.5-18%)	91% (82-96%)	0.28 (0.068-0.73)	0.42% (-0.12-1.3%)	0.36% (-0.27-1.3%)	0.13 (-0.1-0.49)	1.4 (-0.44-4.4)
Qatar	2005	Male	90 to 94	10% (4-21%)	90% (79-96%)	0.29 (0.07-0.76)	0.29% (-0.29-1.3%)	0.19% (-0.51-1.3%)	0.035 (-0.091-0.23)	0.33 (-0.34-1.4)
Qatar	2005	Male	95 plus	12% (4.4-26%)	88% (74-96%)	0.34 (0.063-1)	0.29% (-0.44-1.5%)	0.12% (-0.77-1.5%)	0.0061 (-0.038-0.074)	0.06 (-0.093-0.3)
Qatar	2010	Female	All Ages	12% (6.5-18%)	88% (82-93%)	0.51 (0.33-0.75)	1.1% (0.64-1.7%)	1.8% (0.71-3.5%)	11 (3.9-22)	810 (460-1300)
Qatar	2010	Female	15 to 19	8.7% (2.9-19%)	91% (81-97%)	0.32 (0.073-0.86)	0.75% (0.38-1.4%)	1.7% (0.58-4.2%)	0.12 (0.038-0.29)	20 (9.9-36)
Qatar	2010	Female	20 to 24	11% (3.3-24%)	89% (76-97%)	0.51 (0.13-1.2)	1.3% (0.74-2.2%)	2.5% (0.72-5.8%)	0.37 (0.097-0.91)	66 (35-110)
Qatar	2010	Female	25 to 29	13% (3.7-27%)	87% (73-96%)	0.6 (0.16-1.3)	1.4% (0.82-2.2%)	2.9% (0.94-6%)	0.78 (0.24-1.6)	120 (67-200)
Qatar	2010	Female	30 to 34	14% (3.4-30%)	86% (70-97%)	0.68 (0.22-1.3)	1.4% (0.81-2.3%)	3.3% (1.2-6.3%)	0.97 (0.35-1.9)	120 (69-190)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	2010	Female	35 to 39	16% (4.3-30%)	84% (70-96%)	0.79 (0.28-1.5)	1.5% (0.79-2.5%)	4.1% (1.6-7.2%)	1.3 (0.49-2.4)	110 (58-180)
Qatar	2010	Female	40 to 44	16% (6-30%)	84% (70-94%)	0.88 (0.35-1.7)	1.6% (0.73-2.9%)	4.2% (1.8-7.6%)	1.4 (0.56-2.7)	100 (45-180)
Qatar	2010	Female	45 to 49	17% (5.9-32%)	83% (68-94%)	0.99 (0.3-2.1)	1.7% (0.5-3.6%)	3.9% (1.3-8%)	1.4 (0.46-2.9)	84 (25-180)
Qatar	2010	Female	50 to 54	18% (6.6-34%)	82% (66-93%)	1 (0.27-2.3)	1.7% (0.26-4.1%)	3.5% (0.91-7.6%)	1.6 (0.37-3.4)	80 (13-200)
Qatar	2010	Female	55 to 59	18% (7-36%)	82% (64-93%)	1.1 (0.26-2.6)	1.8% (0.033-5%)	3.4% (0.62-8.2%)	1.3 (0.23-3.3)	56 (1.1-160)
Qatar	2010	Female	60 to 64	19% (7.6-38%)	81% (62-92%)	1 (0.23-2.7)	1.4% (-0.4-4.7%)	2.4% (-0.13-7%)	0.61 (-0.03-1.8)	22 (-5.8-75)
Qatar	2010	Female	65 to 69	20% (8.6-36%)	80% (64-91%)	0.89 (0.23-2.2)	1.1% (-0.56-4.2%)	1.8% (-0.41-6%)	0.54 (-0.11-1.8)	15 (-7.4-56)
Qatar	2010	Female	70 to 74	19% (8.1-35%)	81% (65-92%)	0.71 (0.18-2)	0.63% (-0.91-3.9%)	0.94% (-1.1-5%)	0.41 (-0.46-2.3)	8.8 (-12-54)
Qatar	2010	Female	75 to 79	17% (6.4-34%)	83% (66-94%)	0.58 (0.12-1.9)	0.34% (-1-3.3%)	0.41% (-1.3-4%)	0.14 (-0.45-1.4)	2.5 (-7.6-24)
Qatar	2010	Female	80 to 84	15% (5.3-33%)	85% (67-95%)	0.52 (0.1-1.6)	0.34% (-1-2.9%)	0.37% (-1.2-3.5%)	0.14 (-0.47-1.3)	1.9 (-5.4-16)
Qatar	2010	Female	85 to 89	16% (5.2-36%)	84% (64-95%)	0.52 (0.096-1.7)	0.25% (-1.1-2.8%)	0.23% (-1.4-3.3%)	0.076 (-0.47-1.1)	0.82 (-3.6-9.5)
Qatar	2010	Female	90 to 94	18% (6.1-40%)	82% (60-94%)	0.58 (0.11-1.8)	0.25% (-1.4-2.9%)	0.2% (-1.8-3.2%)	0.037 (-0.32-0.61)	0.3 (-1.6-3.4)
Qatar	2010	Female	95 plus	22% (7-47%)	78% (53-93%)	0.68 (0.11-2.1)	0.27% (-1.7-3.6%)	0.17% (-2.2-4.1%)	0.012 (-0.15-0.28)	0.08 (-0.48-1.1)
Qatar	2010	Male	All Ages	12% (7.2-18%)	88% (82-93%)	0.81 (0.58-1)	2.6% (1.9-3.4%)	3.9% (2.6-5.5%)	86 (51-130)	6000 (4200-8200)
Qatar	2010	Male	15 to 19	6.3% (2.6-12%)	94% (88-97%)	0.21 (0.057-0.52)	1.4% (0.81-2.3%)	2.4% (1-4.9%)	0.47 (0.19-0.98)	57 (33-99)
Qatar	2010	Male	20 to 24	9.7% (3.9-19%)	90% (81-96%)	0.48 (0.15-1.1)	2.5% (1.5-4.1%)	3.4% (1.3-7%)	3.7 (1.4-7.8)	470 (270-810)
Qatar	2010	Male	25 to 29	12% (4.7-23%)	88% (77-95%)	0.74 (0.27-1.4)	2.9% (1.7-4.4%)	4.6% (1.9-8.3%)	8.6 (3.5-16)	940 (540-1500)
Qatar	2010	Male	30 to 34	14% (4.9-26%)	86% (74-95%)	0.93 (0.41-1.6)	2.9% (1.8-4.3%)	5.2% (2.5-8.7%)	10 (4.9-18)	980 (590-1500)
Qatar	2010	Male	35 to 39	15% (5.9-28%)	85% (72-94%)	1 (0.45-1.8)	3% (1.8-4.5%)	5.8% (2.9-9.4%)	13 (6.1-22)	1000 (590-1600)
Qatar	2010	Male	40 to 44	15% (6.8-27%)	85% (73-93%)	1.1 (0.45-2)	3% (1.7-4.9%)	5.6% (2.7-9.6%)	13 (5.6-22)	870 (460-1400)
Qatar	2010	Male	45 to 49	15% (5.8-28%)	85% (72-94%)	1.2 (0.38-2.5)	3% (1.5-5%)	5.2% (2.3-9.1%)	12 (4.8-21)	670 (320-1100)
Qatar	2010	Male	50 to 54	14% (6.2-29%)	86% (71-94%)	1.2 (0.37-2.6)	2.8% (1.2-4.9%)	4.4% (1.7-7.7%)	9.7 (3.6-18)	470 (200-850)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	2010	Male	55 to 59	14% (5.8-27%)	86% (73-94%)	1.1 (0.31-2.7)	2.6% (1.1-4.9%)	3.8% (1.6-7.3%)	6.5 (2.4-13)	270 (110-520)
Qatar	2010	Male	60 to 64	13% (5.7-26%)	87% (74-94%)	1 (0.27-2.6)	2.4% (0.92-4.6%)	3.3% (1.3-6.3%)	3.2 (1.2-6.5)	110 (41-220)
Qatar	2010	Male	65 to 69	13% (5.7-24%)	87% (76-94%)	0.84 (0.24-2)	2.1% (0.76-4%)	2.8% (1-5.3%)	1.8 (0.63-3.7)	50 (18-100)
Qatar	2010	Male	70 to 74	12% (5.1-21%)	88% (79-95%)	0.63 (0.17-1.6)	1.4% (0.39-3.2%)	1.8% (0.47-4%)	2.5 (0.64-5.9)	56 (14-130)
Qatar	2010	Male	75 to 79	10% (4.2-20%)	90% (80-96%)	0.48 (0.11-1.4)	1.2% (0.33-2.9%)	1.3% (0.25-3.3%)	0.65 (0.11-1.6)	12 (3.2-29)
Qatar	2010	Male	80 to 84	8.9% (3.4-18%)	91% (82-97%)	0.4 (0.094-1.2)	1.2% (0.36-2.6%)	1.2% (0.28-2.8%)	0.47 (0.11-1.2)	6.6 (2-15)
Qatar	2010	Male	85 to 89	8.7% (3.4-17%)	91% (83-97%)	0.36 (0.089-0.94)	0.89% (0.19-2.1%)	0.8% (0.033-2.1%)	0.27 (0.011-0.73)	2.8 (0.61-6.8)
Qatar	2010	Male	90 to 94	9.9% (4-20%)	90% (80-96%)	0.39 (0.089-0.99)	0.73% (0.037-1.9%)	0.58% (-0.2-2%)	0.13 (-0.043-0.44)	1 (0.05-2.8)
Qatar	2010	Male	95 plus	12% (4.5-26%)	88% (74-96%)	0.45 (0.084-1.4)	0.75% (-0.073-2.4%)	0.52% (-0.44-2.4%)	0.034 (-0.029-0.15)	0.21 (-0.019-0.65)
Qatar	2016	Female	All Ages	12% (6.3-19%)	88% (81-94%)	0.62 (0.35-1)	2.4% (0.72-2.1%)	2.4% (0.91-4.4%)	19 (6.8-38)	1300 (670-2200)
Qatar	2016	Female	15 to 19	8.8% (3.1-20%)	91% (80-97%)	0.37 (0.082-1)	0.78% (0.39-1.5%)	2% (0.63-4.7%)	0.21 (0.062-0.53)	33 (16-64)
Qatar	2016	Female	20 to 24	11% (3.4-24%)	89% (76-97%)	0.59 (0.14-1.5)	1.4% (0.78-2.4%)	2.9% (0.94-6.6%)	0.42 (0.13-1)	72 (38-120)
Qatar	2016	Female	25 to 29	13% (3.3-27%)	87% (73-97%)	0.69 (0.18-1.5)	1.5% (0.83-2.4%)	3.4% (1.1-7.1%)	0.94 (0.25-2.2)	140 (74-250)
Qatar	2016	Female	30 to 34	14% (3.4-31%)	86% (69-97%)	0.8 (0.23-1.7)	1.5% (0.84-2.6%)	3.9% (1.4-7.4%)	1.8 (0.53-3.8)	220 (110-390)
Qatar	2016	Female	35 to 39	16% (4.5-32%)	84% (68-96%)	0.94 (0.3-2)	1.6% (0.8-2.9%)	4.7% (1.7-8.6%)	2.3 (0.73-4.9)	210 (96-370)
Qatar	2016	Female	40 to 44	17% (6-31%)	83% (69-94%)	1.1 (0.39-2.1)	1.8% (0.71-3.4%)	4.8% (2-8.9%)	2.6 (0.89-5.5)	180 (72-350)
Qatar	2016	Female	45 to 49	17% (6.2-32%)	83% (68-94%)	1.2 (0.32-2.6)	1.8% (0.61-4%)	4.4% (1.5-8.8%)	2.6 (0.79-6)	160 (51-350)
Qatar	2016	Female	50 to 54	18% (6.9-35%)	82% (65-93%)	1.2 (0.3-2.9)	1.9% (0.39-4.7%)	4% (1.2-8.7%)	2.1 (0.57-4.8)	110 (23-260)
Qatar	2016	Female	55 to 59	19% (7.4-38%)	81% (62-93%)	1.3 (0.29-3.2)	2% (0.25-5.2%)	3.9% (0.9-9%)	1.9 (0.4-4.7)	81 (10-210)
Qatar	2016	Female	60 to 64	20% (7.9-39%)	80% (61-92%)	1.2 (0.27-3.2)	1.7% (-0.14-4.8%)	3% (0.2-7.6%)	1.3 (0.093-3.8)	49 (-3.9-140)
Qatar	2016	Female	65 to 69	21% (8.8-37%)	79% (63-91%)	1.1 (0.25-2.7)	1.5% (-0.32-4.5%)	2.5% (-0.068-6.8%)	0.91 (-0.028-2.7)	26 (-5-79)
Qatar	2016	Female	70 to 74	20% (8-36%)	80% (64-92%)	0.88 (0.19-2.5)	1% (-0.65-4.4%)	1.5% (-0.69-5.9%)	0.68 (-0.34-2.7)	15 (-9.7-63)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	2016	Female	75 to 79	17% (6.1-37%)	83% (63-94%)	0.71 (0.13-2.3)	0.65% (-0.92-3.9%)	0.83% (-1.2-4.8%)	0.51 (-0.78-3)	9.1 (-14-54)
Qatar	2016	Female	80 to 84	16% (5.2-37%)	84% (63-95%)	0.63 (0.11-2.1)	0.67% (-0.8-3.8%)	0.8% (-0.99-4.9%)	0.43 (-0.56-2.6)	5.5 (-6.5-32)
Qatar	2016	Female	85 to 89	16% (5.3-38%)	84% (62-95%)	0.62 (0.11-2)	0.59% (-0.95-4%)	0.64% (-1.2-4.9%)	0.22 (-0.44-1.6)	2 (-3.4-13)
Qatar	2016	Female	90 to 94	19% (5.6-43%)	81% (57-94%)	0.69 (0.12-2.2)	0.59% (-1.2-3.7%)	0.61% (-1.5-4.4%)	0.13 (-0.32-0.92)	0.82 (-1.7-5.3)
Qatar	2016	Female	95 plus	23% (7.2-49%)	77% (51-93%)	0.81 (0.13-2.8)	0.63% (-1.5-4.6%)	0.61% (-1.9-5.4%)	0.047 (-0.14-0.41)	0.21 (-0.45-1.5)
Qatar	2016	Male	All Ages	12% (7-17%)	88% (83-93%)	0.86 (0.51-1.3)	2.6% (1.8-3.5%)	3.9% (2.5-5.6%)	100 (55-180)	7100 (4600-10000)
Qatar	2016	Male	15 to 19	6.2% (2.7-12%)	94% (88-97%)	0.24 (0.063-0.64)	1.3% (0.71-2.3%)	2.1% (0.82-4.7%)	0.8 (0.28-2)	110 (59-200)
Qatar	2016	Male	20 to 24	9.6% (3.8-19%)	90% (81-96%)	0.54 (0.16-1.3)	2.6% (1.5-4.3%)	3.7% (1.4-7.5%)	4.4 (1.6-9.8)	580 (310-1000)
Qatar	2016	Male	25 to 29	12% (4.6-23%)	88% (77-95%)	0.82 (0.28-1.6)	2.9% (1.8-4.6%)	4.9% (2.1-9.2%)	11 (4.3-21)	1200 (680-1900)
Qatar	2016	Male	30 to 34	14% (5-26%)	86% (74-95%)	1 (0.43-2)	2.9% (1.8-4.5%)	5.6% (2.7-9.6%)	13 (5.8-23)	1200 (740-1900)
Qatar	2016	Male	35 to 39	15% (5.4-28%)	85% (72-95%)	1.2 (0.47-2.3)	3.1% (1.8-4.8%)	6.2% (3-10%)	12 (5.1-23)	970 (540-1600)
Qatar	2016	Male	40 to 44	15% (6.6-28%)	85% (72-93%)	1.3 (0.46-2.6)	3.1% (1.7-4.9%)	6% (2.9-10%)	13 (5.3-25)	890 (470-1500)
Qatar	2016	Male	45 to 49	15% (5.9-27%)	85% (73-94%)	1.3 (0.39-2.8)	3% (1.5-5.1%)	5.5% (2.4-9.3%)	13 (5.1-26)	750 (360-1400)
Qatar	2016	Male	50 to 54	14% (6-28%)	86% (72-94%)	1.3 (0.39-3.1)	2.8% (1.3-5%)	4.6% (2-8.2%)	11 (4.4-23)	530 (240-1000)
Qatar	2016	Male	55 to 59	14% (5.8-26%)	86% (74-94%)	1.3 (0.35-3.2)	2.6% (1.1-4.8%)	4% (1.7-7.2%)	9.2 (3.6-20)	380 (160-770)
Qatar	2016	Male	60 to 64	13% (5.5-25%)	87% (75-95%)	1.1 (0.3-3)	2.3% (0.92-4.4%)	3.3% (1.4-6.2%)	6.3 (2.4-13)	220 (83-420)
Qatar	2016	Male	65 to 69	12% (5.6-24%)	88% (76-94%)	0.93 (0.24-2.2)	2.1% (0.8-4%)	2.9% (1.1-5.4%)	3.2 (1.1-6.6)	87 (32-180)
Qatar	2016	Male	70 to 74	11% (5.2-20%)	89% (80-95%)	0.69 (0.18-1.7)	1.6% (0.51-3.3%)	2% (0.7-4.2%)	2.4 (0.76-5.6)	54 (17-120)
Qatar	2016	Male	75 to 79	9.8% (4.1-18%)	90% (82-96%)	0.53 (0.12-1.5)	1.1% (0.23-2.6%)	1.3% (0.23-3.2%)	2.4 (0.41-6.1)	42 (8.7-100)
Qatar	2016	Male	80 to 84	8.8% (3.3-18%)	91% (82-97%)	0.44 (0.1-1.3)	1% (0.23-2.5%)	1.1% (0.19-2.8%)	1.1 (0.18-2.9)	14 (3.2-36)
Qatar	2016	Male	85 to 89	8.6% (3.4-17%)	91% (83-97%)	0.41 (0.093-1.1)	0.81% (0.1-2.2%)	0.82% (-0.0031-2.4%)	0.26 (-0.00088-0.78)	2.4 (0.31-6.8)
Qatar	2016	Male	90 to 94	9.7% (3.8-20%)	90% (80-96%)	0.43 (0.094-1.1)	0.68% (-0.067-2%)	0.64% (-0.22-2.2%)	0.11 (-0.042-0.4)	0.75 (-0.071-2.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Qatar	2016	Male	95 plus	12% (4.2-26%)	88% (74-96%)	0.5 (0.094-1.6)	0.7% (-0.17-2.5%)	0.58% (-0.43-2.6%)	0.041 (-0.032-0.19)	0.21 (-0.054-0.75)
Saudi Arabia	1990	Female	All Ages	11% (8.6-13%)	89% (87-91%)	0.062 (0.047-0.085)	0.084% (0.004-0.16%)	-0.13% (-0.38-0.09%)	-39 (-120-28)	2000 (88-3800)
Saudi Arabia	1990	Female	15 to 19	9.8% (5.4-16%)	90% (84-95%)	0.052 (0.024-0.1)	0.31% (0.21-0.44%)	0.3% (0.096-0.58%)	0.97 (0.3-1.9)	280 (170-410)
Saudi Arabia	1990	Female	20 to 24	13% (6.7-21%)	87% (79-93%)	0.085 (0.038-0.16)	0.63% (0.42-0.89%)	0.42% (0.091-0.91%)	1.6 (0.35-3.5)	570 (370-860)
Saudi Arabia	1990	Female	25 to 29	15% (7.8-23%)	85% (77-92%)	0.11 (0.054-0.19)	0.69% (0.49-0.93%)	0.53% (0.14-1.1%)	2.5 (0.61-5.1)	710 (460-1000)
Saudi Arabia	1990	Female	30 to 34	17% (9.2-27%)	83% (73-91%)	0.13 (0.071-0.23)	0.62% (0.42-0.9%)	0.57% (0.1-1.2%)	2.8 (0.47-5.9)	610 (380-910)
Saudi Arabia	1990	Female	35 to 39	19% (11-30%)	81% (70-89%)	0.16 (0.08-0.28)	0.49% (0.3-0.71%)	0.51% (0.051-1.1%)	2.4 (0.24-5.1)	380 (230-560)
Saudi Arabia	1990	Female	40 to 44	20% (12-31%)	80% (69-88%)	0.18 (0.084-0.32)	0.37% (0.15-0.62%)	0.49% (-0.026-1.1%)	2.1 (-0.12-5.1)	220 (89-370)
Saudi Arabia	1990	Female	45 to 49	21% (11-31%)	79% (69-89%)	0.19 (0.086-0.36)	0.21% (-0.025-0.47%)	0.26% (-0.25-0.94%)	1.2 (-1.1-4.4)	110 (-12-240)
Saudi Arabia	1990	Female	50 to 54	21% (13-33%)	79% (67-87%)	0.19 (0.086-0.35)	0.1% (-0.2-0.38%)	0.17% (-0.4-0.78%)	0.88 (-2.1-4.1)	45 (-87-170)
Saudi Arabia	1990	Female	55 to 59	23% (13-36%)	77% (64-87%)	0.19 (0.088-0.34)	-0.055% (-0.42-0.24%)	-0.036% (-0.71-0.53%)	-0.23 (-4.4-3.4)	-24 (-180-100)
Saudi Arabia	1990	Female	60 to 64	24% (14-38%)	76% (62-86%)	0.18 (0.082-0.35)	-0.26% (-0.78-0.15%)	-0.32% (-1.1-0.28%)	-4.4 (-15-3.9)	-160 (-480-89)
Saudi Arabia	1990	Female	65 to 69	25% (16-36%)	75% (64-84%)	0.16 (0.077-0.3)	-0.36% (-0.91-0.039%)	-0.44% (-1.2-0.12%)	-9 (-24-2.6)	-250 (-600-28)
Saudi Arabia	1990	Female	70 to 74	24% (16-35%)	76% (65-84%)	0.13 (0.057-0.25)	-0.34% (-0.92-0.045%)	-0.4% (-1.1-0.11%)	-9.9 (-28-2.7)	-210 (-560-28)
Saudi Arabia	1990	Female	75 to 79	22% (12-34%)	78% (66-88%)	0.11 (0.041-0.22)	-0.33% (-0.88-0.047%)	-0.38% (-1-0.072%)	-8.7 (-25-1.6)	-140 (-370-20)
Saudi Arabia	1990	Female	80 to 84	21% (11-34%)	79% (66-89%)	0.1 (0.039-0.2)	-0.32% (-0.87-0.057%)	-0.37% (-1-0.089%)	-7.1 (-20-1.8)	-80 (-220-14)
Saudi Arabia	1990	Female	85 to 89	22% (12-38%)	78% (62-88%)	0.11 (0.04-0.22)	-0.39% (-0.99--0.022%)	-0.46% (-1.2--0.017%)	-6.9 (-18--0.26)	-54 (-140--2.9)
Saudi Arabia	1990	Female	90 to 94	26% (15-41%)	74% (59-85%)	0.12 (0.048-0.26)	-0.5% (-1.1--0.077%)	-0.6% (-1.3--0.097%)	-5 (-11--0.8)	-26 (-58--4.1)
Saudi Arabia	1990	Female	95 plus	31% (16-49%)	69% (51-84%)	0.15 (0.05-0.33)	-0.61% (-1.4--0.083%)	-0.76% (-1.7--0.12%)	-2.5 (-5.5--0.41)	-8.3 (-19--1.1)
Saudi Arabia	1990	Male	All Ages	7.3% (5.7-9.2%)	93% (91-94%)	0.054 (0.039-0.075)	0.33% (0.23-0.45%)	0.28% (0.14-0.45%)	120 (57-190)	11000 (7400-15000)
Saudi Arabia	1990	Male	15 to 19	5.4% (3.1-8.7%)	95% (91-97%)	0.023 (0.01-0.045)	0.44% (0.28-0.63%)	0.24% (0.069-0.52%)	2.1 (0.59-4.5)	520 (330-760)
Saudi Arabia	1990	Male	20 to 24	8.3% (4.6-13%)	92% (87-95%)	0.05 (0.023-0.087)	0.94% (0.63-1.4%)	0.47% (0.13-1%)	6.8 (1.9-15)	1600 (1000-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	1990	Male	25 to 29	10% (5.7-16%)	90% (84-94%)	0.08 (0.041-0.14)	1.1% (0.75-1.5%)	0.69% (0.24-1.3%)	11 (3.8-20)	2000 (1300-2900)
Saudi Arabia	1990	Male	30 to 34	12% (6.6-20%)	88% (80-93%)	0.11 (0.056-0.18)	1% (0.73-1.5%)	0.8% (0.34-1.5%)	11 (4.7-20)	1700 (1100-2500)
Saudi Arabia	1990	Male	35 to 39	14% (7.9-22%)	86% (78-92%)	0.13 (0.064-0.22)	0.99% (0.7-1.4%)	0.92% (0.43-1.6%)	12 (5.6-22)	1400 (970-2000)
Saudi Arabia	1990	Male	40 to 44	14% (8.5-23%)	86% (77-92%)	0.14 (0.065-0.25)	0.88% (0.6-1.2%)	0.89% (0.45-1.5%)	10 (5-17)	920 (600-1300)
Saudi Arabia	1990	Male	45 to 49	14% (7.3-21%)	86% (79-93%)	0.14 (0.06-0.27)	0.76% (0.5-1.1%)	0.85% (0.42-1.4%)	8.9 (4.4-15)	620 (410-910)
Saudi Arabia	1990	Male	50 to 54	13% (7.4-21%)	87% (79-93%)	0.13 (0.061-0.26)	0.6% (0.36-0.85%)	0.69% (0.31-1.1%)	7.5 (3.2-12)	420 (250-610)
Saudi Arabia	1990	Male	55 to 59	13% (7.2-20%)	87% (80-93%)	0.13 (0.059-0.24)	0.49% (0.25-0.74%)	0.59% (0.22-0.97%)	7.3 (2.6-12)	320 (160-490)
Saudi Arabia	1990	Male	60 to 64	12% (7-20%)	88% (80-93%)	0.12 (0.051-0.24)	0.38% (0.1-0.64%)	0.45% (0.067-0.81%)	9.3 (1.3-16)	300 (81-510)
Saudi Arabia	1990	Male	65 to 69	12% (7.2-19%)	88% (81-93%)	0.1 (0.047-0.18)	0.31% (0.046-0.53%)	0.37% (0.018-0.66%)	8.9 (0.41-16)	230 (35-400)
Saudi Arabia	1990	Male	70 to 74	11% (7.1-17%)	89% (83-93%)	0.075 (0.033-0.15)	0.31% (0.08-0.51%)	0.38% (0.069-0.64%)	9.2 (1.6-15)	180 (46-290)
Saudi Arabia	1990	Male	75 to 79	9.6% (5.6-15%)	90% (85-94%)	0.056 (0.022-0.12)	0.26% (0.05-0.41%)	0.31% (0.038-0.5%)	6.1 (0.79-9.8)	91 (18-140)
Saudi Arabia	1990	Male	80 to 84	8.5% (4.9-13%)	91% (87-95%)	0.047 (0.02-0.089)	0.28% (0.1-0.41%)	0.32% (0.11-0.49%)	4.6 (1.6-7)	50 (18-75)
Saudi Arabia	1990	Male	85 to 89	8.5% (4.8-14%)	92% (86-95%)	0.045 (0.019-0.088)	0.18% (0.022-0.31%)	0.2% (-0.00065-0.35%)	2.1 (-0.0068-3.7)	18 (2-29)
Saudi Arabia	1990	Male	90 to 94	9.7% (5.7-16%)	90% (84-94%)	0.05 (0.02-0.1)	0.094% (-0.076-0.21%)	0.086% (-0.12-0.24%)	0.46 (-0.62-1.3)	3.1 (-2.5-7.2)
Saudi Arabia	1990	Male	95 plus	12% (6-21%)	88% (79-94%)	0.059 (0.018-0.13)	0.059% (-0.15-0.2%)	0.016% (-0.24-0.18%)	0.028 (-0.42-0.32)	0.44 (-1-1.5)
Saudi Arabia	1995	Female	All Ages	11% (9.2-14%)	89% (86-91%)	0.049 (0.036-0.067)	0.092% (-0.0025-0.17%)	-0.15% (-0.44-0.095%)	-48 (-150-33)	2100 (-53-4000)
Saudi Arabia	1995	Female	15 to 19	11% (5.9-17%)	89% (83-94%)	0.042 (0.019-0.086)	0.31% (0.21-0.45%)	0.27% (0.1-0.49%)	1 (0.37-2)	350 (220-510)
Saudi Arabia	1995	Female	20 to 24	14% (7.3-23%)	86% (77-93%)	0.069 (0.031-0.13)	0.63% (0.43-0.89%)	0.37% (0.078-0.78%)	1.3 (0.28-2.8)	600 (370-910)
Saudi Arabia	1995	Female	25 to 29	16% (8.5-25%)	84% (75-91%)	0.087 (0.041-0.15)	0.68% (0.48-0.93%)	0.45% (0.12-0.91%)	1.8 (0.46-3.7)	640 (420-940)
Saudi Arabia	1995	Female	30 to 34	18% (9.7-28%)	82% (72-90%)	0.1 (0.054-0.17)	0.6% (0.4-0.84%)	0.46% (0.086-0.92%)	2.4 (0.44-4.9)	660 (410-990)
Saudi Arabia	1995	Female	35 to 39	20% (11-31%)	80% (69-89%)	0.12 (0.062-0.21)	0.46% (0.31-0.64%)	0.4% (0.043-0.83%)	2.4 (0.26-4.8)	480 (290-670)
Saudi Arabia	1995	Female	40 to 44	20% (12-31%)	80% (69-88%)	0.13 (0.065-0.23)	0.34% (0.17-0.52%)	0.38% (-0.0023-0.83%)	2.1 (-0.013-4.4)	270 (130-430)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	1995	Female	45 to 49	20% (11-31%)	80% (69-89%)	0.14 (0.063-0.26)	0.19% (-0.0026-0.36%)	0.19% (-0.21-0.57%)	0.9 (-0.98-2.8)	100 (-1.3-200)
Saudi Arabia	1995	Female	50 to 54	22% (13-34%)	78% (66-87%)	0.14 (0.065-0.27)	0.076% (-0.2-0.28%)	0.11% (-0.38-0.51%)	0.7 (-2.4-3.3)	42 (-110-150)
Saudi Arabia	1995	Female	55 to 59	24% (14-36%)	76% (64-86%)	0.15 (0.067-0.27)	-0.072% (-0.41-0.18%)	-0.058% (-0.64-0.37%)	-0.41 (-4.5-2.7)	-36 (-210-88)
Saudi Arabia	1995	Female	60 to 64	26% (15-39%)	74% (61-85%)	0.14 (0.063-0.27)	-0.28% (-0.82-0.082%)	-0.34% (-1.1-0.18%)	-5.7 (-18-2.9)	-190 (-560-59)
Saudi Arabia	1995	Female	65 to 69	26% (17-38%)	74% (62-83%)	0.13 (0.056-0.22)	-0.35% (-0.91-0.065%)	-0.41% (-1.1-0.12%)	-10 (-28-3)	-280 (-710-53)
Saudi Arabia	1995	Female	70 to 74	25% (16-36%)	75% (64-84%)	0.096 (0.043-0.19)	-0.3% (-0.85-0.08%)	-0.34% (-1-0.14%)	-12 (-37-4.9)	-260 (-750-71)
Saudi Arabia	1995	Female	75 to 79	22% (13-33%)	78% (67-87%)	0.075 (0.03-0.15)	-0.25% (-0.76-0.082%)	-0.28% (-0.9-0.11%)	-9.9 (-31-4)	-160 (-470-51)
Saudi Arabia	1995	Female	80 to 84	20% (11-31%)	80% (69-89%)	0.07 (0.028-0.14)	-0.24% (-0.71-0.082%)	-0.27% (-0.83-0.1%)	-8 (-24-3.1)	-88 (-250-28)
Saudi Arabia	1995	Female	85 to 89	21% (12-35%)	79% (65-88%)	0.073 (0.029-0.15)	-0.3% (-0.79-0.016%)	-0.34% (-0.91-0.029%)	-6.6 (-17-0.53)	-51 (-130-2.9)
Saudi Arabia	1995	Female	90 to 94	25% (14-39%)	75% (61-86%)	0.086 (0.035-0.18)	-0.38% (-0.91--0.033%)	-0.46% (-1.1--0.041%)	-5 (-12--0.45)	-26 (-61--2.3)
Saudi Arabia	1995	Female	95 plus	30% (16-48%)	70% (52-84%)	0.11 (0.034-0.23)	-0.5% (-1.2--0.044%)	-0.61% (-1.5--0.075%)	-2.6 (-6.3--0.32)	-8.8 (-21--0.8)
Saudi Arabia	1995	Male	All Ages	7.6% (5.9-9.6%)	92% (90-94%)	0.041 (0.03-0.057)	0.39% (0.28-0.5%)	0.32% (0.18-0.49%)	130 (74-200)	11000 (8100-15000)
Saudi Arabia	1995	Male	15 to 19	5.5% (3.2-8.9%)	94% (91-97%)	0.017 (0.0077-0.034)	0.44% (0.3-0.64%)	0.23% (0.069-0.46%)	2 (0.59-4)	550 (360-830)
Saudi Arabia	1995	Male	20 to 24	8.4% (4.7-13%)	92% (87-95%)	0.037 (0.018-0.069)	0.97% (0.65-1.4%)	0.46% (0.11-0.99%)	4.7 (1.2-10)	1200 (790-1900)
Saudi Arabia	1995	Male	25 to 29	11% (5.7-17%)	89% (83-94%)	0.058 (0.029-0.1)	1.1% (0.78-1.5%)	0.64% (0.21-1.3%)	8.7 (2.8-17)	1900 (1300-2800)
Saudi Arabia	1995	Male	30 to 34	12% (6.6-20%)	88% (80-93%)	0.076 (0.039-0.13)	1% (0.74-1.4%)	0.73% (0.31-1.3%)	10 (4.3-19)	2000 (1300-2900)
Saudi Arabia	1995	Male	35 to 39	14% (7.7-22%)	86% (78-92%)	0.088 (0.047-0.15)	0.97% (0.72-1.3%)	0.84% (0.41-1.5%)	12 (5.9-22)	1700 (1200-2400)
Saudi Arabia	1995	Male	40 to 44	14% (8.3-23%)	86% (77-92%)	0.097 (0.047-0.17)	0.84% (0.58-1.1%)	0.81% (0.43-1.3%)	12 (6.3-21)	1300 (850-1800)
Saudi Arabia	1995	Male	45 to 49	14% (7.4-22%)	86% (78-93%)	0.1 (0.045-0.19)	0.73% (0.5-0.99%)	0.8% (0.44-1.3%)	9.5 (5.1-15)	710 (470-990)
Saudi Arabia	1995	Male	50 to 54	13% (7.6-21%)	87% (79-92%)	0.098 (0.045-0.19)	0.58% (0.37-0.78%)	0.66% (0.32-0.99%)	8.8 (4.4-14)	500 (310-700)
Saudi Arabia	1995	Male	55 to 59	13% (7.4-20%)	87% (80-93%)	0.094 (0.043-0.18)	0.47% (0.27-0.67%)	0.57% (0.25-0.88%)	8.3 (3.6-13)	370 (210-530)
Saudi Arabia	1995	Male	60 to 64	13% (7.1-20%)	87% (80-93%)	0.086 (0.037-0.17)	0.37% (0.13-0.58%)	0.43% (0.11-0.72%)	10 (2.7-17)	340 (120-530)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	1995	Male	65 to 69	12% (7.2-19%)	88% (81-93%)	0.072 (0.033-0.13)	0.32% (0.092-0.51%)	0.38% (0.078-0.63%)	11 (2.3-18)	280 (82-450)
Saudi Arabia	1995	Male	70 to 74	11% (7-17%)	89% (83-93%)	0.053 (0.024-0.1)	0.34% (0.14-0.51%)	0.42% (0.15-0.63%)	13 (5-20)	250 (100-380)
Saudi Arabia	1995	Male	75 to 79	9.5% (5.5-15%)	90% (85-94%)	0.039 (0.016-0.081)	0.29% (0.11-0.42%)	0.35% (0.12-0.52%)	9.4 (3.2-14)	140 (54-200)
Saudi Arabia	1995	Male	80 to 84	8.5% (4.9-13%)	91% (87-95%)	0.033 (0.014-0.062)	0.3% (0.15-0.42%)	0.35% (0.17-0.5%)	6.7 (3.2-9.5)	72 (36-100)
Saudi Arabia	1995	Male	85 to 89	8.5% (4.8-14%)	91% (86-95%)	0.032 (0.014-0.064)	0.21% (0.069-0.31%)	0.23% (0.066-0.36%)	2.9 (0.83-4.5)	23 (7.8-35)
Saudi Arabia	1995	Male	90 to 94	9.8% (5.7-16%)	90% (84-94%)	0.035 (0.014-0.076)	0.12% (-0.033-0.22%)	0.12% (-0.055-0.24%)	0.75 (-0.35-1.6)	4.7 (-1.3-8.9)
Saudi Arabia	1995	Male	95 plus	12% (6-21%)	88% (79-94%)	0.042 (0.013-0.094)	0.084% (-0.098-0.2%)	0.05% (-0.17-0.19%)	0.1 (-0.36-0.4)	0.73 (-0.85-1.8)
Saudi Arabia	2000	Female	All Ages	12% (9.8-15%)	88% (85-90%)	0.046 (0.034-0.062)	0.12% (0.0092-0.22%)	-0.14% (-0.46-0.12%)	-54 (-180-46)	2800 (210-5200)
Saudi Arabia	2000	Female	15 to 19	10% (5.8-17%)	90% (83-94%)	0.034 (0.015-0.066)	0.31% (0.21-0.44%)	0.25% (0.11-0.44%)	1 (0.47-1.9)	380 (240-560)
Saudi Arabia	2000	Female	20 to 24	14% (7.2-22%)	86% (78-93%)	0.056 (0.026-0.11)	0.64% (0.43-0.91%)	0.33% (0.098-0.66%)	1.5 (0.43-3)	800 (490-1200)
Saudi Arabia	2000	Female	25 to 29	16% (8.6-25%)	84% (75-91%)	0.072 (0.035-0.13)	0.69% (0.5-0.93%)	0.41% (0.11-0.8%)	2.1 (0.55-4.2)	900 (580-1300)
Saudi Arabia	2000	Female	30 to 34	18% (10-29%)	82% (71-90%)	0.088 (0.046-0.15)	0.6% (0.41-0.83%)	0.43% (0.091-0.82%)	2.5 (0.53-4.8)	760 (480-1100)
Saudi Arabia	2000	Female	35 to 39	20% (12-31%)	80% (69-88%)	0.1 (0.053-0.17)	0.47% (0.32-0.64%)	0.39% (0.089-0.74%)	2.6 (0.57-4.9)	580 (360-820)
Saudi Arabia	2000	Female	40 to 44	21% (13-32%)	79% (68-87%)	0.11 (0.057-0.19)	0.34% (0.18-0.51%)	0.38% (0.045-0.75%)	2.3 (0.26-4.6)	340 (170-520)
Saudi Arabia	2000	Female	45 to 49	21% (12-32%)	79% (68-88%)	0.12 (0.055-0.21)	0.19% (0.0081-0.35%)	0.2% (-0.15-0.52%)	1.2 (-0.92-3.2)	150 (6.7-280)
Saudi Arabia	2000	Female	50 to 54	22% (13-34%)	78% (66-87%)	0.12 (0.056-0.22)	0.077% (-0.19-0.27%)	0.12% (-0.33-0.44%)	0.92 (-2.5-3.5)	53 (-130-180)
Saudi Arabia	2000	Female	55 to 59	24% (14-37%)	76% (63-86%)	0.12 (0.056-0.22)	-0.056% (-0.38-0.17%)	-0.022% (-0.57-0.36%)	-0.17 (-4.5-3)	-34 (-240-110)
Saudi Arabia	2000	Female	60 to 64	25% (15-39%)	75% (61-85%)	0.12 (0.051-0.22)	-0.24% (-0.79-0.13%)	-0.27% (-0.99-0.22%)	-6.3 (-23-4.8)	-210 (-730-110)
Saudi Arabia	2000	Female	65 to 69	26% (17-37%)	74% (63-83%)	0.1 (0.046-0.18)	-0.29% (-0.79-0.1%)	-0.32% (-0.95-0.16%)	-10 (-30-5)	-280 (-750-98)
Saudi Arabia	2000	Female	70 to 74	24% (16-35%)	76% (65-84%)	0.078 (0.034-0.15)	-0.24% (-0.78-0.12%)	-0.26% (-0.89-0.17%)	-13 (-43-8)	-260 (-850-130)
Saudi Arabia	2000	Female	75 to 79	22% (13-33%)	78% (67-87%)	0.062 (0.024-0.12)	-0.2% (-0.7-0.12%)	-0.22% (-0.78-0.15%)	-11 (-40-7.9)	-180 (-610-100)
Saudi Arabia	2000	Female	80 to 84	21% (11-33%)	79% (67-89%)	0.058 (0.022-0.11)	-0.2% (-0.67-0.11%)	-0.22% (-0.73-0.13%)	-11 (-36-6.7)	-120 (-380-69)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	2000	Female	85 to 89	22% (12-36%)	78% (64-88%)	0.061 (0.025-0.12)	-0.26% (-0.72-0.038%)	-0.3% (-0.81-0.043%)	-8.3 (-23-1.2)	-64 (-180-9.4)
Saudi Arabia	2000	Female	90 to 94	25% (15-39%)	75% (61-85%)	0.072 (0.029-0.15)	-0.34% (-0.85--0.014%)	-0.41% (-0.99--0.023%)	-5.5 (-13--0.31)	-29 (-70--1.1)
Saudi Arabia	2000	Female	95 plus	31% (16-48%)	69% (52-84%)	0.089 (0.029-0.19)	-0.45% (-1.1--0.024%)	-0.55% (-1.4--0.042%)	-3.1 (-7.7--0.25)	-11 (-27--0.57)
Saudi Arabia	2000	Male	All Ages	7.8% (6.1-9.8%)	92% (90-94%)	0.035 (0.025-0.048)	0.45% (0.33-0.57%)	0.36% (0.22-0.52%)	150 (95-220)	13000 (9400-17000)
Saudi Arabia	2000	Male	15 to 19	5.4% (3.1-8.7%)	95% (91-97%)	0.014 (0.0059-0.027)	0.42% (0.29-0.59%)	0.2% (0.068-0.39%)	2 (0.7-4)	580 (380-850)
Saudi Arabia	2000	Male	20 to 24	8.2% (4.5-13%)	92% (87-95%)	0.029 (0.014-0.053)	0.93% (0.62-1.3%)	0.38% (0.1-0.81%)	5.1 (1.4-11)	1500 (960-2200)
Saudi Arabia	2000	Male	25 to 29	10% (5.6-16%)	90% (84-94%)	0.046 (0.023-0.08)	1.1% (0.81-1.5%)	0.57% (0.2-1.1%)	8.4 (3-17)	2300 (1500-3200)
Saudi Arabia	2000	Male	30 to 34	12% (6.5-20%)	88% (80-94%)	0.061 (0.032-0.1)	1% (0.76-1.4%)	0.67% (0.29-1.2%)	9.4 (4.1-17)	2100 (1500-3100)
Saudi Arabia	2000	Male	35 to 39	14% (7.9-22%)	86% (78-92%)	0.072 (0.038-0.13)	0.96% (0.71-1.3%)	0.8% (0.41-1.4%)	12 (5.9-20)	1800 (1200-2400)
Saudi Arabia	2000	Male	40 to 44	14% (8.3-23%)	86% (77-92%)	0.079 (0.039-0.14)	0.83% (0.6-1.1%)	0.78% (0.42-1.2%)	13 (6.8-21)	1400 (980-2000)
Saudi Arabia	2000	Male	45 to 49	14% (7.2-21%)	86% (79-93%)	0.08 (0.036-0.15)	0.72% (0.52-0.94%)	0.79% (0.46-1.2%)	13 (7.5-19)	1000 (720-1400)
Saudi Arabia	2000	Male	50 to 54	13% (7.4-21%)	87% (79-93%)	0.077 (0.035-0.15)	0.58% (0.38-0.78%)	0.66% (0.37-0.96%)	9.4 (5.1-14)	540 (350-730)
Saudi Arabia	2000	Male	55 to 59	12% (7.2-19%)	88% (81-93%)	0.074 (0.034-0.14)	0.48% (0.29-0.65%)	0.58% (0.28-0.86%)	7.8 (3.8-12)	350 (210-490)
Saudi Arabia	2000	Male	60 to 64	12% (7-20%)	88% (80-93%)	0.067 (0.028-0.13)	0.39% (0.16-0.58%)	0.45% (0.15-0.7%)	13 (4.4-21)	420 (170-630)
Saudi Arabia	2000	Male	65 to 69	12% (7-18%)	88% (82-93%)	0.056 (0.025-0.1)	0.35% (0.14-0.52%)	0.41% (0.15-0.63%)	14 (5-22)	360 (150-540)
Saudi Arabia	2000	Male	70 to 74	11% (6.8-16%)	89% (84-93%)	0.041 (0.019-0.08)	0.37% (0.19-0.52%)	0.45% (0.22-0.63%)	17 (8.4-24)	330 (160-460)
Saudi Arabia	2000	Male	75 to 79	9.1% (5.3-15%)	91% (85-95%)	0.03 (0.012-0.06)	0.32% (0.18-0.43%)	0.39% (0.2-0.53%)	14 (6.9-19)	200 (110-270)
Saudi Arabia	2000	Male	80 to 84	8.1% (4.7-13%)	92% (87-95%)	0.025 (0.011-0.047)	0.33% (0.2-0.44%)	0.39% (0.23-0.51%)	10 (6.1-14)	110 (67-150)
Saudi Arabia	2000	Male	85 to 89	8.1% (4.5-13%)	92% (87-95%)	0.025 (0.011-0.05)	0.23% (0.11-0.32%)	0.26% (0.12-0.36%)	4.4 (2-6.1)	35 (17-47)
Saudi Arabia	2000	Male	90 to 94	9.5% (5.4-15%)	91% (85-95%)	0.027 (0.011-0.057)	0.14% (0.012-0.23%)	0.15% (-0.0056-0.25%)	1.1 (-0.043-1.9)	6.7 (0.56-11)
Saudi Arabia	2000	Male	95 plus	12% (5.9-21%)	88% (79-94%)	0.033 (0.01-0.071)	0.11% (-0.055-0.21%)	0.083% (-0.12-0.2%)	0.21 (-0.29-0.52)	1.2 (-0.55-2.2)
Saudi Arabia	2005	Female	All Ages	12% (10-15%)	88% (85-90%)	0.049 (0.036-0.067)	0.15% (0.024-0.25%)	-0.15% (-0.48-0.13%)	-59 (-190-50)	3400 (550-6200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	2005	Female	15 to 19	10% (5.7-17%)	90% (83-94%)	0.035 (0.015-0.068)	0.32% (0.22-0.44%)	0.27% (0.13-0.47%)	1.2 (0.56-2.1)	440 (280-650)
Saudi Arabia	2005	Female	20 to 24	14% (7.1-23%)	86% (77-93%)	0.057 (0.026-0.11)	0.65% (0.44-0.9%)	0.36% (0.13-0.7%)	1.7 (0.62-3.4)	900 (570-1300)
Saudi Arabia	2005	Female	25 to 29	16% (8.5-25%)	84% (75-91%)	0.073 (0.035-0.13)	0.7% (0.51-0.94%)	0.42% (0.15-0.8%)	2.6 (0.94-5)	1100 (720-1600)
Saudi Arabia	2005	Female	30 to 34	18% (9.9-28%)	82% (72-90%)	0.088 (0.046-0.15)	0.63% (0.43-0.86%)	0.46% (0.15-0.85%)	2.7 (0.91-5.1)	950 (610-1400)
Saudi Arabia	2005	Female	35 to 39	20% (12-30%)	80% (70-88%)	0.1 (0.053-0.17)	0.48% (0.33-0.66%)	0.42% (0.13-0.76%)	2.6 (0.9-4.9)	600 (370-860)
Saudi Arabia	2005	Female	40 to 44	21% (12-32%)	79% (68-88%)	0.11 (0.058-0.19)	0.35% (0.2-0.52%)	0.41% (0.093-0.76%)	2.9 (0.67-5.4)	430 (240-660)
Saudi Arabia	2005	Female	45 to 49	21% (11-32%)	79% (68-89%)	0.12 (0.055-0.21)	0.2% (0.032-0.35%)	0.24% (-0.094-0.55%)	1.7 (-0.69-3.8)	200 (29-350)
Saudi Arabia	2005	Female	50 to 54	22% (13-34%)	78% (66-87%)	0.12 (0.057-0.22)	0.083% (-0.18-0.27%)	0.15% (-0.28-0.48%)	1.4 (-2.7-4.4)	72 (-140-230)
Saudi Arabia	2005	Female	55 to 59	24% (14-36%)	76% (64-86%)	0.12 (0.056-0.23)	-0.055% (-0.37-0.17%)	0.0026% (-0.52-0.37%)	0.025 (-4.4-3.3)	-38 (-260-120)
Saudi Arabia	2005	Female	60 to 64	25% (15-39%)	75% (61-85%)	0.12 (0.05-0.23)	-0.23% (-0.77-0.15%)	-0.26% (-0.98-0.24%)	-7.2 (-27-6.7)	-250 (-840-150)
Saudi Arabia	2005	Female	65 to 69	26% (17-37%)	74% (63-83%)	0.1 (0.046-0.19)	-0.29% (-0.79-0.11%)	-0.32% (-0.93-0.17%)	-11 (-32-5.7)	-300 (-820-120)
Saudi Arabia	2005	Female	70 to 74	25% (16-36%)	75% (64-84%)	0.081 (0.036-0.16)	-0.24% (-0.78-0.13%)	-0.26% (-0.92-0.17%)	-12 (-42-8)	-250 (-840-130)
Saudi Arabia	2005	Female	75 to 79	22% (13-34%)	78% (66-87%)	0.065 (0.025-0.13)	-0.2% (-0.72-0.12%)	-0.22% (-0.8-0.16%)	-11 (-41-8.2)	-180 (-610-110)
Saudi Arabia	2005	Female	80 to 84	21% (11-33%)	79% (67-89%)	0.061 (0.024-0.11)	-0.21% (-0.66-0.12%)	-0.23% (-0.73-0.14%)	-13 (-41-7.8)	-140 (-440-81)
Saudi Arabia	2005	Female	85 to 89	22% (12-37%)	78% (63-88%)	0.064 (0.026-0.13)	-0.27% (-0.74-0.034%)	-0.3% (-0.86-0.039%)	-11 (-32-1.4)	-88 (-250-12)
Saudi Arabia	2005	Female	90 to 94	26% (15-40%)	74% (60-85%)	0.075 (0.03-0.16)	-0.35% (-0.9-0.015%)	-0.41% (-1--0.021%)	-7.4 (-19--0.39)	-39 (-97--1.8)
Saudi Arabia	2005	Female	95 plus	31% (16-48%)	69% (52-84%)	0.093 (0.03-0.2)	-0.46% (-1.1--0.027%)	-0.56% (-1.4--0.045%)	-3.8 (-9.6--0.31)	-13 (-33--0.72)
Saudi Arabia	2005	Male	All Ages	8% (6.2-10%)	92% (90-94%)	0.036 (0.026-0.049)	0.53% (0.4-0.67%)	0.41% (0.25-0.57%)	190 (120-270)	16000 (12000-21000)
Saudi Arabia	2005	Male	15 to 19	5.2% (3-8.4%)	95% (92-97%)	0.013 (0.0056-0.026)	0.43% (0.3-0.59%)	0.2% (0.078-0.39%)	2.6 (1-5)	760 (490-1100)
Saudi Arabia	2005	Male	20 to 24	8% (4.4-13%)	92% (87-96%)	0.028 (0.013-0.052)	0.89% (0.59-1.3%)	0.37% (0.11-0.81%)	6.1 (1.7-13)	1700 (1100-2500)
Saudi Arabia	2005	Male	25 to 29	10% (5.4-16%)	90% (84-95%)	0.044 (0.021-0.078)	1.1% (0.76-1.5%)	0.56% (0.2-1.1%)	10 (3.6-20)	2400 (1600-3400)
Saudi Arabia	2005	Male	30 to 34	12% (6.3-19%)	88% (81-94%)	0.058 (0.031-0.1)	1.1% (0.76-1.4%)	0.66% (0.3-1.2%)	12 (5.6-22)	2900 (2000-4200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	2005	Male	35 to 39	13% (7.5-21%)	87% (79-92%)	0.069 (0.036-0.12)	0.97% (0.73-1.3%)	0.8% (0.43-1.3%)	16 (8.6-27)	2500 (1800-3500)
Saudi Arabia	2005	Male	40 to 44	14% (8.1-22%)	86% (78-92%)	0.075 (0.038-0.13)	0.84% (0.61-1.1%)	0.79% (0.45-1.2%)	15 (8.4-24)	1700 (1200-2400)
Saudi Arabia	2005	Male	45 to 49	13% (7.1-21%)	87% (79-93%)	0.077 (0.034-0.14)	0.73% (0.53-0.94%)	0.8% (0.47-1.2%)	16 (9.4-24)	1200 (890-1700)
Saudi Arabia	2005	Male	50 to 54	13% (7.2-20%)	87% (80-93%)	0.074 (0.034-0.14)	0.58% (0.39-0.76%)	0.67% (0.36-0.96%)	14 (7.7-21)	850 (560-1100)
Saudi Arabia	2005	Male	55 to 59	12% (7-19%)	88% (81-93%)	0.071 (0.032-0.14)	0.47% (0.29-0.65%)	0.57% (0.28-0.85%)	8.9 (4.2-13)	410 (240-570)
Saudi Arabia	2005	Male	60 to 64	12% (6.8-19%)	88% (81-93%)	0.064 (0.026-0.12)	0.4% (0.16-0.59%)	0.45% (0.15-0.7%)	14 (4.6-21)	440 (180-650)
Saudi Arabia	2005	Male	65 to 69	12% (6.8-18%)	88% (82-93%)	0.053 (0.024-0.097)	0.36% (0.15-0.53%)	0.43% (0.17-0.65%)	16 (6.2-24)	400 (170-590)
Saudi Arabia	2005	Male	70 to 74	10% (6.6-16%)	90% (84-93%)	0.039 (0.018-0.076)	0.38% (0.2-0.53%)	0.47% (0.24-0.64%)	20 (10-28)	380 (210-520)
Saudi Arabia	2005	Male	75 to 79	8.9% (5.2-14%)	91% (86-95%)	0.029 (0.011-0.057)	0.35% (0.2-0.46%)	0.42% (0.24-0.56%)	17 (9.5-22)	240 (140-320)
Saudi Arabia	2005	Male	80 to 84	7.9% (4.6-13%)	92% (87-95%)	0.024 (0.01-0.045)	0.36% (0.23-0.46%)	0.42% (0.27-0.54%)	14 (9.1-19)	150 (99-200)
Saudi Arabia	2005	Male	85 to 89	7.9% (4.4-13%)	92% (87-96%)	0.024 (0.01-0.047)	0.25% (0.12-0.34%)	0.28% (0.13-0.39%)	6.9 (3.2-9.4)	54 (26-73)
Saudi Arabia	2005	Male	90 to 94	9.1% (5.2-15%)	91% (85-95%)	0.026 (0.01-0.054)	0.16% (0.036-0.24%)	0.17% (0.021-0.27%)	1.8 (0.23-2.9)	10 (2.3-16)
Saudi Arabia	2005	Male	95 plus	11% (5.6-20%)	89% (80-94%)	0.031 (0.0098-0.07)	0.13% (-0.038-0.22%)	0.1% (-0.093-0.22%)	0.33 (-0.3-0.71)	1.7 (-0.5-3)
Saudi Arabia	2010	Female	All Ages	13% (11-16%)	87% (84-89%)	0.056 (0.04-0.075)	0.19% (0.066-0.3%)	-0.13% (-0.48-0.16%)	-50 (-190-63)	4500 (1500-7400)
Saudi Arabia	2010	Female	15 to 19	11% (5.7-17%)	89% (83-94%)	0.037 (0.016-0.074)	0.34% (0.23-0.47%)	0.32% (0.17-0.55%)	1.2 (0.61-2)	470 (290-700)
Saudi Arabia	2010	Female	20 to 24	14% (7-22%)	86% (78-93%)	0.06 (0.026-0.12)	0.69% (0.46-0.97%)	0.4% (0.17-0.77%)	1.9 (0.79-3.7)	1000 (670-1600)
Saudi Arabia	2010	Female	25 to 29	15% (8.3-25%)	85% (75-92%)	0.075 (0.034-0.13)	0.74% (0.54-1%)	0.48% (0.2-0.9%)	2.8 (1.2-5.2)	1200 (800-1700)
Saudi Arabia	2010	Female	30 to 34	18% (9.5-28%)	82% (72-90%)	0.091 (0.045-0.15)	0.66% (0.47-0.9%)	0.52% (0.24-0.91%)	3.2 (1.5-5.6)	1200 (790-1800)
Saudi Arabia	2010	Female	35 to 39	20% (11-30%)	80% (70-89%)	0.11 (0.054-0.18)	0.5% (0.35-0.67%)	0.47% (0.2-0.84%)	3.2 (1.3-5.6)	860 (540-1200)
Saudi Arabia	2010	Female	40 to 44	21% (13-32%)	79% (68-87%)	0.12 (0.06-0.2)	0.36% (0.2-0.52%)	0.44% (0.14-0.79%)	2.9 (0.87-5.1)	440 (230-670)
Saudi Arabia	2010	Female	45 to 49	21% (12-32%)	79% (68-88%)	0.12 (0.059-0.22)	0.21% (0.037-0.36%)	0.28% (-0.04-0.61%)	2.1 (-0.31-4.7)	250 (43-430)
Saudi Arabia	2010	Female	50 to 54	22% (13-34%)	78% (66-87%)	0.13 (0.057-0.23)	0.1% (-0.14-0.28%)	0.2% (-0.21-0.52%)	2 (-2.1-5.2)	100 (-130-290)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	2010	Female	55 to 59	24% (14-36%)	76% (64-86%)	0.13 (0.058-0.24)	-0.032% (-0.34-0.18%)	0.051% (-0.46-0.41%)	0.5 (-4.5-4.2)	-27 (-280-150)
Saudi Arabia	2010	Female	60 to 64	25% (15-39%)	75% (61-85%)	0.12 (0.051-0.24)	-0.2% (-0.72-0.15%)	-0.22% (-0.9-0.25%)	-6.3 (-25-7)	-230 (-820-180)
Saudi Arabia	2010	Female	65 to 69	26% (16-37%)	74% (63-84%)	0.11 (0.048-0.19)	-0.25% (-0.74-0.13%)	-0.27% (-0.89-0.22%)	-10 (-33-8)	-290 (-880-160)
Saudi Arabia	2010	Female	70 to 74	24% (15-35%)	76% (65-85%)	0.082 (0.036-0.16)	-0.2% (-0.71-0.14%)	-0.21% (-0.85-0.2%)	-9.7 (-40-9.4)	-220 (-810-160)
Saudi Arabia	2010	Female	75 to 79	21% (12-32%)	79% (68-88%)	0.064 (0.025-0.12)	-0.16% (-0.62-0.14%)	-0.17% (-0.71-0.2%)	-8 (-34-9.5)	-130 (-520-120)
Saudi Arabia	2010	Female	80 to 84	20% (11-32%)	80% (68-89%)	0.06 (0.023-0.11)	-0.18% (-0.62-0.14%)	-0.19% (-0.71-0.16%)	-11 (-37-8.8)	-120 (-410-91)
Saudi Arabia	2010	Female	85 to 89	22% (12-36%)	78% (64-88%)	0.065 (0.026-0.13)	-0.24% (-0.71-0.047%)	-0.27% (-0.82-0.055%)	-11 (-34-2.4)	-89 (-270-17)
Saudi Arabia	2010	Female	90 to 94	25% (14-40%)	75% (60-86%)	0.076 (0.029-0.16)	-0.33% (-0.85--0.0066%)	-0.39% (-1--0.014%)	-9.2 (-24--0.34)	-48 (-120--0.95)
Saudi Arabia	2010	Female	95 plus	30% (16-48%)	70% (52-84%)	0.094 (0.031-0.21)	-0.43% (-1.1--0.011%)	-0.54% (-1.3--0.033%)	-4.8 (-12--0.3)	-16 (-40--0.38)
Saudi Arabia	2010	Male	All Ages	8.7% (6.8-11%)	91% (89-93%)	0.041 (0.029-0.057)	0.61% (0.47-0.76%)	0.47% (0.29-0.65%)	230 (140-330)	20000 (14000-26000)
Saudi Arabia	2010	Male	15 to 19	5.4% (3.1-8.7%)	95% (91-97%)	0.014 (0.0059-0.028)	0.45% (0.3-0.62%)	0.23% (0.094-0.43%)	2.9 (1.2-5.7)	810 (520-1200)
Saudi Arabia	2010	Male	20 to 24	8.2% (4.6-13%)	92% (87-95%)	0.03 (0.014-0.057)	0.92% (0.62-1.3%)	0.4% (0.12-0.87%)	7.9 (2.4-17)	2100 (1300-3200)
Saudi Arabia	2010	Male	25 to 29	10% (5.5-16%)	90% (84-95%)	0.047 (0.022-0.083)	1.1% (0.75-1.4%)	0.59% (0.21-1.1%)	13 (4.7-26)	2700 (1800-3900)
Saudi Arabia	2010	Male	30 to 34	12% (6.4-20%)	88% (80-94%)	0.061 (0.031-0.11)	1.1% (0.77-1.4%)	0.69% (0.31-1.3%)	14 (6.5-26)	3000 (2100-4400)
Saudi Arabia	2010	Male	35 to 39	14% (7.7-21%)	86% (79-92%)	0.072 (0.037-0.12)	1% (0.76-1.3%)	0.87% (0.48-1.4%)	19 (11-31)	3100 (2200-4300)
Saudi Arabia	2010	Male	40 to 44	14% (8.3-23%)	86% (77-92%)	0.078 (0.04-0.14)	0.86% (0.64-1.1%)	0.84% (0.48-1.3%)	20 (12-31)	2400 (1700-3300)
Saudi Arabia	2010	Male	45 to 49	14% (7.3-21%)	86% (79-93%)	0.08 (0.037-0.15)	0.75% (0.56-0.96%)	0.84% (0.5-1.2%)	20 (12-30)	1700 (1200-2200)
Saudi Arabia	2010	Male	50 to 54	13% (7.3-20%)	87% (80-93%)	0.077 (0.035-0.14)	0.6% (0.41-0.78%)	0.7% (0.39-0.99%)	19 (11-28)	1200 (780-1600)
Saudi Arabia	2010	Male	55 to 59	12% (7-19%)	88% (81-93%)	0.074 (0.033-0.14)	0.48% (0.3-0.65%)	0.59% (0.28-0.88%)	15 (7-22)	710 (430-980)
Saudi Arabia	2010	Male	60 to 64	12% (6.7-20%)	88% (80-93%)	0.068 (0.027-0.14)	0.4% (0.17-0.59%)	0.46% (0.16-0.71%)	16 (5.3-25)	500 (220-750)
Saudi Arabia	2010	Male	65 to 69	12% (6.9-18%)	88% (82-93%)	0.057 (0.025-0.11)	0.36% (0.15-0.54%)	0.43% (0.15-0.65%)	15 (5.3-23)	380 (150-560)
Saudi Arabia	2010	Male	70 to 74	11% (6.7-16%)	89% (84-93%)	0.042 (0.018-0.081)	0.38% (0.21-0.54%)	0.47% (0.25-0.66%)	20 (11-28)	390 (210-540)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	2010	Male	75 to 79	9.1% (5.3-15%)	91% (85-95%)	0.031 (0.012-0.06)	0.36% (0.21-0.48%)	0.43% (0.25-0.59%)	18 (10-25)	270 (160-360)
Saudi Arabia	2010	Male	80 to 84	8.2% (4.7-13%)	92% (87-95%)	0.026 (0.011-0.047)	0.36% (0.23-0.47%)	0.42% (0.27-0.55%)	16 (10-21)	170 (110-230)
Saudi Arabia	2010	Male	85 to 89	8.2% (4.5-13%)	92% (87-95%)	0.025 (0.011-0.052)	0.26% (0.13-0.35%)	0.29% (0.14-0.4%)	8.7 (4.3-12)	68 (35-93)
Saudi Arabia	2010	Male	90 to 94	9.5% (5.5-15%)	91% (85-95%)	0.028 (0.011-0.059)	0.16% (0.034-0.26%)	0.17% (0.021-0.29%)	2.6 (0.31-4.3)	15 (3.1-24)
Saudi Arabia	2010	Male	95 plus	12% (5.9-21%)	88% (79-94%)	0.034 (0.011-0.078)	0.13% (-0.044-0.24%)	0.11% (-0.11-0.23%)	0.48 (-0.5-1)	2.5 (-0.81-4.4)
Saudi Arabia	2016	Female	All Ages	14% (12-17%)	86% (83-88%)	0.073 (0.052-0.099)	0.22% (0.089-0.35%)	-0.12% (-0.49-0.18%)	-44 (-170-67)	5400 (2000-8600)
Saudi Arabia	2016	Female	15 to 19	10% (5.6-17%)	90% (83-94%)	0.043 (0.018-0.09)	0.36% (0.23-0.48%)	0.36% (0.19-0.62%)	1.1 (0.58-1.9)	470 (300-700)
Saudi Arabia	2016	Female	20 to 24	13% (7-22%)	87% (78-93%)	0.07 (0.029-0.14)	0.73% (0.5-1%)	0.47% (0.21-0.88%)	1.8 (0.77-3.4)	1100 (670-1600)
Saudi Arabia	2016	Female	25 to 29	15% (8.2-24%)	85% (76-92%)	0.088 (0.038-0.16)	0.77% (0.56-1%)	0.55% (0.24-1%)	2.6 (1.1-4.8)	1300 (850-1900)
Saudi Arabia	2016	Female	30 to 34	18% (9.8-28%)	82% (72-90%)	0.11 (0.052-0.18)	0.69% (0.48-0.95%)	0.63% (0.32-1%)	3.2 (1.6-5.3)	1300 (810-1800)
Saudi Arabia	2016	Female	35 to 39	20% (12-30%)	80% (70-88%)	0.13 (0.062-0.2)	0.53% (0.37-0.71%)	0.57% (0.28-0.98%)	4.3 (2.1-7.3)	1100 (730-1600)
Saudi Arabia	2016	Female	40 to 44	21% (13-32%)	79% (68-87%)	0.14 (0.071-0.23)	0.39% (0.21-0.56%)	0.54% (0.21-0.93%)	5.1 (2-9)	790 (430-1200)
Saudi Arabia	2016	Female	45 to 49	21% (12-32%)	79% (68-88%)	0.15 (0.07-0.25)	0.24% (0.069-0.39%)	0.37% (0.026-0.75%)	3.2 (0.23-6.5)	350 (100-590)
Saudi Arabia	2016	Female	50 to 54	22% (13-34%)	78% (66-87%)	0.15 (0.071-0.27)	0.13% (-0.11-0.32%)	0.27% (-0.14-0.62%)	3.3 (-1.8-7.6)	170 (-140-430)
Saudi Arabia	2016	Female	55 to 59	24% (14-36%)	76% (64-86%)	0.15 (0.07-0.28)	-0.012% (-0.3-0.2%)	0.12% (-0.37-0.51%)	1.4 (-4.2-6.3)	-13 (-330-220)
Saudi Arabia	2016	Female	60 to 64	25% (15-39%)	75% (61-85%)	0.15 (0.063-0.28)	-0.18% (-0.64-0.15%)	-0.18% (-0.83-0.29%)	-5.4 (-25-8.6)	-240 (-880-200)
Saudi Arabia	2016	Female	65 to 69	26% (17-38%)	74% (62-83%)	0.13 (0.058-0.23)	-0.24% (-0.73-0.12%)	-0.26% (-0.9-0.23%)	-8.8 (-31-7.9)	-290 (-860-140)
Saudi Arabia	2016	Female	70 to 74	25% (16-36%)	75% (64-84%)	0.1 (0.046-0.19)	-0.21% (-0.72-0.14%)	-0.22% (-0.86-0.22%)	-9.7 (-38-10)	-240 (-820-160)
Saudi Arabia	2016	Female	75 to 79	22% (13-33%)	78% (67-87%)	0.079 (0.031-0.15)	-0.17% (-0.61-0.13%)	-0.18% (-0.74-0.19%)	-7.9 (-33-8.4)	-140 (-510-110)
Saudi Arabia	2016	Female	80 to 84	21% (11-32%)	79% (68-89%)	0.073 (0.029-0.14)	-0.19% (-0.61-0.12%)	-0.21% (-0.71-0.15%)	-9.6 (-33-7.1)	-110 (-360-71)
Saudi Arabia	2016	Female	85 to 89	22% (12-35%)	78% (65-88%)	0.077 (0.032-0.15)	-0.25% (-0.72-0.046%)	-0.29% (-0.84-0.059%)	-11 (-31-2.2)	-85 (-250-16)
Saudi Arabia	2016	Female	90 to 94	25% (15-38%)	75% (62-85%)	0.091 (0.037-0.18)	-0.35% (-0.85--0.024%)	-0.42% (-1--0.033%)	-11 (-26--0.84)	-56 (-140--4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Saudi Arabia	2016	Female	95 plus	30% (16-48%)	70% (52-84%)	0.11 (0.038-0.25)	-0.46% (-1.1--0.026%)	-0.58% (-1.4--0.057%)	-6.9 (-17--0.66)	-23 (-56--1.2)
Saudi Arabia	2016	Male	All Ages	9% (7.1-11%)	91% (89-93%)	0.051 (0.036-0.072)	0.69% (0.53-0.86%)	0.55% (0.34-0.78%)	300 (180-430)	24000 (18000-31000)
Saudi Arabia	2016	Male	15 to 19	5.2% (3.1-8.5%)	95% (92-97%)	0.016 (0.0063-0.032)	0.49% (0.34-0.69%)	0.26% (0.11-0.51%)	2.7 (1.2-5.1)	820 (530-1200)
Saudi Arabia	2016	Male	20 to 24	7.9% (4.4-13%)	92% (87-96%)	0.033 (0.015-0.066)	1% (0.69-1.5%)	0.46% (0.14-0.99%)	7 (2.2-15)	2000 (1300-3100)
Saudi Arabia	2016	Male	25 to 29	9.9% (5.4-16%)	90% (84-95%)	0.053 (0.024-0.098)	1.2% (0.85-1.6%)	0.67% (0.25-1.3%)	13 (4.7-24)	3000 (2000-4200)
Saudi Arabia	2016	Male	30 to 34	12% (6.3-19%)	88% (81-94%)	0.07 (0.034-0.12)	1.2% (0.85-1.5%)	0.78% (0.35-1.4%)	14 (6.5-25)	3200 (2100-4500)
Saudi Arabia	2016	Male	35 to 39	13% (7.5-21%)	87% (79-92%)	0.082 (0.042-0.14)	1.1% (0.83-1.4%)	0.98% (0.56-1.6%)	20 (11-33)	3200 (2300-4300)
Saudi Arabia	2016	Male	40 to 44	14% (8.2-22%)	86% (78-92%)	0.09 (0.046-0.16)	0.93% (0.68-1.2%)	0.93% (0.54-1.4%)	27 (15-42)	3200 (2200-4400)
Saudi Arabia	2016	Male	45 to 49	13% (7.1-21%)	87% (79-93%)	0.091 (0.042-0.16)	0.8% (0.6-1%)	0.92% (0.56-1.3%)	33 (20-48)	2700 (1900-3600)
Saudi Arabia	2016	Male	50 to 54	12% (7.2-20%)	88% (80-93%)	0.088 (0.04-0.16)	0.64% (0.44-0.83%)	0.76% (0.42-1.1%)	31 (17-44)	1900 (1200-2500)
Saudi Arabia	2016	Male	55 to 59	12% (6.9-19%)	88% (81-93%)	0.084 (0.038-0.17)	0.51% (0.31-0.7%)	0.64% (0.3-0.93%)	25 (12-37)	1200 (740-1700)
Saudi Arabia	2016	Male	60 to 64	12% (6.6-19%)	88% (81-93%)	0.077 (0.032-0.15)	0.44% (0.21-0.64%)	0.51% (0.2-0.8%)	36 (13-55)	1200 (540-1700)
Saudi Arabia	2016	Male	65 to 69	12% (6.8-18%)	88% (82-93%)	0.064 (0.029-0.12)	0.4% (0.17-0.59%)	0.48% (0.19-0.74%)	22 (8.8-34)	560 (250-830)
Saudi Arabia	2016	Male	70 to 74	10% (6.5-16%)	90% (84-93%)	0.047 (0.021-0.089)	0.41% (0.24-0.58%)	0.51% (0.28-0.73%)	20 (11-28)	380 (220-530)
Saudi Arabia	2016	Male	75 to 79	8.9% (5.2-14%)	91% (86-95%)	0.034 (0.013-0.067)	0.39% (0.23-0.53%)	0.48% (0.28-0.66%)	20 (12-27)	290 (170-400)
Saudi Arabia	2016	Male	80 to 84	7.9% (4.6-13%)	92% (87-95%)	0.029 (0.012-0.054)	0.38% (0.24-0.5%)	0.45% (0.28-0.6%)	18 (12-25)	200 (130-270)
Saudi Arabia	2016	Male	85 to 89	8% (4.5-13%)	92% (87-96%)	0.029 (0.012-0.056)	0.28% (0.14-0.39%)	0.32% (0.15-0.46%)	11 (5-15)	82 (42-120)
Saudi Arabia	2016	Male	90 to 94	9.4% (5.4-15%)	91% (85-95%)	0.032 (0.013-0.067)	0.18% (0.029-0.29%)	0.19% (0.014-0.33%)	3.6 (0.26-6.2)	21 (3.5-33)
Saudi Arabia	2016	Male	95 plus	12% (5.9-21%)	88% (79-94%)	0.039 (0.012-0.088)	0.15% (-0.043-0.28%)	0.12% (-0.11-0.3%)	0.78 (-0.71-1.9)	3.8 (-1.1-7.5)
Syria	1990	Female	All Ages	1.5% (1.1-1.9%)	99% (98-99%)	0.034 (0.022-0.048)	0.18% (0.13-0.24%)	0.13% (0.054-0.24%)	33 (14-59)	3300 (2300-4500)
Syria	1990	Female	15 to 19	1.7% (0.86-2.8%)	98% (97-99%)	0.038 (0.015-0.08)	0.32% (0.18-0.5%)	0.18% (0.08-0.33%)	0.78 (0.35-1.4)	300 (160-490)
Syria	1990	Female	20 to 24	2.2% (1.2-3.7%)	98% (96-99%)	0.065 (0.027-0.13)	0.6% (0.39-0.89%)	0.25% (0.094-0.49%)	1.1 (0.42-2.2)	540 (330-840)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	1990	Female	25 to 29	2.4% (1.2-4%)	98% (96-99%)	0.073 (0.034-0.14)	0.66% (0.46-0.92%)	0.33% (0.14-0.61%)	1.5 (0.61-2.7)	540 (340-780)
Syria	1990	Female	30 to 34	2.2% (1.1-4%)	98% (96-99%)	0.072 (0.033-0.13)	0.62% (0.42-0.85%)	0.39% (0.19-0.69%)	1.9 (0.88-3.3)	460 (290-670)
Syria	1990	Female	35 to 39	2.2% (1.1-3.5%)	98% (96-99%)	0.072 (0.033-0.14)	0.49% (0.34-0.68%)	0.37% (0.18-0.7%)	2.1 (0.95-4)	350 (230-490)
Syria	1990	Female	40 to 44	2.1% (1.2-3.7%)	98% (96-99%)	0.074 (0.031-0.15)	0.43% (0.28-0.65%)	0.43% (0.21-0.8%)	2.1 (1-4)	220 (140-340)
Syria	1990	Female	45 to 49	2.1% (1-3.4%)	98% (97-99%)	0.076 (0.029-0.16)	0.38% (0.23-0.62%)	0.41% (0.15-0.83%)	2.3 (0.91-4.6)	180 (110-300)
Syria	1990	Female	50 to 54	2.1% (1.1-3.5%)	98% (96-99%)	0.075 (0.03-0.15)	0.35% (0.18-0.61%)	0.41% (0.15-0.82%)	3.1 (1.1-6.3)	180 (91-300)
Syria	1990	Female	55 to 59	2.1% (1.1-3.7%)	98% (96-99%)	0.07 (0.027-0.14)	0.29% (0.14-0.49%)	0.35% (0.11-0.67%)	3.9 (1.2-7.5)	170 (80-300)
Syria	1990	Female	60 to 64	2.2% (1.1-4%)	98% (96-99%)	0.064 (0.023-0.13)	0.24% (0.084-0.46%)	0.29% (0.066-0.61%)	4 (0.88-8.5)	140 (48-270)
Syria	1990	Female	65 to 69	2.2% (1.1-3.8%)	98% (96-99%)	0.056 (0.022-0.11)	0.18% (0.042-0.35%)	0.22% (0.019-0.45%)	3.6 (0.3-7.4)	98 (21-190)
Syria	1990	Female	70 to 74	2.1% (1.1-3.7%)	98% (96-99%)	0.046 (0.016-0.097)	0.13% (-0.0027-0.29%)	0.16% (-0.028-0.37%)	3 (-0.57-7)	63 (-1.2-140)
Syria	1990	Female	75 to 79	1.9% (0.95-3.4%)	98% (97-99%)	0.038 (0.013-0.085)	0.1% (-0.029-0.24%)	0.11% (-0.056-0.3%)	2.2 (-1.5-6)	35 (-10-86)
Syria	1990	Female	80 to 84	1.7% (0.86-3%)	98% (97-99%)	0.033 (0.011-0.069)	0.08% (-0.017-0.2%)	0.087% (-0.038-0.23%)	1.4 (-0.59-3.6)	16 (-3.5-40)
Syria	1990	Female	85 to 89	1.7% (0.83-2.9%)	98% (97-99%)	0.031 (0.012-0.068)	0.047% (-0.049-0.15%)	0.044% (-0.078-0.17%)	0.49 (-0.84-1.9)	4.7 (-4.9-15)
Syria	1990	Female	90 to 94	1.8% (0.94-3.1%)	98% (97-99%)	0.032 (0.011-0.072)	0.023% (-0.078-0.13%)	0.012% (-0.12-0.14%)	0.062 (-0.62-0.73)	0.75 (-2.5-4.1)
Syria	1990	Female	95 plus	2.2% (1.1-4.2%)	98% (96-99%)	0.037 (0.01-0.092)	0.0044% (-0.14-0.15%)	-0.021% (-0.2-0.15%)	-0.035 (-0.33-0.26)	0.032 (-0.96-1)
Syria	1990	Male	All Ages	4.3% (3.5-5.4%)	96% (95-97%)	0.21 (0.15-0.29)	0.63% (0.4-0.89%)	0.82% (0.38-1.3%)	260 (120-430)	13000 (8600-19000)
Syria	1990	Male	15 to 19	3.9% (2.3-6.2%)	96% (94-98%)	0.11 (0.047-0.2)	0.55% (0.33-0.84%)	0.45% (0.19-0.94%)	3.4 (1.5-7.1)	610 (350-960)
Syria	1990	Male	20 to 24	6.5% (3.9-10%)	93% (90-96%)	0.27 (0.14-0.46)	1.3% (0.86-2.1%)	1.3% (0.57-2.5%)	11 (4.9-22)	1400 (880-2200)
Syria	1990	Male	25 to 29	7.9% (4.6-12%)	92% (88-95%)	0.46 (0.24-0.79)	1.9% (1.1-2.9%)	2.3% (1-4.1%)	19 (8.4-35)	1800 (1000-2800)
Syria	1990	Male	30 to 34	8.3% (4.7-13%)	92% (87-95%)	0.57 (0.3-0.95)	2% (1.2-3.2%)	2.7% (1.3-4.7%)	22 (11-38)	1700 (1000-2700)
Syria	1990	Male	35 to 39	8.8% (5.2-14%)	91% (86-95%)	0.64 (0.33-1.1)	2.1% (1.3-3.3%)	2.9% (1.4-5.1%)	25 (12-42)	1600 (950-2600)
Syria	1990	Male	40 to 44	8.8% (5.3-14%)	91% (86-95%)	0.65 (0.32-1.2)	2% (1-3.3%)	2.7% (1.2-4.9%)	21 (9-37)	1200 (620-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	1990	Male	45 to 49	8.2% (4.5-13%)	92% (87-95%)	0.56 (0.25-1)	1.7% (0.77-2.9%)	2.2% (0.78-4.1%)	19 (6.7-33)	930 (420-1600)
Syria	1990	Male	50 to 54	7.6% (4.6-12%)	92% (88-95%)	0.54 (0.26-0.99)	1.5% (0.58-2.7%)	1.9% (0.62-3.5%)	20 (6.4-38)	860 (330-1500)
Syria	1990	Male	55 to 59	7.1% (4.2-12%)	93% (88-96%)	0.57 (0.26-1)	1.6% (0.64-2.8%)	2% (0.72-3.7%)	30 (11-56)	1100 (440-1900)
Syria	1990	Male	60 to 64	6.6% (3.7-11%)	93% (89-96%)	0.51 (0.22-0.95)	1.4% (0.47-2.5%)	1.7% (0.51-3.2%)	33 (10-63)	980 (330-1800)
Syria	1990	Male	65 to 69	6.3% (3.6-10%)	94% (90-96%)	0.41 (0.19-0.75)	1% (0.25-2%)	1.2% (0.26-2.5%)	29 (5.9-58)	700 (170-1400)
Syria	1990	Male	70 to 74	6% (3.7-9.6%)	94% (90-96%)	0.3 (0.13-0.56)	0.59% (0.067-1.5%)	0.69% (0.03-1.8%)	15 (0.66-39)	300 (33-750)
Syria	1990	Male	75 to 79	5.4% (3.1-8.9%)	95% (91-97%)	0.22 (0.09-0.46)	0.33% (-0.057-1%)	0.36% (-0.12-1.2%)	7.1 (-2.4-24)	110 (-20-360)
Syria	1990	Male	80 to 84	4.9% (2.8-7.9%)	95% (92-97%)	0.18 (0.079-0.36)	0.23% (-0.078-0.79%)	0.24% (-0.14-0.89%)	3.7 (-2.1-14)	45 (-15-150)
Syria	1990	Male	85 to 89	4.6% (2.5-7.3%)	95% (93-98%)	0.16 (0.067-0.31)	0.14% (-0.11-0.56%)	0.12% (-0.18-0.61%)	1.3 (-1.9-6.4)	13 (-10-50)
Syria	1990	Male	90 to 94	4.8% (2.8-7.8%)	95% (92-97%)	0.15 (0.061-0.31)	0.063% (-0.18-0.44%)	0.026% (-0.27-0.48%)	0.12 (-1.2-2.2)	1.8 (-5.1-13)
Syria	1990	Male	95 plus	5.7% (3-10%)	94% (90-97%)	0.16 (0.055-0.35)	0.027% (-0.29-0.46%)	-0.041% (-0.42-0.48%)	-0.054 (-0.57-0.64)	0.15 (-1.6-2.5)
Syria	1995	Female	All Ages	1.3% (1-1.8%)	99% (98-99%)	0.032 (0.021-0.046)	0.23% (0.17-0.29%)	0.16% (0.073-0.27%)	41 (19-69)	4000 (2800-5300)
Syria	1995	Female	15 to 19	1.5% (0.81-2.4%)	99% (98-99%)	0.035 (0.014-0.07)	0.33% (0.2-0.5%)	0.18% (0.086-0.31%)	0.82 (0.4-1.5)	350 (200-560)
Syria	1995	Female	20 to 24	1.9% (1.1-3.1%)	98% (97-99%)	0.058 (0.026-0.11)	0.65% (0.42-0.95%)	0.24% (0.11-0.45%)	1.1 (0.49-2.1)	660 (400-1000)
Syria	1995	Female	25 to 29	2% (1.2-3.3%)	98% (97-99%)	0.066 (0.032-0.12)	0.69% (0.47-0.95%)	0.32% (0.15-0.59%)	1.5 (0.68-2.7)	630 (400-920)
Syria	1995	Female	30 to 34	1.9% (1.1-3.2%)	98% (97-99%)	0.065 (0.033-0.12)	0.64% (0.43-0.88%)	0.4% (0.21-0.67%)	2 (1-3.4)	540 (340-790)
Syria	1995	Female	35 to 39	1.9% (1-3%)	98% (97-99%)	0.064 (0.031-0.12)	0.51% (0.36-0.69%)	0.38% (0.2-0.66%)	2.2 (1.2-3.8)	410 (270-570)
Syria	1995	Female	40 to 44	1.8% (1.1-3.1%)	98% (97-99%)	0.066 (0.03-0.13)	0.44% (0.29-0.62%)	0.43% (0.22-0.75%)	3.1 (1.6-5.3)	340 (210-490)
Syria	1995	Female	45 to 49	1.9% (0.96-3.1%)	98% (97-99%)	0.07 (0.028-0.14)	0.39% (0.24-0.59%)	0.41% (0.17-0.8%)	2.8 (1.2-5.4)	220 (140-340)
Syria	1995	Female	50 to 54	1.9% (1.1-3.2%)	98% (97-99%)	0.071 (0.029-0.14)	0.36% (0.2-0.58%)	0.42% (0.17-0.79%)	3.5 (1.4-6.6)	200 (100-320)
Syria	1995	Female	55 to 59	2% (1-3.5%)	98% (96-99%)	0.069 (0.027-0.13)	0.3% (0.15-0.52%)	0.36% (0.14-0.7%)	4.1 (1.5-7.8)	180 (85-310)
Syria	1995	Female	60 to 64	2.1% (1.1-3.8%)	98% (96-99%)	0.063 (0.024-0.13)	0.25% (0.1-0.47%)	0.31% (0.091-0.62%)	5.2 (1.5-10)	180 (70-320)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	1995	Female	65 to 69	2.1% (1.1-3.5%)	98% (96-99%)	0.054 (0.022-0.1)	0.19% (0.048-0.35%)	0.23% (0.032-0.46%)	4.7 (0.63-9.5)	130 (31-240)
Syria	1995	Female	70 to 74	2% (1.1-3.5%)	98% (97-99%)	0.044 (0.017-0.095)	0.14% (0.02-0.29%)	0.17% (0.0018-0.35%)	4 (0.042-8.4)	82 (12-170)
Syria	1995	Female	75 to 79	1.8% (0.93-3.2%)	98% (97-99%)	0.036 (0.012-0.081)	0.1% (-0.021-0.24%)	0.12% (-0.042-0.29%)	2.9 (-1.1-7.2)	47 (-11-110)
Syria	1995	Female	80 to 84	1.6% (0.83-2.8%)	98% (97-99%)	0.032 (0.011-0.064)	0.084% (-0.014-0.19%)	0.093% (-0.028-0.23%)	1.9 (-0.59-4.6)	22 (-3.7-51)
Syria	1995	Female	85 to 89	1.6% (0.82-2.7%)	98% (97-99%)	0.03 (0.011-0.067)	0.049% (-0.046-0.14%)	0.047% (-0.07-0.16%)	0.67 (-1-2.3)	6.2 (-5.8-18)
Syria	1995	Female	90 to 94	1.7% (0.9-3%)	98% (97-99%)	0.032 (0.011-0.07)	0.025% (-0.073-0.12%)	0.015% (-0.11-0.13%)	0.097 (-0.7-0.87)	1 (-2.9-4.9)
Syria	1995	Female	95 plus	2.1% (1-4%)	98% (96-99%)	0.036 (0.01-0.087)	0.0064% (-0.13-0.14%)	-0.018% (-0.19-0.15%)	-0.037 (-0.38-0.3)	0.054 (-1.1-1.2)
Syria	1995	Male	All Ages	4.6% (3.9-5.6%)	95% (94-96%)	0.24 (0.17-0.33)	0.82% (0.51-1.2%)	1% (0.49-1.6%)	340 (160-540)	17000 (11000-24000)
Syria	1995	Male	15 to 19	3.9% (2.4-6%)	96% (94-98%)	0.11 (0.05-0.2)	0.56% (0.35-0.86%)	0.47% (0.19-0.93%)	4.1 (1.7-8.2)	730 (440-1100)
Syria	1995	Male	20 to 24	6.4% (4.2-9.3%)	94% (91-96%)	0.28 (0.15-0.47)	1.4% (0.86-2.1%)	1.3% (0.58-2.4%)	13 (6.2-26)	1700 (1100-2700)
Syria	1995	Male	25 to 29	7.5% (5-11%)	92% (89-95%)	0.46 (0.26-0.76)	1.9% (1.1-2.9%)	2.3% (1.1-4.1%)	22 (10-39)	2100 (1200-3200)
Syria	1995	Male	30 to 34	8.2% (5.3-12%)	92% (88-95%)	0.59 (0.34-0.97)	2.1% (1.3-3.1%)	2.8% (1.5-4.7%)	26 (13-43)	2000 (1200-3100)
Syria	1995	Male	35 to 39	9.6% (6.2-14%)	90% (86-94%)	0.74 (0.42-1.2)	2.3% (1.4-3.5%)	3.3% (1.7-5.3%)	32 (16-51)	2100 (1200-3100)
Syria	1995	Male	40 to 44	10% (6.7-15%)	90% (85-93%)	0.77 (0.43-1.3)	2.3% (1.2-3.5%)	3.2% (1.5-5.1%)	34 (16-56)	1900 (1000-2900)
Syria	1995	Male	45 to 49	9% (5.5-13%)	91% (87-95%)	0.64 (0.32-1.1)	1.9% (0.91-3.1%)	2.5% (1-4.4%)	27 (10-46)	1300 (610-2100)
Syria	1995	Male	50 to 54	8.1% (5.2-12%)	92% (88-95%)	0.6 (0.3-1)	1.6% (0.71-2.9%)	2.1% (0.76-3.8%)	25 (8.8-46)	1100 (440-1800)
Syria	1995	Male	55 to 59	7.6% (4.3-12%)	92% (88-96%)	0.64 (0.3-1.2)	1.8% (0.72-3%)	2.2% (0.82-4%)	34 (13-62)	1200 (480-2100)
Syria	1995	Male	60 to 64	7.1% (3.9-12%)	93% (88-96%)	0.58 (0.26-1.1)	1.5% (0.54-2.7%)	1.9% (0.59-3.4%)	41 (12-77)	1200 (410-2200)
Syria	1995	Male	65 to 69	6.9% (4-11%)	93% (89-96%)	0.46 (0.22-0.86)	1.2% (0.34-2.3%)	1.4% (0.35-2.9%)	37 (8.9-77)	910 (260-1800)
Syria	1995	Male	70 to 74	6.5% (4-10%)	94% (90-96%)	0.34 (0.15-0.62)	0.69% (0.088-1.7%)	0.81% (0.047-2%)	23 (1.4-58)	460 (59-1100)
Syria	1995	Male	75 to 79	5.7% (3.3-9.3%)	94% (91-97%)	0.25 (0.1-0.49)	0.39% (-0.043-1.2%)	0.43% (-0.11-1.4%)	9.9 (-2.5-32)	160 (-18-470)
Syria	1995	Male	80 to 84	5.1% (2.9-8.1%)	95% (92-97%)	0.2 (0.084-0.39)	0.27% (-0.061-0.86%)	0.29% (-0.13-0.99%)	4.9 (-2.2-17)	59 (-15-190)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	1995	Male	85 to 89	4.8% (2.6-7.8%)	95% (92-97%)	0.17 (0.073-0.33)	0.16% (-0.12-0.61%)	0.15% (-0.2-0.68%)	1.8 (-2.3-8.1)	17 (-13-64)
Syria	1995	Male	90 to 94	5% (2.9-8.1%)	95% (92-97%)	0.17 (0.067-0.33)	0.078% (-0.19-0.47%)	0.04% (-0.28-0.51%)	0.22 (-1.6-2.8)	2.6 (-6.2-16)
Syria	1995	Male	95 plus	6.1% (3-11%)	94% (89-97%)	0.18 (0.059-0.38)	0.03% (-0.35-0.52%)	-0.043% (-0.49-0.54%)	-0.063 (-0.72-0.79)	0.18 (-2-3.1)
Syria	2000	Female	All Ages	1.4% (1.1-1.8%)	99% (98-99%)	0.038 (0.025-0.055)	0.29% (0.22-0.37%)	0.2% (0.088-0.34%)	51 (22-87)	4900 (3600-6700)
Syria	2000	Female	15 to 19	1.5% (0.82-2.4%)	99% (98-99%)	0.04 (0.016-0.076)	0.34% (0.21-0.52%)	0.2% (0.1-0.34%)	1 (0.51-1.7)	410 (250-650)
Syria	2000	Female	20 to 24	1.9% (1.1-3.1%)	98% (97-99%)	0.065 (0.029-0.13)	0.68% (0.43-1%)	0.28% (0.13-0.54%)	1.5 (0.64-2.7)	800 (490-1300)
Syria	2000	Female	25 to 29	2% (1.2-3.3%)	98% (97-99%)	0.074 (0.035-0.13)	0.73% (0.52-1%)	0.37% (0.17-0.66%)	1.9 (0.85-3.4)	780 (500-1100)
Syria	2000	Female	30 to 34	2% (1.1-3.2%)	98% (97-99%)	0.073 (0.038-0.13)	0.67% (0.46-0.93%)	0.45% (0.24-0.76%)	2.4 (1.3-4.1)	650 (420-950)
Syria	2000	Female	35 to 39	1.9% (1.1-3%)	98% (97-99%)	0.072 (0.035-0.13)	0.54% (0.38-0.72%)	0.44% (0.24-0.76%)	2.7 (1.5-4.7)	490 (320-700)
Syria	2000	Female	40 to 44	1.8% (1.1-3.1%)	98% (97-99%)	0.074 (0.033-0.14)	0.46% (0.31-0.66%)	0.49% (0.26-0.84%)	3.8 (1.9-6.4)	410 (260-580)
Syria	2000	Female	45 to 49	1.8% (0.93-3%)	98% (97-99%)	0.078 (0.032-0.15)	0.41% (0.26-0.63%)	0.46% (0.2-0.86%)	4.5 (2-8.4)	350 (210-550)
Syria	2000	Female	50 to 54	1.9% (1.1-3.2%)	98% (97-99%)	0.08 (0.034-0.15)	0.39% (0.22-0.65%)	0.47% (0.2-0.89%)	4.4 (1.9-8.3)	250 (130-410)
Syria	2000	Female	55 to 59	2% (1-3.5%)	98% (96-99%)	0.079 (0.032-0.16)	0.34% (0.17-0.59%)	0.42% (0.16-0.8%)	4.8 (1.9-9.2)	210 (100-360)
Syria	2000	Female	60 to 64	2.1% (1.1-3.8%)	98% (96-99%)	0.073 (0.028-0.15)	0.29% (0.11-0.53%)	0.35% (0.11-0.71%)	5.6 (1.7-11)	190 (75-350)
Syria	2000	Female	65 to 69	2.1% (1.1-3.6%)	98% (96-99%)	0.064 (0.025-0.12)	0.21% (0.06-0.41%)	0.26% (0.057-0.53%)	5.9 (1.3-12)	160 (47-310)
Syria	2000	Female	70 to 74	2.1% (1.1-3.6%)	98% (96-99%)	0.052 (0.019-0.11)	0.16% (0.034-0.33%)	0.19% (0.02-0.41%)	5.2 (0.56-11)	110 (22-220)
Syria	2000	Female	75 to 79	1.9% (0.97-3.3%)	98% (97-99%)	0.042 (0.014-0.092)	0.11% (-0.0095-0.26%)	0.13% (-0.031-0.32%)	3.7 (-0.85-8.9)	59 (-4.8-130)
Syria	2000	Female	80 to 84	1.7% (0.84-3%)	98% (97-99%)	0.037 (0.012-0.074)	0.095% (-0.0096-0.21%)	0.1% (-0.027-0.25%)	2.5 (-0.65-5.9)	29 (-2.9-65)
Syria	2000	Female	85 to 89	1.6% (0.81-2.8%)	98% (97-99%)	0.035 (0.013-0.073)	0.057% (-0.033-0.16%)	0.056% (-0.059-0.18%)	0.93 (-0.96-3)	8.6 (-5.2-24)
Syria	2000	Female	90 to 94	1.8% (0.94-3.1%)	98% (97-99%)	0.037 (0.013-0.077)	0.034% (-0.063-0.15%)	0.025% (-0.097-0.16%)	0.19 (-0.74-1.2)	1.6 (-2.9-6.9)
Syria	2000	Female	95 plus	2.1% (1-4.1%)	98% (96-99%)	0.042 (0.012-0.1)	0.018% (-0.13-0.18%)	-0.0073% (-0.19-0.18%)	-0.017 (-0.44-0.42)	0.17 (-1.2-1.7)
Syria	2000	Male	All Ages	4.8% (4-5.8%)	95% (94-96%)	0.28 (0.2-0.38)	1% (0.69-1.4%)	1.3% (0.63-2%)	430 (210-680)	21000 (14000-29000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	2000	Male	15 to 19	3.9% (2.4-5.8%)	96% (94-98%)	0.12 (0.053-0.21)	0.59% (0.37-0.9%)	0.5% (0.21-1%)	5.3 (2.2-10)	890 (520-1400)
Syria	2000	Male	20 to 24	6.4% (4.1-9.3%)	94% (91-96%)	0.3 (0.16-0.51)	1.5% (0.93-2.2%)	1.4% (0.62-2.6%)	17 (7.4-31)	2100 (1300-3200)
Syria	2000	Male	25 to 29	7.5% (4.9-11%)	93% (89-95%)	0.5 (0.29-0.83)	2% (1.3-3%)	2.6% (1.2-4.4%)	27 (13-45)	2500 (1500-3800)
Syria	2000	Male	30 to 34	8.1% (5.3-12%)	92% (88-95%)	0.65 (0.39-1)	2.2% (1.4-3.2%)	3.1% (1.7-5.1%)	29 (16-49)	2300 (1500-3400)
Syria	2000	Male	35 to 39	9.5% (6.3-14%)	90% (86-94%)	0.81 (0.45-1.3)	2.4% (1.5-3.6%)	3.6% (2-5.7%)	36 (19-57)	2400 (1500-3500)
Syria	2000	Male	40 to 44	9.9% (6.5-15%)	90% (85-94%)	0.84 (0.46-1.4)	2.4% (1.4-3.6%)	3.5% (1.7-5.5%)	40 (19-63)	2300 (1300-3300)
Syria	2000	Male	45 to 49	8.9% (5.4-13%)	91% (87-95%)	0.7 (0.36-1.2)	2% (1-3.2%)	2.8% (1.2-4.7%)	39 (17-68)	1900 (960-3100)
Syria	2000	Male	50 to 54	8.1% (5.3-12%)	92% (88-95%)	0.66 (0.34-1.1)	1.8% (0.84-3%)	2.4% (0.97-4%)	35 (14-61)	1500 (680-2500)
Syria	2000	Male	55 to 59	7.6% (4.2-13%)	92% (87-96%)	0.71 (0.32-1.3)	1.9% (0.88-3.3%)	2.4% (1-4.2%)	43 (18-77)	1500 (660-2600)
Syria	2000	Male	60 to 64	7% (3.9-12%)	93% (88-96%)	0.63 (0.26-1.2)	1.6% (0.65-3%)	2% (0.73-3.7%)	45 (17-84)	1300 (520-2400)
Syria	2000	Male	65 to 69	6.7% (3.9-11%)	93% (89-96%)	0.5 (0.23-0.89)	1.3% (0.4-2.5%)	1.6% (0.42-3.1%)	48 (12-94)	1200 (350-2200)
Syria	2000	Male	70 to 74	6.3% (3.8-9.9%)	94% (90-96%)	0.37 (0.16-0.68)	0.82% (0.14-1.8%)	0.95% (0.12-2.2%)	34 (4.2-77)	660 (110-1400)
Syria	2000	Male	75 to 79	5.6% (3.3-9%)	94% (91-97%)	0.27 (0.11-0.53)	0.47% (-0.028-1.3%)	0.52% (-0.081-1.6%)	18 (-2.7-52)	280 (-16-780)
Syria	2000	Male	80 to 84	5% (2.9-7.9%)	95% (92-97%)	0.22 (0.092-0.41)	0.33% (-0.046-0.91%)	0.36% (-0.094-1%)	8.6 (-2.2-25)	100 (-14-280)
Syria	2000	Male	85 to 89	4.7% (2.5-7.9%)	95% (92-97%)	0.19 (0.081-0.35)	0.21% (-0.12-0.68%)	0.21% (-0.19-0.77%)	3 (-2.7-11)	27 (-15-88)
Syria	2000	Male	90 to 94	5.1% (2.8-8.5%)	95% (92-97%)	0.18 (0.077-0.36)	0.11% (-0.19-0.57%)	0.082% (-0.28-0.62%)	0.54 (-1.9-4)	4.5 (-7.5-23)
Syria	2000	Male	95 plus	6.1% (3.1-11%)	94% (89-97%)	0.19 (0.065-0.41)	0.059% (-0.33-0.63%)	-0.0082% (-0.47-0.66%)	-0.015 (-0.86-1.2)	0.43 (-2.4-4.5)
Syria	2005	Female	All Ages	1.4% (1.1-1.8%)	99% (98-99%)	0.042 (0.028-0.059)	0.32% (0.24-0.4%)	0.21% (0.11-0.35%)	53 (26-89)	5500 (4000-7300)
Syria	2005	Female	15 to 19	1.4% (0.8-2.4%)	99% (98-99%)	0.041 (0.016-0.078)	0.36% (0.22-0.55%)	0.23% (0.12-0.38%)	1.1 (0.58-1.8)	440 (250-720)
Syria	2005	Female	20 to 24	1.8% (1-3%)	98% (97-99%)	0.066 (0.03-0.13)	0.74% (0.47-1.1%)	0.32% (0.16-0.59%)	1.5 (0.73-2.7)	920 (530-1500)
Syria	2005	Female	25 to 29	2% (1.1-3.2%)	98% (97-99%)	0.076 (0.036-0.14)	0.77% (0.53-1.1%)	0.41% (0.21-0.74%)	2.1 (1.1-3.8)	900 (570-1300)
Syria	2005	Female	30 to 34	1.9% (1-3.1%)	98% (97-99%)	0.076 (0.038-0.14)	0.7% (0.49-0.96%)	0.48% (0.27-0.83%)	2.3 (1.3-4)	710 (460-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	2005	Female	35 to 39	1.8% (1-2.9%)	98% (97-99%)	0.076 (0.036-0.14)	0.57% (0.4-0.76%)	0.48% (0.28-0.81%)	2.5 (1.4-4.3)	540 (360-770)
Syria	2005	Female	40 to 44	1.8% (1.1-3%)	98% (97-99%)	0.078 (0.035-0.14)	0.48% (0.33-0.69%)	0.53% (0.3-0.9%)	3.4 (1.9-5.8)	450 (290-650)
Syria	2005	Female	45 to 49	1.8% (0.91-2.9%)	98% (97-99%)	0.082 (0.034-0.16)	0.42% (0.28-0.63%)	0.49% (0.25-0.9%)	4.3 (2.1-7.8)	370 (230-570)
Syria	2005	Female	50 to 54	1.9% (1-3.1%)	98% (97-99%)	0.083 (0.036-0.16)	0.4% (0.24-0.64%)	0.5% (0.24-0.91%)	5.4 (2.6-10)	330 (200-530)
Syria	2005	Female	55 to 59	2% (1-3.4%)	98% (97-99%)	0.081 (0.033-0.16)	0.35% (0.2-0.6%)	0.45% (0.19-0.87%)	5.4 (2.4-10)	240 (130-410)
Syria	2005	Female	60 to 64	2.1% (1-3.7%)	98% (96-99%)	0.076 (0.028-0.15)	0.3% (0.13-0.56%)	0.38% (0.14-0.74%)	6.4 (2.2-13)	220 (95-410)
Syria	2005	Female	65 to 69	2.1% (1.1-3.6%)	98% (96-99%)	0.066 (0.026-0.13)	0.22% (0.081-0.42%)	0.27% (0.072-0.55%)	5.7 (1.5-12)	160 (58-300)
Syria	2005	Female	70 to 74	2% (1.1-3.5%)	98% (96-99%)	0.053 (0.02-0.11)	0.17% (0.041-0.35%)	0.21% (0.035-0.44%)	5.6 (0.93-12)	120 (29-240)
Syria	2005	Female	75 to 79	1.8% (0.94-3.1%)	98% (97-99%)	0.043 (0.015-0.094)	0.12% (-0.0013-0.27%)	0.14% (-0.018-0.33%)	3.6 (-0.5-8.5)	59 (-0.67-130)
Syria	2005	Female	80 to 84	1.6% (0.82-2.7%)	98% (97-99%)	0.038 (0.013-0.075)	0.097% (-0.003-0.21%)	0.11% (-0.02-0.25%)	2.6 (-0.47-6)	31 (-0.96-68)
Syria	2005	Female	85 to 89	1.6% (0.81-2.7%)	98% (97-99%)	0.036 (0.014-0.074)	0.059% (-0.03-0.16%)	0.059% (-0.051-0.18%)	1.2 (-0.99-3.6)	11 (-5.3-29)
Syria	2005	Female	90 to 94	1.7% (0.9-2.9%)	98% (97-99%)	0.037 (0.014-0.078)	0.037% (-0.06-0.15%)	0.027% (-0.094-0.17%)	0.25 (-0.86-1.6)	2.1 (-3.3-8.9)
Syria	2005	Female	95 plus	2% (0.98-3.9%)	98% (96-99%)	0.042 (0.012-0.1)	0.02% (-0.12-0.17%)	-0.0048% (-0.18-0.18%)	-0.014 (-0.5-0.53)	0.24 (-1.4-2)
Syria	2005	Male	All Ages	4.8% (4-5.8%)	95% (94-96%)	0.31 (0.22-0.41)	1.1% (0.74-1.5%)	1.3% (0.68-2%)	460 (240-720)	22000 (15000-31000)
Syria	2005	Male	15 to 19	3.8% (2.4-5.7%)	96% (94-98%)	0.12 (0.054-0.22)	0.62% (0.39-0.91%)	0.55% (0.24-1%)	5.3 (2.4-10)	930 (560-1400)
Syria	2005	Male	20 to 24	6.2% (4.1-9.3%)	94% (91-96%)	0.32 (0.17-0.53)	1.5% (1-2.2%)	1.6% (0.73-2.9%)	14 (6.4-26)	2100 (1300-3100)
Syria	2005	Male	25 to 29	7.3% (4.7-11%)	93% (89-95%)	0.52 (0.29-0.87)	2% (1.3-2.9%)	2.8% (1.3-4.5%)	24 (12-40)	2500 (1600-3700)
Syria	2005	Male	30 to 34	8% (5.2-12%)	92% (88-95%)	0.67 (0.39-1.1)	2.1% (1.4-3%)	3.3% (1.8-5.1%)	24 (13-38)	2100 (1400-3000)
Syria	2005	Male	35 to 39	9.3% (5.9-13%)	91% (87-94%)	0.83 (0.46-1.4)	2.3% (1.5-3.3%)	3.8% (2.1-5.8%)	32 (18-50)	2300 (1400-3300)
Syria	2005	Male	40 to 44	9.6% (6.3-14%)	90% (86-94%)	0.86 (0.46-1.5)	2.3% (1.4-3.4%)	3.6% (1.9-5.7%)	39 (20-63)	2300 (1400-3500)
Syria	2005	Male	45 to 49	8.7% (5.3-13%)	91% (87-95%)	0.72 (0.37-1.2)	2% (1.1-3.2%)	2.9% (1.3-4.7%)	42 (19-69)	2100 (1100-3300)
Syria	2005	Male	50 to 54	7.8% (5-12%)	92% (88-95%)	0.67 (0.34-1.2)	1.8% (0.89-3%)	2.4% (1-4.1%)	45 (20-78)	1900 (950-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	2005	Male	55 to 59	7.4% (4.1-12%)	93% (88-96%)	0.72 (0.33-1.3)	1.9% (0.93-3.3%)	2.4% (1.1-4.3%)	54 (24-96)	1900 (890-3200)
Syria	2005	Male	60 to 64	6.8% (3.8-11%)	93% (89-96%)	0.64 (0.27-1.2)	1.7% (0.72-3%)	2% (0.84-3.7%)	54 (22-98)	1600 (680-2800)
Syria	2005	Male	65 to 69	6.6% (3.8-11%)	93% (89-96%)	0.51 (0.24-0.89)	1.3% (0.46-2.5%)	1.6% (0.5-3.1%)	50 (15-97)	1200 (400-2200)
Syria	2005	Male	70 to 74	6.2% (3.8-9.8%)	94% (90-96%)	0.37 (0.18-0.69)	0.84% (0.16-1.8%)	0.99% (0.14-2.2%)	38 (5.2-84)	750 (150-1600)
Syria	2005	Male	75 to 79	5.5% (3.2-9%)	95% (91-97%)	0.27 (0.12-0.53)	0.49% (-0.012-1.3%)	0.55% (-0.067-1.5%)	20 (-2.4-56)	320 (-7.4-830)
Syria	2005	Male	80 to 84	4.8% (2.6-7.9%)	95% (92-97%)	0.22 (0.09-0.4)	0.35% (-0.029-0.94%)	0.38% (-0.084-1.1%)	11 (-2.4-31)	130 (-11-350)
Syria	2005	Male	85 to 89	4.5% (2.4-7.5%)	95% (93-98%)	0.19 (0.084-0.36)	0.23% (-0.091-0.73%)	0.23% (-0.17-0.82%)	4.4 (-3.1-15)	39 (-16-120)
Syria	2005	Male	90 to 94	4.9% (2.8-8%)	95% (92-97%)	0.19 (0.076-0.36)	0.14% (-0.17-0.62%)	0.11% (-0.27-0.68%)	0.84 (-2.1-5.4)	6.5 (-8.3-30)
Syria	2005	Male	95 plus	5.9% (2.9-11%)	94% (89-97%)	0.2 (0.064-0.44)	0.087% (-0.31-0.63%)	0.02% (-0.47-0.66%)	0.044 (-0.99-1.4)	0.75 (-2.6-5.4)
Syria	2010	Female	All Ages	1.6% (1.2-2.1%)	98% (98-99%)	0.048 (0.03-0.071)	0.34% (0.27-0.43%)	0.23% (0.12-0.39%)	63 (32-110)	6700 (4800-8800)
Syria	2010	Female	15 to 19	1.6% (0.83-2.7%)	98% (97-99%)	0.044 (0.016-0.09)	0.38% (0.22-0.58%)	0.27% (0.16-0.45%)	1.1 (0.64-1.8)	450 (250-740)
Syria	2010	Female	20 to 24	2.1% (1.1-3.6%)	98% (96-99%)	0.074 (0.03-0.15)	0.76% (0.48-1.1%)	0.37% (0.18-0.7%)	1.7 (0.82-3.2)	1000 (610-1600)
Syria	2010	Female	25 to 29	2.3% (1.2-3.9%)	98% (96-99%)	0.083 (0.036-0.16)	0.79% (0.55-1.1%)	0.45% (0.25-0.81%)	2.3 (1.2-4)	1100 (720-1700)
Syria	2010	Female	30 to 34	2.2% (1.1-3.8%)	98% (96-99%)	0.082 (0.038-0.15)	0.73% (0.5-1%)	0.52% (0.31-0.87%)	2.7 (1.6-4.6)	960 (610-1400)
Syria	2010	Female	35 to 39	2.1% (1.1-3.6%)	98% (96-99%)	0.083 (0.036-0.16)	0.59% (0.41-0.79%)	0.52% (0.29-0.93%)	3 (1.6-5.4)	680 (450-970)
Syria	2010	Female	40 to 44	2.1% (1.1-3.6%)	98% (96-99%)	0.086 (0.035-0.16)	0.48% (0.33-0.7%)	0.56% (0.3-0.94%)	3.7 (2-6.4)	520 (330-750)
Syria	2010	Female	45 to 49	2% (1-3.4%)	98% (97-99%)	0.088 (0.035-0.18)	0.43% (0.28-0.65%)	0.52% (0.26-1%)	4.8 (2.4-9.3)	450 (280-690)
Syria	2010	Female	50 to 54	2.1% (1.1-3.4%)	98% (97-99%)	0.088 (0.035-0.17)	0.4% (0.25-0.64%)	0.53% (0.25-0.98%)	6.2 (3-12)	390 (230-620)
Syria	2010	Female	55 to 59	2.1% (1.1-3.8%)	98% (96-99%)	0.083 (0.031-0.17)	0.36% (0.2-0.6%)	0.48% (0.22-0.88%)	7 (3.2-13)	330 (190-550)
Syria	2010	Female	60 to 64	2.1% (1.1-3.9%)	98% (96-99%)	0.075 (0.027-0.16)	0.3% (0.15-0.56%)	0.4% (0.15-0.77%)	6.8 (2.6-13)	240 (110-430)
Syria	2010	Female	65 to 69	2.1% (1-3.7%)	98% (96-99%)	0.065 (0.025-0.13)	0.23% (0.098-0.43%)	0.3% (0.1-0.59%)	6.5 (2.2-13)	180 (76-340)
Syria	2010	Female	70 to 74	2.1% (1.1-3.6%)	98% (96-99%)	0.053 (0.019-0.11)	0.18% (0.057-0.36%)	0.23% (0.055-0.46%)	6.3 (1.5-13)	130 (41-260)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	2010	Female	75 to 79	1.9% (0.93-3.3%)	98% (97-99%)	0.043 (0.014-0.093)	0.13% (0.024-0.29%)	0.16% (0.013-0.37%)	5.3 (0.44-12)	86 (16-190)
Syria	2010	Female	80 to 84	1.7% (0.82-2.9%)	98% (97-99%)	0.038 (0.013-0.078)	0.11% (0.0076-0.23%)	0.13% (-0.0073-0.27%)	3.6 (-0.21-7.7)	42 (2.8-89)
Syria	2010	Female	85 to 89	1.6% (0.79-2.8%)	98% (97-99%)	0.036 (0.012-0.076)	0.068% (-0.023-0.18%)	0.071% (-0.046-0.21%)	1.6 (-1-4.7)	14 (-5-37)
Syria	2010	Female	90 to 94	1.8% (0.89-3%)	98% (97-99%)	0.037 (0.013-0.079)	0.047% (-0.05-0.16%)	0.038% (-0.085-0.18%)	0.42 (-0.95-2)	3.3 (-3.6-11)
Syria	2010	Female	95 plus	2.1% (1-4%)	98% (96-99%)	0.042 (0.011-0.1)	0.025% (-0.12-0.17%)	-0.0035% (-0.18-0.18%)	-0.012 (-0.61-0.61)	0.37 (-1.7-2.6)
Syria	2010	Male	All Ages	4.9% (4-6.2%)	95% (94-96%)	0.31 (0.2-0.43)	1.1% (0.7-1.5%)	1.2% (0.58-1.9%)	470 (220-740)	24000 (16000-33000)
Syria	2010	Male	15 to 19	3.8% (2.2-6%)	96% (94-98%)	0.12 (0.051-0.23)	0.65% (0.4-0.98%)	0.59% (0.26-1.2%)	5.1 (2.2-10)	950 (570-1500)
Syria	2010	Male	20 to 24	6.3% (3.7-9.8%)	94% (90-96%)	0.31 (0.15-0.55)	1.6% (1-2.3%)	1.6% (0.74-3.2%)	16 (7.2-31)	2400 (1500-3600)
Syria	2010	Male	25 to 29	7.6% (4.4-11%)	92% (89-96%)	0.52 (0.26-0.9)	2.1% (1.3-3%)	2.8% (1.3-4.9%)	29 (14-51)	3200 (1900-4600)
Syria	2010	Male	30 to 34	8% (4.5-13%)	92% (87-95%)	0.65 (0.34-1.1)	2.1% (1.3-3.1%)	3.2% (1.6-5.1%)	30 (15-49)	2700 (1700-3900)
Syria	2010	Male	35 to 39	8.6% (5-13%)	91% (87-95%)	0.74 (0.37-1.3)	2.2% (1.3-3.1%)	3.4% (1.7-5.5%)	34 (16-54)	2500 (1500-3600)
Syria	2010	Male	40 to 44	8.6% (5.2-14%)	91% (86-95%)	0.75 (0.37-1.3)	2.1% (1.2-3.1%)	3.2% (1.4-5.1%)	37 (17-60)	2300 (1300-3500)
Syria	2010	Male	45 to 49	8% (4.4-13%)	92% (87-96%)	0.64 (0.28-1.2)	1.8% (0.91-2.8%)	2.6% (1.1-4.3%)	40 (17-67)	2100 (1100-3400)
Syria	2010	Male	50 to 54	7.4% (4.4-12%)	93% (88-96%)	0.62 (0.28-1.1)	1.7% (0.76-2.7%)	2.2% (0.85-3.9%)	45 (17-81)	2000 (900-3300)
Syria	2010	Male	55 to 59	6.9% (3.9-11%)	93% (89-96%)	0.65 (0.29-1.2)	1.7% (0.78-2.8%)	2.2% (0.93-3.8%)	56 (24-98)	2000 (970-3400)
Syria	2010	Male	60 to 64	6.3% (3.6-11%)	94% (89-96%)	0.58 (0.24-1.2)	1.5% (0.62-2.6%)	1.9% (0.71-3.3%)	54 (20-96)	1600 (650-2700)
Syria	2010	Male	65 to 69	6.1% (3.5-9.7%)	94% (90-96%)	0.46 (0.2-0.86)	1.2% (0.39-2.2%)	1.5% (0.44-2.7%)	46 (14-85)	1100 (370-2100)
Syria	2010	Male	70 to 74	5.8% (3.6-9.3%)	94% (91-96%)	0.34 (0.15-0.65)	0.77% (0.15-1.6%)	0.91% (0.14-2%)	32 (4.8-71)	640 (120-1400)
Syria	2010	Male	75 to 79	5.2% (3-8.5%)	95% (92-97%)	0.25 (0.1-0.5)	0.47% (0.013-1.2%)	0.53% (-0.042-1.4%)	21 (-1.7-57)	340 (9-870)
Syria	2010	Male	80 to 84	4.6% (2.6-7.4%)	95% (93-97%)	0.2 (0.083-0.4)	0.35% (-0.023-0.9%)	0.38% (-0.074-1%)	13 (-2.6-35)	150 (-9.2-400)
Syria	2010	Male	85 to 89	4.3% (2.4-7.1%)	96% (93-98%)	0.18 (0.072-0.35)	0.24% (-0.072-0.74%)	0.24% (-0.14-0.86%)	5.9 (-3.4-21)	52 (-16-160)
Syria	2010	Male	90 to 94	4.7% (2.6-7.7%)	95% (92-97%)	0.17 (0.066-0.35)	0.14% (-0.16-0.65%)	0.11% (-0.26-0.71%)	1.2 (-2.8-7.9)	9.3 (-11-43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	2010	Male	95 plus	5.5% (2.9-10%)	94% (90-97%)	0.18 (0.056-0.42)	0.084% (-0.3-0.65%)	0.02% (-0.43-0.67%)	0.054 (-1.1-1.8)	0.91 (-3.2-6.9)
Syria	2016	Female	All Ages	1.6% (1.2-2.1%)	98% (98-99%)	0.045 (0.023-0.081)	0.22% (0.14-0.33%)	0.14% (0.059-0.29%)	59 (26-110)	5700 (4000-7800)
Syria	2016	Female	15 to 19	1.6% (0.85-2.7%)	98% (97-99%)	0.042 (0.015-0.095)	0.18% (0.088-0.32%)	0.045% (0.018-0.11%)	0.93 (0.51-1.7)	440 (250-700)
Syria	2016	Female	20 to 24	2.2% (1.1-3.6%)	98% (96-99%)	0.071 (0.025-0.16)	0.4% (0.21-0.68%)	0.074% (0.025-0.19%)	1.2 (0.53-2.4)	750 (440-1200)
Syria	2016	Female	25 to 29	2.3% (1.2-3.9%)	98% (96-99%)	0.079 (0.029-0.17)	0.51% (0.3-0.8%)	0.13% (0.047-0.31%)	1.4 (0.69-2.8)	750 (470-1100)
Syria	2016	Female	30 to 34	2.2% (1.1-3.8%)	98% (96-99%)	0.077 (0.03-0.15)	0.49% (0.3-0.76%)	0.17% (0.07-0.37%)	2.1 (1.2-3.7)	770 (490-1100)
Syria	2016	Female	35 to 39	2.1% (1.1-3.5%)	98% (97-99%)	0.077 (0.03-0.16)	0.43% (0.28-0.63%)	0.21% (0.093-0.43%)	2.6 (1.4-4.5)	630 (400-900)
Syria	2016	Female	40 to 44	2.1% (1.1-3.7%)	98% (96-99%)	0.08 (0.029-0.17)	0.39% (0.24-0.58%)	0.28% (0.14-0.56%)	3.3 (1.7-5.8)	480 (290-720)
Syria	2016	Female	45 to 49	2% (1-3.4%)	98% (97-99%)	0.083 (0.028-0.19)	0.35% (0.22-0.56%)	0.31% (0.13-0.64%)	4.1 (1.9-8.5)	420 (250-670)
Syria	2016	Female	50 to 54	2.1% (1.1-3.5%)	98% (96-99%)	0.081 (0.029-0.18)	0.33% (0.2-0.54%)	0.35% (0.15-0.7%)	5.9 (2.7-12)	400 (240-670)
Syria	2016	Female	55 to 59	2.1% (1.1-3.7%)	98% (96-99%)	0.076 (0.027-0.16)	0.3% (0.16-0.51%)	0.35% (0.15-0.68%)	6.7 (2.8-13)	330 (180-570)
Syria	2016	Female	60 to 64	2.1% (1.1-3.9%)	98% (96-99%)	0.069 (0.024-0.16)	0.26% (0.11-0.47%)	0.31% (0.11-0.64%)	7.5 (2.7-15)	280 (130-510)
Syria	2016	Female	65 to 69	2.2% (1.1-3.8%)	98% (96-99%)	0.062 (0.02-0.13)	0.2% (0.075-0.4%)	0.25% (0.07-0.52%)	5.9 (1.5-12)	170 (63-330)
Syria	2016	Female	70 to 74	2.1% (1.1-3.6%)	98% (96-99%)	0.051 (0.017-0.12)	0.17% (0.046-0.34%)	0.21% (0.042-0.45%)	5.9 (1.2-12)	130 (36-250)
Syria	2016	Female	75 to 79	1.9% (0.95-3.3%)	98% (97-99%)	0.042 (0.012-0.1)	0.13% (0.02-0.29%)	0.15% (0.0043-0.37%)	5.1 (0.14-12)	84 (14-190)
Syria	2016	Female	80 to 84	1.7% (0.83-3%)	98% (97-99%)	0.037 (0.011-0.083)	0.1% (0.001-0.25%)	0.12% (-0.014-0.29%)	4 (-0.46-10)	48 (0.5-110)
Syria	2016	Female	85 to 89	1.7% (0.81-2.9%)	98% (97-99%)	0.035 (0.011-0.08)	0.065% (-0.031-0.2%)	0.065% (-0.053-0.23%)	2.1 (-1.7-7.5)	20 (-9.2-61)
Syria	2016	Female	90 to 94	1.8% (0.93-3.1%)	98% (97-99%)	0.036 (0.011-0.082)	0.04% (-0.065-0.17%)	0.03% (-0.1-0.19%)	0.48 (-1.6-3.1)	4.1 (-6.4-17)
Syria	2016	Female	95 plus	2.2% (1-4.1%)	98% (96-99%)	0.041 (0.01-0.11)	0.018% (-0.11-0.18%)	-0.009% (-0.17-0.19%)	-0.06 (-1.2-1.2)	0.5 (-3.1-4.8)
Syria	2016	Male	All Ages	5% (4-6.2%)	95% (94-96%)	0.29 (0.16-0.5)	0.51% (0.25-0.9%)	0.56% (0.17-1.1%)	410 (130-720)	20000 (11000-31000)
Syria	2016	Male	15 to 19	3.8% (2.2-6.1%)	96% (94-98%)	0.11 (0.041-0.25)	0.19% (0.088-0.4%)	0.092% (0.026-0.25%)	5.1 (2.1-11)	930 (530-1500)
Syria	2016	Male	20 to 24	6.4% (3.7-10%)	94% (90-96%)	0.29 (0.12-0.59)	0.32% (0.14-0.72%)	0.17% (0.043-0.48%)	13 (5.1-29)	1900 (1100-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Syria	2016	Male	25 to 29	7.7% (4.5-12%)	92% (88-95%)	0.51 (0.21-0.97)	0.53% (0.23-1.1%)	0.35% (0.094-0.97%)	19 (7.1-36)	2200 (1200-3300)
Syria	2016	Male	30 to 34	8.1% (4.6-13%)	92% (87-95%)	0.62 (0.29-1.2)	0.67% (0.31-1.4%)	0.5% (0.16-1.3%)	22 (9.3-39)	2100 (1200-3200)
Syria	2016	Male	35 to 39	8.6% (5-14%)	91% (86-95%)	0.7 (0.3-1.3)	0.88% (0.44-1.6%)	0.78% (0.27-1.8%)	26 (11-46)	2100 (1200-3100)
Syria	2016	Male	40 to 44	8.7% (5.3-14%)	91% (86-95%)	0.71 (0.29-1.4)	1.1% (0.52-1.8%)	1.1% (0.38-2.2%)	29 (11-50)	1900 (960-3000)
Syria	2016	Male	45 to 49	8.1% (4.5-13%)	92% (87-96%)	0.61 (0.24-1.2)	1.1% (0.51-1.9%)	1.3% (0.41-2.4%)	32 (11-59)	1700 (790-3000)
Syria	2016	Male	50 to 54	7.5% (4.5-12%)	93% (88-96%)	0.58 (0.24-1.2)	1.2% (0.45-2%)	1.4% (0.4-2.6%)	39 (12-74)	1800 (680-3100)
Syria	2016	Male	55 to 59	6.9% (4-11%)	93% (89-96%)	0.61 (0.24-1.3)	1.3% (0.46-2.2%)	1.5% (0.44-2.8%)	51 (15-96)	1900 (670-3300)
Syria	2016	Male	60 to 64	6.4% (3.5-11%)	94% (89-96%)	0.55 (0.2-1.1)	1.2% (0.38-2.1%)	1.4% (0.4-2.6%)	56 (17-100)	1700 (570-3000)
Syria	2016	Male	65 to 69	6.1% (3.5-9.7%)	94% (90-96%)	0.43 (0.17-0.88)	0.96% (0.24-1.9%)	1.1% (0.23-2.3%)	45 (9.2-93)	1100 (270-2200)
Syria	2016	Male	70 to 74	5.8% (3.6-9.2%)	94% (91-96%)	0.32 (0.13-0.67)	0.64% (0.038-1.5%)	0.73% (0.0045-1.8%)	30 (0.19-73)	590 (36-1400)
Syria	2016	Male	75 to 79	5.2% (3-8.4%)	95% (92-97%)	0.24 (0.082-0.52)	0.4% (-0.044-1.1%)	0.44% (-0.11-1.3%)	18 (-4.3-55)	280 (-31-800)
Syria	2016	Male	80 to 84	4.6% (2.5-7.5%)	95% (92-97%)	0.19 (0.068-0.4)	0.3% (-0.064-0.92%)	0.32% (-0.12-1%)	10 (-3.7-33)	120 (-26-360)
Syria	2016	Male	85 to 89	4.3% (2.3-7%)	96% (93-98%)	0.17 (0.057-0.36)	0.22% (-0.077-0.81%)	0.21% (-0.14-0.9%)	6.8 (-4.4-28)	60 (-21-220)
Syria	2016	Male	90 to 94	4.6% (2.7-7.6%)	95% (92-97%)	0.16 (0.056-0.35)	0.13% (-0.15-0.65%)	0.1% (-0.23-0.72%)	1.6 (-3.7-11)	12 (-14-62)
Syria	2016	Male	95 plus	5.5% (2.9-9.9%)	94% (90-97%)	0.17 (0.05-0.44)	0.077% (-0.29-0.64%)	0.017% (-0.41-0.67%)	0.076 (-1.9-3)	1.4 (-5-12)
Tunisia	1990	Female	All Ages	2% (1.5-2.6%)	98% (97-98%)	0.029 (0.021-0.038)	0.19% (0.14-0.25%)	0.098% (0.012-0.21%)	18 (2.2-39)	2300 (1600-3100)
Tunisia	1990	Female	15 to 19	2% (1-3.4%)	98% (97-99%)	0.027 (0.011-0.052)	0.34% (0.2-0.52%)	0.18% (0.096-0.31%)	0.39 (0.2-0.67)	170 (98-280)
Tunisia	1990	Female	20 to 24	2.7% (1.3-4.6%)	97% (95-99%)	0.045 (0.019-0.088)	0.68% (0.44-1%)	0.26% (0.13-0.45%)	0.66 (0.34-1.2)	380 (220-600)
Tunisia	1990	Female	25 to 29	2.9% (1.3-5.3%)	97% (95-99%)	0.049 (0.021-0.092)	0.7% (0.5-0.95%)	0.34% (0.2-0.55%)	0.95 (0.54-1.6)	390 (250-570)
Tunisia	1990	Female	30 to 34	2.8% (1.2-5.2%)	97% (95-99%)	0.049 (0.022-0.092)	0.64% (0.45-0.88%)	0.38% (0.23-0.6%)	1.2 (0.7-1.9)	340 (220-510)
Tunisia	1990	Female	35 to 39	2.7% (1.2-4.8%)	97% (95-99%)	0.051 (0.022-0.094)	0.51% (0.36-0.69%)	0.37% (0.23-0.58%)	1.2 (0.75-2)	260 (160-360)
Tunisia	1990	Female	40 to 44	2.6% (1.3-4.8%)	97% (95-99%)	0.052 (0.023-0.095)	0.42% (0.28-0.59%)	0.41% (0.24-0.62%)	1.3 (0.78-2.1)	170 (100-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	1990	Female	45 to 49	2.6% (1.2-4.5%)	97% (96-99%)	0.055 (0.023-0.1)	0.35% (0.23-0.49%)	0.36% (0.2-0.61%)	1.5 (0.79-2.5)	140 (89-200)
Tunisia	1990	Female	50 to 54	2.8% (1.3-5%)	97% (95-99%)	0.057 (0.025-0.12)	0.3% (0.18-0.43%)	0.35% (0.17-0.59%)	2.2 (1.1-3.8)	140 (88-220)
Tunisia	1990	Female	55 to 59	2.9% (1.4-5.5%)	97% (95-99%)	0.057 (0.023-0.11)	0.23% (0.11-0.37%)	0.28% (0.095-0.52%)	2.5 (0.81-4.7)	120 (62-200)
Tunisia	1990	Female	60 to 64	2.9% (1.4-5.4%)	97% (95-99%)	0.051 (0.02-0.1)	0.16% (0.008-0.32%)	0.19% (-0.022-0.47%)	2.1 (-0.24-5.2)	79 (3.4-160)
Tunisia	1990	Female	65 to 69	2.9% (1.4-5%)	97% (95-99%)	0.044 (0.018-0.085)	0.1% (-0.044-0.26%)	0.12% (-0.087-0.38%)	1.8 (-1.3-5.6)	53 (-23-140)
Tunisia	1990	Female	70 to 74	2.7% (1.3-4.9%)	97% (95-99%)	0.036 (0.014-0.079)	0.06% (-0.081-0.22%)	0.07% (-0.12-0.32%)	1.1 (-1.9-5.1)	26 (-34-97)
Tunisia	1990	Female	75 to 79	2.4% (1.1-4.5%)	98% (96-99%)	0.03 (0.011-0.064)	0.043% (-0.12-0.19%)	0.047% (-0.16-0.25%)	0.94 (-3.1-5)	16 (-42-73)
Tunisia	1990	Female	80 to 84	2.2% (1-3.9%)	98% (96-99%)	0.026 (0.0093-0.053)	0.029% (-0.1-0.17%)	0.027% (-0.14-0.21%)	0.41 (-2.1-3.2)	5.9 (-21-34)
Tunisia	1990	Female	85 to 89	2.1% (1-3.9%)	98% (96-99%)	0.025 (0.0098-0.05)	0.007% (-0.12-0.13%)	-0.0021% (-0.17-0.15%)	-0.022 (-1.6-1.5)	0.66 (-11-12)
Tunisia	1990	Female	90 to 94	2.4% (1.2-4.4%)	98% (96-99%)	0.027 (0.01-0.057)	-0.018% (-0.16-0.1%)	-0.038% (-0.21-0.12%)	-0.18 (-0.99-0.55)	-0.52 (-4.5-3)
Tunisia	1990	Female	95 plus	3.1% (1.3-6.3%)	97% (94-99%)	0.033 (0.0098-0.078)	-0.045% (-0.24-0.091%)	-0.08% (-0.32-0.092%)	-0.11 (-0.46-0.13)	-0.26 (-1.4-0.54)
Tunisia	1990	Male	All Ages	9.4% (7.4-12%)	91% (88-93%)	0.37 (0.33-0.42)	0.98% (0.7-1.3%)	1.1% (0.59-1.7%)	280 (150-440)	14000 (9700-19000)
Tunisia	1990	Male	15 to 19	7.7% (4.5-12%)	92% (88-95%)	0.17 (0.08-0.32)	0.94% (0.58-1.5%)	1.1% (0.49-2.1%)	4.5 (2-8.7)	580 (340-920)
Tunisia	1990	Male	20 to 24	13% (7.5-20%)	87% (80-92%)	0.46 (0.24-0.75)	2.3% (1.5-3.5%)	2.8% (1.4-5.1%)	13 (6.5-24)	1400 (900-2200)
Tunisia	1990	Male	25 to 29	16% (9.5-24%)	84% (76-90%)	0.73 (0.41-1.1)	3% (1.9-4.4%)	4.3% (2.2-6.9%)	22 (11-35)	1900 (1100-2800)
Tunisia	1990	Male	30 to 34	17% (9.8-26%)	83% (74-90%)	0.84 (0.49-1.3)	3% (1.9-4.6%)	4.5% (2.5-7.3%)	23 (13-38)	1800 (1100-2600)
Tunisia	1990	Male	35 to 39	18% (10-26%)	82% (74-90%)	0.88 (0.5-1.4)	2.9% (1.7-4.4%)	4.4% (2.3-7.1%)	22 (11-37)	1500 (890-2300)
Tunisia	1990	Male	40 to 44	17% (11-26%)	83% (74-89%)	0.89 (0.46-1.5)	2.8% (1.4-4.4%)	4.1% (1.9-6.9%)	19 (8.7-33)	1100 (570-1800)
Tunisia	1990	Male	45 to 49	15% (8.5-24%)	85% (76-92%)	0.84 (0.4-1.4)	2.6% (1.3-4.3%)	3.8% (1.7-6.3%)	23 (10-40)	1100 (560-1900)
Tunisia	1990	Male	50 to 54	14% (8.5-22%)	86% (78-91%)	0.76 (0.39-1.3)	2.1% (0.95-3.7%)	3% (1.1-5.3%)	28 (11-51)	1200 (530-2100)
Tunisia	1990	Male	55 to 59	13% (7.3-21%)	87% (79-93%)	0.67 (0.32-1.2)	1.8% (0.75-3.6%)	2.4% (0.94-4.8%)	35 (13-70)	1300 (510-2400)
Tunisia	1990	Male	60 to 64	12% (6.5-20%)	88% (80-94%)	0.55 (0.24-1.1)	1.4% (0.39-3%)	1.8% (0.46-3.9%)	30 (8-68)	910 (270-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	1990	Male	65 to 69	11% (6.4-19%)	89% (81-94%)	0.45 (0.21-0.83)	0.92% (0.2-2.1%)	1.1% (0.19-2.6%)	25 (4.2-58)	620 (130-1400)
Tunisia	1990	Male	70 to 74	11% (6.4-18%)	89% (82-94%)	0.35 (0.16-0.69)	0.58% (0.0063-1.6%)	0.67% (-0.059-1.9%)	15 (-1.3-45)	300 (3.2-840)
Tunisia	1990	Male	75 to 79	9.6% (5.5-16%)	90% (84-95%)	0.27 (0.12-0.52)	0.33% (-0.18-1.1%)	0.36% (-0.26-1.3%)	11 (-7.8-38)	160 (-95-550)
Tunisia	1990	Male	80 to 84	8.7% (4.9-14%)	91% (86-95%)	0.23 (0.1-0.41)	0.25% (-0.13-0.75%)	0.26% (-0.2-0.85%)	5 (-4.1-16)	59 (-32-180)
Tunisia	1990	Male	85 to 89	8.2% (4.5-13%)	92% (87-95%)	0.2 (0.088-0.37)	0.15% (-0.16-0.54%)	0.14% (-0.24-0.59%)	1.3 (-2.1-5.3)	12 (-12-42)
Tunisia	1990	Male	90 to 94	8.6% (4.8-14%)	91% (86-95%)	0.19 (0.08-0.37)	0.068% (-0.25-0.43%)	0.031% (-0.36-0.46%)	0.095 (-1.1-1.4)	1.3 (-4.6-8.2)
Tunisia	1990	Male	95 plus	10% (5.1-18%)	90% (82-95%)	0.2 (0.067-0.44)	0.042% (-0.35-0.49%)	-0.035% (-0.51-0.48%)	-0.027 (-0.39-0.37)	0.13 (-1-1.5)
Tunisia	1995	Female	All Ages	2% (1.5-2.6%)	98% (97-98%)	0.031 (0.024-0.041)	0.24% (0.18-0.3%)	0.12% (0.022-0.24%)	22 (4.1-45)	2700 (1900-3700)
Tunisia	1995	Female	15 to 19	1.9% (0.99-3.3%)	98% (97-99%)	0.027 (0.011-0.053)	0.35% (0.21-0.52%)	0.19% (0.1-0.32%)	0.42 (0.23-0.73)	190 (110-300)
Tunisia	1995	Female	20 to 24	2.6% (1.2-4.6%)	97% (95-99%)	0.046 (0.018-0.09)	0.7% (0.45-1%)	0.28% (0.15-0.49%)	0.7 (0.38-1.2)	400 (230-620)
Tunisia	1995	Female	25 to 29	2.7% (1.3-5.1%)	97% (95-99%)	0.051 (0.023-0.096)	0.75% (0.52-1%)	0.36% (0.21-0.57%)	0.97 (0.56-1.5)	450 (290-680)
Tunisia	1995	Female	30 to 34	2.6% (1.2-5%)	97% (95-99%)	0.051 (0.022-0.095)	0.67% (0.47-0.93%)	0.41% (0.26-0.62%)	1.3 (0.82-2.1)	410 (260-610)
Tunisia	1995	Female	35 to 39	2.6% (1.2-4.7%)	97% (95-99%)	0.054 (0.024-0.096)	0.53% (0.38-0.71%)	0.4% (0.26-0.62%)	1.5 (0.89-2.3)	310 (200-440)
Tunisia	1995	Female	40 to 44	2.6% (1.3-4.5%)	97% (95-99%)	0.055 (0.024-0.1)	0.44% (0.3-0.6%)	0.43% (0.26-0.66%)	1.9 (1.2-3)	250 (160-350)
Tunisia	1995	Female	45 to 49	2.5% (1.1-4.2%)	97% (96-99%)	0.057 (0.024-0.11)	0.36% (0.24-0.49%)	0.39% (0.22-0.64%)	1.8 (1-2.9)	160 (110-230)
Tunisia	1995	Female	50 to 54	2.7% (1.2-4.8%)	97% (95-99%)	0.059 (0.024-0.12)	0.31% (0.2-0.44%)	0.38% (0.2-0.63%)	2 (1.1-3.5)	130 (86-200)
Tunisia	1995	Female	55 to 59	2.8% (1.3-5.3%)	97% (95-99%)	0.058 (0.023-0.12)	0.24% (0.13-0.38%)	0.31% (0.13-0.55%)	2.8 (1.1-4.9)	130 (69-210)
Tunisia	1995	Female	60 to 64	2.8% (1.3-5.3%)	97% (95-99%)	0.053 (0.021-0.11)	0.17% (0.035-0.33%)	0.22% (0.0075-0.5%)	3 (0.098-6.8)	110 (22-220)
Tunisia	1995	Female	65 to 69	2.8% (1.3-4.9%)	97% (95-99%)	0.046 (0.019-0.088)	0.11% (-0.025-0.26%)	0.14% (-0.054-0.37%)	2.4 (-0.89-6.2)	66 (-15-150)
Tunisia	1995	Female	70 to 74	2.7% (1.3-4.8%)	97% (95-99%)	0.038 (0.014-0.083)	0.069% (-0.064-0.22%)	0.086% (-0.11-0.32%)	1.8 (-2.4-6.8)	39 (-36-130)
Tunisia	1995	Female	75 to 79	2.4% (1.1-4.5%)	98% (96-99%)	0.031 (0.011-0.067)	0.05% (-0.11-0.19%)	0.058% (-0.14-0.26%)	1.3 (-3.2-5.8)	21 (-44-83)
Tunisia	1995	Female	80 to 84	2.2% (0.97-3.9%)	98% (96-99%)	0.027 (0.0095-0.056)	0.034% (-0.093-0.16%)	0.035% (-0.12-0.19%)	0.71 (-2.6-4)	9.3 (-25-43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	1995	Female	85 to 89	2.1% (0.98-4%)	98% (96-99%)	0.026 (0.01-0.053)	0.012% (-0.11-0.12%)	0.0058% (-0.15-0.15%)	0.089 (-2.2-2.3)	1.8 (-16-17)
Tunisia	1995	Female	90 to 94	2.4% (1.2-4.5%)	98% (95-99%)	0.028 (0.011-0.061)	-0.013% (-0.15-0.1%)	-0.03% (-0.2-0.12%)	-0.2 (-1.4-0.76)	-0.51 (-6.2-4.1)
Tunisia	1995	Female	95 plus	3.1% (1.2-6.5%)	97% (94-99%)	0.035 (0.01-0.085)	-0.041% (-0.24-0.093%)	-0.075% (-0.32-0.091%)	-0.15 (-0.64-0.18)	-0.33 (-1.9-0.77)
Tunisia	1995	Male	All Ages	9.8% (7.7-12%)	90% (88-92%)	0.41 (0.36-0.46)	1.2% (0.87-1.6%)	1.2% (0.69-1.9%)	330 (180-530)	17000 (12000-22000)
Tunisia	1995	Male	15 to 19	7.5% (4.3-12%)	92% (88-96%)	0.18 (0.081-0.31)	0.95% (0.59-1.4%)	1.1% (0.51-2.2%)	4.7 (2.1-9)	630 (380-960)
Tunisia	1995	Male	20 to 24	13% (7.6-20%)	87% (80-92%)	0.47 (0.24-0.77)	2.4% (1.5-3.5%)	2.9% (1.4-5.1%)	15 (7.4-26)	1600 (990-2400)
Tunisia	1995	Male	25 to 29	16% (9.5-24%)	84% (76-90%)	0.74 (0.42-1.2)	3.1% (2-4.7%)	4.4% (2.2-7.2%)	24 (12-40)	2100 (1300-3100)
Tunisia	1995	Male	30 to 34	17% (9.6-26%)	83% (74-90%)	0.86 (0.5-1.3)	3.1% (1.9-4.6%)	4.6% (2.5-7.3%)	27 (15-44)	2000 (1200-3000)
Tunisia	1995	Male	35 to 39	17% (10-26%)	83% (74-90%)	0.89 (0.5-1.4)	2.9% (1.8-4.4%)	4.5% (2.4-7.2%)	26 (14-43)	1800 (1100-2700)
Tunisia	1995	Male	40 to 44	17% (10-26%)	83% (74-90%)	0.91 (0.47-1.5)	2.7% (1.5-4.2%)	4.2% (2-6.8%)	28 (13-47)	1600 (870-2600)
Tunisia	1995	Male	45 to 49	16% (8.7-24%)	84% (76-91%)	0.88 (0.41-1.5)	2.7% (1.4-4.4%)	3.9% (1.8-6.6%)	28 (13-48)	1400 (680-2200)
Tunisia	1995	Male	50 to 54	14% (8.6-21%)	86% (79-91%)	0.79 (0.4-1.4)	2.2% (1-3.9%)	3.1% (1.3-5.5%)	28 (11-48)	1200 (530-2000)
Tunisia	1995	Male	55 to 59	13% (7.2-21%)	87% (79-93%)	0.69 (0.32-1.3)	1.9% (0.82-3.6%)	2.5% (0.98-4.9%)	38 (14-73)	1400 (570-2500)
Tunisia	1995	Male	60 to 64	12% (6.5-20%)	88% (80-93%)	0.57 (0.25-1.1)	1.4% (0.46-2.9%)	1.8% (0.52-3.9%)	40 (12-85)	1200 (390-2500)
Tunisia	1995	Male	65 to 69	11% (6.6-19%)	89% (81-93%)	0.47 (0.22-0.84)	0.97% (0.26-2.2%)	1.2% (0.26-2.8%)	32 (6.9-74)	800 (220-1800)
Tunisia	1995	Male	70 to 74	11% (6.4-18%)	89% (82-94%)	0.37 (0.17-0.69)	0.6% (0.04-1.6%)	0.71% (0.01-1.9%)	22 (0.3-60)	440 (31-1100)
Tunisia	1995	Male	75 to 79	9.6% (5.5-16%)	90% (84-95%)	0.28 (0.12-0.55)	0.36% (-0.16-1.1%)	0.38% (-0.23-1.3%)	12 (-7.3-40)	180 (-80-590)
Tunisia	1995	Male	80 to 84	8.6% (4.8-14%)	91% (86-95%)	0.24 (0.11-0.43)	0.26% (-0.13-0.82%)	0.27% (-0.19-0.91%)	7.2 (-5.3-25)	83 (-43-270)
Tunisia	1995	Male	85 to 89	8% (4.4-13%)	92% (87-96%)	0.2 (0.093-0.37)	0.15% (-0.15-0.53%)	0.14% (-0.22-0.59%)	2.4 (-3.7-9.8)	21 (-22-77)
Tunisia	1995	Male	90 to 94	8.4% (4.7-14%)	92% (86-95%)	0.19 (0.079-0.37)	0.066% (-0.24-0.41%)	0.031% (-0.33-0.43%)	0.16 (-1.7-2.3)	2 (-7.1-13)
Tunisia	1995	Male	95 plus	9.9% (5-18%)	90% (82-95%)	0.2 (0.068-0.43)	0.033% (-0.37-0.49%)	-0.041% (-0.52-0.48%)	-0.043 (-0.55-0.52)	0.14 (-1.5-2.1)
Tunisia	2000	Female	All Ages	2% (1.5-2.7%)	98% (97-98%)	0.03 (0.022-0.039)	0.27% (0.2-0.34%)	0.11% (0.011-0.24%)	22 (2.2-47)	2900 (2100-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	2000	Female	15 to 19	2% (0.99-3.5%)	98% (97-99%)	0.026 (0.011-0.052)	0.36% (0.22-0.55%)	0.19% (0.11-0.31%)	0.39 (0.22-0.65)	210 (120-340)
Tunisia	2000	Female	20 to 24	2.6% (1.2-4.6%)	97% (95-99%)	0.044 (0.018-0.087)	0.73% (0.46-1.1%)	0.27% (0.15-0.47%)	0.63 (0.34-1.1)	430 (260-670)
Tunisia	2000	Female	25 to 29	2.7% (1.3-5.1%)	97% (95-99%)	0.047 (0.02-0.094)	0.78% (0.53-1.1%)	0.35% (0.21-0.56%)	0.83 (0.48-1.3)	450 (280-660)
Tunisia	2000	Female	30 to 34	2.5% (1-4.6%)	98% (95-99%)	0.044 (0.018-0.084)	0.69% (0.47-0.97%)	0.4% (0.25-0.59%)	1.2 (0.75-1.9)	440 (280-680)
Tunisia	2000	Female	35 to 39	2.2% (0.96-4.1%)	98% (96-99%)	0.042 (0.017-0.079)	0.55% (0.38-0.75%)	0.37% (0.25-0.55%)	1.3 (0.85-2)	340 (220-500)
Tunisia	2000	Female	40 to 44	2.2% (1.1-4%)	98% (96-99%)	0.043 (0.019-0.079)	0.44% (0.29-0.6%)	0.4% (0.26-0.58%)	1.8 (1.1-2.7)	270 (170-390)
Tunisia	2000	Female	45 to 49	2.4% (1-4%)	98% (96-99%)	0.05 (0.02-0.094)	0.36% (0.25-0.48%)	0.37% (0.22-0.58%)	2.1 (1.3-3.4)	220 (150-310)
Tunisia	2000	Female	50 to 54	2.7% (1.2-4.9%)	97% (95-99%)	0.056 (0.023-0.11)	0.31% (0.2-0.43%)	0.38% (0.22-0.61%)	2.3 (1.3-3.8)	150 (97-230)
Tunisia	2000	Female	55 to 59	2.9% (1.3-5.4%)	97% (95-99%)	0.057 (0.022-0.11)	0.24% (0.13-0.38%)	0.31% (0.13-0.53%)	2.5 (0.98-4.3)	120 (62-190)
Tunisia	2000	Female	60 to 64	2.9% (1.3-5.5%)	97% (95-99%)	0.052 (0.021-0.11)	0.16% (0.022-0.32%)	0.21% (0.0053-0.47%)	2.8 (0.071-6.2)	100 (15-200)
Tunisia	2000	Female	65 to 69	2.9% (1.3-5%)	97% (95-99%)	0.045 (0.018-0.085)	0.11% (-0.025-0.26%)	0.14% (-0.051-0.38%)	2.7 (-1-7.3)	78 (-18-180)
Tunisia	2000	Female	70 to 74	2.7% (1.3-5%)	97% (95-99%)	0.036 (0.014-0.079)	0.066% (-0.065-0.21%)	0.083% (-0.095-0.3%)	1.7 (-2-6.3)	39 (-38-120)
Tunisia	2000	Female	75 to 79	2.5% (1.1-4.6%)	98% (95-99%)	0.03 (0.011-0.065)	0.05% (-0.11-0.19%)	0.06% (-0.14-0.26%)	1.6 (-3.9-6.9)	26 (-54-100)
Tunisia	2000	Female	80 to 84	2.3% (1-4.1%)	98% (96-99%)	0.027 (0.0095-0.056)	0.033% (-0.1-0.15%)	0.035% (-0.14-0.2%)	0.8 (-3.1-4.6)	11 (-32-50)
Tunisia	2000	Female	85 to 89	2.2% (1-4.1%)	98% (96-99%)	0.025 (0.0099-0.052)	0.0091% (-0.12-0.12%)	0.0012% (-0.16-0.14%)	0.02 (-2.8-2.4)	1.5 (-20-18)
Tunisia	2000	Female	90 to 94	2.5% (1.2-4.4%)	98% (96-99%)	0.028 (0.01-0.057)	-0.016% (-0.14-0.091%)	-0.036% (-0.2-0.1%)	-0.33 (-1.8-0.93)	-0.91 (-8.5-5.4)
Tunisia	2000	Female	95 plus	3.1% (1.3-6.3%)	97% (94-99%)	0.034 (0.0099-0.08)	-0.042% (-0.23-0.087%)	-0.08% (-0.32-0.087%)	-0.22 (-0.86-0.23)	-0.47 (-2.5-0.98)
Tunisia	2000	Male	All Ages	11% (8.7-13%)	89% (87-91%)	0.43 (0.38-0.5)	1.3% (0.94-1.7%)	1.2% (0.7-1.8%)	330 (190-510)	17000 (12000-23000)
Tunisia	2000	Male	15 to 19	8.2% (4.9-13%)	92% (87-95%)	0.18 (0.084-0.32)	0.97% (0.6-1.5%)	1.2% (0.53-2.2%)	5 (2.2-9.4)	670 (400-1000)
Tunisia	2000	Male	20 to 24	15% (8.8-22%)	85% (78-91%)	0.51 (0.27-0.84)	2.5% (1.6-3.6%)	3.1% (1.5-5.4%)	17 (8.4-30)	1800 (1100-2700)
Tunisia	2000	Male	25 to 29	18% (11-26%)	82% (74-89%)	0.8 (0.46-1.3)	3.2% (2-4.7%)	4.6% (2.5-7.5%)	25 (13-43)	2200 (1400-3300)
Tunisia	2000	Male	30 to 34	19% (11-28%)	81% (72-89%)	0.9 (0.54-1.3)	3.1% (2-4.5%)	4.7% (2.5-7.5%)	28 (15-46)	2100 (1300-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	2000	Male	35 to 39	19% (11-28%)	81% (72-89%)	0.91 (0.52-1.4)	2.8% (1.7-4.3%)	4.4% (2.3-7.1%)	30 (16-50)	2100 (1200-3200)
Tunisia	2000	Male	40 to 44	17% (11-26%)	83% (74-89%)	0.87 (0.47-1.4)	2.6% (1.5-4.1%)	3.9% (1.9-6.6%)	33 (15-57)	1900 (1000-3100)
Tunisia	2000	Male	45 to 49	15% (8.5-23%)	85% (77-91%)	0.8 (0.39-1.3)	2.4% (1.2-3.9%)	3.5% (1.6-6%)	36 (16-63)	1800 (910-3000)
Tunisia	2000	Male	50 to 54	13% (8.3-20%)	87% (80-92%)	0.7 (0.35-1.2)	2% (0.9-3.4%)	2.7% (1.1-4.9%)	26 (10-46)	1100 (510-1900)
Tunisia	2000	Male	55 to 59	12% (7-20%)	88% (80-93%)	0.61 (0.29-1.2)	1.7% (0.74-3.2%)	2.2% (0.9-4.3%)	29 (11-57)	1100 (460-2000)
Tunisia	2000	Male	60 to 64	11% (6.1-19%)	89% (81-94%)	0.5 (0.22-0.95)	1.2% (0.4-2.6%)	1.6% (0.46-3.3%)	32 (8.7-69)	990 (310-2000)
Tunisia	2000	Male	65 to 69	11% (6.2-18%)	89% (82-94%)	0.41 (0.2-0.73)	0.81% (0.21-1.8%)	1% (0.22-2.3%)	32 (6.9-72)	810 (210-1800)
Tunisia	2000	Male	70 to 74	10% (6.1-17%)	90% (83-94%)	0.32 (0.15-0.6)	0.5% (0.011-1.3%)	0.58% (-0.014-1.5%)	21 (-0.5-55)	420 (9.8-1100)
Tunisia	2000	Male	75 to 79	9.1% (5.2-15%)	91% (85-95%)	0.25 (0.11-0.47)	0.29% (-0.12-0.85%)	0.32% (-0.18-0.99%)	12 (-7.1-38)	200 (-79-570)
Tunisia	2000	Male	80 to 84	8.2% (4.6-13%)	92% (87-95%)	0.21 (0.094-0.37)	0.21% (-0.11-0.63%)	0.22% (-0.17-0.7%)	5.8 (-4.4-18)	70 (-35-210)
Tunisia	2000	Male	85 to 89	7.7% (4.3-12%)	92% (88-96%)	0.18 (0.08-0.33)	0.13% (-0.14-0.44%)	0.12% (-0.22-0.47%)	1.7 (-3.2-6.8)	17 (-18-55)
Tunisia	2000	Male	90 to 94	8.1% (4.6-13%)	92% (87-95%)	0.17 (0.073-0.34)	0.057% (-0.21-0.34%)	0.016% (-0.3-0.35%)	0.11 (-2-2.3)	2.3 (-8.3-13)
Tunisia	2000	Male	95 plus	9.5% (4.8-17%)	90% (83-95%)	0.18 (0.062-0.39)	0.032% (-0.32-0.38%)	-0.05% (-0.48-0.36%)	-0.066 (-0.64-0.46)	0.18 (-1.8-2)
Tunisia	2005	Female	All Ages	2% (1.5-2.6%)	98% (97-98%)	0.033 (0.024-0.044)	0.29% (0.22-0.36%)	0.12% (0.015-0.24%)	25 (3.1-51)	3200 (2200-4300)
Tunisia	2005	Female	15 to 19	1.9% (0.93-3.3%)	98% (97-99%)	0.026 (0.011-0.051)	0.36% (0.21-0.56%)	0.2% (0.11-0.32%)	0.38 (0.21-0.63)	210 (110-350)
Tunisia	2005	Female	20 to 24	2.5% (1.1-4.3%)	98% (96-99%)	0.044 (0.017-0.087)	0.76% (0.48-1.1%)	0.28% (0.16-0.48%)	0.63 (0.35-1.1)	480 (280-760)
Tunisia	2005	Female	25 to 29	2.6% (1.2-4.8%)	97% (95-99%)	0.047 (0.02-0.094)	0.79% (0.55-1.1%)	0.35% (0.21-0.54%)	0.82 (0.48-1.3)	490 (300-730)
Tunisia	2005	Female	30 to 34	2.4% (0.99-4.5%)	98% (95-99%)	0.045 (0.018-0.086)	0.7% (0.48-0.98%)	0.41% (0.26-0.62%)	1 (0.65-1.6)	420 (260-630)
Tunisia	2005	Female	35 to 39	2.2% (0.93-3.8%)	98% (96-99%)	0.043 (0.018-0.081)	0.56% (0.4-0.75%)	0.39% (0.26-0.58%)	1.4 (0.87-2.1)	380 (250-550)
Tunisia	2005	Female	40 to 44	2.1% (1.1-3.7%)	98% (96-99%)	0.044 (0.019-0.08)	0.45% (0.31-0.62%)	0.42% (0.28-0.6%)	1.9 (1.2-3)	310 (200-450)
Tunisia	2005	Female	45 to 49	2.3% (1.1-3.8%)	98% (96-99%)	0.051 (0.023-0.1)	0.37% (0.25-0.5%)	0.39% (0.25-0.62%)	2.4 (1.4-3.9)	260 (160-360)
Tunisia	2005	Female	50 to 54	2.6% (1.2-4.7%)	97% (95-99%)	0.057 (0.024-0.11)	0.32% (0.21-0.44%)	0.4% (0.23-0.63%)	3.3 (1.7-5.6)	220 (130-330)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	2005	Female	55 to 59	2.8% (1.3-5.2%)	97% (95-99%)	0.058 (0.023-0.11)	0.25% (0.14-0.37%)	0.34% (0.16-0.55%)	3 (1.4-5.3)	150 (80-230)
Tunisia	2005	Female	60 to 64	2.9% (1.4-5.4%)	97% (95-99%)	0.053 (0.021-0.11)	0.18% (0.051-0.32%)	0.23% (0.034-0.47%)	2.8 (0.42-5.8)	110 (30-200)
Tunisia	2005	Female	65 to 69	2.8% (1.3-4.8%)	97% (95-99%)	0.046 (0.019-0.09)	0.12% (-0.0098-0.25%)	0.16% (-0.027-0.37%)	3 (-0.56-7.1)	86 (-7.2-180)
Tunisia	2005	Female	70 to 74	2.7% (1.3-4.8%)	97% (95-99%)	0.038 (0.014-0.078)	0.075% (-0.053-0.21%)	0.096% (-0.085-0.31%)	2.6 (-2.4-8.4)	58 (-41-160)
Tunisia	2005	Female	75 to 79	2.4% (1.1-4.4%)	98% (96-99%)	0.031 (0.011-0.065)	0.053% (-0.089-0.18%)	0.064% (-0.13-0.25%)	1.8 (-3.6-7)	30 (-51-100)
Tunisia	2005	Female	80 to 84	2.2% (0.97-3.9%)	98% (96-99%)	0.027 (0.0093-0.056)	0.037% (-0.089-0.15%)	0.04% (-0.12-0.19%)	1.1 (-3.3-5.1)	15 (-33-58)
Tunisia	2005	Female	85 to 89	2.1% (0.99-4%)	98% (96-99%)	0.026 (0.01-0.053)	0.011% (-0.12-0.11%)	0.0035% (-0.15-0.13%)	0.086 (-3.8-3.1)	2.6 (-26-25)
Tunisia	2005	Female	90 to 94	2.4% (1.1-4.7%)	98% (95-99%)	0.029 (0.01-0.062)	-0.016% (-0.15-0.081%)	-0.036% (-0.21-0.088%)	-0.45 (-2.6-1.1)	-1.2 (-12-6.6)
Tunisia	2005	Female	95 plus	3.2% (1.3-6.5%)	97% (93-99%)	0.036 (0.01-0.085)	-0.043% (-0.23-0.088%)	-0.082% (-0.33-0.08%)	-0.36 (-1.4-0.36)	-0.78 (-4.3-1.6)
Tunisia	2005	Male	All Ages	11% (9.1-14%)	89% (86-91%)	0.47 (0.4-0.55)	1.4% (1-1.8%)	1.3% (0.72-1.9%)	360 (200-570)	19000 (13000-26000)
Tunisia	2005	Male	15 to 19	8.3% (4.9-13%)	92% (87-95%)	0.19 (0.085-0.34)	1% (0.61-1.5%)	1.2% (0.54-2.3%)	5.3 (2.3-9.9)	700 (420-1100)
Tunisia	2005	Male	20 to 24	15% (8.7-22%)	85% (78-91%)	0.52 (0.27-0.88)	2.5% (1.6-3.8%)	3.1% (1.6-5.5%)	18 (8.9-33)	2000 (1200-3000)
Tunisia	2005	Male	25 to 29	18% (11-26%)	82% (74-89%)	0.81 (0.46-1.3)	3.2% (2.4-8%)	4.7% (2.4-7.9%)	27 (13-46)	2300 (1400-3500)
Tunisia	2005	Male	30 to 34	18% (11-27%)	82% (73-89%)	0.92 (0.54-1.4)	3.1% (1.9-4.7%)	4.9% (2.6-8%)	27 (14-45)	2100 (1200-3200)
Tunisia	2005	Male	35 to 39	18% (12-27%)	82% (73-88%)	0.92 (0.52-1.4)	2.9% (1.8-4.4%)	4.5% (2.4-7.4%)	31 (15-52)	2100 (1200-3300)
Tunisia	2005	Male	40 to 44	17% (11-26%)	83% (74-89%)	0.89 (0.48-1.4)	2.6% (1.5-4.2%)	4% (2-6.7%)	36 (17-63)	2100 (1200-3500)
Tunisia	2005	Male	45 to 49	15% (8.4-23%)	85% (77-92%)	0.81 (0.42-1.3)	2.4% (1.3-4%)	3.5% (1.6-6.2%)	41 (18-74)	2100 (1000-3600)
Tunisia	2005	Male	50 to 54	13% (8.1-20%)	87% (80-92%)	0.71 (0.36-1.2)	2% (0.91-3.4%)	2.8% (1.1-5%)	40 (15-73)	1800 (770-3100)
Tunisia	2005	Male	55 to 59	12% (6.8-20%)	88% (80-93%)	0.62 (0.3-1.2)	1.7% (0.74-3.2%)	2.3% (0.92-4.4%)	32 (12-63)	1200 (490-2200)
Tunisia	2005	Male	60 to 64	11% (6-19%)	89% (81-94%)	0.51 (0.23-0.99)	1.3% (0.4-2.7%)	1.6% (0.46-3.5%)	30 (7.8-65)	930 (280-1900)
Tunisia	2005	Male	65 to 69	11% (6.1-18%)	89% (82-94%)	0.42 (0.2-0.78)	0.84% (0.22-1.9%)	1% (0.22-2.4%)	29 (5.9-68)	740 (190-1700)
Tunisia	2005	Male	70 to 74	10% (6-17%)	90% (83-94%)	0.33 (0.15-0.62)	0.51% (0.026-1.4%)	0.6% (-0.00083-1.7%)	24 (-0.033-69)	500 (26-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	2005	Male	75 to 79	8.9% (5.1-14%)	91% (86-95%)	0.25 (0.11-0.49)	0.3% (-0.1-0.89%)	0.34% (-0.17-1%)	13 (-6.7-42)	220 (-72-650)
Tunisia	2005	Male	80 to 84	8% (4.4-13%)	92% (87-96%)	0.21 (0.091-0.37)	0.22% (-0.089-0.62%)	0.23% (-0.14-0.72%)	7.3 (-4.7-23)	90 (-36-260)
Tunisia	2005	Male	85 to 89	7.5% (4.1-12%)	93% (88-96%)	0.18 (0.077-0.34)	0.14% (-0.12-0.45%)	0.13% (-0.19-0.5%)	2.5 (-3.8-9.8)	24 (-22-79)
Tunisia	2005	Male	90 to 94	7.9% (4.5-13%)	92% (87-96%)	0.17 (0.075-0.34)	0.064% (-0.19-0.35%)	0.026% (-0.3-0.37%)	0.19 (-2.1-2.7)	2.9 (-8.6-16)
Tunisia	2005	Male	95 plus	9.3% (4.7-17%)	91% (83-95%)	0.18 (0.062-0.39)	0.039% (-0.3-0.4%)	-0.042% (-0.46-0.39%)	-0.087 (-0.95-0.83)	0.34 (-2.5-3.5)
Tunisia	2010	Female	All Ages	2.3% (1.7-2.9%)	98% (97-98%)	0.041 (0.03-0.054)	0.3% (0.23-0.37%)	0.12% (0.017-0.24%)	28 (4.2-60)	3600 (2500-4800)
Tunisia	2010	Female	15 to 19	2% (0.99-3.5%)	98% (97-99%)	0.03 (0.012-0.06)	0.37% (0.22-0.57%)	0.21% (0.12-0.35%)	0.33 (0.18-0.55)	190 (110-300)
Tunisia	2010	Female	20 to 24	2.7% (1.3-4.7%)	97% (95-99%)	0.05 (0.019-0.1)	0.77% (0.49-1.1%)	0.3% (0.17-0.52%)	0.64 (0.34-1.1)	490 (280-770)
Tunisia	2010	Female	25 to 29	2.9% (1.4-5.2%)	97% (95-99%)	0.055 (0.024-0.11)	0.81% (0.55-1.1%)	0.38% (0.22-0.59%)	0.9 (0.5-1.4)	560 (340-820)
Tunisia	2010	Female	30 to 34	2.8% (1.2-5.1%)	97% (95-99%)	0.056 (0.024-0.11)	0.72% (0.48-1%)	0.45% (0.28-0.68%)	1.2 (0.72-1.9)	500 (300-760)
Tunisia	2010	Female	35 to 39	2.7% (1.3-4.7%)	97% (95-99%)	0.058 (0.025-0.1)	0.57% (0.41-0.79%)	0.45% (0.29-0.7%)	1.4 (0.84-2.2)	380 (250-560)
Tunisia	2010	Female	40 to 44	2.7% (1.4-4.6%)	97% (95-99%)	0.059 (0.027-0.1)	0.46% (0.31-0.63%)	0.48% (0.31-0.71%)	2.2 (1.3-3.5)	350 (220-500)
Tunisia	2010	Female	45 to 49	2.7% (1.2-4.4%)	97% (96-99%)	0.062 (0.027-0.12)	0.38% (0.26-0.53%)	0.44% (0.27-0.68%)	2.8 (1.6-4.6)	300 (200-430)
Tunisia	2010	Female	50 to 54	2.8% (1.3-5%)	97% (95-99%)	0.064 (0.027-0.13)	0.32% (0.21-0.46%)	0.43% (0.25-0.69%)	3.7 (2-6.4)	260 (160-390)
Tunisia	2010	Female	55 to 59	2.9% (1.4-5.4%)	97% (95-99%)	0.064 (0.025-0.13)	0.26% (0.15-0.38%)	0.36% (0.18-0.58%)	4.1 (2-7.2)	210 (120-330)
Tunisia	2010	Female	60 to 64	3% (1.4-5.7%)	97% (94-99%)	0.059 (0.023-0.12)	0.18% (0.059-0.34%)	0.25% (0.057-0.5%)	3.3 (0.74-6.8)	120 (38-230)
Tunisia	2010	Female	65 to 69	3% (1.4-5.2%)	97% (95-99%)	0.052 (0.021-0.097)	0.12% (-0.0048-0.26%)	0.17% (-0.024-0.39%)	2.9 (-0.43-7)	83 (-3-180)
Tunisia	2010	Female	70 to 74	2.8% (1.4-5.2%)	97% (95-99%)	0.042 (0.016-0.091)	0.076% (-0.055-0.21%)	0.099% (-0.086-0.31%)	2.6 (-2.3-8.4)	59 (-42-160)
Tunisia	2010	Female	75 to 79	2.5% (1.1-4.6%)	98% (95-99%)	0.034 (0.012-0.072)	0.052% (-0.086-0.18%)	0.062% (-0.13-0.26%)	2.3 (-5-9.8)	39 (-70-140)
Tunisia	2010	Female	80 to 84	2.2% (0.99-4.1%)	98% (96-99%)	0.03 (0.01-0.062)	0.037% (-0.083-0.14%)	0.038% (-0.11-0.18%)	1.3 (-3.9-6.2)	18 (-40-69)
Tunisia	2010	Female	85 to 89	2.2% (1-4.1%)	98% (96-99%)	0.028 (0.011-0.059)	0.0091% (-0.11-0.098%)	-0.00098% (-0.16-0.12%)	-0.026 (-4.6-3.5)	2.6 (-32-29)
Tunisia	2010	Female	90 to 94	2.5% (1.2-4.8%)	97% (95-99%)	0.031 (0.011-0.065)	-0.018% (-0.15-0.075%)	-0.04% (-0.2-0.08%)	-0.73 (-3.7-1.5)	-2 (-17-9.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	2010	Female	95 plus	3.2% (1.3-6.7%)	97% (93-99%)	0.038 (0.011-0.089)	-0.044% (-0.24-0.078%)	-0.085% (-0.34-0.068%)	-0.52 (-2-0.41)	-1.1 (-6.1-2)
Tunisia	2010	Male	All Ages	11% (8.7-13%)	89% (87-91%)	0.49 (0.41-0.59)	1.5% (1.1-2%)	1.3% (0.73-2.1%)	410 (220-680)	21000 (14000-30000)
Tunisia	2010	Male	15 to 19	7.4% (4.3-12%)	93% (88-96%)	0.18 (0.077-0.34)	0.97% (0.59-1.5%)	1.2% (0.53-2.2%)	4.4 (1.9-8.6)	600 (360-930)
Tunisia	2010	Male	20 to 24	13% (7.3-20%)	87% (80-93%)	0.47 (0.24-0.81)	2.3% (1.5-3.6%)	2.8% (1.4-5.1%)	17 (7.8-31)	1900 (1200-2900)
Tunisia	2010	Male	25 to 29	16% (9.3-23%)	84% (77-91%)	0.75 (0.41-1.2)	3% (1.9-4.6%)	4.3% (2.2-7.3%)	27 (13-48)	2400 (1500-3900)
Tunisia	2010	Male	30 to 34	17% (9.5-25%)	83% (75-90%)	0.88 (0.49-1.4)	3% (1.9-4.7%)	4.7% (2.4-7.7%)	28 (14-49)	2200 (1300-3500)
Tunisia	2010	Male	35 to 39	17% (9.9-26%)	83% (74-90%)	0.92 (0.49-1.5)	2.8% (1.7-4.4%)	4.5% (2.3-7.4%)	29 (14-52)	2100 (1200-3300)
Tunisia	2010	Male	40 to 44	17% (10-26%)	83% (74-90%)	0.92 (0.47-1.5)	2.7% (1.5-4.2%)	4.1% (2-6.9%)	37 (17-64)	2200 (1200-3600)
Tunisia	2010	Male	45 to 49	15% (8.5-24%)	85% (76-92%)	0.86 (0.45-1.5)	2.5% (1.3-4.1%)	3.8% (1.8-6.4%)	47 (21-85)	2400 (1300-4100)
Tunisia	2010	Male	50 to 54	14% (8.3-21%)	86% (79-92%)	0.78 (0.41-1.3)	2.1% (0.99-3.6%)	3% (1.2-5.3%)	50 (20-91)	2200 (990-3800)
Tunisia	2010	Male	55 to 59	13% (7.1-21%)	87% (79-93%)	0.68 (0.33-1.2)	1.8% (0.79-3.4%)	2.5% (1-4.8%)	52 (19-100)	1900 (810-3600)
Tunisia	2010	Male	60 to 64	11% (6.3-20%)	89% (80-94%)	0.56 (0.24-1.1)	1.4% (0.46-2.9%)	1.8% (0.53-3.9%)	37 (10-80)	1100 (350-2400)
Tunisia	2010	Male	65 to 69	11% (6.2-18%)	89% (82-94%)	0.46 (0.21-0.82)	0.95% (0.27-2.1%)	1.2% (0.28-2.6%)	30 (7.1-72)	770 (220-1800)
Tunisia	2010	Male	70 to 74	10% (6.1-17%)	90% (83-94%)	0.36 (0.16-0.68)	0.58% (0.056-1.5%)	0.68% (0.0077-1.9%)	24 (0.34-66)	500 (44-1300)
Tunisia	2010	Male	75 to 79	9.3% (5.4-15%)	91% (85-95%)	0.28 (0.12-0.52)	0.35% (-0.084-1%)	0.38% (-0.15-1.2%)	18 (-7.4-59)	290 (-70-910)
Tunisia	2010	Male	80 to 84	8.3% (4.7-13%)	92% (87-95%)	0.23 (0.1-0.42)	0.25% (-0.082-0.69%)	0.27% (-0.14-0.8%)	9.7 (-5.9-31)	120 (-40-340)
Tunisia	2010	Male	85 to 89	7.8% (4.3-13%)	92% (87-96%)	0.2 (0.089-0.36)	0.16% (-0.14-0.52%)	0.15% (-0.22-0.58%)	4.2 (-6.1-16)	40 (-35-130)
Tunisia	2010	Male	90 to 94	8.2% (4.7-14%)	92% (86-95%)	0.19 (0.082-0.38)	0.088% (-0.2-0.45%)	0.05% (-0.29-0.48%)	0.59 (-3.7-5.8)	6.5 (-14-34)
Tunisia	2010	Male	95 plus	9.8% (4.9-18%)	90% (82-95%)	0.2 (0.068-0.44)	0.065% (-0.29-0.52%)	-0.019% (-0.46-0.51%)	-0.049 (-1.2-1.3)	0.69 (-3.1-5.5)
Tunisia	2016	Female	All Ages	2.3% (1.7-2.9%)	98% (97-98%)	0.044 (0.026-0.069)	0.31% (0.24-0.4%)	0.13% (0.022-0.26%)	34 (5.7-71)	3900 (2700-5200)
Tunisia	2016	Female	15 to 19	2% (1-3.5%)	98% (97-99%)	0.033 (0.012-0.072)	0.38% (0.22-0.58%)	0.22% (0.12-0.39%)	0.27 (0.14-0.48)	160 (87-260)
Tunisia	2016	Female	20 to 24	2.7% (1.3-4.6%)	97% (95-99%)	0.054 (0.018-0.11)	0.79% (0.5-1.1%)	0.32% (0.17-0.59%)	0.55 (0.27-1)	420 (240-660)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	2016	Female	25 to 29	2.8% (1.3-5.1%)	97% (95-99%)	0.058 (0.022-0.12)	0.82% (0.57-1.1%)	0.39% (0.23-0.65%)	0.88 (0.47-1.5)	560 (350-860)
Tunisia	2016	Female	30 to 34	2.7% (1.2-5.1%)	97% (95-99%)	0.059 (0.023-0.13)	0.74% (0.49-1%)	0.47% (0.28-0.75%)	1.3 (0.75-2.2)	570 (360-860)
Tunisia	2016	Female	35 to 39	2.7% (1.2-4.6%)	97% (95-99%)	0.061 (0.023-0.12)	0.59% (0.4-0.82%)	0.47% (0.28-0.77%)	1.6 (0.88-2.7)	460 (280-660)
Tunisia	2016	Female	40 to 44	2.6% (1.4-4.6%)	97% (95-99%)	0.062 (0.025-0.12)	0.48% (0.33-0.68%)	0.5% (0.31-0.8%)	2.2 (1.2-3.7)	360 (230-530)
Tunisia	2016	Female	45 to 49	2.6% (1.2-4.2%)	97% (96-99%)	0.065 (0.027-0.13)	0.4% (0.27-0.55%)	0.46% (0.26-0.79%)	3.1 (1.7-5.6)	350 (220-510)
Tunisia	2016	Female	50 to 54	2.7% (1.3-4.9%)	97% (95-99%)	0.068 (0.026-0.14)	0.34% (0.22-0.53%)	0.46% (0.26-0.83%)	4.4 (2.2-8.2)	320 (190-510)
Tunisia	2016	Female	55 to 59	2.9% (1.4-5.3%)	97% (95-99%)	0.068 (0.024-0.15)	0.28% (0.16-0.43%)	0.39% (0.2-0.69%)	4.9 (2.3-9.1)	250 (140-400)
Tunisia	2016	Female	60 to 64	2.9% (1.4-5.6%)	97% (94-99%)	0.063 (0.023-0.14)	0.2% (0.078-0.36%)	0.28% (0.086-0.56%)	4.8 (1.4-10)	190 (68-350)
Tunisia	2016	Female	65 to 69	2.9% (1.4-4.9%)	97% (95-99%)	0.054 (0.02-0.11)	0.14% (0.014-0.29%)	0.19% (-0.0038-0.43%)	3.7 (-0.085-8.7)	110 (11-230)
Tunisia	2016	Female	70 to 74	2.7% (1.3-4.9%)	97% (95-99%)	0.044 (0.016-0.098)	0.091% (-0.029-0.21%)	0.12% (-0.056-0.31%)	2.8 (-1.3-7.9)	65 (-21-160)
Tunisia	2016	Female	75 to 79	2.5% (1.1-4.6%)	98% (95-99%)	0.036 (0.013-0.085)	0.062% (-0.063-0.18%)	0.073% (-0.1-0.25%)	2.6 (-3.9-9.1)	46 (-52-140)
Tunisia	2016	Female	80 to 84	2.2% (1-4%)	98% (96-99%)	0.032 (0.011-0.073)	0.046% (-0.069-0.15%)	0.048% (-0.1-0.2%)	1.9 (-4-8.1)	26 (-40-89)
Tunisia	2016	Female	85 to 89	2.2% (1.1-4%)	98% (96-99%)	0.03 (0.011-0.063)	0.017% (-0.098-0.11%)	0.0069% (-0.14-0.12%)	0.28 (-5.8-5.3)	6.6 (-39-43)
Tunisia	2016	Female	90 to 94	2.5% (1.2-4.4%)	98% (96-99%)	0.033 (0.011-0.072)	-0.0074% (-0.12-0.088%)	-0.029% (-0.18-0.097%)	-0.72 (-4.4-2.4)	-1.2 (-19-14)
Tunisia	2016	Female	95 plus	3.2% (1.3-6.7%)	97% (93-99%)	0.041 (0.011-0.11)	-0.034% (-0.21-0.082%)	-0.074% (-0.31-0.073%)	-0.79 (-3.2-0.81)	-1.5 (-9.5-3.7)
Tunisia	2016	Male	All Ages	11% (9-13%)	89% (87-91%)	0.54 (0.34-0.81)	1.6% (0.94-2.4%)	1.4% (0.55-2.5%)	490 (190-930)	23000 (13000-36000)
Tunisia	2016	Male	15 to 19	7.3% (4.3-12%)	93% (88-96%)	0.19 (0.072-0.41)	1% (0.6-1.6%)	1.2% (0.52-2.6%)	4 (1.6-8.4)	530 (320-870)
Tunisia	2016	Male	20 to 24	13% (7.2-19%)	87% (81-93%)	0.5 (0.22-0.93)	2.4% (1.5-4%)	3% (1.4-6.1%)	15 (6.3-31)	1700 (970-2800)
Tunisia	2016	Male	25 to 29	16% (9.3-23%)	84% (77-91%)	0.79 (0.39-1.5)	3% (1.9-4.8%)	4.5% (2.2-8.4%)	26 (11-49)	2400 (1400-3900)
Tunisia	2016	Male	30 to 34	17% (9.6-25%)	83% (75-90%)	0.93 (0.44-1.7)	3% (1.8-4.8%)	4.8% (2.4-8.5%)	30 (14-56)	2400 (1400-4100)
Tunisia	2016	Male	35 to 39	17% (10-26%)	83% (74-90%)	0.96 (0.43-1.7)	2.9% (1.6-4.6%)	4.7% (2.1-8.4%)	31 (14-60)	2300 (1300-3900)
Tunisia	2016	Male	40 to 44	17% (10-26%)	83% (74-90%)	0.97 (0.44-1.8)	2.7% (1.4-4.4%)	4.3% (1.8-7.4%)	36 (15-65)	2200 (1100-3600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tunisia	2016	Male	45 to 49	15% (8.5-24%)	85% (76-91%)	0.93 (0.42-1.7)	2.6% (1.3-4.4%)	3.9% (1.7-7%)	49 (20-90)	2500 (1200-4500)
Tunisia	2016	Male	50 to 54	14% (8.4-21%)	86% (79-92%)	0.83 (0.39-1.6)	2.2% (0.91-3.9%)	3.2% (1.1-5.7%)	59 (20-120)	2600 (1100-4800)
Tunisia	2016	Male	55 to 59	13% (7.1-21%)	87% (79-93%)	0.73 (0.32-1.4)	1.9% (0.72-3.6%)	2.6% (0.89-5.1%)	65 (21-130)	2400 (880-4700)
Tunisia	2016	Male	60 to 64	12% (6.3-20%)	88% (80-94%)	0.59 (0.24-1.2)	1.5% (0.42-3%)	1.9% (0.51-4%)	62 (16-140)	1900 (540-4100)
Tunisia	2016	Male	65 to 69	11% (6-18%)	89% (82-94%)	0.48 (0.19-0.92)	1% (0.24-2.4%)	1.3% (0.22-3%)	41 (6.8-100)	1000 (210-2500)
Tunisia	2016	Male	70 to 74	10% (6-17%)	90% (83-94%)	0.37 (0.16-0.78)	0.66% (0.028-1.9%)	0.78% (-0.014-2.3%)	28 (-0.47-84)	570 (21-1600)
Tunisia	2016	Male	75 to 79	9.3% (5.3-15%)	91% (85-95%)	0.29 (0.12-0.61)	0.43% (-0.075-1.5%)	0.48% (-0.14-1.7%)	21 (-6.5-77)	340 (-66-1200)
Tunisia	2016	Male	80 to 84	8.3% (4.6-13%)	92% (87-95%)	0.24 (0.1-0.48)	0.31% (-0.071-1.1%)	0.34% (-0.13-1.3%)	14 (-5.4-52)	170 (-41-570)
Tunisia	2016	Male	85 to 89	7.8% (4.3-13%)	92% (87-96%)	0.21 (0.083-0.41)	0.21% (-0.11-0.78%)	0.21% (-0.19-0.87%)	8.1 (-7.3-35)	74 (-39-280)
Tunisia	2016	Male	90 to 94	8.3% (4.6-14%)	92% (86-95%)	0.2 (0.078-0.43)	0.13% (-0.2-0.62%)	0.099% (-0.28-0.66%)	1.9 (-5.7-13)	16 (-23-77)
Tunisia	2016	Male	95 plus	9.8% (5-18%)	90% (82-95%)	0.22 (0.068-0.49)	0.11% (-0.28-0.71%)	0.032% (-0.46-0.74%)	0.18 (-2.6-4.4)	2.5 (-6.7-17)
Turkey	1990	Female	All Ages	21% (17-24%)	79% (76-83%)	0.67 (0.53-0.83)	0.81% (0.43-1.3%)	0.89% (0.065-1.9%)	1400 (98-3000)	77000 (40000-120000)
Turkey	1990	Female	15 to 19	22% (14-32%)	78% (68-86%)	0.74 (0.4-1.2)	1.4% (0.81-2.2%)	2.5% (1.2-4.3%)	36 (17-64)	4900 (2800-7900)
Turkey	1990	Female	20 to 24	29% (19-40%)	71% (60-81%)	1.2 (0.67-1.8)	2.4% (1.5-3.5%)	3.8% (1.9-6.2%)	64 (32-110)	8700 (5200-13000)
Turkey	1990	Female	25 to 29	29% (20-41%)	71% (59-80%)	1 (0.6-1.6)	2.2% (1.3-3.2%)	3.4% (1.5-5.9%)	64 (28-110)	8100 (4800-12000)
Turkey	1990	Female	30 to 34	27% (17-38%)	73% (62-83%)	0.97 (0.54-1.5)	2% (1.1-3.1%)	3.5% (1.4-6.2%)	76 (31-130)	7700 (4300-12000)
Turkey	1990	Female	35 to 39	25% (16-36%)	75% (64-84%)	1.2 (0.64-1.8)	2.2% (1.3-3.5%)	4.3% (2-7.4%)	110 (49-190)	8600 (4800-14000)
Turkey	1990	Female	40 to 44	24% (16-35%)	76% (65-84%)	1.2 (0.7-1.9)	2.4% (1.3-3.9%)	4.7% (2.3-7.7%)	120 (58-210)	7700 (4100-13000)
Turkey	1990	Female	45 to 49	24% (15-35%)	76% (65-85%)	1.2 (0.67-2)	2.1% (0.93-3.9%)	3.9% (1.6-7.1%)	130 (50-230)	6800 (3000-12000)
Turkey	1990	Female	50 to 54	25% (17-35%)	75% (65-83%)	1.2 (0.67-2)	2% (0.69-3.9%)	3.5% (1.2-6.7%)	140 (46-260)	6400 (2200-12000)
Turkey	1990	Female	55 to 59	25% (16-36%)	75% (64-84%)	1.2 (0.62-2)	2% (0.38-3.9%)	3.2% (0.66-6.3%)	190 (41-380)	7300 (1400-15000)
Turkey	1990	Female	60 to 64	24% (15-36%)	76% (64-85%)	1.1 (0.55-1.8)	1.5% (-0.12-3.6%)	2.2% (-0.13-5.6%)	170 (-9.4-430)	5300 (-420-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	1990	Female	65 to 69	25% (16-36%)	75% (64-84%)	0.92 (0.49-1.5)	0.98% (-0.29-2.7%)	1.4% (-0.45-3.8%)	120 (-38-340)	3100 (-890-8800)
Turkey	1990	Female	70 to 74	24% (16-35%)	76% (65-84%)	0.76 (0.39-1.3)	0.43% (-0.82-2.1%)	0.55% (-1.2-2.8%)	55 (-110-290)	1100 (-2200-5800)
Turkey	1990	Female	75 to 79	22% (14-33%)	78% (67-86%)	0.64 (0.29-1.2)	0.2% (-0.88-1.7%)	0.25% (-1.2-2.2%)	41 (-200-380)	700 (-3100-6100)
Turkey	1990	Female	80 to 84	21% (13-30%)	79% (70-87%)	0.57 (0.27-0.97)	0.17% (-0.78-1.3%)	0.2% (-1-1.7%)	31 (-160-260)	400 (-1800-3100)
Turkey	1990	Female	85 to 89	20% (12-29%)	80% (71-88%)	0.52 (0.25-0.94)	0.1% (-0.74-1.1%)	0.081% (-1-1.4%)	9.8 (-120-170)	120 (-890-1400)
Turkey	1990	Female	90 to 94	21% (13-30%)	79% (70-87%)	0.52 (0.23-0.96)	0.064% (-0.75-1.1%)	0.0068% (-1.1-1.4%)	0.38 (-62-77)	26 (-300-440)
Turkey	1990	Female	95 plus	24% (14-37%)	76% (63-86%)	0.55 (0.21-1.1)	0.065% (-0.96-1.4%)	-0.044% (-1.4-1.6%)	-0.88 (-29-33)	6.2 (-89-120)
Turkey	1990	Male	All Ages	15% (12-18%)	85% (82-88%)	0.63 (0.49-0.79)	1.4% (0.93-1.9%)	1.5% (0.82-2.4%)	3000 (1600-4700)	150000 (100000-220000)
Turkey	1990	Male	15 to 19	12% (7.5-18%)	88% (82-92%)	0.3 (0.15-0.55)	1.4% (0.87-2.2%)	1.6% (0.79-3.1%)	50 (24-95)	6100 (3600-9800)
Turkey	1990	Male	20 to 24	19% (12-28%)	81% (72-88%)	0.78 (0.42-1.2)	3.5% (2.3-5.1%)	4% (2.1-6.9%)	150 (78-240)	15000 (9700-23000)
Turkey	1990	Male	25 to 29	22% (14-31%)	78% (69-86%)	1.2 (0.7-1.8)	4.4% (2.8-6.4%)	5.9% (3.2-9.4%)	220 (120-350)	19000 (11000-28000)
Turkey	1990	Male	30 to 34	23% (14-34%)	77% (66-86%)	1.3 (0.75-2.1)	4.1% (2.5-6.2%)	5.7% (2.9-9.1%)	220 (110-360)	17000 (9700-25000)
Turkey	1990	Male	35 to 39	24% (15-35%)	76% (65-85%)	1.4 (0.74-2.2)	4% (2.3-6.1%)	5.1% (2.5-8.3%)	240 (120-390)	17000 (9600-25000)
Turkey	1990	Male	40 to 44	25% (16-35%)	75% (65-84%)	1.4 (0.8-2.2)	4.8% (2.7-7.1%)	6.3% (2.9-9.9%)	310 (140-490)	18000 (9700-27000)
Turkey	1990	Male	45 to 49	23% (14-33%)	77% (67-86%)	1.5 (0.79-2.5)	4.3% (2.2-6.7%)	5.5% (2.6-8.9%)	360 (160-600)	18000 (9300-28000)
Turkey	1990	Male	50 to 54	22% (14-30%)	78% (70-86%)	1.4 (0.74-2.2)	3% (1.3-5.2%)	3.7% (1.4-6.5%)	340 (120-610)	15000 (6400-25000)
Turkey	1990	Male	55 to 59	20% (12-31%)	80% (69-88%)	1.1 (0.57-1.8)	2% (0.66-3.9%)	2.4% (0.63-4.9%)	330 (87-680)	12000 (4000-24000)
Turkey	1990	Male	60 to 64	18% (11-29%)	82% (71-89%)	0.85 (0.42-1.5)	1.5% (0.36-3.3%)	1.8% (0.3-4%)	260 (41-580)	7900 (1800-17000)
Turkey	1990	Male	65 to 69	18% (11-27%)	82% (73-89%)	0.76 (0.39-1.3)	1.2% (0.28-2.7%)	1.4% (0.22-3.2%)	180 (28-420)	4600 (1100-10000)
Turkey	1990	Male	70 to 74	17% (11-26%)	83% (74-89%)	0.66 (0.33-1.1)	0.94% (0.029-2.3%)	1.1% (-0.038-2.6%)	120 (-4.5-310)	2500 (74-6000)
Turkey	1990	Male	75 to 79	15% (9.3-24%)	85% (76-91%)	0.54 (0.26-1)	0.56% (-0.1-1.6%)	0.61% (-0.23-1.9%)	130 (-46-390)	2200 (-440-6300)
Turkey	1990	Male	80 to 84	14% (8.2-21%)	86% (79-92%)	0.45 (0.22-0.77)	0.47% (-0.042-1.2%)	0.5% (-0.13-1.4%)	63 (-17-180)	800 (-73-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	1990	Male	85 to 89	13% (7.7-20%)	87% (80-92%)	0.38 (0.19-0.67)	0.3% (-0.12-0.87%)	0.28% (-0.22-0.98%)	20 (-16-70)	200 (-77-600)
Turkey	1990	Male	90 to 94	14% (8.3-21%)	86% (79-92%)	0.35 (0.16-0.63)	0.19% (-0.19-0.74%)	0.12% (-0.37-0.79%)	3.1 (-10-21)	35 (-34-130)
Turkey	1990	Male	95 plus	16% (8.7-27%)	84% (73-91%)	0.36 (0.13-0.75)	0.15% (-0.32-0.87%)	-0.0035% (-0.63-0.88%)	-0.018 (-3.3-4.6)	3.7 (-7.3-21)
Turkey	1995	Female	All Ages	21% (17-24%)	79% (76-83%)	0.83 (0.67-1)	1.2% (0.73-1.8%)	1.4% (0.42-2.7%)	2300 (680-4400)	110000 (67000-170000)
Turkey	1995	Female	15 to 19	23% (15-32%)	77% (68-85%)	0.95 (0.53-1.6)	1.6% (0.96-2.5%)	3% (1.5-5%)	48 (24-80)	6100 (3600-9500)
Turkey	1995	Female	20 to 24	30% (20-40%)	70% (60-80%)	1.5 (0.87-2.3)	2.7% (1.8-3.9%)	4.6% (2.5-7.4%)	88 (47-140)	11000 (7000-16000)
Turkey	1995	Female	25 to 29	29% (20-40%)	71% (60-80%)	1.2 (0.75-1.9)	2.5% (1.6-3.5%)	4.2% (2.2-6.7%)	82 (42-130)	9800 (6100-14000)
Turkey	1995	Female	30 to 34	26% (17-36%)	74% (64-83%)	1.1 (0.64-1.8)	2.3% (1.4-3.4%)	4.1% (2.1-6.7%)	95 (48-160)	9400 (5500-14000)
Turkey	1995	Female	35 to 39	23% (15-34%)	77% (66-85%)	1.3 (0.76-2.1)	2.5% (1.5-3.8%)	4.9% (2.5-7.7%)	140 (69-230)	11000 (6200-16000)
Turkey	1995	Female	40 to 44	22% (15-32%)	78% (68-85%)	1.4 (0.79-2.1)	2.7% (1.6-4.3%)	5.3% (2.9-8.3%)	180 (97-290)	11000 (6400-17000)
Turkey	1995	Female	45 to 49	22% (14-32%)	78% (68-86%)	1.3 (0.76-2.2)	2.6% (1.3-4.4%)	4.7% (2.2-8%)	170 (79-290)	9200 (4600-15000)
Turkey	1995	Female	50 to 54	23% (16-32%)	77% (68-84%)	1.4 (0.78-2.2)	2.6% (1.2-4.7%)	4.4% (1.9-8%)	190 (86-350)	8900 (4100-16000)
Turkey	1995	Female	55 to 59	23% (15-34%)	77% (66-85%)	1.4 (0.74-2.2)	2.6% (0.95-4.8%)	4.3% (1.5-7.9%)	240 (89-450)	9300 (3500-17000)
Turkey	1995	Female	60 to 64	24% (15-36%)	76% (64-85%)	1.3 (0.66-2.2)	2.2% (0.45-4.7%)	3.3% (0.65-7.2%)	280 (58-620)	9100 (1900-20000)
Turkey	1995	Female	65 to 69	24% (16-35%)	76% (65-84%)	1.1 (0.59-1.8)	1.6% (0.24-3.6%)	2.3% (0.3-5.1%)	250 (31-560)	6500 (940-14000)
Turkey	1995	Female	70 to 74	23% (15-33%)	77% (67-85%)	0.88 (0.45-1.5)	0.95% (-0.36-2.9%)	1.3% (-0.57-3.9%)	150 (-69-470)	3200 (-1200-9700)
Turkey	1995	Female	75 to 79	22% (14-32%)	78% (68-86%)	0.74 (0.34-1.4)	0.63% (-0.62-2.5%)	0.78% (-0.87-3.2%)	140 (-150-580)	2300 (-2200-9000)
Turkey	1995	Female	80 to 84	20% (13-30%)	80% (70-87%)	0.66 (0.33-1.1)	0.58% (-0.48-2%)	0.69% (-0.69-2.6%)	130 (-130-500)	1600 (-1300-5700)
Turkey	1995	Female	85 to 89	19% (12-29%)	81% (71-88%)	0.61 (0.31-1.1)	0.48% (-0.47-1.8%)	0.54% (-0.65-2.2%)	98 (-120-400)	840 (-840-3200)
Turkey	1995	Female	90 to 94	20% (13-29%)	80% (71-87%)	0.61 (0.28-1.1)	0.41% (-0.55-1.7%)	0.44% (-0.8-2.1%)	40 (-73-190)	250 (-340-1100)
Turkey	1995	Female	95 plus	23% (13-36%)	77% (64-87%)	0.65 (0.26-1.3)	0.44% (-0.74-2.3%)	0.43% (-1.1-2.7%)	13 (-33-83)	59 (-100-310)
Turkey	1995	Male	All Ages	17% (14-20%)	83% (80-86%)	0.87 (0.7-1.1)	2.1% (1.5-2.7%)	2.3% (1.3-3.3%)	4800 (2800-7200)	230000 (160000-310000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	1995	Male	15 to 19	15% (9.3-21%)	85% (79-91%)	0.46 (0.23-0.79)	1.8% (1-2.8%)	2.2% (1.1-4%)	85 (40-160)	8900 (5200-14000)
Turkey	1995	Male	20 to 24	24% (15-32%)	76% (68-85%)	1.2 (0.66-1.8)	4.3% (2.8-6.6%)	5.3% (2.9-8.6%)	270 (150-440)	24000 (15000-36000)
Turkey	1995	Male	25 to 29	24% (17-33%)	76% (67-83%)	1.6 (0.96-2.4)	5.4% (3.3-7.9%)	7.3% (4.1-12%)	360 (200-560)	28000 (17000-41000)
Turkey	1995	Male	30 to 34	23% (16-33%)	77% (67-84%)	1.6 (0.98-2.5)	4.9% (3-7.3%)	6.9% (3.7-11%)	350 (190-540)	25000 (15000-36000)
Turkey	1995	Male	35 to 39	25% (18-35%)	75% (65-82%)	1.7 (1-2.6)	4.9% (3-7%)	6.5% (3.5-9.7%)	370 (200-570)	24000 (14000-35000)
Turkey	1995	Male	40 to 44	26% (19-36%)	74% (64-81%)	1.9 (1.1-2.9)	5.9% (3.4-8.5%)	8.1% (4-12%)	500 (250-770)	28000 (16000-40000)
Turkey	1995	Male	45 to 49	25% (16-34%)	75% (66-84%)	2 (1.1-3.2)	5.4% (3.1-7.8%)	7.1% (3.9-11%)	490 (260-740)	24000 (14000-35000)
Turkey	1995	Male	50 to 54	23% (15-32%)	77% (68-85%)	1.8 (1-2.9)	4% (2-6.2%)	5% (2.2-8%)	440 (200-710)	18000 (9200-29000)
Turkey	1995	Male	55 to 59	21% (13-32%)	79% (68-87%)	1.4 (0.72-2.3)	2.9% (1.1-5%)	3.5% (1.2-6.4%)	440 (150-810)	16000 (6200-28000)
Turkey	1995	Male	60 to 64	19% (11-30%)	81% (70-89%)	1.1 (0.52-1.9)	2.2% (0.66-4.4%)	2.6% (0.71-5.3%)	470 (120-960)	14000 (4300-28000)
Turkey	1995	Male	65 to 69	18% (11-28%)	82% (72-89%)	0.95 (0.49-1.6)	1.8% (0.56-3.7%)	2.1% (0.6-4.3%)	360 (100-760)	9100 (2800-19000)
Turkey	1995	Male	70 to 74	17% (11-27%)	83% (73-89%)	0.82 (0.41-1.5)	1.4% (0.27-3.2%)	1.6% (0.22-3.7%)	260 (36-610)	5200 (1000-12000)
Turkey	1995	Male	75 to 79	16% (9.6-24%)	84% (76-90%)	0.67 (0.32-1.3)	0.9% (0.025-2.3%)	0.99% (-0.059-2.7%)	190 (-11-510)	3100 (86-8000)
Turkey	1995	Male	80 to 84	14% (8.5-22%)	86% (78-92%)	0.55 (0.28-0.94)	0.75% (0.054-1.8%)	0.81% (-0.018-2%)	140 (-3.1-350)	1700 (130-3900)
Turkey	1995	Male	85 to 89	13% (7.7-21%)	87% (79-92%)	0.46 (0.23-0.83)	0.52% (-0.033-1.4%)	0.53% (-0.13-1.5%)	72 (-17-200)	640 (-41-1700)
Turkey	1995	Male	90 to 94	14% (8.4-22%)	86% (78-92%)	0.44 (0.2-0.8)	0.35% (-0.16-1.2%)	0.31% (-0.32-1.4%)	15 (-15-66)	110 (-50-370)
Turkey	1995	Male	95 plus	16% (8.9-28%)	84% (72-91%)	0.45 (0.16-0.91)	0.29% (-0.3-1.4%)	0.17% (-0.58-1.5%)	2.2 (-7.5-19)	16 (-17-77)
Turkey	2000	Female	All Ages	20% (17-23%)	80% (77-83%)	1 (0.82-1.2)	1.5% (1-2.2%)	2% (0.85-3.7%)	3100 (1300-5600)	130000 (83000-190000)
Turkey	2000	Female	15 to 19	22% (15-31%)	78% (69-85%)	1.1 (0.62-1.8)	1.7% (1-2.6%)	3.9% (2-6.4%)	46 (23-76)	6200 (3700-9300)
Turkey	2000	Female	20 to 24	29% (19-39%)	71% (61-81%)	1.8 (1-2.8)	2.9% (1.9-4%)	5.9% (3.3-9%)	87 (48-130)	12000 (7500-16000)
Turkey	2000	Female	25 to 29	28% (19-39%)	72% (61-81%)	1.5 (0.89-2.3)	2.6% (1.7-3.6%)	5.2% (2.9-8.1%)	82 (46-130)	11000 (6600-15000)
Turkey	2000	Female	30 to 34	25% (16-36%)	75% (64-84%)	1.3 (0.77-2.1)	2.4% (1.5-3.5%)	5.1% (2.8-7.9%)	91 (49-140)	9800 (6000-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	2000	Female	35 to 39	22% (14-33%)	78% (67-86%)	1.5 (0.87-2.5)	2.5% (1.6-3.7%)	5.7% (3.2-8.9%)	120 (67-190)	10000 (6300-15000)
Turkey	2000	Female	40 to 44	21% (14-31%)	79% (69-86%)	1.6 (0.96-2.5)	2.8% (1.7-4.2%)	6.2% (3.7-9.4%)	170 (99-260)	12000 (7200-17000)
Turkey	2000	Female	45 to 49	22% (14-30%)	78% (70-86%)	1.6 (0.91-2.6)	2.8% (1.6-4.5%)	5.7% (3.1-9.4%)	200 (110-340)	12000 (6500-19000)
Turkey	2000	Female	50 to 54	22% (15-32%)	78% (68-85%)	1.6 (0.92-2.6)	2.9% (1.5-5%)	5.5% (2.9-9.4%)	220 (110-380)	11000 (5600-18000)
Turkey	2000	Female	55 to 59	23% (15-33%)	77% (67-85%)	1.6 (0.85-2.7)	3.1% (1.4-5.3%)	5.4% (2.4-9.2%)	270 (120-470)	11000 (5100-19000)
Turkey	2000	Female	60 to 64	23% (14-34%)	77% (66-86%)	1.5 (0.77-2.6)	2.8% (0.86-5.3%)	4.3% (1.4-8.5%)	310 (98-610)	10000 (3300-20000)
Turkey	2000	Female	65 to 69	23% (15-34%)	77% (66-85%)	1.3 (0.7-2.2)	2.2% (0.71-4.4%)	3.3% (1-6.7%)	350 (110-720)	9500 (3000-19000)
Turkey	2000	Female	70 to 74	22% (15-33%)	78% (67-85%)	1.1 (0.53-1.8)	1.5% (0.05-3.8%)	2.1% (0.0083-5.3%)	280 (1.2-710)	6000 (200-15000)
Turkey	2000	Female	75 to 79	21% (13-31%)	79% (69-87%)	0.88 (0.4-1.7)	1.1% (-0.26-3.2%)	1.4% (-0.41-4.2%)	280 (-82-850)	4700 (-1200-14000)
Turkey	2000	Female	80 to 84	20% (12-29%)	80% (71-88%)	0.79 (0.38-1.4)	1% (-0.12-2.8%)	1.3% (-0.19-3.6%)	230 (-34-650)	2800 (-340-7400)
Turkey	2000	Female	85 to 89	19% (12-27%)	81% (73-88%)	0.73 (0.36-1.3)	0.92% (-0.13-2.6%)	1.1% (-0.24-3.1%)	170 (-38-500)	1400 (-210-4000)
Turkey	2000	Female	90 to 94	20% (13-28%)	80% (72-87%)	0.73 (0.32-1.4)	0.83% (-0.26-2.5%)	0.95% (-0.42-3.1%)	98 (-44-310)	570 (-170-1700)
Turkey	2000	Female	95 plus	22% (13-35%)	78% (65-87%)	0.78 (0.31-1.6)	0.9% (-0.47-3.3%)	1% (-0.71-4.1%)	36 (-25-150)	140 (-70-520)
Turkey	2000	Male	All Ages	16% (14-19%)	84% (81-86%)	1.1 (0.87-1.3)	2.5% (1.9-3.2%)	2.8% (1.8-3.9%)	5500 (3600-7700)	250000 (190000-330000)
Turkey	2000	Male	15 to 19	14% (8.9-21%)	86% (79-91%)	0.54 (0.26-0.91)	2% (1.1-3.1%)	2.7% (1.3-4.9%)	80 (36-150)	8800 (5200-14000)
Turkey	2000	Male	20 to 24	23% (14-31%)	77% (69-86%)	1.4 (0.79-2.1)	5% (3.2-7.4%)	6.9% (3.8-11%)	270 (150-440)	25000 (16000-37000)
Turkey	2000	Male	25 to 29	23% (16-31%)	77% (69-84%)	1.8 (1.1-2.7)	5.9% (3.9-8.4%)	9% (5.3-13%)	370 (210-570)	29000 (19000-43000)
Turkey	2000	Male	30 to 34	22% (15-31%)	78% (69-85%)	1.9 (1.1-2.9)	5.4% (3.6-7.9%)	8.4% (5.1-13%)	360 (210-540)	26000 (17000-38000)
Turkey	2000	Male	35 to 39	24% (17-34%)	76% (66-83%)	2 (1.2-3.1)	5.2% (3.4-7.4%)	7.7% (4.5-11%)	380 (220-570)	25000 (16000-36000)
Turkey	2000	Male	40 to 44	25% (18-34%)	75% (66-82%)	2.2 (1.3-3.3)	5.9% (3.7-8.4%)	8.7% (4.9-12%)	500 (280-730)	28000 (18000-40000)
Turkey	2000	Male	45 to 49	24% (15-33%)	76% (67-85%)	2.3 (1.3-3.7)	5.5% (3.5-7.6%)	7.6% (4.5-11%)	530 (320-770)	26000 (17000-37000)
Turkey	2000	Male	50 to 54	22% (15-30%)	78% (70-85%)	2.1 (1.2-3.4)	4.4% (2.6-6.3%)	5.8% (3.2-8.5%)	450 (250-670)	19000 (12000-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	2000	Male	55 to 59	20% (12-30%)	80% (70-88%)	1.6 (0.85-2.7)	3.4% (1.6-5.5%)	4.3% (2-7.2%)	450 (200-720)	16000 (7900-26000)
Turkey	2000	Male	60 to 64	18% (11-29%)	82% (71-89%)	1.3 (0.61-2.2)	2.6% (1-4.7%)	3.2% (1.1-5.9%)	480 (170-860)	15000 (5700-26000)
Turkey	2000	Male	65 to 69	17% (11-26%)	83% (74-89%)	1.1 (0.57-1.9)	2.2% (0.86-4%)	2.7% (0.96-4.9%)	530 (190-970)	13000 (5200-24000)
Turkey	2000	Male	70 to 74	16% (10-25%)	84% (75-90%)	0.94 (0.48-1.7)	1.8% (0.55-3.5%)	2.1% (0.59-4.2%)	400 (110-800)	8000 (2600-16000)
Turkey	2000	Male	75 to 79	15% (8.9-23%)	85% (77-91%)	0.77 (0.37-1.4)	1.3% (0.26-2.8%)	1.4% (0.23-3.2%)	370 (60-830)	5900 (1200-13000)
Turkey	2000	Male	80 to 84	13% (8.1-21%)	87% (79-92%)	0.63 (0.31-1.1)	1.1% (0.24-2.2%)	1.2% (0.2-2.6%)	190 (31-410)	2200 (490-4500)
Turkey	2000	Male	85 to 89	13% (7.2-20%)	87% (80-93%)	0.54 (0.26-0.95)	0.79% (0.14-2%)	0.85% (0.096-2.3%)	93 (10-250)	790 (130-2000)
Turkey	2000	Male	90 to 94	13% (7.9-21%)	87% (79-92%)	0.51 (0.23-0.92)	0.58% (-0.025-1.7%)	0.57% (-0.17-1.9%)	33 (-9.7-110)	210 (-9-590)
Turkey	2000	Male	95 plus	16% (8.4-26%)	84% (74-92%)	0.52 (0.18-1.1)	0.5% (-0.21-1.8%)	0.42% (-0.44-2%)	5.6 (-5.9-26)	28 (-11-100)
Turkey	2005	Female	All Ages	19% (16-22%)	81% (78-84%)	0.97 (0.78-1.2)	1.5% (1-2.1%)	1.9% (0.81-3.4%)	2600 (1100-4700)	120000 (78000-170000)
Turkey	2005	Female	15 to 19	21% (14-30%)	79% (70-86%)	1 (0.58-1.7)	1.5% (0.95-2.3%)	3.6% (1.9-6.1%)	35 (18-60)	5200 (3100-7800)
Turkey	2005	Female	20 to 24	27% (19-37%)	73% (63-81%)	1.6 (0.93-2.6)	2.6% (1.8-3.7%)	5.6% (3.1-8.6%)	67 (38-110)	10000 (6700-15000)
Turkey	2005	Female	25 to 29	27% (18-37%)	73% (63-82%)	1.3 (0.81-2.1)	2.3% (1.6-3.2%)	4.7% (2.7-7.3%)	63 (35-99)	9700 (6300-14000)
Turkey	2005	Female	30 to 34	24% (15-34%)	76% (66-85%)	1.2 (0.7-1.9)	2.1% (1.3-2.9%)	4.7% (2.7-7.2%)	72 (41-110)	9200 (5800-13000)
Turkey	2005	Female	35 to 39	21% (13-31%)	79% (69-87%)	1.4 (0.79-2.3)	2.2% (1.4-3.2%)	5.4% (3-8.4%)	95 (54-150)	9300 (5800-14000)
Turkey	2005	Female	40 to 44	20% (13-29%)	80% (71-87%)	1.5 (0.85-2.3)	2.3% (1.5-3.5%)	5.8% (3.4-8.9%)	130 (79-210)	10000 (6200-15000)
Turkey	2005	Female	45 to 49	20% (13-29%)	80% (71-87%)	1.5 (0.81-2.4)	2.4% (1.4-4%)	5.4% (3-8.9%)	170 (97-290)	11000 (6000-17000)
Turkey	2005	Female	50 to 54	21% (14-30%)	79% (70-86%)	1.5 (0.84-2.4)	2.6% (1.3-4.3%)	5.2% (2.7-8.9%)	220 (120-380)	11000 (6100-19000)
Turkey	2005	Female	55 to 59	21% (14-32%)	79% (68-86%)	1.5 (0.8-2.4)	2.7% (1.3-4.6%)	4.9% (2.3-8.4%)	250 (110-420)	10000 (4700-18000)
Turkey	2005	Female	60 to 64	21% (13-32%)	79% (68-87%)	1.4 (0.71-2.4)	2.4% (0.89-4.5%)	4% (1.5-7.6%)	270 (97-520)	9300 (3300-18000)
Turkey	2005	Female	65 to 69	22% (14-32%)	78% (68-86%)	1.2 (0.64-2)	1.9% (0.65-3.8%)	3% (0.96-6%)	270 (88-550)	7600 (2400-15000)
Turkey	2005	Female	70 to 74	21% (14-31%)	79% (69-86%)	0.97 (0.49-1.7)	1.3% (0.094-3.3%)	1.9% (0.13-4.7%)	270 (18-660)	5900 (390-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	2005	Female	75 to 79	20% (12-29%)	80% (71-88%)	0.8 (0.37-1.5)	0.95% (-0.21-2.8%)	1.3% (-0.33-3.8%)	240 (-61-750)	4100 (-910-12000)
Turkey	2005	Female	80 to 84	18% (12-27%)	82% (73-88%)	0.72 (0.35-1.2)	0.89% (-0.1-2.4%)	1.1% (-0.16-3.1%)	190 (-27-530)	2300 (-280-6100)
Turkey	2005	Female	85 to 89	18% (11-26%)	82% (74-89%)	0.66 (0.32-1.2)	0.76% (-0.14-2.2%)	0.9% (-0.23-2.7%)	130 (-35-410)	1100 (-200-3300)
Turkey	2005	Female	90 to 94	18% (12-27%)	82% (73-88%)	0.66 (0.29-1.2)	0.67% (-0.26-2.2%)	0.77% (-0.43-2.7%)	66 (-36-230)	390 (-150-1300)
Turkey	2005	Female	95 plus	21% (12-33%)	79% (67-88%)	0.71 (0.27-1.5)	0.72% (-0.45-2.8%)	0.8% (-0.71-3.5%)	30 (-27-130)	120 (-77-470)
Turkey	2005	Male	All Ages	16% (13-18%)	84% (82-87%)	1 (0.81-1.2)	2.3% (1.8-2.9%)	2.5% (1.6-3.6%)	4500 (2900-6300)	210000 (160000-270000)
Turkey	2005	Male	15 to 19	13% (8.3-20%)	87% (80-92%)	0.49 (0.24-0.83)	1.8% (1.1-2.7%)	2.5% (1.2-4.4%)	59 (29-110)	7200 (4300-11000)
Turkey	2005	Male	20 to 24	21% (13-29%)	79% (71-87%)	1.2 (0.71-1.9)	4.4% (3-6.4%)	6.3% (3.6-10%)	210 (120-340)	21000 (14000-31000)
Turkey	2005	Male	25 to 29	21% (15-29%)	79% (71-85%)	1.6 (1-2.5)	5.2% (3.5-7.4%)	8.3% (5-13%)	300 (180-460)	25000 (17000-36000)
Turkey	2005	Male	30 to 34	20% (14-28%)	80% (72-86%)	1.7 (1-2.6)	4.7% (3.2-6.8%)	7.8% (4.7-12%)	290 (170-430)	22000 (15000-31000)
Turkey	2005	Male	35 to 39	23% (16-31%)	77% (69-84%)	1.8 (1.1-2.8)	4.5% (2.9-6.3%)	7.1% (4.2-11%)	300 (170-450)	21000 (14000-29000)
Turkey	2005	Male	40 to 44	23% (16-32%)	77% (68-84%)	1.9 (1.2-2.9)	4.5% (3-6.4%)	6.9% (4.1-10%)	350 (210-520)	22000 (14000-30000)
Turkey	2005	Male	45 to 49	22% (14-30%)	78% (70-86%)	2.1 (1.2-3.3)	4.5% (2.8-6.1%)	6.4% (3.9-9.1%)	420 (240-600)	22000 (14000-30000)
Turkey	2005	Male	50 to 54	21% (14-29%)	79% (71-86%)	1.9 (1-3)	3.8% (2.2-5.5%)	5.2% (2.9-7.6%)	430 (240-640)	19000 (11000-28000)
Turkey	2005	Male	55 to 59	19% (11-29%)	81% (71-89%)	1.4 (0.75-2.4)	3% (1.5-4.7%)	3.9% (1.8-6.4%)	370 (170-600)	14000 (7000-22000)
Turkey	2005	Male	60 to 64	17% (10-27%)	83% (73-90%)	1.1 (0.54-2)	2.3% (0.92-3.9%)	2.8% (1.1-5%)	360 (140-650)	11000 (4700-20000)
Turkey	2005	Male	65 to 69	16% (10-25%)	84% (75-90%)	1 (0.52-1.8)	1.9% (0.77-3.4%)	2.3% (0.9-4.3%)	390 (150-730)	10000 (4200-19000)
Turkey	2005	Male	70 to 74	15% (9.5-24%)	85% (76-91%)	0.85 (0.42-1.5)	1.5% (0.5-3%)	1.8% (0.56-3.6%)	400 (120-800)	8200 (2700-16000)
Turkey	2005	Male	75 to 79	14% (8.1-21%)	86% (79-92%)	0.7 (0.33-1.3)	1.1% (0.27-2.5%)	1.3% (0.26-3%)	330 (65-740)	5300 (1200-11000)
Turkey	2005	Male	80 to 84	12% (7.4-19%)	88% (81-93%)	0.57 (0.28-1)	0.93% (0.21-2%)	1% (0.19-2.3%)	160 (29-370)	1900 (420-4100)
Turkey	2005	Male	85 to 89	12% (6.7-18%)	88% (82-93%)	0.49 (0.24-0.89)	0.65% (0.098-1.6%)	0.69% (0.044-1.9%)	64 (4.1-170)	550 (81-1400)
Turkey	2005	Male	90 to 94	12% (7.4-19%)	88% (81-93%)	0.46 (0.2-0.84)	0.45% (-0.063-1.3%)	0.44% (-0.19-1.5%)	18 (-7.8-62)	120 (-17-360)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	2005	Male	95 plus	15% (7.7-25%)	85% (75-92%)	0.47 (0.17-0.97)	0.37% (-0.2-1.4%)	0.28% (-0.43-1.5%)	4.2 (-6.3-22)	24 (-13-90)
Turkey	2010	Female	All Ages	19% (16-22%)	81% (78-84%)	0.76 (0.59-0.96)	1.1% (0.7-1.7%)	1.2% (0.28-2.6%)	1600 (390-3400)	89000 (54000-140000)
Turkey	2010	Female	15 to 19	19% (12-28%)	81% (72-88%)	0.74 (0.39-1.3)	1.2% (0.74-1.9%)	2.9% (1.4-5.1%)	21 (10-39)	3900 (2200-6100)
Turkey	2010	Female	20 to 24	25% (16-36%)	75% (64-84%)	1.2 (0.62-1.9)	2.1% (1.4-3.1%)	4.3% (2.1-7.1%)	37 (19-62)	7900 (4800-12000)
Turkey	2010	Female	25 to 29	26% (16-36%)	74% (64-84%)	0.98 (0.56-1.6)	1.9% (1.3-2.6%)	3.5% (1.8-5.8%)	37 (19-60)	7900 (5000-11000)
Turkey	2010	Female	30 to 34	24% (15-34%)	76% (66-85%)	0.94 (0.51-1.5)	1.7% (1.1-2.5%)	3.6% (2-5.7%)	46 (25-75)	7700 (4800-11000)
Turkey	2010	Female	35 to 39	22% (14-32%)	78% (68-86%)	1.1 (0.65-1.8)	1.8% (1.1-2.5%)	4.3% (2.4-6.8%)	69 (38-110)	8000 (5100-12000)
Turkey	2010	Female	40 to 44	21% (14-31%)	79% (69-86%)	1.2 (0.71-1.9)	1.8% (1.1-2.7%)	4.7% (2.7-7.3%)	99 (56-160)	8300 (5100-13000)
Turkey	2010	Female	45 to 49	21% (13-30%)	79% (70-87%)	1.2 (0.67-1.9)	1.9% (1-3%)	4.3% (2.3-7%)	130 (71-220)	8700 (4900-14000)
Turkey	2010	Female	50 to 54	22% (14-31%)	78% (69-86%)	1.2 (0.66-2)	1.8% (0.88-3.2%)	3.9% (1.9-6.7%)	170 (81-300)	8900 (4400-16000)
Turkey	2010	Female	55 to 59	22% (14-32%)	78% (68-86%)	1.2 (0.62-1.9)	1.8% (0.71-3.3%)	3.5% (1.4-6.4%)	210 (83-380)	8900 (3500-17000)
Turkey	2010	Female	60 to 64	21% (13-32%)	79% (68-87%)	1 (0.52-1.8)	1.5% (0.33-3.2%)	2.6% (0.64-5.4%)	190 (46-400)	6700 (1400-14000)
Turkey	2010	Female	65 to 69	21% (14-32%)	79% (68-86%)	0.91 (0.48-1.5)	1.1% (0.14-2.6%)	1.8% (0.28-4.1%)	180 (28-400)	4900 (630-11000)
Turkey	2010	Female	70 to 74	21% (14-31%)	79% (69-86%)	0.75 (0.37-1.3)	0.72% (-0.24-2.3%)	1.1% (-0.32-3.2%)	150 (-43-450)	3200 (-1100-9700)
Turkey	2010	Female	75 to 79	19% (12-29%)	81% (71-88%)	0.63 (0.29-1.2)	0.42% (-0.49-1.8%)	0.57% (-0.71-2.5%)	110 (-130-500)	1900 (-2200-8400)
Turkey	2010	Female	80 to 84	18% (11-28%)	82% (72-89%)	0.56 (0.26-0.97)	0.37% (-0.5-1.7%)	0.45% (-0.67-2.1%)	100 (-150-460)	1200 (-1600-5300)
Turkey	2010	Female	85 to 89	17% (10-25%)	83% (75-90%)	0.5 (0.24-0.9)	0.25% (-0.4-1.2%)	0.29% (-0.58-1.5%)	45 (-94-240)	430 (-700-2000)
Turkey	2010	Female	90 to 94	18% (11-26%)	82% (74-89%)	0.5 (0.23-0.94)	0.2% (-0.38-1.1%)	0.19% (-0.61-1.3%)	14 (-46-100)	120 (-220-640)
Turkey	2010	Female	95 plus	21% (12-33%)	79% (67-88%)	0.54 (0.2-1.1)	0.2% (-0.57-1.4%)	0.14% (-0.92-1.8%)	4.3 (-28-53)	32 (-89-220)
Turkey	2010	Male	All Ages	14% (12-17%)	86% (83-88%)	0.72 (0.55-0.9)	1.7% (1.2-2.3%)	1.6% (0.9-2.6%)	2900 (1500-4500)	150000 (100000-200000)
Turkey	2010	Male	15 to 19	9.9% (6.1-15%)	90% (85-94%)	0.28 (0.14-0.53)	1.3% (0.83-1.9%)	1.5% (0.76-2.8%)	26 (13-50)	4600 (2800-7100)
Turkey	2010	Male	20 to 24	16% (9.9-24%)	84% (76-90%)	0.74 (0.38-1.2)	3.2% (2.2-4.6%)	3.9% (2-6.6%)	85 (43-150)	13000 (8200-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	2010	Male	25 to 29	19% (12-27%)	81% (73-88%)	1.1 (0.64-1.8)	3.7% (2.5-5.3%)	5.8% (3.1-9.5%)	140 (77-230)	16000 (10000-23000)
Turkey	2010	Male	30 to 34	20% (12-29%)	80% (71-88%)	1.3 (0.73-2)	3.5% (2.3-5%)	5.9% (3.3-9.5%)	160 (88-260)	15000 (10000-22000)
Turkey	2010	Male	35 to 39	21% (13-30%)	79% (70-87%)	1.3 (0.72-2.1)	3.2% (2.1-4.7%)	5.3% (3-8.6%)	180 (98-290)	15000 (9700-22000)
Turkey	2010	Male	40 to 44	21% (14-31%)	79% (69-86%)	1.4 (0.75-2.1)	3.1% (1.9-4.6%)	4.9% (2.6-7.6%)	220 (110-340)	15000 (8800-22000)
Turkey	2010	Male	45 to 49	20% (12-29%)	80% (71-88%)	1.4 (0.78-2.4)	3.2% (1.8-4.8%)	4.6% (2.3-7.2%)	300 (150-480)	16000 (9400-25000)
Turkey	2010	Male	50 to 54	19% (12-27%)	81% (73-88%)	1.3 (0.7-2.1)	2.8% (1.5-4.3%)	3.8% (1.9-6%)	350 (170-570)	16000 (8500-25000)
Turkey	2010	Male	55 to 59	17% (10-26%)	83% (74-90%)	1 (0.51-1.8)	2.1% (0.99-3.6%)	2.6% (1.1-4.8%)	320 (130-580)	12000 (5800-22000)
Turkey	2010	Male	60 to 64	16% (9.2-25%)	84% (75-91%)	0.81 (0.38-1.5)	1.5% (0.57-3%)	1.9% (0.61-3.8%)	270 (89-540)	8600 (3100-17000)
Turkey	2010	Male	65 to 69	15% (9.1-23%)	85% (77-91%)	0.72 (0.36-1.3)	1.2% (0.44-2.4%)	1.4% (0.49-2.9%)	250 (82-530)	6500 (2400-13000)
Turkey	2010	Male	70 to 74	14% (8.9-22%)	86% (78-91%)	0.61 (0.3-1.2)	0.94% (0.26-2.1%)	1.1% (0.27-2.6%)	240 (57-560)	5100 (1300-12000)
Turkey	2010	Male	75 to 79	13% (7.8-20%)	87% (80-92%)	0.51 (0.24-0.97)	0.69% (0.11-1.7%)	0.8% (0.088-2.1%)	200 (21-500)	3300 (540-8100)
Turkey	2010	Male	80 to 84	12% (6.8-19%)	88% (81-93%)	0.41 (0.2-0.73)	0.5% (-0.0038-1.3%)	0.53% (-0.052-1.5%)	110 (-11-290)	1200 (-9.1-3200)
Turkey	2010	Male	85 to 89	11% (6.3-17%)	89% (83-94%)	0.35 (0.17-0.63)	0.31% (-0.037-0.89%)	0.31% (-0.13-0.98%)	31 (-12-100)	310 (-39-880)
Turkey	2010	Male	90 to 94	12% (6.7-18%)	88% (82-93%)	0.33 (0.14-0.62)	0.21% (-0.1-0.69%)	0.15% (-0.28-0.76%)	5.1 (-8.9-26)	52 (-25-170)
Turkey	2010	Male	95 plus	14% (7.1-24%)	86% (76-93%)	0.34 (0.12-0.71)	0.18% (-0.21-0.8%)	0.04% (-0.48-0.81%)	0.39 (-4.6-7.9)	8.7 (-9.9-38)
Turkey	2016	Female	All Ages	19% (16-23%)	81% (77-84%)	0.68 (0.41-1)	0.98% (0.43-1.7%)	0.74% (-0.23-2.2%)	1200 (-420-3600)	83000 (36000-150000)
Turkey	2016	Female	15 to 19	19% (12-29%)	81% (71-88%)	0.64 (0.31-1.2)	1.1% (0.66-1.7%)	2.3% (0.99-4.5%)	20 (8.2-40)	3800 (2100-6000)
Turkey	2016	Female	20 to 24	25% (16-36%)	75% (64-84%)	1 (0.46-1.8)	2% (1.3-3%)	3.5% (1.4-6.3%)	33 (13-62)	7600 (4500-11000)
Turkey	2016	Female	25 to 29	26% (17-36%)	74% (64-83%)	0.85 (0.42-1.5)	1.8% (1.3-2.6%)	2.9% (1.3-5.2%)	31 (13-57)	7700 (4900-11000)
Turkey	2016	Female	30 to 34	24% (15-35%)	76% (65-85%)	0.81 (0.39-1.5)	1.7% (1.1-2.4%)	3% (1.4-5.3%)	41 (18-77)	7800 (4800-13000)
Turkey	2016	Female	35 to 39	22% (14-33%)	78% (67-86%)	0.98 (0.49-1.8)	1.6% (1-2.5%)	3.6% (1.7-6.4%)	64 (29-120)	8400 (5100-13000)
Turkey	2016	Female	40 to 44	21% (14-31%)	79% (69-86%)	1.1 (0.54-1.7)	1.6% (0.94-2.6%)	3.9% (1.9-6.8%)	95 (45-170)	8700 (4900-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	2016	Female	45 to 49	21% (13-31%)	79% (69-87%)	1 (0.51-1.8)	1.6% (0.76-2.8%)	3.5% (1.5-6.5%)	120 (49-230)	8300 (4100-15000)
Turkey	2016	Female	50 to 54	22% (14-31%)	78% (69-86%)	1 (0.51-1.9)	1.5% (0.54-2.9%)	3.1% (1.2-6.1%)	140 (54-300)	8100 (3100-16000)
Turkey	2016	Female	55 to 59	22% (14-32%)	78% (68-86%)	1 (0.46-1.8)	1.4% (0.35-3.1%)	2.7% (0.73-5.9%)	180 (47-410)	8100 (2100-18000)
Turkey	2016	Female	60 to 64	21% (13-32%)	79% (68-87%)	0.91 (0.39-1.7)	1.1% (0.0045-2.8%)	1.9% (0.062-4.8%)	190 (5.5-480)	6700 (30-17000)
Turkey	2016	Female	65 to 69	22% (14-32%)	78% (68-86%)	0.79 (0.37-1.5)	0.76% (-0.25-2.3%)	1.3% (-0.31-3.8%)	140 (-34-430)	4000 (-1300-13000)
Turkey	2016	Female	70 to 74	21% (14-31%)	79% (69-86%)	0.65 (0.28-1.3)	0.4% (-0.49-1.9%)	0.61% (-0.68-2.8%)	91 (-100-410)	2000 (-2300-9900)
Turkey	2016	Female	75 to 79	20% (12-30%)	80% (70-88%)	0.54 (0.22-1.1)	0.16% (-0.64-1.5%)	0.21% (-0.91-2%)	41 (-180-420)	750 (-3100-7300)
Turkey	2016	Female	80 to 84	18% (11-27%)	82% (73-89%)	0.48 (0.2-0.93)	0.12% (-0.67-1.3%)	0.14% (-0.89-1.7%)	32 (-220-430)	440 (-2400-5200)
Turkey	2016	Female	85 to 89	18% (11-26%)	82% (74-89%)	0.44 (0.19-0.87)	0.048% (-0.63-1.1%)	0.02% (-0.85-1.3%)	4.7 (-220-330)	130 (-1600-2800)
Turkey	2016	Female	90 to 94	18% (12-27%)	82% (73-88%)	0.44 (0.18-0.88)	0.0085% (-0.67-1.1%)	-0.055% (-0.94-1.3%)	-9.8 (-170-240)	11 (-830-1400)
Turkey	2016	Female	95 plus	21% (12-34%)	79% (66-88%)	0.48 (0.17-1)	-0.011% (-0.92-1.5%)	-0.12% (-1.3-1.8%)	-10 (-100-140)	-4.1 (-330-540)
Turkey	2016	Male	All Ages	14% (12-17%)	86% (83-88%)	0.63 (0.39-0.95)	1.6% (0.97-2.4%)	1.3% (0.49-2.4%)	2700 (1000-5000)	150000 (88000-230000)
Turkey	2016	Male	15 to 19	10% (6.2-16%)	90% (84-94%)	0.24 (0.1-0.5)	1.2% (0.76-1.8%)	1.3% (0.56-2.6%)	29 (12-62)	5000 (3000-7800)
Turkey	2016	Male	20 to 24	16% (9.8-24%)	84% (76-90%)	0.63 (0.28-1.2)	2.9% (1.9-4.4%)	3.2% (1.5-5.9%)	95 (43-180)	13000 (8100-20000)
Turkey	2016	Male	25 to 29	19% (12-28%)	81% (72-88%)	0.94 (0.45-1.7)	3.5% (2.3-5%)	4.7% (2.3-8.2%)	150 (70-270)	17000 (10000-25000)
Turkey	2016	Male	30 to 34	20% (12-29%)	80% (71-88%)	1.1 (0.56-1.9)	3.2% (2.1-4.9%)	4.8% (2.3-8.6%)	170 (82-320)	17000 (10000-26000)
Turkey	2016	Male	35 to 39	21% (13-30%)	79% (70-87%)	1.1 (0.54-2)	2.9% (1.9-4.7%)	4.4% (2.2-8%)	190 (93-360)	16000 (9800-26000)
Turkey	2016	Male	40 to 44	21% (14-31%)	79% (69-86%)	1.2 (0.58-2)	2.8% (1.6-4.3%)	4.1% (1.8-7.1%)	220 (99-400)	16000 (8700-25000)
Turkey	2016	Male	45 to 49	20% (12-29%)	80% (71-88%)	1.2 (0.58-2.2)	2.7% (1.4-4.3%)	3.8% (1.6-6.6%)	280 (110-510)	16000 (8000-27000)
Turkey	2016	Male	50 to 54	18% (12-26%)	82% (74-88%)	1.1 (0.52-2)	2.3% (1-3.9%)	3% (1.2-5.5%)	310 (110-600)	15000 (6600-27000)
Turkey	2016	Male	55 to 59	17% (10-27%)	83% (73-90%)	0.86 (0.4-1.6)	1.6% (0.6-3.2%)	2.1% (0.62-4.2%)	280 (90-590)	11000 (4500-23000)
Turkey	2016	Male	60 to 64	16% (9-25%)	84% (75-91%)	0.68 (0.29-1.3)	1.2% (0.37-2.6%)	1.5% (0.36-3.3%)	260 (61-590)	8600 (2600-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Turkey	2016	Male	65 to 69	15% (9.1-23%)	85% (77-91%)	0.6 (0.27-1.2)	0.93% (0.21-2.2%)	1.1% (0.2-2.8%)	210 (38-530)	5700 (1400-14000)
Turkey	2016	Male	70 to 74	14% (8.8-22%)	86% (78-91%)	0.52 (0.23-1)	0.74% (0.1-1.9%)	0.88% (0.1-2.2%)	190 (21-490)	4100 (590-10000)
Turkey	2016	Male	75 to 79	13% (7.9-20%)	87% (80-92%)	0.43 (0.18-0.85)	0.52% (-0.008-1.4%)	0.59% (-0.055-1.7%)	150 (-14-450)	2500 (-40-7200)
Turkey	2016	Male	80 to 84	12% (6.8-18%)	88% (82-93%)	0.35 (0.15-0.67)	0.37% (-0.093-1.1%)	0.39% (-0.15-1.3%)	92 (-35-310)	1100 (-280-3500)
Turkey	2016	Male	85 to 89	11% (6.2-17%)	89% (83-94%)	0.3 (0.13-0.57)	0.21% (-0.14-0.78%)	0.19% (-0.23-0.87%)	36 (-43-160)	370 (-250-1400)
Turkey	2016	Male	90 to 94	12% (6.9-18%)	88% (82-93%)	0.29 (0.12-0.56)	0.11% (-0.24-0.64%)	0.043% (-0.4-0.68%)	3.9 (-36-64)	65 (-140-390)
Turkey	2016	Male	95 plus	14% (7.1-24%)	86% (76-93%)	0.3 (0.1-0.65)	0.061% (-0.4-0.75%)	-0.071% (-0.64-0.77%)	-1.9 (-17-21)	7.3 (-47-90)
United Arab Emirates	1990	Female	All Ages	7% (5.6-8.8%)	93% (91-94%)	3.3 (2.3-4.6)	1.9% (1.3-2.8%)	3.8% (2.4-5.4%)	64 (39-95)	2900 (1900-4200)
United Arab Emirates	1990	Female	15 to 19	7.8% (4.7-12%)	92% (88-95%)	3.2 (1.3-6.7)	1.9% (1.2-2.9%)	4.2% (2.5-6.8%)	1.3 (0.74-2.2)	130 (79-200)
United Arab Emirates	1990	Female	20 to 24	11% (6.5-16%)	89% (84-93%)	5.8 (2.4-11)	2.8% (1.9-4.1%)	6.3% (3.8-9.4%)	2.2 (1.3-3.6)	210 (140-310)
United Arab Emirates	1990	Female	25 to 29	11% (6.8-17%)	89% (83-93%)	6.7 (3.2-12)	3.4% (2.2-5%)	7.5% (4.6-11%)	3.8 (2.1-6.5)	320 (200-490)
United Arab Emirates	1990	Female	30 to 34	11% (6.3-17%)	89% (83-94%)	6.8 (3.6-12)	3.6% (2.2-5.4%)	7.9% (4.7-12%)	6 (3.2-10)	430 (260-710)
United Arab Emirates	1990	Female	35 to 39	11% (6.1-16%)	89% (84-94%)	6.9 (3.4-12)	3.7% (2.2-5.6%)	7.5% (4.4-11%)	6.4 (3.3-11)	400 (230-670)
United Arab Emirates	1990	Female	40 to 44	10% (6.1-16%)	90% (84-94%)	7.1 (3.2-13)	3.9% (2.3-5.8%)	7.3% (4.3-11%)	6.1 (3.2-10)	330 (190-540)
United Arab Emirates	1990	Female	45 to 49	10% (5.8-16%)	90% (84-94%)	7.4 (3-14)	3.9% (2.3-6.2%)	6.7% (3.9-10%)	5.2 (2.7-9.2)	250 (140-420)
United Arab Emirates	1990	Female	50 to 54	10% (6.4-16%)	90% (84-94%)	7.3 (3-14)	4% (2.4-6.4%)	6.3% (3.8-9.9%)	4 (2.1-6.8)	170 (92-270)
United Arab Emirates	1990	Female	55 to 59	11% (6.2-17%)	89% (83-94%)	6.9 (2.8-14)	4.3% (2.4-6.9%)	6.3% (3.6-10%)	4.8 (2.5-8.2)	170 (92-290)
United Arab Emirates	1990	Female	60 to 64	11% (6.1-17%)	89% (83-94%)	6 (2.2-12)	4% (2.2-6.6%)	5.5% (3.1-9%)	4 (2.1-7.1)	120 (64-210)
United Arab Emirates	1990	Female	65 to 69	11% (6.4-17%)	89% (83-94%)	5.1 (2.1-10)	4.1% (2.2-6.8%)	5.5% (3-9%)	5.3 (2.6-9.2)	130 (67-220)
United Arab Emirates	1990	Female	70 to 74	10% (6.2-16%)	90% (84-94%)	4.1 (1.5-9)	3.9% (2-6.8%)	5% (2.6-8.6%)	4.5 (2.2-8)	88 (45-150)
United Arab Emirates	1990	Female	75 to 79	9.5% (5.7-15%)	90% (85-94%)	3.4 (1.2-7.9)	3.5% (1.9-6.2%)	4.3% (2.2-7.5%)	5.4 (2.6-9.6)	82 (41-140)
United Arab Emirates	1990	Female	80 to 84	8.7% (5.1-14%)	91% (86-95%)	3.1 (1.1-6.5)	3.6% (2-5.8%)	4.3% (2.3-6.9%)	3.6 (1.8-5.8)	39 (20-63)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	1990	Female	85 to 89	8.4% (4.8-13%)	92% (87-95%)	2.9 (1.1-6.1)	3.5% (1.9-5.9%)	4.2% (2.3-7%)	1.2 (0.65-2.1)	9.6 (5.1-16)
United Arab Emirates	1990	Female	90 to 94	9% (5.3-14%)	91% (86-95%)	3 (1.1-6.4)	3.5% (2-5.7%)	4.1% (2.3-7%)	0.21 (0.11-0.36)	1.1 (0.62-1.9)
United Arab Emirates	1990	Female	95 plus	11% (5.6-18%)	89% (82-94%)	3.4 (1-7.8)	3.7% (1.9-6.4%)	4.4% (2.2-7.6%)	0.13 (0.068-0.23)	0.47 (0.24-0.8)
United Arab Emirates	1990	Male	All Ages	13% (10-16%)	87% (84-90%)	3.7 (2.7-5)	4.2% (2.8-6%)	6.5% (4.3-9%)	240 (150-360)	13000 (8300-19000)
United Arab Emirates	1990	Male	15 to 19	7.9% (4.8-12%)	92% (88-95%)	1.2 (0.48-2.4)	2.8% (1.6-4.7%)	4.7% (2.4-8.5%)	3.5 (1.6-6.8)	320 (170-560)
United Arab Emirates	1990	Male	20 to 24	13% (7.9-20%)	87% (80-92%)	3 (1.3-5.4)	5.5% (3.2-8.7%)	8.4% (4.3-14%)	12 (5.4-21)	980 (550-1600)
United Arab Emirates	1990	Male	25 to 29	16% (9.8-24%)	84% (76-90%)	4.7 (2.3-7.9)	6.1% (3.6-9.3%)	9.9% (5.3-16%)	20 (10-32)	1500 (870-2300)
United Arab Emirates	1990	Male	30 to 34	17% (10-26%)	83% (74-90%)	5.5 (3.1-8.8)	5.9% (3.7-8.8%)	9.7% (5.7-15%)	30 (16-51)	2100 (1300-3400)
United Arab Emirates	1990	Male	35 to 39	19% (11-28%)	81% (72-89%)	6 (3-10)	6.2% (3.9-9.4%)	10% (6.3-16%)	38 (20-61)	2400 (1400-3600)
United Arab Emirates	1990	Male	40 to 44	19% (12-29%)	81% (71-88%)	6.5 (2.9-11)	6.3% (4-9.5%)	9.9% (6-15%)	34 (19-55)	1900 (1100-2900)
United Arab Emirates	1990	Male	45 to 49	18% (9.9-27%)	82% (73-90%)	6.5 (2.7-13)	6.2% (4-9.1%)	9.1% (5.7-13%)	31 (18-48)	1500 (880-2300)
United Arab Emirates	1990	Male	50 to 54	17% (10-25%)	83% (75-90%)	6.2 (2.6-12)	5.8% (3.7-8.6%)	8.1% (5.2-12%)	21 (12-35)	890 (520-1400)
United Arab Emirates	1990	Male	55 to 59	16% (9.1-25%)	84% (75-91%)	5.8 (2.3-12)	5.6% (3.5-8.5%)	7.5% (4.6-11%)	14 (7.9-24)	510 (300-830)
United Arab Emirates	1990	Male	60 to 64	14% (8.4-24%)	86% (76-92%)	4.9 (1.8-11)	5.4% (3.4-8.4%)	6.9% (4.3-10%)	8.9 (5-15)	260 (150-440)
United Arab Emirates	1990	Male	65 to 69	14% (8.1-22%)	86% (78-92%)	4 (1.6-8.2)	5.1% (3.1-7.7%)	6.3% (3.9-9.6%)	8.7 (5-14)	210 (130-340)
United Arab Emirates	1990	Male	70 to 74	13% (8-20%)	87% (80-92%)	3 (1.1-6.8)	4.8% (2.9-7.2%)	5.8% (3.6-8.6%)	4.8 (2.7-7.8)	94 (54-150)
United Arab Emirates	1990	Male	75 to 79	12% (7-19%)	88% (81-93%)	2.4 (0.79-5.6)	3.8% (2.1-5.9%)	4.4% (2.5-6.9%)	8.3 (4.3-14)	130 (67-210)
United Arab Emirates	1990	Male	80 to 84	10% (6.1-17%)	90% (83-94%)	2 (0.75-4.3)	3.7% (2.2-5.8%)	4.3% (2.5-6.6%)	5 (2.7-8.1)	56 (32-89)
United Arab Emirates	1990	Male	85 to 89	9.8% (5.6-16%)	90% (84-94%)	1.8 (0.68-3.7)	3.5% (2-5.4%)	3.9% (2.2-6.2%)	1.5 (0.82-2.4)	12 (6.5-19)
United Arab Emirates	1990	Male	90 to 94	10% (6-17%)	90% (83-94%)	1.8 (0.58-3.9)	3.2% (1.8-5.1%)	3.6% (1.9-5.7%)	0.18 (0.096-0.3)	1 (0.54-1.6)
United Arab Emirates	1990	Male	95 plus	12% (6.4-21%)	88% (79-94%)	1.9 (0.47-4.9)	3.2% (1.4-5.4%)	3.5% (1.4-6.1%)	0.072 (0.029-0.13)	0.27 (0.12-0.46)
United Arab Emirates	1995	Female	All Ages	7.5% (6-9.5%)	92% (91-94%)	2.4 (1.7-3.3)	2.2% (1.5-3.1%)	4.1% (2.8-5.8%)	75 (48-110)	3600 (2400-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	1995	Female	15 to 19	8.1% (4.8-12%)	92% (88-95%)	2.2 (0.87-4.6)	1.8% (1.1-2.8%)	4.2% (2.4-6.8%)	1.6 (0.88-2.6)	160 (99-250)
United Arab Emirates	1995	Female	20 to 24	11% (6.6-17%)	89% (83-93%)	4 (1.7-7.7)	2.8% (1.8-4%)	6.1% (3.6-9.3%)	3.1 (1.7-4.9)	300 (190-440)
United Arab Emirates	1995	Female	25 to 29	11% (7-17%)	89% (83-93%)	4.6 (2.2-8)	3.3% (2.1-4.8%)	7.1% (4.4-11%)	5.4 (3-9.1)	460 (300-700)
United Arab Emirates	1995	Female	30 to 34	11% (6.4-17%)	89% (83-94%)	4.6 (2.5-7.7)	3.5% (2.1-5.2%)	7.5% (4.4-12%)	7.3 (3.8-13)	530 (310-850)
United Arab Emirates	1995	Female	35 to 39	11% (6.2-17%)	89% (83-94%)	4.7 (2.3-8)	3.5% (2.1-5.4%)	7.2% (4.1-11%)	8.3 (4.3-14)	520 (300-850)
United Arab Emirates	1995	Female	40 to 44	10% (6.3-17%)	90% (83-94%)	4.8 (2.1-8.5)	3.7% (2.3-5.4%)	7% (4.2-11%)	7.7 (4.2-13)	420 (250-670)
United Arab Emirates	1995	Female	45 to 49	10% (6-16%)	90% (84-94%)	5 (2.1-10)	3.8% (2.3-5.8%)	6.5% (3.9-10%)	6.4 (3.5-11)	310 (180-500)
United Arab Emirates	1995	Female	50 to 54	11% (6.4-16%)	89% (84-94%)	5 (2-10)	3.9% (2.3-6.2%)	6.1% (3.6-9.7%)	5.3 (2.9-8.7)	220 (130-360)
United Arab Emirates	1995	Female	55 to 59	11% (6.5-17%)	89% (83-93%)	4.6 (1.9-9.2)	4.1% (2.3-6.8%)	6.1% (3.5-9.9%)	4.1 (2.2-7)	150 (81-250)
United Arab Emirates	1995	Female	60 to 64	11% (6.3-18%)	89% (82-94%)	4 (1.5-8.6)	3.8% (2.1-6.6%)	5.3% (2.9-8.7%)	4.3 (2.3-7.7)	130 (70-230)
United Arab Emirates	1995	Female	65 to 69	11% (6.4-17%)	89% (83-94%)	3.4 (1.3-6.8)	3.9% (2.1-6.5%)	5.2% (2.8-8.8%)	3.9 (2-6.9)	98 (52-170)
United Arab Emirates	1995	Female	70 to 74	11% (6.4-17%)	89% (83-94%)	2.7 (1-6)	3.7% (1.9-6.4%)	4.7% (2.4-8.3%)	5.6 (2.8-9.9)	110 (58-190)
United Arab Emirates	1995	Female	75 to 79	9.7% (5.7-15%)	90% (85-94%)	2.2 (0.79-5)	3.3% (1.7-5.8%)	4% (2.1-7.2%)	4.1 (2-7.2)	63 (32-110)
United Arab Emirates	1995	Female	80 to 84	8.9% (5.2-14%)	91% (86-95%)	2 (0.75-4.2)	3.3% (1.8-5.4%)	4% (2.1-6.6%)	4.1 (2.1-6.8)	46 (24-75)
United Arab Emirates	1995	Female	85 to 89	8.6% (4.9-13%)	91% (87-95%)	1.9 (0.72-4)	3.3% (1.8-5.4%)	3.9% (2.1-6.5%)	3.3 (1.7-5.5)	25 (14-43)
United Arab Emirates	1995	Female	90 to 94	9.1% (5.4-14%)	91% (86-95%)	2 (0.71-4.2)	3.2% (1.8-5.4%)	3.8% (2.1-6.5%)	0.77 (0.41-1.3)	4.1 (2.2-6.8)
United Arab Emirates	1995	Female	95 plus	11% (5.8-19%)	89% (81-94%)	2.3 (0.66-5.2)	3.4% (1.6-5.9%)	4% (1.8-7.2%)	0.083 (0.038-0.15)	0.29 (0.14-0.51)
United Arab Emirates	1995	Male	All Ages	14% (11-17%)	86% (83-89%)	2.4 (1.7-3.3)	4.4% (3.1-6%)	6.6% (4.6-8.9%)	300 (190-440)	17000 (11000-23000)
United Arab Emirates	1995	Male	15 to 19	7.8% (4.8-12%)	92% (88-95%)	0.75 (0.28-1.5)	2.5% (1.3-4.3%)	4% (1.8-7.3%)	3.2 (1.4-6.1)	290 (150-520)
United Arab Emirates	1995	Male	20 to 24	13% (8.1-20%)	87% (80-92%)	1.8 (0.84-3.3)	5.1% (3.1-7.8%)	7.5% (4-13%)	15 (7.2-26)	1200 (700-2000)
United Arab Emirates	1995	Male	25 to 29	16% (10-25%)	84% (75-90%)	2.9 (1.4-4.8)	5.7% (3.6-8.5%)	9.1% (5.2-14%)	31 (17-51)	2400 (1500-3700)
United Arab Emirates	1995	Male	30 to 34	18% (11-26%)	82% (74-89%)	3.4 (2-5.6)	5.6% (3.5-8.4%)	9.3% (5.7-14%)	37 (20-59)	2600 (1600-3900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	1995	Male	35 to 39	19% (11-28%)	81% (72-89%)	3.7 (1.9-6.3)	5.9% (3.8-8.9%)	9.8% (6.1-15%)	45 (26-72)	2800 (1700-4300)
United Arab Emirates	1995	Male	40 to 44	19% (12-29%)	81% (71-88%)	4 (1.8-7.1)	5.9% (3.7-8.7%)	9.2% (5.7-14%)	46 (26-74)	2600 (1500-4000)
United Arab Emirates	1995	Male	45 to 49	18% (10-27%)	82% (73-90%)	4 (1.7-8)	5.8% (3.7-8.4%)	8.5% (5.3-12%)	39 (23-60)	1900 (1100-2800)
United Arab Emirates	1995	Male	50 to 54	17% (10-25%)	83% (75-90%)	3.8 (1.6-7.5)	5.4% (3.4-7.9%)	7.4% (4.6-11%)	29 (16-46)	1200 (720-1900)
United Arab Emirates	1995	Male	55 to 59	16% (9.2-25%)	84% (75-91%)	3.5 (1.4-7.4)	5.2% (3.2-7.9%)	6.8% (4.2-10%)	17 (9.8-28)	620 (360-990)
United Arab Emirates	1995	Male	60 to 64	14% (8.3-24%)	86% (76-92%)	3 (1.1-6.3)	4.9% (3-7.6%)	6.1% (3.8-9.5%)	12 (7.2-20)	370 (220-590)
United Arab Emirates	1995	Male	65 to 69	14% (8.2-22%)	86% (78-92%)	2.4 (0.98-4.8)	4.8% (2.9-7.4%)	5.8% (3.5-8.9%)	7.9 (4.6-13)	190 (120-300)
United Arab Emirates	1995	Male	70 to 74	13% (8.2-21%)	87% (79-92%)	1.9 (0.68-4.2)	4% (2.1-6.3%)	4.8% (2.4-7.5%)	8.7 (4.2-14)	170 (85-270)
United Arab Emirates	1995	Male	75 to 79	12% (7-19%)	88% (81-93%)	1.4 (0.45-3.3)	3.4% (1.4-5.6%)	3.9% (1.6-6.4%)	3.8 (1.5-6.6)	59 (25-100)
United Arab Emirates	1995	Male	80 to 84	10% (6.2-17%)	90% (83-94%)	1.2 (0.43-2.6)	3.2% (1.4-5.2%)	3.6% (1.5-5.9%)	4.5 (1.9-7.5)	51 (23-84)
United Arab Emirates	1995	Male	85 to 89	9.8% (5.6-16%)	90% (84-94%)	1.1 (0.39-2.4)	2.9% (1.3-4.8%)	3.3% (1.3-5.5%)	4.2 (1.7-7.1)	33 (14-56)
United Arab Emirates	1995	Male	90 to 94	10% (6-17%)	90% (83-94%)	1.1 (0.35-2.4)	2.7% (1-4.5%)	2.9% (1-5.1%)	0.83 (0.28-1.4)	4.6 (1.7-7.9)
United Arab Emirates	1995	Male	95 plus	12% (6.3-21%)	88% (79-94%)	1.2 (0.28-3)	2.6% (0.71-4.8%)	2.8% (0.65-5.3%)	0.061 (0.014-0.12)	0.23 (0.061-0.43)
United Arab Emirates	2000	Female	All Ages	7.2% (5.7-9%)	93% (91-94%)	1.5 (1.1-2.1)	2.1% (1.5-2.9%)	4.1% (2.8-5.7%)	78 (52-110)	3800 (2700-5200)
United Arab Emirates	2000	Female	15 to 19	7.4% (4.4-12%)	93% (88-96%)	1.4 (0.56-2.8)	1.5% (0.94-2.3%)	3.6% (2-6%)	1.6 (0.84-2.6)	170 (100-270)
United Arab Emirates	2000	Female	20 to 24	9.8% (6-15%)	90% (85-94%)	2.4 (1-4.6)	2.4% (1.6-3.5%)	5.3% (3-8.1%)	3 (1.7-4.7)	320 (200-450)
United Arab Emirates	2000	Female	25 to 29	10% (6.5-16%)	90% (84-93%)	2.8 (1.3-4.9)	2.8% (1.9-4.1%)	6.2% (3.7-9.3%)	5.4 (3.1-8.5)	490 (320-710)
United Arab Emirates	2000	Female	30 to 34	10% (6.2-16%)	90% (84-94%)	2.8 (1.6-4.7)	3.1% (1.9-4.5%)	6.7% (4-10%)	7.5 (4.3-13)	580 (370-870)
United Arab Emirates	2000	Female	35 to 39	10% (6.1-16%)	90% (84-94%)	3 (1.5-5)	3.2% (2-4.9%)	6.7% (3.9-10%)	9.1 (4.9-15)	590 (350-930)
United Arab Emirates	2000	Female	40 to 44	10% (6.4-16%)	90% (84-94%)	3.1 (1.4-5.3)	3.4% (2.2-5%)	6.5% (4-9.8%)	8.5 (4.9-14)	480 (290-740)
United Arab Emirates	2000	Female	45 to 49	9.9% (5.8-15%)	90% (85-94%)	3.1 (1.3-6)	3.5% (2.1-5.2%)	6% (3.6-9.4%)	6.3 (3.5-10)	310 (180-480)
United Arab Emirates	2000	Female	50 to 54	10% (6.5-16%)	90% (84-94%)	3.1 (1.3-6)	3.6% (2.1-5.5%)	5.7% (3.3-8.9%)	5.8 (3.3-9.5)	250 (150-400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	2000	Female	55 to 59	11% (6.2-17%)	89% (83-94%)	3 (1.2-6.1)	3.8% (2.1-6.1%)	5.7% (3.2-9.1%)	4.4 (2.4-7.3)	160 (89-260)
United Arab Emirates	2000	Female	60 to 64	11% (6.2-18%)	89% (82-94%)	2.6 (0.98-5.5)	3.4% (1.9-5.9%)	4.8% (2.7-8%)	3.9 (2.2-6.6)	120 (68-210)
United Arab Emirates	2000	Female	65 to 69	11% (6.4-17%)	89% (83-94%)	2.2 (0.86-4.5)	3.5% (1.9-5.9%)	4.8% (2.6-8%)	5.1 (2.7-8.8)	130 (69-220)
United Arab Emirates	2000	Female	70 to 74	10% (6.4-17%)	90% (83-94%)	1.8 (0.68-3.9)	3.3% (1.7-5.6%)	4.3% (2.2-7.4%)	3.8 (1.9-6.7)	78 (41-130)
United Arab Emirates	2000	Female	75 to 79	9.7% (5.7-16%)	90% (84-94%)	1.5 (0.53-3.2)	2.9% (1.4-5.2%)	3.6% (1.7-6.5%)	4.7 (2.3-8.6)	73 (36-130)
United Arab Emirates	2000	Female	80 to 84	8.9% (5.1-14%)	91% (86-95%)	1.3 (0.5-2.7)	3% (1.5-4.9%)	3.6% (1.8-6.1%)	4.2 (2.7-2)	47 (24-80)
United Arab Emirates	2000	Female	85 to 89	8.6% (5-13%)	91% (87-95%)	1.3 (0.49-2.6)	2.9% (1.5-5%)	3.5% (1.7-6%)	3.1 (1.5-5.4)	24 (12-42)
United Arab Emirates	2000	Female	90 to 94	9.1% (5.3-14%)	91% (86-95%)	1.3 (0.47-2.9)	2.9% (1.4-5%)	3.4% (1.6-6%)	1.6 (0.82-2.9)	8.8 (4.4-15)
United Arab Emirates	2000	Female	95 plus	11% (5.7-19%)	89% (81-94%)	1.5 (0.43-3.3)	3% (1.3-5.4%)	3.5% (1.4-6.5%)	0.28 (0.12-0.53)	0.98 (0.42-1.8)
United Arab Emirates	2000	Male	All Ages	14% (11-17%)	86% (83-89%)	1.8 (1.3-2.4)	4.2% (3.1-5.6%)	6.3% (4.5-8.6%)	350 (230-480)	19000 (13000-26000)
United Arab Emirates	2000	Male	15 to 19	7.3% (4.4-11%)	93% (89-96%)	0.49 (0.19-1)	1.9% (1-3.3%)	3% (1.4-5.7%)	3 (1.3-6)	290 (150-510)
United Arab Emirates	2000	Male	20 to 24	12% (7.5-19%)	88% (81-92%)	1.2 (0.53-2.3)	4.1% (2.4-6.4%)	6% (3-10%)	15 (7-27)	1300 (760-2200)
United Arab Emirates	2000	Male	25 to 29	16% (10-23%)	84% (77-90%)	2 (0.96-3.3)	4.9% (3.1-7.2%)	7.9% (4.5-12%)	32 (17-52)	2600 (1500-3900)
United Arab Emirates	2000	Male	30 to 34	18% (11-27%)	82% (73-89%)	2.4 (1.4-4)	5.1% (3.3-7.3%)	8.5% (5.3-12%)	40 (22-63)	2900 (1800-4300)
United Arab Emirates	2000	Male	35 to 39	20% (13-29%)	80% (71-87%)	2.8 (1.4-4.5)	5.5% (3.5-7.9%)	9.3% (5.9-14%)	53 (29-83)	3400 (2000-5100)
United Arab Emirates	2000	Male	40 to 44	20% (13-30%)	80% (70-87%)	3 (1.4-5.2)	5.6% (3.5-8.2%)	8.8% (5.2-13%)	56 (31-88)	3100 (1900-4800)
United Arab Emirates	2000	Male	45 to 49	19% (11-28%)	81% (72-89%)	3 (1.3-5.9)	5.6% (3.5-8.1%)	8.2% (5.1-12%)	45 (27-69)	2200 (1400-3300)
United Arab Emirates	2000	Male	50 to 54	18% (11-26%)	82% (74-89%)	2.9 (1.3-5.5)	5.1% (3.2-7.4%)	7.1% (4.4-10%)	34 (20-52)	1400 (880-2200)
United Arab Emirates	2000	Male	55 to 59	16% (9.8-26%)	84% (74-90%)	2.6 (1-5.3)	4.9% (2.8-7.6%)	6.5% (3.8-9.9%)	21 (12-34)	760 (420-1200)
United Arab Emirates	2000	Male	60 to 64	15% (8.6-25%)	85% (75-91%)	2.2 (0.8-4.5)	4.5% (2.5-7.1%)	5.7% (3.2-9%)	12 (6.5-20)	370 (200-610)
United Arab Emirates	2000	Male	65 to 69	14% (8.3-22%)	86% (78-92%)	1.7 (0.69-3.5)	4.1% (2.1-6.6%)	5% (2.6-8%)	12 (6.2-20)	300 (160-490)
United Arab Emirates	2000	Male	70 to 74	13% (8.3-21%)	87% (79-92%)	1.3 (0.51-2.8)	3.4% (1.4-5.7%)	4.1% (1.6-7%)	6.9 (2.7-12)	140 (56-240)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	2000	Male	75 to 79	12% (7-19%)	88% (81-93%)	1 (0.34-2.3)	2.6% (0.71-4.8%)	3% (0.74-5.5%)	6 (1.5-12)	94 (26-180)
United Arab Emirates	2000	Male	80 to 84	11% (6.3-17%)	89% (83-94%)	0.88 (0.32-1.9)	2.6% (0.8-4.7%)	2.9% (0.83-5.2%)	3.8 (1.2-7.2)	44 (14-81)
United Arab Emirates	2000	Male	85 to 89	10% (5.7-16%)	90% (84-94%)	0.79 (0.3-1.7)	2.3% (0.73-4.3%)	2.6% (0.72-4.7%)	2.5 (0.7-4.7)	21 (6.5-38)
United Arab Emirates	2000	Male	90 to 94	11% (6.1-17%)	89% (83-94%)	0.79 (0.26-1.8)	2.1% (0.64-4%)	2.3% (0.56-4.5%)	1.6 (0.41-3.1)	9.3 (2.8-18)
United Arab Emirates	2000	Male	95 plus	13% (6.5-21%)	87% (79-94%)	0.86 (0.2-2.2)	2.1% (0.45-4.4%)	2.2% (0.3-4.8%)	0.22 (0.03-0.48)	0.86 (0.18-1.8)
United Arab Emirates	2005	Female	All Ages	7.5% (6-9.4%)	92% (91-94%)	1.7 (1.2-2.3)	2.2% (1.6-3%)	4.4% (3.1-5.9%)	94 (65-130)	5000 (3700-6600)
United Arab Emirates	2005	Female	15 to 19	7.2% (4.3-11%)	93% (89-96%)	1.3 (0.51-2.6)	1.4% (0.89-2.1%)	3.6% (2-6%)	1.6 (0.89-2.7)	190 (120-290)
United Arab Emirates	2005	Female	20 to 24	9.5% (5.8-14%)	91% (86-94%)	2.3 (1-4.5)	2.2% (1.5-3.2%)	5.2% (3-8%)	3.7 (2.1-5.8)	420 (280-610)
United Arab Emirates	2005	Female	25 to 29	10% (6.2-15%)	90% (85-94%)	2.7 (1.2-4.7)	2.6% (1.8-3.8%)	6.1% (3.6-9.3%)	6.8 (4-11)	660 (430-940)
United Arab Emirates	2005	Female	30 to 34	10% (6.1-16%)	90% (84-94%)	2.8 (1.5-4.7)	2.8% (1.8-4.3%)	6.6% (3.9-10%)	9.5 (5.5-15)	770 (500-1100)
United Arab Emirates	2005	Female	35 to 39	10% (6.1-16%)	90% (84-94%)	2.9 (1.4-5)	3% (1.9-4.5%)	6.6% (3.9-10%)	10 (6-17)	710 (450-1100)
United Arab Emirates	2005	Female	40 to 44	10% (6.2-16%)	90% (84-94%)	3 (1.3-5.3)	3.2% (2-4.8%)	6.3% (3.9-9.6%)	11 (6.6-17)	650 (410-970)
United Arab Emirates	2005	Female	45 to 49	9.8% (5.8-15%)	90% (85-94%)	3.1 (1.3-6)	3.3% (2-5.1%)	5.9% (3.4-9.2%)	9.7 (5.5-16)	490 (290-750)
United Arab Emirates	2005	Female	50 to 54	10% (6.4-16%)	90% (84-94%)	3.1 (1.2-6.1)	3.4% (2-5.3%)	5.6% (3.2-8.7%)	7.2 (4.2-12)	320 (190-510)
United Arab Emirates	2005	Female	55 to 59	11% (6.2-17%)	89% (83-94%)	3 (1.2-5.8)	3.6% (2-6%)	5.6% (3.2-9%)	6.7 (3.9-11)	250 (140-420)
United Arab Emirates	2005	Female	60 to 64	11% (6.1-18%)	89% (82-94%)	2.6 (0.99-5.6)	3.3% (1.8-5.6%)	4.8% (2.7-7.9%)	4 (2.3-6.6)	130 (69-220)
United Arab Emirates	2005	Female	65 to 69	11% (6.4-17%)	89% (83-94%)	2.2 (0.9-4.5)	3.4% (1.8-5.8%)	4.8% (2.6-7.9%)	4.3 (2.3-7.1)	110 (60-190)
United Arab Emirates	2005	Female	70 to 74	10% (6.3-17%)	90% (83-94%)	1.8 (0.65-3.7)	3.1% (1.6-5.4%)	4.1% (2.1-7%)	5.1 (2.6-9)	110 (55-180)
United Arab Emirates	2005	Female	75 to 79	9.6% (5.7-16%)	90% (84-94%)	1.5 (0.49-3.3)	2.7% (1.3-4.9%)	3.5% (1.7-6.3%)	3.5 (1.6-6.3)	56 (26-97)
United Arab Emirates	2005	Female	80 to 84	8.8% (5.1-14%)	91% (86-95%)	1.3 (0.49-2.7)	2.8% (1.4-4.8%)	3.5% (1.7-5.9%)	4 (1.9-6.9)	46 (23-80)
United Arab Emirates	2005	Female	85 to 89	8.5% (4.9-13%)	92% (87-95%)	1.3 (0.48-2.5)	2.8% (1.4-4.9%)	3.4% (1.7-5.8%)	3.7 (1.8-6.5)	30 (15-50)
United Arab Emirates	2005	Female	90 to 94	8.9% (5.3-14%)	91% (86-95%)	1.3 (0.44-2.7)	2.8% (1.4-4.8%)	3.3% (1.5-5.9%)	1.8 (0.86-3.3)	9.9 (4.7-17)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	2005	Female	95 plus	11% (5.6-18%)	89% (82-94%)	1.5 (0.43-3.2)	2.9% (1.2-5.3%)	3.4% (1.4-6.5%)	0.7 (0.29-1.3)	2.5 (1.1-4.5)
United Arab Emirates	2005	Male	All Ages	14% (11-18%)	86% (82-89%)	1.7 (1.2-2.3)	4.3% (3.2-5.7%)	6.5% (4.8-8.8%)	490 (340-700)	28000 (20000-38000)
United Arab Emirates	2005	Male	15 to 19	7.1% (4.3-11%)	93% (89-96%)	0.45 (0.16-0.93)	1.8% (1-3%)	2.9% (1.3-5.5%)	3.9 (1.7-7.4)	410 (230-690)
United Arab Emirates	2005	Male	20 to 24	12% (7-18%)	88% (82-93%)	1.1 (0.47-2)	3.9% (2.3-6%)	5.7% (2.9-9.8%)	22 (11-39)	2000 (1200-3300)
United Arab Emirates	2005	Male	25 to 29	15% (9.5-23%)	85% (77-90%)	1.8 (0.88-3.1)	4.7% (2.9-6.9%)	7.6% (4.2-12%)	50 (27-81)	4100 (2500-6100)
United Arab Emirates	2005	Male	30 to 34	17% (11-26%)	83% (74-89%)	2.2 (1.3-3.6)	4.8% (3.2-6.9%)	8.3% (5.1-12%)	64 (37-97)	4800 (3100-6800)
United Arab Emirates	2005	Male	35 to 39	19% (12-28%)	81% (72-88%)	2.5 (1.3-4.2)	5.3% (3.4-7.6%)	9% (5.6-13%)	72 (41-110)	4700 (3000-6800)
United Arab Emirates	2005	Male	40 to 44	20% (13-30%)	80% (70-87%)	2.8 (1.3-4.8)	5.3% (3.3-7.8%)	8.6% (5-13%)	74 (40-110)	4200 (2500-6400)
United Arab Emirates	2005	Male	45 to 49	19% (10-28%)	81% (72-90%)	2.8 (1.2-5.4)	5.3% (3.3-7.8%)	7.9% (4.7-12%)	63 (36-96)	3100 (1900-4600)
United Arab Emirates	2005	Male	50 to 54	17% (11-25%)	83% (75-89%)	2.6 (1.1-5.1)	4.9% (2.9-7.2%)	6.9% (4.1-10%)	46 (27-70)	2000 (1200-3000)
United Arab Emirates	2005	Male	55 to 59	16% (9.1-26%)	84% (74-91%)	2.4 (0.91-4.9)	4.6% (2.6-7.2%)	6.2% (3.5-9.7%)	35 (20-56)	1300 (720-2000)
United Arab Emirates	2005	Male	60 to 64	14% (8.1-24%)	86% (76-92%)	2 (0.73-4.4)	4.2% (2.3-6.5%)	5.4% (3-8.6%)	19 (10-30)	590 (320-920)
United Arab Emirates	2005	Male	65 to 69	14% (8.2-21%)	86% (79-92%)	1.6 (0.64-3.3)	3.9% (2.1-6.2%)	4.9% (2.5-7.8%)	12 (6.1-19)	300 (160-480)
United Arab Emirates	2005	Male	70 to 74	13% (7.8-20%)	87% (80-92%)	1.2 (0.44-2.6)	3.1% (1.2-5.2%)	3.7% (1.4-6.3%)	10 (3.8-18)	210 (80-370)
United Arab Emirates	2005	Male	75 to 79	11% (6.6-18%)	89% (82-93%)	0.95 (0.29-2.1)	2.6% (0.82-4.7%)	3% (0.82-5.4%)	5.2 (1.4-9.5)	83 (25-150)
United Arab Emirates	2005	Male	80 to 84	10% (6-17%)	90% (83-94%)	0.81 (0.28-1.7)	2.5% (0.88-4.5%)	2.8% (0.91-5.1%)	4.4 (1.5-8)	51 (18-92)
United Arab Emirates	2005	Male	85 to 89	9.6% (5.4-16%)	90% (84-95%)	0.71 (0.26-1.5)	2.3% (0.77-4.3%)	2.5% (0.8-4.8%)	2.8 (0.85-5.2)	23 (7.9-42)
United Arab Emirates	2005	Male	90 to 94	10% (5.9-16%)	90% (84-94%)	0.71 (0.24-1.6)	2.1% (0.68-4.2%)	2.3% (0.63-4.7%)	1.2 (0.33-2.4)	6.9 (2.2-13)
United Arab Emirates	2005	Male	95 plus	12% (6.5-21%)	88% (79-94%)	0.79 (0.19-2)	2.2% (0.53-4.5%)	2.3% (0.44-4.9%)	0.55 (0.11-1.2)	2.2 (0.54-4.5)
United Arab Emirates	2010	Female	All Ages	8% (6.3-10%)	92% (90-94%)	1.9 (1.2-2.8)	2.2% (1.6-2.9%)	4.4% (3.1-6%)	140 (95-200)	8100 (5600-11000)
United Arab Emirates	2010	Female	15 to 19	7.6% (4.5-12%)	92% (88-95%)	1.4 (0.57-2.9)	1.4% (0.9-2.2%)	3.7% (2-6.1%)	2.7 (1.5-4.7)	320 (190-500)
United Arab Emirates	2010	Female	20 to 24	10% (6.1-16%)	90% (84-94%)	2.5 (1-5)	2.2% (1.5-3.3%)	5.2% (3-8.1%)	7 (3.9-11)	810 (540-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	2010	Female	25 to 29	11% (6.4-16%)	89% (84-94%)	2.9 (1.3-5.3)	2.6% (1.7-3.8%)	6% (3.6-9.3%)	11 (6.2-18)	1100 (730-1600)
United Arab Emirates	2010	Female	30 to 34	10% (6.1-16%)	90% (84-94%)	2.9 (1.6-5.1)	2.8% (1.8-4.2%)	6.5% (3.8-10%)	14 (7.8-24)	1200 (760-1800)
United Arab Emirates	2010	Female	35 to 39	10% (5.9-16%)	90% (84-94%)	3 (1.4-5.2)	2.9% (1.8-4.5%)	6.4% (3.7-9.9%)	18 (9.4-30)	1200 (750-1900)
United Arab Emirates	2010	Female	40 to 44	10% (6.1-16%)	90% (84-94%)	3.1 (1.3-5.4)	3.1% (1.8-4.6%)	6.2% (3.7-9.7%)	18 (10-30)	1100 (630-1700)
United Arab Emirates	2010	Female	45 to 49	10% (5.7-15%)	90% (85-94%)	3.2 (1.3-6.4)	3.2% (1.9-5%)	5.9% (3.4-9.2%)	16 (8.7-26)	830 (480-1300)
United Arab Emirates	2010	Female	50 to 54	10% (6.3-16%)	90% (84-94%)	3.3 (1.3-6.4)	3.3% (1.9-5.3%)	5.5% (3.2-8.7%)	12 (7-21)	550 (310-900)
United Arab Emirates	2010	Female	55 to 59	10% (6.1-16%)	90% (84-94%)	3 (1.2-6.1)	3.5% (1.9-5.7%)	5.5% (3.1-8.8%)	8.7 (4.8-14)	330 (180-550)
United Arab Emirates	2010	Female	60 to 64	11% (6.1-17%)	89% (83-94%)	2.7 (0.98-5.6)	3.2% (1.8-5.4%)	4.7% (2.6-7.6%)	7 (3.9-12)	230 (120-380)
United Arab Emirates	2010	Female	65 to 69	11% (6.3-17%)	89% (83-94%)	2.3 (0.85-4.6)	3.3% (1.7-5.5%)	4.6% (2.6-7.6%)	4.8 (2.6-8)	130 (68-220)
United Arab Emirates	2010	Female	70 to 74	10% (6-16%)	90% (84-94%)	1.8 (0.63-3.9)	3% (1.5-5.3%)	4% (2-6.9%)	4.9 (2.5-8.8)	100 (52-180)
United Arab Emirates	2010	Female	75 to 79	9.3% (5.4-15%)	91% (85-95%)	1.5 (0.46-3.4)	2.7% (1.3-4.8%)	3.4% (1.5-6.1%)	5.5 (2.5-10)	88 (41-160)
United Arab Emirates	2010	Female	80 to 84	8.6% (5.1-14%)	91% (86-95%)	1.3 (0.46-2.9)	2.7% (1.4-4.7%)	3.3% (1.7-5.8%)	4.9 (2.5-8.3)	56 (29-96)
United Arab Emirates	2010	Female	85 to 89	8.2% (4.8-13%)	92% (87-95%)	1.3 (0.48-2.7)	2.7% (1.4-4.7%)	3.2% (1.6-5.7%)	3.8 (1.8-6.7)	30 (15-52)
United Arab Emirates	2010	Female	90 to 94	8.8% (5.2-14%)	91% (86-95%)	1.3 (0.44-2.8)	2.7% (1.3-4.8%)	3.2% (1.5-5.7%)	2.5 (1.1-4.4)	13 (6.3-23)
United Arab Emirates	2010	Female	95 plus	10% (5.5-18%)	90% (82-95%)	1.5 (0.44-3.5)	2.9% (1.3-5.2%)	3.4% (1.4-6.3%)	0.81 (0.33-1.5)	2.9 (1.2-5.2)
United Arab Emirates	2010	Male	All Ages	15% (12-19%)	85% (81-88%)	1.9 (1.3-2.7)	4.4% (3.1-6.1%)	6.7% (4.8-9.3%)	1000 (660-1500)	58000 (41000-83000)
United Arab Emirates	2010	Male	15 to 19	7.6% (4.7-12%)	92% (88-95%)	0.51 (0.19-1)	2.1% (1.2-3.5%)	3.5% (1.6-6.8%)	7.3 (3.1-15)	720 (390-1300)
United Arab Emirates	2010	Male	20 to 24	13% (7.6-19%)	87% (81-92%)	1.2 (0.52-2.2)	4.1% (2.4-6.6%)	6.2% (3-11%)	49 (23-90)	4500 (2500-7400)
United Arab Emirates	2010	Male	25 to 29	16% (9.6-24%)	84% (76-90%)	1.9 (0.92-3.3)	4.9% (3-7.3%)	8% (4.3-12%)	99 (51-170)	8100 (4800-13000)
United Arab Emirates	2010	Male	30 to 34	17% (10-26%)	83% (74-90%)	2.3 (1.3-3.8)	4.8% (3.1-7.2%)	8.2% (4.8-13%)	130 (77-220)	10000 (6200-15000)
United Arab Emirates	2010	Male	35 to 39	18% (11-27%)	82% (73-89%)	2.5 (1.2-4.3)	5.1% (3.2-7.7%)	8.7% (5.2-13%)	170 (91-270)	11000 (6500-17000)
United Arab Emirates	2010	Male	40 to 44	18% (12-27%)	82% (73-88%)	2.7 (1.2-4.7)	5.2% (3.2-7.9%)	8.3% (4.7-13%)	150 (80-240)	8400 (4900-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	2010	Male	45 to 49	17% (9.8-26%)	83% (74-90%)	2.7 (1.2-5.4)	5.1% (3.2-7.6%)	7.6% (4.5-12%)	140 (80-220)	6900 (4200-11000)
United Arab Emirates	2010	Male	50 to 54	16% (10-24%)	84% (76-90%)	2.6 (1-5.2)	4.7% (2.7-7%)	6.6% (3.8-9.9%)	98 (55-160)	4200 (2400-6400)
United Arab Emirates	2010	Male	55 to 59	15% (8.9-24%)	85% (76-91%)	2.4 (0.9-5)	4.6% (2.6-7.2%)	6.1% (3.5-9.5%)	61 (33-99)	2200 (1200-3600)
United Arab Emirates	2010	Male	60 to 64	14% (7.9-24%)	86% (76-92%)	2 (0.71-4.3)	4.1% (2.3-6.5%)	5.3% (2.9-8.4%)	41 (22-69)	1300 (680-2000)
United Arab Emirates	2010	Male	65 to 69	13% (7.7-21%)	87% (79-92%)	1.6 (0.63-3.3)	3.9% (2-6.2%)	4.9% (2.5-7.8%)	24 (12-39)	590 (310-970)
United Arab Emirates	2010	Male	70 to 74	12% (7.7-19%)	88% (81-92%)	1.2 (0.43-2.7)	3.3% (1.4-5.5%)	4% (1.7-6.8%)	13 (5.7-23)	270 (110-460)
United Arab Emirates	2010	Male	75 to 79	11% (6.6-18%)	89% (82-93%)	0.95 (0.3-2.2)	2.7% (0.9-4.9%)	3.1% (0.96-5.6%)	11 (3.3-20)	170 (58-310)
United Arab Emirates	2010	Male	80 to 84	10% (5.8-17%)	90% (83-94%)	0.81 (0.27-1.8)	2.7% (1.1-4.7%)	3% (1.2-5.4%)	7.8 (3-14)	90 (36-160)
United Arab Emirates	2010	Male	85 to 89	9.5% (5.4-15%)	91% (85-95%)	0.72 (0.26-1.6)	2.5% (1-4.4%)	2.8% (1-5%)	3.7 (1.4-6.6)	30 (12-53)
United Arab Emirates	2010	Male	90 to 94	10% (5.8-16%)	90% (84-94%)	0.72 (0.23-1.6)	2.4% (0.85-4.5%)	2.6% (0.85-5%)	1.7 (0.54-3.1)	9.4 (3.2-17)
United Arab Emirates	2010	Male	95 plus	12% (6.2-20%)	88% (80-94%)	0.79 (0.18-2)	2.5% (0.79-4.9%)	2.6% (0.71-5.4%)	0.43 (0.12-0.89)	1.7 (0.55-3.3)
United Arab Emirates	2016	Female	All Ages	8.2% (6.6-10%)	92% (90-93%)	2.1 (1.1-3.5)	2.3% (1.7-3.1%)	4.6% (3.2-6.3%)	200 (130-290)	11000 (7300-15000)
United Arab Emirates	2016	Female	15 to 19	7.8% (4.7-12%)	92% (88-95%)	1.5 (0.53-3.2)	1.4% (0.88-2.1%)	3.6% (2-5.9%)	2.2 (1.2-3.8)	260 (160-400)
United Arab Emirates	2016	Female	20 to 24	10% (6.3-16%)	90% (84-94%)	2.7 (0.97-5.8)	2.2% (1.5-3.2%)	5.1% (2.9-7.9%)	5.5 (3-9.1)	650 (420-940)
United Arab Emirates	2016	Female	25 to 29	11% (6.5-17%)	89% (83-93%)	3.1 (1.2-6.2)	2.5% (1.7-3.6%)	5.8% (3.4-8.9%)	11 (5.9-19)	1100 (740-1700)
United Arab Emirates	2016	Female	30 to 34	10% (6.1-17%)	90% (83-94%)	3.1 (1.4-6)	2.7% (1.8-4%)	6.2% (3.7-9.7%)	18 (9.5-30)	1500 (940-2300)
United Arab Emirates	2016	Female	35 to 39	10% (6-16%)	90% (84-94%)	3.1 (1.3-6)	2.8% (1.7-4.4%)	6.3% (3.6-9.5%)	23 (12-39)	1600 (940-2500)
United Arab Emirates	2016	Female	40 to 44	10% (6.1-16%)	90% (84-94%)	3.2 (1.3-6.1)	3% (1.8-4.5%)	6.1% (3.7-9.3%)	26 (14-43)	1500 (910-2400)
United Arab Emirates	2016	Female	45 to 49	10% (5.9-15%)	90% (85-94%)	3.4 (1.4-7.1)	3.2% (1.9-4.9%)	5.8% (3.4-9.1%)	26 (14-43)	1300 (750-2100)
United Arab Emirates	2016	Female	50 to 54	10% (6.3-16%)	90% (84-94%)	3.4 (1.3-7.2)	3.2% (1.9-5.1%)	5.4% (3.1-8.6%)	23 (13-39)	1000 (560-1700)
United Arab Emirates	2016	Female	55 to 59	11% (6.3-17%)	89% (83-94%)	3.2 (1.1-6.8)	3.4% (1.9-5.6%)	5.3% (3.1-8.6%)	18 (9.9-30)	670 (370-1100)
United Arab Emirates	2016	Female	60 to 64	11% (6-18%)	89% (82-94%)	2.7 (0.94-6.3)	3% (1.7-5.2%)	4.5% (2.5-7.3%)	10 (5.6-18)	340 (180-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	2016	Female	65 to 69	11% (6.5-16%)	89% (84-94%)	2.3 (0.83-5.2)	3.1% (1.7-5.3%)	4.4% (2.4-7.3%)	9.4 (5-16)	250 (130-430)
United Arab Emirates	2016	Female	70 to 74	10% (6.4-16%)	90% (84-94%)	1.9 (0.62-4)	2.8% (1.5-4.9%)	3.8% (1.9-6.5%)	6.2 (3.1-11)	130 (65-230)
United Arab Emirates	2016	Female	75 to 79	9.4% (5.6-15%)	91% (85-94%)	1.5 (0.48-3.6)	2.6% (1.2-4.7%)	3.3% (1.5-5.9%)	5.3 (2.5-9.8)	85 (40-150)
United Arab Emirates	2016	Female	80 to 84	8.7% (5-14%)	91% (86-95%)	1.4 (0.47-3.1)	2.6% (1.4-4.4%)	3.2% (1.7-5.4%)	6.2 (3.1-11)	71 (37-120)
United Arab Emirates	2016	Female	85 to 89	8.4% (4.9-13%)	92% (87-95%)	1.3 (0.46-2.9)	2.6% (1.3-4.4%)	3.1% (1.5-5.4%)	5.4 (2.7-9.7)	43 (22-74)
United Arab Emirates	2016	Female	90 to 94	9% (5.2-14%)	91% (86-95%)	1.4 (0.43-3.3)	2.6% (1.3-4.4%)	3% (1.4-5.3%)	2.6 (1.2-4.7)	14 (6.7-24)
United Arab Emirates	2016	Female	95 plus	11% (5.6-18%)	89% (82-94%)	1.6 (0.41-4)	2.7% (1.1-4.9%)	3.2% (1.2-5.9%)	1.2 (0.48-2.2)	4.4 (1.9-7.9)
United Arab Emirates	2016	Male	All Ages	15% (12-18%)	85% (82-88%)	1.9 (1.1-3.1)	4.3% (3-6%)	6.2% (4.1-8.7%)	1300 (760-2000)	68000 (45000-98000)
United Arab Emirates	2016	Male	15 to 19	7.5% (4.6-12%)	92% (88-95%)	0.49 (0.17-1.1)	2% (1.1-3.4%)	3.5% (1.6-6.8%)	6.7 (2.8-14)	670 (350-1200)
United Arab Emirates	2016	Male	20 to 24	13% (7.6-19%)	87% (81-92%)	1.2 (0.44-2.6)	4.1% (2.4-6.5%)	6.3% (2.9-11%)	36 (16-67)	3300 (1800-5300)
United Arab Emirates	2016	Male	25 to 29	16% (9.5-23%)	84% (77-91%)	1.9 (0.75-3.8)	4.7% (2.8-7.1%)	7.7% (3.9-12%)	82 (39-140)	6800 (3800-11000)
United Arab Emirates	2016	Male	30 to 34	17% (10-26%)	83% (74-90%)	2.3 (1.1-4.3)	4.6% (2.9-7.1%)	7.8% (4.3-12%)	110 (56-190)	8400 (4900-13000)
United Arab Emirates	2016	Male	35 to 39	18% (11-27%)	82% (73-89%)	2.4 (1.1-4.6)	4.9% (3-7.6%)	8.2% (4.7-13%)	160 (82-270)	10000 (6100-16000)
United Arab Emirates	2016	Male	40 to 44	19% (12-28%)	81% (72-88%)	2.6 (1.1-4.9)	5% (3-7.5%)	7.9% (4.4-12%)	210 (110-340)	12000 (6500-18000)
United Arab Emirates	2016	Male	45 to 49	18% (9.7-27%)	82% (73-90%)	2.6 (1.1-5.5)	5% (3-7.6%)	7.3% (4.2-11%)	200 (110-320)	9800 (5500-15000)
United Arab Emirates	2016	Male	50 to 54	16% (9.9-24%)	84% (76-90%)	2.5 (0.95-5.4)	4.5% (2.6-6.9%)	6.3% (3.5-9.6%)	180 (89-300)	7500 (4000-12000)
United Arab Emirates	2016	Male	55 to 59	15% (8.7-23%)	85% (77-91%)	2.3 (0.88-5)	4.3% (2.4-6.9%)	5.7% (3.1-8.8%)	130 (66-230)	4800 (2500-8200)
United Arab Emirates	2016	Male	60 to 64	14% (7.8-23%)	86% (77-92%)	1.9 (0.71-4.5)	3.8% (1.9-6.2%)	4.8% (2.4-8%)	77 (38-130)	2300 (1200-3900)
United Arab Emirates	2016	Male	65 to 69	13% (7.8-21%)	87% (79-92%)	1.6 (0.58-3.4)	3.5% (1.5-5.7%)	4.3% (1.9-7.1%)	53 (22-90)	1300 (580-2200)
United Arab Emirates	2016	Male	70 to 74	12% (7.6-19%)	88% (81-92%)	1.2 (0.4-2.7)	2.8% (0.91-4.9%)	3.3% (1.1-5.8%)	26 (8.2-49)	530 (170-960)
United Arab Emirates	2016	Male	75 to 79	11% (6.5-18%)	89% (82-93%)	0.93 (0.29-2.2)	2.5% (0.76-4.6%)	2.9% (0.81-5.4%)	13 (3.6-26)	210 (62-400)
United Arab Emirates	2016	Male	80 to 84	10% (5.9-16%)	90% (84-94%)	0.79 (0.26-1.8)	2.4% (0.83-4.4%)	2.7% (0.84-5%)	10 (3.3-20)	120 (40-230)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
United Arab Emirates	2016	Male	85 to 89	9.4% (5.4-15%)	91% (85-95%)	0.71 (0.24-1.7)	2.2% (0.77-4.2%)	2.5% (0.8-4.7%)	6.5 (2-12)	53 (18-98)
United Arab Emirates	2016	Male	90 to 94	10% (5.8-16%)	90% (84-94%)	0.71 (0.21-1.7)	2.1% (0.55-4.1%)	2.3% (0.55-4.6%)	1.8 (0.43-3.7)	10 (2.7-20)
United Arab Emirates	2016	Male	95 plus	12% (6.1-20%)	88% (80-94%)	0.78 (0.17-2.1)	2.1% (0.49-4.2%)	2.2% (0.43-4.7%)	0.59 (0.11-1.3)	2.3 (0.55-4.7)
Yemen	1990	Female	All Ages	10% (6.2-16%)	90% (84-94%)	0.15 (0.12-0.17)	0.12% (-0.038-0.29%)	0.045% (-0.39-0.53%)	29 (-250-340)	5400 (-1700-14000)
Yemen	1990	Female	15 to 19	9.8% (3.3-21%)	90% (79-97%)	0.13 (0.039-0.29)	0.34% (0.15-0.67%)	0.38% (0.079-0.95%)	4.2 (0.86-10)	520 (240-990)
Yemen	1990	Female	20 to 24	13% (3.8-28%)	87% (72-96%)	0.21 (0.068-0.43)	0.62% (0.29-1.2%)	0.62% (0.098-1.5%)	7.7 (1.2-19)	930 (420-1800)
Yemen	1990	Female	25 to 29	15% (4.4-30%)	85% (70-96%)	0.26 (0.086-0.51)	0.75% (0.35-1.4%)	0.84% (0.14-1.9%)	9.4 (1.6-22)	940 (410-1700)
Yemen	1990	Female	30 to 34	17% (4.3-34%)	83% (66-96%)	0.31 (0.1-0.59)	0.8% (0.29-1.6%)	1% (0.14-2.4%)	11 (1.4-27)	880 (310-1800)
Yemen	1990	Female	35 to 39	18% (5.2-35%)	82% (65-95%)	0.37 (0.12-0.67)	0.78% (0.18-1.6%)	1.1% (0.052-2.4%)	12 (0.61-29)	810 (180-1700)
Yemen	1990	Female	40 to 44	19% (7-35%)	81% (65-93%)	0.41 (0.16-0.76)	0.84% (0.02-2%)	1.2% (-0.08-3.1%)	17 (-1.1-43)	880 (22-2100)
Yemen	1990	Female	45 to 49	20% (6.8-37%)	80% (63-93%)	0.43 (0.15-0.9)	0.73% (-0.31-2.2%)	1.1% (-0.54-3.4%)	15 (-7.5-48)	690 (-300-2100)
Yemen	1990	Female	50 to 54	20% (8.1-39%)	80% (61-92%)	0.44 (0.14-0.92)	0.59% (-0.68-2.2%)	0.84% (-0.96-3.2%)	14 (-17-55)	550 (-630-2100)
Yemen	1990	Female	55 to 59	21% (8.6-41%)	79% (59-91%)	0.44 (0.13-0.93)	0.32% (-1-1.9%)	0.44% (-1.4-2.7%)	9 (-28-57)	290 (-880-1800)
Yemen	1990	Female	60 to 64	22% (9.1-42%)	78% (58-91%)	0.42 (0.11-1)	-0.0063% (-1.5-1.9%)	0.0054% (-1.9-2.5%)	0.17 (-47-62)	-2.4 (-1300-1700)
Yemen	1990	Female	65 to 69	23% (9.8-41%)	77% (59-90%)	0.37 (0.1-0.81)	-0.34% (-1.8-1.1%)	-0.42% (-2.2-1.5%)	-12 (-67-45)	-280 (-1500-970)
Yemen	1990	Female	70 to 74	22% (9.4-39%)	78% (61-91%)	0.3 (0.079-0.73)	-0.49% (-1.8-0.77%)	-0.59% (-2.2-0.96%)	-19 (-70-29)	-340 (-1300-540)
Yemen	1990	Female	75 to 79	19% (7.6-38%)	81% (62-92%)	0.24 (0.055-0.66)	-0.51% (-1.8-0.51%)	-0.61% (-2.1-0.6%)	-18 (-63-18)	-250 (-880-250)
Yemen	1990	Female	80 to 84	18% (5.9-37%)	82% (63-94%)	0.22 (0.047-0.61)	-0.42% (-1.6-0.46%)	-0.5% (-1.8-0.58%)	-12 (-45-13)	-120 (-460-140)
Yemen	1990	Female	85 to 89	18% (5.9-38%)	82% (62-94%)	0.22 (0.044-0.63)	-0.45% (-1.6-0.39%)	-0.55% (-2-0.48%)	-7 (-26-6.1)	-51 (-190-44)
Yemen	1990	Female	90 to 94	21% (6.5-43%)	79% (57-93%)	0.25 (0.046-0.73)	-0.52% (-1.8-0.36%)	-0.66% (-2.2-0.42%)	-3.1 (-10-2)	-15 (-51-11)
Yemen	1990	Female	95 plus	25% (7.6-51%)	75% (49-92%)	0.3 (0.05-0.89)	-0.61% (-2-0.64%)	-0.78% (-2.4-0.76%)	-1 (-3.1-0.97)	-3.2 (-10-3.4)
Yemen	1990	Male	All Ages	7.6% (5.1-11%)	92% (89-95%)	0.15 (0.13-0.18)	0.32% (0.19-0.46%)	0.44% (0.18-0.77%)	260 (110-460)	15000 (9200-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	1990	Male	15 to 19	7.2% (3.1-14%)	93% (86-97%)	0.094 (0.03-0.2)	0.63% (0.33-1.1%)	0.69% (0.19-1.5%)	7.1 (2-16)	860 (440-1500)
Yemen	1990	Male	20 to 24	11% (4.4-22%)	89% (78-96%)	0.21 (0.076-0.41)	1.4% (0.78-2.4%)	1.5% (0.58-3%)	20 (7.4-39)	2000 (1100-3400)
Yemen	1990	Male	25 to 29	14% (5.6-26%)	86% (74-94%)	0.33 (0.14-0.61)	1.9% (1-3.1%)	2.3% (0.96-4.2%)	26 (11-49)	2200 (1100-3600)
Yemen	1990	Male	30 to 34	16% (5.9-30%)	84% (70-94%)	0.42 (0.18-0.74)	2% (1.1-3.3%)	2.5% (1.2-4.6%)	28 (13-51)	2000 (1100-3300)
Yemen	1990	Male	35 to 39	17% (6.5-32%)	83% (68-93%)	0.48 (0.19-0.87)	2% (1-3.6%)	2.6% (1.1-4.9%)	29 (12-57)	1800 (890-3200)
Yemen	1990	Male	40 to 44	17% (7.7-30%)	83% (70-92%)	0.51 (0.2-0.97)	1.8% (0.75-3.3%)	2.2% (0.75-4.6%)	28 (9.5-57)	1500 (650-2900)
Yemen	1990	Male	45 to 49	17% (6.6-31%)	83% (69-93%)	0.52 (0.17-1)	1.7% (0.56-3.5%)	2.1% (0.49-4.7%)	26 (6-58)	1200 (400-2600)
Yemen	1990	Male	50 to 54	16% (6.7-31%)	84% (69-93%)	0.51 (0.18-1.1)	1.3% (0.27-3%)	1.6% (0.14-4%)	22 (1.8-56)	910 (190-2100)
Yemen	1990	Male	55 to 59	15% (6.9-30%)	85% (70-93%)	0.49 (0.16-1)	1.2% (0.18-2.9%)	1.5% (0.11-3.8%)	23 (1.7-57)	780 (110-1900)
Yemen	1990	Male	60 to 64	15% (6.4-29%)	85% (71-94%)	0.44 (0.13-1)	0.95% (-0.035-2.8%)	1.1% (-0.13-3.5%)	21 (-2.6-64)	610 (-21-1800)
Yemen	1990	Male	65 to 69	14% (6.5-26%)	86% (74-94%)	0.37 (0.12-0.77)	0.56% (-0.2-1.9%)	0.66% (-0.33-2.3%)	13 (-6.6-48)	330 (-120-1100)
Yemen	1990	Male	70 to 74	13% (6.1-23%)	87% (77-94%)	0.27 (0.079-0.64)	0.37% (-0.29-1.5%)	0.42% (-0.38-1.8%)	9 (-8.4-38)	170 (-140-700)
Yemen	1990	Male	75 to 79	11% (5.1-21%)	89% (79-95%)	0.21 (0.055-0.54)	0.22% (-0.32-1%)	0.24% (-0.41-1.2%)	4.4 (-7.7-23)	68 (-100-330)
Yemen	1990	Male	80 to 84	10% (4.2-20%)	90% (80-96%)	0.18 (0.047-0.42)	0.2% (-0.22-0.83%)	0.22% (-0.3-0.98%)	3.4 (-4.7-15)	39 (-44-160)
Yemen	1990	Male	85 to 89	10% (4-20%)	90% (80-96%)	0.16 (0.043-0.4)	0.12% (-0.28-0.61%)	0.11% (-0.38-0.7%)	0.97 (-3.4-6.4)	9.2 (-23-49)
Yemen	1990	Male	90 to 94	11% (4.6-23%)	89% (77-95%)	0.17 (0.046-0.44)	0.041% (-0.36-0.52%)	0.0046% (-0.51-0.59%)	0.015 (-1.7-2)	0.84 (-7.6-11)
Yemen	1990	Male	95 plus	14% (5.2-29%)	86% (71-95%)	0.2 (0.041-0.55)	0.028% (-0.46-0.65%)	-0.039% (-0.67-0.71%)	-0.033 (-0.56-0.59)	0.097 (-1.6-2.2)
Yemen	1995	Female	All Ages	10% (6.4-16%)	90% (84-94%)	0.11 (0.09-0.13)	0.053% (-0.098-0.19%)	-0.13% (-0.56-0.22%)	-89 (-390-170)	2700 (-4800-9800)
Yemen	1995	Female	15 to 19	9.8% (3-23%)	90% (77-97%)	0.1 (0.031-0.24)	0.29% (0.15-0.53%)	0.29% (0.054-0.72%)	3.6 (0.68-9)	520 (260-970)
Yemen	1995	Female	20 to 24	13% (4-27%)	87% (73-96%)	0.16 (0.05-0.35)	0.52% (0.26-0.94%)	0.43% (0.051-1.1%)	5.2 (0.62-14)	790 (390-1400)
Yemen	1995	Female	25 to 29	14% (3.6-30%)	86% (70-96%)	0.2 (0.06-0.41)	0.6% (0.3-1%)	0.56% (0.064-1.3%)	6.9 (0.81-16)	850 (420-1500)
Yemen	1995	Female	30 to 34	16% (4.2-32%)	84% (68-96%)	0.24 (0.074-0.46)	0.57% (0.18-1.1%)	0.61% (-0.049-1.6%)	8.5 (-0.73-22)	850 (270-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	1995	Female	35 to 39	18% (5-33%)	82% (67-95%)	0.27 (0.095-0.49)	0.51% (0.072-1%)	0.6% (-0.15-1.5%)	9.9 (-2.4-25)	760 (99-1600)
Yemen	1995	Female	40 to 44	18% (6.8-34%)	82% (66-93%)	0.3 (0.12-0.56)	0.48% (-0.11-1.2%)	0.64% (-0.32-1.9%)	12 (-5.8-34)	700 (-170-1700)
Yemen	1995	Female	45 to 49	19% (6.9-35%)	81% (65-93%)	0.33 (0.12-0.64)	0.32% (-0.51-1.3%)	0.42% (-0.87-1.9%)	8 (-16-38)	410 (-640-1600)
Yemen	1995	Female	50 to 54	20% (7.8-38%)	80% (62-92%)	0.34 (0.11-0.68)	0.15% (-0.87-1.3%)	0.19% (-1.3-1.8%)	3.9 (-26-38)	170 (-940-1400)
Yemen	1995	Female	55 to 59	21% (8.2-40%)	79% (60-92%)	0.33 (0.096-0.72)	-0.1% (-1.3-1%)	-0.14% (-1.8-1.4%)	-2.8 (-37-30)	-90 (-1200-940)
Yemen	1995	Female	60 to 64	22% (8.6-41%)	78% (59-91%)	0.32 (0.083-0.75)	-0.39% (-1.7-0.89%)	-0.5% (-2.2-1.2%)	-13 (-57-29)	-360 (-1600-780)
Yemen	1995	Female	65 to 69	22% (9.7-39%)	78% (61-90%)	0.27 (0.078-0.59)	-0.61% (-1.9-0.4%)	-0.76% (-2.4-0.49%)	-26 (-82-17)	-600 (-1900-380)
Yemen	1995	Female	70 to 74	21% (8.4-38%)	79% (62-92%)	0.22 (0.059-0.53)	-0.66% (-1.7-0.19%)	-0.8% (-2.1-0.24%)	-32 (-87-9.3)	-590 (-1600-170)
Yemen	1995	Female	75 to 79	19% (6.8-38%)	81% (62-93%)	0.18 (0.039-0.49)	-0.64% (-1.9-0.068%)	-0.76% (-2.3-0.098%)	-31 (-92-3.7)	-430 (-1300-45)
Yemen	1995	Female	80 to 84	17% (6.2-38%)	83% (62-94%)	0.16 (0.037-0.45)	-0.55% (-1.8-0.081%)	-0.66% (-2.1-0.1%)	-24 (-76-3.7)	-250 (-800-38)
Yemen	1995	Female	85 to 89	18% (6.2-39%)	82% (61-94%)	0.17 (0.036-0.49)	-0.58% (-1.8-0.065%)	-0.7% (-2.2-0.081%)	-12 (-39-1.4)	-90 (-280-9.7)
Yemen	1995	Female	90 to 94	21% (7-43%)	79% (57-93%)	0.19 (0.041-0.54)	-0.65% (-2--0.00073%)	-0.81% (-2.3--0.014%)	-5.2 (-15--0.093)	-25 (-75--0.029)
Yemen	1995	Female	95 plus	25% (7.8-49%)	75% (51-92%)	0.23 (0.041-0.67)	-0.76% (-2.1-0.085%)	-0.95% (-2.6-0.1%)	-1.6 (-4.4-0.17)	-5.3 (-14-0.59)
Yemen	1995	Male	All Ages	8% (5.5-11%)	92% (89-94%)	0.12 (0.11-0.15)	0.28% (0.15-0.43%)	0.31% (0.055-0.61%)	230 (37-450)	15000 (7900-21000)
Yemen	1995	Male	15 to 19	7.2% (3.1-15%)	93% (85-97%)	0.072 (0.023-0.16)	0.58% (0.3-0.99%)	0.58% (0.15-1.3%)	7.6 (2-17)	1000 (510-1700)
Yemen	1995	Male	20 to 24	11% (4.3-22%)	89% (78-96%)	0.16 (0.057-0.31)	1.3% (0.71-2.1%)	1.3% (0.44-2.6%)	18 (6.3-38)	2000 (1100-3300)
Yemen	1995	Male	25 to 29	14% (5.6-26%)	86% (74-94%)	0.25 (0.098-0.47)	1.6% (0.88-2.5%)	1.8% (0.68-3.3%)	23 (8.6-43)	2000 (1100-3300)
Yemen	1995	Male	30 to 34	16% (5.9-30%)	84% (70-94%)	0.31 (0.13-0.54)	1.6% (0.89-2.6%)	1.9% (0.84-3.5%)	25 (11-46)	1900 (1000-3000)
Yemen	1995	Male	35 to 39	17% (6.8-31%)	83% (69-93%)	0.36 (0.15-0.62)	1.5% (0.75-2.5%)	1.8% (0.71-3.3%)	30 (12-57)	2000 (990-3400)
Yemen	1995	Male	40 to 44	17% (8.2-31%)	83% (69-92%)	0.38 (0.16-0.7)	1.2% (0.54-2.2%)	1.5% (0.43-2.9%)	29 (8.4-56)	1700 (740-3000)
Yemen	1995	Male	45 to 49	17% (7.1-31%)	83% (69-93%)	0.4 (0.14-0.77)	1.1% (0.27-2.2%)	1.3% (0.13-2.9%)	28 (2.6-62)	1400 (350-2800)
Yemen	1995	Male	50 to 54	16% (7.3-31%)	84% (69-93%)	0.39 (0.14-0.81)	0.82% (0.0062-2%)	0.95% (-0.21-2.6%)	21 (-4.8-55)	920 (6.5-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	1995	Male	55 to 59	16% (7.1-29%)	84% (71-93%)	0.38 (0.13-0.77)	0.68% (-0.053-1.8%)	0.8% (-0.19-2.3%)	17 (-4.1-50)	630 (-52-1700)
Yemen	1995	Male	60 to 64	15% (6.4-28%)	85% (72-94%)	0.34 (0.1-0.77)	0.49% (-0.24-1.7%)	0.56% (-0.41-2.1%)	16 (-11-60)	490 (-240-1700)
Yemen	1995	Male	65 to 69	14% (6.3-27%)	86% (73-94%)	0.27 (0.089-0.59)	0.23% (-0.37-1.1%)	0.24% (-0.52-1.3%)	8.5 (-19-44)	230 (-370-1000)
Yemen	1995	Male	70 to 74	13% (5.9-23%)	87% (77-94%)	0.2 (0.064-0.46)	0.12% (-0.4-0.71%)	0.12% (-0.55-0.86%)	5 (-23-36)	110 (-380-650)
Yemen	1995	Male	75 to 79	11% (4.9-22%)	89% (78-95%)	0.16 (0.042-0.41)	0.049% (-0.43-0.48%)	0.036% (-0.56-0.54%)	1.5 (-22-21)	33 (-290-320)
Yemen	1995	Male	80 to 84	10% (4-20%)	90% (80-96%)	0.13 (0.037-0.31)	0.06% (-0.33-0.42%)	0.051% (-0.4-0.49%)	1.6 (-13-16)	24 (-130-170)
Yemen	1995	Male	85 to 89	9.9% (4-19%)	90% (81-96%)	0.12 (0.034-0.29)	0.0052% (-0.35-0.29%)	-0.021% (-0.45-0.32%)	-0.28 (-6-4.2)	0.63 (-41-34)
Yemen	1995	Male	90 to 94	11% (4.4-23%)	89% (77-96%)	0.13 (0.034-0.33)	-0.058% (-0.45-0.23%)	-0.11% (-0.59-0.25%)	-0.57 (-3.1-1.3)	-1.8 (-14-7.1)
Yemen	1995	Male	95 plus	14% (5.1-30%)	86% (70-95%)	0.14 (0.031-0.4)	-0.075% (-0.55-0.24%)	-0.16% (-0.73-0.23%)	-0.2 (-0.94-0.29)	-0.39 (-2.8-1.2)
Yemen	2000	Female	All Ages	10% (6-16%)	90% (84-94%)	0.066 (0.053-0.079)	-0.0028% (-0.13-0.1%)	-0.24% (-0.63-0.04%)	-170 (-440-30)	-120 (-6800-5200)
Yemen	2000	Female	15 to 19	9.7% (3-21%)	90% (79-97%)	0.06 (0.019-0.14)	0.25% (0.13-0.41%)	0.19% (0.021-0.47%)	2.6 (0.3-6.6)	530 (270-890)
Yemen	2000	Female	20 to 24	12% (3.6-27%)	88% (73-96%)	0.096 (0.031-0.2)	0.43% (0.25-0.67%)	0.24% (-0.012-0.61%)	3.5 (-0.17-9)	840 (470-1300)
Yemen	2000	Female	25 to 29	14% (4.1-30%)	86% (70-96%)	0.12 (0.038-0.24)	0.47% (0.26-0.74%)	0.31% (-0.007-0.76%)	3.7 (-0.085-9.1)	700 (380-1100)
Yemen	2000	Female	30 to 34	16% (4.2-33%)	84% (67-96%)	0.14 (0.046-0.28)	0.41% (0.14-0.68%)	0.28% (-0.19-0.8%)	3.6 (-2.5-10)	590 (180-1000)
Yemen	2000	Female	35 to 39	17% (4.7-34%)	83% (66-95%)	0.16 (0.053-0.3)	0.31% (-0.00062-0.56%)	0.23% (-0.28-0.7%)	3.6 (-4.5-11)	460 (-0.92-860)
Yemen	2000	Female	40 to 44	18% (6.8-35%)	82% (65-93%)	0.18 (0.068-0.34)	0.18% (-0.25-0.57%)	0.14% (-0.58-0.82%)	2.6 (-11-16)	290 (-380-900)
Yemen	2000	Female	45 to 49	19% (7-35%)	81% (65-93%)	0.19 (0.068-0.37)	-0.026% (-0.67-0.43%)	-0.13% (-1.1-0.6%)	-3 (-26-14)	-39 (-1000-690)
Yemen	2000	Female	50 to 54	20% (8-37%)	80% (63-92%)	0.2 (0.068-0.39)	-0.21% (-1-0.33%)	-0.34% (-1.5-0.44%)	-8.8 (-39-12)	-300 (-1400-470)
Yemen	2000	Female	55 to 59	21% (8.6-39%)	79% (61-91%)	0.2 (0.06-0.42)	-0.43% (-1.5-0.21%)	-0.59% (-2.1-0.29%)	-16 (-58-8.1)	-520 (-1900-250)
Yemen	2000	Female	60 to 64	22% (8.4-42%)	78% (58-92%)	0.19 (0.05-0.45)	-0.63% (-1.9-0.063%)	-0.82% (-2.5-0.087%)	-23 (-70-2.6)	-630 (-2000-61)
Yemen	2000	Female	65 to 69	23% (9.2-40%)	77% (60-91%)	0.17 (0.049-0.36)	-0.72% (-1.9--0.0055%)	-0.9% (-2.4-0.0015%)	-30 (-83-0.05)	-700 (-1900--5.7)
Yemen	2000	Female	70 to 74	21% (9-38%)	79% (62-91%)	0.13 (0.037-0.33)	-0.66% (-1.8-0.00012%)	-0.81% (-2.1-0.01%)	-33 (-88-0.4)	-610 (-1600-0.11)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	2000	Female	75 to 79	19% (7.3-39%)	81% (61-93%)	0.11 (0.023-0.3)	-0.59% (-1.9-0.017%)	-0.71% (-2.2-0.026%)	-30 (-92-1)	-420 (-1300-13)
Yemen	2000	Female	80 to 84	17% (6-39%)	83% (61-94%)	0.097 (0.021-0.29)	-0.51% (-1.7-0.033%)	-0.61% (-2-0.037%)	-24 (-78-1.5)	-250 (-840-17)
Yemen	2000	Female	85 to 89	18% (6.1-40%)	82% (60-94%)	0.098 (0.021-0.29)	-0.53% (-1.7-0.0022%)	-0.64% (-2.1-0.00018%)	-14 (-46-0.0036)	-100 (-340-0.41)
Yemen	2000	Female	90 to 94	20% (6.5-43%)	80% (57-93%)	0.11 (0.023-0.32)	-0.59% (-1.8--0.031%)	-0.73% (-2.2--0.046%)	-4.9 (-15--0.32)	-24 (-74--1.2)
Yemen	2000	Female	95 plus	25% (8-50%)	75% (50-92%)	0.14 (0.024-0.39)	-0.7% (-2.2--0.024%)	-0.87% (-2.6--0.061%)	-1.6 (-4.7--0.11)	-5 (-15--0.17)
Yemen	2000	Male	All Ages	7.7% (5.4-11%)	92% (89-95%)	0.072 (0.06-0.085)	0.21% (0.098-0.36%)	0.16% (-0.057-0.39%)	120 (-40-270)	11000 (5100-17000)
Yemen	2000	Male	15 to 19	7.1% (3-14%)	93% (86-97%)	0.042 (0.014-0.088)	0.48% (0.28-0.8%)	0.4% (0.097-0.97%)	5.9 (1.4-14)	990 (550-1700)
Yemen	2000	Male	20 to 24	11% (4.3-21%)	89% (79-96%)	0.093 (0.034-0.18)	1% (0.6-1.7%)	0.85% (0.22-2%)	15 (4-35)	2100 (1200-3500)
Yemen	2000	Male	25 to 29	13% (5.5-26%)	87% (74-94%)	0.15 (0.059-0.28)	1.2% (0.69-1.9%)	1.2% (0.39-2.4%)	17 (6-35)	1900 (1100-3000)
Yemen	2000	Male	30 to 34	15% (5.6-30%)	85% (70-94%)	0.19 (0.078-0.33)	1.2% (0.64-1.9%)	1.2% (0.45-2.5%)	16 (6.1-33)	1500 (800-2400)
Yemen	2000	Male	35 to 39	17% (6.3-31%)	83% (69-94%)	0.21 (0.079-0.38)	1.1% (0.56-1.8%)	1.2% (0.35-2.2%)	17 (5.1-34)	1300 (660-2100)
Yemen	2000	Male	40 to 44	17% (7.6-30%)	83% (70-92%)	0.22 (0.092-0.42)	0.82% (0.34-1.4%)	0.86% (0.16-1.7%)	16 (2.8-32)	1100 (460-1900)
Yemen	2000	Male	45 to 49	16% (6.9-30%)	84% (70-93%)	0.23 (0.085-0.45)	0.63% (0.1-1.2%)	0.66% (-0.17-1.5%)	15 (-3.8-35)	920 (150-1700)
Yemen	2000	Male	50 to 54	16% (6.7-30%)	84% (70-93%)	0.22 (0.082-0.46)	0.4% (-0.18-0.92%)	0.38% (-0.48-1.1%)	9.8 (-12-30)	540 (-260-1300)
Yemen	2000	Male	55 to 59	15% (6.4-29%)	85% (71-94%)	0.22 (0.071-0.45)	0.27% (-0.32-0.8%)	0.25% (-0.56-0.99%)	6.9 (-15-27)	320 (-380-950)
Yemen	2000	Male	60 to 64	15% (6.1-28%)	85% (72-94%)	0.2 (0.059-0.45)	0.14% (-0.47-0.63%)	0.11% (-0.67-0.74%)	3.1 (-19-21)	140 (-460-630)
Yemen	2000	Male	65 to 69	14% (6.3-25%)	86% (75-94%)	0.16 (0.053-0.34)	0.017% (-0.49-0.41%)	-0.022% (-0.68-0.49%)	-0.78 (-24-17)	16 (-500-420)
Yemen	2000	Male	70 to 74	13% (5.7-23%)	87% (77-94%)	0.12 (0.037-0.27)	-0.0019% (-0.45-0.31%)	-0.029% (-0.61-0.36%)	-1.2 (-24-14)	-1.6 (-410-280)
Yemen	2000	Male	75 to 79	11% (4.6-22%)	89% (78-95%)	0.092 (0.023-0.25)	-0.01% (-0.41-0.27%)	-0.034% (-0.53-0.31%)	-1.4 (-21-12)	-6.9 (-280-180)
Yemen	2000	Male	80 to 84	10% (3.9-20%)	90% (80-96%)	0.077 (0.021-0.19)	0.009% (-0.33-0.26%)	-0.0085% (-0.42-0.3%)	-0.31 (-16-11)	4.2 (-150-120)
Yemen	2000	Male	85 to 89	9.7% (3.9-19%)	90% (81-96%)	0.07 (0.02-0.18)	-0.018% (-0.34-0.19%)	-0.048% (-0.44-0.21%)	-0.85 (-7.9-3.8)	-2.9 (-54-30)
Yemen	2000	Male	90 to 94	11% (4.5-22%)	89% (78-95%)	0.074 (0.02-0.19)	-0.064% (-0.38-0.15%)	-0.11% (-0.5-0.16%)	-0.55 (-2.4-0.8)	-1.9 (-11-4.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	2000	Male	95 plus	13% (5-28%)	87% (72-95%)	0.085 (0.018-0.24)	-0.074% (-0.48-0.15%)	-0.15% (-0.66-0.13%)	-0.19 (-0.81-0.16)	-0.37 (-2.4-0.78)
Yemen	2005	Female	All Ages	10% (6-16%)	90% (84-94%)	0.043 (0.035-0.052)	0.014% (-0.1-0.11%)	-0.2% (-0.55-0.046%)	-140 (-350-33)	670 (-4800-5200)
Yemen	2005	Female	15 to 19	9.8% (3.2-21%)	90% (79-97%)	0.038 (0.011-0.085)	0.24% (0.14-0.37%)	0.15% (0.031-0.36%)	2.3 (0.47-5.3)	580 (330-930)
Yemen	2005	Female	20 to 24	12% (3.7-27%)	88% (73-96%)	0.06 (0.019-0.12)	0.42% (0.25-0.62%)	0.18% (-0.0083-0.44%)	2.9 (-0.13-7.1)	990 (560-1500)
Yemen	2005	Female	25 to 29	14% (3.9-30%)	86% (70-96%)	0.072 (0.023-0.14)	0.46% (0.28-0.68%)	0.23% (-0.01-0.58%)	3.3 (-0.15-8.3)	880 (520-1300)
Yemen	2005	Female	30 to 34	15% (4.1-32%)	85% (68-96%)	0.085 (0.027-0.17)	0.4% (0.19-0.6%)	0.21% (-0.13-0.57%)	2.6 (-1.5-7)	590 (270-920)
Yemen	2005	Female	35 to 39	17% (4.8-33%)	83% (67-95%)	0.1 (0.032-0.19)	0.29% (0.076-0.48%)	0.16% (-0.2-0.5%)	2.4 (-2.8-7)	420 (110-740)
Yemen	2005	Female	40 to 44	18% (7-34%)	82% (66-93%)	0.12 (0.044-0.22)	0.16% (-0.16-0.39%)	0.071% (-0.45-0.5%)	1.3 (-8.1-9.3)	250 (-240-640)
Yemen	2005	Female	45 to 49	19% (7.4-35%)	81% (65-93%)	0.12 (0.044-0.25)	-0.035% (-0.51-0.27%)	-0.17% (-0.92-0.35%)	-3.9 (-22-8)	-56 (-830-450)
Yemen	2005	Female	50 to 54	20% (8.2-38%)	80% (62-92%)	0.13 (0.044-0.26)	-0.2% (-0.84-0.2%)	-0.34% (-1.3-0.25%)	-10 (-38-7.5)	-340 (-1400-340)
Yemen	2005	Female	55 to 59	21% (9-40%)	79% (60-91%)	0.13 (0.039-0.27)	-0.37% (-1.3-0.12%)	-0.52% (-1.8-0.15%)	-18 (-60-5.2)	-570 (-2000-180)
Yemen	2005	Female	60 to 64	22% (9.3-42%)	78% (58-91%)	0.12 (0.033-0.27)	-0.51% (-1.6-0.046%)	-0.68% (-2.1-0.048%)	-24 (-71-1.8)	-670 (-2000-61)
Yemen	2005	Female	65 to 69	22% (9.7-40%)	78% (60-90%)	0.1 (0.029-0.22)	-0.53% (-1.5-0.044%)	-0.66% (-1.9-0.051%)	-23 (-68-1.8)	-540 (-1600-45)
Yemen	2005	Female	70 to 74	21% (8.7-38%)	79% (62-91%)	0.081 (0.022-0.2)	-0.45% (-1.4-0.068%)	-0.55% (-1.7-0.079%)	-22 (-67-3.2)	-400 (-1200-57)
Yemen	2005	Female	75 to 79	18% (6.8-37%)	82% (63-93%)	0.066 (0.014-0.19)	-0.4% (-1.4-0.078%)	-0.48% (-1.7-0.093%)	-20 (-71-3.9)	-280 (-1000-57)
Yemen	2005	Female	80 to 84	17% (5.5-37%)	83% (63-94%)	0.059 (0.012-0.17)	-0.34% (-1.3-0.082%)	-0.41% (-1.5-0.095%)	-15 (-53-3.4)	-150 (-570-37)
Yemen	2005	Female	85 to 89	17% (5.7-38%)	83% (62-94%)	0.06 (0.012-0.17)	-0.36% (-1.3-0.049%)	-0.44% (-1.6-0.053%)	-9.7 (-35-1.2)	-70 (-260-9.6)
Yemen	2005	Female	90 to 94	20% (6.9-43%)	80% (57-93%)	0.07 (0.014-0.2)	-0.42% (-1.4-0.024%)	-0.53% (-1.6-0.019%)	-4.1 (-13-0.15)	-20 (-65-1.2)
Yemen	2005	Female	95 plus	25% (7.9-49%)	75% (51-92%)	0.085 (0.015-0.25)	-0.53% (-1.8-0.031%)	-0.66% (-2.2-0.015%)	-1.2 (-3.9-0.028)	-3.8 (-13-0.22)
Yemen	2005	Male	All Ages	7.9% (5.4-11%)	92% (89-95%)	0.046 (0.038-0.056)	0.22% (0.11-0.35%)	0.15% (-0.055-0.35%)	100 (-36-250)	11000 (5300-16000)
Yemen	2005	Male	15 to 19	7% (2.9-13%)	93% (87-97%)	0.026 (0.0084-0.054)	0.42% (0.25-0.68%)	0.27% (0.073-0.65%)	4.8 (1.3-11)	1000 (600-1700)
Yemen	2005	Male	20 to 24	11% (4.3-21%)	89% (79-96%)	0.057 (0.021-0.11)	0.89% (0.54-1.4%)	0.57% (0.14-1.3%)	12 (3.1-29)	2100 (1300-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	2005	Male	25 to 29	13% (5.4-25%)	87% (75-95%)	0.089 (0.036-0.17)	1% (0.61-1.6%)	0.82% (0.24-1.7%)	16 (4.6-32)	2000 (1200-3200)
Yemen	2005	Male	30 to 34	15% (5.8-29%)	85% (71-94%)	0.12 (0.047-0.21)	0.96% (0.57-1.5%)	0.89% (0.29-1.9%)	14 (4.6-30)	1500 (860-2400)
Yemen	2005	Male	35 to 39	17% (6.2-31%)	83% (69-94%)	0.13 (0.049-0.24)	0.9% (0.51-1.4%)	0.89% (0.24-1.8%)	13 (3.6-26)	1200 (650-1800)
Yemen	2005	Male	40 to 44	17% (7.6-30%)	83% (70-92%)	0.14 (0.055-0.27)	0.7% (0.32-1.1%)	0.66% (0.041-1.4%)	11 (0.64-23)	880 (410-1500)
Yemen	2005	Male	45 to 49	17% (6.8-30%)	83% (70-93%)	0.14 (0.052-0.29)	0.53% (0.12-0.96%)	0.5% (-0.13-1.2%)	11 (-2.9-27)	760 (160-1400)
Yemen	2005	Male	50 to 54	16% (7-31%)	84% (69-93%)	0.14 (0.051-0.29)	0.31% (-0.18-0.69%)	0.26% (-0.49-0.82%)	7.4 (-14-23)	480 (-290-1100)
Yemen	2005	Male	55 to 59	15% (6.5-29%)	85% (71-93%)	0.14 (0.046-0.28)	0.19% (-0.35-0.57%)	0.14% (-0.59-0.7%)	4.7 (-19-22)	270 (-500-830)
Yemen	2005	Male	60 to 64	15% (6.2-28%)	85% (72-94%)	0.12 (0.037-0.28)	0.09% (-0.39-0.42%)	0.05% (-0.59-0.5%)	1.7 (-21-18)	110 (-490-540)
Yemen	2005	Male	65 to 69	14% (6.1-25%)	86% (75-94%)	0.099 (0.032-0.21)	0.033% (-0.41-0.33%)	-0.00076% (-0.57-0.39%)	-0.042 (-19-13)	31 (-400-320)
Yemen	2005	Male	70 to 74	12% (5.6-23%)	88% (77-94%)	0.073 (0.022-0.17)	0.047% (-0.32-0.32%)	0.031% (-0.42-0.37%)	1.2 (-17-14)	40 (-270-270)
Yemen	2005	Male	75 to 79	11% (4.4-21%)	89% (79-96%)	0.056 (0.014-0.15)	0.047% (-0.29-0.27%)	0.036% (-0.38-0.31%)	1.4 (-14-12)	30 (-190-180)
Yemen	2005	Male	80 to 84	9.8% (4-19%)	90% (81-96%)	0.047 (0.013-0.12)	0.058% (-0.21-0.28%)	0.052% (-0.28-0.33%)	1.5 (-8.3-9.9)	22 (-77-100)
Yemen	2005	Male	85 to 89	9.6% (4-19%)	90% (81-96%)	0.044 (0.012-0.11)	0.029% (-0.21-0.2%)	0.011% (-0.29-0.23%)	0.22 (-6.4-9)	5.4 (-40-38)
Yemen	2005	Male	90 to 94	11% (4.4-22%)	89% (78-96%)	0.046 (0.012-0.12)	-0.012% (-0.29-0.16%)	-0.049% (-0.38-0.17%)	-0.34 (-2.7-1.2)	-0.51 (-12-7)
Yemen	2005	Male	95 plus	13% (4.8-28%)	87% (72-95%)	0.053 (0.011-0.14)	-0.024% (-0.38-0.16%)	-0.089% (-0.53-0.14%)	-0.11 (-0.64-0.17)	-0.12 (-1.9-0.79)
Yemen	2010	Female	All Ages	11% (6.1-16%)	89% (84-94%)	0.037 (0.027-0.047)	0.039% (-0.079-0.13%)	-0.18% (-0.51-0.054%)	-120 (-330-39)	1900 (-3600-6200)
Yemen	2010	Female	15 to 19	9.8% (3.3-21%)	90% (79-97%)	0.031 (0.0097-0.068)	0.25% (0.14-0.38%)	0.15% (0.046-0.33%)	2.1 (0.63-4.6)	650 (360-1000)
Yemen	2010	Female	20 to 24	13% (4.1-27%)	87% (73-96%)	0.049 (0.017-0.099)	0.46% (0.28-0.67%)	0.18% (0.013-0.4%)	2.8 (0.2-6.3)	1200 (710-1900)
Yemen	2010	Female	25 to 29	14% (3.8-30%)	86% (70-96%)	0.06 (0.019-0.12)	0.49% (0.32-0.69%)	0.22% (0.012-0.53%)	3.4 (0.2-7.9)	1100 (690-1700)
Yemen	2010	Female	30 to 34	16% (3.7-33%)	84% (67-96%)	0.071 (0.021-0.14)	0.42% (0.24-0.63%)	0.21% (-0.071-0.51%)	3.1 (-1-7.5)	820 (420-1300)
Yemen	2010	Female	35 to 39	18% (4.6-35%)	82% (65-95%)	0.084 (0.024-0.16)	0.3% (0.11-0.47%)	0.16% (-0.17-0.46%)	2.3 (-2.2-6.4)	460 (170-760)
Yemen	2010	Female	40 to 44	18% (6.3-34%)	82% (66-94%)	0.094 (0.034-0.19)	0.18% (-0.07-0.38%)	0.085% (-0.36-0.41%)	1.4 (-5.8-7.1)	270 (-110-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	2010	Female	45 to 49	19% (6.8-35%)	81% (65-93%)	0.1 (0.035-0.2)	0.0085% (-0.39-0.25%)	-0.12% (-0.77-0.3%)	-2.6 (-17-6.6)	15 (-670-420)
Yemen	2010	Female	50 to 54	19% (7.2-37%)	81% (63-93%)	0.1 (0.031-0.22)	-0.12% (-0.68-0.22%)	-0.25% (-1.1-0.24%)	-7.5 (-33-7.5)	-220 (-1200-380)
Yemen	2010	Female	55 to 59	20% (7.8-40%)	80% (60-92%)	0.1 (0.029-0.22)	-0.27% (-1.1-0.13%)	-0.4% (-1.5-0.16%)	-16 (-58-6.1)	-500 (-1900-230)
Yemen	2010	Female	60 to 64	21% (8.4-41%)	79% (59-92%)	0.096 (0.025-0.22)	-0.41% (-1.4-0.078%)	-0.55% (-1.8-0.091%)	-23 (-79-4)	-660 (-2200-130)
Yemen	2010	Female	65 to 69	22% (9.4-40%)	78% (60-91%)	0.084 (0.022-0.19)	-0.42% (-1.3-0.071%)	-0.54% (-1.7-0.079%)	-24 (-73-3.5)	-560 (-1700-93)
Yemen	2010	Female	70 to 74	21% (8.5-38%)	79% (62-92%)	0.067 (0.018-0.17)	-0.36% (-1.2-0.077%)	-0.45% (-1.5-0.091%)	-18 (-60-3.7)	-340 (-1100-73)
Yemen	2010	Female	75 to 79	18% (7-38%)	82% (62-93%)	0.054 (0.012-0.15)	-0.32% (-1.2-0.081%)	-0.39% (-1.5-0.096%)	-16 (-59-3.9)	-220 (-850-57)
Yemen	2010	Female	80 to 84	17% (5.9-38%)	83% (62-94%)	0.049 (0.01-0.14)	-0.27% (-1.1-0.091%)	-0.33% (-1.3-0.11%)	-11 (-45-3.7)	-120 (-470-40)
Yemen	2010	Female	85 to 89	17% (5.9-38%)	83% (62-94%)	0.05 (0.0097-0.15)	-0.3% (-1.1-0.052%)	-0.37% (-1.3-0.054%)	-8.3 (-30-1.3)	-60 (-220-10)
Yemen	2010	Female	90 to 94	20% (6.9-43%)	80% (57-93%)	0.057 (0.011-0.16)	-0.35% (-1.1-0.037%)	-0.44% (-1.4-0.034%)	-3.9 (-13-0.31)	-19 (-63-2)
Yemen	2010	Female	95 plus	25% (7.3-50%)	75% (50-93%)	0.068 (0.011-0.2)	-0.43% (-1.5-0.039%)	-0.54% (-1.9-0.024%)	-1.3 (-4.3-0.055)	-4 (-14-0.36)
Yemen	2010	Male	All Ages	8.1% (5.3-11%)	92% (89-95%)	0.039 (0.029-0.05)	0.25% (0.13-0.38%)	0.16% (-0.031-0.35%)	110 (-22-250)	12000 (6800-18000)
Yemen	2010	Male	15 to 19	7% (2.9-14%)	93% (86-97%)	0.021 (0.0067-0.045)	0.43% (0.26-0.65%)	0.25% (0.073-0.56%)	4.6 (1.4-10)	1200 (690-1800)
Yemen	2010	Male	20 to 24	11% (4.2-21%)	89% (79-96%)	0.046 (0.017-0.087)	0.88% (0.55-1.3%)	0.51% (0.14-1.2%)	12 (3.3-28)	2500 (1500-3900)
Yemen	2010	Male	25 to 29	13% (5.3-25%)	87% (75-95%)	0.07 (0.03-0.13)	0.98% (0.63-1.5%)	0.73% (0.23-1.5%)	16 (4.9-33)	2400 (1500-3600)
Yemen	2010	Male	30 to 34	15% (5.7-28%)	85% (72-94%)	0.091 (0.038-0.16)	0.91% (0.57-1.4%)	0.79% (0.25-1.6%)	16 (5.1-33)	1900 (1100-3000)
Yemen	2010	Male	35 to 39	17% (6.5-30%)	83% (70-94%)	0.11 (0.039-0.19)	0.86% (0.51-1.3%)	0.82% (0.26-1.6%)	15 (4.6-28)	1400 (830-2200)
Yemen	2010	Male	40 to 44	17% (7.8-29%)	83% (71-92%)	0.11 (0.044-0.22)	0.69% (0.34-1.1%)	0.63% (0.086-1.2%)	11 (1.5-22)	930 (450-1500)
Yemen	2010	Male	45 to 49	17% (6.5-30%)	83% (70-93%)	0.12 (0.038-0.25)	0.54% (0.16-0.89%)	0.5% (-0.087-1.1%)	10 (-1.7-22)	720 (190-1200)
Yemen	2010	Male	50 to 54	16% (6.7-31%)	84% (69-93%)	0.11 (0.039-0.24)	0.33% (-0.085-0.68%)	0.28% (-0.37-0.77%)	7.7 (-9.9-22)	510 (-130-1000)
Yemen	2010	Male	55 to 59	15% (6.4-29%)	85% (71-94%)	0.11 (0.035-0.23)	0.21% (-0.22-0.53%)	0.16% (-0.48-0.63%)	5.6 (-17-22)	330 (-370-860)
Yemen	2010	Male	60 to 64	14% (6.3-28%)	86% (72-94%)	0.099 (0.029-0.23)	0.11% (-0.33-0.42%)	0.069% (-0.54-0.49%)	2.7 (-22-20)	160 (-480-620)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	2010	Male	65 to 69	14% (6.3-25%)	86% (75-94%)	0.081 (0.026-0.17)	0.063% (-0.34-0.34%)	0.034% (-0.49-0.4%)	1.3 (-19-16)	73 (-400-400)
Yemen	2010	Male	70 to 74	13% (5.9-23%)	87% (77-94%)	0.061 (0.018-0.14)	0.075% (-0.24-0.31%)	0.064% (-0.35-0.38%)	2.2 (-12-13)	60 (-200-250)
Yemen	2010	Male	75 to 79	11% (4.6-21%)	89% (79-95%)	0.046 (0.012-0.12)	0.07% (-0.22-0.27%)	0.063% (-0.29-0.33%)	2.2 (-10-11)	42 (-140-170)
Yemen	2010	Male	80 to 84	9.8% (3.8-20%)	90% (80-96%)	0.039 (0.0097-0.094)	0.077% (-0.16-0.27%)	0.073% (-0.23-0.32%)	2 (-6.3-8.5)	26 (-55-91)
Yemen	2010	Male	85 to 89	9.6% (3.8-19%)	90% (81-96%)	0.036 (0.009-0.091)	0.048% (-0.16-0.2%)	0.032% (-0.23-0.23%)	0.64 (-4.7-4.5)	8.5 (-29-36)
Yemen	2010	Male	90 to 94	11% (4.2-22%)	89% (78-96%)	0.038 (0.0091-0.1)	0.012% (-0.22-0.16%)	-0.022% (-0.32-0.16%)	-0.2 (-3-1.5)	0.69 (-13-9.2)
Yemen	2010	Male	95 plus	13% (4.7-28%)	87% (72-95%)	0.043 (0.0084-0.12)	0.0038% (-0.3-0.16%)	-0.058% (-0.44-0.14%)	-0.12 (-0.9-0.3)	0.033 (-2.5-1.4)
Yemen	2016	Female	All Ages	11% (6.7-17%)	89% (83-93%)	0.043 (0.026-0.067)	0.04% (-0.1-0.15%)	-0.21% (-0.61-0.054%)	-160 (-440-43)	2200 (-5000-7800)
Yemen	2016	Female	15 to 19	9.7% (3.3-22%)	90% (78-97%)	0.035 (0.009-0.085)	0.22% (0.13-0.34%)	0.11% (0.037-0.23%)	2.2 (0.74-4.6)	670 (370-1100)
Yemen	2016	Female	20 to 24	12% (3.9-28%)	88% (72-96%)	0.055 (0.016-0.12)	0.42% (0.27-0.63%)	0.15% (0.028-0.33%)	3.4 (0.6-7.5)	1400 (840-2200)
Yemen	2016	Female	25 to 29	14% (4.2-30%)	86% (70-96%)	0.066 (0.02-0.14)	0.48% (0.31-0.69%)	0.2% (0.031-0.44%)	4.4 (0.66-9.5)	1400 (910-2200)
Yemen	2016	Female	30 to 34	16% (4.7-33%)	84% (67-95%)	0.078 (0.026-0.16)	0.42% (0.24-0.63%)	0.2% (-0.041-0.48%)	4.4 (-0.88-10)	1100 (620-1800)
Yemen	2016	Female	35 to 39	17% (5.1-32%)	83% (68-95%)	0.092 (0.03-0.19)	0.31% (0.12-0.48%)	0.17% (-0.13-0.45%)	3.8 (-3-10)	730 (270-1200)
Yemen	2016	Female	40 to 44	18% (6.9-33%)	82% (67-93%)	0.1 (0.038-0.22)	0.18% (-0.069-0.39%)	0.095% (-0.31-0.44%)	2 (-6.7-9.6)	350 (-130-720)
Yemen	2016	Female	45 to 49	19% (6.8-35%)	81% (65-93%)	0.11 (0.034-0.26)	0.011% (-0.38-0.27%)	-0.11% (-0.74-0.32%)	-2.7 (-19-7.9)	21 (-680-500)
Yemen	2016	Female	50 to 54	20% (7.8-38%)	80% (62-92%)	0.12 (0.032-0.27)	-0.13% (-0.72-0.21%)	-0.24% (-1.1-0.24%)	-8.4 (-37-7.9)	-250 (-1400-410)
Yemen	2016	Female	55 to 59	21% (8.6-39%)	79% (61-91%)	0.11 (0.03-0.27)	-0.28% (-1.1-0.12%)	-0.41% (-1.6-0.14%)	-18 (-70-6.4)	-590 (-2300-270)
Yemen	2016	Female	60 to 64	22% (8.8-41%)	78% (59-91%)	0.11 (0.025-0.26)	-0.42% (-1.4-0.076%)	-0.57% (-1.8-0.076%)	-32 (-110-4.2)	-890 (-3000-160)
Yemen	2016	Female	65 to 69	22% (9.1-39%)	78% (61-91%)	0.096 (0.025-0.23)	-0.46% (-1.4-0.064%)	-0.58% (-1.7-0.066%)	-35 (-110-4)	-830 (-2500-110)
Yemen	2016	Female	70 to 74	21% (8.5-38%)	79% (62-91%)	0.078 (0.019-0.2)	-0.41% (-1.3-0.059%)	-0.51% (-1.6-0.078%)	-29 (-93-4.5)	-540 (-1700-77)
Yemen	2016	Female	75 to 79	19% (7.2-38%)	81% (62-93%)	0.064 (0.012-0.19)	-0.37% (-1.3-0.072%)	-0.46% (-1.7-0.08%)	-21 (-76-3.7)	-300 (-1100-58)
Yemen	2016	Female	80 to 84	18% (5.9-39%)	82% (61-94%)	0.058 (0.01-0.17)	-0.33% (-1.3-0.084%)	-0.4% (-1.5-0.094%)	-14 (-54-3.3)	-150 (-570-37)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Yemen	2016	Female	85 to 89	18% (6-40%)	82% (60-94%)	0.058 (0.011-0.19)	-0.35% (-1.3-0.052%)	-0.43% (-1.6-0.054%)	-10 (-37-1.3)	-73 (-270-11)
Yemen	2016	Female	90 to 94	21% (6.7-43%)	79% (57-93%)	0.065 (0.012-0.2)	-0.4% (-1.3-0.03%)	-0.5% (-1.7-0.024%)	-5.1 (-17-0.25)	-25 (-84-1.9)
Yemen	2016	Female	95 plus	25% (7.8-49%)	75% (51-92%)	0.078 (0.013-0.24)	-0.48% (-1.6-0.031%)	-0.61% (-2-0.014%)	-1.8 (-5.8-0.04)	-5.6 (-19-0.38)
Yemen	2016	Male	All Ages	8.4% (5.6-12%)	92% (88-94%)	0.046 (0.028-0.07)	0.26% (0.15-0.4%)	0.16% (-0.017-0.35%)	140 (-14-310)	16000 (9100-24000)
Yemen	2016	Male	15 to 19	7% (2.9-14%)	93% (86-97%)	0.023 (0.0065-0.055)	0.29% (0.17-0.45%)	0.13% (0.037-0.3%)	5 (1.5-12)	1200 (700-1900)
Yemen	2016	Male	20 to 24	11% (4.2-22%)	89% (78-96%)	0.051 (0.017-0.11)	0.52% (0.31-0.8%)	0.23% (0.061-0.54%)	15 (4.1-37)	3100 (1800-4800)
Yemen	2016	Male	25 to 29	13% (5.3-25%)	87% (75-95%)	0.078 (0.03-0.16)	0.67% (0.41-1%)	0.39% (0.13-0.87%)	22 (7.1-48)	3200 (1900-4900)
Yemen	2016	Male	30 to 34	15% (5.8-28%)	85% (72-94%)	0.099 (0.04-0.19)	0.7% (0.44-1.1%)	0.49% (0.16-1%)	22 (7.2-45)	2600 (1600-4100)
Yemen	2016	Male	35 to 39	16% (6.6-30%)	84% (70-93%)	0.11 (0.043-0.24)	0.73% (0.44-1.1%)	0.6% (0.19-1.2%)	22 (7.2-44)	2100 (1200-3300)
Yemen	2016	Male	40 to 44	17% (7.6-29%)	83% (71-92%)	0.12 (0.044-0.26)	0.62% (0.32-0.96%)	0.52% (0.09-1%)	16 (2.8-31)	1300 (670-2100)
Yemen	2016	Male	45 to 49	16% (6.4-30%)	84% (70-94%)	0.13 (0.038-0.28)	0.51% (0.15-0.87%)	0.46% (-0.043-0.96%)	13 (-1.3-27)	890 (270-1500)
Yemen	2016	Male	50 to 54	16% (6.7-31%)	84% (69-93%)	0.13 (0.037-0.3)	0.33% (-0.08-0.65%)	0.27% (-0.32-0.73%)	8.2 (-10-22)	530 (-120-1000)
Yemen	2016	Male	55 to 59	15% (6.6-29%)	85% (71-93%)	0.12 (0.033-0.28)	0.2% (-0.24-0.51%)	0.15% (-0.44-0.57%)	6 (-17-24)	360 (-400-930)
Yemen	2016	Male	60 to 64	15% (6.2-28%)	85% (72-94%)	0.11 (0.029-0.27)	0.11% (-0.36-0.42%)	0.067% (-0.55-0.47%)	3.4 (-28-25)	200 (-650-760)
Yemen	2016	Male	65 to 69	14% (6-26%)	86% (74-94%)	0.089 (0.026-0.21)	0.056% (-0.35-0.32%)	0.024% (-0.5-0.37%)	1.3 (-28-21)	89 (-570-520)
Yemen	2016	Male	70 to 74	12% (5.7-22%)	88% (78-94%)	0.066 (0.019-0.17)	0.064% (-0.25-0.28%)	0.05% (-0.35-0.33%)	2.5 (-18-16)	73 (-290-320)
Yemen	2016	Male	75 to 79	11% (4.6-21%)	89% (79-95%)	0.05 (0.012-0.14)	0.062% (-0.21-0.25%)	0.052% (-0.3-0.28%)	1.9 (-11-10)	41 (-140-160)
Yemen	2016	Male	80 to 84	9.6% (3.8-19%)	90% (81-96%)	0.042 (0.0096-0.12)	0.071% (-0.18-0.25%)	0.063% (-0.23-0.29%)	1.7 (-6.2-7.7)	24 (-58-84)
Yemen	2016	Male	85 to 89	9.5% (3.9-19%)	91% (81-96%)	0.039 (0.0097-0.11)	0.04% (-0.19-0.19%)	0.02% (-0.27-0.21%)	0.43 (-5.6-4.4)	7.5 (-36-36)
Yemen	2016	Male	90 to 94	11% (4.5-23%)	89% (77-95%)	0.042 (0.0098-0.12)	0.0035% (-0.24-0.16%)	-0.033% (-0.32-0.15%)	-0.34 (-3.4-1.6)	0.23 (-15-10)
Yemen	2016	Male	95 plus	14% (5-28%)	86% (72-95%)	0.049 (0.0097-0.16)	-0.0054% (-0.34-0.16%)	-0.071% (-0.48-0.13%)	-0.22 (-1.5-0.41)	-0.065 (-4.3-2)
Afghanistan	1990	Female	All Ages	12% (7-17%)	88% (83-93%)	0.015 (0.007-0.027)	0.037% (-0.056-0.12%)	-0.0059% (-0.21-0.14%)	-4.7 (-170-110)	2100 (-3200-6400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	1990	Female	15 to 19	11% (3.6-23%)	89% (77-96%)	0.014 (0.0034-0.039)	0.15% (0.045-0.27%)	0.065% (-0.092-0.23%)	0.98 (-1.5-3.4)	270 (80-490)
Afghanistan	1990	Female	20 to 24	14% (4.4-30%)	86% (70-96%)	0.021 (0.0054-0.055)	0.27% (0.039-0.51%)	0.072% (-0.25-0.4%)	1.3 (-4.5-7.3)	490 (74-940)
Afghanistan	1990	Female	25 to 29	16% (4.9-33%)	84% (67-95%)	0.026 (0.0069-0.061)	0.31% (0.0027-0.62%)	0.14% (-0.27-0.6%)	2.6 (-5.4-11)	540 (4.6-1100)
Afghanistan	1990	Female	30 to 34	18% (5.2-36%)	82% (64-95%)	0.031 (0.0086-0.073)	0.29% (-0.069-0.62%)	0.15% (-0.35-0.67%)	3.1 (-7.1-13)	480 (-120-1000)
Afghanistan	1990	Female	35 to 39	20% (5.8-37%)	80% (63-94%)	0.036 (0.01-0.078)	0.22% (-0.085-0.51%)	0.14% (-0.28-0.56%)	3.2 (-6.9-14)	370 (-150-880)
Afghanistan	1990	Female	40 to 44	20% (7.9-37%)	80% (63-92%)	0.04 (0.012-0.093)	0.16% (-0.22-0.48%)	0.12% (-0.37-0.55%)	3.2 (-11-16)	280 (-380-830)
Afghanistan	1990	Female	45 to 49	21% (7.9-39%)	79% (61-92%)	0.043 (0.012-0.11)	0.065% (-0.37-0.36%)	0.027% (-0.53-0.41%)	0.67 (-14-11)	91 (-530-500)
Afghanistan	1990	Female	50 to 54	22% (8.5-41%)	78% (59-92%)	0.044 (0.011-0.1)	-0.0014% (-0.56-0.33%)	-0.025% (-0.74-0.38%)	-0.74 (-20-11)	-1.6 (-730-420)
Afghanistan	1990	Female	55 to 59	23% (8.8-43%)	77% (57-91%)	0.043 (0.011-0.11)	-0.066% (-0.72-0.29%)	-0.085% (-0.84-0.34%)	-2.6 (-26-11)	-78 (-860-350)
Afghanistan	1990	Female	60 to 64	24% (9.5-44%)	76% (56-90%)	0.042 (0.009-0.11)	-0.12% (-0.83-0.27%)	-0.14% (-0.95-0.33%)	-4.9 (-34-11)	-140 (-950-310)
Afghanistan	1990	Female	65 to 69	25% (10-42%)	75% (58-90%)	0.037 (0.0083-0.094)	-0.11% (-0.7-0.26%)	-0.12% (-0.81-0.29%)	-5 (-33-12)	-120 (-780-280)
Afghanistan	1990	Female	70 to 74	23% (9.8-41%)	77% (59-90%)	0.029 (0.0059-0.082)	-0.074% (-0.7-0.24%)	-0.084% (-0.79-0.27%)	-3.3 (-32-11)	-61 (-580-190)
Afghanistan	1990	Female	75 to 79	21% (7.9-42%)	79% (58-92%)	0.024 (0.0041-0.075)	-0.051% (-0.69-0.22%)	-0.058% (-0.78-0.25%)	-1.7 (-23-7.1)	-23 (-320-99)
Afghanistan	1990	Female	80 to 84	19% (6.6-42%)	81% (58-93%)	0.022 (0.0036-0.069)	-0.037% (-0.65-0.23%)	-0.046% (-0.74-0.26%)	-0.6 (-9.7-3.4)	-5.7 (-100-35)
Afghanistan	1990	Female	85 to 89	19% (6.6-41%)	81% (59-93%)	0.021 (0.0037-0.067)	-0.046% (-0.56-0.2%)	-0.06% (-0.65-0.24%)	-0.51 (-5.4-2)	-3.3 (-40-15)
Afghanistan	1990	Female	90 to 94	22% (7.2-44%)	78% (56-93%)	0.024 (0.0039-0.073)	-0.054% (-0.53-0.19%)	-0.076% (-0.65-0.23%)	-0.26 (-2.2-0.8)	-1.1 (-11-4.1)
Afghanistan	1990	Female	95 plus	26% (8.7-50%)	74% (50-91%)	0.028 (0.0042-0.087)	-0.076% (-0.65-0.19%)	-0.11% (-0.77-0.22%)	-0.1 (-0.76-0.21)	-0.3 (-2.5-0.74)
Afghanistan	1990	Male	All Ages	8.7% (5.9-12%)	91% (88-94%)	0.017 (0.008-0.03)	0.13% (0.048-0.22%)	0.11% (-0.036-0.27%)	89 (-27-210)	7300 (2700-12000)
Afghanistan	1990	Male	15 to 19	7.7% (3.2-16%)	92% (84-97%)	0.0094 (0.0025-0.024)	0.24% (0.13-0.39%)	0.089% (0.021-0.25%)	1.7 (0.39-4.8)	480 (260-800)
Afghanistan	1990	Male	20 to 24	12% (4.9-23%)	88% (77-95%)	0.021 (0.0062-0.049)	0.48% (0.28-0.76%)	0.21% (0.041-0.57%)	5.3 (1-14)	1100 (620-1700)
Afghanistan	1990	Male	25 to 29	15% (6-27%)	85% (73-94%)	0.032 (0.0098-0.07)	0.55% (0.3-0.92%)	0.34% (0.052-0.8%)	7.7 (1.2-19)	1100 (600-1800)
Afghanistan	1990	Male	30 to 34	17% (6.4-31%)	83% (69-94%)	0.04 (0.013-0.088)	0.56% (0.27-0.97%)	0.41% (0.026-0.97%)	9.6 (0.66-23)	1000 (480-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	1990	Male	35 to 39	18% (7.4-33%)	82% (67-93%)	0.046 (0.015-0.1)	0.61% (0.22-1.1%)	0.5% (0.013-1.2%)	13 (0.35-33)	1100 (380-2100)
Afghanistan	1990	Male	40 to 44	19% (8.4-33%)	81% (67-92%)	0.049 (0.016-0.11)	0.54% (0.13-1%)	0.45% (-0.093-1.1%)	13 (-2.4-31)	890 (220-1700)
Afghanistan	1990	Male	45 to 49	18% (7-32%)	82% (68-93%)	0.051 (0.015-0.12)	0.45% (-0.041-0.93%)	0.4% (-0.22-1%)	11 (-6.1-30)	670 (-61-1400)
Afghanistan	1990	Male	50 to 54	17% (7.5-33%)	83% (67-92%)	0.05 (0.014-0.12)	0.31% (-0.19-0.7%)	0.27% (-0.33-0.75%)	7.6 (-9.1-22)	390 (-240-900)
Afghanistan	1990	Male	55 to 59	17% (7.4-32%)	83% (68-93%)	0.048 (0.013-0.12)	0.21% (-0.26-0.54%)	0.18% (-0.37-0.58%)	5.1 (-11-16)	220 (-270-580)
Afghanistan	1990	Male	60 to 64	16% (7.1-30%)	84% (70-93%)	0.044 (0.011-0.11)	0.14% (-0.29-0.45%)	0.12% (-0.39-0.5%)	4 (-13-17)	150 (-300-490)
Afghanistan	1990	Male	65 to 69	16% (7-29%)	84% (71-93%)	0.037 (0.0092-0.09)	0.11% (-0.25-0.35%)	0.098% (-0.32-0.38%)	3.5 (-11-14)	99 (-230-320)
Afghanistan	1990	Male	70 to 74	14% (6.4-25%)	86% (75-94%)	0.027 (0.0061-0.074)	0.11% (-0.23-0.32%)	0.11% (-0.29-0.36%)	3.5 (-8.7-11)	72 (-140-210)
Afghanistan	1990	Male	75 to 79	12% (5.2-23%)	88% (77-95%)	0.021 (0.0041-0.058)	0.1% (-0.16-0.27%)	0.11% (-0.21-0.3%)	2.3 (-4.8-6.6)	37 (-60-94)
Afghanistan	1990	Male	80 to 84	11% (4.3-22%)	89% (78-96%)	0.017 (0.0038-0.052)	0.11% (-0.11-0.26%)	0.11% (-0.15-0.29%)	1.2 (-1.5-3.1)	14 (-13-33)
Afghanistan	1990	Male	85 to 89	11% (4.3-22%)	89% (78-96%)	0.016 (0.0036-0.044)	0.087% (-0.098-0.21%)	0.084% (-0.14-0.23%)	0.62 (-1-1.7)	5.5 (-6.2-13)
Afghanistan	1990	Male	90 to 94	12% (4.7-24%)	88% (76-95%)	0.017 (0.0035-0.049)	0.062% (-0.11-0.17%)	0.048% (-0.16-0.19%)	0.16 (-0.53-0.61)	1.2 (-2.1-3.5)
Afghanistan	1990	Male	95 plus	15% (5.4-30%)	85% (70-95%)	0.019 (0.003-0.065)	0.062% (-0.15-0.18%)	0.029% (-0.22-0.17%)	0.024 (-0.19-0.14)	0.21 (-0.49-0.61)
Afghanistan	1995	Female	All Ages	11% (6.8-17%)	89% (83-93%)	0.015 (0.0078-0.025)	0.04% (-0.057-0.12%)	-0.0051% (-0.22-0.16%)	-5.5 (-240-170)	2900 (-4100-9100)
Afghanistan	1995	Female	15 to 19	10% (3.3-22%)	90% (78-97%)	0.013 (0.0032-0.034)	0.15% (0.052-0.26%)	0.062% (-0.077-0.21%)	1.4 (-1.7-4.7)	380 (130-670)
Afghanistan	1995	Female	20 to 24	13% (4-27%)	87% (73-96%)	0.02 (0.0055-0.049)	0.26% (0.032-0.46%)	0.071% (-0.22-0.38%)	1.9 (-6-10)	690 (84-1300)
Afghanistan	1995	Female	25 to 29	15% (4-31%)	85% (69-96%)	0.024 (0.0063-0.056)	0.3% (0.018-0.55%)	0.13% (-0.24-0.53%)	3.7 (-6.8-15)	740 (47-1400)
Afghanistan	1995	Female	30 to 34	17% (4.4-34%)	83% (66-96%)	0.029 (0.0075-0.066)	0.28% (-0.031-0.59%)	0.15% (-0.3-0.59%)	4.4 (-8.5-18)	650 (-68-1400)
Afghanistan	1995	Female	35 to 39	18% (5-35%)	82% (65-95%)	0.034 (0.0095-0.073)	0.22% (-0.086-0.51%)	0.14% (-0.27-0.55%)	4.7 (-9.2-19)	510 (-190-1200)
Afghanistan	1995	Female	40 to 44	19% (7.6-36%)	81% (64-92%)	0.038 (0.012-0.083)	0.16% (-0.19-0.47%)	0.12% (-0.33-0.53%)	4.4 (-12-19)	360 (-420-1000)
Afghanistan	1995	Female	45 to 49	20% (7.4-37%)	80% (63-93%)	0.041 (0.012-0.088)	0.069% (-0.36-0.35%)	0.034% (-0.5-0.4%)	1.5 (-21-18)	160 (-790-830)
Afghanistan	1995	Female	50 to 54	21% (9-39%)	79% (61-91%)	0.043 (0.012-0.1)	0.005% (-0.53-0.32%)	-0.017% (-0.65-0.38%)	-0.77 (-29-16)	9.8 (-1100-640)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	1995	Female	55 to 59	23% (9.3-43%)	77% (57-91%)	0.044 (0.011-0.11)	-0.069% (-0.72-0.3%)	-0.087% (-0.84-0.35%)	-4.3 (-40-17)	-130 (-1300-550)
Afghanistan	1995	Female	60 to 64	24% (9.4-44%)	76% (56-91%)	0.042 (0.0095-0.1)	-0.13% (-0.87-0.26%)	-0.15% (-0.98-0.31%)	-7.6 (-52-16)	-210 (-1400-440)
Afghanistan	1995	Female	65 to 69	24% (10-43%)	76% (57-90%)	0.036 (0.0086-0.092)	-0.1% (-0.68-0.25%)	-0.11% (-0.77-0.29%)	-6.6 (-44-16)	-150 (-1000-370)
Afghanistan	1995	Female	70 to 74	23% (9.4-41%)	77% (59-91%)	0.028 (0.006-0.075)	-0.064% (-0.63-0.24%)	-0.072% (-0.71-0.28%)	-4.3 (-42-17)	-79 (-760-300)
Afghanistan	1995	Female	75 to 79	20% (7.3-39%)	80% (61-93%)	0.023 (0.004-0.066)	-0.04% (-0.62-0.22%)	-0.046% (-0.69-0.25%)	-2.2 (-33-12)	-30 (-470-170)
Afghanistan	1995	Female	80 to 84	19% (6.6-40%)	81% (60-93%)	0.021 (0.0038-0.063)	-0.027% (-0.57-0.23%)	-0.035% (-0.64-0.27%)	-0.83 (-16-6.5)	-7.7 (-160-65)
Afghanistan	1995	Female	85 to 89	19% (7.1-41%)	81% (59-93%)	0.021 (0.0042-0.064)	-0.04% (-0.53-0.21%)	-0.052% (-0.62-0.25%)	-0.51 (-6.1-2.4)	-3.3 (-44-17)
Afghanistan	1995	Female	90 to 94	22% (8-44%)	78% (56-92%)	0.024 (0.0044-0.067)	-0.053% (-0.52-0.2%)	-0.074% (-0.63-0.24%)	-0.3 (-2.6-0.99)	-1.3 (-13-5.1)
Afghanistan	1995	Female	95 plus	27% (8.7-50%)	73% (50-91%)	0.028 (0.0049-0.084)	-0.076% (-0.64-0.2%)	-0.11% (-0.77-0.23%)	-0.15 (-1.1-0.32)	-0.43 (-3.5-1.1)
Afghanistan	1995	Male	All Ages	9.2% (6.3-13%)	91% (87-94%)	0.017 (0.0094-0.029)	0.14% (0.044-0.24%)	0.12% (-0.052-0.29%)	130 (-61-320)	10000 (3400-18000)
Afghanistan	1995	Male	15 to 19	7.8% (3.3-15%)	92% (85-97%)	0.0092 (0.0025-0.023)	0.22% (0.12-0.35%)	0.079% (0.019-0.22%)	2.5 (0.6-7)	690 (380-1100)
Afghanistan	1995	Male	20 to 24	12% (4.9-23%)	88% (77-95%)	0.02 (0.0068-0.046)	0.43% (0.25-0.67%)	0.18% (0.038-0.46%)	7.6 (1.6-19)	1500 (870-2400)
Afghanistan	1995	Male	25 to 29	15% (6.3-28%)	85% (72-94%)	0.032 (0.011-0.07)	0.51% (0.27-0.83%)	0.3% (0.036-0.71%)	11 (1.4-26)	1500 (840-2600)
Afghanistan	1995	Male	30 to 34	17% (6.7-32%)	83% (68-93%)	0.041 (0.015-0.089)	0.53% (0.24-0.92%)	0.37% (0.0056-0.88%)	13 (0.21-32)	1400 (630-2500)
Afghanistan	1995	Male	35 to 39	19% (7.4-34%)	81% (66-93%)	0.046 (0.016-0.098)	0.57% (0.2-1.1%)	0.47% (-0.0081-1.2%)	18 (-0.29-45)	1400 (490-2800)
Afghanistan	1995	Male	40 to 44	19% (8.4-33%)	81% (67-92%)	0.049 (0.017-0.11)	0.5% (0.086-0.98%)	0.42% (-0.11-1.1%)	17 (-4.6-44)	1200 (220-2400)
Afghanistan	1995	Male	45 to 49	19% (7.6-34%)	81% (66-92%)	0.051 (0.015-0.11)	0.43% (-0.1-0.93%)	0.38% (-0.29-1%)	17 (-13-47)	980 (-240-2200)
Afghanistan	1995	Male	50 to 54	18% (7.7-34%)	82% (66-92%)	0.051 (0.016-0.12)	0.29% (-0.21-0.69%)	0.25% (-0.35-0.75%)	12 (-18-36)	620 (-460-1500)
Afghanistan	1995	Male	55 to 59	17% (7.7-32%)	83% (68-92%)	0.049 (0.014-0.12)	0.2% (-0.3-0.56%)	0.17% (-0.41-0.63%)	8.2 (-20-30)	360 (-510-1000)
Afghanistan	1995	Male	60 to 64	16% (7.2-31%)	84% (69-93%)	0.044 (0.011-0.11)	0.13% (-0.32-0.46%)	0.11% (-0.44-0.5%)	5.6 (-22-26)	200 (-490-740)
Afghanistan	1995	Male	65 to 69	16% (6.9-28%)	84% (72-93%)	0.036 (0.0098-0.087)	0.11% (-0.22-0.35%)	0.098% (-0.29-0.39%)	5.2 (-15-21)	150 (-320-490)
Afghanistan	1995	Male	70 to 74	14% (6.5-26%)	86% (74-93%)	0.027 (0.0068-0.067)	0.11% (-0.22-0.33%)	0.11% (-0.27-0.36%)	5.8 (-14-19)	120 (-220-350)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	1995	Male	75 to 79	12% (5.3-23%)	88% (77-95%)	0.02 (0.0045-0.054)	0.1% (-0.15-0.26%)	0.11% (-0.18-0.29%)	3.9 (-7.1-11)	60 (-89-150)
Afghanistan	1995	Male	80 to 84	11% (4.3-21%)	89% (79-96%)	0.017 (0.0042-0.047)	0.11% (-0.1-0.26%)	0.11% (-0.13-0.29%)	2 (-2.4-5)	23 (-21-54)
Afghanistan	1995	Male	85 to 89	11% (4.2-21%)	89% (79-96%)	0.015 (0.0037-0.041)	0.086% (-0.074-0.21%)	0.083% (-0.11-0.23%)	0.68 (-0.94-1.9)	6.1 (-5.3-15)
Afghanistan	1995	Male	90 to 94	12% (4.7-24%)	88% (76-95%)	0.016 (0.0036-0.044)	0.063% (-0.098-0.17%)	0.049% (-0.15-0.18%)	0.19 (-0.58-0.69)	1.4 (-2.2-3.9)
Afghanistan	1995	Male	95 plus	15% (5.3-31%)	85% (69-95%)	0.018 (0.0032-0.058)	0.062% (-0.12-0.18%)	0.03% (-0.2-0.17%)	0.036 (-0.25-0.2)	0.31 (-0.6-0.88)
Afghanistan	2000	Female	All Ages	11% (6.7-17%)	89% (83-93%)	0.018 (0.011-0.028)	0.031% (-0.088-0.14%)	-0.033% (-0.28-0.17%)	-41 (-340-210)	2500 (-6800-11000)
Afghanistan	2000	Female	15 to 19	11% (3.4-24%)	89% (76-97%)	0.016 (0.0044-0.04)	0.16% (0.039-0.28%)	0.066% (-0.096-0.26%)	1.7 (-2.3-6.5)	450 (110-830)
Afghanistan	2000	Female	20 to 24	14% (4.1-30%)	86% (70-96%)	0.026 (0.0076-0.059)	0.26% (0.016-0.51%)	0.074% (-0.28-0.43%)	2.3 (-8.9-13)	800 (47-1600)
Afghanistan	2000	Female	25 to 29	16% (4.5-32%)	84% (68-95%)	0.031 (0.0092-0.066)	0.3% (-0.041-0.61%)	0.13% (-0.37-0.62%)	4.3 (-11-20)	860 (-120-1800)
Afghanistan	2000	Female	30 to 34	17% (5-36%)	83% (64-95%)	0.036 (0.012-0.073)	0.27% (-0.083-0.64%)	0.15% (-0.39-0.7%)	5.1 (-13-24)	740 (-210-1800)
Afghanistan	2000	Female	35 to 39	19% (5.2-37%)	81% (63-95%)	0.043 (0.013-0.087)	0.2% (-0.19-0.57%)	0.13% (-0.36-0.63%)	5.2 (-14-25)	560 (-480-1500)
Afghanistan	2000	Female	40 to 44	20% (7.3-36%)	80% (64-93%)	0.047 (0.016-0.097)	0.14% (-0.29-0.49%)	0.099% (-0.44-0.6%)	4.2 (-18-24)	360 (-710-1300)
Afghanistan	2000	Female	45 to 49	20% (7.1-38%)	80% (62-93%)	0.05 (0.016-0.1)	0.029% (-0.44-0.37%)	-0.008% (-0.58-0.43%)	-0.38 (-28-21)	74 (-1100-930)
Afghanistan	2000	Female	50 to 54	21% (8.1-39%)	79% (61-92%)	0.051 (0.015-0.11)	-0.046% (-0.63-0.32%)	-0.072% (-0.78-0.39%)	-4.3 (-48-24)	-120 (-1700-870)
Afghanistan	2000	Female	55 to 59	22% (9-41%)	78% (59-91%)	0.051 (0.014-0.12)	-0.13% (-0.79-0.26%)	-0.15% (-0.92-0.31%)	-9.1 (-55-19)	-290 (-1800-580)
Afghanistan	2000	Female	60 to 64	23% (9.3-43%)	77% (57-91%)	0.049 (0.012-0.12)	-0.2% (-0.99-0.23%)	-0.22% (-1.1-0.28%)	-14 (-71-18)	-380 (-1900-460)
Afghanistan	2000	Female	65 to 69	24% (10-43%)	76% (57-90%)	0.043 (0.011-0.11)	-0.18% (-0.81-0.21%)	-0.2% (-0.93-0.25%)	-13 (-61-16)	-300 (-1400-350)
Afghanistan	2000	Female	70 to 74	23% (9.4-40%)	77% (60-91%)	0.034 (0.0079-0.089)	-0.14% (-0.76-0.22%)	-0.15% (-0.86-0.25%)	-10 (-58-17)	-190 (-1100-310)
Afghanistan	2000	Female	75 to 79	20% (7.7-40%)	80% (60-92%)	0.028 (0.0054-0.08)	-0.11% (-0.76-0.22%)	-0.12% (-0.86-0.24%)	-7 (-52-14)	-98 (-720-200)
Afghanistan	2000	Female	80 to 84	19% (6.2-39%)	81% (61-94%)	0.025 (0.0049-0.074)	-0.083% (-0.7-0.22%)	-0.097% (-0.78-0.25%)	-3.5 (-28-9.2)	-35 (-290-94)
Afghanistan	2000	Female	85 to 89	19% (6.3-41%)	81% (59-94%)	0.025 (0.005-0.076)	-0.094% (-0.66-0.2%)	-0.11% (-0.77-0.24%)	-1.5 (-10-3.2)	-11 (-74-22)
Afghanistan	2000	Female	90 to 94	22% (6.9-44%)	78% (56-93%)	0.029 (0.0054-0.085)	-0.11% (-0.63-0.19%)	-0.14% (-0.78-0.23%)	-0.5 (-2.8-0.82)	-2.4 (-14-4.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	2000	Female	95 plus	26% (8.2-51%)	74% (49-92%)	0.034 (0.0058-0.1)	-0.14% (-0.79-0.2%)	-0.18% (-0.96-0.23%)	-0.22 (-1.2-0.29)	-0.69 (-4-0.99)
Afghanistan	2000	Male	All Ages	9% (6.2-13%)	91% (87-94%)	0.02 (0.013-0.03)	0.15% (0.032-0.27%)	0.12% (-0.085-0.32%)	150 (-100-400)	12000 (2700-22000)
Afghanistan	2000	Male	15 to 19	7.8% (3.3-16%)	92% (84-97%)	0.011 (0.0032-0.025)	0.24% (0.13-0.38%)	0.099% (0.021-0.26%)	3.5 (0.76-9)	850 (470-1400)
Afghanistan	2000	Male	20 to 24	12% (5-24%)	88% (76-95%)	0.025 (0.009-0.052)	0.47% (0.27-0.75%)	0.22% (0.041-0.57%)	10 (2-26)	1900 (1100-3100)
Afghanistan	2000	Male	25 to 29	15% (6.1-28%)	85% (72-94%)	0.039 (0.014-0.077)	0.56% (0.28-0.96%)	0.35% (0.022-0.82%)	15 (0.94-35)	1900 (930-3300)
Afghanistan	2000	Male	30 to 34	17% (6.4-32%)	83% (68-94%)	0.049 (0.019-0.095)	0.56% (0.22-1%)	0.41% (-0.028-1%)	17 (-1.1-42)	1700 (650-3200)
Afghanistan	2000	Male	35 to 39	19% (7.2-34%)	81% (66-93%)	0.055 (0.02-0.11)	0.59% (0.17-1.2%)	0.5% (-0.059-1.2%)	21 (-2.6-54)	1700 (460-3300)
Afghanistan	2000	Male	40 to 44	19% (8.7-33%)	81% (67-91%)	0.059 (0.022-0.12)	0.5% (0.031-1%)	0.42% (-0.17-1.1%)	20 (-8.3-52)	1400 (87-2900)
Afghanistan	2000	Male	45 to 49	18% (7.5-33%)	82% (67-92%)	0.061 (0.02-0.13)	0.41% (-0.17-0.96%)	0.36% (-0.37-1.1%)	19 (-20-57)	1100 (-480-2600)
Afghanistan	2000	Male	50 to 54	18% (7.8-33%)	82% (67-92%)	0.061 (0.02-0.13)	0.27% (-0.29-0.72%)	0.22% (-0.48-0.77%)	13 (-30-45)	680 (-820-1800)
Afghanistan	2000	Male	55 to 59	17% (7.4-33%)	83% (67-93%)	0.059 (0.017-0.14)	0.17% (-0.39-0.57%)	0.14% (-0.53-0.63%)	8.3 (-32-39)	380 (-870-1300)
Afghanistan	2000	Male	60 to 64	16% (7.3-31%)	84% (69-93%)	0.053 (0.014-0.13)	0.098% (-0.42-0.45%)	0.076% (-0.53-0.5%)	4.6 (-33-30)	190 (-810-870)
Afghanistan	2000	Male	65 to 69	16% (7.2-28%)	84% (72-93%)	0.043 (0.013-0.096)	0.081% (-0.3-0.37%)	0.068% (-0.39-0.4%)	3.9 (-23-23)	120 (-460-550)
Afghanistan	2000	Male	70 to 74	14% (6.6-25%)	86% (75-93%)	0.032 (0.0093-0.077)	0.094% (-0.26-0.34%)	0.091% (-0.31-0.38%)	5.4 (-19-22)	110 (-320-430)
Afghanistan	2000	Male	75 to 79	12% (5.5-23%)	88% (77-94%)	0.024 (0.0062-0.065)	0.086% (-0.2-0.26%)	0.085% (-0.24-0.3%)	4 (-12-13)	64 (-150-190)
Afghanistan	2000	Male	80 to 84	11% (4.6-22%)	89% (78-95%)	0.02 (0.0052-0.053)	0.095% (-0.14-0.27%)	0.095% (-0.18-0.3%)	2.5 (-4.7-7.7)	29 (-45-82)
Afghanistan	2000	Male	85 to 89	11% (4.4-21%)	89% (79-96%)	0.019 (0.0049-0.05)	0.071% (-0.12-0.2%)	0.064% (-0.18-0.22%)	0.68 (-1.8-2.4)	6.4 (-11-18)
Afghanistan	2000	Male	90 to 94	12% (4.9-24%)	88% (76-95%)	0.019 (0.0049-0.052)	0.049% (-0.13-0.17%)	0.032% (-0.19-0.18%)	0.1 (-0.6-0.57)	0.95 (-2.5-3.2)
Afghanistan	2000	Male	95 plus	15% (5.5-30%)	85% (70-94%)	0.022 (0.0046-0.065)	0.046% (-0.16-0.17%)	0.0092% (-0.24-0.16%)	0.0099 (-0.26-0.17)	0.2 (-0.68-0.74)
Afghanistan	2005	Female	All Ages	11% (6.9-17%)	89% (83-93%)	0.03 (0.022-0.041)	-0.0044% (-0.17-0.16%)	-0.14% (-0.48-0.17%)	-180 (-640-230)	-360 (-14000-13000)
Afghanistan	2005	Female	15 to 19	11% (3.5-23%)	89% (77-97%)	0.027 (0.0079-0.061)	0.19% (0.031-0.36%)	0.091% (-0.15-0.36%)	2.5 (-4.1-9.9)	610 (97-1200)
Afghanistan	2005	Female	20 to 24	14% (4.5-29%)	86% (71-95%)	0.042 (0.013-0.091)	0.28% (-0.0067-0.62%)	0.085% (-0.36-0.59%)	3 (-13-21)	1100 (-28-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	2005	Female	25 to 29	15% (4.2-32%)	85% (68-96%)	0.051 (0.016-0.1)	0.31% (-0.12-0.73%)	0.14% (-0.53-0.74%)	5.3 (-19-29)	1100 (-420-2600)
Afghanistan	2005	Female	30 to 34	17% (4.7-34%)	83% (66-95%)	0.06 (0.021-0.11)	0.27% (-0.2-0.75%)	0.14% (-0.56-0.85%)	5.6 (-22-33)	870 (-630-2400)
Afghanistan	2005	Female	35 to 39	19% (5.2-35%)	81% (65-95%)	0.07 (0.023-0.13)	0.19% (-0.28-0.67%)	0.12% (-0.5-0.81%)	5.4 (-23-35)	600 (-890-2100)
Afghanistan	2005	Female	40 to 44	20% (7.4-35%)	80% (65-93%)	0.077 (0.029-0.15)	0.087% (-0.47-0.55%)	0.044% (-0.68-0.7%)	2.1 (-32-32)	260 (-1300-1600)
Afghanistan	2005	Female	45 to 49	20% (7.1-37%)	80% (63-93%)	0.082 (0.027-0.17)	-0.077% (-0.64-0.37%)	-0.12% (-0.86-0.46%)	-6.5 (-44-24)	-210 (-1800-1100)
Afghanistan	2005	Female	50 to 54	21% (8-39%)	79% (61-92%)	0.086 (0.026-0.18)	-0.22% (-1-0.29%)	-0.26% (-1.2-0.36%)	-16 (-74-22)	-620 (-2800-850)
Afghanistan	2005	Female	55 to 59	22% (9.1-42%)	78% (58-91%)	0.086 (0.024-0.2)	-0.36% (-1.3-0.17%)	-0.4% (-1.5-0.25%)	-31 (-120-19)	-1100 (-4000-510)
Afghanistan	2005	Female	60 to 64	23% (9.6-44%)	77% (56-90%)	0.083 (0.021-0.2)	-0.45% (-1.5-0.14%)	-0.5% (-1.7-0.17%)	-37 (-130-12)	-1100 (-3600-320)
Afghanistan	2005	Female	65 to 69	24% (11-42%)	76% (58-89%)	0.073 (0.02-0.16)	-0.43% (-1.4-0.1%)	-0.47% (-1.6-0.13%)	-38 (-120-10)	-890 (-2800-220)
Afghanistan	2005	Female	70 to 74	23% (9.7-40%)	77% (60-90%)	0.058 (0.015-0.15)	-0.37% (-1.3-0.14%)	-0.4% (-1.5-0.16%)	-31 (-110-13)	-580 (-2000-220)
Afghanistan	2005	Female	75 to 79	20% (7.8-39%)	80% (61-92%)	0.047 (0.01-0.13)	-0.31% (-1.3-0.13%)	-0.34% (-1.4-0.15%)	-22 (-92-9.8)	-320 (-1300-130)
Afghanistan	2005	Female	80 to 84	19% (6.4-40%)	81% (60-94%)	0.042 (0.0088-0.12)	-0.25% (-1.1-0.17%)	-0.28% (-1.3-0.2%)	-12 (-55-8.5)	-130 (-580-84)
Afghanistan	2005	Female	85 to 89	19% (6.3-40%)	81% (60-94%)	0.042 (0.0086-0.12)	-0.25% (-1.1-0.16%)	-0.29% (-1.3-0.2%)	-5.4 (-24-3.6)	-40 (-170-25)
Afghanistan	2005	Female	90 to 94	22% (7-44%)	78% (56-93%)	0.047 (0.0093-0.13)	-0.27% (-1.1-0.13%)	-0.33% (-1.3-0.15%)	-1.5 (-6.1-0.72)	-7.6 (-31-3.7)
Afghanistan	2005	Female	95 plus	26% (8.1-50%)	74% (50-92%)	0.056 (0.01-0.16)	-0.33% (-1.3-0.12%)	-0.4% (-1.5-0.14%)	-0.42 (-1.6-0.14)	-1.4 (-5.2-0.51)
Afghanistan	2005	Male	All Ages	8.9% (6-13%)	91% (87-94%)	0.034 (0.025-0.044)	0.19% (0.035-0.35%)	0.14% (-0.13-0.39%)	190 (-180-540)	17000 (3100-31000)
Afghanistan	2005	Male	15 to 19	7.8% (3.2-16%)	92% (84-97%)	0.018 (0.0058-0.038)	0.32% (0.19-0.53%)	0.18% (0.042-0.44%)	7 (1.6-17)	1200 (710-2100)
Afghanistan	2005	Male	20 to 24	12% (4.8-23%)	88% (77-95%)	0.04 (0.015-0.081)	0.63% (0.35-1%)	0.39% (0.066-0.94%)	20 (3.4-47)	2900 (1600-4800)
Afghanistan	2005	Male	25 to 29	15% (6-27%)	85% (73-94%)	0.062 (0.024-0.12)	0.74% (0.36-1.3%)	0.56% (0.055-1.3%)	28 (2.8-64)	3000 (1500-5300)
Afghanistan	2005	Male	30 to 34	17% (6.6-32%)	83% (68-93%)	0.079 (0.034-0.14)	0.73% (0.3-1.4%)	0.62% (0.037-1.4%)	29 (1.7-68)	2600 (1100-4800)
Afghanistan	2005	Male	35 to 39	18% (7.4-33%)	82% (67-93%)	0.09 (0.036-0.17)	0.72% (0.22-1.4%)	0.66% (-0.034-1.6%)	32 (-1.7-78)	2400 (750-4800)
Afghanistan	2005	Male	40 to 44	19% (8.5-32%)	81% (68-91%)	0.096 (0.038-0.19)	0.57% (0.031-1.1%)	0.51% (-0.19-1.3%)	27 (-10-72)	1800 (98-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	2005	Male	45 to 49	18% (7.2-33%)	82% (67-93%)	0.1 (0.033-0.2)	0.43% (-0.19-1%)	0.39% (-0.46-1.1%)	24 (-28-69)	1400 (-630-3200)
Afghanistan	2005	Male	50 to 54	17% (7.6-33%)	83% (67-92%)	0.098 (0.034-0.2)	0.24% (-0.42-0.73%)	0.19% (-0.61-0.81%)	13 (-44-57)	730 (-1300-2300)
Afghanistan	2005	Male	55 to 59	16% (7.3-31%)	84% (69-93%)	0.094 (0.03-0.2)	0.13% (-0.51-0.59%)	0.087% (-0.66-0.64%)	6.3 (-49-48)	350 (-1400-1600)
Afghanistan	2005	Male	60 to 64	16% (7.1-30%)	84% (70-93%)	0.084 (0.023-0.2)	0.043% (-0.58-0.45%)	0.012% (-0.72-0.49%)	0.85 (-57-38)	100 (-1400-1100)
Afghanistan	2005	Male	65 to 69	15% (6.9-28%)	85% (72-93%)	0.069 (0.022-0.15)	0.013% (-0.5-0.35%)	-0.011% (-0.59-0.39%)	-0.77 (-43-29)	24 (-920-670)
Afghanistan	2005	Male	70 to 74	14% (6.2-24%)	86% (76-94%)	0.051 (0.015-0.13)	0.045% (-0.37-0.34%)	0.035% (-0.46-0.37%)	2.3 (-29-24)	60 (-500-440)
Afghanistan	2005	Male	75 to 79	12% (4.9-23%)	88% (77-95%)	0.039 (0.01-0.1)	0.046% (-0.33-0.28%)	0.039% (-0.39-0.3%)	2.1 (-21-16)	39 (-280-230)
Afghanistan	2005	Male	80 to 84	11% (4.3-21%)	89% (79-96%)	0.033 (0.009-0.085)	0.067% (-0.25-0.28%)	0.06% (-0.31-0.31%)	1.9 (-9.7-9.4)	25 (-92-100)
Afghanistan	2005	Male	85 to 89	11% (4.2-21%)	89% (79-96%)	0.03 (0.0081-0.079)	0.041% (-0.21-0.21%)	0.026% (-0.28-0.23%)	0.39 (-4.1-3.4)	5.2 (-27-27)
Afghanistan	2005	Male	90 to 94	12% (4.9-24%)	88% (76-95%)	0.032 (0.0081-0.085)	0.013% (-0.23-0.17%)	-0.015% (-0.31-0.18%)	-0.063 (-1.3-0.72)	0.33 (-5.6-4.2)
Afghanistan	2005	Male	95 plus	15% (5.5-31%)	85% (69-95%)	0.037 (0.0075-0.11)	0.00059% (-0.3-0.17%)	-0.052% (-0.41-0.16%)	-0.047 (-0.37-0.14)	0.0025 (-1.1-0.61)
Afghanistan	2010	Female	All Ages	11% (6.4-17%)	89% (83-94%)	0.05 (0.034-0.07)	-0.037% (-0.26-0.16%)	-0.25% (-0.7-0.13%)	-320 (-930-170)	-2900 (-20000-13000)
Afghanistan	2010	Female	15 to 19	10% (3.1-22%)	90% (78-97%)	0.044 (0.011-0.1)	0.21% (0.063-0.4%)	0.11% (-0.1-0.41%)	3.2 (-3.1-11)	730 (220-1400)
Afghanistan	2010	Female	20 to 24	13% (3.8-27%)	87% (73-96%)	0.069 (0.021-0.15)	0.35% (0.047-0.68%)	0.13% (-0.32-0.66%)	4.7 (-11-22)	1300 (170-2500)
Afghanistan	2010	Female	25 to 29	15% (4.2-31%)	85% (69-96%)	0.084 (0.027-0.17)	0.37% (-0.045-0.79%)	0.21% (-0.45-0.88%)	8.3 (-18-37)	1400 (-170-3000)
Afghanistan	2010	Female	30 to 34	17% (4.7-35%)	83% (65-95%)	0.1 (0.033-0.19)	0.31% (-0.24-0.82%)	0.2% (-0.58-0.98%)	8.6 (-25-43)	1100 (-850-3100)
Afghanistan	2010	Female	35 to 39	19% (5.1-36%)	81% (64-95%)	0.12 (0.038-0.22)	0.22% (-0.32-0.77%)	0.17% (-0.58-0.99%)	8.4 (-27-48)	790 (-1100-2800)
Afghanistan	2010	Female	40 to 44	20% (6.9-37%)	80% (63-93%)	0.13 (0.046-0.26)	0.087% (-0.52-0.65%)	0.062% (-0.73-0.89%)	3.2 (-39-45)	280 (-1700-2100)
Afghanistan	2010	Female	45 to 49	20% (7.1-37%)	80% (63-93%)	0.14 (0.047-0.3)	-0.13% (-0.9-0.4%)	-0.17% (-1.1-0.59%)	-9.7 (-64-31)	-390 (-2800-1200)
Afghanistan	2010	Female	50 to 54	21% (8.6-39%)	79% (61-91%)	0.15 (0.045-0.31)	-0.34% (-1.3-0.27%)	-0.39% (-1.5-0.37%)	-25 (-99-24)	-1000 (-3800-810)
Afghanistan	2010	Female	55 to 59	22% (9.3-40%)	78% (60-91%)	0.15 (0.042-0.31)	-0.54% (-1.7-0.16%)	-0.61% (-2-0.22%)	-46 (-150-16)	-1600 (-4900-430)
Afghanistan	2010	Female	60 to 64	23% (9.2-41%)	77% (59-91%)	0.14 (0.035-0.31)	-0.66% (-1.8-0.048%)	-0.74% (-2.1-0.1%)	-67 (-200-9.6)	-1900 (-5600-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	2010	Female	65 to 69	23% (10-40%)	77% (60-90%)	0.12 (0.034-0.27)	-0.69% (-1.9--0.0019%)	-0.77% (-2.1-0.018%)	-68 (-180-1.6)	-1600 (-4300--4.3)
Afghanistan	2010	Female	70 to 74	22% (10-39%)	78% (61-90%)	0.096 (0.026-0.24)	-0.61% (-1.7-0.049%)	-0.68% (-1.9-0.079%)	-59 (-170-6.7)	-1100 (-3200-82)
Afghanistan	2010	Female	75 to 79	20% (7.6-38%)	80% (62-92%)	0.078 (0.017-0.22)	-0.53% (-1.8-0.056%)	-0.6% (-2-0.077%)	-44 (-150-5.5)	-620 (-2200-65)
Afghanistan	2010	Female	80 to 84	18% (6.3-38%)	82% (62-94%)	0.071 (0.015-0.2)	-0.45% (-1.7-0.11%)	-0.51% (-1.9-0.13%)	-26 (-98-6.5)	-270 (-1000-64)
Afghanistan	2010	Female	85 to 89	19% (5.6-40%)	81% (60-94%)	0.071 (0.014-0.21)	-0.46% (-1.6-0.089%)	-0.53% (-1.8-0.097%)	-12 (-41-2.2)	-86 (-300-17)
Afghanistan	2010	Female	90 to 94	22% (6.8-44%)	78% (56-93%)	0.082 (0.015-0.23)	-0.5% (-1.7-0.062%)	-0.6% (-2-0.065%)	-3.8 (-13-0.41)	-19 (-65-2.3)
Afghanistan	2010	Female	95 plus	26% (8.2-52%)	74% (48-92%)	0.097 (0.017-0.28)	-0.58% (-1.9-0.051%)	-0.7% (-2.3-0.048%)	-0.92 (-3-0.062)	-3 (-10-0.27)
Afghanistan	2010	Male	All Ages	8.8% (6-12%)	91% (88-94%)	0.057 (0.041-0.077)	0.27% (0.082-0.46%)	0.21% (-0.092-0.51%)	300 (-130-700)	24000 (6800-40000)
Afghanistan	2010	Male	15 to 19	7.6% (3.2-14%)	92% (86-97%)	0.03 (0.0095-0.067)	0.37% (0.2-0.6%)	0.24% (0.058-0.56%)	10 (2.5-25)	1600 (850-2700)
Afghanistan	2010	Male	20 to 24	12% (4.6-22%)	88% (78-95%)	0.067 (0.024-0.14)	0.74% (0.39-1.3%)	0.53% (0.12-1.2%)	31 (6.8-71)	3700 (2000-6500)
Afghanistan	2010	Male	25 to 29	14% (6-27%)	86% (73-94%)	0.11 (0.042-0.19)	0.93% (0.45-1.6%)	0.8% (0.2-1.6%)	45 (12-91)	4200 (2000-7200)
Afghanistan	2010	Male	30 to 34	17% (6.3-31%)	83% (69-94%)	0.14 (0.061-0.25)	0.95% (0.46-1.7%)	0.91% (0.2-1.9%)	50 (11-100)	4000 (1900-7100)
Afghanistan	2010	Male	35 to 39	18% (7-33%)	82% (67-93%)	0.15 (0.06-0.3)	0.94% (0.34-1.8%)	0.93% (0.14-2%)	52 (7.5-110)	3500 (1300-6600)
Afghanistan	2010	Male	40 to 44	18% (8.2-32%)	82% (68-92%)	0.16 (0.063-0.32)	0.75% (0.12-1.5%)	0.73% (-0.09-1.7%)	43 (-5.4-100)	2700 (460-5200)
Afghanistan	2010	Male	45 to 49	18% (7.1-32%)	82% (68-93%)	0.17 (0.055-0.35)	0.58% (-0.079-1.3%)	0.57% (-0.26-1.6%)	38 (-19-100)	2000 (-270-4500)
Afghanistan	2010	Male	50 to 54	17% (7.5-33%)	83% (67-93%)	0.17 (0.055-0.35)	0.32% (-0.38-0.93%)	0.29% (-0.57-1%)	21 (-43-77)	1000 (-1200-3100)
Afghanistan	2010	Male	55 to 59	16% (7.1-31%)	84% (69-93%)	0.16 (0.051-0.34)	0.17% (-0.51-0.68%)	0.14% (-0.69-0.77%)	11 (-57-63)	510 (-1700-2100)
Afghanistan	2010	Male	60 to 64	16% (6.7-30%)	84% (70-93%)	0.14 (0.042-0.32)	0.054% (-0.58-0.55%)	0.025% (-0.74-0.6%)	2 (-63-52)	140 (-1600-1500)
Afghanistan	2010	Male	65 to 69	15% (6.5-27%)	85% (73-93%)	0.12 (0.038-0.25)	-0.028% (-0.58-0.37%)	-0.058% (-0.73-0.42%)	-4.8 (-59-35)	-61 (-1200-790)
Afghanistan	2010	Male	70 to 74	14% (6-24%)	86% (76-94%)	0.087 (0.026-0.2)	-0.00078% (-0.45-0.34%)	-0.017% (-0.55-0.38%)	-1.3 (-40-28)	-0.98 (-690-510)
Afghanistan	2010	Male	75 to 79	12% (5.2-22%)	88% (78-95%)	0.066 (0.017-0.17)	0.0057% (-0.44-0.27%)	-0.0078% (-0.52-0.3%)	-0.46 (-30-17)	5.1 (-400-240)
Afghanistan	2010	Male	80 to 84	11% (4.1-21%)	89% (79-96%)	0.055 (0.015-0.14)	0.034% (-0.34-0.28%)	0.02% (-0.4-0.31%)	0.69 (-14-11)	14 (-130-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	2010	Male	85 to 89	10% (4.2-20%)	90% (80-96%)	0.051 (0.014-0.13)	0.0054% (-0.31-0.22%)	-0.02% (-0.4-0.23%)	-0.36 (-7.2-4.2)	0.85 (-48-33)
Afghanistan	2010	Male	90 to 94	12% (4.9-23%)	88% (77-95%)	0.054 (0.014-0.14)	-0.031% (-0.35-0.17%)	-0.074% (-0.46-0.16%)	-0.44 (-2.7-0.95)	-1.1 (-12-5.8)
Afghanistan	2010	Male	95 plus	14% (5.4-30%)	86% (70-95%)	0.062 (0.012-0.18)	-0.052% (-0.44-0.17%)	-0.12% (-0.59-0.14%)	-0.15 (-0.7-0.17)	-0.25 (-2.1-0.84)
Afghanistan	2016	Female	All Ages	11% (6.8-17%)	89% (83-93%)	0.067 (0.038-0.11)	-0.028% (-0.27-0.19%)	-0.28% (-0.77-0.13%)	-400 (-1100-180)	-2400 (-22000-16000)
Afghanistan	2016	Female	15 to 19	10% (3.3-22%)	90% (78-97%)	0.057 (0.015-0.14)	0.23% (0.092-0.41%)	0.14% (-0.053-0.44%)	5.4 (-2-17)	1000 (400-2000)
Afghanistan	2016	Female	20 to 24	13% (4-27%)	87% (73-96%)	0.09 (0.025-0.21)	0.4% (0.11-0.74%)	0.2% (-0.22-0.72%)	8.3 (-8.8-31)	1800 (470-3500)
Afghanistan	2016	Female	25 to 29	15% (4.4-31%)	85% (69-96%)	0.11 (0.031-0.24)	0.44% (0.056-0.89%)	0.28% (-0.28-0.99%)	12 (-12-45)	1900 (230-3800)
Afghanistan	2016	Female	30 to 34	17% (4.3-34%)	83% (66-96%)	0.13 (0.039-0.27)	0.37% (-0.082-0.88%)	0.27% (-0.4-1%)	13 (-21-52)	1600 (-360-3700)
Afghanistan	2016	Female	35 to 39	18% (4.9-34%)	82% (66-95%)	0.15 (0.042-0.31)	0.28% (-0.23-0.79%)	0.25% (-0.42-1%)	14 (-25-58)	1200 (-970-3300)
Afghanistan	2016	Female	40 to 44	19% (7.2-35%)	81% (65-93%)	0.17 (0.055-0.36)	0.15% (-0.48-0.76%)	0.15% (-0.74-1%)	8.8 (-41-59)	570 (-1800-2900)
Afghanistan	2016	Female	45 to 49	20% (6.9-37%)	80% (63-93%)	0.18 (0.054-0.41)	-0.073% (-0.82-0.65%)	-0.092% (-1-0.84%)	-6 (-69-54)	-260 (-2900-2300)
Afghanistan	2016	Female	50 to 54	21% (7.4-39%)	79% (61-93%)	0.19 (0.051-0.44)	-0.29% (-1.3-0.48%)	-0.33% (-1.6-0.64%)	-24 (-110-47)	-1000 (-4500-1600)
Afghanistan	2016	Female	55 to 59	22% (8.1-41%)	78% (59-92%)	0.19 (0.049-0.45)	-0.53% (-1.8-0.21%)	-0.59% (-2.1-0.33%)	-49 (-170-27)	-1700 (-6000-720)
Afghanistan	2016	Female	60 to 64	23% (8.9-43%)	77% (57-91%)	0.18 (0.044-0.45)	-0.71% (-2.1-0.062%)	-0.79% (-2.4-0.12%)	-72 (-220-11)	-2100 (-6400-200)
Afghanistan	2016	Female	65 to 69	23% (9.9-41%)	77% (59-90%)	0.16 (0.042-0.37)	-0.79% (-2--0.014%)	-0.87% (-2.3--0.00082%)	-98 (-260--0.0086)	-2300 (-6000--39)
Afghanistan	2016	Female	70 to 74	22% (9.7-38%)	78% (62-90%)	0.13 (0.03-0.32)	-0.73% (-1.9--0.0092%)	-0.81% (-2.1-0.006%)	-85 (-220-0.58)	-1600 (-4100--19)
Afghanistan	2016	Female	75 to 79	20% (8-40%)	80% (60-92%)	0.1 (0.022-0.3)	-0.66% (-2.1--0.0027%)	-0.74% (-2.3-0.0078%)	-65 (-200-0.71)	-930 (-2900--3.7)
Afghanistan	2016	Female	80 to 84	18% (6.7-40%)	82% (60-93%)	0.094 (0.019-0.28)	-0.58% (-1.9-0.045%)	-0.65% (-2.1-0.054%)	-39 (-130-3.2)	-410 (-1400-32)
Afghanistan	2016	Female	85 to 89	19% (6.6-40%)	81% (60-93%)	0.095 (0.018-0.28)	-0.59% (-1.9-0.0066%)	-0.68% (-2.2-0.0014%)	-18 (-59-0.038)	-130 (-450-1.5)
Afghanistan	2016	Female	90 to 94	22% (6.9-44%)	78% (56-93%)	0.11 (0.02-0.32)	-0.64% (-1.9--0.011%)	-0.76% (-2.2--0.024%)	-6 (-17--0.19)	-30 (-89--0.56)
Afghanistan	2016	Female	95 plus	27% (8.3-51%)	73% (49-92%)	0.13 (0.018-0.35)	-0.73% (-2.1-0.0011%)	-0.88% (-2.5--0.019%)	-1.6 (-4.6--0.034)	-5.2 (-15-0.0076)
Afghanistan	2016	Male	All Ages	9% (6.1-13%)	91% (87-94%)	0.076 (0.044-0.12)	0.35% (0.14-0.6%)	0.3% (-0.015-0.67%)	480 (-23-1100)	35000 (15000-59000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Afghanistan	2016	Male	15 to 19	7.4% (3.2-14%)	93% (86-97%)	0.038 (0.012-0.093)	0.36% (0.19-0.66%)	0.25% (0.058-0.63%)	16 (3.9-39)	2300 (1200-3900)
Afghanistan	2016	Male	20 to 24	12% (4.6-22%)	88% (78-95%)	0.086 (0.029-0.2)	0.71% (0.35-1.3%)	0.5% (0.12-1.2%)	46 (12-110)	5300 (2800-9400)
Afghanistan	2016	Male	25 to 29	14% (5.7-27%)	86% (73-94%)	0.14 (0.049-0.28)	0.96% (0.47-1.7%)	0.83% (0.24-1.7%)	65 (20-130)	5900 (3000-10000)
Afghanistan	2016	Male	30 to 34	16% (5.9-31%)	84% (69-94%)	0.17 (0.065-0.35)	1% (0.49-1.8%)	0.99% (0.31-2.1%)	73 (22-150)	5600 (2800-9900)
Afghanistan	2016	Male	35 to 39	18% (7.1-32%)	82% (68-93%)	0.2 (0.065-0.42)	1.1% (0.44-2%)	1.1% (0.27-2.3%)	80 (20-170)	5300 (2200-9900)
Afghanistan	2016	Male	40 to 44	18% (8.4-32%)	82% (68-92%)	0.21 (0.074-0.44)	0.89% (0.26-1.8%)	0.9% (0.084-2%)	68 (6.7-150)	4000 (1200-8000)
Afghanistan	2016	Male	45 to 49	18% (7.4-32%)	82% (68-93%)	0.22 (0.07-0.47)	0.73% (0.047-1.6%)	0.75% (-0.11-1.9%)	61 (-9.3-160)	3100 (200-7100)
Afghanistan	2016	Male	50 to 54	17% (7.4-32%)	83% (68-93%)	0.21 (0.064-0.46)	0.44% (-0.23-1.3%)	0.44% (-0.43-1.5%)	39 (-36-120)	1800 (-870-4900)
Afghanistan	2016	Male	55 to 59	16% (7.4-30%)	84% (70-93%)	0.2 (0.06-0.47)	0.28% (-0.39-1%)	0.28% (-0.54-1.1%)	26 (-51-110)	1000 (-1400-3700)
Afghanistan	2016	Male	60 to 64	16% (6.8-29%)	84% (71-93%)	0.18 (0.049-0.45)	0.14% (-0.49-0.8%)	0.13% (-0.62-0.9%)	13 (-64-92)	460 (-1600-2600)
Afghanistan	2016	Male	65 to 69	15% (6.7-27%)	85% (73-93%)	0.15 (0.046-0.34)	0.014% (-0.56-0.48%)	-0.0097% (-0.7-0.52%)	-0.99 (-70-52)	38 (-1500-1200)
Afghanistan	2016	Male	70 to 74	13% (6-24%)	87% (76-94%)	0.11 (0.031-0.29)	0.014% (-0.45-0.38%)	-0.00073% (-0.55-0.42%)	-0.068 (-50-38)	27 (-840-710)
Afghanistan	2016	Male	75 to 79	12% (4.9-22%)	88% (78-95%)	0.084 (0.021-0.23)	0.0081% (-0.43-0.26%)	-0.0058% (-0.5-0.29%)	-0.41 (-36-20)	9.4 (-480-290)
Afghanistan	2016	Male	80 to 84	11% (4.2-21%)	89% (79-96%)	0.071 (0.018-0.19)	0.035% (-0.34-0.31%)	0.019% (-0.42-0.34%)	0.79 (-17-14)	17 (-170-150)
Afghanistan	2016	Male	85 to 89	10% (4.2-21%)	90% (79-96%)	0.066 (0.017-0.17)	-0.0019% (-0.33-0.21%)	-0.031% (-0.42-0.21%)	-0.64 (-8.7-4.4)	-0.31 (-57-36)
Afghanistan	2016	Male	90 to 94	12% (4.8-23%)	88% (77-95%)	0.07 (0.016-0.19)	-0.046% (-0.39-0.16%)	-0.096% (-0.5-0.15%)	-0.7 (-3.7-1.1)	-2 (-17-7.1)
Afghanistan	2016	Male	95 plus	15% (5.2-30%)	85% (70-95%)	0.081 (0.015-0.23)	-0.071% (-0.52-0.17%)	-0.15% (-0.7-0.13%)	-0.26 (-1.2-0.22)	-0.48 (-3.5-1.1)
Bangladesh	1990	Female	All Ages	0.29% (0.2-0.4%)	100% (100-100%)	0.021 (0.018-0.024)	0.35% (0.29-0.41%)	0.5% (0.39-0.61%)	2500 (1900-3100)	120000 (98000-140000)
Bangladesh	1990	Female	15 to 19	0.25% (0.098-0.49%)	100% (100-100%)	0.012 (0.004-0.025)	0.31% (0.2-0.46%)	0.095% (0.051-0.16%)	11 (5.8-18)	4300 (2500-6600)
Bangladesh	1990	Female	20 to 24	0.26% (0.087-0.52%)	100% (99-100%)	0.024 (0.0091-0.045)	0.7% (0.48-0.96%)	0.3% (0.17-0.46%)	42 (25-67)	10000 (6800-15000)
Bangladesh	1990	Female	25 to 29	0.27% (0.087-0.56%)	100% (99-100%)	0.035 (0.013-0.066)	0.9% (0.66-1.2%)	0.57% (0.35-0.88%)	73 (44-110)	11000 (8000-16000)
Bangladesh	1990	Female	30 to 34	0.31% (0.087-0.64%)	100% (99-100%)	0.045 (0.016-0.082)	1% (0.77-1.4%)	0.89% (0.6-1.3%)	94 (63-130)	10000 (7200-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	1990	Female	35 to 39	0.35% (0.12-0.6%)	100% (99-100%)	0.051 (0.02-0.088)	1.4% (1-1.9%)	1.6% (1-2.3%)	140 (95-220)	10000 (7700-14000)
Bangladesh	1990	Female	40 to 44	0.37% (0.16-0.65%)	100% (99-100%)	0.051 (0.023-0.094)	1.2% (0.89-1.5%)	1.3% (0.92-1.9%)	160 (110-220)	10000 (7600-14000)
Bangladesh	1990	Female	45 to 49	0.38% (0.16-0.67%)	100% (99-100%)	0.048 (0.019-0.088)	1.5% (1.1-2.1%)	1.9% (1.3-2.8%)	280 (180-410)	14000 (10000-20000)
Bangladesh	1990	Female	50 to 54	0.39% (0.15-0.72%)	100% (99-100%)	0.049 (0.018-0.093)	1.6% (1.1-2.2%)	2% (1.3-2.9%)	350 (230-530)	15000 (10000-21000)
Bangladesh	1990	Female	55 to 59	0.41% (0.17-0.77%)	100% (99-100%)	0.055 (0.02-0.11)	1.2% (0.82-1.8%)	1.5% (0.91-2.2%)	300 (190-460)	11000 (7300-16000)
Bangladesh	1990	Female	60 to 64	0.43% (0.19-0.86%)	100% (99-100%)	0.057 (0.02-0.12)	0.88% (0.58-1.3%)	0.98% (0.61-1.5%)	250 (150-380)	7800 (5100-11000)
Bangladesh	1990	Female	65 to 69	0.46% (0.2-0.8%)	100% (99-100%)	0.052 (0.019-0.098)	0.83% (0.56-1.1%)	0.94% (0.6-1.3%)	270 (170-390)	6700 (4600-9200)
Bangladesh	1990	Female	70 to 74	0.47% (0.22-0.83%)	100% (99-100%)	0.045 (0.016-0.094)	0.71% (0.48-1%)	0.8% (0.52-1.2%)	230 (150-340)	4400 (3000-6500)
Bangladesh	1990	Female	75 to 79	0.49% (0.2-0.88%)	100% (99-100%)	0.04 (0.014-0.085)	0.61% (0.4-0.87%)	0.68% (0.43-1%)	150 (93-220)	2200 (1400-3300)
Bangladesh	1990	Female	80 to 84	0.5% (0.2-0.9%)	100% (99-100%)	0.037 (0.012-0.079)	0.47% (0.31-0.69%)	0.52% (0.32-0.77%)	62 (39-93)	700 (450-1000)
Bangladesh	1990	Female	85 to 89	0.55% (0.23-1.1%)	99% (99-100%)	0.037 (0.012-0.08)	0.39% (0.26-0.56%)	0.41% (0.26-0.61%)	28 (17-41)	230 (150-320)
Bangladesh	1990	Female	90 to 94	0.65% (0.28-1.2%)	99% (99-100%)	0.039 (0.015-0.088)	0.34% (0.22-0.5%)	0.34% (0.21-0.52%)	8.5 (5.2-13)	51 (33-74)
Bangladesh	1990	Female	95 plus	0.82% (0.3-1.7%)	99% (98-100%)	0.044 (0.014-0.11)	0.34% (0.22-0.5%)	0.3% (0.17-0.49%)	1.9 (1.1-3.1)	8.3 (5.4-12)
Bangladesh	1990	Male	All Ages	0.75% (0.58-0.99%)	99% (99-99%)	0.012 (0.0099-0.015)	0.53% (0.43-0.67%)	0.6% (0.47-0.86%)	3300 (2500-4800)	20000 (160000-250000)
Bangladesh	1990	Male	15 to 19	0.48% (0.22-0.86%)	100% (99-100%)	0.0064 (0.0024-0.013)	0.61% (0.4-0.9%)	0.16% (0.11-0.25%)	17 (11-26)	7400 (4600-11000)
Bangladesh	1990	Male	20 to 24	0.78% (0.31-1.4%)	99% (99-100%)	0.014 (0.0055-0.025)	1.5% (1-2%)	0.58% (0.34-0.94%)	82 (45-130)	20000 (13000-29000)
Bangladesh	1990	Male	25 to 29	1.1% (0.55-1.9%)	99% (98-99%)	0.02 (0.01-0.035)	1.9% (1.4-2.6%)	1.1% (0.59-1.7%)	150 (81-240)	24000 (17000-33000)
Bangladesh	1990	Male	30 to 34	1.4% (0.69-2.5%)	99% (98-99%)	0.027 (0.014-0.044)	2.4% (1.8-3.1%)	1.9% (1.2-2.9%)	230 (150-350)	24000 (17000-33000)
Bangladesh	1990	Male	35 to 39	1.7% (0.85-2.6%)	98% (97-99%)	0.032 (0.015-0.057)	2.5% (1.8-3.4%)	2.4% (1.5-3.7%)	270 (160-410)	21000 (15000-28000)
Bangladesh	1990	Male	40 to 44	1.7% (0.92-2.7%)	98% (97-99%)	0.032 (0.015-0.057)	2.5% (1.8-3.5%)	2.7% (1.7-4.1%)	380 (240-580)	23000 (17000-33000)
Bangladesh	1990	Male	45 to 49	1.5% (0.73-2.5%)	98% (98-99%)	0.03 (0.013-0.056)	2.1% (1.5-3.2%)	2.3% (1.4-3.7%)	410 (250-670)	22000 (15000-32000)
Bangladesh	1990	Male	50 to 54	1.4% (0.73-2.3%)	99% (98-99%)	0.029 (0.012-0.053)	1.7% (1.2-2.6%)	1.8% (1.1-3%)	380 (230-650)	17000 (12000-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	1990	Male	55 to 59	1.3% (0.66-2.1%)	99% (98-99%)	0.027 (0.012-0.051)	1.4% (0.92-2.1%)	1.5% (0.88-2.4%)	360 (210-600)	14000 (8800-22000)
Bangladesh	1990	Male	60 to 64	1.2% (0.62-2%)	99% (98-99%)	0.025 (0.01-0.048)	1.1% (0.69-1.6%)	1.1% (0.66-1.8%)	320 (190-520)	10000 (6600-16000)
Bangladesh	1990	Male	65 to 69	1.2% (0.58-1.9%)	99% (98-99%)	0.022 (0.009-0.041)	0.86% (0.57-1.3%)	0.92% (0.57-1.4%)	290 (170-440)	7400 (4900-11000)
Bangladesh	1990	Male	70 to 74	1.1% (0.55-1.7%)	99% (98-99%)	0.018 (0.0065-0.036)	0.65% (0.43-1%)	0.69% (0.43-1.1%)	220 (140-360)	4400 (2900-6800)
Bangladesh	1990	Male	75 to 79	0.95% (0.44-1.7%)	99% (98-100%)	0.014 (0.005-0.03)	0.46% (0.29-0.7%)	0.48% (0.29-0.78%)	120 (71-200)	1900 (1200-2900)
Bangladesh	1990	Male	80 to 84	0.85% (0.42-1.4%)	99% (99-100%)	0.011 (0.0043-0.023)	0.37% (0.23-0.55%)	0.4% (0.24-0.62%)	54 (33-85)	610 (390-910)
Bangladesh	1990	Male	85 to 89	0.83% (0.4-1.5%)	99% (99-100%)	0.01 (0.004-0.02)	0.28% (0.18-0.4%)	0.29% (0.18-0.44%)	24 (15-35)	190 (130-280)
Bangladesh	1990	Male	90 to 94	0.91% (0.45-1.5%)	99% (98-100%)	0.01 (0.004-0.022)	0.22% (0.14-0.32%)	0.21% (0.13-0.33%)	6.2 (3.8-10)	38 (26-57)
Bangladesh	1990	Male	95 plus	1.1% (0.49-2%)	99% (98-100%)	0.011 (0.0033-0.024)	0.21% (0.13-0.32%)	0.17% (0.088-0.3%)	1.2 (0.6-2)	5.7 (3.5-8.7)
Bangladesh	1995	Female	All Ages	0.29% (0.2-0.41%)	100% (100-100%)	0.028 (0.024-0.032)	0.42% (0.35-0.49%)	0.58% (0.48-0.69%)	2600 (2100-3200)	130000 (100000-150000)
Bangladesh	1995	Female	15 to 19	0.25% (0.1-0.48%)	100% (100-100%)	0.016 (0.0053-0.033)	0.37% (0.24-0.52%)	0.12% (0.064-0.22%)	14 (6.9-24)	5100 (3100-7600)
Bangladesh	1995	Female	20 to 24	0.26% (0.088-0.51%)	100% (99-100%)	0.031 (0.011-0.058)	0.79% (0.55-1.1%)	0.33% (0.19-0.52%)	41 (23-65)	11000 (7200-16000)
Bangladesh	1995	Female	25 to 29	0.27% (0.084-0.57%)	100% (99-100%)	0.044 (0.016-0.083)	0.98% (0.73-1.3%)	0.6% (0.37-0.92%)	74 (44-120)	13000 (9000-18000)
Bangladesh	1995	Female	30 to 34	0.3% (0.081-0.63%)	100% (99-100%)	0.057 (0.019-0.1)	1.1% (0.82-1.4%)	0.93% (0.63-1.3%)	110 (73-160)	13000 (9100-17000)
Bangladesh	1995	Female	35 to 39	0.34% (0.11-0.63%)	100% (99-100%)	0.065 (0.022-0.11)	1.4% (1-1.9%)	1.6% (1.1-2.4%)	170 (110-250)	13000 (9700-18000)
Bangladesh	1995	Female	40 to 44	0.36% (0.15-0.63%)	100% (99-100%)	0.065 (0.029-0.12)	1.2% (0.9-1.5%)	1.4% (0.97-1.9%)	130 (89-180)	8700 (6500-11000)
Bangladesh	1995	Female	45 to 49	0.37% (0.16-0.62%)	100% (99-100%)	0.06 (0.023-0.11)	1.5% (1.1-2%)	2% (1.3-2.8%)	250 (170-350)	13000 (9200-18000)
Bangladesh	1995	Female	50 to 54	0.38% (0.15-0.71%)	100% (99-100%)	0.062 (0.022-0.12)	1.5% (1.1-2.1%)	2% (1.3-2.9%)	340 (220-500)	15000 (10000-20000)
Bangladesh	1995	Female	55 to 59	0.4% (0.16-0.76%)	100% (99-100%)	0.07 (0.025-0.14)	1.3% (0.88-1.8%)	1.6% (1-2.3%)	330 (210-480)	12000 (8000-17000)
Bangladesh	1995	Female	60 to 64	0.42% (0.17-0.86%)	100% (99-100%)	0.073 (0.026-0.16)	0.91% (0.62-1.3%)	1% (0.66-1.6%)	260 (170-390)	8300 (5600-12000)
Bangladesh	1995	Female	65 to 69	0.45% (0.18-0.8%)	100% (99-100%)	0.066 (0.023-0.13)	0.85% (0.61-1.2%)	0.98% (0.67-1.4%)	290 (200-430)	7300 (5200-10000)
Bangladesh	1995	Female	70 to 74	0.46% (0.21-0.8%)	100% (99-100%)	0.056 (0.019-0.11)	0.75% (0.53-1.1%)	0.86% (0.59-1.2%)	270 (180-390)	5200 (3600-7400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	1995	Female	75 to 79	0.47% (0.18-0.88%)	100% (99-100%)	0.05 (0.016-0.11)	0.63% (0.44-0.89%)	0.71% (0.48-1%)	170 (120-260)	2600 (1800-3800)
Bangladesh	1995	Female	80 to 84	0.48% (0.2-0.88%)	100% (99-100%)	0.047 (0.016-0.099)	0.51% (0.35-0.71%)	0.56% (0.37-0.81%)	82 (53-120)	930 (630-1300)
Bangladesh	1995	Female	85 to 89	0.53% (0.22-1%)	99% (99-100%)	0.046 (0.016-0.1)	0.43% (0.29-0.61%)	0.46% (0.3-0.67%)	39 (25-58)	320 (220-460)
Bangladesh	1995	Female	90 to 94	0.63% (0.26-1.2%)	99% (99-100%)	0.05 (0.018-0.11)	0.39% (0.26-0.55%)	0.4% (0.25-0.6%)	12 (7.7-18)	73 (49-110)
Bangladesh	1995	Female	95 plus	0.8% (0.31-1.6%)	99% (98-100%)	0.055 (0.017-0.13)	0.39% (0.25-0.58%)	0.36% (0.21-0.6%)	3 (1.7-5)	13 (8.2-19)
Bangladesh	1995	Male	All Ages	0.77% (0.59-1%)	99% (99-99%)	0.017 (0.014-0.021)	0.68% (0.56-0.84%)	0.72% (0.58-1%)	3700 (2900-5300)	220000 (180000-280000)
Bangladesh	1995	Male	15 to 19	0.48% (0.22-0.83%)	100% (99-100%)	0.0084 (0.0031-0.017)	0.67% (0.44-0.95%)	0.18% (0.12-0.28%)	19 (13-28)	8300 (5300-13000)
Bangladesh	1995	Male	20 to 24	0.78% (0.31-1.4%)	99% (99-100%)	0.018 (0.0072-0.034)	1.6% (1.1-2.1%)	0.63% (0.37-0.97%)	82 (48-130)	21000 (14000-30000)
Bangladesh	1995	Male	25 to 29	1.1% (0.52-2%)	99% (98-99%)	0.027 (0.013-0.046)	2.1% (1.6-2.8%)	1.2% (0.69-1.9%)	160 (94-270)	28000 (19000-37000)
Bangladesh	1995	Male	30 to 34	1.4% (0.66-2.5%)	99% (97-99%)	0.036 (0.018-0.058)	2.6% (1.9-3.4%)	2% (1.4-3.1%)	290 (190-430)	32000 (23000-42000)
Bangladesh	1995	Male	35 to 39	1.6% (0.81-2.6%)	98% (97-99%)	0.042 (0.02-0.076)	2.6% (2-3.6%)	2.5% (1.6-3.9%)	340 (210-530)	27000 (20000-37000)
Bangladesh	1995	Male	40 to 44	1.7% (0.88-2.7%)	98% (97-99%)	0.043 (0.019-0.078)	2.6% (1.9-3.6%)	2.8% (1.9-4.3%)	350 (230-520)	22000 (16000-30000)
Bangladesh	1995	Male	45 to 49	1.5% (0.72-2.5%)	98% (98-99%)	0.041 (0.016-0.076)	2.2% (1.6-3.4%)	2.4% (1.5-3.9%)	400 (240-640)	21000 (15000-31000)
Bangladesh	1995	Male	50 to 54	1.4% (0.73-2.3%)	99% (98-99%)	0.038 (0.016-0.071)	1.8% (1.2-2.7%)	1.9% (1.2-3.3%)	410 (250-690)	19000 (13000-29000)
Bangladesh	1995	Male	55 to 59	1.3% (0.66-2.1%)	99% (98-99%)	0.036 (0.016-0.068)	1.5% (1-2.2%)	1.6% (0.98-2.6%)	400 (240-660)	16000 (10000-23000)
Bangladesh	1995	Male	60 to 64	1.2% (0.6-2.1%)	99% (98-99%)	0.033 (0.013-0.062)	1.2% (0.77-1.8%)	1.2% (0.73-2%)	370 (220-610)	12000 (7700-18000)
Bangladesh	1995	Male	65 to 69	1.2% (0.6-1.9%)	99% (98-99%)	0.029 (0.011-0.054)	0.93% (0.64-1.3%)	1% (0.65-1.5%)	330 (210-500)	8500 (5700-12000)
Bangladesh	1995	Male	70 to 74	1.1% (0.56-1.7%)	99% (98-99%)	0.023 (0.0088-0.048)	0.71% (0.47-1.1%)	0.76% (0.47-1.2%)	260 (160-410)	5200 (3500-7800)
Bangladesh	1995	Male	75 to 79	0.94% (0.44-1.7%)	99% (98-100%)	0.019 (0.0066-0.04)	0.5% (0.33-0.75%)	0.54% (0.33-0.87%)	150 (91-240)	2300 (1500-3500)
Bangladesh	1995	Male	80 to 84	0.84% (0.41-1.4%)	99% (99-100%)	0.015 (0.0055-0.03)	0.4% (0.27-0.59%)	0.44% (0.28-0.66%)	74 (47-110)	830 (550-1200)
Bangladesh	1995	Male	85 to 89	0.82% (0.38-1.4%)	99% (99-100%)	0.013 (0.0053-0.027)	0.31% (0.21-0.44%)	0.33% (0.21-0.49%)	33 (21-48)	270 (180-370)
Bangladesh	1995	Male	90 to 94	0.89% (0.45-1.5%)	99% (99-100%)	0.013 (0.0051-0.028)	0.24% (0.16-0.35%)	0.24% (0.15-0.37%)	9.5 (5.8-14)	57 (38-83)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	1995	Male	95 plus	1.1% (0.5-1.9%)	99% (98-99%)	0.014 (0.0044-0.031)	0.24% (0.16-0.35%)	0.2% (0.11-0.34%)	1.9 (1.1-3.2)	9 (5.8-14)
Bangladesh	2000	Female	All Ages	0.29% (0.21-0.42%)	100% (100-100%)	0.037 (0.031-0.042)	0.49% (0.42-0.57%)	0.64% (0.53-0.75%)	2500 (2100-3100)	130000 (110000-160000)
Bangladesh	2000	Female	15 to 19	0.25% (0.1-0.5%)	100% (100-100%)	0.02 (0.0066-0.041)	0.43% (0.28-0.64%)	0.15% (0.081-0.27%)	14 (6.9-24)	5600 (3500-8700)
Bangladesh	2000	Female	20 to 24	0.26% (0.092-0.51%)	100% (99-100%)	0.038 (0.015-0.071)	0.9% (0.61-1.3%)	0.36% (0.22-0.58%)	38 (21-62)	12000 (7900-18000)
Bangladesh	2000	Female	25 to 29	0.27% (0.09-0.59%)	100% (99-100%)	0.055 (0.022-0.11)	1.1% (0.8-1.4%)	0.64% (0.38-0.99%)	61 (37-96)	13000 (8900-18000)
Bangladesh	2000	Female	30 to 34	0.3% (0.084-0.63%)	100% (99-100%)	0.07 (0.024-0.13)	1.1% (0.87-1.5%)	0.97% (0.66-1.4%)	98 (66-140)	13000 (9700-18000)
Bangladesh	2000	Female	35 to 39	0.34% (0.1-0.61%)	100% (99-100%)	0.08 (0.03-0.13)	1.4% (1.1-1.8%)	1.7% (1.1-2.4%)	180 (120-270)	16000 (11000-20000)
Bangladesh	2000	Female	40 to 44	0.36% (0.15-0.64%)	100% (99-100%)	0.081 (0.035-0.15)	1.2% (0.89-1.5%)	1.4% (0.98-1.9%)	140 (97-190)	10000 (7800-14000)
Bangladesh	2000	Female	45 to 49	0.37% (0.16-0.65%)	100% (99-100%)	0.075 (0.029-0.14)	1.4% (1.1-1.9%)	1.9% (1.3-2.7%)	180 (130-260)	10000 (7400-13000)
Bangladesh	2000	Female	50 to 54	0.39% (0.15-0.73%)	100% (99-100%)	0.077 (0.027-0.16)	1.5% (1.1-2%)	2% (1.3-2.9%)	280 (190-400)	13000 (9000-17000)
Bangladesh	2000	Female	55 to 59	0.4% (0.16-0.75%)	100% (99-100%)	0.087 (0.031-0.17)	1.3% (0.91-1.8%)	1.7% (1.1-2.4%)	320 (200-480)	12000 (8300-17000)
Bangladesh	2000	Female	60 to 64	0.43% (0.19-0.82%)	100% (99-100%)	0.09 (0.032-0.18)	0.95% (0.66-1.3%)	1.1% (0.74-1.7%)	260 (170-400)	8700 (6000-12000)
Bangladesh	2000	Female	65 to 69	0.45% (0.19-0.78%)	100% (99-100%)	0.081 (0.029-0.16)	0.86% (0.64-1.2%)	1% (0.72-1.4%)	290 (190-410)	7300 (5300-10000)
Bangladesh	2000	Female	70 to 74	0.46% (0.2-0.8%)	100% (99-100%)	0.069 (0.025-0.14)	0.75% (0.54-1.1%)	0.89% (0.61-1.3%)	280 (180-410)	5500 (3800-7900)
Bangladesh	2000	Female	75 to 79	0.47% (0.2-0.86%)	100% (99-100%)	0.061 (0.02-0.13)	0.59% (0.42-0.84%)	0.68% (0.46-0.99%)	190 (130-270)	2900 (2000-4100)
Bangladesh	2000	Female	80 to 84	0.48% (0.19-0.9%)	100% (99-100%)	0.058 (0.019-0.12)	0.51% (0.35-0.73%)	0.58% (0.38-0.84%)	110 (73-160)	1300 (860-1800)
Bangladesh	2000	Female	85 to 89	0.53% (0.22-1%)	99% (99-100%)	0.058 (0.02-0.13)	0.46% (0.32-0.68%)	0.5% (0.32-0.76%)	57 (37-88)	470 (320-700)
Bangladesh	2000	Female	90 to 94	0.63% (0.26-1.1%)	99% (99-100%)	0.061 (0.023-0.14)	0.43% (0.29-0.63%)	0.45% (0.28-0.69%)	20 (12-30)	120 (78-170)
Bangladesh	2000	Female	95 plus	0.78% (0.28-1.5%)	99% (98-100%)	0.068 (0.02-0.16)	0.45% (0.29-0.69%)	0.42% (0.24-0.71%)	4.9 (2.8-8.2)	20 (13-32)
Bangladesh	2000	Male	All Ages	0.86% (0.66-1.1%)	99% (99-99%)	0.023 (0.018-0.028)	0.88% (0.73-1%)	0.9% (0.73-1.2%)	4300 (3500-5700)	260000 (210000-320000)
Bangladesh	2000	Male	15 to 19	0.53% (0.25-0.9%)	99% (99-100%)	0.011 (0.0042-0.021)	0.75% (0.49-1.1%)	0.21% (0.14-0.3%)	20 (14-29)	9400 (5700-14000)
Bangladesh	2000	Male	20 to 24	0.92% (0.39-1.5%)	99% (98-100%)	0.025 (0.011-0.046)	1.8% (1.3-2.4%)	0.72% (0.44-1.1%)	89 (53-140)	24000 (16000-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	2000	Male	25 to 29	1.3% (0.64-2%)	99% (98-99%)	0.037 (0.018-0.061)	2.4% (1.8-3.1%)	1.4% (0.83-2.2%)	180 (100-280)	30000 (21000-41000)
Bangladesh	2000	Male	30 to 34	1.5% (0.74-2.4%)	99% (98-99%)	0.044 (0.025-0.071)	2.8% (2.2-3.6%)	2.3% (1.6-3.4%)	320 (220-460)	36000 (26000-48000)
Bangladesh	2000	Male	35 to 39	1.7% (0.92-2.5%)	98% (97-99%)	0.051 (0.027-0.089)	2.9% (2.2-3.9%)	2.9% (1.9-4.3%)	440 (280-660)	36000 (26000-48000)
Bangladesh	2000	Male	40 to 44	1.7% (1-2.7%)	98% (97-99%)	0.054 (0.025-0.093)	2.9% (2.2-3.9%)	3.3% (2.2-4.7%)	470 (310-670)	30000 (22000-40000)
Bangladesh	2000	Male	45 to 49	1.5% (0.78-2.4%)	98% (98-99%)	0.048 (0.02-0.089)	2.5% (1.8-3.5%)	2.8% (1.8-4.4%)	380 (240-600)	21000 (15000-30000)
Bangladesh	2000	Male	50 to 54	1.3% (0.72-2.2%)	99% (98-99%)	0.044 (0.019-0.083)	2% (1.4-2.8%)	2.2% (1.5-3.4%)	420 (270-660)	20000 (14000-28000)
Bangladesh	2000	Male	55 to 59	1.4% (0.75-2.2%)	99% (98-99%)	0.048 (0.021-0.086)	1.6% (1.1-2.4%)	1.8% (1.1-2.9%)	450 (280-740)	18000 (12000-26000)
Bangladesh	2000	Male	60 to 64	1.4% (0.79-2.3%)	99% (98-99%)	0.047 (0.02-0.087)	1.3% (0.9-1.9%)	1.5% (0.9-2.3%)	450 (270-700)	14000 (9500-21000)
Bangladesh	2000	Male	65 to 69	1.3% (0.62-2%)	99% (98-99%)	0.037 (0.015-0.07)	1.1% (0.74-1.5%)	1.2% (0.76-1.7%)	400 (260-600)	10000 (7300-15000)
Bangladesh	2000	Male	70 to 74	1.1% (0.56-1.7%)	99% (98-99%)	0.028 (0.011-0.055)	0.76% (0.51-1.1%)	0.83% (0.52-1.3%)	310 (190-470)	6300 (4200-9300)
Bangladesh	2000	Male	75 to 79	0.93% (0.45-1.7%)	99% (98-100%)	0.021 (0.0078-0.046)	0.57% (0.38-0.82%)	0.62% (0.39-0.93%)	210 (130-310)	3200 (2100-4700)
Bangladesh	2000	Male	80 to 84	0.82% (0.4-1.3%)	99% (99-100%)	0.017 (0.0066-0.035)	0.47% (0.32-0.67%)	0.52% (0.34-0.77%)	120 (80-180)	1400 (930-2000)
Bangladesh	2000	Male	85 to 89	0.79% (0.36-1.4%)	99% (99-100%)	0.015 (0.0058-0.032)	0.39% (0.27-0.53%)	0.42% (0.27-0.6%)	57 (37-82)	460 (310-630)
Bangladesh	2000	Male	90 to 94	0.87% (0.43-1.4%)	99% (99-100%)	0.015 (0.006-0.032)	0.32% (0.22-0.46%)	0.33% (0.21-0.5%)	17 (11-26)	100 (70-150)
Bangladesh	2000	Male	95 plus	1% (0.49-1.9%)	99% (98-100%)	0.016 (0.0052-0.036)	0.33% (0.21-0.46%)	0.3% (0.17-0.47%)	4 (2.3-6.3)	17 (11-25)
Bangladesh	2005	Female	All Ages	0.3% (0.21-0.41%)	100% (100-100%)	0.044 (0.037-0.051)	0.58% (0.49-0.67%)	0.72% (0.6-0.86%)	2700 (2200-3300)	140000 (120000-170000)
Bangladesh	2005	Female	15 to 19	0.26% (0.1-0.47%)	100% (100-100%)	0.022 (0.0074-0.045)	0.51% (0.33-0.73%)	0.22% (0.13-0.35%)	16 (8.9-26)	6000 (3600-9200)
Bangladesh	2005	Female	20 to 24	0.26% (0.092-0.52%)	100% (99-100%)	0.042 (0.016-0.08)	1% (0.71-1.4%)	0.45% (0.27-0.7%)	40 (23-64)	13000 (8500-20000)
Bangladesh	2005	Female	25 to 29	0.27% (0.081-0.55%)	100% (99-100%)	0.061 (0.023-0.12)	1.2% (0.89-1.6%)	0.78% (0.48-1.2%)	69 (41-110)	15000 (10000-21000)
Bangladesh	2005	Female	30 to 34	0.31% (0.07-0.64%)	100% (99-100%)	0.078 (0.024-0.14)	1.2% (0.93-1.6%)	1.1% (0.75-1.6%)	98 (66-150)	14000 (10000-19000)
Bangladesh	2005	Female	35 to 39	0.35% (0.1-0.65%)	100% (99-100%)	0.09 (0.033-0.15)	1.4% (1.1-1.9%)	1.8% (1.2-2.6%)	200 (130-290)	17000 (13000-23000)
Bangladesh	2005	Female	40 to 44	0.38% (0.15-0.67%)	100% (99-100%)	0.091 (0.04-0.16)	1.1% (0.87-1.5%)	1.4% (0.96-2%)	160 (110-240)	13000 (9700-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	2005	Female	45 to 49	0.38% (0.16-0.69%)	100% (99-100%)	0.085 (0.033-0.16)	1.4% (1-1.9%)	2% (1.4-2.8%)	240 (160-330)	13000 (9600-18000)
Bangladesh	2005	Female	50 to 54	0.4% (0.16-0.73%)	100% (99-100%)	0.087 (0.03-0.17)	1.5% (1.1-2%)	2% (1.4-2.9%)	230 (150-330)	11000 (7800-15000)
Bangladesh	2005	Female	55 to 59	0.41% (0.17-0.78%)	100% (99-100%)	0.098 (0.035-0.2)	1.3% (0.91-1.7%)	1.7% (1.1-2.4%)	290 (190-420)	11000 (7900-16000)
Bangladesh	2005	Female	60 to 64	0.44% (0.18-0.85%)	100% (99-100%)	0.1 (0.035-0.22)	0.96% (0.67-1.3%)	1.2% (0.75-1.7%)	280 (180-400)	9300 (6400-13000)
Bangladesh	2005	Female	65 to 69	0.46% (0.2-0.81%)	100% (99-100%)	0.092 (0.033-0.18)	0.87% (0.63-1.2%)	1.1% (0.74-1.5%)	290 (200-420)	7600 (5500-11000)
Bangladesh	2005	Female	70 to 74	0.47% (0.21-0.84%)	100% (99-100%)	0.078 (0.027-0.16)	0.75% (0.53-1.1%)	0.92% (0.63-1.3%)	280 (190-420)	5700 (3900-8100)
Bangladesh	2005	Female	75 to 79	0.49% (0.19-0.88%)	100% (99-100%)	0.069 (0.022-0.15)	0.61% (0.42-0.89%)	0.72% (0.48-1.1%)	220 (150-340)	3500 (2400-5100)
Bangladesh	2005	Female	80 to 84	0.5% (0.22-0.94%)	100% (99-100%)	0.065 (0.022-0.14)	0.54% (0.37-0.8%)	0.63% (0.42-0.94%)	160 (110-250)	1900 (1300-2800)
Bangladesh	2005	Female	85 to 89	0.55% (0.22-1%)	99% (99-100%)	0.064 (0.023-0.14)	0.5% (0.35-0.73%)	0.56% (0.36-0.85%)	96 (63-150)	790 (530-1200)
Bangladesh	2005	Female	90 to 94	0.64% (0.27-1.2%)	99% (99-100%)	0.068 (0.025-0.15)	0.48% (0.32-0.7%)	0.52% (0.33-0.79%)	35 (22-54)	200 (140-300)
Bangladesh	2005	Female	95 plus	0.79% (0.31-1.6%)	99% (98-100%)	0.074 (0.024-0.18)	0.51% (0.32-0.8%)	0.5% (0.29-0.85%)	8.8 (5.1-15)	36 (22-57)
Bangladesh	2005	Male	All Ages	0.92% (0.72-1.2%)	99% (99-99%)	0.027 (0.022-0.034)	1% (0.86-1.2%)	0.99% (0.81-1.2%)	4800 (3900-6100)	290000 (240000-360000)
Bangladesh	2005	Male	15 to 19	0.55% (0.26-0.92%)	99% (99-100%)	0.012 (0.0047-0.024)	0.78% (0.53-1.1%)	0.21% (0.14-0.31%)	20 (13-29)	9900 (6200-15000)
Bangladesh	2005	Male	20 to 24	0.95% (0.42-1.6%)	99% (98-100%)	0.029 (0.012-0.053)	1.9% (1.3-2.6%)	0.75% (0.46-1.1%)	93 (57-140)	27000 (18000-39000)
Bangladesh	2005	Male	25 to 29	1.3% (0.68-2.1%)	99% (98-99%)	0.042 (0.021-0.072)	2.5% (1.9-3.3%)	1.5% (0.89-2.3%)	190 (110-290)	34000 (24000-46000)
Bangladesh	2005	Male	30 to 34	1.5% (0.78-2.7%)	98% (97-99%)	0.051 (0.029-0.08)	2.9% (2.2-3.7%)	2.3% (1.6-3.3%)	300 (210-440)	36000 (26000-48000)
Bangladesh	2005	Male	35 to 39	1.8% (0.96-2.7%)	98% (97-99%)	0.059 (0.032-0.1)	2.9% (2.2-3.9%)	2.9% (1.9-4.4%)	450 (290-670)	39000 (28000-52000)
Bangladesh	2005	Male	40 to 44	1.8% (1.1-2.8%)	98% (97-99%)	0.062 (0.03-0.1)	2.8% (2.2-3.8%)	3.1% (2.2-4.5%)	540 (370-770)	37000 (27000-49000)
Bangladesh	2005	Male	45 to 49	1.6% (0.81-2.5%)	98% (97-99%)	0.056 (0.023-0.1)	2.4% (1.8-3.4%)	2.8% (1.8-4.3%)	480 (310-730)	27000 (19000-38000)
Bangladesh	2005	Male	50 to 54	1.4% (0.76-2.3%)	99% (98-99%)	0.051 (0.022-0.095)	2% (1.4-2.9%)	2.3% (1.5-3.6%)	390 (250-620)	19000 (13000-27000)
Bangladesh	2005	Male	55 to 59	1.5% (0.79-2.3%)	99% (98-99%)	0.055 (0.025-0.1)	1.6% (1.1-2.4%)	1.8% (1.2-2.8%)	440 (290-690)	18000 (12000-25000)
Bangladesh	2005	Male	60 to 64	1.5% (0.84-2.5%)	98% (98-99%)	0.055 (0.023-0.1)	1.3% (0.92-1.9%)	1.5% (0.92-2.2%)	480 (300-740)	16000 (11000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	2005	Male	65 to 69	1.3% (0.68-2%)	99% (98-99%)	0.043 (0.017-0.078)	1.1% (0.78-1.5%)	1.2% (0.79-1.7%)	440 (300-650)	12000 (8400-16000)
Bangladesh	2005	Male	70 to 74	1.1% (0.6-1.8%)	99% (98-99%)	0.031 (0.012-0.063)	0.8% (0.56-1.1%)	0.87% (0.57-1.3%)	360 (230-530)	7500 (5100-11000)
Bangladesh	2005	Male	75 to 79	0.95% (0.47-1.7%)	99% (98-100%)	0.024 (0.0089-0.053)	0.63% (0.44-0.9%)	0.7% (0.45-1%)	280 (180-410)	4300 (2900-6000)
Bangladesh	2005	Male	80 to 84	0.84% (0.4-1.4%)	99% (99-100%)	0.02 (0.0072-0.04)	0.57% (0.37-0.81%)	0.64% (0.4-0.94%)	220 (130-310)	2400 (1500-3400)
Bangladesh	2005	Male	85 to 89	0.82% (0.38-1.5%)	99% (99-100%)	0.017 (0.0067-0.036)	0.49% (0.33-0.67%)	0.54% (0.34-0.75%)	110 (67-150)	850 (560-1100)
Bangladesh	2005	Male	90 to 94	0.9% (0.44-1.5%)	99% (99-100%)	0.017 (0.0066-0.035)	0.42% (0.27-0.61%)	0.45% (0.26-0.68%)	34 (19-51)	190 (120-270)
Bangladesh	2005	Male	95 plus	1.1% (0.5-2%)	99% (98-99%)	0.018 (0.006-0.041)	0.44% (0.27-0.62%)	0.43% (0.22-0.65%)	7.9 (4.1-12)	32 (19-46)
Bangladesh	2010	Female	All Ages	0.31% (0.23-0.43%)	100% (100-100%)	0.048 (0.041-0.056)	0.64% (0.54-0.74%)	0.72% (0.6-0.86%)	2600 (2200-3100)	140000 (120000-170000)
Bangladesh	2010	Female	15 to 19	0.26% (0.1-0.49%)	100% (100-100%)	0.022 (0.0075-0.047)	0.56% (0.36-0.84%)	0.26% (0.15-0.41%)	14 (8.1-24)	6000 (3600-9300)
Bangladesh	2010	Female	20 to 24	0.26% (0.09-0.53%)	100% (99-100%)	0.043 (0.016-0.082)	1.1% (0.79-1.6%)	0.47% (0.29-0.72%)	33 (19-51)	14000 (8700-21000)
Bangladesh	2010	Female	25 to 29	0.28% (0.089-0.58%)	100% (99-100%)	0.064 (0.025-0.13)	1.3% (0.94-1.7%)	0.78% (0.5-1.1%)	60 (38-88)	15000 (11000-22000)
Bangladesh	2010	Female	30 to 34	0.32% (0.09-0.65%)	100% (99-100%)	0.083 (0.03-0.14)	1.2% (0.92-1.6%)	1% (0.67-1.4%)	85 (56-120)	15000 (10000-21000)
Bangladesh	2010	Female	35 to 39	0.37% (0.12-0.65%)	100% (99-100%)	0.095 (0.038-0.16)	1.3% (1-1.7%)	1.5% (1-2.2%)	150 (100-210)	15000 (11000-20000)
Bangladesh	2010	Female	40 to 44	0.39% (0.17-0.7%)	100% (99-100%)	0.096 (0.044-0.17)	1.1% (0.82-1.4%)	1.2% (0.82-1.7%)	150 (98-210)	13000 (9700-17000)
Bangladesh	2010	Female	45 to 49	0.4% (0.18-0.7%)	100% (99-100%)	0.09 (0.034-0.17)	1.2% (0.94-1.6%)	1.7% (1.1-2.4%)	240 (160-330)	15000 (11000-19000)
Bangladesh	2010	Female	50 to 54	0.42% (0.17-0.77%)	100% (99-100%)	0.093 (0.034-0.19)	1.3% (0.98-1.8%)	1.8% (1.3-2.6%)	250 (180-370)	12000 (9000-17000)
Bangladesh	2010	Female	55 to 59	0.43% (0.19-0.79%)	100% (99-100%)	0.11 (0.038-0.21)	1.2% (0.84-1.5%)	1.5% (1-2.1%)	210 (140-300)	8800 (6400-12000)
Bangladesh	2010	Female	60 to 64	0.46% (0.2-0.9%)	100% (99-100%)	0.11 (0.039-0.23)	0.9% (0.64-1.2%)	1.1% (0.69-1.6%)	230 (140-330)	8000 (5600-11000)
Bangladesh	2010	Female	65 to 69	0.49% (0.2-0.85%)	100% (99-100%)	0.098 (0.034-0.19)	0.82% (0.6-1.1%)	1% (0.7-1.4%)	280 (190-390)	7400 (5300-10000)
Bangladesh	2010	Female	70 to 74	0.5% (0.22-0.89%)	100% (99-100%)	0.084 (0.029-0.18)	0.71% (0.49-0.99%)	0.88% (0.59-1.2%)	270 (180-380)	5500 (3800-7800)
Bangladesh	2010	Female	75 to 79	0.51% (0.21-0.95%)	99% (99-100%)	0.075 (0.025-0.17)	0.56% (0.39-0.8%)	0.66% (0.44-0.97%)	210 (140-320)	3400 (2400-4900)
Bangladesh	2010	Female	80 to 84	0.52% (0.21-0.94%)	99% (99-100%)	0.07 (0.024-0.15)	0.54% (0.37-0.79%)	0.62% (0.41-0.93%)	210 (130-310)	2400 (1600-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	2010	Female	85 to 89	0.57% (0.25-1%)	99% (99-100%)	0.069 (0.026-0.14)	0.53% (0.37-0.76%)	0.59% (0.4-0.86%)	150 (97-210)	1200 (820-1700)
Bangladesh	2010	Female	90 to 94	0.68% (0.3-1.3%)	99% (99-100%)	0.074 (0.029-0.16)	0.53% (0.36-0.79%)	0.58% (0.37-0.88%)	62 (40-96)	350 (240-530)
Bangladesh	2010	Female	95 plus	0.85% (0.32-1.7%)	99% (98-100%)	0.082 (0.026-0.2)	0.58% (0.36-0.9%)	0.58% (0.33-0.97%)	16 (9.3-27)	65 (41-100)
Bangladesh	2010	Male	All Ages	0.95% (0.73-1.2%)	99% (99-99%)	0.029 (0.023-0.037)	1.2% (1-1.4%)	1.1% (0.92-1.3%)	5400 (4400-6600)	310000 (250000-390000)
Bangladesh	2010	Male	15 to 19	0.51% (0.25-0.91%)	99% (99-100%)	0.012 (0.0045-0.025)	0.84% (0.54-1.2%)	0.21% (0.14-0.3%)	18 (12-26)	10000 (6000-15000)
Bangladesh	2010	Male	20 to 24	0.86% (0.37-1.5%)	99% (99-100%)	0.026 (0.011-0.051)	2% (1.4-2.8%)	0.72% (0.44-1.1%)	83 (51-130)	27000 (18000-39000)
Bangladesh	2010	Male	25 to 29	1.2% (0.61-2.1%)	99% (98-99%)	0.04 (0.019-0.069)	2.6% (1.9-3.4%)	1.4% (0.82-2.2%)	180 (100-280)	35000 (25000-48000)
Bangladesh	2010	Male	30 to 34	1.6% (0.81-2.9%)	98% (97-99%)	0.054 (0.028-0.089)	3% (2.3-3.9%)	2.3% (1.6-3.2%)	310 (210-440)	39000 (29000-53000)
Bangladesh	2010	Male	35 to 39	1.8% (0.94-2.9%)	98% (97-99%)	0.062 (0.031-0.11)	3% (2.3-4%)	3% (1.8-4.5%)	440 (270-660)	39000 (29000-53000)
Bangladesh	2010	Male	40 to 44	1.8% (1-3%)	98% (97-99%)	0.063 (0.03-0.11)	2.9% (2.2-3.8%)	3.1% (2.2-4.5%)	570 (390-840)	40000 (30000-53000)
Bangladesh	2010	Male	45 to 49	1.7% (0.83-2.6%)	98% (97-99%)	0.06 (0.025-0.11)	2.4% (1.8-3.4%)	2.7% (1.8-4.2%)	570 (370-890)	33000 (24000-47000)
Bangladesh	2010	Male	50 to 54	1.5% (0.79-2.5%)	98% (98-99%)	0.056 (0.025-0.1)	2.1% (1.5-2.9%)	2.4% (1.5-3.6%)	520 (320-790)	25000 (18000-34000)
Bangladesh	2010	Male	55 to 59	1.4% (0.74-2.3%)	99% (98-99%)	0.054 (0.024-0.099)	1.7% (1.2-2.3%)	1.9% (1.2-2.8%)	420 (260-610)	17000 (12000-23000)
Bangladesh	2010	Male	60 to 64	1.4% (0.69-2.3%)	99% (98-99%)	0.05 (0.021-0.094)	1.3% (0.89-1.9%)	1.4% (0.85-2.2%)	430 (260-670)	15000 (9900-21000)
Bangladesh	2010	Male	65 to 69	1.3% (0.65-2%)	99% (98-99%)	0.043 (0.018-0.08)	1% (0.74-1.5%)	1.1% (0.75-1.6%)	440 (290-650)	12000 (8500-16000)
Bangladesh	2010	Male	70 to 74	1.2% (0.66-1.8%)	99% (98-99%)	0.035 (0.014-0.072)	0.84% (0.58-1.2%)	0.92% (0.6-1.4%)	400 (260-600)	8400 (5800-12000)
Bangladesh	2010	Male	75 to 79	1% (0.55-1.8%)	99% (98-99%)	0.027 (0.011-0.059)	0.71% (0.47-1%)	0.79% (0.51-1.2%)	350 (220-510)	5300 (3500-7600)
Bangladesh	2010	Male	80 to 84	0.92% (0.45-1.5%)	99% (99-100%)	0.022 (0.0083-0.044)	0.7% (0.42-1%)	0.81% (0.47-1.2%)	340 (200-520)	3700 (2300-5500)
Bangladesh	2010	Male	85 to 89	0.89% (0.42-1.6%)	99% (98-100%)	0.019 (0.0075-0.039)	0.65% (0.35-0.89%)	0.74% (0.37-1%)	200 (100-290)	1600 (840-2200)
Bangladesh	2010	Male	90 to 94	0.99% (0.51-1.6%)	99% (98-99%)	0.019 (0.0076-0.041)	0.6% (0.29-0.86%)	0.67% (0.27-0.99%)	73 (29-110)	400 (190-570)
Bangladesh	2010	Male	95 plus	1.2% (0.59-2.1%)	99% (98-99%)	0.021 (0.0068-0.045)	0.64% (0.3-0.92%)	0.67% (0.25-1%)	17 (6.4-26)	65 (30-94)
Bangladesh	2016	Female	All Ages	0.32% (0.23-0.43%)	100% (100-100%)	0.054 (0.035-0.076)	0.77% (0.67-0.9%)	0.82% (0.68-0.98%)	2900 (2300-3600)	160000 (130000-190000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	2016	Female	15 to 19	0.27% (0.11-0.49%)	100% (100-100%)	0.024 (0.0074-0.054)	0.61% (0.4-0.88%)	0.29% (0.17-0.46%)	13 (7.2-21)	6100 (3700-9200)
Bangladesh	2016	Female	20 to 24	0.27% (0.095-0.53%)	100% (99-100%)	0.047 (0.016-0.097)	1.3% (0.83-1.8%)	0.51% (0.31-0.77%)	28 (17-44)	14000 (8600-21000)
Bangladesh	2016	Female	25 to 29	0.29% (0.087-0.59%)	100% (99-100%)	0.068 (0.023-0.14)	1.4% (1-1.9%)	0.8% (0.52-1.2%)	50 (32-75)	16000 (11000-22000)
Bangladesh	2016	Female	30 to 34	0.32% (0.087-0.67%)	100% (99-100%)	0.087 (0.028-0.17)	1.3% (0.96-1.8%)	0.99% (0.66-1.4%)	75 (49-110)	16000 (11000-22000)
Bangladesh	2016	Female	35 to 39	0.37% (0.12-0.67%)	100% (99-100%)	0.098 (0.034-0.18)	1.3% (1-1.7%)	1.5% (1-2.1%)	140 (92-210)	17000 (12000-23000)
Bangladesh	2016	Female	40 to 44	0.39% (0.16-0.68%)	100% (99-100%)	0.098 (0.04-0.18)	1.1% (0.84-1.5%)	1.2% (0.84-1.8%)	140 (91-210)	14000 (9600-19000)
Bangladesh	2016	Female	45 to 49	0.4% (0.18-0.69%)	100% (99-100%)	0.09 (0.035-0.17)	1.2% (0.9-1.6%)	1.6% (1.1-2.3%)	230 (150-350)	16000 (11000-21000)
Bangladesh	2016	Female	50 to 54	0.41% (0.19-0.76%)	100% (99-100%)	0.093 (0.034-0.2)	1.3% (0.96-1.7%)	1.8% (1.2-2.5%)	320 (210-480)	17000 (12000-23000)
Bangladesh	2016	Female	55 to 59	0.43% (0.19-0.8%)	100% (99-100%)	0.11 (0.038-0.23)	1.2% (0.84-1.6%)	1.5% (1-2.2%)	290 (180-430)	12000 (8700-17000)
Bangladesh	2016	Female	60 to 64	0.47% (0.2-0.91%)	100% (99-100%)	0.11 (0.038-0.25)	0.92% (0.66-1.2%)	1.1% (0.74-1.6%)	210 (130-300)	7400 (5200-10000)
Bangladesh	2016	Female	65 to 69	0.49% (0.21-0.86%)	100% (99-100%)	0.1 (0.035-0.22)	0.82% (0.6-1.1%)	1% (0.72-1.4%)	250 (170-360)	6800 (4900-9600)
Bangladesh	2016	Female	70 to 74	0.5% (0.22-0.89%)	99% (99-100%)	0.088 (0.027-0.2)	0.74% (0.51-1%)	0.95% (0.61-1.4%)	300 (190-460)	6100 (4200-8900)
Bangladesh	2016	Female	75 to 79	0.52% (0.21-0.96%)	99% (99-100%)	0.079 (0.022-0.19)	0.55% (0.38-0.78%)	0.67% (0.45-0.97%)	230 (150-340)	3600 (2500-5200)
Bangladesh	2016	Female	80 to 84	0.53% (0.22-0.94%)	99% (99-100%)	0.073 (0.024-0.17)	0.55% (0.38-0.82%)	0.64% (0.43-0.98%)	260 (170-390)	2900 (2000-4300)
Bangladesh	2016	Female	85 to 89	0.58% (0.25-1.1%)	99% (99-100%)	0.073 (0.024-0.16)	0.56% (0.38-0.82%)	0.64% (0.42-0.96%)	210 (130-310)	1700 (1100-2500)
Bangladesh	2016	Female	90 to 94	0.7% (0.3-1.3%)	99% (99-100%)	0.078 (0.027-0.19)	0.59% (0.41-0.86%)	0.66% (0.43-1%)	110 (71-160)	610 (420-900)
Bangladesh	2016	Female	95 plus	0.88% (0.33-1.7%)	99% (98-100%)	0.088 (0.026-0.23)	0.66% (0.41-1%)	0.69% (0.4-1.1%)	33 (19-53)	130 (79-200)
Bangladesh	2016	Male	All Ages	1% (0.78-1.3%)	99% (99-99%)	0.033 (0.022-0.048)	1.5% (1.3-1.8%)	1.3% (1.1-1.6%)	6600 (5200-8300)	37000 (29000-46000)
Bangladesh	2016	Male	15 to 19	0.51% (0.23-0.9%)	99% (99-100%)	0.013 (0.004-0.027)	0.91% (0.59-1.3%)	0.21% (0.14-0.3%)	17 (11-26)	10000 (6500-16000)
Bangladesh	2016	Male	20 to 24	0.87% (0.36-1.5%)	99% (99-100%)	0.028 (0.01-0.057)	2.1% (1.5-3%)	0.67% (0.42-1%)	76 (45-120)	29000 (19000-42000)
Bangladesh	2016	Male	25 to 29	1.3% (0.63-2.2%)	99% (98-99%)	0.043 (0.019-0.078)	2.8% (2.1-3.5%)	1.3% (0.78-2%)	160 (94-250)	38000 (26000-53000)
Bangladesh	2016	Male	30 to 34	1.6% (0.8-2.9%)	98% (97-99%)	0.058 (0.026-0.11)	3.2% (2.4-4.1%)	2.3% (1.5-3.4%)	320 (200-490)	45000 (32000-61000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bangladesh	2016	Male	35 to 39	1.9% (0.96-3%)	98% (97-99%)	0.067 (0.031-0.12)	3.2% (2.4-4.2%)	3% (1.9-4.6%)	490 (290-740)	47000 (34000-63000)
Bangladesh	2016	Male	40 to 44	1.9% (1-3%)	98% (97-99%)	0.067 (0.028-0.12)	3.1% (2.3-4.1%)	3.2% (2.1-4.7%)	590 (380-880)	44000 (32000-60000)
Bangladesh	2016	Male	45 to 49	1.7% (0.83-2.7%)	98% (97-99%)	0.064 (0.025-0.13)	2.6% (1.9-3.6%)	2.8% (1.8-4.1%)	660 (400-1000)	40000 (28000-55000)
Bangladesh	2016	Male	50 to 54	1.6% (0.82-2.6%)	98% (97-99%)	0.06 (0.024-0.12)	2.4% (1.7-3.2%)	2.7% (1.7-4%)	760 (480-1100)	37000 (26000-51000)
Bangladesh	2016	Male	55 to 59	1.5% (0.77-2.4%)	99% (98-99%)	0.057 (0.023-0.11)	1.9% (1.3-2.6%)	2.1% (1.3-3.1%)	640 (400-950)	26000 (18000-36000)
Bangladesh	2016	Male	60 to 64	1.4% (0.72-2.3%)	99% (98-99%)	0.052 (0.02-0.11)	1.3% (0.92-2%)	1.4% (0.86-2.2%)	430 (250-680)	15000 (9800-22000)
Bangladesh	2016	Male	65 to 69	1.3% (0.68-2.1%)	99% (98-99%)	0.046 (0.017-0.091)	1.1% (0.77-1.5%)	1.2% (0.78-1.7%)	440 (290-660)	12000 (8500-17000)
Bangladesh	2016	Male	70 to 74	1.2% (0.64-1.9%)	99% (98-99%)	0.037 (0.013-0.084)	0.93% (0.64-1.3%)	1% (0.67-1.6%)	490 (300-720)	10000 (6600-15000)
Bangladesh	2016	Male	75 to 79	1.1% (0.52-1.8%)	99% (98-99%)	0.029 (0.0099-0.069)	0.83% (0.53-1.2%)	0.95% (0.57-1.4%)	450 (270-680)	6900 (4400-10000)
Bangladesh	2016	Male	80 to 84	0.93% (0.45-1.5%)	99% (98-100%)	0.023 (0.0081-0.051)	0.85% (0.44-1.2%)	1% (0.48-1.5%)	530 (250-790)	5700 (3000-8300)
Bangladesh	2016	Male	85 to 89	0.9% (0.43-1.6%)	99% (98-100%)	0.02 (0.007-0.045)	0.83% (0.37-1.2%)	0.97% (0.37-1.4%)	360 (140-520)	2700 (1200-3800)
Bangladesh	2016	Male	90 to 94	0.99% (0.51-1.6%)	99% (98-99%)	0.02 (0.0075-0.047)	0.81% (0.29-1.2%)	0.94% (0.27-1.4%)	150 (44-230)	790 (290-1200)
Bangladesh	2016	Male	95 plus	1.2% (0.55-2.1%)	99% (98-99%)	0.021 (0.0065-0.052)	0.87% (0.31-1.3%)	0.96% (0.26-1.5%)	39 (10-61)	140 (49-210)
Bhutan	1990	Female	All Ages	1.8% (1.1-2.6%)	98% (97-99%)	0.23 (0.12-0.41)	0.56% (0.41-0.74%)	0.95% (0.59-1.4%)	25 (15-38)	1100 (770-1400)
Bhutan	1990	Female	15 to 19	1.4% (0.53-2.9%)	99% (97-99%)	0.1 (0.025-0.26)	0.66% (0.42-1%)	0.31% (0.093-0.74%)	0.16 (0.046-0.38)	42 (25-65)
Bhutan	1990	Female	20 to 24	1.6% (0.48-3.3%)	98% (97-100%)	0.22 (0.052-0.58)	1.3% (0.86-1.9%)	0.52% (0.2-1.1%)	0.29 (0.11-0.59)	79 (50-120)
Bhutan	1990	Female	25 to 29	1.7% (0.52-3.8%)	98% (96-99%)	0.34 (0.09-0.85)	1.5% (1-2.1%)	0.77% (0.33-1.5%)	0.39 (0.16-0.81)	76 (49-110)
Bhutan	1990	Female	30 to 34	2.1% (0.58-4.7%)	98% (95-99%)	0.47 (0.13-1.1)	1.6% (1.1-2.4%)	1.3% (0.6-2.3%)	0.62 (0.29-1.2)	75 (49-110)
Bhutan	1990	Female	35 to 39	2.6% (0.65-5.5%)	97% (95-99%)	0.56 (0.15-1.3)	1.9% (1.3-3%)	2% (1-3.5%)	1 (0.5-1.8)	85 (54-130)
Bhutan	1990	Female	40 to 44	2.9% (1-6.1%)	97% (94-99%)	0.58 (0.21-1.3)	2.2% (1.5-3.4%)	2.6% (1.5-4.5%)	1.6 (0.87-2.8)	100 (64-160)
Bhutan	1990	Female	45 to 49	3.2% (1.2-6.9%)	97% (93-99%)	0.57 (0.19-1.3)	2.4% (1.5-3.8%)	3% (1.6-5.1%)	2.1 (1.1-3.7)	110 (67-170)
Bhutan	1990	Female	50 to 54	3.5% (1.3-7.9%)	96% (92-99%)	0.62 (0.17-1.5)	2.5% (1.5-4%)	3.1% (1.7-5.3%)	2.6 (1.3-4.5)	110 (64-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	1990	Female	55 to 59	3.8% (1.3-8.1%)	96% (92-99%)	0.75 (0.21-1.8)	2.4% (1.4-4.3%)	3% (1.5-5.6%)	2.9 (1.4-5.5)	110 (57-190)
Bhutan	1990	Female	60 to 64	4.1% (1.4-8.7%)	96% (91-99%)	0.8 (0.2-2.1)	2.2% (1.2-4%)	2.7% (1.3-5.1%)	3.1 (1.4-5.8)	91 (47-170)
Bhutan	1990	Female	65 to 69	4.5% (1.6-9.2%)	96% (91-98%)	0.73 (0.17-1.8)	2.2% (1-4%)	2.6% (1.2-4.9%)	3.2 (1.4-6.4)	77 (36-150)
Bhutan	1990	Female	70 to 74	4.6% (1.7-9.3%)	95% (91-98%)	0.62 (0.15-1.7)	1.9% (0.86-3.7%)	2.3% (0.97-4.4%)	2.8 (1.2-5.5)	53 (23-100)
Bhutan	1990	Female	75 to 79	4.7% (1.7-10%)	95% (90-98%)	0.54 (0.12-1.6)	1.6% (0.6-3.2%)	1.8% (0.66-3.8%)	1.9 (0.68-3.9)	28 (10-57)
Bhutan	1990	Female	80 to 84	5.1% (1.7-12%)	95% (88-98%)	0.53 (0.12-1.6)	1.5% (0.49-3.5%)	1.7% (0.53-4%)	1.3 (0.4-3)	14 (4.5-31)
Bhutan	1990	Female	85 to 89	6% (1.8-14%)	94% (86-98%)	0.56 (0.12-1.6)	1.4% (0.41-3.5%)	1.6% (0.42-4.1%)	0.64 (0.17-1.7)	4.9 (1.4-12)
Bhutan	1990	Female	90 to 94	7.6% (2.4-17%)	92% (83-98%)	0.63 (0.13-1.6)	1.4% (0.34-3.4%)	1.6% (0.34-4%)	0.22 (0.046-0.54)	1.2 (0.28-2.8)
Bhutan	1990	Female	95 plus	10% (3-23%)	90% (77-97%)	0.74 (0.15-2)	1.5% (0.23-3.7%)	1.6% (0.16-4.3%)	0.054 (0.0056-0.15)	0.2 (0.031-0.51)
Bhutan	1990	Male	All Ages	19% (11-28%)	81% (72-89%)	0.98 (0.65-1.4)	2.3% (1.6-3%)	4.1% (2.6-5.8%)	110 (66-160)	4400 (3100-5800)
Bhutan	1990	Male	15 to 19	5.9% (1.9-14%)	94% (86-98%)	0.23 (0.046-0.68)	1.4% (0.81-2.6%)	1.2% (0.41-3%)	0.49 (0.15-1.2)	70 (39-120)
Bhutan	1990	Male	20 to 24	12% (3.4-28%)	88% (72-97%)	0.61 (0.1-1.7)	4% (2.3-7%)	3.8% (1.3-8.2%)	1.8 (0.6-3.8)	190 (100-340)
Bhutan	1990	Male	25 to 29	21% (4.7-43%)	79% (57-95%)	1.1 (0.2-2.8)	6.4% (3.2-11%)	7% (2.3-14%)	3.2 (1.1-6.4)	290 (140-490)
Bhutan	1990	Male	30 to 34	30% (6.4-55%)	70% (45-94%)	1.7 (0.41-3.8)	8.8% (4.6-14%)	11% (4.5-18%)	5.4 (2.2-9.3)	390 (200-620)
Bhutan	1990	Male	35 to 39	38% (9.6-62%)	62% (38-90%)	2.3 (0.64-4.7)	10% (5.3-15%)	13% (6-20%)	7.3 (3.2-12)	460 (240-700)
Bhutan	1990	Male	40 to 44	42% (13-65%)	58% (35-87%)	2.6 (0.85-5)	11% (6.2-16%)	14% (7.4-21%)	10 (5.4-16)	550 (310-830)
Bhutan	1990	Male	45 to 49	43% (15-65%)	57% (35-85%)	2.8 (0.91-5.8)	11% (6-16%)	14% (7.5-21%)	12 (5.7-19)	550 (290-840)
Bhutan	1990	Male	50 to 54	43% (19-65%)	57% (35-81%)	3 (1-6.4)	11% (5.5-16%)	14% (6.7-21%)	13 (5.8-20)	510 (250-780)
Bhutan	1990	Male	55 to 59	42% (23-62%)	58% (38-77%)	3.1 (0.98-6.9)	10% (4.7-15%)	13% (5.9-20%)	13 (5.6-21)	440 (200-700)
Bhutan	1990	Male	60 to 64	43% (22-63%)	57% (37-78%)	3.1 (0.9-7.2)	8.8% (3.4-14%)	11% (3.9-18%)	12 (4.5-21)	360 (140-600)
Bhutan	1990	Male	65 to 69	44% (23-63%)	56% (37-77%)	2.8 (0.85-6.3)	7.6% (2.7-13%)	9.3% (3.1-16%)	11 (3.5-19)	260 (89-450)
Bhutan	1990	Male	70 to 74	42% (23-61%)	58% (39-77%)	2.4 (0.66-6.2)	5.9% (1.3-12%)	7% (1.4-14%)	8.1 (1.6-17)	150 (35-310)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	1990	Male	75 to 79	39% (19-60%)	61% (40-81%)	2.1 (0.44-6)	4.4% (0.68-10%)	5.1% (0.66-12%)	5 (0.62-11)	74 (11-170)
Bhutan	1990	Male	80 to 84	37% (17-58%)	63% (42-83%)	2 (0.44-5.5)	4% (0.5-9.2%)	4.6% (0.44-11%)	3.1 (0.29-7.2)	34 (4.1-77)
Bhutan	1990	Male	85 to 89	37% (15-57%)	63% (43-85%)	2.1 (0.44-5.9)	3.6% (0.31-8%)	4% (0.21-9.4%)	1.5 (0.083-3.5)	12 (1-27)
Bhutan	1990	Male	90 to 94	39% (17-58%)	61% (42-83%)	2.4 (0.39-7.1)	3.2% (0.11-7.3%)	3.5% (-0.069-8.4%)	0.5 (-0.0099-1.2)	2.7 (0.087-6.3)
Bhutan	1990	Male	95 plus	42% (19-64%)	58% (36-81%)	2.6 (0.35-8.5)	2.9% (0.045-7%)	3.1% (-0.3-7.8%)	0.1 (-0.01-0.26)	0.39 (0.006-0.93)
Bhutan	1995	Female	All Ages	1.7% (1.2-2.6%)	98% (97-99%)	0.28 (0.14-0.49)	0.64% (0.48-0.84%)	1.1% (0.71-1.7%)	24 (15-36)	980 (720-1300)
Bhutan	1995	Female	15 to 19	1.4% (0.53-2.8%)	99% (97-99%)	0.13 (0.035-0.33)	0.73% (0.45-1.1%)	0.39% (0.13-0.9%)	0.18 (0.059-0.41)	43 (25-66)
Bhutan	1995	Female	20 to 24	1.6% (0.47-3.5%)	98% (97-100%)	0.28 (0.069-0.73)	1.4% (0.95-2.1%)	0.6% (0.23-1.2%)	0.28 (0.11-0.59)	78 (49-110)
Bhutan	1995	Female	25 to 29	1.8% (0.5-4%)	98% (96-99%)	0.44 (0.12-1.1)	1.6% (1.1-2.3%)	0.87% (0.38-1.7%)	0.38 (0.16-0.77)	76 (50-110)
Bhutan	1995	Female	30 to 34	2.1% (0.61-4.8%)	98% (95-99%)	0.6 (0.16-1.4)	1.8% (1.2-2.5%)	1.4% (0.66-2.5%)	0.55 (0.25-1)	69 (45-100)
Bhutan	1995	Female	35 to 39	2.7% (0.7-5.5%)	97% (94-99%)	0.73 (0.2-1.6)	2.1% (1.4-3.1%)	2.2% (1.1-3.8%)	1 (0.49-1.7)	83 (53-120)
Bhutan	1995	Female	40 to 44	3% (1.1-6.5%)	97% (93-99%)	0.78 (0.28-1.7)	2.3% (1.5-3.6%)	2.9% (1.6-4.9%)	1.2 (0.69-2.2)	80 (51-130)
Bhutan	1995	Female	45 to 49	3.3% (1.1-7.1%)	97% (93-99%)	0.75 (0.24-1.7)	2.5% (1.5-3.9%)	3.2% (1.7-5.4%)	1.6 (0.85-2.7)	83 (50-130)
Bhutan	1995	Female	50 to 54	3.5% (1.3-7.7%)	96% (92-99%)	0.8 (0.23-1.9)	2.6% (1.6-4.1%)	3.3% (1.8-5.6%)	2 (1.1-3.5)	89 (54-140)
Bhutan	1995	Female	55 to 59	3.7% (1.4-8.3%)	96% (92-99%)	0.95 (0.29-2.3)	2.5% (1.4-4.4%)	3.1% (1.7-5.7%)	2.5 (1.3-4.6)	90 (51-160)
Bhutan	1995	Female	60 to 64	4.1% (1.5-8.5%)	96% (92-99%)	1 (0.27-2.7)	2.4% (1.3-4.2%)	2.9% (1.4-5.3%)	3 (1.5-5.7)	90 (47-160)
Bhutan	1995	Female	65 to 69	4.4% (1.5-9.1%)	96% (91-98%)	0.93 (0.24-2.3)	2.3% (1.1-4.3%)	2.8% (1.3-5.3%)	3.3 (1.6-6.3)	80 (39-150)
Bhutan	1995	Female	70 to 74	4.5% (1.7-9.1%)	95% (91-98%)	0.78 (0.19-2)	2.1% (0.98-3.9%)	2.5% (1.1-4.7%)	3 (1.4-5.6)	57 (28-110)
Bhutan	1995	Female	75 to 79	4.6% (1.6-10%)	95% (90-98%)	0.69 (0.16-2)	1.8% (0.71-3.6%)	2.1% (0.8-4.2%)	2.1 (0.87-4.3)	31 (13-62)
Bhutan	1995	Female	80 to 84	5.1% (1.6-12%)	95% (88-98%)	0.69 (0.15-2)	1.7% (0.63-3.9%)	2% (0.7-4.6%)	1.6 (0.56-3.8)	17 (6.2-41)
Bhutan	1995	Female	85 to 89	6% (1.8-14%)	94% (86-98%)	0.74 (0.17-2.1)	1.7% (0.62-4%)	2% (0.66-4.8%)	0.98 (0.32-2.3)	7.5 (2.6-17)
Bhutan	1995	Female	90 to 94	7.6% (2.4-18%)	92% (82-98%)	0.84 (0.19-2.3)	1.8% (0.53-4.1%)	2.1% (0.54-4.7%)	0.36 (0.096-0.82)	1.9 (0.56-4.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	1995	Female	95 plus	10% (3-24%)	90% (76-97%)	0.98 (0.19-2.8)	1.9% (0.44-4.7%)	2.2% (0.42-5.6%)	0.092 (0.017-0.23)	0.33 (0.075-0.79)
Bhutan	1995	Male	All Ages	18% (11-26%)	82% (74-89%)	1.2 (0.8-1.6)	2.7% (2-3.5%)	4.9% (3.1-6.8%)	110 (69-150)	4200 (3100-5400)
Bhutan	1995	Male	15 to 19	5.7% (1.9-14%)	94% (86-98%)	0.29 (0.055-0.81)	1.6% (0.9-2.8%)	1.4% (0.48-3.4%)	0.51 (0.17-1.2)	72 (40-130)
Bhutan	1995	Male	20 to 24	12% (3.4-28%)	88% (72-97%)	0.74 (0.14-2.1)	4.3% (2.5-7.2%)	4.2% (1.6-8.5%)	1.7 (0.61-3.5)	190 (110-320)
Bhutan	1995	Male	25 to 29	20% (4.9-40%)	80% (60-95%)	1.4 (0.25-3.3)	6.8% (3.6-11%)	7.6% (2.7-14%)	3.1 (1.1-5.8)	280 (150-450)
Bhutan	1995	Male	30 to 34	29% (5.5-54%)	71% (46-95%)	2.2 (0.49-4.6)	9.3% (5-14%)	12% (5.2-19%)	4.7 (1.9-7.7)	340 (180-530)
Bhutan	1995	Male	35 to 39	37% (9.6-62%)	63% (38-90%)	2.8 (0.81-5.8)	11% (5.9-16%)	14% (6.9-21%)	6.7 (3.1-10)	420 (230-620)
Bhutan	1995	Male	40 to 44	41% (13-65%)	59% (35-87%)	3.3 (1.1-6.5)	12% (7-16%)	16% (8.7-22%)	7.7 (4.2-11)	420 (250-600)
Bhutan	1995	Male	45 to 49	43% (15-65%)	57% (35-85%)	3.6 (1.1-7.3)	12% (6.8-17%)	16% (8.6-23%)	9.9 (5.2-15)	460 (260-660)
Bhutan	1995	Male	50 to 54	43% (20-65%)	57% (35-80%)	3.8 (1.3-7.9)	12% (6.6-16%)	16% (8.5-22%)	12 (6.1-17)	470 (260-670)
Bhutan	1995	Male	55 to 59	42% (22-63%)	58% (37-78%)	3.9 (1.2-8.5)	11% (5.5-16%)	15% (6.8-21%)	13 (5.7-19)	440 (210-640)
Bhutan	1995	Male	60 to 64	43% (22-63%)	57% (37-78%)	3.9 (1.1-9)	9.8% (4.2-14%)	12% (5.2-18%)	14 (5.8-21)	390 (180-610)
Bhutan	1995	Male	65 to 69	44% (23-63%)	56% (37-77%)	3.6 (1.1-7.9)	8.8% (3.6-14%)	11% (4.3-17%)	13 (4.9-21)	300 (120-490)
Bhutan	1995	Male	70 to 74	42% (23-61%)	58% (39-77%)	3.1 (0.84-7.7)	6.9% (1.9-12%)	8.4% (2.1-15%)	9.9 (2.4-18)	190 (49-340)
Bhutan	1995	Male	75 to 79	39% (19-60%)	61% (40-81%)	2.7 (0.55-7.4)	5.3% (0.96-11%)	6.3% (1-12%)	6.4 (0.98-13)	95 (17-180)
Bhutan	1995	Male	80 to 84	37% (17-58%)	63% (42-83%)	2.6 (0.58-7)	5% (0.74-10%)	5.7% (0.73-12%)	4.5 (0.56-9.2)	49 (7.3-99)
Bhutan	1995	Male	85 to 89	37% (16-56%)	63% (44-84%)	2.7 (0.57-7.8)	4.4% (0.59-8.7%)	5% (0.57-10%)	2.4 (0.27-5)	19 (2.5-38)
Bhutan	1995	Male	90 to 94	39% (18-58%)	61% (42-82%)	3 (0.53-9.1)	3.9% (0.38-8.1%)	4.4% (0.24-9.3%)	0.8 (0.045-1.7)	4.4 (0.42-9)
Bhutan	1995	Male	95 plus	43% (20-64%)	57% (36-80%)	3.3 (0.48-11)	3.6% (0.2-7.6%)	3.8% (-0.026-8.6%)	0.17 (-0.0011-0.37)	0.62 (0.035-1.3)
Bhutan	2000	Female	All Ages	1.7% (1.1-2.6%)	98% (97-99%)	0.26 (0.13-0.45)	0.68% (0.52-0.87%)	1.1% (0.71-1.7%)	22 (14-33)	960 (720-1200)
Bhutan	2000	Female	15 to 19	1.3% (0.55-2.8%)	99% (97-99%)	0.12 (0.032-0.31)	0.73% (0.46-1.1%)	0.32% (0.11-0.72%)	0.15 (0.049-0.34)	47 (28-70)
Bhutan	2000	Female	20 to 24	1.5% (0.5-3.3%)	98% (97-100%)	0.25 (0.066-0.68)	1.5% (1-2.2%)	0.53% (0.22-1.1%)	0.26 (0.1-0.55)	93 (58-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	2000	Female	25 to 29	1.7% (0.5-3.8%)	98% (96-100%)	0.4 (0.11-1)	1.7% (1.2-2.3%)	0.76% (0.33-1.5%)	0.31 (0.13-0.62)	81 (54-110)
Bhutan	2000	Female	30 to 34	2.1% (0.56-4.6%)	98% (95-99%)	0.54 (0.14-1.3)	1.7% (1.2-2.4%)	1.2% (0.6-2.2%)	0.51 (0.23-0.95)	77 (51-110)
Bhutan	2000	Female	35 to 39	2.6% (0.69-5.3%)	97% (95-99%)	0.64 (0.17-1.5)	2% (1.4-2.8%)	2% (1-3.4%)	0.79 (0.4-1.3)	75 (50-110)
Bhutan	2000	Female	40 to 44	2.9% (1-6.4%)	97% (94-99%)	0.69 (0.24-1.5)	2.1% (1.4-3.1%)	2.6% (1.5-4.2%)	1.2 (0.66-2)	84 (54-130)
Bhutan	2000	Female	45 to 49	3.2% (1.1-7%)	97% (93-99%)	0.68 (0.22-1.6)	2.3% (1.5-3.5%)	2.9% (1.6-4.9%)	1.5 (0.82-2.5)	82 (53-130)
Bhutan	2000	Female	50 to 54	3.5% (1.2-7.6%)	97% (92-99%)	0.72 (0.21-1.7)	2.3% (1.5-3.6%)	3% (1.6-5%)	1.6 (0.86-2.8)	76 (46-120)
Bhutan	2000	Female	55 to 59	3.7% (1.3-8%)	96% (92-99%)	0.85 (0.25-2.2)	2.3% (1.3-3.9%)	2.9% (1.5-5.3%)	2 (1-3.8)	77 (44-130)
Bhutan	2000	Female	60 to 64	4% (1.4-8.6%)	96% (91-99%)	0.91 (0.24-2.5)	2.1% (1.1-3.8%)	2.7% (1.3-5%)	2.4 (1.2-4.5)	75 (39-130)
Bhutan	2000	Female	65 to 69	4.4% (1.5-9.1%)	96% (91-98%)	0.84 (0.22-2.1)	2.1% (1-3.9%)	2.6% (1.2-5%)	3 (1.4-5.7)	74 (37-140)
Bhutan	2000	Female	70 to 74	4.5% (1.7-9%)	95% (91-98%)	0.71 (0.19-1.9)	1.9% (0.93-3.6%)	2.4% (1.1-4.5%)	2.9 (1.3-5.4)	55 (27-100)
Bhutan	2000	Female	75 to 79	4.6% (1.7-9.9%)	95% (90-98%)	0.64 (0.15-1.8)	1.7% (0.68-3.3%)	2% (0.78-4%)	2.1 (0.84-4.4)	32 (13-64)
Bhutan	2000	Female	80 to 84	5% (1.6-12%)	95% (88-98%)	0.63 (0.14-1.9)	1.6% (0.57-3.7%)	1.9% (0.66-4.4%)	1.7 (0.56-4)	18 (6.4-43)
Bhutan	2000	Female	85 to 89	5.9% (1.8-13%)	94% (87-98%)	0.66 (0.15-1.8)	1.6% (0.53-3.9%)	1.9% (0.57-4.8%)	1.1 (0.34-2.8)	8.5 (2.8-21)
Bhutan	2000	Female	90 to 94	7.3% (2.3-17%)	93% (83-98%)	0.74 (0.16-2.1)	1.7% (0.47-3.9%)	1.9% (0.5-4.7%)	0.47 (0.12-1.1)	2.5 (0.71-5.8)
Bhutan	2000	Female	95 plus	9.7% (2.5-23%)	90% (77-97%)	0.86 (0.16-2.4)	1.8% (0.37-4.4%)	2% (0.31-5.1%)	0.12 (0.019-0.3)	0.43 (0.086-1)
Bhutan	2000	Male	All Ages	19% (11-27%)	81% (73-89%)	1.1 (0.74-1.6)	2.7% (2-3.4%)	4.6% (2.8-6.5%)	100 (61-150)	4000 (2900-5200)
Bhutan	2000	Male	15 to 19	5.6% (1.7-14%)	94% (86-98%)	0.25 (0.045-0.78)	1.3% (0.74-2.2%)	0.96% (0.35-2.2%)	0.51 (0.19-1.2)	79 (45-140)
Bhutan	2000	Male	20 to 24	12% (3.2-26%)	88% (74-97%)	0.66 (0.13-1.9)	3.5% (2.1-5.7%)	2.9% (1.1-6.2%)	1.6 (0.62-3.3)	200 (120-340)
Bhutan	2000	Male	25 to 29	20% (4.5-42%)	80% (58-96%)	1.3 (0.23-3.1)	5.6% (3.1-8.9%)	5.5% (2-11%)	2.8 (1-5.2)	270 (150-440)
Bhutan	2000	Male	30 to 34	29% (5.6-55%)	71% (45-94%)	2 (0.46-4.2)	7.7% (4.5-11%)	8.9% (4.1-14%)	4.6 (2.1-7.6)	360 (200-540)
Bhutan	2000	Male	35 to 39	37% (9.3-62%)	63% (38-91%)	2.6 (0.71-5.5)	9.1% (5.1-13%)	11% (5.4-17%)	5.8 (2.7-9.2)	380 (210-550)
Bhutan	2000	Male	40 to 44	41% (13-64%)	59% (36-87%)	3 (0.96-6.1)	9.8% (5.8-14%)	13% (6.7-19%)	7.4 (3.9-11)	420 (250-600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	2000	Male	45 to 49	42% (14-65%)	58% (35-86%)	3.3 (0.98-6.8)	10% (5.5-15%)	13% (6.8-19%)	8.6 (4.3-13)	410 (220-590)
Bhutan	2000	Male	50 to 54	43% (19-64%)	57% (36-81%)	3.5 (1.1-7.1)	10% (5.2-14%)	13% (6.5-19%)	10 (5-15)	420 (220-610)
Bhutan	2000	Male	55 to 59	42% (22-63%)	58% (37-78%)	3.6 (1.1-8)	9.6% (4.6-14%)	13% (5.7-18%)	11 (5.1-17)	400 (190-600)
Bhutan	2000	Male	60 to 64	43% (22-63%)	57% (37-78%)	3.6 (1-8.2)	8.5% (3.4-13%)	11% (4.1-17%)	12 (4.5-19)	360 (140-560)
Bhutan	2000	Male	65 to 69	44% (24-63%)	56% (37-76%)	3.3 (0.98-7.2)	7.7% (2.7-12%)	9.6% (3.2-16%)	12 (4.1-21)	300 (100-500)
Bhutan	2000	Male	70 to 74	42% (24-61%)	58% (39-76%)	2.8 (0.79-7.2)	6% (1.5-11%)	7.3% (1.7-14%)	10 (2.3-19)	190 (48-360)
Bhutan	2000	Male	75 to 79	39% (20-60%)	61% (40-80%)	2.5 (0.51-6.8)	4.7% (0.73-9.6%)	5.5% (0.75-12%)	6.7 (0.92-14)	100 (16-210)
Bhutan	2000	Male	80 to 84	37% (16-58%)	63% (42-84%)	2.4 (0.53-6.4)	4.4% (0.55-9.2%)	5.1% (0.49-11%)	4.8 (0.43-10)	53 (6.2-110)
Bhutan	2000	Male	85 to 89	37% (16-57%)	63% (43-84%)	2.5 (0.52-6.7)	4% (0.42-8.7%)	4.5% (0.33-10%)	2.8 (0.2-6.2)	22 (2.3-47)
Bhutan	2000	Male	90 to 94	40% (17-59%)	60% (41-83%)	2.7 (0.48-8.1)	3.6% (0.16-8%)	4.1% (-0.039-9.2%)	1.1 (-0.011-2.4)	5.8 (0.25-13)
Bhutan	2000	Male	95 plus	42% (19-64%)	58% (36-81%)	3 (0.41-9.5)	3.4% (0.096-7.7%)	3.6% (-0.13-8.6%)	0.22 (-0.0077-0.53)	0.84 (0.024-1.9)
Bhutan	2005	Female	All Ages	1.8% (1.2-2.7%)	98% (97-99%)	0.21 (0.11-0.38)	0.82% (0.63-1%)	1.1% (0.66-1.8%)	20 (11-31)	990 (740-1300)
Bhutan	2005	Female	15 to 19	1.4% (0.53-2.9%)	99% (97-99%)	0.088 (0.021-0.23)	0.91% (0.59-1.3%)	0.3% (0.097-0.69%)	0.1 (0.032-0.24)	55 (34-81)
Bhutan	2005	Female	20 to 24	1.6% (0.5-3.5%)	98% (97-100%)	0.19 (0.048-0.53)	2.1% (1.4-2.9%)	0.54% (0.22-1.1%)	0.19 (0.075-0.42)	120 (73-180)
Bhutan	2005	Female	25 to 29	1.8% (0.54-3.9%)	98% (96-99%)	0.3 (0.086-0.71)	2.1% (1.6-2.9%)	0.78% (0.35-1.5%)	0.26 (0.11-0.51)	110 (70-150)
Bhutan	2005	Female	30 to 34	2.2% (0.64-4.7%)	98% (95-99%)	0.4 (0.12-0.95)	2% (1.4-2.7%)	1.3% (0.63-2.2%)	0.43 (0.2-0.78)	86 (58-120)
Bhutan	2005	Female	35 to 39	2.6% (0.74-5.7%)	97% (94-99%)	0.48 (0.13-1.1)	2% (1.4-2.8%)	2% (1-3.3%)	0.73 (0.38-1.2)	85 (57-120)
Bhutan	2005	Female	40 to 44	3% (1.1-6.2%)	97% (94-99%)	0.51 (0.18-1.2)	2% (1.4-2.9%)	2.5% (1.4-4.1%)	0.97 (0.54-1.6)	79 (52-110)
Bhutan	2005	Female	45 to 49	3.2% (1.2-6.6%)	97% (93-99%)	0.5 (0.16-1.1)	2.1% (1.4-3.2%)	2.7% (1.6-4.5%)	1.5 (0.82-2.4)	88 (58-130)
Bhutan	2005	Female	50 to 54	3.5% (1.3-7.6%)	96% (92-99%)	0.54 (0.15-1.3)	2.1% (1.4-3.2%)	2.8% (1.5-4.8%)	1.5 (0.76-2.4)	73 (45-110)
Bhutan	2005	Female	55 to 59	3.8% (1.4-8.4%)	96% (92-99%)	0.65 (0.18-1.7)	2.1% (1.2-3.5%)	2.7% (1.4-5%)	1.7 (0.84-3.2)	69 (40-120)
Bhutan	2005	Female	60 to 64	4.1% (1.4-8.8%)	96% (91-99%)	0.69 (0.17-2)	2% (1.1-3.5%)	2.5% (1.2-4.9%)	2 (0.96-3.9)	66 (36-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	2005	Female	65 to 69	4.5% (1.5-9.6%)	95% (90-98%)	0.64 (0.16-1.6)	1.9% (0.9-3.4%)	2.4% (1.1-4.6%)	2.4 (1-4.5)	60 (29-110)
Bhutan	2005	Female	70 to 74	4.7% (1.7-9.4%)	95% (91-98%)	0.54 (0.13-1.4)	1.7% (0.72-3.1%)	2.1% (0.87-4.2%)	2.5 (0.99-4.9)	49 (21-93)
Bhutan	2005	Female	75 to 79	4.8% (1.6-11%)	95% (89-98%)	0.48 (0.1-1.4)	1.4% (0.48-2.9%)	1.7% (0.56-3.6%)	2 (0.64-4.3)	30 (11-63)
Bhutan	2005	Female	80 to 84	5.1% (1.7-12%)	95% (88-98%)	0.46 (0.1-1.4)	1.3% (0.4-3.2%)	1.6% (0.44-3.9%)	1.7 (0.44-4.1)	19 (5.6-44)
Bhutan	2005	Female	85 to 89	5.9% (1.9-13%)	94% (87-98%)	0.48 (0.12-1.4)	1.3% (0.31-3.3%)	1.5% (0.32-4%)	1.1 (0.22-2.8)	8.4 (2-21)
Bhutan	2005	Female	90 to 94	7.4% (2.5-18%)	93% (82-98%)	0.55 (0.12-1.5)	1.3% (0.24-3.4%)	1.5% (0.2-4.1%)	0.49 (0.066-1.3)	2.6 (0.49-6.8)
Bhutan	2005	Female	95 plus	9.9% (2.9-23%)	90% (77-97%)	0.64 (0.12-1.9)	1.4% (0.13-3.8%)	1.5% (0.02-4.6%)	0.14 (0.0019-0.43)	0.52 (0.046-1.5)
Bhutan	2005	Male	All Ages	20% (12-29%)	80% (71-88%)	0.86 (0.55-1.3)	2.9% (2-3.9%)	4.4% (2.2-6.7%)	94 (47-150)	4100 (2800-5500)
Bhutan	2005	Male	15 to 19	5.8% (1.8-15%)	94% (85-98%)	0.19 (0.034-0.6)	1.6% (0.99-2.5%)	1.1% (0.45-2.6%)	0.46 (0.18-1.1)	89 (54-150)
Bhutan	2005	Male	20 to 24	12% (3.2-28%)	88% (72-97%)	0.49 (0.088-1.4)	4.3% (2.9-6.5%)	3.2% (1.3-6.7%)	1.8 (0.71-3.8)	290 (180-460)
Bhutan	2005	Male	25 to 29	20% (4.4-42%)	80% (58-96%)	0.9 (0.17-2.2)	6% (3.8-9.1%)	5.6% (2.1-11%)	3.1 (1.2-6.2)	380 (230-590)
Bhutan	2005	Male	30 to 34	29% (5.6-55%)	71% (45-94%)	1.4 (0.32-3)	7.5% (4.7-11%)	8.6% (4.1-14%)	4.4 (2-7.5)	390 (230-580)
Bhutan	2005	Male	35 to 39	36% (9.9-61%)	64% (39-90%)	1.9 (0.53-3.9)	8.3% (4.9-12%)	10% (5.1-17%)	6.5 (3.1-11)	460 (260-700)
Bhutan	2005	Male	40 to 44	40% (15-64%)	60% (36-85%)	2.2 (0.71-4.3)	8.6% (4.9-13%)	11% (5.4-18%)	6.6 (3.1-11)	390 (230-590)
Bhutan	2005	Male	45 to 49	42% (16-65%)	58% (35-84%)	2.4 (0.73-4.9)	8.7% (4.6-13%)	11% (5.3-18%)	8.9 (4-15)	450 (230-700)
Bhutan	2005	Male	50 to 54	42% (20-64%)	58% (36-80%)	2.5 (0.77-5.4)	8.3% (4.1-13%)	11% (4.5-18%)	8.3 (3.4-14)	360 (160-560)
Bhutan	2005	Male	55 to 59	42% (22-62%)	58% (38-78%)	2.6 (0.78-5.8)	7.8% (3.1-13%)	10% (3.6-17%)	9.8 (3.4-17)	360 (140-590)
Bhutan	2005	Male	60 to 64	43% (23-63%)	57% (37-77%)	2.6 (0.73-6)	6.8% (2.4-12%)	8.6% (2.5-15%)	10 (2.7-18)	310 (100-530)
Bhutan	2005	Male	65 to 69	44% (24-62%)	56% (38-76%)	2.4 (0.72-5.6)	5.9% (1.6-11%)	7.4% (1.7-14%)	10 (2.3-20)	250 (68-480)
Bhutan	2005	Male	70 to 74	42% (24-61%)	58% (39-76%)	2.1 (0.57-5.5)	4.5% (0.7-10%)	5.5% (0.59-13%)	8.7 (0.91-20)	170 (26-380)
Bhutan	2005	Male	75 to 79	39% (20-60%)	61% (40-80%)	1.8 (0.36-5.1)	3.5% (0.29-8.4%)	4.1% (0.13-10%)	6.2 (0.21-16)	95 (7.6-230)
Bhutan	2005	Male	80 to 84	37% (17-58%)	63% (42-83%)	1.8 (0.36-4.7)	3.2% (0.09-8%)	3.7% (-0.068-9.4%)	4.9 (-0.081-13)	54 (1.5-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	2005	Male	85 to 89	37% (16-57%)	63% (43-84%)	1.8 (0.36-5.1)	3% (-0.078-7.6%)	3.4% (-0.26-8.8%)	2.7 (-0.21-7.1)	21 (-0.57-54)
Bhutan	2005	Male	90 to 94	39% (18-59%)	61% (41-82%)	2 (0.36-6)	2.8% (-0.17-7.4%)	3.1% (-0.42-8.6%)	1.1 (-0.15-3)	6 (-0.36-16)
Bhutan	2005	Male	95 plus	43% (19-64%)	57% (36-81%)	2.3 (0.32-7.6)	2.7% (-0.23-7.4%)	2.9% (-0.58-8.3%)	0.27 (-0.053-0.76)	1 (-0.086-2.7)
Bhutan	2010	Female	All Ages	1.8% (1.2-2.7%)	98% (97-99%)	0.19 (0.086-0.36)	0.96% (0.73-1.2%)	1.1% (0.58-1.9%)	17 (8.5-29)	1000 (740-1300)
Bhutan	2010	Female	15 to 19	1.3% (0.51-2.7%)	99% (97-99%)	0.072 (0.016-0.21)	0.97% (0.63-1.4%)	0.26% (0.09-0.62%)	0.065 (0.021-0.16)	52 (32-79)
Bhutan	2010	Female	20 to 24	1.5% (0.48-3.4%)	98% (97-100%)	0.15 (0.037-0.44)	2.3% (1.5-3.2%)	0.5% (0.2-1%)	0.15 (0.058-0.32)	130 (85-200)
Bhutan	2010	Female	25 to 29	1.7% (0.5-3.8%)	98% (96-100%)	0.24 (0.064-0.59)	2.3% (1.7-3%)	0.73% (0.34-1.4%)	0.22 (0.097-0.43)	120 (78-170)
Bhutan	2010	Female	30 to 34	2.1% (0.6-4.5%)	98% (95-99%)	0.33 (0.091-0.77)	2% (1.4-2.7%)	1.2% (0.61-2.1%)	0.39 (0.19-0.71)	100 (66-140)
Bhutan	2010	Female	35 to 39	2.5% (0.7-5.5%)	97% (94-99%)	0.39 (0.1-0.91)	2% (1.5-2.6%)	1.9% (1-3.1%)	0.64 (0.34-1.1)	90 (61-120)
Bhutan	2010	Female	40 to 44	2.8% (1-6.3%)	97% (94-99%)	0.41 (0.13-0.97)	1.9% (1.4-2.7%)	2.3% (1.3-3.8%)	0.91 (0.51-1.5)	85 (57-120)
Bhutan	2010	Female	45 to 49	3.1% (1.1-6.6%)	97% (93-99%)	0.41 (0.12-0.98)	1.9% (1.3-2.8%)	2.4% (1.4-4%)	1.2 (0.66-2.1)	81 (53-120)
Bhutan	2010	Female	50 to 54	3.4% (1.2-7.3%)	97% (93-99%)	0.44 (0.11-1.1)	1.9% (1.2-2.8%)	2.5% (1.3-4.2%)	1.4 (0.72-2.4)	77 (48-120)
Bhutan	2010	Female	55 to 59	3.6% (1.3-7.7%)	96% (92-99%)	0.52 (0.13-1.4)	1.8% (1.1-3%)	2.4% (1.2-4.3%)	1.5 (0.72-2.7)	65 (38-110)
Bhutan	2010	Female	60 to 64	3.9% (1.3-8.5%)	96% (92-99%)	0.56 (0.13-1.6)	1.7% (0.93-3.1%)	2.2% (1-4.4%)	1.7 (0.77-3.4)	58 (31-110)
Bhutan	2010	Female	65 to 69	4.3% (1.6-9.1%)	96% (91-98%)	0.51 (0.12-1.4)	1.6% (0.77-2.9%)	2.1% (0.93-4%)	1.9 (0.81-3.7)	52 (25-95)
Bhutan	2010	Female	70 to 74	4.4% (1.8-8.8%)	96% (91-98%)	0.43 (0.1-1.1)	1.4% (0.59-2.6%)	1.8% (0.69-3.5%)	1.9 (0.7-3.7)	39 (16-73)
Bhutan	2010	Female	75 to 79	4.5% (1.6-9.8%)	96% (90-98%)	0.38 (0.079-1.2)	1.1% (0.38-2.5%)	1.4% (0.44-3.2%)	1.8 (0.52-4)	28 (9.1-60)
Bhutan	2010	Female	80 to 84	4.8% (1.6-12%)	95% (88-98%)	0.37 (0.078-1.1)	1.1% (0.29-2.7%)	1.3% (0.32-3.4%)	1.6 (0.38-4.4)	19 (5-48)
Bhutan	2010	Female	85 to 89	5.7% (1.7-14%)	94% (86-98%)	0.4 (0.083-1.2)	1% (0.18-2.8%)	1.2% (0.17-3.5%)	1.2 (0.16-3.3)	9.3 (1.6-26)
Bhutan	2010	Female	90 to 94	7.2% (2.3-17%)	93% (83-98%)	0.45 (0.088-1.3)	1% (0.1-2.9%)	1.2% (0.032-3.5%)	0.54 (0.015-1.6)	3 (0.29-8.6)
Bhutan	2010	Female	95 plus	9.6% (2.8-22%)	90% (78-97%)	0.53 (0.094-1.6)	1.1% (-0.044-3.4%)	1.2% (-0.19-4%)	0.17 (-0.026-0.57)	0.63 (-0.027-2)
Bhutan	2010	Male	All Ages	22% (13-32%)	78% (68-87%)	0.82 (0.47-1.3)	3.3% (2.2-4.5%)	4.2% (1.8-7%)	91 (39-160)	4200 (2700-6000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	2010	Male	15 to 19	5.8% (1.7-15%)	94% (85-98%)	0.16 (0.027-0.48)	1.6% (1-2.4%)	1% (0.43-2.4%)	0.39 (0.16-0.92)	85 (53-140)
Bhutan	2010	Male	20 to 24	12% (3.1-28%)	88% (72-97%)	0.41 (0.069-1.2)	4.2% (2.9-6%)	2.9% (1.2-5.9%)	1.5 (0.63-3.1)	270 (170-420)
Bhutan	2010	Male	25 to 29	20% (4.6-42%)	80% (58-95%)	0.77 (0.15-1.9)	5.7% (3.7-8.4%)	4.9% (2-9.6%)	3.4 (1.3-6.5)	450 (280-690)
Bhutan	2010	Male	30 to 34	29% (6.2-54%)	71% (46-94%)	1.2 (0.28-2.7)	6.8% (4.4-10%)	7.4% (3.5-13%)	5.2 (2.4-9.2)	500 (300-760)
Bhutan	2010	Male	35 to 39	37% (9.8-61%)	63% (39-90%)	1.6 (0.45-3.5)	7.5% (4.5-11%)	9% (4.3-16%)	5.9 (2.7-11)	450 (260-690)
Bhutan	2010	Male	40 to 44	41% (14-64%)	59% (36-86%)	1.9 (0.57-4)	7.6% (4.4-12%)	9.6% (4.3-16%)	7.4 (3.2-13)	470 (270-730)
Bhutan	2010	Male	45 to 49	42% (15-65%)	58% (35-85%)	2.1 (0.57-4.4)	7.6% (3.9-12%)	9.9% (4.4-17%)	7.7 (3.2-14)	410 (200-670)
Bhutan	2010	Male	50 to 54	42% (19-65%)	58% (35-81%)	2.2 (0.61-4.8)	7.1% (3.3-12%)	9.3% (3.6-16%)	9.3 (3.4-16)	410 (190-690)
Bhutan	2010	Male	55 to 59	42% (22-62%)	58% (38-78%)	2.2 (0.61-5)	6.6% (2.5-11%)	8.6% (2.8-16%)	8.5 (2.7-16)	320 (120-570)
Bhutan	2010	Male	60 to 64	43% (22-63%)	57% (37-78%)	2.2 (0.58-5.2)	5.7% (1.8-11%)	7.1% (1.7-14%)	9.5 (2.3-19)	300 (90-570)
Bhutan	2010	Male	65 to 69	44% (24-62%)	56% (38-76%)	2 (0.53-5)	4.8% (1-10%)	5.9% (0.91-13%)	9.1 (1.4-20)	230 (54-480)
Bhutan	2010	Male	70 to 74	43% (24-61%)	57% (39-76%)	1.8 (0.41-4.8)	3.6% (0.39-9%)	4.4% (0.18-11%)	7.8 (0.34-21)	160 (17-400)
Bhutan	2010	Male	75 to 79	40% (20-60%)	60% (40-80%)	1.6 (0.29-4.5)	2.8% (0.087-7.5%)	3.2% (-0.14-9.1%)	6.1 (-0.26-18)	96 (3-260)
Bhutan	2010	Male	80 to 84	37% (18-58%)	63% (42-82%)	1.5 (0.29-4.3)	2.5% (-0.12-7.3%)	2.8% (-0.33-8.7%)	5.1 (-0.59-16)	59 (-2.8-170)
Bhutan	2010	Male	85 to 89	37% (16-56%)	63% (44-84%)	1.5 (0.27-4.5)	2.3% (-0.22-6.8%)	2.5% (-0.46-8%)	3 (-0.53-9.4)	24 (-2.5-73)
Bhutan	2010	Male	90 to 94	39% (17-59%)	61% (41-83%)	1.7 (0.26-5.8)	2.2% (-0.33-6.6%)	2.3% (-0.63-7.7%)	1.1 (-0.31-3.7)	6.4 (-1-20)
Bhutan	2010	Male	95 plus	42% (19-64%)	58% (36-81%)	1.9 (0.24-6.8)	2.2% (-0.44-6.6%)	2.2% (-0.82-7.3%)	0.29 (-0.11-0.95)	1.2 (-0.23-3.4)
Bhutan	2016	Female	All Ages	1.9% (1.3-2.8%)	98% (97-99%)	0.21 (0.081-0.43)	1.2% (0.92-1.5%)	1.2% (0.56-2.2%)	19 (8.3-33)	1100 (830-1500)
Bhutan	2016	Female	15 to 19	1.3% (0.55-2.7%)	99% (97-99%)	0.071 (0.016-0.22)	1% (0.65-1.5%)	0.27% (0.097-0.66%)	0.049 (0.016-0.11)	49 (29-74)
Bhutan	2016	Female	20 to 24	1.5% (0.5-3.4%)	98% (97-99%)	0.15 (0.033-0.45)	2.4% (1.7-3.3%)	0.5% (0.21-1.1%)	0.12 (0.048-0.26)	130 (82-190)
Bhutan	2016	Female	25 to 29	1.7% (0.49-3.7%)	98% (96-100%)	0.24 (0.06-0.64)	2.4% (1.8-3.3%)	0.72% (0.33-1.4%)	0.21 (0.088-0.41)	140 (96-210)
Bhutan	2016	Female	30 to 34	2.1% (0.59-4.7%)	98% (95-99%)	0.33 (0.084-0.81)	2.1% (1.5-2.9%)	1.2% (0.6-2.1%)	0.39 (0.18-0.71)	120 (78-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	2016	Female	35 to 39	2.6% (0.64-5.5%)	97% (94-99%)	0.39 (0.094-0.96)	2% (1.5-2.6%)	1.8% (0.98-3%)	0.7 (0.36-1.2)	110 (75-150)
Bhutan	2016	Female	40 to 44	2.9% (1-6.1%)	97% (94-99%)	0.41 (0.11-1)	1.9% (1.4-2.6%)	2.2% (1.3-3.5%)	0.92 (0.52-1.6)	98 (65-140)
Bhutan	2016	Female	45 to 49	3.1% (1.1-6.7%)	97% (93-99%)	0.39 (0.1-0.96)	1.8% (1.3-2.6%)	2.3% (1.3-3.8%)	1.2 (0.64-2.1)	93 (61-130)
Bhutan	2016	Female	50 to 54	3.3% (1.2-7.3%)	97% (93-99%)	0.43 (0.1-1.1)	1.8% (1.2-2.7%)	2.3% (1.2-3.9%)	1.4 (0.7-2.5)	82 (51-120)
Bhutan	2016	Female	55 to 59	3.6% (1.3-7.9%)	96% (92-99%)	0.52 (0.11-1.5)	1.7% (1.1-2.8%)	2.2% (1.1-4.1%)	1.7 (0.76-3.1)	77 (45-130)
Bhutan	2016	Female	60 to 64	3.9% (1.3-8.4%)	96% (92-99%)	0.55 (0.12-1.7)	1.6% (0.87-2.9%)	2.1% (0.93-4.2%)	1.8 (0.79-3.7)	65 (34-120)
Bhutan	2016	Female	65 to 69	4.2% (1.5-8.9%)	96% (91-98%)	0.5 (0.1-1.4)	1.5% (0.7-2.7%)	2% (0.8-3.7%)	1.9 (0.75-3.8)	54 (25-99)
Bhutan	2016	Female	70 to 74	4.3% (1.6-8.7%)	96% (91-98%)	0.42 (0.085-1.2)	1.3% (0.52-2.4%)	1.7% (0.6-3.3%)	2 (0.68-3.9)	42 (17-78)
Bhutan	2016	Female	75 to 79	4.4% (1.6-9.5%)	96% (91-98%)	0.37 (0.072-1.2)	1% (0.34-2.2%)	1.3% (0.36-2.9%)	1.8 (0.47-4)	28 (8.7-61)
Bhutan	2016	Female	80 to 84	4.7% (1.6-11%)	95% (89-98%)	0.36 (0.07-1.1)	0.96% (0.21-2.3%)	1.2% (0.2-2.9%)	1.9 (0.3-4.8)	22 (4.5-54)
Bhutan	2016	Female	85 to 89	5.6% (1.7-13%)	94% (87-98%)	0.39 (0.072-1.2)	0.95% (0.12-2.6%)	1.1% (0.088-3.2%)	1.7 (0.12-4.8)	13 (1.7-37)
Bhutan	2016	Female	90 to 94	7.3% (2.3-17%)	93% (83-98%)	0.45 (0.085-1.4)	0.96% (0.00036-2.8%)	1.1% (-0.082-3.4%)	0.87 (-0.067-2.7)	4.8 (0.0018-14)
Bhutan	2016	Female	95 plus	9.8% (2.9-23%)	90% (77-97%)	0.54 (0.095-1.8)	1.1% (-0.11-3.3%)	1.1% (-0.29-3.9%)	0.28 (-0.072-1)	1.1 (-0.12-3.5)
Bhutan	2016	Male	All Ages	25% (14-35%)	75% (65-86%)	0.92 (0.43-1.7)	3.9% (2.4-5.7%)	4.4% (1.6-8%)	100 (35-190)	4800 (2800-7200)
Bhutan	2016	Male	15 to 19	5.8% (1.9-14%)	94% (86-98%)	0.16 (0.027-0.51)	1.7% (1.1-2.5%)	1.1% (0.45-2.5%)	0.33 (0.14-0.76)	81 (49-130)
Bhutan	2016	Male	20 to 24	12% (3.3-27%)	88% (73-97%)	0.41 (0.068-1.2)	4.4% (2.9-6.5%)	2.9% (1.2-6.1%)	1.4 (0.57-3)	270 (170-420)
Bhutan	2016	Male	25 to 29	20% (4.5-42%)	80% (58-96%)	0.76 (0.13-2)	5.8% (3.8-8.5%)	4.8% (1.9-9.5%)	2.9 (1.1-5.9)	420 (260-640)
Bhutan	2016	Male	30 to 34	30% (6-55%)	70% (45-94%)	1.2 (0.26-2.9)	6.8% (4.3-9.9%)	7.2% (3.3-13%)	5.8 (2.6-11)	600 (350-930)
Bhutan	2016	Male	35 to 39	38% (9.6-63%)	62% (37-90%)	1.6 (0.38-3.8)	7.2% (4.4-11%)	8.5% (3.8-15%)	7.5 (3.2-14)	610 (360-960)
Bhutan	2016	Male	40 to 44	42% (13-65%)	58% (35-87%)	1.8 (0.52-4.3)	7.3% (4.2-11%)	9.2% (4.2-16%)	7.6 (3.1-14)	500 (280-830)
Bhutan	2016	Male	45 to 49	43% (14-65%)	57% (35-86%)	2 (0.45-4.7)	7.2% (3.8-12%)	9.4% (4-17%)	9.4 (3.8-18)	510 (260-870)
Bhutan	2016	Male	50 to 54	42% (19-64%)	58% (36-81%)	2.1 (0.5-5.1)	6.8% (3-11%)	8.8% (3.3-15%)	9.3 (3.3-18)	420 (180-740)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bhutan	2016	Male	55 to 59	42% (22-62%)	58% (38-78%)	2.2 (0.53-5.5)	6.2% (2.3-11%)	8% (2.4-15%)	11 (3.1-21)	410 (150-760)
Bhutan	2016	Male	60 to 64	43% (22-63%)	57% (37-78%)	2.1 (0.49-5.8)	5.3% (1.5-10%)	6.5% (1.4-13%)	9.8 (2.1-20)	310 (87-620)
Bhutan	2016	Male	65 to 69	43% (23-63%)	57% (37-77%)	2 (0.46-5)	4.4% (0.9-9.7%)	5.4% (0.68-13%)	10 (1.3-23)	260 (50-570)
Bhutan	2016	Male	70 to 74	42% (24-61%)	58% (39-76%)	1.7 (0.36-5.1)	3.4% (0.32-8.6%)	4% (0.086-11%)	8.8 (0.17-24)	180 (16-470)
Bhutan	2016	Male	75 to 79	39% (19-60%)	61% (40-81%)	1.5 (0.25-4.9)	2.5% (0.041-7.2%)	2.9% (-0.18-8.6%)	6.8 (-0.44-20)	110 (1.8-310)
Bhutan	2016	Male	80 to 84	37% (17-58%)	63% (42-83%)	1.4 (0.24-4.5)	2.3% (-0.23-7%)	2.5% (-0.43-8.4%)	6.4 (-1.1-21)	74 (-7.3-230)
Bhutan	2016	Male	85 to 89	37% (16-57%)	63% (43-84%)	1.5 (0.24-4.5)	2.1% (-0.33-6.5%)	2.3% (-0.58-7.7%)	4.2 (-1.14)	34 (-5.3-110)
Bhutan	2016	Male	90 to 94	40% (18-59%)	60% (41-82%)	1.7 (0.26-5.7)	2% (-0.42-6.2%)	2.1% (-0.75-7.1%)	1.7 (-0.6-5.9)	9.7 (-2.1-31)
Bhutan	2016	Male	95 plus	43% (20-64%)	57% (36-80%)	1.9 (0.24-6.7)	2% (-0.49-6.2%)	2% (-0.85-7%)	0.42 (-0.18-1.4)	1.7 (-0.41-5.2)
India	1990	Female	All Ages	4.7% (3.4-6.3%)	95% (94-97%)	0.12 (0.089-0.16)	0.42% (0.31-0.55%)	0.66% (0.43-0.97%)	28000 (18000-41000)	1100000 (800000-1400000)
India	1990	Female	15 to 19	3.3% (1.6-6%)	97% (94-98%)	0.042 (0.017-0.087)	0.2% (0.11-0.34%)	0.13% (0.025-0.32%)	140 (26-330)	23000 (13000-39000)
India	1990	Female	20 to 24	3.7% (1.6-6.6%)	96% (93-98%)	0.088 (0.033-0.18)	0.49% (0.3-0.77%)	0.37% (0.15-0.75%)	430 (170-890)	57000 (34000-92000)
India	1990	Female	25 to 29	4.2% (1.9-7.9%)	96% (92-98%)	0.15 (0.057-0.29)	0.77% (0.49-1.3%)	0.74% (0.34-1.4%)	780 (350-1500)	78000 (48000-130000)
India	1990	Female	30 to 34	5.2% (2.3-9.6%)	95% (90-98%)	0.21 (0.086-0.4)	1.1% (0.68-1.7%)	1.3% (0.68-2.2%)	1200 (650-2100)	97000 (61000-150000)
India	1990	Female	35 to 39	6.1% (2.6-10%)	94% (90-97%)	0.25 (0.1-0.46)	1.3% (0.86-2%)	1.8% (1-2.9%)	1700 (1000-2800)	110000 (73000-170000)
India	1990	Female	40 to 44	6.8% (3.4-12%)	93% (88-97%)	0.27 (0.12-0.49)	1.4% (0.94-2.1%)	2% (1.2-3.2%)	1900 (1200-3000)	110000 (69000-160000)
India	1990	Female	45 to 49	7.2% (3.4-12%)	93% (88-97%)	0.25 (0.099-0.48)	1.3% (0.89-2%)	1.8% (1.1-3%)	2100 (1300-3500)	100000 (68000-160000)
India	1990	Female	50 to 54	7.8% (3.8-13%)	92% (87-96%)	0.27 (0.11-0.53)	1.4% (0.88-2.1%)	1.8% (1.1-2.9%)	3000 (1800-4800)	120000 (78000-180000)
India	1990	Female	55 to 59	8.4% (4.1-16%)	92% (84-96%)	0.34 (0.14-0.68)	1.4% (0.82-2.3%)	1.8% (0.99-3%)	3700 (2100-6300)	130000 (75000-210000)
India	1990	Female	60 to 64	9.4% (4.6-18%)	91% (82-95%)	0.38 (0.15-0.78)	1.2% (0.63-2%)	1.4% (0.75-2.6%)	3700 (1900-6500)	100000 (57000-180000)
India	1990	Female	65 to 69	9.9% (4.9-18%)	90% (82-95%)	0.36 (0.15-0.69)	0.95% (0.51-1.7%)	1.1% (0.59-2%)	3400 (1700-6000)	78000 (42000-140000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	1990	Female	70 to 74	10% (5.1-17%)	90% (83-95%)	0.32 (0.13-0.62)	0.72% (0.34-1.4%)	0.85% (0.39-1.7%)	2600 (1200-5100)	48000 (23000-94000)
India	1990	Female	75 to 79	9.7% (4.5-18%)	90% (82-95%)	0.28 (0.096-0.58)	0.51% (0.2-1.1%)	0.59% (0.22-1.3%)	1600 (570-3200)	22000 (8900-47000)
India	1990	Female	80 to 84	10% (4.6-20%)	90% (80-95%)	0.28 (0.1-0.66)	0.43% (0.14-0.97%)	0.49% (0.15-1.1%)	1000 (320-2300)	11000 (3700-25000)
India	1990	Female	85 to 89	11% (5.1-22%)	89% (78-95%)	0.31 (0.11-0.69)	0.34% (0.091-0.77%)	0.38% (0.087-0.89%)	410 (95-980)	3200 (870-7400)
India	1990	Female	90 to 94	13% (6.1-27%)	87% (73-94%)	0.37 (0.13-0.83)	0.26% (-0.0062-0.68%)	0.28% (-0.04-0.77%)	100 (-14-280)	570 (-13-1500)
India	1990	Female	95 plus	17% (6.7-33%)	83% (67-93%)	0.46 (0.14-1.1)	0.22% (-0.13-0.76%)	0.2% (-0.19-0.81%)	17 (-16-70)	76 (-42-260)
India	1990	Male	All Ages	17% (15-19%)	83% (81-85%)	0.68 (0.56-0.85)	2.4% (1.9-2.9%)	3.2% (2.3-4.1%)	150000 (110000-190000)	6600000 (5200000-8000000)
India	1990	Male	15 to 19	7.3% (5-10%)	93% (90-95%)	0.23 (0.12-0.39)	1.2% (0.79-1.7%)	1.1% (0.61-1.8%)	910 (520-1500)	110000 (72000-160000)
India	1990	Male	20 to 24	16% (11-22%)	84% (78-89%)	0.66 (0.35-1.1)	3.7% (2.6-5.1%)	3.7% (2.3-5.5%)	3900 (2400-5800)	370000 (250000-510000)
India	1990	Male	25 to 29	25% (18-33%)	75% (67-82%)	1.1 (0.68-1.6)	6.1% (4.4-7.9%)	6.5% (4.3-9%)	7300 (4800-9900)	610000 (430000-790000)
India	1990	Male	30 to 34	32% (23-42%)	68% (58-77%)	1.4 (0.92-2.1)	7.9% (5.9-10%)	9% (6.3-12%)	12000 (8100-15000)	810000 (600000-1000000)
India	1990	Male	35 to 39	36% (26-45%)	64% (55-74%)	1.6 (1-2.3)	8.1% (6.1-11%)	9.7% (6.8-13%)	14000 (9900-19000)	850000 (630000-1100000)
India	1990	Male	40 to 44	36% (27-45%)	64% (55-73%)	1.7 (1-2.4)	8% (5.7-11%)	9.7% (6.6-13%)	14000 (9800-19000)	750000 (530000-980000)
India	1990	Male	45 to 49	34% (24-42%)	66% (58-76%)	1.6 (0.96-2.4)	7.4% (5.1-10%)	8.9% (6-12%)	16000 (11000-22000)	730000 (500000-990000)
India	1990	Male	50 to 54	31% (24-40%)	69% (60-76%)	1.5 (0.89-2.2)	6.4% (4.1-9%)	7.5% (4.7-11%)	17000 (11000-25000)	680000 (430000-950000)
India	1990	Male	55 to 59	30% (22-38%)	70% (62-78%)	1.4 (0.84-2.1)	5.5% (3.7-7.7%)	6.5% (4.2-9.1%)	18000 (12000-25000)	610000 (410000-850000)
India	1990	Male	60 to 64	28% (20-37%)	72% (63-80%)	1.3 (0.76-2)	4.3% (2.7-6.6%)	4.9% (2.9-7.7%)	17000 (9800-26000)	480000 (290000-730000)
India	1990	Male	65 to 69	27% (20-35%)	73% (65-80%)	1.2 (0.72-1.8)	3.3% (1.9-4.9%)	3.7% (2.1-5.6%)	13000 (7500-20000)	310000 (180000-460000)
India	1990	Male	70 to 74	25% (19-33%)	75% (67-81%)	0.97 (0.58-1.5)	2.3% (1.2-3.7%)	2.6% (1.3-4.2%)	8800 (4300-14000)	170000 (87000-270000)
India	1990	Male	75 to 79	22% (15-30%)	78% (70-85%)	0.77 (0.42-1.2)	1.5% (0.72-2.6%)	1.7% (0.73-2.9%)	4600 (2000-8100)	69000 (32000-120000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	1990	Male	80 to 84	19% (13-26%)	81% (74-87%)	0.62 (0.36-0.96)	1% (0.41-1.8%)	1.1% (0.4-2%)	1900 (710-3600)	22000 (8800-39000)
India	1990	Male	85 to 89	19% (13-26%)	81% (74-87%)	0.55 (0.31-0.87)	0.67% (0.22-1.3%)	0.69% (0.16-1.4%)	670 (150-1300)	5600 (1800-10000)
India	1990	Male	90 to 94	21% (15-29%)	79% (71-85%)	0.53 (0.29-0.88)	0.43% (0.049-0.93%)	0.39% (-0.049-0.95%)	130 (-16-310)	860 (100-1800)
India	1990	Male	95 plus	24% (16-34%)	76% (66-84%)	0.53 (0.23-0.93)	0.34% (-0.028-0.85%)	0.23% (-0.18-0.79%)	16 (-13-57)	97 (-7.6-240)
India	1995	Female	All Ages	2.5% (2.2-2.9%)	98% (97-98%)	0.081 (0.064-0.1)	0.37% (0.3-0.44%)	0.58% (0.42-0.76%)	24000 (18000-32000)	950000 (760000-1200000)
India	1995	Female	15 to 19	1.3% (0.86-2%)	99% (98-99%)	0.02 (0.0091-0.037)	0.16% (0.11-0.24%)	0.076% (0.031-0.15%)	82 (34-170)	19000 (13000-29000)
India	1995	Female	20 to 24	1.6% (1-2.3%)	98% (98-99%)	0.045 (0.023-0.078)	0.37% (0.26-0.52%)	0.21% (0.12-0.36%)	260 (140-440)	45000 (30000-65000)
India	1995	Female	25 to 29	2.2% (1.4-3.2%)	98% (97-99%)	0.087 (0.045-0.15)	0.57% (0.41-0.78%)	0.46% (0.27-0.75%)	490 (290-800)	59000 (42000-83000)
India	1995	Female	30 to 34	2.9% (1.9-4.2%)	97% (96-98%)	0.13 (0.071-0.22)	0.79% (0.59-1.1%)	0.85% (0.59-1.2%)	860 (580-1200)	76000 (55000-100000)
India	1995	Female	35 to 39	3.5% (2.3-4.7%)	97% (95-98%)	0.16 (0.089-0.25)	1% (0.77-1.3%)	1.3% (0.92-1.8%)	1300 (920-1800)	92000 (69000-120000)
India	1995	Female	40 to 44	3.9% (2.7-5.5%)	96% (95-97%)	0.17 (0.096-0.28)	1.1% (0.84-1.5%)	1.5% (1.1-2%)	1700 (1200-2300)	100000 (75000-130000)
India	1995	Female	45 to 49	4.2% (2.7-5.7%)	96% (94-97%)	0.17 (0.087-0.29)	1.1% (0.86-1.5%)	1.5% (1.1-2%)	1800 (1300-2500)	92000 (70000-120000)
India	1995	Female	50 to 54	5% (3.5-7%)	95% (93-96%)	0.2 (0.11-0.33)	1.2% (0.87-1.6%)	1.5% (1.1-2.1%)	2500 (1800-3500)	100000 (77000-140000)
India	1995	Female	55 to 59	5.2% (3.7-7.3%)	95% (93-96%)	0.24 (0.13-0.4)	1.1% (0.78-1.6%)	1.4% (0.93-2%)	3100 (2100-4500)	110000 (76000-150000)
India	1995	Female	60 to 64	5.3% (3.6-7.5%)	95% (92-96%)	0.25 (0.12-0.43)	0.9% (0.6-1.3%)	1.1% (0.7-1.7%)	3000 (1900-4700)	89000 (59000-130000)
India	1995	Female	65 to 69	5.4% (3.6-7.7%)	95% (92-96%)	0.23 (0.12-0.39)	0.76% (0.5-1.1%)	0.91% (0.59-1.4%)	2900 (1900-4400)	69000 (45000-100000)
India	1995	Female	70 to 74	6.2% (4.2-9.1%)	94% (91-96%)	0.23 (0.12-0.41)	0.62% (0.38-0.96%)	0.73% (0.44-1.2%)	2400 (1400-3800)	45000 (28000-71000)
India	1995	Female	75 to 79	7.5% (4.8-11%)	93% (89-95%)	0.26 (0.12-0.5)	0.51% (0.27-0.9%)	0.6% (0.3-1.1%)	1700 (870-3000)	25000 (13000-44000)
India	1995	Female	80 to 84	9.4% (5.7-14%)	91% (86-94%)	0.32 (0.14-0.65)	0.48% (0.2-0.94%)	0.56% (0.23-1.1%)	1300 (530-2500)	14000 (5900-27000)
India	1995	Female	85 to 89	11% (6.6-18%)	89% (82-93%)	0.4 (0.17-0.79)	0.44% (0.17-0.89%)	0.5% (0.17-1%)	670 (230-1400)	5200 (1900-10000)
India	1995	Female	90 to 94	13% (7.8-22%)	87% (78-92%)	0.49 (0.21-1)	0.4% (0.11-0.91%)	0.44% (0.1-1%)	210 (49-480)	1200 (330-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	1995	Female	95 plus	16% (8.8-28%)	84% (72-91%)	0.62 (0.21-1.3)	0.41% (0.039-1.1%)	0.41% (-0.016-1.2%)	46 (-1.8-130)	180 (17-470)
India	1995	Male	All Ages	16% (15-18%)	84% (82-85%)	0.89 (0.72-1.1)	3% (2.5-3.6%)	4% (3.1-4.9%)	190000 (150000-240000)	820000 (670000-960000)
India	1995	Male	15 to 19	3.9% (2.9-5.3%)	96% (95-97%)	0.16 (0.089-0.27)	0.99% (0.73-1.3%)	0.85% (0.54-1.3%)	730 (460-1100)	96000 (69000-130000)
India	1995	Male	20 to 24	12% (8.7-15%)	88% (85-91%)	0.65 (0.37-1)	3.5% (2.6-4.7%)	3.5% (2.3-5%)	3900 (2600-5600)	380000 (270000-510000)
India	1995	Male	25 to 29	22% (18-27%)	78% (73-82%)	1.3 (0.85-1.9)	6.3% (4.7-7.9%)	6.9% (4.8-9.3%)	8300 (5700-11000)	670000 (500000-860000)
India	1995	Male	30 to 34	31% (25-37%)	69% (63-75%)	1.9 (1.2-2.6)	8.5% (6.6-11%)	10% (7.3-13%)	14000 (10000-18000)	960000 (740000-1200000)
India	1995	Male	35 to 39	35% (29-41%)	65% (59-71%)	2.2 (1.5-3)	9.1% (7.1-11%)	11% (8.3-14%)	17000 (13000-22000)	1000000 (810000-1300000)
India	1995	Male	40 to 44	36% (30-42%)	64% (58-70%)	2.2 (1.5-3.1)	9.1% (7-11%)	11% (8.4-14%)	20000 (15000-26000)	1100000 (820000-1300000)
India	1995	Male	45 to 49	34% (28-40%)	66% (60-72%)	2.1 (1.4-3.1)	8.5% (6.3-11%)	10% (7.5-13%)	20000 (15000-27000)	930000 (690000-1200000)
India	1995	Male	50 to 54	32% (27-38%)	68% (62-73%)	2 (1.3-2.9)	7.5% (5.2-10%)	9% (6-12%)	21000 (14000-28000)	840000 (580000-1100000)
India	1995	Male	55 to 59	30% (25-36%)	70% (64-75%)	1.9 (1.3-2.8)	6.6% (4.7-8.8%)	7.8% (5.3-10%)	23000 (16000-32000)	790000 (560000-1000000)
India	1995	Male	60 to 64	28% (23-34%)	72% (66-77%)	1.7 (1.1-2.5)	5.1% (3.4-7.3%)	5.9% (3.8-8.4%)	21000 (14000-31000)	610000 (410000-870000)
India	1995	Male	65 to 69	26% (20-31%)	74% (69-80%)	1.5 (0.93-2.1)	3.9% (2.6-5.5%)	4.5% (2.9-6.4%)	17000 (11000-25000)	410000 (270000-580000)
India	1995	Male	70 to 74	24% (19-29%)	76% (71-81%)	1.2 (0.77-1.8)	2.8% (1.7-4.3%)	3.1% (1.8-4.9%)	12000 (8000-18000)	230000 (130000-340000)
India	1995	Male	75 to 79	21% (16-26%)	79% (74-84%)	0.97 (0.58-1.5)	1.9% (1-3%)	2.1% (1.1-3.4%)	6600 (3400-11000)	98000 (54000-160000)
India	1995	Male	80 to 84	19% (14-24%)	81% (76-86%)	0.8 (0.49-1.2)	1.3% (0.69-2.2%)	1.5% (0.72-2.4%)	3000 (1500-4900)	33000 (17000-53000)
India	1995	Male	85 to 89	19% (14-25%)	81% (75-86%)	0.74 (0.45-1.1)	0.96% (0.44-1.7%)	1% (0.41-1.8%)	1200 (470-2100)	9600 (4400-16000)
India	1995	Male	90 to 94	21% (16-27%)	79% (73-84%)	0.72 (0.41-1.1)	0.67% (0.23-1.3%)	0.65% (0.12-1.4%)	270 (51-570)	1700 (580-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	1995	Male	95 plus	24% (16-31%)	76% (69-84%)	0.7 (0.32-1.2)	0.53% (0.091-1.2%)	0.43% (-0.064-1.1%)	40 (-6.1-110)	200 (34-430)
India	2000	Female	All Ages	2.4% (2.1-2.7%)	98% (97-98%)	0.081 (0.064-0.1)	0.4% (0.33-0.48%)	0.6% (0.44-0.78%)	26000 (19000-34000)	1000000 (820000-1200000)
India	2000	Female	15 to 19	1.3% (0.9-1.9%)	99% (98-99%)	0.02 (0.01-0.036)	0.17% (0.12-0.25%)	0.081% (0.035-0.15%)	90 (39-170)	22000 (14000-32000)
India	2000	Female	20 to 24	1.6% (1.1-2.3%)	98% (98-99%)	0.045 (0.025-0.076)	0.4% (0.28-0.56%)	0.22% (0.13-0.36%)	280 (160-450)	51000 (33000-73000)
India	2000	Female	25 to 29	2.1% (1.5-2.9%)	98% (97-99%)	0.085 (0.048-0.14)	0.59% (0.43-0.78%)	0.46% (0.29-0.7%)	510 (320-790)	65000 (47000-89000)
India	2000	Female	30 to 34	2.6% (1.9-3.7%)	97% (96-98%)	0.13 (0.073-0.19)	0.79% (0.6-1%)	0.82% (0.6-1.1%)	860 (620-1200)	79000 (59000-110000)
India	2000	Female	35 to 39	3.2% (2.3-4.1%)	97% (96-98%)	0.15 (0.089-0.22)	1% (0.78-1.3%)	1.3% (0.92-1.7%)	1300 (990-1800)	97000 (75000-120000)
India	2000	Female	40 to 44	3.5% (2.5-4.6%)	97% (95-97%)	0.16 (0.094-0.25)	1.1% (0.84-1.4%)	1.4% (1.1-1.9%)	1700 (1300-2200)	100000 (80000-130000)
India	2000	Female	45 to 49	3.8% (2.5-5%)	96% (95-97%)	0.15 (0.085-0.25)	1.1% (0.84-1.4%)	1.4% (1.1-1.9%)	2000 (1500-2700)	110000 (82000-130000)
India	2000	Female	50 to 54	4.5% (3.2-6.3%)	95% (94-97%)	0.19 (0.1-0.3)	1.1% (0.84-1.5%)	1.5% (1.1-2%)	2500 (1800-3500)	110000 (80000-140000)
India	2000	Female	55 to 59	4.9% (3.5-6.8%)	95% (93-97%)	0.24 (0.13-0.39)	1.1% (0.77-1.5%)	1.4% (0.92-2%)	3000 (2000-4300)	110000 (75000-150000)
India	2000	Female	60 to 64	4.8% (3.3-7%)	95% (93-97%)	0.24 (0.12-0.42)	0.88% (0.59-1.3%)	1.1% (0.68-1.6%)	3200 (2000-4800)	94000 (63000-140000)
India	2000	Female	65 to 69	5% (3.3-7%)	95% (93-97%)	0.23 (0.12-0.37)	0.74% (0.5-1.1%)	0.89% (0.59-1.3%)	3200 (2100-4800)	77000 (52000-110000)
India	2000	Female	70 to 74	5.8% (3.9-8.4%)	94% (92-96%)	0.23 (0.12-0.4)	0.6% (0.37-0.92%)	0.72% (0.43-1.1%)	2700 (1600-4200)	51000 (31000-79000)
India	2000	Female	75 to 79	6.9% (4.4-11%)	93% (89-96%)	0.26 (0.12-0.49)	0.49% (0.26-0.84%)	0.58% (0.29-0.99%)	1900 (970-3300)	28000 (15000-47000)
India	2000	Female	80 to 84	8.6% (5.2-13%)	91% (87-95%)	0.32 (0.14-0.64)	0.45% (0.2-0.88%)	0.53% (0.23-1%)	1400 (620-2700)	15000 (6900-28000)
India	2000	Female	85 to 89	10% (5.8-17%)	90% (83-94%)	0.38 (0.16-0.79)	0.42% (0.16-0.85%)	0.48% (0.17-0.98%)	750 (270-1500)	5900 (2200-12000)
India	2000	Female	90 to 94	12% (6.8-21%)	88% (79-93%)	0.48 (0.2-0.97)	0.39% (0.11-0.89%)	0.43% (0.1-1%)	260 (61-610)	1400 (420-3300)
India	2000	Female	95 plus	15% (7.7-27%)	85% (73-92%)	0.61 (0.21-1.3)	0.41% (0.055-1.1%)	0.41% (0.0064-1.2%)	62 (0.96-180)	250 (33-650)
India	2000	Male	All Ages	17% (16-19%)	83% (81-84%)	1.1 (0.86-1.3)	3.4% (2.9-4%)	4.2% (3.3-5.1%)	210000 (160000-260000)	9300000 (7800000-11000000)
India	2000	Male	15 to 19	7.5% (5.6-9.7%)	93% (90-94%)	0.32 (0.19-0.52)	1.4% (1-2%)	1.4% (0.85-2.1%)	1300 (810-2000)	150000 (110000-210000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	2000	Male	20 to 24	19% (15-24%)	81% (76-85%)	1.2 (0.73-1.7)	4.9% (3.7-6.4%)	5.2% (3.5-7.3%)	6200 (4200-8700)	580000 (430000-770000)
India	2000	Male	25 to 29	28% (23-33%)	72% (67-77%)	1.8 (1.2-2.6)	7.4% (5.7-9.1%)	8.1% (5.8-11%)	11000 (7900-14000)	890000 (680000-1100000)
India	2000	Male	30 to 34	33% (27-39%)	67% (61-73%)	2.2 (1.5-3.1)	8.9% (7-11%)	10% (7.8-13%)	16000 (12000-21000)	1100000 (890000-1400000)
India	2000	Male	35 to 39	35% (30-40%)	65% (60-70%)	2.4 (1.6-3.2)	9% (7.1-11%)	11% (8.3-13%)	20000 (15000-24000)	1200000 (930000-1400000)
India	2000	Male	40 to 44	35% (30-41%)	65% (59-70%)	2.4 (1.6-3.3)	8.9% (6.9-11%)	11% (8.2-14%)	22000 (17000-28000)	1200000 (910000-1500000)
India	2000	Male	45 to 49	33% (27-39%)	67% (61-73%)	2.3 (1.5-3.3)	8.3% (6.3-11%)	10% (7.5-13%)	25000 (19000-32000)	1200000 (870000-1500000)
India	2000	Male	50 to 54	30% (24-36%)	70% (64-76%)	2 (1.3-3)	7.2% (5.1-9.4%)	8.7% (5.9-11%)	22000 (15000-30000)	890000 (630000-1200000)
India	2000	Male	55 to 59	26% (20-32%)	74% (68-80%)	1.8 (1.1-2.7)	6.1% (4.3-8%)	7.2% (4.9-9.7%)	22000 (15000-30000)	750000 (530000-1000000)
India	2000	Male	60 to 64	23% (18-30%)	77% (70-82%)	1.5 (0.92-2.4)	4.6% (3.1-6.6%)	5.3% (3.5-7.6%)	20000 (13000-29000)	590000 (400000-830000)
India	2000	Male	65 to 69	21% (16-27%)	79% (73-84%)	1.4 (0.81-2)	3.6% (2.4-5.1%)	4% (2.6-5.8%)	17000 (11000-25000)	410000 (280000-590000)
India	2000	Male	70 to 74	20% (15-25%)	80% (75-85%)	1.1 (0.67-1.7)	2.5% (1.6-3.9%)	2.8% (1.7-4.4%)	12000 (7200-19000)	230000 (140000-350000)
India	2000	Male	75 to 79	16% (12-22%)	84% (78-88%)	0.84 (0.47-1.3)	1.7% (0.98-2.7%)	1.9% (1-3%)	6600 (3600-11000)	99000 (57000-160000)
India	2000	Male	80 to 84	15% (9.9-20%)	85% (80-90%)	0.67 (0.39-1.1)	1.2% (0.64-1.8%)	1.3% (0.65-2.1%)	3000 (1600-4900)	34000 (19000-54000)
India	2000	Male	85 to 89	15% (10-21%)	85% (79-90%)	0.62 (0.35-0.98)	0.84% (0.41-1.4%)	0.88% (0.38-1.6%)	1200 (520-2100)	10000 (4900-17000)
India	2000	Male	90 to 94	16% (11-23%)	84% (77-89%)	0.61 (0.33-1)	0.6% (0.23-1.1%)	0.57% (0.15-1.2%)	290 (77-620)	1800 (730-3500)
India	2000	Male	95 plus	19% (12-27%)	81% (73-88%)	0.6 (0.26-1.1)	0.49% (0.12-1.1%)	1.3% (-0.019-1%)	47 (-2.3-130)	240 (59-510)
India	2005	Female	All Ages	2.5% (2.2-2.8%)	98% (97-98%)	0.09 (0.072-0.11)	0.4% (0.33-0.49%)	0.6% (0.43-0.8%)	26000 (19000-35000)	990000 (800000-1200000)
India	2005	Female	15 to 19	1.3% (0.92-2%)	99% (98-99%)	0.022 (0.011-0.04)	0.17% (0.11-0.24%)	0.089% (0.041-0.17%)	91 (41-170)	21000 (13000-31000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	2005	Female	20 to 24	1.7% (1.1-2.3%)	98% (98-99%)	0.05 (0.028-0.083)	0.37% (0.26-0.52%)	0.23% (0.14-0.38%)	280 (170-470)	49000 (33000-70000)
India	2005	Female	25 to 29	2.1% (1.5-2.9%)	98% (97-98%)	0.093 (0.052-0.15)	0.54% (0.4-0.72%)	0.47% (0.3-0.73%)	530 (330-820)	63000 (45000-86000)
India	2005	Female	30 to 34	2.6% (1.9-3.7%)	97% (96-98%)	0.14 (0.079-0.21)	0.71% (0.54-0.92%)	0.79% (0.56-1.1%)	820 (580-1100)	75000 (56000-100000)
India	2005	Female	35 to 39	3.2% (2.3-4.2%)	97% (96-98%)	0.16 (0.097-0.25)	0.9% (0.7-1.2%)	1.2% (0.88-1.6%)	1300 (940-1700)	92000 (70000-120000)
India	2005	Female	40 to 44	3.6% (2.6-4.7%)	96% (95-97%)	0.17 (0.1-0.27)	0.96% (0.74-1.2%)	1.4% (1-1.8%)	1600 (1200-2100)	98000 (76000-120000)
India	2005	Female	45 to 49	3.8% (2.6-5%)	96% (95-97%)	0.17 (0.093-0.27)	0.94% (0.73-1.2%)	1.3% (0.99-1.8%)	1900 (1400-2600)	99000 (77000-130000)
India	2005	Female	50 to 54	4.3% (3-6%)	96% (94-97%)	0.19 (0.11-0.32)	1% (0.74-1.4%)	1.4% (0.98-1.9%)	2500 (1800-3500)	110000 (81000-150000)
India	2005	Female	55 to 59	4.7% (3.3-6.4%)	95% (94-97%)	0.25 (0.13-0.41)	0.99% (0.68-1.4%)	1.3% (0.85-1.9%)	2900 (1900-4200)	100000 (73000-150000)
India	2005	Female	60 to 64	4.7% (3.2-6.8%)	95% (93-97%)	0.25 (0.13-0.43)	0.83% (0.54-1.2%)	1% (0.65-1.6%)	3000 (1900-4600)	90000 (59000-140000)
India	2005	Female	65 to 69	5% (3.3-7%)	95% (93-97%)	0.24 (0.13-0.39)	0.7% (0.46-1%)	0.85% (0.55-1.3%)	3300 (2100-5000)	79000 (53000-120000)
India	2005	Female	70 to 74	5.6% (3.8-8.2%)	94% (92-96%)	0.24 (0.12-0.41)	0.57% (0.33-0.9%)	0.68% (0.4-1.1%)	2900 (1700-4700)	55000 (33000-87000)
India	2005	Female	75 to 79	6.6% (4.2-10%)	93% (90-96%)	0.26 (0.12-0.49)	0.47% (0.23-0.83%)	0.56% (0.27-0.98%)	2100 (1000-3700)	30000 (15000-53000)
India	2005	Female	80 to 84	8.2% (4.9-13%)	92% (87-95%)	0.32 (0.14-0.61)	0.43% (0.19-0.83%)	0.51% (0.21-0.99%)	1500 (640-3000)	17000 (7300-33000)
India	2005	Female	85 to 89	9.6% (5.7-16%)	90% (84-94%)	0.38 (0.16-0.78)	0.41% (0.15-0.84%)	0.46% (0.15-0.96%)	870 (290-1800)	6900 (2500-14000)
India	2005	Female	90 to 94	11% (6.5-19%)	89% (81-93%)	0.47 (0.2-0.94)	0.39% (0.11-0.89%)	0.42% (0.088-1%)	310 (66-750)	1800 (480-4100)
India	2005	Female	95 plus	14% (7.2-25%)	86% (75-93%)	0.6 (0.22-1.3)	0.41% (0.051-1.1%)	0.42% (-0.0037-1.2%)	81 (-0.73-240)	320 (40-860)
India	2005	Male	All Ages	18% (17-20%)	82% (80-83%)	1.2 (0.94-1.4)	3.8% (3.2-4.4%)	4.5% (3.6-5.4%)	230000 (180000-280000)	10000000 (8700000-12000000)
India	2005	Male	15 to 19	7.5% (5.7-9.7%)	92% (90-94%)	0.34 (0.19-0.54)	1.5% (1.1-2%)	1.5% (0.91-2.2%)	1300 (810-2000)	160000 (110000-220000)
India	2005	Male	20 to 24	19% (15-24%)	81% (76-85%)	1.2 (0.76-1.8)	5% (3.8-6.6%)	5.3% (3.6-7.5%)	6700 (4500-9300)	630000 (460000-820000)
India	2005	Male	25 to 29	28% (23-33%)	72% (67-77%)	1.9 (1.3-2.6)	7.3% (5.7-9.1%)	8.1% (5.8-11%)	12000 (8400-15000)	960000 (730000-1200000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	2005	Male	30 to 34	33% (28-39%)	67% (61-72%)	2.3 (1.6-3.2)	8.7% (6.9-11%)	10% (7.7-13%)	17000 (13000-22000)	1200000 (940000-1500000)
India	2005	Male	35 to 39	36% (31-41%)	64% (59-69%)	2.5 (1.7-3.4)	9% (7.2-11%)	11% (8.3-13%)	21000 (17000-27000)	1300000 (1000000-1600000)
India	2005	Male	40 to 44	36% (30-41%)	64% (59-70%)	2.5 (1.7-3.5)	8.9% (7-11%)	11% (8.4-13%)	24000 (19000-30000)	1300000 (1000000-1600000)
India	2005	Male	45 to 49	34% (28-39%)	66% (61-72%)	2.4 (1.6-3.5)	8.3% (6.4-11%)	10% (7.6-13%)	27000 (20000-34000)	1200000 (940000-1600000)
India	2005	Male	50 to 54	30% (25-36%)	70% (64-75%)	2.1 (1.4-3.2)	7.3% (5.3-9.4%)	8.8% (6.2-11%)	28000 (20000-37000)	1100000 (820000-1400000)
India	2005	Male	55 to 59	27% (21-33%)	73% (67-79%)	1.9 (1.3-2.9)	6.2% (4.5-8.1%)	7.4% (5.3-9.8%)	24000 (17000-32000)	830000 (600000-1100000)
India	2005	Male	60 to 64	25% (19-31%)	75% (69-81%)	1.7 (1-2.6)	4.9% (3.3-6.7%)	5.6% (3.7-7.9%)	22000 (14000-30000)	630000 (430000-860000)
India	2005	Male	65 to 69	23% (17-29%)	77% (71-83%)	1.5 (0.92-2.2)	3.8% (2.5-5.3%)	4.3% (2.8-6.1%)	19000 (13000-28000)	460000 (310000-640000)
India	2005	Male	70 to 74	21% (16-26%)	79% (74-84%)	1.2 (0.74-1.9)	2.7% (1.7-4%)	3% (1.8-4.6%)	14000 (8100-21000)	260000 (160000-400000)
India	2005	Male	75 to 79	18% (13-24%)	82% (76-87%)	0.94 (0.53-1.5)	1.8% (1-2.9%)	2% (1.1-3.3%)	8000 (4300-13000)	120000 (68000-190000)
India	2005	Male	80 to 84	15% (10-21%)	85% (79-90%)	0.73 (0.42-1.1)	1.2% (0.68-2%)	1.4% (0.69-2.2%)	3800 (1900-6200)	43000 (23000-69000)
India	2005	Male	85 to 89	15% (10-21%)	85% (79-90%)	0.66 (0.37-1.1)	0.9% (0.43-1.5%)	0.94% (0.41-1.7%)	1500 (670-2700)	13000 (6200-22000)
India	2005	Male	90 to 94	17% (12-23%)	83% (77-88%)	0.65 (0.35-1.1)	0.64% (0.24-1.2%)	0.61% (0.14-1.3%)	380 (87-800)	2400 (860-4700)
India	2005	Male	95 plus	19% (12-27%)	81% (73-88%)	0.64 (0.29-1.1)	0.53% (0.12-1.2%)	0.41% (-0.051-1.2%)	62 (-7.6-180)	320 (73-710)
India	2010	Female	All Ages	3.5% (2.9-4%)	97% (96-97%)	0.11 (0.085-0.14)	0.47% (0.39-0.58%)	0.67% (0.48-0.89%)	29000 (21000-38000)	1100000 (880000-1400000)
India	2010	Female	15 to 19	1.9% (1.2-2.8%)	98% (97-99%)	0.038 (0.018-0.071)	0.21% (0.14-0.31%)	0.13% (0.054-0.27%)	110 (45-230)	23000 (15000-35000)
India	2010	Female	20 to 24	1.9% (1.2-2.8%)	98% (97-99%)	0.068 (0.037-0.12)	0.45% (0.31-0.62%)	0.29% (0.17-0.5%)	310 (170-520)	55000 (37000-80000)
India	2010	Female	25 to 29	2.3% (1.6-3.3%)	98% (97-98%)	0.11 (0.061-0.19)	0.62% (0.45-0.81%)	0.54% (0.33-0.84%)	560 (350-880)	72000 (52000-100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	2010	Female	30 to 34	2.7% (1.8-3.9%)	97% (96-98%)	0.15 (0.086-0.25)	0.76% (0.57-1%)	0.84% (0.58-1.2%)	830 (570-1200)	81000 (60000-110000)
India	2010	Female	35 to 39	3.1% (2.1-4.2%)	97% (96-98%)	0.18 (0.099-0.29)	0.92% (0.7-1.2%)	1.2% (0.9-1.6%)	1300 (930-1700)	96000 (73000-120000)
India	2010	Female	40 to 44	3.4% (2.4-4.7%)	97% (95-98%)	0.19 (0.1-0.31)	0.95% (0.73-1.2%)	1.4% (1-1.8%)	1600 (1200-2100)	100000 (77000-130000)
India	2010	Female	45 to 49	3.7% (2.4-5.1%)	96% (95-98%)	0.18 (0.099-0.3)	0.94% (0.73-1.2%)	1.3% (0.98-1.8%)	2000 (1400-2700)	100000 (80000-140000)
India	2010	Female	50 to 54	4.2% (2.7-6.2%)	96% (94-97%)	0.22 (0.11-0.39)	1% (0.75-1.4%)	1.4% (1-2%)	2800 (2000-3900)	120000 (88000-160000)
India	2010	Female	55 to 59	4.7% (3-7.1%)	95% (93-97%)	0.29 (0.14-0.51)	1% (0.7-1.5%)	1.4% (0.89-2.1%)	3500 (2300-5300)	130000 (86000-190000)
India	2010	Female	60 to 64	5.1% (3.2-8%)	95% (92-97%)	0.32 (0.16-0.59)	0.89% (0.56-1.4%)	1.1% (0.68-1.8%)	3600 (2200-5800)	110000 (67000-170000)
India	2010	Female	65 to 69	5.3% (3.3-8%)	95% (92-97%)	0.3 (0.15-0.51)	0.76% (0.48-1.1%)	0.95% (0.58-1.5%)	3700 (2200-5700)	88000 (56000-130000)
India	2010	Female	70 to 74	5.4% (3.5-8.1%)	95% (92-97%)	0.27 (0.13-0.49)	0.61% (0.36-1%)	0.74% (0.43-1.2%)	3400 (2000-5600)	64000 (37000-110000)
India	2010	Female	75 to 79	5.5% (3.2-8.9%)	95% (91-97%)	0.24 (0.11-0.47)	0.47% (0.25-0.83%)	0.56% (0.29-0.99%)	2400 (1200-4200)	35000 (18000-60000)
India	2010	Female	80 to 84	6% (3.3-10%)	94% (90-97%)	0.26 (0.11-0.52)	0.4% (0.19-0.78%)	0.47% (0.21-0.92%)	1700 (780-3400)	19000 (9100-36000)
India	2010	Female	85 to 89	6.6% (3.6-12%)	93% (88-96%)	0.29 (0.12-0.58)	0.35% (0.15-0.72%)	0.39% (0.16-0.8%)	920 (370-1900)	7300 (3200-15000)
India	2010	Female	90 to 94	7.6% (4.1-14%)	92% (86-96%)	0.34 (0.14-0.68)	0.32% (0.11-0.72%)	0.34% (0.097-0.83%)	330 (93-790)	1900 (640-4300)
India	2010	Female	95 plus	9.1% (4.2-18%)	91% (82-96%)	0.41 (0.15-0.93)	0.34% (0.071-0.84%)	0.34% (0.03-0.94%)	86 (7.6-240)	340 (73-860)
India	2010	Male	All Ages	21% (20-22%)	79% (78-80%)	1.3 (1.1-1.7)	4.1% (3.5-4.9%)	4.8% (3.8-5.9%)	250000 (200000-310000)	11000000 (9200000-13000000)
India	2010	Male	15 to 19	9.7% (7.7-12%)	90% (88-92%)	0.45 (0.28-0.69)	1.6% (1.2-2.2%)	1.8% (1.1-2.7%)	1500 (920-2200)	160000 (120000-220000)
India	2010	Male	20 to 24	22% (19-25%)	78% (75-81%)	1.4 (0.94-2.1)	5.3% (3.9-6.9%)	6% (4.1-8.3%)	7200 (4900-10000)	650000 (470000-850000)
India	2010	Male	25 to 29	31% (27-35%)	69% (65-73%)	2.2 (1.5-3)	7.6% (5.9-9.4%)	8.9% (6.4-11%)	13000 (9300-17000)	1000000 (780000-1300000)
India	2010	Male	30 to 34	34% (30-39%)	66% (61-70%)	2.5 (1.7-3.4)	8.6% (7-11%)	10% (8-13%)	17000 (13000-22000)	1200000 (950000-1500000)
India	2010	Male	35 to 39	37% (33-40%)	63% (60-67%)	2.7 (1.8-3.6)	8.9% (7.1-11%)	11% (8.3-13%)	22000 (17000-27000)	1300000 (1000000-1600000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	2010	Male	40 to 44	36% (32-40%)	64% (60-68%)	2.6 (1.8-3.7)	8.6% (6.7-11%)	11% (8.1-13%)	24000 (19000-30000)	1300000 (1000000-1600000)
India	2010	Male	45 to 49	34% (30-39%)	66% (61-70%)	2.6 (1.7-3.7)	8.2% (6.2-10%)	10% (7.5-13%)	28000 (21000-35000)	1300000 (980000-1600000)
India	2010	Male	50 to 54	32% (27-37%)	68% (63-73%)	2.4 (1.5-3.5)	7.2% (5.2-9.3%)	8.7% (6.3-11%)	29000 (22000-39000)	1200000 (870000-1500000)
India	2010	Male	55 to 59	29% (24-35%)	71% (65-76%)	2.2 (1.4-3.3)	6.3% (4.4-8.1%)	7.5% (5.3-9.9%)	30000 (21000-40000)	1000000 (750000-1400000)
India	2010	Male	60 to 64	28% (22-35%)	72% (65-78%)	2 (1.3-3.1)	5% (3.4-7%)	5.9% (3.9-8.2%)	25000 (16000-34000)	720000 (480000-980000)
India	2010	Male	65 to 69	27% (21-34%)	73% (66-79%)	1.8 (1.2-2.8)	3.9% (2.6-5.6%)	4.5% (2.9-6.5%)	21000 (14000-30000)	510000 (340000-710000)
India	2010	Male	70 to 74	26% (20-32%)	74% (68-80%)	1.6 (0.97-2.4)	2.9% (1.7-4.3%)	3.3% (1.9-5.1%)	16000 (9400-25000)	320000 (190000-480000)
India	2010	Male	75 to 79	23% (17-30%)	77% (70-83%)	1.3 (0.74-2)	2% (1.1-3.3%)	2.2% (1.1-3.7%)	10000 (4900-17000)	150000 (78000-240000)
India	2010	Male	80 to 84	20% (13-27%)	80% (73-87%)	0.99 (0.66-1.5)	1.4% (0.66-2.2%)	1.5% (0.67-2.5%)	5200 (2300-8900)	59000 (28000-99000)
India	2010	Male	85 to 89	19% (12-28%)	81% (72-88%)	0.85 (0.46-1.4)	0.94% (0.4-1.7%)	0.97% (0.34-1.8%)	2000 (700-3900)	17000 (7200-31000)
India	2010	Male	90 to 94	21% (14-30%)	79% (70-86%)	0.84 (0.43-1.4)	0.64% (0.14-1.4%)	0.59% (0.0034-1.4%)	470 (2.7-1100)	3100 (680-6500)
India	2010	Male	95 plus	25% (15-37%)	75% (63-85%)	0.86 (0.39-1.6)	0.49% (-0.051-1.3%)	0.35% (-0.29-1.3%)	67 (-55-250)	380 (-39-1000)
India	2016	Female	All Ages	4.1% (3.3-5.1%)	96% (95-97%)	0.18 (0.14-0.25)	0.71% (0.57-0.89%)	0.95% (0.68-1.3%)	42000 (30000-58000)	1500000 (1200000-1900000)
India	2016	Female	15 to 19	2.6% (1.5-4.4%)	97% (96-99%)	0.066 (0.027-0.13)	0.27% (0.17-0.44%)	0.22% (0.081-0.52%)	160 (58-370)	28000 (17000-46000)
India	2016	Female	20 to 24	2.7% (1.5-4.4%)	97% (96-98%)	0.12 (0.056-0.24)	0.61% (0.41-0.89%)	0.52% (0.25-1%)	450 (220-880)	69000 (45000-100000)
India	2016	Female	25 to 29	3.2% (1.9-5.2%)	97% (95-98%)	0.2 (0.093-0.39)	0.86% (0.59-1.3%)	0.92% (0.49-1.6%)	820 (440-1500)	96000 (63000-140000)
India	2016	Female	30 to 34	3.6% (2-6.2%)	96% (94-98%)	0.27 (0.13-0.49)	1% (0.72-1.5%)	1.4% (0.81-2.2%)	1200 (740-2000)	110000 (79000-170000)
India	2016	Female	35 to 39	4.1% (2.3-6.3%)	96% (94-98%)	0.31 (0.15-0.54)	1.2% (0.88-1.7%)	1.9% (1.2-2.7%)	1900 (1200-2700)	140000 (95000-180000)
India	2016	Female	40 to 44	4.4% (2.7-7%)	96% (93-97%)	0.32 (0.15-0.55)	1.2% (0.89-1.7%)	2% (1.3-2.8%)	2300 (1500-3200)	140000 (100000-190000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	2016	Female	45 to 49	4.7% (2.7-7.2%)	95% (93-97%)	0.29 (0.14-0.54)	1.2% (0.86-1.7%)	1.8% (1.3-2.6%)	2700 (1900-3900)	140000 (100000-190000)
India	2016	Female	50 to 54	4.9% (3-7.7%)	95% (92-97%)	0.31 (0.15-0.57)	1.3% (0.88-1.8%)	1.8% (1.2-2.7%)	3900 (2600-5800)	160000 (120000-230000)
India	2016	Female	55 to 59	5.3% (3.2-8.9%)	95% (91-97%)	0.39 (0.18-0.74)	1.3% (0.83-1.9%)	1.8% (1.1-2.8%)	4800 (3000-7500)	170000 (110000-260000)
India	2016	Female	60 to 64	5.7% (3.3-9.8%)	94% (90-97%)	0.42 (0.2-0.81)	1.1% (0.67-1.8%)	1.5% (0.86-2.4%)	5500 (3200-9000)	160000 (98000-260000)
India	2016	Female	65 to 69	5.8% (3.4-9.3%)	94% (91-97%)	0.38 (0.18-0.71)	0.95% (0.57-1.5%)	1.2% (0.71-1.9%)	5300 (3100-8400)	130000 (75000-200000)
India	2016	Female	70 to 74	5.8% (3.6-9.2%)	94% (91-96%)	0.33 (0.15-0.65)	0.75% (0.43-1.3%)	0.94% (0.51-1.6%)	4400 (2400-7500)	84000 (47000-140000)
India	2016	Female	75 to 79	5.7% (3.3-9.5%)	94% (91-97%)	0.3 (0.12-0.6)	0.58% (0.3-1%)	0.71% (0.36-1.3%)	3300 (1700-6200)	49000 (26000-90000)
India	2016	Female	80 to 84	6% (3.1-10%)	94% (90-97%)	0.31 (0.13-0.64)	0.5% (0.24-0.98%)	0.6% (0.28-1.2%)	2700 (1200-5200)	29000 (14000-57000)
India	2016	Female	85 to 89	6.6% (3.4-12%)	93% (88-97%)	0.34 (0.13-0.7)	0.44% (0.19-0.9%)	0.5% (0.2-1.1%)	1500 (630-3200)	12000 (5300-25000)
India	2016	Female	90 to 94	7.7% (3.9-14%)	92% (86-96%)	0.4 (0.15-0.83)	0.41% (0.15-0.97%)	0.45% (0.14-1.1%)	600 (190-1500)	3400 (1200-7800)
India	2016	Female	95 plus	9.3% (4-18%)	91% (82-96%)	0.48 (0.16-1.1)	0.43% (0.087-1.1%)	0.45% (0.046-1.2%)	170 (17-440)	650 (130-1600)
India	2016	Male	All Ages	20% (18-22%)	80% (78-82%)	1.5 (1.2-1.8)	4.7% (4-5.5%)	5.4% (4.3-6.5%)	290000 (230000-350000)	12000000 (10000000-14000000)
India	2016	Male	15 to 19	6.6% (4.5-9.6%)	93% (90-95%)	0.36 (0.19-0.6)	1.4% (0.97-1.9%)	1.5% (0.93-2.4%)	1100 (670-1800)	130000 (93000-190000)
India	2016	Male	20 to 24	15% (11-20%)	85% (80-89%)	1.1 (0.7-1.8)	4.4% (3.2-6%)	5.1% (3.3-7.3%)	5600 (3700-8100)	520000 (370000-700000)
India	2016	Male	25 to 29	25% (20-31%)	75% (69-80%)	2 (1.3-2.9)	6.9% (5.2-8.7%)	8.4% (5.9-11%)	11000 (8000-15000)	900000 (680000-1100000)
India	2016	Male	30 to 34	30% (24-38%)	70% (62-76%)	2.5 (1.6-3.5)	8.3% (6.6-10%)	11% (7.9-13%)	17000 (13000-22000)	1200000 (930000-1500000)
India	2016	Male	35 to 39	34% (27-41%)	66% (59-73%)	2.8 (1.8-4)	8.9% (7.1-11%)	11% (8.7-14%)	23000 (17000-28000)	1400000 (1100000-1700000)
India	2016	Male	40 to 44	34% (28-41%)	66% (59-72%)	2.9 (1.9-4.2)	8.8% (6.9-11%)	11% (8.7-14%)	26000 (20000-32000)	1400000 (1100000-1700000)
India	2016	Male	45 to 49	33% (26-40%)	67% (60-74%)	2.8 (1.8-4.2)	8.5% (6.5-11%)	11% (8.1-13%)	30000 (22000-37000)	1400000 (1100000-1700000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
India	2016	Male	50 to 54	32% (25-39%)	68% (61-75%)	2.7 (1.7-4)	7.7% (5.7-9.7%)	9.5% (7-12%)	33000 (24000-42000)	1300000 (1000000-1700000)
India	2016	Male	55 to 59	30% (24-37%)	70% (63-76%)	2.6 (1.7-3.8)	6.8% (5-8.8%)	8.3% (6-11%)	37000 (26000-47000)	1300000 (920000-1600000)
India	2016	Male	60 to 64	29% (23-37%)	71% (63-77%)	2.4 (1.5-3.6)	5.5% (3.8-7.5%)	6.5% (4.4-9.1%)	34000 (23000-47000)	1000000 (670000-1400000)
India	2016	Male	65 to 69	28% (22-35%)	72% (65-78%)	2.2 (1.4-3.2)	4.4% (2.9-6.1%)	5.2% (3.4-7.2%)	27000 (18000-38000)	660000 (430000-910000)
India	2016	Male	70 to 74	26% (21-33%)	74% (67-79%)	1.8 (1.1-2.8)	3.3% (2-4.9%)	3.8% (2.2-5.7%)	20000 (12000-30000)	390000 (230000-570000)
India	2016	Male	75 to 79	24% (18-31%)	76% (69-82%)	1.5 (0.86-2.3)	2.3% (1.2-3.7%)	2.6% (1.3-4.3%)	13000 (6700-21000)	200000 (110000-320000)
India	2016	Male	80 to 84	21% (14-28%)	79% (72-86%)	1.2 (0.68-1.8)	1.6% (0.8-2.6%)	1.8% (0.82-2.9%)	7900 (3600-13000)	89000 (43000-140000)
India	2016	Male	85 to 89	20% (13-28%)	80% (72-87%)	1 (0.56-1.7)	1.1% (0.52-2%)	1.2% (0.49-2.1%)	3300 (1300-5800)	27000 (12000-46000)
India	2016	Male	90 to 94	22% (15-31%)	78% (69-85%)	0.99 (0.53-1.7)	0.79% (0.2-1.7%)	0.76% (0.08-1.8%)	830 (86-1900)	5200 (1300-11000)
India	2016	Male	95 plus	26% (16-37%)	74% (63-84%)	1 (0.46-1.8)	0.6% (-0.029-1.6%)	0.47% (-0.26-1.6%)	130 (-70-420)	640 (-30-1700)
Nepal	1990	Female	All Ages	0.71% (0.53-0.95%)	99% (99-99%)	0.013 (0.0095-0.016)	0.22% (0.17-0.28%)	0.25% (0.19-0.31%)	270 (210-340)	16000 (13000-19000)
Nepal	1990	Female	15 to 19	0.52% (0.24-0.91%)	99% (99-100%)	0.0062 (0.0023-0.013)	0.29% (0.18-0.44%)	0.036% (0.016-0.068%)	0.71 (0.32-1.4)	660 (380-1000)
Nepal	1990	Female	20 to 24	0.61% (0.24-1.1%)	99% (99-100%)	0.014 (0.005-0.026)	0.62% (0.41-0.89%)	0.12% (0.066-0.21%)	2.8 (1.5-4.8)	1400 (880-2200)
Nepal	1990	Female	25 to 29	0.7% (0.3-1.3%)	99% (99-100%)	0.019 (0.0082-0.036)	0.72% (0.51-0.96%)	0.24% (0.14-0.37%)	5.1 (3-8.1)	1500 (980-2000)
Nepal	1990	Female	30 to 34	0.84% (0.38-1.6%)	99% (98-100%)	0.026 (0.011-0.047)	0.8% (0.59-1.1%)	0.48% (0.3-0.72%)	11 (6.8-16)	1600 (1100-2200)
Nepal	1990	Female	35 to 39	1% (0.48-1.7%)	99% (98-100%)	0.032 (0.014-0.053)	0.88% (0.67-1.1%)	0.74% (0.48-1.1%)	18 (12-27)	1700 (1300-2300)
Nepal	1990	Female	40 to 44	1.1% (0.58-1.8%)	99% (98-99%)	0.033 (0.015-0.058)	0.93% (0.69-1.2%)	0.94% (0.62-1.3%)	22 (14-32)	1500 (1100-2000)
Nepal	1990	Female	45 to 49	1.2% (0.53-1.9%)	99% (98-99%)	0.03 (0.013-0.052)	0.91% (0.68-1.2%)	0.97% (0.65-1.4%)	28 (18-40)	1600 (1200-2100)
Nepal	1990	Female	50 to 54	1.3% (0.61-2.2%)	99% (98-99%)	0.029 (0.012-0.054)	0.85% (0.6-1.2%)	0.91% (0.58-1.4%)	34 (21-51)	1600 (1100-2200)
Nepal	1990	Female	55 to 59	1.4% (0.67-2.3%)	99% (98-99%)	0.03 (0.013-0.056)	0.69% (0.48-0.95%)	0.73% (0.47-1.1%)	34 (21-51)	1300 (910-1900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	1990	Female	60 to 64	1.4% (0.69-2.5%)	99% (98-99%)	0.029 (0.012-0.056)	0.52% (0.34-0.74%)	0.54% (0.32-0.83%)	32 (19-49)	1100 (680-1500)
Nepal	1990	Female	65 to 69	1.4% (0.7-2.3%)	99% (98-99%)	0.025 (0.011-0.045)	0.42% (0.28-0.6%)	0.45% (0.27-0.67%)	32 (19-48)	820 (530-1200)
Nepal	1990	Female	70 to 74	1.4% (0.74-2.2%)	99% (98-99%)	0.021 (0.009-0.038)	0.32% (0.19-0.51%)	0.35% (0.19-0.58%)	25 (14-42)	500 (300-800)
Nepal	1990	Female	75 to 79	1.4% (0.7-2.3%)	99% (98-99%)	0.018 (0.0076-0.036)	0.25% (0.15-0.4%)	0.27% (0.15-0.45%)	17 (9-28)	260 (150-410)
Nepal	1990	Female	80 to 84	1.4% (0.72-2.3%)	99% (98-99%)	0.017 (0.0071-0.032)	0.17% (0.091-0.26%)	0.17% (0.084-0.29%)	6.9 (3.3-12)	80 (43-130)
Nepal	1990	Female	85 to 89	1.6% (0.77-2.7%)	98% (97-99%)	0.016 (0.007-0.032)	0.12% (0.067-0.18%)	0.11% (0.053-0.19%)	2.3 (1.1-4)	21 (12-33)
Nepal	1990	Female	90 to 94	1.8% (0.92-3.1%)	98% (97-99%)	0.017 (0.0071-0.032)	0.086% (0.043-0.14%)	0.07% (0.019-0.13%)	0.48 (0.13-0.91)	3.5 (1.7-5.7)
Nepal	1990	Female	95 plus	2.2% (0.97-4.2%)	98% (96-99%)	0.018 (0.0058-0.037)	0.078% (0.026-0.13%)	0.043% (-0.013-0.099%)	0.074 (-0.022-0.17)	0.52 (0.17-0.9)
Nepal	1990	Male	All Ages	18% (15-22%)	82% (78-85%)	0.13 (0.11-0.16)	0.53% (0.28-0.79%)	0.56% (0.11-0.98%)	560 (110-1000)	37000 (20000-55000)
Nepal	1990	Male	15 to 19	8.7% (5.2-14%)	91% (86-95%)	0.044 (0.022-0.081)	0.67% (0.44-0.96%)	0.33% (0.15-0.59%)	5.7 (2.6-10)	1300 (840-2000)
Nepal	1990	Male	20 to 24	23% (14-32%)	77% (68-86%)	0.15 (0.084-0.25)	1.8% (1.1-2.5%)	1.2% (0.44-2.1%)	25 (9.3-46)	3700 (2300-5400)
Nepal	1990	Male	25 to 29	35% (24-47%)	65% (53-76%)	0.26 (0.16-0.38)	2.5% (1.5-3.6%)	1.8% (0.36-3.5%)	38 (7.3-71)	4600 (2600-6800)
Nepal	1990	Male	30 to 34	43% (30-56%)	57% (44-70%)	0.33 (0.21-0.47)	3% (1.5-4.4%)	2.6% (0.56-4.7%)	59 (13-100)	5500 (2800-8200)
Nepal	1990	Male	35 to 39	45% (32-57%)	55% (43-68%)	0.35 (0.22-0.52)	2.9% (1.4-4.3%)	2.8% (0.74-5%)	72 (19-130)	5400 (2700-8300)
Nepal	1990	Male	40 to 44	45% (32-56%)	55% (44-68%)	0.35 (0.21-0.53)	2.7% (1.2-4.2%)	2.7% (0.66-4.9%)	67 (16-120)	4200 (1800-6700)
Nepal	1990	Male	45 to 49	43% (29-55%)	57% (45-71%)	0.34 (0.19-0.53)	2.4% (1-3.7%)	2.5% (0.68-4.3%)	75 (20-130)	4000 (1800-6500)
Nepal	1990	Male	50 to 54	39% (28-51%)	61% (49-72%)	0.32 (0.18-0.51)	1.9% (0.65-3.1%)	2% (0.37-3.7%)	71 (13-130)	3300 (1100-5500)
Nepal	1990	Male	55 to 59	34% (24-45%)	66% (55-76%)	0.29 (0.16-0.47)	1.4% (0.4-2.4%)	1.4% (0.096-2.7%)	57 (3.6-110)	2300 (600-4000)
Nepal	1990	Male	60 to 64	31% (20-42%)	69% (58-80%)	0.26 (0.14-0.43)	0.91% (0.063-1.7%)	0.86% (-0.19-1.9%)	42 (-9.7-93)	1500 (110-2800)
Nepal	1990	Male	65 to 69	28% (18-39%)	72% (61-82%)	0.22 (0.12-0.36)	0.6% (-0.052-1.2%)	0.55% (-0.26-1.3%)	29 (-13-70)	850 (-72-1800)
Nepal	1990	Male	70 to 74	26% (17-36%)	74% (64-83%)	0.18 (0.095-0.3)	0.35% (-0.2-0.86%)	0.31% (-0.37-0.89%)	15 (-19-45)	380 (-220-920)
Nepal	1990	Male	75 to 79	23% (14-34%)	77% (66-86%)	0.14 (0.067-0.26)	0.22% (-0.26-0.59%)	0.18% (-0.4-0.62%)	7.2 (-16-25)	140 (-170-390)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	1990	Male	80 to 84	21% (13-32%)	79% (68-87%)	0.12 (0.056-0.2)	0.12% (-0.22-0.4%)	0.074% (-0.32-0.4%)	1.7 (-7.6-9.8)	33 (-64-120)
Nepal	1990	Male	85 to 89	21% (13-32%)	79% (68-87%)	0.1 (0.052-0.18)	0.06% (-0.21-0.29%)	0.009% (-0.31-0.27%)	0.12 (-4-3.6)	6.9 (-24-33)
Nepal	1990	Male	90 to 94	24% (14-36%)	76% (64-86%)	0.1 (0.048-0.19)	0.016% (-0.24-0.22%)	-0.057% (-0.35-0.18%)	-0.27 (-1.6-0.87)	0.46 (-6.6-6.2)
Nepal	1990	Male	95 plus	28% (15-45%)	72% (55-85%)	0.11 (0.04-0.22)	0.027% (-0.26-0.22%)	-0.081% (-0.4-0.14%)	-0.086 (-0.43-0.15)	0.11 (-1.1-0.93)
Nepal	1995	Female	All Ages	0.71% (0.54-0.94%)	99% (99-99%)	0.025 (0.019-0.031)	0.31% (0.25-0.38%)	0.37% (0.28-0.47%)	390 (290-500)	20000 (16000-25000)
Nepal	1995	Female	15 to 19	0.51% (0.24-0.91%)	99% (99-100%)	0.012 (0.0045-0.025)	0.33% (0.2-0.49%)	0.063% (0.029-0.14%)	1.4 (0.63-3)	810 (470-1300)
Nepal	1995	Female	20 to 24	0.6% (0.23-1.1%)	99% (99-100%)	0.026 (0.0098-0.048)	0.72% (0.47-1%)	0.2% (0.11-0.33%)	4.6 (2.5-7.8)	1800 (1100-2700)
Nepal	1995	Female	25 to 29	0.69% (0.3-1.3%)	99% (99-100%)	0.037 (0.016-0.068)	0.85% (0.61-1.1%)	0.36% (0.21-0.59%)	7.8 (4.4-13)	1800 (1200-2500)
Nepal	1995	Female	30 to 34	0.84% (0.37-1.6%)	99% (98-100%)	0.05 (0.022-0.091)	0.97% (0.7-1.3%)	0.67% (0.42-1%)	14 (8.7-22)	1900 (1300-2600)
Nepal	1995	Female	35 to 39	1% (0.49-1.7%)	99% (98-100%)	0.062 (0.028-0.11)	1.1% (0.79-1.4%)	0.99% (0.65-1.4%)	23 (15-34)	2100 (1500-2800)
Nepal	1995	Female	40 to 44	1.1% (0.58-1.9%)	99% (98-99%)	0.065 (0.03-0.12)	1.2% (0.84-1.5%)	1.3% (0.83-1.8%)	33 (22-48)	2300 (1600-3100)
Nepal	1995	Female	45 to 49	1.2% (0.54-1.9%)	99% (98-99%)	0.059 (0.026-0.11)	1.1% (0.83-1.5%)	1.3% (0.86-1.8%)	35 (23-50)	1900 (1400-2600)
Nepal	1995	Female	50 to 54	1.3% (0.64-2.1%)	99% (98-99%)	0.057 (0.024-0.1)	1.1% (0.76-1.5%)	1.2% (0.8-1.8%)	46 (29-69)	2100 (1400-2900)
Nepal	1995	Female	55 to 59	1.4% (0.68-2.3%)	99% (98-99%)	0.06 (0.027-0.11)	0.91% (0.64-1.3%)	1% (0.65-1.5%)	49 (31-75)	1900 (1300-2700)
Nepal	1995	Female	60 to 64	1.4% (0.69-2.4%)	99% (98-99%)	0.058 (0.025-0.11)	0.71% (0.46-1%)	0.78% (0.46-1.2%)	46 (28-72)	1500 (930-2100)
Nepal	1995	Female	65 to 69	1.4% (0.71-2.3%)	99% (98-99%)	0.05 (0.023-0.09)	0.56% (0.38-0.83%)	0.62% (0.38-0.96%)	47 (29-72)	1200 (770-1700)
Nepal	1995	Female	70 to 74	1.4% (0.76-2.3%)	99% (98-99%)	0.041 (0.019-0.076)	0.43% (0.26-0.64%)	0.47% (0.27-0.75%)	39 (22-62)	760 (450-1200)
Nepal	1995	Female	75 to 79	1.4% (0.71-2.4%)	99% (98-99%)	0.036 (0.014-0.069)	0.32% (0.19-0.51%)	0.36% (0.2-0.58%)	25 (14-41)	380 (220-590)
Nepal	1995	Female	80 to 84	1.5% (0.71-2.4%)	99% (98-99%)	0.033 (0.013-0.063)	0.22% (0.13-0.35%)	0.23% (0.12-0.39%)	11 (5.7-18)	130 (72-200)
Nepal	1995	Female	85 to 89	1.6% (0.79-2.7%)	98% (97-99%)	0.033 (0.015-0.062)	0.15% (0.088-0.23%)	0.15% (0.076-0.25%)	4 (2-6.6)	35 (20-53)
Nepal	1995	Female	90 to 94	1.8% (0.94-3.1%)	98% (97-99%)	0.033 (0.014-0.064)	0.11% (0.061-0.18%)	0.097% (0.038-0.18%)	0.88 (0.35-1.6)	6 (3.2-9.8)
Nepal	1995	Female	95 plus	2.2% (0.95-4%)	98% (96-99%)	0.035 (0.012-0.075)	0.097% (0.038-0.18%)	0.061% (-0.003-0.15%)	0.13 (-0.0065-0.32)	0.83 (0.32-1.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	1995	Male	All Ages	19% (15-23%)	81% (77-85%)	0.26 (0.22-0.3)	0.96% (0.61-1.3%)	1.1% (0.54-1.7%)	1100 (520-1600)	62000 (40000-83000)
Nepal	1995	Male	15 to 19	8.7% (5.1-14%)	91% (86-95%)	0.085 (0.042-0.15)	0.83% (0.54-1.2%)	0.54% (0.24-0.96%)	10 (4.6-18)	1800 (1200-2700)
Nepal	1995	Male	20 to 24	23% (14-33%)	77% (67-86%)	0.3 (0.17-0.49)	2.4% (1.6-3.4%)	2% (1-3.3%)	49 (25-82)	5900 (3900-8400)
Nepal	1995	Male	25 to 29	35% (24-47%)	65% (53-76%)	0.5 (0.31-0.74)	3.5% (2.3-4.8%)	3.3% (1.5-5.2%)	78 (36-120)	7700 (4900-11000)
Nepal	1995	Male	30 to 34	42% (30-56%)	58% (44-70%)	0.64 (0.41-0.9)	4.3% (2.9-6%)	4.5% (2.5-6.8%)	110 (59-160)	8500 (5500-12000)
Nepal	1995	Male	35 to 39	45% (31-57%)	55% (43-69%)	0.67 (0.42-0.99)	4.2% (2.6-5.9%)	4.6% (2.4-7.1%)	120 (61-190)	8200 (5000-12000)
Nepal	1995	Male	40 to 44	45% (32-57%)	55% (43-68%)	0.67 (0.4-1)	4% (2.3-5.9%)	4.5% (2.2-7.2%)	140 (65-220)	7900 (4500-12000)
Nepal	1995	Male	45 to 49	43% (29-56%)	57% (44-71%)	0.67 (0.37-1.1)	3.6% (1.9-5.5%)	4.2% (1.8-6.6%)	130 (55-210)	6300 (3200-9700)
Nepal	1995	Male	50 to 54	39% (28-51%)	61% (49-72%)	0.62 (0.35-1)	3% (1.4-4.8%)	3.4% (1.4-5.8%)	130 (53-230)	5700 (2600-9100)
Nepal	1995	Male	55 to 59	34% (24-46%)	66% (54-76%)	0.57 (0.32-0.91)	2.4% (1.1-4%)	2.6% (0.99-4.7%)	120 (44-210)	4400 (1900-7400)
Nepal	1995	Male	60 to 64	31% (20-43%)	69% (57-80%)	0.5 (0.28-0.87)	1.6% (0.56-3%)	1.7% (0.36-3.5%)	91 (19-180)	2900 (1000-5400)
Nepal	1995	Male	65 to 69	28% (19-39%)	72% (61-81%)	0.42 (0.23-0.68)	1.1% (0.23-2.1%)	1.1% (0.087-2.3%)	65 (5.1-140)	1700 (380-3300)
Nepal	1995	Male	70 to 74	26% (17-36%)	74% (64-83%)	0.35 (0.18-0.57)	0.61% (-0.039-1.4%)	0.61% (-0.15-1.5%)	35 (-8.8-88)	780 (-50-1800)
Nepal	1995	Male	75 to 79	23% (14-35%)	77% (65-86%)	0.28 (0.14-0.5)	0.34% (-0.2-0.9%)	0.31% (-0.32-0.95%)	14 (-15-45)	260 (-160-690)
Nepal	1995	Male	80 to 84	22% (13-32%)	78% (68-87%)	0.23 (0.11-0.41)	0.16% (-0.22-0.51%)	0.11% (-0.33-0.53%)	3.1 (-9.5-15)	54 (-78-180)
Nepal	1995	Male	85 to 89	22% (13-33%)	78% (67-87%)	0.2 (0.097-0.36)	0.05% (-0.3-0.34%)	-0.019% (-0.43-0.31%)	-0.3 (-6.8-5)	7 (-42-46)
Nepal	1995	Male	90 to 94	24% (14-36%)	76% (64-86%)	0.2 (0.09-0.37)	-0.03% (-0.37-0.24%)	-0.13% (-0.56-0.18%)	-0.79 (-3.4-1.1)	-1.1 (-13-8.6)
Nepal	1995	Male	95 plus	28% (15-45%)	72% (55-85%)	0.2 (0.075-0.41)	-0.026% (-0.4-0.25%)	-0.17% (-0.6-0.13%)	-0.23 (-0.83-0.18)	-0.14 (-2.2-1.4)
Nepal	2000	Female	All Ages	1.4% (1.2-1.7%)	99% (98-99%)	0.083 (0.066-0.1)	0.53% (0.42-0.67%)	0.69% (0.49-0.92%)	650 (460-880)	31000 (24000-39000)
Nepal	2000	Female	15 to 19	0.89% (0.47-1.5%)	99% (99-100%)	0.03 (0.013-0.056)	0.41% (0.26-0.6%)	0.14% (0.062-0.31%)	2.9 (1.2-6)	1000 (640-1600)
Nepal	2000	Female	20 to 24	1.4% (0.68-2.1%)	99% (98-99%)	0.083 (0.039-0.14)	0.96% (0.65-1.4%)	0.47% (0.24-0.84%)	10 (5.4-19)	2400 (1600-3700)
Nepal	2000	Female	25 to 29	1.7% (0.94-2.7%)	98% (97-99%)	0.13 (0.069-0.22)	1.2% (0.88-1.7%)	0.86% (0.47-1.5%)	17 (9.5-29)	2700 (1800-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	2000	Female	30 to 34	2.2% (1.2-3.6%)	98% (96-99%)	0.18 (0.096-0.3)	1.4% (1-2%)	1.4% (0.86-2.3%)	28 (17-45)	2900 (2000-4100)
Nepal	2000	Female	35 to 39	2.6% (1.5-3.9%)	97% (96-98%)	0.22 (0.12-0.35)	1.7% (1.2-2.3%)	2% (1.2-3%)	40 (25-61)	3100 (2200-4300)
Nepal	2000	Female	40 to 44	2.8% (1.7-4.2%)	97% (96-98%)	0.23 (0.12-0.37)	1.8% (1.3-2.4%)	2.3% (1.5-3.5%)	52 (33-77)	3200 (2200-4400)
Nepal	2000	Female	45 to 49	2.9% (1.6-4.4%)	97% (96-98%)	0.2 (0.1-0.34)	1.7% (1.2-2.4%)	2.3% (1.5-3.3%)	64 (42-93)	3300 (2300-4500)
Nepal	2000	Female	50 to 54	3% (1.9-4.6%)	97% (95-98%)	0.19 (0.098-0.32)	1.7% (1.1-2.4%)	2.1% (1.3-3.2%)	68 (41-100)	2900 (2000-4200)
Nepal	2000	Female	55 to 59	3.1% (1.8-5%)	97% (95-98%)	0.19 (0.099-0.33)	1.5% (0.99-2.2%)	1.9% (1.1-2.8%)	83 (50-130)	3000 (1900-4500)
Nepal	2000	Female	60 to 64	3.1% (1.8-5%)	97% (95-98%)	0.18 (0.087-0.32)	1.2% (0.71-1.9%)	1.4% (0.79-2.4%)	83 (45-140)	2500 (1500-4000)
Nepal	2000	Female	65 to 69	2.9% (1.6-4.4%)	97% (96-98%)	0.14 (0.072-0.24)	0.91% (0.56-1.4%)	1.1% (0.64-1.6%)	76 (44-120)	1800 (1100-2800)
Nepal	2000	Female	70 to 74	2.6% (1.5-4%)	97% (96-98%)	0.11 (0.055-0.19)	0.65% (0.39-1.1%)	0.76% (0.43-1.3%)	62 (35-100)	1200 (710-1900)
Nepal	2000	Female	75 to 79	2.2% (1.2-3.6%)	98% (96-99%)	0.079 (0.035-0.15)	0.45% (0.26-0.74%)	0.52% (0.28-0.85%)	39 (21-66)	580 (330-950)
Nepal	2000	Female	80 to 84	1.9% (0.95-3%)	98% (97-99%)	0.061 (0.027-0.11)	0.31% (0.17-0.49%)	0.34% (0.18-0.56%)	17 (9.2-29)	200 (110-310)
Nepal	2000	Female	85 to 89	1.9% (0.92-3.2%)	98% (97-99%)	0.054 (0.023-0.1)	0.22% (0.12-0.35%)	0.23% (0.12-0.38%)	7 (3.6-12)	59 (33-94)
Nepal	2000	Female	90 to 94	2.1% (1.1-3.6%)	98% (96-99%)	0.055 (0.024-0.1)	0.16% (0.081-0.27%)	0.15% (0.059-0.29%)	1.7 (0.69-3.3)	11 (5.6-19)
Nepal	2000	Female	95 plus	2.6% (1.2-4.9%)	97% (95-99%)	0.058 (0.021-0.12)	0.14% (0.048-0.28%)	0.099% (-0.000078-0.26%)	0.29 (-0.00021-0.75)	1.6 (0.54-3.2)
Nepal	2000	Male	All Ages	24% (20-29%)	76% (71-80%)	0.47 (0.4-0.54)	1.6% (1.2-2.1%)	2% (1.2-3%)	1900 (1100-2700)	96000 (69000-130000)
Nepal	2000	Male	15 to 19	14% (8.8-20%)	86% (80-91%)	0.19 (0.1-0.32)	1.2% (0.75-1.7%)	1.1% (0.5-1.8%)	20 (9.5-35)	2700 (1700-4000)
Nepal	2000	Male	20 to 24	32% (22-41%)	68% (59-78%)	0.61 (0.35-0.96)	3.5% (2.3-5.1%)	3.6% (1.9-5.8%)	84 (46-140)	8400 (5400-12000)
Nepal	2000	Male	25 to 29	44% (34-53%)	56% (47-66%)	0.93 (0.6-1.3)	5.1% (3.4-7%)	5.6% (3.1-8.5%)	130 (71-200)	11000 (7100-15000)
Nepal	2000	Male	30 to 34	50% (40-60%)	50% (40-60%)	1.1 (0.75-1.5)	6.1% (4.2-8.2%)	7.1% (4.3-10%)	170 (100-250)	13000 (8500-17000)
Nepal	2000	Male	35 to 39	52% (41-61%)	48% (39-59%)	1.1 (0.77-1.6)	6% (4-8.3%)	7.2% (4.3-11%)	190 (110-290)	12000 (8000-17000)
Nepal	2000	Male	40 to 44	52% (42-61%)	48% (39-58%)	1.2 (0.73-1.7)	5.7% (3.6-8%)	7.1% (4.1-10%)	210 (120-320)	11000 (7100-17000)
Nepal	2000	Male	45 to 49	51% (40-60%)	49% (40-60%)	1.2 (0.72-1.7)	5.2% (3.1-7.6%)	6.4% (3.6-9.7%)	240 (130-370)	11000 (6600-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	2000	Male	50 to 54	46% (36-55%)	54% (45-64%)	1.1 (0.65-1.6)	4.5% (2.4-6.8%)	5.3% (2.5-8.4%)	210 (100-330)	8600 (4500-13000)
Nepal	2000	Male	55 to 59	40% (31-50%)	60% (50-69%)	0.98 (0.59-1.5)	3.6% (1.8-6%)	4.2% (1.8-7.3%)	210 (90-360)	7300 (3500-12000)
Nepal	2000	Male	60 to 64	36% (26-47%)	64% (53-74%)	0.87 (0.49-1.4)	2.6% (1.1-4.9%)	3% (1.1-5.8%)	170 (63-350)	5300 (2100-9900)
Nepal	2000	Male	65 to 69	34% (24-45%)	66% (55-76%)	0.74 (0.44-1.2)	1.8% (0.54-3.5%)	1.9% (0.41-4%)	120 (26-250)	3100 (920-6100)
Nepal	2000	Male	70 to 74	31% (21-41%)	69% (59-79%)	0.6 (0.33-0.98)	1.1% (0.091-2.4%)	1.1% (-0.045-2.7%)	73 (-2.9-180)	1500 (130-3500)
Nepal	2000	Male	75 to 79	26% (17-38%)	74% (62-83%)	0.46 (0.22-0.78)	0.58% (-0.19-1.5%)	0.58% (-0.33-1.6%)	31 (-18-90)	530 (-170-1400)
Nepal	2000	Male	80 to 84	24% (14-35%)	76% (65-86%)	0.36 (0.18-0.61)	0.3% (-0.26-0.93%)	0.26% (-0.43-0.99%)	8.8 (-15-33)	130 (-110-390)
Nepal	2000	Male	85 to 89	23% (14-36%)	77% (64-86%)	0.32 (0.16-0.55)	0.11% (-0.37-0.54%)	0.027% (-0.55-0.52%)	0.53 (-11-10)	18 (-63-92)
Nepal	2000	Male	90 to 94	26% (16-39%)	74% (61-84%)	0.31 (0.15-0.55)	-0.052% (-0.55-0.37%)	-0.19% (-0.78-0.28%)	-1.4 (-5.8-2)	-2.3 (-24-16)
Nepal	2000	Male	95 plus	31% (17-48%)	69% (52-83%)	0.31 (0.12-0.62)	-0.088% (-0.61-0.36%)	-0.27% (-0.89-0.23%)	-0.49 (-1.6-0.41)	-0.62 (-4.3-2.6)
Nepal	2005	Female	All Ages	1.9% (1.6-2.3%)	98% (98-98%)	0.15 (0.12-0.18)	0.79% (0.63-1%)	0.98% (0.73-1.3%)	900 (660-1200)	41000 (33000-52000)
Nepal	2005	Female	15 to 19	1.2% (0.7-1.9%)	99% (98-99%)	0.051 (0.024-0.092)	0.49% (0.31-0.72%)	0.25% (0.11-0.5%)	4.6 (2-9.2)	1200 (760-1900)
Nepal	2005	Female	20 to 24	2.1% (1.2-3.1%)	98% (97-99%)	0.16 (0.085-0.26)	1.2% (0.85-1.8%)	0.86% (0.44-1.5%)	17 (8.8-30)	3000 (2000-4400)
Nepal	2005	Female	25 to 29	2.7% (1.7-4%)	97% (96-98%)	0.25 (0.15-0.41)	1.6% (1.1-2.2%)	1.5% (0.8-2.4%)	29 (16-47)	3600 (2400-5100)
Nepal	2005	Female	30 to 34	3.5% (2.1-5.3%)	97% (95-98%)	0.36 (0.21-0.56)	1.9% (1.4-2.7%)	2.3% (1.4-3.7%)	44 (27-72)	4000 (2700-5700)
Nepal	2005	Female	35 to 39	4.2% (2.7-6%)	96% (94-97%)	0.45 (0.26-0.68)	2.3% (1.6-3.2%)	3.1% (1.9-4.7%)	62 (39-94)	4300 (3000-6200)
Nepal	2005	Female	40 to 44	4.5% (3-6.4%)	95% (94-97%)	0.45 (0.26-0.68)	2.4% (1.7-3.4%)	3.5% (2.2-5.1%)	80 (51-120)	4600 (3200-6400)
Nepal	2005	Female	45 to 49	4.1% (2.5-5.9%)	96% (94-98%)	0.36 (0.2-0.56)	2.2% (1.6-3.1%)	3.2% (2.1-4.5%)	84 (54-120)	4200 (2900-5700)
Nepal	2005	Female	50 to 54	3.5% (2.1-5.2%)	97% (95-98%)	0.27 (0.14-0.44)	2% (1.3-2.8%)	2.6% (1.7-3.8%)	97 (61-150)	4100 (2800-5900)
Nepal	2005	Female	55 to 59	3.2% (1.9-4.9%)	97% (95-98%)	0.24 (0.12-0.41)	1.7% (1.1-2.5%)	2.2% (1.3-3.3%)	98 (59-150)	3500 (2200-5200)
Nepal	2005	Female	60 to 64	3.2% (1.8-5%)	97% (95-98%)	0.22 (0.11-0.39)	1.4% (0.8-2.2%)	1.7% (0.93-2.7%)	110 (57-170)	3200 (1800-4800)
Nepal	2005	Female	65 to 69	2.9% (1.7-4.3%)	97% (96-98%)	0.18 (0.092-0.3)	1.1% (0.65-1.6%)	1.3% (0.74-2%)	100 (57-160)	2400 (1500-3700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	2005	Female	70 to 74	2.6% (1.6-4.1%)	97% (96-98%)	0.13 (0.067-0.24)	0.79% (0.48-1.2%)	0.93% (0.55-1.5%)	78 (45-120)	1500 (890-2400)
Nepal	2005	Female	75 to 79	2.2% (1.2-3.7%)	98% (96-99%)	0.098 (0.046-0.18)	0.55% (0.32-0.9%)	0.63% (0.35-1.1%)	56 (31-94)	830 (480-1400)
Nepal	2005	Female	80 to 84	1.9% (0.99-3%)	98% (97-99%)	0.076 (0.034-0.13)	0.38% (0.23-0.61%)	0.43% (0.23-0.71%)	27 (14-44)	290 (170-470)
Nepal	2005	Female	85 to 89	1.9% (0.95-3.3%)	98% (97-99%)	0.068 (0.031-0.13)	0.28% (0.16-0.44%)	0.3% (0.16-0.5%)	12 (6.5-20)	100 (56-160)
Nepal	2005	Female	90 to 94	2.2% (1.1-3.5%)	98% (96-99%)	0.069 (0.03-0.13)	0.21% (0.11-0.38%)	0.21% (0.088-0.41%)	3.4 (1.4-6.7)	21 (11-37)
Nepal	2005	Female	95 plus	2.7% (1.3-4.8%)	97% (95-99%)	0.073 (0.027-0.15)	0.18% (0.064-0.39%)	0.15% (0.017-0.39%)	0.65 (0.075-1.7)	3.1 (1.1-6.6)
Nepal	2005	Male	All Ages	25% (21-30%)	75% (70-79%)	0.58 (0.49-0.68)	2.1% (1.5-2.8%)	2.5% (1.5-3.7%)	2300 (1400-3300)	110000 (80000-150000)
Nepal	2005	Male	15 to 19	14% (8.8-20%)	86% (80-91%)	0.23 (0.12-0.4)	1.3% (0.84-1.9%)	1.2% (0.65-2.1%)	22 (11-38)	2900 (1800-4300)
Nepal	2005	Male	20 to 24	32% (23-42%)	68% (58-77%)	0.74 (0.43-1.2)	3.8% (2.5-5.5%)	3.9% (2.1-6.4%)	80 (43-130)	8000 (5100-12000)
Nepal	2005	Male	25 to 29	45% (35-54%)	55% (46-65%)	1.1 (0.71-1.6)	5.4% (3.4-7.3%)	6% (3-9%)	130 (61-190)	11000 (6800-15000)
Nepal	2005	Male	30 to 34	51% (40-61%)	49% (39-60%)	1.3 (0.92-1.9)	6.4% (4.4-8.5%)	7.7% (4.7-11%)	190 (110-270)	14000 (9200-19000)
Nepal	2005	Male	35 to 39	52% (42-61%)	48% (39-58%)	1.4 (0.93-2)	6.4% (4.3-8.9%)	8% (4.8-12%)	230 (130-350)	14000 (9400-20000)
Nepal	2005	Male	40 to 44	52% (42-61%)	48% (39-58%)	1.4 (0.87-2)	6.2% (3.9-8.6%)	7.9% (4.6-11%)	240 (140-360)	13000 (8400-19000)
Nepal	2005	Male	45 to 49	51% (40-61%)	49% (39-60%)	1.4 (0.88-2)	5.8% (3.6-8.4%)	7.3% (4.2-11%)	250 (140-390)	12000 (7400-18000)
Nepal	2005	Male	50 to 54	47% (37-56%)	53% (44-63%)	1.3 (0.77-1.9)	5% (2.9-7.6%)	6.1% (3.3-9.6%)	280 (150-450)	12000 (6700-18000)
Nepal	2005	Male	55 to 59	41% (31-51%)	59% (49-69%)	1.2 (0.69-1.8)	4.1% (2.1-6.9%)	4.9% (2.2-8.5%)	250 (110-440)	8900 (4500-15000)
Nepal	2005	Male	60 to 64	37% (26-48%)	63% (52-74%)	1 (0.6-1.7)	3.1% (1.2-5.6%)	3.5% (1.2-6.7%)	250 (84-470)	7500 (3000-14000)
Nepal	2005	Male	65 to 69	34% (24-46%)	66% (54-76%)	0.89 (0.52-1.4)	2.1% (0.69-4%)	2.3% (0.59-4.7%)	190 (47-380)	4700 (1500-9000)
Nepal	2005	Male	70 to 74	31% (22-41%)	69% (59-78%)	0.72 (0.41-1.2)	1.3% (0.15-2.8%)	1.4% (-0.012-3.3%)	110 (-0.89-240)	2200 (240-4700)
Nepal	2005	Male	75 to 79	27% (17-38%)	73% (62-83%)	0.55 (0.27-0.93)	0.73% (-0.1-1.9%)	0.75% (-0.27-2.1%)	52 (-19-150)	870 (-120-2300)
Nepal	2005	Male	80 to 84	24% (15-35%)	76% (65-85%)	0.44 (0.22-0.73)	0.42% (-0.24-1.2%)	0.39% (-0.4-1.3%)	17 (-17-54)	220 (-130-630)
Nepal	2005	Male	85 to 89	24% (15-36%)	76% (64-85%)	0.39 (0.19-0.69)	0.18% (-0.41-0.74%)	0.096% (-0.62-0.75%)	2.3 (-15-18)	38 (-88-160)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	2005	Male	90 to 94	27% (16-39%)	73% (61-84%)	0.38 (0.18-0.68)	-0.033% (-0.63-0.53%)	-0.18% (-0.89-0.47%)	-1.7 (-8.5-4.4)	-1.9 (-36-30)
Nepal	2005	Male	95 plus	31% (17-48%)	69% (52-83%)	0.39 (0.15-0.78)	-0.099% (-0.79-0.51%)	-0.31% (-1.1-0.37%)	-0.71 (-2.6-0.86)	-0.9 (-7.2-4.6)
Nepal	2010	Female	All Ages	2% (1.6-2.4%)	98% (98-98%)	0.2 (0.15-0.25)	0.89% (0.74-1.1%)	1% (0.78-1.3%)	890 (680-1200)	43000 (35000-53000)
Nepal	2010	Female	15 to 19	1.5% (0.87-2.3%)	99% (98-99%)	0.077 (0.036-0.14)	0.62% (0.39-0.91%)	0.4% (0.16-0.81%)	6 (2.4-12)	1500 (910-2200)
Nepal	2010	Female	20 to 24	2.4% (1.5-3.5%)	98% (96-99%)	0.24 (0.13-0.38)	1.5% (1-2.1%)	1.2% (0.63-2%)	20 (10-34)	3500 (2200-5000)
Nepal	2010	Female	25 to 29	3% (1.9-4.5%)	97% (96-98%)	0.36 (0.2-0.58)	1.8% (1.3-2.4%)	1.8% (0.99-2.9%)	29 (16-48)	3800 (2600-5300)
Nepal	2010	Female	30 to 34	3.7% (2.3-5.5%)	96% (95-98%)	0.48 (0.29-0.76)	2.1% (1.5-3%)	2.5% (1.5-4%)	45 (27-73)	4300 (3100-6100)
Nepal	2010	Female	35 to 39	4.4% (2.7-6.3%)	96% (94-97%)	0.6 (0.32-0.94)	2.4% (1.7-3.2%)	3.4% (2.2-4.9%)	68 (44-100)	4900 (3500-6700)
Nepal	2010	Female	40 to 44	4.8% (3.2-6.7%)	95% (93-97%)	0.62 (0.34-0.97)	2.6% (1.9-3.6%)	3.8% (2.6-5.5%)	90 (61-130)	5300 (3800-7500)
Nepal	2010	Female	45 to 49	3.7% (2.3-5.4%)	96% (95-98%)	0.41 (0.23-0.68)	2.3% (1.6-3.1%)	3.2% (2.2-4.6%)	95 (63-130)	4800 (3300-6500)
Nepal	2010	Female	50 to 54	2.1% (1.1-3.4%)	98% (97-99%)	0.21 (0.096-0.36)	1.7% (1.2-2.3%)	2.2% (1.5-3.2%)	80 (52-120)	3500 (2500-5000)
Nepal	2010	Female	55 to 59	1.6% (0.81-2.6%)	98% (97-99%)	0.15 (0.069-0.28)	1.3% (0.92-1.9%)	1.7% (1.1-2.5%)	86 (53-130)	3200 (2200-4600)
Nepal	2010	Female	60 to 64	1.5% (0.76-2.6%)	98% (97-99%)	0.14 (0.059-0.26)	1.1% (0.7-1.6%)	1.3% (0.79-2%)	83 (50-130)	2600 (1700-3800)
Nepal	2010	Female	65 to 69	1.5% (0.74-2.4%)	99% (98-99%)	0.12 (0.054-0.22)	0.92% (0.6-1.3%)	1.1% (0.68-1.6%)	92 (57-140)	2300 (1500-3400)
Nepal	2010	Female	70 to 74	1.5% (0.82-2.5%)	99% (98-99%)	0.097 (0.041-0.18)	0.73% (0.46-1.1%)	0.87% (0.52-1.3%)	79 (47-120)	1500 (940-2300)
Nepal	2010	Female	75 to 79	1.5% (0.74-2.5%)	99% (97-99%)	0.085 (0.034-0.16)	0.57% (0.34-0.88%)	0.66% (0.38-1%)	59 (33-94)	880 (520-1400)
Nepal	2010	Female	80 to 84	1.5% (0.73-2.5%)	98% (98-99%)	0.078 (0.033-0.15)	0.45% (0.26-0.73%)	0.5% (0.28-0.82%)	35 (19-58)	380 (220-630)
Nepal	2010	Female	85 to 89	1.7% (0.82-2.8%)	98% (97-99%)	0.077 (0.033-0.15)	0.34% (0.2-0.61%)	0.37% (0.2-0.69%)	20 (10-35)	160 (87-280)
Nepal	2010	Female	90 to 94	1.9% (0.99-3.2%)	98% (97-99%)	0.079 (0.033-0.16)	0.27% (0.14-0.52%)	0.28% (0.13-0.58%)	6.5 (2.9-13)	38 (19-73)
Nepal	2010	Female	95 plus	2.3% (1.1-4.4%)	98% (96-99%)	0.084 (0.029-0.18)	0.25% (0.1-0.54%)	0.23% (0.053-0.55%)	1.5 (0.35-3.6)	6.3 (2.5-14)
Nepal	2010	Male	All Ages	20% (17-24%)	80% (76-83%)	0.6 (0.48-0.74)	2.5% (1.8-3.4%)	2.9% (1.7-4.4%)	2700 (1500-4100)	130000 (87000-170000)
Nepal	2010	Male	15 to 19	8.9% (5.3-14%)	91% (86-95%)	0.19 (0.088-0.34)	1.4% (0.97-2%)	1.1% (0.62-2%)	19 (10-34)	3100 (2100-4500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	2010	Male	20 to 24	24% (15-34%)	76% (66-85%)	0.66 (0.35-1.1)	4.2% (2.9-5.9%)	4% (2.2-6.6%)	80 (42-130)	8900 (6000-13000)
Nepal	2010	Male	25 to 29	36% (25-48%)	64% (52-75%)	1.1 (0.65-1.7)	5.8% (3.6-8%)	6.3% (2.8-9.8%)	120 (54-190)	11000 (6600-16000)
Nepal	2010	Male	30 to 34	44% (31-57%)	56% (43-69%)	1.4 (0.88-2.1)	6.8% (3.9-9.3%)	8.2% (3.6-12%)	170 (72-250)	13000 (7800-18000)
Nepal	2010	Male	35 to 39	46% (33-58%)	54% (42-67%)	1.5 (0.9-2.3)	6.9% (4.5-9.6%)	8.7% (4.7-13%)	220 (120-330)	14000 (8800-20000)
Nepal	2010	Male	40 to 44	46% (34-57%)	54% (43-66%)	1.5 (0.87-2.3)	6.8% (4.4-9.4%)	8.7% (5.1-13%)	250 (150-380)	14000 (9000-20000)
Nepal	2010	Male	45 to 49	44% (29-56%)	56% (44-71%)	1.4 (0.8-2.2)	6.4% (3.8-9.2%)	8.1% (4.5-12%)	300 (160-450)	14000 (8700-21000)
Nepal	2010	Male	50 to 54	40% (29-51%)	60% (49-71%)	1.3 (0.76-2.1)	5.6% (3.3-8.5%)	7% (3.8-11%)	310 (160-490)	13000 (7400-20000)
Nepal	2010	Male	55 to 59	35% (24-47%)	65% (53-76%)	1.2 (0.69-2)	4.8% (2.4-7.9%)	5.8% (2.7-9.8%)	350 (170-610)	13000 (6500-21000)
Nepal	2010	Male	60 to 64	31% (21-43%)	69% (57-79%)	1.1 (0.58-1.9)	3.6% (1.7-6.7%)	4.2% (1.8-8.2%)	310 (130-590)	9300 (4200-17000)
Nepal	2010	Male	65 to 69	29% (19-40%)	71% (60-81%)	0.94 (0.5-1.6)	2.5% (1.1-4.7%)	2.9% (1.1-5.6%)	250 (94-500)	6300 (2600-12000)
Nepal	2010	Male	70 to 74	27% (18-37%)	73% (63-82%)	0.77 (0.4-1.3)	1.7% (0.42-3.4%)	1.8% (0.31-4%)	170 (30-380)	3500 (830-7300)
Nepal	2010	Male	75 to 79	24% (15-35%)	76% (65-85%)	0.61 (0.29-1.1)	1% (0.016-2.4%)	1.1% (-0.084-2.8%)	84 (-6.4-210)	1400 (21-3200)
Nepal	2010	Male	80 to 84	22% (13-32%)	78% (68-87%)	0.5 (0.24-0.87)	0.64% (-0.096-1.6%)	0.64% (-0.22-1.8%)	35 (-12-95)	430 (-63-1100)
Nepal	2010	Male	85 to 89	22% (13-34%)	78% (66-87%)	0.45 (0.21-0.79)	0.35% (-0.27-1.1%)	0.29% (-0.46-1.1%)	9.5 (-15-37)	100 (-78-310)
Nepal	2010	Male	90 to 94	25% (15-37%)	75% (63-85%)	0.44 (0.21-0.84)	0.11% (-0.57-0.82%)	-0.023% (-0.84-0.8%)	-0.29 (-10-10)	8 (-42-63)
Nepal	2010	Male	95 plus	29% (16-45%)	71% (55-84%)	0.45 (0.17-0.93)	0.016% (-0.71-0.84%)	-0.19% (-1-0.73%)	-0.58 (-3.2-2.2)	0.2 (-8.4-11)
Nepal	2016	Female	All Ages	1.5% (1.2-1.9%)	98% (98-99%)	0.16 (0.095-0.25)	0.93% (0.76-1.1%)	1% (0.74-1.3%)	900 (660-1200)	42000 (34000-52000)
Nepal	2016	Female	15 to 19	1.1% (0.6-1.8%)	99% (98-99%)	0.065 (0.024-0.13)	0.65% (0.42-0.97%)	0.36% (0.13-0.76%)	4.3 (1.4-9.2)	1400 (850-2200)
Nepal	2016	Female	20 to 24	1.7% (0.9-2.6%)	98% (97-99%)	0.18 (0.077-0.34)	1.5% (1-2.1%)	0.92% (0.47-1.6%)	13 (6.2-21)	3300 (2100-4800)
Nepal	2016	Female	25 to 29	2.1% (1.1-3.2%)	98% (97-99%)	0.27 (0.12-0.48)	1.7% (1.2-2.2%)	1.3% (0.68-2.1%)	19 (10-31)	3600 (2500-4900)
Nepal	2016	Female	30 to 34	2.4% (1.4-3.9%)	98% (96-99%)	0.35 (0.16-0.64)	1.7% (1.3-2.4%)	1.8% (1.1-2.8%)	30 (18-47)	3700 (2600-5300)
Nepal	2016	Female	35 to 39	2.9% (1.7-4.3%)	97% (96-98%)	0.43 (0.2-0.78)	1.9% (1.4-2.5%)	2.5% (1.7-3.6%)	51 (34-75)	4300 (3200-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	2016	Female	40 to 44	3.1% (1.9-4.6%)	97% (95-98%)	0.43 (0.2-0.78)	2% (1.5-2.7%)	2.9% (2-4%)	72 (49-100)	4700 (3300-6300)
Nepal	2016	Female	45 to 49	2.6% (1.4-3.9%)	97% (96-99%)	0.31 (0.13-0.59)	1.9% (1.4-2.5%)	2.6% (1.8-3.7%)	81 (55-120)	4400 (3100-5900)
Nepal	2016	Female	50 to 54	1.8% (0.99-3%)	98% (97-99%)	0.2 (0.085-0.38)	1.6% (1.2-2.2%)	2.1% (1.4-3%)	84 (55-120)	3900 (2800-5300)
Nepal	2016	Female	55 to 59	1.6% (0.75-2.5%)	98% (97-99%)	0.16 (0.068-0.34)	1.3% (0.96-1.9%)	1.7% (1.1-2.5%)	83 (53-120)	3200 (2200-4600)
Nepal	2016	Female	60 to 64	1.5% (0.75-2.6%)	98% (97-99%)	0.15 (0.059-0.32)	1.1% (0.74-1.6%)	1.3% (0.81-2%)	100 (62-150)	3200 (2100-4700)
Nepal	2016	Female	65 to 69	1.5% (0.76-2.4%)	98% (98-99%)	0.13 (0.055-0.27)	0.96% (0.63-1.4%)	1.2% (0.71-1.8%)	110 (69-170)	2800 (1800-4200)
Nepal	2016	Female	70 to 74	1.5% (0.81-2.5%)	99% (98-99%)	0.11 (0.041-0.22)	0.79% (0.48-1.2%)	0.95% (0.56-1.5%)	97 (57-150)	1900 (1200-2900)
Nepal	2016	Female	75 to 79	1.5% (0.77-2.5%)	99% (97-99%)	0.094 (0.033-0.2)	0.62% (0.35-0.97%)	0.73% (0.4-1.2%)	73 (39-120)	1100 (600-1700)
Nepal	2016	Female	80 to 84	1.5% (0.78-2.5%)	98% (98-99%)	0.088 (0.033-0.17)	0.5% (0.26-0.85%)	0.58% (0.28-0.99%)	44 (22-76)	480 (260-810)
Nepal	2016	Female	85 to 89	1.7% (0.84-2.9%)	98% (97-99%)	0.086 (0.033-0.18)	0.4% (0.2-0.74%)	0.45% (0.21-0.85%)	25 (12-49)	200 (100-380)
Nepal	2016	Female	90 to 94	1.9% (1.1-3.2%)	98% (97-99%)	0.088 (0.032-0.19)	0.33% (0.14-0.67%)	0.34% (0.13-0.75%)	10 (3.8-22)	59 (25-120)
Nepal	2016	Female	95 plus	2.4% (1.1-4.4%)	98% (96-99%)	0.094 (0.03-0.21)	0.3% (0.11-0.68%)	0.28% (0.067-0.72%)	2.6 (0.62-6.7)	11 (4-25)
Nepal	2016	Male	All Ages	21% (17-25%)	79% (75-83%)	0.66 (0.41-0.99)	3% (1.9-4.5%)	3.3% (1.7-5.7%)	3100 (1500-5300)	140000 (90000-210000)
Nepal	2016	Male	15 to 19	9.1% (5.3-14%)	91% (86-95%)	0.2 (0.084-0.39)	1.5% (1-2.2%)	1.2% (0.64-2.3%)	22 (11-40)	3700 (2400-5400)
Nepal	2016	Male	20 to 24	24% (15-34%)	76% (66-85%)	0.69 (0.32-1.3)	4.5% (3-6.7%)	4.4% (2.3-8.1%)	98 (47-180)	11000 (7200-17000)
Nepal	2016	Male	25 to 29	37% (25-48%)	63% (52-75%)	1.2 (0.59-2.1)	6.3% (4.1-9.5%)	7.1% (3.5-12%)	140 (70-250)	13000 (8400-20000)
Nepal	2016	Male	30 to 34	44% (31-58%)	56% (42-69%)	1.5 (0.77-2.6)	7.2% (4.6-11%)	8.9% (4.2-15%)	180 (88-290)	14000 (8400-21000)
Nepal	2016	Male	35 to 39	46% (33-58%)	54% (42-67%)	1.6 (0.81-2.7)	7.3% (4.4-11%)	9.3% (4.6-15%)	210 (110-340)	14000 (8600-21000)
Nepal	2016	Male	40 to 44	46% (33-57%)	54% (43-67%)	1.6 (0.79-2.8)	7.1% (4.3-11%)	9.3% (4.9-15%)	270 (140-430)	15000 (9200-23000)
Nepal	2016	Male	45 to 49	44% (30-56%)	56% (44-70%)	1.5 (0.76-2.7)	6.8% (3.9-11%)	8.8% (4.7-15%)	330 (170-540)	16000 (9300-25000)
Nepal	2016	Male	50 to 54	40% (29-52%)	60% (48-71%)	1.4 (0.72-2.5)	6% (3.2-9.7%)	7.6% (3.7-13%)	360 (180-620)	15000 (8200-25000)
Nepal	2016	Male	55 to 59	35% (25-47%)	65% (53-75%)	1.3 (0.65-2.4)	5.1% (2.5-8.9%)	6.3% (2.8-11%)	370 (160-650)	13000 (6300-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nepal	2016	Male	60 to 64	32% (21-44%)	68% (56-79%)	1.2 (0.53-2.2)	4% (1.6-7.6%)	4.7% (1.7-9.4%)	380 (130-760)	11000 (4700-22000)
Nepal	2016	Male	65 to 69	30% (19-41%)	70% (59-81%)	1 (0.47-1.8)	2.8% (1-5.7%)	3.2% (1-7%)	300 (90-620)	7500 (2600-15000)
Nepal	2016	Male	70 to 74	27% (18-37%)	73% (63-82%)	0.82 (0.36-1.5)	1.9% (0.47-4.5%)	2.2% (0.38-5.4%)	220 (36-530)	4400 (1000-10000)
Nepal	2016	Male	75 to 79	24% (15-36%)	76% (64-85%)	0.65 (0.26-1.3)	1.2% (0.11-3.3%)	1.3% (-0.0018-3.8%)	130 (-0.19-380)	2000 (190-5600)
Nepal	2016	Male	80 to 84	22% (13-33%)	78% (67-87%)	0.53 (0.21-1)	0.77% (-0.11-2.3%)	0.8% (-0.27-2.5%)	52 (-18-170)	630 (-96-1900)
Nepal	2016	Male	85 to 89	22% (13-34%)	78% (66-87%)	0.47 (0.2-0.91)	0.44% (-0.3-1.6%)	0.4% (-0.45-1.7%)	16 (-18-71)	160 (-100-570)
Nepal	2016	Male	90 to 94	25% (15-37%)	75% (63-85%)	0.47 (0.19-0.96)	0.17% (-0.57-1.2%)	0.048% (-0.81-1.1%)	0.86 (-15-21)	18 (-60-130)
Nepal	2016	Male	95 plus	29% (16-46%)	71% (54-84%)	0.47 (0.16-1)	0.067% (-0.76-1.2%)	-0.14% (-1.1-1.1%)	-0.56 (-4.4-4.8)	1.1 (-12-19)
Pakistan	1990	Female	All Ages	1.3% (0.98-1.7%)	99% (98-99%)	0.031 (0.029-0.033)	0.24% (0.19-0.29%)	0.28% (0.18-0.38%)	1300 (880-1900)	81000 (63000-100000)
Pakistan	1990	Female	15 to 19	0.98% (0.49-1.7%)	99% (98-100%)	0.014 (0.0057-0.028)	0.36% (0.22-0.54%)	0.074% (0.02-0.17%)	6 (1.6-14)	3700 (2200-5800)
Pakistan	1990	Female	20 to 24	1.1% (0.51-1.9%)	99% (98-99%)	0.031 (0.013-0.054)	0.78% (0.52-1.1%)	0.26% (0.14-0.46%)	25 (13-44)	8500 (5400-12000)
Pakistan	1990	Female	25 to 29	1.3% (0.65-2.2%)	99% (98-99%)	0.052 (0.025-0.088)	0.93% (0.66-1.3%)	0.44% (0.25-0.74%)	40 (23-66)	8800 (5900-12000)
Pakistan	1990	Female	30 to 34	1.5% (0.73-2.6%)	98% (97-99%)	0.069 (0.036-0.11)	1% (0.76-1.4%)	0.79% (0.49-1.2%)	68 (41-100)	8800 (6200-12000)
Pakistan	1990	Female	35 to 39	1.8% (0.91-2.8%)	98% (97-99%)	0.079 (0.041-0.13)	1.1% (0.77-1.4%)	1% (0.63-1.5%)	85 (52-130)	7900 (5500-11000)
Pakistan	1990	Female	40 to 44	1.9% (1.1-3%)	98% (97-99%)	0.08 (0.041-0.13)	1% (0.73-1.4%)	1.1% (0.67-1.6%)	98 (61-150)	7500 (5200-10000)
Pakistan	1990	Female	45 to 49	1.9% (1-3.1%)	98% (97-99%)	0.073 (0.034-0.13)	0.98% (0.7-1.4%)	1.1% (0.67-1.7%)	130 (77-190)	7600 (5300-10000)
Pakistan	1990	Female	50 to 54	2% (1.1-3.3%)	98% (97-99%)	0.075 (0.036-0.13)	0.96% (0.66-1.3%)	1.1% (0.65-1.6%)	150 (88-230)	7200 (4800-10000)
Pakistan	1990	Female	55 to 59	2.2% (1.2-3.4%)	98% (97-99%)	0.086 (0.043-0.15)	0.88% (0.57-1.3%)	1% (0.57-1.7%)	170 (94-280)	6700 (4200-10000)
Pakistan	1990	Female	60 to 64	2.3% (1.2-3.8%)	98% (96-99%)	0.088 (0.041-0.16)	0.72% (0.4-1.2%)	0.81% (0.38-1.4%)	160 (76-290)	5300 (2900-8600)
Pakistan	1990	Female	65 to 69	2.5% (1.4-4%)	98% (96-99%)	0.081 (0.04-0.14)	0.57% (0.3-0.91%)	0.65% (0.31-1.1%)	160 (74-260)	4000 (2100-6400)
Pakistan	1990	Female	70 to 74	2.6% (1.5-4.2%)	97% (96-98%)	0.07 (0.033-0.12)	0.4% (0.18-0.72%)	0.46% (0.18-0.84%)	120 (48-230)	2500 (1100-4300)
Pakistan	1990	Female	75 to 79	2.7% (1.4-4.6%)	97% (95-99%)	0.062 (0.026-0.12)	0.26% (0.092-0.52%)	0.29% (0.081-0.61%)	72 (20-150)	1100 (390-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	1990	Female	80 to 84	2.9% (1.5-4.8%)	97% (95-99%)	0.06 (0.025-0.11)	0.18% (0.027-0.39%)	0.19% (0.0071-0.45%)	36 (1.3-85)	420 (62-920)
Pakistan	1990	Female	85 to 89	3.2% (1.6-5.4%)	97% (95-98%)	0.06 (0.027-0.11)	0.11% (-0.026-0.28%)	0.1% (-0.063-0.31%)	13 (-7.7-39)	120 (-29-310)
Pakistan	1990	Female	90 to 94	3.7% (1.9-6.3%)	96% (94-98%)	0.061 (0.026-0.12)	0.054% (-0.087-0.21%)	0.025% (-0.14-0.2%)	1.3 (-7.5-11)	17 (-27-67)
Pakistan	1990	Female	95 plus	4.6% (2-8.8%)	95% (91-98%)	0.066 (0.023-0.14)	0.02% (-0.16-0.2%)	-0.038% (-0.26-0.17%)	-0.58 (-3.9-2.5)	1.2 (-9.8-12)
Pakistan	1990	Male	All Ages	0.72% (0.55-0.96%)	99% (99-99%)	0.00097 (0.00073-0.0012)	0.38% (0.28-0.48%)	0.25% (0.16-0.35%)	1300 (840-1900)	130000 (98000-170000)
Pakistan	1990	Male	15 to 19	0.5% (0.21-0.9%)	100% (99-100%)	0.00057 (0.00016-0.0012)	0.71% (0.47-1%)	0.18% (0.088-0.32%)	14 (6.7-25)	6700 (4100-10000)
Pakistan	1990	Male	20 to 24	0.74% (0.28-1.3%)	99% (99-100%)	0.0011 (0.00034-0.0022)	1.6% (1.1-2.3%)	0.47% (0.25-0.78%)	47 (24-77)	17000 (11000-25000)
Pakistan	1990	Male	25 to 29	1% (0.46-1.7%)	99% (98-100%)	0.0016 (0.00058-0.0031)	2% (1.5-2.7%)	0.7% (0.35-1.2%)	70 (35-120)	19000 (13000-27000)
Pakistan	1990	Male	30 to 34	1.3% (0.56-2.3%)	99% (98-99%)	0.0021 (0.00088-0.0039)	2.2% (1.6-2.9%)	1.1% (0.59-1.9%)	110 (58-190)	19000 (13000-26000)
Pakistan	1990	Male	35 to 39	1.5% (0.69-2.4%)	99% (98-99%)	0.0025 (0.001-0.0045)	1.9% (1.4-2.6%)	1.2% (0.57-2.1%)	110 (54-200)	15000 (10000-20000)
Pakistan	1990	Male	40 to 44	1.5% (0.8-2.4%)	99% (98-99%)	0.0025 (0.001-0.0046)	1.7% (1.2-2.3%)	1.1% (0.58-1.9%)	130 (65-220)	13000 (8900-19000)
Pakistan	1990	Male	45 to 49	1.4% (0.66-2.2%)	99% (98-99%)	0.0024 (0.00098-0.0047)	1.4% (1-2%)	1.1% (0.55-1.9%)	160 (77-280)	12000 (8300-18000)
Pakistan	1990	Male	50 to 54	1.4% (0.68-2.3%)	99% (98-99%)	0.0024 (0.001-0.0046)	1.1% (0.77-1.5%)	0.83% (0.44-1.4%)	140 (76-240)	9600 (6400-13000)
Pakistan	1990	Male	55 to 59	1.4% (0.72-2.4%)	99% (98-99%)	0.0025 (0.001-0.0046)	0.83% (0.57-1.1%)	0.62% (0.33-1.1%)	130 (66-210)	7300 (4900-10000)
Pakistan	1990	Male	60 to 64	1.3% (0.66-2.3%)	99% (98-99%)	0.0023 (0.00088-0.0044)	0.59% (0.4-0.81%)	0.43% (0.22-0.69%)	110 (57-180)	5300 (3500-7500)
Pakistan	1990	Male	65 to 69	1.2% (0.59-1.9%)	99% (98-99%)	0.0018 (0.00077-0.0033)	0.44% (0.32-0.58%)	0.34% (0.21-0.52%)	100 (61-150)	3800 (2700-5200)
Pakistan	1990	Male	70 to 74	1% (0.55-1.7%)	99% (98-99%)	0.0014 (0.00051-0.0029)	0.32% (0.22-0.44%)	0.25% (0.15-0.4%)	85 (50-130)	2400 (1600-3300)
Pakistan	1990	Male	75 to 79	0.96% (0.44-1.7%)	99% (98-100%)	0.0012 (0.00039-0.0026)	0.25% (0.17-0.35%)	0.22% (0.13-0.34%)	67 (39-110)	1300 (860-1900)
Pakistan	1990	Male	80 to 84	0.88% (0.41-1.5%)	99% (98-100%)	0.0011 (0.00035-0.0022)	0.17% (0.12-0.23%)	0.14% (0.088-0.23%)	32 (20-51)	470 (320-670)
Pakistan	1990	Male	85 to 89	0.9% (0.42-1.7%)	99% (98-100%)	0.0011 (0.00035-0.0023)	0.14% (0.097-0.19%)	0.11% (0.068-0.17%)	16 (9.8-25)	180 (120-240)
Pakistan	1990	Male	90 to 94	1% (0.49-1.9%)	99% (98-100%)	0.0012 (0.00035-0.0026)	0.13% (0.094-0.18%)	0.094% (0.053-0.15%)	5.8 (3.3-9.2)	50 (35-70)
Pakistan	1990	Male	95 plus	1.3% (0.54-2.6%)	99% (97-99%)	0.0014 (0.00029-0.0037)	0.16% (0.11-0.22%)	0.093% (0.049-0.15%)	1.6 (0.84-2.6)	11 (7.7-16)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	1995	Female	All Ages	1.3% (1-1.7%)	99% (98-99%)	0.034 (0.032-0.037)	0.31% (0.25-0.38%)	0.37% (0.26-0.51%)	2000 (1400-2800)	110000 (86000-130000)
Pakistan	1995	Female	15 to 19	0.99% (0.53-1.7%)	99% (98-99%)	0.015 (0.0062-0.03)	0.35% (0.21-0.52%)	0.089% (0.029-0.19%)	9.8 (3.1-22)	4600 (2700-7200)
Pakistan	1995	Female	20 to 24	1.1% (0.54-1.9%)	99% (98-99%)	0.034 (0.015-0.058)	0.78% (0.52-1.1%)	0.31% (0.16-0.52%)	38 (19-63)	10000 (6600-15000)
Pakistan	1995	Female	25 to 29	1.3% (0.67-2.3%)	99% (98-99%)	0.055 (0.027-0.094)	0.93% (0.67-1.3%)	0.5% (0.28-0.81%)	58 (33-93)	11000 (7300-15000)
Pakistan	1995	Female	30 to 34	1.6% (0.78-2.8%)	98% (97-99%)	0.074 (0.039-0.12)	1.1% (0.81-1.4%)	0.9% (0.57-1.3%)	100 (65-160)	12000 (8400-16000)
Pakistan	1995	Female	35 to 39	1.8% (0.92-2.9%)	98% (97-99%)	0.085 (0.044-0.13)	1.1% (0.83-1.5%)	1.1% (0.73-1.7%)	140 (86-200)	11000 (8200-15000)
Pakistan	1995	Female	40 to 44	2% (1.1-3.1%)	98% (97-99%)	0.086 (0.044-0.14)	1.1% (0.77-1.4%)	1.2% (0.73-1.7%)	130 (84-200)	9300 (6600-13000)
Pakistan	1995	Female	45 to 49	2% (1-3.1%)	98% (97-99%)	0.079 (0.036-0.14)	1.1% (0.75-1.5%)	1.2% (0.75-1.8%)	170 (100-260)	9400 (6400-13000)
Pakistan	1995	Female	50 to 54	2.1% (1.2-3.4%)	98% (97-99%)	0.082 (0.04-0.14)	1% (0.71-1.5%)	1.2% (0.75-1.8%)	220 (140-330)	10000 (6800-14000)
Pakistan	1995	Female	55 to 59	2.2% (1.3-3.6%)	98% (96-99%)	0.095 (0.047-0.16)	0.98% (0.63-1.5%)	1.1% (0.64-1.8%)	250 (150-410)	9500 (6000-15000)
Pakistan	1995	Female	60 to 64	2.3% (1.3-4%)	98% (96-99%)	0.096 (0.045-0.18)	0.8% (0.44-1.3%)	0.9% (0.43-1.5%)	250 (120-430)	7700 (4300-13000)
Pakistan	1995	Female	65 to 69	2.5% (1.4-4.1%)	97% (96-99%)	0.087 (0.043-0.15)	0.64% (0.34-1%)	0.73% (0.35-1.2%)	240 (120-400)	5900 (3200-9600)
Pakistan	1995	Female	70 to 74	2.7% (1.6-4.3%)	97% (96-98%)	0.075 (0.035-0.13)	0.46% (0.21-0.82%)	0.52% (0.22-0.99%)	190 (79-370)	3700 (1700-6700)
Pakistan	1995	Female	75 to 79	2.8% (1.5-4.6%)	97% (95-98%)	0.066 (0.028-0.13)	0.3% (0.1-0.59%)	0.33% (0.098-0.68%)	110 (32-230)	1700 (570-3300)
Pakistan	1995	Female	80 to 84	3% (1.6-5.2%)	97% (95-98%)	0.065 (0.027-0.12)	0.21% (0.03-0.45%)	0.22% (0.012-0.5%)	55 (3-130)	640 (90-1400)
Pakistan	1995	Female	85 to 89	3.3% (1.7-5.7%)	97% (94-98%)	0.065 (0.029-0.12)	0.13% (-0.03-0.34%)	0.12% (-0.062-0.37%)	19 (-9.9-59)	180 (-41-470)
Pakistan	1995	Female	90 to 94	3.8% (2-6.6%)	96% (93-98%)	0.067 (0.028-0.14)	0.063% (-0.1-0.25%)	0.035% (-0.16-0.26%)	2.3 (-11-17)	25 (-38-100)
Pakistan	1995	Female	95 plus	4.6% (2.2-8.6%)	95% (91-98%)	0.071 (0.023-0.15)	0.024% (-0.19-0.24%)	-0.033% (-0.28-0.2%)	-0.64 (-5.3-3.9)	1.8 (-14-18)
Pakistan	1995	Male	All Ages	0.75% (0.58-0.99%)	99% (99-99%)	0.001 (0.00079-0.0013)	0.45% (0.34-0.57%)	0.29% (0.19-0.41%)	1700 (1100-2500)	160000 (120000-210000)
Pakistan	1995	Male	15 to 19	0.51% (0.23-0.91%)	99% (99-100%)	0.00061 (0.00018-0.0014)	0.7% (0.46-1%)	0.19% (0.091-0.32%)	18 (8.5-31)	8000 (4900-12000)
Pakistan	1995	Male	20 to 24	0.76% (0.3-1.4%)	99% (99-100%)	0.0011 (0.00035-0.0024)	1.6% (1.1-2.2%)	0.49% (0.25-0.82%)	58 (30-95)	19000 (12000-27000)
Pakistan	1995	Male	25 to 29	1% (0.48-1.8%)	99% (98-100%)	0.0017 (0.00066-0.0033)	2% (1.5-2.7%)	0.73% (0.37-1.2%)	88 (44-150)	22000 (15000-31000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	1995	Male	30 to 34	1.3% (0.61-2.5%)	99% (98-99%)	0.0022 (0.00099-0.0041)	2.2% (1.6-2.9%)	1.2% (0.6-1.9%)	150 (77-240)	24000 (16000-33000)
Pakistan	1995	Male	35 to 39	1.5% (0.73-2.5%)	98% (98-99%)	0.0026 (0.0011-0.0048)	2% (1.4-2.6%)	1.2% (0.57-2.1%)	160 (75-280)	20000 (14000-27000)
Pakistan	1995	Male	40 to 44	1.5% (0.82-2.5%)	98% (98-99%)	0.0027 (0.0011-0.0049)	1.7% (1.2-2.3%)	1.1% (0.6-1.9%)	150 (78-260)	15000 (10000-21000)
Pakistan	1995	Male	45 to 49	1.4% (0.68-2.3%)	99% (98-99%)	0.0026 (0.001-0.005)	1.4% (0.99-2%)	1.1% (0.57-1.8%)	180 (95-300)	14000 (9200-19000)
Pakistan	1995	Male	50 to 54	1.4% (0.7-2.3%)	99% (98-99%)	0.0026 (0.001-0.0049)	1.1% (0.77-1.5%)	0.83% (0.46-1.4%)	190 (100-300)	12000 (8000-17000)
Pakistan	1995	Male	55 to 59	1.4% (0.72-2.4%)	99% (98-99%)	0.0027 (0.0011-0.0049)	0.82% (0.57-1.1%)	0.63% (0.34-1%)	170 (88-280)	9100 (6100-13000)
Pakistan	1995	Male	60 to 64	1.4% (0.66-2.4%)	99% (98-99%)	0.0024 (0.00095-0.0047)	0.58% (0.39-0.8%)	0.43% (0.23-0.69%)	140 (75-240)	6500 (4300-9200)
Pakistan	1995	Male	65 to 69	1.2% (0.61-2%)	99% (98-99%)	0.0019 (0.00081-0.0036)	0.43% (0.32-0.58%)	0.34% (0.21-0.51%)	130 (78-200)	4600 (3300-6300)
Pakistan	1995	Male	70 to 74	1.1% (0.57-1.8%)	99% (98-99%)	0.0015 (0.00055-0.0031)	0.32% (0.22-0.45%)	0.26% (0.16-0.41%)	110 (67-170)	3000 (2000-4300)
Pakistan	1995	Male	75 to 79	0.99% (0.46-1.7%)	99% (98-100%)	0.0013 (0.00039-0.0028)	0.26% (0.18-0.35%)	0.23% (0.13-0.34%)	91 (53-140)	1700 (1100-2400)
Pakistan	1995	Male	80 to 84	0.9% (0.42-1.6%)	99% (98-100%)	0.0011 (0.00037-0.0023)	0.18% (0.12-0.25%)	0.15% (0.094-0.24%)	45 (28-71)	640 (440-900)
Pakistan	1995	Male	85 to 89	0.91% (0.43-1.6%)	99% (98-100%)	0.0011 (0.00036-0.0024)	0.15% (0.11-0.2%)	0.12% (0.076-0.18%)	22 (14-33)	240 (170-320)
Pakistan	1995	Male	90 to 94	1.1% (0.5-1.9%)	99% (98-99%)	0.0012 (0.00036-0.0027)	0.14% (0.097-0.19%)	0.1% (0.061-0.16%)	7.9 (4.7-12)	65 (45-89)
Pakistan	1995	Male	95 plus	1.3% (0.54-2.7%)	99% (97-99%)	0.0014 (0.00031-0.0038)	0.17% (0.12-0.24%)	0.1% (0.056-0.17%)	2.1 (1.2-3.5)	14 (10-20)
Pakistan	2000	Female	All Ages	1.4% (1.1-1.7%)	99% (98-99%)	0.036 (0.033-0.038)	0.37% (0.31-0.45%)	0.43% (0.3-0.58%)	2300 (1600-3200)	130000 (100000-160000)
Pakistan	2000	Female	15 to 19	1% (0.53-1.7%)	99% (98-99%)	0.015 (0.0064-0.03)	0.35% (0.22-0.54%)	0.095% (0.033-0.21%)	12 (4.2-26)	5400 (3200-8600)
Pakistan	2000	Female	20 to 24	1.2% (0.55-2%)	99% (98-99%)	0.034 (0.015-0.059)	0.8% (0.54-1.1%)	0.33% (0.17-0.55%)	45 (24-76)	12000 (7800-18000)
Pakistan	2000	Female	25 to 29	1.4% (0.68-2.4%)	99% (98-99%)	0.056 (0.027-0.096)	0.98% (0.7-1.3%)	0.54% (0.31-0.86%)	66 (38-100)	12000 (8500-17000)
Pakistan	2000	Female	30 to 34	1.6% (0.78-2.9%)	98% (97-99%)	0.075 (0.04-0.12)	1.1% (0.85-1.5%)	0.96% (0.63-1.4%)	120 (75-170)	13000 (9600-18000)
Pakistan	2000	Female	35 to 39	1.9% (0.93-3%)	98% (97-99%)	0.086 (0.045-0.14)	1.2% (0.87-1.6%)	1.2% (0.8-1.7%)	160 (100-230)	14000 (9900-18000)
Pakistan	2000	Female	40 to 44	2% (1.1-3.2%)	98% (97-99%)	0.086 (0.045-0.14)	1.1% (0.8-1.5%)	1.2% (0.78-1.8%)	170 (110-250)	12000 (8200-16000)
Pakistan	2000	Female	45 to 49	2% (1.1-3.2%)	98% (97-99%)	0.079 (0.036-0.14)	1.1% (0.76-1.5%)	1.2% (0.77-1.9%)	180 (110-280)	10000 (7200-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	2000	Female	50 to 54	2.2% (1.2-3.5%)	98% (96-99%)	0.083 (0.039-0.15)	1.1% (0.72-1.5%)	1.2% (0.76-1.9%)	240 (140-370)	11000 (7200-16000)
Pakistan	2000	Female	55 to 59	2.3% (1.2-3.8%)	98% (96-99%)	0.095 (0.045-0.17)	1% (0.64-1.5%)	1.1% (0.66-1.8%)	300 (170-490)	11000 (7200-17000)
Pakistan	2000	Female	60 to 64	2.4% (1.3-4%)	98% (96-99%)	0.096 (0.044-0.17)	0.81% (0.47-1.3%)	0.91% (0.46-1.5%)	290 (140-490)	9000 (5000-14000)
Pakistan	2000	Female	65 to 69	2.6% (1.4-4.1%)	97% (96-99%)	0.087 (0.042-0.15)	0.66% (0.36-1%)	0.75% (0.38-1.2%)	290 (140-490)	7100 (3900-12000)
Pakistan	2000	Female	70 to 74	2.7% (1.5-4.3%)	97% (96-98%)	0.075 (0.035-0.14)	0.48% (0.22-0.86%)	0.55% (0.23-1%)	230 (100-430)	4500 (2000-8000)
Pakistan	2000	Female	75 to 79	2.9% (1.6-4.7%)	97% (95-98%)	0.067 (0.029-0.13)	0.31% (0.1-0.6%)	0.34% (0.091-0.69%)	130 (37-280)	2000 (700-4100)
Pakistan	2000	Female	80 to 84	3.1% (1.6-5.1%)	97% (95-98%)	0.066 (0.028-0.12)	0.21% (0.019-0.45%)	0.22% (-0.0027-0.52%)	67 (-0.8-160)	770 (71-1700)
Pakistan	2000	Female	85 to 89	3.4% (1.8-5.9%)	97% (94-98%)	0.066 (0.03-0.13)	0.13% (-0.037-0.34%)	0.12% (-0.07-0.38%)	22 (-12-68)	200 (-56-530)
Pakistan	2000	Female	90 to 94	3.9% (2.1-6.8%)	96% (93-98%)	0.068 (0.029-0.13)	0.056% (-0.12-0.25%)	0.026% (-0.18-0.26%)	1.9 (-13-19)	24 (-52-110)
Pakistan	2000	Female	95 plus	4.8% (2.1-9.1%)	95% (91-98%)	0.072 (0.024-0.15)	0.013% (-0.2-0.22%)	-0.049% (-0.31-0.19%)	-1 (-6.5-4.1)	1.1 (-17-19)
Pakistan	2000	Male	All Ages	0.72% (0.56-0.95%)	99% (99-99%)	0.00098 (0.00074-0.0012)	0.54% (0.41-0.69%)	0.34% (0.22-0.47%)	2100 (1400-3000)	190000 (140000-250000)
Pakistan	2000	Male	15 to 19	0.49% (0.22-0.87%)	100% (99-100%)	0.00057 (0.00017-0.0012)	0.72% (0.46-1%)	0.2% (0.094-0.34%)	22 (10-39)	9600 (6000-15000)
Pakistan	2000	Male	20 to 24	0.7% (0.28-1.2%)	99% (99-100%)	0.001 (0.00033-0.0021)	1.7% (1.1-2.3%)	0.52% (0.27-0.89%)	71 (35-120)	23000 (14000-33000)
Pakistan	2000	Male	25 to 29	0.92% (0.42-1.6%)	99% (98-100%)	0.0015 (0.00057-0.0029)	2.1% (1.5-2.8%)	0.79% (0.4-1.3%)	110 (51-170)	26000 (18000-36000)
Pakistan	2000	Male	30 to 34	1.1% (0.51-2%)	99% (98-99%)	0.0019 (0.00082-0.0035)	2.3% (1.6-3.1%)	1.2% (0.66-2%)	180 (92-300)	28000 (19000-39000)
Pakistan	2000	Male	35 to 39	1.3% (0.62-2%)	99% (98-99%)	0.0022 (0.00098-0.004)	2% (1.5-2.8%)	1.3% (0.6-2.2%)	210 (92-360)	25000 (17000-35000)
Pakistan	2000	Male	40 to 44	1.4% (0.75-2.2%)	99% (98-99%)	0.0024 (0.00098-0.0043)	1.7% (1.2-2.4%)	1.2% (0.63-2%)	210 (110-340)	20000 (14000-28000)
Pakistan	2000	Male	45 to 49	1.4% (0.67-2.1%)	99% (98-99%)	0.0025 (0.001-0.0049)	1.5% (1-2.1%)	1.1% (0.59-1.9%)	210 (110-350)	16000 (11000-22000)
Pakistan	2000	Male	50 to 54	1.4% (0.71-2.3%)	99% (98-99%)	0.0025 (0.001-0.0049)	1.1% (0.77-1.5%)	0.87% (0.48-1.4%)	210 (120-340)	13000 (8900-18000)
Pakistan	2000	Male	55 to 59	1.4% (0.73-2.5%)	99% (98-99%)	0.0027 (0.0011-0.0049)	0.85% (0.58-1.2%)	0.66% (0.36-1%)	210 (110-340)	11000 (7500-15000)
Pakistan	2000	Male	60 to 64	1.4% (0.66-2.4%)	99% (98-99%)	0.0024 (0.00096-0.0047)	0.6% (0.4-0.81%)	0.45% (0.24-0.69%)	180 (93-280)	7900 (5200-11000)
Pakistan	2000	Male	65 to 69	1.2% (0.62-2%)	99% (98-99%)	0.002 (0.00082-0.0036)	0.44% (0.32-0.6%)	0.35% (0.21-0.52%)	160 (96-240)	5500 (3900-7700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	2000	Male	70 to 74	1.1% (0.58-1.8%)	99% (98-99%)	0.0016 (0.00057-0.0032)	0.33% (0.23-0.47%)	0.27% (0.17-0.43%)	130 (83-210)	3500 (2500-5000)
Pakistan	2000	Male	75 to 79	1% (0.47-1.8%)	99% (98-100%)	0.0013 (0.00042-0.0029)	0.27% (0.18-0.37%)	0.24% (0.14-0.35%)	110 (66-170)	2100 (1400-2900)
Pakistan	2000	Male	80 to 84	0.94% (0.45-1.6%)	99% (98-100%)	0.0012 (0.0004-0.0024)	0.19% (0.13-0.26%)	0.16% (0.099-0.25%)	58 (35-89)	810 (560-1100)
Pakistan	2000	Male	85 to 89	0.95% (0.45-1.7%)	99% (98-100%)	0.0011 (0.00039-0.0024)	0.16% (0.11-0.21%)	0.13% (0.083-0.19%)	29 (18-43)	300 (210-400)
Pakistan	2000	Male	90 to 94	1.1% (0.52-1.9%)	99% (98-99%)	0.0012 (0.00038-0.0028)	0.15% (0.11-0.2%)	0.11% (0.066-0.17%)	10 (6-16)	82 (58-110)
Pakistan	2000	Male	95 plus	1.3% (0.56-2.7%)	99% (97-99%)	0.0015 (0.0003-0.0039)	0.18% (0.13-0.25%)	0.11% (0.06-0.18%)	2.7 (1.5-4.5)	18 (12-24)
Pakistan	2005	Female	All Ages	1.4% (1.1-1.8%)	99% (98-99%)	0.032 (0.028-0.035)	0.36% (0.29-0.43%)	0.36% (0.25-0.49%)	2100 (1500-2900)	130000 (100000-160000)
Pakistan	2005	Female	15 to 19	1% (0.55-1.7%)	99% (98-99%)	0.013 (0.0056-0.025)	0.33% (0.2-0.49%)	0.07% (0.026-0.14%)	11 (4.1-24)	6200 (3600-9900)
Pakistan	2005	Female	20 to 24	1.2% (0.57-2%)	99% (98-99%)	0.029 (0.013-0.051)	0.76% (0.51-1.1%)	0.26% (0.14-0.44%)	45 (24-73)	14000 (8900-20000)
Pakistan	2005	Female	25 to 29	1.4% (0.69-2.4%)	99% (98-99%)	0.048 (0.023-0.083)	0.91% (0.66-1.2%)	0.43% (0.25-0.69%)	64 (37-100)	14000 (9300-19000)
Pakistan	2005	Female	30 to 34	1.6% (0.82-2.9%)	98% (97-99%)	0.064 (0.034-0.1)	1% (0.77-1.3%)	0.77% (0.51-1.1%)	110 (69-160)	14000 (9700-18000)
Pakistan	2005	Female	35 to 39	1.9% (0.96-3%)	98% (97-99%)	0.073 (0.037-0.11)	1.1% (0.78-1.3%)	0.97% (0.65-1.4%)	150 (95-210)	14000 (9700-18000)
Pakistan	2005	Female	40 to 44	2% (1.1-3.3%)	98% (97-99%)	0.073 (0.037-0.12)	0.96% (0.71-1.3%)	0.98% (0.65-1.4%)	160 (110-240)	12000 (8800-17000)
Pakistan	2005	Female	45 to 49	2.1% (1.1-3.2%)	98% (97-99%)	0.068 (0.03-0.12)	0.95% (0.68-1.3%)	1% (0.68-1.5%)	190 (120-290)	12000 (8200-16000)
Pakistan	2005	Female	50 to 54	2.2% (1.2-3.5%)	98% (97-99%)	0.071 (0.034-0.12)	0.92% (0.65-1.3%)	1% (0.65-1.6%)	220 (130-330)	10000 (6800-15000)
Pakistan	2005	Female	55 to 59	2.3% (1.3-3.8%)	98% (96-99%)	0.081 (0.04-0.15)	0.84% (0.55-1.3%)	0.95% (0.54-1.5%)	260 (140-420)	10000 (6400-15000)
Pakistan	2005	Female	60 to 64	2.4% (1.3-4.2%)	98% (96-99%)	0.083 (0.038-0.16)	0.67% (0.38-1.1%)	0.74% (0.36-1.3%)	270 (130-470)	8600 (4800-14000)
Pakistan	2005	Female	65 to 69	2.6% (1.4-4.2%)	97% (96-99%)	0.075 (0.037-0.13)	0.53% (0.28-0.87%)	0.6% (0.28-1%)	260 (120-450)	6500 (3500-11000)
Pakistan	2005	Female	70 to 74	2.8% (1.7-4.5%)	97% (95-98%)	0.065 (0.031-0.12)	0.39% (0.17-0.7%)	0.44% (0.17-0.82%)	210 (81-400)	4200 (1800-7400)
Pakistan	2005	Female	75 to 79	3% (1.6-5%)	97% (95-98%)	0.058 (0.025-0.11)	0.23% (0.04-0.47%)	0.25% (0.019-0.54%)	110 (9-250)	1800 (310-3700)
Pakistan	2005	Female	80 to 84	3.2% (1.6-5.3%)	97% (95-98%)	0.057 (0.024-0.11)	0.14% (-0.041-0.34%)	0.14% (-0.08-0.38%)	49 (-28-130)	590 (-180-1500)
Pakistan	2005	Female	85 to 89	3.5% (1.9-6.1%)	96% (94-98%)	0.057 (0.026-0.11)	0.065% (-0.11-0.24%)	0.049% (-0.16-0.25%)	10 (-33-53)	120 (-190-420)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	2005	Female	90 to 94	4% (2.1-7%)	96% (93-98%)	0.058 (0.024-0.12)	0.0024% (-0.18-0.16%)	-0.037% (-0.24-0.15%)	-3 (-20-12)	1.3 (-83-77)
Pakistan	2005	Female	95 plus	4.9% (2.2-9.3%)	95% (91-98%)	0.062 (0.021-0.14)	-0.037% (-0.26-0.15%)	-0.1% (-0.37-0.11%)	-2.4 (-8.5-2.6)	-3.3 (-23-13)
Pakistan	2005	Male	All Ages	0.74% (0.58-0.95%)	99% (99-99%)	0.00088 (0.00066-0.0011)	0.57% (0.43-0.73%)	0.35% (0.23-0.49%)	2500 (1600-3500)	230000 (170000-290000)
Pakistan	2005	Male	15 to 19	0.49% (0.23-0.85%)	100% (99-100%)	0.00048 (0.00015-0.001)	0.59% (0.38-0.88%)	0.14% (0.066-0.26%)	27 (12-47)	12000 (6900-17000)
Pakistan	2005	Male	20 to 24	0.71% (0.31-1.3%)	99% (99-100%)	0.00088 (0.00029-0.0018)	1.4% (0.93-2%)	0.4% (0.2-0.69%)	83 (41-150)	27000 (17000-40000)
Pakistan	2005	Male	25 to 29	0.94% (0.45-1.6%)	99% (98-100%)	0.0013 (0.00052-0.0025)	1.8% (1.3-2.5%)	0.64% (0.32-1.1%)	120 (61-220)	31000 (21000-43000)
Pakistan	2005	Male	30 to 34	1.1% (0.53-2%)	99% (98-99%)	0.0016 (0.00073-0.0031)	2.1% (1.5-2.9%)	1.1% (0.56-1.8%)	210 (110-360)	32000 (22000-46000)
Pakistan	2005	Male	35 to 39	1.3% (0.63-2.1%)	99% (98-99%)	0.0019 (0.00082-0.0034)	1.9% (1.4-2.6%)	1.2% (0.55-2%)	250 (110-430)	29000 (19000-41000)
Pakistan	2005	Male	40 to 44	1.4% (0.76-2.2%)	99% (98-99%)	0.0021 (0.00085-0.0037)	1.7% (1.2-2.2%)	1.1% (0.6-1.9%)	260 (130-430)	25000 (17000-34000)
Pakistan	2005	Male	45 to 49	1.4% (0.68-2.2%)	99% (98-99%)	0.0021 (0.00086-0.004)	1.4% (0.99-2%)	1.1% (0.57-1.9%)	280 (140-490)	21000 (14000-29000)
Pakistan	2005	Male	50 to 54	1.4% (0.73-2.3%)	99% (98-99%)	0.0022 (0.00091-0.0042)	1.1% (0.77-1.5%)	0.86% (0.48-1.3%)	240 (130-380)	15000 (9800-20000)
Pakistan	2005	Male	55 to 59	1.5% (0.76-2.5%)	99% (98-99%)	0.0023 (0.00096-0.0042)	0.85% (0.58-1.2%)	0.66% (0.36-1%)	230 (120-370)	12000 (8100-17000)
Pakistan	2005	Male	60 to 64	1.4% (0.7-2.4%)	99% (98-99%)	0.0021 (0.00082-0.0042)	0.6% (0.41-0.81%)	0.45% (0.24-0.71%)	210 (110-330)	9300 (6100-13000)
Pakistan	2005	Male	65 to 69	1.2% (0.64-2%)	99% (98-99%)	0.0017 (0.00073-0.0031)	0.45% (0.32-0.6%)	0.35% (0.22-0.53%)	190 (110-290)	6500 (4600-8900)
Pakistan	2005	Male	70 to 74	1.1% (0.58-1.8%)	99% (98-99%)	0.0013 (0.0005-0.0027)	0.33% (0.24-0.46%)	0.27% (0.17-0.43%)	150 (99-240)	4100 (2900-5800)
Pakistan	2005	Male	75 to 79	1% (0.49-1.8%)	99% (98-100%)	0.0011 (0.00037-0.0025)	0.27% (0.19-0.37%)	0.24% (0.15-0.36%)	130 (79-190)	2400 (1700-3300)
Pakistan	2005	Male	80 to 84	0.94% (0.45-1.6%)	99% (98-100%)	0.00099 (0.00034-0.002)	0.19% (0.13-0.27%)	0.17% (0.1-0.26%)	70 (42-110)	970 (670-1300)
Pakistan	2005	Male	85 to 89	0.95% (0.43-1.8%)	99% (98-100%)	0.00096 (0.00032-0.0021)	0.16% (0.12-0.22%)	0.14% (0.086-0.2%)	36 (22-52)	370 (260-490)
Pakistan	2005	Male	90 to 94	1.1% (0.5-2%)	99% (98-100%)	0.001 (0.00032-0.0024)	0.16% (0.11-0.21%)	0.12% (0.073-0.18%)	13 (7.8-19)	100 (69-140)
Pakistan	2005	Male	95 plus	1.4% (0.56-2.8%)	99% (97-99%)	0.0013 (0.00026-0.0034)	0.19% (0.13-0.26%)	0.12% (0.067-0.2%)	3.4 (1.9-5.6)	21 (15-30)
Pakistan	2010	Female	All Ages	1.5% (1.1-1.8%)	99% (98-99%)	0.026 (0.019-0.034)	0.35% (0.28-0.43%)	0.29% (0.19-0.42%)	1700 (1100-2500)	130000 (95000-160000)
Pakistan	2010	Female	15 to 19	1% (0.56-1.7%)	99% (98-99%)	0.01 (0.0041-0.022)	0.38% (0.23-0.59%)	0.069% (0.027-0.13%)	8.8 (3.4-17)	6500 (3700-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	2010	Female	20 to 24	1.2% (0.57-2%)	99% (98-99%)	0.023 (0.0095-0.041)	0.84% (0.56-1.2%)	0.25% (0.14-0.41%)	38 (20-62)	15000 (9500-22000)
Pakistan	2010	Female	25 to 29	1.4% (0.72-2.5%)	99% (98-99%)	0.038 (0.017-0.068)	0.97% (0.69-1.3%)	0.4% (0.23-0.63%)	54 (31-89)	15000 (10000-21000)
Pakistan	2010	Female	30 to 34	1.7% (0.83-3%)	98% (97-99%)	0.05 (0.024-0.087)	1% (0.76-1.4%)	0.71% (0.45-1.1%)	90 (56-140)	14000 (9900-20000)
Pakistan	2010	Female	35 to 39	1.9% (0.97-3.1%)	98% (97-99%)	0.057 (0.028-0.096)	1% (0.76-1.3%)	0.88% (0.56-1.3%)	120 (74-180)	13000 (9400-18000)
Pakistan	2010	Female	40 to 44	2% (1.2-3.3%)	98% (97-99%)	0.057 (0.026-0.097)	0.92% (0.67-1.2%)	0.88% (0.57-1.3%)	140 (87-200)	12000 (8400-16000)
Pakistan	2010	Female	45 to 49	2.1% (1.1-3.2%)	98% (97-99%)	0.052 (0.022-0.096)	0.86% (0.63-1.2%)	0.91% (0.59-1.3%)	170 (110-260)	12000 (8100-16000)
Pakistan	2010	Female	50 to 54	2.2% (1.2-3.5%)	98% (96-99%)	0.055 (0.025-0.1)	0.82% (0.57-1.1%)	0.9% (0.54-1.4%)	210 (120-330)	11000 (7200-15000)
Pakistan	2010	Female	55 to 59	2.3% (1.3-3.7%)	98% (96-99%)	0.063 (0.029-0.12)	0.71% (0.46-1%)	0.77% (0.44-1.2%)	200 (110-340)	8600 (5400-13000)
Pakistan	2010	Female	60 to 64	2.4% (1.3-4.1%)	98% (96-99%)	0.063 (0.028-0.12)	0.54% (0.31-0.89%)	0.57% (0.27-1%)	200 (89-360)	6800 (3800-11000)
Pakistan	2010	Female	65 to 69	2.7% (1.5-4.1%)	97% (96-99%)	0.058 (0.026-0.1)	0.41% (0.21-0.68%)	0.45% (0.19-0.79%)	210 (84-370)	5500 (2600-9200)
Pakistan	2010	Female	70 to 74	2.8% (1.7-4.5%)	97% (96-98%)	0.05 (0.022-0.091)	0.3% (0.11-0.56%)	0.34% (0.11-0.67%)	170 (51-350)	3500 (1300-6700)
Pakistan	2010	Female	75 to 79	3% (1.6-4.9%)	97% (95-98%)	0.044 (0.018-0.088)	0.17% (-0.0052-0.36%)	0.18% (-0.036-0.42%)	86 (-18-210)	1400 (-48-3100)
Pakistan	2010	Female	80 to 84	3.2% (1.7-5.4%)	97% (95-98%)	0.043 (0.017-0.083)	0.076% (-0.11-0.24%)	0.069% (-0.16-0.27%)	27 (-61-100)	370 (-530-1100)
Pakistan	2010	Female	85 to 89	3.5% (1.8-6.1%)	97% (94-98%)	0.044 (0.018-0.091)	0.018% (-0.15-0.16%)	-0.0053% (-0.21-0.16%)	-1.3 (-51-40)	39 (-320-340)
Pakistan	2010	Female	90 to 94	4% (2.1-6.9%)	96% (93-98%)	0.045 (0.018-0.095)	-0.031% (-0.22-0.11%)	-0.076% (-0.3-0.095%)	-7.1 (-28-8.7)	-17 (-120-64)
Pakistan	2010	Female	95 plus	4.9% (2.2-9.4%)	95% (91-98%)	0.048 (0.016-0.11)	-0.057% (-0.29-0.11%)	-0.13% (-0.41-0.067%)	-3.3 (-10-1.7)	-5.6 (-28-11)
Pakistan	2010	Male	All Ages	0.83% (0.65-1.1%)	99% (99-99%)	0.00075 (0.0005-0.0011)	0.67% (0.5-0.86%)	0.4% (0.26-0.55%)	2800 (1800-4000)	26000 (190000-340000)
Pakistan	2010	Male	15 to 19	0.52% (0.23-0.91%)	99% (99-100%)	0.00039 (0.00011-0.00089)	0.74% (0.49-1.1%)	0.2% (0.093-0.35%)	28 (13-48)	12000 (7700-18000)
Pakistan	2010	Male	20 to 24	0.78% (0.31-1.4%)	99% (99-100%)	0.00073 (0.00022-0.0016)	1.7% (1.1-2.3%)	0.5% (0.24-0.89%)	95 (45-170)	32000 (20000-46000)
Pakistan	2010	Male	25 to 29	1.1% (0.5-1.9%)	99% (98-99%)	0.0011 (0.00042-0.0022)	2.1% (1.5-2.9%)	0.77% (0.39-1.3%)	140 (67-240)	36000 (24000-51000)
Pakistan	2010	Male	30 to 34	1.3% (0.64-2.4%)	99% (98-99%)	0.0015 (0.00063-0.0028)	2.4% (1.7-3.3%)	1.3% (0.7-2.3%)	240 (120-430)	38000 (26000-54000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	2010	Male	35 to 39	1.6% (0.79-2.5%)	98% (98-99%)	0.0017 (0.00068-0.0032)	2.2% (1.5-2.9%)	1.4% (0.64-2.4%)	270 (120-480)	33000 (22000-46000)
Pakistan	2010	Male	40 to 44	1.6% (0.87-2.6%)	98% (97-99%)	0.0018 (0.00068-0.0033)	1.8% (1.3-2.5%)	1.3% (0.69-2.1%)	290 (150-480)	28000 (19000-39000)
Pakistan	2010	Male	45 to 49	1.5% (0.71-2.4%)	99% (98-99%)	0.0017 (0.00064-0.0035)	1.6% (1.1-2.2%)	1.2% (0.62-2.1%)	330 (160-580)	25000 (17000-35000)
Pakistan	2010	Male	50 to 54	1.4% (0.75-2.4%)	99% (98-99%)	0.0017 (0.00066-0.0033)	1.2% (0.83-1.7%)	0.94% (0.53-1.5%)	290 (160-490)	19000 (12000-27000)
Pakistan	2010	Male	55 to 59	1.5% (0.76-2.4%)	99% (98-99%)	0.0017 (0.00067-0.0033)	0.9% (0.63-1.2%)	0.7% (0.38-1.1%)	230 (120-370)	12000 (8400-18000)
Pakistan	2010	Male	60 to 64	1.4% (0.69-2.4%)	99% (98-99%)	0.0016 (0.00058-0.0032)	0.64% (0.44-0.88%)	0.48% (0.25-0.75%)	210 (110-340)	9500 (6300-13000)
Pakistan	2010	Male	65 to 69	1.2% (0.62-2%)	99% (98-99%)	0.0012 (0.00049-0.0024)	0.48% (0.34-0.64%)	0.37% (0.23-0.57%)	210 (120-310)	7300 (5100-10000)
Pakistan	2010	Male	70 to 74	1.1% (0.59-1.8%)	99% (98-99%)	0.001 (0.00036-0.0021)	0.35% (0.25-0.48%)	0.29% (0.19-0.44%)	170 (110-260)	4600 (3200-6300)
Pakistan	2010	Male	75 to 79	1% (0.49-1.8%)	99% (98-100%)	0.00085 (0.00026-0.0018)	0.28% (0.2-0.38%)	0.25% (0.16-0.38%)	140 (88-210)	2700 (1800-3600)
Pakistan	2010	Male	80 to 84	0.94% (0.45-1.6%)	99% (98-100%)	0.00074 (0.00023-0.0016)	0.2% (0.14-0.28%)	0.18% (0.11-0.27%)	79 (48-120)	1100 (780-1500)
Pakistan	2010	Male	85 to 89	0.95% (0.45-1.7%)	99% (98-100%)	0.00071 (0.00024-0.0016)	0.17% (0.12-0.23%)	0.15% (0.093-0.21%)	42 (26-61)	430 (300-570)
Pakistan	2010	Male	90 to 94	1.1% (0.53-2%)	99% (98-99%)	0.00078 (0.00024-0.0018)	0.16% (0.12-0.22%)	0.12% (0.075-0.19%)	15 (9.2-24)	120 (82-160)
Pakistan	2010	Male	95 plus	1.4% (0.56-2.7%)	99% (97-99%)	0.00094 (0.0002-0.0026)	0.2% (0.14-0.27%)	0.12% (0.072-0.21%)	4 (2.3-6.7)	25 (17-35)
Pakistan	2016	Female	All Ages	1.5% (1.2-1.9%)	99% (98-99%)	0.026 (0.015-0.043)	0.43% (0.33-0.55%)	0.32% (0.2-0.51%)	1900 (1100-3000)	140000 (110000-190000)
Pakistan	2016	Female	15 to 19	1% (0.55-1.7%)	99% (98-99%)	0.0099 (0.0035-0.024)	0.43% (0.26-0.65%)	0.076% (0.033-0.15%)	8 (3.4-16)	6700 (3900-11000)
Pakistan	2016	Female	20 to 24	1.2% (0.57-2%)	99% (98-99%)	0.022 (0.0083-0.044)	0.94% (0.61-1.3%)	0.26% (0.15-0.43%)	35 (19-60)	16000 (9900-25000)
Pakistan	2016	Female	25 to 29	1.4% (0.71-2.5%)	99% (98-99%)	0.036 (0.015-0.072)	1.1% (0.76-1.5%)	0.41% (0.24-0.69%)	57 (32-97)	18000 (12000-25000)
Pakistan	2016	Female	30 to 34	1.7% (0.81-2.9%)	98% (97-99%)	0.048 (0.02-0.094)	1.1% (0.8-1.5%)	0.73% (0.45-1.1%)	96 (57-150)	17000 (11000-24000)
Pakistan	2016	Female	35 to 39	1.9% (0.97-3%)	98% (97-99%)	0.054 (0.022-0.1)	1.1% (0.77-1.4%)	0.91% (0.57-1.4%)	130 (79-210)	16000 (11000-21000)
Pakistan	2016	Female	40 to 44	2% (1.2-3.3%)	98% (97-99%)	0.054 (0.021-0.11)	0.94% (0.67-1.3%)	0.9% (0.57-1.4%)	140 (86-230)	13000 (9000-19000)
Pakistan	2016	Female	45 to 49	2.1% (1.1-3.2%)	98% (97-99%)	0.05 (0.019-0.1)	0.87% (0.63-1.2%)	0.9% (0.56-1.4%)	180 (110-280)	13000 (8800-18000)
Pakistan	2016	Female	50 to 54	2.2% (1.2-3.5%)	98% (97-99%)	0.052 (0.021-0.11)	0.81% (0.56-1.2%)	0.88% (0.54-1.4%)	230 (130-380)	12000 (8200-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	2016	Female	55 to 59	2.3% (1.3-3.8%)	98% (96-99%)	0.06 (0.025-0.12)	0.7% (0.43-1.1%)	0.76% (0.4-1.3%)	250 (120-450)	11000 (6300-18000)
Pakistan	2016	Female	60 to 64	2.5% (1.3-4.1%)	98% (96-99%)	0.061 (0.024-0.13)	0.54% (0.3-0.91%)	0.57% (0.25-1.1%)	210 (90-410)	7500 (3900-13000)
Pakistan	2016	Female	65 to 69	2.7% (1.5-4.2%)	97% (96-99%)	0.056 (0.022-0.11)	0.41% (0.2-0.71%)	0.46% (0.2-0.85%)	210 (87-400)	5600 (2700-9900)
Pakistan	2016	Female	70 to 74	2.8% (1.6-4.6%)	97% (95-98%)	0.048 (0.019-0.1)	0.31% (0.12-0.58%)	0.35% (0.11-0.69%)	200 (64-400)	4000 (1500-7700)
Pakistan	2016	Female	75 to 79	2.9% (1.6-4.9%)	97% (95-98%)	0.043 (0.016-0.095)	0.17% (0.013-0.39%)	0.19% (-0.013-0.47%)	100 (-7.2-250)	1700 (130-3700)
Pakistan	2016	Female	80 to 84	3.2% (1.7-5.3%)	97% (95-98%)	0.042 (0.016-0.091)	0.083% (-0.091-0.27%)	0.076% (-0.13-0.3%)	33 (-62-130)	450 (-500-1500)
Pakistan	2016	Female	85 to 89	3.5% (1.8-6%)	96% (94-98%)	0.043 (0.015-0.096)	0.024% (-0.15-0.17%)	-0.000092% (-0.21-0.18%)	0.036 (-62-56)	62 (-390-470)
Pakistan	2016	Female	90 to 94	4.1% (2.2-6.9%)	96% (93-98%)	0.044 (0.016-0.11)	-0.025% (-0.22-0.13%)	-0.071% (-0.3-0.1%)	-8.7 (-37-13)	-19 (-160-92)
Pakistan	2016	Female	95 plus	5% (2.2-9.6%)	95% (90-98%)	0.047 (0.015-0.11)	-0.05% (-0.29-0.13%)	-0.12% (-0.43-0.088%)	-4.1 (-14-2.9)	-6.5 (-38-16)
Pakistan	2016	Male	All Ages	0.86% (0.66-1.1%)	99% (99-99%)	0.00076 (0.0004-0.0013)	0.84% (0.63-1.1%)	0.47% (0.31-0.66%)	3400 (2100-4900)	320000 (230000-410000)
Pakistan	2016	Male	15 to 19	0.52% (0.24-0.94%)	99% (99-100%)	0.00037 (0.000096-0.00098)	0.82% (0.53-1.2%)	0.22% (0.1-0.39%)	28 (13-51)	13000 (8000-20000)
Pakistan	2016	Male	20 to 24	0.79% (0.31-1.4%)	99% (99-100%)	0.00071 (0.00019-0.0017)	1.9% (1.2-2.7%)	0.57% (0.28-0.97%)	100 (50-180)	36000 (22000-52000)
Pakistan	2016	Male	25 to 29	1.1% (0.51-1.9%)	99% (98-99%)	0.0011 (0.00035-0.0023)	2.3% (1.6-3.1%)	0.85% (0.42-1.5%)	170 (81-310)	45000 (30000-64000)
Pakistan	2016	Male	30 to 34	1.4% (0.65-2.5%)	99% (98-99%)	0.0014 (0.00053-0.0031)	2.6% (1.9-3.6%)	1.5% (0.79-2.5%)	300 (150-530)	47000 (32000-67000)
Pakistan	2016	Male	35 to 39	1.6% (0.78-2.5%)	98% (97-99%)	0.0017 (0.0006-0.0035)	2.3% (1.7-3.2%)	1.5% (0.69-2.6%)	340 (150-610)	41000 (27000-57000)
Pakistan	2016	Male	40 to 44	1.6% (0.89-2.5%)	98% (97-99%)	0.0017 (0.00059-0.0036)	2% (1.4-2.7%)	1.4% (0.75-2.3%)	340 (170-570)	33000 (22000-47000)
Pakistan	2016	Male	45 to 49	1.5% (0.7-2.3%)	99% (98-99%)	0.0016 (0.00055-0.0037)	1.7% (1.2-2.4%)	1.3% (0.65-2.3%)	390 (180-700)	29000 (20000-42000)
Pakistan	2016	Male	50 to 54	1.4% (0.75-2.4%)	99% (98-99%)	0.0016 (0.00054-0.0036)	1.3% (0.89-1.8%)	1% (0.57-1.6%)	370 (200-630)	23000 (15000-33000)
Pakistan	2016	Male	55 to 59	1.5% (0.75-2.4%)	99% (98-99%)	0.0017 (0.00057-0.0036)	0.97% (0.66-1.3%)	0.75% (0.41-1.2%)	310 (170-530)	17000 (12000-24000)
Pakistan	2016	Male	60 to 64	1.4% (0.7-2.5%)	99% (98-99%)	0.0015 (0.00051-0.0035)	0.69% (0.48-0.95%)	0.51% (0.28-0.82%)	240 (120-380)	11000 (7200-16000)
Pakistan	2016	Male	65 to 69	1.3% (0.63-2.1%)	99% (98-99%)	0.0012 (0.00042-0.0027)	0.52% (0.38-0.69%)	0.41% (0.26-0.6%)	220 (130-340)	7900 (5600-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Pakistan	2016	Male	70 to 74	1.1% (0.59-1.8%)	99% (98-99%)	0.00097 (0.00033-0.0023)	0.38% (0.27-0.52%)	0.31% (0.2-0.48%)	200 (120-320)	5500 (3800-7600)
Pakistan	2016	Male	75 to 79	1% (0.5-1.8%)	99% (98-99%)	0.00082 (0.00024-0.002)	0.31% (0.22-0.41%)	0.27% (0.17-0.4%)	170 (100-260)	3300 (2300-4500)
Pakistan	2016	Male	80 to 84	0.95% (0.46-1.6%)	99% (98-100%)	0.00072 (0.00022-0.0017)	0.22% (0.15-0.3%)	0.19% (0.12-0.29%)	97 (62-150)	1400 (950-1900)
Pakistan	2016	Male	85 to 89	0.96% (0.46-1.8%)	99% (98-100%)	0.0007 (0.00021-0.0017)	0.18% (0.13-0.24%)	0.16% (0.1-0.22%)	51 (33-74)	520 (380-700)
Pakistan	2016	Male	90 to 94	1.1% (0.54-2%)	99% (98-99%)	0.00076 (0.0002-0.0019)	0.17% (0.12-0.23%)	0.13% (0.085-0.21%)	19 (12-30)	150 (100-200)
Pakistan	2016	Male	95 plus	1.4% (0.59-2.6%)	99% (97-99%)	0.0009 (0.00018-0.0025)	0.21% (0.15-0.29%)	0.14% (0.077-0.22%)	5.2 (2.9-8.5)	32 (22-44)
Angola	1990	Female	All Ages	22% (17-27%)	78% (73-83%)	0.27 (0.21-0.34)	0.54% (0.19-0.96%)	0.8% (0.066-1.6%)	660 (56-1400)	31000 (10000-56000)
Angola	1990	Female	15 to 19	22% (13-35%)	78% (65-87%)	0.24 (0.12-0.42)	0.74% (0.21-1.5%)	0.68% (-0.12-1.7%)	9.5 (-1.5-25)	1100 (310-2300)
Angola	1990	Female	20 to 24	31% (18-46%)	69% (54-82%)	0.45 (0.23-0.75)	1.7% (0.61-3.3%)	1.6% (0.074-3.8%)	27 (1.2-62)	2700 (960-5300)
Angola	1990	Female	25 to 29	35% (20-51%)	65% (49-80%)	0.6 (0.35-0.93)	2.5% (0.96-4.8%)	2.8% (0.58-5.9%)	46 (9.3-100)	3700 (1400-7200)
Angola	1990	Female	30 to 34	37% (22-54%)	63% (46-78%)	0.67 (0.39-1)	3% (1.1-5.4%)	3.6% (0.92-6.9%)	61 (15-120)	4100 (1500-7400)
Angola	1990	Female	35 to 39	39% (24-56%)	61% (44-76%)	0.66 (0.38-1.1)	3.1% (0.92-5.9%)	3.7% (0.81-7.6%)	68 (14-140)	3900 (1100-7400)
Angola	1990	Female	40 to 44	39% (25-57%)	61% (43-75%)	0.65 (0.33-1.1)	3% (0.88-5.9%)	3.8% (0.82-7.7%)	63 (14-130)	3200 (910-6300)
Angola	1990	Female	45 to 49	38% (23-56%)	62% (44-77%)	0.62 (0.31-1.1)	2.9% (0.71-6.1%)	3.7% (0.69-7.8%)	70 (13-150)	3000 (690-6400)
Angola	1990	Female	50 to 54	35% (21-53%)	65% (47-79%)	0.56 (0.29-0.96)	2.9% (0.44-6%)	3.5% (0.37-7.8%)	83 (8.5-190)	3100 (460-6900)
Angola	1990	Female	55 to 59	33% (20-49%)	67% (51-80%)	0.51 (0.25-0.91)	2.5% (0.25-5.7%)	3% (0.21-6.8%)	82 (5.7-200)	2600 (270-6300)
Angola	1990	Female	60 to 64	32% (20-49%)	68% (51-80%)	0.49 (0.24-0.89)	1.8% (-0.085-4.7%)	2.1% (-0.14-5.5%)	68 (-4.5-180)	1800 (-93-4800)
Angola	1990	Female	65 to 69	34% (22-50%)	66% (50-78%)	0.48 (0.25-0.81)	1.2% (-0.5-3.5%)	1.4% (-0.56-4.1%)	53 (-22-150)	1200 (-530-3300)
Angola	1990	Female	70 to 74	34% (22-49%)	66% (51-78%)	0.43 (0.21-0.78)	0.62% (-0.96-2.6%)	0.73% (-1.1-3.1%)	27 (-43-120)	470 (-760-2000)
Angola	1990	Female	75 to 79	31% (19-46%)	69% (54-81%)	0.35 (0.16-0.64)	0.19% (-1-1.8%)	0.23% (-1.2-2%)	6.8 (-37-59)	90 (-510-810)
Angola	1990	Female	80 to 84	28% (17-43%)	72% (57-83%)	0.29 (0.13-0.52)	0.029% (-0.94-1.3%)	0.04% (-1.1-1.5%)	0.7 (-20-25)	6.3 (-210-250)
Angola	1990	Female	85 to 89	27% (16-40%)	73% (60-84%)	0.26 (0.12-0.47)	-0.044% (-0.9-0.89%)	-0.05% (-1-1.1%)	-0.46 (-9.6-9.7)	-3.3 (-70-66)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	1990	Female	90 to 94	28% (18-42%)	72% (58-82%)	0.25 (0.12-0.47)	-0.053% (-0.8-0.83%)	-0.072% (-0.98-0.96%)	-0.22 (-3-2.9)	-0.94 (-15-14)
Angola	1990	Female	95 plus	32% (18-49%)	68% (51-82%)	0.25 (0.1-0.51)	0.034% (-0.69-0.86%)	0.0049% (-0.86-0.98%)	0.0043 (-0.73-0.85)	0.12 (-2.3-2.9)
Angola	1990	Male	All Ages	28% (23-36%)	72% (64-77%)	0.34 (0.27-0.42)	1.1% (0.45-1.7%)	1.6% (0.46-2.8%)	1500 (420-2600)	69000 (29000-110000)
Angola	1990	Male	15 to 19	24% (14-39%)	76% (61-86%)	0.26 (0.12-0.47)	1.3% (0.44-2.5%)	1.2% (0.13-2.8%)	23 (2.3-55)	2400 (780-4400)
Angola	1990	Male	20 to 24	37% (22-53%)	63% (47-78%)	0.53 (0.29-0.88)	3.2% (1.1-5.8%)	3.1% (0.49-6.3%)	76 (13-160)	6600 (2300-12000)
Angola	1990	Male	25 to 29	44% (27-61%)	56% (39-73%)	0.76 (0.45-1.1)	4.9% (1.7-8.6%)	5% (1.1-9.8%)	120 (28-230)	8800 (3000-15000)
Angola	1990	Male	30 to 34	46% (28-65%)	54% (35-72%)	0.85 (0.51-1.3)	5.7% (2.1-9.6%)	6.2% (1.7-11%)	140 (39-260)	9300 (3400-16000)
Angola	1990	Male	35 to 39	46% (29-64%)	54% (36-71%)	0.84 (0.49-1.3)	5.8% (2.1-10%)	6.5% (1.8-12%)	160 (43-290)	8900 (3000-15000)
Angola	1990	Male	40 to 44	48% (32-67%)	52% (33-68%)	0.86 (0.45-1.4)	6% (2-11%)	6.8% (1.9-13%)	160 (41-300)	7700 (2400-14000)
Angola	1990	Male	45 to 49	51% (33-69%)	49% (31-67%)	0.91 (0.46-1.5)	6% (1.7-11%)	6.9% (1.7-13%)	180 (44-350)	7600 (2200-14000)
Angola	1990	Male	50 to 54	51% (33-70%)	49% (30-67%)	0.88 (0.47-1.5)	5.5% (1.1-10%)	6.3% (1-12%)	180 (27-360)	6700 (1400-13000)
Angola	1990	Male	55 to 59	49% (32-68%)	51% (32-68%)	0.78 (0.41-1.3)	4.3% (0.71-8.5%)	4.9% (0.62-10%)	150 (19-320)	4900 (790-10000)
Angola	1990	Male	60 to 64	48% (30-68%)	52% (32-70%)	0.68 (0.34-1.2)	3.2% (0.44-6.7%)	3.6% (0.33-7.7%)	120 (13-260)	3300 (470-7000)
Angola	1990	Male	65 to 69	48% (31-65%)	52% (35-69%)	0.61 (0.32-1)	2.2% (-0.17-5.2%)	2.5% (-0.35-6%)	82 (-12-200)	1900 (-140-4500)
Angola	1990	Male	70 to 74	49% (32-66%)	51% (34-68%)	0.53 (0.28-0.93)	1.4% (-0.7-3.7%)	1.6% (-0.87-4.2%)	48 (-27-130)	880 (-440-2400)
Angola	1990	Male	75 to 79	47% (31-65%)	53% (35-69%)	0.44 (0.2-0.8)	0.92% (-0.73-2.9%)	1% (-0.95-3.4%)	23 (-22-78)	330 (-280-1100)
Angola	1990	Male	80 to 84	42% (25-60%)	58% (40-75%)	0.33 (0.16-0.6)	0.56% (-0.69-1.9%)	0.59% (-0.88-2.2%)	6.9 (-11-26)	77 (-100-270)
Angola	1990	Male	85 to 89	36% (22-53%)	64% (47-78%)	0.25 (0.12-0.46)	0.44% (-0.54-1.5%)	0.44% (-0.72-1.7%)	2.4 (-4-9)	20 (-26-67)
Angola	1990	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.22 (0.099-0.41)	0.26% (-0.47-1%)	0.21% (-0.67-1.1%)	0.37 (-1.2-2)	2.7 (-5-10)
Angola	1990	Male	95 plus	39% (21-59%)	61% (41-79%)	0.2 (0.079-0.41)	0.33% (-0.61-1.4%)	0.26% (-0.88-1.5%)	0.11 (-0.39-0.66)	0.58 (-1-2.4)
Angola	1995	Female	All Ages	22% (17-27%)	78% (73-83%)	0.29 (0.22-0.36)	0.65% (0.24-1.1%)	0.96% (0.16-1.9%)	860 (140-1700)	40000 (14000-67000)
Angola	1995	Female	15 to 19	22% (14-34%)	78% (66-86%)	0.25 (0.12-0.44)	0.8% (0.25-1.5%)	0.77% (-0.061-1.9%)	12 (-0.98-29)	1400 (430-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	1995	Female	20 to 24	31% (19-46%)	69% (54-81%)	0.48 (0.25-0.77)	1.8% (0.74-3.3%)	1.8% (0.22-3.9%)	34 (4.4-76)	3300 (1300-6200)
Angola	1995	Female	25 to 29	35% (21-52%)	65% (48-79%)	0.64 (0.36-0.98)	2.6% (1.1-4.6%)	2.8% (0.78-5.6%)	56 (15-120)	4400 (1800-8200)
Angola	1995	Female	30 to 34	37% (22-55%)	63% (45-78%)	0.71 (0.4-1.1)	3.1% (1.3-5.4%)	3.6% (1.1-6.9%)	75 (23-140)	5000 (2100-8900)
Angola	1995	Female	35 to 39	39% (24-56%)	61% (44-76%)	0.71 (0.39-1.2)	3.2% (1.1-6%)	3.9% (0.99-7.7%)	88 (21-180)	5000 (1700-9600)
Angola	1995	Female	40 to 44	39% (25-57%)	61% (43-75%)	0.7 (0.36-1.1)	3.3% (1-6.3%)	4% (0.98-8%)	96 (24-200)	4800 (1500-9200)
Angola	1995	Female	45 to 49	38% (23-56%)	62% (44-77%)	0.66 (0.34-1.1)	3.2% (0.89-6.5%)	4% (0.93-8.2%)	91 (20-200)	3900 (1100-8100)
Angola	1995	Female	50 to 54	36% (21-54%)	64% (46-79%)	0.6 (0.3-1)	3.1% (0.65-6.5%)	3.8% (0.68-8%)	110 (17-240)	4000 (710-8700)
Angola	1995	Female	55 to 59	33% (20-50%)	67% (50-80%)	0.54 (0.27-0.96)	2.7% (0.31-5.9%)	3.2% (0.31-7.2%)	110 (9.5-250)	3400 (390-7700)
Angola	1995	Female	60 to 64	32% (20-49%)	68% (51-80%)	0.52 (0.24-0.93)	1.9% (-0.014-4.9%)	2.2% (-0.041-5.8%)	86 (-1.8-230)	2300 (-16-6100)
Angola	1995	Female	65 to 69	34% (22-49%)	66% (51-78%)	0.51 (0.25-0.9)	1.3% (-0.46-3.9%)	1.5% (-0.51-4.6%)	67 (-25-190)	1500 (-520-4300)
Angola	1995	Female	70 to 74	34% (22-50%)	66% (50-78%)	0.46 (0.22-0.84)	0.68% (-0.97-2.9%)	0.8% (-1.1-3.4%)	35 (-49-140)	620 (-880-2500)
Angola	1995	Female	75 to 79	31% (20-47%)	69% (53-80%)	0.37 (0.17-0.69)	0.21% (-1.1-1.7%)	0.25% (-1.2-2%)	8.9 (-43-72)	120 (-590-990)
Angola	1995	Female	80 to 84	28% (17-44%)	72% (56-83%)	0.31 (0.14-0.56)	0.049% (-0.94-1.3%)	0.062% (-1.1-1.5%)	1.3 (-25-31)	13 (-250-310)
Angola	1995	Female	85 to 89	27% (16-39%)	73% (61-84%)	0.27 (0.13-0.5)	-0.035% (-0.91-0.85%)	-0.041% (-1.1-1%)	-0.46 (-11-11)	-3.1 (-80-78)
Angola	1995	Female	90 to 94	28% (17-42%)	72% (58-83%)	0.26 (0.12-0.48)	-0.049% (-0.79-0.8%)	-0.069% (-0.96-0.96%)	-0.25 (-3.6-3.5)	-1 (-18-17)
Angola	1995	Female	95 plus	32% (18-49%)	68% (51-82%)	0.27 (0.1-0.55)	0.046% (-0.67-0.88%)	0.018% (-0.85-1.1%)	0.018 (-0.86-1)	0.19 (-2.7-3.6)
Angola	1995	Male	All Ages	29% (23-36%)	71% (64-77%)	0.36 (0.28-0.45)	1.2% (0.57-1.8%)	1.8% (0.67-2.9%)	1700 (620-2900)	81000 (38000-130000)
Angola	1995	Male	15 to 19	24% (14-39%)	76% (61-86%)	0.27 (0.13-0.49)	1.5% (0.53-2.7%)	1.5% (0.26-3.2%)	28 (5.2-61)	2800 (1000-5300)
Angola	1995	Male	20 to 24	37% (23-54%)	63% (46-77%)	0.57 (0.3-0.91)	3.8% (1.6-6.4%)	3.8% (0.96-7.2%)	91 (23-180)	7900 (3400-14000)
Angola	1995	Male	25 to 29	45% (27-62%)	55% (38-73%)	0.82 (0.5-1.2)	5.4% (2.2-8.9%)	5.8% (1.6-11%)	140 (38-260)	10000 (4100-18000)
Angola	1995	Male	30 to 34	47% (29-65%)	53% (35-71%)	0.92 (0.55-1.4)	6% (2.6-9.9%)	6.7% (2.4-12%)	170 (55-310)	11000 (4400-19000)
Angola	1995	Male	35 to 39	47% (30-65%)	53% (35-70%)	0.9 (0.52-1.4)	6% (2.3-10%)	6.8% (2.1-12%)	180 (55-330)	10000 (3800-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	1995	Male	40 to 44	49% (32-67%)	51% (33-68%)	0.92 (0.5-1.5)	6.2% (2-11%)	7.2% (2-13%)	200 (57-380)	9900 (3200-18000)
Angola	1995	Male	45 to 49	52% (32-69%)	48% (31-68%)	0.97 (0.51-1.6)	6.2% (2.1-11%)	7.2% (2.2-13%)	190 (55-370)	8300 (2700-15000)
Angola	1995	Male	50 to 54	52% (34-70%)	48% (30-66%)	0.94 (0.5-1.6)	5.7% (1.5-11%)	6.7% (1.5-13%)	200 (42-390)	7400 (1800-14000)
Angola	1995	Male	55 to 59	50% (32-68%)	50% (32-68%)	0.83 (0.44-1.4)	4.5% (1.2-8.7%)	5.2% (1.2-10%)	170 (36-350)	5500 (1400-11000)
Angola	1995	Male	60 to 64	48% (30-68%)	52% (32-70%)	0.72 (0.36-1.2)	3.3% (0.58-6.8%)	3.8% (0.49-7.9%)	130 (18-280)	3600 (630-7600)
Angola	1995	Male	65 to 69	49% (31-66%)	51% (34-69%)	0.65 (0.33-1)	2.3% (-0.027-5.1%)	2.7% (-0.13-5.9%)	93 (-4.4-210)	2100 (-22-4800)
Angola	1995	Male	70 to 74	49% (32-66%)	51% (34-68%)	0.57 (0.28-0.98)	1.5% (-0.51-3.8%)	1.7% (-0.66-4.4%)	55 (-21-150)	1000 (-340-2600)
Angola	1995	Male	75 to 79	48% (32-66%)	52% (34-68%)	0.47 (0.22-0.84)	0.99% (-0.71-2.9%)	1.1% (-0.87-3.4%)	27 (-23-83)	390 (-280-1200)
Angola	1995	Male	80 to 84	42% (25-61%)	58% (39-75%)	0.36 (0.17-0.65)	0.61% (-0.64-2%)	0.65% (-0.82-2.2%)	8.6 (-11-30)	95 (-110-310)
Angola	1995	Male	85 to 89	37% (22-54%)	63% (46-78%)	0.27 (0.13-0.49)	0.47% (-0.45-1.5%)	0.48% (-0.61-1.7%)	3.1 (-4-11)	26 (-25-81)
Angola	1995	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.23 (0.11-0.43)	0.27% (-0.45-1%)	0.23% (-0.66-1.1%)	0.49 (-1.4-2.3)	3.4 (-5.7-13)
Angola	1995	Male	95 plus	40% (22-60%)	60% (40-78%)	0.22 (0.084-0.45)	0.35% (-0.6-1.4%)	0.29% (-0.86-1.5%)	0.15 (-0.45-0.79)	0.73 (-1.2-2.8)
Angola	2000	Female	All Ages	21% (17-27%)	79% (73-83%)	0.37 (0.29-0.47)	0.88% (0.44-1.4%)	1.4% (0.49-2.5%)	1400 (470-2500)	59000 (28000-94000)
Angola	2000	Female	15 to 19	21% (13-34%)	79% (66-87%)	0.32 (0.15-0.56)	0.96% (0.42-1.8%)	1% (0.19-2.3%)	17 (3.2-41)	1900 (790-3700)
Angola	2000	Female	20 to 24	30% (18-45%)	70% (55-82%)	0.62 (0.32-1)	2.2% (1-3.7%)	2.2% (0.62-4.5%)	49 (13-100)	4600 (2100-8200)
Angola	2000	Female	25 to 29	34% (20-51%)	66% (49-80%)	0.83 (0.48-1.3)	2.9% (1.5-5%)	3.3% (1.3-6.1%)	78 (32-150)	6000 (2900-11000)
Angola	2000	Female	30 to 34	37% (22-54%)	63% (46-78%)	0.93 (0.52-1.4)	3.6% (1.8-5.9%)	4.2% (1.8-7.3%)	100 (42-190)	6800 (3200-12000)
Angola	2000	Female	35 to 39	38% (24-55%)	62% (45-76%)	0.92 (0.5-1.5)	3.8% (1.7-6.6%)	4.6% (1.9-8.3%)	120 (46-230)	6900 (2900-12000)
Angola	2000	Female	40 to 44	38% (24-57%)	62% (43-76%)	0.91 (0.46-1.5)	4% (1.7-7%)	5% (2-8.9%)	140 (51-260)	6700 (2800-12000)
Angola	2000	Female	45 to 49	37% (22-55%)	63% (45-78%)	0.86 (0.44-1.5)	4.2% (1.6-7.8%)	5.2% (1.8-9.8%)	160 (48-310)	6700 (2200-13000)
Angola	2000	Female	50 to 54	35% (20-53%)	65% (47-80%)	0.78 (0.39-1.3)	4.2% (1.4-8.3%)	5.2% (1.6-10%)	160 (45-320)	5900 (1800-12000)
Angola	2000	Female	55 to 59	32% (20-49%)	68% (51-80%)	0.71 (0.34-1.3)	3.8% (1.1-7.7%)	4.6% (1.3-9.3%)	160 (45-350)	5200 (1500-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	2000	Female	60 to 64	31% (19-48%)	69% (52-81%)	0.67 (0.31-1.3)	2.8% (0.59-6.2%)	3.3% (0.64-7.3%)	140 (28-310)	3700 (820-8400)
Angola	2000	Female	65 to 69	33% (22-48%)	67% (52-78%)	0.66 (0.33-1.1)	2.2% (0.03-5.1%)	2.5% (0.019-6%)	120 (0.86-280)	2600 (34-6200)
Angola	2000	Female	70 to 74	33% (22-48%)	67% (52-78%)	0.6 (0.28-1.1)	1.4% (-0.5-4.1%)	1.6% (-0.56-4.8%)	79 (-28-230)	1400 (-500-4000)
Angola	2000	Female	75 to 79	30% (19-45%)	70% (55-81%)	0.48 (0.22-0.88)	0.75% (-0.75-2.9%)	0.87% (-0.86-3.4%)	35 (-35-140)	480 (-480-1900)
Angola	2000	Female	80 to 84	27% (16-42%)	73% (58-84%)	0.4 (0.18-0.72)	0.45% (-0.66-2%)	0.52% (-0.78-2.3%)	14 (-20-61)	140 (-200-630)
Angola	2000	Female	85 to 89	26% (16-39%)	74% (61-84%)	0.35 (0.16-0.65)	0.27% (-0.7-1.5%)	0.31% (-0.83-1.8%)	3.8 (-9.9-22)	28 (-71-160)
Angola	2000	Female	90 to 94	28% (17-41%)	72% (59-83%)	0.34 (0.15-0.63)	0.19% (-0.58-1.3%)	0.21% (-0.72-1.6%)	0.88 (-3-6.5)	4.7 (-14-33)
Angola	2000	Female	95 plus	31% (18-49%)	69% (51-82%)	0.35 (0.13-0.71)	0.24% (-0.54-1.3%)	0.24% (-0.7-1.5%)	0.28 (-0.77-1.7)	1.1 (-2.4-5.9)
Angola	2000	Male	All Ages	28% (22-36%)	72% (64-78%)	0.48 (0.37-0.61)	1.5% (0.85-2.2%)	2.3% (1.2-3.6%)	2300 (1100-3700)	110000 (60000-160000)
Angola	2000	Male	15 to 19	24% (14-39%)	76% (61-86%)	0.37 (0.17-0.66)	1.7% (0.78-3.1%)	1.8% (0.52-3.7%)	35 (10-76)	3500 (1600-6500)
Angola	2000	Male	20 to 24	37% (23-54%)	63% (46-77%)	0.77 (0.41-1.2)	4.4% (2.2-7.4%)	4.6% (1.6-8.5%)	120 (42-230)	10000 (4800-18000)
Angola	2000	Male	25 to 29	44% (27-61%)	56% (39-73%)	1.1 (0.66-1.7)	6.3% (3.3-10%)	7% (3-12%)	190 (76-320)	13000 (6400-22000)
Angola	2000	Male	30 to 34	46% (28-64%)	54% (36-72%)	1.2 (0.72-1.8)	7.1% (3.8-11%)	8.1% (3.9-13%)	220 (99-380)	14000 (7300-23000)
Angola	2000	Male	35 to 39	46% (30-64%)	54% (36-70%)	1.2 (0.68-1.9)	7% (3.5-11%)	8.2% (3.7-13%)	230 (100-400)	13000 (6300-21000)
Angola	2000	Male	40 to 44	48% (31-67%)	52% (33-69%)	1.2 (0.65-2)	7.3% (3.3-12%)	8.7% (3.5-15%)	260 (100-450)	12000 (5500-21000)
Angola	2000	Male	45 to 49	51% (33-68%)	49% (32-67%)	1.3 (0.71-2.2)	7.6% (3.4-12%)	9% (3.8-15%)	280 (110-500)	12000 (5100-21000)
Angola	2000	Male	50 to 54	51% (33-69%)	49% (31-67%)	1.3 (0.67-2.2)	7.3% (3.2-13%)	8.8% (3.7-15%)	250 (92-460)	9400 (3700-17000)
Angola	2000	Male	55 to 59	49% (32-68%)	51% (32-68%)	1.1 (0.59-1.9)	6.1% (2.4-11%)	7.3% (2.7-13%)	220 (79-400)	7200 (2700-13000)
Angola	2000	Male	60 to 64	48% (30-67%)	52% (33-70%)	0.99 (0.49-1.8)	4.5% (1.7-8.4%)	5.3% (1.8-10%)	180 (58-360)	5000 (1700-9700)
Angola	2000	Male	65 to 69	49% (32-66%)	51% (34-68%)	0.89 (0.45-1.5)	3.4% (0.89-6.7%)	4% (0.95-8%)	140 (30-280)	3200 (770-6300)
Angola	2000	Male	70 to 74	49% (32-66%)	51% (34-68%)	0.78 (0.39-1.4)	2.3% (0.095-5.2%)	2.6% (0.021-6%)	89 (0.68-200)	1600 (64-3600)
Angola	2000	Male	75 to 79	48% (31-66%)	52% (34-69%)	0.64 (0.3-1.2)	1.6% (-0.2-4%)	1.8% (-0.29-4.7%)	47 (-7.4-120)	670 (-87-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	2000	Male	80 to 84	42% (25-61%)	58% (39-75%)	0.49 (0.23-0.87)	0.99% (-0.27-2.7%)	1.1% (-0.38-3.1%)	16 (-5.5-47)	180 (-47-480)
Angola	2000	Male	85 to 89	37% (22-54%)	63% (46-78%)	0.37 (0.17-0.67)	0.71% (-0.18-1.9%)	0.76% (-0.31-2.2%)	5.5 (-2.2-16)	44 (-11-120)
Angola	2000	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.32 (0.15-0.58)	0.42% (-0.32-1.3%)	0.4% (-0.51-1.5%)	1 (-1.2-3.7)	6.2 (-4.7-19)
Angola	2000	Male	95 plus	39% (21-61%)	61% (39-79%)	0.3 (0.11-0.61)	0.53% (-0.47-1.6%)	0.48% (-0.72-1.8%)	0.3 (-0.45-1.1)	1.3 (-1.1-3.9)
Angola	2005	Female	All Ages	21% (17-26%)	79% (74-83%)	0.54 (0.41-0.7)	1.4% (0.82-2.1%)	2.4% (1.2-3.8%)	2400 (1100-3900)	94000 (55000-140000)
Angola	2005	Female	15 to 19	21% (13-33%)	79% (67-87%)	0.47 (0.22-0.83)	1.4% (0.66-2.5%)	1.7% (0.56-3.6%)	30 (9.8-61)	2900 (1300-5200)
Angola	2005	Female	20 to 24	30% (18-44%)	70% (56-82%)	0.91 (0.47-1.5)	3% (1.6-4.9%)	3.4% (1.4-6.1%)	77 (31-140)	6800 (3500-11000)
Angola	2005	Female	25 to 29	34% (20-50%)	66% (50-80%)	1.2 (0.7-1.9)	3.8% (2.1-5.9%)	4.4% (2.1-7.5%)	120 (57-200)	8800 (5000-14000)
Angola	2005	Female	30 to 34	36% (21-53%)	64% (47-79%)	1.4 (0.76-2.1)	4.5% (2.5-7.1%)	5.4% (2.8-9.1%)	160 (77-260)	9900 (5500-16000)
Angola	2005	Female	35 to 39	38% (23-56%)	62% (44-77%)	1.4 (0.72-2.2)	4.9% (2.7-7.5%)	6% (3.1-9.4%)	180 (85-310)	9900 (5100-17000)
Angola	2005	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.3 (0.71-2.2)	5.4% (2.7-8.8%)	6.7% (3.3-11%)	210 (94-370)	10000 (4900-17000)
Angola	2005	Female	45 to 49	37% (22-54%)	63% (46-78%)	1.3 (0.65-2.2)	5.8% (2.8-9.8%)	7.4% (3.5-13%)	240 (110-450)	10000 (4700-18000)
Angola	2005	Female	50 to 54	34% (20-51%)	66% (49-80%)	1.2 (0.58-2)	6.1% (2.7-11%)	7.6% (3.1-13%)	290 (110-540)	11000 (4300-20000)
Angola	2005	Female	55 to 59	32% (20-49%)	68% (51-80%)	1 (0.51-1.8)	5.7% (2.4-9.8%)	6.9% (2.9-12%)	260 (100-480)	8200 (3400-15000)
Angola	2005	Female	60 to 64	31% (19-48%)	69% (52-81%)	1 (0.45-1.8)	4.5% (1.6-8.5%)	5.3% (1.9-10%)	230 (79-460)	6300 (2200-12000)
Angola	2005	Female	65 to 69	33% (21-48%)	67% (52-79%)	0.98 (0.48-1.7)	3.8% (1.1-7.5%)	4.5% (1.3-8.9%)	230 (62-450)	5000 (1400-10000)
Angola	2005	Female	70 to 74	32% (21-47%)	68% (53-79%)	0.88 (0.42-1.6)	2.9% (0.41-6.3%)	3.3% (0.46-7.5%)	170 (22-390)	3100 (390-6900)
Angola	2005	Female	75 to 79	30% (18-46%)	70% (54-82%)	0.71 (0.33-1.3)	1.9% (-0.19-5%)	2.2% (-0.22-5.5%)	98 (-9.7-260)	1400 (-130-3600)
Angola	2005	Female	80 to 84	27% (16-43%)	73% (57-84%)	0.59 (0.28-1)	1.3% (-0.23-3.4%)	1.5% (-0.26-3.9%)	47 (-8.1-120)	480 (-79-1200)
Angola	2005	Female	85 to 89	26% (15-39%)	74% (61-85%)	0.52 (0.24-0.94)	0.94% (-0.24-2.6%)	1.1% (-0.29-3.1%)	16 (-4.2-44)	110 (-29-320)
Angola	2005	Female	90 to 94	27% (16-40%)	73% (60-84%)	0.5 (0.22-0.93)	0.74% (-0.27-2.4%)	0.86% (-0.35-2.8%)	4 (-1.6-13)	21 (-7.4-66)
Angola	2005	Female	95 plus	31% (17-48%)	69% (52-83%)	0.51 (0.19-1.1)	0.69% (-0.28-2.4%)	0.76% (-0.39-2.8%)	0.97 (-0.48-3.5)	3.5 (-1.3-12)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	2005	Male	All Ages	28% (22-35%)	72% (65-78%)	0.72 (0.54-0.91)	2.2% (1.4-3.1%)	3.5% (2.1-5.1%)	3400 (2000-5100)	150000 (100000-220000)
Angola	2005	Male	15 to 19	24% (14-39%)	76% (61-86%)	0.55 (0.26-1)	2.5% (1.3-4.4%)	3% (1.3-5.8%)	56 (23-110)	5200 (2700-9000)
Angola	2005	Male	20 to 24	37% (22-54%)	63% (46-78%)	1.1 (0.61-1.8)	6.5% (3.8-10%)	7.5% (3.8-12%)	190 (89-320)	15000 (8400-24000)
Angola	2005	Male	25 to 29	44% (27-62%)	56% (38-73%)	1.6 (0.96-2.5)	8.6% (5.2-13%)	10% (5.5-15%)	280 (140-460)	20000 (11000-31000)
Angola	2005	Male	30 to 34	46% (29-64%)	54% (36-71%)	1.8 (1-2.7)	9.2% (5.5-13%)	11% (6.1-16%)	320 (160-500)	20000 (11000-30000)
Angola	2005	Male	35 to 39	46% (30-63%)	54% (37-70%)	1.8 (0.97-2.8)	9.1% (5.5-14%)	11% (6.2-17%)	330 (170-520)	18000 (10000-28000)
Angola	2005	Male	40 to 44	48% (32-66%)	52% (34-68%)	1.8 (0.96-3)	9.4% (5.4-14%)	12% (6.4-17%)	360 (190-590)	17000 (9600-28000)
Angola	2005	Male	45 to 49	51% (33-69%)	49% (31-67%)	2 (1.1-3.2)	10% (5.7-15%)	12% (6.8-19%)	390 (200-640)	17000 (8800-27000)
Angola	2005	Male	50 to 54	52% (33-70%)	48% (30-67%)	1.9 (1-3.2)	9.9% (5.2-16%)	12% (6.3-19%)	400 (200-670)	15000 (7500-25000)
Angola	2005	Male	55 to 59	49% (32-68%)	51% (32-68%)	1.7 (0.88-2.9)	8.5% (4.2-14%)	10% (5-17%)	320 (150-540)	10000 (5100-17000)
Angola	2005	Male	60 to 64	48% (30-67%)	52% (33-70%)	1.5 (0.7-2.7)	6.6% (3-11%)	7.9% (3.6-13%)	270 (120-490)	7500 (3400-13000)
Angola	2005	Male	65 to 69	48% (31-66%)	52% (34-69%)	1.3 (0.66-2.2)	5.2% (2-9.1%)	6.2% (2.3-11%)	230 (82-400)	5200 (1900-9000)
Angola	2005	Male	70 to 74	49% (32-66%)	51% (34-68%)	1.2 (0.56-2.1)	3.8% (1.1-7.5%)	4.4% (1.2-8.8%)	160 (45-310)	2900 (850-5700)
Angola	2005	Male	75 to 79	48% (31-66%)	52% (34-69%)	0.96 (0.45-1.7)	2.8% (0.62-5.9%)	3.2% (0.66-6.8%)	94 (19-200)	1300 (290-2800)
Angola	2005	Male	80 to 84	42% (25-61%)	58% (39-75%)	0.75 (0.35-1.3)	1.9% (0.34-4.1%)	2.1% (0.31-4.8%)	37 (5.7-81)	390 (75-850)
Angola	2005	Male	85 to 89	37% (22-53%)	63% (47-78%)	0.56 (0.26-1)	1.3% (0.21-2.8%)	1.5% (0.17-3.3%)	13 (1.4-29)	97 (17-210)
Angola	2005	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.48 (0.22-0.89)	0.8% (0.01-2%)	0.84% (-0.12-2.3%)	2.5 (-0.36-6.8)	14 (0.18-35)
Angola	2005	Male	95 plus	39% (21-61%)	61% (39-79%)	0.45 (0.17-0.92)	0.95% (-0.084-2.5%)	0.97% (-0.3-2.8%)	0.7 (-0.22-2)	2.7 (-0.25-7.2)
Angola	2010	Female	All Ages	20% (16-26%)	80% (74-84%)	0.73 (0.52-0.98)	1.9% (1.2-2.8%)	3.3% (1.9-5.3%)	3200 (1700-5200)	120000 (75000-190000)
Angola	2010	Female	15 to 19	20% (12-33%)	80% (67-88%)	0.63 (0.3-1.1)	1.8% (0.91-3%)	2.3% (0.91-4.2%)	40 (16-76)	3900 (1900-6600)
Angola	2010	Female	20 to 24	29% (18-44%)	71% (56-82%)	1.2 (0.63-2)	3.6% (2.1-5.8%)	4.4% (2.1-7.8%)	98 (47-170)	8600 (4800-14000)
Angola	2010	Female	25 to 29	34% (20-50%)	66% (50-80%)	1.7 (0.94-2.7)	4.4% (2.7-6.8%)	5.4% (2.9-8.6%)	140 (76-230)	11000 (6400-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	2010	Female	30 to 34	36% (21-53%)	64% (47-79%)	1.9 (1-2.9)	5.1% (3.1-7.8%)	6.3% (3.6-10%)	190 (99-310)	12000 (7000-19000)
Angola	2010	Female	35 to 39	37% (23-56%)	63% (44-77%)	1.8 (0.92-3.1)	5.5% (3.2-8.3%)	7% (3.9-11%)	210 (110-360)	12000 (6500-19000)
Angola	2010	Female	40 to 44	38% (23-56%)	62% (44-77%)	1.8 (0.87-3)	6.2% (3.4-9.8%)	8.1% (4.3-13%)	240 (120-440)	12000 (6100-21000)
Angola	2010	Female	45 to 49	37% (21-54%)	63% (46-79%)	1.7 (0.83-3)	7% (3.8-11%)	9.2% (4.9-15%)	300 (140-510)	13000 (6100-22000)
Angola	2010	Female	50 to 54	34% (20-52%)	66% (48-80%)	1.6 (0.77-2.8)	7.6% (3.7-12%)	9.7% (4.8-16%)	360 (160-640)	13000 (6100-24000)
Angola	2010	Female	55 to 59	31% (19-49%)	69% (51-81%)	1.4 (0.67-2.5)	7.3% (3.4-12%)	9.1% (4.3-15%)	380 (170-690)	12000 (5500-22000)
Angola	2010	Female	60 to 64	31% (18-48%)	69% (52-82%)	1.3 (0.6-2.4)	5.9% (2.4-11%)	7.2% (2.9-13%)	310 (120-570)	8300 (3300-15000)
Angola	2010	Female	65 to 69	32% (21-47%)	68% (53-79%)	1.3 (0.63-2.4)	5.4% (2-10%)	6.5% (2.4-12%)	320 (110-630)	7100 (2500-14000)
Angola	2010	Female	70 to 74	32% (21-47%)	68% (53-79%)	1.2 (0.54-2.2)	4.3% (1.2-8.5%)	5.1% (1.4-10%)	270 (71-550)	4800 (1300-9900)
Angola	2010	Female	75 to 79	29% (18-44%)	71% (56-82%)	0.96 (0.43-1.9)	3.1% (0.48-6.9%)	3.6% (0.55-8.1%)	170 (26-400)	2300 (360-5600)
Angola	2010	Female	80 to 84	27% (16-42%)	73% (58-84%)	0.78 (0.36-1.4)	2.3% (0.32-5.1%)	2.6% (0.37-6%)	89 (12-210)	920 (120-2200)
Angola	2010	Female	85 to 89	25% (15-39%)	75% (61-85%)	0.7 (0.31-1.3)	1.7% (0.16-4.3%)	2% (0.17-4.9%)	33 (2.8-81)	250 (22-610)
Angola	2010	Female	90 to 94	27% (17-40%)	73% (60-83%)	0.69 (0.3-1.3)	1.4% (0.057-3.7%)	1.6% (0.048-4.3%)	8.9 (0.27-23)	46 (1.8-120)
Angola	2010	Female	95 plus	31% (17-49%)	69% (51-83%)	0.7 (0.26-1.5)	1.3% (-0.051-3.6%)	1.5% (-0.13-4.2%)	2 (-0.17-5.7)	7.1 (-0.29-19)
Angola	2010	Male	All Ages	27% (22-35%)	73% (65-78%)	0.96 (0.7-1.3)	2.9% (2-4.1%)	4.8% (3.1-6.9%)	4600 (2800-6700)	200000 (130000-280000)
Angola	2010	Male	15 to 19	23% (14-38%)	77% (62-86%)	0.72 (0.34-1.4)	3.1% (1.7-5.2%)	3.9% (1.8-7.1%)	75 (32-140)	6900 (3600-12000)
Angola	2010	Male	20 to 24	36% (22-53%)	64% (47-78%)	1.5 (0.83-2.6)	7.9% (4.8-12%)	9.6% (5.4-15%)	240 (120-400)	19000 (11000-30000)
Angola	2010	Male	25 to 29	43% (26-60%)	57% (40-74%)	2.2 (1.3-3.5)	10% (6.6-15%)	13% (7.5-19%)	350 (190-560)	25000 (15000-38000)
Angola	2010	Male	30 to 34	45% (28-64%)	55% (36-72%)	2.5 (1.4-3.8)	11% (7-15%)	13% (8.2-19%)	400 (220-640)	25000 (15000-38000)
Angola	2010	Male	35 to 39	45% (29-63%)	55% (37-71%)	2.4 (1.3-4)	11% (6.7-15%)	13% (8-19%)	410 (230-650)	22000 (13000-35000)
Angola	2010	Male	40 to 44	47% (31-66%)	53% (34-69%)	2.4 (1.3-4.2)	11% (6.9-16%)	14% (8.6-20%)	430 (240-740)	21000 (12000-35000)
Angola	2010	Male	45 to 49	50% (33-68%)	50% (32-67%)	2.6 (1.3-4.5)	12% (7.2-17%)	15% (9-22%)	480 (260-770)	21000 (12000-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	2010	Male	50 to 54	51% (33-69%)	49% (31-67%)	2.6 (1.3-4.4)	12% (6.9-18%)	15% (8.9-23%)	510 (270-820)	19000 (10000-30000)
Angola	2010	Male	55 to 59	48% (31-68%)	52% (32-69%)	2.3 (1.2-3.9)	11% (5.8-16%)	13% (7.2-20%)	480 (240-770)	15000 (7900-25000)
Angola	2010	Male	60 to 64	47% (29-67%)	53% (33-71%)	2 (0.88-3.5)	8.6% (4.3-14%)	11% (5.2-17%)	370 (180-630)	10000 (5000-17000)
Angola	2010	Male	65 to 69	48% (31-66%)	52% (34-69%)	1.8 (0.87-3.1)	7.1% (3.5-12%)	8.5% (4.1-14%)	320 (150-570)	7400 (3400-13000)
Angola	2010	Male	70 to 74	49% (31-66%)	51% (34-69%)	1.5 (0.74-2.8)	5.3% (2.1-10%)	6.3% (2.5-12%)	250 (92-460)	4500 (1700-8400)
Angola	2010	Male	75 to 79	47% (31-65%)	53% (35-69%)	1.3 (0.59-2.4)	4% (1.3-8.1%)	4.7% (1.4-9.6%)	150 (45-310)	2100 (650-4300)
Angola	2010	Male	80 to 84	41% (24-59%)	59% (41-76%)	0.98 (0.45-1.8)	2.7% (0.78-5.7%)	3.2% (0.86-6.6%)	64 (17-140)	680 (180-1400)
Angola	2010	Male	85 to 89	36% (21-53%)	64% (47-79%)	0.74 (0.35-1.4)	2% (0.59-4.1%)	2.2% (0.61-4.7%)	23 (6.3-49)	180 (53-360)
Angola	2010	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.64 (0.28-1.2)	1.3% (0.3-3%)	1.4% (0.23-3.5%)	5.1 (0.84-12)	28 (6.6-66)
Angola	2010	Male	95 plus	39% (20-59%)	61% (41-80%)	0.6 (0.22-1.3)	1.5% (0.2-3.7%)	1.6% (0.048-4.2%)	1.4 (0.041-3.6)	5.1 (0.67-12)
Angola	2016	Female	All Ages	20% (16-25%)	80% (75-84%)	0.84 (0.48-1.4)	2.4% (1.3-3.7%)	4.1% (1.9-6.6%)	3400 (1500-5700)	130000 (68000-200000)
Angola	2016	Female	15 to 19	20% (12-33%)	80% (67-88%)	0.73 (0.31-1.4)	1.7% (0.88-3%)	2.3% (0.86-4.7%)	38 (14-80)	4000 (2000-7200)
Angola	2016	Female	20 to 24	29% (17-44%)	71% (56-83%)	1.4 (0.63-2.6)	3.6% (2-5.6%)	4.5% (1.9-7.9%)	86 (35-160)	8400 (4600-14000)
Angola	2016	Female	25 to 29	33% (19-50%)	67% (50-81%)	1.9 (0.92-3.3)	4.3% (2.4-6.5%)	5.3% (2.5-9%)	130 (58-230)	10000 (5800-17000)
Angola	2016	Female	30 to 34	35% (21-53%)	65% (47-79%)	2.1 (0.97-3.9)	4.9% (2.8-7.6%)	6.3% (3.2-10%)	170 (82-310)	12000 (6300-19000)
Angola	2016	Female	35 to 39	37% (22-54%)	63% (46-78%)	2.1 (0.93-4)	5.5% (3-8.7%)	7.2% (3.7-12%)	210 (96-380)	12000 (6300-21000)
Angola	2016	Female	40 to 44	37% (23-55%)	63% (45-77%)	2.1 (0.9-3.9)	6.2% (3.3-9.8%)	8.4% (4.3-13%)	250 (120-460)	13000 (6200-22000)
Angola	2016	Female	45 to 49	36% (22-53%)	64% (47-78%)	2 (0.86-3.7)	7.2% (3.6-11%)	9.7% (4.8-16%)	310 (140-540)	13000 (6000-23000)
Angola	2016	Female	50 to 54	34% (20-51%)	66% (49-80%)	1.8 (0.75-3.4)	7.6% (3.6-12%)	10% (4.7-16%)	370 (150-650)	14000 (5900-24000)
Angola	2016	Female	55 to 59	31% (19-49%)	69% (51-81%)	1.6 (0.66-3.3)	7.3% (3.1-12%)	9.3% (3.9-16%)	390 (160-700)	13000 (5100-22000)
Angola	2016	Female	60 to 64	31% (18-48%)	69% (52-82%)	1.5 (0.61-3.3)	6.1% (2.4-11%)	7.6% (2.9-14%)	380 (140-720)	10000 (3900-20000)
Angola	2016	Female	65 to 69	32% (21-48%)	68% (52-79%)	1.5 (0.64-3.2)	5.8% (1.9-11%)	7% (2.4-14%)	350 (120-710)	7900 (2600-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	2016	Female	70 to 74	32% (21-47%)	68% (53-79%)	1.4 (0.55-2.7)	4.8% (1.1-10%)	5.7% (1.4-12%)	300 (68-630)	5400 (1200-11000)
Angola	2016	Female	75 to 79	29% (18-45%)	71% (55-82%)	1.1 (0.45-2.3)	3.6% (0.42-8.1%)	4.2% (0.5-9.7%)	210 (25-490)	2900 (330-7000)
Angola	2016	Female	80 to 84	27% (16-41%)	73% (59-84%)	0.91 (0.37-1.8)	2.7% (0.29-6.3%)	3.2% (0.34-7.5%)	120 (14-280)	1200 (150-3000)
Angola	2016	Female	85 to 89	25% (15-38%)	75% (62-85%)	0.81 (0.32-1.6)	2.1% (0.15-5.3%)	2.4% (0.16-6.3%)	50 (3.2-130)	370 (28-940)
Angola	2016	Female	90 to 94	27% (16-40%)	73% (60-84%)	0.79 (0.31-1.6)	1.7% (0.0073-4.7%)	2% (-0.012-5.5%)	14 (-0.076-40)	72 (0.3-200)
Angola	2016	Female	95 plus	31% (17-49%)	69% (51-83%)	0.81 (0.28-1.8)	1.6% (-0.036-4.7%)	1.8% (-0.12-5.4%)	3.1 (-0.2-9.3)	11 (-0.26-32)
Angola	2016	Male	All Ages	27% (22-34%)	73% (66-78%)	1.1 (0.63-1.7)	3.9% (2.3-5.8%)	6.2% (3.6-9.4%)	5300 (2900-8300)	23000 (140000-350000)
Angola	2016	Male	15 to 19	23% (14-38%)	77% (62-86%)	0.82 (0.34-1.6)	3.1% (1.6-5.4%)	4% (1.6-7.6%)	87 (34-170)	8200 (4100-15000)
Angola	2016	Male	20 to 24	36% (22-51%)	64% (49-78%)	1.7 (0.78-3.3)	8.1% (4.6-13%)	9.9% (4.9-17%)	270 (120-490)	22000 (12000-37000)
Angola	2016	Male	25 to 29	43% (26-61%)	57% (39-74%)	2.5 (1.3-4.5)	11% (6.2-15%)	13% (7.1-20%)	390 (190-630)	28000 (15000-43000)
Angola	2016	Male	30 to 34	45% (27-64%)	55% (36-73%)	2.8 (1.3-5.1)	11% (6.5-16%)	14% (7.5-20%)	430 (220-690)	28000 (15000-43000)
Angola	2016	Male	35 to 39	45% (29-64%)	55% (36-71%)	2.8 (1.3-5.1)	11% (6.6-16%)	14% (7.8-21%)	460 (240-760)	26000 (14000-41000)
Angola	2016	Male	40 to 44	47% (31-66%)	53% (34-69%)	2.8 (1.2-5.3)	11% (6.5-17%)	15% (8.2-22%)	490 (260-820)	24000 (13000-40000)
Angola	2016	Male	45 to 49	50% (32-69%)	50% (31-68%)	3 (1.3-5.7)	12% (7-18%)	16% (8.9-23%)	540 (270-870)	23000 (12000-37000)
Angola	2016	Male	50 to 54	50% (32-68%)	50% (32-68%)	2.9 (1.2-5.7)	12% (6.8-18%)	16% (8.4-24%)	570 (280-940)	22000 (11000-35000)
Angola	2016	Male	55 to 59	48% (31-67%)	52% (33-69%)	2.6 (1.1-4.9)	11% (5.7-17%)	14% (7.3-22%)	540 (260-890)	18000 (8500-29000)
Angola	2016	Male	60 to 64	47% (29-66%)	53% (34-71%)	2.2 (0.88-4.6)	9.1% (4.2-15%)	11% (5.3-18%)	490 (220-840)	14000 (6300-23000)
Angola	2016	Male	65 to 69	48% (31-65%)	52% (35-69%)	2 (0.85-4)	7.6% (3.3-13%)	9.3% (4.1-17%)	390 (160-700)	8900 (3800-16000)
Angola	2016	Male	70 to 74	48% (30-65%)	52% (35-70%)	1.8 (0.75-3.3)	6% (2.2-11%)	7.1% (2.6-14%)	300 (100-580)	5400 (1900-11000)
Angola	2016	Male	75 to 79	46% (30-64%)	54% (36-70%)	1.5 (0.6-2.9)	4.5% (1.3-9.6%)	5.3% (1.5-11%)	190 (54-420)	2800 (780-5900)
Angola	2016	Male	80 to 84	41% (24-59%)	59% (41-76%)	1.1 (0.44-2.3)	3.2% (0.83-7.2%)	3.7% (0.89-8.3%)	87 (21-200)	930 (240-2000)
Angola	2016	Male	85 to 89	35% (22-52%)	65% (48-78%)	0.84 (0.35-1.8)	2.3% (0.62-5.1%)	2.6% (0.66-5.9%)	34 (8.8-79)	260 (72-590)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Angola	2016	Male	90 to 94	35% (21-51%)	65% (49-79%)	0.73 (0.28-1.5)	1.5% (0.29-3.9%)	1.7% (0.24-4.4%)	8.2 (1.2-21)	45 (8.4-110)
Angola	2016	Male	95 plus	38% (20-59%)	62% (41-80%)	0.68 (0.21-1.6)	1.7% (0.32-4.6%)	1.9% (0.2-5.2%)	2.1 (0.21-5.9)	7.8 (1.4-21)
Central African Republic	1990	Female	All Ages	24% (19-29%)	76% (71-81%)	1.4 (0.96-1.8)	4.3% (2.9-6%)	7.9% (5-11%)	1900 (1200-2700)	59000 (40000-83000)
Central African Republic	1990	Female	15 to 19	23% (14-35%)	77% (65-86%)	1.1 (0.5-2)	3.5% (1.5-6.4%)	5.2% (2.1-9.6%)	19 (7.4-36)	1500 (640-2700)
Central African Republic	1990	Female	20 to 24	32% (19-47%)	68% (53-81%)	2.1 (1-3.6)	7.3% (3.9-12%)	9.8% (4.8-16%)	49 (24-80)	3500 (1900-5500)
Central African Republic	1990	Female	25 to 29	36% (21-52%)	64% (48-79%)	2.7 (1.4-4.5)	9.1% (5.3-14%)	12% (6.5-19%)	66 (36-100)	4200 (2500-6600)
Central African Republic	1990	Female	30 to 34	38% (23-55%)	62% (45-77%)	3 (1.7-4.9)	11% (6.5-16%)	14% (8.4-21%)	82 (47-120)	4800 (2900-7200)
Central African Republic	1990	Female	35 to 39	39% (25-57%)	61% (43-75%)	3 (1.6-5)	12% (6.8-17%)	15% (8.7-23%)	93 (54-150)	4900 (2900-7400)
Central African Republic	1990	Female	40 to 44	40% (25-58%)	60% (42-75%)	3 (1.4-5)	12% (7.5-18%)	16% (9.6-23%)	100 (60-150)	4700 (2900-7000)
Central African Republic	1990	Female	45 to 49	39% (24-57%)	61% (43-76%)	2.8 (1.5-5.1)	14% (8.3-21%)	18% (11-27%)	140 (80-210)	5600 (3300-8400)
Central African Republic	1990	Female	50 to 54	36% (22-53%)	64% (47-78%)	2.6 (1.3-4.4)	15% (9-22%)	18% (11-28%)	170 (100-270)	6200 (3600-9600)
Central African Republic	1990	Female	55 to 59	34% (21-51%)	66% (49-79%)	2.3 (1.1-4.2)	15% (8.1-23%)	18% (9.7-28%)	210 (110-330)	6500 (3500-10000)
Central African Republic	1990	Female	60 to 64	33% (20-50%)	67% (50-80%)	2.2 (0.99-4)	13% (6.4-20%)	15% (7.5-24%)	210 (100-330)	5600 (2700-8700)
Central African Republic	1990	Female	65 to 69	35% (23-51%)	65% (49-77%)	2.2 (1.1-3.8)	13% (6.3-20%)	15% (7.4-23%)	230 (120-370)	5200 (2600-8100)
Central African Republic	1990	Female	70 to 74	35% (23-50%)	65% (50-77%)	2 (0.91-3.6)	11% (4.7-18%)	13% (5.3-21%)	210 (88-350)	3700 (1600-6200)
Central African Republic	1990	Female	75 to 79	32% (20-48%)	68% (52-80%)	1.6 (0.71-3.1)	8.6% (3.1-15%)	9.8% (3.7-17%)	150 (54-260)	2000 (740-3600)
Central African Republic	1990	Female	80 to 84	29% (17-44%)	71% (56-83%)	1.3 (0.6-2.5)	6.7% (2.4-12%)	7.7% (2.8-14%)	82 (30-150)	840 (310-1600)
Central African Republic	1990	Female	85 to 89	28% (17-40%)	72% (60-83%)	1.2 (0.51-2.2)	5.3% (1.8-10%)	6.2% (2-12%)	32 (11-62)	230 (77-450)
Central African Republic	1990	Female	90 to 94	29% (18-43%)	71% (57-82%)	1.1 (0.49-2.2)	4.3% (1.2-9.2%)	5.2% (1.5-11%)	8.3 (2.3-18)	41 (11-86)
Central African Republic	1990	Female	95 plus	33% (19-50%)	67% (50-81%)	1.2 (0.43-2.3)	3.7% (0.75-8.1%)	4.3% (0.85-9.6%)	1.7 (0.33-3.8)	5.7 (1.2-13)
Central African Republic	1990	Male	All Ages	31% (25-39%)	69% (61-75%)	1.7 (1.2-2.3)	7.2% (5.3-9.3%)	12% (8.2-15%)	3200 (2200-4300)	120000 (86000-150000)
Central African Republic	1990	Male	15 to 19	25% (15-40%)	75% (60-85%)	1.2 (0.51-2.2)	6.3% (2.8-11%)	8.1% (3.4-14%)	44 (18-76)	3300 (1400-5600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	1990	Male	20 to 24	37% (23-54%)	63% (46-77%)	2.4 (1.2-4.1)	14% (8.1-21%)	17% (9.2-25%)	130 (69-190)	8600 (4900-13000)
Central African Republic	1990	Male	25 to 29	45% (28-62%)	55% (38-72%)	3.4 (1.9-5.6)	17% (11-23%)	20% (13-28%)	170 (110-240)	11000 (7000-15000)
Central African Republic	1990	Male	30 to 34	47% (29-65%)	53% (35-71%)	3.8 (2.2-6.1)	19% (13-25%)	22% (15-30%)	200 (130-270)	11000 (7600-15000)
Central African Republic	1990	Male	35 to 39	47% (30-65%)	53% (35-70%)	3.8 (2-6.1)	20% (13-27%)	23% (15-31%)	220 (140-290)	11000 (7300-15000)
Central African Republic	1990	Male	40 to 44	49% (32-68%)	51% (32-68%)	3.9 (2-6.4)	21% (14-29%)	25% (16-33%)	260 (160-360)	12000 (7600-16000)
Central African Republic	1990	Male	45 to 49	52% (33-70%)	48% (30-67%)	4.1 (2.3-6.8)	23% (15-31%)	27% (18-36%)	320 (220-450)	13000 (8900-18000)
Central African Republic	1990	Male	50 to 54	52% (34-71%)	48% (29-66%)	4 (2.1-6.7)	23% (15-31%)	27% (18-36%)	370 (230-510)	13000 (8200-18000)
Central African Republic	1990	Male	55 to 59	50% (33-69%)	50% (31-67%)	3.5 (1.8-6.2)	21% (13-29%)	24% (15-34%)	370 (220-520)	12000 (6800-16000)
Central African Republic	1990	Male	60 to 64	49% (31-68%)	51% (32-69%)	3.1 (1.4-5.6)	18% (9.4-25%)	20% (11-29%)	340 (170-510)	8900 (4700-13000)
Central African Republic	1990	Male	65 to 69	49% (32-66%)	51% (34-68%)	2.8 (1.4-4.8)	15% (8-23%)	17% (8.9-26%)	300 (150-460)	6600 (3400-10000)
Central African Republic	1990	Male	70 to 74	50% (33-67%)	50% (33-67%)	2.4 (1.2-4.3)	13% (5.8-21%)	14% (6.5-23%)	230 (100-380)	4100 (1900-6800)
Central African Republic	1990	Male	75 to 79	48% (31-66%)	52% (34-69%)	2 (0.9-3.7)	9.9% (3.6-18%)	11% (4.1-20%)	150 (54-270)	2000 (760-3800)
Central African Republic	1990	Male	80 to 84	43% (25-61%)	57% (39-75%)	1.5 (0.7-2.8)	7.3% (2.6-13%)	8.2% (2.9-15%)	65 (23-120)	670 (240-1200)
Central African Republic	1990	Male	85 to 89	37% (23-54%)	63% (46-77%)	1.1 (0.52-2.1)	5% (1.7-10%)	5.8% (1.9-11%)	19 (6.5-38)	140 (49-270)
Central African Republic	1990	Male	90 to 94	37% (22-53%)	63% (47-78%)	0.99 (0.43-1.8)	3.4% (0.85-7.1%)	3.9% (0.92-8.3%)	3.6 (0.84-7.7)	18 (4.5-39)
Central African Republic	1990	Male	95 plus	40% (22-60%)	60% (40-78%)	0.92 (0.34-1.9)	3.4% (0.55-8%)	3.9% (0.5-9.3%)	0.81 (0.1-1.9)	2.8 (0.46-6.5)
Central African Republic	1995	Female	All Ages	24% (20-29%)	76% (71-80%)	1.2 (0.84-1.6)	3.7% (2.3-5.2%)	6.6% (3.9-9.5%)	1900 (1100-2800)	61000 (39000-88000)
Central African Republic	1995	Female	15 to 19	22% (14-35%)	78% (65-86%)	0.93 (0.45-1.7)	2.9% (1.1-5.4%)	4.1% (1.4-7.9%)	18 (5.9-36)	1400 (570-2700)
Central African Republic	1995	Female	20 to 24	32% (19-47%)	68% (53-81%)	1.8 (0.88-3)	5.5% (2.7-8.8%)	6.8% (3-12%)	49 (22-81)	3500 (1700-5700)
Central African Republic	1995	Female	25 to 29	36% (21-52%)	64% (48-79%)	2.3 (1.3-3.8)	6.1% (3.1-10%)	7.3% (3.4-13%)	69 (34-110)	4500 (2300-7200)
Central African Republic	1995	Female	30 to 34	38% (23-56%)	62% (44-77%)	2.6 (1.4-4.2)	7.3% (4-11%)	8.8% (4.5-14%)	87 (47-140)	5100 (2900-8000)
Central African Republic	1995	Female	35 to 39	40% (25-57%)	60% (43-75%)	2.6 (1.4-4.4)	8.4% (4.6-13%)	10% (5.3-17%)	100 (55-170)	5400 (3000-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	1995	Female	40 to 44	40% (25-58%)	60% (42-75%)	2.6 (1.3-4.4)	9.2% (5.3-14%)	11% (6.3-18%)	110 (59-180)	5100 (2800-8200)
Central African Republic	1995	Female	45 to 49	39% (24-57%)	61% (43-76%)	2.4 (1.2-4.3)	11% (6.2-17%)	14% (7.7-21%)	130 (73-220)	5500 (3100-9000)
Central African Republic	1995	Female	50 to 54	36% (22-54%)	64% (46-78%)	2.2 (1.1-3.8)	12% (6.8-19%)	15% (8.3-23%)	180 (91-290)	6300 (3300-10000)
Central African Republic	1995	Female	55 to 59	33% (21-51%)	67% (49-79%)	2 (0.99-3.6)	13% (6.7-20%)	15% (7.7-24%)	210 (100-340)	6600 (3300-11000)
Central African Republic	1995	Female	60 to 64	33% (20-50%)	67% (50-80%)	1.9 (0.87-3.5)	11% (5.4-18%)	13% (6.1-21%)	210 (99-340)	5600 (2600-9100)
Central African Republic	1995	Female	65 to 69	35% (22-50%)	65% (50-78%)	1.9 (0.91-3.2)	11% (5.2-18%)	13% (6-21%)	240 (110-400)	5300 (2400-8700)
Central African Republic	1995	Female	70 to 74	35% (23-50%)	65% (50-77%)	1.7 (0.77-3.1)	9.5% (3.7-16%)	11% (4.3-18%)	210 (81-370)	3800 (1400-6600)
Central African Republic	1995	Female	75 to 79	32% (20-48%)	68% (52-80%)	1.4 (0.61-2.7)	7.2% (2.2-14%)	8.2% (2.6-15%)	140 (45-270)	1900 (620-3700)
Central African Republic	1995	Female	80 to 84	29% (18-45%)	71% (55-82%)	1.1 (0.51-2.1)	5.6% (1.7-11%)	6.4% (2-12%)	79 (25-160)	810 (260-1600)
Central African Republic	1995	Female	85 to 89	27% (17-41%)	73% (59-83%)	1 (0.44-1.8)	4.4% (1.4-9.4%)	5.1% (1.6-11%)	30 (8.9-62)	210 (65-440)
Central African Republic	1995	Female	90 to 94	29% (18-43%)	71% (57-82%)	0.97 (0.43-1.8)	3.6% (0.84-8%)	4.3% (1-9.6%)	8.1 (1.9-18)	40 (9.5-89)
Central African Republic	1995	Female	95 plus	33% (19-51%)	67% (49-81%)	0.99 (0.37-1.9)	3.1% (0.45-7.4%)	3.6% (0.48-8.9%)	1.7 (0.23-4.2)	5.9 (0.87-14)
Central African Republic	1995	Male	All Ages	31% (25-39%)	69% (61-75%)	1.5 (1.1-2)	6.4% (4.5-8.5%)	10% (6.6-13%)	3300 (2100-4600)	130000 (86000-170000)
Central African Republic	1995	Male	15 to 19	25% (15-40%)	75% (60-85%)	1 (0.46-1.9)	5.4% (2.4-9.8%)	6.9% (2.7-13%)	45 (18-85)	3400 (1500-6200)
Central African Republic	1995	Male	20 to 24	38% (24-54%)	62% (46-76%)	2.1 (1.1-3.6)	12% (6.5-18%)	14% (7.3-22%)	140 (70-210)	9400 (5100-14000)
Central African Republic	1995	Male	25 to 29	45% (28-63%)	55% (37-72%)	3 (1.7-5)	13% (7.8-19%)	15% (8.6-23%)	190 (110-280)	12000 (7300-17000)
Central African Republic	1995	Male	30 to 34	47% (29-66%)	53% (34-71%)	3.4 (1.9-5.2)	14% (8.5-20%)	16% (9.3-23%)	220 (130-310)	12000 (7900-18000)
Central African Republic	1995	Male	35 to 39	47% (31-66%)	53% (34-69%)	3.3 (1.8-5.3)	15% (9-21%)	17% (10-25%)	240 (150-360)	13000 (7700-18000)
Central African Republic	1995	Male	40 to 44	49% (33-68%)	51% (32-67%)	3.4 (1.7-5.5)	17% (9.7-23%)	19% (11-26%)	290 (160-420)	13000 (7600-19000)
Central African Republic	1995	Male	45 to 49	52% (34-69%)	48% (31-66%)	3.6 (2-5.9)	19% (12-26%)	21% (13-30%)	340 (200-500)	14000 (8100-20000)
Central African Republic	1995	Male	50 to 54	53% (35-71%)	47% (29-65%)	3.5 (1.8-5.9)	20% (12-28%)	23% (14-32%)	390 (220-580)	14000 (8000-21000)
Central African Republic	1995	Male	55 to 59	50% (33-70%)	50% (30-67%)	3.1 (1.6-5.3)	19% (11-26%)	21% (12-30%)	390 (210-590)	12000 (6600-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	1995	Male	60 to 64	49% (31-68%)	51% (32-69%)	2.7 (1.2-4.8)	15% (7.7-23%)	17% (8.7-26%)	340 (160-540)	9100 (4300-14000)
Central African Republic	1995	Male	65 to 69	50% (32-67%)	50% (33-68%)	2.4 (1.2-4)	13% (6.5-21%)	15% (7.2-23%)	300 (150-480)	6700 (3300-11000)
Central African Republic	1995	Male	70 to 74	50% (33-67%)	50% (33-67%)	2.1 (1-3.7)	11% (4.3-18%)	12% (4.9-20%)	230 (93-390)	4100 (1700-7000)
Central African Republic	1995	Male	75 to 79	48% (32-66%)	52% (34-68%)	1.7 (0.78-3.2)	8.4% (3-16%)	9.5% (3.3-18%)	140 (50-270)	2000 (700-3700)
Central African Republic	1995	Male	80 to 84	43% (26-61%)	57% (39-74%)	1.3 (0.61-2.4)	6.2% (2.1-12%)	7% (2.3-14%)	64 (21-130)	650 (220-1300)
Central African Republic	1995	Male	85 to 89	37% (23-55%)	63% (45-77%)	1 (0.46-1.8)	4.2% (1.3-8.9%)	4.9% (1.4-10%)	18 (5.4-38)	130 (40-280)
Central African Republic	1995	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.86 (0.37-1.6)	2.8% (0.62-6.1%)	3.2% (0.64-7.2%)	3.6 (0.72-7.9)	18 (4-39)
Central African Republic	1995	Male	95 plus	40% (22-61%)	60% (39-78%)	0.81 (0.3-1.6)	2.9% (0.35-7%)	3.3% (0.27-8%)	0.84 (0.07-2)	2.9 (0.35-6.9)
Central African Republic	2000	Female	All Ages	23% (19-29%)	77% (71-81%)	0.99 (0.71-1.3)	3% (1.7-4.5%)	5.3% (2.8-8%)	1800 (940-2900)	60000 (34000-92000)
Central African Republic	2000	Female	15 to 19	22% (13-34%)	78% (66-87%)	0.78 (0.37-1.4)	2.5% (1-4.5%)	3.4% (1.2-6.5%)	17 (6.2-34)	1400 (580-2600)
Central African Republic	2000	Female	20 to 24	31% (19-46%)	69% (54-81%)	1.5 (0.75-2.6)	4.6% (2.3-7.8%)	5.6% (2.5-9.9%)	48 (22-85)	3500 (1800-6000)
Central African Republic	2000	Female	25 to 29	35% (21-52%)	65% (48-79%)	2 (1.1-3.3)	4.5% (2.2-7.9%)	5.1% (2.3-9.4%)	70 (33-120)	4500 (2300-7700)
Central African Republic	2000	Female	30 to 34	37% (23-55%)	63% (45-77%)	2.2 (1.2-3.5)	5% (2.6-8.2%)	5.7% (2.8-9.5%)	86 (43-150)	5000 (2600-8400)
Central African Republic	2000	Female	35 to 39	39% (24-57%)	61% (43-76%)	2.2 (1.2-3.8)	5.8% (2.9-9.8%)	6.7% (3.2-12%)	100 (50-180)	5300 (2700-9300)
Central African Republic	2000	Female	40 to 44	39% (25-58%)	61% (42-75%)	2.2 (1.1-3.8)	6.6% (3.5-11%)	7.7% (4-12%)	110 (52-200)	5200 (2500-9000)
Central African Republic	2000	Female	45 to 49	38% (23-56%)	62% (44-77%)	2 (1-3.8)	8.4% (4.3-13%)	9.9% (5.1-16%)	140 (64-240)	5600 (2800-9800)
Central African Republic	2000	Female	50 to 54	36% (21-53%)	64% (47-79%)	1.8 (0.9-3.2)	9.8% (4.8-16%)	12% (5.6-19%)	170 (77-290)	6000 (2800-10000)
Central African Republic	2000	Female	55 to 59	33% (20-50%)	67% (50-80%)	1.7 (0.81-3)	11% (5.2-18%)	13% (6.1-21%)	200 (88-360)	6400 (2800-11000)
Central African Republic	2000	Female	60 to 64	32% (20-49%)	68% (51-80%)	1.6 (0.75-3)	9.6% (3.9-16%)	11% (4.6-19%)	200 (82-360)	5400 (2200-9400)
Central African Republic	2000	Female	65 to 69	34% (22-49%)	66% (51-78%)	1.6 (0.77-2.7)	9.3% (4-16%)	11% (4.6-18%)	220 (92-390)	4900 (2000-8600)
Central African Republic	2000	Female	70 to 74	34% (22-49%)	66% (51-78%)	1.4 (0.66-2.5)	7.9% (2.8-14%)	9% (3.3-16%)	200 (71-370)	3600 (1300-6500)
Central African Republic	2000	Female	75 to 79	31% (19-47%)	69% (53-81%)	1.1 (0.51-2.2)	5.9% (1.7-12%)	6.7% (1.9-13%)	130 (37-270)	1800 (510-3700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	2000	Female	80 to 84	28% (17-44%)	72% (56-83%)	0.94 (0.43-1.7)	4.6% (1.3-9.2%)	5.3% (1.5-11%)	75 (20-150)	760 (200-1500)
Central African Republic	2000	Female	85 to 89	27% (17-40%)	73% (60-83%)	0.83 (0.38-1.6)	3.6% (0.89-7.8%)	4.2% (1-9.1%)	26 (6.5-57)	180 (47-410)
Central African Republic	2000	Female	90 to 94	28% (18-41%)	72% (59-82%)	0.81 (0.35-1.6)	2.9% (0.56-6.7%)	3.5% (0.68-8.2%)	6.9 (1.3-16)	34 (6.4-79)
Central African Republic	2000	Female	95 plus	32% (18-50%)	68% (50-82%)	0.83 (0.32-1.6)	2.6% (0.26-6.3%)	3% (0.24-7.5%)	1.6 (0.13-3.9)	5.3 (0.54-13)
Central African Republic	2000	Male	All Ages	31% (25-39%)	69% (61-75%)	1.3 (0.93-1.7)	5.7% (3.8-7.8%)	8.7% (5.5-12%)	3400 (2000-4900)	130000 (83000-180000)
Central African Republic	2000	Male	15 to 19	25% (14-39%)	75% (61-86%)	0.89 (0.4-1.7)	4.9% (2-8.9%)	6.2% (2.3-11%)	48 (17-91)	3600 (1500-6600)
Central African Republic	2000	Male	20 to 24	37% (23-54%)	63% (46-77%)	1.9 (0.94-3.2)	11% (6.1-18%)	13% (6.8-21%)	140 (73-230)	9900 (5400-16000)
Central African Republic	2000	Male	25 to 29	45% (27-62%)	55% (38-73%)	2.6 (1.5-4.2)	12% (6.8-18%)	14% (7.3-21%)	200 (120-310)	13000 (7600-19000)
Central African Republic	2000	Male	30 to 34	47% (29-65%)	53% (35-71%)	3 (1.7-4.6)	12% (7.1-17%)	13% (7.6-19%)	240 (140-350)	13000 (8100-20000)
Central African Republic	2000	Male	35 to 39	47% (31-65%)	53% (35-69%)	2.9 (1.6-4.7)	12% (6.9-18%)	13% (7.6-20%)	260 (150-400)	13000 (7700-21000)
Central African Republic	2000	Male	40 to 44	49% (32-68%)	51% (32-68%)	2.9 (1.5-5)	14% (8-20%)	15% (8.7-22%)	310 (170-480)	14000 (7900-22000)
Central African Republic	2000	Male	45 to 49	52% (34-69%)	48% (31-66%)	3.1 (1.7-5.3)	16% (9.3-22%)	18% (10-25%)	360 (190-560)	15000 (7900-23000)
Central African Republic	2000	Male	50 to 54	52% (35-71%)	48% (29-65%)	3 (1.6-5.1)	17% (9.6-26%)	19% (11-29%)	390 (200-620)	14000 (7100-22000)
Central African Republic	2000	Male	55 to 59	50% (33-68%)	50% (32-67%)	2.7 (1.4-4.7)	16% (8.8-24%)	19% (9.9-27%)	390 (200-620)	12000 (6200-19000)
Central African Republic	2000	Male	60 to 64	49% (31-68%)	51% (32-69%)	2.3 (1.1-4.3)	14% (6-22%)	15% (6.6-24%)	340 (150-560)	9000 (4000-15000)
Central African Republic	2000	Male	65 to 69	50% (33-67%)	50% (33-67%)	2.1 (1.1-3.6)	12% (5-19%)	13% (5.5-21%)	290 (120-500)	6500 (2600-11000)
Central African Republic	2000	Male	70 to 74	50% (34-66%)	50% (34-66%)	1.8 (0.9-3.2)	9.3% (3.6-16%)	10% (4-18%)	230 (85-400)	4000 (1500-7000)
Central African Republic	2000	Male	75 to 79	48% (32-66%)	52% (34-68%)	1.5 (0.69-2.8)	7.3% (2.2-14%)	8.1% (2.4-15%)	140 (41-270)	1900 (580-3700)
Central African Republic	2000	Male	80 to 84	43% (26-62%)	57% (38-74%)	1.2 (0.54-2.1)	5.3% (1.5-11%)	5.9% (1.6-12%)	62 (17-130)	630 (180-1300)
Central African Republic	2000	Male	85 to 89	38% (23-54%)	62% (46-77%)	0.88 (0.41-1.6)	3.6% (0.9-7.6%)	4.1% (0.99-8.6%)	17 (4.1-35)	120 (31-260)
Central African Republic	2000	Male	90 to 94	37% (23-53%)	63% (47-77%)	0.76 (0.33-1.4)	2.3% (0.41-5.3%)	2.7% (0.38-6.2%)	3.3 (0.47-7.6)	17 (2.9-38)
Central African Republic	2000	Male	95 plus	40% (21-61%)	60% (39-79%)	0.71 (0.26-1.5)	2.4% (0.18-6%)	2.7% (0.068-7.1%)	0.82 (0.02-2.1)	2.8 (0.21-7.3)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	2005	Female	All Ages	23% (19-29%)	77% (71-81%)	0.93 (0.65-1.3)	2.9% (1.6-4.4%)	4.9% (2.6-7.7%)	1800 (910-3000)	59000 (33000-98000)
Central African Republic	2005	Female	15 to 19	21% (13-34%)	79% (66-87%)	0.72 (0.34-1.3)	2.2% (0.97-4.1%)	2.9% (1.1-5.7%)	17 (6.2-36)	1400 (620-2700)
Central African Republic	2005	Female	20 to 24	30% (18-45%)	70% (55-82%)	1.4 (0.72-2.5)	4.8% (2.3-8.2%)	5.9% (2.4-11%)	50 (22-93)	3700 (1800-6500)
Central African Republic	2005	Female	25 to 29	34% (21-51%)	66% (49-79%)	1.9 (1-3.1)	4.5% (2.2-7.7%)	5.2% (2.3-9.4%)	72 (33-130)	4700 (2300-8200)
Central African Republic	2005	Female	30 to 34	37% (22-54%)	63% (46-78%)	2.1 (1.1-3.5)	4.5% (2.3-7.5%)	5% (2.5-8.6%)	87 (41-160)	5100 (2600-8900)
Central African Republic	2005	Female	35 to 39	38% (24-56%)	62% (44-76%)	2.1 (1.1-3.5)	4.8% (2.4-8.9%)	5.5% (2.6-10%)	94 (43-180)	4900 (2400-9200)
Central African Republic	2005	Female	40 to 44	39% (24-56%)	61% (44-76%)	2 (0.99-3.6)	5.6% (2.7-9.1%)	6.4% (3-11%)	110 (47-190)	5000 (2300-9000)
Central African Republic	2005	Female	45 to 49	38% (23-55%)	62% (45-77%)	2 (0.97-3.5)	7.5% (3.8-12%)	8.8% (4.3-15%)	140 (63-270)	5900 (2700-11000)
Central African Republic	2005	Female	50 to 54	35% (21-52%)	65% (48-79%)	1.8 (0.85-3.1)	9.1% (4.3-16%)	11% (5.2-18%)	170 (71-320)	6300 (2600-11000)
Central African Republic	2005	Female	55 to 59	32% (20-49%)	68% (51-80%)	1.6 (0.76-2.8)	10% (4.7-17%)	12% (5.6-20%)	200 (79-360)	6200 (2500-11000)
Central African Republic	2005	Female	60 to 64	32% (19-49%)	68% (51-81%)	1.5 (0.71-2.8)	8.9% (3.8-15%)	10% (4.4-17%)	200 (77-360)	5200 (2100-9500)
Central African Republic	2005	Female	65 to 69	33% (22-49%)	67% (51-78%)	1.5 (0.72-2.6)	8.6% (3.6-15%)	9.9% (4.1-17%)	220 (84-390)	4800 (1900-8600)
Central African Republic	2005	Female	70 to 74	33% (22-48%)	67% (52-78%)	1.3 (0.61-2.4)	7.3% (2.4-13%)	8.3% (2.8-16%)	200 (65-380)	3500 (1100-6800)
Central African Republic	2005	Female	75 to 79	30% (19-46%)	70% (54-81%)	1.1 (0.47-2.1)	5.5% (1.4-11%)	6.2% (1.5-13%)	130 (33-290)	1800 (450-4000)
Central African Republic	2005	Female	80 to 84	28% (17-43%)	72% (57-83%)	0.89 (0.4-1.6)	4.2% (1.1-9.1%)	4.9% (1.2-10%)	76 (20-160)	770 (200-1700)
Central African Republic	2005	Female	85 to 89	26% (16-40%)	74% (60-84%)	0.78 (0.34-1.5)	3.3% (0.62-7.2%)	3.8% (0.73-8.4%)	26 (5.2-59)	190 (38-420)
Central African Republic	2005	Female	90 to 94	28% (17-41%)	72% (59-83%)	0.76 (0.31-1.5)	2.6% (0.41-6.2%)	3.2% (0.46-7.5%)	6.5 (0.95-16)	32 (4.8-77)
Central African Republic	2005	Female	95 plus	31% (17-49%)	69% (51-83%)	0.76 (0.28-1.6)	2.3% (0.2-6.1%)	2.7% (0.18-7.2%)	1.5 (0.094-4)	5 (0.42-13)
Central African Republic	2005	Male	All Ages	31% (25-39%)	69% (61-75%)	1.2 (0.89-1.7)	5.6% (3.6-7.8%)	8.4% (5.2-12%)	3400 (2000-5000)	130000 (84000-190000)
Central African Republic	2005	Male	15 to 19	25% (15-40%)	75% (60-85%)	0.85 (0.37-1.6)	4.6% (1.9-8.5%)	5.6% (2.1-11%)	50 (18-94)	3800 (1600-6900)
Central African Republic	2005	Male	20 to 24	38% (23-55%)	62% (45-77%)	1.8 (0.9-3)	12% (6.3-18%)	14% (6.9-21%)	160 (76-250)	11000 (5600-17000)
Central African Republic	2005	Male	25 to 29	45% (28-62%)	55% (38-72%)	2.5 (1.4-4.1)	13% (7.5-19%)	15% (8.2-23%)	220 (120-330)	14000 (7900-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	2005	Male	30 to 34	47% (30-65%)	53% (35-70%)	2.8 (1.5-4.5)	12% (7.3-17%)	13% (7.9-20%)	250 (140-370)	14000 (8600-21000)
Central African Republic	2005	Male	35 to 39	46% (30-64%)	54% (36-70%)	2.7 (1.4-4.6)	12% (6.5-17%)	13% (7.1-19%)	260 (150-410)	14000 (7700-21000)
Central African Republic	2005	Male	40 to 44	48% (32-67%)	52% (33-68%)	2.8 (1.4-4.8)	13% (7.1-19%)	14% (7.9-21%)	310 (160-490)	14000 (7700-22000)
Central African Republic	2005	Male	45 to 49	52% (33-69%)	48% (31-67%)	3 (1.6-5.1)	15% (8.1-22%)	17% (9.1-24%)	370 (190-570)	15000 (7800-23000)
Central African Republic	2005	Male	50 to 54	52% (34-70%)	48% (30-66%)	2.9 (1.5-5)	16% (8.8-24%)	18% (9.8-27%)	400 (190-640)	14000 (6900-23000)
Central African Republic	2005	Male	55 to 59	50% (33-69%)	50% (31-67%)	2.5 (1.3-4.4)	16% (8-24%)	18% (9-27%)	370 (180-600)	12000 (5700-18000)
Central African Republic	2005	Male	60 to 64	48% (30-68%)	52% (32-70%)	2.2 (1-4)	13% (5.6-21%)	14% (6.3-24%)	330 (140-570)	8800 (3700-15000)
Central African Republic	2005	Male	65 to 69	49% (32-66%)	51% (34-68%)	2 (0.99-3.5)	11% (4.7-18%)	12% (5.2-21%)	290 (120-500)	6400 (2700-11000)
Central African Republic	2005	Male	70 to 74	50% (33-67%)	50% (33-67%)	1.8 (0.85-3.1)	8.7% (3.1-16%)	9.7% (3.4-18%)	220 (77-400)	4000 (1400-7100)
Central African Republic	2005	Male	75 to 79	48% (32-66%)	52% (34-68%)	1.5 (0.65-2.8)	6.8% (1.8-13%)	7.6% (2.1-15%)	140 (37-280)	2000 (520-3800)
Central African Republic	2005	Male	80 to 84	43% (26-61%)	57% (39-74%)	1.1 (0.5-2.1)	4.9% (1.2-10%)	5.5% (1.4-12%)	63 (15-130)	640 (160-1400)
Central African Republic	2005	Male	85 to 89	38% (23-54%)	62% (46-77%)	0.85 (0.37-1.5)	3.3% (0.71-7.1%)	3.8% (0.79-8%)	17 (3.6-37)	130 (27-270)
Central African Republic	2005	Male	90 to 94	37% (22-53%)	63% (47-78%)	0.73 (0.31-1.4)	2.1% (0.25-5%)	2.4% (0.22-5.8%)	3.2 (0.3-7.7)	17 (2-39)
Central African Republic	2005	Male	95 plus	40% (21-62%)	60% (38-79%)	0.68 (0.24-1.4)	2.2% (0.14-5.7%)	2.5% (0.0019-6.7%)	0.83 (0.00064-2.2)	2.9 (0.19-7.4)
Central African Republic	2010	Female	All Ages	23% (18-28%)	77% (72-82%)	0.96 (0.64-1.4)	3% (1.6-4.7%)	5.1% (2.5-8.1%)	1900 (870-3200)	62000 (32000-100000)
Central African Republic	2010	Female	15 to 19	21% (13-33%)	79% (67-87%)	0.74 (0.34-1.4)	2.2% (0.9-4.1%)	2.8% (0.97-5.6%)	19 (6.4-39)	1600 (640-3000)
Central African Republic	2010	Female	20 to 24	30% (19-45%)	70% (55-81%)	1.4 (0.71-2.6)	5.2% (2.4-9.1%)	6.6% (2.7-12%)	56 (24-100)	4100 (2000-7200)
Central African Republic	2010	Female	25 to 29	34% (21-50%)	66% (50-79%)	1.9 (0.99-3.2)	5.4% (2.7-9%)	6.5% (3-11%)	80 (37-140)	5200 (2700-8900)
Central African Republic	2010	Female	30 to 34	37% (22-54%)	63% (46-78%)	2.1 (1.1-3.5)	5.3% (2.7-8.8%)	6.1% (3-10%)	95 (43-170)	5600 (2800-9800)
Central African Republic	2010	Female	35 to 39	38% (24-57%)	62% (43-76%)	2.1 (1-3.6)	5.1% (2.5-9%)	5.9% (2.8-11%)	97 (44-180)	5100 (2400-9500)
Central African Republic	2010	Female	40 to 44	38% (24-57%)	62% (43-76%)	2 (0.99-3.6)	5.9% (3-9.6%)	6.8% (3.4-11%)	110 (48-190)	5000 (2300-8700)
Central African Republic	2010	Female	45 to 49	37% (22-55%)	63% (45-78%)	2 (0.95-3.6)	7.9% (3.8-13%)	9.4% (4.6-16%)	140 (63-270)	5900 (2700-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	2010	Female	50 to 54	35% (20-53%)	65% (47-80%)	1.8 (0.85-3.1)	9.4% (4.6-16%)	11% (5.4-19%)	180 (75-330)	6500 (2800-12000)
Central African Republic	2010	Female	55 to 59	32% (19-49%)	68% (51-81%)	1.6 (0.73-3)	10% (4.6-17%)	12% (5.6-20%)	210 (84-370)	6500 (2600-11000)
Central African Republic	2010	Female	60 to 64	32% (19-48%)	68% (52-81%)	1.5 (0.7-2.9)	9% (3.6-15%)	11% (4.3-18%)	200 (72-360)	5200 (1900-9600)
Central African Republic	2010	Female	65 to 69	33% (22-48%)	67% (52-78%)	1.5 (0.7-2.7)	8.6% (3.3-15%)	9.9% (3.8-18%)	220 (82-410)	4900 (1800-9100)
Central African Republic	2010	Female	70 to 74	33% (21-48%)	67% (52-79%)	1.3 (0.59-2.6)	7.2% (2.2-14%)	8.3% (2.5-16%)	200 (63-390)	3600 (1100-7000)
Central African Republic	2010	Female	75 to 79	30% (19-46%)	70% (54-81%)	1.1 (0.48-2.2)	5.4% (1.2-12%)	6.2% (1.4-14%)	140 (29-300)	1900 (400-4100)
Central African Republic	2010	Female	80 to 84	27% (16-42%)	73% (58-84%)	0.89 (0.39-1.7)	4.2% (0.88-9.1%)	4.9% (1-11%)	83 (18-190)	840 (180-1900)
Central African Republic	2010	Female	85 to 89	26% (16-39%)	74% (61-84%)	0.79 (0.34-1.5)	3.3% (0.59-7.5%)	3.8% (0.68-8.7%)	30 (5.5-69)	220 (39-500)
Central African Republic	2010	Female	90 to 94	28% (17-41%)	72% (59-83%)	0.77 (0.33-1.5)	2.7% (0.41-6.4%)	3.2% (0.48-7.7%)	7.7 (1.2-19)	38 (5.9-91)
Central African Republic	2010	Female	95 plus	32% (18-50%)	68% (50-82%)	0.79 (0.29-1.6)	2.4% (0.21-6.1%)	2.8% (0.19-7.2%)	1.7 (0.12-4.3)	5.6 (0.5-15)
Central African Republic	2010	Male	All Ages	31% (25-39%)	69% (61-75%)	1.3 (0.86-1.8)	5.9% (3.7-8.4%)	8.9% (5.2-13%)	3600 (2000-5400)	140000 (86000-200000)
Central African Republic	2010	Male	15 to 19	24% (14-39%)	76% (61-86%)	0.85 (0.36-1.6)	4.4% (1.8-8%)	5.3% (2-10%)	52 (19-99)	4000 (1600-7200)
Central African Republic	2010	Male	20 to 24	37% (23-54%)	63% (46-77%)	1.8 (0.88-3.2)	12% (6.2-18%)	14% (6.8-22%)	170 (82-280)	12000 (6000-19000)
Central African Republic	2010	Male	25 to 29	44% (27-61%)	56% (39-73%)	2.5 (1.4-4.2)	15% (8.7-21%)	17% (9.8-25%)	240 (140-380)	15000 (8900-23000)
Central African Republic	2010	Male	30 to 34	46% (28-65%)	54% (35-72%)	2.8 (1.5-4.6)	14% (8.6-20%)	16% (9.5-24%)	270 (160-410)	16000 (9200-23000)
Central African Republic	2010	Male	35 to 39	46% (30-64%)	54% (36-70%)	2.7 (1.4-4.6)	13% (7.4-20%)	15% (8.3-22%)	290 (150-450)	15000 (8000-23000)
Central African Republic	2010	Male	40 to 44	48% (32-67%)	52% (33-68%)	2.8 (1.4-4.8)	14% (7.9-21%)	16% (8.8-23%)	320 (170-520)	15000 (7800-24000)
Central African Republic	2010	Male	45 to 49	51% (34-69%)	49% (31-66%)	3 (1.5-5)	16% (8.7-23%)	18% (9.9-26%)	370 (190-590)	15000 (7700-24000)
Central African Republic	2010	Male	50 to 54	52% (33-70%)	48% (30-67%)	2.9 (1.5-5.1)	17% (9.3-25%)	19% (11-28%)	410 (200-660)	15000 (7300-23000)
Central African Republic	2010	Male	55 to 59	49% (32-68%)	51% (32-68%)	2.6 (1.2-4.6)	16% (8.1-25%)	18% (9-28%)	390 (180-630)	12000 (5800-20000)
Central African Republic	2010	Male	60 to 64	48% (30-68%)	52% (32-70%)	2.2 (1-4.2)	13% (5.9-21%)	15% (6.6-24%)	330 (140-560)	8800 (3800-15000)
Central African Republic	2010	Male	65 to 69	49% (32-67%)	51% (33-68%)	2 (0.98-3.5)	11% (4.6-19%)	13% (5-21%)	300 (110-540)	6700 (2500-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	2010	Male	70 to 74	50% (32-66%)	50% (34-68%)	1.8 (0.8-3.2)	8.8% (2.9-16%)	9.8% (3.3-18%)	230 (76-430)	4100 (1400-7700)
Central African Republic	2010	Male	75 to 79	48% (32-65%)	52% (35-68%)	1.5 (0.64-2.8)	6.7% (1.7-13%)	7.5% (1.8-15%)	150 (37-300)	2000 (520-4100)
Central African Republic	2010	Male	80 to 84	42% (25-60%)	58% (40-75%)	1.1 (0.49-2.1)	4.7% (1.1-9.8%)	5.4% (1.2-11%)	67 (14-140)	680 (150-1400)
Central African Republic	2010	Male	85 to 89	37% (22-54%)	63% (46-78%)	0.84 (0.36-1.6)	3.2% (0.64-7%)	3.7% (0.71-8%)	19 (3.7-42)	140 (28-310)
Central African Republic	2010	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.72 (0.3-1.4)	2.1% (0.23-5%)	2.4% (0.18-5.9%)	3.7 (0.28-8.8)	19 (2.1-44)
Central African Republic	2010	Male	95 plus	40% (21-60%)	60% (40-79%)	0.68 (0.24-1.4)	2.3% (0.1-5.7%)	2.6% (-0.0031-6.7%)	0.95 (-0.0011-2.5)	3.3 (0.15-8.4)
Central African Republic	2016	Female	All Ages	23% (19-29%)	77% (71-81%)	1 (0.54-1.7)	3.4% (1.7-5.7%)	5.7% (2.5-9.5%)	2100 (920-3700)	71000 (34000-120000)
Central African Republic	2016	Female	15 to 19	21% (13-34%)	79% (66-87%)	0.76 (0.3-1.5)	2.2% (0.78-4.5%)	2.8% (0.78-6.2%)	21 (6.2-47)	1700 (660-3600)
Central African Republic	2016	Female	20 to 24	30% (18-45%)	70% (55-82%)	1.5 (0.6-2.8)	5.4% (2.2-9.7%)	6.9% (2.5-13%)	59 (21-110)	4400 (1900-7900)
Central African Republic	2016	Female	25 to 29	34% (20-51%)	66% (49-80%)	1.9 (0.87-3.7)	6% (2.9-10%)	7.4% (3.2-13%)	89 (36-160)	5900 (2700-10000)
Central African Republic	2016	Female	30 to 34	36% (22-53%)	64% (47-78%)	2.1 (0.94-3.9)	6.4% (3.1-11%)	7.7% (3.6-13%)	120 (51-210)	6800 (3300-12000)
Central African Republic	2016	Female	35 to 39	38% (24-55%)	62% (45-76%)	2.1 (0.95-4)	6.2% (2.9-11%)	7.4% (3.3-13%)	120 (52-230)	6400 (2800-12000)
Central African Republic	2016	Female	40 to 44	38% (24-56%)	62% (44-76%)	2.1 (0.92-4.1)	6.7% (3.2-11%)	8% (3.7-13%)	130 (53-230)	5900 (2600-11000)
Central African Republic	2016	Female	45 to 49	37% (22-54%)	63% (46-78%)	2 (0.85-4.1)	8.5% (3.8-15%)	10% (4.6-18%)	150 (64-290)	6300 (2700-12000)
Central African Republic	2016	Female	50 to 54	35% (20-53%)	65% (47-80%)	1.8 (0.75-3.7)	9.9% (4.5-17%)	12% (5.4-20%)	190 (77-350)	6900 (2900-13000)
Central African Republic	2016	Female	55 to 59	32% (20-50%)	68% (50-80%)	1.7 (0.67-3.6)	11% (4.3-18%)	13% (5.1-22%)	240 (88-450)	7400 (2800-14000)
Central African Republic	2016	Female	60 to 64	32% (19-48%)	68% (52-81%)	1.6 (0.63-3.4)	9.4% (3.3-17%)	11% (3.9-19%)	230 (76-430)	6100 (2000-11000)
Central African Republic	2016	Female	65 to 69	33% (21-49%)	67% (51-79%)	1.6 (0.64-3.3)	8.9% (2.8-16%)	10% (3.2-19%)	240 (73-470)	5300 (1600-10000)
Central African Republic	2016	Female	70 to 74	33% (22-48%)	67% (52-78%)	1.4 (0.53-3)	7.5% (1.8-15%)	8.7% (2.1-17%)	230 (58-460)	4000 (1000-8200)
Central African Republic	2016	Female	75 to 79	30% (19-45%)	70% (55-81%)	1.1 (0.42-2.4)	5.7% (1-12%)	6.5% (1.2-14%)	160 (30-360)	2200 (410-5000)
Central African Republic	2016	Female	80 to 84	28% (17-43%)	72% (57-83%)	0.93 (0.36-1.9)	4.5% (0.74-10%)	5.2% (0.86-12%)	99 (17-230)	1000 (160-2300)
Central African Republic	2016	Female	85 to 89	26% (16-40%)	74% (60-84%)	0.82 (0.32-1.7)	3.5% (0.38-8%)	4.1% (0.46-9.4%)	38 (4.5-90)	270 (33-640)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Central African Republic	2016	Female	90 to 94	28% (17-41%)	72% (59-83%)	0.8 (0.3-1.7)	2.9% (0.28-7.2%)	3.4% (0.33-8.7%)	10 (0.97-27)	51 (5.1-130)
Central African Republic	2016	Female	95 plus	31% (17-50%)	69% (50-83%)	0.81 (0.27-1.8)	2.5% (0.14-6.8%)	3% (0.12-8%)	2.3 (0.092-6.1)	7.6 (0.41-20)
Central African Republic	2016	Male	All Ages	31% (25-39%)	69% (61-75%)	1.3 (0.72-2.2)	6.8% (3.9-10%)	9.9% (5.3-15%)	4200 (2100-6500)	160000 (92000-240000)
Central African Republic	2016	Male	15 to 19	24% (14-39%)	76% (61-86%)	0.87 (0.32-1.8)	4.4% (1.5-8.6%)	5.3% (1.6-11%)	58 (18-120)	4500 (1600-8800)
Central African Republic	2016	Male	20 to 24	37% (22-53%)	63% (47-78%)	1.8 (0.76-3.5)	12% (5.6-19%)	14% (6.2-23%)	180 (79-320)	13000 (6000-21000)
Central African Republic	2016	Male	25 to 29	44% (27-62%)	56% (38-73%)	2.6 (1.2-4.7)	16% (8.7-23%)	18% (9.7-28%)	280 (150-430)	18000 (9600-27000)
Central African Republic	2016	Male	30 to 34	46% (28-64%)	54% (36-72%)	2.9 (1.3-5.1)	16% (9-24%)	19% (10-28%)	330 (180-500)	19000 (10000-28000)
Central African Republic	2016	Male	35 to 39	46% (29-64%)	54% (36-71%)	2.8 (1.3-5.3)	16% (8.5-23%)	18% (9.7-27%)	350 (180-550)	18000 (9700-28000)
Central African Republic	2016	Male	40 to 44	48% (32-67%)	52% (33-68%)	2.9 (1.3-5.3)	16% (8.6-24%)	18% (9.8-27%)	390 (190-630)	18000 (9100-29000)
Central African Republic	2016	Male	45 to 49	51% (33-70%)	49% (30-67%)	3.1 (1.4-5.9)	17% (9.2-25%)	20% (10-29%)	430 (210-680)	18000 (8900-28000)
Central African Republic	2016	Male	50 to 54	52% (33-69%)	48% (31-67%)	3 (1.3-6.1)	18% (9-27%)	20% (10-30%)	460 (210-740)	16000 (7400-26000)
Central African Republic	2016	Male	55 to 59	49% (31-68%)	51% (32-69%)	2.6 (1.1-5.3)	17% (7.5-27%)	19% (8.5-30%)	440 (190-730)	14000 (6000-23000)
Central African Republic	2016	Male	60 to 64	48% (30-68%)	52% (32-70%)	2.3 (0.92-4.8)	13% (5.3-23%)	15% (5.9-26%)	370 (140-640)	10000 (3800-17000)
Central African Republic	2016	Male	65 to 69	49% (31-66%)	51% (34-69%)	2 (0.86-4.1)	12% (3.7-21%)	13% (4.1-23%)	330 (110-590)	7200 (2400-13000)
Central African Republic	2016	Male	70 to 74	49% (31-66%)	51% (34-69%)	1.8 (0.72-3.7)	9.2% (2.4-18%)	10% (2.7-20%)	260 (65-510)	4700 (1200-9000)
Central African Republic	2016	Male	75 to 79	47% (31-65%)	53% (35-69%)	1.5 (0.56-3)	7% (1.5-15%)	7.9% (1.6-17%)	170 (34-370)	2400 (470-5100)
Central African Republic	2016	Male	80 to 84	42% (26-60%)	58% (40-74%)	1.1 (0.45-2.3)	5% (0.84-11%)	5.6% (0.9-13%)	79 (13-180)	810 (130-1800)
Central African Republic	2016	Male	85 to 89	36% (22-53%)	64% (47-78%)	0.84 (0.34-1.8)	3.5% (0.54-8.1%)	3.9% (0.55-9.3%)	24 (3.2-57)	170 (26-410)
Central African Republic	2016	Male	90 to 94	36% (22-52%)	64% (48-78%)	0.72 (0.27-1.6)	2.3% (0.23-5.9%)	2.6% (0.19-6.8%)	4.8 (0.33-12)	25 (2.5-63)
Central African Republic	2016	Male	95 plus	39% (21-60%)	61% (40-79%)	0.67 (0.22-1.5)	2.4% (0.1-6.4%)	2.7% (0.013-7.4%)	1.2 (0.0059-3.4)	4.3 (0.18-11)
Congo	1990	Female	All Ages	25% (20-29%)	75% (71-80%)	1 (0.81-1.3)	3.5% (2.4-4.8%)	5.7% (3.8-8%)	850 (550-1200)	29000 (20000-40000)
Congo	1990	Female	15 to 19	22% (15-32%)	78% (68-85%)	0.78 (0.43-1.3)	2% (0.95-3.6%)	2.7% (0.96-5.2%)	6.9 (2.4-13)	610 (290-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	1990	Female	20 to 24	34% (22-46%)	66% (54-78%)	1.6 (0.91-2.4)	4.3% (2.2-7%)	5.2% (2.1-9.1%)	19 (7.5-33)	1500 (770-2500)
Congo	1990	Female	25 to 29	40% (28-52%)	60% (48-72%)	2.3 (1.5-3.4)	6.2% (3.2-9.4%)	7.5% (3.4-12%)	32 (14-52)	2200 (1100-3400)
Congo	1990	Female	30 to 34	44% (30-57%)	56% (43-70%)	2.7 (1.7-3.9)	7.8% (4.1-12%)	9.8% (4.7-15%)	44 (20-69)	2700 (1300-4100)
Congo	1990	Female	35 to 39	44% (31-57%)	56% (43-69%)	2.6 (1.5-3.9)	8.8% (4.8-13%)	11% (5.8-16%)	55 (28-88)	2900 (1500-4700)
Congo	1990	Female	40 to 44	43% (31-57%)	57% (43-69%)	2.4 (1.4-3.8)	9.1% (5.1-14%)	11% (6.1-17%)	59 (31-91)	2800 (1500-4300)
Congo	1990	Female	45 to 49	43% (28-55%)	57% (45-72%)	2.2 (1.3-3.4)	9.4% (5.2-14%)	12% (6.3-17%)	68 (37-100)	2900 (1600-4400)
Congo	1990	Female	50 to 54	39% (27-52%)	61% (48-73%)	1.9 (1.1-3.1)	9.5% (5.3-14%)	11% (6.3-17%)	81 (43-130)	3000 (1600-4600)
Congo	1990	Female	55 to 59	35% (24-47%)	65% (53-76%)	1.7 (0.93-2.9)	8.9% (4.8-14%)	11% (5.6-17%)	88 (45-150)	2800 (1400-4600)
Congo	1990	Female	60 to 64	33% (22-46%)	67% (54-78%)	1.7 (0.86-2.9)	7.7% (3.9-13%)	9% (4.5-15%)	87 (44-150)	2300 (1200-3900)
Congo	1990	Female	65 to 69	34% (24-45%)	66% (55-76%)	1.7 (0.96-2.7)	7.5% (4-12%)	8.6% (4.4-14%)	98 (50-160)	2200 (1100-3600)
Congo	1990	Female	70 to 74	34% (24-45%)	66% (55-76%)	1.6 (0.88-2.6)	6.7% (3.1-11%)	7.6% (3.5-13%)	95 (43-160)	1700 (770-2900)
Congo	1990	Female	75 to 79	33% (22-48%)	67% (52-78%)	1.4 (0.71-2.3)	5.2% (1.8-9.8%)	5.9% (2.1-11%)	68 (23-130)	940 (320-1800)
Congo	1990	Female	80 to 84	31% (19-45%)	69% (55-81%)	1.1 (0.57-1.9)	3.8% (1.2-7.6%)	4.3% (1.3-8.6%)	40 (12-78)	410 (130-800)
Congo	1990	Female	85 to 89	29% (18-41%)	71% (59-82%)	0.96 (0.49-1.7)	2.7% (0.68-5.8%)	3.1% (0.75-6.7%)	11 (2.8-25)	84 (21-180)
Congo	1990	Female	90 to 94	30% (19-42%)	70% (58-81%)	0.9 (0.45-1.6)	2% (0.29-4.6%)	2.3% (0.31-5.4%)	2.3 (0.31-5.4)	12 (1.7-27)
Congo	1990	Female	95 plus	33% (19-49%)	67% (51-81%)	0.87 (0.35-1.7)	1.6% (-0.0072-4.2%)	1.8% (-0.072-4.8%)	0.4 (-0.017-1.1)	1.4 (-0.0059-3.7)
Congo	1990	Male	All Ages	33% (28-40%)	67% (60-72%)	1.5 (1.2-1.8)	6.1% (4.4-7.9%)	9% (6.3-12%)	1500 (1000-2100)	57000 (40000-76000)
Congo	1990	Male	15 to 19	26% (17-37%)	74% (63-83%)	1 (0.52-1.7)	4% (1.9-6.7%)	5% (2.1-8.9%)	18 (7.2-32)	1400 (680-2500)
Congo	1990	Male	20 to 24	40% (27-52%)	60% (48-73%)	2.2 (1.3-3.4)	9.4% (5-14%)	11% (5.2-17%)	57 (27-93)	4200 (2200-6500)
Congo	1990	Male	25 to 29	50% (36-63%)	50% (37-64%)	3.4 (2.1-4.8)	12% (6.3-17%)	14% (6.7-21%)	88 (44-130)	5800 (3100-8300)
Congo	1990	Male	30 to 34	53% (37-65%)	47% (35-63%)	3.7 (2.3-5.4)	13% (7.5-18%)	15% (8-22%)	100 (56-150)	6000 (3500-8500)
Congo	1990	Male	35 to 39	49% (35-63%)	51% (37-65%)	3.4 (2-5.1)	13% (7.8-19%)	15% (8.5-23%)	110 (58-160)	5600 (3200-8200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	1990	Male	40 to 44	51% (38-64%)	49% (36-62%)	3.3 (1.9-5.1)	15% (8.8-21%)	17% (9.9-24%)	130 (71-180)	5900 (3400-8600)
Congo	1990	Male	45 to 49	56% (40-69%)	44% (31-60%)	3.6 (2.2-5.6)	17% (10-23%)	19% (12-27%)	150 (89-220)	6400 (3800-9100)
Congo	1990	Male	50 to 54	58% (44-70%)	42% (30-56%)	3.5 (2.1-5.5)	17% (11-24%)	20% (12-27%)	170 (100-250)	6400 (3800-9100)
Congo	1990	Male	55 to 59	55% (40-68%)	45% (32-60%)	3 (1.7-4.8)	15% (8.9-21%)	17% (10-25%)	160 (93-250)	5200 (3000-7800)
Congo	1990	Male	60 to 64	52% (37-67%)	48% (33-63%)	2.6 (1.5-4.5)	12% (6.9-18%)	14% (7.8-21%)	150 (79-230)	3900 (2200-6300)
Congo	1990	Male	65 to 69	53% (38-66%)	47% (34-62%)	2.4 (1.4-3.7)	11% (5.9-16%)	12% (6.7-19%)	130 (73-210)	3000 (1600-4700)
Congo	1990	Male	70 to 74	54% (40-68%)	46% (32-60%)	2.2 (1.3-3.4)	8.7% (4.2-14%)	9.7% (4.6-16%)	110 (51-180)	1900 (920-3200)
Congo	1990	Male	75 to 79	54% (37-70%)	46% (30-63%)	1.9 (1-3.1)	6.8% (2.6-12%)	7.7% (2.9-13%)	73 (28-130)	1000 (390-1800)
Congo	1990	Male	80 to 84	49% (31-66%)	51% (34-69%)	1.5 (0.79-2.4)	4.7% (1.8-8.6%)	5.3% (2-9.7%)	33 (12-62)	340 (130-640)
Congo	1990	Male	85 to 89	42% (27-59%)	58% (41-73%)	1.1 (0.55-1.8)	3% (1-5.6%)	3.4% (1.1-6.4%)	9.9 (3.2-19)	74 (25-140)
Congo	1990	Male	90 to 94	41% (27-57%)	59% (43-73%)	0.86 (0.43-1.5)	1.8% (0.32-3.7%)	2% (0.25-4.2%)	1.8 (0.23-3.9)	9.7 (1.8-20)
Congo	1990	Male	95 plus	44% (25-63%)	56% (37-75%)	0.74 (0.29-1.4)	1.7% (0.11-4.2%)	1.8% (-0.065-4.8%)	0.4 (-0.015-1.1)	1.5 (0.093-3.8)
Congo	1995	Female	All Ages	25% (21-29%)	75% (71-79%)	0.86 (0.68-1)	2.7% (1.8-3.8%)	4.2% (2.6-6.1%)	780 (480-1200)	28000 (19000-39000)
Congo	1995	Female	15 to 19	23% (15-32%)	77% (68-85%)	0.63 (0.36-1)	1.7% (0.75-3%)	2.1% (0.66-4.2%)	6.5 (2.1-13)	600 (270-1100)
Congo	1995	Female	20 to 24	34% (23-45%)	66% (55-77%)	1.3 (0.76-2)	3.4% (1.7-5.6%)	4% (1.6-7%)	19 (7.8-34)	1600 (780-2600)
Congo	1995	Female	25 to 29	41% (28-53%)	59% (47-72%)	1.9 (1.2-2.7)	4.4% (2.5-7%)	5% (2.4-8.5%)	34 (17-55)	2300 (1300-3600)
Congo	1995	Female	30 to 34	43% (30-57%)	57% (43-70%)	2.1 (1.3-3.1)	5.5% (3.2-8.4%)	6.4% (3.4-10%)	46 (24-71)	2800 (1600-4200)
Congo	1995	Female	35 to 39	44% (30-57%)	56% (43-70%)	2 (1.2-3.1)	6.4% (3.5-9.9%)	7.6% (3.9-12%)	56 (28-91)	3000 (1600-4800)
Congo	1995	Female	40 to 44	43% (31-56%)	57% (44-69%)	2 (1.1-3)	6.8% (3.6-11%)	8.2% (4.3-13%)	60 (30-95)	2800 (1500-4400)
Congo	1995	Female	45 to 49	42% (28-55%)	58% (45-72%)	1.8 (1.1-2.7)	7.2% (4.1-11%)	8.6% (4.8-13%)	66 (36-100)	2800 (1600-4200)
Congo	1995	Female	50 to 54	39% (27-52%)	61% (48-73%)	1.5 (0.87-2.4)	7.4% (3.9-12%)	8.7% (4.7-14%)	77 (40-120)	2800 (1500-4400)
Congo	1995	Female	55 to 59	35% (24-47%)	65% (53-76%)	1.4 (0.74-2.2)	7% (3.3-12%)	8.2% (3.9-14%)	81 (39-140)	2600 (1200-4400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	1995	Female	60 to 64	33% (22-46%)	67% (54-78%)	1.3 (0.71-2.3)	5.9% (2.8-10%)	6.8% (3.3-12%)	77 (36-140)	2100 (970-3700)
Congo	1995	Female	65 to 69	34% (24-45%)	66% (55-76%)	1.3 (0.77-2.1)	5.5% (2.5-9.5%)	6.3% (2.9-11%)	84 (38-150)	1900 (850-3200)
Congo	1995	Female	70 to 74	34% (24-46%)	66% (54-76%)	1.3 (0.72-2.1)	4.9% (1.7-9%)	5.5% (1.9-10%)	80 (29-150)	1400 (520-2700)
Congo	1995	Female	75 to 79	33% (22-48%)	67% (52-78%)	1.1 (0.58-1.9)	3.7% (0.76-7.5%)	4.1% (0.89-8.4%)	56 (12-110)	770 (170-1600)
Congo	1995	Female	80 to 84	31% (19-45%)	69% (55-81%)	0.9 (0.47-1.5)	2.6% (0.5-5.6%)	2.9% (0.57-6.4%)	32 (6.3-69)	320 (64-710)
Congo	1995	Female	85 to 89	29% (18-41%)	71% (59-82%)	0.77 (0.4-1.3)	1.8% (0.21-4%)	2% (0.23-4.6%)	8.9 (0.99-20)	65 (7.9-140)
Congo	1995	Female	90 to 94	30% (19-42%)	70% (58-81%)	0.72 (0.35-1.3)	1.3% (-0.087-3.4%)	1.5% (-0.12-4%)	1.7 (-0.15-4.8)	8.9 (-0.62-24)
Congo	1995	Female	95 plus	33% (19-49%)	67% (51-81%)	0.69 (0.28-1.3)	0.99% (-0.26-3.1%)	1.1% (-0.38-3.6%)	0.29 (-0.098-0.94)	1 (-0.27-3.2)
Congo	1995	Male	All Ages	34% (29-40%)	66% (60-71%)	1.2 (0.99-1.5)	4.8% (3.4-6.3%)	6.9% (4.7-9.3%)	1400 (900-1900)	54000 (38000-72000)
Congo	1995	Male	15 to 19	27% (18-38%)	73% (62-82%)	0.83 (0.45-1.3)	3.5% (1.7-5.7%)	4.3% (1.8-7.5%)	17 (7.2-31)	1400 (710-2400)
Congo	1995	Male	20 to 24	40% (28-53%)	60% (47-72%)	1.8 (1.1-2.7)	8.4% (4.6-13%)	9.7% (4.8-15%)	58 (29-90)	4300 (2400-6500)
Congo	1995	Male	25 to 29	51% (37-64%)	49% (36-63%)	2.8 (1.7-3.9)	9.8% (5.2-15%)	11% (5.2-17%)	89 (47-130)	5900 (3300-8500)
Congo	1995	Male	30 to 34	53% (38-66%)	47% (34-62%)	3.1 (1.9-4.4)	9.8% (5.7-14%)	11% (5.9-16%)	100 (59-150)	6100 (3700-8600)
Congo	1995	Male	35 to 39	50% (36-63%)	50% (37-64%)	2.7 (1.6-4.1)	9.7% (5.7-14%)	11% (6.1-16%)	100 (60-150)	5500 (3300-7900)
Congo	1995	Male	40 to 44	51% (38-64%)	49% (36-62%)	2.7 (1.5-4)	11% (6.4-16%)	12% (6.9-18%)	120 (68-170)	5600 (3300-8200)
Congo	1995	Male	45 to 49	57% (41-69%)	43% (31-59%)	3 (1.8-4.5)	13% (7.8-18%)	14% (8.7-21%)	140 (83-210)	5900 (3500-8600)
Congo	1995	Male	50 to 54	58% (45-71%)	42% (29-55%)	2.9 (1.7-4.5)	14% (8.1-20%)	16% (9.4-23%)	160 (87-240)	5800 (3200-8600)
Congo	1995	Male	55 to 59	55% (41-68%)	45% (32-59%)	2.4 (1.4-3.8)	12% (7-18%)	14% (7.8-21%)	140 (81-220)	4600 (2600-6900)
Congo	1995	Male	60 to 64	53% (38-68%)	47% (32-62%)	2.1 (1.2-3.6)	9.5% (5-15%)	11% (5.6-17%)	130 (65-210)	3400 (1800-5500)
Congo	1995	Male	65 to 69	53% (39-66%)	47% (34-61%)	2 (1.2-3)	8.2% (4.3-13%)	9.3% (4.9-15%)	110 (57-180)	2500 (1300-4100)
Congo	1995	Male	70 to 74	54% (40-67%)	46% (33-60%)	1.8 (1.1-2.7)	6.6% (2.8-11%)	7.5% (3.2-13%)	92 (39-160)	1700 (700-2800)
Congo	1995	Male	75 to 79	54% (38-69%)	46% (31-62%)	1.5 (0.87-2.5)	5.1% (1.5-9.5%)	5.8% (1.7-11%)	61 (18-120)	860 (250-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	1995	Male	80 to 84	49% (32-67%)	51% (33-68%)	1.2 (0.62-1.9)	3.4% (0.86-6.8%)	3.9% (0.91-7.8%)	28 (6.6-55)	290 (73-570)
Congo	1995	Male	85 to 89	43% (28-59%)	57% (41-72%)	0.85 (0.46-1.4)	2.2% (0.49-4.3%)	2.5% (0.51-4.9%)	8.2 (1.7-16)	61 (14-120)
Congo	1995	Male	90 to 94	41% (27-57%)	59% (43-73%)	0.68 (0.36-1.2)	1.2% (0.016-2.8%)	1.3% (-0.11-3.2%)	1.5 (-0.12-3.5)	7.9 (0.1-18)
Congo	1995	Male	95 plus	44% (25-64%)	56% (36-75%)	0.59 (0.23-1.1)	1.2% (-0.15-3.1%)	1.2% (-0.37-3.6%)	0.33 (-0.1-0.98)	1.3 (-0.17-3.4)
Congo	2000	Female	All Ages	24% (20-27%)	76% (73-80%)	0.78 (0.62-0.95)	2.2% (1.5-3.1%)	3.4% (2.1-4.9%)	740 (440-1100)	27000 (18000-37000)
Congo	2000	Female	15 to 19	23% (16-30%)	77% (70-84%)	0.59 (0.36-0.91)	1.5% (0.77-2.5%)	1.8% (0.73-3.5%)	6.5 (2.7-12)	610 (330-1100)
Congo	2000	Female	20 to 24	33% (25-42%)	67% (58-75%)	1.2 (0.75-1.8)	3.1% (1.7-5%)	3.5% (1.7-6.1%)	20 (9.3-33)	1600 (910-2600)
Congo	2000	Female	25 to 29	39% (30-48%)	61% (52-70%)	1.7 (1.1-2.4)	3.6% (2.1-5.7%)	4% (2.1-6.8%)	34 (18-54)	2400 (1500-3700)
Congo	2000	Female	30 to 34	42% (31-52%)	58% (48-69%)	1.9 (1.3-2.7)	4.2% (2.4-6.3%)	4.6% (2.4-7.2%)	45 (23-71)	2800 (1500-4200)
Congo	2000	Female	35 to 39	41% (31-53%)	59% (47-69%)	1.8 (1.1-2.7)	4.8% (2.8-7.5%)	5.5% (2.9-8.9%)	54 (30-85)	2900 (1700-4500)
Congo	2000	Female	40 to 44	40% (30-51%)	60% (49-70%)	1.7 (1-2.6)	5.2% (3-8%)	6% (3.4-9.5%)	57 (31-90)	2700 (1500-4200)
Congo	2000	Female	45 to 49	40% (28-51%)	60% (49-72%)	1.6 (0.96-2.4)	5.8% (3.4-8.9%)	6.9% (3.9-11%)	64 (36-100)	2700 (1500-4200)
Congo	2000	Female	50 to 54	36% (25-47%)	64% (53-75%)	1.3 (0.77-2.1)	6.1% (3.3-9.6%)	7.2% (3.8-11%)	72 (37-120)	2700 (1400-4200)
Congo	2000	Female	55 to 59	30% (21-41%)	70% (59-79%)	1.1 (0.61-1.8)	5.6% (2.8-9.4%)	6.6% (3.3-11%)	73 (35-130)	2300 (1100-4000)
Congo	2000	Female	60 to 64	28% (18-39%)	72% (61-82%)	1 (0.56-1.8)	4.7% (2.1-8.4%)	5.4% (2.4-9.7%)	67 (30-120)	1800 (810-3200)
Congo	2000	Female	65 to 69	30% (21-40%)	70% (60-79%)	1.1 (0.66-1.8)	4.5% (2-7.8%)	5.2% (2.4-8.9%)	75 (34-130)	1700 (760-2900)
Congo	2000	Female	70 to 74	33% (24-43%)	67% (57-76%)	1.2 (0.66-1.8)	4.1% (1.4-7.9%)	4.7% (1.6-8.9%)	74 (24-140)	1300 (430-2500)
Congo	2000	Female	75 to 79	33% (22-47%)	67% (53-78%)	1 (0.57-1.7)	3.2% (0.54-6.5%)	3.6% (0.62-7.4%)	53 (9.2-110)	730 (130-1500)
Congo	2000	Female	80 to 84	31% (19-44%)	69% (56-81%)	0.84 (0.45-1.4)	2.2% (0.38-4.9%)	2.5% (0.46-5.6%)	30 (5.5-66)	310 (54-680)
Congo	2000	Female	85 to 89	28% (17-41%)	72% (59-83%)	0.7 (0.36-1.2)	1.5% (0.13-3.5%)	1.7% (0.15-4.1%)	8.5 (0.72-20)	62 (5.6-150)
Congo	2000	Female	90 to 94	29% (19-42%)	71% (58-81%)	0.65 (0.33-1.1)	1% (-0.12-2.9%)	1.2% (-0.17-3.4%)	1.7 (-0.24-4.7)	8.5 (-0.97-23)
Congo	2000	Female	95 plus	32% (19-48%)	68% (52-81%)	0.63 (0.25-1.2)	0.82% (-0.32-2.8%)	0.9% (-0.42-3.2%)	0.28 (-0.13-1)	1 (-0.38-3.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	2000	Male	All Ages	36% (30-42%)	64% (58-70%)	1.3 (1-1.5)	4.1% (2.9-5.3%)	5.9% (4.1-8%)	1300 (850-1700)	51000 (37000-68000)
Congo	2000	Male	15 to 19	28% (19-37%)	72% (63-81%)	0.82 (0.47-1.3)	3% (1.6-4.9%)	3.7% (1.6-6.2%)	16 (7.4-29)	1400 (740-2300)
Congo	2000	Male	20 to 24	43% (32-54%)	57% (46-68%)	1.8 (1.1-2.8)	7.9% (4.7-12%)	9.1% (4.9-14%)	62 (34-95)	4700 (2700-6900)
Congo	2000	Male	25 to 29	56% (44-65%)	44% (35-56%)	2.9 (1.9-4)	9.9% (5.9-14%)	11% (6.2-17%)	98 (58-140)	6600 (4100-9200)
Congo	2000	Male	30 to 34	55% (43-66%)	45% (34-57%)	3 (2-4.2)	9.1% (5.5-13%)	10% (5.7-15%)	110 (64-150)	6500 (4000-8800)
Congo	2000	Male	35 to 39	48% (38-60%)	52% (40-62%)	2.5 (1.6-3.7)	8% (4.8-12%)	8.9% (5.1-14%)	99 (60-150)	5300 (3400-7700)
Congo	2000	Male	40 to 44	50% (40-62%)	50% (38-60%)	2.5 (1.5-3.8)	8.8% (5.3-13%)	9.9% (5.9-15%)	110 (64-160)	5100 (3100-7400)
Congo	2000	Male	45 to 49	60% (47-70%)	40% (30-53%)	3 (1.9-4.4)	11% (6.8-15%)	12% (7.6-18%)	130 (77-190)	5400 (3300-7900)
Congo	2000	Male	50 to 54	61% (49-72%)	39% (28-51%)	2.9 (1.8-4.4)	12% (7.1-17%)	14% (8.2-19%)	130 (76-200)	5000 (2800-7300)
Congo	2000	Male	55 to 59	55% (41-67%)	45% (33-59%)	2.3 (1.4-3.7)	10% (5.8-16%)	12% (6.9-18%)	120 (65-190)	3800 (2100-5900)
Congo	2000	Male	60 to 64	51% (37-66%)	49% (34-63%)	2 (1.1-3.2)	8.3% (4.5-13%)	9.7% (5.1-16%)	100 (54-180)	2800 (1500-4700)
Congo	2000	Male	65 to 69	53% (40-66%)	47% (34-60%)	1.9 (1.2-2.9)	7.3% (3.9-12%)	8.4% (4.5-13%)	96 (49-160)	2200 (1100-3600)
Congo	2000	Male	70 to 74	57% (44-69%)	43% (31-56%)	1.8 (1.1-2.7)	6.2% (2.7-10%)	7.1% (3-12%)	84 (36-150)	1500 (650-2600)
Congo	2000	Male	75 to 79	60% (45-73%)	40% (27-55%)	1.7 (0.98-2.6)	4.9% (1.5-9%)	5.6% (1.7-10%)	58 (18-110)	810 (240-1500)
Congo	2000	Male	80 to 84	52% (35-69%)	48% (31-65%)	1.2 (0.68-2)	3.2% (0.76-6.3%)	3.7% (0.81-7.3%)	26 (5.8-54)	280 (65-550)
Congo	2000	Male	85 to 89	42% (27-59%)	58% (41-73%)	0.81 (0.44-1.4)	2% (0.47-4.1%)	2.2% (0.48-4.7%)	7.7 (1.7-16)	58 (15-120)
Congo	2000	Male	90 to 94	41% (26-58%)	59% (42-74%)	0.66 (0.34-1.1)	1.1% (0.021-2.6%)	1.2% (-0.09-3%)	1.4 (-0.11-3.5)	7.8 (0.15-19)
Congo	2000	Male	95 plus	43% (24-65%)	57% (35-76%)	0.57 (0.23-1.1)	1.1% (-0.13-3%)	1.1% (-0.35-3.5%)	0.34 (-0.1-1)	1.3 (-0.16-3.7)
Congo	2005	Female	All Ages	23% (19-26%)	77% (74-81%)	0.81 (0.64-0.99)	2.1% (1.4-2.9%)	3.2% (1.9-4.7%)	680 (390-1000)	25000 (16000-34000)
Congo	2005	Female	15 to 19	22% (16-29%)	78% (71-84%)	0.61 (0.35-0.95)	1.3% (0.73-2.1%)	1.5% (0.71-2.8%)	5.7 (2.6-10)	570 (320-920)
Congo	2005	Female	20 to 24	32% (24-41%)	68% (59-76%)	1.2 (0.77-1.8)	3% (1.8-4.6%)	3.4% (1.8-5.7%)	16 (8.5-27)	1500 (900-2200)
Congo	2005	Female	25 to 29	38% (29-47%)	62% (53-71%)	1.8 (1.2-2.5)	3.5% (2.1-5.3%)	3.9% (2-6.4%)	30 (17-47)	2200 (1400-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	2005	Female	30 to 34	40% (30-51%)	60% (49-70%)	2 (1.4-2.8)	3.7% (2.4-5.4%)	4.1% (2.4-6.3%)	40 (24-61)	2600 (1600-3700)
Congo	2005	Female	35 to 39	40% (29-52%)	60% (48-71%)	1.9 (1.1-2.8)	3.9% (2.4-5.9%)	4.4% (2.5-7%)	46 (26-70)	2500 (1500-3800)
Congo	2005	Female	40 to 44	39% (28-50%)	61% (50-72%)	1.8 (1.1-2.7)	4.4% (2.7-6.6%)	5.1% (3-7.8%)	50 (28-77)	2400 (1400-3700)
Congo	2005	Female	45 to 49	39% (27-50%)	61% (50-73%)	1.6 (0.98-2.5)	5.1% (3-7.6%)	6% (3.6-9.2%)	58 (33-90)	2500 (1400-3800)
Congo	2005	Female	50 to 54	35% (25-47%)	65% (53-75%)	1.4 (0.79-2.2)	5.5% (3.1-8.8%)	6.6% (3.6-10%)	67 (36-110)	2500 (1300-4000)
Congo	2005	Female	55 to 59	29% (20-40%)	71% (60-80%)	1.2 (0.61-1.9)	5.2% (2.6-8.7%)	6.2% (3.1-10%)	68 (34-120)	2200 (1100-3600)
Congo	2005	Female	60 to 64	27% (18-37%)	73% (63-82%)	1.1 (0.56-1.9)	4.4% (2.1-7.7%)	5.1% (2.4-9%)	64 (29-110)	1700 (780-3000)
Congo	2005	Female	65 to 69	29% (20-39%)	71% (61-80%)	1.1 (0.67-1.8)	4.3% (2-7.4%)	5% (2.4-8.7%)	72 (34-130)	1600 (760-2800)
Congo	2005	Female	70 to 74	31% (22-42%)	69% (58-78%)	1.2 (0.67-1.9)	4% (1.3-7.5%)	4.6% (1.6-8.6%)	72 (23-130)	1300 (420-2400)
Congo	2005	Female	75 to 79	32% (21-45%)	68% (55-79%)	1.1 (0.56-1.8)	3.1% (0.59-6.5%)	3.5% (0.68-7.4%)	53 (9.8-110)	730 (130-1500)
Congo	2005	Female	80 to 84	29% (18-43%)	71% (57-82%)	0.86 (0.44-1.5)	2.2% (0.36-4.8%)	2.5% (0.43-5.5%)	32 (5.4-71)	330 (56-730)
Congo	2005	Female	85 to 89	27% (16-40%)	73% (60-84%)	0.71 (0.36-1.2)	1.4% (0.15-3.4%)	1.6% (0.18-3.8%)	8.8 (0.99-21)	65 (7.1-150)
Congo	2005	Female	90 to 94	28% (18-40%)	72% (60-82%)	0.66 (0.33-1.2)	0.99% (-0.12-2.7%)	1.1% (-0.17-3.1%)	1.7 (-0.26-4.7)	8.8 (-1.1-24)
Congo	2005	Female	95 plus	30% (18-47%)	70% (53-82%)	0.63 (0.26-1.2)	0.8% (-0.21-2.5%)	0.87% (-0.31-2.9%)	0.29 (-0.1-0.97)	1 (-0.27-3.4)
Congo	2005	Male	All Ages	35% (30-41%)	65% (59-70%)	1.3 (1.1-1.6)	3.5% (2.5-4.6%)	5.2% (3.6-7%)	960 (650-1300)	40000 (29000-53000)
Congo	2005	Male	15 to 19	27% (19-36%)	73% (64-81%)	0.85 (0.46-1.4)	2.7% (1.6-4.3%)	3.3% (1.7-5.6%)	13 (6.2-22)	1200 (660-1800)
Congo	2005	Male	20 to 24	42% (31-53%)	58% (47-69%)	1.9 (1.2-2.9)	7.9% (5-11%)	9.6% (5.5-15%)	46 (26-73)	3600 (2300-5300)
Congo	2005	Male	25 to 29	55% (44-65%)	45% (35-56%)	3 (2-4.2)	9.9% (6.3-14%)	12% (6.8-18%)	77 (47-110)	5300 (3500-7600)
Congo	2005	Male	30 to 34	54% (42-65%)	46% (35-58%)	3.1 (2.1-4.4)	8.7% (5.6-12%)	10% (6.1-15%)	84 (51-120)	5200 (3400-7400)
Congo	2005	Male	35 to 39	47% (36-58%)	53% (42-64%)	2.6 (1.6-3.8)	7.1% (4.4-10%)	8% (4.8-12%)	76 (45-110)	4200 (2600-6100)
Congo	2005	Male	40 to 44	49% (38-61%)	51% (39-62%)	2.6 (1.6-3.9)	7.2% (4.7-10%)	8.2% (5.2-12%)	79 (48-120)	3800 (2400-5600)
Congo	2005	Male	45 to 49	59% (47-69%)	41% (31-53%)	3.1 (1.9-4.7)	9% (5.8-13%)	11% (6.7-15%)	93 (58-140)	4000 (2600-5800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	2005	Male	50 to 54	60% (48-71%)	40% (29-52%)	3 (1.8-4.7)	10% (6.3-14%)	12% (7.6-18%)	100 (61-150)	3800 (2300-5400)
Congo	2005	Male	55 to 59	54% (40-67%)	46% (33-60%)	2.4 (1.4-3.8)	9.1% (5.5-14%)	11% (6.6-17%)	89 (52-140)	2900 (1700-4400)
Congo	2005	Male	60 to 64	51% (37-65%)	49% (35-63%)	2 (1.2-3.4)	7.4% (4.1-12%)	8.9% (4.8-14%)	80 (43-130)	2200 (1200-3500)
Congo	2005	Male	65 to 69	52% (39-65%)	48% (35-61%)	2 (1.2-3)	6.6% (3.6-10%)	7.8% (4.2-12%)	76 (42-120)	1700 (960-2800)
Congo	2005	Male	70 to 74	57% (43-68%)	43% (32-57%)	1.9 (1.2-3)	5.6% (2.7-9.3%)	6.6% (3.1-11%)	67 (31-110)	1200 (570-2000)
Congo	2005	Male	75 to 79	59% (44-72%)	41% (28-56%)	1.8 (1-2.8)	4.5% (1.6-8.3%)	5.2% (1.8-9.7%)	48 (17-89)	680 (240-1200)
Congo	2005	Male	80 to 84	52% (34-69%)	48% (31-66%)	1.3 (0.7-2.1)	2.9% (0.9-5.7%)	3.4% (0.97-6.6%)	23 (6.7-47)	240 (76-480)
Congo	2005	Male	85 to 89	42% (26-59%)	58% (41-74%)	0.86 (0.45-1.5)	1.8% (0.51-3.6%)	2.1% (0.51-4.2%)	7.3 (1.8-15)	56 (16-110)
Congo	2005	Male	90 to 94	40% (26-57%)	60% (43-74%)	0.69 (0.36-1.2)	1% (0.078-2.4%)	1.1% (-0.022-2.8%)	1.4 (-0.029-3.4)	7.8 (0.59-18)
Congo	2005	Male	95 plus	43% (25-63%)	57% (37-75%)	0.6 (0.24-1.2)	1% (-0.0042-2.9%)	1.1% (-0.17-3.3%)	0.33 (-0.052-0.99)	1.3 (-0.0051-3.5)
Congo	2010	Female	All Ages	23% (19-27%)	77% (73-81%)	0.88 (0.69-1.1)	2.4% (1.6-3.3%)	3.6% (2.2-5.2%)	760 (420-1100)	27000 (17000-38000)
Congo	2010	Female	15 to 19	20% (13-29%)	80% (71-87%)	0.61 (0.33-1)	1.3% (0.72-2%)	1.4% (0.68-2.7%)	5.6 (2.6-10)	590 (330-940)
Congo	2010	Female	20 to 24	31% (20-43%)	69% (57-80%)	1.3 (0.72-2)	3% (1.9-4.5%)	3.5% (1.8-5.9%)	16 (8.2-26)	1500 (890-2200)
Congo	2010	Female	25 to 29	37% (25-50%)	63% (50-75%)	1.9 (1.2-2.7)	3.8% (2.3-5.5%)	4.4% (2.4-7%)	30 (16-47)	2300 (1400-3300)
Congo	2010	Female	30 to 34	40% (28-54%)	60% (46-72%)	2.1 (1.3-3.1)	4.1% (2.6-6.1%)	4.8% (2.8-7.5%)	43 (24-67)	2800 (1700-4100)
Congo	2010	Female	35 to 39	41% (28-54%)	59% (46-72%)	2 (1.2-3.2)	4.4% (2.6-6.6%)	5.1% (2.9-8%)	51 (27-83)	2900 (1600-4400)
Congo	2010	Female	40 to 44	40% (28-53%)	60% (47-72%)	2 (1.1-3.1)	5% (2.9-7.4%)	5.9% (3.4-8.9%)	58 (31-89)	2800 (1600-4300)
Congo	2010	Female	45 to 49	39% (25-53%)	61% (47-75%)	1.8 (1-2.8)	5.6% (3.3-8.5%)	6.9% (4-11%)	64 (35-100)	2800 (1600-4400)
Congo	2010	Female	50 to 54	36% (24-49%)	64% (51-76%)	1.5 (0.86-2.5)	6% (3.3-9.5%)	7.3% (4-11%)	73 (39-120)	2700 (1500-4500)
Congo	2010	Female	55 to 59	32% (21-44%)	68% (56-79%)	1.4 (0.69-2.3)	5.9% (2.9-9.7%)	7.1% (3.6-12%)	78 (38-140)	2500 (1200-4400)
Congo	2010	Female	60 to 64	31% (20-43%)	69% (57-80%)	1.3 (0.66-2.3)	5% (2.3-8.9%)	6% (2.8-11%)	75 (34-130)	2000 (910-3700)
Congo	2010	Female	65 to 69	31% (21-42%)	69% (58-79%)	1.3 (0.72-2.1)	4.8% (2.1-8.1%)	5.6% (2.6-9.6%)	82 (36-140)	1800 (810-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	2010	Female	70 to 74	31% (21-42%)	69% (58-79%)	1.2 (0.69-2)	4.1% (1.4-7.6%)	4.8% (1.6-8.8%)	78 (25-150)	1400 (460-2700)
Congo	2010	Female	75 to 79	30% (19-44%)	70% (56-81%)	1.1 (0.54-1.9)	3.2% (0.7-6.4%)	3.6% (0.84-7.4%)	57 (13-120)	800 (170-1700)
Congo	2010	Female	80 to 84	28% (17-41%)	72% (59-83%)	0.86 (0.42-1.5)	2.3% (0.42-4.9%)	2.6% (0.5-5.7%)	36 (6.5-77)	370 (69-800)
Congo	2010	Female	85 to 89	26% (16-38%)	74% (62-84%)	0.73 (0.38-1.3)	1.5% (0.23-3.6%)	1.8% (0.23-4.1%)	11 (1.4-26)	83 (11-190)
Congo	2010	Female	90 to 94	27% (17-39%)	73% (61-83%)	0.69 (0.33-1.2)	1.1% (-0.0058-2.7%)	1.3% (-0.045-3.1%)	2.3 (-0.076-5.6)	12 (-0.06-29)
Congo	2010	Female	95 plus	30% (17-46%)	70% (54-83%)	0.67 (0.27-1.3)	0.92% (-0.17-2.6%)	1% (-0.27-2.9%)	0.38 (-0.11-1.2)	1.4 (-0.26-4)
Congo	2010	Male	All Ages	33% (27-39%)	67% (61-73%)	1.3 (1-1.6)	3.6% (2.6-4.9%)	5.4% (3.7-7.4%)	940 (610-1400)	40000 (28000-55000)
Congo	2010	Male	15 to 19	25% (16-35%)	75% (65-84%)	0.84 (0.44-1.4)	2.5% (1.5-3.9%)	3% (1.5-5.1%)	12 (5.7-21)	1100 (660-1800)
Congo	2010	Male	20 to 24	37% (26-50%)	63% (50-74%)	1.8 (1-2.8)	7.4% (4.7-11%)	9% (5.2-14%)	40 (22-64)	3300 (2000-5000)
Congo	2010	Male	25 to 29	49% (35-62%)	51% (38-65%)	2.8 (1.8-4.1)	9.6% (6.5-14%)	12% (7.3-17%)	67 (41-100)	4900 (3200-7000)
Congo	2010	Male	30 to 34	51% (37-65%)	49% (35-63%)	3.2 (2-4.6)	9.3% (6.1-13%)	11% (6.9-17%)	80 (48-120)	5100 (3300-7500)
Congo	2010	Male	35 to 39	47% (34-60%)	53% (40-66%)	2.8 (1.6-4.2)	8.1% (5.3-11%)	9.7% (6-14%)	78 (45-120)	4400 (2800-6400)
Congo	2010	Male	40 to 44	49% (36-62%)	51% (38-64%)	2.8 (1.6-4.2)	8% (5.2-11%)	9.7% (6.1-14%)	80 (47-120)	4000 (2400-5900)
Congo	2010	Male	45 to 49	54% (39-67%)	46% (33-61%)	3.1 (1.8-4.8)	9.3% (5.9-13%)	11% (7-16%)	89 (54-140)	3900 (2400-5900)
Congo	2010	Male	50 to 54	56% (42-69%)	44% (31-58%)	3 (1.7-4.8)	10% (6.3-15%)	13% (8-18%)	96 (56-140)	3600 (2200-5400)
Congo	2010	Male	55 to 59	53% (37-66%)	47% (34-63%)	2.5 (1.4-4)	9.4% (5.7-14%)	12% (7.2-18%)	90 (50-140)	3000 (1700-4500)
Congo	2010	Male	60 to 64	51% (36-66%)	49% (34-64%)	2.2 (1.2-3.8)	7.8% (4.3-12%)	9.6% (5.2-15%)	83 (43-130)	2300 (1200-3700)
Congo	2010	Male	65 to 69	51% (36-65%)	49% (35-64%)	2 (1.2-3.2)	6.8% (3.8-11%)	8.2% (4.5-13%)	79 (42-130)	1800 (970-2900)
Congo	2010	Male	70 to 74	52% (38-65%)	48% (35-62%)	1.9 (1.1-3)	5.4% (2.5-9.2%)	6.5% (3-11%)	67 (31-120)	1200 (570-2100)
Congo	2010	Male	75 to 79	52% (36-67%)	48% (33-64%)	1.6 (0.87-2.6)	4.2% (1.6-7.7%)	5% (1.9-9%)	47 (17-87)	670 (240-1200)
Congo	2010	Male	80 to 84	47% (29-64%)	53% (36-71%)	1.2 (0.63-2)	2.8% (0.9-5.3%)	3.2% (0.98-6.2%)	24 (7-46)	250 (78-480)
Congo	2010	Male	85 to 89	40% (25-57%)	60% (43-75%)	0.86 (0.45-1.5)	1.8% (0.55-3.5%)	2.1% (0.57-4.1%)	8.4 (2.4-17)	65 (19-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	2010	Male	90 to 94	39% (24-56%)	61% (44-76%)	0.7 (0.35-1.2)	1.1% (0.16-2.3%)	1.2% (0.07-2.6%)	1.7 (0.1-3.9)	9.6 (1.4-21)
Congo	2010	Male	95 plus	42% (23-62%)	58% (38-77%)	0.61 (0.24-1.2)	1.1% (0.084-2.7%)	1.1% (-0.074-3%)	0.38 (-0.026-1.1)	1.5 (0.12-3.8)
Congo	2016	Female	All Ages	23% (19-27%)	77% (73-81%)	0.99 (0.62-1.5)	2.7% (1.6-4.1%)	4.2% (2.1-6.7%)	860 (400-1400)	30000 (16000-47000)
Congo	2016	Female	15 to 19	20% (13-29%)	80% (71-87%)	0.7 (0.33-1.2)	1.3% (0.76-2.2%)	1.5% (0.66-2.9%)	6 (2.4-11)	660 (350-1100)
Congo	2016	Female	20 to 24	31% (20-42%)	69% (58-80%)	1.4 (0.72-2.4)	3.1% (1.8-4.8%)	3.5% (1.7-6.2%)	15 (6.3-26)	1500 (840-2300)
Congo	2016	Female	25 to 29	37% (25-49%)	63% (51-75%)	2.1 (1.1-3.4)	3.9% (2.3-5.9%)	4.6% (2.3-7.6%)	27 (13-47)	2200 (1200-3400)
Congo	2016	Female	30 to 34	40% (27-54%)	60% (46-73%)	2.4 (1.3-3.9)	4.4% (2.6-6.9%)	5.3% (2.8-8.5%)	42 (21-72)	2800 (1600-4600)
Congo	2016	Female	35 to 39	41% (28-53%)	59% (47-72%)	2.3 (1.2-3.9)	4.7% (2.7-7.4%)	5.7% (3-9%)	55 (26-95)	3100 (1600-5100)
Congo	2016	Female	40 to 44	40% (28-53%)	60% (47-72%)	2.2 (1.2-3.8)	5.2% (3-8.2%)	6.4% (3.6-10%)	65 (32-110)	3200 (1700-5400)
Congo	2016	Female	45 to 49	39% (25-52%)	61% (48-75%)	2 (1-3.5)	5.8% (3.1-9%)	7.2% (3.9-11%)	72 (36-120)	3100 (1600-5200)
Congo	2016	Female	50 to 54	36% (25-49%)	64% (51-75%)	1.7 (0.86-3.1)	6.2% (3.2-10%)	7.6% (4-13%)	79 (37-140)	3000 (1400-5100)
Congo	2016	Female	55 to 59	32% (21-44%)	68% (56-79%)	1.6 (0.7-2.9)	6.1% (2.8-11%)	7.5% (3.4-13%)	85 (36-160)	2700 (1200-5000)
Congo	2016	Female	60 to 64	30% (20-42%)	70% (58-80%)	1.5 (0.67-2.9)	5.4% (2.1-9.8%)	6.5% (2.6-12%)	83 (32-150)	2300 (880-4200)
Congo	2016	Female	65 to 69	31% (21-41%)	69% (59-79%)	1.5 (0.73-2.6)	5.2% (2.1-9.6%)	6.2% (2.5-11%)	94 (36-180)	2100 (830-4000)
Congo	2016	Female	70 to 74	31% (22-42%)	69% (58-78%)	1.4 (0.73-2.5)	4.7% (1.4-9.1%)	5.5% (1.7-11%)	94 (27-190)	1700 (500-3400)
Congo	2016	Female	75 to 79	30% (19-45%)	70% (55-81%)	1.2 (0.6-2.3)	3.7% (0.75-8%)	4.3% (0.9-9.4%)	73 (15-160)	1000 (210-2200)
Congo	2016	Female	80 to 84	28% (17-42%)	72% (58-83%)	1 (0.45-1.9)	2.7% (0.45-6.2%)	3.2% (0.55-7.2%)	49 (7.8-120)	510 (80-1200)
Congo	2016	Female	85 to 89	26% (16-38%)	74% (62-84%)	0.84 (0.39-1.6)	1.9% (0.23-4.6%)	2.2% (0.28-5.2%)	16 (1.9-40)	120 (15-300)
Congo	2016	Female	90 to 94	27% (17-39%)	73% (61-83%)	0.78 (0.34-1.5)	1.4% (-0.077-3.5%)	1.6% (-0.098-4.1%)	3.7 (-0.23-9.4)	19 (-1.1-48)
Congo	2016	Female	95 plus	30% (17-46%)	70% (54-83%)	0.75 (0.28-1.5)	1.2% (-0.12-3.3%)	1.3% (-0.21-3.7%)	0.64 (-0.11-1.9)	2.3 (-0.24-6.4)
Congo	2016	Male	All Ages	32% (27-38%)	68% (62-73%)	1.5 (0.91-2.2)	4.3% (2.7-6.2%)	6.2% (3.7-9.1%)	1100 (600-1700)	46000 (28000-70000)
Congo	2016	Male	15 to 19	25% (16-35%)	75% (65-84%)	0.93 (0.45-1.7)	2.7% (1.5-4.3%)	3.1% (1.5-5.7%)	14 (5.9-26)	1400 (720-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Congo	2016	Male	20 to 24	37% (25-49%)	63% (51-75%)	2 (1-3.7)	7.6% (4.7-12%)	9.1% (5-15%)	44 (21-78)	3700 (2100-6100)
Congo	2016	Male	25 to 29	48% (35-61%)	52% (39-65%)	3.2 (1.7-5.2)	10% (6.3-14%)	12% (6.9-19%)	70 (38-120)	5100 (3100-7800)
Congo	2016	Male	30 to 34	50% (36-64%)	50% (36-64%)	3.5 (1.9-5.8)	9.9% (6.2-14%)	12% (7.1-17%)	84 (47-140)	5500 (3300-8400)
Congo	2016	Male	35 to 39	47% (33-60%)	53% (40-67%)	3.1 (1.7-5.2)	9% (5.6-13%)	11% (6.3-16%)	90 (47-150)	5200 (2900-8100)
Congo	2016	Male	40 to 44	48% (35-61%)	52% (39-65%)	3.1 (1.7-5.3)	8.9% (5.4-13%)	11% (6.4-15%)	97 (51-160)	4900 (2800-7600)
Congo	2016	Male	45 to 49	54% (39-67%)	46% (33-61%)	3.4 (1.8-5.9)	9.9% (6.1-14%)	12% (7.3-18%)	110 (58-180)	4800 (2700-7500)
Congo	2016	Male	50 to 54	56% (42-69%)	44% (31-58%)	3.3 (1.7-6)	11% (6-15%)	13% (7.5-19%)	110 (58-180)	4300 (2300-6700)
Congo	2016	Male	55 to 59	52% (38-66%)	48% (34-62%)	2.8 (1.4-5.1)	9.7% (5.3-15%)	12% (6.7-18%)	100 (50-180)	3500 (1700-5600)
Congo	2016	Male	60 to 64	50% (35-66%)	50% (34-65%)	2.4 (1.2-4.5)	8.1% (4.2-13%)	10% (5.2-16%)	98 (45-170)	2700 (1300-4700)
Congo	2016	Male	65 to 69	51% (36-64%)	49% (36-64%)	2.3 (1.2-3.8)	7.2% (3.4-12%)	8.8% (4.3-15%)	94 (40-160)	2200 (960-3700)
Congo	2016	Male	70 to 74	52% (38-65%)	48% (35-62%)	2.1 (1.1-3.5)	6% (2.5-10%)	7.1% (3-12%)	83 (33-150)	1500 (620-2800)
Congo	2016	Male	75 to 79	52% (36-67%)	48% (33-64%)	1.8 (0.87-3.2)	4.6% (1.7-9%)	5.5% (1.9-11%)	61 (20-120)	870 (300-1700)
Congo	2016	Male	80 to 84	47% (30-64%)	53% (36-70%)	1.4 (0.64-2.5)	3.1% (0.97-6.3%)	3.7% (1.1-7.6%)	32 (9-67)	340 (100-720)
Congo	2016	Male	85 to 89	40% (25-57%)	60% (43-75%)	0.97 (0.47-1.7)	2.1% (0.62-4.3%)	2.4% (0.65-5%)	12 (3.2-25)	91 (27-190)
Congo	2016	Male	90 to 94	39% (25-55%)	61% (45-75%)	0.78 (0.34-1.5)	1.3% (0.2-2.9%)	1.4% (0.13-3.3%)	2.6 (0.26-6.3)	15 (2.4-34)
Congo	2016	Male	95 plus	41% (24-62%)	59% (38-76%)	0.68 (0.25-1.4)	1.2% (0.14-3.1%)	1.3% (-0.0029-3.5%)	0.6 (-0.0013-1.6)	2.3 (0.26-5.8)
Democratic Republic of the Congo	1990	Female	All Ages	24% (19-29%)	76% (71-81%)	0.54 (0.43-0.68)	1.3% (0.69-2.1%)	2.3% (1.1-3.9%)	5000 (2300-8200)	190000 (110000-290000)
Democratic Republic of the Congo	1990	Female	15 to 19	23% (14-36%)	77% (64-86%)	0.45 (0.22-0.77)	1.2% (0.53-2.3%)	1.6% (0.45-3.4%)	57 (16-120)	5500 (2400-10000)
Democratic Republic of the Congo	1990	Female	20 to 24	32% (20-48%)	68% (52-80%)	0.86 (0.47-1.4)	3.1% (1.5-5.6%)	3.9% (1.4-7.7%)	180 (64-350)	15000 (7100-27000)
Democratic Republic of the Congo	1990	Female	25 to 29	36% (22-52%)	64% (48-78%)	1.1 (0.65-1.8)	4.2% (2.1-7%)	5.4% (2.3-9.7%)	260 (110-470)	19000 (9500-31000)
Democratic Republic of the Congo	1990	Female	30 to 34	39% (24-56%)	61% (44-76%)	1.3 (0.75-2)	5% (2.6-8%)	6.7% (3.2-11%)	330 (150-550)	21000 (11000-33000)
Democratic Republic of the Congo	1990	Female	35 to 39	40% (25-57%)	60% (43-75%)	1.3 (0.7-2)	5% (2.4-8.2%)	6.8% (2.9-11%)	340 (150-590)	19000 (9200-31000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	1990	Female	40 to 44	40% (26-59%)	60% (41-74%)	1.2 (0.66-2)	5% (2.3-8.4%)	6.9% (3.1-12%)	370 (170-640)	18000 (8600-31000)
Democratic Republic of the Congo	1990	Female	45 to 49	39% (24-57%)	61% (43-76%)	1.2 (0.6-2)	5.5% (2.6-9.5%)	7.6% (3.5-13%)	450 (200-800)	19000 (9000-33000)
Democratic Republic of the Congo	1990	Female	50 to 54	37% (23-54%)	63% (46-77%)	1.1 (0.56-1.9)	5.7% (2.3-10%)	7.6% (2.9-14%)	530 (200-970)	20000 (7800-35000)
Democratic Republic of the Congo	1990	Female	55 to 59	34% (21-51%)	66% (49-79%)	0.99 (0.5-1.7)	5.6% (2.1-10%)	7.4% (2.7-13%)	600 (220-1100)	19000 (7100-34000)
Democratic Republic of the Congo	1990	Female	60 to 64	34% (20-51%)	66% (49-80%)	0.94 (0.45-1.7)	4.5% (1.5-9.2%)	5.7% (1.9-12%)	550 (180-1100)	15000 (4900-30000)
Democratic Republic of the Congo	1990	Female	65 to 69	35% (23-51%)	65% (49-77%)	0.92 (0.49-1.6)	3.9% (0.96-7.9%)	4.8% (1.2-9.6%)	540 (130-1100)	12000 (2900-24000)
Democratic Republic of the Congo	1990	Female	70 to 74	35% (23-51%)	65% (49-77%)	0.83 (0.4-1.5)	2.9% (0.22-6.9%)	3.5% (0.28-8.4%)	410 (33-980)	7300 (540-17000)
Democratic Republic of the Congo	1990	Female	75 to 79	32% (20-48%)	68% (52-80%)	0.67 (0.3-1.2)	1.8% (-0.35-5%)	2.2% (-0.42-6%)	230 (-42-600)	3100 (-610-8300)
Democratic Republic of the Congo	1990	Female	80 to 84	29% (18-45%)	71% (55-82%)	0.55 (0.26-0.96)	1.2% (-0.35-3.6%)	1.5% (-0.4-4.3%)	98 (-27-290)	990 (-280-2900)
Democratic Republic of the Congo	1990	Female	85 to 89	28% (17-42%)	72% (58-83%)	0.49 (0.24-0.84)	0.82% (-0.37-2.7%)	1% (-0.47-3.2%)	38 (-18-130)	280 (-130-900)
Democratic Republic of the Congo	1990	Female	90 to 94	30% (18-43%)	70% (57-82%)	0.48 (0.22-0.87)	0.6% (-0.45-2.3%)	0.73% (-0.55-2.9%)	11 (-8-41)	53 (-40-200)
Democratic Republic of the Congo	1990	Female	95 plus	33% (19-50%)	67% (50-81%)	0.48 (0.19-0.98)	0.55% (-0.48-2.3%)	0.63% (-0.61-2.8%)	2.5 (-2.4-11)	8.9 (-7.4-37)
Democratic Republic of the Congo	1990	Male	All Ages	31% (25-38%)	69% (62-75%)	0.67 (0.53-0.84)	1.8% (1.1-2.9%)	3% (1.7-4.7%)	6800 (4100-10000)	30000 (20000-42000)
Democratic Republic of the Congo	1990	Male	15 to 19	25% (15-40%)	75% (60-85%)	0.49 (0.24-0.87)	2% (1-3.4%)	2.4% (0.91-4.8%)	93 (35-180)	8800 (4500-15000)
Democratic Republic of the Congo	1990	Male	20 to 24	38% (23-54%)	62% (46-77%)	1 (0.56-1.6)	5.2% (2.8-8.6%)	6.1% (2.7-11%)	330 (140-570)	26000 (14000-43000)
Democratic Republic of the Congo	1990	Male	25 to 29	45% (28-63%)	55% (37-72%)	1.4 (0.84-2.2)	6.9% (3.8-11%)	8.1% (3.9-13%)	510 (250-820)	36000 (21000-55000)
Democratic Republic of the Congo	1990	Male	30 to 34	47% (29-66%)	53% (34-71%)	1.6 (0.93-2.5)	7.5% (4.2-12%)	8.9% (4.5-14%)	610 (310-950)	38000 (22000-58000)
Democratic Republic of the Congo	1990	Male	35 to 39	48% (31-66%)	52% (34-69%)	1.6 (0.91-2.5)	7.5% (4.2-12%)	9% (4.7-14%)	640 (340-1000)	35000 (20000-55000)
Democratic Republic of the Congo	1990	Male	40 to 44	50% (34-67%)	50% (33-66%)	1.6 (0.89-2.6)	7.9% (4.2-12%)	9.7% (4.8-15%)	730 (360-1200)	35000 (18000-56000)
Democratic Republic of the Congo	1990	Male	45 to 49	53% (34-70%)	47% (30-66%)	1.7 (0.96-2.7)	8.4% (4.6-13%)	10% (5.5-16%)	800 (420-1300)	34000 (19000-53000)
Democratic Republic of the Congo	1990	Male	50 to 54	53% (34-71%)	47% (29-66%)	1.7 (0.91-2.8)	8.3% (4.1-14%)	10% (5-17%)	810 (380-1300)	30000 (15000-49000)
Democratic Republic of the Congo	1990	Male	55 to 59	51% (33-70%)	49% (30-67%)	1.5 (0.79-2.5)	7.3% (3.4-12%)	9% (4.1-15%)	720 (320-1200)	23000 (11000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	1990	Male	60 to 64	49% (31-69%)	51% (31-69%)	1.3 (0.66-2.3)	5.5% (2.3-10%)	6.8% (2.7-12%)	580 (240-1100)	16000 (6800-29000)
Democratic Republic of the Congo	1990	Male	65 to 69	50% (33-67%)	50% (33-67%)	1.2 (0.61-1.9)	4.3% (1.5-8%)	5.2% (1.9-9.5%)	450 (170-850)	10000 (3900-19000)
Democratic Republic of the Congo	1990	Male	70 to 74	50% (33-67%)	50% (33-67%)	1 (0.5-1.7)	3.1% (0.67-6.4%)	3.7% (0.75-7.7%)	310 (64-660)	5700 (1200-12000)
Democratic Republic of the Congo	1990	Male	75 to 79	49% (32-66%)	51% (34-68%)	0.82 (0.39-1.4)	2.1% (0.1-5%)	2.5% (0.064-5.9%)	170 (4.4-410)	2400 (110-5700)
Democratic Republic of the Congo	1990	Male	80 to 84	43% (26-62%)	57% (38-74%)	0.64 (0.3-1.1)	1.4% (-0.0082-3.3%)	1.6% (-0.055-3.9%)	65 (-2.4-160)	690 (-4.1-1700)
Democratic Republic of the Congo	1990	Male	85 to 89	38% (24-55%)	62% (45-76%)	0.48 (0.23-0.84)	0.89% (-0.11-2.3%)	1% (-0.2-2.7%)	19 (-3.8-53)	150 (-19-390)
Democratic Republic of the Congo	1990	Male	90 to 94	37% (23-55%)	63% (45-77%)	0.41 (0.19-0.77)	0.52% (-0.27-1.6%)	0.53% (-0.43-1.9%)	3.5 (-2.8-12)	20 (-11-63)
Democratic Republic of the Congo	1990	Male	95 plus	41% (22-61%)	59% (39-78%)	0.39 (0.15-0.79)	0.6% (-0.33-1.9%)	0.58% (-0.56-2.3%)	0.95 (-0.92-3.7)	4 (-2.2-13)
Democratic Republic of the Congo	1995	Female	All Ages	23% (19-29%)	77% (71-81%)	0.49 (0.39-0.61)	1.1% (0.6-1.9%)	2% (0.86-3.4%)	5300 (2100-9100)	200000 (110000-310000)
Democratic Republic of the Congo	1995	Female	15 to 19	23% (14-35%)	77% (65-86%)	0.41 (0.2-0.71)	1.1% (0.45-2.1%)	1.3% (0.29-3.1%)	57 (12-140)	5700 (2400-11000)
Democratic Republic of the Congo	1995	Female	20 to 24	32% (19-47%)	68% (53-81%)	0.78 (0.43-1.3)	2.7% (1.2-5.1%)	3.2% (0.95-6.9%)	190 (55-400)	16000 (6800-30000)
Democratic Republic of the Congo	1995	Female	25 to 29	36% (22-53%)	64% (47-78%)	1 (0.6-1.6)	3.5% (1.6-5.9%)	4.3% (1.6-7.8%)	280 (110-500)	20000 (9500-34000)
Democratic Republic of the Congo	1995	Female	30 to 34	39% (24-57%)	61% (43-76%)	1.2 (0.67-1.8)	4% (1.8-6.7%)	5.1% (2-9%)	350 (130-610)	22000 (10000-36000)
Democratic Republic of the Congo	1995	Female	35 to 39	40% (26-58%)	60% (42-74%)	1.2 (0.66-1.9)	4.1% (1.9-7%)	5.4% (2.2-9.6%)	370 (150-650)	21000 (9200-35000)
Democratic Republic of the Congo	1995	Female	40 to 44	40% (26-59%)	60% (41-74%)	1.1 (0.6-1.9)	4.1% (1.8-7.3%)	5.5% (2.2-10%)	380 (150-690)	18000 (7900-33000)
Democratic Republic of the Congo	1995	Female	45 to 49	39% (24-57%)	61% (43-76%)	1.1 (0.55-1.8)	4.7% (2-8.2%)	6.3% (2.7-11%)	470 (200-860)	20000 (8700-36000)
Democratic Republic of the Congo	1995	Female	50 to 54	37% (22-54%)	63% (46-78%)	0.97 (0.5-1.7)	4.9% (1.8-9.1%)	6.5% (2.3-12%)	570 (200-1100)	21000 (7700-38000)
Democratic Republic of the Congo	1995	Female	55 to 59	34% (21-51%)	66% (49-79%)	0.88 (0.45-1.5)	4.9% (1.7-9%)	6.3% (2.2-12%)	640 (220-1200)	20000 (7000-38000)
Democratic Republic of the Congo	1995	Female	60 to 64	33% (20-50%)	67% (50-80%)	0.84 (0.4-1.5)	4% (1.1-8.2%)	4.9% (1.4-10%)	590 (160-1200)	16000 (4400-32000)
Democratic Republic of the Congo	1995	Female	65 to 69	35% (23-50%)	65% (50-77%)	0.82 (0.43-1.4)	3.4% (0.69-7.3%)	4.1% (0.85-8.9%)	570 (120-1300)	12000 (2600-28000)
Democratic Republic of the Congo	1995	Female	70 to 74	35% (23-51%)	65% (49-77%)	0.74 (0.35-1.3)	2.5% (-0.0054-6%)	3% (0.027-7.3%)	430 (3.9-1100)	7500 (-17-19000)
Democratic Republic of the Congo	1995	Female	75 to 79	32% (20-49%)	68% (51-80%)	0.61 (0.28-1.1)	1.5% (-0.51-4.3%)	1.8% (-0.57-5.1%)	220 (-71-650)	3000 (-1000-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	1995	Female	80 to 84	29% (18-44%)	71% (56-82%)	0.5 (0.23-0.87)	0.94% (-0.55-3.1%)	1.1% (-0.62-3.7%)	94 (-52-300)	940 (-520-3000)
Democratic Republic of the Congo	1995	Female	85 to 89	28% (17-41%)	72% (59-83%)	0.44 (0.21-0.76)	0.64% (-0.51-2.4%)	0.77% (-0.61-2.8%)	36 (-28-130)	260 (-210-960)
Democratic Republic of the Congo	1995	Female	90 to 94	29% (18-43%)	71% (57-82%)	0.43 (0.2-0.79)	0.46% (-0.55-2.1%)	0.56% (-0.67-2.6%)	9.8 (-12-45)	48 (-57-220)
Democratic Republic of the Congo	1995	Female	95 plus	33% (19-51%)	67% (49-81%)	0.44 (0.17-0.87)	0.44% (-0.49-2%)	0.5% (-0.64-2.4%)	2.5 (-3.2-12)	8.6 (-9.7-40)
Democratic Republic of the Congo	1995	Male	All Ages	31% (25-38%)	69% (62-75%)	0.62 (0.49-0.76)	1.7% (1-2.7%)	2.7% (1.5-4.3%)	7700 (4300-11000)	340000 (220000-480000)
Democratic Republic of the Congo	1995	Male	15 to 19	25% (15-40%)	75% (60-85%)	0.45 (0.22-0.8)	1.8% (0.87-3.2%)	2.1% (0.7-4.3%)	100 (33-200)	9800 (4700-17000)
Democratic Republic of the Congo	1995	Male	20 to 24	38% (24-54%)	62% (46-76%)	0.93 (0.51-1.5)	4.7% (2.3-8.2%)	5.4% (1.9-10%)	360 (130-690)	29000 (14000-50000)
Democratic Republic of the Congo	1995	Male	25 to 29	45% (28-63%)	55% (37-72%)	1.3 (0.76-2)	6.3% (3.2-10%)	7.4% (3.2-13%)	570 (250-960)	41000 (21000-64000)
Democratic Republic of the Congo	1995	Male	30 to 34	48% (30-66%)	52% (34-70%)	1.5 (0.86-2.2)	6.5% (3.3-11%)	7.6% (3.4-13%)	680 (320-1200)	43000 (22000-69000)
Democratic Republic of the Congo	1995	Male	35 to 39	48% (32-66%)	52% (34-68%)	1.5 (0.84-2.3)	6.5% (3.5-10%)	7.6% (3.8-12%)	720 (360-1200)	40000 (22000-63000)
Democratic Republic of the Congo	1995	Male	40 to 44	50% (33-68%)	50% (32-67%)	1.5 (0.82-2.4)	6.9% (3.3-11%)	8.2% (3.7-13%)	800 (360-1300)	39000 (19000-63000)
Democratic Republic of the Congo	1995	Male	45 to 49	53% (34-71%)	47% (29-66%)	1.6 (0.88-2.5)	7.4% (3.8-12%)	9% (4.3-14%)	900 (440-1500)	39000 (20000-62000)
Democratic Republic of the Congo	1995	Male	50 to 54	53% (35-71%)	47% (29-65%)	1.5 (0.82-2.5)	7.5% (3.5-12%)	9.2% (4.1-15%)	920 (410-1500)	34000 (16000-57000)
Democratic Republic of the Congo	1995	Male	55 to 59	51% (33-70%)	49% (30-67%)	1.3 (0.7-2.2)	6.6% (2.7-11%)	8% (3.2-14%)	810 (320-1400)	26000 (11000-45000)
Democratic Republic of the Congo	1995	Male	60 to 64	49% (31-68%)	51% (32-69%)	1.2 (0.58-2)	5% (1.9-9.5%)	6.1% (2.2-12%)	650 (240-1200)	18000 (6900-33000)
Democratic Republic of the Congo	1995	Male	65 to 69	50% (32-67%)	50% (33-68%)	1.1 (0.56-1.7)	3.9% (1.3-7.4%)	4.6% (1.5-8.8%)	500 (160-970)	11000 (3700-22000)
Democratic Republic of the Congo	1995	Male	70 to 74	51% (33-68%)	49% (32-67%)	0.92 (0.47-1.6)	2.8% (0.45-6%)	3.2% (0.46-7.1%)	340 (49-750)	6200 (1000-13000)
Democratic Republic of the Congo	1995	Male	75 to 79	49% (33-66%)	51% (34-67%)	0.76 (0.36-1.3)	1.9% (-0.0074-4.6%)	2.2% (-0.035-5.4%)	190 (-3-470)	2600 (-10-6500)
Democratic Republic of the Congo	1995	Male	80 to 84	44% (27-62%)	56% (38-73%)	0.59 (0.28-1)	1.2% (-0.17-3%)	1.3% (-0.24-3.6%)	70 (-13-190)	740 (-110-1900)
Democratic Republic of the Congo	1995	Male	85 to 89	38% (23-55%)	62% (45-77%)	0.44 (0.23-0.77)	0.77% (-0.22-2.1%)	0.86% (-0.33-2.5%)	20 (-7.8-58)	160 (-43-430)
Democratic Republic of the Congo	1995	Male	90 to 94	38% (24-55%)	62% (45-76%)	0.38 (0.18-0.71)	0.45% (-0.36-1.5%)	0.45% (-0.55-1.7%)	3.6 (-4.3-13)	21 (-17-67)
Democratic Republic of the Congo	1995	Male	95 plus	41% (22-62%)	59% (38-78%)	0.36 (0.13-0.72)	0.52% (-0.4-1.7%)	0.5% (-0.63-2%)	1 (-1.3-3.9)	4.2 (-3.2-14)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	2000	Female	All Ages	22% (18-28%)	78% (72-82%)	0.48 (0.37-0.6)	1.2% (0.63-2%)	2.2% (0.91-3.8%)	6500 (2700-11000)	250000 (130000-380000)
Democratic Republic of the Congo	2000	Female	15 to 19	22% (14-35%)	78% (65-86%)	0.4 (0.19-0.71)	1.1% (0.43-2.2%)	1.4% (0.3-3.1%)	70 (15-160)	6900 (2600-13000)
Democratic Republic of the Congo	2000	Female	20 to 24	31% (19-45%)	69% (55-81%)	0.77 (0.41-1.3)	2.8% (1.2-5.2%)	3.5% (1-7%)	220 (65-460)	19000 (8100-34000)
Democratic Republic of the Congo	2000	Female	25 to 29	35% (21-51%)	65% (49-79%)	1 (0.58-1.6)	3.5% (1.6-6%)	4.3% (1.6-7.9%)	330 (130-600)	24000 (11000-41000)
Democratic Republic of the Congo	2000	Female	30 to 34	38% (23-55%)	62% (45-77%)	1.2 (0.66-1.8)	4% (1.9-6.7%)	5% (2.1-8.8%)	420 (180-740)	27000 (13000-44000)
Democratic Republic of the Congo	2000	Female	35 to 39	39% (25-57%)	61% (43-75%)	1.1 (0.63-1.8)	4.1% (1.8-7%)	5.2% (2.1-9.2%)	450 (190-790)	25000 (11000-42000)
Democratic Republic of the Congo	2000	Female	40 to 44	39% (25-57%)	61% (43-75%)	1.1 (0.59-1.8)	4.2% (1.7-7.2%)	5.4% (2.1-9.6%)	460 (180-830)	22000 (9300-39000)
Democratic Republic of the Congo	2000	Female	45 to 49	38% (23-55%)	62% (45-77%)	1.1 (0.54-1.8)	4.9% (2-8.6%)	6.5% (2.5-12%)	560 (220-1000)	24000 (9700-42000)
Democratic Republic of the Congo	2000	Female	50 to 54	36% (21-54%)	64% (46-79%)	0.96 (0.49-1.7)	5.2% (2-9.5%)	6.8% (2.4-13%)	700 (240-1300)	25000 (9100-48000)
Democratic Republic of the Congo	2000	Female	55 to 59	33% (20-50%)	67% (50-80%)	0.88 (0.44-1.5)	5.3% (1.8-9.9%)	6.8% (2.3-13%)	800 (260-1500)	25000 (8600-47000)
Democratic Republic of the Congo	2000	Female	60 to 64	32% (20-49%)	68% (51-80%)	0.83 (0.4-1.5)	4.3% (1.3-8.6%)	5.4% (1.5-11%)	730 (210-1500)	19000 (5600-39000)
Democratic Republic of the Congo	2000	Female	65 to 69	34% (22-49%)	66% (51-78%)	0.82 (0.43-1.4)	3.7% (0.76-7.8%)	4.6% (0.92-9.6%)	710 (140-1600)	16000 (3200-34000)
Democratic Republic of the Congo	2000	Female	70 to 74	34% (22-50%)	66% (50-78%)	0.73 (0.35-1.3)	2.8% (0.099-6.7%)	3.3% (0.16-7.9%)	540 (25-1300)	9500 (360-24000)
Democratic Republic of the Congo	2000	Female	75 to 79	31% (19-46%)	69% (54-81%)	0.59 (0.27-1.1)	1.7% (-0.39-4.6%)	2.1% (-0.45-5.5%)	290 (-63-790)	4000 (-920-11000)
Democratic Republic of the Congo	2000	Female	80 to 84	28% (17-44%)	72% (56-83%)	0.48 (0.23-0.86)	1.1% (-0.43-3.4%)	1.3% (-0.5-4%)	130 (-46-400)	1300 (-500-4000)
Democratic Republic of the Congo	2000	Female	85 to 89	27% (17-41%)	73% (59-83%)	0.43 (0.21-0.77)	0.8% (-0.47-2.8%)	0.96% (-0.55-3.4%)	53 (-30-180)	380 (-220-1300)
Democratic Republic of the Congo	2000	Female	90 to 94	29% (18-42%)	71% (58-82%)	0.43 (0.19-0.79)	0.6% (-0.5-2.3%)	0.73% (-0.63-2.8%)	15 (-13-57)	74 (-63-280)
Democratic Republic of the Congo	2000	Female	95 plus	32% (18-50%)	68% (50-82%)	0.43 (0.17-0.86)	0.58% (-0.49-2.4%)	0.66% (-0.64-2.9%)	3.8 (-3.7-17)	13 (-11-56)
Democratic Republic of the Congo	2000	Male	All Ages	30% (24-38%)	70% (62-76%)	0.63 (0.49-0.78)	2% (1.2-3%)	3.2% (1.7-4.9%)	10000 (5500-15000)	440000 (270000-610000)
Democratic Republic of the Congo	2000	Male	15 to 19	25% (15-39%)	75% (61-85%)	0.46 (0.22-0.84)	1.9% (0.87-3.5%)	2.3% (0.7-4.7%)	140 (43-280)	13000 (5800-23000)
Democratic Republic of the Congo	2000	Male	20 to 24	38% (23-55%)	62% (45-77%)	0.96 (0.53-1.6)	5.2% (2.6-8.7%)	6% (2.4-11%)	480 (200-880)	38000 (19000-65000)
Democratic Republic of the Congo	2000	Male	25 to 29	45% (28-62%)	55% (38-72%)	1.4 (0.8-2.1)	7.1% (3.7-11%)	8.4% (3.7-14%)	760 (350-1200)	53000 (27000-81000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	2000	Male	30 to 34	47% (29-65%)	53% (35-71%)	1.5 (0.89-2.4)	7.4% (3.9-12%)	8.7% (4-14%)	900 (410-1400)	55000 (29000-85000)
Democratic Republic of the Congo	2000	Male	35 to 39	47% (31-64%)	53% (36-69%)	1.5 (0.86-2.4)	7% (3.7-11%)	8.3% (4.1-13%)	930 (460-1500)	51000 (27000-79000)
Democratic Republic of the Congo	2000	Male	40 to 44	49% (32-68%)	51% (32-68%)	1.5 (0.83-2.4)	7.5% (3.7-12%)	8.9% (4.2-14%)	1000 (490-1700)	50000 (25000-77000)
Democratic Republic of the Congo	2000	Male	45 to 49	52% (34-69%)	48% (31-66%)	1.6 (0.92-2.6)	8.1% (4.3-13%)	9.8% (5-16%)	1100 (570-1800)	48000 (25000-76000)
Democratic Republic of the Congo	2000	Male	50 to 54	52% (35-70%)	48% (30-65%)	1.6 (0.87-2.6)	8.3% (3.7-14%)	10% (4.4-17%)	1200 (520-2000)	44000 (20000-73000)
Democratic Republic of the Congo	2000	Male	55 to 59	50% (33-68%)	50% (32-67%)	1.4 (0.75-2.3)	7.4% (3.2-12%)	9% (3.8-15%)	1100 (440-1800)	35000 (15000-58000)
Democratic Republic of the Congo	2000	Male	60 to 64	49% (31-68%)	51% (32-69%)	1.2 (0.6-2.1)	5.6% (2.1-11%)	6.8% (2.5-13%)	860 (320-1600)	23000 (9000-43000)
Democratic Republic of the Congo	2000	Male	65 to 69	50% (32-66%)	50% (34-68%)	1.1 (0.58-1.8)	4.4% (1.3-8.3%)	5.2% (1.6-10%)	670 (190-1300)	15000 (4700-28000)
Democratic Republic of the Congo	2000	Male	70 to 74	50% (33-67%)	50% (33-67%)	0.95 (0.49-1.6)	3.1% (0.52-6.8%)	3.7% (0.55-8%)	460 (70-1000)	8200 (1300-18000)
Democratic Republic of the Congo	2000	Male	75 to 79	48% (32-67%)	52% (33-68%)	0.79 (0.37-1.4)	2.2% (-0.044-5.3%)	2.5% (-0.097-6.3%)	250 (-9.6-620)	3500 (-70-8500)
Democratic Republic of the Congo	2000	Male	80 to 84	43% (25-61%)	57% (39-75%)	0.61 (0.28-1)	1.4% (-0.12-3.5%)	1.6% (-0.22-4.1%)	98 (-13-250)	1000 (-92-2600)
Democratic Republic of the Congo	2000	Male	85 to 89	38% (23-54%)	62% (46-77%)	0.46 (0.22-0.83)	0.91% (-0.19-2.4%)	1% (-0.29-2.8%)	28 (-8-79)	220 (-44-580)
Democratic Republic of the Congo	2000	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.4 (0.18-0.72)	0.55% (-0.31-1.7%)	0.58% (-0.47-2%)	5.3 (-4.2-18)	30 (-16-92)
Democratic Republic of the Congo	2000	Male	95 plus	40% (22-61%)	60% (39-78%)	0.37 (0.13-0.78)	0.64% (-0.45-2.1%)	0.65% (-0.7-2.4%)	1.5 (-1.6-5.4)	5.8 (-4-19)
Democratic Republic of the Congo	2005	Female	All Ages	22% (17-27%)	78% (73-83%)	0.49 (0.37-0.64)	1.4% (0.68-2.1%)	2.4% (1-4.1%)	7600 (3200-13000)	280000 (150000-440000)
Democratic Republic of the Congo	2005	Female	15 to 19	21% (13-33%)	79% (67-87%)	0.41 (0.2-0.72)	1.1% (0.49-2.2%)	1.4% (0.35-3.1%)	79 (21-180)	7900 (3300-15000)
Democratic Republic of the Congo	2005	Female	20 to 24	30% (18-45%)	70% (55-82%)	0.8 (0.42-1.3)	2.9% (1.3-5.2%)	3.6% (1.2-7.1%)	240 (80-480)	21000 (9600-37000)
Democratic Republic of the Congo	2005	Female	25 to 29	34% (20-51%)	66% (49-80%)	1.1 (0.6-1.7)	3.4% (1.7-5.9%)	4.2% (1.6-8.1%)	350 (140-630)	25000 (12000-43000)
Democratic Republic of the Congo	2005	Female	30 to 34	37% (22-54%)	63% (46-78%)	1.2 (0.68-1.9)	3.8% (1.8-6.5%)	4.7% (1.9-8.6%)	440 (190-780)	28000 (14000-47000)
Democratic Republic of the Congo	2005	Female	35 to 39	38% (24-56%)	62% (44-76%)	1.2 (0.64-1.9)	3.9% (1.9-6.6%)	4.9% (2.1-8.6%)	500 (220-850)	28000 (13000-45000)
Democratic Republic of the Congo	2005	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.2 (0.61-2)	4% (1.8-7%)	5.2% (2.1-9.3%)	530 (220-950)	25000 (11000-45000)
Democratic Republic of the Congo	2005	Female	45 to 49	37% (23-55%)	63% (45-77%)	1.1 (0.55-1.9)	4.9% (2.2-8.8%)	6.5% (2.8-12%)	660 (290-1200)	28000 (13000-50000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	2005	Female	50 to 54	35% (21-53%)	65% (47-79%)	1 (0.5-1.8)	5.4% (2.1-9.7%)	7% (2.6-13%)	790 (290-1500)	29000 (11000-53000)
Democratic Republic of the Congo	2005	Female	55 to 59	32% (20-50%)	68% (50-80%)	0.92 (0.46-1.6)	5.7% (2-11%)	7.3% (2.4-14%)	970 (320-1800)	30000 (11000-58000)
Democratic Republic of the Congo	2005	Female	60 to 64	32% (19-49%)	68% (51-81%)	0.87 (0.42-1.6)	4.6% (1.4-9%)	5.7% (1.7-11%)	890 (260-1700)	24000 (6900-47000)
Democratic Republic of the Congo	2005	Female	65 to 69	33% (22-49%)	67% (51-78%)	0.85 (0.44-1.5)	4% (0.96-8.4%)	4.9% (1.2-10%)	870 (210-1800)	19000 (4600-40000)
Democratic Republic of the Congo	2005	Female	70 to 74	33% (22-48%)	67% (52-78%)	0.76 (0.36-1.4)	3% (0.26-7.2%)	3.6% (0.31-8.4%)	670 (57-1600)	12000 (1000-28000)
Democratic Republic of the Congo	2005	Female	75 to 79	30% (19-47%)	70% (53-81%)	0.62 (0.28-1.1)	2% (-0.32-5.3%)	2.3% (-0.32-6.3%)	390 (-54-1000)	5200 (-830-14000)
Democratic Republic of the Congo	2005	Female	80 to 84	28% (16-43%)	72% (57-84%)	0.51 (0.24-0.91)	1.3% (-0.34-3.6%)	1.5% (-0.36-4.4%)	170 (-41-480)	1700 (-450-4900)
Democratic Republic of the Congo	2005	Female	85 to 89	26% (16-39%)	74% (61-84%)	0.45 (0.21-0.82)	0.92% (-0.34-2.9%)	1.1% (-0.41-3.5%)	73 (-27-230)	520 (-200-1700)
Democratic Republic of the Congo	2005	Female	90 to 94	28% (17-41%)	72% (59-83%)	0.43 (0.19-0.83)	0.7% (-0.38-2.5%)	0.85% (-0.48-3.1%)	21 (-12-77)	110 (-56-380)
Democratic Republic of the Congo	2005	Female	95 plus	31% (17-48%)	69% (52-83%)	0.44 (0.17-0.91)	0.68% (-0.4-2.6%)	0.78% (-0.51-3.2%)	5.4 (-3.5-22)	19 (-11-73)
Democratic Republic of the Congo	2005	Male	All Ages	29% (24-37%)	71% (63-76%)	0.66 (0.5-0.84)	2.2% (1.3-3.2%)	3.5% (1.9-5.3%)	11000 (6400-16000)	500000 (310000-690000)
Democratic Republic of the Congo	2005	Male	15 to 19	25% (15-39%)	75% (61-85%)	0.49 (0.23-0.89)	2% (0.98-3.6%)	2.3% (0.86-4.7%)	160 (58-330)	15000 (7400-28000)
Democratic Republic of the Congo	2005	Male	20 to 24	37% (23-55%)	63% (45-77%)	1 (0.55-1.7)	5.4% (2.8-8.8%)	6.3% (2.7-11%)	570 (260-1000)	45000 (23000-74000)
Democratic Republic of the Congo	2005	Male	25 to 29	45% (27-62%)	55% (38-73%)	1.4 (0.83-2.2)	7.5% (4-12%)	9.1% (4.3-15%)	860 (410-1400)	59000 (32000-90000)
Democratic Republic of the Congo	2005	Male	30 to 34	46% (29-65%)	54% (35-71%)	1.6 (0.93-2.5)	7.8% (4.3-12%)	9.4% (4.7-15%)	1000 (520-1600)	62000 (35000-95000)
Democratic Republic of the Congo	2005	Male	35 to 39	46% (30-64%)	54% (36-70%)	1.6 (0.89-2.5)	7.5% (4.1-12%)	8.9% (4.5-14%)	1000 (540-1700)	57000 (32000-88000)
Democratic Republic of the Congo	2005	Male	40 to 44	48% (32-67%)	52% (33-68%)	1.6 (0.83-2.6)	7.7% (4-12%)	9.3% (4.5-15%)	1200 (560-1900)	56000 (29000-88000)
Democratic Republic of the Congo	2005	Male	45 to 49	52% (33-69%)	48% (31-67%)	1.7 (0.93-2.8)	8.4% (4.5-13%)	10% (5.3-16%)	1300 (660-2000)	54000 (29000-85000)
Democratic Republic of the Congo	2005	Male	50 to 54	52% (34-70%)	48% (30-66%)	1.7 (0.87-2.8)	8.6% (4.2-14%)	11% (4.9-17%)	1300 (610-2200)	48000 (23000-80000)
Democratic Republic of the Congo	2005	Male	55 to 59	50% (33-69%)	50% (31-67%)	1.5 (0.78-2.5)	7.6% (3.4-13%)	9.4% (4-16%)	1200 (520-2000)	39000 (17000-65000)
Democratic Republic of the Congo	2005	Male	60 to 64	48% (30-68%)	52% (32-70%)	1.3 (0.61-2.3)	5.9% (2.3-11%)	7.1% (2.6-13%)	1000 (370-1800)	27000 (10000-49000)
Democratic Republic of the Congo	2005	Male	65 to 69	49% (32-66%)	51% (34-68%)	1.1 (0.61-1.9)	4.6% (1.6-8.7%)	5.6% (1.8-10%)	790 (250-1500)	18000 (5900-33000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	2005	Male	70 to 74	50% (33-67%)	50% (33-67%)	1 (0.51-1.8)	3.4% (0.65-7.2%)	4% (0.72-8.5%)	550 (98-1200)	9900 (1900-21000)
Democratic Republic of the Congo	2005	Male	75 to 79	48% (31-66%)	52% (34-69%)	0.84 (0.41-1.5)	2.4% (0.07-5.7%)	2.8% (0.051-6.7%)	310 (5.7-760)	4400 (130-10000)
Democratic Republic of the Congo	2005	Male	80 to 84	43% (26-62%)	57% (38-74%)	0.65 (0.31-1.2)	1.5% (-0.066-3.8%)	1.8% (-0.13-4.5%)	130 (-9.5-320)	1300 (-58-3300)
Democratic Republic of the Congo	2005	Male	85 to 89	38% (23-54%)	62% (46-77%)	0.49 (0.24-0.88)	1% (-0.084-2.4%)	1.1% (-0.16-2.9%)	38 (-5.3-96)	290 (-24-720)
Democratic Republic of the Congo	2005	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.42 (0.19-0.77)	0.6% (-0.26-1.8%)	0.64% (-0.4-2.1%)	6.9 (-4.4-22)	39 (-17-110)
Democratic Republic of the Congo	2005	Male	95 plus	40% (22-61%)	60% (39-78%)	0.39 (0.14-0.8)	0.69% (-0.38-2.2%)	0.71% (-0.62-2.6%)	1.8 (-1.6-6.6)	7.2 (-4-23)
Democratic Republic of the Congo	2010	Female	All Ages	22% (17-27%)	78% (73-83%)	0.54 (0.38-0.72)	1.5% (0.77-2.5%)	2.7% (1.1-4.8%)	9200 (3900-16000)	340000 (170000-530000)
Democratic Republic of the Congo	2010	Female	15 to 19	21% (13-34%)	79% (66-87%)	0.45 (0.21-0.83)	1.2% (0.51-2.2%)	1.4% (0.39-3.1%)	90 (25-200)	9200 (4000-18000)
Democratic Republic of the Congo	2010	Female	20 to 24	30% (18-45%)	70% (55-82%)	0.87 (0.45-1.5)	3% (1.4-5.3%)	3.7% (1.2-7.5%)	260 (89-530)	23000 (11000-42000)
Democratic Republic of the Congo	2010	Female	25 to 29	35% (21-51%)	65% (49-79%)	1.2 (0.63-2)	3.6% (1.7-6.3%)	4.6% (1.8-8.6%)	380 (150-700)	28000 (14000-48000)
Democratic Republic of the Congo	2010	Female	30 to 34	37% (23-54%)	63% (46-77%)	1.3 (0.73-2.1)	4.1% (2-6.9%)	5.2% (2.2-9.2%)	500 (220-880)	32000 (16000-54000)
Democratic Republic of the Congo	2010	Female	35 to 39	39% (24-57%)	61% (43-76%)	1.3 (0.69-2.2)	4.1% (1.9-7.1%)	5.2% (2.2-9.3%)	560 (240-970)	31000 (15000-52000)
Democratic Republic of the Congo	2010	Female	40 to 44	39% (24-58%)	61% (42-76%)	1.3 (0.64-2.2)	4.2% (1.9-7.3%)	5.4% (2.3-10%)	600 (260-1100)	29000 (14000-52000)
Democratic Republic of the Congo	2010	Female	45 to 49	38% (22-56%)	62% (44-78%)	1.2 (0.58-2.2)	5.3% (2.3-9.5%)	6.9% (2.9-13%)	810 (330-1500)	34000 (14000-62000)
Democratic Republic of the Congo	2010	Female	50 to 54	35% (21-53%)	65% (47-79%)	1.1 (0.51-1.9)	5.8% (2.3-11%)	7.5% (2.9-14%)	960 (370-1800)	35000 (14000-65000)
Democratic Republic of the Congo	2010	Female	55 to 59	33% (20-49%)	67% (51-80%)	0.99 (0.46-1.7)	6.2% (2.1-12%)	7.9% (2.7-15%)	1100 (380-2100)	36000 (12000-67000)
Democratic Republic of the Congo	2010	Female	60 to 64	32% (20-49%)	68% (51-80%)	0.95 (0.43-1.7)	5% (1.5-9.9%)	6.2% (1.9-12%)	1100 (320-2100)	29000 (8800-56000)
Democratic Republic of the Congo	2010	Female	65 to 69	34% (22-49%)	66% (51-78%)	0.93 (0.46-1.6)	4.5% (1.1-9.1%)	5.4% (1.4-11%)	1100 (270-2200)	24000 (5900-49000)
Democratic Republic of the Congo	2010	Female	70 to 74	33% (22-49%)	67% (51-78%)	0.83 (0.38-1.5)	3.4% (0.43-8%)	4.1% (0.54-9.6%)	860 (110-2000)	15000 (1900-36000)
Democratic Republic of the Congo	2010	Female	75 to 79	30% (19-46%)	70% (54-81%)	0.67 (0.3-1.3)	2.3% (-0.074-5.9%)	2.7% (-0.07-7%)	520 (-13-1400)	7100 (-210-19000)
Democratic Republic of the Congo	2010	Female	80 to 84	28% (16-43%)	72% (57-84%)	0.55 (0.25-1)	1.5% (-0.2-4.3%)	1.8% (-0.21-5.1%)	250 (-28-690)	2500 (-330-7000)
Democratic Republic of the Congo	2010	Female	85 to 89	26% (16-40%)	74% (60-84%)	0.49 (0.23-0.92)	1.1% (-0.21-3.3%)	1.3% (-0.25-4%)	100 (-20-310)	730 (-140-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	2010	Female	90 to 94	28% (17-41%)	72% (59-83%)	0.48 (0.22-0.91)	0.86% (-0.39-2.9%)	1% (-0.49-3.4%)	31 (-15-100)	150 (-72-510)
Democratic Republic of the Congo	2010	Female	95 plus	32% (18-50%)	68% (50-82%)	0.49 (0.19-1)	0.8% (-0.4-3%)	0.92% (-0.53-3.7%)	7.8 (-4.4-31)	26 (-13-100)
Democratic Republic of the Congo	2010	Male	All Ages	29% (23-38%)	71% (62-77%)	0.71 (0.51-0.95)	2.6% (1.5-3.9%)	4.1% (2.3-6.3%)	14000 (8000-21000)	610000 (380000-860000)
Democratic Republic of the Congo	2010	Male	15 to 19	24% (14-40%)	76% (60-86%)	0.52 (0.24-0.98)	2.2% (1.1-3.9%)	2.6% (0.96-5.2%)	200 (74-400)	19000 (8900-33000)
Democratic Republic of the Congo	2010	Male	20 to 24	37% (23-54%)	63% (46-77%)	1.1 (0.55-1.9)	6.2% (3.2-10%)	7.5% (3.2-13%)	710 (310-1200)	56000 (29000-93000)
Democratic Republic of the Congo	2010	Male	25 to 29	45% (28-62%)	55% (38-72%)	1.6 (0.82-2.5)	8.8% (4.7-14%)	11% (5-18%)	1100 (510-1700)	75000 (39000-120000)
Democratic Republic of the Congo	2010	Male	30 to 34	47% (29-65%)	53% (35-71%)	1.7 (0.97-2.8)	9.2% (5.1-14%)	11% (5.9-18%)	1200 (640-1900)	76000 (42000-120000)
Democratic Republic of the Congo	2010	Male	35 to 39	47% (30-65%)	53% (35-70%)	1.7 (0.9-2.8)	8.6% (4.7-13%)	11% (5.3-17%)	1300 (650-2000)	69000 (39000-110000)
Democratic Republic of the Congo	2010	Male	40 to 44	48% (32-67%)	52% (33-68%)	1.7 (0.89-2.9)	8.9% (4.6-14%)	11% (5.3-17%)	1400 (690-2200)	68000 (35000-110000)
Democratic Republic of the Congo	2010	Male	45 to 49	52% (34-69%)	48% (31-66%)	1.8 (0.96-3.1)	9.4% (5.1-15%)	12% (6.1-19%)	1500 (800-2600)	66000 (35000-110000)
Democratic Republic of the Congo	2010	Male	50 to 54	52% (34-70%)	48% (30-66%)	1.8 (0.91-3.1)	9.5% (4.7-16%)	12% (5.7-20%)	1600 (740-2700)	59000 (28000-98000)
Democratic Republic of the Congo	2010	Male	55 to 59	50% (32-69%)	50% (31-68%)	1.6 (0.78-2.7)	8.3% (3.6-14%)	10% (4.4-17%)	1400 (610-2400)	46000 (20000-77000)
Democratic Republic of the Congo	2010	Male	60 to 64	48% (30-68%)	52% (32-70%)	1.4 (0.63-2.5)	6.4% (2.4-12%)	7.8% (2.9-14%)	1200 (460-2200)	34000 (13000-60000)
Democratic Republic of the Congo	2010	Male	65 to 69	49% (32-66%)	51% (34-68%)	1.2 (0.62-2.1)	5.1% (1.7-9.6%)	6.1% (2-12%)	1000 (320-1900)	23000 (7600-43000)
Democratic Republic of the Congo	2010	Male	70 to 74	50% (33-66%)	50% (34-67%)	1.1 (0.52-2)	3.7% (0.83-7.9%)	4.4% (0.92-9.4%)	710 (150-1500)	13000 (2900-27000)
Democratic Republic of the Congo	2010	Male	75 to 79	48% (32-66%)	52% (34-68%)	0.89 (0.41-1.7)	2.6% (0.21-6.3%)	3.1% (0.19-7.4%)	410 (25-980)	5700 (400-14000)
Democratic Republic of the Congo	2010	Male	80 to 84	43% (26-62%)	57% (38-74%)	0.68 (0.31-1.2)	1.7% (-0.026-4.1%)	2% (-0.11-4.8%)	170 (-9.5-420)	1800 (-29-4300)
Democratic Republic of the Congo	2010	Male	85 to 89	37% (23-54%)	63% (46-77%)	0.52 (0.24-0.97)	1.1% (-0.086-2.8%)	1.3% (-0.16-3.3%)	52 (-6.5-130)	400 (-30-980)
Democratic Republic of the Congo	2010	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.44 (0.2-0.84)	0.69% (-0.21-2%)	0.75% (-0.34-2.4%)	10 (-4.5-32)	56 (-17-170)
Democratic Republic of the Congo	2010	Male	95 plus	40% (21-61%)	60% (39-79%)	0.42 (0.15-0.91)	0.79% (-0.28-2.6%)	0.83% (-0.48-3%)	2.6 (-1.5-9.5)	9.9 (-3.4-32)
Democratic Republic of the Congo	2016	Female	All Ages	21% (17-27%)	79% (73-83%)	0.63 (0.35-1.1)	1.9% (0.85-3.3%)	3.3% (1.2-6.3%)	11000 (4000-20000)	390000 (180000-660000)
Democratic Republic of the Congo	2016	Female	15 to 19	21% (13-34%)	79% (66-87%)	0.53 (0.21-1.1)	1.3% (0.57-2.6%)	1.7% (0.48-3.9%)	110 (31-250)	11000 (4900-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	2016	Female	20 to 24	30% (18-45%)	70% (55-82%)	1 (0.42-2)	3.3% (1.6-6%)	4.3% (1.4-8.9%)	290 (94-610)	27000 (13000-48000)
Democratic Republic of the Congo	2016	Female	25 to 29	34% (20-51%)	66% (49-80%)	1.4 (0.6-2.6)	4.1% (1.9-7%)	5.6% (2.1-10%)	410 (150-770)	32000 (15000-55000)
Democratic Republic of the Congo	2016	Female	30 to 34	37% (22-53%)	63% (47-78%)	1.5 (0.67-2.8)	4.7% (2.3-8.3%)	6.5% (2.6-12%)	550 (220-990)	36000 (17000-61000)
Democratic Republic of the Congo	2016	Female	35 to 39	38% (24-55%)	62% (45-76%)	1.5 (0.69-2.8)	4.7% (2.2-8.3%)	6.5% (2.8-12%)	600 (250-1100)	34000 (16000-59000)
Democratic Republic of the Congo	2016	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.5 (0.64-2.8)	4.9% (2.1-8.8%)	6.7% (2.6-12%)	670 (260-1200)	33000 (14000-61000)
Democratic Republic of the Congo	2016	Female	45 to 49	37% (23-55%)	63% (45-77%)	1.4 (0.56-2.8)	6.1% (2.4-11%)	8.4% (3.2-16%)	920 (350-1700)	39000 (16000-72000)
Democratic Republic of the Congo	2016	Female	50 to 54	35% (21-53%)	65% (47-79%)	1.3 (0.49-2.5)	6.6% (2.3-13%)	8.8% (3-17%)	1100 (400-2200)	42000 (15000-79000)
Democratic Republic of the Congo	2016	Female	55 to 59	32% (20-50%)	68% (50-80%)	1.2 (0.45-2.3)	7% (2.2-14%)	9.1% (2.9-18%)	1300 (430-2600)	42000 (14000-82000)
Democratic Republic of the Congo	2016	Female	60 to 64	32% (19-49%)	68% (51-81%)	1.1 (0.45-2.2)	5.6% (1.6-12%)	7.1% (2-15%)	1200 (350-2500)	32000 (9400-66000)
Democratic Republic of the Congo	2016	Female	65 to 69	33% (21-48%)	67% (52-79%)	1.1 (0.45-2.1)	5.1% (1.2-11%)	6.3% (1.5-13%)	1300 (320-2700)	29000 (7300-61000)
Democratic Republic of the Congo	2016	Female	70 to 74	33% (21-48%)	67% (52-79%)	0.97 (0.38-1.9)	4% (0.58-9.8%)	4.9% (0.73-12%)	1100 (170-2600)	19000 (2900-47000)
Democratic Republic of the Congo	2016	Female	75 to 79	30% (19-46%)	70% (54-81%)	0.8 (0.31-1.7)	2.8% (0.087-7.7%)	3.3% (0.13-9.3%)	710 (27-1900)	9700 (330-27000)
Democratic Republic of the Congo	2016	Female	80 to 84	28% (17-43%)	72% (57-83%)	0.65 (0.26-1.3)	2% (-0.022-5.7%)	2.4% (-0.011-6.6%)	380 (-1.7-1100)	3800 (-42-11000)
Democratic Republic of the Congo	2016	Female	85 to 89	26% (16-39%)	74% (61-84%)	0.57 (0.22-1.2)	1.5% (-0.13-4.6%)	1.7% (-0.14-5.5%)	160 (-13-490)	1100 (-99-3500)
Democratic Republic of the Congo	2016	Female	90 to 94	28% (17-42%)	72% (58-83%)	0.56 (0.23-1.2)	1.2% (-0.3-3.8%)	1.4% (-0.35-4.6%)	48 (-12-160)	240 (-59-800)
Democratic Republic of the Congo	2016	Female	95 plus	32% (17-50%)	68% (50-83%)	0.57 (0.19-1.3)	1.1% (-0.3-3.8%)	1.3% (-0.41-4.5%)	12 (-4.1-44)	42 (-12-150)
Democratic Republic of the Congo	2016	Male	All Ages	29% (23-36%)	71% (64-77%)	0.82 (0.45-1.4)	3.2% (1.6-5.1%)	5% (2.3-8.4%)	17000 (8300-29000)	740000 (400000-1200000)
Democratic Republic of the Congo	2016	Male	15 to 19	24% (14-39%)	76% (61-86%)	0.6 (0.23-1.3)	2.4% (1.1-4.5%)	3% (0.99-6.2%)	250 (86-520)	24000 (11000-42000)
Democratic Republic of the Congo	2016	Male	20 to 24	37% (22-53%)	63% (47-78%)	1.3 (0.52-2.4)	6.9% (3.3-12%)	8.4% (3.3-15%)	870 (330-1500)	69000 (33000-120000)
Democratic Republic of the Congo	2016	Male	25 to 29	44% (27-61%)	56% (39-73%)	1.8 (0.83-3.3)	9.9% (5.1-16%)	13% (5.7-21%)	1300 (580-2200)	91000 (46000-150000)
Democratic Republic of the Congo	2016	Male	30 to 34	46% (28-64%)	54% (36-72%)	2 (0.93-3.7)	11% (5.4-17%)	14% (6.3-22%)	1500 (680-2500)	93000 (47000-150000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Democratic Republic of the Congo	2016	Male	35 to 39	46% (29-64%)	54% (36-71%)	2 (0.9-3.6)	10% (5.1-16%)	13% (6-21%)	1500 (700-2500)	83000 (42000-130000)
Democratic Republic of the Congo	2016	Male	40 to 44	48% (32-67%)	52% (33-68%)	2 (0.87-3.9)	10% (4.9-16%)	13% (5.8-21%)	1600 (740-2700)	80000 (37000-130000)
Democratic Republic of the Congo	2016	Male	45 to 49	51% (32-70%)	49% (30-68%)	2.1 (0.89-4)	11% (5-18%)	14% (6.2-23%)	1800 (810-3000)	78000 (36000-130000)
Democratic Republic of the Congo	2016	Male	50 to 54	52% (34-70%)	48% (30-66%)	2.1 (0.85-4)	11% (4.5-18%)	13% (5.4-23%)	1900 (750-3300)	71000 (29000-120000)
Democratic Republic of the Congo	2016	Male	55 to 59	50% (32-69%)	50% (31-68%)	1.8 (0.76-3.4)	9.2% (3.6-16%)	12% (4.5-20%)	1700 (670-3100)	56000 (22000-98000)
Democratic Republic of the Congo	2016	Male	60 to 64	48% (30-68%)	52% (32-70%)	1.6 (0.64-3.2)	7.2% (2.6-14%)	8.8% (3.1-17%)	1500 (540-2800)	40000 (15000-76000)
Democratic Republic of the Congo	2016	Male	65 to 69	49% (32-66%)	51% (34-68%)	1.4 (0.63-2.8)	5.8% (1.9-12%)	7% (2.2-14%)	1300 (400-2600)	29000 (9300-59000)
Democratic Republic of the Congo	2016	Male	70 to 74	49% (32-66%)	51% (34-68%)	1.2 (0.52-2.5)	4.3% (0.96-9.7%)	5.1% (1.1-12%)	960 (210-2200)	17000 (4000-40000)
Democratic Republic of the Congo	2016	Male	75 to 79	47% (31-65%)	53% (35-69%)	1 (0.42-2.1)	3.1% (0.49-7.8%)	3.7% (0.5-9.4%)	580 (80-1500)	8200 (1300-21000)
Democratic Republic of the Congo	2016	Male	80 to 84	42% (25-60%)	58% (40-75%)	0.79 (0.32-1.6)	2.1% (0.15-5.4%)	2.4% (0.11-6.3%)	260 (12-680)	2700 (200-7100)
Democratic Republic of the Congo	2016	Male	85 to 89	37% (22-54%)	63% (46-78%)	0.6 (0.24-1.2)	1.4% (0.082-4%)	1.6% (0.027-4.8%)	84 (1.4-250)	640 (36-1800)
Democratic Republic of the Congo	2016	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.51 (0.21-1.1)	0.91% (-0.11-2.7%)	0.99% (-0.23-3.1%)	17 (-4.1-55)	93 (-11-280)
Democratic Republic of the Congo	2016	Male	95 plus	39% (21-60%)	61% (40-79%)	0.48 (0.17-1.1)	1% (-0.16-3.3%)	1.1% (-0.33-3.9%)	4.3 (-1.3-15)	16 (-2.4-52)
Equatorial Guinea	1990	Female	All Ages	25% (20-31%)	75% (69-80%)	0.75 (0.56-1)	2.3% (1.3-3.5%)	3.9% (1.9-6.5%)	140 (66-230)	4700 (2700-7200)
Equatorial Guinea	1990	Female	15 to 19	23% (14-36%)	77% (64-86%)	0.56 (0.26-1.1)	2% (0.69-3.9%)	2.6% (0.68-5.5%)	1.1 (0.28-2.3)	89 (31-180)
Equatorial Guinea	1990	Female	20 to 24	32% (20-47%)	68% (53-80%)	1.1 (0.56-1.8)	4.7% (2.1-7.9%)	5.9% (2.3-10%)	4.4 (1.7-7.9)	330 (150-580)
Equatorial Guinea	1990	Female	25 to 29	36% (21-52%)	64% (48-79%)	1.4 (0.85-2.2)	6.8% (3.7-11%)	8.7% (4.4-14%)	8.2 (4-14)	540 (290-900)
Equatorial Guinea	1990	Female	30 to 34	38% (23-55%)	62% (45-77%)	1.6 (0.92-2.4)	8.5% (4.4-13%)	11% (5.4-17%)	10 (5-17)	610 (320-990)
Equatorial Guinea	1990	Female	35 to 39	40% (25-58%)	60% (42-75%)	1.6 (0.83-2.7)	8.6% (4.5-14%)	11% (5.6-18%)	8.7 (4.1-15)	460 (230-770)
Equatorial Guinea	1990	Female	40 to 44	40% (25-58%)	60% (42-75%)	1.5 (0.72-2.8)	8.5% (3.7-14%)	11% (4.5-18%)	5.7 (2.5-10)	270 (120-470)
Equatorial Guinea	1990	Female	45 to 49	39% (24-56%)	61% (44-76%)	1.5 (0.69-2.8)	9.1% (4.2-16%)	11% (5.1-20%)	7.4 (3.3-13)	310 (140-530)
Equatorial Guinea	1990	Female	50 to 54	37% (22-54%)	63% (46-78%)	1.4 (0.63-2.5)	9.2% (4-16%)	11% (4.6-20%)	11 (4.3-19)	390 (160-680)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	1990	Female	55 to 59	34% (21-51%)	66% (49-79%)	1.2 (0.6-2.1)	9% (3.4-15%)	11% (4.1-19%)	15 (5.4-27)	460 (170-830)
Equatorial Guinea	1990	Female	60 to 64	34% (21-51%)	66% (49-79%)	1.2 (0.56-2.2)	7.3% (2.4-13%)	8.5% (2.9-16%)	16 (5.1-29)	420 (140-770)
Equatorial Guinea	1990	Female	65 to 69	35% (23-51%)	65% (49-77%)	1.1 (0.55-2)	6.5% (2.1-12%)	7.4% (2.5-14%)	17 (5.7-34)	380 (120-750)
Equatorial Guinea	1990	Female	70 to 74	35% (23-50%)	65% (50-77%)	1 (0.48-1.9)	5.1% (1-11%)	5.8% (1.2-13%)	15 (3-32)	270 (53-560)
Equatorial Guinea	1990	Female	75 to 79	32% (20-48%)	68% (52-80%)	0.84 (0.38-1.6)	3.5% (0.16-8.5%)	4% (0.18-9.7%)	9.1 (0.42-22)	130 (5.9-300)
Equatorial Guinea	1990	Female	80 to 84	29% (18-44%)	71% (56-82%)	0.69 (0.31-1.3)	2.5% (-0.073-6.2%)	2.9% (-0.074-7.2%)	4.6 (-0.12-12)	47 (-1.4-120)
Equatorial Guinea	1990	Female	85 to 89	28% (17-42%)	72% (58-83%)	0.61 (0.29-1.1)	1.9% (-0.17-4.9%)	2.2% (-0.19-5.8%)	1.5 (-0.14-4.1)	11 (-1-30)
Equatorial Guinea	1990	Female	90 to 94	29% (18-43%)	71% (57-82%)	0.6 (0.26-1.1)	1.5% (-0.25-4.4%)	1.7% (-0.32-5.3%)	0.37 (-0.068-1.1)	1.9 (-0.31-5.5)
Equatorial Guinea	1990	Female	95 plus	33% (19-50%)	67% (50-81%)	0.6 (0.24-1.2)	1.2% (-0.41-4%)	1.4% (-0.55-4.7%)	0.09 (-0.035-0.3)	0.31 (-0.1-0.99)
Equatorial Guinea	1990	Male	All Ages	34% (27-42%)	66% (58-73%)	1 (0.75-1.3)	4% (2.6-5.8%)	6.2% (3.6-9.2%)	230 (130-350)	9500 (6100-13000)
Equatorial Guinea	1990	Male	15 to 19	25% (15-40%)	75% (60-85%)	0.6 (0.27-1.1)	3.3% (1.3-6.3%)	4% (1.3-8%)	2.2 (0.71-4.6)	180 (69-340)
Equatorial Guinea	1990	Male	20 to 24	38% (23-54%)	62% (46-77%)	1.2 (0.67-2.1)	9.1% (4.5-14%)	11% (4.8-17%)	11 (5-19)	800 (390-1300)
Equatorial Guinea	1990	Male	25 to 29	45% (28-62%)	55% (38-72%)	1.8 (1.1-2.7)	14% (7.8-20%)	16% (8.7-24%)	21 (11-32)	1300 (750-2000)
Equatorial Guinea	1990	Male	30 to 34	47% (29-65%)	53% (35-71%)	2 (1.2-3.1)	15% (9.4-22%)	19% (11-27%)	25 (15-38)	1500 (880-2200)
Equatorial Guinea	1990	Male	35 to 39	47% (31-66%)	53% (34-69%)	2 (1.1-3.3)	15% (9.2-23%)	19% (11-28%)	22 (12-34)	1100 (670-1800)
Equatorial Guinea	1990	Male	40 to 44	49% (33-68%)	51% (32-67%)	2 (1-3.6)	16% (8.3-24%)	19% (9.5-29%)	17 (8.4-27)	800 (400-1300)
Equatorial Guinea	1990	Male	45 to 49	52% (34-69%)	48% (31-66%)	2.1 (1-3.9)	16% (8.3-25%)	19% (9.6-30%)	19 (9.1-30)	770 (390-1200)
Equatorial Guinea	1990	Male	50 to 54	53% (34-70%)	47% (30-66%)	2.1 (1-3.6)	15% (7.3-24%)	18% (8.3-28%)	22 (10-36)	800 (370-1300)
Equatorial Guinea	1990	Male	55 to 59	50% (33-69%)	50% (31-67%)	1.8 (0.92-3.2)	13% (6.3-21%)	15% (7.3-25%)	24 (11-40)	760 (360-1300)
Equatorial Guinea	1990	Male	60 to 64	49% (31-68%)	51% (32-69%)	1.6 (0.8-2.9)	10% (4.1-17%)	12% (4.6-20%)	22 (8.6-40)	590 (230-1100)
Equatorial Guinea	1990	Male	65 to 69	49% (32-67%)	51% (33-68%)	1.4 (0.7-2.5)	8.2% (2.9-15%)	9.3% (3.3-17%)	19 (6.4-36)	430 (150-800)
Equatorial Guinea	1990	Male	70 to 74	50% (33-67%)	50% (33-67%)	1.3 (0.62-2.2)	6.1% (1.4-12%)	6.9% (1.5-14%)	15 (3.3-30)	270 (61-530)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	1990	Male	75 to 79	48% (32-66%)	52% (34-68%)	1 (0.49-1.9)	4.4% (0.43-9.9%)	5% (0.44-11%)	8.9 (0.76-21)	120 (11-280)
Equatorial Guinea	1990	Male	80 to 84	43% (26-61%)	57% (39-74%)	0.8 (0.37-1.5)	3% (0.09-7.1%)	3.4% (0.016-8.2%)	3.7 (0.016-9.1)	38 (1.1-92)
Equatorial Guinea	1990	Male	85 to 89	37% (23-54%)	63% (46-77%)	0.6 (0.28-1.1)	2% (-0.035-4.8%)	2.2% (-0.12-5.6%)	1 (-0.056-2.6)	7.7 (-0.14-19)
Equatorial Guinea	1990	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.52 (0.24-0.99)	1.2% (-0.31-3.3%)	1.3% (-0.49-3.8%)	0.19 (-0.07-0.55)	1 (-0.27-2.8)
Equatorial Guinea	1990	Male	95 plus	40% (22-60%)	60% (40-78%)	0.48 (0.19-1)	1.2% (-0.45-3.8%)	1.3% (-0.73-4.3%)	0.059 (-0.032-0.19)	0.22 (-0.08-0.65)
Equatorial Guinea	1995	Female	All Ages	25% (20-31%)	75% (69-80%)	0.75 (0.56-1)	2.2% (1.3-3.5%)	3.8% (1.9-6.3%)	140 (70-240)	5200 (3000-8100)
Equatorial Guinea	1995	Female	15 to 19	23% (14-36%)	77% (64-86%)	0.56 (0.25-1)	1.9% (0.71-3.8%)	2.4% (0.71-5.3%)	0.9 (0.26-1.9)	76 (29-150)
Equatorial Guinea	1995	Female	20 to 24	32% (19-47%)	68% (53-81%)	1.1 (0.54-1.8)	4.3% (1.9-7.6%)	5.2% (2-9.6%)	3.6 (1.3-6.7)	270 (120-480)
Equatorial Guinea	1995	Female	25 to 29	36% (22-53%)	64% (47-78%)	1.4 (0.86-2.3)	6.2% (3.3-9.8%)	7.6% (3.9-12%)	8.3 (4-14)	550 (290-900)
Equatorial Guinea	1995	Female	30 to 34	39% (24-56%)	61% (44-76%)	1.6 (0.94-2.4)	7.8% (4.4-12%)	9.7% (5.2-15%)	13 (6.7-21)	760 (420-1200)
Equatorial Guinea	1995	Female	35 to 39	40% (26-58%)	60% (42-74%)	1.6 (0.89-2.6)	8% (4.3-13%)	10% (5.2-16%)	14 (6.8-23)	720 (380-1200)
Equatorial Guinea	1995	Female	40 to 44	40% (26-59%)	60% (41-74%)	1.5 (0.79-2.6)	8.1% (4-13%)	10% (4.9-16%)	11 (4.9-18)	490 (240-820)
Equatorial Guinea	1995	Female	45 to 49	39% (24-57%)	61% (43-76%)	1.5 (0.73-2.6)	8.6% (4-15%)	11% (4.9-18%)	7.8 (3.5-14)	320 (150-560)
Equatorial Guinea	1995	Female	50 to 54	37% (22-55%)	63% (45-78%)	1.4 (0.66-2.5)	8.8% (3.9-16%)	11% (4.7-19%)	9.9 (4.1-18)	360 (150-660)
Equatorial Guinea	1995	Female	55 to 59	34% (21-52%)	66% (48-79%)	1.2 (0.59-2.2)	8.6% (3.3-15%)	10% (4-18%)	13 (4.8-25)	410 (150-780)
Equatorial Guinea	1995	Female	60 to 64	33% (20-50%)	67% (50-80%)	1.1 (0.54-2.1)	7% (2.4-13%)	8.1% (2.8-15%)	15 (4.8-29)	390 (130-770)
Equatorial Guinea	1995	Female	65 to 69	35% (23-50%)	65% (50-77%)	1.1 (0.54-2)	6.2% (1.9-12%)	7.2% (2.3-14%)	17 (5.1-34)	370 (110-750)
Equatorial Guinea	1995	Female	70 to 74	35% (23-50%)	65% (50-77%)	1 (0.49-1.9)	4.9% (0.86-11%)	5.6% (1-12%)	15 (2.6-34)	270 (46-590)
Equatorial Guinea	1995	Female	75 to 79	32% (20-48%)	68% (52-80%)	0.84 (0.37-1.6)	3.4% (0.0089-8.1%)	3.9% (-0.0015-9.4%)	9.8 (-0.0039-23)	130 (0.39-320)
Equatorial Guinea	1995	Female	80 to 84	29% (18-45%)	71% (55-82%)	0.69 (0.32-1.3)	2.5% (-0.02-6.3%)	2.8% (-0.028-7.2%)	5 (-0.05-13)	51 (-0.42-130)
Equatorial Guinea	1995	Female	85 to 89	28% (17-42%)	72% (58-83%)	0.61 (0.28-1.1)	1.8% (-0.14-5%)	2.1% (-0.17-6%)	1.7 (-0.14-4.8)	12 (-1-34)
Equatorial Guinea	1995	Female	90 to 94	30% (19-43%)	70% (57-81%)	0.6 (0.28-1.1)	1.4% (-0.29-4.1%)	1.7% (-0.36-5.1%)	0.44 (-0.093-1.3)	2.2 (-0.44-6.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	1995	Female	95 plus	33% (19-51%)	67% (49-81%)	0.61 (0.25-1.2)	1.2% (-0.43-4%)	1.4% (-0.57-4.7%)	0.097 (-0.039-0.32)	0.33 (-0.11-1.1)
Equatorial Guinea	1995	Male	All Ages	34% (27-42%)	66% (58-73%)	1 (0.76-1.4)	3.8% (2.4-5.4%)	5.9% (3.5-8.6%)	240 (140-370)	10000 (6300-14000)
Equatorial Guinea	1995	Male	15 to 19	25% (15-40%)	75% (60-85%)	0.62 (0.28-1.2)	3.3% (1.4-6.1%)	4.1% (1.4-7.9%)	1.8 (0.63-3.6)	150 (59-270)
Equatorial Guinea	1995	Male	20 to 24	38% (24-55%)	62% (45-76%)	1.3 (0.67-2.1)	8.7% (4.3-14%)	10% (4.6-17%)	8.5 (3.8-14)	620 (300-1000)
Equatorial Guinea	1995	Male	25 to 29	46% (29-62%)	54% (38-71%)	1.8 (1.1-2.8)	12% (7.1-18%)	15% (7.9-22%)	19 (9.9-29)	1200 (700-1800)
Equatorial Guinea	1995	Male	30 to 34	48% (30-66%)	52% (34-70%)	2 (1.2-3)	14% (8.6-20%)	16% (9.9-23%)	28 (16-41)	1600 (970-2400)
Equatorial Guinea	1995	Male	35 to 39	48% (32-66%)	52% (34-68%)	2 (1.2-3.2)	14% (8.2-21%)	17% (9.4-25%)	30 (17-46)	1600 (920-2400)
Equatorial Guinea	1995	Male	40 to 44	50% (34-68%)	50% (32-66%)	2 (1.1-3.4)	15% (8-22%)	17% (9.2-26%)	27 (14-43)	1300 (690-2000)
Equatorial Guinea	1995	Male	45 to 49	53% (34-70%)	47% (30-66%)	2.2 (1.1-3.7)	15% (8-22%)	18% (9.2-27%)	20 (9.9-33)	850 (420-1300)
Equatorial Guinea	1995	Male	50 to 54	53% (35-71%)	47% (29-65%)	2.1 (1.1-3.8)	14% (7.1-23%)	17% (8.2-27%)	20 (9.1-34)	730 (340-1200)
Equatorial Guinea	1995	Male	55 to 59	51% (33-70%)	49% (30-67%)	1.8 (0.95-3.2)	12% (5.5-20%)	15% (6.6-24%)	21 (8.9-36)	660 (290-1100)
Equatorial Guinea	1995	Male	60 to 64	49% (31-69%)	51% (31-69%)	1.6 (0.77-2.9)	9.6% (3.8-17%)	11% (4.4-20%)	20 (7.2-37)	540 (200-1000)
Equatorial Guinea	1995	Male	65 to 69	50% (33-67%)	50% (33-67%)	1.4 (0.73-2.5)	7.8% (2.9-14%)	9% (3.1-16%)	18 (6.4-34)	400 (140-740)
Equatorial Guinea	1995	Male	70 to 74	51% (33-68%)	49% (32-67%)	1.3 (0.64-2.2)	5.9% (1.3-12%)	6.7% (1.4-13%)	14 (3.4-29)	260 (63-520)
Equatorial Guinea	1995	Male	75 to 79	49% (33-67%)	51% (33-67%)	1 (0.5-1.9)	4.3% (0.33-9.4%)	4.9% (0.34-11%)	9.2 (0.65-20)	130 (11-280)
Equatorial Guinea	1995	Male	80 to 84	44% (27-61%)	56% (39-73%)	0.81 (0.38-1.5)	3% (0.14-6.7%)	3.4% (0.13-7.8%)	4 (0.14-9.1)	41 (1.8-93)
Equatorial Guinea	1995	Male	85 to 89	38% (23-55%)	62% (45-77%)	0.62 (0.28-1.1)	2% (-0.04-5%)	2.2% (-0.12-5.8%)	1.2 (-0.066-3.1)	9.1 (-0.18-23)
Equatorial Guinea	1995	Male	90 to 94	38% (23-54%)	62% (46-77%)	0.53 (0.24-0.99)	1.2% (-0.21-3.4%)	1.4% (-0.36-3.9%)	0.23 (-0.061-0.67)	1.2 (-0.21-3.4)
Equatorial Guinea	1995	Male	95 plus	41% (22-62%)	59% (38-78%)	0.49 (0.19-0.99)	1.3% (-0.43-3.8%)	1.4% (-0.67-4.5%)	0.058 (-0.029-0.19)	0.21 (-0.071-0.64)
Equatorial Guinea	2000	Female	All Ages	24% (19-30%)	76% (70-81%)	0.98 (0.75-1.3)	2.2% (1.4-3.2%)	3.8% (2.2-5.8%)	120 (63-190)	4200 (2500-6300)
Equatorial Guinea	2000	Female	15 to 19	22% (13-34%)	78% (66-87%)	0.74 (0.34-1.3)	1.9% (0.95-3.3%)	2.5% (1-4.6%)	1.1 (0.41-2)	97 (46-170)
Equatorial Guinea	2000	Female	20 to 24	31% (19-47%)	69% (53-81%)	1.4 (0.71-2.4)	3.4% (1.8-5.6%)	4% (1.8-7%)	2.1 (0.94-3.8)	180 (93-290)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	2000	Female	25 to 29	35% (21-51%)	65% (49-79%)	1.9 (1.1-3.1)	4.2% (2.4-6.5%)	4.8% (2.4-8%)	4.4 (2.3-7.4)	320 (180-500)
Equatorial Guinea	2000	Female	30 to 34	38% (23-55%)	62% (45-77%)	2.1 (1.2-3.3)	5.3% (3.1-8.2%)	6.3% (3.4-10%)	8.4 (4.6-14)	520 (310-810)
Equatorial Guinea	2000	Female	35 to 39	39% (24-57%)	61% (43-76%)	2.1 (1.2-3.2)	5.6% (3.3-8.8%)	6.9% (3.9-11%)	11 (5.6-18)	590 (330-950)
Equatorial Guinea	2000	Female	40 to 44	39% (25-58%)	61% (42-75%)	2.1 (1.1-3.2)	6.3% (3.7-9.6%)	7.9% (4.4-12%)	11 (5.7-18)	540 (290-860)
Equatorial Guinea	2000	Female	45 to 49	38% (22-55%)	62% (45-78%)	2 (1-3.4)	7.2% (4-11%)	9.2% (5-14%)	9.7 (4.7-17)	420 (210-700)
Equatorial Guinea	2000	Female	50 to 54	36% (21-53%)	64% (47-79%)	1.8 (0.87-3.1)	7.5% (4-12%)	9.5% (5-15%)	7.3 (3.4-13)	270 (130-480)
Equatorial Guinea	2000	Female	55 to 59	33% (20-50%)	67% (50-80%)	1.6 (0.8-2.9)	7.9% (3.7-13%)	9.8% (4.6-16%)	8.9 (3.8-16)	280 (120-510)
Equatorial Guinea	2000	Female	60 to 64	32% (19-49%)	68% (51-81%)	1.5 (0.74-2.7)	6.7% (2.8-12%)	8.1% (3.3-14%)	10 (3.7-19)	270 (99-510)
Equatorial Guinea	2000	Female	65 to 69	34% (22-50%)	66% (50-78%)	1.5 (0.77-2.6)	6.4% (2.5-12%)	7.6% (3-13%)	13 (4.9-25)	300 (110-570)
Equatorial Guinea	2000	Female	70 to 74	34% (22-50%)	66% (50-78%)	1.4 (0.69-2.4)	5.4% (1.6-10%)	6.3% (1.8-12%)	13 (3.9-27)	240 (70-480)
Equatorial Guinea	2000	Female	75 to 79	31% (20-47%)	69% (53-80%)	1.1 (0.51-2.1)	3.9% (0.59-8.5%)	4.6% (0.7-9.9%)	9.5 (1.6-21)	130 (21-290)
Equatorial Guinea	2000	Female	80 to 84	28% (17-43%)	72% (57-83%)	0.92 (0.43-1.7)	2.9% (0.51-6.4%)	3.4% (0.6-7.5%)	5.7 (0.98-13)	59 (10-130)
Equatorial Guinea	2000	Female	85 to 89	27% (17-40%)	73% (60-83%)	0.81 (0.37-1.4)	2.2% (0.27-4.9%)	2.6% (0.34-5.7%)	2 (0.26-4.5)	15 (1.9-33)
Equatorial Guinea	2000	Female	90 to 94	28% (18-42%)	72% (58-82%)	0.79 (0.36-1.4)	1.8% (0.1-4.1%)	2.1% (0.097-4.9%)	0.56 (0.027-1.3)	2.8 (0.17-6.7)
Equatorial Guinea	2000	Female	95 plus	32% (18-50%)	68% (50-82%)	0.8 (0.31-1.6)	1.6% (-0.11-4.3%)	1.8% (-0.23-5%)	0.13 (-0.018-0.37)	0.46 (-0.032-1.3)
Equatorial Guinea	2000	Male	All Ages	33% (26-41%)	67% (59-74%)	1.4 (1.1-1.7)	3.9% (2.7-5.5%)	6% (4-8.5%)	200 (120-300)	8500 (5500-12000)
Equatorial Guinea	2000	Male	15 to 19	25% (14-40%)	75% (60-86%)	0.84 (0.38-1.6)	3.3% (1.7-5.7%)	4.3% (1.9-7.9%)	1.9 (0.77-3.7)	170 (81-290)
Equatorial Guinea	2000	Male	20 to 24	37% (23-55%)	63% (45-77%)	1.8 (0.89-3)	7.7% (4.5-12%)	9.2% (4.7-15%)	5.2 (2.5-9.2)	400 (220-660)
Equatorial Guinea	2000	Male	25 to 29	45% (28-62%)	55% (38-72%)	2.5 (1.4-4)	9.8% (5.8-14%)	11% (6-17%)	11 (6-18)	780 (440-1200)
Equatorial Guinea	2000	Male	30 to 34	47% (29-66%)	53% (34-71%)	2.8 (1.7-4.1)	11% (6.6-15%)	12% (7.3-18%)	19 (11-30)	1200 (710-1800)
Equatorial Guinea	2000	Male	35 to 39	47% (31-65%)	53% (35-69%)	2.7 (1.6-4.2)	11% (6.8-15%)	13% (7.5-19%)	25 (14-39)	1300 (800-2000)
Equatorial Guinea	2000	Male	40 to 44	49% (32-67%)	51% (33-68%)	2.8 (1.6-4.3)	12% (7.1-17%)	14% (8.2-20%)	27 (15-44)	1300 (750-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	2000	Male	45 to 49	52% (34-69%)	48% (31-66%)	2.9 (1.6-4.9)	13% (7.8-18%)	15% (9.3-22%)	24 (13-39)	1000 (570-1600)
Equatorial Guinea	2000	Male	50 to 54	52% (34-70%)	48% (30-66%)	2.9 (1.5-4.9)	13% (7.4-19%)	16% (9.3-24%)	17 (8.5-28)	630 (320-1000)
Equatorial Guinea	2000	Male	55 to 59	50% (33-68%)	50% (32-67%)	2.6 (1.3-4.4)	12% (6.4-18%)	14% (8.1-22%)	15 (7.4-26)	490 (250-820)
Equatorial Guinea	2000	Male	60 to 64	49% (31-68%)	51% (32-69%)	2.2 (1.1-4.1)	9.6% (4.9-15%)	12% (5.8-18%)	15 (6.8-27)	410 (190-730)
Equatorial Guinea	2000	Male	65 to 69	49% (32-67%)	51% (33-68%)	2 (1.1-3.4)	8% (3.9-13%)	9.6% (4.7-16%)	15 (7.1-26)	340 (160-580)
Equatorial Guinea	2000	Male	70 to 74	50% (33-67%)	50% (33-67%)	1.8 (0.93-3.1)	6.3% (2.5-11%)	7.4% (2.9-13%)	13 (5-24)	230 (93-430)
Equatorial Guinea	2000	Male	75 to 79	48% (32-67%)	52% (33-68%)	1.5 (0.72-2.5)	4.8% (1.2-9.2%)	5.6% (1.4-11%)	9 (2.3-18)	130 (33-250)
Equatorial Guinea	2000	Male	80 to 84	43% (26-62%)	57% (38-74%)	1.1 (0.54-2)	3.4% (0.79-6.7%)	3.9% (0.88-7.8%)	4.5 (1-9.2)	48 (11-97)
Equatorial Guinea	2000	Male	85 to 89	38% (23-54%)	62% (46-77%)	0.86 (0.4-1.5)	2.3% (0.58-4.6%)	2.7% (0.58-5.3%)	1.5 (0.34-3.1)	11 (2.8-23)
Equatorial Guinea	2000	Male	90 to 94	37% (23-54%)	63% (46-77%)	0.74 (0.34-1.4)	1.5% (0.14-3.4%)	1.7% (0.045-4%)	0.34 (0.0087-0.79)	1.8 (0.17-4)
Equatorial Guinea	2000	Male	95 plus	40% (22-61%)	60% (39-78%)	0.69 (0.26-1.4)	1.6% (0.087-4.1%)	1.8% (-0.065-4.6%)	0.083 (-0.0031-0.22)	0.31 (0.017-0.76)
Equatorial Guinea	2005	Female	All Ages	23% (19-29%)	77% (71-81%)	1.3 (1-1.6)	2.2% (1.3-3.2%)	3.7% (2.2-5.6%)	110 (54-190)	3900 (2200-6400)
Equatorial Guinea	2005	Female	15 to 19	21% (13-33%)	79% (67-87%)	0.97 (0.47-1.8)	1.9% (1.1-3.3%)	2.6% (1.2-4.7%)	1.3 (0.48-2.6)	130 (62-220)
Equatorial Guinea	2005	Female	20 to 24	30% (18-45%)	70% (55-82%)	1.9 (0.97-3.2)	3.1% (1.7-5%)	3.4% (1.5-6.2%)	2.3 (0.89-4.4)	220 (110-350)
Equatorial Guinea	2005	Female	25 to 29	34% (20-50%)	66% (50-80%)	2.6 (1.4-4.2)	2.9% (1.5-4.8%)	3% (1.2-5.5%)	2.9 (1.2-5)	220 (120-360)
Equatorial Guinea	2005	Female	30 to 34	36% (22-53%)	64% (47-78%)	2.8 (1.6-4.4)	3.3% (1.8-5.2%)	3.5% (1.7-6%)	5 (2.4-8.6)	340 (190-550)
Equatorial Guinea	2005	Female	35 to 39	38% (24-55%)	62% (45-76%)	2.8 (1.6-4.3)	3.8% (2.1-5.9%)	4.2% (2.1-6.9%)	7.7 (3.5-13)	450 (240-730)
Equatorial Guinea	2005	Female	40 to 44	38% (24-56%)	62% (44-76%)	2.7 (1.5-4.3)	4.5% (2.6-6.7%)	5.3% (2.9-8%)	10 (4.7-18)	520 (270-870)
Equatorial Guinea	2005	Female	45 to 49	37% (23-54%)	63% (46-77%)	2.6 (1.4-4.3)	5.7% (3.3-8.8%)	7% (3.8-11%)	12 (5.5-22)	520 (260-930)
Equatorial Guinea	2005	Female	50 to 54	34% (20-51%)	66% (49-80%)	2.4 (1.2-3.9)	6.4% (3.6-10%)	8% (4.5-13%)	9.9 (4.2-19)	380 (180-710)
Equatorial Guinea	2005	Female	55 to 59	32% (19-49%)	68% (51-81%)	2.2 (1.1-3.9)	7% (3.7-11%)	8.9% (4.8-14%)	7.4 (3.2-14)	240 (110-460)
Equatorial Guinea	2005	Female	60 to 64	31% (19-48%)	69% (52-81%)	2.1 (0.98-3.8)	6.5% (3.2-10%)	8.1% (4-13%)	7.7 (3.1-16)	220 (87-420)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	2005	Female	65 to 69	33% (22-48%)	67% (52-78%)	2 (1.1-3.4)	6.8% (3.3-11%)	8.3% (4.3-13%)	11 (4.3-21)	240 (100-460)
Equatorial Guinea	2005	Female	70 to 74	33% (21-48%)	67% (52-79%)	1.8 (0.91-3.2)	6.1% (2.5-11%)	7.4% (3-13%)	12 (4.5-24)	220 (81-430)
Equatorial Guinea	2005	Female	75 to 79	30% (19-46%)	70% (54-81%)	1.5 (0.69-2.8)	4.8% (1.4-9.2%)	5.8% (1.7-11%)	10 (3-21)	140 (42-300)
Equatorial Guinea	2005	Female	80 to 84	27% (16-43%)	73% (57-84%)	1.2 (0.59-2.1)	3.8% (1.2-7.7%)	4.6% (1.4-9%)	6.9 (2-14)	72 (21-150)
Equatorial Guinea	2005	Female	85 to 89	26% (15-39%)	74% (61-85%)	1.1 (0.5-1.9)	3% (0.77-5.9%)	3.5% (0.89-7.1%)	3 (0.77-6.1)	22 (5.9-46)
Equatorial Guinea	2005	Female	90 to 94	27% (17-40%)	73% (60-83%)	1.1 (0.48-2)	2.4% (0.52-5.2%)	2.9% (0.58-6.3%)	0.82 (0.16-1.8)	4.2 (0.88-9.4)
Equatorial Guinea	2005	Female	95 plus	31% (17-48%)	69% (52-83%)	1.1 (0.4-2.1)	2.3% (0.23-5.7%)	2.7% (0.2-6.7%)	0.19 (0.015-0.5)	0.68 (0.065-1.7)
Equatorial Guinea	2005	Male	All Ages	33% (26-40%)	67% (60-74%)	1.9 (1.5-2.3)	4% (2.8-5.6%)	6.1% (4.1-8.4%)	200 (120-300)	8200 (5300-12000)
Equatorial Guinea	2005	Male	15 to 19	24% (14-39%)	76% (61-86%)	1.1 (0.55-2.1)	3.6% (2-6.1%)	4.8% (2.3-8.9%)	2.6 (1.2-5.2)	240 (130-430)
Equatorial Guinea	2005	Male	20 to 24	37% (22-54%)	63% (46-78%)	2.4 (1.2-3.9)	8% (4.8-12%)	9.6% (5.1-16%)	6.7 (3.2-12)	530 (300-880)
Equatorial Guinea	2005	Male	25 to 29	44% (27-62%)	56% (38-73%)	3.4 (1.9-5.4)	8.2% (4.7-12%)	9.3% (4.6-15%)	8.6 (4.4-14)	610 (360-940)
Equatorial Guinea	2005	Male	30 to 34	46% (29-65%)	54% (35-71%)	3.8 (2.2-5.7)	7.8% (4.7-11%)	8.6% (4.7-13%)	13 (7.2-21)	810 (500-1300)
Equatorial Guinea	2005	Male	35 to 39	46% (29-63%)	54% (37-71%)	3.7 (2.1-5.7)	8% (4.9-12%)	9% (5.2-14%)	18 (10-30)	1000 (620-1600)
Equatorial Guinea	2005	Male	40 to 44	48% (32-66%)	52% (34-68%)	3.7 (2.1-5.7)	8.7% (5.4-12%)	10% (6.1-15%)	24 (13-40)	1200 (690-1900)
Equatorial Guinea	2005	Male	45 to 49	51% (32-69%)	49% (31-68%)	4 (2.2-6.3)	10% (6.7-15%)	12% (7.9-17%)	27 (15-44)	1200 (690-1900)
Equatorial Guinea	2005	Male	50 to 54	52% (33-69%)	48% (31-67%)	3.9 (2.1-6.5)	11% (7.3-16%)	14% (8.9-20%)	23 (12-37)	880 (490-1400)
Equatorial Guinea	2005	Male	55 to 59	50% (32-68%)	50% (32-68%)	3.5 (1.8-5.8)	11% (6.9-16%)	14% (8.6-20%)	15 (7.9-25)	500 (270-810)
Equatorial Guinea	2005	Male	60 to 64	48% (30-68%)	52% (32-70%)	3.1 (1.5-5.5)	9.8% (5.5-15%)	12% (6.8-18%)	13 (6.6-23)	380 (190-640)
Equatorial Guinea	2005	Male	65 to 69	49% (32-65%)	51% (35-68%)	2.7 (1.5-4.6)	8.7% (4.9-13%)	11% (6-16%)	14 (6.7-23)	320 (160-530)
Equatorial Guinea	2005	Male	70 to 74	49% (32-66%)	51% (34-68%)	2.4 (1.3-4)	7.3% (3.4-12%)	8.8% (4-14%)	13 (5.6-23)	250 (110-430)
Equatorial Guinea	2005	Male	75 to 79	48% (31-66%)	52% (34-69%)	2 (0.97-3.5)	5.8% (2.2-10%)	6.9% (2.6-12%)	9.9 (3.5-19)	140 (51-270)
Equatorial Guinea	2005	Male	80 to 84	43% (25-61%)	57% (39-75%)	1.6 (0.74-2.6)	4.3% (1.6-8%)	5.1% (1.8-9.2%)	5.6 (1.9-11)	60 (21-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	2005	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.2 (0.56-2.1)	3% (1.1-5.5%)	3.5% (1.2-6.4%)	2.1 (0.71-4)	16 (5.8-30)
Equatorial Guinea	2005	Male	90 to 94	37% (22-53%)	63% (47-78%)	1 (0.45-1.9)	2.1% (0.59-4.4%)	2.4% (0.56-5.1%)	0.51 (0.12-1.1)	2.8 (0.83-6)
Equatorial Guinea	2005	Male	95 plus	40% (21-61%)	60% (39-79%)	0.96 (0.36-1.9)	2.2% (0.48-5.1%)	2.4% (0.35-5.9%)	0.12 (0.018-0.31)	0.45 (0.1-1.1)
Equatorial Guinea	2010	Female	All Ages	23% (19-29%)	77% (71-81%)	1.6 (1.1-2.1)	2.3% (1.4-3.5%)	4% (2.3-6%)	120 (60-200)	4300 (2400-6900)
Equatorial Guinea	2010	Female	15 to 19	21% (13-33%)	79% (67-87%)	1.1 (0.56-2.1)	1.9% (1.1-3.2%)	2.5% (1.2-4.7%)	1.3 (0.48-2.6)	130 (66-240)
Equatorial Guinea	2010	Female	20 to 24	30% (18-44%)	70% (56-82%)	2.2 (1.1-3.9)	3.3% (1.8-5.2%)	3.6% (1.5-6.5%)	2.9 (1.2-5.4)	280 (150-460)
Equatorial Guinea	2010	Female	25 to 29	34% (20-51%)	66% (49-80%)	3 (1.6-4.9)	2.8% (1.4-4.7%)	2.9% (1.1-5.4%)	3.4 (1.5-6.2)	280 (150-460)
Equatorial Guinea	2010	Female	30 to 34	36% (22-53%)	64% (47-78%)	3.4 (1.8-5.5)	2.8% (1.5-4.5%)	2.8% (1.3-5%)	3.8 (1.7-6.8)	270 (150-430)
Equatorial Guinea	2010	Female	35 to 39	38% (24-56%)	62% (44-76%)	3.3 (1.8-5.6)	3.1% (1.7-4.9%)	3.3% (1.6-5.8%)	5.7 (2.5-9.7)	340 (180-550)
Equatorial Guinea	2010	Female	40 to 44	38% (23-56%)	62% (44-77%)	3.2 (1.7-5.3)	3.8% (2.2-5.9%)	4.3% (2.4-7%)	9.1 (4.3-17)	480 (250-820)
Equatorial Guinea	2010	Female	45 to 49	37% (21-54%)	63% (46-79%)	3.1 (1.6-5.2)	5.1% (3-7.8%)	6.1% (3.4-9.4%)	13 (6-24)	590 (300-1000)
Equatorial Guinea	2010	Female	50 to 54	34% (20-53%)	66% (47-80%)	2.8 (1.4-4.7)	6.1% (3.5-9.4%)	7.6% (4.4-12%)	14 (6.5-27)	550 (260-1000)
Equatorial Guinea	2010	Female	55 to 59	31% (19-49%)	69% (51-81%)	2.6 (1.3-4.5)	7.1% (3.9-11%)	9% (5-14%)	12 (5.3-23)	410 (180-750)
Equatorial Guinea	2010	Female	60 to 64	31% (19-48%)	69% (52-81%)	2.5 (1.1-4.6)	6.7% (3.4-11%)	8.4% (4.4-13%)	8.1 (3.4-16)	230 (100-440)
Equatorial Guinea	2010	Female	65 to 69	33% (21-47%)	67% (53-79%)	2.4 (1.2-4.1)	7.2% (3.6-12%)	9% (4.7-14%)	10 (4.3-19)	230 (100-440)
Equatorial Guinea	2010	Female	70 to 74	32% (21-48%)	68% (52-79%)	2.1 (1-3.9)	6.7% (2.9-12%)	8.2% (3.6-14%)	11 (4.5-22)	210 (84-400)
Equatorial Guinea	2010	Female	75 to 79	30% (19-45%)	70% (55-81%)	1.8 (0.77-3.2)	5.5% (1.8-10%)	6.7% (2.2-12%)	11 (3.4-22)	150 (48-310)
Equatorial Guinea	2010	Female	80 to 84	27% (16-42%)	73% (58-84%)	1.4 (0.66-2.5)	4.5% (1.5-8.5%)	5.4% (1.9-10%)	7.8 (2.6-16)	83 (28-170)
Equatorial Guinea	2010	Female	85 to 89	26% (15-39%)	74% (61-85%)	1.3 (0.59-2.3)	3.6% (1.1-7.4%)	4.3% (1.3-8.7%)	4.1 (1.2-8.9)	31 (9.2-67)
Equatorial Guinea	2010	Female	90 to 94	27% (17-40%)	73% (60-83%)	1.3 (0.57-2.4)	3.1% (0.79-6.6%)	3.6% (0.91-7.9%)	1.3 (0.34-3)	7 (1.8-15)
Equatorial Guinea	2010	Female	95 plus	31% (17-50%)	69% (50-83%)	1.3 (0.49-2.7)	2.9% (0.53-7%)	3.4% (0.55-8.5%)	0.31 (0.05-0.75)	1.1 (0.19-2.6)
Equatorial Guinea	2010	Male	All Ages	32% (26-41%)	68% (59-74%)	2.2 (1.6-2.9)	4.2% (2.9-5.9%)	6.4% (4.3-9.1%)	210 (120-330)	8400 (5300-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	2010	Male	15 to 19	24% (14-38%)	76% (62-86%)	1.3 (0.64-2.5)	3.4% (1.9-5.5%)	4.4% (2.1-8%)	2.4 (1-4.8)	230 (120-410)
Equatorial Guinea	2010	Male	20 to 24	36% (22-54%)	64% (46-78%)	2.8 (1.4-4.8)	8% (4.9-12%)	9.9% (5.2-16%)	7.3 (3.5-13)	620 (350-1000)
Equatorial Guinea	2010	Male	25 to 29	44% (27-61%)	56% (39-73%)	4 (2.2-6.4)	7.9% (4.6-12%)	9% (4.5-15%)	9.2 (5.1-16)	690 (420-1100)
Equatorial Guinea	2010	Male	30 to 34	45% (28-64%)	55% (36-72%)	4.4 (2.5-7)	6.8% (4.2-10%)	7.4% (4-12%)	9.2 (5-16)	620 (380-950)
Equatorial Guinea	2010	Male	35 to 39	46% (29-64%)	54% (36-71%)	4.3 (2.4-7)	6.7% (4.2-10%)	7.4% (4.2-12%)	13 (7.1-22)	750 (450-1200)
Equatorial Guinea	2010	Male	40 to 44	47% (31-66%)	53% (34-69%)	4.4 (2.5-7)	7.6% (5-11%)	8.7% (5.4-13%)	20 (11-33)	1000 (590-1600)
Equatorial Guinea	2010	Male	45 to 49	51% (33-68%)	49% (32-67%)	4.7 (2.5-7.6)	9.2% (6-13%)	11% (7-16%)	27 (15-44)	1200 (710-1900)
Equatorial Guinea	2010	Male	50 to 54	51% (33-69%)	49% (31-67%)	4.6 (2.4-7.6)	11% (7.2-15%)	14% (9-19%)	29 (16-49)	1100 (660-1800)
Equatorial Guinea	2010	Male	55 to 59	49% (31-68%)	51% (32-69%)	4.1 (2.1-6.8)	11% (7.1-16%)	14% (9.2-20%)	24 (13-39)	800 (440-1300)
Equatorial Guinea	2010	Male	60 to 64	47% (29-66%)	53% (34-71%)	3.6 (1.7-6.5)	10% (6.1-15%)	13% (7.7-18%)	16 (8.3-27)	460 (250-740)
Equatorial Guinea	2010	Male	65 to 69	48% (31-66%)	52% (34-69%)	3.2 (1.7-5.5)	9.5% (5.5-14%)	12% (6.8-17%)	15 (7.2-25)	340 (180-580)
Equatorial Guinea	2010	Male	70 to 74	49% (32-65%)	51% (35-68%)	2.8 (1.4-4.9)	8.1% (3.9-13%)	9.9% (4.9-16%)	14 (6.1-24)	260 (120-450)
Equatorial Guinea	2010	Male	75 to 79	47% (30-65%)	53% (35-70%)	2.3 (1.1-4.2)	6.5% (2.6-11%)	7.8% (3-14%)	11 (4-21)	160 (61-300)
Equatorial Guinea	2010	Male	80 to 84	42% (25-60%)	58% (40-75%)	1.8 (0.86-3.2)	5% (1.8-9.4%)	5.8% (2.2-11%)	6.7 (2.4-13)	72 (26-140)
Equatorial Guinea	2010	Male	85 to 89	36% (22-53%)	64% (47-78%)	1.4 (0.63-2.4)	3.6% (1.3-6.8%)	4.1% (1.5-7.9%)	2.9 (0.98-5.7)	22 (7.9-44)
Equatorial Guinea	2010	Male	90 to 94	36% (22-53%)	64% (47-78%)	1.2 (0.51-2.3)	2.6% (0.79-5.7%)	2.9% (0.77-6.7%)	0.79 (0.22-1.8)	4.3 (1.3-9.7)
Equatorial Guinea	2010	Male	95 plus	39% (20-60%)	61% (40-80%)	1.1 (0.42-2.3)	2.6% (0.7-6.1%)	2.8% (0.62-6.9%)	0.18 (0.038-0.44)	0.66 (0.18-1.6)
Equatorial Guinea	2016	Female	All Ages	23% (19-29%)	77% (71-81%)	1.7 (1-2.8)	2.6% (1.5-4%)	4.4% (2.3-6.8%)	120 (51-210)	4300 (2200-7100)
Equatorial Guinea	2016	Female	15 to 19	21% (13-32%)	79% (68-87%)	1.2 (0.52-2.5)	1.7% (0.88-3%)	2% (0.76-4.1%)	1 (0.34-2.3)	120 (60-220)
Equatorial Guinea	2016	Female	20 to 24	29% (18-44%)	71% (56-82%)	2.4 (1.1-4.6)	3.3% (1.9-5.1%)	3.8% (1.6-6.9%)	2.3 (0.88-4.7)	260 (140-430)
Equatorial Guinea	2016	Female	25 to 29	33% (19-50%)	67% (50-81%)	3.2 (1.5-5.7)	3.2% (1.7-5.1%)	3.4% (1.2-6.6%)	3.5 (1.4-6.7)	320 (170-520)
Equatorial Guinea	2016	Female	30 to 34	36% (21-52%)	64% (48-79%)	3.6 (1.7-6.6)	3% (1.6-4.9%)	3.2% (1.4-5.8%)	4.1 (1.7-7.4)	310 (170-510)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	2016	Female	35 to 39	37% (23-54%)	63% (46-77%)	3.6 (1.6-6.9)	3% (1.6-4.8%)	3.3% (1.4-5.9%)	3.9 (1.6-7.1)	250 (130-420)
Equatorial Guinea	2016	Female	40 to 44	37% (23-55%)	63% (45-77%)	3.5 (1.5-6.6)	3.6% (2-5.7%)	4.2% (2.1-6.9%)	5.9 (2.3-11)	320 (150-540)
Equatorial Guinea	2016	Female	45 to 49	36% (21-54%)	64% (46-79%)	3.3 (1.5-6.4)	4.9% (2.7-7.7%)	6.2% (3.1-9.8%)	10 (4.3-19)	490 (220-830)
Equatorial Guinea	2016	Female	50 to 54	34% (20-51%)	66% (49-80%)	3 (1.4-5.6)	5.9% (3.1-9.1%)	7.5% (4.1-12%)	14 (6-27)	580 (260-1000)
Equatorial Guinea	2016	Female	55 to 59	31% (19-48%)	69% (52-81%)	2.8 (1.2-5.2)	6.8% (3.6-10%)	9% (5-14%)	17 (6.6-32)	560 (240-1100)
Equatorial Guinea	2016	Female	60 to 64	31% (19-49%)	69% (51-81%)	2.6 (1.1-5.3)	6.6% (3.2-10%)	8.5% (4.3-13%)	13 (5.1-26)	390 (160-710)
Equatorial Guinea	2016	Female	65 to 69	33% (21-48%)	67% (52-79%)	2.6 (1.2-5.1)	6.8% (3.3-11%)	8.7% (4.2-14%)	10 (3.9-20)	240 (95-450)
Equatorial Guinea	2016	Female	70 to 74	32% (21-48%)	68% (52-79%)	2.3 (1-4.7)	6.4% (2.6-11%)	8% (3.4-13%)	9.5 (3.4-19)	180 (64-350)
Equatorial Guinea	2016	Female	75 to 79	30% (18-45%)	70% (55-82%)	1.9 (0.74-4.1)	5.4% (1.6-9.8%)	6.6% (2.1-12%)	9 (2.5-18)	130 (38-260)
Equatorial Guinea	2016	Female	80 to 84	27% (16-42%)	73% (58-84%)	1.6 (0.64-3.2)	4.6% (1.4-9.1%)	5.5% (1.7-11%)	7.4 (2.1-16)	79 (23-170)
Equatorial Guinea	2016	Female	85 to 89	26% (15-39%)	74% (61-85%)	1.4 (0.58-2.9)	3.8% (1.1-7.9%)	4.5% (1.2-9.4%)	4.7 (1.2-11)	36 (9.2-79)
Equatorial Guinea	2016	Female	90 to 94	27% (17-41%)	73% (59-83%)	1.4 (0.53-2.8)	3.3% (0.72-7.1%)	3.9% (0.84-8.4%)	1.9 (0.42-4.3)	9.8 (2.3-22)
Equatorial Guinea	2016	Female	95 plus	31% (17-49%)	69% (51-83%)	1.4 (0.48-3)	3.1% (0.5-7.5%)	3.7% (0.54-9.1%)	0.51 (0.075-1.2)	1.8 (0.28-4.2)
Equatorial Guinea	2016	Male	All Ages	32% (26-40%)	68% (60-74%)	2.4 (1.4-3.9)	4.7% (3-6.8%)	7.1% (4.5-10%)	230 (120-380)	9100 (5600-14000)
Equatorial Guinea	2016	Male	15 to 19	24% (14-38%)	76% (62-86%)	1.4 (0.58-2.9)	3% (1.6-5.1%)	3.5% (1.5-6.8%)	2.4 (0.89-5.1)	240 (120-440)
Equatorial Guinea	2016	Male	20 to 24	36% (22-52%)	64% (48-78%)	3 (1.3-5.6)	7.9% (4.8-12%)	9.7% (4.9-16%)	7.3 (3.2-14)	650 (350-1100)
Equatorial Guinea	2016	Male	25 to 29	43% (26-61%)	57% (39-74%)	4.2 (2-7.4)	8.3% (4.9-12%)	9.7% (4.8-16%)	11 (5.6-20)	860 (520-1400)
Equatorial Guinea	2016	Male	30 to 34	45% (27-64%)	55% (36-73%)	4.7 (2.3-8.4)	7.1% (4.2-11%)	7.9% (4-13%)	11 (6.1-19)	780 (480-1200)
Equatorial Guinea	2016	Male	35 to 39	45% (29-64%)	55% (36-71%)	4.7 (2.1-8.5)	6.5% (3.9-9.8%)	7.2% (3.8-12%)	10 (5.4-18)	640 (380-1000)
Equatorial Guinea	2016	Male	40 to 44	47% (31-66%)	53% (34-69%)	4.8 (2.2-8.7)	7.1% (4.5-10%)	8.1% (4.9-12%)	14 (7.5-25)	760 (430-1200)
Equatorial Guinea	2016	Male	45 to 49	50% (32-68%)	50% (32-68%)	5 (2.3-9.4)	8.8% (5.7-13%)	11% (6.7-16%)	23 (12-39)	1100 (600-1700)
Equatorial Guinea	2016	Male	50 to 54	51% (32-69%)	49% (31-68%)	4.9 (2.2-9)	10% (6.6-14%)	13% (8.4-18%)	31 (16-52)	1200 (680-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Equatorial Guinea	2016	Male	55 to 59	49% (31-67%)	51% (33-69%)	4.3 (1.9-8.3)	11% (6.9-15%)	14% (9.1-19%)	33 (17-55)	1100 (590-1800)
Equatorial Guinea	2016	Male	60 to 64	47% (30-67%)	53% (33-70%)	3.8 (1.7-7.6)	10% (5.8-14%)	13% (7.6-18%)	27 (13-47)	780 (390-1300)
Equatorial Guinea	2016	Male	65 to 69	48% (31-66%)	52% (34-69%)	3.4 (1.6-6.4)	9.3% (5-14%)	12% (6.3-17%)	19 (8.4-32)	440 (210-740)
Equatorial Guinea	2016	Male	70 to 74	49% (31-66%)	51% (34-69%)	3 (1.3-5.8)	8% (3.6-13%)	9.8% (4.5-16%)	14 (5.6-25)	270 (110-480)
Equatorial Guinea	2016	Male	75 to 79	47% (30-65%)	53% (35-70%)	2.5 (1-5.1)	6.6% (2.4-12%)	8% (2.8-14%)	11 (3.8-22)	170 (57-320)
Equatorial Guinea	2016	Male	80 to 84	42% (25-59%)	58% (41-75%)	1.9 (0.8-3.8)	5.1% (1.7-9.8%)	6.1% (2-12%)	7.3 (2.3-15)	79 (26-160)
Equatorial Guinea	2016	Male	85 to 89	36% (21-52%)	64% (48-79%)	1.5 (0.61-2.9)	3.8% (1.2-7.7%)	4.4% (1.4-9.1%)	3.6 (1.1-7.7)	28 (9.1-60)
Equatorial Guinea	2016	Male	90 to 94	35% (21-52%)	65% (48-79%)	1.2 (0.51-2.5)	2.7% (0.71-6.1%)	3.1% (0.69-7.1%)	1.1 (0.26-2.7)	6.1 (1.6-14)
Equatorial Guinea	2016	Male	95 plus	38% (20-59%)	62% (41-80%)	1.2 (0.4-2.5)	2.6% (0.57-6.3%)	2.9% (0.47-7.2%)	0.26 (0.043-0.67)	0.99 (0.22-2.4)
Gabon	1990	Female	All Ages	24% (20-30%)	76% (70-80%)	1.8 (1.5-2.2)	4.7% (3.5-6.1%)	8.1% (5.7-11%)	430 (300-590)	12000 (9100-16000)
Gabon	1990	Female	15 to 19	22% (14-35%)	78% (65-86%)	1.4 (0.7-2.4)	2.7% (1.4-4.5%)	3.9% (1.6-7.1%)	2.4 (0.96-4.5)	230 (120-390)
Gabon	1990	Female	20 to 24	31% (19-46%)	69% (54-81%)	2.7 (1.5-4.3)	4.7% (2.8-7%)	6.2% (3.1-9.8%)	4.8 (2.4-7.7)	430 (250-630)
Gabon	1990	Female	25 to 29	35% (21-52%)	65% (48-79%)	3.6 (2-5.5)	6.2% (3.8-8.8%)	8.4% (4.4-13%)	6.5 (3.5-9.9)	500 (310-730)
Gabon	1990	Female	30 to 34	38% (23-55%)	62% (45-77%)	4 (2.3-6)	7.7% (5.1-11%)	11% (6.6-16%)	9 (5.4-14)	590 (380-850)
Gabon	1990	Female	35 to 39	39% (25-57%)	61% (43-75%)	3.9 (2.2-6.3)	8.3% (5.2-12%)	12% (6.8-16%)	11 (6.4-17)	650 (390-940)
Gabon	1990	Female	40 to 44	40% (25-57%)	60% (43-75%)	3.8 (2-6.2)	9.2% (6-13%)	13% (8-18%)	15 (9-22)	740 (480-1100)
Gabon	1990	Female	45 to 49	39% (24-56%)	61% (44-76%)	3.7 (2-6.1)	11% (7-15%)	15% (9.5-20%)	23 (14-34)	990 (640-1400)
Gabon	1990	Female	50 to 54	36% (22-54%)	64% (46-78%)	3.4 (1.8-5.6)	11% (7.5-16%)	15% (9.9-21%)	33 (21-49)	1200 (790-1800)
Gabon	1990	Female	55 to 59	34% (21-51%)	66% (49-79%)	3.1 (1.6-5.2)	12% (7.3-16%)	15% (9.2-21%)	46 (28-67)	1500 (910-2200)
Gabon	1990	Female	60 to 64	33% (20-50%)	67% (50-80%)	2.9 (1.4-5)	11% (6.4-15%)	13% (7.7-19%)	54 (31-79)	1500 (860-2100)
Gabon	1990	Female	65 to 69	35% (22-50%)	65% (50-78%)	2.9 (1.5-4.6)	11% (6.3-15%)	13% (7.6-18%)	64 (38-93)	1500 (860-2100)
Gabon	1990	Female	70 to 74	34% (23-49%)	66% (51-77%)	2.6 (1.3-4.4)	9.5% (4.9-14%)	11% (5.8-17%)	65 (34-100)	1200 (610-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	1990	Female	75 to 79	32% (20-47%)	68% (53-80%)	2.1 (1-3.8)	7.6% (3.5-13%)	8.8% (4.1-15%)	48 (22-81)	670 (310-1100)
Gabon	1990	Female	80 to 84	29% (17-44%)	71% (56-83%)	1.8 (0.88-3)	6.1% (2.7-11%)	7.1% (3.1-12%)	32 (14-55)	330 (140-580)
Gabon	1990	Female	85 to 89	27% (17-41%)	73% (59-83%)	1.6 (0.79-2.7)	4.8% (2-8.7%)	5.6% (2.3-10%)	15 (6.3-28)	110 (47-200)
Gabon	1990	Female	90 to 94	29% (18-43%)	71% (57-82%)	1.5 (0.74-2.7)	3.9% (1.5-7.3%)	4.6% (1.7-8.6%)	4.7 (1.7-8.7)	24 (9.3-45)
Gabon	1990	Female	95 plus	33% (18-50%)	67% (50-82%)	1.5 (0.6-3)	3.6% (1-7.4%)	4.1% (1.1-8.8%)	1.1 (0.31-2.4)	3.9 (1.1-8.1)
Gabon	1990	Male	All Ages	32% (26-39%)	68% (61-74%)	2.2 (1.9-2.6)	8.1% (6.5-10%)	12% (9.3-15%)	770 (570-980)	26000 (20000-32000)
Gabon	1990	Male	15 to 19	25% (15-39%)	75% (61-85%)	1.5 (0.76-2.7)	5.2% (2.7-8.7%)	6.6% (3.2-12%)	7.1 (3.4-13)	580 (300-980)
Gabon	1990	Male	20 to 24	37% (23-54%)	63% (46-77%)	3.1 (1.7-5.1)	12% (7.6-17%)	15% (8.4-21%)	21 (12-32)	1600 (950-2300)
Gabon	1990	Male	25 to 29	45% (27-62%)	55% (38-73%)	4.5 (2.6-6.7)	16% (10-22%)	19% (11-27%)	28 (17-40)	1900 (1200-2600)
Gabon	1990	Male	30 to 34	47% (29-64%)	53% (36-71%)	5 (2.9-7.3)	17% (12-23%)	21% (14-28%)	32 (21-44)	1900 (1300-2600)
Gabon	1990	Male	35 to 39	47% (30-65%)	53% (35-70%)	4.9 (2.8-7.7)	18% (12-24%)	22% (15-29%)	38 (25-50)	2000 (1400-2700)
Gabon	1990	Male	40 to 44	49% (32-67%)	51% (33-68%)	5 (2.8-8)	20% (14-25%)	23% (17-31%)	49 (34-67)	2300 (1600-3200)
Gabon	1990	Male	45 to 49	52% (33-70%)	48% (30-67%)	5.3 (3-8.2)	21% (15-26%)	25% (18-31%)	63 (45-83)	2600 (1900-3500)
Gabon	1990	Male	50 to 54	52% (34-70%)	48% (30-66%)	5.1 (2.9-8.2)	21% (15-27%)	24% (17-31%)	79 (54-110)	2900 (2000-3900)
Gabon	1990	Male	55 to 59	50% (32-69%)	50% (31-68%)	4.6 (2.6-7.2)	19% (13-25%)	22% (16-29%)	89 (60-120)	2800 (1900-3800)
Gabon	1990	Male	60 to 64	48% (31-68%)	52% (32-69%)	4 (2.1-6.6)	16% (11-22%)	18% (12-25%)	91 (57-130)	2500 (1600-3400)
Gabon	1990	Male	65 to 69	49% (32-67%)	51% (33-68%)	3.6 (2-5.5)	14% (9.1-20%)	16% (10-22%)	88 (55-120)	2000 (1300-2800)
Gabon	1990	Male	70 to 74	50% (32-67%)	50% (33-68%)	3.1 (1.7-5.2)	12% (6.6-18%)	13% (7.4-20%)	80 (44-120)	1400 (800-2200)
Gabon	1990	Male	75 to 79	48% (31-66%)	52% (34-69%)	2.6 (1.3-4.4)	9.4% (4.5-15%)	11% (5-17%)	58 (27-95)	810 (390-1300)
Gabon	1990	Male	80 to 84	43% (25-61%)	57% (39-75%)	2 (0.99-3.4)	6.8% (3-11%)	7.7% (3.3-13%)	31 (13-54)	320 (140-560)
Gabon	1990	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.5 (0.8-2.7)	4.8% (2.1-8.6%)	5.4% (2.3-9.8%)	10 (4.5-19)	77 (34-140)
Gabon	1990	Male	90 to 94	37% (23-54%)	63% (46-77%)	1.3 (0.66-2.4)	3.3% (1.2-6.2%)	3.6% (1.3-7.1%)	2.2 (0.76-4.3)	12 (4.4-23)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	1990	Male	95 plus	40% (21-60%)	60% (40-79%)	1.3 (0.47-2.4)	3.2% (0.97-6.7%)	3.5% (0.96-7.6%)	0.49 (0.13-1.1)	1.8 (0.54-3.8)
Gabon	1995	Female	All Ages	24% (20-29%)	76% (71-80%)	1.4 (1.2-1.6)	3.9% (2.8-5.2%)	6.3% (4.2-8.8%)	380 (250-540)	11000 (8000-14000)
Gabon	1995	Female	15 to 19	22% (14-35%)	78% (65-86%)	1.1 (0.54-1.8)	2.2% (1.2-3.6%)	3.1% (1.5-5.5%)	2.2 (1-3.9)	220 (120-350)
Gabon	1995	Female	20 to 24	31% (19-46%)	69% (54-81%)	2 (1.1-3.3)	3.7% (2.4-5.3%)	4.5% (2.5-7%)	4.5 (2.5-7.1)	420 (260-610)
Gabon	1995	Female	25 to 29	36% (22-52%)	64% (48-78%)	2.7 (1.6-4.1)	4.8% (3.2-6.6%)	5.9% (3.6-8.7%)	6.8 (4.2-10)	540 (360-740)
Gabon	1995	Female	30 to 34	38% (23-56%)	62% (44-77%)	3.1 (1.8-4.7)	6% (4.1-8.4%)	7.8% (4.9-12%)	9.3 (5.9-14)	620 (410-880)
Gabon	1995	Female	35 to 39	39% (25-57%)	61% (43-75%)	3 (1.7-4.7)	6.7% (4.3-9.3%)	8.8% (5.5-13%)	11 (6.9-16)	650 (410-910)
Gabon	1995	Female	40 to 44	40% (25-57%)	60% (43-75%)	3 (1.6-4.8)	7.6% (4.9-11%)	10% (6.5-14%)	14 (8.9-21)	730 (460-1000)
Gabon	1995	Female	45 to 49	39% (24-56%)	61% (44-76%)	2.8 (1.5-4.7)	8.8% (5.6-12%)	12% (7.4-16%)	20 (12-29)	880 (550-1300)
Gabon	1995	Female	50 to 54	36% (22-54%)	64% (46-78%)	2.6 (1.4-4.2)	9.4% (5.9-13%)	12% (7.6-17%)	29 (18-42)	1100 (690-1600)
Gabon	1995	Female	55 to 59	33% (20-51%)	67% (49-80%)	2.3 (1.2-3.8)	9.5% (5.8-14%)	12% (7.3-18%)	39 (23-57)	1200 (750-1800)
Gabon	1995	Female	60 to 64	33% (20-49%)	67% (51-80%)	2.2 (1.1-3.7)	8.5% (4.7-13%)	10% (5.6-16%)	46 (25-71)	1300 (690-1900)
Gabon	1995	Female	65 to 69	34% (22-50%)	66% (50-78%)	2.2 (1.2-3.5)	8.3% (4.7-13%)	10% (5.6-15%)	58 (31-90)	1300 (690-2000)
Gabon	1995	Female	70 to 74	34% (22-50%)	66% (50-78%)	2 (1-3.4)	7.2% (3.2-12%)	8.5% (3.9-14%)	54 (24-93)	990 (440-1700)
Gabon	1995	Female	75 to 79	31% (20-47%)	69% (53-80%)	1.6 (0.75-2.9)	5.5% (2-10%)	6.4% (2.2-12%)	42 (15-79)	590 (210-1100)
Gabon	1995	Female	80 to 84	28% (17-44%)	72% (56-83%)	1.3 (0.67-2.3)	4.2% (1.6-8.1%)	4.9% (1.9-9.5%)	26 (10-51)	280 (110-530)
Gabon	1995	Female	85 to 89	27% (17-40%)	73% (60-83%)	1.2 (0.6-2.1)	3.3% (1.2-6.4%)	3.8% (1.4-7.4%)	11 (4.1-22)	83 (31-170)
Gabon	1995	Female	90 to 94	29% (18-42%)	71% (58-82%)	1.2 (0.56-2.1)	2.6% (0.77-5.5%)	3% (0.87-6.4%)	3.5 (1-7.3)	18 (5.4-38)
Gabon	1995	Female	95 plus	33% (19-50%)	67% (50-81%)	1.2 (0.46-2.3)	2.4% (0.5-5.8%)	2.8% (0.52-6.7%)	0.87 (0.16-2.1)	3 (0.61-7)
Gabon	1995	Male	All Ages	32% (25-39%)	68% (61-75%)	1.7 (1.4-2)	6.8% (5.3-8.6%)	9.7% (7.1-13%)	670 (480-890)	24000 (18000-30000)
Gabon	1995	Male	15 to 19	25% (15-40%)	75% (60-85%)	1.2 (0.59-2.1)	4.3% (2.5-7.3%)	5.5% (2.8-9.6%)	6.7 (3.4-12)	560 (320-950)
Gabon	1995	Male	20 to 24	37% (23-55%)	63% (45-77%)	2.4 (1.4-3.8)	10% (6.5-15%)	12% (7.1-18%)	21 (13-32)	1600 (1000-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	1995	Male	25 to 29	45% (28-62%)	55% (38-72%)	3.5 (2.1-5.1)	13% (8.7-17%)	15% (9.3-21%)	30 (20-41)	2000 (1400-2700)
Gabon	1995	Male	30 to 34	47% (29-66%)	53% (34-71%)	3.9 (2.4-5.6)	14% (9.6-18%)	16% (11-21%)	33 (23-45)	2000 (1500-2700)
Gabon	1995	Male	35 to 39	47% (31-65%)	53% (35-69%)	3.8 (2.2-5.9)	14% (10-19%)	17% (11-23%)	36 (25-48)	1900 (1400-2600)
Gabon	1995	Male	40 to 44	49% (32-67%)	51% (33-68%)	3.9 (2.2-6.1)	15% (11-20%)	18% (13-24%)	45 (30-60)	2100 (1500-2800)
Gabon	1995	Male	45 to 49	52% (33-69%)	48% (31-67%)	4.1 (2.3-6.6)	17% (12-22%)	20% (14-26%)	58 (39-79)	2400 (1700-3300)
Gabon	1995	Male	50 to 54	52% (34-70%)	48% (30-66%)	4 (2.2-6.5)	17% (12-23%)	20% (14-27%)	67 (45-94)	2500 (1700-3400)
Gabon	1995	Male	55 to 59	50% (32-69%)	50% (31-68%)	3.5 (1.9-5.7)	16% (10-21%)	18% (12-25%)	74 (46-100)	2400 (1500-3300)
Gabon	1995	Male	60 to 64	49% (30-68%)	51% (32-70%)	3.1 (1.6-5.1)	13% (7.6-19%)	15% (8.7-22%)	77 (44-120)	2100 (1200-3100)
Gabon	1995	Male	65 to 69	49% (32-67%)	51% (33-68%)	2.8 (1.5-4.4)	11% (6.5-17%)	13% (7.5-19%)	76 (44-120)	1700 (990-2600)
Gabon	1995	Male	70 to 74	50% (33-67%)	50% (33-67%)	2.4 (1.3-4.1)	9% (4.6-15%)	10% (5.1-16%)	64 (31-110)	1200 (560-1900)
Gabon	1995	Male	75 to 79	48% (32-66%)	52% (34-68%)	2 (1-3.4)	7% (3-12%)	7.9% (3.4-14%)	46 (20-86)	650 (280-1200)
Gabon	1995	Male	80 to 84	43% (25-61%)	57% (39-75%)	1.6 (0.78-2.6)	4.8% (2-8.8%)	5.5% (2.2-10%)	25 (9.7-45)	260 (100-470)
Gabon	1995	Male	85 to 89	37% (23-55%)	63% (45-77%)	1.2 (0.61-2.1)	3.4% (1.3-6.3%)	3.8% (1.5-7.1%)	8.4 (3.2-16)	64 (25-120)
Gabon	1995	Male	90 to 94	37% (23-54%)	63% (46-77%)	1 (0.5-1.8)	2.2% (0.77-4.4%)	2.4% (0.78-4.9%)	1.7 (0.55-3.6)	9.3 (3.3-19)
Gabon	1995	Male	95 plus	40% (22-61%)	60% (39-78%)	0.95 (0.37-1.8)	2.2% (0.61-4.8%)	2.4% (0.54-5.5%)	0.39 (0.088-0.85)	1.4 (0.4-3.1)
Gabon	2000	Female	All Ages	23% (19-29%)	77% (71-81%)	1.2 (0.99-1.4)	3.5% (2.4-4.7%)	5.3% (3.4-7.7%)	390 (250-580)	12000 (8100-16000)
Gabon	2000	Female	15 to 19	21% (13-34%)	79% (66-87%)	0.9 (0.46-1.6)	1.8% (1-3.1%)	2.5% (1.1-4.6%)	2.2 (0.95-4.1)	220 (120-380)
Gabon	2000	Female	20 to 24	30% (18-44%)	70% (56-82%)	1.7 (0.96-2.8)	3.1% (1.9-4.7%)	3.5% (1.8-5.8%)	4.7 (2.4-7.6)	450 (270-670)
Gabon	2000	Female	25 to 29	34% (20-51%)	66% (49-80%)	2.3 (1.3-3.5)	3.6% (2.2-5.3%)	4% (2.1-6.5%)	7.7 (4.3-12)	610 (380-850)
Gabon	2000	Female	30 to 34	37% (22-54%)	63% (46-78%)	2.6 (1.5-4)	4.7% (3-6.9%)	5.4% (3.1-8.3%)	12 (7.1-18)	770 (500-1100)
Gabon	2000	Female	35 to 39	38% (24-56%)	62% (44-76%)	2.6 (1.5-4.1)	5.3% (3.3-7.6%)	6.3% (3.7-9.5%)	14 (8.5-21)	810 (510-1200)
Gabon	2000	Female	40 to 44	38% (24-56%)	62% (44-76%)	2.5 (1.3-4.2)	6.2% (3.9-8.9%)	7.7% (4.7-11%)	17 (10-25)	850 (530-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	2000	Female	45 to 49	38% (22-55%)	62% (45-78%)	2.4 (1.3-4.1)	7.4% (4.6-11%)	9.3% (5.6-13%)	23 (14-34)	1000 (600-1500)
Gabon	2000	Female	50 to 54	35% (21-53%)	65% (47-79%)	2.2 (1.2-3.6)	8% (4.8-12%)	9.9% (6-15%)	29 (17-45)	1100 (660-1700)
Gabon	2000	Female	55 to 59	32% (20-49%)	68% (51-80%)	2 (1-3.3)	8.2% (4.7-12%)	10% (5.8-15%)	39 (22-60)	1300 (710-1900)
Gabon	2000	Female	60 to 64	32% (19-48%)	68% (52-81%)	1.9 (0.94-3.2)	7.3% (3.6-12%)	8.7% (4.3-14%)	44 (22-72)	1200 (610-1900)
Gabon	2000	Female	65 to 69	33% (22-49%)	67% (51-78%)	1.9 (0.99-3)	7.1% (3.5-11%)	8.3% (4.2-13%)	57 (29-90)	1300 (670-2100)
Gabon	2000	Female	70 to 74	33% (22-49%)	67% (51-78%)	1.7 (0.85-2.9)	6.1% (2.4-11%)	7.1% (2.9-13%)	59 (24-110)	1100 (430-1900)
Gabon	2000	Female	75 to 79	31% (19-46%)	69% (54-81%)	1.4 (0.64-2.5)	4.6% (1.5-9%)	5.3% (1.7-10%)	43 (14-83)	600 (200-1200)
Gabon	2000	Female	80 to 84	28% (16-42%)	72% (58-84%)	1.1 (0.57-2)	3.5% (1.2-6.9%)	4% (1.3-7.9%)	28 (9.3-56)	290 (97-590)
Gabon	2000	Female	85 to 89	26% (16-40%)	74% (60-84%)	1 (0.51-1.7)	2.6% (0.83-5.2%)	3% (0.94-6%)	11 (3.6-23)	84 (27-170)
Gabon	2000	Female	90 to 94	28% (17-41%)	72% (59-83%)	0.98 (0.47-1.7)	2.1% (0.52-4.3%)	2.4% (0.55-5.1%)	3.1 (0.73-6.7)	16 (4-34)
Gabon	2000	Female	95 plus	31% (18-49%)	69% (51-82%)	0.99 (0.4-1.9)	1.9% (0.33-4.7%)	2.2% (0.3-5.3%)	0.79 (0.11-1.9)	2.8 (0.45-6.6)
Gabon	2000	Male	All Ages	31% (25-39%)	69% (61-75%)	1.6 (1.3-1.8)	5.3% (4-6.8%)	7.5% (5.5-10%)	540 (370-750)	20000 (15000-26000)
Gabon	2000	Male	15 to 19	24% (14-39%)	76% (61-86%)	1 (0.53-1.9)	3.6% (2-6%)	4.5% (2.1-8%)	5.6 (2.6-10)	500 (280-830)
Gabon	2000	Male	20 to 24	37% (23-54%)	63% (46-77%)	2.2 (1.2-3.4)	8.3% (5.1-12%)	9.5% (5.3-15%)	18 (10-29)	1400 (880-2100)
Gabon	2000	Male	25 to 29	44% (27-62%)	56% (38-73%)	3.1 (1.8-4.5)	9.4% (6-13%)	10% (6.1-16%)	27 (16-39)	1900 (1200-2600)
Gabon	2000	Male	30 to 34	46% (28-65%)	54% (35-72%)	3.5 (2-5.1)	9.5% (6.2-13%)	10% (6.3-15%)	31 (19-44)	1900 (1200-2700)
Gabon	2000	Male	35 to 39	46% (30-64%)	54% (36-70%)	3.4 (2-5.2)	9.6% (6.3-14%)	11% (6.7-16%)	33 (20-45)	1800 (1200-2500)
Gabon	2000	Male	40 to 44	48% (32-67%)	52% (33-68%)	3.4 (1.9-5.4)	11% (7.2-15%)	12% (7.9-17%)	38 (24-53)	1800 (1200-2500)
Gabon	2000	Male	45 to 49	51% (33-69%)	49% (31-67%)	3.7 (2-5.9)	12% (8-17%)	14% (9.4-20%)	45 (28-65)	1900 (1200-2800)
Gabon	2000	Male	50 to 54	52% (34-70%)	48% (30-66%)	3.6 (2-5.8)	13% (8.6-19%)	16% (10-22%)	53 (32-79)	2000 (1200-2900)
Gabon	2000	Male	55 to 59	49% (32-68%)	51% (32-68%)	3.2 (1.7-5.2)	13% (7.9-18%)	15% (9.4-21%)	55 (33-83)	1800 (1100-2600)
Gabon	2000	Male	60 to 64	48% (30-68%)	52% (32-70%)	2.8 (1.4-4.7)	11% (6.1-16%)	13% (6.9-19%)	57 (31-91)	1600 (880-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	2000	Male	65 to 69	49% (32-66%)	51% (34-68%)	2.5 (1.4-3.9)	9.3% (5.4-14%)	11% (6.2-16%)	59 (33-96)	1300 (750-2200)
Gabon	2000	Male	70 to 74	50% (33-66%)	50% (34-67%)	2.2 (1.2-3.6)	7.4% (3.7-12%)	8.6% (4.3-14%)	52 (26-88)	950 (470-1600)
Gabon	2000	Male	75 to 79	48% (31-66%)	52% (34-69%)	1.8 (0.88-3.1)	5.7% (2.4-10%)	6.5% (2.7-12%)	36 (14-67)	520 (210-950)
Gabon	2000	Male	80 to 84	42% (25-61%)	58% (39-75%)	1.4 (0.68-2.4)	4% (1.6-7.4%)	4.6% (1.8-8.4%)	20 (7.8-38)	220 (84-410)
Gabon	2000	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.1 (0.55-1.8)	2.8% (1.1-5.2%)	3.2% (1.3-6%)	7.5 (2.8-14)	56 (22-100)
Gabon	2000	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.92 (0.44-1.6)	1.9% (0.6-3.7%)	2.1% (0.59-4.3%)	1.7 (0.46-3.4)	8.9 (2.9-18)
Gabon	2000	Male	95 plus	39% (21-61%)	61% (39-79%)	0.87 (0.32-1.7)	2% (0.55-4.2%)	2.1% (0.47-4.7%)	0.38 (0.08-0.86)	1.4 (0.38-3)
Gabon	2005	Female	All Ages	22% (18-28%)	78% (72-82%)	1.2 (1-1.4)	3.2% (2.3-4.4%)	5% (3.1-7.4%)	400 (250-610)	12000 (8500-16000)
Gabon	2005	Female	15 to 19	20% (12-32%)	80% (68-88%)	0.88 (0.45-1.5)	1.8% (1-3%)	2.5% (1.1-4.4%)	2.4 (1-4.4)	250 (130-420)
Gabon	2005	Female	20 to 24	29% (18-44%)	71% (56-82%)	1.7 (0.96-2.8)	3.1% (1.9-4.7%)	3.6% (1.6-6.1%)	5.2 (2.6-8.8)	510 (300-770)
Gabon	2005	Female	25 to 29	33% (20-49%)	67% (51-80%)	2.3 (1.3-3.5)	3.4% (2.1-5.1%)	3.7% (1.9-6.2%)	8.4 (4.7-13)	670 (420-950)
Gabon	2005	Female	30 to 34	36% (21-52%)	64% (48-79%)	2.6 (1.5-4)	4% (2.5-6%)	4.6% (2.6-7.3%)	12 (7.3-19)	820 (530-1200)
Gabon	2005	Female	35 to 39	37% (23-55%)	63% (45-77%)	2.6 (1.5-4)	4.6% (2.8-6.7%)	5.4% (3.1-8.3%)	16 (9.2-23)	890 (560-1300)
Gabon	2005	Female	40 to 44	37% (23-55%)	63% (45-77%)	2.5 (1.3-4.1)	5.4% (3.4-7.8%)	6.5% (4-9.6%)	19 (11-27)	940 (570-1300)
Gabon	2005	Female	45 to 49	37% (22-54%)	63% (46-78%)	2.4 (1.3-4)	6.7% (4.1-10%)	8.3% (4.9-12%)	24 (14-37)	1000 (630-1600)
Gabon	2005	Female	50 to 54	34% (20-51%)	66% (49-80%)	2.2 (1.1-3.6)	7.5% (4.4-11%)	9.1% (5.3-14%)	30 (17-47)	1100 (660-1700)
Gabon	2005	Female	55 to 59	31% (18-48%)	69% (52-82%)	2 (1-3.4)	7.8% (4.4-12%)	9.5% (5.3-14%)	36 (20-56)	1200 (650-1800)
Gabon	2005	Female	60 to 64	30% (18-47%)	70% (53-82%)	1.9 (0.92-3.3)	7% (3.6-11%)	8.3% (4.3-13%)	42 (21-67)	1100 (580-1900)
Gabon	2005	Female	65 to 69	32% (21-47%)	68% (53-79%)	1.8 (0.94-3)	6.8% (3.4-11%)	8.1% (4.1-13%)	54 (27-87)	1200 (620-2000)
Gabon	2005	Female	70 to 74	32% (21-47%)	68% (53-79%)	1.6 (0.84-2.9)	5.9% (2.4-11%)	6.9% (2.9-12%)	58 (23-100)	1100 (420-1900)
Gabon	2005	Female	75 to 79	29% (18-45%)	71% (55-82%)	1.4 (0.63-2.5)	4.5% (1.5-8.8%)	5.2% (1.8-10%)	47 (15-94)	660 (210-1300)
Gabon	2005	Female	80 to 84	27% (16-42%)	73% (58-84%)	1.1 (0.55-2)	3.4% (1.2-6.8%)	4% (1.4-7.8%)	32 (11-63)	330 (110-660)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	2005	Female	85 to 89	25% (15-38%)	75% (62-85%)	0.97 (0.48-1.7)	2.6% (0.84-5.2%)	3% (0.95-6%)	12 (4-26)	93 (30-190)
Gabon	2005	Female	90 to 94	27% (16-40%)	73% (60-84%)	0.95 (0.43-1.7)	2% (0.55-4.4%)	2.3% (0.59-5.1%)	3.5 (0.89-7.5)	18 (4.7-39)
Gabon	2005	Female	95 plus	30% (16-47%)	70% (53-84%)	0.96 (0.38-1.8)	1.9% (0.35-4.7%)	2.2% (0.34-5.5%)	0.82 (0.13-2)	2.9 (0.52-6.9)
Gabon	2005	Male	All Ages	31% (25-38%)	69% (62-75%)	1.6 (1.4-1.8)	4.7% (3.5-6%)	6.7% (4.8-8.9%)	500 (340-700)	19000 (14000-25000)
Gabon	2005	Male	15 to 19	24% (14-39%)	76% (61-86%)	1 (0.51-1.9)	3.4% (1.9-5.6%)	4.3% (2-7.6%)	5.4 (2.4-10)	500 (270-840)
Gabon	2005	Male	20 to 24	36% (22-53%)	64% (47-78%)	2.1 (1.2-3.4)	7.9% (4.8-12%)	9.2% (5-14%)	17 (9.1-29)	1400 (850-2200)
Gabon	2005	Male	25 to 29	43% (26-61%)	57% (39-74%)	3.1 (1.8-4.4)	8.4% (5-12%)	9.3% (5.1-14%)	25 (14-38)	1800 (1100-2600)
Gabon	2005	Male	30 to 34	45% (28-63%)	55% (37-72%)	3.4 (2.1-5)	7.8% (5.1-11%)	8.4% (5-13%)	29 (17-43)	1800 (1200-2700)
Gabon	2005	Male	35 to 39	45% (29-63%)	55% (37-71%)	3.3 (1.9-5.1)	7.8% (5.1-11%)	8.6% (5.5-13%)	32 (20-46)	1800 (1200-2500)
Gabon	2005	Male	40 to 44	47% (31-66%)	53% (34-69%)	3.4 (1.9-5.4)	8.5% (5.5-12%)	9.6% (6.1-14%)	35 (22-51)	1700 (1100-2500)
Gabon	2005	Male	45 to 49	50% (32-68%)	50% (32-68%)	3.7 (2-5.8)	10% (6.9-14%)	12% (7.8-17%)	41 (25-60)	1800 (1100-2500)
Gabon	2005	Male	50 to 54	51% (32-69%)	49% (31-68%)	3.6 (2-5.9)	12% (7.6-16%)	14% (9-19%)	47 (29-68)	1800 (1100-2500)
Gabon	2005	Male	55 to 59	49% (31-67%)	51% (33-69%)	3.2 (1.7-5.1)	12% (7.2-16%)	14% (8.7-20%)	50 (30-73)	1600 (1000-2400)
Gabon	2005	Male	60 to 64	47% (29-67%)	53% (33-71%)	2.8 (1.4-4.8)	10% (5.9-15%)	12% (7-18%)	50 (29-77)	1400 (810-2100)
Gabon	2005	Male	65 to 69	48% (31-65%)	52% (35-69%)	2.5 (1.3-4)	8.8% (5-13%)	11% (5.9-16%)	52 (29-82)	1200 (670-1900)
Gabon	2005	Male	70 to 74	49% (31-65%)	51% (35-69%)	2.2 (1.2-3.7)	7.2% (3.7-12%)	8.4% (4.2-14%)	49 (24-82)	890 (440-1500)
Gabon	2005	Male	75 to 79	47% (30-64%)	53% (36-70%)	1.8 (0.9-3.2)	5.5% (2.4-10%)	6.4% (2.8-12%)	36 (15-67)	520 (220-950)
Gabon	2005	Male	80 to 84	42% (25-60%)	58% (40-75%)	1.4 (0.7-2.4)	3.9% (1.6-7%)	4.5% (1.8-8.2%)	20 (8.3-38)	220 (88-400)
Gabon	2005	Male	85 to 89	36% (22-53%)	64% (47-78%)	1.1 (0.55-1.9)	2.8% (1.1-5.3%)	3.2% (1.2-6.1%)	8.2 (3.2-16)	63 (25-120)
Gabon	2005	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.93 (0.43-1.7)	1.9% (0.6-3.9%)	2.1% (0.6-4.5%)	1.9 (0.56-4.1)	10 (3.4-22)
Gabon	2005	Male	95 plus	39% (21-60%)	61% (40-79%)	0.86 (0.34-1.7)	2% (0.57-4.4%)	2.1% (0.5-5%)	0.44 (0.11-1)	1.6 (0.47-3.6)
Gabon	2010	Female	All Ages	22% (18-27%)	78% (73-82%)	1.2 (1-1.5)	3.1% (2.2-4.1%)	4.8% (3-7.1%)	360 (210-530)	11000 (7300-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	2010	Female	15 to 19	20% (12-32%)	80% (68-88%)	0.9 (0.46-1.6)	1.8% (1-2.9%)	2.5% (1.1-4.5%)	2.3 (1-4.3)	250 (140-410)
Gabon	2010	Female	20 to 24	29% (17-43%)	71% (57-83%)	1.8 (0.98-2.8)	3.3% (2.1-4.8%)	4% (2.1-6.5%)	5.2 (2.7-8.7)	530 (320-790)
Gabon	2010	Female	25 to 29	33% (19-50%)	67% (50-81%)	2.4 (1.4-3.6)	3.6% (2.3-5.2%)	4.2% (2.3-6.7%)	8 (4.4-12)	670 (440-960)
Gabon	2010	Female	30 to 34	35% (21-52%)	65% (48-79%)	2.6 (1.5-4.1)	4% (2.6-5.7%)	4.6% (2.7-7.3%)	11 (6.7-18)	780 (510-1100)
Gabon	2010	Female	35 to 39	37% (22-55%)	63% (45-78%)	2.6 (1.5-4.1)	4.2% (2.7-6.2%)	5.1% (3-7.8%)	13 (7.9-21)	800 (500-1200)
Gabon	2010	Female	40 to 44	37% (23-55%)	63% (45-77%)	2.6 (1.3-4.2)	5% (3.2-7.2%)	6.2% (3.9-9%)	17 (10-25)	870 (540-1300)
Gabon	2010	Female	45 to 49	36% (21-53%)	64% (47-79%)	2.5 (1.3-4)	6.1% (3.7-9.3%)	7.8% (4.8-12%)	22 (12-33)	960 (570-1400)
Gabon	2010	Female	50 to 54	33% (19-51%)	67% (49-81%)	2.2 (1.1-3.7)	7% (4.2-10%)	8.8% (5.2-13%)	26 (15-41)	1000 (580-1600)
Gabon	2010	Female	55 to 59	31% (18-48%)	69% (52-82%)	2 (0.99-3.5)	7.3% (4.1-11%)	9.2% (5.1-14%)	32 (17-51)	1000 (560-1600)
Gabon	2010	Female	60 to 64	30% (18-47%)	70% (53-82%)	1.9 (0.92-3.4)	6.7% (3.4-10%)	8.2% (4.2-13%)	35 (17-56)	960 (470-1500)
Gabon	2010	Female	65 to 69	32% (20-46%)	68% (54-80%)	1.9 (0.95-3.1)	6.6% (3.3-11%)	8% (4.1-13%)	45 (22-75)	1000 (500-1700)
Gabon	2010	Female	70 to 74	31% (20-46%)	69% (54-80%)	1.7 (0.86-3)	5.7% (2.3-10%)	6.8% (2.7-12%)	48 (19-86)	880 (340-1600)
Gabon	2010	Female	75 to 79	29% (18-44%)	71% (56-82%)	1.4 (0.65-2.5)	4.3% (1.4-8.6%)	5.1% (1.7-10%)	42 (13-85)	600 (190-1200)
Gabon	2010	Female	80 to 84	26% (15-41%)	74% (59-85%)	1.1 (0.52-2)	3.4% (1.1-6.8%)	3.9% (1.3-7.9%)	32 (10-63)	330 (110-660)
Gabon	2010	Female	85 to 89	25% (15-37%)	75% (63-85%)	0.99 (0.47-1.7)	2.5% (0.78-5.1%)	3% (0.91-5.9%)	13 (4-26)	98 (30-200)
Gabon	2010	Female	90 to 94	26% (16-39%)	74% (61-84%)	0.97 (0.45-1.7)	2% (0.51-4.4%)	2.4% (0.57-5.2%)	3.6 (0.88-8.1)	19 (4.9-41)
Gabon	2010	Female	95 plus	30% (17-48%)	70% (52-83%)	1 (0.39-2)	2% (0.4-4.7%)	2.3% (0.39-5.4%)	0.89 (0.15-2.2)	3.1 (0.61-7.3)
Gabon	2010	Male	All Ages	31% (25-38%)	69% (62-75%)	1.7 (1.4-2)	4.7% (3.5-6%)	6.6% (4.7-8.9%)	490 (330-690)	19000 (14000-25000)
Gabon	2010	Male	15 to 19	23% (13-37%)	77% (63-87%)	1 (0.51-1.8)	3.2% (1.8-5.2%)	4% (2-7%)	5.3 (2.5-9.7)	500 (280-850)
Gabon	2010	Male	20 to 24	35% (21-51%)	65% (49-79%)	2.2 (1.2-3.4)	7.9% (5.1-11%)	9.3% (5.5-14%)	17 (9.2-29)	1400 (880-2200)
Gabon	2010	Male	25 to 29	42% (25-60%)	58% (40-75%)	3.1 (1.9-4.6)	8.8% (5.8-12%)	10% (6-15%)	26 (16-40)	1900 (1200-2700)
Gabon	2010	Male	30 to 34	44% (27-63%)	56% (37-73%)	3.5 (2.1-5.2)	8% (5.2-11%)	8.8% (5.4-13%)	29 (17-44)	1900 (1200-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	2010	Male	35 to 39	45% (28-63%)	55% (37-72%)	3.4 (1.9-5.4)	7.6% (5.1-11%)	8.5% (5.5-12%)	31 (19-47)	1800 (1200-2600)
Gabon	2010	Male	40 to 44	46% (29-65%)	54% (35-71%)	3.5 (1.9-5.6)	8.1% (5.5-11%)	9.3% (6-13%)	36 (22-54)	1800 (1100-2600)
Gabon	2010	Male	45 to 49	50% (32-68%)	50% (32-68%)	3.7 (2-5.8)	9.6% (6.3-13%)	11% (7.1-16%)	41 (25-60)	1800 (1100-2600)
Gabon	2010	Male	50 to 54	50% (32-68%)	50% (32-68%)	3.6 (2-6)	11% (7.4-15%)	13% (8.7-18%)	45 (27-66)	1700 (1100-2500)
Gabon	2010	Male	55 to 59	48% (31-67%)	52% (33-69%)	3.2 (1.7-5.4)	11% (7-15%)	14% (8.6-19%)	47 (29-70)	1600 (970-2300)
Gabon	2010	Male	60 to 64	46% (28-66%)	54% (34-72%)	2.8 (1.4-4.9)	9.9% (6-15%)	12% (7.2-18%)	49 (28-75)	1400 (800-2100)
Gabon	2010	Male	65 to 69	47% (30-64%)	53% (36-70%)	2.5 (1.4-4.1)	8.7% (4.9-13%)	10% (5.8-16%)	49 (26-77)	1100 (610-1800)
Gabon	2010	Male	70 to 74	48% (30-65%)	52% (35-70%)	2.2 (1.2-3.8)	7% (3.4-12%)	8.3% (4-14%)	45 (22-79)	840 (400-1400)
Gabon	2010	Male	75 to 79	46% (30-64%)	54% (36-70%)	1.8 (0.88-3.3)	5.3% (2.2-9.7%)	6.3% (2.6-12%)	36 (14-67)	510 (200-940)
Gabon	2010	Male	80 to 84	41% (25-59%)	59% (41-75%)	1.4 (0.68-2.5)	3.8% (1.5-7.1%)	4.4% (1.7-8.2%)	21 (7.9-40)	220 (84-420)
Gabon	2010	Male	85 to 89	35% (21-52%)	65% (48-79%)	1.1 (0.53-1.9)	2.7% (1.1-5.3%)	3.1% (1.2-6.2%)	9.3 (3.6-19)	71 (28-140)
Gabon	2010	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.92 (0.42-1.7)	1.8% (0.57-3.8%)	2% (0.56-4.3%)	2.4 (0.67-5.1)	13 (4.1-27)
Gabon	2010	Male	95 plus	38% (20-59%)	62% (41-80%)	0.87 (0.33-1.7)	1.9% (0.56-4.3%)	2.1% (0.48-4.9%)	0.56 (0.13-1.3)	2.1 (0.61-4.7)
Gabon	2016	Female	All Ages	22% (18-28%)	78% (72-82%)	1.3 (0.87-2)	3.3% (2.1-4.9%)	5.3% (2.9-8.4%)	350 (180-570)	11000 (6400-15000)
Gabon	2016	Female	15 to 19	20% (12-32%)	80% (68-88%)	0.96 (0.43-1.9)	1.8% (1-3.1%)	2.7% (1.2-5.1%)	2.2 (0.94-4.3)	250 (140-430)
Gabon	2016	Female	20 to 24	29% (17-43%)	71% (57-83%)	1.9 (0.9-3.4)	3.5% (2.2-5.2%)	4.5% (2.2-7.6%)	4.8 (2.2-8.2)	530 (320-790)
Gabon	2016	Female	25 to 29	33% (19-49%)	67% (51-81%)	2.5 (1.3-4.3)	4.2% (2.7-5.9%)	5.5% (3.1-8.5%)	7.5 (4.1-12)	680 (450-970)
Gabon	2016	Female	30 to 34	35% (21-52%)	65% (48-79%)	2.8 (1.4-4.8)	4.8% (3.1-6.7%)	6.5% (3.7-10%)	11 (6-17)	800 (510-1200)
Gabon	2016	Female	35 to 39	36% (22-54%)	64% (46-78%)	2.8 (1.3-4.9)	5% (3.2-7%)	6.9% (4.1-10%)	13 (7-21)	800 (480-1200)
Gabon	2016	Female	40 to 44	36% (23-54%)	64% (46-77%)	2.7 (1.3-5)	5.6% (3.5-8.3%)	7.9% (4.8-12%)	16 (9-26)	850 (490-1300)
Gabon	2016	Female	45 to 49	36% (21-53%)	64% (47-79%)	2.6 (1.2-4.8)	6.6% (3.9-9.9%)	9.3% (5.5-14%)	21 (12-35)	960 (540-1500)
Gabon	2016	Female	50 to 54	33% (20-51%)	67% (49-80%)	2.4 (1-4.4)	7.1% (4-11%)	9.6% (5.5-14%)	26 (14-42)	1000 (550-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	2016	Female	55 to 59	31% (18-48%)	69% (52-82%)	2.2 (0.96-4)	7.3% (3.9-11%)	9.7% (5.1-15%)	30 (16-50)	1000 (520-1600)
Gabon	2016	Female	60 to 64	30% (18-47%)	70% (53-82%)	2.1 (0.89-3.9)	6.7% (3.2-11%)	8.5% (4.1-14%)	33 (15-54)	920 (430-1500)
Gabon	2016	Female	65 to 69	32% (20-47%)	68% (53-80%)	2 (0.92-3.8)	6.7% (3.1-11%)	8.3% (3.9-14%)	41 (18-69)	930 (410-1600)
Gabon	2016	Female	70 to 74	31% (20-47%)	69% (53-80%)	1.8 (0.79-3.6)	5.8% (2.3-11%)	7.1% (2.9-13%)	44 (17-83)	820 (320-1500)
Gabon	2016	Female	75 to 79	29% (18-45%)	71% (55-82%)	1.5 (0.62-3.1)	4.5% (1.3-9.2%)	5.4% (1.5-11%)	40 (11-83)	570 (160-1200)
Gabon	2016	Female	80 to 84	26% (16-41%)	74% (59-84%)	1.2 (0.53-2.5)	3.6% (0.93-7.8%)	4.3% (1.1-9.2%)	33 (8.7-72)	350 (90-770)
Gabon	2016	Female	85 to 89	25% (15-38%)	75% (62-85%)	1.1 (0.48-2.2)	2.8% (0.66-6.5%)	3.3% (0.79-7.7%)	18 (4-40)	130 (31-300)
Gabon	2016	Female	90 to 94	26% (16-39%)	74% (61-84%)	1.1 (0.44-2.1)	2.3% (0.38-5.5%)	2.7% (0.42-6.6%)	5.4 (0.83-13)	28 (4.3-68)
Gabon	2016	Female	95 plus	30% (16-48%)	70% (52-84%)	1.1 (0.38-2.4)	2.3% (0.32-6%)	2.6% (0.31-7.1%)	1.3 (0.15-3.5)	4.4 (0.63-12)
Gabon	2016	Male	All Ages	31% (25-38%)	69% (62-75%)	1.8 (1.2-2.7)	5.2% (3.5-7.2%)	7.3% (4.6-11%)	520 (300-830)	20000 (13000-30000)
Gabon	2016	Male	15 to 19	23% (13-38%)	77% (62-87%)	1.1 (0.48-2.1)	3.2% (1.8-5.5%)	4.1% (1.9-7.8%)	5.5 (2.3-11)	530 (280-940)
Gabon	2016	Male	20 to 24	35% (21-51%)	65% (49-79%)	2.3 (1.1-4.1)	8.1% (5-12%)	9.8% (5.2-16%)	18 (8.7-32)	1500 (880-2500)
Gabon	2016	Male	25 to 29	42% (26-59%)	58% (41-74%)	3.3 (1.7-5.5)	9.6% (6.4-14%)	12% (6.8-17%)	27 (14-44)	2000 (1200-3100)
Gabon	2016	Male	30 to 34	44% (27-63%)	56% (37-73%)	3.7 (1.9-6.1)	9.1% (5.9-13%)	11% (6.4-16%)	31 (17-50)	2100 (1300-3100)
Gabon	2016	Male	35 to 39	45% (28-64%)	55% (36-72%)	3.6 (1.8-6.3)	8.8% (5.7-12%)	10% (6.4-15%)	34 (19-55)	2000 (1200-3100)
Gabon	2016	Male	40 to 44	47% (30-66%)	53% (34-70%)	3.7 (1.9-6.8)	9% (5.7-13%)	11% (6.5-15%)	38 (21-62)	1900 (1100-3000)
Gabon	2016	Male	45 to 49	50% (31-68%)	50% (32-69%)	4 (1.8-7.3)	10% (6.2-14%)	12% (7.6-17%)	46 (24-72)	2000 (1100-3100)
Gabon	2016	Male	50 to 54	50% (32-68%)	50% (32-68%)	3.9 (1.8-6.9)	11% (6.9-15%)	14% (8.4-18%)	49 (27-79)	1900 (1100-3000)
Gabon	2016	Male	55 to 59	48% (30-67%)	52% (33-70%)	3.4 (1.6-6.2)	11% (6.5-16%)	14% (8.2-19%)	49 (27-78)	1600 (930-2600)
Gabon	2016	Male	60 to 64	46% (29-66%)	54% (34-71%)	3 (1.4-5.6)	9.8% (5.5-14%)	12% (6.8-18%)	49 (25-81)	1400 (740-2200)
Gabon	2016	Male	65 to 69	47% (31-65%)	53% (35-69%)	2.7 (1.2-5)	8.7% (4.6-14%)	11% (5.5-17%)	51 (25-91)	1200 (590-2100)
Gabon	2016	Male	70 to 74	48% (30-65%)	52% (35-70%)	2.3 (1.1-4.5)	7.2% (3.2-13%)	8.6% (3.9-15%)	46 (19-88)	860 (370-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Gabon	2016	Male	75 to 79	46% (29-64%)	54% (36-71%)	1.9 (0.85-3.9)	5.5% (2-11%)	6.6% (2.4-13%)	37 (13-76)	530 (190-1100)
Gabon	2016	Male	80 to 84	41% (24-59%)	59% (41-76%)	1.5 (0.65-3)	4% (1.4-8.3%)	4.7% (1.6-9.8%)	23 (7.9-51)	250 (84-540)
Gabon	2016	Male	85 to 89	35% (21-52%)	65% (48-79%)	1.1 (0.51-2.3)	2.9% (0.96-6.4%)	3.3% (1.1-7.3%)	12 (3.8-26)	89 (30-190)
Gabon	2016	Male	90 to 94	34% (21-51%)	66% (49-79%)	0.96 (0.4-1.9)	2% (0.54-4.6%)	2.2% (0.53-5.2%)	3.4 (0.82-8)	19 (5-42)
Gabon	2016	Male	95 plus	38% (20-58%)	62% (42-80%)	0.9 (0.3-1.9)	2% (0.54-5%)	2.2% (0.47-5.7%)	0.86 (0.18-2.2)	3.2 (0.82-7.9)
Burundi	1990	Female	All Ages	21% (17-27%)	79% (73-83%)	2.3 (1.8-2.9)	3.9% (2.8-5.1%)	7.3% (5.4-9.6%)	3500 (2500-4800)	110000 (82000-160000)
Burundi	1990	Female	15 to 19	22% (13-36%)	78% (64-87%)	2.7 (1.3-4.8)	6.2% (3.2-10%)	8.4% (4.1-14%)	58 (28-97)	4400 (2300-7200)
Burundi	1990	Female	20 to 24	27% (15-43%)	73% (57-85%)	3.8 (1.9-6.4)	7.5% (4-11%)	9.1% (4.6-15%)	91 (47-150)	6500 (3600-10000)
Burundi	1990	Female	25 to 29	29% (16-46%)	71% (54-84%)	4.6 (2.5-7.4)	9.4% (5.2-15%)	11% (6-18%)	140 (71-230)	8900 (4800-15000)
Burundi	1990	Female	30 to 34	32% (18-50%)	68% (50-82%)	5 (2.7-8)	10% (6.2-16%)	13% (7.2-20%)	170 (99-290)	10000 (6000-16000)
Burundi	1990	Female	35 to 39	36% (21-54%)	64% (46-79%)	5 (2.6-8.1)	12% (7.3-17%)	14% (8.4-21%)	220 (130-340)	11000 (6900-18000)
Burundi	1990	Female	40 to 44	38% (23-56%)	62% (44-77%)	5 (2.5-8.6)	12% (8.4-18%)	15% (9.6-21%)	190 (120-280)	8800 (5600-13000)
Burundi	1990	Female	45 to 49	38% (21-55%)	62% (45-79%)	5.2 (2.4-9)	14% (8.7-19%)	16% (10-23%)	220 (130-330)	9100 (5600-13000)
Burundi	1990	Female	50 to 54	36% (20-56%)	64% (44-80%)	5.2 (2.5-9.4)	15% (9.6-22%)	18% (11-25%)	300 (180-430)	11000 (6800-16000)
Burundi	1990	Female	55 to 59	35% (20-54%)	65% (46-80%)	5.2 (2.4-9.6)	15% (9.3-22%)	17% (10-25%)	340 (210-510)	11000 (6500-16000)
Burundi	1990	Female	60 to 64	35% (20-54%)	65% (46-80%)	5.2 (2.4-9.6)	14% (8.8-20%)	16% (9.8-22%)	360 (230-520)	9600 (6100-14000)
Burundi	1990	Female	65 to 69	36% (21-54%)	64% (46-79%)	4.8 (2.2-8.6)	14% (9.1-21%)	16% (10-23%)	430 (270-640)	9700 (6100-14000)
Burundi	1990	Female	70 to 74	35% (21-53%)	65% (47-79%)	4.1 (1.8-7.8)	13% (8-19%)	14% (8.7-21%)	410 (250-610)	7400 (4500-11000)
Burundi	1990	Female	75 to 79	32% (18-50%)	68% (50-82%)	3.5 (1.4-6.8)	11% (6.1-17%)	12% (6.7-18%)	310 (170-460)	4300 (2300-6400)
Burundi	1990	Female	80 to 84	31% (16-49%)	69% (51-84%)	3.1 (1.3-6.1)	9.8% (5.1-15%)	11% (5.6-16%)	180 (93-280)	1900 (970-2900)
Burundi	1990	Female	85 to 89	31% (17-48%)	69% (52-83%)	3 (1.2-5.7)	8.3% (4.2-13%)	9.3% (4.6-15%)	65 (33-100)	480 (240-760)
Burundi	1990	Female	90 to 94	33% (19-50%)	67% (50-81%)	3.1 (1.3-5.8)	7.3% (3.6-12%)	8.4% (4-14%)	18 (8.4-29)	90 (43-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	1990	Female	95 plus	38% (19-60%)	62% (40-81%)	3.3 (1.2-6.9)	7.3% (3.2-12%)	8.2% (3.5-14%)	4.3 (1.8-7.4)	15 (6.4-25)
Burundi	1990	Male	All Ages	30% (24-38%)	70% (62-76%)	3.2 (2.6-3.9)	6.2% (4.6-8.1%)	10% (7.8-13%)	5100 (3600-6600)	200000 (140000-260000)
Burundi	1990	Male	15 to 19	28% (17-44%)	72% (56-83%)	3.4 (1.7-5.9)	10% (6.4-16%)	13% (7.6-20%)	110 (65-180)	8600 (5100-13000)
Burundi	1990	Male	20 to 24	39% (24-57%)	61% (43-76%)	4.9 (2.6-8.2)	18% (11-25%)	20% (13-29%)	250 (150-370)	18000 (11000-25000)
Burundi	1990	Male	25 to 29	45% (27-62%)	55% (38-73%)	6.3 (3.6-9.5)	21% (13-29%)	24% (15-34%)	340 (220-490)	22000 (14000-31000)
Burundi	1990	Male	30 to 34	49% (30-67%)	51% (33-70%)	7.3 (4.4-11)	23% (15-31%)	26% (17-36%)	410 (260-570)	24000 (16000-32000)
Burundi	1990	Male	35 to 39	51% (34-69%)	49% (31-66%)	8.1 (4.7-12)	24% (16-33%)	28% (18-38%)	460 (300-630)	24000 (16000-32000)
Burundi	1990	Male	40 to 44	52% (35-71%)	48% (29-65%)	8.4 (4.5-13)	25% (17-32%)	28% (19-37%)	410 (270-550)	19000 (13000-25000)
Burundi	1990	Male	45 to 49	53% (34-71%)	47% (29-66%)	8.4 (4.4-14)	25% (18-32%)	28% (20-37%)	430 (290-610)	18000 (12000-25000)
Burundi	1990	Male	50 to 54	53% (34-72%)	47% (28-66%)	8.3 (4.3-14)	25% (18-33%)	28% (20-37%)	490 (320-690)	18000 (12000-25000)
Burundi	1990	Male	55 to 59	53% (35-71%)	47% (29-65%)	8.1 (4.3-14)	24% (17-31%)	26% (19-34%)	510 (340-720)	16000 (11000-23000)
Burundi	1990	Male	60 to 64	51% (32-70%)	49% (30-68%)	7.1 (3.5-13)	20% (15-26%)	23% (16-29%)	470 (330-660)	13000 (8800-18000)
Burundi	1990	Male	65 to 69	48% (31-67%)	52% (33-69%)	5.7 (3-10)	18% (12-23%)	19% (14-26%)	420 (280-590)	9400 (6300-13000)
Burundi	1990	Male	70 to 74	46% (28-63%)	54% (37-72%)	4.5 (2.2-8.2)	15% (9.4-20%)	16% (10-22%)	350 (220-500)	6300 (3900-8900)
Burundi	1990	Male	75 to 79	43% (27-61%)	57% (39-73%)	3.7 (1.8-6.9)	12% (7-18%)	14% (7.7-20%)	240 (130-350)	3300 (1800-4800)
Burundi	1990	Male	80 to 84	40% (24-60%)	60% (40-76%)	3.1 (1.5-5.7)	10% (5.5-15%)	11% (6.2-17%)	120 (62-180)	1200 (650-1800)
Burundi	1990	Male	85 to 89	38% (22-55%)	62% (45-78%)	2.7 (1.3-4.7)	7.8% (4.2-12%)	8.7% (4.5-13%)	40 (21-60)	300 (160-450)
Burundi	1990	Male	90 to 94	39% (23-57%)	61% (43-77%)	2.5 (1.1-4.6)	5.9% (2.7-9.6%)	6.6% (2.8-11%)	9.7 (4.1-16)	51 (22-84)
Burundi	1990	Male	95 plus	42% (22-64%)	58% (36-78%)	2.4 (0.93-5)	5.6% (2.2-9.7%)	6.1% (2.3-11%)	2.1 (0.79-3.7)	7.5 (3.1-13)
Burundi	1995	Female	All Ages	21% (17-27%)	79% (73-83%)	2.4 (1.9-2.9)	4.1% (2.9-5.5%)	7.7% (5.6-10%)	4000 (2900-5500)	130000 (93000-180000)
Burundi	1995	Female	15 to 19	22% (13-36%)	78% (64-87%)	2.8 (1.4-5)	6.3% (3.3-10%)	8.4% (4.1-14%)	77 (37-130)	5800 (3000-9600)
Burundi	1995	Female	20 to 24	27% (15-43%)	73% (57-85%)	4 (2-6.7)	6.9% (3.8-11%)	8.3% (4.2-14%)	94 (47-160)	6700 (3600-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	1995	Female	25 to 29	30% (16-47%)	70% (53-84%)	4.8 (2.5-8.1)	8.3% (4.6-13%)	9.8% (5.1-16%)	140 (72-220)	8800 (4800-14000)
Burundi	1995	Female	30 to 34	33% (18-51%)	67% (49-82%)	5.3 (2.9-8.6)	9.3% (5.2-15%)	11% (5.8-18%)	180 (97-290)	10000 (5800-17000)
Burundi	1995	Female	35 to 39	36% (21-55%)	64% (45-79%)	5.3 (2.8-8.4)	11% (6.5-17%)	13% (7.6-20%)	250 (150-400)	13000 (7800-21000)
Burundi	1995	Female	40 to 44	38% (22-57%)	62% (43-78%)	5.3 (2.7-8.8)	12% (8-18%)	14% (9.2-21%)	270 (170-410)	13000 (8000-19000)
Burundi	1995	Female	45 to 49	38% (20-56%)	62% (44-80%)	5.5 (2.6-9.6)	13% (8.5-19%)	15% (9.7-23%)	250 (150-380)	10000 (6500-16000)
Burundi	1995	Female	50 to 54	36% (20-56%)	64% (44-80%)	5.4 (2.5-9.7)	15% (9.5-22%)	18% (11-25%)	330 (200-490)	12000 (7300-18000)
Burundi	1995	Female	55 to 59	35% (20-54%)	65% (46-80%)	5.4 (2.6-9.8)	15% (9.4-23%)	18% (11-26%)	380 (230-580)	12000 (7200-18000)
Burundi	1995	Female	60 to 64	35% (20-54%)	65% (46-80%)	5.4 (2.5-10)	14% (9.1-21%)	16% (10-24%)	410 (260-600)	11000 (6900-16000)
Burundi	1995	Female	65 to 69	37% (21-55%)	63% (45-79%)	5.1 (2.4-9)	15% (9.7-22%)	17% (11-24%)	480 (300-700)	11000 (6800-16000)
Burundi	1995	Female	70 to 74	35% (21-53%)	65% (47-79%)	4.4 (1.9-8.4)	14% (8.5-19%)	15% (9.3-21%)	470 (280-690)	8400 (5100-12000)
Burundi	1995	Female	75 to 79	32% (18-51%)	68% (49-82%)	3.7 (1.5-7.3)	12% (6.7-18%)	13% (7.3-19%)	360 (200-530)	5000 (2700-7400)
Burundi	1995	Female	80 to 84	31% (16-49%)	69% (51-84%)	3.3 (1.4-6.4)	10% (5.6-16%)	12% (6.2-17%)	210 (110-310)	2100 (1100-3200)
Burundi	1995	Female	85 to 89	31% (17-49%)	69% (51-83%)	3.2 (1.3-5.9)	8.9% (4.7-14%)	10% (5.2-15%)	86 (45-130)	630 (330-980)
Burundi	1995	Female	90 to 94	33% (19-51%)	67% (49-81%)	3.3 (1.3-6.2)	7.8% (3.9-12%)	9% (4.4-14%)	22 (11-34)	110 (55-170)
Burundi	1995	Female	95 plus	38% (19-60%)	62% (40-81%)	3.5 (1.2-7.4)	7.7% (3.5-13%)	8.8% (4-15%)	5.7 (2.6-9.6)	20 (8.9-33)
Burundi	1995	Male	All Ages	30% (24-37%)	70% (63-76%)	3.2 (2.6-3.9)	6.4% (4.7-8.3%)	11% (7.9-14%)	5700 (4100-7400)	22000 (16000-29000)
Burundi	1995	Male	15 to 19	28% (17-43%)	72% (57-83%)	3.5 (1.8-6)	10% (6.1-15%)	12% (7.1-18%)	150 (87-240)	11000 (6700-18000)
Burundi	1995	Male	20 to 24	39% (24-58%)	61% (42-76%)	5.2 (2.7-8.6)	16% (9.9-22%)	18% (11-25%)	260 (160-380)	18000 (11000-26000)
Burundi	1995	Male	25 to 29	45% (27-63%)	55% (37-73%)	6.6 (3.7-10)	18% (11-26%)	20% (12-29%)	330 (210-470)	21000 (14000-29000)
Burundi	1995	Male	30 to 34	49% (30-67%)	51% (33-70%)	7.7 (4.6-11)	19% (12-26%)	22% (13-30%)	410 (270-570)	24000 (16000-32000)
Burundi	1995	Male	35 to 39	51% (34-69%)	49% (31-66%)	8.4 (4.9-13)	21% (14-29%)	24% (15-33%)	520 (340-710)	27000 (18000-36000)
Burundi	1995	Male	40 to 44	52% (36-70%)	48% (30-64%)	8.7 (4.8-14)	23% (16-30%)	26% (17-34%)	590 (400-800)	27000 (19000-37000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	1995	Male	45 to 49	53% (35-71%)	47% (29-65%)	8.8 (4.8-15)	24% (16-31%)	27% (18-35%)	500 (330-700)	21000 (14000-29000)
Burundi	1995	Male	50 to 54	54% (34-72%)	46% (28-66%)	8.7 (4.6-15)	25% (18-33%)	28% (20-36%)	540 (360-750)	20000 (13000-27000)
Burundi	1995	Male	55 to 59	53% (35-72%)	47% (28-65%)	8.4 (4.4-14)	24% (17-31%)	27% (19-35%)	560 (380-760)	18000 (12000-24000)
Burundi	1995	Male	60 to 64	51% (32-71%)	49% (29-68%)	7.4 (3.7-13)	21% (15-27%)	23% (17-30%)	540 (380-730)	14000 (10000-20000)
Burundi	1995	Male	65 to 69	49% (31-67%)	51% (33-69%)	6 (3.2-10)	18% (13-24%)	20% (14-26%)	480 (320-660)	11000 (7200-15000)
Burundi	1995	Male	70 to 74	46% (29-64%)	54% (36-71%)	4.7 (2.4-8.6)	16% (10-21%)	17% (11-23%)	400 (250-550)	7100 (4500-9900)
Burundi	1995	Male	75 to 79	43% (27-62%)	57% (38-73%)	3.8 (1.8-7.1)	13% (7.7-19%)	14% (8.5-21%)	270 (150-390)	3700 (2100-5300)
Burundi	1995	Male	80 to 84	41% (24-59%)	59% (41-76%)	3.3 (1.6-6)	11% (6.1-16%)	12% (6.7-18%)	130 (71-190)	1400 (740-2000)
Burundi	1995	Male	85 to 89	38% (23-55%)	62% (45-77%)	2.8 (1.4-4.8)	8.5% (4.7-13%)	9.5% (5.1-14%)	52 (28-78)	390 (210-580)
Burundi	1995	Male	90 to 94	38% (23-56%)	62% (44-77%)	2.5 (1.1-4.8)	6.3% (2.9-10%)	7.1% (3.1-12%)	11 (5.1-19)	59 (27-96)
Burundi	1995	Male	95 plus	42% (22-64%)	58% (36-78%)	2.5 (0.93-5.3)	6.1% (2.6-10%)	6.6% (2.8-12%)	3 (1.3-5.1)	11 (4.6-18)
Burundi	2000	Female	All Ages	21% (17-27%)	79% (73-83%)	2.2 (1.7-2.7)	4.3% (3-5.9%)	8.2% (5.8-11%)	4300 (3000-6000)	140000 (96000-190000)
Burundi	2000	Female	15 to 19	22% (12-35%)	78% (65-88%)	2.6 (1.3-4.5)	6.7% (3.7-11%)	9.1% (4.7-15%)	97 (49-170)	7300 (3900-12000)
Burundi	2000	Female	20 to 24	27% (14-42%)	73% (58-86%)	3.7 (1.8-6.1)	7% (3.8-11%)	8.4% (4.2-14%)	110 (53-180)	7800 (4100-12000)
Burundi	2000	Female	25 to 29	29% (17-46%)	71% (54-83%)	4.5 (2.3-7.5)	7.7% (4.1-13%)	9% (4.5-15%)	130 (69-210)	8400 (4600-13000)
Burundi	2000	Female	30 to 34	32% (18-50%)	68% (50-82%)	5 (2.6-8.1)	8.2% (4.7-13%)	9.5% (5.2-16%)	160 (88-260)	9300 (5300-15000)
Burundi	2000	Female	35 to 39	36% (20-55%)	64% (45-80%)	5 (2.6-8.1)	10% (6-16%)	12% (6.7-18%)	230 (130-360)	12000 (7000-18000)
Burundi	2000	Female	40 to 44	37% (22-56%)	63% (44-78%)	5 (2.5-8.5)	12% (7.3-17%)	13% (8.3-20%)	290 (180-440)	13000 (8300-20000)
Burundi	2000	Female	45 to 49	37% (20-55%)	63% (45-80%)	5.1 (2.5-8.8)	13% (8.4-20%)	16% (9.8-24%)	330 (200-500)	13000 (8200-20000)
Burundi	2000	Female	50 to 54	36% (20-56%)	64% (44-80%)	5 (2.4-9.2)	16% (9.8-24%)	18% (11-28%)	340 (200-520)	12000 (7500-19000)
Burundi	2000	Female	55 to 59	34% (19-54%)	66% (46-81%)	5 (2.3-9.4)	16% (9.8-25%)	19% (11-28%)	380 (230-600)	12000 (7100-19000)
Burundi	2000	Female	60 to 64	35% (19-54%)	65% (46-81%)	5 (2.2-9.4)	15% (9.6-22%)	17% (11-25%)	410 (260-620)	11000 (6800-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	2000	Female	65 to 69	36% (21-54%)	64% (46-79%)	4.7 (2.2-8.4)	16% (10-23%)	18% (11-26%)	520 (320-760)	12000 (7200-17000)
Burundi	2000	Female	70 to 74	35% (21-53%)	65% (47-79%)	4 (1.8-7.7)	15% (9.1-21%)	16% (9.9-23%)	510 (310-750)	9100 (5600-13000)
Burundi	2000	Female	75 to 79	32% (18-51%)	68% (49-82%)	3.4 (1.4-6.7)	13% (7.2-19%)	14% (7.9-21%)	420 (240-640)	5900 (3300-8800)
Burundi	2000	Female	80 to 84	31% (16-50%)	69% (50-84%)	3.1 (1.2-5.9)	11% (5.9-17%)	12% (6.5-19%)	250 (140-390)	2600 (1400-4000)
Burundi	2000	Female	85 to 89	31% (17-48%)	69% (52-83%)	2.9 (1.2-5.4)	9.7% (4.9-15%)	11% (5.6-17%)	96 (50-150)	700 (370-1100)
Burundi	2000	Female	90 to 94	33% (19-51%)	67% (49-81%)	3 (1.3-5.7)	8.7% (4.5-14%)	10% (5.1-16%)	30 (15-48)	150 (79-240)
Burundi	2000	Female	95 plus	38% (19-60%)	62% (40-81%)	3.3 (1.2-7)	8.7% (4.1-15%)	9.9% (4.6-17%)	7.3 (3.3-12)	25 (11-41)
Burundi	2000	Male	All Ages	30% (24-37%)	70% (63-76%)	3.1 (2.5-3.7)	6.4% (4.9-8.2%)	11% (8.2-14%)	5700 (4300-7400)	220000 (170000-290000)
Burundi	2000	Male	15 to 19	27% (17-43%)	73% (57-83%)	3.3 (1.8-5.6)	10% (6.5-15%)	12% (7.6-19%)	180 (100-260)	13000 (8100-20000)
Burundi	2000	Male	20 to 24	39% (24-55%)	61% (45-76%)	4.9 (2.6-7.7)	16% (11-22%)	18% (11-25%)	300 (190-420)	21000 (14000-29000)
Burundi	2000	Male	25 to 29	45% (27-63%)	55% (37-73%)	6.3 (3.6-9.6)	17% (11-24%)	19% (12-28%)	300 (200-410)	19000 (13000-26000)
Burundi	2000	Male	30 to 34	49% (30-67%)	51% (33-70%)	7.3 (4.3-11)	17% (11-24%)	19% (13-27%)	350 (230-470)	20000 (14000-27000)
Burundi	2000	Male	35 to 39	51% (34-70%)	49% (30-66%)	8 (4.5-12)	19% (13-25%)	21% (14-29%)	450 (310-600)	23000 (16000-31000)
Burundi	2000	Male	40 to 44	52% (36-71%)	48% (29-64%)	8.3 (4.5-13)	21% (14-27%)	23% (16-31%)	570 (400-770)	27000 (19000-36000)
Burundi	2000	Male	45 to 49	53% (34-72%)	47% (28-66%)	8.3 (4.5-14)	22% (15-29%)	25% (17-33%)	600 (400-830)	25000 (17000-34000)
Burundi	2000	Male	50 to 54	53% (35-71%)	47% (29-65%)	8.2 (4.4-14)	24% (17-32%)	27% (20-36%)	510 (350-710)	19000 (13000-26000)
Burundi	2000	Male	55 to 59	53% (35-71%)	47% (29-65%)	7.9 (4.2-13)	24% (18-31%)	27% (20-36%)	510 (350-690)	16000 (11000-22000)
Burundi	2000	Male	60 to 64	51% (32-70%)	49% (30-68%)	7 (3.5-13)	21% (15-27%)	24% (17-31%)	490 (340-670)	13000 (9300-18000)
Burundi	2000	Male	65 to 69	48% (31-66%)	52% (34-69%)	5.6 (3-9.7)	19% (13-25%)	21% (15-28%)	480 (320-670)	11000 (7300-15000)
Burundi	2000	Male	70 to 74	45% (28-63%)	55% (37-72%)	4.3 (2.2-7.8)	16% (10-22%)	18% (11-24%)	420 (260-590)	7600 (4700-11000)
Burundi	2000	Male	75 to 79	43% (26-62%)	57% (38-74%)	3.6 (1.7-6.6)	14% (7.7-19%)	15% (8.8-22%)	290 (160-420)	4000 (2200-5900)
Burundi	2000	Male	80 to 84	40% (23-59%)	60% (41-77%)	3.1 (1.4-5.5)	12% (6.4-17%)	13% (7.2-19%)	150 (78-220)	1500 (810-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	2000	Male	85 to 89	38% (23-55%)	62% (45-77%)	2.6 (1.2-4.5)	9% (5-14%)	10% (5.5-15%)	51 (28-78)	380 (210-570)
Burundi	2000	Male	90 to 94	38% (23-56%)	62% (44-77%)	2.4 (1.1-4.5)	7% (3.3-11%)	7.9% (3.7-13%)	15 (7.1-25)	78 (37-130)
Burundi	2000	Male	95 plus	42% (21-63%)	58% (37-79%)	2.4 (0.9-4.8)	7.2% (3.3-12%)	8.1% (3.5-14%)	3.7 (1.6-6.3)	13 (5.7-22)
Burundi	2005	Female	All Ages	22% (17-28%)	78% (72-83%)	2.3 (1.8-2.8)	4.2% (3-5.6%)	8.1% (5.8-11%)	4000 (2800-5400)	130000 (91000-170000)
Burundi	2005	Female	15 to 19	22% (12-36%)	78% (64-88%)	2.6 (1.3-4.5)	6.3% (3.4-9.9%)	9.1% (4.7-15%)	92 (47-150)	7100 (3700-11000)
Burundi	2005	Female	20 to 24	27% (15-43%)	73% (57-85%)	3.7 (1.8-6.1)	6.9% (3.9-11%)	8.8% (4.6-15%)	110 (59-180)	8100 (4600-13000)
Burundi	2005	Female	25 to 29	29% (16-46%)	71% (54-84%)	4.5 (2.4-7.4)	7.4% (4.2-12%)	9% (4.7-15%)	130 (70-210)	8700 (4900-14000)
Burundi	2005	Female	30 to 34	32% (17-51%)	68% (49-83%)	4.9 (2.5-8)	7.1% (4.2-11%)	8.3% (4.7-14%)	130 (74-210)	7800 (4700-12000)
Burundi	2005	Female	35 to 39	36% (20-54%)	64% (46-80%)	4.9 (2.5-7.9)	8.7% (5.3-13%)	10% (6-16%)	180 (110-280)	9500 (5700-15000)
Burundi	2005	Female	40 to 44	37% (23-57%)	63% (43-77%)	4.9 (2.5-8.5)	10% (6.4-15%)	12% (7.4-18%)	230 (140-340)	11000 (6700-16000)
Burundi	2005	Female	45 to 49	37% (20-56%)	63% (44-80%)	5.1 (2.4-8.8)	12% (7.8-18%)	15% (9.4-22%)	300 (190-460)	13000 (8000-19000)
Burundi	2005	Female	50 to 54	36% (20-56%)	64% (44-80%)	5 (2.3-9.2)	15% (9.2-22%)	18% (11-26%)	390 (230-570)	14000 (8400-21000)
Burundi	2005	Female	55 to 59	34% (20-54%)	66% (46-80%)	5 (2.3-9.2)	16% (9.7-23%)	18% (11-28%)	340 (210-520)	11000 (6600-16000)
Burundi	2005	Female	60 to 64	35% (20-53%)	65% (47-80%)	5 (2.2-9.5)	15% (9.4-21%)	17% (11-24%)	360 (230-520)	9600 (6200-14000)
Burundi	2005	Female	65 to 69	36% (22-54%)	64% (46-78%)	4.7 (2.2-8.3)	16% (9.9-22%)	18% (11-25%)	450 (280-650)	9900 (6200-14000)
Burundi	2005	Female	70 to 74	35% (21-53%)	65% (47-79%)	4.1 (1.8-7.8)	14% (8.8-21%)	16% (9.8-23%)	460 (280-680)	8300 (5100-12000)
Burundi	2005	Female	75 to 79	32% (18-51%)	68% (49-82%)	3.4 (1.4-6.8)	13% (7.3-19%)	14% (8.1-21%)	390 (220-590)	5400 (3100-8300)
Burundi	2005	Female	80 to 84	31% (16-48%)	69% (52-84%)	3.1 (1.2-5.8)	11% (6.1-17%)	13% (6.9-19%)	270 (150-420)	2800 (1500-4300)
Burundi	2005	Female	85 to 89	31% (17-48%)	69% (52-83%)	3 (1.2-5.6)	9.9% (5.2-16%)	11% (5.8-18%)	110 (59-180)	830 (430-1300)
Burundi	2005	Female	90 to 94	34% (19-51%)	66% (49-81%)	3.1 (1.3-5.8)	9% (4.5-14%)	10% (5.2-17%)	33 (16-52)	160 (82-260)
Burundi	2005	Female	95 plus	38% (19-60%)	62% (40-81%)	3.3 (1.2-6.9)	9.1% (4.4-15%)	10% (5-17%)	9.4 (4.5-16)	32 (15-52)
Burundi	2005	Male	All Ages	30% (24-38%)	70% (62-76%)	3.2 (2.6-3.9)	6.6% (5.1-8.4%)	11% (8.6-14%)	5400 (4200-7000)	220000 (170000-270000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	2005	Male	15 to 19	27% (17-43%)	73% (57-83%)	3.2 (1.7-5.4)	10% (6.5-15%)	13% (8.2-19%)	170 (100-250)	13000 (8100-19000)
Burundi	2005	Male	20 to 24	39% (24-56%)	61% (44-76%)	4.8 (2.6-7.6)	18% (12-24%)	21% (14-29%)	340 (220-470)	24000 (16000-33000)
Burundi	2005	Male	25 to 29	45% (28-62%)	55% (38-72%)	6.1 (3.5-9.3)	19% (13-26%)	23% (15-32%)	360 (230-490)	23000 (15000-31000)
Burundi	2005	Male	30 to 34	49% (30-67%)	51% (33-70%)	7.2 (4.2-11)	18% (12-24%)	20% (14-28%)	330 (220-450)	19000 (14000-26000)
Burundi	2005	Male	35 to 39	51% (34-69%)	49% (31-66%)	7.8 (4.4-12)	18% (13-24%)	21% (14-28%)	390 (280-520)	20000 (15000-27000)
Burundi	2005	Male	40 to 44	52% (35-71%)	48% (29-65%)	8.1 (4.4-13)	19% (14-25%)	22% (16-28%)	480 (340-640)	22000 (16000-30000)
Burundi	2005	Male	45 to 49	53% (34-71%)	47% (29-66%)	8.1 (4.3-13)	20% (15-27%)	24% (17-31%)	570 (400-790)	24000 (17000-32000)
Burundi	2005	Male	50 to 54	53% (34-72%)	47% (28-66%)	8 (4.2-14)	22% (16-29%)	26% (19-34%)	600 (420-820)	22000 (16000-30000)
Burundi	2005	Male	55 to 59	52% (34-72%)	48% (28-66%)	7.7 (4-13)	22% (17-29%)	26% (19-34%)	470 (340-650)	15000 (11000-20000)
Burundi	2005	Male	60 to 64	50% (32-71%)	50% (29-68%)	6.8 (3.3-12)	20% (14-26%)	23% (17-30%)	430 (300-590)	12000 (8200-16000)
Burundi	2005	Male	65 to 69	48% (31-66%)	52% (34-69%)	5.4 (2.9-9.2)	17% (12-23%)	20% (14-27%)	400 (270-560)	9000 (6100-12000)
Burundi	2005	Male	70 to 74	45% (28-62%)	55% (38-72%)	4.2 (2-7.6)	15% (9.3-20%)	17% (10-23%)	380 (240-540)	6800 (4300-9600)
Burundi	2005	Male	75 to 79	42% (26-61%)	58% (39-74%)	3.5 (1.6-6.4)	13% (7.2-18%)	14% (8.1-21%)	290 (160-420)	4000 (2300-5900)
Burundi	2005	Male	80 to 84	40% (23-58%)	60% (42-77%)	3 (1.4-5.4)	11% (6-16%)	13% (6.8-19%)	160 (86-240)	1700 (890-2500)
Burundi	2005	Male	85 to 89	37% (23-56%)	63% (44-77%)	2.5 (1.2-4.5)	8.7% (4.7-13%)	9.9% (5.3-15%)	56 (30-87)	420 (230-640)
Burundi	2005	Male	90 to 94	38% (23-56%)	62% (44-77%)	2.3 (1-4.3)	6.8% (3.1-11%)	7.7% (3.5-13%)	14 (6.5-24)	74 (34-120)
Burundi	2005	Male	95 plus	42% (22-63%)	58% (37-78%)	2.3 (0.85-4.6)	7.3% (3.2-12%)	8.2% (3.5-14%)	4.6 (2-8)	16 (7-27)
Burundi	2010	Female	All Ages	22% (17-28%)	78% (72-83%)	2.3 (1.8-2.8)	4.3% (3.1-5.7%)	8.2% (6-11%)	3900 (2700-5300)	130000 (90000-170000)
Burundi	2010	Female	15 to 19	22% (12-36%)	78% (64-88%)	2.5 (1.2-4.4)	5.4% (3-8.6%)	7.8% (4.1-13%)	77 (41-130)	6200 (3400-9900)
Burundi	2010	Female	20 to 24	27% (15-43%)	73% (57-85%)	3.6 (1.8-5.9)	7% (4.1-11%)	9.4% (4.9-15%)	110 (58-180)	8500 (4800-13000)
Burundi	2010	Female	25 to 29	30% (17-46%)	70% (54-83%)	4.3 (2.3-7)	8.2% (4.6-13%)	11% (5.6-17%)	140 (78-230)	9700 (5600-15000)
Burundi	2010	Female	30 to 34	32% (18-50%)	68% (50-82%)	4.8 (2.6-7.7)	7.9% (4.8-12%)	9.8% (5.6-16%)	140 (81-220)	8600 (5200-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	2010	Female	35 to 39	36% (21-54%)	64% (46-79%)	4.7 (2.4-7.8)	9.2% (5.7-14%)	11% (6.8-17%)	170 (100-270)	9100 (5500-14000)
Burundi	2010	Female	40 to 44	38% (23-56%)	62% (44-77%)	4.8 (2.3-8.1)	10% (6.4-15%)	13% (8-18%)	210 (120-320)	10000 (6100-15000)
Burundi	2010	Female	45 to 49	38% (21-56%)	62% (44-79%)	4.9 (2.4-8.5)	12% (8.1-18%)	15% (9.8-22%)	270 (170-420)	12000 (7200-17000)
Burundi	2010	Female	50 to 54	36% (20-56%)	64% (44-80%)	4.9 (2.3-8.6)	15% (9.1-21%)	18% (11-26%)	390 (240-580)	14000 (8800-21000)
Burundi	2010	Female	55 to 59	35% (20-54%)	65% (46-80%)	4.9 (2.3-9)	15% (9.5-22%)	18% (12-27%)	410 (250-620)	13000 (8000-19000)
Burundi	2010	Female	60 to 64	36% (20-54%)	64% (46-80%)	4.9 (2.2-9.1)	14% (8.9-20%)	16% (10-24%)	340 (210-500)	9100 (5700-13000)
Burundi	2010	Female	65 to 69	36% (21-55%)	64% (45-79%)	4.6 (2.1-8.2)	15% (9.1-21%)	17% (10-24%)	400 (250-590)	9000 (5500-13000)
Burundi	2010	Female	70 to 74	35% (20-52%)	65% (48-80%)	3.9 (1.7-7.4)	13% (8.1-20%)	15% (9.2-22%)	400 (240-600)	7200 (4300-11000)
Burundi	2010	Female	75 to 79	32% (18-51%)	68% (49-82%)	3.3 (1.3-6.7)	12% (6.4-18%)	14% (7.3-21%)	360 (200-550)	5000 (2700-7600)
Burundi	2010	Female	80 to 84	31% (17-49%)	69% (51-83%)	3 (1.2-5.7)	11% (5.6-16%)	12% (6.4-18%)	280 (140-420)	2800 (1500-4300)
Burundi	2010	Female	85 to 89	31% (17-48%)	69% (52-83%)	2.8 (1.2-5.4)	9.4% (4.6-15%)	11% (5.3-17%)	120 (58-190)	860 (430-1400)
Burundi	2010	Female	90 to 94	34% (19-51%)	66% (49-81%)	2.9 (1.2-5.6)	8.6% (4.3-14%)	9.9% (4.9-16%)	36 (18-57)	180 (91-280)
Burundi	2010	Female	95 plus	38% (20-60%)	62% (40-80%)	3.1 (1.1-6.4)	8.6% (4-14%)	9.9% (4.4-17%)	8.7 (4-15)	29 (14-49)
Burundi	2010	Male	All Ages	31% (25-39%)	69% (61-75%)	3.2 (2.5-3.9)	6.9% (5.2-8.6%)	11% (8.7-14%)	5600 (4200-7200)	220000 (170000-290000)
Burundi	2010	Male	15 to 19	28% (16-43%)	72% (57-84%)	3.1 (1.6-5.4)	9.2% (6-13%)	12% (7.2-18%)	150 (86-220)	11000 (7100-17000)
Burundi	2010	Male	20 to 24	39% (24-56%)	61% (44-76%)	4.6 (2.5-7.4)	17% (12-24%)	21% (14-29%)	330 (210-470)	24000 (16000-33000)
Burundi	2010	Male	25 to 29	45% (27-63%)	55% (37-73%)	5.9 (3.5-8.8)	20% (14-27%)	25% (17-33%)	400 (260-540)	26000 (18000-35000)
Burundi	2010	Male	30 to 34	49% (29-67%)	51% (33-71%)	6.9 (4.1-10)	20% (14-26%)	23% (16-32%)	390 (270-530)	23000 (16000-31000)
Burundi	2010	Male	35 to 39	51% (34-69%)	49% (31-66%)	7.6 (4.2-12)	19% (13-25%)	22% (16-30%)	380 (260-500)	20000 (14000-26000)
Burundi	2010	Male	40 to 44	52% (36-71%)	48% (29-64%)	7.9 (4.1-13)	19% (14-25%)	23% (17-30%)	440 (300-600)	21000 (15000-28000)
Burundi	2010	Male	45 to 49	53% (34-71%)	47% (29-66%)	7.8 (4.2-13)	20% (15-26%)	24% (18-31%)	520 (360-710)	22000 (15000-30000)
Burundi	2010	Male	50 to 54	53% (34-72%)	47% (28-66%)	7.7 (4.1-13)	22% (16-28%)	26% (19-34%)	620 (430-860)	23000 (16000-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	2010	Male	55 to 59	52% (34-71%)	48% (29-66%)	7.4 (3.9-12)	22% (16-28%)	26% (19-33%)	610 (430-840)	19000 (14000-27000)
Burundi	2010	Male	60 to 64	51% (32-70%)	49% (30-68%)	6.5 (3.3-12)	19% (13-24%)	22% (16-28%)	440 (310-620)	12000 (8500-17000)
Burundi	2010	Male	65 to 69	48% (30-67%)	52% (33-70%)	5.3 (2.7-8.9)	16% (11-22%)	19% (13-25%)	390 (260-550)	8700 (5800-12000)
Burundi	2010	Male	70 to 74	45% (28-62%)	55% (38-72%)	4.1 (2-7.4)	14% (8.2-19%)	16% (9.4-22%)	340 (200-490)	6100 (3600-8700)
Burundi	2010	Male	75 to 79	43% (26-61%)	57% (39-74%)	3.4 (1.6-6.3)	12% (6.5-17%)	14% (7.4-20%)	280 (160-420)	3900 (2200-5900)
Burundi	2010	Male	80 to 84	40% (24-59%)	60% (41-76%)	2.9 (1.3-5.3)	10% (5.5-15%)	12% (6.2-18%)	180 (94-270)	1900 (1000-2800)
Burundi	2010	Male	85 to 89	38% (22-55%)	62% (45-78%)	2.4 (1.2-4.3)	8.2% (4.4-13%)	9.3% (4.8-15%)	65 (34-100)	480 (250-760)
Burundi	2010	Male	90 to 94	38% (23-56%)	62% (44-77%)	2.2 (0.99-4.2)	6.4% (2.9-11%)	7.3% (3.2-12%)	16 (7.1-27)	83 (38-140)
Burundi	2010	Male	95 plus	42% (22-64%)	58% (36-78%)	2.2 (0.82-4.5)	6.9% (3-12%)	7.8% (3.1-14%)	4.1 (1.6-7.3)	14 (6-25)
Burundi	2016	Female	All Ages	22% (17-28%)	78% (72-83%)	2.2 (1.4-3.2)	4.7% (3.2-6.5%)	8.8% (6.1-12%)	4300 (3000-6000)	140000 (100000-200000)
Burundi	2016	Female	15 to 19	22% (13-36%)	78% (64-87%)	2.5 (1.1-4.7)	5.2% (2.8-8.5%)	7.7% (3.7-13%)	73 (35-130)	5900 (3000-9600)
Burundi	2016	Female	20 to 24	27% (15-43%)	73% (57-85%)	3.5 (1.6-6.3)	7.2% (4.1-11%)	9.9% (5-16%)	110 (55-180)	8600 (4800-13000)
Burundi	2016	Female	25 to 29	30% (16-46%)	70% (54-84%)	4.3 (2.1-7.5)	8.9% (5.3-14%)	12% (6.8-20%)	170 (89-270)	11000 (6800-18000)
Burundi	2016	Female	30 to 34	32% (17-50%)	68% (50-83%)	4.7 (2.3-8.1)	9.5% (6-14%)	13% (7.7-20%)	190 (110-310)	12000 (7100-18000)
Burundi	2016	Female	35 to 39	36% (20-53%)	64% (47-80%)	4.6 (2.2-8.4)	11% (7-17%)	15% (9.1-22%)	220 (130-350)	12000 (7000-18000)
Burundi	2016	Female	40 to 44	38% (23-56%)	62% (44-77%)	4.7 (2.1-8.7)	12% (7.4-17%)	15% (9.7-22%)	220 (130-340)	11000 (6300-16000)
Burundi	2016	Female	45 to 49	38% (20-56%)	62% (44-80%)	4.9 (2-9.1)	13% (8.4-19%)	17% (11-25%)	290 (170-450)	12000 (7400-19000)
Burundi	2016	Female	50 to 54	36% (20-56%)	64% (44-80%)	4.8 (2-9.3)	15% (9.4-22%)	19% (12-27%)	400 (240-590)	15000 (8700-22000)
Burundi	2016	Female	55 to 59	35% (20-54%)	65% (46-80%)	4.8 (2.1-9.4)	15% (9.3-22%)	19% (11-26%)	490 (280-730)	15000 (8900-23000)
Burundi	2016	Female	60 to 64	35% (19-55%)	65% (45-81%)	4.8 (2-8.9)	14% (8.8-20%)	17% (10-24%)	490 (290-720)	13000 (7800-19000)
Burundi	2016	Female	65 to 69	37% (20-54%)	63% (46-80%)	4.5 (1.8-8.8)	14% (9.1-21%)	17% (10-24%)	460 (280-680)	10000 (6300-15000)
Burundi	2016	Female	70 to 74	35% (21-53%)	65% (47-79%)	3.9 (1.6-7.8)	13% (7.8-19%)	15% (8.8-22%)	420 (240-620)	7500 (4500-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	2016	Female	75 to 79	33% (19-51%)	67% (49-81%)	3.3 (1.2-6.9)	12% (6.2-18%)	13% (7.1-20%)	370 (190-580)	5100 (2600-7900)
Burundi	2016	Female	80 to 84	31% (16-49%)	69% (51-84%)	3 (1.1-5.9)	10% (5.3-16%)	12% (6-18%)	290 (150-450)	2900 (1500-4600)
Burundi	2016	Female	85 to 89	31% (18-49%)	69% (51-82%)	2.8 (1.1-5.7)	9.1% (4.5-14%)	10% (5.2-17%)	130 (63-200)	940 (470-1500)
Burundi	2016	Female	90 to 94	34% (19-51%)	66% (49-81%)	2.9 (1.1-5.9)	8.3% (3.9-13%)	9.6% (4.6-16%)	39 (19-63)	200 (95-320)
Burundi	2016	Female	95 plus	38% (19-59%)	62% (41-81%)	3.1 (1.1-6.9)	8.5% (3.8-14%)	9.8% (4.3-16%)	10 (4.5-17)	34 (16-56)
Burundi	2016	Male	All Ages	31% (25-39%)	69% (61-75%)	3.1 (2-4.5)	7.6% (5.4-10%)	12% (8.9-16%)	6500 (4700-8600)	260000 (190000-340000)
Burundi	2016	Male	15 to 19	28% (16-43%)	72% (57-84%)	3.1 (1.4-5.7)	9.1% (5.7-13%)	12% (6.9-18%)	150 (84-230)	12000 (7300-18000)
Burundi	2016	Male	20 to 24	39% (23-57%)	61% (43-77%)	4.5 (2.2-7.8)	17% (12-23%)	21% (13-29%)	350 (220-500)	25000 (16000-36000)
Burundi	2016	Male	25 to 29	44% (27-62%)	56% (38-73%)	5.8 (3.1-9.6)	21% (15-28%)	26% (18-35%)	470 (300-650)	31000 (21000-42000)
Burundi	2016	Male	30 to 34	48% (29-67%)	52% (33-71%)	6.8 (3.6-11)	22% (15-28%)	27% (19-35%)	510 (330-710)	31000 (21000-42000)
Burundi	2016	Male	35 to 39	51% (34-69%)	49% (31-66%)	7.4 (3.8-13)	22% (16-28%)	27% (19-35%)	510 (340-690)	27000 (19000-36000)
Burundi	2016	Male	40 to 44	52% (35-71%)	48% (29-65%)	7.7 (3.6-13)	22% (16-28%)	27% (19-34%)	480 (330-650)	23000 (16000-31000)
Burundi	2016	Male	45 to 49	53% (34-71%)	47% (29-66%)	7.7 (3.5-14)	22% (16-29%)	27% (20-35%)	540 (360-750)	23000 (15000-31000)
Burundi	2016	Male	50 to 54	53% (35-72%)	47% (28-65%)	7.7 (3.5-14)	23% (17-30%)	28% (20-36%)	650 (440-900)	24000 (16000-33000)
Burundi	2016	Male	55 to 59	52% (35-71%)	48% (29-65%)	7.3 (3.5-13)	22% (16-28%)	26% (19-34%)	730 (510-1000)	23000 (16000-32000)
Burundi	2016	Male	60 to 64	50% (32-70%)	50% (30-68%)	6.4 (2.9-12)	19% (13-25%)	22% (15-29%)	660 (460-930)	18000 (13000-25000)
Burundi	2016	Male	65 to 69	48% (30-66%)	52% (34-70%)	5.2 (2.4-9.2)	16% (11-22%)	19% (12-25%)	480 (310-690)	11000 (7000-16000)
Burundi	2016	Male	70 to 74	45% (28-63%)	55% (37-72%)	4 (1.8-7.7)	14% (7.8-19%)	16% (9-22%)	370 (210-540)	6700 (3800-9700)
Burundi	2016	Male	75 to 79	43% (26-61%)	57% (39-74%)	3.3 (1.4-6.5)	12% (5.8-18%)	13% (6.7-20%)	280 (140-440)	4000 (2000-6100)
Burundi	2016	Male	80 to 84	40% (23-59%)	60% (41-77%)	2.8 (1.2-5.3)	10% (4.9-15%)	11% (5.6-17%)	190 (94-300)	2000 (980-3100)
Burundi	2016	Male	85 to 89	37% (22-55%)	63% (45-78%)	2.4 (1-4.6)	8% (3.7-13%)	9.1% (4.2-15%)	79 (37-130)	590 (280-950)
Burundi	2016	Male	90 to 94	38% (23-56%)	62% (44-77%)	2.2 (0.89-4.4)	6.3% (2.7-11%)	7.2% (3-12%)	20 (8.3-35)	110 (44-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burundi	2016	Male	95 plus	42% (21-63%)	58% (37-79%)	2.1 (0.74-4.7)	6.9% (2.7-12%)	7.8% (2.9-14%)	5 (1.9-8.8)	17 (6.7-30)
Comoros	1990	Female	All Ages	23% (17-30%)	77% (70-83%)	0.12 (0.086-0.15)	0.22% (0.042-0.38%)	0.13% (-0.21-0.46%)	2.7 (-4.2-9.2)	270 (53-490)
Comoros	1990	Female	15 to 19	22% (9.8-39%)	78% (61-90%)	0.13 (0.054-0.24)	0.48% (0.15-0.91%)	0.26% (-0.25-0.94%)	0.093 (-0.089-0.33)	21 (6.2-40)
Comoros	1990	Female	20 to 24	27% (12-47%)	73% (53-88%)	0.18 (0.08-0.31)	0.88% (0.34-1.5%)	0.37% (-0.36-1.2%)	0.16 (-0.16-0.52)	40 (15-67)
Comoros	1990	Female	25 to 29	28% (12-47%)	72% (53-88%)	0.21 (0.094-0.37)	1% (0.5-1.5%)	0.56% (-0.14-1.3%)	0.24 (-0.062-0.55)	41 (19-65)
Comoros	1990	Female	30 to 34	31% (12-48%)	69% (52-88%)	0.24 (0.11-0.39)	0.93% (0.39-1.5%)	0.62% (-0.046-1.3%)	0.28 (-0.02-0.59)	37 (15-59)
Comoros	1990	Female	35 to 39	35% (16-52%)	65% (48-84%)	0.24 (0.11-0.41)	1% (0.35-1.7%)	0.97% (-0.04-1.9%)	0.45 (-0.018-0.93)	37 (12-62)
Comoros	1990	Female	40 to 44	36% (20-53%)	64% (47-80%)	0.24 (0.11-0.43)	1.1% (0.43-1.7%)	1.2% (0.3-2.1%)	0.5 (0.12-0.93)	31 (13-50)
Comoros	1990	Female	45 to 49	35% (18-50%)	65% (50-82%)	0.24 (0.11-0.43)	0.92% (0.27-1.6%)	1% (0.12-2%)	0.55 (0.058-1.1)	28 (7.4-48)
Comoros	1990	Female	50 to 54	34% (18-49%)	66% (51-82%)	0.24 (0.11-0.44)	0.78% (0.12-1.4%)	0.9% (0.044-1.7%)	0.62 (0.029-1.3)	26 (3.8-47)
Comoros	1990	Female	55 to 59	34% (19-50%)	66% (50-81%)	0.26 (0.11-0.48)	0.33% (-0.36-0.98%)	0.38% (-0.5-1.2%)	0.32 (-0.41-1)	11 (-12-34)
Comoros	1990	Female	60 to 64	36% (20-54%)	64% (46-80%)	0.27 (0.11-0.48)	0.2% (-0.61-0.88%)	0.25% (-0.73-1.1%)	0.26 (-0.74-1.1)	6.9 (-21-31)
Comoros	1990	Female	65 to 69	38% (20-55%)	62% (45-80%)	0.26 (0.1-0.45)	0.17% (-0.69-0.86%)	0.18% (-0.84-1%)	0.23 (-1-1.3)	5.8 (-23-30)
Comoros	1990	Female	70 to 74	36% (20-53%)	64% (47-80%)	0.22 (0.086-0.42)	-0.19% (-0.99-0.49%)	-0.27% (-1.2-0.51%)	-0.38 (-1.6-0.68)	-5.5 (-28-14)
Comoros	1990	Female	75 to 79	33% (16-50%)	67% (50-84%)	0.19 (0.065-0.4)	-0.07% (-1-0.61%)	-0.096% (-1.2-0.67%)	-0.12 (-1.4-0.87)	-1.5 (-20-13)
Comoros	1990	Female	80 to 84	30% (15-48%)	70% (52-85%)	0.16 (0.058-0.35)	-0.27% (-1.1-0.28%)	-0.31% (-1.2-0.31%)	-0.27 (-1.1-0.27)	-2.7 (-11-2.8)
Comoros	1990	Female	85 to 89	30% (15-46%)	70% (54-85%)	0.15 (0.062-0.32)	-0.32% (-1.1-0.15%)	-0.39% (-1.2-0.16%)	-0.16 (-0.52-0.066)	-1.1 (-3.8-0.53)
Comoros	1990	Female	90 to 94	32% (17-47%)	68% (53-83%)	0.16 (0.064-0.32)	-0.4% (-1.1-0.056%)	-0.51% (-1.3-0.027%)	-0.068 (-0.18-0.0036)	-0.31 (-0.89-0.043)
Comoros	1990	Female	95 plus	35% (17-51%)	65% (49-83%)	0.16 (0.06-0.36)	-0.15% (-0.88-0.4%)	-0.26% (-1.1-0.41%)	-0.0085 (-0.036-0.014)	-0.019 (-0.11-0.053)
Comoros	1990	Male	All Ages	8.2% (6.9-9.9%)	92% (90-93%)	0.017 (0.012-0.024)	0.46% (0.3-0.64%)	0.45% (0.17-0.77%)	9.3 (3.4-16)	650 (410-900)
Comoros	1990	Male	15 to 19	7.4% (4.8-11%)	93% (89-95%)	0.015 (0.0077-0.026)	0.82% (0.51-1.2%)	0.13% (0.038-0.24%)	0.052 (0.015-0.1)	38 (23-57)
Comoros	1990	Male	20 to 24	11% (6.6-15%)	89% (85-93%)	0.024 (0.012-0.042)	1.6% (1-2.4%)	0.24% (0.026-0.51%)	0.13 (0.014-0.29)	81 (49-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	1990	Male	25 to 29	13% (8-19%)	87% (81-92%)	0.032 (0.017-0.052)	1.9% (1.3-2.6%)	0.64% (0.19-1.2%)	0.34 (0.1-0.66)	88 (58-130)
Comoros	1990	Male	30 to 34	14% (8.7-22%)	86% (78-91%)	0.038 (0.02-0.063)	1.8% (1.1-2.5%)	0.86% (0.21-1.6%)	0.48 (0.11-0.92)	78 (47-110)
Comoros	1990	Male	35 to 39	15% (9.3-21%)	85% (79-91%)	0.042 (0.022-0.075)	1.7% (1-2.5%)	1.2% (0.33-2.2%)	0.66 (0.18-1.3)	67 (38-100)
Comoros	1990	Male	40 to 44	15% (9.6-21%)	85% (79-90%)	0.044 (0.021-0.079)	1.7% (0.93-2.6%)	1.5% (0.51-2.8%)	0.77 (0.26-1.5)	55 (29-87)
Comoros	1990	Male	45 to 49	15% (8.9-21%)	85% (79-91%)	0.044 (0.021-0.077)	1.7% (0.88-2.8%)	1.7% (0.6-3.1%)	1 (0.37-2)	58 (28-94)
Comoros	1990	Male	50 to 54	15% (9.9-21%)	85% (79-90%)	0.044 (0.021-0.08)	1.5% (0.74-2.6%)	1.5% (0.52-2.9%)	1.1 (0.39-2.2)	52 (25-89)
Comoros	1990	Male	55 to 59	15% (9.7-21%)	85% (79-90%)	0.043 (0.021-0.079)	1.3% (0.63-2.2%)	1.3% (0.48-2.5%)	1.1 (0.38-2)	43 (20-73)
Comoros	1990	Male	60 to 64	14% (9-21%)	86% (79-91%)	0.037 (0.017-0.073)	1.3% (0.59-2.1%)	1.3% (0.49-2.4%)	1.2 (0.44-2.2)	39 (18-66)
Comoros	1990	Male	65 to 69	13% (8.7-19%)	87% (81-91%)	0.029 (0.014-0.054)	0.77% (0.34-1.3%)	0.81% (0.26-1.5%)	0.81 (0.27-1.5)	21 (9.2-36)
Comoros	1990	Male	70 to 74	12% (8.2-17%)	88% (83-92%)	0.022 (0.01-0.042)	0.58% (0.25-0.98%)	0.63% (0.22-1.1%)	0.65 (0.23-1.2)	13 (5.4-21)
Comoros	1990	Male	75 to 79	12% (7.4-17%)	88% (83-93%)	0.018 (0.0077-0.036)	0.55% (0.23-0.95%)	0.6% (0.19-1.1%)	0.52 (0.16-0.93)	7.8 (3.1-13)
Comoros	1990	Male	80 to 84	10% (6.4-15%)	90% (85-94%)	0.016 (0.0069-0.03)	0.4% (0.2-0.65%)	0.41% (0.16-0.72%)	0.22 (0.086-0.38)	2.6 (1.2-4.1)
Comoros	1990	Male	85 to 89	9.6% (5.7-14%)	90% (86-94%)	0.013 (0.006-0.026)	0.42% (0.2-0.68%)	0.42% (0.15-0.75%)	0.1 (0.036-0.18)	0.89 (0.42-1.5)
Comoros	1990	Male	90 to 94	9.6% (6.1-15%)	90% (85-94%)	0.013 (0.0051-0.026)	0.32% (0.17-0.5%)	0.28% (0.11-0.5%)	0.022 (0.0082-0.039)	0.15 (0.079-0.23)
Comoros	1990	Male	95 plus	11% (6.2-17%)	89% (83-94%)	0.013 (0.0044-0.028)	0.43% (0.21-0.68%)	0.39% (0.12-0.7%)	0.0075 (0.0023-0.014)	0.034 (0.016-0.054)
Comoros	1995	Female	All Ages	23% (16-29%)	77% (71-84%)	0.096 (0.076-0.12)	0.24% (0.06-0.41%)	0.17% (-0.18-0.5%)	3.5 (-3.9-11)	310 (75-540)
Comoros	1995	Female	15 to 19	22% (10-38%)	78% (62-90%)	0.11 (0.049-0.2)	0.47% (0.15-0.88%)	0.22% (-0.29-0.92%)	0.087 (-0.11-0.36)	23 (7-44)
Comoros	1995	Female	20 to 24	27% (12-45%)	73% (55-88%)	0.15 (0.068-0.25)	0.88% (0.3-1.5%)	0.33% (-0.48-1.2%)	0.15 (-0.23-0.55)	43 (14-78)
Comoros	1995	Female	25 to 29	28% (12-46%)	72% (54-88%)	0.18 (0.082-0.3)	0.99% (0.51-1.5%)	0.5% (-0.22-1.2%)	0.23 (-0.1-0.54)	44 (23-69)
Comoros	1995	Female	30 to 34	31% (13-48%)	69% (52-87%)	0.2 (0.088-0.33)	0.92% (0.42-1.4%)	0.57% (-0.17-1.3%)	0.27 (-0.086-0.61)	39 (17-63)
Comoros	1995	Female	35 to 39	35% (16-51%)	65% (49-84%)	0.2 (0.092-0.34)	1% (0.28-1.6%)	0.94% (-0.14-1.9%)	0.5 (-0.074-1)	42 (11-71)
Comoros	1995	Female	40 to 44	36% (20-52%)	64% (48-80%)	0.2 (0.095-0.35)	1.1% (0.43-1.7%)	1.2% (0.23-2.1%)	0.65 (0.13-1.2)	40 (15-66)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	1995	Female	45 to 49	34% (18-48%)	66% (52-82%)	0.2 (0.085-0.34)	0.91% (0.26-1.6%)	1% (0.13-2%)	0.56 (0.067-1.1)	29 (7.7-50)
Comoros	1995	Female	50 to 54	33% (18-49%)	67% (51-82%)	0.2 (0.091-0.35)	0.76% (0.11-1.4%)	0.88% (0.035-1.7%)	0.64 (0.026-1.3)	27 (3.8-50)
Comoros	1995	Female	55 to 59	34% (19-50%)	66% (50-81%)	0.21 (0.096-0.38)	0.31% (-0.36-0.94%)	0.35% (-0.49-1.2%)	0.32 (-0.45-1)	11 (-13-34)
Comoros	1995	Female	60 to 64	36% (20-53%)	64% (47-80%)	0.22 (0.093-0.4)	0.22% (-0.53-0.87%)	0.27% (-0.61-1%)	0.31 (-0.65-1.2)	8.4 (-19-33)
Comoros	1995	Female	65 to 69	37% (20-54%)	63% (46-80%)	0.21 (0.087-0.37)	0.24% (-0.56-0.93%)	0.26% (-0.67-1.1%)	0.35 (-0.91-1.5)	8.9 (-20-35)
Comoros	1995	Female	70 to 74	36% (19-52%)	64% (48-81%)	0.18 (0.074-0.33)	-0.095% (-0.88-0.53%)	-0.17% (-1.1-0.55%)	-0.26 (-1.6-0.81)	-3 (-27-16)
Comoros	1995	Female	75 to 79	32% (16-48%)	68% (52-84%)	0.15 (0.054-0.32)	0.059% (-0.8-0.69%)	0.053% (-0.91-0.77%)	0.075 (-1.3-1.1)	1.4 (-18-16)
Comoros	1995	Female	80 to 84	29% (15-46%)	71% (54-85%)	0.13 (0.048-0.28)	-0.16% (-0.9-0.34%)	-0.19% (-1-0.38%)	-0.19 (-1.1-0.38)	-2 (-11-4.1)
Comoros	1995	Female	85 to 89	29% (15-46%)	71% (54-85%)	0.13 (0.054-0.26)	-0.22% (-0.91-0.18%)	-0.26% (-1-0.2%)	-0.14 (-0.55-0.1)	-0.96 (-4.1-0.81)
Comoros	1995	Female	90 to 94	31% (16-46%)	69% (54-84%)	0.13 (0.05-0.25)	-0.29% (-0.9-0.1%)	-0.38% (-1.1-0.085%)	-0.063 (-0.18-0.015)	-0.29 (-0.87-0.11)
Comoros	1995	Female	95 plus	34% (18-51%)	66% (49-82%)	0.13 (0.049-0.27)	-0.023% (-0.65-0.48%)	-0.1% (-0.86-0.52%)	-0.004 (-0.034-0.02)	-0.0034 (-0.1-0.076)
Comoros	1995	Male	All Ages	8.3% (6.9-9.9%)	92% (90-93%)	0.014 (0.01-0.02)	0.52% (0.34-0.71%)	0.51% (0.2-0.86%)	11 (4.3-18)	750 (470-1000)
Comoros	1995	Male	15 to 19	7.3% (4.7-11%)	93% (89-95%)	0.013 (0.0065-0.022)	0.85% (0.54-1.2%)	0.12% (0.04-0.22%)	0.057 (0.018-0.11)	46 (28-70)
Comoros	1995	Male	20 to 24	11% (6.7-16%)	89% (84-93%)	0.02 (0.01-0.033)	1.7% (1.1-2.5%)	0.24% (0.045-0.46%)	0.14 (0.027-0.3)	93 (57-140)
Comoros	1995	Male	25 to 29	13% (7.9-19%)	87% (81-92%)	0.026 (0.014-0.043)	1.9% (1.4-2.6%)	0.64% (0.22-1.2%)	0.38 (0.13-0.71)	99 (67-140)
Comoros	1995	Male	30 to 34	14% (8.8-21%)	86% (79-91%)	0.031 (0.017-0.051)	1.8% (1.2-2.5%)	0.88% (0.28-1.6%)	0.54 (0.17-1)	89 (57-130)
Comoros	1995	Male	35 to 39	15% (9.2-21%)	85% (79-91%)	0.035 (0.018-0.06)	1.7% (1.1-2.5%)	1.2% (0.41-2.3%)	0.8 (0.26-1.5)	80 (47-120)
Comoros	1995	Male	40 to 44	15% (9.5-21%)	85% (79-90%)	0.036 (0.018-0.061)	1.7% (0.94-2.6%)	1.5% (0.52-2.8%)	1 (0.32-1.9)	71 (38-110)
Comoros	1995	Male	45 to 49	15% (9-21%)	85% (79-91%)	0.036 (0.017-0.064)	1.8% (0.91-2.8%)	1.7% (0.62-3.1%)	1.1 (0.39-2.1)	62 (32-100)
Comoros	1995	Male	50 to 54	15% (9.8-21%)	85% (79-90%)	0.037 (0.017-0.067)	1.6% (0.77-2.6%)	1.6% (0.56-3%)	1.2 (0.45-2.3)	57 (28-94)
Comoros	1995	Male	55 to 59	15% (9.6-21%)	85% (79-90%)	0.036 (0.018-0.064)	1.4% (0.68-2.2%)	1.4% (0.53-2.6%)	1.3 (0.47-2.3)	49 (24-81)
Comoros	1995	Male	60 to 64	14% (9.1-21%)	86% (79-91%)	0.031 (0.014-0.059)	1.3% (0.62-2.1%)	1.4% (0.53-2.4%)	1.4 (0.52-2.5)	44 (20-73)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	1995	Male	65 to 69	13% (8.5-19%)	87% (81-92%)	0.025 (0.012-0.043)	0.79% (0.36-1.4%)	0.83% (0.29-1.6%)	0.91 (0.31-1.7)	24 (11-42)
Comoros	1995	Male	70 to 74	12% (8.1-17%)	88% (83-92%)	0.019 (0.0089-0.034)	0.6% (0.27-1%)	0.65% (0.25-1.2%)	0.73 (0.27-1.3)	14 (6.4-24)
Comoros	1995	Male	75 to 79	11% (7.4-17%)	89% (83-93%)	0.015 (0.0068-0.029)	0.58% (0.25-0.99%)	0.64% (0.23-1.1%)	0.62 (0.22-1.1)	9.3 (4-16)
Comoros	1995	Male	80 to 84	11% (6.6-15%)	89% (85-93%)	0.013 (0.0058-0.024)	0.42% (0.21-0.68%)	0.43% (0.18-0.75%)	0.27 (0.11-0.46)	3.2 (1.6-5.1)
Comoros	1995	Male	85 to 89	9.6% (5.7-14%)	90% (86-94%)	0.011 (0.0051-0.021)	0.43% (0.21-0.7%)	0.44% (0.17-0.77%)	0.13 (0.052-0.23)	1.1 (0.54-1.8)
Comoros	1995	Male	90 to 94	9.5% (5.9-15%)	91% (85-94%)	0.01 (0.0044-0.019)	0.33% (0.18-0.5%)	0.3% (0.13-0.5%)	0.029 (0.013-0.049)	0.19 (0.11-0.29)
Comoros	1995	Male	95 plus	10% (6-16%)	90% (84-94%)	0.01 (0.0038-0.021)	0.45% (0.24-0.7%)	0.41% (0.14-0.72%)	0.0087 (0.003-0.015)	0.039 (0.02-0.061)
Comoros	2000	Female	All Ages	22% (15-29%)	78% (71-85%)	0.11 (0.084-0.14)	0.26% (0.049-0.44%)	0.18% (-0.24-0.56%)	3.9 (-5.3-12)	360 (65-640)
Comoros	2000	Female	15 to 19	22% (10-39%)	78% (61-90%)	0.12 (0.053-0.22)	0.49% (0.1-0.96%)	0.24% (-0.4-1.1%)	0.11 (-0.17-0.47)	28 (6.1-55)
Comoros	2000	Female	20 to 24	26% (11-46%)	74% (54-89%)	0.17 (0.077-0.3)	0.93% (0.25-1.6%)	0.37% (-0.58-1.3%)	0.19 (-0.29-0.68)	53 (13-93)
Comoros	2000	Female	25 to 29	28% (12-45%)	72% (55-88%)	0.2 (0.092-0.34)	1.1% (0.48-1.7%)	0.56% (-0.29-1.4%)	0.27 (-0.14-0.65)	52 (23-83)
Comoros	2000	Female	30 to 34	30% (12-47%)	70% (53-88%)	0.22 (0.096-0.36)	0.96% (0.4-1.5%)	0.61% (-0.26-1.4%)	0.31 (-0.13-0.72)	45 (18-70)
Comoros	2000	Female	35 to 39	34% (15-51%)	66% (49-85%)	0.22 (0.099-0.37)	1.1% (0.22-1.8%)	1% (-0.29-2.2%)	0.57 (-0.15-1.2)	48 (9.1-82)
Comoros	2000	Female	40 to 44	33% (18-50%)	67% (50-82%)	0.21 (0.1-0.38)	1.1% (0.35-1.7%)	1.2% (0.13-2.2%)	0.73 (0.081-1.4)	46 (15-75)
Comoros	2000	Female	45 to 49	29% (14-44%)	71% (56-86%)	0.19 (0.077-0.34)	0.92% (0.23-1.6%)	1% (0.04-2%)	0.71 (0.03-1.4)	37 (9.4-65)
Comoros	2000	Female	50 to 54	26% (14-41%)	74% (59-86%)	0.18 (0.072-0.33)	0.8% (0.15-1.5%)	0.91% (0.059-1.9%)	0.66 (0.036-1.4)	28 (5-52)
Comoros	2000	Female	55 to 59	29% (15-46%)	71% (54-85%)	0.21 (0.087-0.41)	0.38% (-0.37-1%)	0.43% (-0.51-1.3%)	0.4 (-0.47-1.2)	14 (-13-39)
Comoros	2000	Female	60 to 64	34% (18-51%)	66% (49-82%)	0.24 (0.095-0.44)	0.26% (-0.55-1%)	0.32% (-0.66-1.2%)	0.38 (-0.8-1.5)	10 (-22-39)
Comoros	2000	Female	65 to 69	36% (19-54%)	64% (46-81%)	0.24 (0.093-0.42)	0.24% (-0.61-0.98%)	0.26% (-0.74-1.2%)	0.37 (-1.1-1.7)	9 (-23-39)
Comoros	2000	Female	70 to 74	35% (19-52%)	65% (48-81%)	0.21 (0.082-0.39)	-0.11% (-0.91-0.56%)	-0.18% (-1.1-0.59%)	-0.29 (-1.8-0.94)	-3.8 (-31-18)
Comoros	2000	Female	75 to 79	32% (16-49%)	68% (51-84%)	0.17 (0.062-0.36)	0.028% (-0.85-0.69%)	0.022% (-0.98-0.78%)	0.034 (-1.5-1.2)	0.68 (-21-17)
Comoros	2000	Female	80 to 84	29% (14-47%)	71% (53-86%)	0.15 (0.055-0.32)	-0.19% (-0.95-0.35%)	-0.22% (-1.1-0.4%)	-0.25 (-1.3-0.46)	-2.6 (-13-4.7)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	2000	Female	85 to 89	28% (15-46%)	72% (54-85%)	0.14 (0.06-0.3)	-0.24% (-0.98-0.2%)	-0.29% (-1.1-0.21%)	-0.18 (-0.71-0.13)	-1.3 (-5.2-1)
Comoros	2000	Female	90 to 94	30% (17-47%)	70% (53-83%)	0.15 (0.06-0.3)	-0.32% (-0.96-0.11%)	-0.42% (-1.2-0.086%)	-0.086 (-0.24-0.018)	-0.39 (-1.2-0.13)
Comoros	2000	Female	95 plus	34% (17-52%)	66% (48-83%)	0.15 (0.054-0.34)	-0.037% (-0.73-0.52%)	-0.13% (-0.98-0.54%)	-0.0062 (-0.048-0.026)	-0.007 (-0.14-0.097)
Comoros	2000	Male	All Ages	7.4% (6.1-8.9%)	93% (91-94%)	0.015 (0.01-0.02)	0.55% (0.36-0.73%)	0.5% (0.18-0.83%)	11 (4.1-18)	820 (540-1100)
Comoros	2000	Male	15 to 19	6.9% (4.5-9.9%)	93% (90-95%)	0.014 (0.0069-0.023)	0.91% (0.58-1.3%)	0.12% (0.018-0.24%)	0.059 (0.0096-0.12)	55 (33-85)
Comoros	2000	Male	20 to 24	9.2% (5.8-13%)	91% (87-94%)	0.02 (0.01-0.031)	1.8% (1.1-2.7%)	0.23% (0.0073-0.48%)	0.14 (0.0042-0.29)	110 (67-170)
Comoros	2000	Male	25 to 29	11% (6.7-16%)	89% (84-93%)	0.025 (0.014-0.042)	2.1% (1.4-2.8%)	0.62% (0.16-1.2%)	0.37 (0.093-0.68)	110 (73-160)
Comoros	2000	Male	30 to 34	12% (7.4-18%)	88% (82-93%)	0.031 (0.017-0.051)	1.9% (1.3-2.6%)	0.85% (0.24-1.6%)	0.51 (0.15-0.99)	95 (62-140)
Comoros	2000	Male	35 to 39	13% (7.6-18%)	87% (82-92%)	0.034 (0.017-0.059)	1.7% (1.1-2.5%)	1.2% (0.38-2.2%)	0.78 (0.24-1.5)	86 (52-130)
Comoros	2000	Male	40 to 44	12% (7.9-18%)	88% (82-92%)	0.035 (0.017-0.06)	1.7% (0.95-2.5%)	1.5% (0.49-2.7%)	1 (0.33-1.9)	78 (43-120)
Comoros	2000	Male	45 to 49	13% (7.8-18%)	87% (82-92%)	0.036 (0.016-0.064)	1.7% (0.93-2.7%)	1.7% (0.6-3.1%)	1.3 (0.45-2.4)	74 (39-120)
Comoros	2000	Male	50 to 54	13% (8.9-19%)	87% (81-91%)	0.037 (0.018-0.069)	1.5% (0.76-2.6%)	1.6% (0.55-2.9%)	1.2 (0.4-2.1)	56 (27-90)
Comoros	2000	Male	55 to 59	14% (9.2-20%)	86% (80-91%)	0.038 (0.018-0.068)	1.3% (0.68-2.2%)	1.4% (0.48-2.5%)	1.2 (0.45-2.2)	49 (24-82)
Comoros	2000	Male	60 to 64	13% (8.5-20%)	87% (80-92%)	0.034 (0.016-0.065)	1.3% (0.59-2.1%)	1.4% (0.5-2.4%)	1.5 (0.52-2.6)	47 (22-77)
Comoros	2000	Male	65 to 69	12% (8.1-17%)	88% (83-92%)	0.026 (0.013-0.046)	0.78% (0.34-1.3%)	0.82% (0.26-1.5%)	0.95 (0.31-1.8)	25 (11-42)
Comoros	2000	Male	70 to 74	12% (7.9-16%)	88% (84-92%)	0.02 (0.0096-0.036)	0.59% (0.25-0.99%)	0.64% (0.23-1.1%)	0.77 (0.28-1.4)	15 (6.2-25)
Comoros	2000	Male	75 to 79	11% (7-16%)	89% (84-93%)	0.017 (0.0071-0.032)	0.58% (0.24-1%)	0.63% (0.22-1.2%)	0.66 (0.24-1.2)	9.9 (4.2-18)
Comoros	2000	Male	80 to 84	10% (6.2-14%)	90% (86-94%)	0.014 (0.0064-0.026)	0.41% (0.2-0.66%)	0.42% (0.17-0.73%)	0.3 (0.13-0.51)	3.6 (1.7-5.7)
Comoros	2000	Male	85 to 89	9.1% (5.4-14%)	91% (86-95%)	0.012 (0.0056-0.022)	0.43% (0.2-0.69%)	0.43% (0.15-0.74%)	0.15 (0.054-0.26)	1.3 (0.61-2.1)
Comoros	2000	Male	90 to 94	9.1% (5.7-13%)	91% (87-94%)	0.011 (0.0048-0.022)	0.33% (0.18-0.5%)	0.3% (0.12-0.51%)	0.034 (0.014-0.059)	0.23 (0.13-0.35)
Comoros	2000	Male	95 plus	9.9% (5.9-15%)	90% (85-94%)	0.011 (0.004-0.023)	0.45% (0.23-0.73%)	0.4% (0.13-0.74%)	0.01 (0.0034-0.019)	0.046 (0.024-0.075)
Comoros	2005	Female	All Ages	22% (16-29%)	78% (71-84%)	0.13 (0.1-0.17)	0.32% (0.092-0.53%)	0.23% (-0.23-0.65%)	4.8 (-4.8-14)	440 (120-730)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	2005	Female	15 to 19	22% (10-38%)	78% (62-90%)	0.15 (0.063-0.27)	0.58% (0.18-1.1%)	0.38% (-0.3-1.2%)	0.17 (-0.14-0.52)	35 (11-65)
Comoros	2005	Female	20 to 24	26% (12-45%)	74% (55-88%)	0.2 (0.094-0.35)	1.1% (0.46-1.8%)	0.54% (-0.41-1.6%)	0.28 (-0.21-0.84)	67 (27-110)
Comoros	2005	Female	25 to 29	27% (12-45%)	73% (55-88%)	0.24 (0.11-0.4)	1.2% (0.66-1.8%)	0.75% (-0.029-1.6%)	0.37 (-0.014-0.8)	65 (35-99)
Comoros	2005	Female	30 to 34	30% (12-47%)	70% (53-88%)	0.26 (0.12-0.44)	1.1% (0.57-1.7%)	0.82% (-0.075-1.7%)	0.4 (-0.037-0.82)	54 (26-82)
Comoros	2005	Female	35 to 39	34% (14-50%)	66% (50-86%)	0.26 (0.12-0.44)	1.2% (0.41-1.9%)	1.2% (-0.0036-2.4%)	0.63 (-0.0021-1.3)	54 (19-87)
Comoros	2005	Female	40 to 44	33% (18-50%)	67% (50-82%)	0.25 (0.11-0.45)	1.1% (0.45-1.8%)	1.3% (0.2-2.3%)	0.76 (0.12-1.4)	50 (19-78)
Comoros	2005	Female	45 to 49	28% (13-43%)	72% (57-87%)	0.23 (0.091-0.43)	0.97% (0.28-1.6%)	1.1% (0.13-2.2%)	0.78 (0.093-1.5)	42 (12-72)
Comoros	2005	Female	50 to 54	25% (13-41%)	75% (59-87%)	0.21 (0.083-0.4)	0.86% (0.18-1.5%)	1% (0.11-2%)	0.83 (0.092-1.6)	35 (6.8-64)
Comoros	2005	Female	55 to 59	29% (15-45%)	71% (55-85%)	0.25 (0.1-0.5)	0.46% (-0.33-1.2%)	0.55% (-0.49-1.5%)	0.45 (-0.41-1.3)	16 (-11-40)
Comoros	2005	Female	60 to 64	34% (18-52%)	66% (48-82%)	0.29 (0.11-0.54)	0.33% (-0.45-1.1%)	0.42% (-0.54-1.4%)	0.46 (-0.6-1.6)	12 (-17-41)
Comoros	2005	Female	65 to 69	37% (20-54%)	63% (46-80%)	0.29 (0.11-0.51)	0.29% (-0.64-1.1%)	0.33% (-0.77-1.3%)	0.44 (-1-1.8)	10 (-24-40)
Comoros	2005	Female	70 to 74	35% (19-52%)	65% (48-81%)	0.25 (0.097-0.47)	-0.082% (-0.95-0.63%)	-0.15% (-1.1-0.69%)	-0.23 (-1.8-1.1)	-2.7 (-31-21)
Comoros	2005	Female	75 to 79	32% (16-49%)	68% (51-84%)	0.21 (0.076-0.44)	0.035% (-0.92-0.7%)	0.036% (-1.1-0.81%)	0.052 (-1.6-1.2)	0.86 (-22-17)
Comoros	2005	Female	80 to 84	29% (14-47%)	71% (53-86%)	0.18 (0.066-0.39)	-0.2% (-1-0.35%)	-0.24% (-1.2-0.42%)	-0.27 (-1.3-0.47)	-2.8 (-14-4.8)
Comoros	2005	Female	85 to 89	29% (14-46%)	71% (54-86%)	0.17 (0.068-0.35)	-0.27% (-1.1-0.22%)	-0.33% (-1.3-0.24%)	-0.22 (-0.85-0.16)	-1.6 (-6.2-1.3)
Comoros	2005	Female	90 to 94	31% (16-46%)	69% (54-84%)	0.18 (0.069-0.35)	-0.37% (-1.1-0.11%)	-0.48% (-1.3-0.083%)	-0.12 (-0.32-0.019)	-0.53 (-1.5-0.15)
Comoros	2005	Female	95 plus	34% (18-52%)	66% (48-82%)	0.19 (0.067-0.4)	-0.067% (-0.82-0.53%)	-0.16% (-1.1-0.55%)	-0.0094 (-0.062-0.031)	-0.015 (-0.19-0.12)
Comoros	2005	Male	All Ages	7.2% (6-8.6%)	93% (91-94%)	0.017 (0.012-0.024)	0.63% (0.42-0.85%)	0.54% (0.22-0.91%)	12 (4.7-19)	920 (610-1200)
Comoros	2005	Male	15 to 19	6.5% (4.2-9.5%)	93% (91-96%)	0.016 (0.0079-0.027)	0.99% (0.64-1.4%)	0.14% (0.035-0.27%)	0.069 (0.017-0.14)	62 (38-93)
Comoros	2005	Male	20 to 24	8.7% (5.5-12%)	91% (88-95%)	0.022 (0.012-0.037)	2% (1.3-2.9%)	0.25% (0.048-0.5%)	0.16 (0.031-0.34)	130 (80-200)
Comoros	2005	Male	25 to 29	10% (6.7-15%)	90% (85-93%)	0.029 (0.016-0.047)	2.2% (1.6-3%)	0.65% (0.22-1.2%)	0.39 (0.13-0.73)	130 (85-190)
Comoros	2005	Male	30 to 34	11% (7-17%)	89% (83-93%)	0.035 (0.02-0.057)	2% (1.4-2.8%)	0.88% (0.26-1.6%)	0.52 (0.16-0.97)	100 (68-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	2005	Male	35 to 39	12% (7.2-17%)	88% (83-93%)	0.039 (0.02-0.069)	1.8% (1.2-2.6%)	1.2% (0.4-2.3%)	0.78 (0.24-1.5)	91 (56-130)
Comoros	2005	Male	40 to 44	12% (7.5-17%)	88% (83-92%)	0.04 (0.018-0.07)	1.8% (1.1-2.7%)	1.5% (0.56-2.9%)	1.1 (0.36-2.1)	84 (49-130)
Comoros	2005	Male	45 to 49	12% (7.5-17%)	88% (83-92%)	0.041 (0.019-0.076)	1.8% (0.96-2.8%)	1.8% (0.64-3.2%)	1.4 (0.5-2.6)	83 (45-130)
Comoros	2005	Male	50 to 54	13% (8.7-18%)	87% (82-91%)	0.043 (0.02-0.079)	1.6% (0.8-2.5%)	1.6% (0.61-3%)	1.4 (0.52-2.6)	69 (34-110)
Comoros	2005	Male	55 to 59	13% (8.5-19%)	87% (81-91%)	0.044 (0.021-0.082)	1.4% (0.69-2.3%)	1.4% (0.5-2.6%)	1.2 (0.43-2.2)	49 (24-81)
Comoros	2005	Male	60 to 64	13% (8.2-19%)	87% (81-92%)	0.039 (0.017-0.074)	1.3% (0.65-2.2%)	1.4% (0.52-2.6%)	1.5 (0.53-2.7)	48 (24-80)
Comoros	2005	Male	65 to 69	12% (7.7-16%)	88% (84-92%)	0.03 (0.014-0.053)	0.82% (0.37-1.4%)	0.87% (0.28-1.6%)	1 (0.34-1.9)	27 (12-46)
Comoros	2005	Male	70 to 74	11% (7.4-16%)	89% (84-93%)	0.023 (0.01-0.041)	0.62% (0.27-1.1%)	0.68% (0.24-1.2%)	0.83 (0.29-1.5)	16 (7-28)
Comoros	2005	Male	75 to 79	11% (6.7-16%)	89% (84-93%)	0.019 (0.0081-0.036)	0.6% (0.26-1%)	0.67% (0.24-1.2%)	0.71 (0.26-1.3)	11 (4.5-19)
Comoros	2005	Male	80 to 84	9.5% (5.9-14%)	90% (86-94%)	0.016 (0.0073-0.029)	0.43% (0.22-0.69%)	0.45% (0.18-0.77%)	0.34 (0.14-0.57)	4 (2-6.3)
Comoros	2005	Male	85 to 89	8.7% (5.2-13%)	91% (87-95%)	0.014 (0.0064-0.026)	0.45% (0.22-0.73%)	0.45% (0.16-0.79%)	0.19 (0.067-0.33)	1.6 (0.78-2.6)
Comoros	2005	Male	90 to 94	8.6% (5.4-13%)	91% (87-95%)	0.013 (0.0054-0.025)	0.34% (0.19-0.52%)	0.31% (0.13-0.53%)	0.043 (0.018-0.074)	0.28 (0.16-0.44)
Comoros	2005	Male	95 plus	9.4% (5.6-15%)	91% (85-94%)	0.013 (0.0046-0.026)	0.46% (0.24-0.74%)	0.42% (0.14-0.76%)	0.012 (0.0041-0.022)	0.055 (0.027-0.089)
Comoros	2010	Female	All Ages	23% (17-29%)	77% (71-83%)	0.14 (0.1-0.2)	0.38% (0.13-0.61%)	0.28% (-0.21-0.74%)	5.8 (-4.5-16)	500 (170-830)
Comoros	2010	Female	15 to 19	21% (10-38%)	79% (62-90%)	0.15 (0.062-0.28)	0.6% (0.25-1%)	0.38% (-0.22-1.1%)	0.16 (-0.096-0.48)	36 (15-64)
Comoros	2010	Female	20 to 24	26% (12-45%)	74% (55-88%)	0.21 (0.096-0.36)	1.1% (0.57-1.8%)	0.56% (-0.26-1.5%)	0.29 (-0.13-0.8)	73 (34-120)
Comoros	2010	Female	25 to 29	27% (12-46%)	73% (54-88%)	0.24 (0.11-0.43)	1.3% (0.78-1.9%)	0.81% (0.035-1.7%)	0.41 (0.018-0.86)	76 (42-110)
Comoros	2010	Female	30 to 34	30% (12-47%)	70% (53-88%)	0.27 (0.12-0.47)	1.2% (0.66-1.8%)	0.91% (0.0067-1.8%)	0.46 (0.0035-0.92)	64 (32-96)
Comoros	2010	Female	35 to 39	34% (15-51%)	66% (49-85%)	0.28 (0.12-0.49)	1.3% (0.6-2%)	1.3% (0.14-2.5%)	0.7 (0.078-1.4)	61 (27-96)
Comoros	2010	Female	40 to 44	35% (19-52%)	65% (48-81%)	0.27 (0.12-0.49)	1.2% (0.46-1.9%)	1.4% (0.2-2.5%)	0.84 (0.13-1.5)	55 (21-88)
Comoros	2010	Female	45 to 49	34% (17-48%)	66% (52-83%)	0.28 (0.12-0.52)	1% (0.27-1.8%)	1.2% (0.11-2.4%)	0.89 (0.082-1.8)	47 (12-82)
Comoros	2010	Female	50 to 54	33% (18-48%)	67% (52-82%)	0.28 (0.12-0.52)	0.88% (0.079-1.7%)	1.1% (-0.0088-2.3%)	0.99 (-0.008-2)	41 (3.2-79)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	2010	Female	55 to 59	33% (18-49%)	67% (51-82%)	0.3 (0.12-0.58)	0.45% (-0.38-1.3%)	0.58% (-0.54-1.7%)	0.6 (-0.58-1.8)	20 (-18-59)
Comoros	2010	Female	60 to 64	35% (19-52%)	65% (48-81%)	0.31 (0.12-0.62)	0.36% (-0.43-1.2%)	0.48% (-0.51-1.6%)	0.51 (-0.55-1.7)	13 (-15-45)
Comoros	2010	Female	65 to 69	37% (20-54%)	63% (46-80%)	0.3 (0.11-0.57)	0.35% (-0.55-1.2%)	0.42% (-0.67-1.4%)	0.56 (-0.89-2)	13 (-20-45)
Comoros	2010	Female	70 to 74	35% (19-52%)	65% (48-81%)	0.26 (0.097-0.5)	-0.032% (-0.83-0.69%)	-0.089% (-1-0.78%)	-0.14 (-1.6-1.2)	-1.1 (-29-23)
Comoros	2010	Female	75 to 79	31% (16-49%)	69% (51-84%)	0.21 (0.076-0.47)	0.086% (-0.83-0.77%)	0.099% (-0.94-0.91%)	0.15 (-1.5-1.4)	2.2 (-22-20)
Comoros	2010	Female	80 to 84	28% (14-46%)	72% (54-86%)	0.18 (0.067-0.39)	-0.16% (-0.98-0.39%)	-0.18% (-1.1-0.46%)	-0.21 (-1.4-0.53)	-2.3 (-15-5.6)
Comoros	2010	Female	85 to 89	28% (15-44%)	72% (56-85%)	0.17 (0.071-0.37)	-0.23% (-0.98-0.24%)	-0.28% (-1.1-0.28%)	-0.22 (-0.89-0.22)	-1.6 (-6.5-1.7)
Comoros	2010	Female	90 to 94	30% (16-45%)	70% (55-84%)	0.18 (0.07-0.35)	-0.34% (-1-0.14%)	-0.45% (-1.3-0.11%)	-0.13 (-0.4-0.034)	-0.61 (-1.9-0.25)
Comoros	2010	Female	95 plus	33% (18-53%)	67% (47-82%)	0.19 (0.066-0.39)	-0.0089% (-0.78-0.56%)	-0.1% (-1-0.6%)	-0.0076 (-0.078-0.045)	-0.0026 (-0.24-0.16)
Comoros	2010	Male	All Ages	8.1% (6.7-9.6%)	92% (90-93%)	0.02 (0.013-0.03)	0.75% (0.51-1%)	0.64% (0.25-1.1%)	14 (5.4-23)	1100 (720-1500)
Comoros	2010	Male	15 to 19	6.8% (4.3-9.9%)	93% (90-96%)	0.016 (0.0079-0.03)	1% (0.68-1.5%)	0.15% (0.041-0.29%)	0.073 (0.02-0.14)	67 (41-100)
Comoros	2010	Male	20 to 24	9.9% (6.1-14%)	90% (86-94%)	0.026 (0.013-0.045)	2.1% (1.4-3.1%)	0.27% (0.037-0.55%)	0.18 (0.023-0.38)	150 (92-230)
Comoros	2010	Male	25 to 29	12% (7.3-17%)	88% (83-93%)	0.034 (0.018-0.058)	2.4% (1.7-3.2%)	0.7% (0.2-1.3%)	0.48 (0.13-0.89)	160 (110-220)
Comoros	2010	Male	30 to 34	13% (7.9-20%)	87% (80-92%)	0.041 (0.021-0.069)	2.2% (1.5-3%)	0.94% (0.31-1.8%)	0.62 (0.2-1.2)	130 (83-180)
Comoros	2010	Male	35 to 39	14% (8.5-19%)	86% (81-91%)	0.046 (0.022-0.083)	1.9% (1.3-2.7%)	1.3% (0.41-2.3%)	0.88 (0.27-1.6)	100 (65-150)
Comoros	2010	Male	40 to 44	14% (9-19%)	86% (81-91%)	0.047 (0.022-0.083)	1.9% (1.1-2.8%)	1.7% (0.57-3%)	1.2 (0.42-2.2)	96 (55-150)
Comoros	2010	Male	45 to 49	14% (8.1-19%)	86% (81-92%)	0.048 (0.021-0.091)	1.9% (1.1-2.9%)	1.9% (0.71-3.4%)	1.7 (0.63-3)	97 (53-150)
Comoros	2010	Male	50 to 54	14% (9.1-20%)	86% (80-91%)	0.048 (0.022-0.092)	1.7% (0.87-2.7%)	1.7% (0.63-3.1%)	1.8 (0.65-3.2)	85 (43-140)
Comoros	2010	Male	55 to 59	14% (8.9-19%)	86% (81-91%)	0.046 (0.021-0.087)	1.5% (0.76-2.3%)	1.5% (0.56-2.7%)	1.6 (0.59-2.9)	67 (34-110)
Comoros	2010	Male	60 to 64	13% (8.4-19%)	87% (81-92%)	0.04 (0.017-0.078)	1.4% (0.69-2.3%)	1.5% (0.57-2.7%)	1.7 (0.6-3)	53 (26-90)
Comoros	2010	Male	65 to 69	12% (7.7-17%)	88% (83-92%)	0.032 (0.014-0.059)	0.86% (0.4-1.4%)	0.92% (0.31-1.6%)	1.1 (0.38-2.1)	30 (14-51)
Comoros	2010	Male	70 to 74	11% (7.3-16%)	89% (84-93%)	0.024 (0.011-0.046)	0.65% (0.27-1.1%)	0.72% (0.25-1.3%)	0.96 (0.33-1.7)	19 (7.9-32)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	2010	Male	75 to 79	10% (6.8-16%)	90% (84-93%)	0.019 (0.0083-0.038)	0.63% (0.27-1.1%)	0.71% (0.26-1.3%)	0.82 (0.29-1.5)	12 (5-21)
Comoros	2010	Male	80 to 84	9.5% (5.9-13%)	91% (87-94%)	0.016 (0.0073-0.032)	0.46% (0.23-0.73%)	0.48% (0.19-0.8%)	0.37 (0.16-0.63)	4.4 (2.2-7)
Comoros	2010	Male	85 to 89	8.6% (5-13%)	91% (87-95%)	0.014 (0.0064-0.028)	0.47% (0.24-0.76%)	0.48% (0.17-0.83%)	0.24 (0.083-0.41)	2 (1-3.3)
Comoros	2010	Male	90 to 94	8.6% (5.5-13%)	91% (87-94%)	0.013 (0.0054-0.026)	0.35% (0.2-0.54%)	0.32% (0.13-0.55%)	0.059 (0.025-0.1)	0.39 (0.22-0.59)
Comoros	2010	Male	95 plus	9.5% (5.5-15%)	90% (85-95%)	0.013 (0.0046-0.027)	0.48% (0.25-0.76%)	0.43% (0.15-0.79%)	0.017 (0.0056-0.031)	0.075 (0.038-0.12)
Comoros	2016	Female	All Ages	24% (17-30%)	76% (70-83%)	0.17 (0.089-0.28)	0.53% (0.21-0.9%)	0.43% (-0.13-1.1%)	9 (-2.8-24)	660 (270-1100)
Comoros	2016	Female	15 to 19	21% (9.9-38%)	79% (62-90%)	0.17 (0.058-0.35)	0.67% (0.34-1.2%)	0.46% (-0.13-1.3%)	0.18 (-0.052-0.5)	41 (19-71)
Comoros	2016	Female	20 to 24	26% (11-44%)	74% (56-89%)	0.23 (0.084-0.45)	1.3% (0.7-2%)	0.69% (-0.15-1.9%)	0.32 (-0.068-0.84)	82 (42-130)
Comoros	2016	Female	25 to 29	27% (12-46%)	73% (54-88%)	0.27 (0.11-0.53)	1.5% (0.89-2.2%)	1% (0.08-2.2%)	0.51 (0.033-1.1)	91 (51-140)
Comoros	2016	Female	30 to 34	30% (13-47%)	70% (53-87%)	0.3 (0.12-0.57)	1.4% (0.76-2.1%)	1.2% (0.19-2.3%)	0.65 (0.1-1.3)	84 (43-130)
Comoros	2016	Female	35 to 39	35% (16-51%)	65% (49-84%)	0.31 (0.13-0.59)	1.4% (0.68-2.3%)	1.6% (0.27-3.1%)	0.95 (0.17-1.9)	81 (38-130)
Comoros	2016	Female	40 to 44	36% (20-52%)	64% (48-80%)	0.31 (0.12-0.63)	1.3% (0.62-2.3%)	1.6% (0.48-3.1%)	1.1 (0.31-2.1)	72 (32-120)
Comoros	2016	Female	45 to 49	34% (17-48%)	66% (52-83%)	0.31 (0.11-0.67)	1.2% (0.41-2.1%)	1.5% (0.28-2.9%)	1.2 (0.22-2.4)	62 (21-110)
Comoros	2016	Female	50 to 54	33% (18-48%)	67% (52-82%)	0.31 (0.11-0.66)	1.1% (0.19-2.1%)	1.3% (0.13-2.9%)	1.4 (0.11-3)	58 (11-110)
Comoros	2016	Female	55 to 59	33% (19-50%)	67% (50-81%)	0.33 (0.12-0.72)	0.62% (-0.29-1.8%)	0.79% (-0.39-2.4%)	1 (-0.52-3.2)	34 (-16-100)
Comoros	2016	Female	60 to 64	35% (20-53%)	65% (47-80%)	0.34 (0.11-0.74)	0.51% (-0.31-1.6%)	0.67% (-0.32-2.1%)	0.99 (-0.48-3.2)	27 (-16-87)
Comoros	2016	Female	65 to 69	37% (20-53%)	63% (47-80%)	0.33 (0.11-0.68)	0.48% (-0.38-1.4%)	0.58% (-0.51-1.8%)	0.86 (-0.7-2.7)	20 (-16-61)
Comoros	2016	Female	70 to 74	35% (19-53%)	65% (47-81%)	0.28 (0.099-0.63)	0.07% (-0.72-0.89%)	0.025% (-0.96-1%)	0.041 (-1.6-1.6)	2.6 (-27-31)
Comoros	2016	Female	75 to 79	31% (15-49%)	69% (51-85%)	0.23 (0.075-0.55)	0.16% (-0.75-0.89%)	0.19% (-0.88-1.1%)	0.32 (-1.4-1.9)	4.7 (-21-26)
Comoros	2016	Female	80 to 84	29% (13-47%)	71% (53-87%)	0.21 (0.064-0.48)	-0.12% (-0.93-0.47%)	-0.14% (-1.1-0.55%)	-0.18 (-1.5-0.73)	-1.9 (-15-7.6)
Comoros	2016	Female	85 to 89	28% (14-46%)	72% (54-86%)	0.19 (0.068-0.43)	-0.21% (-0.97-0.27%)	-0.26% (-1.1-0.31%)	-0.24 (-1-0.28)	-1.7 (-7.6-2.1)
Comoros	2016	Female	90 to 94	30% (16-47%)	70% (53-84%)	0.2 (0.07-0.45)	-0.33% (-1-0.19%)	-0.44% (-1.3-0.18%)	-0.18 (-0.52-0.072)	-0.79 (-2.5-0.43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Comoros	2016	Female	95 plus	34% (18-52%)	66% (48-82%)	0.21 (0.069-0.48)	0.003% (-0.78-0.64%)	-0.084% (-1-0.68%)	-0.0092 (-0.12-0.073)	0.0014 (-0.35-0.27)
Comoros	2016	Male	All Ages	8.3% (6.9-9.9%)	92% (90-93%)	0.024 (0.012-0.041)	0.96% (0.64-1.3%)	0.79% (0.33-1.3%)	17 (6.9-29)	1300 (890-1800)
Comoros	2016	Male	15 to 19	6.8% (4.3-9.8%)	93% (90-96%)	0.018 (0.0075-0.037)	1.1% (0.7-1.5%)	0.16% (0.049-0.33%)	0.086 (0.025-0.18)	76 (47-110)
Comoros	2016	Male	20 to 24	9.8% (6.2-14%)	90% (86-94%)	0.029 (0.012-0.057)	2.2% (1.4-3.2%)	0.3% (0.044-0.62%)	0.22 (0.034-0.49)	170 (110-260)
Comoros	2016	Male	25 to 29	12% (7.3-17%)	88% (83-93%)	0.038 (0.017-0.073)	2.5% (1.7-3.4%)	0.76% (0.23-1.4%)	0.6 (0.18-1.1)	190 (130-270)
Comoros	2016	Male	30 to 34	13% (7.8-19%)	87% (81-92%)	0.045 (0.02-0.085)	2.2% (1.6-3.1%)	1% (0.35-1.9%)	0.83 (0.26-1.5)	160 (110-240)
Comoros	2016	Male	35 to 39	14% (8.5-20%)	86% (80-92%)	0.051 (0.021-0.1)	2% (1.3-2.9%)	1.4% (0.45-2.5%)	1.2 (0.39-2.2)	140 (87-190)
Comoros	2016	Male	40 to 44	14% (8.8-20%)	86% (80-91%)	0.053 (0.021-0.11)	2% (1.2-2.9%)	1.8% (0.66-3.1%)	1.6 (0.57-2.9)	120 (69-180)
Comoros	2016	Male	45 to 49	14% (7.8-20%)	86% (80-92%)	0.053 (0.02-0.12)	2% (1.1-3.1%)	2% (0.76-3.6%)	2 (0.77-3.7)	120 (65-190)
Comoros	2016	Male	50 to 54	14% (9.1-20%)	86% (80-91%)	0.053 (0.021-0.11)	1.7% (0.9-2.7%)	1.8% (0.64-3.1%)	2.2 (0.79-3.9)	100 (53-160)
Comoros	2016	Male	55 to 59	14% (8.9-19%)	86% (81-91%)	0.052 (0.02-0.11)	1.5% (0.8-2.4%)	1.6% (0.6-2.8%)	2.1 (0.78-3.9)	87 (42-140)
Comoros	2016	Male	60 to 64	13% (8.3-19%)	87% (81-92%)	0.045 (0.016-0.098)	1.4% (0.73-2.4%)	1.6% (0.62-2.8%)	2.4 (0.9-4.2)	75 (37-120)
Comoros	2016	Male	65 to 69	12% (7.8-17%)	88% (83-92%)	0.035 (0.014-0.074)	0.89% (0.41-1.5%)	0.96% (0.31-1.7%)	1.3 (0.45-2.5)	35 (15-60)
Comoros	2016	Male	70 to 74	11% (7.5-16%)	89% (84-93%)	0.027 (0.01-0.056)	0.67% (0.3-1.1%)	0.75% (0.27-1.3%)	1.1 (0.39-1.9)	21 (9.5-36)
Comoros	2016	Male	75 to 79	11% (6.7-16%)	89% (84-93%)	0.022 (0.0085-0.046)	0.66% (0.3-1.1%)	0.74% (0.27-1.3%)	0.97 (0.34-1.8)	15 (6.4-25)
Comoros	2016	Male	80 to 84	9.5% (5.8-14%)	90% (86-94%)	0.018 (0.0073-0.038)	0.47% (0.25-0.75%)	0.49% (0.21-0.83%)	0.43 (0.19-0.73)	5.1 (2.6-8.1)
Comoros	2016	Male	85 to 89	8.7% (5.1-13%)	91% (87-95%)	0.016 (0.0063-0.033)	0.49% (0.23-0.79%)	0.5% (0.18-0.88%)	0.28 (0.1-0.49)	2.4 (1.1-3.8)
Comoros	2016	Male	90 to 94	8.7% (5.5-14%)	91% (86-94%)	0.015 (0.0057-0.031)	0.37% (0.21-0.55%)	0.33% (0.14-0.56%)	0.081 (0.033-0.14)	0.53 (0.3-0.8)
Comoros	2016	Male	95 plus	9.5% (5.5-15%)	91% (85-94%)	0.015 (0.0045-0.034)	0.49% (0.26-0.78%)	0.45% (0.16-0.81%)	0.026 (0.009-0.048)	0.12 (0.058-0.19)
Djibouti	1990	Female	All Ages	32% (27-39%)	68% (61-73%)	0.25 (0.18-0.34)	0.23% (0.1-0.39%)	0.21% (-0.1-0.55%)	5.1 (-2.5-14)	430 (190-720)
Djibouti	1990	Female	15 to 19	34% (21-48%)	66% (52-79%)	0.3 (0.15-0.52)	0.63% (0.31-1.1%)	0.4% (-0.11-1.1%)	0.16 (-0.039-0.43)	34 (16-59)
Djibouti	1990	Female	20 to 24	40% (25-56%)	60% (44-75%)	0.41 (0.21-0.69)	1.1% (0.58-1.7%)	0.49% (-0.12-1.1%)	0.23 (-0.053-0.56)	60 (30-94)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	1990	Female	25 to 29	43% (26-59%)	57% (41-74%)	0.49 (0.27-0.79)	1.3% (0.75-2%)	0.86% (0.12-1.8%)	0.39 (0.053-0.82)	65 (36-100)
Djibouti	1990	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.52 (0.29-0.84)	1.3% (0.7-2%)	1% (0.1-2.1%)	0.46 (0.043-0.99)	56 (29-89)
Djibouti	1990	Female	35 to 39	48% (33-63%)	52% (37-67%)	0.5 (0.27-0.8)	1.2% (0.61-2%)	1.2% (0.25-2.4%)	0.57 (0.11-1.1)	48 (24-78)
Djibouti	1990	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.49 (0.25-0.83)	1.2% (0.57-2%)	1.5% (0.46-2.6%)	0.76 (0.24-1.4)	48 (23-77)
Djibouti	1990	Female	45 to 49	49% (32-65%)	51% (35-68%)	0.5 (0.26-0.86)	1.1% (0.39-2.1%)	1.4% (0.33-2.8%)	0.82 (0.19-1.6)	41 (13-75)
Djibouti	1990	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.51 (0.27-0.86)	1.1% (0.28-2.1%)	1.5% (0.3-2.8%)	0.92 (0.19-1.8)	37 (8.8-70)
Djibouti	1990	Female	55 to 59	46% (31-62%)	54% (38-69%)	0.52 (0.26-0.9)	0.72% (-0.11-1.8%)	0.98% (-0.16-2.5%)	0.72 (-0.12-1.9)	23 (-3.5-60)
Djibouti	1990	Female	60 to 64	46% (30-64%)	54% (36-70%)	0.51 (0.25-0.9)	0.54% (-0.3-1.6%)	0.74% (-0.34-2.1%)	0.65 (-0.32-1.9)	17 (-9.5-51)
Djibouti	1990	Female	65 to 69	48% (32-64%)	52% (36-68%)	0.47 (0.25-0.82)	0.4% (-0.56-1.4%)	0.51% (-0.66-1.7%)	0.52 (-0.67-1.7)	12 (-16-39)
Djibouti	1990	Female	70 to 74	46% (32-62%)	54% (38-68%)	0.42 (0.2-0.76)	-0.084% (-1-0.77%)	-0.14% (-1.3-0.91%)	-0.14 (-1.4-1)	-2 (-25-20)
Djibouti	1990	Female	75 to 79	44% (28-60%)	56% (40-72%)	0.36 (0.16-0.66)	-0.14% (-1.1-0.67%)	-0.16% (-1.3-0.83%)	-0.15 (-1.3-0.79)	-2.3 (-17-11)
Djibouti	1990	Female	80 to 84	43% (27-60%)	57% (40-73%)	0.33 (0.15-0.58)	-0.44% (-1.3-0.23%)	-0.51% (-1.6-0.33%)	-0.35 (-1.1-0.22)	-3.8 (-11-2)
Djibouti	1990	Female	85 to 89	43% (27-58%)	57% (42-73%)	0.32 (0.14-0.57)	-0.56% (-1.5-0.095%)	-0.66% (-1.7-0.13%)	-0.27 (-0.71-0.053)	-2 (-5.3-0.36)
Djibouti	1990	Female	90 to 94	45% (29-60%)	55% (40-71%)	0.31 (0.15-0.6)	-0.67% (-1.5--0.03%)	-0.83% (-1.9--0.06%)	-0.13 (-0.3--0.0097)	-0.65 (-1.4--0.03)
Djibouti	1990	Female	95 plus	47% (29-64%)	53% (36-71%)	0.31 (0.13-0.61)	-0.36% (-1.3-0.29%)	-0.52% (-1.6-0.28%)	-0.02 (-0.064-0.011)	-0.057 (-0.2-0.047)
Djibouti	1990	Male	All Ages	41% (35-48%)	59% (52-65%)	0.3 (0.22-0.41)	0.64% (0.27-1%)	0.68% (0.066-1.3%)	21 (2.3-39)	1400 (630-2300)
Djibouti	1990	Male	15 to 19	41% (28-56%)	59% (44-72%)	0.36 (0.19-0.6)	1.5% (0.84-2.4%)	1.1% (0.2-2.4%)	0.67 (0.12-1.4)	100 (54-160)
Djibouti	1990	Male	20 to 24	51% (37-66%)	49% (34-63%)	0.48 (0.27-0.77)	2.7% (1.3-4.4%)	1.6% (-0.022-3.5%)	1.4 (-0.018-3)	210 (99-340)
Djibouti	1990	Male	25 to 29	55% (40-70%)	45% (30-60%)	0.57 (0.34-0.9)	3.2% (1.4-5.1%)	2.2% (-0.13-4.7%)	1.9 (-0.12-4)	230 (96-380)
Djibouti	1990	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.64 (0.39-1)	3.1% (1.1-5.2%)	2.5% (-0.19-5.4%)	2 (-0.14-4.2)	190 (61-320)
Djibouti	1990	Male	35 to 39	59% (45-74%)	41% (26-55%)	0.68 (0.4-1.1)	3.1% (0.91-5.4%)	2.9% (-0.035-6%)	2 (-0.023-4.2)	150 (42-260)
Djibouti	1990	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.71 (0.4-1.1)	2.9% (1-4.9%)	3% (0.51-5.7%)	2.4 (0.41-4.7)	150 (51-250)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	1990	Male	45 to 49	60% (44-75%)	40% (25-56%)	0.7 (0.38-1.1)	2.6% (0.81-4.4%)	2.8% (0.39-5.1%)	2.4 (0.33-4.5)	120 (37-210)
Djibouti	1990	Male	50 to 54	60% (45-75%)	40% (25-55%)	0.69 (0.38-1.1)	2.4% (0.62-4.2%)	2.6% (0.34-4.8%)	2.5 (0.33-4.8)	110 (28-190)
Djibouti	1990	Male	55 to 59	60% (45-75%)	40% (25-55%)	0.67 (0.37-1.1)	2% (0.37-3.6%)	2.1% (0.12-4.2%)	2.2 (0.12-4.5)	82 (16-150)
Djibouti	1990	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.61 (0.32-1)	1.6% (0.3-2.9%)	1.7% (0.12-3.3%)	2 (0.13-4)	60 (11-120)
Djibouti	1990	Male	65 to 69	57% (42-73%)	43% (27-58%)	0.51 (0.27-0.85)	0.75% (-0.29-1.8%)	0.77% (-0.54-2%)	0.93 (-0.64-2.5)	24 (-9.6-58)
Djibouti	1990	Male	70 to 74	56% (41-70%)	44% (30-59%)	0.41 (0.21-0.71)	0.41% (-0.47-1.2%)	0.43% (-0.57-1.4%)	0.51 (-0.71-1.7)	11 (-12-32)
Djibouti	1990	Male	75 to 79	54% (40-69%)	46% (31-60%)	0.35 (0.16-0.62)	0.28% (-0.63-1%)	0.27% (-0.79-1.1%)	0.27 (-0.8-1.1)	4.5 (-10-16)
Djibouti	1990	Male	80 to 84	52% (35-69%)	48% (31-65%)	0.31 (0.15-0.52)	0.14% (-0.58-0.74%)	0.087% (-0.76-0.79%)	0.054 (-0.47-0.48)	1 (-4.3-5.6)
Djibouti	1990	Male	85 to 89	50% (35-64%)	50% (36-65%)	0.27 (0.14-0.45)	0.13% (-0.54-0.67%)	0.057% (-0.75-0.67%)	0.017 (-0.23-0.2)	0.34 (-1.4-1.7)
Djibouti	1990	Male	90 to 94	51% (35-65%)	49% (35-65%)	0.25 (0.12-0.44)	-0.021% (-0.61-0.44%)	-0.16% (-0.87-0.38%)	-0.016 (-0.086-0.038)	-0.012 (-0.36-0.26)
Djibouti	1990	Male	95 plus	52% (33-71%)	48% (29-67%)	0.23 (0.098-0.44)	0.078% (-0.63-0.58%)	-0.11% (-0.96-0.48%)	-0.0022 (-0.019-0.0095)	0.0062 (-0.048-0.045)
Djibouti	1995	Female	All Ages	32% (27-39%)	68% (61-73%)	0.19 (0.13-0.26)	0.21% (0.071-0.36%)	0.17% (-0.14-0.49%)	4.8 (-4-14)	440 (150-750)
Djibouti	1995	Female	15 to 19	34% (22-49%)	66% (51-78%)	0.24 (0.12-0.4)	0.55% (0.21-0.96%)	0.29% (-0.25-0.89%)	0.14 (-0.12-0.45)	36 (13-63)
Djibouti	1995	Female	20 to 24	40% (25-56%)	60% (44-75%)	0.32 (0.17-0.55)	0.97% (0.47-1.5%)	0.34% (-0.28-0.98%)	0.2 (-0.16-0.58)	62 (30-100)
Djibouti	1995	Female	25 to 29	43% (27-59%)	57% (41-73%)	0.39 (0.21-0.63)	1.2% (0.62-1.8%)	0.62% (-0.17-1.4%)	0.35 (-0.098-0.8)	67 (34-110)
Djibouti	1995	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.4 (0.22-0.65)	1.1% (0.52-1.8%)	0.78% (-0.13-1.8%)	0.45 (-0.07-1)	61 (28-98)
Djibouti	1995	Female	35 to 39	48% (32-63%)	52% (37-68%)	0.38 (0.21-0.61)	1.1% (0.41-1.8%)	0.99% (0.0046-2.1%)	0.59 (0.003-1.2)	53 (21-88)
Djibouti	1995	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.38 (0.2-0.64)	1.1% (0.47-1.8%)	1.3% (0.32-2.3%)	0.73 (0.19-1.3)	47 (20-77)
Djibouti	1995	Female	45 to 49	49% (32-66%)	51% (34-68%)	0.39 (0.2-0.68)	1% (0.25-1.8%)	1.2% (0.17-2.5%)	0.84 (0.1-1.7)	43 (11-77)
Djibouti	1995	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.39 (0.19-0.67)	0.94% (0.21-1.8%)	1.2% (0.19-2.4%)	0.93 (0.14-1.9)	38 (8.3-74)
Djibouti	1995	Female	55 to 59	46% (31-63%)	54% (37-69%)	0.4 (0.2-0.7)	0.54% (-0.23-1.4%)	0.74% (-0.3-1.9%)	0.65 (-0.26-1.8)	21 (-8.4-56)
Djibouti	1995	Female	60 to 64	47% (31-64%)	53% (36-69%)	0.4 (0.2-0.7)	0.4% (-0.36-1.3%)	0.56% (-0.42-1.6%)	0.59 (-0.44-1.8)	15 (-13-47)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	1995	Female	65 to 69	48% (32-63%)	52% (37-68%)	0.37 (0.19-0.65)	0.33% (-0.45-1.1%)	0.42% (-0.57-1.4%)	0.51 (-0.72-1.7)	12 (-16-37)
Djibouti	1995	Female	70 to 74	46% (33-61%)	54% (39-67%)	0.32 (0.16-0.61)	-0.099% (-0.92-0.59%)	-0.16% (-1.2-0.69%)	-0.2 (-1.6-0.92)	-2.9 (-28-18)
Djibouti	1995	Female	75 to 79	44% (29-60%)	56% (40-71%)	0.29 (0.12-0.53)	-0.094% (-1-0.63%)	-0.098% (-1.2-0.77%)	-0.11 (-1.4-0.93)	-1.9 (-20-13)
Djibouti	1995	Female	80 to 84	43% (27-60%)	57% (40-73%)	0.26 (0.12-0.47)	-0.38% (-1.2-0.24%)	-0.44% (-1.4-0.3%)	-0.38 (-1.2-0.27)	-4.2 (-13-2.5)
Djibouti	1995	Female	85 to 89	43% (27-59%)	57% (41-73%)	0.25 (0.11-0.46)	-0.47% (-1.3-0.1%)	-0.55% (-1.5-0.14%)	-0.3 (-0.83-0.074)	-2.2 (-6.1-0.48)
Djibouti	1995	Female	90 to 94	45% (29-60%)	55% (40-71%)	0.24 (0.11-0.48)	-0.54% (-1.3-0.015%)	-0.68% (-1.6-0.0017%)	-0.15 (-0.35-0.0037)	-0.71 (-1.7-0.021)
Djibouti	1995	Female	95 plus	47% (29-64%)	53% (36-71%)	0.24 (0.099-0.48)	-0.22% (-1-0.38%)	-0.34% (-1.3-0.39%)	-0.02 (-0.078-0.022)	-0.05 (-0.23-0.088)
Djibouti	1995	Male	All Ages	41% (35-48%)	59% (52-65%)	0.24 (0.17-0.32)	0.58% (0.2-0.96%)	0.59% (-0.028-1.2%)	21 (-1-43)	1500 (520-2400)
Djibouti	1995	Male	15 to 19	41% (28-57%)	59% (43-72%)	0.28 (0.15-0.47)	1.4% (0.7-2.2%)	0.9% (-0.00096-2%)	0.63 (-0.00068-1.4)	110 (53-170)
Djibouti	1995	Male	20 to 24	52% (37-66%)	48% (34-63%)	0.37 (0.21-0.6)	2.5% (1.1-4.1%)	1.3% (-0.47-3.2%)	1.2 (-0.43-2.9)	200 (87-330)
Djibouti	1995	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.45 (0.26-0.73)	2.8% (1-4.7%)	1.7% (-0.74-4.2%)	1.6 (-0.69-3.8)	220 (78-370)
Djibouti	1995	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.5 (0.3-0.79)	2.7% (0.62-4.8%)	1.9% (-0.88-4.6%)	1.9 (-0.86-4.6)	200 (46-370)
Djibouti	1995	Male	35 to 39	59% (46-74%)	41% (26-54%)	0.53 (0.31-0.86)	2.6% (0.55-4.6%)	2.2% (-0.47-4.8%)	2.1 (-0.43-4.6)	170 (33-300)
Djibouti	1995	Male	40 to 44	60% (46-76%)	40% (24-54%)	0.55 (0.3-0.87)	2.4% (0.58-4.3%)	2.4% (-0.073-5%)	2 (-0.062-4.2)	130 (30-230)
Djibouti	1995	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.55 (0.29-0.89)	2.3% (0.51-4%)	2.3% (-0.012-4.5%)	2.4 (-0.011-4.7)	130 (29-220)
Djibouti	1995	Male	50 to 54	61% (45-76%)	39% (24-55%)	0.54 (0.28-0.88)	2.1% (0.35-3.8%)	2.2% (0.038-4.4%)	2.5 (0.041-4.9)	110 (18-200)
Djibouti	1995	Male	55 to 59	60% (45-75%)	40% (25-55%)	0.53 (0.29-0.88)	1.7% (0.055-3.2%)	1.8% (-0.26-3.7%)	2.2 (-0.32-4.6)	82 (2.7-160)
Djibouti	1995	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.47 (0.24-0.83)	1.4% (0.21-2.6%)	1.5% (-0.026-3%)	2 (-0.035-4.1)	64 (9.4-120)
Djibouti	1995	Male	65 to 69	58% (42-74%)	42% (26-58%)	0.39 (0.21-0.68)	0.72% (-0.29-1.7%)	0.73% (-0.5-1.9%)	1.1 (-0.73-2.7)	28 (-12-64)
Djibouti	1995	Male	70 to 74	56% (41-71%)	44% (29-59%)	0.32 (0.17-0.55)	0.44% (-0.46-1.2%)	0.46% (-0.6-1.4%)	0.67 (-0.87-2)	14 (-14-36)
Djibouti	1995	Male	75 to 79	54% (40-70%)	46% (30-60%)	0.27 (0.13-0.48)	0.36% (-0.44-1%)	0.36% (-0.63-1.1%)	0.45 (-0.77-1.4)	7.2 (-9.4-21)
Djibouti	1995	Male	80 to 84	53% (36-70%)	47% (30-64%)	0.24 (0.12-0.43)	0.21% (-0.49-0.77%)	0.17% (-0.65-0.83%)	0.14 (-0.53-0.67)	2 (-4.9-7.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	1995	Male	85 to 89	51% (35-65%)	49% (35-65%)	0.21 (0.11-0.36)	0.22% (-0.37-0.71%)	0.17% (-0.52-0.72%)	0.07 (-0.21-0.3)	0.79 (-1.3-2.5)
Djibouti	1995	Male	90 to 94	51% (35-65%)	49% (35-65%)	0.19 (0.095-0.36)	0.079% (-0.42-0.48%)	-0.027% (-0.6-0.43%)	-0.0037 (-0.081-0.059)	0.064 (-0.33-0.39)
Djibouti	1995	Male	95 plus	53% (33-71%)	47% (29-67%)	0.18 (0.075-0.35)	0.19% (-0.37-0.62%)	0.047% (-0.66-0.55%)	0.0014 (-0.019-0.016)	0.023 (-0.043-0.073)
Djibouti	2000	Female	All Ages	33% (27-39%)	67% (61-73%)	0.19 (0.13-0.25)	0.22% (0.013-0.43%)	0.16% (-0.28-0.58%)	5.1 (-8.9-19)	490 (28-970)
Djibouti	2000	Female	15 to 19	34% (21-49%)	66% (51-79%)	0.23 (0.12-0.39)	0.55% (0.11-1%)	0.3% (-0.44-1.1%)	0.14 (-0.21-0.53)	36 (6.9-68)
Djibouti	2000	Female	20 to 24	40% (25-57%)	60% (43-75%)	0.32 (0.17-0.52)	0.88% (0.34-1.5%)	0.29% (-0.42-0.96%)	0.22 (-0.31-0.77)	72 (27-130)
Djibouti	2000	Female	25 to 29	43% (27-60%)	57% (40-73%)	0.38 (0.21-0.62)	0.94% (0.38-1.6%)	0.46% (-0.33-1.3%)	0.39 (-0.29-1.1)	74 (28-120)
Djibouti	2000	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.4 (0.22-0.66)	0.92% (0.29-1.6%)	0.58% (-0.32-1.6%)	0.51 (-0.28-1.3)	67 (22-120)
Djibouti	2000	Female	35 to 39	48% (32-63%)	52% (37-68%)	0.38 (0.21-0.61)	0.89% (0.099-1.7%)	0.74% (-0.37-1.9%)	0.66 (-0.35-1.7)	60 (6.6-120)
Djibouti	2000	Female	40 to 44	49% (34-65%)	51% (35-66%)	0.38 (0.2-0.61)	0.94% (0.16-1.7%)	1% (-0.1-2.2%)	0.83 (-0.076-1.8)	54 (8.6-100)
Djibouti	2000	Female	45 to 49	49% (32-65%)	51% (35-68%)	0.38 (0.19-0.64)	0.88% (-0.081-1.9%)	1% (-0.37-2.5%)	0.84 (-0.3-2)	43 (-4.2-92)
Djibouti	2000	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.39 (0.2-0.66)	0.86% (-0.13-1.9%)	1.1% (-0.3-2.6%)	1.1 (-0.28-2.5)	43 (-6.8-94)
Djibouti	2000	Female	55 to 59	46% (30-63%)	54% (37-70%)	0.4 (0.2-0.68)	0.51% (-0.55-1.6%)	0.68% (-0.69-2.2%)	0.73 (-0.8-2.6)	23 (-24-80)
Djibouti	2000	Female	60 to 64	47% (31-64%)	53% (36-69%)	0.39 (0.2-0.7)	0.39% (-0.53-1.4%)	0.55% (-0.62-1.8%)	0.7 (-0.76-2.4)	17 (-22-62)
Djibouti	2000	Female	65 to 69	48% (32-64%)	52% (36-68%)	0.37 (0.19-0.64)	0.3% (-0.69-1.2%)	0.4% (-0.83-1.6%)	0.57 (-1.2-2.2)	12 (-27-49)
Djibouti	2000	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.32 (0.16-0.6)	-0.13% (-1.1-0.67%)	-0.19% (-1.3-0.81%)	-0.29 (-2.1-1.2)	-4.5 (-39-23)
Djibouti	2000	Female	75 to 79	44% (29-61%)	56% (39-71%)	0.28 (0.12-0.52)	-0.12% (-1.1-0.65%)	-0.13% (-1.3-0.83%)	-0.18 (-1.8-1.2)	-3 (-27-16)
Djibouti	2000	Female	80 to 84	43% (27-60%)	57% (40-73%)	0.26 (0.11-0.45)	-0.4% (-1.3-0.28%)	-0.45% (-1.5-0.34%)	-0.48 (-1.7-0.36)	-5.3 (-17-3.9)
Djibouti	2000	Female	85 to 89	43% (27-58%)	57% (42-73%)	0.24 (0.11-0.45)	-0.47% (-1.3-0.15%)	-0.55% (-1.6-0.17%)	-0.4 (-1.1-0.13)	-3 (-8.5-0.98)
Djibouti	2000	Female	90 to 94	45% (29-59%)	55% (41-71%)	0.24 (0.11-0.45)	-0.55% (-1.4-0.06%)	-0.69% (-1.7-0.03%)	-0.21 (-0.51-0.0093)	-0.99 (-2.5-0.11)
Djibouti	2000	Female	95 plus	47% (29-65%)	53% (35-71%)	0.24 (0.095-0.46)	-0.18% (-0.98-0.43%)	-0.31% (-1.3-0.45%)	-0.025 (-0.1-0.036)	-0.059 (-0.31-0.14)
Djibouti	2000	Male	All Ages	41% (35-49%)	59% (51-65%)	0.24 (0.17-0.31)	0.62% (0.12-1.1%)	0.61% (-0.21-1.4%)	24 (-8.1-55)	1700 (340-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	2000	Male	15 to 19	41% (28-57%)	59% (43-72%)	0.27 (0.15-0.45)	1.5% (0.61-2.5%)	1% (-0.31-2.6%)	0.64 (-0.21-1.6)	110 (43-180)
Djibouti	2000	Male	20 to 24	52% (38-66%)	48% (34-62%)	0.37 (0.21-0.59)	2.5% (0.77-4.3%)	1.3% (-0.94-3.7%)	1.4 (-0.94-3.7)	240 (70-420)
Djibouti	2000	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.44 (0.25-0.71)	2.5% (0.41-4.6%)	1.5% (-1.2-4.1%)	1.7 (-1.4-4.5)	230 (43-420)
Djibouti	2000	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.49 (0.3-0.76)	2.3% (0.065-4.5%)	1.6% (-1.2-4.4%)	2 (-1.6-5.4)	210 (6-420)
Djibouti	2000	Male	35 to 39	59% (45-74%)	41% (26-55%)	0.53 (0.31-0.84)	2.2% (0.067-4.3%)	1.9% (-0.86-4.7%)	2.6 (-1.3-6.4)	210 (5-400)
Djibouti	2000	Male	40 to 44	60% (45-75%)	40% (25-55%)	0.54 (0.3-0.86)	2.2% (0.18-4.2%)	2.1% (-0.48-4.7%)	2.7 (-0.63-6)	170 (14-320)
Djibouti	2000	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.54 (0.3-0.85)	2.1% (0.14-4%)	2.1% (-0.45-4.6%)	2.4 (-0.58-5.2)	130 (9.7-250)
Djibouti	2000	Male	50 to 54	61% (45-75%)	39% (25-55%)	0.53 (0.29-0.87)	2% (-0.096-3.9%)	2.1% (-0.58-4.6%)	2.8 (-0.73-6.4)	120 (-5.4-250)
Djibouti	2000	Male	55 to 59	60% (45-75%)	40% (25-55%)	0.52 (0.28-0.86)	1.6% (-0.34-3.5%)	1.7% (-0.7-4.1%)	2.5 (-1.1-6)	95 (-21-210)
Djibouti	2000	Male	60 to 64	59% (44-76%)	41% (24-56%)	0.47 (0.25-0.81)	1.4% (-0.1-2.7%)	1.5% (-0.41-3.2%)	2.4 (-0.66-5.1)	74 (-5.8-150)
Djibouti	2000	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.39 (0.22-0.67)	0.69% (-0.5-1.8%)	0.7% (-0.73-2%)	1.2 (-1.2-3.5)	32 (-23-83)
Djibouti	2000	Male	70 to 74	56% (41-71%)	44% (29-59%)	0.31 (0.17-0.54)	0.43% (-0.71-1.4%)	0.45% (-0.9-1.6%)	0.77 (-1.5-2.7)	16 (-26-51)
Djibouti	2000	Male	75 to 79	54% (39-70%)	46% (30-61%)	0.26 (0.13-0.48)	0.36% (-0.64-1.1%)	0.37% (-0.81-1.3%)	0.55 (-1.2-1.9)	8.8 (-16-28)
Djibouti	2000	Male	80 to 84	52% (35-69%)	48% (31-65%)	0.24 (0.11-0.4)	0.22% (-0.59-0.89%)	0.18% (-0.76-0.96%)	0.18 (-0.74-0.94)	2.5 (-7.1-11)
Djibouti	2000	Male	85 to 89	51% (34-65%)	49% (35-66%)	0.21 (0.1-0.36)	0.24% (-0.42-0.83%)	0.2% (-0.58-0.86%)	0.11 (-0.31-0.47)	1.1 (-2-3.9)
Djibouti	2000	Male	90 to 94	51% (36-66%)	49% (34-64%)	0.19 (0.095-0.33)	0.1% (-0.44-0.53%)	0.0042% (-0.63-0.51%)	0.0008 (-0.12-0.095)	0.12 (-0.49-0.6)
Djibouti	2000	Male	95 plus	53% (33-71%)	47% (29-67%)	0.18 (0.077-0.33)	0.24% (-0.36-0.77%)	0.1% (-0.62-0.73%)	0.0043 (-0.026-0.03)	0.04 (-0.061-0.13)
Djibouti	2005	Female	All Ages	33% (28-40%)	67% (60-72%)	0.22 (0.16-0.29)	0.26% (-0.0054-0.54%)	0.18% (-0.35-0.71%)	5.9 (-12-25)	570 (-12-1200)
Djibouti	2005	Female	15 to 19	34% (21-49%)	66% (51-79%)	0.26 (0.13-0.43)	0.62% (0.13-1.2%)	0.37% (-0.49-1.3%)	0.2 (-0.24-0.74)	47 (10-92)
Djibouti	2005	Female	20 to 24	41% (25-56%)	59% (44-75%)	0.35 (0.19-0.58)	0.97% (0.39-1.7%)	0.36% (-0.44-1.3%)	0.27 (-0.3-0.93)	77 (30-130)
Djibouti	2005	Female	25 to 29	43% (27-60%)	57% (40-73%)	0.42 (0.23-0.67)	0.88% (0.33-1.6%)	0.44% (-0.24-1.3%)	0.51 (-0.3-1.5)	91 (35-160)
Djibouti	2005	Female	30 to 34	45% (29-62%)	55% (38-71%)	0.44 (0.25-0.71)	0.8% (0.25-1.5%)	0.49% (-0.21-1.4%)	0.6 (-0.28-1.6)	77 (23-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	2005	Female	35 to 39	48% (32-63%)	52% (37-68%)	0.42 (0.23-0.68)	0.78% (0.084-1.5%)	0.62% (-0.32-1.6%)	0.76 (-0.39-2)	68 (7.1-130)
Djibouti	2005	Female	40 to 44	49% (35-66%)	51% (34-65%)	0.41 (0.21-0.68)	0.82% (0.17-1.6%)	0.84% (-0.095-1.9%)	1 (-0.11-2.2)	64 (14-120)
Djibouti	2005	Female	45 to 49	49% (32-66%)	51% (34-68%)	0.43 (0.21-0.7)	0.82% (-0.17-1.8%)	0.95% (-0.44-2.4%)	1.1 (-0.53-2.7)	53 (-11-120)
Djibouti	2005	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.43 (0.22-0.71)	0.84% (-0.25-2%)	1.1% (-0.4-2.7%)	1.2 (-0.45-3)	46 (-14-110)
Djibouti	2005	Female	55 to 59	46% (31-62%)	54% (38-69%)	0.44 (0.23-0.74)	0.54% (-0.67-1.9%)	0.75% (-0.9-2.6%)	0.98 (-1.1-3.3)	30 (-35-100)
Djibouti	2005	Female	60 to 64	47% (31-63%)	53% (37-69%)	0.43 (0.22-0.75)	0.42% (-0.65-1.5%)	0.61% (-0.75-2.1%)	0.9 (-1.1-3.1)	22 (-34-81)
Djibouti	2005	Female	65 to 69	48% (33-64%)	52% (36-67%)	0.4 (0.21-0.67)	0.3% (-0.83-1.3%)	0.41% (-1-1.7%)	0.68 (-1.7-2.9)	14 (-38-64)
Djibouti	2005	Female	70 to 74	47% (32-62%)	53% (38-68%)	0.36 (0.18-0.64)	-0.16% (-1.3-0.79%)	-0.21% (-1.5-0.99%)	-0.37 (-2.9-1.8)	-6.3 (-52-33)
Djibouti	2005	Female	75 to 79	44% (28-61%)	56% (39-72%)	0.31 (0.14-0.56)	-0.18% (-1.2-0.7%)	-0.19% (-1.4-0.88%)	-0.32 (-2.4-1.6)	-5.1 (-34-20)
Djibouti	2005	Female	80 to 84	43% (26-61%)	57% (39-74%)	0.29 (0.13-0.5)	-0.45% (-1.4-0.3%)	-0.51% (-1.7-0.4%)	-0.65 (-2.2-0.51)	-7.2 (-23-4.9)
Djibouti	2005	Female	85 to 89	43% (27-58%)	57% (42-73%)	0.27 (0.13-0.5)	-0.53% (-1.5-0.17%)	-0.63% (-1.8-0.23%)	-0.55 (-1.6-0.19)	-4.1 (-12-1.3)
Djibouti	2005	Female	90 to 94	45% (30-59%)	55% (41-70%)	0.27 (0.12-0.49)	-0.61% (-1.5-0.072%)	-0.77% (-1.9-0.053%)	-0.31 (-0.74-0.02)	-1.5 (-3.6-0.16)
Djibouti	2005	Female	95 plus	47% (29-65%)	53% (35-71%)	0.26 (0.11-0.52)	-0.22% (-1.1-0.46%)	-0.34% (-1.4-0.5%)	-0.039 (-0.16-0.054)	-0.098 (-0.48-0.22)
Djibouti	2005	Male	All Ages	42% (36-50%)	58% (50-64%)	0.27 (0.2-0.36)	0.76% (0.14-1.4%)	0.74% (-0.23-1.7%)	30 (-9.4-69)	2000 (370-3700)
Djibouti	2005	Male	15 to 19	41% (28-57%)	59% (43-72%)	0.3 (0.16-0.48)	1.6% (0.63-2.7%)	1.2% (-0.37-2.9%)	0.88 (-0.3-2.3)	140 (54-250)
Djibouti	2005	Male	20 to 24	52% (37-66%)	48% (34-63%)	0.4 (0.22-0.64)	2.7% (0.84-4.8%)	1.6% (-1-4.2%)	1.6 (-1-4.3)	250 (74-460)
Djibouti	2005	Male	25 to 29	55% (40-70%)	45% (30-60%)	0.49 (0.28-0.74)	2.5% (0.35-4.6%)	1.6% (-1.1-4.2%)	2.2 (-1.6-6)	290 (43-530)
Djibouti	2005	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.54 (0.32-0.83)	2.2% (0.14-4.4%)	1.5% (-1-4.1%)	2.4 (-1.5-6.3)	240 (15-480)
Djibouti	2005	Male	35 to 39	59% (45-74%)	41% (26-55%)	0.58 (0.34-0.92)	2.2% (0.24-4.2%)	1.8% (-0.61-4.5%)	3 (-1.1-7.3)	230 (25-450)
Djibouti	2005	Male	40 to 44	60% (45-75%)	40% (25-55%)	0.59 (0.34-0.95)	2.1% (0.14-4.1%)	2% (-0.46-4.6%)	3.6 (-0.81-8.2)	220 (16-440)
Djibouti	2005	Male	45 to 49	61% (45-75%)	39% (25-55%)	0.59 (0.33-0.94)	2.1% (0.05-4.1%)	2.1% (-0.48-4.7%)	3.4 (-0.82-7.8)	180 (4-360)
Djibouti	2005	Male	50 to 54	61% (45-75%)	39% (25-55%)	0.58 (0.32-0.93)	2% (-0.19-4.1%)	2.1% (-0.67-4.7%)	3.1 (-0.94-6.7)	130 (-13-270)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	2005	Male	55 to 59	60% (45-74%)	40% (26-55%)	0.57 (0.31-0.91)	1.7% (-0.43-3.7%)	1.9% (-0.91-4.4%)	3.2 (-1.6-7.8)	120 (-29-260)
Djibouti	2005	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.52 (0.27-0.89)	1.5% (-0.16-3%)	1.6% (-0.5-3.4%)	3 (-0.96-6.9)	92 (-11-200)
Djibouti	2005	Male	65 to 69	57% (42-73%)	43% (27-58%)	0.43 (0.24-0.71)	0.72% (-0.67-2%)	0.75% (-0.99-2.4%)	1.5 (-2-4.7)	39 (-36-110)
Djibouti	2005	Male	70 to 74	56% (41-71%)	44% (29-59%)	0.34 (0.19-0.59)	0.43% (-0.81-1.5%)	0.45% (-1-1.7%)	0.91 (-2.1-3.6)	19 (-35-66)
Djibouti	2005	Male	75 to 79	54% (40-70%)	46% (30-60%)	0.29 (0.14-0.51)	0.36% (-0.77-1.3%)	0.38% (-0.97-1.5%)	0.66 (-1.7-2.6)	11 (-22-38)
Djibouti	2005	Male	80 to 84	52% (36-69%)	48% (31-64%)	0.26 (0.13-0.44)	0.22% (-0.75-1%)	0.18% (-0.96-1.1%)	0.22 (-1.1-1.3)	3.1 (-11-14)
Djibouti	2005	Male	85 to 89	50% (34-65%)	50% (35-66%)	0.23 (0.12-0.39)	0.24% (-0.56-0.95%)	0.19% (-0.76-1%)	0.13 (-0.49-0.67)	1.4 (-3.1-5.3)
Djibouti	2005	Male	90 to 94	51% (35-65%)	49% (35-65%)	0.21 (0.099-0.37)	0.1% (-0.51-0.62%)	-0.0014% (-0.74-0.6%)	-0.00034 (-0.19-0.15)	0.15 (-0.77-0.94)
Djibouti	2005	Male	95 plus	52% (33-71%)	48% (29-67%)	0.19 (0.081-0.36)	0.25% (-0.51-0.89%)	0.11% (-0.8-0.89%)	0.0065 (-0.047-0.052)	0.058 (-0.12-0.21)
Djibouti	2010	Female	All Ages	34% (29-41%)	66% (59-71%)	0.24 (0.17-0.35)	0.31% (0.0055-0.59%)	0.19% (-0.39-0.76%)	6.4 (-13-26)	640 (10-1300)
Djibouti	2010	Female	15 to 19	34% (22-50%)	66% (50-78%)	0.27 (0.13-0.49)	0.66% (0.24-1.2%)	0.42% (-0.35-1.3%)	0.2 (-0.19-0.68)	48 (17-86)
Djibouti	2010	Female	20 to 24	41% (25-57%)	59% (43-75%)	0.37 (0.18-0.65)	1.1% (0.53-1.8%)	0.47% (-0.39-1.5%)	0.34 (-0.29-1.1)	97 (44-160)
Djibouti	2010	Female	25 to 29	43% (28-60%)	57% (40-72%)	0.44 (0.23-0.73)	1% (0.44-1.7%)	0.53% (-0.14-1.5%)	0.52 (-0.14-1.4)	93 (41-150)
Djibouti	2010	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.46 (0.24-0.78)	0.82% (0.29-1.5%)	0.5% (-0.11-1.4%)	0.7 (-0.17-1.8)	92 (33-160)
Djibouti	2010	Female	35 to 39	48% (32-63%)	52% (37-68%)	0.44 (0.24-0.74)	0.78% (0.17-1.5%)	0.63% (-0.16-1.6%)	0.86 (-0.23-2.1)	77 (17-150)
Djibouti	2010	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.43 (0.21-0.75)	0.82% (0.22-1.5%)	0.84% (-0.0019-1.8%)	1.1 (-0.0027-2.4)	73 (21-130)
Djibouti	2010	Female	45 to 49	49% (32-65%)	51% (35-68%)	0.44 (0.22-0.77)	0.84% (-0.056-1.8%)	0.99% (-0.22-2.4%)	1.3 (-0.29-3)	65 (-4.4-140)
Djibouti	2010	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.45 (0.23-0.78)	0.87% (-0.15-2%)	1.1% (-0.27-2.7%)	1.5 (-0.34-3.8)	58 (-9.7-140)
Djibouti	2010	Female	55 to 59	46% (30-62%)	54% (38-70%)	0.46 (0.22-0.84)	0.57% (-0.65-1.8%)	0.81% (-0.83-2.5%)	1.1 (-1.1-3.4)	33 (-36-110)
Djibouti	2010	Female	60 to 64	47% (31-64%)	53% (36-69%)	0.45 (0.21-0.85)	0.42% (-0.66-1.6%)	0.63% (-0.79-2.2%)	1.1 (-1.3-3.8)	26 (-39-95)
Djibouti	2010	Female	65 to 69	48% (33-63%)	52% (37-67%)	0.42 (0.21-0.74)	0.29% (-0.92-1.3%)	0.41% (-1.1-1.8%)	0.79 (-2.1-3.3)	16 (-49-72)
Djibouti	2010	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.37 (0.18-0.66)	-0.21% (-1.3-0.72%)	-0.27% (-1.6-0.9%)	-0.54 (-3.4-1.9)	-9.8 (-60-34)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	2010	Female	75 to 79	45% (29-61%)	55% (39-71%)	0.33 (0.14-0.62)	-0.23% (-1.3-0.71%)	-0.25% (-1.5-0.88%)	-0.49 (-3.3-1.8)	-7.9 (-45-24)
Djibouti	2010	Female	80 to 84	43% (27-60%)	57% (40-73%)	0.3 (0.13-0.57)	-0.49% (-1.5-0.31%)	-0.55% (-1.7-0.4%)	-0.84 (-2.7-0.65)	-9.3 (-29-5.9)
Djibouti	2010	Female	85 to 89	43% (27-58%)	57% (42-73%)	0.28 (0.13-0.54)	-0.57% (-1.5-0.16%)	-0.67% (-1.8-0.21%)	-0.72 (-2-0.24)	-5.4 (-14-1.6)
Djibouti	2010	Female	90 to 94	45% (29-59%)	55% (41-71%)	0.28 (0.12-0.56)	-0.67% (-1.6-0.036%)	-0.84% (-2-0.019%)	-0.41 (-0.98-0.0099)	-2 (-4.8-0.11)
Djibouti	2010	Female	95 plus	47% (29-65%)	53% (35-71%)	0.27 (0.11-0.55)	-0.24% (-1.2-0.5%)	-0.38% (-1.5-0.51%)	-0.057 (-0.23-0.079)	-0.14 (-0.71-0.3)
Djibouti	2010	Male	All Ages	43% (37-50%)	57% (50-63%)	0.3 (0.2-0.44)	0.94% (0.24-1.6%)	0.91% (-0.15-2%)	36 (-6.7-79)	2300 (560-4200)
Djibouti	2010	Male	15 to 19	40% (27-55%)	60% (45-73%)	0.31 (0.16-0.54)	1.7% (0.71-2.8%)	1.2% (-0.27-2.9%)	0.93 (-0.21-2.3)	150 (60-260)
Djibouti	2010	Male	20 to 24	51% (37-65%)	49% (35-63%)	0.42 (0.22-0.71)	3% (1.2-5.2%)	1.8% (-0.68-4.5%)	2.1 (-0.79-5.4)	330 (120-590)
Djibouti	2010	Male	25 to 29	55% (40-70%)	45% (30-60%)	0.5 (0.28-0.79)	3% (0.67-5.2%)	2% (-1-4.8%)	2.4 (-1.2-6)	310 (72-540)
Djibouti	2010	Male	30 to 34	58% (42-73%)	42% (27-58%)	0.56 (0.32-0.88)	2.5% (0.43-4.6%)	1.8% (-0.72-4.5%)	3 (-1.3-7.7)	300 (51-560)
Djibouti	2010	Male	35 to 39	59% (45-74%)	41% (26-55%)	0.6 (0.34-1)	2.3% (0.37-4.4%)	2% (-0.45-4.7%)	3.5 (-0.77-8.1)	260 (40-500)
Djibouti	2010	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.62 (0.33-1)	2.3% (0.32-4.2%)	2.2% (-0.18-4.7%)	4.1 (-0.36-8.8)	250 (34-450)
Djibouti	2010	Male	45 to 49	60% (44-75%)	40% (25-56%)	0.61 (0.32-1)	2.2% (0.29-4.1%)	2.3% (-0.087-4.6%)	4.6 (-0.2-9.6)	230 (32-440)
Djibouti	2010	Male	50 to 54	60% (45-74%)	40% (26-55%)	0.6 (0.31-1)	2.2% (0.04-4.2%)	2.3% (-0.4-4.9%)	4.3 (-0.77-9.4)	190 (3.4-370)
Djibouti	2010	Male	55 to 59	60% (45-74%)	40% (26-55%)	0.59 (0.31-1)	1.9% (-0.21-4%)	2.1% (-0.61-4.8%)	3.4 (-0.99-8.1)	130 (-13-270)
Djibouti	2010	Male	60 to 64	59% (44-76%)	41% (24-56%)	0.53 (0.26-0.95)	1.6% (-0.055-3.2%)	1.7% (-0.31-3.7%)	3.7 (-0.66-8.2)	110 (-3.9-230)
Djibouti	2010	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.44 (0.23-0.77)	0.79% (-0.6-2.1%)	0.83% (-0.84-2.4%)	1.9 (-2.5-5.7)	48 (-37-130)
Djibouti	2010	Male	70 to 74	56% (41-70%)	44% (30-59%)	0.36 (0.18-0.62)	0.45% (-0.73-1.5%)	0.48% (-0.92-1.8%)	1.1 (-2.1-4.1)	22 (-36-75)
Djibouti	2010	Male	75 to 79	54% (40-70%)	46% (30-60%)	0.3 (0.15-0.55)	0.39% (-0.77-1.3%)	0.41% (-0.97-1.5%)	0.82 (-1.9-3.1)	13 (-25-44)
Djibouti	2010	Male	80 to 84	52% (35-69%)	48% (31-65%)	0.27 (0.13-0.48)	0.23% (-0.73-1.1%)	0.2% (-0.93-1.2%)	0.28 (-1.3-1.7)	4 (-13-18)
Djibouti	2010	Male	85 to 89	50% (33-64%)	50% (36-67%)	0.23 (0.12-0.42)	0.25% (-0.57-0.96%)	0.2% (-0.79-1%)	0.16 (-0.62-0.83)	1.7 (-4-6.7)
Djibouti	2010	Male	90 to 94	50% (35-66%)	50% (34-65%)	0.21 (0.097-0.4)	0.1% (-0.54-0.65%)	-0.0018% (-0.78-0.64%)	-0.00052 (-0.24-0.2)	0.19 (-1-1.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	2010	Male	95 plus	52% (33-71%)	48% (29-67%)	0.2 (0.078-0.4)	0.27% (-0.52-0.93%)	0.13% (-0.8-0.93%)	0.01 (-0.062-0.073)	0.082 (-0.16-0.29)
Djibouti	2016	Female	All Ages	35% (29-41%)	65% (59-71%)	0.27 (0.14-0.44)	0.43% (0.092-0.86%)	0.3% (-0.38-1%)	9.5 (-12-33)	790 (170-1600)
Djibouti	2016	Female	15 to 19	34% (22-50%)	66% (50-78%)	0.29 (0.12-0.59)	0.68% (0.29-1.2%)	0.42% (-0.26-1.3%)	0.18 (-0.11-0.58)	44 (18-81)
Djibouti	2016	Female	20 to 24	40% (25-57%)	60% (43-75%)	0.39 (0.17-0.78)	1.3% (0.69-2.1%)	0.59% (-0.23-1.8%)	0.32 (-0.14-0.99)	96 (49-160)
Djibouti	2016	Female	25 to 29	43% (27-59%)	57% (41-73%)	0.46 (0.22-0.88)	1.3% (0.67-2.1%)	0.75% (-0.072-2%)	0.63 (-0.058-1.6)	120 (63-190)
Djibouti	2016	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.49 (0.23-0.9)	1.1% (0.48-1.8%)	0.72% (-0.037-1.8%)	0.74 (-0.043-1.8)	98 (43-170)
Djibouti	2016	Female	35 to 39	48% (33-63%)	52% (37-67%)	0.47 (0.21-0.87)	0.95% (0.34-1.9%)	0.82% (-0.022-2.1%)	1.1 (-0.025-2.7)	97 (33-180)
Djibouti	2016	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.46 (0.2-0.9)	0.96% (0.36-1.8%)	1% (0.19-2.2%)	1.4 (0.25-3)	90 (34-170)
Djibouti	2016	Female	45 to 49	49% (32-64%)	51% (36-68%)	0.47 (0.2-0.89)	0.98% (0.15-2%)	1.2% (0.03-2.8%)	1.6 (0.034-3.8)	81 (12-170)
Djibouti	2016	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.47 (0.21-0.9)	1% (-0.013-2.2%)	1.4% (-0.12-3.1%)	2.1 (-0.17-4.9)	80 (-1-180)
Djibouti	2016	Female	55 to 59	46% (31-62%)	54% (38-69%)	0.48 (0.2-0.97)	0.71% (-0.5-2.2%)	1% (-0.66-3%)	1.6 (-1.5-5.1)	50 (-35-160)
Djibouti	2016	Female	60 to 64	46% (30-63%)	54% (37-70%)	0.48 (0.19-1)	0.56% (-0.5-2%)	0.82% (-0.57-2.6%)	1.4 (-0.91-4.6)	35 (-30-120)
Djibouti	2016	Female	65 to 69	48% (32-64%)	52% (36-68%)	0.45 (0.19-0.88)	0.43% (-0.72-1.6%)	0.59% (-0.84-2.1%)	1.3 (-1.8-4.6)	27 (-41-100)
Djibouti	2016	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.39 (0.16-0.79)	-0.1% (-1.2-0.97%)	-0.14% (-1.4-1.2%)	-0.34 (-3.4-2.9)	-5.5 (-63-52)
Djibouti	2016	Female	75 to 79	45% (29-61%)	55% (39-71%)	0.35 (0.13-0.73)	-0.13% (-1.2-0.79%)	-0.12% (-1.4-1%)	-0.28 (-3.3-2.4)	-5.1 (-48-32)
Djibouti	2016	Female	80 to 84	43% (27-61%)	57% (39-73%)	0.32 (0.13-0.69)	-0.42% (-1.5-0.33%)	-0.47% (-1.7-0.47%)	-0.87 (-3.1-0.87)	-9.8 (-33-8.3)
Djibouti	2016	Female	85 to 89	43% (27-58%)	57% (42-73%)	0.3 (0.12-0.64)	-0.52% (-1.5-0.18%)	-0.61% (-1.7-0.25%)	-0.8 (-2.3-0.31)	-6 (-17-2)
Djibouti	2016	Female	90 to 94	45% (29-60%)	55% (40-71%)	0.3 (0.11-0.62)	-0.63% (-1.6-0.072%)	-0.79% (-1.9-0.068%)	-0.49 (-1.2-0.044)	-2.4 (-5.9-0.28)
Djibouti	2016	Female	95 plus	47% (29-65%)	53% (35-71%)	0.29 (0.099-0.65)	-0.18% (-1.1-0.52%)	-0.31% (-1.5-0.55%)	-0.06 (-0.27-0.11)	-0.14 (-0.83-0.41)
Djibouti	2016	Male	All Ages	44% (38-51%)	56% (49-62%)	0.33 (0.18-0.58)	1.3% (0.42-2.2%)	1.2% (-0.043-2.5%)	45 (-1.4-93)	2800 (960-4900)
Djibouti	2016	Male	15 to 19	40% (27-56%)	60% (44-73%)	0.33 (0.14-0.67)	1.7% (0.83-2.9%)	1.2% (-0.056-3%)	0.88 (-0.035-2.1)	140 (64-240)
Djibouti	2016	Male	20 to 24	51% (37-66%)	49% (34-63%)	0.44 (0.2-0.83)	3.3% (1.5-5.5%)	2.1% (-0.39-4.9%)	2.1 (-0.4-5.3)	330 (140-580)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Djibouti	2016	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.54 (0.27-0.99)	3.5% (1.2-6%)	2.5% (-0.6-5.8%)	3.3 (-0.82-7.8)	410 (140-710)
Djibouti	2016	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.6 (0.3-1.1)	3.1% (0.99-5.7%)	2.4% (-0.43-5.7%)	3.3 (-0.6-7.7)	330 (100-610)
Djibouti	2016	Male	35 to 39	59% (45-74%)	41% (26-55%)	0.64 (0.3-1.2)	2.8% (0.71-5.2%)	2.5% (-0.2-5.7%)	4.3 (-0.35-9.9)	330 (80-620)
Djibouti	2016	Male	40 to 44	60% (45-75%)	40% (25-55%)	0.66 (0.3-1.2)	2.7% (0.74-4.9%)	2.7% (0.22-5.5%)	4.8 (0.37-9.8)	290 (76-530)
Djibouti	2016	Male	45 to 49	60% (45-76%)	40% (24-55%)	0.65 (0.29-1.3)	2.6% (0.65-4.8%)	2.7% (0.25-5.6%)	5.2 (0.46-11)	260 (69-490)
Djibouti	2016	Male	50 to 54	61% (45-75%)	39% (25-55%)	0.64 (0.29-1.2)	2.5% (0.4-4.7%)	2.7% (0.054-5.6%)	5.9 (0.12-13)	250 (42-500)
Djibouti	2016	Male	55 to 59	60% (44-74%)	40% (26-56%)	0.63 (0.28-1.2)	2.2% (0.19-4.4%)	2.5% (-0.096-5.3%)	5.3 (-0.2-12)	190 (17-400)
Djibouti	2016	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.57 (0.23-1.1)	1.8% (0.26-3.6%)	2% (0.073-4.2%)	4.2 (0.16-9)	130 (18-260)
Djibouti	2016	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.47 (0.2-0.89)	0.95% (-0.33-2.2%)	1% (-0.58-2.5%)	2.6 (-1.4-6.4)	65 (-22-150)
Djibouti	2016	Male	70 to 74	56% (40-70%)	44% (30-60%)	0.38 (0.16-0.73)	0.54% (-0.52-1.6%)	0.6% (-0.7-1.8%)	1.6 (-1.9-4.9)	31 (-30-92)
Djibouti	2016	Male	75 to 79	54% (39-70%)	46% (30-61%)	0.32 (0.13-0.66)	0.46% (-0.6-1.4%)	0.5% (-0.82-1.6%)	1.2 (-1.9-3.8)	18 (-24-55)
Djibouti	2016	Male	80 to 84	52% (35-69%)	48% (31-65%)	0.28 (0.12-0.59)	0.27% (-0.64-1%)	0.25% (-0.78-1.1%)	0.42 (-1.4-2)	5.7 (-14-22)
Djibouti	2016	Male	85 to 89	50% (34-65%)	50% (35-66%)	0.25 (0.1-0.51)	0.28% (-0.54-0.93%)	0.23% (-0.74-1%)	0.23 (-0.76-1)	2.4 (-4.6-8)
Djibouti	2016	Male	90 to 94	51% (35-66%)	49% (34-65%)	0.23 (0.095-0.45)	0.11% (-0.57-0.65%)	0.0065% (-0.78-0.65%)	0.0024 (-0.31-0.26)	0.26 (-1.3-1.5)
Djibouti	2016	Male	95 plus	53% (33-71%)	47% (29-67%)	0.21 (0.072-0.46)	0.29% (-0.47-0.95%)	0.16% (-0.76-0.95%)	0.015 (-0.074-0.093)	0.11 (-0.18-0.37)
Eritrea	1990	Female	All Ages	22% (17-27%)	78% (73-83%)	0.29 (0.19-0.44)	0.6% (0.31-1%)	0.93% (0.37-1.8%)	170 (67-330)	7600 (4000-13000)
Eritrea	1990	Female	15 to 19	22% (13-36%)	78% (64-87%)	0.34 (0.15-0.65)	0.69% (0.33-1.3%)	0.62% (0.12-1.6%)	2.3 (0.47-5.8)	290 (130-550)
Eritrea	1990	Female	20 to 24	28% (15-44%)	72% (56-85%)	0.49 (0.22-0.89)	1.4% (0.67-2.6%)	1.2% (0.21-2.9%)	5.6 (1-13)	610 (280-1200)
Eritrea	1990	Female	25 to 29	30% (17-46%)	70% (54-83%)	0.59 (0.28-1.1)	1.8% (0.88-3.3%)	1.8% (0.57-3.9%)	8.7 (2.8-19)	740 (350-1400)
Eritrea	1990	Female	30 to 34	33% (18-50%)	67% (50-82%)	0.64 (0.3-1.2)	1.9% (0.9-3.6%)	2.1% (0.73-4.3%)	11 (3.8-22)	760 (340-1400)
Eritrea	1990	Female	35 to 39	36% (21-55%)	64% (45-79%)	0.63 (0.3-1.2)	2.1% (0.93-4.1%)	2.4% (0.92-4.9%)	14 (5.2-29)	820 (360-1600)
Eritrea	1990	Female	40 to 44	38% (23-56%)	62% (44-77%)	0.64 (0.3-1.2)	2.2% (1-4.1%)	2.6% (1.1-5%)	18 (7-34)	890 (390-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	1990	Female	45 to 49	38% (21-55%)	62% (45-79%)	0.65 (0.28-1.3)	2.4% (1.1-4.7%)	2.9% (1.2-5.7%)	20 (8.2-40)	870 (380-1700)
Eritrea	1990	Female	50 to 54	36% (20-56%)	64% (44-80%)	0.64 (0.27-1.2)	2.5% (1-5.1%)	2.9% (1.1-6%)	23 (8.8-49)	870 (340-1800)
Eritrea	1990	Female	55 to 59	35% (20-55%)	65% (45-80%)	0.63 (0.26-1.2)	2.2% (0.63-4.8%)	2.5% (0.72-5.7%)	22 (5.9-51)	690 (200-1600)
Eritrea	1990	Female	60 to 64	35% (19-55%)	65% (45-81%)	0.62 (0.25-1.2)	1.9% (0.44-4.6%)	2.2% (0.51-5.3%)	19 (4.5-47)	510 (120-1300)
Eritrea	1990	Female	65 to 69	37% (21-55%)	63% (45-79%)	0.58 (0.24-1.1)	1.6% (0.28-3.9%)	1.8% (0.28-4.5%)	16 (2.6-41)	370 (60-900)
Eritrea	1990	Female	70 to 74	35% (20-53%)	65% (47-80%)	0.5 (0.2-1)	0.93% (-0.22-3.1%)	1% (-0.31-3.4%)	8.1 (-2.6-28)	150 (-37-510)
Eritrea	1990	Female	75 to 79	33% (18-51%)	67% (49-82%)	0.43 (0.15-0.88)	0.69% (-0.37-2.2%)	0.76% (-0.45-2.5%)	4.6 (-2.7-15)	66 (-35-210)
Eritrea	1990	Female	80 to 84	31% (16-49%)	69% (51-84%)	0.39 (0.14-0.79)	0.32% (-0.48-1.4%)	0.35% (-0.54-1.6%)	1.3 (-2.1-6)	14 (-21-63)
Eritrea	1990	Female	85 to 89	31% (17-48%)	69% (52-83%)	0.37 (0.15-0.77)	0.11% (-0.58-0.98%)	0.11% (-0.66-1.1%)	0.17 (-1.1-7)	1.5 (-7.6-13)
Eritrea	1990	Female	90 to 94	34% (19-51%)	66% (49-81%)	0.38 (0.15-0.78)	-0.063% (-0.72-0.72%)	-0.12% (-0.89-0.78%)	-0.053 (-0.39-0.35)	-0.16 (-1.9-1.9)
Eritrea	1990	Female	95 plus	38% (20-60%)	62% (40-80%)	0.42 (0.15-0.88)	0.13% (-0.69-0.97%)	0.072% (-0.87-1%)	0.0078 (-0.096-0.11)	0.057 (-0.29-0.42)
Eritrea	1990	Male	All Ages	31% (25-39%)	69% (61-75%)	0.42 (0.28-0.61)	1.2% (0.66-1.9%)	1.8% (0.79-3%)	350 (160-600)	18000 (9700-28000)
Eritrea	1990	Male	15 to 19	28% (17-45%)	72% (55-83%)	0.45 (0.22-0.8)	1.8% (0.91-3.2%)	1.7% (0.53-3.5%)	8 (2.5-17)	860 (440-1500)
Eritrea	1990	Male	20 to 24	40% (24-58%)	60% (42-76%)	0.65 (0.32-1.1)	3.6% (1.6-6.4%)	3.1% (0.8-6.5%)	19 (4.8-39)	1800 (820-3200)
Eritrea	1990	Male	25 to 29	46% (28-62%)	54% (38-72%)	0.82 (0.42-1.4)	4.7% (2.1-8.2%)	4.6% (1.4-9%)	27 (8.2-54)	2200 (960-3800)
Eritrea	1990	Male	30 to 34	49% (30-67%)	51% (33-70%)	0.95 (0.48-1.6)	5.3% (2.4-9%)	5.5% (1.9-10%)	34 (11-64)	2300 (990-4000)
Eritrea	1990	Male	35 to 39	52% (34-69%)	48% (31-66%)	1 (0.53-1.8)	5.6% (2.5-9.8%)	6.1% (2.3-11%)	40 (15-74)	2300 (1000-4000)
Eritrea	1990	Male	40 to 44	53% (36-71%)	47% (29-64%)	1.1 (0.54-1.9)	5.5% (2.4-9.3%)	6.1% (2.4-11%)	49 (19-90)	2500 (1000-4300)
Eritrea	1990	Male	45 to 49	54% (34-71%)	46% (29-66%)	1.1 (0.53-1.9)	5.1% (2.3-9.1%)	5.7% (2.4-10%)	47 (19-86)	2000 (900-3600)
Eritrea	1990	Male	50 to 54	54% (35-72%)	46% (28-65%)	1 (0.51-1.9)	4.9% (2.2-8.8%)	5.4% (2.2-9.9%)	45 (18-82)	1700 (710-3000)
Eritrea	1990	Male	55 to 59	53% (35-72%)	47% (28-65%)	1 (0.47-1.8)	4.2% (1.7-7.6%)	4.7% (1.7-8.8%)	37 (13-69)	1200 (470-2200)
Eritrea	1990	Male	60 to 64	51% (33-70%)	49% (30-67%)	0.87 (0.39-1.6)	3.2% (1.2-6.2%)	3.6% (1.2-7%)	26 (8.9-51)	720 (270-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	1990	Male	65 to 69	49% (31-67%)	51% (33-69%)	0.7 (0.33-1.3)	1.9% (0.47-4%)	2.1% (0.44-4.5%)	12 (2.6-27)	290 (72-610)
Eritrea	1990	Male	70 to 74	46% (29-64%)	54% (36-71%)	0.55 (0.24-1.1)	1.1% (0.12-2.8%)	1.3% (0.076-3%)	5.6 (0.35-13)	100 (10-240)
Eritrea	1990	Male	75 to 79	44% (27-62%)	56% (38-73%)	0.45 (0.19-0.86)	0.84% (0.01-2%)	0.91% (-0.049-2.2%)	2.4 (-0.14-6)	35 (0.42-85)
Eritrea	1990	Male	80 to 84	41% (25-60%)	59% (40-75%)	0.38 (0.16-0.71)	0.61% (-0.023-1.4%)	0.62% (-0.11-1.6%)	0.73 (-0.13-1.9)	8.5 (-0.32-20)
Eritrea	1990	Male	85 to 89	39% (23-56%)	61% (44-77%)	0.32 (0.14-0.62)	0.49% (-0.041-1.1%)	0.47% (-0.15-1.1%)	0.23 (-0.069-0.55)	2 (-0.16-4.3)
Eritrea	1990	Male	90 to 94	39% (24-58%)	61% (42-76%)	0.29 (0.12-0.56)	0.26% (-0.18-0.68%)	0.17% (-0.36-0.67%)	0.026 (-0.054-0.099)	0.24 (-0.16-0.61)
Eritrea	1990	Male	95 plus	43% (22-65%)	57% (35-78%)	0.29 (0.099-0.6)	0.33% (-0.18-0.76%)	0.18% (-0.43-0.69%)	0.007 (-0.016-0.027)	0.05 (-0.028-0.12)
Eritrea	1995	Female	All Ages	21% (16-27%)	79% (73-84%)	0.39 (0.27-0.56)	0.9% (0.52-1.5%)	1.5% (0.73-2.7%)	230 (110-410)	9100 (5200-14000)
Eritrea	1995	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.49 (0.22-0.9)	0.91% (0.47-1.6%)	0.91% (0.31-2%)	2.9 (0.99-6.4)	350 (180-620)
Eritrea	1995	Female	20 to 24	27% (15-43%)	73% (57-85%)	0.69 (0.31-1.2)	1.7% (0.87-2.9%)	1.6% (0.5-3.2%)	5.6 (1.8-12)	590 (300-1000)
Eritrea	1995	Female	25 to 29	30% (16-47%)	70% (53-84%)	0.83 (0.41-1.5)	2.1% (1.2-3.6%)	2.1% (0.89-4.2%)	7.8 (3.2-15)	670 (360-1100)
Eritrea	1995	Female	30 to 34	33% (18-51%)	67% (49-82%)	0.91 (0.44-1.7)	2.3% (1.2-3.9%)	2.5% (1.1-4.7%)	9.7 (4.3-18)	700 (360-1100)
Eritrea	1995	Female	35 to 39	36% (21-55%)	64% (45-79%)	0.9 (0.44-1.6)	2.6% (1.4-4.4%)	3.1% (1.5-5.4%)	14 (6.7-24)	820 (430-1400)
Eritrea	1995	Female	40 to 44	38% (22-57%)	62% (43-78%)	0.92 (0.44-1.6)	2.9% (1.5-4.9%)	3.5% (1.7-6.1%)	18 (8.2-31)	880 (440-1500)
Eritrea	1995	Female	45 to 49	38% (21-56%)	62% (44-79%)	0.94 (0.44-1.8)	3.3% (1.6-5.9%)	4% (1.9-7.3%)	26 (12-47)	1100 (520-2000)
Eritrea	1995	Female	50 to 54	36% (20-56%)	64% (44-80%)	0.91 (0.4-1.7)	3.6% (1.7-6.7%)	4.3% (2-8.1%)	31 (14-58)	1100 (530-2200)
Eritrea	1995	Female	55 to 59	35% (20-55%)	65% (45-80%)	0.9 (0.4-1.7)	3.3% (1.2-6.6%)	3.9% (1.4-8%)	31 (11-63)	970 (350-2000)
Eritrea	1995	Female	60 to 64	35% (20-54%)	65% (46-80%)	0.89 (0.37-1.7)	3% (1-6.4%)	3.5% (1.2-7.3%)	29 (9.9-63)	780 (270-1700)
Eritrea	1995	Female	65 to 69	36% (21-54%)	64% (46-79%)	0.83 (0.36-1.5)	2.7% (0.87-5.7%)	3.1% (0.96-6.5%)	26 (8-55)	580 (190-1200)
Eritrea	1995	Female	70 to 74	35% (21-52%)	65% (48-79%)	0.72 (0.3-1.4)	1.8% (0.18-4.5%)	2% (0.14-5.1%)	16 (1.1-40)	290 (28-720)
Eritrea	1995	Female	75 to 79	33% (18-50%)	67% (50-82%)	0.61 (0.24-1.2)	1.4% (0.0022-3.6%)	1.6% (-0.0053-4%)	9.3 (-0.029-24)	130 (0.21-330)
Eritrea	1995	Female	80 to 84	31% (17-49%)	69% (51-83%)	0.55 (0.22-1.1)	0.9% (-0.16-2.5%)	1% (-0.19-2.8%)	3.7 (-0.67-10)	39 (-7.1-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	1995	Female	85 to 89	31% (17-48%)	69% (52-83%)	0.53 (0.21-1.1)	0.56% (-0.3-1.8%)	0.62% (-0.36-2.1%)	1 (-0.58-3.4)	7.7 (-4.2-25)
Eritrea	1995	Female	90 to 94	33% (19-51%)	67% (49-81%)	0.54 (0.22-1)	0.32% (-0.52-1.5%)	0.32% (-0.63-1.6%)	0.15 (-0.29-0.74)	0.85 (-1.3-3.9)
Eritrea	1995	Female	95 plus	38% (19-60%)	62% (40-81%)	0.59 (0.21-1.3)	0.53% (-0.36-1.8%)	0.53% (-0.53-2%)	0.055 (-0.055-0.2)	0.21 (-0.15-0.72)
Eritrea	1995	Male	All Ages	29% (23-37%)	71% (63-77%)	0.55 (0.38-0.78)	1.7% (1-2.4%)	2.5% (1.4-3.8%)	380 (210-610)	19000 (12000-28000)
Eritrea	1995	Male	15 to 19	28% (17-44%)	72% (56-83%)	0.62 (0.31-1.1)	2.4% (1.4-4%)	2.4% (1-4.7%)	9.8 (4.1-19)	1000 (570-1700)
Eritrea	1995	Male	20 to 24	40% (24-58%)	60% (42-76%)	0.93 (0.45-1.6)	4.7% (2.6-7.6%)	4.4% (1.7-8.1%)	20 (8-37)	1900 (1000-3100)
Eritrea	1995	Male	25 to 29	45% (27-64%)	55% (36-73%)	1.2 (0.59-2)	5.9% (3.2-9.5%)	6% (2.5-11%)	26 (11-47)	2100 (1100-3300)
Eritrea	1995	Male	30 to 34	49% (29-68%)	51% (32-71%)	1.3 (0.72-2.3)	6.4% (3.5-10%)	6.9% (3.3-12%)	33 (15-56)	2200 (1200-3600)
Eritrea	1995	Male	35 to 39	51% (34-69%)	49% (31-66%)	1.5 (0.79-2.4)	6.9% (3.8-11%)	7.7% (3.8-13%)	41 (20-68)	2300 (1300-3700)
Eritrea	1995	Male	40 to 44	53% (36-71%)	47% (29-64%)	1.5 (0.8-2.6)	6.8% (3.6-11%)	7.8% (3.9-13%)	45 (22-74)	2200 (1200-3500)
Eritrea	1995	Male	45 to 49	53% (35-71%)	47% (29-65%)	1.5 (0.79-2.8)	6.5% (3.5-11%)	7.4% (3.8-13%)	52 (27-91)	2300 (1200-3900)
Eritrea	1995	Male	50 to 54	54% (34-72%)	46% (28-66%)	1.5 (0.76-2.6)	6.3% (3.2-11%)	7.3% (3.6-13%)	51 (24-89)	1900 (980-3400)
Eritrea	1995	Male	55 to 59	53% (35-72%)	47% (28-65%)	1.4 (0.69-2.5)	5.7% (2.7-9.7%)	6.5% (2.9-11%)	44 (19-79)	1500 (670-2600)
Eritrea	1995	Male	60 to 64	51% (33-71%)	49% (29-67%)	1.2 (0.58-2.2)	4.5% (2-8.1%)	5.1% (2.2-9.3%)	33 (14-60)	910 (400-1700)
Eritrea	1995	Male	65 to 69	49% (31-67%)	51% (33-69%)	1 (0.49-1.8)	2.8% (1.1-5.3%)	3.2% (1.2-6.2%)	17 (6.3-33)	390 (150-750)
Eritrea	1995	Male	70 to 74	46% (29-64%)	54% (36-71%)	0.79 (0.37-1.4)	1.8% (0.47-3.9%)	2% (0.5-4.4%)	8 (1.9-18)	150 (37-320)
Eritrea	1995	Male	75 to 79	44% (27-62%)	56% (38-73%)	0.64 (0.29-1.2)	1.3% (0.31-2.8%)	1.5% (0.31-3.2%)	3.3 (0.67-7.2)	48 (11-100)
Eritrea	1995	Male	80 to 84	41% (24-61%)	59% (39-76%)	0.55 (0.24-0.98)	0.98% (0.23-2.1%)	1% (0.18-2.3%)	1 (0.17-2.2)	11 (2.7-24)
Eritrea	1995	Male	85 to 89	38% (23-55%)	62% (45-77%)	0.46 (0.2-0.86)	0.75% (0.17-1.5%)	0.76% (0.09-1.6%)	0.35 (0.042-0.73)	3 (0.66-5.8)
Eritrea	1995	Male	90 to 94	39% (24-55%)	61% (45-76%)	0.41 (0.18-0.78)	0.43% (-0.096-0.99%)	0.35% (-0.27-1%)	0.049 (-0.037-0.14)	0.36 (-0.081-0.82)
Eritrea	1995	Male	95 plus	42% (22-64%)	58% (36-78%)	0.41 (0.15-0.85)	0.51% (-0.035-1.2%)	0.36% (-0.32-1.1%)	0.012 (-0.011-0.035)	0.066 (-0.0046-0.15)
Eritrea	2000	Female	All Ages	21% (17-28%)	79% (72-83%)	0.45 (0.31-0.64)	1.2% (0.7-1.9%)	2% (1-3.4%)	300 (150-530)	12000 (6700-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	2000	Female	15 to 19	22% (13-35%)	78% (65-87%)	0.53 (0.25-0.94)	0.97% (0.51-1.7%)	0.98% (0.35-2%)	4.2 (1.4-8.6)	480 (250-820)
Eritrea	2000	Female	20 to 24	27% (15-42%)	73% (58-85%)	0.75 (0.36-1.3)	1.7% (0.98-2.9%)	1.6% (0.63-3.2%)	7.4 (2.8-15)	800 (430-1300)
Eritrea	2000	Female	25 to 29	29% (16-45%)	71% (55-84%)	0.92 (0.45-1.6)	2.1% (1.2-3.6%)	2.2% (0.93-4%)	9.6 (4.1-18)	820 (450-1300)
Eritrea	2000	Female	30 to 34	32% (18-50%)	68% (50-82%)	1 (0.48-1.8)	2.4% (1.3-4.1%)	2.6% (1.2-5%)	12 (5.2-22)	820 (430-1400)
Eritrea	2000	Female	35 to 39	36% (20-55%)	64% (45-80%)	0.99 (0.5-1.7)	2.8% (1.5-4.7%)	3.3% (1.6-5.8%)	16 (7.6-28)	920 (490-1500)
Eritrea	2000	Female	40 to 44	38% (22-57%)	62% (43-78%)	1 (0.48-1.7)	3.1% (1.6-5.2%)	3.8% (1.8-6.6%)	20 (9.6-35)	1000 (510-1700)
Eritrea	2000	Female	45 to 49	38% (21-56%)	62% (44-79%)	1 (0.46-1.9)	3.8% (1.8-6.6%)	4.7% (2.1-8.3%)	28 (12-51)	1200 (550-2200)
Eritrea	2000	Female	50 to 54	36% (21-56%)	64% (44-79%)	1 (0.45-1.8)	4.3% (2-7.9%)	5.2% (2.4-9.6%)	41 (18-77)	1500 (680-2800)
Eritrea	2000	Female	55 to 59	35% (20-55%)	65% (45-80%)	1 (0.43-2)	4.2% (1.7-8.1%)	5% (2-9.8%)	42 (16-84)	1300 (520-2600)
Eritrea	2000	Female	60 to 64	36% (19-55%)	64% (45-81%)	1 (0.42-2)	3.8% (1.4-7.8%)	4.5% (1.6-9.2%)	40 (14-82)	1100 (380-2200)
Eritrea	2000	Female	65 to 69	37% (21-54%)	63% (46-79%)	0.95 (0.4-1.7)	3.5% (1.3-7.1%)	4.1% (1.5-8.1%)	38 (13-76)	840 (300-1700)
Eritrea	2000	Female	70 to 74	35% (21-54%)	65% (46-79%)	0.81 (0.34-1.6)	2.5% (0.47-5.6%)	2.8% (0.49-6.5%)	24 (4.2-56)	440 (82-1000)
Eritrea	2000	Female	75 to 79	32% (18-51%)	68% (49-82%)	0.68 (0.25-1.4)	2% (0.39-4.7%)	2.3% (0.4-5.3%)	15 (2.7-37)	210 (39-520)
Eritrea	2000	Female	80 to 84	31% (16-50%)	69% (50-84%)	0.61 (0.24-1.2)	1.3% (0.053-3.3%)	1.5% (0.055-3.8%)	6.4 (0.25-16)	66 (2.7-160)
Eritrea	2000	Female	85 to 89	31% (17-48%)	69% (52-83%)	0.59 (0.25-1.2)	0.93% (-0.11-2.4%)	1.1% (-0.14-2.9%)	2.2 (-0.31-6)	17 (-1.9-44)
Eritrea	2000	Female	90 to 94	33% (19-50%)	67% (50-81%)	0.61 (0.25-1.2)	0.64% (-0.28-2.2%)	0.71% (-0.37-2.4%)	0.44 (-0.24-1.6)	2.4 (-1-8)
Eritrea	2000	Female	95 plus	38% (20-59%)	62% (41-80%)	0.66 (0.24-1.4)	0.91% (-0.12-2.6%)	0.97% (-0.22-2.9%)	0.13 (-0.03-0.39)	0.47 (-0.061-1.3)
Eritrea	2000	Male	All Ages	30% (24-38%)	70% (62-76%)	0.64 (0.45-0.88)	2.1% (1.4-3.1%)	3.1% (1.8-4.6%)	480 (290-750)	24000 (15000-34000)
Eritrea	2000	Male	15 to 19	28% (17-43%)	72% (57-83%)	0.68 (0.35-1.2)	2.2% (1.3-3.6%)	2.1% (0.93-3.9%)	14 (6.1-26)	1400 (790-2300)
Eritrea	2000	Male	20 to 24	39% (24-56%)	61% (44-76%)	1 (0.51-1.7)	4% (2.2-6.4%)	3.6% (1.5-6.3%)	30 (13-53)	2700 (1500-4400)
Eritrea	2000	Male	25 to 29	45% (28-63%)	55% (37-72%)	1.3 (0.69-2.2)	5.5% (3-8.6%)	5.3% (2.4-9.1%)	34 (16-59)	2700 (1500-4100)
Eritrea	2000	Male	30 to 34	49% (30-67%)	51% (33-70%)	1.5 (0.8-2.6)	6.2% (3.6-9.8%)	6.5% (3.3-11%)	38 (19-63)	2500 (1400-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	2000	Male	35 to 39	51% (34-69%)	49% (31-66%)	1.6 (0.89-2.8)	6.8% (3.9-11%)	7.6% (4-13%)	45 (24-77)	2600 (1500-4200)
Eritrea	2000	Male	40 to 44	53% (36-71%)	47% (29-64%)	1.7 (0.89-2.9)	7.1% (4-11%)	8.2% (4.3-13%)	52 (27-85)	2600 (1400-4100)
Eritrea	2000	Male	45 to 49	53% (34-72%)	47% (28-66%)	1.7 (0.85-3)	7.2% (4.1-11%)	8.3% (4.5-13%)	55 (29-94)	2400 (1300-4000)
Eritrea	2000	Male	50 to 54	54% (35-71%)	46% (29-65%)	1.7 (0.82-2.9)	7.3% (3.8-12%)	8.5% (4.3-14%)	67 (33-120)	2600 (1400-4300)
Eritrea	2000	Male	55 to 59	53% (35-72%)	47% (28-65%)	1.6 (0.77-2.8)	6.7% (3.3-11%)	7.8% (3.7-14%)	59 (27-100)	2000 (920-3300)
Eritrea	2000	Male	60 to 64	51% (32-70%)	49% (30-68%)	1.4 (0.63-2.5)	5.3% (2.4-9.4%)	6.1% (2.7-11%)	44 (19-83)	1200 (560-2300)
Eritrea	2000	Male	65 to 69	48% (30-66%)	52% (34-70%)	1.1 (0.54-1.9)	3.4% (1.5-6.4%)	3.9% (1.7-7.4%)	24 (10-45)	560 (240-1000)
Eritrea	2000	Male	70 to 74	45% (29-63%)	55% (37-71%)	0.86 (0.41-1.6)	2.2% (0.73-4.5%)	2.5% (0.82-5.2%)	12 (3.9-25)	220 (73-460)
Eritrea	2000	Male	75 to 79	43% (27-62%)	57% (38-73%)	0.71 (0.31-1.4)	1.7% (0.49-3.6%)	1.9% (0.51-4.1%)	5.5 (1.5-12)	79 (23-170)
Eritrea	2000	Male	80 to 84	40% (24-58%)	60% (42-76%)	0.6 (0.26-1.1)	1.3% (0.33-2.5%)	1.4% (0.3-2.9%)	1.7 (0.37-3.6)	19 (4.8-38)
Eritrea	2000	Male	85 to 89	38% (23-55%)	62% (45-77%)	0.5 (0.22-0.94)	0.96% (0.31-1.9%)	1% (0.24-2.1%)	0.58 (0.14-1.2)	4.8 (1.5-9.2)
Eritrea	2000	Male	90 to 94	39% (23-57%)	61% (43-77%)	0.45 (0.2-0.85)	0.6% (0.049-1.3%)	0.55% (-0.11-1.4%)	0.1 (-0.02-0.27)	0.68 (0.054-1.5)
Eritrea	2000	Male	95 plus	42% (21-64%)	58% (36-79%)	0.45 (0.16-0.93)	0.74% (0.077-1.6%)	0.64% (-0.15-1.6%)	0.027 (-0.0063-0.068)	0.13 (0.013-0.27)
Eritrea	2005	Female	All Ages	22% (17-29%)	78% (71-83%)	0.41 (0.29-0.57)	1.2% (0.67-1.9%)	1.9% (0.88-3.4%)	310 (140-560)	12000 (7000-20000)
Eritrea	2005	Female	15 to 19	22% (13-36%)	78% (64-87%)	0.44 (0.21-0.79)	1% (0.52-1.8%)	1.1% (0.32-2.4%)	4.1 (1.2-9.5)	500 (250-900)
Eritrea	2005	Female	20 to 24	28% (15-43%)	72% (57-85%)	0.63 (0.31-1.1)	1.8% (0.99-3.1%)	1.7% (0.59-3.5%)	8.6 (2.9-18)	980 (530-1600)
Eritrea	2005	Female	25 to 29	30% (17-46%)	70% (54-83%)	0.76 (0.38-1.3)	2.1% (1.2-3.6%)	2.2% (0.8-4.4%)	12 (4.6-24)	1100 (580-1800)
Eritrea	2005	Female	30 to 34	33% (18-51%)	67% (49-82%)	0.83 (0.4-1.5)	2.3% (1.2-3.9%)	2.5% (1-4.7%)	14 (5.9-27)	1000 (500-1700)
Eritrea	2005	Female	35 to 39	36% (21-54%)	64% (46-79%)	0.82 (0.41-1.5)	2.6% (1.3-4.5%)	3% (1.3-5.6%)	17 (7.3-31)	1000 (490-1700)
Eritrea	2005	Female	40 to 44	38% (22-56%)	62% (44-78%)	0.83 (0.4-1.5)	2.8% (1.3-4.9%)	3.4% (1.6-6.1%)	20 (9-38)	1000 (490-1900)
Eritrea	2005	Female	45 to 49	38% (21-56%)	62% (44-79%)	0.85 (0.36-1.6)	3.4% (1.5-6.2%)	4.2% (1.8-7.8%)	28 (12-53)	1200 (530-2200)
Eritrea	2005	Female	50 to 54	36% (20-56%)	64% (44-80%)	0.82 (0.36-1.5)	3.8% (1.7-7.1%)	4.7% (2-8.8%)	37 (15-70)	1400 (570-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	2005	Female	55 to 59	35% (20-54%)	65% (46-80%)	0.82 (0.35-1.6)	3.7% (1.4-7.4%)	4.5% (1.7-9%)	44 (16-94)	1400 (510-2900)
Eritrea	2005	Female	60 to 64	35% (20-54%)	65% (46-80%)	0.81 (0.33-1.6)	3.2% (1.2-6.9%)	3.9% (1.4-8.2%)	40 (14-88)	1100 (380-2300)
Eritrea	2005	Female	65 to 69	36% (21-54%)	64% (46-79%)	0.76 (0.32-1.3)	2.9% (1-6.1%)	3.4% (1.2-7.1%)	37 (13-79)	840 (280-1800)
Eritrea	2005	Female	70 to 74	35% (20-53%)	65% (47-80%)	0.65 (0.27-1.3)	2% (0.25-4.9%)	2.3% (0.25-5.6%)	24 (2.6-57)	430 (55-1000)
Eritrea	2005	Female	75 to 79	32% (18-51%)	68% (49-82%)	0.55 (0.21-1.1)	1.6% (0.11-3.8%)	1.8% (0.088-4.4%)	16 (0.78-39)	220 (14-530)
Eritrea	2005	Female	80 to 84	31% (16-49%)	69% (51-84%)	0.5 (0.19-0.96)	1% (-0.2-2.7%)	1.2% (-0.23-3.1%)	6.8 (-1.3-18)	69 (-13-190)
Eritrea	2005	Female	85 to 89	31% (17-49%)	69% (51-83%)	0.49 (0.2-1)	0.7% (-0.29-2.2%)	0.8% (-0.34-2.5%)	2.2 (-0.98-7.1)	17 (-7-52)
Eritrea	2005	Female	90 to 94	34% (19-52%)	66% (48-81%)	0.51 (0.2-1)	0.44% (-0.48-1.7%)	0.47% (-0.59-2.1%)	0.44 (-0.57-1.9)	2.4 (-2.6-9.8)
Eritrea	2005	Female	95 plus	39% (20-59%)	61% (41-80%)	0.55 (0.19-1.2)	0.77% (-0.24-2.4%)	0.82% (-0.38-2.8%)	0.17 (-0.078-0.57)	0.61 (-0.19-1.9)
Eritrea	2005	Male	All Ages	31% (25-39%)	69% (61-75%)	0.56 (0.39-0.76)	2.2% (1.4-3.2%)	3.1% (1.6-4.7%)	510 (270-780)	26000 (16000-37000)
Eritrea	2005	Male	15 to 19	28% (17-44%)	72% (56-83%)	0.56 (0.28-0.94)	2.5% (1.4-4.2%)	2.6% (1.1-4.8%)	14 (5.6-26)	1500 (810-2400)
Eritrea	2005	Male	20 to 24	39% (24-56%)	61% (44-76%)	0.81 (0.43-1.4)	4.9% (2.7-8%)	4.7% (1.7-8.8%)	35 (13-65)	3300 (1800-5500)
Eritrea	2005	Male	25 to 29	45% (27-62%)	55% (38-73%)	1 (0.53-1.7)	6.2% (3.3-10%)	6.4% (2.5-11%)	45 (17-79)	3500 (1800-5600)
Eritrea	2005	Male	30 to 34	49% (30-67%)	51% (33-70%)	1.2 (0.66-2)	6.4% (3.4-10%)	6.9% (3-12%)	43 (19-74)	3000 (1500-4600)
Eritrea	2005	Male	35 to 39	51% (34-69%)	49% (31-66%)	1.3 (0.72-2.3)	6.7% (3.5-11%)	7.5% (3.5-13%)	46 (22-81)	2700 (1400-4500)
Eritrea	2005	Male	40 to 44	52% (35-71%)	48% (29-65%)	1.4 (0.71-2.4)	6.6% (3.5-11%)	7.6% (3.7-13%)	50 (24-87)	2500 (1300-4200)
Eritrea	2005	Male	45 to 49	53% (34-71%)	47% (29-66%)	1.4 (0.67-2.4)	6.6% (3.5-11%)	7.7% (3.8-13%)	55 (27-94)	2400 (1300-4100)
Eritrea	2005	Male	50 to 54	54% (35-72%)	46% (28-65%)	1.3 (0.65-2.3)	6.6% (3.2-11%)	7.8% (3.6-13%)	59 (26-100)	2300 (1100-3800)
Eritrea	2005	Male	55 to 59	53% (34-72%)	47% (28-66%)	1.3 (0.64-2.3)	6.1% (2.9-10%)	7.1% (3.3-13%)	64 (29-110)	2100 (1000-3700)
Eritrea	2005	Male	60 to 64	51% (32-71%)	49% (29-68%)	1.1 (0.51-2)	4.6% (2.2-8.4%)	5.4% (2.3-9.9%)	47 (21-85)	1300 (610-2400)
Eritrea	2005	Male	65 to 69	48% (30-67%)	52% (33-70%)	0.88 (0.42-1.5)	2.9% (1.2-5.6%)	3.4% (1.3-6.5%)	26 (9.8-50)	600 (240-1100)
Eritrea	2005	Male	70 to 74	45% (29-63%)	55% (37-71%)	0.69 (0.32-1.3)	1.9% (0.54-3.8%)	2.1% (0.57-4.4%)	13 (3.5-27)	240 (70-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	2005	Male	75 to 79	43% (26-61%)	57% (39-74%)	0.56 (0.24-1)	1.4% (0.32-3%)	1.6% (0.31-3.5%)	6.4 (1.2-14)	92 (20-200)
Eritrea	2005	Male	80 to 84	40% (23-58%)	60% (42-77%)	0.47 (0.2-0.86)	1% (0.2-2.2%)	1.1% (0.13-2.5%)	2.2 (0.27-4.9)	25 (4.8-52)
Eritrea	2005	Male	85 to 89	38% (22-55%)	62% (45-78%)	0.4 (0.17-0.77)	0.83% (0.21-1.7%)	0.87% (0.12-1.9%)	0.68 (0.09-1.5)	5.6 (1.4-11)
Eritrea	2005	Male	90 to 94	38% (23-56%)	62% (44-77%)	0.36 (0.16-0.68)	0.52% (0.013-1.2%)	0.47% (-0.11-1.3%)	0.12 (-0.029-0.32)	0.78 (0.019-1.8)
Eritrea	2005	Male	95 plus	42% (22-64%)	58% (36-78%)	0.36 (0.13-0.77)	0.71% (0.0083-1.6%)	0.63% (-0.23-1.6%)	0.037 (-0.013-0.096)	0.17 (0.0019-0.37)
Eritrea	2010	Female	All Ages	23% (18-29%)	77% (71-82%)	0.35 (0.23-0.5)	1.2% (0.6-2%)	1.8% (0.69-3.3%)	310 (120-600)	12000 (6300-21000)
Eritrea	2010	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.37 (0.17-0.7)	0.91% (0.46-1.7%)	0.92% (0.19-2.2%)	3.2 (0.67-7.8)	420 (200-770)
Eritrea	2010	Female	20 to 24	27% (15-43%)	73% (57-85%)	0.53 (0.25-0.93)	1.7% (0.89-2.9%)	1.6% (0.37-3.5%)	7.8 (1.9-17)	960 (480-1600)
Eritrea	2010	Female	25 to 29	30% (16-46%)	70% (54-84%)	0.65 (0.32-1.2)	2.1% (1.1-3.7%)	2.2% (0.74-4.6%)	14 (4.6-28)	1200 (630-2100)
Eritrea	2010	Female	30 to 34	33% (18-51%)	67% (49-82%)	0.71 (0.35-1.3)	2.3% (1.1-4.1%)	2.5% (0.94-5.1%)	16 (5.9-34)	1200 (580-2200)
Eritrea	2010	Female	35 to 39	36% (20-55%)	64% (45-80%)	0.7 (0.32-1.3)	2.5% (1.1-4.6%)	3% (1.1-5.7%)	19 (6.4-36)	1100 (470-2000)
Eritrea	2010	Female	40 to 44	38% (23-57%)	62% (43-77%)	0.7 (0.31-1.3)	2.7% (1.2-4.8%)	3.3% (1.3-6.2%)	20 (7.8-39)	1000 (440-1900)
Eritrea	2010	Female	45 to 49	38% (21-56%)	62% (44-79%)	0.73 (0.3-1.4)	3.1% (1.3-6.1%)	3.9% (1.4-7.7%)	27 (9.6-54)	1200 (440-2300)
Eritrea	2010	Female	50 to 54	37% (21-57%)	63% (43-79%)	0.71 (0.3-1.3)	3.5% (1.3-6.9%)	4.3% (1.6-8.5%)	36 (13-74)	1300 (500-2700)
Eritrea	2010	Female	55 to 59	35% (20-55%)	65% (45-80%)	0.71 (0.31-1.4)	3.3% (1-7.2%)	4% (1.3-8.6%)	39 (12-88)	1200 (390-2800)
Eritrea	2010	Female	60 to 64	36% (21-54%)	64% (46-79%)	0.7 (0.3-1.4)	2.8% (0.86-6.3%)	3.4% (1.1-7.4%)	42 (13-94)	1100 (340-2500)
Eritrea	2010	Female	65 to 69	37% (21-55%)	63% (45-79%)	0.66 (0.28-1.2)	2.5% (0.72-5.4%)	3% (0.84-6.3%)	38 (11-84)	840 (250-1900)
Eritrea	2010	Female	70 to 74	35% (21-53%)	65% (47-79%)	0.57 (0.23-1.1)	1.6% (0.06-4.2%)	1.8% (0.016-4.9%)	23 (0.2-61)	420 (16-1100)
Eritrea	2010	Female	75 to 79	33% (19-51%)	67% (49-81%)	0.48 (0.17-0.98)	1.3% (-0.095-3.2%)	1.5% (-0.11-3.8%)	16 (-1.3-41)	220 (-16-570)
Eritrea	2010	Female	80 to 84	31% (17-49%)	69% (51-83%)	0.43 (0.16-0.87)	0.77% (-0.35-2.3%)	0.88% (-0.4-2.6%)	6.8 (-3.1-21)	69 (-30-210)
Eritrea	2010	Female	85 to 89	31% (17-49%)	69% (51-83%)	0.41 (0.17-0.86)	0.51% (-0.42-1.8%)	0.58% (-0.48-2.2%)	2.2 (-1.9-8.3)	16 (-13-60)
Eritrea	2010	Female	90 to 94	34% (20-52%)	66% (48-80%)	0.43 (0.17-0.87)	0.26% (-0.57-1.5%)	0.27% (-0.69-1.7%)	0.33 (-0.85-2.1)	1.9 (-4-11)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	2010	Female	95 plus	39% (20-60%)	61% (40-80%)	0.47 (0.16-1)	0.6% (-0.38-2%)	0.62% (-0.54-2.3%)	0.19 (-0.16-0.69)	0.7 (-0.44-2.4)
Eritrea	2010	Male	All Ages	31% (25-39%)	69% (61-75%)	0.48 (0.32-0.68)	2.2% (1.2-3.3%)	2.9% (1.4-4.7%)	510 (240-840)	26000 (15000-39000)
Eritrea	2010	Male	15 to 19	28% (17-43%)	72% (57-83%)	0.47 (0.23-0.84)	2.3% (1.2-3.8%)	2.2% (0.72-4.5%)	11 (3.7-21)	1200 (650-2100)
Eritrea	2010	Male	20 to 24	39% (23-56%)	61% (44-77%)	0.69 (0.34-1.2)	4.6% (2.3-7.6%)	4.2% (1.2-8%)	32 (9.6-63)	3200 (1600-5300)
Eritrea	2010	Male	25 to 29	45% (27-63%)	55% (37-73%)	0.88 (0.45-1.4)	5.9% (3-9.9%)	6% (2.1-11%)	48 (17-92)	4000 (2000-6600)
Eritrea	2010	Male	30 to 34	49% (30-67%)	51% (33-70%)	1 (0.56-1.7)	6.2% (3-10%)	6.7% (2.6-12%)	51 (19-92)	3500 (1700-5800)
Eritrea	2010	Male	35 to 39	51% (34-69%)	49% (31-66%)	1.1 (0.58-2.1)	6.4% (3.1-11%)	7.3% (3-13%)	49 (20-87)	2900 (1400-4800)
Eritrea	2010	Male	40 to 44	52% (36-71%)	48% (29-64%)	1.2 (0.58-2.1)	6.4% (3-11%)	7.4% (3-13%)	48 (19-88)	2500 (1100-4200)
Eritrea	2010	Male	45 to 49	53% (33-72%)	47% (28-67%)	1.2 (0.55-2.1)	6.2% (2.9-11%)	7.3% (3.1-13%)	52 (22-93)	2300 (1000-4000)
Eritrea	2010	Male	50 to 54	54% (35-72%)	46% (28-65%)	1.1 (0.54-2)	6.2% (2.7-11%)	7.3% (2.9-14%)	58 (23-110)	2200 (970-4100)
Eritrea	2010	Male	55 to 59	53% (34-71%)	47% (29-66%)	1.1 (0.53-2)	5.5% (2.5-10%)	6.5% (2.7-12%)	56 (24-100)	1800 (830-3300)
Eritrea	2010	Male	60 to 64	51% (32-71%)	49% (29-68%)	0.93 (0.44-1.8)	4.2% (1.9-7.9%)	4.8% (2.1-9.5%)	50 (21-98)	1400 (620-2700)
Eritrea	2010	Male	65 to 69	48% (30-66%)	52% (34-70%)	0.75 (0.36-1.3)	2.6% (0.88-5.1%)	3% (0.93-6%)	28 (9-56)	640 (220-1300)
Eritrea	2010	Male	70 to 74	45% (28-63%)	55% (37-72%)	0.58 (0.26-1.1)	1.6% (0.36-3.5%)	1.8% (0.38-4.1%)	14 (3-32)	260 (61-590)
Eritrea	2010	Male	75 to 79	43% (26-62%)	57% (38-74%)	0.48 (0.2-0.92)	1.3% (0.22-2.7%)	1.4% (0.19-3.1%)	7.4 (1.1-17)	110 (19-240)
Eritrea	2010	Male	80 to 84	40% (24-60%)	60% (40-76%)	0.4 (0.17-0.76)	0.87% (0.087-2%)	0.94% (0.0021-2.3%)	2.6 (0.0056-6.5)	29 (2.8-68)
Eritrea	2010	Male	85 to 89	38% (23-56%)	62% (44-77%)	0.34 (0.14-0.66)	0.74% (0.058-1.6%)	0.77% (-0.033-1.8%)	0.91 (-0.04-2.1)	7.5 (0.58-16)
Eritrea	2010	Male	90 to 94	39% (24-56%)	61% (44-76%)	0.31 (0.14-0.61)	0.44% (-0.066-1.1%)	0.39% (-0.2-1.1%)	0.12 (-0.063-0.37)	0.84 (-0.12-2)
Eritrea	2010	Male	95 plus	42% (21-63%)	58% (37-79%)	0.31 (0.11-0.66)	0.63% (-0.1-1.5%)	0.56% (-0.34-1.6%)	0.04 (-0.025-0.11)	0.18 (-0.03-0.43)
Eritrea	2016	Female	All Ages	23% (18-29%)	77% (71-82%)	0.38 (0.2-0.65)	1.5% (0.65-3%)	2.3% (0.65-4.8%)	400 (120-840)	15000 (6400-29000)
Eritrea	2016	Female	15 to 19	22% (13-36%)	78% (64-87%)	0.41 (0.15-0.85)	1% (0.48-2%)	1.1% (0.21-2.8%)	3.6 (0.66-9.2)	460 (200-890)
Eritrea	2016	Female	20 to 24	28% (15-43%)	72% (57-85%)	0.58 (0.23-1.2)	1.9% (0.94-3.5%)	1.9% (0.39-4.6%)	7.6 (1.6-18)	920 (430-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	2016	Female	25 to 29	30% (17-46%)	70% (54-83%)	0.7 (0.28-1.4)	2.5% (1.1-4.6%)	2.7% (0.74-6%)	15 (4.4-36)	1400 (640-2600)
Eritrea	2016	Female	30 to 34	33% (18-50%)	67% (50-82%)	0.76 (0.33-1.5)	2.7% (1.2-5%)	3.1% (1.1-6.6%)	23 (7.5-49)	1700 (730-3200)
Eritrea	2016	Female	35 to 39	36% (21-54%)	64% (46-79%)	0.75 (0.31-1.5)	3% (1.2-5.9%)	3.7% (1.2-7.6%)	29 (8.5-60)	1700 (650-3300)
Eritrea	2016	Female	40 to 44	38% (23-56%)	62% (44-77%)	0.76 (0.3-1.5)	3.1% (1.3-6%)	3.9% (1.4-8%)	28 (9.8-59)	1400 (580-2800)
Eritrea	2016	Female	45 to 49	38% (21-57%)	62% (43-79%)	0.79 (0.29-1.7)	3.6% (1.2-7.4%)	4.6% (1.5-9.6%)	34 (10-74)	1400 (490-3200)
Eritrea	2016	Female	50 to 54	36% (20-57%)	64% (43-80%)	0.76 (0.27-1.6)	3.9% (1.4-8.4%)	4.8% (1.6-10%)	42 (14-92)	1500 (530-3400)
Eritrea	2016	Female	55 to 59	35% (20-53%)	65% (47-80%)	0.75 (0.28-1.7)	3.6% (0.87-8.5%)	4.4% (1.1-10%)	46 (11-110)	1400 (340-3400)
Eritrea	2016	Female	60 to 64	36% (20-54%)	64% (46-80%)	0.75 (0.28-1.6)	3.1% (0.83-7.6%)	3.8% (1-9.2%)	45 (12-110)	1200 (320-2900)
Eritrea	2016	Female	65 to 69	37% (21-55%)	63% (45-79%)	0.71 (0.27-1.5)	2.9% (0.6-6.7%)	3.4% (0.71-7.7%)	50 (10-120)	1100 (240-2600)
Eritrea	2016	Female	70 to 74	36% (21-54%)	64% (46-79%)	0.61 (0.22-1.3)	2% (0.057-5.6%)	2.2% (0.0011-6.3%)	34 (0.016-100)	620 (18-1800)
Eritrea	2016	Female	75 to 79	33% (19-53%)	67% (47-81%)	0.52 (0.18-1.2)	1.6% (0.018-4.5%)	1.8% (0.028-5.3%)	24 (0.36-71)	340 (3.8-1000)
Eritrea	2016	Female	80 to 84	31% (16-50%)	69% (50-84%)	0.46 (0.16-1)	0.99% (-0.26-3.1%)	1.1% (-0.3-3.6%)	12 (-3.1-37)	120 (-32-380)
Eritrea	2016	Female	85 to 89	31% (17-49%)	69% (51-83%)	0.45 (0.15-1)	0.72% (-0.38-2.6%)	0.82% (-0.47-3%)	4.3 (-2.3-15)	31 (-17-110)
Eritrea	2016	Female	90 to 94	34% (19-51%)	66% (49-81%)	0.46 (0.17-1)	0.46% (-0.55-2.3%)	0.5% (-0.68-2.7%)	0.87 (-1.2-4.6)	4.7 (-5.4-23)
Eritrea	2016	Female	95 plus	38% (19-60%)	62% (40-81%)	0.51 (0.15-1.2)	0.82% (-0.38-3%)	0.88% (-0.51-3.4%)	0.35 (-0.2-1.3)	1.3 (-0.57-4.6)
Eritrea	2016	Male	All Ages	32% (26-40%)	68% (60-74%)	0.53 (0.28-0.91)	2.9% (1.4-5%)	3.7% (1.4-6.8%)	640 (240-1200)	32000 (15000-54000)
Eritrea	2016	Male	15 to 19	28% (17-44%)	72% (56-83%)	0.51 (0.21-1.1)	2.5% (1.3-4.4%)	2.6% (0.75-5.5%)	13 (3.5-28)	1400 (670-2500)
Eritrea	2016	Male	20 to 24	39% (24-56%)	61% (44-76%)	0.75 (0.3-1.5)	5.1% (2.2-9.2%)	4.8% (1.2-10%)	33 (7.8-71)	3200 (1400-5800)
Eritrea	2016	Male	25 to 29	45% (27-62%)	55% (38-73%)	0.95 (0.4-1.8)	6.5% (2.8-12%)	6.8% (1.9-14%)	56 (15-110)	4500 (2000-7900)
Eritrea	2016	Male	30 to 34	49% (30-68%)	51% (32-70%)	1.1 (0.52-2.1)	6.9% (2.9-12%)	7.7% (2.3-15%)	70 (21-140)	4800 (2100-8600)
Eritrea	2016	Male	35 to 39	52% (34-70%)	48% (30-66%)	1.2 (0.54-2.4)	7.1% (3-13%)	8.3% (2.8-15%)	73 (25-140)	4300 (1700-7500)
Eritrea	2016	Male	40 to 44	52% (35-71%)	48% (29-65%)	1.3 (0.55-2.4)	7% (2.9-12%)	8.4% (2.9-16%)	65 (22-120)	3300 (1300-6100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Eritrea	2016	Male	45 to 49	53% (34-71%)	47% (29-66%)	1.3 (0.51-2.5)	6.9% (2.8-13%)	8.2% (3-15%)	61 (21-120)	2700 (1100-5200)
Eritrea	2016	Male	50 to 54	54% (35-72%)	46% (28-65%)	1.2 (0.51-2.5)	6.8% (2.6-13%)	8.1% (2.9-16%)	67 (23-130)	2600 (990-4900)
Eritrea	2016	Male	55 to 59	53% (35-71%)	47% (29-65%)	1.2 (0.48-2.3)	6% (2.3-12%)	7.2% (2.6-14%)	67 (23-140)	2200 (830-4400)
Eritrea	2016	Male	60 to 64	51% (32-71%)	49% (29-68%)	1 (0.4-2.1)	4.6% (1.7-9.2%)	5.3% (2-11%)	55 (19-110)	1500 (600-3000)
Eritrea	2016	Male	65 to 69	49% (30-67%)	51% (33-70%)	0.81 (0.35-1.6)	2.9% (0.85-6.5%)	3.3% (0.9-7.5%)	39 (10-87)	900 (260-2000)
Eritrea	2016	Male	70 to 74	46% (29-64%)	54% (36-71%)	0.64 (0.24-1.4)	1.8% (0.43-4.5%)	2.1% (0.44-5.2%)	22 (4.4-54)	400 (93-990)
Eritrea	2016	Male	75 to 79	43% (27-62%)	57% (38-73%)	0.52 (0.19-1.1)	1.4% (0.27-3.7%)	1.6% (0.26-4.2%)	12 (1.7-31)	170 (31-430)
Eritrea	2016	Male	80 to 84	40% (24-60%)	60% (40-76%)	0.43 (0.17-0.89)	0.99% (0.11-2.5%)	1.1% (0.03-2.8%)	4.3 (0.12-12)	47 (5.4-120)
Eritrea	2016	Male	85 to 89	38% (22-55%)	62% (45-78%)	0.36 (0.14-0.78)	0.82% (0.1-2%)	0.87% (0.0094-2.2%)	1.5 (0.015-3.8)	12 (1.4-29)
Eritrea	2016	Male	90 to 94	38% (23-57%)	62% (43-77%)	0.33 (0.13-0.71)	0.52% (-0.053-1.4%)	0.48% (-0.22-1.5%)	0.25 (-0.11-0.82)	1.6 (-0.17-4.4)
Eritrea	2016	Male	95 plus	42% (21-64%)	58% (36-79%)	0.33 (0.11-0.77)	0.73% (-0.073-1.9%)	0.67% (-0.28-2%)	0.065 (-0.029-0.2)	0.28 (-0.028-0.74)
Ethiopia	1990	Female	All Ages	29% (25-34%)	71% (66-75%)	0.33 (0.29-0.37)	0.68% (0.083-1.3%)	1.1% (-0.011-2.3%)	4200 (-45-8900)	160000 (20000-320000)
Ethiopia	1990	Female	15 to 19	29% (21-38%)	71% (62-79%)	0.36 (0.24-0.52)	0.7% (0.046-1.5%)	0.69% (-0.18-1.8%)	55 (-15-140)	5400 (350-12000)
Ethiopia	1990	Female	20 to 24	37% (26-48%)	63% (52-74%)	0.49 (0.31-0.68)	1.2% (0.072-2.6%)	1.2% (-0.41-2.9%)	110 (-40-270)	10000 (540-21000)
Ethiopia	1990	Female	25 to 29	39% (28-51%)	61% (49-72%)	0.62 (0.43-0.86)	1.9% (0.46-3.6%)	2% (0.1-4.1%)	200 (9.3-390)	14000 (3300-26000)
Ethiopia	1990	Female	30 to 34	42% (30-54%)	58% (46-70%)	0.69 (0.47-0.96)	2.3% (0.6-4.2%)	2.5% (0.43-5%)	260 (44-540)	17000 (4500-32000)
Ethiopia	1990	Female	35 to 39	45% (33-56%)	55% (44-67%)	0.69 (0.46-0.95)	2.3% (0.36-4.6%)	2.6% (0.2-5.6%)	310 (24-660)	17000 (2600-34000)
Ethiopia	1990	Female	40 to 44	47% (36-57%)	53% (43-64%)	0.71 (0.49-0.95)	2.3% (0.47-4.3%)	2.7% (0.44-5.2%)	340 (53-660)	17000 (3200-31000)
Ethiopia	1990	Female	45 to 49	47% (35-58%)	53% (42-65%)	0.72 (0.5-0.99)	2.4% (0.015-5%)	2.9% (-0.09-6.1%)	390 (-13-820)	16000 (91-34000)
Ethiopia	1990	Female	50 to 54	45% (34-56%)	55% (44-66%)	0.69 (0.47-0.97)	2.9% (0.13-6.1%)	3.4% (0.081-7.3%)	520 (13-1100)	19000 (810-41000)
Ethiopia	1990	Female	55 to 59	43% (31-56%)	57% (44-69%)	0.71 (0.45-1)	2.8% (-0.059-6.3%)	3.3% (-0.046-7.4%)	610 (-7.9-1400)	19000 (-360-42000)
Ethiopia	1990	Female	60 to 64	43% (30-56%)	57% (44-70%)	0.71 (0.46-1)	2.3% (0.11-5.2%)	2.7% (0.15-6%)	520 (30-1100)	14000 (660-30000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	1990	Female	65 to 69	44% (33-55%)	56% (45-67%)	0.67 (0.45-0.93)	1.8% (-0.3-4.4%)	2.1% (-0.32-4.9%)	480 (-73-1200)	10000 (-1600-26000)
Ethiopia	1990	Female	70 to 74	42% (31-54%)	58% (46-69%)	0.58 (0.37-0.85)	0.9% (-1.1-3%)	1% (-1.2-3.4%)	220 (-270-740)	3900 (-4700-13000)
Ethiopia	1990	Female	75 to 79	39% (26-54%)	61% (46-74%)	0.48 (0.28-0.74)	0.62% (-1-2.4%)	0.7% (-1.1-2.7%)	130 (-210-510)	1800 (-3000-6900)
Ethiopia	1990	Female	80 to 84	38% (24-52%)	62% (48-76%)	0.43 (0.26-0.65)	0.18% (-1.2-1.6%)	0.22% (-1.4-1.8%)	26 (-160-210)	250 (-1700-2100)
Ethiopia	1990	Female	85 to 89	38% (25-51%)	62% (49-75%)	0.41 (0.24-0.63)	-0.0026% (-1.2-1.3%)	0.011% (-1.4-1.5%)	0.59 (-74-80)	-0.52 (-530-560)
Ethiopia	1990	Female	90 to 94	39% (27-53%)	61% (47-73%)	0.4 (0.23-0.62)	-0.19% (-1.3-0.97%)	-0.23% (-1.5-1.2%)	-4 (-26-20)	-19 (-130-100)
Ethiopia	1990	Female	95 plus	42% (26-61%)	58% (39-74%)	0.4 (0.19-0.68)	0.1% (-1.1-1.2%)	0.079% (-1.4-1.5%)	0.31 (-5.3-5.7)	1.6 (-17-19)
Ethiopia	1990	Male	All Ages	34% (28-39%)	66% (61-72%)	0.63 (0.57-0.7)	2.3% (1.5-3.2%)	3.5% (2-5.1%)	16000 (9300-23000)	670000 (420000-940000)
Ethiopia	1990	Male	15 to 19	30% (21-41%)	70% (59-79%)	0.56 (0.35-0.84)	2.1% (1-3.5%)	1.9% (0.59-3.5%)	250 (76-460)	24000 (11000-39000)
Ethiopia	1990	Male	20 to 24	43% (31-54%)	57% (46-69%)	0.86 (0.58-1.2)	3.8% (1.8-6.2%)	3.3% (1-6%)	560 (180-1000)	49000 (23000-80000)
Ethiopia	1990	Male	25 to 29	48% (35-59%)	52% (41-65%)	1.2 (0.83-1.6)	6.3% (3.6-9.6%)	6.1% (2.9-10%)	900 (440-1500)	65000 (37000-100000)
Ethiopia	1990	Male	30 to 34	51% (38-62%)	49% (38-62%)	1.4 (0.98-1.9)	8.2% (4.8-12%)	8.5% (4.5-13%)	1100 (610-1800)	72000 (42000-110000)
Ethiopia	1990	Male	35 to 39	53% (40-63%)	47% (37-60%)	1.5 (1-2)	9.4% (5.6-14%)	10% (5.7-15%)	1400 (770-2100)	77000 (45000-110000)
Ethiopia	1990	Male	40 to 44	55% (43-64%)	45% (36-57%)	1.5 (1.1-2)	9.8% (5.8-14%)	11% (6.1-16%)	1700 (920-2600)	81000 (47000-120000)
Ethiopia	1990	Male	45 to 49	57% (44-66%)	43% (34-56%)	1.5 (1-2)	8.9% (4.8-14%)	9.8% (5.2-15%)	1600 (850-2600)	70000 (37000-110000)
Ethiopia	1990	Male	50 to 54	59% (48-69%)	41% (31-52%)	1.5 (1.1-2)	9.3% (4.7-15%)	10% (5.2-16%)	1800 (900-2900)	68000 (34000-110000)
Ethiopia	1990	Male	55 to 59	60% (49-70%)	40% (30-51%)	1.7 (1.1-2.2)	9.6% (5.1-14%)	11% (5.6-16%)	2100 (1100-3200)	67000 (35000-100000)
Ethiopia	1990	Male	60 to 64	56% (43-67%)	44% (33-57%)	1.5 (1-2.1)	7.6% (4-12%)	8.4% (4.4-13%)	1700 (860-2600)	45000 (24000-70000)
Ethiopia	1990	Male	65 to 69	49% (36-61%)	51% (39-64%)	1.2 (0.76-1.6)	5.1% (2.4-8.2%)	5.6% (2.6-9.2%)	1200 (530-2000)	27000 (13000-45000)
Ethiopia	1990	Male	70 to 74	45% (32-57%)	55% (43-68%)	0.97 (0.61-1.4)	3.6% (1.1-6.4%)	4% (1.2-7.2%)	870 (250-1600)	16000 (4700-29000)
Ethiopia	1990	Male	75 to 79	44% (30-58%)	56% (42-70%)	0.85 (0.5-1.3)	2.8% (0.56-5.5%)	3.1% (0.59-6.2%)	580 (110-1200)	8100 (1600-16000)
Ethiopia	1990	Male	80 to 84	42% (28-56%)	58% (44-72%)	0.69 (0.42-1)	1.9% (0.2-4%)	2.1% (0.12-4.5%)	220 (13-470)	2300 (230-4900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	1990	Male	85 to 89	40% (27-54%)	60% (46-73%)	0.55 (0.31-0.85)	1.3% (-0.037-2.9%)	1.4% (-0.15-3.3%)	49 (-5.3-120)	380 (-11-880)
Ethiopia	1990	Male	90 to 94	41% (27-56%)	59% (44-73%)	0.47 (0.27-0.74)	0.73% (-0.3-1.8%)	0.7% (-0.5-2%)	6.9 (-4.9-20)	42 (-18-110)
Ethiopia	1990	Male	95 plus	45% (26-64%)	55% (36-74%)	0.42 (0.18-0.75)	0.81% (-0.69-2.3%)	0.75% (-1.1-2.6%)	1.3 (-1.1-4.3)	5.5 (-4.7-16)
Ethiopia	1995	Female	All Ages	29% (24-33%)	71% (67-76%)	0.34 (0.3-0.38)	0.74% (0.18-1.3%)	1.1% (0.11-2.2%)	4700 (470-9400)	190000 (44000-340000)
Ethiopia	1995	Female	15 to 19	29% (20-38%)	71% (62-80%)	0.37 (0.25-0.55)	0.83% (0.16-1.7%)	0.89% (-0.093-2.1%)	64 (-6.5-150)	6400 (1300-13000)
Ethiopia	1995	Female	20 to 24	36% (25-47%)	64% (53-75%)	0.51 (0.34-0.7)	1.3% (0.19-2.5%)	1.2% (-0.22-2.9%)	120 (-22-290)	11000 (1700-22000)
Ethiopia	1995	Female	25 to 29	39% (28-51%)	61% (49-72%)	0.65 (0.44-0.89)	1.8% (0.56-3.2%)	1.9% (0.27-3.7%)	220 (33-440)	17000 (5100-30000)
Ethiopia	1995	Female	30 to 34	42% (29-54%)	58% (46-71%)	0.72 (0.47-0.98)	2.1% (0.74-3.8%)	2.3% (0.49-4.4%)	280 (64-570)	18000 (6300-34000)
Ethiopia	1995	Female	35 to 39	45% (32-55%)	55% (45-68%)	0.72 (0.48-1)	2.2% (0.59-4.1%)	2.5% (0.45-4.8%)	350 (58-690)	19000 (4900-37000)
Ethiopia	1995	Female	40 to 44	47% (36-57%)	53% (43-64%)	0.74 (0.5-1)	2.3% (0.59-4%)	2.6% (0.55-4.9%)	370 (82-690)	18000 (4800-32000)
Ethiopia	1995	Female	45 to 49	47% (35-58%)	53% (42-65%)	0.75 (0.52-1.1)	2.5% (0.37-4.7%)	2.9% (0.39-5.7%)	470 (61-960)	20000 (3000-39000)
Ethiopia	1995	Female	50 to 54	45% (34-56%)	55% (44-66%)	0.72 (0.48-1)	2.9% (0.35-5.7%)	3.4% (0.35-6.7%)	620 (64-1300)	23000 (2700-45000)
Ethiopia	1995	Female	55 to 59	42% (30-55%)	58% (45-70%)	0.73 (0.46-1)	2.8% (0.22-5.9%)	3.3% (0.24-6.7%)	640 (42-1400)	20000 (1500-42000)
Ethiopia	1995	Female	60 to 64	43% (30-55%)	57% (45-70%)	0.74 (0.47-1.1)	2.4% (0.25-5%)	2.7% (0.31-5.9%)	630 (74-1300)	16000 (1800-35000)
Ethiopia	1995	Female	65 to 69	44% (32-55%)	56% (45-68%)	0.7 (0.47-0.96)	1.9% (-0.08-4.1%)	2.1% (-0.09-4.7%)	510 (-20-1200)	11000 (-440-25000)
Ethiopia	1995	Female	70 to 74	42% (31-54%)	58% (46-69%)	0.6 (0.38-0.87)	0.94% (-0.92-2.9%)	1% (-1-3.3%)	270 (-270-870)	4900 (-4700-15000)
Ethiopia	1995	Female	75 to 79	39% (26-53%)	61% (47-74%)	0.5 (0.3-0.76)	0.65% (-0.83-2.3%)	0.73% (-0.94-2.6%)	150 (-190-530)	2000 (-2600-7300)
Ethiopia	1995	Female	80 to 84	38% (24-51%)	62% (49-76%)	0.45 (0.28-0.68)	0.2% (-1.1-1.5%)	0.24% (-1.2-1.7%)	33 (-170-230)	320 (-1700-2300)
Ethiopia	1995	Female	85 to 89	37% (25-50%)	63% (50-75%)	0.42 (0.24-0.63)	-0.0021% (-1.2-1.1%)	0.0082% (-1.3-1.3%)	0.55 (-83-80)	-0.33 (-610-570)
Ethiopia	1995	Female	90 to 94	39% (26-53%)	61% (47-74%)	0.41 (0.24-0.65)	-0.2% (-1.2-0.81%)	-0.25% (-1.4-0.98%)	-4.8 (-28-19)	-23 (-130-92)
Ethiopia	1995	Female	95 plus	42% (25-60%)	58% (40-75%)	0.41 (0.21-0.72)	0.1% (-1.1-1.2%)	0.075% (-1.2-1.4%)	0.41 (-6.7-7.7)	2.2 (-22-26)
Ethiopia	1995	Male	All Ages	33% (28-38%)	67% (62-72%)	0.64 (0.59-0.7)	2.3% (1.6-3.1%)	3.5% (2.2-4.9%)	16000 (9600-22000)	660000 (440000-900000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	1995	Male	15 to 19	30% (21-41%)	70% (59-79%)	0.58 (0.37-0.85)	3% (1.6-4.8%)	2.9% (1-5.2%)	250 (86-460)	25000 (13000-40000)
Ethiopia	1995	Male	20 to 24	42% (30-54%)	58% (46-70%)	0.88 (0.6-1.2)	5.6% (3.1-8.7%)	5.2% (2.1-8.9%)	620 (240-1100)	56000 (31000-86000)
Ethiopia	1995	Male	25 to 29	47% (35-59%)	53% (41-65%)	1.2 (0.86-1.6)	7.5% (4.6-11%)	7.6% (4-12%)	930 (500-1500)	69000 (43000-100000)
Ethiopia	1995	Male	30 to 34	51% (38-62%)	49% (38-62%)	1.5 (1-1.9)	8.5% (5.3-12%)	9% (5-14%)	1100 (620-1700)	72000 (44000-100000)
Ethiopia	1995	Male	35 to 39	53% (40-63%)	47% (37-60%)	1.6 (1.1-2.2)	9.1% (5.6-13%)	9.9% (5.7-15%)	1300 (760-2000)	75000 (46000-110000)
Ethiopia	1995	Male	40 to 44	55% (43-64%)	45% (36-57%)	1.6 (1.1-2.1)	9.1% (5.4-13%)	10% (5.7-15%)	1500 (860-2200)	74000 (44000-110000)
Ethiopia	1995	Male	45 to 49	56% (43-66%)	44% (34-57%)	1.5 (1.1-2.1)	8.3% (4.8-12%)	9.3% (5.1-14%)	1500 (830-2400)	65000 (37000-99000)
Ethiopia	1995	Male	50 to 54	59% (48-69%)	41% (31-52%)	1.6 (1.1-2.2)	8.8% (4.7-14%)	10% (5.1-15%)	1700 (860-2700)	65000 (34000-100000)
Ethiopia	1995	Male	55 to 59	60% (49-70%)	40% (30-51%)	1.7 (1.2-2.3)	9.2% (5.2-13%)	10% (5.7-15%)	1900 (1000-2800)	61000 (34000-91000)
Ethiopia	1995	Male	60 to 64	56% (43-68%)	44% (32-57%)	1.6 (1.1-2.2)	7.2% (4.2-11%)	8.1% (4.6-12%)	1800 (980-2800)	48000 (28000-75000)
Ethiopia	1995	Male	65 to 69	48% (36-61%)	52% (39-64%)	1.2 (0.79-1.6)	4.9% (2.4-7.7%)	5.5% (2.6-8.7%)	1100 (480-1700)	24000 (12000-39000)
Ethiopia	1995	Male	70 to 74	44% (32-56%)	56% (44-68%)	1 (0.63-1.4)	3.5% (1.3-6.2%)	3.9% (1.4-7.1%)	850 (310-1500)	15000 (5800-28000)
Ethiopia	1995	Male	75 to 79	43% (30-57%)	57% (43-70%)	0.88 (0.53-1.3)	2.8% (0.75-5.2%)	3.1% (0.79-6%)	680 (170-1300)	9500 (2600-18000)
Ethiopia	1995	Male	80 to 84	42% (28-56%)	58% (44-72%)	0.72 (0.43-1.1)	1.9% (0.37-3.8%)	2.1% (0.32-4.2%)	290 (42-590)	3100 (590-6100)
Ethiopia	1995	Male	85 to 89	40% (27-55%)	60% (45-73%)	0.57 (0.33-0.89)	1.3% (0.14-2.8%)	1.4% (0.052-3.1%)	66 (2.4-150)	520 (54-1100)
Ethiopia	1995	Male	90 to 94	41% (27-56%)	59% (44-73%)	0.48 (0.28-0.78)	0.73% (-0.2-1.8%)	0.7% (-0.39-2%)	8.1 (-4.6-23)	49 (-14-120)
Ethiopia	1995	Male	95 plus	44% (25-63%)	56% (37-75%)	0.42 (0.19-0.74)	0.83% (-0.55-2.3%)	0.77% (-0.91-2.6%)	1.9 (-2.3-6.5)	8.3 (-5.5-23)
Ethiopia	2000	Female	All Ages	24% (21-28%)	76% (72-79%)	0.36 (0.32-0.41)	0.91% (0.47-1.4%)	1.5% (0.61-2.4%)	6500 (2700-11000)	240000 (130000-370000)
Ethiopia	2000	Female	15 to 19	25% (20-31%)	75% (69-80%)	0.4 (0.28-0.56)	0.84% (0.33-1.5%)	0.87% (0.16-1.7%)	83 (16-170)	8100 (3100-14000)
Ethiopia	2000	Female	20 to 24	33% (27-40%)	67% (60-73%)	0.58 (0.42-0.77)	1.4% (0.51-2.4%)	1.3% (0.17-2.7%)	160 (24-320)	14000 (5500-25000)
Ethiopia	2000	Female	25 to 29	35% (29-41%)	65% (59-71%)	0.71 (0.54-0.91)	1.8% (0.86-3%)	1.8% (0.67-3.3%)	270 (100-480)	20000 (9700-33000)
Ethiopia	2000	Female	30 to 34	35% (28-42%)	65% (58-72%)	0.73 (0.54-0.96)	1.9% (0.97-3.2%)	2% (0.89-3.6%)	330 (140-570)	21000 (11000-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	2000	Female	35 to 39	36% (29-43%)	64% (57-71%)	0.7 (0.5-0.93)	2.1% (0.99-3.4%)	2.3% (0.98-3.9%)	380 (150-650)	21000 (9600-35000)
Ethiopia	2000	Female	40 to 44	37% (29-45%)	63% (55-71%)	0.71 (0.5-0.96)	2.2% (1.1-3.7%)	2.5% (1.2-4.3%)	430 (190-750)	21000 (9900-35000)
Ethiopia	2000	Female	45 to 49	40% (32-47%)	60% (53-68%)	0.78 (0.56-1.1)	2.8% (1.2-4.7%)	3.2% (1.3-5.6%)	550 (220-970)	23000 (9500-40000)
Ethiopia	2000	Female	50 to 54	41% (33-49%)	59% (51-67%)	0.81 (0.58-1.1)	3.5% (1.4-6%)	4.1% (1.6-7.1%)	850 (340-1500)	31000 (12000-52000)
Ethiopia	2000	Female	55 to 59	41% (32-51%)	59% (49-68%)	0.89 (0.62-1.2)	3.6% (1.5-6.3%)	4.3% (1.7-7.5%)	940 (380-1600)	29000 (12000-51000)
Ethiopia	2000	Female	60 to 64	42% (31-52%)	58% (48-69%)	0.89 (0.61-1.2)	3.1% (1.2-5.4%)	3.5% (1.3-6.2%)	830 (310-1500)	22000 (8100-39000)
Ethiopia	2000	Female	65 to 69	40% (31-50%)	60% (50-69%)	0.79 (0.56-1.1)	2.5% (0.82-4.6%)	2.9% (0.94-5.2%)	790 (250-1500)	17000 (5600-32000)
Ethiopia	2000	Female	70 to 74	37% (28-48%)	63% (52-72%)	0.65 (0.42-0.94)	1.5% (-0.2-3.4%)	1.6% (-0.27-3.9%)	430 (-69-1000)	7700 (-1100-18000)
Ethiopia	2000	Female	75 to 79	36% (23-49%)	64% (51-77%)	0.55 (0.33-0.85)	1.1% (-0.31-2.7%)	1.2% (-0.36-3%)	300 (-84-740)	4100 (-1200-10000)
Ethiopia	2000	Female	80 to 84	37% (24-51%)	63% (49-76%)	0.53 (0.33-0.81)	0.55% (-0.66-1.9%)	0.64% (-0.74-2.2%)	100 (-120-360)	1000 (-1300-3600)
Ethiopia	2000	Female	85 to 89	37% (25-51%)	63% (49-75%)	0.52 (0.32-0.78)	0.27% (-0.86-1.4%)	0.32% (-0.96-1.7%)	23 (-69-120)	160 (-510-870)
Ethiopia	2000	Female	90 to 94	39% (27-53%)	61% (47-73%)	0.51 (0.3-0.81)	0.0094% (-1.1-1.1%)	-0.0044% (-1.2-1.3%)	-0.1 (-28-30)	1.4 (-140-150)
Ethiopia	2000	Female	95 plus	42% (26-60%)	58% (40-74%)	0.5 (0.25-0.9)	0.33% (-0.78-1.5%)	0.33% (-0.99-1.8%)	1.9 (-5.9-10)	7.5 (-18-35)
Ethiopia	2000	Male	All Ages	32% (28-36%)	68% (64-72%)	0.77 (0.69-0.84)	2.5% (1.9-3.2%)	3.8% (2.7-4.9%)	18000 (12000-23000)	750000 (550000-950000)
Ethiopia	2000	Male	15 to 19	26% (19-33%)	74% (67-81%)	0.61 (0.4-0.86)	2.4% (1.5-3.6%)	2.2% (1-3.7%)	290 (140-490)	29000 (18000-44000)
Ethiopia	2000	Male	20 to 24	40% (33-48%)	60% (52-67%)	1 (0.74-1.4)	4.6% (2.8-6.6%)	3.9% (2.1-6.3%)	730 (390-1200)	66000 (40000-96000)
Ethiopia	2000	Male	25 to 29	47% (40-54%)	53% (46-60%)	1.5 (1.1-1.9)	6.4% (4.2-8.8%)	6.1% (3.6-8.9%)	1200 (690-1700)	87000 (57000-120000)
Ethiopia	2000	Male	30 to 34	49% (42-56%)	51% (44-58%)	1.7 (1.3-2.2)	7.1% (4.9-9.6%)	7.2% (4.5-10%)	1300 (810-1800)	84000 (57000-110000)
Ethiopia	2000	Male	35 to 39	51% (43-58%)	49% (42-57%)	1.9 (1.4-2.4)	7.8% (5.2-11%)	8.4% (5.3-12%)	1400 (890-2000)	79000 (53000-110000)
Ethiopia	2000	Male	40 to 44	54% (47-62%)	46% (38-53%)	2 (1.4-2.5)	8.6% (5.7-11%)	9.4% (6-13%)	1600 (1000-2200)	78000 (51000-110000)
Ethiopia	2000	Male	45 to 49	57% (50-63%)	43% (37-50%)	1.9 (1.4-2.5)	8.4% (5.6-12%)	9.4% (6-13%)	1600 (1000-2300)	69000 (45000-98000)
Ethiopia	2000	Male	50 to 54	60% (53-66%)	40% (34-47%)	2 (1.5-2.6)	9.6% (6.2-13%)	11% (7-16%)	1900 (1200-2800)	72000 (46000-100000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	2000	Male	55 to 59	63% (56-70%)	37% (30-44%)	2.2 (1.7-2.9)	10% (6.8-14%)	12% (7.7-16%)	2200 (1400-3100)	71000 (46000-100000)
Ethiopia	2000	Male	60 to 64	56% (47-64%)	44% (36-53%)	1.9 (1.4-2.5)	8% (5.3-11%)	9.1% (5.8-13%)	1800 (1100-2600)	49000 (32000-70000)
Ethiopia	2000	Male	65 to 69	46% (36-56%)	54% (44-64%)	1.4 (0.97-1.8)	5.4% (3.1-8.1%)	6.1% (3.5-9.2%)	1400 (770-2100)	31000 (18000-47000)
Ethiopia	2000	Male	70 to 74	45% (35-55%)	55% (45-65%)	1.2 (0.82-1.7)	4.2% (2-6.7%)	4.8% (2.2-7.7%)	900 (430-1500)	16000 (7900-26000)
Ethiopia	2000	Male	75 to 79	44% (32-56%)	56% (44-68%)	1.1 (0.68-1.6)	3.4% (1.3-5.8%)	3.9% (1.4-6.6%)	700 (250-1200)	9900 (3600-17000)
Ethiopia	2000	Male	80 to 84	41% (28-56%)	59% (44-72%)	0.87 (0.52-1.3)	2.4% (0.84-4.4%)	2.7% (0.9-5%)	400 (130-730)	4200 (1400-7500)
Ethiopia	2000	Male	85 to 89	39% (26-53%)	61% (47-74%)	0.69 (0.41-1.1)	1.7% (0.53-3.3%)	1.9% (0.48-3.8%)	130 (33-260)	1000 (320-1900)
Ethiopia	2000	Male	90 to 94	40% (27-55%)	60% (45-73%)	0.59 (0.34-0.95)	1% (0.12-2.1%)	1% (-0.02-2.4%)	17 (-0.33-39)	98 (12-210)
Ethiopia	2000	Male	95 plus	44% (26-63%)	56% (37-74%)	0.52 (0.24-0.9)	1.1% (-0.19-2.7%)	1.1% (-0.5-2.9%)	3.1 (-1.4-8.4)	13 (-2-30)
Ethiopia	2005	Female	All Ages	27% (24-31%)	73% (69-76%)	0.41 (0.37-0.45)	0.99% (0.51-1.5%)	1.5% (0.6-2.5%)	6200 (2500-10000)	240000 (120000-360000)
Ethiopia	2005	Female	15 to 19	30% (25-36%)	70% (64-75%)	0.5 (0.37-0.66)	1.1% (0.43-1.9%)	1.3% (0.25-2.6%)	94 (18-190)	9300 (3600-16000)
Ethiopia	2005	Female	20 to 24	38% (31-44%)	62% (56-69%)	0.67 (0.49-0.88)	1.7% (0.68-3%)	1.8% (0.3-3.6%)	170 (30-340)	16000 (6600-28000)
Ethiopia	2005	Female	25 to 29	38% (31-44%)	62% (56-69%)	0.79 (0.6-1)	2% (0.97-3.2%)	2.1% (0.79-3.8%)	280 (100-490)	21000 (10000-34000)
Ethiopia	2005	Female	30 to 34	39% (33-46%)	61% (54-67%)	0.85 (0.64-1.1)	2% (1-3.1%)	2.1% (0.94-3.6%)	330 (150-570)	22000 (11000-35000)
Ethiopia	2005	Female	35 to 39	41% (35-47%)	59% (53-65%)	0.82 (0.61-1.1)	2.1% (1-3.4%)	2.3% (1-4%)	400 (160-690)	22000 (10000-37000)
Ethiopia	2005	Female	40 to 44	42% (35-49%)	58% (51-65%)	0.82 (0.59-1.1)	2.2% (1-3.5%)	2.5% (1.1-4.1%)	420 (170-690)	20000 (9100-33000)
Ethiopia	2005	Female	45 to 49	44% (36-50%)	56% (50-64%)	0.86 (0.63-1.1)	2.7% (1.1-4.5%)	3.2% (1.3-5.4%)	550 (220-980)	23000 (9800-41000)
Ethiopia	2005	Female	50 to 54	42% (33-51%)	58% (49-67%)	0.84 (0.6-1.2)	3.2% (1.3-5.4%)	3.9% (1.6-6.6%)	720 (280-1200)	26000 (11000-45000)
Ethiopia	2005	Female	55 to 59	41% (31-51%)	59% (49-69%)	0.89 (0.61-1.2)	3.3% (1.3-5.8%)	4% (1.6-7%)	880 (350-1500)	27000 (11000-48000)
Ethiopia	2005	Female	60 to 64	42% (31-53%)	58% (47-69%)	0.9 (0.6-1.3)	2.8% (1-5%)	3.4% (1.2-5.9%)	810 (300-1400)	21000 (7800-38000)
Ethiopia	2005	Female	65 to 69	41% (31-51%)	59% (49-69%)	0.82 (0.57-1.1)	2.4% (0.75-4.3%)	2.8% (0.87-5%)	720 (220-1300)	16000 (4800-28000)
Ethiopia	2005	Female	70 to 74	38% (28-49%)	62% (51-72%)	0.68 (0.43-0.98)	1.4% (-0.19-3.4%)	1.6% (-0.24-3.9%)	460 (-66-1100)	8300 (-1100-20000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	2005	Female	75 to 79	36% (24-50%)	64% (50-76%)	0.57 (0.34-0.89)	1% (-0.28-2.7%)	1.2% (-0.32-3.1%)	280 (-75-730)	3800 (-1000-10000)
Ethiopia	2005	Female	80 to 84	36% (24-51%)	64% (49-76%)	0.54 (0.33-0.82)	0.47% (-0.7-1.7%)	0.55% (-0.8-2%)	100 (-150-390)	1000 (-1500-3900)
Ethiopia	2005	Female	85 to 89	37% (25-51%)	63% (49-75%)	0.53 (0.31-0.79)	0.21% (-0.92-1.4%)	0.25% (-1-1.6%)	22 (-90-140)	160 (-670-980)
Ethiopia	2005	Female	90 to 94	39% (26-52%)	61% (48-74%)	0.51 (0.3-0.82)	-0.05% (-1.1-0.99%)	-0.076% (-1.3-1.1%)	-2 (-35-30)	-7.7 (-170-150)
Ethiopia	2005	Female	95 plus	42% (25-60%)	58% (40-75%)	0.51 (0.26-0.89)	0.36% (-0.79-1.6%)	0.36% (-0.97-1.8%)	2.5 (-6.6-13)	9.5 (-21-42)
Ethiopia	2005	Male	All Ages	34% (30-38%)	66% (62-70%)	0.82 (0.76-0.89)	3% (2.3-3.8%)	4.4% (3.2-5.7%)	18000 (13000-24000)	780000 (590000-1000000)
Ethiopia	2005	Male	15 to 19	37% (30-43%)	63% (57-70%)	0.91 (0.66-1.2)	4.3% (2.8-6.1%)	4.8% (2.4-7.4%)	400 (200-630)	39000 (24000-56000)
Ethiopia	2005	Male	20 to 24	45% (40-51%)	55% (49-60%)	1.2 (0.91-1.5)	7.8% (5.1-11%)	7.8% (4.3-12%)	870 (480-1400)	78000 (50000-110000)
Ethiopia	2005	Male	25 to 29	47% (42-53%)	53% (47-58%)	1.5 (1.2-2)	8.8% (6-12%)	9.2% (5.5-13%)	1200 (730-1700)	89000 (60000-120000)
Ethiopia	2005	Male	30 to 34	49% (44-55%)	51% (45-56%)	1.8 (1.3-2.2)	8.4% (5.9-11%)	8.9% (5.7-13%)	1400 (920-2000)	93000 (64000-130000)
Ethiopia	2005	Male	35 to 39	50% (44-56%)	50% (44-56%)	1.8 (1.4-2.4)	8.2% (5.6-11%)	8.9% (5.7-12%)	1400 (910-2000)	82000 (54000-110000)
Ethiopia	2005	Male	40 to 44	54% (48-60%)	46% (40-52%)	2 (1.5-2.5)	8.6% (5.9-12%)	9.7% (6.4-13%)	1500 (980-2100)	75000 (50000-100000)
Ethiopia	2005	Male	45 to 49	57% (52-62%)	43% (38-48%)	2 (1.5-2.5)	8.5% (5.8-11%)	9.7% (6.4-13%)	1600 (1000-2200)	68000 (45000-93000)
Ethiopia	2005	Male	50 to 54	60% (54-66%)	40% (34-46%)	2 (1.5-2.7)	9.5% (6.2-13%)	11% (7.1-15%)	1800 (1100-2500)	68000 (44000-93000)
Ethiopia	2005	Male	55 to 59	64% (57-69%)	36% (31-43%)	2.3 (1.7-3)	11% (7-14%)	12% (8.2-17%)	2200 (1400-3000)	71000 (46000-96000)
Ethiopia	2005	Male	60 to 64	56% (48-65%)	44% (35-52%)	2 (1.4-2.6)	8.2% (5.3-11%)	9.5% (6-13%)	1900 (1200-2700)	52000 (33000-73000)
Ethiopia	2005	Male	65 to 69	46% (35-56%)	54% (44-65%)	1.4 (0.99-1.9)	5.5% (3.2-8.1%)	6.4% (3.7-9.4%)	1300 (770-2000)	30000 (18000-44000)
Ethiopia	2005	Male	70 to 74	44% (34-54%)	56% (46-66%)	1.3 (0.83-1.7)	4.2% (2-6.5%)	4.8% (2.3-7.5%)	1100 (520-1800)	20000 (9900-32000)
Ethiopia	2005	Male	75 to 79	44% (33-55%)	56% (45-67%)	1.1 (0.69-1.6)	3.5% (1.4-5.7%)	4% (1.5-6.7%)	670 (270-1100)	9400 (3900-16000)
Ethiopia	2005	Male	80 to 84	41% (28-55%)	59% (45-72%)	0.88 (0.53-1.3)	2.4% (0.88-4.2%)	2.8% (0.91-4.9%)	370 (120-660)	3900 (1400-6800)
Ethiopia	2005	Male	85 to 89	39% (25-53%)	61% (47-75%)	0.69 (0.41-1.1)	1.7% (0.55-3.1%)	1.9% (0.5-3.5%)	160 (43-300)	1300 (400-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	2005	Male	90 to 94	40% (27-54%)	60% (46-73%)	0.59 (0.34-0.94)	1% (0.11-2.1%)	1% (-0.054-2.3%)	29 (-1.5-65)	170 (18-350)
Ethiopia	2005	Male	95 plus	44% (25-62%)	56% (38-75%)	0.53 (0.24-0.93)	1.2% (-0.14-2.7%)	1.2% (-0.42-3.1%)	5.4 (-1.9-14)	22 (-2.6-50)
Ethiopia	2010	Female	All Ages	30% (27-34%)	70% (66-73%)	0.42 (0.37-0.48)	1% (0.49-1.5%)	1.4% (0.55-2.4%)	5200 (1900-8700)	210000 (100000-320000)
Ethiopia	2010	Female	15 to 19	35% (29-39%)	65% (61-71%)	0.55 (0.41-0.7)	1% (0.43-1.7%)	1.2% (0.25-2.3%)	90 (18-170)	9500 (3900-16000)
Ethiopia	2010	Female	20 to 24	41% (36-46%)	59% (54-64%)	0.69 (0.52-0.9)	1.7% (0.78-2.7%)	1.8% (0.34-3.3%)	140 (25-250)	14000 (6500-22000)
Ethiopia	2010	Female	25 to 29	41% (36-45%)	59% (55-64%)	0.8 (0.62-1)	2.2% (1.1-3.4%)	2.5% (0.97-4.4%)	240 (93-430)	20000 (10000-32000)
Ethiopia	2010	Female	30 to 34	42% (37-47%)	58% (53-63%)	0.86 (0.67-1.1)	2.1% (1.1-3.4%)	2.5% (1.1-4.1%)	280 (120-490)	20000 (10000-31000)
Ethiopia	2010	Female	35 to 39	43% (39-48%)	57% (52-61%)	0.82 (0.62-1.1)	2.2% (1.1-3.6%)	2.6% (1.1-4.4%)	340 (140-610)	20000 (9400-33000)
Ethiopia	2010	Female	40 to 44	44% (38-49%)	56% (51-62%)	0.81 (0.59-1.1)	2.2% (1.1-3.4%)	2.7% (1.2-4.3%)	360 (150-610)	18000 (8500-30000)
Ethiopia	2010	Female	45 to 49	46% (40-52%)	54% (48-60%)	0.87 (0.65-1.1)	2.6% (1.1-4.4%)	3.3% (1.3-5.5%)	450 (160-810)	19000 (7400-34000)
Ethiopia	2010	Female	50 to 54	46% (36-56%)	54% (44-64%)	0.88 (0.6-1.2)	3% (1.3-5.1%)	3.8% (1.7-6.4%)	640 (260-1100)	23000 (9600-41000)
Ethiopia	2010	Female	55 to 59	42% (30-55%)	58% (45-70%)	0.88 (0.56-1.3)	2.9% (1.1-5.1%)	3.6% (1.4-6.5%)	670 (240-1200)	21000 (7700-38000)
Ethiopia	2010	Female	60 to 64	43% (30-56%)	57% (44-70%)	0.89 (0.56-1.3)	2.4% (0.82-4.5%)	2.9% (1.1-5.4%)	690 (240-1300)	18000 (6100-34000)
Ethiopia	2010	Female	65 to 69	44% (33-55%)	56% (45-67%)	0.83 (0.55-1.2)	2% (0.58-3.7%)	2.4% (0.75-4.5%)	630 (200-1200)	14000 (4200-26000)
Ethiopia	2010	Female	70 to 74	42% (31-54%)	58% (46-69%)	0.72 (0.46-1)	1.1% (-0.32-2.7%)	1.3% (-0.38-3.2%)	340 (-100-850)	6200 (-1900-15000)
Ethiopia	2010	Female	75 to 79	39% (26-53%)	61% (47-74%)	0.59 (0.36-0.91)	0.77% (-0.44-2.2%)	0.91% (-0.49-2.6%)	250 (-130-720)	3300 (-1900-9700)
Ethiopia	2010	Female	80 to 84	38% (25-52%)	62% (48-75%)	0.53 (0.33-0.79)	0.23% (-0.82-1.3%)	0.29% (-0.93-1.6%)	64 (-190-340)	600 (-2000-3400)
Ethiopia	2010	Female	85 to 89	37% (25-51%)	63% (49-75%)	0.5 (0.3-0.76)	0.036% (-0.97-0.98%)	0.053% (-1.1-1.2%)	5.5 (-110-120)	32 (-830-850)
Ethiopia	2010	Female	90 to 94	39% (27-52%)	61% (48-73%)	0.49 (0.3-0.79)	-0.21% (-1.1-0.67%)	-0.27% (-1.4-0.75%)	-9 (-46-25)	-41 (-230-130)
Ethiopia	2010	Female	95 plus	42% (26-60%)	58% (40-74%)	0.48 (0.24-0.86)	0.23% (-0.85-1.2%)	0.2% (-1.1-1.4%)	1.5 (-7.9-10)	6.7 (-25-37)
Ethiopia	2010	Male	All Ages	35% (31-39%)	65% (61-69%)	0.81 (0.72-0.91)	3.4% (2.5-4.3%)	4.5% (3.2-6%)	17000 (11000-23000)	770000 (570000-1000000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	2010	Male	15 to 19	39% (33-45%)	61% (55-67%)	0.92 (0.67-1.2)	4.2% (2.7-5.8%)	4.3% (2.2-6.6%)	410 (200-630)	44000 (27000-61000)
Ethiopia	2010	Male	20 to 24	47% (42-53%)	53% (47-58%)	1.2 (0.89-1.5)	8.2% (5.6-11%)	7.5% (4.1-11%)	810 (430-1300)	83000 (56000-120000)
Ethiopia	2010	Male	25 to 29	48% (42-53%)	52% (47-58%)	1.5 (1.1-1.9)	9.7% (6.8-13%)	9.9% (6-14%)	1100 (670-1700)	95000 (64000-130000)
Ethiopia	2010	Male	30 to 34	50% (44-56%)	50% (44-56%)	1.7 (1.3-2.2)	9.3% (6.7-12%)	10% (6.3-14%)	1300 (770-1900)	88000 (59000-120000)
Ethiopia	2010	Male	35 to 39	50% (44-57%)	50% (43-56%)	1.8 (1.3-2.3)	8.7% (6.2-11%)	9.6% (6.5-13%)	1400 (880-2100)	84000 (57000-120000)
Ethiopia	2010	Male	40 to 44	54% (48-60%)	46% (40-52%)	1.9 (1.4-2.4)	8.8% (6.2-12%)	10% (6.7-14%)	1500 (910-2100)	75000 (50000-110000)
Ethiopia	2010	Male	45 to 49	57% (52-63%)	43% (37-48%)	1.9 (1.4-2.4)	8.6% (5.9-12%)	10% (6.7-14%)	1400 (890-2000)	63000 (41000-88000)
Ethiopia	2010	Male	50 to 54	61% (55-67%)	39% (33-45%)	2 (1.4-2.5)	9.3% (6.1-13%)	11% (7-15%)	1700 (1000-2400)	64000 (41000-92000)
Ethiopia	2010	Male	55 to 59	64% (57-70%)	36% (30-43%)	2.2 (1.6-2.9)	9.7% (6.4-13%)	12% (7.6-16%)	1900 (1200-2800)	64000 (42000-91000)
Ethiopia	2010	Male	60 to 64	58% (47-68%)	42% (32-53%)	1.9 (1.3-2.6)	7.5% (4.8-11%)	8.8% (5.6-12%)	1800 (1100-2600)	49000 (31000-71000)
Ethiopia	2010	Male	65 to 69	48% (35-60%)	52% (40-65%)	1.4 (0.92-2)	5.1% (2.9-7.4%)	5.9% (3.3-8.7%)	1300 (720-2000)	30000 (17000-46000)
Ethiopia	2010	Male	70 to 74	44% (32-56%)	56% (44-68%)	1.2 (0.73-1.7)	3.5% (1.7-5.6%)	4.1% (1.9-6.5%)	930 (420-1500)	17000 (8100-28000)
Ethiopia	2010	Male	75 to 79	43% (29-57%)	57% (43-71%)	1 (0.61-1.6)	3% (1.2-5%)	3.4% (1.4-5.9%)	750 (300-1300)	11000 (4300-19000)
Ethiopia	2010	Male	80 to 84	41% (28-55%)	59% (45-72%)	0.84 (0.49-1.3)	2.1% (0.76-3.6%)	2.4% (0.77-4.3%)	390 (130-710)	4100 (1500-7400)
Ethiopia	2010	Male	85 to 89	39% (26-54%)	61% (46-74%)	0.67 (0.39-1)	1.5% (0.5-2.7%)	1.6% (0.45-3.1%)	130 (35-240)	990 (330-1800)
Ethiopia	2010	Male	90 to 94	40% (26-56%)	60% (44-74%)	0.57 (0.33-0.93)	0.86% (0.057-1.8%)	0.84% (-0.13-2%)	29 (-4.6-68)	170 (12-380)
Ethiopia	2010	Male	95 plus	44% (26-62%)	56% (38-74%)	0.5 (0.23-0.88)	1% (-0.094-2.3%)	0.98% (-0.37-2.6%)	7.5 (-2.8-19)	31 (-2.9-73)
Ethiopia	2016	Female	All Ages	31% (28-34%)	69% (66-72%)	0.45 (0.3-0.66)	1.2% (0.5-2.1%)	1.6% (0.42-3.2%)	5300 (1300-11000)	210000 (89000-380000)
Ethiopia	2016	Female	15 to 19	34% (29-39%)	66% (61-71%)	0.55 (0.34-0.85)	1% (0.48-1.7%)	1.2% (0.31-2.5%)	85 (23-170)	9900 (4400-17000)
Ethiopia	2016	Female	20 to 24	41% (35-46%)	59% (54-65%)	0.7 (0.44-1.1)	1.7% (0.89-2.8%)	1.8% (0.46-3.5%)	130 (33-270)	15000 (7800-25000)
Ethiopia	2016	Female	25 to 29	40% (35-46%)	60% (54-65%)	0.82 (0.51-1.2)	2.3% (1.3-3.6%)	2.7% (1.1-4.9%)	220 (83-430)	19000 (10000-33000)
Ethiopia	2016	Female	30 to 34	42% (36-47%)	58% (53-64%)	0.87 (0.54-1.3)	2.4% (1.3-3.8%)	3% (1.3-5.2%)	290 (120-530)	21000 (11000-35000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	2016	Female	35 to 39	43% (38-48%)	57% (52-62%)	0.84 (0.51-1.3)	2.5% (1.3-4.1%)	3.2% (1.4-5.5%)	350 (150-660)	21000 (10000-36000)
Ethiopia	2016	Female	40 to 44	44% (38-50%)	56% (50-62%)	0.83 (0.51-1.3)	2.4% (1.3-4%)	3.2% (1.5-5.4%)	390 (180-690)	20000 (9800-34000)
Ethiopia	2016	Female	45 to 49	46% (41-52%)	54% (48-59%)	0.89 (0.54-1.4)	2.7% (1.2-4.9%)	3.6% (1.4-6.5%)	490 (180-910)	21000 (8300-38000)
Ethiopia	2016	Female	50 to 54	46% (35-56%)	54% (44-65%)	0.9 (0.52-1.4)	2.9% (1.1-5.3%)	3.8% (1.4-6.9%)	620 (220-1200)	23000 (8300-42000)
Ethiopia	2016	Female	55 to 59	42% (31-54%)	58% (46-69%)	0.89 (0.48-1.5)	2.7% (0.72-5.3%)	3.4% (0.99-6.9%)	660 (180-1400)	21000 (5800-43000)
Ethiopia	2016	Female	60 to 64	43% (30-55%)	57% (45-70%)	0.9 (0.49-1.5)	2.2% (0.46-4.8%)	2.8% (0.66-6.1%)	630 (150-1400)	17000 (3500-36000)
Ethiopia	2016	Female	65 to 69	44% (32-56%)	56% (44-68%)	0.85 (0.49-1.4)	2% (0.37-4.2%)	2.4% (0.5-5.1%)	670 (140-1500)	15000 (2700-33000)
Ethiopia	2016	Female	70 to 74	42% (31-54%)	58% (46-69%)	0.74 (0.4-1.2)	1.1% (-0.47-3.2%)	1.3% (-0.55-3.7%)	390 (-160-1100)	7000 (-2900-21000)
Ethiopia	2016	Female	75 to 79	39% (26-54%)	61% (46-74%)	0.61 (0.32-1)	0.82% (-0.52-2.7%)	0.98% (-0.58-3.2%)	280 (-160-900)	3800 (-2300-12000)
Ethiopia	2016	Female	80 to 84	38% (24-52%)	62% (48-76%)	0.55 (0.28-0.92)	0.29% (-0.86-1.7%)	0.37% (-0.95-2%)	94 (-240-510)	890 (-2600-5100)
Ethiopia	2016	Female	85 to 89	38% (25-53%)	62% (47-75%)	0.52 (0.27-0.93)	0.074% (-0.95-1.4%)	0.1% (-1.1-1.6%)	14 (-150-220)	88 (-1100-1600)
Ethiopia	2016	Female	90 to 94	39% (26-54%)	61% (46-74%)	0.51 (0.27-0.93)	-0.19% (-1.1-1.1%)	-0.25% (-1.3-1.2%)	-10 (-58-52)	-48 (-290-270)
Ethiopia	2016	Female	95 plus	42% (26-60%)	58% (40-74%)	0.51 (0.22-1)	0.31% (-0.69-1.6%)	0.3% (-0.88-2%)	3.1 (-9.2-19)	12 (-28-67)
Ethiopia	2016	Male	All Ages	36% (32-40%)	64% (60-68%)	0.88 (0.59-1.3)	4.2% (2.8-5.8%)	5% (2.9-7.5%)	18000 (9600-28000)	850000 (540000-1200000)
Ethiopia	2016	Male	15 to 19	40% (33-46%)	60% (54-67%)	0.95 (0.57-1.5)	4.1% (2.6-5.9%)	4.1% (1.9-6.7%)	440 (190-780)	50000 (30000-76000)
Ethiopia	2016	Male	20 to 24	48% (42-54%)	52% (46-58%)	1.2 (0.77-1.9)	8.1% (5.4-11%)	7.1% (3.5-12%)	930 (440-1600)	100000 (63000-150000)
Ethiopia	2016	Male	25 to 29	48% (42-53%)	52% (47-58%)	1.5 (0.93-2.4)	9.8% (6.6-14%)	9.8% (5.4-15%)	1200 (610-2000)	110000 (68000-160000)
Ethiopia	2016	Male	30 to 34	50% (44-56%)	50% (44-56%)	1.8 (1.1-2.7)	10% (6.8-14%)	11% (6.3-16%)	1300 (710-2100)	99000 (63000-150000)
Ethiopia	2016	Male	35 to 39	50% (44-57%)	50% (43-56%)	1.8 (1.1-2.8)	9.6% (6.2-14%)	11% (6.5-16%)	1400 (760-2200)	86000 (54000-130000)
Ethiopia	2016	Male	40 to 44	54% (48-60%)	46% (40-52%)	1.9 (1.2-2.9)	9.4% (6-14%)	11% (6.7-17%)	1500 (880-2500)	82000 (51000-130000)
Ethiopia	2016	Male	45 to 49	58% (52-63%)	42% (37-48%)	1.9 (1.2-3)	8.8% (5.5-13%)	11% (6.3-16%)	1500 (840-2500)	71000 (42000-110000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ethiopia	2016	Male	50 to 54	61% (55-67%)	39% (33-45%)	2 (1.2-3.2)	9.1% (5.5-14%)	11% (6.3-17%)	1700 (890-2800)	66000 (36000-100000)
Ethiopia	2016	Male	55 to 59	64% (57-70%)	36% (30-43%)	2.3 (1.3-3.5)	9.3% (5.3-14%)	11% (6.1-17%)	1900 (1000-3100)	65000 (36000-100000)
Ethiopia	2016	Male	60 to 64	58% (46-68%)	42% (32-54%)	2 (1.2-3.3)	7.2% (3.9-11%)	8.4% (4.4-14%)	1700 (840-2900)	49000 (25000-82000)
Ethiopia	2016	Male	65 to 69	48% (36-60%)	52% (40-64%)	1.5 (0.83-2.3)	4.7% (2.3-8.2%)	5.5% (2.5-9.4%)	1300 (580-2400)	31000 (14000-55000)
Ethiopia	2016	Male	70 to 74	44% (31-55%)	56% (45-69%)	1.2 (0.64-2)	3.3% (1.2-6%)	3.8% (1.3-6.9%)	1000 (330-1900)	19000 (6600-36000)
Ethiopia	2016	Male	75 to 79	43% (29-57%)	57% (43-71%)	1.1 (0.53-1.8)	2.8% (0.95-5.4%)	3.2% (1.1-6.3%)	790 (250-1600)	11000 (3700-23000)
Ethiopia	2016	Male	80 to 84	41% (27-55%)	59% (45-73%)	0.86 (0.44-1.5)	2% (0.54-3.9%)	2.2% (0.55-4.4%)	450 (110-930)	4800 (1300-9700)
Ethiopia	2016	Male	85 to 89	39% (26-54%)	61% (46-74%)	0.68 (0.35-1.2)	1.4% (0.38-2.9%)	1.5% (0.31-3.3%)	170 (32-360)	1300 (340-2700)
Ethiopia	2016	Male	90 to 94	40% (27-54%)	60% (46-73%)	0.58 (0.29-1.1)	0.85% (0.054-2%)	0.82% (-0.11-2.2%)	27 (-3.7-73)	170 (11-410)
Ethiopia	2016	Male	95 plus	44% (26-62%)	56% (38-74%)	0.52 (0.21-1)	0.99% (-0.0056-2.4%)	0.93% (-0.24-2.6%)	8.7 (-2.2-25)	37 (-0.21-92)
Kenya	1990	Female	All Ages	11% (9.2-13%)	89% (87-91%)	0.45 (0.38-0.53)	0.63% (0.46-0.83%)	1.1% (0.71-1.7%)	1100 (660-1600)	39000 (28000-53000)
Kenya	1990	Female	15 to 19	11% (7.1-17%)	89% (83-93%)	0.56 (0.31-0.93)	0.84% (0.56-1.2%)	0.79% (0.41-1.3%)	15 (7.8-25)	2000 (1300-3000)
Kenya	1990	Female	20 to 24	15% (9-22%)	85% (78-91%)	0.77 (0.43-1.2)	1.2% (0.82-1.6%)	0.79% (0.41-1.3%)	20 (11-34)	3100 (2000-4400)
Kenya	1990	Female	25 to 29	16% (9.9-24%)	84% (76-90%)	0.89 (0.51-1.4)	1.3% (0.95-1.8%)	1% (0.58-1.6%)	27 (15-42)	3300 (2300-4500)
Kenya	1990	Female	30 to 34	18% (11-28%)	82% (72-89%)	0.99 (0.58-1.5)	1.5% (1.1-2.1%)	1.4% (0.86-2.1%)	35 (22-54)	3300 (2300-4500)
Kenya	1990	Female	35 to 39	21% (12-30%)	79% (70-88%)	1.1 (0.59-1.7)	1.8% (1.2-2.5%)	1.9% (1.2-2.9%)	47 (28-71)	3300 (2300-4600)
Kenya	1990	Female	40 to 44	22% (14-32%)	78% (68-86%)	1.1 (0.59-1.7)	2% (1.3-2.8%)	2.4% (1.4-3.5%)	52 (31-78)	3000 (1900-4200)
Kenya	1990	Female	45 to 49	21% (12-30%)	79% (70-88%)	1.1 (0.58-1.8)	2.3% (1.4-3.4%)	2.8% (1.6-4.3%)	56 (33-89)	2700 (1700-4100)
Kenya	1990	Female	50 to 54	20% (13-30%)	80% (70-87%)	1.1 (0.57-1.8)	2.6% (1.5-4%)	3.2% (1.9-5.1%)	84 (48-140)	3300 (2000-5200)
Kenya	1990	Female	55 to 59	20% (12-29%)	80% (71-88%)	1.1 (0.56-1.8)	2.6% (1.4-4.1%)	3.1% (1.7-5.2%)	110 (57-170)	3500 (2000-5700)
Kenya	1990	Female	60 to 64	20% (12-30%)	80% (70-88%)	1.1 (0.54-1.8)	2.6% (1.4-4.3%)	3% (1.6-5.2%)	120 (65-200)	3400 (1900-5700)
Kenya	1990	Female	65 to 69	21% (14-30%)	79% (70-86%)	1 (0.57-1.7)	2.7% (1.6-4.3%)	3.1% (1.8-5%)	140 (80-220)	3300 (1900-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	1990	Female	70 to 74	20% (13-28%)	80% (72-87%)	0.9 (0.48-1.6)	2.2% (1.1-3.7%)	2.5% (1.2-4.2%)	120 (61-210)	2400 (1200-4100)
Kenya	1990	Female	75 to 79	18% (11-27%)	82% (73-89%)	0.77 (0.37-1.4)	1.9% (0.94-3.3%)	2.1% (1-3.8%)	110 (51-190)	1500 (760-2700)
Kenya	1990	Female	80 to 84	17% (10-25%)	83% (75-90%)	0.71 (0.36-1.2)	1.6% (0.72-2.9%)	1.8% (0.76-3.2%)	74 (32-130)	800 (350-1400)
Kenya	1990	Female	85 to 89	17% (11-26%)	83% (74-89%)	0.72 (0.37-1.2)	1.4% (0.56-2.5%)	1.5% (0.57-2.8%)	41 (16-78)	320 (130-590)
Kenya	1990	Female	90 to 94	19% (12-29%)	81% (71-88%)	0.79 (0.38-1.4)	1.2% (0.42-2.5%)	1.4% (0.41-2.9%)	16 (4.8-33)	85 (29-170)
Kenya	1990	Female	95 plus	23% (12-36%)	77% (64-88%)	0.9 (0.36-1.8)	1.4% (0.49-3.1%)	1.6% (0.46-3.5%)	5.1 (1.5-11)	18 (6.1-40)
Kenya	1990	Male	All Ages	26% (22-31%)	74% (69-78%)	0.82 (0.7-0.96)	1.2% (0.89-1.6%)	1.9% (1.2-2.6%)	2000 (1300-2800)	88000 (63000-110000)
Kenya	1990	Male	15 to 19	23% (15-33%)	77% (67-85%)	0.89 (0.53-1.4)	2% (1.3-3%)	2.2% (1.2-3.8%)	40 (21-70)	4500 (2800-6700)
Kenya	1990	Male	20 to 24	35% (23-48%)	65% (52-77%)	1.3 (0.72-2)	3.1% (2.1-4.5%)	2.8% (1.5-4.7%)	84 (46-140)	8500 (5500-12000)
Kenya	1990	Male	25 to 29	42% (28-57%)	58% (43-72%)	1.6 (0.93-2.3)	3.1% (2.1-4.3%)	2.9% (1.6-4.3%)	110 (61-160)	9300 (6100-13000)
Kenya	1990	Male	30 to 34	47% (31-62%)	53% (38-69%)	1.9 (1.1-2.7)	3.5% (2.4-4.8%)	3.5% (2-5.2%)	130 (75-190)	9300 (6200-13000)
Kenya	1990	Male	35 to 39	50% (34-64%)	50% (36-66%)	2.3 (1.4-3.6)	4.3% (2.8-6%)	4.7% (2.9-6.8%)	160 (99-230)	9700 (6600-14000)
Kenya	1990	Male	40 to 44	50% (36-65%)	50% (35-64%)	2.5 (1.5-3.8)	4.8% (3.2-6.7%)	5.6% (3.5-8%)	160 (100-240)	8600 (5700-12000)
Kenya	1990	Male	45 to 49	50% (34-63%)	50% (37-66%)	2.4 (1.3-3.8)	5.3% (3.4-7.4%)	6.3% (3.9-9%)	160 (95-220)	7000 (4500-9900)
Kenya	1990	Male	50 to 54	51% (37-66%)	49% (34-63%)	2.3 (1.3-3.7)	5.8% (3.7-8.3%)	7.1% (4.2-10%)	190 (110-280)	7600 (4500-11000)
Kenya	1990	Male	55 to 59	54% (41-68%)	46% (32-59%)	2.2 (1.3-3.6)	5.5% (3.4-8%)	6.6% (4-9.9%)	220 (130-330)	7500 (4700-11000)
Kenya	1990	Male	60 to 64	54% (39-69%)	46% (31-61%)	1.9 (1.1-3.3)	5% (2.9-7.7%)	5.9% (3.4-9.2%)	230 (130-360)	6500 (3900-10000)
Kenya	1990	Male	65 to 69	52% (37-65%)	48% (35-63%)	1.6 (0.93-2.5)	3.6% (2-5.7%)	4.2% (2.3-6.9%)	180 (99-290)	4200 (2400-6700)
Kenya	1990	Male	70 to 74	47% (34-60%)	53% (40-66%)	1.2 (0.63-1.9)	2.5% (1.3-4.3%)	2.9% (1.5-5%)	140 (69-230)	2500 (1300-4300)
Kenya	1990	Male	75 to 79	40% (28-55%)	60% (45-72%)	0.87 (0.43-1.5)	1.8% (0.84-3.2%)	2.1% (0.93-3.7%)	92 (41-170)	1300 (610-2400)
Kenya	1990	Male	80 to 84	36% (24-49%)	64% (51-76%)	0.72 (0.39-1.2)	1.3% (0.58-2.3%)	1.5% (0.62-2.6%)	48 (20-84)	520 (230-890)
Kenya	1990	Male	85 to 89	34% (23-48%)	66% (52-77%)	0.63 (0.34-1.1)	1.1% (0.51-1.9%)	1.2% (0.51-2.1%)	25 (11-43)	200 (88-340)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	1990	Male	90 to 94	35% (24-48%)	65% (52-76%)	0.61 (0.3-1)	0.74% (0.27-1.4%)	0.74% (0.2-1.5%)	6.5 (1.8-13)	38 (14-75)
Kenya	1990	Male	95 plus	39% (24-55%)	61% (45-76%)	0.62 (0.26-1.2)	0.88% (0.33-1.7%)	0.82% (0.19-1.8%)	1.9 (0.44-4.1)	8.2 (3-16)
Kenya	1995	Female	All Ages	11% (9.4-14%)	89% (86-91%)	0.48 (0.4-0.56)	0.62% (0.46-0.8%)	1.1% (0.7-1.5%)	1400 (880-2000)	51000 (37000-67000)
Kenya	1995	Female	15 to 19	11% (7.1-17%)	89% (83-93%)	0.57 (0.32-0.95)	0.84% (0.56-1.2%)	0.8% (0.42-1.3%)	20 (10-33)	2600 (1700-3700)
Kenya	1995	Female	20 to 24	15% (9-22%)	85% (78-91%)	0.78 (0.44-1.2)	0.89% (0.61-1.2%)	0.53% (0.27-0.87%)	25 (13-42)	3900 (2600-5400)
Kenya	1995	Female	25 to 29	16% (10-24%)	84% (76-90%)	0.91 (0.52-1.4)	0.79% (0.56-1.1%)	0.51% (0.3-0.8%)	34 (20-52)	4000 (2800-5500)
Kenya	1995	Female	30 to 34	18% (11-27%)	82% (73-89%)	1 (0.59-1.6)	0.9% (0.66-1.2%)	0.69% (0.43-1%)	45 (28-68)	4100 (2900-5600)
Kenya	1995	Female	35 to 39	21% (13-30%)	79% (70-87%)	1.1 (0.6-1.7)	1.1% (0.8-1.6%)	1.1% (0.67-1.6%)	62 (38-91)	4300 (2900-5900)
Kenya	1995	Female	40 to 44	22% (14-32%)	78% (68-86%)	1.1 (0.6-1.8)	1.5% (0.98-2%)	1.5% (0.95-2.2%)	78 (48-110)	4400 (2900-6100)
Kenya	1995	Female	45 to 49	21% (12-30%)	79% (70-88%)	1.1 (0.58-1.9)	1.8% (1.2-2.7%)	2.1% (1.2-3.1%)	89 (53-130)	4200 (2600-6100)
Kenya	1995	Female	50 to 54	20% (12-29%)	80% (71-88%)	1.1 (0.59-1.8)	2.4% (1.4-3.6%)	2.8% (1.6-4.3%)	100 (59-160)	4000 (2400-6100)
Kenya	1995	Female	55 to 59	20% (13-29%)	80% (71-87%)	1.1 (0.57-1.9)	2.5% (1.4-3.9%)	2.9% (1.7-4.8%)	130 (70-200)	4200 (2400-6700)
Kenya	1995	Female	60 to 64	20% (12-30%)	80% (70-88%)	1.1 (0.54-1.9)	2.6% (1.5-4.2%)	3% (1.7-5%)	160 (86-270)	4500 (2500-7300)
Kenya	1995	Female	65 to 69	21% (14-30%)	79% (70-86%)	1.1 (0.57-1.8)	2.8% (1.7-4.3%)	3.2% (1.9-5%)	190 (110-300)	4400 (2600-6800)
Kenya	1995	Female	70 to 74	20% (13-29%)	80% (71-87%)	0.92 (0.47-1.6)	2.3% (1.2-3.9%)	2.6% (1.3-4.4%)	160 (79-270)	3000 (1500-5100)
Kenya	1995	Female	75 to 79	18% (11-27%)	82% (73-89%)	0.79 (0.38-1.4)	2.1% (1.1-3.5%)	2.3% (1.2-4%)	140 (66-230)	2000 (990-3300)
Kenya	1995	Female	80 to 84	17% (10-25%)	83% (75-90%)	0.73 (0.37-1.2)	1.7% (0.79-3%)	1.9% (0.85-3.4%)	92 (41-160)	980 (450-1700)
Kenya	1995	Female	85 to 89	17% (11-26%)	83% (74-89%)	0.74 (0.38-1.3)	1.4% (0.61-2.5%)	1.6% (0.64-2.9%)	47 (19-86)	360 (150-640)
Kenya	1995	Female	90 to 94	19% (12-28%)	81% (72-88%)	0.81 (0.39-1.5)	1.3% (0.46-2.6%)	1.4% (0.46-3%)	18 (5.9-38)	98 (35-190)
Kenya	1995	Female	95 plus	23% (13-36%)	77% (64-87%)	0.92 (0.37-1.8)	1.5% (0.54-3.2%)	1.7% (0.5-3.6%)	6.3 (1.9-13)	23 (7.7-47)
Kenya	1995	Male	All Ages	27% (22-32%)	73% (68-78%)	0.87 (0.73-1)	1.3% (0.94-1.6%)	1.9% (1.3-2.5%)	2700 (1800-3600)	120000 (88000-150000)
Kenya	1995	Male	15 to 19	23% (15-33%)	77% (67-85%)	0.91 (0.53-1.4)	2.1% (1.4-3.1%)	2.4% (1.3-4.1%)	58 (30-97)	6000 (3800-8900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	1995	Male	20 to 24	34% (22-48%)	66% (52-78%)	1.3 (0.73-2)	2.9% (1.9-4.2%)	2.6% (1.5-4.3%)	120 (68-200)	12000 (7700-17000)
Kenya	1995	Male	25 to 29	42% (27-56%)	58% (44-73%)	1.6 (0.95-2.4)	2.3% (1.5-3.1%)	2% (1.2-3%)	160 (92-230)	13000 (8600-18000)
Kenya	1995	Male	30 to 34	47% (31-62%)	53% (38-69%)	1.9 (1.2-2.8)	2.2% (1.5-3%)	2% (1.2-2.9%)	170 (110-250)	12000 (8400-17000)
Kenya	1995	Male	35 to 39	50% (34-64%)	50% (36-66%)	2.4 (1.4-3.6)	2.9% (1.9-3.9%)	2.9% (1.8-4.1%)	220 (140-320)	13000 (8900-18000)
Kenya	1995	Male	40 to 44	50% (36-65%)	50% (35-64%)	2.6 (1.5-3.9)	3.3% (2.2-4.5%)	3.5% (2.2-4.9%)	250 (160-350)	13000 (8600-17000)
Kenya	1995	Male	45 to 49	50% (34-64%)	50% (36-66%)	2.5 (1.4-3.9)	4% (2.7-5.5%)	4.4% (2.8-6.1%)	250 (160-350)	11000 (7500-15000)
Kenya	1995	Male	50 to 54	51% (36-66%)	49% (34-64%)	2.4 (1.3-3.8)	5.1% (3.2-7%)	5.8% (3.6-8.2%)	240 (150-340)	9400 (6000-13000)
Kenya	1995	Male	55 to 59	54% (40-68%)	46% (32-60%)	2.3 (1.3-3.6)	5.3% (3.3-7.5%)	6.1% (3.8-8.8%)	270 (160-390)	9000 (5600-13000)
Kenya	1995	Male	60 to 64	54% (40-69%)	46% (31-60%)	2 (1.1-3.4)	4.9% (3-7.4%)	5.8% (3.4-8.7%)	310 (180-470)	8700 (5200-13000)
Kenya	1995	Male	65 to 69	51% (37-65%)	49% (35-63%)	1.6 (0.94-2.5)	3.7% (2.1-5.7%)	4.3% (2.4-6.7%)	240 (130-380)	5700 (3200-8800)
Kenya	1995	Male	70 to 74	46% (33-60%)	54% (40-67%)	1.2 (0.65-1.9)	2.6% (1.4-4.4%)	3% (1.6-5.1%)	180 (93-300)	3300 (1700-5600)
Kenya	1995	Male	75 to 79	40% (27-54%)	60% (46-73%)	0.89 (0.44-1.5)	1.9% (0.96-3.4%)	2.2% (1.1-3.9%)	120 (57-210)	1700 (840-3000)
Kenya	1995	Male	80 to 84	36% (24-49%)	64% (51-76%)	0.74 (0.4-1.2)	1.5% (0.67-2.4%)	1.6% (0.71-2.7%)	62 (27-110)	680 (310-1100)
Kenya	1995	Male	85 to 89	34% (23-48%)	66% (52-77%)	0.65 (0.35-1.1)	1.2% (0.56-2%)	1.3% (0.57-2.2%)	32 (14-55)	250 (120-420)
Kenya	1995	Male	90 to 94	35% (24-48%)	65% (52-76%)	0.62 (0.3-1.1)	0.81% (0.31-1.6%)	0.82% (0.25-1.7%)	8.8 (2.7-18)	51 (19-98)
Kenya	1995	Male	95 plus	39% (23-54%)	61% (46-77%)	0.63 (0.27-1.2)	0.97% (0.39-1.9%)	0.91% (0.27-1.9%)	2.5 (0.74-5.2)	11 (4.2-20)
Kenya	2000	Female	All Ages	13% (11-16%)	87% (84-89%)	0.5 (0.42-0.59)	0.61% (0.46-0.79%)	0.99% (0.63-1.4%)	1500 (980-2200)	60000 (44000-78000)
Kenya	2000	Female	15 to 19	14% (9.1-21%)	86% (79-91%)	0.64 (0.37-1)	1% (0.65-1.4%)	1% (0.57-1.8%)	27 (14-45)	3400 (2200-4900)
Kenya	2000	Female	20 to 24	17% (11-25%)	83% (75-89%)	0.81 (0.46-1.3)	1% (0.71-1.4%)	0.67% (0.35-1.1%)	34 (18-57)	4900 (3300-6900)
Kenya	2000	Female	25 to 29	18% (11-27%)	82% (73-89%)	0.92 (0.53-1.4)	0.65% (0.46-0.88%)	0.42% (0.24-0.65%)	43 (25-66)	4900 (3400-6700)
Kenya	2000	Female	30 to 34	21% (13-30%)	79% (70-87%)	1 (0.6-1.6)	0.67% (0.47-0.91%)	0.5% (0.32-0.75%)	55 (34-83)	4800 (3300-6600)
Kenya	2000	Female	35 to 39	23% (15-34%)	77% (66-85%)	1.1 (0.62-1.7)	0.84% (0.57-1.2%)	0.74% (0.46-1.1%)	75 (46-110)	5000 (3400-7000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	2000	Female	40 to 44	24% (15-35%)	76% (65-85%)	1.1 (0.59-1.8)	1.1% (0.72-1.5%)	1.1% (0.65-1.6%)	96 (57-140)	5300 (3400-7400)
Kenya	2000	Female	45 to 49	23% (13-32%)	77% (68-87%)	1.1 (0.54-1.8)	1.5% (0.91-2.2%)	1.6% (0.93-2.4%)	120 (70-180)	5600 (3500-8400)
Kenya	2000	Female	50 to 54	20% (13-30%)	80% (70-87%)	0.99 (0.52-1.7)	2% (1.2-3.1%)	2.3% (1.3-3.5%)	140 (82-220)	5500 (3300-8400)
Kenya	2000	Female	55 to 59	19% (12-28%)	81% (72-88%)	0.94 (0.48-1.6)	2.2% (1.2-3.5%)	2.6% (1.4-4.2%)	130 (69-200)	4200 (2400-6700)
Kenya	2000	Female	60 to 64	19% (12-28%)	81% (72-88%)	0.92 (0.46-1.6)	2.3% (1.2-3.7%)	2.6% (1.4-4.4%)	150 (82-250)	4300 (2400-7000)
Kenya	2000	Female	65 to 69	20% (13-29%)	80% (71-87%)	0.91 (0.49-1.5)	2.5% (1.5-3.9%)	2.9% (1.6-4.6%)	200 (110-320)	4800 (2700-7400)
Kenya	2000	Female	70 to 74	19% (13-28%)	81% (72-87%)	0.8 (0.41-1.4)	2.1% (1.1-3.5%)	2.4% (1.2-4%)	170 (85-290)	3300 (1700-5600)
Kenya	2000	Female	75 to 79	18% (11-26%)	82% (74-89%)	0.7 (0.33-1.2)	1.9% (0.96-3.2%)	2.1% (1.1-3.7%)	140 (70-240)	2000 (1000-3400)
Kenya	2000	Female	80 to 84	17% (10-25%)	83% (75-90%)	0.65 (0.33-1.1)	1.5% (0.69-2.7%)	1.7% (0.75-3%)	95 (41-160)	1000 (460-1700)
Kenya	2000	Female	85 to 89	17% (11-26%)	83% (74-89%)	0.66 (0.34-1.1)	1.3% (0.51-2.3%)	1.4% (0.56-2.7%)	46 (17-86)	350 (140-640)
Kenya	2000	Female	90 to 94	20% (12-29%)	80% (71-88%)	0.73 (0.35-1.3)	1.2% (0.4-2.4%)	1.3% (0.4-2.7%)	17 (5.1-36)	89 (31-180)
Kenya	2000	Female	95 plus	23% (13-36%)	77% (64-87%)	0.83 (0.33-1.6)	1.4% (0.51-3%)	1.5% (0.48-3.4%)	5.9 (1.8-13)	21 (7.4-44)
Kenya	2000	Male	All Ages	30% (25-35%)	70% (65-75%)	0.86 (0.72-1)	1.4% (1.1-1.8%)	2% (1.4-2.6%)	3400 (2300-4500)	150000 (120000-190000)
Kenya	2000	Male	15 to 19	28% (19-39%)	72% (61-81%)	0.96 (0.58-1.5)	2.4% (1.6-3.5%)	2.9% (1.6-4.5%)	85 (47-140)	8400 (5400-12000)
Kenya	2000	Male	20 to 24	38% (26-53%)	62% (47-74%)	1.3 (0.74-2)	3.7% (2.4-5.2%)	3.6% (2.1-5.7%)	180 (100-280)	16000 (11000-23000)
Kenya	2000	Male	25 to 29	45% (30-59%)	55% (41-70%)	1.5 (0.9-2.2)	2.5% (1.6-3.3%)	2.2% (1.3-3.2%)	220 (120-310)	17000 (11000-23000)
Kenya	2000	Male	30 to 34	49% (33-63%)	51% (37-67%)	1.8 (1.1-2.6)	2.1% (1.4-2.8%)	1.9% (1.2-2.8%)	240 (140-340)	16000 (11000-22000)
Kenya	2000	Male	35 to 39	52% (37-66%)	48% (34-63%)	2.2 (1.3-3.3)	2.3% (1.6-3.2%)	2.3% (1.5-3.3%)	280 (180-390)	16000 (11000-22000)
Kenya	2000	Male	40 to 44	52% (39-67%)	48% (33-61%)	2.4 (1.4-3.5)	2.9% (1.9-3.9%)	3% (1.9-4.1%)	320 (210-450)	16000 (11000-22000)
Kenya	2000	Male	45 to 49	52% (36-65%)	48% (35-64%)	2.3 (1.3-3.5)	3.4% (2.3-4.7%)	3.7% (2.4-5.1%)	350 (230-490)	16000 (10000-21000)
Kenya	2000	Male	50 to 54	53% (39-67%)	47% (33-61%)	2.2 (1.2-3.5)	4.6% (2.9-6.4%)	5.2% (3.2-7.2%)	370 (230-510)	14000 (8800-20000)
Kenya	2000	Male	55 to 59	55% (41-69%)	45% (31-59%)	2.1 (1.2-3.3)	5.1% (3.2-7.3%)	5.9% (3.6-8.5%)	320 (190-460)	10000 (6600-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	2000	Male	60 to 64	55% (40-70%)	45% (30-60%)	1.8 (0.98-3)	4.7% (2.9-7.2%)	5.4% (3.3-8.2%)	330 (200-500)	9200 (5600-14000)
Kenya	2000	Male	65 to 69	52% (37-66%)	48% (34-63%)	1.4 (0.84-2.2)	3.5% (2-5.5%)	4% (2.2-6.3%)	280 (150-450)	6600 (3700-10000)
Kenya	2000	Male	70 to 74	45% (32-59%)	55% (41-68%)	1 (0.55-1.6)	2.4% (1.2-4%)	2.7% (1.4-4.6%)	200 (100-330)	3600 (1900-6000)
Kenya	2000	Male	75 to 79	38% (26-52%)	62% (48-74%)	0.73 (0.36-1.3)	1.7% (0.9-3%)	2% (1-3.5%)	130 (64-220)	1800 (930-3100)
Kenya	2000	Male	80 to 84	34% (22-47%)	66% (53-78%)	0.61 (0.33-0.96)	1.3% (0.59-2.2%)	1.4% (0.6-2.5%)	66 (28-120)	710 (330-1200)
Kenya	2000	Male	85 to 89	33% (22-46%)	67% (54-78%)	0.54 (0.29-0.9)	1.1% (0.52-1.8%)	1.2% (0.54-2%)	32 (14-54)	250 (120-420)
Kenya	2000	Male	90 to 94	34% (22-46%)	66% (54-78%)	0.52 (0.26-0.91)	0.74% (0.3-1.4%)	0.75% (0.24-1.5%)	8.8 (2.8-17)	51 (21-96)
Kenya	2000	Male	95 plus	38% (22-53%)	62% (47-78%)	0.54 (0.22-1)	0.94% (0.43-1.8%)	0.91% (0.32-1.9%)	2.8 (0.99-5.8)	12 (5.3-22)
Kenya	2005	Female	All Ages	14% (11-16%)	86% (84-89%)	0.53 (0.45-0.62)	0.7% (0.52-0.9%)	1.1% (0.71-1.6%)	1800 (1100-2500)	68000 (50000-88000)
Kenya	2005	Female	15 to 19	14% (9.3-21%)	86% (79-91%)	0.65 (0.38-1)	0.93% (0.61-1.3%)	0.89% (0.47-1.5%)	28 (15-47)	3700 (2400-5300)
Kenya	2005	Female	20 to 24	18% (11-26%)	82% (74-89%)	0.85 (0.47-1.3)	1.3% (0.9-1.8%)	0.92% (0.48-1.5%)	39 (21-64)	5800 (3900-8100)
Kenya	2005	Female	25 to 29	19% (12-27%)	81% (73-88%)	0.96 (0.56-1.5)	0.88% (0.63-1.2%)	0.58% (0.33-0.9%)	50 (29-75)	5900 (4100-8000)
Kenya	2005	Female	30 to 34	21% (13-31%)	79% (69-87%)	1.1 (0.63-1.7)	0.71% (0.51-0.97%)	0.52% (0.32-0.79%)	60 (37-90)	5400 (3800-7500)
Kenya	2005	Female	35 to 39	24% (15-34%)	76% (66-85%)	1.1 (0.63-1.8)	0.83% (0.57-1.2%)	0.72% (0.45-1.1%)	77 (47-120)	5300 (3600-7400)
Kenya	2005	Female	40 to 44	24% (16-35%)	76% (65-84%)	1.1 (0.62-1.9)	1% (0.67-1.4%)	0.99% (0.61-1.5%)	98 (59-150)	5500 (3600-7800)
Kenya	2005	Female	45 to 49	23% (14-33%)	77% (67-86%)	1.1 (0.57-1.9)	1.4% (0.87-2.1%)	1.5% (0.88-2.3%)	130 (77-200)	6200 (3800-9200)
Kenya	2005	Female	50 to 54	21% (13-31%)	79% (69-87%)	1 (0.55-1.7)	2% (1.2-3.1%)	2.3% (1.3-3.7%)	180 (100-280)	6900 (4100-11000)
Kenya	2005	Female	55 to 59	19% (12-29%)	81% (71-88%)	0.98 (0.49-1.7)	2.3% (1.3-3.6%)	2.7% (1.5-4.3%)	170 (91-270)	5600 (3100-8800)
Kenya	2005	Female	60 to 64	20% (12-29%)	80% (71-88%)	0.96 (0.48-1.7)	2.4% (1.3-3.8%)	2.8% (1.5-4.5%)	160 (83-260)	4400 (2400-7100)
Kenya	2005	Female	65 to 69	20% (13-30%)	80% (70-87%)	0.94 (0.51-1.6)	2.6% (1.5-4.1%)	3% (1.7-4.8%)	210 (120-320)	4900 (2800-7500)
Kenya	2005	Female	70 to 74	20% (13-29%)	80% (71-87%)	0.84 (0.43-1.4)	2.2% (1.1-3.8%)	2.5% (1.2-4.3%)	200 (97-350)	3800 (1900-6500)
Kenya	2005	Female	75 to 79	18% (11-27%)	82% (73-89%)	0.72 (0.34-1.3)	2% (0.99-3.4%)	2.3% (1.1-3.9%)	180 (86-300)	2500 (1300-4300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	2005	Female	80 to 84	17% (11-26%)	83% (74-89%)	0.66 (0.33-1.1)	1.6% (0.74-2.8%)	1.9% (0.81-3.2%)	110 (49-200)	1200 (550-2100)
Kenya	2005	Female	85 to 89	18% (11-27%)	82% (73-89%)	0.68 (0.35-1.2)	1.4% (0.57-2.5%)	1.5% (0.61-2.9%)	56 (22-100)	420 (170-770)
Kenya	2005	Female	90 to 94	20% (12-29%)	80% (71-88%)	0.74 (0.35-1.3)	1.3% (0.46-2.5%)	1.4% (0.47-2.9%)	20 (6.7-42)	110 (38-220)
Kenya	2005	Female	95 plus	23% (13-36%)	77% (64-87%)	0.84 (0.33-1.6)	1.5% (0.55-3.1%)	1.7% (0.55-3.5%)	6.6 (2.2-14)	24 (8.5-48)
Kenya	2005	Male	All Ages	30% (25-35%)	70% (65-75%)	0.88 (0.74-1)	1.7% (1.3-2.2%)	2.4% (1.7-3.1%)	4100 (2900-5400)	190000 (140000-240000)
Kenya	2005	Male	15 to 19	27% (19-38%)	73% (62-81%)	0.93 (0.55-1.4)	2.2% (1.4-3.2%)	2.5% (1.4-3.9%)	100 (56-160)	9800 (6200-14000)
Kenya	2005	Male	20 to 24	38% (25-51%)	62% (49-75%)	1.2 (0.71-1.9)	4.3% (2.8-6.1%)	4.4% (2.5-6.9%)	230 (130-360)	20000 (13000-30000)
Kenya	2005	Male	25 to 29	44% (30-58%)	56% (42-70%)	1.5 (0.9-2.2)	3.7% (2.5-5%)	3.6% (2.2-5.2%)	280 (170-410)	22000 (15000-30000)
Kenya	2005	Male	30 to 34	49% (33-63%)	51% (37-67%)	1.8 (1.1-2.6)	2.7% (1.9-3.7%)	2.6% (1.6-3.7%)	300 (180-430)	21000 (14000-28000)
Kenya	2005	Male	35 to 39	52% (37-66%)	48% (34-63%)	2.2 (1.3-3.4)	2.8% (1.9-3.8%)	2.8% (1.8-4%)	340 (220-480)	20000 (14000-27000)
Kenya	2005	Male	40 to 44	52% (38-67%)	48% (33-62%)	2.3 (1.3-3.6)	3% (2.1-4.1%)	3.1% (2.1-4.4%)	360 (240-510)	18000 (12000-25000)
Kenya	2005	Male	45 to 49	51% (35-65%)	49% (35-65%)	2.2 (1.2-3.5)	3.8% (2.5-5.2%)	4.1% (2.7-5.6%)	420 (280-580)	18000 (12000-25000)
Kenya	2005	Male	50 to 54	52% (38-67%)	48% (33-62%)	2.1 (1.2-3.4)	4.9% (3.1-6.7%)	5.5% (3.4-7.6%)	490 (310-680)	19000 (12000-26000)
Kenya	2005	Male	55 to 59	55% (41-69%)	45% (31-59%)	2 (1.2-3.2)	5.5% (3.5-7.8%)	6.4% (3.9-9.2%)	450 (280-640)	15000 (9200-21000)
Kenya	2005	Male	60 to 64	55% (40-70%)	45% (30-60%)	1.8 (0.96-2.9)	5% (3-7.6%)	5.8% (3.4-8.9%)	350 (210-540)	9900 (6000-15000)
Kenya	2005	Male	65 to 69	52% (37-65%)	48% (35-63%)	1.4 (0.81-2.2)	3.7% (2.1-5.7%)	4.2% (2.4-6.7%)	290 (160-460)	6800 (3900-11000)
Kenya	2005	Male	70 to 74	45% (32-58%)	55% (42-68%)	0.98 (0.54-1.6)	2.4% (1.3-4%)	2.8% (1.4-4.6%)	220 (110-370)	4100 (2100-6700)
Kenya	2005	Male	75 to 79	38% (25-52%)	62% (48-75%)	0.72 (0.36-1.2)	1.8% (0.92-3.1%)	2.1% (1-3.6%)	150 (74-260)	2100 (1100-3600)
Kenya	2005	Male	80 to 84	34% (22-46%)	66% (54-78%)	0.59 (0.3-0.94)	1.3% (0.64-2.2%)	1.5% (0.64-2.5%)	78 (34-140)	850 (390-1400)
Kenya	2005	Male	85 to 89	32% (21-45%)	68% (55-79%)	0.52 (0.28-0.87)	1.1% (0.56-1.9%)	1.2% (0.58-2.1%)	37 (17-65)	290 (140-490)
Kenya	2005	Male	90 to 94	33% (22-46%)	67% (54-78%)	0.5 (0.24-0.87)	0.79% (0.33-1.5%)	0.81% (0.29-1.6%)	9.8 (3.5-19)	57 (24-110)
Kenya	2005	Male	95 plus	37% (21-52%)	63% (48-79%)	0.51 (0.21-0.97)	1% (0.48-1.9%)	1% (0.4-2.1%)	3.3 (1.3-6.7)	13 (6.1-25)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	2010	Female	All Ages	11% (8.9-13%)	89% (87-91%)	0.51 (0.43-0.6)	0.92% (0.69-1.2%)	1.6% (1.1-2.2%)	2200 (1400-3100)	79000 (58000-100000)
Kenya	2010	Female	15 to 19	9.5% (5.9-15%)	91% (85-94%)	0.52 (0.28-0.86)	0.72% (0.48-1%)	0.58% (0.29-0.97%)	20 (10-34)	3100 (2000-4500)
Kenya	2010	Female	20 to 24	13% (7.4-19%)	87% (81-93%)	0.73 (0.39-1.2)	1.3% (0.9-1.8%)	0.87% (0.44-1.4%)	34 (17-55)	5800 (3900-8100)
Kenya	2010	Female	25 to 29	14% (8.4-21%)	86% (79-92%)	0.86 (0.48-1.3)	1.1% (0.83-1.5%)	0.76% (0.45-1.2%)	48 (28-74)	6400 (4500-8600)
Kenya	2010	Female	30 to 34	15% (9.2-23%)	85% (77-91%)	0.93 (0.54-1.5)	1% (0.73-1.4%)	0.77% (0.47-1.2%)	61 (38-92)	6000 (4300-8300)
Kenya	2010	Female	35 to 39	17% (10-25%)	83% (75-90%)	0.97 (0.55-1.6)	1.1% (0.75-1.5%)	0.96% (0.6-1.4%)	78 (47-120)	5600 (3900-7700)
Kenya	2010	Female	40 to 44	18% (11-28%)	82% (72-89%)	1 (0.55-1.7)	1.3% (0.89-1.8%)	1.3% (0.83-2%)	94 (58-140)	5400 (3600-7600)
Kenya	2010	Female	45 to 49	19% (11-27%)	81% (73-89%)	1.1 (0.55-1.9)	1.8% (1.2-2.6%)	2% (1.2-3%)	130 (81-200)	6200 (4000-9200)
Kenya	2010	Female	50 to 54	19% (12-28%)	81% (72-88%)	1.1 (0.61-1.9)	2.6% (1.6-4%)	3.1% (1.9-4.9%)	200 (120-310)	7800 (4800-12000)
Kenya	2010	Female	55 to 59	20% (13-30%)	80% (70-87%)	1.2 (0.65-2.1)	3.1% (1.7-4.7%)	3.8% (2.1-5.9%)	240 (130-370)	8000 (4500-12000)
Kenya	2010	Female	60 to 64	22% (14-32%)	78% (68-86%)	1.3 (0.68-2.2)	3.2% (1.8-5.1%)	3.9% (2.2-6.3%)	250 (140-390)	7000 (4000-11000)
Kenya	2010	Female	65 to 69	23% (15-33%)	77% (67-85%)	1.3 (0.71-2.1)	3.5% (2-5.3%)	4.2% (2.4-6.5%)	260 (150-390)	6000 (3500-9000)
Kenya	2010	Female	70 to 74	21% (14-31%)	79% (69-86%)	1.1 (0.59-1.9)	2.9% (1.5-4.8%)	3.4% (1.7-5.6%)	250 (130-420)	4800 (2500-7900)
Kenya	2010	Female	75 to 79	19% (12-28%)	81% (72-88%)	0.93 (0.46-1.6)	2.6% (1.3-4.2%)	3% (1.5-4.9%)	250 (120-410)	3600 (1800-5900)
Kenya	2010	Female	80 to 84	18% (11-26%)	82% (74-89%)	0.84 (0.43-1.4)	2.1% (1-3.5%)	2.4% (1.2-4.1%)	170 (79-280)	1800 (870-3000)
Kenya	2010	Female	85 to 89	19% (11-28%)	81% (72-89%)	0.86 (0.44-1.4)	1.8% (0.8-3.2%)	2.1% (0.86-3.7%)	89 (37-160)	680 (300-1200)
Kenya	2010	Female	90 to 94	21% (13-30%)	79% (70-87%)	0.94 (0.44-1.7)	1.7% (0.66-3.2%)	1.9% (0.69-3.8%)	34 (12-66)	180 (68-340)
Kenya	2010	Female	95 plus	25% (13-39%)	75% (61-87%)	1.1 (0.43-2)	2% (0.78-4%)	2.2% (0.79-4.5%)	11 (3.9-22)	38 (15-77)
Kenya	2010	Male	All Ages	27% (23-32%)	73% (68-77%)	0.96 (0.81-1.2)	2.4% (1.9-3%)	3.4% (2.5-4.4%)	5300 (3900-6800)	230000 (180000-290000)
Kenya	2010	Male	15 to 19	18% (11-27%)	82% (73-89%)	0.73 (0.41-1.2)	1.7% (1.1-2.5%)	1.8% (0.95-2.9%)	82 (44-130)	8400 (5200-12000)
Kenya	2010	Male	20 to 24	31% (20-44%)	69% (56-80%)	1.2 (0.64-1.9)	4.3% (2.7-6.2%)	4.5% (2.6-7.2%)	240 (140-380)	22000 (14000-32000)
Kenya	2010	Male	25 to 29	41% (27-55%)	59% (45-73%)	1.6 (0.98-2.4)	4.8% (3.3-6.4%)	5.1% (3.1-7.2%)	360 (220-500)	28000 (19000-37000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	2010	Male	30 to 34	47% (31-62%)	53% (38-69%)	2 (1.2-2.9)	4.6% (3.2-6.1%)	4.8% (3.1-6.8%)	410 (260-570)	27000 (19000-37000)
Kenya	2010	Male	35 to 39	50% (34-64%)	50% (36-66%)	2.5 (1.5-3.8)	4.5% (3.3-6.1%)	4.9% (3.4-6.8%)	460 (320-640)	27000 (19000-36000)
Kenya	2010	Male	40 to 44	49% (36-64%)	51% (36-64%)	2.6 (1.5-4.1)	4.8% (3.5-6.6%)	5.4% (3.7-7.4%)	470 (320-650)	24000 (17000-32000)
Kenya	2010	Male	45 to 49	48% (32-63%)	52% (37-68%)	2.5 (1.4-4)	5.5% (3.8-7.4%)	6.3% (4.2-8.4%)	490 (330-660)	22000 (15000-29000)
Kenya	2010	Male	50 to 54	50% (35-66%)	50% (34-65%)	2.4 (1.4-3.8)	6.9% (4.5-9.4%)	8.1% (5.2-11%)	590 (380-810)	23000 (15000-31000)
Kenya	2010	Male	55 to 59	55% (41-69%)	45% (31-59%)	2.4 (1.4-3.7)	7.3% (4.8-10%)	8.8% (5.7-12%)	630 (410-880)	21000 (14000-29000)
Kenya	2010	Male	60 to 64	56% (41-70%)	44% (30-59%)	2.1 (1.2-3.5)	6.4% (4-9.5%)	7.7% (4.8-11%)	540 (340-800)	15000 (9600-22000)
Kenya	2010	Male	65 to 69	53% (38-66%)	47% (34-62%)	1.7 (0.99-2.7)	4.8% (2.8-7.2%)	5.6% (3.3-8.6%)	370 (220-560)	8500 (5100-13000)
Kenya	2010	Male	70 to 74	46% (33-59%)	54% (41-67%)	1.2 (0.67-1.9)	3.2% (1.6-5.1%)	3.7% (1.9-6%)	270 (140-450)	5100 (2700-8200)
Kenya	2010	Male	75 to 79	39% (26-53%)	61% (47-74%)	0.88 (0.45-1.5)	2.3% (1.2-4%)	2.7% (1.4-4.6%)	210 (100-350)	3000 (1500-5000)
Kenya	2010	Male	80 to 84	34% (23-47%)	66% (53-77%)	0.71 (0.38-1.1)	1.7% (0.82-2.8%)	1.9% (0.89-3.3%)	120 (53-200)	1200 (600-2100)
Kenya	2010	Male	85 to 89	33% (22-46%)	67% (54-78%)	0.63 (0.33-1.1)	1.4% (0.72-2.5%)	1.6% (0.77-2.8%)	55 (26-95)	430 (220-720)
Kenya	2010	Male	90 to 94	34% (23-46%)	66% (54-77%)	0.6 (0.29-1.1)	1% (0.46-2%)	1.1% (0.43-2.1%)	15 (5.6-28)	82 (35-150)
Kenya	2010	Male	95 plus	37% (22-54%)	63% (46-78%)	0.62 (0.26-1.2)	1.3% (0.6-2.5%)	1.4% (0.54-2.7%)	4.5 (1.8-9.1)	18 (8-33)
Kenya	2016	Female	All Ages	11% (9.2-13%)	89% (87-91%)	0.55 (0.47-0.66)	1.2% (0.94-1.6%)	2.2% (1.4-3.1%)	2700 (1700-3700)	94000 (69000-120000)
Kenya	2016	Female	15 to 19	9.6% (6-15%)	90% (85-94%)	0.55 (0.29-0.92)	0.82% (0.55-1.1%)	0.71% (0.35-1.2%)	21 (10-36)	3400 (2200-4900)
Kenya	2016	Female	20 to 24	13% (7.7-19%)	87% (81-92%)	0.77 (0.4-1.3)	1.4% (0.99-1.9%)	0.94% (0.48-1.6%)	31 (15-51)	5800 (3900-8200)
Kenya	2016	Female	25 to 29	14% (8.4-21%)	86% (79-92%)	0.91 (0.51-1.4)	1.4% (1.1-1.9%)	1.1% (0.63-1.6%)	52 (31-80)	7200 (5000-9800)
Kenya	2016	Female	30 to 34	15% (9.5-23%)	85% (77-91%)	0.99 (0.58-1.5)	1.5% (1.1-2%)	1.3% (0.83-2%)	77 (47-120)	7800 (5500-11000)
Kenya	2016	Female	35 to 39	17% (10-26%)	83% (74-90%)	1 (0.6-1.6)	1.7% (1.2-2.4%)	1.9% (1.2-2.8%)	110 (67-160)	7900 (5500-11000)
Kenya	2016	Female	40 to 44	18% (12-28%)	82% (72-88%)	1.1 (0.6-1.8)	2% (1.3-2.8%)	2.3% (1.4-3.5%)	120 (76-180)	7100 (4800-10000)
Kenya	2016	Female	45 to 49	19% (11-28%)	81% (72-89%)	1.2 (0.58-2)	2.4% (1.6-3.5%)	3.1% (1.9-4.6%)	150 (92-220)	7100 (4700-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	2016	Female	50 to 54	20% (12-29%)	80% (71-88%)	1.2 (0.65-2)	3% (1.9-4.6%)	4% (2.4-6.1%)	210 (130-320)	8300 (5200-12000)
Kenya	2016	Female	55 to 59	21% (13-30%)	79% (70-87%)	1.3 (0.68-2.2)	3.4% (2-5.1%)	4.4% (2.5-6.8%)	270 (150-420)	9000 (5300-14000)
Kenya	2016	Female	60 to 64	22% (14-33%)	78% (67-86%)	1.4 (0.71-2.4)	3.4% (1.9-5.2%)	4.3% (2.4-6.8%)	310 (170-500)	9000 (5100-14000)
Kenya	2016	Female	65 to 69	23% (15-33%)	77% (67-85%)	1.4 (0.74-2.2)	3.7% (2.2-5.4%)	4.5% (2.6-6.8%)	350 (210-530)	8400 (5000-12000)
Kenya	2016	Female	70 to 74	22% (15-31%)	78% (69-85%)	1.2 (0.63-2)	3% (1.6-4.9%)	3.6% (1.8-5.9%)	270 (140-440)	5200 (2800-8300)
Kenya	2016	Female	75 to 79	19% (12-29%)	81% (71-88%)	0.99 (0.49-1.7)	2.7% (1.4-4.4%)	3.2% (1.7-5.2%)	260 (130-430)	3800 (2000-6200)
Kenya	2016	Female	80 to 84	18% (11-27%)	82% (73-89%)	0.89 (0.45-1.5)	2.4% (1.2-3.8%)	2.8% (1.4-4.5%)	210 (110-340)	2200 (1100-3600)
Kenya	2016	Female	85 to 89	19% (11-28%)	81% (72-89%)	0.91 (0.47-1.5)	2.1% (1-3.5%)	2.4% (1.2-4.1%)	140 (65-230)	1000 (500-1700)
Kenya	2016	Female	90 to 94	21% (13-31%)	79% (69-87%)	1 (0.47-1.8)	2% (0.85-3.6%)	2.3% (0.93-4.2%)	57 (23-100)	300 (130-530)
Kenya	2016	Female	95 plus	25% (14-39%)	75% (61-86%)	1.1 (0.46-2.2)	2.4% (1-4.3%)	2.7% (1.1-5.1%)	19 (7.8-37)	67 (29-130)
Kenya	2016	Male	All Ages	28% (23-33%)	72% (67-77%)	1 (0.87-1.3)	3% (2.4-3.7%)	4.2% (3.1-5.4%)	6200 (4500-8000)	270000 (210000-340000)
Kenya	2016	Male	15 to 19	18% (11-27%)	82% (73-89%)	0.75 (0.41-1.2)	1.8% (1.2-2.7%)	2% (1-3.1%)	88 (47-140)	9200 (5900-13000)
Kenya	2016	Male	20 to 24	31% (20-44%)	69% (56-80%)	1.2 (0.66-2)	4.3% (2.8-6.3%)	4.5% (2.5-7.1%)	230 (130-370)	22000 (14000-31000)
Kenya	2016	Male	25 to 29	41% (27-55%)	59% (45-73%)	1.7 (0.98-2.5)	5.1% (3.5-6.7%)	5.5% (3.4-7.7%)	380 (230-530)	30000 (20000-40000)
Kenya	2016	Male	30 to 34	47% (31-61%)	53% (39-69%)	2.1 (1.3-3)	5.2% (3.6-6.9%)	5.7% (3.7-7.9%)	470 (300-650)	33000 (23000-43000)
Kenya	2016	Male	35 to 39	49% (34-64%)	51% (36-66%)	2.5 (1.5-3.8)	5.8% (4.3-7.8%)	6.7% (4.7-9.2%)	580 (400-780)	34000 (24000-45000)
Kenya	2016	Male	40 to 44	49% (35-64%)	51% (36-65%)	2.7 (1.5-4.1)	6.1% (4.3-8.2%)	7.2% (5-9.9%)	600 (410-820)	31000 (22000-41000)
Kenya	2016	Male	45 to 49	48% (33-63%)	52% (37-67%)	2.6 (1.4-4.1)	6.6% (4.6-8.8%)	7.9% (5.5-11%)	600 (400-810)	27000 (19000-35000)
Kenya	2016	Male	50 to 54	50% (35-65%)	50% (35-65%)	2.5 (1.4-3.9)	7.3% (4.8-10%)	9% (5.8-12%)	640 (410-890)	25000 (16000-34000)
Kenya	2016	Male	55 to 59	54% (40-68%)	46% (32-60%)	2.5 (1.4-3.8)	7.6% (5.1-11%)	9.5% (6.2-13%)	690 (460-960)	23000 (15000-32000)
Kenya	2016	Male	60 to 64	55% (40-70%)	45% (30-60%)	2.2 (1.2-3.6)	6.6% (4.1-9.6%)	8% (5-12%)	680 (420-1000)	19000 (12000-28000)
Kenya	2016	Male	65 to 69	53% (38-66%)	47% (34-62%)	1.7 (1-2.7)	4.8% (2.9-7.3%)	5.8% (3.4-8.8%)	500 (300-760)	12000 (6900-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Kenya	2016	Male	70 to 74	46% (33-59%)	54% (41-67%)	1.2 (0.7-2)	3.2% (1.7-5.1%)	3.8% (2-6.1%)	300 (150-480)	5500 (2900-8800)
Kenya	2016	Male	75 to 79	39% (26-53%)	61% (47-74%)	0.9 (0.46-1.5)	2.4% (1.2-4%)	2.8% (1.4-4.7%)	210 (100-360)	3000 (1500-5100)
Kenya	2016	Male	80 to 84	34% (22-47%)	66% (53-78%)	0.73 (0.38-1.2)	1.8% (0.88-3%)	2% (0.97-3.5%)	130 (63-230)	1400 (710-2500)
Kenya	2016	Male	85 to 89	33% (22-45%)	67% (55-78%)	0.64 (0.34-1)	1.5% (0.79-2.5%)	1.7% (0.84-2.9%)	74 (37-130)	570 (300-950)
Kenya	2016	Male	90 to 94	34% (23-46%)	66% (54-77%)	0.62 (0.29-1.1)	1.1% (0.51-2%)	1.2% (0.48-2.2%)	20 (8.4-39)	110 (53-210)
Kenya	2016	Male	95 plus	37% (23-54%)	63% (46-77%)	0.64 (0.27-1.2)	1.4% (0.68-2.6%)	1.5% (0.63-2.8%)	6.2 (2.6-12)	24 (11-44)
Madagascar	1990	Female	All Ages	20% (16-26%)	80% (74-84%)	0.64 (0.49-0.81)	1.1% (0.72-1.5%)	2% (1.2-3.1%)	1400 (790-2100)	52000 (34000-71000)
Madagascar	1990	Female	15 to 19	21% (12-34%)	79% (66-88%)	0.73 (0.36-1.3)	1.5% (0.77-2.6%)	1.9% (0.78-3.5%)	23 (9.3-42)	2100 (1100-3600)
Madagascar	1990	Female	20 to 24	26% (14-42%)	74% (58-86%)	1 (0.51-1.7)	2.6% (1.5-4.1%)	3% (1.4-5.1%)	42 (19-73)	3800 (2100-5900)
Madagascar	1990	Female	25 to 29	28% (15-45%)	72% (55-85%)	1.3 (0.67-2.2)	3.7% (2.2-5.7%)	4.5% (2.4-7.3%)	63 (33-100)	4700 (2900-7400)
Madagascar	1990	Female	30 to 34	31% (17-49%)	69% (51-83%)	1.4 (0.7-2.3)	4.3% (2.5-6.6%)	5.5% (2.8-8.7%)	78 (41-120)	5100 (2900-7700)
Madagascar	1990	Female	35 to 39	34% (20-52%)	66% (48-80%)	1.4 (0.73-2.3)	4.3% (2.5-6.6%)	5.5% (3-8.7%)	83 (45-130)	4700 (2700-7200)
Madagascar	1990	Female	40 to 44	36% (21-54%)	64% (46-79%)	1.4 (0.68-2.5)	4.4% (2.4-7.1%)	5.7% (2.9-9.4%)	79 (40-130)	4000 (2200-6500)
Madagascar	1990	Female	45 to 49	36% (20-54%)	64% (46-80%)	1.4 (0.66-2.6)	4.8% (2.4-8.5%)	6.2% (3-11%)	98 (46-180)	4300 (2100-7300)
Madagascar	1990	Female	50 to 54	35% (19-54%)	65% (46-81%)	1.4 (0.65-2.6)	5.2% (2.5-8.9%)	6.6% (3.2-11%)	140 (62-240)	5100 (2400-8800)
Madagascar	1990	Female	55 to 59	33% (19-52%)	67% (48-81%)	1.4 (0.62-2.6)	5.1% (2.3-9.3%)	6.3% (2.8-11%)	160 (69-280)	5100 (2200-9000)
Madagascar	1990	Female	60 to 64	34% (18-52%)	66% (48-82%)	1.4 (0.6-2.6)	4.9% (2-8.8%)	5.9% (2.4-10%)	170 (67-300)	4600 (1800-8300)
Madagascar	1990	Female	65 to 69	35% (20-53%)	65% (47-80%)	1.3 (0.57-2.3)	4.7% (2-8.2%)	5.5% (2.3-9.8%)	170 (70-310)	3900 (1600-6900)
Madagascar	1990	Female	70 to 74	33% (20-51%)	67% (49-80%)	1.1 (0.47-2.1)	3.7% (1.1-7.5%)	4.2% (1.2-8.6%)	120 (34-260)	2200 (630-4600)
Madagascar	1990	Female	75 to 79	30% (16-49%)	70% (51-84%)	0.91 (0.36-1.9)	2.8% (0.65-6.2%)	3.2% (0.73-7.1%)	93 (21-210)	1300 (310-2900)
Madagascar	1990	Female	80 to 84	29% (15-47%)	71% (53-85%)	0.82 (0.33-1.6)	2% (0.47-4.8%)	2.3% (0.51-5.5%)	57 (12-130)	590 (140-1400)
Madagascar	1990	Female	85 to 89	29% (16-46%)	71% (54-84%)	0.8 (0.32-1.6)	1.5% (0.2-3.8%)	1.7% (0.2-4.4%)	25 (3.1-64)	190 (25-470)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	1990	Female	90 to 94	32% (18-50%)	68% (50-82%)	0.83 (0.34-1.7)	1.1% (-0.094-3.4%)	1.3% (-0.16-3.9%)	6.3 (-0.83-20)	34 (-2.6-100)
Madagascar	1990	Female	95 plus	37% (18-58%)	63% (42-82%)	0.92 (0.32-2)	1.2% (-0.16-4%)	1.3% (-0.27-4.4%)	1.6 (-0.33-5.5)	5.8 (-0.74-19)
Madagascar	1990	Male	All Ages	30% (24-37%)	70% (63-76%)	0.94 (0.73-1.2)	1.9% (1.3-2.5%)	2.9% (1.9-4.1%)	2300 (1500-3200)	100000 (72000-130000)
Madagascar	1990	Male	15 to 19	27% (16-42%)	73% (58-84%)	0.96 (0.49-1.7)	3% (1.8-4.7%)	3.3% (1.7-5.7%)	45 (22-77)	4400 (2600-6800)
Madagascar	1990	Male	20 to 24	38% (23-55%)	62% (45-77%)	1.4 (0.75-2.3)	5.9% (3.6-8.7%)	6.1% (3.1-9.7%)	110 (53-170)	9300 (5600-14000)
Madagascar	1990	Male	25 to 29	43% (26-61%)	57% (39-74%)	1.8 (1-2.8)	7.8% (4.9-11%)	8.6% (4.8-13%)	150 (81-220)	11000 (6800-16000)
Madagascar	1990	Male	30 to 34	47% (29-65%)	53% (35-71%)	2.1 (1.2-3.2)	9% (5.8-13%)	10% (6-16%)	180 (100-280)	12000 (7400-17000)
Madagascar	1990	Male	35 to 39	50% (32-68%)	50% (32-68%)	2.3 (1.3-3.6)	9.5% (5.9-14%)	11% (6.7-17%)	210 (120-310)	12000 (7200-17000)
Madagascar	1990	Male	40 to 44	51% (34-70%)	49% (30-66%)	2.4 (1.2-3.8)	9.6% (5.8-14%)	12% (6.5-17%)	190 (110-290)	9600 (5700-14000)
Madagascar	1990	Male	45 to 49	52% (32-70%)	48% (30-68%)	2.4 (1.2-4)	9.4% (5.4-14%)	11% (6.2-18%)	210 (110-330)	8900 (4900-14000)
Madagascar	1990	Male	50 to 54	52% (33-70%)	48% (30-67%)	2.3 (1.2-3.9)	9% (4.9-14%)	11% (5.8-17%)	240 (130-380)	9300 (4900-14000)
Madagascar	1990	Male	55 to 59	51% (33-70%)	49% (30-67%)	2.2 (1.2-3.8)	8.1% (4.3-13%)	9.6% (4.9-15%)	260 (130-430)	8700 (4500-14000)
Madagascar	1990	Male	60 to 64	50% (31-69%)	50% (31-69%)	1.9 (0.93-3.4)	6.8% (3.3-12%)	7.9% (3.7-14%)	260 (120-450)	7100 (3500-12000)
Madagascar	1990	Male	65 to 69	47% (29-65%)	53% (35-71%)	1.6 (0.76-2.7)	4.5% (1.9-8.1%)	5.1% (2.2-9.3%)	180 (78-340)	4200 (1900-7700)
Madagascar	1990	Male	70 to 74	44% (27-62%)	56% (38-73%)	1.2 (0.57-2.1)	2.9% (0.98-6.1%)	3.3% (1.1-7%)	120 (38-250)	2200 (730-4600)
Madagascar	1990	Male	75 to 79	42% (25-60%)	58% (40-75%)	0.99 (0.45-1.9)	2% (0.49-4.7%)	2.3% (0.52-5.3%)	72 (17-170)	1000 (250-2300)
Madagascar	1990	Male	80 to 84	39% (23-58%)	61% (42-77%)	0.84 (0.39-1.5)	1.5% (0.31-3.3%)	1.6% (0.29-3.6%)	34 (6-76)	370 (80-800)
Madagascar	1990	Male	85 to 89	37% (21-54%)	63% (46-79%)	0.7 (0.32-1.3)	1% (0.11-2.2%)	1% (0.028-2.4%)	13 (0.34-29)	100 (11-230)
Madagascar	1990	Male	90 to 94	37% (23-55%)	63% (45-77%)	0.64 (0.27-1.2)	0.48% (-0.24-1.5%)	0.4% (-0.43-1.5%)	1.6 (-1.7-6)	11 (-5.2-33)
Madagascar	1990	Male	95 plus	41% (21-62%)	59% (38-79%)	0.64 (0.23-1.4)	0.44% (-0.42-1.7%)	0.27% (-0.71-1.6%)	0.23 (-0.61-1.4)	1.5 (-1.4-5.5)
Madagascar	1995	Female	All Ages	20% (16-26%)	80% (74-84%)	0.57 (0.43-0.72)	1% (0.64-1.4%)	1.8% (0.97-2.9%)	1400 (710-2200)	51000 (33000-72000)
Madagascar	1995	Female	15 to 19	20% (11-33%)	80% (67-89%)	0.64 (0.31-1.1)	1.3% (0.64-2.2%)	1.6% (0.57-2.9%)	21 (7.4-39)	2100 (1000-3500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	1995	Female	20 to 24	26% (14-41%)	74% (59-86%)	0.91 (0.44-1.5)	2.2% (1.3-3.4%)	2.4% (1-4.1%)	38 (17-66)	3700 (2000-5700)
Madagascar	1995	Female	25 to 29	28% (15-45%)	72% (55-85%)	1.1 (0.57-1.9)	3.2% (1.9-4.8%)	3.7% (1.9-6%)	58 (31-95)	4600 (2800-7000)
Madagascar	1995	Female	30 to 34	31% (17-49%)	69% (51-83%)	1.2 (0.64-2)	3.7% (2.1-5.6%)	4.7% (2.3-7.4%)	74 (38-120)	5000 (2900-7500)
Madagascar	1995	Female	35 to 39	34% (20-53%)	66% (47-80%)	1.2 (0.63-2)	3.7% (2.1-5.7%)	4.7% (2.5-7.7%)	81 (42-130)	4700 (2700-7500)
Madagascar	1995	Female	40 to 44	36% (21-55%)	64% (45-79%)	1.2 (0.59-2.2)	3.8% (2-6.2%)	4.9% (2.4-8.2%)	89 (43-150)	4500 (2400-7400)
Madagascar	1995	Female	45 to 49	36% (20-54%)	64% (46-80%)	1.2 (0.57-2.3)	4.1% (2-7.2%)	5.3% (2.4-9.6%)	94 (42-170)	4100 (1900-7200)
Madagascar	1995	Female	50 to 54	34% (19-54%)	66% (46-81%)	1.2 (0.55-2.3)	4.5% (2-8%)	5.7% (2.5-10%)	120 (53-220)	4700 (2100-8200)
Madagascar	1995	Female	55 to 59	33% (19-53%)	67% (47-81%)	1.2 (0.54-2.3)	4.4% (1.8-8.4%)	5.4% (2.3-10%)	150 (59-290)	4900 (1900-9200)
Madagascar	1995	Female	60 to 64	34% (19-52%)	66% (48-81%)	1.2 (0.52-2.3)	4.2% (1.5-8%)	5.1% (1.8-9.6%)	180 (64-330)	4800 (1800-9000)
Madagascar	1995	Female	65 to 69	35% (19-53%)	65% (47-81%)	1.1 (0.49-2)	4% (1.5-7.3%)	4.7% (1.7-8.5%)	180 (64-330)	4000 (1500-7400)
Madagascar	1995	Female	70 to 74	34% (19-52%)	66% (48-81%)	0.97 (0.42-1.9)	3.1% (0.7-6.8%)	3.5% (0.74-7.7%)	130 (28-290)	2400 (550-5300)
Madagascar	1995	Female	75 to 79	31% (17-49%)	69% (51-83%)	0.82 (0.32-1.6)	2.3% (0.37-5.5%)	2.6% (0.4-6.2%)	78 (12-180)	1100 (180-2600)
Madagascar	1995	Female	80 to 84	29% (15-47%)	71% (53-85%)	0.73 (0.29-1.4)	1.6% (0.14-4%)	1.8% (0.14-4.5%)	45 (3.3-110)	470 (38-1100)
Madagascar	1995	Female	85 to 89	29% (16-47%)	71% (53-84%)	0.71 (0.3-1.4)	1.2% (-0.053-3.1%)	1.3% (-0.11-3.6%)	20 (-1.6-56)	150 (-6.8-410)
Madagascar	1995	Female	90 to 94	32% (18-49%)	68% (51-82%)	0.74 (0.3-1.5)	0.86% (-0.29-3%)	0.95% (-0.4-3.5%)	6 (-2.5-22)	32 (-11-110)
Madagascar	1995	Female	95 plus	37% (18-58%)	63% (42-82%)	0.81 (0.29-1.8)	0.99% (-0.28-3.6%)	1% (-0.45-4%)	1.7 (-0.74-6.5)	6.2 (-1.8-22)
Madagascar	1995	Male	All Ages	30% (24-37%)	70% (63-76%)	0.83 (0.64-1)	1.7% (1.2-2.3%)	2.6% (1.6-3.7%)	2100 (1300-3100)	97000 (67000-130000)
Madagascar	1995	Male	15 to 19	26% (15-42%)	74% (58-85%)	0.83 (0.42-1.4)	2.7% (1.6-4.2%)	2.9% (1.4-5.1%)	42 (20-76)	4400 (2600-6900)
Madagascar	1995	Male	20 to 24	38% (23-55%)	62% (45-77%)	1.2 (0.65-2)	5.3% (3.2-7.8%)	5.3% (2.7-8.6%)	100 (50-170)	9600 (5700-14000)
Madagascar	1995	Male	25 to 29	43% (26-61%)	57% (39-74%)	1.6 (0.87-2.4)	6.8% (4.4-9.9%)	7.3% (4-11%)	140 (76-220)	11000 (6700-16000)
Madagascar	1995	Male	30 to 34	47% (27-66%)	53% (34-73%)	1.8 (1-2.8)	7.8% (4.9-11%)	8.9% (5-14%)	170 (96-270)	12000 (7100-17000)
Madagascar	1995	Male	35 to 39	50% (32-67%)	50% (33-68%)	2 (1.1-3.2)	8.3% (5.1-12%)	9.8% (5.7-15%)	200 (110-310)	12000 (6900-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	1995	Male	40 to 44	51% (34-69%)	49% (31-66%)	2.1 (1.1-3.3)	8.3% (4.7-12%)	10% (5.5-15%)	220 (120-340)	11000 (6100-17000)
Madagascar	1995	Male	45 to 49	52% (33-70%)	48% (30-67%)	2.1 (1-3.6)	8% (4.4-13%)	9.7% (5-16%)	200 (100-320)	8600 (4600-14000)
Madagascar	1995	Male	50 to 54	52% (33-70%)	48% (30-67%)	2 (1-3.4)	7.8% (4-13%)	9.3% (4.7-15%)	210 (100-340)	7900 (4000-13000)
Madagascar	1995	Male	55 to 59	51% (34-71%)	49% (29-66%)	1.9 (1-3.3)	7% (3.5-12%)	8.3% (4-14%)	230 (110-390)	7500 (3800-13000)
Madagascar	1995	Male	60 to 64	49% (31-69%)	51% (31-69%)	1.7 (0.82-3)	5.7% (2.7-10%)	6.7% (3.1-12%)	230 (100-410)	6400 (3000-11000)
Madagascar	1995	Male	65 to 69	47% (30-65%)	53% (35-70%)	1.4 (0.67-2.4)	3.8% (1.5-6.9%)	4.3% (1.7-8%)	170 (62-310)	3800 (1500-7100)
Madagascar	1995	Male	70 to 74	44% (28-62%)	56% (38-72%)	1.1 (0.5-1.9)	2.5% (0.78-5.2%)	2.8% (0.86-6%)	110 (35-250)	2100 (690-4400)
Madagascar	1995	Male	75 to 79	42% (25-60%)	58% (40-75%)	0.86 (0.39-1.6)	1.7% (0.38-3.9%)	1.9% (0.37-4.5%)	65 (12-150)	930 (200-2200)
Madagascar	1995	Male	80 to 84	39% (22-58%)	61% (42-78%)	0.73 (0.33-1.3)	1.2% (0.16-2.8%)	1.3% (0.11-3.2%)	30 (2.4-71)	320 (44-750)
Madagascar	1995	Male	85 to 89	36% (21-53%)	64% (47-79%)	0.61 (0.28-1.1)	0.82% (0.04-1.8%)	0.85% (-0.049-2%)	12 (-0.69-28)	98 (4.8-220)
Madagascar	1995	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.55 (0.24-1.1)	0.37% (-0.28-1.2%)	0.29% (-0.48-1.2%)	1.6 (-2.6-6.5)	12 (-9.1-38)
Madagascar	1995	Male	95 plus	40% (21-62%)	60% (38-79%)	0.54 (0.2-1.2)	0.37% (-0.44-1.3%)	0.21% (-0.78-1.2%)	0.25 (-0.92-1.5)	1.7 (-2.1-5.8)
Madagascar	2000	Female	All Ages	20% (16-26%)	80% (74-84%)	0.43 (0.32-0.55)	0.78% (0.47-1.2%)	1.3% (0.6-2.2%)	1000 (470-1800)	41000 (24000-61000)
Madagascar	2000	Female	15 to 19	20% (12-33%)	80% (67-88%)	0.49 (0.24-0.86)	0.97% (0.51-1.6%)	1.1% (0.36-2.1%)	14 (4.5-27)	1500 (790-2600)
Madagascar	2000	Female	20 to 24	26% (14-41%)	74% (59-86%)	0.7 (0.34-1.2)	1.7% (0.99-2.6%)	1.6% (0.61-2.9%)	27 (10-50)	3100 (1700-4700)
Madagascar	2000	Female	25 to 29	28% (15-44%)	72% (56-85%)	0.84 (0.44-1.4)	2.4% (1.5-3.6%)	2.6% (1.3-4.3%)	45 (22-75)	3900 (2400-6000)
Madagascar	2000	Female	30 to 34	31% (16-48%)	69% (52-84%)	0.92 (0.46-1.5)	2.7% (1.6-4.1%)	3.3% (1.6-5.4%)	58 (28-95)	4100 (2400-6400)
Madagascar	2000	Female	35 to 39	34% (19-53%)	66% (47-81%)	0.91 (0.47-1.5)	2.7% (1.6-4.3%)	3.3% (1.7-5.6%)	63 (33-110)	3900 (2200-6200)
Madagascar	2000	Female	40 to 44	36% (21-55%)	64% (45-79%)	0.92 (0.45-1.6)	2.8% (1.5-4.6%)	3.5% (1.8-6.1%)	73 (36-130)	3800 (2000-6400)
Madagascar	2000	Female	45 to 49	36% (19-54%)	64% (46-81%)	0.95 (0.44-1.8)	2.9% (1.4-5.4%)	3.8% (1.6-7.1%)	88 (37-160)	3900 (1800-7100)
Madagascar	2000	Female	50 to 54	34% (19-54%)	66% (46-81%)	0.93 (0.43-1.7)	3.3% (1.4-6.1%)	4.1% (1.7-7.7%)	100 (40-190)	3800 (1600-7100)
Madagascar	2000	Female	55 to 59	33% (19-52%)	67% (48-81%)	0.92 (0.42-1.8)	3.1% (1.1-6.3%)	3.8% (1.3-7.7%)	110 (37-230)	3600 (1200-7300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	2000	Female	60 to 64	33% (18-53%)	67% (47-82%)	0.91 (0.41-1.7)	2.9% (0.82-6.1%)	3.5% (0.99-7.2%)	130 (36-280)	3500 (1000-7500)
Madagascar	2000	Female	65 to 69	35% (20-52%)	65% (48-80%)	0.86 (0.38-1.5)	2.6% (0.74-5.4%)	3.1% (0.82-6.3%)	140 (37-290)	3100 (860-6500)
Madagascar	2000	Female	70 to 74	33% (19-52%)	67% (48-81%)	0.74 (0.32-1.4)	1.8% (0.15-4.3%)	2% (0.13-4.9%)	93 (5.7-230)	1700 (140-4100)
Madagascar	2000	Female	75 to 79	31% (17-49%)	69% (51-83%)	0.62 (0.24-1.2)	1.3% (-0.15-3.5%)	1.5% (-0.19-4%)	57 (-7.1-150)	810 (-88-2100)
Madagascar	2000	Female	80 to 84	29% (15-47%)	71% (53-85%)	0.55 (0.22-1.1)	0.81% (-0.3-2.5%)	0.92% (-0.36-2.9%)	26 (-9.9-81)	270 (-99-840)
Madagascar	2000	Female	85 to 89	29% (16-46%)	71% (54-84%)	0.53 (0.22-1.1)	0.51% (-0.47-1.8%)	0.56% (-0.55-2.1%)	7.9 (-7.7-29)	60 (-55-220)
Madagascar	2000	Female	90 to 94	32% (17-50%)	68% (50-83%)	0.56 (0.23-1.1)	0.27% (-0.62-1.7%)	0.26% (-0.78-2%)	1.5 (-4.7-11)	9.3 (-21-59)
Madagascar	2000	Female	95 plus	37% (18-59%)	63% (41-82%)	0.62 (0.21-1.4)	0.41% (-0.69-2%)	0.37% (-0.94-2.2%)	0.65 (-1.7-3.9)	2.8 (-4.7-14)
Madagascar	2000	Male	All Ages	30% (24-37%)	70% (63-76%)	0.63 (0.48-0.79)	1.4% (0.97-1.9%)	2% (1.2-2.9%)	1700 (980-2500)	82000 (56000-110000)
Madagascar	2000	Male	15 to 19	26% (16-40%)	74% (60-84%)	0.63 (0.32-1.1)	2.2% (1.4-3.4%)	2.2% (1.1-3.8%)	32 (16-57)	3700 (2200-5700)
Madagascar	2000	Male	20 to 24	37% (23-54%)	63% (46-77%)	0.94 (0.49-1.5)	4.3% (2.7-6.2%)	3.9% (1.8-6.4%)	86 (39-150)	8800 (5400-13000)
Madagascar	2000	Male	25 to 29	43% (26-61%)	57% (39-74%)	1.2 (0.68-1.9)	5.3% (3.4-7.6%)	5.3% (2.8-8.4%)	120 (63-190)	10000 (6200-15000)
Madagascar	2000	Male	30 to 34	47% (28-66%)	53% (34-72%)	1.4 (0.8-2.1)	5.9% (3.7-8.5%)	6.4% (3.5-10%)	140 (77-230)	10000 (6300-15000)
Madagascar	2000	Male	35 to 39	50% (33-68%)	50% (32-67%)	1.5 (0.85-2.5)	6.1% (3.7-9%)	7% (3.9-11%)	160 (91-260)	9800 (5900-14000)
Madagascar	2000	Male	40 to 44	51% (35-69%)	49% (31-65%)	1.6 (0.85-2.5)	6.1% (3.4-9.3%)	7.1% (3.7-11%)	180 (92-280)	9200 (5100-14000)
Madagascar	2000	Male	45 to 49	51% (32-70%)	49% (30-68%)	1.6 (0.81-2.7)	5.9% (3.1-9.7%)	7% (3.4-12%)	190 (90-310)	8400 (4400-14000)
Madagascar	2000	Male	50 to 54	52% (33-70%)	48% (30-67%)	1.5 (0.78-2.6)	5.7% (2.8-9.3%)	6.7% (3.2-11%)	170 (77-290)	6500 (3300-11000)
Madagascar	2000	Male	55 to 59	51% (33-70%)	49% (30-67%)	1.5 (0.77-2.6)	5.1% (2.5-8.6%)	5.9% (2.8-10%)	160 (75-280)	5500 (2700-9300)
Madagascar	2000	Male	60 to 64	49% (30-68%)	51% (32-70%)	1.3 (0.6-2.3)	4.1% (1.8-7.7%)	4.7% (2.1-9%)	160 (70-310)	4600 (2100-8700)
Madagascar	2000	Male	65 to 69	47% (29-64%)	53% (36-71%)	1 (0.51-1.8)	2.6% (0.93-4.9%)	2.9% (0.97-5.6%)	120 (39-230)	2800 (980-5200)
Madagascar	2000	Male	70 to 74	44% (27-61%)	56% (39-73%)	0.8 (0.38-1.4)	1.6% (0.33-3.5%)	1.8% (0.34-4%)	78 (15-170)	1500 (290-3100)
Madagascar	2000	Male	75 to 79	41% (25-60%)	59% (40-75%)	0.65 (0.29-1.2)	1.1% (0.098-2.6%)	1.2% (0.062-2.9%)	47 (2.5-110)	690 (62-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	2000	Male	80 to 84	39% (22-57%)	61% (43-78%)	0.55 (0.25-0.99)	0.74% (-0.14-1.8%)	0.79% (-0.21-2%)	21 (-5.6-54)	230 (-44-570)
Madagascar	2000	Male	85 to 89	36% (21-54%)	64% (46-79%)	0.46 (0.21-0.86)	0.52% (-0.14-1.3%)	0.52% (-0.26-1.4%)	7.6 (-3.8-21)	65 (-17-160)
Madagascar	2000	Male	90 to 94	37% (22-55%)	63% (45-78%)	0.42 (0.18-0.84)	0.18% (-0.42-0.78%)	0.083% (-0.64-0.79%)	0.5 (-3.9-4.8)	6.7 (-15-28)
Madagascar	2000	Male	95 plus	40% (20-62%)	60% (38-80%)	0.42 (0.15-0.9)	0.23% (-0.56-0.89%)	0.054% (-0.88-0.85%)	0.086 (-1.4-1.4)	1.4 (-3.4-5.7)
Madagascar	2005	Female	All Ages	20% (16-26%)	80% (74-84%)	0.36 (0.26-0.45)	0.66% (0.36-1%)	1% (0.38-1.9%)	830 (320-1500)	35000 (20000-55000)
Madagascar	2005	Female	15 to 19	20% (11-34%)	80% (66-89%)	0.4 (0.19-0.72)	0.79% (0.43-1.3%)	0.76% (0.23-1.6%)	11 (3.3-24)	1500 (780-2500)
Madagascar	2005	Female	20 to 24	26% (14-40%)	74% (60-86%)	0.57 (0.28-0.98)	1.4% (0.84-2.2%)	1.1% (0.37-2.3%)	19 (6.3-37)	2500 (1400-4000)
Madagascar	2005	Female	25 to 29	28% (15-44%)	72% (56-85%)	0.69 (0.34-1.2)	2% (1.2-3%)	2% (0.95-3.5%)	37 (18-64)	3600 (2200-5500)
Madagascar	2005	Female	30 to 34	30% (16-47%)	70% (53-84%)	0.75 (0.38-1.2)	2.2% (1.3-3.2%)	2.4% (1.2-4%)	48 (23-81)	3800 (2200-5700)
Madagascar	2005	Female	35 to 39	34% (19-52%)	66% (48-81%)	0.74 (0.4-1.3)	2.1% (1.2-3.3%)	2.5% (1.2-4.2%)	53 (26-90)	3400 (1900-5300)
Madagascar	2005	Female	40 to 44	36% (21-55%)	64% (45-79%)	0.76 (0.36-1.4)	2.2% (1.2-3.7%)	2.7% (1.3-4.9%)	63 (30-110)	3400 (1700-5600)
Madagascar	2005	Female	45 to 49	36% (19-53%)	64% (47-81%)	0.78 (0.35-1.5)	2.3% (1.1-4.3%)	2.9% (1.2-5.7%)	76 (32-160)	3400 (1600-6600)
Madagascar	2005	Female	50 to 54	34% (18-54%)	66% (46-82%)	0.76 (0.35-1.4)	2.4% (0.98-4.8%)	3.1% (1.2-6.1%)	97 (36-190)	3700 (1400-7200)
Madagascar	2005	Female	55 to 59	33% (19-52%)	67% (48-81%)	0.76 (0.34-1.5)	2.2% (0.64-4.7%)	2.7% (0.78-5.9%)	89 (25-200)	2900 (800-6300)
Madagascar	2005	Female	60 to 64	34% (18-53%)	66% (47-82%)	0.76 (0.33-1.4)	2.1% (0.48-4.4%)	2.5% (0.57-5.3%)	96 (21-220)	2600 (590-5800)
Madagascar	2005	Female	65 to 69	35% (20-52%)	65% (48-80%)	0.71 (0.32-1.3)	1.9% (0.36-4.1%)	2.2% (0.4-4.7%)	100 (18-230)	2400 (440-5200)
Madagascar	2005	Female	70 to 74	33% (20-51%)	67% (49-80%)	0.61 (0.27-1.2)	1.2% (-0.15-3.2%)	1.3% (-0.25-3.7%)	69 (-13-200)	1300 (-150-3600)
Madagascar	2005	Female	75 to 79	31% (17-49%)	69% (51-83%)	0.51 (0.21-1.1)	0.85% (-0.39-2.5%)	0.95% (-0.5-2.9%)	44 (-22-130)	620 (-300-1900)
Madagascar	2005	Female	80 to 84	29% (15-48%)	71% (52-85%)	0.46 (0.18-0.89)	0.44% (-0.6-1.8%)	0.5% (-0.69-2%)	17 (-23-71)	180 (-240-720)
Madagascar	2005	Female	85 to 89	30% (16-47%)	70% (53-84%)	0.45 (0.18-0.91)	0.2% (-0.68-1.2%)	0.21% (-0.81-1.4%)	3.3 (-13-23)	27 (-91-170)
Madagascar	2005	Female	90 to 94	32% (18-50%)	68% (50-82%)	0.47 (0.18-0.97)	-0.018% (-0.86-1.1%)	-0.072% (-1.1-1.2%)	-0.38 (-5.6-6.3)	-0.55 (-27-33)
Madagascar	2005	Female	95 plus	36% (18-58%)	64% (42-82%)	0.51 (0.17-1.1)	0.14% (-0.9-1.3%)	0.066% (-1.2-1.5%)	0.11 (-1.9-2.3)	0.87 (-5.7-8.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	2005	Male	All Ages	29% (23-37%)	71% (63-77%)	0.52 (0.39-0.65)	1.2% (0.81-1.7%)	1.6% (0.85-2.4%)	1400 (760-2300)	74000 (47000-100000)
Madagascar	2005	Male	15 to 19	26% (16-41%)	74% (59-84%)	0.52 (0.26-0.91)	1.9% (1.2-2.8%)	1.6% (0.77-3%)	29 (13-54)	3700 (2200-5900)
Madagascar	2005	Male	20 to 24	37% (22-55%)	63% (45-78%)	0.76 (0.4-1.2)	3.5% (2.3-5.1%)	2.8% (1.3-4.9%)	63 (27-110)	7400 (4400-11000)
Madagascar	2005	Male	25 to 29	43% (25-60%)	57% (40-75%)	0.97 (0.55-1.5)	4.2% (2.7-6.1%)	3.8% (1.9-6.3%)	97 (49-160)	9200 (5800-13000)
Madagascar	2005	Male	30 to 34	47% (28-65%)	53% (35-72%)	1.1 (0.63-1.7)	4.5% (2.8-6.6%)	4.6% (2.4-7.2%)	120 (63-200)	9300 (5600-14000)
Madagascar	2005	Male	35 to 39	49% (32-67%)	51% (33-68%)	1.2 (0.69-2)	4.7% (2.8-7%)	5.1% (2.6-8.2%)	140 (69-230)	8800 (5000-13000)
Madagascar	2005	Male	40 to 44	51% (33-69%)	49% (31-67%)	1.3 (0.67-2.1)	4.7% (2.6-7.1%)	5.4% (2.7-8.5%)	150 (72-260)	8200 (4400-13000)
Madagascar	2005	Male	45 to 49	51% (33-71%)	49% (29-67%)	1.3 (0.67-2.2)	4.5% (2.5-7.4%)	5.3% (2.6-8.8%)	160 (78-280)	7600 (4000-12000)
Madagascar	2005	Male	50 to 54	52% (33-70%)	48% (30-67%)	1.3 (0.64-2.1)	4.4% (2.1-7.2%)	5.1% (2.2-8.6%)	170 (75-290)	6800 (3300-11000)
Madagascar	2005	Male	55 to 59	51% (33-71%)	49% (29-67%)	1.2 (0.62-2)	3.9% (1.9-6.9%)	4.5% (2-8.1%)	140 (60-260)	4800 (2200-8700)
Madagascar	2005	Male	60 to 64	49% (30-69%)	51% (31-70%)	1 (0.49-1.9)	3.2% (1.4-6%)	3.6% (1.5-6.9%)	130 (50-260)	3700 (1600-7100)
Madagascar	2005	Male	65 to 69	46% (29-64%)	54% (36-71%)	0.83 (0.39-1.5)	1.9% (0.61-3.7%)	2.2% (0.58-4.3%)	92 (24-180)	2200 (670-4100)
Madagascar	2005	Male	70 to 74	43% (27-61%)	57% (39-73%)	0.65 (0.31-1.2)	1.2% (0.12-2.5%)	1.3% (0.093-2.9%)	62 (4.2-140)	1200 (120-2500)
Madagascar	2005	Male	75 to 79	41% (25-59%)	59% (41-75%)	0.52 (0.23-1)	0.82% (-0.06-2%)	0.91% (-0.11-2.2%)	38 (-4.5-93)	560 (-39-1300)
Madagascar	2005	Male	80 to 84	38% (22-57%)	62% (43-78%)	0.44 (0.2-0.81)	0.53% (-0.17-1.4%)	0.55% (-0.25-1.5%)	17 (-7.7-46)	190 (-60-500)
Madagascar	2005	Male	85 to 89	35% (21-53%)	65% (47-79%)	0.37 (0.17-0.69)	0.4% (-0.19-1%)	0.38% (-0.32-1.1%)	6.8 (-5.6-19)	61 (-29-150)
Madagascar	2005	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.34 (0.14-0.67)	0.12% (-0.42-0.59%)	0.013% (-0.61-0.56%)	0.089 (-4.1-3.7)	4.6 (-16-23)
Madagascar	2005	Male	95 plus	40% (20-61%)	60% (39-80%)	0.34 (0.12-0.74)	0.18% (-0.5-0.73%)	0.02% (-0.8-0.7%)	0.035 (-1.4-1.3)	1.3 (-3.5-5.1)
Madagascar	2010	Female	All Ages	21% (16-27%)	79% (73-84%)	0.38 (0.28-0.51)	0.73% (0.4-1.2%)	1.1% (0.41-2.1%)	1000 (360-1900)	42000 (23000-67000)
Madagascar	2010	Female	15 to 19	20% (11-34%)	80% (66-89%)	0.42 (0.2-0.77)	0.85% (0.45-1.5%)	0.84% (0.26-1.8%)	14 (4.4-32)	1900 (950-3200)
Madagascar	2010	Female	20 to 24	26% (14-41%)	74% (59-86%)	0.6 (0.29-1.1)	1.5% (0.88-2.4%)	1.3% (0.44-2.5%)	25 (8.8-50)	3300 (1900-5200)
Madagascar	2010	Female	25 to 29	28% (15-44%)	72% (56-85%)	0.73 (0.37-1.3)	2.1% (1.3-3.2%)	2.1% (0.95-3.8%)	40 (18-74)	3900 (2300-6100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	2010	Female	30 to 34	31% (17-48%)	69% (52-83%)	0.79 (0.39-1.4)	2.2% (1.3-3.5%)	2.5% (1.2-4.5%)	56 (25-100)	4300 (2500-6700)
Madagascar	2010	Female	35 to 39	34% (19-52%)	66% (48-81%)	0.78 (0.4-1.3)	2.2% (1.3-3.5%)	2.6% (1.3-4.6%)	64 (32-110)	4100 (2400-6700)
Madagascar	2010	Female	40 to 44	36% (21-54%)	64% (46-79%)	0.79 (0.36-1.4)	2.2% (1.2-3.8%)	2.8% (1.3-4.9%)	73 (34-130)	3900 (2000-6600)
Madagascar	2010	Female	45 to 49	36% (19-54%)	64% (46-81%)	0.81 (0.38-1.5)	2.4% (1.1-4.3%)	3% (1.3-5.8%)	90 (36-180)	4000 (1800-7700)
Madagascar	2010	Female	50 to 54	35% (19-55%)	65% (45-81%)	0.8 (0.36-1.5)	2.6% (1-4.9%)	3.2% (1.2-6.3%)	120 (41-240)	4400 (1700-9000)
Madagascar	2010	Female	55 to 59	33% (19-53%)	67% (47-81%)	0.8 (0.35-1.5)	2.4% (0.71-5%)	2.9% (0.87-6.4%)	120 (34-280)	4000 (1100-8800)
Madagascar	2010	Female	60 to 64	34% (19-52%)	66% (48-81%)	0.79 (0.34-1.6)	2.2% (0.54-5%)	2.7% (0.65-6.1%)	110 (27-270)	3100 (730-7400)
Madagascar	2010	Female	65 to 69	35% (19-53%)	65% (47-81%)	0.74 (0.32-1.4)	2% (0.48-4.2%)	2.3% (0.53-4.9%)	120 (26-270)	2700 (630-6000)
Madagascar	2010	Female	70 to 74	33% (19-51%)	67% (49-81%)	0.63 (0.26-1.2)	1.3% (-0.18-3.5%)	1.4% (-0.25-4.1%)	81 (-14-230)	1500 (-200-4200)
Madagascar	2010	Female	75 to 79	30% (17-48%)	70% (52-83%)	0.53 (0.2-1.1)	0.91% (-0.31-2.9%)	1% (-0.38-3.3%)	56 (-21-180)	800 (-290-2500)
Madagascar	2010	Female	80 to 84	29% (16-47%)	71% (53-84%)	0.48 (0.19-0.93)	0.49% (-0.51-1.9%)	0.56% (-0.57-2.2%)	23 (-23-94)	250 (-240-980)
Madagascar	2010	Female	85 to 89	30% (16-46%)	70% (54-84%)	0.47 (0.2-0.97)	0.24% (-0.6-1.4%)	0.26% (-0.74-1.6%)	5.4 (-15-35)	43 (-110-250)
Madagascar	2010	Female	90 to 94	32% (18-49%)	68% (51-82%)	0.49 (0.19-0.99)	0.027% (-0.81-1.2%)	-0.023% (-1-1.4%)	-0.15 (-6.9-8.9)	1.1 (-33-47)
Madagascar	2010	Female	95 plus	37% (18-58%)	63% (42-82%)	0.53 (0.18-1.2)	0.18% (-0.87-1.5%)	0.11% (-1.2-1.7%)	0.16 (-1.7-2.5)	1.1 (-5.1-8.9)
Madagascar	2010	Male	All Ages	30% (24-37%)	70% (63-76%)	0.55 (0.41-0.72)	1.3% (0.88-1.9%)	1.7% (0.9-2.6%)	1700 (860-2600)	87000 (55000-120000)
Madagascar	2010	Male	15 to 19	26% (16-41%)	74% (59-84%)	0.53 (0.27-0.95)	1.9% (1.2-2.9%)	1.7% (0.8-3.2%)	35 (15-68)	4500 (2700-7100)
Madagascar	2010	Male	20 to 24	37% (21-55%)	63% (45-79%)	0.79 (0.4-1.3)	3.7% (2.3-5.5%)	3% (1.4-5.2%)	78 (31-150)	9200 (5400-14000)
Madagascar	2010	Male	25 to 29	43% (25-61%)	57% (39-75%)	1 (0.55-1.6)	4.4% (2.8-6.3%)	3.9% (2-6.6%)	100 (49-180)	9700 (5900-15000)
Madagascar	2010	Male	30 to 34	47% (28-65%)	53% (35-72%)	1.2 (0.65-1.8)	4.6% (2.8-6.8%)	4.6% (2.4-7.4%)	140 (68-250)	11000 (6500-17000)
Madagascar	2010	Male	35 to 39	49% (31-67%)	51% (33-69%)	1.3 (0.71-2.1)	4.7% (2.9-6.9%)	5.1% (2.8-8%)	160 (83-280)	10000 (6000-16000)
Madagascar	2010	Male	40 to 44	50% (33-69%)	50% (31-67%)	1.3 (0.7-2.2)	4.7% (2.7-7.2%)	5.4% (2.8-8.5%)	180 (86-300)	9500 (5100-15000)
Madagascar	2010	Male	45 to 49	51% (32-70%)	49% (30-68%)	1.3 (0.7-2.3)	4.6% (2.5-7.7%)	5.4% (2.7-9.3%)	190 (91-330)	8800 (4600-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	2010	Male	50 to 54	51% (33-71%)	49% (29-67%)	1.3 (0.65-2.2)	4.5% (2.3-7.4%)	5.2% (2.4-8.9%)	200 (95-350)	8100 (4100-14000)
Madagascar	2010	Male	55 to 59	51% (33-70%)	49% (30-67%)	1.2 (0.63-2.2)	4% (1.9-7%)	4.7% (2.1-8.2%)	200 (83-370)	6700 (3000-12000)
Madagascar	2010	Male	60 to 64	49% (30-69%)	51% (31-70%)	1.1 (0.52-2)	3.2% (1.4-5.9%)	3.7% (1.5-7%)	150 (58-290)	4300 (1800-7900)
Madagascar	2010	Male	65 to 69	46% (29-64%)	54% (36-71%)	0.86 (0.42-1.5)	2% (0.61-3.7%)	2.2% (0.6-4.3%)	96 (27-190)	2300 (710-4400)
Madagascar	2010	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.67 (0.31-1.2)	1.2% (0.16-2.6%)	1.4% (0.12-3%)	66 (6.3-150)	1200 (160-2700)
Madagascar	2010	Male	75 to 79	41% (25-60%)	59% (40-75%)	0.55 (0.24-1.1)	0.82% (-0.052-2%)	0.91% (-0.11-2.3%)	42 (-5.1-100)	620 (-38-1500)
Madagascar	2010	Male	80 to 84	38% (21-58%)	62% (42-79%)	0.46 (0.21-0.84)	0.54% (-0.15-1.4%)	0.56% (-0.21-1.5%)	20 (-7.6-55)	230 (-61-580)
Madagascar	2010	Male	85 to 89	36% (21-53%)	64% (47-79%)	0.38 (0.17-0.73)	0.39% (-0.16-1%)	0.37% (-0.3-1.1%)	7.6 (-6.2-23)	70 (-28-180)
Madagascar	2010	Male	90 to 94	36% (21-54%)	64% (46-79%)	0.35 (0.14-0.69)	0.11% (-0.41-0.59%)	0.0012% (-0.61-0.54%)	0.011 (-5-4.4)	5.4 (-20-28)
Madagascar	2010	Male	95 plus	40% (20-62%)	60% (38-80%)	0.35 (0.13-0.74)	0.15% (-0.5-0.67%)	-0.017% (-0.8-0.61%)	-0.034 (-1.6-1.2)	1.2 (-3.9-5.4)
Madagascar	2016	Female	All Ages	21% (17-27%)	79% (73-83%)	0.48 (0.28-0.76)	1% (0.47-1.8%)	1.6% (0.5-3.2%)	1500 (430-3300)	60000 (28000-110000)
Madagascar	2016	Female	15 to 19	20% (11-33%)	80% (67-89%)	0.5 (0.2-1.1)	1% (0.49-1.9%)	1.1% (0.3-2.5%)	19 (5.5-46)	2400 (1100-4500)
Madagascar	2016	Female	20 to 24	25% (14-41%)	75% (59-86%)	0.71 (0.31-1.4)	1.8% (0.99-3%)	1.6% (0.51-3.4%)	33 (10-72)	4300 (2300-7100)
Madagascar	2016	Female	25 to 29	28% (15-44%)	72% (56-85%)	0.86 (0.39-1.7)	2.3% (1.4-3.9%)	2.4% (1.1-4.9%)	52 (21-100)	5100 (2800-8400)
Madagascar	2016	Female	30 to 34	31% (16-48%)	69% (52-84%)	0.95 (0.43-1.8)	2.5% (1.4-4.3%)	3% (1.3-5.6%)	63 (26-130)	5000 (2700-8700)
Madagascar	2016	Female	35 to 39	34% (19-52%)	66% (48-81%)	0.94 (0.41-1.8)	2.5% (1.4-4.4%)	3.2% (1.5-5.9%)	82 (35-160)	5300 (2700-9400)
Madagascar	2016	Female	40 to 44	36% (21-54%)	64% (46-79%)	0.96 (0.4-1.8)	2.6% (1.2-4.8%)	3.4% (1.5-6.4%)	100 (40-210)	5500 (2500-10000)
Madagascar	2016	Female	45 to 49	36% (19-55%)	64% (45-81%)	0.98 (0.41-2)	2.9% (1.3-5.8%)	3.8% (1.5-7.9%)	130 (47-270)	5800 (2400-12000)
Madagascar	2016	Female	50 to 54	35% (19-54%)	65% (46-81%)	0.96 (0.37-1.9)	3.1% (1.2-6.4%)	4% (1.5-8.1%)	170 (55-360)	6300 (2300-14000)
Madagascar	2016	Female	55 to 59	33% (19-52%)	67% (48-81%)	0.94 (0.38-2)	3% (0.78-6.6%)	3.7% (0.91-8.2%)	180 (43-430)	5900 (1500-14000)
Madagascar	2016	Female	60 to 64	34% (19-53%)	66% (47-81%)	0.94 (0.36-2.1)	2.8% (0.62-6.6%)	3.4% (0.73-8%)	200 (42-490)	5400 (1200-13000)
Madagascar	2016	Female	65 to 69	35% (20-52%)	65% (48-80%)	0.89 (0.34-1.8)	2.6% (0.6-6.3%)	3.1% (0.68-7.4%)	180 (36-450)	4200 (880-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	2016	Female	70 to 74	34% (19-51%)	66% (49-81%)	0.77 (0.29-1.6)	1.9% (0.03-5.2%)	2.1% (-0.03-6%)	130 (-1.9-380)	2400 (41-6900)
Madagascar	2016	Female	75 to 79	31% (17-50%)	69% (50-83%)	0.64 (0.22-1.4)	1.4% (-0.12-4.3%)	1.6% (-0.17-4.9%)	99 (-10-310)	1400 (-120-4400)
Madagascar	2016	Female	80 to 84	29% (16-47%)	71% (53-84%)	0.58 (0.21-1.3)	0.91% (-0.4-3.4%)	1% (-0.46-3.9%)	53 (-24-200)	560 (-240-2000)
Madagascar	2016	Female	85 to 89	29% (16-46%)	71% (54-84%)	0.56 (0.2-1.3)	0.6% (-0.47-2.6%)	0.66% (-0.56-2.9%)	19 (-16-83)	150 (-110-620)
Madagascar	2016	Female	90 to 94	32% (18-50%)	68% (50-82%)	0.58 (0.21-1.3)	0.35% (-0.72-2.4%)	0.35% (-0.89-2.7%)	3.4 (-8.6-25)	20 (-40-130)
Madagascar	2016	Female	95 plus	36% (18-58%)	64% (42-82%)	0.64 (0.21-1.6)	0.48% (-0.74-2.8%)	0.45% (-0.96-3.2%)	0.98 (-2.1-7.1)	4.1 (-6.2-24)
Madagascar	2016	Male	All Ages	30% (24-38%)	70% (62-76%)	0.68 (0.41-1.1)	1.8% (1.1-2.9%)	2.3% (1.1-4%)	2400 (1100-4400)	120000 (69000-200000)
Madagascar	2016	Male	15 to 19	26% (15-42%)	74% (58-85%)	0.64 (0.28-1.3)	2.2% (1.3-3.6%)	2.1% (0.82-4.1%)	48 (19-100)	5900 (3300-10000)
Madagascar	2016	Male	20 to 24	37% (22-54%)	63% (46-78%)	0.94 (0.43-1.7)	4.3% (2.5-6.5%)	3.6% (1.5-6.8%)	110 (41-220)	13000 (7100-20000)
Madagascar	2016	Male	25 to 29	43% (25-60%)	57% (40-75%)	1.2 (0.58-2.2)	5.1% (3.1-8%)	4.8% (2.2-8.6%)	150 (64-290)	14000 (8000-23000)
Madagascar	2016	Male	30 to 34	47% (27-66%)	53% (34-73%)	1.4 (0.66-2.6)	5.4% (3.1-8.6%)	5.6% (2.6-9.9%)	170 (74-320)	13000 (7300-22000)
Madagascar	2016	Male	35 to 39	49% (32-68%)	51% (32-68%)	1.5 (0.71-2.8)	5.4% (3.1-8.9%)	6% (2.9-10%)	220 (96-410)	14000 (7400-23000)
Madagascar	2016	Male	40 to 44	50% (33-70%)	50% (30-67%)	1.6 (0.76-2.9)	5.5% (2.9-9.4%)	6.4% (3-11%)	260 (110-480)	14000 (6800-25000)
Madagascar	2016	Male	45 to 49	51% (32-69%)	49% (31-68%)	1.6 (0.74-3)	5.4% (2.7-9.9%)	6.4% (2.9-12%)	270 (110-540)	12000 (5800-23000)
Madagascar	2016	Male	50 to 54	51% (33-70%)	49% (30-67%)	1.6 (0.7-3)	5.3% (2.5-9.8%)	6.3% (2.7-12%)	290 (110-580)	11000 (5000-22000)
Madagascar	2016	Male	55 to 59	50% (32-69%)	50% (31-68%)	1.5 (0.63-2.8)	4.8% (2.1-9.4%)	5.6% (2.3-11%)	280 (110-580)	9600 (4100-19000)
Madagascar	2016	Male	60 to 64	49% (30-69%)	51% (31-70%)	1.3 (0.57-2.6)	3.9% (1.5-8.5%)	4.6% (1.7-9.8%)	260 (95-580)	7500 (2900-16000)
Madagascar	2016	Male	65 to 69	46% (28-64%)	54% (36-72%)	1 (0.45-2)	2.4% (0.72-5.5%)	2.8% (0.73-6.3%)	150 (39-340)	3500 (1000-8000)
Madagascar	2016	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.8 (0.33-1.7)	1.5% (0.21-3.9%)	1.7% (0.19-4.5%)	87 (8.8-220)	1600 (230-4200)
Madagascar	2016	Male	75 to 79	41% (25-60%)	59% (40-75%)	0.65 (0.25-1.4)	1% (0.013-2.8%)	1.1% (-0.038-3.1%)	56 (-2-160)	820 (10-2300)
Madagascar	2016	Male	80 to 84	38% (22-57%)	62% (43-78%)	0.55 (0.23-1.1)	0.72% (-0.077-2%)	0.77% (-0.17-2.3%)	30 (-6.4-92)	340 (-38-980)
Madagascar	2016	Male	85 to 89	35% (21-52%)	65% (48-79%)	0.46 (0.18-1)	0.49% (-0.11-1.4%)	0.47% (-0.26-1.5%)	12 (-6-37)	100 (-23-290)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Madagascar	2016	Male	90 to 94	36% (21-54%)	64% (46-79%)	0.41 (0.16-0.89)	0.19% (-0.35-0.86%)	0.084% (-0.55-0.85%)	0.83 (-5.7-8.6)	11 (-21-52)
Madagascar	2016	Male	95 plus	39% (19-61%)	61% (39-81%)	0.41 (0.13-0.99)	0.21% (-0.46-0.92%)	0.029% (-0.79-0.85%)	0.074 (-2.1-2.2)	2.1 (-4.7-9.5)
Malawi	1990	Female	All Ages	8.6% (7-11%)	91% (89-93%)	0.16 (0.12-0.23)	0.25% (0.17-0.36%)	0.41% (0.24-0.66%)	320 (180-510)	14000 (9700-21000)
Malawi	1990	Female	15 to 19	8.7% (5.4-13%)	91% (87-95%)	0.16 (0.08-0.29)	0.5% (0.28-0.84%)	0.37% (0.11-0.82%)	4.5 (1.3-10)	660 (360-1100)
Malawi	1990	Female	20 to 24	11% (6.8-17%)	89% (83-93%)	0.25 (0.12-0.42)	0.75% (0.43-1.2%)	0.45% (0.16-0.96%)	8.6 (2.9-18)	1300 (720-2000)
Malawi	1990	Female	25 to 29	12% (7.3-19%)	88% (81-93%)	0.33 (0.17-0.55)	0.95% (0.57-1.5%)	0.72% (0.33-1.4%)	15 (7.2-29)	1600 (980-2500)
Malawi	1990	Female	30 to 34	14% (8.4-22%)	86% (78-92%)	0.37 (0.19-0.64)	1.1% (0.65-1.7%)	0.97% (0.45-1.7%)	20 (9.4-34)	1600 (970-2500)
Malawi	1990	Female	35 to 39	17% (9.8-25%)	83% (75-90%)	0.35 (0.18-0.59)	1.1% (0.67-1.7%)	1% (0.55-1.8%)	20 (11-34)	1400 (850-2100)
Malawi	1990	Female	40 to 44	18% (11-26%)	82% (74-89%)	0.34 (0.17-0.59)	1.2% (0.74-1.8%)	1.3% (0.71-2.1%)	25 (14-41)	1400 (880-2200)
Malawi	1990	Female	45 to 49	17% (9.7-26%)	83% (74-90%)	0.36 (0.17-0.68)	1.3% (0.73-2.1%)	1.4% (0.73-2.5%)	26 (14-47)	1200 (720-2100)
Malawi	1990	Female	50 to 54	15% (9.3-23%)	85% (77-91%)	0.36 (0.18-0.67)	1.5% (0.91-2.5%)	1.8% (1-3%)	33 (17-57)	1300 (720-2100)
Malawi	1990	Female	55 to 59	13% (8.1-20%)	87% (80-92%)	0.35 (0.18-0.64)	1.4% (0.66-2.4%)	1.6% (0.75-2.8%)	31 (14-56)	1000 (480-1800)
Malawi	1990	Female	60 to 64	13% (7.9-20%)	87% (80-92%)	0.35 (0.17-0.65)	1.4% (0.68-2.5%)	1.6% (0.77-3%)	38 (17-70)	1000 (480-1900)
Malawi	1990	Female	65 to 69	14% (8.8-21%)	86% (79-91%)	0.34 (0.17-0.57)	1.4% (0.71-2.3%)	1.6% (0.78-2.7%)	43 (20-77)	990 (480-1700)
Malawi	1990	Female	70 to 74	14% (8.6-20%)	86% (80-91%)	0.29 (0.14-0.51)	0.87% (0.29-1.7%)	0.92% (0.27-1.9%)	29 (8.3-62)	570 (200-1200)
Malawi	1990	Female	75 to 79	12% (7.2-20%)	88% (80-93%)	0.24 (0.11-0.46)	0.82% (0.33-1.5%)	0.92% (0.35-1.7%)	22 (8-43)	320 (120-600)
Malawi	1990	Female	80 to 84	12% (6.6-18%)	88% (82-93%)	0.22 (0.097-0.39)	0.47% (0.11-0.94%)	0.53% (0.12-1.1%)	6.6 (1.4-14)	70 (16-140)
Malawi	1990	Female	85 to 89	12% (6.6-19%)	88% (81-93%)	0.21 (0.097-0.38)	0.28% (-0.051-0.66%)	0.3% (-0.088-0.74%)	1.8 (-0.51-4.4)	14 (-2.7-33)
Malawi	1990	Female	90 to 94	13% (7.5-20%)	87% (80-92%)	0.21 (0.099-0.4)	0.15% (-0.17-0.49%)	0.13% (-0.26-0.54%)	0.31 (-0.62-1.3)	2.1 (-2.4-7.1)
Malawi	1990	Female	95 plus	14% (7.5-24%)	86% (76-93%)	0.22 (0.09-0.45)	0.44% (0.058-0.89%)	0.44% (-0.011-0.98%)	0.26 (-0.0065-0.57)	1 (0.13-2.1)
Malawi	1990	Male	All Ages	29% (24-35%)	71% (65-76%)	0.93 (0.68-1.2)	0.89% (0.54-1.3%)	1.4% (0.77-2.2%)	1200 (620-1800)	55000 (33000-82000)
Malawi	1990	Male	15 to 19	26% (17-37%)	74% (63-83%)	0.93 (0.49-1.6)	2.7% (1.6-4.1%)	2.9% (1.4-5%)	27 (11-51)	2800 (1600-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	1990	Male	20 to 24	39% (26-52%)	61% (48-74%)	1.5 (0.82-2.5)	3.8% (2.1-6%)	3.5% (1.5-6.4%)	62 (26-120)	5800 (3200-9600)
Malawi	1990	Male	25 to 29	46% (32-59%)	54% (41-68%)	2 (1.2-3)	3.7% (1.8-6.1%)	3.5% (1.4-6.4%)	91 (41-160)	7100 (3800-11000)
Malawi	1990	Male	30 to 34	50% (35-64%)	50% (36-65%)	2.2 (1.3-3.3)	3.9% (1.9-6.5%)	3.8% (1.6-7.1%)	110 (47-180)	7100 (3800-11000)
Malawi	1990	Male	35 to 39	52% (37-65%)	48% (35-63%)	2.2 (1.3-3.4)	4.1% (1.9-6.9%)	4.3% (1.7-7.8%)	110 (50-180)	6400 (3400-10000)
Malawi	1990	Male	40 to 44	52% (39-65%)	48% (35-61%)	2.3 (1.2-3.4)	4.4% (2.2-7.3%)	4.9% (2.2-8.8%)	120 (54-210)	6200 (3100-10000)
Malawi	1990	Male	45 to 49	54% (38-67%)	46% (33-62%)	2.2 (1.2-3.5)	4.9% (2.6-8.1%)	5.6% (2.6-9.7%)	110 (56-200)	5100 (2700-8500)
Malawi	1990	Male	50 to 54	53% (39-67%)	47% (33-61%)	2.1 (1.2-3.4)	5.7% (3.1-9.1%)	6.8% (3.5-11%)	110 (54-190)	4300 (2200-7300)
Malawi	1990	Male	55 to 59	50% (36-66%)	50% (34-64%)	2.1 (1.2-3.4)	6% (3.4-9.4%)	7.2% (3.9-11%)	110 (52-200)	3800 (1900-6400)
Malawi	1990	Male	60 to 64	47% (31-64%)	53% (36-69%)	1.9 (0.99-3.2)	5.4% (2.9-9%)	6.5% (3.5-11%)	110 (51-200)	3100 (1500-5500)
Malawi	1990	Male	65 to 69	44% (29-60%)	56% (40-71%)	1.4 (0.77-2.4)	3.7% (1.9-6.4%)	4.4% (2.2-7.7%)	77 (35-140)	1800 (840-3200)
Malawi	1990	Male	70 to 74	41% (27-56%)	59% (44-73%)	1.1 (0.58-1.8)	2.6% (1.1-4.6%)	3% (1.3-5.4%)	61 (25-120)	1100 (460-2100)
Malawi	1990	Male	75 to 79	39% (25-55%)	61% (45-75%)	0.95 (0.47-1.7)	2% (0.73-3.9%)	2.3% (0.81-4.5%)	36 (12-71)	510 (180-1000)
Malawi	1990	Male	80 to 84	37% (23-54%)	63% (46-77%)	0.81 (0.4-1.4)	1.5% (0.56-2.9%)	1.7% (0.58-3.3%)	14 (4.5-27)	150 (54-290)
Malawi	1990	Male	85 to 89	34% (21-49%)	66% (51-79%)	0.67 (0.32-1.1)	1.2% (0.49-2.1%)	1.3% (0.48-2.4%)	4.6 (1.7-8.7)	37 (15-67)
Malawi	1990	Male	90 to 94	34% (22-48%)	66% (52-78%)	0.63 (0.31-1.2)	0.78% (0.21-1.7%)	0.79% (0.081-1.8%)	1.1 (0.12-2.7)	6.9 (1.7-15)
Malawi	1990	Male	95 plus	36% (20-55%)	64% (45-80%)	0.63 (0.26-1.2)	0.95% (0.3-2%)	0.92% (0.14-2.1%)	0.3 (0.045-0.69)	1.3 (0.4-2.7)
Malawi	1995	Female	All Ages	11% (8.8-13%)	89% (87-91%)	0.17 (0.12-0.23)	0.25% (0.16-0.38%)	0.39% (0.21-0.63%)	340 (180-570)	15000 (9200-22000)
Malawi	1995	Female	15 to 19	12% (7.7-18%)	88% (82-92%)	0.19 (0.098-0.32)	0.48% (0.25-0.85%)	0.36% (0.065-0.85%)	4.8 (0.97-11)	700 (350-1300)
Malawi	1995	Female	20 to 24	16% (10-23%)	84% (77-90%)	0.29 (0.16-0.5)	0.61% (0.32-1%)	0.36% (0.084-0.82%)	9.5 (2.4-22)	1300 (750-2200)
Malawi	1995	Female	25 to 29	17% (11-25%)	83% (75-89%)	0.38 (0.21-0.63)	0.6% (0.32-1%)	0.43% (0.16-0.93%)	16 (6.6-32)	1600 (970-2500)
Malawi	1995	Female	30 to 34	20% (13-29%)	80% (71-87%)	0.43 (0.23-0.72)	0.66% (0.34-1.1%)	0.55% (0.21-1.1%)	19 (8.2-37)	1500 (850-2500)
Malawi	1995	Female	35 to 39	22% (14-32%)	78% (68-86%)	0.39 (0.21-0.63)	0.68% (0.37-1.1%)	0.61% (0.26-1.1%)	19 (8.5-35)	1300 (710-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	1995	Female	40 to 44	24% (16-34%)	76% (66-84%)	0.39 (0.2-0.66)	0.79% (0.44-1.3%)	0.79% (0.38-1.4%)	22 (11-39)	1200 (660-2000)
Malawi	1995	Female	45 to 49	23% (14-32%)	77% (68-86%)	0.4 (0.2-0.7)	0.91% (0.45-1.6%)	0.97% (0.42-1.8%)	29 (13-56)	1400 (650-2400)
Malawi	1995	Female	50 to 54	17% (11-26%)	83% (74-89%)	0.33 (0.16-0.6)	1.2% (0.65-2%)	1.3% (0.68-2.3%)	39 (18-73)	1500 (760-2700)
Malawi	1995	Female	55 to 59	13% (8.2-21%)	87% (79-92%)	0.3 (0.15-0.51)	1% (0.53-1.9%)	1.2% (0.56-2.1%)	34 (14-65)	1100 (490-2100)
Malawi	1995	Female	60 to 64	14% (7.9-21%)	86% (79-92%)	0.3 (0.14-0.53)	1.1% (0.54-2%)	1.3% (0.61-2.3%)	38 (16-72)	1000 (460-2000)
Malawi	1995	Female	65 to 69	14% (8.9-22%)	86% (78-91%)	0.29 (0.15-0.48)	1.1% (0.56-1.9%)	1.3% (0.58-2.2%)	43 (17-80)	990 (440-1800)
Malawi	1995	Female	70 to 74	14% (8.5-21%)	86% (79-91%)	0.25 (0.12-0.44)	0.68% (0.15-1.4%)	0.71% (0.1-1.5%)	27 (4.2-60)	530 (120-1100)
Malawi	1995	Female	75 to 79	13% (7.4-21%)	87% (79-93%)	0.2 (0.093-0.38)	0.7% (0.24-1.3%)	0.78% (0.25-1.4%)	29 (8.3-56)	410 (130-780)
Malawi	1995	Female	80 to 84	12% (6.9-20%)	88% (80-93%)	0.19 (0.085-0.34)	0.38% (0.0014-0.77%)	0.42% (-0.017-0.88%)	8.9 (-0.34-20)	94 (0.33-210)
Malawi	1995	Female	85 to 89	12% (6.7-19%)	88% (81-93%)	0.18 (0.083-0.32)	0.19% (-0.14-0.51%)	0.21% (-0.19-0.57%)	1.7 (-1.5-4.8)	14 (-9.9-36)
Malawi	1995	Female	90 to 94	13% (7.6-20%)	87% (80-92%)	0.18 (0.084-0.33)	0.072% (-0.25-0.36%)	0.045% (-0.34-0.38%)	0.11 (-0.86-0.97)	1.1 (-3.9-5.3)
Malawi	1995	Female	95 plus	14% (7.4-24%)	86% (76-93%)	0.19 (0.075-0.39)	0.38% (-0.013-0.78%)	0.38% (-0.1-0.84%)	0.29 (-0.077-0.66)	1.2 (-0.04-2.4)
Malawi	1995	Male	All Ages	33% (27-39%)	67% (61-73%)	0.86 (0.63-1.2)	1% (0.63-1.5%)	1.5% (0.88-2.4%)	1400 (770-2200)	65000 (39000-99000)
Malawi	1995	Male	15 to 19	33% (23-44%)	67% (56-77%)	0.98 (0.55-1.6)	2.9% (1.7-4.5%)	3.3% (1.6-5.7%)	36 (16-66)	3500 (2000-5700)
Malawi	1995	Male	20 to 24	47% (34-60%)	53% (40-66%)	1.5 (0.89-2.5)	4% (2.2-6.3%)	3.8% (1.6-6.6%)	84 (39-150)	7400 (4200-12000)
Malawi	1995	Male	25 to 29	53% (40-64%)	47% (36-60%)	1.9 (1.2-2.8)	3% (1.6-5.1%)	2.8% (1.2-5.3%)	110 (54-200)	8500 (4900-14000)
Malawi	1995	Male	30 to 34	56% (42-68%)	44% (32-58%)	2.1 (1.3-3)	2.7% (1.4-4.6%)	2.6% (1.2-4.8%)	120 (58-210)	7900 (4300-13000)
Malawi	1995	Male	35 to 39	57% (44-68%)	43% (32-56%)	2 (1.2-3.1)	2.9% (1.4-5.1%)	3% (1.3-5.5%)	120 (62-210)	7000 (3800-11000)
Malawi	1995	Male	40 to 44	58% (46-69%)	42% (31-54%)	2.1 (1.2-3.2)	3.1% (1.6-5.4%)	3.3% (1.6-5.9%)	130 (63-220)	6300 (3400-10000)
Malawi	1995	Male	45 to 49	59% (45-70%)	41% (30-55%)	2 (1.2-3.1)	3.6% (1.9-6%)	3.9% (1.9-6.7%)	150 (73-250)	6500 (3500-11000)
Malawi	1995	Male	50 to 54	57% (44-69%)	43% (31-56%)	1.9 (1-2.9)	4.5% (2.5-7.2%)	5% (2.7-8.1%)	150 (73-260)	5800 (2900-9600)
Malawi	1995	Male	55 to 59	52% (37-68%)	48% (32-63%)	1.8 (0.98-2.9)	4.9% (2.8-8.1%)	5.7% (3-9.5%)	140 (72-250)	4700 (2500-8100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	1995	Male	60 to 64	48% (32-65%)	52% (35-68%)	1.6 (0.84-2.6)	4.5% (2.5-7.3%)	5.1% (2.8-8.5%)	130 (63-230)	3700 (1800-6400)
Malawi	1995	Male	65 to 69	44% (29-60%)	56% (40-71%)	1.2 (0.66-2)	3% (1.4-5.1%)	3.5% (1.6-5.9%)	90 (41-170)	2100 (970-3800)
Malawi	1995	Male	70 to 74	42% (27-57%)	58% (43-73%)	0.93 (0.5-1.6)	2.1% (0.83-3.9%)	2.4% (0.94-4.5%)	66 (25-130)	1200 (470-2300)
Malawi	1995	Male	75 to 79	40% (26-56%)	60% (44-74%)	0.8 (0.42-1.4)	1.6% (0.56-3.1%)	1.9% (0.64-3.6%)	48 (16-94)	680 (230-1300)
Malawi	1995	Male	80 to 84	37% (23-54%)	63% (46-77%)	0.68 (0.35-1.2)	1.2% (0.35-2.3%)	1.4% (0.34-2.7%)	19 (4.7-38)	210 (59-400)
Malawi	1995	Male	85 to 89	34% (21-49%)	66% (51-79%)	0.56 (0.27-0.96)	0.94% (0.34-1.7%)	1% (0.32-1.9%)	5.4 (1.7-10)	43 (16-79)
Malawi	1995	Male	90 to 94	34% (21-49%)	66% (51-79%)	0.52 (0.25-0.95)	0.6% (0.095-1.3%)	0.59% (-0.0088-1.4%)	0.94 (-0.014-2.2)	5.7 (0.91-12)
Malawi	1995	Male	95 plus	36% (20-54%)	64% (46-80%)	0.51 (0.21-1)	0.78% (0.2-1.7%)	0.74% (0.03-1.7%)	0.29 (0.012-0.69)	1.3 (0.3-2.7)
Malawi	2000	Female	All Ages	7.7% (6.3-9.4%)	92% (91-94%)	0.14 (0.098-0.19)	0.28% (0.16-0.45%)	0.44% (0.22-0.78%)	440 (210-810)	17000 (10000-28000)
Malawi	2000	Female	15 to 19	7.8% (5-11%)	92% (89-95%)	0.14 (0.074-0.25)	0.47% (0.25-0.79%)	0.32% (0.079-0.74%)	4.5 (1-11)	720 (380-1200)
Malawi	2000	Female	20 to 24	9.4% (6.1-14%)	91% (86-94%)	0.2 (0.11-0.35)	0.56% (0.29-0.94%)	0.29% (0.079-0.68%)	8.4 (2.4-19)	1400 (820-2200)
Malawi	2000	Female	25 to 29	11% (6.7-15%)	89% (85-93%)	0.27 (0.15-0.44)	0.43% (0.21-0.75%)	0.27% (0.093-0.62%)	15 (6.2-31)	1600 (980-2600)
Malawi	2000	Female	30 to 34	13% (8.2-19%)	87% (81-92%)	0.33 (0.18-0.55)	0.42% (0.23-0.75%)	0.32% (0.12-0.66%)	19 (8.1-36)	1600 (900-2600)
Malawi	2000	Female	35 to 39	16% (11-23%)	84% (77-89%)	0.33 (0.18-0.56)	0.46% (0.22-0.85%)	0.39% (0.16-0.83%)	19 (8.4-40)	1300 (710-2300)
Malawi	2000	Female	40 to 44	18% (13-26%)	82% (74-87%)	0.35 (0.17-0.57)	0.57% (0.3-1%)	0.54% (0.24-1.1%)	23 (10-45)	1300 (670-2300)
Malawi	2000	Female	45 to 49	18% (12-25%)	82% (75-88%)	0.37 (0.18-0.68)	0.77% (0.36-1.5%)	0.79% (0.32-1.6%)	31 (13-67)	1400 (640-2900)
Malawi	2000	Female	50 to 54	15% (9.3-22%)	85% (78-91%)	0.34 (0.17-0.6)	1.1% (0.57-2.1%)	1.3% (0.58-2.4%)	53 (22-110)	2000 (910-4000)
Malawi	2000	Female	55 to 59	12% (7.4-19%)	88% (81-93%)	0.32 (0.16-0.56)	1.2% (0.55-2.1%)	1.3% (0.59-2.4%)	51 (19-110)	1700 (670-3400)
Malawi	2000	Female	60 to 64	12% (7.5-19%)	88% (81-93%)	0.32 (0.16-0.57)	1.3% (0.65-2.4%)	1.5% (0.72-2.8%)	56 (23-120)	1500 (660-3100)
Malawi	2000	Female	65 to 69	13% (8.6-20%)	87% (80-91%)	0.31 (0.16-0.54)	1.4% (0.68-2.3%)	1.5% (0.75-2.7%)	59 (24-120)	1300 (570-2600)
Malawi	2000	Female	70 to 74	13% (8.3-19%)	87% (81-92%)	0.27 (0.14-0.49)	0.92% (0.29-1.9%)	0.99% (0.26-2.1%)	40 (10-90)	770 (230-1600)
Malawi	2000	Female	75 to 79	12% (7.1-19%)	88% (81-93%)	0.23 (0.1-0.43)	0.89% (0.34-1.7%)	0.99% (0.36-1.9%)	37 (13-76)	530 (190-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	2000	Female	80 to 84	12% (6.6-19%)	88% (81-93%)	0.21 (0.099-0.4)	0.54% (0.12-1.1%)	0.62% (0.12-1.3%)	16 (3-35)	170 (35-360)
Malawi	2000	Female	85 to 89	12% (6.6-18%)	88% (82-93%)	0.2 (0.093-0.36)	0.33% (-0.019-0.78%)	0.37% (-0.05-0.89%)	4.6 (-0.56-11)	35 (-1.9-81)
Malawi	2000	Female	90 to 94	12% (7.5-19%)	88% (81-93%)	0.2 (0.096-0.37)	0.19% (-0.16-0.59%)	0.18% (-0.22-0.64%)	0.59 (-0.7-2.2)	3.6 (-2.9-11)
Malawi	2000	Female	95 plus	14% (7.5-24%)	86% (76-92%)	0.21 (0.083-0.45)	0.5% (0.054-0.97%)	0.51% (-0.03-1.1%)	0.39 (-0.022-0.82)	1.5 (0.17-2.9)
Malawi	2000	Male	All Ages	28% (24-34%)	72% (66-76%)	0.88 (0.63-1.2)	1.6% (0.98-2.3%)	2.3% (1.4-3.5%)	2400 (1400-3700)	110000 (65000-150000)
Malawi	2000	Male	15 to 19	26% (19-36%)	74% (64-81%)	0.92 (0.51-1.5)	3.3% (1.9-5.2%)	3.8% (1.8-6.5%)	54 (24-97)	4900 (2700-8100)
Malawi	2000	Male	20 to 24	39% (28-50%)	61% (50-72%)	1.4 (0.83-2.4)	5.3% (3-8.4%)	5.5% (2.6-9.4%)	140 (65-240)	11000 (6400-18000)
Malawi	2000	Male	25 to 29	46% (35-58%)	54% (42-65%)	1.9 (1.2-2.9)	3.9% (2-6.7%)	3.9% (1.7-7.1%)	190 (99-310)	14000 (7900-21000)
Malawi	2000	Male	30 to 34	51% (38-63%)	49% (37-62%)	2.2 (1.4-3.3)	3.2% (1.7-5.3%)	3.2% (1.5-5.5%)	200 (100-330)	13000 (7200-20000)
Malawi	2000	Male	35 to 39	52% (41-64%)	48% (36-59%)	2.2 (1.3-3.3)	3.2% (1.6-5.4%)	3.3% (1.6-5.7%)	200 (100-330)	11000 (6000-17000)
Malawi	2000	Male	40 to 44	53% (41-64%)	47% (36-59%)	2.2 (1.3-3.4)	3.5% (1.8-5.7%)	3.6% (1.8-6.2%)	200 (110-330)	9700 (5200-15000)
Malawi	2000	Male	45 to 49	54% (42-66%)	46% (34-58%)	2.2 (1.2-3.4)	4.1% (2.2-6.7%)	4.4% (2.2-7.4%)	210 (110-350)	9100 (4900-15000)
Malawi	2000	Male	50 to 54	53% (41-65%)	47% (35-59%)	2 (1.1-3.2)	5.5% (3.1-8.8%)	6.1% (3.4-9.6%)	270 (140-450)	10000 (5500-16000)
Malawi	2000	Male	55 to 59	49% (34-64%)	51% (36-66%)	1.9 (1.1-3.2)	6.5% (3.6-10%)	7.3% (4-12%)	280 (140-460)	8900 (4700-15000)
Malawi	2000	Male	60 to 64	45% (30-61%)	55% (39-70%)	1.7 (0.9-2.9)	5.9% (3.2-9.7%)	6.7% (3.6-11%)	230 (120-400)	6300 (3300-11000)
Malawi	2000	Male	65 to 69	41% (27-57%)	59% (43-73%)	1.3 (0.7-2.1)	4.2% (2.1-7%)	4.8% (2.4-8.1%)	160 (73-280)	3600 (1700-6200)
Malawi	2000	Male	70 to 74	39% (26-55%)	61% (45-74%)	1 (0.54-1.8)	3% (1.2-5.6%)	3.5% (1.4-6.4%)	120 (48-230)	2100 (880-4100)
Malawi	2000	Male	75 to 79	39% (26-55%)	61% (45-74%)	0.93 (0.47-1.6)	2.4% (0.86-4.8%)	2.8% (0.97-5.4%)	80 (27-160)	1100 (400-2200)
Malawi	2000	Male	80 to 84	37% (23-53%)	63% (47-77%)	0.79 (0.39-1.4)	1.9% (0.63-3.8%)	2.2% (0.68-4.3%)	39 (12-74)	410 (130-780)
Malawi	2000	Male	85 to 89	33% (21-49%)	67% (51-79%)	0.65 (0.31-1.1)	1.5% (0.56-2.8%)	1.6% (0.57-3.2%)	15 (5.2-28)	110 (43-210)
Malawi	2000	Male	90 to 94	33% (21-49%)	67% (51-79%)	0.6 (0.29-1.1)	0.97% (0.27-2%)	1% (0.2-2.3%)	2.4 (0.44-5.2)	13 (3.6-28)
Malawi	2000	Male	95 plus	36% (20-56%)	64% (44-80%)	0.6 (0.24-1.2)	1.2% (0.39-2.7%)	1.3% (0.3-3%)	0.54 (0.13-1.3)	2.1 (0.66-4.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	2005	Female	All Ages	7.4% (6.1-9%)	93% (91-94%)	0.14 (0.099-0.2)	0.29% (0.17-0.47%)	0.46% (0.22-0.82%)	440 (200-800)	17000 (10000-28000)
Malawi	2005	Female	15 to 19	7.5% (4.9-11%)	93% (89-95%)	0.15 (0.076-0.25)	0.43% (0.23-0.7%)	0.25% (0.061-0.55%)	4.2 (1-9.6)	790 (430-1300)
Malawi	2005	Female	20 to 24	9.2% (5.8-13%)	91% (87-94%)	0.21 (0.11-0.36)	0.64% (0.33-1%)	0.29% (0.083-0.66%)	7.3 (2.4-16)	1400 (860-2200)
Malawi	2005	Female	25 to 29	10% (6.6-15%)	90% (85-93%)	0.28 (0.15-0.45)	0.43% (0.22-0.78%)	0.24% (0.081-0.57%)	13 (5.3-26)	1700 (1000-2600)
Malawi	2005	Female	30 to 34	13% (7.9-18%)	87% (82-92%)	0.35 (0.18-0.57)	0.35% (0.19-0.61%)	0.23% (0.093-0.5%)	16 (6.8-32)	1500 (910-2400)
Malawi	2005	Female	35 to 39	16% (10-22%)	84% (78-90%)	0.34 (0.18-0.57)	0.36% (0.19-0.66%)	0.28% (0.11-0.61%)	16 (7.3-33)	1200 (680-2000)
Malawi	2005	Female	40 to 44	18% (12-25%)	82% (75-88%)	0.36 (0.18-0.63)	0.43% (0.23-0.78%)	0.39% (0.18-0.78%)	19 (8.5-37)	1100 (570-1900)
Malawi	2005	Female	45 to 49	18% (11-25%)	82% (75-89%)	0.39 (0.19-0.73)	0.64% (0.31-1.2%)	0.64% (0.26-1.3%)	26 (11-56)	1200 (600-2400)
Malawi	2005	Female	50 to 54	14% (8.8-21%)	86% (79-91%)	0.35 (0.17-0.64)	0.99% (0.49-1.8%)	1.1% (0.49-2%)	41 (17-86)	1600 (710-3200)
Malawi	2005	Female	55 to 59	12% (7.3-18%)	88% (82-93%)	0.33 (0.16-0.58)	1.2% (0.55-2.2%)	1.3% (0.59-2.5%)	53 (20-110)	1700 (720-3600)
Malawi	2005	Female	60 to 64	12% (7.2-19%)	88% (81-93%)	0.34 (0.16-0.62)	1.3% (0.66-2.4%)	1.5% (0.76-2.8%)	62 (24-130)	1700 (720-3500)
Malawi	2005	Female	65 to 69	13% (8.5-20%)	87% (80-92%)	0.33 (0.17-0.57)	1.5% (0.76-2.5%)	1.7% (0.83-2.9%)	67 (28-130)	1500 (680-3000)
Malawi	2005	Female	70 to 74	13% (8.2-19%)	87% (81-92%)	0.29 (0.14-0.5)	1% (0.41-2.1%)	1.2% (0.41-2.3%)	45 (14-98)	860 (310-1800)
Malawi	2005	Female	75 to 79	12% (6.8-19%)	88% (81-93%)	0.24 (0.11-0.45)	1% (0.46-1.8%)	1.1% (0.5-2.1%)	41 (16-81)	580 (240-1100)
Malawi	2005	Female	80 to 84	11% (6.6-18%)	89% (82-93%)	0.22 (0.1-0.41)	0.65% (0.19-1.3%)	0.76% (0.2-1.5%)	20 (5.2-43)	210 (59-450)
Malawi	2005	Female	85 to 89	11% (6.5-18%)	89% (82-93%)	0.21 (0.097-0.39)	0.42% (0.028-0.98%)	0.48% (0.018-1.1%)	6.5 (0.23-15)	49 (3.1-110)
Malawi	2005	Female	90 to 94	12% (7.3-20%)	88% (80-93%)	0.21 (0.099-0.4)	0.27% (-0.094-0.75%)	0.28% (-0.16-0.86%)	1.2 (-0.71-3.8)	7 (-2.5-20)
Malawi	2005	Female	95 plus	14% (7.2-24%)	86% (76-93%)	0.23 (0.087-0.48)	0.59% (0.12-1.2%)	0.62% (0.083-1.3%)	0.54 (0.074-1.2)	2 (0.42-4)
Malawi	2005	Male	All Ages	28% (23-33%)	72% (67-77%)	0.92 (0.65-1.2)	1.9% (1.2-2.7%)	2.8% (1.7-4.1%)	2800 (1600-4200)	120000 (75000-170000)
Malawi	2005	Male	15 to 19	25% (18-35%)	75% (65-82%)	0.94 (0.52-1.5)	2.9% (1.7-4.7%)	3.2% (1.5-5.5%)	64 (29-110)	5900 (3200-9400)
Malawi	2005	Male	20 to 24	38% (28-49%)	62% (51-72%)	1.5 (0.86-2.3)	6% (3.4-9.3%)	6.2% (3.1-11%)	160 (78-280)	13000 (7400-21000)
Malawi	2005	Male	25 to 29	46% (34-56%)	54% (44-66%)	2 (1.2-3)	4.9% (2.6-8.2%)	5% (2.2-9.1%)	230 (120-360)	16000 (9500-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	2005	Male	30 to 34	50% (37-63%)	50% (37-63%)	2.3 (1.4-3.5)	3.6% (1.9-5.8%)	3.5% (1.7-6%)	240 (120-370)	15000 (8400-22000)
Malawi	2005	Male	35 to 39	51% (40-63%)	49% (37-60%)	2.3 (1.4-3.6)	3.3% (1.8-5.3%)	3.4% (1.7-5.6%)	230 (120-350)	12000 (6800-19000)
Malawi	2005	Male	40 to 44	52% (41-64%)	48% (36-59%)	2.3 (1.3-3.6)	3.5% (1.9-5.5%)	3.7% (1.8-5.9%)	220 (110-350)	11000 (5700-17000)
Malawi	2005	Male	45 to 49	54% (41-65%)	46% (35-59%)	2.3 (1.3-3.8)	4.3% (2.3-6.9%)	4.6% (2.4-7.6%)	230 (120-380)	9900 (5300-16000)
Malawi	2005	Male	50 to 54	53% (41-65%)	47% (35-59%)	2.1 (1.2-3.4)	5.8% (3.3-9%)	6.3% (3.5-10%)	270 (140-440)	10000 (5400-16000)
Malawi	2005	Male	55 to 59	48% (34-64%)	52% (36-66%)	2 (1.1-3.3)	7.2% (4.1-11%)	8.1% (4.6-13%)	350 (170-570)	11000 (5800-18000)
Malawi	2005	Male	60 to 64	44% (29-60%)	56% (40-71%)	1.8 (0.93-3.1)	6.6% (3.6-11%)	7.5% (4.1-12%)	310 (150-520)	8400 (4200-14000)
Malawi	2005	Male	65 to 69	40% (27-55%)	60% (45-73%)	1.3 (0.72-2.2)	4.7% (2.5-8%)	5.4% (2.8-9.2%)	210 (97-360)	4700 (2300-8100)
Malawi	2005	Male	70 to 74	38% (25-54%)	62% (46-75%)	1.1 (0.56-1.9)	3.5% (1.5-6.5%)	4% (1.6-7.7%)	150 (61-290)	2700 (1100-5200)
Malawi	2005	Male	75 to 79	39% (25-54%)	61% (46-75%)	0.97 (0.48-1.8)	2.9% (1-5.5%)	3.3% (1.2-6.3%)	100 (37-200)	1400 (530-2700)
Malawi	2005	Male	80 to 84	36% (22-53%)	64% (47-78%)	0.83 (0.4-1.5)	2.2% (0.8-4.3%)	2.5% (0.88-4.9%)	49 (18-96)	510 (190-990)
Malawi	2005	Male	85 to 89	33% (20-47%)	67% (53-80%)	0.67 (0.32-1.2)	1.7% (0.68-3.3%)	1.9% (0.73-3.8%)	20 (7.7-39)	150 (60-290)
Malawi	2005	Male	90 to 94	32% (20-48%)	68% (52-80%)	0.62 (0.29-1.2)	1.2% (0.37-2.5%)	1.3% (0.31-2.8%)	4.6 (1.1-10)	25 (7.9-54)
Malawi	2005	Male	95 plus	35% (19-54%)	65% (46-81%)	0.61 (0.25-1.2)	1.5% (0.49-3.2%)	1.6% (0.4-3.7%)	0.97 (0.24-2.2)	3.6 (1.2-7.8)
Malawi	2010	Female	All Ages	6.6% (5.4-8.1%)	93% (92-95%)	0.15 (0.098-0.22)	0.37% (0.22-0.56%)	0.62% (0.32-1.1%)	540 (260-970)	20000 (12000-31000)
Malawi	2010	Female	15 to 19	6.1% (3.6-9.3%)	94% (91-96%)	0.14 (0.067-0.25)	0.39% (0.22-0.63%)	0.2% (0.059-0.47%)	4.4 (1.3-10)	870 (480-1400)
Malawi	2010	Female	20 to 24	7.5% (4.5-11%)	92% (89-95%)	0.2 (0.1-0.36)	0.73% (0.41-1.1%)	0.32% (0.1-0.73%)	8.1 (2.6-17)	1700 (1000-2600)
Malawi	2010	Female	25 to 29	8.6% (5.1-13%)	91% (87-95%)	0.28 (0.14-0.49)	0.56% (0.3-0.95%)	0.31% (0.1-0.67%)	13 (5.4-25)	1800 (1100-2600)
Malawi	2010	Female	30 to 34	10% (6.1-16%)	90% (84-94%)	0.33 (0.16-0.6)	0.46% (0.27-0.75%)	0.3% (0.12-0.59%)	16 (7.2-31)	1700 (1000-2500)
Malawi	2010	Female	35 to 39	13% (7.9-19%)	87% (81-92%)	0.34 (0.16-0.6)	0.46% (0.26-0.75%)	0.36% (0.16-0.69%)	18 (8.5-32)	1400 (820-2100)
Malawi	2010	Female	40 to 44	15% (9.5-21%)	85% (79-91%)	0.35 (0.16-0.65)	0.56% (0.33-0.93%)	0.51% (0.25-0.92%)	19 (9.2-35)	1100 (670-1900)
Malawi	2010	Female	45 to 49	15% (8.7-23%)	85% (77-91%)	0.4 (0.19-0.76)	0.85% (0.45-1.5%)	0.89% (0.41-1.7%)	26 (12-50)	1200 (630-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	2010	Female	50 to 54	14% (9-22%)	86% (78-91%)	0.42 (0.2-0.8)	1.4% (0.73-2.6%)	1.6% (0.76-3%)	43 (18-86)	1700 (780-3200)
Malawi	2010	Female	55 to 59	13% (7.8-20%)	87% (80-92%)	0.43 (0.21-0.79)	1.7% (0.8-3.1%)	2% (0.93-3.9%)	55 (22-110)	1800 (760-3600)
Malawi	2010	Female	60 to 64	13% (7.7-21%)	87% (79-92%)	0.43 (0.21-0.78)	1.8% (0.9-3.3%)	2.2% (1.1-4%)	80 (33-160)	2200 (960-4300)
Malawi	2010	Female	65 to 69	14% (8.5-21%)	86% (79-91%)	0.42 (0.21-0.77)	1.9% (1-3.4%)	2.3% (1.2-4.1%)	93 (40-180)	2100 (970-4100)
Malawi	2010	Female	70 to 74	14% (8.4-20%)	86% (80-92%)	0.36 (0.17-0.67)	1.4% (0.55-2.8%)	1.6% (0.55-3.3%)	66 (21-140)	1200 (440-2500)
Malawi	2010	Female	75 to 79	12% (7.2-20%)	88% (80-93%)	0.3 (0.13-0.58)	1.2% (0.53-2.3%)	1.4% (0.61-2.7%)	55 (21-110)	780 (300-1500)
Malawi	2010	Female	80 to 84	12% (6.7-18%)	88% (82-93%)	0.27 (0.12-0.53)	0.83% (0.27-1.7%)	0.97% (0.3-1.9%)	29 (8.3-60)	300 (89-610)
Malawi	2010	Female	85 to 89	12% (6.8-19%)	88% (81-93%)	0.26 (0.11-0.52)	0.58% (0.11-1.3%)	0.66% (0.11-1.5%)	11 (1.8-25)	80 (15-190)
Malawi	2010	Female	90 to 94	13% (7.5-20%)	87% (80-93%)	0.27 (0.12-0.52)	0.4% (-0.062-1.1%)	0.43% (-0.11-1.3%)	2.3 (-0.63-6.7)	13 (-1.8-34)
Malawi	2010	Female	95 plus	14% (7.5-25%)	86% (75-93%)	0.28 (0.1-0.61)	0.78% (0.19-1.5%)	0.78% (0.13-1.8%)	0.99 (0.17-2.2)	3.6 (0.92-7.7)
Malawi	2010	Male	All Ages	25% (21-30%)	75% (70-79%)	1 (0.67-1.4)	2.4% (1.6-3.4%)	3.7% (2.2-5.4%)	3500 (2000-5200)	150000 (93000-210000)
Malawi	2010	Male	15 to 19	19% (12-28%)	81% (72-88%)	0.82 (0.41-1.5)	2.4% (1.3-4%)	2.3% (1-4.4%)	63 (27-120)	6000 (3300-9700)
Malawi	2010	Male	20 to 24	33% (24-43%)	67% (57-76%)	1.5 (0.85-2.5)	6.4% (3.7-9.8%)	6.7% (3.3-11%)	190 (94-330)	16000 (9000-25000)
Malawi	2010	Male	25 to 29	42% (32-53%)	58% (47-68%)	2.2 (1.3-3.5)	6.2% (3.3-9.9%)	6.5% (3-11%)	270 (140-420)	19000 (11000-29000)
Malawi	2010	Male	30 to 34	47% (36-59%)	53% (41-64%)	2.5 (1.5-3.9)	5.1% (2.8-8%)	5.2% (2.6-8.8%)	300 (160-460)	19000 (11000-28000)
Malawi	2010	Male	35 to 39	48% (38-60%)	52% (40-62%)	2.5 (1.5-4.1)	4.5% (2.6-6.8%)	4.7% (2.5-7.4%)	280 (160-430)	16000 (9400-23000)
Malawi	2010	Male	40 to 44	47% (36-59%)	53% (41-64%)	2.4 (1.3-4)	4.7% (2.7-7.1%)	5% (2.8-7.7%)	260 (140-410)	13000 (7300-20000)
Malawi	2010	Male	45 to 49	50% (38-60%)	50% (40-62%)	2.5 (1.4-4.2)	5.7% (3.2-8.8%)	6.2% (3.4-9.9%)	260 (140-410)	11000 (6400-17000)
Malawi	2010	Male	50 to 54	52% (41-63%)	48% (37-59%)	2.5 (1.4-4.1)	7.7% (4.3-12%)	8.7% (4.8-13%)	300 (160-480)	11000 (6200-18000)
Malawi	2010	Male	55 to 59	50% (35-65%)	50% (35-65%)	2.5 (1.3-4.4)	9.4% (5.4-14%)	11% (6.2-16%)	360 (200-560)	12000 (6600-18000)
Malawi	2010	Male	60 to 64	47% (30-64%)	53% (36-70%)	2.3 (1.2-4)	8.6% (4.7-14%)	10% (5.4-16%)	410 (210-660)	11000 (5900-18000)
Malawi	2010	Male	65 to 69	43% (29-59%)	57% (41-71%)	1.7 (0.91-3)	6.3% (3.1-11%)	7.3% (3.6-13%)	310 (150-530)	7000 (3400-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	2010	Male	70 to 74	40% (27-56%)	60% (44-73%)	1.4 (0.71-2.4)	4.6% (1.9-8.4%)	5.3% (2.2-9.7%)	210 (90-420)	3900 (1600-7600)
Malawi	2010	Male	75 to 79	39% (25-55%)	61% (45-75%)	1.2 (0.56-2.1)	3.6% (1.4-6.9%)	4.1% (1.6-7.9%)	140 (54-290)	2000 (750-4000)
Malawi	2010	Male	80 to 84	36% (21-52%)	64% (48-79%)	0.97 (0.46-1.8)	2.8% (0.94-5.3%)	3.2% (1-6.2%)	69 (22-140)	720 (250-1400)
Malawi	2010	Male	85 to 89	33% (20-48%)	67% (52-80%)	0.81 (0.37-1.5)	2.1% (0.83-4.2%)	2.4% (0.88-4.8%)	28 (10-57)	210 (82-430)
Malawi	2010	Male	90 to 94	33% (21-48%)	67% (52-79%)	0.75 (0.34-1.4)	1.5% (0.42-3.3%)	1.6% (0.38-3.8%)	7.1 (1.6-16)	38 (11-85)
Malawi	2010	Male	95 plus	35% (20-55%)	65% (45-80%)	0.75 (0.29-1.5)	1.9% (0.6-4.1%)	2.1% (0.52-4.7%)	2.1 (0.53-4.7)	7.6 (2.4-16)
Malawi	2016	Female	All Ages	8.3% (6.7-10%)	92% (90-93%)	0.18 (0.096-0.31)	0.48% (0.28-0.82%)	0.76% (0.31-1.5%)	570 (210-1100)	22000 (12000-38000)
Malawi	2016	Female	15 to 19	8.6% (5.2-13%)	91% (87-95%)	0.19 (0.078-0.35)	0.46% (0.26-0.77%)	0.26% (0.058-0.66%)	5.3 (1.2-14)	1100 (600-1800)
Malawi	2016	Female	20 to 24	11% (6.6-17%)	89% (83-93%)	0.28 (0.12-0.54)	0.9% (0.54-1.4%)	0.41% (0.1-0.97%)	9.5 (2.5-22)	2100 (1200-3200)
Malawi	2016	Female	25 to 29	12% (7.1-18%)	88% (82-93%)	0.37 (0.16-0.73)	0.85% (0.49-1.4%)	0.49% (0.16-1.1%)	16 (6-35)	2300 (1400-3500)
Malawi	2016	Female	30 to 34	14% (8.2-22%)	86% (78-92%)	0.42 (0.18-0.85)	0.79% (0.46-1.3%)	0.56% (0.21-1.2%)	19 (8-39)	2000 (1200-3200)
Malawi	2016	Female	35 to 39	16% (9.6-25%)	84% (75-90%)	0.4 (0.16-0.79)	0.83% (0.49-1.4%)	0.71% (0.31-1.4%)	22 (9.6-45)	1700 (1000-2900)
Malawi	2016	Female	40 to 44	18% (11-26%)	82% (74-89%)	0.4 (0.16-0.84)	0.95% (0.54-1.6%)	0.76% (0.44-1.9%)	24 (11-48)	1500 (820-2600)
Malawi	2016	Female	45 to 49	17% (9.6-26%)	83% (74-90%)	0.42 (0.16-0.93)	1.2% (0.62-2.3%)	1.4% (0.6-2.9%)	30 (12-64)	1400 (700-2800)
Malawi	2016	Female	50 to 54	15% (8.9-23%)	85% (77-91%)	0.41 (0.16-0.87)	1.6% (0.75-3.2%)	2% (0.85-4%)	41 (16-86)	1600 (690-3200)
Malawi	2016	Female	55 to 59	13% (7.8-20%)	87% (80-92%)	0.4 (0.17-0.85)	1.7% (0.69-3.5%)	2.2% (0.8-4.5%)	49 (17-110)	1600 (580-3500)
Malawi	2016	Female	60 to 64	13% (7.8-21%)	87% (79-92%)	0.4 (0.16-0.81)	1.7% (0.69-3.5%)	2.2% (0.83-4.4%)	65 (23-140)	1800 (670-3800)
Malawi	2016	Female	65 to 69	14% (8.9-21%)	86% (79-91%)	0.39 (0.16-0.77)	1.8% (0.74-3.5%)	2.2% (0.82-4.3%)	96 (34-200)	2200 (870-4500)
Malawi	2016	Female	70 to 74	14% (8.5-20%)	86% (80-92%)	0.33 (0.14-0.69)	1.2% (0.33-2.9%)	1.4% (0.29-3.4%)	74 (16-180)	1400 (360-3300)
Malawi	2016	Female	75 to 79	12% (7.3-20%)	88% (80-93%)	0.28 (0.11-0.59)	1.2% (0.42-2.4%)	1.3% (0.45-2.8%)	67 (21-140)	960 (320-2000)
Malawi	2016	Female	80 to 84	12% (6.7-19%)	88% (81-93%)	0.25 (0.099-0.53)	0.74% (0.12-1.8%)	0.87% (0.12-2.1%)	32 (4.2-78)	330 (49-810)
Malawi	2016	Female	85 to 89	12% (6.7-19%)	88% (81-93%)	0.24 (0.094-0.52)	0.51% (-0.0016-1.5%)	0.59% (-0.022-1.7%)	13 (-0.51-38)	97 (-0.31-280)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Malawi	2016	Female	90 to 94	13% (7.5-20%)	87% (80-92%)	0.25 (0.098-0.55)	0.35% (-0.12-1.2%)	0.37% (-0.19-1.4%)	3 (-1.6-11)	17 (-6.2-59)
Malawi	2016	Female	95 plus	15% (7.7-25%)	85% (75-92%)	0.26 (0.087-0.61)	0.66% (0.1-1.6%)	0.7% (0.01-1.8%)	1.4 (0.021-3.5)	5.1 (0.77-12)
Malawi	2016	Male	All Ages	28% (23-34%)	72% (66-77%)	1.1 (0.58-1.8)	2.8% (1.6-4.3%)	4% (2.1-6.5%)	3400 (1700-5700)	150000 (85000-230000)
Malawi	2016	Male	15 to 19	25% (16-37%)	75% (63-84%)	1 (0.46-2)	2.6% (1.3-4.4%)	2.5% (1-5%)	72 (29-140)	7000 (3600-12000)
Malawi	2016	Male	20 to 24	38% (26-51%)	62% (49-74%)	1.7 (0.76-3.1)	6.3% (3.4-10%)	6.4% (2.8-11%)	190 (81-350)	17000 (8600-27000)
Malawi	2016	Male	25 to 29	45% (31-59%)	55% (41-69%)	2.2 (1.1-4)	6.5% (3.5-11%)	6.7% (3.2-12%)	280 (130-470)	21000 (11000-33000)
Malawi	2016	Male	30 to 34	49% (34-63%)	51% (37-66%)	2.5 (1.2-4.5)	5.6% (3.1-9%)	5.7% (2.8-9.8%)	290 (140-480)	19000 (10000-29000)
Malawi	2016	Male	35 to 39	51% (37-64%)	49% (36-63%)	2.5 (1.2-4.8)	5.6% (3-8.9%)	6% (2.9-10%)	300 (150-500)	17000 (9200-27000)
Malawi	2016	Male	40 to 44	52% (39-65%)	48% (35-61%)	2.6 (1.2-4.8)	6.2% (3.4-9.9%)	6.9% (3.5-11%)	300 (150-510)	15000 (7900-25000)
Malawi	2016	Male	45 to 49	53% (38-66%)	47% (34-62%)	2.5 (1.2-5.1)	7.2% (3.8-12%)	8.3% (4.1-14%)	290 (130-480)	12000 (6100-20000)
Malawi	2016	Male	50 to 54	53% (39-66%)	47% (34-61%)	2.4 (1-4.7)	8.4% (4.3-13%)	9.9% (4.9-16%)	290 (140-490)	11000 (5400-18000)
Malawi	2016	Male	55 to 59	50% (35-65%)	50% (35-65%)	2.4 (1.1-4.8)	9.4% (4.8-15%)	11% (5.7-18%)	320 (150-520)	10000 (5000-17000)
Malawi	2016	Male	60 to 64	46% (31-63%)	54% (37-69%)	2.1 (0.93-4.2)	8.1% (3.9-14%)	9.6% (4.5-16%)	310 (140-570)	8700 (4000-15000)
Malawi	2016	Male	65 to 69	43% (29-59%)	57% (41-71%)	1.6 (0.73-3.1)	5.8% (2.3-11%)	6.8% (2.7-13%)	300 (110-580)	6700 (2700-13000)
Malawi	2016	Male	70 to 74	40% (26-55%)	60% (45-74%)	1.3 (0.57-2.5)	4.1% (1.3-8.6%)	4.9% (1.5-10%)	230 (73-490)	4200 (1300-8800)
Malawi	2016	Male	75 to 79	39% (25-55%)	61% (45-75%)	1.1 (0.46-2.1)	3.2% (0.92-7.1%)	3.7% (1.1-8.2%)	150 (41-350)	2100 (590-4900)
Malawi	2016	Male	80 to 84	36% (22-53%)	64% (47-78%)	0.92 (0.38-1.8)	2.5% (0.65-5.6%)	2.8% (0.68-6.6%)	74 (17-170)	780 (200-1800)
Malawi	2016	Male	85 to 89	33% (20-48%)	67% (52-80%)	0.75 (0.3-1.6)	1.9% (0.53-4.2%)	2.1% (0.52-4.9%)	30 (7.7-71)	230 (66-520)
Malawi	2016	Male	90 to 94	33% (21-47%)	67% (53-79%)	0.69 (0.27-1.5)	1.3% (0.24-3.5%)	1.4% (0.15-4%)	7.7 (0.83-21)	42 (7.6-110)
Malawi	2016	Male	95 plus	35% (19-54%)	65% (46-81%)	0.68 (0.24-1.5)	1.7% (0.36-4.3%)	1.8% (0.25-4.7%)	2.3 (0.33-6.1)	8.5 (1.9-22)
Mauritius	1990	Female	All Ages	26% (23-31%)	74% (69-77%)	0.29 (0.23-0.38)	0.17% (-0.29-0.7%)	-0.45% (-1.5-0.71%)	-13 (-42-20)	230 (-390-950)
Mauritius	1990	Female	15 to 19	22% (15-31%)	78% (69-85%)	0.26 (0.15-0.43)	0.81% (0.42-1.3%)	1.4% (0.39-2.9%)	0.46 (0.12-0.93)	57 (30-93)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	1990	Female	20 to 24	28% (18-38%)	72% (62-82%)	0.33 (0.19-0.53)	1.2% (0.68-1.9%)	1.5% (0.36-3.1%)	0.65 (0.15-1.3)	97 (53-150)
Mauritius	1990	Female	25 to 29	32% (21-43%)	68% (57-79%)	0.37 (0.22-0.56)	1.2% (0.71-1.6%)	1.3% (0.45-2.5%)	0.56 (0.19-1)	82 (48-120)
Mauritius	1990	Female	30 to 34	34% (23-46%)	66% (54-77%)	0.39 (0.23-0.62)	1% (0.64-1.5%)	1.3% (0.51-2.2%)	0.68 (0.27-1.2)	76 (46-110)
Mauritius	1990	Female	35 to 39	34% (23-45%)	66% (55-77%)	0.42 (0.24-0.65)	0.89% (0.5-1.4%)	1.2% (0.41-2.2%)	0.66 (0.23-1.2)	64 (35-98)
Mauritius	1990	Female	40 to 44	34% (23-45%)	66% (55-77%)	0.43 (0.25-0.68)	0.88% (0.4-1.4%)	1.3% (0.38-2.4%)	0.94 (0.28-1.7)	65 (28-100)
Mauritius	1990	Female	45 to 49	36% (24-46%)	64% (54-76%)	0.46 (0.27-0.74)	0.59% (-0.018-1.3%)	0.8% (-0.23-2.1%)	0.69 (-0.2-1.8)	42 (-1.2-90)
Mauritius	1990	Female	50 to 54	36% (26-48%)	64% (52-74%)	0.53 (0.32-0.83)	0.11% (-0.74-1.1%)	0.0046% (-1.4-1.5%)	0.005 (-1.5-1.7)	8 (-52-72)
Mauritius	1990	Female	55 to 59	35% (24-46%)	65% (54-76%)	0.6 (0.34-0.96)	-0.16% (-1.3-1.2%)	-0.33% (-2.1-6%)	-0.52 (-3.1-2.4)	-12 (-98-88)
Mauritius	1990	Female	60 to 64	33% (22-44%)	67% (56-78%)	0.61 (0.34-1)	-0.59% (-1.9-0.83%)	-0.84% (-2.6-1%)	-2 (-6.2-2.5)	-52 (-170-75)
Mauritius	1990	Female	65 to 69	32% (22-42%)	68% (58-78%)	0.58 (0.33-0.89)	-0.68% (-1.9-0.59%)	-0.91% (-2.5-0.71%)	-2.9 (-7.9-2.3)	-64 (-180-54)
Mauritius	1990	Female	70 to 74	31% (22-41%)	69% (59-78%)	0.56 (0.31-0.9)	-0.53% (-1.8-0.8%)	-0.69% (-2.2-0.95%)	-2.4 (-7.8-3.3)	-42 (-140-62)
Mauritius	1990	Female	75 to 79	32% (21-44%)	68% (56-79%)	0.57 (0.31-0.95)	-0.69% (-2-0.71%)	-0.88% (-2.5-0.81%)	-3.2 (-9.3-3.1)	-44 (-130-45)
Mauritius	1990	Female	80 to 84	32% (21-44%)	68% (56-79%)	0.56 (0.31-0.85)	-0.78% (-1.9-0.45%)	-0.98% (-2.4-0.55%)	-3.2 (-7.9-1.8)	-32 (-80-19)
Mauritius	1990	Female	85 to 89	33% (22-44%)	67% (56-78%)	0.55 (0.3-0.89)	-0.77% (-1.9-0.39%)	-0.98% (-2.4-0.43%)	-2 (-5-0.89)	-14 (-35-7.4)
Mauritius	1990	Female	90 to 94	34% (23-47%)	66% (53-77%)	0.56 (0.3-0.97)	-0.77% (-1.9-0.43%)	-1% (-2.3-0.46%)	-0.76 (-1.7-0.35)	-3.6 (-8.5-2)
Mauritius	1990	Female	95 plus	36% (22-53%)	64% (47-78%)	0.58 (0.27-1.1)	-0.74% (-2.1-0.78%)	-0.97% (-2.6-0.8%)	-0.16 (-0.43-0.13)	-0.48 (-1.4-0.52)
Mauritius	1990	Male	All Ages	55% (48-61%)	45% (39-52%)	2.1 (1.8-2.5)	8.6% (6.5-11%)	10% (7.1-14%)	400 (280-530)	15000 (11000-19000)
Mauritius	1990	Male	15 to 19	45% (32-59%)	55% (41-68%)	1 (0.61-1.6)	2.8% (1.8-4.2%)	5.2% (3-8%)	2 (1.2-3.1)	190 (130-290)
Mauritius	1990	Male	20 to 24	65% (51-76%)	35% (24-49%)	2.2 (1.5-3.1)	7.1% (4.9-9.9%)	12% (7.3-17%)	6.7 (4.1-9.5)	570 (390-790)
Mauritius	1990	Male	25 to 29	69% (57-79%)	31% (21-43%)	2.8 (2-3.7)	9.6% (6.8-13%)	14% (9.1-19%)	11 (7.6-16)	870 (620-1100)
Mauritius	1990	Male	30 to 34	71% (59-80%)	29% (20-41%)	3.4 (2.4-4.4)	12% (8.9-16%)	16% (12-22%)	18 (13-24)	1200 (890-1600)
Mauritius	1990	Male	35 to 39	73% (62-81%)	27% (19-38%)	3.9 (2.7-5.1)	14% (11-19%)	19% (13-24%)	28 (21-37)	1600 (1200-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	1990	Male	40 to 44	73% (62-82%)	27% (18-38%)	4.2 (3-5.4)	15% (11-19%)	19% (13-24%)	35 (25-45)	1800 (1300-2300)
Mauritius	1990	Male	45 to 49	72% (60-82%)	28% (18-40%)	4.3 (3-5.8)	15% (11-19%)	19% (13-24%)	37 (26-49)	1700 (1200-2100)
Mauritius	1990	Male	50 to 54	72% (60-81%)	28% (19-40%)	4.4 (3-5.9)	14% (9.3-19%)	17% (11-22%)	45 (29-60)	1700 (1100-2300)
Mauritius	1990	Male	55 to 59	73% (59-83%)	27% (17-41%)	4.3 (2.8-5.9)	13% (8.2-18%)	15% (9.4-22%)	51 (31-72)	1700 (1100-2400)
Mauritius	1990	Male	60 to 64	68% (53-80%)	32% (20-47%)	3.9 (2.5-5.5)	12% (6.9-17%)	13% (7.5-19%)	61 (34-90)	1700 (970-2500)
Mauritius	1990	Male	65 to 69	59% (43-72%)	41% (28-57%)	3.1 (2-4.4)	8.8% (4.6-14%)	9.8% (4.9-15%)	51 (26-79)	1200 (610-1800)
Mauritius	1990	Male	70 to 74	53% (39-67%)	47% (33-61%)	2.5 (1.5-3.6)	6.2% (2.4-10%)	6.8% (2.5-12%)	30 (11-51)	560 (210-940)
Mauritius	1990	Male	75 to 79	50% (35-66%)	50% (34-65%)	2 (1.2-3.2)	4.2% (1-8.1%)	4.6% (0.97-8.9%)	17 (3.6-33)	250 (61-480)
Mauritius	1990	Male	80 to 84	47% (32-62%)	53% (38-68%)	1.7 (1-2.5)	2.7% (0.49-5.6%)	2.9% (0.38-6.2%)	6.6 (0.88-14)	73 (13-150)
Mauritius	1990	Male	85 to 89	44% (29-60%)	56% (40-71%)	1.4 (0.8-2.2)	1.6% (-0.075-4%)	1.7% (-0.26-4.4%)	1.5 (-0.23-3.9)	12 (-0.57-31)
Mauritius	1990	Male	90 to 94	45% (30-60%)	55% (40-70%)	1.2 (0.7-2.1)	0.78% (-0.73-3%)	0.72% (-1.1-3.3%)	0.14 (-0.21-0.65)	0.92 (-0.86-3.5)
Mauritius	1990	Male	95 plus	48% (29-67%)	52% (33-71%)	1.1 (0.5-2.1)	0.43% (-1.2-2.9%)	0.22% (-1.7-3%)	0.0063 (-0.048-0.086)	0.048 (-0.13-0.32)
Mauritius	1995	Female	All Ages	27% (23-31%)	73% (69-77%)	0.31 (0.24-0.41)	0.22% (-0.29-0.79%)	-0.36% (-1.4-0.89%)	-11 (-44-26)	310 (-410-1100)
Mauritius	1995	Female	15 to 19	22% (15-31%)	78% (69-85%)	0.27 (0.15-0.45)	0.79% (0.44-1.2%)	1.4% (0.39-3%)	0.34 (0.097-0.7)	48 (27-79)
Mauritius	1995	Female	20 to 24	27% (18-37%)	73% (63-82%)	0.34 (0.19-0.57)	1.2% (0.72-1.8%)	1.5% (0.36-3.2%)	0.49 (0.12-1)	86 (50-130)
Mauritius	1995	Female	25 to 29	32% (21-43%)	68% (57-79%)	0.38 (0.23-0.58)	1.2% (0.77-1.6%)	1.4% (0.5-2.5%)	0.53 (0.2-0.97)	88 (55-130)
Mauritius	1995	Female	30 to 34	33% (22-47%)	67% (53-78%)	0.4 (0.23-0.63)	1.1% (0.67-1.5%)	1.4% (0.59-2.3%)	0.68 (0.29-1.1)	81 (49-120)
Mauritius	1995	Female	35 to 39	33% (22-45%)	67% (55-78%)	0.43 (0.24-0.68)	1% (0.62-1.6%)	1.5% (0.73-2.6%)	0.95 (0.45-1.6)	81 (47-120)
Mauritius	1995	Female	40 to 44	33% (23-44%)	67% (56-77%)	0.44 (0.25-0.69)	1% (0.53-1.6%)	1.7% (0.76-2.8%)	1.3 (0.61-2.3)	85 (43-130)
Mauritius	1995	Female	45 to 49	35% (24-46%)	65% (54-76%)	0.47 (0.27-0.76)	0.68% (0.043-1.4%)	0.99% (-0.16-2.3%)	1.1 (-0.17-2.4)	59 (3.6-120)
Mauritius	1995	Female	50 to 54	36% (25-48%)	64% (52-75%)	0.54 (0.32-0.85)	0.2% (-0.67-1.1%)	0.18% (-1.2-1.7%)	0.25 (-1.6-2.3)	17 (-54-99)
Mauritius	1995	Female	55 to 59	34% (24-45%)	66% (55-76%)	0.61 (0.35-0.97)	-0.05% (-1.2-1.3%)	-0.14% (-1.8-1.8%)	-0.25 (-3.2-3.2)	-4.5 (-100-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	1995	Female	60 to 64	32% (22-44%)	68% (56-78%)	0.62 (0.35-1)	-0.44% (-1.7-1.1%)	-0.63% (-2.4-1.3%)	-1.6 (-6.1-3.4)	-42 (-170-110)
Mauritius	1995	Female	65 to 69	31% (22-42%)	69% (58-78%)	0.59 (0.34-0.94)	-0.55% (-1.8-0.82%)	-0.74% (-2.3-1%)	-2.5 (-7.8-3.3)	-54 (-180-80)
Mauritius	1995	Female	70 to 74	31% (22-41%)	69% (59-78%)	0.58 (0.32-0.97)	-0.43% (-1.8-1%)	-0.56% (-2.3-1.3%)	-2.2 (-8.8-5)	-39 (-160-96)
Mauritius	1995	Female	75 to 79	31% (20-43%)	69% (57-80%)	0.59 (0.32-1)	-0.62% (-2-0.86%)	-0.82% (-2.6-1%)	-3 (-9.5-3.8)	-42 (-130-55)
Mauritius	1995	Female	80 to 84	31% (21-44%)	69% (56-79%)	0.58 (0.33-0.91)	-0.72% (-1.9-0.6%)	-0.93% (-2.3-0.68%)	-3.3 (-8.3-2.4)	-33 (-85-26)
Mauritius	1995	Female	85 to 89	32% (21-44%)	68% (56-79%)	0.56 (0.31-0.92)	-0.73% (-1.9-0.45%)	-0.95% (-2.4-0.48%)	-2.3 (-6-1.2)	-16 (-43-10)
Mauritius	1995	Female	90 to 94	33% (22-46%)	67% (54-78%)	0.57 (0.31-1)	-0.73% (-1.8-0.47%)	-0.97% (-2.3-0.49%)	-1 (-2.4-0.5)	-4.6 (-12-3)
Mauritius	1995	Female	95 plus	36% (22-51%)	64% (49-78%)	0.59 (0.27-1.1)	-0.71% (-2.1-0.86%)	-0.95% (-2.7-0.84%)	-0.25 (-0.7-0.22)	-0.74 (-2.2-0.9)
Mauritius	1995	Male	All Ages	56% (49-62%)	44% (38-51%)	2.3 (1.9-2.7)	9.4% (7.3-12%)	11% (7.7-14%)	460 (320-600)	17000 (13000-22000)
Mauritius	1995	Male	15 to 19	45% (32-59%)	55% (41-68%)	1.1 (0.64-1.6)	3.2% (2-4.9%)	5.5% (3.2-8.6%)	2.5 (1.4-3.8)	220 (140-330)
Mauritius	1995	Male	20 to 24	64% (50-76%)	36% (24-50%)	2.3 (1.6-3.3)	7.8% (5.2-11%)	12% (7.5-18%)	7.8 (4.8-12)	640 (430-900)
Mauritius	1995	Male	25 to 29	69% (56-79%)	31% (21-44%)	2.9 (2-3.8)	10% (7.2-13%)	15% (10-20%)	12 (8.1-17)	930 (660-1200)
Mauritius	1995	Male	30 to 34	71% (58-81%)	29% (19-42%)	3.5 (2.5-4.6)	13% (9.4-16%)	17% (12-22%)	23 (16-30)	1500 (1100-1900)
Mauritius	1995	Male	35 to 39	72% (60-81%)	28% (19-40%)	4 (2.7-5.3)	15% (11-19%)	20% (14-25%)	35 (25-44)	2000 (1500-2500)
Mauritius	1995	Male	40 to 44	72% (61-82%)	28% (18-39%)	4.3 (3.1-5.6)	16% (12-20%)	20% (15-25%)	43 (32-54)	2200 (1600-2700)
Mauritius	1995	Male	45 to 49	72% (58-81%)	28% (19-42%)	4.4 (3.1-5.9)	16% (11-20%)	19% (13-25%)	50 (35-64)	2200 (1600-2800)
Mauritius	1995	Male	50 to 54	72% (60-81%)	28% (19-40%)	4.5 (3.1-6.1)	15% (10-19%)	18% (12-23%)	50 (33-66)	1900 (1300-2500)
Mauritius	1995	Male	55 to 59	72% (59-82%)	28% (18-41%)	4.4 (3-6.1)	14% (8.6-19%)	16% (9.8-22%)	56 (35-78)	1900 (1200-2500)
Mauritius	1995	Male	60 to 64	68% (53-80%)	32% (20-47%)	4 (2.6-5.7)	12% (7-17%)	14% (7.9-20%)	57 (32-83)	1600 (940-2300)
Mauritius	1995	Male	65 to 69	59% (44-72%)	41% (28-56%)	3.2 (2.1-4.6)	9.3% (4.9-14%)	10% (5.3-16%)	53 (27-83)	1200 (640-1900)
Mauritius	1995	Male	70 to 74	53% (39-65%)	47% (35-61%)	2.5 (1.6-3.9)	6.7% (2.7-11%)	7.4% (2.8-13%)	40 (15-68)	740 (300-1300)
Mauritius	1995	Male	75 to 79	50% (35-65%)	50% (35-65%)	2.1 (1.2-3.3)	4.5% (1.2-8.7%)	4.9% (1.1-9.8%)	20 (4.5-39)	290 (75-560)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	1995	Male	80 to 84	46% (30-62%)	54% (38-70%)	1.7 (1-2.6)	3.1% (0.7-6.3%)	3.3% (0.56-7.1%)	9.1 (1.5-19)	100 (23-210)
Mauritius	1995	Male	85 to 89	44% (29-59%)	56% (41-71%)	1.4 (0.81-2.3)	1.8% (-0.032-4.5%)	1.9% (-0.23-5%)	2.4 (-0.3-6.4)	19 (-0.35-49)
Mauritius	1995	Male	90 to 94	45% (31-60%)	55% (40-69%)	1.3 (0.7-2.2)	0.88% (-0.76-3.3%)	0.82% (-1.1-3.6%)	0.28 (-0.39-1.2)	1.8 (-1.6-6.7)
Mauritius	1995	Male	95 plus	47% (29-66%)	53% (34-71%)	1.2 (0.51-2.2)	0.48% (-1.2-3.3%)	0.26% (-1.7-3.4%)	0.014 (-0.092-0.18)	0.1 (-0.26-0.69)
Mauritius	2000	Female	All Ages	22% (19-26%)	78% (74-81%)	0.25 (0.19-0.33)	0.18% (-0.27-0.65%)	-0.4% (-1.3-0.66%)	-14 (-47-23)	270 (-390-1000)
Mauritius	2000	Female	15 to 19	18% (12-26%)	82% (74-88%)	0.21 (0.12-0.34)	0.67% (0.36-1.1%)	1.1% (0.24-2.3%)	0.24 (0.054-0.51)	39 (21-62)
Mauritius	2000	Female	20 to 24	21% (14-29%)	79% (71-86%)	0.25 (0.14-0.41)	1.1% (0.67-1.6%)	1.2% (0.25-2.4%)	0.36 (0.076-0.75)	77 (43-120)
Mauritius	2000	Female	25 to 29	26% (18-34%)	74% (66-82%)	0.29 (0.18-0.43)	1.1% (0.71-1.5%)	1.1% (0.4-2%)	0.39 (0.14-0.7)	71 (45-100)
Mauritius	2000	Female	30 to 34	28% (20-38%)	72% (62-80%)	0.32 (0.2-0.48)	1% (0.67-1.4%)	1.3% (0.66-2.1%)	0.57 (0.29-0.93)	78 (50-110)
Mauritius	2000	Female	35 to 39	27% (20-37%)	73% (63-80%)	0.33 (0.2-0.52)	0.93% (0.59-1.3%)	1.3% (0.62-2.1%)	0.87 (0.42-1.4)	84 (51-120)
Mauritius	2000	Female	40 to 44	27% (19-36%)	73% (64-81%)	0.33 (0.2-0.51)	0.92% (0.51-1.4%)	1.4% (0.65-2.3%)	1.3 (0.61-2.2)	91 (50-140)
Mauritius	2000	Female	45 to 49	30% (20-38%)	70% (62-80%)	0.37 (0.23-0.57)	0.64% (0.12-1.2%)	0.9% (0.02-2%)	1.2 (0.027-2.7)	68 (14-130)
Mauritius	2000	Female	50 to 54	32% (24-41%)	68% (59-76%)	0.45 (0.28-0.7)	0.17% (-0.58-0.99%)	0.15% (-1.1-1.4%)	0.24 (-1.7-2.3)	17 (-56-97)
Mauritius	2000	Female	55 to 59	29% (21-39%)	71% (61-79%)	0.5 (0.3-0.78)	-0.054% (-0.96-1%)	-0.13% (-1.4-1.4%)	-0.24 (-2.8-2.7)	-5.1 (-87-96)
Mauritius	2000	Female	60 to 64	27% (19-36%)	73% (64-81%)	0.49 (0.28-0.79)	-0.36% (-1.4-0.82%)	-0.51% (-1.8-1.1%)	-1.4 (-5.1-2.9)	-37 (-140-89)
Mauritius	2000	Female	65 to 69	26% (18-35%)	74% (65-82%)	0.46 (0.27-0.73)	-0.45% (-1.4-0.55%)	-0.61% (-1.9-0.7%)	-2 (-6.4-2.3)	-45 (-140-56)
Mauritius	2000	Female	70 to 74	26% (18-34%)	74% (66-82%)	0.46 (0.26-0.75)	-0.44% (-1.5-0.73%)	-0.58% (-1.9-0.91%)	-2.6 (-8.6-4.1)	-47 (-160-78)
Mauritius	2000	Female	75 to 79	26% (18-37%)	74% (63-82%)	0.47 (0.25-0.78)	-0.65% (-1.8-0.48%)	-0.85% (-2.3-0.61%)	-4.3 (-12-3)	-59 (-160-44)
Mauritius	2000	Female	80 to 84	27% (18-38%)	73% (62-82%)	0.47 (0.27-0.73)	-0.71% (-1.8-0.34%)	-0.91% (-2.2-0.4%)	-4.1 (-9.8-1.8)	-42 (-100-21)
Mauritius	2000	Female	85 to 89	28% (19-39%)	72% (61-81%)	0.47 (0.26-0.76)	-0.7% (-1.7-0.34%)	-0.91% (-2.1-0.36%)	-2.9 (-6.8-1.2)	-20 (-49-9.8)
Mauritius	2000	Female	90 to 94	30% (19-41%)	70% (59-81%)	0.48 (0.26-0.83)	-0.67% (-1.7-0.38%)	-0.88% (-2.1-0.39%)	-1.3 (-3.1-0.57)	-6 (-15-3.5)
Mauritius	2000	Female	95 plus	32% (19-48%)	68% (52-81%)	0.51 (0.23-0.98)	-0.68% (-1.8-0.63%)	-0.91% (-2.3-0.59%)	-0.37 (-0.95-0.24)	-1.1 (-3-1.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	2000	Male	All Ages	59% (54-64%)	41% (36-46%)	2.5 (2.1-2.9)	9% (6.7-11%)	10% (6.9-14%)	450 (300-610)	17000 (13000-21000)
Mauritius	2000	Male	15 to 19	48% (38-59%)	52% (41-62%)	1.1 (0.71-1.6)	3% (1.9-4.3%)	5.6% (3.1-8.4%)	1.9 (1.1-2.9)	180 (120-260)
Mauritius	2000	Male	20 to 24	66% (55-74%)	34% (26-45%)	2.3 (1.7-3.1)	7.5% (5-10%)	12% (7.4-17%)	7.3 (4.6-10)	610 (400-820)
Mauritius	2000	Male	25 to 29	71% (62-78%)	29% (22-38%)	2.8 (2.1-3.7)	9.4% (6.6-13%)	14% (9.2-19%)	10 (6.5-14)	760 (540-1000)
Mauritius	2000	Male	30 to 34	75% (66-81%)	25% (19-34%)	3.6 (2.7-4.6)	11% (8.1-14%)	16% (11-21%)	16 (11-22)	1100 (800-1400)
Mauritius	2000	Male	35 to 39	75% (68-82%)	25% (18-32%)	4 (3-5.2)	13% (9.7-17%)	18% (13-23%)	31 (22-40)	1800 (1300-2300)
Mauritius	2000	Male	40 to 44	76% (69-82%)	24% (18-31%)	4.4 (3.2-5.6)	14% (10-18%)	18% (13-23%)	44 (31-57)	2200 (1600-2900)
Mauritius	2000	Male	45 to 49	76% (67-82%)	24% (18-33%)	4.6 (3.4-5.9)	14% (10-18%)	18% (12-23%)	54 (36-71)	2400 (1700-3200)
Mauritius	2000	Male	50 to 54	74% (66-81%)	26% (19-34%)	4.5 (3.3-5.9)	13% (8.8-18%)	16% (10-22%)	52 (34-71)	2000 (1400-2800)
Mauritius	2000	Male	55 to 59	75% (65-83%)	25% (17-35%)	4.5 (3.2-5.9)	13% (7.5-17%)	15% (8.5-21%)	49 (28-69)	1700 (990-2300)
Mauritius	2000	Male	60 to 64	74% (64-82%)	26% (18-36%)	4.4 (3.1-6)	12% (6.7-17%)	14% (7.6-20%)	59 (32-86)	1700 (930-2400)
Mauritius	2000	Male	65 to 69	65% (53-75%)	35% (25-47%)	3.5 (2.5-4.8)	9.4% (5-14%)	11% (5.6-17%)	52 (27-81)	1200 (650-1900)
Mauritius	2000	Male	70 to 74	53% (40-65%)	47% (35-60%)	2.4 (1.6-3.6)	6.1% (2.4-10%)	6.9% (2.7-12%)	40 (15-68)	750 (300-1300)
Mauritius	2000	Male	75 to 79	50% (37-63%)	50% (37-63%)	2 (1.2-3)	4.1% (0.97-8%)	4.5% (0.9-9%)	23 (4.8-47)	350 (85-690)
Mauritius	2000	Male	80 to 84	48% (31-63%)	52% (37-69%)	1.7 (1-2.6)	2.9% (0.57-6%)	3.1% (0.47-6.7%)	11 (1.7-24)	130 (25-260)
Mauritius	2000	Male	85 to 89	44% (29-59%)	56% (41-71%)	1.4 (0.78-2.2)	1.6% (-0.13-3.9%)	1.7% (-0.36-4.4%)	2.6 (-0.55-6.8)	22 (-1.7-54)
Mauritius	2000	Male	90 to 94	44% (30-59%)	56% (41-70%)	1.2 (0.66-2)	0.68% (-0.8-3%)	0.58% (-1.2-3.2%)	0.29 (-0.58-1.6)	2.1 (-2.4-8.9)
Mauritius	2000	Male	95 plus	47% (28-65%)	53% (35-72%)	1.1 (0.48-2)	0.27% (-1.3-2.7%)	0.011% (-1.9-2.8%)	0.001 (-0.17-0.26)	0.099 (-0.46-0.99)
Mauritius	2005	Female	All Ages	24% (21-27%)	76% (73-79%)	0.26 (0.2-0.32)	0.024% (-0.43-0.46%)	-0.69% (-1.6-0.27%)	-26 (-58-10)	38 (-620-720)
Mauritius	2005	Female	15 to 19	19% (12-27%)	81% (73-88%)	0.2 (0.12-0.34)	0.66% (0.38-1%)	1.1% (0.35-2.2%)	0.2 (0.067-0.42)	35 (20-55)
Mauritius	2005	Female	20 to 24	22% (15-30%)	78% (70-85%)	0.24 (0.14-0.4)	1.1% (0.67-1.5%)	1.1% (0.33-2.2%)	0.28 (0.086-0.56)	64 (39-98)
Mauritius	2005	Female	25 to 29	27% (19-35%)	73% (65-81%)	0.28 (0.18-0.42)	1.1% (0.71-1.4%)	1.1% (0.44-1.8%)	0.31 (0.13-0.53)	71 (46-100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	2005	Female	30 to 34	29% (22-39%)	71% (61-78%)	0.31 (0.2-0.47)	0.98% (0.64-1.3%)	1.3% (0.69-1.9%)	0.49 (0.26-0.74)	67 (43-96)
Mauritius	2005	Female	35 to 39	29% (21-38%)	71% (62-79%)	0.32 (0.19-0.5)	0.87% (0.53-1.2%)	1.2% (0.61-2%)	0.78 (0.39-1.3)	74 (45-110)
Mauritius	2005	Female	40 to 44	28% (21-37%)	72% (63-79%)	0.32 (0.19-0.49)	0.9% (0.5-1.3%)	1.5% (0.82-2.3%)	1.5 (0.79-2.3)	96 (51-140)
Mauritius	2005	Female	45 to 49	31% (22-40%)	69% (60-78%)	0.36 (0.22-0.56)	0.59% (0.078-1.1%)	0.95% (0.097-1.9%)	1.3 (0.14-2.7)	69 (9.3-130)
Mauritius	2005	Female	50 to 54	33% (24-42%)	67% (58-76%)	0.44 (0.28-0.67)	0.085% (-0.58-0.76%)	0.13% (-0.9-1.2%)	0.26 (-1.7-2.4)	11 (-71-97)
Mauritius	2005	Female	55 to 59	31% (23-40%)	69% (60-77%)	0.49 (0.29-0.76)	-0.27% (-1.1-0.75%)	-0.38% (-1.6-1%)	-0.92 (-3.8-2.6)	-34 (-140-95)
Mauritius	2005	Female	60 to 64	28% (20-37%)	72% (63-80%)	0.48 (0.28-0.75)	-0.63% (-1.6-0.45%)	-0.85% (-2.1-0.63%)	-2.4 (-6-1.8)	-70 (-180-51)
Mauritius	2005	Female	65 to 69	27% (18-36%)	73% (64-82%)	0.45 (0.27-0.68)	-0.69% (-1.6-0.21%)	-0.9% (-2.1-0.25%)	-3.1 (-7.3-0.89)	-74 (-170-23)
Mauritius	2005	Female	70 to 74	27% (19-36%)	73% (64-81%)	0.45 (0.26-0.7)	-0.74% (-1.7-0.22%)	-0.96% (-2.2-0.29%)	-3.8 (-8.8-1.1)	-72 (-170-22)
Mauritius	2005	Female	75 to 79	28% (19-39%)	72% (61-81%)	0.46 (0.25-0.74)	-0.91% (-2-0.14%)	-1.2% (-2.6-0.16%)	-5.9 (-13-0.82)	-83 (-180-12)
Mauritius	2005	Female	80 to 84	29% (19-39%)	71% (61-81%)	0.46 (0.26-0.73)	-1% (-2.1-0.011%)	-1.3% (-2.6--0.017%)	-7 (-14--0.095)	-72 (-150-0.84)
Mauritius	2005	Female	85 to 89	30% (19-41%)	70% (59-81%)	0.46 (0.26-0.75)	-0.97% (-2.1-0.032%)	-1.2% (-2.6--0.015%)	-5.1 (-11--0.061)	-36 (-78-1.2)
Mauritius	2005	Female	90 to 94	31% (21-42%)	69% (58-79%)	0.46 (0.26-0.79)	-0.87% (-1.8-0.12%)	-1.1% (-2.4-0.051%)	-2.2 (-4.5-0.1)	-10 (-22-1.5)
Mauritius	2005	Female	95 plus	33% (20-49%)	67% (51-80%)	0.48 (0.23-0.9)	-0.81% (-2-0.48%)	-1.1% (-2.5-0.43%)	-0.63 (-1.5-0.26)	-1.9 (-4.6-1.2)
Mauritius	2005	Male	All Ages	61% (56-66%)	39% (34-44%)	2.4 (1.9-2.8)	8.2% (6-10%)	8.8% (5.9-12%)	420 (270-560)	16000 (12000-20000)
Mauritius	2005	Male	15 to 19	49% (38-60%)	51% (40-62%)	1 (0.66-1.5)	2.8% (1.8-4.1%)	5.2% (3-7.9%)	1.6 (0.94-2.5)	160 (100-230)
Mauritius	2005	Male	20 to 24	66% (56-75%)	34% (25-44%)	2.1 (1.5-2.9)	7% (4.7-9.5%)	11% (6.9-16%)	5.8 (3.6-8.4)	490 (330-670)
Mauritius	2005	Male	25 to 29	71% (63-78%)	29% (22-37%)	2.6 (1.9-3.5)	8.6% (6-11%)	13% (8.1-17%)	10 (6.5-14)	780 (530-1000)
Mauritius	2005	Male	30 to 34	75% (67-81%)	25% (19-33%)	3.3 (2.5-4.3)	10% (7.4-13%)	14% (10-19%)	13 (8.9-18)	910 (660-1200)
Mauritius	2005	Male	35 to 39	76% (69-82%)	24% (18-31%)	3.7 (2.7-4.8)	12% (8.7-15%)	16% (11-21%)	25 (18-33)	1500 (1100-1900)
Mauritius	2005	Male	40 to 44	76% (69-83%)	24% (17-31%)	4 (3-5.2)	13% (9.4-16%)	17% (12-21%)	40 (28-52)	2100 (1500-2700)
Mauritius	2005	Male	45 to 49	76% (68-83%)	24% (17-32%)	4.2 (3.1-5.5)	13% (9.3-17%)	17% (11-21%)	52 (36-68)	2400 (1700-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	2005	Male	50 to 54	75% (66-82%)	25% (18-34%)	4.2 (3-5.4)	12% (7.7-16%)	14% (9.2-20%)	57 (37-78)	2300 (1500-3100)
Mauritius	2005	Male	55 to 59	75% (65-83%)	25% (17-35%)	4.1 (3-5.4)	11% (6.6-15%)	13% (7.6-18%)	56 (33-79)	1900 (1100-2600)
Mauritius	2005	Male	60 to 64	74% (64-82%)	26% (18-36%)	4.1 (2.9-5.6)	10% (5.8-15%)	12% (6.6-18%)	50 (28-74)	1500 (820-2100)
Mauritius	2005	Male	65 to 69	66% (55-75%)	34% (25-45%)	3.3 (2.3-4.5)	7.9% (4-12%)	9% (4.5-14%)	42 (21-67)	1000 (520-1600)
Mauritius	2005	Male	70 to 74	54% (41-65%)	46% (35-59%)	2.3 (1.4-3.3)	5% (1.9-8.7%)	5.6% (2-10%)	29 (10-51)	550 (210-970)
Mauritius	2005	Male	75 to 79	51% (37-64%)	49% (36-63%)	1.9 (1.1-2.8)	3.3% (0.65-6.7%)	3.7% (0.64-7.6%)	19 (3.3-40)	290 (57-590)
Mauritius	2005	Male	80 to 84	48% (32-63%)	52% (37-68%)	1.6 (0.94-2.3)	2.2% (0.22-4.9%)	2.4% (0.027-5.5%)	11 (0.13-26)	130 (12-290)
Mauritius	2005	Male	85 to 89	44% (29-59%)	56% (41-71%)	1.3 (0.72-2.1)	1.2% (-0.26-3.2%)	1.2% (-0.51-3.5%)	2.9 (-1.2-8.5)	25 (-5.4-68)
Mauritius	2005	Male	90 to 94	45% (30-60%)	55% (40-70%)	1.1 (0.62-1.8)	0.42% (-0.89-2.2%)	0.29% (-1.3-2.3%)	0.2 (-0.9-1.6)	1.8 (-3.8-9.4)
Mauritius	2005	Male	95 plus	47% (29-67%)	53% (33-71%)	1 (0.47-1.9)	0.17% (-1.2-2.1%)	-0.096% (-1.7-2.1%)	-0.014 (-0.25-0.32)	0.099 (-0.69-1.3)
Mauritius	2010	Female	All Ages	28% (24-32%)	72% (68-76%)	0.31 (0.23-0.41)	-0.13% (-0.62-0.42%)	-0.94% (-1.9-0.13%)	-38 (-75-5.2)	-210 (-990-670)
Mauritius	2010	Female	15 to 19	21% (14-30%)	79% (70-86%)	0.24 (0.13-0.39)	0.69% (0.41-1%)	1.2% (0.49-2.3%)	0.21 (0.082-0.39)	38 (22-58)
Mauritius	2010	Female	20 to 24	27% (18-37%)	73% (63-82%)	0.3 (0.17-0.5)	1.1% (0.71-1.6%)	1.2% (0.36-2.4%)	0.28 (0.086-0.56)	64 (39-94)
Mauritius	2010	Female	25 to 29	31% (21-42%)	69% (58-79%)	0.32 (0.19-0.52)	1.1% (0.7-1.5%)	1.1% (0.4-2%)	0.31 (0.12-0.56)	63 (39-92)
Mauritius	2010	Female	30 to 34	33% (22-45%)	67% (55-78%)	0.35 (0.2-0.55)	0.97% (0.59-1.4%)	1.3% (0.67-2%)	0.55 (0.29-0.87)	74 (45-110)
Mauritius	2010	Female	35 to 39	33% (22-44%)	67% (56-78%)	0.37 (0.21-0.59)	0.85% (0.49-1.3%)	1.3% (0.59-2.1%)	0.75 (0.34-1.2)	68 (38-100)
Mauritius	2010	Female	40 to 44	33% (23-44%)	67% (56-77%)	0.39 (0.21-0.61)	0.86% (0.41-1.4%)	1.5% (0.73-2.4%)	1.4 (0.66-2.2)	83 (40-130)
Mauritius	2010	Female	45 to 49	35% (23-46%)	65% (54-77%)	0.41 (0.23-0.67)	0.51% (-0.067-1.1%)	0.92% (0.013-2%)	1.4 (0.021-3.1)	67 (-9.1-150)
Mauritius	2010	Female	50 to 54	35% (24-46%)	65% (54-76%)	0.47 (0.28-0.74)	-0.023% (-0.73-0.7%)	0.037% (-1-1.1%)	0.073 (-2-2.3)	-3.3 (-98-94)
Mauritius	2010	Female	55 to 59	33% (23-44%)	67% (56-77%)	0.53 (0.31-0.86)	-0.48% (-1.4-0.55%)	-0.63% (-2-0.87%)	-1.7 (-5.6-2.4)	-69 (-210-82)
Mauritius	2010	Female	60 to 64	32% (22-43%)	68% (57-78%)	0.54 (0.3-0.88)	-0.83% (-1.9-0.38%)	-1.1% (-2.6-0.46%)	-3.8 (-9-1.6)	-120 (-280-53)
Mauritius	2010	Female	65 to 69	31% (21-41%)	69% (59-79%)	0.52 (0.3-0.82)	-0.96% (-2-0.12%)	-1.3% (-2.7-0.15%)	-4.5 (-9.4-0.55)	-110 (-240-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	2010	Female	70 to 74	30% (21-41%)	70% (59-79%)	0.51 (0.28-0.82)	-1% (-2.2-0.095%)	-1.4% (-2.9-0.038%)	-5.7 (-12-0.15)	-110 (-230-9.5)
Mauritius	2010	Female	75 to 79	30% (20-42%)	70% (58-80%)	0.51 (0.27-0.86)	-1.1% (-2.3-0.038%)	-1.5% (-3.1-0.017%)	-7.1 (-15-0.079)	-100 (-210-3.2)
Mauritius	2010	Female	80 to 84	31% (20-42%)	69% (58-80%)	0.5 (0.28-0.8)	-1.2% (-2.2--0.096%)	-1.5% (-2.8--0.21%)	-8.3 (-16--1.2)	-87 (-170--7.3)
Mauritius	2010	Female	85 to 89	32% (21-43%)	68% (57-79%)	0.49 (0.26-0.8)	-1.1% (-2.2--0.084%)	-1.4% (-2.8--0.17%)	-7.3 (-14--0.85)	-52 (-110--3.9)
Mauritius	2010	Female	90 to 94	33% (23-45%)	67% (55-77%)	0.5 (0.26-0.87)	-0.94% (-2-0.15%)	-1.2% (-2.5-0.14%)	-3.3 (-6.8-0.37)	-16 (-33-2.6)
Mauritius	2010	Female	95 plus	36% (22-51%)	64% (49-78%)	0.53 (0.23-0.97)	-0.81% (-2-0.58%)	-1.1% (-2.6-0.55%)	-0.89 (-2.1-0.45)	-2.7 (-6.7-1.9)
Mauritius	2010	Male	All Ages	58% (51-63%)	42% (37-49%)	2.2 (1.8-2.8)	7.8% (5.6-10%)	8.1% (5.3-11%)	420 (270-580)	16000 (11000-21000)
Mauritius	2010	Male	15 to 19	44% (31-58%)	56% (42-69%)	0.9 (0.51-1.4)	2.7% (1.7-4.1%)	4.7% (2.6-7.5%)	1.8 (1-2.9)	170 (110-260)
Mauritius	2010	Male	20 to 24	64% (49-76%)	36% (24-51%)	2 (1.2-2.9)	6.7% (4.4-9.5%)	10% (6.1-15%)	5.6 (3.3-8.1)	470 (310-670)
Mauritius	2010	Male	25 to 29	68% (55-79%)	32% (21-45%)	2.5 (1.6-3.4)	8.3% (5.7-11%)	12% (7.3-16%)	9.1 (5.8-13)	690 (460-940)
Mauritius	2010	Male	30 to 34	71% (58-80%)	29% (20-42%)	3 (2.1-4)	9.3% (6.6-12%)	13% (8.5-17%)	14 (9.8-20)	1000 (710-1300)
Mauritius	2010	Male	35 to 39	72% (59-81%)	28% (19-41%)	3.4 (2.3-4.7)	11% (8-14%)	14% (10-19%)	21 (15-29)	1300 (930-1700)
Mauritius	2010	Male	40 to 44	72% (61-81%)	28% (19-39%)	3.7 (2.5-5)	12% (8.3-15%)	15% (11-20%)	33 (23-44)	1800 (1300-2300)
Mauritius	2010	Male	45 to 49	71% (58-81%)	29% (19-42%)	3.8 (2.6-5.2)	12% (8.1-15%)	15% (10-20%)	52 (34-69)	2400 (1600-3100)
Mauritius	2010	Male	50 to 54	71% (59-81%)	29% (19-41%)	3.8 (2.6-5.2)	11% (6.8-15%)	13% (8.3-18%)	61 (37-83)	2400 (1500-3300)
Mauritius	2010	Male	55 to 59	72% (58-82%)	28% (18-42%)	3.8 (2.5-5.2)	10% (5.9-14%)	12% (6.9-17%)	62 (36-89)	2200 (1300-3100)
Mauritius	2010	Male	60 to 64	67% (52-79%)	33% (21-48%)	3.4 (2.2-5)	8.7% (5-13%)	10% (5.7-15%)	58 (32-87)	1700 (960-2500)
Mauritius	2010	Male	65 to 69	58% (43-71%)	42% (29-57%)	2.8 (1.8-4)	6.6% (3.3-10%)	7.7% (3.7-12%)	39 (19-62)	950 (480-1500)
Mauritius	2010	Male	70 to 74	52% (38-65%)	48% (35-62%)	2.2 (1.3-3.3)	4.6% (1.8-8.3%)	5.3% (2-9.6%)	28 (10-49)	550 (210-960)
Mauritius	2010	Male	75 to 79	49% (34-64%)	51% (36-66%)	1.8 (1-2.8)	3.1% (0.69-6.4%)	3.5% (0.62-7.3%)	18 (3.2-37)	270 (60-550)
Mauritius	2010	Male	80 to 84	45% (30-61%)	55% (39-70%)	1.5 (0.83-2.3)	2.2% (0.25-4.8%)	2.3% (0.12-5.5%)	11 (0.56-25)	120 (14-280)
Mauritius	2010	Male	85 to 89	43% (29-58%)	57% (42-71%)	1.2 (0.67-2)	1.2% (-0.14-3.3%)	1.3% (-0.35-3.7%)	4.1 (-1.1-12)	35 (-4-92)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	2010	Male	90 to 94	43% (29-59%)	57% (41-71%)	1.1 (0.57-1.8)	0.47% (-0.77-2.2%)	0.36% (-1.1-2.4%)	0.43 (-1.3-2.9)	3.4 (-5.5-16)
Mauritius	2010	Male	95 plus	46% (27-65%)	54% (35-73%)	0.95 (0.42-1.8)	0.26% (-0.97-2.1%)	0.017% (-1.5-2%)	0.0038 (-0.33-0.44)	0.23 (-0.86-1.8)
Mauritius	2016	Female	All Ages	28% (24-32%)	72% (68-76%)	0.33 (0.2-0.49)	-0.24% (-0.8-0.49%)	-1% (-2.1-0.3%)	-48 (-96-15)	-410 (-1400-790)
Mauritius	2016	Female	15 to 19	21% (14-30%)	79% (70-86%)	0.23 (0.12-0.42)	0.69% (0.4-1%)	1.2% (0.4-2.3%)	0.19 (0.064-0.38)	35 (19-54)
Mauritius	2016	Female	20 to 24	26% (17-36%)	74% (64-83%)	0.29 (0.15-0.52)	1.1% (0.71-1.6%)	1.1% (0.35-2.4%)	0.27 (0.077-0.56)	68 (40-100)
Mauritius	2016	Female	25 to 29	31% (21-41%)	69% (59-79%)	0.32 (0.17-0.55)	1% (0.66-1.5%)	0.93% (0.28-1.8%)	0.24 (0.074-0.48)	62 (37-92)
Mauritius	2016	Female	30 to 34	32% (22-44%)	68% (56-78%)	0.34 (0.18-0.59)	0.93% (0.56-1.4%)	1.1% (0.47-1.9%)	0.36 (0.15-0.65)	60 (34-91)
Mauritius	2016	Female	35 to 39	32% (21-43%)	68% (57-79%)	0.37 (0.19-0.63)	0.8% (0.43-1.3%)	1.1% (0.38-2.1%)	0.63 (0.23-1.2)	69 (34-110)
Mauritius	2016	Female	40 to 44	32% (22-43%)	68% (57-78%)	0.38 (0.19-0.65)	0.75% (0.29-1.3%)	1.2% (0.43-2.3%)	0.98 (0.32-1.8)	69 (27-120)
Mauritius	2016	Female	45 to 49	34% (23-45%)	66% (55-77%)	0.41 (0.21-0.71)	0.48% (-0.091-1.1%)	0.81% (-0.14-1.9%)	0.97 (-0.17-2.3)	51 (-11-120)
Mauritius	2016	Female	50 to 54	34% (24-46%)	66% (54-76%)	0.47 (0.25-0.8)	-0.043% (-0.79-0.81%)	-0.052% (-1.2-1.3%)	-0.11 (-2.6-2.8)	-6.6 (-110-120)
Mauritius	2016	Female	55 to 59	33% (23-45%)	67% (55-77%)	0.53 (0.28-0.94)	-0.52% (-1.6-0.69%)	-0.73% (-2.2-1%)	-2.2 (-6.7-2.9)	-84 (-250-100)
Mauritius	2016	Female	60 to 64	31% (21-43%)	69% (57-79%)	0.54 (0.27-0.93)	-0.85% (-2.1-0.52%)	-1.2% (-2.8-0.74%)	-4.6 (-11-2.7)	-140 (-350-85)
Mauritius	2016	Female	65 to 69	30% (20-40%)	70% (60-80%)	0.51 (0.26-0.84)	-0.97% (-2.1-0.3%)	-1.3% (-2.9-0.39%)	-6.4 (-14-1.9)	-160 (-360-49)
Mauritius	2016	Female	70 to 74	30% (20-40%)	70% (60-80%)	0.5 (0.25-0.85)	-1% (-2.2-0.37%)	-1.3% (-2.9-0.39%)	-6.6 (-15-2)	-130 (-290-43)
Mauritius	2016	Female	75 to 79	30% (19-42%)	70% (58-81%)	0.51 (0.24-0.9)	-1.1% (-2.3-0.29%)	-1.5% (-3.1-0.33%)	-7.6 (-16-1.5)	-110 (-240-28)
Mauritius	2016	Female	80 to 84	30% (20-42%)	70% (58-80%)	0.5 (0.25-0.84)	-1.1% (-2.2-0.096%)	-1.5% (-2.9-0.036%)	-9.5 (-19-0.24)	-100 (-200-8)
Mauritius	2016	Female	85 to 89	31% (21-42%)	69% (58-79%)	0.49 (0.24-0.84)	-1.1% (-2.1-0.11%)	-1.4% (-2.8-0.062%)	-8.3 (-16-0.37)	-59 (-120-6.2)
Mauritius	2016	Female	90 to 94	33% (22-45%)	67% (55-78%)	0.51 (0.24-0.93)	-0.93% (-1.9-0.31%)	-1.2% (-2.5-0.27%)	-4.5 (-9-0.91)	-21 (-44-6.6)
Mauritius	2016	Female	95 plus	36% (21-53%)	64% (47-79%)	0.54 (0.22-1)	-0.79% (-2-0.67%)	-1.1% (-2.6-0.68%)	-1.4 (-3.4-0.83)	-4.2 (-11-3.3)
Mauritius	2016	Male	All Ages	58% (52-64%)	42% (36-48%)	2.3 (1.6-3.3)	7.3% (4.4-11%)	7.5% (3.9-12%)	420 (210-660)	15000 (9100-22000)
Mauritius	2016	Male	15 to 19	43% (30-57%)	57% (43-70%)	0.88 (0.46-1.5)	2.6% (1.5-4%)	4.6% (2.5-7.6%)	1.5 (0.76-2.5)	150 (87-230)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritius	2016	Male	20 to 24	63% (49-75%)	37% (25-51%)	2 (1.1-3.1)	6.5% (4-9.4%)	10% (5.4-16%)	5.4 (2.8-8.7)	470 (280-710)
Mauritius	2016	Male	25 to 29	68% (55-78%)	32% (22-45%)	2.4 (1.4-3.7)	8% (5.1-11%)	12% (6.8-17%)	8 (4.4-13)	630 (400-930)
Mauritius	2016	Male	30 to 34	70% (56-80%)	30% (20-44%)	3 (1.8-4.6)	9% (5.9-13%)	13% (8-18%)	11 (6.4-17)	770 (490-1100)
Mauritius	2016	Male	35 to 39	72% (59-80%)	28% (20-41%)	3.4 (2-5.2)	10% (6.9-14%)	14% (9-20%)	20 (12-29)	1200 (810-1800)
Mauritius	2016	Male	40 to 44	72% (61-81%)	28% (19-39%)	3.7 (2.2-5.6)	11% (7.3-15%)	15% (9.4-21%)	27 (16-38)	1500 (950-2100)
Mauritius	2016	Male	45 to 49	71% (57-81%)	29% (19-43%)	3.8 (2.2-5.7)	11% (7-16%)	15% (9.1-21%)	38 (22-54)	1800 (1100-2500)
Mauritius	2016	Male	50 to 54	71% (59-80%)	29% (20-41%)	3.8 (2.3-5.8)	11% (5.9-15%)	13% (7.3-19%)	55 (30-85)	2300 (1200-3400)
Mauritius	2016	Male	55 to 59	71% (57-82%)	29% (18-43%)	3.7 (2.3-5.8)	9.5% (5-14%)	12% (6-17%)	62 (32-94)	2200 (1200-3300)
Mauritius	2016	Male	60 to 64	66% (50-79%)	34% (21-50%)	3.4 (2-5.3)	8.4% (4-13%)	10% (4.7-16%)	64 (30-100)	1900 (930-3000)
Mauritius	2016	Male	65 to 69	57% (42-71%)	43% (29-58%)	2.7 (1.6-4.3)	6.4% (2.7-11%)	7.5% (3.1-13%)	54 (22-94)	1300 (560-2300)
Mauritius	2016	Male	70 to 74	51% (38-64%)	49% (36-62%)	2.1 (1.2-3.5)	4.6% (1.4-8.6%)	5.3% (1.5-10%)	33 (9.4-63)	650 (200-1200)
Mauritius	2016	Male	75 to 79	48% (33-64%)	52% (36-67%)	1.8 (0.88-3)	3.1% (0.49-7%)	3.5% (0.4-8%)	20 (2.1-46)	310 (41-700)
Mauritius	2016	Male	80 to 84	45% (30-61%)	55% (39-70%)	1.4 (0.72-2.4)	2.1% (0.078-5.3%)	2.3% (-0.057-6.1%)	12 (-0.3-32)	140 (5.3-350)
Mauritius	2016	Male	85 to 89	42% (28-57%)	58% (43-72%)	1.2 (0.6-2)	1.2% (-0.32-3.7%)	1.3% (-0.53-4.1%)	4.7 (-1.9-15)	41 (-10-120)
Mauritius	2016	Male	90 to 94	42% (28-58%)	58% (42-72%)	1 (0.54-1.9)	0.53% (-0.81-2.6%)	0.42% (-1.2-2.8%)	0.71 (-2-4.7)	5.4 (-8.3-26)
Mauritius	2016	Male	95 plus	45% (28-64%)	55% (36-72%)	0.95 (0.38-1.9)	0.35% (-0.91-2.3%)	0.12% (-1.4-2.3%)	0.052 (-0.6-0.99)	0.61 (-1.6-4)
Mozambique	1990	Female	All Ages	35% (30-42%)	65% (58-70%)	0.25 (0.19-0.33)	0.19% (-0.024-0.4%)	0.18% (-0.23-0.61%)	180 (-220-580)	13000 (-1600-27000)
Mozambique	1990	Female	15 to 19	38% (26-52%)	62% (48-74%)	0.32 (0.18-0.52)	0.63% (-0.057-1.4%)	0.56% (-0.5-1.7%)	9.4 (-8.9-29)	1200 (-100-2600)
Mozambique	1990	Female	20 to 24	46% (32-60%)	54% (40-68%)	0.44 (0.25-0.71)	0.97% (0.1-1.9%)	0.68% (-0.52-1.9%)	13 (-9.7-37)	1800 (180-3500)
Mozambique	1990	Female	25 to 29	46% (33-59%)	54% (41-67%)	0.5 (0.29-0.81)	1.4% (0.22-2.6%)	1.2% (-0.43-3%)	22 (-7.5-53)	2100 (290-4100)
Mozambique	1990	Female	30 to 34	49% (35-61%)	51% (39-65%)	0.53 (0.29-0.88)	1.4% (0.11-2.8%)	1.4% (-0.47-3.5%)	24 (-8.3-61)	1900 (140-4000)
Mozambique	1990	Female	35 to 39	54% (40-65%)	46% (35-60%)	0.53 (0.29-0.87)	1.3% (-0.23-2.6%)	1.4% (-0.84-3.4%)	25 (-15-64)	1700 (-320-3700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	1990	Female	40 to 44	56% (43-67%)	44% (33-57%)	0.53 (0.29-0.84)	0.99% (-0.11-2.2%)	1.1% (-0.46-2.9%)	22 (-9.1-56)	1300 (-140-2900)
Mozambique	1990	Female	45 to 49	54% (40-66%)	46% (34-60%)	0.52 (0.27-0.86)	0.87% (-0.17-2%)	1.1% (-0.45-2.8%)	23 (-9.7-58)	1100 (-210-2500)
Mozambique	1990	Female	50 to 54	50% (38-62%)	50% (38-62%)	0.49 (0.25-0.81)	0.88% (-0.24-2.1%)	1.1% (-0.41-2.8%)	28 (-10-71)	1100 (-290-2600)
Mozambique	1990	Female	55 to 59	46% (33-58%)	54% (42-67%)	0.47 (0.24-0.8)	0.57% (-0.56-1.8%)	0.72% (-0.73-2.2%)	22 (-22-68)	700 (-700-2100)
Mozambique	1990	Female	60 to 64	46% (32-60%)	54% (40-68%)	0.46 (0.23-0.8)	0.39% (-0.59-1.5%)	0.51% (-0.74-1.9%)	19 (-27-70)	490 (-730-1900)
Mozambique	1990	Female	65 to 69	48% (34-63%)	52% (37-66%)	0.44 (0.24-0.74)	0.27% (-0.78-1.4%)	0.32% (-0.92-1.6%)	14 (-39-69)	310 (-880-1500)
Mozambique	1990	Female	70 to 74	47% (33-64%)	53% (36-67%)	0.38 (0.2-0.65)	-0.15% (-1.2-0.84%)	-0.2% (-1.4-0.99%)	-9.2 (-67-44)	-140 (-1200-810)
Mozambique	1990	Female	75 to 79	45% (30-62%)	55% (38-70%)	0.32 (0.15-0.57)	-0.18% (-1.2-0.6%)	-0.21% (-1.4-0.75%)	-7.8 (-54-28)	-110 (-750-380)
Mozambique	1990	Female	80 to 84	43% (27-61%)	57% (39-73%)	0.3 (0.15-0.51)	-0.4% (-1.3-0.25%)	-0.46% (-1.5-0.31%)	-10 (-35-7.1)	-110 (-370-68)
Mozambique	1990	Female	85 to 89	43% (29-58%)	57% (42-71%)	0.28 (0.14-0.5)	-0.54% (-1.4-0.07%)	-0.64% (-1.7-0.077%)	-9.1 (-24-1.1)	-66 (-170-8.5)
Mozambique	1990	Female	90 to 94	45% (31-60%)	55% (40-69%)	0.28 (0.14-0.51)	-0.63% (-1.5--0.027%)	-0.79% (-1.8--0.064%)	-4.5 (-10--0.37)	-22 (-52--0.94)
Mozambique	1990	Female	95 plus	47% (30-66%)	53% (34-70%)	0.28 (0.13-0.55)	-0.32% (-1.2-0.34%)	-0.46% (-1.5-0.33%)	-0.72 (-2.3-0.52)	-2 (-7.4-2.1)
Mozambique	1990	Male	All Ages	45% (37-52%)	55% (48-63%)	0.29 (0.22-0.39)	0.49% (0.12-0.83%)	0.65% (0.017-1.2%)	680 (18-1300)	38000 (9400-64000)
Mozambique	1990	Male	15 to 19	46% (32-61%)	54% (39-68%)	0.38 (0.22-0.6)	1.5% (0.5-2.7%)	1.4% (-0.11-3%)	26 (-2.1-57)	3000 (930-5300)
Mozambique	1990	Male	20 to 24	61% (46-73%)	39% (27-54%)	0.53 (0.31-0.83)	2.6% (0.62-4.6%)	2% (-0.46-4.6%)	44 (-11-100)	5100 (1200-8900)
Mozambique	1990	Male	25 to 29	62% (49-72%)	38% (28-51%)	0.61 (0.36-0.93)	2.8% (0.66-5.2%)	2.3% (-0.55-5.4%)	48 (-11-110)	4600 (1000-8300)
Mozambique	1990	Male	30 to 34	63% (50-73%)	37% (27-50%)	0.67 (0.39-1)	2.9% (0.26-5.4%)	2.6% (-0.67-5.9%)	51 (-13-110)	4000 (380-7600)
Mozambique	1990	Male	35 to 39	66% (54-75%)	34% (25-46%)	0.72 (0.42-1.1)	2.9% (0.25-5.9%)	2.9% (-0.54-6.7%)	60 (-10-140)	3900 (340-7700)
Mozambique	1990	Male	40 to 44	68% (57-77%)	32% (23-43%)	0.75 (0.43-1.2)	2.9% (0.24-5.5%)	3.1% (-0.3-6.4%)	71 (-7.6-150)	3900 (350-7300)
Mozambique	1990	Male	45 to 49	70% (59-79%)	30% (21-41%)	0.76 (0.42-1.2)	2.8% (0.5-5.3%)	3.1% (0.13-6.2%)	79 (3-160)	3700 (600-7100)
Mozambique	1990	Male	50 to 54	70% (60-79%)	30% (21-40%)	0.73 (0.4-1.2)	2.6% (0.36-4.9%)	2.9% (0.098-5.8%)	82 (2.9-160)	3300 (470-6200)
Mozambique	1990	Male	55 to 59	67% (56-76%)	33% (24-44%)	0.69 (0.38-1.1)	2.2% (0.17-4.2%)	2.4% (-0.027-4.9%)	76 (-0.9-160)	2600 (200-5200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	1990	Male	60 to 64	65% (52-76%)	35% (24-48%)	0.6 (0.32-0.99)	1.5% (-0.051-3.1%)	1.6% (-0.18-3.6%)	58 (-6.6-130)	1700 (-56-3500)
Mozambique	1990	Male	65 to 69	64% (48-77%)	36% (23-52%)	0.49 (0.27-0.8)	0.78% (-0.53-2%)	0.84% (-0.73-2.3%)	32 (-27-87)	780 (-520-2000)
Mozambique	1990	Male	70 to 74	61% (43-75%)	39% (25-57%)	0.39 (0.21-0.66)	0.59% (-0.42-1.5%)	0.67% (-0.53-1.8%)	25 (-20-66)	470 (-340-1200)
Mozambique	1990	Male	75 to 79	57% (40-74%)	43% (26-60%)	0.32 (0.16-0.54)	0.52% (-0.41-1.3%)	0.57% (-0.49-1.5%)	17 (-15-44)	250 (-190-630)
Mozambique	1990	Male	80 to 84	54% (36-72%)	46% (28-64%)	0.27 (0.14-0.44)	0.5% (-0.24-1.1%)	0.52% (-0.35-1.2%)	8 (-5.3-19)	91 (-44-210)
Mozambique	1990	Male	85 to 89	51% (34-67%)	49% (33-66%)	0.23 (0.11-0.39)	0.43% (-0.13-0.89%)	0.43% (-0.23-0.97%)	3.8 (-2-8.7)	33 (-10-68)
Mozambique	1990	Male	90 to 94	51% (35-68%)	49% (32-65%)	0.2 (0.098-0.38)	0.24% (-0.24-0.62%)	0.18% (-0.4-0.62%)	0.63 (-1.4-2.1)	5.1 (-4.9-13)
Mozambique	1990	Male	95 plus	54% (32-74%)	46% (26-68%)	0.2 (0.087-0.38)	0.38% (-0.14-0.77%)	0.29% (-0.34-0.77%)	0.25 (-0.28-0.65)	1.3 (-0.46-2.6)
Mozambique	1995	Female	All Ages	36% (31-42%)	64% (58-69%)	0.34 (0.25-0.45)	0.32% (0.058-0.57%)	0.39% (-0.11-0.9%)	430 (-130-1000)	25000 (4700-45000)
Mozambique	1995	Female	15 to 19	38% (26-51%)	62% (49-74%)	0.42 (0.22-0.69)	0.86% (0.08-1.8%)	0.93% (-0.3-2.5%)	17 (-4.9-46)	1800 (170-3900)
Mozambique	1995	Female	20 to 24	46% (32-60%)	54% (40-68%)	0.58 (0.32-0.93)	1.2% (0.31-2.2%)	0.99% (-0.19-2.5%)	27 (-5.8-66)	3000 (870-5600)
Mozambique	1995	Female	25 to 29	46% (32-59%)	54% (41-68%)	0.66 (0.37-1)	1.5% (0.44-2.9%)	1.4% (0.061-3.3%)	46 (1.9-100)	3900 (1200-7300)
Mozambique	1995	Female	30 to 34	49% (34-61%)	51% (39-66%)	0.69 (0.37-1.1)	1.5% (0.39-3%)	1.5% (0.073-3.6%)	48 (2.5-110)	3500 (940-6900)
Mozambique	1995	Female	35 to 39	54% (40-65%)	46% (35-60%)	0.67 (0.38-1.1)	1.5% (0.096-3.1%)	1.6% (-0.18-3.8%)	48 (-5.4-110)	3000 (200-6200)
Mozambique	1995	Female	40 to 44	56% (44-67%)	44% (33-56%)	0.67 (0.36-1.1)	1.2% (0.11-2.5%)	1.4% (-0.079-3.2%)	39 (-2.2-91)	2100 (190-4400)
Mozambique	1995	Female	45 to 49	54% (40-66%)	46% (34-60%)	0.67 (0.35-1.1)	1.2% (0.047-2.6%)	1.5% (-0.086-3.5%)	42 (-2.4-100)	1900 (76-4400)
Mozambique	1995	Female	50 to 54	50% (37-62%)	50% (38-63%)	0.64 (0.33-1.1)	1.3% (0.013-3.1%)	1.6% (-0.083-4%)	52 (-2.3-130)	2000 (19-4600)
Mozambique	1995	Female	55 to 59	46% (33-59%)	54% (41-67%)	0.6 (0.32-1)	1% (-0.33-2.6%)	1.3% (-0.37-3.4%)	48 (-14-120)	1500 (-490-3900)
Mozambique	1995	Female	60 to 64	46% (32-59%)	54% (41-68%)	0.59 (0.3-1)	0.81% (-0.45-2.3%)	1% (-0.55-3%)	44 (-23-130)	1100 (-640-3300)
Mozambique	1995	Female	65 to 69	48% (35-63%)	52% (37-65%)	0.56 (0.31-0.95)	0.64% (-0.54-2%)	0.78% (-0.65-2.4%)	37 (-30-120)	820 (-670-2600)
Mozambique	1995	Female	70 to 74	47% (33-63%)	53% (37-67%)	0.49 (0.26-0.84)	0.14% (-1.1-1.4%)	0.14% (-1.3-1.7%)	7 (-64-83)	140 (-1100-1500)
Mozambique	1995	Female	75 to 79	45% (30-61%)	55% (39-70%)	0.42 (0.2-0.74)	-0.0052% (-1.1-0.96%)	0.0048% (-1.3-1.2%)	0.14 (-53-49)	-3.4 (-750-670)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	1995	Female	80 to 84	44% (28-60%)	56% (40-72%)	0.39 (0.19-0.65)	-0.29% (-1.2-0.5%)	-0.32% (-1.4-0.6%)	-8.4 (-38-16)	-91 (-390-160)
Mozambique	1995	Female	85 to 89	43% (29-59%)	57% (41-71%)	0.37 (0.19-0.64)	-0.47% (-1.4-0.25%)	-0.56% (-1.7-0.3%)	-9.2 (-27-4.9)	-67 (-200-35)
Mozambique	1995	Female	90 to 94	44% (31-60%)	56% (40-69%)	0.36 (0.17-0.65)	-0.59% (-1.5-0.079%)	-0.74% (-1.8-0.06%)	-5.1 (-12-0.42)	-24 (-60-3.3)
Mozambique	1995	Female	95 plus	47% (30-66%)	53% (34-70%)	0.37 (0.16-0.72)	-0.3% (-1.2-0.42%)	-0.43% (-1.5-0.42%)	-0.87 (-3-0.87)	-2.4 (-9.1-3.4)
Mozambique	1995	Male	All Ages	46% (39-54%)	54% (46-61%)	0.41 (0.3-0.53)	0.71% (0.25-1.1%)	0.96% (0.18-1.7%)	1200 (210-2000)	61000 (21000-98000)
Mozambique	1995	Male	15 to 19	46% (32-60%)	54% (40-68%)	0.49 (0.28-0.79)	2% (0.86-3.4%)	2% (0.3-4.1%)	36 (5.8-74)	3900 (1600-6700)
Mozambique	1995	Male	20 to 24	60% (46-72%)	40% (28-54%)	0.68 (0.39-1.1)	3.2% (1.1-5.4%)	2.8% (0.0073-5.7%)	70 (0.18-140)	7100 (2400-12000)
Mozambique	1995	Male	25 to 29	62% (48-72%)	38% (28-52%)	0.78 (0.46-1.2)	3.1% (0.96-5.4%)	2.7% (0.17-5.7%)	85 (5-170)	7400 (2300-13000)
Mozambique	1995	Male	30 to 34	63% (51-73%)	37% (27-49%)	0.85 (0.52-1.3)	3% (0.69-5.6%)	2.8% (0.12-6%)	100 (4.6-210)	7400 (1800-13000)
Mozambique	1995	Male	35 to 39	66% (54-75%)	34% (25-46%)	0.92 (0.55-1.5)	3.1% (0.68-5.9%)	3.2% (0.24-6.7%)	120 (8.4-240)	7200 (1600-13000)
Mozambique	1995	Male	40 to 44	68% (57-77%)	32% (23-43%)	0.97 (0.55-1.6)	3.2% (0.94-5.8%)	3.4% (0.65-6.7%)	130 (24-260)	6700 (2000-13000)
Mozambique	1995	Male	45 to 49	70% (58-79%)	30% (21-42%)	0.98 (0.55-1.6)	3.2% (0.79-5.9%)	3.6% (0.56-6.9%)	130 (20-250)	6000 (1500-11000)
Mozambique	1995	Male	50 to 54	70% (59-78%)	30% (22-41%)	0.95 (0.52-1.5)	3.2% (0.72-6%)	3.7% (0.55-7.2%)	140 (19-260)	5300 (1100-10000)
Mozambique	1995	Male	55 to 59	67% (56-77%)	33% (23-44%)	0.89 (0.51-1.4)	2.8% (0.62-5.4%)	3.3% (0.54-6.3%)	120 (19-240)	4100 (880-8000)
Mozambique	1995	Male	60 to 64	65% (52-75%)	35% (25-48%)	0.77 (0.42-1.3)	2% (0.33-4%)	2.2% (0.22-4.6%)	93 (9.4-200)	2700 (440-5400)
Mozambique	1995	Male	65 to 69	63% (48-77%)	37% (23-52%)	0.63 (0.35-1)	1.1% (-0.21-2.6%)	1.2% (-0.34-3%)	53 (-14-130)	1300 (-240-2900)
Mozambique	1995	Male	70 to 74	60% (44-75%)	40% (25-56%)	0.5 (0.27-0.87)	0.78% (-0.38-1.9%)	0.89% (-0.48-2.3%)	37 (-21-95)	690 (-340-1700)
Mozambique	1995	Male	75 to 79	57% (40-73%)	43% (27-60%)	0.41 (0.2-0.7)	0.64% (-0.31-1.5%)	0.72% (-0.39-1.8%)	24 (-14-59)	350 (-180-830)
Mozambique	1995	Male	80 to 84	54% (35-71%)	46% (29-65%)	0.35 (0.18-0.58)	0.57% (-0.21-1.2%)	0.61% (-0.31-1.4%)	11 (-5.8-25)	120 (-46-270)
Mozambique	1995	Male	85 to 89	51% (33-67%)	49% (33-67%)	0.29 (0.14-0.5)	0.46% (-0.15-0.98%)	0.46% (-0.25-1.1%)	4.6 (-2.5-11)	39 (-12-84)
Mozambique	1995	Male	90 to 94	50% (34-67%)	50% (33-66%)	0.26 (0.13-0.48)	0.24% (-0.3-0.66%)	0.18% (-0.46-0.67%)	0.69 (-1.8-2.6)	5.7 (-6.9-16)
Mozambique	1995	Male	95 plus	53% (32-73%)	47% (27-68%)	0.25 (0.11-0.49)	0.39% (-0.17-0.85%)	0.29% (-0.38-0.86%)	0.31 (-0.4-0.88)	1.6 (-0.7-3.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	2000	Female	All Ages	35% (30-41%)	65% (59-70%)	0.48 (0.35-0.63)	0.57% (0.22-0.98%)	0.88% (0.18-1.7%)	1100 (210-2200)	48000 (19000-84000)
Mozambique	2000	Female	15 to 19	41% (29-55%)	59% (45-71%)	0.65 (0.37-1.1)	1.4% (0.39-2.8%)	1.8% (0.14-4.1%)	34 (2.8-79)	3200 (860-6500)
Mozambique	2000	Female	20 to 24	48% (35-60%)	52% (40-65%)	0.86 (0.49-1.4)	1.6% (0.59-3%)	1.5% (0.23-3.6%)	53 (7.6-120)	5000 (1800-9700)
Mozambique	2000	Female	25 to 29	44% (33-55%)	56% (45-67%)	0.89 (0.52-1.4)	1.5% (0.6-2.9%)	1.5% (0.4-3.3%)	78 (22-160)	6000 (2500-11000)
Mozambique	2000	Female	30 to 34	46% (35-57%)	54% (43-65%)	0.92 (0.51-1.5)	1.5% (0.61-3.1%)	1.6% (0.47-3.5%)	86 (27-180)	5800 (2500-11000)
Mozambique	2000	Female	35 to 39	51% (41-62%)	49% (38-59%)	0.92 (0.51-1.5)	1.6% (0.51-3.2%)	1.8% (0.39-3.9%)	88 (20-180)	5100 (1500-9900)
Mozambique	2000	Female	40 to 44	54% (43-64%)	46% (36-57%)	0.93 (0.49-1.5)	1.5% (0.41-3.1%)	1.7% (0.34-3.7%)	75 (15-160)	3800 (1100-7600)
Mozambique	2000	Female	45 to 49	52% (40-63%)	48% (37-60%)	0.92 (0.5-1.5)	1.7% (0.43-3.7%)	2.1% (0.42-4.6%)	86 (17-190)	3800 (910-8000)
Mozambique	2000	Female	50 to 54	47% (36-58%)	53% (42-64%)	0.86 (0.46-1.5)	2.1% (0.53-4.5%)	2.7% (0.6-5.8%)	110 (24-240)	4000 (930-8900)
Mozambique	2000	Female	55 to 59	43% (32-55%)	57% (45-68%)	0.82 (0.43-1.4)	2.3% (0.38-5.1%)	2.9% (0.44-6.4%)	120 (18-280)	3900 (630-8700)
Mozambique	2000	Female	60 to 64	44% (31-58%)	56% (42-69%)	0.82 (0.41-1.4)	2% (0.21-4.6%)	2.5% (0.27-5.8%)	120 (13-290)	3200 (340-7800)
Mozambique	2000	Female	65 to 69	48% (34-63%)	52% (37-66%)	0.8 (0.43-1.3)	1.8% (0.079-4.2%)	2.2% (0.11-5.1%)	120 (6-280)	2600 (130-6200)
Mozambique	2000	Female	70 to 74	48% (33-64%)	52% (36-67%)	0.71 (0.37-1.2)	1.1% (-0.64-3.5%)	1.4% (-0.79-4.2%)	74 (-45-230)	1300 (-760-4200)
Mozambique	2000	Female	75 to 79	45% (31-63%)	55% (37-69%)	0.61 (0.29-1.1)	0.77% (-0.71-2.7%)	0.94% (-0.84-3.2%)	44 (-39-150)	590 (-540-2000)
Mozambique	2000	Female	80 to 84	44% (29-61%)	56% (39-71%)	0.56 (0.28-0.95)	0.31% (-0.94-1.8%)	0.4% (-1.1-2.2%)	12 (-33-67)	120 (-350-680)
Mozambique	2000	Female	85 to 89	43% (29-59%)	57% (41-71%)	0.53 (0.27-0.95)	0.013% (-1.1-1.2%)	0.018% (-1.4-1.4%)	0.34 (-25-26)	2.3 (-180-180)
Mozambique	2000	Female	90 to 94	45% (31-60%)	55% (40-69%)	0.53 (0.25-0.96)	-0.2% (-1.2-0.92%)	-0.28% (-1.5-1.1%)	-2.2 (-12-8.6)	-9.6 (-58-44)
Mozambique	2000	Female	95 plus	48% (31-66%)	52% (34-69%)	0.54 (0.23-1)	0.053% (-1-1.3%)	-0.023% (-1.3-1.4%)	-0.052 (-3.1-3.3)	0.52 (-9.3-12)
Mozambique	2000	Male	All Ages	48% (41-56%)	52% (44-59%)	0.62 (0.46-0.81)	1.3% (0.62-2%)	1.8% (0.72-3%)	2400 (960-4100)	120000 (57000-180000)
Mozambique	2000	Male	15 to 19	54% (40-66%)	46% (34-60%)	0.85 (0.5-1.3)	3.2% (1.4-5.4%)	3.7% (1.1-6.8%)	82 (23-160)	7500 (3200-13000)
Mozambique	2000	Male	20 to 24	66% (54-75%)	34% (25-46%)	1.1 (0.68-1.6)	4.8% (2-8.1%)	4.7% (1.1-9.3%)	170 (37-320)	14000 (5700-25000)
Mozambique	2000	Male	25 to 29	63% (54-71%)	37% (29-46%)	1.2 (0.72-1.8)	4% (1.8-7.1%)	3.9% (1.2-7.7%)	180 (55-340)	14000 (6200-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	2000	Male	30 to 34	64% (54-72%)	36% (28-46%)	1.3 (0.77-1.9)	3.8% (1.6-7%)	3.8% (1.3-7.6%)	200 (72-370)	13000 (5700-23000)
Mozambique	2000	Male	35 to 39	66% (57-74%)	34% (26-43%)	1.3 (0.79-2.1)	4% (1.5-7.1%)	4.2% (1.3-8.1%)	230 (70-440)	13000 (4800-23000)
Mozambique	2000	Male	40 to 44	68% (60-76%)	32% (24-40%)	1.4 (0.8-2.2)	4.2% (1.6-7.4%)	4.6% (1.5-8.3%)	260 (88-460)	13000 (5000-22000)
Mozambique	2000	Male	45 to 49	72% (63-79%)	28% (21-37%)	1.5 (0.87-2.3)	4.5% (1.8-8%)	5% (1.9-9.4%)	270 (97-500)	12000 (4700-21000)
Mozambique	2000	Male	50 to 54	71% (63-78%)	29% (22-37%)	1.4 (0.79-2.3)	5% (1.9-8.8%)	5.7% (2.1-10%)	280 (95-530)	11000 (3800-20000)
Mozambique	2000	Male	55 to 59	68% (59-76%)	32% (24-41%)	1.3 (0.75-2.1)	4.7% (1.7-8.6%)	5.5% (1.8-10%)	270 (77-510)	8700 (2800-16000)
Mozambique	2000	Male	60 to 64	66% (56-75%)	34% (25-44%)	1.1 (0.64-1.8)	3.5% (1-6.9%)	4% (1-8.1%)	200 (49-410)	5600 (1600-11000)
Mozambique	2000	Male	65 to 69	64% (49-77%)	36% (23-51%)	0.93 (0.54-1.5)	2.2% (0.3-4.6%)	2.5% (0.23-5.4%)	130 (13-280)	2900 (400-6300)
Mozambique	2000	Male	70 to 74	61% (43-75%)	39% (25-57%)	0.73 (0.39-1.2)	1.4% (-0.16-3.3%)	1.6% (-0.23-4%)	83 (-12-190)	1500 (-170-3500)
Mozambique	2000	Male	75 to 79	58% (40-74%)	42% (26-60%)	0.6 (0.3-1)	1.1% (-0.24-2.7%)	1.3% (-0.32-3.2%)	51 (-13-130)	710 (-160-1800)
Mozambique	2000	Male	80 to 84	54% (35-72%)	46% (28-65%)	0.51 (0.26-0.84)	0.9% (-0.15-2.2%)	1% (-0.24-2.5%)	23 (-5.5-58)	250 (-42-610)
Mozambique	2000	Male	85 to 89	51% (33-67%)	49% (33-67%)	0.43 (0.21-0.74)	0.65% (-0.19-1.5%)	0.69% (-0.32-1.7%)	8.3 (-3.8-21)	67 (-19-160)
Mozambique	2000	Male	90 to 94	51% (34-67%)	49% (33-66%)	0.38 (0.19-0.71)	0.35% (-0.28-1.1%)	0.29% (-0.48-1.2%)	1.3 (-2.2-5.2)	9.3 (-7.6-29)
Mozambique	2000	Male	95 plus	53% (32-73%)	47% (27-68%)	0.37 (0.15-0.71)	0.52% (-0.27-1.3%)	0.44% (-0.49-1.4%)	0.5 (-0.57-1.6)	2.4 (-1.2-6.2)
Mozambique	2005	Female	All Ages	35% (29-40%)	65% (60-71%)	0.56 (0.41-0.74)	0.63% (0.3-1.1%)	1% (0.36-2%)	1400 (460-2800)	56000 (26000-100000)
Mozambique	2005	Female	15 to 19	41% (29-54%)	59% (46-71%)	0.77 (0.43-1.2)	1.5% (0.6-2.7%)	1.9% (0.44-4%)	33 (7.3-76)	3300 (1200-6500)
Mozambique	2005	Female	20 to 24	47% (34-60%)	53% (40-66%)	1 (0.6-1.6)	1.6% (0.69-3%)	1.5% (0.39-3.5%)	55 (13-120)	5300 (2400-10000)
Mozambique	2005	Female	25 to 29	44% (33-54%)	56% (46-67%)	1.1 (0.6-1.7)	1.2% (0.47-2.4%)	1.1% (0.28-2.6%)	75 (24-160)	6100 (2800-11000)
Mozambique	2005	Female	30 to 34	45% (34-57%)	55% (43-66%)	1.1 (0.61-1.8)	1.1% (0.45-2.2%)	1% (0.31-2.3%)	82 (28-180)	5700 (2600-11000)
Mozambique	2005	Female	35 to 39	51% (40-61%)	49% (39-60%)	1.1 (0.6-1.7)	1.1% (0.42-2.4%)	1.2% (0.31-2.7%)	90 (25-200)	5300 (2000-11000)
Mozambique	2005	Female	40 to 44	53% (43-64%)	47% (36-57%)	1.1 (0.6-1.8)	1.2% (0.4-2.4%)	1.2% (0.35-2.7%)	83 (23-180)	4300 (1500-8700)
Mozambique	2005	Female	45 to 49	52% (40-62%)	48% (38-60%)	1.1 (0.62-1.8)	1.6% (0.56-3.4%)	1.9% (0.57-4.2%)	110 (33-230)	4600 (1600-9600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	2005	Female	50 to 54	47% (35-58%)	53% (42-65%)	1 (0.54-1.7)	2.2% (0.73-4.3%)	2.6% (0.84-5.4%)	130 (38-270)	4800 (1500-10000)
Mozambique	2005	Female	55 to 59	43% (31-55%)	57% (45-69%)	0.97 (0.51-1.7)	2.6% (0.74-5.4%)	3.3% (0.93-7%)	150 (40-340)	4800 (1300-11000)
Mozambique	2005	Female	60 to 64	44% (31-57%)	56% (43-69%)	0.97 (0.5-1.7)	2.5% (0.66-5.4%)	3.3% (0.84-6.8%)	160 (38-370)	4300 (1000-9800)
Mozambique	2005	Female	65 to 69	47% (34-63%)	53% (37-66%)	0.96 (0.52-1.6)	2.5% (0.71-5.1%)	3.2% (0.85-6.4%)	170 (43-370)	3700 (930-8200)
Mozambique	2005	Female	70 to 74	47% (33-63%)	53% (37-67%)	0.85 (0.42-1.5)	1.9% (-0.12-4.7%)	2.4% (-0.16-5.8%)	130 (-7.5-350)	2300 (-130-6200)
Mozambique	2005	Female	75 to 79	45% (30-62%)	55% (38-70%)	0.72 (0.35-1.3)	1.4% (-0.28-3.9%)	1.8% (-0.34-4.8%)	88 (-16-240)	1200 (-240-3200)
Mozambique	2005	Female	80 to 84	43% (27-61%)	57% (39-73%)	0.67 (0.34-1.1)	0.87% (-0.57-2.7%)	1.1% (-0.65-3.3%)	39 (-21-120)	390 (-240-1200)
Mozambique	2005	Female	85 to 89	43% (28-59%)	57% (41-72%)	0.64 (0.32-1.1)	0.49% (-0.78-2%)	0.6% (-0.92-2.5%)	13 (-19-53)	90 (-140-380)
Mozambique	2005	Female	90 to 94	45% (31-61%)	55% (39-69%)	0.63 (0.31-1.2)	0.19% (-0.96-1.7%)	0.21% (-1.2-2%)	1.8 (-10-17)	10 (-49-87)
Mozambique	2005	Female	95 plus	48% (30-66%)	52% (34-70%)	0.64 (0.28-1.3)	0.46% (-0.75-2.1%)	0.47% (-1-2.5%)	1.2 (-2.5-6.3)	4.6 (-7.5-21)
Mozambique	2005	Male	All Ages	48% (41-55%)	52% (45-59%)	0.72 (0.53-0.95)	1.6% (0.89-2.5%)	2.3% (1.1-3.7%)	3400 (1500-5500)	160000 (84000-240000)
Mozambique	2005	Male	15 to 19	53% (40-66%)	47% (34-60%)	1 (0.6-1.6)	3.7% (1.8-6.1%)	4.5% (1.7-8%)	110 (38-200)	9700 (4500-17000)
Mozambique	2005	Male	20 to 24	66% (54-76%)	34% (24-46%)	1.3 (0.77-2)	5.6% (2.6-9.6%)	5.9% (2-11%)	230 (77-440)	19000 (8500-33000)
Mozambique	2005	Male	25 to 29	63% (54-72%)	37% (28-46%)	1.4 (0.84-2.1)	4.4% (2-7.8%)	4.3% (1.6-8.4%)	270 (100-490)	20000 (9500-33000)
Mozambique	2005	Male	30 to 34	64% (55-72%)	36% (28-45%)	1.5 (0.9-2.3)	3.8% (1.8-6.9%)	3.8% (1.5-7.6%)	280 (110-490)	18000 (8400-29000)
Mozambique	2005	Male	35 to 39	66% (57-74%)	34% (26-43%)	1.6 (0.94-2.5)	3.9% (1.7-7%)	4.1% (1.6-7.8%)	290 (110-510)	16000 (7200-27000)
Mozambique	2005	Male	40 to 44	68% (60-77%)	32% (23-40%)	1.7 (1-2.6)	4.2% (1.9-7.4%)	4.5% (1.8-8.2%)	320 (130-570)	16000 (7000-27000)
Mozambique	2005	Male	45 to 49	72% (63-79%)	28% (21-37%)	1.7 (1-2.8)	4.7% (2.1-8.1%)	5.2% (2.2-9.2%)	350 (150-620)	15000 (6800-26000)
Mozambique	2005	Male	50 to 54	71% (63-78%)	29% (22-37%)	1.7 (0.93-2.7)	5.5% (2.4-9.5%)	6.3% (2.6-11%)	380 (150-690)	14000 (6200-25000)
Mozambique	2005	Male	55 to 59	68% (59-76%)	32% (24-41%)	1.5 (0.91-2.4)	5.6% (2.4-9.8%)	6.6% (2.6-12%)	370 (130-700)	12000 (4500-22000)
Mozambique	2005	Male	60 to 64	66% (56-75%)	34% (25-44%)	1.4 (0.76-2.2)	4.3% (1.6-8.2%)	5% (1.8-9.5%)	290 (94-570)	8100 (2700-16000)
Mozambique	2005	Male	65 to 69	64% (48-77%)	36% (23-52%)	1.1 (0.63-1.8)	2.9% (0.67-5.9%)	3.4% (0.73-6.9%)	200 (39-410)	4500 (950-9000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	2005	Male	70 to 74	60% (43-75%)	40% (25-57%)	0.86 (0.45-1.5)	1.9% (0.15-4.4%)	2.3% (0.089-5.1%)	130 (5-290)	2400 (190-5200)
Mozambique	2005	Male	75 to 79	57% (40-73%)	43% (27-60%)	0.7 (0.35-1.2)	1.5% (-0.033-3.6%)	1.8% (-0.073-4.3%)	84 (-3.6-200)	1200 (-22-2800)
Mozambique	2005	Male	80 to 84	53% (34-70%)	47% (30-66%)	0.6 (0.3-0.98)	1.2% (-0.0079-2.9%)	1.4% (-0.073-3.3%)	41 (-2-99)	430 (-2.7-1000)
Mozambique	2005	Male	85 to 89	50% (33-66%)	50% (34-67%)	0.5 (0.25-0.87)	0.89% (-0.058-2%)	0.97% (-0.18-2.3%)	14 (-2.7-35)	110 (-7.9-260)
Mozambique	2005	Male	90 to 94	50% (34-67%)	50% (33-66%)	0.45 (0.22-0.81)	0.5% (-0.23-1.4%)	0.47% (-0.43-1.5%)	2.4 (-2.2-7.8)	15 (-7.3-42)
Mozambique	2005	Male	95 plus	53% (32-72%)	47% (28-68%)	0.43 (0.18-0.82)	0.71% (-0.23-1.9%)	0.66% (-0.47-2%)	0.83 (-0.6-2.5)	3.6 (-1.1-9.3)
Mozambique	2010	Female	All Ages	36% (30-42%)	64% (58-70%)	0.57 (0.4-0.77)	0.66% (0.29-1.1%)	1.1% (0.3-2.1%)	1500 (390-3000)	59000 (25000-100000)
Mozambique	2010	Female	15 to 19	38% (26-52%)	62% (48-74%)	0.71 (0.39-1.2)	1.2% (0.51-2.3%)	1.5% (0.29-3.3%)	32 (6.6-76)	3400 (1300-6800)
Mozambique	2010	Female	20 to 24	46% (32-60%)	54% (40-68%)	0.99 (0.54-1.6)	1.5% (0.66-2.8%)	1.4% (0.32-3.3%)	49 (12-110)	5100 (2400-9600)
Mozambique	2010	Female	25 to 29	47% (33-59%)	53% (41-67%)	1.1 (0.61-1.8)	1.1% (0.49-2.3%)	1% (0.29-2.5%)	72 (24-160)	6100 (2800-11000)
Mozambique	2010	Female	30 to 34	49% (36-61%)	51% (39-64%)	1.2 (0.65-2)	0.99% (0.42-2%)	0.92% (0.29-2.1%)	82 (27-170)	5900 (2600-11000)
Mozambique	2010	Female	35 to 39	54% (41-65%)	46% (35-59%)	1.2 (0.65-1.9)	1% (0.39-2.1%)	1% (0.29-2.3%)	87 (25-190)	5200 (2100-10000)
Mozambique	2010	Female	40 to 44	56% (44-67%)	44% (33-56%)	1.2 (0.63-1.9)	1.1% (0.38-2.1%)	1.1% (0.3-2.4%)	85 (22-180)	4400 (1500-8900)
Mozambique	2010	Female	45 to 49	54% (40-65%)	46% (35-60%)	1.2 (0.61-2)	1.6% (0.49-3.2%)	1.9% (0.51-3.9%)	110 (32-240)	5000 (1600-10000)
Mozambique	2010	Female	50 to 54	50% (37-62%)	50% (38-63%)	1.1 (0.58-1.9)	2.3% (0.65-4.5%)	2.8% (0.78-5.7%)	150 (37-310)	5400 (1500-12000)
Mozambique	2010	Female	55 to 59	46% (34-59%)	54% (41-66%)	1.1 (0.55-1.8)	2.7% (0.63-5.4%)	3.6% (0.83-7.1%)	170 (35-360)	5300 (1200-11000)
Mozambique	2010	Female	60 to 64	47% (32-60%)	53% (40-68%)	1 (0.51-1.8)	2.6% (0.5-5.6%)	3.4% (0.72-7.3%)	170 (30-390)	4600 (770-11000)
Mozambique	2010	Female	65 to 69	49% (35-64%)	51% (36-65%)	0.98 (0.51-1.7)	2.5% (0.43-5.3%)	3.3% (0.56-6.7%)	180 (30-410)	4100 (640-9000)
Mozambique	2010	Female	70 to 74	48% (33-64%)	52% (36-67%)	0.85 (0.43-1.5)	1.9% (-0.16-4.9%)	2.4% (-0.21-6.1%)	140 (-10-400)	2600 (-190-7300)
Mozambique	2010	Female	75 to 79	45% (31-63%)	55% (37-69%)	0.73 (0.35-1.3)	1.4% (-0.36-3.9%)	1.8% (-0.43-4.8%)	100 (-23-280)	1400 (-320-3900)
Mozambique	2010	Female	80 to 84	44% (29-62%)	56% (38-71%)	0.67 (0.33-1.1)	0.87% (-0.56-2.7%)	1.1% (-0.66-3.4%)	47 (-29-140)	470 (-300-1500)
Mozambique	2010	Female	85 to 89	44% (29-59%)	56% (41-71%)	0.64 (0.32-1.1)	0.5% (-0.85-2%)	0.62% (-1-2.4%)	16 (-26-64)	120 (-200-460)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	2010	Female	90 to 94	45% (31-61%)	55% (39-69%)	0.63 (0.3-1.2)	0.2% (-1-1.8%)	0.21% (-1.2-2.3%)	2.2 (-13-23)	12 (-62-110)
Mozambique	2010	Female	95 plus	48% (31-67%)	52% (33-69%)	0.64 (0.27-1.3)	0.49% (-0.73-2.4%)	0.51% (-1-2.8%)	1.4 (-2.8-8)	5.6 (-8.2-27)
Mozambique	2010	Male	All Ages	45% (38-53%)	55% (47-62%)	0.67 (0.48-0.92)	1.7% (0.92-2.7%)	2.4% (1.1-3.9%)	3700 (1600-6000)	170000 (89000-260000)
Mozambique	2010	Male	15 to 19	46% (32-61%)	54% (39-68%)	0.84 (0.47-1.3)	3.1% (1.5-5.1%)	3.5% (1.3-6.4%)	110 (35-210)	10000 (4600-17000)
Mozambique	2010	Male	20 to 24	61% (46-73%)	39% (27-54%)	1.2 (0.66-1.9)	5.5% (2.6-9.1%)	5.6% (2-10%)	240 (84-440)	20000 (9500-34000)
Mozambique	2010	Male	25 to 29	62% (49-72%)	38% (28-51%)	1.3 (0.78-2.1)	4.6% (2.1-7.9%)	4.5% (1.6-8.8%)	310 (110-550)	23000 (11000-37000)
Mozambique	2010	Male	30 to 34	64% (52-74%)	36% (26-48%)	1.5 (0.87-2.3)	3.9% (1.9-7.1%)	3.9% (1.6-7.7%)	340 (130-590)	22000 (10000-36000)
Mozambique	2010	Male	35 to 39	66% (55-75%)	34% (25-45%)	1.6 (0.92-2.6)	3.8% (1.6-6.8%)	4% (1.5-7.6%)	320 (120-580)	18000 (8100-31000)
Mozambique	2010	Male	40 to 44	68% (57-77%)	32% (23-43%)	1.7 (0.9-2.6)	4.1% (1.8-7.1%)	4.4% (1.7-7.8%)	320 (130-570)	16000 (6800-27000)
Mozambique	2010	Male	45 to 49	70% (57-79%)	30% (21-43%)	1.7 (0.93-2.7)	4.6% (2.1-7.9%)	5.1% (2.1-9%)	360 (140-620)	15000 (6800-26000)
Mozambique	2010	Male	50 to 54	70% (60-78%)	30% (22-40%)	1.6 (0.88-2.7)	5.5% (2.3-9.6%)	6.3% (2.5-11%)	400 (150-740)	15000 (6000-27000)
Mozambique	2010	Male	55 to 59	68% (56-77%)	32% (23-44%)	1.5 (0.83-2.4)	5.7% (2.4-9.9%)	6.7% (2.7-12%)	410 (160-780)	13000 (5200-25000)
Mozambique	2010	Male	60 to 64	65% (53-76%)	35% (24-47%)	1.3 (0.72-2.2)	4.3% (1.5-8.3%)	5% (1.7-9.7%)	320 (99-660)	8900 (2900-18000)
Mozambique	2010	Male	65 to 69	64% (48-77%)	36% (23-52%)	1.1 (0.6-1.8)	2.9% (0.61-6%)	3.4% (0.66-6.9%)	220 (44-460)	5100 (1100-10000)
Mozambique	2010	Male	70 to 74	61% (43-75%)	39% (25-57%)	0.86 (0.46-1.5)	2% (0.14-4.4%)	2.3% (0.12-5.2%)	150 (9.4-340)	2700 (200-6000)
Mozambique	2010	Male	75 to 79	57% (40-73%)	43% (27-60%)	0.7 (0.35-1.2)	1.5% (-0.083-3.7%)	1.8% (-0.16-4.3%)	100 (-8.4-240)	1400 (-76-3400)
Mozambique	2010	Male	80 to 84	54% (35-72%)	46% (28-65%)	0.6 (0.3-1)	1.2% (-0.026-2.8%)	1.4% (-0.086-3.3%)	52 (-3-120)	550 (-11-1300)
Mozambique	2010	Male	85 to 89	51% (33-67%)	49% (33-67%)	0.5 (0.25-0.87)	0.92% (-0.1-2.2%)	1% (-0.2-2.5%)	18 (-3.9-46)	150 (-16-340)
Mozambique	2010	Male	90 to 94	50% (34-67%)	50% (33-66%)	0.45 (0.21-0.85)	0.52% (-0.26-1.4%)	0.5% (-0.43-1.6%)	3.1 (-2.7-10)	20 (-9.4-55)
Mozambique	2010	Male	95 plus	53% (33-73%)	47% (27-67%)	0.43 (0.17-0.89)	0.76% (-0.24-2.1%)	0.72% (-0.5-2.3%)	1 (-0.74-3.4)	4.4 (-1.4-12)
Mozambique	2016	Female	All Ages	36% (30-42%)	64% (58-70%)	0.6 (0.35-0.96)	0.95% (0.33-1.7%)	1.6% (0.32-3.3%)	1900 (380-4100)	74000 (25000-140000)
Mozambique	2016	Female	15 to 19	39% (27-52%)	61% (48-73%)	0.76 (0.37-1.4)	1.3% (0.5-2.5%)	1.5% (0.28-3.3%)	43 (8.2-97)	4500 (1700-8600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	2016	Female	20 to 24	47% (33-60%)	53% (40-67%)	1 (0.5-1.9)	1.8% (0.8-3.4%)	1.8% (0.37-3.9%)	63 (14-140)	6700 (3000-12000)
Mozambique	2016	Female	25 to 29	47% (33-59%)	53% (41-67%)	1.2 (0.58-2.2)	1.6% (0.72-3%)	1.6% (0.43-3.4%)	82 (26-170)	7100 (3300-12000)
Mozambique	2016	Female	30 to 34	49% (36-62%)	51% (38-64%)	1.3 (0.58-2.3)	1.6% (0.69-3.1%)	1.7% (0.47-3.6%)	96 (31-200)	6900 (3000-13000)
Mozambique	2016	Female	35 to 39	54% (41-66%)	46% (34-59%)	1.2 (0.58-2.3)	1.8% (0.63-3.4%)	2% (0.52-4.2%)	110 (27-220)	6400 (2300-12000)
Mozambique	2016	Female	40 to 44	56% (45-67%)	44% (33-55%)	1.2 (0.56-2.3)	1.9% (0.58-3.8%)	2.3% (0.53-4.8%)	100 (24-220)	5300 (1700-11000)
Mozambique	2016	Female	45 to 49	55% (40-66%)	45% (34-60%)	1.2 (0.54-2.4)	2.5% (0.67-5.5%)	3.3% (0.79-7.5%)	140 (32-320)	6000 (1600-14000)
Mozambique	2016	Female	50 to 54	51% (38-62%)	49% (38-62%)	1.2 (0.51-2.3)	3.2% (0.84-6.6%)	4.3% (1-9.2%)	180 (45-410)	6900 (1800-15000)
Mozambique	2016	Female	55 to 59	47% (34-59%)	53% (41-66%)	1.1 (0.5-2.2)	3.5% (0.61-7.4%)	4.9% (0.8-10%)	210 (35-470)	6700 (1100-15000)
Mozambique	2016	Female	60 to 64	47% (33-60%)	53% (40-67%)	1.1 (0.49-2)	3.1% (0.44-7.2%)	4.2% (0.59-9.4%)	220 (32-510)	5900 (840-14000)
Mozambique	2016	Female	65 to 69	49% (35-65%)	51% (35-65%)	1 (0.48-1.9)	2.9% (0.45-7.1%)	3.8% (0.58-9%)	240 (35-590)	5300 (790-13000)
Mozambique	2016	Female	70 to 74	48% (34-64%)	52% (36-66%)	0.9 (0.38-1.8)	2.1% (-0.22-6.1%)	2.7% (-0.25-7.6%)	190 (-18-570)	3500 (-330-10000)
Mozambique	2016	Female	75 to 79	45% (31-62%)	55% (38-69%)	0.78 (0.33-1.5)	1.6% (-0.35-4.9%)	2.1% (-0.42-6.1%)	150 (-27-430)	2000 (-410-6000)
Mozambique	2016	Female	80 to 84	44% (29-62%)	56% (38-71%)	0.71 (0.33-1.4)	0.97% (-0.61-3.4%)	1.2% (-0.7-4%)	70 (-40-230)	700 (-430-2400)
Mozambique	2016	Female	85 to 89	44% (29-59%)	56% (41-71%)	0.68 (0.3-1.3)	0.59% (-0.77-2.7%)	0.72% (-0.93-3.2%)	26 (-33-110)	180 (-240-810)
Mozambique	2016	Female	90 to 94	46% (32-62%)	54% (38-68%)	0.68 (0.29-1.4)	0.29% (-1-2.3%)	0.33% (-1.3-2.9%)	4.7 (-19-41)	26 (-93-210)
Mozambique	2016	Female	95 plus	49% (32-68%)	51% (32-68%)	0.69 (0.26-1.6)	0.58% (-0.76-3%)	0.62% (-1-3.5%)	2.3 (-4-13)	8.8 (-12-45)
Mozambique	2016	Male	All Ages	46% (39-54%)	54% (46-61%)	0.72 (0.41-1.1)	2.1% (0.98-3.5%)	2.8% (1.1-5%)	4000 (1500-7200)	190000 (87000-320000)
Mozambique	2016	Male	15 to 19	46% (32-61%)	54% (39-68%)	0.89 (0.43-1.6)	2.8% (1.3-4.9%)	3% (0.96-5.8%)	120 (37-250)	12000 (5100-21000)
Mozambique	2016	Male	20 to 24	61% (46-73%)	39% (27-54%)	1.2 (0.61-2.2)	5.7% (2.6-9.8%)	5.7% (1.8-11%)	260 (77-530)	23000 (9800-41000)
Mozambique	2016	Male	25 to 29	62% (48-73%)	38% (27-52%)	1.4 (0.71-2.5)	5% (2.2-8.9%)	5% (1.6-9.9%)	320 (100-610)	25000 (11000-42000)
Mozambique	2016	Male	30 to 34	64% (51-74%)	36% (26-49%)	1.5 (0.77-2.7)	4.5% (1.9-8%)	4.5% (1.5-8.6%)	360 (120-680)	24000 (10000-42000)
Mozambique	2016	Male	35 to 39	66% (54-75%)	34% (25-46%)	1.7 (0.79-3)	4.5% (1.8-8.2%)	4.7% (1.5-9.2%)	380 (120-730)	22000 (8500-40000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mozambique	2016	Male	40 to 44	68% (57-77%)	32% (23-43%)	1.8 (0.82-3.3)	4.9% (2-8.8%)	5.3% (1.9-9.9%)	350 (120-670)	18000 (6900-33000)
Mozambique	2016	Male	45 to 49	70% (58-79%)	30% (21-42%)	1.8 (0.82-3.4)	5.6% (2.2-10%)	6.4% (2.3-12%)	340 (120-660)	15000 (5900-28000)
Mozambique	2016	Male	50 to 54	70% (60-78%)	30% (22-40%)	1.7 (0.77-3.2)	6.4% (2.3-12%)	7.6% (2.6-14%)	400 (130-760)	15000 (5200-28000)
Mozambique	2016	Male	55 to 59	68% (56-77%)	32% (23-44%)	1.6 (0.76-2.8)	6.4% (2.4-12%)	7.8% (2.7-14%)	420 (140-790)	14000 (4800-25000)
Mozambique	2016	Male	60 to 64	65% (53-76%)	35% (24-47%)	1.4 (0.64-2.6)	4.8% (1.4-10%)	5.7% (1.5-12%)	360 (97-750)	9900 (2900-21000)
Mozambique	2016	Male	65 to 69	63% (48-76%)	37% (24-52%)	1.1 (0.52-2.1)	3.1% (0.59-7.1%)	3.8% (0.63-8.6%)	250 (41-580)	5800 (1100-13000)
Mozambique	2016	Male	70 to 74	60% (43-74%)	40% (26-57%)	0.89 (0.39-1.8)	2.1% (0.21-5.5%)	2.6% (0.22-6.6%)	180 (16-450)	3300 (340-8100)
Mozambique	2016	Male	75 to 79	57% (39-73%)	43% (27-61%)	0.73 (0.32-1.4)	1.6% (0.0000038-4.2%)	1.9% (-0.044-5.1%)	120 (-2.6-320)	1700 (0.004-4500)
Mozambique	2016	Male	80 to 84	54% (34-71%)	46% (29-66%)	0.62 (0.29-1.2)	1.3% (0.021-3.1%)	1.5% (-0.039-3.6%)	68 (-1.8-170)	720 (12-1700)
Mozambique	2016	Male	85 to 89	51% (33-68%)	49% (32-67%)	0.52 (0.23-1.1)	0.96% (-0.052-2.5%)	1.1% (-0.17-2.8%)	24 (-3.5-65)	190 (-10-490)
Mozambique	2016	Male	90 to 94	51% (35-68%)	49% (32-65%)	0.47 (0.2-0.98)	0.57% (-0.25-1.8%)	0.55% (-0.42-2%)	4.3 (-3.2-16)	27 (-12-83)
Mozambique	2016	Male	95 plus	54% (33-73%)	46% (27-67%)	0.45 (0.17-1)	0.78% (-0.25-2.3%)	0.74% (-0.52-2.5%)	1.3 (-0.93-4.6)	5.7 (-1.8-17)
Rwanda	1990	Female	All Ages	39% (34-44%)	61% (56-66%)	1.6 (1.2-2.2)	3.6% (2.5-5%)	6.7% (4.4-9.5%)	3600 (2300-5100)	130000 (89000-180000)
Rwanda	1990	Female	15 to 19	41% (31-50%)	59% (50-69%)	2.1 (1.2-3.4)	7.5% (3.8-13%)	10% (5-17%)	100 (50-180)	7800 (3900-13000)
Rwanda	1990	Female	20 to 24	52% (41-62%)	48% (38-59%)	3 (1.8-4.8)	9.1% (5-14%)	11% (5.9-18%)	150 (77-240)	10000 (5600-16000)
Rwanda	1990	Female	25 to 29	56% (45-66%)	44% (34-55%)	3.7 (2.3-5.8)	11% (6-16%)	13% (7-20%)	190 (100-290)	12000 (6900-18000)
Rwanda	1990	Female	30 to 34	59% (48-68%)	41% (32-52%)	3.9 (2.4-5.9)	11% (6.9-17%)	14% (8.3-22%)	230 (130-360)	13000 (7700-20000)
Rwanda	1990	Female	35 to 39	62% (51-69%)	38% (31-49%)	3.6 (2.2-5.4)	12% (7.5-18%)	15% (8.9-23%)	250 (150-380)	13000 (7800-19000)
Rwanda	1990	Female	40 to 44	62% (53-69%)	38% (31-47%)	3.5 (2.1-5.5)	12% (7.1-18%)	15% (8.7-22%)	220 (130-340)	10000 (5900-16000)
Rwanda	1990	Female	45 to 49	61% (51-68%)	39% (32-49%)	3.6 (2-5.7)	14% (8.2-21%)	17% (9.9-26%)	220 (130-330)	9100 (5300-14000)
Rwanda	1990	Female	50 to 54	58% (47-67%)	42% (33-53%)	3.5 (1.9-5.6)	17% (9.4-24%)	20% (11-29%)	300 (170-450)	11000 (6100-16000)
Rwanda	1990	Female	55 to 59	55% (39-71%)	45% (29-61%)	3.5 (1.8-5.9)	17% (9.5-24%)	19% (11-29%)	400 (230-610)	13000 (7200-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	1990	Female	60 to 64	56% (38-73%)	44% (27-62%)	3.4 (1.8-6)	15% (8.4-22%)	17% (9.7-26%)	430 (240-650)	12000 (6300-17000)
Rwanda	1990	Female	65 to 69	57% (40-73%)	43% (27-60%)	3.1 (1.8-5.2)	14% (8.1-21%)	16% (9.1-24%)	410 (230-620)	9000 (5000-14000)
Rwanda	1990	Female	70 to 74	55% (39-71%)	45% (29-61%)	2.7 (1.5-4.6)	12% (5.9-20%)	14% (6.5-22%)	300 (140-480)	5300 (2500-8600)
Rwanda	1990	Female	75 to 79	53% (38-70%)	47% (30-62%)	2.4 (1.2-4.3)	10% (4.3-18%)	11% (4.9-20%)	240 (100-420)	3300 (1400-5800)
Rwanda	1990	Female	80 to 84	52% (35-69%)	48% (31-65%)	2.2 (1.2-4)	8.2% (3.4-15%)	9.3% (3.9-17%)	120 (48-210)	1200 (490-2200)
Rwanda	1990	Female	85 to 89	51% (36-66%)	49% (34-64%)	2.1 (1.1-3.7)	6.6% (2.4-12%)	7.5% (2.8-14%)	48 (18-90)	350 (130-660)
Rwanda	1990	Female	90 to 94	53% (37-68%)	47% (32-63%)	2 (1-3.7)	5.1% (1.5-11%)	6% (1.7-12%)	14 (3.8-28)	69 (20-140)
Rwanda	1990	Female	95 plus	55% (36-73%)	45% (27-64%)	2 (0.88-3.7)	4.9% (0.95-11%)	5.6% (0.99-12%)	3.7 (0.65-8.1)	13 (2.5-27)
Rwanda	1990	Male	All Ages	59% (53-65%)	41% (35-47%)	2.1 (1.6-2.8)	5.2% (3.8-6.7%)	8.5% (6.1-11%)	4900 (3400-6400)	210000 (150000-270000)
Rwanda	1990	Male	15 to 19	59% (48-68%)	41% (32-52%)	2.9 (1.8-4.5)	11% (6.7-16%)	14% (7.6-20%)	180 (96-260)	14000 (8000-20000)
Rwanda	1990	Male	20 to 24	68% (58-76%)	32% (24-42%)	3.6 (2.2-5.5)	15% (9.5-22%)	16% (9.6-24%)	300 (190-440)	22000 (14000-32000)
Rwanda	1990	Male	25 to 29	75% (66-81%)	25% (19-34%)	4.4 (2.7-6.5)	16% (10-23%)	17% (11-26%)	370 (240-510)	24000 (16000-33000)
Rwanda	1990	Male	30 to 34	78% (70-83%)	22% (17-30%)	4.9 (3.1-7.2)	17% (12-24%)	19% (13-27%)	420 (290-570)	25000 (17000-33000)
Rwanda	1990	Male	35 to 39	78% (71-83%)	22% (17-29%)	5.1 (3.1-7.8)	19% (13-26%)	21% (14-30%)	450 (310-610)	24000 (17000-32000)
Rwanda	1990	Male	40 to 44	79% (72-84%)	21% (16-28%)	5.2 (3.1-8.1)	21% (14-28%)	23% (16-32%)	460 (310-630)	21000 (14000-29000)
Rwanda	1990	Male	45 to 49	81% (74-86%)	19% (14-26%)	5.3 (3-8.2)	22% (14-29%)	25% (16-33%)	390 (250-550)	16000 (11000-23000)
Rwanda	1990	Male	50 to 54	82% (77-86%)	18% (14-23%)	5.2 (3-8.2)	24% (15-32%)	27% (17-36%)	450 (280-640)	17000 (10000-23000)
Rwanda	1990	Male	55 to 59	81% (74-86%)	19% (14-26%)	5 (3-8)	23% (15-31%)	26% (18-35%)	550 (350-780)	18000 (11000-25000)
Rwanda	1990	Male	60 to 64	79% (69-86%)	21% (14-31%)	4.4 (2.6-7.3)	19% (12-27%)	22% (14-30%)	520 (310-760)	14000 (8300-20000)
Rwanda	1990	Male	65 to 69	75% (59-87%)	25% (13-41%)	3.6 (2.2-6)	16% (9.2-23%)	18% (10-26%)	360 (200-540)	7900 (4500-12000)
Rwanda	1990	Male	70 to 74	73% (57-84%)	27% (16-43%)	2.9 (1.7-4.8)	12% (5.9-19%)	13% (6.8-21%)	200 (94-330)	3500 (1700-5800)
Rwanda	1990	Male	75 to 79	71% (56-84%)	29% (16-44%)	2.5 (1.3-4.3)	9.4% (3.6-17%)	11% (4.2-19%)	120 (47-210)	1700 (650-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	1990	Male	80 to 84	70% (52-83%)	30% (17-48%)	2.2 (1.2-3.7)	7.8% (3.4-13%)	8.8% (3.7-15%)	57 (24-100)	590 (250-1000)
Rwanda	1990	Male	85 to 89	67% (50-80%)	33% (20-50%)	1.9 (1.1-3.2)	5.9% (2.4-10%)	6.7% (2.7-12%)	24 (9.6-44)	180 (74-320)
Rwanda	1990	Male	90 to 94	67% (52-81%)	33% (19-48%)	1.7 (0.93-2.9)	3.9% (1.3-7.1%)	4.4% (1.3-8.2%)	5.9 (1.7-11)	31 (10-57)
Rwanda	1990	Male	95 plus	68% (49-84%)	32% (16-51%)	1.6 (0.72-3)	4.1% (0.93-8.7%)	4.5% (0.76-9.9%)	1.6 (0.26-3.4)	5.7 (1.3-12)
Rwanda	1995	Female	All Ages	41% (37-45%)	59% (55-63%)	1.6 (1.2-2.2)	3.7% (2.2-5.3%)	6.6% (3.9-9.7%)	3500 (2000-5200)	130000 (80000-190000)
Rwanda	1995	Female	15 to 19	44% (39-48%)	56% (52-61%)	2.1 (1.4-3.3)	7.3% (3.4-13%)	9.8% (4.3-18%)	120 (51-200)	8600 (4000-15000)
Rwanda	1995	Female	20 to 24	55% (51-59%)	45% (41-49%)	3.1 (1.9-4.5)	8.7% (4.6-14%)	11% (5.2-18%)	170 (82-280)	12000 (6000-19000)
Rwanda	1995	Female	25 to 29	58% (54-62%)	42% (38-46%)	3.6 (2.3-5.5)	9.1% (4.6-14%)	11% (5.1-17%)	210 (110-330)	13000 (6900-20000)
Rwanda	1995	Female	30 to 34	60% (56-64%)	40% (36-44%)	3.7 (2.4-5.6)	9.5% (5.2-15%)	11% (5.7-18%)	230 (120-390)	13000 (7400-22000)
Rwanda	1995	Female	35 to 39	60% (57-65%)	40% (35-43%)	3.3 (2.1-5)	10% (5.5-16%)	12% (6.3-20%)	260 (140-420)	13000 (7400-21000)
Rwanda	1995	Female	40 to 44	60% (56-65%)	40% (35-44%)	3.2 (1.9-5)	10% (5.5-17%)	12% (6.4-20%)	230 (120-370)	11000 (5700-17000)
Rwanda	1995	Female	45 to 49	59% (54-63%)	41% (37-46%)	3.2 (1.8-5.1)	13% (6.6-20%)	15% (7.9-24%)	270 (140-430)	11000 (5900-18000)
Rwanda	1995	Female	50 to 54	57% (47-66%)	43% (34-53%)	3.2 (1.8-5.3)	15% (8-24%)	18% (9.4-28%)	320 (160-500)	11000 (5700-18000)
Rwanda	1995	Female	55 to 59	57% (41-73%)	43% (27-59%)	3.4 (1.8-5.8)	17% (8.5-26%)	19% (10-30%)	310 (160-500)	9700 (5000-16000)
Rwanda	1995	Female	60 to 64	57% (41-75%)	43% (25-59%)	3.3 (1.7-5.7)	15% (7.7-24%)	18% (8.9-28%)	410 (200-630)	11000 (5400-17000)
Rwanda	1995	Female	65 to 69	58% (41-74%)	42% (26-59%)	3 (1.7-5.1)	15% (7.5-23%)	16% (8.5-25%)	380 (190-590)	8400 (4300-13000)
Rwanda	1995	Female	70 to 74	56% (39-72%)	44% (28-61%)	2.6 (1.4-4.5)	13% (5.5-21%)	14% (6.2-23%)	260 (110-440)	4700 (2000-7800)
Rwanda	1995	Female	75 to 79	54% (38-70%)	46% (30-62%)	2.3 (1.2-4.2)	11% (3.9-19%)	12% (4.4-21%)	170 (64-300)	2300 (880-4200)
Rwanda	1995	Female	80 to 84	54% (37-70%)	46% (30-63%)	2.2 (1.1-3.8)	8.7% (3.4-16%)	9.8% (3.8-18%)	97 (38-180)	1000 (390-1800)
Rwanda	1995	Female	85 to 89	53% (37-67%)	47% (33-63%)	2.1 (1-3.6)	7.1% (2.4-13%)	8.1% (2.8-15%)	47 (16-89)	340 (120-650)
Rwanda	1995	Female	90 to 94	54% (39-69%)	46% (31-61%)	2 (1-3.5)	5.5% (1.4-11%)	6.5% (1.5-14%)	14 (3.4-30)	71 (18-150)
Rwanda	1995	Female	95 plus	56% (37-74%)	44% (26-63%)	1.9 (0.85-3.6)	5.2% (0.95-11%)	6% (0.96-13%)	3.7 (0.61-8.2)	13 (2.3-28)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	1995	Male	All Ages	64% (59-69%)	36% (31-41%)	2.1 (1.6-2.9)	5.8% (4.1-7.6%)	9.2% (6.3-12%)	5500 (3700-7300)	240000 (170000-310000)
Rwanda	1995	Male	15 to 19	69% (63-74%)	31% (26-37%)	3.2 (2.1-4.8)	11% (6.2-17%)	13% (6.7-20%)	240 (130-350)	18000 (10000-26000)
Rwanda	1995	Male	20 to 24	72% (68-77%)	28% (23-32%)	3.6 (2.2-5.5)	14% (7.9-19%)	14% (7.8-21%)	370 (220-520)	26000 (16000-36000)
Rwanda	1995	Male	25 to 29	77% (73-80%)	23% (20-27%)	4.3 (2.7-6.4)	14% (8.6-20%)	15% (8.7-22%)	430 (270-600)	28000 (18000-38000)
Rwanda	1995	Male	30 to 34	80% (76-83%)	20% (17-24%)	4.8 (3-7.2)	15% (9.9-20%)	16% (10-22%)	480 (310-650)	28000 (19000-37000)
Rwanda	1995	Male	35 to 39	79% (76-82%)	21% (18-24%)	4.9 (3-7.3)	17% (11-23%)	18% (12-25%)	520 (350-710)	27000 (18000-36000)
Rwanda	1995	Male	40 to 44	81% (78-84%)	19% (16-22%)	5.1 (3.2-7.8)	19% (13-26%)	21% (14-28%)	520 (340-700)	24000 (16000-33000)
Rwanda	1995	Male	45 to 49	84% (80-86%)	16% (14-20%)	5.2 (3.1-7.9)	21% (14-28%)	23% (15-32%)	540 (340-740)	22000 (14000-30000)
Rwanda	1995	Male	50 to 54	84% (80-87%)	16% (13-20%)	5.1 (2.9-8)	24% (15-32%)	27% (17-36%)	510 (310-690)	18000 (11000-25000)
Rwanda	1995	Male	55 to 59	82% (77-87%)	18% (13-23%)	4.9 (2.8-8)	24% (15-32%)	27% (17-36%)	490 (290-680)	15000 (9200-21000)
Rwanda	1995	Male	60 to 64	79% (71-86%)	21% (14-29%)	4.3 (2.4-7.2)	21% (12-29%)	23% (13-32%)	530 (300-750)	14000 (8000-20000)
Rwanda	1995	Male	65 to 69	76% (61-87%)	24% (13-39%)	3.5 (2.1-5.7)	17% (8.7-25%)	19% (9.7-28%)	450 (230-670)	9900 (5100-15000)
Rwanda	1995	Male	70 to 74	73% (58-84%)	27% (16-42%)	2.8 (1.6-4.6)	13% (5.6-21%)	14% (6.3-23%)	240 (110-400)	4300 (1900-7100)
Rwanda	1995	Male	75 to 79	72% (58-84%)	28% (16-42%)	2.4 (1.2-4)	10% (3.6-18%)	12% (4-20%)	130 (46-230)	1800 (640-3200)
Rwanda	1995	Male	80 to 84	70% (54-83%)	30% (17-46%)	2.1 (1.2-3.5)	8.6% (3.4-15%)	9.7% (3.7-17%)	57 (22-100)	590 (230-1000)
Rwanda	1995	Male	85 to 89	68% (51-80%)	32% (20-49%)	1.8 (1-3)	6.5% (2.4-12%)	7.4% (2.6-13%)	24 (8.3-44)	180 (65-320)
Rwanda	1995	Male	90 to 94	67% (52-81%)	33% (19-48%)	1.6 (0.88-2.8)	4.3% (1.3-8%)	4.9% (1.2-9.3%)	6.5 (1.7-12)	34 (10-64)
Rwanda	1995	Male	95 plus	69% (48-85%)	31% (15-52%)	1.5 (0.67-2.8)	4.6% (0.81-9.5%)	5.1% (0.66-11%)	1.8 (0.23-3.8)	6.3 (1.1-13)
Rwanda	2000	Female	All Ages	41% (37-45%)	59% (55-63%)	1.6 (1.2-2.1)	3.2% (2-4.6%)	5.9% (3.4-8.5%)	3400 (2000-5100)	120000 (74000-170000)
Rwanda	2000	Female	15 to 19	43% (39-47%)	57% (53-61%)	2 (1.3-2.9)	6.7% (3.3-11%)	10% (4.6-18%)	110 (52-200)	8600 (4300-15000)
Rwanda	2000	Female	20 to 24	54% (50-58%)	46% (42-50%)	2.8 (1.8-4.2)	7.5% (4.1-12%)	9.7% (4.9-17%)	130 (69-220)	9700 (5300-15000)
Rwanda	2000	Female	25 to 29	58% (54-61%)	42% (39-46%)	3.4 (2.1-5.1)	6.3% (3.2-10%)	7.4% (3.5-13%)	150 (81-250)	9900 (5500-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	2000	Female	30 to 34	59% (56-63%)	41% (37-44%)	3.5 (2.1-5.3)	5.9% (3.2-9.4%)	6.8% (3.4-11%)	150 (82-250)	8900 (5000-14000)
Rwanda	2000	Female	35 to 39	60% (56-63%)	40% (37-44%)	3.1 (2-4.8)	6.3% (3.5-11%)	7.4% (3.8-13%)	210 (110-350)	11000 (6200-18000)
Rwanda	2000	Female	40 to 44	60% (56-63%)	40% (37-44%)	3 (1.8-4.7)	6.5% (3.5-11%)	7.7% (4.1-13%)	190 (99-310)	8800 (4800-14000)
Rwanda	2000	Female	45 to 49	58% (53-62%)	42% (38-47%)	3 (1.8-4.9)	8.9% (4.8-14%)	11% (5.7-17%)	240 (120-400)	10000 (5300-16000)
Rwanda	2000	Female	50 to 54	56% (46-66%)	44% (34-54%)	3 (1.7-5.1)	12% (6.3-18%)	14% (7.6-22%)	290 (150-450)	11000 (5400-16000)
Rwanda	2000	Female	55 to 59	55% (39-72%)	45% (28-61%)	3.1 (1.6-5.3)	13% (7-21%)	16% (8.3-24%)	330 (170-530)	10000 (5300-16000)
Rwanda	2000	Female	60 to 64	56% (39-73%)	44% (27-61%)	3.1 (1.6-5.3)	13% (6.5-19%)	15% (7.6-23%)	260 (130-410)	6900 (3600-11000)
Rwanda	2000	Female	65 to 69	57% (40-74%)	43% (26-60%)	2.8 (1.6-4.7)	12% (6-19%)	14% (7.1-21%)	460 (230-720)	10000 (5100-16000)
Rwanda	2000	Female	70 to 74	56% (39-71%)	44% (29-61%)	2.5 (1.3-4.2)	10% (4.2-17%)	12% (4.8-20%)	230 (95-390)	4100 (1700-6900)
Rwanda	2000	Female	75 to 79	53% (37-71%)	47% (29-63%)	2.2 (1.1-3.8)	8.7% (3.1-16%)	9.9% (3.5-18%)	430 (150-780)	6000 (2100-11000)
Rwanda	2000	Female	80 to 84	52% (35-69%)	48% (31-65%)	2 (1-3.4)	7% (2.4-13%)	8.1% (2.8-15%)	180 (63-330)	1900 (650-3400)
Rwanda	2000	Female	85 to 89	52% (36-66%)	48% (34-64%)	1.9 (0.95-3.2)	5.7% (1.9-11%)	6.5% (2.2-12%)	39 (13-74)	290 (93-540)
Rwanda	2000	Female	90 to 94	53% (37-68%)	47% (32-63%)	1.8 (0.93-3.3)	4.4% (1-9.1%)	5.2% (1.2-11%)	13 (2.9-27)	66 (15-140)
Rwanda	2000	Female	95 plus	55% (36-74%)	45% (26-64%)	1.8 (0.78-3.5)	4.3% (0.76-9.7%)	5% (0.78-11%)	4.1 (0.64-9.3)	14 (2.4-32)
Rwanda	2000	Male	All Ages	65% (60-69%)	35% (31-40%)	2.1 (1.6-2.8)	5.5% (4.1-7.2%)	8.5% (5.9-11%)	5200 (3600-6900)	220000 (160000-280000)
Rwanda	2000	Male	15 to 19	70% (65-74%)	30% (26-35%)	3.1 (2.1-4.6)	13% (8.1-19%)	17% (10-24%)	230 (130-330)	18000 (11000-25000)
Rwanda	2000	Male	20 to 24	73% (69-77%)	27% (23-31%)	3.5 (2.2-5.1)	18% (12-25%)	21% (13-30%)	340 (210-480)	25000 (16000-34000)
Rwanda	2000	Male	25 to 29	77% (73-80%)	23% (20-27%)	4.1 (2.5-6.3)	16% (11-23%)	18% (11-27%)	360 (230-510)	24000 (16000-33000)
Rwanda	2000	Male	30 to 34	80% (77-83%)	20% (17-23%)	4.6 (2.8-7.1)	14% (9.6-20%)	16% (10-22%)	390 (250-520)	23000 (15000-31000)
Rwanda	2000	Male	35 to 39	79% (76-82%)	21% (18-24%)	4.7 (3-7)	14% (9.3-19%)	15% (9.9-22%)	530 (360-720)	28000 (19000-38000)
Rwanda	2000	Male	40 to 44	81% (78-84%)	19% (16-22%)	4.9 (3-7.5)	15% (10-20%)	17% (11-22%)	470 (310-650)	22000 (15000-30000)
Rwanda	2000	Male	45 to 49	84% (81-86%)	16% (14-19%)	5 (3-7.6)	17% (11-23%)	19% (12-26%)	470 (310-670)	20000 (13000-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	2000	Male	50 to 54	84% (81-87%)	16% (13-19%)	4.9 (2.8-7.7)	19% (13-27%)	22% (15-30%)	430 (270-600)	16000 (10000-22000)
Rwanda	2000	Male	55 to 59	82% (77-87%)	18% (13-23%)	4.7 (2.8-7.5)	20% (13-28%)	24% (15-32%)	440 (270-610)	14000 (8700-19000)
Rwanda	2000	Male	60 to 64	79% (70-86%)	21% (14-30%)	4.1 (2.3-6.5)	17% (9.7-24%)	20% (11-28%)	310 (170-450)	8300 (4600-12000)
Rwanda	2000	Male	65 to 69	75% (60-87%)	25% (13-40%)	3.3 (1.9-5.1)	14% (7.4-20%)	16% (8.4-23%)	460 (240-700)	10000 (5300-15000)
Rwanda	2000	Male	70 to 74	73% (57-84%)	27% (16-43%)	2.6 (1.5-4.3)	10% (4.4-17%)	12% (5-19%)	190 (77-310)	3300 (1400-5500)
Rwanda	2000	Male	75 to 79	71% (55-83%)	29% (17-45%)	2.2 (1.2-3.8)	8.2% (2.8-15%)	9.3% (3.2-16%)	380 (130-680)	5300 (1800-9500)
Rwanda	2000	Male	80 to 84	69% (51-82%)	31% (18-49%)	2 (1.1-3.1)	6.8% (2.5-12%)	7.8% (2.9-14%)	130 (50-240)	1400 (530-2400)
Rwanda	2000	Male	85 to 89	67% (50-80%)	33% (20-50%)	1.7 (0.91-2.8)	5.2% (1.8-9.3%)	6% (2-11%)	23 (7.5-41)	170 (59-300)
Rwanda	2000	Male	90 to 94	67% (52-81%)	33% (19-48%)	1.5 (0.8-2.6)	3.5% (1-6.9%)	4% (1-7.9%)	6.3 (1.6-13)	33 (10-65)
Rwanda	2000	Male	95 plus	68% (48-84%)	32% (16-52%)	1.4 (0.63-2.7)	3.9% (0.83-8.4%)	4.4% (0.66-9.7%)	2.4 (0.35-5.3)	8.5 (1.8-18)
Rwanda	2005	Female	All Ages	40% (36-44%)	60% (56-64%)	1.6 (1.2-2.2)	2.3% (1.6-3.2%)	4.3% (2.7-6.1%)	2000 (1200-2900)	72000 (49000-98000)
Rwanda	2005	Female	15 to 19	41% (37-45%)	59% (55-63%)	1.9 (1.2-2.8)	4.6% (2.5-6.9%)	6.7% (3.4-11%)	72 (37-110)	5800 (3200-8800)
Rwanda	2005	Female	20 to 24	52% (48-56%)	48% (44-52%)	2.8 (1.7-4.1)	5.7% (3.4-8.6%)	7.7% (4.2-12%)	95 (52-150)	7400 (4400-11000)
Rwanda	2005	Female	25 to 29	56% (52-60%)	44% (40-48%)	3.3 (2-4.9)	5% (2.9-7.7%)	6.1% (3.1-10%)	92 (52-140)	6400 (3900-9500)
Rwanda	2005	Female	30 to 34	58% (54-62%)	42% (38-46%)	3.4 (2.1-5.2)	4.1% (2.4-6.2%)	4.7% (2.6-7.6%)	90 (53-140)	5700 (3400-8400)
Rwanda	2005	Female	35 to 39	58% (54-63%)	42% (37-46%)	3.1 (1.9-4.8)	4.1% (2.4-6.3%)	4.7% (2.6-7.5%)	89 (51-140)	4900 (2900-7500)
Rwanda	2005	Female	40 to 44	58% (54-63%)	42% (37-46%)	2.9 (1.7-4.7)	4% (2.2-6%)	4.7% (2.5-7.3%)	110 (59-170)	5300 (3000-8100)
Rwanda	2005	Female	45 to 49	56% (51-61%)	44% (39-49%)	2.9 (1.6-4.8)	5.9% (3.3-9.1%)	7.4% (4.1-11%)	140 (79-220)	5900 (3400-9300)
Rwanda	2005	Female	50 to 54	55% (44-65%)	45% (35-56%)	2.9 (1.6-4.9)	8.2% (4.5-12%)	10% (5.7-16%)	180 (100-280)	6700 (3700-10000)
Rwanda	2005	Female	55 to 59	54% (38-71%)	46% (29-62%)	3 (1.6-5.2)	9.6% (5.2-15%)	12% (6.6-19%)	190 (100-300)	6100 (3300-9600)
Rwanda	2005	Female	60 to 64	54% (37-73%)	46% (27-63%)	3 (1.5-5.2)	9.4% (4.7-15%)	12% (6-19%)	200 (100-320)	5400 (2800-8800)
Rwanda	2005	Female	65 to 69	55% (39-72%)	45% (28-61%)	2.8 (1.5-4.5)	9.3% (4.7-14%)	11% (5.8-17%)	170 (83-270)	3800 (1900-5900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	2005	Female	70 to 74	54% (37-70%)	46% (30-63%)	2.4 (1.2-4.2)	8% (3.2-13%)	9.5% (3.9-16%)	250 (110-430)	4600 (2000-7800)
Rwanda	2005	Female	75 to 79	52% (36-68%)	48% (32-64%)	2.1 (1-3.7)	6.7% (2.4-12%)	8% (2.8-15%)	110 (37-210)	1500 (520-2900)
Rwanda	2005	Female	80 to 84	51% (33-68%)	49% (32-67%)	2 (1-3.3)	5.4% (1.9-10%)	6.4% (2.2-12%)	140 (49-270)	1500 (500-2800)
Rwanda	2005	Female	85 to 89	50% (34-66%)	50% (34-66%)	1.8 (0.93-3.2)	4.3% (1.4-8.4%)	5.1% (1.6-9.7%)	57 (18-110)	410 (130-790)
Rwanda	2005	Female	90 to 94	52% (36-67%)	48% (33-64%)	1.8 (0.88-3.3)	3.5% (0.73-7.8%)	4.1% (0.8-9.1%)	8.1 (1.6-18)	41 (8.7-90)
Rwanda	2005	Female	95 plus	54% (35-74%)	46% (26-65%)	1.7 (0.76-3.5)	3.5% (0.59-8%)	4% (0.58-9.5%)	2.5 (0.36-6.1)	8.6 (1.5-20)
Rwanda	2005	Male	All Ages	63% (58-68%)	37% (32-42%)	2.1 (1.5-2.8)	4.4% (3.3-5.7%)	6.6% (4.7-8.8%)	3100 (2200-4100)	140000 (100000-180000)
Rwanda	2005	Male	15 to 19	68% (63-72%)	32% (28-37%)	3.1 (2-4.5)	9.3% (6.3-13%)	12% (7.3-17%)	140 (88-200)	12000 (8000-17000)
Rwanda	2005	Male	20 to 24	72% (67-76%)	28% (24-33%)	3.4 (2.1-5)	16% (11-21%)	18% (11-26%)	230 (140-330)	19000 (13000-26000)
Rwanda	2005	Male	25 to 29	76% (72-79%)	24% (21-28%)	4 (2.5-6.1)	16% (10-22%)	18% (11-26%)	240 (150-360)	18000 (12000-24000)
Rwanda	2005	Male	30 to 34	79% (75-82%)	21% (18-25%)	4.5 (2.8-6.9)	13% (8.7-18%)	14% (9.1-20%)	230 (150-320)	15000 (10000-20000)
Rwanda	2005	Male	35 to 39	78% (74-81%)	22% (19-26%)	4.6 (2.9-7.1)	11% (7.9-16%)	13% (8.4-18%)	230 (150-320)	13000 (8700-17000)
Rwanda	2005	Male	40 to 44	80% (76-83%)	20% (17-24%)	4.8 (2.9-7.5)	11% (7.6-15%)	13% (8.2-18%)	300 (190-420)	15000 (9700-20000)
Rwanda	2005	Male	45 to 49	83% (79-86%)	17% (14-21%)	4.9 (2.8-7.6)	13% (8.3-18%)	15% (9.2-20%)	270 (180-370)	12000 (7700-16000)
Rwanda	2005	Male	50 to 54	83% (80-86%)	17% (14-20%)	4.8 (2.8-7.6)	15% (9.8-21%)	18% (11-25%)	300 (190-410)	11000 (7300-15000)
Rwanda	2005	Male	55 to 59	82% (76-86%)	18% (14-24%)	4.6 (2.7-7.1)	16% (10-22%)	20% (13-27%)	260 (170-370)	8600 (5500-12000)
Rwanda	2005	Male	60 to 64	78% (68-86%)	22% (14-32%)	4 (2.3-6.6)	14% (8-20%)	16% (9.4-23%)	240 (140-350)	6600 (3900-9500)
Rwanda	2005	Male	65 to 69	74% (58-86%)	26% (14-42%)	3.2 (1.9-5.2)	11% (5.9-16%)	13% (6.9-19%)	160 (86-240)	3600 (2000-5600)
Rwanda	2005	Male	70 to 74	72% (55-84%)	28% (16-45%)	2.5 (1.4-4.2)	7.8% (3.6-13%)	9.1% (4.1-15%)	210 (98-360)	3900 (1800-6500)
Rwanda	2005	Male	75 to 79	70% (54-83%)	30% (17-46%)	2.1 (1.1-3.6)	6.3% (2.3-12%)	7.3% (2.7-14%)	83 (31-160)	1200 (440-2200)
Rwanda	2005	Male	80 to 84	68% (50-81%)	32% (19-50%)	1.9 (1-3.1)	5.1% (1.9-9.3%)	5.9% (2.1-11%)	110 (42-210)	1200 (450-2200)
Rwanda	2005	Male	85 to 89	65% (48-79%)	35% (21-52%)	1.6 (0.87-2.6)	3.9% (1.3-7.1%)	4.4% (1.4-8.2%)	41 (13-75)	310 (110-560)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	2005	Male	90 to 94	65% (47-79%)	35% (21-53%)	1.5 (0.76-2.5)	2.7% (0.69-5.1%)	2.9% (0.59-5.9%)	4 (0.82-8.1)	22 (5.5-43)
Rwanda	2005	Male	95 plus	66% (45-83%)	34% (17-55%)	1.4 (0.61-2.6)	3% (0.74-6.5%)	3.2% (0.57-7.2%)	1.3 (0.22-2.8)	4.7 (1.1-9.9)
Rwanda	2010	Female	All Ages	41% (36-47%)	59% (53-64%)	1.6 (1.1-2.3)	2.7% (1.7-3.9%)	5% (2.7-7.6%)	1800 (960-2800)	63000 (39000-91000)
Rwanda	2010	Female	15 to 19	39% (30-50%)	61% (50-70%)	1.8 (0.98-2.9)	3.4% (1.7-5.5%)	4.9% (2.1-8.3%)	41 (19-68)	3500 (1800-5600)
Rwanda	2010	Female	20 to 24	51% (40-62%)	49% (38-60%)	2.6 (1.5-4.2)	4.8% (2.6-7.4%)	6.5% (3.2-11%)	67 (33-110)	5700 (3100-9000)
Rwanda	2010	Female	25 to 29	55% (43-66%)	45% (34-57%)	3.2 (1.8-5.1)	4.9% (2.8-7.5%)	6.5% (3.2-11%)	82 (44-130)	6200 (3700-9300)
Rwanda	2010	Female	30 to 34	58% (47-67%)	42% (33-53%)	3.3 (1.9-5.4)	4.7% (2.7-7.2%)	6% (3.2-9.8%)	75 (41-120)	5000 (2900-7400)
Rwanda	2010	Female	35 to 39	61% (50-69%)	39% (31-50%)	3.1 (1.7-5.3)	4.9% (2.8-7.7%)	6.4% (3.4-10%)	80 (44-130)	4600 (2600-7200)
Rwanda	2010	Female	40 to 44	62% (51-69%)	38% (31-49%)	3 (1.6-5)	5.1% (2.7-8%)	6.8% (3.5-11%)	75 (39-120)	3800 (2000-6000)
Rwanda	2010	Female	45 to 49	60% (49-68%)	40% (32-51%)	3.1 (1.6-5.2)	6.8% (3.6-11%)	9.5% (4.9-15%)	130 (71-210)	5800 (3100-9300)
Rwanda	2010	Female	50 to 54	57% (46-67%)	43% (33-54%)	3 (1.6-5.3)	8.3% (4.2-13%)	12% (6.1-17%)	160 (83-250)	6100 (3100-9600)
Rwanda	2010	Female	55 to 59	54% (38-71%)	46% (29-62%)	3 (1.5-5.5)	9% (4.4-14%)	12% (6.1-19%)	190 (90-310)	6100 (2900-10000)
Rwanda	2010	Female	60 to 64	55% (38-73%)	45% (27-62%)	2.9 (1.4-5.3)	8.6% (4-14%)	11% (5.3-18%)	180 (84-310)	5100 (2300-8500)
Rwanda	2010	Female	65 to 69	56% (39-72%)	44% (28-61%)	2.7 (1.4-4.7)	8.4% (3.9-14%)	11% (4.9-17%)	200 (93-340)	4500 (2100-7800)
Rwanda	2010	Female	70 to 74	55% (38-70%)	45% (30-62%)	2.4 (1.2-4.3)	7.1% (2.4-13%)	8.5% (3-16%)	140 (47-260)	2500 (850-4700)
Rwanda	2010	Female	75 to 79	52% (36-69%)	48% (31-64%)	2.1 (0.96-3.8)	5.9% (1.6-12%)	7.1% (1.9-14%)	190 (49-370)	2600 (690-5200)
Rwanda	2010	Female	80 to 84	51% (34-68%)	49% (32-66%)	1.9 (0.97-3.4)	4.8% (1.1-9.5%)	5.7% (1.3-11%)	110 (25-220)	1100 (260-2300)
Rwanda	2010	Female	85 to 89	51% (35-65%)	49% (35-65%)	1.8 (0.89-3.3)	3.8% (0.91-8%)	4.5% (1.1-9.6%)	60 (15-130)	440 (110-960)
Rwanda	2010	Female	90 to 94	52% (36-68%)	48% (32-64%)	1.7 (0.85-3.3)	3% (0.32-7.2%)	3.5% (0.27-8.4%)	17 (1.3-41)	86 (8.8-210)
Rwanda	2010	Female	95 plus	54% (35-74%)	46% (26-65%)	1.7 (0.73-3.4)	3% (0.28-7.8%)	3.5% (0.18-9.4%)	2 (0.1-5.1)	6.7 (0.6-17)
Rwanda	2010	Male	All Ages	65% (60-69%)	35% (31-40%)	2.1 (1.4-3)	5.1% (3.7-6.7%)	7.3% (4.8-10%)	2700 (1800-3700)	130000 (90000-160000)
Rwanda	2010	Male	15 to 19	69% (62-74%)	31% (26-38%)	3 (1.7-4.7)	7.8% (5-11%)	9.3% (5.2-14%)	91 (51-140)	8400 (5300-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	2010	Male	20 to 24	73% (68-77%)	27% (23-32%)	3.4 (1.9-5.4)	14% (9.2-19%)	15% (8.9-22%)	200 (110-300)	18000 (12000-25000)
Rwanda	2010	Male	25 to 29	77% (72-80%)	23% (20-28%)	3.9 (2.3-6.2)	15% (10-20%)	17% (10-25%)	230 (140-340)	18000 (12000-25000)
Rwanda	2010	Male	30 to 34	80% (75-83%)	20% (17-25%)	4.4 (2.6-6.9)	14% (9.5-19%)	16% (10-24%)	210 (130-300)	14000 (9600-19000)
Rwanda	2010	Male	35 to 39	79% (76-82%)	21% (18-24%)	4.6 (2.6-7.7)	13% (8.7-18%)	16% (9.7-22%)	190 (120-280)	11000 (7300-15000)
Rwanda	2010	Male	40 to 44	81% (77-84%)	19% (16-23%)	4.7 (2.7-7.6)	13% (8.6-18%)	16% (10-22%)	190 (120-270)	9500 (6000-13000)
Rwanda	2010	Male	45 to 49	83% (80-86%)	17% (14-20%)	4.8 (2.6-7.9)	14% (9.1-20%)	18% (11-25%)	270 (160-390)	12000 (7400-17000)
Rwanda	2010	Male	50 to 54	84% (80-87%)	16% (13-20%)	4.7 (2.6-7.9)	16% (9.9-22%)	20% (12-28%)	260 (160-370)	10000 (6200-14000)
Rwanda	2010	Male	55 to 59	82% (77-87%)	18% (13-23%)	4.5 (2.5-7.4)	16% (9.8-22%)	20% (12-28%)	280 (170-390)	9100 (5600-13000)
Rwanda	2010	Male	60 to 64	79% (70-86%)	21% (14-30%)	3.9 (2-6.8)	13% (7.1-19%)	16% (8.5-23%)	220 (120-330)	6100 (3300-9100)
Rwanda	2010	Male	65 to 69	74% (57-87%)	26% (13-43%)	3.1 (1.7-5.3)	9.6% (4.9-15%)	12% (5.8-19%)	180 (87-290)	4100 (2000-6700)
Rwanda	2010	Male	70 to 74	71% (55-84%)	29% (16-45%)	2.5 (1.3-4.2)	6.9% (2.9-12%)	8.2% (3.5-15%)	110 (45-190)	1900 (840-3500)
Rwanda	2010	Male	75 to 79	70% (54-82%)	30% (18-46%)	2.1 (1.1-3.7)	5.5% (1.6-11%)	6.4% (1.9-13%)	140 (41-280)	2000 (610-4000)
Rwanda	2010	Male	80 to 84	68% (50-82%)	32% (18-50%)	1.9 (0.98-3.2)	4.5% (1.4-8.5%)	5.2% (1.6-10%)	77 (23-150)	810 (250-1600)
Rwanda	2010	Male	85 to 89	65% (49-79%)	35% (21-51%)	1.6 (0.84-2.8)	3.4% (1.1-6.6%)	3.9% (1.1-7.6%)	44 (12-87)	330 (100-650)
Rwanda	2010	Male	90 to 94	65% (49-79%)	35% (21-51%)	1.4 (0.73-2.6)	2.3% (0.53-5%)	2.5% (0.38-5.7%)	10 (1.5-23)	56 (13-120)
Rwanda	2010	Male	95 plus	67% (46-83%)	33% (17-54%)	1.3 (0.6-2.6)	2.6% (0.6-5.9%)	2.8% (0.29-6.7%)	1 (0.11-2.4)	3.9 (0.88-8.6)
Rwanda	2016	Female	All Ages	42% (37-47%)	58% (53-63%)	1.7 (0.93-2.9)	3.3% (1.7-5.1%)	5.8% (2.6-9.7%)	2000 (850-3400)	69000 (36000-110000)
Rwanda	2016	Female	15 to 19	40% (30-49%)	60% (51-70%)	1.8 (0.79-3.4)	3.4% (1.5-5.9%)	5.5% (2.1-10%)	39 (14-72)	3500 (1500-6000)
Rwanda	2016	Female	20 to 24	51% (39-61%)	49% (39-61%)	2.6 (1.2-4.9)	5% (2.6-8%)	7.5% (3.2-13%)	50 (21-88)	4500 (2300-7200)
Rwanda	2016	Female	25 to 29	55% (43-65%)	45% (35-57%)	3.2 (1.5-5.7)	5.7% (3.2-8.9%)	8.8% (4.3-15%)	77 (38-130)	6000 (3300-9400)
Rwanda	2016	Female	30 to 34	58% (47-67%)	42% (33-53%)	3.4 (1.7-6.2)	5.8% (3.2-9%)	8.9% (4.4-15%)	95 (47-160)	6500 (3400-10000)
Rwanda	2016	Female	35 to 39	61% (50-68%)	39% (32-50%)	3.2 (1.5-6)	6% (3.1-9.7%)	9.2% (4.5-15%)	97 (46-160)	5700 (2900-9100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	2016	Female	40 to 44	62% (52-69%)	38% (31-48%)	3.1 (1.4-5.7)	5.7% (2.7-9.1%)	8.4% (4-13%)	95 (44-160)	4900 (2300-7900)
Rwanda	2016	Female	45 to 49	60% (48-68%)	40% (32-52%)	3.1 (1.4-6)	6.8% (3.1-11%)	9.9% (4.7-16%)	120 (53-190)	5100 (2300-8200)
Rwanda	2016	Female	50 to 54	57% (45-66%)	43% (34-55%)	3 (1.3-5.8)	7.8% (3.6-12%)	11% (5.1-17%)	190 (86-310)	7000 (3200-12000)
Rwanda	2016	Female	55 to 59	54% (39-71%)	46% (29-61%)	2.9 (1.3-5.8)	8.3% (3.4-14%)	11% (4.8-19%)	210 (90-370)	6900 (2900-12000)
Rwanda	2016	Female	60 to 64	55% (38-72%)	45% (28-62%)	2.9 (1.2-5.5)	8% (3.3-13%)	10% (4.3-18%)	230 (92-400)	6300 (2400-11000)
Rwanda	2016	Female	65 to 69	56% (39-72%)	44% (28-61%)	2.7 (1.2-5.2)	7.9% (2.9-14%)	9.9% (3.8-17%)	230 (90-410)	5200 (2000-9400)
Rwanda	2016	Female	70 to 74	55% (38-71%)	45% (29-62%)	2.4 (1.1-5)	6.6% (1.7-13%)	8% (2.1-16%)	200 (53-410)	3700 (950-7600)
Rwanda	2016	Female	75 to 79	53% (36-70%)	47% (30-64%)	2.1 (0.86-4.4)	5.6% (1.1-12%)	6.8% (1.4-14%)	130 (28-290)	1900 (370-4100)
Rwanda	2016	Female	80 to 84	51% (34-69%)	49% (31-66%)	1.9 (0.81-3.9)	4.6% (0.83-10%)	5.5% (0.99-12%)	110 (21-250)	1200 (210-2700)
Rwanda	2016	Female	85 to 89	51% (35-66%)	49% (34-65%)	1.8 (0.79-3.8)	3.8% (0.48-9.5%)	4.5% (0.56-11%)	82 (11-210)	610 (76-1500)
Rwanda	2016	Female	90 to 94	52% (36-68%)	48% (32-64%)	1.8 (0.74-3.7)	3.1% (0.0068-8.4%)	3.6% (-0.025-10%)	23 (-0.16-65)	120 (0.26-330)
Rwanda	2016	Female	95 plus	54% (35-74%)	46% (26-65%)	1.7 (0.63-4.2)	3.2% (0.14-8.9%)	3.6% (0.055-11%)	9.5 (0.14-27)	33 (1.4-92)
Rwanda	2016	Male	All Ages	61% (55-66%)	39% (34-45%)	2.1 (1.1-3.5)	6.3% (3.9-8.8%)	8.6% (4.8-13%)	3000 (1600-4600)	140000 (89000-200000)
Rwanda	2016	Male	15 to 19	57% (45-67%)	43% (33-55%)	2.5 (1.1-4.7)	7.6% (4.4-12%)	9% (4.2-15%)	91 (41-160)	8900 (4900-14000)
Rwanda	2016	Male	20 to 24	67% (56-75%)	33% (25-44%)	3.1 (1.5-5.9)	14% (8.5-20%)	15% (7.4-24%)	160 (77-260)	15000 (8900-22000)
Rwanda	2016	Male	25 to 29	74% (65-80%)	26% (20-35%)	3.8 (1.9-7)	16% (10-22%)	19% (10-27%)	260 (140-410)	21000 (13000-30000)
Rwanda	2016	Male	30 to 34	77% (69-83%)	23% (17-31%)	4.2 (2.1-7.5)	16% (9.9-22%)	19% (11-28%)	280 (150-420)	19000 (12000-28000)
Rwanda	2016	Male	35 to 39	77% (69-82%)	23% (18-31%)	4.4 (2.1-7.8)	15% (9.2-21%)	19% (11-28%)	260 (140-390)	15000 (8800-22000)
Rwanda	2016	Male	40 to 44	78% (72-83%)	22% (17-28%)	4.5 (2-8)	15% (8.7-21%)	19% (11-27%)	220 (120-330)	12000 (6600-16000)
Rwanda	2016	Male	45 to 49	81% (73-85%)	19% (15-27%)	4.6 (2.1-8.6)	15% (8.5-21%)	19% (10-28%)	220 (120-330)	9800 (5400-14000)
Rwanda	2016	Male	50 to 54	82% (75-86%)	18% (14-25%)	4.5 (2.1-8.5)	15% (8.2-21%)	19% (10-27%)	310 (160-480)	12000 (6700-18000)
Rwanda	2016	Male	55 to 59	80% (73-86%)	20% (14-27%)	4.3 (2.1-7.9)	15% (8.2-21%)	19% (10-27%)	300 (160-450)	10000 (5500-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Rwanda	2016	Male	60 to 64	78% (67-86%)	22% (14-33%)	3.9 (1.7-7.3)	12% (5.9-19%)	15% (7.2-23%)	270 (120-430)	7600 (3600-12000)
Rwanda	2016	Male	65 to 69	75% (59-86%)	25% (14-41%)	3.1 (1.5-5.8)	9.2% (3.8-16%)	11% (4.5-20%)	200 (80-350)	4500 (1900-8100)
Rwanda	2016	Male	70 to 74	72% (56-84%)	28% (16-44%)	2.5 (1.2-5.1)	6.6% (2.2-13%)	7.9% (2.6-16%)	150 (47-300)	2700 (890-5400)
Rwanda	2016	Male	75 to 79	70% (55-83%)	30% (17-45%)	2.1 (0.94-4.4)	5.3% (1.2-12%)	6.4% (1.4-14%)	88 (19-200)	1300 (290-2800)
Rwanda	2016	Male	80 to 84	68% (49-81%)	32% (19-51%)	1.9 (0.84-3.7)	4.4% (0.94-10%)	5.1% (1-12%)	74 (15-170)	790 (170-1800)
Rwanda	2016	Male	85 to 89	65% (48-79%)	35% (21-52%)	1.6 (0.74-3.3)	3.4% (0.72-7.9%)	3.8% (0.72-9.1%)	50 (9.4-120)	390 (85-900)
Rwanda	2016	Male	90 to 94	65% (49-79%)	35% (21-51%)	1.4 (0.65-2.9)	2.4% (0.36-6.2%)	2.6% (0.17-6.9%)	12 (0.78-32)	67 (10-170)
Rwanda	2016	Male	95 plus	67% (47-83%)	33% (17-53%)	1.3 (0.5-2.9)	2.6% (0.47-7%)	2.7% (0.22-7.8%)	5.3 (0.45-15)	20 (3.8-54)
Seychelles	1990	Female	All Ages	8.2% (6-11%)	92% (89-94%)	0.41 (0.3-0.54)	2.3% (1.6-3.2%)	4% (2.3-6.3%)	9.8 (5.4-15)	200 (130-290)
Seychelles	1990	Female	15 to 19	7.7% (3.5-14%)	92% (86-96%)	0.33 (0.12-0.71)	1% (0.57-1.7%)	1.5% (0.72-3.1%)	0.033 (0.015-0.067)	4.2 (2.4-7.1)
Seychelles	1990	Female	20 to 24	9.2% (3.8-18%)	91% (82-96%)	0.45 (0.16-0.92)	1.5% (0.92-2.4%)	1.9% (0.84-3.6%)	0.046 (0.02-0.091)	6.6 (3.8-10)
Seychelles	1990	Female	25 to 29	9.8% (4.2-19%)	90% (81-96%)	0.5 (0.19-1)	1.7% (1.1-2.6%)	2.2% (1.1-4.1%)	0.068 (0.033-0.13)	7.9 (4.8-12)
Seychelles	1990	Female	30 to 34	11% (4.5-20%)	89% (80-96%)	0.57 (0.21-1.1)	1.9% (1.2-3%)	2.8% (1.5-4.9%)	0.097 (0.048-0.18)	8.5 (5.1-14)
Seychelles	1990	Female	35 to 39	12% (4.8-22%)	88% (78-95%)	0.64 (0.24-1.3)	2.2% (1.3-3.6%)	3.4% (1.7-5.9%)	0.13 (0.062-0.23)	8.7 (4.9-14)
Seychelles	1990	Female	40 to 44	12% (5.4-21%)	88% (79-95%)	0.72 (0.26-1.5)	2.6% (1.4-4.4%)	4% (2-7.1%)	0.1 (0.05-0.18)	5.6 (3-9.5)
Seychelles	1990	Female	45 to 49	12% (5.2-23%)	88% (77-95%)	0.83 (0.27-1.9)	3% (1.5-5.2%)	4.4% (2.2-7.8%)	0.19 (0.095-0.35)	9.2 (4.7-16)
Seychelles	1990	Female	50 to 54	13% (5.6-24%)	87% (76-94%)	0.93 (0.32-2)	3.5% (1.7-6.1%)	4.9% (2.3-8.8%)	0.39 (0.19-0.7)	16 (7.8-28)
Seychelles	1990	Female	55 to 59	13% (5.7-24%)	87% (76-94%)	1 (0.33-2.1)	3.7% (1.7-6.6%)	5.1% (2.3-8.9%)	0.53 (0.24-0.94)	18 (8.2-31)
Seychelles	1990	Female	60 to 64	13% (5.4-24%)	87% (76-95%)	1 (0.32-2.2)	3.7% (1.7-6.9%)	4.9% (2.2-8.9%)	0.75 (0.35-1.4)	21 (9.9-39)
Seychelles	1990	Female	65 to 69	12% (5.5-23%)	88% (77-95%)	0.94 (0.33-2)	3.9% (1.8-7.2%)	5.1% (2.3-9.1%)	1 (0.47-1.8)	24 (11-42)
Seychelles	1990	Female	70 to 74	12% (5.4-22%)	88% (78-95%)	0.85 (0.28-1.9)	3.7% (1.5-7.2%)	4.7% (1.9-9.2%)	1.1 (0.45-2.1)	20 (8.3-39)
Seychelles	1990	Female	75 to 79	11% (4.8-22%)	89% (78-95%)	0.79 (0.23-1.8)	3.4% (1.2-6.6%)	4.1% (1.5-8.1%)	1.4 (0.51-2.8)	20 (7.2-39)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	1990	Female	80 to 84	11% (5-22%)	89% (78-95%)	0.79 (0.25-1.8)	3.4% (1.3-6.6%)	4.2% (1.5-8%)	1.7 (0.63-3.4)	18 (6.5-35)
Seychelles	1990	Female	85 to 89	12% (5.3-22%)	88% (78-95%)	0.83 (0.25-1.9)	3.3% (1.2-6.8%)	4% (1.4-8.3%)	1.3 (0.45-2.7)	9.8 (3.3-20)
Seychelles	1990	Female	90 to 94	14% (5.7-26%)	86% (74-94%)	0.95 (0.25-2.3)	3.4% (1.1-7%)	4% (1.2-8.4%)	0.64 (0.19-1.3)	3.3 (1-6.8)
Seychelles	1990	Female	95 plus	17% (6.6-34%)	83% (66-93%)	1.1 (0.24-3.2)	3.8% (1.1-8.5%)	4.5% (1.3-10%)	0.23 (0.064-0.51)	0.76 (0.22-1.7)
Seychelles	1990	Male	All Ages	25% (19-32%)	75% (68-81%)	1.2 (1-1.4)	6.9% (5.2-8.5%)	7.9% (5.4-10%)	25 (17-34)	870 (660-1100)
Seychelles	1990	Male	15 to 19	21% (12-36%)	79% (64-88%)	0.6 (0.27-1.1)	2.5% (1.5-4%)	3.3% (1.7-5.6%)	0.19 (0.1-0.33)	17 (9.7-27)
Seychelles	1990	Male	20 to 24	33% (18-51%)	67% (49-82%)	1.4 (0.73-2.3)	6.2% (3.7-9.5%)	7.5% (4.1-12%)	0.63 (0.35-1)	48 (29-73)
Seychelles	1990	Male	25 to 29	38% (20-56%)	62% (44-80%)	2 (1.1-3.1)	8.4% (5.4-12%)	10% (6.1-15%)	1 (0.63-1.6)	72 (46-110)
Seychelles	1990	Male	30 to 34	41% (22-60%)	59% (40-78%)	2.5 (1.4-3.7)	10% (6.8-14%)	12% (7.9-17%)	1.5 (0.92-2.1)	90 (59-130)
Seychelles	1990	Male	35 to 39	44% (26-63%)	56% (37-74%)	2.8 (1.5-4.5)	12% (7.6-17%)	14% (8.9-20%)	1.8 (1.1-2.6)	96 (60-140)
Seychelles	1990	Male	40 to 44	43% (26-64%)	57% (36-74%)	3 (1.6-4.9)	13% (7.9-18%)	15% (9.2-21%)	1.2 (0.76-1.8)	60 (37-85)
Seychelles	1990	Male	45 to 49	42% (23-61%)	58% (39-77%)	2.9 (1.5-4.9)	13% (7.8-18%)	15% (9.2-21%)	1.8 (1.1-2.6)	78 (48-110)
Seychelles	1990	Male	50 to 54	40% (22-59%)	60% (41-78%)	2.8 (1.4-4.6)	12% (7-17%)	14% (8-20%)	2.3 (1.3-3.3)	86 (49-120)
Seychelles	1990	Male	55 to 59	37% (21-58%)	63% (42-79%)	2.5 (1.2-4.1)	11% (6.1-16%)	12% (7-18%)	2.9 (1.7-4.4)	95 (54-140)
Seychelles	1990	Male	60 to 64	34% (19-54%)	66% (46-81%)	2.1 (0.99-3.8)	8.9% (4.6-14%)	10% (5.2-16%)	3 (1.5-4.8)	82 (42-130)
Seychelles	1990	Male	65 to 69	31% (17-49%)	69% (51-83%)	1.8 (0.84-3.1)	7.4% (3.8-12%)	8.4% (4.3-13%)	2.7 (1.4-4.4)	62 (31-99)
Seychelles	1990	Male	70 to 74	28% (15-44%)	72% (56-85%)	1.5 (0.67-2.7)	5.8% (2.7-10%)	6.6% (3-12%)	2 (0.92-3.7)	38 (17-68)
Seychelles	1990	Male	75 to 79	25% (14-42%)	75% (58-86%)	1.2 (0.52-2.3)	4.4% (1.8-8.3%)	4.9% (1.9-9.4%)	1.5 (0.59-2.9)	22 (8.7-41)
Seychelles	1990	Male	80 to 84	23% (11-40%)	77% (60-89%)	1.1 (0.45-2)	3.6% (1.5-6.9%)	4.1% (1.6-7.9%)	1.3 (0.5-2.5)	14 (5.5-26)
Seychelles	1990	Male	85 to 89	22% (11-36%)	78% (64-89%)	0.94 (0.4-1.8)	2.8% (1-5.6%)	3.1% (1.1-6.4%)	0.62 (0.22-1.3)	4.8 (1.8-9.6)
Seychelles	1990	Male	90 to 94	23% (12-38%)	77% (62-88%)	0.91 (0.4-1.8)	2.3% (0.79-5.2%)	2.6% (0.83-5.9%)	0.19 (0.061-0.43)	1 (0.34-2.3)
Seychelles	1990	Male	95 plus	26% (12-45%)	74% (55-88%)	0.94 (0.32-2)	2.1% (0.5-5.6%)	2.3% (0.42-6%)	0.041 (0.0075-0.11)	0.15 (0.036-0.38)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	1995	Female	All Ages	8.3% (6-11%)	92% (89-94%)	0.52 (0.39-0.69)	2.6% (1.8-3.6%)	4.5% (2.7-6.9%)	11 (6.4-17)	230 (160-320)
Seychelles	1995	Female	15 to 19	7.4% (3.3-14%)	93% (86-97%)	0.4 (0.13-0.9)	1.1% (0.61-1.9%)	1.8% (0.83-3.7%)	0.035 (0.016-0.072)	4.5 (2.5-8)
Seychelles	1995	Female	20 to 24	9% (3.8-17%)	91% (83-96%)	0.53 (0.18-1.1)	1.7% (1-2.6%)	2.2% (0.95-4.2%)	0.049 (0.021-0.092)	7.3 (4.3-12)
Seychelles	1995	Female	25 to 29	9.7% (4.2-18%)	90% (82-96%)	0.6 (0.22-1.3)	1.9% (1.2-2.8%)	2.6% (1.2-4.7%)	0.063 (0.028-0.12)	7.9 (4.8-12)
Seychelles	1995	Female	30 to 34	11% (4.2-21%)	89% (79-96%)	0.68 (0.26-1.4)	2.1% (1.3-3.1%)	3.2% (1.7-5.5%)	0.1 (0.051-0.18)	9.7 (5.9-15)
Seychelles	1995	Female	35 to 39	12% (4.5-22%)	88% (78-95%)	0.77 (0.3-1.5)	2.3% (1.4-3.6%)	3.9% (2.1-6.6%)	0.15 (0.079-0.27)	11 (6.5-18)
Seychelles	1995	Female	40 to 44	12% (5.2-22%)	88% (78-95%)	0.87 (0.33-1.7)	2.7% (1.5-4.6%)	4.6% (2.4-7.7%)	0.2 (0.1-0.36)	12 (6.6-20)
Seychelles	1995	Female	45 to 49	12% (4.9-22%)	88% (78-95%)	1 (0.33-2.2)	3.2% (1.7-5.4%)	5% (2.6-8.5%)	0.17 (0.085-0.3)	8.2 (4.4-14)
Seychelles	1995	Female	50 to 54	12% (5.2-24%)	88% (76-95%)	1.1 (0.39-2.5)	3.7% (2-6.4%)	5.5% (2.8-9.6%)	0.31 (0.16-0.56)	13 (6.8-22)
Seychelles	1995	Female	55 to 59	12% (5.5-23%)	88% (77-95%)	1.2 (0.4-2.6)	4% (2-6.9%)	5.7% (2.8-9.8%)	0.58 (0.29-1)	20 (10-34)
Seychelles	1995	Female	60 to 64	12% (5.3-23%)	88% (77-95%)	1.2 (0.39-2.7)	4.1% (2-7.4%)	5.5% (2.7-10%)	0.8 (0.4-1.5)	23 (11-42)
Seychelles	1995	Female	65 to 69	12% (5.3-22%)	88% (78-95%)	1.1 (0.38-2.3)	4.2% (2-7.4%)	5.6% (2.7-9.8%)	1.2 (0.55-2)	27 (13-48)
Seychelles	1995	Female	70 to 74	11% (5-21%)	89% (79-95%)	1 (0.31-2.3)	4% (1.8-7.4%)	5.1% (2.3-9.7%)	1.4 (0.62-2.6)	26 (12-49)
Seychelles	1995	Female	75 to 79	11% (4.6-22%)	89% (78-95%)	0.94 (0.27-2.2)	3.6% (1.5-7%)	4.6% (1.8-8.8%)	1.3 (0.54-2.7)	19 (7.7-38)
Seychelles	1995	Female	80 to 84	11% (4.7-21%)	89% (79-95%)	0.93 (0.28-2.2)	3.7% (1.5-7.2%)	4.6% (1.8-9%)	1.8 (0.71-3.5)	19 (7.4-37)
Seychelles	1995	Female	85 to 89	12% (5.3-22%)	88% (78-95%)	0.99 (0.31-2.3)	3.7% (1.4-7.3%)	4.5% (1.7-8.8%)	1.7 (0.6-3.2)	12 (4.6-24)
Seychelles	1995	Female	90 to 94	14% (5.9-26%)	86% (74-94%)	1.1 (0.3-2.7)	3.8% (1.3-7.5%)	4.5% (1.6-9.1%)	0.89 (0.3-1.8)	4.6 (1.6-9)
Seychelles	1995	Female	95 plus	17% (6.6-35%)	83% (65-93%)	1.4 (0.28-3.8)	4.3% (1.3-9.2%)	5.1% (1.5-11%)	0.32 (0.099-0.7)	1.1 (0.34-2.3)
Seychelles	1995	Male	All Ages	26% (21-34%)	74% (66-79%)	1.6 (1.4-1.9)	8.8% (7-11%)	9.8% (7.1-13%)	35 (25-45)	1200 (970-1500)
Seychelles	1995	Male	15 to 19	21% (12-35%)	79% (65-88%)	0.7 (0.3-1.4)	2.8% (1.7-4.6%)	3.8% (2-6.7%)	0.21 (0.11-0.37)	18 (11-31)
Seychelles	1995	Male	20 to 24	32% (18-50%)	68% (50-82%)	1.7 (0.82-2.8)	7.2% (4.2-11%)	8.8% (4.8-14%)	0.8 (0.43-1.3)	61 (36-93)
Seychelles	1995	Male	25 to 29	38% (20-56%)	62% (44-80%)	2.4 (1.3-3.9)	9.8% (6.2-14%)	12% (7.1-18%)	1.3 (0.73-1.9)	86 (53-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	1995	Male	30 to 34	41% (22-61%)	59% (39-78%)	3 (1.7-4.5)	12% (8.1-16%)	14% (9.5-19%)	2 (1.3-2.7)	120 (83-160)
Seychelles	1995	Male	35 to 39	43% (26-63%)	57% (37-74%)	3.4 (2-5.3)	13% (9.2-18%)	16% (11-22%)	2.8 (1.8-3.8)	150 (100-200)
Seychelles	1995	Male	40 to 44	43% (26-63%)	57% (37-74%)	3.6 (2-5.5)	14% (9.6-19%)	17% (11-23%)	3.3 (2.2-4.5)	160 (110-220)
Seychelles	1995	Male	45 to 49	42% (24-62%)	58% (38-76%)	3.5 (1.8-5.9)	15% (9.7-20%)	17% (11-23%)	2.5 (1.6-3.4)	110 (69-150)
Seychelles	1995	Male	50 to 54	40% (22-58%)	60% (42-78%)	3.3 (1.7-5.6)	14% (8.5-19%)	16% (9.9-21%)	3.1 (1.9-4.2)	110 (73-160)
Seychelles	1995	Male	55 to 59	37% (22-57%)	63% (43-78%)	3 (1.4-5)	12% (7.5-17%)	14% (8.7-20%)	3.2 (1.9-4.5)	100 (63-140)
Seychelles	1995	Male	60 to 64	34% (19-54%)	66% (46-81%)	2.6 (1.2-4.6)	10% (5.8-16%)	12% (6.6-18%)	3.7 (2-5.7)	100 (56-160)
Seychelles	1995	Male	65 to 69	31% (17-49%)	69% (51-83%)	2.2 (1-3.8)	8.9% (4.7-14%)	10% (5.3-16%)	3.7 (2-5.8)	86 (46-130)
Seychelles	1995	Male	70 to 74	28% (15-44%)	72% (56-85%)	1.8 (0.79-3.3)	7.1% (3.4-12%)	8.1% (3.8-13%)	3.1 (1.5-5.1)	58 (27-94)
Seychelles	1995	Male	75 to 79	25% (14-41%)	75% (59-86%)	1.5 (0.63-2.8)	5.4% (2.3-9.6%)	6.1% (2.6-11%)	2.1 (0.91-3.7)	30 (13-53)
Seychelles	1995	Male	80 to 84	23% (12-39%)	77% (61-88%)	1.3 (0.53-2.4)	4.6% (2-8.3%)	5.2% (2.2-9.4%)	1.6 (0.65-2.8)	17 (7.1-30)
Seychelles	1995	Male	85 to 89	22% (11-36%)	78% (64-89%)	1.1 (0.48-2.2)	3.7% (1.5-7%)	4.1% (1.7-8%)	0.89 (0.36-1.7)	6.9 (2.8-13)
Seychelles	1995	Male	90 to 94	23% (12-38%)	77% (62-88%)	1.1 (0.48-2.1)	3.2% (1.2-6.6%)	3.6% (1.3-7.6%)	0.33 (0.12-0.7)	1.8 (0.67-3.7)
Seychelles	1995	Male	95 plus	27% (12-46%)	73% (54-88%)	1.2 (0.38-2.6)	3.2% (0.95-7.3%)	3.4% (0.93-8.2%)	0.08 (0.022-0.19)	0.29 (0.086-0.68)
Seychelles	2000	Female	All Ages	8.7% (6.4-12%)	91% (88-94%)	0.72 (0.52-0.96)	3.2% (2.3-4.3%)	5.3% (3.4-7.9%)	15 (9.5-22)	310 (220-420)
Seychelles	2000	Female	15 to 19	7.5% (3.4-14%)	93% (86-97%)	0.52 (0.18-1.2)	1.3% (0.71-2.3%)	2.3% (1-4.6%)	0.048 (0.021-0.096)	5.7 (3.1-10)
Seychelles	2000	Female	20 to 24	9.1% (3.6-18%)	91% (82-96%)	0.7 (0.24-1.4)	1.9% (1.2-3%)	2.7% (1.2-4.9%)	0.067 (0.03-0.12)	8.8 (5.2-14)
Seychelles	2000	Female	25 to 29	9.9% (4.1-19%)	90% (81-96%)	0.8 (0.28-1.7)	2.2% (1.4-3.4%)	3.1% (1.5-5.4%)	0.094 (0.044-0.18)	10 (6.1-16)
Seychelles	2000	Female	30 to 34	11% (4.3-21%)	89% (79-96%)	0.9 (0.34-1.9)	2.5% (1.5-3.8%)	3.9% (2.1-6.4%)	0.15 (0.071-0.26)	12 (7.2-20)
Seychelles	2000	Female	35 to 39	12% (4.8-21%)	88% (79-95%)	1 (0.4-1.9)	2.9% (1.7-4.5%)	4.7% (2.6-7.8%)	0.25 (0.12-0.42)	17 (9.2-27)
Seychelles	2000	Female	40 to 44	12% (5.4-22%)	88% (78-95%)	1.1 (0.41-2.2)	3.3% (1.9-5.3%)	5.4% (2.9-8.6%)	0.37 (0.19-0.63)	21 (12-35)
Seychelles	2000	Female	45 to 49	12% (5-22%)	88% (78-95%)	1.3 (0.43-2.7)	3.7% (2-6.3%)	5.7% (3-9.6%)	0.46 (0.24-0.8)	22 (12-37)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	2000	Female	50 to 54	12% (5.2-24%)	88% (76-95%)	1.4 (0.5-3.1)	4.3% (2.3-7.2%)	6.2% (3.3-11%)	0.35 (0.18-0.6)	14 (7.6-24)
Seychelles	2000	Female	55 to 59	12% (5.6-23%)	88% (77-94%)	1.6 (0.52-3.4)	4.5% (2.4-7.5%)	6.4% (3.4-11%)	0.54 (0.29-0.92)	19 (10-31)
Seychelles	2000	Female	60 to 64	12% (5.4-23%)	88% (77-95%)	1.6 (0.5-3.6)	4.6% (2.4-8.1%)	6.2% (3.2-11%)	0.98 (0.51-1.7)	28 (15-50)
Seychelles	2000	Female	65 to 69	12% (5.4-22%)	88% (78-95%)	1.5 (0.51-3.1)	4.9% (2.5-8.3%)	6.4% (3.4-11%)	1.4 (0.75-2.5)	34 (18-59)
Seychelles	2000	Female	70 to 74	12% (5.2-22%)	88% (78-95%)	1.3 (0.43-3)	4.6% (2.2-8.1%)	5.9% (2.9-10%)	1.8 (0.9-3.2)	35 (17-61)
Seychelles	2000	Female	75 to 79	11% (4.8-21%)	89% (79-95%)	1.2 (0.37-2.8)	4.2% (1.8-7.6%)	5.3% (2.3-9.7%)	2 (0.86-3.6)	28 (12-51)
Seychelles	2000	Female	80 to 84	11% (5.2-22%)	89% (78-95%)	1.2 (0.4-2.8)	4.3% (2-8%)	5.3% (2.4-9.9%)	2.5 (1.1-4.6)	26 (12-49)
Seychelles	2000	Female	85 to 89	12% (5.6-23%)	88% (77-94%)	1.3 (0.41-3)	4.4% (1.9-8.5%)	5.3% (2.3-10%)	2.1 (0.89-4.1)	15 (6.8-31)
Seychelles	2000	Female	90 to 94	14% (6.2-27%)	86% (73-94%)	1.5 (0.41-3.6)	4.5% (1.8-8.5%)	5.3% (2.2-10%)	1.3 (0.52-2.4)	6.6 (2.7-12)
Seychelles	2000	Female	95 plus	18% (6.7-34%)	82% (66-93%)	1.8 (0.39-5.2)	5% (1.9-10%)	5.9% (2.2-12%)	0.5 (0.18-1)	1.7 (0.61-3.4)
Seychelles	2000	Male	All Ages	27% (22-35%)	73% (65-78%)	2.2 (1.9-2.6)	10% (8.6-12%)	12% (9.2-14%)	41 (32-50)	1400 (1200-1700)
Seychelles	2000	Male	15 to 19	21% (12-35%)	79% (65-88%)	0.92 (0.4-1.8)	3.4% (1.9-5.7%)	4.7% (2.5-8.4%)	0.27 (0.14-0.49)	23 (13-39)
Seychelles	2000	Male	20 to 24	33% (18-51%)	67% (49-82%)	2.2 (1.1-3.8)	8.7% (5.1-13%)	11% (6.1-17%)	0.94 (0.51-1.4)	71 (42-100)
Seychelles	2000	Male	25 to 29	38% (21-56%)	62% (44-79%)	3.1 (1.6-5)	11% (7.4-16%)	14% (9.1-20%)	1.4 (0.89-2)	98 (63-140)
Seychelles	2000	Male	30 to 34	41% (23-61%)	59% (39-77%)	3.8 (2.1-5.8)	13% (9.5-17%)	17% (12-22%)	2 (1.4-2.6)	120 (86-160)
Seychelles	2000	Male	35 to 39	43% (26-63%)	57% (37-74%)	4.3 (2.5-6.5)	15% (11-19%)	18% (14-24%)	2.8 (2-3.7)	150 (110-200)
Seychelles	2000	Male	40 to 44	43% (26-63%)	57% (37-74%)	4.5 (2.6-6.9)	16% (12-20%)	19% (14-25%)	3.7 (2.7-4.9)	180 (130-230)
Seychelles	2000	Male	45 to 49	41% (23-62%)	59% (38-77%)	4.5 (2.3-7.3)	16% (11-21%)	19% (14-25%)	4.3 (3-5.6)	180 (130-240)
Seychelles	2000	Male	50 to 54	39% (22-59%)	61% (41-78%)	4.2 (2.2-7.1)	15% (11-20%)	18% (13-24%)	2.9 (2-3.9)	110 (78-150)
Seychelles	2000	Male	55 to 59	37% (21-58%)	63% (42-79%)	3.8 (1.9-6.5)	14% (9.5-19%)	17% (11-22%)	3.3 (2.3-4.5)	110 (73-150)
Seychelles	2000	Male	60 to 64	34% (19-54%)	66% (46-81%)	3.3 (1.5-6.1)	12% (7.5-17%)	14% (8.8-20%)	3.5 (2.2-5)	98 (60-140)
Seychelles	2000	Male	65 to 69	31% (17-49%)	69% (51-83%)	2.8 (1.3-4.9)	11% (6.4-15%)	12% (7.4-17%)	4.4 (2.7-6.3)	100 (61-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	2000	Male	70 to 74	27% (15-44%)	73% (56-85%)	2.2 (1-4.2)	8.7% (4.8-13%)	10% (5.5-15%)	4.1 (2.2-6)	76 (42-110)
Seychelles	2000	Male	75 to 79	25% (14-41%)	75% (59-86%)	1.9 (0.79-3.6)	6.9% (3.4-11%)	7.9% (3.8-13%)	3.1 (1.5-5)	45 (22-72)
Seychelles	2000	Male	80 to 84	23% (11-38%)	77% (62-89%)	1.6 (0.69-3.1)	6% (2.8-10%)	6.9% (3.2-12%)	2.5 (1.2-4.2)	27 (13-45)
Seychelles	2000	Male	85 to 89	22% (11-36%)	78% (64-89%)	1.4 (0.64-2.8)	5% (2.2-8.6%)	5.7% (2.5-9.8%)	1.1 (0.5-1.9)	8.6 (3.8-15)
Seychelles	2000	Male	90 to 94	23% (12-38%)	77% (62-88%)	1.4 (0.61-2.8)	4.4% (1.9-8.1%)	5.1% (2.1-9.4%)	0.49 (0.2-0.9)	2.6 (1.1-4.7)
Seychelles	2000	Male	95 plus	26% (12-45%)	74% (55-88%)	1.5 (0.5-3.2)	4.4% (1.5-9%)	4.9% (1.6-10%)	0.14 (0.045-0.29)	0.5 (0.17-1)
Seychelles	2005	Female	All Ages	8.9% (6.3-12%)	91% (88-94%)	0.81 (0.6-1.1)	3.2% (2.3-4.3%)	5.3% (3.4-8%)	16 (10-24)	340 (250-470)
Seychelles	2005	Female	15 to 19	7.3% (3.3-14%)	93% (86-97%)	0.54 (0.18-1.2)	1.3% (0.71-2.2%)	2.3% (0.99-4.7%)	0.043 (0.018-0.087)	5.1 (2.8-8.8)
Seychelles	2005	Female	20 to 24	8.9% (3.8-17%)	91% (83-96%)	0.73 (0.25-1.5)	1.9% (1.1-2.9%)	2.7% (1.2-5%)	0.068 (0.029-0.13)	9.2 (5.3-15)
Seychelles	2005	Female	25 to 29	9.7% (4.1-18%)	90% (82-96%)	0.83 (0.28-1.8)	2.1% (1.3-3.2%)	3% (1.4-5.3%)	0.097 (0.043-0.17)	11 (6.8-18)
Seychelles	2005	Female	30 to 34	11% (4.3-21%)	89% (79-96%)	0.93 (0.34-1.9)	2.4% (1.5-3.6%)	3.8% (2-6.1%)	0.16 (0.08-0.27)	14 (8.2-22)
Seychelles	2005	Female	35 to 39	11% (4.6-22%)	89% (78-95%)	1 (0.39-2)	2.8% (1.6-4.3%)	4.6% (2.5-7.6%)	0.26 (0.13-0.45)	18 (11-29)
Seychelles	2005	Female	40 to 44	12% (5.2-21%)	88% (79-95%)	1.2 (0.45-2.3)	3.2% (1.9-5.2%)	5.3% (3-8.6%)	0.42 (0.22-0.71)	24 (14-39)
Seychelles	2005	Female	45 to 49	12% (5.1-22%)	88% (78-95%)	1.3 (0.48-2.8)	3.7% (2-6.3%)	5.8% (3.1-9.7%)	0.59 (0.31-1)	28 (15-46)
Seychelles	2005	Female	50 to 54	12% (5.2-24%)	88% (76-95%)	1.5 (0.55-3.2)	4.3% (2.3-7.4%)	6.3% (3.4-11%)	0.73 (0.39-1.3)	30 (16-50)
Seychelles	2005	Female	55 to 59	13% (5.6-24%)	87% (76-94%)	1.7 (0.55-3.7)	4.6% (2.4-7.8%)	6.5% (3.4-11%)	0.52 (0.27-0.92)	18 (9.4-31)
Seychelles	2005	Female	60 to 64	13% (5.5-24%)	87% (76-95%)	1.7 (0.54-3.9)	4.7% (2.5-8.3%)	6.3% (3.3-11%)	0.78 (0.41-1.4)	23 (12-40)
Seychelles	2005	Female	65 to 69	12% (5.4-23%)	88% (77-95%)	1.6 (0.55-3.2)	4.9% (2.5-8.3%)	6.5% (3.3-11%)	1.5 (0.75-2.5)	35 (18-59)
Seychelles	2005	Female	70 to 74	12% (5.4-22%)	88% (78-95%)	1.4 (0.47-3.3)	4.6% (2.3-8%)	6% (3-10%)	1.9 (0.95-3.3)	36 (18-62)
Seychelles	2005	Female	75 to 79	11% (5-21%)	89% (79-95%)	1.3 (0.4-3.1)	4.2% (1.9-7.6%)	5.3% (2.4-9.6%)	2.2 (1-4)	32 (15-57)
Seychelles	2005	Female	80 to 84	11% (5.1-22%)	89% (78-95%)	1.3 (0.42-3)	4.4% (2-7.9%)	5.4% (2.5-9.8%)	2.8 (1.3-5.1)	29 (14-53)
Seychelles	2005	Female	85 to 89	12% (5.4-23%)	88% (77-95%)	1.4 (0.44-3.2)	4.4% (1.9-8.4%)	5.3% (2.2-10%)	2.3 (0.96-4.4)	17 (7.2-33)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	2005	Female	90 to 94	14% (6.1-26%)	86% (74-94%)	1.6 (0.45-3.8)	4.5% (1.9-8.2%)	5.4% (2.2-10%)	1.2 (0.51-2.3)	6.4 (2.6-12)
Seychelles	2005	Female	95 plus	17% (6.9-34%)	83% (66-93%)	1.9 (0.41-5.4)	5% (1.8-9.8%)	5.8% (2.1-12%)	0.55 (0.2-1.1)	1.9 (0.7-3.7)
Seychelles	2005	Male	All Ages	29% (23-37%)	71% (63-77%)	2.5 (2.1-3)	10% (8.8-12%)	12% (9.7-14%)	42 (34-51)	1500 (1200-1800)
Seychelles	2005	Male	15 to 19	21% (12-35%)	79% (65-88%)	0.95 (0.4-1.9)	3.5% (1.9-5.9%)	4.9% (2.5-8.6%)	0.27 (0.14-0.48)	22 (12-39)
Seychelles	2005	Male	20 to 24	32% (17-50%)	68% (50-83%)	2.3 (1.1-4)	8.8% (5.1-13%)	11% (6.1-17%)	1 (0.55-1.6)	77 (44-120)
Seychelles	2005	Male	25 to 29	37% (20-56%)	63% (44-80%)	3.2 (1.6-5.2)	11% (7.3-15%)	14% (9-20%)	1.5 (0.9-2.2)	100 (67-150)
Seychelles	2005	Male	30 to 34	41% (22-60%)	59% (40-78%)	4 (2.2-6.1)	13% (9.4-17%)	17% (12-22%)	2.1 (1.4-2.8)	130 (92-170)
Seychelles	2005	Male	35 to 39	43% (25-63%)	57% (37-75%)	4.5 (2.6-7)	14% (11-19%)	19% (14-24%)	2.8 (2-3.7)	160 (110-200)
Seychelles	2005	Male	40 to 44	43% (26-63%)	57% (37-74%)	4.8 (2.7-7.2)	15% (11-20%)	20% (15-25%)	3.7 (2.7-4.9)	180 (140-240)
Seychelles	2005	Male	45 to 49	41% (23-60%)	59% (40-77%)	4.7 (2.5-7.5)	16% (12-20%)	20% (15-25%)	4.5 (3.2-5.9)	200 (140-250)
Seychelles	2005	Male	50 to 54	39% (22-58%)	61% (42-78%)	4.4 (2.3-7.3)	15% (11-20%)	19% (14-24%)	4.8 (3.5-6.3)	180 (130-240)
Seychelles	2005	Male	55 to 59	37% (21-56%)	63% (44-79%)	4 (1.9-6.7)	14% (9.6-18%)	17% (12-22%)	3.2 (2.2-4.4)	110 (73-140)
Seychelles	2005	Male	60 to 64	34% (19-53%)	66% (47-81%)	3.5 (1.5-6.3)	12% (7.7-17%)	14% (9.1-20%)	3.3 (2.1-4.5)	91 (58-130)
Seychelles	2005	Male	65 to 69	31% (17-48%)	69% (52-83%)	2.9 (1.3-5)	11% (6.5-15%)	12% (7.7-17%)	3.2 (1.9-4.4)	73 (46-100)
Seychelles	2005	Male	70 to 74	27% (15-43%)	73% (57-85%)	2.4 (1.1-4.3)	8.8% (5-13%)	10% (5.8-15%)	3.9 (2.2-5.6)	72 (41-110)
Seychelles	2005	Male	75 to 79	25% (14-41%)	75% (59-86%)	2 (0.81-3.8)	7% (3.4-11%)	8.1% (4-13%)	3.3 (1.6-5.2)	48 (24-75)
Seychelles	2005	Male	80 to 84	23% (11-38%)	77% (62-89%)	1.7 (0.71-3.2)	6.2% (2.9-10%)	7.1% (3.3-12%)	2.6 (1.2-4.3)	28 (13-46)
Seychelles	2005	Male	85 to 89	21% (11-35%)	79% (65-89%)	1.5 (0.66-2.9)	5.2% (2.5-8.9%)	5.9% (2.7-10%)	1.4 (0.65-2.5)	11 (5-19)
Seychelles	2005	Male	90 to 94	23% (12-37%)	77% (63-88%)	1.5 (0.63-2.8)	4.6% (2-8.2%)	5.3% (2.2-9.6%)	0.46 (0.19-0.82)	2.4 (1-4.3)
Seychelles	2005	Male	95 plus	26% (12-45%)	74% (55-88%)	1.5 (0.51-3.3)	4.6% (1.7-9.5%)	5.1% (1.7-11%)	0.14 (0.047-0.29)	0.5 (0.18-1)
Seychelles	2010	Female	All Ages	8.9% (6.4-12%)	91% (88-94%)	0.74 (0.52-1)	2.9% (2-3.9%)	4.8% (3-7.1%)	14 (8.8-22)	310 (220-420)
Seychelles	2010	Female	15 to 19	7.3% (3.2-14%)	93% (86-97%)	0.48 (0.15-1.1)	1.1% (0.64-1.9%)	1.9% (0.78-4%)	0.033 (0.013-0.07)	4.4 (2.4-7.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	2010	Female	20 to 24	8.8% (3.7-17%)	91% (83-96%)	0.64 (0.2-1.4)	1.7% (1.1-2.7%)	2.2% (0.92-4.3%)	0.045 (0.019-0.088)	7.3 (4.3-12)
Seychelles	2010	Female	25 to 29	9.6% (4.2-19%)	90% (81-96%)	0.73 (0.24-1.6)	1.9% (1.2-2.9%)	2.5% (1.1-4.4%)	0.061 (0.026-0.11)	8.7 (5.4-14)
Seychelles	2010	Female	30 to 34	11% (4.2-20%)	89% (80-96%)	0.83 (0.29-1.7)	2% (1.3-3.1%)	3.2% (1.6-5.4%)	0.1 (0.05-0.18)	11 (6.3-17)
Seychelles	2010	Female	35 to 39	11% (4.6-22%)	89% (78-95%)	0.92 (0.36-1.8)	2.3% (1.4-3.6%)	4% (2.1-6.7%)	0.18 (0.093-0.31)	14 (8.1-22)
Seychelles	2010	Female	40 to 44	12% (5.3-21%)	88% (79-95%)	1 (0.38-2.1)	2.8% (1.6-4.5%)	4.7% (2.6-7.9%)	0.31 (0.16-0.52)	19 (11-31)
Seychelles	2010	Female	45 to 49	12% (4.8-23%)	88% (77-95%)	1.2 (0.4-2.5)	3.3% (1.8-5.5%)	5.3% (2.7-9%)	0.52 (0.26-0.9)	26 (14-44)
Seychelles	2010	Female	50 to 54	12% (5.2-24%)	88% (76-95%)	1.3 (0.47-2.8)	3.8% (2.1-6.6%)	5.8% (3-10%)	0.76 (0.39-1.3)	32 (17-54)
Seychelles	2010	Female	55 to 59	12% (5.4-23%)	88% (77-95%)	1.4 (0.48-3)	4.1% (2.2-7%)	5.9% (3.1-10%)	0.91 (0.47-1.6)	32 (17-55)
Seychelles	2010	Female	60 to 64	12% (5.2-23%)	88% (77-95%)	1.5 (0.45-3.2)	4.2% (2.2-7.6%)	5.8% (3-10%)	0.69 (0.36-1.3)	20 (11-37)
Seychelles	2010	Female	65 to 69	12% (5.3-23%)	88% (77-95%)	1.4 (0.45-2.8)	4.4% (2.2-7.6%)	5.9% (3-10%)	1.1 (0.53-1.9)	26 (13-45)
Seychelles	2010	Female	70 to 74	11% (5.2-21%)	89% (79-95%)	1.2 (0.38-2.9)	4.1% (1.9-7.3%)	5.4% (2.5-9.7%)	1.6 (0.73-2.8)	30 (14-53)
Seychelles	2010	Female	75 to 79	11% (4.9-21%)	89% (79-95%)	1.1 (0.33-2.6)	3.7% (1.5-7%)	4.7% (1.9-8.9%)	1.8 (0.73-3.4)	26 (11-49)
Seychelles	2010	Female	80 to 84	11% (4.9-21%)	89% (79-95%)	1.1 (0.36-2.7)	3.8% (1.6-7.1%)	4.7% (2-8.8%)	2.4 (1-4.5)	26 (11-48)
Seychelles	2010	Female	85 to 89	12% (5.5-22%)	88% (78-95%)	1.2 (0.38-2.8)	3.9% (1.6-7.3%)	4.7% (1.9-9%)	2.2 (0.88-4.1)	16 (6.7-31)
Seychelles	2010	Female	90 to 94	14% (6.3-26%)	86% (74-94%)	1.4 (0.41-3.4)	4% (1.5-7.6%)	4.8% (1.8-9.3%)	1.2 (0.47-2.4)	6.4 (2.5-12)
Seychelles	2010	Female	95 plus	17% (6.4-34%)	83% (66-94%)	1.7 (0.35-5.1)	4.5% (1.4-9.2%)	5.3% (1.7-11%)	0.49 (0.16-0.99)	1.7 (0.55-3.4)
Seychelles	2010	Male	All Ages	29% (23-36%)	71% (64-77%)	2.3 (1.8-2.8)	9.4% (7.7-11%)	11% (8.5-14%)	40 (31-50)	1400 (1200-1700)
Seychelles	2010	Male	15 to 19	21% (11-34%)	79% (66-89%)	0.81 (0.34-1.6)	3% (1.7-5.1%)	4.2% (2.2-7.4%)	0.21 (0.11-0.38)	18 (10-31)
Seychelles	2010	Male	20 to 24	32% (17-49%)	68% (51-83%)	2 (0.92-3.5)	7.5% (4.3-11%)	9.8% (5.1-16%)	0.76 (0.4-1.2)	60 (34-91)
Seychelles	2010	Male	25 to 29	37% (20-55%)	63% (45-80%)	2.8 (1.4-4.7)	9.5% (6-13%)	13% (7.6-18%)	1.1 (0.68-1.7)	82 (52-120)
Seychelles	2010	Male	30 to 34	40% (22-60%)	60% (40-78%)	3.4 (1.8-5.6)	11% (7.6-15%)	15% (9.9-20%)	1.6 (1-2.2)	100 (70-140)
Seychelles	2010	Male	35 to 39	42% (25-62%)	58% (38-75%)	3.9 (2.2-6.3)	12% (8.8-16%)	17% (12-22%)	2.3 (1.5-3)	130 (91-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	2010	Male	40 to 44	42% (25-62%)	58% (38-75%)	4.1 (2.2-6.3)	14% (9.7-17%)	18% (13-23%)	3.1 (2.2-4)	160 (110-200)
Seychelles	2010	Male	45 to 49	41% (23-61%)	59% (39-77%)	4.1 (2.1-6.5)	14% (10-18%)	18% (13-24%)	4.2 (2.9-5.4)	180 (130-240)
Seychelles	2010	Male	50 to 54	39% (21-58%)	61% (42-79%)	3.8 (2-6.3)	14% (9.8-18%)	17% (12-22%)	5.1 (3.6-6.8)	200 (140-260)
Seychelles	2010	Male	55 to 59	37% (21-57%)	63% (43-79%)	3.4 (1.7-5.8)	13% (8.5-17%)	16% (10-21%)	5.2 (3.5-7.1)	170 (120-230)
Seychelles	2010	Male	60 to 64	34% (18-54%)	66% (46-82%)	3 (1.3-5.6)	11% (6.7-16%)	13% (8-19%)	3.2 (2-4.6)	91 (56-130)
Seychelles	2010	Male	65 to 69	30% (17-47%)	70% (53-83%)	2.5 (1.2-4.5)	9.6% (5.5-14%)	11% (6.5-16%)	3.6 (2.1-5.3)	84 (49-120)
Seychelles	2010	Male	70 to 74	27% (15-43%)	73% (57-85%)	2 (0.9-3.8)	7.7% (4.1-12%)	9.1% (4.8-14%)	3.1 (1.7-4.8)	59 (31-89)
Seychelles	2010	Male	75 to 79	24% (13-41%)	76% (59-87%)	1.7 (0.69-3.3)	6.1% (2.9-10%)	7.1% (3.3-12%)	2.7 (1.2-4.5)	40 (18-66)
Seychelles	2010	Male	80 to 84	22% (11-37%)	78% (63-89%)	1.5 (0.61-2.7)	5.3% (2.3-9.1%)	6.1% (2.6-10%)	2.3 (0.95-3.9)	24 (11-42)
Seychelles	2010	Male	85 to 89	21% (11-34%)	79% (66-89%)	1.3 (0.57-2.5)	4.3% (1.9-7.8%)	4.9% (2.1-9.1%)	1.3 (0.54-2.3)	9.7 (4.3-18)
Seychelles	2010	Male	90 to 94	22% (12-38%)	78% (62-88%)	1.2 (0.52-2.4)	3.8% (1.5-7.3%)	4.3% (1.7-8.5%)	0.48 (0.19-0.95)	2.6 (1-5)
Seychelles	2010	Male	95 plus	26% (11-46%)	74% (54-89%)	1.3 (0.43-2.8)	3.8% (1.2-8.3%)	4.1% (1.3-9.5%)	0.11 (0.032-0.24)	0.38 (0.13-0.85)
Seychelles	2016	Female	All Ages	9.3% (6.8-12%)	91% (88-93%)	0.71 (0.41-1.1)	2.8% (1.9-3.9%)	4.5% (2.7-6.8%)	15 (8.5-22)	320 (210-460)
Seychelles	2016	Female	15 to 19	7.6% (3.5-14%)	92% (86-97%)	0.44 (0.13-1.1)	1.1% (0.58-1.9%)	1.8% (0.73-3.8%)	0.026 (0.01-0.056)	3.6 (1.9-6.5)
Seychelles	2016	Female	20 to 24	9.1% (3.8-17%)	91% (83-96%)	0.59 (0.17-1.4)	1.7% (1-2.5%)	2% (0.81-3.8%)	0.038 (0.015-0.077)	6.5 (3.8-10)
Seychelles	2016	Female	25 to 29	9.8% (4.1-19%)	90% (81-96%)	0.67 (0.2-1.5)	1.7% (1.1-2.7%)	2.1% (0.97-3.9%)	0.051 (0.022-0.1)	7.7 (4.6-12)
Seychelles	2016	Female	30 to 34	11% (4.3-20%)	89% (80-96%)	0.76 (0.25-1.7)	1.9% (1.2-3%)	2.8% (1.4-5%)	0.094 (0.043-0.19)	10 (5.8-16)
Seychelles	2016	Female	35 to 39	12% (4.7-22%)	88% (78-95%)	0.84 (0.3-1.8)	2.2% (1.3-3.4%)	3.6% (1.9-6.2%)	0.17 (0.084-0.32)	13 (7.6-22)
Seychelles	2016	Female	40 to 44	12% (5.3-22%)	88% (78-95%)	0.93 (0.31-2)	2.6% (1.5-4.2%)	4.3% (2.3-7.2%)	0.3 (0.15-0.55)	19 (9.7-31)
Seychelles	2016	Female	45 to 49	12% (5.2-23%)	88% (77-95%)	1.1 (0.35-2.3)	3% (1.6-5.2%)	4.9% (2.5-8.4%)	0.51 (0.24-0.89)	26 (13-44)
Seychelles	2016	Female	50 to 54	13% (5.5-24%)	87% (76-94%)	1.2 (0.4-2.5)	3.6% (1.9-6.2%)	5.4% (2.7-9.4%)	0.83 (0.41-1.4)	35 (17-60)
Seychelles	2016	Female	55 to 59	13% (5.7-24%)	87% (76-94%)	1.3 (0.42-2.8)	3.8% (2-6.4%)	5.5% (2.9-9.4%)	1.1 (0.55-2)	40 (20-69)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	2016	Female	60 to 64	13% (5.6-25%)	87% (75-94%)	1.3 (0.41-3.1)	3.9% (1.9-6.8%)	5.4% (2.7-9.4%)	1.3 (0.63-2.3)	38 (19-69)
Seychelles	2016	Female	65 to 69	12% (5.7-23%)	88% (77-94%)	1.3 (0.39-2.7)	4% (1.9-7%)	5.4% (2.7-9.3%)	1 (0.47-1.8)	25 (12-44)
Seychelles	2016	Female	70 to 74	12% (5.4-21%)	88% (79-95%)	1.1 (0.33-2.7)	3.8% (1.6-6.7%)	5% (2.2-9%)	1.2 (0.52-2.2)	23 (10-42)
Seychelles	2016	Female	75 to 79	11% (4.8-22%)	89% (78-95%)	1.1 (0.29-2.8)	3.4% (1.3-6.5%)	4.4% (1.7-8.5%)	1.7 (0.64-3.3)	24 (9.4-47)
Seychelles	2016	Female	80 to 84	11% (4.9-21%)	89% (79-95%)	1.1 (0.3-2.7)	3.5% (1.3-6.6%)	4.4% (1.6-8.3%)	2.2 (0.82-4.3)	24 (8.9-46)
Seychelles	2016	Female	85 to 89	12% (5.4-23%)	88% (77-95%)	1.1 (0.33-2.8)	3.5% (1.3-6.7%)	4.3% (1.6-8.3%)	2.2 (0.8-4.3)	17 (6.2-33)
Seychelles	2016	Female	90 to 94	14% (6-27%)	86% (73-94%)	1.3 (0.33-3.3)	3.6% (1.3-7.1%)	4.3% (1.5-8.5%)	1.4 (0.5-2.7)	7.1 (2.6-14)
Seychelles	2016	Female	95 plus	18% (6.5-34%)	82% (66-93%)	1.6 (0.29-4.7)	4.1% (1.1-8.6%)	4.8% (1.3-10%)	0.56 (0.15-1.2)	1.9 (0.52-4.1)
Seychelles	2016	Male	All Ages	29% (23-37%)	71% (63-77%)	2.1 (1.3-3.1)	8.8% (6.2-11%)	10% (6.7-13%)	41 (26-56)	1400 (950-1900)
Seychelles	2016	Male	15 to 19	21% (11-34%)	79% (66-89%)	0.74 (0.27-1.6)	2.8% (1.6-5%)	4% (2-7.4%)	0.16 (0.082-0.31)	14 (8-26)
Seychelles	2016	Male	20 to 24	32% (17-51%)	68% (49-83%)	1.8 (0.73-3.6)	6.8% (3.7-11%)	8.8% (4.4-15%)	0.6 (0.3-1)	48 (26-79)
Seychelles	2016	Male	25 to 29	38% (20-56%)	62% (44-80%)	2.5 (1.1-4.7)	8.5% (5-13%)	11% (6.1-17%)	0.88 (0.46-1.4)	64 (37-99)
Seychelles	2016	Male	30 to 34	41% (22-60%)	59% (40-78%)	3.1 (1.5-5.7)	9.9% (6.1-14%)	13% (7.8-19%)	1.3 (0.75-2)	87 (52-130)
Seychelles	2016	Male	35 to 39	43% (26-62%)	57% (38-74%)	3.5 (1.7-6.3)	11% (7.1-15%)	15% (9.3-21%)	1.9 (1.1-2.8)	110 (68-160)
Seychelles	2016	Male	40 to 44	43% (26-63%)	57% (37-74%)	3.7 (1.8-6.3)	12% (8.1-16%)	16% (11-22%)	2.9 (1.7-4.1)	150 (92-200)
Seychelles	2016	Male	45 to 49	41% (23-61%)	59% (39-77%)	3.7 (1.7-6.4)	13% (8.6-18%)	17% (11-23%)	4 (2.4-5.8)	180 (110-250)
Seychelles	2016	Male	50 to 54	39% (22-59%)	61% (41-78%)	3.4 (1.6-5.9)	13% (7.9-17%)	16% (10-22%)	5 (2.9-7.4)	190 (120-280)
Seychelles	2016	Male	55 to 59	37% (21-57%)	63% (43-79%)	3.1 (1.4-5.5)	12% (7.1-16%)	15% (8.8-20%)	5.8 (3.4-8.5)	190 (110-280)
Seychelles	2016	Male	60 to 64	34% (19-54%)	66% (46-81%)	2.7 (1.1-5.1)	10% (5.4-15%)	12% (6.5-18%)	5.4 (2.9-8.2)	150 (83-230)
Seychelles	2016	Male	65 to 69	31% (17-49%)	69% (51-83%)	2.2 (0.94-4.2)	8.6% (4.5-13%)	10% (5.4-15%)	3.3 (1.7-5.2)	78 (39-120)
Seychelles	2016	Male	70 to 74	27% (15-44%)	73% (56-85%)	1.8 (0.72-3.6)	7% (3.3-11%)	8.2% (3.9-13%)	2.9 (1.4-4.7)	54 (26-87)
Seychelles	2016	Male	75 to 79	25% (13-41%)	75% (59-87%)	1.5 (0.58-3.2)	5.4% (2.3-9.6%)	6.3% (2.6-11%)	2.3 (0.94-4.2)	34 (14-61)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Seychelles	2016	Male	80 to 84	23% (11-38%)	77% (62-89%)	1.3 (0.49-2.6)	4.7% (1.8-8.4%)	5.4% (2.1-9.6%)	2.1 (0.8-3.8)	23 (9-42)
Seychelles	2016	Male	85 to 89	21% (11-35%)	79% (65-89%)	1.2 (0.45-2.4)	3.8% (1.4-7.3%)	4.3% (1.6-8.4%)	1.3 (0.48-2.7)	10 (3.8-20)
Seychelles	2016	Male	90 to 94	22% (12-37%)	78% (63-88%)	1.1 (0.43-2.3)	3.3% (1.1-6.7%)	3.8% (1.3-7.8%)	0.56 (0.18-1.2)	3 (1-6.1)
Seychelles	2016	Male	95 plus	26% (12-44%)	74% (56-88%)	1.2 (0.36-2.8)	3.3% (0.98-7.7%)	3.6% (1-8.6%)	0.15 (0.042-0.35)	0.55 (0.16-1.3)
Somalia	1990	Female	All Ages	35% (29-41%)	65% (59-71%)	0.073 (0.067-0.078)	0.23% (0.084-0.4%)	0.26% (-0.013-0.56%)	120 (-5.6-260)	6200 (2200-11000)
Somalia	1990	Female	15 to 19	36% (23-51%)	64% (49-77%)	0.087 (0.05-0.13)	0.3% (0.052-0.57%)	0.091% (-0.26-0.46%)	0.76 (-2.1-3.7)	260 (46-490)
Somalia	1990	Female	20 to 24	42% (27-58%)	58% (42-73%)	0.12 (0.071-0.18)	0.56% (0.19-0.98%)	0.13% (-0.33-0.63%)	1.3 (-3.4-6.6)	520 (170-950)
Somalia	1990	Female	25 to 29	44% (29-61%)	56% (39-71%)	0.14 (0.084-0.2)	0.69% (0.19-1.2%)	0.33% (-0.33-1%)	3.5 (-3.8-11)	610 (160-1100)
Somalia	1990	Female	30 to 34	46% (31-61%)	54% (39-69%)	0.14 (0.088-0.21)	0.75% (0.2-1.3%)	0.51% (-0.18-1.3%)	5.8 (-2.1-14)	640 (160-1100)
Somalia	1990	Female	35 to 39	49% (34-64%)	51% (36-66%)	0.14 (0.085-0.2)	0.83% (0.31-1.5%)	0.74% (0.064-1.6%)	9.5 (0.84-21)	710 (270-1300)
Somalia	1990	Female	40 to 44	50% (36-66%)	50% (34-64%)	0.14 (0.079-0.2)	1% (0.48-1.6%)	1% (0.41-1.8%)	14 (5.4-27)	790 (370-1300)
Somalia	1990	Female	45 to 49	50% (33-67%)	50% (33-67%)	0.14 (0.078-0.22)	0.9% (0.34-1.5%)	0.96% (0.27-1.7%)	14 (3.8-26)	670 (250-1200)
Somalia	1990	Female	50 to 54	48% (33-65%)	52% (35-67%)	0.14 (0.082-0.21)	0.89% (0.29-1.6%)	0.99% (0.26-1.9%)	18 (4.7-36)	720 (230-1400)
Somalia	1990	Female	55 to 59	47% (32-63%)	53% (37-68%)	0.14 (0.079-0.23)	0.46% (-0.11-1.1%)	0.51% (-0.16-1.2%)	10 (-3.2-26)	350 (-85-820)
Somalia	1990	Female	60 to 64	48% (31-64%)	52% (36-69%)	0.14 (0.077-0.23)	0.45% (-0.2-1.1%)	0.52% (-0.23-1.2%)	12 (-5.6-30)	330 (-150-790)
Somalia	1990	Female	65 to 69	49% (34-65%)	51% (35-66%)	0.13 (0.078-0.2)	0.5% (-0.14-1.2%)	0.54% (-0.18-1.3%)	15 (-4.9-36)	350 (-100-830)
Somalia	1990	Female	70 to 74	48% (33-63%)	52% (37-67%)	0.12 (0.062-0.2)	0.17% (-0.45-0.71%)	0.14% (-0.54-0.74%)	4.2 (-16-22)	100 (-260-410)
Somalia	1990	Female	75 to 79	45% (29-61%)	55% (39-71%)	0.1 (0.049-0.17)	0.23% (-0.38-0.71%)	0.24% (-0.45-0.78%)	6.1 (-11-20)	92 (-150-290)
Somalia	1990	Female	80 to 84	44% (28-61%)	56% (39-72%)	0.095 (0.048-0.15)	0.016% (-0.51-0.41%)	0.011% (-0.57-0.45%)	0.18 (-9.2-7.2)	3.1 (-94-74)
Somalia	1990	Female	85 to 89	44% (29-59%)	56% (41-71%)	0.091 (0.047-0.14)	-0.051% (-0.5-0.27%)	-0.072% (-0.6-0.31%)	-0.59 (-4.9-2.5)	-3.5 (-35-19)
Somalia	1990	Female	90 to 94	46% (30-60%)	54% (40-70%)	0.089 (0.045-0.15)	-0.11% (-0.53-0.18%)	-0.17% (-0.66-0.19%)	-0.38 (-1.5-0.44)	-1.5 (-7.4-2.5)
Somalia	1990	Female	95 plus	48% (30-65%)	52% (35-70%)	0.086 (0.039-0.16)	0.22% (-0.24-0.63%)	0.19% (-0.34-0.68%)	0.091 (-0.17-0.33)	0.41 (-0.45-1.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	1990	Male	All Ages	43% (37-51%)	57% (49-63%)	0.087 (0.081-0.093)	0.42% (0.083-0.77%)	0.49% (-0.059-1%)	220 (-28-480)	13000 (2700-23000)
Somalia	1990	Male	15 to 19	42% (30-57%)	58% (43-70%)	0.1 (0.062-0.15)	0.7% (0.28-1.1%)	0.24% (-0.21-0.76%)	2.4 (-2.2-7.6)	680 (260-1100)
Somalia	1990	Male	20 to 24	53% (39-67%)	47% (33-61%)	0.13 (0.086-0.19)	1.2% (0.28-2.1%)	0.33% (-0.6-1.3%)	4.7 (-9-19)	1400 (310-2500)
Somalia	1990	Male	25 to 29	56% (41-71%)	44% (29-59%)	0.16 (0.1-0.21)	1.5% (0.12-2.9%)	0.61% (-1-2.3%)	8 (-14-29)	1500 (120-2800)
Somalia	1990	Male	30 to 34	58% (43-73%)	42% (27-57%)	0.18 (0.12-0.24)	1.5% (-0.14-3.1%)	0.81% (-1.2-2.9%)	10 (-15-35)	1300 (-130-2800)
Somalia	1990	Male	35 to 39	60% (47-75%)	40% (25-53%)	0.19 (0.12-0.27)	1.6% (-0.15-3.4%)	1.2% (-1-3.4%)	16 (-13-45)	1300 (-130-2800)
Somalia	1990	Male	40 to 44	61% (47-76%)	39% (24-53%)	0.19 (0.12-0.29)	1.8% (0.079-3.4%)	1.6% (-0.46-3.7%)	22 (-6.9-51)	1400 (56-2700)
Somalia	1990	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.19 (0.12-0.28)	1.8% (0.26-3.4%)	1.8% (-0.17-3.7%)	25 (-2.2-54)	1300 (170-2400)
Somalia	1990	Male	50 to 54	61% (45-76%)	39% (24-55%)	0.19 (0.11-0.28)	1.7% (0.12-3.3%)	1.7% (-0.18-3.7%)	27 (-2.6-59)	1200 (81-2300)
Somalia	1990	Male	55 to 59	61% (46-76%)	39% (24-54%)	0.18 (0.11-0.28)	1.4% (0.055-2.7%)	1.4% (-0.22-3%)	24 (-3.6-53)	900 (32-1800)
Somalia	1990	Male	60 to 64	60% (44-76%)	40% (24-56%)	0.17 (0.098-0.27)	1.5% (0.43-2.5%)	1.5% (0.34-2.8%)	29 (6-56)	870 (260-1600)
Somalia	1990	Male	65 to 69	58% (44-74%)	42% (26-56%)	0.14 (0.086-0.21)	0.93% (0.22-1.7%)	0.99% (0.15-1.9%)	20 (3-39)	490 (110-910)
Somalia	1990	Male	70 to 74	57% (42-71%)	43% (29-58%)	0.11 (0.067-0.18)	0.69% (0.13-1.2%)	0.76% (0.1-1.4%)	15 (2.1-29)	290 (55-540)
Somalia	1990	Male	75 to 79	55% (41-70%)	45% (30-59%)	0.096 (0.053-0.15)	0.65% (0.18-1.1%)	0.7% (0.16-1.3%)	12 (2.6-21)	170 (45-300)
Somalia	1990	Male	80 to 84	54% (37-70%)	46% (30-63%)	0.086 (0.049-0.13)	0.42% (0.084-0.75%)	0.43% (0.048-0.83%)	4 (0.43-7.7)	46 (9.3-83)
Somalia	1990	Male	85 to 89	52% (36-66%)	48% (34-64%)	0.076 (0.044-0.12)	0.45% (0.16-0.75%)	0.45% (0.13-0.8%)	2.4 (0.69-4.3)	20 (7.3-33)
Somalia	1990	Male	90 to 94	52% (37-66%)	48% (34-63%)	0.07 (0.039-0.11)	0.29% (0.062-0.51%)	0.25% (-0.011-0.51%)	0.4 (-0.018-0.82)	2.8 (0.59-4.8)
Somalia	1990	Male	95 plus	54% (35-72%)	46% (28-65%)	0.065 (0.03-0.11)	0.44% (0.16-0.71%)	0.39% (0.033-0.73%)	0.12 (0.01-0.23)	0.55 (0.19-0.9)
Somalia	1995	Female	All Ages	35% (29-41%)	65% (59-71%)	0.072 (0.063-0.081)	0.24% (0.065-0.45%)	0.27% (-0.061-0.62%)	120 (-25-270)	6200 (1700-12000)
Somalia	1995	Female	15 to 19	36% (23-51%)	64% (49-77%)	0.087 (0.05-0.13)	0.31% (0.00096-0.65%)	0.097% (-0.35-0.56%)	0.73 (-2.4-4.1)	260 (0.78-530)
Somalia	1995	Female	20 to 24	42% (27-57%)	58% (43-73%)	0.12 (0.072-0.18)	0.57% (0.12-1.1%)	0.14% (-0.45-0.77%)	1.3 (-4.5-7.5)	510 (100-950)
Somalia	1995	Female	25 to 29	44% (29-60%)	56% (40-71%)	0.14 (0.086-0.2)	0.7% (0.11-1.3%)	0.33% (-0.44-1.1%)	3.5 (-5-12)	610 (99-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	1995	Female	30 to 34	46% (31-62%)	54% (38-69%)	0.15 (0.087-0.21)	0.75% (0.13-1.5%)	0.52% (-0.29-1.5%)	5.9 (-3.5-16)	640 (100-1200)
Somalia	1995	Female	35 to 39	49% (33-64%)	51% (36-67%)	0.14 (0.082-0.21)	0.84% (0.24-1.6%)	0.75% (-0.0048-1.7%)	9.7 (-0.05-23)	710 (200-1400)
Somalia	1995	Female	40 to 44	50% (36-65%)	50% (35-64%)	0.13 (0.075-0.2)	1% (0.42-1.8%)	1.1% (0.36-2%)	15 (4.9-29)	830 (350-1500)
Somalia	1995	Female	45 to 49	50% (34-66%)	50% (34-66%)	0.14 (0.078-0.21)	0.91% (0.27-1.6%)	0.98% (0.2-1.9%)	15 (3-29)	700 (200-1300)
Somalia	1995	Female	50 to 54	48% (33-64%)	52% (36-67%)	0.14 (0.079-0.22)	0.91% (0.21-1.7%)	1% (0.13-2%)	18 (2.2-36)	690 (150-1300)
Somalia	1995	Female	55 to 59	47% (32-64%)	53% (36-68%)	0.14 (0.078-0.23)	0.46% (-0.26-1.1%)	0.51% (-0.33-1.3%)	11 (-7.3-29)	360 (-200-930)
Somalia	1995	Female	60 to 64	48% (32-65%)	52% (35-68%)	0.14 (0.078-0.23)	0.46% (-0.28-1.1%)	0.52% (-0.3-1.3%)	12 (-7.3-31)	330 (-190-810)
Somalia	1995	Female	65 to 69	49% (34-64%)	51% (36-66%)	0.13 (0.077-0.21)	0.52% (-0.13-1.2%)	0.55% (-0.14-1.3%)	15 (-4-36)	350 (-87-790)
Somalia	1995	Female	70 to 74	48% (34-63%)	52% (37-66%)	0.12 (0.062-0.2)	0.18% (-0.46-0.76%)	0.15% (-0.56-0.8%)	4.3 (-16-23)	100 (-270-450)
Somalia	1995	Female	75 to 79	46% (30-61%)	54% (39-70%)	0.1 (0.05-0.17)	0.25% (-0.42-0.75%)	0.26% (-0.47-0.81%)	6.4 (-12-20)	95 (-160-290)
Somalia	1995	Female	80 to 84	44% (28-61%)	56% (39-72%)	0.095 (0.049-0.15)	0.024% (-0.51-0.42%)	0.021% (-0.58-0.46%)	0.35 (-9.6-7.7)	4.7 (-99-79)
Somalia	1995	Female	85 to 89	44% (28-59%)	56% (41-72%)	0.09 (0.046-0.15)	-0.043% (-0.5-0.3%)	-0.062% (-0.59-0.32%)	-0.52 (-4.9-2.7)	-3 (-35-20)
Somalia	1995	Female	90 to 94	46% (30-61%)	54% (39-70%)	0.089 (0.044-0.15)	-0.1% (-0.52-0.21%)	-0.16% (-0.65-0.21%)	-0.47 (-1.9-0.63)	-1.8 (-9.1-3.6)
Somalia	1995	Female	95 plus	48% (30-65%)	52% (35-70%)	0.086 (0.037-0.16)	0.24% (-0.2-0.66%)	0.21% (-0.32-0.72%)	0.16 (-0.24-0.53)	0.68 (-0.61-1.9)
Somalia	1995	Male	All Ages	43% (37-50%)	57% (50-63%)	0.086 (0.076-0.097)	0.45% (0.059-0.85%)	0.52% (-0.1-1.2%)	220 (-44-500)	13000 (1800-24000)
Somalia	1995	Male	15 to 19	42% (29-58%)	58% (42-71%)	0.099 (0.061-0.15)	0.79% (0.29-1.3%)	0.29% (-0.34-0.98%)	2.5 (-3.1-8.6)	680 (250-1200)
Somalia	1995	Male	20 to 24	53% (39-67%)	47% (33-61%)	0.13 (0.083-0.19)	1.4% (0.26-2.5%)	0.41% (-0.94-1.7%)	4.8 (-11-20)	1400 (260-2500)
Somalia	1995	Male	25 to 29	56% (42-70%)	44% (30-58%)	0.16 (0.1-0.22)	1.6% (-0.011-3.3%)	0.7% (-1.3-2.7%)	8 (-15-32)	1500 (-8.6-3000)
Somalia	1995	Male	30 to 34	59% (43-73%)	41% (27-57%)	0.18 (0.12-0.24)	1.6% (-0.27-3.3%)	0.87% (-1.5-3.1%)	10 (-17-37)	1300 (-220-2800)
Somalia	1995	Male	35 to 39	60% (46-74%)	40% (26-54%)	0.19 (0.12-0.28)	1.7% (-0.34-3.6%)	1.3% (-1.2-3.7%)	16 (-15-47)	1300 (-260-2900)
Somalia	1995	Male	40 to 44	61% (46-76%)	39% (24-54%)	0.19 (0.12-0.29)	1.8% (-0.036-3.5%)	1.6% (-0.65-3.8%)	22 (-9.6-53)	1300 (-28-2700)
Somalia	1995	Male	45 to 49	61% (46-77%)	39% (23-54%)	0.19 (0.11-0.29)	1.8% (0.073-3.6%)	1.8% (-0.35-3.9%)	26 (-5-58)	1300 (48-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	1995	Male	50 to 54	61% (46-76%)	39% (24-54%)	0.19 (0.11-0.28)	1.7% (-0.089-3.3%)	1.7% (-0.45-3.6%)	25 (-6.9-58)	1100 (-56-2300)
Somalia	1995	Male	55 to 59	61% (46-75%)	39% (25-54%)	0.19 (0.11-0.29)	1.4% (-0.069-2.9%)	1.4% (-0.38-3.2%)	24 (-6.4-56)	890 (-42-1900)
Somalia	1995	Male	60 to 64	60% (44-76%)	40% (24-56%)	0.17 (0.098-0.27)	1.4% (0.34-2.7%)	1.5% (0.2-2.9%)	28 (3.7-56)	830 (190-1600)
Somalia	1995	Male	65 to 69	58% (44-75%)	42% (25-56%)	0.14 (0.085-0.21)	0.94% (0.15-1.7%)	1% (0.059-1.9%)	19 (1.2-39)	470 (73-910)
Somalia	1995	Male	70 to 74	57% (42-71%)	43% (29-58%)	0.11 (0.067-0.18)	0.7% (0.095-1.3%)	0.76% (0.061-1.5%)	15 (1.2-29)	280 (39-530)
Somalia	1995	Male	75 to 79	55% (41-71%)	45% (29-59%)	0.095 (0.052-0.15)	0.66% (0.15-1.2%)	0.71% (0.13-1.3%)	11 (2.1-21)	170 (38-300)
Somalia	1995	Male	80 to 84	54% (38-70%)	46% (30-62%)	0.086 (0.047-0.13)	0.42% (0.069-0.77%)	0.44% (0.03-0.84%)	4.2 (0.28-8.2)	48 (7.8-88)
Somalia	1995	Male	85 to 89	52% (37-67%)	48% (33-63%)	0.076 (0.042-0.12)	0.46% (0.15-0.75%)	0.47% (0.098-0.82%)	2.6 (0.54-4.5)	21 (6.5-35)
Somalia	1995	Male	90 to 94	52% (37-66%)	48% (34-63%)	0.07 (0.037-0.11)	0.29% (0.05-0.51%)	0.26% (-0.028-0.53%)	0.55 (-0.06-1.1)	3.7 (0.64-6.6)
Somalia	1995	Male	95 plus	54% (35-72%)	46% (28-65%)	0.065 (0.029-0.12)	0.45% (0.15-0.74%)	0.4% (0.052-0.76%)	0.2 (0.025-0.37)	0.87 (0.28-1.4)
Somalia	2000	Female	All Ages	35% (29-41%)	65% (59-71%)	0.075 (0.05-0.11)	0.25% (-0.043-0.54%)	0.27% (-0.24-0.77%)	120 (-110-370)	6700 (-1200-15000)
Somalia	2000	Female	15 to 19	36% (23-51%)	64% (49-77%)	0.089 (0.043-0.15)	0.31% (-0.11-0.74%)	0.093% (-0.54-0.78%)	0.79 (-4.6-6.5)	290 (-97-700)
Somalia	2000	Female	20 to 24	42% (27-57%)	58% (43-73%)	0.12 (0.062-0.21)	0.56% (-0.052-1.2%)	0.13% (-0.71-0.98%)	1.4 (-7.6-11)	570 (-51-1200)
Somalia	2000	Female	25 to 29	44% (29-60%)	56% (40-71%)	0.14 (0.078-0.25)	0.67% (-0.055-1.4%)	0.3% (-0.64-1.3%)	3.7 (-8.1-16)	660 (-50-1500)
Somalia	2000	Female	30 to 34	46% (31-62%)	54% (38-69%)	0.15 (0.079-0.25)	0.71% (-0.13-1.6%)	0.47% (-0.66-1.6%)	6.3 (-8.4-22)	700 (-120-1500)
Somalia	2000	Female	35 to 39	49% (34-64%)	51% (36-66%)	0.14 (0.075-0.24)	0.77% (-0.078-1.7%)	0.68% (-0.43-1.9%)	10 (-6-30)	760 (-85-1700)
Somalia	2000	Female	40 to 44	50% (35-65%)	50% (35-65%)	0.14 (0.069-0.25)	0.95% (0.15-1.8%)	0.99% (-0.031-2.1%)	16 (-0.47-35)	890 (130-1700)
Somalia	2000	Female	45 to 49	50% (32-66%)	50% (34-68%)	0.15 (0.071-0.25)	0.85% (-0.068-1.8%)	0.91% (-0.22-2.1%)	16 (-3.9-38)	770 (-62-1600)
Somalia	2000	Female	50 to 54	48% (33-65%)	52% (35-67%)	0.15 (0.073-0.25)	0.84% (-0.24-2%)	0.93% (-0.34-2.3%)	20 (-7.5-48)	760 (-220-1700)
Somalia	2000	Female	55 to 59	47% (33-64%)	53% (36-67%)	0.15 (0.074-0.27)	0.4% (-0.61-1.4%)	0.44% (-0.74-1.6%)	10 (-17-38)	340 (-520-1200)
Somalia	2000	Female	60 to 64	48% (31-65%)	52% (35-69%)	0.15 (0.071-0.27)	0.42% (-0.56-1.2%)	0.48% (-0.61-1.5%)	14 (-17-41)	360 (-470-1100)
Somalia	2000	Female	65 to 69	49% (34-65%)	51% (35-66%)	0.14 (0.07-0.24)	0.48% (-0.36-1.3%)	0.52% (-0.45-1.4%)	15 (-13-42)	360 (-270-940)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	2000	Female	70 to 74	48% (34-62%)	52% (38-66%)	0.12 (0.059-0.21)	0.15% (-0.66-0.83%)	0.12% (-0.77-0.89%)	3.9 (-25-28)	94 (-430-520)
Somalia	2000	Female	75 to 79	46% (29-62%)	54% (38-71%)	0.11 (0.049-0.2)	0.22% (-0.65-0.84%)	0.23% (-0.72-0.93%)	6.4 (-20-26)	95 (-280-360)
Somalia	2000	Female	80 to 84	44% (28-62%)	56% (38-72%)	0.099 (0.045-0.18)	0.0067% (-0.65-0.47%)	0.0027% (-0.73-0.54%)	0.05 (-14-9.9)	1.5 (-150-100)
Somalia	2000	Female	85 to 89	44% (29-59%)	56% (41-71%)	0.093 (0.044-0.17)	-0.051% (-0.61-0.32%)	-0.071% (-0.73-0.35%)	-0.65 (-6.5-3.2)	-3.9 (-46-24)
Somalia	2000	Female	90 to 94	46% (30-60%)	54% (40-70%)	0.091 (0.042-0.17)	-0.1% (-0.63-0.22%)	-0.16% (-0.76-0.24%)	-0.51 (-2.5-0.76)	-2 (-12-4.3)
Somalia	2000	Female	95 plus	48% (30-65%)	52% (35-70%)	0.088 (0.035-0.18)	0.25% (-0.24-0.68%)	0.22% (-0.35-0.73%)	0.22 (-0.36-0.74)	0.98 (-0.94-2.7)
Somalia	2000	Male	All Ages	43% (37-50%)	57% (50-63%)	0.09 (0.06-0.13)	0.47% (-0.058-1%)	0.51% (-0.36-1.4%)	230 (-170-620)	14000 (-1800-29000)
Somalia	2000	Male	15 to 19	42% (29-57%)	58% (43-71%)	0.1 (0.055-0.17)	0.8% (0.12-1.6%)	0.28% (-0.62-1.2%)	2.8 (-6.2-13)	790 (120-1600)
Somalia	2000	Male	20 to 24	53% (39-67%)	47% (33-61%)	0.14 (0.074-0.23)	1.4% (0.038-2.8%)	0.39% (-1.3-2.1%)	5.1 (-17-27)	1600 (45-3200)
Somalia	2000	Male	25 to 29	56% (41-71%)	44% (29-59%)	0.16 (0.093-0.26)	1.6% (-0.29-3.6%)	0.63% (-1.7-3.1%)	8.2 (-23-38)	1600 (-280-3500)
Somalia	2000	Male	30 to 34	58% (43-73%)	42% (27-57%)	0.18 (0.1-0.3)	1.5% (-0.62-3.7%)	0.77% (-1.8-3.4%)	11 (-25-47)	1400 (-580-3500)
Somalia	2000	Male	35 to 39	60% (45-74%)	40% (26-55%)	0.2 (0.11-0.33)	1.5% (-0.7-3.7%)	1.1% (-1.6-3.9%)	16 (-24-56)	1400 (-630-3500)
Somalia	2000	Male	40 to 44	61% (46-76%)	39% (24-54%)	0.2 (0.11-0.35)	1.6% (-0.54-3.8%)	1.4% (-1.2-4.2%)	22 (-20-68)	1400 (-470-3400)
Somalia	2000	Male	45 to 49	61% (46-76%)	39% (24-54%)	0.2 (0.11-0.34)	1.7% (-0.57-3.6%)	1.6% (-1-4%)	27 (-18-70)	1400 (-480-3100)
Somalia	2000	Male	50 to 54	61% (46-76%)	39% (24-54%)	0.2 (0.11-0.34)	1.5% (-0.66-3.5%)	1.5% (-1.1-4%)	27 (-21-73)	1200 (-530-2800)
Somalia	2000	Male	55 to 59	61% (46-75%)	39% (25-54%)	0.19 (0.1-0.33)	1.3% (-0.74-3.1%)	1.3% (-1.1-3.5%)	23 (-21-67)	880 (-490-2200)
Somalia	2000	Male	60 to 64	60% (44-75%)	40% (25-56%)	0.18 (0.091-0.31)	1.3% (-0.033-2.7%)	1.4% (-0.22-3.1%)	30 (-4.7-68)	890 (-23-1900)
Somalia	2000	Male	65 to 69	58% (44-73%)	42% (27-56%)	0.15 (0.077-0.24)	0.86% (-0.25-1.9%)	0.91% (-0.4-2.2%)	20 (-8.3-49)	480 (-140-1100)
Somalia	2000	Male	70 to 74	57% (42-71%)	43% (29-58%)	0.12 (0.061-0.2)	0.65% (-0.16-1.4%)	0.71% (-0.25-1.6%)	16 (-5.1-37)	300 (-72-670)
Somalia	2000	Male	75 to 79	55% (40-71%)	45% (29-60%)	0.099 (0.05-0.17)	0.61% (-0.092-1.2%)	0.66% (-0.14-1.4%)	12 (-2.7-26)	180 (-27-360)
Somalia	2000	Male	80 to 84	54% (37-70%)	46% (30-63%)	0.089 (0.045-0.15)	0.4% (-0.11-0.87%)	0.41% (-0.2-0.97%)	4.5 (-2.2-11)	52 (-14-110)
Somalia	2000	Male	85 to 89	52% (36-66%)	48% (34-64%)	0.079 (0.039-0.14)	0.44% (0.022-0.82%)	0.44% (-0.051-0.89%)	2.7 (-0.31-5.5)	23 (1.1-42)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	2000	Male	90 to 94	52% (37-67%)	48% (33-63%)	0.073 (0.035-0.13)	0.28% (-0.028-0.55%)	0.24% (-0.12-0.57%)	0.59 (-0.28-1.4)	4 (-0.4-8)
Somalia	2000	Male	95 plus	54% (34-71%)	46% (29-66%)	0.068 (0.027-0.14)	0.44% (0.073-0.77%)	0.39% (-0.059-0.79%)	0.27 (-0.042-0.56)	1.2 (0.2-2.2)
Somalia	2005	Female	All Ages	34% (29-41%)	66% (59-71%)	0.081 (0.042-0.14)	0.26% (-0.11-0.62%)	0.26% (-0.39-0.89%)	130 (-180-430)	7200 (-2900-18000)
Somalia	2005	Female	15 to 19	36% (23-51%)	64% (49-77%)	0.099 (0.043-0.19)	0.34% (-0.13-0.87%)	0.11% (-0.65-0.93%)	1 (-5.8-8.9)	350 (-130-910)
Somalia	2005	Female	20 to 24	42% (27-57%)	58% (43-73%)	0.13 (0.059-0.25)	0.62% (-0.063-1.4%)	0.15% (-0.8-1.2%)	1.7 (-9.2-14)	660 (-67-1500)
Somalia	2005	Female	25 to 29	44% (28-61%)	56% (39-72%)	0.16 (0.074-0.31)	0.7% (-0.1-1.6%)	0.33% (-0.73-1.5%)	4.2 (-9.2-18)	750 (-100-1700)
Somalia	2005	Female	30 to 34	46% (30-62%)	54% (38-70%)	0.17 (0.072-0.32)	0.74% (-0.23-1.7%)	0.5% (-0.78-1.8%)	7.1 (-10-24)	780 (-230-1800)
Somalia	2005	Female	35 to 39	48% (33-64%)	52% (36-67%)	0.16 (0.067-0.31)	0.78% (-0.19-1.7%)	0.68% (-0.61-1.9%)	11 (-9.2-32)	840 (-190-1900)
Somalia	2005	Female	40 to 44	50% (36-65%)	50% (35-64%)	0.16 (0.066-0.32)	0.95% (0.031-1.9%)	1% (-0.18-2.2%)	17 (-3.1-40)	960 (30-2000)
Somalia	2005	Female	45 to 49	50% (33-67%)	50% (33-67%)	0.16 (0.067-0.33)	0.83% (-0.3-1.9%)	0.89% (-0.49-2.3%)	17 (-9.7-45)	810 (-300-1900)
Somalia	2005	Female	50 to 54	48% (33-64%)	52% (36-67%)	0.16 (0.068-0.32)	0.81% (-0.49-2.1%)	0.9% (-0.66-2.5%)	21 (-15-56)	810 (-480-2100)
Somalia	2005	Female	55 to 59	47% (31-63%)	53% (37-69%)	0.17 (0.069-0.34)	0.37% (-0.85-1.5%)	0.41% (-1-1.8%)	11 (-26-46)	350 (-800-1400)
Somalia	2005	Female	60 to 64	48% (32-65%)	52% (35-68%)	0.16 (0.067-0.32)	0.38% (-0.69-1.3%)	0.44% (-0.76-1.6%)	13 (-23-43)	330 (-600-1100)
Somalia	2005	Female	65 to 69	49% (34-64%)	51% (36-66%)	0.15 (0.064-0.29)	0.44% (-0.6-1.4%)	0.48% (-0.71-1.6%)	16 (-24-53)	380 (-520-1200)
Somalia	2005	Female	70 to 74	48% (33-63%)	52% (37-67%)	0.13 (0.056-0.26)	0.11% (-0.85-0.94%)	0.075% (-0.97-1%)	2.5 (-32-33)	71 (-560-610)
Somalia	2005	Female	75 to 79	45% (30-62%)	55% (38-70%)	0.12 (0.048-0.24)	0.17% (-0.88-0.88%)	0.17% (-0.98-0.97%)	5.2 (-29-29)	79 (-400-410)
Somalia	2005	Female	80 to 84	44% (28-62%)	56% (38-72%)	0.11 (0.043-0.22)	-0.031% (-0.91-0.55%)	-0.039% (-1-0.6%)	-0.87 (-23-13)	-7.9 (-230-130)
Somalia	2005	Female	85 to 89	44% (29-58%)	56% (42-71%)	0.1 (0.042-0.21)	-0.093% (-0.88-0.4%)	-0.12% (-1-0.44%)	-1.1 (-10-4.3)	-7.4 (-72-33)
Somalia	2005	Female	90 to 94	46% (31-60%)	54% (40-69%)	0.1 (0.039-0.2)	-0.15% (-0.84-0.27%)	-0.21% (-1-0.29%)	-0.7 (-3.4-0.96)	-3 (-17-5.4)
Somalia	2005	Female	95 plus	48% (31-66%)	52% (34-69%)	0.098 (0.034-0.22)	0.22% (-0.46-0.71%)	0.2% (-0.61-0.77%)	0.2 (-0.63-0.78)	0.89 (-1.8-2.8)
Somalia	2005	Male	All Ages	42% (36-50%)	58% (50-64%)	0.097 (0.052-0.17)	0.53% (-0.13-1.2%)	0.56% (-0.53-1.6%)	260 (-250-770)	15000 (-3800-35000)
Somalia	2005	Male	15 to 19	42% (29-58%)	58% (42-71%)	0.11 (0.052-0.22)	0.85% (0.053-1.8%)	0.33% (-0.8-1.5%)	3.7 (-8.8-17)	950 (57-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	2005	Male	20 to 24	53% (39-66%)	47% (34-61%)	0.15 (0.069-0.29)	1.5% (-0.05-3.2%)	0.47% (-1.5-2.7%)	6.8 (-22-37)	1800 (-61-3900)
Somalia	2005	Male	25 to 29	56% (41-70%)	44% (30-59%)	0.18 (0.085-0.34)	1.7% (-0.39-3.8%)	0.71% (-1.9-3.3%)	10 (-27-48)	1800 (-450-4300)
Somalia	2005	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.2 (0.093-0.39)	1.5% (-0.75-3.9%)	0.85% (-2-3.7%)	13 (-30-55)	1600 (-750-4000)
Somalia	2005	Male	35 to 39	60% (46-74%)	40% (26-54%)	0.22 (0.097-0.42)	1.6% (-0.73-3.9%)	1.2% (-1.6-4.1%)	19 (-27-67)	1600 (-750-4100)
Somalia	2005	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.22 (0.096-0.44)	1.6% (-0.72-3.9%)	1.5% (-1.5-4.4%)	25 (-23-76)	1600 (-650-3900)
Somalia	2005	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.22 (0.095-0.46)	1.7% (-0.71-3.9%)	1.6% (-1.4-4.4%)	29 (-25-80)	1500 (-720-3600)
Somalia	2005	Male	50 to 54	61% (46-76%)	39% (24-54%)	0.22 (0.096-0.44)	1.5% (-0.98-3.9%)	1.5% (-1.5-4.3%)	30 (-31-87)	1300 (-880-3300)
Somalia	2005	Male	55 to 59	61% (46-75%)	39% (25-54%)	0.21 (0.094-0.41)	1.3% (-0.99-3.3%)	1.3% (-1.4-3.7%)	27 (-32-84)	1000 (-810-2800)
Somalia	2005	Male	60 to 64	60% (45-75%)	40% (25-55%)	0.19 (0.081-0.37)	1.4% (-0.33-3%)	1.4% (-0.57-3.3%)	32 (-12-76)	950 (-220-2100)
Somalia	2005	Male	65 to 69	58% (44-73%)	42% (27-56%)	0.16 (0.07-0.3)	0.88% (-0.49-2.2%)	0.93% (-0.66-2.4%)	22 (-17-60)	550 (-300-1400)
Somalia	2005	Male	70 to 74	57% (41-71%)	43% (29-59%)	0.13 (0.056-0.25)	0.66% (-0.34-1.7%)	0.72% (-0.44-1.9%)	17 (-11-46)	320 (-170-830)
Somalia	2005	Male	75 to 79	55% (40-70%)	45% (30-60%)	0.11 (0.045-0.21)	0.62% (-0.28-1.5%)	0.67% (-0.38-1.7%)	14 (-7.4-33)	200 (-93-460)
Somalia	2005	Male	80 to 84	53% (36-69%)	47% (31-64%)	0.096 (0.04-0.19)	0.41% (-0.3-1%)	0.42% (-0.41-1.2%)	5.4 (-5.2-15)	63 (-45-160)
Somalia	2005	Male	85 to 89	51% (36-65%)	49% (35-64%)	0.085 (0.037-0.17)	0.45% (-0.12-0.97%)	0.46% (-0.23-1.1%)	3 (-1.5-7)	25 (-6.8-54)
Somalia	2005	Male	90 to 94	52% (37-67%)	48% (33-63%)	0.079 (0.031-0.16)	0.29% (-0.13-0.68%)	0.26% (-0.26-0.72%)	0.65 (-0.64-1.8)	4.4 (-1.8-10)
Somalia	2005	Male	95 plus	53% (34-71%)	47% (29-66%)	0.073 (0.026-0.15)	0.45% (-0.06-0.92%)	0.4% (-0.21-1%)	0.29 (-0.15-0.74)	1.3 (-0.17-2.7)
Somalia	2010	Female	All Ages	35% (29-41%)	65% (59-71%)	0.09 (0.042-0.17)	0.25% (-0.091-0.57%)	0.24% (-0.37-0.82%)	130 (-200-440)	7900 (-2800-18000)
Somalia	2010	Female	15 to 19	36% (23-52%)	64% (48-77%)	0.11 (0.043-0.23)	0.35% (-0.11-0.87%)	0.12% (-0.6-0.95%)	1.3 (-6.4-10)	420 (-130-1100)
Somalia	2010	Female	20 to 24	42% (26-58%)	58% (42-74%)	0.15 (0.058-0.31)	0.64% (0.005-1.3%)	0.17% (-0.73-1.2%)	2.1 (-9-15)	780 (6-1600)
Somalia	2010	Female	25 to 29	44% (29-60%)	56% (40-71%)	0.18 (0.072-0.35)	0.75% (-0.045-1.6%)	0.37% (-0.68-1.5%)	5.1 (-9.2-21)	870 (-48-1900)
Somalia	2010	Female	30 to 34	46% (30-62%)	54% (38-70%)	0.18 (0.071-0.38)	0.77% (-0.17-1.7%)	0.54% (-0.7-1.8%)	8.1 (-9.7-27)	880 (-200-2000)
Somalia	2010	Female	35 to 39	48% (34-64%)	52% (36-66%)	0.17 (0.067-0.37)	0.82% (-0.18-1.8%)	0.72% (-0.56-2%)	12 (-9.4-35)	930 (-200-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	2010	Female	40 to 44	50% (36-66%)	50% (34-64%)	0.17 (0.063-0.37)	0.97% (0.095-1.9%)	1% (-0.12-2.2%)	19 (-2.1-41)	1100 (99-2100)
Somalia	2010	Female	45 to 49	50% (33-66%)	50% (34-67%)	0.18 (0.065-0.39)	0.85% (-0.28-1.9%)	0.91% (-0.51-2.2%)	19 (-11-45)	880 (-310-2000)
Somalia	2010	Female	50 to 54	48% (33-65%)	52% (35-67%)	0.18 (0.066-0.38)	0.82% (-0.48-2.1%)	0.92% (-0.67-2.5%)	22 (-16-60)	870 (-490-2200)
Somalia	2010	Female	55 to 59	47% (32-63%)	53% (37-68%)	0.18 (0.069-0.41)	0.37% (-0.84-1.5%)	0.42% (-0.94-1.8%)	12 (-28-48)	390 (-840-1500)
Somalia	2010	Female	60 to 64	48% (31-65%)	52% (35-69%)	0.18 (0.066-0.38)	0.38% (-0.68-1.3%)	0.44% (-0.79-1.5%)	14 (-24-50)	360 (-670-1300)
Somalia	2010	Female	65 to 69	49% (34-65%)	51% (35-66%)	0.17 (0.065-0.35)	0.41% (-0.68-1.3%)	0.45% (-0.8-1.5%)	16 (-28-52)	360 (-580-1100)
Somalia	2010	Female	70 to 74	48% (34-62%)	52% (38-66%)	0.15 (0.053-0.32)	0.071% (-1-0.87%)	0.042% (-1.1-0.96%)	1.6 (-43-36)	55 (-740-650)
Somalia	2010	Female	75 to 79	46% (30-62%)	54% (38-70%)	0.13 (0.046-0.28)	0.13% (-0.95-0.9%)	0.14% (-1.1-1%)	4.4 (-35-32)	66 (-470-440)
Somalia	2010	Female	80 to 84	45% (28-61%)	55% (39-72%)	0.12 (0.043-0.27)	-0.077% (-1-0.52%)	-0.088% (-1.2-0.57%)	-2.2 (-29-14)	-22 (-290-150)
Somalia	2010	Female	85 to 89	44% (28-59%)	56% (41-72%)	0.11 (0.043-0.25)	-0.13% (-0.96-0.37%)	-0.16% (-1.1-0.41%)	-1.9 (-13-4.7)	-13 (-91-35)
Somalia	2010	Female	90 to 94	46% (30-60%)	54% (40-70%)	0.11 (0.039-0.24)	-0.17% (-0.9-0.26%)	-0.23% (-1.1-0.26%)	-0.86 (-4-0.97)	-3.7 (-20-5.4)
Somalia	2010	Female	95 plus	48% (30-65%)	52% (35-70%)	0.11 (0.034-0.25)	0.17% (-0.52-0.66%)	0.14% (-0.65-0.69%)	0.16 (-0.75-0.77)	0.76 (-2.2-2.9)
Somalia	2010	Male	All Ages	42% (36-50%)	58% (50-64%)	0.11 (0.049-0.21)	0.51% (-0.073-1.1%)	0.55% (-0.44-1.5%)	290 (-240-830)	18000 (-2500-39000)
Somalia	2010	Male	15 to 19	42% (29-57%)	58% (43-71%)	0.12 (0.049-0.25)	0.79% (0.04-1.6%)	0.3% (-0.67-1.4%)	4.3 (-9.5-20)	1100 (58-2400)
Somalia	2010	Male	20 to 24	52% (38-66%)	48% (34-62%)	0.16 (0.068-0.33)	1.4% (-0.13-2.9%)	0.42% (-1.4-2.3%)	7.9 (-26-43)	2100 (-220-4500)
Somalia	2010	Male	25 to 29	56% (41-71%)	44% (29-59%)	0.2 (0.084-0.39)	1.6% (-0.38-3.5%)	0.69% (-1.8-3%)	12 (-31-53)	2100 (-480-4700)
Somalia	2010	Male	30 to 34	58% (43-73%)	42% (27-57%)	0.22 (0.091-0.45)	1.5% (-0.61-3.7%)	0.83% (-1.9-3.5%)	14 (-33-60)	1800 (-740-4500)
Somalia	2010	Male	35 to 39	60% (46-74%)	40% (26-54%)	0.24 (0.096-0.51)	1.6% (-0.59-3.9%)	1.2% (-1.5-4.1%)	22 (-26-73)	1800 (-680-4300)
Somalia	2010	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.24 (0.092-0.52)	1.6% (-0.54-4%)	1.5% (-1.2-4.4%)	29 (-23-83)	1800 (-590-4200)
Somalia	2010	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.24 (0.092-0.52)	1.7% (-0.51-3.7%)	1.7% (-1-4.2%)	33 (-20-85)	1700 (-520-3800)
Somalia	2010	Male	50 to 54	61% (45-75%)	39% (25-55%)	0.24 (0.091-0.5)	1.6% (-0.76-3.7%)	1.6% (-1.2-4.3%)	34 (-28-91)	1500 (-680-3500)
Somalia	2010	Male	55 to 59	61% (46-75%)	39% (25-54%)	0.23 (0.088-0.49)	1.3% (-0.79-3.3%)	1.4% (-1.2-3.7%)	31 (-28-91)	1200 (-670-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	2010	Male	60 to 64	60% (44-76%)	40% (24-56%)	0.21 (0.077-0.45)	1.4% (-0.19-2.9%)	1.5% (-0.46-3.3%)	38 (-11-89)	1100 (-170-2500)
Somalia	2010	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.17 (0.069-0.37)	0.89% (-0.42-2.1%)	0.95% (-0.63-2.4%)	24 (-16-61)	590 (-290-1400)
Somalia	2010	Male	70 to 74	56% (42-70%)	44% (30-58%)	0.14 (0.054-0.29)	0.64% (-0.37-1.6%)	0.7% (-0.47-1.8%)	19 (-13-50)	360 (-210-890)
Somalia	2010	Male	75 to 79	55% (41-70%)	45% (30-59%)	0.12 (0.043-0.25)	0.6% (-0.29-1.4%)	0.66% (-0.4-1.6%)	15 (-8.6-36)	210 (-100-510)
Somalia	2010	Male	80 to 84	53% (36-70%)	47% (30-64%)	0.11 (0.04-0.23)	0.38% (-0.3-1%)	0.39% (-0.41-1.1%)	5.9 (-5.9-17)	69 (-52-180)
Somalia	2010	Male	85 to 89	51% (34-66%)	49% (34-66%)	0.093 (0.035-0.2)	0.41% (-0.19-0.91%)	0.41% (-0.29-1%)	3.4 (-2.5-8.3)	29 (-14-64)
Somalia	2010	Male	90 to 94	52% (36-66%)	48% (34-64%)	0.087 (0.032-0.19)	0.25% (-0.17-0.6%)	0.21% (-0.27-0.61%)	0.65 (-0.83-1.8)	4.4 (-2.9-11)
Somalia	2010	Male	95 plus	53% (33-72%)	47% (28-67%)	0.081 (0.027-0.19)	0.35% (-0.11-0.77%)	0.3% (-0.24-0.79%)	0.28 (-0.22-0.74)	1.3 (-0.38-2.8)
Somalia	2016	Female	All Ages	34% (29-41%)	66% (59-71%)	0.098 (0.041-0.19)	0.32% (-0.1-0.75%)	0.29% (-0.43-0.96%)	150 (-230-500)	9400 (-2800-21000)
Somalia	2016	Female	15 to 19	35% (22-51%)	65% (49-78%)	0.12 (0.043-0.26)	0.37% (-0.11-0.91%)	0.14% (-0.61-1%)	1.7 (-7.4-12)	520 (-150-1300)
Somalia	2016	Female	20 to 24	42% (26-58%)	58% (42-74%)	0.16 (0.059-0.34)	0.72% (0.037-1.5%)	0.21% (-0.81-1.3%)	2.7 (-11-17)	970 (49-2000)
Somalia	2016	Female	25 to 29	44% (28-60%)	56% (40-72%)	0.19 (0.076-0.4)	0.85% (0.024-1.8%)	0.45% (-0.74-1.7%)	6.5 (-11-25)	1100 (30-2300)
Somalia	2016	Female	30 to 34	46% (30-61%)	54% (39-70%)	0.2 (0.073-0.42)	0.88% (-0.069-1.9%)	0.64% (-0.68-2%)	10 (-11-32)	1100 (-84-2300)
Somalia	2016	Female	35 to 39	48% (33-63%)	52% (37-67%)	0.19 (0.066-0.42)	0.91% (-0.21-2%)	0.83% (-0.63-2.3%)	15 (-10-42)	1100 (-250-2400)
Somalia	2016	Female	40 to 44	50% (36-65%)	50% (35-64%)	0.19 (0.062-0.43)	1.1% (0.099-2.1%)	1.1% (-0.12-2.5%)	22 (-2.1-50)	1200 (100-2500)
Somalia	2016	Female	45 to 49	50% (33-65%)	50% (35-67%)	0.19 (0.065-0.42)	0.92% (-0.24-2%)	1% (-0.44-2.4%)	22 (-9.6-53)	1000 (-250-2300)
Somalia	2016	Female	50 to 54	48% (33-64%)	52% (36-67%)	0.19 (0.064-0.42)	0.89% (-0.45-2.2%)	1% (-0.64-2.7%)	26 (-16-66)	1000 (-530-2400)
Somalia	2016	Female	55 to 59	47% (32-62%)	53% (38-68%)	0.2 (0.067-0.44)	0.41% (-0.9-1.7%)	0.47% (-1.1-2%)	14 (-31-58)	450 (-970-1800)
Somalia	2016	Female	60 to 64	47% (31-64%)	53% (36-69%)	0.2 (0.066-0.46)	0.4% (-0.72-1.4%)	0.47% (-0.84-1.7%)	16 (-28-56)	420 (-760-1500)
Somalia	2016	Female	65 to 69	49% (33-65%)	51% (35-67%)	0.18 (0.063-0.41)	0.41% (-0.67-1.3%)	0.45% (-0.77-1.5%)	17 (-29-57)	400 (-650-1300)
Somalia	2016	Female	70 to 74	48% (33-63%)	52% (37-67%)	0.16 (0.055-0.36)	0.053% (-1.1-0.9%)	0.019% (-1.3-0.97%)	0.68 (-51-37)	41 (-870-700)
Somalia	2016	Female	75 to 79	46% (29-62%)	54% (38-71%)	0.14 (0.045-0.32)	0.11% (-1.1-0.9%)	0.11% (-1.2-1%)	4 (-47-38)	62 (-640-510)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Somalia	2016	Female	80 to 84	44% (28-61%)	56% (39-72%)	0.13 (0.042-0.29)	-0.11% (-1.1-0.54%)	-0.13% (-1.3-0.6%)	-3.5 (-34-17)	-36 (-360-170)
Somalia	2016	Female	85 to 89	44% (29-59%)	56% (41-71%)	0.12 (0.042-0.3)	-0.17% (-1-0.39%)	-0.2% (-1.2-0.45%)	-2.4 (-14-5.2)	-17 (-110-39)
Somalia	2016	Female	90 to 94	46% (30-60%)	54% (40-70%)	0.12 (0.037-0.29)	-0.24% (-1.1-0.27%)	-0.31% (-1.3-0.28%)	-1.2 (-4.9-1.1)	-5.3 (-24-6.3)
Somalia	2016	Female	95 plus	48% (30-65%)	52% (35-70%)	0.12 (0.035-0.3)	0.16% (-0.66-0.76%)	0.12% (-0.84-0.81%)	0.12 (-0.79-0.77)	0.61 (-2.4-2.8)
Somalia	2016	Male	All Ages	42% (36-50%)	58% (50-64%)	0.12 (0.049-0.23)	0.7% (-0.051-1.5%)	0.72% (-0.47-1.9%)	360 (-220-990)	22000 (-1600-45000)
Somalia	2016	Male	15 to 19	42% (29-56%)	58% (44-71%)	0.14 (0.048-0.29)	0.88% (0.066-1.7%)	0.35% (-0.68-1.5%)	5.7 (-11-24)	1400 (110-2900)
Somalia	2016	Male	20 to 24	52% (38-66%)	48% (34-62%)	0.18 (0.069-0.37)	1.5% (0.032-3.1%)	0.51% (-1.4-2.4%)	11 (-30-52)	2700 (55-5600)
Somalia	2016	Male	25 to 29	56% (41-70%)	44% (30-59%)	0.22 (0.086-0.44)	1.8% (-0.27-3.8%)	0.83% (-1.8-3.5%)	16 (-35-68)	2700 (-430-5800)
Somalia	2016	Male	30 to 34	58% (43-72%)	42% (28-57%)	0.24 (0.094-0.5)	1.7% (-0.5-4%)	1% (-1.9-4%)	19 (-36-75)	2300 (-640-5400)
Somalia	2016	Male	35 to 39	60% (46-74%)	40% (26-54%)	0.26 (0.094-0.55)	1.8% (-0.52-4.2%)	1.4% (-1.5-4.6%)	27 (-27-85)	2200 (-580-5100)
Somalia	2016	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.27 (0.088-0.58)	1.9% (-0.36-4.2%)	1.8% (-1.1-4.7%)	35 (-21-94)	2100 (-390-4800)
Somalia	2016	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.27 (0.087-0.58)	1.9% (-0.33-4.2%)	1.9% (-0.96-4.8%)	41 (-20-100)	2100 (-330-4500)
Somalia	2016	Male	50 to 54	61% (46-75%)	39% (25-54%)	0.26 (0.086-0.56)	1.8% (-0.6-4.2%)	1.8% (-1.1-4.8%)	41 (-26-110)	1800 (-600-4200)
Somalia	2016	Male	55 to 59	60% (46-75%)	40% (25-54%)	0.25 (0.088-0.55)	1.5% (-0.64-3.7%)	1.6% (-1.1-4.3%)	38 (-27-110)	1400 (-620-3500)
Somalia	2016	Male	60 to 64	60% (44-76%)	40% (24-56%)	0.23 (0.079-0.5)	1.5% (-0.037-3.2%)	1.6% (-0.25-3.6%)	45 (-6.9-100)	1300 (-31-2800)
Somalia	2016	Male	65 to 69	58% (44-74%)	42% (26-56%)	0.19 (0.067-0.42)	0.96% (-0.44-2.3%)	1% (-0.65-2.6%)	30 (-18-75)	730 (-330-1700)
Somalia	2016	Male	70 to 74	57% (42-71%)	43% (29-58%)	0.15 (0.054-0.32)	0.69% (-0.44-1.7%)	0.75% (-0.59-1.9%)	21 (-17-56)	400 (-270-1000)
Somalia	2016	Male	75 to 79	55% (41-70%)	45% (30-59%)	0.13 (0.044-0.28)	0.65% (-0.34-1.5%)	0.71% (-0.43-1.7%)	17 (-11-43)	260 (-130-600)
Somalia	2016	Male	80 to 84	53% (36-69%)	47% (31-64%)	0.12 (0.039-0.25)	0.41% (-0.4-1.1%)	0.43% (-0.5-1.2%)	7 (-8.5-21)	81 (-80-220)
Somalia	2016	Male	85 to 89	51% (35-66%)	49% (34-65%)	0.1 (0.035-0.23)	0.45% (-0.22-1%)	0.46% (-0.34-1.1%)	3.9 (-2.8-9.5)	32 (-15-72)
Somalia	2016	Male	90 to 94	52% (36-67%)	48% (33-64%)	0.094 (0.031-0.22)	0.3% (-0.2-0.73%)	0.26% (-0.34-0.78%)	0.8 (-1.1-2.4)	5.4 (-3.6-14)
Somalia	2016	Male	95 plus	53% (33-72%)	47% (28-67%)	0.087 (0.027-0.21)	0.45% (-0.18-1%)	0.4% (-0.34-1.1%)	0.31 (-0.26-0.83)	1.4 (-0.53-3.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	1990	Female	All Ages	20% (16-26%)	80% (74-84%)	1.4 (1-1.9)	1.5% (1.1-2%)	3.2% (2.2-4.5%)	5200 (3500-7300)	170000 (130000-220000)
Tanzania	1990	Female	15 to 19	21% (12-34%)	79% (66-88%)	1.6 (0.75-2.9)	2.7% (1.4-4.4%)	3.6% (1.7-6.1%)	100 (48-170)	8700 (4600-14000)
Tanzania	1990	Female	20 to 24	26% (14-42%)	74% (58-86%)	2.3 (1.1-4.1)	3.1% (1.8-4.9%)	3.5% (1.8-5.9%)	130 (66-220)	11000 (6500-17000)
Tanzania	1990	Female	25 to 29	28% (16-45%)	72% (55-84%)	2.8 (1.5-4.8)	3.7% (2.5-5.4%)	4.3% (2.6-6.6%)	170 (100-250)	13000 (8300-18000)
Tanzania	1990	Female	30 to 34	31% (17-48%)	69% (52-83%)	3.1 (1.6-5.4)	4.5% (3-6.4%)	5.4% (3.3-8%)	200 (120-300)	13000 (8900-19000)
Tanzania	1990	Female	35 to 39	34% (19-52%)	66% (48-81%)	3.1 (1.5-5.2)	5.1% (3.2-7.3%)	6.4% (3.9-9.5%)	240 (140-340)	13000 (8500-19000)
Tanzania	1990	Female	40 to 44	36% (22-54%)	64% (46-78%)	3.1 (1.5-5.4)	5.5% (3.4-7.8%)	7.1% (4.2-10%)	270 (160-370)	13000 (8200-18000)
Tanzania	1990	Female	45 to 49	36% (20-53%)	64% (47-80%)	3.2 (1.5-5.8)	6.6% (4.1-9.6%)	8.6% (5.2-13%)	330 (200-490)	14000 (8900-21000)
Tanzania	1990	Female	50 to 54	35% (19-55%)	65% (45-81%)	3.2 (1.4-5.8)	7.8% (4.7-11%)	9.9% (6-14%)	430 (260-630)	16000 (10000-24000)
Tanzania	1990	Female	55 to 59	33% (19-52%)	67% (48-81%)	3.1 (1.4-5.8)	8% (4.6-12%)	9.9% (5.7-15%)	500 (290-760)	16000 (9300-24000)
Tanzania	1990	Female	60 to 64	34% (19-53%)	66% (47-81%)	3.1 (1.3-6)	8.2% (4.6-12%)	9.8% (5.6-15%)	580 (320-900)	16000 (9000-24000)
Tanzania	1990	Female	65 to 69	35% (20-53%)	65% (47-80%)	2.9 (1.3-5.4)	8.8% (4.9-13%)	10% (5.6-16%)	700 (390-1100)	16000 (8900-25000)
Tanzania	1990	Female	70 to 74	34% (20-52%)	66% (48-80%)	2.5 (1.1-5)	7.7% (3.6-13%)	8.8% (4.1-15%)	630 (280-1100)	12000 (5300-19000)
Tanzania	1990	Female	75 to 79	31% (18-50%)	69% (50-82%)	2.1 (0.79-4.2)	6.5% (2.7-11%)	7.4% (3-13%)	460 (180-820)	6500 (2600-12000)
Tanzania	1990	Female	80 to 84	29% (15-48%)	71% (52-85%)	1.9 (0.73-3.7)	5.5% (2.2-10%)	6.3% (2.6-11%)	260 (100-480)	2700 (1100-5000)
Tanzania	1990	Female	85 to 89	29% (16-46%)	71% (54-84%)	1.8 (0.73-3.6)	4.6% (1.6-9%)	5.2% (1.8-10%)	130 (45-260)	970 (340-1900)
Tanzania	1990	Female	90 to 94	32% (18-50%)	68% (50-82%)	1.9 (0.75-4)	4% (1.3-8%)	4.7% (1.4-9.3%)	46 (14-91)	230 (73-460)
Tanzania	1990	Female	95 plus	36% (18-59%)	64% (41-82%)	2 (0.66-4.5)	4.3% (1.3-9%)	4.9% (1.4-10%)	13 (3.7-28)	45 (13-95)
Tanzania	1990	Male	All Ages	29% (23-37%)	71% (63-77%)	2 (1.5-2.7)	2.6% (2-3.2%)	4.4% (3.2-5.6%)	7500 (5500-9600)	320000 (240000-390000)
Tanzania	1990	Male	15 to 19	27% (16-43%)	73% (57-84%)	2.1 (1.1-3.6)	5.6% (3.3-8.5%)	7.3% (3.9-12%)	190 (100-310)	16000 (10000-26000)
Tanzania	1990	Male	20 to 24	38% (23-55%)	62% (45-77%)	3.1 (1.6-5.2)	8.7% (5.4-12%)	9.7% (5.5-15%)	390 (230-590)	31000 (20000-45000)
Tanzania	1990	Male	25 to 29	44% (26-61%)	56% (39-74%)	3.9 (2.2-6.3)	9% (6-13%)	9.9% (6.1-15%)	510 (330-710)	36000 (25000-48000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	1990	Male	30 to 34	48% (28-66%)	52% (34-72%)	4.6 (2.6-7.3)	9.6% (6.6-13%)	11% (6.8-15%)	550 (370-760)	35000 (24000-47000)
Tanzania	1990	Male	35 to 39	50% (32-68%)	50% (32-68%)	5.1 (2.8-8.3)	11% (7.1-15%)	12% (7.8-18%)	610 (410-820)	34000 (23000-44000)
Tanzania	1990	Male	40 to 44	51% (34-70%)	49% (30-66%)	5.3 (2.6-8.8)	11% (7.8-15%)	13% (8.9-18%)	650 (440-880)	32000 (22000-43000)
Tanzania	1990	Male	45 to 49	52% (33-70%)	48% (30-67%)	5.3 (2.7-8.9)	12% (8.3-16%)	14% (9.8-19%)	690 (470-920)	30000 (21000-40000)
Tanzania	1990	Male	50 to 54	52% (33-70%)	48% (30-67%)	5.2 (2.7-9)	13% (8.8-17%)	16% (11-21%)	750 (490-1000)	28000 (19000-38000)
Tanzania	1990	Male	55 to 59	52% (34-70%)	48% (30-66%)	5 (2.5-8.6)	13% (8.3-17%)	15% (10-20%)	750 (470-1000)	25000 (16000-33000)
Tanzania	1990	Male	60 to 64	50% (31-69%)	50% (31-69%)	4.4 (2-8)	11% (7-16%)	13% (8.1-18%)	720 (430-1000)	20000 (12000-28000)
Tanzania	1990	Male	65 to 69	47% (29-66%)	53% (34-71%)	3.5 (1.7-6.2)	9.3% (5.1-13%)	11% (6-16%)	610 (320-880)	14000 (7600-20000)
Tanzania	1990	Male	70 to 74	44% (27-62%)	56% (38-73%)	2.8 (1.3-5)	7.4% (3.5-12%)	8.5% (4-14%)	480 (220-800)	8700 (4100-15000)
Tanzania	1990	Male	75 to 79	42% (26-60%)	58% (40-74%)	2.3 (1-4.3)	5.9% (2.4-11%)	6.8% (2.8-12%)	320 (130-570)	4500 (1800-8100)
Tanzania	1990	Male	80 to 84	39% (23-58%)	61% (42-77%)	1.9 (0.86-3.6)	4.9% (1.9-9.2%)	5.6% (2.1-11%)	160 (59-300)	1700 (640-3100)
Tanzania	1990	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.6 (0.74-3)	3.7% (1.5-7.3%)	4.2% (1.6-8.2%)	68 (26-130)	520 (210-1000)
Tanzania	1990	Male	90 to 94	38% (23-56%)	62% (44-77%)	1.5 (0.66-2.9)	2.8% (0.88-5.9%)	3.1% (0.86-6.7%)	19 (5.3-41)	100 (32-210)
Tanzania	1990	Male	95 plus	41% (21-63%)	59% (37-79%)	1.5 (0.52-3)	2.8% (0.89-6.2%)	3% (0.78-6.8%)	4.3 (1.1-9.9)	16 (5.2-36)
Tanzania	1995	Female	All Ages	21% (16-27%)	79% (73-84%)	1.3 (0.94-1.8)	1.5% (1-2%)	3.1% (1.9-4.3%)	6100 (3700-8600)	200000 (140000-270000)
Tanzania	1995	Female	15 to 19	21% (12-35%)	79% (65-88%)	1.5 (0.67-2.7)	2.4% (1.2-4%)	3.2% (1.4-5.6%)	100 (45-180)	9200 (4600-15000)
Tanzania	1995	Female	20 to 24	27% (14-42%)	73% (58-86%)	2.1 (1-3.7)	2.5% (1.4-3.9%)	2.6% (1.2-4.5%)	140 (67-240)	12000 (7100-19000)
Tanzania	1995	Female	25 to 29	29% (16-46%)	71% (54-84%)	2.6 (1.3-4.3)	2.6% (1.6-4%)	2.6% (1.5-4.5%)	190 (110-290)	14000 (9300-21000)
Tanzania	1995	Female	30 to 34	32% (17-50%)	68% (50-83%)	2.8 (1.5-4.9)	2.9% (1.8-4.3%)	3.1% (1.8-4.9%)	230 (140-350)	15000 (9700-22000)
Tanzania	1995	Female	35 to 39	35% (20-54%)	65% (46-80%)	2.8 (1.4-4.8)	3.5% (2-5.3%)	3.9% (2.1-6.3%)	260 (150-400)	15000 (9400-22000)
Tanzania	1995	Female	40 to 44	37% (22-55%)	63% (45-78%)	2.8 (1.3-4.9)	4% (2.4-6%)	4.7% (2.7-7.3%)	290 (170-430)	15000 (8700-21000)
Tanzania	1995	Female	45 to 49	37% (20-55%)	63% (45-80%)	2.9 (1.3-5.1)	5.3% (3.1-7.9%)	6.4% (3.6-9.6%)	400 (230-620)	17000 (10000-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	1995	Female	50 to 54	35% (19-55%)	65% (45-81%)	2.8 (1.3-5)	6.6% (3.8-9.9%)	8% (4.6-12%)	510 (280-790)	19000 (11000-30000)
Tanzania	1995	Female	55 to 59	34% (19-54%)	66% (46-81%)	2.8 (1.3-5.4)	7.2% (4.1-11%)	8.7% (4.8-13%)	600 (330-920)	19000 (11000-29000)
Tanzania	1995	Female	60 to 64	35% (19-53%)	65% (47-81%)	2.8 (1.2-5.4)	7.6% (4-12%)	8.9% (4.8-14%)	700 (370-1100)	19000 (10000-30000)
Tanzania	1995	Female	65 to 69	36% (20-54%)	64% (46-80%)	2.7 (1.2-4.9)	8.2% (4.3-13%)	9.5% (5-15%)	830 (420-1300)	19000 (9700-30000)
Tanzania	1995	Female	70 to 74	35% (20-52%)	65% (48-80%)	2.3 (0.96-4.4)	7.2% (3.1-12%)	8.1% (3.5-14%)	730 (300-1300)	13000 (5700-23000)
Tanzania	1995	Female	75 to 79	32% (18-49%)	68% (51-82%)	1.9 (0.74-3.8)	6.1% (2.3-11%)	6.9% (2.5-13%)	550 (190-1000)	7700 (2800-14000)
Tanzania	1995	Female	80 to 84	30% (16-48%)	70% (52-84%)	1.7 (0.66-3.4)	5.1% (1.8-9.6%)	5.8% (2.1-11%)	320 (110-600)	3300 (1200-6200)
Tanzania	1995	Female	85 to 89	30% (16-48%)	70% (52-84%)	1.7 (0.67-3.3)	4.1% (1.3-8.5%)	4.7% (1.5-9.8%)	140 (43-300)	1000 (330-2200)
Tanzania	1995	Female	90 to 94	33% (18-50%)	67% (50-82%)	1.7 (0.68-3.4)	3.6% (0.95-7.6%)	4.1% (1.1-9%)	48 (12-100)	250 (65-530)
Tanzania	1995	Female	95 plus	37% (18-59%)	63% (41-82%)	1.8 (0.59-4.2)	3.8% (1.1-8.5%)	4.3% (1.2-9.7%)	14 (3.9-32)	49 (14-110)
Tanzania	1995	Male	All Ages	30% (24-38%)	70% (62-76%)	1.8 (1.3-2.4)	2.5% (1.8-3.2%)	4% (2.8-5.4%)	8300 (5800-11000)	350000 (260000-460000)
Tanzania	1995	Male	15 to 19	27% (16-43%)	73% (57-84%)	1.9 (0.92-3.3)	5.1% (3-7.9%)	6.6% (3.4-11%)	200 (100-340)	18000 (10000-28000)
Tanzania	1995	Male	20 to 24	39% (23-56%)	61% (44-77%)	2.8 (1.5-4.7)	7.7% (4.5-11%)	8.4% (4.4-13%)	440 (240-680)	35000 (22000-52000)
Tanzania	1995	Male	25 to 29	44% (26-62%)	56% (38-74%)	3.5 (2-5.6)	6.7% (4.1-9.9%)	7% (3.8-11%)	570 (340-830)	40000 (26000-56000)
Tanzania	1995	Male	30 to 34	48% (29-67%)	52% (33-71%)	4.1 (2.4-6.6)	6.3% (3.9-9.1%)	6.5% (3.8-10%)	630 (390-900)	39000 (26000-54000)
Tanzania	1995	Male	35 to 39	51% (33-68%)	49% (32-67%)	4.6 (2.5-7.5)	7.1% (4.4-11%)	7.7% (4.5-12%)	690 (430-980)	38000 (25000-53000)
Tanzania	1995	Male	40 to 44	52% (35-70%)	48% (30-65%)	4.7 (2.4-7.8)	7.7% (5-11%)	8.5% (5.3-12%)	720 (460-1000)	35000 (23000-50000)
Tanzania	1995	Male	45 to 49	52% (33-70%)	48% (30-67%)	4.7 (2.3-7.9)	8.8% (5.6-12%)	9.9% (6.2-14%)	780 (490-1100)	34000 (22000-47000)
Tanzania	1995	Male	50 to 54	53% (33-71%)	47% (29-67%)	4.6 (2.3-7.9)	11% (6.6-14%)	12% (7.7-17%)	840 (510-1200)	32000 (20000-45000)
Tanzania	1995	Male	55 to 59	52% (34-72%)	48% (28-66%)	4.5 (2.2-7.5)	11% (7-15%)	13% (8.2-18%)	850 (510-1200)	28000 (17000-39000)
Tanzania	1995	Male	60 to 64	50% (32-70%)	50% (30-68%)	3.9 (1.9-7.2)	10% (5.7-14%)	12% (6.7-16%)	810 (460-1200)	22000 (13000-33000)
Tanzania	1995	Male	65 to 69	48% (30-66%)	52% (34-70%)	3.2 (1.5-5.5)	8.2% (4.3-12%)	9.5% (4.9-14%)	670 (340-1000)	15000 (8000-24000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	1995	Male	70 to 74	45% (28-62%)	55% (38-72%)	2.5 (1.2-4.6)	6.4% (2.8-11%)	7.3% (3.2-13%)	510 (220-890)	9300 (4000-16000)
Tanzania	1995	Male	75 to 79	42% (26-61%)	58% (39-74%)	2 (0.87-3.8)	5.1% (1.9-9.6%)	5.8% (2.2-11%)	340 (130-650)	4800 (1800-9100)
Tanzania	1995	Male	80 to 84	40% (23-59%)	60% (41-77%)	1.7 (0.75-3.2)	4.2% (1.5-8.2%)	4.8% (1.7-9.2%)	170 (60-330)	1800 (650-3400)
Tanzania	1995	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.4 (0.65-2.7)	3.1% (1.2-6.5%)	3.5% (1.3-7.3%)	69 (25-140)	530 (200-1100)
Tanzania	1995	Male	90 to 94	38% (23-55%)	62% (45-77%)	1.3 (0.57-2.5)	2.3% (0.67-5.2%)	2.5% (0.62-5.9%)	19 (4.7-45)	100 (31-240)
Tanzania	1995	Male	95 plus	41% (21-63%)	59% (37-79%)	1.3 (0.44-2.7)	2.3% (0.69-5.6%)	2.5% (0.53-6.3%)	4.5 (0.98-11)	17 (5-41)
Tanzania	2000	Female	All Ages	21% (17-27%)	79% (73-83%)	1.2 (0.88-1.7)	1.6% (1-2.2%)	3.1% (1.8-4.5%)	6700 (3900-9800)	220000 (150000-310000)
Tanzania	2000	Female	15 to 19	21% (12-35%)	79% (65-88%)	1.4 (0.65-2.6)	2.4% (1.3-4.1%)	3.3% (1.4-5.9%)	110 (50-200)	10000 (5300-17000)
Tanzania	2000	Female	20 to 24	27% (14-42%)	73% (58-86%)	2 (0.96-3.6)	2.6% (1.4-4.1%)	2.8% (1.2-4.7%)	160 (78-260)	14000 (8100-21000)
Tanzania	2000	Female	25 to 29	29% (17-45%)	71% (55-83%)	2.5 (1.2-4.1)	2.2% (1.3-3.6%)	2.2% (1.1-4%)	220 (120-340)	16000 (10000-24000)
Tanzania	2000	Female	30 to 34	32% (18-50%)	68% (50-82%)	2.7 (1.3-4.7)	2.3% (1.3-3.5%)	2.3% (1.3-3.8%)	260 (150-400)	17000 (11000-25000)
Tanzania	2000	Female	35 to 39	35% (20-54%)	65% (46-80%)	2.7 (1.3-4.5)	2.6% (1.4-4.2%)	2.8% (1.4-4.7%)	290 (160-450)	16000 (9700-25000)
Tanzania	2000	Female	40 to 44	37% (22-56%)	63% (44-78%)	2.7 (1.2-4.6)	3.2% (1.7-4.8%)	3.5% (1.8-5.5%)	320 (170-500)	16000 (8800-25000)
Tanzania	2000	Female	45 to 49	37% (21-55%)	63% (45-79%)	2.8 (1.2-5)	4.5% (2.5-7%)	5.2% (2.8-8.2%)	420 (230-680)	18000 (10000-29000)
Tanzania	2000	Female	50 to 54	36% (20-56%)	64% (44-80%)	2.7 (1.2-4.9)	6% (3.4-9.2%)	7.1% (3.9-11%)	580 (300-950)	22000 (11000-35000)
Tanzania	2000	Female	55 to 59	34% (20-54%)	66% (46-80%)	2.7 (1.2-5)	7% (3.7-11%)	8.3% (4.4-13%)	680 (340-1100)	22000 (11000-35000)
Tanzania	2000	Female	60 to 64	35% (19-54%)	65% (46-81%)	2.7 (1.2-5.4)	7.3% (3.9-11%)	8.6% (4.6-14%)	760 (380-1300)	21000 (11000-34000)
Tanzania	2000	Female	65 to 69	36% (21-54%)	64% (46-79%)	2.5 (1.1-4.7)	7.9% (4.2-13%)	9.2% (4.8-15%)	910 (450-1500)	21000 (10000-33000)
Tanzania	2000	Female	70 to 74	34% (20-53%)	66% (47-80%)	2.2 (0.91-4.2)	6.9% (3-12%)	7.8% (3.3-13%)	800 (340-1400)	15000 (6300-26000)
Tanzania	2000	Female	75 to 79	32% (18-51%)	68% (49-82%)	1.8 (0.71-3.6)	5.9% (2.2-11%)	6.7% (2.5-12%)	610 (220-1100)	8600 (3200-16000)
Tanzania	2000	Female	80 to 84	30% (16-49%)	70% (51-84%)	1.6 (0.61-3.2)	4.9% (1.7-9.3%)	5.6% (2-11%)	380 (130-730)	4000 (1300-7600)
Tanzania	2000	Female	85 to 89	31% (17-47%)	69% (53-83%)	1.6 (0.64-3.1)	4% (1.3-7.9%)	4.6% (1.4-9.2%)	160 (49-320)	1200 (370-2400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	2000	Female	90 to 94	33% (19-51%)	67% (49-81%)	1.6 (0.63-3.2)	3.5% (0.9-7.4%)	4% (0.96-8.7%)	50 (12-110)	260 (65-550)
Tanzania	2000	Female	95 plus	38% (19-59%)	62% (41-81%)	1.7 (0.59-3.9)	3.7% (1-8.6%)	4.2% (1.1-9.8%)	15 (3.7-35)	52 (14-120)
Tanzania	2000	Male	All Ages	30% (24-38%)	70% (62-76%)	1.8 (1.3-2.4)	2.8% (2.1-3.6%)	4.4% (3.1-5.8%)	9700 (6800-13000)	410000 (300000-540000)
Tanzania	2000	Male	15 to 19	27% (16-42%)	73% (58-84%)	1.8 (0.9-3.1)	5.2% (3.1-8.1%)	6.7% (3.5-11%)	230 (120-380)	20000 (12000-32000)
Tanzania	2000	Male	20 to 24	39% (24-55%)	61% (45-76%)	2.6 (1.4-4.4)	8.6% (5.4-12%)	9.6% (5.3-15%)	520 (300-820)	42000 (26000-62000)
Tanzania	2000	Male	25 to 29	45% (27-63%)	55% (37-73%)	3.4 (1.8-5.2)	7.1% (4.3-11%)	7.4% (4-12%)	680 (430-1000)	48000 (32000-68000)
Tanzania	2000	Male	30 to 34	48% (29-67%)	52% (33-71%)	3.9 (2.2-6.2)	6% (3.9-8.8%)	6.2% (3.7-9.5%)	750 (480-1100)	47000 (32000-65000)
Tanzania	2000	Male	35 to 39	51% (34-70%)	49% (30-66%)	4.3 (2.3-7.2)	6.1% (3.8-9.1%)	6.5% (3.8-10%)	800 (510-1200)	44000 (29000-62000)
Tanzania	2000	Male	40 to 44	52% (36-71%)	48% (29-64%)	4.5 (2.2-7.4)	6.7% (4.4-9.8%)	7.2% (4.6-11%)	840 (540-1200)	41000 (26000-58000)
Tanzania	2000	Male	45 to 49	53% (34-71%)	47% (29-66%)	4.5 (2.1-7.6)	7.7% (4.9-11%)	8.6% (5.2-12%)	870 (550-1300)	37000 (24000-54000)
Tanzania	2000	Male	50 to 54	53% (35-71%)	47% (29-65%)	4.4 (2.2-7.6)	9.9% (6.4-13%)	11% (7.2-15%)	980 (620-1400)	37000 (24000-52000)
Tanzania	2000	Male	55 to 59	52% (35-71%)	48% (29-65%)	4.3 (2.1-7.4)	11% (6.8-15%)	13% (7.9-18%)	1000 (600-1400)	32000 (20000-46000)
Tanzania	2000	Male	60 to 64	50% (32-69%)	50% (31-68%)	3.7 (1.7-6.9)	9.9% (5.6-14%)	11% (6.4-17%)	930 (500-1400)	26000 (14000-38000)
Tanzania	2000	Male	65 to 69	48% (30-66%)	52% (34-70%)	3 (1.5-5.3)	8.2% (4.3-12%)	9.5% (5-15%)	780 (400-1200)	18000 (9300-28000)
Tanzania	2000	Male	70 to 74	45% (28-62%)	55% (38-72%)	2.4 (1.2-4.2)	6.3% (2.8-11%)	7.3% (3.2-13%)	590 (260-1100)	11000 (4800-19000)
Tanzania	2000	Male	75 to 79	43% (26-61%)	57% (39-74%)	1.9 (0.84-3.5)	5.1% (1.9-9.9%)	5.9% (2.1-11%)	400 (150-790)	5700 (2200-11000)
Tanzania	2000	Male	80 to 84	40% (23-58%)	60% (42-77%)	1.6 (0.72-3)	4.3% (1.5-8.3%)	4.9% (1.7-9.4%)	210 (73-410)	2200 (790-4300)
Tanzania	2000	Male	85 to 89	38% (23-54%)	62% (46-77%)	1.4 (0.64-2.6)	3.3% (1.2-6.6%)	3.7% (1.3-7.5%)	84 (30-170)	640 (240-1300)
Tanzania	2000	Male	90 to 94	38% (23-57%)	62% (43-77%)	1.3 (0.54-2.4)	2.4% (0.73-5.3%)	2.7% (0.72-6%)	23 (6.2-52)	120 (38-270)
Tanzania	2000	Male	95 plus	42% (21-64%)	58% (36-79%)	1.3 (0.43-2.7)	2.6% (0.83-5.8%)	2.8% (0.73-6.7%)	6 (1.6-14)	22 (7.3-51)
Tanzania	2005	Female	All Ages	21% (16-27%)	79% (73-84%)	1.3 (0.88-1.7)	1.6% (1-2.2%)	3.1% (1.8-4.4%)	6600 (3700-9700)	220000 (140000-300000)
Tanzania	2005	Female	15 to 19	21% (12-35%)	79% (65-88%)	1.4 (0.64-2.7)	2.2% (1.2-3.7%)	2.9% (1.3-5%)	110 (50-200)	10000 (5300-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	2005	Female	20 to 24	27% (14-42%)	73% (58-86%)	2 (0.97-3.6)	2.6% (1.5-4.1%)	2.8% (1.3-5%)	150 (74-260)	14000 (8300-22000)
Tanzania	2005	Female	25 to 29	29% (16-45%)	71% (55-84%)	2.4 (1.2-4.2)	2.1% (1.2-3.4%)	2.1% (1-3.8%)	200 (110-320)	16000 (10000-23000)
Tanzania	2005	Female	30 to 34	31% (17-49%)	69% (51-83%)	2.7 (1.3-4.9)	1.9% (1.2-3%)	1.9% (1.1-3.2%)	240 (140-370)	16000 (10000-24000)
Tanzania	2005	Female	35 to 39	35% (20-53%)	65% (47-80%)	2.7 (1.3-4.7)	2.1% (1.2-3.3%)	2.2% (1.2-3.7%)	270 (150-410)	15000 (9300-23000)
Tanzania	2005	Female	40 to 44	36% (22-55%)	64% (45-78%)	2.7 (1.2-4.7)	2.5% (1.4-3.9%)	2.7% (1.5-4.3%)	290 (160-460)	15000 (8400-23000)
Tanzania	2005	Female	45 to 49	36% (20-54%)	64% (46-80%)	2.8 (1.2-5)	3.7% (1.9-5.8%)	4.2% (2.2-6.8%)	390 (200-640)	17000 (9300-27000)
Tanzania	2005	Female	50 to 54	35% (19-54%)	65% (46-81%)	2.7 (1.2-5)	5.4% (2.9-8.4%)	6.4% (3.5-9.9%)	550 (280-900)	21000 (11000-33000)
Tanzania	2005	Female	55 to 59	33% (19-53%)	67% (47-81%)	2.7 (1.2-5.1)	6.7% (3.6-10%)	8% (4.4-12%)	680 (340-1100)	22000 (11000-36000)
Tanzania	2005	Female	60 to 64	34% (19-53%)	66% (47-81%)	2.7 (1.2-5.4)	7.1% (3.8-11%)	8.5% (4.6-13%)	790 (390-1300)	22000 (11000-36000)
Tanzania	2005	Female	65 to 69	35% (20-53%)	65% (47-80%)	2.5 (1.1-4.7)	7.8% (4-12%)	9.2% (4.8-14%)	910 (430-1500)	21000 (10000-34000)
Tanzania	2005	Female	70 to 74	34% (20-52%)	66% (48-80%)	2.2 (0.9-4.3)	6.9% (3-11%)	7.9% (3.4-13%)	820 (330-1400)	15000 (6200-27000)
Tanzania	2005	Female	75 to 79	32% (17-50%)	68% (50-83%)	1.8 (0.68-3.8)	6% (2.4-10%)	6.9% (2.8-12%)	520 (190-980)	7300 (2700-14000)
Tanzania	2005	Female	80 to 84	30% (16-49%)	70% (51-84%)	1.7 (0.6-3.3)	5% (1.8-9.4%)	5.8% (2.1-11%)	420 (150-810)	4400 (1600-8300)
Tanzania	2005	Female	85 to 89	30% (16-49%)	70% (51-84%)	1.6 (0.64-3.2)	4.2% (1.4-8.4%)	4.8% (1.6-9.9%)	200 (68-420)	1500 (520-3100)
Tanzania	2005	Female	90 to 94	33% (18-50%)	67% (50-82%)	1.7 (0.65-3.3)	3.7% (1-7.9%)	4.3% (1.2-9.3%)	62 (17-140)	320 (91-690)
Tanzania	2005	Female	95 plus	38% (19-60%)	62% (40-81%)	1.8 (0.61-4.1)	4% (1.1-8.9%)	4.6% (1.2-10%)	17 (4.3-38)	58 (16-130)
Tanzania	2005	Male	All Ages	30% (24-38%)	70% (62-76%)	1.8 (1.3-2.4)	3.2% (2.4-4.1%)	4.9% (3.5-6.4%)	11000 (7800-15000)	470000 (350000-600000)
Tanzania	2005	Male	15 to 19	27% (16-42%)	73% (58-84%)	1.8 (0.86-3.2)	4.9% (2.9-7.7%)	6.2% (3.2-10%)	250 (120-420)	22000 (13000-35000)
Tanzania	2005	Male	20 to 24	38% (23-56%)	62% (44-77%)	2.6 (1.4-4.4)	9.2% (5.7-13%)	11% (5.8-16%)	550 (320-870)	45000 (29000-66000)
Tanzania	2005	Male	25 to 29	44% (26-61%)	56% (39-74%)	3.4 (1.8-5.3)	8.4% (5.1-12%)	9.2% (5-15%)	750 (450-1100)	53000 (35000-75000)
Tanzania	2005	Male	30 to 34	48% (29-66%)	52% (34-71%)	3.9 (2.2-6.3)	6.7% (4.4-9.9%)	7% (4.3-11%)	840 (530-1200)	53000 (35000-73000)
Tanzania	2005	Male	35 to 39	50% (33-69%)	50% (31-67%)	4.3 (2.3-7.3)	6.4% (4.1-9.4%)	6.8% (4-10%)	900 (580-1300)	50000 (33000-69000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	2005	Male	40 to 44	51% (35-70%)	49% (30-65%)	4.5 (2.2-7.5)	6.6% (4.4-9.1%)	7% (4.6-10%)	920 (600-1300)	45000 (30000-62000)
Tanzania	2005	Male	45 to 49	52% (34-70%)	48% (30-66%)	4.5 (2.2-7.8)	7.8% (4.9-11%)	8.7% (5.2-12%)	980 (610-1400)	42000 (27000-60000)
Tanzania	2005	Male	50 to 54	52% (33-71%)	48% (29-67%)	4.4 (2.2-7.8)	9.9% (6.4-14%)	11% (7.3-16%)	1000 (650-1500)	39000 (25000-55000)
Tanzania	2005	Male	55 to 59	51% (33-71%)	49% (29-67%)	4.2 (2-7.5)	11% (7-16%)	13% (8.2-18%)	1100 (680-1600)	37000 (22000-52000)
Tanzania	2005	Male	60 to 64	50% (31-70%)	50% (30-69%)	3.7 (1.7-6.7)	10% (5.7-15%)	12% (6.6-17%)	1000 (560-1500)	29000 (16000-42000)
Tanzania	2005	Male	65 to 69	47% (30-66%)	53% (34-70%)	3 (1.5-5.3)	8.4% (4.3-13%)	9.8% (5-15%)	950 (490-1500)	22000 (11000-34000)
Tanzania	2005	Male	70 to 74	44% (27-62%)	56% (38-73%)	2.3 (1.1-4.4)	6.5% (2.9-11%)	7.5% (3.3-13%)	760 (330-1400)	14000 (6100-25000)
Tanzania	2005	Male	75 to 79	42% (26-60%)	58% (40-74%)	1.9 (0.82-3.8)	5.3% (1.9-10%)	6.1% (2.3-12%)	640 (230-1200)	9100 (3300-18000)
Tanzania	2005	Male	80 to 84	39% (23-58%)	61% (42-77%)	1.6 (0.72-3)	4.4% (1.8-8.5%)	5% (1.9-9.9%)	310 (120-630)	3300 (1300-6500)
Tanzania	2005	Male	85 to 89	36% (22-53%)	64% (47-78%)	1.4 (0.62-2.6)	3.4% (1.3-6.8%)	3.8% (1.5-7.6%)	110 (41-210)	810 (320-1600)
Tanzania	2005	Male	90 to 94	37% (22-55%)	63% (45-78%)	1.2 (0.53-2.4)	2.5% (0.78-5.4%)	2.8% (0.78-6.3%)	27 (7.9-62)	150 (46-320)
Tanzania	2005	Male	95 plus	41% (21-63%)	59% (37-79%)	1.2 (0.43-2.6)	2.8% (0.88-6.4%)	3.1% (0.82-7.2%)	7.3 (1.9-17)	27 (8.4-61)
Tanzania	2010	Female	All Ages	21% (16-27%)	79% (73-84%)	1.3 (0.8-1.9)	1.8% (1.1-2.4%)	3.5% (1.9-5.1%)	7200 (3700-11000)	230000 (140000-320000)
Tanzania	2010	Female	15 to 19	21% (12-35%)	79% (65-88%)	1.4 (0.6-2.7)	1.9% (1-3.2%)	2.4% (1-4.3%)	110 (46-190)	10000 (5300-17000)
Tanzania	2010	Female	20 to 24	26% (14-41%)	74% (59-86%)	2 (0.91-3.6)	2.5% (1.5-3.9%)	2.6% (1.2-4.7%)	140 (64-220)	14000 (8000-21000)
Tanzania	2010	Female	25 to 29	29% (16-45%)	71% (55-84%)	2.4 (1.2-4.3)	2.3% (1.3-3.6%)	2.3% (1.1-4.3%)	180 (98-290)	15000 (9600-22000)
Tanzania	2010	Female	30 to 34	31% (17-49%)	69% (51-83%)	2.7 (1.3-5.1)	2.1% (1.3-3.3%)	2.2% (1.2-3.7%)	230 (130-360)	16000 (10000-24000)
Tanzania	2010	Female	35 to 39	35% (20-53%)	65% (47-80%)	2.7 (1.3-4.8)	2.2% (1.2-3.5%)	2.4% (1.2-4%)	260 (140-410)	16000 (9100-23000)
Tanzania	2010	Female	40 to 44	36% (21-55%)	64% (45-79%)	2.7 (1.2-4.8)	2.6% (1.5-4%)	2.9% (1.6-4.6%)	290 (150-450)	15000 (8500-23000)
Tanzania	2010	Female	45 to 49	37% (20-54%)	63% (46-80%)	2.8 (1.2-5.2)	3.8% (2-5.8%)	4.4% (2.2-7%)	370 (190-600)	17000 (8800-26000)
Tanzania	2010	Female	50 to 54	35% (19-55%)	65% (45-81%)	2.8 (1.2-5.3)	5.7% (3-8.9%)	7% (3.6-11%)	550 (260-910)	21000 (10000-34000)
Tanzania	2010	Female	55 to 59	34% (19-53%)	66% (47-81%)	2.8 (1.2-5.4)	6.9% (3.6-11%)	8.5% (4.4-13%)	670 (320-1100)	22000 (10000-36000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	2010	Female	60 to 64	35% (19-53%)	65% (47-81%)	2.8 (1.1-5.6)	7.3% (3.5-12%)	8.9% (4.5-14%)	820 (370-1400)	23000 (11000-38000)
Tanzania	2010	Female	65 to 69	36% (20-54%)	64% (46-80%)	2.6 (1.1-5.2)	7.9% (3.8-13%)	9.5% (4.5-15%)	1000 (450-1700)	23000 (10000-38000)
Tanzania	2010	Female	70 to 74	34% (20-52%)	66% (48-80%)	2.2 (0.87-4.4)	6.9% (2.7-12%)	8% (3-14%)	910 (330-1600)	17000 (6200-29000)
Tanzania	2010	Female	75 to 79	31% (17-50%)	69% (50-83%)	1.9 (0.67-3.8)	6% (2.1-10%)	6.9% (2.5-12%)	770 (270-1400)	11000 (3900-20000)
Tanzania	2010	Female	80 to 84	30% (16-48%)	70% (52-84%)	1.7 (0.59-3.4)	5% (1.7-9.2%)	5.8% (2-11%)	490 (160-970)	5100 (1600-10000)
Tanzania	2010	Female	85 to 89	30% (17-47%)	70% (53-83%)	1.6 (0.62-3.4)	4.2% (1.3-8.5%)	4.9% (1.5-9.9%)	250 (77-500)	1800 (580-3700)
Tanzania	2010	Female	90 to 94	33% (18-51%)	67% (49-82%)	1.7 (0.6-3.5)	3.7% (0.96-8.3%)	4.3% (1.1-9.7%)	84 (21-190)	430 (110-950)
Tanzania	2010	Female	95 plus	37% (18-60%)	63% (40-82%)	1.8 (0.56-4.1)	4% (1.1-9.4%)	4.6% (1.1-11%)	21 (5.2-51)	74 (19-170)
Tanzania	2010	Male	All Ages	30% (24-38%)	70% (62-76%)	1.8 (1.2-2.6)	3.8% (2.7-4.9%)	5.7% (3.9-7.9%)	13000 (8200-17000)	520000 (370000-680000)
Tanzania	2010	Male	15 to 19	27% (16-42%)	73% (58-84%)	1.7 (0.79-3.2)	4.5% (2.6-7.1%)	5.4% (2.7-8.9%)	280 (130-470)	25000 (13000-39000)
Tanzania	2010	Male	20 to 24	38% (23-55%)	62% (45-77%)	2.6 (1.3-4.4)	9.3% (5.7-13%)	11% (5.9-16%)	600 (330-960)	49000 (30000-73000)
Tanzania	2010	Male	25 to 29	43% (26-62%)	57% (38-74%)	3.3 (1.7-5.5)	9.5% (5.9-14%)	11% (5.9-17%)	810 (460-1200)	58000 (37000-83000)
Tanzania	2010	Male	30 to 34	47% (28-66%)	53% (34-72%)	3.9 (2.1-6.7)	8.3% (5.3-12%)	9.1% (5.4-14%)	920 (560-1300)	59000 (38000-83000)
Tanzania	2010	Male	35 to 39	50% (32-68%)	50% (32-68%)	4.3 (2.1-7.6)	7.8% (5-11%)	8.6% (5.1-13%)	1000 (630-1500)	57000 (37000-79000)
Tanzania	2010	Male	40 to 44	51% (34-70%)	49% (30-66%)	4.4 (2.2-7.9)	8% (5.3-11%)	8.8% (5.8-12%)	1100 (670-1500)	52000 (34000-72000)
Tanzania	2010	Male	45 to 49	52% (32-71%)	48% (29-68%)	4.5 (2-8)	9% (5.6-13%)	10% (6.1-15%)	1100 (670-1600)	48000 (30000-68000)
Tanzania	2010	Male	50 to 54	52% (33-71%)	48% (29-67%)	4.4 (2.1-8.2)	11% (6.8-15%)	13% (8-18%)	1200 (760-1700)	47000 (29000-66000)
Tanzania	2010	Male	55 to 59	52% (33-70%)	48% (30-67%)	4.2 (2-7.7)	12% (7.1-17%)	14% (8.4-20%)	1200 (720-1700)	40000 (24000-55000)
Tanzania	2010	Male	60 to 64	50% (31-70%)	50% (30-69%)	3.7 (1.7-7.4)	10% (5.6-15%)	12% (6.5-18%)	1200 (640-1800)	34000 (18000-49000)
Tanzania	2010	Male	65 to 69	47% (29-66%)	53% (34-71%)	3 (1.4-5.7)	8.5% (4.1-13%)	10% (4.8-16%)	1000 (470-1600)	23000 (11000-36000)
Tanzania	2010	Male	70 to 74	44% (27-61%)	56% (39-73%)	2.3 (1-4.5)	6.5% (2.7-12%)	7.6% (3.1-13%)	860 (350-1500)	16000 (6500-28000)
Tanzania	2010	Male	75 to 79	42% (25-60%)	58% (40-75%)	1.9 (0.77-3.7)	5.4% (1.8-11%)	6.2% (2-12%)	660 (210-1300)	9300 (3100-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	2010	Male	80 to 84	39% (23-59%)	61% (41-77%)	1.6 (0.66-3.1)	4.4% (1.5-8.8%)	5% (1.7-10%)	440 (150-890)	4700 (1600-9400)
Tanzania	2010	Male	85 to 89	36% (22-55%)	64% (45-78%)	1.4 (0.58-2.7)	3.4% (1.2-7.1%)	3.9% (1.3-8.2%)	160 (56-340)	1200 (440-2600)
Tanzania	2010	Male	90 to 94	37% (22-55%)	63% (45-78%)	1.2 (0.47-2.5)	2.5% (0.76-5.7%)	2.8% (0.74-6.5%)	35 (9.3-81)	190 (58-430)
Tanzania	2010	Male	95 plus	41% (20-62%)	59% (38-80%)	1.2 (0.4-2.7)	2.8% (0.82-6.6%)	3.1% (0.76-7.5%)	8.7 (2.2-21)	32 (9.4-74)
Tanzania	2016	Female	All Ages	21% (17-27%)	79% (73-83%)	1.2 (0.64-2.1)	2.2% (1.2-3.2%)	4.3% (2-6.6%)	7700 (3500-12000)	240000 (140000-350000)
Tanzania	2016	Female	15 to 19	21% (12-35%)	79% (65-88%)	1.4 (0.52-3)	2% (0.9-3.3%)	2.5% (0.9-4.8%)	100 (38-190)	11000 (4800-18000)
Tanzania	2016	Female	20 to 24	27% (15-42%)	73% (58-85%)	2 (0.76-4.1)	2.8% (1.5-4.1%)	3% (1.3-5.1%)	130 (60-230)	14000 (8100-21000)
Tanzania	2016	Female	25 to 29	29% (16-44%)	71% (56-84%)	2.4 (0.96-4.9)	2.8% (1.7-4.2%)	3% (1.5-5.1%)	180 (90-290)	16000 (9800-24000)
Tanzania	2016	Female	30 to 34	32% (17-49%)	68% (51-83%)	2.6 (1.1-5.3)	3% (1.8-4.6%)	3.5% (1.8-5.7%)	230 (120-380)	17000 (10000-26000)
Tanzania	2016	Female	35 to 39	35% (20-53%)	65% (47-80%)	2.6 (1-5.2)	3.5% (2-5.2%)	4.3% (2.3-6.6%)	290 (150-470)	18000 (10000-28000)
Tanzania	2016	Female	40 to 44	37% (22-55%)	63% (45-78%)	2.6 (1-5.2)	4% (2.1-6.1%)	5.1% (2.5-7.9%)	330 (160-530)	17000 (8800-27000)
Tanzania	2016	Female	45 to 49	37% (20-54%)	63% (46-80%)	2.7 (0.99-5.7)	5.1% (2.4-8%)	6.7% (3.2-11%)	420 (180-690)	19000 (8700-30000)
Tanzania	2016	Female	50 to 54	35% (20-56%)	65% (44-80%)	2.7 (0.94-6)	6.4% (3-10%)	8.4% (4-13%)	570 (260-960)	22000 (10000-36000)
Tanzania	2016	Female	55 to 59	34% (20-52%)	66% (48-80%)	2.7 (0.89-5.5)	7% (3.2-11%)	9% (4.2-15%)	680 (300-1100)	22000 (9800-37000)
Tanzania	2016	Female	60 to 64	35% (19-53%)	65% (47-81%)	2.7 (0.9-5.9)	7.1% (3.1-12%)	8.8% (3.9-15%)	820 (350-1400)	23000 (9700-39000)
Tanzania	2016	Female	65 to 69	36% (20-53%)	64% (47-80%)	2.5 (0.9-5.6)	7.5% (3.1-12%)	9.1% (3.7-15%)	1100 (410-1800)	24000 (9500-41000)
Tanzania	2016	Female	70 to 74	35% (20-52%)	65% (48-80%)	2.2 (0.71-4.9)	6.4% (1.9-12%)	7.5% (2.2-14%)	990 (300-1800)	18000 (5800-33000)
Tanzania	2016	Female	75 to 79	32% (18-50%)	68% (50-82%)	1.8 (0.58-4.1)	5.6% (1.6-10%)	6.6% (1.9-12%)	870 (250-1600)	12000 (3600-23000)
Tanzania	2016	Female	80 to 84	30% (16-48%)	70% (52-84%)	1.7 (0.55-3.8)	4.7% (1.2-9.3%)	5.4% (1.4-11%)	610 (150-1200)	6400 (1600-13000)
Tanzania	2016	Female	85 to 89	30% (17-48%)	70% (52-83%)	1.6 (0.52-3.7)	3.9% (0.91-8.2%)	4.6% (1-9.5%)	300 (67-640)	2200 (520-4800)
Tanzania	2016	Female	90 to 94	33% (18-50%)	67% (50-82%)	1.6 (0.49-3.9)	3.5% (0.59-8.1%)	4.1% (0.64-9.5%)	95 (16-230)	490 (80-1100)
Tanzania	2016	Female	95 plus	37% (19-59%)	63% (41-81%)	1.7 (0.48-4.3)	3.8% (0.73-8.6%)	4.4% (0.76-10%)	30 (5.2-69)	100 (19-230)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Tanzania	2016	Male	All Ages	30% (24-38%)	70% (62-76%)	1.7 (0.9-3)	4.6% (2.9-6.3%)	6.6% (3.7-9.4%)	14000 (7500-20000)	580000 (360000-800000)
Tanzania	2016	Male	15 to 19	27% (16-43%)	73% (57-84%)	1.7 (0.68-3.4)	4.6% (2.4-7.5%)	5.5% (2.4-9.9%)	310 (130-570)	28000 (15000-48000)
Tanzania	2016	Male	20 to 24	38% (23-56%)	62% (44-77%)	2.5 (1.1-4.9)	9.4% (5.5-14%)	11% (5.4-17%)	680 (330-1100)	56000 (31000-87000)
Tanzania	2016	Male	25 to 29	44% (26-62%)	56% (38-74%)	3.3 (1.4-6.6)	10% (6.2-15%)	12% (6.2-18%)	880 (440-1400)	64000 (37000-96000)
Tanzania	2016	Male	30 to 34	48% (28-67%)	52% (33-72%)	3.8 (1.7-7.4)	9.8% (6.1-14%)	11% (6.3-16%)	970 (540-1500)	63000 (38000-91000)
Tanzania	2016	Male	35 to 39	50% (32-68%)	50% (32-68%)	4.1 (1.7-8.2)	9.8% (6.1-14%)	11% (6.5-16%)	1100 (630-1700)	64000 (38000-91000)
Tanzania	2016	Male	40 to 44	51% (34-70%)	49% (30-66%)	4.3 (1.8-8.2)	10% (6.1-14%)	12% (6.9-16%)	1200 (670-1700)	60000 (35000-84000)
Tanzania	2016	Male	45 to 49	52% (33-71%)	48% (29-67%)	4.3 (1.7-9)	11% (6.1-15%)	13% (7.2-18%)	1300 (670-1800)	55000 (31000-79000)
Tanzania	2016	Male	50 to 54	52% (34-71%)	48% (29-66%)	4.3 (1.7-8.9)	12% (6.6-17%)	14% (7.8-21%)	1300 (710-2000)	51000 (28000-75000)
Tanzania	2016	Male	55 to 59	51% (34-70%)	49% (30-66%)	4.1 (1.6-8.4)	12% (6.2-17%)	15% (7.5-21%)	1400 (690-2000)	45000 (23000-64000)
Tanzania	2016	Male	60 to 64	50% (31-70%)	50% (30-69%)	3.6 (1.4-7.6)	10% (4.7-15%)	12% (5.5-18%)	1200 (530-1800)	33000 (16000-50000)
Tanzania	2016	Male	65 to 69	47% (29-65%)	53% (35-71%)	2.9 (1.1-6.1)	8% (3.2-13%)	9.4% (3.7-16%)	1100 (420-1800)	25000 (9900-41000)
Tanzania	2016	Male	70 to 74	44% (27-62%)	56% (38-73%)	2.2 (0.8-4.7)	6.1% (1.9-11%)	7.1% (2.2-13%)	850 (260-1600)	16000 (4900-30000)
Tanzania	2016	Male	75 to 79	42% (26-60%)	58% (40-74%)	1.8 (0.66-4.2)	5% (1.4-10%)	5.9% (1.6-12%)	690 (180-1500)	9900 (2600-21000)
Tanzania	2016	Male	80 to 84	39% (23-59%)	61% (41-77%)	1.6 (0.56-3.3)	4.2% (1.1-9%)	4.8% (1.2-10%)	480 (120-1100)	5100 (1300-11000)
Tanzania	2016	Male	85 to 89	37% (22-54%)	63% (46-78%)	1.3 (0.48-3)	3.2% (0.84-7.7%)	3.7% (0.88-8.9%)	230 (53-540)	1800 (440-4100)
Tanzania	2016	Male	90 to 94	37% (22-55%)	63% (45-78%)	1.2 (0.39-2.8)	2.4% (0.47-6.2%)	2.7% (0.42-7%)	63 (9.7-170)	340 (64-870)
Tanzania	2016	Male	95 plus	41% (20-63%)	59% (37-80%)	1.2 (0.33-2.9)	2.7% (0.6-6.7%)	3% (0.5-7.6%)	12 (2-32)	45 (9.8-110)
Uganda	1990	Female	All Ages	40% (33-47%)	60% (53-67%)	1.4 (1.1-1.9)	2% (1.4-2.6%)	3.9% (2.5-5.4%)	5100 (3200-7000)	180000 (120000-230000)
Uganda	1990	Female	15 to 19	40% (28-53%)	60% (47-72%)	1.9 (1.1-3)	3.4% (1.6-5.6%)	4.6% (1.7-8.3%)	96 (34-170)	8200 (3900-14000)
Uganda	1990	Female	20 to 24	49% (35-61%)	51% (39-65%)	2.5 (1.4-4)	4% (2-6.7%)	4.6% (1.8-8.4%)	150 (62-270)	12000 (6300-20000)
Uganda	1990	Female	25 to 29	53% (39-65%)	47% (35-61%)	3.1 (1.8-4.9)	4.5% (2.4-7.5%)	5.2% (2.3-9.1%)	200 (94-330)	14000 (7500-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	1990	Female	30 to 34	57% (44-69%)	43% (31-56%)	3.4 (2-5.1)	5.4% (2.9-8.4%)	6.4% (3-11%)	240 (120-370)	15000 (8300-22000)
Uganda	1990	Female	35 to 39	62% (49-73%)	38% (27-51%)	3.2 (1.9-5.1)	6.2% (3.2-9.7%)	7.5% (3.5-12%)	280 (140-440)	15000 (8200-23000)
Uganda	1990	Female	40 to 44	65% (53-76%)	35% (24-47%)	3.3 (1.9-5.2)	6.9% (3.6-11%)	8.5% (4.3-13%)	290 (150-450)	14000 (7500-21000)
Uganda	1990	Female	45 to 49	64% (50-75%)	36% (25-50%)	3.3 (1.8-5.3)	7.9% (4.6-12%)	9.8% (5.6-15%)	340 (190-520)	15000 (8300-22000)
Uganda	1990	Female	50 to 54	63% (48-74%)	37% (26-52%)	3.2 (1.7-5.2)	9.6% (5.5-14%)	12% (6.6-17%)	420 (230-620)	16000 (8600-23000)
Uganda	1990	Female	55 to 59	62% (46-76%)	38% (24-54%)	3.1 (1.6-5.1)	10% (5.6-15%)	12% (6.9-18%)	590 (330-880)	19000 (10000-28000)
Uganda	1990	Female	60 to 64	60% (43-75%)	40% (25-57%)	2.9 (1.6-5.1)	9.9% (5.1-15%)	12% (6.2-18%)	620 (330-960)	17000 (8900-26000)
Uganda	1990	Female	65 to 69	57% (41-72%)	43% (28-59%)	2.6 (1.4-4.2)	9.6% (5-15%)	11% (5.9-17%)	660 (350-1000)	15000 (7900-23000)
Uganda	1990	Female	70 to 74	54% (38-69%)	46% (31-62%)	2.2 (1.1-3.7)	7.8% (3.5-13%)	8.8% (3.9-15%)	520 (230-870)	9400 (4200-16000)
Uganda	1990	Female	75 to 79	52% (35-69%)	48% (31-65%)	1.8 (0.9-3.1)	6.1% (2.3-11%)	6.9% (2.6-13%)	340 (130-630)	4800 (1800-8800)
Uganda	1990	Female	80 to 84	51% (33-69%)	49% (31-67%)	1.7 (0.9-2.8)	5% (1.9-9.3%)	5.6% (2.1-10%)	190 (73-360)	2000 (770-3700)
Uganda	1990	Female	85 to 89	51% (33-67%)	49% (33-67%)	1.7 (0.85-2.8)	3.9% (1.3-7.7%)	4.4% (1.5-8.8%)	70 (23-140)	520 (170-1000)
Uganda	1990	Female	90 to 94	52% (36-68%)	48% (32-64%)	1.7 (0.83-2.9)	3.2% (0.72-6.9%)	3.6% (0.75-8.1%)	17 (3.5-37)	87 (20-190)
Uganda	1990	Female	95 plus	55% (34-75%)	45% (25-66%)	1.7 (0.74-3.2)	3% (0.54-7.1%)	3.4% (0.51-8.3%)	3.5 (0.52-8.4)	12 (2.2-29)
Uganda	1990	Male	All Ages	46% (38-53%)	54% (47-62%)	1.6 (1.2-2.1)	2.3% (1.7-3.2%)	4% (2.8-5.4%)	6200 (4300-8500)	260000 (180000-350000)
Uganda	1990	Male	15 to 19	40% (27-54%)	60% (46-73%)	1.8 (1-3)	4.5% (2.1-7.6%)	5.9% (2.3-10%)	140 (60-260)	11000 (5400-19000)
Uganda	1990	Male	20 to 24	52% (38-64%)	48% (36-62%)	2.5 (1.4-3.8)	6.1% (2.9-10%)	7% (2.9-12%)	310 (130-520)	22000 (11000-37000)
Uganda	1990	Male	25 to 29	62% (50-72%)	38% (28-50%)	3.3 (2-5.1)	6.2% (3-10%)	6.9% (2.9-12%)	430 (200-700)	28000 (15000-44000)
Uganda	1990	Male	30 to 34	69% (56-79%)	31% (21-44%)	4 (2.4-5.9)	7.2% (3.7-11%)	7.9% (3.8-13%)	510 (260-770)	30000 (16000-44000)
Uganda	1990	Male	35 to 39	72% (61-81%)	28% (19-39%)	4.4 (2.6-7)	8.3% (4.3-13%)	9.4% (4.5-15%)	570 (300-850)	30000 (17000-44000)
Uganda	1990	Male	40 to 44	75% (64-83%)	25% (17-36%)	4.6 (2.7-7)	9.2% (5.4-14%)	10% (5.9-16%)	590 (330-870)	28000 (16000-40000)
Uganda	1990	Male	45 to 49	76% (63-85%)	24% (15-37%)	4.4 (2.5-6.9)	10% (6.3-15%)	12% (6.9-18%)	620 (360-910)	26000 (16000-38000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	1990	Male	50 to 54	76% (63-85%)	24% (15-37%)	4.2 (2.3-6.8)	12% (7.1-17%)	14% (8.2-20%)	600 (360-870)	22000 (13000-32000)
Uganda	1990	Male	55 to 59	75% (62-85%)	25% (15-38%)	4 (2.3-6.4)	12% (7.4-17%)	14% (8.6-20%)	710 (420-1100)	23000 (14000-34000)
Uganda	1990	Male	60 to 64	72% (59-84%)	28% (16-41%)	3.4 (1.9-5.8)	11% (6-16%)	12% (7-18%)	630 (360-980)	17000 (9800-26000)
Uganda	1990	Male	65 to 69	69% (54-81%)	31% (19-46%)	2.8 (1.6-4.4)	8% (4.3-13%)	9.3% (5.1-15%)	470 (260-750)	11000 (5900-17000)
Uganda	1990	Male	70 to 74	66% (50-78%)	34% (22-50%)	2.2 (1.2-3.6)	5.8% (2.6-10%)	6.7% (3.1-12%)	320 (150-570)	5800 (2700-10000)
Uganda	1990	Male	75 to 79	62% (46-78%)	38% (22-54%)	1.8 (0.91-3)	4.4% (1.7-8.2%)	5.1% (2-9.4%)	190 (74-360)	2700 (1000-5000)
Uganda	1990	Male	80 to 84	59% (40-75%)	41% (25-60%)	1.5 (0.8-2.5)	3.6% (1.5-6.6%)	4.1% (1.6-7.5%)	91 (36-160)	950 (380-1700)
Uganda	1990	Male	85 to 89	55% (37-71%)	45% (29-63%)	1.2 (0.63-2.1)	2.5% (1.1-4.6%)	2.8% (1.1-5.2%)	31 (13-58)	230 (100-430)
Uganda	1990	Male	90 to 94	54% (38-71%)	46% (29-62%)	1.1 (0.54-2)	1.6% (0.47-3.6%)	1.7% (0.39-4.2%)	6.4 (1.4-15)	35 (10-79)
Uganda	1990	Male	95 plus	56% (35-76%)	44% (24-65%)	1.1 (0.45-2.1)	1.6% (0.34-3.9%)	1.6% (0.18-4.2%)	1.3 (0.14-3.3)	5 (1.1-12)
Uganda	1995	Female	All Ages	40% (34-46%)	60% (54-66%)	1.4 (1-1.9)	2% (1.4-2.8%)	3.8% (2.4-5.5%)	6000 (3700-8600)	210000 (140000-290000)
Uganda	1995	Female	15 to 19	45% (35-55%)	55% (45-65%)	2.1 (1.3-3.2)	4.3% (2.3-6.9%)	6.1% (3-9.8%)	140 (68-230)	12000 (6400-18000)
Uganda	1995	Female	20 to 24	51% (42-59%)	49% (41-58%)	2.7 (1.6-4.1)	4.1% (2.3-6.6%)	4.7% (2.2-8.1%)	200 (100-330)	16000 (8900-25000)
Uganda	1995	Female	25 to 29	54% (46-61%)	46% (39-54%)	3.2 (2-4.6)	3.5% (1.9-5.9%)	3.7% (1.8-7%)	260 (140-400)	18000 (11000-27000)
Uganda	1995	Female	30 to 34	56% (48-63%)	44% (37-52%)	3.3 (2.1-4.9)	3.6% (2-5.7%)	3.9% (2-6.4%)	280 (160-430)	17000 (10000-26000)
Uganda	1995	Female	35 to 39	59% (50-67%)	41% (33-50%)	3 (1.8-4.6)	4.2% (2.4-6.7%)	4.7% (2.5-7.9%)	310 (170-470)	17000 (9700-25000)
Uganda	1995	Female	40 to 44	60% (51-70%)	40% (30-49%)	3 (1.7-4.7)	4.9% (2.8-7.6%)	5.6% (3.1-8.7%)	340 (190-520)	16000 (9300-25000)
Uganda	1995	Female	45 to 49	61% (51-71%)	39% (29-49%)	3.1 (1.7-5)	6.1% (3.5-9.4%)	7.2% (4-11%)	390 (220-600)	17000 (9500-25000)
Uganda	1995	Female	50 to 54	62% (48-74%)	38% (26-52%)	3.1 (1.7-5.2)	8.5% (4.8-12%)	10% (5.7-15%)	540 (290-810)	20000 (11000-30000)
Uganda	1995	Female	55 to 59	61% (45-76%)	39% (24-55%)	3.1 (1.6-5.3)	9.7% (5.3-14%)	11% (6.2-17%)	580 (310-880)	18000 (9900-28000)
Uganda	1995	Female	60 to 64	60% (44-75%)	40% (25-56%)	2.9 (1.6-5.1)	9.7% (5.2-14%)	11% (6.1-17%)	730 (390-1100)	20000 (10000-30000)
Uganda	1995	Female	65 to 69	57% (41-71%)	43% (29-59%)	2.7 (1.4-4.3)	9.6% (5.1-15%)	11% (5.8-17%)	780 (400-1200)	17000 (9000-27000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	1995	Female	70 to 74	54% (38-68%)	46% (32-62%)	2.2 (1.1-3.7)	8% (3.6-13%)	9% (4.1-15%)	640 (270-1100)	12000 (4900-20000)
Uganda	1995	Female	75 to 79	52% (36-69%)	48% (31-64%)	1.8 (0.92-3.1)	6.4% (2.4-12%)	7.2% (2.8-13%)	430 (160-800)	6000 (2200-11000)
Uganda	1995	Female	80 to 84	51% (33-69%)	49% (31-67%)	1.7 (0.88-2.9)	5.2% (2.1-9.7%)	5.9% (2.5-11%)	240 (94-450)	2500 (990-4600)
Uganda	1995	Female	85 to 89	50% (33-66%)	50% (34-67%)	1.7 (0.87-2.9)	4.1% (1.5-8.1%)	4.6% (1.7-9.4%)	86 (30-170)	640 (230-1300)
Uganda	1995	Female	90 to 94	52% (36-68%)	48% (32-64%)	1.7 (0.84-2.9)	3.3% (0.81-7.2%)	3.8% (0.87-8.4%)	22 (5-49)	110 (28-250)
Uganda	1995	Female	95 plus	54% (34-75%)	46% (25-66%)	1.7 (0.72-3.1)	3.2% (0.66-7.5%)	3.6% (0.6-8.6%)	4.8 (0.82-12)	17 (3.5-39)
Uganda	1995	Male	All Ages	50% (43-57%)	50% (43-57%)	1.8 (1.3-2.4)	2.6% (1.8-3.5%)	4.2% (2.9-5.7%)	7900 (5300-11000)	330000 (230000-440000)
Uganda	1995	Male	15 to 19	50% (40-60%)	50% (40-60%)	2.4 (1.5-3.5)	6.4% (3.7-9.7%)	8.6% (4.7-13%)	240 (130-380)	19000 (11000-29000)
Uganda	1995	Male	20 to 24	59% (51-67%)	41% (33-49%)	2.9 (1.8-4.4)	8.4% (4.8-13%)	9.8% (5.2-16%)	470 (250-760)	33000 (19000-52000)
Uganda	1995	Male	25 to 29	67% (60-73%)	33% (27-40%)	3.7 (2.3-5.4)	6.5% (3.4-11%)	7.1% (3.4-12%)	620 (350-930)	40000 (24000-58000)
Uganda	1995	Male	30 to 34	74% (66-80%)	26% (20-34%)	4.4 (2.8-6.4)	5.8% (3.3-9.1%)	6.1% (3.3-9.9%)	680 (410-980)	39000 (25000-56000)
Uganda	1995	Male	35 to 39	76% (69-82%)	24% (18-31%)	4.7 (2.8-7.3)	6.5% (3.7-10%)	7.1% (3.9-11%)	690 (430-970)	36000 (23000-50000)
Uganda	1995	Male	40 to 44	78% (71-84%)	22% (16-29%)	4.9 (2.8-7.6)	7.2% (4.4-10%)	7.9% (4.7-11%)	730 (460-1000)	34000 (22000-47000)
Uganda	1995	Male	45 to 49	79% (69-86%)	21% (14-31%)	4.7 (2.7-7.4)	8.4% (4.9-12%)	9.3% (5.4-14%)	760 (450-1100)	31000 (19000-45000)
Uganda	1995	Male	50 to 54	78% (68-85%)	22% (15-32%)	4.4 (2.5-7)	11% (6.4-15%)	12% (7.2-17%)	830 (480-1200)	30000 (18000-43000)
Uganda	1995	Male	55 to 59	76% (63-86%)	24% (14-37%)	4.1 (2.3-6.6)	12% (7.2-17%)	14% (8.4-19%)	750 (450-1100)	24000 (15000-35000)
Uganda	1995	Male	60 to 64	72% (58-84%)	28% (16-42%)	3.4 (1.9-5.7)	10% (6.1-15%)	12% (6.8-17%)	760 (430-1200)	21000 (12000-32000)
Uganda	1995	Male	65 to 69	69% (53-82%)	31% (18-47%)	2.8 (1.6-4.4)	7.9% (4.3-12%)	9.1% (4.9-14%)	570 (310-930)	13000 (7100-21000)
Uganda	1995	Male	70 to 74	66% (50-78%)	34% (22-50%)	2.2 (1.2-3.5)	5.9% (2.7-10%)	6.7% (3.1-11%)	400 (190-690)	7300 (3400-12000)
Uganda	1995	Male	75 to 79	62% (46-77%)	38% (23-54%)	1.8 (0.92-3)	4.6% (1.7-8.3%)	5.2% (1.9-9.5%)	250 (89-450)	3500 (1300-6300)
Uganda	1995	Male	80 to 84	59% (40-75%)	41% (25-60%)	1.5 (0.81-2.5)	3.8% (1.5-6.7%)	4.3% (1.7-7.7%)	120 (46-210)	1200 (490-2200)
Uganda	1995	Male	85 to 89	55% (37-71%)	45% (29-63%)	1.3 (0.66-2.2)	2.6% (1.1-4.8%)	3% (1.2-5.4%)	39 (16-71)	300 (130-550)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	1995	Male	90 to 94	54% (38-71%)	46% (29-62%)	1.1 (0.57-2)	1.7% (0.56-3.6%)	1.9% (0.54-4%)	8.3 (2.4-18)	45 (15-95)
Uganda	1995	Male	95 plus	56% (35-76%)	44% (24-65%)	1.1 (0.44-2.1)	1.7% (0.44-3.8%)	1.8% (0.3-4.2%)	1.7 (0.29-4.1)	6.7 (1.7-15)
Uganda	2000	Female	All Ages	39% (33-45%)	61% (55-67%)	1.5 (1.1-1.9)	2.4% (1.7-3.3%)	4.6% (3-6.5%)	7600 (4900-11000)	260000 (180000-360000)
Uganda	2000	Female	15 to 19	45% (36-54%)	55% (46-64%)	2.3 (1.4-3.4)	4.8% (2.6-7.5%)	6.9% (3.6-12%)	180 (95-300)	15000 (8600-24000)
Uganda	2000	Female	20 to 24	50% (41-57%)	50% (43-59%)	2.8 (1.7-4.3)	5.5% (3.1-8.5%)	6.9% (3.5-11%)	250 (130-420)	19000 (11000-30000)
Uganda	2000	Female	25 to 29	52% (46-59%)	48% (41-54%)	3.3 (2.1-4.8)	4.2% (2.3-7%)	4.7% (2.3-8.3%)	320 (180-490)	22000 (13000-32000)
Uganda	2000	Female	30 to 34	55% (48-62%)	45% (38-52%)	3.4 (2.2-5.1)	3.7% (2.1-5.8%)	3.9% (2.1-6.5%)	340 (190-530)	21000 (13000-31000)
Uganda	2000	Female	35 to 39	58% (50-66%)	42% (34-50%)	3.1 (1.9-4.7)	4% (2.4-6.6%)	4.4% (2.4-7.4%)	340 (200-530)	18000 (11000-28000)
Uganda	2000	Female	40 to 44	59% (51-68%)	41% (32-49%)	3.1 (1.8-4.8)	4.7% (2.8-7.2%)	5.3% (3.1-8.1%)	380 (220-570)	18000 (11000-27000)
Uganda	2000	Female	45 to 49	60% (51-69%)	40% (31-49%)	3.2 (1.8-5.2)	6.5% (3.8-9.7%)	7.5% (4.3-11%)	500 (290-770)	21000 (13000-32000)
Uganda	2000	Female	50 to 54	61% (50-71%)	39% (29-50%)	3.3 (1.9-5.2)	9.5% (5.9-14%)	11% (6.9-16%)	710 (410-1000)	26000 (15000-37000)
Uganda	2000	Female	55 to 59	60% (46-74%)	40% (26-54%)	3.1 (1.7-5.3)	11% (6.7-17%)	14% (8-20%)	830 (470-1200)	26000 (15000-39000)
Uganda	2000	Female	60 to 64	58% (41-73%)	42% (27-59%)	2.9 (1.5-5)	11% (6.2-16%)	13% (7.3-19%)	760 (410-1200)	20000 (11000-31000)
Uganda	2000	Female	65 to 69	55% (40-69%)	45% (31-60%)	2.7 (1.5-4.4)	11% (6.2-16%)	13% (7.2-19%)	980 (550-1500)	22000 (12000-34000)
Uganda	2000	Female	70 to 74	53% (37-67%)	47% (33-63%)	2.3 (1.2-3.8)	9.3% (4.6-15%)	11% (5.4-17%)	840 (400-1400)	15000 (7400-25000)
Uganda	2000	Female	75 to 79	51% (35-68%)	49% (32-65%)	1.9 (0.96-3.3)	7.7% (3.4-14%)	8.8% (3.9-15%)	620 (260-1100)	8600 (3800-15000)
Uganda	2000	Female	80 to 84	51% (33-69%)	49% (31-67%)	1.8 (0.97-3)	6.4% (2.8-11%)	7.4% (3.2-13%)	370 (160-660)	3800 (1600-6800)
Uganda	2000	Female	85 to 89	51% (34-66%)	49% (34-66%)	1.8 (0.93-3.1)	5.3% (2.1-10%)	6.1% (2.5-11%)	130 (55-250)	980 (410-1800)
Uganda	2000	Female	90 to 94	52% (36-68%)	48% (32-64%)	1.8 (0.88-3.1)	4.4% (1.5-8.9%)	5.1% (1.8-10%)	34 (12-69)	170 (61-350)
Uganda	2000	Female	95 plus	55% (35-73%)	45% (27-65%)	1.8 (0.76-3.3)	4.3% (1.3-9.1%)	4.9% (1.3-11%)	7.9 (2.2-17)	27 (7.9-57)
Uganda	2000	Male	All Ages	47% (39-54%)	53% (46-61%)	1.8 (1.3-2.3)	3.3% (2.5-4.2%)	5.3% (3.8-6.8%)	10000 (7400-13000)	420000 (320000-540000)
Uganda	2000	Male	15 to 19	45% (35-56%)	55% (44-65%)	2.2 (1.4-3.4)	6.8% (4-10%)	9% (5-14%)	320 (180-490)	25000 (14000-37000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	2000	Male	20 to 24	53% (44-63%)	47% (37-56%)	2.7 (1.6-4.2)	11% (6.4-16%)	13% (7.5-19%)	610 (350-920)	43000 (26000-63000)
Uganda	2000	Male	25 to 29	63% (55-71%)	37% (29-45%)	3.6 (2.2-5.3)	10% (5.8-16%)	11% (6.1-18%)	800 (500-1200)	51000 (34000-73000)
Uganda	2000	Male	30 to 34	72% (64-78%)	28% (22-36%)	4.5 (2.9-6.5)	7.8% (5-11%)	8.5% (5.2-13%)	900 (600-1200)	52000 (36000-69000)
Uganda	2000	Male	35 to 39	75% (67-81%)	25% (19-33%)	4.9 (3-7.4)	7.3% (4.6-11%)	7.9% (4.8-12%)	880 (590-1200)	46000 (31000-61000)
Uganda	2000	Male	40 to 44	76% (69-83%)	24% (17-31%)	5 (2.9-7.5)	8% (5.4-11%)	8.7% (5.8-12%)	860 (570-1200)	40000 (27000-54000)
Uganda	2000	Male	45 to 49	78% (68-85%)	22% (15-32%)	4.9 (2.8-7.8)	9.4% (5.7-13%)	10% (6.3-15%)	910 (590-1200)	38000 (25000-51000)
Uganda	2000	Male	50 to 54	78% (68-86%)	22% (14-32%)	4.7 (2.8-7.6)	12% (7.8-17%)	14% (8.7-19%)	1000 (650-1400)	37000 (24000-51000)
Uganda	2000	Male	55 to 59	76% (63-85%)	24% (15-37%)	4.3 (2.5-6.8)	14% (9.3-19%)	16% (11-22%)	1100 (690-1500)	34000 (22000-46000)
Uganda	2000	Male	60 to 64	71% (56-83%)	29% (17-44%)	3.5 (1.9-5.9)	12% (7-17%)	14% (8.1-20%)	820 (480-1200)	22000 (13000-32000)
Uganda	2000	Male	65 to 69	68% (52-80%)	32% (20-48%)	2.8 (1.6-4.5)	9.5% (5.2-15%)	11% (6-17%)	750 (420-1200)	17000 (9400-27000)
Uganda	2000	Male	70 to 74	65% (49-77%)	35% (23-51%)	2.2 (1.3-3.6)	7.2% (3.5-12%)	8.2% (4-14%)	550 (270-930)	9900 (4900-17000)
Uganda	2000	Male	75 to 79	62% (45-77%)	38% (23-55%)	1.9 (0.98-3.2)	5.8% (2.3-10%)	6.6% (2.7-12%)	360 (150-660)	5000 (2100-9200)
Uganda	2000	Male	80 to 84	59% (41-75%)	41% (25-59%)	1.6 (0.9-2.7)	4.7% (2-8.5%)	5.4% (2.3-9.8%)	180 (76-330)	1900 (820-3400)
Uganda	2000	Male	85 to 89	55% (38-71%)	45% (29-62%)	1.3 (0.69-2.3)	3.4% (1.5-6.1%)	3.9% (1.6-7%)	63 (27-110)	470 (200-840)
Uganda	2000	Male	90 to 94	55% (39-70%)	45% (30-61%)	1.2 (0.6-2.1)	2.4% (0.79-4.9%)	2.7% (0.85-5.6%)	14 (4.5-30)	75 (25-150)
Uganda	2000	Male	95 plus	57% (36-76%)	43% (24-64%)	1.2 (0.49-2.2)	2.6% (0.77-5.5%)	2.8% (0.75-6%)	3.3 (0.87-7.1)	12 (3.5-25)
Uganda	2005	Female	All Ages	38% (33-45%)	62% (55-67%)	1.4 (1-1.9)	2.6% (1.8-3.5%)	5% (3.3-7%)	8200 (5200-12000)	280000 (190000-370000)
Uganda	2005	Female	15 to 19	44% (35-53%)	56% (47-65%)	2.2 (1.3-3.3)	3.9% (2.2-6.4%)	5.1% (2.5-9%)	190 (98-330)	16000 (9000-26000)
Uganda	2005	Female	20 to 24	49% (40-56%)	51% (44-60%)	2.7 (1.6-4.1)	5.3% (3-8.5%)	6.8% (3.3-12%)	260 (130-440)	21000 (12000-33000)
Uganda	2005	Female	25 to 29	52% (46-58%)	48% (42-54%)	3.2 (2-4.8)	5.4% (3-8.5%)	6.5% (3.1-11%)	340 (200-510)	24000 (15000-34000)
Uganda	2005	Female	30 to 34	55% (47-62%)	45% (38-53%)	3.3 (2.1-5)	4.5% (2.7-7.1%)	5.2% (2.8-8.6%)	370 (210-570)	23000 (14000-33000)
Uganda	2005	Female	35 to 39	57% (50-66%)	43% (34-50%)	3 (1.9-4.6)	4.4% (2.6-6.9%)	4.9% (2.8-7.9%)	360 (210-550)	20000 (12000-29000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	2005	Female	40 to 44	59% (51-67%)	41% (33-49%)	3 (1.8-4.8)	4.9% (3-7.4%)	5.6% (3.4-8.6%)	360 (220-550)	18000 (11000-26000)
Uganda	2005	Female	45 to 49	60% (50-69%)	40% (31-50%)	3.1 (1.7-5)	6.9% (4.1-10%)	8.2% (4.7-12%)	480 (280-730)	20000 (12000-30000)
Uganda	2005	Female	50 to 54	61% (49-72%)	39% (28-51%)	3.1 (1.8-5)	10% (6-15%)	13% (7.2-18%)	770 (440-1100)	28000 (16000-42000)
Uganda	2005	Female	55 to 59	60% (44-74%)	40% (26-56%)	3 (1.7-5.2)	12% (7.1-18%)	15% (8.6-21%)	900 (520-1400)	28000 (16000-43000)
Uganda	2005	Female	60 to 64	57% (42-72%)	43% (28-58%)	2.8 (1.5-5)	11% (6.5-17%)	14% (7.8-20%)	910 (490-1400)	24000 (13000-38000)
Uganda	2005	Female	65 to 69	55% (39-69%)	45% (31-61%)	2.6 (1.4-4.3)	11% (6.4-17%)	13% (7.6-20%)	900 (510-1400)	20000 (11000-31000)
Uganda	2005	Female	70 to 74	52% (37-67%)	48% (33-63%)	2.2 (1.1-3.7)	9.8% (4.6-16%)	11% (5.4-18%)	950 (430-1500)	17000 (7800-28000)
Uganda	2005	Female	75 to 79	51% (35-68%)	49% (32-65%)	1.9 (0.93-3.3)	8.2% (3.4-14%)	9.4% (3.9-16%)	730 (300-1200)	10000 (4100-17000)
Uganda	2005	Female	80 to 84	51% (33-68%)	49% (32-67%)	1.8 (0.91-3)	6.8% (2.9-12%)	7.9% (3.3-14%)	470 (200-810)	4800 (2000-8300)
Uganda	2005	Female	85 to 89	50% (33-67%)	50% (33-67%)	1.7 (0.87-3)	5.7% (2.3-10%)	6.6% (2.6-12%)	180 (71-330)	1300 (520-2400)
Uganda	2005	Female	90 to 94	52% (36-69%)	48% (31-64%)	1.7 (0.84-3)	4.9% (1.8-9.3%)	5.7% (2-11%)	46 (16-89)	230 (82-440)
Uganda	2005	Female	95 plus	55% (35-74%)	45% (26-65%)	1.8 (0.78-3.2)	4.8% (1.4-9.7%)	5.5% (1.5-11%)	10 (2.7-21)	34 (10-70)
Uganda	2005	Male	All Ages	47% (40-54%)	53% (46-60%)	1.7 (1.2-2.3)	3.8% (2.8-4.9%)	6.1% (4.4-8%)	11000 (8000-15000)	480000 (350000-610000)
Uganda	2005	Male	15 to 19	45% (35-56%)	55% (44-65%)	2.2 (1.3-3.3)	6% (3.5-9.5%)	7.6% (4.1-12%)	360 (200-570)	28000 (16000-44000)
Uganda	2005	Male	20 to 24	53% (44-63%)	47% (37-56%)	2.7 (1.5-4.1)	11% (6.3-16%)	13% (7.3-20%)	690 (390-1100)	49000 (29000-73000)
Uganda	2005	Male	25 to 29	63% (55-70%)	37% (30-45%)	3.4 (2.1-5.1)	13% (7.6-19%)	15% (8.3-23%)	920 (560-1300)	59000 (38000-82000)
Uganda	2005	Male	30 to 34	72% (64-78%)	28% (22-36%)	4.3 (2.8-6.4)	12% (7.3-16%)	13% (7.9-20%)	1100 (670-1400)	61000 (40000-82000)
Uganda	2005	Male	35 to 39	75% (68-81%)	25% (19-32%)	4.8 (2.9-7.3)	10% (6.5-14%)	11% (7-16%)	1000 (690-1400)	55000 (37000-74000)
Uganda	2005	Male	40 to 44	76% (69-83%)	24% (17-31%)	4.9 (2.9-7.6)	9.7% (6.6-13%)	11% (7.3-15%)	980 (650-1400)	46000 (31000-63000)
Uganda	2005	Male	45 to 49	78% (68-85%)	22% (15-32%)	4.8 (2.6-7.4)	11% (7.1-16%)	13% (7.9-18%)	950 (600-1300)	39000 (25000-53000)
Uganda	2005	Male	50 to 54	78% (69-86%)	22% (14-31%)	4.6 (2.5-7.1)	14% (8.4-19%)	16% (9.7-23%)	1100 (660-1500)	39000 (24000-53000)
Uganda	2005	Male	55 to 59	76% (65-85%)	24% (15-35%)	4.2 (2.4-6.9)	15% (9.6-21%)	18% (11-25%)	1100 (710-1600)	36000 (23000-49000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	2005	Male	60 to 64	71% (57-83%)	29% (17-43%)	3.4 (1.9-5.8)	13% (7.8-19%)	15% (9.1-22%)	1000 (590-1500)	27000 (16000-40000)
Uganda	2005	Male	65 to 69	68% (52-80%)	32% (20-48%)	2.7 (1.5-4.5)	10% (5.6-16%)	12% (6.5-18%)	740 (390-1200)	17000 (8900-26000)
Uganda	2005	Male	70 to 74	64% (49-77%)	36% (23-51%)	2.2 (1.2-3.6)	7.7% (3.7-13%)	8.9% (4.4-15%)	650 (310-1100)	12000 (5700-20000)
Uganda	2005	Male	75 to 79	62% (45-76%)	38% (24-55%)	1.8 (0.95-3.1)	6.2% (2.4-11%)	7.1% (2.9-13%)	430 (180-760)	6100 (2500-11000)
Uganda	2005	Male	80 to 84	59% (40-75%)	41% (25-60%)	1.6 (0.83-2.6)	5% (2.1-8.9%)	5.8% (2.4-10%)	230 (94-410)	2400 (990-4300)
Uganda	2005	Male	85 to 89	55% (38-70%)	45% (30-62%)	1.3 (0.64-2.2)	3.7% (1.6-6.5%)	4.2% (1.8-7.5%)	84 (35-150)	630 (270-1100)
Uganda	2005	Male	90 to 94	54% (38-70%)	46% (30-62%)	1.1 (0.56-2)	2.6% (0.88-5.1%)	3% (0.95-5.9%)	19 (6.3-39)	100 (35-200)
Uganda	2005	Male	95 plus	57% (35-76%)	43% (24-65%)	1.1 (0.48-2.1)	2.9% (0.84-6.3%)	3.3% (0.84-7.1%)	4.7 (1.2-10)	17 (4.9-36)
Uganda	2010	Female	All Ages	39% (32-46%)	61% (54-68%)	1.4 (0.92-2)	2.5% (1.6-3.5%)	4.9% (2.8-7.1%)	7700 (4300-11000)	260000 (160000-370000)
Uganda	2010	Female	15 to 19	40% (28-52%)	60% (48-72%)	1.8 (0.94-3.2)	3.2% (1.7-5.3%)	4% (1.9-7.1%)	170 (81-300)	15000 (7900-25000)
Uganda	2010	Female	20 to 24	48% (35-61%)	52% (39-65%)	2.5 (1.3-4.3)	4.5% (2.5-7%)	5.4% (2.4-9.4%)	240 (120-420)	20000 (11000-32000)
Uganda	2010	Female	25 to 29	52% (39-64%)	48% (36-61%)	3 (1.7-4.9)	4.4% (2.4-7%)	5% (2.3-8.9%)	310 (170-480)	22000 (14000-33000)
Uganda	2010	Female	30 to 34	57% (43-68%)	43% (32-57%)	3.2 (1.8-5.3)	4.5% (2.5-7.2%)	5.2% (2.6-9.1%)	350 (200-560)	22000 (14000-35000)
Uganda	2010	Female	35 to 39	62% (48-73%)	38% (27-52%)	3.1 (1.7-5.3)	4.7% (2.6-7.5%)	5.5% (2.9-9.1%)	380 (210-590)	21000 (12000-32000)
Uganda	2010	Female	40 to 44	64% (52-75%)	36% (25-48%)	3.2 (1.7-5.3)	5.2% (3-8%)	6.1% (3.4-9.4%)	370 (210-580)	18000 (10000-28000)
Uganda	2010	Female	45 to 49	64% (49-75%)	36% (25-51%)	3.2 (1.7-5.6)	6.9% (3.7-10%)	8.4% (4.6-13%)	440 (240-680)	19000 (10000-28000)
Uganda	2010	Female	50 to 54	62% (48-75%)	38% (25-52%)	3.1 (1.5-5.4)	9.7% (5.3-14%)	12% (6.5-18%)	620 (330-960)	23000 (12000-36000)
Uganda	2010	Female	55 to 59	62% (46-77%)	38% (23-54%)	3 (1.5-5.5)	11% (6.1-17%)	14% (7.7-21%)	830 (430-1300)	26000 (14000-41000)
Uganda	2010	Female	60 to 64	60% (44-75%)	40% (25-56%)	2.9 (1.4-5.1)	11% (5.7-17%)	13% (7.1-20%)	860 (430-1400)	23000 (12000-37000)
Uganda	2010	Female	65 to 69	57% (40-71%)	43% (29-60%)	2.6 (1.3-4.5)	11% (5.4-17%)	13% (6.6-20%)	940 (470-1500)	21000 (10000-33000)
Uganda	2010	Female	70 to 74	53% (38-69%)	47% (31-62%)	2.1 (1.1-3.7)	8.9% (3.8-15%)	10% (4.5-17%)	750 (320-1200)	14000 (5900-22000)
Uganda	2010	Female	75 to 79	52% (35-68%)	48% (32-65%)	1.8 (0.82-3.2)	7.4% (2.6-13%)	8.7% (3.1-15%)	700 (250-1200)	9800 (3600-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	2010	Female	80 to 84	51% (32-68%)	49% (32-68%)	1.7 (0.8-2.9)	6.1% (2.3-11%)	7.1% (2.6-13%)	480 (180-870)	5000 (1900-9000)
Uganda	2010	Female	85 to 89	50% (33-66%)	50% (34-67%)	1.6 (0.77-2.9)	5.1% (1.8-10%)	6% (2.1-12%)	190 (67-380)	1400 (500-2800)
Uganda	2010	Female	90 to 94	52% (35-68%)	48% (32-65%)	1.6 (0.74-2.9)	4.3% (1.2-9%)	5.1% (1.4-11%)	51 (14-100)	260 (73-530)
Uganda	2010	Female	95 plus	54% (34-73%)	46% (27-66%)	1.6 (0.68-3)	4.4% (1.1-9.2%)	5.1% (1.2-11%)	11 (2.5-23)	37 (9.3-78)
Uganda	2010	Male	All Ages	44% (37-52%)	56% (48-63%)	1.5 (1-2.2)	4.1% (2.8-5.5%)	6.5% (4.4-9%)	12000 (7700-16000)	490000 (340000-660000)
Uganda	2010	Male	15 to 19	39% (27-52%)	61% (48-73%)	1.7 (0.91-3)	5.1% (2.7-8.3%)	6.3% (3.1-11%)	340 (170-580)	27000 (14000-45000)
Uganda	2010	Male	20 to 24	51% (38-63%)	49% (37-62%)	2.4 (1.3-4)	9.9% (5.6-15%)	12% (6.4-19%)	700 (380-1100)	50000 (29000-77000)
Uganda	2010	Male	25 to 29	62% (49-72%)	38% (28-51%)	3.1 (1.8-5.1)	12% (6.4-17%)	14% (7.3-21%)	950 (550-1400)	62000 (37000-90000)
Uganda	2010	Male	30 to 34	69% (56-79%)	31% (21-44%)	3.8 (2.3-6.1)	12% (7.4-18%)	14% (8.2-22%)	1100 (710-1600)	65000 (41000-90000)
Uganda	2010	Male	35 to 39	72% (61-81%)	28% (19-39%)	4.2 (2.5-7)	12% (7.6-17%)	14% (8.4-20%)	1100 (740-1600)	60000 (40000-82000)
Uganda	2010	Male	40 to 44	74% (63-83%)	26% (17-37%)	4.4 (2.5-7.2)	12% (7.6-16%)	13% (8.8-19%)	1100 (700-1600)	52000 (34000-72000)
Uganda	2010	Male	45 to 49	76% (63-85%)	24% (15-37%)	4.2 (2.1-7)	12% (7.6-17%)	14% (8.7-20%)	1000 (610-1400)	42000 (26000-58000)
Uganda	2010	Male	50 to 54	76% (64-85%)	24% (15-36%)	4.1 (2.1-7)	14% (8.7-20%)	17% (10-24%)	1000 (600-1400)	37000 (22000-50000)
Uganda	2010	Male	55 to 59	75% (61-84%)	25% (16-39%)	3.9 (2.1-6.7)	15% (9.1-21%)	18% (11-26%)	1100 (640-1500)	34000 (21000-47000)
Uganda	2010	Male	60 to 64	72% (58-84%)	28% (16-42%)	3.3 (1.7-5.9)	13% (7-19%)	15% (8.4-23%)	970 (520-1500)	26000 (14000-39000)
Uganda	2010	Male	65 to 69	68% (53-81%)	32% (19-47%)	2.6 (1.4-4.7)	9.9% (5.1-16%)	12% (6.2-19%)	820 (420-1400)	18000 (9500-30000)
Uganda	2010	Male	70 to 74	65% (49-77%)	35% (23-51%)	2.1 (1.1-3.6)	7.2% (3.1-13%)	8.4% (3.7-15%)	560 (250-980)	10000 (4400-18000)
Uganda	2010	Male	75 to 79	61% (44-77%)	39% (23-56%)	1.7 (0.81-3.1)	5.7% (2.1-11%)	6.6% (2.4-13%)	440 (160-840)	6200 (2300-12000)
Uganda	2010	Male	80 to 84	57% (38-74%)	43% (26-62%)	1.4 (0.7-2.5)	4.5% (1.7-8.5%)	5.2% (1.9-9.7%)	250 (89-480)	2600 (950-5000)
Uganda	2010	Male	85 to 89	54% (37-70%)	46% (30-63%)	1.2 (0.56-2.1)	3.4% (1.2-6.5%)	3.9% (1.3-7.5%)	92 (30-180)	690 (230-1300)
Uganda	2010	Male	90 to 94	53% (37-70%)	47% (30-63%)	1 (0.49-1.9)	2.4% (0.62-5%)	2.7% (0.63-5.7%)	22 (5-46)	110 (30-240)
Uganda	2010	Male	95 plus	55% (35-75%)	45% (25-65%)	1 (0.42-1.9)	2.7% (0.7-5.7%)	3% (0.64-6.6%)	5.3 (1.2-12)	19 (5-41)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	2016	Female	All Ages	39% (32-47%)	61% (53-68%)	1.2 (0.64-2.1)	2.9% (1.5-4.3%)	5.3% (2.4-8.5%)	7200 (3200-12000)	250000 (130000-380000)
Uganda	2016	Female	15 to 19	40% (27-52%)	60% (48-73%)	1.6 (0.71-3.1)	3.3% (1.5-5.7%)	4.6% (1.6-8.8%)	160 (58-290)	14000 (6600-25000)
Uganda	2016	Female	20 to 24	48% (35-61%)	52% (39-65%)	2.2 (0.97-4.2)	4.5% (2.2-7.5%)	5.5% (2.1-10%)	220 (88-410)	20000 (10000-34000)
Uganda	2016	Female	25 to 29	52% (39-64%)	48% (36-61%)	2.7 (1.2-4.9)	4.7% (2.5-7.7%)	5.8% (2.4-11%)	280 (130-490)	22000 (12000-34000)
Uganda	2016	Female	30 to 34	57% (43-68%)	43% (32-57%)	2.9 (1.4-5.2)	5.4% (2.8-8.9%)	7% (3.1-13%)	340 (160-590)	23000 (12000-37000)
Uganda	2016	Female	35 to 39	62% (48-73%)	38% (27-52%)	2.8 (1.3-5.1)	6.4% (3.3-10%)	8.7% (4.3-14%)	400 (200-670)	23000 (12000-37000)
Uganda	2016	Female	40 to 44	64% (52-75%)	36% (25-48%)	2.8 (1.3-5.5)	7% (3.6-11%)	9.7% (4.9-16%)	420 (200-700)	21000 (10000-34000)
Uganda	2016	Female	45 to 49	64% (49-75%)	36% (25-51%)	2.9 (1.2-5.7)	8.4% (4-14%)	12% (5.4-18%)	470 (210-800)	20000 (9200-34000)
Uganda	2016	Female	50 to 54	62% (48-75%)	38% (25-52%)	2.7 (1.1-5.4)	9.7% (4.5-15%)	13% (6.1-20%)	560 (250-920)	21000 (9200-34000)
Uganda	2016	Female	55 to 59	61% (46-76%)	39% (24-54%)	2.6 (1.1-5.2)	11% (4.8-17%)	14% (6.4-22%)	670 (290-1100)	21000 (9200-36000)
Uganda	2016	Female	60 to 64	60% (43-74%)	40% (26-57%)	2.5 (1-5)	9.6% (3.9-16%)	12% (5.1-20%)	780 (310-1300)	21000 (8200-35000)
Uganda	2016	Female	65 to 69	57% (40-71%)	43% (29-60%)	2.3 (0.97-4.5)	9.2% (3.6-16%)	11% (4.5-19%)	870 (330-1500)	20000 (7400-34000)
Uganda	2016	Female	70 to 74	53% (38-68%)	47% (32-62%)	1.9 (0.77-3.8)	7.5% (2.2-14%)	8.8% (2.7-17%)	750 (220-1400)	14000 (4100-26000)
Uganda	2016	Female	75 to 79	52% (35-69%)	48% (31-65%)	1.6 (0.63-3.2)	6.2% (1.5-13%)	7.3% (1.8-15%)	560 (140-1200)	7800 (2000-16000)
Uganda	2016	Female	80 to 84	51% (33-68%)	49% (32-67%)	1.5 (0.6-2.8)	5.1% (1.1-11%)	5.9% (1.2-13%)	400 (81-830)	4100 (830-8600)
Uganda	2016	Female	85 to 89	50% (34-67%)	50% (33-66%)	1.5 (0.57-2.8)	4.3% (0.78-9.4%)	5.1% (0.92-11%)	250 (46-540)	1800 (330-4000)
Uganda	2016	Female	90 to 94	51% (35-68%)	49% (32-65%)	1.4 (0.56-3)	3.6% (0.46-8.6%)	4.2% (0.48-10%)	71 (8-170)	360 (45-870)
Uganda	2016	Female	95 plus	54% (34-74%)	46% (26-66%)	1.5 (0.51-3.1)	3.8% (0.48-9.1%)	4.4% (0.46-11%)	16 (1.7-39)	55 (6.8-130)
Uganda	2016	Male	All Ages	45% (37-53%)	55% (47-63%)	1.4 (0.71-2.4)	4.6% (2.6-6.8%)	7% (3.8-11%)	12000 (6200-18000)	500000 (280000-740000)
Uganda	2016	Male	15 to 19	39% (27-52%)	61% (48-73%)	1.5 (0.69-2.9)	4.9% (2.3-8.5%)	6.3% (2.6-11%)	330 (140-610)	27000 (12000-47000)
Uganda	2016	Male	20 to 24	51% (37-63%)	49% (37-63%)	2.1 (0.92-4)	9.1% (4.3-15%)	11% (4.8-18%)	700 (300-1200)	51000 (24000-84000)
Uganda	2016	Male	25 to 29	62% (49-72%)	38% (28-51%)	2.8 (1.2-5)	11% (5.3-17%)	13% (5.8-21%)	950 (440-1500)	63000 (31000-96000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Uganda	2016	Male	30 to 34	69% (56-78%)	31% (22-44%)	3.4 (1.6-6.1)	11% (5.9-17%)	13% (6.6-21%)	1100 (590-1600)	65000 (36000-97000)
Uganda	2016	Male	35 to 39	72% (60-81%)	28% (19-40%)	3.7 (1.8-7)	12% (7.1-18%)	15% (8.1-22%)	1200 (680-1700)	63000 (36000-90000)
Uganda	2016	Male	40 to 44	74% (63-83%)	26% (17-37%)	3.9 (1.9-7.2)	14% (7.6-20%)	17% (9.3-24%)	1200 (670-1800)	57000 (32000-83000)
Uganda	2016	Male	45 to 49	76% (63-85%)	24% (15-37%)	3.8 (1.7-7.1)	14% (7.5-20%)	17% (9.2-25%)	1100 (590-1700)	48000 (25000-70000)
Uganda	2016	Male	50 to 54	76% (63-85%)	24% (15-37%)	3.6 (1.6-6.9)	15% (7.6-21%)	18% (9.5-26%)	1100 (550-1600)	39000 (21000-58000)
Uganda	2016	Male	55 to 59	75% (61-85%)	25% (15-39%)	3.4 (1.5-6.4)	14% (7.2-21%)	18% (8.8-26%)	980 (500-1500)	31000 (16000-47000)
Uganda	2016	Male	60 to 64	72% (57-83%)	28% (17-43%)	2.9 (1.2-5.8)	11% (5.2-18%)	14% (6.4-22%)	860 (400-1400)	23000 (11000-38000)
Uganda	2016	Male	65 to 69	69% (53-81%)	31% (19-47%)	2.3 (0.98-4.5)	8.5% (3.3-15%)	10% (3.9-18%)	720 (280-1300)	16000 (6300-30000)
Uganda	2016	Male	70 to 74	65% (49-77%)	35% (23-51%)	1.8 (0.78-3.6)	6.1% (2-12%)	7.2% (2.3-15%)	570 (190-1200)	10000 (3400-21000)
Uganda	2016	Male	75 to 79	62% (44-76%)	38% (24-56%)	1.5 (0.62-3)	4.8% (1.2-10%)	5.6% (1.4-12%)	380 (90-820)	5300 (1300-11000)
Uganda	2016	Male	80 to 84	58% (39-74%)	42% (26-61%)	1.3 (0.53-2.5)	3.8% (0.84-8.5%)	4.4% (0.93-9.9%)	230 (49-520)	2400 (530-5400)
Uganda	2016	Male	85 to 89	54% (36-70%)	46% (30-64%)	1 (0.41-2.1)	2.9% (0.75-6.6%)	3.3% (0.79-7.7%)	110 (25-240)	820 (210-1800)
Uganda	2016	Male	90 to 94	53% (37-69%)	47% (31-63%)	0.92 (0.35-1.9)	2% (0.39-4.9%)	2.2% (0.34-5.8%)	25 (4.1-65)	140 (27-340)
Uganda	2016	Male	95 plus	56% (35-75%)	44% (25-65%)	0.89 (0.3-1.8)	2.3% (0.4-5.7%)	2.5% (0.29-6.5%)	6.5 (0.73-16)	24 (3.9-57)
Zambia	1990	Female	All Ages	11% (9.6-14%)	89% (86-90%)	0.57 (0.41-0.77)	0.94% (0.63-1.3%)	1.7% (1.1-2.6%)	1000 (600-1600)	38000 (25000-54000)
Zambia	1990	Female	15 to 19	11% (7.5-16%)	89% (84-93%)	0.64 (0.33-1.1)	1.7% (0.86-3%)	2% (0.83-3.9%)	21 (9.1-41)	1900 (980-3300)
Zambia	1990	Female	20 to 24	15% (9.4-21%)	85% (79-91%)	0.95 (0.51-1.6)	2.1% (1.1-3.4%)	2% (0.9-3.7%)	35 (15-62)	3000 (1600-4900)
Zambia	1990	Female	25 to 29	16% (10-23%)	84% (77-90%)	1.2 (0.67-2)	2.5% (1.4-4%)	2.6% (1.3-4.6%)	47 (25-78)	3500 (2100-5500)
Zambia	1990	Female	30 to 34	18% (12-27%)	82% (73-88%)	1.4 (0.72-2.3)	3% (1.7-4.8%)	3.2% (1.6-5.5%)	54 (28-91)	3500 (2000-5600)
Zambia	1990	Female	35 to 39	21% (14-29%)	79% (71-86%)	1.3 (0.67-2.2)	3.4% (1.9-5.4%)	3.8% (2-6.5%)	61 (32-98)	3500 (2000-5400)
Zambia	1990	Female	40 to 44	22% (15-31%)	78% (69-85%)	1.3 (0.68-2.1)	3.6% (2.1-5.7%)	4.2% (2.2-6.9%)	62 (33-110)	3100 (1700-5200)
Zambia	1990	Female	45 to 49	22% (13-31%)	78% (69-87%)	1.3 (0.69-2.2)	3.9% (2.3-6.2%)	4.7% (2.6-7.5%)	72 (39-120)	3100 (1800-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	1990	Female	50 to 54	20% (13-29%)	80% (71-87%)	1.2 (0.68-2)	4.5% (2.6-7.3%)	5.4% (3.1-8.7%)	91 (49-150)	3400 (1900-5700)
Zambia	1990	Female	55 to 59	19% (12-27%)	81% (73-88%)	1.2 (0.63-2.1)	4.4% (2.3-7.1%)	5.1% (2.7-8.3%)	97 (47-170)	3100 (1600-5400)
Zambia	1990	Female	60 to 64	20% (12-29%)	80% (71-88%)	1.2 (0.59-2.2)	4.4% (2.3-7.4%)	5.1% (2.7-8.5%)	110 (53-190)	3000 (1500-5200)
Zambia	1990	Female	65 to 69	20% (13-28%)	80% (72-87%)	1.1 (0.59-1.9)	4.5% (2.5-7.5%)	5.1% (2.8-8.4%)	130 (63-210)	2900 (1500-4800)
Zambia	1990	Female	70 to 74	19% (13-27%)	81% (73-87%)	1 (0.52-1.8)	3.6% (1.5-6.6%)	4% (1.7-7.3%)	100 (40-190)	1900 (760-3500)
Zambia	1990	Female	75 to 79	17% (10-26%)	83% (74-90%)	0.85 (0.43-1.6)	3.1% (1.3-5.8%)	3.5% (1.4-6.5%)	76 (31-150)	1100 (450-2100)
Zambia	1990	Female	80 to 84	16% (9.9-24%)	84% (76-90%)	0.79 (0.42-1.3)	2.5% (1.1-4.7%)	2.8% (1.2-5.2%)	41 (17-76)	430 (190-800)
Zambia	1990	Female	85 to 89	17% (10-25%)	83% (75-90%)	0.8 (0.39-1.4)	2% (0.77-3.9%)	2.2% (0.85-4.4%)	19 (7.1-38)	140 (55-280)
Zambia	1990	Female	90 to 94	18% (12-27%)	82% (73-88%)	0.86 (0.42-1.6)	1.8% (0.55-3.9%)	2% (0.58-4.4%)	6.3 (1.9-14)	33 (10-72)
Zambia	1990	Female	95 plus	21% (12-35%)	79% (65-88%)	0.97 (0.4-2)	2.2% (0.7-4.8%)	2.4% (0.71-5.6%)	2 (0.6-4.6)	7 (2.3-16)
Zambia	1990	Male	All Ages	29% (23-34%)	71% (66-77%)	1.5 (1.1-1.9)	1.3% (0.92-1.8%)	2.1% (1.4-2.9%)	1200 (770-1700)	56000 (39000-77000)
Zambia	1990	Male	15 to 19	25% (17-35%)	75% (65-83%)	1.4 (0.78-2.1)	3.6% (2.3-5.5%)	4.2% (2.2-6.9%)	30 (16-50)	3100 (1900-4600)
Zambia	1990	Male	20 to 24	40% (28-52%)	60% (48-72%)	2.3 (1.4-3.4)	4.6% (2.9-7%)	4.3% (2.3-7.2%)	67 (37-110)	6200 (3900-9100)
Zambia	1990	Male	25 to 29	47% (34-59%)	53% (41-66%)	3.3 (2.1-4.7)	4.3% (2.6-6.7%)	4% (2.1-6.7%)	93 (55-140)	7200 (4700-10000)
Zambia	1990	Male	30 to 34	52% (39-64%)	48% (36-61%)	3.9 (2.5-5.8)	4.6% (2.8-6.9%)	4.4% (2.4-7.2%)	97 (59-150)	6600 (4300-9400)
Zambia	1990	Male	35 to 39	55% (41-66%)	45% (34-59%)	4 (2.4-5.9)	5% (3-7.6%)	5.1% (2.8-8.5%)	99 (61-150)	5800 (3700-8600)
Zambia	1990	Male	40 to 44	54% (42-65%)	46% (35-58%)	3.9 (2.4-5.7)	5.4% (3.3-8.1%)	5.7% (3.3-9%)	100 (63-150)	5300 (3400-7700)
Zambia	1990	Male	45 to 49	52% (38-63%)	48% (37-62%)	3.8 (2.2-5.7)	6.4% (3.9-9.3%)	7.1% (4.1-11%)	110 (67-160)	5000 (3300-7000)
Zambia	1990	Male	50 to 54	51% (40-63%)	49% (37-60%)	3.7 (2.2-5.7)	7.9% (5-11%)	9.2% (5.5-14%)	120 (75-180)	4900 (3100-7100)
Zambia	1990	Male	55 to 59	50% (39-62%)	50% (38-61%)	3.4 (2.1-5.2)	7.7% (4.7-11%)	9.1% (5.4-13%)	120 (68-180)	4000 (2500-5800)
Zambia	1990	Male	60 to 64	48% (34-62%)	52% (38-66%)	2.7 (1.6-4.3)	6.8% (4-10%)	8.1% (4.6-12%)	110 (61-170)	3200 (1900-4900)
Zambia	1990	Male	65 to 69	46% (32-61%)	54% (39-68%)	2.2 (1.2-3.5)	5.2% (2.8-8.2%)	6.1% (3.3-9.9%)	84 (44-140)	2000 (1100-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	1990	Male	70 to 74	44% (31-58%)	56% (42-69%)	1.8 (1-2.9)	4.1% (2-6.8%)	4.8% (2.3-8.2%)	68 (33-120)	1300 (630-2200)
Zambia	1990	Male	75 to 79	41% (27-57%)	59% (43-73%)	1.5 (0.79-2.6)	3.4% (1.5-6.1%)	3.9% (1.7-7.2%)	48 (21-88)	700 (310-1300)
Zambia	1990	Male	80 to 84	38% (23-54%)	62% (46-77%)	1.3 (0.67-2.1)	2.8% (1.3-5%)	3.2% (1.4-5.8%)	25 (11-47)	270 (120-510)
Zambia	1990	Male	85 to 89	36% (23-50%)	64% (50-77%)	1.1 (0.56-1.8)	2.1% (0.97-3.8%)	2.4% (1-4.3%)	10 (4.7-19)	83 (39-150)
Zambia	1990	Male	90 to 94	36% (23-52%)	64% (48-77%)	1 (0.51-1.8)	1.6% (0.6-3.2%)	1.7% (0.53-3.7%)	2.9 (0.91-6.1)	16 (6.1-33)
Zambia	1990	Male	95 plus	39% (21-59%)	61% (41-79%)	1 (0.42-1.9)	1.8% (0.67-3.8%)	1.8% (0.56-4.1%)	0.72 (0.21-1.6)	2.9 (1.1-6.1)
Zambia	1995	Female	All Ages	12% (10-14%)	88% (86-90%)	0.55 (0.4-0.74)	1.2% (0.77-1.7%)	2% (1.3-3%)	1600 (960-2400)	57000 (37000-84000)
Zambia	1995	Female	15 to 19	12% (7.9-17%)	88% (83-92%)	0.62 (0.33-1)	1.8% (0.88-3.1%)	2.1% (0.88-3.9%)	29 (13-54)	2500 (1300-4400)
Zambia	1995	Female	20 to 24	15% (9.9-22%)	85% (78-90%)	0.93 (0.49-1.5)	1.8% (0.95-3.3%)	1.8% (0.75-3.6%)	51 (22-95)	4100 (2200-6900)
Zambia	1995	Female	25 to 29	17% (11-24%)	83% (76-89%)	1.2 (0.65-1.9)	2% (0.97-3.5%)	2% (0.83-3.8%)	76 (38-140)	5300 (3000-8900)
Zambia	1995	Female	30 to 34	19% (12-28%)	81% (72-88%)	1.3 (0.71-2.2)	2.3% (1.2-3.8%)	2.3% (1.1-4.1%)	86 (43-150)	5300 (2900-8700)
Zambia	1995	Female	35 to 39	22% (14-29%)	78% (71-86%)	1.2 (0.66-2.1)	2.8% (1.4-4.8%)	3% (1.4-5.4%)	100 (52-170)	5400 (2900-9000)
Zambia	1995	Female	40 to 44	23% (15-32%)	77% (68-85%)	1.2 (0.63-2.1)	3.1% (1.7-5.1%)	3.3% (1.7-5.7%)	100 (53-170)	4900 (2700-8000)
Zambia	1995	Female	45 to 49	23% (14-32%)	77% (68-86%)	1.3 (0.67-2.1)	3.6% (1.9-5.8%)	4% (2.2-6.6%)	120 (64-200)	5200 (2800-8500)
Zambia	1995	Female	50 to 54	21% (14-30%)	79% (70-86%)	1.2 (0.66-2)	4.4% (2.5-6.9%)	5% (2.7-7.8%)	150 (82-250)	5700 (3100-9200)
Zambia	1995	Female	55 to 59	20% (13-29%)	80% (71-87%)	1.2 (0.61-2)	4.4% (2.4-7.2%)	4.9% (2.6-8%)	160 (79-260)	5000 (2600-8100)
Zambia	1995	Female	60 to 64	20% (13-30%)	80% (70-87%)	1.1 (0.57-2.1)	4.4% (2.3-7.6%)	4.9% (2.5-8.5%)	170 (84-290)	4500 (2300-7800)
Zambia	1995	Female	65 to 69	21% (14-29%)	79% (71-86%)	1.1 (0.59-1.8)	4.5% (2.4-7.5%)	5% (2.7-8.4%)	190 (96-320)	4200 (2200-7200)
Zambia	1995	Female	70 to 74	20% (13-27%)	80% (73-87%)	0.97 (0.5-1.7)	3.6% (1.5-6.8%)	4% (1.6-7.5%)	150 (62-280)	2800 (1200-5100)
Zambia	1995	Female	75 to 79	18% (12-26%)	82% (74-88%)	0.83 (0.41-1.5)	3.1% (1.3-5.9%)	3.4% (1.4-6.4%)	110 (45-210)	1500 (640-3000)
Zambia	1995	Female	80 to 84	17% (11-25%)	83% (75-89%)	0.77 (0.41-1.3)	2.5% (1-4.6%)	2.7% (1.1-5.1%)	58 (24-110)	610 (250-1100)
Zambia	1995	Female	85 to 89	17% (10-26%)	83% (74-90%)	0.78 (0.4-1.3)	2% (0.76-3.8%)	2.2% (0.82-4.3%)	23 (8.3-44)	170 (63-330)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	1995	Female	90 to 94	19% (12-28%)	81% (72-88%)	0.83 (0.41-1.5)	1.7% (0.52-3.7%)	1.9% (0.52-4.1%)	7.1 (2-16)	37 (11-81)
Zambia	1995	Female	95 plus	22% (12-35%)	78% (65-88%)	0.93 (0.37-1.8)	2.1% (0.69-4.7%)	2.3% (0.7-5.2%)	2.5 (0.75-5.8)	8.7 (2.8-20)
Zambia	1995	Male	All Ages	29% (24-35%)	71% (65-76%)	1.4 (1.1-1.8)	2% (1.4-2.8%)	3% (2-4.3%)	2300 (1500-3400)	100000 (72000-150000)
Zambia	1995	Male	15 to 19	26% (18-37%)	74% (63-82%)	1.3 (0.76-2)	4.4% (2.7-6.6%)	5.3% (2.8-8.3%)	57 (29-93)	5200 (3100-7900)
Zambia	1995	Male	20 to 24	41% (28-53%)	59% (47-72%)	2.2 (1.4-3.3)	6% (3.6-9.2%)	6.2% (3.2-10%)	140 (80-230)	11000 (7300-17000)
Zambia	1995	Male	25 to 29	49% (35-60%)	51% (40-65%)	3.2 (2.1-4.6)	4.9% (2.8-7.7%)	4.8% (2.5-8.1%)	200 (120-310)	14000 (9300-20000)
Zambia	1995	Male	30 to 34	53% (40-65%)	47% (35-60%)	3.7 (2.4-5.5)	4.5% (2.6-7.2%)	4.5% (2.4-7.7%)	210 (130-320)	13000 (8600-19000)
Zambia	1995	Male	35 to 39	56% (43-67%)	44% (33-57%)	3.8 (2.4-5.7)	5.1% (2.9-8.2%)	5.3% (2.9-8.9%)	210 (130-310)	11000 (7400-16000)
Zambia	1995	Male	40 to 44	55% (43-66%)	45% (34-57%)	3.7 (2.3-5.5)	5.3% (3.2-8.2%)	5.6% (3.2-9%)	200 (120-290)	9500 (6100-14000)
Zambia	1995	Male	45 to 49	53% (39-64%)	47% (36-61%)	3.7 (2.2-5.5)	6.4% (4-9.4%)	6.8% (4.1-10%)	220 (140-330)	9400 (6100-14000)
Zambia	1995	Male	50 to 54	52% (41-64%)	48% (36-59%)	3.6 (2.2-5.5)	8.3% (5.3-12%)	9.2% (5.7-13%)	250 (150-360)	9300 (5900-14000)
Zambia	1995	Male	55 to 59	51% (39-63%)	49% (37-61%)	3.2 (2-5)	8.6% (5.4-12%)	9.7% (6-14%)	240 (140-360)	7800 (4800-12000)
Zambia	1995	Male	60 to 64	50% (36-63%)	50% (37-64%)	2.6 (1.5-4.2)	7.4% (4.3-11%)	8.3% (4.7-13%)	200 (110-320)	5700 (3200-8900)
Zambia	1995	Male	65 to 69	48% (33-62%)	52% (38-67%)	2.1 (1.2-3.4)	5.7% (3.1-9.2%)	6.5% (3.5-10%)	150 (80-260)	3500 (1900-5800)
Zambia	1995	Male	70 to 74	45% (31-59%)	55% (41-69%)	1.8 (0.98-2.8)	4.5% (2.2-7.4%)	5.1% (2.4-8.4%)	120 (52-200)	2100 (1000-3600)
Zambia	1995	Male	75 to 79	43% (28-59%)	57% (41-72%)	1.5 (0.79-2.5)	3.6% (1.6-6.4%)	4.1% (1.8-7.3%)	80 (35-140)	1100 (500-2000)
Zambia	1995	Male	80 to 84	40% (25-55%)	60% (45-75%)	1.2 (0.67-2.1)	2.9% (1.3-5.1%)	3.3% (1.4-5.8%)	40 (17-71)	430 (190-760)
Zambia	1995	Male	85 to 89	37% (24-52%)	63% (48-76%)	1.1 (0.57-1.7)	2.2% (1-4%)	2.4% (1-4.4%)	15 (6-27)	120 (51-210)
Zambia	1995	Male	90 to 94	37% (23-52%)	63% (48-77%)	0.97 (0.49-1.7)	1.6% (0.56-3.2%)	1.7% (0.48-3.5%)	3.7 (1.1-7.6)	21 (7.5-41)
Zambia	1995	Male	95 plus	39% (22-59%)	61% (41-78%)	0.95 (0.4-1.8)	1.8% (0.67-3.8%)	1.9% (0.55-4.1%)	0.94 (0.28-2.1)	3.7 (1.3-7.7)
Zambia	2000	Female	All Ages	12% (11-15%)	88% (85-89%)	0.49 (0.35-0.67)	1.3% (0.78-1.9%)	2% (1.2-3.1%)	1900 (1100-2900)	72000 (42000-110000)
Zambia	2000	Female	15 to 19	13% (9-18%)	87% (82-91%)	0.57 (0.3-0.93)	1.9% (0.92-3.3%)	2.4% (0.94-4.5%)	36 (14-69)	3100 (1400-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	2000	Female	20 to 24	17% (12-23%)	83% (77-88%)	0.84 (0.46-1.4)	2.1% (1-3.9%)	2.2% (0.85-4.6%)	68 (27-130)	5400 (2700-9600)
Zambia	2000	Female	25 to 29	18% (13-25%)	82% (75-87%)	1 (0.61-1.7)	1.8% (0.88-3.4%)	1.9% (0.79-3.7%)	110 (51-200)	7500 (3900-13000)
Zambia	2000	Female	30 to 34	20% (14-28%)	80% (72-86%)	1.1 (0.62-1.9)	1.9% (0.95-3.3%)	2% (0.88-3.5%)	120 (57-220)	7400 (3800-13000)
Zambia	2000	Female	35 to 39	22% (15-29%)	78% (71-85%)	1.1 (0.59-1.8)	2.4% (1.1-4.3%)	2.5% (1.1-4.6%)	130 (60-230)	7000 (3400-12000)
Zambia	2000	Female	40 to 44	24% (18-32%)	76% (68-82%)	1.1 (0.58-1.8)	2.5% (1.3-4.4%)	2.7% (1.3-4.7%)	130 (60-220)	6000 (3000-10000)
Zambia	2000	Female	45 to 49	25% (17-32%)	75% (68-83%)	1.2 (0.64-1.9)	3.4% (1.7-5.7%)	3.7% (1.9-6.2%)	160 (82-280)	6800 (3500-11000)
Zambia	2000	Female	50 to 54	22% (15-29%)	78% (71-85%)	1 (0.58-1.7)	4% (2.1-6.5%)	4.4% (2.3-7.2%)	190 (96-330)	7100 (3600-12000)
Zambia	2000	Female	55 to 59	19% (12-27%)	81% (73-88%)	0.91 (0.48-1.6)	3.9% (1.9-6.8%)	4.4% (2.1-7.6%)	190 (89-340)	5900 (2800-11000)
Zambia	2000	Female	60 to 64	19% (12-28%)	81% (72-88%)	0.89 (0.47-1.6)	3.8% (1.9-6.7%)	4.3% (2.1-7.4%)	190 (90-330)	5000 (2500-8800)
Zambia	2000	Female	65 to 69	20% (14-27%)	80% (73-86%)	0.88 (0.48-1.4)	3.9% (2.1-6.6%)	4.3% (2.3-7.3%)	200 (110-360)	4600 (2400-7900)
Zambia	2000	Female	70 to 74	19% (13-27%)	81% (73-87%)	0.78 (0.41-1.3)	3.2% (1.3-5.9%)	3.4% (1.4-6.5%)	160 (61-320)	3000 (1100-5700)
Zambia	2000	Female	75 to 79	18% (11-26%)	82% (74-89%)	0.69 (0.33-1.2)	2.8% (1.1-5.3%)	3.1% (1.2-5.8%)	130 (49-240)	1800 (700-3400)
Zambia	2000	Female	80 to 84	17% (11-25%)	83% (75-89%)	0.66 (0.35-1.1)	2.2% (0.9-4.1%)	2.4% (0.99-4.6%)	69 (28-130)	720 (290-1400)
Zambia	2000	Female	85 to 89	18% (11-26%)	82% (74-89%)	0.66 (0.33-1.1)	1.7% (0.63-3.3%)	1.9% (0.67-3.7%)	22 (7.9-42)	160 (60-310)
Zambia	2000	Female	90 to 94	19% (12-29%)	81% (71-88%)	0.71 (0.34-1.3)	1.5% (0.37-3.2%)	1.6% (0.38-3.6%)	6.2 (1.4-14)	32 (8.1-71)
Zambia	2000	Female	95 plus	23% (13-36%)	77% (64-87%)	0.79 (0.32-1.5)	1.8% (0.55-4.1%)	2% (0.55-4.5%)	2.2 (0.6-4.9)	7.7 (2.4-17)
Zambia	2000	Male	All Ages	32% (28-38%)	68% (62-72%)	1.3 (0.97-1.7)	3% (2-4.1%)	4.2% (2.7-5.9%)	4000 (2500-5700)	180000 (120000-240000)
Zambia	2000	Male	15 to 19	31% (23-39%)	69% (61-77%)	1.3 (0.78-2)	5.5% (3.3-8.4%)	6.5% (3.5-10%)	110 (57-180)	8900 (5200-14000)
Zambia	2000	Male	20 to 24	46% (36-56%)	54% (44-64%)	2.1 (1.4-3.1)	8.9% (5.3-13%)	9.7% (5.4-15%)	280 (160-420)	21000 (13000-30000)
Zambia	2000	Male	25 to 29	53% (43-62%)	47% (38-57%)	2.9 (2-4.1)	6.9% (3.9-11%)	7.2% (3.7-12%)	380 (230-560)	25000 (17000-36000)
Zambia	2000	Male	30 to 34	57% (46-66%)	43% (34-54%)	3.4 (2.3-4.7)	5.7% (3.3-8.8%)	5.9% (3.2-9.4%)	400 (250-560)	24000 (15000-32000)
Zambia	2000	Male	35 to 39	58% (49-67%)	42% (33-51%)	3.4 (2.2-5)	5.8% (3.4-9%)	6.1% (3.4-9.8%)	370 (240-530)	20000 (13000-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	2000	Male	40 to 44	58% (48-66%)	42% (34-52%)	3.3 (2.1-4.8)	6.3% (3.9-9.2%)	6.6% (4-9.8%)	350 (210-510)	16000 (10000-24000)
Zambia	2000	Male	45 to 49	54% (43-64%)	46% (36-57%)	3.1 (1.9-4.6)	7.2% (4.4-11%)	7.7% (4.5-12%)	350 (200-510)	15000 (8900-21000)
Zambia	2000	Male	50 to 54	53% (44-63%)	47% (37-56%)	3.1 (1.9-4.6)	9.5% (5.8-14%)	10% (6.3-15%)	430 (240-640)	16000 (9200-23000)
Zambia	2000	Male	55 to 59	52% (42-62%)	48% (38-58%)	2.7 (1.8-4.1)	10% (6.1-15%)	11% (6.7-16%)	420 (230-640)	13000 (7600-20000)
Zambia	2000	Male	60 to 64	48% (36-61%)	52% (39-64%)	2.1 (1.3-3.4)	7.9% (4.6-12%)	8.8% (5-14%)	340 (180-550)	9100 (5100-15000)
Zambia	2000	Male	65 to 69	46% (32-60%)	54% (40-68%)	1.7 (0.97-2.7)	6% (3.2-9.7%)	6.7% (3.6-11%)	240 (120-420)	5500 (2800-9300)
Zambia	2000	Male	70 to 74	44% (30-58%)	56% (42-70%)	1.4 (0.78-2.3)	4.5% (2-7.7%)	5% (2.2-8.6%)	170 (76-310)	3100 (1400-5500)
Zambia	2000	Male	75 to 79	42% (28-56%)	58% (44-72%)	1.2 (0.66-2)	3.6% (1.5-6.6%)	4% (1.6-7.3%)	110 (44-200)	1500 (630-2800)
Zambia	2000	Male	80 to 84	39% (24-55%)	61% (45-76%)	1 (0.56-1.7)	2.8% (1.2-5.2%)	3.2% (1.3-5.9%)	56 (22-110)	590 (250-1100)
Zambia	2000	Male	85 to 89	37% (23-51%)	63% (49-77%)	0.87 (0.48-1.4)	2.2% (0.9-3.9%)	2.4% (0.93-4.4%)	18 (7-34)	140 (58-260)
Zambia	2000	Male	90 to 94	37% (23-51%)	63% (49-77%)	0.81 (0.4-1.4)	1.5% (0.52-3.2%)	1.6% (0.48-3.5%)	4.1 (1.2-8.8)	23 (7.9-47)
Zambia	2000	Male	95 plus	40% (22-59%)	60% (41-78%)	0.8 (0.34-1.6)	1.9% (0.65-4.2%)	2% (0.52-4.6%)	1.2 (0.31-2.7)	4.5 (1.5-9.7)
Zambia	2005	Female	All Ages	12% (10-13%)	88% (87-90%)	0.42 (0.29-0.57)	1.2% (0.72-1.8%)	1.9% (1.1-3%)	1800 (970-2900)	66000 (38000-100000)
Zambia	2005	Female	15 to 19	11% (7.9-15%)	89% (85-92%)	0.44 (0.23-0.73)	1.4% (0.68-2.5%)	1.6% (0.6-3.2%)	29 (11-61)	2700 (1300-5100)
Zambia	2005	Female	20 to 24	15% (11-20%)	85% (80-89%)	0.68 (0.37-1.1)	2% (0.97-3.6%)	2.1% (0.74-4.2%)	57 (23-110)	4700 (2400-8400)
Zambia	2005	Female	25 to 29	17% (12-22%)	83% (78-88%)	0.88 (0.51-1.4)	1.8% (0.84-3.4%)	1.8% (0.7-3.9%)	99 (43-180)	7000 (3600-12000)
Zambia	2005	Female	30 to 34	18% (13-26%)	82% (74-87%)	0.97 (0.53-1.6)	1.6% (0.76-2.9%)	1.6% (0.68-3.1%)	110 (51-220)	7000 (3500-13000)
Zambia	2005	Female	35 to 39	21% (15-27%)	79% (73-85%)	0.92 (0.51-1.6)	1.8% (0.86-3.4%)	1.9% (0.82-3.7%)	120 (52-210)	6300 (3100-11000)
Zambia	2005	Female	40 to 44	24% (17-30%)	76% (70-83%)	0.96 (0.52-1.5)	2% (0.97-3.5%)	2.1% (0.98-3.7%)	110 (48-190)	5100 (2500-8800)
Zambia	2005	Female	45 to 49	24% (17-31%)	76% (69-83%)	1 (0.56-1.7)	2.8% (1.4-4.8%)	3.1% (1.4-5.3%)	140 (61-230)	5700 (2700-9700)
Zambia	2005	Female	50 to 54	22% (15-29%)	78% (71-85%)	0.95 (0.53-1.6)	3.6% (1.8-6.1%)	4% (2-6.8%)	170 (76-310)	6300 (2900-11000)
Zambia	2005	Female	55 to 59	19% (12-26%)	81% (74-88%)	0.83 (0.44-1.4)	3.7% (1.7-6.4%)	4.2% (1.9-7.3%)	180 (78-330)	5800 (2500-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	2005	Female	60 to 64	19% (12-27%)	81% (73-88%)	0.81 (0.41-1.4)	3.6% (1.7-6.4%)	4.1% (1.9-7.4%)	180 (85-330)	4900 (2300-8900)
Zambia	2005	Female	65 to 69	20% (14-27%)	80% (73-86%)	0.81 (0.45-1.3)	3.8% (2-6.4%)	4.2% (2.2-7.1%)	200 (100-360)	4500 (2300-8000)
Zambia	2005	Female	70 to 74	19% (13-27%)	81% (73-87%)	0.72 (0.37-1.2)	3% (1.2-5.8%)	3.3% (1.2-6.5%)	170 (65-330)	3000 (1200-6100)
Zambia	2005	Female	75 to 79	18% (12-26%)	82% (74-88%)	0.64 (0.31-1.1)	2.8% (1.1-5.2%)	3.1% (1.2-5.9%)	140 (53-270)	1900 (750-3700)
Zambia	2005	Female	80 to 84	17% (11-26%)	83% (74-89%)	0.61 (0.32-1)	2.2% (0.86-4.1%)	2.4% (0.94-4.6%)	83 (32-160)	860 (340-1700)
Zambia	2005	Female	85 to 89	18% (11-27%)	82% (73-89%)	0.61 (0.31-1)	1.8% (0.65-3.5%)	2% (0.71-4%)	26 (9.2-53)	190 (70-390)
Zambia	2005	Female	90 to 94	20% (12-29%)	80% (71-88%)	0.65 (0.32-1.2)	1.5% (0.42-3.3%)	1.7% (0.43-3.8%)	6.3 (1.5-14)	32 (8.6-70)
Zambia	2005	Female	95 plus	22% (12-36%)	78% (64-88%)	0.73 (0.31-1.4)	1.9% (0.55-4.3%)	2.1% (0.56-4.9%)	2 (0.54-4.7)	7.1 (2-16)
Zambia	2005	Male	All Ages	32% (28-38%)	68% (62-72%)	1.2 (0.88-1.6)	3.9% (2.7-5.4%)	5.4% (3.6-7.4%)	5500 (3500-7700)	240000 (160000-330000)
Zambia	2005	Male	15 to 19	31% (23-39%)	69% (61-77%)	1.2 (0.7-1.9)	5.1% (2.9-7.9%)	5.8% (3-9.4%)	140 (73-230)	11000 (6400-18000)
Zambia	2005	Male	20 to 24	47% (36-56%)	53% (44-64%)	2 (1.3-2.9)	10% (6.1-15%)	11% (6.2-17%)	350 (190-520)	26000 (15000-37000)
Zambia	2005	Male	25 to 29	53% (44-62%)	47% (38-56%)	2.7 (1.9-3.8)	9.6% (5.5-15%)	10% (5.4-17%)	510 (320-730)	34000 (22000-47000)
Zambia	2005	Male	30 to 34	57% (47-66%)	43% (34-53%)	3.1 (2.1-4.4)	7.7% (4.6-11%)	8% (4.5-12%)	550 (360-780)	33000 (21000-45000)
Zambia	2005	Male	35 to 39	59% (49-68%)	41% (32-51%)	3.1 (2-4.6)	7.2% (4.5-11%)	7.6% (4.5-12%)	530 (340-770)	28000 (18000-40000)
Zambia	2005	Male	40 to 44	58% (48-67%)	42% (33-52%)	3 (1.9-4.4)	7.4% (4.8-11%)	7.8% (5-11%)	490 (310-720)	23000 (15000-33000)
Zambia	2005	Male	45 to 49	54% (43-64%)	46% (36-57%)	2.9 (1.8-4.3)	8.8% (5.3-13%)	9.5% (5.5-14%)	480 (290-720)	20000 (12000-30000)
Zambia	2005	Male	50 to 54	53% (44-63%)	47% (37-56%)	2.8 (1.8-4.2)	11% (6.8-16%)	12% (7.4-17%)	550 (320-800)	20000 (12000-29000)
Zambia	2005	Male	55 to 59	53% (43-63%)	47% (37-57%)	2.5 (1.6-3.7)	12% (6.9-17%)	13% (7.6-19%)	570 (330-870)	18000 (10000-27000)
Zambia	2005	Male	60 to 64	49% (36-62%)	51% (38-64%)	2 (1.1-3)	8.9% (4.9-14%)	9.7% (5.4-15%)	460 (250-740)	13000 (7000-20000)
Zambia	2005	Male	65 to 69	46% (32-60%)	54% (40-68%)	1.6 (0.9-2.4)	6.6% (3.4-11%)	7.3% (3.8-12%)	350 (180-580)	7900 (4100-13000)
Zambia	2005	Male	70 to 74	44% (30-58%)	56% (42-70%)	1.3 (0.7-2.1)	4.8% (2.1-8.3%)	5.3% (2.3-9.1%)	240 (100-420)	4300 (1800-7500)
Zambia	2005	Male	75 to 79	42% (27-57%)	58% (43-73%)	1.1 (0.57-1.8)	3.8% (1.4-7%)	4.2% (1.5-7.8%)	150 (51-290)	2200 (740-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	2005	Male	80 to 84	39% (24-55%)	61% (45-76%)	0.93 (0.49-1.5)	2.9% (1.1-5.4%)	3.2% (1.1-6.1%)	79 (28-150)	830 (310-1600)
Zambia	2005	Male	85 to 89	36% (23-52%)	64% (48-77%)	0.79 (0.41-1.4)	2.3% (0.85-4.3%)	2.5% (0.86-4.8%)	24 (8-45)	180 (66-340)
Zambia	2005	Male	90 to 94	37% (23-52%)	63% (48-77%)	0.73 (0.37-1.3)	1.6% (0.48-3.3%)	1.7% (0.42-3.7%)	4.6 (1.1-10)	25 (7.7-53)
Zambia	2005	Male	95 plus	40% (22-59%)	60% (41-78%)	0.72 (0.31-1.4)	2.1% (0.59-4.5%)	2.2% (0.47-5%)	1.4 (0.29-3.2)	5.2 (1.5-11)
Zambia	2010	Female	All Ages	9% (7.9-10%)	91% (90-92%)	0.37 (0.25-0.55)	1.3% (0.77-2.1%)	2.3% (1.3-3.7%)	1900 (970-3100)	63000 (35000-99000)
Zambia	2010	Female	15 to 19	5.3% (3.9-8.4%)	95% (92-96%)	0.25 (0.12-0.44)	0.79% (0.4-1.4%)	0.74% (0.27-1.6%)	16 (5.4-34)	1700 (850-3100)
Zambia	2010	Female	20 to 24	9.6% (7.6-13%)	90% (87-92%)	0.51 (0.27-0.87)	1.6% (0.81-2.9%)	1.6% (0.55-3.2%)	41 (16-82)	3800 (2000-6700)
Zambia	2010	Female	25 to 29	13% (10-15%)	87% (85-90%)	0.76 (0.44-1.2)	1.7% (0.83-3.2%)	1.7% (0.66-3.5%)	76 (33-140)	5700 (3100-9700)
Zambia	2010	Female	30 to 34	14% (12-17%)	86% (83-88%)	0.83 (0.47-1.3)	1.7% (0.86-3%)	1.7% (0.74-3.4%)	97 (46-190)	6200 (3400-11000)
Zambia	2010	Female	35 to 39	17% (14-20%)	83% (80-86%)	0.84 (0.49-1.4)	1.9% (0.91-3.4%)	2% (0.85-3.7%)	110 (50-200)	6000 (3000-11000)
Zambia	2010	Female	40 to 44	19% (16-22%)	81% (78-84%)	0.88 (0.5-1.5)	2.2% (1.1-3.7%)	2.3% (1.1-4.1%)	97 (44-180)	4800 (2300-8500)
Zambia	2010	Female	45 to 49	20% (16-24%)	80% (76-84%)	1 (0.55-1.6)	3.2% (1.6-5.3%)	3.6% (1.7-6.2%)	120 (51-200)	4900 (2300-8500)
Zambia	2010	Female	50 to 54	21% (15-28%)	79% (72-85%)	1 (0.55-1.8)	4.4% (2.2-7.7%)	5.2% (2.6-9.1%)	150 (67-280)	5700 (2600-10000)
Zambia	2010	Female	55 to 59	20% (13-29%)	80% (71-87%)	1 (0.52-1.8)	4.9% (2.4-8.5%)	5.8% (2.8-10%)	190 (82-350)	6000 (2700-11000)
Zambia	2010	Female	60 to 64	21% (13-30%)	79% (70-87%)	1 (0.51-1.8)	4.8% (2.3-8.4%)	5.6% (2.7-10%)	210 (94-380)	5600 (2600-10000)
Zambia	2010	Female	65 to 69	21% (14-29%)	79% (71-86%)	1 (0.52-1.7)	4.8% (2.5-8.5%)	5.5% (2.9-9.9%)	240 (120-430)	5300 (2700-9500)
Zambia	2010	Female	70 to 74	20% (14-28%)	80% (72-86%)	0.87 (0.45-1.5)	3.8% (1.7-7.3%)	4.3% (1.8-8.4%)	200 (82-400)	3700 (1500-7200)
Zambia	2010	Female	75 to 79	18% (11-27%)	82% (73-89%)	0.75 (0.36-1.4)	3.4% (1.4-6.6%)	3.8% (1.6-7.5%)	170 (71-350)	2400 (1000-4800)
Zambia	2010	Female	80 to 84	17% (11-25%)	83% (75-89%)	0.7 (0.34-1.2)	2.7% (1-5.1%)	3% (1.2-5.7%)	110 (42-220)	1200 (450-2300)
Zambia	2010	Female	85 to 89	18% (11-27%)	82% (73-89%)	0.71 (0.33-1.3)	2.2% (0.79-4.6%)	2.5% (0.88-5.2%)	40 (14-85)	290 (100-630)
Zambia	2010	Female	90 to 94	20% (12-29%)	80% (71-88%)	0.75 (0.35-1.4)	2% (0.51-4.5%)	2.3% (0.55-5.3%)	9.4 (2.3-22)	49 (12-110)
Zambia	2010	Female	95 plus	22% (12-35%)	78% (65-88%)	0.83 (0.32-1.7)	2.4% (0.71-5.5%)	2.7% (0.76-6.2%)	2.4 (0.68-5.4)	8.3 (2.5-18)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	2010	Male	All Ages	28% (24-33%)	72% (67-76%)	1.3 (0.86-1.8)	5.1% (3.4-6.9%)	7.1% (4.6-9.9%)	6600 (4100-9400)	280000 (190000-390000)
Zambia	2010	Male	15 to 19	17% (11-27%)	83% (73-89%)	0.76 (0.38-1.4)	3.4% (1.8-5.9%)	3.6% (1.6-6.6%)	100 (45-190)	8800 (4400-15000)
Zambia	2010	Male	20 to 24	37% (29-46%)	63% (54-71%)	1.8 (1.1-2.8)	9.8% (5.7-15%)	11% (5.6-16%)	360 (180-550)	27000 (15000-39000)
Zambia	2010	Male	25 to 29	50% (41-57%)	50% (43-59%)	2.9 (1.9-4.3)	12% (6.7-17%)	13% (6.8-20%)	570 (340-830)	38000 (24000-53000)
Zambia	2010	Male	30 to 34	54% (45-63%)	46% (37-55%)	3.4 (2.2-4.9)	11% (6.5-16%)	12% (6.7-18%)	670 (430-950)	40000 (26000-55000)
Zambia	2010	Male	35 to 39	56% (49-64%)	44% (36-51%)	3.5 (2.2-5.1)	10% (6.7-15%)	11% (7.1-16%)	680 (430-1000)	36000 (23000-51000)
Zambia	2010	Male	40 to 44	56% (47-64%)	44% (36-53%)	3.3 (2.1-5.1)	11% (6.9-16%)	12% (7.4-17%)	650 (390-930)	30000 (18000-44000)
Zambia	2010	Male	45 to 49	53% (44-61%)	47% (39-56%)	3.3 (1.9-4.9)	13% (7.9-18%)	14% (8.7-20%)	600 (350-900)	25000 (15000-37000)
Zambia	2010	Male	50 to 54	51% (42-60%)	49% (40-58%)	3.1 (1.8-4.8)	15% (9.2-21%)	17% (10-24%)	620 (360-920)	22000 (13000-33000)
Zambia	2010	Male	55 to 59	51% (42-59%)	49% (41-58%)	2.8 (1.7-4.3)	15% (8.9-21%)	17% (9.8-24%)	600 (330-890)	19000 (11000-28000)
Zambia	2010	Male	60 to 64	50% (37-62%)	50% (38-63%)	2.3 (1.3-3.8)	11% (6.2-17%)	13% (6.9-19%)	550 (290-870)	15000 (8100-23000)
Zambia	2010	Male	65 to 69	48% (33-62%)	52% (38-67%)	1.9 (1-3.1)	8.6% (4.4-14%)	9.6% (4.8-15%)	460 (220-750)	10000 (5100-17000)
Zambia	2010	Male	70 to 74	45% (31-59%)	55% (41-69%)	1.6 (0.84-2.6)	6.2% (2.8-11%)	6.9% (3.1-12%)	330 (150-600)	6000 (2600-11000)
Zambia	2010	Male	75 to 79	42% (28-59%)	58% (41-72%)	1.3 (0.66-2.2)	4.9% (2-9%)	5.4% (2.2-10%)	220 (90-410)	3100 (1200-5700)
Zambia	2010	Male	80 to 84	40% (25-56%)	60% (44-75%)	1.1 (0.54-1.8)	3.7% (1.3-7%)	4.1% (1.4-7.7%)	120 (42-230)	1300 (460-2400)
Zambia	2010	Male	85 to 89	37% (23-52%)	63% (48-77%)	0.92 (0.46-1.6)	2.9% (1.1-5.5%)	3.2% (1.2-6.1%)	36 (13-69)	270 (100-510)
Zambia	2010	Male	90 to 94	37% (23-52%)	63% (48-77%)	0.84 (0.4-1.5)	2% (0.62-4.3%)	2.2% (0.6-4.8%)	6.7 (1.8-15)	36 (11-76)
Zambia	2010	Male	95 plus	39% (23-59%)	61% (41-77%)	0.83 (0.33-1.7)	2.6% (0.78-5.7%)	2.8% (0.72-6.3%)	1.9 (0.49-4.4)	7.1 (2.1-15)
Zambia	2016	Female	All Ages	10% (8.5-12%)	90% (88-91%)	0.43 (0.22-0.73)	1.5% (0.77-2.6%)	2.6% (1.1-4.5%)	1700 (700-3300)	60000 (29000-100000)
Zambia	2016	Female	15 to 19	8.6% (5.1-13%)	91% (87-95%)	0.4 (0.15-0.82)	1% (0.48-2%)	1% (0.28-2.5%)	20 (5.4-47)	2200 (1000-4300)
Zambia	2016	Female	20 to 24	12% (8-17%)	88% (83-92%)	0.63 (0.26-1.3)	1.8% (0.92-3.3%)	1.7% (0.52-3.8%)	37 (12-80)	3900 (2000-7000)
Zambia	2016	Female	25 to 29	14% (10-20%)	86% (80-90%)	0.85 (0.39-1.6)	2.1% (1-3.7%)	2.1% (0.73-4.4%)	66 (25-130)	5300 (2700-9600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	2016	Female	30 to 34	16% (12-23%)	84% (77-88%)	0.97 (0.43-1.9)	2.4% (1.2-4.3%)	2.7% (1-5.3%)	89 (34-190)	6000 (2800-11000)
Zambia	2016	Female	35 to 39	18% (13-25%)	82% (75-87%)	0.94 (0.41-1.8)	2.8% (1.4-5.2%)	3.3% (1.4-6.3%)	110 (41-240)	6200 (2800-12000)
Zambia	2016	Female	40 to 44	20% (14-26%)	80% (74-86%)	0.93 (0.39-1.8)	2.9% (1.4-5%)	3.5% (1.6-6.1%)	100 (40-210)	5200 (2300-10000)
Zambia	2016	Female	45 to 49	21% (14-27%)	79% (73-86%)	1 (0.44-2)	3.6% (1.7-6.6%)	4.5% (1.9-8.2%)	110 (43-240)	4900 (2000-9900)
Zambia	2016	Female	50 to 54	20% (14-29%)	80% (71-86%)	1 (0.43-2.1)	4.3% (2-7.6%)	5.3% (2.4-9.4%)	130 (48-260)	4800 (1900-9300)
Zambia	2016	Female	55 to 59	20% (13-29%)	80% (71-87%)	1 (0.42-2.1)	4.5% (1.8-8.4%)	5.7% (2.3-10%)	140 (49-290)	4500 (1600-9000)
Zambia	2016	Female	60 to 64	20% (13-30%)	80% (70-87%)	1 (0.41-2)	4.4% (1.7-8.1%)	5.4% (2.1-10%)	170 (61-330)	4600 (1700-9000)
Zambia	2016	Female	65 to 69	21% (14-29%)	79% (71-86%)	0.99 (0.44-2)	4.5% (1.9-8.2%)	5.3% (2.3-9.8%)	210 (81-410)	4800 (1900-9200)
Zambia	2016	Female	70 to 74	20% (14-28%)	80% (72-86%)	0.88 (0.39-1.8)	3.6% (1.3-7.3%)	4.1% (1.4-8.6%)	190 (59-410)	3400 (1100-7500)
Zambia	2016	Female	75 to 79	18% (12-28%)	82% (72-88%)	0.76 (0.32-1.6)	3.3% (1.2-6.8%)	3.8% (1.3-7.7%)	170 (57-360)	2400 (810-5100)
Zambia	2016	Female	80 to 84	17% (11-26%)	83% (74-89%)	0.71 (0.29-1.4)	2.6% (0.75-5.7%)	3% (0.88-6.6%)	120 (35-270)	1200 (360-2800)
Zambia	2016	Female	85 to 89	18% (11-26%)	82% (74-89%)	0.71 (0.27-1.4)	2.2% (0.58-5%)	2.5% (0.65-5.8%)	50 (13-120)	370 (98-900)
Zambia	2016	Female	90 to 94	19% (12-29%)	81% (71-88%)	0.76 (0.3-1.6)	1.9% (0.29-5%)	2.2% (0.28-5.8%)	13 (1.7-35)	68 (10-180)
Zambia	2016	Female	95 plus	22% (13-36%)	78% (64-87%)	0.85 (0.29-1.9)	2.4% (0.63-6%)	2.7% (0.65-6.8%)	3.2 (0.75-7.9)	11 (2.8-27)
Zambia	2016	Male	All Ages	29% (25-33%)	71% (67-75%)	1.3 (0.68-2.2)	5.4% (3.1-7.9%)	7.3% (4-11%)	6200 (3100-10000)	270000 (150000-410000)
Zambia	2016	Male	15 to 19	17% (11-26%)	83% (74-89%)	0.76 (0.32-1.5)	3.1% (1.5-5.4%)	3.2% (1.2-6.1%)	93 (32-180)	8400 (4000-15000)
Zambia	2016	Male	20 to 24	37% (29-45%)	63% (55-71%)	1.8 (0.87-3.4)	9% (4.7-14%)	9.7% (4.5-16%)	320 (140-540)	25000 (13000-40000)
Zambia	2016	Male	25 to 29	50% (41-57%)	50% (43-59%)	2.9 (1.5-5.2)	11% (6.1-17%)	13% (6.1-20%)	530 (270-820)	36000 (20000-54000)
Zambia	2016	Male	30 to 34	54% (45-62%)	46% (38-55%)	3.4 (1.7-5.8)	12% (6.6-17%)	13% (6.7-19%)	610 (320-920)	37000 (21000-55000)
Zambia	2016	Male	35 to 39	56% (48-65%)	44% (35-52%)	3.5 (1.8-5.9)	12% (6.8-17%)	13% (7.4-20%)	660 (350-1000)	36000 (19000-55000)
Zambia	2016	Male	40 to 44	56% (47-64%)	44% (36-53%)	3.3 (1.7-5.8)	12% (7-18%)	14% (7.7-21%)	670 (340-1100)	32000 (17000-51000)
Zambia	2016	Male	45 to 49	53% (44-61%)	47% (39-56%)	3.3 (1.6-5.8)	13% (7.3-19%)	15% (8.2-22%)	640 (310-1100)	27000 (14000-44000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zambia	2016	Male	50 to 54	51% (42-60%)	49% (40-58%)	3.1 (1.4-5.7)	14% (7.3-20%)	16% (8.4-23%)	610 (290-1000)	23000 (11000-37000)
Zambia	2016	Male	55 to 59	51% (42-60%)	49% (40-58%)	2.9 (1.3-5.3)	13% (6.8-20%)	15% (8-23%)	530 (230-870)	17000 (7700-27000)
Zambia	2016	Male	60 to 64	49% (37-62%)	51% (38-63%)	2.4 (1.1-4.4)	10% (4.9-17%)	12% (5.5-19%)	440 (190-740)	12000 (5300-20000)
Zambia	2016	Male	65 to 69	47% (34-62%)	53% (38-66%)	1.9 (0.85-3.5)	7.7% (3.3-14%)	8.8% (3.7-16%)	390 (150-750)	8800 (3500-17000)
Zambia	2016	Male	70 to 74	45% (31-59%)	55% (41-69%)	1.6 (0.69-3)	5.7% (2.1-11%)	6.4% (2.5-13%)	310 (110-650)	5700 (2100-12000)
Zambia	2016	Male	75 to 79	42% (28-58%)	58% (42-72%)	1.3 (0.55-2.6)	4.5% (1.4-9.6%)	5.1% (1.6-11%)	230 (69-500)	3300 (980-7000)
Zambia	2016	Male	80 to 84	39% (25-55%)	61% (45-75%)	1.1 (0.44-2.2)	3.5% (0.99-7.7%)	3.9% (1.1-8.6%)	130 (35-300)	1400 (380-3200)
Zambia	2016	Male	85 to 89	36% (23-51%)	64% (49-77%)	0.92 (0.37-1.9)	2.7% (0.81-6.1%)	3% (0.85-7%)	42 (11-96)	320 (93-710)
Zambia	2016	Male	90 to 94	36% (23-51%)	64% (49-77%)	0.84 (0.34-1.8)	2% (0.49-4.8%)	2.1% (0.44-5.4%)	8.1 (1.7-21)	44 (11-110)
Zambia	2016	Male	95 plus	39% (22-59%)	61% (41-78%)	0.83 (0.29-1.8)	2.5% (0.72-5.9%)	2.7% (0.61-6.7%)	2.1 (0.49-5.4)	7.8 (2.2-19)
Botswana	1990	Female	All Ages	16% (12-20%)	84% (80-88%)	0.83 (0.63-1.1)	2.6% (1.6-3.8%)	4.1% (2-6.6%)	160 (78-250)	6000 (3800-8700)
Botswana	1990	Female	15 to 19	14% (7.7-24%)	86% (76-92%)	0.65 (0.27-1.2)	2% (0.98-3.3%)	2.8% (1.1-5.5%)	2.1 (0.75-4)	230 (110-380)
Botswana	1990	Female	20 to 24	19% (9.6-30%)	81% (70-90%)	1.2 (0.56-2.1)	3.1% (1.7-5.3%)	3.5% (1.3-7.2%)	4.1 (1.6-8.1)	430 (240-700)
Botswana	1990	Female	25 to 29	21% (11-35%)	79% (65-89%)	1.5 (0.74-2.6)	3.6% (2-5.8%)	4.2% (1.7-7.8%)	5.5 (2.5-9.8)	470 (270-760)
Botswana	1990	Female	30 to 34	24% (12-40%)	76% (60-88%)	1.8 (0.91-3)	4% (2.2-6.5%)	5.2% (2.3-9.5%)	7.2 (3.3-12)	520 (300-840)
Botswana	1990	Female	35 to 39	27% (15-43%)	73% (57-85%)	1.9 (0.92-3.2)	4.9% (2.7-7.8%)	6.7% (3.2-11%)	9.6 (4.8-17)	580 (320-950)
Botswana	1990	Female	40 to 44	29% (16-45%)	71% (55-84%)	1.9 (0.86-3.4)	5.8% (3-9.7%)	8.2% (3.9-14%)	10 (4.7-18)	520 (260-890)
Botswana	1990	Female	45 to 49	29% (14-45%)	71% (55-86%)	1.9 (0.8-3.7)	6.4% (3.2-11%)	8.9% (4.1-15%)	13 (6-23)	570 (280-990)
Botswana	1990	Female	50 to 54	29% (15-46%)	71% (54-85%)	1.9 (0.8-3.7)	6.6% (3-11%)	8.7% (3.9-15%)	16 (6.9-29)	620 (280-1100)
Botswana	1990	Female	55 to 59	28% (16-45%)	72% (55-84%)	1.8 (0.8-3.4)	6.5% (2.8-11%)	8.3% (3.5-15%)	18 (7.3-32)	590 (250-1100)
Botswana	1990	Female	60 to 64	29% (16-46%)	71% (54-84%)	1.8 (0.73-3.5)	6% (2-11%)	7.6% (2.5-14%)	18 (5.7-35)	500 (160-960)
Botswana	1990	Female	65 to 69	30% (17-45%)	70% (55-83%)	1.7 (0.78-3)	5.5% (1.7-11%)	6.7% (2.1-13%)	21 (6.4-41)	480 (150-930)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	1990	Female	70 to 74	28% (16-43%)	72% (57-84%)	1.4 (0.59-2.5)	4% (0.69-8.5%)	4.8% (0.85-10%)	14 (2.5-30)	260 (46-560)
Botswana	1990	Female	75 to 79	25% (14-41%)	75% (59-86%)	1.1 (0.43-2.3)	2.8% (-0.077-7.2%)	3.3% (-0.093-8.4%)	8.2 (-0.21-20)	120 (-2.9-290)
Botswana	1990	Female	80 to 84	24% (13-39%)	76% (61-87%)	1 (0.41-2)	2.3% (-0.11-5.9%)	2.6% (-0.15-6.9%)	5.5 (-0.31-15)	58 (-2.7-160)
Botswana	1990	Female	85 to 89	25% (13-40%)	75% (60-87%)	1 (0.4-1.9)	1.9% (-0.21-5.2%)	2.1% (-0.3-6%)	3.1 (-0.44-9)	24 (-2.7-67)
Botswana	1990	Female	90 to 94	27% (15-43%)	73% (57-85%)	1 (0.42-2)	1.5% (-0.57-4.9%)	1.7% (-0.75-5.7%)	1.1 (-0.47-3.5)	5.6 (-2.1-18)
Botswana	1990	Female	95 plus	32% (16-51%)	68% (49-84%)	1.1 (0.38-2.5)	1.5% (-1.5-8%)	1.6% (-1.3-6.6%)	0.25 (-0.2-1)	0.89 (-0.6-3.4)
Botswana	1990	Male	All Ages	30% (24-38%)	70% (62-76%)	1.8 (1.4-2.3)	6.2% (4.6-8.3%)	8.6% (6-12%)	400 (270-560)	18000 (13000-24000)
Botswana	1990	Male	15 to 19	26% (16-40%)	74% (60-84%)	1.3 (0.66-2.3)	5.5% (3-8.9%)	7.1% (3.5-12%)	9.4 (4.4-17)	820 (450-1400)
Botswana	1990	Male	20 to 24	39% (25-56%)	61% (44-75%)	2.8 (1.5-4.5)	10% (6.3-15%)	12% (6.5-18%)	26 (14-42)	2100 (1200-3100)
Botswana	1990	Male	25 to 29	46% (29-63%)	54% (37-71%)	3.9 (2.3-5.9)	11% (6.5-16%)	12% (6.5-19%)	33 (20-48)	2300 (1500-3200)
Botswana	1990	Male	30 to 34	49% (31-66%)	51% (34-69%)	4.3 (2.6-6.7)	11% (6.7-15%)	12% (7.1-19%)	34 (21-50)	2200 (1400-3100)
Botswana	1990	Male	35 to 39	51% (34-68%)	49% (32-66%)	4.3 (2.5-6.7)	12% (7.2-17%)	14% (8-20%)	38 (23-56)	2100 (1400-3000)
Botswana	1990	Male	40 to 44	51% (35-69%)	49% (31-65%)	4.2 (2.2-6.7)	13% (7.9-18%)	15% (9.1-22%)	35 (20-52)	1700 (1000-2500)
Botswana	1990	Male	45 to 49	51% (33-68%)	49% (32-67%)	4.2 (2.2-7.3)	15% (9.6-21%)	18% (12-25%)	41 (25-61)	1800 (1100-2600)
Botswana	1990	Male	50 to 54	51% (32-69%)	49% (31-68%)	4.1 (2-7)	15% (9.5-21%)	18% (11-26%)	44 (26-66)	1700 (1000-2500)
Botswana	1990	Male	55 to 59	50% (33-69%)	50% (31-67%)	3.8 (1.9-6.4)	14% (8.7-20%)	17% (10-24%)	42 (23-65)	1400 (780-2100)
Botswana	1990	Male	60 to 64	48% (32-67%)	52% (33-68%)	3.2 (1.5-5.7)	12% (6.1-17%)	14% (7-21%)	34 (17-55)	940 (470-1500)
Botswana	1990	Male	65 to 69	45% (30-62%)	55% (38-70%)	2.5 (1.2-4.2)	9% (4.4-15%)	11% (5.2-17%)	31 (14-52)	710 (330-1200)
Botswana	1990	Male	70 to 74	42% (27-58%)	58% (42-73%)	1.8 (0.87-3)	5.5% (2.1-9.5%)	6.2% (2.5-11%)	16 (5.9-30)	300 (120-550)
Botswana	1990	Male	75 to 79	41% (27-58%)	59% (42-73%)	1.4 (0.63-2.6)	3.7% (1-7.7%)	4.2% (1.1-8.6%)	8 (2.2-17)	120 (33-240)
Botswana	1990	Male	80 to 84	39% (25-57%)	61% (43-75%)	1.2 (0.58-2.3)	2.8% (0.71-6.1%)	3.1% (0.71-6.9%)	4.2 (0.93-9)	45 (11-97)
Botswana	1990	Male	85 to 89	36% (23-51%)	64% (49-77%)	1 (0.47-1.9)	2% (0.37-4.3%)	2.1% (0.32-4.8%)	1.7 (0.25-3.9)	14 (2.6-30)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	1990	Male	90 to 94	35% (22-51%)	65% (49-78%)	0.91 (0.39-1.7)	1.3% (0.11-3.2%)	1.4% (-0.013-3.6%)	0.43 (-0.0039-1.1)	2.4 (0.21-5.6)
Botswana	1990	Male	95 plus	37% (21-56%)	63% (44-79%)	0.85 (0.29-1.7)	1.2% (-0.034-3.4%)	1.1% (-0.28-3.6%)	0.076 (-0.018-0.24)	0.31 (-0.009-0.9)
Botswana	1995	Female	All Ages	17% (13-21%)	83% (79-87%)	0.9 (0.7-1.1)	1.8% (0.73-3.3%)	2.6% (0.91-5.1%)	180 (48-430)	7200 (2700-15000)
Botswana	1995	Female	15 to 19	14% (7.8-23%)	86% (77-92%)	0.68 (0.3-1.3)	1.5% (0.7-2.9%)	1.8% (0.48-4%)	2.4 (0.56-6.2)	270 (120-550)
Botswana	1995	Female	20 to 24	19% (10-31%)	81% (69-90%)	1.2 (0.6-2.1)	1.6% (0.65-3.2%)	1.3% (0.3-3.3%)	4.8 (1.1-12)	500 (220-1000)
Botswana	1995	Female	25 to 29	22% (12-36%)	78% (64-88%)	1.6 (0.79-2.8)	1.3% (0.47-2.9%)	1.2% (0.23-3.1%)	6.6 (1.5-18)	560 (230-1300)
Botswana	1995	Female	30 to 34	25% (13-40%)	75% (60-87%)	1.9 (0.97-3.1)	1.6% (0.55-3.4%)	1.5% (0.35-3.8%)	8.6 (1.9-24)	610 (220-1500)
Botswana	1995	Female	35 to 39	28% (16-44%)	72% (56-84%)	1.9 (0.98-3.3)	2.2% (0.67-4.9%)	2.3% (0.47-5.6%)	12 (2.5-34)	720 (230-1800)
Botswana	1995	Female	40 to 44	29% (17-47%)	71% (53-83%)	2 (0.93-3.4)	3% (0.98-6.5%)	3.3% (0.82-7.5%)	15 (3.4-42)	790 (230-2000)
Botswana	1995	Female	45 to 49	30% (15-47%)	70% (53-85%)	2 (0.89-3.8)	3.7% (1.2-7.7%)	4.2% (1.1-9.2%)	15 (3.1-38)	680 (180-1600)
Botswana	1995	Female	50 to 54	29% (16-48%)	71% (52-84%)	1.9 (0.87-3.9)	4.5% (1.5-8.8%)	5.2% (1.6-11%)	18 (3.8-44)	690 (180-1600)
Botswana	1995	Female	55 to 59	29% (17-46%)	71% (54-83%)	1.9 (0.85-3.6)	4.9% (1.8-9.5%)	5.9% (2-11%)	20 (4.4-50)	680 (180-1600)
Botswana	1995	Female	60 to 64	30% (16-48%)	70% (52-84%)	1.9 (0.77-3.6)	5% (1.5-10%)	6.1% (1.8-12%)	21 (4.3-55)	600 (140-1500)
Botswana	1995	Female	65 to 69	30% (17-47%)	70% (53-83%)	1.7 (0.79-3.1)	4.8% (1.4-10%)	5.9% (1.8-12%)	21 (4.3-53)	480 (110-1200)
Botswana	1995	Female	70 to 74	28% (17-44%)	72% (56-83%)	1.4 (0.64-2.6)	3.8% (0.68-8.4%)	4.6% (0.87-10%)	18 (2.4-51)	340 (49-920)
Botswana	1995	Female	75 to 79	25% (14-41%)	75% (59-86%)	1.2 (0.46-2.3)	2.8% (0.0081-7%)	3.3% (-0.025-8.4%)	11 (-0.077-33)	150 (0.035-470)
Botswana	1995	Female	80 to 84	24% (13-39%)	76% (61-87%)	1.1 (0.43-2.1)	2.3% (-0.0067-6.1%)	2.7% (-0.023-7.4%)	6.1 (-0.059-19)	65 (-0.25-200)
Botswana	1995	Female	85 to 89	25% (14-40%)	75% (60-86%)	1 (0.43-2)	1.9% (-0.15-5.2%)	2.2% (-0.26-6.1%)	3 (-0.31-9.4)	23 (-2-69)
Botswana	1995	Female	90 to 94	28% (16-44%)	72% (56-84%)	1.1 (0.44-2.1)	1.6% (-0.54-5.1%)	1.9% (-0.71-6.1%)	1.2 (-0.45-4.4)	6.4 (-2-22)
Botswana	1995	Female	95 plus	33% (17-54%)	67% (46-83%)	1.2 (0.4-2.6)	1.6% (-0.98-6.3%)	1.8% (-1.3-7.1%)	0.33 (-0.25-1.5)	1.2 (-0.76-4.9)
Botswana	1995	Male	All Ages	32% (26-39%)	68% (61-74%)	1.9 (1.5-2.4)	5% (2.5-8%)	6.4% (3-10%)	570 (210-1100)	26000 (11000-46000)
Botswana	1995	Male	15 to 19	26% (17-40%)	74% (60-83%)	1.4 (0.68-2.4)	5.4% (2.9-8.8%)	6.7% (3.3-12%)	12 (4.3-25)	1100 (480-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	1995	Male	20 to 24	40% (26-56%)	60% (44-74%)	2.9 (1.6-4.6)	7.4% (3.6-12%)	7.8% (3.1-14%)	35 (12-65)	2700 (1200-4600)
Botswana	1995	Male	25 to 29	46% (29-64%)	54% (36-71%)	4 (2.4-6.1)	5.7% (2.4-10%)	5.8% (2-11%)	43 (15-80)	3000 (1300-5100)
Botswana	1995	Male	30 to 34	49% (31-67%)	51% (33-69%)	4.4 (2.7-6.6)	5.5% (2.3-9.8%)	5.6% (1.9-11%)	47 (16-90)	2900 (1200-5200)
Botswana	1995	Male	35 to 39	51% (35-69%)	49% (31-65%)	4.3 (2.5-6.7)	6.2% (2.4-11%)	6.5% (2.2-12%)	54 (17-110)	3000 (1100-5700)
Botswana	1995	Male	40 to 44	51% (35-69%)	49% (31-65%)	4.3 (2.3-6.8)	7.3% (2.9-13%)	7.8% (2.8-15%)	63 (19-140)	3000 (1100-6300)
Botswana	1995	Male	45 to 49	51% (33-68%)	49% (32-67%)	4.2 (2.3-7)	9.5% (4.2-16%)	10% (4.2-18%)	61 (19-130)	2600 (940-5400)
Botswana	1995	Male	50 to 54	51% (33-69%)	49% (31-67%)	4.1 (2.2-7)	12% (6-18%)	13% (6.5-20%)	65 (20-130)	2500 (890-4900)
Botswana	1995	Male	55 to 59	51% (34-69%)	49% (31-66%)	3.9 (2-6.5)	12% (6.5-18%)	13% (7.2-20%)	63 (21-130)	2100 (780-4000)
Botswana	1995	Male	60 to 64	49% (32-68%)	51% (32-68%)	3.3 (1.6-5.9)	10% (5.2-16%)	12% (5.9-18%)	51 (17-100)	1400 (520-2800)
Botswana	1995	Male	65 to 69	45% (31-62%)	55% (38-69%)	2.5 (1.3-4.3)	8.2% (4-14%)	9.4% (4.7-15%)	36 (12-71)	820 (300-1600)
Botswana	1995	Male	70 to 74	42% (27-59%)	58% (41-73%)	1.8 (0.88-3.1)	5.2% (2-9.4%)	5.9% (2.3-11%)	24 (7.6-49)	440 (150-890)
Botswana	1995	Male	75 to 79	41% (27-58%)	59% (42-73%)	1.5 (0.65-2.8)	3.6% (1-7.5%)	4% (1-8.5%)	12 (2.8-26)	170 (45-370)
Botswana	1995	Male	80 to 84	40% (24-57%)	60% (43-76%)	1.3 (0.58-2.3)	2.9% (0.73-6.2%)	3.2% (0.75-7%)	5.3 (1.2-12)	58 (14-130)
Botswana	1995	Male	85 to 89	37% (24-53%)	63% (47-76%)	1.1 (0.49-1.9)	2.1% (0.39-4.6%)	2.2% (0.31-5.1%)	1.9 (0.26-4.6)	15 (2.9-35)
Botswana	1995	Male	90 to 94	36% (23-51%)	64% (49-77%)	0.94 (0.4-1.8)	1.4% (0.18-3.6%)	1.5% (0.081-4%)	0.54 (0.03-1.4)	3 (0.36-7.6)
Botswana	1995	Male	95 plus	38% (21-57%)	62% (43-79%)	0.87 (0.3-1.8)	1.3% (0.038-3.7%)	1.3% (-0.17-4%)	0.11 (-0.014-0.36)	0.46 (0.013-1.3)
Botswana	2000	Female	All Ages	17% (13-22%)	83% (78-87%)	1.1 (0.87-1.4)	1.8% (0.83-3.8%)	2.6% (1-5.4%)	360 (120-950)	13000 (5500-33000)
Botswana	2000	Female	15 to 19	14% (8.1-24%)	86% (76-92%)	0.81 (0.35-1.5)	2.1% (0.92-4.2%)	2.6% (0.78-5.9%)	4.8 (1.4-12)	470 (190-990)
Botswana	2000	Female	20 to 24	19% (10-32%)	81% (68-90%)	1.4 (0.69-2.6)	1.6% (0.63-3.5%)	1.4% (0.4-3.7%)	10 (2.8-29)	900 (370-2100)
Botswana	2000	Female	25 to 29	22% (12-35%)	78% (65-88%)	1.9 (0.93-3.2)	1.1% (0.36-2.8%)	0.95% (0.23-2.9%)	14 (3.6-44)	1000 (390-2800)
Botswana	2000	Female	30 to 34	25% (13-40%)	75% (60-87%)	2.2 (1.1-3.7)	1.1% (0.38-2.9%)	1.1% (0.28-3.1%)	17 (4.5-53)	1100 (390-3000)
Botswana	2000	Female	35 to 39	28% (15-44%)	72% (56-85%)	2.3 (1.2-4)	1.7% (0.52-4.5%)	1.7% (0.43-4.9%)	24 (6.1-77)	1300 (420-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	2000	Female	40 to 44	29% (17-46%)	71% (54-83%)	2.4 (1.1-4.2)	2.2% (0.77-5.6%)	2.3% (0.68-6.1%)	30 (7.9-88)	1500 (440-4000)
Botswana	2000	Female	45 to 49	30% (16-46%)	70% (54-84%)	2.4 (1-4.5)	3% (1.1-7%)	3.2% (1-7.6%)	35 (10-100)	1500 (490-4200)
Botswana	2000	Female	50 to 54	30% (16-49%)	70% (51-84%)	2.3 (1-4.7)	3.9% (1.5-8.2%)	4.2% (1.5-9%)	31 (9.5-89)	1200 (390-3200)
Botswana	2000	Female	55 to 59	30% (17-46%)	70% (54-83%)	2.3 (0.99-4.3)	5% (1.9-10%)	5.6% (2-11%)	36 (9.8-99)	1200 (350-3200)
Botswana	2000	Female	60 to 64	30% (17-48%)	70% (52-83%)	2.2 (0.89-4.4)	5.5% (1.8-11%)	6.3% (2.1-12%)	40 (10-110)	1100 (290-3000)
Botswana	2000	Female	65 to 69	31% (17-47%)	69% (53-83%)	2 (0.94-3.8)	5.8% (2-11%)	6.7% (2.3-13%)	42 (10-110)	960 (230-2400)
Botswana	2000	Female	70 to 74	29% (17-45%)	71% (55-83%)	1.7 (0.71-3.2)	4.9% (1.2-10%)	5.7% (1.5-12%)	32 (6.1-88)	590 (120-1600)
Botswana	2000	Female	75 to 79	26% (14-41%)	74% (59-86%)	1.4 (0.52-2.8)	3.9% (0.39-9.1%)	4.5% (0.46-10%)	25 (2.1-72)	350 (31-1000)
Botswana	2000	Female	80 to 84	25% (13-40%)	75% (60-87%)	1.3 (0.52-2.5)	3.5% (0.36-8.5%)	4% (0.41-9.9%)	14 (1.3-41)	150 (15-430)
Botswana	2000	Female	85 to 89	25% (14-41%)	75% (59-86%)	1.2 (0.51-2.4)	3% (0.27-7.4%)	3.5% (0.26-8.6%)	6.4 (0.41-17)	48 (3.4-130)
Botswana	2000	Female	90 to 94	28% (16-44%)	72% (56-84%)	1.3 (0.5-2.6)	2.7% (-0.14-7.4%)	3.1% (-0.24-8.7%)	2.3 (-0.14-6.4)	12 (-0.55-32)
Botswana	2000	Female	95 plus	33% (17-52%)	67% (48-83%)	1.4 (0.47-3.1)	2.8% (-0.55-8.7%)	3.1% (-0.75-9.9%)	0.74 (-0.19-2.4)	2.6 (-0.54-8)
Botswana	2000	Male	All Ages	33% (27-40%)	67% (60-73%)	2.3 (1.8-2.9)	5.2% (2.8-8.8%)	6.4% (3.3-11%)	1000 (430-2100)	44000 (21000-86000)
Botswana	2000	Male	15 to 19	27% (17-41%)	73% (59-83%)	1.6 (0.78-2.7)	7% (3.8-12%)	8.9% (4.3-15%)	21 (8.2-42)	1800 (820-3300)
Botswana	2000	Male	20 to 24	40% (26-56%)	60% (44-74%)	3.3 (1.8-5.3)	8.9% (4.6-14%)	9.6% (4.3-16%)	62 (26-120)	4600 (2200-8200)
Botswana	2000	Male	25 to 29	47% (29-64%)	53% (36-71%)	4.5 (2.7-6.6)	5.3% (2.3-9.9%)	5.3% (2.1-11%)	75 (32-140)	5000 (2400-9100)
Botswana	2000	Male	30 to 34	50% (31-68%)	50% (32-69%)	5 (3.1-7.5)	4.4% (2-8.2%)	4.5% (1.8-8.7%)	80 (32-160)	4800 (2100-9200)
Botswana	2000	Male	35 to 39	51% (35-69%)	49% (31-65%)	5 (2.9-7.5)	5.1% (2.1-10%)	5.3% (2-11%)	91 (33-190)	4800 (1900-10000)
Botswana	2000	Male	40 to 44	51% (35-70%)	49% (30-65%)	4.9 (2.6-7.8)	6% (2.5-12%)	6.2% (2.4-12%)	110 (38-240)	5100 (1900-11000)
Botswana	2000	Male	45 to 49	51% (33-69%)	49% (31-67%)	4.8 (2.5-7.9)	8% (3.5-15%)	8.4% (3.4-16%)	130 (45-300)	5400 (2000-12000)
Botswana	2000	Male	50 to 54	51% (33-69%)	49% (31-67%)	4.7 (2.4-7.6)	10% (5.4-17%)	11% (5.5-19%)	110 (42-250)	4200 (1700-8900)
Botswana	2000	Male	55 to 59	51% (34-69%)	49% (31-66%)	4.4 (2.3-7.4)	12% (6.7-19%)	13% (7.3-20%)	110 (43-230)	3500 (1500-7300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	2000	Male	60 to 64	49% (33-68%)	51% (32-67%)	3.8 (1.8-6.6)	11% (5.6-17%)	12% (6.2-18%)	91 (35-190)	2500 (1000-5100)
Botswana	2000	Male	65 to 69	46% (31-63%)	54% (37-69%)	2.9 (1.5-4.9)	9.3% (4.6-15%)	10% (5.1-17%)	66 (27-140)	1500 (630-3000)
Botswana	2000	Male	70 to 74	43% (28-60%)	57% (40-72%)	2.1 (1-3.6)	6.2% (2.6-11%)	6.9% (2.9-12%)	35 (12-72)	640 (230-1300)
Botswana	2000	Male	75 to 79	42% (28-59%)	58% (41-72%)	1.7 (0.76-3.1)	4.5% (1.4-8.9%)	5% (1.5-10%)	23 (6.3-49)	320 (92-690)
Botswana	2000	Male	80 to 84	40% (25-57%)	60% (43-75%)	1.5 (0.68-2.6)	3.8% (1.1-7.5%)	4.2% (1.1-8.4%)	10 (2.8-22)	110 (32-230)
Botswana	2000	Male	85 to 89	37% (24-53%)	63% (47-76%)	1.2 (0.58-2.1)	2.7% (0.68-5.6%)	3% (0.68-6.3%)	3.1 (0.71-6.8)	24 (6.2-51)
Botswana	2000	Male	90 to 94	36% (23-52%)	64% (48-77%)	1.1 (0.47-2)	1.9% (0.39-4.5%)	2.1% (0.34-4.9%)	0.78 (0.12-1.8)	4.3 (0.83-9.8)
Botswana	2000	Male	95 plus	38% (21-57%)	62% (43-79%)	0.99 (0.36-2)	1.8% (0.16-4.4%)	1.8% (-0.019-5%)	0.19 (-0.0019-0.51)	0.72 (0.066-1.8)
Botswana	2005	Female	All Ages	17% (14-22%)	83% (78-86%)	1.3 (0.99-1.7)	1.9% (0.99-3.8%)	2.8% (1.3-5.4%)	420 (170-1000)	15000 (7200-34000)
Botswana	2005	Female	15 to 19	14% (7.9-23%)	86% (77-92%)	0.9 (0.39-1.7)	2.6% (1.2-4.9%)	3.6% (1.3-7.7%)	5.2 (1.8-13)	520 (220-1100)
Botswana	2005	Female	20 to 24	19% (10-31%)	81% (69-90%)	1.6 (0.78-2.9)	2.3% (0.88-4.6%)	2.1% (0.6-5.1%)	11 (3.7-31)	1000 (480-2300)
Botswana	2005	Female	25 to 29	22% (12-36%)	78% (64-88%)	2.1 (1-3.8)	1.2% (0.43-3%)	1.1% (0.3-3.2%)	16 (5.1-47)	1200 (510-3000)
Botswana	2005	Female	30 to 34	24% (13-40%)	76% (60-87%)	2.4 (1.2-4.2)	1% (0.4-2.5%)	0.95% (0.3-2.5%)	19 (6.1-52)	1200 (490-3100)
Botswana	2005	Female	35 to 39	27% (15-43%)	73% (57-85%)	2.6 (1.3-4.5)	1.4% (0.53-3.5%)	1.4% (0.45-3.8%)	24 (7.8-72)	1400 (500-3700)
Botswana	2005	Female	40 to 44	29% (17-46%)	71% (54-83%)	2.7 (1.1-4.9)	1.9% (0.76-4.4%)	1.9% (0.69-4.7%)	29 (9.4-81)	1400 (520-3800)
Botswana	2005	Female	45 to 49	30% (15-46%)	70% (54-85%)	2.7 (1.2-5)	2.7% (1.1-5.9%)	2.8% (1.1-6.3%)	36 (12-94)	1500 (580-3900)
Botswana	2005	Female	50 to 54	30% (16-48%)	70% (52-84%)	2.6 (1.2-5.1)	3.8% (1.6-7.5%)	4% (1.6-8.1%)	41 (14-110)	1600 (580-3900)
Botswana	2005	Female	55 to 59	29% (17-47%)	71% (53-83%)	2.6 (1.1-4.8)	5.2% (2.2-9.8%)	5.8% (2.4-11%)	38 (13-99)	1200 (450-3200)
Botswana	2005	Female	60 to 64	30% (16-47%)	70% (53-84%)	2.5 (1-4.9)	6% (2.3-11%)	7% (2.7-13%)	42 (13-110)	1200 (380-3000)
Botswana	2005	Female	65 to 69	30% (17-47%)	70% (53-83%)	2.3 (1-4.2)	6.6% (2.6-12%)	7.7% (3.1-14%)	50 (16-120)	1100 (370-2800)
Botswana	2005	Female	70 to 74	28% (16-43%)	72% (57-84%)	1.9 (0.79-3.5)	5.6% (1.7-11%)	6.6% (2-13%)	42 (11-110)	780 (200-2000)
Botswana	2005	Female	75 to 79	25% (14-41%)	75% (59-86%)	1.5 (0.57-3.1)	4.6% (0.81-10%)	5.4% (0.98-12%)	29 (4.3-78)	420 (64-1100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	2005	Female	80 to 84	24% (13-39%)	76% (61-87%)	1.4 (0.57-2.7)	4.3% (0.8-9.6%)	5% (0.9-11%)	21 (3.2-54)	220 (36-560)
Botswana	2005	Female	85 to 89	25% (14-40%)	75% (60-86%)	1.4 (0.59-2.6)	3.9% (0.56-8.9%)	4.5% (0.64-10%)	11 (1.4-27)	83 (11-200)
Botswana	2005	Female	90 to 94	28% (15-44%)	72% (56-85%)	1.5 (0.58-2.9)	3.6% (0.18-8.5%)	4.1% (0.14-10%)	3.6 (0.11-9.4)	19 (1-47)
Botswana	2005	Female	95 plus	32% (17-54%)	68% (46-83%)	1.6 (0.49-3.6)	3.7% (-0.28-9.8%)	4.2% (-0.42-11%)	1 (-0.093-2.9)	3.6 (-0.25-9.9)
Botswana	2005	Male	All Ages	33% (27-41%)	67% (59-73%)	2.8 (2.3-3.5)	6.2% (3.6-9.2%)	7.5% (4.4-11%)	1200 (610-2100)	53000 (28000-85000)
Botswana	2005	Male	15 to 19	27% (17-41%)	73% (59-83%)	1.8 (0.92-3.3)	8.2% (4.5-13%)	11% (5.5-18%)	25 (12-44)	2000 (1100-3400)
Botswana	2005	Male	20 to 24	40% (25-56%)	60% (44-75%)	3.9 (2.1-6.1)	12% (7.2-18%)	14% (7.2-22%)	76 (39-120)	5600 (3200-8800)
Botswana	2005	Male	25 to 29	46% (30-62%)	54% (38-70%)	5.3 (3.1-7.9)	7.1% (3.6-12%)	7.4% (3.4-13%)	95 (48-160)	6300 (3400-9900)
Botswana	2005	Male	30 to 34	49% (32-67%)	51% (33-68%)	5.9 (3.5-8.6)	4.9% (2.5-8%)	5% (2.3-8.4%)	96 (44-160)	5700 (2900-9400)
Botswana	2005	Male	35 to 39	51% (34-68%)	49% (32-66%)	5.8 (3.3-9)	5.2% (2.4-8.8%)	5.4% (2.3-9.3%)	100 (43-180)	5400 (2400-9500)
Botswana	2005	Male	40 to 44	52% (35-69%)	48% (31-65%)	5.7 (2.9-9.3)	6.1% (2.9-10%)	6.3% (2.8-11%)	110 (45-210)	5400 (2300-9700)
Botswana	2005	Male	45 to 49	51% (33-69%)	49% (31-67%)	5.7 (3-9.6)	8% (3.9-13%)	8.4% (3.9-14%)	140 (56-260)	5800 (2500-11000)
Botswana	2005	Male	50 to 54	51% (33-69%)	49% (31-67%)	5.5 (2.9-9.2)	11% (6.1-17%)	12% (6.3-18%)	150 (64-270)	5600 (2500-10000)
Botswana	2005	Male	55 to 59	51% (34-69%)	49% (31-66%)	5.1 (2.6-8.8)	13% (7.8-19%)	15% (8.6-21%)	130 (59-230)	4100 (2000-7200)
Botswana	2005	Male	60 to 64	49% (33-67%)	51% (33-67%)	4.4 (2.1-7.6)	13% (7-19%)	14% (7.6-20%)	110 (50-200)	3100 (1400-5300)
Botswana	2005	Male	65 to 69	46% (30-63%)	54% (37-70%)	3.3 (1.7-5.6)	11% (6-17%)	13% (6.6-19%)	91 (42-160)	2100 (980-3500)
Botswana	2005	Male	70 to 74	42% (27-59%)	58% (41-73%)	2.4 (1.2-4.2)	7.8% (3.2-13%)	8.7% (3.6-14%)	52 (20-96)	960 (390-1700)
Botswana	2005	Male	75 to 79	41% (27-58%)	59% (42-73%)	2 (0.84-3.7)	5.7% (1.8-10%)	6.4% (1.9-12%)	28 (7.9-54)	400 (120-770)
Botswana	2005	Male	80 to 84	40% (24-57%)	60% (43-76%)	1.7 (0.75-3.1)	4.8% (1.5-9.5%)	5.4% (1.6-11%)	16 (4.6-33)	170 (52-340)
Botswana	2005	Male	85 to 89	37% (24-53%)	63% (47-76%)	1.4 (0.64-2.6)	3.7% (1-7.3%)	4.1% (1.1-8.3%)	6 (1.6-12)	46 (13-94)
Botswana	2005	Male	90 to 94	36% (23-52%)	64% (48-77%)	1.3 (0.54-2.5)	2.8% (0.63-6.2%)	3.1% (0.63-7%)	1.4 (0.26-3.1)	7.3 (1.7-16)
Botswana	2005	Male	95 plus	38% (21-58%)	62% (42-79%)	1.2 (0.42-2.5)	2.7% (0.44-6.8%)	2.9% (0.32-7.5%)	0.31 (0.033-0.82)	1.2 (0.19-2.9)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	2010	Female	All Ages	17% (14-22%)	83% (78-86%)	1.4 (0.99-1.9)	2.3% (1.1-4%)	3.3% (1.4-5.9%)	340 (110-750)	12000 (5500-26000)
Botswana	2010	Female	15 to 19	14% (7.6-23%)	86% (77-92%)	0.91 (0.38-1.8)	2.2% (1.1-4.1%)	2.9% (1-5.8%)	3.6 (1-8.6)	400 (180-790)
Botswana	2010	Female	20 to 24	19% (9.8-31%)	81% (69-90%)	1.6 (0.75-3.1)	2.7% (1.3-5.1%)	2.6% (0.72-6.5%)	7.7 (2.3-19)	840 (400-1700)
Botswana	2010	Female	25 to 29	21% (11-35%)	79% (65-89%)	2.1 (0.97-3.8)	1.8% (0.67-3.9%)	1.6% (0.38-4.4%)	12 (3.4-33)	1000 (460-2300)
Botswana	2010	Female	30 to 34	24% (13-38%)	76% (62-87%)	2.4 (1.2-4.3)	1.4% (0.55-2.9%)	1.2% (0.34-3.1%)	15 (4.3-41)	1100 (430-2500)
Botswana	2010	Female	35 to 39	27% (15-42%)	73% (58-85%)	2.6 (1.3-5)	1.6% (0.59-3.8%)	1.6% (0.43-4.1%)	19 (4.9-53)	1100 (400-2800)
Botswana	2010	Female	40 to 44	28% (16-45%)	72% (55-84%)	2.7 (1.1-5)	2.1% (0.8-4.7%)	2.1% (0.69-5.1%)	22 (6.2-58)	1100 (390-2800)
Botswana	2010	Female	45 to 49	29% (15-46%)	71% (54-85%)	2.7 (1.1-5.5)	2.9% (1.1-6.5%)	3.2% (1-7.2%)	26 (7.5-65)	1200 (400-2800)
Botswana	2010	Female	50 to 54	29% (16-46%)	71% (54-84%)	2.7 (1.1-5.5)	4% (1.6-8.1%)	4.5% (1.6-9.4%)	31 (8.6-77)	1200 (410-2900)
Botswana	2010	Female	55 to 59	28% (16-45%)	72% (55-84%)	2.6 (1.1-4.9)	5.5% (2.2-10%)	6.6% (2.5-12%)	36 (9.4-89)	1200 (370-2900)
Botswana	2010	Female	60 to 64	29% (16-46%)	71% (54-84%)	2.5 (1-5)	6.1% (2.3-11%)	7.6% (2.9-14%)	32 (9.2-76)	910 (280-2100)
Botswana	2010	Female	65 to 69	29% (16-45%)	71% (55-84%)	2.3 (0.97-4.5)	6.6% (2.4-12%)	8.2% (3.2-15%)	37 (10-88)	880 (260-2000)
Botswana	2010	Female	70 to 74	27% (16-42%)	73% (58-84%)	1.9 (0.73-3.8)	5.4% (1.4-11%)	6.7% (1.7-13%)	34 (6.9-84)	650 (140-1500)
Botswana	2010	Female	75 to 79	25% (13-40%)	75% (60-87%)	1.6 (0.55-3.2)	4.3% (0.46-9.5%)	5.3% (0.59-12%)	26 (2.4-69)	380 (37-980)
Botswana	2010	Female	80 to 84	24% (13-39%)	76% (61-87%)	1.4 (0.54-2.8)	4.1% (0.51-9.1%)	4.9% (0.56-11%)	20 (2.3-52)	210 (26-550)
Botswana	2010	Female	85 to 89	25% (13-40%)	75% (60-87%)	1.4 (0.53-2.9)	3.8% (0.43-8.9%)	4.5% (0.47-10%)	11 (0.95-27)	82 (8.2-210)
Botswana	2010	Female	90 to 94	27% (15-43%)	73% (57-85%)	1.5 (0.56-3)	3.5% (0.11-8.7%)	4.2% (0.058-10%)	4.3 (0.06-12)	22 (0.73-59)
Botswana	2010	Female	95 plus	32% (16-52%)	68% (48-84%)	1.6 (0.5-3.8)	3.6% (-0.36-9.7%)	4.2% (-0.59-11%)	1.1 (-0.16-3.3)	3.8 (-0.41-11)
Botswana	2010	Male	All Ages	34% (28-42%)	66% (58-72%)	3 (2.2-4.1)	7.5% (4.7-10%)	8.9% (5.6-12%)	1200 (640-1700)	50000 (28000-73000)
Botswana	2010	Male	15 to 19	26% (16-41%)	74% (59-84%)	1.9 (0.89-3.4)	7.2% (4-11%)	9.2% (4.7-15%)	21 (9.7-36)	1700 (930-2900)
Botswana	2010	Male	20 to 24	39% (25-56%)	61% (44-75%)	3.9 (2-6.6)	13% (8.3-19%)	15% (8.7-23%)	64 (34-95)	4900 (2900-7000)
Botswana	2010	Male	25 to 29	46% (29-63%)	54% (37-71%)	5.3 (3-8.2)	10% (5.7-15%)	11% (5.5-18%)	87 (46-130)	6000 (3400-8500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	2010	Male	30 to 34	49% (32-66%)	51% (34-68%)	6 (3.5-9.3)	6.9% (3.7-11%)	7.2% (3.5-12%)	94 (45-150)	5800 (3200-8700)
Botswana	2010	Male	35 to 39	51% (34-68%)	49% (32-66%)	5.9 (3.3-9.5)	6.3% (3.2-9.7%)	6.6% (3-11%)	97 (44-160)	5300 (2600-8400)
Botswana	2010	Male	40 to 44	51% (34-69%)	49% (31-66%)	5.8 (2.9-9.6)	6.9% (3.6-11%)	7.3% (3.6-12%)	100 (46-180)	5000 (2400-8300)
Botswana	2010	Male	45 to 49	50% (32-68%)	50% (32-68%)	5.7 (2.8-9.8)	9% (4.7-14%)	9.7% (4.8-15%)	120 (53-200)	5000 (2400-8300)
Botswana	2010	Male	50 to 54	50% (31-68%)	50% (32-69%)	5.5 (2.7-9.4)	11% (6.5-17%)	12% (6.8-18%)	130 (62-220)	5000 (2500-8100)
Botswana	2010	Male	55 to 59	50% (33-68%)	50% (32-67%)	5.2 (2.5-8.9)	14% (8.5-19%)	16% (9.6-22%)	140 (67-230)	4500 (2300-7200)
Botswana	2010	Male	60 to 64	48% (32-67%)	52% (33-68%)	4.4 (2-8)	13% (7.6-19%)	15% (8.6-21%)	100 (50-170)	2900 (1400-4700)
Botswana	2010	Male	65 to 69	45% (30-62%)	55% (38-70%)	3.4 (1.6-5.8)	11% (5.8-17%)	13% (6.8-19%)	88 (39-140)	2000 (920-3300)
Botswana	2010	Male	70 to 74	42% (27-58%)	58% (42-73%)	2.4 (1.1-4.4)	7.8% (3.1-13%)	8.8% (3.5-15%)	55 (21-97)	1000 (400-1800)
Botswana	2010	Male	75 to 79	40% (26-57%)	60% (43-74%)	1.9 (0.81-3.7)	5.6% (1.7-11%)	6.3% (1.8-12%)	31 (8.6-59)	440 (130-840)
Botswana	2010	Male	80 to 84	39% (24-57%)	61% (43-76%)	1.7 (0.72-3.1)	4.6% (1.3-9.1%)	5.2% (1.4-10%)	17 (4.4-33)	180 (51-350)
Botswana	2010	Male	85 to 89	36% (23-52%)	64% (48-77%)	1.4 (0.62-2.7)	3.5% (0.92-7.4%)	4% (0.97-8.4%)	6.1 (1.5-13)	47 (12-98)
Botswana	2010	Male	90 to 94	35% (22-51%)	65% (49-78%)	1.3 (0.53-2.4)	2.7% (0.5-6.1%)	3.1% (0.46-7%)	1.7 (0.26-3.9)	9 (1.6-20)
Botswana	2010	Male	95 plus	37% (21-57%)	63% (43-79%)	1.2 (0.41-2.5)	2.6% (0.41-6.5%)	2.8% (0.26-7.3%)	0.34 (0.031-0.86)	1.2 (0.19-3)
Botswana	2016	Female	All Ages	18% (14-22%)	82% (78-86%)	1.4 (0.81-2.3)	3.6% (1.5-5.9%)	5.2% (1.8-9%)	380 (89-760)	14000 (4900-26000)
Botswana	2016	Female	15 to 19	14% (7.6-23%)	86% (77-92%)	0.92 (0.34-2)	2.3% (1.1-4.2%)	3.3% (1.1-6.7%)	3.3 (0.82-7.3)	380 (160-700)
Botswana	2016	Female	20 to 24	18% (9.8-30%)	82% (70-90%)	1.7 (0.64-3.4)	3.3% (1.7-5.6%)	3.5% (1.1-7%)	6.9 (1.8-15)	780 (380-1300)
Botswana	2016	Female	25 to 29	21% (11-34%)	79% (66-89%)	2.1 (0.89-4.1)	3.1% (1.4-5.5%)	3.4% (0.77-7.1%)	12 (2.7-26)	1000 (460-1900)
Botswana	2016	Female	30 to 34	24% (12-39%)	76% (61-88%)	2.4 (1.1-4.8)	3% (1.3-5.6%)	3.4% (0.82-7%)	17 (3.3-36)	1200 (460-2400)
Botswana	2016	Female	35 to 39	27% (14-42%)	73% (58-86%)	2.6 (1.2-5.2)	3.4% (1.2-6.4%)	4% (0.89-8.1%)	24 (3.9-53)	1400 (430-2900)
Botswana	2016	Female	40 to 44	28% (16-44%)	72% (56-84%)	2.6 (1-5.3)	3.8% (1.2-7.3%)	4.6% (1.1-9.1%)	26 (4.7-58)	1400 (380-2800)
Botswana	2016	Female	45 to 49	28% (15-45%)	72% (55-85%)	2.6 (1-5.7)	4.6% (1.5-8.9%)	5.7% (1.5-11%)	30 (5.4-63)	1300 (340-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	2016	Female	50 to 54	28% (15-47%)	72% (53-85%)	2.6 (1-5.6)	5.4% (1.9-10%)	6.7% (2-13%)	33 (6.3-73)	1300 (330-2700)
Botswana	2016	Female	55 to 59	28% (16-45%)	72% (55-84%)	2.6 (1-5.4)	6.7% (2.4-12%)	8.9% (3.3-15%)	38 (7.3-86)	1300 (330-2800)
Botswana	2016	Female	60 to 64	29% (16-46%)	71% (54-84%)	2.5 (0.91-5.3)	6.8% (2.1-13%)	9% (3-16%)	42 (8-92)	1200 (280-2600)
Botswana	2016	Female	65 to 69	29% (17-46%)	71% (54-83%)	2.3 (0.87-4.6)	6.8% (2-13%)	8.9% (2.8-16%)	40 (7.3-90)	930 (200-2100)
Botswana	2016	Female	70 to 74	27% (16-43%)	73% (57-84%)	1.9 (0.69-3.9)	5.5% (1-11%)	7% (1.3-14%)	35 (4.5-82)	670 (94-1500)
Botswana	2016	Female	75 to 79	25% (14-41%)	75% (59-86%)	1.6 (0.51-3.5)	4.3% (0.36-9.4%)	5.3% (0.44-12%)	30 (2.1-74)	440 (30-1100)
Botswana	2016	Female	80 to 84	24% (12-39%)	76% (61-88%)	1.4 (0.47-3)	3.9% (0.11-9%)	4.7% (0.076-11%)	23 (0.42-57)	240 (7.5-600)
Botswana	2016	Female	85 to 89	25% (13-39%)	75% (61-87%)	1.4 (0.47-3.1)	3.6% (-0.098-8.8%)	4.3% (-0.18-10%)	14 (-0.56-36)	100 (-2.8-270)
Botswana	2016	Female	90 to 94	27% (15-43%)	73% (57-85%)	1.5 (0.47-3.3)	3.3% (-0.31-8.9%)	3.9% (-0.47-10%)	5 (-0.56-14)	26 (-2.3-71)
Botswana	2016	Female	95 plus	32% (16-51%)	68% (49-84%)	1.6 (0.43-3.8)	3.4% (-0.62-9.6%)	3.9% (-0.87-11%)	1.5 (-0.31-4.6)	5.1 (-0.89-15)
Botswana	2016	Male	All Ages	35% (29-43%)	65% (57-71%)	3.2 (1.9-5.2)	9.3% (5.4-13%)	11% (6.5-15%)	1100 (520-1800)	48000 (24000-71000)
Botswana	2016	Male	15 to 19	26% (16-40%)	74% (60-84%)	1.9 (0.75-3.8)	6.8% (3.5-11%)	9.1% (4.1-16%)	16 (6.9-29)	1400 (710-2400)
Botswana	2016	Male	20 to 24	39% (25-55%)	61% (45-75%)	4 (1.8-7.8)	13% (7.5-18%)	15% (8-22%)	49 (23-77)	3900 (2100-5700)
Botswana	2016	Male	25 to 29	45% (29-63%)	55% (37-71%)	5.4 (2.6-10)	12% (6.6-17%)	13% (6.8-20%)	69 (31-110)	4900 (2500-7300)
Botswana	2016	Male	30 to 34	48% (31-66%)	52% (34-69%)	6 (3-11)	10% (5.5-15%)	11% (5.4-17%)	85 (35-140)	5400 (2700-8300)
Botswana	2016	Male	35 to 39	50% (34-68%)	50% (32-66%)	5.9 (2.9-12)	9.3% (4.7-15%)	10% (4.6-17%)	98 (36-170)	5500 (2300-8900)
Botswana	2016	Male	40 to 44	51% (34-69%)	49% (31-66%)	5.9 (2.5-12)	9.6% (4.6-15%)	11% (4.6-17%)	110 (39-180)	5200 (2200-8700)
Botswana	2016	Male	45 to 49	50% (32-68%)	50% (32-68%)	5.8 (2.5-11)	11% (5.6-17%)	13% (5.8-20%)	120 (42-200)	5100 (2000-8400)
Botswana	2016	Male	50 to 54	50% (32-68%)	50% (32-68%)	5.6 (2.3-11)	13% (6.8-19%)	15% (7.4-22%)	120 (46-200)	4700 (2000-7600)
Botswana	2016	Male	55 to 59	50% (33-67%)	50% (33-67%)	5.3 (2.3-10)	15% (8.3-21%)	17% (9.4-24%)	130 (56-210)	4300 (2000-6900)
Botswana	2016	Male	60 to 64	48% (32-67%)	52% (33-68%)	4.5 (1.8-8.9)	13% (6.9-19%)	16% (8.3-23%)	120 (53-200)	3400 (1500-5500)
Botswana	2016	Male	65 to 69	45% (30-62%)	55% (38-70%)	3.4 (1.4-6.5)	11% (5.3-17%)	13% (6.2-20%)	90 (35-150)	2100 (830-3500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Botswana	2016	Male	70 to 74	42% (27-58%)	58% (42-73%)	2.5 (0.96-4.8)	7.8% (2.8-13%)	9% (3.2-15%)	57 (18-100)	1100 (350-1900)
Botswana	2016	Male	75 to 79	40% (26-58%)	60% (42-74%)	2 (0.71-4.2)	5.7% (1.3-11%)	6.5% (1.5-13%)	36 (7.8-73)	520 (120-1100)
Botswana	2016	Male	80 to 84	38% (24-55%)	62% (45-76%)	1.7 (0.66-3.7)	4.7% (1.9-7.7%)	5.3% (1.1-11%)	19 (3.8-42)	210 (45-450)
Botswana	2016	Male	85 to 89	35% (23-51%)	65% (49-77%)	1.5 (0.53-3)	3.6% (0.72-8.3%)	4.1% (0.75-9.5%)	7.6 (1.5-17)	59 (12-130)
Botswana	2016	Male	90 to 94	35% (22-50%)	65% (50-78%)	1.3 (0.45-2.8)	2.9% (0.41-6.9%)	3.2% (0.37-8%)	2 (0.22-4.9)	11 (1.5-25)
Botswana	2016	Male	95 plus	37% (20-56%)	63% (44-80%)	1.2 (0.35-3)	2.8% (0.31-7.4%)	3% (0.18-8.3%)	0.47 (0.028-1.3)	1.7 (0.19-4.6)
Lesotho	1990	Female	All Ages	16% (13-19%)	84% (81-87%)	1.1 (0.77-1.5)	3.5% (2.2-5%)	6% (3.4-9%)	400 (220-620)	13000 (8400-19000)
Lesotho	1990	Female	15 to 19	14% (8.2-22%)	86% (78-92%)	0.84 (0.37-1.7)	3.5% (1.6-6.3%)	5.8% (2.3-11%)	6.7 (2.5-13)	560 (260-1000)
Lesotho	1990	Female	20 to 24	18% (11-29%)	82% (71-89%)	1.5 (0.75-2.8)	5.7% (2.9-9.9%)	8.3% (3.5-15%)	12 (5-21)	940 (480-1600)
Lesotho	1990	Female	25 to 29	21% (12-33%)	79% (67-88%)	2 (0.96-3.4)	6.3% (3.4-10%)	9.4% (4.4-16%)	13 (6-21)	910 (490-1400)
Lesotho	1990	Female	30 to 34	24% (14-37%)	76% (63-86%)	2.3 (1.1-3.9)	6.9% (3.8-11%)	10% (5.3-18%)	15 (7.6-24)	940 (530-1500)
Lesotho	1990	Female	35 to 39	26% (15-40%)	74% (60-85%)	2.4 (1.1-4.2)	7.6% (4.4-12%)	12% (6.4-18%)	18 (10-29)	1000 (600-1600)
Lesotho	1990	Female	40 to 44	28% (17-43%)	72% (57-83%)	2.4 (1.1-4.3)	8.4% (4.5-13%)	13% (6.6-19%)	21 (11-34)	1000 (550-1600)
Lesotho	1990	Female	45 to 49	28% (16-42%)	72% (58-84%)	2.4 (1.1-4.6)	9% (4.7-14%)	13% (6.8-20%)	20 (11-33)	880 (470-1400)
Lesotho	1990	Female	50 to 54	27% (16-43%)	73% (57-84%)	2.3 (1.1-4.3)	9.3% (4.9-15%)	13% (6.6-20%)	33 (16-55)	1200 (610-2000)
Lesotho	1990	Female	55 to 59	27% (16-42%)	73% (58-84%)	2.3 (1-4.2)	9.2% (4.5-15%)	12% (6-20%)	46 (21-79)	1500 (710-2500)
Lesotho	1990	Female	60 to 64	28% (16-43%)	72% (57-84%)	2.2 (0.98-4.2)	8.7% (3.9-15%)	11% (4.9-19%)	47 (20-82)	1300 (570-2200)
Lesotho	1990	Female	65 to 69	28% (18-42%)	72% (58-82%)	2 (0.95-3.6)	8.4% (3.5-14%)	10% (4.4-17%)	59 (23-100)	1300 (530-2200)
Lesotho	1990	Female	70 to 74	26% (17-39%)	74% (61-83%)	1.6 (0.77-3)	6.3% (2.3-11%)	7.7% (2.7-13%)	48 (16-86)	860 (300-1600)
Lesotho	1990	Female	75 to 79	23% (14-36%)	77% (64-86%)	1.3 (0.55-2.5)	4.7% (1.3-9.1%)	5.6% (1.5-11%)	32 (8.4-64)	450 (120-880)
Lesotho	1990	Female	80 to 84	22% (13-34%)	78% (66-87%)	1.2 (0.55-2.2)	3.9% (1-7.8%)	4.7% (1.2-9.2%)	20 (5.3-42)	210 (55-430)
Lesotho	1990	Female	85 to 89	23% (13-34%)	77% (66-87%)	1.2 (0.5-2.2)	3.4% (0.72-7.3%)	4% (0.83-8.6%)	11 (2.2-23)	79 (16-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	1990	Female	90 to 94	24% (15-36%)	76% (64-85%)	1.2 (0.53-2.3)	2.9% (0.49-6.7%)	3.5% (0.55-7.9%)	3.6 (0.55-8.1)	18 (3-40)
Lesotho	1990	Female	95 plus	28% (16-43%)	72% (57-84%)	1.3 (0.47-2.7)	2.8% (0.078-7.2%)	3.2% (0.002-8.5%)	0.92 (0.00059-2.4)	3.1 (0.087-8)
Lesotho	1990	Male	All Ages	36% (31-40%)	64% (60-69%)	2.8 (2-3.6)	8.8% (6.7-12%)	13% (9.4-17%)	1100 (750-1400)	41000 (30000-53000)
Lesotho	1990	Male	15 to 19	30% (22-39%)	70% (61-78%)	2 (1-3.3)	8.9% (5.2-14%)	12% (6.6-19%)	25 (13-41)	2000 (1100-3000)
Lesotho	1990	Male	20 to 24	48% (37-59%)	52% (41-63%)	4.6 (2.6-7.3)	18% (12-25%)	22% (15-32%)	65 (42-92)	4600 (3100-6400)
Lesotho	1990	Male	25 to 29	56% (45-66%)	44% (34-55%)	6.3 (3.6-9.7)	20% (15-27%)	25% (18-33%)	64 (46-87)	4100 (3000-5500)
Lesotho	1990	Male	30 to 34	61% (49-69%)	39% (31-51%)	7 (4.3-10)	21% (15-27%)	25% (18-34%)	57 (41-77)	3300 (2400-4400)
Lesotho	1990	Male	35 to 39	62% (50-71%)	38% (29-50%)	6.9 (4-11)	21% (15-28%)	26% (19-35%)	63 (44-85)	3300 (2300-4400)
Lesotho	1990	Male	40 to 44	64% (53-72%)	36% (28-47%)	6.9 (3.8-11)	23% (17-29%)	28% (21-36%)	73 (51-97)	3400 (2400-4500)
Lesotho	1990	Male	45 to 49	66% (54-74%)	34% (26-46%)	7.1 (3.9-12)	24% (18-30%)	29% (22-37%)	81 (57-110)	3400 (2400-4500)
Lesotho	1990	Male	50 to 54	66% (56-74%)	34% (26-44%)	6.9 (3.7-11)	24% (17-30%)	28% (20-37%)	110 (76-150)	4100 (2800-5600)
Lesotho	1990	Male	55 to 59	64% (52-72%)	36% (28-48%)	6.3 (3.5-9.9)	22% (16-29%)	26% (18-34%)	130 (88-180)	4200 (2800-5800)
Lesotho	1990	Male	60 to 64	60% (47-70%)	40% (30-53%)	5.2 (2.7-8.7)	19% (12-25%)	19% (13-29%)	130 (75-180)	3400 (2000-4900)
Lesotho	1990	Male	65 to 69	54% (37-70%)	46% (30-63%)	3.8 (1.9-6.4)	15% (8.4-22%)	17% (9.7-25%)	110 (61-170)	2600 (1400-3800)
Lesotho	1990	Male	70 to 74	50% (33-67%)	50% (33-67%)	2.8 (1.4-4.9)	9.9% (4.7-16%)	11% (5.3-18%)	71 (32-120)	1300 (580-2100)
Lesotho	1990	Male	75 to 79	49% (32-66%)	51% (34-68%)	2.2 (1-4.1)	7.1% (2.7-13%)	8% (3-14%)	44 (16-83)	620 (230-1100)
Lesotho	1990	Male	80 to 84	47% (30-64%)	53% (36-70%)	1.9 (0.92-3.4)	5.5% (2-10%)	6.1% (2.2-12%)	23 (8.1-43)	240 (86-450)
Lesotho	1990	Male	85 to 89	44% (29-60%)	56% (40-71%)	1.6 (0.76-2.9)	4% (1.3-7.8%)	4.5% (1.5-8.9%)	8.7 (2.8-18)	65 (22-130)
Lesotho	1990	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.4 (0.67-2.6)	2.9% (0.85-6.2%)	3.3% (0.89-7.2%)	1.9 (0.53-4.3)	10 (3-22)
Lesotho	1990	Male	95 plus	45% (26-67%)	55% (33-74%)	1.3 (0.49-2.6)	2.5% (0.5-5.9%)	2.7% (0.43-6.7%)	0.34 (0.054-0.85)	1.2 (0.25-2.9)
Lesotho	1995	Female	All Ages	16% (13-20%)	84% (80-87%)	1.1 (0.79-1.5)	3% (1.8-4.5%)	5.2% (2.8-8%)	430 (220-680)	14000 (8300-21000)
Lesotho	1995	Female	15 to 19	14% (8.2-22%)	86% (78-92%)	0.85 (0.37-1.7)	3% (1.4-5.7%)	4.8% (1.8-9.3%)	6.6 (2.5-13)	560 (250-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	1995	Female	20 to 24	18% (11-29%)	82% (71-89%)	1.5 (0.77-2.8)	3.7% (1.9-6.5%)	4.5% (1.9-8.6%)	12 (4.9-22)	960 (480-1600)
Lesotho	1995	Female	25 to 29	21% (12-34%)	79% (66-88%)	2 (0.98-3.5)	3.5% (1.7-6%)	4.1% (1.6-7.7%)	13 (5.8-23)	980 (510-1600)
Lesotho	1995	Female	30 to 34	24% (14-37%)	76% (63-86%)	2.3 (1.1-3.9)	4% (2.1-7%)	4.9% (2.2-9.2%)	14 (7-25)	930 (490-1600)
Lesotho	1995	Female	35 to 39	26% (15-41%)	74% (59-85%)	2.4 (1.2-4.2)	4.9% (2.5-8%)	6.2% (2.9-11%)	18 (8.9-30)	980 (550-1600)
Lesotho	1995	Female	40 to 44	28% (17-43%)	72% (57-83%)	2.4 (1.1-4.4)	6% (3.3-9.7%)	7.9% (3.9-13%)	23 (12-39)	1100 (610-1900)
Lesotho	1995	Female	45 to 49	28% (15-44%)	72% (56-85%)	2.4 (1.1-4.5)	7% (3.8-11%)	9.3% (4.8-15%)	27 (13-46)	1200 (600-1900)
Lesotho	1995	Female	50 to 54	27% (16-43%)	73% (57-84%)	2.3 (1.1-4.4)	7.9% (4.1-13%)	10% (5.4-17%)	27 (13-48)	1000 (500-1800)
Lesotho	1995	Female	55 to 59	27% (16-41%)	73% (59-84%)	2.3 (1.1-4.2)	8.2% (4.1-14%)	11% (5.2-17%)	41 (19-73)	1300 (640-2400)
Lesotho	1995	Female	60 to 64	28% (16-43%)	72% (57-84%)	2.2 (0.99-4.3)	8.1% (3.8-14%)	10% (4.8-18%)	57 (24-100)	1500 (670-2700)
Lesotho	1995	Female	65 to 69	28% (18-42%)	72% (58-82%)	2 (0.98-3.6)	8% (3.4-13%)	9.9% (4.2-16%)	59 (23-100)	1300 (520-2300)
Lesotho	1995	Female	70 to 74	26% (16-39%)	74% (61-84%)	1.7 (0.77-3)	6.2% (2.1-11%)	7.5% (2.5-13%)	52 (18-93)	940 (320-1700)
Lesotho	1995	Female	75 to 79	23% (14-36%)	77% (64-86%)	1.3 (0.55-2.6)	4.6% (1.2-9%)	5.5% (1.4-10%)	37 (8.6-74)	520 (120-1100)
Lesotho	1995	Female	80 to 84	22% (13-34%)	78% (66-87%)	1.2 (0.55-2.2)	3.9% (0.98-7.9%)	4.6% (1.2-9.3%)	23 (5.1-48)	240 (54-490)
Lesotho	1995	Female	85 to 89	23% (14-34%)	77% (66-86%)	1.2 (0.53-2.2)	3.4% (0.76-7.1%)	4% (0.87-8.4%)	13 (2.6-26)	92 (19-190)
Lesotho	1995	Female	90 to 94	24% (15-37%)	76% (63-85%)	1.2 (0.54-2.3)	2.9% (0.52-6.8%)	3.5% (0.57-8%)	4.4 (0.7-10)	22 (3.9-51)
Lesotho	1995	Female	95 plus	28% (15-44%)	72% (56-85%)	1.3 (0.45-2.7)	2.8% (0.085-7.2%)	3.3% (0.028-8.5%)	1.1 (0.0098-2.9)	3.7 (0.096-9.6)
Lesotho	1995	Male	All Ages	37% (32-41%)	63% (59-68%)	2.9 (2.1-3.7)	8.4% (6.4-11%)	12% (8.7-15%)	1200 (860-1600)	48000 (36000-62000)
Lesotho	1995	Male	15 to 19	30% (21-39%)	70% (61-79%)	2 (1-3.4)	8.8% (5.2-13%)	12% (6.4-19%)	30 (16-48)	2300 (1300-3600)
Lesotho	1995	Male	20 to 24	48% (37-59%)	52% (41-63%)	4.6 (2.5-7.3)	15% (9.7-21%)	18% (11-26%)	78 (49-110)	5500 (3600-7700)
Lesotho	1995	Male	25 to 29	56% (45-66%)	44% (34-55%)	6.3 (3.6-9.4)	15% (9.4-21%)	17% (10-24%)	85 (59-110)	5400 (3900-7200)
Lesotho	1995	Male	30 to 34	60% (49-69%)	40% (31-51%)	7 (4.3-10)	14% (9.6-20%)	16% (10-24%)	73 (51-99)	4300 (3000-5700)
Lesotho	1995	Male	35 to 39	62% (51-70%)	38% (30-49%)	6.9 (4-11)	16% (10-22%)	18% (11-26%)	75 (52-100)	3900 (2800-5300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	1995	Male	40 to 44	64% (53-72%)	36% (28-47%)	6.9 (3.8-11)	18% (12-23%)	20% (14-27%)	86 (60-120)	4000 (2800-5400)
Lesotho	1995	Male	45 to 49	66% (54-74%)	34% (26-46%)	7 (3.8-12)	20% (15-26%)	24% (17-30%)	100 (70-140)	4200 (3000-5600)
Lesotho	1995	Male	50 to 54	66% (56-74%)	34% (26-44%)	6.8 (3.6-11)	22% (15-28%)	25% (18-32%)	110 (73-150)	4100 (2700-5500)
Lesotho	1995	Male	55 to 59	64% (53-72%)	36% (28-47%)	6.2 (3.4-10)	21% (14-27%)	24% (16-31%)	140 (88-190)	4400 (2900-5900)
Lesotho	1995	Male	60 to 64	60% (47-70%)	40% (30-53%)	5.2 (2.6-8.7)	18% (11-24%)	21% (12-27%)	150 (86-210)	4100 (2300-5600)
Lesotho	1995	Male	65 to 69	54% (37-71%)	46% (29-63%)	3.8 (1.9-6.2)	15% (8-21%)	17% (9.1-24%)	130 (66-190)	2800 (1500-4200)
Lesotho	1995	Male	70 to 74	50% (33-67%)	50% (33-67%)	2.7 (1.3-4.9)	9.6% (4.3-16%)	11% (4.8-18%)	87 (37-150)	1600 (670-2600)
Lesotho	1995	Male	75 to 79	48% (33-66%)	52% (34-67%)	2.2 (0.97-4.1)	6.9% (2.4-13%)	7.7% (2.7-14%)	50 (17-94)	700 (240-1300)
Lesotho	1995	Male	80 to 84	47% (30-65%)	53% (35-70%)	1.9 (0.88-3.4)	5.4% (1.8-10%)	6.1% (2-11%)	26 (8.7-50)	280 (94-520)
Lesotho	1995	Male	85 to 89	44% (29-60%)	56% (40-71%)	1.6 (0.76-2.9)	4% (1.3-8.1%)	4.6% (1.5-9.2%)	10 (3.3-21)	77 (26-160)
Lesotho	1995	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.4 (0.67-2.7)	3% (0.92-6.4%)	3.3% (0.99-7.4%)	2.6 (0.75-5.8)	14 (4.2-29)
Lesotho	1995	Male	95 plus	45% (26-67%)	55% (33-74%)	1.3 (0.49-2.7)	2.6% (0.53-6.1%)	2.8% (0.45-7%)	0.48 (0.076-1.2)	1.7 (0.35-4.2)
Lesotho	2000	Female	All Ages	16% (13-20%)	84% (80-87%)	1.2 (0.86-1.7)	3% (1.6-4.6%)	4.9% (2.6-7.6%)	690 (340-1100)	22000 (12000-33000)
Lesotho	2000	Female	15 to 19	14% (8.2-22%)	86% (78-92%)	0.92 (0.41-1.8)	3.4% (1.5-6.3%)	5% (1.8-9.8%)	10 (4-20)	840 (370-1500)
Lesotho	2000	Female	20 to 24	18% (11-29%)	82% (71-89%)	1.7 (0.79-3.1)	3% (1.3-5.5%)	3.3% (1.2-6.9%)	18 (8-33)	1400 (690-2400)
Lesotho	2000	Female	25 to 29	21% (12-33%)	79% (67-88%)	2.2 (1-3.9)	2.2% (0.97-4.3%)	2.2% (0.83-4.8%)	20 (8.7-37)	1400 (670-2400)
Lesotho	2000	Female	30 to 34	24% (14-37%)	76% (63-86%)	2.5 (1.2-4.4)	2.5% (1.1-4.4%)	2.6% (1.1-4.8%)	24 (10-43)	1500 (690-2500)
Lesotho	2000	Female	35 to 39	26% (16-40%)	74% (60-84%)	2.6 (1.3-4.6)	3.2% (1.4-5.8%)	3.5% (1.4-6.7%)	30 (14-53)	1600 (780-2800)
Lesotho	2000	Female	40 to 44	28% (17-43%)	72% (57-83%)	2.7 (1.2-4.8)	4.2% (2-7.4%)	4.8% (2.1-8.5%)	38 (17-67)	1800 (840-3100)
Lesotho	2000	Female	45 to 49	28% (16-43%)	72% (57-84%)	2.6 (1.2-5.1)	5.5% (2.5-9.3%)	6.4% (2.9-11%)	49 (21-87)	2100 (910-3600)
Lesotho	2000	Female	50 to 54	27% (16-42%)	73% (58-84%)	2.5 (1.1-4.8)	7.1% (3.4-12%)	8.3% (3.9-14%)	57 (24-100)	2100 (930-3700)
Lesotho	2000	Female	55 to 59	27% (17-42%)	73% (58-83%)	2.5 (1.1-4.7)	8.2% (4.1-13%)	9.7% (4.9-16%)	54 (23-96)	1700 (760-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	2000	Female	60 to 64	28% (17-44%)	72% (56-83%)	2.4 (1.1-4.8)	8.6% (4-15%)	10% (4.8-17%)	76 (32-130)	2000 (860-3500)
Lesotho	2000	Female	65 to 69	28% (18-42%)	72% (58-82%)	2.2 (1.1-3.9)	8.9% (4-15%)	11% (4.8-17%)	100 (42-180)	2300 (920-4000)
Lesotho	2000	Female	70 to 74	26% (16-39%)	74% (61-84%)	1.8 (0.82-3.3)	7.1% (2.6-12%)	8.3% (3.1-14%)	79 (26-140)	1400 (480-2500)
Lesotho	2000	Female	75 to 79	23% (14-35%)	77% (65-86%)	1.4 (0.6-2.8)	5.4% (1.4-10%)	6.3% (1.7-12%)	59 (15-120)	820 (210-1600)
Lesotho	2000	Female	80 to 84	22% (13-35%)	78% (65-87%)	1.3 (0.6-2.4)	4.8% (1.4-9.1%)	5.6% (1.6-11%)	40 (10-78)	410 (110-810)
Lesotho	2000	Female	85 to 89	23% (14-35%)	77% (65-86%)	1.3 (0.59-2.5)	4.3% (1.1-8.5%)	5% (1.3-9.9%)	21 (5.1-42)	150 (37-310)
Lesotho	2000	Female	90 to 94	25% (15-37%)	75% (63-85%)	1.4 (0.6-2.5)	3.8% (0.86-8%)	4.4% (0.97-9.4%)	7.2 (1.5-16)	36 (8-78)
Lesotho	2000	Female	95 plus	28% (15-43%)	72% (57-85%)	1.4 (0.49-2.9)	3.6% (0.29-8.5%)	4.2% (0.27-10%)	1.9 (0.12-4.5)	6.4 (0.5-15)
Lesotho	2000	Male	All Ages	38% (34-41%)	62% (59-66%)	3.1 (2.2-4.1)	8.7% (6.3-11%)	12% (8.4-15%)	1800 (1300-2300)	70000 (50000-89000)
Lesotho	2000	Male	15 to 19	29% (25-33%)	71% (67-75%)	2 (1.1-3.2)	10% (6.1-15%)	13% (7.5-19%)	43 (24-65)	3200 (1900-4900)
Lesotho	2000	Male	20 to 24	47% (42-52%)	53% (48-58%)	4.6 (2.7-7)	15% (9.4-21%)	17% (10-24%)	110 (71-150)	7400 (5000-10000)
Lesotho	2000	Male	25 to 29	58% (53-62%)	42% (38-47%)	6.7 (4.1-10)	12% (7.1-17%)	12% (7.3-20%)	120 (84-160)	7600 (5400-10000)
Lesotho	2000	Male	30 to 34	62% (57-66%)	38% (34-43%)	7.6 (4.8-11)	11% (6.7-15%)	11% (6.8-17%)	110 (78-150)	6500 (4600-8700)
Lesotho	2000	Male	35 to 39	63% (58-68%)	37% (32-42%)	7.3 (4.5-11)	12% (7.3-17%)	13% (7.7-19%)	120 (79-160)	6100 (4100-8200)
Lesotho	2000	Male	40 to 44	67% (62-71%)	33% (29-38%)	7.6 (4.3-12)	14% (9.6-20%)	16% (10-22%)	140 (90-190)	6300 (4200-8600)
Lesotho	2000	Male	45 to 49	70% (66-74%)	30% (26-34%)	8 (4.5-13)	18% (13-24%)	20% (14-26%)	170 (110-230)	6900 (4700-9300)
Lesotho	2000	Male	50 to 54	71% (66-74%)	29% (26-34%)	7.8 (4.4-12)	22% (16-28%)	24% (17-31%)	200 (130-270)	7200 (4800-9700)
Lesotho	2000	Male	55 to 59	67% (62-71%)	33% (29-38%)	7 (3.9-11)	22% (15-28%)	24% (17-31%)	180 (120-240)	5600 (3700-7500)
Lesotho	2000	Male	60 to 64	60% (48-70%)	40% (30-52%)	5.5 (2.8-9.2)	19% (12-25%)	21% (14-28%)	190 (110-250)	5000 (3000-6700)
Lesotho	2000	Male	65 to 69	55% (37-71%)	45% (29-63%)	4.1 (2.1-6.8)	16% (9.3-23%)	18% (10-25%)	180 (100-260)	4000 (2300-5700)
Lesotho	2000	Male	70 to 74	50% (33-68%)	50% (32-67%)	2.9 (1.5-5.1)	11% (5.1-17%)	12% (5.7-19%)	120 (54-190)	2100 (1000-3400)
Lesotho	2000	Male	75 to 79	49% (32-66%)	51% (34-68%)	2.4 (1.1-4.3)	8% (3.1-14%)	8.9% (3.4-16%)	81 (31-150)	1100 (430-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	2000	Male	80 to 84	47% (30-64%)	53% (36-70%)	2.1 (0.98-3.5)	6.5% (2.3-11%)	7.3% (2.6-13%)	44 (15-80)	450 (160-820)
Lesotho	2000	Male	85 to 89	44% (29-61%)	56% (39-71%)	1.7 (0.85-3)	4.8% (1.9-9.1%)	5.4% (2.1-10%)	14 (5.3-27)	100 (40-200)
Lesotho	2000	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.5 (0.71-2.7)	3.5% (1.2-7%)	3.9% (1.3-8.1%)	3.5 (1.1-7.1)	18 (6.1-36)
Lesotho	2000	Male	95 plus	46% (26-67%)	54% (33-74%)	1.4 (0.54-2.6)	3% (0.79-6.6%)	3.4% (0.76-7.6%)	0.74 (0.17-1.7)	2.6 (0.69-5.7)
Lesotho	2005	Female	All Ages	16% (13-20%)	84% (80-87%)	1.3 (0.89-1.8)	3% (1.7-4.6%)	4.8% (2.6-7.3%)	920 (480-1400)	29000 (16000-43000)
Lesotho	2005	Female	15 to 19	14% (8-22%)	86% (78-92%)	0.93 (0.41-1.8)	4.1% (1.8-7.8%)	5.8% (2.2-12%)	14 (5.6-29)	1100 (520-2200)
Lesotho	2005	Female	20 to 24	19% (11-29%)	81% (71-89%)	1.7 (0.83-3.2)	3.5% (1.6-6.6%)	3.9% (1.4-8%)	27 (12-50)	2000 (1000-3500)
Lesotho	2005	Female	25 to 29	21% (12-34%)	79% (66-88%)	2.3 (1.1-4)	2% (0.88-4.1%)	2.1% (0.79-4.4%)	29 (13-54)	1900 (960-3400)
Lesotho	2005	Female	30 to 34	24% (14-37%)	76% (63-86%)	2.6 (1.2-4.6)	2% (0.97-3.8%)	2% (0.9-4.1%)	31 (14-56)	1800 (910-3200)
Lesotho	2005	Female	35 to 39	27% (15-40%)	73% (60-85%)	2.7 (1.3-5)	2.7% (1.2-5.1%)	2.9% (1.2-5.6%)	39 (18-71)	2000 (980-3700)
Lesotho	2005	Female	40 to 44	28% (17-43%)	72% (57-83%)	2.8 (1.2-5.2)	3.4% (1.5-5.9%)	3.6% (1.6-6.5%)	46 (21-83)	2200 (1000-3900)
Lesotho	2005	Female	45 to 49	28% (16-43%)	72% (57-84%)	2.7 (1.2-5.4)	4.7% (2.2-8.2%)	5.2% (2.4-9.1%)	64 (29-110)	2700 (1200-4600)
Lesotho	2005	Female	50 to 54	28% (16-43%)	72% (57-84%)	2.6 (1.2-5)	6.6% (3.2-11%)	7.4% (3.6-12%)	85 (38-150)	3100 (1400-5300)
Lesotho	2005	Female	55 to 59	27% (17-42%)	73% (58-83%)	2.6 (1.1-4.9)	8.2% (4.2-13%)	9.4% (4.8-15%)	91 (43-160)	2900 (1400-5000)
Lesotho	2005	Female	60 to 64	28% (17-43%)	72% (57-83%)	2.5 (1.1-4.8)	8.8% (4.1-14%)	10% (4.7-17%)	79 (34-140)	2100 (920-3600)
Lesotho	2005	Female	65 to 69	28% (18-42%)	72% (58-82%)	2.2 (1.1-4.1)	9.4% (4.3-15%)	11% (5.2-17%)	120 (51-200)	2600 (1100-4500)
Lesotho	2005	Female	70 to 74	26% (17-39%)	74% (61-83%)	1.8 (0.81-3.4)	7.6% (2.9-13%)	8.8% (3.4-15%)	120 (44-210)	2100 (790-3800)
Lesotho	2005	Female	75 to 79	23% (14-36%)	77% (64-86%)	1.5 (0.57-2.8)	6.1% (1.7-11%)	7% (2-13%)	80 (21-160)	1100 (300-2200)
Lesotho	2005	Female	80 to 84	22% (13-35%)	78% (65-87%)	1.4 (0.6-2.4)	5.5% (1.8-10%)	6.4% (2-12%)	57 (17-110)	580 (180-1100)
Lesotho	2005	Female	85 to 89	23% (14-35%)	77% (65-86%)	1.4 (0.6-2.6)	4.9% (1.4-9.5%)	5.7% (1.7-11%)	28 (7.7-58)	210 (56-420)
Lesotho	2005	Female	90 to 94	25% (15-37%)	75% (63-85%)	1.4 (0.59-2.6)	4.3% (1-8.8%)	5% (1.2-10%)	9.5 (2.2-20)	48 (11-100)
Lesotho	2005	Female	95 plus	27% (15-43%)	73% (57-85%)	1.5 (0.47-3.1)	4.2% (0.37-9.6%)	4.9% (0.37-11%)	2.5 (0.19-6)	8.6 (0.77-20)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	2005	Male	All Ages	38% (35-41%)	62% (59-65%)	3.4 (2.5-4.4)	9.6% (7.1-12%)	13% (9.3-16%)	2600 (1800-3400)	99000 (73000-130000)
Lesotho	2005	Male	15 to 19	30% (27-33%)	70% (67-73%)	2.2 (1.3-3.6)	12% (7.8-17%)	15% (9.4-22%)	65 (38-95)	4800 (3000-7000)
Lesotho	2005	Male	20 to 24	49% (46-52%)	51% (48-54%)	5.2 (3.1-7.7)	18% (12-26%)	21% (13-30%)	170 (110-230)	11000 (7800-15000)
Lesotho	2005	Male	25 to 29	58% (55-61%)	42% (39-45%)	7.2 (4.4-11)	13% (7.7-19%)	14% (7.9-21%)	180 (130-250)	11000 (8000-15000)
Lesotho	2005	Male	30 to 34	62% (59-65%)	38% (35-41%)	7.9 (5-12)	10% (6.6-15%)	11% (6.7-17%)	170 (120-230)	9600 (6600-13000)
Lesotho	2005	Male	35 to 39	63% (60-66%)	37% (34-40%)	7.6 (4.5-12)	12% (7.1-17%)	12% (7.4-18%)	170 (110-230)	8800 (6000-12000)
Lesotho	2005	Male	40 to 44	66% (62-69%)	34% (31-38%)	7.8 (4.3-13)	14% (9-19%)	14% (9.5-20%)	180 (120-240)	8300 (5500-11000)
Lesotho	2005	Male	45 to 49	69% (65-72%)	31% (28-35%)	8.1 (4.5-13)	17% (12-23%)	19% (13-25%)	220 (140-300)	9000 (6000-12000)
Lesotho	2005	Male	50 to 54	70% (66-73%)	30% (27-34%)	8.1 (4.3-13)	22% (16-28%)	24% (17-30%)	280 (190-380)	10000 (6900-14000)
Lesotho	2005	Male	55 to 59	66% (62-70%)	34% (30-38%)	7.2 (4-11)	23% (17-29%)	25% (18-32%)	290 (200-390)	9100 (6200-12000)
Lesotho	2005	Male	60 to 64	59% (48-70%)	41% (30-52%)	5.7 (2.8-9.5)	20% (13-27%)	23% (15-30%)	250 (150-330)	6600 (4100-8800)
Lesotho	2005	Male	65 to 69	55% (37-70%)	45% (30-63%)	4.3 (2.1-7)	18% (11-25%)	20% (12-28%)	240 (140-340)	5400 (3200-7600)
Lesotho	2005	Male	70 to 74	50% (33-68%)	50% (32-67%)	3.1 (1.5-5.3)	13% (6.3-19%)	14% (6.9-21%)	150 (77-240)	2800 (1400-4300)
Lesotho	2005	Male	75 to 79	49% (33-66%)	51% (34-67%)	2.5 (1.1-4.6)	9.4% (3.6-16%)	10% (4-18%)	99 (38-170)	1400 (540-2300)
Lesotho	2005	Male	80 to 84	47% (30-64%)	53% (36-70%)	2.2 (1-3.8)	7.9% (3-14%)	8.8% (3.4-15%)	68 (27-120)	710 (270-1200)
Lesotho	2005	Male	85 to 89	44% (29-61%)	56% (39-71%)	1.8 (0.87-3.3)	6% (2.3-11%)	6.8% (2.6-13%)	22 (8.7-41)	170 (64-300)
Lesotho	2005	Male	90 to 94	44% (28-60%)	56% (40-72%)	1.6 (0.75-2.9)	4.4% (1.5-8.9%)	5.1% (1.7-10%)	5 (1.6-9.9)	25 (8.5-51)
Lesotho	2005	Male	95 plus	46% (27-67%)	54% (33-73%)	1.5 (0.53-3)	4.1% (0.97-8.8%)	4.7% (0.98-10%)	1.2 (0.25-2.5)	4.1 (0.97-8.8)
Lesotho	2010	Female	All Ages	16% (13-20%)	84% (80-87%)	1.3 (0.91-1.9)	3.5% (2.1-5.2%)	5.6% (3.2-8.4%)	970 (520-1500)	30000 (18000-45000)
Lesotho	2010	Female	15 to 19	14% (8.2-23%)	86% (77-92%)	0.97 (0.41-1.9)	3.8% (1.6-7%)	5% (1.9-9.8%)	15 (5.7-29)	1200 (530-2300)
Lesotho	2010	Female	20 to 24	19% (11-29%)	81% (71-89%)	1.8 (0.84-3.3)	4.2% (1.9-7.9%)	4.9% (1.8-9.7%)	29 (13-53)	2200 (1100-3800)
Lesotho	2010	Female	25 to 29	22% (12-34%)	78% (66-88%)	2.3 (1.1-4)	2.6% (1.2-5%)	2.8% (1.1-5.7%)	36 (16-67)	2400 (1200-4300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	2010	Female	30 to 34	24% (14-37%)	76% (63-86%)	2.6 (1.3-4.5)	2.4% (1.2-4.4%)	2.5% (1.1-4.7%)	37 (16-67)	2200 (1100-3900)
Lesotho	2010	Female	35 to 39	27% (15-41%)	73% (59-85%)	2.8 (1.3-5.2)	3.2% (1.5-5.7%)	3.5% (1.5-6.3%)	38 (17-68)	2000 (930-3500)
Lesotho	2010	Female	40 to 44	28% (17-45%)	72% (55-83%)	2.9 (1.2-5.4)	4.2% (2.1-7%)	4.6% (2.2-7.8%)	42 (19-72)	2000 (950-3300)
Lesotho	2010	Female	45 to 49	29% (16-44%)	71% (56-84%)	2.9 (1.2-5.7)	6.1% (3-10%)	6.9% (3.3-11%)	60 (27-100)	2500 (1200-4200)
Lesotho	2010	Female	50 to 54	28% (16-43%)	72% (57-84%)	2.8 (1.2-5.5)	8.3% (4.2-13%)	9.6% (5-16%)	86 (42-140)	3200 (1600-5200)
Lesotho	2010	Female	55 to 59	27% (16-42%)	73% (58-84%)	2.7 (1.2-5.2)	10% (5.1-16%)	12% (6.1-19%)	110 (49-180)	3300 (1600-5600)
Lesotho	2010	Female	60 to 64	28% (17-43%)	72% (57-83%)	2.6 (1.1-5)	10% (4.7-16%)	12% (5.6-19%)	110 (45-170)	2800 (1200-4700)
Lesotho	2010	Female	65 to 69	28% (18-42%)	72% (58-82%)	2.4 (1-4.3)	10% (4.7-17%)	12% (5.8-19%)	100 (43-170)	2200 (960-3800)
Lesotho	2010	Female	70 to 74	26% (17-39%)	74% (61-83%)	1.9 (0.83-3.6)	8.2% (3-14%)	9.5% (3.6-16%)	110 (38-190)	2000 (670-3500)
Lesotho	2010	Female	75 to 79	23% (14-36%)	77% (64-86%)	1.5 (0.6-3.1)	6.3% (1.6-12%)	7.3% (1.9-14%)	98 (24-190)	1400 (330-2700)
Lesotho	2010	Female	80 to 84	22% (14-35%)	78% (65-86%)	1.4 (0.61-2.6)	5.7% (1.5-11%)	6.5% (1.9-12%)	72 (19-140)	740 (200-1400)
Lesotho	2010	Female	85 to 89	23% (14-34%)	77% (66-86%)	1.4 (0.6-2.7)	5.1% (1.3-10%)	5.9% (1.5-12%)	28 (7-56)	200 (51-410)
Lesotho	2010	Female	90 to 94	25% (15-37%)	75% (63-85%)	1.4 (0.58-2.8)	4.4% (0.82-9.2%)	5.2% (0.91-11%)	9.1 (1.6-19)	46 (8.5-95)
Lesotho	2010	Female	95 plus	27% (15-44%)	73% (56-85%)	1.5 (0.48-3.4)	4.4% (0.3-10%)	5.1% (0.29-12%)	2.5 (0.15-5.8)	8.4 (0.59-19)
Lesotho	2010	Male	All Ages	39% (36-42%)	61% (58-64%)	3.6 (2.5-4.9)	11% (8.4-14%)	14% (11-18%)	2600 (1900-3400)	110000 (80000-140000)
Lesotho	2010	Male	15 to 19	31% (27-34%)	69% (66-73%)	2.3 (1.3-3.7)	11% (7-16%)	14% (8.1-20%)	67 (40-99)	5000 (3000-7300)
Lesotho	2010	Male	20 to 24	50% (46-53%)	50% (47-54%)	5.3 (3.1-8.1)	20% (14-27%)	23% (16-32%)	180 (130-250)	13000 (9000-17000)
Lesotho	2010	Male	25 to 29	58% (55-62%)	42% (38-45%)	7.3 (4.4-11)	16% (9.9-22%)	17% (10-26%)	230 (160-300)	14000 (10000-19000)
Lesotho	2010	Male	30 to 34	62% (59-65%)	38% (35-41%)	8.1 (5.2-12)	12% (8.3-17%)	13% (8.7-19%)	220 (150-290)	12000 (8900-16000)
Lesotho	2010	Male	35 to 39	63% (60-67%)	37% (33-40%)	7.9 (4.6-13)	13% (8.7-18%)	14% (9.3-20%)	200 (140-270)	10000 (7400-14000)
Lesotho	2010	Male	40 to 44	66% (63-70%)	34% (30-37%)	8.1 (4.4-13)	16% (11-21%)	17% (12-22%)	190 (130-250)	8700 (6000-12000)
Lesotho	2010	Male	45 to 49	69% (66-72%)	31% (28-34%)	8.4 (4.5-14)	20% (15-26%)	22% (16-28%)	200 (140-270)	8300 (5700-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	2010	Male	50 to 54	70% (67-73%)	30% (27-33%)	8.3 (4.2-13)	25% (18-30%)	27% (20-33%)	250 (170-330)	9100 (6300-12000)
Lesotho	2010	Male	55 to 59	66% (62-70%)	34% (30-38%)	7.4 (3.9-12)	26% (19-32%)	29% (21-35%)	280 (200-370)	8900 (6200-12000)
Lesotho	2010	Male	60 to 64	59% (48-70%)	41% (30-52%)	5.8 (2.8-10)	22% (15-29%)	25% (16-32%)	260 (160-350)	6900 (4300-9200)
Lesotho	2010	Male	65 to 69	54% (37-71%)	46% (29-63%)	4.3 (2.1-7.3)	19% (11-26%)	21% (13-30%)	210 (120-290)	4600 (2600-6500)
Lesotho	2010	Male	70 to 74	50% (33-67%)	50% (33-67%)	3.1 (1.4-5.4)	13% (6.2-20%)	15% (7-22%)	150 (71-240)	2700 (1300-4200)
Lesotho	2010	Male	75 to 79	48% (33-65%)	52% (35-67%)	2.5 (1.1-4.7)	9.7% (3.3-16%)	11% (3.7-18%)	97 (34-170)	1300 (470-2300)
Lesotho	2010	Male	80 to 84	47% (30-64%)	53% (36-70%)	2.2 (1-4)	7.9% (3-14%)	8.9% (3.4-16%)	70 (26-130)	730 (270-1300)
Lesotho	2010	Male	85 to 89	44% (28-60%)	56% (40-72%)	1.8 (0.82-3.4)	6% (2.1-11%)	6.9% (2.4-13%)	24 (8.6-45)	180 (64-330)
Lesotho	2010	Male	90 to 94	43% (28-60%)	57% (40-72%)	1.6 (0.7-3)	4.5% (1.4-9%)	5.2% (1.7-11%)	5.5 (1.7-11)	28 (9.5-57)
Lesotho	2010	Male	95 plus	45% (26-66%)	55% (34-74%)	1.5 (0.53-3.1)	4.3% (1-9.1%)	4.9% (1.1-11%)	1.3 (0.29-2.8)	4.5 (1.1-9.6)
Lesotho	2016	Female	All Ages	17% (13-21%)	83% (79-87%)	1.5 (0.83-2.4)	3.7% (2-5.9%)	5.8% (2.9-9.2%)	880 (380-1500)	28000 (14000-47000)
Lesotho	2016	Female	15 to 19	14% (8.2-23%)	86% (77-92%)	1 (0.41-2.1)	2.9% (1.2-5.6%)	3.4% (1.2-6.8%)	13 (4.8-27)	1100 (460-2200)
Lesotho	2016	Female	20 to 24	19% (11-29%)	81% (71-89%)	1.9 (0.81-3.7)	4.1% (1.9-7.7%)	4.6% (1.7-9.4%)	26 (10-50)	2000 (980-3600)
Lesotho	2016	Female	25 to 29	22% (13-34%)	78% (66-87%)	2.4 (1.1-4.5)	3% (1.3-6.2%)	3.2% (1.2-7.2%)	35 (14-73)	2400 (1100-4700)
Lesotho	2016	Female	30 to 34	24% (14-38%)	76% (62-86%)	2.8 (1.3-5.3)	2.7% (1.2-5.3%)	2.8% (1.1-5.8%)	43 (17-85)	2600 (1100-4900)
Lesotho	2016	Female	35 to 39	27% (16-41%)	73% (59-84%)	2.9 (1.3-5.6)	3.4% (1.5-6.2%)	3.7% (1.5-6.8%)	45 (17-90)	2400 (1000-4700)
Lesotho	2016	Female	40 to 44	28% (17-43%)	72% (57-83%)	2.9 (1.1-5.8)	4.5% (2.1-7.8%)	5% (2.2-8.9%)	39 (16-76)	1800 (790-3500)
Lesotho	2016	Female	45 to 49	28% (16-44%)	72% (56-84%)	2.9 (1.1-6.1)	6.4% (2.9-11%)	7.4% (3.3-13%)	43 (17-78)	1800 (750-3200)
Lesotho	2016	Female	50 to 54	28% (16-44%)	72% (56-84%)	2.8 (1-5.7)	8.2% (3.9-13%)	9.8% (4.6-16%)	58 (23-110)	2200 (890-3900)
Lesotho	2016	Female	55 to 59	27% (16-42%)	73% (58-84%)	2.7 (1.1-5.8)	10% (4.8-16%)	12% (6-20%)	83 (33-150)	2600 (1100-4800)
Lesotho	2016	Female	60 to 64	28% (17-43%)	72% (57-83%)	2.7 (1-5.4)	10% (4.2-16%)	12% (5.3-19%)	100 (38-180)	2800 (1100-5000)
Lesotho	2016	Female	65 to 69	28% (18-43%)	72% (57-82%)	2.4 (0.99-4.9)	10% (4.3-17%)	12% (5.3-20%)	120 (43-210)	2600 (970-4700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	2016	Female	70 to 74	26% (17-40%)	74% (60-83%)	2 (0.77-4.1)	8% (2.5-14%)	9.5% (3.2-16%)	83 (24-150)	1500 (430-2700)
Lesotho	2016	Female	75 to 79	23% (14-36%)	77% (64-86%)	1.6 (0.63-3.5)	6.3% (1.6-12%)	7.4% (1.9-13%)	75 (16-150)	1000 (220-2100)
Lesotho	2016	Female	80 to 84	22% (13-35%)	78% (65-87%)	1.5 (0.6-3.1)	5.6% (1.4-11%)	6.5% (1.6-13%)	70 (17-140)	720 (170-1400)
Lesotho	2016	Female	85 to 89	23% (14-35%)	77% (65-86%)	1.5 (0.58-3.1)	5.1% (1.1-10%)	5.9% (1.3-12%)	36 (8.4-75)	270 (61-540)
Lesotho	2016	Female	90 to 94	25% (16-37%)	75% (63-84%)	1.5 (0.58-3.3)	4.5% (0.78-9.5%)	5.3% (0.92-11%)	9.2 (1.6-20)	46 (7.9-98)
Lesotho	2016	Female	95 plus	28% (16-44%)	72% (56-84%)	1.6 (0.52-3.5)	4.4% (0.37-10%)	5.2% (0.43-12%)	2.4 (0.19-5.6)	7.9 (0.71-19)
Lesotho	2016	Male	All Ages	40% (35-46%)	60% (54-65%)	4.1 (2.4-6.7)	12% (8.5-15%)	15% (11-19%)	2600 (1800-3500)	110000 (76000-140000)
Lesotho	2016	Male	15 to 19	30% (21-39%)	70% (61-79%)	2.4 (1.1-4.6)	9% (4.9-14%)	10% (5.4-16%)	61 (32-97)	4600 (2500-7200)
Lesotho	2016	Male	20 to 24	49% (37-59%)	51% (41-63%)	5.6 (2.8-9.9)	19% (13-26%)	22% (14-30%)	170 (110-240)	12000 (8100-17000)
Lesotho	2016	Male	25 to 29	57% (45-67%)	43% (33-55%)	7.6 (3.6-13)	16% (10-23%)	18% (11-26%)	230 (150-310)	14000 (9700-19000)
Lesotho	2016	Male	30 to 34	61% (49-70%)	39% (30-51%)	8.4 (4.3-15)	14% (9-19%)	15% (9.4-21%)	260 (170-350)	15000 (10000-20000)
Lesotho	2016	Male	35 to 39	63% (51-71%)	37% (29-49%)	8.2 (4.2-15)	14% (8.9-19%)	15% (9.5-21%)	260 (170-360)	13000 (8600-19000)
Lesotho	2016	Male	40 to 44	64% (53-73%)	36% (27-47%)	8.3 (3.8-16)	17% (11-22%)	18% (12-24%)	240 (150-340)	11000 (6800-15000)
Lesotho	2016	Male	45 to 49	66% (53-74%)	34% (26-47%)	8.5 (3.7-17)	21% (15-27%)	23% (16-30%)	220 (130-300)	8800 (5600-12000)
Lesotho	2016	Male	50 to 54	66% (56-74%)	34% (26-44%)	8.2 (3.6-16)	24% (17-30%)	26% (18-33%)	200 (120-270)	7200 (4500-9700)
Lesotho	2016	Male	55 to 59	64% (53-72%)	36% (28-47%)	7.4 (3.4-14)	25% (17-31%)	28% (19-34%)	220 (140-300)	6800 (4400-9300)
Lesotho	2016	Male	60 to 64	60% (48-70%)	40% (30-52%)	6.1 (2.6-12)	22% (14-29%)	25% (15-32%)	240 (140-330)	6300 (3700-8700)
Lesotho	2016	Male	65 to 69	55% (38-71%)	45% (29-62%)	4.6 (1.9-9)	19% (10-26%)	21% (11-29%)	220 (120-310)	4800 (2600-6900)
Lesotho	2016	Male	70 to 74	50% (34-67%)	50% (33-66%)	3.3 (1.4-6.5)	13% (5.7-21%)	15% (6.4-23%)	140 (59-220)	2500 (1100-4000)
Lesotho	2016	Male	75 to 79	49% (33-66%)	51% (34-67%)	2.7 (1-5.6)	10% (3.3-17%)	11% (3.7-19%)	96 (32-170)	1300 (460-2300)
Lesotho	2016	Male	80 to 84	47% (30-65%)	53% (35-70%)	2.3 (0.97-4.8)	8.3% (2.8-15%)	9.3% (3.1-17%)	64 (21-120)	660 (220-1200)
Lesotho	2016	Male	85 to 89	44% (29-60%)	56% (40-71%)	2 (0.81-4.3)	6.4% (2-13%)	7.3% (2.3-14%)	23 (7-44)	170 (53-330)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Lesotho	2016	Male	90 to 94	43% (27-61%)	57% (39-73%)	1.8 (0.68-3.8)	4.9% (1.3-10%)	5.7% (1.5-12%)	5.5 (1.4-11)	28 (7.6-58)
Lesotho	2016	Male	95 plus	45% (26-66%)	55% (34-74%)	1.6 (0.55-3.7)	4.7% (0.95-10%)	5.4% (0.97-12%)	1.4 (0.25-3)	4.8 (0.96-10)
Namibia	1990	Female	All Ages	27% (23-31%)	73% (69-77%)	0.14 (0.079-0.23)	0.19% (-0.16-0.53%)	-0.16% (-0.8-0.45%)	-8.8 (-44-24)	610 (-510-1700)
Namibia	1990	Female	15 to 19	23% (16-31%)	77% (69-84%)	0.12 (0.06-0.23)	0.63% (0.18-1.2%)	0.42% (-0.38-1.4%)	0.4 (-0.4-1.3)	86 (22-170)
Namibia	1990	Female	20 to 24	33% (23-44%)	67% (56-77%)	0.24 (0.12-0.44)	1.2% (0.33-2.3%)	0.69% (-0.72-2.2%)	0.87 (-0.94-2.8)	180 (43-330)
Namibia	1990	Female	25 to 29	37% (27-49%)	63% (51-73%)	0.27 (0.13-0.46)	1.2% (0.29-2.1%)	0.7% (-0.62-2.3%)	0.92 (-0.8-3)	160 (40-300)
Namibia	1990	Female	30 to 34	41% (29-53%)	59% (47-71%)	0.3 (0.14-0.51)	1.1% (0.16-2.1%)	0.77% (-0.76-2.4%)	1 (-0.99-3.3)	130 (20-260)
Namibia	1990	Female	35 to 39	44% (32-54%)	56% (46-68%)	0.33 (0.16-0.56)	1% (-0.11-2.2%)	0.92% (-0.88-2.7%)	1.3 (-1.2-3.7)	110 (-13-250)
Namibia	1990	Female	40 to 44	46% (35-56%)	54% (44-65%)	0.34 (0.17-0.6)	0.9% (-0.27-2.1%)	0.93% (-0.84-2.8%)	1.3 (-1.2-4)	92 (-27-210)
Namibia	1990	Female	45 to 49	46% (33-56%)	54% (44-67%)	0.31 (0.15-0.54)	0.71% (-0.42-1.9%)	0.73% (-0.9-2.4%)	1.2 (-1.5-3.8)	71 (-41-180)
Namibia	1990	Female	50 to 54	44% (33-56%)	56% (44-67%)	0.27 (0.13-0.48)	0.33% (-0.85-1.4%)	0.24% (-1.4-1.7%)	0.49 (-2.7-3.4)	33 (-84-140)
Namibia	1990	Female	55 to 59	43% (31-54%)	57% (46-69%)	0.23 (0.11-0.42)	-0.065% (-1.2-0.91%)	-0.18% (-1.6-1.1%)	-0.47 (-4.3-2.7)	-6.8 (-120-93)
Namibia	1990	Female	60 to 64	42% (30-55%)	58% (45-70%)	0.21 (0.095-0.39)	-0.48% (-1.6-0.48%)	-0.65% (-2-0.55%)	-2.2 (-6.8-1.9)	-55 (-180-55)
Namibia	1990	Female	65 to 69	41% (30-52%)	59% (48-70%)	0.19 (0.095-0.35)	-0.64% (-1.8-0.31%)	-0.79% (-2.2-0.38%)	-3.2 (-8.6-1.5)	-70 (-190-33)
Namibia	1990	Female	70 to 74	38% (28-48%)	62% (52-72%)	0.16 (0.076-0.3)	-0.6% (-1.7-0.13%)	-0.71% (-2-0.15%)	-3.3 (-9.1-0.66)	-59 (-160-12)
Namibia	1990	Female	75 to 79	33% (23-46%)	67% (54-77%)	0.13 (0.059-0.24)	-0.64% (-1.5-0.029%)	-0.75% (-1.8-0.023%)	-3.3 (-7.9-0.1)	-45 (-110-2.1)
Namibia	1990	Female	80 to 84	33% (22-44%)	67% (56-78%)	0.12 (0.056-0.22)	-0.58% (-1.4-0.026%)	-0.68% (-1.7-0.023%)	-2.2 (-5.4-0.072)	-23 (-56--1)
Namibia	1990	Female	85 to 89	35% (24-47%)	65% (53-76%)	0.12 (0.056-0.21)	-0.58% (-1.3-0.023%)	-0.68% (-1.6-0.032%)	-1.3 (-3-0.061)	-9.2 (-22--0.37)
Namibia	1990	Female	90 to 94	38% (27-51%)	62% (49-73%)	0.12 (0.054-0.22)	-0.57% (-1.4-0.034%)	-0.69% (-1.7-0.054%)	-0.47 (-1.1-0.037)	-2.3 (-5.7--0.14)
Namibia	1990	Female	95 plus	44% (28-62%)	56% (38-72%)	0.13 (0.05-0.25)	-0.54% (-1.5-0.027%)	-0.68% (-1.8-0.0099%)	-0.11 (-0.28--0.0016)	-0.33 (-0.91-0.017)
Namibia	1990	Male	All Ages	39% (34-44%)	61% (56-66%)	0.27 (0.16-0.43)	1.1% (0.34-1.9%)	1% (-0.065-2.1%)	68 (-4.3-140)	4400 (1300-7500)
Namibia	1990	Male	15 to 19	34% (24-44%)	66% (56-76%)	0.21 (0.11-0.36)	1.7% (0.79-2.8%)	1.4% (0.25-3.1%)	2.3 (0.41-5.1)	290 (130-510)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	1990	Male	20 to 24	56% (44-67%)	44% (33-56%)	0.53 (0.3-0.88)	3.7% (1.7-6.3%)	3.3% (0.64-6.5%)	7.7 (1.4-15)	780 (340-1300)
Namibia	1990	Male	25 to 29	62% (51-72%)	38% (28-49%)	0.63 (0.35-1.1)	3.7% (1.6-6.2%)	3.3% (0.61-6.5%)	8.2 (1.5-16)	740 (300-1200)
Namibia	1990	Male	30 to 34	65% (53-74%)	35% (26-47%)	0.65 (0.37-1.1)	3.2% (1.2-5.5%)	3% (0.28-5.9%)	6.9 (0.7-13)	550 (200-930)
Namibia	1990	Male	35 to 39	66% (55-74%)	34% (26-45%)	0.62 (0.34-1)	2.8% (0.75-5.2%)	2.7% (0.14-5.9%)	6.2 (0.34-13)	410 (110-770)
Namibia	1990	Male	40 to 44	65% (54-73%)	35% (27-46%)	0.59 (0.32-0.97)	2.5% (0.29-4.9%)	2.5% (-0.28-5.5%)	6.2 (-0.69-14)	350 (40-680)
Namibia	1990	Male	45 to 49	62% (48-70%)	38% (30-52%)	0.57 (0.32-0.94)	3% (0.51-5.8%)	3.1% (-0.03-6.5%)	8.4 (-0.087-18)	410 (73-790)
Namibia	1990	Male	50 to 54	62% (51-70%)	38% (30-49%)	0.56 (0.29-0.93)	2.6% (0.28-5.1%)	2.6% (-0.23-5.8%)	7.7 (-0.73-17)	350 (38-670)
Namibia	1990	Male	55 to 59	65% (54-73%)	35% (27-46%)	0.56 (0.29-0.92)	1.8% (-0.4-4.1%)	1.8% (-0.88-4.6%)	6.3 (-3.2-16)	250 (-53-550)
Namibia	1990	Male	60 to 64	62% (50-71%)	38% (29-50%)	0.45 (0.25-0.78)	0.8% (-1.2-4%)	0.69% (-1.5-2.6%)	3 (-6.7-11)	110 (-150-340)
Namibia	1990	Male	65 to 69	52% (40-63%)	48% (37-60%)	0.29 (0.16-0.51)	0.51% (-0.92-1.8%)	0.44% (-1.2-1.9%)	2 (-5.9-8.9)	62 (-110-210)
Namibia	1990	Male	70 to 74	48% (37-60%)	52% (40-63%)	0.18 (0.096-0.33)	0.38% (-0.58-1.2%)	0.34% (-0.78-1.2%)	1.5 (-3.5-5.8)	35 (-54-110)
Namibia	1990	Male	75 to 79	50% (35-64%)	50% (36-65%)	0.14 (0.065-0.25)	0.21% (-0.59-0.81%)	0.16% (-0.74-0.85%)	0.58 (-2.7-3.2)	12 (-34-48)
Namibia	1990	Male	80 to 84	50% (34-65%)	50% (35-66%)	0.12 (0.058-0.22)	0.17% (-0.45-0.65%)	0.13% (-0.6-0.68%)	0.3 (-1.4-1.6)	4.7 (-12-18)
Namibia	1990	Male	85 to 89	47% (33-61%)	53% (39-67%)	0.1 (0.049-0.18)	0.15% (-0.36-0.52%)	0.11% (-0.46-0.54%)	0.12 (-0.52-0.61)	1.4 (-3.3-4.9)
Namibia	1990	Male	90 to 94	46% (32-61%)	54% (39-68%)	0.089 (0.042-0.17)	0.15% (-0.25-0.45%)	0.11% (-0.35-0.46%)	0.045 (-0.14-0.19)	0.35 (-0.61-1.1)
Namibia	1990	Male	95 plus	48% (30-68%)	52% (32-70%)	0.082 (0.032-0.17)	0.2% (-0.16-0.45%)	0.12% (-0.31-0.42%)	0.012 (-0.03-0.041)	0.076 (-0.06-0.18)
Namibia	1995	Female	All Ages	23% (21-27%)	77% (73-79%)	0.27 (0.17-0.41)	0.36% (-0.083-0.88%)	-0.12% (-0.89-0.7%)	-8 (-61-45)	1400 (-320-3400)
Namibia	1995	Female	15 to 19	18% (14-24%)	82% (76-86%)	0.21 (0.11-0.37)	0.85% (0.38-1.5%)	0.81% (0.0072-1.9%)	0.91 (0.0096-2.2)	130 (58-240)
Namibia	1995	Female	20 to 24	27% (21-35%)	73% (65-79%)	0.43 (0.22-0.72)	1.4% (0.58-2.5%)	1.1% (0.067-2.6%)	2.5 (0.16-5.8)	320 (140-550)
Namibia	1995	Female	25 to 29	32% (26-39%)	68% (61-74%)	0.51 (0.29-0.82)	1.2% (0.49-2.3%)	0.95% (0.07-2.4%)	2.9 (0.21-6.5)	310 (130-550)
Namibia	1995	Female	30 to 34	36% (30-44%)	64% (56-70%)	0.57 (0.31-0.93)	1.2% (0.46-2.4%)	1.1% (0.1-2.8%)	3.5 (0.3-7.7)	300 (110-550)
Namibia	1995	Female	35 to 39	38% (32-47%)	62% (53-68%)	0.63 (0.35-1)	1.4% (0.41-2.9%)	1.5% (0.16-3.5%)	4.4 (0.58-9.8)	290 (87-570)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	1995	Female	40 to 44	39% (32-49%)	61% (51-68%)	0.63 (0.34-1.1)	1.5% (0.34-3.1%)	1.7% (0.19-4%)	4.7 (0.52-11)	260 (56-540)
Namibia	1995	Female	45 to 49	41% (33-49%)	59% (51-67%)	0.59 (0.33-1)	1.3% (0.053-3%)	1.5% (-0.14-3.7%)	4 (-0.36-10)	190 (8.3-440)
Namibia	1995	Female	50 to 54	42% (33-53%)	58% (47-67%)	0.56 (0.29-0.97)	0.8% (-0.64-2.7%)	0.9% (-0.96-3.4%)	2.6 (-2.7-9.4)	110 (-81-360)
Namibia	1995	Female	55 to 59	43% (31-54%)	57% (46-69%)	0.52 (0.26-0.9)	0.094% (-1.3-1.7%)	0.073% (-1.7-2.1%)	0.2 (-5.6-6.5)	11 (-170-210)
Namibia	1995	Female	60 to 64	43% (30-55%)	57% (45-70%)	0.47 (0.23-0.86)	-0.64% (-2.1-0.87%)	-0.79% (-2.5-1.1%)	-3 (-10-4.2)	-81 (-270-110)
Namibia	1995	Female	65 to 69	42% (31-52%)	58% (48-69%)	0.42 (0.23-0.75)	-1.1% (-2.4-0.18%)	-1.2% (-2.9-0.31%)	-6 (-15-1.5)	-140 (-330-25)
Namibia	1995	Female	70 to 74	38% (28-49%)	62% (51-72%)	0.35 (0.18-0.62)	-1.1% (-2.4--0.12%)	-1.3% (-2.8--0.084%)	-6.8 (-15--0.42)	-130 (-270--12)
Namibia	1995	Female	75 to 79	34% (23-46%)	66% (54-77%)	0.29 (0.14-0.51)	-1.3% (-2.4--0.3%)	-1.5% (-2.8--0.35%)	-7.4 (-14--1.8)	-100 (-200--25)
Namibia	1995	Female	80 to 84	33% (23-44%)	67% (56-77%)	0.26 (0.14-0.44)	-1.2% (-2.4--0.35%)	-1.4% (-2.7--0.39%)	-5.5 (-11--1.5)	-58 (-110--17)
Namibia	1995	Female	85 to 89	35% (24-47%)	65% (53-76%)	0.27 (0.13-0.45)	-1.3% (-2.5--0.38%)	-1.5% (-2.8--0.45%)	-3.3 (-6.4--0.96)	-24 (-48--7.3)
Namibia	1995	Female	90 to 94	39% (27-52%)	61% (48-73%)	0.27 (0.13-0.48)	-1.3% (-2.6--0.37%)	-1.6% (-3--0.45%)	-1.3 (-2.6--0.36)	-6.4 (-13--1.8)
Namibia	1995	Female	95 plus	44% (28-61%)	56% (39-72%)	0.29 (0.12-0.55)	-1.3% (-2.6--0.27%)	-1.6% (-3.1--0.38%)	-0.33 (-0.66--0.076)	-1.1 (-2.1--0.22)
Namibia	1995	Male	All Ages	38% (34-41%)	62% (59-66%)	0.59 (0.38-0.88)	2.2% (1.1-3.5%)	2.3% (0.84-4.1%)	190 (68-330)	10000 (5200-17000)
Namibia	1995	Male	15 to 19	32% (27-37%)	68% (63-73%)	0.44 (0.27-0.69)	2.6% (1.4-4%)	2.9% (1.1-4.9%)	5.4 (2.1-9.3)	540 (290-830)
Namibia	1995	Male	20 to 24	53% (47-58%)	47% (42-53%)	1.1 (0.7-1.7)	5.5% (3.1-8.8%)	5.6% (2.7-9.9%)	20 (9.7-32)	1600 (950-2500)
Namibia	1995	Male	25 to 29	60% (55-64%)	40% (36-45%)	1.3 (0.83-2)	5% (2.7-8.1%)	5% (2.3-8.9%)	22 (11-36)	1700 (960-2500)
Namibia	1995	Male	30 to 34	62% (57-67%)	38% (33-43%)	1.4 (0.87-2.1)	4.4% (2.3-7.2%)	4.4% (2-8%)	21 (9.6-35)	1400 (730-2200)
Namibia	1995	Male	35 to 39	63% (58-68%)	37% (32-42%)	1.3 (0.8-2)	4.2% (2.1-7.2%)	4.5% (2-8.1%)	19 (8.1-33)	1100 (530-1800)
Namibia	1995	Male	40 to 44	62% (57-67%)	38% (33-43%)	1.2 (0.76-1.9)	4.3% (1.9-7.5%)	4.6% (1.8-8.5%)	19 (7.1-34)	950 (400-1600)
Namibia	1995	Male	45 to 49	59% (54-65%)	41% (35-46%)	1.2 (0.73-1.9)	5% (2.3-8.6%)	5.5% (2.2-9.7%)	22 (8.9-39)	990 (450-1700)
Namibia	1995	Male	50 to 54	59% (53-65%)	41% (35-47%)	1.2 (0.7-1.8)	4.9% (2-8.7%)	5.4% (1.8-9.7%)	22 (6.6-41)	870 (330-1500)
Namibia	1995	Male	55 to 59	62% (55-69%)	38% (31-45%)	1.2 (0.71-1.7)	4.1% (1.2-7.6%)	4.5% (1.1-8.6%)	19 (4.4-38)	660 (190-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	1995	Male	60 to 64	61% (49-70%)	39% (30-51%)	0.99 (0.58-1.5)	2.3% (-0.095-5.2%)	2.4% (-0.36-5.8%)	12 (-1.8-29)	350 (-14-810)
Namibia	1995	Male	65 to 69	53% (41-64%)	47% (36-59%)	0.66 (0.39-1.1)	1.1% (-0.65-3.1%)	1.1% (-0.95-3.4%)	6.2 (-5.4-18)	160 (-98-420)
Namibia	1995	Male	70 to 74	49% (37-60%)	51% (40-63%)	0.42 (0.23-0.7)	0.35% (-0.84-1.5%)	0.31% (-1.1-1.6%)	1.7 (-5.7-8.9)	39 (-92-170)
Namibia	1995	Male	75 to 79	50% (36-65%)	50% (35-64%)	0.32 (0.16-0.56)	0.0076% (-1.1-0.91%)	-0.069% (-1.4-0.96%)	-0.31 (-6-4.2)	0.56 (-78-65)
Namibia	1995	Male	80 to 84	50% (34-65%)	50% (35-66%)	0.28 (0.14-0.47)	-0.055% (-0.99-0.72%)	-0.13% (-1.2-0.76%)	-0.38 (-3.5-2.2)	-1.9 (-34-25)
Namibia	1995	Male	85 to 89	47% (33-62%)	53% (38-67%)	0.23 (0.12-0.39)	-0.062% (-0.82-0.58%)	-0.14% (-1-0.61%)	-0.18 (-1.4-0.8)	-0.69 (-9.2-6.3)
Namibia	1995	Male	90 to 94	46% (32-61%)	54% (39-68%)	0.2 (0.1-0.36)	-0.037% (-0.69-0.5%)	-0.12% (-0.86-0.5%)	-0.051 (-0.37-0.22)	-0.092 (-1.7-1.3)
Namibia	1995	Male	95 plus	49% (30-67%)	51% (33-70%)	0.19 (0.076-0.37)	0.068% (-0.49-0.54%)	-0.047% (-0.71-0.48%)	-0.0051 (-0.078-0.053)	0.029 (-0.21-0.23)
Namibia	2000	Female	All Ages	24% (22-27%)	76% (73-78%)	0.78 (0.52-1.1)	1.6% (0.66-3%)	1.9% (0.47-3.9%)	200 (47-450)	9700 (3500-18000)
Namibia	2000	Female	15 to 19	20% (16-25%)	80% (75-84%)	0.61 (0.35-0.96)	2.1% (1-3.7%)	2.8% (1-5.6%)	4.3 (1.5-8.8)	410 (190-730)
Namibia	2000	Female	20 to 24	29% (24-35%)	71% (65-76%)	1.3 (0.73-2)	2.7% (1.1-5%)	2.8% (0.81-6%)	12 (4.1-24)	1000 (460-1900)
Namibia	2000	Female	25 to 29	34% (30-40%)	66% (60-70%)	1.5 (0.9-2.2)	1.8% (0.65-3.6%)	1.8% (0.47-4%)	15 (4.8-30)	1100 (470-2000)
Namibia	2000	Female	30 to 34	37% (31-42%)	63% (58-69%)	1.6 (0.96-2.4)	1.9% (0.71-3.9%)	2% (0.53-4.4%)	17 (5.3-37)	1100 (430-2200)
Namibia	2000	Female	35 to 39	39% (34-44%)	61% (56-66%)	1.7 (1.1-2.6)	2.5% (0.86-5%)	2.7% (0.77-5.7%)	23 (6.9-46)	1300 (450-2400)
Namibia	2000	Female	40 to 44	41% (36-47%)	59% (53-64%)	1.8 (1-2.7)	3.2% (1.2-6%)	3.5% (1.2-6.9%)	25 (8-49)	1200 (440-2300)
Namibia	2000	Female	45 to 49	41% (35-48%)	59% (52-65%)	1.6 (0.96-2.6)	3.6% (1.3-7.1%)	4.2% (1.4-8.3%)	26 (7.3-54)	1100 (350-2200)
Namibia	2000	Female	50 to 54	41% (33-49%)	59% (51-67%)	1.4 (0.82-2.3)	3.9% (1.3-7.8%)	4.5% (1.4-9.2%)	24 (6.6-52)	910 (260-1900)
Namibia	2000	Female	55 to 59	43% (34-53%)	57% (47-66%)	1.4 (0.77-2.3)	3.5% (0.61-7.6%)	4.2% (0.71-9%)	21 (3.2-51)	680 (110-1600)
Namibia	2000	Female	60 to 64	41% (32-50%)	59% (50-68%)	1.2 (0.63-2.1)	2.7% (-0.15-7.1%)	3.2% (-0.22-8.3%)	16 (-0.93-47)	440 (-25-1300)
Namibia	2000	Female	65 to 69	35% (26-44%)	65% (56-74%)	0.93 (0.5-1.6)	1.8% (-0.63-5%)	2.2% (-0.75-6.1%)	13 (-3.8-39)	280 (-93-850)
Namibia	2000	Female	70 to 74	31% (23-41%)	69% (59-77%)	0.77 (0.42-1.3)	0.96% (-1.1-3.9%)	1.2% (-1.3-4.7%)	7.7 (-9-35)	140 (-160-630)
Namibia	2000	Female	75 to 79	30% (21-40%)	70% (60-79%)	0.68 (0.35-1.2)	0.23% (-1.5-2.7%)	0.29% (-1.8-3.2%)	1.8 (-12-21)	23 (-170-290)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	2000	Female	80 to 84	32% (22-42%)	68% (58-78%)	0.67 (0.36-1.1)	-0.087% (-1.8-2.1%)	-0.087% (-2.1-2.5%)	-0.45 (-11-12)	-5.2 (-110-120)
Namibia	2000	Female	85 to 89	35% (24-47%)	65% (53-76%)	0.7 (0.36-1.2)	-0.45% (-2.1-1.5%)	-0.53% (-2.4-1.7%)	-1.4 (-6.9-4.8)	-10 (-49-35)
Namibia	2000	Female	90 to 94	38% (27-50%)	62% (50-73%)	0.72 (0.37-1.3)	-0.73% (-2.3-1.2%)	-0.9% (-2.8-1.4%)	-0.88 (-2.7-1.4)	-4.2 (-13-7.1)
Namibia	2000	Female	95 plus	44% (27-62%)	56% (38-73%)	0.75 (0.33-1.4)	-0.84% (-2.6-1.4%)	-1.1% (-3.1-1.5%)	-0.27 (-0.8-0.41)	-0.82 (-2.6-1.4)
Namibia	2000	Male	All Ages	41% (37-45%)	59% (55-63%)	1.3 (0.94-1.9)	4.7% (2.9-6.7%)	5.5% (3.3-8.3%)	680 (390-1000)	32000 (20000-47000)
Namibia	2000	Male	15 to 19	36% (30-43%)	64% (57-70%)	1 (0.57-1.6)	5.2% (3.7-9.9%)	6.4% (3.3-10%)	15 (7.5-25)	1300 (700-2000)
Namibia	2000	Male	20 to 24	57% (50-63%)	43% (37-50%)	2.3 (1.4-3.6)	9.1% (5.3-14%)	10% (5.2-16%)	51 (29-77)	3800 (2300-5600)
Namibia	2000	Male	25 to 29	63% (57-68%)	37% (32-43%)	2.9 (1.7-4.3)	6.9% (3.7-11%)	7.2% (3.6-13%)	64 (36-97)	4300 (2600-6200)
Namibia	2000	Male	30 to 34	66% (61-72%)	34% (28-39%)	3.1 (1.9-4.6)	6.3% (3.3-9.7%)	6.5% (3.2-10%)	67 (38-100)	4000 (2400-6000)
Namibia	2000	Male	35 to 39	66% (61-71%)	34% (29-39%)	2.9 (1.7-4.4)	6.6% (3.5-11%)	7% (3.5-12%)	70 (38-110)	3700 (2100-5800)
Namibia	2000	Male	40 to 44	65% (59-70%)	35% (30-41%)	2.8 (1.6-4.3)	7.3% (3.9-12%)	7.9% (4.1-13%)	71 (38-120)	3400 (1900-5400)
Namibia	2000	Male	45 to 49	61% (54-67%)	39% (33-46%)	2.6 (1.5-4.1)	8.8% (4.7-14%)	9.6% (4.9-16%)	74 (38-120)	3100 (1700-5100)
Namibia	2000	Male	50 to 54	60% (53-67%)	40% (33-47%)	2.5 (1.4-4)	10% (5.1-16%)	11% (5.5-18%)	76 (36-130)	2900 (1400-4700)
Namibia	2000	Male	55 to 59	65% (59-72%)	35% (28-41%)	2.6 (1.6-4.2)	10% (5.5-16%)	12% (5.9-19%)	77 (38-130)	2500 (1300-4100)
Namibia	2000	Male	60 to 64	63% (55-71%)	37% (29-45%)	2.2 (1.3-3.6)	7.8% (3.5-14%)	8.6% (3.7-15%)	58 (24-110)	1600 (690-2900)
Namibia	2000	Male	65 to 69	52% (42-62%)	48% (38-58%)	1.4 (0.84-2.3)	4.7% (1.5-9.1%)	5.3% (1.5-10%)	37 (11-73)	830 (260-1600)
Namibia	2000	Male	70 to 74	48% (37-58%)	52% (42-63%)	0.91 (0.53-1.5)	2% (-0.037-4.7%)	2.2% (-0.11-5.2%)	16 (-0.82-39)	290 (-5.1-720)
Namibia	2000	Male	75 to 79	51% (38-64%)	49% (36-62%)	0.74 (0.39-1.2)	0.74% (-0.93-2.9%)	0.78% (-1.1-3.2%)	4.6 (-6.7-18)	69 (-86-260)
Namibia	2000	Male	80 to 84	51% (35-65%)	49% (35-65%)	0.65 (0.34-1.1)	0.36% (-1-2%)	0.35% (-1.3-2.2%)	1.3 (-4.8-8.2)	16 (-46-87)
Namibia	2000	Male	85 to 89	47% (32-62%)	53% (38-68%)	0.53 (0.27-0.88)	0.12% (-0.95-1.2%)	0.065% (-1.2-1.3%)	0.11 (-1.9-2.1)	1.7 (-13-17)
Namibia	2000	Male	90 to 94	47% (33-61%)	53% (39-67%)	0.46 (0.23-0.81)	0.0097% (-0.88-0.9%)	-0.09% (-1.1-0.95%)	-0.044 (-0.56-0.47)	0.03 (-2.6-2.6)
Namibia	2000	Male	95 plus	49% (30-68%)	51% (32-70%)	0.42 (0.19-0.79)	0.15% (-0.7-1%)	0.013% (-0.98-1%)	0.0015 (-0.11-0.12)	0.07 (-0.32-0.46)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	2005	Female	All Ages	25% (22-27%)	75% (73-78%)	1.2 (0.81-1.6)	1.8% (0.9-3.2%)	2.4% (0.98-4.4%)	310 (120-620)	13000 (6200-23000)
Namibia	2005	Female	15 to 19	21% (17-24%)	79% (76-83%)	0.96 (0.58-1.4)	2.9% (1.6-4.8%)	4.2% (1.8-7.4%)	6.3 (2.7-12)	600 (310-1000)
Namibia	2005	Female	20 to 24	31% (28-36%)	69% (64-72%)	2 (1.2-2.9)	3.4% (1.5-6.2%)	3.6% (1.2-7.7%)	15 (6.4-28)	1200 (660-2100)
Namibia	2005	Female	25 to 29	35% (31-39%)	65% (61-69%)	2.2 (1.4-3.2)	1.6% (0.69-3.2%)	1.5% (0.53-3.4%)	16 (6.9-33)	1200 (610-2200)
Namibia	2005	Female	30 to 34	36% (32-40%)	64% (60-68%)	2.2 (1.5-3.2)	1.4% (0.62-2.8%)	1.3% (0.51-3%)	18 (7.5-37)	1200 (580-2200)
Namibia	2005	Female	35 to 39	38% (34-43%)	62% (57-66%)	2.5 (1.6-3.7)	1.8% (0.81-3.6%)	1.9% (0.73-3.9%)	23 (10-46)	1300 (640-2500)
Namibia	2005	Female	40 to 44	41% (36-45%)	59% (55-64%)	2.6 (1.6-3.8)	2.5% (1.2-4.6%)	2.6% (1.1-5.1%)	29 (13-56)	1400 (670-2700)
Namibia	2005	Female	45 to 49	40% (35-45%)	60% (55-65%)	2.3 (1.4-3.5)	3.3% (1.5-6.2%)	3.6% (1.5-6.9%)	31 (13-62)	1400 (620-2600)
Namibia	2005	Female	50 to 54	40% (33-48%)	60% (52-67%)	2.1 (1.2-3.3)	4.3% (1.9-7.8%)	4.9% (2.1-9%)	34 (14-70)	1300 (540-2600)
Namibia	2005	Female	55 to 59	43% (34-52%)	57% (48-66%)	2 (1.1-3.3)	4.8% (1.9-8.9%)	5.7% (2.1-10%)	32 (11-67)	1000 (370-2100)
Namibia	2005	Female	60 to 64	40% (31-50%)	60% (50-69%)	1.7 (0.91-2.9)	4.6% (1.1-9.6%)	5.6% (1.3-11%)	30 (6.6-69)	830 (180-1900)
Namibia	2005	Female	65 to 69	34% (25-43%)	66% (57-75%)	1.3 (0.74-2.2)	3.7% (0.79-7.9%)	4.5% (0.99-9.6%)	25 (4.9-59)	570 (100-1300)
Namibia	2005	Female	70 to 74	31% (23-40%)	69% (60-77%)	1.1 (0.63-1.9)	2.7% (0.078-6.4%)	3.3% (0.095-7.5%)	20 (0.59-51)	360 (8.8-920)
Namibia	2005	Female	75 to 79	29% (21-40%)	71% (60-79%)	0.97 (0.51-1.6)	1.8% (-0.62-5.1%)	2.1% (-0.7-6%)	13 (-4.2-41)	190 (-60-570)
Namibia	2005	Female	80 to 84	31% (22-41%)	69% (59-78%)	0.96 (0.53-1.6)	1.4% (-0.76-4.5%)	1.7% (-0.93-5.2%)	8.9 (-4.6-30)	92 (-48-320)
Namibia	2005	Female	85 to 89	34% (23-46%)	66% (54-77%)	1 (0.54-1.7)	0.97% (-1.1-4%)	1.1% (-1.3-4.7%)	2.9 (-3.4-12)	22 (-25-91)
Namibia	2005	Female	90 to 94	38% (26-51%)	62% (49-74%)	1 (0.55-1.8)	0.55% (-1.5-3.7%)	0.6% (-1.8-4.3%)	0.57 (-1.8-4.2)	3.1 (-8.4-21)
Namibia	2005	Female	95 plus	43% (27-60%)	57% (40-73%)	1.1 (0.49-2)	0.45% (-1.9-4.5%)	0.44% (-2.4-5.1%)	0.11 (-0.56-1.2)	0.42 (-1.8-4)
Namibia	2005	Male	All Ages	42% (39-46%)	58% (54-61%)	2.5 (1.8-3.5)	6.8% (4.5-9.3%)	8.2% (5.2-12%)	1200 (730-1700)	52000 (34000-71000)
Namibia	2005	Male	15 to 19	38% (33-44%)	62% (56-67%)	2.1 (1.4-3)	9.2% (6.1-13%)	12% (7.5-17%)	31 (19-45)	2400 (1600-3500)
Namibia	2005	Male	20 to 24	60% (54-65%)	40% (35-46%)	4.7 (3.2-6.5)	16% (10-22%)	19% (11-27%)	85 (56-120)	6100 (4200-8300)
Namibia	2005	Male	25 to 29	64% (59-69%)	36% (31-41%)	5.4 (3.7-7.5)	10% (6-16%)	11% (5.9-18%)	100 (66-130)	6500 (4500-8600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	2005	Male	30 to 34	67% (62-72%)	33% (28-38%)	5.6 (3.8-8)	7.6% (4.6-12%)	7.9% (4.6-12%)	100 (68-140)	6100 (4100-8200)
Namibia	2005	Male	35 to 39	67% (62-71%)	33% (29-38%)	5.2 (3.5-7.5)	7.7% (4.6-12%)	8.2% (4.6-13%)	110 (66-150)	5700 (3600-7800)
Namibia	2005	Male	40 to 44	66% (60-70%)	34% (30-40%)	5 (3.3-7.2)	8.7% (5.4-13%)	9.2% (5.6-14%)	120 (72-170)	5500 (3500-7800)
Namibia	2005	Male	45 to 49	62% (56-67%)	38% (33-44%)	4.8 (3.2-6.9)	11% (6.8-16%)	12% (7.2-17%)	120 (75-180)	5200 (3200-7500)
Namibia	2005	Male	50 to 54	60% (52-66%)	40% (34-48%)	4.4 (2.9-6.4)	14% (9-19%)	15% (9.9-21%)	120 (73-180)	4600 (2800-6600)
Namibia	2005	Male	55 to 59	64% (57-71%)	36% (29-43%)	4.5 (3.1-6.6)	16% (11-22%)	18% (12-24%)	130 (75-180)	4100 (2500-5900)
Namibia	2005	Male	60 to 64	63% (54-70%)	37% (30-46%)	3.8 (2.6-5.5)	14% (8.4-19%)	15% (9.3-22%)	110 (61-170)	3000 (1700-4600)
Namibia	2005	Male	65 to 69	52% (42-62%)	48% (38-58%)	2.4 (1.6-3.7)	9.3% (5-15%)	10% (5.6-17%)	74 (37-120)	1700 (840-2800)
Namibia	2005	Male	70 to 74	47% (37-58%)	53% (42-63%)	1.6 (0.93-2.5)	4.9% (1.8-9.1%)	5.4% (2-10%)	38 (14-73)	700 (260-1300)
Namibia	2005	Male	75 to 79	50% (37-64%)	50% (36-63%)	1.3 (0.67-2.1)	2.8% (0.34-6.2%)	3.1% (0.29-6.8%)	20 (1.9-46)	280 (34-640)
Namibia	2005	Male	80 to 84	50% (35-65%)	50% (35-65%)	1.1 (0.62-1.9)	2.1% (0.0054-5%)	2.3% (-0.086-5.5%)	9.8 (-0.38-24)	110 (0.28-250)
Namibia	2005	Male	85 to 89	46% (31-61%)	54% (39-69%)	0.94 (0.5-1.6)	1.3% (-0.21-3.4%)	1.4% (-0.36-3.9%)	2.7 (-0.63-7.1)	21 (-3.3-54)
Namibia	2005	Male	90 to 94	46% (32-60%)	54% (40-68%)	0.83 (0.44-1.4)	0.9% (-0.35-2.6%)	0.93% (-0.54-2.9%)	0.52 (-0.28-1.6)	3 (-1.1-8.5)
Namibia	2005	Male	95 plus	48% (30-67%)	52% (33-70%)	0.77 (0.33-1.4)	0.98% (-0.27-3%)	0.95% (-0.5-3.3%)	0.12 (-0.06-0.41)	0.48 (-0.13-1.5)
Namibia	2010	Female	All Ages	27% (24-30%)	73% (70-76%)	1.6 (0.99-2.3)	2.1% (1-3.6%)	3% (1.2-5.5%)	300 (110-610)	12000 (5700-21000)
Namibia	2010	Female	15 to 19	20% (17-24%)	80% (76-83%)	1.2 (0.67-1.7)	2.7% (1.4-4.5%)	3.7% (1.6-6.6%)	5.8 (2.3-12)	590 (300-1000)
Namibia	2010	Female	20 to 24	35% (31-39%)	65% (61-69%)	2.7 (1.6-4.3)	4.2% (2-7.1%)	4.7% (1.6-9.4%)	13 (5.3-25)	1200 (660-2000)
Namibia	2010	Female	25 to 29	39% (35-44%)	61% (56-65%)	3 (1.8-4.5)	2.2% (0.94-4%)	2.1% (0.65-4.5%)	13 (5.1-24)	1100 (540-1800)
Namibia	2010	Female	30 to 34	39% (34-44%)	61% (56-66%)	2.9 (1.8-4.4)	1.6% (0.74-2.9%)	1.5% (0.54-3.1%)	14 (5.6-26)	990 (500-1700)
Namibia	2010	Female	35 to 39	40% (35-45%)	60% (55-65%)	3.2 (1.9-4.9)	1.9% (0.91-3.3%)	1.9% (0.76-3.6%)	18 (7.4-32)	1100 (530-1800)
Namibia	2010	Female	40 to 44	41% (36-46%)	59% (54-64%)	3.2 (1.9-5)	2.7% (1.3-4.6%)	2.9% (1.2-5.1%)	23 (9.7-40)	1200 (560-2000)
Namibia	2010	Female	45 to 49	41% (36-47%)	59% (53-64%)	2.9 (1.6-4.7)	3.7% (1.7-6.5%)	4.2% (1.7-7.8%)	27 (11-50)	1200 (550-2200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	2010	Female	50 to 54	42% (32-52%)	58% (48-68%)	2.7 (1.4-4.4)	5% (2.2-8.8%)	6% (2.6-11%)	31 (12-62)	1200 (500-2300)
Namibia	2010	Female	55 to 59	43% (31-55%)	57% (45-69%)	2.4 (1.2-4.2)	5.8% (2.2-11%)	7.4% (2.8-13%)	32 (9.9-67)	1100 (350-2100)
Namibia	2010	Female	60 to 64	42% (30-54%)	58% (46-70%)	2.2 (1-3.8)	5.8% (1.5-11%)	7.5% (2-14%)	31 (6.9-68)	870 (210-1900)
Namibia	2010	Female	65 to 69	41% (30-51%)	59% (49-70%)	1.9 (1-3.3)	5.4% (1.3-11%)	7% (1.7-14%)	31 (7.1-72)	720 (160-1600)
Namibia	2010	Female	70 to 74	37% (27-47%)	63% (53-73%)	1.6 (0.83-2.8)	4.1% (0.55-9.1%)	5.2% (0.78-11%)	25 (3.4-63)	460 (58-1200)
Namibia	2010	Female	75 to 79	33% (22-44%)	67% (56-78%)	1.3 (0.64-2.3)	2.8% (-0.27-7%)	3.5% (-0.33-8.9%)	18 (-1.8-50)	250 (-24-720)
Namibia	2010	Female	80 to 84	32% (22-43%)	68% (57-78%)	1.2 (0.62-2.1)	2.3% (-0.39-6.1%)	2.8% (-0.51-7.5%)	13 (-2.4-38)	140 (-25-410)
Namibia	2010	Female	85 to 89	34% (23-46%)	66% (54-77%)	1.2 (0.62-2.2)	1.8% (-0.78-5.8%)	2.1% (-0.97-6.9%)	6 (-2.8-21)	45 (-20-150)
Namibia	2010	Female	90 to 94	38% (26-50%)	62% (50-74%)	1.2 (0.62-2.3)	1.4% (-1.2-5.3%)	1.6% (-1.6-6.4%)	1.5 (-1.5-5.9)	7.8 (-7.3-31)
Namibia	2010	Female	95 plus	43% (27-61%)	57% (39-73%)	1.3 (0.55-2.6)	1.3% (-1.6-6.4%)	1.4% (-2-7.5%)	0.34 (-0.47-1.7)	1.2 (-1.4-5.9)
Namibia	2010	Male	All Ages	46% (42-49%)	54% (51-58%)	3.2 (2-4.8)	7.8% (5.3-10%)	9.8% (6.4-13%)	1100 (710-1500)	50000 (34000-66000)
Namibia	2010	Male	15 to 19	40% (35-44%)	60% (56-65%)	2.5 (1.5-3.8)	9.2% (5.9-13%)	12% (7-18%)	34 (20-51)	2700 (1700-4000)
Namibia	2010	Male	20 to 24	64% (61-68%)	36% (32-39%)	6 (3.7-9)	19% (13-25%)	22% (14-31%)	91 (60-120)	6600 (4500-8700)
Namibia	2010	Male	25 to 29	68% (65-72%)	32% (28-35%)	6.8 (4.2-10)	13% (8.1-19%)	15% (8.5-23%)	93 (63-120)	6200 (4300-8100)
Namibia	2010	Male	30 to 34	69% (65-72%)	31% (28-35%)	6.8 (4.2-10)	9.4% (6.1-14%)	10% (6.3-15%)	91 (62-120)	5500 (3900-7100)
Namibia	2010	Male	35 to 39	69% (65-72%)	31% (28-35%)	6.3 (3.9-9.8)	9.1% (5.9-13%)	9.9% (6.2-15%)	96 (65-130)	5200 (3600-6700)
Namibia	2010	Male	40 to 44	67% (63-71%)	33% (29-37%)	6 (3.8-9.3)	10% (7.1-14%)	12% (7.7-16%)	100 (68-140)	4900 (3300-6500)
Namibia	2010	Male	45 to 49	63% (59-68%)	37% (32-41%)	5.7 (3.6-9)	13% (9.2-18%)	15% (10-21%)	110 (75-160)	4800 (3300-6500)
Namibia	2010	Male	50 to 54	61% (54-67%)	39% (33-46%)	5.4 (3.3-8.5)	16% (11-22%)	19% (13-25%)	110 (72-160)	4300 (2800-5900)
Namibia	2010	Male	55 to 59	67% (61-73%)	33% (27-39%)	5.6 (3.4-8.6)	19% (13-24%)	22% (15-29%)	110 (71-160)	3700 (2400-5200)
Namibia	2010	Male	60 to 64	67% (60-72%)	33% (28-40%)	4.8 (3-7.4)	16% (9.9-23%)	19% (11-27%)	110 (59-160)	2900 (1600-4400)
Namibia	2010	Male	65 to 69	55% (45-65%)	45% (35-55%)	3.1 (1.8-4.8)	11% (5.8-18%)	13% (6.6-20%)	77 (37-130)	1800 (860-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	2010	Male	70 to 74	48% (37-59%)	52% (41-63%)	1.9 (1-3.2)	5.8% (2.2-11%)	6.6% (2.4-12%)	39 (14-75)	730 (260-1400)
Namibia	2010	Male	75 to 79	49% (35-63%)	51% (37-65%)	1.5 (0.72-2.6)	3.4% (0.58-7.6%)	3.8% (0.61-8.6%)	21 (3.3-47)	300 (48-670)
Namibia	2010	Male	80 to 84	49% (33-64%)	51% (36-67%)	1.3 (0.64-2.2)	2.5% (0.25-5.9%)	2.8% (0.17-6.7%)	12 (0.75-28)	130 (12-300)
Namibia	2010	Male	85 to 89	46% (32-61%)	54% (39-68%)	1.1 (0.57-1.9)	1.7% (-0.01-4.3%)	1.9% (-0.12-4.9%)	3.9 (-0.27-9.9)	31 (-0.19-75)
Namibia	2010	Male	90 to 94	45% (31-60%)	55% (40-69%)	0.96 (0.48-1.8)	1.2% (-0.15-3.6%)	1.3% (-0.29-4%)	0.81 (-0.18-2.5)	4.5 (-0.54-13)
Namibia	2010	Male	95 plus	48% (29-67%)	52% (33-71%)	0.9 (0.37-1.8)	1.3% (-0.12-3.9%)	1.3% (-0.35-4.4%)	0.17 (-0.045-0.58)	0.69 (-0.063-2.1)
Namibia	2016	Female	All Ages	28% (25-31%)	72% (69-75%)	1.6 (0.81-2.9)	2.6% (1-4.7%)	3.7% (1-7.5%)	290 (68-630)	12000 (4400-22000)
Namibia	2016	Female	15 to 19	19% (15-25%)	81% (75-85%)	1 (0.47-2)	2.4% (1.2-4.1%)	3.3% (1.1-6.6%)	4.2 (1.3-8.7)	470 (210-850)
Namibia	2016	Female	20 to 24	35% (28-41%)	65% (59-72%)	2.7 (1.2-4.8)	4.2% (2.1-7%)	4.9% (1.5-9.6%)	11 (3.6-21)	1100 (550-1900)
Namibia	2016	Female	25 to 29	42% (34-47%)	58% (53-66%)	3.1 (1.5-5.6)	3.1% (1.3-5.4%)	3.2% (0.82-6.7%)	12 (3.7-23)	1100 (510-1800)
Namibia	2016	Female	30 to 34	41% (34-47%)	59% (53-66%)	3.1 (1.5-5.6)	2.4% (1-4.4%)	2.5% (0.66-5.3%)	12 (3.8-25)	940 (430-1700)
Namibia	2016	Female	35 to 39	42% (35-49%)	58% (51-65%)	3.2 (1.6-6)	2.7% (1.1-4.7%)	3% (0.86-5.9%)	16 (4.8-31)	1000 (410-1800)
Namibia	2016	Female	40 to 44	42% (36-50%)	58% (50-64%)	3.2 (1.5-5.8)	3.5% (1.3-6.4%)	4.2% (1.3-8.2%)	21 (6.2-42)	1100 (420-2100)
Namibia	2016	Female	45 to 49	43% (35-49%)	57% (51-65%)	2.9 (1.3-5.4)	4.6% (1.8-8.5%)	5.9% (2-11%)	26 (7.5-51)	1200 (410-2300)
Namibia	2016	Female	50 to 54	42% (33-52%)	58% (48-67%)	2.6 (1.1-4.9)	5.5% (1.8-10%)	7.3% (2.5-14%)	30 (8.1-63)	1200 (360-2400)
Namibia	2016	Female	55 to 59	42% (30-54%)	58% (46-70%)	2.3 (0.95-4.5)	6% (1.8-12%)	8.6% (2.4-16%)	31 (6.6-70)	1000 (250-2300)
Namibia	2016	Female	60 to 64	41% (29-54%)	59% (46-71%)	2.1 (0.84-4)	5.6% (0.91-12%)	7.9% (1.3-16%)	31 (4.3-74)	870 (130-2100)
Namibia	2016	Female	65 to 69	40% (29-51%)	60% (49-71%)	1.9 (0.83-3.5)	5.1% (0.41-11%)	7% (0.59-15%)	30 (2.2-80)	700 (56-1800)
Namibia	2016	Female	70 to 74	37% (27-47%)	63% (53-73%)	1.6 (0.7-3.1)	3.8% (-0.26-9.7%)	5.1% (-0.3-13%)	24 (-1.4-68)	450 (-26-1300)
Namibia	2016	Female	75 to 79	33% (23-45%)	67% (55-77%)	1.3 (0.54-2.5)	2.7% (-0.74-7.8%)	3.4% (-0.96-9.8%)	17 (-4.4-53)	240 (-65-780)
Namibia	2016	Female	80 to 84	32% (22-43%)	68% (57-78%)	1.2 (0.5-2.2)	2.1% (-0.83-6.9%)	2.7% (-1-8.7%)	12 (-4.6-41)	130 (-49-420)
Namibia	2016	Female	85 to 89	34% (23-46%)	66% (54-77%)	1.2 (0.5-2.4)	1.7% (-1.1-6.5%)	2.1% (-1.4-7.7%)	6.9 (-4.9-27)	52 (-34-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Namibia	2016	Female	90 to 94	38% (26-50%)	62% (50-74%)	1.2 (0.52-2.5)	1.3% (-1.4-6.1%)	1.6% (-1.8-7.1%)	2.2 (-2.8-10)	12 (-13-53)
Namibia	2016	Female	95 plus	43% (27-60%)	57% (40-73%)	1.3 (0.47-2.8)	1.3% (-1.7-7.3%)	1.4% (-2.2-8.6%)	0.44 (-0.73-2.7)	1.6 (-2.3-9)
Namibia	2016	Male	All Ages	46% (43-50%)	54% (50-57%)	3.6 (1.8-6.2)	8.9% (5.5-12%)	11% (6.5-15%)	1100 (640-1600)	51000 (31000-71000)
Namibia	2016	Male	15 to 19	35% (30-40%)	65% (60-70%)	2.4 (1.1-4.2)	8.4% (4.9-13%)	11% (5.9-17%)	30 (15-48)	2400 (1400-3700)
Namibia	2016	Male	20 to 24	64% (58-68%)	36% (32-42%)	6.4 (3.1-11)	18% (12-25%)	22% (14-31%)	95 (55-130)	7000 (4400-9500)
Namibia	2016	Male	25 to 29	70% (65-73%)	30% (27-35%)	7.5 (3.7-13)	16% (10-21%)	18% (11-26%)	100 (59-140)	6800 (4200-9000)
Namibia	2016	Male	30 to 34	69% (64-73%)	31% (27-36%)	7.3 (3.7-13)	11% (7.2-16%)	13% (7.5-19%)	88 (53-120)	5400 (3400-7300)
Namibia	2016	Male	35 to 39	69% (65-73%)	31% (27-35%)	6.8 (3.4-12)	10% (6.3-14%)	11% (6.8-17%)	91 (54-130)	5000 (3100-6900)
Namibia	2016	Male	40 to 44	67% (62-71%)	33% (29-38%)	6.5 (3.2-11)	11% (6.7-15%)	12% (7.4-17%)	99 (58-140)	4800 (2900-6800)
Namibia	2016	Male	45 to 49	63% (57-68%)	37% (32-43%)	6.1 (3-11)	13% (8.4-18%)	15% (9.5-21%)	110 (67-160)	4800 (3000-6700)
Namibia	2016	Male	50 to 54	60% (54-66%)	40% (34-46%)	5.7 (2.8-10)	15% (9.5-20%)	18% (11-24%)	120 (68-170)	4500 (2700-6400)
Namibia	2016	Male	55 to 59	67% (60-73%)	33% (27-40%)	6 (3-10)	18% (11-24%)	21% (13-28%)	120 (70-180)	4100 (2400-5700)
Namibia	2016	Male	60 to 64	67% (60-72%)	33% (28-40%)	5.2 (2.6-9)	16% (8.2-23%)	19% (9.6-27%)	110 (55-170)	3000 (1600-4600)
Namibia	2016	Male	65 to 69	55% (44-65%)	45% (35-56%)	3.3 (1.6-6.2)	11% (4.9-18%)	13% (5.7-22%)	82 (34-140)	1900 (800-3100)
Namibia	2016	Male	70 to 74	48% (36-60%)	52% (40-64%)	2 (0.91-3.8)	6.2% (1.7-12%)	7.1% (1.9-14%)	46 (12-92)	850 (230-1700)
Namibia	2016	Male	75 to 79	49% (34-63%)	51% (37-66%)	1.6 (0.67-3)	3.8% (0.47-9.2%)	4.2% (0.45-11%)	24 (2.6-60)	350 (44-860)
Namibia	2016	Male	80 to 84	49% (33-63%)	51% (37-67%)	1.4 (0.59-2.6)	2.9% (0.2-7.3%)	3.2% (0.11-8.2%)	14 (0.52-36)	150 (11-380)
Namibia	2016	Male	85 to 89	46% (31-61%)	54% (39-69%)	1.2 (0.5-2.3)	2% (-0.036-5.6%)	2.2% (-0.17-6.4%)	5.4 (-0.42-15)	42 (-0.78-120)
Namibia	2016	Male	90 to 94	45% (31-60%)	55% (40-69%)	1 (0.42-2.1)	1.5% (-0.18-4.7%)	1.6% (-0.33-5.4%)	1.4 (-0.28-4.5)	7.6 (-0.91-24)
Namibia	2016	Male	95 plus	47% (29-66%)	53% (34-71%)	0.96 (0.34-2.1)	1.6% (-0.066-5%)	1.6% (-0.27-5.6%)	0.28 (-0.046-0.94)	1.1 (-0.046-3.3)
South Africa	1990	Female	All Ages	21% (18-25%)	79% (75-82%)	1.7 (1.2-2.3)	4.7% (3.7-5.9%)	7.3% (5.5-9.2%)	9300 (7000-12000)	380000 (300000-470000)
South Africa	1990	Female	15 to 19	17% (11-24%)	83% (76-89%)	1.2 (0.6-2.2)	4.2% (2.4-6.3%)	7.1% (3.9-11%)	150 (83-230)	13000 (7700-19000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	1990	Female	20 to 24	22% (15-32%)	78% (68-85%)	2.2 (1.1-3.7)	7.7% (4.9-11%)	10% (6.3-14%)	540 (330-780)	41000 (26000-58000)
South Africa	1990	Female	25 to 29	26% (17-36%)	74% (64-83%)	2.7 (1.5-4.4)	9.4% (6.3-13%)	12% (7.7-16%)	850 (550-1200)	57000 (38000-76000)
South Africa	1990	Female	30 to 34	29% (19-41%)	71% (59-81%)	3.1 (1.7-4.9)	9.6% (6.7-13%)	12% (8.6-17%)	940 (640-1300)	57000 (40000-77000)
South Africa	1990	Female	35 to 39	32% (22-45%)	68% (55-78%)	3.1 (1.8-5)	9.5% (6.8-13%)	13% (9.3-18%)	730 (510-980)	41000 (29000-54000)
South Africa	1990	Female	40 to 44	34% (24-47%)	66% (53-76%)	3.1 (1.6-5.1)	9.4% (6.4-13%)	14% (9.3-18%)	620 (420-830)	31000 (22000-42000)
South Africa	1990	Female	45 to 49	34% (22-46%)	66% (54-78%)	3.2 (1.6-5.8)	9.5% (6.4-13%)	14% (9.3-18%)	560 (380-760)	25000 (17000-34000)
South Africa	1990	Female	50 to 54	33% (23-45%)	67% (55-77%)	3.3 (1.7-5.7)	9.2% (6.1-13%)	13% (8.8-18%)	500 (330-690)	20000 (13000-27000)
South Africa	1990	Female	55 to 59	33% (23-45%)	67% (55-77%)	3.4 (1.7-5.8)	9.1% (5.9-13%)	13% (8.3-18%)	630 (400-880)	21000 (14000-30000)
South Africa	1990	Female	60 to 64	34% (23-46%)	66% (54-77%)	3.4 (1.7-6)	8.5% (5.2-12%)	12% (7.1-17%)	720 (440-1000)	21000 (13000-29000)
South Africa	1990	Female	65 to 69	35% (25-46%)	65% (54-75%)	3.1 (1.7-5.4)	8.3% (5.2-12%)	11% (7.2-16%)	900 (570-1300)	22000 (14000-31000)
South Africa	1990	Female	70 to 74	33% (24-44%)	67% (56-76%)	2.7 (1.4-4.5)	6.7% (3.8-10%)	9% (5-13%)	550 (310-820)	11000 (6100-16000)
South Africa	1990	Female	75 to 79	29% (20-40%)	71% (60-80%)	2.2 (1.1-3.8)	5.2% (2.5-8.2%)	6.8% (3.3-11%)	570 (280-920)	8800 (4200-14000)
South Africa	1990	Female	80 to 84	28% (19-39%)	72% (61-81%)	2 (0.97-3.4)	4.8% (2.3-7.8%)	6.1% (2.9-9.8%)	610 (290-1000)	6800 (3200-11000)
South Africa	1990	Female	85 to 89	28% (19-39%)	72% (61-81%)	1.9 (0.96-3.4)	4.5% (2-7.5%)	5.5% (2.4-9.2%)	320 (140-530)	2400 (1100-4000)
South Africa	1990	Female	90 to 94	30% (20-42%)	70% (58-80%)	1.9 (0.91-3.6)	4.2% (1.7-7.3%)	5.1% (2-9%)	110 (41-190)	560 (230-980)
South Africa	1990	Female	95 plus	34% (21-48%)	66% (52-79%)	2 (0.8-4.2)	4.1% (1.2-8.2%)	4.9% (1.2-10%)	23 (6-48)	80 (23-160)
South Africa	1990	Male	All Ages	31% (27-36%)	69% (64-73%)	2.4 (1.7-3.3)	8.4% (6.8-10%)	11% (8.8-13%)	18000 (14000-21000)	810000 (650000-960000)
South Africa	1990	Male	15 to 19	25% (17-37%)	75% (63-83%)	1.4 (0.68-2.5)	5.7% (3.5-8.9%)	8% (4.4-13%)	240 (130-380)	21000 (12000-31000)
South Africa	1990	Male	20 to 24	37% (25-50%)	63% (50-75%)	3.1 (1.7-5.1)	13% (8.5-17%)	15% (9.9-22%)	950 (600-1300)	70000 (47000-96000)
South Africa	1990	Male	25 to 29	43% (30-57%)	57% (43-70%)	4.9 (2.7-7.5)	17% (13-22%)	21% (15-27%)	1800 (1300-2300)	120000 (85000-150000)
South Africa	1990	Male	30 to 34	46% (32-60%)	54% (40-68%)	5.6 (3.3-8.6)	19% (14-24%)	22% (16-28%)	2300 (1700-3000)	140000 (100000-170000)
South Africa	1990	Male	35 to 39	48% (33-61%)	52% (39-67%)	5.5 (3.2-8.7)	19% (14-23%)	22% (17-28%)	2200 (1700-2800)	120000 (92000-150000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	1990	Male	40 to 44	47% (34-61%)	53% (39-66%)	5 (2.6-8)	18% (13-23%)	21% (16-27%)	1900 (1400-2400)	91000 (68000-110000)
South Africa	1990	Male	45 to 49	46% (31-60%)	54% (40-69%)	4.2 (2.2-7.3)	17% (13-22%)	21% (15-26%)	1800 (1300-2300)	77000 (55000-97000)
South Africa	1990	Male	50 to 54	45% (32-59%)	55% (41-68%)	3.9 (2-6.7)	16% (11-20%)	19% (13-25%)	1400 (940-1800)	53000 (37000-67000)
South Africa	1990	Male	55 to 59	46% (34-59%)	54% (41-66%)	3.9 (2.1-6.5)	15% (11-19%)	18% (13-23%)	1400 (990-1800)	47000 (34000-61000)
South Africa	1990	Male	60 to 64	45% (32-59%)	55% (41-68%)	3.5 (1.8-6.1)	12% (7.9-16%)	15% (9.5-20%)	1000 (670-1400)	29000 (19000-39000)
South Africa	1990	Male	65 to 69	43% (32-56%)	57% (44-68%)	2.8 (1.6-4.8)	9.8% (6.4-14%)	12% (7.8-17%)	1100 (690-1500)	25000 (16000-35000)
South Africa	1990	Male	70 to 74	41% (29-53%)	59% (47-71%)	2.1 (1.1-3.4)	6.7% (3.9-9.9%)	8.1% (4.7-12%)	610 (350-890)	12000 (6700-17000)
South Africa	1990	Male	75 to 79	37% (26-50%)	63% (50-74%)	1.6 (0.82-2.8)	4.7% (2.3-7.5%)	5.6% (2.7-9.1%)	470 (230-760)	7100 (3500-11000)
South Africa	1990	Male	80 to 84	35% (24-47%)	65% (53-76%)	1.4 (0.68-2.3)	3.7% (1.8-6.1%)	4.3% (2-7.1%)	290 (130-490)	3300 (1500-5400)
South Africa	1990	Male	85 to 89	33% (23-46%)	67% (54-77%)	1.2 (0.58-2)	2.9% (1.3-5.1%)	3.3% (1.5-5.9%)	150 (63-260)	1200 (530-2000)
South Africa	1990	Male	90 to 94	33% (23-45%)	67% (55-77%)	1 (0.47-1.9)	2.3% (0.88-4.4%)	2.6% (0.91-5.2%)	53 (18-100)	290 (110-550)
South Africa	1990	Male	95 plus	36% (22-52%)	64% (48-78%)	0.93 (0.36-2)	2.1% (0.74-4.4%)	2.3% (0.65-5%)	13 (3.6-27)	48 (17-100)
South Africa	1995	Female	All Ages	23% (19-26%)	77% (74-81%)	1.8 (1.3-2.5)	4.2% (3.3-5.2%)	6.7% (4.9-8.6%)	9400 (6900-12000)	360000 (280000-440000)
South Africa	1995	Female	15 to 19	17% (11-25%)	83% (75-89%)	1.3 (0.63-2.3)	3.6% (2.1-5.5%)	6.2% (3.4-9.6%)	140 (78-220)	13000 (7500-19000)
South Africa	1995	Female	20 to 24	23% (15-33%)	77% (67-85%)	2.2 (1.2-3.8)	5.6% (3.6-8%)	7.2% (4.4-11%)	410 (250-610)	33000 (21000-47000)
South Africa	1995	Female	25 to 29	26% (17-37%)	74% (63-83%)	2.9 (1.6-4.7)	6.8% (4.5-9.5%)	8.5% (5.4-12%)	680 (430-960)	47000 (31000-64000)
South Africa	1995	Female	30 to 34	30% (20-43%)	70% (57-80%)	3.3 (1.8-5.2)	7.3% (5.1-9.9%)	9.4% (6.3-13%)	740 (500-1000)	47000 (32000-62000)
South Africa	1995	Female	35 to 39	34% (23-46%)	66% (54-77%)	3.2 (1.8-5.3)	7% (5-9.5%)	9.5% (6.6-13%)	590 (410-810)	34000 (24000-46000)
South Africa	1995	Female	40 to 44	35% (24-49%)	65% (51-76%)	3.2 (1.7-5.4)	7.5% (5.1-10%)	11% (7.1-14%)	570 (390-770)	29000 (21000-39000)
South Africa	1995	Female	45 to 49	35% (23-47%)	65% (53-77%)	3.4 (1.7-6)	8% (5.4-11%)	11% (7.6-15%)	580 (390-780)	26000 (18000-35000)
South Africa	1995	Female	50 to 54	34% (24-47%)	66% (53-76%)	3.4 (1.8-6.1)	8.3% (5.5-11%)	12% (8-16%)	570 (380-770)	23000 (16000-31000)
South Africa	1995	Female	55 to 59	34% (23-46%)	66% (54-77%)	3.5 (1.8-6)	8.5% (5.5-12%)	12% (7.6-16%)	710 (460-980)	24000 (16000-33000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	1995	Female	60 to 64	35% (23-47%)	65% (53-77%)	3.4 (1.7-6.1)	8.1% (5.1-11%)	11% (7-16%)	800 (500-1100)	23000 (15000-33000)
South Africa	1995	Female	65 to 69	35% (25-47%)	65% (53-75%)	3.2 (1.7-5.4)	8.1% (5.1-11%)	11% (6.9-16%)	970 (620-1400)	23000 (15000-33000)
South Africa	1995	Female	70 to 74	33% (24-45%)	67% (55-76%)	2.7 (1.4-4.6)	6.6% (3.8-9.9%)	8.9% (5.1-13%)	670 (390-990)	13000 (7500-20000)
South Africa	1995	Female	75 to 79	30% (20-42%)	70% (58-80%)	2.3 (1.1-4)	5.2% (2.5-8.2%)	6.9% (3.4-11%)	650 (320-1000)	9900 (4800-16000)
South Africa	1995	Female	80 to 84	28% (19-39%)	72% (61-81%)	2 (1-3.5)	4.8% (2.3-7.8%)	6.1% (2.9-9.9%)	740 (360-1200)	8100 (4000-13000)
South Africa	1995	Female	85 to 89	29% (19-40%)	71% (60-81%)	2 (1-3.4)	4.6% (2.1-7.5%)	5.6% (2.6-9.2%)	420 (190-700)	3300 (1500-5300)
South Africa	1995	Female	90 to 94	31% (21-43%)	69% (57-79%)	2 (0.98-3.8)	4.3% (1.7-7.5%)	5.2% (2-9.2%)	150 (57-260)	780 (310-1300)
South Africa	1995	Female	95 plus	35% (22-49%)	65% (51-78%)	2.1 (0.83-4.3)	4.2% (1.3-8.2%)	5% (1.4-10%)	33 (9.1-66)	110 (33-220)
South Africa	1995	Male	All Ages	33% (28-38%)	67% (62-72%)	2.4 (1.7-3.3)	8.2% (6.7-9.7%)	10% (8.3-12%)	20000 (16000-24000)	880000 (720000-1100000)
South Africa	1995	Male	15 to 19	26% (17-36%)	74% (64-83%)	1.3 (0.65-2.3)	5.7% (3.4-8.8%)	8% (4.2-13%)	290 (150-460)	24000 (14000-37000)
South Africa	1995	Male	20 to 24	38% (25-51%)	62% (49-75%)	3 (1.6-4.9)	11% (7.5-15%)	13% (8.6-19%)	1000 (660-1400)	75000 (52000-100000)
South Africa	1995	Male	25 to 29	44% (31-57%)	56% (43-69%)	4.7 (2.6-7.5)	14% (10-18%)	16% (11-21%)	1800 (1300-2400)	120000 (85000-150000)
South Africa	1995	Male	30 to 34	47% (33-61%)	53% (39-67%)	5.5 (3.2-8.4)	15% (12-19%)	18% (13-23%)	2300 (1600-2900)	140000 (100000-170000)
South Africa	1995	Male	35 to 39	49% (35-62%)	51% (38-65%)	5.3 (3.2-8.4)	15% (12-19%)	18% (14-23%)	2200 (1700-2800)	120000 (91000-150000)
South Africa	1995	Male	40 to 44	48% (35-61%)	52% (39-65%)	4.8 (2.6-7.8)	15% (11-19%)	18% (13-22%)	2000 (1500-2500)	96000 (72000-120000)
South Africa	1995	Male	45 to 49	46% (32-60%)	54% (40-68%)	4.1 (2.2-6.9)	16% (12-20%)	19% (14-23%)	2100 (1500-2600)	90000 (67000-110000)
South Africa	1995	Male	50 to 54	46% (33-60%)	54% (40-67%)	3.7 (1.9-6.6)	15% (11-19%)	18% (12-23%)	1700 (1200-2200)	67000 (47000-85000)
South Africa	1995	Male	55 to 59	47% (35-59%)	53% (41-65%)	3.7 (2.1-6.2)	14% (10-18%)	17% (12-21%)	1800 (1300-2300)	60000 (44000-77000)
South Africa	1995	Male	60 to 64	46% (33-60%)	54% (40-67%)	3.4 (1.8-5.8)	12% (7.8-16%)	14% (9.2-19%)	1300 (890-1800)	38000 (25000-51000)
South Africa	1995	Male	65 to 69	44% (33-57%)	56% (43-67%)	2.7 (1.5-4.5)	9.5% (6.2-13%)	11% (7.3-16%)	1300 (820-1800)	30000 (20000-41000)
South Africa	1995	Male	70 to 74	42% (30-53%)	58% (47-70%)	2 (1.1-3.3)	6.5% (3.8-9.6%)	7.7% (4.4-12%)	770 (440-1200)	15000 (8500-22000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	1995	Male	75 to 79	38% (27-52%)	62% (48-73%)	1.6 (0.79-2.7)	4.5% (2.2-7.3%)	5.3% (2.6-8.6%)	540 (260-870)	8100 (3900-13000)
South Africa	1995	Male	80 to 84	36% (25-48%)	64% (52-75%)	1.3 (0.69-2.2)	3.5% (1.7-5.7%)	4.1% (1.9-6.7%)	350 (160-580)	3900 (1800-6300)
South Africa	1995	Male	85 to 89	34% (23-46%)	66% (54-77%)	1.1 (0.55-1.9)	2.7% (1.2-4.8%)	3.1% (1.3-5.5%)	180 (74-320)	1400 (620-2500)
South Africa	1995	Male	90 to 94	34% (23-47%)	66% (53-77%)	0.96 (0.45-1.8)	2.1% (0.83-4.1%)	2.4% (0.86-4.8%)	61 (22-120)	330 (130-650)
South Africa	1995	Male	95 plus	37% (23-52%)	63% (48-77%)	0.88 (0.35-1.8)	1.9% (0.65-4.2%)	2.1% (0.57-4.6%)	15 (4.2-34)	58 (19-130)
South Africa	2000	Female	All Ages	21% (18-25%)	79% (75-82%)	1.6 (1.1-2.2)	3.6% (2.7-4.6%)	5.4% (3.8-7.1%)	14000 (10000-19000)	510000 (390000-650000)
South Africa	2000	Female	15 to 19	15% (10-23%)	85% (77-90%)	1.1 (0.52-1.8)	3.8% (2.2-5.8%)	5.4% (2.8-8.5%)	260 (140-400)	21000 (12000-32000)
South Africa	2000	Female	20 to 24	19% (12-29%)	81% (71-88%)	1.8 (0.91-3)	4.2% (2.4-6.4%)	4.6% (2.5-7.4%)	650 (370-980)	48000 (29000-71000)
South Africa	2000	Female	25 to 29	22% (14-32%)	78% (68-86%)	2.3 (1.2-3.8)	4.2% (2.5-6.3%)	4.5% (2.5-6.9%)	1100 (600-1600)	69000 (41000-100000)
South Africa	2000	Female	30 to 34	26% (17-37%)	74% (63-83%)	2.6 (1.4-4.3)	4.2% (2.6-6.4%)	4.6% (2.7-7.1%)	1000 (640-1500)	63000 (40000-91000)
South Africa	2000	Female	35 to 39	30% (19-43%)	70% (57-81%)	2.7 (1.5-4.5)	4.1% (2.4-6.1%)	4.5% (2.5-7%)	870 (500-1300)	48000 (29000-71000)
South Africa	2000	Female	40 to 44	32% (22-45%)	68% (55-78%)	2.8 (1.4-4.6)	4% (2.5-6%)	4.6% (2.7-7%)	710 (430-1100)	36000 (23000-53000)
South Africa	2000	Female	45 to 49	32% (20-43%)	68% (57-80%)	2.9 (1.5-5)	4.9% (3.2-7.1%)	5.8% (3.6-8.5%)	750 (470-1100)	34000 (22000-48000)
South Africa	2000	Female	50 to 54	32% (22-45%)	68% (55-78%)	3 (1.6-5)	6% (3.9-8.4%)	7.2% (4.6-10%)	830 (530-1200)	32000 (21000-46000)
South Africa	2000	Female	55 to 59	34% (24-46%)	66% (54-76%)	3.2 (1.7-5.6)	7.2% (4.6-10%)	8.9% (5.7-13%)	970 (620-1400)	32000 (21000-45000)
South Africa	2000	Female	60 to 64	36% (24-48%)	64% (52-76%)	3.3 (1.7-5.8)	7.8% (4.7-11%)	9.7% (5.8-14%)	1300 (780-1900)	37000 (22000-53000)
South Africa	2000	Female	65 to 69	36% (26-47%)	64% (53-74%)	3.1 (1.7-5.2)	8.1% (4.9-11%)	10% (6.2-14%)	1500 (890-2100)	34000 (21000-48000)
South Africa	2000	Female	70 to 74	34% (24-45%)	66% (55-76%)	2.6 (1.3-4.3)	7.1% (3.9-11%)	8.7% (4.9-13%)	1400 (790-2100)	27000 (15000-40000)
South Africa	2000	Female	75 to 79	31% (21-43%)	69% (57-79%)	2.2 (1.1-3.8)	5.6% (2.6-9%)	6.9% (3.2-11%)	940 (440-1500)	14000 (6500-23000)
South Africa	2000	Female	80 to 84	30% (20-41%)	70% (59-80%)	2 (1-3.5)	5.2% (2.4-8.7%)	6.3% (3-11%)	1200 (550-1900)	12000 (5900-21000)
South Africa	2000	Female	85 to 89	31% (21-42%)	69% (58-79%)	2 (1-3.5)	4.9% (2.1-8.2%)	5.9% (2.5-9.9%)	680 (290-1100)	5100 (2200-8600)
South Africa	2000	Female	90 to 94	33% (23-45%)	67% (55-77%)	2 (1-3.7)	4.5% (1.8-7.9%)	5.4% (2.1-9.5%)	230 (88-410)	1200 (460-2100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	2000	Female	95 plus	37% (24-51%)	63% (49-76%)	2.1 (0.85-4.1)	4.4% (1.2-8.3%)	5.1% (1.3-9.9%)	52 (13-100)	180 (48-340)
South Africa	2000	Male	All Ages	34% (29-39%)	66% (61-71%)	2.4 (1.7-3.2)	7.3% (5.8-8.9%)	9% (7-11%)	27000 (21000-33000)	1200000 (930000-1400000)
South Africa	2000	Male	15 to 19	27% (18-38%)	73% (62-82%)	1.3 (0.65-2.3)	6.5% (3.7-9.9%)	8.3% (4.3-13%)	500 (260-790)	40000 (22000-61000)
South Africa	2000	Male	20 to 24	39% (27-52%)	61% (48-73%)	3 (1.6-4.9)	9.6% (6.4-14%)	11% (6.9-16%)	1300 (830-1900)	95000 (64000-130000)
South Africa	2000	Male	25 to 29	46% (32-59%)	54% (41-68%)	4.7 (2.7-7.5)	9.4% (6-13%)	10% (6.1-15%)	2100 (1300-3000)	140000 (90000-190000)
South Africa	2000	Male	30 to 34	48% (34-62%)	52% (38-66%)	5.4 (3.2-8.2)	10% (7.3-14%)	11% (7.7-15%)	2700 (1900-3700)	160000 (120000-210000)
South Africa	2000	Male	35 to 39	49% (36-63%)	51% (37-64%)	5.1 (3-8)	11% (7.7-14%)	12% (8.2-16%)	2900 (2100-3900)	150000 (110000-200000)
South Africa	2000	Male	40 to 44	48% (35-61%)	52% (39-65%)	4.6 (2.4-7.6)	10% (7.2-13%)	11% (7.8-15%)	2600 (1800-3500)	120000 (88000-160000)
South Africa	2000	Male	45 to 49	46% (32-59%)	54% (41-68%)	3.8 (2-6.4)	12% (8.6-15%)	13% (9.6-17%)	2900 (2100-3700)	120000 (88000-160000)
South Africa	2000	Male	50 to 54	45% (31-58%)	55% (42-69%)	3.5 (1.8-5.8)	13% (9.1-17%)	14% (10-19%)	2800 (2000-3600)	110000 (75000-140000)
South Africa	2000	Male	55 to 59	46% (34-59%)	54% (41-66%)	3.5 (2-5.9)	13% (9-16%)	14% (10-19%)	2500 (1800-3300)	83000 (59000-110000)
South Africa	2000	Male	60 to 64	47% (35-61%)	53% (39-65%)	3.3 (1.7-5.6)	11% (7.4-15%)	13% (8.4-17%)	2200 (1400-3000)	61000 (40000-82000)
South Africa	2000	Male	65 to 69	47% (36-59%)	53% (41-64%)	2.7 (1.5-4.5)	9.5% (6.1-13%)	11% (7-15%)	1700 (1100-2300)	39000 (25000-53000)
South Africa	2000	Male	70 to 74	45% (33-56%)	55% (44-67%)	2.1 (1.1-3.4)	6.8% (3.8-10%)	7.8% (4.3-12%)	1200 (670-1800)	23000 (12000-34000)
South Africa	2000	Male	75 to 79	42% (29-54%)	58% (46-71%)	1.6 (0.83-2.7)	4.6% (2.2-7.6%)	5.3% (2.5-8.7%)	660 (310-1100)	9700 (4500-16000)
South Africa	2000	Male	80 to 84	40% (28-52%)	60% (48-72%)	1.4 (0.75-2.3)	3.8% (1.8-6.3%)	4.3% (1.9-7.1%)	500 (230-820)	5400 (2500-8900)
South Africa	2000	Male	85 to 89	38% (27-50%)	62% (50-73%)	1.2 (0.6-2.1)	2.9% (1.3-5.1%)	3.2% (1.3-5.8%)	230 (95-410)	1700 (750-3100)
South Africa	2000	Male	90 to 94	37% (26-49%)	63% (51-74%)	0.99 (0.5-1.8)	2.1% (0.82-4.1%)	2.4% (0.86-4.7%)	69 (25-140)	380 (150-730)
South Africa	2000	Male	95 plus	38% (23-53%)	62% (47-77%)	0.87 (0.35-1.8)	2% (0.66-4.2%)	2.1% (0.59-4.7%)	18 (5.1-41)	67 (23-150)
South Africa	2005	Female	All Ages	21% (18-24%)	79% (76-82%)	1.7 (1.2-2.3)	3% (2.2-4%)	4.2% (2.9-5.6%)	16000 (12000-21000)	600000 (440000-780000)
South Africa	2005	Female	15 to 19	16% (11-24%)	84% (76-89%)	1.2 (0.59-2.1)	4.7% (2.7-7.1%)	6.5% (3.5-10%)	340 (190-550)	28000 (16000-42000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	2005	Female	20 to 24	20% (14-29%)	80% (71-86%)	2 (1-3.3)	5% (3-7.4%)	5.5% (3.1-8.4%)	1000 (610-1500)	75000 (46000-110000)
South Africa	2005	Female	25 to 29	22% (14-31%)	78% (69-86%)	2.4 (1.3-3.9)	3.3% (1.8-5.1%)	3.4% (1.8-5.4%)	1300 (720-2000)	84000 (48000-130000)
South Africa	2005	Female	30 to 34	25% (16-36%)	75% (64-84%)	2.7 (1.5-4.4)	3.5% (2.1-5.3%)	3.6% (2.2-5.6%)	1500 (950-2300)	90000 (56000-130000)
South Africa	2005	Female	35 to 39	28% (19-39%)	72% (61-81%)	2.7 (1.4-4.4)	2.5% (1.2-4.3%)	2.6% (1.1-4.6%)	960 (430-1600)	52000 (26000-87000)
South Africa	2005	Female	40 to 44	30% (21-42%)	70% (58-79%)	2.7 (1.4-4.6)	2.7% (1.6-4.2%)	2.9% (1.6-4.6%)	850 (480-1300)	42000 (25000-66000)
South Africa	2005	Female	45 to 49	30% (20-41%)	70% (59-80%)	2.9 (1.5-4.8)	2.7% (1.6-4.2%)	2.9% (1.5-4.7%)	680 (370-1100)	31000 (18000-48000)
South Africa	2005	Female	50 to 54	30% (21-41%)	70% (59-79%)	2.9 (1.6-5.1)	4.1% (2.6-6.1%)	4.7% (2.9-6.9%)	900 (560-1300)	35000 (22000-52000)
South Africa	2005	Female	55 to 59	31% (22-41%)	69% (59-78%)	3.1 (1.6-5.5)	5.4% (3.5-7.6%)	6.3% (4-9%)	1000 (650-1400)	34000 (22000-49000)
South Africa	2005	Female	60 to 64	32% (22-43%)	68% (57-78%)	3.1 (1.6-5.5)	6% (3.7-8.8%)	7.2% (4.4-11%)	1100 (690-1700)	32000 (20000-48000)
South Africa	2005	Female	65 to 69	32% (23-43%)	68% (57-77%)	2.9 (1.5-4.8)	6.8% (4.2-9.8%)	8.3% (5-12%)	1400 (860-2000)	33000 (20000-48000)
South Africa	2005	Female	70 to 74	30% (21-41%)	70% (59-79%)	2.4 (1.2-4)	5.9% (3.3-9%)	7.3% (4.1-11%)	1300 (710-1900)	24000 (13000-36000)
South Africa	2005	Female	75 to 79	28% (19-38%)	72% (62-81%)	2.1 (1-3.6)	5.4% (2.6-8.7%)	6.5% (3.2-10%)	1100 (530-1800)	16000 (7700-26000)
South Africa	2005	Female	80 to 84	28% (20-38%)	72% (62-80%)	2 (1-3.4)	5.3% (2.6-8.7%)	6.3% (3.1-10%)	1500 (710-2400)	16000 (7600-26000)
South Africa	2005	Female	85 to 89	29% (20-40%)	71% (60-80%)	2 (1-3.5)	5.1% (2.3-8.4%)	6.1% (2.7-10%)	830 (370-1400)	6200 (2800-10000)
South Africa	2005	Female	90 to 94	31% (22-42%)	69% (58-78%)	2 (1-3.7)	4.8% (2-8.1%)	5.7% (2.3-9.7%)	310 (120-530)	1600 (640-2700)
South Africa	2005	Female	95 plus	34% (22-48%)	66% (52-78%)	2.1 (0.87-4.1)	4.7% (1.4-8.8%)	5.5% (1.6-10%)	75 (22-140)	250 (79-470)
South Africa	2005	Male	All Ages	32% (28-37%)	68% (63-72%)	2.6 (1.8-3.6)	6.3% (5-8%)	7.6% (6-9.5%)	29000 (23000-36000)	1300000 (1000000-1600000)
South Africa	2005	Male	15 to 19	27% (18-38%)	73% (62-82%)	1.5 (0.73-2.6)	7% (4-10%)	9.2% (5-14%)	530 (290-810)	42000 (24000-63000)
South Africa	2005	Male	20 to 24	38% (26-51%)	62% (49-74%)	3.3 (1.8-5.3)	11% (6.9-15%)	12% (7.6-17%)	1600 (1000-2200)	110000 (75000-160000)
South Africa	2005	Male	25 to 29	43% (30-57%)	57% (43-70%)	5 (2.7-8)	6.9% (3.8-11%)	7.1% (3.7-12%)	1900 (1000-2900)	130000 (73000-190000)
South Africa	2005	Male	30 to 34	46% (32-59%)	54% (41-68%)	5.8 (3.3-9)	8.1% (5.5-11%)	8.5% (5.7-12%)	3200 (2200-4500)	190000 (130000-260000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	2005	Male	35 to 39	47% (33-60%)	53% (40-67%)	5.4 (3.1-8.6)	7.5% (4.9-11%)	8% (5.1-12%)	3100 (2000-4400)	160000 (110000-230000)
South Africa	2005	Male	40 to 44	45% (32-59%)	55% (41-68%)	4.8 (2.5-8.1)	7.8% (5.5-11%)	8.4% (5.8-12%)	3000 (2100-4100)	140000 (100000-190000)
South Africa	2005	Male	45 to 49	42% (28-56%)	58% (44-72%)	4 (2.1-6.5)	8.4% (6-11%)	9.1% (6.3-12%)	2900 (2000-3900)	120000 (86000-160000)
South Africa	2005	Male	50 to 54	41% (28-55%)	59% (45-72%)	3.6 (1.9-6)	11% (7.6-14%)	12% (8.2-15%)	3100 (2200-4100)	120000 (83000-150000)
South Africa	2005	Male	55 to 59	42% (30-56%)	58% (44-70%)	3.6 (1.9-6.1)	12% (8.3-15%)	13% (9.2-17%)	3000 (2100-3900)	96000 (68000-120000)
South Africa	2005	Male	60 to 64	42% (30-56%)	58% (44-70%)	3.3 (1.7-5.5)	9.9% (6.5-13%)	11% (7.4-15%)	2200 (1400-3000)	62000 (41000-83000)
South Africa	2005	Male	65 to 69	41% (30-54%)	59% (46-70%)	2.6 (1.4-4.3)	9.1% (5.8-12%)	10% (6.5-14%)	1900 (1200-2600)	44000 (28000-60000)
South Africa	2005	Male	70 to 74	38% (28-51%)	62% (49-72%)	2 (1-3.3)	5.9% (3.4-8.8%)	6.9% (3.8-10%)	1000 (580-1600)	20000 (11000-29000)
South Africa	2005	Male	75 to 79	36% (25-49%)	64% (51-75%)	1.5 (0.75-2.6)	4.5% (2.1-7.2%)	5.1% (2.4-8.2%)	690 (330-1100)	10000 (4800-16000)
South Africa	2005	Male	80 to 84	34% (24-47%)	66% (53-76%)	1.3 (0.67-2.1)	3.7% (1.7-6%)	4.2% (1.9-6.8%)	590 (260-970)	6400 (3000-10000)
South Africa	2005	Male	85 to 89	33% (23-45%)	67% (55-77%)	1.1 (0.56-1.9)	2.8% (1.2-4.9%)	3.2% (1.3-5.6%)	230 (96-400)	1800 (760-3100)
South Africa	2005	Male	90 to 94	32% (22-44%)	68% (56-78%)	0.93 (0.45-1.7)	2.1% (0.86-4.1%)	2.4% (0.9-4.7%)	69 (26-130)	370 (150-710)
South Africa	2005	Male	95 plus	34% (20-48%)	66% (52-80%)	0.83 (0.34-1.6)	2.1% (0.7-4.3%)	2.2% (0.7-4.8%)	17 (5.5-38)	64 (23-130)
South Africa	2010	Female	All Ages	22% (19-25%)	78% (75-81%)	1.9 (1.3-2.6)	2.8% (2.1-3.6%)	4.1% (2.9-5.4%)	15000 (10000-19000)	510000 (390000-650000)
South Africa	2010	Female	15 to 19	18% (12-25%)	82% (75-88%)	1.4 (0.67-2.4)	4.1% (2.4-6.1%)	5.5% (3.1-8.6%)	270 (160-430)	23000 (14000-35000)
South Africa	2010	Female	20 to 24	23% (15-32%)	77% (68-85%)	2.3 (1.2-3.9)	6% (3.8-8.5%)	6.9% (4.1-10%)	950 (590-1400)	70000 (45000-98000)
South Africa	2010	Female	25 to 29	25% (17-35%)	75% (65-83%)	2.8 (1.6-4.7)	3.6% (2.1-5.5%)	3.8% (2-6%)	1100 (640-1600)	74000 (45000-110000)
South Africa	2010	Female	30 to 34	27% (18-39%)	73% (61-82%)	3.1 (1.8-5)	2.3% (1.3-3.5%)	2.3% (1.3-3.7%)	790 (450-1200)	49000 (30000-73000)
South Africa	2010	Female	35 to 39	30% (20-42%)	70% (58-80%)	3 (1.6-4.9)	1.9% (0.96-3.1%)	1.9% (0.86-3.3%)	610 (290-1000)	35000 (19000-57000)
South Africa	2010	Female	40 to 44	32% (22-43%)	68% (57-78%)	3 (1.5-5.2)	1.4% (0.79-2.3%)	1.3% (0.62-2.3%)	360 (180-640)	20000 (12000-34000)
South Africa	2010	Female	45 to 49	31% (21-43%)	69% (57-79%)	3.1 (1.6-5.2)	2.5% (1.5-3.8%)	2.7% (1.6-4.2%)	660 (380-1000)	31000 (18000-45000)
South Africa	2010	Female	50 to 54	30% (21-42%)	70% (58-79%)	3.1 (1.6-5.3)	3.8% (2.4-5.5%)	4.2% (2.6-6.1%)	890 (570-1300)	35000 (23000-51000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	2010	Female	55 to 59	29% (20-39%)	71% (61-80%)	3.1 (1.6-5.5)	5.6% (3.6-8%)	6.6% (4.3-9.4%)	1200 (790-1700)	40000 (26000-57000)
South Africa	2010	Female	60 to 64	29% (20-41%)	71% (59-80%)	3 (1.5-5.2)	6.1% (3.6-8.9%)	7.3% (4.4-11%)	1300 (770-1900)	36000 (22000-53000)
South Africa	2010	Female	65 to 69	30% (21-40%)	70% (60-79%)	2.7 (1.5-4.4)	6.6% (4-9.7%)	8.1% (4.9-12%)	1400 (830-2000)	32000 (19000-47000)
South Africa	2010	Female	70 to 74	28% (20-38%)	72% (62-80%)	2.4 (1.2-4)	5.9% (3.1-8.8%)	7.2% (3.9-11%)	1300 (720-2000)	25000 (13000-38000)
South Africa	2010	Female	75 to 79	26% (18-36%)	74% (64-82%)	2 (0.99-3.5)	5.1% (2.4-8.3%)	6.3% (3-10%)	1100 (550-1800)	17000 (8000-27000)
South Africa	2010	Female	80 to 84	26% (18-36%)	74% (64-82%)	1.9 (0.98-3.2)	5% (2.5-8.2%)	6.1% (3-9.8%)	1600 (770-2500)	17000 (8200-27000)
South Africa	2010	Female	85 to 89	27% (19-37%)	73% (63-81%)	1.9 (0.95-3.4)	4.8% (2.2-8%)	5.7% (2.6-9.6%)	850 (380-1400)	6400 (2900-11000)
South Africa	2010	Female	90 to 94	29% (20-39%)	71% (61-80%)	1.9 (0.94-3.5)	4.5% (1.7-7.7%)	5.3% (1.9-9.1%)	300 (110-520)	1600 (590-2700)
South Africa	2010	Female	95 plus	32% (20-45%)	68% (55-80%)	2 (0.82-3.9)	4.5% (1.3-8.3%)	5.3% (1.4-10%)	80 (21-150)	270 (77-500)
South Africa	2010	Male	All Ages	32% (27-36%)	68% (64-73%)	2.8 (2-3.9)	5.8% (4.6-7.2%)	6.9% (5.3-8.5%)	25000 (19000-31000)	1100000 (830000-1300000)
South Africa	2010	Male	15 to 19	25% (17-35%)	75% (65-83%)	1.4 (0.71-2.5)	5.6% (3.3-8.3%)	7% (3.8-11%)	390 (210-590)	33000 (19000-49000)
South Africa	2010	Male	20 to 24	36% (24-48%)	64% (52-76%)	3.2 (1.8-5.2)	12% (8-16%)	13% (8.7-19%)	1500 (990-2100)	110000 (76000-150000)
South Africa	2010	Male	25 to 29	41% (28-55%)	59% (45-72%)	5 (2.8-8)	8.2% (4.9-12%)	8.7% (4.7-13%)	1900 (1100-2800)	130000 (80000-180000)
South Africa	2010	Male	30 to 34	45% (30-58%)	55% (42-70%)	5.9 (3.4-9.3)	6.2% (4-9.1%)	6.5% (3.9-9.7%)	2000 (1200-2900)	120000 (79000-170000)
South Africa	2010	Male	35 to 39	45% (32-58%)	55% (42-68%)	5.6 (3-9.2)	5.9% (3.8-8.7%)	6.2% (3.8-9.3%)	2100 (1300-3100)	110000 (76000-160000)
South Africa	2010	Male	40 to 44	43% (31-57%)	57% (43-69%)	4.9 (2.5-8.2)	5.3% (3.6-7.5%)	5.6% (3.7-7.9%)	1800 (1200-2500)	88000 (59000-120000)
South Africa	2010	Male	45 to 49	41% (27-55%)	59% (45-73%)	4.1 (2.2-6.7)	7.2% (5.1-9.7%)	7.8% (5.5-11%)	2300 (1600-3100)	100000 (71000-130000)
South Africa	2010	Male	50 to 54	41% (29-54%)	59% (46-71%)	3.7 (1.9-6.4)	9.2% (6.6-12%)	10% (7.2-13%)	2700 (2000-3600)	100000 (75000-140000)
South Africa	2010	Male	55 to 59	41% (29-55%)	59% (45-71%)	3.7 (2-6.3)	11% (8-14%)	13% (9-16%)	3100 (2200-4000)	100000 (72000-130000)
South Africa	2010	Male	60 to 64	40% (28-54%)	60% (46-72%)	3.2 (1.6-5.6)	9.9% (6.5-13%)	11% (7.4-15%)	2500 (1600-3400)	70000 (46000-94000)
South Africa	2010	Male	65 to 69	38% (27-50%)	62% (50-73%)	2.5 (1.3-4.1)	8.5% (5.4-12%)	9.8% (6.2-14%)	1900 (1200-2600)	43000 (28000-60000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	2010	Male	70 to 74	35% (25-47%)	65% (53-75%)	1.9 (0.95-3.2)	5.8% (3.3-8.9%)	6.8% (3.8-10%)	1200 (660-1800)	22000 (12000-33000)
South Africa	2010	Male	75 to 79	33% (23-45%)	67% (55-77%)	1.5 (0.7-2.5)	4.2% (2.1-6.7%)	4.8% (2.4-7.7%)	630 (300-1000)	9200 (4600-15000)
South Africa	2010	Male	80 to 84	32% (22-43%)	68% (57-78%)	1.2 (0.63-2.1)	3.4% (1.6-5.5%)	3.9% (1.7-6.3%)	540 (250-880)	5900 (2700-9500)
South Africa	2010	Male	85 to 89	30% (20-42%)	70% (58-80%)	1 (0.52-1.8)	2.6% (1.1-4.5%)	2.9% (1.2-5.1%)	250 (100-430)	1900 (820-3400)
South Africa	2010	Male	90 to 94	30% (20-41%)	70% (59-80%)	0.91 (0.43-1.7)	2% (0.78-3.9%)	2.3% (0.82-4.5%)	70 (25-140)	380 (140-720)
South Africa	2010	Male	95 plus	33% (20-47%)	67% (53-80%)	0.84 (0.34-1.7)	2% (0.72-4.2%)	2.2% (0.67-4.8%)	17 (5.3-37)	63 (23-130)
South Africa	2016	Female	All Ages	23% (20-27%)	77% (73-80%)	2 (1.4-2.8)	3.7% (2.7-4.6%)	5.4% (3.8-7.2%)	14000 (9600-18000)	470000 (360000-600000)
South Africa	2016	Female	15 to 19	16% (11-24%)	84% (76-89%)	1.2 (0.6-2.2)	2.1% (1.3-3.3%)	2.4% (1.3-4.1%)	99 (53-160)	11000 (6600-17000)
South Africa	2016	Female	20 to 24	22% (14-31%)	78% (69-86%)	2.2 (1.1-3.7)	4.6% (3-6.5%)	5.5% (3.3-8.2%)	430 (270-640)	37000 (24000-53000)
South Africa	2016	Female	25 to 29	25% (16-35%)	75% (65-84%)	2.8 (1.6-4.6)	4.7% (3-6.7%)	5.5% (3.3-8.2%)	790 (510-1100)	56000 (38000-78000)
South Africa	2016	Female	30 to 34	28% (19-40%)	72% (60-81%)	3.1 (1.8-5.1)	4.3% (2.9-6.2%)	4.9% (3.1-7.2%)	840 (560-1200)	52000 (36000-73000)
South Africa	2016	Female	35 to 39	31% (21-44%)	69% (56-79%)	3.1 (1.6-4.9)	3.6% (2.2-5.4%)	4.2% (2.4-6.2%)	700 (420-1000)	40000 (25000-56000)
South Africa	2016	Female	40 to 44	33% (23-46%)	67% (54-77%)	3.1 (1.5-5.2)	3.7% (2.3-5.4%)	4.4% (2.6-6.3%)	700 (440-990)	36000 (23000-50000)
South Africa	2016	Female	45 to 49	33% (21-44%)	67% (56-79%)	3.2 (1.7-5.4)	4.1% (2.7-5.9%)	5% (3.2-7.2%)	790 (520-1100)	37000 (24000-51000)
South Africa	2016	Female	50 to 54	32% (23-44%)	68% (56-77%)	3.2 (1.8-5.7)	5.4% (3.5-7.7%)	6.9% (4.5-9.6%)	1000 (670-1400)	41000 (27000-57000)
South Africa	2016	Female	55 to 59	33% (23-44%)	67% (56-77%)	3.4 (1.8-5.8)	6.5% (4.1-9.3%)	8.6% (5.4-12%)	1200 (740-1700)	40000 (25000-58000)
South Africa	2016	Female	60 to 64	33% (23-46%)	67% (54-77%)	3.4 (1.7-5.8)	7.1% (4.3-10%)	9.4% (5.7-14%)	1300 (760-1900)	36000 (22000-54000)
South Africa	2016	Female	65 to 69	34% (25-45%)	66% (55-75%)	3.1 (1.7-5.1)	7.1% (4.2-11%)	9.4% (5.6-14%)	1200 (690-1700)	28000 (16000-42000)
South Africa	2016	Female	70 to 74	32% (23-43%)	68% (57-77%)	2.6 (1.3-4.4)	6% (3.2-9.3%)	7.9% (4.2-12%)	1100 (600-1700)	22000 (11000-34000)
South Africa	2016	Female	75 to 79	28% (20-39%)	72% (61-80%)	2.2 (1.1-3.8)	4.9% (2.3-7.9%)	6.4% (3.1-10%)	900 (430-1400)	14000 (6400-22000)
South Africa	2016	Female	80 to 84	27% (18-37%)	73% (63-82%)	2 (1-3.3)	4.7% (2.2-7.6%)	5.9% (2.8-9.6%)	1200 (580-2000)	13000 (6500-22000)
South Africa	2016	Female	85 to 89	28% (19-38%)	72% (62-81%)	1.9 (0.97-3.3)	4.5% (2.1-7.6%)	5.5% (2.5-9.2%)	890 (400-1500)	6800 (3100-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Africa	2016	Female	90 to 94	31% (22-41%)	69% (59-78%)	2 (0.98-3.5)	4.3% (1.7-7.6%)	5.2% (2.1-9.1%)	360 (140-630)	1900 (740-3300)
South Africa	2016	Female	95 plus	34% (22-47%)	66% (53-78%)	2.1 (0.86-3.9)	4.4% (1.3-8.3%)	5.3% (1.4-10%)	94 (25-180)	320 (92-600)
South Africa	2016	Male	All Ages	33% (28-38%)	67% (62-72%)	2.7 (1.9-3.8)	6.8% (5.5-8.2%)	8.1% (6.3-9.8%)	23000 (18000-28000)	990000 (800000-1200000)
South Africa	2016	Male	15 to 19	24% (16-35%)	76% (65-84%)	1.3 (0.61-2.2)	3.2% (1.9-4.7%)	3.3% (1.7-5.4%)	170 (89-280)	17000 (11000-26000)
South Africa	2016	Male	20 to 24	35% (23-48%)	65% (52-77%)	2.9 (1.5-4.8)	10% (6.9-14%)	12% (7.5-17%)	1000 (660-1500)	83000 (57000-110000)
South Africa	2016	Male	25 to 29	41% (28-55%)	59% (45-72%)	4.6 (2.6-7.4)	11% (7.3-14%)	12% (7.8-16%)	1900 (1300-2500)	130000 (92000-170000)
South Africa	2016	Male	30 to 34	44% (30-58%)	56% (42-70%)	5.4 (3.1-8.3)	9.6% (6.8-13%)	11% (7.2-14%)	2200 (1500-3000)	130000 (96000-180000)
South Africa	2016	Male	35 to 39	46% (32-59%)	54% (41-68%)	5.2 (2.8-8.1)	8.5% (5.9-12%)	9.5% (6.4-13%)	2100 (1500-2900)	120000 (84000-150000)
South Africa	2016	Male	40 to 44	45% (33-59%)	55% (41-67%)	4.7 (2.4-7.8)	8% (5.6-10%)	8.9% (6.2-12%)	2000 (1400-2600)	96000 (68000-120000)
South Africa	2016	Male	45 to 49	44% (29-58%)	56% (42-71%)	4 (2.1-6.7)	8.3% (5.8-11%)	9.4% (6.6-12%)	2000 (1400-2600)	88000 (62000-110000)
South Africa	2016	Male	50 to 54	43% (30-57%)	57% (43-70%)	3.7 (2-6.4)	9.8% (6.9-13%)	11% (7.9-15%)	2400 (1700-3100)	93000 (66000-120000)
South Africa	2016	Male	55 to 59	44% (31-57%)	56% (43-69%)	3.7 (2-6.2)	11% (7.7-14%)	13% (9.1-16%)	2600 (1900-3400)	86000 (62000-110000)
South Africa	2016	Male	60 to 64	43% (31-58%)	57% (42-69%)	3.3 (1.6-5.6)	10% (6.5-14%)	12% (7.6-16%)	2400 (1500-3300)	68000 (43000-93000)
South Africa	2016	Male	65 to 69	42% (30-54%)	58% (46-70%)	2.6 (1.4-4.4)	8.4% (5.2-12%)	10% (6.2-14%)	1700 (1000-2300)	38000 (23000-54000)
South Africa	2016	Male	70 to 74	39% (28-51%)	61% (49-72%)	1.9 (0.98-3.3)	5.7% (3-8.6%)	6.8% (3.6-10%)	1000 (560-1600)	20000 (11000-30000)
South Africa	2016	Male	75 to 79	36% (25-49%)	64% (51-75%)	1.5 (0.73-2.5)	4% (1.8-6.4%)	4.7% (2.2-7.6%)	530 (250-870)	7900 (3800-13000)
South Africa	2016	Male	80 to 84	33% (23-46%)	67% (54-77%)	1.2 (0.61-2)	3.1% (1.4-5.1%)	3.6% (1.6-6%)	460 (200-770)	5000 (2300-8400)
South Africa	2016	Male	85 to 89	32% (22-43%)	68% (57-78%)	1 (0.53-1.7)	2.4% (0.99-4.2%)	2.7% (1.1-4.9%)	250 (100-450)	1900 (820-3500)
South Africa	2016	Male	90 to 94	32% (22-43%)	68% (57-78%)	0.89 (0.43-1.6)	1.8% (0.67-3.6%)	2% (0.68-4.1%)	86 (29-170)	470 (180-910)
South Africa	2016	Male	95 plus	34% (21-49%)	66% (51-79%)	0.83 (0.33-1.6)	1.8% (0.59-3.9%)	1.9% (0.53-4.3%)	19 (5.4-44)	74 (24-160)
Swaziland	1990	Female	All Ages	9.3% (7.6-11%)	91% (89-92%)	0.71 (0.52-0.95)	2.7% (1.9-3.8%)	4.6% (3-6.6%)	130 (83-200)	4900 (3400-6800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	1990	Female	15 to 19	8.2% (5.1-12%)	92% (88-95%)	0.54 (0.27-0.95)	2.5% (1.3-4.2%)	3.8% (1.7-7%)	2.2 (0.98-4.1)	210 (110-350)
Swaziland	1990	Female	20 to 24	12% (7.3-18%)	88% (82-93%)	1.1 (0.55-1.8)	4.5% (2.6-7.1%)	6.1% (2.9-11%)	4.3 (2.1-7.5)	380 (220-610)
Swaziland	1990	Female	25 to 29	13% (8.1-19%)	87% (81-92%)	1.3 (0.72-2.1)	5.1% (3-7.8%)	7.2% (3.7-12%)	5 (2.6-8.5)	390 (230-600)
Swaziland	1990	Female	30 to 34	14% (8.9-23%)	86% (77-91%)	1.5 (0.8-2.4)	5.3% (3.2-8%)	7.7% (4.1-12%)	5.5 (2.8-9.1)	370 (210-570)
Swaziland	1990	Female	35 to 39	17% (10-24%)	83% (76-90%)	1.6 (0.89-2.7)	6% (3.8-8.8%)	8.7% (5.2-13%)	7 (4-11)	410 (240-630)
Swaziland	1990	Female	40 to 44	18% (12-26%)	82% (74-88%)	1.7 (0.86-2.8)	7.1% (4.4-10%)	10% (6.1-15%)	9.1 (5.1-15)	460 (270-710)
Swaziland	1990	Female	45 to 49	19% (11-27%)	81% (73-89%)	1.6 (0.86-2.7)	7.7% (4.8-12%)	11% (6.5-16%)	11 (6.1-17)	470 (280-740)
Swaziland	1990	Female	50 to 54	19% (12-27%)	81% (73-88%)	1.6 (0.82-2.8)	7.7% (4.6-12%)	10% (6-15%)	12 (6.7-20)	470 (260-750)
Swaziland	1990	Female	55 to 59	19% (12-28%)	81% (72-88%)	1.7 (0.83-3)	7.8% (4.3-12%)	9.8% (5.5-15%)	14 (7.4-23)	470 (240-750)
Swaziland	1990	Female	60 to 64	19% (12-29%)	81% (71-88%)	1.7 (0.83-3)	7.4% (3.8-12%)	9.1% (4.7-14%)	16 (7.9-26)	440 (220-730)
Swaziland	1990	Female	65 to 69	19% (13-27%)	81% (73-87%)	1.5 (0.79-2.5)	6.8% (3.4-11%)	8.2% (4.1-13%)	17 (8.2-28)	390 (190-630)
Swaziland	1990	Female	70 to 74	17% (11-25%)	83% (75-89%)	1.2 (0.59-2.2)	5% (2.3-9%)	5.9% (2.6-11%)	13 (5.9-24)	240 (110-440)
Swaziland	1990	Female	75 to 79	14% (8.7-22%)	86% (78-91%)	0.89 (0.4-1.7)	3.6% (1.3-7.1%)	4.2% (1.5-8.2%)	8.5 (3-17)	120 (43-240)
Swaziland	1990	Female	80 to 84	13% (7.9-20%)	87% (80-92%)	0.76 (0.36-1.4)	2.8% (1-5.8%)	3.2% (1.2-6.8%)	4.9 (1.7-10)	52 (18-110)
Swaziland	1990	Female	85 to 89	13% (7.9-20%)	87% (80-92%)	0.71 (0.37-1.3)	2.3% (0.7-4.9%)	2.6% (0.8-5.7%)	2 (0.61-4.4)	15 (4.7-33)
Swaziland	1990	Female	90 to 94	14% (8.8-21%)	86% (79-91%)	0.68 (0.33-1.3)	1.8% (0.42-4.4%)	2.1% (0.43-5.1%)	0.54 (0.11-1.3)	2.8 (0.63-6.7)
Swaziland	1990	Female	95 plus	15% (8.5-24%)	85% (76-92%)	0.67 (0.27-1.3)	1.6% (0.15-4.3%)	1.8% (0.12-4.9%)	0.11 (0.0071-0.29)	0.38 (0.036-1)
Swaziland	1990	Male	All Ages	17% (14-20%)	83% (80-86%)	1.7 (1.3-2.3)	6.6% (5.1-8.2%)	10% (7.9-13%)	350 (260-450)	14000 (11000-18000)
Swaziland	1990	Male	15 to 19	15% (9.4-21%)	85% (79-91%)	1.2 (0.6-2.1)	5.8% (3.4-9.1%)	7.3% (3.9-12%)	7.3 (4-12)	610 (360-950)
Swaziland	1990	Male	20 to 24	25% (16-34%)	75% (66-84%)	2.8 (1.5-4.6)	13% (8.6-18%)	16% (9.7-23%)	18 (11-27)	1400 (890-2000)
Swaziland	1990	Male	25 to 29	30% (20-41%)	70% (59-80%)	4 (2.3-6.4)	16% (11-21%)	19% (13-27%)	20 (13-27)	1300 (910-1800)
Swaziland	1990	Male	30 to 34	34% (22-47%)	66% (53-78%)	4.7 (2.6-7.5)	17% (11-22%)	21% (14-27%)	19 (13-26)	1200 (810-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	1990	Male	35 to 39	37% (25-50%)	63% (50-75%)	4.9 (2.7-7.9)	18% (13-23%)	22% (16-29%)	22 (15-29)	1200 (830-1600)
Swaziland	1990	Male	40 to 44	37% (26-50%)	63% (50-74%)	4.9 (2.5-7.8)	19% (14-25%)	23% (17-30%)	27 (19-37)	1300 (900-1700)
Swaziland	1990	Male	45 to 49	36% (22-48%)	64% (52-78%)	4.7 (2.5-7.7)	20% (15-25%)	24% (18-31%)	35 (24-48)	1500 (1000-2000)
Swaziland	1990	Male	50 to 54	36% (24-49%)	64% (51-76%)	4.7 (2.5-7.7)	20% (14-25%)	23% (17-30%)	45 (31-63)	1700 (1200-2400)
Swaziland	1990	Male	55 to 59	39% (27-53%)	61% (47-73%)	4.9 (2.5-8.4)	19% (14-24%)	23% (16-29%)	46 (32-64)	1500 (1000-2000)
Swaziland	1990	Male	60 to 64	36% (24-50%)	64% (50-76%)	4.1 (2-7.2)	16% (11-21%)	19% (13-25%)	39 (25-55)	1100 (690-1500)
Swaziland	1990	Male	65 to 69	28% (19-38%)	72% (62-81%)	2.7 (1.4-4.4)	12% (7.9-17%)	14% (9-19%)	31 (19-45)	710 (430-1000)
Swaziland	1990	Male	70 to 74	24% (17-34%)	76% (66-83%)	1.8 (0.94-3.1)	8% (4.6-12%)	9.1% (5.2-14%)	20 (11-32)	370 (200-590)
Swaziland	1990	Male	75 to 79	24% (15-35%)	76% (65-85%)	1.4 (0.69-2.6)	5.8% (2.8-10%)	6.6% (3.1-11%)	11 (4.9-19)	160 (71-270)
Swaziland	1990	Male	80 to 84	23% (14-35%)	77% (65-86%)	1.3 (0.62-2.2)	4.6% (2.1-8.2%)	5.2% (2.4-9.2%)	5.2 (2.3-9.4)	55 (25-99)
Swaziland	1990	Male	85 to 89	22% (13-32%)	78% (68-87%)	1 (0.53-1.9)	3.4% (1.5-6.4%)	3.8% (1.7-7.3%)	1.7 (0.75-3.3)	13 (5.8-25)
Swaziland	1990	Male	90 to 94	22% (14-32%)	78% (68-86%)	0.92 (0.44-1.7)	2.6% (1.1-5.3%)	2.9% (1.2-6.1%)	0.41 (0.16-0.89)	2.2 (0.92-4.7)
Swaziland	1990	Male	95 plus	24% (13-38%)	76% (62-87%)	0.88 (0.33-1.9)	2.3% (0.8-4.9%)	2.4% (0.77-5.5%)	0.083 (0.026-0.19)	0.31 (0.11-0.66)
Swaziland	1995	Female	All Ages	9% (7.4-11%)	91% (89-93%)	0.69 (0.5-0.92)	1.9% (1.1-2.9%)	3.2% (1.7-5.1%)	120 (57-190)	4300 (2400-6800)
Swaziland	1995	Female	15 to 19	8.1% (5-12%)	92% (88-95%)	0.54 (0.26-0.95)	1.8% (0.89-3.1%)	2.3% (0.85-4.6%)	1.9 (0.66-3.8)	190 (94-330)
Swaziland	1995	Female	20 to 24	12% (7.2-18%)	88% (82-93%)	1.1 (0.55-1.8)	2% (1-3.5%)	2% (0.67-4.2%)	3.4 (1.3-6.6)	320 (180-530)
Swaziland	1995	Female	25 to 29	13% (7.9-19%)	87% (81-92%)	1.3 (0.68-2.2)	2% (0.97-3.8%)	2% (0.72-4.5%)	4.1 (1.7-7.8)	340 (180-570)
Swaziland	1995	Female	30 to 34	14% (8.9-22%)	86% (78-91%)	1.5 (0.79-2.4)	2.3% (1.1-4.1%)	2.4% (0.88-5%)	4.4 (1.9-8.3)	320 (170-530)
Swaziland	1995	Female	35 to 39	16% (9.9-24%)	84% (76-90%)	1.6 (0.83-2.6)	3% (1.4-5.4%)	3.4% (1.3-6.9%)	5.6 (2.5-11)	340 (180-590)
Swaziland	1995	Female	40 to 44	18% (12-26%)	82% (74-88%)	1.7 (0.82-2.7)	4.1% (2.1-6.7%)	5.1% (2.4-9.2%)	7.5 (3.3-14)	390 (190-660)
Swaziland	1995	Female	45 to 49	19% (11-26%)	81% (74-89%)	1.7 (0.82-2.8)	5.1% (2.7-8.5%)	6.4% (3.3-11%)	9.3 (4.2-17)	420 (210-740)
Swaziland	1995	Female	50 to 54	19% (12-26%)	81% (74-88%)	1.6 (0.84-2.9)	5.7% (3-9.4%)	7.2% (3.7-12%)	10 (4.3-20)	400 (190-740)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	1995	Female	55 to 59	18% (12-27%)	82% (73-88%)	1.7 (0.84-2.9)	6.1% (3.3-10%)	7.7% (4.1-12%)	12 (4.9-23)	400 (180-760)
Swaziland	1995	Female	60 to 64	19% (11-29%)	81% (71-89%)	1.7 (0.82-3.1)	6.2% (3-11%)	7.8% (3.9-13%)	14 (5.5-26)	380 (160-710)
Swaziland	1995	Female	65 to 69	19% (12-27%)	81% (73-88%)	1.5 (0.78-2.6)	6% (3.1-9.9%)	7.5% (3.8-12%)	15 (6-28)	350 (140-630)
Swaziland	1995	Female	70 to 74	17% (11-25%)	83% (75-89%)	1.2 (0.61-2.2)	4.5% (1.9-8%)	5.5% (2.4-9.6%)	12 (4.3-24)	220 (83-440)
Swaziland	1995	Female	75 to 79	14% (8.6-22%)	86% (78-91%)	0.89 (0.41-1.7)	3.3% (1.1-6.5%)	4% (1.4-7.8%)	8.2 (2.5-17)	120 (38-240)
Swaziland	1995	Female	80 to 84	13% (7.8-19%)	87% (81-92%)	0.75 (0.37-1.4)	2.6% (0.85-5.6%)	3.1% (1-6.5%)	4.9 (1.6-11)	52 (17-120)
Swaziland	1995	Female	85 to 89	13% (7.9-19%)	87% (81-92%)	0.71 (0.35-1.3)	2.2% (0.72-4.7%)	2.6% (0.83-5.5%)	2.3 (0.68-4.8)	17 (5.2-36)
Swaziland	1995	Female	90 to 94	14% (8.7-20%)	86% (80-91%)	0.68 (0.33-1.3)	1.8% (0.44-4.2%)	2.1% (0.49-5%)	0.62 (0.14-1.5)	3.2 (0.73-7.5)
Swaziland	1995	Female	95 plus	15% (8.8-23%)	85% (77-91%)	0.68 (0.27-1.3)	1.6% (0.15-4.2%)	1.9% (0.09-4.9%)	0.13 (0.0056-0.34)	0.44 (0.037-1.1)
Swaziland	1995	Male	All Ages	17% (13-20%)	83% (80-87%)	1.7 (1.3-2.3)	5.7% (4.3-7.4%)	8.5% (6.3-11%)	380 (270-520)	15000 (11000-20000)
Swaziland	1995	Male	15 to 19	14% (9.3-21%)	86% (79-91%)	1.2 (0.6-2.1)	5.4% (3.2-8.5%)	6.9% (3.6-11%)	7.8 (4.1-13)	660 (380-1000)
Swaziland	1995	Male	20 to 24	24% (16-34%)	76% (66-84%)	2.8 (1.5-4.5)	8.7% (5.3-13%)	9.6% (5.4-15%)	19 (12-29)	1400 (930-2100)
Swaziland	1995	Male	25 to 29	30% (19-41%)	70% (59-81%)	4 (2.2-6.5)	8.7% (5.2-13%)	9.5% (5.2-16%)	21 (13-29)	1400 (930-1900)
Swaziland	1995	Male	30 to 34	34% (22-47%)	66% (53-78%)	4.6 (2.5-7.2)	9.2% (5.6-14%)	10% (5.8-16%)	21 (14-30)	1300 (860-1800)
Swaziland	1995	Male	35 to 39	37% (24-49%)	63% (51-76%)	4.8 (2.6-7.5)	11% (6.3-16%)	12% (6.7-18%)	26 (17-37)	1400 (930-1900)
Swaziland	1995	Male	40 to 44	37% (26-51%)	63% (49-74%)	4.8 (2.5-8)	12% (7.8-17%)	14% (8.7-20%)	31 (20-45)	1500 (1000-2100)
Swaziland	1995	Male	45 to 49	35% (22-48%)	65% (52-78%)	4.6 (2.4-7.8)	15% (10-21%)	17% (12-24%)	40 (26-59)	1700 (1100-2500)
Swaziland	1995	Male	50 to 54	36% (25-48%)	64% (52-75%)	4.6 (2.5-7.6)	16% (12-21%)	19% (13-24%)	45 (29-67)	1700 (1100-2500)
Swaziland	1995	Male	55 to 59	39% (27-52%)	61% (48-73%)	4.8 (2.5-8.2)	17% (12-22%)	19% (14-25%)	52 (34-76)	1700 (1100-2400)
Swaziland	1995	Male	60 to 64	36% (23-49%)	64% (51-77%)	4 (2-7.2)	14% (9.8-19%)	17% (11-22%)	45 (28-66)	1200 (780-1800)
Swaziland	1995	Male	65 to 69	28% (19-38%)	72% (62-81%)	2.6 (1.4-4.5)	11% (7-15%)	13% (8-18%)	32 (18-48)	720 (420-1100)
Swaziland	1995	Male	70 to 74	24% (16-33%)	76% (67-84%)	1.8 (0.93-3.1)	7.4% (4.1-11%)	8.5% (4.7-13%)	21 (11-34)	390 (200-620)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	1995	Male	75 to 79	23% (15-34%)	77% (66-85%)	1.4 (0.67-2.7)	5.5% (2.6-9.6%)	6.2% (2.9-11%)	13 (5.8-23)	190 (82-330)
Swaziland	1995	Male	80 to 84	23% (14-33%)	77% (67-86%)	1.2 (0.61-2.2)	4.5% (2.1-7.8%)	5.1% (2.4-8.8%)	6.3 (2.8-12)	67 (31-120)
Swaziland	1995	Male	85 to 89	21% (13-31%)	79% (69-87%)	1 (0.51-1.9)	3.3% (1.5-6.4%)	3.8% (1.6-7.2%)	2.1 (0.89-3.9)	16 (6.9-30)
Swaziland	1995	Male	90 to 94	21% (13-31%)	79% (69-87%)	0.91 (0.43-1.7)	2.5% (1.1-5.2%)	2.9% (1.1-5.8%)	0.48 (0.19-0.98)	2.5 (1.1-5.1)
Swaziland	1995	Male	95 plus	23% (13-38%)	77% (62-87%)	0.86 (0.33-1.8)	2.3% (0.83-4.9%)	2.4% (0.8-5.5%)	0.09 (0.028-0.2)	0.33 (0.12-0.71)
Swaziland	2000	Female	All Ages	8.6% (7.1-10%)	91% (90-93%)	0.68 (0.51-0.89)	1.5% (0.89-2.8%)	2.5% (1.3-4.4%)	180 (94-370)	6500 (3800-13000)
Swaziland	2000	Female	15 to 19	7.7% (4.9-11%)	92% (89-95%)	0.55 (0.27-0.92)	2% (0.97-3.6%)	2.7% (0.96-5.3%)	3.5 (1.3-7.2)	310 (160-580)
Swaziland	2000	Female	20 to 24	10% (6-16%)	90% (84-94%)	0.96 (0.48-1.7)	1.5% (0.71-3%)	1.5% (0.51-3.3%)	5.5 (2.4-11)	470 (250-870)
Swaziland	2000	Female	25 to 29	11% (6.3-16%)	89% (84-94%)	1.1 (0.59-1.8)	1% (0.43-2.3%)	0.92% (0.32-2.4%)	6 (2.6-14)	450 (230-930)
Swaziland	2000	Female	30 to 34	12% (7.3-19%)	88% (81-93%)	1.3 (0.7-2.2)	1.1% (0.49-2.5%)	1.1% (0.39-2.6%)	7 (3-16)	460 (230-970)
Swaziland	2000	Female	35 to 39	15% (9.2-22%)	85% (78-91%)	1.5 (0.8-2.5)	1.6% (0.66-3.6%)	1.6% (0.58-3.9%)	9.4 (4.2-23)	540 (270-1200)
Swaziland	2000	Female	40 to 44	16% (11-23%)	84% (77-89%)	1.6 (0.8-2.6)	2.2% (1-4.7%)	2.3% (1-5.3%)	12 (5.4-28)	600 (290-1300)
Swaziland	2000	Female	45 to 49	17% (11-25%)	83% (75-89%)	1.6 (0.81-2.7)	3% (1.4-6.2%)	3.3% (1.5-7.2%)	15 (6.8-35)	660 (310-1500)
Swaziland	2000	Female	50 to 54	19% (12-26%)	81% (74-88%)	1.7 (0.91-2.8)	4% (1.9-7.7%)	4.4% (2.1-8.6%)	17 (7.4-40)	650 (300-1500)
Swaziland	2000	Female	55 to 59	19% (12-27%)	81% (73-88%)	1.7 (0.91-3)	5% (2.6-8.9%)	5.7% (2.9-10%)	18 (8.1-41)	600 (280-1300)
Swaziland	2000	Female	60 to 64	20% (13-28%)	80% (72-87%)	1.8 (0.91-3.1)	5.8% (2.9-9.9%)	6.9% (3.4-11%)	21 (9-45)	580 (260-1200)
Swaziland	2000	Female	65 to 69	21% (14-29%)	79% (71-86%)	1.7 (0.95-2.9)	6.3% (3.2-10%)	7.4% (3.8-12%)	24 (10-50)	540 (230-1100)
Swaziland	2000	Female	70 to 74	19% (13-26%)	81% (74-87%)	1.4 (0.73-2.4)	5.1% (2.3-8.9%)	6% (2.7-10%)	19 (7-41)	350 (130-730)
Swaziland	2000	Female	75 to 79	15% (9.1-22%)	85% (78-91%)	0.98 (0.49-1.8)	3.8% (1.3-7%)	4.4% (1.6-8.4%)	13 (3.9-29)	180 (56-400)
Swaziland	2000	Female	80 to 84	13% (8.1-19%)	87% (81-92%)	0.8 (0.39-1.4)	3% (1.1-5.7%)	3.5% (1.3-6.7%)	7.1 (2.4-15)	75 (25-160)
Swaziland	2000	Female	85 to 89	13% (7.7-19%)	87% (81-92%)	0.74 (0.36-1.4)	2.5% (0.82-5%)	2.9% (0.96-5.9%)	3.5 (1.1-7.1)	26 (8.4-53)
Swaziland	2000	Female	90 to 94	14% (8.7-20%)	86% (80-91%)	0.71 (0.35-1.3)	2% (0.52-4.5%)	2.3% (0.59-5.3%)	1 (0.24-2.2)	5.1 (1.3-11)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	2000	Female	95 plus	15% (8.5-24%)	85% (76-92%)	0.7 (0.27-1.4)	1.9% (0.17-4.4%)	2.1% (0.1-5.2%)	0.2 (0.01-0.5)	0.7 (0.061-1.7)
Swaziland	2000	Male	All Ages	17% (14-20%)	83% (80-86%)	1.8 (1.4-2.4)	5.4% (3.5-7.9%)	7.7% (5.1-11%)	650 (380-990)	25000 (16000-38000)
Swaziland	2000	Male	15 to 19	13% (8.1-18%)	87% (82-92%)	1.1 (0.55-1.8)	5.9% (3.4-9.1%)	7.3% (3.8-12%)	13 (5.9-22)	1000 (550-1700)
Swaziland	2000	Male	20 to 24	21% (13-29%)	79% (71-87%)	2.5 (1.3-3.9)	7.9% (4.4-12%)	8.5% (4.4-14%)	30 (16-49)	2200 (1300-3400)
Swaziland	2000	Male	25 to 29	27% (18-36%)	73% (64-82%)	3.7 (2.2-5.7)	5.6% (3.1-9.7%)	5.7% (2.9-10%)	33 (19-52)	2200 (1300-3300)
Swaziland	2000	Male	30 to 34	31% (21-42%)	69% (58-79%)	4.4 (2.6-6.7)	5.6% (3.1-9.6%)	5.8% (3-10%)	34 (20-54)	2000 (1200-3100)
Swaziland	2000	Male	35 to 39	38% (28-50%)	62% (50-72%)	5.3 (3.1-8.2)	7.2% (3.8-12%)	7.6% (3.8-14%)	44 (24-73)	2300 (1300-3700)
Swaziland	2000	Male	40 to 44	42% (31-53%)	58% (47-69%)	5.7 (3.2-9.1)	8.7% (4.8-14%)	9.3% (5-16%)	57 (31-96)	2700 (1500-4400)
Swaziland	2000	Male	45 to 49	37% (25-50%)	63% (50-75%)	5.2 (2.8-8.3)	11% (6.4-17%)	12% (6.6-19%)	66 (35-120)	2800 (1500-4800)
Swaziland	2000	Male	50 to 54	37% (26-48%)	63% (52-74%)	5 (2.9-7.9)	14% (8.9-20%)	15% (9.5-22%)	77 (39-130)	2800 (1500-4600)
Swaziland	2000	Male	55 to 59	43% (31-54%)	57% (46-69%)	5.7 (3.1-9)	16% (11-22%)	18% (12-25%)	88 (48-140)	2800 (1500-4400)
Swaziland	2000	Male	60 to 64	40% (28-51%)	60% (49-72%)	4.8 (2.5-8.1)	15% (10-20%)	17% (12-23%)	86 (48-130)	2300 (1300-3500)
Swaziland	2000	Male	65 to 69	29% (20-39%)	71% (61-80%)	2.8 (1.5-4.7)	12% (7.5-16%)	13% (8.4-18%)	57 (32-90)	1300 (720-2000)
Swaziland	2000	Male	70 to 74	24% (16-33%)	76% (67-84%)	1.9 (1-3.1)	7.9% (4.5-12%)	8.8% (5.1-13%)	31 (16-50)	570 (300-900)
Swaziland	2000	Male	75 to 79	24% (15-34%)	76% (66-85%)	1.5 (0.76-2.7)	6.1% (2.9-9.9%)	6.8% (3.3-11%)	20 (8.8-33)	280 (120-460)
Swaziland	2000	Male	80 to 84	23% (15-34%)	77% (66-85%)	1.4 (0.68-2.3)	5.3% (2.5-8.9%)	6% (2.8-10%)	10 (4.6-17)	110 (50-180)
Swaziland	2000	Male	85 to 89	22% (14-32%)	78% (68-86%)	1.2 (0.58-2)	4% (1.9-7%)	4.6% (2-8%)	3.6 (1.6-6.5)	27 (12-48)
Swaziland	2000	Male	90 to 94	22% (14-33%)	78% (67-86%)	1 (0.5-1.9)	3% (1.3-5.9%)	3.5% (1.4-6.9%)	0.8 (0.32-1.6)	4.2 (1.8-8.1)
Swaziland	2000	Male	95 plus	24% (13-39%)	76% (61-87%)	0.96 (0.38-2)	2.8% (1-5.9%)	3.1% (1-6.5%)	0.15 (0.048-0.31)	0.53 (0.19-1.1)
Swaziland	2005	Female	All Ages	8.8% (7.3-11%)	91% (89-93%)	0.62 (0.48-0.79)	1.5% (0.91-2.7%)	2.3% (1.3-4%)	240 (140-460)	8600 (5200-16000)
Swaziland	2005	Female	15 to 19	7.8% (5-12%)	92% (88-95%)	0.49 (0.24-0.84)	2.5% (1.2-4.4%)	3.3% (1.3-6.4%)	4.8 (1.9-9.8)	410 (210-780)
Swaziland	2005	Female	20 to 24	10% (6.2-16%)	90% (84-94%)	0.86 (0.46-1.4)	2% (0.87-3.9%)	2% (0.7-4.5%)	8.2 (3.8-18)	670 (360-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	2005	Female	25 to 29	11% (6.5-16%)	89% (84-94%)	0.98 (0.56-1.6)	1% (0.46-2.4%)	0.97% (0.36-2.5%)	8.6 (3.9-19)	620 (330-1300)
Swaziland	2005	Female	30 to 34	13% (7.8-20%)	87% (80-92%)	1.2 (0.64-1.9)	0.91% (0.42-2%)	0.86% (0.35-2%)	9.1 (4-20)	570 (290-1200)
Swaziland	2005	Female	35 to 39	15% (9.6-23%)	85% (77-90%)	1.4 (0.76-2.2)	1.3% (0.59-3%)	1.3% (0.53-3.2%)	12 (5.6-29)	670 (340-1500)
Swaziland	2005	Female	40 to 44	16% (11-23%)	84% (77-89%)	1.4 (0.74-2.3)	1.7% (0.83-3.6%)	1.8% (0.81-3.9%)	16 (7-36)	760 (360-1700)
Swaziland	2005	Female	45 to 49	18% (11-25%)	82% (75-89%)	1.4 (0.72-2.4)	2.5% (1.2-5.1%)	2.7% (1.2-5.6%)	20 (9.1-45)	850 (400-1900)
Swaziland	2005	Female	50 to 54	19% (13-26%)	81% (74-87%)	1.5 (0.82-2.5)	3.5% (1.8-6.8%)	3.8% (2-7.4%)	24 (11-53)	900 (440-1900)
Swaziland	2005	Female	55 to 59	19% (12-27%)	81% (73-88%)	1.6 (0.86-2.7)	4.8% (2.5-8.3%)	5.4% (2.7-9.4%)	27 (12-55)	860 (400-1700)
Swaziland	2005	Female	60 to 64	20% (13-30%)	80% (70-87%)	1.6 (0.82-2.9)	5.6% (2.7-9.7%)	6.4% (3-11%)	29 (12-57)	780 (340-1600)
Swaziland	2005	Female	65 to 69	22% (15-29%)	78% (71-85%)	1.6 (0.83-2.6)	6.1% (3-10%)	7% (3.4-11%)	32 (15-63)	730 (330-1400)
Swaziland	2005	Female	70 to 74	19% (13-26%)	81% (74-87%)	1.2 (0.65-2)	4.7% (2-8.5%)	5.4% (2.3-9.6%)	24 (9.2-50)	440 (170-890)
Swaziland	2005	Female	75 to 79	15% (9.4-22%)	85% (78-91%)	0.87 (0.42-1.6)	3.5% (1.1-6.9%)	4% (1.2-7.9%)	16 (4.5-33)	220 (63-470)
Swaziland	2005	Female	80 to 84	13% (8.2-19%)	87% (81-92%)	0.71 (0.35-1.2)	2.7% (0.86-5.4%)	3.1% (0.99-6.3%)	8.7 (2.7-18)	91 (29-180)
Swaziland	2005	Female	85 to 89	13% (8-19%)	87% (81-92%)	0.65 (0.32-1.1)	2.1% (0.63-4.5%)	2.5% (0.73-5.2%)	3.7 (1.1-7.8)	27 (8-57)
Swaziland	2005	Female	90 to 94	14% (8.8-20%)	86% (80-91%)	0.62 (0.3-1.1)	1.7% (0.33-4%)	1.9% (0.34-4.6%)	1.1 (0.21-2.7)	5.8 (1.2-14)
Swaziland	2005	Female	95 plus	15% (8.5-23%)	85% (77-91%)	0.61 (0.24-1.2)	1.6% (0.0034-4.1%)	1.8% (-0.068-4.7%)	0.25 (-0.0092-0.64)	0.86 (0.002-2.2)
Swaziland	2005	Male	All Ages	17% (14-20%)	83% (80-86%)	1.6 (1.3-2)	5.3% (3.6-7.6%)	7.3% (5-10%)	800 (520-1200)	32000 (22000-47000)
Swaziland	2005	Male	15 to 19	13% (8.3-19%)	87% (81-92%)	0.95 (0.5-1.6)	6.2% (3.5-9.6%)	7.5% (3.9-12%)	16 (7.9-28)	1300 (700-2200)
Swaziland	2005	Male	20 to 24	21% (14-29%)	79% (71-86%)	2.2 (1.2-3.5)	9.8% (5.7-15%)	11% (5.8-17%)	42 (24-68)	3000 (1800-4700)
Swaziland	2005	Male	25 to 29	27% (19-36%)	73% (64-81%)	3.3 (2-4.9)	6.4% (3.5-11%)	6.6% (3.5-12%)	46 (27-72)	3000 (1800-4500)
Swaziland	2005	Male	30 to 34	31% (21-43%)	69% (57-79%)	3.9 (2.3-5.8)	5.1% (3-8.3%)	5.2% (2.9-8.8%)	48 (29-75)	2800 (1700-4300)
Swaziland	2005	Male	35 to 39	38% (28-50%)	62% (50-72%)	4.5 (2.8-6.8)	6.4% (3.6-11%)	6.7% (3.6-11%)	61 (36-97)	3200 (1900-5000)
Swaziland	2005	Male	40 to 44	42% (31-53%)	58% (47-69%)	4.9 (2.7-7.6)	7.7% (4.6-12%)	8.1% (4.7-13%)	72 (42-120)	3400 (2000-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	2005	Male	45 to 49	37% (25-50%)	63% (50-75%)	4.4 (2.4-7)	9.4% (5.6-15%)	9.9% (5.7-16%)	82 (46-140)	3400 (2000-5700)
Swaziland	2005	Male	50 to 54	37% (26-49%)	63% (51-74%)	4.3 (2.4-6.7)	13% (8.3-18%)	14% (8.8-19%)	93 (53-150)	3400 (2000-5500)
Swaziland	2005	Male	55 to 59	43% (32-54%)	57% (46-68%)	4.8 (2.8-7.5)	16% (11-22%)	18% (12-24%)	110 (64-170)	3300 (2100-5200)
Swaziland	2005	Male	60 to 64	40% (29-51%)	60% (49-71%)	4.1 (2.2-6.8)	15% (10-20%)	17% (11-23%)	92 (57-140)	2500 (1600-3700)
Swaziland	2005	Male	65 to 69	29% (20-39%)	71% (61-80%)	2.4 (1.3-4)	11% (7-16%)	13% (7.7-18%)	68 (38-100)	1500 (870-2300)
Swaziland	2005	Male	70 to 74	24% (17-33%)	76% (67-83%)	1.6 (0.85-2.5)	7.5% (4-12%)	8.3% (4.5-13%)	37 (19-60)	670 (350-1100)
Swaziland	2005	Male	75 to 79	24% (16-35%)	76% (65-84%)	1.3 (0.65-2.3)	5.6% (2.6-9.4%)	6.2% (2.9-11%)	21 (9.5-36)	290 (130-500)
Swaziland	2005	Male	80 to 84	24% (15-35%)	76% (65-85%)	1.2 (0.59-2)	4.7% (2.2-8.2%)	5.3% (2.4-9.2%)	11 (5.3-20)	120 (56-200)
Swaziland	2005	Male	85 to 89	23% (14-33%)	77% (67-86%)	0.99 (0.51-1.7)	3.6% (1.6-6.3%)	4% (1.8-7.1%)	3.7 (1.6-6.7)	28 (13-50)
Swaziland	2005	Male	90 to 94	22% (14-33%)	78% (67-86%)	0.87 (0.42-1.6)	2.6% (1.1-5%)	3% (1.2-5.8%)	0.91 (0.37-1.8)	4.8 (2-9.2)
Swaziland	2005	Male	95 plus	25% (14-40%)	75% (60-86%)	0.82 (0.32-1.7)	2.5% (0.89-5.2%)	2.8% (0.87-6%)	0.18 (0.055-0.38)	0.63 (0.22-1.3)
Swaziland	2010	Female	All Ages	9.5% (7.7-11%)	90% (89-92%)	0.57 (0.42-0.76)	1.2% (0.68-2.1%)	1.8% (0.85-3.2%)	170 (72-330)	6400 (3400-12000)
Swaziland	2010	Female	15 to 19	8.1% (5-12%)	92% (88-95%)	0.42 (0.21-0.73)	1.6% (0.78-2.8%)	1.8% (0.73-3.6%)	3.3 (1.3-6.6)	310 (150-570)
Swaziland	2010	Female	20 to 24	12% (7-18%)	88% (82-93%)	0.82 (0.43-1.3)	1.9% (0.86-3.7%)	1.9% (0.65-4.2%)	7 (3-15)	630 (330-1200)
Swaziland	2010	Female	25 to 29	13% (7.9-19%)	87% (81-92%)	0.99 (0.54-1.6)	1.1% (0.49-2.5%)	1.1% (0.37-2.6%)	7.7 (3.4-17)	590 (310-1100)
Swaziland	2010	Female	30 to 34	14% (8.8-22%)	86% (78-91%)	1.1 (0.6-1.9)	0.89% (0.4-1.9%)	0.82% (0.3-2%)	7.2 (3-17)	480 (230-1000)
Swaziland	2010	Female	35 to 39	16% (9.9-24%)	84% (76-90%)	1.2 (0.64-2.1)	1.1% (0.46-2.3%)	1% (0.39-2.4%)	8.1 (3.3-19)	460 (210-1000)
Swaziland	2010	Female	40 to 44	18% (11-26%)	82% (74-89%)	1.3 (0.63-2.2)	1.5% (0.67-3%)	1.5% (0.63-3.3%)	11 (4.2-24)	530 (240-1100)
Swaziland	2010	Female	45 to 49	19% (11-26%)	81% (74-89%)	1.3 (0.61-2.2)	2.1% (0.94-4.2%)	2.2% (0.94-4.6%)	14 (5.7-30)	600 (270-1300)
Swaziland	2010	Female	50 to 54	18% (12-26%)	82% (74-88%)	1.2 (0.62-2.1)	2.8% (1.2-5.4%)	3% (1.3-6%)	16 (6.3-37)	630 (250-1300)
Swaziland	2010	Female	55 to 59	19% (12-27%)	81% (73-88%)	1.3 (0.63-2.3)	3.8% (1.6-7%)	4.4% (1.8-8%)	19 (7-41)	640 (240-1300)
Swaziland	2010	Female	60 to 64	19% (11-28%)	81% (72-89%)	1.3 (0.61-2.3)	4.2% (1.6-8%)	5% (1.9-9.4%)	20 (6.8-42)	560 (190-1200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	2010	Female	65 to 69	19% (13-27%)	81% (73-87%)	1.1 (0.57-1.9)	4.2% (1.5-7.7%)	5% (1.9-9.2%)	20 (6.3-41)	450 (140-930)
Swaziland	2010	Female	70 to 74	17% (11-24%)	83% (76-89%)	0.9 (0.46-1.6)	3.1% (0.81-6.2%)	3.7% (0.97-7.3%)	15 (3.5-35)	280 (64-630)
Swaziland	2010	Female	75 to 79	14% (8.8-21%)	86% (79-91%)	0.67 (0.3-1.3)	2.2% (0.32-5.2%)	2.6% (0.38-6.1%)	9.8 (1.2-24)	140 (18-340)
Swaziland	2010	Female	80 to 84	13% (8-19%)	87% (81-92%)	0.57 (0.27-1)	1.7% (0.23-4%)	2% (0.27-4.6%)	6 (0.74-15)	63 (7.5-150)
Swaziland	2010	Female	85 to 89	13% (8-19%)	87% (81-92%)	0.53 (0.26-0.95)	1.3% (0.034-3.2%)	1.5% (0.023-3.7%)	2.1 (0.031-5.1)	16 (0.38-38)
Swaziland	2010	Female	90 to 94	14% (8.8-20%)	86% (80-91%)	0.51 (0.25-0.97)	0.99% (-0.17-2.9%)	1.1% (-0.23-3.4%)	0.58 (-0.12-1.7)	3 (-0.52-8.7)
Swaziland	2010	Female	95 plus	15% (8.6-24%)	85% (76-91%)	0.5 (0.2-1)	0.9% (-0.35-2.9%)	0.99% (-0.51-3.3%)	0.14 (-0.07-0.49)	0.51 (-0.21-1.7)
Swaziland	2010	Male	All Ages	18% (15-22%)	82% (78-85%)	1.5 (1.1-2)	5.1% (3.5-6.9%)	6.8% (4.4-9.1%)	700 (440-970)	29000 (19000-40000)
Swaziland	2010	Male	15 to 19	15% (9.2-22%)	85% (78-91%)	0.89 (0.46-1.5)	4.7% (2.6-7.6%)	5.4% (2.7-9.1%)	14 (6.8-24)	1200 (630-1900)
Swaziland	2010	Male	20 to 24	25% (16-34%)	75% (66-84%)	2.1 (1.2-3.4)	9.5% (5.5-14%)	11% (5.6-17%)	46 (26-71)	3400 (2100-5100)
Swaziland	2010	Male	25 to 29	31% (20-41%)	69% (59-80%)	3 (1.7-4.7)	7.3% (3.9-12%)	7.7% (3.8-14%)	54 (32-81)	3600 (2200-5100)
Swaziland	2010	Male	30 to 34	34% (22-47%)	66% (53-78%)	3.5 (2-5.6)	5.6% (3.1-8.9%)	5.8% (3.1-9.6%)	49 (28-72)	2900 (1700-4200)
Swaziland	2010	Male	35 to 39	37% (25-50%)	63% (50-75%)	3.6 (2-5.9)	5.8% (3.1-8.9%)	6.1% (3.1-9.5%)	51 (28-76)	2700 (1600-3900)
Swaziland	2010	Male	40 to 44	37% (26-50%)	63% (50-74%)	3.6 (1.9-6)	6.8% (3.8-9.9%)	7.2% (4-11%)	58 (31-89)	2700 (1500-4100)
Swaziland	2010	Male	45 to 49	36% (23-48%)	64% (52-77%)	3.5 (1.7-5.9)	8.5% (4.9-13%)	9.1% (5.1-14%)	66 (35-110)	2800 (1500-4400)
Swaziland	2010	Male	50 to 54	36% (25-48%)	64% (52-75%)	3.5 (1.8-5.8)	11% (6.7-16%)	12% (7.2-17%)	77 (41-120)	2800 (1600-4400)
Swaziland	2010	Male	55 to 59	39% (27-53%)	61% (47-73%)	3.7 (1.8-6.1)	14% (8.7-19%)	15% (9.6-21%)	86 (48-130)	2800 (1600-4100)
Swaziland	2010	Male	60 to 64	36% (25-50%)	64% (50-75%)	3.1 (1.5-5.3)	13% (7.2-18%)	14% (7.9-20%)	78 (40-120)	2100 (1100-3200)
Swaziland	2010	Male	65 to 69	29% (19-39%)	71% (61-81%)	2 (0.99-3.3)	9.6% (5-14%)	11% (5.6-16%)	57 (28-88)	1300 (650-2000)
Swaziland	2010	Male	70 to 74	24% (17-33%)	76% (67-83%)	1.3 (0.65-2.2)	6% (2.8-9.8%)	6.7% (3.1-11%)	34 (15-56)	610 (270-1000)
Swaziland	2010	Male	75 to 79	24% (15-34%)	76% (66-85%)	1.1 (0.48-1.9)	4.2% (1.7-7.8%)	4.7% (1.9-8.7%)	17 (6.7-33)	240 (97-460)
Swaziland	2010	Male	80 to 84	23% (14-33%)	77% (67-86%)	0.91 (0.43-1.6)	3.3% (1.3-6.3%)	3.7% (1.4-7.1%)	8.8 (3.5-17)	94 (37-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	2010	Male	85 to 89	21% (13-31%)	79% (69-87%)	0.76 (0.36-1.3)	2.4% (0.99-4.7%)	2.7% (1.1-5.3%)	2.5 (1-5)	19 (7.9-37)
Swaziland	2010	Male	90 to 94	21% (13-32%)	79% (68-87%)	0.67 (0.31-1.3)	1.8% (0.67-3.9%)	2% (0.69-4.5%)	0.59 (0.2-1.3)	3.2 (1.2-6.8)
Swaziland	2010	Male	95 plus	24% (13-38%)	76% (62-87%)	0.64 (0.25-1.3)	1.8% (0.54-4%)	1.9% (0.47-4.5%)	0.14 (0.033-0.33)	0.52 (0.15-1.2)
Swaziland	2016	Female	All Ages	9.5% (7.8-11%)	90% (89-92%)	0.6 (0.34-0.95)	1.6% (0.68-2.9%)	2.4% (0.82-4.6%)	140 (38-300)	5400 (2000-11000)
Swaziland	2016	Female	15 to 19	8% (5-12%)	92% (88-95%)	0.43 (0.19-0.86)	1.4% (0.64-2.6%)	1.5% (0.44-3.3%)	2.1 (0.54-4.8)	220 (97-430)
Swaziland	2016	Female	20 to 24	12% (7-18%)	88% (82-93%)	0.83 (0.38-1.5)	1.9% (0.9-3.6%)	1.7% (0.48-4%)	4.8 (1.4-11)	500 (230-940)
Swaziland	2016	Female	25 to 29	13% (7.8-19%)	87% (81-92%)	1 (0.46-1.8)	1.5% (0.64-3%)	1.4% (0.38-3.3%)	6.3 (1.8-15)	550 (250-1100)
Swaziland	2016	Female	30 to 34	14% (8.7-22%)	86% (78-91%)	1.1 (0.53-2)	1.5% (0.64-3%)	1.5% (0.43-3.5%)	6.6 (1.7-16)	480 (200-1000)
Swaziland	2016	Female	35 to 39	16% (9.8-23%)	84% (77-90%)	1.2 (0.56-2.4)	1.8% (0.73-3.8%)	2% (0.57-4.4%)	6.7 (1.7-16)	410 (140-890)
Swaziland	2016	Female	40 to 44	18% (11-25%)	82% (75-89%)	1.3 (0.55-2.5)	2.3% (0.83-4.7%)	2.6% (0.8-5.6%)	7.2 (1.8-17)	370 (120-810)
Swaziland	2016	Female	45 to 49	18% (11-26%)	82% (74-89%)	1.3 (0.54-2.6)	3.1% (1.2-6.3%)	3.7% (1.3-7.7%)	9.3 (2.4-22)	420 (130-950)
Swaziland	2016	Female	50 to 54	18% (12-26%)	82% (74-88%)	1.3 (0.53-2.4)	3.5% (1.3-7%)	4.3% (1.5-8.6%)	12 (3.1-28)	470 (140-1100)
Swaziland	2016	Female	55 to 59	18% (12-27%)	82% (73-88%)	1.3 (0.53-2.5)	4.5% (1.7-8.5%)	5.9% (2.1-11%)	15 (4-36)	520 (150-1200)
Swaziland	2016	Female	60 to 64	19% (11-28%)	81% (72-89%)	1.3 (0.52-2.6)	4.5% (1.4-9.2%)	5.9% (1.9-12%)	18 (4.3-41)	500 (130-1100)
Swaziland	2016	Female	65 to 69	19% (12-27%)	81% (73-88%)	1.2 (0.49-2.3)	4.3% (1.1-8.6%)	5.6% (1.5-11%)	18 (3.6-43)	430 (84-1000)
Swaziland	2016	Female	70 to 74	17% (11-25%)	83% (75-89%)	0.94 (0.39-1.9)	3.2% (0.47-7.1%)	4% (0.63-8.7%)	14 (1.8-33)	260 (30-600)
Swaziland	2016	Female	75 to 79	14% (8.4-21%)	86% (79-92%)	0.7 (0.27-1.5)	2.3% (0.12-5.8%)	2.8% (0.16-7%)	9.6 (0.5-26)	140 (6.5-370)
Swaziland	2016	Female	80 to 84	13% (7.9-19%)	87% (81-92%)	0.6 (0.23-1.2)	1.8% (-0.029-4.8%)	2.1% (-0.048-5.7%)	6.3 (-0.14-19)	67 (-0.95-200)
Swaziland	2016	Female	85 to 89	13% (7.8-19%)	87% (81-92%)	0.56 (0.23-1.1)	1.4% (-0.085-4%)	1.7% (-0.11-4.7%)	2.6 (-0.12-7.9)	20 (-0.9-59)
Swaziland	2016	Female	90 to 94	13% (8.7-20%)	87% (80-91%)	0.54 (0.2-1.1)	1.1% (-0.21-3.8%)	1.3% (-0.28-4.5%)	0.64 (-0.13-2.3)	3.4 (-0.58-11)
Swaziland	2016	Female	95 plus	15% (8.2-23%)	85% (77-92%)	0.53 (0.18-1.2)	1% (-0.4-3.8%)	1.2% (-0.55-4.4%)	0.14 (-0.067-0.54)	0.5 (-0.19-1.8)
Swaziland	2016	Male	All Ages	19% (15-23%)	81% (77-85%)	1.6 (0.87-2.6)	5.7% (3.5-7.9%)	7.4% (4.4-10%)	590 (320-890)	25000 (15000-37000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Swaziland	2016	Male	15 to 19	15% (9.3-21%)	85% (79-91%)	0.88 (0.39-1.7)	4% (2-6.7%)	4.4% (1.9-8%)	10 (4.3-19)	870 (440-1500)
Swaziland	2016	Male	20 to 24	24% (15-34%)	76% (66-85%)	2.1 (0.99-3.8)	8.7% (5.1-13%)	9.5% (5-15%)	37 (19-59)	2800 (1600-4300)
Swaziland	2016	Male	25 to 29	30% (19-42%)	70% (58-81%)	3 (1.4-5.5)	7.5% (4.1-12%)	8% (3.9-13%)	53 (29-82)	3600 (2100-5300)
Swaziland	2016	Male	30 to 34	34% (22-47%)	66% (53-78%)	3.5 (1.7-6.4)	6.7% (3.7-10%)	7.1% (3.7-11%)	52 (28-80)	3200 (1800-4700)
Swaziland	2016	Male	35 to 39	37% (24-49%)	63% (51-76%)	3.6 (1.7-6.7)	7.3% (3.9-11%)	8% (4.1-12%)	47 (23-74)	2500 (1300-3900)
Swaziland	2016	Male	40 to 44	37% (26-50%)	63% (50-74%)	3.6 (1.6-6.9)	8.9% (4.9-13%)	9.8% (5.3-15%)	45 (23-75)	2200 (1100-3500)
Swaziland	2016	Male	45 to 49	35% (22-48%)	65% (52-78%)	3.5 (1.5-6.8)	11% (6.5-16%)	12% (7.1-18%)	51 (26-83)	2100 (1100-3400)
Swaziland	2016	Male	50 to 54	36% (24-48%)	64% (52-76%)	3.4 (1.5-6.4)	12% (7.4-18%)	14% (8.2-20%)	56 (28-92)	2100 (1100-3400)
Swaziland	2016	Male	55 to 59	39% (27-52%)	61% (48-73%)	3.6 (1.6-6.6)	14% (8-20%)	17% (9.1-23%)	63 (31-100)	2000 (1100-3300)
Swaziland	2016	Male	60 to 64	36% (24-49%)	64% (51-76%)	3 (1.2-5.9)	12% (6-18%)	14% (7-21%)	62 (27-100)	1700 (770-2700)
Swaziland	2016	Male	65 to 69	28% (19-39%)	72% (61-81%)	1.9 (0.81-3.7)	9.2% (4.1-15%)	11% (4.8-17%)	50 (21-82)	1100 (470-1900)
Swaziland	2016	Male	70 to 74	24% (16-34%)	76% (66-84%)	1.3 (0.53-2.5)	5.8% (2.2-10%)	6.6% (2.5-12%)	30 (11-55)	550 (200-1000)
Swaziland	2016	Male	75 to 79	24% (15-34%)	76% (66-85%)	1 (0.42-2.2)	4.1% (1.2-8.3%)	4.6% (1.4-9.4%)	18 (5.1-39)	260 (74-550)
Swaziland	2016	Male	80 to 84	23% (14-33%)	77% (67-86%)	0.91 (0.36-1.8)	3.2% (0.88-6.5%)	3.6% (0.96-7.4%)	9.4 (2.5-20)	100 (27-210)
Swaziland	2016	Male	85 to 89	21% (13-31%)	79% (69-87%)	0.75 (0.29-1.5)	2.4% (0.64-5.2%)	2.6% (0.68-5.9%)	2.7 (0.71-6.1)	21 (5.7-46)
Swaziland	2016	Male	90 to 94	21% (13-31%)	79% (69-87%)	0.67 (0.24-1.4)	1.8% (0.45-4.4%)	2% (0.44-5%)	0.58 (0.13-1.4)	3.1 (0.78-7.4)
Swaziland	2016	Male	95 plus	23% (13-37%)	77% (63-87%)	0.64 (0.2-1.4)	1.8% (0.4-4.7%)	1.9% (0.35-5.3%)	0.13 (0.024-0.36)	0.49 (0.11-1.3)
Zimbabwe	1990	Female	All Ages	8.3% (7-9.7%)	92% (90-93%)	0.31 (0.24-0.39)	0.99% (0.7-1.3%)	1.6% (1-2.3%)	580 (340-850)	22000 (15000-30000)
Zimbabwe	1990	Female	15 to 19	7.2% (4.6-11%)	93% (89-95%)	0.26 (0.14-0.43)	0.91% (0.56-1.4%)	0.87% (0.38-1.6%)	6.3 (2.9-12)	900 (550-1400)
Zimbabwe	1990	Female	20 to 24	9.8% (6.2-14%)	90% (86-94%)	0.47 (0.26-0.77)	1.3% (0.78-2%)	0.92% (0.41-1.7%)	13 (6.4-22)	1800 (1200-2600)
Zimbabwe	1990	Female	25 to 29	12% (7.2-17%)	88% (83-93%)	0.6 (0.34-0.96)	1.3% (0.76-2%)	1% (0.44-2%)	17 (8.8-29)	1900 (1200-2800)
Zimbabwe	1990	Female	30 to 34	13% (8.1-20%)	87% (80-92%)	0.68 (0.37-1.1)	1.5% (0.87-2.3%)	1.4% (0.65-2.5%)	22 (12-37)	1900 (1200-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	1990	Female	35 to 39	15% (9.3-21%)	85% (79-91%)	0.71 (0.38-1.1)	1.8% (1.1-2.8%)	1.9% (0.94-3.3%)	29 (16-48)	1900 (1200-2900)
Zimbabwe	1990	Female	40 to 44	16% (10-23%)	84% (77-90%)	0.73 (0.4-1.2)	2.2% (1.3-3.3%)	2.6% (1.4-4.2%)	31 (18-50)	1700 (1000-2500)
Zimbabwe	1990	Female	45 to 49	17% (9.8-23%)	83% (77-90%)	0.74 (0.39-1.2)	2.7% (1.6-4%)	3.3% (1.9-5.2%)	43 (23-67)	1900 (1100-2900)
Zimbabwe	1990	Female	50 to 54	16% (11-23%)	84% (77-89%)	0.73 (0.39-1.2)	3.1% (1.8-4.8%)	3.9% (2.2-6.1%)	52 (27-87)	2000 (1100-3300)
Zimbabwe	1990	Female	55 to 59	15% (9.8-22%)	85% (78-90%)	0.72 (0.38-1.2)	3.3% (2-5%)	4.1% (2.5-6.3%)	62 (31-100)	2100 (1100-3400)
Zimbabwe	1990	Female	60 to 64	16% (10-23%)	84% (77-90%)	0.73 (0.39-1.2)	3.2% (1.8-5.2%)	3.9% (2.2-6.4%)	72 (36-120)	2000 (1000-3300)
Zimbabwe	1990	Female	65 to 69	17% (11-24%)	83% (76-89%)	0.72 (0.41-1.2)	3.2% (1.9-5.2%)	4% (2.4-6.3%)	81 (41-140)	1800 (940-3000)
Zimbabwe	1990	Female	70 to 74	17% (11-24%)	83% (76-89%)	0.61 (0.33-1)	2.4% (1.2-4%)	2.9% (1.4-4.8%)	72 (34-120)	1300 (630-2200)
Zimbabwe	1990	Female	75 to 79	15% (9.1-23%)	85% (77-91%)	0.49 (0.24-0.87)	1.4% (0.46-2.9%)	1.7% (0.54-3.4%)	38 (12-80)	540 (170-1100)
Zimbabwe	1990	Female	80 to 84	14% (8.8-22%)	86% (78-91%)	0.44 (0.24-0.76)	1% (0.24-2.1%)	1.2% (0.27-2.5%)	23 (5.2-50)	240 (57-510)
Zimbabwe	1990	Female	85 to 89	15% (9-23%)	85% (77-91%)	0.44 (0.22-0.75)	0.75% (0.039-1.8%)	0.86% (0.018-2.1%)	9.9 (0.21-24)	75 (3.8-180)
Zimbabwe	1990	Female	90 to 94	16% (10-25%)	84% (75-90%)	0.45 (0.22-0.82)	0.52% (-0.2-1.6%)	0.57% (-0.31-1.8%)	2.5 (-1.3-8.2)	14 (-5.5-42)
Zimbabwe	1990	Female	95 plus	19% (11-31%)	81% (69-89%)	0.47 (0.19-0.95)	0.42% (-0.52-1.8%)	0.42% (-0.71-2%)	0.46 (-0.79-2.2)	1.8 (-2.3-7.6)
Zimbabwe	1990	Male	All Ages	31% (26-36%)	69% (64-74%)	1.6 (1.4-2)	3.4% (1.8-4.8%)	5% (2.5-7.1%)	2200 (980-3300)	96000 (49000-140000)
Zimbabwe	1990	Male	15 to 19	27% (19-36%)	73% (64-81%)	1.3 (0.8-2)	4.4% (2.8-6.4%)	5.7% (3.2-8.9%)	45 (19-76)	4300 (2300-6700)
Zimbabwe	1990	Male	20 to 24	41% (29-52%)	59% (48-71%)	2.9 (1.9-4.2)	7% (3.7-11%)	7.5% (3.2-13%)	140 (61-230)	12000 (6300-17000)
Zimbabwe	1990	Male	25 to 29	47% (34-59%)	53% (41-66%)	3.5 (2.4-4.8)	5.7% (2.7-9.5%)	5.8% (2.2-11%)	170 (75-270)	12000 (6600-18000)
Zimbabwe	1990	Male	30 to 34	50% (37-61%)	50% (39-63%)	3.8 (2.6-5.3)	5.4% (2.5-9%)	5.7% (2.2-10%)	170 (72-260)	11000 (5500-16000)
Zimbabwe	1990	Male	35 to 39	53% (39-65%)	47% (35-61%)	3.9 (2.6-5.6)	6% (2.7-9.9%)	6.6% (2.5-12%)	170 (70-280)	9700 (4600-15000)
Zimbabwe	1990	Male	40 to 44	56% (43-67%)	44% (33-57%)	4.1 (2.6-5.8)	6.5% (3-11%)	7.3% (2.9-13%)	160 (65-250)	7900 (3600-12000)
Zimbabwe	1990	Male	45 to 49	59% (44-70%)	41% (30-56%)	4.5 (2.8-6.4)	9% (4.1-14%)	10% (4.2-17%)	210 (84-330)	9400 (4100-14000)
Zimbabwe	1990	Male	50 to 54	59% (46-70%)	41% (30-54%)	4.2 (2.6-6.1)	10% (5.6-15%)	12% (6.2-18%)	220 (88-340)	8500 (3900-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	1990	Male	55 to 59	56% (43-67%)	44% (33-57%)	3.2 (2-4.6)	9.5% (5.3-14%)	12% (6.3-17%)	200 (79-320)	6700 (3000-10000)
Zimbabwe	1990	Male	60 to 64	54% (40-65%)	46% (35-60%)	2.5 (1.5-3.8)	8.2% (4.5-12%)	10% (5.6-15%)	190 (78-310)	5300 (2300-8300)
Zimbabwe	1990	Male	65 to 69	53% (39-64%)	47% (36-61%)	2.1 (1.3-3)	7.9% (4.7-11%)	9.6% (5.9-14%)	190 (77-290)	4300 (1900-6700)
Zimbabwe	1990	Male	70 to 74	49% (36-62%)	51% (38-64%)	1.7 (1.1-2.6)	6.5% (3.8-9.6%)	7.9% (4.7-12%)	170 (73-280)	3200 (1400-5000)
Zimbabwe	1990	Male	75 to 79	47% (32-63%)	53% (37-68%)	1.5 (0.85-2.3)	4.9% (2.4-7.7%)	5.8% (2.9-9.2%)	110 (43-180)	1500 (620-2500)
Zimbabwe	1990	Male	80 to 84	43% (29-58%)	57% (42-71%)	1.2 (0.7-1.8)	3.5% (1.6-5.6%)	4.1% (2-6.6%)	55 (24-96)	570 (260-990)
Zimbabwe	1990	Male	85 to 89	39% (26-52%)	61% (48-74%)	0.91 (0.5-1.4)	2.1% (0.81-3.6%)	2.4% (0.88-4.4%)	19 (7-35)	140 (55-260)
Zimbabwe	1990	Male	90 to 94	36% (25-49%)	64% (51-75%)	0.71 (0.38-1.2)	1.1% (0.27-2.3%)	1.2% (0.21-2.6%)	4 (0.62-8.7)	22 (5.3-46)
Zimbabwe	1990	Male	95 plus	37% (23-53%)	63% (47-77%)	0.58 (0.25-1.1)	0.76% (-0.053-2%)	0.73% (-0.24-2.2%)	0.58 (-0.2-1.7)	2.5 (-0.17-6.4)
Zimbabwe	1995	Female	All Ages	8.8% (7.8-10%)	91% (90-92%)	0.27 (0.21-0.33)	0.53% (0.31-0.84%)	0.72% (0.34-1.3%)	420 (190-780)	18000 (11000-29000)
Zimbabwe	1995	Female	15 to 19	6.2% (4.6-8.4%)	94% (92-95%)	0.18 (0.1-0.28)	0.68% (0.4-1%)	0.48% (0.17-1%)	4.5 (1.7-9.7)	850 (500-1300)
Zimbabwe	1995	Female	20 to 24	9.7% (7.4-12%)	90% (88-93%)	0.37 (0.23-0.56)	0.69% (0.34-1.1%)	0.35% (0.11-0.83%)	10 (3.9-20)	1700 (1000-2500)
Zimbabwe	1995	Female	25 to 29	12% (9.2-15%)	88% (85-91%)	0.5 (0.32-0.76)	0.48% (0.24-0.87%)	0.28% (0.089-0.67%)	14 (5.4-28)	1700 (1000-2600)
Zimbabwe	1995	Female	30 to 34	15% (11-19%)	85% (81-89%)	0.62 (0.38-0.93)	0.54% (0.27-0.95%)	0.39% (0.13-0.86%)	19 (7.3-38)	1700 (920-2900)
Zimbabwe	1995	Female	35 to 39	18% (14-22%)	82% (78-86%)	0.68 (0.41-1)	0.71% (0.32-1.4%)	0.62% (0.22-1.4%)	26 (11-54)	1800 (950-3200)
Zimbabwe	1995	Female	40 to 44	19% (15-23%)	81% (77-85%)	0.69 (0.42-1)	0.9% (0.43-1.6%)	0.88% (0.34-1.8%)	32 (13-66)	1800 (890-3300)
Zimbabwe	1995	Female	45 to 49	20% (15-25%)	80% (75-85%)	0.72 (0.43-1.1)	1.1% (0.51-2.1%)	1.2% (0.47-2.3%)	32 (13-65)	1500 (700-2800)
Zimbabwe	1995	Female	50 to 54	18% (12-25%)	82% (75-88%)	0.66 (0.36-1.1)	1.5% (0.7-2.7%)	1.6% (0.71-3.1%)	41 (16-84)	1600 (710-3200)
Zimbabwe	1995	Female	55 to 59	15% (9.7-22%)	85% (78-90%)	0.58 (0.31-0.95)	1.7% (0.85-3%)	2% (0.91-3.5%)	44 (18-86)	1500 (670-2800)
Zimbabwe	1995	Female	60 to 64	16% (9.9-24%)	84% (76-90%)	0.59 (0.31-1)	1.8% (0.84-3.2%)	2.1% (1-3.8%)	48 (18-99)	1400 (570-2800)
Zimbabwe	1995	Female	65 to 69	18% (12-24%)	82% (76-88%)	0.58 (0.34-0.92)	1.9% (0.91-3.2%)	2.3% (1.2-3.8%)	58 (22-120)	1300 (530-2600)
Zimbabwe	1995	Female	70 to 74	17% (11-24%)	83% (76-89%)	0.49 (0.27-0.79)	1.4% (0.63-2.5%)	1.7% (0.77-3%)	43 (16-89)	790 (290-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	1995	Female	75 to 79	15% (9.2-23%)	85% (77-91%)	0.39 (0.2-0.69)	0.84% (0.14-1.8%)	1% (0.17-2.1%)	28 (4-65)	390 (55-910)
Zimbabwe	1995	Female	80 to 84	15% (8.4-22%)	85% (78-92%)	0.35 (0.18-0.6)	0.56% (-0.05-1.3%)	0.67% (-0.063-1.5%)	15 (-1.2-37)	160 (-12-380)
Zimbabwe	1995	Female	85 to 89	15% (9.1-23%)	85% (77-91%)	0.35 (0.18-0.61)	0.32% (-0.23-0.95%)	0.36% (-0.29-1.1%)	4.7 (-3.7-15)	36 (-27-120)
Zimbabwe	1995	Female	90 to 94	16% (10-24%)	84% (76-90%)	0.35 (0.18-0.64)	0.13% (-0.44-0.8%)	0.1% (-0.58-0.92%)	0.52 (-3.1-4.8)	3.9 (-14-25)
Zimbabwe	1995	Female	95 plus	19% (10-31%)	81% (69-90%)	0.37 (0.16-0.72)	0.038% (-0.74-0.84%)	-0.041% (-0.95-0.89%)	-0.058 (-1.3-1.2)	0.19 (-3.9-4.4)
Zimbabwe	1995	Male	All Ages	36% (32-40%)	64% (60-68%)	1.5 (1.2-1.8)	2.3% (1.3-3.8%)	3% (1.6-4.9%)	2200 (1100-3700)	99000 (55000-160000)
Zimbabwe	1995	Male	15 to 19	35% (28-42%)	65% (58-72%)	1.4 (0.9-1.9)	4.5% (2.9-6.3%)	5.7% (3.4-8.6%)	57 (25-100)	5400 (3000-8600)
Zimbabwe	1995	Male	20 to 24	49% (41-56%)	51% (44-59%)	2.8 (1.9-3.8)	5.8% (2.9-9.9%)	6% (2.5-11%)	170 (79-300)	14000 (7600-23000)
Zimbabwe	1995	Male	25 to 29	54% (47-61%)	46% (39-53%)	3.2 (2.4-4.4)	3.2% (1.5-6.1%)	3.1% (1.2-6.4%)	180 (85-320)	13000 (7100-22000)
Zimbabwe	1995	Male	30 to 34	56% (49-62%)	44% (38-51%)	3.4 (2.5-4.6)	2.7% (1.3-5%)	2.6% (1.1-5.2%)	180 (85-340)	12000 (6200-20000)
Zimbabwe	1995	Male	35 to 39	56% (49-64%)	44% (36-51%)	3.3 (2.3-4.5)	2.9% (1.3-5.7%)	2.9% (1.1-6.2%)	180 (81-340)	10000 (5100-18000)
Zimbabwe	1995	Male	40 to 44	58% (50-67%)	42% (33-50%)	3.4 (2.3-4.7)	3.3% (1.5-6.2%)	3.4% (1.4-6.9%)	200 (86-360)	9600 (4500-17000)
Zimbabwe	1995	Male	45 to 49	62% (53-70%)	38% (30-47%)	3.8 (2.5-5.2)	4.3% (2-7.7%)	4.5% (1.9-8.3%)	200 (86-350)	8500 (4100-15000)
Zimbabwe	1995	Male	50 to 54	61% (52-69%)	39% (31-48%)	3.5 (2.3-5)	5.6% (3-9.2%)	6% (3-10%)	220 (96-390)	8300 (4100-15000)
Zimbabwe	1995	Male	55 to 59	56% (44-67%)	44% (33-56%)	2.5 (1.6-3.7)	5.3% (2.7-8.6%)	5.9% (2.9-9.8%)	180 (79-310)	5900 (2900-10000)
Zimbabwe	1995	Male	60 to 64	54% (39-65%)	46% (35-61%)	1.9 (1.2-3)	4.9% (2.5-8%)	5.6% (2.7-9.3%)	160 (65-280)	4400 (2000-7700)
Zimbabwe	1995	Male	65 to 69	53% (39-64%)	47% (36-61%)	1.6 (1-2.4)	4.9% (2.7-7.5%)	5.7% (3.1-8.8%)	160 (72-280)	3700 (1700-6400)
Zimbabwe	1995	Male	70 to 74	49% (37-62%)	51% (38-63%)	1.4 (0.83-2)	4.2% (2.2-6.4%)	4.9% (2.6-7.5%)	130 (58-230)	2400 (1100-4200)
Zimbabwe	1995	Male	75 to 79	46% (32-62%)	54% (38-68%)	1.2 (0.65-1.8)	3.2% (1.5-5.2%)	3.8% (1.7-6.2%)	100 (40-180)	1400 (580-2500)
Zimbabwe	1995	Male	80 to 84	43% (29-58%)	57% (42-71%)	0.93 (0.54-1.4)	2.4% (1-4.2%)	2.9% (1.2-4.9%)	51 (20-93)	530 (210-960)
Zimbabwe	1995	Male	85 to 89	39% (26-52%)	61% (48-74%)	0.71 (0.4-1.1)	1.4% (0.44-2.6%)	1.6% (0.45-3%)	16 (4.4-30)	120 (37-220)
Zimbabwe	1995	Male	90 to 94	36% (25-50%)	64% (50-75%)	0.56 (0.3-0.95)	0.72% (0.028-1.6%)	0.77% (-0.067-1.8%)	3 (-0.27-7.2)	17 (0.58-38)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	1995	Male	95 plus	37% (23-54%)	63% (46-77%)	0.46 (0.21-0.89)	0.52% (-0.15-1.2%)	0.45% (-0.35-1.3%)	0.43 (-0.35-1.3)	2 (-0.59-5)
Zimbabwe	2000	Female	All Ages	8.2% (7.1-9.6%)	92% (90-93%)	0.25 (0.2-0.32)	0.55% (0.32-0.91%)	0.74% (0.36-1.3%)	670 (300-1200)	27000 (15000-45000)
Zimbabwe	2000	Female	15 to 19	6.8% (4.6-9.7%)	93% (90-95%)	0.2 (0.11-0.33)	0.83% (0.46-1.4%)	0.74% (0.27-1.6%)	9.8 (3.7-22)	1300 (750-2300)
Zimbabwe	2000	Female	20 to 24	8.9% (5.9-12%)	91% (88-94%)	0.36 (0.2-0.56)	0.81% (0.39-1.4%)	0.52% (0.17-1.2%)	19 (7.8-39)	2500 (1500-4000)
Zimbabwe	2000	Female	25 to 29	11% (7.5-15%)	89% (85-93%)	0.47 (0.27-0.73)	0.43% (0.21-0.88%)	0.29% (0.092-0.71%)	24 (9.5-51)	2400 (1300-4200)
Zimbabwe	2000	Female	30 to 34	13% (9-18%)	87% (82-91%)	0.56 (0.32-0.89)	0.41% (0.2-0.79%)	0.32% (0.11-0.73%)	29 (11-63)	2300 (1200-4100)
Zimbabwe	2000	Female	35 to 39	15% (10-20%)	85% (80-90%)	0.59 (0.34-0.92)	0.57% (0.25-1.2%)	0.51% (0.18-1.2%)	40 (16-91)	2500 (1200-4900)
Zimbabwe	2000	Female	40 to 44	17% (12-21%)	83% (79-88%)	0.62 (0.35-0.97)	0.75% (0.34-1.4%)	0.73% (0.29-1.5%)	50 (21-110)	2600 (1200-5100)
Zimbabwe	2000	Female	45 to 49	17% (12-23%)	83% (77-88%)	0.64 (0.36-1)	1% (0.48-2%)	1.1% (0.43-2.2%)	60 (25-120)	2700 (1200-5300)
Zimbabwe	2000	Female	50 to 54	16% (11-22%)	84% (78-89%)	0.61 (0.34-1)	1.3% (0.63-2.4%)	1.4% (0.62-2.6%)	56 (23-110)	2200 (990-4200)
Zimbabwe	2000	Female	55 to 59	14% (9.2-20%)	86% (80-91%)	0.56 (0.3-0.91)	1.7% (0.89-3%)	1.9% (0.92-3.3%)	70 (30-140)	2300 (1000-4400)
Zimbabwe	2000	Female	60 to 64	15% (9.7-22%)	85% (78-90%)	0.58 (0.31-0.97)	1.8% (0.87-3.2%)	2% (0.95-3.7%)	73 (29-140)	2000 (860-3900)
Zimbabwe	2000	Female	65 to 69	18% (12-25%)	82% (75-88%)	0.61 (0.35-0.97)	2.1% (0.95-3.6%)	2.4% (1.1-4.2%)	92 (37-180)	2100 (870-4100)
Zimbabwe	2000	Female	70 to 74	17% (12-24%)	83% (76-88%)	0.52 (0.29-0.87)	1.5% (0.58-2.9%)	1.7% (0.66-3.3%)	73 (24-160)	1300 (440-2800)
Zimbabwe	2000	Female	75 to 79	15% (9.3-23%)	85% (77-91%)	0.41 (0.2-0.72)	0.88% (0.057-2%)	1% (0.074-2.3%)	39 (2.7-95)	550 (36-1300)
Zimbabwe	2000	Female	80 to 84	15% (9-22%)	85% (78-91%)	0.38 (0.2-0.65)	0.58% (-0.072-1.5%)	0.67% (-0.092-1.7%)	22 (-3.2-60)	230 (-33-620)
Zimbabwe	2000	Female	85 to 89	16% (9.4-23%)	84% (77-91%)	0.38 (0.19-0.65)	0.33% (-0.31-1.1%)	0.37% (-0.38-1.3%)	7.4 (-7.1-26)	57 (-49-190)
Zimbabwe	2000	Female	90 to 94	17% (11-25%)	83% (75-89%)	0.38 (0.19-0.67)	0.14% (-0.5-0.96%)	0.12% (-0.64-1.1%)	0.84 (-4.5-7.7)	5.9 (-21-40)
Zimbabwe	2000	Female	95 plus	20% (11-32%)	80% (68-89%)	0.4 (0.17-0.79)	0.056% (-0.78-1%)	-0.015% (-1-1.1%)	-0.03 (-1.9-2.1)	0.41 (-5.6-7.6)
Zimbabwe	2000	Male	All Ages	34% (30-39%)	66% (61-70%)	1.5 (1.2-1.8)	3.6% (2.4-5%)	4.5% (2.9-6.3%)	4600 (2900-6600)	200000 (130000-270000)
Zimbabwe	2000	Male	15 to 19	30% (23-37%)	70% (63-77%)	1.2 (0.82-1.8)	5.2% (3.4-7.4%)	6.4% (4-9.5%)	110 (61-170)	9500 (5900-14000)
Zimbabwe	2000	Male	20 to 24	45% (36-54%)	55% (46-64%)	2.6 (1.8-3.5)	9.4% (5.5-14%)	10% (5.5-16%)	370 (230-540)	27000 (18000-38000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	2000	Male	25 to 29	51% (42-59%)	49% (41-58%)	3.2 (2.2-4.2)	5% (2.7-8.9%)	5.1% (2.5-9.6%)	390 (240-570)	26000 (17000-37000)
Zimbabwe	2000	Male	30 to 34	53% (44-61%)	47% (39-56%)	3.4 (2.3-4.5)	3.6% (2-6%)	3.6% (1.9-6.2%)	370 (220-550)	22000 (14000-32000)
Zimbabwe	2000	Male	35 to 39	55% (47-65%)	45% (35-53%)	3.4 (2.4-4.7)	3.9% (2.2-6.7%)	4% (2.1-7.2%)	390 (240-590)	21000 (13000-31000)
Zimbabwe	2000	Male	40 to 44	58% (48-68%)	42% (32-52%)	3.6 (2.4-5)	4.6% (2.6-7.2%)	4.8% (2.7-7.7%)	430 (240-660)	20000 (12000-31000)
Zimbabwe	2000	Male	45 to 49	62% (52-71%)	38% (29-48%)	4 (2.6-5.6)	6.3% (3.6-10%)	6.7% (3.7-11%)	500 (280-770)	21000 (12000-32000)
Zimbabwe	2000	Male	50 to 54	63% (54-72%)	37% (28-46%)	3.8 (2.4-5.5)	7.5% (4.5-11%)	8% (4.7-12%)	420 (230-660)	16000 (9000-24000)
Zimbabwe	2000	Male	55 to 59	58% (47-67%)	42% (33-53%)	2.8 (1.8-4)	7.7% (4.7-12%)	8.5% (5.1-13%)	400 (210-630)	13000 (7100-20000)
Zimbabwe	2000	Male	60 to 64	53% (41-64%)	47% (36-59%)	2 (1.2-3.1)	6.4% (3.6-10%)	7.1% (3.9-11%)	330 (160-560)	8900 (4500-15000)
Zimbabwe	2000	Male	65 to 69	51% (38-62%)	49% (38-62%)	1.7 (1.1-2.5)	6.3% (3.5-9.5%)	7% (4-11%)	320 (160-510)	7100 (3700-11000)
Zimbabwe	2000	Male	70 to 74	50% (38-62%)	50% (38-62%)	1.4 (0.89-2.2)	5.3% (2.7-8.2%)	6% (3-9.4%)	270 (130-450)	4800 (2300-8000)
Zimbabwe	2000	Male	75 to 79	49% (35-63%)	51% (37-65%)	1.3 (0.72-1.9)	4% (1.7-6.7%)	4.6% (1.9-7.8%)	180 (65-310)	2400 (940-4200)
Zimbabwe	2000	Male	80 to 84	44% (30-60%)	56% (40-70%)	1 (0.57-1.6)	3% (0.99-5.2%)	3.4% (1.1-5.9%)	95 (30-170)	980 (330-1800)
Zimbabwe	2000	Male	85 to 89	39% (26-52%)	61% (48-74%)	0.76 (0.42-1.2)	1.7% (0.48-3.3%)	1.9% (0.48-3.7%)	30 (7.7-60)	230 (64-440)
Zimbabwe	2000	Male	90 to 94	37% (25-50%)	63% (50-75%)	0.6 (0.31-0.99)	0.9% (0.078-1.9%)	0.97% (0.0033-2.1%)	5.3 (0.017-12)	29 (2.6-62)
Zimbabwe	2000	Male	95 plus	37% (23-53%)	63% (47-77%)	0.49 (0.21-0.94)	0.69% (-0.057-1.6%)	0.66% (-0.24-1.8%)	0.84 (-0.3-2.3)	3.5 (-0.29-8.5)
Zimbabwe	2005	Female	All Ages	8% (6.7-9.4%)	92% (91-93%)	0.28 (0.22-0.35)	0.68% (0.38-1.1%)	0.97% (0.5-1.6%)	1000 (480-1700)	38000 (21000-60000)
Zimbabwe	2005	Female	15 to 19	7.1% (4.5-10%)	93% (90-95%)	0.24 (0.13-0.39)	0.86% (0.47-1.5%)	0.79% (0.31-1.6%)	16 (5.9-33)	1800 (960-3100)
Zimbabwe	2005	Female	20 to 24	8.8% (5.6-13%)	91% (87-94%)	0.4 (0.22-0.64)	1.2% (0.57-2.1%)	0.92% (0.31-2.1%)	33 (13-68)	3600 (2000-6100)
Zimbabwe	2005	Female	25 to 29	10% (6.7-15%)	90% (85-93%)	0.5 (0.29-0.81)	0.62% (0.28-1.2%)	0.48% (0.15-1.1%)	40 (15-85)	3600 (1800-6200)
Zimbabwe	2005	Female	30 to 34	12% (7.7-18%)	88% (82-92%)	0.58 (0.32-0.94)	0.47% (0.21-0.94%)	0.39% (0.13-0.89%)	41 (14-87)	3000 (1400-5600)
Zimbabwe	2005	Female	35 to 39	14% (8.4-19%)	86% (81-92%)	0.61 (0.33-0.95)	0.6% (0.24-1.2%)	0.56% (0.18-1.2%)	50 (18-110)	3000 (1300-6000)
Zimbabwe	2005	Female	40 to 44	15% (10-21%)	85% (79-90%)	0.63 (0.35-1)	0.79% (0.34-1.5%)	0.78% (0.28-1.6%)	60 (22-120)	3000 (1300-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	2005	Female	45 to 49	16% (11-22%)	84% (78-89%)	0.67 (0.36-1.1)	1.2% (0.5-2.3%)	1.2% (0.47-2.4%)	76 (30-150)	3300 (1500-6200)
Zimbabwe	2005	Female	50 to 54	16% (11-22%)	84% (78-89%)	0.67 (0.37-1.1)	1.7% (0.79-3%)	1.8% (0.79-3.2%)	91 (37-170)	3500 (1500-6400)
Zimbabwe	2005	Female	55 to 59	14% (9.3-20%)	86% (80-91%)	0.64 (0.35-1.1)	2.2% (1.1-3.7%)	2.4% (1.2-4.1%)	89 (38-170)	2900 (1300-5400)
Zimbabwe	2005	Female	60 to 64	15% (9.8-22%)	85% (78-90%)	0.67 (0.36-1.1)	2.5% (1.2-4.2%)	2.8% (1.4-4.8%)	110 (46-220)	3100 (1300-5900)
Zimbabwe	2005	Female	65 to 69	18% (12-25%)	82% (75-88%)	0.7 (0.4-1.1)	2.8% (1.4-4.8%)	3.2% (1.6-5.5%)	140 (62-260)	3200 (1400-5700)
Zimbabwe	2005	Female	70 to 74	18% (12-25%)	82% (75-88%)	0.6 (0.33-1)	2.1% (0.91-4%)	2.4% (1.1-4.6%)	120 (44-240)	2100 (790-4200)
Zimbabwe	2005	Female	75 to 79	16% (9.2-24%)	84% (76-91%)	0.48 (0.24-0.85)	1.3% (0.28-2.8%)	1.5% (0.32-3.2%)	74 (14-170)	1000 (200-2400)
Zimbabwe	2005	Female	80 to 84	15% (9.3-23%)	85% (77-91%)	0.44 (0.23-0.77)	0.91% (0.16-2%)	1% (0.19-2.2%)	45 (7.8-100)	460 (79-1000)
Zimbabwe	2005	Female	85 to 89	16% (9.7-24%)	84% (76-90%)	0.43 (0.22-0.73)	0.63% (-0.038-1.6%)	0.71% (-0.053-1.8%)	16 (-1.3-41)	120 (-6.5-300)
Zimbabwe	2005	Female	90 to 94	17% (11-26%)	83% (74-89%)	0.44 (0.22-0.75)	0.39% (-0.24-1.3%)	0.42% (-0.33-1.5%)	3.6 (-2.9-13)	20 (-12-65)
Zimbabwe	2005	Female	95 plus	20% (11-32%)	80% (68-89%)	0.45 (0.2-0.89)	0.29% (-0.48-1.4%)	0.27% (-0.66-1.5%)	0.56 (-1.4-3.3)	2.4 (-4-12)
Zimbabwe	2005	Male	All Ages	31% (27-35%)	69% (65-73%)	1.6 (1.3-1.8)	5% (3.7-6.4%)	6.5% (4.8-8.5%)	7300 (5300-9600)	300000 (230000-390000)
Zimbabwe	2005	Male	15 to 19	24% (17-32%)	76% (68-83%)	1.1 (0.69-1.6)	4.9% (3.1-7.3%)	5.6% (3.2-8.6%)	150 (83-220)	12000 (7500-18000)
Zimbabwe	2005	Male	20 to 24	38% (29-48%)	62% (52-71%)	2.6 (1.8-3.5)	13% (8.2-18%)	15% (8.7-21%)	540 (360-750)	39000 (26000-53000)
Zimbabwe	2005	Male	25 to 29	46% (37-54%)	54% (46-63%)	3.3 (2.3-4.3)	8.6% (5-14%)	9.1% (4.9-16%)	610 (420-820)	40000 (28000-53000)
Zimbabwe	2005	Male	30 to 34	48% (38-58%)	52% (42-62%)	3.4 (2.4-4.7)	5.6% (3.4-8.9%)	5.7% (3.4-9.4%)	560 (380-770)	33000 (23000-44000)
Zimbabwe	2005	Male	35 to 39	52% (44-62%)	48% (38-56%)	3.6 (2.5-5)	5.4% (3.2-8.6%)	5.7% (3.2-9.3%)	560 (360-800)	29000 (19000-41000)
Zimbabwe	2005	Male	40 to 44	56% (46-66%)	44% (34-54%)	3.9 (2.6-5.4)	6.4% (4.2-9.2%)	6.8% (4.3-9.8%)	620 (410-870)	29000 (19000-40000)
Zimbabwe	2005	Male	45 to 49	62% (51-71%)	38% (29-49%)	4.5 (3-6.4)	9% (5.9-13%)	9.6% (6.2-14%)	750 (510-1100)	31000 (21000-43000)
Zimbabwe	2005	Male	50 to 54	65% (55-73%)	35% (27-45%)	4.4 (2.8-6.3)	12% (7.9-16%)	13% (8.5-17%)	790 (510-1100)	29000 (19000-40000)
Zimbabwe	2005	Male	55 to 59	59% (49-68%)	41% (32-51%)	3.2 (2.1-4.6)	11% (7.5-16%)	13% (8.3-18%)	580 (370-840)	18000 (12000-26000)
Zimbabwe	2005	Male	60 to 64	53% (42-64%)	47% (36-58%)	2.3 (1.4-3.5)	9.9% (5.9-15%)	11% (6.4-17%)	560 (330-870)	15000 (9000-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	2005	Male	65 to 69	51% (40-63%)	49% (37-60%)	1.9 (1.2-2.8)	9.3% (5.6-14%)	10% (6.1-15%)	540 (320-820)	12000 (7200-18000)
Zimbabwe	2005	Male	70 to 74	50% (38-61%)	50% (39-62%)	1.6 (1-2.4)	7.7% (4.1-12%)	8.7% (4.7-13%)	460 (240-700)	8100 (4300-13000)
Zimbabwe	2005	Male	75 to 79	49% (36-63%)	51% (37-64%)	1.4 (0.82-2.3)	6% (2.5-9.9%)	6.7% (2.8-11%)	320 (140-530)	4400 (1900-7400)
Zimbabwe	2005	Male	80 to 84	45% (31-60%)	55% (40-69%)	1.1 (0.66-1.8)	4.3% (1.7-7.4%)	4.9% (1.9-8.4%)	170 (69-300)	1800 (720-3000)
Zimbabwe	2005	Male	85 to 89	39% (26-52%)	61% (48-74%)	0.85 (0.47-1.3)	2.5% (0.88-4.6%)	2.9% (0.98-5.3%)	51 (17-92)	370 (130-670)
Zimbabwe	2005	Male	90 to 94	37% (25-49%)	63% (51-75%)	0.66 (0.36-1.1)	1.4% (0.34-2.7%)	1.5% (0.32-3.1%)	10 (2.1-20)	53 (13-100)
Zimbabwe	2005	Male	95 plus	37% (23-53%)	63% (47-77%)	0.54 (0.23-1)	1% (0.2-2.4%)	1.1% (0.072-2.6%)	1.6 (0.1-3.8)	6 (1.1-13)
Zimbabwe	2010	Female	All Ages	9.1% (7.6-11%)	91% (89-92%)	0.3 (0.22-0.39)	0.73% (0.4-1.2%)	1% (0.51-1.7%)	950 (410-1700)	38000 (20000-62000)
Zimbabwe	2010	Female	15 to 19	7.7% (5.1-11%)	92% (89-95%)	0.24 (0.12-0.39)	0.71% (0.36-1.3%)	0.59% (0.2-1.3%)	15 (5-32)	1800 (890-3100)
Zimbabwe	2010	Female	20 to 24	11% (6.5-15%)	89% (85-94%)	0.43 (0.23-0.68)	1.4% (0.67-2.4%)	1.1% (0.39-2.4%)	36 (13-73)	4000 (2200-6700)
Zimbabwe	2010	Female	25 to 29	13% (7.7-18%)	87% (82-92%)	0.54 (0.31-0.89)	0.9% (0.4-1.7%)	0.74% (0.23-1.7%)	49 (18-100)	4300 (2300-7600)
Zimbabwe	2010	Female	30 to 34	14% (8.7-22%)	86% (78-91%)	0.62 (0.35-1)	0.69% (0.3-1.3%)	0.59% (0.19-1.3%)	51 (17-110)	3700 (1700-6800)
Zimbabwe	2010	Female	35 to 39	16% (9.8-23%)	84% (77-90%)	0.65 (0.35-1.1)	0.76% (0.32-1.6%)	0.72% (0.24-1.6%)	55 (19-120)	3300 (1500-6600)
Zimbabwe	2010	Female	40 to 44	17% (11-24%)	83% (76-89%)	0.67 (0.35-1.1)	0.98% (0.42-1.9%)	0.98% (0.36-2%)	57 (20-120)	2900 (1200-5600)
Zimbabwe	2010	Female	45 to 49	18% (11-25%)	82% (75-89%)	0.67 (0.35-1.1)	1.4% (0.6-2.8%)	1.5% (0.56-3%)	68 (24-140)	3000 (1200-5900)
Zimbabwe	2010	Female	50 to 54	17% (11-24%)	83% (76-89%)	0.65 (0.35-1.1)	2% (0.9-3.6%)	2.1% (0.93-4%)	83 (31-170)	3200 (1300-6200)
Zimbabwe	2010	Female	55 to 59	16% (11-22%)	84% (78-89%)	0.65 (0.35-1.1)	2.5% (1.3-4.4%)	2.9% (1.4-5%)	100 (37-200)	3300 (1300-6300)
Zimbabwe	2010	Female	60 to 64	17% (11-24%)	83% (76-89%)	0.66 (0.35-1.1)	2.6% (1.3-4.7%)	3% (1.4-5.5%)	94 (34-180)	2600 (1000-5000)
Zimbabwe	2010	Female	65 to 69	19% (12-26%)	81% (74-88%)	0.65 (0.34-1.1)	2.8% (1.3-4.9%)	3.2% (1.6-5.7%)	130 (49-260)	2900 (1200-5700)
Zimbabwe	2010	Female	70 to 74	18% (12-25%)	82% (75-88%)	0.55 (0.28-0.93)	1.9% (0.76-3.7%)	2.2% (0.88-4.2%)	100 (32-210)	1800 (590-3800)
Zimbabwe	2010	Female	75 to 79	16% (9.7-24%)	84% (76-90%)	0.44 (0.21-0.77)	1.1% (0.17-2.5%)	1.3% (0.19-2.9%)	62 (8-150)	860 (120-2100)
Zimbabwe	2010	Female	80 to 84	15% (9.3-23%)	85% (77-91%)	0.39 (0.2-0.69)	0.72% (0.0076-1.7%)	0.84% (0.018-2%)	39 (0.81-94)	400 (3.5-970)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	2010	Female	85 to 89	16% (9.6-24%)	84% (76-90%)	0.39 (0.2-0.67)	0.46% (-0.17-1.3%)	0.53% (-0.23-1.5%)	12 (-5.3-36)	91 (-35-270)
Zimbabwe	2010	Female	90 to 94	17% (11-26%)	83% (74-89%)	0.39 (0.19-0.67)	0.24% (-0.38-1%)	0.24% (-0.49-1.2%)	1.9 (-4-9.3)	11 (-19-49)
Zimbabwe	2010	Female	95 plus	20% (11-32%)	80% (68-89%)	0.4 (0.17-0.81)	0.15% (-0.57-1.2%)	0.11% (-0.74-1.3%)	0.24 (-1.6-2.9)	1.3 (-4.8-10)
Zimbabwe	2010	Male	All Ages	33% (28-38%)	67% (62-72%)	1.5 (1.2-1.9)	5.4% (3.9-7%)	7.1% (5-9.3%)	7200 (5000-9700)	310000 (220000-400000)
Zimbabwe	2010	Male	15 to 19	28% (20-38%)	72% (62-80%)	1.2 (0.7-1.8)	4.4% (2.6-6.6%)	4.8% (2.6-7.6%)	160 (85-240)	13000 (7700-19000)
Zimbabwe	2010	Male	20 to 24	42% (30-53%)	58% (47-70%)	2.6 (1.7-3.7)	13% (8.4-19%)	15% (9.5-22%)	570 (360-800)	41000 (27000-57000)
Zimbabwe	2010	Male	25 to 29	49% (36-59%)	51% (41-64%)	3.2 (2.2-4.3)	11% (6.6-17%)	12% (6.6-20%)	690 (470-950)	45000 (31000-60000)
Zimbabwe	2010	Male	30 to 34	52% (39-63%)	48% (37-61%)	3.4 (2.3-4.7)	8.1% (5.1-12%)	8.6% (5.1-14%)	660 (430-910)	38000 (26000-53000)
Zimbabwe	2010	Male	35 to 39	54% (42-65%)	46% (35-58%)	3.4 (2.2-5)	7.1% (4.4-11%)	7.5% (4.5-12%)	600 (380-870)	31000 (20000-45000)
Zimbabwe	2010	Male	40 to 44	57% (45-68%)	43% (32-55%)	3.6 (2.3-5.3)	7.7% (5-11%)	8.3% (5.3-12%)	590 (370-870)	27000 (18000-40000)
Zimbabwe	2010	Male	45 to 49	60% (46-71%)	40% (29-54%)	4 (2.4-5.7)	11% (6.7-16%)	12% (7-17%)	670 (420-960)	28000 (18000-39000)
Zimbabwe	2010	Male	50 to 54	60% (48-71%)	40% (29-52%)	3.7 (2.3-5.4)	13% (8-18%)	14% (8.6-20%)	680 (420-990)	25000 (15000-36000)
Zimbabwe	2010	Male	55 to 59	57% (45-68%)	43% (32-55%)	2.8 (1.8-4.1)	12% (7.8-18%)	14% (8.5-20%)	610 (370-930)	19000 (12000-29000)
Zimbabwe	2010	Male	60 to 64	56% (42-67%)	44% (33-58%)	2.2 (1.3-3.5)	10% (6.1-17%)	12% (6.7-18%)	470 (260-750)	13000 (7100-20000)
Zimbabwe	2010	Male	65 to 69	54% (42-65%)	46% (35-58%)	1.8 (1.1-2.8)	9.8% (5.7-15%)	11% (6.4-17%)	540 (300-840)	12000 (6800-19000)
Zimbabwe	2010	Male	70 to 74	51% (38-63%)	49% (37-62%)	1.5 (0.91-2.3)	7.6% (3.7-12%)	8.6% (4.1-13%)	440 (210-710)	7900 (3800-13000)
Zimbabwe	2010	Male	75 to 79	48% (34-63%)	52% (37-66%)	1.3 (0.72-2)	5.5% (1.9-9.3%)	6.2% (2.1-11%)	300 (100-500)	4100 (1400-7000)
Zimbabwe	2010	Male	80 to 84	45% (30-59%)	55% (41-70%)	1 (0.57-1.6)	3.9% (1.3-7%)	4.4% (1.4-7.9%)	160 (53-310)	1700 (550-3100)
Zimbabwe	2010	Male	85 to 89	40% (28-54%)	60% (46-72%)	0.79 (0.44-1.3)	2.3% (0.72-4.5%)	2.6% (0.76-5.1%)	48 (14-95)	350 (110-690)
Zimbabwe	2010	Male	90 to 94	38% (26-51%)	62% (49-74%)	0.62 (0.32-1)	1.2% (0.14-2.6%)	1.4% (0.079-3%)	8.4 (0.5-19)	44 (5.3-94)
Zimbabwe	2010	Male	95 plus	38% (24-55%)	62% (45-76%)	0.51 (0.21-0.97)	0.94% (0.078-2.1%)	0.96% (-0.065-2.4%)	1.6 (-0.11-3.9)	6 (0.5-14)
Zimbabwe	2016	Female	All Ages	9.1% (7.6-11%)	91% (89-92%)	0.32 (0.19-0.52)	0.97% (0.55-1.6%)	1.3% (0.61-2.6%)	890 (380-1800)	37000 (20000-63000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	2016	Female	15 to 19	7.5% (4.9-11%)	93% (89-95%)	0.24 (0.11-0.47)	0.78% (0.4-1.4%)	0.65% (0.21-1.5%)	12 (3.6-25)	1500 (800-2700)
Zimbabwe	2016	Female	20 to 24	10% (6.5-15%)	90% (85-93%)	0.44 (0.2-0.84)	1.5% (0.82-2.5%)	1.1% (0.37-2.4%)	26 (9.2-52)	3400 (1900-5600)
Zimbabwe	2016	Female	25 to 29	12% (7.6-18%)	88% (82-92%)	0.57 (0.27-1.1)	1.3% (0.69-2.3%)	1.1% (0.37-2.3%)	39 (15-77)	3900 (2200-6500)
Zimbabwe	2016	Female	30 to 34	14% (8.7-21%)	86% (79-91%)	0.65 (0.31-1.3)	1.3% (0.67-2.2%)	1.1% (0.44-2.3%)	50 (19-100)	4000 (2200-7000)
Zimbabwe	2016	Female	35 to 39	16% (9.6-22%)	84% (78-90%)	0.68 (0.32-1.3)	1.5% (0.77-2.7%)	1.5% (0.63-3%)	61 (25-130)	3900 (2000-7300)
Zimbabwe	2016	Female	40 to 44	17% (11-24%)	83% (76-89%)	0.7 (0.32-1.4)	1.8% (0.87-3.3%)	2% (0.82-3.9%)	61 (25-130)	3200 (1500-6000)
Zimbabwe	2016	Female	45 to 49	17% (10-25%)	83% (75-90%)	0.7 (0.31-1.4)	2.2% (1-3.9%)	2.5% (1.1-4.8%)	61 (25-120)	2700 (1200-5200)
Zimbabwe	2016	Female	50 to 54	17% (11-24%)	83% (76-89%)	0.68 (0.32-1.3)	2.5% (1.2-4.5%)	2.9% (1.3-5.4%)	70 (27-140)	2700 (1200-5400)
Zimbabwe	2016	Female	55 to 59	16% (10-22%)	84% (78-90%)	0.68 (0.32-1.3)	2.9% (1.4-5.4%)	3.5% (1.7-6.6%)	87 (37-170)	2900 (1300-5600)
Zimbabwe	2016	Female	60 to 64	16% (11-24%)	84% (76-89%)	0.69 (0.32-1.4)	2.8% (1.2-5.5%)	3.4% (1.4-6.8%)	100 (39-220)	2900 (1200-5900)
Zimbabwe	2016	Female	65 to 69	18% (12-25%)	82% (75-88%)	0.68 (0.32-1.3)	2.8% (1.1-5.6%)	3.3% (1.3-6.7%)	110 (38-220)	2400 (860-5000)
Zimbabwe	2016	Female	70 to 74	17% (12-25%)	83% (75-88%)	0.57 (0.27-1.1)	1.9% (0.6-4.3%)	2.3% (0.71-5%)	91 (27-210)	1600 (490-3800)
Zimbabwe	2016	Female	75 to 79	15% (9.7-23%)	85% (77-90%)	0.46 (0.2-0.89)	1.2% (0.1-3%)	1.4% (0.12-3.4%)	60 (5.2-160)	840 (72-2300)
Zimbabwe	2016	Female	80 to 84	15% (9.2-23%)	85% (77-91%)	0.42 (0.19-0.79)	0.78% (-0.056-2.2%)	0.91% (-0.069-2.6%)	40 (-2.8-110)	410 (-28-1200)
Zimbabwe	2016	Female	85 to 89	16% (9.3-24%)	84% (76-91%)	0.41 (0.18-0.81)	0.54% (-0.24-1.8%)	0.62% (-0.31-2.1%)	17 (-8-60)	130 (-55-450)
Zimbabwe	2016	Female	90 to 94	17% (11-25%)	83% (75-89%)	0.42 (0.19-0.84)	0.33% (-0.39-1.6%)	0.35% (-0.5-1.8%)	3.4 (-4.9-17)	19 (-22-87)
Zimbabwe	2016	Female	95 plus	19% (11-31%)	81% (69-89%)	0.43 (0.16-0.94)	0.27% (-0.55-1.7%)	0.24% (-0.74-1.9%)	0.54 (-1.6-4.2)	2.3 (-4.7-15)
Zimbabwe	2016	Male	All Ages	33% (28-39%)	67% (61-72%)	1.7 (1-2.6)	6% (4-8.3%)	7.8% (4.9-11%)	6100 (3700-9200)	270000 (170000-380000)
Zimbabwe	2016	Male	15 to 19	27% (19-37%)	73% (63-81%)	1.2 (0.61-2.3)	4.3% (2.3-7.1%)	4.7% (2.3-8.3%)	120 (57-220)	10000 (5500-17000)
Zimbabwe	2016	Male	20 to 24	42% (30-54%)	58% (46-70%)	2.7 (1.3-4.6)	12% (7.1-18%)	14% (7.5-21%)	440 (240-660)	33000 (19000-48000)
Zimbabwe	2016	Male	25 to 29	48% (35-59%)	52% (41-65%)	3.3 (1.8-5.6)	11% (7-17%)	13% (7.2-20%)	570 (340-820)	38000 (24000-53000)
Zimbabwe	2016	Male	30 to 34	51% (38-62%)	49% (38-62%)	3.6 (2-6.2)	10% (6.3-15%)	12% (6.7-18%)	620 (370-910)	37000 (23000-53000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Zimbabwe	2016	Male	35 to 39	54% (41-65%)	46% (35-59%)	3.6 (2-6.2)	10% (6.5-15%)	12% (7.1-18%)	610 (370-900)	32000 (20000-47000)
Zimbabwe	2016	Male	40 to 44	57% (44-68%)	43% (32-56%)	3.8 (2.1-6.7)	11% (6.9-16%)	13% (7.8-19%)	570 (330-860)	27000 (16000-40000)
Zimbabwe	2016	Male	45 to 49	60% (45-71%)	40% (29-55%)	4.2 (2.1-7.3)	14% (8.4-20%)	16% (9.4-23%)	570 (330-870)	24000 (14000-36000)
Zimbabwe	2016	Male	50 to 54	60% (47-70%)	40% (30-53%)	3.9 (2-6.9)	14% (8.3-21%)	16% (9.2-24%)	530 (300-840)	20000 (11000-31000)
Zimbabwe	2016	Male	55 to 59	57% (44-68%)	43% (32-56%)	2.9 (1.5-5.2)	13% (7.1-19%)	15% (8.2-22%)	460 (240-760)	15000 (7800-24000)
Zimbabwe	2016	Male	60 to 64	55% (41-66%)	45% (34-59%)	2.3 (1.2-4.4)	10% (5-17%)	12% (5.8-20%)	420 (200-780)	12000 (5500-21000)
Zimbabwe	2016	Male	65 to 69	54% (41-64%)	46% (36-59%)	1.9 (0.98-3.6)	9% (4.4-16%)	10% (5.1-18%)	370 (180-680)	8400 (4000-15000)
Zimbabwe	2016	Male	70 to 74	51% (38-63%)	49% (37-62%)	1.6 (0.8-2.8)	6.8% (2.9-12%)	7.8% (3.3-14%)	350 (150-650)	6300 (2600-12000)
Zimbabwe	2016	Male	75 to 79	48% (33-62%)	52% (38-67%)	1.3 (0.64-2.4)	4.8% (1.5-9.3%)	5.5% (1.7-11%)	250 (76-500)	3400 (1100-7000)
Zimbabwe	2016	Male	80 to 84	44% (30-59%)	56% (41-70%)	1.1 (0.51-2)	3.3% (0.97-6.6%)	3.8% (1.1-7.5%)	140 (39-290)	1400 (420-3000)
Zimbabwe	2016	Male	85 to 89	39% (27-53%)	61% (47-73%)	0.82 (0.38-1.5)	2% (0.48-4.6%)	2.3% (0.49-5.2%)	47 (10-110)	350 (81-790)
Zimbabwe	2016	Male	90 to 94	37% (26-50%)	63% (50-74%)	0.64 (0.29-1.2)	1.1% (0.15-2.8%)	1.3% (0.1-3.1%)	9 (0.74-23)	48 (6.4-120)
Zimbabwe	2016	Male	95 plus	38% (24-55%)	62% (45-76%)	0.53 (0.2-1.1)	0.92% (0.092-2.4%)	0.94% (-0.015-2.7%)	1.5 (-0.024-4.4)	5.9 (0.55-16)
Benin	1990	Female	All Ages	58% (51-65%)	42% (35-49%)	0.72 (0.57-0.91)	0.59% (0.3-0.93%)	0.95% (0.2-1.9%)	290 (61-560)	13000 (6500-20000)
Benin	1990	Female	15 to 19	60% (46-73%)	40% (27-54%)	0.82 (0.41-1.4)	1.1% (0.51-1.9%)	1.2% (0.2-2.4%)	5 (0.88-10)	640 (280-1100)
Benin	1990	Female	20 to 24	67% (54-79%)	33% (21-46%)	1.2 (0.64-1.9)	2% (1-3.3%)	1.9% (0.47-3.9%)	10 (2.4-20)	1200 (610-2000)
Benin	1990	Female	25 to 29	69% (55-79%)	31% (21-45%)	1.4 (0.74-2.2)	2.4% (1.2-3.7%)	2.6% (0.83-4.6%)	13 (4-23)	1200 (620-1900)
Benin	1990	Female	30 to 34	71% (57-80%)	29% (20-43%)	1.4 (0.81-2.3)	2.4% (1.3-3.8%)	2.9% (1.2-5%)	14 (5.6-25)	1100 (610-1800)
Benin	1990	Female	35 to 39	74% (63-83%)	26% (17-37%)	1.5 (0.8-2.4)	2.7% (1.3-4.5%)	3.7% (1.5-6.4%)	19 (7.5-33)	1200 (580-1900)
Benin	1990	Female	40 to 44	75% (65-84%)	25% (16-35%)	1.5 (0.75-2.5)	3% (1.4-5.1%)	4.3% (1.8-7.6%)	24 (10-43)	1300 (580-2200)
Benin	1990	Female	45 to 49	74% (62-84%)	26% (16-38%)	1.5 (0.76-2.6)	3.2% (1.2-5.9%)	4.6% (1.6-8.6%)	31 (11-58)	1400 (510-2500)
Benin	1990	Female	50 to 54	73% (60-83%)	27% (17-40%)	1.5 (0.74-2.6)	3.4% (1.2-6.5%)	4.9% (1.7-9.3%)	36 (12-69)	1400 (480-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	1990	Female	55 to 59	71% (54-84%)	29% (16-46%)	1.6 (0.72-2.8)	3.5% (1-6.7%)	4.9% (1.5-9.4%)	43 (13-81)	1400 (390-2600)
Benin	1990	Female	60 to 64	71% (53-84%)	29% (16-47%)	1.6 (0.73-2.7)	2.8% (0.34-6.2%)	3.7% (0.5-8.2%)	40 (5.4-85)	1100 (130-2300)
Benin	1990	Female	65 to 69	72% (57-85%)	28% (15-43%)	1.4 (0.74-2.3)	1.9% (-0.37-5%)	2.4% (-0.51-6.5%)	32 (-6.6-85)	710 (-140-1900)
Benin	1990	Female	70 to 74	71% (55-83%)	29% (17-45%)	1.2 (0.55-2)	1.2% (-1.4-4%)	1.5% (-1.2-5.4%)	24 (-19-86)	430 (-360-1600)
Benin	1990	Female	75 to 79	68% (53-81%)	32% (19-47%)	1 (0.43-1.9)	0.34% (-1.7-3.3%)	0.42% (-2-4%)	5.9 (-29-56)	82 (-400-780)
Benin	1990	Female	80 to 84	66% (48-81%)	34% (19-52%)	0.93 (0.41-1.6)	-0.065% (-1.9-2%)	-0.086% (-2.3-2.4%)	-0.75 (-20-21)	-7 (-200-220)
Benin	1990	Female	85 to 89	66% (49-79%)	34% (21-51%)	0.91 (0.39-1.6)	-0.39% (-2-1.6%)	-0.51% (-2.5-1.8%)	-2.8 (-14-9.8)	-19 (-100-75)
Benin	1990	Female	90 to 94	68% (54-80%)	32% (20-46%)	0.92 (0.37-1.7)	-0.62% (-2.3-1.4%)	-0.85% (-2.8-1.6%)	-1.8 (-6.1-3.6)	-8.1 (-29-19)
Benin	1990	Female	95 plus	69% (50-84%)	31% (16-50%)	0.91 (0.3-1.9)	-0.58% (-2.2-1.7%)	-0.86% (-2.8-1.8%)	-0.5 (-1.6-1.1)	-1.4 (-5-3.9)
Benin	1990	Male	All Ages	9.5% (7.4-12%)	91% (88-93%)	0.21 (0.15-0.28)	0.66% (0.49-0.9%)	1.1% (0.78-1.6%)	370 (260-540)	16000 (12000-22000)
Benin	1990	Male	15 to 19	7.5% (4-13%)	92% (87-96%)	0.12 (0.049-0.23)	0.68% (0.44-0.99%)	0.48% (0.23-0.86%)	2.4 (1.2-4.4)	400 (250-590)
Benin	1990	Male	20 to 24	13% (6.6-21%)	87% (79-93%)	0.27 (0.12-0.51)	1.8% (1.1-2.6%)	1.4% (0.67-2.5%)	9.2 (4.4-16)	1100 (670-1600)
Benin	1990	Male	25 to 29	16% (8.6-27%)	84% (73-91%)	0.42 (0.18-0.79)	2.8% (1.7-4.3%)	2.7% (1.4-4.8%)	17 (8.7-31)	1600 (970-2400)
Benin	1990	Male	30 to 34	18% (9.9-31%)	82% (69-90%)	0.51 (0.23-0.94)	3.5% (2.3-5.4%)	3.9% (2.2-6.6%)	25 (14-41)	1800 (1100-2700)
Benin	1990	Male	35 to 39	20% (11-33%)	80% (67-89%)	0.56 (0.23-1)	4.1% (2.7-6.2%)	5% (3-7.9%)	32 (20-51)	1900 (1200-2900)
Benin	1990	Male	40 to 44	21% (11-35%)	79% (65-89%)	0.6 (0.24-1.2)	4.5% (2.8-7.3%)	5.7% (3.3-9.3%)	36 (21-58)	1800 (1100-2800)
Benin	1990	Male	45 to 49	21% (11-35%)	79% (65-89%)	0.62 (0.24-1.3)	4.9% (3.1-7.9%)	6.2% (3.8-10%)	47 (28-77)	2000 (1300-3300)
Benin	1990	Male	50 to 54	21% (12-35%)	79% (65-88%)	0.62 (0.25-1.2)	4.3% (2.5-7.2%)	5.4% (3-9.1%)	43 (23-74)	1600 (950-2700)
Benin	1990	Male	55 to 59	21% (12-35%)	79% (65-88%)	0.59 (0.23-1.1)	3.9% (2.3-6.4%)	4.8% (2.8-8.1%)	41 (23-67)	1300 (780-2200)
Benin	1990	Male	60 to 64	21% (11-35%)	79% (65-89%)	0.51 (0.19-1)	3.1% (1.8-5.5%)	3.8% (2.1-6.7%)	38 (21-67)	1100 (610-1800)
Benin	1990	Male	65 to 69	21% (11-34%)	79% (66-89%)	0.42 (0.17-0.84)	2.3% (1.3-3.9%)	2.7% (1.5-4.7%)	31 (17-55)	720 (420-1200)
Benin	1990	Male	70 to 74	19% (11-32%)	81% (68-89%)	0.32 (0.14-0.63)	1.6% (0.84-3%)	1.9% (0.97-3.6%)	24 (12-45)	440 (230-810)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	1990	Male	75 to 79	17% (9.8-29%)	83% (71-90%)	0.24 (0.086-0.49)	1.1% (0.56-2.1%)	1.3% (0.62-2.5%)	14 (7-28)	210 (100-390)
Benin	1990	Male	80 to 84	15% (8.1-26%)	85% (74-92%)	0.19 (0.071-0.38)	0.89% (0.46-1.7%)	1% (0.49-2%)	6.2 (3-12)	66 (34-130)
Benin	1990	Male	85 to 89	14% (7.7-22%)	86% (78-92%)	0.15 (0.055-0.33)	0.67% (0.34-1.2%)	0.75% (0.35-1.4%)	2.8 (1.3-5.2)	21 (11-38)
Benin	1990	Male	90 to 94	14% (7.9-23%)	86% (77-92%)	0.14 (0.047-0.3)	0.38% (0.13-0.77%)	0.39% (0.076-0.86%)	0.55 (0.11-1.2)	3.3 (1.1-6.8)
Benin	1990	Male	95 plus	16% (7.7-29%)	84% (71-92%)	0.14 (0.034-0.36)	0.35% (0.027-0.83%)	0.3% (-0.095-0.92%)	0.11 (-0.035-0.33)	0.52 (0.04-1.3)
Benin	1995	Female	All Ages	59% (52-66%)	41% (34-48%)	0.71 (0.56-0.9)	0.65% (0.33-1%)	0.99% (0.18-2%)	330 (56-660)	15000 (7800-24000)
Benin	1995	Female	15 to 19	61% (47-72%)	39% (28-53%)	0.81 (0.41-1.3)	1.2% (0.49-2%)	1.2% (0.17-2.5%)	6.3 (0.88-13)	780 (320-1400)
Benin	1995	Female	20 to 24	68% (54-79%)	32% (21-46%)	1.2 (0.62-1.8)	2.1% (1-3.5%)	2% (0.44-3.9%)	13 (2.7-25)	1500 (720-2400)
Benin	1995	Female	25 to 29	70% (56-79%)	30% (21-44%)	1.3 (0.72-2.1)	2.4% (1.2-3.8%)	2.5% (0.79-4.7%)	17 (5.2-31)	1600 (820-2500)
Benin	1995	Female	30 to 34	72% (58-81%)	28% (19-42%)	1.4 (0.81-2.2)	2.4% (1.3-3.9%)	2.8% (1.2-5%)	19 (7.6-33)	1500 (780-2300)
Benin	1995	Female	35 to 39	75% (65-83%)	25% (17-35%)	1.4 (0.77-2.3)	2.8% (1.2-4.6%)	3.6% (1.3-6.5%)	24 (8.4-43)	1500 (670-2500)
Benin	1995	Female	40 to 44	76% (66-84%)	24% (16-34%)	1.5 (0.74-2.5)	3.1% (1.4-5.2%)	4.3% (1.7-7.4%)	28 (11-50)	1500 (660-2500)
Benin	1995	Female	45 to 49	75% (62-84%)	25% (16-38%)	1.5 (0.74-2.6)	3.2% (1.1-6%)	4.5% (1.4-8.8%)	33 (10-64)	1500 (500-2800)
Benin	1995	Female	50 to 54	73% (61-84%)	27% (16-39%)	1.5 (0.73-2.6)	3.4% (1.1-6.4%)	4.8% (1.4-9%)	43 (12-83)	1600 (510-3100)
Benin	1995	Female	55 to 59	72% (55-84%)	28% (16-45%)	1.5 (0.7-2.7)	3.4% (0.92-6.7%)	4.7% (1.3-9.2%)	48 (13-94)	1500 (420-3000)
Benin	1995	Female	60 to 64	72% (54-85%)	28% (15-46%)	1.5 (0.71-2.6)	2.7% (0.35-5.9%)	3.5% (0.46-7.9%)	43 (5.7-97)	1200 (150-2600)
Benin	1995	Female	65 to 69	72% (56-85%)	28% (15-44%)	1.4 (0.69-2.3)	1.7% (-0.46-5.1%)	2.2% (-0.6-6.3%)	33 (-8.3-91)	730 (-200-2100)
Benin	1995	Female	70 to 74	71% (56-84%)	29% (16-44%)	1.1 (0.53-2)	1.1% (-1.1-4.5%)	1.4% (-1.4-5.4%)	24 (-23-90)	420 (-430-1600)
Benin	1995	Female	75 to 79	68% (53-82%)	32% (18-47%)	0.97 (0.41-1.8)	0.31% (-1.7-3.2%)	0.38% (-2.1-3.9%)	6 (-34-62)	84 (-460-870)
Benin	1995	Female	80 to 84	67% (49-80%)	33% (20-51%)	0.91 (0.41-1.6)	-0.088% (-2-2.1%)	-0.12% (-2.4-2.6%)	-1.2 (-23-25)	-11 (-240-260)
Benin	1995	Female	85 to 89	67% (50-79%)	33% (21-50%)	0.89 (0.36-1.6)	-0.4% (-2.2-1.6%)	-0.53% (-2.7-1.9%)	-3.4 (-18-12)	-23 (-120-94)
Benin	1995	Female	90 to 94	68% (54-80%)	32% (20-46%)	0.89 (0.36-1.7)	-0.61% (-2.3-1.5%)	-0.84% (-2.9-1.6%)	-2.3 (-7.8-4.5)	-10 (-37-24)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	1995	Female	95 plus	69% (51-84%)	31% (16-49%)	0.88 (0.3-1.9)	-0.56% (-2.1-1.7%)	-0.84% (-2.7-1.8%)	-0.62 (-2-1.3)	-1.7 (-6.2-5)
Benin	1995	Male	All Ages	10% (7.7-13%)	90% (87-92%)	0.22 (0.16-0.3)	0.76% (0.56-1.1%)	1.2% (0.86-1.8%)	450 (310-670)	20000 (15000-27000)
Benin	1995	Male	15 to 19	7.8% (4.3-14%)	92% (86-96%)	0.13 (0.051-0.25)	0.7% (0.45-1%)	0.5% (0.23-0.92%)	3 (1.4-5.5)	490 (310-740)
Benin	1995	Male	20 to 24	13% (7-21%)	87% (79-93%)	0.28 (0.12-0.51)	1.8% (1.1-2.6%)	1.4% (0.66-2.5%)	12 (5.3-21)	1300 (830-2000)
Benin	1995	Male	25 to 29	17% (8.9-28%)	83% (72-91%)	0.43 (0.19-0.81)	2.7% (1.7-4.2%)	2.6% (1.3-4.7%)	23 (11-41)	2000 (1200-3200)
Benin	1995	Male	30 to 34	19% (10-33%)	81% (67-90%)	0.53 (0.24-0.96)	3.4% (2.1-5.3%)	3.7% (1.9-6.3%)	33 (18-56)	2400 (1500-3700)
Benin	1995	Male	35 to 39	21% (11-34%)	79% (66-89%)	0.57 (0.25-1.1)	4% (2.5-6.1%)	4.7% (2.7-7.6%)	42 (25-70)	2500 (1600-3900)
Benin	1995	Male	40 to 44	21% (12-36%)	79% (64-88%)	0.61 (0.25-1.2)	4.3% (2.6-6.9%)	5.3% (3-8.9%)	48 (27-82)	2400 (1500-3900)
Benin	1995	Male	45 to 49	22% (11-36%)	78% (64-89%)	0.63 (0.25-1.3)	4.7% (2.9-7.7%)	5.8% (3.5-9.8%)	52 (31-89)	2300 (1400-3800)
Benin	1995	Male	50 to 54	22% (12-37%)	78% (63-88%)	0.63 (0.26-1.3)	4.3% (2.5-7.2%)	5.3% (3-9%)	55 (31-97)	2100 (1200-3700)
Benin	1995	Male	55 to 59	22% (13-37%)	78% (63-87%)	0.61 (0.23-1.2)	3.9% (2.3-6.6%)	4.8% (2.7-8.1%)	51 (28-87)	1700 (960-2800)
Benin	1995	Male	60 to 64	22% (12-37%)	78% (63-88%)	0.53 (0.19-1.1)	3.1% (1.7-5.6%)	3.7% (2-6.8%)	44 (24-80)	1200 (700-2200)
Benin	1995	Male	65 to 69	21% (12-35%)	79% (65-88%)	0.43 (0.17-0.84)	2.3% (1.3-4%)	2.7% (1.5-4.8%)	35 (20-62)	800 (460-1400)
Benin	1995	Male	70 to 74	20% (12-33%)	80% (67-88%)	0.33 (0.13-0.67)	1.6% (0.85-3%)	1.9% (0.98-3.5%)	27 (14-50)	490 (260-910)
Benin	1995	Male	75 to 79	18% (10-30%)	82% (70-90%)	0.24 (0.083-0.5)	1.1% (0.53-2.2%)	1.3% (0.6-2.5%)	16 (7.5-32)	230 (110-450)
Benin	1995	Male	80 to 84	16% (8.5-27%)	84% (73-92%)	0.19 (0.069-0.4)	0.89% (0.46-1.7%)	1% (0.49-2%)	7.1 (3.4-14)	76 (38-150)
Benin	1995	Male	85 to 89	15% (8-24%)	85% (76-92%)	0.16 (0.053-0.33)	0.68% (0.34-1.3%)	0.75% (0.33-1.5%)	3.2 (1.4-6.3)	25 (12-47)
Benin	1995	Male	90 to 94	15% (8.3-24%)	85% (76-92%)	0.14 (0.046-0.32)	0.41% (0.13-0.87%)	0.41% (0.084-0.97%)	0.72 (0.15-1.7)	4.3 (1.3-8.9)
Benin	1995	Male	95 plus	17% (8-30%)	83% (70-92%)	0.15 (0.035-0.37)	0.41% (0.043-1%)	0.37% (-0.051-1.1%)	0.17 (-0.023-0.5)	0.73 (0.078-1.8)
Benin	2000	Female	All Ages	64% (56-72%)	36% (28-44%)	0.78 (0.61-0.98)	0.74% (0.37-1.2%)	1.1% (0.19-2.1%)	390 (69-750)	19000 (9100-29000)
Benin	2000	Female	15 to 19	70% (57-82%)	30% (18-43%)	0.99 (0.53-1.6)	1.4% (0.52-2.3%)	1.6% (0.15-3.3%)	8.9 (0.74-19)	1000 (390-1800)
Benin	2000	Female	20 to 24	76% (66-83%)	24% (17-34%)	1.4 (0.75-2.1)	2.3% (1-3.7%)	2.2% (0.45-4.4%)	17 (3.2-33)	1900 (840-3000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	2000	Female	25 to 29	75% (65-83%)	25% (17-35%)	1.5 (0.84-2.3)	2.5% (1.2-4.1%)	2.6% (0.82-5%)	23 (6.8-43)	2100 (1000-3400)
Benin	2000	Female	30 to 34	76% (66-83%)	24% (17-34%)	1.5 (0.9-2.4)	2.4% (1.3-3.9%)	2.7% (1.1-4.8%)	24 (9.1-44)	1800 (970-3000)
Benin	2000	Female	35 to 39	78% (69-85%)	22% (15-31%)	1.5 (0.87-2.4)	2.7% (1.3-4.6%)	3.4% (1.3-6%)	31 (11-54)	1900 (870-3200)
Benin	2000	Female	40 to 44	78% (70-85%)	22% (15-30%)	1.6 (0.8-2.6)	3% (1.3-5.1%)	4% (1.6-7%)	34 (14-59)	1800 (780-3000)
Benin	2000	Female	45 to 49	78% (68-85%)	22% (15-32%)	1.6 (0.82-2.8)	3.3% (1-6.1%)	4.6% (1.3-8.6%)	39 (12-74)	1700 (570-3200)
Benin	2000	Female	50 to 54	75% (63-84%)	25% (16-37%)	1.5 (0.76-2.6)	3.5% (1-6.6%)	4.8% (1.3-9.2%)	48 (13-94)	1800 (520-3500)
Benin	2000	Female	55 to 59	72% (56-84%)	28% (16-44%)	1.6 (0.7-2.6)	3.5% (0.94-6.9%)	4.8% (1.3-9.3%)	59 (16-120)	1900 (510-3800)
Benin	2000	Female	60 to 64	72% (55-85%)	28% (15-45%)	1.5 (0.7-2.7)	2.7% (0.22-6.3%)	3.5% (0.28-8.2%)	50 (4-120)	1300 (100-3200)
Benin	2000	Female	65 to 69	73% (58-85%)	27% (15-42%)	1.4 (0.7-2.3)	1.7% (-0.64-5%)	2.2% (-0.81-6.5%)	36 (-13-110)	810 (-290-2400)
Benin	2000	Female	70 to 74	71% (56-83%)	29% (17-44%)	1.1 (0.53-2)	1% (-1.3-4.4%)	1.3% (-1.6-5.4%)	24 (-28-97)	420 (-510-1700)
Benin	2000	Female	75 to 79	68% (52-81%)	32% (19-48%)	0.98 (0.42-1.8)	0.27% (-1.8-3.1%)	0.33% (-2.1-3.8%)	5.5 (-36-62)	76 (-520-880)
Benin	2000	Female	80 to 84	67% (48-81%)	33% (19-52%)	0.91 (0.4-1.6)	-0.091% (-1.9-2.3%)	-0.12% (-2.3-2.8%)	-1.3 (-25-29)	-12 (-260-310)
Benin	2000	Female	85 to 89	67% (49-79%)	33% (21-51%)	0.89 (0.37-1.6)	-0.38% (-2.1-1.8%)	-0.5% (-2.7-2.1%)	-3.7 (-20-16)	-24 (-140-120)
Benin	2000	Female	90 to 94	68% (54-80%)	32% (20-46%)	0.89 (0.35-1.7)	-0.55% (-2.3-1.7%)	-0.77% (-2.9-2%)	-2.4 (-9.2-6.3)	-11 (-44-33)
Benin	2000	Female	95 plus	69% (50-84%)	31% (16-50%)	0.88 (0.29-1.9)	-0.45% (-2.1-2.1%)	-0.73% (-2.7-2.4%)	-0.67 (-2.5-2.2)	-1.7 (-7.7-7.8)
Benin	2000	Male	All Ages	10% (8.1-14%)	90% (86-92%)	0.24 (0.17-0.32)	0.87% (0.64-1.2%)	1.4% (0.95-2.1%)	540 (370-820)	24000 (18000-33000)
Benin	2000	Male	15 to 19	8.2% (4.6-14%)	92% (86-95%)	0.13 (0.054-0.26)	0.74% (0.45-1.1%)	0.54% (0.23-1%)	3.7 (1.6-7)	580 (350-890)
Benin	2000	Male	20 to 24	14% (7.3-22%)	86% (78-93%)	0.3 (0.13-0.54)	1.8% (1.2-2.8%)	1.5% (0.71-2.8%)	14 (6.7-25)	1600 (980-2400)
Benin	2000	Male	25 to 29	18% (9.7-30%)	82% (70-90%)	0.46 (0.2-0.85)	2.6% (1.6-4.2%)	2.5% (1.2-4.7%)	27 (13-49)	2300 (1400-3800)
Benin	2000	Male	30 to 34	20% (11-33%)	80% (67-89%)	0.56 (0.25-1)	3.3% (2-5.1%)	3.5% (1.9-5.9%)	41 (21-68)	2900 (1800-4500)
Benin	2000	Male	35 to 39	21% (12-35%)	79% (65-88%)	0.6 (0.27-1.1)	3.8% (2.4-5.9%)	4.3% (2.5-6.8%)	52 (30-87)	3100 (1900-4800)
Benin	2000	Male	40 to 44	22% (12-37%)	78% (63-88%)	0.63 (0.26-1.2)	4.1% (2.6-6.6%)	4.9% (2.9-8%)	60 (34-99)	3000 (1800-4800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	2000	Male	45 to 49	23% (12-37%)	77% (63-88%)	0.67 (0.27-1.3)	4.6% (2.8-8%)	5.6% (3.2-9.8%)	68 (39-120)	3000 (1800-5100)
Benin	2000	Male	50 to 54	23% (12-38%)	77% (62-88%)	0.67 (0.27-1.3)	4.3% (2.4-7.5%)	5.2% (2.8-9.2%)	62 (33-110)	2300 (1300-4000)
Benin	2000	Male	55 to 59	23% (13-38%)	77% (62-87%)	0.64 (0.24-1.2)	4% (2.2-6.9%)	4.8% (2.5-8.5%)	66 (34-110)	2200 (1200-3700)
Benin	2000	Male	60 to 64	23% (12-38%)	77% (62-88%)	0.56 (0.21-1.1)	3.1% (1.7-5.8%)	3.7% (1.9-7.1%)	54 (28-100)	1500 (810-2700)
Benin	2000	Male	65 to 69	22% (13-36%)	78% (64-87%)	0.46 (0.19-0.87)	2.3% (1.2-4%)	2.7% (1.4-4.9%)	39 (21-71)	900 (490-1600)
Benin	2000	Male	70 to 74	21% (12-34%)	79% (66-88%)	0.35 (0.14-0.73)	1.6% (0.8-3.2%)	1.9% (0.91-3.8%)	29 (14-57)	520 (260-1000)
Benin	2000	Male	75 to 79	19% (11-31%)	81% (69-89%)	0.26 (0.091-0.54)	1.1% (0.51-2.2%)	1.3% (0.55-2.6%)	17 (7.4-35)	240 (110-500)
Benin	2000	Male	80 to 84	17% (9.2-28%)	83% (72-91%)	0.21 (0.076-0.42)	0.9% (0.42-1.7%)	1% (0.46-2%)	8 (3.6-16)	85 (40-170)
Benin	2000	Male	85 to 89	15% (8.6-25%)	85% (75-91%)	0.17 (0.058-0.35)	0.69% (0.29-1.4%)	0.77% (0.29-1.6%)	3.6 (1.4-7.6)	28 (12-56)
Benin	2000	Male	90 to 94	16% (8.9-26%)	84% (74-91%)	0.16 (0.051-0.35)	0.43% (0.12-0.94%)	0.44% (0.057-1.1%)	0.84 (0.11-2)	5 (1.3-11)
Benin	2000	Male	95 plus	18% (8.8-32%)	82% (68-91%)	0.16 (0.039-0.4)	0.46% (0.056-1.2%)	0.44% (-0.062-1.3%)	0.22 (-0.032-0.66)	0.95 (0.12-2.5)
Benin	2005	Female	All Ages	59% (52-66%)	41% (34-48%)	0.79 (0.62-0.99)	0.83% (0.44-1.3%)	1.2% (0.3-2.3%)	460 (110-860)	21000 (11000-33000)
Benin	2005	Female	15 to 19	62% (51-74%)	38% (26-49%)	0.93 (0.47-1.5)	1.3% (0.58-2.2%)	1.5% (0.25-3%)	9.2 (1.5-19)	1100 (470-1900)
Benin	2005	Female	20 to 24	69% (58-78%)	31% (22-42%)	1.3 (0.69-2.1)	2.3% (1.2-3.7%)	2.2% (0.56-4.4%)	17 (4.3-36)	2000 (1000-3300)
Benin	2005	Female	25 to 29	69% (57-78%)	31% (22-43%)	1.5 (0.81-2.3)	2.4% (1.3-3.8%)	2.4% (0.88-4.4%)	24 (8.6-44)	2200 (1200-3600)
Benin	2005	Female	30 to 34	71% (59-80%)	29% (20-41%)	1.5 (0.89-2.4)	2.2% (1.2-3.4%)	2.3% (1-4%)	25 (10-44)	2000 (1100-3200)
Benin	2005	Female	35 to 39	74% (64-82%)	26% (18-36%)	1.6 (0.87-2.5)	2.5% (1.3-4.1%)	3.1% (1.3-5.3%)	35 (15-60)	2200 (1100-3500)
Benin	2005	Female	40 to 44	75% (66-83%)	25% (17-34%)	1.6 (0.82-2.6)	2.8% (1.4-4.6%)	3.6% (1.6-6%)	40 (18-67)	2100 (1000-3400)
Benin	2005	Female	45 to 49	75% (64-83%)	25% (17-36%)	1.6 (0.85-2.9)	3.3% (1.3-6.2%)	4.5% (1.6-8.6%)	47 (17-89)	2100 (840-3900)
Benin	2005	Female	50 to 54	74% (62-83%)	26% (17-38%)	1.7 (0.82-2.9)	3.6% (1.2-7%)	4.9% (1.6-9.4%)	53 (17-100)	2000 (690-3900)
Benin	2005	Female	55 to 59	72% (56-84%)	28% (16-44%)	1.7 (0.8-2.9)	3.7% (1-7.5%)	5.1% (1.4-9.8%)	66 (18-130)	2100 (580-4200)
Benin	2005	Female	60 to 64	72% (55-85%)	28% (15-45%)	1.7 (0.78-2.9)	2.9% (0.21-6.6%)	3.8% (0.35-8.5%)	62 (5.8-140)	1700 (130-3900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	2005	Female	65 to 69	72% (57-85%)	28% (15-43%)	1.5 (0.77-2.5)	1.9% (-0.57-5.4%)	2.5% (-0.74-6.8%)	46 (-13-120)	1000 (-290-2900)
Benin	2005	Female	70 to 74	71% (55-83%)	29% (17-45%)	1.3 (0.59-2.2)	1.2% (-1.2-4.9%)	1.5% (-1.5-6.1%)	30 (-29-110)	530 (-530-2100)
Benin	2005	Female	75 to 79	69% (52-82%)	31% (18-48%)	1.1 (0.46-2)	0.41% (-1.8-3.7%)	0.5% (-2.2-4.4%)	8.5 (-38-76)	120 (-520-1100)
Benin	2005	Female	80 to 84	67% (49-81%)	33% (19-51%)	1 (0.45-1.8)	0.084% (-1.9-2.7%)	0.08% (-2.2-3.2%)	0.84 (-26-35)	11 (-260-370)
Benin	2005	Female	85 to 89	67% (50-79%)	33% (21-50%)	0.98 (0.4-1.8)	-0.19% (-2-2.2%)	-0.29% (-2.6-2.5%)	-2.3 (-20-20)	-13 (-140-150)
Benin	2005	Female	90 to 94	69% (54-81%)	31% (19-46%)	0.99 (0.39-2)	-0.39% (-2.3-2.1%)	-0.59% (-2.9-2.4%)	-2.1 (-10-8.7)	-8.5 (-49-46)
Benin	2005	Female	95 plus	70% (52-84%)	30% (16-48%)	0.98 (0.33-2.1)	-0.3% (-2.1-2.7%)	-0.55% (-2.7-3.1%)	-0.59 (-2.9-3.4)	-1.3 (-8.9-12)
Benin	2005	Male	All Ages	11% (8.2-14%)	89% (86-92%)	0.27 (0.19-0.35)	1% (0.75-1.4%)	1.6% (1.1-2.4%)	670 (450-1000)	30000 (22000-41000)
Benin	2005	Male	15 to 19	8.3% (4.6-15%)	92% (85-95%)	0.15 (0.062-0.3)	0.79% (0.5-1.2%)	0.58% (0.26-1.1%)	4.5 (2.1-8.8)	710 (440-1100)
Benin	2005	Male	20 to 24	14% (7.3-23%)	86% (77-93%)	0.33 (0.14-0.62)	2% (1.3-2.9%)	1.6% (0.78-2.9%)	17 (8.1-31)	1900 (1200-3000)
Benin	2005	Male	25 to 29	18% (9.4-29%)	82% (71-91%)	0.5 (0.22-0.94)	2.8% (1.8-4.3%)	2.7% (1.4-4.7%)	32 (16-57)	2800 (1800-4300)
Benin	2005	Male	30 to 34	20% (11-33%)	80% (67-89%)	0.61 (0.28-1.1)	3.3% (2.1-5.2%)	3.5% (1.9-5.9%)	48 (27-82)	3400 (2100-5300)
Benin	2005	Male	35 to 39	21% (12-34%)	79% (66-88%)	0.65 (0.29-1.2)	3.8% (2.4-5.8%)	4.3% (2.5-6.9%)	64 (36-100)	3700 (2300-5800)
Benin	2005	Male	40 to 44	22% (13-37%)	78% (63-87%)	0.7 (0.29-1.3)	4.1% (2.6-6.6%)	4.9% (2.9-7.8%)	75 (43-120)	3800 (2300-6000)
Benin	2005	Male	45 to 49	23% (12-38%)	77% (62-88%)	0.73 (0.3-1.5)	4.7% (2.9-7.9%)	5.7% (3.3-9.7%)	86 (49-150)	3800 (2300-6200)
Benin	2005	Male	50 to 54	23% (13-38%)	77% (62-87%)	0.74 (0.31-1.5)	4.5% (2.5-7.7%)	5.5% (2.8-9.5%)	82 (42-140)	3100 (1700-5300)
Benin	2005	Male	55 to 59	23% (13-38%)	77% (62-87%)	0.71 (0.27-1.4)	4.3% (2.3-7.2%)	5.1% (2.7-8.9%)	75 (39-130)	2500 (1300-4200)
Benin	2005	Male	60 to 64	23% (12-37%)	77% (63-88%)	0.62 (0.23-1.2)	3.4% (1.8-6.1%)	4% (2-7.4%)	70 (35-130)	2000 (1000-3500)
Benin	2005	Male	65 to 69	22% (13-36%)	78% (64-87%)	0.51 (0.22-0.94)	2.4% (1.3-4.3%)	2.8% (1.5-5%)	48 (25-88)	1100 (600-2000)
Benin	2005	Male	70 to 74	21% (12-34%)	79% (66-88%)	0.38 (0.16-0.79)	1.7% (0.82-3.3%)	2% (0.94-3.9%)	33 (16-64)	610 (300-1200)
Benin	2005	Male	75 to 79	19% (11-31%)	81% (69-89%)	0.29 (0.1-0.56)	1.2% (0.54-2.3%)	1.3% (0.59-2.7%)	19 (8.4-39)	270 (120-550)
Benin	2005	Male	80 to 84	17% (9.2-28%)	83% (72-91%)	0.23 (0.085-0.46)	0.95% (0.47-1.8%)	1.1% (0.5-2.1%)	8.9 (4.1-17)	95 (46-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	2005	Male	85 to 89	15% (8.7-25%)	85% (75-91%)	0.19 (0.067-0.4)	0.73% (0.33-1.4%)	0.81% (0.33-1.6%)	4 (1.6-8.2)	32 (14-61)
Benin	2005	Male	90 to 94	16% (9.1-25%)	84% (75-91%)	0.17 (0.058-0.38)	0.47% (0.13-1.1%)	0.48% (0.064-1.2%)	0.98 (0.13-2.5)	5.8 (1.6-13)
Benin	2005	Male	95 plus	18% (8.8-32%)	82% (68-91%)	0.18 (0.043-0.47)	0.5% (0.082-1.3%)	0.47% (-0.052-1.4%)	0.26 (-0.029-0.77)	1.1 (0.18-2.8)
Benin	2010	Female	All Ages	56% (48-63%)	44% (37-52%)	0.78 (0.59-1)	0.97% (0.52-1.5%)	1.5% (0.32-2.9%)	560 (120-1100)	25000 (13000-40000)
Benin	2010	Female	15 to 19	57% (44-70%)	43% (30-56%)	0.86 (0.41-1.5)	1.3% (0.56-2.2%)	1.4% (0.3-2.8%)	10 (2.1-21)	1300 (550-2200)
Benin	2010	Female	20 to 24	64% (51-75%)	36% (25-49%)	1.2 (0.64-2.1)	2.3% (1.2-3.8%)	2.2% (0.65-4.3%)	19 (5.4-38)	2300 (1200-3800)
Benin	2010	Female	25 to 29	65% (52-76%)	35% (24-48%)	1.4 (0.75-2.3)	2.5% (1.5-3.9%)	2.6% (0.98-4.7%)	25 (9.3-45)	2400 (1400-3800)
Benin	2010	Female	30 to 34	67% (53-78%)	33% (22-47%)	1.5 (0.84-2.4)	2.4% (1.4-3.7%)	2.6% (1.2-4.5%)	28 (13-47)	2300 (1300-3600)
Benin	2010	Female	35 to 39	71% (59-80%)	29% (20-41%)	1.6 (0.81-2.5)	2.7% (1.4-4.3%)	3.4% (1.6-5.7%)	38 (16-63)	2400 (1300-3800)
Benin	2010	Female	40 to 44	72% (62-82%)	28% (18-38%)	1.6 (0.8-2.7)	3.2% (1.6-5.1%)	4.3% (1.9-7.3%)	47 (21-80)	2500 (1200-4100)
Benin	2010	Female	45 to 49	73% (60-83%)	27% (17-40%)	1.7 (0.83-2.9)	3.6% (1.5-6.9%)	5.2% (1.9-9.8%)	57 (21-110)	2600 (1000-4800)
Benin	2010	Female	50 to 54	73% (61-83%)	27% (17-39%)	1.7 (0.83-3.1)	4% (1.4-7.9%)	5.7% (2-11%)	68 (24-130)	2600 (940-5000)
Benin	2010	Female	55 to 59	72% (56-84%)	28% (16-44%)	1.8 (0.84-3.2)	4.2% (1.2-8.2%)	5.7% (1.6-11%)	79 (22-160)	2500 (730-5200)
Benin	2010	Female	60 to 64	72% (55-85%)	28% (15-45%)	1.8 (0.82-3.2)	3.2% (0.28-7.3%)	4.2% (0.44-9.4%)	74 (7.3-170)	2000 (180-4600)
Benin	2010	Female	65 to 69	73% (57-85%)	27% (15-43%)	1.6 (0.8-2.7)	2.2% (-0.57-6%)	2.8% (-0.68-7.5%)	62 (-15-170)	1400 (-340-3900)
Benin	2010	Female	70 to 74	71% (56-83%)	29% (17-44%)	1.4 (0.61-2.5)	1.3% (-1.2-5.2%)	1.7% (-1.5-6.5%)	39 (-35-150)	700 (-630-2700)
Benin	2010	Female	75 to 79	69% (53-81%)	31% (19-47%)	1.2 (0.47-2.2)	0.54% (-1.8-4.1%)	0.65% (-2.1-5%)	13 (-44-96)	180 (-630-1400)
Benin	2010	Female	80 to 84	67% (50-81%)	33% (19-50%)	1.1 (0.46-1.9)	0.2% (-1.8-3%)	0.22% (-2.2-3.6%)	2.7 (-28-44)	32 (-290-470)
Benin	2010	Female	85 to 89	67% (50-80%)	33% (20-50%)	1 (0.42-1.9)	-0.057% (-2.1-2.5%)	-0.13% (-2.6-2.9%)	-1.1 (-22-24)	-4.2 (-160-180)
Benin	2010	Female	90 to 94	68% (54-80%)	32% (20-46%)	1 (0.4-2.1)	-0.24% (-2.1-2.7%)	-0.42% (-2.7-3.1%)	-1.6 (-11-12)	-5.8 (-50-63)
Benin	2010	Female	95 plus	69% (52-84%)	31% (16-48%)	1 (0.34-2.2)	-0.17% (-2-3.1%)	-0.4% (-2.7-3.6%)	-0.49 (-3.2-4.4)	-0.81 (-10-15)
Benin	2010	Male	All Ages	11% (8.4-14%)	89% (86-92%)	0.29 (0.2-0.39)	1.2% (0.88-1.7%)	1.9% (1.3-2.8%)	810 (540-1200)	36000 (26000-50000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	2010	Male	15 to 19	8.4% (4.7-15%)	92% (85-95%)	0.16 (0.067-0.31)	0.81% (0.52-1.2%)	0.6% (0.27-1.1%)	5.5 (2.4-10)	850 (520-1300)
Benin	2010	Male	20 to 24	14% (7.5-23%)	86% (77-93%)	0.35 (0.15-0.66)	2.1% (1.3-3.2%)	1.7% (0.83-3%)	20 (9.5-36)	2300 (1400-3600)
Benin	2010	Male	25 to 29	18% (9.7-29%)	82% (71-90%)	0.54 (0.24-1)	3.1% (1.9-4.8%)	3.1% (1.5-5.4%)	38 (19-69)	3400 (2100-5400)
Benin	2010	Male	30 to 34	20% (11-34%)	80% (66-89%)	0.65 (0.3-1.2)	3.7% (2.4-5.7%)	4.1% (2.3-6.7%)	56 (30-94)	4000 (2500-6200)
Benin	2010	Male	35 to 39	22% (12-36%)	78% (64-88%)	0.69 (0.3-1.3)	4.2% (2.6-6.3%)	4.9% (2.8-7.7%)	73 (41-120)	4300 (2600-6500)
Benin	2010	Male	40 to 44	22% (13-37%)	78% (63-87%)	0.74 (0.31-1.4)	4.7% (2.8-7.4%)	5.7% (3.2-9.2%)	89 (50-140)	4500 (2700-7100)
Benin	2010	Male	45 to 49	23% (12-37%)	77% (63-88%)	0.78 (0.3-1.6)	5.3% (3.1-8.8%)	6.6% (3.8-11%)	110 (61-180)	4700 (2800-7900)
Benin	2010	Male	50 to 54	23% (12-38%)	77% (62-88%)	0.78 (0.31-1.6)	5% (2.7-8.3%)	6.1% (3.2-10%)	100 (53-180)	3900 (2100-6600)
Benin	2010	Male	55 to 59	23% (13-38%)	77% (62-87%)	0.75 (0.28-1.5)	4.6% (2.5-7.9%)	5.6% (2.9-9.8%)	100 (52-170)	3300 (1800-5500)
Benin	2010	Male	60 to 64	23% (12-38%)	77% (62-88%)	0.65 (0.25-1.4)	3.5% (1.9-6.3%)	4.2% (2.1-7.6%)	77 (39-140)	2200 (1100-3800)
Benin	2010	Male	65 to 69	22% (13-37%)	78% (63-87%)	0.54 (0.22-1)	2.5% (1.3-4.5%)	2.9% (1.5-5.3%)	62 (32-110)	1400 (770-2500)
Benin	2010	Male	70 to 74	21% (12-34%)	79% (66-88%)	0.41 (0.16-0.88)	1.8% (0.88-3.6%)	2.1% (1-4.2%)	41 (20-85)	750 (370-1500)
Benin	2010	Male	75 to 79	19% (11-31%)	81% (69-89%)	0.31 (0.1-0.64)	1.2% (0.56-2.4%)	1.4% (0.6-2.9%)	22 (9.5-45)	310 (140-630)
Benin	2010	Male	80 to 84	17% (9.1-28%)	83% (72-91%)	0.25 (0.087-0.51)	0.99% (0.47-1.8%)	1.1% (0.51-2.2%)	10 (4.5-19)	110 (50-200)
Benin	2010	Male	85 to 89	16% (8.3-25%)	84% (75-92%)	0.2 (0.07-0.44)	0.75% (0.35-1.4%)	0.83% (0.36-1.6%)	4.3 (1.9-8.6)	34 (16-65)
Benin	2010	Male	90 to 94	16% (9-26%)	84% (74-91%)	0.19 (0.058-0.42)	0.49% (0.13-1.1%)	0.5% (0.071-1.2%)	1.1 (0.15-2.6)	6.4 (1.8-14)
Benin	2010	Male	95 plus	18% (8.9-33%)	82% (67-91%)	0.19 (0.045-0.51)	0.51% (0.091-1.3%)	0.48% (-0.04-1.4%)	0.28 (-0.023-0.83)	1.2 (0.21-3)
Benin	2016	Female	All Ages	60% (53-67%)	40% (33-47%)	0.86 (0.5-1.3)	1.1% (0.45-2.1%)	1.7% (0.13-3.9%)	640 (51-1500)	29000 (12000-52000)
Benin	2016	Female	15 to 19	61% (48-74%)	39% (26-52%)	0.95 (0.42-1.8)	1.3% (0.58-2.4%)	1.4% (0.26-3.1%)	11 (2-24)	1400 (600-2600)
Benin	2016	Female	20 to 24	69% (55-79%)	31% (21-45%)	1.4 (0.64-2.6)	2.4% (1.3-4.2%)	2.3% (0.58-5.1%)	20 (4.9-44)	2600 (1300-4500)
Benin	2016	Female	25 to 29	70% (57-79%)	30% (21-43%)	1.6 (0.72-2.9)	2.7% (1.5-4.4%)	2.8% (0.99-5.5%)	26 (9.2-53)	2700 (1500-4400)
Benin	2016	Female	30 to 34	72% (60-81%)	28% (19-40%)	1.7 (0.79-2.9)	2.7% (1.4-4.4%)	3.1% (1.2-5.8%)	30 (12-57)	2600 (1400-4200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	2016	Female	35 to 39	75% (65-82%)	25% (18-35%)	1.7 (0.78-3.1)	3% (1.5-5.1%)	3.9% (1.5-7.1%)	41 (16-75)	2700 (1300-4600)
Benin	2016	Female	40 to 44	76% (66-84%)	24% (16-34%)	1.7 (0.73-3.3)	3.3% (1.5-6%)	4.8% (1.9-8.7%)	54 (21-98)	2900 (1300-5100)
Benin	2016	Female	45 to 49	75% (63-84%)	25% (16-37%)	1.7 (0.75-3.4)	3.8% (1.3-7.4%)	5.5% (1.8-11%)	68 (22-140)	3000 (1100-6000)
Benin	2016	Female	50 to 54	74% (62-84%)	26% (16-38%)	1.8 (0.74-3.4)	4.1% (1.2-8.4%)	5.8% (1.7-12%)	83 (24-180)	3200 (920-6700)
Benin	2016	Female	55 to 59	72% (57-85%)	28% (15-43%)	1.8 (0.76-3.7)	4.2% (0.82-9.1%)	5.8% (1.2-12%)	96 (20-210)	3100 (620-6700)
Benin	2016	Female	60 to 64	72% (56-85%)	28% (15-44%)	1.8 (0.73-3.6)	3.2% (-0.023-8.3%)	4.2% (0.011-11%)	81 (0.19-200)	2200 (-16-5700)
Benin	2016	Female	65 to 69	72% (55-85%)	28% (15-45%)	1.6 (0.69-3.1)	2.2% (-0.82-7%)	2.7% (-1-8.7%)	65 (-23-210)	1500 (-550-4900)
Benin	2016	Female	70 to 74	71% (55-83%)	29% (17-45%)	1.3 (0.52-2.6)	1.3% (-1.4-6.1%)	1.6% (-1.7-7.5%)	45 (-48-210)	820 (-890-3800)
Benin	2016	Female	75 to 79	68% (53-82%)	32% (18-47%)	1.2 (0.42-2.4)	0.54% (-1.8-5%)	0.63% (-2.3-5.9%)	16 (-54-140)	230 (-740-2000)
Benin	2016	Female	80 to 84	67% (48-81%)	33% (19-52%)	1.1 (0.39-2.1)	0.21% (-2-3.9%)	0.22% (-2.5-4.6%)	3.2 (-37-68)	39 (-370-730)
Benin	2016	Female	85 to 89	67% (49-80%)	33% (20-51%)	1.1 (0.39-2.2)	-0.019% (-2.2-3.4%)	-0.091% (-2.7-3.9%)	-0.87 (-26-38)	-1.6 (-180-290)
Benin	2016	Female	90 to 94	69% (55-81%)	31% (19-45%)	1.1 (0.35-2.3)	-0.19% (-2.2-3.4%)	-0.36% (-2.9-3.8%)	-1.5 (-12-16)	-4.9 (-57-86)
Benin	2016	Female	95 plus	70% (52-85%)	30% (15-48%)	1.1 (0.3-2.4)	-0.11% (-2.2-3.9%)	-0.34% (-2.9-4.4%)	-0.46 (-3.8-5.9)	-0.59 (-12-20)
Benin	2016	Male	All Ages	11% (8.4-14%)	89% (86-92%)	0.29 (0.16-0.49)	1.4% (0.91-2%)	2.1% (1.3-3.2%)	900 (550-1400)	41000 (27000-61000)
Benin	2016	Male	15 to 19	8.4% (4.7-15%)	92% (85-95%)	0.16 (0.06-0.34)	0.84% (0.51-1.3%)	0.6% (0.24-1.2%)	6.1 (2.5-12)	990 (590-1600)
Benin	2016	Male	20 to 24	14% (7.7-23%)	86% (77-92%)	0.36 (0.13-0.74)	2.1% (1.3-3.4%)	1.7% (0.76-3.3%)	22 (9.7-44)	2600 (1600-4300)
Benin	2016	Male	25 to 29	18% (9.7-29%)	82% (71-90%)	0.54 (0.21-1.1)	3.1% (1.9-5.2%)	3.1% (1.5-5.9%)	42 (19-82)	3800 (2200-6300)
Benin	2016	Male	30 to 34	20% (11-33%)	80% (67-89%)	0.65 (0.26-1.3)	3.8% (2.4-6.1%)	4.2% (2.2-7.4%)	60 (30-110)	4400 (2600-7100)
Benin	2016	Male	35 to 39	21% (12-35%)	79% (65-88%)	0.69 (0.26-1.5)	4.3% (2.6-6.7%)	5.1% (2.8-8.4%)	78 (42-130)	4700 (2800-7500)
Benin	2016	Male	40 to 44	22% (13-37%)	78% (63-87%)	0.74 (0.26-1.6)	4.7% (2.7-7.9%)	5.8% (3.1-10%)	95 (50-160)	4900 (2700-8100)
Benin	2016	Male	45 to 49	23% (12-36%)	77% (64-88%)	0.78 (0.28-1.7)	5.3% (2.9-9.2%)	6.6% (3.5-12%)	120 (62-220)	5300 (2900-9400)
Benin	2016	Male	50 to 54	23% (12-37%)	77% (63-88%)	0.78 (0.28-1.8)	4.8% (2.5-8.5%)	5.8% (2.9-10%)	120 (59-210)	4500 (2400-8100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Benin	2016	Male	55 to 59	23% (13-37%)	77% (63-87%)	0.75 (0.25-1.7)	4.4% (2.3-8%)	5.4% (2.7-9.7%)	110 (57-210)	3800 (2000-6700)
Benin	2016	Male	60 to 64	23% (12-38%)	77% (62-88%)	0.67 (0.22-1.5)	3.4% (1.7-6.4%)	4% (1.9-7.7%)	94 (45-180)	2600 (1300-4800)
Benin	2016	Male	65 to 69	22% (13-36%)	78% (64-87%)	0.55 (0.19-1.2)	2.4% (1.1-4.7%)	2.8% (1.3-5.5%)	62 (28-120)	1500 (690-2900)
Benin	2016	Male	70 to 74	21% (12-34%)	79% (66-88%)	0.41 (0.14-0.97)	1.7% (0.76-3.5%)	2% (0.87-4.1%)	46 (20-95)	850 (390-1800)
Benin	2016	Male	75 to 79	19% (11-31%)	81% (69-89%)	0.31 (0.088-0.68)	1.1% (0.51-2.4%)	1.3% (0.56-2.7%)	24 (10-51)	350 (160-740)
Benin	2016	Male	80 to 84	17% (9.3-29%)	83% (71-91%)	0.25 (0.073-0.54)	0.92% (0.41-1.8%)	1% (0.43-2.1%)	10 (4.3-21)	110 (50-220)
Benin	2016	Male	85 to 89	16% (8.9-25%)	84% (75-91%)	0.2 (0.062-0.49)	0.69% (0.3-1.4%)	0.76% (0.29-1.6%)	4.3 (1.6-9)	34 (15-69)
Benin	2016	Male	90 to 94	16% (8.9-26%)	84% (74-91%)	0.18 (0.051-0.46)	0.46% (0.12-1%)	0.46% (0.048-1.2%)	1 (0.11-2.6)	6.3 (1.6-14)
Benin	2016	Male	95 plus	18% (9-32%)	82% (68-91%)	0.19 (0.039-0.5)	0.48% (0.096-1.3%)	0.43% (-0.05-1.4%)	0.27 (-0.031-0.86)	1.2 (0.23-3.2)
Burkina Faso	1990	Female	All Ages	33% (29-38%)	67% (62-71%)	1.2 (0.9-1.6)	1.3% (0.84-1.8%)	2.7% (1.6-4%)	1900 (1200-2800)	63000 (43000-88000)
Burkina Faso	1990	Female	15 to 19	32% (23-42%)	68% (58-77%)	1.3 (0.7-2)	1.8% (0.98-2.9%)	2.3% (1-4.1%)	25 (11-43)	2200 (1200-3500)
Burkina Faso	1990	Female	20 to 24	38% (27-49%)	62% (51-73%)	1.8 (1.1-2.8)	3.1% (1.9-4.6%)	3.5% (1.8-5.6%)	48 (25-79)	4200 (2500-6300)
Burkina Faso	1990	Female	25 to 29	41% (30-52%)	59% (48-70%)	2.1 (1.3-3.2)	4% (2.4-6.1%)	4.8% (2.5-7.8%)	68 (36-110)	4900 (3000-7500)
Burkina Faso	1990	Female	30 to 34	44% (33-55%)	56% (45-67%)	2.3 (1.4-3.5)	4.3% (2.7-6.4%)	5.5% (3.2-8.4%)	73 (43-110)	4700 (2900-7000)
Burkina Faso	1990	Female	35 to 39	48% (36-57%)	52% (43-64%)	2.5 (1.5-3.7)	4.7% (2.9-6.8%)	6.1% (3.6-9.1%)	81 (49-120)	4500 (2900-6600)
Burkina Faso	1990	Female	40 to 44	50% (39-60%)	50% (40-61%)	2.7 (1.6-4.1)	5.5% (3.4-8%)	7.5% (4.5-11%)	97 (58-140)	4700 (2900-6900)
Burkina Faso	1990	Female	45 to 49	52% (39-62%)	48% (38-61%)	3 (1.8-4.5)	7% (4.2-10%)	9.6% (5.7-14%)	140 (83-200)	5800 (3600-8600)
Burkina Faso	1990	Female	50 to 54	53% (42-63%)	47% (37-58%)	3.2 (1.9-4.9)	8% (4.5-12%)	11% (6.1-16%)	170 (97-260)	6500 (3700-9700)
Burkina Faso	1990	Female	55 to 59	55% (44-66%)	45% (34-56%)	3.2 (2-5)	8.1% (4.6-12%)	11% (6.1-16%)	210 (120-310)	6600 (3800-9900)
Burkina Faso	1990	Female	60 to 64	56% (44-66%)	44% (34-56%)	3.1 (1.8-4.8)	7% (3.8-11%)	8.7% (4.7-14%)	220 (110-340)	5900 (3100-9300)
Burkina Faso	1990	Female	65 to 69	55% (45-65%)	45% (35-55%)	2.7 (1.7-4)	6.7% (3.3-11%)	8.1% (4.1-13%)	250 (130-410)	5600 (2800-9100)
Burkina Faso	1990	Female	70 to 74	53% (43-63%)	47% (37-57%)	2.2 (1.3-3.5)	6.2% (2.7-11%)	7.4% (3.2-13%)	250 (110-440)	4400 (1900-7900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	1990	Female	75 to 79	51% (37-64%)	49% (36-63%)	1.9 (1.1-3.2)	4.8% (1.5-9.2%)	5.6% (1.8-11%)	150 (49-300)	2100 (690-4200)
Burkina Faso	1990	Female	80 to 84	50% (34-65%)	50% (35-66%)	1.8 (0.99-2.9)	4% (1.2-7.9%)	4.7% (1.5-9.4%)	65 (21-130)	670 (220-1300)
Burkina Faso	1990	Female	85 to 89	51% (36-65%)	49% (35-64%)	1.8 (1.1-2.9)	3.5% (1-7.2%)	4.2% (1.1-8.5%)	30 (8.3-62)	220 (63-450)
Burkina Faso	1990	Female	90 to 94	53% (39-67%)	47% (33-61%)	1.9 (1.1-3.1)	3.1% (0.57-6.7%)	3.7% (0.58-8%)	8.9 (1.4-20)	46 (8.2-99)
Burkina Faso	1990	Female	95 plus	56% (37-73%)	44% (27-63%)	1.9 (0.94-3.2)	2.7% (0.15-6.9%)	3.1% (0.021-8.3%)	1.9 (0.013-5)	6.6 (0.37-17)
Burkina Faso	1990	Male	All Ages	25% (21-29%)	75% (71-79%)	1.4 (1-1.8)	1.8% (1.3-2.3%)	3.6% (2.5-4.8%)	2600 (1800-3400)	97000 (71000-120000)
Burkina Faso	1990	Male	15 to 19	20% (13-29%)	80% (71-87%)	1.1 (0.6-1.8)	2.6% (1.5-4.2%)	3.5% (1.8-5.7%)	36 (18-62)	3100 (1700-5000)
Burkina Faso	1990	Male	20 to 24	31% (20-41%)	69% (59-80%)	2.2 (1.2-3.5)	6.2% (3.8-9.3%)	7.6% (4.2-12%)	110 (63-170)	8200 (5000-12000)
Burkina Faso	1990	Male	25 to 29	37% (27-49%)	63% (51-73%)	2.7 (1.6-4.1)	6.9% (4.2-10%)	8.3% (4.5-13%)	140 (84-200)	9300 (6000-13000)
Burkina Faso	1990	Male	30 to 34	41% (30-53%)	59% (47-70%)	3.1 (1.9-4.6)	7.8% (4.9-11%)	9.4% (5.6-14%)	150 (95-220)	9300 (5900-13000)
Burkina Faso	1990	Male	35 to 39	44% (31-55%)	56% (45-69%)	3.6 (2.1-5.5)	9.2% (6.1-13%)	11% (7.2-16%)	170 (110-240)	9200 (6200-12000)
Burkina Faso	1990	Male	40 to 44	45% (33-55%)	55% (45-67%)	3.8 (2.4-5.7)	10% (7-14%)	13% (8.6-18%)	190 (130-260)	9100 (6200-12000)
Burkina Faso	1990	Male	45 to 49	46% (33-56%)	54% (44-67%)	3.9 (2.3-5.8)	12% (7.9-16%)	15% (9.9-20%)	230 (150-310)	9500 (6400-13000)
Burkina Faso	1990	Male	50 to 54	48% (37-59%)	52% (41-63%)	4.1 (2.4-6.4)	13% (8.5-17%)	16% (11-22%)	250 (160-340)	9100 (5900-12000)
Burkina Faso	1990	Male	55 to 59	50% (38-61%)	50% (39-62%)	4.4 (2.6-6.8)	13% (8.7-17%)	16% (11-21%)	260 (170-360)	8400 (5600-11000)
Burkina Faso	1990	Male	60 to 64	51% (37-63%)	49% (37-63%)	4.2 (2.4-6.4)	11% (6.7-15%)	13% (8.1-18%)	260 (160-360)	7200 (4400-9900)
Burkina Faso	1990	Male	65 to 69	50% (38-60%)	50% (40-62%)	3.6 (2.1-5.5)	9.7% (5.8-14%)	12% (7-17%)	270 (160-400)	6200 (3700-8900)
Burkina Faso	1990	Male	70 to 74	49% (38-60%)	51% (40-62%)	3 (1.7-4.6)	8.6% (4.7-13%)	10% (5.5-16%)	260 (140-410)	4700 (2500-7300)
Burkina Faso	1990	Male	75 to 79	49% (35-63%)	51% (37-65%)	2.5 (1.4-4.1)	6.7% (3-12%)	7.9% (3.5-14%)	160 (72-290)	2300 (1000-4000)
Burkina Faso	1990	Male	80 to 84	46% (32-60%)	54% (40-68%)	2 (1.1-3.2)	5.3% (2.3-9.1%)	6.2% (2.7-11%)	61 (26-110)	630 (270-1100)
Burkina Faso	1990	Male	85 to 89	41% (28-55%)	59% (45-72%)	1.5 (0.83-2.4)	3.8% (1.6-6.5%)	4.4% (1.8-7.8%)	23 (9.3-41)	170 (72-300)
Burkina Faso	1990	Male	90 to 94	40% (28-54%)	60% (46-72%)	1.2 (0.63-1.9)	2.3% (0.71-4.4%)	2.7% (0.72-5.2%)	4.8 (1.3-9.4)	25 (7.6-48)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	1990	Male	95 plus	42% (27-60%)	58% (40-73%)	0.94 (0.43-1.8)	2% (0.069-4.6%)	2.2% (-0.11-5.5%)	0.91 (-0.047-2.3)	3.3 (0.12-7.9)
Burkina Faso	1995	Female	All Ages	33% (29-38%)	67% (62-71%)	1.3 (0.97-1.7)	1.3% (0.88-1.9%)	2.8% (1.7-4.1%)	2100 (1300-3100)	70000 (47000-97000)
Burkina Faso	1995	Female	15 to 19	32% (23-42%)	68% (58-77%)	1.3 (0.77-2.1)	1.8% (1-2.9%)	2.4% (1.1-4.1%)	28 (13-49)	2600 (1500-4200)
Burkina Faso	1995	Female	20 to 24	38% (27-49%)	62% (51-73%)	2 (1.2-3)	2.9% (1.8-4.3%)	3.2% (1.7-5.1%)	50 (27-83)	4400 (2800-6700)
Burkina Faso	1995	Female	25 to 29	41% (30-53%)	59% (47-70%)	2.2 (1.4-3.4)	3.2% (1.9-5%)	3.6% (1.9-6%)	70 (40-110)	5200 (3200-8000)
Burkina Faso	1995	Female	30 to 34	45% (33-55%)	55% (45-67%)	2.5 (1.5-3.8)	3.4% (2.1-5.2%)	3.9% (2.2-6.4%)	77 (45-120)	5000 (3100-7600)
Burkina Faso	1995	Female	35 to 39	48% (36-58%)	52% (42-64%)	2.6 (1.6-3.9)	3.8% (2.4-5.5%)	4.6% (2.8-6.9%)	86 (52-130)	4900 (3100-7000)
Burkina Faso	1995	Female	40 to 44	51% (40-60%)	49% (40-60%)	2.9 (1.8-4.3)	4.8% (3-7%)	6.2% (3.8-9.1%)	110 (67-150)	5300 (3300-7500)
Burkina Faso	1995	Female	45 to 49	52% (39-62%)	48% (38-61%)	3.2 (1.9-4.9)	6.4% (4-9.3%)	8.4% (5.1-12%)	140 (88-220)	6200 (3800-9100)
Burkina Faso	1995	Female	50 to 54	54% (43-63%)	46% (37-57%)	3.4 (2-5.2)	7.8% (4.5-12%)	10% (5.9-15%)	200 (110-300)	7300 (4100-11000)
Burkina Faso	1995	Female	55 to 59	55% (43-65%)	45% (35-57%)	3.4 (2.1-5.4)	8.2% (4.9-12%)	10% (6.3-16%)	230 (130-350)	7400 (4200-11000)
Burkina Faso	1995	Female	60 to 64	56% (44-66%)	44% (34-56%)	3.3 (2-5.1)	7.2% (3.9-11%)	9% (4.9-14%)	240 (130-370)	6400 (3500-10000)
Burkina Faso	1995	Female	65 to 69	55% (44-65%)	45% (35-56%)	2.9 (1.8-4.4)	7.1% (3.6-11%)	8.6% (4.4-14%)	270 (140-440)	6200 (3100-9900)
Burkina Faso	1995	Female	70 to 74	53% (43-64%)	47% (36-57%)	2.4 (1.3-3.6)	6.6% (2.8-11%)	7.8% (3.4-13%)	280 (120-490)	5000 (2100-8700)
Burkina Faso	1995	Female	75 to 79	51% (37-64%)	49% (36-63%)	2 (1.1-3.2)	5.1% (1.9-9.6%)	6.1% (2.1-11%)	180 (66-340)	2500 (920-4800)
Burkina Faso	1995	Female	80 to 84	50% (34-65%)	50% (35-66%)	1.9 (1.1-3.1)	4.3% (1.5-8.2%)	5.1% (1.7-9.5%)	84 (28-160)	870 (300-1600)
Burkina Faso	1995	Female	85 to 89	50% (36-65%)	50% (35-64%)	1.9 (1.1-3.1)	3.8% (1.2-7.8%)	4.5% (1.4-9.1%)	42 (13-83)	310 (95-610)
Burkina Faso	1995	Female	90 to 94	53% (39-67%)	47% (33-61%)	1.9 (1.1-3.2)	3.4% (0.81-7.4%)	4.1% (0.9-8.8%)	13 (2.9-28)	66 (15-140)
Burkina Faso	1995	Female	95 plus	56% (37-73%)	44% (27-63%)	2 (0.98-3.5)	3.1% (0.21-7.5%)	3.6% (0.12-8.8%)	2.8 (0.095-7.1)	9.8 (0.67-24)
Burkina Faso	1995	Male	All Ages	25% (21-29%)	75% (71-79%)	1.5 (1.1-2)	1.9% (1.4-2.5%)	3.8% (2.7-5%)	3000 (2200-3900)	110000 (85000-140000)
Burkina Faso	1995	Male	15 to 19	20% (14-29%)	80% (71-86%)	1.2 (0.67-1.9)	2.9% (1.7-4.6%)	3.9% (2.1-6.4%)	47 (25-78)	4000 (2300-6200)
Burkina Faso	1995	Male	20 to 24	31% (21-41%)	69% (59-79%)	2.4 (1.3-3.7)	6.7% (4.2-9.8%)	8.3% (4.9-12%)	130 (77-200)	9900 (6200-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	1995	Male	25 to 29	38% (27-49%)	62% (51-73%)	2.9 (1.8-4.4)	6.9% (4.3-10%)	8.1% (4.7-12%)	170 (110-250)	11000 (7400-16000)
Burkina Faso	1995	Male	30 to 34	42% (30-53%)	58% (47-70%)	3.4 (2-5.1)	6.9% (4.5-9.8%)	8% (5-12%)	190 (120-270)	11000 (7600-16000)
Burkina Faso	1995	Male	35 to 39	44% (31-55%)	56% (45-69%)	3.9 (2.4-5.9)	8.1% (5.4-11%)	9.6% (6.2-14%)	210 (150-300)	11000 (8000-15000)
Burkina Faso	1995	Male	40 to 44	45% (34-56%)	55% (44-66%)	4.2 (2.5-6.2)	9.1% (6.3-12%)	11% (7.5-15%)	220 (150-290)	11000 (7400-14000)
Burkina Faso	1995	Male	45 to 49	46% (33-56%)	54% (44-67%)	4.3 (2.4-6.4)	11% (7.4-14%)	13% (8.8-18%)	250 (170-340)	11000 (7300-14000)
Burkina Faso	1995	Male	50 to 54	48% (37-59%)	52% (41-63%)	4.5 (2.5-6.9)	12% (8.6-16%)	15% (11-20%)	290 (190-390)	11000 (7100-14000)
Burkina Faso	1995	Male	55 to 59	51% (39-61%)	49% (39-61%)	4.8 (2.9-7.3)	13% (9.1-17%)	16% (11-21%)	300 (200-400)	9500 (6500-13000)
Burkina Faso	1995	Male	60 to 64	51% (37-63%)	49% (37-63%)	4.5 (2.7-7)	11% (7.2-15%)	13% (8.7-18%)	280 (170-380)	7600 (4700-10000)
Burkina Faso	1995	Male	65 to 69	50% (38-61%)	50% (39-62%)	3.9 (2.3-5.9)	10% (6.3-14%)	12% (7.6-17%)	290 (170-410)	6500 (4000-9300)
Burkina Faso	1995	Male	70 to 74	50% (39-61%)	50% (39-61%)	3.3 (1.8-5)	9.4% (5.1-14%)	11% (6.1-17%)	290 (160-450)	5300 (2900-8000)
Burkina Faso	1995	Male	75 to 79	49% (35-63%)	51% (37-65%)	2.7 (1.5-4.5)	7.4% (3.5-13%)	8.7% (4-15%)	190 (91-330)	2700 (1300-4600)
Burkina Faso	1995	Male	80 to 84	46% (32-61%)	54% (39-68%)	2.2 (1.2-3.6)	5.9% (2.7-10%)	6.9% (3.2-12%)	78 (36-140)	810 (370-1400)
Burkina Faso	1995	Male	85 to 89	42% (29-56%)	58% (44-71%)	1.6 (0.89-2.7)	4.3% (2-7.7%)	5% (2.3-9%)	32 (15-58)	240 (110-430)
Burkina Faso	1995	Male	90 to 94	41% (28-54%)	59% (46-72%)	1.3 (0.69-2.2)	2.6% (0.9-5.2%)	3% (0.98-5.9%)	6.9 (2.2-14)	36 (12-71)
Burkina Faso	1995	Male	95 plus	43% (26-61%)	57% (39-74%)	1 (0.44-2)	2.2% (0.25-5.4%)	2.5% (0.079-6.3%)	1.3 (0.042-3.3)	4.7 (0.52-11)
Burkina Faso	2000	Female	All Ages	31% (27-35%)	69% (65-73%)	1.3 (0.93-1.6)	1.4% (0.97-2%)	3% (1.9-4.4%)	2400 (1500-3500)	81000 (56000-110000)
Burkina Faso	2000	Female	15 to 19	30% (23-37%)	70% (63-77%)	1.3 (0.79-2)	1.9% (1.1-3%)	2.4% (1.3-4.2%)	33 (17-56)	3000 (1800-4800)
Burkina Faso	2000	Female	20 to 24	33% (26-42%)	67% (58-74%)	1.8 (1.1-2.8)	3.2% (2.1-4.7%)	3.6% (1.9-5.9%)	60 (32-99)	5300 (3300-7800)
Burkina Faso	2000	Female	25 to 29	36% (28-44%)	64% (56-72%)	2.1 (1.3-3.1)	3.3% (2.1-4.9%)	3.7% (2.1-6%)	78 (46-120)	5800 (3700-8500)
Burkina Faso	2000	Female	30 to 34	37% (30-47%)	63% (53-70%)	2.2 (1.4-3.3)	3.2% (2-4.8%)	3.6% (2.1-5.8%)	86 (52-140)	5600 (3600-8600)
Burkina Faso	2000	Female	35 to 39	43% (36-51%)	57% (49-64%)	2.5 (1.6-3.7)	3.5% (2.2-5.2%)	4.2% (2.4-6.4%)	99 (61-140)	5600 (3600-7900)
Burkina Faso	2000	Female	40 to 44	47% (39-55%)	53% (45-61%)	2.9 (1.8-4.2)	4.9% (3.1-6.8%)	6.1% (3.8-8.7%)	130 (83-190)	6500 (4200-9100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	2000	Female	45 to 49	50% (40-58%)	50% (42-60%)	3.3 (2-5)	6.7% (4.2-9.6%)	8.7% (5.4-12%)	180 (110-250)	7600 (4700-11000)
Burkina Faso	2000	Female	50 to 54	54% (45-62%)	46% (38-55%)	3.7 (2.2-5.5)	8.4% (5.1-12%)	11% (6.6-15%)	230 (130-330)	8500 (5100-12000)
Burkina Faso	2000	Female	55 to 59	56% (47-65%)	44% (35-53%)	3.7 (2.3-5.7)	9.1% (5.6-13%)	12% (7.2-17%)	280 (170-410)	8900 (5300-13000)
Burkina Faso	2000	Female	60 to 64	57% (47-65%)	43% (35-53%)	3.5 (2.2-5.4)	8% (4.5-12%)	9.9% (5.7-15%)	280 (160-430)	7700 (4200-12000)
Burkina Faso	2000	Female	65 to 69	55% (45-64%)	45% (36-55%)	3.1 (1.9-4.6)	7.6% (4.1-12%)	9.2% (5-15%)	300 (160-500)	6800 (3600-11000)
Burkina Faso	2000	Female	70 to 74	52% (42-61%)	48% (39-58%)	2.5 (1.4-3.8)	6.5% (2.7-11%)	7.7% (3.2-13%)	280 (110-500)	5100 (2000-9100)
Burkina Faso	2000	Female	75 to 79	49% (36-62%)	51% (38-64%)	2.1 (1.1-3.5)	5% (1.5-9.6%)	5.9% (1.8-11%)	190 (57-350)	2600 (790-5000)
Burkina Faso	2000	Female	80 to 84	50% (34-65%)	50% (35-66%)	2.1 (1.1-3.3)	4.4% (1.4-8.4%)	5.2% (1.6-9.8%)	95 (29-180)	990 (300-1800)
Burkina Faso	2000	Female	85 to 89	52% (37-66%)	48% (34-63%)	2.1 (1.2-3.3)	4.1% (1.3-7.9%)	4.8% (1.4-9.3%)	53 (16-100)	390 (120-750)
Burkina Faso	2000	Female	90 to 94	55% (41-68%)	45% (32-59%)	2.2 (1.2-3.5)	3.7% (0.86-7.7%)	4.4% (0.97-9.3%)	18 (3.9-37)	89 (21-190)
Burkina Faso	2000	Female	95 plus	57% (39-73%)	43% (27-61%)	2.2 (1.1-3.7)	3.4% (0.36-8.2%)	3.9% (0.23-9.6%)	4 (0.23-9.9)	14 (1.5-33)
Burkina Faso	2000	Male	All Ages	24% (21-28%)	76% (72-79%)	1.5 (1.2-2)	2.1% (1.6-2.7%)	4.2% (3.1-5.5%)	3400 (2600-4300)	130000 (100000-160000)
Burkina Faso	2000	Male	15 to 19	21% (16-27%)	79% (73-84%)	1.3 (0.76-2)	3.2% (1.9-4.8%)	4.3% (2.4-6.7%)	59 (32-94)	5000 (3000-7300)
Burkina Faso	2000	Male	20 to 24	29% (23-36%)	71% (64-77%)	2.4 (1.4-3.5)	7.1% (4.6-9.8%)	8.9% (5.6-13%)	150 (95-220)	11000 (7500-16000)
Burkina Faso	2000	Male	25 to 29	32% (26-39%)	68% (61-74%)	2.6 (1.7-3.9)	7.4% (5-10%)	8.8% (5.5-13%)	180 (120-250)	12000 (8300-16000)
Burkina Faso	2000	Male	30 to 34	38% (31-46%)	62% (54-69%)	3.3 (2.1-4.9)	7.4% (5.2-10%)	8.7% (5.9-12%)	220 (150-290)	13000 (9400-17000)
Burkina Faso	2000	Male	35 to 39	43% (37-50%)	57% (50-63%)	4.1 (2.7-5.9)	8.1% (5.8-11%)	9.5% (6.6-13%)	250 (180-330)	13000 (9900-17000)
Burkina Faso	2000	Male	40 to 44	43% (36-50%)	57% (50-64%)	4.2 (2.7-6.1)	8.9% (6.5-11%)	11% (7.6-14%)	260 (190-340)	12000 (9200-16000)
Burkina Faso	2000	Male	45 to 49	45% (36-52%)	55% (48-64%)	4.4 (2.7-6.5)	11% (7.6-14%)	13% (9.2-17%)	280 (210-360)	12000 (8800-15000)
Burkina Faso	2000	Male	50 to 54	49% (41-56%)	51% (44-59%)	4.9 (3-7.3)	13% (9.3-16%)	16% (11-20%)	320 (230-400)	12000 (8500-15000)
Burkina Faso	2000	Male	55 to 59	53% (45-61%)	47% (39-55%)	5.3 (3.4-8)	14% (10-18%)	17% (13-22%)	350 (250-450)	11000 (8200-14000)
Burkina Faso	2000	Male	60 to 64	54% (44-63%)	46% (37-56%)	5.1 (3.2-7.6)	13% (8.6-16%)	15% (10-19%)	330 (230-430)	9100 (6300-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	2000	Male	65 to 69	53% (45-61%)	47% (39-55%)	4.4 (2.8-6.4)	12% (7.6-15%)	14% (9-18%)	320 (200-430)	7200 (4600-9500)
Burkina Faso	2000	Male	70 to 74	55% (46-63%)	45% (37-54%)	3.9 (2.3-5.7)	11% (5.8-15%)	13% (6.8-18%)	320 (170-460)	5700 (3100-8400)
Burkina Faso	2000	Male	75 to 79	57% (46-68%)	43% (32-54%)	3.4 (1.9-5.4)	8.5% (3.9-14%)	10% (4.6-16%)	220 (100-360)	3100 (1500-5100)
Burkina Faso	2000	Male	80 to 84	54% (39-67%)	46% (33-61%)	2.7 (1.5-4.4)	6.8% (3.1-11%)	7.9% (3.5-13%)	95 (43-160)	1000 (450-1700)
Burkina Faso	2000	Male	85 to 89	47% (33-61%)	53% (39-67%)	2 (1.1-3.1)	4.9% (2.1-8.2%)	5.7% (2.3-9.6%)	41 (17-71)	310 (130-530)
Burkina Faso	2000	Male	90 to 94	46% (33-59%)	54% (41-67%)	1.6 (0.86-2.6)	3% (0.95-5.9%)	3.5% (0.94-6.9%)	9.7 (2.6-19)	51 (16-98)
Burkina Faso	2000	Male	95 plus	47% (31-64%)	53% (36-69%)	1.2 (0.56-2.3)	2.7% (0.37-6%)	3% (0.23-7.2%)	1.9 (0.15-4.6)	6.9 (0.98-16)
Burkina Faso	2005	Female	All Ages	30% (27-34%)	70% (66-73%)	1.2 (0.9-1.6)	1.5% (1-2.2%)	3.1% (1.9-4.6%)	2500 (1500-3700)	87000 (61000-120000)
Burkina Faso	2005	Female	15 to 19	30% (23-37%)	70% (63-77%)	1.3 (0.78-2)	1.8% (1-2.8%)	2.2% (1-3.8%)	34 (16-59)	3200 (1800-5000)
Burkina Faso	2005	Female	20 to 24	33% (25-41%)	67% (59-75%)	1.8 (1.1-2.7)	3.3% (2.1-4.9%)	3.9% (2.1-6.1%)	66 (35-100)	5900 (3700-8600)
Burkina Faso	2005	Female	25 to 29	35% (28-43%)	65% (57-72%)	2 (1.3-3)	3.9% (2.5-5.8%)	4.7% (2.6-7.4%)	88 (52-140)	6700 (4400-9900)
Burkina Faso	2005	Female	30 to 34	37% (29-46%)	63% (54-71%)	2.1 (1.3-3.3)	3.7% (2.4-5.3%)	4.4% (2.7-6.7%)	96 (59-150)	6300 (4200-9200)
Burkina Faso	2005	Female	35 to 39	43% (36-51%)	57% (49-64%)	2.5 (1.6-3.7)	4.1% (2.6-5.9%)	5% (3-7.6%)	120 (72-170)	6700 (4400-9600)
Burkina Faso	2005	Female	40 to 44	47% (38-55%)	53% (45-62%)	2.8 (1.7-4.1)	5.3% (3.4-7.4%)	6.8% (4.3-9.7%)	150 (99-220)	7600 (4900-11000)
Burkina Faso	2005	Female	45 to 49	49% (40-58%)	51% (42-60%)	3.2 (1.9-4.9)	7.3% (4.5-10%)	9.6% (5.9-14%)	210 (130-300)	8900 (5500-13000)
Burkina Faso	2005	Female	50 to 54	53% (44-62%)	47% (38-56%)	3.6 (2.1-5.5)	8.6% (5-12%)	11% (6.5-16%)	250 (140-360)	9200 (5400-13000)
Burkina Faso	2005	Female	55 to 59	56% (47-64%)	44% (36-53%)	3.6 (2.3-5.6)	9.1% (5.4-13%)	12% (7.1-17%)	290 (170-420)	9200 (5500-14000)
Burkina Faso	2005	Female	60 to 64	56% (47-65%)	44% (35-53%)	3.4 (2.1-5.4)	7.9% (4.4-12%)	9.8% (5.6-15%)	300 (160-450)	8000 (4400-12000)
Burkina Faso	2005	Female	65 to 69	55% (46-64%)	45% (36-54%)	3 (1.8-4.6)	7.5% (3.7-12%)	9% (4.4-15%)	310 (150-510)	7100 (3400-11000)
Burkina Faso	2005	Female	70 to 74	52% (42-61%)	48% (39-58%)	2.4 (1.4-3.8)	5.9% (2-11%)	7% (2.4-13%)	260 (90-480)	4800 (1600-8700)
Burkina Faso	2005	Female	75 to 79	49% (36-62%)	51% (38-64%)	2 (1.1-3.4)	4.5% (0.98-8.7%)	5.3% (1.2-10%)	170 (38-340)	2400 (530-4700)
Burkina Faso	2005	Female	80 to 84	50% (34-65%)	50% (35-66%)	2 (1.1-3.2)	3.9% (0.95-7.7%)	4.6% (1.1-9.1%)	89 (20-170)	930 (220-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	2005	Female	85 to 89	52% (37-66%)	48% (34-63%)	2.1 (1.1-3.2)	3.6% (0.69-7.3%)	4.2% (0.73-8.6%)	51 (9-100)	380 (75-770)
Burkina Faso	2005	Female	90 to 94	55% (41-68%)	45% (32-59%)	2.1 (1.2-3.4)	3.3% (0.31-7.3%)	3.8% (0.27-8.8%)	18 (1.3-42)	94 (9-210)
Burkina Faso	2005	Female	95 plus	57% (39-73%)	43% (27-61%)	2.1 (1.1-3.6)	3% (-0.022-7.8%)	3.5% (-0.19-9%)	4.4 (-0.25-11)	15 (-0.11-39)
Burkina Faso	2005	Male	All Ages	24% (21-27%)	76% (73-79%)	1.5 (1.1-2)	2.1% (1.6-2.8%)	4.2% (3-5.6%)	3400 (2500-4400)	130000 (100000-160000)
Burkina Faso	2005	Male	15 to 19	21% (15-27%)	79% (73-85%)	1.3 (0.75-2.1)	2.7% (1.6-4.2%)	3.5% (1.9-5.7%)	55 (30-89)	4800 (2900-7100)
Burkina Faso	2005	Male	20 to 24	29% (23-36%)	71% (64-77%)	2.3 (1.4-3.5)	6.6% (4.4-9.3%)	8.1% (5.1-12%)	150 (92-220)	11000 (7500-16000)
Burkina Faso	2005	Male	25 to 29	32% (25-39%)	68% (61-75%)	2.6 (1.6-3.8)	7.9% (5.4-11%)	9.8% (6.3-14%)	190 (120-260)	13000 (8900-17000)
Burkina Faso	2005	Male	30 to 34	38% (31-46%)	62% (54-69%)	3.2 (2.1-4.8)	8.5% (5.9-11%)	10% (7-14%)	220 (150-300)	14000 (9400-18000)
Burkina Faso	2005	Male	35 to 39	43% (36-49%)	57% (51-64%)	4 (2.6-5.8)	9% (6.7-12%)	11% (7.8-15%)	270 (190-340)	15000 (11000-18000)
Burkina Faso	2005	Male	40 to 44	43% (36-50%)	57% (50-64%)	4.1 (2.6-6)	9.5% (7-12%)	12% (8.4-15%)	290 (210-370)	14000 (10000-17000)
Burkina Faso	2005	Male	45 to 49	44% (37-52%)	56% (48-63%)	4.3 (2.7-6.3)	11% (8.1-14%)	14% (9.9-18%)	320 (230-400)	13000 (9700-17000)
Burkina Faso	2005	Male	50 to 54	49% (41-57%)	51% (43-59%)	4.8 (2.9-7)	13% (9.2-16%)	16% (11-20%)	320 (230-410)	12000 (8600-15000)
Burkina Faso	2005	Male	55 to 59	53% (45-61%)	47% (39-55%)	5.3 (3.3-7.9)	14% (10-18%)	17% (12-22%)	350 (250-440)	11000 (8200-14000)
Burkina Faso	2005	Male	60 to 64	53% (44-63%)	47% (37-56%)	5 (3.1-7.6)	13% (8.7-16%)	15% (10-19%)	350 (230-440)	9500 (6400-12000)
Burkina Faso	2005	Male	65 to 69	53% (44-61%)	47% (39-56%)	4.3 (2.6-6.3)	11% (7.2-15%)	14% (8.6-18%)	320 (200-430)	7200 (4600-9700)
Burkina Faso	2005	Male	70 to 74	54% (45-63%)	46% (37-55%)	3.8 (2.3-5.7)	10% (5.3-15%)	12% (6.3-18%)	290 (150-430)	5200 (2800-7700)
Burkina Faso	2005	Male	75 to 79	57% (46-69%)	43% (31-54%)	3.4 (1.8-5.4)	8.1% (3.4-14%)	9.4% (4.1-16%)	200 (88-330)	2800 (1200-4700)
Burkina Faso	2005	Male	80 to 84	53% (38-67%)	47% (33-62%)	2.7 (1.4-4.3)	6.3% (2.7-11%)	7.3% (3.1-13%)	88 (37-160)	920 (400-1600)
Burkina Faso	2005	Male	85 to 89	46% (33-60%)	54% (40-67%)	1.9 (1.1-3.2)	4.4% (1.8-7.8%)	5.1% (2-9.1%)	40 (16-72)	300 (120-540)
Burkina Faso	2005	Male	90 to 94	45% (32-58%)	55% (42-68%)	1.5 (0.85-2.5)	2.7% (0.74-5.4%)	3.1% (0.71-6.3%)	10 (2.3-20)	53 (14-100)
Burkina Faso	2005	Male	95 plus	47% (31-64%)	53% (36-69%)	1.2 (0.54-2.2)	2.3% (0.32-5.5%)	2.6% (0.14-6.3%)	2.1 (0.11-5)	7.5 (1-17)
Burkina Faso	2010	Female	All Ages	33% (28-38%)	67% (62-72%)	1.2 (0.83-1.7)	1.6% (0.92-2.3%)	3.1% (1.5-4.8%)	2500 (1300-3800)	89000 (54000-130000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	2010	Female	15 to 19	32% (23-42%)	68% (58-77%)	1.2 (0.67-2)	1.7% (0.93-2.8%)	2% (0.87-3.6%)	35 (15-62)	3400 (1800-5600)
Burkina Faso	2010	Female	20 to 24	38% (27-49%)	62% (51-73%)	1.8 (1-3.1)	3.3% (2-5.1%)	3.8% (1.9-6.4%)	68 (34-120)	6300 (3700-9500)
Burkina Faso	2010	Female	25 to 29	41% (30-53%)	59% (47-70%)	2.1 (1.2-3.4)	4.1% (2.5-6.1%)	5% (2.7-8.1%)	92 (50-150)	7200 (4300-11000)
Burkina Faso	2010	Female	30 to 34	44% (32-55%)	56% (45-68%)	2.3 (1.3-3.7)	4.2% (2.6-6.4%)	5.4% (3-8.5%)	100 (57-160)	7000 (4200-10000)
Burkina Faso	2010	Female	35 to 39	48% (36-58%)	52% (42-64%)	2.5 (1.4-4)	4.5% (2.7-6.8%)	5.9% (3.3-9.2%)	120 (67-190)	7000 (4200-10000)
Burkina Faso	2010	Female	40 to 44	50% (40-60%)	50% (40-60%)	2.7 (1.5-4.4)	5.5% (3.3-8.1%)	7.5% (4.3-11%)	160 (91-230)	7800 (4500-11000)
Burkina Faso	2010	Female	45 to 49	52% (40-62%)	48% (38-60%)	3 (1.6-4.9)	7.4% (4.1-11%)	10% (5.6-15%)	210 (120-320)	9100 (5100-14000)
Burkina Faso	2010	Female	50 to 54	53% (43-63%)	47% (37-57%)	3.2 (1.8-5.1)	8.4% (4.7-13%)	11% (6.3-17%)	260 (140-390)	9700 (5400-15000)
Burkina Faso	2010	Female	55 to 59	55% (44-65%)	45% (35-56%)	3.2 (1.9-5.3)	8.7% (4.6-14%)	11% (6-17%)	300 (160-460)	9500 (5100-15000)
Burkina Faso	2010	Female	60 to 64	56% (44-66%)	44% (34-56%)	3.1 (1.7-5.1)	7.3% (3.1-12%)	9% (4-15%)	280 (130-470)	7800 (3500-13000)
Burkina Faso	2010	Female	65 to 69	55% (45-64%)	45% (36-55%)	2.7 (1.6-4.3)	6.6% (2.6-11%)	8% (3.3-14%)	310 (120-530)	6900 (2700-12000)
Burkina Faso	2010	Female	70 to 74	53% (43-62%)	47% (38-57%)	2.2 (1.2-3.5)	5% (0.98-9.9%)	6% (1.2-12%)	240 (46-470)	4400 (850-8500)
Burkina Faso	2010	Female	75 to 79	51% (37-64%)	49% (36-63%)	1.9 (0.94-3.2)	3.7% (0.17-8.4%)	4.4% (0.16-9.9%)	150 (5.6-340)	2200 (95-4800)
Burkina Faso	2010	Female	80 to 84	49% (34-64%)	51% (36-66%)	1.8 (0.88-2.9)	3.2% (0.13-6.9%)	3.7% (0.099-8.2%)	76 (2-170)	800 (34-1800)
Burkina Faso	2010	Female	85 to 89	50% (35-64%)	50% (36-65%)	1.7 (0.91-2.9)	2.7% (-0.037-6.4%)	3.2% (-0.083-7.6%)	43 (-1.2-100)	320 (-4.5-760)
Burkina Faso	2010	Female	90 to 94	53% (38-67%)	47% (33-62%)	1.8 (0.91-3)	2.4% (-0.39-6.2%)	2.8% (-0.62-7.3%)	16 (-3.6-41)	82 (-13-210)
Burkina Faso	2010	Female	95 plus	55% (36-73%)	45% (27-64%)	1.8 (0.86-3.3)	2.2% (-0.52-6.7%)	2.5% (-0.8-7.8%)	3.9 (-1.3-12)	14 (-3.3-42)
Burkina Faso	2010	Male	All Ages	25% (21-30%)	75% (70-79%)	1.4 (0.98-2)	2.3% (1.5-3.1%)	4.3% (2.8-5.8%)	3600 (2400-4700)	140000 (100000-180000)
Burkina Faso	2010	Male	15 to 19	20% (14-28%)	80% (72-86%)	1.1 (0.58-1.8)	2.5% (1.4-3.9%)	3.1% (1.5-5.2%)	53 (27-91)	4800 (2600-7500)
Burkina Faso	2010	Male	20 to 24	31% (21-42%)	69% (58-79%)	2.2 (1.2-3.6)	6.5% (4-9.5%)	7.8% (4.3-12%)	150 (86-240)	12000 (7400-18000)
Burkina Faso	2010	Male	25 to 29	38% (27-49%)	62% (51-73%)	2.7 (1.5-4.2)	8.7% (5.6-12%)	11% (6.6-16%)	220 (130-320)	15000 (9900-21000)
Burkina Faso	2010	Male	30 to 34	41% (30-52%)	59% (48-70%)	3.1 (1.8-5)	9.6% (6.4-13%)	12% (7.7-17%)	260 (160-350)	16000 (11000-21000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	2010	Male	35 to 39	44% (31-54%)	56% (46-69%)	3.6 (2-5.8)	10% (6.8-14%)	13% (8.4-18%)	290 (190-390)	16000 (11000-21000)
Burkina Faso	2010	Male	40 to 44	45% (34-55%)	55% (45-66%)	3.9 (2.2-6.3)	11% (7.4-15%)	14% (9.2-19%)	330 (220-430)	16000 (11000-21000)
Burkina Faso	2010	Male	45 to 49	46% (32-56%)	54% (44-68%)	3.9 (2.2-6.3)	13% (8.1-16%)	16% (10-21%)	360 (230-470)	15000 (9800-20000)
Burkina Faso	2010	Male	50 to 54	48% (36-59%)	52% (41-64%)	4.2 (2.2-6.7)	13% (8.3-17%)	16% (10-22%)	360 (230-480)	13000 (8600-18000)
Burkina Faso	2010	Male	55 to 59	50% (39-61%)	50% (39-61%)	4.4 (2.5-7.4)	13% (8.5-18%)	17% (11-22%)	360 (230-480)	11000 (7500-15000)
Burkina Faso	2010	Male	60 to 64	51% (37-63%)	49% (37-63%)	4.2 (2.3-7)	12% (6.7-16%)	14% (8.1-19%)	330 (190-470)	9200 (5300-13000)
Burkina Faso	2010	Male	65 to 69	50% (38-61%)	50% (39-62%)	3.6 (2-5.8)	10% (5.5-14%)	12% (6.5-17%)	310 (170-460)	7100 (3800-10000)
Burkina Faso	2010	Male	70 to 74	50% (39-60%)	50% (40-61%)	3 (1.7-4.9)	8.4% (3.7-14%)	9.9% (4.4-16%)	260 (120-420)	4700 (2100-7700)
Burkina Faso	2010	Male	75 to 79	49% (35-63%)	51% (37-65%)	2.6 (1.3-4.4)	6.4% (2.3-12%)	7.4% (2.7-14%)	160 (57-300)	2300 (820-4200)
Burkina Faso	2010	Male	80 to 84	47% (32-60%)	53% (40-68%)	2.1 (1-3.6)	4.9% (1.7-9.7%)	5.7% (1.9-11%)	70 (23-140)	740 (250-1400)
Burkina Faso	2010	Male	85 to 89	42% (29-56%)	58% (44-71%)	1.6 (0.8-2.6)	3.5% (1.1-6.8%)	4% (1.1-8%)	33 (9.5-66)	250 (77-490)
Burkina Faso	2010	Male	90 to 94	41% (28-55%)	59% (45-72%)	1.2 (0.61-2.1)	2.1% (0.39-4.6%)	2.3% (0.32-5.3%)	8.4 (1.1-19)	45 (8.2-100)
Burkina Faso	2010	Male	95 plus	43% (27-60%)	57% (40-73%)	0.97 (0.41-1.8)	1.8% (-0.051-4.6%)	1.9% (-0.31-5.3%)	1.8 (-0.3-5.1)	6.8 (-0.2-18)
Burkina Faso	2016	Female	All Ages	33% (29-38%)	67% (62-71%)	1.1 (0.63-2)	1.7% (0.79-2.9%)	3.3% (1.2-5.9%)	2600 (890-4700)	94000 (45000-150000)
Burkina Faso	2016	Female	15 to 19	32% (23-41%)	68% (59-77%)	1.2 (0.56-2.2)	1.7% (0.84-3%)	2% (0.74-4.1%)	35 (12-69)	3500 (1800-6100)
Burkina Faso	2016	Female	20 to 24	38% (27-49%)	62% (51-73%)	1.7 (0.81-3.3)	3.3% (1.8-5.3%)	3.7% (1.6-6.8%)	68 (29-130)	6600 (3700-11000)
Burkina Faso	2016	Female	25 to 29	41% (30-53%)	59% (47-70%)	2 (0.95-3.7)	4% (2.2-6.5%)	5% (2.2-8.8%)	89 (42-160)	7300 (4000-12000)
Burkina Faso	2016	Female	30 to 34	45% (33-55%)	55% (45-67%)	2.2 (1-3.9)	4.3% (2.3-7%)	5.8% (2.7-9.7%)	110 (50-180)	7300 (3900-12000)
Burkina Faso	2016	Female	35 to 39	48% (36-58%)	52% (42-64%)	2.3 (1.1-4.3)	4.9% (2.6-8%)	6.9% (3.3-12%)	130 (63-230)	7800 (4100-13000)
Burkina Faso	2016	Female	40 to 44	51% (39-60%)	49% (40-61%)	2.5 (1.2-4.7)	5.8% (3-9.3%)	8.4% (4-13%)	160 (79-260)	8200 (4200-13000)
Burkina Faso	2016	Female	45 to 49	52% (40-62%)	48% (38-60%)	2.8 (1.3-5.4)	7.2% (3.3-12%)	10% (4.7-17%)	220 (100-370)	9700 (4500-16000)
Burkina Faso	2016	Female	50 to 54	53% (42-63%)	47% (37-58%)	3 (1.4-5.6)	8% (3.4-13%)	11% (4.8-18%)	280 (120-450)	10000 (4500-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	2016	Female	55 to 59	55% (44-65%)	45% (35-56%)	3.1 (1.4-5.9)	8% (3.2-14%)	11% (4.3-18%)	320 (130-540)	10000 (4100-17000)
Burkina Faso	2016	Female	60 to 64	56% (44-66%)	44% (34-56%)	2.9 (1.4-5.8)	6.6% (2.1-12%)	8.3% (2.7-15%)	300 (96-540)	8100 (2500-15000)
Burkina Faso	2016	Female	65 to 69	55% (45-65%)	45% (35-55%)	2.5 (1.2-4.6)	6.1% (1.3-12%)	7.4% (1.7-15%)	300 (69-600)	6900 (1500-14000)
Burkina Faso	2016	Female	70 to 74	53% (43-64%)	47% (36-57%)	2.1 (0.97-3.8)	4.6% (0.28-11%)	5.4% (0.32-13%)	250 (15-600)	4500 (260-11000)
Burkina Faso	2016	Female	75 to 79	51% (36-64%)	49% (36-64%)	1.8 (0.73-3.4)	3.3% (-0.45-9.3%)	3.9% (-0.58-11%)	160 (-23-440)	2200 (-290-6200)
Burkina Faso	2016	Female	80 to 84	49% (33-64%)	51% (36-67%)	1.7 (0.74-3.1)	2.8% (-0.37-7.9%)	3.3% (-0.44-9.3%)	76 (-10-220)	800 (-100-2200)
Burkina Faso	2016	Female	85 to 89	50% (35-65%)	50% (35-65%)	1.7 (0.74-3.2)	2.4% (-0.58-7.2%)	2.8% (-0.73-8.4%)	41 (-11-120)	310 (-72-910)
Burkina Faso	2016	Female	90 to 94	53% (38-66%)	47% (34-62%)	1.7 (0.75-3.2)	2.1% (-0.86-7.1%)	2.4% (-1.1-8.3%)	15 (-7.3-53)	80 (-32-270)
Burkina Faso	2016	Female	95 plus	55% (37-72%)	45% (28-63%)	1.7 (0.72-3.6)	2% (-0.91-7%)	2.2% (-1.3-8.2%)	4.1 (-2.5-16)	15 (-6.9-53)
Burkina Faso	2016	Male	All Ages	26% (22-30%)	74% (70-78%)	1.4 (0.77-2.4)	2.6% (1.6-3.8%)	4.7% (2.7-6.9%)	3800 (2200-5500)	160000 (95000-220000)
Burkina Faso	2016	Male	15 to 19	21% (14-29%)	79% (71-86%)	1.1 (0.49-2.1)	2.5% (1.3-4.3%)	3.1% (1.4-5.9%)	56 (25-110)	5200 (2600-8900)
Burkina Faso	2016	Male	20 to 24	31% (21-41%)	69% (59-79%)	2.1 (0.99-4)	6.2% (3.5-9.5%)	7.3% (3.6-12%)	160 (80-260)	13000 (7200-20000)
Burkina Faso	2016	Male	25 to 29	38% (27-49%)	62% (51-73%)	2.6 (1.2-4.7)	8.3% (4.8-12%)	10% (5.5-16%)	230 (120-350)	16000 (9400-24000)
Burkina Faso	2016	Male	30 to 34	42% (30-53%)	58% (47-70%)	3 (1.5-5.3)	9.4% (5.7-13%)	12% (6.6-17%)	280 (150-400)	17000 (10000-24000)
Burkina Faso	2016	Male	35 to 39	44% (32-55%)	56% (45-68%)	3.4 (1.7-6.1)	11% (6.4-15%)	13% (7.8-19%)	330 (190-460)	18000 (11000-25000)
Burkina Faso	2016	Male	40 to 44	45% (34-56%)	55% (44-66%)	3.7 (1.7-6.8)	11% (7.2-16%)	15% (8.8-20%)	360 (210-490)	17000 (11000-24000)
Burkina Faso	2016	Male	45 to 49	46% (33-56%)	54% (44-67%)	3.8 (1.8-7)	13% (7.1-17%)	16% (8.9-22%)	410 (220-570)	17000 (9700-24000)
Burkina Faso	2016	Male	50 to 54	48% (37-58%)	52% (42-63%)	4 (1.7-7.1)	12% (6.8-17%)	16% (8.4-22%)	410 (220-570)	15000 (8400-21000)
Burkina Faso	2016	Male	55 to 59	51% (39-61%)	49% (39-61%)	4.2 (2-8)	13% (6.8-17%)	16% (8.6-21%)	410 (230-570)	13000 (7400-18000)
Burkina Faso	2016	Male	60 to 64	51% (37-63%)	49% (37-63%)	4 (1.9-7.7)	11% (5.1-16%)	13% (6.1-19%)	360 (170-520)	9800 (4800-14000)
Burkina Faso	2016	Male	65 to 69	50% (39-61%)	50% (39-61%)	3.4 (1.6-6.4)	9.1% (4-14%)	11% (4.8-17%)	310 (140-500)	7200 (3200-11000)
Burkina Faso	2016	Male	70 to 74	50% (39-60%)	50% (40-61%)	2.9 (1.4-5.1)	7.6% (2.7-13%)	9% (3.1-16%)	270 (94-490)	4900 (1700-8800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Burkina Faso	2016	Male	75 to 79	50% (36-63%)	50% (37-64%)	2.4 (1.1-4.8)	5.9% (1.7-12%)	6.9% (2-14%)	170 (50-350)	2400 (730-4900)
Burkina Faso	2016	Male	80 to 84	47% (32-61%)	53% (39-68%)	2 (0.84-3.6)	4.6% (1.2-10%)	5.3% (1.3-12%)	70 (17-150)	730 (190-1600)
Burkina Faso	2016	Male	85 to 89	43% (30-56%)	57% (44-70%)	1.5 (0.66-2.9)	3.2% (0.73-7.4%)	3.7% (0.75-8.6%)	31 (6.3-72)	230 (53-540)
Burkina Faso	2016	Male	90 to 94	41% (29-55%)	59% (45-71%)	1.2 (0.49-2.2)	1.9% (0.16-4.9%)	2.1% (-0.022-5.7%)	8.1 (-0.082-22)	44 (3.6-110)
Burkina Faso	2016	Male	95 plus	43% (27-61%)	57% (39-73%)	0.92 (0.34-2)	1.6% (-0.14-4.9%)	1.7% (-0.44-5.6%)	1.9 (-0.48-6.2)	7.1 (-0.58-21)
Cameroon	1990	Female	All Ages	14% (11-19%)	86% (81-89%)	1.3 (0.94-1.6)	1.7% (1.4-2.2%)	3.8% (2.7-5.1%)	2400 (1700-3300)	74000 (57000-94000)
Cameroon	1990	Female	15 to 19	13% (6.4-24%)	87% (76-94%)	1.1 (0.4-2.2)	1.7% (0.89-2.8%)	2.1% (0.87-3.9%)	22 (9.2-42)	2300 (1200-3900)
Cameroon	1990	Female	20 to 24	17% (7.6-30%)	83% (70-92%)	1.7 (0.67-3.3)	2.6% (1.6-4%)	2.7% (1.3-4.8%)	35 (16-63)	3600 (2200-5800)
Cameroon	1990	Female	25 to 29	19% (8.3-33%)	81% (67-92%)	2.2 (0.89-4)	3.2% (2-4.9%)	3.8% (2-6.2%)	48 (25-78)	4000 (2500-5900)
Cameroon	1990	Female	30 to 34	21% (9.4-38%)	79% (62-91%)	2.5 (1.1-4.6)	3.7% (2.4-5.3%)	4.7% (2.7-7.4%)	57 (33-89)	4100 (2600-6000)
Cameroon	1990	Female	35 to 39	24% (11-41%)	76% (59-89%)	2.7 (1.1-4.9)	4.7% (3-6.9%)	6.5% (3.9-9.7%)	84 (51-130)	4900 (3100-7100)
Cameroon	1990	Female	40 to 44	25% (13-43%)	75% (57-87%)	2.9 (1.2-5.3)	5.6% (3.7-8.1%)	8% (5.2-12%)	110 (70-160)	5500 (3600-7900)
Cameroon	1990	Female	45 to 49	26% (12-43%)	74% (57-88%)	3 (1.1-5.7)	6.7% (4.2-9.3%)	9.5% (6-13%)	140 (90-210)	6300 (4000-9000)
Cameroon	1990	Female	50 to 54	27% (13-46%)	73% (54-87%)	3.2 (1.3-6.1)	7.7% (4.9-11%)	11% (6.7-15%)	190 (120-280)	7300 (4600-11000)
Cameroon	1990	Female	55 to 59	27% (14-46%)	73% (54-86%)	3.4 (1.4-6.9)	8% (5-12%)	11% (6.6-15%)	240 (150-350)	7700 (4800-11000)
Cameroon	1990	Female	60 to 64	28% (14-48%)	72% (52-86%)	3.6 (1.3-7.2)	7.6% (4.9-11%)	9.7% (6.2-14%)	270 (170-410)	7500 (4600-11000)
Cameroon	1990	Female	65 to 69	29% (16-48%)	71% (52-84%)	3.3 (1.2-6.6)	8.3% (5-12%)	10% (6.1-15%)	350 (200-520)	7800 (4500-12000)
Cameroon	1990	Female	70 to 74	27% (14-44%)	73% (56-86%)	2.6 (0.92-5.9)	7.7% (4.1-12%)	9.3% (4.8-15%)	350 (190-560)	6400 (3400-10000)
Cameroon	1990	Female	75 to 79	24% (12-41%)	76% (59-88%)	2.1 (0.64-4.8)	6.1% (2.6-10%)	7.2% (3.1-12%)	250 (110-420)	3500 (1500-5900)
Cameroon	1990	Female	80 to 84	22% (9.8-39%)	78% (61-90%)	1.9 (0.55-4)	5.3% (2-9.2%)	6.3% (2.4-11%)	160 (61-280)	1700 (650-2900)
Cameroon	1990	Female	85 to 89	23% (11-39%)	77% (61-89%)	1.9 (0.57-4.4)	4.7% (1.7-8.6%)	5.6% (2-10%)	88 (32-160)	650 (240-1200)
Cameroon	1990	Female	90 to 94	25% (12-43%)	75% (57-88%)	2 (0.55-4.9)	4.4% (1.3-8.4%)	5.2% (1.5-10%)	31 (8.9-60)	160 (47-310)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	1990	Female	95 plus	30% (13-53%)	70% (47-87%)	2.3 (0.54-6.1)	4.3% (0.94-8.6%)	5% (0.97-10%)	7.8 (1.5-16)	27 (5.7-54)
Cameroon	1990	Male	All Ages	11% (8.3-15%)	89% (85-92%)	1.4 (1.1-1.9)	2.1% (1.7-2.6%)	4.1% (3.2-5.1%)	2800 (2100-3500)	100000 (80000-120000)
Cameroon	1990	Male	15 to 19	8.5% (4-16%)	91% (84-96%)	0.82 (0.29-1.8)	1.9% (1-3.2%)	2.4% (1.1-4.4%)	25 (11-46)	2400 (1300-4100)
Cameroon	1990	Male	20 to 24	13% (6-24%)	87% (76-94%)	1.7 (0.61-3.3)	4.2% (2.5-6.3%)	4.8% (2.6-7.8%)	70 (36-110)	5900 (3500-8900)
Cameroon	1990	Male	25 to 29	17% (8-30%)	83% (70-92%)	2.6 (1-4.8)	5.8% (3.6-8.9%)	7% (4-11%)	110 (62-170)	7900 (5000-12000)
Cameroon	1990	Male	30 to 34	20% (8.5-36%)	80% (64-91%)	3.2 (1.4-5.9)	7.1% (4.6-10%)	8.9% (5.2-14%)	130 (81-210)	8600 (5600-13000)
Cameroon	1990	Male	35 to 39	22% (10-40%)	78% (60-90%)	3.7 (1.5-6.9)	8.5% (5.6-12%)	11% (7-16%)	170 (120-250)	9600 (6600-13000)
Cameroon	1990	Male	40 to 44	23% (12-40%)	77% (60-88%)	3.9 (1.6-7.6)	9.6% (6.6-13%)	13% (8.6-17%)	210 (150-290)	10000 (7200-14000)
Cameroon	1990	Male	45 to 49	24% (11-39%)	76% (61-89%)	4.2 (1.5-8.5)	11% (7.6-15%)	14% (9.9-20%)	260 (180-370)	11000 (7700-16000)
Cameroon	1990	Male	50 to 54	24% (11-42%)	76% (58-89%)	4.2 (1.6-8.7)	10% (6.9-14%)	13% (9-18%)	270 (180-380)	10000 (6800-14000)
Cameroon	1990	Male	55 to 59	24% (12-42%)	76% (58-88%)	4.1 (1.5-8.3)	10% (6.9-14%)	13% (8.8-17%)	290 (200-410)	9600 (6500-13000)
Cameroon	1990	Male	60 to 64	24% (12-41%)	76% (59-88%)	3.7 (1.3-7.8)	8.7% (5.9-12%)	11% (7.4-15%)	300 (200-410)	8100 (5400-11000)
Cameroon	1990	Male	65 to 69	24% (12-40%)	76% (60-88%)	3.1 (1.1-6.4)	8% (5.1-11%)	9.8% (6.3-14%)	290 (180-420)	6700 (4200-9600)
Cameroon	1990	Male	70 to 74	22% (12-38%)	78% (62-88%)	2.3 (0.79-5.3)	7.1% (4-11%)	8.5% (4.9-13%)	270 (150-420)	5000 (2800-7700)
Cameroon	1990	Male	75 to 79	20% (9.7-35%)	80% (65-90%)	1.7 (0.51-4)	5.4% (2.5-9.1%)	6.4% (3-11%)	180 (85-300)	2600 (1200-4200)
Cameroon	1990	Male	80 to 84	18% (8.2-33%)	82% (67-92%)	1.4 (0.41-3.2)	4.7% (2.1-8%)	5.6% (2.5-9.4%)	100 (46-170)	1100 (490-1800)
Cameroon	1990	Male	85 to 89	17% (8.1-32%)	83% (68-92%)	1.2 (0.37-2.9)	4% (1.6-7%)	4.7% (1.8-8.3%)	53 (21-93)	390 (160-690)
Cameroon	1990	Male	90 to 94	18% (8.4-34%)	82% (66-92%)	1.1 (0.3-2.9)	3.1% (1-6.1%)	3.6% (1.1-7.2%)	16 (5-32)	83 (28-160)
Cameroon	1990	Male	95 plus	22% (8.6-42%)	78% (58-91%)	1.2 (0.23-3.5)	2.8% (0.69-6.2%)	3.1% (0.67-7.3%)	3.6 (0.76-8)	13 (3.1-28)
Cameroon	1995	Female	All Ages	15% (11-19%)	85% (81-89%)	1.1 (0.8-1.4)	1.6% (1.2-2.1%)	3.5% (2.3-4.9%)	2700 (1700-3800)	83000 (61000-110000)
Cameroon	1995	Female	15 to 19	14% (6.6-24%)	86% (76-93%)	0.92 (0.32-1.9)	1.6% (0.82-2.8%)	2% (0.74-3.8%)	26 (9.7-50)	2700 (1400-4700)
Cameroon	1995	Female	20 to 24	17% (7.9-31%)	83% (69-92%)	1.5 (0.56-2.8)	2.3% (1.3-3.6%)	2.3% (0.99-4%)	42 (18-73)	4400 (2500-6700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	1995	Female	25 to 29	20% (8.4-34%)	80% (66-92%)	1.8 (0.74-3.4)	2.7% (1.6-4.1%)	2.8% (1.3-5%)	58 (30-97)	4900 (3000-7400)
Cameroon	1995	Female	30 to 34	22% (9.7-39%)	78% (61-90%)	2.1 (0.93-3.9)	3% (1.9-4.4%)	3.4% (1.8-5.4%)	68 (37-110)	4900 (3000-7000)
Cameroon	1995	Female	35 to 39	25% (12-42%)	75% (58-88%)	2.3 (0.95-4.2)	4% (2.5-6.1%)	5% (2.9-8%)	96 (57-150)	5700 (3600-8200)
Cameroon	1995	Female	40 to 44	26% (14-43%)	74% (57-86%)	2.5 (1-4.5)	4.9% (3.1-7.1%)	6.5% (3.9-9.8%)	120 (75-180)	6200 (3900-8900)
Cameroon	1995	Female	45 to 49	27% (12-44%)	73% (56-88%)	2.5 (0.92-4.9)	6% (3.7-8.7%)	8% (4.7-12%)	160 (94-240)	7000 (4300-10000)
Cameroon	1995	Female	50 to 54	27% (13-47%)	73% (53-87%)	2.6 (1-5.2)	7% (4.3-10%)	9.3% (5.6-14%)	210 (130-320)	8100 (4800-12000)
Cameroon	1995	Female	55 to 59	28% (14-48%)	72% (52-86%)	2.9 (1.1-5.8)	7.6% (4.5-11%)	9.8% (5.7-15%)	260 (150-410)	8500 (4900-13000)
Cameroon	1995	Female	60 to 64	30% (15-49%)	70% (51-85%)	3 (1.1-6.4)	7.3% (4.2-11%)	9.1% (5.3-14%)	300 (170-460)	8200 (4700-13000)
Cameroon	1995	Female	65 to 69	30% (16-49%)	70% (51-84%)	2.8 (1-5.7)	7.9% (4.4-12%)	9.7% (5.3-15%)	380 (210-600)	8600 (4700-13000)
Cameroon	1995	Female	70 to 74	28% (15-45%)	72% (55-85%)	2.2 (0.78-5)	7.2% (3.3-12%)	8.6% (3.9-14%)	380 (170-650)	6900 (3100-12000)
Cameroon	1995	Female	75 to 79	24% (12-42%)	76% (58-88%)	1.7 (0.5-4)	5.6% (2.1-9.7%)	6.6% (2.5-12%)	270 (98-480)	3800 (1400-6700)
Cameroon	1995	Female	80 to 84	22% (11-39%)	78% (61-89%)	1.5 (0.46-3.3)	4.9% (1.7-9.2%)	5.7% (1.9-11%)	170 (59-330)	1800 (620-3400)
Cameroon	1995	Female	85 to 89	23% (11-39%)	77% (61-89%)	1.5 (0.47-3.5)	4.3% (1.3-8.3%)	5.1% (1.5-9.9%)	96 (29-190)	710 (220-1400)
Cameroon	1995	Female	90 to 94	26% (12-44%)	74% (56-88%)	1.7 (0.46-3.9)	4% (1-8.2%)	4.8% (1.2-10%)	36 (8.8-76)	180 (46-390)
Cameroon	1995	Female	95 plus	31% (13-54%)	69% (46-87%)	1.9 (0.44-4.8)	4% (0.58-8.6%)	4.6% (0.58-10%)	9.3 (1.2-20)	32 (4.6-69)
Cameroon	1995	Male	All Ages	12% (8.8-16%)	88% (84-91%)	1.3 (0.95-1.6)	2.2% (1.7-2.7%)	4.1% (3.1-5.2%)	3400 (2600-4400)	120000 (98000-150000)
Cameroon	1995	Male	15 to 19	9.2% (4.4-17%)	91% (83-96%)	0.73 (0.25-1.5)	2% (1-3.4%)	2.5% (1.1-4.6%)	34 (14-64)	3200 (1600-5500)
Cameroon	1995	Male	20 to 24	14% (6.3-25%)	86% (75-94%)	1.5 (0.56-3)	4.2% (2.4-6.4%)	4.8% (2.4-7.8%)	97 (48-160)	7900 (4500-12000)
Cameroon	1995	Male	25 to 29	18% (8.4-32%)	82% (68-92%)	2.3 (0.91-4.3)	5.4% (3.1-8.4%)	6.1% (3.2-10%)	150 (84-240)	11000 (6500-16000)
Cameroon	1995	Male	30 to 34	21% (9.6-37%)	79% (63-90%)	2.9 (1.3-5.3)	6.2% (3.8-9.1%)	7.2% (4.1-11%)	190 (110-280)	12000 (7300-17000)
Cameroon	1995	Male	35 to 39	24% (11-41%)	76% (59-89%)	3.2 (1.4-6)	7.3% (4.7-11%)	8.9% (5.4-13%)	220 (140-320)	12000 (8000-17000)
Cameroon	1995	Male	40 to 44	24% (12-41%)	76% (59-88%)	3.5 (1.4-6.4)	8.5% (5.7-12%)	10% (6.8-15%)	260 (170-370)	13000 (8500-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	1995	Male	45 to 49	25% (11-41%)	75% (59-89%)	3.6 (1.3-7.1)	10% (6.6-14%)	12% (8.2-18%)	320 (210-460)	14000 (9100-19000)
Cameroon	1995	Male	50 to 54	25% (12-45%)	75% (55-88%)	3.7 (1.4-7.6)	9.9% (6.5-14%)	12% (8.1-17%)	330 (220-480)	12000 (8200-18000)
Cameroon	1995	Male	55 to 59	25% (13-45%)	75% (55-87%)	3.6 (1.3-7.2)	9.9% (6.6-14%)	12% (8.3-17%)	360 (230-510)	12000 (7600-16000)
Cameroon	1995	Male	60 to 64	25% (13-44%)	75% (56-87%)	3.2 (1.1-7.1)	8.5% (5.6-12%)	10% (6.8-15%)	350 (220-510)	9600 (6100-14000)
Cameroon	1995	Male	65 to 69	25% (13-42%)	75% (58-87%)	2.7 (1-5.6)	7.8% (4.8-11%)	9.4% (5.7-14%)	350 (210-520)	7900 (4700-12000)
Cameroon	1995	Male	70 to 74	24% (12-39%)	76% (61-88%)	2 (0.68-4.6)	6.8% (3.6-11%)	8.1% (4.3-13%)	320 (160-510)	5800 (3000-9200)
Cameroon	1995	Male	75 to 79	21% (10-38%)	79% (62-90%)	1.5 (0.46-3.5)	5.2% (2.2-9%)	6.1% (2.6-11%)	210 (89-370)	3000 (1200-5200)
Cameroon	1995	Male	80 to 84	20% (8.6-35%)	80% (65-91%)	1.3 (0.37-2.9)	4.6% (1.8-8.1%)	5.3% (2.1-9.4%)	120 (48-210)	1300 (510-2200)
Cameroon	1995	Male	85 to 89	19% (8.4-34%)	81% (66-92%)	1.1 (0.31-2.5)	3.8% (1.4-7.1%)	4.5% (1.5-8.3%)	60 (20-110)	450 (160-820)
Cameroon	1995	Male	90 to 94	20% (8.8-36%)	80% (64-91%)	1 (0.27-2.6)	3% (0.88-6.3%)	3.4% (0.97-7.4%)	19 (5.2-40)	96 (29-210)
Cameroon	1995	Male	95 plus	23% (9.6-45%)	77% (55-90%)	1 (0.2-3)	2.8% (0.63-6.7%)	3.2% (0.59-7.8%)	4.4 (0.82-11)	16 (3.5-37)
Cameroon	2000	Female	All Ages	15% (11-20%)	85% (80-89%)	1.1 (0.82-1.4)	1.8% (1.3-2.4%)	3.7% (2.4-5.3%)	3500 (2100-5100)	110000 (74000-140000)
Cameroon	2000	Female	15 to 19	14% (7-25%)	86% (75-93%)	0.95 (0.35-2)	1.7% (0.86-3%)	2.1% (0.78-4.1%)	34 (13-64)	3400 (1700-5900)
Cameroon	2000	Female	20 to 24	18% (8.3-32%)	82% (68-92%)	1.5 (0.59-2.9)	2.3% (1.3-3.6%)	2.2% (0.94-4.1%)	54 (23-98)	5500 (3000-8500)
Cameroon	2000	Female	25 to 29	20% (9.6-36%)	80% (64-90%)	1.9 (0.79-3.5)	2.3% (1.2-3.6%)	2.3% (1-4%)	76 (38-120)	6300 (3800-9400)
Cameroon	2000	Female	30 to 34	23% (10-41%)	77% (59-90%)	2.2 (0.96-4)	2.5% (1.4-3.8%)	2.6% (1.3-4.2%)	89 (48-140)	6300 (3800-9400)
Cameroon	2000	Female	35 to 39	26% (12-44%)	74% (56-88%)	2.4 (1-4.4)	3.4% (2-5.3%)	3.9% (2-6.4%)	130 (72-200)	7500 (4500-11000)
Cameroon	2000	Female	40 to 44	27% (14-46%)	73% (54-86%)	2.6 (1-4.9)	4.3% (2.7-6.5%)	5.2% (3.2-8.2%)	150 (92-240)	7800 (4900-12000)
Cameroon	2000	Female	45 to 49	28% (13-46%)	72% (54-87%)	2.7 (0.94-5.2)	5.6% (3.3-8.4%)	7.1% (4-11%)	200 (120-320)	8900 (5200-14000)
Cameroon	2000	Female	50 to 54	28% (13-49%)	72% (51-87%)	2.8 (1.1-5.4)	7% (4.1-10%)	8.8% (5.3-13%)	270 (150-430)	10000 (5700-16000)
Cameroon	2000	Female	55 to 59	29% (14-49%)	71% (51-86%)	3 (1.2-6.2)	7.9% (4.5-12%)	9.8% (5.6-15%)	340 (180-540)	11000 (5900-17000)
Cameroon	2000	Female	60 to 64	30% (15-50%)	70% (50-85%)	3.1 (1.1-6.4)	7.6% (4.5-12%)	9.2% (5.4-14%)	380 (210-600)	10000 (5600-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	2000	Female	65 to 69	31% (16-49%)	69% (51-84%)	2.9 (1.1-5.7)	8.4% (4.6-13%)	10% (5.5-15%)	490 (250-780)	11000 (5800-18000)
Cameroon	2000	Female	70 to 74	29% (16-47%)	71% (53-84%)	2.3 (0.82-5.2)	7.6% (3.4-13%)	9% (3.9-15%)	480 (200-840)	8600 (3700-15000)
Cameroon	2000	Female	75 to 79	25% (12-43%)	75% (57-88%)	1.8 (0.53-4.1)	6% (2.3-11%)	7.1% (2.6-13%)	350 (130-650)	4900 (1800-9000)
Cameroon	2000	Female	80 to 84	23% (11-40%)	77% (60-89%)	1.6 (0.45-3.4)	5.4% (1.9-10%)	6.3% (2.2-12%)	230 (76-450)	2400 (790-4600)
Cameroon	2000	Female	85 to 89	23% (11-40%)	77% (60-89%)	1.6 (0.48-3.6)	4.8% (1.4-9.3%)	5.6% (1.6-11%)	130 (36-250)	920 (280-1800)
Cameroon	2000	Female	90 to 94	26% (12-45%)	74% (55-88%)	1.7 (0.46-4.1)	4.5% (1.1-9%)	5.4% (1.2-11%)	49 (11-100)	250 (60-510)
Cameroon	2000	Female	95 plus	31% (13-55%)	69% (45-87%)	1.9 (0.45-5.1)	4.5% (0.73-9.6%)	5.2% (0.72-12%)	14 (1.8-30)	46 (7.5-100)
Cameroon	2000	Male	All Ages	12% (9.3-17%)	88% (83-91%)	1.3 (1-1.7)	2.5% (1.9-3.1%)	4.5% (3.4-5.7%)	4400 (3300-5700)	160000 (130000-200000)
Cameroon	2000	Male	15 to 19	9.7% (4.7-18%)	90% (82-95%)	0.79 (0.29-1.7)	2.2% (1.1-3.7%)	2.7% (1.1-4.9%)	48 (20-87)	4400 (2200-7300)
Cameroon	2000	Male	20 to 24	15% (6.7-26%)	85% (74-93%)	1.6 (0.6-3.1)	4.5% (2.6-7%)	5.1% (2.5-8.2%)	130 (66-220)	10000 (5800-16000)
Cameroon	2000	Male	25 to 29	19% (8.9-33%)	81% (67-91%)	2.4 (1-4.5)	5.2% (2.9-8.1%)	5.8% (2.9-9.4%)	200 (120-330)	14000 (8700-22000)
Cameroon	2000	Male	30 to 34	22% (10-39%)	78% (61-90%)	3 (1.3-5.6)	5.8% (3.5-8.7%)	6.5% (3.7-10%)	260 (150-400)	16000 (9900-24000)
Cameroon	2000	Male	35 to 39	25% (12-42%)	75% (58-88%)	3.4 (1.4-6.4)	6.7% (4.3-9.8%)	7.7% (4.7-12%)	300 (190-440)	16000 (11000-23000)
Cameroon	2000	Male	40 to 44	25% (13-44%)	75% (56-87%)	3.7 (1.5-6.9)	7.8% (5.2-11%)	9.1% (5.9-13%)	340 (220-490)	16000 (11000-23000)
Cameroon	2000	Male	45 to 49	26% (12-43%)	74% (57-88%)	3.9 (1.4-7.6)	9.5% (6.3-13%)	11% (7.5-16%)	420 (270-590)	18000 (11000-25000)
Cameroon	2000	Male	50 to 54	27% (13-46%)	73% (54-87%)	3.9 (1.5-7.9)	9.9% (6.4-14%)	12% (7.7-16%)	430 (280-610)	16000 (10000-23000)
Cameroon	2000	Male	55 to 59	27% (14-45%)	73% (55-86%)	3.8 (1.4-7.7)	10% (6.8-14%)	12% (8.2-17%)	470 (310-670)	15000 (10000-21000)
Cameroon	2000	Male	60 to 64	27% (13-45%)	73% (55-87%)	3.5 (1.2-7.4)	8.9% (5.8-13%)	11% (6.9-15%)	450 (290-660)	12000 (7900-18000)
Cameroon	2000	Male	65 to 69	26% (13-43%)	74% (57-87%)	2.9 (1-6)	8.3% (4.9-12%)	9.8% (5.8-15%)	440 (260-670)	10000 (5900-15000)
Cameroon	2000	Male	70 to 74	25% (13-41%)	75% (59-87%)	2.2 (0.74-4.8)	7.3% (3.6-12%)	8.5% (4.4-14%)	400 (200-660)	7300 (3700-12000)
Cameroon	2000	Male	75 to 79	23% (11-39%)	77% (61-89%)	1.7 (0.51-3.8)	5.6% (2.2-9.7%)	6.5% (2.5-11%)	270 (110-480)	3800 (1500-6700)
Cameroon	2000	Male	80 to 84	21% (9.9-36%)	79% (64-90%)	1.4 (0.4-3)	5% (1.9-8.9%)	5.8% (2.2-10%)	160 (61-280)	1600 (640-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	2000	Male	85 to 89	20% (9.2-35%)	80% (65-91%)	1.2 (0.33-2.7)	4.2% (1.5-7.8%)	4.9% (1.7-9.2%)	78 (26-150)	580 (200-1100)
Cameroon	2000	Male	90 to 94	21% (9.9-37%)	79% (63-90%)	1.1 (0.29-2.8)	3.4% (0.94-7.1%)	3.9% (1-8.3%)	24 (6.4-52)	130 (35-260)
Cameroon	2000	Male	95 plus	25% (10-47%)	75% (53-90%)	1.2 (0.23-3.2)	3.3% (0.78-7.6%)	3.8% (0.75-9%)	6.2 (1.2-15)	22 (4.9-50)
Cameroon	2005	Female	All Ages	15% (11-20%)	85% (80-89%)	1.3 (0.96-1.7)	1.8% (1.3-2.4%)	3.8% (2.5-5.3%)	3900 (2400-5600)	120000 (83000-160000)
Cameroon	2005	Female	15 to 19	14% (7-25%)	86% (75-93%)	1.1 (0.42-2.3)	1.8% (0.92-3%)	2.2% (0.89-4.1%)	39 (16-77)	4000 (2100-6600)
Cameroon	2005	Female	20 to 24	18% (8.7-32%)	82% (68-91%)	1.8 (0.7-3.4)	2.4% (1.4-3.6%)	2.3% (1.1-4%)	63 (29-110)	6300 (3800-9700)
Cameroon	2005	Female	25 to 29	21% (9.7-36%)	79% (64-90%)	2.2 (0.93-4.1)	2.2% (1.2-3.4%)	2.1% (0.93-3.8%)	83 (44-130)	7000 (4300-10000)
Cameroon	2005	Female	30 to 34	23% (10-42%)	77% (58-90%)	2.5 (1.1-4.6)	2.2% (1.3-3.3%)	2.2% (1.1-3.6%)	100 (53-150)	7100 (4300-10000)
Cameroon	2005	Female	35 to 39	26% (12-44%)	74% (56-88%)	2.8 (1.2-5)	3% (1.8-4.5%)	3.3% (1.8-5.3%)	140 (80-220)	8400 (5200-12000)
Cameroon	2005	Female	40 to 44	27% (14-45%)	73% (55-86%)	3 (1.2-5.7)	3.8% (2.4-5.7%)	4.4% (2.6-6.7%)	170 (100-270)	8700 (5500-13000)
Cameroon	2005	Female	45 to 49	28% (13-45%)	72% (55-87%)	3.1 (1.1-6)	5.2% (3.1-7.7%)	6.4% (3.7-9.7%)	220 (120-340)	9700 (5600-15000)
Cameroon	2005	Female	50 to 54	29% (13-48%)	71% (52-87%)	3.3 (1.2-6.5)	6.8% (4.1-10%)	8.5% (5.1-13%)	300 (160-470)	11000 (6300-17000)
Cameroon	2005	Female	55 to 59	29% (15-48%)	71% (52-85%)	3.6 (1.3-7.2)	7.9% (4.7-12%)	9.8% (6-15%)	370 (200-590)	12000 (6600-19000)
Cameroon	2005	Female	60 to 64	30% (15-51%)	70% (49-85%)	3.6 (1.3-7.7)	7.7% (4.7-12%)	9.4% (5.8-14%)	410 (230-660)	11000 (6400-18000)
Cameroon	2005	Female	65 to 69	31% (16-50%)	69% (50-84%)	3.3 (1.3-6.5)	8.8% (5.1-13%)	11% (6.1-16%)	540 (290-860)	12000 (6600-19000)
Cameroon	2005	Female	70 to 74	29% (16-47%)	71% (53-84%)	2.6 (0.97-5.8)	8% (3.9-13%)	9.5% (4.6-15%)	540 (250-930)	9800 (4600-17000)
Cameroon	2005	Female	75 to 79	25% (12-44%)	75% (56-88%)	2.1 (0.59-4.8)	6.5% (2.6-11%)	7.6% (3-13%)	410 (160-730)	5700 (2300-10000)
Cameroon	2005	Female	80 to 84	23% (11-40%)	77% (60-89%)	1.8 (0.53-4)	5.8% (2.1-11%)	6.9% (2.5-12%)	280 (100-520)	2900 (1100-5300)
Cameroon	2005	Female	85 to 89	24% (12-41%)	76% (59-88%)	1.8 (0.56-4.3)	5.3% (1.8-10%)	6.2% (2-12%)	150 (50-290)	1100 (370-2100)
Cameroon	2005	Female	90 to 94	27% (13-46%)	73% (54-87%)	2 (0.53-4.9)	5% (1.4-9.7%)	6% (1.5-12%)	58 (15-110)	300 (82-580)
Cameroon	2005	Female	95 plus	32% (14-54%)	68% (46-86%)	2.3 (0.55-6.1)	5% (0.91-10%)	5.9% (0.95-12%)	17 (2.8-36)	57 (10-120)
Cameroon	2005	Male	All Ages	13% (9.3-17%)	87% (83-91%)	1.6 (1.2-2.1)	2.6% (2.1-3.3%)	4.7% (3.7-5.9%)	5100 (4000-6500)	190000 (150000-240000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	2005	Male	15 to 19	9.8% (4.8-18%)	90% (82-95%)	0.91 (0.34-1.9)	2.3% (1.2-3.8%)	2.8% (1.3-5%)	59 (26-100)	5300 (2800-8700)
Cameroon	2005	Male	20 to 24	15% (7-27%)	85% (73-93%)	1.9 (0.71-3.6)	4.8% (2.8-7.3%)	5.5% (2.8-8.8%)	160 (83-270)	13000 (7400-20000)
Cameroon	2005	Male	25 to 29	19% (9-34%)	81% (66-91%)	2.8 (1.2-5.3)	5.6% (3.3-8.5%)	6.2% (3.3-10%)	240 (140-390)	17000 (11000-26000)
Cameroon	2005	Male	30 to 34	22% (10-39%)	78% (61-90%)	3.5 (1.5-6.5)	5.7% (3.5-8.6%)	6.3% (3.7-9.6%)	300 (180-460)	19000 (12000-28000)
Cameroon	2005	Male	35 to 39	24% (12-41%)	76% (59-88%)	3.9 (1.6-7.6)	6.4% (4.3-9.3%)	7.3% (4.6-11%)	350 (230-510)	19000 (13000-27000)
Cameroon	2005	Male	40 to 44	25% (13-44%)	75% (56-87%)	4.3 (1.7-8.4)	7.3% (5.1-10%)	8.4% (5.7-12%)	400 (270-560)	19000 (13000-26000)
Cameroon	2005	Male	45 to 49	26% (12-43%)	74% (57-88%)	4.6 (1.7-9)	9.2% (6.2-13%)	11% (7.2-15%)	470 (320-660)	20000 (13000-28000)
Cameroon	2005	Male	50 to 54	27% (13-46%)	73% (54-87%)	4.7 (1.7-9.4)	9.9% (6.4-13%)	12% (7.8-16%)	480 (310-660)	18000 (12000-24000)
Cameroon	2005	Male	55 to 59	27% (13-46%)	73% (54-87%)	4.5 (1.6-9.1)	11% (7.3-14%)	13% (8.8-17%)	530 (350-740)	17000 (11000-23000)
Cameroon	2005	Male	60 to 64	27% (13-45%)	73% (55-87%)	4 (1.4-8.9)	9.2% (6-13%)	11% (7.2-16%)	510 (320-740)	14000 (9000-20000)
Cameroon	2005	Male	65 to 69	26% (13-44%)	74% (56-87%)	3.3 (1.2-7.1)	8.5% (5.2-12%)	10% (6.2-15%)	500 (300-730)	11000 (6900-17000)
Cameroon	2005	Male	70 to 74	25% (13-41%)	75% (59-87%)	2.5 (0.9-5.5)	7.6% (4.2-12%)	8.9% (5-13%)	470 (260-710)	8500 (4700-13000)
Cameroon	2005	Male	75 to 79	22% (11-40%)	78% (60-89%)	1.9 (0.56-4.3)	5.9% (2.5-10%)	6.9% (2.9-12%)	320 (130-540)	4500 (1900-7600)
Cameroon	2005	Male	80 to 84	20% (9.4-36%)	80% (64-91%)	1.5 (0.45-3.5)	5.3% (2.1-9.1%)	6.1% (2.5-11%)	190 (77-330)	2000 (810-3400)
Cameroon	2005	Male	85 to 89	19% (9.4-34%)	81% (66-91%)	1.3 (0.4-3)	4.5% (1.7-8.1%)	5.3% (1.9-9.4%)	94 (34-170)	700 (250-1200)
Cameroon	2005	Male	90 to 94	21% (10-36%)	79% (64-90%)	1.3 (0.35-3.2)	3.7% (1.1-7.2%)	4.3% (1.2-8.5%)	30 (8.5-59)	150 (45-300)
Cameroon	2005	Male	95 plus	25% (10-46%)	75% (54-90%)	1.3 (0.27-3.9)	3.6% (0.84-7.9%)	4.1% (0.82-9.4%)	7.4 (1.5-17)	26 (5.9-57)
Cameroon	2010	Female	All Ages	16% (11-21%)	84% (79-89%)	1.4 (0.96-2)	1.8% (1.3-2.5%)	3.8% (2.4-5.3%)	4000 (2400-5900)	120000 (85000-170000)
Cameroon	2010	Female	15 to 19	14% (6.9-25%)	86% (75-93%)	1.2 (0.45-2.6)	1.7% (0.91-2.9%)	2% (0.81-3.7%)	40 (16-77)	4200 (2200-7400)
Cameroon	2010	Female	20 to 24	18% (8.6-33%)	82% (67-91%)	1.9 (0.71-3.6)	2.6% (1.5-3.8%)	2.5% (1.1-4.4%)	66 (29-110)	7000 (4100-11000)
Cameroon	2010	Female	25 to 29	21% (9.1-37%)	79% (63-91%)	2.3 (0.94-4.4)	2.4% (1.4-3.7%)	2.3% (1.1-4.1%)	89 (46-150)	7800 (4800-12000)
Cameroon	2010	Female	30 to 34	23% (11-41%)	77% (59-89%)	2.7 (1.1-5.1)	2.3% (1.3-3.5%)	2.2% (1.1-3.8%)	100 (55-170)	7700 (4700-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	2010	Female	35 to 39	26% (12-44%)	74% (56-88%)	3 (1.2-5.7)	3% (1.8-4.5%)	3.2% (1.7-5.1%)	150 (82-230)	8900 (5500-13000)
Cameroon	2010	Female	40 to 44	27% (14-46%)	73% (54-86%)	3.2 (1.2-6.1)	3.8% (2.4-5.6%)	4.4% (2.6-6.8%)	180 (110-280)	9300 (5800-14000)
Cameroon	2010	Female	45 to 49	28% (13-45%)	72% (55-87%)	3.3 (1.1-6.4)	5.2% (3.1-7.6%)	6.4% (3.6-9.7%)	230 (130-370)	10000 (6100-16000)
Cameroon	2010	Female	50 to 54	28% (14-48%)	72% (52-86%)	3.4 (1.3-6.8)	6.8% (4.1-10%)	8.6% (5.1-13%)	300 (160-490)	11000 (6300-18000)
Cameroon	2010	Female	55 to 59	29% (15-49%)	71% (51-85%)	3.7 (1.3-7.5)	8% (4.6-12%)	10% (6-15%)	370 (190-600)	12000 (6400-19000)
Cameroon	2010	Female	60 to 64	30% (16-50%)	70% (50-84%)	3.8 (1.3-8.1)	7.9% (4.6-12%)	9.8% (5.8-15%)	420 (230-690)	12000 (6300-19000)
Cameroon	2010	Female	65 to 69	31% (16-50%)	69% (50-84%)	3.5 (1.2-7)	8.9% (5.1-13%)	11% (6.1-16%)	550 (280-890)	13000 (6500-20000)
Cameroon	2010	Female	70 to 74	29% (15-46%)	71% (54-85%)	2.8 (0.95-6.3)	8.1% (3.9-13%)	9.7% (4.6-16%)	550 (240-930)	10000 (4500-17000)
Cameroon	2010	Female	75 to 79	25% (13-43%)	75% (57-87%)	2.2 (0.63-5.1)	6.5% (2.6-11%)	7.7% (3-13%)	420 (170-770)	6000 (2400-11000)
Cameroon	2010	Female	80 to 84	23% (11-39%)	77% (61-89%)	1.9 (0.54-4.4)	5.9% (2.2-11%)	6.9% (2.5-13%)	300 (110-560)	3200 (1200-5800)
Cameroon	2010	Female	85 to 89	23% (11-40%)	77% (60-89%)	1.9 (0.59-4.3)	5.3% (1.8-10%)	6.3% (2-12%)	170 (55-330)	1200 (410-2400)
Cameroon	2010	Female	90 to 94	26% (12-45%)	74% (55-88%)	2 (0.55-5)	5.1% (1.4-9.8%)	6% (1.6-12%)	63 (17-120)	320 (91-630)
Cameroon	2010	Female	95 plus	31% (13-55%)	69% (45-87%)	2.3 (0.55-6.4)	5% (0.83-10%)	5.9% (0.88-12%)	18 (2.6-37)	60 (9.7-120)
Cameroon	2010	Male	All Ages	13% (9.5-17%)	87% (83-91%)	1.7 (1.2-2.4)	2.8% (2.2-3.5%)	5% (3.9-6.3%)	5600 (4300-7100)	210000 (160000-260000)
Cameroon	2010	Male	15 to 19	9.8% (4.7-18%)	90% (82-95%)	0.95 (0.34-2.1)	2.3% (1.2-3.8%)	2.7% (1.2-4.8%)	64 (28-110)	5800 (3100-9900)
Cameroon	2010	Male	20 to 24	15% (7-27%)	85% (73-93%)	1.9 (0.74-3.9)	5% (3.1-7.6%)	5.8% (3.1-9.3%)	180 (92-300)	15000 (8300-22000)
Cameroon	2010	Male	25 to 29	19% (9.2-34%)	81% (66-91%)	2.9 (1.2-5.6)	6.2% (3.7-9.1%)	7.1% (3.9-11%)	280 (160-440)	20000 (12000-30000)
Cameroon	2010	Male	30 to 34	22% (10-39%)	78% (61-90%)	3.7 (1.6-6.9)	6.5% (3.9-9.5%)	7.3% (4.2-11%)	340 (210-520)	21000 (14000-32000)
Cameroon	2010	Male	35 to 39	25% (11-43%)	75% (57-89%)	4.2 (1.6-8.4)	7% (4.7-10%)	8.1% (5.2-12%)	400 (260-580)	22000 (15000-31000)
Cameroon	2010	Male	40 to 44	26% (13-44%)	74% (56-87%)	4.5 (1.6-8.9)	7.9% (5.6-11%)	9.3% (6.5-13%)	450 (300-630)	22000 (15000-30000)
Cameroon	2010	Male	45 to 49	26% (13-43%)	74% (57-87%)	4.8 (1.6-9.7)	9.7% (6.5-13%)	12% (7.7-16%)	520 (340-730)	22000 (15000-31000)
Cameroon	2010	Male	50 to 54	27% (13-47%)	73% (53-87%)	4.8 (1.7-10)	10% (6.6-14%)	12% (8.1-17%)	510 (330-710)	19000 (12000-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	2010	Male	55 to 59	27% (14-45%)	73% (55-86%)	4.7 (1.6-9.2)	11% (7.4-15%)	13% (9.1-18%)	560 (370-780)	18000 (12000-25000)
Cameroon	2010	Male	60 to 64	27% (13-46%)	73% (54-87%)	4.2 (1.3-8.9)	9.6% (6.4-14%)	11% (7.6-16%)	550 (350-790)	15000 (9600-21000)
Cameroon	2010	Male	65 to 69	27% (13-44%)	73% (56-87%)	3.5 (1.2-7.2)	8.8% (5.4-13%)	10% (6.5-15%)	550 (330-790)	12000 (7600-18000)
Cameroon	2010	Male	70 to 74	25% (13-41%)	75% (59-87%)	2.6 (0.92-5.7)	7.8% (4.4-12%)	9.2% (5.1-14%)	510 (280-780)	9200 (5100-14000)
Cameroon	2010	Male	75 to 79	23% (11-39%)	77% (61-89%)	2 (0.56-4.7)	6% (2.6-10%)	7% (3-12%)	350 (140-590)	4900 (2000-8300)
Cameroon	2010	Male	80 to 84	21% (9.4-36%)	79% (64-91%)	1.6 (0.47-3.7)	5.3% (2.2-9.1%)	6.2% (2.5-11%)	220 (89-360)	2200 (940-3800)
Cameroon	2010	Male	85 to 89	20% (9.3-34%)	80% (66-91%)	1.4 (0.4-3.3)	4.6% (1.7-8.1%)	5.4% (2-9.5%)	110 (40-190)	810 (300-1400)
Cameroon	2010	Male	90 to 94	21% (10-37%)	79% (63-90%)	1.3 (0.34-3.3)	3.8% (1.1-7.4%)	4.4% (1.2-8.7%)	34 (9.1-67)	180 (50-340)
Cameroon	2010	Male	95 plus	25% (11-47%)	75% (53-89%)	1.4 (0.26-3.9)	3.7% (0.81-8.1%)	4.3% (0.83-9.6%)	8.4 (1.6-19)	29 (6.5-64)
Cameroon	2016	Female	All Ages	16% (11-21%)	84% (79-89%)	1.4 (0.79-2.4)	1.9% (1.3-2.8%)	4% (2.4-5.7%)	4100 (2300-6200)	130000 (82000-180000)
Cameroon	2016	Female	15 to 19	14% (6.9-25%)	86% (75-93%)	1.2 (0.38-2.8)	1.6% (0.8-2.7%)	1.7% (0.61-3.3%)	37 (13-75)	4200 (2000-7300)
Cameroon	2016	Female	20 to 24	18% (8.4-32%)	82% (68-92%)	1.9 (0.65-4.2)	2.5% (1.6-3.8%)	2.3% (1.1-4.2%)	61 (24-110)	7100 (4100-11000)
Cameroon	2016	Female	25 to 29	21% (9.5-36%)	79% (64-91%)	2.4 (0.87-5.1)	2.5% (1.5-3.8%)	2.4% (1.1-4.3%)	86 (44-150)	8100 (5000-12000)
Cameroon	2016	Female	30 to 34	23% (10-41%)	77% (59-90%)	2.7 (1-5.6)	2.5% (1.5-3.8%)	2.5% (1.2-4.2%)	110 (55-180)	8400 (5200-13000)
Cameroon	2016	Female	35 to 39	26% (12-44%)	74% (56-88%)	3 (1.1-6.4)	3.1% (1.9-4.9%)	3.5% (1.9-5.7%)	160 (84-250)	9600 (5700-14000)
Cameroon	2016	Female	40 to 44	27% (14-45%)	73% (55-86%)	3.2 (1.1-6.9)	3.9% (2.4-5.9%)	4.7% (2.7-7.4%)	190 (110-320)	10000 (5900-15000)
Cameroon	2016	Female	45 to 49	28% (13-46%)	72% (54-87%)	3.3 (1-7.3)	5.3% (3.1-8%)	6.7% (3.9-10%)	250 (140-400)	11000 (6300-17000)
Cameroon	2016	Female	50 to 54	29% (13-48%)	71% (52-87%)	3.5 (1.1-7.3)	6.8% (4-10%)	8.9% (5.2-13%)	310 (160-530)	12000 (6500-20000)
Cameroon	2016	Female	55 to 59	29% (14-49%)	71% (51-86%)	3.8 (1.3-8.5)	7.8% (4.6-12%)	10% (6.1-15%)	370 (190-620)	12000 (6200-20000)
Cameroon	2016	Female	60 to 64	30% (15-50%)	70% (50-85%)	3.9 (1.2-8.7)	7.6% (4.2-12%)	9.6% (5.4-15%)	410 (210-700)	11000 (6000-19000)
Cameroon	2016	Female	65 to 69	31% (16-49%)	69% (51-84%)	3.5 (1.1-7.6)	8.4% (4.5-13%)	10% (5.5-16%)	540 (250-900)	12000 (5800-20000)
Cameroon	2016	Female	70 to 74	29% (15-46%)	71% (54-85%)	2.8 (0.86-6.8)	7.5% (3.3-12%)	9% (4-15%)	530 (230-930)	9800 (4200-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	2016	Female	75 to 79	25% (12-43%)	75% (57-88%)	2.2 (0.57-5.3)	6% (2.3-10%)	7.2% (2.7-12%)	420 (150-770)	6000 (2200-11000)
Cameroon	2016	Female	80 to 84	23% (11-40%)	77% (60-89%)	1.9 (0.52-4.7)	5.4% (1.9-9.8%)	6.4% (2.2-12%)	300 (99-580)	3200 (1000-6000)
Cameroon	2016	Female	85 to 89	23% (11-40%)	77% (60-89%)	1.9 (0.53-4.7)	5% (1.6-9.3%)	5.9% (1.8-11%)	180 (56-370)	1400 (410-2700)
Cameroon	2016	Female	90 to 94	27% (12-46%)	73% (54-88%)	2.2 (0.52-5.6)	4.9% (1.2-9.5%)	5.8% (1.3-12%)	74 (17-150)	380 (90-740)
Cameroon	2016	Female	95 plus	32% (14-56%)	68% (44-86%)	2.5 (0.46-7)	5% (0.66-10%)	5.9% (0.64-12%)	21 (2.3-43)	70 (9.6-140)
Cameroon	2016	Male	All Ages	13% (9.6-18%)	87% (82-90%)	1.7 (0.94-2.9)	3% (2.3-3.9%)	5.1% (3.8-6.6%)	5900 (4200-7700)	220000 (170000-290000)
Cameroon	2016	Male	15 to 19	9.8% (4.7-18%)	90% (82-95%)	0.98 (0.28-2.4)	2.1% (1.1-3.4%)	2.4% (0.94-4.3%)	64 (26-120)	6000 (3000-10000)
Cameroon	2016	Male	20 to 24	15% (7-27%)	85% (73-93%)	2 (0.65-4.3)	4.7% (2.8-7%)	5.2% (2.6-8.4%)	180 (90-290)	15000 (8400-22000)
Cameroon	2016	Male	25 to 29	19% (9.3-33%)	81% (67-91%)	3 (1-6.6)	6% (3.8-8.7%)	6.8% (3.9-10%)	290 (160-470)	21000 (13000-32000)
Cameroon	2016	Male	30 to 34	22% (10-38%)	78% (62-90%)	3.8 (1.4-8.1)	6.6% (4.1-9.8%)	7.4% (4.3-11%)	380 (220-590)	24000 (15000-36000)
Cameroon	2016	Male	35 to 39	25% (12-41%)	75% (59-88%)	4.2 (1.5-8.9)	7.1% (4.7-10%)	8.2% (5.2-12%)	440 (280-650)	25000 (16000-35000)
Cameroon	2016	Male	40 to 44	26% (13-44%)	74% (56-87%)	4.5 (1.6-9.9)	7.7% (5.3-11%)	9% (6.1-13%)	490 (310-710)	24000 (16000-34000)
Cameroon	2016	Male	45 to 49	26% (12-43%)	74% (57-88%)	4.7 (1.5-10)	9.3% (6.1-13%)	11% (7.3-15%)	570 (360-810)	24000 (16000-34000)
Cameroon	2016	Male	50 to 54	26% (12-46%)	74% (54-88%)	4.7 (1.4-10)	9.3% (6.2-13%)	11% (7.3-15%)	550 (340-780)	20000 (13000-29000)
Cameroon	2016	Male	55 to 59	27% (14-45%)	73% (55-86%)	4.6 (1.5-10)	10% (6.8-14%)	12% (8.3-17%)	580 (360-810)	19000 (12000-26000)
Cameroon	2016	Male	60 to 64	27% (13-44%)	73% (56-87%)	4.2 (1.1-9.6)	9% (5.7-13%)	11% (6.7-15%)	550 (340-810)	15000 (9400-22000)
Cameroon	2016	Male	65 to 69	26% (13-43%)	74% (57-87%)	3.5 (1.1-8.2)	8.1% (4.7-12%)	9.7% (5.7-14%)	550 (320-810)	12000 (7200-18000)
Cameroon	2016	Male	70 to 74	25% (13-42%)	75% (58-87%)	2.7 (0.79-6.4)	7.2% (3.7-11%)	8.5% (4.5-13%)	510 (270-810)	9300 (4900-15000)
Cameroon	2016	Male	75 to 79	23% (11-39%)	77% (61-89%)	2.1 (0.51-5.1)	5.6% (2.3-9.3%)	6.5% (2.7-11%)	360 (150-600)	5100 (2100-8400)
Cameroon	2016	Male	80 to 84	21% (9.5-37%)	79% (63-91%)	1.7 (0.45-4.2)	5% (2-8.4%)	5.8% (2.3-9.9%)	220 (91-380)	2300 (960-3900)
Cameroon	2016	Male	85 to 89	20% (8.8-35%)	80% (65-91%)	1.4 (0.35-3.6)	4.3% (1.5-8%)	5% (1.8-9.3%)	120 (40-220)	880 (310-1600)
Cameroon	2016	Male	90 to 94	21% (9.7-38%)	79% (62-90%)	1.4 (0.32-3.6)	3.6% (0.99-7.1%)	4.1% (1.1-8.4%)	38 (9.9-78)	200 (54-400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cameroon	2016	Male	95 plus	25% (10-46%)	75% (54-90%)	1.4 (0.24-4.3)	3.5% (0.7-7.8%)	4% (0.68-9%)	9.2 (1.6-21)	32 (6.4-72)
Cape Verde	1990	Female	All Ages	13% (9.5-17%)	87% (83-90%)	0.89 (0.78-1)	2.1% (1.6-2.6%)	4.4% (2.7-6.2%)	47 (28-65)	1400 (1100-1700)
Cape Verde	1990	Female	15 to 19	12% (5.8-22%)	88% (78-94%)	0.75 (0.3-1.5)	1.8% (1-3%)	2.9% (1.1-5.5%)	0.35 (0.13-0.68)	45 (25-74)
Cape Verde	1990	Female	20 to 24	15% (6.8-28%)	85% (72-93%)	1.2 (0.5-2.1)	3.5% (2.2-5.3%)	4.7% (2.2-8.3%)	0.74 (0.33-1.3)	92 (56-140)
Cape Verde	1990	Female	25 to 29	17% (7.7-31%)	83% (69-92%)	1.4 (0.65-2.5)	4% (2.6-5.7%)	5.9% (2.9-9.4%)	1.1 (0.53-1.8)	110 (68-160)
Cape Verde	1990	Female	30 to 34	20% (8.4-35%)	80% (65-92%)	1.7 (0.74-3)	4.4% (2.8-6.5%)	7.1% (3.9-11%)	1.1 (0.61-1.8)	91 (57-140)
Cape Verde	1990	Female	35 to 39	22% (10-38%)	78% (62-90%)	1.9 (0.77-3.4)	5.1% (3.2-7.5%)	8.8% (5-13%)	1.5 (0.83-2.3)	95 (59-140)
Cape Verde	1990	Female	40 to 44	23% (12-40%)	77% (60-88%)	2 (0.83-3.9)	5.5% (3.3-8.3%)	9.9% (5.4-15%)	1.1 (0.61-1.7)	60 (35-89)
Cape Verde	1990	Female	45 to 49	24% (11-40%)	76% (60-89%)	2 (0.72-4.2)	6.2% (3.4-9.5%)	11% (5.5-16%)	1.8 (0.9-2.8)	81 (45-120)
Cape Verde	1990	Female	50 to 54	24% (11-43%)	76% (57-89%)	2.1 (0.82-4.2)	6.4% (3.6-9.6%)	10% (5.7-15%)	3.2 (1.8-5)	130 (71-190)
Cape Verde	1990	Female	55 to 59	25% (12-43%)	75% (57-88%)	2.3 (0.96-4.4)	6.7% (3.8-10%)	10% (5.6-16%)	4.2 (2.3-6.6)	140 (80-220)
Cape Verde	1990	Female	60 to 64	26% (13-45%)	74% (55-87%)	2.4 (0.91-4.6)	6.4% (3.4-10%)	9.1% (4.8-15%)	5.1 (2.7-8.3)	150 (77-240)
Cape Verde	1990	Female	65 to 69	27% (14-44%)	73% (56-86%)	2.2 (0.88-4.1)	6.9% (3.5-11%)	9.5% (4.8-15%)	4.6 (2.3-7.3)	110 (54-170)
Cape Verde	1990	Female	70 to 74	25% (13-42%)	75% (58-87%)	1.8 (0.66-3.8)	6.2% (2.7-11%)	8.1% (3.4-14%)	6 (2.6-10)	110 (47-190)
Cape Verde	1990	Female	75 to 79	22% (11-39%)	78% (61-89%)	1.4 (0.43-3.2)	4.7% (1.5-8.7%)	6% (1.9-11%)	6 (1.9-11)	86 (28-160)
Cape Verde	1990	Female	80 to 84	20% (9.1-35%)	80% (65-91%)	1.2 (0.39-2.6)	4.3% (1.3-8.2%)	5.3% (1.6-10%)	4.7 (1.3-9)	49 (15-95)
Cape Verde	1990	Female	85 to 89	21% (9.7-36%)	79% (64-90%)	1.2 (0.41-2.8)	4% (1.1-8.5%)	4.9% (1.3-10%)	3.4 (0.92-7.3)	26 (7-54)
Cape Verde	1990	Female	90 to 94	24% (11-42%)	76% (58-89%)	1.4 (0.39-3.3)	3.8% (0.82-8.3%)	4.6% (0.91-10%)	1.3 (0.25-2.9)	6.7 (1.4-15)
Cape Verde	1990	Female	95 plus	28% (12-50%)	72% (50-88%)	1.6 (0.38-4.1)	3.7% (0.32-8.8%)	4.4% (0.26-11%)	0.42 (0.025-1)	1.5 (0.13-3.5)
Cape Verde	1990	Male	All Ages	9.3% (6.9-13%)	91% (87-93%)	0.86 (0.75-0.98)	2.4% (2-3%)	4.2% (3.1-5.3%)	54 (40-71)	1900 (1500-2300)
Cape Verde	1990	Male	15 to 19	7.7% (3.5-14%)	92% (86-96%)	0.54 (0.2-1.1)	1.9% (1-3.3%)	2.3% (0.94-4.5%)	0.66 (0.27-1.2)	67 (35-110)
Cape Verde	1990	Male	20 to 24	12% (5.4-21%)	88% (79-95%)	1.1 (0.46-2.1)	4.4% (2.6-7%)	5% (2.5-8.7%)	1.8 (0.85-3.1)	160 (89-250)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	1990	Male	25 to 29	15% (7.4-27%)	85% (73-93%)	1.7 (0.74-3.1)	6% (3.8-8.8%)	7.3% (4.2-11%)	2.4 (1.4-3.9)	190 (110-280)
Cape Verde	1990	Male	30 to 34	18% (7.8-33%)	82% (67-92%)	2.2 (0.97-3.9)	7.1% (4.6-10%)	8.9% (5.2-13%)	2.1 (1.2-3.2)	140 (90-210)
Cape Verde	1990	Male	35 to 39	20% (9.3-36%)	80% (64-91%)	2.5 (0.97-4.6)	8.2% (5.4-12%)	11% (6.7-16%)	2.8 (1.7-4.2)	160 (100-230)
Cape Verde	1990	Male	40 to 44	21% (11-38%)	79% (62-89%)	2.7 (1.1-5.4)	8.7% (5.9-12%)	11% (7.5-16%)	2 (1.3-3)	100 (66-150)
Cape Verde	1990	Male	45 to 49	22% (9.9-37%)	78% (63-90%)	2.9 (1.1-6.1)	10% (6.3-14%)	13% (8.1-18%)	2.5 (1.5-3.6)	110 (69-150)
Cape Verde	1990	Male	50 to 54	22% (10-40%)	78% (60-90%)	2.9 (1.1-5.9)	9.1% (5.9-13%)	11% (7.4-16%)	5.1 (3.2-7.3)	190 (120-270)
Cape Verde	1990	Male	55 to 59	22% (11-40%)	78% (60-89%)	2.8 (1.1-5.3)	8.7% (5.6-12%)	11% (6.9-15%)	6.8 (4.4-9.7)	220 (150-320)
Cape Verde	1990	Male	60 to 64	22% (11-39%)	78% (61-89%)	2.5 (0.89-4.9)	7.1% (4.3-11%)	8.6% (5.1-13%)	7.5 (4.4-11)	210 (120-310)
Cape Verde	1990	Male	65 to 69	22% (11-37%)	78% (63-89%)	2.1 (0.79-4.2)	6.2% (3.4-9.3%)	7.3% (4.1-11%)	5.5 (3-8.6)	130 (71-200)
Cape Verde	1990	Male	70 to 74	21% (10-35%)	79% (65-90%)	1.6 (0.56-3.3)	5% (2.3-8.3%)	5.9% (2.7-9.9%)	6.1 (2.8-10)	110 (51-190)
Cape Verde	1990	Male	75 to 79	19% (8.9-34%)	81% (66-91%)	1.2 (0.35-2.7)	3.5% (1.2-6.5%)	4% (1.4-7.5%)	4.4 (1.4-8.1)	63 (21-120)
Cape Verde	1990	Male	80 to 84	17% (7.6-31%)	83% (69-92%)	0.97 (0.29-2.1)	2.9% (1-5.4%)	3.3% (1.1-6.2%)	2.7 (0.9-5.1)	29 (10-54)
Cape Verde	1990	Male	85 to 89	16% (7.2-28%)	84% (72-93%)	0.82 (0.25-1.9)	2.4% (0.71-4.9%)	2.7% (0.76-5.6%)	1.4 (0.38-2.9)	10 (3.2-22)
Cape Verde	1990	Male	90 to 94	17% (7.7-31%)	83% (69-92%)	0.77 (0.2-2)	1.7% (0.31-4.4%)	1.9% (0.25-5%)	0.31 (0.04-0.8)	1.7 (0.29-4.3)
Cape Verde	1990	Male	95 plus	20% (7.7-40%)	80% (60-92%)	0.8 (0.16-2.3)	1.4% (-0.036-4.1%)	1.4% (-0.27-4.7%)	0.062 (-0.011-0.2)	0.24 (-0.0059-0.72)
Cape Verde	1995	Female	All Ages	14% (10-18%)	86% (82-90%)	0.99 (0.87-1.1)	2.1% (1.5-2.8%)	4.6% (2.8-6.5%)	55 (33-79)	1600 (1200-2100)
Cape Verde	1995	Female	15 to 19	12% (6-22%)	88% (78-94%)	0.82 (0.32-1.7)	1.9% (1-3.2%)	3% (1.2-5.6%)	0.41 (0.16-0.82)	52 (27-86)
Cape Verde	1995	Female	20 to 24	16% (7-28%)	84% (72-93%)	1.3 (0.53-2.5)	3.6% (2.2-5.4%)	4.8% (2.2-8.2%)	0.76 (0.34-1.4)	94 (56-140)
Cape Verde	1995	Female	25 to 29	18% (7.5-31%)	82% (69-92%)	1.6 (0.68-2.8)	4% (2.6-5.7%)	5.6% (2.8-9%)	1.1 (0.55-1.9)	110 (69-160)
Cape Verde	1995	Female	30 to 34	20% (8.8-35%)	80% (65-91%)	1.9 (0.85-3.2)	4.4% (2.8-6.3%)	6.8% (3.8-10%)	1.6 (0.89-2.7)	130 (81-200)
Cape Verde	1995	Female	35 to 39	23% (11-39%)	77% (61-89%)	2.1 (0.95-3.6)	5.1% (3.2-7.3%)	8.7% (5.1-13%)	1.9 (1-3)	120 (75-180)
Cape Verde	1995	Female	40 to 44	24% (12-41%)	76% (59-88%)	2.2 (0.94-4.2)	5.7% (3.6-8.5%)	10% (5.7-15%)	2.2 (1.2-3.5)	120 (72-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	1995	Female	45 to 49	25% (11-41%)	75% (59-89%)	2.3 (0.83-4.8)	6.4% (3.6-9.6%)	11% (5.8-16%)	1.7 (0.87-2.6)	76 (42-120)
Cape Verde	1995	Female	50 to 54	25% (12-44%)	75% (56-88%)	2.4 (0.95-4.9)	6.7% (3.8-10%)	11% (6.2-16%)	2.4 (1.3-3.9)	96 (53-150)
Cape Verde	1995	Female	55 to 59	26% (13-45%)	74% (55-87%)	2.6 (1-5)	7.1% (3.9-11%)	11% (5.9-17%)	4.6 (2.4-7.5)	160 (86-250)
Cape Verde	1995	Female	60 to 64	27% (13-46%)	73% (54-87%)	2.7 (1-5.2)	6.9% (3.7-11%)	9.9% (5.2-16%)	5.9 (3.2-9.7)	170 (91-270)
Cape Verde	1995	Female	65 to 69	28% (14-46%)	72% (54-86%)	2.5 (0.99-4.6)	7.5% (3.9-12%)	10% (5.4-16%)	8.2 (4.1-13)	190 (98-300)
Cape Verde	1995	Female	70 to 74	26% (13-42%)	74% (58-87%)	2 (0.72-4.1)	6.6% (2.8-11%)	8.7% (3.7-15%)	6 (2.5-11)	110 (48-200)
Cape Verde	1995	Female	75 to 79	22% (11-39%)	78% (61-89%)	1.5 (0.45-3.6)	5.1% (1.6-9.4%)	6.5% (2-12%)	5.7 (1.7-11)	83 (25-160)
Cape Verde	1995	Female	80 to 84	20% (9.2-37%)	80% (63-91%)	1.4 (0.41-2.9)	4.6% (1.4-9%)	5.7% (1.7-11%)	4.7 (1.4-9.3)	50 (15-97)
Cape Verde	1995	Female	85 to 89	21% (9.5-37%)	79% (63-90%)	1.3 (0.44-3)	4.4% (1.2-9.2%)	5.4% (1.4-11%)	4.6 (1.2-9.8)	35 (9.6-72)
Cape Verde	1995	Female	90 to 94	23% (11-41%)	77% (59-89%)	1.5 (0.41-3.6)	4.2% (0.91-8.8%)	5.1% (1-11%)	2.4 (0.48-5)	12 (2.6-25)
Cape Verde	1995	Female	95 plus	28% (12-51%)	72% (49-88%)	1.7 (0.38-4.5)	4.1% (0.43-9.7%)	4.9% (0.42-12%)	0.6 (0.052-1.4)	2.1 (0.22-4.8)
Cape Verde	1995	Male	All Ages	10% (7.5-14%)	90% (86-93%)	1 (0.89-1.2)	2.8% (2.1-3.5%)	4.7% (3.4-6%)	72 (52-94)	2600 (2000-3300)
Cape Verde	1995	Male	15 to 19	8.1% (3.8-15%)	92% (85-96%)	0.63 (0.22-1.3)	2.1% (1.1-3.7%)	2.6% (1.1-5%)	0.87 (0.35-1.7)	84 (43-150)
Cape Verde	1995	Male	20 to 24	13% (5.6-23%)	87% (77-94%)	1.3 (0.52-2.5)	4.9% (2.8-7.7%)	5.6% (2.7-9.6%)	2.4 (1.2-4.2)	210 (110-330)
Cape Verde	1995	Male	25 to 29	16% (7.3-30%)	84% (70-93%)	2 (0.89-3.5)	6.5% (4.1-9.6%)	7.9% (4.5-12%)	3.6 (1.9-5.8)	260 (160-400)
Cape Verde	1995	Male	30 to 34	19% (8.3-35%)	81% (65-92%)	2.5 (1.2-4.3)	7.7% (5-11%)	9.7% (5.7-14%)	4.3 (2.4-6.7)	280 (170-420)
Cape Verde	1995	Male	35 to 39	21% (9.9-37%)	79% (63-90%)	2.8 (1.2-5.2)	8.8% (5.8-13%)	11% (7.1-16%)	3.7 (2.2-5.6)	210 (130-300)
Cape Verde	1995	Male	40 to 44	22% (11-39%)	78% (61-89%)	3.1 (1.3-5.8)	9.7% (6.5-13%)	13% (8.4-17%)	4.8 (3-7.3)	240 (150-360)
Cape Verde	1995	Male	45 to 49	23% (10-38%)	77% (62-90%)	3.3 (1.2-7.1)	11% (6.9-15%)	14% (8.9-19%)	3.7 (2.2-5.5)	160 (98-230)
Cape Verde	1995	Male	50 to 54	23% (11-41%)	77% (59-89%)	3.3 (1.2-6.9)	10% (6.5-14%)	13% (8.2-18%)	3.5 (2.2-5.1)	130 (84-190)
Cape Verde	1995	Male	55 to 59	23% (12-40%)	77% (60-88%)	3.2 (1.2-6.1)	9.9% (6.5-14%)	12% (7.9-17%)	7.8 (4.9-11)	250 (160-370)
Cape Verde	1995	Male	60 to 64	23% (12-40%)	77% (60-88%)	2.8 (1-5.6)	8% (5-12%)	9.6% (5.9-14%)	9.1 (5.5-14)	250 (150-380)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	1995	Male	65 to 69	23% (11-39%)	77% (61-89%)	2.4 (0.91-4.6)	7% (3.9-10%)	8.3% (4.8-13%)	10 (5.6-16)	230 (130-360)
Cape Verde	1995	Male	70 to 74	21% (11-37%)	79% (63-89%)	1.8 (0.63-3.8)	5.7% (2.7-9.3%)	6.8% (3.1-11%)	6.9 (3.1-11)	130 (59-210)
Cape Verde	1995	Male	75 to 79	19% (9.2-35%)	81% (65-91%)	1.3 (0.38-3.1)	4% (1.3-7.4%)	4.6% (1.5-8.5%)	5.1 (1.6-9.4)	74 (24-130)
Cape Verde	1995	Male	80 to 84	18% (7.8-32%)	82% (68-92%)	1.1 (0.33-2.5)	3.4% (1.2-6.4%)	3.9% (1.3-7.4%)	3.3 (1.1-6.4)	36 (12-67)
Cape Verde	1995	Male	85 to 89	16% (7.6-30%)	84% (70-92%)	0.93 (0.28-2.1)	2.8% (0.85-5.7%)	3.2% (0.92-6.6%)	2.1 (0.59-4.2)	16 (4.8-32)
Cape Verde	1995	Male	90 to 94	18% (8.2-32%)	82% (68-92%)	0.88 (0.23-2.2)	2.2% (0.45-5.1%)	2.5% (0.42-5.9%)	0.69 (0.11-1.7)	3.7 (0.74-8.7)
Cape Verde	1995	Male	95 plus	21% (8.4-41%)	79% (59-92%)	0.93 (0.18-2.7)	2% (0.17-5.4%)	2.1% (0.017-6.1%)	0.11 (0.00088-0.33)	0.42 (0.037-1.2)
Cape Verde	2000	Female	All Ages	14% (10-19%)	86% (81-90%)	1.1 (0.94-1.2)	2.6% (1.9-3.5%)	5.6% (3.4-7.9%)	60 (34-89)	1700 (1200-2400)
Cape Verde	2000	Female	15 to 19	13% (6.1-23%)	87% (77-94%)	0.85 (0.33-1.7)	2% (1.1-3.2%)	3.5% (1.4-6.8%)	0.46 (0.16-0.89)	62 (35-100)
Cape Verde	2000	Female	20 to 24	16% (7.3-29%)	84% (71-93%)	1.4 (0.57-2.6)	3.7% (2.3-5.4%)	5.2% (2.4-8.9%)	0.74 (0.33-1.4)	98 (60-150)
Cape Verde	2000	Female	25 to 29	18% (8.3-33%)	82% (67-92%)	1.7 (0.73-3)	3.8% (2.4-5.4%)	5.3% (2.6-8.9%)	0.91 (0.44-1.5)	98 (61-150)
Cape Verde	2000	Female	30 to 34	21% (9.1-37%)	79% (63-91%)	2 (0.89-3.4)	4% (2.6-5.7%)	5.9% (3.1-9.3%)	1.3 (0.71-2.1)	110 (72-170)
Cape Verde	2000	Female	35 to 39	23% (11-39%)	77% (61-89%)	2.2 (0.98-3.6)	4.6% (2.9-6.7%)	7.7% (4.3-12%)	2.3 (1.2-3.5)	150 (92-220)
Cape Verde	2000	Female	40 to 44	25% (12-43%)	75% (57-88%)	2.3 (1-4.1)	5.2% (3.2-7.7%)	9% (5.2-14%)	2.4 (1.3-3.9)	130 (80-200)
Cape Verde	2000	Female	45 to 49	25% (11-42%)	75% (58-89%)	2.4 (0.91-4.9)	6% (3.5-9.2%)	10% (5.8-16%)	2.8 (1.5-4.5)	130 (72-200)
Cape Verde	2000	Female	50 to 54	26% (12-45%)	74% (55-88%)	2.6 (1-5.2)	6.5% (3.7-9.8%)	11% (6-16%)	2 (1.1-3.3)	79 (43-130)
Cape Verde	2000	Female	55 to 59	26% (13-45%)	74% (55-87%)	2.8 (1.1-5.4)	7.1% (3.9-11%)	11% (6.1-17%)	3.4 (1.8-5.6)	120 (63-190)
Cape Verde	2000	Female	60 to 64	27% (14-46%)	73% (54-86%)	2.8 (1.1-5.6)	6.9% (3.6-11%)	10% (5.3-16%)	6.4 (3.3-11)	180 (95-310)
Cape Verde	2000	Female	65 to 69	28% (14-46%)	72% (54-86%)	2.6 (1-4.8)	7.5% (3.8-12%)	11% (5.4-17%)	9 (4.3-15)	210 (110-350)
Cape Verde	2000	Female	70 to 74	26% (13-42%)	74% (58-87%)	2 (0.75-4.2)	6.6% (2.7-11%)	8.8% (3.5-15%)	9.5 (3.7-17)	180 (72-330)
Cape Verde	2000	Female	75 to 79	22% (11-40%)	78% (60-89%)	1.6 (0.46-3.7)	5.1% (1.5-9.7%)	6.6% (1.9-13%)	5.5 (1.6-11)	80 (24-160)
Cape Verde	2000	Female	80 to 84	21% (9.2-37%)	79% (63-91%)	1.4 (0.41-3)	4.8% (1.3-9.4%)	6% (1.6-12%)	4.3 (1.1-8.5)	46 (13-90)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	2000	Female	85 to 89	21% (10-37%)	79% (63-90%)	1.4 (0.46-3.2)	4.6% (1.2-9.6%)	5.7% (1.4-12%)	4.4 (1.1-9.3)	33 (8.3-68)
Cape Verde	2000	Female	90 to 94	24% (11-42%)	76% (58-89%)	1.6 (0.44-3.7)	4.6% (1-9.7%)	5.6% (1.1-12%)	3 (0.61-6.7)	16 (3.4-34)
Cape Verde	2000	Female	95 plus	29% (12-51%)	71% (49-88%)	1.8 (0.42-4.8)	4.6% (0.44-11%)	5.5% (0.35-13%)	1.1 (0.071-2.6)	3.8 (0.33-8.5)
Cape Verde	2000	Male	All Ages	11% (7.9-15%)	89% (85-92%)	1.1 (0.99-1.3)	3.8% (3-4.8%)	6% (4.5-7.5%)	90 (61-120)	3200 (2400-4200)
Cape Verde	2000	Male	15 to 19	8.6% (4.1-16%)	91% (84-96%)	0.69 (0.26-1.4)	2.6% (1.3-4.4%)	3.4% (1.4-6.2%)	1.2 (0.47-2.4)	120 (59-210)
Cape Verde	2000	Male	20 to 24	13% (6-24%)	87% (76-94%)	1.4 (0.57-2.7)	5.4% (3.2-8.6%)	6.6% (3.3-11%)	2.7 (1.3-4.8)	230 (130-370)
Cape Verde	2000	Male	25 to 29	17% (7.5-31%)	83% (69-93%)	2.1 (0.9-3.9)	7% (4.3-10%)	8.7% (4.9-14%)	3.9 (2-6.4)	290 (170-450)
Cape Verde	2000	Male	30 to 34	20% (8.5-36%)	80% (64-91%)	2.7 (1.2-4.7)	7.9% (4.9-11%)	9.8% (5.7-15%)	5 (2.8-8.1)	320 (200-500)
Cape Verde	2000	Male	35 to 39	22% (10-38%)	78% (62-90%)	3 (1.3-5.3)	9% (5.8-13%)	11% (7.2-16%)	6.4 (3.7-9.8)	360 (220-540)
Cape Verde	2000	Male	40 to 44	23% (12-40%)	77% (60-88%)	3.3 (1.4-6.1)	9.8% (6.7-14%)	12% (8.6-17%)	5.4 (3.4-8.3)	270 (170-400)
Cape Verde	2000	Male	45 to 49	24% (11-39%)	76% (61-89%)	3.5 (1.3-7.5)	12% (7.7-16%)	15% (9.9-21%)	7.4 (4.4-11)	320 (200-470)
Cape Verde	2000	Male	50 to 54	24% (11-43%)	76% (57-89%)	3.6 (1.4-7.6)	11% (7-15%)	14% (8.9-19%)	4.9 (3-7.5)	190 (120-280)
Cape Verde	2000	Male	55 to 59	24% (12-42%)	76% (58-88%)	3.4 (1.3-6.9)	11% (7.2-15%)	13% (8.8-19%)	5.3 (3.2-7.7)	170 (110-250)
Cape Verde	2000	Male	60 to 64	24% (12-41%)	76% (59-88%)	3.1 (1.1-6.3)	8.7% (5.4-13%)	11% (6.5-15%)	9.8 (5.7-15)	270 (160-420)
Cape Verde	2000	Male	65 to 69	24% (12-40%)	76% (60-88%)	2.5 (1-4.9)	7.6% (4.4-11%)	9.1% (5.2-14%)	12 (6.8-19)	280 (160-440)
Cape Verde	2000	Male	70 to 74	22% (12-38%)	78% (62-88%)	1.9 (0.7-4)	6.3% (3-10%)	7.4% (3.5-12%)	12 (5.2-20)	220 (97-370)
Cape Verde	2000	Male	75 to 79	20% (9.9-36%)	80% (64-90%)	1.4 (0.4-3.3)	4.3% (1.3-8.1%)	5% (1.5-9.4%)	6.1 (1.7-12)	88 (26-170)
Cape Verde	2000	Male	80 to 84	18% (8.4-33%)	82% (67-92%)	1.2 (0.35-2.7)	3.7% (1.1-7.1%)	4.3% (1.3-8.3%)	3.9 (1.1-7.7)	42 (12-81)
Cape Verde	2000	Male	85 to 89	17% (8-31%)	83% (69-92%)	1 (0.31-2.4)	3.2% (0.88-6.4%)	3.7% (0.94-7.4%)	2.4 (0.63-4.8)	18 (5.1-37)
Cape Verde	2000	Male	90 to 94	19% (8.8-34%)	81% (66-91%)	0.97 (0.26-2.4)	2.7% (0.57-6%)	3% (0.55-7.2%)	0.96 (0.17-2.3)	5.1 (1.1-12)
Cape Verde	2000	Male	95 plus	23% (9.1-44%)	77% (56-91%)	1 (0.2-2.9)	2.5% (0.39-6.7%)	2.8% (0.29-7.7%)	0.25 (0.026-0.68)	0.9 (0.14-2.4)
Cape Verde	2005	Female	All Ages	15% (11-19%)	85% (81-89%)	1.2 (1.1-1.4)	2.9% (2.2-3.9%)	6.1% (3.8-8.8%)	65 (40-97)	1900 (1400-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	2005	Female	15 to 19	13% (5.9-22%)	87% (78-94%)	0.95 (0.37-1.9)	1.9% (1.1-3.1%)	3.4% (1.4-6.4%)	0.49 (0.19-0.9)	71 (40-110)
Cape Verde	2005	Female	20 to 24	16% (7.5-29%)	84% (71-92%)	1.5 (0.65-2.9)	3.6% (2.4-5.2%)	5.2% (2.4-8.9%)	0.8 (0.38-1.4)	120 (73-170)
Cape Verde	2005	Female	25 to 29	18% (8.2-33%)	82% (67-92%)	1.9 (0.81-3.4)	3.7% (2.4-5.3%)	5% (2.4-8.5%)	0.86 (0.43-1.4)	100 (64-150)
Cape Verde	2005	Female	30 to 34	21% (8.8-38%)	79% (62-91%)	2.2 (0.97-3.9)	3.7% (2.4-5.3%)	5.4% (2.8-8.4%)	1.1 (0.58-1.7)	96 (63-140)
Cape Verde	2005	Female	35 to 39	23% (10-40%)	77% (60-90%)	2.4 (1-4.1)	4.2% (2.7-6%)	6.9% (3.8-11%)	1.7 (0.96-2.6)	120 (76-170)
Cape Verde	2005	Female	40 to 44	24% (12-42%)	76% (58-88%)	2.6 (1.1-4.6)	4.8% (3.1-7.1%)	8.3% (5-13%)	2.8 (1.6-4.4)	160 (99-240)
Cape Verde	2005	Female	45 to 49	25% (11-42%)	75% (58-89%)	2.7 (1-5.2)	5.6% (3.4-8.4%)	9.7% (5.7-15%)	3.1 (1.8-4.9)	150 (89-230)
Cape Verde	2005	Female	50 to 54	26% (12-45%)	74% (55-88%)	2.8 (1.2-5.5)	6.3% (3.7-9.6%)	11% (6.2-16%)	3.5 (1.9-5.7)	140 (79-230)
Cape Verde	2005	Female	55 to 59	26% (13-46%)	74% (54-87%)	3.1 (1.2-6.2)	6.9% (3.9-11%)	11% (6.1-17%)	2.5 (1.4-4.2)	88 (48-140)
Cape Verde	2005	Female	60 to 64	28% (13-47%)	72% (53-87%)	3.2 (1.2-6.6)	6.9% (3.8-11%)	10% (5.8-16%)	4.5 (2.4-7.6)	130 (71-220)
Cape Verde	2005	Female	65 to 69	28% (14-47%)	72% (53-86%)	2.9 (1.1-5.5)	7.6% (4-12%)	11% (5.8-17%)	9.5 (4.7-16)	230 (110-380)
Cape Verde	2005	Female	70 to 74	26% (14-43%)	74% (57-86%)	2.3 (0.8-4.9)	6.8% (2.9-11%)	9.3% (3.9-16%)	11 (4.7-20)	210 (91-380)
Cape Verde	2005	Female	75 to 79	22% (11-40%)	78% (60-89%)	1.8 (0.51-4.1)	5.3% (1.7-9.7%)	7% (2.3-13%)	9.3 (3-17)	140 (45-250)
Cape Verde	2005	Female	80 to 84	20% (9.2-37%)	80% (63-91%)	1.5 (0.44-3.4)	5% (1.5-9.3%)	6.3% (1.9-12%)	5.8 (1.7-11)	62 (19-120)
Cape Verde	2005	Female	85 to 89	21% (9.8-37%)	79% (63-90%)	1.5 (0.5-3.5)	4.9% (1.4-9.8%)	6.1% (1.8-12%)	4 (1.2-8)	30 (9-60)
Cape Verde	2005	Female	90 to 94	24% (11-42%)	76% (58-89%)	1.7 (0.49-4.1)	4.8% (1.2-9.9%)	5.9% (1.3-12%)	2.9 (0.63-6)	15 (3.5-30)
Cape Verde	2005	Female	95 plus	29% (12-52%)	71% (48-88%)	2 (0.47-5.2)	4.9% (0.66-11%)	5.9% (0.63-13%)	1.3 (0.14-2.9)	4.5 (0.61-9.8)
Cape Verde	2005	Male	All Ages	11% (8.3-15%)	89% (85-92%)	1.4 (1.2-1.5)	4.6% (3.7-5.7%)	6.7% (5.1-8.5%)	110 (78-150)	4000 (3000-5200)
Cape Verde	2005	Male	15 to 19	8.5% (3.9-16%)	92% (84-96%)	0.76 (0.28-1.5)	2.7% (1.4-4.7%)	3.5% (1.5-6.6%)	1.6 (0.63-3.2)	150 (75-270)
Cape Verde	2005	Male	20 to 24	13% (5.9-24%)	87% (76-94%)	1.6 (0.64-3)	5.7% (3.3-9%)	6.9% (3.5-12%)	3.9 (1.8-6.9)	320 (170-530)
Cape Verde	2005	Male	25 to 29	17% (8.1-30%)	83% (70-92%)	2.4 (1-4.4)	7.3% (4.5-11%)	9.2% (5.2-14%)	4.5 (2.4-7.6)	330 (200-530)
Cape Verde	2005	Male	30 to 34	20% (9-36%)	80% (64-91%)	3.1 (1.4-5.2)	8.2% (5.2-12%)	10% (6.2-16%)	5.5 (3-8.7)	360 (210-550)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	2005	Male	35 to 39	22% (10-38%)	78% (62-90%)	3.4 (1.4-6)	9.1% (6-13%)	12% (7.4-17%)	7 (4.1-11)	400 (250-600)
Cape Verde	2005	Male	40 to 44	23% (12-40%)	77% (60-88%)	3.6 (1.6-6.5)	10% (6.8-14%)	13% (9-18%)	8.5 (5.4-13)	420 (280-630)
Cape Verde	2005	Male	45 to 49	24% (11-39%)	76% (61-89%)	3.9 (1.5-7.6)	12% (7.8-16%)	15% (9.7-20%)	7.9 (5-12)	340 (220-500)
Cape Verde	2005	Male	50 to 54	24% (11-43%)	76% (57-89%)	3.9 (1.6-8.1)	11% (7.5-16%)	14% (9.4-19%)	9.1 (5.7-13)	350 (220-500)
Cape Verde	2005	Male	55 to 59	24% (12-42%)	76% (58-88%)	3.8 (1.4-7.6)	11% (7.4-15%)	13% (9.1-19%)	6.5 (4.1-9.4)	210 (140-310)
Cape Verde	2005	Male	60 to 64	24% (11-41%)	76% (59-89%)	3.4 (1.2-7.2)	8.9% (5.7-13%)	11% (6.8-16%)	6.6 (4-10)	190 (110-280)
Cape Verde	2005	Male	65 to 69	24% (12-40%)	76% (60-88%)	2.8 (1.1-5.7)	7.9% (4.7-12%)	9.4% (5.7-14%)	14 (7.9-21)	320 (190-480)
Cape Verde	2005	Male	70 to 74	22% (11-37%)	78% (63-89%)	2.1 (0.74-4.4)	6.7% (3.3-10%)	7.9% (3.8-12%)	16 (7.7-26)	300 (140-490)
Cape Verde	2005	Male	75 to 79	20% (9.7-36%)	80% (64-90%)	1.6 (0.46-3.7)	4.8% (1.7-8.5%)	5.5% (1.9-9.6%)	11 (3.6-20)	160 (52-290)
Cape Verde	2005	Male	80 to 84	18% (8.2-33%)	82% (67-92%)	1.3 (0.38-2.9)	4.1% (1.3-7.5%)	4.8% (1.5-8.6%)	6.1 (2-11)	65 (21-120)
Cape Verde	2005	Male	85 to 89	17% (8.1-32%)	83% (68-92%)	1.1 (0.34-2.7)	3.6% (1.1-7%)	4.2% (1.1-8.2%)	2.5 (0.69-4.9)	19 (5.5-37)
Cape Verde	2005	Male	90 to 94	19% (8.8-33%)	81% (67-91%)	1.1 (0.29-2.6)	3% (0.73-6.4%)	3.4% (0.76-7.4%)	0.96 (0.22-2.1)	5.1 (1.3-11)
Cape Verde	2005	Male	95 plus	23% (9.1-43%)	77% (57-91%)	1.2 (0.24-3.2)	2.8% (0.5-6.9%)	3.1% (0.39-7.8%)	0.28 (0.035-0.71)	1 (0.17-2.5)
Cape Verde	2010	Female	All Ages	15% (11-20%)	85% (80-89%)	1.3 (1.1-1.5)	3.1% (2.3-4.1%)	6.1% (3.5-9%)	65 (37-94)	1800 (1300-2400)
Cape Verde	2010	Female	15 to 19	12% (5.8-22%)	88% (78-94%)	0.94 (0.35-2)	1.8% (1.1-2.9%)	3.1% (1.2-5.8%)	0.41 (0.16-0.78)	65 (38-100)
Cape Verde	2010	Female	20 to 24	16% (6.9-29%)	84% (71-93%)	1.5 (0.6-2.9)	3.5% (2.3-4.9%)	4.8% (2.2-8.3%)	0.72 (0.33-1.2)	120 (76-180)
Cape Verde	2010	Female	25 to 29	18% (8-33%)	82% (67-92%)	1.9 (0.78-3.4)	3.5% (2.4-4.9%)	4.6% (2.3-7.8%)	0.84 (0.46-1.3)	110 (76-160)
Cape Verde	2010	Female	30 to 34	20% (8.9-37%)	80% (63-91%)	2.2 (0.94-3.8)	3.4% (2.3-4.8%)	5% (2.6-7.9%)	0.97 (0.57-1.5)	99 (66-140)
Cape Verde	2010	Female	35 to 39	23% (11-39%)	77% (61-89%)	2.4 (1-4.1)	3.9% (2.6-5.4%)	6.6% (3.6-10%)	1.4 (0.8-2)	100 (67-140)
Cape Verde	2010	Female	40 to 44	24% (12-41%)	76% (59-88%)	2.5 (1-4.6)	4.4% (2.9-6.4%)	8.1% (4.9-12%)	2.2 (1.3-3.3)	130 (86-190)
Cape Verde	2010	Female	45 to 49	25% (11-41%)	75% (59-89%)	2.6 (1-5.1)	5.3% (3.2-7.8%)	9.5% (5.4-14%)	3.6 (2-5.5)	180 (100-260)
Cape Verde	2010	Female	50 to 54	25% (12-44%)	75% (56-88%)	2.8 (1.1-5.5)	5.9% (3.4-8.8%)	10% (5.8-15%)	3.9 (2.2-6.1)	170 (95-250)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	2010	Female	55 to 59	26% (13-44%)	74% (56-87%)	3 (1.2-6.1)	6.5% (3.7-10%)	11% (5.9-16%)	4.1 (2.2-6.7)	150 (79-230)
Cape Verde	2010	Female	60 to 64	27% (14-46%)	73% (54-86%)	3.1 (1.1-6.5)	6.5% (3.6-11%)	9.7% (5.3-15%)	3 (1.6-4.8)	88 (47-140)
Cape Verde	2010	Female	65 to 69	28% (15-46%)	72% (54-85%)	2.8 (1.1-5.5)	7.2% (3.7-12%)	10% (5.5-17%)	6.3 (3.2-10)	150 (77-250)
Cape Verde	2010	Female	70 to 74	26% (14-43%)	74% (57-86%)	2.3 (0.76-4.9)	6.3% (2.8-11%)	8.7% (3.6-15%)	10 (4.1-17)	190 (79-330)
Cape Verde	2010	Female	75 to 79	22% (11-39%)	78% (61-89%)	1.7 (0.47-4.3)	4.9% (1.5-9.1%)	6.6% (2-12%)	9.9 (3-18)	150 (46-280)
Cape Verde	2010	Female	80 to 84	20% (9-36%)	80% (64-91%)	1.5 (0.44-3.3)	4.7% (1.3-8.8%)	6% (1.7-11%)	7.7 (2.1-14)	84 (24-150)
Cape Verde	2010	Female	85 to 89	21% (10-36%)	79% (64-90%)	1.5 (0.51-3.4)	4.7% (1.4-9.4%)	5.8% (1.6-12%)	5.6 (1.6-12)	42 (13-86)
Cape Verde	2010	Female	90 to 94	24% (11-41%)	76% (59-89%)	1.7 (0.45-4)	4.6% (1-9.5%)	5.7% (1.1-12%)	2.7 (0.52-5.7)	14 (2.9-29)
Cape Verde	2010	Female	95 plus	28% (12-51%)	72% (49-88%)	1.9 (0.44-5.1)	4.7% (0.66-10%)	5.6% (0.62-13%)	1.2 (0.14-2.8)	4.3 (0.59-9.6)
Cape Verde	2010	Male	All Ages	12% (9-16%)	88% (84-91%)	1.5 (1.3-1.8)	5.2% (4.3-6.4%)	7.1% (5.5-9%)	120 (92-160)	4400 (3500-5500)
Cape Verde	2010	Male	15 to 19	8.6% (4.1-16%)	91% (84-96%)	0.78 (0.29-1.6)	2.7% (1.4-4.7%)	3.6% (1.5-6.7%)	1.6 (0.64-3.1)	150 (74-260)
Cape Verde	2010	Male	20 to 24	13% (6.1-24%)	87% (76-94%)	1.6 (0.63-3.1)	5.7% (3.3-9%)	6.9% (3.4-12%)	4.5 (2.1-8)	370 (200-620)
Cape Verde	2010	Male	25 to 29	17% (7.8-30%)	83% (70-92%)	2.4 (1-4.2)	7.2% (4.5-11%)	9.1% (5.2-14%)	5.9 (3.1-9.7)	440 (260-680)
Cape Verde	2010	Male	30 to 34	20% (8.8-35%)	80% (65-91%)	3 (1.4-5.3)	8.1% (5.1-12%)	10% (6.2-16%)	6.1 (3.3-9.6)	400 (240-600)
Cape Verde	2010	Male	35 to 39	22% (10-38%)	78% (62-90%)	3.4 (1.4-6)	9.1% (6-13%)	12% (7.6-17%)	7 (4.1-10)	400 (250-590)
Cape Verde	2010	Male	40 to 44	23% (11-40%)	77% (60-89%)	3.6 (1.5-6.7)	9.9% (6.9-14%)	13% (9.1-18%)	8.4 (5.5-12)	420 (280-610)
Cape Verde	2010	Male	45 to 49	23% (11-39%)	77% (61-89%)	3.9 (1.5-7.7)	12% (7.9-16%)	15% (10-21%)	12 (7.5-16)	510 (330-710)
Cape Verde	2010	Male	50 to 54	24% (11-42%)	76% (58-89%)	3.9 (1.5-8.1)	11% (7.5-15%)	14% (9.4-19%)	9.7 (6.3-14)	370 (240-510)
Cape Verde	2010	Male	55 to 59	24% (12-42%)	76% (58-88%)	3.9 (1.4-7.9)	11% (7.5-15%)	14% (9.2-19%)	11 (7.2-15)	360 (240-500)
Cape Verde	2010	Male	60 to 64	24% (11-42%)	76% (58-89%)	3.5 (1.2-7.4)	8.8% (5.7-13%)	11% (6.7-16%)	6.4 (4.1-9.5)	180 (110-270)
Cape Verde	2010	Male	65 to 69	24% (12-40%)	76% (60-88%)	2.8 (1-5.8)	7.9% (4.7-12%)	9.3% (5.6-14%)	8.8 (5.2-14)	200 (120-310)
Cape Verde	2010	Male	70 to 74	22% (11-37%)	78% (63-89%)	2.1 (0.72-4.6)	6.6% (3.1-10%)	7.7% (3.6-12%)	16 (7.9-27)	300 (150-490)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	2010	Male	75 to 79	20% (9.4-36%)	80% (64-91%)	1.6 (0.44-3.8)	4.6% (1.6-8.4%)	5.3% (1.8-9.7%)	14 (4.6-26)	200 (69-370)
Cape Verde	2010	Male	80 to 84	18% (8.3-33%)	82% (67-92%)	1.3 (0.37-2.9)	4% (1.3-7.2%)	4.6% (1.4-8.3%)	8.4 (2.7-15)	90 (29-160)
Cape Verde	2010	Male	85 to 89	17% (8.1-31%)	83% (69-92%)	1.1 (0.34-2.5)	3.5% (1-6.7%)	4% (1-7.8%)	3.4 (0.9-6.7)	26 (7.3-50)
Cape Verde	2010	Male	90 to 94	19% (8.7-33%)	81% (67-91%)	1.1 (0.28-2.7)	2.9% (0.64-6.4%)	3.3% (0.65-7.4%)	0.86 (0.17-1.9)	4.6 (1-10)
Cape Verde	2010	Male	95 plus	22% (9.1-43%)	78% (57-91%)	1.1 (0.23-3.3)	2.7% (0.43-6.7%)	2.9% (0.32-7.7%)	0.23 (0.025-0.6)	0.83 (0.13-2.1)
Cape Verde	2016	Female	All Ages	15% (11-20%)	85% (80-89%)	1.4 (0.86-2.1)	3.2% (2.3-4.2%)	5.8% (2.9-8.9%)	65 (32-100)	1900 (1300-2500)
Cape Verde	2016	Female	15 to 19	12% (5.8-22%)	88% (78-94%)	0.92 (0.3-2.1)	1.7% (0.99-2.8%)	2.9% (1-5.8%)	0.3 (0.1-0.61)	51 (29-87)
Cape Verde	2016	Female	20 to 24	16% (6.8-29%)	84% (71-93%)	1.5 (0.5-3.1)	3.3% (2.1-4.8%)	4.2% (1.8-7.5%)	0.64 (0.27-1.2)	120 (72-180)
Cape Verde	2016	Female	25 to 29	18% (7.9-32%)	82% (68-92%)	1.8 (0.73-3.7)	3.4% (2.3-4.6%)	4.3% (2.1-7.2%)	0.87 (0.46-1.4)	130 (87-190)
Cape Verde	2016	Female	30 to 34	20% (9-37%)	80% (63-91%)	2.1 (0.9-4.2)	3.3% (2.3-4.6%)	4.8% (2.7-7.7%)	1.1 (0.63-1.7)	120 (80-170)
Cape Verde	2016	Female	35 to 39	23% (11-40%)	77% (60-89%)	2.4 (0.9-4.6)	3.7% (2.4-5.3%)	6.4% (3.6-10%)	1.4 (0.81-2.2)	110 (72-160)
Cape Verde	2016	Female	40 to 44	24% (12-42%)	76% (58-88%)	2.5 (0.94-5.2)	4.1% (2.6-6%)	7.7% (4.6-12%)	1.9 (1.1-2.8)	120 (74-170)
Cape Verde	2016	Female	45 to 49	25% (11-41%)	75% (59-89%)	2.5 (0.95-5.5)	4.8% (2.9-7.2%)	8.9% (5-13%)	2.9 (1.6-4.6)	150 (84-220)
Cape Verde	2016	Female	50 to 54	25% (12-44%)	75% (56-88%)	2.7 (1-5.5)	5.4% (3.2-8.1%)	9.4% (5.4-14%)	4.6 (2.4-7.4)	190 (110-310)
Cape Verde	2016	Female	55 to 59	25% (13-44%)	75% (56-87%)	2.9 (1-6.1)	5.9% (3.2-9.1%)	9.6% (5.2-15%)	5.3 (2.8-8.7)	190 (100-300)
Cape Verde	2016	Female	60 to 64	27% (13-45%)	73% (55-87%)	3 (1-6.8)	5.9% (3.1-9.5%)	8.8% (4.4-14%)	5.4 (2.6-9)	160 (80-270)
Cape Verde	2016	Female	65 to 69	27% (14-45%)	73% (55-86%)	2.7 (0.95-5.9)	6.5% (3.1-11%)	9.3% (4.3-15%)	4.2 (1.9-7.1)	100 (48-170)
Cape Verde	2016	Female	70 to 74	25% (13-42%)	75% (58-87%)	2.2 (0.69-5.2)	5.6% (2-9.9%)	7.7% (2.7-14%)	5.8 (1.9-10)	110 (39-200)
Cape Verde	2016	Female	75 to 79	22% (10-39%)	78% (61-90%)	1.7 (0.42-4.5)	4.5% (1.3-8.6%)	5.9% (1.6-11%)	8.3 (2.2-16)	120 (34-240)
Cape Verde	2016	Female	80 to 84	20% (9.2-36%)	80% (64-91%)	1.5 (0.41-3.6)	4.3% (1.1-8.4%)	5.4% (1.4-11%)	8.5 (2.1-17)	92 (24-180)
Cape Verde	2016	Female	85 to 89	21% (9.6-37%)	79% (63-90%)	1.5 (0.42-3.7)	4.2% (1.1-8.7%)	5.2% (1.2-11%)	8.5 (1.9-18)	64 (16-130)
Cape Verde	2016	Female	90 to 94	23% (11-40%)	77% (60-89%)	1.6 (0.43-4.3)	4.1% (0.72-9%)	5% (0.78-11%)	4.6 (0.72-10)	24 (4.1-52)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cape Verde	2016	Female	95 plus	28% (11-51%)	72% (49-89%)	1.9 (0.41-5.3)	4.3% (0.46-10%)	5.1% (0.33-12%)	1.3 (0.082-3.1)	4.5 (0.47-10)
Cape Verde	2016	Male	All Ages	13% (9.3-17%)	87% (83-91%)	1.6 (1-2.5)	5.5% (4.4-6.9%)	7% (5.3-8.9%)	120 (86-160)	4500 (3500-5900)
Cape Verde	2016	Male	15 to 19	8.3% (4-16%)	92% (84-96%)	0.75 (0.24-1.7)	2.5% (1.3-4.5%)	3.3% (1.3-6.2%)	1.3 (0.49-2.5)	120 (57-220)
Cape Verde	2016	Male	20 to 24	13% (5.7-23%)	87% (77-94%)	1.5 (0.51-3.3)	5.2% (2.9-8.4%)	6.2% (2.7-11%)	3.7 (1.6-6.8)	320 (170-530)
Cape Verde	2016	Male	25 to 29	16% (7.7-29%)	84% (71-92%)	2.3 (0.88-4.6)	6.5% (4.1-9.6%)	8.2% (4.4-13%)	5.4 (2.9-8.8)	420 (260-650)
Cape Verde	2016	Male	30 to 34	19% (8.7-34%)	81% (66-91%)	2.9 (1.2-5.5)	7.4% (4.8-11%)	9.6% (5.8-14%)	6.9 (3.9-11)	470 (290-700)
Cape Verde	2016	Male	35 to 39	22% (10-37%)	78% (63-90%)	3.3 (1.3-6.4)	8.4% (5.6-12%)	11% (7.1-16%)	7.6 (4.3-12)	440 (270-660)
Cape Verde	2016	Male	40 to 44	23% (11-39%)	77% (61-89%)	3.5 (1.3-7.6)	9.4% (6.4-13%)	13% (8.6-17%)	8.6 (5.3-13)	440 (280-640)
Cape Verde	2016	Male	45 to 49	23% (11-38%)	77% (62-89%)	3.7 (1.3-8.3)	11% (7.4-15%)	15% (9.8-20%)	12 (7.2-17)	510 (320-730)
Cape Verde	2016	Male	50 to 54	23% (11-42%)	77% (58-89%)	3.7 (1.3-8.2)	11% (7.1-15%)	14% (9.1-19%)	13 (8.5-20)	510 (330-740)
Cape Verde	2016	Male	55 to 59	23% (12-41%)	77% (59-88%)	3.6 (1.2-7.5)	10% (6.8-14%)	13% (8.5-18%)	12 (7.9-18)	400 (260-580)
Cape Verde	2016	Male	60 to 64	23% (11-40%)	77% (60-89%)	3.3 (1-7.4)	8.4% (5.2-12%)	10% (6.2-15%)	11 (6.5-17)	310 (190-460)
Cape Verde	2016	Male	65 to 69	23% (11-39%)	77% (61-89%)	2.7 (0.88-5.9)	7.1% (4-11%)	8.5% (4.8-13%)	7.1 (3.9-11)	160 (91-260)
Cape Verde	2016	Male	70 to 74	22% (11-38%)	78% (62-89%)	2.1 (0.66-5)	5.8% (2.6-9.3%)	6.9% (3-11%)	6.7 (2.9-11)	130 (55-210)
Cape Verde	2016	Male	75 to 79	20% (9.5-34%)	80% (66-90%)	1.6 (0.39-3.9)	4.2% (1.3-7.8%)	4.9% (1.5-9%)	9.5 (3-18)	140 (44-260)
Cape Verde	2016	Male	80 to 84	18% (7.7-32%)	82% (68-92%)	1.3 (0.31-3.1)	3.6% (1.1-6.9%)	4.1% (1.2-8%)	8.6 (2.5-17)	92 (28-180)
Cape Verde	2016	Male	85 to 89	17% (7.9-30%)	83% (70-92%)	1.1 (0.3-2.7)	3.1% (0.84-6.4%)	3.5% (0.88-7.3%)	5 (1.3-10)	38 (10-77)
Cape Verde	2016	Male	90 to 94	18% (8.3-34%)	82% (66-92%)	1 (0.27-2.7)	2.6% (0.51-6%)	2.9% (0.49-7%)	1.4 (0.23-3.2)	7.3 (1.4-17)
Cape Verde	2016	Male	95 plus	22% (8.9-41%)	78% (59-91%)	1.1 (0.2-3.2)	2.4% (0.32-6.1%)	2.6% (0.21-7%)	0.22 (0.018-0.59)	0.82 (0.11-2.1)
Chad	1990	Female	All Ages	15% (13-18%)	85% (82-87%)	0.33 (0.29-0.37)	0.46% (0.32-0.63%)	0.85% (0.48-1.3%)	350 (200-530)	14000 (9500-19000)
Chad	1990	Female	15 to 19	14% (9.4-20%)	86% (80-91%)	0.26 (0.15-0.43)	0.59% (0.35-0.93%)	0.37% (0.085-0.82%)	2.3 (0.49-5.1)	430 (250-690)
Chad	1990	Female	20 to 24	18% (12-26%)	82% (74-88%)	0.42 (0.25-0.67)	1.4% (0.81-2.2%)	0.96% (0.27-2%)	6.9 (2-15)	990 (570-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	1990	Female	25 to 29	21% (14-30%)	79% (70-86%)	0.58 (0.36-0.86)	1.8% (1.1-2.7%)	1.6% (0.73-2.9%)	11 (5-21)	1200 (690-1800)
Chad	1990	Female	30 to 34	23% (15-33%)	77% (67-85%)	0.71 (0.47-1)	2.1% (1.3-3.1%)	2.3% (1.2-3.8%)	17 (8.9-28)	1300 (830-1900)
Chad	1990	Female	35 to 39	26% (17-35%)	74% (65-83%)	0.8 (0.5-1.1)	2.5% (1.5-3.8%)	3.1% (1.6-5%)	24 (13-39)	1500 (900-2300)
Chad	1990	Female	40 to 44	26% (18-35%)	74% (65-82%)	0.8 (0.5-1.1)	2.6% (1.6-3.9%)	3.4% (1.9-5.4%)	23 (13-38)	1200 (740-1800)
Chad	1990	Female	45 to 49	27% (17-35%)	73% (65-83%)	0.72 (0.45-1)	2.6% (1.6-4%)	3.5% (1.9-5.5%)	26 (15-42)	1200 (690-1800)
Chad	1990	Female	50 to 54	26% (18-36%)	74% (64-82%)	0.74 (0.47-1.1)	2.7% (1.6-4.4%)	3.7% (2-5.7%)	33 (19-52)	1300 (730-1900)
Chad	1990	Female	55 to 59	26% (18-34%)	74% (66-82%)	0.86 (0.53-1.3)	3% (1.7-4.6%)	4% (2.2-6.1%)	44 (24-69)	1400 (790-2200)
Chad	1990	Female	60 to 64	27% (19-37%)	73% (63-81%)	0.93 (0.56-1.4)	2.8% (1.5-4.6%)	3.6% (1.9-5.8%)	49 (26-82)	1300 (710-2200)
Chad	1990	Female	65 to 69	29% (21-38%)	71% (62-79%)	0.89 (0.55-1.3)	2.3% (1-4%)	2.8% (1.2-4.9%)	47 (20-83)	1100 (450-1800)
Chad	1990	Female	70 to 74	27% (20-36%)	73% (64-80%)	0.72 (0.43-1.1)	1.8% (0.44-3.8%)	2.2% (0.49-4.6%)	40 (9.1-83)	720 (170-1500)
Chad	1990	Female	75 to 79	22% (15-32%)	78% (68-85%)	0.53 (0.29-0.88)	0.97% (-0.072-2.4%)	1.1% (-0.11-2.9%)	18 (-1.7-46)	260 (-20-640)
Chad	1990	Female	80 to 84	19% (13-27%)	81% (73-87%)	0.45 (0.26-0.71)	0.56% (-0.21-1.5%)	0.65% (-0.27-1.8%)	6.9 (-2.9-20)	73 (-27-200)
Chad	1990	Female	85 to 89	20% (13-28%)	80% (72-87%)	0.45 (0.26-0.69)	0.28% (-0.42-1.2%)	0.31% (-0.55-1.4%)	2.1 (-3.8-9.3)	17 (-25-69)
Chad	1990	Female	90 to 94	21% (14-30%)	79% (70-86%)	0.46 (0.25-0.75)	0.037% (-0.64-0.81%)	-0.01% (-0.85-0.94%)	-0.03 (-2.5-2.8)	0.67 (-11-15)
Chad	1990	Female	95 plus	23% (15-33%)	77% (67-85%)	0.48 (0.22-0.82)	-0.076% (-0.89-0.86%)	-0.19% (-1.2-0.93%)	-0.16 (-1-0.82)	-0.27 (-3.2-3)
Chad	1990	Male	All Ages	15% (12-17%)	85% (83-88%)	0.65 (0.6-0.7)	1.1% (0.9-1.3%)	1.9% (1.5-2.4%)	840 (650-1000)	36000 (29000-43000)
Chad	1990	Male	15 to 19	11% (7.2-16%)	89% (84-93%)	0.33 (0.19-0.53)	0.98% (0.61-1.5%)	0.92% (0.47-1.6%)	6.2 (3.2-11)	720 (450-1100)
Chad	1990	Male	20 to 24	19% (12-27%)	81% (73-88%)	0.85 (0.54-1.3)	3% (2-4.2%)	3% (1.7-4.7%)	26 (16-41)	2300 (1500-3200)
Chad	1990	Male	25 to 29	26% (17-35%)	74% (65-83%)	1.5 (0.97-2.1)	5.3% (3.5-7.3%)	6% (3.7-8.9%)	52 (33-75)	3700 (2500-5100)
Chad	1990	Male	30 to 34	29% (20-41%)	71% (59-80%)	1.8 (1.2-2.5)	6.7% (4.5-9%)	8.2% (5.2-11%)	69 (45-96)	4400 (3000-5900)
Chad	1990	Male	35 to 39	31% (20-41%)	69% (59-80%)	1.8 (1.1-2.5)	7.2% (4.9-9.8%)	9% (6-13%)	79 (54-110)	4400 (3100-5800)
Chad	1990	Male	40 to 44	30% (21-40%)	70% (60-79%)	1.8 (1.1-2.5)	7.7% (5.3-10%)	9.9% (6.6-14%)	82 (55-110)	4000 (2800-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	1990	Male	45 to 49	30% (19-39%)	70% (61-81%)	1.7 (1.1-2.4)	8.2% (5.5-11%)	10% (7-14%)	97 (65-130)	4100 (2800-5500)
Chad	1990	Male	50 to 54	29% (21-39%)	71% (61-79%)	1.6 (1.1-2.5)	7.7% (5-11%)	9.8% (6.3-14%)	100 (65-140)	3700 (2500-5300)
Chad	1990	Male	55 to 59	29% (20-38%)	71% (62-80%)	1.5 (0.95-2.3)	6.6% (4.1-9.5%)	8.2% (5.1-12%)	93 (57-140)	3000 (1900-4300)
Chad	1990	Male	60 to 64	28% (19-39%)	72% (61-81%)	1.3 (0.78-2.2)	5% (3-7.9%)	6.1% (3.6-9.7%)	81 (48-130)	2200 (1400-3500)
Chad	1990	Male	65 to 69	28% (20-36%)	72% (64-80%)	1.1 (0.68-1.7)	3.8% (2.3-5.8%)	4.6% (2.7-7.1%)	67 (40-100)	1500 (930-2300)
Chad	1990	Male	70 to 74	26% (19-35%)	74% (65-81%)	0.83 (0.51-1.3)	2.6% (1.3-4.2%)	3.1% (1.6-5.1%)	49 (24-81)	890 (450-1500)
Chad	1990	Male	75 to 79	23% (16-32%)	77% (68-84%)	0.61 (0.35-0.99)	1.7% (0.6-3%)	1.9% (0.66-3.6%)	25 (8.8-48)	360 (130-670)
Chad	1990	Male	80 to 84	20% (13-27%)	80% (73-87%)	0.47 (0.27-0.74)	1.2% (0.4-2.2%)	1.3% (0.43-2.5%)	10 (3.4-20)	110 (39-210)
Chad	1990	Male	85 to 89	19% (12-27%)	81% (73-88%)	0.38 (0.22-0.6)	0.74% (0.13-1.4%)	0.83% (0.079-1.7%)	3.8 (0.37-7.9)	30 (5.5-59)
Chad	1990	Male	90 to 94	19% (13-28%)	81% (72-87%)	0.33 (0.18-0.55)	0.28% (-0.26-0.81%)	0.24% (-0.42-0.91%)	0.46 (-0.8-1.7)	3.2 (-2.9-9.3)
Chad	1990	Male	95 plus	22% (13-32%)	78% (68-87%)	0.31 (0.14-0.55)	0.11% (-0.53-0.71%)	-0.017% (-0.8-0.75%)	-0.0081 (-0.38-0.36)	0.2 (-0.99-1.4)
Chad	1995	Female	All Ages	15% (13-18%)	85% (82-87%)	0.35 (0.31-0.39)	0.5% (0.35-0.68%)	0.92% (0.52-1.4%)	430 (250-630)	17000 (12000-23000)
Chad	1995	Female	15 to 19	14% (9.6-20%)	86% (80-90%)	0.29 (0.17-0.46)	0.62% (0.37-0.97%)	0.42% (0.11-0.91%)	2.9 (0.73-6.3)	530 (310-840)
Chad	1995	Female	20 to 24	19% (12-26%)	81% (74-88%)	0.46 (0.27-0.73)	1.4% (0.82-2.2%)	1% (0.32-2.1%)	9.2 (2.9-20)	1300 (730-2000)
Chad	1995	Female	25 to 29	21% (14-30%)	79% (70-86%)	0.64 (0.4-0.94)	1.7% (0.99-2.5%)	1.4% (0.64-2.7%)	14 (5.9-25)	1400 (830-2100)
Chad	1995	Female	30 to 34	24% (16-34%)	76% (66-84%)	0.79 (0.52-1.1)	1.9% (1.2-2.9%)	2% (1-3.5%)	19 (9.7-33)	1500 (920-2300)
Chad	1995	Female	35 to 39	26% (17-35%)	74% (65-83%)	0.89 (0.56-1.3)	2.4% (1.4-3.7%)	2.8% (1.5-4.8%)	28 (15-46)	1700 (1000-2600)
Chad	1995	Female	40 to 44	27% (19-37%)	73% (63-81%)	0.89 (0.56-1.3)	2.5% (1.5-3.8%)	3.2% (1.7-5.1%)	32 (17-51)	1700 (960-2500)
Chad	1995	Female	45 to 49	27% (17-36%)	73% (64-83%)	0.8 (0.5-1.2)	2.6% (1.5-4%)	3.4% (1.9-5.5%)	30 (17-49)	1300 (800-2100)
Chad	1995	Female	50 to 54	27% (19-36%)	73% (64-81%)	0.82 (0.52-1.2)	2.9% (1.6-4.5%)	3.8% (2-5.9%)	39 (21-62)	1500 (820-2300)
Chad	1995	Female	55 to 59	27% (18-35%)	73% (65-82%)	0.96 (0.58-1.4)	3.2% (1.8-5.1%)	4.2% (2.3-6.7%)	52 (28-83)	1700 (940-2600)
Chad	1995	Female	60 to 64	28% (19-38%)	72% (62-81%)	1 (0.62-1.6)	3.1% (1.6-5%)	3.9% (2-6.2%)	59 (30-97)	1600 (840-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	1995	Female	65 to 69	30% (22-39%)	70% (61-78%)	0.97 (0.6-1.4)	2.6% (1.1-4.3%)	3.1% (1.3-5.3%)	57 (23-98)	1300 (530-2200)
Chad	1995	Female	70 to 74	28% (20-36%)	72% (64-80%)	0.77 (0.46-1.2)	2% (0.48-3.8%)	2.4% (0.55-4.7%)	48 (11-95)	880 (210-1700)
Chad	1995	Female	75 to 79	22% (15-32%)	78% (68-85%)	0.57 (0.31-0.92)	1.1% (-0.026-2.6%)	1.3% (-0.061-3.2%)	23 (-1.1-56)	330 (-8-790)
Chad	1995	Female	80 to 84	20% (13-27%)	80% (73-87%)	0.48 (0.28-0.75)	0.67% (-0.16-1.7%)	0.78% (-0.2-2%)	9.5 (-2.4-24)	100 (-23-250)
Chad	1995	Female	85 to 89	20% (13-28%)	80% (72-87%)	0.47 (0.27-0.75)	0.4% (-0.32-1.4%)	0.44% (-0.43-1.6%)	3.5 (-3.4-13)	27 (-22-93)
Chad	1995	Female	90 to 94	22% (14-30%)	78% (70-86%)	0.49 (0.26-0.81)	0.15% (-0.57-1.1%)	0.13% (-0.77-1.3%)	0.43 (-2.5-4.2)	3.1 (-12-22)
Chad	1995	Female	95 plus	24% (15-33%)	76% (67-85%)	0.51 (0.25-0.87)	0.042% (-0.84-1.1%)	-0.053% (-1.1-1.3%)	-0.054 (-1.2-1.3)	0.17 (-3.5-4.8)
Chad	1995	Male	All Ages	15% (13-18%)	85% (82-87%)	0.69 (0.64-0.74)	1.2% (0.97-1.5%)	2.1% (1.6-2.6%)	1000 (790-1300)	43000 (35000-53000)
Chad	1995	Male	15 to 19	11% (7.4-16%)	89% (84-93%)	0.36 (0.21-0.57)	1.1% (0.69-1.7%)	1.1% (0.52-2%)	8 (3.7-15)	910 (560-1400)
Chad	1995	Male	20 to 24	20% (13-27%)	80% (73-87%)	0.93 (0.6-1.3)	3.3% (2.1-4.9%)	3.5% (1.9-5.6%)	34 (18-55)	2900 (1900-4400)
Chad	1995	Male	25 to 29	27% (18-36%)	73% (64-82%)	1.6 (1.1-2.2)	5.3% (3.2-7.8%)	6% (3.2-9.3%)	64 (36-97)	4600 (2800-6500)
Chad	1995	Male	30 to 34	30% (21-42%)	70% (58-79%)	2 (1.3-2.7)	6.3% (4-8.8%)	7.4% (4.2-11%)	83 (49-120)	5200 (3300-7300)
Chad	1995	Male	35 to 39	31% (21-41%)	69% (59-79%)	1.9 (1.3-2.7)	6.7% (4.1-9.4%)	8.1% (4.6-12%)	94 (57-130)	5200 (3300-7200)
Chad	1995	Male	40 to 44	31% (22-42%)	69% (58-78%)	1.9 (1.2-2.6)	7.3% (4.8-10%)	9% (5.7-13%)	110 (68-150)	5200 (3400-7200)
Chad	1995	Male	45 to 49	30% (20-40%)	70% (60-80%)	1.9 (1.2-2.6)	8% (5.2-11%)	9.9% (6.3-14%)	110 (72-160)	4800 (3200-6600)
Chad	1995	Male	50 to 54	30% (21-40%)	70% (60-79%)	1.8 (1.2-2.7)	7.8% (5.1-11%)	9.7% (6.2-14%)	120 (78-180)	4500 (3000-6500)
Chad	1995	Male	55 to 59	30% (21-40%)	70% (60-79%)	1.7 (1-2.5)	6.9% (4.3-10%)	8.5% (5.2-12%)	120 (70-170)	3700 (2300-5500)
Chad	1995	Male	60 to 64	29% (20-40%)	71% (60-80%)	1.4 (0.85-2.3)	5.3% (3.2-8.3%)	6.3% (3.8-10%)	98 (57-160)	2700 (1600-4300)
Chad	1995	Male	65 to 69	29% (21-38%)	71% (62-79%)	1.2 (0.74-1.8)	4% (2.4-6.1%)	4.8% (2.8-7.4%)	80 (47-130)	1800 (1100-2800)
Chad	1995	Male	70 to 74	27% (19-36%)	73% (64-81%)	0.91 (0.56-1.4)	2.8% (1.4-4.7%)	3.4% (1.7-5.5%)	60 (30-100)	1100 (550-1800)
Chad	1995	Male	75 to 79	24% (16-34%)	76% (66-84%)	0.67 (0.39-1.1)	1.8% (0.64-3.3%)	2.1% (0.72-3.9%)	33 (11-61)	460 (170-860)
Chad	1995	Male	80 to 84	21% (14-28%)	79% (72-86%)	0.52 (0.3-0.81)	1.3% (0.45-2.5%)	1.5% (0.47-2.9%)	14 (4.4-28)	150 (51-290)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	1995	Male	85 to 89	19% (12-28%)	81% (72-88%)	0.42 (0.24-0.65)	0.86% (0.18-1.6%)	0.96% (0.12-1.9%)	5.2 (0.66-10)	40 (8.7-77)
Chad	1995	Male	90 to 94	20% (13-29%)	80% (71-87%)	0.36 (0.2-0.6)	0.38% (-0.19-1%)	0.36% (-0.33-1.1%)	0.75 (-0.7-2.4)	4.7 (-2.4-13)
Chad	1995	Male	95 plus	23% (14-33%)	77% (67-86%)	0.34 (0.15-0.61)	0.23% (-0.51-1%)	0.14% (-0.77-1.1%)	0.075 (-0.43-0.63)	0.52 (-1.1-2.3)
Chad	2000	Female	All Ages	15% (14-17%)	85% (83-86%)	0.34 (0.3-0.38)	0.5% (0.34-0.69%)	0.88% (0.48-1.3%)	490 (270-740)	20000 (13000-27000)
Chad	2000	Female	15 to 19	15% (11-19%)	85% (81-89%)	0.28 (0.18-0.42)	0.63% (0.36-0.99%)	0.44% (0.09-0.96%)	3.9 (0.81-8.7)	660 (370-1100)
Chad	2000	Female	20 to 24	19% (15-24%)	81% (76-85%)	0.46 (0.3-0.68)	1.4% (0.79-2.3%)	1.1% (0.33-2.3%)	13 (3.8-28)	1600 (930-2700)
Chad	2000	Female	25 to 29	22% (17-28%)	78% (72-83%)	0.64 (0.46-0.87)	1.5% (0.89-2.3%)	1.3% (0.58-2.4%)	19 (8.4-35)	1800 (1100-2800)
Chad	2000	Female	30 to 34	25% (19-32%)	75% (68-81%)	0.8 (0.58-1.1)	1.7% (1-2.7%)	1.7% (0.88-3%)	24 (13-42)	1800 (1100-2800)
Chad	2000	Female	35 to 39	27% (21-33%)	73% (67-79%)	0.88 (0.63-1.2)	2.1% (1.2-3.2%)	2.4% (1.2-4%)	33 (17-53)	2000 (1200-3100)
Chad	2000	Female	40 to 44	29% (23-35%)	71% (65-77%)	0.9 (0.61-1.2)	2.3% (1.3-3.5%)	2.8% (1.5-4.4%)	37 (20-58)	1900 (1100-2900)
Chad	2000	Female	45 to 49	31% (24-37%)	69% (63-76%)	0.86 (0.6-1.2)	2.6% (1.5-4%)	3.3% (1.8-5.5%)	45 (25-71)	2000 (1100-3100)
Chad	2000	Female	50 to 54	27% (21-34%)	73% (66-79%)	0.8 (0.53-1.1)	2.8% (1.5-4.2%)	3.6% (1.9-5.6%)	45 (24-69)	1700 (940-2600)
Chad	2000	Female	55 to 59	23% (17-30%)	77% (70-83%)	0.78 (0.5-1.1)	2.9% (1.7-4.6%)	3.8% (2-6%)	54 (29-87)	1700 (970-2800)
Chad	2000	Female	60 to 64	25% (19-32%)	75% (68-81%)	0.86 (0.56-1.3)	2.9% (1.5-4.6%)	3.6% (1.9-5.7%)	62 (33-100)	1700 (910-2700)
Chad	2000	Female	65 to 69	28% (22-35%)	72% (65-78%)	0.86 (0.58-1.2)	2.4% (1-4.1%)	2.9% (1.2-5.1%)	60 (26-110)	1400 (580-2400)
Chad	2000	Female	70 to 74	25% (19-33%)	75% (67-81%)	0.68 (0.42-1)	1.8% (0.42-3.7%)	2.2% (0.48-4.6%)	50 (10-100)	900 (200-1900)
Chad	2000	Female	75 to 79	20% (14-29%)	80% (71-86%)	0.49 (0.28-0.79)	1% (0.023-2.5%)	1.2% (-0.012-3%)	25 (-0.24-62)	360 (8.4-860)
Chad	2000	Female	80 to 84	19% (12-25%)	81% (75-88%)	0.43 (0.26-0.66)	0.66% (-0.16-1.8%)	0.77% (-0.25-2.1%)	11 (-3.4-30)	120 (-27-310)
Chad	2000	Female	85 to 89	19% (12-26%)	81% (74-88%)	0.42 (0.25-0.66)	0.41% (-0.35-1.4%)	0.47% (-0.46-1.7%)	4.2 (-4.1-15)	32 (-27-110)
Chad	2000	Female	90 to 94	20% (14-28%)	80% (72-86%)	0.43 (0.24-0.7)	0.21% (-0.57-1.3%)	0.2% (-0.77-1.5%)	0.73 (-2.9-5.7)	4.7 (-13-29)
Chad	2000	Female	95 plus	22% (14-31%)	78% (69-86%)	0.45 (0.22-0.77)	0.12% (-0.75-1.2%)	0.034% (-1-1.4%)	0.039 (-1.2-1.5)	0.53 (-3.3-5.5)
Chad	2000	Male	All Ages	19% (16-22%)	81% (78-84%)	0.79 (0.74-0.84)	1.5% (1.2-1.8%)	2.4% (1.9-3%)	1400 (1100-1800)	63000 (50000-77000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	2000	Male	15 to 19	15% (10-20%)	85% (80-90%)	0.45 (0.28-0.66)	1.4% (0.85-2.1%)	1.5% (0.78-2.5%)	15 (7.9-26)	1500 (910-2300)
Chad	2000	Male	20 to 24	28% (20-36%)	72% (64-80%)	1.2 (0.86-1.7)	4.2% (2.9-6%)	4.6% (2.9-6.9%)	66 (41-98)	5200 (3500-7400)
Chad	2000	Male	25 to 29	36% (27-43%)	64% (57-73%)	2 (1.4-2.6)	6.3% (4.3-8.8%)	7.2% (4.5-11%)	120 (76-160)	7900 (5500-11000)
Chad	2000	Male	30 to 34	38% (28-45%)	62% (55-72%)	2.3 (1.6-2.9)	7% (4.8-9.5%)	8.1% (5.3-11%)	130 (93-190)	8100 (5800-11000)
Chad	2000	Male	35 to 39	37% (28-44%)	63% (56-72%)	2.2 (1.5-2.9)	7% (4.9-9.7%)	8.3% (5.5-12%)	140 (97-190)	7500 (5400-10000)
Chad	2000	Male	40 to 44	37% (30-44%)	63% (56-70%)	2.1 (1.5-2.8)	7.7% (5.2-10%)	9.2% (6.1-13%)	160 (110-210)	7500 (5200-10000)
Chad	2000	Male	45 to 49	37% (29-44%)	63% (56-71%)	2.1 (1.5-2.8)	8.6% (5.9-12%)	10% (7.1-14%)	180 (120-240)	7500 (5100-10000)
Chad	2000	Male	50 to 54	35% (27-43%)	65% (57-73%)	1.9 (1.3-2.7)	8.4% (5.6-12%)	10% (6.7-14%)	160 (100-230)	6000 (3900-8400)
Chad	2000	Male	55 to 59	30% (23-37%)	70% (63-77%)	1.5 (1-2.2)	6.8% (4.4-10%)	8.3% (5.3-12%)	140 (90-210)	4600 (3000-6800)
Chad	2000	Male	60 to 64	26% (18-34%)	74% (66-82%)	1.2 (0.72-1.9)	4.8% (2.8-7.5%)	5.7% (3.3-9%)	110 (63-180)	3000 (1800-4900)
Chad	2000	Male	65 to 69	28% (21-36%)	72% (64-79%)	1.1 (0.72-1.6)	3.9% (2.2-5.9%)	4.6% (2.6-7.1%)	93 (53-150)	2100 (1200-3300)
Chad	2000	Male	70 to 74	28% (22-36%)	72% (64-78%)	0.88 (0.57-1.3)	2.8% (1.3-4.8%)	3.3% (1.5-5.6%)	70 (31-120)	1300 (580-2100)
Chad	2000	Male	75 to 79	23% (16-32%)	77% (68-84%)	0.61 (0.36-0.98)	1.7% (0.55-3.3%)	2% (0.59-3.9%)	37 (11-72)	520 (170-1000)
Chad	2000	Male	80 to 84	21% (14-28%)	79% (72-86%)	0.48 (0.29-0.74)	1.3% (0.33-2.4%)	1.4% (0.33-2.8%)	17 (3.8-32)	180 (46-330)
Chad	2000	Male	85 to 89	20% (13-28%)	80% (72-87%)	0.4 (0.24-0.62)	0.86% (0.13-1.7%)	0.96% (0.07-2%)	6.2 (0.47-13)	48 (7.1-96)
Chad	2000	Male	90 to 94	21% (14-29%)	79% (71-86%)	0.35 (0.2-0.56)	0.4% (-0.24-1.1%)	0.39% (-0.39-1.3%)	0.95 (-0.96-3)	5.9 (-3.5-16)
Chad	2000	Male	95 plus	23% (15-33%)	77% (67-85%)	0.33 (0.15-0.57)	0.31% (-0.51-1.2%)	0.24% (-0.78-1.3%)	0.15 (-0.5-0.83)	0.8 (-1.3-3)
Chad	2005	Female	All Ages	15% (14-17%)	85% (83-86%)	0.37 (0.33-0.41)	0.53% (0.37-0.72%)	0.93% (0.55-1.4%)	570 (330-870)	23000 (16000-31000)
Chad	2005	Female	15 to 19	15% (11-19%)	85% (81-89%)	0.31 (0.2-0.45)	0.68% (0.4-1%)	0.49% (0.13-1%)	4.8 (1.2-11)	820 (480-1300)
Chad	2005	Female	20 to 24	19% (15-24%)	81% (76-85%)	0.5 (0.34-0.74)	1.5% (0.87-2.3%)	1.1% (0.43-2.3%)	15 (5.4-31)	1900 (1100-3000)
Chad	2005	Female	25 to 29	22% (17-28%)	78% (72-83%)	0.71 (0.5-0.94)	1.5% (0.89-2.3%)	1.2% (0.56-2.3%)	21 (10-38)	2100 (1300-3200)
Chad	2005	Female	30 to 34	25% (19-31%)	75% (69-81%)	0.89 (0.65-1.2)	1.5% (0.96-2.3%)	1.4% (0.75-2.5%)	27 (14-45)	2100 (1300-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	2005	Female	35 to 39	27% (21-33%)	73% (67-79%)	0.99 (0.69-1.3)	1.8% (1.1-2.8%)	2% (1.1-3.3%)	35 (20-56)	2200 (1400-3200)
Chad	2005	Female	40 to 44	29% (22-35%)	71% (65-78%)	1 (0.68-1.4)	2% (1.3-3.1%)	2.4% (1.4-3.8%)	40 (23-61)	2100 (1300-3100)
Chad	2005	Female	45 to 49	31% (23-36%)	69% (64-77%)	0.96 (0.68-1.3)	2.5% (1.5-3.7%)	3.1% (1.8-4.8%)	49 (29-76)	2200 (1300-3200)
Chad	2005	Female	50 to 54	27% (21-34%)	73% (66-79%)	0.88 (0.6-1.2)	2.8% (1.6-4.2%)	3.6% (2.1-5.4%)	60 (33-91)	2300 (1300-3400)
Chad	2005	Female	55 to 59	23% (17-30%)	77% (70-83%)	0.86 (0.56-1.3)	3.1% (1.7-4.8%)	4% (2.2-6.1%)	61 (33-100)	2000 (1100-3200)
Chad	2005	Female	60 to 64	25% (18-32%)	75% (68-82%)	0.96 (0.63-1.4)	3% (1.7-4.8%)	3.8% (2.1-6.1%)	71 (38-120)	1900 (1100-3200)
Chad	2005	Female	65 to 69	28% (21-35%)	72% (65-79%)	0.96 (0.64-1.4)	2.6% (1.2-4.4%)	3.3% (1.5-5.4%)	71 (31-120)	1600 (710-2800)
Chad	2005	Female	70 to 74	25% (19-33%)	75% (67-81%)	0.75 (0.46-1.1)	2% (0.55-3.9%)	2.4% (0.65-4.9%)	59 (15-120)	1100 (280-2100)
Chad	2005	Female	75 to 79	20% (14-29%)	80% (71-86%)	0.54 (0.31-0.87)	1.2% (0.13-2.8%)	1.4% (0.13-3.4%)	31 (2.8-74)	450 (47-1000)
Chad	2005	Female	80 to 84	18% (12-25%)	82% (75-88%)	0.47 (0.28-0.73)	0.8% (-0.055-2%)	0.94% (-0.1-2.4%)	15 (-1.6-38)	160 (-11-400)
Chad	2005	Female	85 to 89	19% (13-26%)	81% (74-87%)	0.47 (0.28-0.74)	0.55% (-0.22-1.6%)	0.62% (-0.31-1.9%)	6.2 (-3-19)	48 (-19-140)
Chad	2005	Female	90 to 94	20% (13-28%)	80% (72-87%)	0.48 (0.26-0.76)	0.33% (-0.5-1.4%)	0.34% (-0.67-1.7%)	1.4 (-2.7-6.8)	8.1 (-12-35)
Chad	2005	Female	95 plus	22% (14-31%)	78% (69-86%)	0.49 (0.24-0.83)	0.23% (-0.65-1.4%)	0.17% (-0.88-1.6%)	0.2 (-1.1-1.9)	1.1 (-3-6.7)
Chad	2005	Male	All Ages	18% (15-21%)	82% (79-85%)	0.85 (0.79-0.9)	1.5% (1.2-1.9%)	2.5% (2-3.2%)	1700 (1300-2100)	73000 (58000-89000)
Chad	2005	Male	15 to 19	15% (11-20%)	85% (80-89%)	0.48 (0.31-0.71)	1.5% (0.95-2.2%)	1.7% (0.92-2.7%)	18 (10-30)	1800 (1100-2700)
Chad	2005	Male	20 to 24	28% (20-35%)	72% (65-80%)	1.3 (0.94-1.8)	4.7% (3.2-6.4%)	5.4% (3.5-7.9%)	77 (49-110)	6100 (4200-8500)
Chad	2005	Male	25 to 29	36% (27-43%)	64% (57-73%)	2.2 (1.6-2.8)	6.7% (4.6-9.3%)	7.9% (5.1-12%)	140 (90-190)	9300 (6400-13000)
Chad	2005	Male	30 to 34	37% (28-44%)	63% (56-72%)	2.5 (1.8-3.1)	7% (5-9.4%)	8.1% (5.4-11%)	160 (110-220)	9700 (6900-13000)
Chad	2005	Male	35 to 39	37% (30-43%)	63% (57-70%)	2.3 (1.6-3.1)	6.9% (4.9-9.4%)	8.1% (5.5-11%)	160 (110-210)	8500 (6100-11000)
Chad	2005	Male	40 to 44	37% (29-44%)	63% (56-71%)	2.3 (1.6-3)	7.3% (5.2-9.9%)	8.7% (6-12%)	170 (120-230)	8200 (5800-11000)
Chad	2005	Male	45 to 49	37% (28-44%)	63% (56-72%)	2.3 (1.6-3)	8.5% (6-11%)	10% (7.2-14%)	200 (140-270)	8400 (5900-11000)
Chad	2005	Male	50 to 54	35% (27-42%)	65% (58-73%)	2.1 (1.5-2.9)	8.6% (5.9-12%)	10% (7.2-14%)	200 (130-280)	7400 (5000-10000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	2005	Male	55 to 59	29% (22-37%)	71% (63-78%)	1.7 (1.1-2.4)	7.2% (4.8-10%)	8.8% (5.9-13%)	160 (100-230)	5100 (3400-7400)
Chad	2005	Male	60 to 64	25% (18-34%)	75% (66-82%)	1.3 (0.77-2)	5.1% (3.2-7.9%)	6.1% (3.7-9.5%)	130 (78-200)	3500 (2200-5600)
Chad	2005	Male	65 to 69	28% (21-35%)	72% (65-79%)	1.2 (0.76-1.8)	4.1% (2.5-6.2%)	4.9% (2.9-7.5%)	110 (64-170)	2500 (1500-3900)
Chad	2005	Male	70 to 74	28% (21-35%)	72% (65-79%)	0.96 (0.63-1.4)	3% (1.4-4.9%)	3.5% (1.7-5.8%)	81 (38-130)	1500 (710-2400)
Chad	2005	Male	75 to 79	23% (16-32%)	77% (68-84%)	0.66 (0.4-1.1)	1.8% (0.66-3.4%)	2.1% (0.73-4%)	43 (15-80)	610 (220-1100)
Chad	2005	Male	80 to 84	20% (14-27%)	80% (73-86%)	0.52 (0.31-0.79)	1.4% (0.47-2.5%)	1.6% (0.49-3%)	21 (6.5-39)	220 (77-410)
Chad	2005	Male	85 to 89	20% (13-27%)	80% (73-87%)	0.43 (0.25-0.67)	0.96% (0.23-1.8%)	1.1% (0.2-2.2%)	7.9 (1.5-16)	61 (15-120)
Chad	2005	Male	90 to 94	20% (14-28%)	80% (72-86%)	0.38 (0.22-0.61)	0.48% (-0.14-1.2%)	0.48% (-0.29-1.3%)	1.3 (-0.81-3.8)	8 (-2.4-20)
Chad	2005	Male	95 plus	23% (15-33%)	77% (67-85%)	0.36 (0.17-0.63)	0.41% (-0.39-1.3%)	0.35% (-0.65-1.5%)	0.24 (-0.45-1)	1.1 (-1.1-3.7)
Chad	2010	Female	All Ages	12% (11-14%)	88% (86-89%)	0.36 (0.32-0.4)	0.59% (0.44-0.78%)	1.1% (0.71-1.5%)	690 (440-1000)	27000 (20000-35000)
Chad	2010	Female	15 to 19	11% (8.7-14%)	89% (86-91%)	0.27 (0.18-0.4)	0.67% (0.42-0.97%)	0.44% (0.17-0.89%)	5.2 (1.8-11)	950 (600-1400)
Chad	2010	Female	20 to 24	15% (12-19%)	85% (81-88%)	0.46 (0.32-0.69)	1.5% (0.99-2.2%)	1.1% (0.5-2.1%)	16 (6.7-30)	2200 (1400-3400)
Chad	2010	Female	25 to 29	18% (15-22%)	82% (78-85%)	0.7 (0.53-0.91)	1.6% (1-2.3%)	1.3% (0.65-2.2%)	22 (12-37)	2400 (1600-3300)
Chad	2010	Female	30 to 34	22% (18-25%)	78% (75-82%)	0.92 (0.71-1.2)	1.6% (1.1-2.3%)	1.5% (0.89-2.5%)	29 (18-46)	2300 (1600-3400)
Chad	2010	Female	35 to 39	24% (20-28%)	76% (72-80%)	1 (0.76-1.4)	1.9% (1.2-2.6%)	2% (1.3-3.1%)	38 (24-56)	2400 (1700-3300)
Chad	2010	Female	40 to 44	23% (19-28%)	77% (72-81%)	0.96 (0.68-1.3)	2.1% (1.4-2.9%)	2.4% (1.6-3.6%)	40 (26-59)	2200 (1500-3000)
Chad	2010	Female	45 to 49	22% (18-28%)	78% (72-82%)	0.84 (0.6-1.2)	2.5% (1.6-3.6%)	3.1% (2-4.7%)	49 (32-73)	2200 (1500-3200)
Chad	2010	Female	50 to 54	23% (19-28%)	77% (72-81%)	0.92 (0.65-1.3)	3.2% (2-4.6%)	4.2% (2.6-6.3%)	70 (43-100)	2600 (1700-3900)
Chad	2010	Female	55 to 59	22% (18-27%)	78% (73-82%)	1 (0.71-1.4)	3.7% (2.3-5.7%)	4.9% (3-7.5%)	92 (54-140)	3000 (1800-4600)
Chad	2010	Female	60 to 64	20% (16-26%)	80% (74-84%)	0.95 (0.64-1.4)	3.4% (2.1-5.2%)	4.3% (2.6-6.5%)	82 (48-130)	2200 (1300-3500)
Chad	2010	Female	65 to 69	23% (19-29%)	77% (71-81%)	0.97 (0.67-1.4)	3.1% (1.7-4.7%)	3.8% (2.1-5.9%)	87 (48-140)	2000 (1100-3200)
Chad	2010	Female	70 to 74	23% (18-29%)	77% (71-82%)	0.81 (0.52-1.2)	2.4% (0.98-4.4%)	3% (1.1-5.3%)	76 (30-140)	1400 (540-2500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	2010	Female	75 to 79	19% (14-25%)	81% (75-86%)	0.6 (0.36-0.92)	1.6% (0.37-3.2%)	1.9% (0.43-3.9%)	44 (9.8-90)	620 (150-1300)
Chad	2010	Female	80 to 84	18% (12-24%)	82% (76-88%)	0.55 (0.33-0.84)	1.1% (0.17-2.4%)	1.3% (0.18-2.9%)	23 (3.1-52)	240 (38-540)
Chad	2010	Female	85 to 89	19% (12-26%)	81% (74-88%)	0.56 (0.32-0.85)	0.85% (-0.0008-2%)	0.99% (-0.057-2.4%)	11 (-0.65-28)	85 (-0.079-200)
Chad	2010	Female	90 to 94	20% (14-28%)	80% (72-86%)	0.57 (0.31-0.91)	0.62% (-0.27-1.9%)	0.68% (-0.4-2.2%)	3.2 (-1.9-10)	17 (-7.8-54)
Chad	2010	Female	95 plus	22% (14-31%)	78% (69-86%)	0.59 (0.29-1)	0.5% (-0.52-1.9%)	0.49% (-0.75-2.2%)	0.65 (-1-2.9)	2.7 (-2.8-10)
Chad	2010	Male	All Ages	15% (14-17%)	85% (83-86%)	0.88 (0.82-0.94)	1.8% (1.4-2.2%)	2.9% (2.3-3.6%)	2000 (1600-2500)	88000 (72000-110000)
Chad	2010	Male	15 to 19	11% (8.3-13%)	89% (87-92%)	0.42 (0.27-0.6)	1.4% (0.89-1.9%)	1.4% (0.84-2.3%)	19 (11-30)	2000 (1300-2800)
Chad	2010	Male	20 to 24	22% (18-26%)	78% (74-82%)	1.3 (0.93-1.7)	4.6% (3.2-6.5%)	5.3% (3.6-7.6%)	84 (55-120)	6800 (4800-9400)
Chad	2010	Male	25 to 29	31% (26-35%)	69% (65-74%)	2.3 (1.7-2.8)	7.4% (5.2-9.8%)	8.9% (5.9-12%)	160 (110-210)	11000 (7800-14000)
Chad	2010	Male	30 to 34	34% (28-39%)	66% (61-72%)	2.7 (2.1-3.4)	8.1% (5.8-10%)	9.6% (6.7-13%)	190 (140-260)	12000 (8700-15000)
Chad	2010	Male	35 to 39	33% (29-38%)	67% (62-71%)	2.6 (1.9-3.3)	7.9% (5.8-11%)	9.5% (6.7-13%)	200 (140-260)	11000 (8000-14000)
Chad	2010	Male	40 to 44	33% (28-38%)	67% (62-72%)	2.5 (1.8-3.2)	8.5% (6.1-11%)	10% (7.4-14%)	200 (150-270)	9800 (7100-13000)
Chad	2010	Male	45 to 49	30% (26-35%)	70% (65-74%)	2.3 (1.7-3)	9.4% (6.9-12%)	12% (8.4-15%)	220 (160-310)	9500 (6900-13000)
Chad	2010	Male	50 to 54	27% (22-32%)	73% (68-78%)	2 (1.4-2.8)	8.8% (6.2-12%)	11% (7.7-15%)	220 (150-290)	8000 (5600-11000)
Chad	2010	Male	55 to 59	28% (24-33%)	72% (67-76%)	1.9 (1.4-2.7)	8.5% (5.8-12%)	11% (7.3-15%)	220 (150-300)	7000 (4800-9800)
Chad	2010	Male	60 to 64	27% (22-32%)	73% (68-78%)	1.6 (1.1-2.4)	6.6% (4.3-9.7%)	8% (5.1-12%)	170 (110-260)	4700 (3000-7000)
Chad	2010	Male	65 to 69	26% (21-31%)	74% (69-79%)	1.3 (0.93-1.9)	4.9% (3.1-7.2%)	5.8% (3.8-8.7%)	140 (88-210)	3200 (2000-4700)
Chad	2010	Male	70 to 74	23% (19-29%)	77% (71-81%)	0.96 (0.65-1.4)	3.3% (1.8-5.1%)	3.9% (2.2-6.1%)	99 (52-150)	1800 (990-2800)
Chad	2010	Male	75 to 79	20% (15-26%)	80% (74-85%)	0.68 (0.42-1.1)	2.1% (0.95-3.7%)	2.4% (1.1-4.3%)	54 (23-95)	760 (340-1400)
Chad	2010	Male	80 to 84	19% (13-25%)	81% (75-87%)	0.59 (0.36-0.9)	1.7% (0.7-2.8%)	1.9% (0.76-3.3%)	28 (11-48)	290 (120-500)
Chad	2010	Male	85 to 89	19% (12-25%)	81% (75-88%)	0.5 (0.29-0.76)	1.2% (0.47-2.1%)	1.4% (0.46-2.5%)	11 (3.8-21)	88 (34-160)
Chad	2010	Male	90 to 94	19% (13-27%)	81% (73-87%)	0.44 (0.25-0.69)	0.66% (0.059-1.4%)	0.69% (-0.058-1.7%)	2.2 (-0.18-5.2)	13 (1.1-28)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	2010	Male	95 plus	22% (14-31%)	78% (69-86%)	0.42 (0.2-0.72)	0.58% (-0.17-1.6%)	0.55% (-0.39-1.8%)	0.43 (-0.3-1.4)	1.9 (-0.54-5.1)
Chad	2016	Female	All Ages	12% (11-14%)	88% (86-89%)	0.35 (0.24-0.47)	0.6% (0.39-0.87%)	1.1% (0.57-1.8%)	690 (370-1100)	28000 (19000-41000)
Chad	2016	Female	15 to 19	11% (8.6-14%)	89% (86-91%)	0.26 (0.15-0.43)	0.65% (0.4-0.99%)	0.39% (0.13-0.83%)	5.3 (1.6-12)	1100 (650-1700)
Chad	2016	Female	20 to 24	15% (12-18%)	85% (82-88%)	0.45 (0.27-0.73)	1.5% (0.95-2.3%)	0.99% (0.37-2.1%)	15 (5.4-32)	2500 (1600-4000)
Chad	2016	Female	25 to 29	18% (15-22%)	82% (78-85%)	0.68 (0.42-1)	1.6% (1-2.3%)	1.3% (0.61-2.3%)	22 (11-39)	2700 (1700-3800)
Chad	2016	Female	30 to 34	21% (18-25%)	79% (75-82%)	0.89 (0.57-1.3)	1.7% (1.1-2.4%)	1.5% (0.81-2.6%)	29 (16-49)	2600 (1700-3900)
Chad	2016	Female	35 to 39	23% (20-27%)	77% (73-80%)	0.99 (0.64-1.4)	1.9% (1.2-2.8%)	2% (1.2-3.2%)	40 (23-63)	2600 (1700-3900)
Chad	2016	Female	40 to 44	23% (19-27%)	77% (73-81%)	0.92 (0.56-1.4)	2% (1.3-3%)	2.5% (1.5-3.8%)	43 (26-67)	2400 (1500-3500)
Chad	2016	Female	45 to 49	22% (18-27%)	78% (73-82%)	0.8 (0.49-1.2)	2.4% (1.5-3.7%)	3.1% (1.8-5%)	49 (28-79)	2300 (1400-3500)
Chad	2016	Female	50 to 54	23% (19-28%)	77% (72-81%)	0.88 (0.53-1.3)	3.1% (1.7-4.9%)	4.1% (2.3-6.7%)	67 (37-110)	2600 (1500-4100)
Chad	2016	Female	55 to 59	22% (18-26%)	78% (74-82%)	0.96 (0.6-1.4)	3.5% (1.9-5.9%)	4.8% (2.6-8.2%)	88 (47-150)	2900 (1600-4900)
Chad	2016	Female	60 to 64	20% (16-25%)	80% (75-84%)	0.9 (0.54-1.4)	3.1% (1.7-5.2%)	4% (2.1-6.7%)	94 (49-160)	2600 (1400-4400)
Chad	2016	Female	65 to 69	23% (19-28%)	77% (72-81%)	0.92 (0.57-1.4)	2.8% (1.2-5.2%)	3.5% (1.5-6.5%)	85 (35-160)	2000 (840-3600)
Chad	2016	Female	70 to 74	22% (18-28%)	78% (72-82%)	0.77 (0.45-1.3)	2.2% (0.62-4.5%)	2.7% (0.75-5.4%)	71 (20-150)	1300 (360-2700)
Chad	2016	Female	75 to 79	18% (14-24%)	82% (76-86%)	0.57 (0.31-0.98)	1.4% (0.16-3.2%)	1.7% (0.19-3.9%)	42 (4.5-98)	610 (78-1400)
Chad	2016	Female	80 to 84	18% (12-24%)	82% (76-88%)	0.53 (0.29-0.92)	0.99% (0.0095-2.5%)	1.2% (-0.023-3%)	22 (-0.41-57)	240 (2.2-600)
Chad	2016	Female	85 to 89	19% (12-27%)	81% (73-88%)	0.54 (0.29-0.92)	0.72% (-0.19-2.1%)	0.83% (-0.28-2.5%)	11 (-3.5-31)	82 (-21-230)
Chad	2016	Female	90 to 94	20% (13-28%)	80% (72-87%)	0.55 (0.29-0.94)	0.5% (-0.4-1.9%)	0.54% (-0.57-2.2%)	3 (-3-12)	17 (-13-62)
Chad	2016	Female	95 plus	22% (14-31%)	78% (69-86%)	0.57 (0.25-1)	0.4% (-0.58-1.8%)	0.36% (-0.83-2.1%)	0.58 (-1.4-3.3)	2.6 (-3.8-12)
Chad	2016	Male	All Ages	15% (14-17%)	85% (83-86%)	0.86 (0.6-1.2)	1.8% (1.3-2.5%)	3% (2-4.2%)	2200 (1500-3000)	96000 (68000-130000)
Chad	2016	Male	15 to 19	10% (8.5-13%)	90% (87-92%)	0.41 (0.23-0.65)	1.3% (0.82-1.9%)	1.3% (0.68-2.3%)	20 (10-35)	2200 (1400-3400)
Chad	2016	Male	20 to 24	22% (18-26%)	78% (74-82%)	1.2 (0.78-1.9)	4.5% (3-6.5%)	5.1% (3.1-7.9%)	91 (53-140)	7700 (5000-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Chad	2016	Male	25 to 29	30% (25-35%)	70% (65-75%)	2.2 (1.4-3.2)	7.3% (4.8-10%)	8.9% (5.5-13%)	170 (100-250)	12000 (7900-17000)
Chad	2016	Male	30 to 34	34% (28-38%)	66% (62-72%)	2.6 (1.7-3.8)	8.1% (5.6-11%)	9.9% (6.4-14%)	210 (140-290)	13000 (8800-17000)
Chad	2016	Male	35 to 39	33% (29-37%)	67% (63-71%)	2.5 (1.6-3.6)	8% (5.5-11%)	9.7% (6.5-14%)	220 (150-300)	12000 (8400-16000)
Chad	2016	Male	40 to 44	33% (28-38%)	67% (62-72%)	2.4 (1.5-3.6)	8.3% (5.5-12%)	10% (6.8-14%)	230 (150-320)	11000 (7500-15000)
Chad	2016	Male	45 to 49	30% (26-35%)	70% (65-74%)	2.2 (1.4-3.3)	9.1% (5.9-13%)	11% (7.3-16%)	240 (150-350)	10000 (6300-15000)
Chad	2016	Male	50 to 54	27% (22-33%)	73% (67-78%)	1.9 (1.2-3)	8.4% (5.5-12%)	11% (6.7-15%)	220 (140-310)	8200 (5200-12000)
Chad	2016	Male	55 to 59	28% (23-33%)	72% (67-77%)	1.9 (1.1-2.9)	8.2% (5.1-12%)	10% (6.3-15%)	220 (140-340)	7200 (4400-11000)
Chad	2016	Male	60 to 64	27% (22-33%)	73% (67-78%)	1.6 (0.92-2.6)	6.3% (3.7-9.9%)	7.7% (4.4-12%)	200 (110-320)	5400 (3100-8700)
Chad	2016	Male	65 to 69	26% (21-31%)	74% (69-79%)	1.3 (0.8-2)	4.7% (2.6-7.7%)	5.6% (3.2-9.3%)	140 (79-240)	3300 (1800-5500)
Chad	2016	Male	70 to 74	23% (18-29%)	77% (71-82%)	0.93 (0.56-1.5)	3.1% (1.6-5.5%)	3.8% (1.9-6.7%)	100 (50-190)	1900 (940-3400)
Chad	2016	Male	75 to 79	20% (15-26%)	80% (74-85%)	0.66 (0.38-1.1)	2% (0.8-3.7%)	2.3% (0.9-4.4%)	58 (22-110)	820 (320-1600)
Chad	2016	Male	80 to 84	19% (13-26%)	81% (74-87%)	0.57 (0.31-0.95)	1.6% (0.64-2.8%)	1.8% (0.69-3.3%)	29 (11-53)	310 (120-560)
Chad	2016	Male	85 to 89	19% (12-26%)	81% (74-88%)	0.48 (0.27-0.81)	1.1% (0.37-2.2%)	1.3% (0.35-2.6%)	12 (3.4-25)	94 (31-190)
Chad	2016	Male	90 to 94	19% (13-27%)	81% (73-87%)	0.42 (0.22-0.72)	0.61% (-0.013-1.5%)	0.62% (-0.15-1.7%)	2.4 (-0.56-6.3)	14 (-0.3-34)
Chad	2016	Male	95 plus	22% (14-31%)	78% (69-86%)	0.4 (0.17-0.74)	0.51% (-0.22-1.5%)	0.46% (-0.44-1.7%)	0.44 (-0.42-1.6)	2 (-0.84-5.9)
Cote d'Ivoire	1990	Female	All Ages	16% (14-19%)	84% (81-86%)	0.45 (0.31-0.62)	0.82% (0.51-1.2%)	1.5% (0.77-2.5%)	930 (470-1500)	36000 (23000-53000)
Cote d'Ivoire	1990	Female	15 to 19	16% (11-22%)	84% (78-89%)	0.46 (0.24-0.77)	0.94% (0.51-1.6%)	0.94% (0.33-1.9%)	11 (3.7-22)	1300 (740-2200)
Cote d'Ivoire	1990	Female	20 to 24	22% (14-30%)	78% (70-86%)	0.72 (0.37-1.2)	1.8% (1.1-2.9%)	1.6% (0.64-3.1%)	23 (9.4-44)	2700 (1600-4200)
Cote d'Ivoire	1990	Female	25 to 29	23% (15-32%)	77% (68-85%)	0.75 (0.4-1.2)	1.9% (1.1-3%)	1.8% (0.81-3.4%)	26 (12-50)	2600 (1600-4200)
Cote d'Ivoire	1990	Female	30 to 34	25% (16-37%)	75% (63-84%)	0.82 (0.43-1.4)	2% (1.3-3.2%)	2.2% (1.1-3.9%)	32 (16-57)	2500 (1600-4000)
Cote d'Ivoire	1990	Female	35 to 39	29% (19-39%)	71% (61-81%)	0.93 (0.5-1.5)	2.4% (1.4-4%)	3% (1.6-5.3%)	46 (24-80)	2800 (1700-4500)
Cote d'Ivoire	1990	Female	40 to 44	30% (20-40%)	70% (60-80%)	1 (0.55-1.6)	3.1% (1.8-4.7%)	4.2% (2.4-6.4%)	65 (37-100)	3300 (2000-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	1990	Female	45 to 49	29% (18-39%)	71% (61-82%)	1.1 (0.57-1.8)	3.5% (1.9-5.8%)	4.7% (2.4-8.1%)	83 (43-140)	3600 (2000-6200)
Cote d'Ivoire	1990	Female	50 to 54	28% (20-39%)	72% (61-80%)	1.1 (0.62-1.9)	4% (2.1-6.9%)	5.3% (2.8-9%)	110 (56-180)	4100 (2200-7000)
Cote d'Ivoire	1990	Female	55 to 59	29% (20-39%)	71% (61-80%)	1.2 (0.65-2.1)	4.2% (2-7.4%)	5.4% (2.6-9.5%)	120 (57-220)	4000 (1900-7000)
Cote d'Ivoire	1990	Female	60 to 64	30% (20-42%)	70% (58-80%)	1.2 (0.63-2.1)	3.7% (1.6-6.9%)	4.6% (2-8.6%)	120 (51-230)	3300 (1400-6200)
Cote d'Ivoire	1990	Female	65 to 69	32% (22-42%)	68% (58-78%)	1.2 (0.66-2)	3.3% (1.1-6.3%)	4% (1.3-7.6%)	110 (38-220)	2600 (850-5000)
Cote d'Ivoire	1990	Female	70 to 74	29% (20-39%)	71% (61-80%)	0.95 (0.5-1.6)	2.8% (0.64-6%)	3.3% (0.73-7%)	95 (21-200)	1700 (380-3600)
Cote d'Ivoire	1990	Female	75 to 79	24% (16-34%)	76% (66-84%)	0.71 (0.35-1.3)	1.7% (-0.02-4.2%)	2% (-0.045-5%)	49 (-1.1-130)	700 (-7.7-1800)
Cote d'Ivoire	1990	Female	80 to 84	22% (14-31%)	78% (69-86%)	0.62 (0.32-1)	1.1% (-0.1-2.8%)	1.3% (-0.15-3.3%)	21 (-2.5-57)	230 (-22-600)
Cote d'Ivoire	1990	Female	85 to 89	24% (15-35%)	76% (65-85%)	0.64 (0.31-1.1)	0.74% (-0.34-2.3%)	0.84% (-0.46-2.8%)	7.7 (-4.1-26)	59 (-27-190)
Cote d'Ivoire	1990	Female	90 to 94	27% (18-39%)	73% (61-82%)	0.69 (0.33-1.2)	0.43% (-0.7-2.2%)	0.45% (-0.94-2.5%)	1.4 (-3-7.9)	8.4 (-13-42)
Cote d'Ivoire	1990	Female	95 plus	32% (18-48%)	68% (52-82%)	0.77 (0.3-1.5)	0.27% (-1-2.4%)	0.19% (-1.4-2.7%)	0.15 (-1.1-2.1)	0.85 (-3.2-7.5)
Cote d'Ivoire	1990	Male	All Ages	29% (24-34%)	71% (66-76%)	1.3 (0.93-1.8)	2.6% (1.8-3.5%)	4.2% (2.8-6%)	3600 (2400-5000)	150000 (100000-200000)
Cote d'Ivoire	1990	Male	15 to 19	22% (14-31%)	78% (69-86%)	0.78 (0.39-1.3)	2% (1.1-4.2%)	2.4% (1.1-4.2%)	32 (15-57)	2900 (1700-4800)
Cote d'Ivoire	1990	Male	20 to 24	35% (24-47%)	65% (53-76%)	1.8 (0.94-2.9)	5.2% (3.2-7.9%)	5.9% (3.3-9.4%)	120 (64-190)	9200 (5500-14000)
Cote d'Ivoire	1990	Male	25 to 29	43% (31-55%)	57% (45-69%)	2.4 (1.4-3.8)	6.8% (4.3-10%)	7.8% (4.5-12%)	200 (120-300)	14000 (8500-20000)
Cote d'Ivoire	1990	Male	30 to 34	47% (34-59%)	53% (41-66%)	2.9 (1.7-4.3)	8% (5.1-11%)	9.3% (5.6-14%)	270 (160-390)	16000 (10000-23000)
Cote d'Ivoire	1990	Male	35 to 39	48% (35-60%)	52% (40-65%)	3.2 (1.8-5)	9.4% (5.9-13%)	11% (6.8-17%)	340 (210-490)	18000 (12000-26000)
Cote d'Ivoire	1990	Male	40 to 44	47% (35-58%)	53% (42-65%)	3.2 (1.8-5.1)	10% (6.6-15%)	13% (7.8-18%)	420 (260-610)	20000 (13000-29000)
Cote d'Ivoire	1990	Male	45 to 49	45% (31-57%)	55% (43-69%)	2.9 (1.7-4.5)	11% (6.7-16%)	13% (8.1-19%)	470 (280-680)	20000 (12000-28000)
Cote d'Ivoire	1990	Male	50 to 54	45% (33-57%)	55% (43-67%)	2.6 (1.6-4.2)	9.7% (5.5-15%)	12% (6.7-18%)	430 (250-660)	16000 (9400-24000)
Cote d'Ivoire	1990	Male	55 to 59	47% (34-59%)	53% (41-66%)	2.5 (1.5-4.2)	8.8% (5-14%)	11% (5.9-17%)	400 (220-620)	13000 (7300-20000)
Cote d'Ivoire	1990	Male	60 to 64	46% (33-60%)	54% (40-67%)	2.3 (1.2-3.9)	7.1% (3.5-12%)	8.4% (4.1-14%)	320 (160-540)	8900 (4400-15000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	1990	Male	65 to 69	45% (33-56%)	55% (44-67%)	2 (1.2-3.3)	5.7% (2.8-9.5%)	6.7% (3.3-11%)	250 (120-410)	5600 (2800-9400)
Cote d'Ivoire	1990	Male	70 to 74	43% (32-55%)	57% (45-68%)	1.7 (0.94-2.8)	4.6% (1.9-8.4%)	5.4% (2.1-9.8%)	180 (76-340)	3400 (1400-6200)
Cote d'Ivoire	1990	Male	75 to 79	40% (28-55%)	60% (45-72%)	1.4 (0.69-2.3)	3.3% (1.1-6.5%)	3.8% (1.3-7.5%)	100 (34-210)	1500 (500-2900)
Cote d'Ivoire	1990	Male	80 to 84	37% (25-50%)	63% (50-75%)	1.1 (0.55-1.8)	2.4% (0.7-4.7%)	2.7% (0.73-5.5%)	45 (12-91)	480 (140-960)
Cote d'Ivoire	1990	Male	85 to 89	34% (22-48%)	66% (52-78%)	0.82 (0.41-1.4)	1.5% (0.31-3.2%)	1.6% (0.26-3.7%)	14 (2.3-32)	110 (24-240)
Cote d'Ivoire	1990	Male	90 to 94	34% (23-48%)	66% (52-77%)	0.67 (0.33-1.2)	0.56% (-0.34-1.8%)	0.52% (-0.54-2%)	1.5 (-1.6-5.7)	9.8 (-5.8-31)
Cote d'Ivoire	1990	Male	95 plus	38% (22-56%)	62% (44-78%)	0.57 (0.22-1.2)	0.17% (-0.8-1.4%)	-0.013% (-1.2-1.5%)	-0.0082 (-0.74-0.89)	0.44 (-2-3.5)
Cote d'Ivoire	1995	Female	All Ages	17% (15-21%)	83% (79-85%)	0.54 (0.37-0.77)	1.1% (0.63-1.6%)	1.9% (0.97-3.2%)	1500 (730-2500)	54000 (33000-82000)
Cote d'Ivoire	1995	Female	15 to 19	17% (11-23%)	83% (77-89%)	0.55 (0.28-0.95)	1.2% (0.63-2%)	1.3% (0.45-2.7%)	18 (6.4-38)	2000 (1100-3500)
Cote d'Ivoire	1995	Female	20 to 24	22% (15-31%)	78% (69-85%)	0.86 (0.44-1.5)	2% (1.1-3.3%)	1.9% (0.76-3.6%)	39 (16-78)	4000 (2300-6600)
Cote d'Ivoire	1995	Female	25 to 29	24% (16-33%)	76% (67-84%)	0.9 (0.49-1.5)	1.8% (1-3.1%)	1.7% (0.73-3.3%)	43 (20-81)	3800 (2300-6100)
Cote d'Ivoire	1995	Female	30 to 34	26% (17-37%)	74% (63-83%)	0.98 (0.5-1.7)	1.9% (1.1-3.2%)	2% (0.95-3.7%)	50 (25-91)	3700 (2200-6000)
Cote d'Ivoire	1995	Female	35 to 39	30% (20-40%)	70% (60-80%)	1.1 (0.59-1.8)	2.4% (1.3-3.9%)	2.8% (1.4-4.8%)	69 (35-120)	4100 (2400-6600)
Cote d'Ivoire	1995	Female	40 to 44	31% (21-41%)	69% (59-79%)	1.2 (0.66-2)	3.1% (1.8-4.8%)	3.8% (2.1-6.1%)	91 (53-140)	4600 (2700-7000)
Cote d'Ivoire	1995	Female	45 to 49	30% (19-40%)	70% (60-81%)	1.3 (0.68-2.2)	3.8% (2-6.4%)	4.9% (2.4-8.3%)	120 (59-210)	5200 (2700-8900)
Cote d'Ivoire	1995	Female	50 to 54	29% (20-40%)	71% (60-80%)	1.3 (0.73-2.2)	4.6% (2.4-7.6%)	5.9% (3-9.8%)	170 (84-280)	6200 (3200-10000)
Cote d'Ivoire	1995	Female	55 to 59	30% (21-40%)	70% (60-79%)	1.4 (0.76-2.4)	5% (2.4-8.5%)	6.3% (3.1-11%)	200 (93-340)	6300 (3100-11000)
Cote d'Ivoire	1995	Female	60 to 64	31% (21-43%)	69% (57-79%)	1.4 (0.74-2.5)	4.6% (2-8.1%)	5.6% (2.4-9.8%)	190 (82-350)	5300 (2300-9700)
Cote d'Ivoire	1995	Female	65 to 69	33% (23-43%)	67% (57-77%)	1.4 (0.74-2.2)	4.2% (1.5-7.6%)	5% (1.8-9.1%)	190 (69-350)	4200 (1600-7700)
Cote d'Ivoire	1995	Female	70 to 74	30% (21-40%)	70% (60-79%)	1.1 (0.57-1.9)	3.5% (1-7.2%)	4.2% (1.2-8.5%)	150 (43-310)	2800 (790-5700)
Cote d'Ivoire	1995	Female	75 to 79	24% (16-36%)	76% (64-84%)	0.82 (0.4-1.4)	2.3% (0.26-5.3%)	2.7% (0.29-6.2%)	83 (9.3-200)	1200 (140-2800)
Cote d'Ivoire	1995	Female	80 to 84	22% (14-32%)	78% (68-86%)	0.71 (0.37-1.2)	1.5% (0.032-3.7%)	1.8% (-0.0017-4.3%)	37 (-0.037-92)	390 (7.9-970)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	1995	Female	85 to 89	24% (15-35%)	76% (65-85%)	0.73 (0.37-1.3)	1.2% (-0.2-3.2%)	1.3% (-0.3-3.7%)	15 (-3.2-42)	110 (-20-310)
Cote d'Ivoire	1995	Female	90 to 94	28% (18-40%)	72% (60-82%)	0.8 (0.38-1.4)	0.86% (-0.53-3%)	0.94% (-0.74-3.6%)	3.7 (-2.8-14)	20 (-12-71)
Cote d'Ivoire	1995	Female	95 plus	33% (19-50%)	67% (50-81%)	0.87 (0.34-1.7)	0.68% (-0.9-3.4%)	0.67% (-1.2-3.9%)	0.65 (-1.2-3.9)	2.7 (-3.5-14)
Cote d'Ivoire	1995	Male	All Ages	30% (25-35%)	70% (65-75%)	1.6 (1.1-2.1)	3.5% (2.4-4.6%)	5.4% (3.6-7.5%)	5900 (3900-8100)	240000 (170000-320000)
Cote d'Ivoire	1995	Male	15 to 19	23% (15-32%)	77% (68-85%)	0.93 (0.47-1.6)	2.7% (1.4-4.3%)	3.3% (1.5-5.6%)	62 (30-110)	5200 (2800-8600)
Cote d'Ivoire	1995	Male	20 to 24	37% (25-48%)	63% (52-75%)	2.1 (1.1-3.4)	6.5% (3.7-10%)	7.5% (3.9-12%)	220 (110-350)	16000 (9100-25000)
Cote d'Ivoire	1995	Male	25 to 29	45% (32-57%)	55% (43-68%)	2.9 (1.6-4.4)	7.7% (4.5-12%)	8.6% (4.8-13%)	340 (200-500)	22000 (14000-32000)
Cote d'Ivoire	1995	Male	30 to 34	48% (35-60%)	52% (40-65%)	3.4 (1.9-5.1)	8.3% (5.2-12%)	9.3% (5.6-14%)	440 (280-620)	26000 (17000-36000)
Cote d'Ivoire	1995	Male	35 to 39	49% (36-62%)	51% (38-64%)	3.8 (2.1-5.9)	9.5% (5.9-14%)	11% (6.5-16%)	540 (340-740)	29000 (18000-39000)
Cote d'Ivoire	1995	Male	40 to 44	49% (36-60%)	51% (40-64%)	3.8 (2.1-6.1)	11% (6.6-15%)	12% (7.5-17%)	640 (410-890)	30000 (19000-42000)
Cote d'Ivoire	1995	Male	45 to 49	46% (32-58%)	54% (42-68%)	3.3 (1.9-5.4)	11% (7.1-16%)	13% (8.2-19%)	720 (450-1000)	30000 (19000-43000)
Cote d'Ivoire	1995	Male	50 to 54	46% (35-58%)	54% (42-65%)	3.1 (1.8-4.9)	11% (6.3-16%)	13% (7.5-19%)	700 (400-1100)	26000 (15000-38000)
Cote d'Ivoire	1995	Male	55 to 59	48% (36-60%)	52% (40-64%)	3 (1.7-4.8)	11% (5.9-16%)	12% (6.8-18%)	670 (370-1000)	21000 (12000-32000)
Cote d'Ivoire	1995	Male	60 to 64	48% (34-60%)	52% (40-66%)	2.7 (1.5-4.4)	8.5% (4.3-14%)	9.9% (4.9-16%)	540 (270-880)	15000 (7500-24000)
Cote d'Ivoire	1995	Male	65 to 69	46% (35-58%)	54% (42-65%)	2.4 (1.4-3.8)	7% (3.4-11%)	8.1% (4-13%)	420 (200-700)	9600 (4700-16000)
Cote d'Ivoire	1995	Male	70 to 74	45% (34-56%)	55% (44-66%)	2 (1.1-3.2)	5.8% (2.4-10%)	6.7% (2.8-12%)	320 (130-570)	5800 (2500-10000)
Cote d'Ivoire	1995	Male	75 to 79	42% (29-55%)	58% (45-71%)	1.6 (0.81-2.7)	4.3% (1.5-8.3%)	4.9% (1.6-9.5%)	180 (60-340)	2500 (880-4800)
Cote d'Ivoire	1995	Male	80 to 84	38% (26-52%)	62% (48-74%)	1.2 (0.64-2)	3.2% (1-6.1%)	3.6% (1.1-7.2%)	75 (23-150)	790 (250-1600)
Cote d'Ivoire	1995	Male	85 to 89	35% (24-49%)	65% (51-76%)	0.97 (0.5-1.7)	2.1% (0.52-4.3%)	2.3% (0.52-4.9%)	24 (5.5-52)	180 (47-380)
Cote d'Ivoire	1995	Male	90 to 94	36% (24-49%)	64% (51-76%)	0.81 (0.4-1.4)	1% (-0.17-2.8%)	1% (-0.37-3.1%)	3.7 (-1.3-11)	22 (-3.6-59)
Cote d'Ivoire	1995	Male	95 plus	39% (24-56%)	61% (44-76%)	0.69 (0.28-1.4)	0.62% (-0.62-2.5%)	0.51% (-1-2.8%)	0.41 (-0.82-2.3)	2 (-2-8.3)
Cote d'Ivoire	2000	Female	All Ages	15% (13-18%)	85% (82-87%)	0.49 (0.35-0.69)	1.1% (0.63-1.6%)	1.9% (0.96-3.2%)	1700 (830-2800)	62000 (36000-94000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	2000	Female	15 to 19	14% (9.9-19%)	86% (81-90%)	0.46 (0.25-0.76)	1.1% (0.63-1.8%)	1.1% (0.44-2.3%)	18 (6.8-37)	2100 (1200-3600)
Cote d'Ivoire	2000	Female	20 to 24	18% (13-24%)	82% (76-87%)	0.7 (0.39-1.1)	1.8% (1.1-2.9%)	1.6% (0.66-3%)	39 (17-73)	4300 (2600-6700)
Cote d'Ivoire	2000	Female	25 to 29	19% (14-25%)	81% (75-86%)	0.74 (0.42-1.2)	1.5% (0.85-2.5%)	1.2% (0.55-2.5%)	45 (21-83)	4200 (2600-6800)
Cote d'Ivoire	2000	Female	30 to 34	22% (16-29%)	78% (71-84%)	0.85 (0.48-1.4)	1.5% (0.85-2.4%)	1.4% (0.67-2.6%)	55 (28-98)	4200 (2500-6800)
Cote d'Ivoire	2000	Female	35 to 39	27% (20-33%)	73% (67-80%)	1 (0.59-1.7)	1.9% (1.1-3.1%)	2.1% (1-3.6%)	79 (41-140)	4700 (2800-7600)
Cote d'Ivoire	2000	Female	40 to 44	29% (23-36%)	71% (64-77%)	1.2 (0.69-1.9)	2.6% (1.5-4.1%)	3% (1.7-4.9%)	110 (61-180)	5500 (3200-8700)
Cote d'Ivoire	2000	Female	45 to 49	28% (20-36%)	72% (64-80%)	1.2 (0.7-2.1)	3.3% (1.8-5.6%)	4% (2-6.9%)	130 (66-230)	5800 (3000-9900)
Cote d'Ivoire	2000	Female	50 to 54	27% (20-35%)	73% (65-80%)	1.3 (0.7-2.1)	4.3% (2.2-7.2%)	5.3% (2.7-8.8%)	190 (89-320)	6900 (3400-12000)
Cote d'Ivoire	2000	Female	55 to 59	27% (20-36%)	73% (64-80%)	1.3 (0.74-2.2)	4.8% (2.4-8.3%)	5.9% (2.9-10%)	230 (100-410)	7300 (3400-13000)
Cote d'Ivoire	2000	Female	60 to 64	29% (21-39%)	71% (61-79%)	1.4 (0.74-2.4)	4.6% (1.9-8.4%)	5.6% (2.3-10%)	230 (91-440)	6400 (2500-12000)
Cote d'Ivoire	2000	Female	65 to 69	31% (23-40%)	69% (60-77%)	1.3 (0.76-2.2)	4.3% (1.6-8%)	5.1% (1.9-9.4%)	230 (78-430)	5100 (1800-9800)
Cote d'Ivoire	2000	Female	70 to 74	28% (21-37%)	72% (63-79%)	1.1 (0.58-1.9)	3.5% (0.99-7.1%)	4.1% (1.2-8.3%)	170 (48-360)	3100 (900-6500)
Cote d'Ivoire	2000	Female	75 to 79	24% (15-34%)	76% (66-85%)	0.8 (0.4-1.5)	2.3% (0.31-5.5%)	2.7% (0.34-6.3%)	91 (11-220)	1300 (170-3100)
Cote d'Ivoire	2000	Female	80 to 84	22% (14-31%)	78% (69-86%)	0.71 (0.37-1.2)	1.6% (0.15-3.8%)	1.8% (0.12-4.3%)	41 (2.7-100)	430 (38-1000)
Cote d'Ivoire	2000	Female	85 to 89	24% (16-34%)	76% (66-84%)	0.73 (0.37-1.2)	1.3% (-0.14-3.3%)	1.4% (-0.2-3.9%)	17 (-2.4-48)	130 (-14-360)
Cote d'Ivoire	2000	Female	90 to 94	27% (18-39%)	73% (61-82%)	0.79 (0.39-1.4)	0.98% (-0.44-3.4%)	1.1% (-0.62-3.9%)	4.6 (-2.5-17)	25 (-11-84)
Cote d'Ivoire	2000	Female	95 plus	32% (18-48%)	68% (52-82%)	0.86 (0.34-1.7)	0.84% (-0.79-3.6%)	0.85% (-1.1-4.2%)	0.92 (-1.2-4.6)	3.6 (-3.4-16)
Cote d'Ivoire	2000	Male	All Ages	33% (29-38%)	67% (62-71%)	1.7 (1.2-2.4)	3.6% (2.5-4.9%)	5.6% (3.7-7.8%)	6700 (4500-9600)	28000 (19000-37000)
Cote d'Ivoire	2000	Male	15 to 19	29% (22-37%)	71% (63-78%)	1.2 (0.67-2)	3.3% (1.8-5.3%)	4.2% (2.1-6.9%)	87 (43-150)	7200 (4000-12000)
Cote d'Ivoire	2000	Male	20 to 24	46% (38-54%)	54% (46-62%)	2.7 (1.6-4.2)	7.9% (4.7-12%)	9.2% (5.2-14%)	290 (160-460)	21000 (13000-32000)
Cote d'Ivoire	2000	Male	25 to 29	50% (42-58%)	50% (42-58%)	3.3 (2-5.1)	8.3% (5.1-12%)	9.3% (5.4-14%)	410 (250-600)	27000 (17000-39000)
Cote d'Ivoire	2000	Male	30 to 34	52% (44-59%)	48% (41-56%)	3.7 (2.3-5.6)	8.1% (5.3-12%)	9% (5.6-13%)	500 (330-710)	30000 (20000-41000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	2000	Male	35 to 39	52% (45-60%)	48% (40-55%)	4 (2.4-6.1)	8.7% (5.6-12%)	9.8% (5.9-14%)	600 (390-820)	32000 (21000-43000)
Cote d'Ivoire	2000	Male	40 to 44	50% (43-58%)	50% (42-57%)	3.9 (2.3-5.9)	9.5% (6.1-13%)	11% (6.8-15%)	700 (440-960)	33000 (21000-44000)
Cote d'Ivoire	2000	Male	45 to 49	48% (39-56%)	52% (44-61%)	3.5 (2.1-5.4)	11% (6.8-15%)	12% (7.6-17%)	770 (490-1100)	32000 (20000-46000)
Cote d'Ivoire	2000	Male	50 to 54	45% (37-54%)	55% (46-63%)	3.1 (1.8-4.8)	10% (6-15%)	12% (6.9-18%)	740 (430-1100)	27000 (16000-41000)
Cote d'Ivoire	2000	Male	55 to 59	48% (38-58%)	52% (42-62%)	3 (1.7-4.9)	10% (6-15%)	12% (6.8-18%)	750 (410-1100)	24000 (13000-36000)
Cote d'Ivoire	2000	Male	60 to 64	50% (39-60%)	50% (40-61%)	2.9 (1.7-4.8)	8.9% (4.3-14%)	10% (5-16%)	660 (310-1000)	18000 (8600-29000)
Cote d'Ivoire	2000	Male	65 to 69	49% (40-58%)	51% (42-60%)	2.6 (1.6-4)	7.3% (3.5-12%)	8.5% (4-14%)	520 (250-880)	12000 (5700-20000)
Cote d'Ivoire	2000	Male	70 to 74	47% (38-56%)	53% (44-62%)	2.2 (1.2-3.4)	6.1% (2.6-11%)	7% (2.9-13%)	380 (160-700)	7000 (3000-13000)
Cote d'Ivoire	2000	Male	75 to 79	43% (31-56%)	57% (44-69%)	1.7 (0.87-2.9)	4.4% (1.5-8.7%)	5% (1.7-10%)	200 (68-400)	2900 (970-5700)
Cote d'Ivoire	2000	Male	80 to 84	39% (27-53%)	61% (47-73%)	1.3 (0.7-2.2)	3.3% (1.1-6.5%)	3.8% (1.2-7.4%)	87 (27-170)	920 (310-1800)
Cote d'Ivoire	2000	Male	85 to 89	38% (25-52%)	62% (48-75%)	1.1 (0.53-1.8)	2.3% (0.58-4.6%)	2.5% (0.56-5.3%)	28 (6.3-57)	220 (52-440)
Cote d'Ivoire	2000	Male	90 to 94	38% (26-52%)	62% (48-74%)	0.89 (0.46-1.6)	1.2% (-0.096-3%)	1.3% (-0.28-3.4%)	4.7 (-1.1-13)	27 (-2.2-69)
Cote d'Ivoire	2000	Male	95 plus	42% (26-59%)	58% (41-74%)	0.77 (0.32-1.5)	0.89% (-0.52-3.2%)	0.83% (-0.88-3.6%)	0.73 (-0.78-3.2)	3.2 (-1.8-11)
Cote d'Ivoire	2005	Female	All Ages	15% (13-17%)	85% (83-87%)	0.48 (0.32-0.69)	1% (0.58-1.6%)	1.8% (0.85-3%)	1700 (790-2900)	62000 (36000-96000)
Cote d'Ivoire	2005	Female	15 to 19	14% (9.7-19%)	86% (81-90%)	0.45 (0.25-0.75)	0.98% (0.58-1.6%)	0.95% (0.37-1.8%)	17 (6.4-33)	2100 (1200-3500)
Cote d'Ivoire	2005	Female	20 to 24	18% (13-24%)	82% (76-87%)	0.7 (0.39-1.1)	1.7% (1-2.7%)	1.5% (0.6-2.7%)	35 (15-67)	4100 (2500-6500)
Cote d'Ivoire	2005	Female	25 to 29	19% (14-25%)	81% (75-86%)	0.72 (0.4-1.2)	1.4% (0.79-2.2%)	1.1% (0.5-2.2%)	42 (20-78)	4200 (2600-6500)
Cote d'Ivoire	2005	Female	30 to 34	22% (16-28%)	78% (72-84%)	0.83 (0.45-1.4)	1.3% (0.74-2.1%)	1.1% (0.55-2.1%)	53 (27-92)	4200 (2600-6300)
Cote d'Ivoire	2005	Female	35 to 39	27% (20-33%)	73% (67-80%)	1 (0.57-1.6)	1.6% (0.93-2.7%)	1.7% (0.84-3%)	78 (41-130)	4800 (2800-7500)
Cote d'Ivoire	2005	Female	40 to 44	29% (22-36%)	71% (64-78%)	1.1 (0.66-1.9)	2.3% (1.3-3.4%)	2.6% (1.4-4%)	110 (61-170)	5600 (3300-8700)
Cote d'Ivoire	2005	Female	45 to 49	28% (20-36%)	72% (64-80%)	1.2 (0.66-2)	2.9% (1.6-4.9%)	3.4% (1.8-5.9%)	130 (68-230)	5900 (3100-10000)
Cote d'Ivoire	2005	Female	50 to 54	27% (20-35%)	73% (65-80%)	1.2 (0.66-2.1)	3.9% (1.9-6.8%)	4.8% (2.3-8.2%)	180 (84-330)	6900 (3200-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	2005	Female	55 to 59	27% (20-36%)	73% (64-80%)	1.3 (0.7-2.2)	4.6% (2.2-7.9%)	5.6% (2.6-9.6%)	230 (100-400)	7300 (3300-13000)
Cote d'Ivoire	2005	Female	60 to 64	29% (21-38%)	71% (62-79%)	1.4 (0.73-2.3)	4.4% (1.8-8.1%)	5.2% (2.2-9.5%)	240 (92-450)	6500 (2500-12000)
Cote d'Ivoire	2005	Female	65 to 69	31% (23-40%)	69% (60-77%)	1.3 (0.74-2.2)	4.1% (1.4-7.8%)	4.8% (1.6-9.1%)	230 (80-460)	5300 (1800-10000)
Cote d'Ivoire	2005	Female	70 to 74	28% (20-37%)	72% (63-80%)	1.1 (0.54-1.9)	3.2% (0.82-6.6%)	3.7% (0.98-7.7%)	170 (43-360)	3100 (800-6500)
Cote d'Ivoire	2005	Female	75 to 79	24% (16-34%)	76% (66-84%)	0.8 (0.38-1.5)	2.1% (0.21-5.1%)	2.4% (0.2-5.9%)	88 (7.5-210)	1200 (120-2900)
Cote d'Ivoire	2005	Female	80 to 84	22% (14-31%)	78% (69-86%)	0.7 (0.37-1.2)	1.5% (0.0065-3.8%)	1.7% (-0.069-4.4%)	38 (-1.5-98)	400 (1.7-1000)
Cote d'Ivoire	2005	Female	85 to 89	24% (15-34%)	76% (66-85%)	0.71 (0.36-1.3)	1.2% (-0.21-3.2%)	1.3% (-0.28-3.7%)	16 (-3.6-45)	120 (-21-340)
Cote d'Ivoire	2005	Female	90 to 94	27% (18-39%)	73% (61-82%)	0.77 (0.37-1.4)	0.9% (-0.55-3.1%)	0.98% (-0.76-3.6%)	4.3 (-3.2-16)	24 (-14-81)
Cote d'Ivoire	2005	Female	95 plus	32% (18-49%)	68% (51-82%)	0.86 (0.35-1.7)	0.77% (-0.85-3.5%)	0.76% (-1.2-4%)	0.86 (-1.3-4.7)	3.5 (-3.9-16)
Cote d'Ivoire	2005	Male	All Ages	32% (28-37%)	68% (63-72%)	1.7 (1.1-2.3)	3.5% (2.3-4.8%)	5.4% (3.4-7.6%)	6700 (4300-9500)	270000 (180000-360000)
Cote d'Ivoire	2005	Male	15 to 19	28% (21-37%)	72% (63-79%)	1.2 (0.67-1.9)	3% (1.7-4.8%)	3.6% (1.9-6%)	84 (44-140)	7000 (4000-12000)
Cote d'Ivoire	2005	Male	20 to 24	45% (37-53%)	55% (47-63%)	2.6 (1.5-4)	7.5% (4.5-11%)	8.7% (5.1-13%)	270 (150-410)	20000 (12000-30000)
Cote d'Ivoire	2005	Male	25 to 29	50% (41-57%)	50% (43-59%)	3.2 (1.9-5)	8.3% (5-12%)	9.4% (5.4-14%)	390 (240-570)	26000 (17000-37000)
Cote d'Ivoire	2005	Male	30 to 34	52% (44-59%)	48% (41-56%)	3.6 (2.2-5.6)	8.1% (5.2-11%)	9% (5.5-13%)	480 (320-670)	29000 (19000-39000)
Cote d'Ivoire	2005	Male	35 to 39	52% (44-59%)	48% (41-56%)	4 (2.4-6.1)	8.4% (5.6-12%)	9.4% (6.1-13%)	580 (390-790)	31000 (21000-41000)
Cote d'Ivoire	2005	Male	40 to 44	50% (42-57%)	50% (43-58%)	3.8 (2.3-5.9)	9% (5.9-12%)	10% (6.6-14%)	680 (430-930)	32000 (21000-43000)
Cote d'Ivoire	2005	Male	45 to 49	48% (39-56%)	52% (44-61%)	3.5 (2.1-5.4)	10% (6.4-14%)	12% (7.2-16%)	760 (460-1100)	32000 (20000-45000)
Cote d'Ivoire	2005	Male	50 to 54	45% (36-54%)	55% (46-64%)	3 (1.7-4.9)	9.9% (5.9-14%)	11% (6.5-17%)	710 (410-1100)	26000 (15000-39000)
Cote d'Ivoire	2005	Male	55 to 59	48% (38-57%)	52% (43-62%)	3 (1.7-4.7)	10% (5.6-15%)	12% (6.6-17%)	750 (420-1200)	24000 (14000-37000)
Cote d'Ivoire	2005	Male	60 to 64	50% (40-60%)	50% (40-60%)	2.8 (1.6-4.5)	8.7% (4.4-14%)	10% (5-16%)	680 (340-1100)	19000 (9400-29000)
Cote d'Ivoire	2005	Male	65 to 69	48% (39-57%)	52% (43-61%)	2.5 (1.5-3.9)	7.1% (3.5-12%)	8.2% (3.9-14%)	540 (260-930)	12000 (6000-21000)
Cote d'Ivoire	2005	Male	70 to 74	46% (37-55%)	54% (45-63%)	2.1 (1.2-3.4)	5.9% (2.4-11%)	6.7% (2.7-12%)	410 (170-740)	7400 (3000-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	2005	Male	75 to 79	42% (30-55%)	58% (45-70%)	1.7 (0.83-2.9)	4.2% (1.3-8.4%)	4.8% (1.4-9.7%)	210 (59-420)	2900 (880-5900)
Cote d'Ivoire	2005	Male	80 to 84	39% (26-52%)	61% (48-74%)	1.3 (0.66-2.2)	3.1% (0.94-6.5%)	3.6% (1-7.4%)	83 (23-170)	890 (270-1800)
Cote d'Ivoire	2005	Male	85 to 89	37% (25-51%)	63% (49-75%)	1 (0.53-1.8)	2.1% (0.51-4.5%)	2.4% (0.46-5.2%)	29 (5.5-62)	220 (54-470)
Cote d'Ivoire	2005	Male	90 to 94	38% (26-52%)	62% (48-74%)	0.88 (0.43-1.5)	1.1% (-0.12-3%)	1.1% (-0.3-3.3%)	4.5 (-1.2-14)	26 (-2.9-71)
Cote d'Ivoire	2005	Male	95 plus	42% (26-59%)	58% (41-74%)	0.75 (0.31-1.5)	0.79% (-0.56-2.8%)	0.71% (-0.94-3.1%)	0.64 (-0.86-2.8)	2.9 (-2-10)
Cote d'Ivoire	2010	Female	All Ages	17% (14-20%)	83% (80-86%)	0.53 (0.33-0.8)	1.1% (0.58-1.7%)	1.9% (0.82-3.3%)	1700 (740-3100)	65000 (36000-100000)
Cote d'Ivoire	2010	Female	15 to 19	16% (11-23%)	84% (77-89%)	0.53 (0.27-0.9)	1% (0.54-1.6%)	0.96% (0.36-2%)	19 (7.1-39)	2400 (1300-3900)
Cote d'Ivoire	2010	Female	20 to 24	22% (14-31%)	78% (69-86%)	0.84 (0.43-1.5)	1.9% (1.1-3%)	1.6% (0.63-3.2%)	39 (15-77)	4600 (2700-7500)
Cote d'Ivoire	2010	Female	25 to 29	23% (15-32%)	77% (68-85%)	0.87 (0.45-1.5)	1.6% (0.9-2.6%)	1.4% (0.55-2.6%)	45 (21-83)	4500 (2800-7000)
Cote d'Ivoire	2010	Female	30 to 34	25% (17-36%)	75% (64-83%)	0.95 (0.47-1.7)	1.5% (0.86-2.4%)	1.4% (0.63-2.6%)	57 (28-100)	4600 (2800-7200)
Cote d'Ivoire	2010	Female	35 to 39	29% (19-40%)	71% (60-81%)	1.1 (0.53-2)	1.8% (0.98-2.9%)	1.9% (0.89-3.3%)	82 (39-140)	5100 (2800-8200)
Cote d'Ivoire	2010	Female	40 to 44	30% (21-41%)	70% (59-79%)	1.2 (0.6-2.1)	2.4% (1.3-3.9%)	2.8% (1.4-4.7%)	120 (60-190)	6000 (3200-9700)
Cote d'Ivoire	2010	Female	45 to 49	29% (18-39%)	71% (61-82%)	1.2 (0.61-2.2)	3% (1.5-5.1%)	3.6% (1.7-6.4%)	140 (65-260)	6200 (3100-11000)
Cote d'Ivoire	2010	Female	50 to 54	29% (20-39%)	71% (61-80%)	1.3 (0.64-2.3)	4% (1.9-6.9%)	5% (2.3-8.7%)	190 (83-340)	7300 (3300-13000)
Cote d'Ivoire	2010	Female	55 to 59	29% (20-40%)	71% (60-80%)	1.4 (0.68-2.4)	4.6% (2-8.1%)	5.7% (2.6-10%)	230 (96-420)	7400 (3100-13000)
Cote d'Ivoire	2010	Female	60 to 64	30% (20-42%)	70% (58-80%)	1.4 (0.69-2.4)	4.2% (1.5-8%)	5.1% (1.9-9.4%)	230 (81-450)	6300 (2300-12000)
Cote d'Ivoire	2010	Female	65 to 69	32% (23-42%)	68% (58-77%)	1.3 (0.68-2.2)	3.8% (1.2-7.7%)	4.6% (1.4-9.1%)	230 (67-480)	5300 (1500-11000)
Cote d'Ivoire	2010	Female	70 to 74	29% (21-39%)	71% (61-79%)	1.1 (0.54-2)	3% (0.53-7.1%)	3.5% (0.6-8.4%)	170 (29-410)	3200 (580-7500)
Cote d'Ivoire	2010	Female	75 to 79	24% (16-35%)	76% (65-84%)	0.78 (0.37-1.5)	1.9% (0.021-5.2%)	2.3% (-0.0033-6%)	90 (-0.13-250)	1300 (14-3500)
Cote d'Ivoire	2010	Female	80 to 84	22% (14-31%)	78% (69-86%)	0.67 (0.33-1.2)	1.3% (-0.17-3.9%)	1.5% (-0.23-4.4%)	36 (-5.8-110)	390 (-49-1100)
Cote d'Ivoire	2010	Female	85 to 89	23% (15-34%)	77% (66-85%)	0.69 (0.32-1.2)	1% (-0.32-3.3%)	1.1% (-0.43-3.7%)	15 (-5.8-50)	120 (-37-370)
Cote d'Ivoire	2010	Female	90 to 94	27% (17-39%)	73% (61-83%)	0.75 (0.34-1.4)	0.76% (-0.63-3.1%)	0.81% (-0.86-3.6%)	3.9 (-4.2-17)	22 (-18-90)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	2010	Female	95 plus	32% (17-48%)	68% (52-83%)	0.82 (0.3-1.6)	0.62% (-0.95-3.2%)	0.58% (-1.3-3.6%)	0.72 (-1.6-4.6)	3.1 (-4.8-16)
Cote d'Ivoire	2010	Male	All Ages	30% (24-35%)	70% (65-76%)	1.5 (0.9-2.2)	3.3% (2.2-4.7%)	5.2% (3.2-7.5%)	6400 (3800-9200)	260000 (170000-350000)
Cote d'Ivoire	2010	Male	15 to 19	23% (15-32%)	77% (68-85%)	0.93 (0.45-1.6)	2.3% (1.2-3.9%)	2.7% (1.2-4.7%)	71 (32-130)	6200 (3300-11000)
Cote d'Ivoire	2010	Male	20 to 24	37% (25-48%)	63% (52-75%)	2 (1.1-3.4)	6.1% (3.5-9.4%)	7% (3.6-11%)	230 (120-370)	17000 (9800-26000)
Cote d'Ivoire	2010	Male	25 to 29	44% (32-57%)	56% (43-68%)	2.8 (1.5-4.6)	7.6% (4.5-11%)	8.7% (4.8-13%)	340 (190-500)	23000 (14000-34000)
Cote d'Ivoire	2010	Male	30 to 34	48% (35-61%)	52% (39-65%)	3.3 (1.8-5.5)	8% (4.9-12%)	9.1% (5.4-14%)	450 (270-640)	27000 (17000-38000)
Cote d'Ivoire	2010	Male	35 to 39	49% (36-61%)	51% (39-64%)	3.7 (1.9-6.4)	8.5% (5.4-12%)	9.7% (5.9-14%)	540 (340-750)	29000 (19000-40000)
Cote d'Ivoire	2010	Male	40 to 44	48% (36-60%)	52% (40-64%)	3.6 (2-6)	9.2% (5.8-13%)	11% (6.6-15%)	640 (400-910)	30000 (19000-43000)
Cote d'Ivoire	2010	Male	45 to 49	46% (32-57%)	54% (43-68%)	3.3 (1.8-5.4)	10% (6.3-14%)	12% (7.3-17%)	720 (430-1000)	30000 (19000-44000)
Cote d'Ivoire	2010	Male	50 to 54	46% (35-58%)	54% (42-65%)	3 (1.6-5.1)	10% (5.7-15%)	12% (6.6-17%)	700 (390-1100)	26000 (15000-39000)
Cote d'Ivoire	2010	Male	55 to 59	48% (36-60%)	52% (40-64%)	2.9 (1.5-5)	9.9% (5.4-15%)	12% (6.4-18%)	710 (370-1100)	23000 (12000-35000)
Cote d'Ivoire	2010	Male	60 to 64	48% (35-60%)	52% (40-65%)	2.6 (1.4-4.7)	8.3% (3.9-13%)	9.6% (4.5-16%)	660 (300-1100)	18000 (8600-29000)
Cote d'Ivoire	2010	Male	65 to 69	46% (35-58%)	54% (42-65%)	2.3 (1.2-3.8)	6.6% (2.9-12%)	7.7% (3.3-14%)	540 (240-960)	12000 (5500-22000)
Cote d'Ivoire	2010	Male	70 to 74	45% (34-56%)	55% (44-66%)	2 (1-3.4)	5.4% (2-10%)	6.3% (2.3-12%)	420 (150-810)	7600 (2800-14000)
Cote d'Ivoire	2010	Male	75 to 79	42% (28-55%)	58% (45-72%)	1.6 (0.79-3)	3.9% (1.1-8.1%)	4.5% (1.2-9.4%)	220 (58-470)	3100 (860-6600)
Cote d'Ivoire	2010	Male	80 to 84	38% (26-52%)	62% (48-74%)	1.3 (0.62-2.3)	2.9% (0.79-6.1%)	3.3% (0.82-7%)	84 (20-180)	900 (240-1900)
Cote d'Ivoire	2010	Male	85 to 89	36% (24-50%)	64% (50-76%)	0.99 (0.48-1.8)	1.9% (0.39-4.3%)	2.1% (0.35-5%)	29 (4.8-68)	230 (48-520)
Cote d'Ivoire	2010	Male	90 to 94	37% (25-50%)	63% (50-75%)	0.82 (0.39-1.5)	0.98% (-0.18-2.9%)	1% (-0.39-3.3%)	4.8 (-1.8-16)	28 (-5.1-85)
Cote d'Ivoire	2010	Male	95 plus	40% (25-58%)	60% (42-75%)	0.7 (0.27-1.4)	0.67% (-0.59-2.6%)	0.57% (-0.96-3%)	0.61 (-1-3.1)	2.9 (-2.6-11)
Cote d'Ivoire	2016	Female	All Ages	18% (15-21%)	82% (79-85%)	0.52 (0.27-0.9)	1.2% (0.53-2.1%)	2% (0.61-4.2%)	1800 (550-3700)	69000 (32000-130000)
Cote d'Ivoire	2016	Female	15 to 19	17% (11-23%)	83% (77-89%)	0.51 (0.21-0.98)	1% (0.54-1.8%)	0.97% (0.28-2.2%)	20 (5.6-44)	2600 (1400-4800)
Cote d'Ivoire	2016	Female	20 to 24	22% (14-31%)	78% (69-86%)	0.81 (0.34-1.5)	2% (1.1-3.2%)	1.7% (0.55-3.6%)	39 (13-84)	5100 (2900-8400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	2016	Female	25 to 29	23% (15-33%)	77% (67-85%)	0.83 (0.37-1.6)	1.9% (1.1-3%)	1.6% (0.58-3.2%)	45 (18-89)	4900 (2900-7800)
Cote d'Ivoire	2016	Female	30 to 34	26% (17-37%)	74% (63-83%)	0.91 (0.39-1.8)	1.8% (1-2.9%)	1.8% (0.74-3.3%)	57 (24-100)	4800 (2800-7700)
Cote d'Ivoire	2016	Female	35 to 39	29% (19-40%)	71% (60-81%)	1 (0.43-2)	2.2% (1.1-3.7%)	2.5% (1-4.7%)	86 (34-160)	5500 (2700-9700)
Cote d'Ivoire	2016	Female	40 to 44	31% (21-41%)	69% (59-79%)	1.1 (0.48-2.2)	2.9% (1.4-4.8%)	3.6% (1.6-6.3%)	130 (56-230)	6600 (3200-12000)
Cote d'Ivoire	2016	Female	45 to 49	30% (19-40%)	70% (60-81%)	1.2 (0.46-2.4)	3.4% (1.4-6.3%)	4.4% (1.8-8.4%)	160 (60-310)	7000 (2800-13000)
Cote d'Ivoire	2016	Female	50 to 54	29% (20-39%)	71% (61-80%)	1.3 (0.51-2.4)	4.2% (1.6-7.9%)	5.4% (2-10%)	210 (75-410)	8000 (3000-15000)
Cote d'Ivoire	2016	Female	55 to 59	29% (20-40%)	71% (60-80%)	1.3 (0.58-2.7)	4.6% (1.5-8.9%)	5.8% (2-11%)	250 (78-490)	7900 (2600-16000)
Cote d'Ivoire	2016	Female	60 to 64	31% (20-43%)	69% (57-80%)	1.3 (0.57-2.7)	4.1% (1.1-8.3%)	5% (1.4-10%)	240 (65-490)	6500 (1800-13000)
Cote d'Ivoire	2016	Female	65 to 69	32% (23-43%)	68% (57-77%)	1.3 (0.56-2.5)	3.6% (0.62-8.3%)	4.3% (0.74-9.9%)	230 (39-530)	5200 (900-12000)
Cote d'Ivoire	2016	Female	70 to 74	30% (21-40%)	70% (60-79%)	1 (0.44-2.1)	2.7% (0.078-7.6%)	3.2% (0.076-8.9%)	180 (4.4-480)	3200 (92-8800)
Cote d'Ivoire	2016	Female	75 to 79	24% (16-35%)	76% (65-84%)	0.76 (0.29-1.6)	1.8% (-0.26-5.6%)	2% (-0.31-6.7%)	96 (-15-310)	1400 (-200-4500)
Cote d'Ivoire	2016	Female	80 to 84	22% (14-32%)	78% (68-86%)	0.65 (0.26-1.3)	1.2% (-0.4-4.3%)	1.4% (-0.53-4.9%)	39 (-14-140)	410 (-130-1500)
Cote d'Ivoire	2016	Female	85 to 89	24% (15-35%)	76% (65-85%)	0.67 (0.27-1.4)	0.92% (-0.59-3.8%)	1% (-0.73-4.4%)	17 (-12-71)	130 (-82-540)
Cote d'Ivoire	2016	Female	90 to 94	27% (18-39%)	73% (61-82%)	0.74 (0.28-1.5)	0.69% (-0.92-3.6%)	0.73% (-1.2-4.3%)	4.3 (-7-25)	24 (-32-130)
Cote d'Ivoire	2016	Female	95 plus	32% (18-49%)	68% (51-82%)	0.81 (0.26-1.8)	0.58% (-1.2-4%)	0.53% (-1.6-4.7%)	0.82 (-2.4-7.2)	3.6 (-7.3-25)
Cote d'Ivoire	2016	Male	All Ages	30% (25-35%)	70% (65-75%)	1.4 (0.72-2.4)	3.4% (1.9-5.2%)	5.2% (2.6-8.4%)	6200 (3100-10000)	250000 (140000-380000)
Cote d'Ivoire	2016	Male	15 to 19	23% (15-32%)	77% (68-85%)	0.88 (0.37-1.7)	2.2% (1.1-4%)	2.6% (1-5.1%)	75 (29-150)	6700 (3200-12000)
Cote d'Ivoire	2016	Male	20 to 24	37% (25-48%)	63% (52-75%)	1.9 (0.87-3.5)	5.9% (3-9.7%)	6.7% (3-12%)	240 (110-420)	19000 (9500-31000)
Cote d'Ivoire	2016	Male	25 to 29	44% (32-56%)	56% (44-68%)	2.6 (1.2-4.9)	7.6% (4.1-12%)	8.8% (4.4-14%)	350 (180-560)	24000 (13000-37000)
Cote d'Ivoire	2016	Male	30 to 34	48% (35-60%)	52% (40-65%)	3.1 (1.5-5.9)	8.2% (4.7-12%)	9.5% (5.1-14%)	440 (240-660)	27000 (15000-40000)
Cote d'Ivoire	2016	Male	35 to 39	49% (37-61%)	51% (39-63%)	3.5 (1.5-6.9)	8.8% (5.1-13%)	10% (5.8-16%)	540 (290-790)	29000 (16000-42000)
Cote d'Ivoire	2016	Male	40 to 44	48% (37-60%)	52% (40-63%)	3.5 (1.5-6.5)	9.4% (5.2-13%)	11% (5.9-16%)	620 (320-940)	30000 (16000-45000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Cote d'Ivoire	2016	Male	45 to 49	47% (32-58%)	53% (42-68%)	3.1 (1.4-6)	10% (5.2-15%)	12% (6-18%)	690 (350-1100)	30000 (15000-45000)
Cote d'Ivoire	2016	Male	50 to 54	47% (35-59%)	53% (41-65%)	2.9 (1.3-5.5)	9.4% (4.4-15%)	11% (5.2-18%)	670 (300-1100)	25000 (12000-39000)
Cote d'Ivoire	2016	Male	55 to 59	48% (36-61%)	52% (39-64%)	2.8 (1.2-5.2)	9.3% (4.1-15%)	11% (4.7-18%)	690 (290-1100)	22000 (9700-36000)
Cote d'Ivoire	2016	Male	60 to 64	48% (34-61%)	52% (39-66%)	2.5 (1.1-4.9)	7.7% (2.9-13%)	9% (3.3-16%)	610 (220-1100)	17000 (6300-29000)
Cote d'Ivoire	2016	Male	65 to 69	47% (36-58%)	53% (42-64%)	2.2 (0.99-4.1)	6.1% (2-12%)	7.1% (2.3-14%)	530 (160-1000)	12000 (3900-23000)
Cote d'Ivoire	2016	Male	70 to 74	45% (34-56%)	55% (44-66%)	1.9 (0.85-3.7)	4.9% (1.3-10%)	5.7% (1.4-12%)	420 (100-890)	7700 (2000-16000)
Cote d'Ivoire	2016	Male	75 to 79	42% (29-56%)	58% (44-71%)	1.5 (0.63-3.1)	3.5% (0.58-8.6%)	4% (0.61-10%)	240 (36-590)	3400 (560-8300)
Cote d'Ivoire	2016	Male	80 to 84	38% (26-52%)	62% (48-74%)	1.2 (0.5-2.4)	2.6% (0.39-6.4%)	2.9% (0.38-7.5%)	90 (11-230)	970 (150-2400)
Cote d'Ivoire	2016	Male	85 to 89	36% (24-50%)	64% (50-76%)	0.93 (0.39-1.9)	1.7% (0.12-4.7%)	1.9% (0.046-5.4%)	31 (0.77-92)	250 (17-680)
Cote d'Ivoire	2016	Male	90 to 94	36% (24-51%)	64% (49-76%)	0.77 (0.3-1.5)	0.86% (-0.33-3.1%)	0.85% (-0.57-3.4%)	5.2 (-3.4-21)	32 (-12-110)
Cote d'Ivoire	2016	Male	95 plus	40% (25-58%)	60% (42-75%)	0.66 (0.22-1.5)	0.58% (-0.58-2.8%)	0.45% (-0.94-3%)	0.67 (-1.4-4.4)	3.4 (-3.5-16)
The Gambia	1990	Female	All Ages	22% (16-28%)	78% (72-84%)	0.39 (0.27-0.53)	0.67% (0.38-1%)	1.1% (0.43-2.2%)	44 (17-86)	2000 (1100-3100)
The Gambia	1990	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.38 (0.15-0.71)	0.89% (0.44-1.6%)	0.85% (0.1-2.1%)	0.59 (0.073-1.5)	84 (39-150)
The Gambia	1990	Female	20 to 24	27% (13-43%)	73% (57-87%)	0.59 (0.24-1.1)	1.9% (0.99-3.3%)	1.7% (0.38-3.9%)	1.7 (0.38-4.1)	220 (110-400)
The Gambia	1990	Female	25 to 29	30% (15-47%)	70% (53-85%)	0.73 (0.32-1.3)	2.3% (1.2-3.9%)	2.5% (0.78-5%)	2.5 (0.77-5)	250 (120-400)
The Gambia	1990	Female	30 to 34	32% (16-51%)	68% (49-84%)	0.82 (0.39-1.4)	2.5% (1.3-4%)	3.1% (1.2-5.7%)	2.9 (1.2-5.5)	230 (120-370)
The Gambia	1990	Female	35 to 39	36% (19-54%)	64% (46-81%)	0.88 (0.4-1.5)	2.8% (1.4-4.8%)	3.9% (1.6-7.4%)	3.3 (1.3-6.1)	200 (98-350)
The Gambia	1990	Female	40 to 44	37% (22-56%)	63% (44-78%)	0.93 (0.37-1.7)	3.1% (1.6-5.4%)	4.6% (2-8.2%)	2.9 (1.2-5.2)	150 (73-260)
The Gambia	1990	Female	45 to 49	38% (20-58%)	62% (42-80%)	0.97 (0.37-1.8)	3.3% (1.4-6.4%)	4.7% (1.8-9.5%)	3.9 (1.5-7.9)	170 (72-340)
The Gambia	1990	Female	50 to 54	39% (21-59%)	61% (41-79%)	1 (0.41-1.9)	3.5% (1.4-6.3%)	4.8% (1.9-8.8%)	3.9 (1.5-7.4)	150 (61-280)
The Gambia	1990	Female	55 to 59	39% (23-58%)	61% (42-77%)	1.1 (0.44-2.2)	3.6% (1.1-7.1%)	4.8% (1.4-9.6%)	5 (1.4-11)	160 (47-340)
The Gambia	1990	Female	60 to 64	40% (22-59%)	60% (41-78%)	1.1 (0.41-2.2)	3% (0.61-6.6%)	3.8% (0.78-8.5%)	5.2 (0.99-12)	140 (28-320)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	1990	Female	65 to 69	41% (24-59%)	59% (41-76%)	1 (0.41-2)	2.2% (0.078-5.7%)	2.8% (0.074-7.1%)	4.6 (0.11-12)	100 (3.1-280)
The Gambia	1990	Female	70 to 74	39% (24-57%)	61% (43-76%)	0.83 (0.32-1.7)	1.7% (-0.33-5.4%)	2.1% (-0.43-6.7%)	4.7 (-1-15)	84 (-17-260)
The Gambia	1990	Female	75 to 79	35% (19-54%)	65% (46-81%)	0.68 (0.22-1.5)	0.91% (-0.77-3.9%)	1.1% (-0.96-4.6%)	1.9 (-1.7-8.1)	27 (-23-110)
The Gambia	1990	Female	80 to 84	33% (17-52%)	67% (48-83%)	0.62 (0.2-1.3)	0.54% (-0.88-2.6%)	0.63% (-1.1-3.1%)	0.56 (-0.95-3)	6 (-9.6-31)
The Gambia	1990	Female	85 to 89	33% (18-52%)	67% (48-82%)	0.61 (0.2-1.3)	0.25% (-1.2-2.2%)	0.26% (-1.4-2.6%)	0.12 (-0.64-1.2)	1 (-4.6-8.9)
The Gambia	1990	Female	90 to 94	37% (21-55%)	63% (45-79%)	0.65 (0.2-1.4)	0.0086% (-1.4-1.9%)	-0.059% (-1.7-2.3%)	-0.0084 (-0.25-0.33)	0.0099 (-1.2-1.7)
The Gambia	1990	Female	95 plus	41% (22-61%)	59% (39-78%)	0.7 (0.18-1.6)	-0.067% (-1.5-2.2%)	-0.2% (-1.9-2.5%)	-0.0075 (-0.072-0.092)	-0.0097 (-0.22-0.33)
The Gambia	1990	Male	All Ages	18% (14-24%)	82% (76-86%)	0.47 (0.34-0.63)	1.1% (0.73-1.5%)	1.8% (1.1-2.8%)	84 (53-130)	3600 (2500-5100)
The Gambia	1990	Male	15 to 19	14% (7.6-24%)	86% (76-92%)	0.29 (0.11-0.58)	1% (0.59-1.7%)	0.94% (0.33-1.9%)	0.63 (0.22-1.3)	87 (48-150)
The Gambia	1990	Male	20 to 24	22% (11-35%)	78% (65-89%)	0.59 (0.23-1.1)	2.7% (1.5-4.4%)	2.7% (1-5.2%)	2.7 (1.1-5.2)	280 (150-460)
The Gambia	1990	Male	25 to 29	27% (14-43%)	73% (57-86%)	0.87 (0.37-1.5)	4.2% (2.4-6.5%)	4.8% (2.1-8.4%)	4.6 (2-8.1)	370 (210-590)
The Gambia	1990	Male	30 to 34	31% (16-49%)	69% (51-84%)	1.1 (0.52-1.8)	5.2% (3.1-7.9%)	6.5% (3.5-10%)	6.1 (3.2-10)	420 (250-650)
The Gambia	1990	Male	35 to 39	34% (18-52%)	66% (48-82%)	1.2 (0.52-2.2)	5.9% (3.6-9.1%)	7.8% (4.5-12%)	7.2 (4-12)	420 (250-650)
The Gambia	1990	Male	40 to 44	35% (20-54%)	65% (46-80%)	1.3 (0.5-2.5)	6.4% (3.8-10%)	8.8% (5-14%)	7.6 (4.2-13)	380 (220-620)
The Gambia	1990	Male	45 to 49	36% (19-54%)	64% (46-81%)	1.4 (0.52-2.8)	6.9% (3.9-11%)	9.3% (5.2-15%)	9.8 (5.4-16)	430 (240-700)
The Gambia	1990	Male	50 to 54	36% (20-57%)	64% (43-80%)	1.4 (0.52-2.8)	6.4% (3.2-11%)	8.5% (4.1-15%)	8.9 (4.3-16)	340 (170-590)
The Gambia	1990	Male	55 to 59	36% (21-56%)	64% (44-79%)	1.4 (0.52-2.7)	5.8% (2.9-10%)	7.5% (3.7-13%)	9.2 (4.5-16)	300 (150-530)
The Gambia	1990	Male	60 to 64	36% (20-55%)	64% (45-80%)	1.2 (0.43-2.5)	4.5% (2.1-8.5%)	5.6% (2.5-11%)	8.5 (3.8-16)	240 (110-440)
The Gambia	1990	Male	65 to 69	36% (21-54%)	64% (46-79%)	1 (0.39-2.1)	3.4% (1.4-7.3%)	4.2% (1.7-8.9%)	8.2 (3.3-17)	190 (79-390)
The Gambia	1990	Male	70 to 74	34% (20-52%)	66% (48-80%)	0.78 (0.29-1.8)	2.4% (0.67-5.5%)	2.9% (0.78-6.8%)	5.8 (1.6-14)	110 (32-250)
The Gambia	1990	Male	75 to 79	31% (17-50%)	69% (50-83%)	0.6 (0.18-1.3)	1.5% (0.28-3.8%)	1.8% (0.29-4.5%)	3 (0.51-7.6)	43 (8.1-110)
The Gambia	1990	Male	80 to 84	28% (15-46%)	72% (54-85%)	0.49 (0.16-1)	1.2% (0.2-2.8%)	1.4% (0.16-3.4%)	1.2 (0.14-2.9)	13 (2.1-30)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	1990	Male	85 to 89	27% (14-43%)	73% (57-86%)	0.42 (0.13-0.92)	0.81% (-0.032-2.1%)	0.91% (-0.12-2.4%)	0.45 (-0.06-1.2)	3.5 (-0.13-8.9)
The Gambia	1990	Male	90 to 94	28% (15-46%)	72% (54-85%)	0.4 (0.11-0.97)	0.36% (-0.34-1.4%)	0.34% (-0.53-1.6%)	0.05 (-0.082-0.24)	0.33 (-0.31-1.3)
The Gambia	1990	Male	95 plus	33% (15-55%)	67% (45-85%)	0.4 (0.086-1.1)	0.27% (-0.65-1.8%)	0.16% (-0.99-2%)	0.0048 (-0.03-0.06)	0.033 (-0.079-0.22)
The Gambia	1995	Female	All Ages	22% (16-28%)	78% (72-84%)	0.38 (0.28-0.51)	0.71% (0.41-1.1%)	1.2% (0.44-2.2%)	51 (19-96)	2300 (1300-3600)
The Gambia	1995	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.38 (0.15-0.73)	0.92% (0.43-1.8%)	0.9% (0.079-2.3%)	0.75 (0.065-1.9)	110 (48-200)
The Gambia	1995	Female	20 to 24	27% (14-44%)	73% (56-86%)	0.59 (0.24-1.1)	1.9% (0.96-3.3%)	1.7% (0.34-4%)	1.6 (0.32-3.9)	210 (100-380)
The Gambia	1995	Female	25 to 29	30% (15-47%)	70% (53-85%)	0.72 (0.32-1.3)	2.3% (1.2-3.8%)	2.5% (0.72-5.1%)	2.7 (0.77-5.5)	270 (140-450)
The Gambia	1995	Female	30 to 34	33% (17-53%)	67% (47-83%)	0.82 (0.39-1.4)	2.5% (1.3-4%)	3% (1.2-5.6%)	3.2 (1.3-6)	260 (130-410)
The Gambia	1995	Female	35 to 39	36% (20-55%)	64% (45-80%)	0.88 (0.41-1.5)	2.8% (1.4-4.9%)	3.8% (1.6-7.2%)	4 (1.6-7.6)	250 (130-430)
The Gambia	1995	Female	40 to 44	38% (22-57%)	62% (43-78%)	0.93 (0.38-1.7)	3.1% (1.5-5.4%)	4.5% (2-8.3%)	4.3 (1.9-7.9)	220 (110-390)
The Gambia	1995	Female	45 to 49	39% (20-58%)	61% (42-80%)	0.96 (0.38-1.8)	3.2% (1.4-5.9%)	4.6% (1.8-9.1%)	3.6 (1.4-7)	160 (68-300)
The Gambia	1995	Female	50 to 54	39% (21-59%)	61% (41-79%)	1 (0.39-1.9)	3.5% (1.5-6.6%)	4.8% (1.9-9.2%)	5.5 (2.1-11)	210 (83-390)
The Gambia	1995	Female	55 to 59	39% (22-58%)	61% (42-78%)	1.1 (0.43-2.1)	3.5% (1.1-7.1%)	4.7% (1.4-9.8%)	5.4 (1.6-11)	170 (55-360)
The Gambia	1995	Female	60 to 64	40% (22-60%)	60% (40-78%)	1.1 (0.41-2.2)	2.9% (0.61-6.5%)	3.8% (0.76-8.3%)	5.8 (1.3-13)	160 (35-350)
The Gambia	1995	Female	65 to 69	42% (25-59%)	58% (41-75%)	1 (0.42-2)	2.2% (0.028-5.7%)	2.8% (0.025-7%)	5.5 (0.054-14)	120 (1.8-320)
The Gambia	1995	Female	70 to 74	40% (24-57%)	60% (43-76%)	0.84 (0.32-1.8)	1.7% (-0.45-5.1%)	2.1% (-0.56-6.3%)	4.9 (-1.3-15)	89 (-24-270)
The Gambia	1995	Female	75 to 79	35% (19-55%)	65% (45-81%)	0.67 (0.21-1.5)	0.91% (-0.91-3.8%)	1.1% (-1.1-4.4%)	2.5 (-2.6-11)	35 (-35-150)
The Gambia	1995	Female	80 to 84	33% (17-51%)	67% (49-83%)	0.6 (0.2-1.2)	0.53% (-0.91-2.7%)	0.62% (-1.1-3.1%)	0.75 (-1.4-3.9)	8 (-14-41)
The Gambia	1995	Female	85 to 89	33% (18-51%)	67% (49-82%)	0.6 (0.19-1.3)	0.25% (-1.1-2.2%)	0.27% (-1.4-2.5%)	0.17 (-0.83-1.6)	1.4 (-5.8-12)
The Gambia	1995	Female	90 to 94	36% (20-54%)	64% (46-80%)	0.63 (0.19-1.4)	0.02% (-1.4-2%)	-0.043% (-1.7-2.5%)	-0.0089 (-0.36-0.53)	0.028 (-1.7-2.7)
The Gambia	1995	Female	95 plus	41% (22-62%)	59% (38-78%)	0.67 (0.18-1.6)	-0.044% (-1.5-2.3%)	-0.17% (-1.9-2.6%)	-0.0083 (-0.092-0.13)	-0.008 (-0.29-0.44)
The Gambia	1995	Male	All Ages	18% (14-24%)	82% (76-86%)	0.47 (0.35-0.63)	1.1% (0.78-1.6%)	1.9% (1.2-3%)	95 (59-140)	4100 (2900-5600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	1995	Male	15 to 19	15% (7.6-25%)	85% (75-92%)	0.3 (0.11-0.62)	1.1% (0.62-1.8%)	1% (0.31-2.2%)	0.78 (0.24-1.7)	110 (59-180)
The Gambia	1995	Male	20 to 24	23% (12-37%)	77% (63-88%)	0.61 (0.24-1.1)	2.7% (1.5-4.4%)	2.6% (0.99-5.1%)	2.6 (0.95-5.1)	270 (150-450)
The Gambia	1995	Male	25 to 29	28% (15-44%)	72% (56-85%)	0.91 (0.37-1.6)	4% (2.2-6.3%)	4.5% (2-7.9%)	5.1 (2.2-9.1)	420 (230-670)
The Gambia	1995	Male	30 to 34	32% (16-50%)	68% (50-84%)	1.1 (0.54-1.9)	4.8% (3-7.3%)	6% (3.2-9.6%)	6.6 (3.5-11)	460 (260-710)
The Gambia	1995	Male	35 to 39	34% (19-53%)	66% (47-81%)	1.2 (0.56-2.2)	5.6% (3.6-8.7%)	7.3% (4.3-12%)	8.4 (4.8-13)	480 (300-740)
The Gambia	1995	Male	40 to 44	36% (21-54%)	64% (46-79%)	1.3 (0.53-2.5)	6.1% (3.7-9.6%)	8.4% (4.9-13%)	9.3 (5.3-15)	470 (270-730)
The Gambia	1995	Male	45 to 49	36% (19-55%)	64% (45-81%)	1.4 (0.54-2.8)	6.6% (3.8-11%)	9% (5-15%)	9.7 (5.2-16)	420 (230-690)
The Gambia	1995	Male	50 to 54	37% (20-57%)	63% (43-80%)	1.4 (0.55-2.7)	6.3% (3.2-11%)	8.3% (4.1-14%)	11 (5.5-19)	420 (220-720)
The Gambia	1995	Male	55 to 59	37% (21-57%)	63% (43-79%)	1.4 (0.55-2.7)	5.8% (2.9-9.9%)	7.5% (3.7-13%)	10 (5-18)	340 (170-580)
The Gambia	1995	Male	60 to 64	37% (20-57%)	63% (43-80%)	1.3 (0.44-2.6)	4.6% (2-8.7%)	5.7% (2.5-11%)	9.7 (4.2-19)	270 (120-510)
The Gambia	1995	Male	65 to 69	37% (21-55%)	63% (45-79%)	1 (0.4-2.1)	3.4% (1.4-6.9%)	4.2% (1.7-8.6%)	8.4 (3.4-17)	190 (83-400)
The Gambia	1995	Male	70 to 74	35% (20-53%)	65% (47-80%)	0.8 (0.3-1.7)	2.4% (0.67-5.3%)	2.9% (0.78-6.5%)	7.5 (1.9-18)	140 (38-320)
The Gambia	1995	Male	75 to 79	32% (18-51%)	68% (49-82%)	0.61 (0.18-1.4)	1.5% (0.25-3.9%)	1.8% (0.24-4.7%)	3.7 (0.47-9.6)	53 (8.1-130)
The Gambia	1995	Male	80 to 84	29% (15-49%)	71% (51-85%)	0.51 (0.16-1.1)	1.2% (0.2-3%)	1.4% (0.18-3.5%)	1.5 (0.18-3.6)	16 (2.6-38)
The Gambia	1995	Male	85 to 89	28% (15-45%)	72% (55-85%)	0.44 (0.14-0.98)	0.85% (-0.075-2.3%)	0.95% (-0.16-2.7%)	0.6 (-0.1-1.7)	4.7 (-0.42-13)
The Gambia	1995	Male	90 to 94	30% (16-48%)	70% (52-84%)	0.42 (0.12-1)	0.41% (-0.36-1.5%)	0.39% (-0.57-1.7%)	0.096 (-0.14-0.42)	0.61 (-0.54-2.2)
The Gambia	1995	Male	95 plus	34% (16-57%)	66% (43-84%)	0.42 (0.093-1.1)	0.38% (-0.6-2.1%)	0.29% (-0.9-2.4%)	0.014 (-0.044-0.11)	0.073 (-0.12-0.39)
The Gambia	2000	Female	All Ages	22% (17-29%)	78% (71-83%)	0.43 (0.32-0.57)	0.85% (0.49-1.3%)	1.4% (0.54-2.7%)	66 (26-130)	2900 (1700-4500)
The Gambia	2000	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.42 (0.17-0.81)	1% (0.48-2%)	1.1% (0.15-2.7%)	1 (0.14-2.6)	140 (63-260)
The Gambia	2000	Female	20 to 24	28% (14-45%)	72% (55-86%)	0.67 (0.29-1.2)	2.1% (1.1-3.6%)	2% (0.44-4.5%)	2.4 (0.54-5.6)	300 (150-530)
The Gambia	2000	Female	25 to 29	30% (15-48%)	70% (52-85%)	0.82 (0.36-1.4)	2.4% (1.3-3.9%)	2.6% (0.85-5.1%)	3.1 (1-6.2)	310 (160-510)
The Gambia	2000	Female	30 to 34	33% (17-52%)	67% (48-83%)	0.93 (0.44-1.6)	2.5% (1.4-4.2%)	3% (1.3-5.7%)	3.9 (1.7-7.4)	310 (170-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	2000	Female	35 to 39	36% (19-55%)	64% (45-81%)	1 (0.45-1.8)	2.9% (1.5-5%)	4% (1.8-7.4%)	4.9 (2.1-9.2)	300 (150-530)
The Gambia	2000	Female	40 to 44	38% (22-57%)	62% (43-78%)	1.1 (0.43-1.9)	3.3% (1.7-5.6%)	4.8% (2.2-8.5%)	5.4 (2.4-9.8)	280 (140-480)
The Gambia	2000	Female	45 to 49	39% (20-57%)	61% (43-80%)	1.1 (0.44-2.1)	3.5% (1.5-6.3%)	5.1% (2.1-9.5%)	5.6 (2.2-10)	250 (100-450)
The Gambia	2000	Female	50 to 54	39% (21-60%)	61% (40-79%)	1.1 (0.46-2.1)	3.9% (1.6-7.3%)	5.4% (2.2-10%)	5.4 (2.1-10)	200 (82-390)
The Gambia	2000	Female	55 to 59	39% (23-58%)	61% (42-77%)	1.2 (0.47-2.3)	4% (1.4-8.1%)	5.5% (1.8-11%)	8.1 (2.6-17)	260 (86-530)
The Gambia	2000	Female	60 to 64	40% (23-59%)	60% (41-77%)	1.2 (0.47-2.4)	3.4% (0.83-7.5%)	4.4% (1.1-9.5%)	7.1 (1.7-16)	190 (45-430)
The Gambia	2000	Female	65 to 69	41% (25-59%)	59% (41-75%)	1.1 (0.47-2.2)	2.7% (0.16-6.6%)	3.4% (0.14-8.3%)	7 (0.3-18)	160 (10-400)
The Gambia	2000	Female	70 to 74	39% (24-57%)	61% (43-76%)	0.93 (0.36-1.9)	2% (-0.46-5.9%)	2.5% (-0.62-7.3%)	6.6 (-1.7-19)	120 (-27-350)
The Gambia	2000	Female	75 to 79	35% (19-54%)	65% (46-81%)	0.75 (0.24-1.7)	1.2% (-0.85-4.7%)	1.4% (-1.1-5.6%)	3.4 (-2.5-14)	49 (-35-200)
The Gambia	2000	Female	80 to 84	33% (17-53%)	67% (47-83%)	0.68 (0.23-1.4)	0.78% (-0.84-3.4%)	0.92% (-1.1-4.1%)	1.3 (-1.5-6)	14 (-15-62)
The Gambia	2000	Female	85 to 89	33% (18-51%)	67% (49-82%)	0.68 (0.22-1.5)	0.51% (-1-3.1%)	0.58% (-1.3-3.6%)	0.5 (-1.1-3)	3.9 (-7.9-22)
The Gambia	2000	Female	90 to 94	36% (19-54%)	64% (46-81%)	0.71 (0.22-1.5)	0.3% (-1.4-3.2%)	0.29% (-1.8-3.8%)	0.084 (-0.5-1.1)	0.53 (-2.4-5.5)
The Gambia	2000	Female	95 plus	40% (21-61%)	60% (39-79%)	0.74 (0.2-1.7)	0.24% (-1.5-3.4%)	0.16% (-1.9-4%)	0.011 (-0.13-0.28)	0.068 (-0.39-0.96)
The Gambia	2000	Male	All Ages	19% (14-25%)	81% (75-86%)	0.53 (0.4-0.72)	1.3% (0.93-1.9%)	2.3% (1.4-3.6%)	120 (74-200)	5200 (3600-7100)
The Gambia	2000	Male	15 to 19	16% (8.1-26%)	84% (74-92%)	0.35 (0.13-0.7)	1.2% (0.64-2%)	1.2% (0.4-2.6%)	1.1 (0.36-2.4)	140 (77-250)
The Gambia	2000	Male	20 to 24	23% (12-39%)	77% (61-88%)	0.71 (0.29-1.3)	2.8% (1.6-4.7%)	2.8% (1.1-5.3%)	3.7 (1.4-7.1)	380 (200-620)
The Gambia	2000	Male	25 to 29	29% (15-46%)	71% (54-85%)	1 (0.45-1.9)	3.9% (2.1-6.3%)	4.3% (1.9-7.8%)	5.9 (2.5-10)	470 (250-750)
The Gambia	2000	Male	30 to 34	33% (17-52%)	67% (48-83%)	1.3 (0.62-2.2)	4.7% (2.7-7.3%)	5.6% (2.9-9.3%)	8.3 (4.4-14)	560 (340-880)
The Gambia	2000	Male	35 to 39	35% (20-54%)	65% (46-80%)	1.4 (0.63-2.6)	5.4% (3.3-8.3%)	6.8% (3.9-11%)	9.8 (5.6-15)	560 (340-870)
The Gambia	2000	Male	40 to 44	36% (21-56%)	64% (44-79%)	1.5 (0.6-2.8)	6.1% (3.5-9.4%)	8% (4.4-12%)	12 (6.3-18)	570 (330-890)
The Gambia	2000	Male	45 to 49	37% (19-55%)	63% (45-81%)	1.6 (0.64-3.1)	7% (3.8-11%)	9.2% (4.7-15%)	14 (7.1-22)	590 (320-960)
The Gambia	2000	Male	50 to 54	38% (21-58%)	62% (42-79%)	1.6 (0.64-3.2)	6.9% (3.4-11%)	8.9% (4.4-15%)	13 (6.4-22)	490 (250-830)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	2000	Male	55 to 59	38% (22-57%)	62% (43-78%)	1.6 (0.61-3.1)	6.5% (3.1-11%)	8.2% (3.9-14%)	15 (7.1-26)	490 (240-840)
The Gambia	2000	Male	60 to 64	37% (21-57%)	63% (43-79%)	1.4 (0.51-3)	5.1% (2.2-9.5%)	6.3% (2.7-12%)	13 (5.4-24)	350 (150-660)
The Gambia	2000	Male	65 to 69	37% (21-55%)	63% (45-79%)	1.2 (0.46-2.3)	3.9% (1.5-8%)	4.7% (1.8-9.6%)	11 (4.3-23)	260 (100-520)
The Gambia	2000	Male	70 to 74	36% (21-53%)	64% (47-79%)	0.92 (0.35-1.9)	2.8% (0.73-6.6%)	3.3% (0.85-7.9%)	9.4 (2.4-23)	170 (45-410)
The Gambia	2000	Male	75 to 79	32% (18-51%)	68% (49-82%)	0.7 (0.22-1.7)	1.8% (0.24-4.8%)	2.1% (0.21-5.8%)	5.6 (0.59-16)	80 (11-220)
The Gambia	2000	Male	80 to 84	30% (16-49%)	70% (51-84%)	0.59 (0.18-1.2)	1.5% (0.22-3.8%)	1.7% (0.21-4.5%)	2.3 (0.28-6.2)	25 (3.7-63)
The Gambia	2000	Male	85 to 89	29% (15-47%)	71% (53-85%)	0.52 (0.16-1.1)	1.1% (0.0087-3.1%)	1.2% (-0.081-3.7%)	0.9 (-0.063-2.7)	6.9 (0.057-20)
The Gambia	2000	Male	90 to 94	31% (16-49%)	69% (51-84%)	0.49 (0.15-1.2)	0.61% (-0.37-2.3%)	0.63% (-0.55-2.7%)	0.19 (-0.17-0.82)	1.1 (-0.67-4.3)
The Gambia	2000	Male	95 plus	35% (17-58%)	65% (42-83%)	0.5 (0.11-1.3)	0.67% (-0.58-3%)	0.65% (-0.92-3.5%)	0.049 (-0.07-0.27)	0.21 (-0.18-0.91)
The Gambia	2005	Female	All Ages	22% (17-29%)	78% (71-83%)	0.48 (0.36-0.65)	0.93% (0.54-1.4%)	1.6% (0.59-3.1%)	75 (28-140)	3200 (1900-5100)
The Gambia	2005	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.46 (0.19-0.87)	1% (0.52-1.8%)	1.1% (0.19-2.8%)	1 (0.17-2.5)	140 (70-250)
The Gambia	2005	Female	20 to 24	27% (14-44%)	73% (56-86%)	0.73 (0.32-1.3)	2% (1.1-3.3%)	1.8% (0.44-3.9%)	2.2 (0.57-4.9)	310 (170-510)
The Gambia	2005	Female	25 to 29	30% (15-47%)	70% (53-85%)	0.89 (0.4-1.6)	2.1% (1.2-3.3%)	2.1% (0.71-4.2%)	3.2 (1.1-6.3)	350 (200-550)
The Gambia	2005	Female	30 to 34	33% (17-52%)	67% (48-83%)	1 (0.49-1.8)	2.1% (1.2-3.4%)	2.3% (1-4.3%)	3.4 (1.5-6.4)	290 (160-470)
The Gambia	2005	Female	35 to 39	36% (20-56%)	64% (44-80%)	1.1 (0.5-1.9)	2.6% (1.4-4.4%)	3.4% (1.5-6.2%)	5.1 (2.3-9.4)	330 (180-550)
The Gambia	2005	Female	40 to 44	38% (22-57%)	62% (43-78%)	1.2 (0.49-2.1)	3% (1.6-5%)	4.3% (2-7.5%)	6.1 (2.8-11)	320 (160-550)
The Gambia	2005	Female	45 to 49	39% (21-58%)	61% (42-79%)	1.2 (0.48-2.3)	3.3% (1.5-6.2%)	4.9% (2-9.1%)	6.5 (2.6-12)	290 (130-540)
The Gambia	2005	Female	50 to 54	39% (21-60%)	61% (40-79%)	1.3 (0.52-2.5)	4.1% (1.8-7.4%)	5.8% (2.4-11%)	8.1 (3.3-15)	310 (130-570)
The Gambia	2005	Female	55 to 59	39% (23-58%)	61% (42-77%)	1.4 (0.54-2.6)	4.3% (1.5-8.3%)	5.9% (2-12%)	7.4 (2.5-15)	240 (80-470)
The Gambia	2005	Female	60 to 64	40% (23-59%)	60% (41-77%)	1.4 (0.52-2.7)	3.7% (0.85-8.2%)	5% (1.2-11%)	9.5 (2.3-21)	260 (61-570)
The Gambia	2005	Female	65 to 69	41% (24-59%)	59% (41-76%)	1.2 (0.51-2.3)	3% (0.27-7.1%)	3.8% (0.35-9.1%)	7.9 (0.71-19)	180 (15-440)
The Gambia	2005	Female	70 to 74	39% (24-57%)	61% (43-76%)	1 (0.39-2.2)	2.3% (-0.36-6.5%)	2.9% (-0.43-8.1%)	7.1 (-1.2-21)	130 (-20-370)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	2005	Female	75 to 79	35% (19-54%)	65% (46-81%)	0.84 (0.27-1.8)	1.4% (-0.85-5.4%)	1.7% (-1.1-6.3%)	4.3 (-2.8-16)	62 (-38-230)
The Gambia	2005	Female	80 to 84	33% (18-53%)	67% (47-82%)	0.75 (0.26-1.6)	0.95% (-0.85-4%)	1.1% (-1.1-4.8%)	1.7 (-1.7-7.6)	19 (-16-79)
The Gambia	2005	Female	85 to 89	33% (18-52%)	67% (48-82%)	0.74 (0.24-1.6)	0.68% (-1-3.6%)	0.78% (-1.3-4.3%)	0.77 (-1.3-4.4)	6 (-9.1-32)
The Gambia	2005	Female	90 to 94	36% (20-55%)	64% (45-80%)	0.78 (0.24-1.8)	0.47% (-1.4-3.5%)	0.49% (-1.8-4.3%)	0.19 (-0.68-1.6)	1.1 (-3.2-8.3)
The Gambia	2005	Female	95 plus	40% (20-61%)	60% (39-80%)	0.82 (0.21-1.9)	0.43% (-1.5-3.9%)	0.38% (-1.9-4.6%)	0.035 (-0.18-0.42)	0.16 (-0.53-1.4)
The Gambia	2005	Male	All Ages	19% (14-24%)	81% (76-86%)	0.59 (0.43-0.79)	1.6% (1.1-2.3%)	2.7% (1.7-4.2%)	160 (95-240)	6500 (4500-9000)
The Gambia	2005	Male	15 to 19	15% (7.9-26%)	85% (74-92%)	0.38 (0.14-0.75)	1.3% (0.72-2.2%)	1.3% (0.45-2.8%)	1.4 (0.47-3)	170 (93-290)
The Gambia	2005	Male	20 to 24	23% (12-39%)	77% (61-88%)	0.78 (0.32-1.4)	3% (1.6-4.9%)	3% (1.2-5.6%)	4.7 (1.9-9)	460 (250-770)
The Gambia	2005	Male	25 to 29	29% (15-46%)	71% (54-85%)	1.2 (0.5-2.1)	3.8% (2-6.4%)	4.2% (1.8-7.8%)	7.9 (3.5-14)	630 (350-1000)
The Gambia	2005	Male	30 to 34	32% (17-52%)	68% (48-83%)	1.4 (0.69-2.5)	4.3% (2.5-6.8%)	4.9% (2.6-8.3%)	9.5 (5.3-15)	640 (380-960)
The Gambia	2005	Male	35 to 39	35% (19-55%)	65% (45-81%)	1.6 (0.68-2.8)	5.1% (3.1-7.9%)	6.2% (3.6-10%)	12 (7.2-20)	710 (430-1100)
The Gambia	2005	Male	40 to 44	36% (21-55%)	64% (45-79%)	1.7 (0.66-3)	5.7% (3.3-8.8%)	7.2% (4-11%)	14 (7.8-22)	690 (400-1100)
The Gambia	2005	Male	45 to 49	37% (19-57%)	63% (43-81%)	1.8 (0.67-3.5)	6.9% (3.8-11%)	8.8% (4.7-14%)	17 (9.3-27)	740 (410-1200)
The Gambia	2005	Male	50 to 54	37% (21-59%)	63% (41-79%)	1.8 (0.72-3.6)	7.1% (3.6-12%)	9.2% (4.6-15%)	18 (9-30)	680 (350-1100)
The Gambia	2005	Male	55 to 59	37% (21-57%)	63% (43-79%)	1.7 (0.65-3.3)	7% (3.4-11%)	8.8% (4.2-15%)	17 (7.8-29)	560 (260-930)
The Gambia	2005	Male	60 to 64	37% (21-56%)	63% (44-79%)	1.6 (0.53-3.4)	5.7% (2.5-11%)	6.9% (3-13%)	18 (7.7-34)	490 (220-940)
The Gambia	2005	Male	65 to 69	37% (21-55%)	63% (45-79%)	1.3 (0.52-2.7)	4.4% (1.7-9%)	5.3% (2.1-11%)	14 (5.5-30)	320 (130-670)
The Gambia	2005	Male	70 to 74	35% (21-53%)	65% (47-79%)	1 (0.39-2.2)	3.1% (0.87-7.4%)	3.8% (1-9%)	11 (3.1-28)	210 (59-500)
The Gambia	2005	Male	75 to 79	32% (18-51%)	68% (49-82%)	0.79 (0.24-1.8)	2% (0.32-5.7%)	2.4% (0.34-6.8%)	6.7 (0.92-18)	95 (15-260)
The Gambia	2005	Male	80 to 84	30% (16-49%)	70% (51-84%)	0.66 (0.22-1.4)	1.7% (0.25-4.4%)	1.9% (0.24-5.2%)	3.1 (0.36-8.1)	33 (5.1-86)
The Gambia	2005	Male	85 to 89	28% (15-47%)	72% (53-85%)	0.56 (0.18-1.3)	1.2% (0.043-3.4%)	1.4% (-0.032-4%)	1.4 (-0.031-4)	11 (0.37-30)
The Gambia	2005	Male	90 to 94	30% (15-49%)	70% (51-85%)	0.53 (0.16-1.2)	0.71% (-0.33-2.7%)	0.75% (-0.51-3.1%)	0.27 (-0.18-1.2)	1.6 (-0.72-6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	2005	Male	95 plus	35% (16-56%)	65% (44-84%)	0.53 (0.11-1.4)	0.77% (-0.53-3.3%)	0.76% (-0.85-3.9%)	0.072 (-0.08-0.36)	0.29 (-0.2-1.3)
The Gambia	2010	Female	All Ages	22% (17-29%)	78% (71-83%)	0.49 (0.32-0.73)	0.99% (0.55-1.6%)	1.6% (0.52-3.3%)	80 (25-160)	3500 (1900-5700)
The Gambia	2010	Female	15 to 19	22% (11-36%)	78% (64-89%)	0.46 (0.17-0.94)	1% (0.52-1.8%)	1.1% (0.17-2.7%)	0.98 (0.16-2.4)	150 (78-280)
The Gambia	2010	Female	20 to 24	27% (14-44%)	73% (56-86%)	0.73 (0.3-1.4)	2% (1.1-3.2%)	1.7% (0.39-3.8%)	2 (0.46-4.6)	320 (180-510)
The Gambia	2010	Female	25 to 29	30% (16-47%)	70% (53-84%)	0.89 (0.39-1.7)	2% (1.2-3.2%)	1.8% (0.61-3.8%)	3 (1-6.1)	350 (200-560)
The Gambia	2010	Female	30 to 34	33% (17-52%)	67% (48-83%)	1 (0.48-1.9)	1.9% (1.1-3%)	1.9% (0.88-3.6%)	3.6 (1.6-6.8)	330 (190-530)
The Gambia	2010	Female	35 to 39	36% (20-55%)	64% (45-80%)	1.1 (0.46-2.1)	2.3% (1.2-3.9%)	2.9% (1.2-5.4%)	4.8 (2.1-8.8)	320 (170-540)
The Gambia	2010	Female	40 to 44	38% (22-57%)	62% (43-78%)	1.2 (0.49-2.2)	2.7% (1.4-4.8%)	3.9% (1.7-7%)	6.7 (3.1-12)	360 (180-620)
The Gambia	2010	Female	45 to 49	39% (20-57%)	61% (43-80%)	1.2 (0.44-2.4)	3% (1.4-5.7%)	4.4% (1.8-8.3%)	7.3 (3.1-14)	330 (150-620)
The Gambia	2010	Female	50 to 54	39% (21-60%)	61% (40-79%)	1.3 (0.48-2.5)	3.9% (1.6-7.4%)	5.6% (2.3-11%)	9.6 (3.9-19)	370 (150-700)
The Gambia	2010	Female	55 to 59	39% (22-58%)	61% (42-78%)	1.4 (0.49-2.6)	4.2% (1.3-8.4%)	5.8% (1.8-11%)	10 (3.3-21)	340 (110-690)
The Gambia	2010	Female	60 to 64	40% (23-59%)	60% (41-77%)	1.4 (0.49-2.8)	3.6% (0.68-8%)	4.8% (0.9-11%)	8.1 (1.7-19)	220 (43-520)
The Gambia	2010	Female	65 to 69	41% (24-60%)	59% (40-76%)	1.3 (0.5-2.5)	2.9% (0.016-7.2%)	3.7% (0.01-9.2%)	9.5 (0.027-25)	220 (1.3-560)
The Gambia	2010	Female	70 to 74	39% (24-57%)	61% (43-76%)	1 (0.38-2.2)	2.1% (-0.49-6.5%)	2.6% (-0.62-8.3%)	7 (-1.8-22)	130 (-30-400)
The Gambia	2010	Female	75 to 79	35% (19-53%)	65% (47-81%)	0.82 (0.26-1.8)	1.3% (-0.96-5%)	1.5% (-1.2-6.2%)	4 (-3.2-16)	57 (-44-230)
The Gambia	2010	Female	80 to 84	32% (17-51%)	68% (49-83%)	0.73 (0.24-1.6)	0.83% (-0.98-3.9%)	0.97% (-1.2-4.7%)	1.6 (-2.2-8)	18 (-22-84)
The Gambia	2010	Female	85 to 89	33% (18-50%)	67% (50-82%)	0.72 (0.22-1.5)	0.57% (-1.1-3.4%)	0.64% (-1.4-4.1%)	0.76 (-1.7-4.8)	6 (-12-36)
The Gambia	2010	Female	90 to 94	36% (20-54%)	64% (46-80%)	0.77 (0.23-1.7)	0.38% (-1.4-3.8%)	0.38% (-1.8-4.5%)	0.19 (-0.88-2.3)	1.2 (-4.1-12)
The Gambia	2010	Female	95 plus	41% (21-62%)	59% (38-79%)	0.82 (0.2-2.1)	0.35% (-1.6-4.5%)	0.27% (-2.1-5.4%)	0.036 (-0.29-0.71)	0.19 (-0.86-2.5)
The Gambia	2010	Male	All Ages	19% (14-24%)	81% (76-86%)	0.59 (0.39-0.87)	1.7% (1.1-2.5%)	2.9% (1.7-4.6%)	170 (99-280)	7100 (4700-10000)
The Gambia	2010	Male	15 to 19	15% (7.9-27%)	85% (73-92%)	0.38 (0.14-0.78)	1.3% (0.7-2.3%)	1.4% (0.44-2.9%)	1.6 (0.51-3.4)	200 (110-350)
The Gambia	2010	Male	20 to 24	23% (12-38%)	77% (62-88%)	0.77 (0.3-1.5)	3.1% (1.7-5.1%)	3.1% (1.2-6.2%)	5 (1.9-9.8)	510 (280-850)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	2010	Male	25 to 29	28% (15-45%)	72% (55-85%)	1.1 (0.47-2.2)	3.8% (2-6.3%)	4.1% (1.7-7.5%)	8.5 (3.8-16)	700 (380-1200)
The Gambia	2010	Male	30 to 34	32% (17-51%)	68% (49-83%)	1.4 (0.65-2.6)	4.1% (2.4-6.4%)	4.6% (2.4-7.9%)	11 (6-18)	770 (450-1200)
The Gambia	2010	Male	35 to 39	35% (19-53%)	65% (47-81%)	1.5 (0.64-2.9)	4.6% (2.8-7.3%)	5.5% (3-9.2%)	13 (7.2-20)	730 (440-1100)
The Gambia	2010	Male	40 to 44	36% (21-54%)	64% (46-79%)	1.6 (0.68-3.1)	5.3% (3.2-8.2%)	6.6% (3.8-10%)	16 (9-25)	800 (470-1200)
The Gambia	2010	Male	45 to 49	37% (19-56%)	63% (44-81%)	1.8 (0.66-3.7)	6.3% (3.6-10%)	8% (4.4-13%)	19 (10-31)	810 (450-1300)
The Gambia	2010	Male	50 to 54	37% (21-58%)	63% (42-79%)	1.8 (0.67-3.8)	6.8% (3.5-11%)	8.6% (4.5-14%)	20 (10-34)	750 (390-1300)
The Gambia	2010	Male	55 to 59	38% (22-58%)	62% (42-78%)	1.7 (0.61-3.5)	6.7% (3.1-11%)	8.5% (3.9-14%)	21 (9.3-35)	670 (310-1100)
The Gambia	2010	Male	60 to 64	37% (21-57%)	63% (43-79%)	1.6 (0.55-3.4)	5.5% (2.5-10%)	6.8% (3-13%)	17 (7.5-32)	470 (210-880)
The Gambia	2010	Male	65 to 69	37% (21-56%)	63% (44-79%)	1.3 (0.53-2.7)	4.2% (1.6-8.8%)	5.2% (2-11%)	16 (6.3-35)	380 (150-800)
The Gambia	2010	Male	70 to 74	35% (21-54%)	65% (46-79%)	1 (0.39-2.3)	3% (0.84-7.3%)	3.7% (0.99-8.8%)	12 (3.2-28)	210 (59-510)
The Gambia	2010	Male	75 to 79	32% (17-51%)	68% (49-83%)	0.78 (0.24-1.9)	2% (0.28-5.3%)	2.3% (0.27-6.2%)	6.6 (0.74-18)	94 (14-250)
The Gambia	2010	Male	80 to 84	30% (15-48%)	70% (52-85%)	0.65 (0.21-1.4)	1.6% (0.29-4.2%)	1.9% (0.32-5%)	3.2 (0.53-8.4)	34 (6-88)
The Gambia	2010	Male	85 to 89	28% (15-45%)	72% (55-85%)	0.56 (0.17-1.2)	1.2% (0.0087-3.2%)	1.3% (-0.081-3.7%)	1.5 (-0.092-4.4)	12 (0.088-33)
The Gambia	2010	Male	90 to 94	30% (15-48%)	70% (52-85%)	0.53 (0.16-1.3)	0.7% (-0.32-2.8%)	0.72% (-0.53-3.2%)	0.35 (-0.26-1.6)	2.1 (-0.96-8.1)
The Gambia	2010	Male	95 plus	35% (17-56%)	65% (44-83%)	0.53 (0.11-1.4)	0.75% (-0.47-3.3%)	0.74% (-0.77-3.8%)	0.083 (-0.085-0.43)	0.34 (-0.21-1.5)
The Gambia	2016	Female	All Ages	23% (17-29%)	77% (71-83%)	0.45 (0.23-0.78)	1.1% (0.51-1.9%)	1.7% (0.34-4%)	83 (18-200)	3800 (1800-6900)
The Gambia	2016	Female	15 to 19	21% (12-36%)	79% (64-88%)	0.42 (0.14-0.94)	0.96% (0.49-1.7%)	0.91% (0.093-2.5%)	0.93 (0.097-2.6)	170 (81-310)
The Gambia	2016	Female	20 to 24	27% (14-43%)	73% (57-86%)	0.66 (0.24-1.4)	2% (1.1-3.3%)	1.5% (0.26-3.9%)	1.9 (0.31-4.8)	340 (190-570)
The Gambia	2016	Female	25 to 29	30% (15-47%)	70% (53-85%)	0.81 (0.3-1.7)	2% (1.2-3.3%)	1.8% (0.52-4%)	2.7 (0.78-5.9)	360 (200-580)
The Gambia	2016	Female	30 to 34	33% (16-53%)	67% (47-84%)	0.91 (0.37-1.8)	2% (1.2-3.3%)	2.1% (0.79-4.3%)	3.6 (1.4-7.3)	350 (200-600)
The Gambia	2016	Female	35 to 39	36% (20-55%)	64% (45-80%)	0.99 (0.38-2.1)	2.3% (1.3-4.2%)	3.1% (1.2-6.2%)	5.5 (2.3-11)	380 (200-700)
The Gambia	2016	Female	40 to 44	38% (22-57%)	62% (43-78%)	1 (0.37-2.4)	2.7% (1.3-5%)	4% (1.7-7.9%)	6.7 (2.8-13)	370 (180-690)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	2016	Female	45 to 49	39% (20-58%)	61% (42-80%)	1.1 (0.36-2.5)	3% (1.3-6.2%)	4.6% (1.7-9.6%)	8.4 (3.1-17)	380 (170-760)
The Gambia	2016	Female	50 to 54	39% (22-60%)	61% (40-78%)	1.1 (0.38-2.6)	3.6% (1.4-7.4%)	5.3% (1.9-11%)	11 (4-23)	420 (160-860)
The Gambia	2016	Female	55 to 59	39% (23-58%)	61% (42-77%)	1.2 (0.4-2.7)	3.7% (0.99-8.3%)	5.3% (1.3-12%)	12 (3-27)	390 (100-870)
The Gambia	2016	Female	60 to 64	41% (23-60%)	59% (40-77%)	1.3 (0.38-2.9)	3.1% (0.34-7.7%)	4.2% (0.5-10%)	11 (1.3-26)	290 (34-720)
The Gambia	2016	Female	65 to 69	41% (24-59%)	59% (41-76%)	1.1 (0.39-2.5)	2.4% (-0.39-7.6%)	3.1% (-0.53-9.6%)	7.3 (-1.2-24)	170 (-27-540)
The Gambia	2016	Female	70 to 74	39% (23-58%)	61% (42-77%)	0.93 (0.3-2.2)	1.7% (-0.71-6.4%)	2.1% (-0.91-7.8%)	6.7 (-2.9-26)	120 (-53-460)
The Gambia	2016	Female	75 to 79	35% (19-54%)	65% (46-81%)	0.75 (0.22-1.8)	0.98% (-1.1-5%)	1.2% (-1.4-6%)	3.4 (-4.3-18)	50 (-58-260)
The Gambia	2016	Female	80 to 84	32% (17-52%)	68% (48-83%)	0.68 (0.19-1.6)	0.63% (-1.2-4.2%)	0.72% (-1.4-4.9%)	1.3 (-2.7-9.1)	14 (-27-95)
The Gambia	2016	Female	85 to 89	33% (18-51%)	67% (49-82%)	0.68 (0.19-1.7)	0.41% (-1.2-3.7%)	0.44% (-1.6-4.4%)	0.58 (-2.1-5.7)	4.9 (-15-42)
The Gambia	2016	Female	90 to 94	36% (20-54%)	64% (46-80%)	0.71 (0.19-1.8)	0.24% (-1.4-3.8%)	0.19% (-1.8-4.4%)	0.12 (-1.2-2.9)	0.93 (-5.5-14)
The Gambia	2016	Female	95 plus	40% (20-62%)	60% (38-80%)	0.76 (0.16-2)	0.23% (-1.7-4.2%)	0.12% (-2.1-4.8%)	0.022 (-0.4-0.9)	0.17 (-1.2-3.1)
The Gambia	2016	Male	All Ages	19% (14-25%)	81% (75-86%)	0.54 (0.28-0.93)	1.8% (1.1-2.8%)	2.9% (1.5-4.8%)	180 (94-300)	7700 (4600-12000)
The Gambia	2016	Male	15 to 19	15% (8.1-26%)	85% (74-92%)	0.35 (0.11-0.79)	1.2% (0.65-2.3%)	1.2% (0.33-2.8%)	1.7 (0.47-4)	230 (120-430)
The Gambia	2016	Male	20 to 24	23% (12-38%)	77% (62-88%)	0.7 (0.24-1.5)	2.9% (1.6-5.3%)	2.9% (1-6.4%)	5.3 (1.9-12)	560 (310-1000)
The Gambia	2016	Male	25 to 29	28% (15-45%)	72% (55-85%)	1 (0.37-2.2)	3.7% (2-6.6%)	4% (1.6-8%)	8.8 (3.6-17)	740 (390-1300)
The Gambia	2016	Male	30 to 34	32% (17-51%)	68% (49-83%)	1.3 (0.51-2.5)	4% (2.3-6.4%)	4.5% (2.2-7.8%)	12 (6-21)	840 (480-1300)
The Gambia	2016	Male	35 to 39	35% (19-53%)	65% (47-81%)	1.4 (0.51-2.9)	4.5% (2.7-7%)	5.3% (2.9-8.9%)	15 (8.5-25)	890 (530-1400)
The Gambia	2016	Male	40 to 44	36% (21-55%)	64% (45-79%)	1.5 (0.53-3.3)	5% (2.9-7.9%)	6.1% (3.4-9.7%)	16 (9-26)	820 (470-1300)
The Gambia	2016	Male	45 to 49	37% (19-56%)	63% (44-81%)	1.6 (0.5-3.7)	6% (3.2-9.8%)	7.5% (4-13%)	20 (11-35)	900 (480-1500)
The Gambia	2016	Male	50 to 54	37% (20-58%)	63% (42-80%)	1.6 (0.54-3.6)	6% (2.9-10%)	7.6% (3.7-13%)	21 (9.7-36)	780 (380-1400)
The Gambia	2016	Male	55 to 59	37% (21-57%)	63% (43-79%)	1.6 (0.49-3.6)	6% (2.8-11%)	7.7% (3.5-13%)	21 (9.4-38)	700 (320-1200)
The Gambia	2016	Male	60 to 64	37% (20-57%)	63% (43-80%)	1.4 (0.42-3.4)	5% (2-9.7%)	6.2% (2.5-12%)	19 (7.6-37)	540 (220-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
The Gambia	2016	Male	65 to 69	37% (21-55%)	63% (45-79%)	1.2 (0.39-2.9)	3.8% (1.3-8.2%)	4.7% (1.6-10%)	15 (5.1-33)	340 (120-750)
The Gambia	2016	Male	70 to 74	36% (21-53%)	64% (47-79%)	0.92 (0.28-2.3)	2.7% (0.73-6.9%)	3.3% (0.87-8.4%)	12 (3.2-32)	220 (60-580)
The Gambia	2016	Male	75 to 79	33% (18-52%)	67% (48-82%)	0.71 (0.2-1.8)	1.7% (0.26-5%)	2.1% (0.25-6%)	6.4 (0.79-18)	92 (14-260)
The Gambia	2016	Male	80 to 84	30% (16-48%)	70% (52-84%)	0.59 (0.17-1.4)	1.5% (0.23-3.9%)	1.7% (0.2-4.6%)	2.9 (0.33-8.2)	32 (4.9-86)
The Gambia	2016	Male	85 to 89	29% (15-46%)	71% (54-85%)	0.51 (0.14-1.3)	1.1% (0.013-3.2%)	1.2% (-0.087-3.7%)	1.5 (-0.11-4.5)	12 (0.14-34)
The Gambia	2016	Male	90 to 94	30% (15-49%)	70% (51-85%)	0.49 (0.12-1.3)	0.62% (-0.34-2.7%)	0.63% (-0.55-3.1%)	0.36 (-0.31-1.8)	2.1 (-1.2-9.2)
The Gambia	2016	Male	95 plus	35% (17-58%)	65% (42-83%)	0.49 (0.092-1.5)	0.66% (-0.46-3.3%)	0.62% (-0.74-3.8%)	0.098 (-0.12-0.6)	0.42 (-0.3-2.1)
Ghana	1990	Female	All Ages	18% (15-21%)	82% (79-85%)	0.93 (0.64-1.3)	2.3% (1.5-3.2%)	4.6% (2.8-6.5%)	3100 (1900-4400)	110000 (72000-150000)
Ghana	1990	Female	15 to 19	16% (11-22%)	84% (78-89%)	0.74 (0.38-1.3)	1.7% (0.97-2.9%)	2.1% (0.9-3.9%)	28 (12-52)	2900 (1600-4800)
Ghana	1990	Female	20 to 24	21% (14-29%)	79% (71-86%)	1.2 (0.63-2.1)	3.7% (2.1-5.9%)	4.3% (2-7.7%)	72 (34-130)	6500 (3700-10000)
Ghana	1990	Female	25 to 29	24% (16-33%)	76% (67-84%)	1.5 (0.79-2.5)	5.1% (3-7.8%)	6.4% (3.3-11%)	110 (57-180)	8000 (4700-12000)
Ghana	1990	Female	30 to 34	27% (17-38%)	73% (62-83%)	1.9 (0.99-3.1)	5.8% (3.6-8.4%)	7.8% (4.3-12%)	130 (72-190)	8200 (5100-12000)
Ghana	1990	Female	35 to 39	29% (20-39%)	71% (61-80%)	2 (1-3.4)	7.3% (4.4-11%)	10% (5.8-16%)	160 (95-260)	9200 (5500-14000)
Ghana	1990	Female	40 to 44	31% (21-41%)	69% (59-79%)	2.2 (1.1-3.5)	7.9% (4.6-12%)	11% (6.4-17%)	210 (120-310)	10000 (5900-15000)
Ghana	1990	Female	45 to 49	32% (21-43%)	68% (57-79%)	2.3 (1.2-3.8)	8.9% (5.2-13%)	12% (7-18%)	250 (140-370)	11000 (6000-16000)
Ghana	1990	Female	50 to 54	34% (24-45%)	66% (55-76%)	2.5 (1.3-4.1)	9.5% (5.3-14%)	13% (7.1-18%)	300 (160-450)	11000 (6200-17000)
Ghana	1990	Female	55 to 59	37% (26-48%)	63% (52-74%)	2.6 (1.4-4.4)	9.9% (5.6-14%)	13% (7.2-18%)	350 (190-530)	11000 (6300-17000)
Ghana	1990	Female	60 to 64	39% (27-51%)	61% (49-73%)	2.6 (1.3-4.5)	8.9% (4.7-14%)	11% (5.6-17%)	370 (180-570)	10000 (5000-15000)
Ghana	1990	Female	65 to 69	40% (29-50%)	60% (50-71%)	2.4 (1.3-4)	8.7% (4.3-14%)	10% (5-16%)	400 (190-650)	9100 (4300-15000)
Ghana	1990	Female	70 to 74	36% (26-47%)	64% (53-74%)	1.9 (1-3.3)	7.7% (3-14%)	9.1% (3.6-16%)	360 (140-640)	6500 (2600-11000)
Ghana	1990	Female	75 to 79	30% (20-42%)	70% (58-80%)	1.5 (0.71-2.6)	5.7% (1.8-11%)	6.6% (2.1-12%)	220 (71-410)	3000 (1000-5700)
Ghana	1990	Female	80 to 84	27% (18-37%)	73% (63-82%)	1.3 (0.66-2.2)	4.8% (1.6-9.1%)	5.5% (1.8-11%)	100 (32-190)	1000 (330-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	1990	Female	85 to 89	27% (18-38%)	73% (62-82%)	1.3 (0.65-2.3)	4.3% (1.3-8.5%)	5% (1.4-9.9%)	51 (15-100)	380 (110-750)
Ghana	1990	Female	90 to 94	29% (20-41%)	71% (59-80%)	1.4 (0.65-2.5)	4% (0.83-8.5%)	4.6% (0.89-10%)	18 (3.4-38)	91 (19-190)
Ghana	1990	Female	95 plus	32% (19-48%)	68% (52-81%)	1.5 (0.61-2.9)	3.7% (0.27-9%)	4.2% (0.14-11%)	4.5 (0.14-11)	16 (1.1-38)
Ghana	1990	Male	All Ages	27% (23-31%)	73% (69-77%)	2.1 (1.5-2.8)	4% (3.1-5.1%)	7.9% (5.9-10%)	6000 (4400-7800)	220000 (160000-270000)
Ghana	1990	Male	15 to 19	18% (12-25%)	82% (75-88%)	1.1 (0.55-1.8)	2.7% (1.5-4.3%)	3.4% (1.6-6%)	47 (22-83)	4300 (2400-7000)
Ghana	1990	Male	20 to 24	31% (21-41%)	69% (59-79%)	2.4 (1.3-4)	7.4% (4.8-11%)	9% (5.4-14%)	180 (100-270)	14000 (8700-20000)
Ghana	1990	Male	25 to 29	42% (30-53%)	58% (47-70%)	3.9 (2.3-6.1)	11% (7.2-15%)	13% (8.5-18%)	290 (190-390)	19000 (13000-26000)
Ghana	1990	Male	30 to 34	48% (36-60%)	52% (40-64%)	5 (3-7.9)	12% (8.3-16%)	15% (9.8-20%)	320 (220-430)	19000 (14000-25000)
Ghana	1990	Male	35 to 39	51% (38-61%)	49% (39-62%)	5.4 (3.1-8.7)	12% (8.7-17%)	16% (11-21%)	330 (230-450)	18000 (13000-24000)
Ghana	1990	Male	40 to 44	52% (40-62%)	48% (38-60%)	5.3 (2.9-8.2)	14% (10-19%)	19% (13-25%)	450 (310-610)	22000 (15000-29000)
Ghana	1990	Male	45 to 49	52% (39-62%)	48% (38-61%)	5 (2.8-7.9)	17% (12-22%)	22% (16-29%)	540 (380-720)	23000 (16000-30000)
Ghana	1990	Male	50 to 54	54% (44-65%)	46% (35-56%)	5.2 (2.9-8.2)	18% (13-24%)	23% (16-30%)	610 (420-820)	22000 (16000-30000)
Ghana	1990	Male	55 to 59	60% (49-69%)	40% (31-51%)	5.7 (3.3-9)	19% (14-24%)	24% (17-31%)	680 (480-900)	22000 (15000-29000)
Ghana	1990	Male	60 to 64	61% (49-72%)	39% (28-51%)	5.4 (3-8.7)	17% (12-23%)	21% (14-28%)	680 (460-920)	19000 (13000-25000)
Ghana	1990	Male	65 to 69	58% (46-67%)	42% (33-54%)	4.7 (2.6-7.4)	16% (10-22%)	19% (13-26%)	650 (410-900)	15000 (9500-20000)
Ghana	1990	Male	70 to 74	54% (43-64%)	46% (36-57%)	3.9 (2.2-6.3)	15% (9-21%)	18% (11-26%)	590 (340-840)	11000 (6100-15000)
Ghana	1990	Male	75 to 79	50% (37-63%)	50% (37-63%)	3.1 (1.5-5.4)	13% (5.9-19%)	15% (6.9-23%)	390 (180-600)	5400 (2500-8400)
Ghana	1990	Male	80 to 84	46% (34-58%)	54% (42-66%)	2.5 (1.3-4.1)	10% (4.9-16%)	12% (5.7-19%)	170 (81-270)	1800 (840-2800)
Ghana	1990	Male	85 to 89	44% (32-57%)	56% (43-68%)	2 (1.1-3.6)	7.9% (3.7-13%)	9.3% (4.3-15%)	71 (33-120)	520 (250-870)
Ghana	1990	Male	90 to 94	46% (33-59%)	54% (41-67%)	1.8 (0.86-3.2)	5.5% (2-11%)	6.4% (2.2-13%)	19 (6.4-37)	96 (33-180)
Ghana	1990	Male	95 plus	50% (33-68%)	50% (32-67%)	1.6 (0.66-3.2)	5.2% (1.1-12%)	6% (1.1-14%)	4.4 (0.83-9.9)	15 (3.4-34)
Ghana	1995	Female	All Ages	18% (16-21%)	82% (79-84%)	0.91 (0.62-1.3)	2.3% (1.5-3.2%)	4.3% (2.5-6.3%)	3200 (1800-4700)	110000 (72000-150000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	1995	Female	15 to 19	16% (10-22%)	84% (78-90%)	0.7 (0.37-1.2)	1.6% (0.95-2.6%)	1.9% (0.79-3.4%)	27 (11-50)	2900 (1700-4700)
Ghana	1995	Female	20 to 24	21% (14-29%)	79% (71-86%)	1.1 (0.6-2)	3.2% (1.9-5.2%)	3.5% (1.7-6.5%)	70 (34-130)	6700 (3900-11000)
Ghana	1995	Female	25 to 29	24% (16-33%)	76% (67-84%)	1.5 (0.74-2.4)	4.1% (2.4-6.3%)	4.7% (2.3-8.1%)	110 (56-180)	8500 (5100-13000)
Ghana	1995	Female	30 to 34	27% (18-38%)	73% (62-82%)	1.8 (0.96-3)	4.6% (2.8-7.1%)	5.7% (3.1-9.1%)	130 (72-210)	8900 (5300-13000)
Ghana	1995	Female	35 to 39	29% (20-39%)	71% (61-80%)	1.9 (0.98-3.2)	5.9% (3.4-9.1%)	7.7% (4.2-12%)	170 (98-270)	9800 (5800-15000)
Ghana	1995	Female	40 to 44	31% (22-41%)	69% (59-78%)	2.1 (1.1-3.4)	6.6% (3.8-9.6%)	8.9% (5.1-13%)	190 (110-290)	9400 (5600-14000)
Ghana	1995	Female	45 to 49	32% (21-43%)	68% (57-79%)	2.3 (1.1-3.6)	7.7% (4.5-11%)	10% (5.9-16%)	260 (140-390)	11000 (6400-16000)
Ghana	1995	Female	50 to 54	34% (24-45%)	66% (55-76%)	2.4 (1.2-4)	8.5% (4.6-12%)	11% (6.1-16%)	310 (160-470)	12000 (6200-17000)
Ghana	1995	Female	55 to 59	37% (26-48%)	63% (52-74%)	2.5 (1.3-4.2)	8.9% (4.9-13%)	11% (6.2-17%)	360 (190-550)	12000 (6100-18000)
Ghana	1995	Female	60 to 64	39% (27-52%)	61% (48-73%)	2.4 (1.3-4.2)	8% (3.9-13%)	9.8% (4.8-15%)	370 (170-590)	10000 (4800-16000)
Ghana	1995	Female	65 to 69	40% (29-50%)	60% (50-71%)	2.3 (1.2-3.8)	7.9% (3.5-13%)	9.4% (4.2-15%)	410 (180-680)	9200 (4100-15000)
Ghana	1995	Female	70 to 74	36% (27-47%)	64% (53-73%)	1.8 (0.98-3.1)	6.8% (2.6-12%)	8% (3-14%)	360 (140-660)	6600 (2500-12000)
Ghana	1995	Female	75 to 79	30% (20-42%)	70% (58-80%)	1.4 (0.67-2.4)	5% (1.4-9.8%)	5.8% (1.6-11%)	220 (62-450)	3100 (890-6200)
Ghana	1995	Female	80 to 84	27% (18-37%)	73% (63-82%)	1.2 (0.59-2.1)	4.1% (1.2-8.2%)	4.8% (1.3-9.5%)	100 (28-210)	1100 (300-2100)
Ghana	1995	Female	85 to 89	27% (17-38%)	73% (62-83%)	1.2 (0.61-2.2)	3.7% (0.86-8%)	4.3% (0.97-9.3%)	54 (12-120)	400 (94-860)
Ghana	1995	Female	90 to 94	29% (19-40%)	71% (60-81%)	1.3 (0.61-2.3)	3.3% (0.36-7.7%)	3.9% (0.35-9.1%)	18 (1.7-42)	92 (9.9-210)
Ghana	1995	Female	95 plus	32% (19-46%)	68% (54-81%)	1.3 (0.53-2.6)	3.1% (-0.0043-7.8%)	3.5% (-0.17-9.2%)	4.4 (-0.22-12)	15 (-0.022-40)
Ghana	1995	Male	All Ages	28% (24-32%)	72% (68-76%)	2 (1.4-2.8)	4.1% (3-5.1%)	7.8% (5.6-10%)	6400 (4600-8300)	23000 (17000-29000)
Ghana	1995	Male	15 to 19	18% (12-25%)	82% (75-88%)	1 (0.53-1.7)	2.5% (1.4-3.9%)	3.1% (1.5-5.3%)	47 (23-80)	4400 (2500-7000)
Ghana	1995	Male	20 to 24	31% (21-41%)	69% (59-79%)	2.3 (1.3-3.8)	6.6% (4.2-9.7%)	7.8% (4.5-12%)	170 (97-260)	14000 (8400-20000)
Ghana	1995	Male	25 to 29	42% (30-53%)	58% (47-70%)	3.8 (2.1-5.9)	9.2% (6.1-13%)	11% (6.9-16%)	300 (190-410)	20000 (14000-27000)
Ghana	1995	Male	30 to 34	48% (36-60%)	52% (40-64%)	4.8 (2.8-7.4)	9.6% (6.7-13%)	11% (7.4-16%)	340 (230-450)	21000 (15000-28000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	1995	Male	35 to 39	51% (40-62%)	49% (38-60%)	5.1 (2.9-8.1)	10% (6.9-14%)	12% (8.2-17%)	360 (250-480)	20000 (14000-26000)
Ghana	1995	Male	40 to 44	52% (40-62%)	48% (38-60%)	5 (2.8-7.8)	12% (8.3-16%)	15% (10-20%)	410 (280-550)	20000 (14000-26000)
Ghana	1995	Male	45 to 49	52% (38-62%)	48% (38-62%)	4.7 (2.6-7.4)	15% (10-20%)	19% (13-25%)	580 (390-770)	25000 (17000-32000)
Ghana	1995	Male	50 to 54	54% (43-65%)	46% (35-57%)	5 (2.8-7.9)	16% (11-21%)	20% (14-26%)	630 (440-850)	24000 (17000-31000)
Ghana	1995	Male	55 to 59	60% (49-69%)	40% (31-51%)	5.5 (3.1-8.7)	17% (12-22%)	22% (15-28%)	700 (480-920)	22000 (16000-29000)
Ghana	1995	Male	60 to 64	61% (48-72%)	39% (28-52%)	5.2 (2.9-8.5)	16% (11-21%)	20% (13-26%)	700 (460-940)	19000 (13000-25000)
Ghana	1995	Male	65 to 69	58% (46-68%)	42% (32-54%)	4.5 (2.5-7.2)	15% (9.3-20%)	18% (11-24%)	690 (420-970)	16000 (9700-22000)
Ghana	1995	Male	70 to 74	55% (43-65%)	45% (35-57%)	3.7 (2.1-6)	14% (7.8-20%)	16% (9.4-23%)	670 (380-980)	12000 (6900-18000)
Ghana	1995	Male	75 to 79	51% (37-63%)	49% (37-63%)	3 (1.5-5.1)	11% (5.3-18%)	14% (6.2-21%)	480 (220-750)	6700 (3000-10000)
Ghana	1995	Male	80 to 84	47% (34-59%)	53% (41-66%)	2.4 (1.3-4.1)	9.2% (4.2-15%)	11% (5.1-18%)	220 (99-360)	2200 (1000-3700)
Ghana	1995	Male	85 to 89	45% (32-58%)	55% (42-68%)	2 (1-3.5)	7% (3-12%)	8.1% (3.5-15%)	85 (36-150)	630 (280-1100)
Ghana	1995	Male	90 to 94	46% (33-59%)	54% (41-67%)	1.7 (0.81-3)	4.8% (1.6-9.7%)	5.5% (1.7-11%)	21 (6.7-43)	110 (36-220)
Ghana	1995	Male	95 plus	50% (33-67%)	50% (33-67%)	1.5 (0.59-3.1)	4.5% (0.9-10%)	5.1% (0.8-12%)	4.7 (0.73-11)	16 (3.1-38)
Ghana	2000	Female	All Ages	16% (14-18%)	84% (82-86%)	0.75 (0.51-1)	2% (1.3-2.8%)	3.6% (2-5.5%)	2900 (1600-4500)	100000 (65000-150000)
Ghana	2000	Female	15 to 19	13% (9.3-18%)	87% (82-91%)	0.56 (0.3-0.94)	1.3% (0.78-2.1%)	1.4% (0.62-2.6%)	22 (9.3-39)	2600 (1500-4200)
Ghana	2000	Female	20 to 24	17% (12-23%)	83% (77-88%)	0.87 (0.49-1.5)	2.4% (1.5-3.7%)	2.3% (1.1-4.1%)	49 (25-87)	5500 (3400-8300)
Ghana	2000	Female	25 to 29	19% (14-25%)	81% (75-86%)	1.1 (0.6-1.8)	2.7% (1.7-4.2%)	2.8% (1.4-4.6%)	84 (46-140)	7200 (4500-11000)
Ghana	2000	Female	30 to 34	21% (16-28%)	79% (72-84%)	1.3 (0.74-2.1)	3.1% (1.8-4.6%)	3.3% (1.8-5.4%)	110 (64-180)	7900 (4900-12000)
Ghana	2000	Female	35 to 39	25% (20-31%)	75% (69-80%)	1.6 (0.88-2.5)	4% (2.4-6%)	4.8% (2.7-7.7%)	160 (92-250)	9200 (5700-14000)
Ghana	2000	Female	40 to 44	28% (23-34%)	72% (66-77%)	1.8 (0.99-2.7)	4.8% (2.9-7%)	6.1% (3.6-9.1%)	190 (110-280)	9400 (5800-14000)
Ghana	2000	Female	45 to 49	28% (21-35%)	72% (65-79%)	1.8 (0.99-2.9)	5.7% (3.3-8.8%)	7.3% (4.1-11%)	210 (120-340)	9400 (5300-14000)
Ghana	2000	Female	50 to 54	29% (22-36%)	71% (64-78%)	1.9 (1-3.1)	6.6% (3.6-10%)	8.4% (4.6-13%)	290 (150-470)	11000 (5800-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	2000	Female	55 to 59	33% (26-40%)	67% (60-74%)	2 (1.1-3.4)	7.3% (3.8-11%)	9.1% (4.8-14%)	340 (170-550)	11000 (5500-18000)
Ghana	2000	Female	60 to 64	34% (25-43%)	66% (57-75%)	2 (1.1-3.4)	6.6% (3.1-11%)	8% (3.8-13%)	350 (160-580)	9600 (4600-16000)
Ghana	2000	Female	65 to 69	35% (26-44%)	65% (56-74%)	1.8 (1-3.1)	6.5% (2.9-11%)	7.7% (3.5-13%)	380 (170-670)	8600 (3800-15000)
Ghana	2000	Female	70 to 74	33% (25-41%)	67% (59-75%)	1.6 (0.84-2.6)	5.6% (2-10%)	6.6% (2.3-12%)	330 (110-620)	6100 (2200-11000)
Ghana	2000	Female	75 to 79	28% (20-37%)	72% (63-80%)	1.2 (0.6-2.1)	4.2% (1.1-8.2%)	4.9% (1.2-9.7%)	210 (54-430)	3000 (800-6100)
Ghana	2000	Female	80 to 84	25% (17-34%)	75% (66-83%)	1.1 (0.55-1.8)	3.5% (0.95-7.2%)	4% (1-8.4%)	110 (28-230)	1100 (300-2400)
Ghana	2000	Female	85 to 89	25% (17-35%)	75% (65-83%)	1.1 (0.52-1.9)	3.1% (0.61-7.1%)	3.6% (0.66-8.2%)	58 (11-130)	430 (87-960)
Ghana	2000	Female	90 to 94	27% (18-37%)	73% (63-82%)	1.1 (0.55-2)	2.8% (0.27-6.6%)	3.2% (0.23-7.7%)	20 (1.4-48)	100 (9.9-240)
Ghana	2000	Female	95 plus	29% (17-43%)	71% (57-83%)	1.2 (0.48-2.2)	2.6% (-0.12-7%)	2.9% (-0.29-8%)	4.7 (-0.47-13)	17 (-0.73-45)
Ghana	2000	Male	All Ages	28% (25-31%)	72% (69-75%)	2 (1.4-2.7)	4.4% (3.3-5.6%)	8% (5.7-10%)	7300 (5200-9500)	260000 (190000-330000)
Ghana	2000	Male	15 to 19	14% (9.9-19%)	86% (81-90%)	0.75 (0.41-1.2)	2.1% (1.3-3.3%)	2.4% (1.2-4.3%)	42 (21-75)	4200 (2500-6600)
Ghana	2000	Male	20 to 24	25% (19-32%)	75% (68-81%)	1.7 (0.99-2.8)	5.6% (3.5-8.2%)	6.4% (3.7-9.8%)	160 (89-250)	13000 (7900-19000)
Ghana	2000	Male	25 to 29	40% (33-46%)	60% (54-67%)	3.4 (2-5.2)	8.5% (5.8-12%)	10% (6.3-14%)	320 (210-460)	22000 (15000-30000)
Ghana	2000	Male	30 to 34	50% (43-56%)	50% (44-57%)	4.7 (3-7)	9.3% (6.6-12%)	11% (7.3-15%)	410 (280-560)	25000 (18000-34000)
Ghana	2000	Male	35 to 39	56% (50-61%)	44% (39-50%)	5.4 (3.3-8.1)	9.9% (6.9-13%)	12% (7.8-16%)	460 (320-600)	25000 (18000-32000)
Ghana	2000	Male	40 to 44	57% (51-62%)	43% (38-49%)	5.3 (3.2-8.1)	11% (8.3-15%)	14% (9.7-17%)	530 (370-690)	25000 (18000-33000)
Ghana	2000	Male	45 to 49	54% (47-60%)	46% (40-53%)	4.7 (2.8-7.3)	14% (9.5-18%)	17% (12-22%)	620 (430-810)	26000 (18000-34000)
Ghana	2000	Male	50 to 54	57% (50-63%)	43% (37-50%)	5 (2.9-7.8)	15% (11-20%)	19% (13-24%)	770 (520-1000)	29000 (20000-38000)
Ghana	2000	Male	55 to 59	63% (57-69%)	37% (31-43%)	5.6 (3.3-8.9)	17% (12-22%)	21% (14-26%)	860 (580-1100)	28000 (19000-36000)
Ghana	2000	Male	60 to 64	63% (55-70%)	37% (30-45%)	5.1 (3-8.1)	16% (9.8-21%)	19% (12-25%)	830 (510-1100)	23000 (14000-30000)
Ghana	2000	Male	65 to 69	56% (47-65%)	44% (35-53%)	4.1 (2.5-6.5)	14% (8.4-18%)	16% (9.9-22%)	740 (450-1000)	17000 (10000-23000)
Ghana	2000	Male	70 to 74	51% (41-60%)	49% (40-59%)	3.2 (1.8-5.2)	12% (6.4-18%)	14% (7.5-21%)	680 (350-1000)	12000 (6500-18000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	2000	Male	75 to 79	48% (37-60%)	52% (40-63%)	2.7 (1.4-4.4)	10% (4.5-16%)	12% (5.3-19%)	500 (220-800)	7000 (3200-11000)
Ghana	2000	Male	80 to 84	47% (35-59%)	53% (41-65%)	2.3 (1.2-3.8)	8.3% (3.8-14%)	9.7% (4.4-17%)	260 (120-440)	2600 (1200-4500)
Ghana	2000	Male	85 to 89	46% (33-59%)	54% (41-67%)	1.9 (1-3.4)	6.4% (2.5-12%)	7.4% (2.8-14%)	110 (40-200)	780 (310-1400)
Ghana	2000	Male	90 to 94	47% (34-60%)	53% (40-66%)	1.7 (0.84-3)	4.4% (1.3-9.3%)	5.1% (1.4-11%)	25 (6.8-54)	130 (38-270)
Ghana	2000	Male	95 plus	51% (34-68%)	49% (32-66%)	1.5 (0.63-2.9)	4.1% (0.82-9.6%)	4.6% (0.69-11%)	5.3 (0.79-13)	19 (3.6-44)
Ghana	2005	Female	All Ages	16% (15-18%)	84% (82-85%)	0.81 (0.55-1.1)	2% (1.3-2.8%)	3.6% (1.9-5.5%)	3300 (1700-5200)	110000 (70000-160000)
Ghana	2005	Female	15 to 19	12% (8.9-17%)	88% (83-91%)	0.52 (0.27-0.88)	1.2% (0.73-1.9%)	1.2% (0.53-2.2%)	20 (8.8-35)	2600 (1600-4100)
Ghana	2005	Female	20 to 24	17% (14-22%)	83% (78-86%)	0.91 (0.51-1.5)	2.3% (1.5-3.6%)	2.1% (1-3.8%)	48 (25-82)	5800 (3700-8500)
Ghana	2005	Female	25 to 29	20% (16-24%)	80% (76-84%)	1.1 (0.65-1.9)	2.5% (1.6-3.6%)	2.4% (1.2-4.1%)	79 (44-120)	7200 (4800-10000)
Ghana	2005	Female	30 to 34	23% (19-28%)	77% (72-81%)	1.5 (0.86-2.2)	2.7% (1.7-3.9%)	2.8% (1.5-4.4%)	120 (65-170)	8400 (5500-12000)
Ghana	2005	Female	35 to 39	27% (22-31%)	73% (69-78%)	1.7 (1-2.7)	3.4% (2.2-5.1%)	4% (2.3-6.1%)	170 (100-260)	10000 (6300-14000)
Ghana	2005	Female	40 to 44	28% (23-33%)	72% (67-77%)	1.8 (0.99-2.8)	4.1% (2.5-5.9%)	4.9% (3-7.4%)	200 (120-300)	10000 (6400-15000)
Ghana	2005	Female	45 to 49	29% (24-35%)	71% (65-76%)	1.9 (1.1-3)	5.2% (3-7.7%)	6.5% (3.6-9.8%)	250 (130-390)	11000 (6300-17000)
Ghana	2005	Female	50 to 54	32% (26-39%)	68% (61-74%)	2.1 (1.2-3.4)	6.5% (3.5-9.8%)	8.2% (4.4-13%)	300 (150-480)	12000 (6000-18000)
Ghana	2005	Female	55 to 59	37% (30-44%)	63% (56-70%)	2.4 (1.4-3.9)	7.5% (4.1-11%)	9.5% (5.1-14%)	400 (200-640)	13000 (6500-21000)
Ghana	2005	Female	60 to 64	36% (29-43%)	64% (57-71%)	2.2 (1.2-3.7)	6.5% (3.1-10%)	8% (3.7-12%)	380 (170-630)	10000 (4700-17000)
Ghana	2005	Female	65 to 69	35% (28-43%)	65% (57-72%)	1.9 (1.1-3.1)	6.2% (2.6-10%)	7.5% (3.1-13%)	430 (170-730)	9800 (3900-17000)
Ghana	2005	Female	70 to 74	32% (26-40%)	68% (60-74%)	1.5 (0.84-2.5)	5.3% (1.7-10%)	6.2% (2-12%)	410 (130-760)	7500 (2300-14000)
Ghana	2005	Female	75 to 79	25% (19-33%)	75% (67-81%)	1.1 (0.55-1.9)	3.7% (0.88-7.3%)	4.3% (1-8.5%)	270 (57-540)	3800 (860-7600)
Ghana	2005	Female	80 to 84	24% (16-32%)	76% (68-84%)	1 (0.52-1.7)	3.1% (0.74-6.6%)	3.6% (0.84-7.7%)	150 (33-310)	1600 (370-3200)
Ghana	2005	Female	85 to 89	25% (17-34%)	75% (66-83%)	1.1 (0.54-1.9)	2.9% (0.58-6.4%)	3.3% (0.63-7.5%)	78 (15-170)	580 (120-1300)
Ghana	2005	Female	90 to 94	27% (18-36%)	73% (64-82%)	1.1 (0.56-2)	2.6% (0.21-6.5%)	3% (0.18-7.7%)	28 (1.6-73)	150 (12-370)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	2005	Female	95 plus	29% (18-42%)	71% (58-82%)	1.2 (0.5-2.3)	2.5% (-0.16-6.8%)	2.7% (-0.34-7.9%)	6.9 (-0.84-20)	24 (-1.4-69)
Ghana	2005	Male	All Ages	29% (26-31%)	71% (69-74%)	2.1 (1.4-2.8)	4.6% (3.4-5.8%)	8.3% (5.8-11%)	8400 (5900-11000)	300000 (220000-370000)
Ghana	2005	Male	15 to 19	11% (8.4-15%)	89% (85-92%)	0.61 (0.34-1)	1.8% (1.1-2.7%)	1.9% (1-3.3%)	39 (19-70)	4100 (2400-6400)
Ghana	2005	Male	20 to 24	27% (22-32%)	73% (68-78%)	1.9 (1.1-3.1)	5.8% (3.8-8.1%)	6.6% (3.9-9.8%)	180 (100-280)	15000 (9300-21000)
Ghana	2005	Male	25 to 29	44% (39-49%)	56% (51-61%)	3.9 (2.4-5.9)	9.1% (6.2-12%)	11% (6.9-15%)	370 (240-500)	25000 (17000-34000)
Ghana	2005	Male	30 to 34	50% (45-56%)	50% (44-55%)	4.9 (3.1-7.1)	9.1% (6.5-12%)	11% (7.3-14%)	440 (310-590)	27000 (20000-36000)
Ghana	2005	Male	35 to 39	53% (48-58%)	47% (42-52%)	5.1 (3.1-7.7)	9.5% (6.9-12%)	11% (7.9-15%)	510 (370-670)	28000 (21000-36000)
Ghana	2005	Male	40 to 44	55% (50-60%)	45% (40-50%)	5.1 (3-7.8)	11% (7.8-14%)	13% (9.2-16%)	600 (420-790)	29000 (21000-38000)
Ghana	2005	Male	45 to 49	53% (49-58%)	47% (42-51%)	4.7 (2.9-7.1)	13% (9.3-17%)	16% (11-21%)	730 (500-960)	31000 (22000-40000)
Ghana	2005	Male	50 to 54	57% (53-62%)	43% (38-47%)	5.2 (3-7.7)	15% (10-19%)	18% (12-23%)	770 (520-1000)	29000 (20000-37000)
Ghana	2005	Male	55 to 59	65% (61-70%)	35% (30-39%)	5.9 (3.5-9.1)	17% (12-21%)	20% (14-26%)	1000 (690-1300)	32000 (22000-42000)
Ghana	2005	Male	60 to 64	65% (58-71%)	35% (29-42%)	5.4 (3.2-8.7)	16% (10-20%)	18% (12-24%)	960 (620-1200)	26000 (17000-34000)
Ghana	2005	Male	65 to 69	60% (53-66%)	40% (34-47%)	4.5 (2.7-6.9)	14% (8.5-19%)	16% (10-22%)	890 (540-1200)	20000 (12000-28000)
Ghana	2005	Male	70 to 74	55% (47-62%)	45% (38-53%)	3.6 (2.1-5.7)	12% (6.8-18%)	15% (8-21%)	830 (440-1200)	15000 (8400-22000)
Ghana	2005	Male	75 to 79	52% (43-61%)	48% (39-57%)	3 (1.6-4.8)	10% (4.5-16%)	12% (5.3-19%)	610 (270-960)	8500 (3800-13000)
Ghana	2005	Male	80 to 84	50% (38-61%)	50% (39-62%)	2.6 (1.4-4.1)	8.5% (3.8-14%)	9.9% (4.5-17%)	310 (140-530)	3300 (1500-5400)
Ghana	2005	Male	85 to 89	48% (36-60%)	52% (40-64%)	2.1 (1.1-3.6)	6.4% (2.5-12%)	7.4% (2.9-14%)	140 (54-260)	1000 (420-1900)
Ghana	2005	Male	90 to 94	50% (37-62%)	50% (38-63%)	1.8 (0.94-3.2)	4.4% (1.3-9.8%)	5.1% (1.5-11%)	35 (10-79)	180 (54-400)
Ghana	2005	Male	95 plus	54% (37-70%)	46% (30-63%)	1.6 (0.69-3.1)	4% (0.68-9.5%)	4.5% (0.85-11%)	6.8 (0.85-17)	24 (4.1-57)
Ghana	2010	Female	All Ages	18% (16-21%)	82% (79-84%)	1 (0.67-1.4)	2.4% (1.6-3.4%)	4.4% (2.3-6.6%)	4500 (2400-7000)	150000 (93000-210000)
Ghana	2010	Female	15 to 19	14% (9.8-19%)	86% (81-90%)	0.67 (0.35-1.2)	1.3% (0.8-2%)	1.3% (0.61-2.3%)	23 (11-41)	3000 (1800-4800)
Ghana	2010	Female	20 to 24	19% (14-25%)	81% (75-86%)	1.1 (0.6-2)	2.6% (1.7-3.7%)	2.4% (1.2-4.1%)	52 (27-87)	6500 (4100-9400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	2010	Female	25 to 29	21% (16-28%)	79% (72-84%)	1.4 (0.76-2.3)	2.9% (1.8-4.1%)	2.9% (1.5-4.8%)	89 (50-140)	8400 (5500-12000)
Ghana	2010	Female	30 to 34	25% (19-32%)	75% (68-81%)	1.8 (0.99-2.8)	3.1% (2-4.5%)	3.4% (1.9-5.2%)	140 (80-210)	10000 (6500-14000)
Ghana	2010	Female	35 to 39	28% (22-34%)	72% (66-78%)	2 (1.1-3.3)	3.8% (2.4-5.4%)	4.4% (2.6-6.7%)	200 (120-290)	12000 (7700-17000)
Ghana	2010	Female	40 to 44	29% (23-36%)	71% (64-77%)	2.1 (1.1-3.3)	4.5% (2.8-6.4%)	5.6% (3.4-8.1%)	250 (150-360)	13000 (8000-18000)
Ghana	2010	Female	45 to 49	31% (24-38%)	69% (62-76%)	2.3 (1.2-3.7)	5.8% (3.3-8.4%)	7.5% (4.2-11%)	320 (180-470)	14000 (8000-21000)
Ghana	2010	Female	50 to 54	34% (26-42%)	66% (58-74%)	2.6 (1.4-4.3)	7.5% (4.2-11%)	9.6% (5.4-14%)	430 (220-660)	16000 (8900-25000)
Ghana	2010	Female	55 to 59	38% (30-47%)	62% (53-70%)	2.8 (1.6-4.8)	8.5% (4.7-13%)	11% (6-16%)	490 (260-760)	16000 (8700-24000)
Ghana	2010	Female	60 to 64	39% (30-48%)	61% (52-70%)	2.7 (1.4-4.8)	7.7% (3.7-12%)	9.3% (4.5-15%)	500 (240-800)	14000 (6700-22000)
Ghana	2010	Female	65 to 69	38% (29-47%)	62% (53-71%)	2.3 (1.3-4)	7.4% (3.2-12%)	8.9% (3.8-14%)	520 (220-880)	12000 (5100-20000)
Ghana	2010	Female	70 to 74	34% (26-42%)	66% (58-74%)	1.8 (0.96-3)	6.2% (2.2-11%)	7.3% (2.5-13%)	560 (190-1000)	10000 (3600-18000)
Ghana	2010	Female	75 to 79	27% (19-37%)	73% (63-81%)	1.3 (0.63-2.4)	4.5% (1.2-8.6%)	5.2% (1.4-9.9%)	450 (120-870)	6500 (1700-12000)
Ghana	2010	Female	80 to 84	25% (17-34%)	75% (66-83%)	1.2 (0.61-2.1)	3.9% (0.91-7.8%)	4.5% (1-9%)	290 (65-610)	3100 (720-6300)
Ghana	2010	Female	85 to 89	26% (17-36%)	74% (64-83%)	1.3 (0.62-2.3)	3.6% (0.71-7.6%)	4.2% (0.76-8.9%)	140 (25-300)	1000 (200-2200)
Ghana	2010	Female	90 to 94	28% (19-38%)	72% (62-81%)	1.3 (0.64-2.5)	3.3% (0.36-7.7%)	3.9% (0.33-9.2%)	47 (3.9-110)	240 (26-570)
Ghana	2010	Female	95 plus	31% (19-45%)	69% (55-81%)	1.4 (0.59-2.8)	3.2% (0.03-8%)	3.6% (-0.12-9.4%)	12 (-0.39-33)	43 (0.39-110)
Ghana	2010	Male	All Ages	29% (26-32%)	71% (68-74%)	2.4 (1.6-3.3)	5% (3.8-6.2%)	8.9% (6.5-11%)	9500 (6900-12000)	330000 (250000-410000)
Ghana	2010	Male	15 to 19	13% (9.2-18%)	87% (82-91%)	0.79 (0.42-1.4)	2% (1.2-3.2%)	2.2% (1.1-4%)	51 (24-95)	5200 (3000-8400)
Ghana	2010	Male	20 to 24	29% (24-36%)	71% (64-76%)	2.4 (1.3-4)	6.4% (4.2-8.9%)	7.4% (4.2-11%)	220 (120-330)	18000 (11000-25000)
Ghana	2010	Male	25 to 29	45% (39-52%)	55% (48-61%)	4.5 (2.7-6.9)	9.8% (7-13%)	12% (8-16%)	400 (260-550)	28000 (19000-37000)
Ghana	2010	Male	30 to 34	50% (44-57%)	50% (43-56%)	5.5 (3.4-8.2)	10% (7.6-13%)	12% (8.7-15%)	470 (340-630)	29000 (21000-39000)
Ghana	2010	Male	35 to 39	52% (46-58%)	48% (42-54%)	5.6 (3.3-8.7)	10% (7.6-13%)	12% (8.8-16%)	540 (400-720)	30000 (22000-39000)
Ghana	2010	Male	40 to 44	54% (47-59%)	46% (41-53%)	5.6 (3.2-8.7)	12% (8.6-14%)	14% (10-17%)	670 (480-870)	33000 (24000-42000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	2010	Male	45 to 49	53% (47-59%)	47% (41-53%)	5.3 (3.1-8.2)	14% (11-18%)	17% (13-22%)	840 (610-1100)	36000 (26000-46000)
Ghana	2010	Male	50 to 54	57% (51-63%)	43% (37-49%)	5.8 (3.4-9.2)	15% (11-19%)	19% (14-23%)	910 (640-1200)	34000 (24000-43000)
Ghana	2010	Male	55 to 59	65% (59-70%)	35% (30-41%)	6.6 (3.9-10)	17% (13-21%)	21% (15-25%)	1000 (710-1300)	32000 (24000-41000)
Ghana	2010	Male	60 to 64	65% (57-72%)	35% (28-43%)	6.1 (3.5-10)	16% (11-20%)	19% (13-24%)	1100 (780-1400)	31000 (22000-39000)
Ghana	2010	Male	65 to 69	61% (53-68%)	39% (32-47%)	5.2 (3-8.1)	15% (9.6-19%)	17% (11-23%)	1000 (660-1300)	23000 (15000-30000)
Ghana	2010	Male	70 to 74	57% (48-65%)	43% (35-52%)	4.3 (2.4-6.9)	13% (7.4-19%)	16% (8.8-22%)	950 (520-1400)	17000 (9600-25000)
Ghana	2010	Male	75 to 79	54% (44-63%)	46% (37-56%)	3.5 (1.8-5.9)	11% (5-16%)	13% (5.7-19%)	720 (330-1100)	10000 (4700-15000)
Ghana	2010	Male	80 to 84	51% (39-61%)	49% (39-61%)	2.9 (1.5-4.8)	9.1% (4-15%)	11% (4.6-17%)	390 (170-640)	4100 (1700-6700)
Ghana	2010	Male	85 to 89	48% (36-60%)	52% (40-64%)	2.3 (1.2-3.9)	6.9% (2.7-13%)	8% (3.1-15%)	170 (66-310)	1300 (500-2300)
Ghana	2010	Male	90 to 94	49% (36-62%)	51% (38-64%)	2 (0.97-3.6)	4.9% (1.4-10%)	5.6% (1.5-12%)	48 (13-100)	250 (74-530)
Ghana	2010	Male	95 plus	54% (36-69%)	46% (31-64%)	1.8 (0.73-3.6)	4.3% (0.73-10%)	4.8% (0.6-12%)	9.8 (1.2-24)	35 (5.8-82)
Ghana	2016	Female	All Ages	20% (17-23%)	80% (77-83%)	1.3 (0.68-2.1)	3.1% (1.8-4.4%)	5.5% (2.7-8.4%)	5000 (2400-7700)	160000 (95000-230000)
Ghana	2016	Female	15 to 19	16% (11-22%)	84% (78-89%)	0.88 (0.38-1.7)	1.5% (0.82-2.3%)	1.5% (0.61-2.9%)	24 (9.2-45)	3300 (1800-5300)
Ghana	2016	Female	20 to 24	21% (14-30%)	79% (70-86%)	1.4 (0.62-2.8)	3% (1.9-4.3%)	2.8% (1.3-4.8%)	50 (21-86)	6800 (4100-9800)
Ghana	2016	Female	25 to 29	24% (16-33%)	76% (67-84%)	1.8 (0.8-3.5)	3.4% (2.1-4.9%)	3.6% (1.8-5.9%)	84 (42-130)	8500 (5400-12000)
Ghana	2016	Female	30 to 34	27% (18-39%)	73% (61-82%)	2.2 (1-4.1)	3.7% (2.4-5.2%)	4.3% (2.4-6.6%)	130 (74-200)	10000 (6700-15000)
Ghana	2016	Female	35 to 39	30% (20-40%)	70% (60-80%)	2.4 (1-4.5)	4.4% (2.8-6.3%)	5.7% (3.2-8.4%)	210 (120-310)	13000 (8000-18000)
Ghana	2016	Female	40 to 44	31% (22-41%)	69% (59-78%)	2.6 (1.2-4.7)	5.2% (3.1-7.4%)	7% (3.9-10%)	280 (150-410)	14000 (8400-21000)
Ghana	2016	Female	45 to 49	33% (22-43%)	67% (57-78%)	2.8 (1.2-5.3)	6.6% (3.8-9.5%)	8.9% (5.1-13%)	370 (200-550)	16000 (9400-24000)
Ghana	2016	Female	50 to 54	35% (25-46%)	65% (54-75%)	3 (1.3-5.8)	7.8% (4.3-11%)	10% (5.8-15%)	470 (250-710)	18000 (9900-27000)
Ghana	2016	Female	55 to 59	37% (27-49%)	63% (51-73%)	3.1 (1.4-6)	8.5% (4.3-13%)	11% (5.6-16%)	560 (270-880)	18000 (9100-29000)
Ghana	2016	Female	60 to 64	39% (28-52%)	61% (48-72%)	3.1 (1.4-6.2)	8% (3.3-12%)	9.9% (4.2-15%)	550 (230-890)	15000 (6500-25000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	2016	Female	65 to 69	40% (29-51%)	60% (49-71%)	2.9 (1.3-5.4)	8.2% (3.2-13%)	10% (3.9-16%)	640 (250-1100)	15000 (5800-25000)
Ghana	2016	Female	70 to 74	36% (27-47%)	64% (53-73%)	2.3 (1-4.4)	7.1% (2.2-12%)	8.4% (2.6-15%)	540 (160-980)	10000 (2900-18000)
Ghana	2016	Female	75 to 79	30% (20-41%)	70% (59-80%)	1.7 (0.68-3.4)	5.4% (1.2-10%)	6.3% (1.3-12%)	440 (87-850)	6300 (1300-12000)
Ghana	2016	Female	80 to 84	27% (18-37%)	73% (63-82%)	1.5 (0.65-2.9)	4.7% (0.95-9%)	5.5% (1.1-11%)	320 (59-630)	3400 (640-6600)
Ghana	2016	Female	85 to 89	27% (18-38%)	73% (62-82%)	1.5 (0.62-2.9)	4.3% (0.76-9%)	5.1% (0.81-10%)	220 (35-470)	1600 (290-3400)
Ghana	2016	Female	90 to 94	29% (19-41%)	71% (59-81%)	1.6 (0.66-3.3)	4.1% (0.31-8.9%)	4.7% (0.23-11%)	71 (3.5-160)	370 (28-810)
Ghana	2016	Female	95 plus	32% (20-47%)	68% (53-80%)	1.7 (0.61-3.8)	4% (0.011-9.5%)	4.5% (-0.17-11%)	17 (-0.66-42)	58 (0.17-140)
Ghana	2016	Male	All Ages	30% (26-34%)	70% (66-74%)	2.8 (1.5-4.7)	5.7% (4.2-7.2%)	9.9% (6.7-13%)	9700 (6600-13000)	350000 (250000-440000)
Ghana	2016	Male	15 to 19	19% (12-26%)	81% (74-88%)	1.3 (0.56-2.6)	2.7% (1.4-4.5%)	3.2% (1.4-5.9%)	75 (30-140)	7300 (3700-12000)
Ghana	2016	Male	20 to 24	31% (21-42%)	69% (58-79%)	2.9 (1.3-5.5)	6.9% (4.3-9.8%)	8.1% (4.4-12%)	240 (130-380)	20000 (12000-29000)
Ghana	2016	Male	25 to 29	42% (31-53%)	58% (47-69%)	4.7 (2.3-8.5)	9.7% (6.6-13%)	12% (7.8-16%)	400 (260-560)	28000 (19000-39000)
Ghana	2016	Male	30 to 34	49% (37-60%)	51% (40-63%)	6.1 (3.1-11)	10% (7.6-13%)	13% (9.1-17%)	470 (330-640)	30000 (22000-39000)
Ghana	2016	Male	35 to 39	52% (39-62%)	48% (38-61%)	6.5 (3.1-11)	11% (8-14%)	14% (9.7-18%)	570 (390-760)	32000 (23000-42000)
Ghana	2016	Male	40 to 44	53% (41-63%)	47% (37-59%)	6.4 (2.9-12)	12% (8.6-15%)	15% (10-18%)	670 (460-900)	33000 (23000-44000)
Ghana	2016	Male	45 to 49	52% (39-62%)	48% (38-61%)	6 (2.8-11)	14% (10-18%)	18% (13-22%)	900 (620-1200)	39000 (27000-51000)
Ghana	2016	Male	50 to 54	55% (44-65%)	45% (35-56%)	6.4 (3.1-12)	15% (11-19%)	18% (13-23%)	950 (660-1200)	36000 (25000-46000)
Ghana	2016	Male	55 to 59	61% (50-70%)	39% (30-50%)	7 (3.3-13)	16% (11-20%)	20% (14-24%)	1100 (740-1400)	35000 (24000-45000)
Ghana	2016	Male	60 to 64	62% (49-72%)	38% (28-51%)	6.6 (3-12)	15% (9.9-19%)	18% (12-23%)	1000 (650-1300)	29000 (18000-37000)
Ghana	2016	Male	65 to 69	58% (46-68%)	42% (32-54%)	5.7 (2.7-10)	14% (8.3-18%)	16% (9.9-22%)	1100 (640-1500)	25000 (15000-33000)
Ghana	2016	Male	70 to 74	55% (44-65%)	45% (35-56%)	4.7 (2.2-8.8)	13% (6.7-18%)	15% (7.9-21%)	970 (500-1400)	18000 (9200-26000)
Ghana	2016	Male	75 to 79	52% (39-65%)	48% (35-61%)	3.9 (1.6-7.5)	11% (4.5-16%)	12% (5.1-19%)	660 (280-1000)	9400 (3900-14000)
Ghana	2016	Male	80 to 84	48% (35-61%)	52% (39-65%)	3.1 (1.3-5.8)	9% (3.6-14%)	11% (4.2-17%)	370 (150-590)	3900 (1600-6200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Ghana	2016	Male	85 to 89	46% (33-59%)	54% (41-67%)	2.5 (1.1-4.8)	7.3% (2.4-13%)	8.5% (2.8-15%)	190 (64-350)	1500 (500-2600)
Ghana	2016	Male	90 to 94	47% (34-61%)	53% (39-66%)	2.2 (0.94-4.5)	5.4% (1.3-11%)	6.2% (1.3-13%)	57 (12-120)	300 (70-620)
Ghana	2016	Male	95 plus	52% (34-68%)	48% (32-66%)	2 (0.69-4.4)	4.6% (0.61-11%)	5.2% (0.45-13%)	12 (1-30)	43 (5.6-100)
Guinea	1990	Female	All Ages	24% (19-30%)	76% (70-81%)	0.11 (0.094-0.12)	0.18% (0.048-0.32%)	0.19% (-0.12-0.49%)	86 (-52-220)	5800 (1600-10000)
Guinea	1990	Female	15 to 19	22% (12-35%)	78% (65-88%)	0.094 (0.039-0.18)	0.38% (0.16-0.65%)	0.082% (-0.17-0.38%)	0.53 (-1.1-2.4)	290 (120-500)
Guinea	1990	Female	20 to 24	27% (14-44%)	73% (56-86%)	0.15 (0.064-0.26)	0.8% (0.39-1.3%)	0.17% (-0.25-0.65%)	1.3 (-2.1-5)	620 (300-1000)
Guinea	1990	Female	25 to 29	30% (16-47%)	70% (53-84%)	0.18 (0.081-0.31)	0.89% (0.38-1.5%)	0.35% (-0.33-1.1%)	2.7 (-2.6-8.6)	660 (280-1100)
Guinea	1990	Female	30 to 34	33% (17-51%)	67% (49-83%)	0.2 (0.1-0.33)	0.94% (0.38-1.5%)	0.62% (-0.21-1.5%)	5.1 (-1.7-12)	670 (250-1100)
Guinea	1990	Female	35 to 39	36% (20-56%)	64% (44-80%)	0.22 (0.11-0.36)	0.95% (0.36-1.6%)	0.84% (-0.069-1.8%)	7.7 (-0.63-16)	660 (240-1100)
Guinea	1990	Female	40 to 44	38% (22-57%)	62% (43-78%)	0.23 (0.1-0.38)	1.2% (0.45-1.9%)	1.4% (0.31-2.5%)	11 (2.5-20)	670 (250-1100)
Guinea	1990	Female	45 to 49	38% (20-58%)	62% (42-80%)	0.23 (0.1-0.42)	1% (0.2-1.8%)	1.2% (-0.028-2.4%)	12 (-0.27-23)	580 (110-1000)
Guinea	1990	Female	50 to 54	39% (21-60%)	61% (40-79%)	0.24 (0.11-0.44)	1.2% (0.3-2.1%)	1.5% (0.26-2.7%)	17 (3.1-31)	680 (170-1200)
Guinea	1990	Female	55 to 59	39% (23-58%)	61% (42-77%)	0.27 (0.12-0.47)	0.82% (-0.14-1.7%)	1% (-0.21-2.2%)	14 (-3-32)	480 (-78-1000)
Guinea	1990	Female	60 to 64	41% (22-60%)	59% (40-78%)	0.27 (0.11-0.49)	0.6% (-0.37-1.4%)	0.74% (-0.48-1.8%)	13 (-8.1-33)	370 (-220-890)
Guinea	1990	Female	65 to 69	41% (24-59%)	59% (41-76%)	0.24 (0.11-0.43)	0.17% (-0.88-1%)	0.19% (-1.1-1.2%)	4.1 (-24-27)	99 (-510-610)
Guinea	1990	Female	70 to 74	39% (24-57%)	61% (43-76%)	0.2 (0.081-0.4)	0.12% (-0.96-0.95%)	0.13% (-1.2-1.1%)	2.9 (-26-26)	57 (-460-460)
Guinea	1990	Female	75 to 79	35% (20-55%)	65% (45-80%)	0.16 (0.054-0.35)	-0.079% (-1.2-0.64%)	-0.11% (-1.3-0.75%)	-1.9 (-24-13)	-23 (-320-190)
Guinea	1990	Female	80 to 84	33% (17-52%)	67% (48-83%)	0.15 (0.048-0.3)	-0.18% (-1-0.41%)	-0.22% (-1.2-0.47%)	-2 (-11-4.3)	-19 (-110-45)
Guinea	1990	Female	85 to 89	33% (18-51%)	67% (49-82%)	0.14 (0.052-0.31)	-0.28% (-1.1-0.24%)	-0.35% (-1.3-0.27%)	-1.6 (-5.8-1.2)	-11 (-41-9.2)
Guinea	1990	Female	90 to 94	36% (20-54%)	64% (46-80%)	0.15 (0.05-0.32)	-0.38% (-1.2-0.13%)	-0.5% (-1.5-0.13%)	-0.79 (-2.3-0.2)	-3.6 (-11-1.3)
Guinea	1990	Female	95 plus	40% (21-61%)	60% (39-79%)	0.16 (0.045-0.36)	-0.38% (-1.3-0.17%)	-0.53% (-1.6-0.15%)	-0.24 (-0.72-0.07)	-0.69 (-2.3-0.31)
Guinea	1990	Male	All Ages	20% (15-25%)	80% (75-85%)	0.13 (0.11-0.14)	0.37% (0.24-0.54%)	0.59% (0.32-0.93%)	260 (130-410)	12000 (7800-17000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	1990	Male	15 to 19	15% (7.8-25%)	85% (75-92%)	0.072 (0.028-0.14)	0.58% (0.36-0.88%)	0.27% (0.066-0.57%)	1.4 (0.33-2.9)	350 (210-550)
Guinea	1990	Male	20 to 24	22% (11-36%)	78% (64-89%)	0.15 (0.059-0.27)	1.3% (0.78-2%)	0.7% (0.1-1.4%)	4.4 (0.67-8.9)	830 (490-1300)
Guinea	1990	Male	25 to 29	27% (15-44%)	73% (56-85%)	0.22 (0.098-0.37)	1.8% (1-2.7%)	1.3% (0.23-2.5%)	8.3 (1.4-17)	1100 (600-1600)
Guinea	1990	Male	30 to 34	31% (16-49%)	69% (51-84%)	0.26 (0.13-0.43)	2.2% (1.2-3.3%)	2% (0.69-3.6%)	14 (4.7-24)	1300 (700-1900)
Guinea	1990	Male	35 to 39	34% (18-53%)	66% (47-82%)	0.29 (0.14-0.49)	2.7% (1.6-4%)	3% (1.5-4.9%)	23 (11-37)	1500 (940-2300)
Guinea	1990	Male	40 to 44	35% (20-54%)	65% (46-80%)	0.31 (0.14-0.55)	3% (1.8-4.4%)	3.7% (2-5.7%)	27 (14-43)	1400 (850-2100)
Guinea	1990	Male	45 to 49	36% (19-55%)	64% (45-81%)	0.33 (0.14-0.63)	3% (1.8-4.6%)	3.8% (2.2-6.1%)	32 (18-53)	1500 (880-2300)
Guinea	1990	Male	50 to 54	36% (19-57%)	64% (43-81%)	0.33 (0.14-0.63)	2.6% (1.4-4.1%)	3.3% (1.6-5.3%)	32 (15-52)	1300 (670-2000)
Guinea	1990	Male	55 to 59	36% (21-55%)	64% (45-79%)	0.32 (0.13-0.59)	2.4% (1.2-3.7%)	2.9% (1.4-4.7%)	31 (15-51)	1100 (550-1700)
Guinea	1990	Male	60 to 64	36% (20-56%)	64% (44-80%)	0.29 (0.11-0.57)	1.9% (0.92-3.1%)	2.3% (1-4%)	31 (14-54)	890 (440-1500)
Guinea	1990	Male	65 to 69	36% (21-54%)	64% (46-79%)	0.24 (0.1-0.46)	1.3% (0.5-2.2%)	1.6% (0.5-2.8%)	24 (7.3-41)	550 (210-940)
Guinea	1990	Male	70 to 74	34% (21-52%)	66% (48-79%)	0.19 (0.074-0.39)	0.91% (0.16-1.8%)	1.1% (0.14-2.2%)	17 (2.1-34)	310 (52-610)
Guinea	1990	Male	75 to 79	31% (17-50%)	69% (50-83%)	0.14 (0.046-0.32)	0.64% (-0.015-1.3%)	0.74% (-0.071-1.6%)	8.9 (-0.88-19)	130 (-3-270)
Guinea	1990	Male	80 to 84	28% (15-47%)	72% (53-85%)	0.12 (0.038-0.25)	0.5% (-0.095-1.1%)	0.56% (-0.15-1.3%)	3.3 (-0.91-7.6)	36 (-7-79)
Guinea	1990	Male	85 to 89	27% (14-44%)	73% (56-86%)	0.1 (0.032-0.22)	0.37% (-0.092-0.86%)	0.4% (-0.17-0.99%)	1.3 (-0.56-3.3)	11 (-2.6-25)
Guinea	1990	Male	90 to 94	29% (15-46%)	71% (54-85%)	0.096 (0.028-0.23)	0.18% (-0.23-0.5%)	0.14% (-0.38-0.56%)	0.2 (-0.5-0.76)	1.5 (-1.9-4.2)
Guinea	1990	Male	95 plus	33% (15-55%)	67% (45-85%)	0.098 (0.022-0.26)	0.15% (-0.38-0.53%)	0.069% (-0.63-0.54%)	0.026 (-0.24-0.2)	0.24 (-0.6-0.81)
Guinea	1995	Female	All Ages	24% (19-31%)	76% (69-81%)	0.12 (0.11-0.14)	0.2% (0.044-0.36%)	0.19% (-0.16-0.54%)	100 (-84-290)	7300 (1700-13000)
Guinea	1995	Female	15 to 19	22% (12-37%)	78% (63-88%)	0.11 (0.044-0.2)	0.4% (0.17-0.68%)	0.11% (-0.19-0.46%)	0.9 (-1.5-3.9)	390 (170-680)
Guinea	1995	Female	20 to 24	28% (14-44%)	72% (56-86%)	0.17 (0.073-0.31)	0.82% (0.38-1.3%)	0.19% (-0.32-0.72%)	2 (-3.1-7.4)	820 (360-1400)
Guinea	1995	Female	25 to 29	30% (16-48%)	70% (52-84%)	0.2 (0.095-0.35)	0.92% (0.4-1.5%)	0.38% (-0.34-1.2%)	3.8 (-3.4-12)	840 (360-1400)
Guinea	1995	Female	30 to 34	33% (17-52%)	67% (48-83%)	0.23 (0.12-0.38)	0.96% (0.36-1.6%)	0.63% (-0.26-1.6%)	6.5 (-2.7-16)	840 (320-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	1995	Female	35 to 39	37% (20-56%)	63% (44-80%)	0.25 (0.12-0.4)	0.97% (0.33-1.7%)	0.84% (-0.086-1.8%)	9.6 (-0.98-21)	830 (280-1500)
Guinea	1995	Female	40 to 44	38% (22-57%)	62% (43-78%)	0.26 (0.12-0.43)	1.2% (0.46-2%)	1.4% (0.27-2.7%)	16 (3.2-32)	990 (360-1700)
Guinea	1995	Female	45 to 49	39% (20-58%)	61% (42-80%)	0.26 (0.12-0.48)	1.1% (0.19-1.9%)	1.2% (0.03-2.5%)	14 (0.36-28)	720 (130-1300)
Guinea	1995	Female	50 to 54	39% (21-59%)	61% (41-79%)	0.28 (0.12-0.5)	1.2% (0.21-2.1%)	1.5% (0.14-2.7%)	21 (2-39)	820 (140-1400)
Guinea	1995	Female	55 to 59	39% (22-58%)	61% (42-78%)	0.3 (0.13-0.53)	0.81% (-0.22-1.8%)	1% (-0.33-2.3%)	18 (-5.7-41)	580 (-170-1300)
Guinea	1995	Female	60 to 64	41% (23-60%)	59% (40-77%)	0.3 (0.13-0.56)	0.58% (-0.38-1.5%)	0.71% (-0.5-1.9%)	16 (-11-41)	430 (-290-1100)
Guinea	1995	Female	65 to 69	42% (25-59%)	58% (41-75%)	0.28 (0.12-0.5)	0.12% (-0.99-1%)	0.13% (-1.2-1.2%)	3.6 (-32-34)	89 (-710-760)
Guinea	1995	Female	70 to 74	40% (25-58%)	60% (42-75%)	0.23 (0.093-0.46)	0.063% (-0.99-0.93%)	0.06% (-1.2-1.1%)	1.8 (-37-33)	40 (-640-590)
Guinea	1995	Female	75 to 79	36% (19-55%)	64% (45-81%)	0.18 (0.061-0.4)	-0.11% (-1.1-0.62%)	-0.14% (-1.4-0.71%)	-3.4 (-33-17)	-42 (-450-240)
Guinea	1995	Female	80 to 84	33% (17-53%)	67% (47-83%)	0.17 (0.055-0.34)	-0.23% (-1.2-0.38%)	-0.29% (-1.4-0.46%)	-3.7 (-18-5.9)	-36 (-190-62)
Guinea	1995	Female	85 to 89	33% (18-52%)	67% (48-82%)	0.16 (0.055-0.34)	-0.35% (-1.3-0.23%)	-0.44% (-1.5-0.26%)	-2.7 (-9.5-1.6)	-18 (-67-12)
Guinea	1995	Female	90 to 94	36% (19-54%)	64% (46-81%)	0.17 (0.056-0.36)	-0.44% (-1.3-0.11%)	-0.58% (-1.6-0.1%)	-1.2 (-3.2-0.21)	-5.3 (-15-1.4)
Guinea	1995	Female	95 plus	41% (21-61%)	59% (39-79%)	0.18 (0.05-0.42)	-0.43% (-1.4-0.16%)	-0.6% (-1.8-0.11%)	-0.32 (-0.94-0.059)	-0.92 (-2.9-0.34)
Guinea	1995	Male	All Ages	20% (16-26%)	80% (74-84%)	0.15 (0.13-0.17)	0.44% (0.27-0.64%)	0.68% (0.34-1.1%)	360 (180-570)	17000 (10000-24000)
Guinea	1995	Male	15 to 19	15% (7.9-26%)	85% (74-92%)	0.085 (0.033-0.16)	0.61% (0.36-0.92%)	0.32% (0.062-0.69%)	2.3 (0.46-5)	510 (290-810)
Guinea	1995	Male	20 to 24	23% (12-39%)	77% (61-88%)	0.17 (0.069-0.33)	1.4% (0.81-2.1%)	0.8% (0.15-1.6%)	7.1 (1.4-14)	1200 (650-1800)
Guinea	1995	Male	25 to 29	28% (14-45%)	72% (55-86%)	0.25 (0.11-0.44)	1.8% (0.95-2.8%)	1.4% (0.2-2.7%)	13 (1.8-25)	1500 (790-2300)
Guinea	1995	Male	30 to 34	32% (17-51%)	68% (49-83%)	0.31 (0.15-0.52)	2.2% (1.1-3.3%)	2% (0.62-3.7%)	20 (6.2-37)	1700 (870-2700)
Guinea	1995	Male	35 to 39	35% (20-54%)	65% (46-80%)	0.34 (0.16-0.57)	2.7% (1.6-4%)	3% (1.5-4.9%)	33 (16-53)	2100 (1200-3100)
Guinea	1995	Male	40 to 44	36% (21-56%)	64% (44-79%)	0.37 (0.17-0.63)	2.9% (1.8-4.5%)	3.6% (1.9-5.7%)	43 (23-67)	2200 (1300-3400)
Guinea	1995	Male	45 to 49	37% (20-55%)	63% (45-80%)	0.39 (0.17-0.73)	3% (1.8-4.8%)	3.7% (2-6%)	43 (24-71)	2000 (1100-3000)
Guinea	1995	Male	50 to 54	38% (21-59%)	62% (41-79%)	0.39 (0.17-0.75)	2.7% (1.4-4.3%)	3.3% (1.6-5.4%)	43 (21-76)	1700 (920-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	1995	Male	55 to 59	38% (22-58%)	62% (42-78%)	0.38 (0.16-0.72)	2.4% (1.2-3.9%)	2.9% (1.4-4.9%)	43 (20-75)	1500 (740-2500)
Guinea	1995	Male	60 to 64	38% (20-57%)	62% (43-80%)	0.34 (0.13-0.69)	1.9% (0.91-3.2%)	2.3% (0.96-4%)	40 (17-73)	1200 (560-2000)
Guinea	1995	Male	65 to 69	37% (21-55%)	63% (45-79%)	0.29 (0.12-0.53)	1.3% (0.4-2.3%)	1.5% (0.42-2.8%)	30 (8.4-56)	720 (230-1300)
Guinea	1995	Male	70 to 74	35% (21-53%)	65% (47-79%)	0.22 (0.086-0.46)	0.88% (0.09-1.8%)	1% (0.061-2.2%)	22 (1.3-47)	410 (42-850)
Guinea	1995	Male	75 to 79	33% (18-52%)	67% (48-82%)	0.17 (0.054-0.37)	0.6% (-0.15-1.3%)	0.68% (-0.25-1.6%)	12 (-4.3-28)	180 (-45-400)
Guinea	1995	Male	80 to 84	30% (15-48%)	70% (52-85%)	0.14 (0.046-0.29)	0.47% (-0.15-1.1%)	0.52% (-0.26-1.3%)	4.7 (-2.4-12)	53 (-17-130)
Guinea	1995	Male	85 to 89	29% (14-46%)	71% (54-86%)	0.12 (0.038-0.26)	0.35% (-0.19-0.91%)	0.36% (-0.29-1%)	1.7 (-1.4-4.7)	14 (-7.7-37)
Guinea	1995	Male	90 to 94	30% (16-49%)	70% (51-84%)	0.11 (0.034-0.27)	0.16% (-0.35-0.53%)	0.12% (-0.51-0.57%)	0.2 (-0.89-0.98)	1.6 (-3.6-5.5)
Guinea	1995	Male	95 plus	35% (17-57%)	65% (43-83%)	0.11 (0.027-0.29)	0.15% (-0.44-0.63%)	0.065% (-0.71-0.66%)	0.03 (-0.31-0.3)	0.28 (-0.78-1.2)
Guinea	2000	Female	All Ages	24% (19-30%)	76% (70-81%)	0.11 (0.092-0.12)	0.2% (0.025-0.37%)	0.16% (-0.23-0.54%)	88 (-130-300)	7300 (920-14000)
Guinea	2000	Female	15 to 19	23% (12-38%)	77% (62-88%)	0.095 (0.04-0.18)	0.4% (0.16-0.65%)	0.078% (-0.23-0.43%)	0.73 (-2.2-3.9)	430 (170-750)
Guinea	2000	Female	20 to 24	28% (14-45%)	72% (55-86%)	0.15 (0.065-0.26)	0.8% (0.35-1.3%)	0.14% (-0.37-0.69%)	1.6 (-4.5-7.8)	910 (380-1500)
Guinea	2000	Female	25 to 29	31% (15-48%)	69% (52-85%)	0.18 (0.081-0.31)	0.85% (0.35-1.4%)	0.3% (-0.36-1%)	3.5 (-4.1-12)	900 (360-1500)
Guinea	2000	Female	30 to 34	34% (17-53%)	66% (47-83%)	0.2 (0.099-0.34)	0.86% (0.24-1.5%)	0.5% (-0.36-1.3%)	5.9 (-4.2-15)	850 (240-1500)
Guinea	2000	Female	35 to 39	37% (20-56%)	63% (44-80%)	0.22 (0.11-0.36)	0.87% (0.25-1.5%)	0.7% (-0.16-1.6%)	8.9 (-2-21)	830 (250-1500)
Guinea	2000	Female	40 to 44	39% (23-57%)	61% (43-77%)	0.23 (0.11-0.39)	1.1% (0.33-1.8%)	1.2% (0.066-2.3%)	16 (0.84-31)	970 (290-1700)
Guinea	2000	Female	45 to 49	39% (21-58%)	61% (42-79%)	0.24 (0.11-0.44)	0.94% (0.016-1.8%)	1.1% (-0.22-2.3%)	16 (-3.4-34)	810 (14-1500)
Guinea	2000	Female	50 to 54	39% (22-60%)	61% (40-78%)	0.25 (0.11-0.44)	1% (0.036-2%)	1.3% (-0.063-2.6%)	19 (-0.92-38)	780 (27-1400)
Guinea	2000	Female	55 to 59	40% (23-58%)	60% (42-77%)	0.27 (0.12-0.48)	0.67% (-0.43-1.7%)	0.81% (-0.62-2.2%)	15 (-12-40)	520 (-330-1300)
Guinea	2000	Female	60 to 64	41% (24-60%)	59% (40-76%)	0.27 (0.11-0.51)	0.45% (-0.52-1.3%)	0.54% (-0.65-1.7%)	13 (-16-40)	360 (-410-1100)
Guinea	2000	Female	65 to 69	42% (25-60%)	58% (40-75%)	0.25 (0.11-0.44)	0.046% (-1-0.94%)	0.038% (-1.3-1.1%)	1.1 (-36-33)	37 (-810-720)
Guinea	2000	Female	70 to 74	40% (24-58%)	60% (42-76%)	0.2 (0.082-0.4)	0.013% (-1.1-0.9%)	-0.0046% (-1.3-1.1%)	-0.089 (-41-34)	10 (-720-600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	2000	Female	75 to 79	36% (20-54%)	64% (46-80%)	0.16 (0.055-0.35)	-0.089% (-1.2-0.6%)	-0.12% (-1.4-0.69%)	-3.3 (-36-19)	-38 (-500-270)
Guinea	2000	Female	80 to 84	33% (18-52%)	67% (48-82%)	0.15 (0.051-0.3)	-0.22% (-1.2-0.37%)	-0.28% (-1.4-0.43%)	-4.3 (-21-6.7)	-41 (-220-71)
Guinea	2000	Female	85 to 89	34% (18-51%)	66% (49-82%)	0.14 (0.05-0.29)	-0.34% (-1.2-0.26%)	-0.43% (-1.5-0.28%)	-3.5 (-12-2.4)	-24 (-85-19)
Guinea	2000	Female	90 to 94	37% (20-54%)	63% (46-80%)	0.15 (0.049-0.32)	-0.42% (-1.3-0.15%)	-0.56% (-1.6-0.15%)	-1.4 (-4.1-0.37)	-6.4 (-19-2.4)
Guinea	2000	Female	95 plus	41% (22-62%)	59% (38-78%)	0.16 (0.045-0.37)	-0.4% (-1.3-0.17%)	-0.57% (-1.7-0.12%)	-0.35 (-1-0.074)	-0.96 (-3.2-0.4)
Guinea	2000	Male	All Ages	21% (16-27%)	79% (73-84%)	0.13 (0.12-0.15)	0.45% (0.26-0.69%)	0.68% (0.28-1.2%)	380 (160-660)	18000 (10000-27000)
Guinea	2000	Male	15 to 19	16% (8.1-27%)	84% (73-92%)	0.08 (0.031-0.15)	0.56% (0.34-0.87%)	0.26% (0.02-0.62%)	2.3 (0.18-5.4)	560 (330-880)
Guinea	2000	Male	20 to 24	24% (12-39%)	76% (61-88%)	0.16 (0.066-0.29)	1.2% (0.68-1.9%)	0.64% (0.043-1.3%)	7.3 (0.48-16)	1300 (700-2000)
Guinea	2000	Male	25 to 29	29% (16-45%)	71% (55-84%)	0.23 (0.1-0.4)	1.6% (0.82-2.5%)	1.1% (0.079-2.1%)	13 (0.94-26)	1600 (800-2400)
Guinea	2000	Male	30 to 34	33% (17-51%)	67% (49-83%)	0.29 (0.14-0.47)	1.9% (0.91-2.9%)	1.6% (0.45-3%)	20 (5.7-38)	1800 (860-2800)
Guinea	2000	Male	35 to 39	36% (20-55%)	64% (45-80%)	0.31 (0.15-0.53)	2.3% (1.3-3.6%)	2.5% (1.1-4.3%)	34 (16-58)	2200 (1200-3300)
Guinea	2000	Male	40 to 44	37% (22-56%)	63% (44-78%)	0.34 (0.16-0.58)	2.6% (1.4-4.1%)	3% (1.4-4.9%)	45 (21-74)	2400 (1300-3700)
Guinea	2000	Male	45 to 49	38% (20-56%)	62% (44-80%)	0.35 (0.16-0.66)	2.7% (1.5-4.4%)	3.2% (1.6-5.4%)	53 (26-90)	2400 (1300-3900)
Guinea	2000	Male	50 to 54	38% (21-59%)	62% (41-79%)	0.36 (0.15-0.66)	2.4% (1.1-3.9%)	2.8% (1.1-4.8%)	45 (18-79)	1800 (810-3000)
Guinea	2000	Male	55 to 59	38% (23-58%)	62% (42-77%)	0.35 (0.15-0.64)	2.1% (0.93-3.6%)	2.5% (0.95-4.5%)	46 (17-83)	1500 (670-2700)
Guinea	2000	Male	60 to 64	38% (21-58%)	62% (42-79%)	0.31 (0.12-0.62)	1.7% (0.68-3.1%)	2% (0.7-3.7%)	42 (15-80)	1200 (490-2200)
Guinea	2000	Male	65 to 69	38% (22-56%)	62% (44-78%)	0.26 (0.11-0.49)	1.2% (0.26-2.2%)	1.3% (0.22-2.7%)	31 (5.3-63)	730 (170-1400)
Guinea	2000	Male	70 to 74	37% (22-54%)	63% (46-78%)	0.2 (0.078-0.41)	0.79% (-0.065-1.8%)	0.92% (-0.11-2.1%)	23 (-2.7-52)	430 (-33-940)
Guinea	2000	Male	75 to 79	33% (19-52%)	67% (48-81%)	0.16 (0.051-0.34)	0.54% (-0.21-1.3%)	0.6% (-0.28-1.5%)	13 (-6.1-32)	190 (-73-450)
Guinea	2000	Male	80 to 84	31% (16-49%)	69% (51-84%)	0.13 (0.043-0.26)	0.43% (-0.22-1.1%)	0.47% (-0.31-1.2%)	5.5 (-3.7-15)	62 (-31-150)
Guinea	2000	Male	85 to 89	30% (15-47%)	70% (53-85%)	0.11 (0.036-0.25)	0.32% (-0.25-0.88%)	0.32% (-0.34-0.99%)	2 (-2.1-6.1)	17 (-13-46)
Guinea	2000	Male	90 to 94	32% (16-50%)	68% (50-84%)	0.11 (0.033-0.25)	0.16% (-0.36-0.56%)	0.11% (-0.53-0.59%)	0.24 (-1.1-1.2)	1.9 (-4.5-6.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	2000	Male	95 plus	36% (17-58%)	64% (42-83%)	0.11 (0.026-0.28)	0.16% (-0.5-0.67%)	0.078% (-0.73-0.68%)	0.038 (-0.36-0.33)	0.32 (-0.98-1.3)
Guinea	2005	Female	All Ages	24% (19-31%)	76% (69-81%)	0.094 (0.083-0.11)	0.2% (0.04-0.37%)	0.15% (-0.23-0.52%)	82 (-120-280)	7600 (1500-14000)
Guinea	2005	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.083 (0.035-0.16)	0.4% (0.17-0.66%)	0.066% (-0.21-0.37%)	0.64 (-2-3.7)	470 (200-790)
Guinea	2005	Female	20 to 24	28% (14-45%)	72% (55-86%)	0.13 (0.057-0.23)	0.81% (0.38-1.4%)	0.12% (-0.39-0.68%)	1.4 (-4.7-7.9)	1000 (460-1700)
Guinea	2005	Female	25 to 29	31% (16-49%)	69% (51-84%)	0.16 (0.074-0.28)	0.81% (0.36-1.3%)	0.25% (-0.33-0.86%)	3.4 (-4.5-12)	990 (420-1600)
Guinea	2005	Female	30 to 34	34% (17-53%)	66% (47-83%)	0.18 (0.09-0.3)	0.79% (0.28-1.3%)	0.42% (-0.27-1.1%)	5.9 (-3.6-15)	900 (330-1500)
Guinea	2005	Female	35 to 39	37% (20-57%)	63% (43-80%)	0.2 (0.096-0.33)	0.81% (0.28-1.3%)	0.61% (-0.058-1.4%)	8.8 (-0.87-20)	840 (290-1400)
Guinea	2005	Female	40 to 44	39% (22-58%)	61% (42-78%)	0.21 (0.098-0.35)	1% (0.39-1.6%)	1.1% (0.21-1.9%)	16 (3.2-29)	970 (370-1600)
Guinea	2005	Female	45 to 49	40% (21-59%)	60% (41-79%)	0.21 (0.093-0.39)	0.88% (0.12-1.6%)	0.98% (-0.084-2%)	16 (-1.4-32)	820 (110-1500)
Guinea	2005	Female	50 to 54	40% (22-60%)	60% (40-78%)	0.22 (0.098-0.41)	0.98% (0.1-1.8%)	1.2% (0.018-2.4%)	22 (0.35-45)	900 (95-1700)
Guinea	2005	Female	55 to 59	40% (23-58%)	60% (42-77%)	0.24 (0.1-0.43)	0.59% (-0.39-1.5%)	0.7% (-0.57-1.8%)	13 (-11-37)	470 (-300-1200)
Guinea	2005	Female	60 to 64	41% (23-60%)	59% (40-77%)	0.24 (0.099-0.45)	0.37% (-0.56-1.2%)	0.43% (-0.67-1.5%)	10 (-17-36)	290 (-450-980)
Guinea	2005	Female	65 to 69	42% (25-60%)	58% (40-75%)	0.22 (0.095-0.39)	0.0095% (-1-0.87%)	-0.013% (-1.3-1%)	-0.35 (-37-30)	8.7 (-810-670)
Guinea	2005	Female	70 to 74	40% (24-58%)	60% (42-76%)	0.18 (0.073-0.35)	-0.019% (-1-0.8%)	-0.049% (-1.3-0.94%)	-1.4 (-39-30)	-11 (-700-550)
Guinea	2005	Female	75 to 79	36% (20-55%)	64% (45-80%)	0.14 (0.047-0.31)	-0.088% (-1.2-0.59%)	-0.13% (-1.4-0.68%)	-3.4 (-36-19)	-38 (-500-270)
Guinea	2005	Female	80 to 84	33% (18-52%)	67% (48-82%)	0.13 (0.045-0.26)	-0.21% (-1.1-0.36%)	-0.27% (-1.3-0.41%)	-4.2 (-21-6)	-39 (-210-65)
Guinea	2005	Female	85 to 89	34% (19-52%)	66% (48-81%)	0.13 (0.045-0.26)	-0.31% (-1.2-0.24%)	-0.4% (-1.4-0.26%)	-3.9 (-14-2.4)	-26 (-97-20)
Guinea	2005	Female	90 to 94	37% (20-55%)	63% (45-80%)	0.14 (0.044-0.28)	-0.41% (-1.2-0.15%)	-0.54% (-1.6-0.13%)	-1.8 (-5.3-0.46)	-8.3 (-25-2.9)
Guinea	2005	Female	95 plus	41% (21-62%)	59% (38-79%)	0.14 (0.039-0.32)	-0.38% (-1.3-0.18%)	-0.54% (-1.6-0.14%)	-0.42 (-1.3-0.11)	-1.2 (-3.9-0.58)
Guinea	2005	Male	All Ages	21% (16-26%)	79% (74-84%)	0.12 (0.1-0.13)	0.49% (0.28-0.75%)	0.71% (0.31-1.2%)	410 (180-710)	20000 (12000-30000)
Guinea	2005	Male	15 to 19	16% (8.3-27%)	84% (73-92%)	0.071 (0.028-0.13)	0.58% (0.36-0.88%)	0.26% (0.044-0.58%)	2.4 (0.41-5.5)	620 (360-980)
Guinea	2005	Male	20 to 24	24% (12-40%)	76% (60-88%)	0.14 (0.059-0.26)	1.2% (0.73-1.9%)	0.64% (0.094-1.3%)	7.9 (1.1-17)	1400 (820-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	2005	Male	25 to 29	29% (15-47%)	71% (53-85%)	0.21 (0.095-0.35)	1.5% (0.83-2.3%)	1% (0.15-2%)	14 (2-27)	1700 (920-2700)
Guinea	2005	Male	30 to 34	33% (17-51%)	67% (49-83%)	0.26 (0.12-0.42)	1.8% (0.91-2.7%)	1.5% (0.48-2.8%)	22 (7-41)	1900 (980-3000)
Guinea	2005	Male	35 to 39	36% (19-55%)	64% (45-81%)	0.28 (0.13-0.47)	2.2% (1.3-3.4%)	2.3% (1.1-3.8%)	36 (17-62)	2300 (1300-3600)
Guinea	2005	Male	40 to 44	37% (22-56%)	63% (44-78%)	0.3 (0.14-0.51)	2.4% (1.4-3.7%)	2.8% (1.3-4.5%)	47 (22-76)	2500 (1300-3800)
Guinea	2005	Male	45 to 49	38% (20-58%)	62% (42-80%)	0.31 (0.14-0.6)	2.5% (1.5-4.1%)	3% (1.5-5%)	55 (28-93)	2500 (1400-4000)
Guinea	2005	Male	50 to 54	39% (21-59%)	61% (41-79%)	0.32 (0.14-0.58)	2.3% (1.2-3.7%)	2.7% (1.2-4.6%)	55 (24-96)	2200 (1100-3600)
Guinea	2005	Male	55 to 59	38% (22-58%)	62% (42-78%)	0.31 (0.13-0.58)	2.1% (0.95-3.5%)	2.4% (0.99-4.3%)	47 (19-85)	1600 (730-2800)
Guinea	2005	Male	60 to 64	38% (21-58%)	62% (42-79%)	0.28 (0.11-0.56)	1.7% (0.7-2.9%)	2% (0.72-3.5%)	45 (17-81)	1300 (540-2200)
Guinea	2005	Male	65 to 69	38% (21-56%)	62% (44-79%)	0.23 (0.1-0.44)	1.1% (0.26-2.2%)	1.3% (0.22-2.6%)	32 (5.3-65)	760 (170-1500)
Guinea	2005	Male	70 to 74	36% (22-54%)	64% (46-78%)	0.18 (0.07-0.37)	0.78% (0.037-1.7%)	0.9% (-0.01-1.9%)	23 (-0.26-50)	430 (21-910)
Guinea	2005	Male	75 to 79	33% (18-53%)	67% (47-82%)	0.14 (0.044-0.3)	0.54% (-0.12-1.3%)	0.6% (-0.17-1.5%)	13 (-3.9-34)	200 (-45-470)
Guinea	2005	Male	80 to 84	31% (16-49%)	69% (51-84%)	0.11 (0.037-0.23)	0.45% (-0.14-1%)	0.48% (-0.22-1.2%)	6 (-2.7-15)	66 (-21-150)
Guinea	2005	Male	85 to 89	29% (16-47%)	71% (53-84%)	0.099 (0.033-0.21)	0.33% (-0.2-0.85%)	0.34% (-0.28-0.99%)	2.5 (-2.2-7.5)	21 (-12-57)
Guinea	2005	Male	90 to 94	31% (17-49%)	69% (51-83%)	0.093 (0.029-0.22)	0.17% (-0.29-0.57%)	0.14% (-0.45-0.62%)	0.36 (-1.2-1.6)	2.7 (-4.6-9)
Guinea	2005	Male	95 plus	36% (18-57%)	64% (43-82%)	0.094 (0.023-0.24)	0.18% (-0.41-0.64%)	0.097% (-0.65-0.66%)	0.056 (-0.37-0.38)	0.41 (-0.95-1.5)
Guinea	2010	Female	All Ages	24% (19-31%)	76% (69-81%)	0.093 (0.078-0.11)	0.23% (0.045-0.42%)	0.15% (-0.24-0.55%)	85 (-140-310)	8500 (1700-16000)
Guinea	2010	Female	15 to 19	23% (12-36%)	77% (64-88%)	0.082 (0.034-0.16)	0.43% (0.19-0.69%)	0.066% (-0.2-0.36%)	0.69 (-2.1-3.7)	550 (250-910)
Guinea	2010	Female	20 to 24	28% (14-45%)	72% (55-86%)	0.13 (0.058-0.23)	0.87% (0.41-1.4%)	0.12% (-0.38-0.64%)	1.6 (-5-8.6)	1200 (530-2000)
Guinea	2010	Female	25 to 29	31% (16-48%)	69% (52-84%)	0.16 (0.072-0.27)	0.86% (0.4-1.4%)	0.26% (-0.35-0.87%)	3.8 (-5.1-12)	1100 (520-1900)
Guinea	2010	Female	30 to 34	34% (17-52%)	66% (48-83%)	0.18 (0.09-0.3)	0.82% (0.3-1.3%)	0.43% (-0.23-1.1%)	6.7 (-3.4-17)	1000 (380-1700)
Guinea	2010	Female	35 to 39	37% (20-56%)	63% (44-80%)	0.19 (0.092-0.32)	0.82% (0.34-1.4%)	0.63% (-0.044-1.4%)	9.9 (-0.69-22)	950 (390-1600)
Guinea	2010	Female	40 to 44	39% (23-58%)	61% (42-77%)	0.2 (0.096-0.35)	1% (0.44-1.6%)	1.1% (0.27-2%)	17 (4-31)	1100 (450-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	2010	Female	45 to 49	40% (21-58%)	60% (42-79%)	0.21 (0.093-0.38)	0.9% (0.2-1.6%)	1% (0.0059-1.9%)	17 (0.11-35)	890 (200-1600)
Guinea	2010	Female	50 to 54	40% (22-60%)	60% (40-78%)	0.22 (0.097-0.38)	0.96% (0.16-1.8%)	1.2% (0.028-2.3%)	24 (0.59-49)	960 (150-1900)
Guinea	2010	Female	55 to 59	40% (23-59%)	60% (41-77%)	0.24 (0.1-0.42)	0.58% (-0.35-1.4%)	0.69% (-0.52-1.8%)	17 (-12-44)	580 (-350-1400)
Guinea	2010	Female	60 to 64	41% (23-60%)	59% (40-77%)	0.24 (0.097-0.44)	0.32% (-0.65-1.2%)	0.37% (-0.84-1.4%)	9.5 (-22-39)	270 (-590-1000)
Guinea	2010	Female	65 to 69	42% (25-60%)	58% (40-75%)	0.21 (0.093-0.39)	-0.039% (-1.1-0.83%)	-0.071% (-1.4-0.99%)	-2.2 (-43-32)	-32 (-940-720)
Guinea	2010	Female	70 to 74	40% (24-57%)	60% (43-76%)	0.17 (0.068-0.34)	-0.05% (-1.1-0.79%)	-0.085% (-1.3-0.93%)	-2.8 (-48-33)	-35 (-840-590)
Guinea	2010	Female	75 to 79	35% (20-54%)	65% (46-80%)	0.14 (0.044-0.3)	-0.1% (-1.3-0.57%)	-0.15% (-1.5-0.67%)	-4.3 (-44-20)	-50 (-600-280)
Guinea	2010	Female	80 to 84	33% (17-52%)	67% (48-83%)	0.13 (0.041-0.25)	-0.23% (-1.2-0.37%)	-0.29% (-1.4-0.42%)	-4.8 (-23-7.1)	-45 (-230-76)
Guinea	2010	Female	85 to 89	34% (18-52%)	66% (48-82%)	0.12 (0.043-0.26)	-0.32% (-1.2-0.24%)	-0.41% (-1.4-0.25%)	-4.4 (-16-2.7)	-29 (-110-22)
Guinea	2010	Female	90 to 94	37% (20-54%)	63% (46-80%)	0.13 (0.042-0.27)	-0.41% (-1.3-0.14%)	-0.55% (-1.6-0.13%)	-2.4 (-7.1-0.53)	-11 (-34-3.8)
Guinea	2010	Female	95 plus	41% (22-62%)	59% (38-78%)	0.14 (0.038-0.32)	-0.4% (-1.3-0.18%)	-0.57% (-1.7-0.13%)	-0.64 (-1.9-0.15)	-1.8 (-5.9-0.82)
Guinea	2010	Male	All Ages	21% (16-27%)	79% (73-84%)	0.11 (0.098-0.13)	0.58% (0.34-0.88%)	0.83% (0.37-1.4%)	490 (210-850)	23000 (13000-35000)
Guinea	2010	Male	15 to 19	16% (8.4-27%)	84% (73-92%)	0.067 (0.028-0.13)	0.59% (0.37-0.89%)	0.27% (0.055-0.58%)	2.9 (0.56-6.4)	730 (440-1100)
Guinea	2010	Male	20 to 24	24% (12-39%)	76% (61-88%)	0.14 (0.058-0.25)	1.3% (0.77-1.9%)	0.68% (0.13-1.4%)	9.8 (1.8-20)	1700 (990-2700)
Guinea	2010	Male	25 to 29	29% (15-46%)	71% (54-85%)	0.2 (0.09-0.34)	1.6% (0.89-2.4%)	1.1% (0.2-2.1%)	17 (2.9-34)	2100 (1100-3300)
Guinea	2010	Male	30 to 34	33% (17-51%)	67% (49-83%)	0.25 (0.12-0.41)	1.9% (1-2.9%)	1.6% (0.56-3%)	28 (9.5-51)	2400 (1300-3800)
Guinea	2010	Male	35 to 39	36% (20-56%)	64% (44-80%)	0.27 (0.13-0.46)	2.3% (1.4-3.6%)	2.5% (1.2-4.1%)	44 (22-74)	2800 (1600-4300)
Guinea	2010	Male	40 to 44	37% (22-57%)	63% (43-78%)	0.29 (0.13-0.51)	2.6% (1.6-4%)	3% (1.6-4.8%)	55 (29-90)	2900 (1700-4500)
Guinea	2010	Male	45 to 49	38% (20-57%)	62% (43-80%)	0.31 (0.13-0.58)	2.7% (1.5-4.3%)	3.1% (1.6-5.2%)	63 (31-110)	2800 (1500-4700)
Guinea	2010	Male	50 to 54	39% (21-60%)	61% (40-79%)	0.31 (0.13-0.59)	2.4% (1.1-3.9%)	2.8% (1.1-4.7%)	63 (25-110)	2500 (1100-4100)
Guinea	2010	Male	55 to 59	39% (22-59%)	61% (41-78%)	0.3 (0.13-0.55)	2.2% (0.99-3.7%)	2.6% (1-4.5%)	64 (24-120)	2200 (930-3700)
Guinea	2010	Male	60 to 64	39% (22-58%)	61% (42-78%)	0.27 (0.11-0.55)	1.8% (0.71-3%)	2% (0.74-3.7%)	52 (19-98)	1500 (590-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	2010	Male	65 to 69	39% (22-57%)	61% (43-78%)	0.23 (0.097-0.43)	1.2% (0.27-2.2%)	1.3% (0.21-2.7%)	38 (5.7-76)	880 (190-1700)
Guinea	2010	Male	70 to 74	37% (22-55%)	63% (45-78%)	0.18 (0.065-0.36)	0.8% (0.024-1.7%)	0.93% (-0.0016-2%)	27 (-0.043-60)	490 (14-1100)
Guinea	2010	Male	75 to 79	34% (18-53%)	66% (47-82%)	0.13 (0.042-0.29)	0.55% (-0.11-1.2%)	0.62% (-0.2-1.5%)	15 (-4.5-36)	220 (-43-500)
Guinea	2010	Male	80 to 84	31% (16-50%)	69% (50-84%)	0.11 (0.034-0.23)	0.47% (-0.13-1%)	0.51% (-0.22-1.2%)	6.7 (-2.8-16)	75 (-22-170)
Guinea	2010	Male	85 to 89	30% (16-47%)	70% (53-84%)	0.096 (0.031-0.2)	0.35% (-0.17-0.87%)	0.36% (-0.27-0.98%)	3 (-2.2-8.2)	25 (-12-62)
Guinea	2010	Male	90 to 94	31% (17-49%)	69% (51-83%)	0.091 (0.027-0.21)	0.19% (-0.29-0.6%)	0.15% (-0.45-0.64%)	0.51 (-1.5-2.2)	3.8 (-6-12)
Guinea	2010	Male	95 plus	36% (18-57%)	64% (43-82%)	0.092 (0.022-0.23)	0.19% (-0.43-0.67%)	0.11% (-0.66-0.7%)	0.087 (-0.52-0.56)	0.59 (-1.3-2.1)
Guinea	2016	Female	All Ages	25% (19-31%)	75% (69-81%)	0.1 (0.065-0.15)	0.28% (0.075-0.48%)	0.16% (-0.3-0.57%)	90 (-170-320)	10000 (2800-18000)
Guinea	2016	Female	15 to 19	23% (12-38%)	77% (62-88%)	0.088 (0.033-0.19)	0.46% (0.22-0.73%)	0.078% (-0.19-0.37%)	0.84 (-2.1-4.1)	640 (300-1000)
Guinea	2016	Female	20 to 24	28% (15-46%)	72% (54-85%)	0.14 (0.055-0.27)	0.99% (0.52-1.6%)	0.15% (-0.32-0.7%)	1.9 (-3.9-9.2)	1400 (710-2300)
Guinea	2016	Female	25 to 29	31% (16-49%)	69% (51-84%)	0.17 (0.065-0.32)	1% (0.54-1.5%)	0.31% (-0.29-0.94%)	4.5 (-4.4-13)	1400 (720-2300)
Guinea	2016	Female	30 to 34	34% (17-53%)	66% (47-83%)	0.19 (0.083-0.34)	0.91% (0.45-1.4%)	0.48% (-0.16-1.1%)	7.9 (-2.7-18)	1300 (620-2000)
Guinea	2016	Female	35 to 39	38% (20-56%)	62% (44-80%)	0.21 (0.088-0.38)	0.89% (0.41-1.4%)	0.68% (-0.011-1.5%)	12 (-0.18-25)	1200 (530-1900)
Guinea	2016	Female	40 to 44	39% (23-59%)	61% (41-77%)	0.22 (0.088-0.43)	1.1% (0.51-1.7%)	1.1% (0.32-2%)	20 (5.7-37)	1300 (580-2000)
Guinea	2016	Female	45 to 49	40% (22-59%)	60% (41-78%)	0.22 (0.086-0.43)	0.96% (0.26-1.7%)	1.1% (0.088-2.1%)	20 (1.4-40)	1000 (270-1800)
Guinea	2016	Female	50 to 54	40% (22-61%)	60% (39-78%)	0.23 (0.086-0.45)	1% (0.16-1.7%)	1.2% (0.079-2.2%)	27 (1.8-53)	1100 (180-2000)
Guinea	2016	Female	55 to 59	40% (24-59%)	60% (41-76%)	0.25 (0.097-0.49)	0.61% (-0.31-1.4%)	0.73% (-0.45-1.8%)	20 (-13-50)	670 (-350-1600)
Guinea	2016	Female	60 to 64	41% (24-61%)	59% (39-76%)	0.25 (0.093-0.49)	0.3% (-0.66-1.1%)	0.35% (-0.85-1.4%)	12 (-28-48)	340 (-750-1300)
Guinea	2016	Female	65 to 69	42% (25-60%)	58% (40-75%)	0.23 (0.087-0.45)	-0.093% (-1.2-0.75%)	-0.14% (-1.5-0.89%)	-4.9 (-51-32)	-88 (-1100-730)
Guinea	2016	Female	70 to 74	40% (24-58%)	60% (42-76%)	0.19 (0.065-0.39)	-0.11% (-1.2-0.71%)	-0.16% (-1.4-0.84%)	-5.8 (-53-30)	-85 (-940-550)
Guinea	2016	Female	75 to 79	36% (20-54%)	64% (46-80%)	0.15 (0.044-0.35)	-0.17% (-1.4-0.54%)	-0.22% (-1.6-0.61%)	-7.4 (-57-21)	-92 (-760-300)
Guinea	2016	Female	80 to 84	34% (18-53%)	66% (47-82%)	0.14 (0.041-0.3)	-0.3% (-1.3-0.33%)	-0.38% (-1.5-0.35%)	-7.4 (-30-7)	-70 (-310-77)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea	2016	Female	85 to 89	34% (19-52%)	66% (48-81%)	0.14 (0.042-0.3)	-0.38% (-1.4-0.22%)	-0.49% (-1.7-0.23%)	-5.8 (-20-2.6)	-39 (-140-22)
Guinea	2016	Female	90 to 94	37% (20-55%)	63% (45-80%)	0.14 (0.042-0.32)	-0.48% (-1.4-0.12%)	-0.63% (-1.8-0.091%)	-3.1 (-8.7-0.44)	-14 (-42-3.6)
Guinea	2016	Female	95 plus	41% (22-62%)	59% (38-78%)	0.15 (0.039-0.39)	-0.45% (-1.5-0.14%)	-0.64% (-1.9-0.081%)	-0.94 (-2.8-0.12)	-2.6 (-8.7-0.85)
Guinea	2016	Male	All Ages	21% (16-27%)	79% (73-84%)	0.12 (0.082-0.18)	0.7% (0.44-0.99%)	0.94% (0.47-1.5%)	560 (280-890)	27000 (18000-39000)
Guinea	2016	Male	15 to 19	16% (8.2-28%)	84% (72-92%)	0.074 (0.026-0.16)	0.62% (0.4-0.92%)	0.28% (0.072-0.62%)	3.4 (0.86-7.4)	860 (520-1300)
Guinea	2016	Male	20 to 24	24% (13-40%)	76% (60-87%)	0.15 (0.058-0.29)	1.4% (0.86-2%)	0.74% (0.17-1.5%)	12 (2.6-25)	2100 (1200-3200)
Guinea	2016	Male	25 to 29	30% (16-47%)	70% (53-84%)	0.22 (0.087-0.4)	1.7% (1-2.6%)	1.2% (0.33-2.3%)	21 (5.3-41)	2600 (1500-3900)
Guinea	2016	Male	30 to 34	33% (18-52%)	67% (48-82%)	0.26 (0.12-0.47)	2% (1.2-2.9%)	1.8% (0.71-2.9%)	34 (13-57)	2900 (1700-4400)
Guinea	2016	Male	35 to 39	36% (20-54%)	64% (46-80%)	0.29 (0.12-0.55)	2.4% (1.5-3.5%)	2.6% (1.4-3.9%)	52 (28-85)	3400 (2100-5000)
Guinea	2016	Male	40 to 44	38% (22-57%)	62% (43-78%)	0.31 (0.12-0.6)	2.6% (1.7-3.8%)	3% (1.8-4.5%)	64 (35-100)	3400 (2100-5100)
Guinea	2016	Male	45 to 49	38% (20-57%)	62% (43-80%)	0.33 (0.12-0.67)	2.7% (1.6-4%)	3.2% (1.8-5%)	71 (37-110)	3200 (1900-4900)
Guinea	2016	Male	50 to 54	39% (21-59%)	61% (41-79%)	0.33 (0.12-0.66)	2.4% (1.2-3.8%)	2.9% (1.3-4.5%)	70 (32-110)	2800 (1300-4300)
Guinea	2016	Male	55 to 59	39% (23-58%)	61% (42-77%)	0.32 (0.12-0.64)	2.2% (1.1-3.6%)	2.6% (1.2-4.4%)	72 (33-120)	2400 (1200-4000)
Guinea	2016	Male	60 to 64	39% (22-58%)	61% (42-78%)	0.29 (0.1-0.6)	1.8% (0.84-2.8%)	2.1% (0.9-3.4%)	67 (29-110)	1900 (910-3100)
Guinea	2016	Male	65 to 69	39% (23-57%)	61% (43-77%)	0.25 (0.092-0.49)	1.1% (0.32-2%)	1.3% (0.3-2.4%)	41 (8.9-77)	970 (270-1800)
Guinea	2016	Male	70 to 74	37% (22-55%)	63% (45-78%)	0.19 (0.063-0.42)	0.79% (0.063-1.6%)	0.91% (0.019-1.9%)	29 (0.62-58)	540 (43-1100)
Guinea	2016	Male	75 to 79	34% (19-52%)	66% (48-81%)	0.15 (0.043-0.34)	0.52% (-0.13-1.1%)	0.58% (-0.21-1.2%)	16 (-5.7-34)	230 (-57-480)
Guinea	2016	Male	80 to 84	31% (16-50%)	69% (50-84%)	0.12 (0.034-0.26)	0.43% (-0.15-0.94%)	0.47% (-0.25-1.1%)	6.9 (-3.6-16)	78 (-28-170)
Guinea	2016	Male	85 to 89	29% (15-47%)	71% (53-85%)	0.1 (0.031-0.24)	0.31% (-0.17-0.73%)	0.32% (-0.27-0.82%)	2.8 (-2.4-7.3)	24 (-13-57)
Guinea	2016	Male	90 to 94	31% (16-49%)	69% (51-84%)	0.097 (0.026-0.23)	0.16% (-0.31-0.5%)	0.12% (-0.46-0.53%)	0.44 (-1.7-2)	3.7 (-7-11)
Guinea	2016	Male	95 plus	36% (17-58%)	64% (42-83%)	0.098 (0.022-0.27)	0.17% (-0.43-0.58%)	0.08% (-0.64-0.58%)	0.082 (-0.67-0.62)	0.69 (-1.8-2.4)
Guinea-Bissau	1990	Female	All Ages	25% (20-32%)	75% (68-80%)	0.77 (0.57-1)	1.5% (0.86-2.2%)	2.8% (1.3-4.7%)	230 (110-390)	7700 (4500-12000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	1990	Female	15 to 19	24% (13-39%)	76% (61-87%)	0.7 (0.28-1.3)	1.6% (0.63-2.9%)	1.9% (0.5-3.9%)	2.2 (0.59-4.6)	210 (80-380)
Guinea-Bissau	1990	Female	20 to 24	29% (15-46%)	71% (54-85%)	1.1 (0.46-2)	3% (1.4-5.2%)	3.3% (1.1-6.4%)	4.7 (1.6-9.3)	410 (200-720)
Guinea-Bissau	1990	Female	25 to 29	32% (16-51%)	68% (49-84%)	1.3 (0.56-2.4)	3.9% (2-6.5%)	4.7% (2-8.4%)	6.9 (2.9-12)	510 (260-830)
Guinea-Bissau	1990	Female	30 to 34	34% (18-53%)	66% (47-82%)	1.5 (0.69-2.5)	4.5% (2.5-7.5%)	5.7% (2.8-10%)	9 (4.4-16)	570 (310-960)
Guinea-Bissau	1990	Female	35 to 39	38% (21-57%)	62% (43-79%)	1.6 (0.74-2.7)	5.3% (2.8-8.6%)	6.9% (3.5-11%)	13 (6.4-21)	700 (380-1100)
Guinea-Bissau	1990	Female	40 to 44	40% (23-59%)	60% (41-77%)	1.6 (0.78-3)	6% (3.1-9.8%)	8.1% (4-13%)	14 (6.9-24)	690 (340-1100)
Guinea-Bissau	1990	Female	45 to 49	40% (21-60%)	60% (40-79%)	1.7 (0.71-3)	6.6% (3.1-11%)	8.7% (4.1-15%)	18 (7.8-31)	740 (340-1300)
Guinea-Bissau	1990	Female	50 to 54	40% (23-61%)	60% (39-77%)	1.8 (0.75-3.3)	6.6% (2.9-11%)	8.4% (3.6-14%)	20 (8.1-34)	720 (300-1300)
Guinea-Bissau	1990	Female	55 to 59	41% (24-60%)	59% (40-76%)	1.9 (0.82-3.6)	7.1% (2.6-13%)	8.7% (3.1-15%)	26 (9-47)	810 (280-1500)
Guinea-Bissau	1990	Female	60 to 64	42% (24-61%)	58% (39-76%)	1.9 (0.77-3.7)	6.3% (1.9-11%)	7.5% (2.3-14%)	29 (8.8-58)	780 (240-1500)
Guinea-Bissau	1990	Female	65 to 69	43% (26-61%)	57% (39-74%)	1.8 (0.78-3.2)	5.4% (1.3-11%)	6.3% (1.4-13%)	32 (7.8-65)	720 (170-1500)
Guinea-Bissau	1990	Female	70 to 74	41% (25-59%)	59% (41-75%)	1.5 (0.6-2.9)	4.5% (0.41-11%)	5.2% (0.48-12%)	29 (2.7-71)	510 (49-1300)
Guinea-Bissau	1990	Female	75 to 79	37% (20-56%)	63% (44-80%)	1.2 (0.38-2.6)	3% (-0.35-8.6%)	3.4% (-0.43-10%)	16 (-1.9-47)	220 (-25-650)
Guinea-Bissau	1990	Female	80 to 84	35% (19-53%)	65% (47-81%)	1.1 (0.36-2.2)	2.1% (-0.44-6.4%)	2.4% (-0.51-7.3%)	5.6 (-1.2-17)	59 (-12-180)
Guinea-Bissau	1990	Female	85 to 89	35% (20-53%)	65% (47-80%)	1.1 (0.38-2.2)	1.7% (-0.64-5.4%)	1.9% (-0.79-6.2%)	1.9 (-0.78-6)	14 (-5.2-44)
Guinea-Bissau	1990	Female	90 to 94	38% (21-56%)	62% (44-79%)	1.1 (0.37-2.4)	1.3% (-0.92-5.1%)	1.5% (-1.2-6.1%)	0.43 (-0.34-1.8)	2.3 (-1.5-9.1)
Guinea-Bissau	1990	Female	95 plus	42% (23-62%)	58% (38-77%)	1.2 (0.32-2.8)	1.2% (-1.3-5.6%)	1.3% (-1.7-6.4%)	0.1 (-0.14-0.53)	0.39 (-0.42-1.8)
Guinea-Bissau	1990	Male	All Ages	21% (16-27%)	79% (73-84%)	0.91 (0.68-1.2)	2.3% (1.6-3%)	4% (2.6-5.5%)	370 (240-520)	14000 (10000-19000)
Guinea-Bissau	1990	Male	15 to 19	16% (8.3-27%)	84% (73-92%)	0.53 (0.19-1.1)	1.8% (0.83-3.5%)	2% (0.74-4.3%)	3.3 (1.1-6.7)	280 (120-540)
Guinea-Bissau	1990	Male	20 to 24	24% (13-39%)	76% (61-87%)	1.1 (0.44-2)	4.9% (2.4-8.5%)	5.4% (2.3-9.9%)	11 (4.4-20)	810 (390-1400)
Guinea-Bissau	1990	Male	25 to 29	29% (15-46%)	71% (54-85%)	1.6 (0.67-2.8)	7.9% (4-13%)	9.2% (4.1-15%)	18 (8.3-30)	1200 (600-1900)
Guinea-Bissau	1990	Male	30 to 34	33% (17-52%)	67% (48-83%)	2 (0.93-3.4)	9.9% (5.7-15%)	12% (6.3-18%)	25 (14-39)	1500 (850-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	1990	Male	35 to 39	36% (20-55%)	64% (45-80%)	2.2 (0.96-3.8)	11% (6.2-17%)	13% (7.2-20%)	33 (18-50)	1700 (970-2600)
Guinea-Bissau	1990	Male	40 to 44	37% (21-57%)	63% (43-79%)	2.3 (1-4.2)	12% (6.7-17%)	14% (7.8-21%)	37 (19-56)	1700 (930-2600)
Guinea-Bissau	1990	Male	45 to 49	38% (20-56%)	62% (44-80%)	2.4 (0.98-4.5)	12% (6.7-19%)	15% (7.8-22%)	43 (22-67)	1800 (950-2800)
Guinea-Bissau	1990	Male	50 to 54	38% (21-59%)	62% (41-79%)	2.4 (1-4.7)	12% (5.8-18%)	14% (6.7-21%)	41 (19-64)	1500 (720-2300)
Guinea-Bissau	1990	Male	55 to 59	38% (22-58%)	62% (42-78%)	2.4 (0.91-4.5)	11% (5.1-17%)	12% (5.8-19%)	42 (19-69)	1300 (620-2200)
Guinea-Bissau	1990	Male	60 to 64	38% (21-58%)	62% (42-79%)	2.1 (0.8-4.2)	8.1% (3.3-14%)	9.4% (3.8-16%)	38 (15-66)	1000 (420-1800)
Guinea-Bissau	1990	Male	65 to 69	38% (22-56%)	62% (44-78%)	1.8 (0.74-3.4)	6.5% (2.5-12%)	7.4% (2.8-13%)	34 (13-64)	770 (300-1400)
Guinea-Bissau	1990	Male	70 to 74	36% (22-54%)	64% (46-78%)	1.4 (0.52-2.8)	4.8% (1.4-10%)	5.5% (1.5-12%)	26 (6.9-56)	460 (130-1000)
Guinea-Bissau	1990	Male	75 to 79	33% (19-52%)	67% (48-81%)	1.1 (0.33-2.3)	3.3% (0.62-7.7%)	3.7% (0.66-8.8%)	13 (2.3-30)	180 (33-430)
Guinea-Bissau	1990	Male	80 to 84	31% (16-49%)	69% (51-84%)	0.89 (0.27-1.8)	2.5% (0.46-6.1%)	2.9% (0.48-7.1%)	4.5 (0.73-11)	47 (8.6-110)
Guinea-Bissau	1990	Male	85 to 89	29% (15-47%)	71% (53-85%)	0.77 (0.25-1.6)	1.9% (0.22-4.8%)	2.1% (0.17-5.5%)	1.4 (0.11-3.6)	10 (1.2-26)
Guinea-Bissau	1990	Male	90 to 94	31% (16-49%)	69% (51-84%)	0.72 (0.22-1.6)	1.1% (-0.15-3.6%)	1.2% (-0.32-4.2%)	0.23 (-0.061-0.8)	1.3 (-0.17-4.2)
Guinea-Bissau	1990	Male	95 plus	35% (18-56%)	65% (44-82%)	0.72 (0.16-1.8)	1% (-0.45-4%)	1% (-0.75-4.6%)	0.05 (-0.037-0.23)	0.2 (-0.091-0.79)
Guinea-Bissau	1995	Female	All Ages	25% (20-32%)	75% (68-80%)	0.73 (0.54-0.96)	1.5% (0.84-2.2%)	2.7% (1.3-4.6%)	220 (100-390)	7700 (4400-12000)
Guinea-Bissau	1995	Female	15 to 19	24% (13-38%)	76% (62-87%)	0.66 (0.26-1.3)	1.5% (0.62-2.9%)	1.8% (0.45-3.9%)	2.3 (0.58-5.1)	220 (86-420)
Guinea-Bissau	1995	Female	20 to 24	29% (16-46%)	71% (54-84%)	1 (0.44-1.9)	2.9% (1.3-4.9%)	3.1% (1-6%)	5 (1.6-9.8)	440 (210-760)
Guinea-Bissau	1995	Female	25 to 29	32% (16-50%)	68% (50-84%)	1.3 (0.54-2.3)	3.7% (1.8-6.1%)	4.3% (1.7-7.7%)	7.1 (2.8-13)	530 (260-880)
Guinea-Bissau	1995	Female	30 to 34	35% (18-54%)	65% (46-82%)	1.4 (0.68-2.5)	4.2% (2.2-6.9%)	5.2% (2.5-9%)	8.8 (4.2-16)	570 (300-950)
Guinea-Bissau	1995	Female	35 to 39	38% (21-58%)	62% (42-79%)	1.5 (0.72-2.6)	4.9% (2.5-8%)	6.3% (3-11%)	12 (5.7-21)	670 (340-1100)
Guinea-Bissau	1995	Female	40 to 44	40% (23-59%)	60% (41-77%)	1.6 (0.75-2.7)	5.6% (2.9-9.2%)	7.5% (3.6-12%)	17 (7.9-28)	800 (390-1300)
Guinea-Bissau	1995	Female	45 to 49	41% (21-59%)	59% (41-79%)	1.6 (0.69-3)	6.2% (2.9-10%)	8.1% (3.7-14%)	18 (8-32)	760 (350-1300)
Guinea-Bissau	1995	Female	50 to 54	41% (23-61%)	59% (39-77%)	1.7 (0.74-3.2)	6.3% (2.5-11%)	7.9% (3.2-14%)	22 (8.5-39)	810 (310-1400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	1995	Female	55 to 59	41% (24-59%)	59% (41-76%)	1.9 (0.77-3.4)	6.7% (2.4-12%)	8.2% (2.8-15%)	25 (8.2-47)	800 (260-1500)
Guinea-Bissau	1995	Female	60 to 64	42% (24-61%)	58% (39-76%)	1.9 (0.74-3.6)	5.8% (1.7-11%)	6.9% (2-13%)	27 (7.7-52)	720 (210-1400)
Guinea-Bissau	1995	Female	65 to 69	43% (26-61%)	57% (39-74%)	1.7 (0.77-3.1)	5% (0.97-10%)	5.8% (1.1-12%)	29 (5.4-60)	640 (120-1400)
Guinea-Bissau	1995	Female	70 to 74	41% (26-59%)	59% (41-74%)	1.4 (0.56-2.8)	4% (0.3-9.8%)	4.6% (0.35-11%)	27 (2-69)	480 (36-1200)
Guinea-Bissau	1995	Female	75 to 79	37% (21-57%)	63% (43-79%)	1.1 (0.38-2.5)	2.6% (-0.55-8.1%)	3% (-0.64-9.2%)	15 (-3.4-49)	210 (-44-680)
Guinea-Bissau	1995	Female	80 to 84	35% (19-53%)	65% (47-81%)	1 (0.35-2.1)	1.9% (-0.55-6.1%)	2.2% (-0.67-7%)	6.2 (-2-21)	65 (-19-210)
Guinea-Bissau	1995	Female	85 to 89	35% (20-53%)	65% (47-80%)	1 (0.37-2.1)	1.5% (-0.69-5.5%)	1.7% (-0.84-6.2%)	2.3 (-1.1-8.2)	17 (-7.7-60)
Guinea-Bissau	1995	Female	90 to 94	38% (22-55%)	62% (45-78%)	1.1 (0.35-2.2)	1.2% (-1-5.1%)	1.4% (-1.3-6%)	0.52 (-0.49-2.3)	2.8 (-2.2-12)
Guinea-Bissau	1995	Female	95 plus	42% (23-62%)	58% (38-77%)	1.1 (0.29-2.6)	1.1% (-1.3-5.6%)	1.2% (-1.7-6.5%)	0.11 (-0.16-0.59)	0.41 (-0.47-2)
Guinea-Bissau	1995	Male	All Ages	21% (16-27%)	79% (73-84%)	0.88 (0.65-1.1)	4.1% (1.7-3.2%)	4.1% (2.7-5.7%)	390 (250-550)	15000 (11000-20000)
Guinea-Bissau	1995	Male	15 to 19	16% (8.5-27%)	84% (73-91%)	0.53 (0.19-1.1)	1.8% (0.81-3.5%)	2.1% (0.74-4.3%)	3.7 (1.3-7.8)	320 (140-610)
Guinea-Bissau	1995	Male	20 to 24	24% (13-40%)	76% (60-87%)	1.1 (0.45-2)	4.8% (2.4-8.4%)	5.3% (2.2-9.8%)	12 (5-22)	910 (440-1600)
Guinea-Bissau	1995	Male	25 to 29	30% (16-47%)	70% (53-84%)	1.6 (0.64-2.8)	7.5% (3.7-12%)	8.7% (4-15%)	20 (8.8-33)	1300 (640-2100)
Guinea-Bissau	1995	Male	30 to 34	34% (18-52%)	66% (48-82%)	1.9 (0.95-3.3)	9.2% (5.2-14%)	11% (5.7-17%)	26 (14-40)	1500 (860-2300)
Guinea-Bissau	1995	Male	35 to 39	37% (20-56%)	63% (44-80%)	2.1 (0.94-3.9)	10% (5.5-15%)	12% (6.6-19%)	32 (17-49)	1700 (940-2600)
Guinea-Bissau	1995	Male	40 to 44	38% (22-57%)	62% (43-78%)	2.2 (0.98-4.1)	11% (6-17%)	13% (7.1-20%)	42 (22-65)	2000 (1000-3000)
Guinea-Bissau	1995	Male	45 to 49	39% (21-57%)	61% (43-79%)	2.4 (0.99-4.4)	12% (6.6-18%)	14% (7.6-21%)	45 (24-70)	1900 (1000-2900)
Guinea-Bissau	1995	Male	50 to 54	39% (22-60%)	61% (40-78%)	2.4 (1-4.6)	11% (5.4-17%)	13% (6.4-20%)	47 (22-74)	1700 (840-2700)
Guinea-Bissau	1995	Male	55 to 59	39% (23-59%)	61% (41-77%)	2.4 (0.94-4.5)	10% (4.9-16%)	12% (5.6-19%)	43 (20-72)	1400 (630-2300)
Guinea-Bissau	1995	Male	60 to 64	39% (22-59%)	61% (41-78%)	2.1 (0.79-4.4)	7.9% (3.1-14%)	9.1% (3.6-16%)	37 (15-67)	1000 (410-1800)
Guinea-Bissau	1995	Male	65 to 69	39% (22-57%)	61% (43-78%)	1.8 (0.74-3.5)	6.2% (2.4-12%)	7.1% (2.8-14%)	33 (12-63)	740 (280-1400)
Guinea-Bissau	1995	Male	70 to 74	37% (22-55%)	63% (45-78%)	1.4 (0.53-2.8)	4.6% (1.3-10%)	5.2% (1.4-12%)	26 (6.7-58)	470 (130-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	1995	Male	75 to 79	34% (20-54%)	66% (46-80%)	1.1 (0.35-2.3)	3.1% (0.5-8%)	3.5% (0.54-9.1%)	13 (2-35)	190 (30-480)
Guinea-Bissau	1995	Male	80 to 84	32% (16-51%)	68% (49-84%)	0.89 (0.29-1.9)	2.4% (0.4-5.7%)	2.7% (0.39-6.5%)	5.1 (0.78-12)	54 (9.4-130)
Guinea-Bissau	1995	Male	85 to 89	30% (15-48%)	70% (52-85%)	0.76 (0.25-1.6)	1.8% (0.21-4.8%)	2% (0.19-5.5%)	1.7 (0.16-4.8)	13 (1.6-35)
Guinea-Bissau	1995	Male	90 to 94	32% (16-50%)	68% (50-84%)	0.73 (0.22-1.7)	1.2% (-0.2-3.9%)	1.3% (-0.36-4.5%)	0.31 (-0.089-1.1)	1.7 (-0.29-5.8)
Guinea-Bissau	1995	Male	95 plus	37% (18-58%)	63% (42-82%)	0.73 (0.16-1.9)	1.2% (-0.44-4.8%)	1.2% (-0.71-5.6%)	0.068 (-0.039-0.31)	0.26 (-0.094-1.1)
Guinea-Bissau	2000	Female	All Ages	25% (20-32%)	75% (68-80%)	0.71 (0.52-0.92)	1.5% (0.87-2.4%)	2.7% (1.2-4.8%)	230 (100-410)	8200 (4600-13000)
Guinea-Bissau	2000	Female	15 to 19	24% (13-39%)	76% (61-87%)	0.64 (0.26-1.2)	1.5% (0.63-3%)	1.8% (0.43-4.2%)	2.7 (0.64-6.2)	260 (100-500)
Guinea-Bissau	2000	Female	20 to 24	29% (15-46%)	71% (54-85%)	1 (0.44-1.9)	2.7% (1.3-4.7%)	2.9% (0.93-5.7%)	5.4 (1.9-11)	490 (230-860)
Guinea-Bissau	2000	Female	25 to 29	32% (17-50%)	68% (50-83%)	1.2 (0.54-2.1)	3.2% (1.6-5.4%)	3.6% (1.3-6.7%)	7.6 (3-14)	570 (280-970)
Guinea-Bissau	2000	Female	30 to 34	35% (19-53%)	65% (47-81%)	1.4 (0.66-2.4)	3.7% (1.9-6.1%)	4.4% (2-7.7%)	9.3 (4.3-16)	600 (320-990)
Guinea-Bissau	2000	Female	35 to 39	39% (22-57%)	61% (43-78%)	1.5 (0.68-2.5)	4.3% (2.2-7.2%)	5.4% (2.5-9.3%)	12 (5.6-22)	690 (330-1200)
Guinea-Bissau	2000	Female	40 to 44	40% (24-59%)	60% (41-76%)	1.5 (0.7-2.7)	5.1% (2.5-8.6%)	6.7% (3.2-11%)	16 (7.2-28)	790 (380-1400)
Guinea-Bissau	2000	Female	45 to 49	41% (22-59%)	59% (41-78%)	1.6 (0.68-2.8)	5.8% (2.6-10%)	7.5% (3.2-13%)	22 (9.4-40)	930 (400-1700)
Guinea-Bissau	2000	Female	50 to 54	41% (23-61%)	59% (39-77%)	1.6 (0.73-3)	6% (2.4-11%)	7.5% (3-14%)	24 (9-44)	860 (340-1600)
Guinea-Bissau	2000	Female	55 to 59	41% (25-60%)	59% (40-75%)	1.8 (0.75-3.3)	6.5% (2.2-12%)	8% (2.6-15%)	30 (9.6-57)	930 (310-1800)
Guinea-Bissau	2000	Female	60 to 64	42% (24-61%)	58% (39-76%)	1.8 (0.7-3.6)	5.5% (1.5-11%)	6.6% (1.8-13%)	27 (7.1-55)	720 (190-1500)
Guinea-Bissau	2000	Female	65 to 69	44% (26-60%)	56% (40-74%)	1.6 (0.73-3)	4.7% (0.8-10%)	5.4% (0.92-12%)	27 (4.7-59)	600 (100-1300)
Guinea-Bissau	2000	Female	70 to 74	41% (25-59%)	59% (41-75%)	1.4 (0.55-2.7)	3.6% (-0.071-9.6%)	4.2% (-0.057-11%)	24 (-0.3-64)	420 (-8.3-1100)
Guinea-Bissau	2000	Female	75 to 79	37% (20-55%)	63% (45-80%)	1.1 (0.36-2.4)	2.4% (-0.63-7.6%)	2.7% (-0.75-8.7%)	14 (-4.1-46)	200 (-55-640)
Guinea-Bissau	2000	Female	80 to 84	35% (18-53%)	65% (47-82%)	1 (0.33-2)	1.8% (-0.71-6.2%)	2.1% (-0.83-7.2%)	6.6 (-2.7-23)	69 (-27-240)
Guinea-Bissau	2000	Female	85 to 89	35% (20-52%)	65% (48-80%)	0.98 (0.36-2.1)	1.5% (-0.79-5.4%)	1.7% (-0.98-6.3%)	2.6 (-1.5-9.7)	19 (-10-71)
Guinea-Bissau	2000	Female	90 to 94	38% (22-54%)	62% (46-78%)	1 (0.35-2.2)	1.2% (-1.1-5.5%)	1.4% (-1.3-6.6%)	0.67 (-0.63-3.2)	3.6 (-3-16)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	2000	Female	95 plus	42% (22-62%)	58% (38-78%)	1.1 (0.3-2.5)	1.2% (-1.4-5.9%)	1.2% (-1.8-6.8%)	0.14 (-0.21-0.75)	0.51 (-0.63-2.5)
Guinea-Bissau	2000	Male	All Ages	22% (17-28%)	78% (72-83%)	0.87 (0.64-1.1)	2.6% (1.8-3.4%)	4.3% (2.7-6%)	420 (270-600)	17000 (11000-22000)
Guinea-Bissau	2000	Male	15 to 19	17% (8.6-29%)	83% (71-91%)	0.54 (0.2-1.1)	2% (0.85-3.7%)	2.2% (0.8-4.7%)	4.6 (1.5-9.5)	390 (170-750)
Guinea-Bissau	2000	Male	20 to 24	25% (13-41%)	75% (59-87%)	1.1 (0.44-2)	4.8% (2.3-8.6%)	5.4% (2.2-10%)	14 (5.6-26)	1000 (480-1900)
Guinea-Bissau	2000	Male	25 to 29	31% (17-48%)	69% (52-83%)	1.6 (0.65-2.8)	7.1% (3.5-12%)	8.1% (3.6-14%)	22 (10-38)	1500 (740-2500)
Guinea-Bissau	2000	Male	30 to 34	35% (18-53%)	65% (47-82%)	1.9 (0.92-3.3)	8.5% (4.6-13%)	9.9% (5.2-16%)	28 (14-45)	1700 (920-2600)
Guinea-Bissau	2000	Male	35 to 39	38% (21-57%)	62% (43-79%)	2.1 (0.91-3.7)	9.3% (5.1-14%)	11% (5.8-17%)	34 (18-52)	1800 (960-2700)
Guinea-Bissau	2000	Male	40 to 44	39% (23-58%)	61% (42-77%)	2.2 (0.96-3.9)	10% (5.6-16%)	12% (6.3-19%)	42 (21-66)	2000 (1000-3100)
Guinea-Bissau	2000	Male	45 to 49	39% (21-57%)	61% (43-79%)	2.3 (0.97-4.4)	11% (5.9-17%)	13% (6.9-20%)	54 (27-84)	2200 (1100-3500)
Guinea-Bissau	2000	Male	50 to 54	39% (22-60%)	61% (40-78%)	2.3 (0.96-4.6)	11% (5.2-17%)	13% (6-20%)	51 (23-83)	1900 (870-3000)
Guinea-Bissau	2000	Male	55 to 59	39% (23-58%)	61% (42-77%)	2.3 (0.9-4.5)	10% (4.5-16%)	12% (5.1-19%)	51 (23-84)	1600 (750-2600)
Guinea-Bissau	2000	Male	60 to 64	40% (22-59%)	60% (41-78%)	2.1 (0.78-4.4)	7.7% (3-14%)	8.9% (3.4-16%)	39 (14-71)	1000 (390-1900)
Guinea-Bissau	2000	Male	65 to 69	40% (23-58%)	60% (42-77%)	1.8 (0.73-3.5)	6.1% (2.3-12%)	7% (2.5-14%)	33 (11-63)	740 (260-1400)
Guinea-Bissau	2000	Male	70 to 74	38% (23-56%)	62% (44-77%)	1.4 (0.53-2.7)	4.5% (1.1-10%)	5.1% (1.2-11%)	25 (5.7-58)	450 (110-1000)
Guinea-Bissau	2000	Male	75 to 79	35% (20-54%)	65% (46-80%)	1 (0.34-2.3)	3% (0.38-7.9%)	3.4% (0.4-8.9%)	14 (1.7-38)	200 (26-530)
Guinea-Bissau	2000	Male	80 to 84	32% (17-51%)	68% (49-83%)	0.88 (0.28-1.9)	2.4% (0.31-5.8%)	2.7% (0.31-6.7%)	5.9 (0.65-15)	62 (8.2-150)
Guinea-Bissau	2000	Male	85 to 89	31% (16-50%)	69% (50-84%)	0.76 (0.26-1.6)	1.8% (0.13-4.8%)	2.1% (0.062-5.5%)	2.1 (0.062-5.7)	16 (1.1-42)
Guinea-Bissau	2000	Male	90 to 94	33% (17-51%)	67% (49-83%)	0.72 (0.22-1.6)	1.2% (-0.19-3.9%)	1.3% (-0.36-4.5%)	0.43 (-0.12-1.5)	2.4 (-0.36-7.6)
Guinea-Bissau	2000	Male	95 plus	37% (18-59%)	63% (41-82%)	0.72 (0.17-1.8)	1.4% (-0.43-5.1%)	1.5% (-0.67-6%)	0.1 (-0.047-0.42)	0.38 (-0.12-1.5)
Guinea-Bissau	2005	Female	All Ages	25% (20-32%)	75% (68-80%)	0.72 (0.53-0.96)	1.6% (0.88-2.5%)	2.8% (1.2-4.9%)	250 (110-450)	8800 (4900-14000)
Guinea-Bissau	2005	Female	15 to 19	23% (13-38%)	77% (62-87%)	0.63 (0.25-1.2)	1.5% (0.62-2.9%)	1.8% (0.47-4.2%)	2.7 (0.71-6.3)	270 (110-530)
Guinea-Bissau	2005	Female	20 to 24	29% (15-47%)	71% (53-85%)	1 (0.44-1.8)	2.7% (1.2-4.8%)	2.8% (0.84-5.9%)	6.2 (1.9-13)	560 (270-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	2005	Female	25 to 29	32% (16-50%)	68% (50-84%)	1.2 (0.55-2.2)	2.8% (1.3-4.8%)	3% (1.1-6%)	8.3 (3.3-16)	640 (320-1100)
Guinea-Bissau	2005	Female	30 to 34	35% (18-53%)	65% (47-82%)	1.4 (0.68-2.5)	3% (1.6-5.2%)	3.4% (1.5-6.2%)	10 (4.5-18)	660 (340-1100)
Guinea-Bissau	2005	Female	35 to 39	38% (21-57%)	62% (43-79%)	1.5 (0.72-2.5)	3.7% (1.8-6.3%)	4.5% (2-7.8%)	13 (6-23)	730 (360-1200)
Guinea-Bissau	2005	Female	40 to 44	40% (24-58%)	60% (42-76%)	1.6 (0.73-2.7)	4.5% (2.2-7.7%)	5.7% (2.5-9.9%)	17 (7.3-30)	810 (360-1400)
Guinea-Bissau	2005	Female	45 to 49	41% (22-59%)	59% (41-78%)	1.6 (0.68-2.9)	5.4% (2.2-9.3%)	6.8% (2.8-12%)	22 (8.6-40)	920 (380-1700)
Guinea-Bissau	2005	Female	50 to 54	41% (23-61%)	59% (39-77%)	1.7 (0.73-3.1)	5.8% (2.2-11%)	7.3% (2.7-13%)	29 (10-55)	1100 (400-2000)
Guinea-Bissau	2005	Female	55 to 59	41% (24-59%)	59% (41-76%)	1.8 (0.75-3.3)	6.5% (2.1-12%)	8% (2.6-15%)	31 (10-62)	990 (320-1900)
Guinea-Bissau	2005	Female	60 to 64	42% (25-61%)	58% (39-75%)	1.8 (0.73-3.5)	5.6% (1.2-11%)	6.7% (1.4-13%)	31 (6.3-65)	830 (170-1800)
Guinea-Bissau	2005	Female	65 to 69	43% (26-60%)	57% (40-74%)	1.7 (0.72-3)	4.7% (0.69-10%)	5.5% (0.83-13%)	27 (4.1-62)	600 (83-1400)
Guinea-Bissau	2005	Female	70 to 74	41% (26-58%)	59% (42-74%)	1.4 (0.54-2.8)	3.8% (-0.083-9.9%)	4.4% (-0.075-11%)	23 (-0.41-62)	410 (-9.5-1100)
Guinea-Bissau	2005	Female	75 to 79	37% (21-56%)	63% (44-79%)	1.1 (0.37-2.5)	2.5% (-0.66-8.2%)	2.8% (-0.77-9.3%)	14 (-3.7-46)	190 (-50-630)
Guinea-Bissau	2005	Female	80 to 84	35% (19-54%)	65% (46-81%)	1 (0.34-2.1)	2% (-0.64-6.7%)	2.3% (-0.75-7.8%)	7.2 (-2.4-25)	75 (-23-260)
Guinea-Bissau	2005	Female	85 to 89	35% (19-54%)	65% (46-81%)	1 (0.36-2.1)	1.7% (-0.8-6.2%)	1.9% (-0.97-7.2%)	3.3 (-1.6-12)	24 (-11-91)
Guinea-Bissau	2005	Female	90 to 94	38% (21-55%)	62% (45-79%)	1 (0.35-2.3)	1.5% (-1-6.1%)	1.7% (-1.3-7.3%)	0.91 (-0.73-4)	4.7 (-3.4-20)
Guinea-Bissau	2005	Female	95 plus	42% (22-63%)	58% (37-78%)	1.1 (0.29-2.6)	1.4% (-1.4-6.3%)	1.5% (-1.8-7.2%)	0.2 (-0.24-0.95)	0.72 (-0.7-3.2)
Guinea-Bissau	2005	Male	All Ages	21% (16-27%)	79% (73-84%)	0.89 (0.67-1.2)	2.9% (2-3.9%)	4.7% (3-6.6%)	490 (310-700)	19000 (13000-26000)
Guinea-Bissau	2005	Male	15 to 19	17% (8.6-28%)	83% (72-91%)	0.53 (0.2-1.1)	2% (0.89-3.8%)	2.3% (0.8-4.8%)	5.1 (1.7-11)	430 (190-820)
Guinea-Bissau	2005	Male	20 to 24	25% (13-41%)	75% (59-87%)	1.1 (0.44-2)	5% (2.4-8.6%)	5.5% (2.3-9.9%)	17 (7.3-30)	1300 (620-2200)
Guinea-Bissau	2005	Male	25 to 29	30% (16-48%)	70% (52-84%)	1.6 (0.68-2.9)	6.9% (3.3-11%)	7.9% (3.5-13%)	26 (12-44)	1700 (870-2800)
Guinea-Bissau	2005	Male	30 to 34	34% (18-52%)	66% (48-82%)	1.9 (0.93-3.3)	8% (4.6-12%)	9.2% (5-14%)	33 (18-51)	2000 (1100-3000)
Guinea-Bissau	2005	Male	35 to 39	37% (21-56%)	63% (44-79%)	2.1 (0.95-3.7)	8.9% (4.9-14%)	10% (5.5-16%)	39 (21-60)	2100 (1100-3100)
Guinea-Bissau	2005	Male	40 to 44	38% (22-57%)	62% (43-78%)	2.2 (0.96-4.1)	9.8% (5-15%)	11% (5.8-18%)	47 (24-73)	2200 (1100-3400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	2005	Male	45 to 49	39% (21-58%)	61% (42-79%)	2.3 (0.99-4.5)	11% (5.8-17%)	13% (6.6-20%)	58 (29-91)	2400 (1200-3700)
Guinea-Bissau	2005	Male	50 to 54	39% (22-60%)	61% (40-78%)	2.4 (0.97-4.6)	11% (5.5-18%)	13% (6.4-20%)	65 (31-110)	2400 (1200-3900)
Guinea-Bissau	2005	Male	55 to 59	39% (24-58%)	61% (42-76%)	2.3 (0.94-4.4)	10% (4.7-17%)	12% (5.3-19%)	59 (25-97)	1900 (800-3000)
Guinea-Bissau	2005	Male	60 to 64	39% (23-58%)	61% (42-77%)	2.1 (0.77-4.2)	8.1% (3.1-14%)	9.3% (3.5-16%)	50 (18-90)	1300 (500-2400)
Guinea-Bissau	2005	Male	65 to 69	39% (22-57%)	61% (43-78%)	1.8 (0.73-3.5)	6.4% (2.3-12%)	7.3% (2.6-14%)	37 (12-72)	830 (280-1600)
Guinea-Bissau	2005	Male	70 to 74	38% (22-55%)	62% (45-78%)	1.4 (0.53-2.9)	4.8% (1.1-11%)	5.4% (1.2-12%)	28 (6.2-65)	500 (110-1200)
Guinea-Bissau	2005	Male	75 to 79	34% (19-54%)	66% (46-81%)	1.1 (0.33-2.4)	3.2% (0.37-8.5%)	3.6% (0.38-9.6%)	16 (1.6-42)	220 (24-580)
Guinea-Bissau	2005	Male	80 to 84	32% (16-50%)	68% (50-84%)	0.89 (0.28-1.9)	2.6% (0.36-6.4%)	2.9% (0.36-7.3%)	7.1 (0.9-18)	74 (11-190)
Guinea-Bissau	2005	Male	85 to 89	30% (16-48%)	70% (52-84%)	0.77 (0.25-1.7)	2% (0.17-5.3%)	2.3% (0.1-6.1%)	2.7 (0.12-7.4)	20 (1.7-55)
Guinea-Bissau	2005	Male	90 to 94	32% (17-50%)	68% (50-83%)	0.73 (0.22-1.7)	1.4% (-0.16-4.4%)	1.5% (-0.31-5.2%)	0.58 (-0.12-2)	3.1 (-0.36-10)
Guinea-Bissau	2005	Male	95 plus	37% (18-58%)	63% (42-82%)	0.73 (0.17-1.8)	1.5% (-0.37-5.6%)	1.7% (-0.64-6.6%)	0.15 (-0.056-0.58)	0.54 (-0.13-2)
Guinea-Bissau	2010	Female	All Ages	25% (20-32%)	75% (68-80%)	0.73 (0.52-1)	1.7% (0.92-2.8%)	2.9% (1.2-5.3%)	250 (100-480)	9200 (4800-15000)
Guinea-Bissau	2010	Female	15 to 19	24% (13-37%)	76% (63-87%)	0.64 (0.25-1.3)	1.5% (0.61-2.9%)	1.8% (0.41-4%)	2.7 (0.63-6.2)	280 (110-540)
Guinea-Bissau	2010	Female	20 to 24	29% (15-47%)	71% (53-85%)	1 (0.4-1.9)	2.6% (1.3-4.8%)	2.8% (0.82-5.9%)	5.9 (1.8-12)	570 (270-1000)
Guinea-Bissau	2010	Female	25 to 29	32% (16-51%)	68% (49-84%)	1.2 (0.53-2.2)	2.6% (1.3-4.5%)	2.7% (1-5.4%)	8.7 (3.5-16)	690 (360-1200)
Guinea-Bissau	2010	Female	30 to 34	35% (17-53%)	65% (47-83%)	1.4 (0.64-2.5)	2.7% (1.4-4.6%)	2.9% (1.3-5.5%)	11 (4.9-19)	700 (370-1200)
Guinea-Bissau	2010	Female	35 to 39	38% (21-57%)	62% (43-79%)	1.5 (0.7-2.7)	3.2% (1.5-5.7%)	3.8% (1.7-6.8%)	14 (6.1-25)	780 (380-1300)
Guinea-Bissau	2010	Female	40 to 44	40% (23-59%)	60% (41-77%)	1.6 (0.69-2.8)	4.2% (1.9-7.3%)	5.1% (2.3-8.9%)	18 (7.8-32)	870 (400-1500)
Guinea-Bissau	2010	Female	45 to 49	41% (22-59%)	59% (41-78%)	1.6 (0.67-3)	4.9% (2-8.9%)	6.2% (2.5-11%)	22 (8.8-41)	940 (390-1700)
Guinea-Bissau	2010	Female	50 to 54	41% (23-62%)	59% (38-77%)	1.7 (0.7-3.3)	5.5% (2-11%)	6.9% (2.5-13%)	28 (9.9-56)	1000 (370-2000)
Guinea-Bissau	2010	Female	55 to 59	41% (24-60%)	59% (40-76%)	1.8 (0.71-3.5)	6.2% (1.9-12%)	7.6% (2.3-15%)	37 (11-76)	1200 (340-2400)
Guinea-Bissau	2010	Female	60 to 64	42% (24-61%)	58% (39-76%)	1.8 (0.69-3.7)	5.3% (0.95-11%)	6.3% (1.1-14%)	31 (5.6-67)	840 (150-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	2010	Female	65 to 69	43% (26-61%)	57% (39-74%)	1.6 (0.69-3.2)	4.4% (0.43-11%)	5.2% (0.48-12%)	30 (2.8-72)	660 (61-1600)
Guinea-Bissau	2010	Female	70 to 74	41% (25-58%)	59% (42-75%)	1.4 (0.51-2.8)	3.5% (-0.33-9.3%)	4.1% (-0.44-11%)	22 (-2.3-60)	390 (-40-1100)
Guinea-Bissau	2010	Female	75 to 79	37% (21-56%)	63% (44-79%)	1.1 (0.36-2.4)	2.3% (-0.78-7.8%)	2.7% (-0.95-9.1%)	13 (-4.4-44)	180 (-61-630)
Guinea-Bissau	2010	Female	80 to 84	35% (18-53%)	65% (47-82%)	1 (0.36-2.1)	1.8% (-0.87-6.5%)	2.1% (-1-7.5%)	6.5 (-3.1-23)	68 (-31-240)
Guinea-Bissau	2010	Female	85 to 89	35% (19-53%)	65% (47-81%)	0.99 (0.35-2.1)	1.5% (-0.96-6.1%)	1.8% (-1.2-7.2%)	3.2 (-2.1-13)	24 (-14-95)
Guinea-Bissau	2010	Female	90 to 94	38% (21-55%)	62% (45-79%)	1 (0.33-2.2)	1.3% (-1.2-5.8%)	1.5% (-1.5-6.9%)	0.95 (-0.98-4.5)	5 (-4.7-23)
Guinea-Bissau	2010	Female	95 plus	42% (22-63%)	58% (37-78%)	1.1 (0.27-2.5)	1.2% (-1.4-6.4%)	1.3% (-1.8-7.4%)	0.2 (-0.28-1.1)	0.76 (-0.87-3.9)
Guinea-Bissau	2010	Male	All Ages	22% (17-28%)	78% (72-83%)	0.9 (0.63-1.2)	3.3% (2.2-4.4%)	5.1% (3.1-7.3%)	530 (310-770)	21000 (14000-28000)
Guinea-Bissau	2010	Male	15 to 19	17% (8.9-28%)	83% (72-91%)	0.53 (0.19-1.1)	2% (0.88-3.8%)	2.3% (0.78-4.8%)	5.2 (1.9-11)	450 (200-860)
Guinea-Bissau	2010	Male	20 to 24	25% (13-40%)	75% (60-87%)	1.1 (0.42-2)	4.9% (2.3-8.4%)	5.4% (2.1-10%)	17 (6.6-31)	1300 (600-2200)
Guinea-Bissau	2010	Male	25 to 29	30% (16-47%)	70% (53-84%)	1.5 (0.68-2.8)	6.6% (3.2-11%)	7.5% (3.3-13%)	29 (13-49)	1900 (940-3200)
Guinea-Bissau	2010	Male	30 to 34	34% (18-53%)	66% (47-82%)	1.9 (0.9-3.4)	7.5% (4-12%)	8.6% (4.3-14%)	36 (19-56)	2100 (1200-3300)
Guinea-Bissau	2010	Male	35 to 39	37% (20-57%)	63% (43-80%)	2.1 (0.93-3.9)	8.2% (4.4-13%)	9.5% (5-15%)	42 (22-66)	2200 (1200-3400)
Guinea-Bissau	2010	Male	40 to 44	38% (22-58%)	62% (42-78%)	2.2 (0.93-4.2)	9.4% (4.7-15%)	11% (5.3-17%)	50 (23-79)	2300 (1100-3600)
Guinea-Bissau	2010	Male	45 to 49	39% (21-57%)	61% (43-79%)	2.3 (0.91-4.6)	11% (5.4-16%)	12% (6.2-19%)	60 (29-97)	2500 (1200-4000)
Guinea-Bissau	2010	Male	50 to 54	40% (22-60%)	60% (40-78%)	2.4 (0.98-4.7)	11% (5.2-17%)	13% (6-20%)	65 (30-110)	2400 (1100-3900)
Guinea-Bissau	2010	Male	55 to 59	40% (24-60%)	60% (40-76%)	2.3 (0.91-4.6)	10% (4.5-17%)	12% (5.2-20%)	71 (30-120)	2300 (950-3700)
Guinea-Bissau	2010	Male	60 to 64	40% (23-59%)	60% (41-77%)	2.1 (0.76-4.4)	8.1% (3-14%)	9.3% (3.4-17%)	55 (19-100)	1500 (520-2700)
Guinea-Bissau	2010	Male	65 to 69	40% (24-57%)	60% (43-76%)	1.8 (0.68-3.6)	6.4% (2.2-12%)	7.3% (2.4-14%)	44 (14-86)	990 (330-1900)
Guinea-Bissau	2010	Male	70 to 74	38% (23-56%)	62% (44-77%)	1.4 (0.52-2.9)	4.6% (1.1-10%)	5.3% (1.2-12%)	29 (6.3-66)	520 (120-1200)
Guinea-Bissau	2010	Male	75 to 79	35% (19-53%)	65% (47-81%)	1.1 (0.32-2.3)	3.1% (0.44-8.1%)	3.5% (0.45-9.3%)	16 (2-41)	220 (29-570)
Guinea-Bissau	2010	Male	80 to 84	32% (16-51%)	68% (49-84%)	0.89 (0.28-1.9)	2.5% (0.38-6.2%)	2.8% (0.38-7.1%)	7.3 (0.95-19)	76 (12-190)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	2010	Male	85 to 89	31% (16-49%)	69% (51-84%)	0.76 (0.25-1.7)	2% (0.11-5.5%)	2.2% (0.019-6.2%)	2.9 (0.024-8.5)	22 (1.3-62)
Guinea-Bissau	2010	Male	90 to 94	32% (17-50%)	68% (50-83%)	0.71 (0.21-1.7)	1.3% (-0.17-4.5%)	1.5% (-0.33-5.3%)	0.65 (-0.14-2.4)	3.5 (-0.45-12)
Guinea-Bissau	2010	Male	95 plus	36% (18-58%)	64% (42-82%)	0.71 (0.16-1.9)	1.5% (-0.4-5.3%)	1.6% (-0.66-6.3%)	0.16 (-0.067-0.64)	0.6 (-0.16-2.2)
Guinea-Bissau	2016	Female	All Ages	26% (20-32%)	74% (68-80%)	0.74 (0.41-1.2)	2% (0.83-3.5%)	3.2% (0.89-6.7%)	270 (74-560)	9800 (4100-18000)
Guinea-Bissau	2016	Female	15 to 19	23% (13-39%)	77% (61-87%)	0.64 (0.22-1.4)	1.4% (0.58-2.7%)	1.6% (0.29-3.8%)	2.5 (0.46-6.1)	270 (110-550)
Guinea-Bissau	2016	Female	20 to 24	29% (15-46%)	71% (54-85%)	0.99 (0.35-2.1)	2.7% (1.3-4.8%)	2.8% (0.75-6.1%)	5.5 (1.5-12)	570 (280-1000)
Guinea-Bissau	2016	Female	25 to 29	32% (16-51%)	68% (49-84%)	1.2 (0.47-2.4)	2.7% (1.3-4.7%)	2.9% (0.91-5.9%)	8.4 (2.9-17)	700 (330-1200)
Guinea-Bissau	2016	Female	30 to 34	35% (18-53%)	65% (47-82%)	1.4 (0.57-2.7)	2.8% (1.3-4.9%)	3.1% (1.2-6.2%)	12 (4.7-22)	790 (390-1400)
Guinea-Bissau	2016	Female	35 to 39	38% (21-57%)	62% (43-79%)	1.5 (0.6-2.9)	3.3% (1.4-5.9%)	4% (1.5-7.4%)	15 (5.4-29)	870 (360-1600)
Guinea-Bissau	2016	Female	40 to 44	40% (24-58%)	60% (42-76%)	1.5 (0.59-3.1)	4.2% (1.7-7.6%)	5.4% (2-9.8%)	19 (7.4-36)	950 (380-1700)
Guinea-Bissau	2016	Female	45 to 49	41% (21-60%)	59% (40-79%)	1.6 (0.58-3.4)	5.1% (1.8-9.8%)	6.6% (2.2-13%)	24 (7.7-46)	1000 (340-1900)
Guinea-Bissau	2016	Female	50 to 54	41% (23-62%)	59% (38-77%)	1.7 (0.62-3.6)	5.6% (1.7-11%)	7.1% (2.2-15%)	29 (8.5-58)	1100 (320-2100)
Guinea-Bissau	2016	Female	55 to 59	41% (24-60%)	59% (40-76%)	1.8 (0.62-4.2)	6.2% (1.4-13%)	7.8% (1.8-16%)	36 (8.5-76)	1100 (270-2400)
Guinea-Bissau	2016	Female	60 to 64	42% (24-61%)	58% (39-76%)	1.8 (0.63-4.2)	5.2% (0.68-11%)	6.3% (0.84-14%)	37 (5-83)	990 (140-2200)
Guinea-Bissau	2016	Female	65 to 69	43% (25-61%)	57% (39-75%)	1.7 (0.59-3.7)	4.4% (-0.042-11%)	5.2% (-0.041-13%)	31 (-0.25-81)	700 (-7-1800)
Guinea-Bissau	2016	Female	70 to 74	41% (25-59%)	59% (41-75%)	1.4 (0.45-3.1)	3.4% (-0.57-10%)	4% (-0.68-12%)	24 (-4-76)	440 (-74-1400)
Guinea-Bissau	2016	Female	75 to 79	37% (21-55%)	63% (45-79%)	1.1 (0.32-2.6)	2.3% (-1-8.6%)	2.6% (-1.2-9.9%)	13 (-6.3-52)	190 (-86-720)
Guinea-Bissau	2016	Female	80 to 84	34% (18-54%)	66% (46-82%)	1 (0.32-2.3)	1.8% (-1-7%)	2.1% (-1.2-8.1%)	6.5 (-3.8-26)	68 (-39-270)
Guinea-Bissau	2016	Female	85 to 89	35% (19-53%)	65% (47-81%)	0.98 (0.3-2.3)	1.5% (-1.2-6.4%)	1.7% (-1.4-7.5%)	3.2 (-2.5-13)	24 (-18-100)
Guinea-Bissau	2016	Female	90 to 94	38% (21-55%)	62% (45-79%)	1 (0.29-2.3)	1.3% (-1.3-6.1%)	1.5% (-1.7-7.3%)	1 (-1.2-5.1)	5.3 (-5.5-26)
Guinea-Bissau	2016	Female	95 plus	42% (22-63%)	58% (37-78%)	1.1 (0.26-2.7)	1.2% (-1.5-6.2%)	1.3% (-2-7.2%)	0.24 (-0.38-1.3)	0.9 (-1.1-4.6)
Guinea-Bissau	2016	Male	All Ages	22% (17-29%)	78% (71-83%)	0.91 (0.5-1.5)	3.7% (2.1-5.4%)	5.5% (2.8-8.6%)	570 (280-910)	22000 (13000-34000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guinea-Bissau	2016	Male	15 to 19	17% (8.8-29%)	83% (71-91%)	0.53 (0.17-1.2)	1.8% (0.75-3.6%)	2% (0.63-4.4%)	5.1 (1.5-11)	460 (190-910)
Guinea-Bissau	2016	Male	20 to 24	25% (13-40%)	75% (60-87%)	1.1 (0.37-2.3)	4.7% (2-8.3%)	5.2% (1.8-10%)	17 (5.8-33)	1300 (580-2400)
Guinea-Bissau	2016	Male	25 to 29	30% (17-47%)	70% (53-83%)	1.5 (0.57-3.1)	6.5% (3-11%)	7.4% (3-13%)	29 (12-51)	2000 (920-3300)
Guinea-Bissau	2016	Male	30 to 34	34% (18-52%)	66% (48-82%)	1.9 (0.78-3.7)	7.2% (3.8-12%)	8.3% (4-14%)	40 (19-64)	2400 (1200-3800)
Guinea-Bissau	2016	Male	35 to 39	37% (21-56%)	63% (44-79%)	2.1 (0.83-4.2)	7.8% (3.9-12%)	9% (4.3-14%)	46 (21-72)	2400 (1200-3800)
Guinea-Bissau	2016	Male	40 to 44	38% (22-58%)	62% (42-78%)	2.2 (0.8-4.5)	8.7% (3.9-14%)	10% (4.5-16%)	54 (23-87)	2500 (1100-4100)
Guinea-Bissau	2016	Male	45 to 49	39% (21-58%)	61% (42-79%)	2.3 (0.8-5)	10% (4.7-16%)	12% (5.4-19%)	64 (28-110)	2700 (1200-4400)
Guinea-Bissau	2016	Male	50 to 54	39% (22-59%)	61% (41-78%)	2.4 (0.84-5.3)	10% (4.4-17%)	12% (5-20%)	67 (27-120)	2400 (1000-4200)
Guinea-Bissau	2016	Male	55 to 59	39% (23-59%)	61% (41-77%)	2.3 (0.76-5.1)	10% (3.9-17%)	12% (4.6-20%)	70 (26-120)	2200 (850-3800)
Guinea-Bissau	2016	Male	60 to 64	39% (22-58%)	61% (42-78%)	2.1 (0.67-4.8)	7.9% (2.6-14%)	9% (2.9-17%)	64 (20-120)	1700 (540-3200)
Guinea-Bissau	2016	Male	65 to 69	39% (23-57%)	61% (43-77%)	1.8 (0.6-4)	6.1% (1.8-13%)	7% (2-15%)	48 (13-100)	1100 (320-2200)
Guinea-Bissau	2016	Male	70 to 74	38% (23-56%)	62% (44-77%)	1.4 (0.44-3.1)	4.5% (0.97-11%)	5.2% (1.1-13%)	34 (6.8-86)	610 (130-1500)
Guinea-Bissau	2016	Male	75 to 79	35% (20-53%)	65% (47-80%)	1.1 (0.29-2.7)	3.1% (0.4-8.6%)	3.5% (0.42-9.8%)	17 (2.1-50)	240 (30-690)
Guinea-Bissau	2016	Male	80 to 84	32% (17-51%)	68% (49-83%)	0.89 (0.26-2.1)	2.5% (0.29-7%)	2.8% (0.25-8%)	7.7 (0.67-22)	80 (9.1-230)
Guinea-Bissau	2016	Male	85 to 89	31% (16-49%)	69% (51-84%)	0.76 (0.23-1.8)	2% (0.084-5.9%)	2.2% (0.026-6.8%)	3.1 (0.036-9.4)	23 (1-69)
Guinea-Bissau	2016	Male	90 to 94	33% (17-51%)	67% (49-83%)	0.72 (0.19-1.7)	1.3% (-0.15-4.7%)	1.5% (-0.31-5.4%)	0.72 (-0.15-2.6)	3.9 (-0.44-14)
Guinea-Bissau	2016	Male	95 plus	37% (19-60%)	63% (40-81%)	0.73 (0.15-2)	1.5% (-0.43-5.8%)	1.6% (-0.68-6.8%)	0.19 (-0.083-0.82)	0.72 (-0.21-2.8)
Liberia	1990	Female	All Ages	20% (17-23%)	80% (77-83%)	1.3 (1-1.5)	1.5% (1.2-1.8%)	3.5% (2.6-4.4%)	560 (420-710)	18000 (14000-23000)
Liberia	1990	Female	15 to 19	18% (12-25%)	82% (75-88%)	1.3 (0.79-1.9)	1.9% (1.2-2.9%)	2.2% (1.2-3.6%)	7.6 (4.1-13)	690 (420-1000)
Liberia	1990	Female	20 to 24	23% (15-32%)	77% (68-85%)	1.9 (1.1-2.8)	3.4% (2.3-5%)	3.9% (2.3-6%)	14 (8.2-22)	1200 (770-1700)
Liberia	1990	Female	25 to 29	27% (17-38%)	73% (62-83%)	2.2 (1.4-3.2)	4.2% (2.8-5.9%)	5.1% (3.2-7.6%)	16 (10-24)	1200 (820-1700)
Liberia	1990	Female	30 to 34	30% (19-43%)	70% (57-81%)	2.4 (1.5-3.6)	4.3% (3-6.1%)	5.6% (3.7-8.2%)	17 (11-25)	1100 (790-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	1990	Female	35 to 39	33% (21-47%)	67% (53-79%)	2.6 (1.6-3.9)	5.4% (3.8-7.5%)	7.3% (5-10%)	23 (15-32)	1300 (890-1800)
Liberia	1990	Female	40 to 44	35% (23-50%)	65% (50-77%)	2.6 (1.6-3.8)	6% (4.1-8.1%)	8.4% (5.8-12%)	24 (16-33)	1200 (840-1600)
Liberia	1990	Female	45 to 49	35% (21-51%)	65% (49-79%)	2.3 (1.3-3.6)	6.7% (4.5-9.4%)	9.3% (6.2-13%)	31 (20-45)	1400 (900-1900)
Liberia	1990	Female	50 to 54	36% (23-51%)	64% (49-77%)	2.7 (1.6-4.2)	8.9% (6-12%)	12% (8.4-17%)	47 (32-65)	1800 (1200-2400)
Liberia	1990	Female	55 to 59	36% (24-51%)	64% (49-76%)	3.7 (2-6.2)	11% (7.4-15%)	15% (9.8-20%)	65 (43-89)	2100 (1400-2800)
Liberia	1990	Female	60 to 64	37% (24-53%)	63% (47-76%)	4 (2.2-6.9)	11% (7.6-15%)	14% (9.8-19%)	75 (51-100)	2000 (1400-2800)
Liberia	1990	Female	65 to 69	39% (27-53%)	61% (47-73%)	3.7 (2.1-5.8)	12% (7.9-16%)	15% (9.8-20%)	88 (58-120)	2000 (1300-2700)
Liberia	1990	Female	70 to 74	36% (25-51%)	64% (49-75%)	2.9 (1.6-4.9)	11% (6.4-16%)	13% (7.7-19%)	78 (45-110)	1400 (830-2000)
Liberia	1990	Female	75 to 79	31% (20-45%)	69% (55-80%)	2.2 (1-3.9)	8.2% (4.1-13%)	9.9% (5-15%)	44 (23-69)	620 (320-970)
Liberia	1990	Female	80 to 84	29% (18-42%)	71% (58-82%)	1.9 (0.95-3.3)	6.9% (3.6-11%)	8.2% (4.1-13%)	19 (9.5-30)	200 (99-310)
Liberia	1990	Female	85 to 89	29% (19-42%)	71% (58-81%)	1.9 (0.94-3.2)	6.1% (2.8-10%)	7.2% (3.3-12%)	9 (4.2-15)	66 (31-110)
Liberia	1990	Female	90 to 94	32% (21-45%)	68% (55-79%)	1.9 (0.93-3.5)	5.4% (2.2-10%)	6.4% (2.5-12%)	2.7 (1.1-5.3)	14 (5.6-26)
Liberia	1990	Female	95 plus	35% (21-52%)	65% (48-79%)	2 (0.84-4)	4.8% (1.4-9.9%)	5.6% (1.6-12%)	0.66 (0.18-1.4)	2.3 (0.69-4.7)
Liberia	1990	Male	All Ages	27% (23-31%)	73% (69-77%)	1.7 (1.4-2.1)	1.9% (1.5-2.3%)	3.6% (2.8-4.5%)	670 (520-840)	27000 (22000-32000)
Liberia	1990	Male	15 to 19	20% (13-27%)	80% (73-87%)	1.1 (0.67-1.7)	1.7% (1-2.6%)	1.8% (1-2.9%)	9.7 (5.5-15)	830 (510-1200)
Liberia	1990	Male	20 to 24	34% (24-45%)	66% (55-76%)	2.7 (1.7-3.8)	3.7% (2.6-5.1%)	3.8% (2.5-5.4%)	30 (20-42)	2200 (1500-3000)
Liberia	1990	Male	25 to 29	48% (34-61%)	52% (39-66%)	4.1 (2.8-5.6)	6.6% (4.8-8.5%)	7.3% (5.2-9.7%)	44 (32-58)	2900 (2200-3800)
Liberia	1990	Male	30 to 34	52% (36-67%)	48% (33-64%)	4.6 (3.1-6.3)	8.5% (6.4-11%)	10% (7.5-13%)	48 (36-62)	2900 (2200-3700)
Liberia	1990	Male	35 to 39	48% (32-65%)	52% (35-68%)	4.2 (2.7-5.8)	9.9% (7.5-13%)	13% (9.4-16%)	50 (37-65)	2700 (2000-3400)
Liberia	1990	Male	40 to 44	47% (32-65%)	53% (35-68%)	4 (2.6-5.7)	12% (9.1-15%)	16% (12-19%)	54 (41-68)	2600 (2000-3200)
Liberia	1990	Male	45 to 49	48% (31-65%)	52% (35-69%)	4.1 (2.5-6)	14% (10-17%)	18% (14-23%)	70 (53-90)	3000 (2300-3800)
Liberia	1990	Male	50 to 54	48% (33-66%)	52% (34-67%)	3.9 (2.5-5.9)	14% (10-18%)	18% (14-24%)	73 (53-95)	2700 (2000-3500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	1990	Male	55 to 59	48% (33-65%)	52% (35-67%)	3.6 (2.1-5.6)	13% (9-17%)	17% (12-22%)	70 (48-94)	2300 (1600-3000)
Liberia	1990	Male	60 to 64	47% (31-65%)	53% (35-69%)	3.2 (1.8-5.1)	11% (7.2-15%)	14% (9.3-19%)	66 (42-92)	1800 (1100-2500)
Liberia	1990	Male	65 to 69	47% (32-64%)	53% (36-68%)	2.7 (1.6-4.3)	9.8% (6.1-14%)	12% (7.8-18%)	62 (39-91)	1400 (890-2100)
Liberia	1990	Male	70 to 74	45% (32-62%)	55% (38-68%)	2.2 (1.2-3.5)	8.1% (4.5-13%)	10% (5.5-16%)	50 (28-80)	910 (500-1400)
Liberia	1990	Male	75 to 79	42% (29-59%)	58% (41-71%)	1.7 (0.86-3)	6.1% (2.9-10%)	7.4% (3.5-13%)	28 (13-49)	400 (190-680)
Liberia	1990	Male	80 to 84	39% (25-56%)	61% (44-75%)	1.4 (0.7-2.3)	4.7% (2.1-8.1%)	5.7% (2.5-9.9%)	10 (4.6-18)	110 (48-190)
Liberia	1990	Male	85 to 89	36% (24-52%)	64% (48-76%)	1.1 (0.55-1.8)	3.3% (1.4-5.7%)	3.9% (1.6-6.9%)	4.2 (1.7-7.6)	31 (13-56)
Liberia	1990	Male	90 to 94	37% (24-53%)	63% (47-76%)	0.88 (0.43-1.5)	1.8% (0.46-3.7%)	2.1% (0.43-4.4%)	0.8 (0.16-1.7)	4.2 (1.1-8.9)
Liberia	1990	Male	95 plus	41% (24-60%)	59% (40-76%)	0.76 (0.32-1.6)	1.2% (-0.19-3.5%)	1.3% (-0.48-4.2%)	0.12 (-0.045-0.39)	0.47 (-0.076-1.4)
Liberia	1995	Female	All Ages	20% (17-24%)	80% (76-83%)	1.3 (1-1.5)	1.7% (1.3-2.1%)	3.8% (2.8-4.9%)	610 (440-780)	20000 (15000-24000)
Liberia	1995	Female	15 to 19	18% (12-25%)	82% (75-88%)	1.2 (0.79-1.8)	2.1% (1.3-3.3%)	2.7% (1.4-4.6%)	8.6 (4.4-14)	750 (450-1200)
Liberia	1995	Female	20 to 24	23% (16-32%)	77% (68-84%)	1.9 (1.2-2.7)	3.8% (2.4-5.7%)	4.5% (2.6-7%)	17 (9.6-26)	1400 (850-2000)
Liberia	1995	Female	25 to 29	27% (17-39%)	73% (61-83%)	2.2 (1.3-3.1)	4.3% (2.9-6.3%)	5.3% (3.2-8.2%)	19 (11-29)	1400 (880-2000)
Liberia	1995	Female	30 to 34	30% (19-45%)	70% (55-81%)	2.4 (1.5-3.5)	4.4% (2.9-6.3%)	5.6% (3.4-8.3%)	19 (12-29)	1200 (820-1800)
Liberia	1995	Female	35 to 39	34% (21-50%)	66% (50-79%)	2.6 (1.6-3.9)	5.4% (3.7-7.8%)	7.3% (4.7-11%)	25 (16-36)	1400 (950-2000)
Liberia	1995	Female	40 to 44	36% (23-52%)	64% (48-77%)	2.6 (1.6-3.9)	6% (4-8.5%)	8.4% (5.6-12%)	28 (18-39)	1400 (910-1900)
Liberia	1995	Female	45 to 49	36% (21-52%)	64% (48-79%)	2.4 (1.3-3.6)	6.8% (4.4-9.7%)	9.4% (6-14%)	30 (19-43)	1300 (820-1800)
Liberia	1995	Female	50 to 54	36% (23-52%)	64% (48-77%)	2.6 (1.6-4.1)	9% (6-12%)	12% (8.2-17%)	50 (33-70)	1800 (1200-2600)
Liberia	1995	Female	55 to 59	36% (24-52%)	64% (48-76%)	3.5 (1.9-6)	11% (7.3-16%)	15% (9.6-20%)	70 (45-97)	2200 (1400-3100)
Liberia	1995	Female	60 to 64	38% (24-54%)	62% (46-76%)	3.9 (2.1-6.7)	11% (7.4-15%)	14% (9.5-20%)	79 (52-110)	2100 (1400-3000)
Liberia	1995	Female	65 to 69	39% (27-55%)	61% (45-73%)	3.6 (2-5.5)	12% (7.6-17%)	15% (9.5-20%)	94 (61-130)	2100 (1400-2900)
Liberia	1995	Female	70 to 74	37% (25-52%)	63% (48-75%)	2.8 (1.5-4.7)	11% (6.2-16%)	13% (7.3-19%)	89 (49-130)	1600 (890-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	1995	Female	75 to 79	31% (20-46%)	69% (54-80%)	2.1 (1-3.6)	8.1% (4.1-13%)	9.6% (4.9-15%)	48 (24-78)	670 (340-1100)
Liberia	1995	Female	80 to 84	29% (18-42%)	71% (58-82%)	1.8 (0.93-3)	6.7% (3.3-11%)	7.9% (3.8-13%)	21 (10-34)	220 (110-350)
Liberia	1995	Female	85 to 89	29% (19-42%)	71% (58-81%)	1.8 (0.9-3)	5.8% (2.7-10%)	6.9% (3.1-12%)	12 (5.4-20)	85 (40-140)
Liberia	1995	Female	90 to 94	32% (21-46%)	68% (54-79%)	1.8 (0.88-3.3)	5.1% (1.9-9.9%)	6.1% (2.2-12%)	3.6 (1.3-7)	18 (6.8-35)
Liberia	1995	Female	95 plus	35% (21-53%)	65% (47-79%)	1.9 (0.81-3.8)	4.6% (1.3-9.9%)	5.3% (1.4-12%)	0.8 (0.21-1.7)	2.7 (0.77-6)
Liberia	1995	Male	All Ages	27% (23-32%)	73% (68-77%)	1.7 (1.4-2)	2% (1.6-2.5%)	3.8% (2.9-4.8%)	670 (510-850)	26000 (21000-33000)
Liberia	1995	Male	15 to 19	20% (13-27%)	80% (73-87%)	1.1 (0.66-1.6)	1.9% (1.1-3%)	2.1% (1.1-3.6%)	8.8 (4.4-14)	760 (450-1200)
Liberia	1995	Male	20 to 24	35% (24-45%)	65% (55-76%)	2.6 (1.7-3.7)	4.4% (2.8-6.3%)	4.8% (2.8-7.1%)	29 (17-43)	2200 (1400-3100)
Liberia	1995	Male	25 to 29	48% (34-62%)	52% (38-66%)	4 (2.8-5.4)	7.2% (4.8-9.8%)	8.4% (5.4-12%)	43 (28-59)	2900 (1900-3800)
Liberia	1995	Male	30 to 34	52% (37-67%)	48% (33-63%)	4.5 (3.1-6)	8.8% (6.4-12%)	11% (7.5-14%)	48 (33-64)	2900 (2100-3700)
Liberia	1995	Male	35 to 39	49% (33-66%)	51% (34-67%)	4 (2.6-5.7)	9.8% (7.1-13%)	13% (9-17%)	50 (36-66)	2700 (1900-3500)
Liberia	1995	Male	40 to 44	47% (33-64%)	53% (36-67%)	3.8 (2.5-5.5)	11% (8.4-15%)	15% (11-19%)	55 (39-71)	2600 (1900-3300)
Liberia	1995	Male	45 to 49	48% (31-65%)	52% (35-69%)	3.9 (2.4-5.7)	13% (9.6-17%)	18% (13-23%)	60 (44-78)	2500 (1900-3300)
Liberia	1995	Male	50 to 54	49% (33-67%)	51% (33-67%)	3.8 (2.4-5.6)	14% (9.9-18%)	18% (13-24%)	75 (53-99)	2800 (2000-3600)
Liberia	1995	Male	55 to 59	49% (33-65%)	51% (35-67%)	3.6 (2-5.5)	13% (8.8-17%)	17% (12-23%)	72 (48-99)	2300 (1600-3200)
Liberia	1995	Male	60 to 64	48% (33-65%)	52% (35-67%)	3.1 (1.8-5)	11% (6.8-16%)	14% (8.9-20%)	66 (41-93)	1800 (1100-2500)
Liberia	1995	Male	65 to 69	48% (33-65%)	52% (35-67%)	2.7 (1.6-4.1)	9.8% (6-14%)	12% (7.6-18%)	60 (37-89)	1400 (830-2000)
Liberia	1995	Male	70 to 74	46% (33-63%)	54% (37-67%)	2.2 (1.2-3.5)	8.2% (4.6-13%)	10% (5.6-16%)	52 (29-82)	940 (530-1500)
Liberia	1995	Male	75 to 79	43% (29-59%)	57% (41-71%)	1.7 (0.89-2.8)	6.1% (2.8-11%)	7.4% (3.3-13%)	30 (13-53)	420 (190-740)
Liberia	1995	Male	80 to 84	40% (25-56%)	60% (44-75%)	1.3 (0.72-2.2)	4.7% (2.1-8.4%)	5.7% (2.5-10%)	11 (5-21)	120 (52-210)
Liberia	1995	Male	85 to 89	37% (25-52%)	63% (48-75%)	1.1 (0.55-1.8)	3.3% (1.4-6.1%)	4% (1.6-7.2%)	5.6 (2.2-10)	41 (17-75)
Liberia	1995	Male	90 to 94	38% (24-53%)	62% (47-76%)	0.88 (0.45-1.5)	1.9% (0.48-3.9%)	2.2% (0.43-4.7%)	1.2 (0.24-2.6)	6.4 (1.6-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	1995	Male	95 plus	41% (25-60%)	59% (40-75%)	0.76 (0.32-1.6)	1.4% (-0.074-3.9%)	1.6% (-0.3-4.8%)	0.21 (-0.039-0.6)	0.76 (-0.039-2.1)
Liberia	2000	Female	All Ages	21% (18-24%)	79% (76-82%)	1.1 (0.91-1.3)	1.8% (1.3-2.3%)	3.9% (2.7-5.2%)	700 (490-960)	23000 (17000-29000)
Liberia	2000	Female	15 to 19	18% (12-25%)	82% (75-88%)	1.1 (0.68-1.6)	2.1% (1.3-3.5%)	2.9% (1.5-5.1%)	9.2 (4.6-17)	840 (480-1400)
Liberia	2000	Female	20 to 24	23% (15-32%)	77% (68-85%)	1.7 (1-2.4)	3.7% (2.4-5.5%)	4.4% (2.4-7.1%)	18 (10-30)	1600 (1000-2400)
Liberia	2000	Female	25 to 29	27% (17-38%)	73% (62-83%)	1.9 (1.2-2.7)	3.6% (2.3-5.3%)	4.2% (2.4-6.6%)	21 (12-32)	1600 (1000-2200)
Liberia	2000	Female	30 to 34	30% (20-44%)	70% (56-80%)	2.1 (1.3-3)	3.5% (2.3-5.1%)	4.2% (2.5-6.4%)	21 (13-32)	1400 (930-2000)
Liberia	2000	Female	35 to 39	34% (21-49%)	66% (51-79%)	2.3 (1.4-3.3)	4.3% (2.8-6.3%)	5.5% (3.4-8.5%)	28 (17-41)	1600 (1000-2300)
Liberia	2000	Female	40 to 44	35% (23-51%)	65% (49-77%)	2.2 (1.4-3.3)	4.8% (3.2-7.1%)	6.4% (4.1-9.7%)	31 (20-47)	1500 (1000-2200)
Liberia	2000	Female	45 to 49	36% (22-51%)	64% (49-78%)	2 (1.2-3)	5.5% (3.4-8.2%)	7.5% (4.6-11%)	36 (22-55)	1600 (970-2300)
Liberia	2000	Female	50 to 54	36% (23-51%)	64% (49-77%)	2.2 (1.3-3.5)	7.6% (4.9-11%)	10% (6.5-14%)	51 (31-75)	1900 (1200-2800)
Liberia	2000	Female	55 to 59	36% (24-51%)	64% (49-76%)	3 (1.6-5)	10% (6.2-14%)	13% (8-19%)	83 (50-120)	2700 (1600-3900)
Liberia	2000	Female	60 to 64	37% (24-53%)	63% (47-76%)	3.3 (1.8-5.4)	10% (6.5-14%)	13% (8.2-18%)	96 (60-140)	2600 (1600-3800)
Liberia	2000	Female	65 to 69	39% (27-53%)	61% (47-73%)	3 (1.8-4.6)	10% (6.3-15%)	13% (7.9-19%)	110 (67-160)	2500 (1500-3700)
Liberia	2000	Female	70 to 74	36% (25-51%)	64% (49-75%)	2.4 (1.3-3.9)	9.2% (4.6-14%)	11% (5.6-17%)	100 (51-160)	1800 (920-2900)
Liberia	2000	Female	75 to 79	31% (20-45%)	69% (55-80%)	1.8 (0.86-3.1)	6.6% (3-11%)	7.9% (3.5-14%)	60 (27-100)	830 (380-1400)
Liberia	2000	Female	80 to 84	28% (18-42%)	72% (58-82%)	1.5 (0.78-2.6)	5.3% (2.3-9.3%)	6.3% (2.7-11%)	24 (10-43)	250 (110-440)
Liberia	2000	Female	85 to 89	29% (18-41%)	71% (59-82%)	1.5 (0.78-2.5)	4.5% (1.8-8.1%)	5.3% (2.2-9.6%)	10 (4.3-19)	77 (32-140)
Liberia	2000	Female	90 to 94	31% (20-45%)	69% (55-80%)	1.5 (0.75-2.7)	3.9% (1.4-8.3%)	4.7% (1.6-10%)	3.8 (1.3-8.3)	19 (6.8-41)
Liberia	2000	Female	95 plus	35% (20-52%)	65% (48-80%)	1.6 (0.67-3.1)	3.5% (0.85-8.3%)	4.1% (0.84-10%)	0.87 (0.18-2.1)	3 (0.7-7)
Liberia	2000	Male	All Ages	28% (24-33%)	72% (67-76%)	1.5 (1.2-1.8)	2.3% (1.8-2.9%)	4.2% (3.1-5.5%)	800 (580-1100)	32000 (25000-40000)
Liberia	2000	Male	15 to 19	20% (14-28%)	80% (72-86%)	0.93 (0.57-1.4)	2.6% (1.6-3.9%)	3.8% (2-6.1%)	11 (5.8-17)	950 (560-1400)
Liberia	2000	Male	20 to 24	35% (24-46%)	65% (54-76%)	2.2 (1.4-3.1)	7% (4.7-9.9%)	9.3% (5.8-13%)	36 (23-53)	2700 (1800-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	2000	Male	25 to 29	48% (33-62%)	52% (38-67%)	3.3 (2.3-4.6)	9.4% (6.4-13%)	12% (7.9-17%)	56 (37-76)	3700 (2600-4900)
Liberia	2000	Male	30 to 34	53% (37-68%)	47% (32-63%)	3.8 (2.6-5.1)	9.8% (7-13%)	13% (8.8-18%)	62 (44-84)	3700 (2700-5000)
Liberia	2000	Male	35 to 39	49% (33-66%)	51% (34-67%)	3.4 (2.3-4.8)	9.7% (6.9-13%)	13% (8.8-18%)	64 (45-85)	3400 (2500-4500)
Liberia	2000	Male	40 to 44	47% (32-63%)	53% (37-68%)	3.3 (2.1-4.7)	10% (7.6-14%)	14% (10-18%)	69 (50-92)	3300 (2400-4400)
Liberia	2000	Male	45 to 49	48% (32-66%)	52% (34-68%)	3.3 (2-4.8)	12% (8.4-16%)	16% (11-21%)	77 (55-100)	3300 (2300-4400)
Liberia	2000	Male	50 to 54	49% (34-66%)	51% (34-66%)	3.2 (2-4.8)	12% (8.5-16%)	16% (11-22%)	78 (54-110)	2900 (2000-3900)
Liberia	2000	Male	55 to 59	49% (33-66%)	51% (34-67%)	3 (1.8-4.7)	12% (7.6-17%)	15% (9.9-21%)	88 (56-130)	2800 (1800-4000)
Liberia	2000	Male	60 to 64	48% (33-66%)	52% (34-67%)	2.7 (1.5-4.2)	9.6% (5.7-14%)	12% (7.1-18%)	80 (46-120)	2200 (1300-3300)
Liberia	2000	Male	65 to 69	48% (33-64%)	52% (36-67%)	2.3 (1.4-3.5)	8.1% (4.7-13%)	10% (5.9-16%)	69 (40-110)	1600 (920-2500)
Liberia	2000	Male	70 to 74	46% (33-62%)	54% (38-67%)	1.9 (1.1-3)	6.6% (3.4-11%)	8.1% (4.2-14%)	57 (29-97)	1000 (530-1700)
Liberia	2000	Male	75 to 79	44% (30-60%)	56% (40-70%)	1.5 (0.77-2.5)	4.9% (1.9-9%)	5.9% (2.3-11%)	35 (13-65)	490 (190-910)
Liberia	2000	Male	80 to 84	41% (26-57%)	59% (43-74%)	1.2 (0.62-2)	3.7% (1.5-7%)	4.4% (1.6-8.5%)	13 (4.7-25)	130 (51-260)
Liberia	2000	Male	85 to 89	38% (25-53%)	62% (47-75%)	0.93 (0.49-1.6)	2.5% (0.86-5.1%)	3% (0.91-6%)	4.8 (1.5-9.8)	36 (12-71)
Liberia	2000	Male	90 to 94	39% (25-54%)	61% (46-75%)	0.77 (0.39-1.3)	1.4% (0.12-3.2%)	1.6% (-0.0035-3.7%)	1.2 (-0.0026-2.8)	6.2 (0.54-14)
Liberia	2000	Male	95 plus	42% (26-60%)	58% (40-74%)	0.67 (0.28-1.4)	1% (-0.27-3.2%)	1.1% (-0.57-3.8%)	0.21 (-0.11-0.7)	0.81 (-0.22-2.5)
Liberia	2005	Female	All Ages	19% (16-22%)	81% (78-84%)	0.93 (0.76-1.1)	1.7% (1.3-2.3%)	3.7% (2.5-5.1%)	630 (420-880)	20000 (15000-27000)
Liberia	2005	Female	15 to 19	15% (10-21%)	85% (79-90%)	0.77 (0.48-1.2)	1.5% (0.93-2.3%)	1.9% (0.98-3.3%)	5.8 (2.9-10)	600 (370-950)
Liberia	2005	Female	20 to 24	19% (13-26%)	81% (74-87%)	1.2 (0.75-1.8)	2.7% (1.8-4%)	3.1% (1.7-5.1%)	12 (7-20)	1200 (780-1700)
Liberia	2005	Female	25 to 29	25% (16-35%)	75% (65-84%)	1.5 (0.96-2.2)	2.8% (1.8-4%)	3.1% (1.7-4.9%)	16 (9.5-25)	1300 (870-1900)
Liberia	2005	Female	30 to 34	30% (18-43%)	70% (57-82%)	1.8 (1.1-2.7)	2.6% (1.8-3.8%)	2.9% (1.8-4.6%)	17 (11-26)	1200 (850-1700)
Liberia	2005	Female	35 to 39	33% (21-48%)	67% (52-79%)	2 (1.2-2.9)	3.3% (2.1-4.8%)	4% (2.4-6.2%)	23 (15-35)	1400 (930-2000)
Liberia	2005	Female	40 to 44	35% (23-51%)	65% (49-77%)	2 (1.3-2.9)	3.7% (2.5-5.5%)	4.7% (3.1-7.2%)	27 (18-41)	1400 (930-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	2005	Female	45 to 49	35% (21-51%)	65% (49-79%)	1.8 (1-2.8)	4.3% (2.8-6.4%)	5.8% (3.6-8.6%)	33 (20-50)	1400 (910-2100)
Liberia	2005	Female	50 to 54	35% (23-51%)	65% (49-77%)	2 (1.2-3.2)	6.3% (4-9.2%)	8.4% (5.4-12%)	51 (31-80)	1900 (1200-3000)
Liberia	2005	Female	55 to 59	36% (23-51%)	64% (49-77%)	2.7 (1.4-4.5)	8.8% (5.3-13%)	11% (6.9-16%)	73 (42-110)	2300 (1400-3500)
Liberia	2005	Female	60 to 64	37% (24-53%)	63% (47-76%)	3 (1.6-5)	8.9% (5.4-13%)	11% (6.9-16%)	91 (53-140)	2500 (1500-3700)
Liberia	2005	Female	65 to 69	38% (26-53%)	62% (47-74%)	2.7 (1.5-4.2)	9.1% (5.3-14%)	11% (6.6-17%)	100 (59-160)	2300 (1300-3500)
Liberia	2005	Female	70 to 74	36% (25-50%)	64% (50-75%)	2.1 (1.1-3.4)	7.7% (3.7-13%)	9.3% (4.5-15%)	88 (42-150)	1600 (760-2700)
Liberia	2005	Female	75 to 79	31% (20-46%)	69% (54-80%)	1.6 (0.74-2.7)	5.5% (2.4-9.8%)	6.6% (2.8-12%)	53 (22-95)	740 (320-1300)
Liberia	2005	Female	80 to 84	28% (18-42%)	72% (58-82%)	1.4 (0.69-2.3)	4.3% (1.8-7.8%)	5.2% (2.2-9.4%)	24 (9.9-44)	250 (110-460)
Liberia	2005	Female	85 to 89	29% (18-42%)	71% (58-82%)	1.3 (0.69-2.2)	3.6% (1.5-6.5%)	4.2% (1.7-7.9%)	11 (4.2-20)	79 (32-150)
Liberia	2005	Female	90 to 94	31% (20-45%)	69% (55-80%)	1.3 (0.66-2.4)	3.1% (0.93-6.8%)	3.7% (1-8.2%)	3.1 (0.87-6.9)	16 (4.8-35)
Liberia	2005	Female	95 plus	34% (20-51%)	66% (49-80%)	1.4 (0.6-2.7)	2.8% (0.49-6.7%)	3.2% (0.45-8.2%)	0.82 (0.11-2.1)	2.8 (0.5-7)
Liberia	2005	Male	All Ages	28% (24-32%)	72% (68-76%)	1.3 (1.1-1.6)	2.4% (1.9-3.1%)	4.2% (3.1-5.5%)	720 (530-960)	30000 (23000-38000)
Liberia	2005	Male	15 to 19	18% (14-24%)	82% (76-86%)	0.74 (0.49-1.1)	2% (1.3-2.9%)	2.8% (1.6-4.3%)	7.6 (4.3-12)	730 (480-1100)
Liberia	2005	Male	20 to 24	35% (27-43%)	65% (57-73%)	2 (1.4-2.6)	6.1% (4.2-8.4%)	8.2% (5.3-12%)	31 (20-46)	2500 (1700-3400)
Liberia	2005	Male	25 to 29	50% (38-62%)	50% (38-62%)	3.2 (2.3-4.1)	8.4% (5.8-12%)	11% (7.4-16%)	52 (36-72)	3600 (2600-4800)
Liberia	2005	Male	30 to 34	54% (39-68%)	46% (32-61%)	3.5 (2.4-4.6)	8.3% (6.1-11%)	11% (7.5-15%)	59 (42-78)	3600 (2700-4700)
Liberia	2005	Male	35 to 39	49% (33-66%)	51% (34-67%)	3.1 (2-4.2)	8% (5.6-11%)	10% (7.1-14%)	59 (41-79)	3200 (2300-4200)
Liberia	2005	Male	40 to 44	47% (32-64%)	53% (36-68%)	2.9 (2-4.1)	8.4% (6.1-11%)	11% (7.8-15%)	64 (45-85)	3100 (2200-4100)
Liberia	2005	Male	45 to 49	48% (32-65%)	52% (35-68%)	3 (1.8-4.3)	9.8% (7-13%)	13% (9.1-18%)	74 (52-100)	3100 (2200-4200)
Liberia	2005	Male	50 to 54	49% (33-66%)	51% (34-67%)	2.9 (1.8-4.3)	10% (6.9-14%)	14% (9.2-18%)	75 (50-100)	2800 (1900-3800)
Liberia	2005	Male	55 to 59	49% (34-66%)	51% (34-66%)	2.7 (1.6-4.2)	9.7% (6.1-14%)	13% (7.9-18%)	70 (43-100)	2300 (1400-3300)
Liberia	2005	Male	60 to 64	48% (32-66%)	52% (34-68%)	2.4 (1.3-3.8)	8.1% (4.6-12%)	10% (5.8-16%)	73 (40-110)	2000 (1100-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	2005	Male	65 to 69	47% (33-63%)	53% (37-67%)	2 (1.2-3.2)	6.5% (3.9-10%)	8.2% (4.8-13%)	61 (36-96)	1400 (810-2200)
Liberia	2005	Male	70 to 74	46% (32-62%)	54% (38-68%)	1.7 (0.94-2.6)	5.2% (2.7-8.8%)	6.4% (3.2-11%)	47 (24-81)	860 (450-1500)
Liberia	2005	Male	75 to 79	43% (30-60%)	57% (40-70%)	1.3 (0.69-2.2)	3.7% (1.4-7%)	4.4% (1.6-8.5%)	27 (9.7-51)	380 (140-710)
Liberia	2005	Male	80 to 84	40% (26-57%)	60% (43-74%)	1 (0.56-1.7)	2.8% (0.99-5.5%)	3.3% (1.1-6.7%)	12 (3.9-24)	120 (45-250)
Liberia	2005	Male	85 to 89	38% (24-53%)	62% (47-76%)	0.82 (0.43-1.3)	1.8% (0.51-3.8%)	2.2% (0.52-4.5%)	4.1 (1-8.5)	31 (8.8-63)
Liberia	2005	Male	90 to 94	38% (25-54%)	62% (46-75%)	0.68 (0.34-1.2)	0.93% (-0.12-2.4%)	1% (-0.3-2.8%)	0.7 (-0.2-2)	3.9 (-0.5-10)
Liberia	2005	Male	95 plus	42% (26-61%)	58% (39-74%)	0.6 (0.25-1.2)	0.68% (-0.4-2.2%)	0.65% (-0.7-2.6%)	0.13 (-0.14-0.56)	0.57 (-0.34-1.9)
Liberia	2010	Female	All Ages	19% (17-22%)	81% (78-83%)	0.83 (0.66-1)	1.8% (1.3-2.4%)	3.7% (2.4-5.3%)	640 (410-930)	21000 (15000-28000)
Liberia	2010	Female	15 to 19	17% (13-21%)	83% (79-87%)	0.77 (0.52-1.1)	1.4% (0.88-2%)	1.7% (0.92-2.8%)	5.4 (2.9-8.9)	600 (370-900)
Liberia	2010	Female	20 to 24	21% (16-27%)	79% (73-84%)	1.2 (0.8-1.7)	2.7% (1.8-3.7%)	3% (1.7-4.8%)	11 (6.6-18)	1200 (800-1700)
Liberia	2010	Female	25 to 29	26% (18-35%)	74% (65-82%)	1.4 (0.95-2.1)	2.7% (1.9-3.9%)	3% (1.8-5%)	15 (9-23)	1300 (920-1900)
Liberia	2010	Female	30 to 34	29% (20-40%)	71% (60-80%)	1.6 (1-2.2)	2.4% (1.7-3.4%)	2.7% (1.7-4.2%)	16 (10-24)	1200 (850-1700)
Liberia	2010	Female	35 to 39	30% (21-44%)	70% (56-79%)	1.6 (1-2.3)	2.8% (1.9-4.1%)	3.4% (2.1-5.1%)	22 (14-32)	1300 (930-1900)
Liberia	2010	Female	40 to 44	32% (22-46%)	68% (54-78%)	1.6 (1-2.3)	3.2% (2.1-4.5%)	4% (2.6-6%)	26 (17-39)	1400 (900-2000)
Liberia	2010	Female	45 to 49	34% (22-50%)	66% (50-78%)	1.5 (0.94-2.3)	3.8% (2.3-5.7%)	5% (3.1-7.7%)	33 (19-50)	1400 (910-2200)
Liberia	2010	Female	50 to 54	35% (23-51%)	65% (49-77%)	1.7 (1-2.8)	5.6% (3.5-8.3%)	7.5% (4.7-11%)	54 (32-83)	2000 (1200-3100)
Liberia	2010	Female	55 to 59	36% (23-51%)	64% (49-77%)	2.4 (1.3-4)	8% (4.6-12%)	10% (6-16%)	84 (47-130)	2700 (1500-4100)
Liberia	2010	Female	60 to 64	37% (24-54%)	63% (46-76%)	2.6 (1.4-4.3)	8% (4.4-12%)	10% (5.7-15%)	89 (50-140)	2400 (1400-3700)
Liberia	2010	Female	65 to 69	38% (26-54%)	62% (46-74%)	2.4 (1.3-3.8)	7.9% (4.3-12%)	9.7% (5.3-15%)	110 (59-170)	2400 (1300-3800)
Liberia	2010	Female	70 to 74	36% (25-51%)	64% (49-75%)	1.9 (0.98-3.1)	6.4% (2.9-11%)	7.7% (3.4-13%)	90 (39-160)	1600 (700-2900)
Liberia	2010	Female	75 to 79	31% (20-46%)	69% (54-80%)	1.4 (0.67-2.5)	4.6% (1.6-8.8%)	5.4% (1.8-10%)	53 (18-100)	750 (260-1500)
Liberia	2010	Female	80 to 84	28% (18-42%)	72% (58-82%)	1.2 (0.61-2.1)	3.5% (1.3-6.6%)	4.1% (1.5-8%)	23 (8.1-44)	240 (87-450)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	2010	Female	85 to 89	29% (18-41%)	71% (59-82%)	1.2 (0.6-2)	2.9% (0.96-5.6%)	3.4% (1.1-6.6%)	12 (3.5-23)	86 (28-170)
Liberia	2010	Female	90 to 94	31% (20-45%)	69% (55-80%)	1.2 (0.57-2.1)	2.4% (0.46-5.6%)	2.8% (0.43-6.5%)	3.4 (0.52-8.1)	18 (3.2-40)
Liberia	2010	Female	95 plus	35% (20-52%)	65% (48-80%)	1.2 (0.52-2.5)	2.2% (0.059-5.8%)	2.5% (-0.094-7%)	0.7 (-0.027-2)	2.5 (0.069-6.8)
Liberia	2010	Male	All Ages	30% (26-33%)	70% (67-74%)	1.3 (1-1.5)	2.7% (2.1-3.5%)	4.5% (3.2-6.1%)	780 (560-1100)	33000 (25000-42000)
Liberia	2010	Male	15 to 19	21% (17-26%)	79% (74-83%)	0.75 (0.51-1.1)	2% (1.3-2.8%)	2.6% (1.6-4%)	8.4 (5-13)	820 (540-1200)
Liberia	2010	Male	20 to 24	39% (32-45%)	61% (55-68%)	1.9 (1.4-2.6)	6% (4.3-8%)	8% (5.4-11%)	34 (22-47)	2700 (1900-3600)
Liberia	2010	Male	25 to 29	53% (42-63%)	47% (37-58%)	3 (2.2-3.9)	8.4% (6-11%)	11% (7.9-16%)	56 (39-78)	3900 (2900-5300)
Liberia	2010	Male	30 to 34	55% (42-67%)	45% (33-58%)	3.2 (2.3-4.1)	8.4% (6.2-11%)	11% (7.9-15%)	64 (46-86)	4000 (3000-5300)
Liberia	2010	Male	35 to 39	50% (38-66%)	50% (34-62%)	2.7 (1.9-3.7)	7.8% (5.5-10%)	10% (6.9-14%)	65 (47-87)	3600 (2600-4700)
Liberia	2010	Male	40 to 44	48% (36-64%)	52% (36-64%)	2.6 (1.8-3.6)	8.2% (6-11%)	11% (7.7-14%)	71 (51-96)	3400 (2500-4600)
Liberia	2010	Male	45 to 49	50% (36-66%)	50% (34-64%)	2.7 (1.8-3.9)	9.4% (6.7-13%)	12% (8.7-17%)	83 (57-120)	3600 (2500-4900)
Liberia	2010	Male	50 to 54	49% (34-67%)	51% (33-66%)	2.6 (1.6-3.8)	9.5% (6.4-13%)	13% (8.5-18%)	85 (57-120)	3200 (2200-4500)
Liberia	2010	Male	55 to 59	49% (33-66%)	51% (34-67%)	2.4 (1.4-3.7)	8.9% (5.6-13%)	12% (7.3-17%)	81 (50-120)	2600 (1600-3900)
Liberia	2010	Male	60 to 64	48% (33-66%)	52% (34-67%)	2.1 (1.2-3.4)	7.3% (4.1-12%)	9.1% (5-15%)	70 (39-120)	1900 (1100-3100)
Liberia	2010	Male	65 to 69	48% (33-64%)	52% (36-67%)	1.8 (1.1-2.9)	5.7% (3.2-9.4%)	7% (3.9-11%)	66 (36-110)	1500 (830-2500)
Liberia	2010	Male	70 to 74	46% (32-63%)	54% (37-68%)	1.5 (0.84-2.3)	4.4% (2-8%)	5.3% (2.4-9.6%)	50 (23-92)	910 (420-1600)
Liberia	2010	Male	75 to 79	43% (30-59%)	57% (41-70%)	1.2 (0.6-2)	3% (0.95-5.9%)	3.6% (1.1-7.1%)	27 (8.5-53)	380 (120-750)
Liberia	2010	Male	80 to 84	40% (26-57%)	60% (43-74%)	0.92 (0.47-1.5)	2.2% (0.63-4.6%)	2.6% (0.68-5.6%)	11 (2.7-23)	110 (31-240)
Liberia	2010	Male	85 to 89	38% (24-53%)	62% (47-76%)	0.72 (0.38-1.2)	1.4% (0.27-3.1%)	1.7% (0.22-3.8%)	4.2 (0.55-9.4)	32 (5.8-70)
Liberia	2010	Male	90 to 94	38% (25-53%)	62% (47-75%)	0.6 (0.3-1)	0.65% (-0.32-1.9%)	0.67% (-0.51-2.2%)	0.6 (-0.46-2)	3.5 (-1.7-11)
Liberia	2010	Male	95 plus	41% (25-60%)	59% (40-75%)	0.52 (0.22-1)	0.43% (-0.54-1.8%)	0.35% (-0.85-2.1%)	0.071 (-0.17-0.43)	0.36 (-0.44-1.5)
Liberia	2016	Female	All Ages	20% (18-22%)	80% (78-82%)	0.87 (0.52-1.3)	2.2% (1.3-3.2%)	4.2% (2.2-6.8%)	680 (350-1100)	22000 (13000-32000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	2016	Female	15 to 19	20% (16-23%)	80% (77-84%)	0.93 (0.54-1.5)	1.5% (0.87-2.4%)	1.8% (0.85-3.4%)	5.9 (2.7-11)	700 (400-1100)
Liberia	2016	Female	20 to 24	25% (21-29%)	75% (71-79%)	1.4 (0.8-2.2)	2.8% (1.8-4.1%)	3.2% (1.6-5.4%)	11 (5.1-18)	1200 (760-1800)
Liberia	2016	Female	25 to 29	26% (21-33%)	74% (67-79%)	1.5 (0.83-2.3)	2.8% (1.8-4.1%)	3.2% (1.7-5.5%)	13 (6.4-22)	1300 (800-1900)
Liberia	2016	Female	30 to 34	27% (21-36%)	73% (64-79%)	1.5 (0.81-2.3)	2.6% (1.6-3.7%)	3% (1.6-4.7%)	15 (8.3-24)	1200 (800-1800)
Liberia	2016	Female	35 to 39	28% (21-39%)	72% (61-79%)	1.5 (0.8-2.4)	2.9% (1.8-4.3%)	3.6% (2-5.8%)	21 (12-33)	1300 (840-2000)
Liberia	2016	Female	40 to 44	30% (22-43%)	70% (57-78%)	1.5 (0.8-2.4)	3.2% (1.9-4.8%)	4.3% (2.4-6.5%)	27 (15-41)	1400 (860-2100)
Liberia	2016	Female	45 to 49	33% (24-46%)	67% (54-76%)	1.5 (0.77-2.4)	3.9% (2.1-6.2%)	5.3% (2.8-8.6%)	36 (18-58)	1600 (860-2500)
Liberia	2016	Female	50 to 54	35% (23-50%)	65% (50-77%)	1.8 (0.89-3)	5.6% (2.9-9%)	7.7% (4-12%)	59 (30-96)	2200 (1200-3600)
Liberia	2016	Female	55 to 59	36% (23-51%)	64% (49-77%)	2.4 (1.1-4.3)	7.7% (3.9-12%)	10% (5.1-16%)	90 (45-140)	2900 (1500-4600)
Liberia	2016	Female	60 to 64	37% (24-53%)	63% (47-76%)	2.6 (1.3-4.6)	9.7% (3.7-12%)	9.7% (4.7-15%)	100 (49-160)	2800 (1400-4300)
Liberia	2016	Female	65 to 69	39% (26-53%)	61% (47-74%)	2.4 (1.2-4.2)	7.6% (3.3-13%)	9.4% (4.1-15%)	100 (45-180)	2400 (1000-4000)
Liberia	2016	Female	70 to 74	36% (24-50%)	64% (50-76%)	1.9 (0.87-3.5)	6.2% (2.1-12%)	7.4% (2.5-14%)	92 (31-180)	1700 (570-3300)
Liberia	2016	Female	75 to 79	31% (20-45%)	69% (55-80%)	1.4 (0.6-2.8)	4.4% (1.1-9.6%)	5.2% (1.2-11%)	60 (14-130)	840 (200-1800)
Liberia	2016	Female	80 to 84	28% (18-42%)	72% (58-82%)	1.2 (0.53-2.3)	3.5% (0.72-7.5%)	4.1% (0.83-8.8%)	28 (5.5-60)	290 (60-620)
Liberia	2016	Female	85 to 89	29% (18-41%)	71% (59-82%)	1.2 (0.52-2.3)	2.9% (0.45-6.7%)	3.4% (0.45-7.9%)	13 (1.7-31)	98 (15-220)
Liberia	2016	Female	90 to 94	31% (20-45%)	69% (55-80%)	1.2 (0.5-2.4)	2.5% (0.021-6.9%)	2.9% (-0.09-8.1%)	4.4 (-0.14-12)	23 (0.19-62)
Liberia	2016	Female	95 plus	35% (20-52%)	65% (48-80%)	1.3 (0.49-2.7)	2.3% (-0.22-6.8%)	2.6% (-0.44-8%)	1.1 (-0.19-3.4)	3.8 (-0.38-11)
Liberia	2016	Male	All Ages	31% (28-35%)	69% (65-72%)	1.3 (0.82-2)	3.2% (2.1-4.6%)	5.1% (3.1-7.5%)	830 (490-1200)	35000 (23000-50000)
Liberia	2016	Male	15 to 19	24% (19-29%)	76% (71-81%)	0.89 (0.51-1.4)	2.2% (1.3-3.5%)	2.8% (1.4-4.8%)	11 (5.6-19)	1100 (630-1700)
Liberia	2016	Male	20 to 24	43% (36-49%)	57% (51-64%)	2.2 (1.2-3.3)	6.2% (4-9%)	8.2% (4.9-12%)	36 (21-56)	2900 (1900-4300)
Liberia	2016	Male	25 to 29	54% (44-64%)	46% (36-56%)	3 (1.7-4.8)	8.4% (5.5-12%)	12% (7.1-17%)	55 (34-81)	4000 (2600-5600)
Liberia	2016	Male	30 to 34	56% (44-68%)	44% (32-56%)	3.2 (1.9-5)	8.6% (5.8-12%)	12% (7.7-17%)	66 (42-93)	4200 (2800-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Liberia	2016	Male	35 to 39	50% (38-64%)	50% (36-62%)	2.8 (1.5-4.4)	8.2% (5.4-11%)	11% (6.9-16%)	70 (44-99)	3900 (2600-5400)
Liberia	2016	Male	40 to 44	48% (37-63%)	52% (37-63%)	2.7 (1.5-4.3)	8.5% (5.4-12%)	11% (7.2-17%)	77 (47-110)	3800 (2400-5600)
Liberia	2016	Male	45 to 49	51% (38-65%)	49% (35-62%)	2.8 (1.5-4.6)	9.7% (6-14%)	13% (7.9-19%)	93 (56-140)	4000 (2500-5800)
Liberia	2016	Male	50 to 54	50% (35-66%)	50% (34-65%)	2.7 (1.4-4.5)	9.3% (5.6-14%)	12% (7.5-18%)	93 (54-140)	3500 (2100-5200)
Liberia	2016	Male	55 to 59	49% (33-66%)	51% (34-67%)	2.5 (1.2-4.4)	8.6% (4.7-14%)	11% (6.1-17%)	91 (49-140)	3000 (1600-4600)
Liberia	2016	Male	60 to 64	48% (32-67%)	52% (33-68%)	2.2 (1-3.8)	7% (3.4-12%)	8.8% (4.2-15%)	79 (38-130)	2200 (1100-3700)
Liberia	2016	Male	65 to 69	48% (34-65%)	52% (35-66%)	1.9 (0.94-3.1)	5.4% (2.5-9.6%)	6.7% (3-12%)	62 (27-110)	1400 (640-2400)
Liberia	2016	Male	70 to 74	46% (32-62%)	54% (38-68%)	1.5 (0.73-2.7)	4.1% (1.6-8.2%)	5% (1.9-10%)	51 (20-100)	940 (370-1900)
Liberia	2016	Male	75 to 79	43% (29-60%)	57% (40-71%)	1.2 (0.54-2.3)	2.7% (0.71-6.1%)	3.3% (0.82-7.3%)	28 (6.9-66)	410 (100-940)
Liberia	2016	Male	80 to 84	40% (26-57%)	60% (43-74%)	0.94 (0.42-1.8)	2% (0.42-4.6%)	2.4% (0.41-5.5%)	11 (2-27)	120 (25-280)
Liberia	2016	Male	85 to 89	38% (24-53%)	62% (47-76%)	0.74 (0.33-1.4)	1.3% (0.13-3%)	1.4% (0.044-3.6%)	4 (0.12-10)	32 (3.2-76)
Liberia	2016	Male	90 to 94	38% (25-55%)	62% (45-75%)	0.61 (0.25-1.2)	0.55% (-0.43-1.9%)	0.53% (-0.68-2.1%)	0.6 (-0.77-2.4)	3.8 (-2.9-13)
Liberia	2016	Male	95 plus	42% (25-61%)	58% (39-75%)	0.53 (0.2-1.2)	0.33% (-0.67-1.6%)	0.19% (-1.1-1.7%)	0.055 (-0.3-0.49)	0.38 (-0.78-1.8)
Mali	1990	Female	All Ages	5.3% (4.4-6.2%)	95% (94-96%)	0.35 (0.26-0.46)	0.7% (0.5-0.93%)	1.6% (1-2.2%)	1200 (780-1700)	37000 (26000-50000)
Mali	1990	Female	15 to 19	5% (3.2-7.3%)	95% (93-97%)	0.32 (0.15-0.55)	0.72% (0.45-1.1%)	0.62% (0.25-1.2%)	7.1 (2.9-13)	910 (540-1400)
Mali	1990	Female	20 to 24	6.3% (3.8-9.4%)	94% (91-96%)	0.51 (0.26-0.88)	1.5% (0.96-2.1%)	1.2% (0.59-2.1%)	15 (7.7-27)	1800 (1100-2700)
Mali	1990	Female	25 to 29	7% (4.2-10%)	93% (90-96%)	0.62 (0.32-1)	1.9% (1.2-2.8%)	1.8% (1-3%)	22 (12-36)	2000 (1300-2900)
Mali	1990	Female	30 to 34	7.7% (4.7-12%)	92% (88-95%)	0.71 (0.38-1.2)	2.2% (1.5-3.2%)	2.5% (1.5-3.8%)	29 (17-45)	2100 (1400-3100)
Mali	1990	Female	35 to 39	8.9% (5.1-13%)	91% (87-95%)	0.8 (0.42-1.4)	2.8% (1.8-4%)	3.4% (2.1-5.1%)	44 (27-67)	2600 (1700-3700)
Mali	1990	Female	40 to 44	9.4% (5.8-14%)	91% (86-94%)	0.82 (0.41-1.4)	3.1% (2-4.4%)	3.9% (2.5-5.8%)	50 (32-75)	2500 (1700-3700)
Mali	1990	Female	45 to 49	9.2% (5.6-13%)	91% (87-94%)	0.76 (0.38-1.3)	3.6% (2.2-5.5%)	4.7% (2.7-7.2%)	73 (43-110)	3200 (1900-4800)
Mali	1990	Female	50 to 54	9.2% (5.9-13%)	91% (87-94%)	0.79 (0.4-1.4)	3.9% (2.3-5.8%)	4.8% (2.9-7.2%)	88 (52-130)	3300 (2000-5000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	1990	Female	55 to 59	9.6% (6.1-14%)	90% (86-94%)	0.91 (0.46-1.6)	4.3% (2.6-6.3%)	5.3% (3.1-7.8%)	120 (70-180)	3800 (2300-5600)
Mali	1990	Female	60 to 64	11% (6.4-16%)	89% (84-94%)	0.99 (0.48-1.8)	4.3% (2.5-6.5%)	5.2% (3-7.8%)	150 (86-230)	4200 (2400-6300)
Mali	1990	Female	65 to 69	12% (7.6-17%)	88% (83-92%)	1 (0.48-1.7)	4.6% (2.5-7.3%)	5.4% (2.9-8.6%)	200 (110-320)	4600 (2400-7200)
Mali	1990	Female	70 to 74	11% (7.1-16%)	89% (84-93%)	0.81 (0.39-1.4)	4.4% (2.1-7.4%)	5.1% (2.4-8.5%)	210 (100-360)	3800 (1800-6500)
Mali	1990	Female	75 to 79	8.5% (5.1-13%)	91% (87-95%)	0.59 (0.26-1.1)	3% (1.2-5.3%)	3.5% (1.4-6.1%)	110 (48-200)	1600 (670-2800)
Mali	1990	Female	80 to 84	7.6% (4.5-11%)	92% (89-95%)	0.51 (0.23-0.89)	2.2% (0.96-3.8%)	2.5% (1.1-4.4%)	37 (16-67)	390 (170-680)
Mali	1990	Female	85 to 89	7.9% (4.6-12%)	92% (88-95%)	0.51 (0.25-0.94)	1.7% (0.7-3.2%)	1.9% (0.78-3.7%)	13 (5.3-25)	97 (41-180)
Mali	1990	Female	90 to 94	8.7% (5.5-13%)	91% (87-95%)	0.55 (0.25-1)	1.3% (0.48-2.8%)	1.5% (0.53-3.3%)	2.8 (0.97-6.1)	15 (5.2-31)
Mali	1990	Female	95 plus	10% (5.8-16%)	90% (84-94%)	0.59 (0.23-1.2)	1.2% (0.26-2.9%)	1.3% (0.23-3.4%)	0.52 (0.089-1.3)	1.9 (0.4-4.5)
Mali	1990	Male	All Ages	2.4% (2-2.9%)	98% (97-98%)	0.16 (0.12-0.21)	0.52% (0.4-0.69%)	1% (0.74-1.5%)	770 (550-1100)	29000 (22000-39000)
Mali	1990	Male	15 to 19	1.9% (1.1-3%)	98% (97-99%)	0.1 (0.045-0.18)	0.58% (0.38-0.81%)	0.35% (0.19-0.63%)	3.5 (1.9-6.4)	650 (420-940)
Mali	1990	Male	20 to 24	3% (1.7-4.5%)	97% (95-98%)	0.21 (0.11-0.38)	1.5% (1-2.1%)	1.1% (0.64-1.9%)	13 (7.5-23)	1600 (1100-2400)
Mali	1990	Male	25 to 29	4% (2.3-6.2%)	96% (94-98%)	0.34 (0.17-0.56)	2.5% (1.7-3.5%)	2.4% (1.4-3.8%)	25 (14-40)	2200 (1500-3200)
Mali	1990	Male	30 to 34	4.5% (2.5-7.2%)	96% (93-97%)	0.41 (0.21-0.68)	3.2% (2.2-4.6%)	3.5% (2.2-5.5%)	36 (22-55)	2500 (1700-3600)
Mali	1990	Male	35 to 39	4.7% (2.6-7.1%)	95% (93-97%)	0.44 (0.22-0.77)	3.9% (2.6-5.6%)	4.7% (3-7.1%)	52 (33-79)	3000 (2100-4400)
Mali	1990	Male	40 to 44	4.9% (2.9-7.5%)	95% (93-97%)	0.45 (0.22-0.79)	4.3% (2.9-6.5%)	5.5% (3.5-8.4%)	59 (38-88)	2900 (2000-4300)
Mali	1990	Male	45 to 49	5.1% (2.8-7.8%)	95% (92-97%)	0.46 (0.21-0.81)	4.7% (3.1-6.7%)	6% (3.9-8.8%)	77 (50-120)	3300 (2200-4900)
Mali	1990	Male	50 to 54	5.2% (3.1-7.9%)	95% (92-97%)	0.47 (0.22-0.85)	4.3% (2.8-6.5%)	5.4% (3.5-8.3%)	74 (47-110)	2800 (1900-4200)
Mali	1990	Male	55 to 59	5.1% (3-7.6%)	95% (92-97%)	0.44 (0.22-0.79)	4% (2.6-6%)	5% (3.2-7.4%)	82 (51-120)	2700 (1800-4000)
Mali	1990	Male	60 to 64	4.9% (2.9-7.8%)	95% (92-97%)	0.4 (0.18-0.75)	3.5% (2.2-5.3%)	4.2% (2.7-6.5%)	91 (56-140)	2500 (1600-3900)
Mali	1990	Male	65 to 69	4.9% (2.9-7.4%)	95% (93-97%)	0.36 (0.17-0.62)	2.9% (1.8-4.5%)	3.5% (2.2-5.5%)	94 (58-150)	2100 (1300-3300)
Mali	1990	Male	70 to 74	4.8% (2.9-7.1%)	95% (93-97%)	0.3 (0.14-0.54)	2.5% (1.4-4%)	2.9% (1.7-4.8%)	89 (51-150)	1600 (930-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	1990	Male	75 to 79	4.3% (2.5-6.9%)	96% (93-98%)	0.24 (0.1-0.46)	1.9% (1-3.2%)	2.2% (1.2-3.8%)	53 (29-93)	740 (410-1300)
Mali	1990	Male	80 to 84	3.8% (2.1-5.6%)	96% (94-98%)	0.18 (0.083-0.33)	1.5% (0.82-2.5%)	1.7% (0.95-2.9%)	18 (10-31)	190 (110-320)
Mali	1990	Male	85 to 89	3.4% (1.9-5.5%)	97% (94-98%)	0.14 (0.068-0.28)	1% (0.59-1.9%)	1.2% (0.65-2.2%)	5.8 (3.1-11)	43 (24-78)
Mali	1990	Male	90 to 94	3.5% (2-5.7%)	97% (94-98%)	0.13 (0.054-0.26)	0.59% (0.31-1.1%)	0.65% (0.31-1.3%)	0.93 (0.44-1.9)	5.1 (2.6-9.7)
Mali	1990	Male	95 plus	4% (2.2-6.6%)	96% (93-98%)	0.12 (0.043-0.25)	0.45% (0.19-1%)	0.45% (0.14-1.1%)	0.12 (0.038-0.31)	0.51 (0.21-1.1)
Mali	1995	Female	All Ages	5.2% (4.3-6.1%)	95% (94-96%)	0.34 (0.25-0.46)	0.67% (0.47-0.89%)	1.5% (0.9-2.1%)	1100 (710-1600)	36000 (25000-49000)
Mali	1995	Female	15 to 19	4.9% (3.1-7.1%)	95% (93-97%)	0.31 (0.15-0.52)	0.72% (0.44-1.1%)	0.6% (0.26-1.1%)	7.4 (3.2-14)	1000 (600-1500)
Mali	1995	Female	20 to 24	6.3% (3.7-9.4%)	94% (91-96%)	0.5 (0.26-0.85)	1.4% (0.94-2.1%)	1.1% (0.56-1.9%)	17 (8.3-28)	2100 (1300-3000)
Mali	1995	Female	25 to 29	6.9% (4.2-11%)	93% (89-96%)	0.6 (0.31-1)	1.8% (1.2-2.5%)	1.6% (0.91-2.7%)	22 (13-37)	2100 (1400-3100)
Mali	1995	Female	30 to 34	7.7% (4.5-12%)	92% (88-96%)	0.7 (0.37-1.1)	2.1% (1.4-3%)	2.2% (1.3-3.4%)	29 (17-45)	2100 (1400-3100)
Mali	1995	Female	35 to 39	8.9% (5.3-13%)	91% (87-95%)	0.79 (0.41-1.3)	2.6% (1.7-3.8%)	3.1% (1.9-4.7%)	41 (25-64)	2500 (1600-3600)
Mali	1995	Female	40 to 44	9.3% (5.8-13%)	91% (87-94%)	0.81 (0.41-1.4)	2.8% (1.9-4.1%)	3.6% (2.3-5.2%)	52 (32-77)	2600 (1700-3800)
Mali	1995	Female	45 to 49	9.1% (5.3-13%)	91% (87-95%)	0.76 (0.36-1.3)	3.3% (2-5.1%)	4.3% (2.4-6.6%)	65 (36-100)	2800 (1600-4400)
Mali	1995	Female	50 to 54	9.1% (5.7-14%)	91% (86-94%)	0.78 (0.38-1.4)	3.6% (2.2-5.5%)	4.5% (2.7-6.9%)	91 (50-140)	3400 (1900-5300)
Mali	1995	Female	55 to 59	9.5% (5.9-14%)	90% (86-94%)	0.89 (0.44-1.6)	4% (2.4-5.9%)	4.9% (2.9-7.3%)	110 (64-170)	3600 (2100-5500)
Mali	1995	Female	60 to 64	10% (6.5-16%)	90% (84-94%)	0.98 (0.45-1.8)	4% (2.3-6.1%)	4.8% (2.7-7.2%)	140 (77-210)	3800 (2100-5800)
Mali	1995	Female	65 to 69	12% (7.6-17%)	88% (83-92%)	0.98 (0.48-1.7)	4.4% (2.2-7%)	5.1% (2.6-8.1%)	190 (93-310)	4300 (2100-6900)
Mali	1995	Female	70 to 74	11% (6.9-16%)	89% (84-93%)	0.8 (0.38-1.4)	4.1% (1.9-7%)	4.7% (2.2-8%)	200 (91-340)	3600 (1700-6100)
Mali	1995	Female	75 to 79	8.4% (5-13%)	92% (87-95%)	0.57 (0.26-1)	2.7% (1.2-4.9%)	3.2% (1.3-5.5%)	110 (49-210)	1600 (690-2800)
Mali	1995	Female	80 to 84	7.4% (4.5-11%)	93% (89-96%)	0.49 (0.23-0.86)	2% (0.85-3.6%)	2.3% (0.96-4.1%)	41 (17-75)	430 (190-780)
Mali	1995	Female	85 to 89	7.6% (4.5-12%)	92% (88-95%)	0.48 (0.23-0.92)	1.5% (0.6-2.9%)	1.8% (0.67-3.4%)	17 (6.3-32)	130 (49-240)
Mali	1995	Female	90 to 94	8.5% (5.4-13%)	92% (87-95%)	0.52 (0.24-1)	1.2% (0.38-2.7%)	1.4% (0.41-3.2%)	4.1 (1.2-9.1)	22 (6.6-47)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	1995	Female	95 plus	9.7% (5.6-15%)	90% (85-94%)	0.56 (0.21-1.1)	1.1% (0.24-2.8%)	1.2% (0.18-3.2%)	0.76 (0.11-2)	2.7 (0.58-6.8)
Mali	1995	Male	All Ages	2.5% (2-3%)	98% (97-98%)	0.17 (0.12-0.23)	0.5% (0.38-0.67%)	0.97% (0.69-1.4%)	750 (530-1100)	29000 (22000-39000)
Mali	1995	Male	15 to 19	2% (1.1-3.1%)	98% (97-99%)	0.1 (0.048-0.19)	0.61% (0.4-0.86%)	0.36% (0.19-0.64%)	3.7 (1.9-6.6)	740 (480-1100)
Mali	1995	Male	20 to 24	3.1% (1.7-4.9%)	97% (95-98%)	0.23 (0.11-0.38)	1.5% (1.1-2.2%)	1.1% (0.63-1.9%)	15 (8.1-26)	1900 (1300-2800)
Mali	1995	Male	25 to 29	4.1% (2.3-6.4%)	96% (94-98%)	0.35 (0.18-0.61)	2.3% (1.5-3.3%)	2.1% (1.2-3.5%)	28 (16-45)	2500 (1700-3600)
Mali	1995	Male	30 to 34	4.7% (2.6-7.7%)	95% (92-97%)	0.43 (0.22-0.75)	2.8% (1.9-4.1%)	3% (1.8-4.8%)	36 (22-58)	2600 (1800-3800)
Mali	1995	Male	35 to 39	4.9% (2.7-7.5%)	95% (92-97%)	0.46 (0.23-0.81)	3.4% (2.3-4.9%)	4% (2.5-6.3%)	48 (30-72)	2800 (1900-4100)
Mali	1995	Male	40 to 44	5.1% (3-7.7%)	95% (92-97%)	0.47 (0.24-0.85)	3.8% (2.6-5.8%)	4.8% (3.1-7.5%)	62 (39-96)	3100 (2000-4700)
Mali	1995	Male	45 to 49	5.3% (2.9-8.2%)	95% (92-97%)	0.49 (0.22-0.89)	4.2% (2.8-6.2%)	5.3% (3.4-8.1%)	67 (42-100)	2900 (1900-4300)
Mali	1995	Male	50 to 54	5.4% (3.3-8.1%)	95% (92-97%)	0.49 (0.23-0.91)	4% (2.6-6.1%)	5% (3.1-7.9%)	75 (47-120)	2900 (1900-4500)
Mali	1995	Male	55 to 59	5.3% (3.2-7.9%)	95% (92-97%)	0.47 (0.23-0.85)	3.8% (2.5-5.6%)	4.7% (3.1-7.2%)	75 (48-120)	2500 (1600-3700)
Mali	1995	Male	60 to 64	5.1% (3-8.2%)	95% (92-97%)	0.43 (0.19-0.82)	3.3% (2.1-5%)	4% (2.5-6.2%)	82 (50-130)	2300 (1400-3500)
Mali	1995	Male	65 to 69	5.1% (3.1-7.6%)	95% (92-97%)	0.38 (0.17-0.65)	2.8% (1.7-4.4%)	3.4% (2-5.5%)	84 (52-140)	1900 (1200-3100)
Mali	1995	Male	70 to 74	4.9% (3.1-7.3%)	95% (93-97%)	0.31 (0.15-0.55)	2.3% (1.4-3.7%)	2.8% (1.6-4.6%)	82 (48-140)	1500 (890-2400)
Mali	1995	Male	75 to 79	4.5% (2.6-7.2%)	96% (93-97%)	0.25 (0.11-0.46)	1.8% (1-3.1%)	2.1% (1.2-3.7%)	58 (32-100)	820 (460-1400)
Mali	1995	Male	80 to 84	3.9% (2.2-5.9%)	96% (94-98%)	0.19 (0.086-0.35)	1.4% (0.81-2.5%)	1.7% (0.92-3%)	24 (13-43)	250 (140-440)
Mali	1995	Male	85 to 89	3.6% (1.9-5.6%)	96% (94-98%)	0.15 (0.072-0.3)	1% (0.6-1.9%)	1.2% (0.67-2.3%)	8.3 (4.6-16)	62 (36-120)
Mali	1995	Male	90 to 94	3.7% (2.2-5.8%)	96% (94-98%)	0.13 (0.06-0.27)	0.6% (0.31-1.2%)	0.65% (0.31-1.4%)	1.4 (0.66-3)	7.7 (3.9-16)
Mali	1995	Male	95 plus	4.2% (2.3-6.9%)	96% (93-98%)	0.13 (0.045-0.27)	0.47% (0.2-1%)	0.46% (0.14-1.2%)	0.19 (0.057-0.49)	0.79 (0.34-1.8)
Mali	2000	Female	All Ages	4.7% (3.9-5.6%)	95% (94-96%)	0.33 (0.24-0.44)	0.62% (0.45-0.81%)	1.3% (0.85-1.9%)	1100 (670-1500)	35000 (25000-46000)
Mali	2000	Female	15 to 19	4.6% (2.9-6.9%)	95% (93-97%)	0.31 (0.15-0.51)	0.7% (0.44-1%)	0.55% (0.25-0.96%)	7.6 (3.3-13)	1100 (660-1600)
Mali	2000	Female	20 to 24	5.5% (3.3-8.2%)	94% (92-97%)	0.47 (0.25-0.8)	1.3% (0.9-1.9%)	0.92% (0.49-1.5%)	15 (8.1-25)	2100 (1400-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	2000	Female	25 to 29	5.9% (3.7-8.8%)	94% (91-96%)	0.56 (0.31-0.91)	1.6% (1.1-2.2%)	1.3% (0.76-2.1%)	22 (13-35)	2300 (1600-3200)
Mali	2000	Female	30 to 34	6.7% (4-10%)	93% (90-96%)	0.65 (0.35-1)	1.8% (1.2-2.5%)	1.8% (1.1-2.7%)	28 (17-42)	2200 (1500-3100)
Mali	2000	Female	35 to 39	7.8% (4.8-12%)	92% (88-95%)	0.75 (0.39-1.2)	2.2% (1.5-3.2%)	2.6% (1.6-3.8%)	38 (23-57)	2400 (1600-3400)
Mali	2000	Female	40 to 44	8.1% (5.2-12%)	92% (88-95%)	0.76 (0.37-1.3)	2.5% (1.7-3.5%)	3.1% (2-4.5%)	46 (29-67)	2400 (1600-3400)
Mali	2000	Female	45 to 49	7.9% (4.7-11%)	92% (89-95%)	0.7 (0.34-1.2)	2.8% (1.8-4.2%)	3.6% (2.2-5.5%)	59 (36-93)	2600 (1700-4000)
Mali	2000	Female	50 to 54	8% (5.3-12%)	92% (88-95%)	0.73 (0.38-1.3)	3.2% (2-4.7%)	4% (2.4-6%)	74 (44-110)	2800 (1700-4300)
Mali	2000	Female	55 to 59	8.8% (5.6-12%)	91% (88-94%)	0.88 (0.46-1.5)	3.6% (2.3-5.4%)	4.5% (2.8-6.7%)	110 (64-160)	3500 (2100-5300)
Mali	2000	Female	60 to 64	11% (6.9-15%)	89% (85-93%)	1.1 (0.54-1.9)	3.9% (2.3-5.9%)	4.7% (2.8-7.1%)	130 (78-210)	3600 (2100-5600)
Mali	2000	Female	65 to 69	12% (8.4-17%)	88% (83-92%)	1.1 (0.58-1.9)	4.5% (2.4-6.9%)	5.4% (2.8-8.3%)	180 (95-280)	4100 (2200-6300)
Mali	2000	Female	70 to 74	11% (7.5-16%)	89% (84-93%)	0.89 (0.45-1.5)	4% (1.9-6.9%)	4.7% (2.2-8%)	180 (87-310)	3300 (1600-5600)
Mali	2000	Female	75 to 79	8.4% (5.1-13%)	92% (87-95%)	0.61 (0.28-1.1)	2.7% (1.2-4.7%)	3.1% (1.4-5.4%)	110 (46-190)	1500 (660-2700)
Mali	2000	Female	80 to 84	7.3% (4.3-11%)	93% (89-96%)	0.51 (0.24-0.91)	2% (0.87-3.7%)	2.3% (0.97-4.2%)	43 (18-81)	450 (200-830)
Mali	2000	Female	85 to 89	7.6% (4.3-12%)	92% (88-96%)	0.51 (0.25-0.97)	1.6% (0.66-3%)	1.8% (0.72-3.5%)	21 (8.5-42)	160 (66-310)
Mali	2000	Female	90 to 94	8.4% (5.2-13%)	92% (87-95%)	0.55 (0.26-1)	1.3% (0.42-2.9%)	1.5% (0.45-3.3%)	6.3 (1.9-14)	33 (11-72)
Mali	2000	Female	95 plus	9.6% (5.3-15%)	90% (85-95%)	0.6 (0.23-1.2)	1.2% (0.29-3%)	1.4% (0.25-3.4%)	1.3 (0.24-3.4)	4.8 (1.1-12)
Mali	2000	Male	All Ages	2.4% (1.9-2.9%)	98% (97-98%)	0.17 (0.12-0.23)	0.46% (0.35-0.63%)	0.89% (0.63-1.3%)	720 (510-1000)	28000 (22000-38000)
Mali	2000	Male	15 to 19	2% (1.2-3%)	98% (97-99%)	0.11 (0.054-0.2)	0.58% (0.38-0.81%)	0.31% (0.17-0.54%)	3.7 (2-6.5)	810 (510-1100)
Mali	2000	Male	20 to 24	3% (1.7-4.5%)	97% (95-98%)	0.23 (0.12-0.39)	1.3% (0.9-1.9%)	0.85% (0.47-1.4%)	14 (8.1-23)	2000 (1400-2900)
Mali	2000	Male	25 to 29	3.6% (2.2-5.4%)	96% (95-98%)	0.33 (0.18-0.54)	1.8% (1.3-2.5%)	1.6% (0.91-2.5%)	27 (16-42)	2600 (1900-3700)
Mali	2000	Male	30 to 34	3.9% (2.3-6.3%)	96% (94-98%)	0.39 (0.2-0.67)	2.1% (1.5-3.1%)	2.1% (1.3-3.4%)	34 (21-54)	2700 (1900-3700)
Mali	2000	Male	35 to 39	4% (2.3-6.2%)	96% (94-98%)	0.41 (0.2-0.69)	2.6% (1.8-3.8%)	2.9% (1.8-4.5%)	43 (27-67)	2600 (1800-3800)
Mali	2000	Male	40 to 44	4.4% (2.7-6.6%)	96% (93-97%)	0.45 (0.23-0.78)	3.1% (2.1-4.6%)	3.8% (2.5-5.8%)	53 (35-81)	2700 (1900-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	2000	Male	45 to 49	5.3% (3.1-7.7%)	95% (92-97%)	0.53 (0.26-0.94)	3.7% (2.5-5.5%)	4.7% (3-7%)	69 (45-100)	3100 (2100-4500)
Mali	2000	Male	50 to 54	5.6% (3.6-8.1%)	94% (92-96%)	0.56 (0.27-0.97)	3.7% (2.5-5.5%)	4.7% (3.1-7.1%)	64 (42-98)	2500 (1700-3700)
Mali	2000	Male	55 to 59	5.7% (3.5-8.1%)	94% (92-96%)	0.55 (0.28-0.97)	3.7% (2.5-5.4%)	4.7% (3.1-7%)	76 (50-110)	2500 (1700-3700)
Mali	2000	Male	60 to 64	5.7% (3.7-8.3%)	94% (92-96%)	0.52 (0.26-0.96)	3.3% (2.2-4.9%)	4.1% (2.7-6.3%)	75 (48-120)	2100 (1400-3200)
Mali	2000	Male	65 to 69	5.9% (3.7-8.4%)	94% (92-96%)	0.47 (0.24-0.79)	2.8% (1.8-4.4%)	3.5% (2.2-5.5%)	77 (47-120)	1800 (1100-2800)
Mali	2000	Male	70 to 74	6.1% (4-8.8%)	94% (91-96%)	0.42 (0.21-0.72)	2.5% (1.5-4%)	3% (1.8-5%)	80 (47-130)	1500 (860-2400)
Mali	2000	Male	75 to 79	5.6% (3.4-8.8%)	94% (91-97%)	0.34 (0.16-0.61)	2% (1.1-3.4%)	2.3% (1.3-4.1%)	59 (32-110)	840 (450-1500)
Mali	2000	Male	80 to 84	4.4% (2.5-6.8%)	96% (93-97%)	0.24 (0.11-0.43)	1.5% (0.86-2.5%)	1.8% (1-3.1%)	27 (15-46)	280 (160-480)
Mali	2000	Male	85 to 89	3.8% (2.1-5.9%)	96% (94-98%)	0.18 (0.084-0.34)	1.1% (0.62-2%)	1.2% (0.69-2.4%)	11 (6.4-22)	86 (50-160)
Mali	2000	Male	90 to 94	3.8% (2.3-6.1%)	96% (94-98%)	0.15 (0.072-0.3)	0.64% (0.34-1.3%)	0.7% (0.34-1.5%)	2.2 (1-4.6)	12 (6.3-24)
Mali	2000	Male	95 plus	4.4% (2.4-7.3%)	96% (93-98%)	0.15 (0.054-0.31)	0.51% (0.22-1.1%)	0.51% (0.15-1.3%)	0.32 (0.093-0.8)	1.3 (0.56-2.9)
Mali	2005	Female	All Ages	4.6% (3.8-5.5%)	95% (95-96%)	0.33 (0.24-0.43)	0.58% (0.42-0.75%)	1.2% (0.78-1.7%)	940 (580-1300)	32000 (23000-42000)
Mali	2005	Female	15 to 19	4.5% (2.8-6.7%)	95% (93-97%)	0.31 (0.15-0.52)	0.71% (0.45-1%)	0.54% (0.25-0.95%)	6.5 (2.9-12)	1100 (650-1600)
Mali	2005	Female	20 to 24	5.4% (3.3-8.1%)	95% (92-97%)	0.47 (0.24-0.81)	1.4% (0.96-1.9%)	0.86% (0.46-1.4%)	13 (6.9-22)	2100 (1500-3000)
Mali	2005	Female	25 to 29	5.8% (3.6-8.7%)	94% (91-96%)	0.56 (0.3-0.89)	1.5% (1.1-2.1%)	1.2% (0.71-1.9%)	20 (11-31)	2300 (1600-3200)
Mali	2005	Female	30 to 34	6.6% (4.1-10%)	93% (90-96%)	0.65 (0.36-1)	1.7% (1.2-2.3%)	1.6% (1-2.5%)	27 (16-41)	2400 (1600-3200)
Mali	2005	Female	35 to 39	7.7% (4.7-11%)	92% (89-95%)	0.75 (0.4-1.2)	2% (1.4-2.8%)	2.3% (1.4-3.4%)	37 (23-55)	2400 (1700-3300)
Mali	2005	Female	40 to 44	8% (5.1-11%)	92% (89-95%)	0.76 (0.39-1.2)	2.3% (1.6-3.1%)	2.9% (1.8-4.1%)	43 (27-62)	2300 (1600-3200)
Mali	2005	Female	45 to 49	7.8% (4.5-11%)	92% (89-95%)	0.7 (0.34-1.2)	2.6% (1.6-3.7%)	3.3% (2-5%)	51 (30-79)	2300 (1500-3500)
Mali	2005	Female	50 to 54	7.9% (5.2-12%)	92% (88-95%)	0.73 (0.37-1.3)	2.9% (1.9-4.3%)	3.8% (2.4-5.7%)	69 (42-100)	2700 (1700-4000)
Mali	2005	Female	55 to 59	8.6% (5.4-13%)	91% (87-95%)	0.89 (0.46-1.6)	3.3% (2.1-4.9%)	4.3% (2.6-6.4%)	87 (51-130)	2900 (1800-4300)
Mali	2005	Female	60 to 64	10% (6.7-15%)	90% (85-93%)	1.1 (0.55-1.9)	3.6% (2.1-5.4%)	4.5% (2.6-6.7%)	120 (72-190)	3400 (2000-5100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	2005	Female	65 to 69	12% (8.2-17%)	88% (83-92%)	1.1 (0.59-1.9)	4.2% (2.3-6.3%)	5% (2.8-7.6%)	150 (80-240)	3500 (1900-5400)
Mali	2005	Female	70 to 74	11% (7.4-16%)	89% (84-93%)	0.9 (0.45-1.6)	3.7% (1.7-6.2%)	4.4% (2-7.4%)	150 (67-260)	2700 (1300-4700)
Mali	2005	Female	75 to 79	8.4% (5-13%)	92% (87-95%)	0.62 (0.28-1.1)	2.5% (1.1-4.4%)	2.9% (1.2-5.2%)	90 (38-160)	1300 (540-2300)
Mali	2005	Female	80 to 84	7.3% (4.3-11%)	93% (89-96%)	0.52 (0.24-0.94)	1.9% (0.81-3.3%)	2.2% (0.91-3.9%)	39 (17-71)	410 (180-740)
Mali	2005	Female	85 to 89	7.6% (4.5-12%)	92% (88-96%)	0.52 (0.26-1)	1.5% (0.61-2.8%)	1.8% (0.67-3.3%)	22 (8.4-40)	160 (66-300)
Mali	2005	Female	90 to 94	8.3% (5.2-13%)	92% (87-95%)	0.55 (0.26-1)	1.3% (0.39-2.6%)	1.5% (0.41-3%)	7.6 (2.1-16)	40 (12-82)
Mali	2005	Female	95 plus	9.5% (5.5-15%)	91% (85-95%)	0.59 (0.23-1.2)	1.2% (0.29-2.8%)	1.4% (0.25-3.3%)	1.9 (0.34-4.5)	6.7 (1.6-15)
Mali	2005	Male	All Ages	2.3% (1.9-2.9%)	98% (97-98%)	0.17 (0.13-0.23)	0.5% (0.38-0.67%)	0.92% (0.65-1.3%)	750 (520-1100)	31000 (23000-42000)
Mali	2005	Male	15 to 19	2% (1.2-2.9%)	98% (97-99%)	0.11 (0.056-0.19)	0.66% (0.45-0.93%)	0.37% (0.2-0.68%)	4 (2.1-7.3)	900 (610-1300)
Mali	2005	Male	20 to 24	2.9% (1.7-4.4%)	97% (96-98%)	0.23 (0.12-0.41)	1.6% (1.1-2.2%)	1% (0.59-1.7%)	15 (8.3-25)	2300 (1500-3200)
Mali	2005	Male	25 to 29	3.6% (2.1-5.4%)	96% (95-98%)	0.34 (0.18-0.55)	2% (1.4-2.8%)	1.7% (1-2.8%)	29 (18-48)	3000 (2100-4100)
Mali	2005	Male	30 to 34	3.9% (2.3-6.2%)	96% (94-98%)	0.4 (0.2-0.66)	2.2% (1.6-3.3%)	2.2% (1.3-3.5%)	41 (25-65)	3300 (2300-4600)
Mali	2005	Male	35 to 39	4% (2.2-6.1%)	96% (94-98%)	0.42 (0.21-0.71)	2.6% (1.8-3.8%)	2.9% (1.9-4.7%)	52 (33-82)	3200 (2200-4800)
Mali	2005	Male	40 to 44	4.4% (2.6-6.5%)	96% (93-97%)	0.46 (0.23-0.78)	3% (2.1-4.4%)	3.6% (2.5-5.6%)	59 (39-94)	3100 (2100-4600)
Mali	2005	Male	45 to 49	5.2% (3-7.3%)	95% (93-97%)	0.53 (0.26-0.92)	3.6% (2.5-5.3%)	4.5% (3-7%)	71 (46-110)	3200 (2200-4600)
Mali	2005	Male	50 to 54	5.6% (3.6-8.2%)	94% (92-96%)	0.56 (0.28-0.97)	3.6% (2.5-5.2%)	4.6% (3-6.9%)	75 (49-110)	2900 (2000-4300)
Mali	2005	Male	55 to 59	5.6% (3.5-8%)	94% (92-96%)	0.55 (0.29-0.98)	3.6% (2.4-5.4%)	4.6% (3-6.9%)	72 (47-110)	2400 (1600-3500)
Mali	2005	Male	60 to 64	5.6% (3.7-8.2%)	94% (92-96%)	0.52 (0.25-0.95)	3.3% (2.2-4.8%)	4.1% (2.7-6.2%)	83 (52-130)	2300 (1500-3500)
Mali	2005	Male	65 to 69	5.8% (3.7-8.3%)	94% (92-96%)	0.47 (0.24-0.81)	2.7% (1.8-4.2%)	3.4% (2.2-5.3%)	73 (46-110)	1700 (1100-2600)
Mali	2005	Male	70 to 74	6.1% (4-8.8%)	94% (91-96%)	0.43 (0.22-0.74)	2.4% (1.5-3.9%)	3% (1.8-4.8%)	75 (44-120)	1400 (820-2200)
Mali	2005	Male	75 to 79	5.6% (3.4-8.7%)	94% (91-97%)	0.35 (0.15-0.64)	1.9% (1.1-3.3%)	2.3% (1.2-4.1%)	56 (30-100)	790 (430-1400)
Mali	2005	Male	80 to 84	4.4% (2.6-6.8%)	96% (93-97%)	0.24 (0.11-0.44)	1.5% (0.87-2.6%)	1.7% (1-3.1%)	27 (15-48)	280 (160-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	2005	Male	85 to 89	3.8% (2-5.9%)	96% (94-98%)	0.18 (0.084-0.34)	1% (0.62-1.9%)	1.2% (0.68-2.2%)	13 (7.3-24)	97 (58-180)
Mali	2005	Male	90 to 94	3.8% (2.3-5.9%)	96% (94-98%)	0.16 (0.071-0.3)	0.63% (0.33-1.2%)	0.69% (0.33-1.4%)	3 (1.4-6.1)	17 (8.9-32)
Mali	2005	Male	95 plus	4.4% (2.4-7.3%)	96% (93-98%)	0.15 (0.054-0.32)	0.5% (0.23-1.1%)	0.49% (0.15-1.3%)	0.46 (0.14-1.2)	1.9 (0.85-4.2)
Mali	2010	Female	All Ages	5% (4.1-5.9%)	95% (94-96%)	0.33 (0.22-0.46)	0.56% (0.39-0.75%)	1.1% (0.64-1.6%)	820 (480-1200)	31000 (21000-41000)
Mali	2010	Female	15 to 19	4.8% (3-7.1%)	95% (93-97%)	0.3 (0.14-0.56)	0.69% (0.43-1%)	0.46% (0.2-0.88%)	5.8 (2.2-12)	1100 (680-1700)
Mali	2010	Female	20 to 24	6.2% (3.8-9.3%)	94% (91-96%)	0.5 (0.24-0.89)	1.4% (0.93-2%)	0.78% (0.38-1.3%)	11 (5.4-20)	2200 (1400-3200)
Mali	2010	Female	25 to 29	6.9% (4.1-10%)	93% (90-96%)	0.6 (0.3-0.99)	1.6% (1.1-2.1%)	1.1% (0.6-1.8%)	18 (9.8-30)	2400 (1600-3400)
Mali	2010	Female	30 to 34	7.7% (4.5-12%)	92% (88-95%)	0.69 (0.37-1.2)	1.7% (1.2-2.3%)	1.6% (0.92-2.3%)	26 (15-40)	2500 (1700-3400)
Mali	2010	Female	35 to 39	8.8% (5.2-13%)	91% (87-95%)	0.79 (0.38-1.3)	2% (1.3-2.7%)	2.2% (1.3-3.3%)	37 (22-56)	2600 (1700-3600)
Mali	2010	Female	40 to 44	9.2% (5.6-13%)	91% (87-94%)	0.8 (0.38-1.4)	2.3% (1.5-3.2%)	3% (1.8-4.4%)	45 (27-68)	2500 (1600-3600)
Mali	2010	Female	45 to 49	9% (5.2-13%)	91% (87-95%)	0.75 (0.34-1.3)	2.6% (1.5-3.9%)	3.4% (1.9-5.3%)	51 (28-81)	2400 (1400-3600)
Mali	2010	Female	50 to 54	9% (5.7-13%)	91% (87-94%)	0.76 (0.35-1.4)	2.8% (1.7-4.3%)	3.8% (2.2-5.8%)	63 (35-99)	2500 (1500-3800)
Mali	2010	Female	55 to 59	9.3% (5.8-14%)	91% (86-94%)	0.88 (0.42-1.6)	3.1% (1.7-4.7%)	4% (2.2-6.3%)	84 (46-140)	2800 (1600-4500)
Mali	2010	Female	60 to 64	10% (6.1-15%)	90% (85-94%)	0.96 (0.44-1.8)	3.1% (1.7-4.9%)	3.9% (2.1-6.1%)	96 (51-150)	2700 (1500-4200)
Mali	2010	Female	65 to 69	11% (7.3-16%)	89% (84-93%)	0.97 (0.48-1.7)	3.4% (1.7-5.4%)	4.2% (2-6.7%)	130 (64-220)	3100 (1500-5000)
Mali	2010	Female	70 to 74	10% (7-15%)	90% (85-93%)	0.78 (0.36-1.4)	3% (1.2-5.4%)	3.6% (1.4-6.5%)	120 (47-220)	2200 (900-4000)
Mali	2010	Female	75 to 79	8.2% (4.9-13%)	92% (87-95%)	0.56 (0.25-1.1)	2.1% (0.77-3.9%)	2.5% (0.89-4.6%)	73 (26-140)	1000 (390-2000)
Mali	2010	Female	80 to 84	7.2% (4.3-11%)	93% (89-96%)	0.47 (0.2-0.89)	1.6% (0.56-3%)	1.8% (0.63-3.4%)	32 (11-63)	350 (130-660)
Mali	2010	Female	85 to 89	7.5% (4.4-11%)	93% (89-96%)	0.47 (0.22-0.92)	1.3% (0.42-2.5%)	1.5% (0.44-3%)	19 (5.8-38)	140 (47-280)
Mali	2010	Female	90 to 94	8.3% (5.1-13%)	92% (87-95%)	0.5 (0.22-0.98)	1.1% (0.24-2.4%)	1.2% (0.2-2.8%)	7 (1.1-16)	37 (8.2-83)
Mali	2010	Female	95 plus	9.5% (5.3-15%)	91% (85-95%)	0.54 (0.2-1.1)	1% (0.13-2.5%)	1.1% (0.039-2.9%)	1.9 (0.068-5.2)	7.1 (0.91-18)
Mali	2010	Male	All Ages	2.6% (2-3.2%)	97% (97-98%)	0.18 (0.12-0.26)	0.52% (0.4-0.67%)	0.88% (0.63-1.2%)	710 (510-980)	32000 (25000-42000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	2010	Male	15 to 19	2% (1.1-3.2%)	98% (97-99%)	0.11 (0.05-0.21)	0.67% (0.45-0.94%)	0.34% (0.18-0.61%)	4.1 (2.1-7.5)	1000 (650-1500)
Mali	2010	Male	20 to 24	3.3% (1.7-5%)	97% (95-98%)	0.24 (0.11-0.43)	1.6% (1.1-2.2%)	0.99% (0.53-1.6%)	15 (8-25)	2500 (1700-3600)
Mali	2010	Male	25 to 29	4.2% (2.4-6.7%)	96% (93-98%)	0.37 (0.18-0.65)	2.1% (1.5-2.8%)	1.8% (1-2.8%)	31 (18-50)	3400 (2300-4700)
Mali	2010	Male	30 to 34	4.8% (2.7-8%)	95% (92-97%)	0.45 (0.23-0.78)	2.4% (1.7-3.2%)	2.3% (1.4-3.4%)	46 (28-69)	3700 (2600-5200)
Mali	2010	Male	35 to 39	5% (2.8-7.6%)	95% (92-97%)	0.48 (0.24-0.85)	2.7% (1.9-3.7%)	3% (1.9-4.4%)	62 (40-91)	3900 (2700-5500)
Mali	2010	Male	40 to 44	5.2% (3.1-7.8%)	95% (92-97%)	0.49 (0.23-0.86)	3% (2.1-4.1%)	3.6% (2.5-5.1%)	69 (45-99)	3600 (2500-5000)
Mali	2010	Male	45 to 49	5.4% (3-8.1%)	95% (92-97%)	0.51 (0.22-0.94)	3.4% (2.3-4.9%)	4.3% (2.7-6.3%)	72 (46-110)	3300 (2200-4700)
Mali	2010	Male	50 to 54	5.5% (3.3-8.1%)	94% (92-97%)	0.51 (0.23-0.95)	3.3% (2.2-4.4%)	4.1% (2.8-5.7%)	69 (44-100)	2700 (1800-3800)
Mali	2010	Male	55 to 59	5.4% (3.1-8.1%)	95% (92-97%)	0.49 (0.22-0.91)	3.3% (2.2-4.6%)	4.1% (2.7-5.9%)	76 (48-110)	2600 (1700-3700)
Mali	2010	Male	60 to 64	5.2% (3-8.3%)	95% (92-97%)	0.45 (0.19-0.88)	2.8% (1.8-4.1%)	3.5% (2.2-5.1%)	69 (43-100)	2000 (1300-2800)
Mali	2010	Male	65 to 69	5.3% (3.1-7.8%)	95% (92-97%)	0.4 (0.18-0.72)	2.3% (1.5-3.4%)	2.8% (1.8-4.1%)	68 (43-100)	1600 (1000-2300)
Mali	2010	Male	70 to 74	5.1% (3.2-7.6%)	95% (92-97%)	0.33 (0.15-0.62)	1.9% (1.1-2.9%)	2.3% (1.4-3.6%)	58 (34-92)	1100 (640-1700)
Mali	2010	Male	75 to 79	4.6% (2.7-7.2%)	95% (93-97%)	0.26 (0.11-0.5)	1.4% (0.77-2.3%)	1.7% (0.89-2.9%)	41 (21-70)	590 (320-990)
Mali	2010	Male	80 to 84	4% (2.3-5.9%)	96% (94-98%)	0.2 (0.085-0.38)	1.2% (0.69-1.9%)	1.4% (0.78-2.2%)	22 (12-35)	230 (140-370)
Mali	2010	Male	85 to 89	3.7% (2-5.8%)	96% (94-98%)	0.16 (0.073-0.31)	0.84% (0.49-1.4%)	0.96% (0.53-1.6%)	10 (5.8-18)	81 (47-130)
Mali	2010	Male	90 to 94	3.8% (2.3-6%)	96% (94-98%)	0.14 (0.063-0.28)	0.5% (0.26-0.92%)	0.53% (0.23-1.1%)	2.7 (1.1-5.2)	15 (7.8-28)
Mali	2010	Male	95 plus	4.4% (2.3-7.4%)	96% (93-98%)	0.14 (0.047-0.3)	0.39% (0.16-0.82%)	0.35% (0.079-0.87%)	0.46 (0.1-1.1)	2.1 (0.89-4.3)
Mali	2016	Female	All Ages	5.1% (4.2-6.1%)	95% (94-96%)	0.36 (0.19-0.61)	0.63% (0.4-0.89%)	1.2% (0.59-1.9%)	900 (440-1500)	34000 (22000-49000)
Mali	2016	Female	15 to 19	4.9% (3.1-7.3%)	95% (93-97%)	0.33 (0.14-0.68)	0.73% (0.46-1%)	0.48% (0.18-0.93%)	6.7 (2.2-14)	1300 (810-2000)
Mali	2016	Female	20 to 24	6.3% (3.8-9.5%)	94% (91-96%)	0.55 (0.23-1.1)	1.5% (0.98-2%)	0.77% (0.34-1.3%)	12 (4.8-23)	2600 (1600-3700)
Mali	2016	Female	25 to 29	6.9% (4.1-11%)	93% (89-96%)	0.65 (0.28-1.3)	1.6% (1.2-2.2%)	1.1% (0.58-1.8%)	18 (8.5-30)	2600 (1700-3700)
Mali	2016	Female	30 to 34	7.8% (4.6-12%)	92% (88-95%)	0.76 (0.33-1.5)	1.7% (1.2-2.4%)	1.6% (0.87-2.5%)	26 (14-43)	2700 (1700-3800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	2016	Female	35 to 39	9% (5.3-13%)	91% (87-95%)	0.86 (0.35-1.6)	2% (1.3-2.8%)	2.3% (1.3-3.4%)	40 (22-62)	2900 (1900-4100)
Mali	2016	Female	40 to 44	9.4% (5.8-14%)	91% (86-94%)	0.88 (0.34-1.8)	2.3% (1.4-3.3%)	3% (1.7-4.5%)	53 (29-83)	3000 (1900-4500)
Mali	2016	Female	45 to 49	9.2% (5.3-14%)	91% (86-95%)	0.81 (0.31-1.7)	2.5% (1.5-3.8%)	3.4% (1.8-5.5%)	60 (31-100)	2900 (1600-4500)
Mali	2016	Female	50 to 54	9.2% (5.8-13%)	91% (87-94%)	0.83 (0.33-1.7)	2.8% (1.6-4.4%)	3.7% (2-5.9%)	70 (37-120)	2800 (1600-4500)
Mali	2016	Female	55 to 59	9.5% (6-14%)	90% (86-94%)	0.97 (0.4-2)	3% (1.6-4.6%)	3.9% (2-6.2%)	86 (42-150)	2900 (1500-4800)
Mali	2016	Female	60 to 64	10% (6.2-16%)	90% (84-94%)	1.1 (0.42-2.2)	3% (1.6-4.8%)	3.8% (1.9-6.1%)	110 (50-180)	3100 (1500-5100)
Mali	2016	Female	65 to 69	12% (7.5-17%)	88% (83-93%)	1.1 (0.44-2.2)	3.4% (1.5-5.4%)	4.1% (1.8-6.8%)	130 (54-240)	3100 (1300-5400)
Mali	2016	Female	70 to 74	11% (7-16%)	89% (84-93%)	0.86 (0.34-1.8)	3% (1.1-5.4%)	3.6% (1.2-6.4%)	140 (45-260)	2500 (870-4800)
Mali	2016	Female	75 to 79	8.4% (5.1-13%)	92% (87-95%)	0.62 (0.23-1.3)	2.1% (0.67-4%)	2.5% (0.75-4.9%)	83 (24-170)	1200 (370-2400)
Mali	2016	Female	80 to 84	7.4% (4.4-11%)	93% (89-96%)	0.52 (0.2-1.1)	1.6% (0.51-3.1%)	1.9% (0.58-3.8%)	35 (10-71)	380 (120-750)
Mali	2016	Female	85 to 89	7.6% (4.5-11%)	92% (89-95%)	0.52 (0.2-1.1)	1.3% (0.36-2.8%)	1.5% (0.39-3.3%)	21 (5.3-46)	160 (43-340)
Mali	2016	Female	90 to 94	8.3% (5.2-13%)	92% (87-95%)	0.54 (0.21-1.2)	1.2% (0.2-2.7%)	1.3% (0.16-3.2%)	8.6 (1-21)	46 (7.4-110)
Mali	2016	Female	95 plus	9.5% (5.4-15%)	90% (85-95%)	0.59 (0.19-1.4)	1.1% (0.079-2.7%)	1.2% (-0.013-3.2%)	2.5 (-0.026-6.4)	9.1 (0.66-22)
Mali	2016	Male	All Ages	2.5% (2-3.2%)	97% (97-98%)	0.19 (0.098-0.33)	0.59% (0.42-0.82%)	0.96% (0.63-1.4%)	780 (510-1200)	37000 (26000-51000)
Mali	2016	Male	15 to 19	2% (1.2-3.3%)	98% (97-99%)	0.12 (0.048-0.25)	0.71% (0.46-0.99%)	0.36% (0.18-0.66%)	4.8 (2.3-8.9)	1200 (760-1800)
Mali	2016	Male	20 to 24	3.2% (1.8-5%)	97% (95-98%)	0.25 (0.096-0.49)	1.7% (1.2-2.3%)	0.96% (0.49-1.6%)	16 (8-28)	2800 (1800-4100)
Mali	2016	Male	25 to 29	4.2% (2.4-6.6%)	96% (93-98%)	0.39 (0.16-0.78)	2.2% (1.5-2.9%)	1.7% (0.95-2.7%)	31 (17-51)	3600 (2400-5000)
Mali	2016	Male	30 to 34	4.7% (2.7-7.7%)	95% (92-97%)	0.47 (0.2-0.94)	2.4% (1.7-3.3%)	2.3% (1.4-3.5%)	47 (28-75)	4000 (2700-5700)
Mali	2016	Male	35 to 39	5% (2.8-7.6%)	95% (92-97%)	0.51 (0.21-0.98)	2.7% (1.9-3.8%)	3% (1.9-4.5%)	68 (40-110)	4400 (2900-6400)
Mali	2016	Male	40 to 44	5.1% (3.1-7.7%)	95% (92-97%)	0.52 (0.21-1.1)	3% (2.1-4.2%)	3.7% (2.4-5.3%)	84 (52-130)	4500 (3000-6700)
Mali	2016	Male	45 to 49	5.4% (3-8%)	95% (92-97%)	0.54 (0.2-1.1)	3.4% (2.2-4.9%)	4.2% (2.6-6.2%)	90 (52-140)	4100 (2500-6300)
Mali	2016	Male	50 to 54	5.5% (3.4-8.3%)	94% (92-97%)	0.55 (0.2-1.1)	3.2% (2.1-4.4%)	4% (2.7-5.6%)	78 (46-120)	3100 (1900-4600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mali	2016	Male	55 to 59	5.4% (3.2-8%)	95% (92-97%)	0.52 (0.2-1.1)	3.2% (2.1-4.5%)	4.1% (2.6-5.9%)	77 (47-120)	2600 (1600-4000)
Mali	2016	Male	60 to 64	5.2% (3-8.2%)	95% (92-97%)	0.47 (0.17-1)	2.8% (1.7-4%)	3.4% (2.1-5%)	80 (47-130)	2300 (1400-3600)
Mali	2016	Male	65 to 69	5.2% (3.1-7.7%)	95% (92-97%)	0.42 (0.16-0.86)	2.2% (1.4-3.2%)	2.8% (1.7-4%)	66 (40-100)	1600 (960-2400)
Mali	2016	Male	70 to 74	5.1% (3.2-7.3%)	95% (93-97%)	0.35 (0.14-0.73)	1.9% (1.1-2.9%)	2.3% (1.3-3.6%)	64 (35-100)	1200 (660-1900)
Mali	2016	Male	75 to 79	4.6% (2.7-7.2%)	95% (93-97%)	0.28 (0.097-0.59)	1.4% (0.73-2.4%)	1.7% (0.84-2.9%)	42 (20-74)	600 (300-1000)
Mali	2016	Male	80 to 84	4% (2.3-6%)	96% (94-98%)	0.21 (0.08-0.45)	1.2% (0.65-2%)	1.4% (0.74-2.4%)	21 (11-37)	230 (120-390)
Mali	2016	Male	85 to 89	3.7% (1.9-5.9%)	96% (94-98%)	0.17 (0.063-0.37)	0.87% (0.44-1.6%)	0.99% (0.47-1.8%)	11 (5-20)	85 (42-150)
Mali	2016	Male	90 to 94	3.8% (2.1-5.9%)	96% (94-98%)	0.15 (0.055-0.33)	0.54% (0.24-1.1%)	0.57% (0.21-1.2%)	3 (1.1-6.3)	17 (7.5-34)
Mali	2016	Male	95 plus	4.3% (2.3-7.1%)	96% (93-98%)	0.14 (0.043-0.33)	0.43% (0.16-0.93%)	0.39% (0.062-1%)	0.58 (0.089-1.5)	2.6 (0.93-5.6)
Mauritania	1990	Female	All Ages	4.3% (3.6-5.2%)	96% (95-96%)	0.041 (0.034-0.049)	0.38% (0.3-0.47%)	0.49% (0.34-0.67%)	45 (31-61)	2200 (1700-2800)
Mauritania	1990	Female	15 to 19	4.1% (2.5-6.1%)	96% (94-98%)	0.037 (0.019-0.065)	0.44% (0.28-0.66%)	0.068% (0.015-0.14%)	0.12 (0.027-0.24)	100 (60-160)
Mauritania	1990	Female	20 to 24	5.1% (3-7.7%)	95% (92-97%)	0.059 (0.03-0.097)	0.95% (0.6-1.4%)	0.15% (0.062-0.25%)	0.32 (0.13-0.53)	220 (130-350)
Mauritania	1990	Female	25 to 29	5.7% (3.4-8.6%)	94% (91-97%)	0.07 (0.038-0.11)	1% (0.73-1.4%)	0.32% (0.17-0.48%)	0.65 (0.35-0.99)	210 (140-300)
Mauritania	1990	Female	30 to 34	6.3% (3.7-10%)	94% (90-96%)	0.079 (0.044-0.12)	1% (0.74-1.4%)	0.59% (0.34-0.87%)	1.2 (0.7-1.8)	190 (130-260)
Mauritania	1990	Female	35 to 39	7% (4-10%)	93% (90-96%)	0.085 (0.047-0.14)	0.99% (0.72-1.3%)	0.79% (0.49-1.1%)	1.7 (1-2.5)	170 (120-220)
Mauritania	1990	Female	40 to 44	7.3% (4.4-11%)	93% (89-96%)	0.087 (0.044-0.14)	1.2% (0.86-1.7%)	1.4% (0.86-2%)	3.2 (2-4.8)	200 (140-280)
Mauritania	1990	Female	45 to 49	7.3% (4-11%)	93% (89-96%)	0.084 (0.042-0.14)	1.1% (0.75-1.5%)	1.2% (0.74-1.9%)	3.5 (2-5.5)	180 (120-260)
Mauritania	1990	Female	50 to 54	7.3% (4.5-11%)	93% (89-95%)	0.089 (0.046-0.15)	1.4% (0.89-2%)	1.7% (0.99-2.5%)	6 (3.5-9.3)	240 (150-360)
Mauritania	1990	Female	55 to 59	7.6% (4.6-11%)	92% (89-95%)	0.1 (0.053-0.18)	1.1% (0.68-1.6%)	1.3% (0.78-2%)	5.7 (3.3-9)	190 (120-300)
Mauritania	1990	Female	60 to 64	8% (4.8-12%)	92% (88-95%)	0.11 (0.053-0.19)	1.1% (0.63-1.7%)	1.3% (0.73-2%)	7.2 (4-11)	200 (120-300)
Mauritania	1990	Female	65 to 69	8.2% (5.3-12%)	92% (88-95%)	0.1 (0.051-0.17)	0.81% (0.44-1.2%)	0.94% (0.48-1.5%)	5.9 (2.9-9.4)	140 (72-220)
Mauritania	1990	Female	70 to 74	7.5% (4.9-11%)	93% (89-95%)	0.081 (0.041-0.14)	0.68% (0.31-1.1%)	0.77% (0.3-1.3%)	5.2 (2.1-9.1)	97 (44-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	1990	Female	75 to 79	6.3% (3.8-9.7%)	94% (90-96%)	0.063 (0.029-0.11)	0.47% (0.16-0.8%)	0.53% (0.16-0.93%)	3.1 (0.97-5.5)	46 (15-79)
Mauritania	1990	Female	80 to 84	5.8% (3.4-8.3%)	94% (92-97%)	0.055 (0.026-0.096)	0.28% (0.048-0.51%)	0.31% (0.032-0.59%)	1 (0.11-2)	11 (2-21)
Mauritania	1990	Female	85 to 89	6% (3.6-9.3%)	94% (91-96%)	0.056 (0.027-0.11)	0.12% (-0.089-0.31%)	0.13% (-0.13-0.35%)	0.23 (-0.24-0.64)	2 (-1.3-4.9)
Mauritania	1990	Female	90 to 94	6.8% (4.2-10%)	93% (90-96%)	0.06 (0.028-0.12)	0.0099% (-0.21-0.19%)	-0.022% (-0.29-0.2%)	-0.014 (-0.19-0.13)	0.039 (-0.82-0.74)
Mauritania	1990	Female	95 plus	7.9% (4.5-13%)	92% (87-96%)	0.068 (0.025-0.14)	0.011% (-0.26-0.22%)	-0.047% (-0.36-0.21%)	-0.0072 (-0.057-0.033)	0.0066 (-0.16-0.14)
Mauritania	1990	Male	All Ages	2.6% (2.1-3.2%)	97% (97-98%)	0.099 (0.085-0.12)	0.74% (0.57-0.93%)	1.1% (0.81-1.5%)	100 (70-130)	4600 (3500-5800)
Mauritania	1990	Male	15 to 19	2.1% (1.1-3.1%)	98% (97-99%)	0.059 (0.028-0.11)	0.59% (0.39-0.85%)	0.27% (0.16-0.44%)	0.56 (0.32-0.88)	140 (90-210)
Mauritania	1990	Male	20 to 24	3.3% (1.8-5%)	97% (95-98%)	0.13 (0.063-0.21)	1.3% (0.92-1.9%)	0.74% (0.42-1.1%)	2 (1.2-3.2)	340 (240-480)
Mauritania	1990	Male	25 to 29	4% (2.4-6.3%)	96% (94-98%)	0.19 (0.099-0.29)	2% (1.4-2.6%)	1.5% (0.85-2.3%)	3.9 (2.1-6.2)	440 (300-600)
Mauritania	1990	Male	30 to 34	4.6% (2.7-7.4%)	95% (93-97%)	0.23 (0.13-0.36)	2.5% (1.8-3.4%)	2.4% (1.4-3.6%)	5.9 (3.3-9)	500 (330-700)
Mauritania	1990	Male	35 to 39	5.1% (2.9-7.7%)	95% (92-97%)	0.25 (0.14-0.41)	3.1% (2.1-4.1%)	3.4% (2.1-4.9%)	8.2 (4.8-12)	530 (340-720)
Mauritania	1990	Male	40 to 44	5.2% (3.1-8%)	95% (92-97%)	0.26 (0.13-0.42)	3.3% (2.3-4.5%)	3.9% (2.6-5.6%)	8.9 (5.5-13)	470 (320-650)
Mauritania	1990	Male	45 to 49	5.2% (2.9-8%)	95% (92-97%)	0.26 (0.13-0.44)	3.5% (2.4-5%)	4.3% (2.8-6.3%)	11 (6.7-16)	490 (320-710)
Mauritania	1990	Male	50 to 54	5.3% (3.1-7.9%)	95% (92-97%)	0.27 (0.13-0.46)	3.1% (2.1-4.4%)	3.7% (2.4-5.4%)	11 (6.8-16)	440 (280-620)
Mauritania	1990	Male	55 to 59	5.3% (3.1-8.2%)	95% (92-97%)	0.26 (0.13-0.43)	2.9% (1.9-4.1%)	3.4% (2.2-5.1%)	12 (7.4-18)	410 (270-590)
Mauritania	1990	Male	60 to 64	5.2% (3-8.2%)	95% (92-97%)	0.23 (0.11-0.42)	2.4% (1.6-3.4%)	2.8% (1.8-4.1%)	12 (7.6-18)	350 (230-500)
Mauritania	1990	Male	65 to 69	5.3% (3.2-7.7%)	95% (92-97%)	0.2 (0.1-0.35)	1.8% (1.2-2.5%)	2.1% (1.4-3%)	9.8 (6.2-15)	230 (150-340)
Mauritania	1990	Male	70 to 74	5.1% (3.2-7.5%)	95% (92-97%)	0.17 (0.083-0.29)	1.3% (0.83-1.9%)	1.5% (0.95-2.3%)	7.3 (4.6-11)	130 (87-210)
Mauritania	1990	Male	75 to 79	4.7% (2.7-7.4%)	95% (93-97%)	0.13 (0.059-0.24)	0.92% (0.59-1.4%)	1.1% (0.66-1.7%)	4.2 (2.6-6.5)	61 (39-92)
Mauritania	1990	Male	80 to 84	4% (2.2-6.1%)	96% (94-98%)	0.098 (0.046-0.17)	0.74% (0.48-1.1%)	0.84% (0.52-1.3%)	1.9 (1.2-2.8)	20 (13-30)
Mauritania	1990	Male	85 to 89	3.6% (2-5.7%)	96% (94-98%)	0.076 (0.036-0.14)	0.56% (0.36-0.81%)	0.62% (0.38-0.93%)	0.74 (0.45-1.1)	5.8 (3.7-8.5)
Mauritania	1990	Male	90 to 94	3.7% (2.2-6%)	96% (94-98%)	0.066 (0.03-0.12)	0.33% (0.19-0.49%)	0.33% (0.16-0.53%)	0.14 (0.063-0.22)	0.81 (0.46-1.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	1990	Male	95 plus	4.2% (2.3-6.9%)	96% (93-98%)	0.062 (0.023-0.13)	0.3% (0.13-0.47%)	0.26% (0.041-0.47%)	0.023 (0.0035-0.041)	0.11 (0.043-0.17)
Mauritania	1995	Female	All Ages	4.3% (3.5-5.1%)	96% (95-97%)	0.03 (0.025-0.036)	0.38% (0.3-0.48%)	0.47% (0.32-0.64%)	46 (31-63)	2400 (1700-3000)
Mauritania	1995	Female	15 to 19	4% (2.4-5.9%)	96% (94-98%)	0.027 (0.013-0.046)	0.46% (0.28-0.68%)	0.059% (0.008-0.13%)	0.11 (0.015-0.23)	110 (66-170)
Mauritania	1995	Female	20 to 24	5.1% (2.9-7.6%)	95% (92-97%)	0.043 (0.022-0.07)	0.98% (0.62-1.4%)	0.12% (0.041-0.21%)	0.28 (0.091-0.47)	240 (140-370)
Mauritania	1995	Female	25 to 29	5.6% (3.3-8.7%)	94% (91-97%)	0.051 (0.029-0.083)	1% (0.74-1.4%)	0.28% (0.13-0.44%)	0.61 (0.28-0.96)	240 (150-340)
Mauritania	1995	Female	30 to 34	6.2% (3.6-10%)	94% (90-96%)	0.058 (0.033-0.092)	1% (0.72-1.4%)	0.51% (0.24-0.79%)	1.2 (0.52-1.8)	210 (140-300)
Mauritania	1995	Female	35 to 39	6.8% (3.9-10%)	93% (90-96%)	0.062 (0.033-0.099)	0.95% (0.66-1.2%)	0.7% (0.35-1%)	1.7 (0.86-2.6)	190 (130-250)
Mauritania	1995	Female	40 to 44	7% (4.2-11%)	93% (89-96%)	0.062 (0.032-0.1)	1.1% (0.76-1.5%)	1.2% (0.69-1.8%)	3.1 (1.7-4.6)	200 (140-280)
Mauritania	1995	Female	45 to 49	7.1% (3.9-11%)	93% (89-96%)	0.061 (0.03-0.1)	1% (0.67-1.4%)	1.1% (0.64-1.7%)	3.3 (1.8-5.2)	180 (120-260)
Mauritania	1995	Female	50 to 54	7.1% (4.3-11%)	93% (89-96%)	0.064 (0.032-0.11)	1.3% (0.8-1.8%)	1.5% (0.88-2.3%)	5.9 (3.3-9.1)	240 (150-360)
Mauritania	1995	Female	55 to 59	7.4% (4.6-11%)	93% (89-95%)	0.074 (0.037-0.13)	0.97% (0.59-1.4%)	1.1% (0.65-1.8%)	5.5 (3.1-8.5)	190 (110-290)
Mauritania	1995	Female	60 to 64	7.8% (4.6-12%)	92% (88-95%)	0.078 (0.039-0.14)	1% (0.6-1.5%)	1.2% (0.68-1.8%)	7.2 (4-11)	200 (120-310)
Mauritania	1995	Female	65 to 69	8% (5.1-12%)	92% (88-95%)	0.072 (0.037-0.12)	0.79% (0.42-1.2%)	0.91% (0.45-1.4%)	6.4 (3.1-10)	150 (78-230)
Mauritania	1995	Female	70 to 74	7.3% (4.8-11%)	93% (89-95%)	0.058 (0.028-0.1)	0.66% (0.3-1.1%)	0.74% (0.31-1.3%)	5.7 (2.4-9.7)	110 (49-180)
Mauritania	1995	Female	75 to 79	6.2% (3.6-9.5%)	94% (91-96%)	0.045 (0.021-0.082)	0.5% (0.21-0.8%)	0.56% (0.2-0.92%)	3.8 (1.4-6.5)	56 (23-92)
Mauritania	1995	Female	80 to 84	5.6% (3.3-8.3%)	94% (92-97%)	0.04 (0.018-0.07)	0.29% (0.069-0.54%)	0.32% (0.061-0.61%)	1.3 (0.23-2.5)	14 (3.3-27)
Mauritania	1995	Female	85 to 89	5.9% (3.5-8.9%)	94% (91-97%)	0.04 (0.019-0.075)	0.14% (-0.052-0.31%)	0.14% (-0.091-0.35%)	0.33 (-0.21-0.81)	2.9 (-1.1-6.2)
Mauritania	1995	Female	90 to 94	6.5% (4-9.9%)	93% (90-96%)	0.043 (0.019-0.081)	0.038% (-0.16-0.2%)	0.008% (-0.24-0.21%)	0.0068 (-0.2-0.18)	0.19 (-0.81-1.1)
Mauritania	1995	Female	95 plus	7.7% (4.2-12%)	92% (88-96%)	0.048 (0.018-0.1)	0.049% (-0.2-0.24%)	-0.0052% (-0.3-0.24%)	-0.0011 (-0.066-0.052)	0.042 (-0.17-0.21)
Mauritania	1995	Male	All Ages	2.9% (2.3-3.5%)	97% (96-98%)	0.08 (0.068-0.093)	0.66% (0.45-0.86%)	0.95% (0.62-1.3%)	88 (55-120)	4300 (2800-5500)
Mauritania	1995	Male	15 to 19	2.2% (1.2-3.4%)	98% (97-99%)	0.046 (0.023-0.082)	0.63% (0.42-0.89%)	0.26% (0.16-0.41%)	0.5 (0.3-0.79)	150 (96-220)
Mauritania	1995	Male	20 to 24	3.5% (2-5.2%)	97% (95-98%)	0.1 (0.051-0.16)	1.4% (1-2%)	0.71% (0.45-1.1%)	1.8 (1.1-2.7)	350 (240-500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	1995	Male	25 to 29	4.3% (2.4-6.5%)	96% (93-98%)	0.15 (0.079-0.23)	1.9% (1.3-2.5%)	1.3% (0.45-2.1%)	3.2 (1-5.1)	430 (280-590)
Mauritania	1995	Male	30 to 34	4.9% (2.8-8.1%)	95% (92-97%)	0.18 (0.1-0.28)	2.3% (1.2-3.1%)	2% (0.26-3.2%)	4.9 (0.63-7.9)	460 (240-650)
Mauritania	1995	Male	35 to 39	5.4% (3.3-8.2%)	95% (92-97%)	0.2 (0.11-0.31)	2.6% (0.99-3.7%)	2.9% (0.38-4.4%)	7.2 (0.95-12)	490 (190-700)
Mauritania	1995	Male	40 to 44	5.6% (3.4-8.4%)	94% (92-97%)	0.2 (0.11-0.33)	2.7% (0.86-3.9%)	3.3% (0.43-4.9%)	8.2 (1.1-13)	460 (150-660)
Mauritania	1995	Male	45 to 49	5.6% (3.1-8.3%)	94% (92-97%)	0.21 (0.1-0.36)	3% (1.5-4.3%)	3.6% (1.6-5.5%)	9.2 (4.2-14)	430 (220-620)
Mauritania	1995	Male	50 to 54	5.7% (3.5-8.4%)	94% (92-97%)	0.21 (0.1-0.38)	2.6% (1.6-3.7%)	3.1% (1.9-4.6%)	9.2 (5.3-14)	370 (230-530)
Mauritania	1995	Male	55 to 59	5.6% (3.3-8.3%)	94% (92-97%)	0.2 (0.1-0.34)	2.5% (1.5-3.5%)	3% (1.8-4.5%)	10 (6.1-15)	350 (220-510)
Mauritania	1995	Male	60 to 64	5.6% (3.3-8.6%)	94% (91-97%)	0.19 (0.089-0.33)	2% (1.3-3%)	2.4% (1.5-3.7%)	11 (6.6-17)	310 (200-460)
Mauritania	1995	Male	65 to 69	5.6% (3.4-8.3%)	94% (92-97%)	0.16 (0.082-0.28)	1.5% (1-2.2%)	1.8% (1.2-2.7%)	9 (5.9-14)	210 (140-320)
Mauritania	1995	Male	70 to 74	5.4% (3.5-8%)	95% (92-97%)	0.13 (0.067-0.23)	1.1% (0.71-1.6%)	1.3% (0.81-2%)	6.9 (4.3-11)	130 (82-200)
Mauritania	1995	Male	75 to 79	4.9% (2.9-7.9%)	95% (92-97%)	0.1 (0.048-0.19)	0.8% (0.52-1.2%)	0.93% (0.59-1.4%)	4.2 (2.6-6.4)	60 (39-91)
Mauritania	1995	Male	80 to 84	4.3% (2.4-6.7%)	96% (93-98%)	0.078 (0.036-0.14)	0.66% (0.42-0.96%)	0.75% (0.45-1.1%)	1.9 (1.2-2.9)	21 (13-31)
Mauritania	1995	Male	85 to 89	3.9% (2.1-6.1%)	96% (94-98%)	0.061 (0.028-0.11)	0.51% (0.34-0.74%)	0.56% (0.35-0.85%)	0.83 (0.52-1.3)	6.6 (4.3-9.7)
Mauritania	1995	Male	90 to 94	4% (2.3-6.3%)	96% (94-98%)	0.052 (0.024-0.099)	0.31% (0.16-0.48%)	0.31% (0.14-0.52%)	0.17 (0.072-0.28)	1 (0.53-1.5)
Mauritania	1995	Male	95 plus	4.5% (2.5-7.2%)	96% (93-98%)	0.048 (0.018-0.099)	0.31% (0.14-0.49%)	0.27% (0.053-0.5%)	0.031 (0.0065-0.06)	0.15 (0.065-0.24)
Mauritania	2000	Female	All Ages	4% (3.3-4.9%)	96% (95-97%)	0.017 (0.014-0.02)	0.4% (0.31-0.5%)	0.48% (0.33-0.63%)	49 (34-67)	2600 (1900-3300)
Mauritania	2000	Female	15 to 19	3.9% (2.4-5.8%)	96% (94-98%)	0.016 (0.0081-0.027)	0.47% (0.3-0.72%)	0.049% (0.0053-0.098%)	0.099 (0.011-0.2)	130 (76-200)
Mauritania	2000	Female	20 to 24	4.9% (3-7.5%)	95% (93-97%)	0.025 (0.013-0.041)	1% (0.64-1.5%)	0.11% (0.036-0.19%)	0.26 (0.087-0.46)	280 (170-440)
Mauritania	2000	Female	25 to 29	5.1% (3-7.9%)	95% (92-97%)	0.029 (0.016-0.045)	1.1% (0.76-1.5%)	0.28% (0.13-0.45%)	0.66 (0.3-1.1)	280 (180-390)
Mauritania	2000	Female	30 to 34	5.3% (3-8.6%)	95% (91-97%)	0.03 (0.016-0.048)	1.1% (0.74-1.5%)	0.52% (0.19-0.82%)	1.3 (0.47-2)	250 (160-340)
Mauritania	2000	Female	35 to 39	6.2% (3.5-9.1%)	94% (91-96%)	0.035 (0.019-0.056)	1% (0.68-1.3%)	0.72% (0.25-1.1%)	1.9 (0.63-2.9)	220 (150-300)
Mauritania	2000	Female	40 to 44	6.8% (4.2-9.8%)	93% (90-96%)	0.037 (0.02-0.06)	1.1% (0.73-1.6%)	1.2% (0.56-1.8%)	3.4 (1.6-5.3)	240 (140-330)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	2000	Female	45 to 49	6.4% (3.8-9.2%)	94% (91-96%)	0.034 (0.017-0.056)	1% (0.66-1.4%)	1.1% (0.61-1.7%)	3.4 (1.8-5.3)	190 (120-280)
Mauritania	2000	Female	50 to 54	6.2% (4-9.2%)	94% (91-96%)	0.034 (0.018-0.059)	1.2% (0.75-1.7%)	1.5% (0.81-2.2%)	5.7 (3.1-8.9)	240 (150-360)
Mauritania	2000	Female	55 to 59	6.8% (4.2-10%)	93% (90-96%)	0.041 (0.021-0.071)	0.91% (0.57-1.3%)	1.1% (0.62-1.6%)	5.2 (2.9-8.1)	190 (110-280)
Mauritania	2000	Female	60 to 64	7.1% (4.3-11%)	93% (89-96%)	0.044 (0.021-0.081)	0.98% (0.59-1.4%)	1.2% (0.68-1.7%)	7.1 (4.1-11)	200 (120-310)
Mauritania	2000	Female	65 to 69	7.5% (4.8-10%)	93% (90-95%)	0.041 (0.021-0.068)	0.81% (0.47-1.2%)	0.94% (0.53-1.4%)	6.9 (3.9-11)	160 (95-250)
Mauritania	2000	Female	70 to 74	7% (4.6-10%)	93% (90-95%)	0.034 (0.017-0.06)	0.66% (0.35-1%)	0.75% (0.37-1.2%)	6.1 (3-10)	120 (62-190)
Mauritania	2000	Female	75 to 79	5.9% (3.5-9.2%)	94% (91-96%)	0.026 (0.012-0.05)	0.56% (0.31-0.84%)	0.63% (0.34-0.98%)	4.7 (2.5-7.4)	69 (38-110)
Mauritania	2000	Female	80 to 84	5.4% (3.2-8.1%)	95% (92-97%)	0.023 (0.011-0.041)	0.34% (0.15-0.53%)	0.37% (0.14-0.61%)	1.8 (0.67-2.9)	20 (8.5-31)
Mauritania	2000	Female	85 to 89	5.7% (3.3-8.7%)	94% (91-97%)	0.023 (0.011-0.041)	0.19% (0.041-0.33%)	0.2% (0.013-0.37%)	0.54 (0.035-1)	4.5 (0.98-7.9)
Mauritania	2000	Female	90 to 94	6.4% (3.9-9.8%)	94% (90-96%)	0.025 (0.011-0.048)	0.1% (-0.05-0.23%)	0.078% (-0.11-0.24%)	0.079 (-0.11-0.24)	0.62 (-0.31-1.4)
Mauritania	2000	Female	95 plus	7.4% (4.2-12%)	93% (88-96%)	0.028 (0.011-0.058)	0.12% (-0.069-0.28%)	0.074% (-0.16-0.26%)	0.02 (-0.043-0.072)	0.13 (-0.074-0.29)
Mauritania	2000	Male	All Ages	3% (2.5-3.7%)	97% (96-98%)	0.053 (0.045-0.062)	0.57% (0.38-0.74%)	0.76% (0.43-1.1%)	72 (40-100)	3800 (2500-5000)
Mauritania	2000	Male	15 to 19	2.3% (1.4-3.5%)	98% (96-99%)	0.031 (0.015-0.053)	0.62% (0.41-0.92%)	0.2% (0.12-0.32%)	0.38 (0.22-0.6)	160 (96-230)
Mauritania	2000	Male	20 to 24	3.7% (2.1-5.6%)	96% (94-98%)	0.066 (0.033-0.11)	1.4% (0.94-1.9%)	0.52% (0.29-0.79%)	1.2 (0.69-1.9)	360 (230-510)
Mauritania	2000	Male	25 to 29	4.5% (2.6-7.1%)	95% (93-97%)	0.096 (0.053-0.15)	1.7% (1.2-2.3%)	0.95% (0.22-1.5%)	2.3 (0.5-3.7)	410 (270-560)
Mauritania	2000	Male	30 to 34	5.1% (3-8.1%)	95% (92-97%)	0.12 (0.067-0.19)	1.9% (1.1-2.6%)	1.5% (0.21-2.5%)	3.6 (0.49-6.1)	420 (250-580)
Mauritania	2000	Male	35 to 39	5.5% (3.2-8.3%)	94% (92-97%)	0.12 (0.069-0.19)	2.1% (0.95-3.1%)	2.2% (0.29-3.8%)	5.5 (0.69-9.5)	430 (190-620)
Mauritania	2000	Male	40 to 44	5.5% (3.3-8.3%)	95% (92-97%)	0.12 (0.063-0.2)	2.2% (0.77-3.2%)	2.7% (0.34-4.3%)	6.9 (0.86-11)	410 (140-610)
Mauritania	2000	Male	45 to 49	5.7% (3.2-8.3%)	94% (92-97%)	0.13 (0.066-0.23)	2.4% (0.75-3.6%)	3% (0.38-4.7%)	8.1 (1-13)	400 (120-600)
Mauritania	2000	Male	50 to 54	5.9% (3.6-8.9%)	94% (91-96%)	0.14 (0.067-0.24)	2.2% (1.2-3.2%)	2.7% (1.1-4.1%)	7.4 (3.2-12)	310 (170-470)
Mauritania	2000	Male	55 to 59	5.9% (3.6-9.2%)	94% (91-96%)	0.13 (0.064-0.22)	2.1% (1.3-3%)	2.5% (1.5-3.9%)	8.1 (4.8-13)	290 (180-430)
Mauritania	2000	Male	60 to 64	5.9% (3.6-9.2%)	94% (91-96%)	0.12 (0.056-0.23)	1.7% (1.2-2.5%)	2.1% (1.3-3.2%)	8.4 (5.1-13)	250 (160-370)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	2000	Male	65 to 69	6% (3.6-8.8%)	94% (91-96%)	0.11 (0.054-0.18)	1.3% (0.86-1.8%)	1.5% (1-2.3%)	7.2 (4.7-11)	170 (120-250)
Mauritania	2000	Male	70 to 74	5.7% (3.6-8.3%)	94% (92-96%)	0.086 (0.042-0.15)	0.93% (0.61-1.4%)	1.1% (0.73-1.7%)	5.9 (3.7-8.7)	110 (73-160)
Mauritania	2000	Male	75 to 79	5.3% (3-8.4%)	95% (92-97%)	0.067 (0.031-0.13)	0.68% (0.45-1%)	0.8% (0.51-1.2%)	3.7 (2.3-5.6)	55 (35-80)
Mauritania	2000	Male	80 to 84	4.6% (2.6-7.1%)	95% (93-97%)	0.052 (0.024-0.091)	0.59% (0.38-0.85%)	0.68% (0.42-1%)	1.9 (1.2-2.9)	21 (13-30)
Mauritania	2000	Male	85 to 89	4.2% (2.3-6.6%)	96% (93-98%)	0.04 (0.02-0.072)	0.47% (0.31-0.69%)	0.52% (0.31-0.79%)	0.88 (0.53-1.3)	7.1 (4.6-10)
Mauritania	2000	Male	90 to 94	4.2% (2.5-6.6%)	96% (93-98%)	0.035 (0.016-0.064)	0.31% (0.17-0.45%)	0.3% (0.13-0.49%)	0.2 (0.086-0.32)	1.2 (0.68-1.8)
Mauritania	2000	Male	95 plus	4.8% (2.6-7.9%)	95% (92-97%)	0.032 (0.012-0.067)	0.32% (0.16-0.52%)	0.29% (0.089-0.52%)	0.044 (0.014-0.08)	0.2 (0.1-0.33)
Mauritania	2005	Female	All Ages	4% (3.3-4.8%)	96% (95-97%)	0.012 (0.0092-0.014)	0.42% (0.33-0.51%)	0.49% (0.34-0.62%)	52 (36-69)	2900 (2200-3600)
Mauritania	2005	Female	15 to 19	3.8% (2.3-5.8%)	96% (94-98%)	0.01 (0.0052-0.018)	0.49% (0.29-0.74%)	0.045% (0.017-0.077%)	0.096 (0.035-0.17)	140 (82-230)
Mauritania	2005	Female	20 to 24	4.9% (2.9-7.2%)	95% (93-97%)	0.017 (0.0088-0.028)	1.1% (0.7-1.6%)	0.1% (0.042-0.17%)	0.25 (0.11-0.43)	320 (190-500)
Mauritania	2005	Female	25 to 29	5.1% (2.9-7.9%)	95% (92-97%)	0.019 (0.01-0.03)	1.2% (0.82-1.6%)	0.27% (0.12-0.44%)	0.67 (0.3-1.1)	320 (210-470)
Mauritania	2005	Female	30 to 34	5.3% (3.1-8.5%)	95% (92-97%)	0.02 (0.011-0.033)	1.1% (0.78-1.5%)	0.52% (0.23-0.82%)	1.3 (0.57-2.1)	290 (190-410)
Mauritania	2005	Female	35 to 39	6.2% (3.5-9.2%)	94% (91-97%)	0.023 (0.012-0.037)	1% (0.73-1.3%)	0.72% (0.26-1.1%)	2 (0.73-3.1)	250 (170-340)
Mauritania	2005	Female	40 to 44	6.7% (4.1-9.8%)	93% (90-96%)	0.024 (0.012-0.04)	1.1% (0.73-1.6%)	1.2% (0.5-1.8%)	3.6 (1.6-5.7)	260 (170-370)
Mauritania	2005	Female	45 to 49	6.3% (3.6-9.1%)	94% (91-96%)	0.022 (0.011-0.038)	1% (0.69-1.4%)	1.1% (0.64-1.6%)	3.8 (2.1-5.8)	220 (150-310)
Mauritania	2005	Female	50 to 54	6.1% (3.8-9%)	94% (91-96%)	0.023 (0.012-0.038)	1.2% (0.79-1.7%)	1.5% (0.87-2.2%)	5.8 (3.1-8.9)	250 (160-360)
Mauritania	2005	Female	55 to 59	6.6% (4.1-9.8%)	93% (90-96%)	0.027 (0.014-0.047)	0.91% (0.6-1.3%)	1.1% (0.65-1.6%)	5.3 (3.1-8)	190 (120-280)
Mauritania	2005	Female	60 to 64	7% (4.3-10%)	93% (90-96%)	0.029 (0.014-0.052)	0.97% (0.63-1.4%)	1.2% (0.73-1.7%)	7.2 (4.2-11)	210 (130-300)
Mauritania	2005	Female	65 to 69	7.3% (4.7-10%)	93% (90-95%)	0.027 (0.013-0.046)	0.82% (0.53-1.2%)	0.96% (0.6-1.4%)	7.2 (4.2-11)	170 (110-260)
Mauritania	2005	Female	70 to 74	6.9% (4.5-10%)	93% (90-96%)	0.022 (0.011-0.04)	0.68% (0.41-1%)	0.78% (0.44-1.2%)	6.6 (3.6-10)	130 (75-190)
Mauritania	2005	Female	75 to 79	5.9% (3.6-9.3%)	94% (91-96%)	0.017 (0.0081-0.033)	0.58% (0.34-0.86%)	0.67% (0.37-1%)	5.2 (2.9-8.4)	78 (46-120)
Mauritania	2005	Female	80 to 84	5.4% (3.2-7.9%)	95% (92-97%)	0.015 (0.0073-0.027)	0.36% (0.2-0.55%)	0.4% (0.21-0.64%)	2.1 (1.1-3.2)	23 (13-35)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	2005	Female	85 to 89	5.7% (3.4-8.8%)	94% (91-97%)	0.015 (0.0075-0.027)	0.23% (0.11-0.35%)	0.24% (0.099-0.39%)	0.77 (0.31-1.3)	6.4 (3.1-10)
Mauritania	2005	Female	90 to 94	6.4% (3.8-10%)	94% (90-96%)	0.017 (0.0074-0.032)	0.14% (0.034-0.25%)	0.13% (-0.016-0.26%)	0.15 (-0.019-0.31)	1 (0.24-1.8)
Mauritania	2005	Female	95 plus	7.5% (4.1-12%)	93% (88-96%)	0.019 (0.007-0.039)	0.17% (0.023-0.29%)	0.12% (-0.041-0.28%)	0.04 (-0.013-0.089)	0.21 (0.03-0.38)
Mauritania	2005	Male	All Ages	3.1% (2.5-3.8%)	97% (96-97%)	0.037 (0.03-0.045)	0.56% (0.38-0.74%)	0.73% (0.41-1.1%)	72 (40-110)	4000 (2600-5500)
Mauritania	2005	Male	15 to 19	2.3% (1.3-3.5%)	98% (96-99%)	0.021 (0.0098-0.037)	0.64% (0.41-0.92%)	0.17% (0.093-0.27%)	0.33 (0.17-0.54)	170 (110-250)
Mauritania	2005	Male	20 to 24	3.7% (2.1-5.6%)	96% (94-98%)	0.044 (0.022-0.075)	1.4% (0.94-2%)	0.44% (0.23-0.67%)	1.1 (0.56-1.8)	390 (250-580)
Mauritania	2005	Male	25 to 29	4.5% (2.6-7%)	95% (93-97%)	0.065 (0.035-0.1)	1.6% (1.1-2.2%)	0.79% (0.13-1.4%)	2 (0.33-3.5)	440 (280-610)
Mauritania	2005	Male	30 to 34	5.1% (3-8.3%)	95% (92-97%)	0.079 (0.044-0.13)	1.8% (1.1-2.5%)	1.3% (0.18-2.4%)	3.3 (0.48-6.1)	440 (260-630)
Mauritania	2005	Male	35 to 39	5.5% (3.2-8.3%)	94% (92-97%)	0.084 (0.045-0.13)	2% (0.93-2.9%)	2% (0.27-3.6%)	5.3 (0.73-9.4)	450 (200-660)
Mauritania	2005	Male	40 to 44	5.5% (3.4-8.2%)	95% (92-97%)	0.083 (0.041-0.13)	2% (0.78-3.1%)	2.4% (0.32-4.2%)	6.7 (0.85-12)	420 (160-650)
Mauritania	2005	Male	45 to 49	5.7% (3-8.6%)	94% (91-97%)	0.087 (0.044-0.15)	2.2% (0.76-3.4%)	2.8% (0.38-4.6%)	8.5 (1.1-14)	430 (150-680)
Mauritania	2005	Male	50 to 54	6% (3.6-9%)	94% (91-96%)	0.092 (0.047-0.16)	2% (1.1-3%)	2.5% (1.1-4.1%)	8.2 (3.4-13)	350 (190-530)
Mauritania	2005	Male	55 to 59	5.9% (3.6-9.2%)	94% (91-96%)	0.089 (0.043-0.15)	1.9% (1.3-2.9%)	2.4% (1.4-3.8%)	8 (4.6-13)	290 (180-440)
Mauritania	2005	Male	60 to 64	5.8% (3.4-8.9%)	94% (91-97%)	0.081 (0.037-0.15)	1.6% (1.1-2.4%)	2% (1.3-3.1%)	8.1 (4.9-13)	240 (160-370)
Mauritania	2005	Male	65 to 69	5.9% (3.5-8.7%)	94% (91-97%)	0.07 (0.035-0.12)	1.2% (0.79-1.7%)	1.4% (0.94-2.2%)	6.7 (4.2-10)	160 (110-240)
Mauritania	2005	Male	70 to 74	5.7% (3.7-8.4%)	94% (92-96%)	0.057 (0.029-0.1)	0.9% (0.58-1.3%)	1.1% (0.71-1.6%)	5.9 (3.6-8.8)	110 (70-160)
Mauritania	2005	Male	75 to 79	5.3% (3.1-8.3%)	95% (92-97%)	0.045 (0.021-0.083)	0.67% (0.45-0.98%)	0.8% (0.52-1.2%)	3.9 (2.4-6)	58 (37-84)
Mauritania	2005	Male	80 to 84	4.6% (2.6-7%)	95% (93-97%)	0.035 (0.016-0.061)	0.59% (0.39-0.85%)	0.69% (0.44-1%)	2.1 (1.3-3.2)	23 (16-34)
Mauritania	2005	Male	85 to 89	4.2% (2.4-6.8%)	96% (93-98%)	0.027 (0.013-0.049)	0.48% (0.33-0.67%)	0.54% (0.35-0.78%)	1 (0.66-1.5)	8.3 (5.6-12)
Mauritania	2005	Male	90 to 94	4.2% (2.5-6.9%)	96% (93-98%)	0.023 (0.011-0.044)	0.32% (0.2-0.47%)	0.33% (0.17-0.51%)	0.25 (0.13-0.4)	1.5 (0.95-2.2)
Mauritania	2005	Male	95 plus	4.8% (2.6-7.9%)	95% (92-97%)	0.021 (0.008-0.045)	0.34% (0.21-0.52%)	0.31% (0.14-0.52%)	0.059 (0.027-0.1)	0.27 (0.16-0.41)
Mauritania	2010	Female	All Ages	4.4% (3.6-5.2%)	96% (95-96%)	0.012 (0.0078-0.016)	0.48% (0.38-0.59%)	0.53% (0.38-0.68%)	55 (37-74)	3200 (2400-4000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	2010	Female	15 to 19	3.9% (2.5-5.9%)	96% (94-98%)	0.0096 (0.0043-0.018)	0.54% (0.33-0.82%)	0.05% (0.022-0.081%)	0.094 (0.04-0.16)	160 (91-250)
Mauritania	2010	Female	20 to 24	5% (2.9-7.6%)	95% (92-97%)	0.016 (0.0072-0.029)	1.2% (0.76-1.8%)	0.11% (0.047-0.19%)	0.25 (0.1-0.42)	360 (210-550)
Mauritania	2010	Female	25 to 29	5.6% (3.3-8.5%)	94% (91-97%)	0.019 (0.0091-0.032)	1.3% (0.9-1.8%)	0.29% (0.098-0.48%)	0.65 (0.22-1.1)	360 (240-530)
Mauritania	2010	Female	30 to 34	6.2% (3.6-9.8%)	94% (90-96%)	0.021 (0.011-0.037)	1.2% (0.85-1.6%)	0.56% (0.25-0.87%)	1.3 (0.55-2.1)	330 (220-460)
Mauritania	2010	Female	35 to 39	6.8% (3.9-10%)	93% (90-96%)	0.023 (0.011-0.039)	1.1% (0.77-1.4%)	0.78% (0.35-1.2%)	2.1 (0.91-3.2)	290 (200-380)
Mauritania	2010	Female	40 to 44	7.1% (4.3-10%)	93% (90-96%)	0.023 (0.011-0.04)	1.2% (0.78-1.6%)	1.3% (0.63-1.9%)	3.9 (1.9-6.1)	300 (190-410)
Mauritania	2010	Female	45 to 49	7.1% (4-10%)	93% (90-96%)	0.022 (0.01-0.041)	1% (0.72-1.4%)	1.2% (0.71-1.8%)	4.3 (2.4-6.7)	250 (170-360)
Mauritania	2010	Female	50 to 54	7.1% (4.5-11%)	93% (89-96%)	0.023 (0.011-0.043)	1.2% (0.81-1.6%)	1.5% (0.92-2.1%)	6.5 (3.7-9.9)	280 (180-400)
Mauritania	2010	Female	55 to 59	7.4% (4.5-11%)	93% (89-96%)	0.027 (0.012-0.05)	0.9% (0.59-1.3%)	1.1% (0.64-1.7%)	5.4 (3.1-8.5)	200 (120-290)
Mauritania	2010	Female	60 to 64	7.8% (4.7-12%)	92% (88-95%)	0.029 (0.013-0.055)	0.94% (0.61-1.3%)	1.2% (0.72-1.7%)	7.3 (4.2-11)	220 (130-320)
Mauritania	2010	Female	65 to 69	8% (5-11%)	92% (89-95%)	0.026 (0.012-0.048)	0.8% (0.52-1.1%)	0.95% (0.59-1.4%)	7.3 (4.3-11)	180 (110-260)
Mauritania	2010	Female	70 to 74	7.3% (4.7-10%)	93% (90-95%)	0.021 (0.0094-0.039)	0.67% (0.41-1%)	0.77% (0.44-1.2%)	6.8 (3.6-11)	130 (76-210)
Mauritania	2010	Female	75 to 79	6.1% (3.7-9.4%)	94% (91-96%)	0.016 (0.0065-0.031)	0.58% (0.35-0.84%)	0.67% (0.39-1%)	5.6 (3.1-8.9)	84 (49-130)
Mauritania	2010	Female	80 to 84	5.5% (3.1-8%)	94% (92-97%)	0.014 (0.0062-0.026)	0.37% (0.21-0.55%)	0.41% (0.21-0.64%)	2.3 (1.1-3.6)	26 (14-39)
Mauritania	2010	Female	85 to 89	5.8% (3.4-8.9%)	94% (91-97%)	0.014 (0.0064-0.027)	0.24% (0.12-0.36%)	0.26% (0.1-0.4%)	0.96 (0.39-1.6)	8 (4-12)
Mauritania	2010	Female	90 to 94	6.4% (4-10%)	94% (90-96%)	0.015 (0.0067-0.03)	0.15% (0.039-0.25%)	0.14% (0.0052-0.27%)	0.22 (0.0088-0.41)	1.4 (0.4-2.4)
Mauritania	2010	Female	95 plus	7.6% (4.3-12%)	92% (88-96%)	0.017 (0.0062-0.038)	0.18% (0.043-0.3%)	0.14% (-0.022-0.29%)	0.057 (-0.0094-0.12)	0.3 (0.07-0.5)
Mauritania	2010	Male	All Ages	3.2% (2.6-3.9%)	97% (96-97%)	0.035 (0.023-0.048)	0.65% (0.45-0.86%)	0.82% (0.49-1.2%)	78 (44-120)	4500 (3000-6100)
Mauritania	2010	Male	15 to 19	2.3% (1.4-3.5%)	98% (96-99%)	0.019 (0.0084-0.036)	0.68% (0.43-1%)	0.17% (0.09-0.27%)	0.31 (0.15-0.51)	190 (110-310)
Mauritania	2010	Male	20 to 24	3.7% (2.1-5.6%)	96% (94-98%)	0.04 (0.018-0.073)	1.5% (1-2.2%)	0.45% (0.24-0.7%)	1.1 (0.53-1.7)	430 (280-650)
Mauritania	2010	Male	25 to 29	4.5% (2.6-7%)	95% (93-97%)	0.059 (0.028-0.1)	1.8% (1.2-2.4%)	0.82% (0.1-1.5%)	2 (0.25-3.6)	480 (300-670)
Mauritania	2010	Male	30 to 34	5.1% (3-8.4%)	95% (92-97%)	0.071 (0.036-0.12)	1.9% (1.2-2.6%)	1.4% (0.17-2.5%)	3.4 (0.42-6.4)	490 (280-700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	2010	Male	35 to 39	5.7% (3.3-8.5%)	94% (91-97%)	0.078 (0.04-0.14)	2.1% (1.1-3%)	2.1% (0.31-3.8%)	5.8 (0.83-10)	500 (260-750)
Mauritania	2010	Male	40 to 44	5.9% (3.6-8.8%)	94% (91-96%)	0.081 (0.038-0.14)	2.1% (0.85-3.1%)	2.6% (0.36-4.3%)	7.5 (0.96-13)	480 (190-740)
Mauritania	2010	Male	45 to 49	5.9% (3.2-8.8%)	94% (91-97%)	0.083 (0.038-0.15)	2.3% (0.85-3.5%)	3% (0.47-5%)	9.5 (1.5-17)	490 (180-780)
Mauritania	2010	Male	50 to 54	5.9% (3.6-8.9%)	94% (91-96%)	0.083 (0.037-0.16)	2% (1.2-3%)	2.6% (1.4-4.2%)	9.5 (4.5-15)	410 (240-630)
Mauritania	2010	Male	55 to 59	5.9% (3.5-8.7%)	94% (91-96%)	0.08 (0.037-0.15)	1.9% (1.3-2.8%)	2.5% (1.5-3.8%)	9.7 (5.4-15)	360 (220-530)
Mauritania	2010	Male	60 to 64	5.8% (3.5-8.9%)	94% (91-97%)	0.073 (0.032-0.14)	1.6% (1.1-2.4%)	2% (1.3-3.1%)	8.6 (5.1-13)	260 (170-390)
Mauritania	2010	Male	65 to 69	5.9% (3.6-8.7%)	94% (91-96%)	0.064 (0.031-0.12)	1.2% (0.79-1.7%)	1.4% (0.95-2.1%)	6.9 (4.2-10)	170 (110-250)
Mauritania	2010	Male	70 to 74	5.7% (3.7-8.5%)	94% (91-96%)	0.052 (0.024-0.095)	0.89% (0.6-1.2%)	1.1% (0.74-1.6%)	6 (3.7-8.8)	110 (73-160)
Mauritania	2010	Male	75 to 79	5.2% (3.1-8.3%)	95% (92-97%)	0.041 (0.017-0.079)	0.66% (0.45-0.94%)	0.81% (0.52-1.2%)	4.1 (2.6-6.2)	61 (40-88)
Mauritania	2010	Male	80 to 84	4.6% (2.7-7%)	95% (93-97%)	0.032 (0.014-0.058)	0.59% (0.41-0.84%)	0.7% (0.47-1%)	2.4 (1.5-3.5)	26 (17-37)
Mauritania	2010	Male	85 to 89	4.2% (2.3-6.5%)	96% (94-98%)	0.024 (0.011-0.047)	0.48% (0.33-0.65%)	0.54% (0.34-0.76%)	1.2 (0.73-1.7)	9.6 (6.4-13)
Mauritania	2010	Male	90 to 94	4.2% (2.5-6.6%)	96% (93-98%)	0.021 (0.0092-0.041)	0.33% (0.21-0.45%)	0.33% (0.19-0.49%)	0.31 (0.17-0.47)	1.9 (1.2-2.7)
Mauritania	2010	Male	95 plus	4.7% (2.6-7.4%)	95% (93-97%)	0.019 (0.007-0.042)	0.35% (0.22-0.49%)	0.31% (0.15-0.49%)	0.074 (0.035-0.12)	0.34 (0.21-0.49)
Mauritania	2016	Female	All Ages	4.4% (3.7-5.2%)	96% (95-96%)	0.015 (0.0079-0.026)	0.58% (0.46-0.71%)	0.59% (0.43-0.76%)	62 (40-88)	3700 (2700-4700)
Mauritania	2016	Female	15 to 19	3.9% (2.4-5.8%)	96% (94-98%)	0.012 (0.0049-0.028)	0.62% (0.37-0.92%)	0.059% (0.03-0.095%)	0.1 (0.046-0.17)	190 (110-300)
Mauritania	2016	Female	20 to 24	5% (2.9-7.5%)	95% (93-97%)	0.02 (0.0076-0.041)	1.4% (0.86-2.1%)	0.13% (0.061-0.23%)	0.26 (0.12-0.45)	430 (250-670)
Mauritania	2016	Female	25 to 29	5.5% (3.2-8.6%)	94% (91-97%)	0.023 (0.0093-0.047)	1.5% (1-2%)	0.34% (0.15-0.55%)	0.7 (0.3-1.2)	420 (270-610)
Mauritania	2016	Female	30 to 34	6.1% (3.5-9.8%)	94% (90-96%)	0.026 (0.011-0.052)	1.3% (0.92-1.8%)	0.61% (0.23-0.94%)	1.4 (0.5-2.4)	370 (250-540)
Mauritania	2016	Female	35 to 39	6.7% (3.8-10%)	93% (90-96%)	0.029 (0.012-0.057)	1.2% (0.83-1.5%)	0.85% (0.44-1.3%)	2.4 (1.1-3.8)	340 (230-460)
Mauritania	2016	Female	40 to 44	7% (4.2-10%)	93% (90-96%)	0.029 (0.011-0.055)	1.2% (0.85-1.7%)	1.4% (0.8-2.1%)	4.6 (2.4-7.3)	360 (230-500)
Mauritania	2016	Female	45 to 49	6.9% (3.8-10%)	93% (90-96%)	0.028 (0.011-0.055)	1.1% (0.77-1.5%)	1.3% (0.77-1.9%)	5.1 (2.9-8.3)	310 (210-440)
Mauritania	2016	Female	50 to 54	7% (4.4-10%)	93% (90-96%)	0.029 (0.012-0.061)	1.2% (0.86-1.7%)	1.5% (0.98-2.2%)	7.6 (4.3-12)	340 (220-510)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	2016	Female	55 to 59	7.2% (4.4-11%)	93% (89-96%)	0.034 (0.014-0.072)	0.91% (0.59-1.3%)	1.1% (0.66-1.7%)	6.7 (3.5-11)	250 (150-390)
Mauritania	2016	Female	60 to 64	7.7% (4.6-12%)	92% (88-95%)	0.036 (0.014-0.08)	0.94% (0.59-1.4%)	1.2% (0.7-1.8%)	8.1 (4.2-13)	240 (140-370)
Mauritania	2016	Female	65 to 69	7.9% (5-11%)	92% (89-95%)	0.033 (0.014-0.071)	0.79% (0.49-1.1%)	0.94% (0.56-1.4%)	7.8 (4.1-12)	190 (110-290)
Mauritania	2016	Female	70 to 74	7.1% (4.6-10%)	93% (90-95%)	0.026 (0.01-0.056)	0.66% (0.36-1%)	0.76% (0.38-1.2%)	7.2 (3.4-12)	140 (75-220)
Mauritania	2016	Female	75 to 79	5.9% (3.5-9.2%)	94% (91-97%)	0.02 (0.007-0.044)	0.57% (0.33-0.84%)	0.67% (0.37-1%)	6.2 (3.2-10)	93 (52-140)
Mauritania	2016	Female	80 to 84	5.4% (3.1-7.9%)	95% (92-97%)	0.018 (0.0068-0.038)	0.35% (0.18-0.55%)	0.39% (0.17-0.64%)	2.4 (1-4.1)	28 (13-44)
Mauritania	2016	Female	85 to 89	5.6% (3.2-8.7%)	94% (91-97%)	0.018 (0.0068-0.038)	0.23% (0.076-0.36%)	0.24% (0.053-0.41%)	1.1 (0.24-1.8)	9.1 (3-14)
Mauritania	2016	Female	90 to 94	6.3% (3.8-9.9%)	94% (90-96%)	0.019 (0.007-0.043)	0.14% (-0.0074-0.25%)	0.12% (-0.061-0.26%)	0.24 (-0.14-0.52)	1.7 (-0.091-3.1)
Mauritania	2016	Female	95 plus	7.4% (4.1-12%)	93% (88-96%)	0.022 (0.0073-0.051)	0.17% (-0.017-0.3%)	0.11% (-0.11-0.28%)	0.067 (-0.065-0.17)	0.39 (-0.04-0.74)
Mauritania	2016	Male	All Ages	3.2% (2.7-3.9%)	97% (96-97%)	0.045 (0.024-0.077)	0.84% (0.59-1.1%)	1% (0.65-1.5%)	100 (57-150)	5700 (3900-7800)
Mauritania	2016	Male	15 to 19	2.3% (1.3-3.6%)	98% (96-99%)	0.024 (0.0093-0.054)	0.73% (0.47-1.1%)	0.19% (0.099-0.31%)	0.38 (0.19-0.65)	220 (130-350)
Mauritania	2016	Male	20 to 24	3.6% (2-5.6%)	96% (94-98%)	0.051 (0.02-0.11)	1.6% (1.1-2.4%)	0.52% (0.29-0.84%)	1.3 (0.69-2.3)	510 (330-770)
Mauritania	2016	Male	25 to 29	4.5% (2.6-6.9%)	96% (93-97%)	0.074 (0.03-0.14)	2% (1.4-2.6%)	0.98% (0.14-1.7%)	2.5 (0.37-4.6)	570 (380-820)
Mauritania	2016	Male	30 to 34	5.1% (2.9-8.1%)	95% (92-97%)	0.09 (0.039-0.17)	2.1% (1.3-2.9%)	1.6% (0.19-2.8%)	4.3 (0.49-8.1)	590 (340-840)
Mauritania	2016	Male	35 to 39	5.7% (3.2-8.7%)	94% (91-97%)	0.099 (0.041-0.2)	2.3% (1.2-3.2%)	2.4% (0.34-4.2%)	7.2 (0.9-13)	630 (320-930)
Mauritania	2016	Male	40 to 44	5.9% (3.5-8.7%)	94% (91-96%)	0.1 (0.04-0.2)	2.3% (1.2-3.4%)	2.9% (0.98-4.7%)	9.8 (2.9-16)	620 (320-930)
Mauritania	2016	Male	45 to 49	5.8% (3.2-8.6%)	94% (91-97%)	0.1 (0.041-0.22)	2.5% (1.5-3.6%)	3.3% (1.6-5.2%)	13 (5.9-21)	650 (350-980)
Mauritania	2016	Male	50 to 54	5.9% (3.6-8.8%)	94% (91-96%)	0.11 (0.041-0.22)	2.2% (1.4-3.1%)	2.8% (1.6-4.4%)	12 (6.3-20)	540 (310-810)
Mauritania	2016	Male	55 to 59	5.9% (3.5-8.7%)	94% (91-96%)	0.1 (0.041-0.21)	2.1% (1.4-3.1%)	2.7% (1.7-4.2%)	13 (7.5-22)	490 (290-750)
Mauritania	2016	Male	60 to 64	5.8% (3.4-9%)	94% (91-97%)	0.093 (0.037-0.21)	1.7% (1.1-2.5%)	2.2% (1.4-3.3%)	12 (7-20)	370 (230-570)
Mauritania	2016	Male	65 to 69	5.9% (3.5-8.7%)	94% (91-97%)	0.082 (0.034-0.18)	1.2% (0.82-1.7%)	1.5% (0.98-2.2%)	8.6 (5-14)	210 (130-320)
Mauritania	2016	Male	70 to 74	5.7% (3.6-8.4%)	94% (92-96%)	0.066 (0.026-0.14)	0.92% (0.62-1.3%)	1.2% (0.75-1.7%)	7 (4.2-11)	130 (84-200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Mauritania	2016	Male	75 to 79	5.2% (3-8.2%)	95% (92-97%)	0.052 (0.018-0.11)	0.68% (0.46-0.99%)	0.83% (0.53-1.2%)	4.7 (2.8-7.3)	70 (44-100)
Mauritania	2016	Male	80 to 84	4.5% (2.6-6.7%)	95% (93-97%)	0.039 (0.015-0.083)	0.61% (0.4-0.86%)	0.72% (0.45-1%)	2.7 (1.6-4.1)	30 (19-43)
Mauritania	2016	Male	85 to 89	4.1% (2.3-6.5%)	96% (94-98%)	0.031 (0.012-0.064)	0.48% (0.33-0.64%)	0.54% (0.34-0.76%)	1.4 (0.89-2)	12 (7.6-16)
Mauritania	2016	Male	90 to 94	4.1% (2.5-6.6%)	96% (93-97%)	0.026 (0.0099-0.058)	0.33% (0.2-0.46%)	0.33% (0.18-0.5%)	0.39 (0.21-0.61)	2.4 (1.5-3.4)
Mauritania	2016	Male	95 plus	4.7% (2.6-7.7%)	95% (92-97%)	0.024 (0.0083-0.057)	0.35% (0.21-0.5%)	0.31% (0.14-0.5%)	0.097 (0.04-0.16)	0.45 (0.26-0.66)
Niger	1990	Female	All Ages	23% (18-30%)	77% (70-82%)	0.094 (0.063-0.13)	0.1% (0.023-0.19%)	0.09% (-0.088-0.27%)	66 (-65-200)	6000 (1300-11000)
Niger	1990	Female	15 to 19	23% (12-37%)	77% (63-88%)	0.09 (0.033-0.18)	0.42% (0.19-0.7%)	0.075% (-0.23-0.42%)	0.6 (-1.8-3.4)	400 (160-680)
Niger	1990	Female	20 to 24	28% (14-45%)	72% (55-86%)	0.14 (0.054-0.26)	0.89% (0.36-1.5%)	0.15% (-0.47-0.82%)	1.4 (-4.3-7.5)	840 (310-1500)
Niger	1990	Female	25 to 29	30% (15-49%)	70% (51-85%)	0.17 (0.074-0.31)	0.94% (0.32-1.6%)	0.31% (-0.54-1.2%)	3 (-5-11)	840 (280-1500)
Niger	1990	Female	30 to 34	33% (17-52%)	67% (48-83%)	0.19 (0.086-0.33)	0.91% (0.26-1.6%)	0.49% (-0.47-1.5%)	4.7 (-4.5-14)	730 (190-1300)
Niger	1990	Female	35 to 39	36% (20-56%)	64% (44-80%)	0.2 (0.089-0.36)	0.88% (0.2-1.6%)	0.68% (-0.36-1.7%)	6.5 (-3.4-17)	640 (140-1200)
Niger	1990	Female	40 to 44	38% (22-57%)	62% (43-78%)	0.21 (0.088-0.4)	1% (0.28-1.8%)	1.1% (-0.049-2.3%)	12 (-0.47-24)	750 (200-1300)
Niger	1990	Female	45 to 49	39% (20-59%)	61% (41-80%)	0.22 (0.086-0.44)	0.86% (-0.00056-1.8%)	0.94% (-0.33-2.3%)	11 (-3.6-27)	590 (-0.38-1200)
Niger	1990	Female	50 to 54	40% (22-60%)	60% (40-78%)	0.24 (0.093-0.47)	0.91% (0.054-1.7%)	1.1% (-0.089-2.2%)	13 (-1.1-28)	560 (34-1100)
Niger	1990	Female	55 to 59	40% (24-59%)	60% (41-76%)	0.25 (0.099-0.5)	0.62% (-0.28-1.5%)	0.74% (-0.45-1.9%)	10 (-6-26)	350 (-150-850)
Niger	1990	Female	60 to 64	41% (23-60%)	59% (40-77%)	0.26 (0.097-0.52)	0.44% (-0.38-1.2%)	0.53% (-0.56-1.5%)	8.2 (-8.4-24)	230 (-200-660)
Niger	1990	Female	65 to 69	42% (25-60%)	58% (40-75%)	0.23 (0.09-0.47)	0.11% (-0.87-0.85%)	0.11% (-1.1-1.1%)	1.8 (-19-19)	52 (-410-410)
Niger	1990	Female	70 to 74	40% (24-58%)	60% (42-76%)	0.19 (0.069-0.45)	0.033% (-0.99-0.8%)	0.017% (-1.2-0.96%)	0.29 (-22-17)	13 (-380-310)
Niger	1990	Female	75 to 79	36% (19-55%)	64% (45-81%)	0.16 (0.046-0.36)	-0.13% (-1.1-0.53%)	-0.17% (-1.3-0.61%)	-2.4 (-18-8.3)	-30 (-250-120)
Niger	1990	Female	80 to 84	34% (18-53%)	66% (47-82%)	0.14 (0.044-0.3)	-0.22% (-1-0.31%)	-0.27% (-1.2-0.35%)	-1.8 (-8-2.3)	-17 (-82-25)
Niger	1990	Female	85 to 89	34% (19-52%)	66% (48-81%)	0.14 (0.044-0.3)	-0.31% (-1.1-0.17%)	-0.39% (-1.3-0.18%)	-1.3 (-4.3-0.59)	-8.6 (-30-4.6)
Niger	1990	Female	90 to 94	37% (21-55%)	63% (45-79%)	0.15 (0.042-0.32)	-0.4% (-1.1-0.1%)	-0.52% (-1.4-0.091%)	-0.56 (-1.6-0.1)	-2.6 (-7.4-0.67)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	1990	Female	95 plus	41% (22-62%)	59% (38-78%)	0.16 (0.038-0.38)	-0.4% (-1.2-0.097%)	-0.54% (-1.5-0.057%)	-0.17 (-0.47-0.018)	-0.48 (-1.5-0.12)
Niger	1990	Male	All Ages	19% (14-25%)	81% (75-86%)	0.11 (0.074-0.15)	0.21% (0.11-0.32%)	0.32% (0.12-0.54%)	250 (92-420)	13000 (7000-20000)
Niger	1990	Male	15 to 19	15% (7.8-26%)	85% (74-92%)	0.068 (0.024-0.14)	0.52% (0.3-0.81%)	0.23% (0.013-0.55%)	1.7 (0.093-4.2)	430 (240-680)
Niger	1990	Male	20 to 24	23% (12-37%)	77% (63-88%)	0.14 (0.056-0.28)	1.2% (0.65-1.9%)	0.58% (0.035-1.3%)	4.7 (0.27-10)	860 (450-1400)
Niger	1990	Male	25 to 29	28% (14-45%)	72% (55-86%)	0.21 (0.085-0.38)	1.6% (0.75-2.5%)	1.1% (-0.02-2.3%)	12 (-0.26-26)	1500 (750-2400)
Niger	1990	Male	30 to 34	32% (16-51%)	68% (49-84%)	0.25 (0.12-0.46)	1.9% (0.88-3.1%)	1.7% (0.34-3.2%)	18 (3.6-35)	1600 (740-2700)
Niger	1990	Male	35 to 39	35% (19-54%)	65% (46-81%)	0.28 (0.12-0.54)	2.3% (1.2-3.5%)	2.3% (0.88-4.1%)	25 (9.2-45)	1700 (900-2700)
Niger	1990	Male	40 to 44	36% (20-55%)	64% (45-80%)	0.3 (0.12-0.59)	2.3% (1.1-3.8%)	2.7% (0.98-4.7%)	29 (11-52)	1600 (760-2600)
Niger	1990	Male	45 to 49	36% (20-55%)	64% (45-80%)	0.32 (0.12-0.67)	2.4% (1.2-4.1%)	2.9% (1.2-5%)	34 (14-58)	1600 (790-2600)
Niger	1990	Male	50 to 54	37% (21-57%)	63% (43-79%)	0.32 (0.13-0.66)	2% (0.8-3.5%)	2.4% (0.75-4.4%)	29 (9-56)	1200 (440-2100)
Niger	1990	Male	55 to 59	37% (21-57%)	63% (43-79%)	0.31 (0.12-0.62)	1.8% (0.75-3.1%)	2.1% (0.74-3.8%)	29 (9.8-53)	1000 (430-1800)
Niger	1990	Male	60 to 64	37% (20-57%)	63% (43-80%)	0.28 (0.097-0.61)	1.5% (0.57-2.6%)	1.7% (0.54-3%)	27 (8.7-50)	800 (300-1400)
Niger	1990	Male	65 to 69	37% (21-55%)	63% (45-79%)	0.23 (0.086-0.51)	0.98% (0.19-1.8%)	1.1% (0.093-2.2%)	19 (1.6-36)	440 (80-840)
Niger	1990	Male	70 to 74	35% (21-52%)	65% (48-79%)	0.18 (0.058-0.42)	0.67% (-0.12-1.4%)	0.76% (-0.18-1.6%)	11 (-2.7-24)	210 (-37-440)
Niger	1990	Male	75 to 79	32% (17-51%)	68% (49-83%)	0.14 (0.039-0.31)	0.45% (-0.2-1%)	0.49% (-0.28-1.2%)	5.4 (-3.2-13)	80 (-37-180)
Niger	1990	Male	80 to 84	29% (15-47%)	71% (53-85%)	0.11 (0.031-0.24)	0.37% (-0.2-0.82%)	0.4% (-0.28-0.94%)	2.1 (-1.5-5)	24 (-13-53)
Niger	1990	Male	85 to 89	28% (15-45%)	72% (55-85%)	0.095 (0.029-0.22)	0.27% (-0.19-0.64%)	0.27% (-0.28-0.72%)	0.64 (-0.67-1.8)	5.6 (-3.8-13)
Niger	1990	Male	90 to 94	30% (15-48%)	70% (52-85%)	0.09 (0.024-0.22)	0.11% (-0.28-0.39%)	0.067% (-0.42-0.42%)	0.045 (-0.28-0.28)	0.46 (-1.1-1.6)
Niger	1990	Male	95 plus	34% (16-56%)	66% (44-84%)	0.092 (0.019-0.24)	0.086% (-0.4-0.38%)	-0.006% (-0.63-0.36%)	-0.00074 (-0.077-0.044)	0.042 (-0.2-0.18)
Niger	1995	Female	All Ages	24% (18-30%)	76% (70-82%)	0.092 (0.062-0.13)	0.11% (0.014-0.2%)	0.088% (-0.12-0.3%)	72 (-100-240)	6800 (860-13000)
Niger	1995	Female	15 to 19	23% (12-37%)	77% (63-88%)	0.087 (0.033-0.18)	0.4% (0.16-0.68%)	0.07% (-0.26-0.42%)	0.7 (-2.5-4.4)	460 (180-820)
Niger	1995	Female	20 to 24	28% (15-45%)	72% (55-85%)	0.14 (0.055-0.26)	0.83% (0.29-1.4%)	0.14% (-0.52-0.82%)	1.6 (-6.1-10)	980 (340-1700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	1995	Female	25 to 29	31% (16-49%)	69% (51-84%)	0.17 (0.07-0.32)	0.87% (0.28-1.5%)	0.28% (-0.52-1.1%)	3.4 (-6.5-14)	960 (300-1700)
Niger	1995	Female	30 to 34	34% (17-52%)	66% (48-83%)	0.18 (0.083-0.33)	0.84% (0.17-1.6%)	0.43% (-0.53-1.4%)	5.4 (-6.3-17)	870 (180-1600)
Niger	1995	Female	35 to 39	37% (20-57%)	63% (43-80%)	0.2 (0.089-0.36)	0.82% (0.13-1.5%)	0.59% (-0.41-1.6%)	7.5 (-5.4-21)	760 (120-1400)
Niger	1995	Female	40 to 44	39% (22-58%)	61% (42-78%)	0.21 (0.086-0.39)	0.95% (0.22-1.7%)	0.96% (-0.068-2.1%)	12 (-0.84-25)	770 (180-1400)
Niger	1995	Female	45 to 49	39% (21-58%)	61% (42-79%)	0.22 (0.086-0.43)	0.79% (-0.15-1.7%)	0.84% (-0.54-2.2%)	12 (-7.7-31)	640 (-120-1400)
Niger	1995	Female	50 to 54	39% (22-60%)	61% (40-78%)	0.22 (0.09-0.45)	0.85% (-0.041-1.7%)	0.99% (-0.23-2.1%)	16 (-3.6-35)	670 (-30-1300)
Niger	1995	Female	55 to 59	40% (23-59%)	60% (41-77%)	0.24 (0.096-0.47)	0.56% (-0.31-1.4%)	0.65% (-0.48-1.8%)	11 (-8.4-31)	400 (-220-1000)
Niger	1995	Female	60 to 64	41% (23-60%)	59% (40-77%)	0.24 (0.092-0.5)	0.4% (-0.43-1.2%)	0.47% (-0.61-1.4%)	9.1 (-12-27)	260 (-280-760)
Niger	1995	Female	65 to 69	42% (25-60%)	58% (40-75%)	0.22 (0.088-0.45)	0.091% (-0.86-0.82%)	0.082% (-1.1-0.99%)	1.7 (-23-20)	53 (-490-460)
Niger	1995	Female	70 to 74	40% (24-57%)	60% (43-76%)	0.18 (0.066-0.43)	0.019% (-0.98-0.77%)	-0.0042% (-1.2-0.94%)	-0.11 (-28-20)	9.4 (-480-370)
Niger	1995	Female	75 to 79	36% (19-56%)	64% (44-81%)	0.15 (0.047-0.36)	-0.1% (-1.1-0.56%)	-0.14% (-1.3-0.64%)	-2.6 (-24-11)	-30 (-320-160)
Niger	1995	Female	80 to 84	34% (18-53%)	66% (47-82%)	0.13 (0.041-0.29)	-0.21% (-1.1-0.33%)	-0.26% (-1.3-0.38%)	-2.4 (-12-3.4)	-23 (-120-37)
Niger	1995	Female	85 to 89	34% (18-52%)	66% (48-82%)	0.13 (0.042-0.3)	-0.31% (-1.1-0.17%)	-0.4% (-1.4-0.2%)	-1.9 (-6.7-0.92)	-13 (-47-7.1)
Niger	1995	Female	90 to 94	37% (20-54%)	63% (46-80%)	0.14 (0.041-0.32)	-0.39% (-1.2-0.11%)	-0.52% (-1.5-0.096%)	-0.79 (-2.3-0.15)	-3.6 (-11-1)
Niger	1995	Female	95 plus	41% (21-62%)	59% (38-79%)	0.15 (0.036-0.37)	-0.39% (-1.2-0.11%)	-0.54% (-1.6-0.069%)	-0.2 (-0.6-0.026)	-0.58 (-1.9-0.17)
Niger	1995	Male	All Ages	19% (15-25%)	81% (75-85%)	0.11 (0.074-0.15)	0.21% (0.11-0.32%)	0.31% (0.11-0.53%)	270 (92-460)	14000 (7400-21000)
Niger	1995	Male	15 to 19	15% (8-26%)	85% (74-92%)	0.07 (0.025-0.15)	0.53% (0.3-0.81%)	0.21% (-0.0042-0.55%)	2.1 (-0.04-5.4)	570 (320-900)
Niger	1995	Male	20 to 24	23% (12-38%)	77% (62-88%)	0.14 (0.055-0.29)	1.2% (0.64-1.8%)	0.55% (-0.001-1.3%)	4.8 (-0.0089-11)	950 (510-1500)
Niger	1995	Male	25 to 29	29% (15-46%)	71% (54-85%)	0.21 (0.086-0.4)	1.6% (0.77-2.4%)	0.99% (-0.033-2.1%)	8 (-0.26-17)	1100 (520-1700)
Niger	1995	Male	30 to 34	32% (17-51%)	68% (49-83%)	0.25 (0.11-0.45)	1.8% (0.84-2.8%)	1.5% (0.2-2.9%)	19 (2.7-37)	1800 (830-2900)
Niger	1995	Male	35 to 39	35% (19-55%)	65% (45-81%)	0.28 (0.12-0.52)	2.1% (1.1-3.2%)	2.1% (0.69-3.7%)	29 (9.1-52)	2000 (1000-3200)
Niger	1995	Male	40 to 44	36% (21-55%)	64% (45-79%)	0.3 (0.12-0.58)	2.2% (1-3.5%)	2.4% (0.87-4.3%)	34 (12-61)	1900 (880-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	1995	Male	45 to 49	37% (20-55%)	63% (45-80%)	0.31 (0.12-0.64)	2.2% (1.1-3.7%)	2.6% (0.96-4.6%)	36 (14-64)	1700 (810-2900)
Niger	1995	Male	50 to 54	38% (21-58%)	62% (42-79%)	0.32 (0.13-0.64)	1.9% (0.7-3.3%)	2.2% (0.64-4.2%)	33 (9.3-62)	1300 (480-2400)
Niger	1995	Male	55 to 59	38% (22-57%)	62% (43-78%)	0.31 (0.12-0.62)	1.7% (0.67-2.9%)	2% (0.67-3.6%)	30 (9.4-55)	1000 (410-1800)
Niger	1995	Male	60 to 64	38% (22-57%)	62% (43-78%)	0.28 (0.096-0.59)	1.4% (0.53-2.4%)	1.6% (0.47-2.9%)	27 (8.2-49)	790 (300-1400)
Niger	1995	Male	65 to 69	37% (22-55%)	63% (45-78%)	0.23 (0.087-0.49)	0.94% (0.14-1.7%)	1.1% (0.059-2.1%)	20 (1.2-39)	480 (71-900)
Niger	1995	Male	70 to 74	36% (21-53%)	64% (47-79%)	0.18 (0.06-0.41)	0.65% (-0.11-1.4%)	0.75% (-0.17-1.6%)	14 (-3.1-29)	260 (-41-540)
Niger	1995	Male	75 to 79	33% (18-52%)	67% (48-82%)	0.13 (0.041-0.31)	0.44% (-0.17-0.99%)	0.48% (-0.25-1.2%)	6.4 (-3.3-16)	96 (-36-220)
Niger	1995	Male	80 to 84	30% (16-49%)	70% (51-84%)	0.11 (0.03-0.25)	0.36% (-0.24-0.84%)	0.39% (-0.35-0.97%)	2.6 (-2.3-6.6)	30 (-19-69)
Niger	1995	Male	85 to 89	29% (15-46%)	71% (54-85%)	0.097 (0.029-0.22)	0.26% (-0.19-0.66%)	0.26% (-0.31-0.75%)	0.86 (-1-2.5)	7.6 (-5.5-19)
Niger	1995	Male	90 to 94	30% (15-49%)	70% (51-85%)	0.091 (0.024-0.23)	0.12% (-0.3-0.41%)	0.074% (-0.45-0.43%)	0.076 (-0.47-0.45)	0.75 (-1.9-2.5)
Niger	1995	Male	95 plus	35% (16-57%)	65% (43-84%)	0.094 (0.019-0.26)	0.11% (-0.43-0.43%)	0.017% (-0.66-0.43%)	0.0033 (-0.13-0.082)	0.083 (-0.34-0.34)
Niger	2000	Female	All Ages	23% (18-29%)	77% (71-82%)	0.11 (0.073-0.15)	0.12% (0.01-0.22%)	0.079% (-0.16-0.3%)	66 (-140-250)	7500 (670-14000)
Niger	2000	Female	15 to 19	22% (12-37%)	78% (63-88%)	0.1 (0.038-0.2)	0.44% (0.17-0.74%)	0.081% (-0.27-0.49%)	0.82 (-2.8-5)	540 (200-940)
Niger	2000	Female	20 to 24	28% (14-45%)	72% (55-86%)	0.16 (0.064-0.31)	0.87% (0.38-1.5%)	0.15% (-0.49-0.8%)	2 (-6.1-11)	1200 (460-2000)
Niger	2000	Female	25 to 29	30% (16-48%)	70% (52-84%)	0.2 (0.08-0.38)	0.87% (0.33-1.4%)	0.28% (-0.46-1%)	4.1 (-6.4-15)	1100 (430-2000)
Niger	2000	Female	30 to 34	33% (18-52%)	67% (48-82%)	0.22 (0.1-0.41)	0.81% (0.21-1.4%)	0.39% (-0.4-1.1%)	5.9 (-6-17)	980 (250-1700)
Niger	2000	Female	35 to 39	37% (21-56%)	63% (44-79%)	0.24 (0.11-0.43)	0.78% (0.16-1.4%)	0.55% (-0.32-1.4%)	8.5 (-5-22)	890 (170-1600)
Niger	2000	Female	40 to 44	39% (23-57%)	61% (43-77%)	0.25 (0.1-0.47)	0.88% (0.24-1.5%)	0.86% (-0.082-1.9%)	13 (-1.2-26)	870 (230-1500)
Niger	2000	Female	45 to 49	39% (21-58%)	61% (42-79%)	0.26 (0.1-0.51)	0.76% (-0.11-1.6%)	0.8% (-0.46-2.1%)	12 (-6-31)	650 (-96-1400)
Niger	2000	Female	50 to 54	40% (22-61%)	60% (39-78%)	0.27 (0.11-0.53)	0.8% (-0.091-1.7%)	0.94% (-0.34-2.2%)	16 (-6-38)	690 (-79-1500)
Niger	2000	Female	55 to 59	40% (23-59%)	60% (41-77%)	0.29 (0.12-0.58)	0.53% (-0.35-1.4%)	0.62% (-0.56-1.9%)	12 (-11-38)	440 (-300-1200)
Niger	2000	Female	60 to 64	41% (24-60%)	59% (40-76%)	0.29 (0.11-0.61)	0.32% (-0.58-1.1%)	0.38% (-0.75-1.4%)	8.4 (-17-32)	250 (-460-880)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	2000	Female	65 to 69	42% (25-59%)	58% (41-75%)	0.27 (0.1-0.53)	0.0079% (-0.96-0.77%)	-0.014% (-1.2-0.95%)	-0.37 (-29-23)	5.6 (-630-510)
Niger	2000	Female	70 to 74	40% (25-57%)	60% (43-75%)	0.22 (0.081-0.5)	-0.079% (-1.1-0.7%)	-0.12% (-1.3-0.81%)	-2.9 (-33-20)	-42 (-570-370)
Niger	2000	Female	75 to 79	35% (20-55%)	65% (45-80%)	0.18 (0.054-0.43)	-0.18% (-1.2-0.52%)	-0.23% (-1.5-0.61%)	-4.7 (-30-12)	-59 (-400-180)
Niger	2000	Female	80 to 84	33% (18-53%)	67% (47-82%)	0.16 (0.049-0.35)	-0.28% (-1.2-0.28%)	-0.35% (-1.4-0.31%)	-3.9 (-16-3.5)	-38 (-160-37)
Niger	2000	Female	85 to 89	34% (18-51%)	66% (49-82%)	0.16 (0.05-0.35)	-0.39% (-1.3-0.16%)	-0.49% (-1.6-0.16%)	-3.1 (-10-1.1)	-21 (-70-8.9)
Niger	2000	Female	90 to 94	37% (20-55%)	63% (45-80%)	0.17 (0.047-0.39)	-0.47% (-1.3-0.11%)	-0.62% (-1.7-0.1%)	-1.3 (-3.7-0.22)	-6.1 (-18-1.4)
Niger	2000	Female	95 plus	41% (21-62%)	59% (38-79%)	0.18 (0.044-0.45)	-0.45% (-1.4-0.13%)	-0.63% (-1.8-0.061%)	-0.33 (-0.9-0.032)	-0.92 (-2.8-0.27)
Niger	2000	Male	All Ages	20% (15-26%)	80% (74-85%)	0.14 (0.095-0.19)	0.26% (0.14-0.39%)	0.38% (0.14-0.65%)	340 (120-570)	18000 (9100-27000)
Niger	2000	Male	15 to 19	16% (8-27%)	84% (73-92%)	0.088 (0.031-0.18)	0.58% (0.34-0.91%)	0.26% (0.015-0.65%)	2.4 (0.16-6.1)	620 (340-1000)
Niger	2000	Male	20 to 24	24% (12-39%)	76% (61-88%)	0.18 (0.068-0.35)	1.3% (0.7-2%)	0.65% (0.026-1.5%)	6.8 (0.28-15)	1300 (660-2000)
Niger	2000	Male	25 to 29	29% (15-46%)	71% (54-85%)	0.26 (0.11-0.48)	1.6% (0.85-2.5%)	1.1% (0.078-2.2%)	12 (0.83-25)	1500 (800-2400)
Niger	2000	Male	30 to 34	33% (17-52%)	67% (48-83%)	0.31 (0.14-0.57)	1.8% (0.88-2.8%)	1.5% (0.28-2.9%)	20 (3.7-37)	1800 (880-2900)
Niger	2000	Male	35 to 39	36% (20-56%)	64% (44-80%)	0.34 (0.14-0.62)	2% (1-3.2%)	2.3% (0.71-3.6%)	40 (13-72)	2700 (1400-4300)
Niger	2000	Male	40 to 44	37% (22-56%)	63% (44-78%)	0.37 (0.15-0.69)	2.1% (0.96-3.5%)	2.4% (0.82-4.3%)	47 (15-83)	2600 (1200-4400)
Niger	2000	Male	45 to 49	38% (20-56%)	62% (44-80%)	0.38 (0.15-0.78)	2.2% (1-3.9%)	2.6% (0.94-4.8%)	49 (17-90)	2300 (1000-4000)
Niger	2000	Male	50 to 54	38% (21-58%)	62% (42-79%)	0.39 (0.16-0.78)	2% (0.78-3.6%)	2.3% (0.69-4.5%)	43 (13-82)	1700 (690-3200)
Niger	2000	Male	55 to 59	38% (22-58%)	62% (42-78%)	0.37 (0.14-0.75)	1.8% (0.68-3.1%)	2.1% (0.61-3.9%)	38 (11-72)	1300 (500-2400)
Niger	2000	Male	60 to 64	38% (21-57%)	62% (43-79%)	0.33 (0.12-0.72)	1.4% (0.51-2.4%)	1.6% (0.5-3%)	30 (10-58)	900 (340-1600)
Niger	2000	Male	65 to 69	38% (21-56%)	62% (44-79%)	0.28 (0.1-0.59)	0.92% (0.13-1.8%)	1.1% (0.042-2.1%)	21 (0.84-43)	520 (80-1000)
Niger	2000	Male	70 to 74	36% (21-54%)	64% (46-79%)	0.22 (0.075-0.49)	0.63% (-0.18-1.4%)	0.73% (-0.27-1.6%)	16 (-6-37)	300 (-83-670)
Niger	2000	Male	75 to 79	33% (18-51%)	67% (49-82%)	0.17 (0.049-0.38)	0.41% (-0.26-0.98%)	0.45% (-0.37-1.2%)	8 (-6.4-20)	120 (-77-290)
Niger	2000	Male	80 to 84	31% (16-49%)	69% (51-84%)	0.14 (0.039-0.31)	0.34% (-0.25-0.85%)	0.36% (-0.36-0.97%)	3.2 (-3.2-8.8)	37 (-27-91)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	2000	Male	85 to 89	29% (15-47%)	71% (53-85%)	0.12 (0.036-0.27)	0.24% (-0.28-0.67%)	0.23% (-0.41-0.75%)	1 (-1.8-3.3)	9.1 (-11-25)
Niger	2000	Male	90 to 94	31% (17-49%)	69% (51-83%)	0.12 (0.031-0.29)	0.097% (-0.38-0.46%)	0.043% (-0.54-0.48%)	0.063 (-0.79-0.71)	0.86 (-3.4-4)
Niger	2000	Male	95 plus	36% (18-58%)	64% (42-82%)	0.12 (0.024-0.32)	0.095% (-0.51-0.51%)	-0.0017% (-0.75-0.53%)	-0.00048 (-0.23-0.16)	0.11 (-0.6-0.62)
Niger	2005	Female	All Ages	23% (17-29%)	77% (71-83%)	0.12 (0.08-0.16)	0.13% (0.028-0.24%)	0.071% (-0.18-0.29%)	58 (-150-240)	8200 (1600-14000)
Niger	2005	Female	15 to 19	22% (12-36%)	78% (64-88%)	0.11 (0.043-0.23)	0.47% (0.21-0.77%)	0.092% (-0.25-0.44%)	0.96 (-2.6-4.9)	630 (270-1100)
Niger	2005	Female	20 to 24	27% (14-45%)	73% (55-86%)	0.18 (0.072-0.35)	0.97% (0.43-1.6%)	0.17% (-0.4-0.74%)	2.2 (-5.1-10)	1300 (580-2200)
Niger	2005	Female	25 to 29	30% (15-47%)	70% (53-85%)	0.22 (0.09-0.43)	0.92% (0.46-1.4%)	0.28% (-0.31-0.85%)	4.5 (-4.3-13)	1300 (650-2100)
Niger	2005	Female	30 to 34	33% (17-51%)	67% (49-83%)	0.25 (0.11-0.46)	0.8% (0.34-1.3%)	0.37% (-0.22-0.92%)	6.4 (-3.9-16)	1100 (480-1800)
Niger	2005	Female	35 to 39	37% (20-56%)	63% (44-80%)	0.27 (0.12-0.49)	0.76% (0.25-1.3%)	0.51% (-0.13-1.2%)	9.1 (-2.4-21)	980 (330-1600)
Niger	2005	Female	40 to 44	38% (22-57%)	62% (43-78%)	0.28 (0.12-0.53)	0.85% (0.3-1.4%)	0.82% (-0.0019-1.6%)	15 (-0.034-29)	1000 (340-1700)
Niger	2005	Female	45 to 49	39% (21-58%)	61% (42-79%)	0.3 (0.11-0.58)	0.75% (0.025-1.5%)	0.79% (-0.2-1.8%)	14 (-3.6-32)	760 (25-1500)
Niger	2005	Female	50 to 54	40% (22-60%)	60% (40-78%)	0.31 (0.13-0.61)	0.77% (-0.042-1.6%)	0.91% (-0.18-2.1%)	16 (-3.1-36)	690 (-32-1400)
Niger	2005	Female	55 to 59	40% (23-58%)	60% (42-77%)	0.33 (0.13-0.65)	0.52% (-0.32-1.4%)	0.62% (-0.55-1.9%)	13 (-11-40)	470 (-280-1300)
Niger	2005	Female	60 to 64	41% (23-59%)	59% (41-77%)	0.33 (0.12-0.67)	0.26% (-0.64-1.1%)	0.3% (-0.84-1.4%)	7.7 (-22-36)	230 (-570-980)
Niger	2005	Female	65 to 69	42% (25-59%)	58% (41-75%)	0.3 (0.11-0.61)	-0.09% (-1.1-0.73%)	-0.14% (-1.4-0.88%)	-3.8 (-38-24)	-68 (-820-580)
Niger	2005	Female	70 to 74	40% (24-58%)	60% (42-76%)	0.25 (0.094-0.57)	-0.17% (-1.2-0.58%)	-0.23% (-1.5-0.69%)	-6.5 (-40-20)	-100 (-710-350)
Niger	2005	Female	75 to 79	36% (20-55%)	64% (45-80%)	0.2 (0.061-0.46)	-0.26% (-1.3-0.44%)	-0.33% (-1.6-0.5%)	-7.5 (-36-11)	-97 (-500-160)
Niger	2005	Female	80 to 84	33% (18-53%)	67% (47-82%)	0.18 (0.056-0.39)	-0.35% (-1.3-0.22%)	-0.44% (-1.5-0.25%)	-5.6 (-20-3.2)	-54 (-190-35)
Niger	2005	Female	85 to 89	34% (18-53%)	66% (47-82%)	0.18 (0.056-0.39)	-0.44% (-1.3-0.12%)	-0.56% (-1.6-0.11%)	-4.3 (-13-0.88)	-30 (-89-8)
Niger	2005	Female	90 to 94	37% (20-55%)	63% (45-80%)	0.19 (0.054-0.43)	-0.53% (-1.4-0.047%)	-0.69% (-1.7-0.014%)	-2 (-5-0.04)	-9.3 (-24-0.82)
Niger	2005	Female	95 plus	41% (21-62%)	59% (38-79%)	0.2 (0.05-0.51)	-0.51% (-1.5-0.081%)	-0.71% (-1.9-0.024%)	-0.52 (-1.4-0.017)	-1.5 (-4.2-0.25)
Niger	2005	Male	All Ages	19% (15-25%)	81% (75-85%)	0.15 (0.1-0.21)	0.34% (0.18-0.49%)	0.49% (0.19-0.81%)	440 (160-730)	23000 (12000-33000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	2005	Male	15 to 19	16% (8-27%)	84% (73-92%)	0.096 (0.035-0.2)	0.63% (0.36-0.99%)	0.3% (0.052-0.7%)	3.3 (0.58-7.8)	800 (450-1300)
Niger	2005	Male	20 to 24	23% (12-39%)	77% (61-88%)	0.19 (0.078-0.38)	1.4% (0.83-2.1%)	0.76% (0.14-1.6%)	8.6 (1.7-19)	1500 (870-2400)
Niger	2005	Male	25 to 29	29% (15-46%)	71% (54-85%)	0.29 (0.12-0.55)	1.7% (0.96-2.6%)	1.2% (0.23-2.4%)	17 (3.4-33)	2000 (1100-3100)
Niger	2005	Male	30 to 34	32% (17-50%)	68% (50-83%)	0.35 (0.16-0.63)	1.9% (1-2.9%)	1.6% (0.48-2.9%)	27 (7.8-48)	2400 (1200-3600)
Niger	2005	Male	35 to 39	35% (19-55%)	65% (45-81%)	0.38 (0.16-0.71)	2.1% (1.2-3.4%)	2.2% (0.89-3.9%)	37 (15-65)	2500 (1400-3900)
Niger	2005	Male	40 to 44	36% (21-55%)	64% (45-79%)	0.41 (0.16-0.76)	2.2% (1.1-3.5%)	2.5% (0.97-4.2%)	62 (23-110)	3400 (1500-5400)
Niger	2005	Male	45 to 49	37% (19-57%)	63% (43-81%)	0.42 (0.17-0.83)	2.3% (1.1-4%)	2.7% (1.1-4.9%)	68 (27-120)	3200 (1500-5400)
Niger	2005	Male	50 to 54	38% (21-59%)	62% (41-79%)	0.43 (0.18-0.84)	2.1% (0.83-3.8%)	2.5% (0.81-4.7%)	60 (19-110)	2400 (930-4300)
Niger	2005	Male	55 to 59	38% (22-57%)	62% (43-78%)	0.42 (0.16-0.84)	1.9% (0.73-3.4%)	2.2% (0.73-4.1%)	52 (18-100)	1800 (710-3300)
Niger	2005	Male	60 to 64	38% (22-57%)	62% (43-78%)	0.38 (0.13-0.82)	1.4% (0.56-2.6%)	1.7% (0.5-3.1%)	41 (13-80)	1200 (470-2300)
Niger	2005	Male	65 to 69	38% (21-56%)	62% (44-79%)	0.32 (0.12-0.68)	0.93% (0.15-1.8%)	1.1% (0.082-2.2%)	26 (1.9-53)	620 (100-1200)
Niger	2005	Male	70 to 74	36% (21-54%)	64% (46-79%)	0.25 (0.085-0.56)	0.63% (-0.12-1.4%)	0.72% (-0.21-1.7%)	18 (-5-43)	340 (-68-780)
Niger	2005	Male	75 to 79	33% (18-52%)	67% (48-82%)	0.19 (0.055-0.44)	0.41% (-0.28-1%)	0.45% (-0.39-1.2%)	9.8 (-8-25)	150 (-99-360)
Niger	2005	Male	80 to 84	30% (16-49%)	70% (51-84%)	0.16 (0.044-0.34)	0.34% (-0.24-0.82%)	0.36% (-0.35-0.93%)	4.2 (-4.4-11)	49 (-34-120)
Niger	2005	Male	85 to 89	29% (15-46%)	71% (54-85%)	0.13 (0.042-0.3)	0.23% (-0.29-0.64%)	0.22% (-0.42-0.71%)	1.4 (-2.6-4.4)	12 (-15-34)
Niger	2005	Male	90 to 94	31% (16-49%)	69% (51-84%)	0.13 (0.035-0.32)	0.089% (-0.39-0.44%)	0.031% (-0.56-0.46%)	0.063 (-1.1-0.94)	1.1 (-4.7-5.4)
Niger	2005	Male	95 plus	35% (17-57%)	65% (43-83%)	0.13 (0.027-0.34)	0.094% (-0.48-0.51%)	-0.0087% (-0.72-0.5%)	-0.0039 (-0.32-0.22)	0.17 (-0.84-0.89)
Niger	2010	Female	All Ages	22% (17-28%)	78% (72-83%)	0.11 (0.072-0.15)	0.15% (0.034-0.25%)	0.049% (-0.23-0.27%)	39 (-190-220)	8700 (2000-15000)
Niger	2010	Female	15 to 19	22% (12-35%)	78% (65-88%)	0.11 (0.04-0.22)	0.48% (0.23-0.76%)	0.078% (-0.2-0.39%)	0.97 (-2.7-4.9)	760 (360-1300)
Niger	2010	Female	20 to 24	27% (14-44%)	73% (56-86%)	0.17 (0.067-0.33)	1% (0.53-1.6%)	0.15% (-0.34-0.68%)	2 (-4.9-9.4)	1500 (790-2500)
Niger	2010	Female	25 to 29	30% (16-48%)	70% (52-84%)	0.2 (0.084-0.39)	1% (0.57-1.5%)	0.27% (-0.26-0.8%)	4.1 (-3.7-12)	1500 (770-2300)
Niger	2010	Female	30 to 34	33% (17-52%)	67% (48-83%)	0.23 (0.1-0.42)	0.85% (0.46-1.3%)	0.35% (-0.16-0.87%)	6.1 (-2.6-15)	1200 (640-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	2010	Female	35 to 39	37% (19-56%)	63% (44-81%)	0.25 (0.11-0.45)	0.76% (0.32-1.2%)	0.47% (-0.12-1%)	8.7 (-2.3-20)	1100 (440-1600)
Niger	2010	Female	40 to 44	38% (22-57%)	62% (43-78%)	0.26 (0.1-0.5)	0.82% (0.35-1.3%)	0.76% (0.058-1.5%)	14 (1.1-27)	1000 (410-1700)
Niger	2010	Female	45 to 49	39% (20-57%)	61% (43-80%)	0.27 (0.1-0.51)	0.72% (0.14-1.3%)	0.72% (-0.1-1.6%)	14 (-2.1-30)	820 (150-1500)
Niger	2010	Female	50 to 54	39% (21-60%)	61% (40-79%)	0.28 (0.12-0.54)	0.7% (-0.015-1.4%)	0.8% (-0.21-1.8%)	16 (-4.4-36)	710 (-16-1400)
Niger	2010	Female	55 to 59	40% (23-59%)	60% (41-77%)	0.3 (0.12-0.59)	0.44% (-0.41-1.2%)	0.52% (-0.66-1.7%)	11 (-14-36)	410 (-410-1200)
Niger	2010	Female	60 to 64	41% (23-59%)	59% (41-77%)	0.3 (0.11-0.62)	0.17% (-0.7-0.91%)	0.18% (-0.91-1.2%)	5 (-26-33)	160 (-670-910)
Niger	2010	Female	65 to 69	41% (24-59%)	59% (41-76%)	0.28 (0.1-0.55)	-0.17% (-1.2-0.56%)	-0.24% (-1.6-0.69%)	-7.8 (-49-23)	-150 (-1100-530)
Niger	2010	Female	70 to 74	39% (24-56%)	61% (44-76%)	0.23 (0.082-0.52)	-0.22% (-1.2-0.45%)	-0.3% (-1.5-0.52%)	-9.9 (-51-17)	-160 (-890-320)
Niger	2010	Female	75 to 79	35% (19-54%)	65% (46-81%)	0.18 (0.055-0.44)	-0.27% (-1.2-0.37%)	-0.36% (-1.6-0.41%)	-9.5 (-40-11)	-120 (-550-170)
Niger	2010	Female	80 to 84	33% (18-51%)	67% (49-82%)	0.16 (0.049-0.35)	-0.36% (-1.3-0.19%)	-0.46% (-1.5-0.2%)	-6.9 (-23-3.1)	-66 (-230-36)
Niger	2010	Female	85 to 89	33% (18-51%)	67% (49-82%)	0.16 (0.053-0.36)	-0.42% (-1.2-0.083%)	-0.54% (-1.5-0.066%)	-5.1 (-14-0.61)	-35 (-100-6.6)
Niger	2010	Female	90 to 94	36% (19-54%)	64% (46-81%)	0.17 (0.053-0.4)	-0.51% (-1.4-0.021%)	-0.68% (-1.7-0.016%)	-2.6 (-6.4-0.062)	-12 (-31-0.49)
Niger	2010	Female	95 plus	41% (21-62%)	59% (38-79%)	0.19 (0.045-0.46)	-0.49% (-1.5-0.078%)	-0.69% (-1.9-0.0058%)	-0.7 (-1.9-0.006)	-2 (-5.8-0.32)
Niger	2010	Male	All Ages	19% (15-24%)	81% (76-85%)	0.14 (0.091-0.19)	0.38% (0.21-0.58%)	0.53% (0.19-0.93%)	470 (170-820)	25000 (14000-38000)
Niger	2010	Male	15 to 19	16% (8.2-27%)	84% (73-92%)	0.09 (0.031-0.19)	0.62% (0.35-0.95%)	0.28% (0.053-0.64%)	3.8 (0.74-8.6)	970 (520-1500)
Niger	2010	Male	20 to 24	23% (12-39%)	77% (61-88%)	0.18 (0.07-0.37)	1.4% (0.83-2.1%)	0.71% (0.17-1.5%)	9.9 (2.5-22)	1900 (1100-2900)
Niger	2010	Male	25 to 29	29% (15-45%)	71% (55-85%)	0.26 (0.11-0.5)	1.8% (1-2.7%)	1.2% (0.26-2.3%)	17 (3.6-33)	2200 (1300-3300)
Niger	2010	Male	30 to 34	32% (17-52%)	68% (48-83%)	0.32 (0.15-0.57)	1.9% (1.1-2.9%)	1.6% (0.45-2.9%)	29 (8-55)	2700 (1500-4200)
Niger	2010	Male	35 to 39	35% (19-54%)	65% (46-81%)	0.35 (0.14-0.67)	2.1% (1.1-3.2%)	2.1% (0.82-3.6%)	41 (16-73)	2800 (1500-4400)
Niger	2010	Male	40 to 44	36% (21-55%)	64% (45-79%)	0.38 (0.14-0.71)	2.1% (1-3.5%)	2.4% (0.88-4.2%)	48 (18-86)	2700 (1300-4400)
Niger	2010	Male	45 to 49	37% (20-56%)	63% (44-80%)	0.4 (0.15-0.78)	2.3% (1.1-3.8%)	2.6% (1-4.7%)	77 (30-140)	3700 (1700-6300)
Niger	2010	Male	50 to 54	38% (21-59%)	62% (41-79%)	0.4 (0.16-0.8)	2% (0.79-3.6%)	2.3% (0.7-4.5%)	72 (21-140)	2900 (1200-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	2010	Male	55 to 59	38% (22-58%)	62% (42-78%)	0.39 (0.15-0.78)	1.8% (0.63-3.2%)	2.1% (0.55-3.9%)	62 (16-120)	2200 (800-4000)
Niger	2010	Male	60 to 64	38% (21-58%)	62% (42-79%)	0.35 (0.12-0.77)	1.3% (0.45-2.5%)	1.5% (0.37-3%)	47 (12-96)	1400 (470-2700)
Niger	2010	Male	65 to 69	38% (22-56%)	62% (44-78%)	0.3 (0.11-0.63)	0.83% (0.085-1.6%)	0.93% (-0.018-1.9%)	29 (-0.55-60)	720 (69-1400)
Niger	2010	Male	70 to 74	36% (21-53%)	64% (47-79%)	0.23 (0.079-0.54)	0.56% (-0.12-1.3%)	0.64% (-0.22-1.5%)	19 (-6.2-44)	360 (-78-800)
Niger	2010	Male	75 to 79	33% (18-51%)	67% (49-82%)	0.17 (0.052-0.4)	0.36% (-0.28-0.9%)	0.39% (-0.38-1.1%)	9.8 (-9.7-26)	150 (-110-370)
Niger	2010	Male	80 to 84	30% (16-49%)	70% (51-84%)	0.14 (0.039-0.3)	0.31% (-0.26-0.76%)	0.32% (-0.36-0.87%)	4.6 (-5.1-13)	54 (-44-130)
Niger	2010	Male	85 to 89	29% (15-46%)	71% (54-85%)	0.12 (0.038-0.29)	0.21% (-0.28-0.58%)	0.19% (-0.4-0.63%)	1.6 (-3.3-5.4)	15 (-20-44)
Niger	2010	Male	90 to 94	31% (16-48%)	69% (52-84%)	0.12 (0.034-0.29)	0.08% (-0.35-0.39%)	0.019% (-0.5-0.4%)	0.057 (-1.5-1.2)	1.4 (-6.3-7.1)
Niger	2010	Male	95 plus	35% (17-57%)	65% (43-83%)	0.12 (0.026-0.33)	0.081% (-0.45-0.48%)	-0.023% (-0.7-0.45%)	-0.015 (-0.46-0.29)	0.21 (-1.2-1.2)
Niger	2016	Female	All Ages	22% (17-28%)	78% (72-83%)	0.11 (0.058-0.17)	0.18% (0.042-0.32%)	0.053% (-0.28-0.32%)	44 (-230-270)	11000 (2600-19000)
Niger	2016	Female	15 to 19	22% (12-37%)	78% (63-88%)	0.11 (0.037-0.24)	0.5% (0.26-0.8%)	0.086% (-0.16-0.41%)	1.4 (-2.7-6.7)	1000 (510-1700)
Niger	2016	Female	20 to 24	28% (14-44%)	72% (56-86%)	0.17 (0.061-0.34)	1.1% (0.56-1.7%)	0.16% (-0.31-0.73%)	2.7 (-4.8-12)	2000 (1000-3200)
Niger	2016	Female	25 to 29	30% (16-49%)	70% (51-84%)	0.2 (0.078-0.42)	1.2% (0.69-1.7%)	0.33% (-0.25-0.96%)	5 (-4-15)	1800 (1000-2800)
Niger	2016	Female	30 to 34	33% (17-53%)	67% (47-83%)	0.23 (0.089-0.47)	1% (0.55-1.6%)	0.46% (-0.18-1.1%)	7.4 (-2.5-18)	1500 (800-2400)
Niger	2016	Female	35 to 39	37% (20-56%)	63% (44-80%)	0.25 (0.1-0.49)	0.9% (0.42-1.4%)	0.59% (-0.093-1.3%)	11 (-1.6-23)	1300 (600-2000)
Niger	2016	Female	40 to 44	39% (22-58%)	61% (42-78%)	0.26 (0.098-0.54)	0.94% (0.41-1.5%)	0.89% (0.088-1.8%)	17 (1.6-34)	1300 (560-2100)
Niger	2016	Female	45 to 49	39% (20-59%)	61% (41-80%)	0.27 (0.095-0.57)	0.8% (0.17-1.5%)	0.84% (-0.053-1.9%)	17 (-1.2-39)	1000 (210-1900)
Niger	2016	Female	50 to 54	40% (22-60%)	60% (40-78%)	0.28 (0.1-0.61)	0.76% (-0.0073-1.5%)	0.89% (-0.18-2%)	21 (-4.3-48)	920 (-9.4-1900)
Niger	2016	Female	55 to 59	40% (23-58%)	60% (42-77%)	0.31 (0.11-0.63)	0.48% (-0.34-1.3%)	0.56% (-0.53-1.7%)	15 (-15-49)	550 (-410-1600)
Niger	2016	Female	60 to 64	41% (24-61%)	59% (39-76%)	0.31 (0.1-0.67)	0.17% (-0.71-0.95%)	0.18% (-0.95-1.2%)	5.4 (-30-37)	180 (-780-1100)
Niger	2016	Female	65 to 69	42% (24-60%)	58% (40-76%)	0.28 (0.1-0.6)	-0.18% (-1.2-0.57%)	-0.26% (-1.6-0.67%)	-10 (-61-26)	-200 (-1400-600)
Niger	2016	Female	70 to 74	39% (23-57%)	61% (43-77%)	0.23 (0.073-0.54)	-0.24% (-1.2-0.43%)	-0.33% (-1.5-0.48%)	-14 (-67-21)	-230 (-1200-410)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	2016	Female	75 to 79	35% (20-54%)	65% (46-80%)	0.18 (0.052-0.45)	-0.27% (-1.2-0.36%)	-0.36% (-1.6-0.4%)	-13 (-56-14)	-170 (-760-210)
Niger	2016	Female	80 to 84	33% (17-51%)	67% (49-83%)	0.16 (0.047-0.37)	-0.36% (-1.2-0.19%)	-0.46% (-1.5-0.19%)	-9.3 (-31-3.7)	-90 (-310-48)
Niger	2016	Female	85 to 89	33% (19-51%)	67% (49-81%)	0.16 (0.05-0.41)	-0.43% (-1.3-0.088%)	-0.56% (-1.6-0.076%)	-7 (-19-0.91)	-47 (-140-9.3)
Niger	2016	Female	90 to 94	37% (19-54%)	63% (46-81%)	0.18 (0.048-0.45)	-0.53% (-1.4-0.021%)	-0.7% (-1.7--0.023%)	-3.5 (-8.7--0.11)	-16 (-41-0.64)
Niger	2016	Female	95 plus	41% (21-62%)	59% (38-79%)	0.19 (0.043-0.51)	-0.51% (-1.5-0.068%)	-0.71% (-1.9-0.0092%)	-1 (-2.7-0.013)	-2.9 (-8.3-0.39)
Niger	2016	Male	All Ages	19% (14-24%)	81% (76-86%)	0.13 (0.074-0.21)	0.46% (0.24-0.71%)	0.62% (0.2-1.1%)	590 (200-1100)	31000 (16000-49000)
Niger	2016	Male	15 to 19	16% (8.1-26%)	84% (74-92%)	0.09 (0.029-0.2)	0.64% (0.38-0.99%)	0.28% (0.064-0.66%)	5 (1.2-11)	1300 (760-2100)
Niger	2016	Male	20 to 24	23% (12-38%)	77% (62-88%)	0.18 (0.062-0.38)	1.5% (0.89-2.2%)	0.74% (0.19-1.6%)	13 (3.5-29)	2600 (1500-4000)
Niger	2016	Male	25 to 29	29% (15-46%)	71% (54-85%)	0.26 (0.1-0.56)	1.9% (1.1-2.9%)	1.3% (0.33-2.6%)	22 (5.4-46)	2900 (1700-4500)
Niger	2016	Male	30 to 34	33% (17-51%)	67% (49-83%)	0.32 (0.13-0.65)	2.1% (1.2-3.4%)	1.8% (0.58-3.6%)	34 (11-65)	3100 (1700-5000)
Niger	2016	Male	35 to 39	36% (20-54%)	64% (46-80%)	0.36 (0.14-0.74)	2.3% (1.2-3.6%)	2.3% (0.92-4.1%)	51 (20-90)	3500 (1900-5600)
Niger	2016	Male	40 to 44	37% (21-55%)	63% (45-79%)	0.38 (0.14-0.81)	2.3% (1.1-3.7%)	2.5% (0.98-4.6%)	59 (24-110)	3300 (1700-5700)
Niger	2016	Male	45 to 49	37% (19-56%)	63% (44-81%)	0.4 (0.13-0.9)	2.4% (1.2-4.2%)	2.8% (1.1-5.2%)	68 (27-130)	3200 (1500-5800)
Niger	2016	Male	50 to 54	38% (20-59%)	62% (41-80%)	0.4 (0.14-0.91)	2.1% (0.86-3.9%)	2.5% (0.82-4.8%)	89 (30-180)	3600 (1500-7000)
Niger	2016	Male	55 to 59	38% (22-57%)	62% (43-78%)	0.39 (0.13-0.87)	1.9% (0.74-3.6%)	2.2% (0.68-4.3%)	90 (27-170)	3100 (1200-5800)
Niger	2016	Male	60 to 64	38% (21-57%)	62% (43-79%)	0.35 (0.11-0.83)	1.4% (0.42-2.7%)	1.6% (0.36-3.4%)	68 (16-140)	2000 (620-3900)
Niger	2016	Male	65 to 69	38% (21-56%)	62% (44-79%)	0.3 (0.097-0.67)	0.87% (0.078-1.8%)	0.98% (-0.02-2.1%)	41 (-0.95-91)	1000 (87-2100)
Niger	2016	Male	70 to 74	36% (21-54%)	64% (46-79%)	0.23 (0.072-0.56)	0.58% (-0.099-1.3%)	0.66% (-0.17-1.5%)	26 (-6.9-64)	510 (-84-1200)
Niger	2016	Male	75 to 79	33% (19-51%)	67% (49-81%)	0.18 (0.048-0.46)	0.37% (-0.26-0.9%)	0.4% (-0.34-1%)	13 (-11-33)	200 (-130-480)
Niger	2016	Male	80 to 84	30% (16-49%)	70% (51-84%)	0.14 (0.039-0.34)	0.32% (-0.21-0.77%)	0.33% (-0.32-0.88%)	5.7 (-5.5-15)	67 (-45-160)
Niger	2016	Male	85 to 89	29% (15-47%)	71% (53-85%)	0.12 (0.035-0.32)	0.22% (-0.23-0.59%)	0.2% (-0.33-0.66%)	2.1 (-3.6-6.9)	20 (-21-55)
Niger	2016	Male	90 to 94	31% (16-49%)	69% (51-84%)	0.12 (0.029-0.32)	0.088% (-0.33-0.41%)	0.026% (-0.5-0.4%)	0.11 (-2.1-1.8)	2.3 (-8.6-10)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Niger	2016	Male	95 plus	36% (17-59%)	64% (41-83%)	0.12 (0.025-0.34)	0.085% (-0.47-0.46%)	-0.021% (-0.73-0.45%)	-0.022 (-0.74-0.45)	0.34 (-1.9-1.9)
Nigeria	1990	Female	All Ages	39% (33-45%)	61% (55-67%)	0.67 (0.48-0.93)	0.8% (0.48-1.2%)	1.5% (0.78-2.6%)	8800 (4400-15000)	340000 (200000-530000)
Nigeria	1990	Female	15 to 19	37% (26-49%)	63% (51-74%)	0.77 (0.42-1.3)	1.4% (0.65-2.4%)	1.7% (0.45-3.5%)	130 (35-270)	14000 (6400-25000)
Nigeria	1990	Female	20 to 24	45% (32-58%)	55% (42-68%)	1.1 (0.63-1.9)	2.6% (1.4-4.2%)	2.7% (1-5.3%)	250 (89-510)	27000 (14000-45000)
Nigeria	1990	Female	25 to 29	49% (36-62%)	51% (38-64%)	1.3 (0.72-2.1)	3.2% (1.7-5.2%)	3.9% (1.6-7%)	370 (140-690)	31000 (16000-52000)
Nigeria	1990	Female	30 to 34	53% (40-65%)	47% (35-60%)	1.3 (0.78-2.2)	3.4% (1.9-5.6%)	4.4% (2.1-7.6%)	430 (190-770)	30000 (17000-50000)
Nigeria	1990	Female	35 to 39	55% (41-67%)	45% (33-59%)	1.3 (0.78-2.1)	3.8% (2.1-6.1%)	5.2% (2.7-8.8%)	540 (260-930)	32000 (17000-52000)
Nigeria	1990	Female	40 to 44	57% (44-68%)	43% (32-56%)	1.3 (0.73-2.1)	4.1% (2.2-6.6%)	6% (3-9.8%)	660 (320-1100)	33000 (17000-53000)
Nigeria	1990	Female	45 to 49	58% (44-69%)	42% (31-56%)	1.1 (0.57-1.8)	3.7% (1.6-6.6%)	5.4% (2.2-9.5%)	660 (260-1200)	29000 (12000-51000)
Nigeria	1990	Female	50 to 54	58% (44-70%)	42% (30-56%)	1.1 (0.59-1.9)	4% (1.8-6.9%)	5.6% (2.5-9.8%)	770 (350-1400)	29000 (14000-51000)
Nigeria	1990	Female	55 to 59	56% (42-70%)	44% (30-58%)	1.5 (0.79-2.5)	5.3% (2.5-8.9%)	7.2% (3.4-12%)	1200 (530-2000)	37000 (17000-65000)
Nigeria	1990	Female	60 to 64	56% (40-70%)	44% (30-60%)	1.6 (0.84-2.7)	4.9% (2.1-8.7%)	6.5% (2.8-11%)	1300 (540-2200)	34000 (15000-60000)
Nigeria	1990	Female	65 to 69	56% (40-71%)	44% (29-60%)	1.4 (0.8-2.3)	3.9% (1.5-7.1%)	5% (1.8-9.1%)	1100 (410-2100)	25000 (9400-47000)
Nigeria	1990	Female	70 to 74	53% (38-68%)	47% (32-62%)	1.2 (0.62-2)	3% (0.38-6.5%)	3.7% (0.4-8%)	860 (94-1900)	15000 (1900-34000)
Nigeria	1990	Female	75 to 79	49% (33-66%)	51% (34-67%)	0.96 (0.46-1.7)	1.8% (-0.48-4.9%)	2.1% (-0.59-6%)	390 (-110-1100)	5500 (-1500-15000)
Nigeria	1990	Female	80 to 84	47% (30-65%)	53% (35-70%)	0.89 (0.46-1.6)	1.1% (-0.68-3.6%)	1.3% (-0.82-4.4%)	120 (-75-400)	1300 (-750-4100)
Nigeria	1990	Female	85 to 89	47% (31-64%)	53% (36-69%)	0.9 (0.46-1.5)	0.76% (-0.82-3.1%)	0.88% (-1-3.7%)	52 (-64-210)	400 (-430-1600)
Nigeria	1990	Female	90 to 94	50% (35-67%)	50% (33-65%)	0.94 (0.46-1.6)	0.51% (-1-2.9%)	0.54% (-1.4-3.4%)	13 (-34-85)	78 (-160-440)
Nigeria	1990	Female	95 plus	53% (34-73%)	47% (27-66%)	0.97 (0.4-1.9)	0.44% (-1.1-3%)	0.39% (-1.5-3.4%)	3 (-11-26)	13 (-33-90)
Nigeria	1990	Male	All Ages	46% (39-53%)	54% (47-61%)	2.7 (2-3.6)	2.9% (2.2-3.8%)	5.8% (4.2-7.7%)	37000 (26000-50000)	140000 (110000-190000)
Nigeria	1990	Male	15 to 19	39% (27-53%)	61% (47-73%)	1.8 (0.97-3)	4.3% (2.4-6.7%)	6.4% (3.4-10%)	500 (260-830)	43000 (24000-66000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	1990	Male	20 to 24	55% (40-68%)	45% (32-60%)	3.7 (2.2-5.9)	11% (7.1-15%)	14% (9.4-21%)	1400 (870-2100)	110000 (69000-150000)
Nigeria	1990	Male	25 to 29	62% (47-74%)	38% (26-53%)	5.4 (3.3-8.2)	15% (9.7-21%)	20% (12-30%)	2100 (1300-3100)	140000 (93000-200000)
Nigeria	1990	Male	30 to 34	65% (50-76%)	35% (24-50%)	6.1 (3.7-9.1)	16% (11-22%)	22% (14-31%)	2300 (1500-3300)	140000 (97000-200000)
Nigeria	1990	Male	35 to 39	66% (52-77%)	34% (23-48%)	6 (3.7-9.3)	16% (11-22%)	23% (15-31%)	2500 (1700-3500)	140000 (94000-190000)
Nigeria	1990	Male	40 to 44	67% (54-78%)	33% (22-46%)	6.3 (3.8-9.6)	18% (14-24%)	26% (19-34%)	3100 (2200-4200)	150000 (110000-200000)
Nigeria	1990	Male	45 to 49	68% (55-78%)	32% (22-45%)	7 (4.2-11)	21% (16-27%)	29% (22-37%)	3800 (2700-5100)	160000 (120000-220000)
Nigeria	1990	Male	50 to 54	70% (57-80%)	30% (20-43%)	7 (4.1-11)	21% (16-28%)	29% (21-38%)	4100 (2900-5600)	150000 (110000-200000)
Nigeria	1990	Male	55 to 59	69% (56-80%)	31% (20-44%)	6.3 (3.7-9.7)	20% (14-26%)	26% (19-35%)	4000 (2700-5500)	130000 (89000-180000)
Nigeria	1990	Male	60 to 64	68% (54-81%)	32% (19-46%)	5.2 (2.8-8.5)	16% (11-22%)	21% (14-28%)	3700 (2400-5200)	100000 (66000-140000)
Nigeria	1990	Male	65 to 69	66% (49-79%)	34% (21-51%)	4.4 (2.6-7)	15% (9.4-20%)	19% (12-26%)	3500 (2200-5000)	80000 (51000-110000)
Nigeria	1990	Male	70 to 74	63% (47-77%)	37% (23-53%)	3.6 (2.1-5.9)	13% (7.7-20%)	16% (9.5-24%)	3000 (1700-4600)	54000 (32000-84000)
Nigeria	1990	Male	75 to 79	60% (43-74%)	40% (26-57%)	2.8 (1.4-4.8)	10% (5.2-17%)	13% (6.1-20%)	1700 (810-2900)	24000 (12000-40000)
Nigeria	1990	Male	80 to 84	57% (38-73%)	43% (27-62%)	2.2 (1.2-3.6)	8.2% (3.9-14%)	9.9% (4.8-17%)	640 (300-1100)	6600 (3200-11000)
Nigeria	1990	Male	85 to 89	53% (35-69%)	47% (31-65%)	1.7 (0.9-2.8)	5.9% (2.7-11%)	7.1% (3.2-13%)	280 (130-510)	2100 (970-3800)
Nigeria	1990	Male	90 to 94	53% (37-70%)	47% (30-63%)	1.3 (0.68-2.3)	3.6% (1.2-7.1%)	4.2% (1.2-8.4%)	71 (21-140)	370 (120-730)
Nigeria	1990	Male	95 plus	57% (35-75%)	43% (25-65%)	1.1 (0.45-2.1)	2.7% (0.086-6.3%)	3% (-0.16-7.4%)	14 (-0.75-35)	51 (1.5-120)
Nigeria	1995	Female	All Ages	39% (33-45%)	61% (55-67%)	0.75 (0.55-1)	0.91% (0.57-1.3%)	1.8% (0.92-2.8%)	12000 (5800-19000)	440000 (270000-660000)
Nigeria	1995	Female	15 to 19	38% (27-50%)	62% (50-73%)	0.87 (0.46-1.4)	1.5% (0.73-2.7%)	1.9% (0.58-3.9%)	180 (55-390)	19000 (8200-35000)
Nigeria	1995	Female	20 to 24	46% (33-59%)	54% (41-67%)	1.3 (0.72-2.1)	2.6% (1.4-4.2%)	2.8% (1.1-5.1%)	360 (130-660)	35000 (19000-56000)
Nigeria	1995	Female	25 to 29	50% (36-62%)	50% (38-64%)	1.4 (0.85-2.3)	3.1% (1.6-4.9%)	3.4% (1.4-6.2%)	480 (200-880)	39000 (21000-63000)
Nigeria	1995	Female	30 to 34	53% (39-66%)	47% (34-61%)	1.5 (0.88-2.3)	3.2% (1.7-5%)	3.8% (1.8-6.5%)	550 (260-950)	38000 (21000-61000)
Nigeria	1995	Female	35 to 39	56% (42-67%)	44% (33-58%)	1.5 (0.89-2.4)	3.7% (1.9-5.9%)	4.8% (2.3-8%)	700 (350-1200)	40000 (22000-65000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	1995	Female	40 to 44	57% (46-69%)	43% (31-54%)	1.5 (0.81-2.3)	4.2% (2.2-6.5%)	5.8% (2.9-9.1%)	820 (400-1300)	41000 (21000-65000)
Nigeria	1995	Female	45 to 49	58% (44-70%)	42% (30-56%)	1.2 (0.65-2)	3.9% (1.7-6.8%)	5.4% (2.2-9.4%)	820 (350-1500)	36000 (16000-62000)
Nigeria	1995	Female	50 to 54	58% (45-70%)	42% (30-55%)	1.2 (0.67-2.1)	4.3% (2.1-7.4%)	5.9% (2.8-10%)	1000 (450-1800)	37000 (17000-65000)
Nigeria	1995	Female	55 to 59	57% (42-70%)	43% (30-58%)	1.7 (0.92-2.8)	5.8% (2.8-9.6%)	7.8% (4-13%)	1500 (730-2600)	48000 (23000-83000)
Nigeria	1995	Female	60 to 64	56% (41-70%)	44% (30-59%)	1.8 (1-2.9)	5.6% (2.7-9.3%)	7.2% (3.6-12%)	1600 (750-2800)	44000 (20000-76000)
Nigeria	1995	Female	65 to 69	56% (40-71%)	44% (29-60%)	1.6 (0.92-2.5)	4.6% (1.8-8%)	5.8% (2.2-9.9%)	1500 (590-2600)	33000 (13000-60000)
Nigeria	1995	Female	70 to 74	54% (39-69%)	46% (31-61%)	1.3 (0.69-2.2)	3.5% (0.57-7.3%)	4.3% (0.73-9.2%)	1200 (180-2400)	21000 (3300-44000)
Nigeria	1995	Female	75 to 79	50% (35-67%)	50% (33-65%)	1.1 (0.52-1.9)	2.2% (-0.22-5.5%)	2.7% (-0.29-6.6%)	570 (-61-1500)	8000 (-800-20000)
Nigeria	1995	Female	80 to 84	48% (31-66%)	52% (34-69%)	1 (0.51-1.7)	1.5% (-0.5-4.1%)	1.7% (-0.62-5%)	180 (-65-530)	1900 (-660-5500)
Nigeria	1995	Female	85 to 89	48% (32-64%)	52% (36-68%)	0.99 (0.53-1.6)	1.1% (-0.68-3.3%)	1.2% (-0.87-4%)	82 (-57-270)	630 (-380-2000)
Nigeria	1995	Female	90 to 94	50% (35-66%)	50% (34-65%)	1 (0.51-1.7)	0.77% (-0.9-3.4%)	0.84% (-1.2-4%)	24 (-35-110)	130 (-160-580)
Nigeria	1995	Female	95 plus	53% (35-72%)	47% (28-65%)	1.1 (0.45-1.9)	0.67% (-1.1-3.5%)	0.65% (-1.4-4.1%)	5.5 (-12-35)	23 (-36-120)
Nigeria	1995	Male	All Ages	47% (40-54%)	53% (46-60%)	3 (2.2-3.9)	3.1% (2.2-4.1%)	6.1% (4.4-8.1%)	43000 (30000-58000)	1700000 (1200000-2200000)
Nigeria	1995	Male	15 to 19	40% (27-55%)	60% (45-73%)	2 (1.1-3.3)	4.8% (2.7-7.4%)	7.4% (3.9-12%)	670 (350-1100)	55000 (31000-86000)
Nigeria	1995	Male	20 to 24	55% (40-69%)	45% (31-60%)	4.2 (2.5-6.5)	11% (7.3-16%)	15% (9.5-22%)	1800 (1100-2700)	130000 (85000-190000)
Nigeria	1995	Male	25 to 29	63% (48-74%)	37% (26-52%)	6 (3.7-8.8)	13% (8.3-19%)	17% (10-26%)	2500 (1500-3600)	160000 (110000-230000)
Nigeria	1995	Male	30 to 34	65% (52-76%)	35% (24-48%)	6.7 (4.3-10)	13% (8.6-19%)	17% (10-26%)	2700 (1700-3800)	160000 (110000-230000)
Nigeria	1995	Male	35 to 39	66% (53-77%)	34% (23-47%)	6.6 (4-10)	14% (9-20%)	19% (11-27%)	2900 (1900-4000)	160000 (110000-210000)
Nigeria	1995	Male	40 to 44	67% (55-79%)	33% (21-45%)	7 (4.2-10)	16% (11-22%)	21% (14-30%)	3300 (2200-4700)	160000 (110000-220000)
Nigeria	1995	Male	45 to 49	69% (55-79%)	31% (21-45%)	7.8 (4.8-12)	19% (14-25%)	25% (18-34%)	4200 (2900-5800)	180000 (120000-240000)
Nigeria	1995	Male	50 to 54	70% (58-80%)	30% (20-42%)	7.9 (4.7-12)	20% (15-26%)	27% (20-36%)	4700 (3300-6300)	180000 (120000-230000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	1995	Male	55 to 59	70% (58-80%)	30% (20-42%)	7.1 (4.2-11)	20% (14-26%)	26% (19-35%)	4700 (3100-6500)	150000 (100000-200000)
Nigeria	1995	Male	60 to 64	69% (55-81%)	31% (19-45%)	5.9 (3.3-9.4)	17% (12-23%)	22% (15-29%)	4400 (2900-6000)	120000 (79000-160000)
Nigeria	1995	Male	65 to 69	67% (52-80%)	33% (20-48%)	5 (3-7.8)	16% (10-21%)	20% (13-27%)	4200 (2700-5800)	95000 (62000-130000)
Nigeria	1995	Male	70 to 74	65% (49-77%)	35% (23-51%)	4.1 (2.3-6.5)	15% (8.7-21%)	18% (11-26%)	3700 (2100-5500)	68000 (39000-98000)
Nigeria	1995	Male	75 to 79	61% (44-76%)	39% (24-56%)	3.2 (1.6-5.4)	12% (5.7-18%)	14% (6.7-22%)	2200 (1000-3600)	31000 (15000-49000)
Nigeria	1995	Male	80 to 84	58% (39-74%)	42% (26-61%)	2.5 (1.3-4.2)	9.3% (4.5-15%)	11% (5.4-18%)	850 (410-1400)	8800 (4300-14000)
Nigeria	1995	Male	85 to 89	54% (37-71%)	46% (29-63%)	1.9 (1-3.2)	6.9% (3.2-12%)	8.2% (3.7-14%)	380 (170-650)	2800 (1300-4700)
Nigeria	1995	Male	90 to 94	54% (38-71%)	46% (29-62%)	1.5 (0.78-2.6)	4.3% (1.5-8.3%)	5.1% (1.7-9.9%)	97 (32-190)	500 (180-960)
Nigeria	1995	Male	95 plus	58% (36-77%)	42% (23-64%)	1.3 (0.52-2.3)	3.3% (0.42-7.4%)	3.7% (0.26-8.7%)	19 (1.3-45)	67 (8.2-150)
Nigeria	2000	Female	All Ages	38% (32-44%)	62% (56-68%)	0.77 (0.57-1)	0.86% (0.5-1.3%)	1.6% (0.81-2.7%)	12000 (5800-20000)	460000 (270000-690000)
Nigeria	2000	Female	15 to 19	39% (28-50%)	61% (50-72%)	0.92 (0.52-1.5)	1.5% (0.77-2.5%)	1.9% (0.65-3.6%)	190 (58-400)	20000 (9800-36000)
Nigeria	2000	Female	20 to 24	45% (34-57%)	55% (43-66%)	1.4 (0.78-2.1)	2.4% (1.4-3.8%)	2.5% (0.97-4.5%)	370 (130-720)	38000 (20000-63000)
Nigeria	2000	Female	25 to 29	49% (38-59%)	51% (41-62%)	1.5 (0.89-2.2)	2.4% (1.3-3.9%)	2.5% (1-4.6%)	500 (210-910)	42000 (23000-67000)
Nigeria	2000	Female	30 to 34	52% (40-64%)	48% (36-60%)	1.5 (0.92-2.4)	2.4% (1.3-3.8%)	2.6% (1.2-4.5%)	540 (260-940)	39000 (22000-62000)
Nigeria	2000	Female	35 to 39	54% (43-65%)	46% (35-57%)	1.5 (0.91-2.5)	2.8% (1.5-4.5%)	3.3% (1.6-5.7%)	690 (340-1200)	41000 (23000-65000)
Nigeria	2000	Female	40 to 44	56% (45-66%)	44% (34-55%)	1.5 (0.84-2.4)	3.3% (1.8-5.4%)	4.3% (2.1-7.2%)	840 (410-1400)	42000 (22000-67000)
Nigeria	2000	Female	45 to 49	57% (45-68%)	43% (32-55%)	1.2 (0.68-2)	3.3% (1.5-5.7%)	4.4% (1.9-7.7%)	830 (350-1500)	36000 (16000-65000)
Nigeria	2000	Female	50 to 54	56% (43-68%)	44% (32-57%)	1.2 (0.69-2.1)	3.8% (1.8-6.5%)	5.1% (2.4-8.8%)	1000 (470-1800)	38000 (18000-67000)
Nigeria	2000	Female	55 to 59	54% (40-68%)	46% (32-60%)	1.7 (0.91-2.7)	5.3% (2.6-8.8%)	7.2% (3.5-12%)	1600 (750-2700)	50000 (24000-85000)
Nigeria	2000	Female	60 to 64	54% (39-69%)	46% (31-61%)	1.8 (0.97-2.9)	5.3% (2.3-9%)	6.8% (3-12%)	1700 (720-3000)	46000 (19000-80000)
Nigeria	2000	Female	65 to 69	55% (40-70%)	45% (30-60%)	1.6 (0.93-2.6)	4.4% (1.8-8.1%)	5.6% (2.3-10%)	1500 (600-2900)	34000 (14000-64000)
Nigeria	2000	Female	70 to 74	54% (38-69%)	46% (31-62%)	1.4 (0.74-2.3)	3.4% (0.48-7.3%)	4.2% (0.57-9.1%)	1200 (160-2600)	21000 (2900-47000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	2000	Female	75 to 79	50% (33-66%)	50% (34-67%)	1.1 (0.56-2)	2.1% (-0.3-5.6%)	2.5% (-0.37-6.7%)	580 (-91-1500)	8100 (-1200-21000)
Nigeria	2000	Female	80 to 84	47% (30-65%)	53% (35-70%)	1 (0.53-1.7)	1.4% (-0.52-4%)	1.6% (-0.68-4.9%)	190 (-80-550)	2000 (-750-5800)
Nigeria	2000	Female	85 to 89	48% (32-63%)	52% (37-68%)	1 (0.53-1.7)	0.99% (-0.83-3.3%)	1.1% (-1-4%)	85 (-79-300)	660 (-540-2300)
Nigeria	2000	Female	90 to 94	50% (35-66%)	50% (34-65%)	1.1 (0.54-1.8)	0.71% (-1.1-3.3%)	0.75% (-1.4-3.9%)	24 (-44-120)	140 (-210-630)
Nigeria	2000	Female	95 plus	53% (34-71%)	47% (29-66%)	1.1 (0.46-2.1)	0.62% (-1.2-3.7%)	0.56% (-1.6-4.2%)	5.2 (-15-39)	23 (-44-140)
Nigeria	2000	Male	All Ages	49% (43-56%)	51% (44-57%)	3.3 (2.5-4.3)	3.2% (2.3-4.3%)	6.2% (4.4-8.2%)	49000 (35000-65000)	1900000 (1400000-2500000)
Nigeria	2000	Male	15 to 19	47% (34-60%)	53% (40-66%)	2.6 (1.5-4)	5.5% (3.2-8.2%)	8.4% (4.6-13%)	870 (470-1400)	72000 (43000-110000)
Nigeria	2000	Male	20 to 24	61% (49-72%)	39% (28-51%)	4.8 (3-7.5)	11% (7.8-16%)	15% (10-22%)	2200 (1400-3300)	160000 (110000-230000)
Nigeria	2000	Male	25 to 29	65% (54-75%)	35% (25-46%)	6.5 (4.2-9.4)	12% (7.9-18%)	15% (9.3-23%)	2900 (1900-4300)	190000 (130000-280000)
Nigeria	2000	Male	30 to 34	66% (55-76%)	34% (24-45%)	7.2 (4.8-10)	11% (7.7-16%)	14% (8.7-21%)	3000 (2000-4400)	180000 (130000-260000)
Nigeria	2000	Male	35 to 39	67% (56-77%)	33% (23-44%)	7 (4.5-11)	12% (7.5-17%)	15% (8.7-22%)	3300 (2200-4600)	180000 (120000-250000)
Nigeria	2000	Male	40 to 44	68% (57-78%)	32% (22-43%)	7.4 (4.6-11)	13% (9.3-19%)	17% (11-24%)	3800 (2600-5300)	180000 (130000-250000)
Nigeria	2000	Male	45 to 49	70% (59-79%)	30% (21-41%)	8.2 (5.1-12)	16% (12-22%)	21% (14-29%)	4500 (3100-6300)	190000 (130000-260000)
Nigeria	2000	Male	50 to 54	71% (60-80%)	29% (20-40%)	8.3 (5-13)	19% (14-25%)	25% (18-32%)	5200 (3600-7100)	190000 (140000-260000)
Nigeria	2000	Male	55 to 59	70% (60-80%)	30% (20-40%)	7.4 (4.5-11)	19% (14-25%)	25% (18-32%)	5300 (3600-7400)	170000 (120000-240000)
Nigeria	2000	Male	60 to 64	68% (54-80%)	32% (20-46%)	6.1 (3.4-9.6)	16% (11-21%)	21% (15-27%)	4900 (3300-6800)	130000 (92000-190000)
Nigeria	2000	Male	65 to 69	67% (52-80%)	33% (20-48%)	5.2 (3.1-7.9)	16% (11-21%)	20% (13-26%)	4600 (3000-6400)	110000 (70000-150000)
Nigeria	2000	Male	70 to 74	65% (50-77%)	35% (23-50%)	4.3 (2.5-6.8)	14% (8.6-20%)	18% (11-25%)	4200 (2400-6100)	76000 (45000-110000)
Nigeria	2000	Male	75 to 79	61% (45-76%)	39% (24-55%)	3.3 (1.8-5.6)	12% (6.1-18%)	14% (7.4-22%)	2500 (1300-4000)	36000 (18000-55000)
Nigeria	2000	Male	80 to 84	58% (40-73%)	42% (27-60%)	2.6 (1.5-4.3)	9.4% (4.8-15%)	11% (5.7-18%)	980 (490-1600)	10000 (5200-17000)
Nigeria	2000	Male	85 to 89	55% (37-71%)	45% (29-63%)	2 (1.1-3.3)	6.9% (3.2-12%)	8.2% (3.8-14%)	440 (200-770)	3300 (1500-5600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	2000	Male	90 to 94	55% (39-72%)	45% (28-61%)	1.6 (0.84-2.7)	4.4% (1.6-8.2%)	5.1% (1.7-9.8%)	110 (37-210)	580 (210-1100)
Nigeria	2000	Male	95 plus	58% (37-77%)	42% (23-63%)	1.3 (0.56-2.4)	3.3% (0.35-7.6%)	3.7% (0.082-8.9%)	21 (0.47-51)	75 (7.8-170)
Nigeria	2005	Female	All Ages	38% (32-44%)	62% (56-68%)	0.73 (0.53-0.98)	0.69% (0.42-1.1%)	1.2% (0.6-2.1%)	9400 (4300-16000)	380000 (220000-580000)
Nigeria	2005	Female	15 to 19	38% (28-50%)	62% (50-72%)	0.87 (0.49-1.4)	1.2% (0.68-2.9%)	1.4% (0.47-2.9%)	130 (38-290)	17000 (8500-28000)
Nigeria	2005	Female	20 to 24	45% (34-57%)	55% (43-66%)	1.3 (0.71-2)	2.1% (1.2-3.1%)	1.9% (0.74-3.5%)	260 (94-500)	32000 (18000-51000)
Nigeria	2005	Female	25 to 29	49% (37-59%)	51% (41-63%)	1.4 (0.85-2.1)	1.8% (1-2.9%)	1.7% (0.64-3.2%)	370 (150-660)	36000 (20000-54000)
Nigeria	2005	Female	30 to 34	52% (40-63%)	48% (37-60%)	1.5 (0.87-2.2)	1.7% (0.92-2.7%)	1.6% (0.71-3%)	430 (190-770)	34000 (19000-53000)
Nigeria	2005	Female	35 to 39	54% (42-65%)	46% (35-58%)	1.5 (0.87-2.3)	1.9% (1.1-3.1%)	2.1% (0.98-3.7%)	540 (270-950)	34000 (19000-55000)
Nigeria	2005	Female	40 to 44	55% (44-66%)	45% (34-56%)	1.4 (0.79-2.3)	2.3% (1.2-3.7%)	2.8% (1.4-4.7%)	690 (330-1100)	36000 (19000-57000)
Nigeria	2005	Female	45 to 49	57% (44-67%)	43% (33-56%)	1.1 (0.64-1.9)	2.5% (1.1-4.2%)	3.2% (1.4-5.6%)	710 (290-1200)	32000 (14000-54000)
Nigeria	2005	Female	50 to 54	56% (43-68%)	44% (32-57%)	1.2 (0.63-1.9)	3% (1.4-5.1%)	4% (1.8-6.9%)	840 (370-1500)	32000 (15000-55000)
Nigeria	2005	Female	55 to 59	54% (40-68%)	46% (32-60%)	1.6 (0.87-2.6)	4.3% (2.1-7.2%)	5.9% (2.9-9.7%)	1300 (600-2200)	41000 (19000-72000)
Nigeria	2005	Female	60 to 64	54% (39-69%)	46% (31-61%)	1.7 (0.93-2.7)	4.3% (1.8-7.4%)	5.7% (2.5-9.9%)	1400 (580-2400)	38000 (16000-66000)
Nigeria	2005	Female	65 to 69	55% (40-71%)	45% (29-60%)	1.6 (0.9-2.4)	3.6% (1.3-6.8%)	4.7% (1.7-8.7%)	1200 (450-2400)	28000 (10000-53000)
Nigeria	2005	Female	70 to 74	53% (38-69%)	47% (31-62%)	1.3 (0.7-2.2)	2.7% (0.11-6.1%)	3.3% (0.084-7.6%)	900 (24-2100)	16000 (700-39000)
Nigeria	2005	Female	75 to 79	49% (35-66%)	51% (34-65%)	1.1 (0.54-1.9)	1.6% (-0.6-4.6%)	1.9% (-0.76-5.5%)	420 (-170-1200)	6000 (-2300-17000)
Nigeria	2005	Female	80 to 84	47% (31-64%)	53% (36-69%)	0.98 (0.52-1.7)	0.95% (-0.81-3.3%)	1.1% (-0.99-3.9%)	130 (-110-440)	1400 (-1100-4600)
Nigeria	2005	Female	85 to 89	47% (32-63%)	53% (37-68%)	0.98 (0.52-1.7)	0.61% (-1-2.8%)	0.66% (-1.3-3.2%)	53 (-110-260)	440 (-740-1900)
Nigeria	2005	Female	90 to 94	50% (35-66%)	50% (34-65%)	1 (0.52-1.8)	0.35% (-1.4-2.5%)	0.3% (-1.8-2.9%)	11 (-63-100)	75 (-280-530)
Nigeria	2005	Female	95 plus	53% (34-73%)	47% (27-66%)	1 (0.46-2)	0.32% (-1.5-3%)	0.19% (-2-3.3%)	2 (-20-34)	13 (-58-120)
Nigeria	2005	Male	All Ages	49% (42-56%)	51% (44-58%)	3.1 (2.4-4.1)	2.9% (2.2-3.8%)	5.5% (4.2-7.4%)	46000 (33000-61000)	180000 (140000-230000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	2005	Male	15 to 19	47% (34-60%)	53% (40-66%)	2.4 (1.4-3.9)	4.6% (2.7-6.9%)	7.1% (3.9-11%)	740 (400-1200)	64000 (38000-97000)
Nigeria	2005	Male	20 to 24	60% (48-71%)	40% (29-52%)	4.6 (2.9-7.1)	10% (7.1-14%)	14% (9.2-20%)	2000 (1300-2900)	150000 (100000-210000)
Nigeria	2005	Male	25 to 29	65% (52-74%)	35% (26-48%)	6.2 (4-8.7)	11% (7.4-15%)	14% (8.6-21%)	2800 (1800-4000)	190000 (130000-260000)
Nigeria	2005	Male	30 to 34	66% (54-76%)	34% (24-46%)	6.8 (4.6-9.6)	9.7% (6.8-13%)	12% (7.6-17%)	2900 (2100-4200)	180000 (130000-250000)
Nigeria	2005	Male	35 to 39	67% (55-76%)	33% (24-45%)	6.6 (4.1-10)	9.7% (6.8-14%)	12% (7.8-17%)	3000 (2100-4200)	170000 (120000-230000)
Nigeria	2005	Male	40 to 44	68% (57-78%)	32% (22-43%)	7 (4.4-11)	11% (8-15%)	14% (9.5-19%)	3600 (2500-4900)	170000 (130000-230000)
Nigeria	2005	Male	45 to 49	70% (59-79%)	30% (21-41%)	7.9 (5.1-12)	14% (10-19%)	18% (13-24%)	4300 (3100-6000)	190000 (140000-250000)
Nigeria	2005	Male	50 to 54	71% (61-80%)	29% (20-39%)	8 (4.8-12)	16% (12-22%)	21% (16-28%)	4700 (3400-6400)	180000 (130000-240000)
Nigeria	2005	Male	55 to 59	70% (60-80%)	30% (20-40%)	7.1 (4.2-10)	17% (13-22%)	22% (17-29%)	4900 (3500-6700)	160000 (120000-210000)
Nigeria	2005	Male	60 to 64	68% (54-80%)	32% (20-46%)	5.7 (3.3-9.2)	15% (10-19%)	19% (13-25%)	4700 (3300-6500)	130000 (93000-180000)
Nigeria	2005	Male	65 to 69	66% (50-79%)	34% (21-50%)	5 (3-7.5)	14% (9.4-18%)	18% (12-23%)	4400 (2900-6100)	100000 (67000-140000)
Nigeria	2005	Male	70 to 74	65% (49-77%)	35% (23-51%)	4.1 (2.4-6.6)	13% (7.6-18%)	16% (9.5-22%)	3800 (2200-5600)	69000 (41000-100000)
Nigeria	2005	Male	75 to 79	61% (44-76%)	39% (24-56%)	3.2 (1.8-5.4)	10% (4.9-16%)	12% (5.9-19%)	2300 (1100-3700)	32000 (16000-52000)
Nigeria	2005	Male	80 to 84	58% (40-74%)	42% (26-60%)	2.5 (1.4-4.3)	8.1% (4-13%)	9.8% (4.6-16%)	890 (420-1500)	9400 (4400-16000)
Nigeria	2005	Male	85 to 89	55% (37-71%)	45% (29-63%)	2 (1.1-3.2)	5.8% (2.6-10%)	6.9% (3.1-12%)	420 (190-760)	3100 (1500-5600)
Nigeria	2005	Male	90 to 94	55% (38-72%)	45% (28-62%)	1.6 (0.83-2.6)	3.6% (1.2-7.2%)	4.2% (1.2-8.5%)	110 (30-220)	560 (180-1100)
Nigeria	2005	Male	95 plus	58% (37-77%)	42% (23-63%)	1.3 (0.54-2.4)	2.6% (0.25-6.3%)	2.9% (-0.043-7.2%)	18 (-0.28-46)	69 (6.7-160)
Nigeria	2010	Female	All Ages	39% (33-45%)	61% (55-67%)	0.68 (0.43-0.98)	0.55% (0.3-0.87%)	0.92% (0.33-1.7%)	6600 (2400-13000)	300000 (160000-480000)
Nigeria	2010	Female	15 to 19	38% (26-50%)	62% (50-74%)	0.78 (0.39-1.4)	1% (0.58-1.6%)	0.9% (0.27-1.9%)	76 (23-170)	14000 (7300-22000)
Nigeria	2010	Female	20 to 24	46% (32-59%)	54% (41-68%)	1.2 (0.61-2.1)	1.8% (1.1-2.8%)	1.3% (0.44-2.8%)	150 (49-300)	26000 (15000-40000)
Nigeria	2010	Female	25 to 29	50% (36-62%)	50% (38-64%)	1.3 (0.68-2.1)	1.6% (0.9-2.5%)	1.2% (0.41-2.4%)	220 (86-430)	29000 (17000-43000)
Nigeria	2010	Female	30 to 34	53% (39-65%)	47% (35-61%)	1.4 (0.74-2.2)	1.3% (0.76-2.1%)	1.1% (0.47-2%)	280 (120-520)	27000 (16000-43000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	2010	Female	35 to 39	56% (42-67%)	44% (33-58%)	1.4 (0.74-2.3)	1.4% (0.78-2.3%)	1.4% (0.62-2.5%)	370 (170-670)	26000 (15000-42000)
Nigeria	2010	Female	40 to 44	57% (45-68%)	43% (32-55%)	1.3 (0.68-2.3)	1.7% (0.96-2.8%)	2% (0.96-3.5%)	490 (230-860)	27000 (15000-45000)
Nigeria	2010	Female	45 to 49	58% (43-70%)	42% (30-57%)	1.1 (0.56-1.9)	1.9% (0.9-3.4%)	2.4% (1.1-4.4%)	530 (230-960)	25000 (12000-43000)
Nigeria	2010	Female	50 to 54	58% (45-70%)	42% (30-55%)	1.1 (0.56-1.9)	2.4% (1.2-4.2%)	3.4% (1.5-5.9%)	660 (290-1200)	26000 (12000-46000)
Nigeria	2010	Female	55 to 59	57% (43-70%)	43% (30-57%)	1.5 (0.79-2.6)	3.5% (1.5-6.1%)	5% (2.3-8.6%)	960 (410-1700)	32000 (14000-57000)
Nigeria	2010	Female	60 to 64	56% (41-70%)	44% (30-59%)	1.6 (0.86-2.8)	3.4% (1.3-6.5%)	4.7% (1.8-8.8%)	1000 (370-2100)	29000 (10000-58000)
Nigeria	2010	Female	65 to 69	56% (40-72%)	44% (28-60%)	1.4 (0.78-2.4)	2.8% (0.74-5.8%)	3.7% (0.99-7.7%)	900 (230-2000)	21000 (5500-45000)
Nigeria	2010	Female	70 to 74	53% (38-69%)	47% (31-62%)	1.2 (0.58-2.1)	1.9% (-0.23-4.9%)	2.4% (-0.34-6.3%)	610 (-85-1600)	11000 (-1500-30000)
Nigeria	2010	Female	75 to 79	49% (34-65%)	51% (35-66%)	0.95 (0.44-1.8)	1% (-0.81-3.7%)	1.2% (-1-4.5%)	260 (-220-980)	3800 (-3100-14000)
Nigeria	2010	Female	80 to 84	47% (31-65%)	53% (35-69%)	0.87 (0.42-1.5)	0.49% (-1-2.6%)	0.55% (-1.3-3.1%)	59 (-150-350)	680 (-1500-3700)
Nigeria	2010	Female	85 to 89	47% (32-64%)	53% (36-68%)	0.87 (0.44-1.6)	0.2% (-1.2-2.1%)	0.16% (-1.6-2.6%)	13 (-130-210)	150 (-930-1600)
Nigeria	2010	Female	90 to 94	50% (34-65%)	50% (35-66%)	0.9 (0.43-1.7)	-0.022% (-1.5-2.2%)	-0.16% (-2-2.6%)	-6.3 (-78-98)	-4.9 (-370-540)
Nigeria	2010	Female	95 plus	53% (34-72%)	47% (28-66%)	0.93 (0.37-1.9)	0.024% (-1.5-2.6%)	-0.18% (-2-2.8%)	-2.1 (-24-33)	1.2 (-70-120)
Nigeria	2010	Male	All Ages	47% (40-54%)	53% (46-60%)	2.7 (1.8-3.8)	2.5% (1.9-3.4%)	4.8% (3.4-6.4%)	38000 (27000-53000)	1600000 (1200000-2100000)
Nigeria	2010	Male	15 to 19	40% (27-55%)	60% (45-73%)	1.8 (0.91-3.4)	3.3% (1.8-5.4%)	4.6% (2.3-8.4%)	520 (250-950)	49000 (27000-80000)
Nigeria	2010	Male	20 to 24	56% (41-69%)	44% (31-59%)	3.7 (2.1-6.2)	8% (5.2-11%)	11% (6.5-16%)	1400 (860-2200)	120000 (75000-170000)
Nigeria	2010	Male	25 to 29	63% (49-74%)	37% (26-51%)	5.4 (3-8.6)	9.5% (6.3-13%)	12% (7.5-18%)	2300 (1400-3300)	160000 (110000-220000)
Nigeria	2010	Male	30 to 34	65% (51-76%)	35% (24-49%)	6 (3.5-9.6)	8.8% (6.2-12%)	11% (6.9-15%)	2500 (1700-3600)	160000 (120000-220000)
Nigeria	2010	Male	35 to 39	66% (54-77%)	34% (23-46%)	6 (3.4-9.4)	8.5% (5.9-12%)	10% (6.8-15%)	2700 (1800-3800)	150000 (110000-210000)
Nigeria	2010	Male	40 to 44	67% (55-78%)	33% (22-45%)	6.3 (3.7-10)	9.6% (7-13%)	12% (8.5-16%)	3000 (2100-4200)	150000 (110000-200000)
Nigeria	2010	Male	45 to 49	69% (56-80%)	31% (20-44%)	7 (3.9-11)	12% (8.6-16%)	16% (11-21%)	3800 (2600-5200)	160000 (120000-220000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	2010	Male	50 to 54	70% (59-80%)	30% (20-41%)	7 (3.9-11)	14% (10-18%)	19% (13-24%)	4100 (2900-5700)	160000 (110000-210000)
Nigeria	2010	Male	55 to 59	70% (58-81%)	30% (19-42%)	6.4 (3.5-10)	15% (10-19%)	20% (14-25%)	4100 (2800-5500)	140000 (96000-180000)
Nigeria	2010	Male	60 to 64	69% (55-80%)	31% (20-45%)	5.3 (2.9-8.9)	13% (8.3-17%)	17% (11-23%)	4100 (2500-5800)	120000 (73000-160000)
Nigeria	2010	Male	65 to 69	67% (51-80%)	33% (20-49%)	4.5 (2.5-7.4)	12% (7-16%)	15% (8.9-21%)	3800 (2300-5500)	88000 (54000-130000)
Nigeria	2010	Male	70 to 74	64% (48-77%)	36% (23-52%)	3.7 (2-6.2)	10% (5.4-16%)	13% (6.8-20%)	3100 (1600-5000)	58000 (30000-92000)
Nigeria	2010	Male	75 to 79	61% (44-76%)	39% (24-56%)	2.9 (1.4-4.9)	7.7% (3.3-13%)	9.6% (4-17%)	1800 (760-3200)	26000 (11000-45000)
Nigeria	2010	Male	80 to 84	58% (39-75%)	42% (25-61%)	2.2 (1.1-4)	6.2% (2.6-12%)	7.4% (3-14%)	700 (280-1300)	7500 (3000-14000)
Nigeria	2010	Male	85 to 89	55% (36-71%)	45% (29-64%)	1.7 (0.89-3.1)	4.4% (1.7-8.8%)	5.2% (1.9-11%)	340 (130-710)	2600 (1000-5300)
Nigeria	2010	Male	90 to 94	55% (38-71%)	45% (29-62%)	1.4 (0.7-2.5)	2.6% (0.46-6.1%)	2.9% (0.38-7.3%)	87 (11-210)	480 (81-1100)
Nigeria	2010	Male	95 plus	57% (37-76%)	43% (24-63%)	1.1 (0.45-2.3)	1.8% (-0.24-5.1%)	1.9% (-0.63-5.8%)	14 (-4.8-46)	57 (-7.8-160)
Nigeria	2016	Female	All Ages	39% (33-45%)	61% (55-67%)	0.58 (0.31-1)	0.58% (0.3-1.1%)	0.9% (0.24-2.1%)	5600 (1500-13000)	280000 (140000-510000)
Nigeria	2016	Female	15 to 19	38% (27-50%)	62% (50-73%)	0.67 (0.27-1.4)	0.92% (0.52-1.6%)	0.67% (0.13-1.7%)	63 (12-170)	14000 (7400-24000)
Nigeria	2016	Female	20 to 24	46% (32-59%)	54% (41-68%)	1 (0.44-2)	1.9% (1.1-2.9%)	1.1% (0.26-2.7%)	120 (27-280)	27000 (15000-43000)
Nigeria	2016	Female	25 to 29	50% (36-62%)	50% (38-64%)	1.1 (0.48-2)	1.8% (1-2.8%)	1.2% (0.37-2.7%)	180 (57-390)	28000 (16000-44000)
Nigeria	2016	Female	30 to 34	53% (40-65%)	47% (35-60%)	1.2 (0.51-2.2)	1.6% (0.92-2.5%)	1.3% (0.53-2.6%)	240 (95-500)	27000 (16000-45000)
Nigeria	2016	Female	35 to 39	56% (42-68%)	44% (32-58%)	1.2 (0.55-2.2)	1.7% (0.97-2.8%)	1.8% (0.77-3.4%)	350 (150-700)	27000 (15000-46000)
Nigeria	2016	Female	40 to 44	57% (45-69%)	43% (31-55%)	1.1 (0.53-2.1)	2% (1.1-3.3%)	2.5% (1.2-4.6%)	470 (210-900)	28000 (14000-49000)
Nigeria	2016	Female	45 to 49	58% (44-70%)	42% (30-56%)	0.92 (0.41-1.9)	2% (1-3.8%)	2.9% (1.2-5.6%)	490 (200-1000)	24000 (11000-45000)
Nigeria	2016	Female	50 to 54	58% (45-70%)	42% (30-55%)	0.96 (0.41-1.9)	2.4% (1.2-4.6%)	3.6% (1.7-6.9%)	610 (270-1200)	24000 (11000-46000)
Nigeria	2016	Female	55 to 59	57% (42-70%)	43% (30-58%)	1.3 (0.61-2.5)	3.1% (1.2-6.2%)	4.8% (1.8-9.3%)	850 (300-1700)	28000 (10000-57000)
Nigeria	2016	Female	60 to 64	56% (41-71%)	44% (29-59%)	1.4 (0.65-2.7)	2.9% (0.83-6.3%)	4.2% (1.3-9.3%)	870 (250-2000)	24000 (6700-56000)
Nigeria	2016	Female	65 to 69	56% (41-71%)	44% (29-59%)	1.2 (0.55-2.4)	2.2% (0.22-5.7%)	3% (0.36-7.6%)	730 (87-1900)	17000 (1700-45000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	2016	Female	70 to 74	54% (39-68%)	46% (32-61%)	1 (0.43-2)	1.4% (-0.55-5%)	1.8% (-0.69-6.5%)	480 (-200-1700)	8900 (-3600-33000)
Nigeria	2016	Female	75 to 79	49% (33-66%)	51% (34-67%)	0.83 (0.32-1.7)	0.63% (-1-3.6%)	0.78% (-1.3-4.4%)	180 (-310-1100)	2700 (-4300-15000)
Nigeria	2016	Female	80 to 84	47% (30-66%)	53% (34-70%)	0.76 (0.31-1.6)	0.18% (-1.2-2.7%)	0.19% (-1.6-3.3%)	22 (-180-370)	280 (-1900-4000)
Nigeria	2016	Female	85 to 89	47% (31-63%)	53% (37-69%)	0.76 (0.31-1.5)	-0.066% (-1.5-2.2%)	-0.16% (-1.8-2.5%)	-13 (-170-220)	-51 (-1200-1800)
Nigeria	2016	Female	90 to 94	50% (35-66%)	50% (34-65%)	0.79 (0.32-1.7)	-0.26% (-1.6-2.1%)	-0.45% (-2.1-2.4%)	-21 (-97-110)	-75 (-460-620)
Nigeria	2016	Female	95 plus	53% (34-72%)	47% (28-66%)	0.81 (0.27-1.8)	-0.18% (-1.6-2.4%)	-0.42% (-2.2-2.7%)	-6.5 (-34-43)	-11 (-100-160)
Nigeria	2016	Male	All Ages	47% (40-55%)	53% (45-60%)	2.4 (1.2-4.2)	2.5% (1.7-3.5%)	4.7% (2.9-6.7%)	34000 (21000-50000)	1400000 (930000-2000000)
Nigeria	2016	Male	15 to 19	40% (28-54%)	60% (46-72%)	1.6 (0.66-3.2)	2.7% (1.4-4.8%)	3.5% (1.4-7%)	450 (180-910)	45000 (23000-82000)
Nigeria	2016	Male	20 to 24	56% (41-69%)	44% (31-59%)	3.2 (1.5-6.3)	6.7% (3.9-9.9%)	8.7% (4.2-14%)	1200 (580-2000)	100000 (59000-160000)
Nigeria	2016	Male	25 to 29	63% (49-74%)	37% (26-51%)	4.7 (2.2-8.6)	8.4% (5-12%)	11% (5.8-17%)	1900 (1000-2900)	140000 (84000-200000)
Nigeria	2016	Male	30 to 34	66% (51-76%)	34% (24-49%)	5.2 (2.5-9.7)	8.3% (5.4-12%)	10% (6.2-15%)	2300 (1400-3300)	150000 (97000-210000)
Nigeria	2016	Male	35 to 39	66% (53-77%)	34% (23-47%)	5.2 (2.6-9.4)	8.3% (5.5-12%)	11% (6.6-15%)	2500 (1600-3700)	150000 (95000-210000)
Nigeria	2016	Male	40 to 44	67% (55-78%)	33% (22-45%)	5.5 (2.8-9.8)	9.1% (6.2-12%)	12% (7.8-16%)	2900 (1900-4300)	150000 (97000-210000)
Nigeria	2016	Male	45 to 49	69% (55-79%)	31% (21-45%)	6 (2.8-11)	11% (7.5-16%)	15% (9.9-20%)	3500 (2200-5000)	150000 (98000-220000)
Nigeria	2016	Male	50 to 54	70% (58-80%)	30% (20-42%)	6.1 (2.9-11)	13% (8.3-17%)	17% (11-22%)	3800 (2300-5300)	150000 (92000-200000)
Nigeria	2016	Male	55 to 59	70% (57-81%)	30% (19-43%)	5.5 (2.7-10)	13% (8-17%)	17% (11-23%)	3700 (2300-5300)	130000 (78000-180000)
Nigeria	2016	Male	60 to 64	69% (55-81%)	31% (19-45%)	4.6 (2.1-9.2)	11% (6.1-15%)	14% (8.1-20%)	3400 (1800-5200)	98000 (53000-140000)
Nigeria	2016	Male	65 to 69	67% (51-80%)	33% (20-49%)	3.9 (1.8-7.5)	9.4% (4.8-14%)	12% (6.1-19%)	3200 (1500-5000)	75000 (36000-120000)
Nigeria	2016	Male	70 to 74	64% (48-77%)	36% (23-52%)	3.2 (1.5-6.1)	8.1% (3.4-14%)	10% (4.4-18%)	2700 (1200-4800)	51000 (22000-90000)
Nigeria	2016	Male	75 to 79	61% (44-76%)	39% (24-56%)	2.5 (1-4.7)	6% (2.1-12%)	7.5% (2.6-15%)	1600 (540-3200)	23000 (8000-46000)
Nigeria	2016	Male	80 to 84	58% (40-74%)	42% (26-60%)	1.9 (0.82-3.8)	4.8% (1.5-10%)	5.8% (1.8-13%)	600 (180-1300)	6400 (2000-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Nigeria	2016	Male	85 to 89	55% (37-71%)	45% (29-63%)	1.5 (0.65-2.9)	3.4% (0.85-7.8%)	3.9% (0.9-9.5%)	290 (65-690)	2200 (570-5200)
Nigeria	2016	Male	90 to 94	55% (38-72%)	45% (28-62%)	1.2 (0.49-2.4)	1.9% (-0.025-5.7%)	2.1% (-0.27-6.7%)	73 (-9.4-230)	410 (-5.5-1200)
Nigeria	2016	Male	95 plus	58% (37-77%)	42% (23-63%)	0.99 (0.35-2.2)	1.2% (-0.51-4.5%)	1.2% (-0.94-5.2%)	11 (-9.1-50)	50 (-20-180)
Sao Tome and Principe	1990	Female	All Ages	12% (8.9-16%)	88% (84-91%)	0.94 (0.71-1.2)	1.6% (1.2-2.1%)	3.6% (2.3-5%)	19 (12-27)	510 (380-660)
Sao Tome and Principe	1990	Female	15 to 19	11% (5.3-21%)	89% (79-95%)	0.85 (0.31-1.7)	1.6% (0.87-2.6%)	1.9% (0.71-3.5%)	0.14 (0.053-0.28)	17 (8.8-28)
Sao Tome and Principe	1990	Female	20 to 24	15% (6.5-26%)	85% (74-93%)	1.3 (0.52-2.6)	2.9% (1.8-4.2%)	2.9% (1.3-5%)	0.25 (0.11-0.44)	29 (17-44)
Sao Tome and Principe	1990	Female	25 to 29	17% (7.3-29%)	83% (71-93%)	1.7 (0.69-3.1)	3.2% (2.1-4.6%)	3.6% (1.9-5.6%)	0.3 (0.15-0.48)	29 (18-42)
Sao Tome and Principe	1990	Female	30 to 34	19% (8.2-33%)	81% (67-92%)	1.9 (0.81-3.4)	3.8% (2.5-5.3%)	4.8% (2.8-7.2%)	0.37 (0.22-0.58)	29 (19-43)
Sao Tome and Principe	1990	Female	35 to 39	21% (9.3-36%)	79% (64-91%)	2.1 (0.83-3.9)	4.5% (2.7-6.3%)	6.2% (3.5-9%)	0.44 (0.25-0.65)	27 (17-39)
Sao Tome and Principe	1990	Female	40 to 44	22% (11-39%)	78% (61-89%)	2.3 (0.86-4.5)	5.2% (3.3-7.4%)	7.5% (4.7-11%)	0.48 (0.29-0.7)	25 (15-36)
Sao Tome and Principe	1990	Female	45 to 49	23% (9.9-39%)	77% (61-90%)	2.4 (0.85-4.9)	5.8% (3.5-8.5%)	8.4% (4.9-12%)	0.66 (0.38-0.98)	29 (18-43)
Sao Tome and Principe	1990	Female	50 to 54	24% (11-41%)	76% (59-89%)	2.5 (0.99-4.9)	6.5% (4-9.4%)	9% (5.6-13%)	1.2 (0.7-1.7)	45 (28-66)
Sao Tome and Principe	1990	Female	55 to 59	24% (12-42%)	76% (58-88%)	2.7 (1-5.4)	6.7% (4.1-9.9%)	9% (5.4-13%)	1.6 (0.97-2.5)	53 (32-80)
Sao Tome and Principe	1990	Female	60 to 64	25% (12-44%)	75% (56-88%)	2.8 (1-5.7)	6.5% (3.9-10%)	8.3% (4.9-13%)	2.1 (1.2-3.2)	58 (34-89)
Sao Tome and Principe	1990	Female	65 to 69	26% (13-43%)	74% (57-87%)	2.6 (1-5.1)	6.8% (3.8-11%)	8.4% (4.6-13%)	2.7 (1.4-4.3)	62 (34-99)
Sao Tome and Principe	1990	Female	70 to 74	24% (13-41%)	76% (59-87%)	2.1 (0.72-4.6)	6.2% (3-10%)	7.5% (3.6-12%)	2.6 (1.2-4.5)	49 (22-82)
Sao Tome and Principe	1990	Female	75 to 79	21% (9.8-39%)	79% (61-90%)	1.6 (0.47-3.9)	4.8% (1.8-8.6%)	5.7% (2.2-10%)	2 (0.74-3.6)	28 (11-51)
Sao Tome and Principe	1990	Female	80 to 84	19% (8.4-35%)	81% (65-92%)	1.4 (0.44-3.2)	4.3% (1.5-8%)	5.1% (1.8-9.6%)	1.7 (0.61-3.3)	18 (6.5-35)
Sao Tome and Principe	1990	Female	85 to 89	20% (9.2-35%)	80% (65-91%)	1.4 (0.44-3.5)	4% (1.3-8%)	4.7% (1.4-9.4%)	1.4 (0.42-2.8)	10 (3.2-20)
Sao Tome and Principe	1990	Female	90 to 94	22% (10-39%)	78% (61-90%)	1.6 (0.44-3.7)	3.8% (0.93-7.7%)	4.4% (1-9.3%)	0.6 (0.14-1.3)	3.1 (0.75-6.3)
Sao Tome and Principe	1990	Female	95 plus	27% (10-49%)	73% (51-90%)	1.8 (0.4-4.7)	3.8% (0.55-8.7%)	4.5% (0.55-10%)	0.17 (0.021-0.39)	0.57 (0.083-1.3)
Sao Tome and Principe	1990	Male	All Ages	9.2% (6.8-12%)	91% (88-93%)	1 (0.77-1.3)	2% (1.5-2.4%)	3.8% (2.8-4.8%)	21 (15-27)	700 (550-870)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	1990	Male	15 to 19	7.5% (3.5-14%)	93% (86-97%)	0.64 (0.22-1.3)	2% (1.1-3.3%)	2.4% (1.1-4.4%)	0.21 (0.092-0.39)	22 (12-36)
Sao Tome and Principe	1990	Male	20 to 24	12% (5.1-21%)	88% (79-95%)	1.3 (0.51-2.6)	4.6% (2.8-7%)	5.2% (2.7-8.7%)	0.56 (0.29-0.95)	51 (30-79)
Sao Tome and Principe	1990	Male	25 to 29	15% (6.9-27%)	85% (73-93%)	2 (0.83-3.8)	6.5% (3.9-9.4%)	8% (4.1-12%)	0.76 (0.38-1.2)	59 (35-86)
Sao Tome and Principe	1990	Male	30 to 34	18% (7.6-32%)	82% (68-92%)	2.6 (1.1-4.9)	7.7% (4.6-11%)	10% (5.3-15%)	0.81 (0.4-1.2)	55 (31-78)
Sao Tome and Principe	1990	Male	35 to 39	20% (9-35%)	80% (65-91%)	2.9 (1.1-5.5)	8.8% (5.1-12%)	12% (6.3-17%)	0.87 (0.46-1.3)	50 (29-71)
Sao Tome and Principe	1990	Male	40 to 44	21% (10-37%)	79% (63-90%)	3.2 (1.2-6.2)	9.7% (6.2-13%)	14% (8.5-18%)	0.85 (0.52-1.2)	42 (27-58)
Sao Tome and Principe	1990	Male	45 to 49	21% (9.4-36%)	79% (64-91%)	3.3 (1.2-7)	11% (7.3-15%)	15% (9.8-20%)	1.2 (0.75-1.6)	51 (33-70)
Sao Tome and Principe	1990	Male	50 to 54	22% (10-38%)	78% (62-90%)	3.3 (1.3-7)	9.4% (6.2-13%)	13% (8.4-17%)	2 (1.3-2.9)	76 (50-110)
Sao Tome and Principe	1990	Male	55 to 59	22% (10-38%)	78% (62-90%)	3.2 (1.2-6.5)	9.1% (6.2-12%)	12% (8.1-16%)	2.4 (1.6-3.5)	80 (54-110)
Sao Tome and Principe	1990	Male	60 to 64	21% (10-38%)	79% (62-90%)	2.9 (1-5.9)	7.7% (5.1-11%)	9.7% (6.4-14%)	2.6 (1.6-3.7)	71 (46-100)
Sao Tome and Principe	1990	Male	65 to 69	21% (10-36%)	79% (64-90%)	2.4 (0.88-4.9)	6.7% (4.1-9.8%)	8.3% (5.1-12%)	2.6 (1.6-3.9)	61 (37-90)
Sao Tome and Principe	1990	Male	70 to 74	20% (10-34%)	80% (66-90%)	1.8 (0.62-4)	5.6% (3-8.8%)	6.8% (3.6-11%)	2.3 (1.2-3.7)	43 (22-68)
Sao Tome and Principe	1990	Male	75 to 79	18% (8.6-32%)	82% (68-91%)	1.4 (0.39-3.2)	4.2% (1.8-7.3%)	5% (2.1-8.6%)	1.7 (0.68-2.9)	24 (9.7-42)
Sao Tome and Principe	1990	Male	80 to 84	16% (7.1-30%)	84% (70-93%)	1.1 (0.33-2.4)	3.5% (1.4-6.3%)	4.1% (1.7-7.4%)	1.1 (0.46-2)	12 (4.9-21)
Sao Tome and Principe	1990	Male	85 to 89	15% (6.7-28%)	85% (72-93%)	0.93 (0.27-2.2)	3% (1-5.8%)	3.5% (1.2-6.8%)	0.62 (0.21-1.2)	4.6 (1.7-9)
Sao Tome and Principe	1990	Male	90 to 94	16% (7.3-30%)	84% (70-93%)	0.89 (0.24-2.3)	2.3% (0.63-4.9%)	2.7% (0.67-5.9%)	0.18 (0.046-0.4)	0.96 (0.26-2.1)
Sao Tome and Principe	1990	Male	95 plus	20% (7.6-39%)	80% (61-92%)	0.93 (0.19-2.6)	2% (0.33-4.8%)	2.2% (0.25-5.5%)	0.035 (0.004-0.088)	0.13 (0.021-0.31)
Sao Tome and Principe	1995	Female	All Ages	12% (9.2-17%)	88% (83-91%)	1.3 (1-1.7)	2.1% (1.5-2.7%)	4.6% (3.1-6.2%)	26 (17-36)	690 (480-890)
Sao Tome and Principe	1995	Female	15 to 19	12% (5.8-21%)	88% (79-94%)	1.2 (0.45-2.3)	1.9% (1-3%)	2.4% (1-4.1%)	0.26 (0.11-0.46)	28 (15-45)
Sao Tome and Principe	1995	Female	20 to 24	15% (6.7-28%)	85% (72-93%)	1.9 (0.73-3.6)	3.2% (1.9-4.7%)	3.3% (1.6-5.7%)	0.34 (0.16-0.58)	37 (22-55)
Sao Tome and Principe	1995	Female	25 to 29	17% (7.7-30%)	83% (70-92%)	2.3 (1-4.3)	3.4% (1.6-5%)	3.8% (0.58-6.2%)	0.35 (0.053-0.57)	34 (17-50)
Sao Tome and Principe	1995	Female	30 to 34	19% (8.6-35%)	81% (65-91%)	2.7 (1.2-4.9)	4% (1.4-5.9%)	5.1% (0.65-7.8%)	0.49 (0.063-0.79)	37 (14-56)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	1995	Female	35 to 39	22% (10-39%)	78% (61-90%)	3 (1.2-5.4)	4.8% (1.4-7%)	6.6% (0.85-10%)	0.65 (0.086-1)	40 (12-59)
Sao Tome and Principe	1995	Female	40 to 44	23% (11-39%)	77% (61-89%)	3.2 (1.2-6.2)	5.5% (1.5-8.1%)	7.9% (1.1-12%)	0.69 (0.096-1)	36 (10-52)
Sao Tome and Principe	1995	Female	45 to 49	24% (11-40%)	76% (60-89%)	3.3 (1.2-7.2)	6.4% (2.7-9.5%)	9% (3.3-14%)	0.76 (0.29-1.2)	34 (15-52)
Sao Tome and Principe	1995	Female	50 to 54	24% (11-43%)	76% (57-89%)	3.5 (1.4-7.3)	7.3% (3.9-11%)	9.8% (5-14%)	1.2 (0.58-1.8)	44 (24-66)
Sao Tome and Principe	1995	Female	55 to 59	25% (12-44%)	75% (56-88%)	3.8 (1.4-7.7)	7.7% (4.6-11%)	9.9% (6.1-15%)	2 (1.2-3.1)	67 (40-100)
Sao Tome and Principe	1995	Female	60 to 64	26% (13-45%)	74% (55-87%)	3.9 (1.4-8.2)	7.5% (4.8-11%)	9.3% (5.9-14%)	2.7 (1.6-4.2)	76 (46-120)
Sao Tome and Principe	1995	Female	65 to 69	27% (14-45%)	73% (55-86%)	3.6 (1.4-7)	8% (4.7-12%)	9.7% (5.6-15%)	3.8 (2.2-6)	88 (50-140)
Sao Tome and Principe	1995	Female	70 to 74	25% (13-42%)	75% (58-87%)	2.9 (0.98-6.3)	7.6% (4-12%)	9% (4.8-14%)	4.4 (2.3-7)	80 (43-130)
Sao Tome and Principe	1995	Female	75 to 79	22% (9.9-39%)	78% (61-90%)	2.3 (0.67-5.3)	6% (2.9-9.8%)	7% (3.3-12%)	3.2 (1.4-5.3)	45 (21-75)
Sao Tome and Principe	1995	Female	80 to 84	20% (8.8-35%)	80% (65-91%)	2 (0.59-4.4)	5.4% (2.3-9.3%)	6.4% (2.7-11%)	2.4 (0.98-4.3)	25 (10-45)
Sao Tome and Principe	1995	Female	85 to 89	20% (9.3-36%)	80% (64-91%)	1.9 (0.59-4.6)	4.9% (2-9.1%)	5.8% (2.3-11%)	1.7 (0.67-3.2)	13 (5-24)
Sao Tome and Principe	1995	Female	90 to 94	23% (10-40%)	77% (60-90%)	2.1 (0.59-5.2)	4.6% (1.7-8.9%)	5.5% (1.9-11%)	0.85 (0.29-1.7)	4.3 (1.5-8.5)
Sao Tome and Principe	1995	Female	95 plus	27% (11-50%)	73% (50-89%)	2.4 (0.51-6.3)	4.7% (1.1-9.6%)	5.5% (1.2-11%)	0.27 (0.061-0.58)	0.94 (0.21-1.9)
Sao Tome and Principe	1995	Male	All Ages	9.6% (7.1-13%)	90% (87-93%)	1.5 (1.1-1.8)	2.2% (1.2-2.8%)	4.3% (2.6-5.6%)	24 (14-32)	770 (420-1000)
Sao Tome and Principe	1995	Male	15 to 19	7.8% (3.7-14%)	92% (86-96%)	0.9 (0.33-1.8)	2.3% (1.3-3.6%)	2.9% (1.4-5%)	0.32 (0.15-0.56)	32 (18-50)
Sao Tome and Principe	1995	Male	20 to 24	12% (5.4-22%)	88% (78-95%)	1.9 (0.74-3.6)	4.7% (2.7-7.2%)	5.5% (2.4-9.2%)	0.62 (0.26-1.1)	56 (31-89)
Sao Tome and Principe	1995	Male	25 to 29	16% (7.1-28%)	84% (72-93%)	2.8 (1.2-5.2)	6% (2.1-9.5%)	7.3% (0.91-12%)	0.79 (0.098-1.4)	63 (22-100)
Sao Tome and Principe	1995	Male	30 to 34	18% (8.3-33%)	82% (67-92%)	3.6 (1.6-6.5)	6.8% (2-11%)	8.7% (1.1-15%)	0.85 (0.1-1.5)	60 (18-95)
Sao Tome and Principe	1995	Male	35 to 39	20% (9.4-36%)	80% (64-91%)	4.1 (1.7-7.6)	7.9% (1.9-12%)	11% (1.3-17%)	0.95 (0.11-1.6)	56 (14-88)
Sao Tome and Principe	1995	Male	40 to 44	21% (11-38%)	79% (62-89%)	4.5 (1.7-8.9)	8.9% (1.9-13%)	12% (1.5-19%)	0.99 (0.12-1.5)	50 (11-76)
Sao Tome and Principe	1995	Male	45 to 49	22% (9.8-37%)	78% (63-90%)	4.8 (1.8-11)	10% (2.1-15%)	14% (1.8-21%)	1.1 (0.13-1.6)	47 (9.4-69)
Sao Tome and Principe	1995	Male	50 to 54	22% (10-40%)	78% (60-90%)	4.9 (1.8-11)	9.6% (2.8-14%)	13% (2.8-19%)	1.2 (0.27-1.9)	48 (14-70)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	1995	Male	55 to 59	23% (11-40%)	77% (60-89%)	4.7 (1.7-9.6)	9.7% (4.9-14%)	13% (6-18%)	2.6 (1.2-3.7)	84 (41-120)
Sao Tome and Principe	1995	Male	60 to 64	23% (11-40%)	77% (60-89%)	4.2 (1.5-9)	8.6% (5.4-12%)	11% (6.7-15%)	3.1 (1.8-4.4)	86 (50-120)
Sao Tome and Principe	1995	Male	65 to 69	22% (11-38%)	78% (62-89%)	3.5 (1.3-7.2)	7.6% (4.7-11%)	9.4% (5.8-14%)	3.2 (1.9-4.7)	73 (44-110)
Sao Tome and Principe	1995	Male	70 to 74	21% (11-36%)	79% (64-89%)	2.6 (0.89-5.6)	6.7% (3.9-10%)	8.1% (4.7-12%)	3.2 (1.9-5)	60 (35-92)
Sao Tome and Principe	1995	Male	75 to 79	19% (9.1-34%)	81% (66-91%)	2 (0.57-4.7)	5.2% (2.5-8.6%)	6.2% (2.9-10%)	2.3 (1.1-3.8)	32 (16-53)
Sao Tome and Principe	1995	Male	80 to 84	17% (7.3-32%)	83% (68-93%)	1.6 (0.49-3.6)	4.5% (2.1-7.5%)	5.3% (2.4-8.8%)	1.5 (0.68-2.5)	16 (7.3-26)
Sao Tome and Principe	1995	Male	85 to 89	16% (7.1-30%)	84% (70-93%)	1.4 (0.4-3.3)	4% (1.6-7.1%)	4.6% (1.9-8.4%)	0.9 (0.36-1.6)	6.7 (2.8-12)
Sao Tome and Principe	1995	Male	90 to 94	17% (7.9-32%)	83% (68-92%)	1.3 (0.35-3.3)	3.2% (1-6.2%)	3.7% (1.1-7.3%)	0.31 (0.096-0.61)	1.6 (0.53-3.1)
Sao Tome and Principe	1995	Male	95 plus	21% (8.2-41%)	79% (59-92%)	1.4 (0.27-3.8)	2.8% (0.62-5.9%)	3.1% (0.61-6.9%)	0.066 (0.013-0.14)	0.23 (0.053-0.5)
Sao Tome and Principe	2000	Female	All Ages	13% (9.4-17%)	87% (83-91%)	1.6 (1.2-2)	2.5% (1.8-3.3%)	5.4% (3.7-7.3%)	32 (21-43)	820 (560-1100)
Sao Tome and Principe	2000	Female	15 to 19	12% (5.8-22%)	88% (78-94%)	1.4 (0.53-2.8)	2% (1.2-3.1%)	2.7% (1.2-4.6%)	0.23 (0.11-0.41)	26 (15-41)
Sao Tome and Principe	2000	Female	20 to 24	16% (7-28%)	84% (72-93%)	2.2 (0.89-4.1)	3.3% (2.1-4.9%)	3.6% (1.7-6.1%)	0.45 (0.2-0.76)	49 (30-74)
Sao Tome and Principe	2000	Female	25 to 29	18% (7.6-32%)	82% (68-92%)	2.8 (1.2-5.2)	3.4% (1.7-5%)	3.8% (0.56-6.4%)	0.36 (0.052-0.61)	36 (18-53)
Sao Tome and Principe	2000	Female	30 to 34	20% (8.6-37%)	80% (63-91%)	3.3 (1.4-5.9)	4% (1.5-6%)	5.2% (0.64-8.3%)	0.51 (0.061-0.82)	39 (14-59)
Sao Tome and Principe	2000	Female	35 to 39	23% (10-39%)	77% (61-90%)	3.6 (1.5-6.5)	4.9% (1.4-7.4%)	6.9% (0.83-11%)	0.78 (0.094-1.3)	48 (14-74)
Sao Tome and Principe	2000	Female	40 to 44	24% (12-42%)	76% (58-88%)	3.8 (1.5-7.4)	5.7% (1.5-8.4%)	8.1% (1-12%)	0.98 (0.13-1.5)	51 (13-76)
Sao Tome and Principe	2000	Female	45 to 49	25% (11-42%)	75% (58-89%)	4 (1.5-8.6)	6.8% (1.8-10%)	9.5% (1.7-14%)	1.1 (0.21-1.8)	50 (13-75)
Sao Tome and Principe	2000	Female	50 to 54	25% (12-44%)	75% (56-88%)	4.2 (1.6-8.9)	8% (4.2-12%)	11% (5.6-15%)	1.4 (0.65-2)	51 (26-76)
Sao Tome and Principe	2000	Female	55 to 59	26% (12-44%)	74% (56-88%)	4.7 (1.7-9.3)	8.4% (5.2-12%)	11% (6.4-16%)	1.9 (1.1-2.8)	62 (37-92)
Sao Tome and Principe	2000	Female	60 to 64	27% (13-45%)	73% (55-87%)	4.7 (1.7-9.8)	8.2% (5.3-12%)	10% (6.4-15%)	3.2 (2-4.8)	88 (56-130)
Sao Tome and Principe	2000	Female	65 to 69	27% (14-44%)	73% (56-86%)	4.3 (1.6-8.5)	8.7% (5.3-13%)	10% (6.3-15%)	4.6 (2.7-6.8)	100 (62-150)
Sao Tome and Principe	2000	Female	70 to 74	25% (13-42%)	75% (58-87%)	3.4 (1.1-7.7)	8.2% (4.5-13%)	9.6% (5.4-15%)	5.4 (2.9-8.3)	98 (53-150)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	2000	Female	75 to 79	22% (11-39%)	78% (61-89%)	2.6 (0.76-6.4)	6.5% (3.2-11%)	7.6% (3.8-13%)	4.5 (2.2-7.5)	64 (31-110)
Sao Tome and Principe	2000	Female	80 to 84	20% (9.1-36%)	80% (64-91%)	2.3 (0.67-5.2)	6% (2.8-10%)	7% (3.2-12%)	3.3 (1.5-5.7)	35 (16-60)
Sao Tome and Principe	2000	Female	85 to 89	20% (9.4-37%)	80% (63-91%)	2.3 (0.72-5.6)	5.5% (2.5-9.7%)	6.4% (2.9-11%)	1.9 (0.84-3.3)	14 (6.2-24)
Sao Tome and Principe	2000	Female	90 to 94	23% (10-40%)	77% (60-90%)	2.5 (0.69-6.1)	5.3% (2.1-9.5%)	6.2% (2.4-11%)	0.88 (0.35-1.6)	4.5 (1.8-8.2)
Sao Tome and Principe	2000	Female	95 plus	28% (11-50%)	72% (50-89%)	2.9 (0.65-7.6)	5.4% (1.7-10%)	6.3% (1.9-12%)	0.33 (0.099-0.65)	1.1 (0.36-2.2)
Sao Tome and Principe	2000	Male	All Ages	10% (7.6-14%)	90% (86-92%)	1.9 (1.4-2.3)	2.6% (1.5-3.4%)	5% (3.2-6.6%)	29 (18-38)	920 (530-1200)
Sao Tome and Principe	2000	Male	15 to 19	8.3% (4.1-15%)	92% (85-96%)	1.1 (0.43-2.3)	2.6% (1.6-4.3%)	3.5% (1.7-6.2%)	0.35 (0.17-0.63)	34 (19-56)
Sao Tome and Principe	2000	Male	20 to 24	13% (5.8-24%)	87% (76-94%)	2.4 (0.96-4.5)	5.1% (2.9-8%)	6.1% (2.8-10%)	0.93 (0.41-1.6)	82 (45-130)
Sao Tome and Principe	2000	Male	25 to 29	16% (7.7-30%)	84% (70-92%)	3.6 (1.5-6.5)	6.1% (2.2-9.6%)	7.5% (0.99-13%)	0.89 (0.12-1.6)	71 (25-110)
Sao Tome and Principe	2000	Male	30 to 34	19% (8.7-35%)	81% (65-91%)	4.6 (2-8)	6.6% (2.1-11%)	8.3% (1.1-15%)	0.98 (0.13-1.8)	69 (21-110)
Sao Tome and Principe	2000	Male	35 to 39	22% (10-38%)	78% (62-90%)	5.1 (2.2-9.4)	7.6% (2-13%)	10% (1.4-18%)	1.2 (0.16-2.1)	70 (18-120)
Sao Tome and Principe	2000	Male	40 to 44	22% (11-39%)	78% (61-89%)	5.6 (2.2-11)	8.8% (2-14%)	12% (1.6-20%)	1.3 (0.17-2.2)	66 (15-110)
Sao Tome and Principe	2000	Male	45 to 49	23% (11-39%)	77% (61-89%)	5.9 (2.2-13)	11% (2.1-16%)	15% (1.8-22%)	1.5 (0.18-2.3)	67 (13-100)
Sao Tome and Principe	2000	Male	50 to 54	23% (11-41%)	77% (59-89%)	6 (2.2-13)	10% (2.2-15%)	14% (1.8-20%)	1.4 (0.19-2)	52 (11-77)
Sao Tome and Principe	2000	Male	55 to 59	23% (11-41%)	77% (59-89%)	5.8 (2-12)	11% (5.5-15%)	14% (7.1-19%)	1.8 (0.87-2.6)	60 (30-83)
Sao Tome and Principe	2000	Male	60 to 64	23% (11-40%)	77% (60-89%)	5.2 (1.7-11)	9.4% (5.7-13%)	12% (7.1-16%)	3.6 (2.1-5)	99 (59-140)
Sao Tome and Principe	2000	Male	65 to 69	23% (12-39%)	77% (61-88%)	4.3 (1.6-9.2)	8.5% (5.4-12%)	10% (6.6-15%)	4.1 (2.5-5.9)	94 (59-140)
Sao Tome and Principe	2000	Male	70 to 74	22% (11-38%)	78% (62-89%)	3.3 (1.1-7.2)	7.5% (4.4-11%)	9% (5.3-13%)	4 (2.4-6)	74 (44-110)
Sao Tome and Principe	2000	Male	75 to 79	20% (9.9-35%)	80% (65-90%)	2.5 (0.69-5.8)	5.9% (3-9.3%)	7% (3.6-11%)	3.2 (1.7-5.1)	46 (24-73)
Sao Tome and Principe	2000	Male	80 to 84	18% (8.1-32%)	82% (68-92%)	2 (0.59-4.7)	5.1% (2.6-8.4%)	5.9% (2.9-9.7%)	2 (1-3.3)	21 (11-35)
Sao Tome and Principe	2000	Male	85 to 89	17% (7.7-31%)	83% (69-92%)	1.7 (0.5-4.1)	4.5% (2-7.8%)	5.3% (2.4-9.2%)	1.1 (0.5-1.9)	8.3 (3.8-14)
Sao Tome and Principe	2000	Male	90 to 94	18% (8.5-33%)	82% (67-91%)	1.6 (0.42-4)	3.7% (1.5-6.8%)	4.3% (1.7-8%)	0.4 (0.15-0.75)	2.1 (0.82-3.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	2000	Male	95 plus	22% (8.6-41%)	78% (59-91%)	1.7 (0.34-4.6)	3.2% (0.86-6.6%)	3.7% (0.88-7.6%)	0.095 (0.023-0.2)	0.34 (0.089-0.67)
Sao Tome and Principe	2005	Female	All Ages	13% (9.8-18%)	87% (82-90%)	1.7 (1.3-2.1)	2.7% (1.8-3.4%)	5.6% (3.6-7.5%)	31 (20-43)	820 (550-1000)
Sao Tome and Principe	2005	Female	15 to 19	12% (5.9-23%)	88% (77-94%)	1.4 (0.56-2.9)	2% (1.1-3.3%)	2.7% (1.1-4.9%)	0.24 (0.093-0.43)	28 (16-45)
Sao Tome and Principe	2005	Female	20 to 24	16% (7.2-29%)	84% (71-93%)	2.3 (0.89-4.2)	3.4% (2.1-5%)	3.9% (1.5-6.5%)	0.35 (0.13-0.59)	41 (25-62)
Sao Tome and Principe	2005	Female	25 to 29	18% (8.1-32%)	82% (68-92%)	2.8 (1.2-5.2)	3.5% (1.8-5.2%)	3.9% (0.58-6.8%)	0.44 (0.065-0.78)	47 (23-71)
Sao Tome and Principe	2005	Female	30 to 34	21% (9.2-36%)	79% (64-91%)	3.3 (1.4-5.7)	4% (1.5-6.1%)	5.3% (0.72-8.7%)	0.5 (0.067-0.82)	40 (16-61)
Sao Tome and Principe	2005	Female	35 to 39	23% (11-39%)	77% (61-89%)	3.7 (1.6-6.5)	4.8% (1.5-7.3%)	7% (0.89-11%)	0.73 (0.094-1.2)	46 (15-71)
Sao Tome and Principe	2005	Female	40 to 44	24% (13-42%)	76% (58-87%)	3.9 (1.5-7.5)	5.7% (1.5-8.3%)	8.4% (1.1-13%)	1.1 (0.14-1.7)	56 (15-84)
Sao Tome and Principe	2005	Female	45 to 49	25% (12-41%)	75% (59-88%)	4.1 (1.5-8.4)	6.8% (2.3-10%)	9.7% (2.8-14%)	1.4 (0.39-2.2)	64 (22-95)
Sao Tome and Principe	2005	Female	50 to 54	26% (12-45%)	74% (55-88%)	4.3 (1.7-8.5)	7.9% (4.1-12%)	11% (5.5-16%)	1.6 (0.83-2.4)	63 (34-92)
Sao Tome and Principe	2005	Female	55 to 59	26% (13-45%)	74% (55-87%)	4.7 (1.8-9.4)	8.3% (5.1-12%)	11% (6.6-16%)	1.7 (1-2.6)	57 (35-83)
Sao Tome and Principe	2005	Female	60 to 64	27% (14-47%)	73% (53-86%)	4.8 (1.7-10)	8% (5.1-12%)	10% (6.2-15%)	2.3 (1.4-3.4)	63 (39-94)
Sao Tome and Principe	2005	Female	65 to 69	28% (15-46%)	72% (54-85%)	4.4 (1.7-9.2)	8.6% (5.2-13%)	10% (6.2-15%)	4.2 (2.5-6.3)	97 (57-140)
Sao Tome and Principe	2005	Female	70 to 74	26% (14-43%)	74% (57-86%)	3.5 (1.2-8)	8.1% (4.4-13%)	9.7% (5.3-15%)	5 (2.7-7.8)	92 (51-140)
Sao Tome and Principe	2005	Female	75 to 79	22% (11-40%)	78% (60-89%)	2.7 (0.79-6.7)	6.5% (3.2-11%)	7.6% (3.7-13%)	4.6 (2.2-7.6)	64 (32-110)
Sao Tome and Principe	2005	Female	80 to 84	20% (9.1-37%)	80% (63-91%)	2.4 (0.69-5.3)	6% (2.8-10%)	7% (3.3-12%)	3.8 (1.8-6.4)	39 (18-67)
Sao Tome and Principe	2005	Female	85 to 89	21% (9.6-37%)	79% (63-90%)	2.4 (0.71-5.7)	5.6% (2.5-9.7%)	6.5% (2.8-11%)	2.1 (0.93-3.7)	16 (6.9-28)
Sao Tome and Principe	2005	Female	90 to 94	24% (11-41%)	76% (59-89%)	2.6 (0.69-6.5)	5.3% (2.2-9.6%)	6.2% (2.5-11%)	0.82 (0.33-1.5)	4.2 (1.7-7.6)
Sao Tome and Principe	2005	Female	95 plus	29% (12-51%)	71% (49-88%)	3 (0.64-8)	5.4% (1.8-11%)	6.3% (2.1-12%)	0.28 (0.092-0.55)	0.96 (0.32-1.9)
Sao Tome and Principe	2005	Male	All Ages	11% (7.9-15%)	89% (85-92%)	1.9 (1.5-2.4)	2.9% (1.6-3.9%)	5.4% (3.3-7%)	31 (19-41)	1000 (570-1400)
Sao Tome and Principe	2005	Male	15 to 19	8.6% (4-16%)	91% (84-96%)	1.2 (0.43-2.5)	2.7% (1.5-4.4%)	3.5% (1.7-6.4%)	0.45 (0.21-0.8)	43 (24-70)
Sao Tome and Principe	2005	Male	20 to 24	13% (5.9-24%)	87% (76-94%)	2.4 (0.96-4.6)	5.4% (2.9-8.2%)	6.4% (2.9-11%)	0.93 (0.4-1.6)	80 (42-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	2005	Male	25 to 29	17% (7.8-30%)	83% (70-92%)	3.6 (1.5-6.6)	6.3% (2.2-10%)	7.6% (1.1-13%)	1.3 (0.19-2.3)	100 (37-170)
Sao Tome and Principe	2005	Male	30 to 34	20% (8.7-35%)	80% (65-91%)	4.5 (2-8)	6.7% (2.1-11%)	8.5% (1.2-15%)	1.1 (0.15-2.1)	81 (25-130)
Sao Tome and Principe	2005	Male	35 to 39	22% (11-38%)	78% (62-89%)	5.1 (2.1-9.2)	7.7% (2.1-13%)	10% (1.4-17%)	1.5 (0.2-2.6)	88 (23-150)
Sao Tome and Principe	2005	Male	40 to 44	23% (12-40%)	77% (60-88%)	5.6 (2.3-11)	8.8% (2.1-14%)	12% (1.7-20%)	1.7 (0.24-3)	88 (20-140)
Sao Tome and Principe	2005	Male	45 to 49	24% (11-40%)	76% (60-89%)	6 (2.3-13)	11% (2.2-16%)	15% (1.9-22%)	2.2 (0.27-3.3)	94 (18-140)
Sao Tome and Principe	2005	Male	50 to 54	24% (12-44%)	76% (56-88%)	6.1 (2.2-13)	11% (2.4-15%)	14% (2-20%)	2 (0.31-2.9)	74 (16-110)
Sao Tome and Principe	2005	Male	55 to 59	24% (12-42%)	76% (58-88%)	5.9 (2.1-12)	11% (6.2-15%)	14% (7.6-19%)	1.8 (0.98-2.5)	60 (33-83)
Sao Tome and Principe	2005	Male	60 to 64	24% (11-41%)	76% (59-89%)	5.3 (1.7-12)	9.5% (6.2-13%)	12% (7.5-16%)	2.2 (1.4-3)	60 (38-84)
Sao Tome and Principe	2005	Male	65 to 69	24% (12-40%)	76% (60-88%)	4.4 (1.6-9.6)	8.5% (5.5-12%)	10% (6.6-15%)	4 (2.6-5.8)	93 (60-130)
Sao Tome and Principe	2005	Male	70 to 74	22% (11-37%)	78% (63-89%)	3.3 (1.1-7.3)	7.4% (4.4-11%)	8.9% (5.3-13%)	4.3 (2.5-6.5)	79 (47-120)
Sao Tome and Principe	2005	Male	75 to 79	20% (9.9-36%)	80% (64-90%)	2.5 (0.7-6.1)	5.9% (3-9.3%)	7% (3.6-11%)	3.3 (1.7-5.2)	47 (24-74)
Sao Tome and Principe	2005	Male	80 to 84	18% (8.5-33%)	82% (67-92%)	2 (0.61-4.6)	5.1% (2.6-8.1%)	6% (3-9.6%)	2.3 (1.2-3.7)	24 (12-39)
Sao Tome and Principe	2005	Male	85 to 89	17% (8-31%)	83% (69-92%)	1.7 (0.5-4.1)	4.6% (2.1-7.8%)	5.4% (2.4-9.3%)	1.3 (0.58-2.2)	9.7 (4.4-16)
Sao Tome and Principe	2005	Male	90 to 94	19% (8.5-33%)	81% (67-92%)	1.7 (0.41-4.1)	3.8% (1.4-6.8%)	4.4% (1.6-8.1%)	0.43 (0.16-0.79)	2.2 (0.85-4)
Sao Tome and Principe	2005	Male	95 plus	23% (8.6-43%)	77% (57-91%)	1.7 (0.35-4.7)	3.3% (0.87-6.8%)	3.7% (0.88-7.9%)	0.1 (0.025-0.22)	0.37 (0.094-0.75)
Sao Tome and Principe	2010	Female	All Ages	13% (9.8-18%)	87% (82-90%)	1.5 (1.1-2)	3% (2.1-3.8%)	5.8% (3.9-7.9%)	29 (19-40)	800 (580-1000)
Sao Tome and Principe	2010	Female	15 to 19	13% (6-22%)	87% (78-94%)	1.3 (0.47-2.7)	1.9% (1.1-2.9%)	2.6% (0.76-4.8%)	0.17 (0.051-0.33)	23 (13-38)
Sao Tome and Principe	2010	Female	20 to 24	16% (7.5-29%)	84% (71-93%)	2.1 (0.83-4)	3.4% (2.2-4.8%)	3.9% (1.8-6.6%)	0.34 (0.15-0.57)	45 (29-66)
Sao Tome and Principe	2010	Female	25 to 29	18% (8-33%)	82% (67-92%)	2.6 (1.1-4.7)	3.6% (1.9-5.2%)	4.3% (0.57-7.2%)	0.35 (0.046-0.6)	40 (21-59)
Sao Tome and Principe	2010	Female	30 to 34	21% (9.2-37%)	79% (63-91%)	3 (1.3-5.2)	4.1% (1.7-5.9%)	5.8% (0.77-9.1%)	0.65 (0.084-1.1)	54 (23-81)
Sao Tome and Principe	2010	Female	35 to 39	23% (11-40%)	77% (60-89%)	3.3 (1.4-5.9)	5% (2.1-7.3%)	7.7% (1.9-11%)	0.79 (0.18-1.2)	51 (21-75)
Sao Tome and Principe	2010	Female	40 to 44	24% (12-41%)	76% (59-88%)	3.5 (1.5-6.5)	5.9% (2.3-8.4%)	9.1% (2.5-13%)	1.1 (0.32-1.6)	58 (22-85)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	2010	Female	45 to 49	25% (12-42%)	75% (58-88%)	3.6 (1.4-7.3)	6.8% (3.6-10%)	10% (5.1-15%)	1.6 (0.82-2.5)	71 (38-110)
Sao Tome and Principe	2010	Female	50 to 54	25% (12-44%)	75% (56-88%)	3.8 (1.5-7.6)	7.7% (4.6-11%)	11% (6.3-15%)	2 (1.2-3)	78 (47-110)
Sao Tome and Principe	2010	Female	55 to 59	26% (13-46%)	74% (54-87%)	4.2 (1.5-8.6)	7.9% (5-12%)	10% (6.5-15%)	2 (1.2-3)	66 (41-99)
Sao Tome and Principe	2010	Female	60 to 64	27% (14-46%)	73% (54-86%)	4.3 (1.5-9.3)	7.6% (4.8-11%)	9.7% (6.1-14%)	2 (1.2-3)	54 (33-82)
Sao Tome and Principe	2010	Female	65 to 69	28% (14-46%)	72% (54-86%)	3.9 (1.4-8)	8.1% (4.8-12%)	10% (5.7-15%)	2.8 (1.7-4.2)	65 (38-97)
Sao Tome and Principe	2010	Female	70 to 74	26% (13-43%)	74% (57-87%)	3.1 (1-7.2)	7.5% (4.1-12%)	9.1% (4.9-14%)	4.3 (2.2-6.7)	79 (42-120)
Sao Tome and Principe	2010	Female	75 to 79	22% (10-39%)	78% (61-90%)	2.5 (0.69-6.3)	6.1% (2.9-10%)	7.2% (3.3-12%)	4 (1.9-6.7)	57 (27-95)
Sao Tome and Principe	2010	Female	80 to 84	20% (9-37%)	80% (63-91%)	2.2 (0.64-4.9)	5.6% (2.5-9.4%)	6.6% (2.9-11%)	3.6 (1.6-6.2)	38 (17-64)
Sao Tome and Principe	2010	Female	85 to 89	21% (9.9-37%)	79% (63-90%)	2.1 (0.66-5)	5.2% (2.2-9.3%)	6.1% (2.6-11%)	2.3 (0.99-4.2)	17 (7.4-31)
Sao Tome and Principe	2010	Female	90 to 94	24% (11-42%)	76% (58-89%)	2.3 (0.61-5.7)	5% (1.8-9.4%)	5.8% (2.1-11%)	0.9 (0.33-1.7)	4.6 (1.7-8.7)
Sao Tome and Principe	2010	Female	95 plus	28% (11-51%)	72% (49-89%)	2.6 (0.57-7.2)	5.1% (1.4-10%)	5.9% (1.6-12%)	0.25 (0.065-0.51)	0.85 (0.23-1.7)
Sao Tome and Principe	2010	Male	All Ages	11% (8-15%)	89% (85-92%)	1.8 (1.4-2.4)	3.4% (1.9-4.3%)	5.8% (3.4-7.3%)	30 (17-39)	1100 (600-1400)
Sao Tome and Principe	2010	Male	15 to 19	8.5% (4.1-16%)	91% (84-96%)	1.1 (0.38-2.2)	2.5% (1.4-4.1%)	3.4% (1.4-6.1%)	0.36 (0.15-0.65)	37 (20-59)
Sao Tome and Principe	2010	Male	20 to 24	13% (6-23%)	87% (77-94%)	2.2 (0.87-4.3)	5.4% (3.3-8%)	6.5% (3.3-11%)	1 (0.49-1.7)	90 (53-140)
Sao Tome and Principe	2010	Male	25 to 29	17% (7.9-30%)	83% (70-92%)	3.3 (1.4-5.9)	6.6% (2.5-10%)	8.2% (1.1-14%)	1.1 (0.16-1.9)	91 (33-140)
Sao Tome and Principe	2010	Male	30 to 34	20% (8.7-36%)	80% (64-91%)	4.2 (1.8-7.4)	7.1% (2.3-11%)	9.2% (1.2-16%)	1.6 (0.2-2.8)	110 (36-180)
Sao Tome and Principe	2010	Male	35 to 39	22% (10-39%)	78% (61-90%)	4.7 (2-8.9)	8.3% (2.2-13%)	11% (1.5-19%)	1.6 (0.2-2.7)	98 (26-150)
Sao Tome and Principe	2010	Male	40 to 44	23% (12-39%)	77% (61-88%)	5.1 (2-9.8)	9.3% (2.2-14%)	13% (1.7-20%)	2 (0.25-3.3)	100 (23-160)
Sao Tome and Principe	2010	Male	45 to 49	23% (11-39%)	77% (61-89%)	5.3 (2-11)	11% (2.4-16%)	16% (2-22%)	2.6 (0.32-3.9)	110 (23-170)
Sao Tome and Principe	2010	Male	50 to 54	24% (11-43%)	76% (57-89%)	5.4 (2-12)	10% (3.3-15%)	14% (3.6-20%)	2.3 (0.57-3.4)	88 (27-130)
Sao Tome and Principe	2010	Male	55 to 59	24% (12-42%)	76% (58-88%)	5.3 (1.8-11)	11% (6.6-15%)	14% (8.5-19%)	2.3 (1.3-3.2)	75 (45-100)
Sao Tome and Principe	2010	Male	60 to 64	24% (12-42%)	76% (58-88%)	4.8 (1.6-11)	9.3% (6-13%)	12% (7.5-16%)	1.8 (1.1-2.6)	51 (33-71)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	2010	Male	65 to 69	24% (12-40%)	76% (60-88%)	4 (1.4-8.5)	8.1% (5.3-11%)	10% (6.4-14%)	2.1 (1.3-3)	48 (31-69)
Sao Tome and Principe	2010	Male	70 to 74	22% (11-38%)	78% (62-89%)	3 (1-6.7)	7% (4.1-10%)	8.5% (5-13%)	3.7 (2.1-5.5)	68 (40-100)
Sao Tome and Principe	2010	Male	75 to 79	20% (9.9-35%)	80% (65-90%)	2.2 (0.6-5.4)	5.4% (2.7-8.6%)	6.5% (3.2-10%)	3.1 (1.5-5)	44 (22-71)
Sao Tome and Principe	2010	Male	80 to 84	18% (8.2-32%)	82% (68-92%)	1.8 (0.51-4.1)	4.7% (2.2-7.6%)	5.5% (2.7-8.8%)	2.2 (1.1-3.6)	24 (11-38)
Sao Tome and Principe	2010	Male	85 to 89	17% (7.9-31%)	83% (69-92%)	1.6 (0.45-3.7)	4.2% (1.9-7.2%)	4.9% (2.2-8.6%)	1.3 (0.57-2.2)	9.7 (4.3-17)
Sao Tome and Principe	2010	Male	90 to 94	19% (8.6-34%)	81% (66-91%)	1.5 (0.38-3.8)	3.4% (1.2-6.4%)	4% (1.3-7.6%)	0.45 (0.15-0.84)	2.3 (0.79-4.3)
Sao Tome and Principe	2010	Male	95 plus	22% (9.2-42%)	78% (58-91%)	1.6 (0.31-4.3)	3% (0.77-6.4%)	3.4% (0.72-7.6%)	0.098 (0.021-0.22)	0.35 (0.087-0.76)
Sao Tome and Principe	2016	Female	All Ages	14% (9.9-18%)	86% (82-90%)	1.5 (0.89-2.4)	3.1% (2.3-3.8%)	5.7% (3.9-7.6%)	27 (18-37)	800 (610-1000)
Sao Tome and Principe	2016	Female	15 to 19	12% (6-22%)	88% (78-94%)	1.3 (0.4-3)	1.8% (1-2.7%)	2.3% (0.87-4.3%)	0.15 (0.058-0.29)	24 (13-38)
Sao Tome and Principe	2016	Female	20 to 24	16% (7.1-28%)	84% (72-93%)	2 (0.67-4.2)	3.3% (2.2-4.6%)	3.7% (1.9-6%)	0.26 (0.13-0.44)	40 (26-59)
Sao Tome and Principe	2016	Female	25 to 29	18% (8-31%)	82% (69-92%)	2.5 (0.93-5)	3.6% (2.5-5%)	4.4% (2.5-6.9%)	0.38 (0.2-0.61)	46 (31-66)
Sao Tome and Principe	2016	Female	30 to 34	21% (8.8-37%)	79% (63-91%)	2.9 (1.2-5.9)	4.1% (2.7-5.5%)	5.9% (3.4-8.6%)	0.52 (0.29-0.81)	46 (30-66)
Sao Tome and Principe	2016	Female	35 to 39	23% (11-40%)	77% (60-89%)	3.2 (1.2-6.1)	4.9% (3.3-6.8%)	7.9% (5.1-11%)	1 (0.61-1.6)	68 (44-99)
Sao Tome and Principe	2016	Female	40 to 44	24% (12-42%)	76% (58-88%)	3.3 (1.4-6.8)	5.7% (3.9-7.9%)	9.2% (6.2-13%)	1.2 (0.82-1.9)	67 (45-98)
Sao Tome and Principe	2016	Female	45 to 49	25% (11-42%)	75% (58-89%)	3.5 (1.2-7.4)	6.5% (4.3-9.2%)	9.8% (6.4-14%)	1.5 (0.95-2.3)	69 (45-100)
Sao Tome and Principe	2016	Female	50 to 54	25% (12-45%)	75% (55-88%)	3.7 (1.2-8.4)	7.1% (4.7-10%)	10% (6.6-14%)	2.1 (1.3-3.2)	83 (53-120)
Sao Tome and Principe	2016	Female	55 to 59	26% (13-44%)	74% (56-87%)	4 (1.4-8.7)	7.3% (4.7-11%)	9.9% (6.3-14%)	2.5 (1.5-3.7)	83 (51-120)
Sao Tome and Principe	2016	Female	60 to 64	27% (13-46%)	73% (54-87%)	4.1 (1.4-9.5)	7.1% (4.4-11%)	9.1% (5.7-14%)	2.4 (1.4-3.6)	66 (41-99)
Sao Tome and Principe	2016	Female	65 to 69	28% (14-45%)	72% (55-86%)	3.8 (1.3-8.4)	7.4% (4.1-11%)	9.2% (5.1-14%)	2.4 (1.3-3.7)	55 (31-86)
Sao Tome and Principe	2016	Female	70 to 74	26% (13-42%)	74% (58-87%)	3 (0.89-7.6)	6.8% (3.5-11%)	8.3% (4.1-13%)	2.7 (1.4-4.2)	49 (26-78)
Sao Tome and Principe	2016	Female	75 to 79	22% (11-38%)	78% (62-89%)	2.3 (0.64-6)	5.4% (2.6-9.2%)	6.5% (3-11%)	3.2 (1.5-5.4)	46 (22-78)
Sao Tome and Principe	2016	Female	80 to 84	20% (9.2-37%)	80% (63-91%)	2 (0.61-5.3)	5% (2.2-8.5%)	5.9% (2.6-10%)	3.2 (1.4-5.5)	34 (15-58)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sao Tome and Principe	2016	Female	85 to 89	21% (9.8-36%)	79% (64-90%)	2 (0.56-5.2)	4.6% (1.9-8.4%)	5.4% (2.2-9.9%)	2.2 (0.88-4.1)	16 (6.7-30)
Sao Tome and Principe	2016	Female	90 to 94	24% (11-41%)	76% (59-89%)	2.2 (0.56-6)	4.4% (1.5-8.4%)	5.2% (1.7-10%)	0.99 (0.32-1.9)	5 (1.7-9.7)
Sao Tome and Principe	2016	Female	95 plus	29% (12-51%)	71% (49-88%)	2.6 (0.53-7.4)	4.6% (1.2-9.3%)	5.3% (1.3-11%)	0.28 (0.071-0.59)	0.97 (0.25-2)
Sao Tome and Principe	2016	Male	All Ages	11% (8.1-15%)	89% (85-92%)	1.8 (1.1-2.8)	4.1% (2.8-5%)	6.5% (4.7-8.1%)	33 (23-44)	1300 (870-1600)
Sao Tome and Principe	2016	Male	15 to 19	8.4% (4-16%)	92% (84-96%)	1 (0.31-2.4)	2.4% (1.4-3.9%)	3.1% (1.4-5.7%)	0.38 (0.17-0.69)	39 (23-66)
Sao Tome and Principe	2016	Male	20 to 24	13% (5.9-24%)	87% (76-94%)	2.1 (0.71-4.5)	5.3% (3.4-7.8%)	6.3% (3.5-10%)	0.95 (0.5-1.6)	86 (53-130)
Sao Tome and Principe	2016	Male	25 to 29	17% (7.9-30%)	83% (70-92%)	3.1 (1.1-6.5)	7% (4.3-10%)	9% (4.7-14%)	1.5 (0.72-2.4)	120 (70-180)
Sao Tome and Principe	2016	Male	30 to 34	20% (8.6-35%)	80% (65-91%)	3.9 (1.6-7.9)	7.8% (3.4-11%)	11% (2.5-16%)	1.6 (0.35-2.5)	110 (45-160)
Sao Tome and Principe	2016	Male	35 to 39	22% (9.9-37%)	78% (63-90%)	4.4 (1.7-9)	9% (4.5-13%)	13% (5.3-18%)	2.5 (0.98-3.8)	150 (69-210)
Sao Tome and Principe	2016	Male	40 to 44	23% (12-40%)	77% (60-88%)	4.8 (1.8-9.7)	10% (6-14%)	15% (8.2-20%)	2.7 (1.4-3.8)	140 (77-190)
Sao Tome and Principe	2016	Male	45 to 49	24% (11-39%)	76% (61-89%)	5.1 (1.7-12)	12% (7.5-16%)	17% (11-22%)	3.4 (1.9-4.8)	150 (90-210)
Sao Tome and Principe	2016	Male	50 to 54	24% (11-43%)	76% (57-89%)	5.2 (1.7-12)	11% (6.9-14%)	14% (9.6-19%)	3.3 (2-4.7)	130 (80-180)
Sao Tome and Principe	2016	Male	55 to 59	24% (12-42%)	76% (58-88%)	5.1 (1.6-12)	11% (7.6-14%)	14% (9.9-18%)	3.3 (2.1-4.6)	110 (71-150)
Sao Tome and Principe	2016	Male	60 to 64	24% (11-40%)	76% (60-89%)	4.6 (1.4-11)	9.1% (6.4-12%)	11% (8-16%)	2.7 (1.8-4)	77 (50-110)
Sao Tome and Principe	2016	Male	65 to 69	24% (12-40%)	76% (60-88%)	3.8 (1.2-9)	7.7% (5-11%)	9.5% (6.1-13%)	2 (1.2-3.1)	47 (29-71)
Sao Tome and Principe	2016	Male	70 to 74	22% (12-39%)	78% (61-88%)	2.9 (0.87-7)	6.4% (3.7-9.6%)	7.8% (4.5-11%)	1.9 (1-3)	35 (19-56)
Sao Tome and Principe	2016	Male	75 to 79	20% (9.5-35%)	80% (65-90%)	2.2 (0.57-5.5)	5.1% (2.5-8.1%)	6.1% (3-9.8%)	2.7 (1.3-4.5)	39 (18-64)
Sao Tome and Principe	2016	Male	80 to 84	18% (8.2-33%)	82% (67-92%)	1.8 (0.47-4.5)	4.3% (2.1-7.1%)	5.1% (2.4-8.3%)	2.4 (1.1-4)	25 (12-43)
Sao Tome and Principe	2016	Male	85 to 89	17% (7.9-31%)	83% (69-92%)	1.5 (0.41-3.9)	3.8% (1.6-6.7%)	4.5% (1.9-7.9%)	1.4 (0.55-2.4)	10 (4.2-18)
Sao Tome and Principe	2016	Male	90 to 94	19% (8.5-34%)	81% (66-92%)	1.5 (0.34-4.1)	3.1% (1-5.9%)	3.6% (1.1-7.1%)	0.46 (0.14-0.91)	2.4 (0.77-4.8)
Sao Tome and Principe	2016	Male	95 plus	23% (8.9-42%)	77% (58-91%)	1.5 (0.27-4.8)	2.7% (0.63-5.8%)	3.1% (0.57-6.8%)	0.11 (0.019-0.24)	0.38 (0.083-0.83)
Senegal	1990	Female	All Ages	2.9% (2.4-3.5%)	97% (97-98%)	0.2 (0.15-0.25)	0.74% (0.57-0.97%)	1.5% (1-2%)	560 (380-770)	20000 (15000-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	1990	Female	15 to 19	2.7% (1.6-4.1%)	97% (96-98%)	0.18 (0.09-0.31)	0.84% (0.51-1.3%)	0.7% (0.3-1.4%)	5.1 (2.1-9.9)	730 (430-1100)
Senegal	1990	Female	20 to 24	3.5% (2-5.3%)	96% (95-98%)	0.29 (0.14-0.5)	1.7% (1.1-2.5%)	1.3% (0.62-2.5%)	10 (4.9-20)	1400 (870-2100)
Senegal	1990	Female	25 to 29	4% (2.3-6.2%)	96% (94-98%)	0.35 (0.18-0.6)	2% (1.4-3%)	1.9% (0.98-3.2%)	14 (7.1-25)	1500 (970-2200)
Senegal	1990	Female	30 to 34	4.4% (2.5-7.2%)	96% (93-97%)	0.4 (0.2-0.69)	2.2% (1.5-3.3%)	2.4% (1.3-4%)	17 (9.6-30)	1400 (950-2200)
Senegal	1990	Female	35 to 39	4.9% (2.7-7.4%)	95% (93-97%)	0.43 (0.22-0.75)	2.4% (1.6-3.5%)	2.9% (1.7-4.6%)	22 (13-35)	1400 (930-2100)
Senegal	1990	Female	40 to 44	5.1% (3-7.7%)	95% (92-97%)	0.44 (0.21-0.76)	2.8% (1.8-4.1%)	3.6% (2.2-5.6%)	27 (17-41)	1500 (970-2100)
Senegal	1990	Female	45 to 49	5% (2.8-7.6%)	95% (92-97%)	0.43 (0.21-0.77)	3% (1.9-4.5%)	4% (2.5-6.4%)	36 (22-56)	1600 (1000-2500)
Senegal	1990	Female	50 to 54	5% (3-7.5%)	95% (92-97%)	0.46 (0.23-0.77)	3.4% (2.2-5.2%)	4.5% (2.7-6.9%)	50 (30-77)	1900 (1200-2900)
Senegal	1990	Female	55 to 59	5.2% (3.1-8%)	95% (92-97%)	0.54 (0.28-0.96)	3.7% (2.2-5.6%)	4.7% (2.8-7.4%)	65 (38-100)	2100 (1300-3300)
Senegal	1990	Female	60 to 64	5.7% (3.3-8.8%)	94% (91-97%)	0.59 (0.28-1.1)	3.5% (2.1-5.5%)	4.3% (2.6-6.7%)	73 (43-110)	2000 (1200-3100)
Senegal	1990	Female	65 to 69	6.1% (3.7-8.9%)	94% (91-96%)	0.56 (0.28-0.98)	3.4% (2-5.4%)	4.2% (2.4-6.6%)	83 (48-130)	1900 (1100-3000)
Senegal	1990	Female	70 to 74	5.5% (3.5-8.2%)	94% (92-96%)	0.45 (0.22-0.78)	3.2% (1.7-5.4%)	3.8% (2-6.4%)	80 (43-130)	1500 (790-2400)
Senegal	1990	Female	75 to 79	4.5% (2.6-7.1%)	96% (93-97%)	0.33 (0.15-0.62)	2.2% (1.1-3.8%)	2.6% (1.2-4.6%)	44 (21-79)	620 (310-1100)
Senegal	1990	Female	80 to 84	4% (2.2-5.9%)	96% (94-98%)	0.28 (0.13-0.51)	1.6% (0.78-2.9%)	1.9% (0.9-3.4%)	18 (8.5-33)	190 (91-340)
Senegal	1990	Female	85 to 89	4.2% (2.4-6.6%)	96% (93-98%)	0.28 (0.14-0.5)	1.4% (0.61-2.5%)	1.6% (0.7-3%)	8.3 (3.7-16)	62 (28-110)
Senegal	1990	Female	90 to 94	4.7% (2.9-7.5%)	95% (92-97%)	0.3 (0.14-0.56)	1.2% (0.49-2.4%)	1.4% (0.53-2.9%)	2.5 (0.95-5.1)	13 (5.2-26)
Senegal	1990	Female	95 plus	5.5% (3.1-9%)	94% (91-97%)	0.33 (0.13-0.69)	1.1% (0.39-2.6%)	1.3% (0.38-3%)	0.56 (0.16-1.3)	2 (0.69-4.6)
Senegal	1990	Male	All Ages	2.9% (2.4-3.6%)	97% (96-98%)	0.085 (0.064-0.11)	0.61% (0.46-0.79%)	0.96% (0.66-1.3%)	410 (280-560)	18000 (14000-24000)
Senegal	1990	Male	15 to 19	2.2% (1.3-3.5%)	98% (97-99%)	0.048 (0.023-0.086)	0.61% (0.39-0.86%)	0.31% (0.16-0.56%)	2.4 (1.2-4.2)	530 (330-770)
Senegal	1990	Male	20 to 24	3.6% (2.1-5.4%)	96% (95-98%)	0.11 (0.054-0.19)	1.4% (1-2%)	0.89% (0.46-1.5%)	8.4 (4.3-14)	1200 (840-1800)
Senegal	1990	Male	25 to 29	4.7% (2.7-7.3%)	95% (93-97%)	0.17 (0.085-0.29)	2.2% (1.5-3.1%)	1.8% (0.98-3%)	16 (8.7-27)	1700 (1100-2400)
Senegal	1990	Male	30 to 34	5.4% (3.1-8.7%)	95% (91-97%)	0.21 (0.11-0.35)	2.7% (1.8-3.9%)	2.7% (1.5-4.3%)	24 (13-37)	1900 (1200-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	1990	Male	35 to 39	5.9% (3.3-8.8%)	94% (91-97%)	0.22 (0.11-0.39)	3.2% (2.1-4.5%)	3.5% (2.1-5.3%)	32 (19-49)	2000 (1300-2900)
Senegal	1990	Male	40 to 44	6% (3.5-9%)	94% (91-96%)	0.23 (0.11-0.4)	3.3% (2.2-4.7%)	3.9% (2.4-5.7%)	36 (22-54)	1900 (1300-2700)
Senegal	1990	Male	45 to 49	6.1% (3.3-9.3%)	94% (91-97%)	0.24 (0.12-0.43)	3.5% (2.3-5%)	4.2% (2.5-6.2%)	48 (29-72)	2200 (1400-3100)
Senegal	1990	Male	50 to 54	6.3% (3.8-9.4%)	94% (91-96%)	0.25 (0.12-0.42)	3.1% (1.9-4.5%)	3.7% (2.2-5.5%)	51 (30-77)	2000 (1200-2900)
Senegal	1990	Male	55 to 59	6.4% (3.8-9.8%)	94% (90-96%)	0.24 (0.13-0.42)	2.8% (1.8-4.2%)	3.4% (2.1-5.1%)	53 (33-82)	1800 (1200-2700)
Senegal	1990	Male	60 to 64	6.2% (3.7-9.8%)	94% (90-96%)	0.22 (0.11-0.39)	2.3% (1.4-3.4%)	2.7% (1.6-4.1%)	50 (30-76)	1400 (870-2100)
Senegal	1990	Male	65 to 69	6% (3.7-8.8%)	94% (91-96%)	0.18 (0.088-0.31)	1.7% (1.1-2.4%)	1.9% (1.2-2.9%)	38 (23-56)	880 (560-1300)
Senegal	1990	Male	70 to 74	5.7% (3.7-8.5%)	94% (92-96%)	0.14 (0.069-0.24)	1.2% (0.72-1.9%)	1.4% (0.8-2.2%)	26 (15-42)	480 (290-760)
Senegal	1990	Male	75 to 79	5.1% (3-8%)	95% (92-97%)	0.1 (0.049-0.19)	0.86% (0.5-1.3%)	0.97% (0.54-1.5%)	14 (7.7-22)	210 (120-310)
Senegal	1990	Male	80 to 84	4.4% (2.5-6.6%)	96% (93-98%)	0.078 (0.034-0.15)	0.7% (0.41-1%)	0.78% (0.44-1.2%)	5.7 (3.2-8.7)	61 (36-91)
Senegal	1990	Male	85 to 89	4% (2.3-6.3%)	96% (94-98%)	0.061 (0.029-0.11)	0.53% (0.32-0.76%)	0.58% (0.32-0.85%)	2.2 (1.2-3.2)	17 (10-24)
Senegal	1990	Male	90 to 94	4.1% (2.4-6.4%)	96% (94-98%)	0.052 (0.023-0.098)	0.32% (0.15-0.48%)	0.32% (0.11-0.52%)	0.41 (0.14-0.66)	2.4 (1.1-3.7)
Senegal	1990	Male	95 plus	4.7% (2.5-7.6%)	95% (92-97%)	0.048 (0.018-0.1)	0.31% (0.12-0.5%)	0.27% (0.044-0.49%)	0.074 (0.012-0.14)	0.34 (0.13-0.56)
Senegal	1995	Female	All Ages	3% (2.5-3.6%)	97% (96-98%)	0.17 (0.13-0.21)	0.71% (0.53-0.91%)	1.3% (0.89-1.9%)	560 (370-790)	20000 (15000-27000)
Senegal	1995	Female	15 to 19	2.8% (1.6-4.2%)	97% (96-98%)	0.15 (0.073-0.25)	0.74% (0.46-1.1%)	0.56% (0.23-1.1%)	4.8 (2-9.3)	780 (460-1200)
Senegal	1995	Female	20 to 24	3.6% (2.1-5.5%)	96% (94-98%)	0.24 (0.12-0.41)	1.6% (0.99-2.3%)	1.1% (0.49-2%)	11 (4.8-20)	1600 (1000-2400)
Senegal	1995	Female	25 to 29	4.1% (2.4-6.4%)	96% (94-98%)	0.29 (0.15-0.49)	1.9% (1.3-2.6%)	1.6% (0.83-2.7%)	14 (7.3-25)	1600 (1000-2400)
Senegal	1995	Female	30 to 34	4.6% (2.6-7.4%)	95% (93-97%)	0.34 (0.17-0.55)	2% (1.3-2.9%)	2% (1.2-3.3%)	18 (9.9-29)	1500 (1000-2200)
Senegal	1995	Female	35 to 39	5.1% (2.8-7.5%)	95% (92-97%)	0.36 (0.18-0.63)	2.1% (1.4-3.2%)	2.5% (1.5-4%)	22 (13-36)	1500 (980-2200)
Senegal	1995	Female	40 to 44	5.2% (3.1-7.9%)	95% (92-97%)	0.37 (0.18-0.63)	2.4% (1.6-3.5%)	3.1% (1.9-4.7%)	28 (17-43)	1600 (1000-2200)
Senegal	1995	Female	45 to 49	5.2% (2.8-7.8%)	95% (92-97%)	0.36 (0.18-0.63)	2.6% (1.6-4%)	3.4% (2-5.5%)	33 (19-53)	1500 (940-2300)
Senegal	1995	Female	50 to 54	5.2% (3.2-7.9%)	95% (92-97%)	0.39 (0.2-0.65)	3% (1.8-4.5%)	3.9% (2.3-5.9%)	48 (28-74)	1900 (1100-2800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	1995	Female	55 to 59	5.4% (3.3-8.2%)	95% (92-97%)	0.45 (0.23-0.8)	3.2% (2-5%)	4.1% (2.4-6.6%)	65 (38-100)	2100 (1300-3400)
Senegal	1995	Female	60 to 64	5.8% (3.5-9%)	94% (91-97%)	0.49 (0.24-0.86)	3% (1.8-4.8%)	3.7% (2.2-5.9%)	74 (42-120)	2100 (1200-3300)
Senegal	1995	Female	65 to 69	6.2% (3.9-9.1%)	94% (91-96%)	0.47 (0.24-0.81)	2.9% (1.7-4.7%)	3.5% (1.9-5.7%)	83 (46-140)	1900 (1100-3100)
Senegal	1995	Female	70 to 74	5.7% (3.6-8.4%)	94% (92-96%)	0.37 (0.18-0.66)	2.6% (1.3-4.6%)	3.1% (1.5-5.5%)	79 (39-140)	1400 (720-2500)
Senegal	1995	Female	75 to 79	4.6% (2.6-7%)	95% (93-97%)	0.27 (0.12-0.51)	1.8% (0.88-3.3%)	2.1% (1-3.7%)	45 (22-84)	640 (320-1200)
Senegal	1995	Female	80 to 84	4.1% (2.2-6%)	96% (94-98%)	0.23 (0.1-0.41)	1.3% (0.6-2.3%)	1.5% (0.66-2.7%)	19 (8.4-34)	200 (91-350)
Senegal	1995	Female	85 to 89	4.3% (2.4-6.6%)	96% (93-98%)	0.23 (0.11-0.42)	1% (0.45-2%)	1.2% (0.5-2.3%)	8.6 (3.5-16)	65 (28-120)
Senegal	1995	Female	90 to 94	4.8% (2.9-7.6%)	95% (92-97%)	0.25 (0.12-0.46)	0.89% (0.31-2%)	1% (0.32-2.3%)	2.6 (0.82-5.7)	14 (4.7-29)
Senegal	1995	Female	95 plus	5.6% (3.2-9.1%)	94% (91-97%)	0.27 (0.11-0.57)	0.85% (0.25-2%)	0.93% (0.21-2.3%)	0.57 (0.13-1.5)	2.1 (0.61-5)
Senegal	1995	Male	All Ages	3.1% (2.5-3.7%)	97% (96-97%)	0.074 (0.056-0.097)	0.57% (0.43-0.75%)	0.88% (0.6-1.2%)	400 (270-570)	18000 (14000-24000)
Senegal	1995	Male	15 to 19	2.3% (1.4-3.6%)	98% (96-99%)	0.042 (0.021-0.075)	0.56% (0.37-0.8%)	0.24% (0.13-0.44%)	2.2 (1.2-4)	580 (370-880)
Senegal	1995	Male	20 to 24	3.8% (2.2-5.8%)	96% (94-98%)	0.095 (0.047-0.16)	1.3% (0.88-1.8%)	0.71% (0.37-1.2%)	8.4 (4.4-14)	1400 (910-2000)
Senegal	1995	Male	25 to 29	4.9% (2.9-7.8%)	95% (92-97%)	0.15 (0.073-0.26)	1.9% (1.3-2.6%)	1.5% (0.77-2.4%)	16 (8.4-27)	1800 (1200-2500)
Senegal	1995	Male	30 to 34	5.6% (3.3-9.4%)	94% (91-97%)	0.18 (0.092-0.3)	2.3% (1.6-3.3%)	2.2% (1.2-3.4%)	24 (13-37)	1900 (1300-2700)
Senegal	1995	Male	35 to 39	6.1% (3.5-9.1%)	94% (91-97%)	0.19 (0.096-0.34)	2.7% (1.8-3.9%)	3% (1.8-4.5%)	32 (20-50)	2000 (1400-2900)
Senegal	1995	Male	40 to 44	6.3% (3.8-9.3%)	94% (91-96%)	0.2 (0.095-0.35)	2.8% (1.9-4%)	3.3% (2-4.8%)	37 (23-55)	1900 (1300-2800)
Senegal	1995	Male	45 to 49	6.4% (3.5-9.5%)	94% (91-97%)	0.21 (0.11-0.37)	3% (2-4.3%)	3.6% (2.2-5.3%)	42 (26-64)	1900 (1200-2800)
Senegal	1995	Male	50 to 54	6.5% (4-9.9%)	93% (90-96%)	0.21 (0.11-0.37)	2.7% (1.7-3.9%)	3.2% (1.9-4.8%)	47 (28-70)	1800 (1200-2700)
Senegal	1995	Male	55 to 59	6.6% (4-9.9%)	93% (90-96%)	0.21 (0.11-0.37)	2.5% (1.6-3.8%)	3% (1.8-4.6%)	52 (32-80)	1800 (1100-2600)
Senegal	1995	Male	60 to 64	6.5% (3.9-10%)	93% (90-96%)	0.19 (0.093-0.35)	2.1% (1.3-3.2%)	2.4% (1.4-3.8%)	51 (29-80)	1400 (880-2200)
Senegal	1995	Male	65 to 69	6.3% (3.9-9.4%)	94% (91-96%)	0.16 (0.079-0.28)	1.5% (0.95-2.2%)	1.7% (1-2.6%)	40 (24-60)	930 (580-1400)
Senegal	1995	Male	70 to 74	6% (3.8-8.7%)	94% (91-96%)	0.12 (0.061-0.22)	1.1% (0.62-1.7%)	1.2% (0.69-2%)	28 (16-44)	520 (300-810)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	1995	Male	75 to 79	5.4% (3.2-8.3%)	95% (92-97%)	0.091 (0.043-0.17)	0.76% (0.44-1.2%)	0.87% (0.47-1.4%)	16 (8.5-25)	230 (130-360)
Senegal	1995	Male	80 to 84	4.6% (2.7-6.8%)	95% (93-97%)	0.069 (0.031-0.13)	0.63% (0.37-0.94%)	0.71% (0.39-1.1%)	6.6 (3.7-10)	72 (43-110)
Senegal	1995	Male	85 to 89	4.2% (2.4-6.5%)	96% (93-98%)	0.054 (0.026-0.098)	0.49% (0.27-0.72%)	0.53% (0.27-0.81%)	2.6 (1.3-4)	21 (11-31)
Senegal	1995	Male	90 to 94	4.3% (2.5-6.7%)	96% (93-97%)	0.046 (0.02-0.086)	0.3% (0.13-0.46%)	0.3% (0.088-0.5%)	0.51 (0.15-0.85)	3.1 (1.3-4.8)
Senegal	1995	Male	95 plus	4.9% (2.7-8.1%)	95% (92-97%)	0.042 (0.016-0.088)	0.3% (0.12-0.48%)	0.26% (0.037-0.49%)	0.098 (0.014-0.18)	0.46 (0.18-0.73)
Senegal	2000	Female	All Ages	2.9% (2.4-3.5%)	97% (97-98%)	0.15 (0.11-0.18)	0.74% (0.56-0.94%)	1.3% (0.86-1.8%)	570 (370-810)	21000 (16000-28000)
Senegal	2000	Female	15 to 19	2.7% (1.6-4%)	97% (96-98%)	0.13 (0.064-0.22)	0.71% (0.46-1.1%)	0.48% (0.21-0.94%)	4.4 (1.8-8.5)	800 (490-1200)
Senegal	2000	Female	20 to 24	3.4% (2-5.2%)	97% (95-98%)	0.21 (0.11-0.35)	1.5% (0.96-2.1%)	0.91% (0.43-1.6%)	9.9 (4.7-18)	1700 (1100-2500)
Senegal	2000	Female	25 to 29	3.8% (2.2-5.9%)	96% (94-98%)	0.25 (0.13-0.41)	1.7% (1.2-2.4%)	1.4% (0.73-2.4%)	14 (7.7-25)	1800 (1200-2600)
Senegal	2000	Female	30 to 34	4.1% (2.4-6.6%)	96% (93-98%)	0.28 (0.14-0.46)	1.8% (1.2-2.5%)	1.7% (0.98-2.7%)	17 (9.9-28)	1600 (1100-2300)
Senegal	2000	Female	35 to 39	4.4% (2.5-6.7%)	96% (93-97%)	0.29 (0.15-0.5)	1.9% (1.3-2.7%)	2.1% (1.3-3.3%)	22 (13-35)	1500 (1100-2200)
Senegal	2000	Female	40 to 44	4.9% (3-7.2%)	95% (93-97%)	0.32 (0.15-0.54)	2.2% (1.5-3.1%)	2.8% (1.7-4.2%)	30 (18-46)	1700 (1100-2400)
Senegal	2000	Female	45 to 49	5.1% (3-7.5%)	95% (92-97%)	0.33 (0.17-0.57)	2.4% (1.5-3.6%)	3.1% (1.8-4.9%)	36 (21-57)	1700 (1100-2600)
Senegal	2000	Female	50 to 54	5% (3.1-7.5%)	95% (92-97%)	0.34 (0.17-0.58)	2.7% (1.6-4.1%)	3.5% (2-5.3%)	47 (26-70)	1800 (1100-2700)
Senegal	2000	Female	55 to 59	5.1% (3.1-7.6%)	95% (92-97%)	0.39 (0.2-0.66)	2.9% (1.7-4.6%)	3.7% (2.1-5.9%)	64 (36-100)	2100 (1200-3300)
Senegal	2000	Female	60 to 64	5.7% (3.4-8.5%)	94% (91-97%)	0.43 (0.21-0.78)	2.8% (1.6-4.4%)	3.4% (2-5.4%)	76 (42-120)	2100 (1200-3300)
Senegal	2000	Female	65 to 69	6.2% (3.8-8.9%)	94% (91-96%)	0.42 (0.21-0.72)	2.7% (1.5-4.3%)	3.2% (1.8-5.1%)	87 (48-140)	2000 (1100-3200)
Senegal	2000	Female	70 to 74	5.6% (3.7-8.2%)	94% (92-96%)	0.34 (0.16-0.59)	2.4% (1.2-4.1%)	2.8% (1.4-4.9%)	82 (41-150)	1500 (770-2700)
Senegal	2000	Female	75 to 79	4.5% (2.6-7.1%)	95% (93-97%)	0.25 (0.11-0.47)	1.6% (0.81-2.9%)	1.9% (0.91-3.4%)	49 (23-91)	700 (340-1300)
Senegal	2000	Female	80 to 84	4.1% (2.2-6%)	96% (94-98%)	0.21 (0.094-0.36)	1.1% (0.53-2%)	1.3% (0.59-2.4%)	21 (9.2-39)	220 (99-400)
Senegal	2000	Female	85 to 89	4.3% (2.5-6.8%)	96% (93-97%)	0.21 (0.1-0.38)	0.92% (0.38-1.8%)	1.1% (0.41-2%)	9.7 (3.8-19)	73 (30-140)
Senegal	2000	Female	90 to 94	4.8% (2.9-7.4%)	95% (93-97%)	0.22 (0.11-0.41)	0.77% (0.23-1.7%)	0.87% (0.22-2%)	2.9 (0.75-6.6)	15 (4.6-34)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	2000	Female	95 plus	5.5% (3.1-8.9%)	94% (91-97%)	0.24 (0.093-0.5)	0.73% (0.18-1.9%)	0.78% (0.12-2.1%)	0.66 (0.1-1.8)	2.5 (0.62-6.2)
Senegal	2000	Male	All Ages	3.3% (2.7-3.9%)	97% (96-97%)	0.072 (0.056-0.09)	0.6% (0.45-0.77%)	0.87% (0.6-1.2%)	410 (280-580)	19000 (15000-25000)
Senegal	2000	Male	15 to 19	2.7% (1.7-4%)	97% (96-98%)	0.045 (0.023-0.078)	0.58% (0.38-0.83%)	0.25% (0.12-0.44%)	2.4 (1.2-4.3)	660 (420-980)
Senegal	2000	Male	20 to 24	4% (2.4-6%)	96% (94-98%)	0.094 (0.049-0.15)	1.3% (0.9-1.8%)	0.69% (0.37-1.2%)	9 (4.9-15)	1600 (1100-2300)
Senegal	2000	Male	25 to 29	5.6% (3.4-8.1%)	94% (92-97%)	0.16 (0.083-0.26)	1.9% (1.3-2.6%)	1.4% (0.76-2.4%)	18 (9.8-30)	2000 (1400-2800)
Senegal	2000	Male	30 to 34	6.3% (3.9-9.4%)	94% (91-96%)	0.19 (0.1-0.3)	2.2% (1.5-3.2%)	2.1% (1.2-3.2%)	26 (14-41)	2100 (1400-3000)
Senegal	2000	Male	35 to 39	5.8% (3.4-8.7%)	94% (91-97%)	0.17 (0.087-0.29)	2.5% (1.7-3.4%)	2.6% (1.6-3.8%)	32 (20-48)	2100 (1500-2900)
Senegal	2000	Male	40 to 44	5.9% (3.7-8.5%)	94% (92-96%)	0.17 (0.089-0.3)	2.6% (1.8-3.5%)	2.9% (1.9-4.1%)	37 (24-53)	2000 (1300-2800)
Senegal	2000	Male	45 to 49	6.5% (4.9-1%)	94% (91-96%)	0.2 (0.11-0.34)	2.9% (1.9-4.1%)	3.3% (2-4.9%)	45 (27-67)	2100 (1400-3000)
Senegal	2000	Male	50 to 54	6.6% (4.3-9.7%)	93% (90-96%)	0.2 (0.1-0.35)	2.5% (1.7-3.7%)	3% (1.8-4.4%)	43 (27-65)	1700 (1100-2500)
Senegal	2000	Male	55 to 59	6.4% (3.9-9.5%)	94% (91-96%)	0.19 (0.1-0.33)	2.4% (1.5-3.5%)	2.8% (1.6-4.2%)	50 (30-76)	1700 (1100-2500)
Senegal	2000	Male	60 to 64	6.4% (3.9-9.7%)	94% (90-96%)	0.17 (0.085-0.31)	2% (1.2-2.9%)	2.3% (1.3-3.5%)	51 (29-81)	1500 (890-2300)
Senegal	2000	Male	65 to 69	6.8% (4.4-9.7%)	93% (90-96%)	0.16 (0.081-0.26)	1.4% (0.9-2.1%)	1.6% (0.99-2.5%)	41 (24-63)	970 (610-1500)
Senegal	2000	Male	70 to 74	6.7% (4.5-9.7%)	93% (90-95%)	0.13 (0.066-0.22)	1% (0.6-1.6%)	1.2% (0.66-1.8%)	31 (17-48)	570 (330-880)
Senegal	2000	Male	75 to 79	5.8% (3.5-8.9%)	94% (91-97%)	0.09 (0.043-0.17)	0.72% (0.41-1.1%)	0.81% (0.44-1.3%)	17 (9.5-28)	250 (140-400)
Senegal	2000	Male	80 to 84	4.8% (2.8-7.1%)	95% (93-97%)	0.066 (0.03-0.12)	0.61% (0.36-0.9%)	0.68% (0.38-1%)	7.8 (4.3-12)	85 (50-120)
Senegal	2000	Male	85 to 89	4.4% (2.5-6.9%)	96% (93-98%)	0.052 (0.025-0.094)	0.47% (0.27-0.7%)	0.51% (0.26-0.77%)	3.2 (1.6-4.8)	26 (14-37)
Senegal	2000	Male	90 to 94	4.4% (2.7-6.8%)	96% (93-97%)	0.044 (0.02-0.086)	0.3% (0.14-0.48%)	0.3% (0.1-0.5%)	0.65 (0.22-1.1)	4 (1.9-6.2)
Senegal	2000	Male	95 plus	5% (2.8-8.2%)	95% (92-97%)	0.04 (0.015-0.083)	0.3% (0.12-0.49%)	0.26% (0.021-0.49%)	0.13 (0.01-0.24)	0.61 (0.23-0.97)
Senegal	2005	Female	All Ages	2.9% (2.4-3.5%)	97% (96-98%)	0.14 (0.11-0.17)	0.74% (0.57-0.94%)	1.2% (0.84-1.8%)	560 (380-800)	22000 (17000-28000)
Senegal	2005	Female	15 to 19	2.7% (1.6-4.2%)	97% (96-98%)	0.12 (0.06-0.2)	0.68% (0.44-1%)	0.41% (0.17-0.79%)	3.7 (1.5-7.2)	820 (510-1200)
Senegal	2005	Female	20 to 24	3.4% (2-5.2%)	97% (95-98%)	0.19 (0.1-0.31)	1.4% (0.94-2%)	0.75% (0.36-1.3%)	8.1 (3.8-14)	1700 (1100-2600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	2005	Female	25 to 29	3.8% (2.2-6%)	96% (94-98%)	0.23 (0.12-0.38)	1.6% (1.1-2.2%)	1.1% (0.63-1.9%)	13 (7.2-22)	1900 (1200-2700)
Senegal	2005	Female	30 to 34	4.1% (2.3-6.7%)	96% (93-98%)	0.25 (0.13-0.42)	1.6% (1.1-2.2%)	1.4% (0.84-2.2%)	16 (9.8-26)	1700 (1100-2400)
Senegal	2005	Female	35 to 39	4.5% (2.5-6.8%)	96% (93-97%)	0.27 (0.14-0.45)	1.7% (1.2-2.4%)	1.8% (1.1-2.8%)	21 (13-33)	1600 (1100-2200)
Senegal	2005	Female	40 to 44	4.9% (3-7.2%)	95% (93-97%)	0.29 (0.15-0.5)	2% (1.3-2.9%)	2.5% (1.6-3.8%)	30 (18-46)	1700 (1200-2500)
Senegal	2005	Female	45 to 49	5.2% (3-7.5%)	95% (92-97%)	0.3 (0.15-0.52)	2.2% (1.4-3.2%)	2.8% (1.7-4.2%)	37 (22-57)	1700 (1100-2600)
Senegal	2005	Female	50 to 54	5.1% (3.1-7.6%)	95% (92-97%)	0.31 (0.15-0.54)	2.5% (1.6-3.6%)	3.2% (1.9-4.7%)	50 (29-77)	2000 (1200-2900)
Senegal	2005	Female	55 to 59	5.2% (3.1-7.8%)	95% (92-97%)	0.36 (0.18-0.62)	2.6% (1.5-4%)	3.3% (1.9-5.1%)	60 (33-97)	2000 (1200-3100)
Senegal	2005	Female	60 to 64	5.7% (3.4-8.9%)	94% (91-97%)	0.4 (0.2-0.73)	2.5% (1.4-4.2%)	3.1% (1.8-5.2%)	74 (42-120)	2100 (1200-3400)
Senegal	2005	Female	65 to 69	6.3% (3.8-9.3%)	94% (91-96%)	0.39 (0.2-0.66)	2.4% (1.4-4%)	2.9% (1.6-4.8%)	85 (47-150)	2000 (1100-3300)
Senegal	2005	Female	70 to 74	5.7% (3.6-8.5%)	94% (92-96%)	0.31 (0.15-0.56)	2% (1-3.5%)	2.4% (1.2-4.2%)	78 (39-140)	1400 (740-2600)
Senegal	2005	Female	75 to 79	4.6% (2.6-7.1%)	95% (93-97%)	0.22 (0.1-0.44)	1.4% (0.72-2.7%)	1.6% (0.81-3.1%)	49 (24-92)	710 (360-1300)
Senegal	2005	Female	80 to 84	4.1% (2.3-6%)	96% (94-98%)	0.19 (0.089-0.34)	1% (0.48-1.8%)	1.1% (0.53-2.1%)	22 (9.7-41)	230 (110-430)
Senegal	2005	Female	85 to 89	4.3% (2.6-6.7%)	96% (93-97%)	0.19 (0.09-0.35)	0.8% (0.34-1.6%)	0.9% (0.36-1.8%)	10 (4-21)	77 (32-150)
Senegal	2005	Female	90 to 94	4.8% (3-7.5%)	95% (92-97%)	0.2 (0.097-0.38)	0.65% (0.19-1.5%)	0.72% (0.17-1.7%)	3 (0.69-7.1)	16 (4.7-37)
Senegal	2005	Female	95 plus	5.6% (3.2-9%)	94% (91-97%)	0.22 (0.09-0.46)	0.63% (0.13-1.5%)	0.66% (0.061-1.7%)	0.72 (0.066-1.9)	2.7 (0.57-6.6)
Senegal	2005	Male	All Ages	3.3% (2.7-3.9%)	97% (96-97%)	0.066 (0.051-0.084)	0.64% (0.48-0.81%)	0.91% (0.62-1.2%)	440 (310-610)	21000 (16000-27000)
Senegal	2005	Male	15 to 19	2.7% (1.6-4%)	97% (96-98%)	0.041 (0.021-0.072)	0.6% (0.4-0.86%)	0.24% (0.13-0.42%)	2.6 (1.4-4.3)	740 (480-1100)
Senegal	2005	Male	20 to 24	4.1% (2.4-6%)	96% (94-98%)	0.086 (0.045-0.14)	1.3% (0.95-1.9%)	0.68% (0.38-1.1%)	9.4 (5.3-15)	1800 (1200-2500)
Senegal	2005	Male	25 to 29	5.6% (3.5-8.1%)	94% (92-97%)	0.14 (0.078-0.23)	1.8% (1.3-2.5%)	1.3% (0.76-2.1%)	20 (11-32)	2300 (1600-3100)
Senegal	2005	Male	30 to 34	6.2% (3.9-9.4%)	94% (91-96%)	0.17 (0.095-0.27)	2.1% (1.5-2.9%)	1.9% (1.1-3%)	29 (17-43)	2400 (1700-3300)
Senegal	2005	Male	35 to 39	5.8% (3.3-8.7%)	94% (91-97%)	0.16 (0.082-0.26)	2.4% (1.7-3.3%)	2.5% (1.6-3.7%)	35 (23-53)	2300 (1600-3200)
Senegal	2005	Male	40 to 44	5.9% (3.7-8.6%)	94% (91-96%)	0.16 (0.08-0.27)	2.5% (1.7-3.5%)	2.8% (1.8-4.1%)	40 (26-60)	2200 (1500-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	2005	Male	45 to 49	6.5% (3.9-9.3%)	94% (91-96%)	0.18 (0.096-0.31)	2.8% (1.8-3.9%)	3.2% (2-4.7%)	48 (30-70)	2200 (1500-3100)
Senegal	2005	Male	50 to 54	6.6% (4.2-9.8%)	93% (90-96%)	0.18 (0.093-0.31)	2.5% (1.6-3.6%)	2.9% (1.8-4.4%)	47 (28-71)	1900 (1200-2700)
Senegal	2005	Male	55 to 59	6.4% (4-9.4%)	94% (91-96%)	0.17 (0.089-0.29)	2.4% (1.5-3.5%)	2.8% (1.6-4.1%)	48 (29-72)	1600 (1000-2400)
Senegal	2005	Male	60 to 64	6.3% (3.9-9.6%)	94% (90-96%)	0.16 (0.079-0.29)	1.9% (1.2-3%)	2.3% (1.4-3.5%)	52 (31-82)	1500 (930-2300)
Senegal	2005	Male	65 to 69	6.7% (4.4-9.5%)	93% (90-96%)	0.14 (0.07-0.24)	1.4% (0.88-2%)	1.6% (0.98-2.4%)	43 (26-64)	1000 (650-1500)
Senegal	2005	Male	70 to 74	6.7% (4.5-9.7%)	93% (90-96%)	0.12 (0.058-0.2)	1% (0.61-1.5%)	1.2% (0.69-1.8%)	33 (19-52)	620 (370-940)
Senegal	2005	Male	75 to 79	5.8% (3.5-8.9%)	94% (91-96%)	0.082 (0.039-0.15)	0.72% (0.43-1.1%)	0.82% (0.47-1.3%)	20 (11-30)	290 (170-430)
Senegal	2005	Male	80 to 84	4.8% (2.9-7.1%)	95% (93-97%)	0.061 (0.029-0.11)	0.63% (0.38-0.92%)	0.71% (0.41-1.1%)	9.4 (5.3-14)	100 (61-150)
Senegal	2005	Male	85 to 89	4.4% (2.5-6.8%)	96% (93-97%)	0.047 (0.023-0.086)	0.49% (0.3-0.71%)	0.53% (0.29-0.8%)	4.1 (2.2-6)	32 (19-46)
Senegal	2005	Male	90 to 94	4.5% (2.7-7%)	96% (93-97%)	0.04 (0.019-0.078)	0.31% (0.16-0.48%)	0.31% (0.11-0.52%)	0.87 (0.31-1.5)	5.3 (2.6-8.1)
Senegal	2005	Male	95 plus	5.1% (2.8-8.3%)	95% (92-97%)	0.037 (0.014-0.078)	0.32% (0.13-0.51%)	0.28% (0.048-0.5%)	0.18 (0.032-0.32)	0.84 (0.33-1.3)
Senegal	2010	Female	All Ages	2.6% (2.1-3.2%)	97% (97-98%)	0.12 (0.086-0.15)	0.77% (0.6-0.98%)	1.2% (0.82-1.7%)	550 (370-790)	22000 (17000-28000)
Senegal	2010	Female	15 to 19	2.2% (1.4-3.4%)	98% (97-99%)	0.093 (0.045-0.16)	0.66% (0.43-0.95%)	0.31% (0.13-0.62%)	2.5 (1-5.2)	800 (490-1200)
Senegal	2010	Female	20 to 24	2.9% (1.8-4.4%)	97% (96-98%)	0.16 (0.078-0.27)	1.4% (0.95-2.1%)	0.6% (0.28-1.1%)	6 (2.7-11)	1800 (1100-2700)
Senegal	2010	Female	25 to 29	3.6% (2.3-5.2%)	96% (95-98%)	0.2 (0.11-0.33)	1.6% (1.1-2.1%)	1% (0.54-1.7%)	11 (5.9-18)	1900 (1300-2600)
Senegal	2010	Female	30 to 34	3.9% (2.4-6.1%)	96% (94-98%)	0.23 (0.12-0.36)	1.6% (1.2-2.1%)	1.3% (0.83-2.1%)	16 (9.5-25)	1800 (1300-2500)
Senegal	2010	Female	35 to 39	4% (2.4-6.1%)	96% (94-98%)	0.23 (0.12-0.38)	1.6% (1.1-2.2%)	1.7% (1.1-2.5%)	21 (13-31)	1700 (1200-2300)
Senegal	2010	Female	40 to 44	4.1% (2.6-5.8%)	96% (94-97%)	0.23 (0.11-0.38)	1.9% (1.3-2.6%)	2.3% (1.6-3.5%)	30 (19-45)	1800 (1200-2500)
Senegal	2010	Female	45 to 49	3.9% (2.4-5.7%)	96% (94-98%)	0.21 (0.11-0.38)	1.9% (1.3-2.6%)	2.4% (1.5-3.5%)	35 (22-52)	1700 (1100-2400)
Senegal	2010	Female	50 to 54	4% (2.6-5.9%)	96% (94-97%)	0.24 (0.12-0.4)	2.1% (1.4-3%)	2.7% (1.7-4%)	50 (30-74)	2000 (1300-2800)
Senegal	2010	Female	55 to 59	4.7% (3.1-6.6%)	95% (93-97%)	0.31 (0.16-0.53)	2.3% (1.4-3.6%)	2.9% (1.7-4.5%)	62 (36-100)	2100 (1200-3200)
Senegal	2010	Female	60 to 64	5.6% (3.8-8%)	94% (92-96%)	0.37 (0.19-0.65)	2.3% (1.4-3.7%)	2.9% (1.6-4.5%)	72 (41-120)	2000 (1200-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	2010	Female	65 to 69	6.6% (4.5-9%)	93% (91-96%)	0.39 (0.21-0.65)	2.3% (1.3-3.7%)	2.7% (1.5-4.4%)	88 (48-150)	2000 (1100-3400)
Senegal	2010	Female	70 to 74	5.8% (4-8.5%)	94% (92-96%)	0.3 (0.15-0.52)	1.8% (0.87-3.2%)	2.1% (1-3.8%)	77 (37-140)	1400 (720-2600)
Senegal	2010	Female	75 to 79	4.1% (2.5-6.4%)	96% (94-97%)	0.19 (0.088-0.36)	1.2% (0.62-2.2%)	1.4% (0.69-2.6%)	49 (24-90)	700 (360-1300)
Senegal	2010	Female	80 to 84	3.7% (2.2-5.5%)	96% (95-98%)	0.16 (0.078-0.29)	0.87% (0.42-1.6%)	0.98% (0.46-1.8%)	22 (10-41)	240 (120-430)
Senegal	2010	Female	85 to 89	4% (2.3-6.1%)	96% (94-98%)	0.17 (0.078-0.31)	0.69% (0.29-1.4%)	0.77% (0.31-1.5%)	11 (4.2-22)	82 (35-160)
Senegal	2010	Female	90 to 94	4.5% (2.7-6.8%)	96% (93-97%)	0.18 (0.083-0.33)	0.55% (0.16-1.2%)	0.6% (0.12-1.4%)	3.2 (0.66-7.5)	18 (4.9-40)
Senegal	2010	Female	95 plus	5.2% (2.9-8.3%)	95% (92-97%)	0.19 (0.077-0.38)	0.53% (0.096-1.3%)	0.54% (0.021-1.4%)	0.77 (0.029-2.1)	3 (0.54-7.3)
Senegal	2010	Male	All Ages	3.1% (2.7-3.7%)	97% (96-97%)	0.063 (0.047-0.083)	0.74% (0.56-0.94%)	1% (0.69-1.4%)	490 (330-680)	24000 (18000-31000)
Senegal	2010	Male	15 to 19	2% (1.3-2.9%)	98% (97-99%)	0.03 (0.015-0.052)	0.58% (0.39-0.83%)	0.21% (0.12-0.34%)	2.3 (1.3-3.8)	790 (500-1200)
Senegal	2010	Male	20 to 24	3% (2.1-4.3%)	97% (96-98%)	0.061 (0.033-0.1)	1.3% (0.91-1.8%)	0.58% (0.35-0.9%)	8.8 (5.3-14)	1900 (1300-2700)
Senegal	2010	Male	25 to 29	3.9% (2.7-5.6%)	96% (94-97%)	0.096 (0.052-0.15)	1.7% (1.2-2.3%)	1.1% (0.68-1.8%)	18 (11-28)	2400 (1700-3300)
Senegal	2010	Male	30 to 34	5% (3.3-7.1%)	95% (93-97%)	0.13 (0.074-0.21)	2.1% (1.5-2.8%)	1.8% (1.1-2.7%)	30 (18-44)	2700 (1900-3600)
Senegal	2010	Male	35 to 39	6.3% (4.3-8.5%)	94% (91-96%)	0.16 (0.089-0.26)	2.4% (1.8-3.3%)	2.6% (1.6-3.8%)	43 (26-65)	2800 (2000-3900)
Senegal	2010	Male	40 to 44	7.9% (5.7-10%)	92% (90-94%)	0.2 (0.11-0.33)	2.7% (1.8-3.8%)	3.1% (2-4.5%)	50 (31-73)	2700 (1800-3700)
Senegal	2010	Male	45 to 49	8.3% (5.8-11%)	92% (89-94%)	0.22 (0.12-0.37)	2.9% (2-4.1%)	3.4% (2.2-5.1%)	58 (35-86)	2700 (1700-3800)
Senegal	2010	Male	50 to 54	8.3% (5.9-11%)	92% (89-94%)	0.22 (0.12-0.36)	2.6% (1.7-3.8%)	3.1% (1.9-4.6%)	55 (33-84)	2200 (1400-3200)
Senegal	2010	Male	55 to 59	9.1% (6.4-12%)	91% (88-94%)	0.24 (0.13-0.38)	2.5% (1.6-3.6%)	3% (1.7-4.4%)	59 (34-87)	2000 (1200-2900)
Senegal	2010	Male	60 to 64	8.8% (6.2-12%)	91% (88-94%)	0.21 (0.11-0.35)	2% (1.3-3.2%)	2.3% (1.4-3.8%)	52 (31-82)	1500 (920-2300)
Senegal	2010	Male	65 to 69	7.9% (5.6-10%)	92% (90-94%)	0.16 (0.088-0.26)	1.4% (0.85-2%)	1.6% (0.93-2.4%)	44 (26-66)	1000 (640-1500)
Senegal	2010	Male	70 to 74	7.2% (5.1-10%)	93% (90-95%)	0.12 (0.062-0.2)	0.99% (0.59-1.5%)	1.2% (0.66-1.8%)	35 (20-53)	650 (390-980)
Senegal	2010	Male	75 to 79	6.3% (4.1-9.1%)	94% (91-96%)	0.086 (0.041-0.16)	0.71% (0.41-1.1%)	0.81% (0.45-1.2%)	21 (12-33)	310 (180-460)
Senegal	2010	Male	80 to 84	5.1% (3.1-7.4%)	95% (93-97%)	0.062 (0.029-0.11)	0.63% (0.37-0.9%)	0.71% (0.4-1%)	11 (5.9-16)	110 (68-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	2010	Male	85 to 89	4.4% (2.5-6.7%)	96% (93-98%)	0.046 (0.022-0.083)	0.49% (0.3-0.71%)	0.53% (0.29-0.8%)	4.8 (2.7-7.3)	39 (23-56)
Senegal	2010	Male	90 to 94	4.5% (2.8-6.9%)	96% (93-97%)	0.039 (0.018-0.074)	0.31% (0.16-0.47%)	0.31% (0.12-0.5%)	1.1 (0.42-1.8)	6.7 (3.4-10)
Senegal	2010	Male	95 plus	5.1% (2.9-8.1%)	95% (92-97%)	0.036 (0.013-0.075)	0.32% (0.14-0.5%)	0.27% (0.058-0.49%)	0.23 (0.05-0.42)	1.1 (0.48-1.7)
Senegal	2016	Female	All Ages	2.6% (2.2-3.2%)	97% (97-98%)	0.12 (0.068-0.19)	0.92% (0.65-1.2%)	1.4% (0.77-2.1%)	630 (340-980)	26000 (17000-36000)
Senegal	2016	Female	15 to 19	2.2% (1.4-3.4%)	98% (97-99%)	0.098 (0.042-0.19)	0.72% (0.46-1%)	0.32% (0.12-0.72%)	2.5 (0.89-5.5)	890 (530-1300)
Senegal	2016	Female	20 to 24	2.9% (1.8-4.4%)	97% (96-98%)	0.16 (0.075-0.32)	1.6% (1-2.2%)	0.63% (0.27-1.3%)	5.8 (2.4-11)	2000 (1200-3000)
Senegal	2016	Female	25 to 29	3.6% (2.2-5.3%)	96% (95-98%)	0.21 (0.1-0.38)	1.7% (1.2-2.3%)	1% (0.53-1.9%)	11 (5.4-20)	2200 (1400-3100)
Senegal	2016	Female	30 to 34	3.9% (2.4-5.9%)	96% (94-98%)	0.23 (0.11-0.44)	1.7% (1.2-2.3%)	1.4% (0.77-2.3%)	17 (9.2-28)	2100 (1400-3000)
Senegal	2016	Female	35 to 39	4% (2.4-6%)	96% (94-98%)	0.24 (0.1-0.44)	1.6% (1.2-2.3%)	1.7% (1-2.7%)	25 (14-40)	2100 (1400-2900)
Senegal	2016	Female	40 to 44	4.1% (2.7-6%)	96% (94-97%)	0.24 (0.1-0.46)	1.9% (1.3-2.7%)	2.4% (1.4-3.6%)	38 (22-58)	2300 (1500-3300)
Senegal	2016	Female	45 to 49	3.9% (2.5-5.7%)	96% (94-98%)	0.22 (0.096-0.42)	1.9% (1.2-2.8%)	2.4% (1.4-3.7%)	43 (24-70)	2100 (1300-3200)
Senegal	2016	Female	50 to 54	4% (2.6-5.9%)	96% (94-97%)	0.24 (0.11-0.45)	2.1% (1.3-3.1%)	2.7% (1.5-4%)	58 (32-90)	2300 (1400-3500)
Senegal	2016	Female	55 to 59	4.7% (3.1-6.8%)	95% (93-97%)	0.32 (0.15-0.61)	2.3% (1.2-3.6%)	2.8% (1.5-4.6%)	72 (36-120)	2400 (1300-3800)
Senegal	2016	Female	60 to 64	5.6% (3.8-7.9%)	94% (92-96%)	0.39 (0.17-0.72)	2.3% (1.2-3.7%)	2.8% (1.5-4.6%)	84 (43-140)	2400 (1200-3800)
Senegal	2016	Female	65 to 69	6.6% (4.5-8.9%)	93% (91-96%)	0.4 (0.19-0.75)	2.2% (1-3.8%)	2.7% (1.2-4.6%)	90 (40-150)	2100 (960-3600)
Senegal	2016	Female	70 to 74	5.8% (4-8.2%)	94% (92-96%)	0.31 (0.14-0.62)	1.8% (0.78-3.4%)	2.1% (0.87-4.1%)	82 (34-160)	1500 (660-3000)
Senegal	2016	Female	75 to 79	4.1% (2.6-6.4%)	96% (94-97%)	0.2 (0.081-0.41)	1.2% (0.54-2.3%)	1.4% (0.6-2.7%)	55 (23-110)	790 (350-1600)
Senegal	2016	Female	80 to 84	3.7% (2.1-5.5%)	96% (94-98%)	0.17 (0.07-0.33)	0.88% (0.36-1.7%)	1% (0.38-2%)	27 (10-54)	290 (120-570)
Senegal	2016	Female	85 to 89	4% (2.3-6.2%)	96% (94-98%)	0.18 (0.067-0.35)	0.71% (0.23-1.6%)	0.8% (0.24-1.8%)	13 (4-30)	100 (34-220)
Senegal	2016	Female	90 to 94	4.5% (2.8-7.1%)	95% (93-97%)	0.19 (0.075-0.39)	0.58% (0.12-1.5%)	0.63% (0.089-1.7%)	4.3 (0.61-12)	24 (5-61)
Senegal	2016	Female	95 plus	5.2% (2.9-8.2%)	95% (92-97%)	0.2 (0.07-0.46)	0.57% (0.088-1.6%)	0.57% (0.0054-1.8%)	1.1 (0.0098-3.4)	4.2 (0.66-12)
Senegal	2016	Male	All Ages	3.2% (2.8-3.8%)	97% (96-97%)	0.067 (0.039-0.11)	0.9% (0.63-1.2%)	1.2% (0.68-1.7%)	570 (320-840)	29000 (19000-39000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Senegal	2016	Male	15 to 19	2% (1.3-2.9%)	98% (97-99%)	0.031 (0.014-0.059)	0.61% (0.4-0.88%)	0.21% (0.11-0.36%)	2.6 (1.4-4.3)	900 (570-1400)
Senegal	2016	Male	20 to 24	3.1% (2-4.4%)	97% (96-98%)	0.064 (0.031-0.13)	1.4% (0.93-1.9%)	0.59% (0.33-0.97%)	10 (5.4-17)	2200 (1500-3300)
Senegal	2016	Male	25 to 29	4% (2.8-5.6%)	96% (94-97%)	0.1 (0.048-0.18)	1.8% (1.3-2.5%)	1.2% (0.69-2%)	22 (13-37)	2900 (2000-4100)
Senegal	2016	Male	30 to 34	5% (3.4-7.3%)	95% (93-97%)	0.13 (0.068-0.25)	2.2% (1.5-3%)	2% (1.1-3%)	37 (20-59)	3300 (2200-4700)
Senegal	2016	Male	35 to 39	6.3% (4.2-8.5%)	94% (91-96%)	0.17 (0.078-0.3)	2.6% (1.7-3.7%)	2.7% (1.5-4.3%)	54 (29-85)	3600 (2300-5200)
Senegal	2016	Male	40 to 44	7.9% (5.8-11%)	92% (89-94%)	0.21 (0.1-0.39)	2.8% (1.7-4%)	3.2% (1.8-5%)	64 (35-100)	3500 (2100-5200)
Senegal	2016	Male	45 to 49	8.4% (5.6-11%)	92% (89-94%)	0.23 (0.11-0.43)	3% (1.8-4.5%)	3.6% (1.9-5.6%)	71 (37-110)	3300 (1900-5000)
Senegal	2016	Male	50 to 54	8.4% (5.9-11%)	92% (89-94%)	0.23 (0.11-0.42)	2.6% (1.6-4%)	3.1% (1.7-4.9%)	65 (34-100)	2600 (1500-3900)
Senegal	2016	Male	55 to 59	9.2% (6.5-12%)	91% (88-94%)	0.25 (0.11-0.45)	2.5% (1.4-4%)	3% (1.6-4.8%)	67 (34-110)	2300 (1300-3500)
Senegal	2016	Male	60 to 64	8.9% (6.2-12%)	91% (88-94%)	0.22 (0.097-0.4)	2% (1.1-3.2%)	2.3% (1.2-3.9%)	59 (28-100)	1700 (900-2800)
Senegal	2016	Male	65 to 69	8% (5.7-10%)	92% (90-94%)	0.17 (0.079-0.31)	1.3% (0.75-2.1%)	1.5% (0.82-2.5%)	42 (21-68)	990 (540-1600)
Senegal	2016	Male	70 to 74	7.2% (5.2-10%)	93% (90-95%)	0.12 (0.058-0.24)	0.95% (0.52-1.5%)	1.1% (0.56-1.8%)	35 (17-56)	650 (350-1000)
Senegal	2016	Male	75 to 79	6.3% (4.2-9.1%)	94% (91-96%)	0.09 (0.038-0.18)	0.69% (0.36-1.1%)	0.78% (0.37-1.3%)	23 (11-38)	340 (180-530)
Senegal	2016	Male	80 to 84	5.2% (3.2-7.6%)	95% (92-97%)	0.064 (0.027-0.13)	0.61% (0.31-0.95%)	0.68% (0.31-1.1%)	12 (5.6-20)	130 (68-210)
Senegal	2016	Male	85 to 89	4.4% (2.5-6.7%)	96% (93-97%)	0.048 (0.018-0.094)	0.47% (0.24-0.72%)	0.51% (0.22-0.82%)	5.7 (2.5-9.2)	46 (23-70)
Senegal	2016	Male	90 to 94	4.5% (2.8-6.8%)	96% (93-97%)	0.04 (0.016-0.086)	0.31% (0.14-0.49%)	0.3% (0.079-0.52%)	1.4 (0.35-2.4)	8.5 (3.6-14)
Senegal	2016	Male	95 plus	5.1% (2.8-8%)	95% (92-97%)	0.036 (0.012-0.085)	0.31% (0.12-0.51%)	0.26% (0.026-0.5%)	0.31 (0.03-0.59)	1.5 (0.56-2.4)
Sierra Leone	1990	Female	All Ages	25% (19-31%)	75% (69-81%)	1.4 (1-1.9)	1.3% (0.88-1.7%)	2.9% (1.7-4%)	780 (470-1100)	27000 (18000-35000)
Sierra Leone	1990	Female	15 to 19	23% (12-37%)	77% (63-88%)	1.2 (0.5-2.4)	1.8% (0.85-3.1%)	2.2% (0.77-4.2%)	7.3 (2.5-15)	740 (350-1300)
Sierra Leone	1990	Female	20 to 24	28% (15-45%)	72% (55-85%)	1.9 (0.79-3.8)	3.4% (1.9-5.3%)	4% (1.6-6.9%)	15 (6.1-27)	1400 (740-2300)
Sierra Leone	1990	Female	25 to 29	31% (16-49%)	69% (51-84%)	2.4 (0.99-4.4)	4.4% (2.6-6.6%)	5.7% (2.9-9.2%)	22 (11-37)	1700 (980-2600)
Sierra Leone	1990	Female	30 to 34	34% (17-53%)	66% (47-83%)	2.7 (1.2-4.8)	4.9% (3-7.3%)	6.7% (3.7-10%)	27 (14-43)	1800 (1100-2700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	1990	Female	35 to 39	37% (20-56%)	63% (44-80%)	2.8 (1.3-4.9)	5.8% (3.4-8.4%)	8.2% (4.5-12%)	37 (20-56)	2100 (1200-3100)
Sierra Leone	1990	Female	40 to 44	39% (22-57%)	61% (43-78%)	2.9 (1.3-5.3)	6.6% (3.9-9.5%)	9.6% (5.4-14%)	47 (26-68)	2300 (1400-3400)
Sierra Leone	1990	Female	45 to 49	39% (21-59%)	61% (41-79%)	3 (1.3-5.9)	7.9% (4.5-12%)	11% (6.2-16%)	61 (33-93)	2600 (1500-4000)
Sierra Leone	1990	Female	50 to 54	40% (22-61%)	60% (39-78%)	3.2 (1.3-5.9)	8.9% (4.8-13%)	12% (6.6-18%)	77 (40-120)	2900 (1500-4300)
Sierra Leone	1990	Female	55 to 59	40% (23-59%)	60% (41-77%)	3.5 (1.5-6.6)	9.6% (5.1-14%)	13% (6.8-19%)	95 (50-140)	3100 (1600-4500)
Sierra Leone	1990	Female	60 to 64	41% (23-61%)	59% (39-77%)	3.5 (1.4-7)	8.8% (4.4-13%)	11% (5.8-17%)	100 (50-160)	2800 (1400-4200)
Sierra Leone	1990	Female	65 to 69	42% (25-60%)	58% (40-75%)	3.2 (1.4-5.9)	9% (3.9-14%)	11% (4.9-18%)	110 (48-180)	2500 (1100-4000)
Sierra Leone	1990	Female	70 to 74	40% (25-58%)	60% (42-75%)	2.6 (1-5.2)	8.1% (2.7-15%)	9.8% (3.3-18%)	96 (31-180)	1700 (570-3200)
Sierra Leone	1990	Female	75 to 79	36% (20-55%)	64% (45-80%)	2.2 (0.74-4.7)	6.3% (1.3-12%)	7.5% (1.6-15%)	52 (11-99)	720 (150-1400)
Sierra Leone	1990	Female	80 to 84	34% (18-52%)	66% (48-82%)	2 (0.65-4.3)	5.4% (1-11%)	6.4% (1.2-13%)	17 (3.2-35)	170 (34-360)
Sierra Leone	1990	Female	85 to 89	34% (18-52%)	66% (48-82%)	2 (0.68-4.3)	4.7% (0.76-10%)	5.6% (0.78-12%)	9.8 (1.4-21)	73 (12-160)
Sierra Leone	1990	Female	90 to 94	37% (20-55%)	63% (45-80%)	2.1 (0.66-4.5)	4.3% (0.35-9.6%)	5.1% (0.31-12%)	3.6 (0.23-8.3)	19 (1.5-42)
Sierra Leone	1990	Female	95 plus	41% (21-62%)	59% (38-79%)	2.2 (0.59-5.2)	4.6% (-0.18-9.7%)	4.6% (-0.35-12%)	0.97 (-0.073-2.5)	3.4 (-0.15-8.3)
Sierra Leone	1990	Male	All Ages	20% (16-26%)	80% (74-84%)	1.7 (1.2-2.3)	2.2% (1.7-2.8%)	4.3% (3.2-5.6%)	1400 (980-1800)	52000 (40000-65000)
Sierra Leone	1990	Male	15 to 19	15% (8-26%)	85% (74-92%)	0.93 (0.35-1.9)	2.3% (1.1-4%)	2.8% (1.1-5.4%)	12 (4.6-24)	1100 (490-2000)
Sierra Leone	1990	Male	20 to 24	23% (12-38%)	77% (62-88%)	1.9 (0.74-3.8)	6% (3.2-9.3%)	7.2% (3.4-12%)	37 (17-61)	2800 (1500-4400)
Sierra Leone	1990	Male	25 to 29	28% (15-45%)	72% (55-85%)	2.8 (1.1-5.1)	9.2% (5.2-13%)	12% (6.2-17%)	57 (30-86)	3900 (2200-5600)
Sierra Leone	1990	Male	30 to 34	32% (17-51%)	68% (49-83%)	3.5 (1.5-6.1)	11% (7.3-16%)	15% (9.1-20%)	75 (46-110)	4500 (2900-6300)
Sierra Leone	1990	Male	35 to 39	35% (19-54%)	65% (46-81%)	3.8 (1.7-6.9)	13% (8.5-17%)	16% (11-22%)	94 (62-130)	5000 (3400-7000)
Sierra Leone	1990	Male	40 to 44	36% (21-55%)	64% (45-79%)	4.1 (1.7-7.8)	14% (8.9-19%)	18% (11-24%)	120 (80-170)	5900 (3900-8000)
Sierra Leone	1990	Male	45 to 49	37% (19-56%)	63% (44-81%)	4.3 (1.7-8.5)	15% (9.5-20%)	19% (12-26%)	150 (99-210)	6500 (4200-8800)
Sierra Leone	1990	Male	50 to 54	37% (20-58%)	63% (42-80%)	4.4 (1.8-8.4)	14% (9.1-19%)	18% (11-24%)	160 (100-230)	6000 (3800-8300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	1990	Male	55 to 59	38% (22-57%)	62% (43-78%)	4.3 (1.8-8.1)	13% (8.2-17%)	16% (10-22%)	160 (100-230)	5300 (3400-7300)
Sierra Leone	1990	Male	60 to 64	37% (21-57%)	63% (43-79%)	3.9 (1.4-8)	11% (6.3-16%)	13% (7.6-19%)	150 (90-230)	4200 (2500-6100)
Sierra Leone	1990	Male	65 to 69	37% (21-55%)	63% (45-79%)	3.2 (1.3-6.5)	9.8% (5.1-15%)	12% (6.1-18%)	140 (73-220)	3300 (1700-5000)
Sierra Leone	1990	Male	70 to 74	35% (21-53%)	65% (47-79%)	2.5 (0.92-5.1)	8.2% (3.1-14%)	9.7% (3.7-16%)	110 (42-190)	2100 (770-3500)
Sierra Leone	1990	Male	75 to 79	32% (18-52%)	68% (48-82%)	1.9 (0.61-4.3)	6.1% (1.7-11%)	7.1% (2-13%)	60 (16-110)	840 (230-1600)
Sierra Leone	1990	Male	80 to 84	30% (15-48%)	70% (52-85%)	1.6 (0.51-3.5)	5.1% (1.4-10%)	5.9% (1.6-12%)	21 (5.7-43)	220 (62-440)
Sierra Leone	1990	Male	85 to 89	28% (15-46%)	72% (54-85%)	1.4 (0.44-3.1)	4.1% (1.1-8.9%)	4.8% (1.2-11%)	9.4 (2.3-21)	70 (18-150)
Sierra Leone	1990	Male	90 to 94	30% (15-48%)	70% (52-85%)	1.3 (0.39-3.2)	3% (0.49-7.4%)	3.5% (0.48-8.7%)	2.4 (0.32-6)	12 (2-31)
Sierra Leone	1990	Male	95 plus	34% (16-56%)	66% (44-84%)	1.3 (0.29-3.7)	2.6% (0.13-7.4%)	3% (-0.077-8.7%)	0.47 (-0.012-1.4)	1.7 (0.078-4.7)
Sierra Leone	1995	Female	All Ages	25% (19-31%)	75% (69-81%)	1.5 (1.1-2.1)	1.4% (1-1.9%)	3.1% (2-4.2%)	830 (530-1100)	29000 (21000-38000)
Sierra Leone	1995	Female	15 to 19	23% (12-38%)	77% (62-88%)	1.4 (0.58-2.8)	1.7% (0.83-3%)	2% (0.71-3.8%)	9.5 (3.3-18)	910 (420-1600)
Sierra Leone	1995	Female	20 to 24	29% (15-46%)	71% (54-85%)	2.2 (0.89-4.2)	3.4% (1.8-5.3%)	3.8% (1.7-6.5%)	19 (8.2-34)	1700 (900-2700)
Sierra Leone	1995	Female	25 to 29	31% (16-49%)	69% (51-84%)	2.6 (1.1-4.8)	4.5% (2.7-6.6%)	5.5% (2.9-8.7%)	26 (13-41)	1900 (1100-2900)
Sierra Leone	1995	Female	30 to 34	34% (19-53%)	66% (47-81%)	3 (1.4-5.3)	5% (3.1-7.2%)	6.5% (3.6-9.7%)	30 (17-46)	2000 (1200-2900)
Sierra Leone	1995	Female	35 to 39	38% (21-56%)	62% (44-79%)	3.2 (1.5-5.6)	6.1% (3.6-8.8%)	8.2% (4.8-12%)	41 (23-62)	2300 (1400-3400)
Sierra Leone	1995	Female	40 to 44	39% (23-58%)	61% (42-77%)	3.4 (1.5-5.9)	7% (4.3-9.8%)	9.8% (5.8-14%)	50 (30-72)	2500 (1500-3500)
Sierra Leone	1995	Female	45 to 49	40% (21-59%)	60% (41-79%)	3.4 (1.5-6.5)	8.4% (5-12%)	12% (6.9-17%)	66 (39-98)	2800 (1700-4200)
Sierra Leone	1995	Female	50 to 54	40% (22-60%)	60% (40-78%)	3.6 (1.6-6.8)	9.6% (5.5-14%)	13% (7.5-19%)	84 (46-120)	3100 (1700-4500)
Sierra Leone	1995	Female	55 to 59	41% (23-59%)	59% (41-77%)	3.9 (1.6-7.2)	10% (5.6-15%)	14% (7.4-20%)	100 (55-150)	3300 (1800-4800)
Sierra Leone	1995	Female	60 to 64	42% (24-61%)	58% (39-76%)	3.9 (1.6-7.8)	9.5% (4.9-14%)	12% (6.2-18%)	100 (53-160)	2900 (1500-4300)
Sierra Leone	1995	Female	65 to 69	43% (25-60%)	57% (40-75%)	3.6 (1.6-6.5)	9.9% (4.6-15%)	12% (5.8-19%)	120 (53-180)	2600 (1200-4100)
Sierra Leone	1995	Female	70 to 74	40% (25-58%)	60% (42-75%)	3 (1.2-5.9)	9% (3.1-16%)	11% (3.7-19%)	98 (32-170)	1800 (590-3100)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	1995	Female	75 to 79	36% (20-56%)	64% (44-80%)	2.4 (0.8-5.3)	7% (1.7-13%)	8.4% (2-16%)	50 (12-95)	710 (170-1300)
Sierra Leone	1995	Female	80 to 84	34% (18-53%)	66% (47-82%)	2.2 (0.74-4.6)	6.1% (1.4-12%)	7.3% (1.7-14%)	16 (3.6-31)	170 (39-320)
Sierra Leone	1995	Female	85 to 89	35% (20-52%)	65% (48-80%)	2.2 (0.77-4.5)	5.6% (1.2-11%)	6.6% (1.3-13%)	11 (2.2-23)	83 (17-170)
Sierra Leone	1995	Female	90 to 94	38% (21-55%)	62% (45-79%)	2.3 (0.74-4.9)	5.2% (0.78-11%)	6.2% (0.79-13%)	4.5 (0.58-10)	23 (3.5-50)
Sierra Leone	1995	Female	95 plus	42% (23-62%)	58% (38-77%)	2.4 (0.68-6)	4.7% (0.18-11%)	5.5% (0.013-13%)	1.2 (0.0028-2.8)	4.2 (0.15-9.5)
Sierra Leone	1995	Male	All Ages	21% (16-26%)	79% (74-84%)	1.9 (1.4-2.6)	2.3% (1.8-2.9%)	4.5% (3.4-5.7%)	1400 (1100-1800)	54000 (43000-67000)
Sierra Leone	1995	Male	15 to 19	16% (8.1-27%)	84% (73-92%)	1.1 (0.42-2.2)	2% (0.92-3.4%)	2.2% (0.86-4.1%)	16 (6.2-30)	1300 (620-2400)
Sierra Leone	1995	Male	20 to 24	24% (13-40%)	76% (60-87%)	2.3 (0.88-4.3)	4.3% (2.4-6.6%)	4.5% (2.3-7.4%)	45 (23-72)	3400 (1900-5200)
Sierra Leone	1995	Male	25 to 29	30% (16-47%)	70% (53-84%)	3.4 (1.3-6.1)	7.2% (4.2-11%)	8.1% (4.6-12%)	65 (37-97)	4400 (2500-6200)
Sierra Leone	1995	Male	30 to 34	33% (17-52%)	67% (48-83%)	4.1 (1.8-7.3)	9.4% (6.2-13%)	11% (7-16%)	81 (51-110)	4900 (3200-6700)
Sierra Leone	1995	Male	35 to 39	36% (20-54%)	64% (46-80%)	4.4 (2-8.1)	11% (7.8-15%)	14% (9.6-19%)	99 (66-140)	5300 (3600-7200)
Sierra Leone	1995	Male	40 to 44	37% (22-56%)	63% (44-78%)	4.7 (2-8.7)	13% (8.8-18%)	16% (11-22%)	120 (76-160)	5600 (3600-7500)
Sierra Leone	1995	Male	45 to 49	38% (20-56%)	62% (44-80%)	5 (2-9.5)	15% (9.9-20%)	18% (12-24%)	160 (100-210)	6500 (4300-8900)
Sierra Leone	1995	Male	50 to 54	39% (22-59%)	61% (41-78%)	5.1 (2.1-10)	14% (9.6-19%)	18% (12-24%)	170 (110-230)	6300 (4200-8600)
Sierra Leone	1995	Male	55 to 59	39% (23-58%)	61% (42-77%)	5 (2-9.5)	14% (9-18%)	17% (11-23%)	170 (110-230)	5500 (3600-7400)
Sierra Leone	1995	Male	60 to 64	39% (21-58%)	61% (42-79%)	4.5 (1.7-9.4)	12% (7-16%)	14% (8.4-19%)	160 (95-220)	4300 (2600-6000)
Sierra Leone	1995	Male	65 to 69	38% (23-56%)	62% (44-77%)	3.8 (1.5-7.4)	11% (5.9-16%)	13% (7-19%)	150 (80-210)	3300 (1800-4800)
Sierra Leone	1995	Male	70 to 74	37% (22-54%)	63% (46-78%)	2.9 (1.1-6)	9.2% (3.7-15%)	11% (4.3-18%)	110 (45-190)	2100 (810-3400)
Sierra Leone	1995	Male	75 to 79	34% (19-54%)	66% (46-81%)	2.2 (0.71-5)	7% (2.1-12%)	8.1% (2.5-15%)	58 (18-110)	820 (250-1500)
Sierra Leone	1995	Male	80 to 84	31% (17-50%)	69% (50-83%)	1.9 (0.61-4.1)	6% (1.9-11%)	7% (2.1-13%)	21 (6.4-40)	220 (69-410)
Sierra Leone	1995	Male	85 to 89	29% (15-46%)	71% (54-85%)	1.6 (0.52-3.5)	5% (1.4-9.9%)	5.9% (1.6-12%)	11 (3-22)	81 (24-160)
Sierra Leone	1995	Male	90 to 94	31% (16-50%)	69% (50-84%)	1.5 (0.47-3.5)	3.9% (0.78-8.5%)	4.5% (0.76-10%)	3 (0.51-6.9)	16 (3.2-35)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	1995	Male	95 plus	36% (17-58%)	64% (42-83%)	1.6 (0.36-4.1)	3.5% (0.36-8.6%)	4% (0.26-10%)	0.66 (0.043-1.7)	2.3 (0.25-5.8)
Sierra Leone	2000	Female	All Ages	25% (19-32%)	75% (68-81%)	1.8 (1.3-2.4)	1.9% (1.4-2.5%)	4.2% (2.9-5.5%)	1100 (750-1500)	38000 (28000-49000)
Sierra Leone	2000	Female	15 to 19	23% (13-38%)	77% (62-87%)	1.6 (0.66-3.1)	2.4% (1.1-4%)	3% (1.2-5.5%)	15 (5.7-27)	1300 (590-2300)
Sierra Leone	2000	Female	20 to 24	29% (15-47%)	71% (53-85%)	2.5 (1.1-4.6)	4.3% (2.5-6.6%)	5.1% (2.5-8.4%)	30 (14-50)	2500 (1400-3800)
Sierra Leone	2000	Female	25 to 29	32% (16-50%)	68% (50-84%)	3 (1.3-5.4)	5.4% (3.2-8%)	6.7% (3.6-11%)	39 (21-61)	2800 (1700-4200)
Sierra Leone	2000	Female	30 to 34	34% (17-53%)	66% (47-83%)	3.4 (1.6-6.1)	5.8% (3.6-8.4%)	7.6% (4.4-11%)	42 (24-62)	2600 (1600-3800)
Sierra Leone	2000	Female	35 to 39	38% (21-57%)	62% (43-79%)	3.7 (1.7-6.3)	7% (4.4-10%)	9.5% (5.6-14%)	54 (32-79)	3000 (1800-4300)
Sierra Leone	2000	Female	40 to 44	40% (23-58%)	60% (42-77%)	3.8 (1.7-6.8)	8% (4.9-11%)	11% (6.7-16%)	64 (39-92)	3100 (1900-4400)
Sierra Leone	2000	Female	45 to 49	40% (22-59%)	60% (41-78%)	3.9 (1.6-7.4)	9.8% (6.1-14%)	13% (8.2-19%)	85 (52-120)	3600 (2200-5200)
Sierra Leone	2000	Female	50 to 54	40% (23-61%)	60% (39-77%)	4.1 (1.8-7.9)	11% (6.6-16%)	15% (8.9-21%)	110 (66-160)	4100 (2500-5800)
Sierra Leone	2000	Female	55 to 59	41% (24-59%)	59% (41-76%)	4.5 (1.9-8.5)	12% (7.1-17%)	16% (9.2-23%)	140 (79-200)	4400 (2500-6300)
Sierra Leone	2000	Female	60 to 64	42% (24-61%)	58% (39-76%)	4.5 (1.8-8.7)	11% (6.3-16%)	14% (7.9-20%)	140 (77-210)	3800 (2100-5600)
Sierra Leone	2000	Female	65 to 69	43% (26-60%)	57% (40-74%)	4.1 (1.8-7.5)	12% (6.4-18%)	14% (7.7-21%)	150 (79-230)	3400 (1800-5100)
Sierra Leone	2000	Female	70 to 74	41% (25-58%)	59% (42-75%)	3.4 (1.3-7)	11% (4.4-18%)	13% (5.1-21%)	120 (48-200)	2200 (860-3700)
Sierra Leone	2000	Female	75 to 79	37% (20-55%)	63% (45-80%)	2.8 (0.9-6.2)	8.4% (2.4-15%)	10% (2.8-18%)	61 (17-110)	850 (240-1500)
Sierra Leone	2000	Female	80 to 84	34% (19-53%)	66% (47-81%)	2.5 (0.85-5.1)	7.5% (2.1-13%)	8.8% (2.4-16%)	18 (5-33)	190 (53-330)
Sierra Leone	2000	Female	85 to 89	35% (19-53%)	65% (47-81%)	2.5 (0.84-5.2)	6.9% (1.8-13%)	8.2% (2-15%)	14 (3.5-26)	100 (27-190)
Sierra Leone	2000	Female	90 to 94	38% (20-55%)	62% (45-80%)	2.6 (0.83-5.7)	6.6% (1.3-13%)	7.9% (1.4-16%)	6.6 (1.2-13)	33 (6.5-66)
Sierra Leone	2000	Female	95 plus	42% (22-62%)	58% (38-78%)	2.7 (0.73-6.7)	6% (0.6-13%)	7% (0.52-15%)	1.9 (0.14-3.9)	6.3 (0.63-13)
Sierra Leone	2000	Male	All Ages	21% (17-27%)	79% (73-83%)	2.2 (1.6-2.9)	2.9% (2.3-3.5%)	5.5% (4.3-6.9%)	1700 (1300-2100)	63000 (50000-79000)
Sierra Leone	2000	Male	15 to 19	17% (8.6-28%)	83% (72-91%)	1.3 (0.49-2.7)	2.9% (1.4-5%)	3.6% (1.5-6.5%)	22 (8.7-38)	1800 (840-3000)
Sierra Leone	2000	Male	20 to 24	25% (13-40%)	75% (60-87%)	2.6 (1.1-4.9)	6.5% (3.8-10%)	7.6% (4-12%)	59 (32-92)	4300 (2500-6600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	2000	Male	25 to 29	30% (17-48%)	70% (52-83%)	3.9 (1.6-7.1)	9.7% (6-14%)	12% (6.8-17%)	85 (49-120)	5600 (3400-8000)
Sierra Leone	2000	Male	30 to 34	34% (18-53%)	66% (47-82%)	4.7 (2.1-8.3)	12% (7.9-16%)	15% (9.6-20%)	100 (65-140)	6000 (4000-8100)
Sierra Leone	2000	Male	35 to 39	37% (20-56%)	63% (44-80%)	5.2 (2.3-9.6)	13% (9.3-18%)	17% (12-23%)	120 (80-160)	6300 (4300-8600)
Sierra Leone	2000	Male	40 to 44	38% (22-58%)	62% (42-78%)	5.5 (2.4-10)	15% (10-20%)	19% (13-25%)	140 (94-190)	6600 (4500-8800)
Sierra Leone	2000	Male	45 to 49	39% (21-58%)	61% (42-79%)	5.8 (2.3-11)	16% (11-22%)	21% (14-27%)	170 (110-220)	6900 (4600-9200)
Sierra Leone	2000	Male	50 to 54	39% (22-59%)	61% (41-78%)	5.9 (2.3-12)	16% (11-22%)	20% (14-27%)	200 (130-270)	7200 (4800-9700)
Sierra Leone	2000	Male	55 to 59	39% (23-59%)	61% (41-77%)	5.7 (2.3-11)	15% (10-21%)	19% (13-25%)	200 (140-280)	6500 (4400-8800)
Sierra Leone	2000	Male	60 to 64	39% (22-58%)	61% (42-78%)	5.2 (1.9-11)	13% (8.3-18%)	16% (9.9-21%)	180 (120-260)	4900 (3200-6900)
Sierra Leone	2000	Male	65 to 69	39% (22-56%)	61% (44-78%)	4.4 (1.8-8.3)	12% (7.4-17%)	15% (8.8-20%)	170 (100-240)	3800 (2300-5300)
Sierra Leone	2000	Male	70 to 74	37% (22-55%)	63% (45-78%)	3.4 (1.3-7.2)	11% (4.8-17%)	13% (5.7-19%)	130 (59-200)	2300 (1100-3500)
Sierra Leone	2000	Male	75 to 79	34% (19-53%)	66% (47-81%)	2.6 (0.83-5.8)	8.3% (2.8-14%)	9.7% (3.3-16%)	65 (22-110)	910 (300-1500)
Sierra Leone	2000	Male	80 to 84	32% (17-50%)	68% (50-83%)	2.2 (0.69-4.8)	7.4% (2.6-13%)	8.6% (3-15%)	22 (7.5-38)	230 (78-390)
Sierra Leone	2000	Male	85 to 89	30% (16-48%)	70% (52-84%)	1.9 (0.59-4.1)	6.4% (2-12%)	7.4% (2.3-14%)	13 (4-24)	96 (31-180)
Sierra Leone	2000	Male	90 to 94	32% (17-50%)	68% (50-83%)	1.9 (0.56-4.3)	5.2% (1.2-11%)	6.1% (1.2-13%)	4.4 (0.89-9.1)	22 (5-46)
Sierra Leone	2000	Male	95 plus	37% (17-59%)	63% (41-83%)	1.9 (0.42-5)	5% (0.85-11%)	5.9% (0.82-13%)	1.1 (0.15-2.3)	3.6 (0.62-8)
Sierra Leone	2005	Female	All Ages	25% (19-32%)	75% (68-81%)	1.8 (1.3-2.5)	2.3% (1.7-2.9%)	4.7% (3.2-6.2%)	1400 (970-1900)	50000 (37000-64000)
Sierra Leone	2005	Female	15 to 19	23% (13-37%)	77% (63-87%)	1.6 (0.63-3.2)	2.5% (1.1-4.3%)	3.3% (1.2-6%)	20 (7.1-36)	1700 (760-3000)
Sierra Leone	2005	Female	20 to 24	29% (15-46%)	71% (54-85%)	2.5 (1.1-4.6)	4.4% (2.5-6.8%)	5.2% (2.4-8.6%)	41 (19-68)	3300 (1800-5100)
Sierra Leone	2005	Female	25 to 29	31% (16-49%)	69% (51-84%)	3.1 (1.4-5.4)	5.2% (3.1-7.7%)	6.3% (3.5-9.7%)	54 (30-83)	3800 (2300-5700)
Sierra Leone	2005	Female	30 to 34	35% (17-52%)	65% (48-83%)	3.5 (1.7-6.1)	5.6% (3.6-8%)	6.9% (4.1-10%)	58 (35-88)	3700 (2300-5400)
Sierra Leone	2005	Female	35 to 39	38% (21-57%)	62% (43-79%)	3.7 (1.8-6.6)	6.8% (4.2-9.7%)	8.8% (5.2-13%)	71 (43-100)	3900 (2500-5600)
Sierra Leone	2005	Female	40 to 44	40% (23-58%)	60% (42-77%)	3.9 (1.7-7.2)	7.8% (4.7-11%)	10% (6.2-15%)	82 (48-120)	4000 (2400-5700)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	2005	Female	45 to 49	40% (22-59%)	60% (41-78%)	4.1 (1.6-7.9)	9.6% (5.7-14%)	13% (7.5-18%)	110 (62-150)	4600 (2700-6600)
Sierra Leone	2005	Female	50 to 54	41% (23-61%)	59% (39-77%)	4.3 (1.7-8.4)	11% (6.5-16%)	15% (8.5-21%)	140 (80-210)	5300 (3000-7600)
Sierra Leone	2005	Female	55 to 59	41% (24-59%)	59% (41-76%)	4.6 (1.9-8.7)	12% (7.5-18%)	16% (9.2-22%)	180 (100-260)	5700 (3300-8200)
Sierra Leone	2005	Female	60 to 64	42% (24-60%)	58% (40-76%)	4.6 (1.8-8.9)	11% (6.3-16%)	14% (7.7-20%)	180 (100-270)	5000 (2800-7200)
Sierra Leone	2005	Female	65 to 69	43% (26-60%)	57% (40-74%)	4.2 (1.8-7.8)	12% (6.1-18%)	14% (7.4-21%)	200 (100-300)	4600 (2300-6700)
Sierra Leone	2005	Female	70 to 74	41% (25-58%)	59% (42-75%)	3.5 (1.4-7.3)	11% (4.4-17%)	13% (5.1-20%)	170 (68-280)	3000 (1200-4900)
Sierra Leone	2005	Female	75 to 79	37% (20-56%)	63% (44-80%)	2.9 (0.93-6.3)	8.6% (2.5-15%)	10% (2.9-18%)	88 (25-150)	1200 (350-2200)
Sierra Leone	2005	Female	80 to 84	34% (19-53%)	66% (47-81%)	2.6 (0.86-5.6)	7.6% (2.2-14%)	8.9% (2.4-16%)	26 (7.3-48)	280 (79-490)
Sierra Leone	2005	Female	85 to 89	35% (19-53%)	65% (47-81%)	2.5 (0.81-5.5)	6.9% (1.8-13%)	8.1% (2.1-15%)	15 (3.9-29)	110 (29-210)
Sierra Leone	2005	Female	90 to 94	38% (20-55%)	62% (45-80%)	2.7 (0.82-5.8)	6.7% (1.4-13%)	8% (1.6-15%)	8 (1.6-15)	41 (8.4-77)
Sierra Leone	2005	Female	95 plus	42% (22-62%)	58% (38-78%)	2.8 (0.72-6.6)	6.1% (0.71-13%)	7.2% (0.58-15%)	2.6 (0.21-5.6)	9 (1-19)
Sierra Leone	2005	Male	All Ages	21% (16-27%)	79% (73-84%)	2.3 (1.6-3.1)	3.2% (2.6-4%)	6.1% (4.7-7.6%)	2100 (1600-2600)	78000 (62000-96000)
Sierra Leone	2005	Male	15 to 19	16% (8.6-28%)	84% (72-91%)	1.3 (0.47-2.8)	3.2% (1.5-5.5%)	4.2% (1.7-7.4%)	26 (11-45)	2100 (1000-3600)
Sierra Leone	2005	Male	20 to 24	24% (13-41%)	76% (59-87%)	2.7 (1.1-5.3)	7.4% (4.4-11%)	9.1% (4.9-14%)	74 (39-120)	5500 (3100-8200)
Sierra Leone	2005	Male	25 to 29	30% (15-47%)	70% (53-85%)	3.9 (1.7-7)	10% (6.4-15%)	13% (7.7-18%)	110 (68-160)	7300 (4700-10000)
Sierra Leone	2005	Male	30 to 34	34% (18-52%)	66% (48-82%)	4.8 (2.3-8.1)	12% (8-16%)	15% (9.7-20%)	130 (85-180)	7800 (5200-11000)
Sierra Leone	2005	Male	35 to 39	37% (20-56%)	63% (44-80%)	5.3 (2.5-9.9)	13% (9-18%)	16% (11-22%)	150 (100-200)	7800 (5400-11000)
Sierra Leone	2005	Male	40 to 44	38% (22-57%)	62% (43-78%)	5.6 (2.4-11)	14% (9.8-19%)	18% (13-24%)	170 (110-230)	8100 (5500-11000)
Sierra Leone	2005	Male	45 to 49	39% (20-57%)	61% (43-80%)	6 (2.3-12)	16% (11-21%)	20% (14-27%)	200 (140-270)	8500 (5900-11000)
Sierra Leone	2005	Male	50 to 54	39% (22-60%)	61% (40-78%)	6.1 (2.3-13)	16% (11-22%)	20% (14-27%)	220 (150-290)	8000 (5300-11000)
Sierra Leone	2005	Male	55 to 59	39% (23-59%)	61% (41-77%)	6 (2.4-12)	16% (11-21%)	19% (13-25%)	250 (170-340)	7800 (5400-11000)
Sierra Leone	2005	Male	60 to 64	39% (23-59%)	61% (41-77%)	5.4 (2.1-11)	13% (8.5-18%)	16% (10-21%)	230 (150-310)	6200 (4000-8500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	2005	Male	65 to 69	39% (22-57%)	61% (43-78%)	4.5 (1.8-9)	12% (7.5-17%)	15% (8.7-20%)	210 (130-300)	4600 (2800-6600)
Sierra Leone	2005	Male	70 to 74	37% (22-55%)	63% (45-78%)	3.5 (1.3-7.2)	11% (5.2-16%)	13% (6-19%)	160 (76-240)	2900 (1400-4300)
Sierra Leone	2005	Male	75 to 79	34% (19-53%)	66% (47-81%)	2.7 (0.83-6.1)	8.2% (2.7-14%)	9.6% (3.2-16%)	80 (27-140)	1100 (380-1900)
Sierra Leone	2005	Male	80 to 84	32% (17-50%)	68% (50-83%)	2.3 (0.71-5)	7.4% (2.5-13%)	8.6% (2.9-15%)	26 (8.9-45)	280 (95-470)
Sierra Leone	2005	Male	85 to 89	30% (16-49%)	70% (51-84%)	2 (0.6-4.4)	6.3% (2-11%)	7.3% (2.2-13%)	13 (3.9-23)	94 (29-170)
Sierra Leone	2005	Male	90 to 94	32% (17-49%)	68% (51-83%)	1.9 (0.57-4.5)	5.2% (1.2-10%)	6.1% (1.4-13%)	4.9 (1.1-9.9)	25 (5.9-50)
Sierra Leone	2005	Male	95 plus	36% (18-57%)	64% (43-82%)	1.9 (0.42-5)	5.1% (0.99-11%)	5.9% (1-13%)	1.3 (0.22-2.9)	4.6 (0.9-9.6)
Sierra Leone	2010	Female	All Ages	25% (19-31%)	75% (69-81%)	1.7 (1.1-2.5)	2.3% (1.5-3%)	4.6% (2.8-6.4%)	1500 (890-2100)	51000 (34000-68000)
Sierra Leone	2010	Female	15 to 19	23% (13-37%)	77% (63-87%)	1.5 (0.56-3.2)	2.3% (1-3.8%)	2.9% (1-5.3%)	18 (6.5-33)	1600 (750-2800)
Sierra Leone	2010	Female	20 to 24	29% (15-46%)	71% (54-85%)	2.4 (0.94-4.6)	4.1% (2.2-6.3%)	4.7% (2.1-7.8%)	38 (17-64)	3200 (1700-5000)
Sierra Leone	2010	Female	25 to 29	31% (16-49%)	69% (51-84%)	2.9 (1.3-5.3)	4.6% (2.7-6.8%)	5.5% (2.9-8.6%)	52 (28-81)	3800 (2200-5600)
Sierra Leone	2010	Female	30 to 34	34% (17-53%)	66% (47-83%)	3.2 (1.5-5.9)	4.8% (3-6.9%)	5.7% (3.3-8.7%)	57 (34-86)	3700 (2300-5400)
Sierra Leone	2010	Female	35 to 39	38% (21-56%)	62% (44-79%)	3.5 (1.6-6.4)	5.9% (3.6-8.4%)	7.4% (4.3-11%)	73 (44-110)	4100 (2500-5900)
Sierra Leone	2010	Female	40 to 44	39% (22-59%)	61% (41-78%)	3.7 (1.5-7.2)	6.9% (4-9.9%)	9% (5.2-13%)	84 (47-120)	4100 (2400-5900)
Sierra Leone	2010	Female	45 to 49	40% (21-58%)	60% (42-79%)	3.8 (1.4-7.6)	8.6% (4.8-12%)	11% (6.2-17%)	110 (58-160)	4600 (2500-6800)
Sierra Leone	2010	Female	50 to 54	40% (22-61%)	60% (39-78%)	4 (1.5-8.2)	10% (5.3-15%)	13% (6.9-19%)	140 (72-220)	5300 (2700-8000)
Sierra Leone	2010	Female	55 to 59	40% (24-59%)	60% (41-76%)	4.3 (1.7-8.7)	11% (6.1-16%)	14% (7.6-21%)	180 (90-270)	5600 (2900-8400)
Sierra Leone	2010	Female	60 to 64	41% (24-61%)	59% (39-76%)	4.3 (1.6-8.7)	10% (5.1-15%)	13% (6.1-19%)	180 (88-280)	5000 (2400-7700)
Sierra Leone	2010	Female	65 to 69	42% (25-60%)	58% (40-75%)	3.9 (1.6-7.6)	11% (4.9-17%)	13% (5.6-20%)	210 (92-320)	4700 (2100-7200)
Sierra Leone	2010	Female	70 to 74	40% (24-58%)	60% (42-76%)	3.2 (1.2-6.9)	9.7% (3.4-16%)	11% (3.8-19%)	170 (59-300)	3100 (1100-5400)
Sierra Leone	2010	Female	75 to 79	36% (20-55%)	64% (45-80%)	2.6 (0.78-5.9)	7.7% (1.6-14%)	8.9% (1.8-17%)	95 (20-180)	1300 (280-2500)
Sierra Leone	2010	Female	80 to 84	34% (17-52%)	66% (48-83%)	2.4 (0.7-5)	6.9% (1.5-13%)	8% (1.7-15%)	36 (6.8-69)	370 (74-710)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	2010	Female	85 to 89	34% (18-52%)	66% (48-82%)	2.4 (0.7-5.2)	6.1% (1.2-12%)	7.1% (1.3-14%)	16 (2.9-32)	120 (23-230)
Sierra Leone	2010	Female	90 to 94	37% (21-54%)	63% (46-79%)	2.5 (0.75-5.6)	5.9% (0.77-12%)	7% (0.8-14%)	6.3 (0.72-13)	32 (4.2-65)
Sierra Leone	2010	Female	95 plus	41% (22-62%)	59% (38-78%)	2.6 (0.68-6.4)	5.5% (0.4-12%)	6.3% (0.27-14%)	2.3 (0.095-5.1)	7.8 (0.57-17)
Sierra Leone	2010	Male	All Ages	21% (16-27%)	79% (73-84%)	2.1 (1.3-3.1)	3.1% (2.4-3.9%)	5.8% (4.2-7.3%)	2000 (1400-2500)	76000 (58000-95000)
Sierra Leone	2010	Male	15 to 19	16% (8.3-28%)	84% (72-92%)	1.2 (0.43-2.6)	2.9% (1.3-4.9%)	3.7% (1.4-6.8%)	23 (8.2-41)	1900 (880-3400)
Sierra Leone	2010	Male	20 to 24	24% (13-41%)	76% (59-87%)	2.5 (0.98-4.8)	6.9% (3.8-11%)	8.2% (4.2-13%)	69 (35-110)	5200 (2900-8000)
Sierra Leone	2010	Male	25 to 29	30% (16-47%)	70% (53-84%)	3.7 (1.6-6.8)	9.4% (5.7-13%)	12% (6.8-17%)	110 (65-160)	7300 (4600-10000)
Sierra Leone	2010	Male	30 to 34	34% (18-52%)	66% (48-82%)	4.5 (2-8.2)	11% (7.1-15%)	13% (8.4-18%)	130 (84-190)	8100 (5200-11000)
Sierra Leone	2010	Male	35 to 39	37% (20-56%)	63% (44-80%)	4.9 (2.2-9.6)	12% (8-16%)	14% (9.7-19%)	150 (100-210)	8100 (5600-11000)
Sierra Leone	2010	Male	40 to 44	38% (23-57%)	62% (43-77%)	5.3 (2.1-11)	13% (8.7-17%)	16% (11-21%)	160 (110-220)	7800 (5300-11000)
Sierra Leone	2010	Male	45 to 49	39% (21-57%)	61% (43-79%)	5.6 (2.1-12)	15% (9.4-19%)	18% (12-24%)	190 (120-260)	8200 (5300-11000)
Sierra Leone	2010	Male	50 to 54	39% (22-60%)	61% (40-78%)	5.7 (2.1-12)	15% (9.5-20%)	18% (12-24%)	210 (130-280)	7600 (4800-10000)
Sierra Leone	2010	Male	55 to 59	39% (23-59%)	61% (41-77%)	5.5 (2.1-11)	14% (9.3-19%)	17% (11-23%)	210 (140-290)	6800 (4400-9300)
Sierra Leone	2010	Male	60 to 64	39% (22-59%)	61% (41-78%)	5 (1.8-11)	12% (7.1-17%)	14% (8.5-20%)	210 (130-300)	5800 (3400-8100)
Sierra Leone	2010	Male	65 to 69	39% (23-57%)	61% (43-77%)	4.2 (1.6-8.8)	11% (5.9-16%)	13% (7.1-19%)	200 (110-290)	4500 (2500-6500)
Sierra Leone	2010	Male	70 to 74	37% (23-55%)	63% (45-77%)	3.2 (1.1-7.2)	9.5% (3.8-15%)	11% (4.4-18%)	160 (63-250)	2800 (1100-4600)
Sierra Leone	2010	Male	75 to 79	34% (19-53%)	66% (47-81%)	2.5 (0.73-6)	7.2% (1.9-13%)	8.4% (2.2-15%)	82 (21-140)	1100 (300-2000)
Sierra Leone	2010	Male	80 to 84	31% (16-50%)	69% (50-84%)	2.1 (0.61-4.8)	6.3% (1.7-12%)	7.4% (2-13%)	30 (7.9-55)	310 (83-570)
Sierra Leone	2010	Male	85 to 89	30% (16-48%)	70% (52-84%)	1.8 (0.53-4.2)	5.3% (1.4-10%)	6.1% (1.6-12%)	12 (3.1-24)	92 (24-180)
Sierra Leone	2010	Male	90 to 94	32% (17-49%)	68% (51-83%)	1.7 (0.46-4.3)	4.3% (0.83-9.3%)	5.1% (0.84-11%)	3.8 (0.64-8.4)	20 (3.7-43)
Sierra Leone	2010	Male	95 plus	36% (18-58%)	64% (42-82%)	1.7 (0.38-4.6)	4.2% (0.55-9.7%)	4.9% (0.46-11%)	1.1 (0.11-2.7)	3.9 (0.51-9.2)
Sierra Leone	2016	Female	All Ages	25% (19-31%)	75% (69-81%)	1.5 (0.78-2.6)	2.2% (1.2-3.1%)	4.5% (2.1-6.6%)	1300 (600-2000)	46000 (26000-64000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	2016	Female	15 to 19	23% (13-37%)	77% (63-87%)	1.3 (0.42-3)	1.9% (0.81-3.3%)	2.3% (0.67-4.5%)	14 (4.2-29)	1400 (610-2600)
Sierra Leone	2016	Female	20 to 24	28% (15-45%)	72% (55-85%)	2.1 (0.72-4.5)	3.6% (1.8-5.6%)	4% (1.5-7%)	30 (11-52)	2800 (1400-4400)
Sierra Leone	2016	Female	25 to 29	31% (16-49%)	69% (51-84%)	2.5 (0.9-5.1)	4.2% (2.3-6.3%)	5% (2.3-8.2%)	41 (18-67)	3300 (1700-4900)
Sierra Leone	2016	Female	30 to 34	34% (17-53%)	66% (47-83%)	2.8 (1.1-5.6)	4.4% (2.5-6.7%)	5.4% (2.7-8.4%)	49 (25-79)	3400 (1900-5200)
Sierra Leone	2016	Female	35 to 39	38% (20-57%)	62% (43-80%)	3.1 (1.2-6.4)	5.3% (3-7.9%)	6.9% (3.7-10%)	67 (35-100)	3800 (2100-5600)
Sierra Leone	2016	Female	40 to 44	39% (23-59%)	61% (41-77%)	3.2 (1.2-7)	6.2% (3.3-9.2%)	8.4% (4.4-12%)	79 (41-120)	4000 (2100-5900)
Sierra Leone	2016	Female	45 to 49	40% (22-59%)	60% (41-78%)	3.3 (1.2-7.5)	7.7% (3.6-12%)	10% (4.8-16%)	98 (44-160)	4200 (2000-6700)
Sierra Leone	2016	Female	50 to 54	40% (23-61%)	60% (39-77%)	3.5 (1.2-8.1)	9% (3.8-14%)	12% (5.2-18%)	130 (52-200)	4700 (1900-7400)
Sierra Leone	2016	Female	55 to 59	41% (24-59%)	59% (41-76%)	3.9 (1.4-8.4)	9.8% (4.3-15%)	13% (5.5-19%)	160 (64-250)	5000 (2100-7900)
Sierra Leone	2016	Female	60 to 64	42% (25-61%)	58% (39-75%)	3.9 (1.3-8.5)	8.9% (3.7-14%)	11% (4.4-18%)	160 (63-260)	4400 (1700-7100)
Sierra Leone	2016	Female	65 to 69	43% (25-61%)	57% (39-75%)	3.5 (1.3-7.6)	9.2% (3-15%)	11% (3.6-18%)	180 (58-310)	4100 (1300-6900)
Sierra Leone	2016	Female	70 to 74	41% (25-59%)	59% (41-75%)	2.9 (0.94-6.5)	8% (1.6-15%)	9.4% (1.8-18%)	160 (28-290)	2800 (530-5300)
Sierra Leone	2016	Female	75 to 79	36% (21-55%)	64% (45-79%)	2.4 (0.63-5.5)	6.3% (0.59-13%)	7.3% (0.69-15%)	92 (8.1-190)	1300 (120-2700)
Sierra Leone	2016	Female	80 to 84	34% (19-53%)	66% (47-81%)	2.1 (0.56-4.9)	5.6% (0.5-12%)	6.5% (0.55-14%)	37 (2.9-81)	390 (34-840)
Sierra Leone	2016	Female	85 to 89	34% (19-53%)	66% (47-81%)	2.1 (0.57-5.1)	5% (0.26-11%)	5.8% (0.24-13%)	18 (0.72-39)	130 (7-290)
Sierra Leone	2016	Female	90 to 94	37% (21-55%)	63% (45-79%)	2.3 (0.57-5.6)	4.7% (0.071-11%)	5.5% (-0.05-13%)	5.4 (-0.05-13)	28 (0.41-64)
Sierra Leone	2016	Female	95 plus	41% (22-62%)	59% (38-78%)	2.4 (0.53-6.3)	4.3% (-0.2-10%)	4.9% (-0.42-12%)	1.3 (-0.11-3.2)	4.5 (-0.21-11)
Sierra Leone	2016	Male	All Ages	21% (16-28%)	79% (72-84%)	1.9 (0.99-3.2)	3.1% (2.2-4%)	5.5% (3.7-7.3%)	1800 (1200-2400)	70000 (49000-91000)
Sierra Leone	2016	Male	15 to 19	17% (8.5-28%)	83% (72-92%)	1.1 (0.36-2.6)	2.5% (1.1-4.4%)	3% (1.1-5.8%)	19 (6.8-37)	1800 (800-3100)
Sierra Leone	2016	Male	20 to 24	24% (13-41%)	76% (59-87%)	2.2 (0.71-4.9)	5.9% (2.9-9.1%)	6.9% (2.9-11%)	56 (24-95)	4500 (2200-7100)
Sierra Leone	2016	Male	25 to 29	30% (16-47%)	70% (53-84%)	3.2 (1.1-6.5)	8.3% (4.9-12%)	10% (5.5-15%)	94 (49-140)	6500 (3700-9600)
Sierra Leone	2016	Male	30 to 34	34% (18-52%)	66% (48-82%)	4 (1.5-7.8)	9.5% (5.8-13%)	12% (6.8-17%)	120 (71-180)	7600 (4600-11000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sierra Leone	2016	Male	35 to 39	37% (20-55%)	63% (45-80%)	4.4 (1.7-9.2)	10% (6.5-14%)	13% (7.8-17%)	140 (89-200)	7800 (4900-11000)
Sierra Leone	2016	Male	40 to 44	38% (22-57%)	62% (43-78%)	4.7 (1.6-10)	11% (7.2-15%)	14% (8.9-19%)	160 (97-220)	7600 (4700-11000)
Sierra Leone	2016	Male	45 to 49	39% (20-58%)	61% (42-80%)	5 (1.6-12)	13% (7.8-18%)	16% (9.8-22%)	180 (110-260)	7700 (4600-11000)
Sierra Leone	2016	Male	50 to 54	39% (21-60%)	61% (40-79%)	5.1 (1.6-12)	12% (7.3-17%)	16% (8.9-21%)	180 (110-260)	6800 (4000-9600)
Sierra Leone	2016	Male	55 to 59	39% (23-58%)	61% (42-77%)	4.9 (1.7-11)	12% (7.1-17%)	15% (8.7-21%)	190 (110-270)	6200 (3500-8800)
Sierra Leone	2016	Male	60 to 64	39% (22-58%)	61% (42-78%)	4.4 (1.4-10)	10% (5.3-15%)	12% (6.3-18%)	170 (89-260)	4800 (2500-7100)
Sierra Leone	2016	Male	65 to 69	39% (22-57%)	61% (43-78%)	3.7 (1.3-8.5)	9.1% (4.2-14%)	11% (5-16%)	180 (80-270)	4000 (1800-6000)
Sierra Leone	2016	Male	70 to 74	37% (22-55%)	63% (45-78%)	2.9 (0.89-6.7)	7.8% (2.5-13%)	9.3% (3-16%)	150 (47-250)	2700 (860-4600)
Sierra Leone	2016	Male	75 to 79	34% (19-53%)	66% (47-81%)	2.3 (0.55-5.6)	5.8% (1.2-11%)	6.8% (1.4-13%)	81 (16-160)	1100 (230-2200)
Sierra Leone	2016	Male	80 to 84	31% (16-50%)	69% (50-84%)	1.9 (0.47-4.6)	5.1% (1.1-10%)	5.9% (1.2-12%)	30 (5.9-60)	310 (66-630)
Sierra Leone	2016	Male	85 to 89	30% (16-48%)	70% (52-84%)	1.7 (0.44-4.2)	4.3% (0.8-9.4%)	5% (0.86-11%)	14 (2.4-30)	100 (19-230)
Sierra Leone	2016	Male	90 to 94	32% (17-51%)	68% (49-83%)	1.6 (0.39-4.2)	3.5% (0.38-8.3%)	4% (0.3-9.8%)	3.8 (0.28-9.2)	20 (2.1-48)
Sierra Leone	2016	Male	95 plus	37% (18-59%)	63% (41-82%)	1.6 (0.32-4.5)	3.3% (0.24-8.4%)	3.7% (0.082-9.9%)	0.82 (0.018-2.2)	2.9 (0.21-7.3)
Togo	1990	Female	All Ages	14% (10-19%)	86% (81-90%)	0.82 (0.61-1.1)	1.5% (1.1-2%)	3.2% (2.1-4.5%)	640 (400-900)	21000 (16000-28000)
Togo	1990	Female	15 to 19	13% (6.2-23%)	87% (77-94%)	0.73 (0.25-1.5)	1.7% (0.8-3%)	2% (0.62-4.1%)	7.6 (2.3-15)	770 (350-1300)
Togo	1990	Female	20 to 24	17% (7.4-30%)	83% (70-93%)	1.2 (0.45-2.3)	3% (1.7-5%)	3.3% (1.4-6%)	14 (6-27)	1400 (750-2200)
Togo	1990	Female	25 to 29	19% (8.7-33%)	81% (67-91%)	1.4 (0.56-2.7)	3.7% (2.1-5.7%)	4.5% (2.1-7.5%)	18 (9-31)	1500 (870-2300)
Togo	1990	Female	30 to 34	21% (9.7-37%)	79% (63-90%)	1.7 (0.71-3.1)	4.5% (2.7-6.9%)	6% (3.2-9.7%)	24 (13-39)	1600 (970-2500)
Togo	1990	Female	35 to 39	24% (11-41%)	76% (59-89%)	1.8 (0.75-3.3)	4.7% (2.8-7.1%)	6.3% (3.5-9.8%)	27 (15-41)	1500 (950-2300)
Togo	1990	Female	40 to 44	25% (13-42%)	75% (58-87%)	1.9 (0.78-3.6)	5.6% (3.3-8.2%)	7.8% (4.5-12%)	34 (20-51)	1700 (1000-2500)
Togo	1990	Female	45 to 49	26% (12-43%)	74% (57-88%)	2 (0.77-3.9)	6.5% (3.7-9.7%)	9.1% (5.1-14%)	44 (24-67)	1900 (1100-2900)
Togo	1990	Female	50 to 54	26% (12-45%)	74% (55-88%)	2 (0.78-4)	7.1% (3.9-11%)	9.5% (5.3-14%)	55 (30-86)	2100 (1200-3200)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	1990	Female	55 to 59	27% (14-46%)	73% (54-86%)	2.3 (0.82-4.5)	7.5% (3.8-12%)	9.7% (4.9-15%)	67 (34-110)	2200 (1100-3400)
Togo	1990	Female	60 to 64	28% (14-47%)	72% (53-86%)	2.3 (0.8-4.9)	6.9% (3.5-11%)	8.5% (4.4-14%)	74 (37-120)	2000 (1000-3200)
Togo	1990	Female	65 to 69	29% (15-46%)	71% (54-85%)	2.1 (0.8-4.3)	7.2% (3.4-12%)	8.7% (4.1-14%)	90 (42-150)	2000 (970-3400)
Togo	1990	Female	70 to 74	27% (14-44%)	73% (56-86%)	1.7 (0.56-3.8)	6.5% (2.6-12%)	7.7% (3.1-14%)	87 (34-160)	1600 (620-2900)
Togo	1990	Female	75 to 79	23% (11-41%)	77% (59-89%)	1.4 (0.37-3.3)	4.9% (1.5-9.6%)	5.7% (1.8-11%)	57 (17-110)	800 (240-1600)
Togo	1990	Female	80 to 84	21% (9.6-38%)	79% (62-90%)	1.2 (0.33-2.8)	4.1% (1.2-8.3%)	4.8% (1.3-9.9%)	27 (7.3-54)	280 (77-560)
Togo	1990	Female	85 to 89	22% (10-37%)	78% (63-90%)	1.2 (0.34-2.8)	3.7% (0.83-7.7%)	4.3% (0.93-9.1%)	13 (2.9-28)	97 (22-200)
Togo	1990	Female	90 to 94	25% (11-43%)	75% (57-89%)	1.3 (0.36-3.1)	3.4% (0.58-7.2%)	4% (0.63-8.6%)	4.4 (0.71-9.8)	23 (3.8-50)
Togo	1990	Female	95 plus	30% (12-53%)	70% (47-88%)	1.6 (0.36-3.9)	3.3% (0.26-7.6%)	3.8% (0.21-9%)	1.2 (0.065-2.8)	4.1 (0.32-9.6)
Togo	1990	Male	All Ages	11% (8.1-15%)	89% (85-92%)	0.94 (0.7-1.2)	1.9% (1.5-2.4%)	3.6% (2.8-4.6%)	760 (580-960)	30000 (24000-36000)
Togo	1990	Male	15 to 19	8.4% (4-16%)	92% (84-96%)	0.54 (0.19-1.2)	1.8% (0.9-3.1%)	2.1% (0.82-4%)	7.6 (2.9-15)	750 (380-1300)
Togo	1990	Male	20 to 24	13% (6-23%)	87% (77-94%)	1.1 (0.42-2.2)	4.3% (2.4-6.8%)	4.8% (2.3-8.2%)	23 (11-39)	1900 (1100-3000)
Togo	1990	Male	25 to 29	17% (7.8-29%)	83% (71-92%)	1.7 (0.66-3.3)	6.5% (3.9-9.7%)	7.7% (4.2-12%)	37 (20-59)	2700 (1600-4000)
Togo	1990	Male	30 to 34	20% (8.8-36%)	80% (64-91%)	2.2 (0.93-4)	8% (5.1-12%)	9.9% (5.9-15%)	47 (28-70)	3000 (1900-4300)
Togo	1990	Male	35 to 39	22% (10-38%)	78% (62-90%)	2.5 (0.97-4.6)	9.1% (6.1-13%)	12% (7.5-16%)	58 (37-82)	3200 (2100-4400)
Togo	1990	Male	40 to 44	23% (12-40%)	77% (60-88%)	2.6 (1.1-5.1)	10% (6.7-14%)	13% (8.5-18%)	68 (44-95)	3300 (2200-4600)
Togo	1990	Male	45 to 49	24% (11-40%)	76% (60-89%)	2.8 (1.1-5.5)	11% (7.1-15%)	14% (9-19%)	80 (52-110)	3400 (2200-4700)
Togo	1990	Male	50 to 54	24% (11-43%)	76% (57-89%)	2.8 (1.1-5.7)	10% (6.5-14%)	13% (8.1-18%)	80 (51-110)	3000 (1900-4200)
Togo	1990	Male	55 to 59	24% (12-42%)	76% (58-88%)	2.7 (0.97-5.6)	9.4% (5.9-13%)	12% (7.4-16%)	82 (51-120)	2700 (1700-3800)
Togo	1990	Male	60 to 64	24% (12-42%)	76% (58-88%)	2.4 (0.79-5.2)	7.8% (4.6-11%)	9.4% (5.4-14%)	76 (43-120)	2100 (1200-3100)
Togo	1990	Male	65 to 69	24% (12-39%)	76% (61-88%)	2 (0.7-4.1)	6.9% (3.7-11%)	8.3% (4.5-13%)	73 (38-110)	1700 (890-2600)
Togo	1990	Male	70 to 74	22% (12-38%)	78% (62-88%)	1.5 (0.48-3.6)	5.8% (2.4-10%)	6.8% (2.9-12%)	62 (27-110)	1100 (490-2000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	1990	Male	75 to 79	20% (9.7-35%)	80% (65-90%)	1.1 (0.29-2.8)	4.3% (1.4-8.2%)	5% (1.7-9.6%)	37 (12-71)	520 (170-990)
Togo	1990	Male	80 to 84	18% (8.3-33%)	82% (67-92%)	0.92 (0.25-2.2)	3.6% (1.3-7%)	4.2% (1.5-8%)	16 (5.6-30)	170 (59-320)
Togo	1990	Male	85 to 89	17% (7.9-30%)	83% (70-92%)	0.79 (0.22-1.9)	2.9% (0.95-6%)	3.4% (1-7%)	7.1 (2.2-14)	53 (17-110)
Togo	1990	Male	90 to 94	18% (8.6-33%)	82% (67-91%)	0.75 (0.18-1.9)	2.2% (0.54-5%)	2.5% (0.55-5.9%)	2 (0.44-4.7)	10 (2.6-23)
Togo	1990	Male	95 plus	22% (8.4-42%)	78% (58-92%)	0.79 (0.15-2.2)	2% (0.33-5.3%)	2.2% (0.22-6.2%)	0.46 (0.045-1.3)	1.7 (0.26-4.4)
Togo	1995	Female	All Ages	14% (10-19%)	86% (81-90%)	0.63 (0.47-0.81)	1.4% (0.95-1.8%)	2.7% (1.6-4.1%)	590 (330-870)	20000 (14000-27000)
Togo	1995	Female	15 to 19	13% (6.5-23%)	87% (77-94%)	0.55 (0.21-1.1)	1.4% (0.66-2.6%)	1.6% (0.51-3.5%)	6.9 (2.2-15)	740 (350-1400)
Togo	1995	Female	20 to 24	17% (7.7-30%)	83% (70-92%)	0.88 (0.34-1.7)	2.4% (1.3-4%)	2.5% (0.94-4.6%)	14 (5.4-26)	1400 (760-2300)
Togo	1995	Female	25 to 29	19% (8.3-34%)	81% (66-92%)	1.1 (0.44-2)	2.8% (1.6-4.4%)	3.1% (1.3-5.4%)	18 (8-32)	1500 (850-2400)
Togo	1995	Female	30 to 34	21% (9.2-37%)	79% (63-91%)	1.3 (0.54-2.3)	3.4% (2-5.5%)	4.2% (2.1-7.2%)	24 (12-40)	1600 (930-2600)
Togo	1995	Female	35 to 39	24% (11-41%)	76% (59-89%)	1.4 (0.56-2.6)	3.6% (2.1-5.8%)	4.6% (2.3-7.8%)	25 (13-42)	1500 (860-2400)
Togo	1995	Female	40 to 44	25% (13-43%)	75% (57-87%)	1.5 (0.59-2.8)	4.4% (2.5-6.8%)	6% (3.2-9.4%)	32 (18-51)	1600 (950-2500)
Togo	1995	Female	45 to 49	26% (12-42%)	74% (58-88%)	1.5 (0.61-2.8)	5.2% (2.7-8.3%)	7.1% (3.5-11%)	41 (20-67)	1800 (940-2900)
Togo	1995	Female	50 to 54	26% (12-46%)	74% (54-88%)	1.5 (0.61-3)	5.8% (3-9.4%)	7.7% (4-13%)	52 (26-85)	2000 (1000-3200)
Togo	1995	Female	55 to 59	27% (13-45%)	73% (55-87%)	1.7 (0.63-3.5)	6.3% (2.9-10%)	8.1% (3.8-13%)	64 (29-110)	2100 (960-3400)
Togo	1995	Female	60 to 64	28% (14-47%)	72% (53-86%)	1.8 (0.66-3.7)	5.9% (2.7-9.8%)	7.3% (3.4-12%)	70 (29-120)	1900 (820-3200)
Togo	1995	Female	65 to 69	29% (15-47%)	71% (53-85%)	1.6 (0.58-3.3)	5.9% (2.3-10%)	7.1% (2.9-13%)	80 (31-140)	1800 (720-3300)
Togo	1995	Female	70 to 74	27% (14-44%)	73% (56-86%)	1.3 (0.44-2.8)	5.1% (1.7-9.8%)	6% (2-12%)	74 (25-150)	1300 (450-2700)
Togo	1995	Female	75 to 79	23% (11-40%)	77% (60-89%)	1 (0.28-2.5)	3.7% (0.95-8.1%)	4.4% (1.1-9.8%)	47 (11-110)	660 (170-1500)
Togo	1995	Female	80 to 84	22% (10-38%)	78% (62-90%)	0.9 (0.25-2.1)	3.1% (0.67-7%)	3.6% (0.75-8.4%)	23 (4.8-53)	240 (53-550)
Togo	1995	Female	85 to 89	22% (11-38%)	78% (62-89%)	0.91 (0.27-2.1)	2.7% (0.49-6.6%)	3.2% (0.53-7.9%)	12 (1.9-30)	90 (15-220)
Togo	1995	Female	90 to 94	25% (12-43%)	75% (57-88%)	0.99 (0.27-2.4)	2.5% (0.2-6.6%)	2.9% (0.18-7.7%)	4 (0.24-10)	20 (1.6-52)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	1995	Female	95 plus	29% (12-53%)	71% (47-88%)	1.1 (0.26-2.9)	2.4% (0.0086-6.6%)	2.8% (-0.098-7.7%)	0.99 (-0.034-2.8)	3.4 (0.012-9.2)
Togo	1995	Male	All Ages	11% (8.1-15%)	89% (85-92%)	0.72 (0.54-0.92)	1.9% (1.5-2.4%)	3.4% (2.5-4.5%)	810 (600-1000)	32000 (25000-39000)
Togo	1995	Male	15 to 19	8.7% (4.1-16%)	91% (84-96%)	0.43 (0.15-0.92)	1.6% (0.79-2.7%)	1.8% (0.65-3.6%)	8.1 (3-16)	810 (400-1400)
Togo	1995	Male	20 to 24	13% (6.2-24%)	87% (76-94%)	0.9 (0.34-1.8)	3.8% (2-6%)	4.1% (1.8-7%)	26 (11-45)	2100 (1100-3400)
Togo	1995	Male	25 to 29	17% (7.5-31%)	83% (69-92%)	1.4 (0.5-2.5)	5.4% (3-8.5%)	6% (3-10%)	40 (20-67)	2900 (1600-4500)
Togo	1995	Male	30 to 34	20% (8.7-36%)	80% (64-91%)	1.7 (0.75-3.1)	6.5% (3.9-9.7%)	7.6% (4.2-12%)	51 (30-79)	3200 (2000-4800)
Togo	1995	Male	35 to 39	22% (11-38%)	78% (62-89%)	1.9 (0.74-3.4)	7.4% (4.6-11%)	8.9% (5.3-13%)	61 (37-92)	3400 (2100-4900)
Togo	1995	Male	40 to 44	23% (12-40%)	77% (60-88%)	2 (0.81-3.9)	8.3% (5.2-12%)	10% (6.2-15%)	73 (42-110)	3500 (2100-5000)
Togo	1995	Male	45 to 49	24% (11-40%)	76% (60-89%)	2.1 (0.8-4.1)	9.4% (5.8-13%)	12% (7.1-16%)	87 (51-120)	3700 (2200-5300)
Togo	1995	Male	50 to 54	24% (11-43%)	76% (57-89%)	2.2 (0.82-4.4)	9% (5.4-13%)	11% (6.6-16%)	89 (53-130)	3300 (2000-4900)
Togo	1995	Male	55 to 59	25% (13-42%)	75% (58-87%)	2.1 (0.76-4.3)	8.6% (4.9-13%)	10% (5.9-15%)	91 (51-140)	2900 (1700-4300)
Togo	1995	Male	60 to 64	24% (12-42%)	76% (58-88%)	1.9 (0.66-4.1)	7% (3.6-11%)	8.3% (4.1-13%)	82 (42-130)	2300 (1200-3500)
Togo	1995	Male	65 to 69	24% (12-40%)	76% (60-88%)	1.6 (0.54-3.2)	6% (2.8-9.9%)	7.1% (3.3-12%)	75 (35-120)	1700 (790-2800)
Togo	1995	Male	70 to 74	23% (12-39%)	77% (61-88%)	1.2 (0.38-2.8)	4.8% (1.8-9.1%)	5.6% (2.2-11%)	61 (24-120)	1100 (430-2100)
Togo	1995	Male	75 to 79	20% (9.9-37%)	80% (63-90%)	0.89 (0.24-2.3)	3.5% (1.1-7.3%)	4.1% (1.2-8.5%)	36 (11-76)	510 (150-1100)
Togo	1995	Male	80 to 84	19% (8.3-34%)	81% (66-92%)	0.73 (0.2-1.8)	2.9% (0.98-6.2%)	3.4% (1.1-7%)	16 (5.1-32)	160 (55-340)
Togo	1995	Male	85 to 89	18% (8.1-31%)	82% (69-92%)	0.63 (0.17-1.6)	2.4% (0.71-5.4%)	2.7% (0.79-6.2%)	7.2 (2.1-16)	54 (16-120)
Togo	1995	Male	90 to 94	19% (8.6-34%)	81% (66-91%)	0.61 (0.14-1.6)	1.8% (0.37-4.6%)	2% (0.34-5.4%)	1.9 (0.33-5.2)	10 (2.1-26)
Togo	1995	Male	95 plus	23% (9.1-43%)	77% (57-91%)	0.64 (0.12-1.8)	1.7% (0.21-4.8%)	1.9% (0.11-5.7%)	0.45 (0.026-1.3)	1.6 (0.2-4.5)
Togo	2000	Female	All Ages	14% (10-19%)	86% (81-90%)	0.53 (0.4-0.69)	1.2% (0.78-1.7%)	2.3% (1.2-3.7%)	550 (280-910)	20000 (13000-28000)
Togo	2000	Female	15 to 19	13% (6.3-24%)	87% (76-94%)	0.45 (0.16-0.93)	1.2% (0.58-2.2%)	1.3% (0.35-2.8%)	5.9 (1.7-13)	700 (340-1300)
Togo	2000	Female	20 to 24	17% (7.7-30%)	83% (70-92%)	0.71 (0.28-1.4)	1.9% (1-3.3%)	1.8% (0.61-3.5%)	13 (4.4-25)	1400 (750-2300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	2000	Female	25 to 29	19% (8.8-34%)	81% (66-91%)	0.88 (0.37-1.6)	1.9% (1-3.1%)	1.8% (0.68-3.4%)	17 (7.1-32)	1500 (850-2500)
Togo	2000	Female	30 to 34	22% (9.6-38%)	78% (62-90%)	1 (0.44-1.9)	2.2% (1.2-3.7%)	2.3% (1.1-4.4%)	23 (11-41)	1600 (910-2700)
Togo	2000	Female	35 to 39	24% (11-42%)	76% (58-89%)	1.1 (0.46-2.1)	2.4% (1.3-4.1%)	2.7% (1.3-5%)	25 (12-43)	1500 (850-2500)
Togo	2000	Female	40 to 44	26% (13-44%)	74% (56-87%)	1.2 (0.49-2.3)	3.1% (1.7-5.2%)	3.8% (1.9-6.7%)	32 (16-55)	1600 (870-2700)
Togo	2000	Female	45 to 49	26% (12-44%)	74% (56-88%)	1.2 (0.48-2.4)	3.9% (1.9-6.5%)	5% (2.3-8.4%)	40 (19-69)	1800 (900-3000)
Togo	2000	Female	50 to 54	27% (12-47%)	73% (53-88%)	1.3 (0.52-2.5)	4.7% (2.3-7.9%)	6% (2.9-10%)	52 (23-90)	2000 (930-3300)
Togo	2000	Female	55 to 59	28% (14-47%)	72% (53-86%)	1.4 (0.53-3)	5.3% (2.3-9.5%)	6.8% (2.9-12%)	65 (26-120)	2100 (850-3800)
Togo	2000	Female	60 to 64	29% (14-48%)	71% (52-86%)	1.5 (0.54-3.1)	5% (2-9%)	6.2% (2.5-11%)	69 (25-130)	1900 (700-3500)
Togo	2000	Female	65 to 69	29% (15-47%)	71% (53-85%)	1.3 (0.47-2.7)	4.9% (1.7-9.3%)	5.9% (2.1-11%)	74 (24-150)	1700 (550-3300)
Togo	2000	Female	70 to 74	27% (14-45%)	73% (55-86%)	1.1 (0.35-2.3)	4.1% (1.1-8.7%)	4.8% (1.3-10%)	63 (17-140)	1200 (310-2500)
Togo	2000	Female	75 to 79	24% (12-41%)	76% (59-88%)	0.83 (0.25-2)	3% (0.57-7.3%)	3.5% (0.64-8.7%)	40 (7.3-99)	560 (110-1400)
Togo	2000	Female	80 to 84	22% (10-38%)	78% (62-90%)	0.74 (0.2-1.6)	2.4% (0.4-6.1%)	2.9% (0.47-7.3%)	19 (3.1-49)	200 (33-510)
Togo	2000	Female	85 to 89	22% (10-38%)	78% (62-90%)	0.74 (0.22-1.7)	2.1% (0.26-5.7%)	2.3% (0.26-6.8%)	11 (1.1-30)	79 (9.1-220)
Togo	2000	Female	90 to 94	25% (12-43%)	75% (57-88%)	0.82 (0.22-2)	2% (0.041-5.9%)	2.3% (-0.033-7%)	3.9 (-0.055-12)	20 (0.41-60)
Togo	2000	Female	95 plus	30% (13-53%)	70% (47-87%)	0.94 (0.22-2.5)	2% (-0.22-6.3%)	2.3% (-0.39-7.3%)	0.96 (-0.17-3.2)	3.4 (-0.36-11)
Togo	2000	Male	All Ages	12% (8.4-16%)	88% (84-92%)	0.62 (0.47-0.8)	1.9% (1.5-2.5%)	3.2% (2.3-4.3%)	870 (620-1200)	35000 (27000-45000)
Togo	2000	Male	15 to 19	9% (4.3-16%)	91% (84-96%)	0.37 (0.13-0.77)	1.4% (0.73-2.6%)	1.6% (0.55-3.2%)	8.8 (3.1-18)	900 (440-1600)
Togo	2000	Male	20 to 24	14% (6.3-25%)	86% (75-94%)	0.75 (0.29-1.5)	3.4% (1.8-5.6%)	3.5% (1.5-6.3%)	30 (13-54)	2500 (1300-4300)
Togo	2000	Male	25 to 29	18% (8.3-32%)	82% (68-92%)	1.1 (0.44-2.1)	4.4% (2.3-7%)	4.7% (2.1-8%)	48 (22-82)	3400 (1800-5500)
Togo	2000	Male	30 to 34	21% (9.5-38%)	79% (62-91%)	1.4 (0.62-2.6)	5.1% (3-7.9%)	5.6% (3.1-9%)	60 (32-96)	3800 (2200-5800)
Togo	2000	Male	35 to 39	23% (11-40%)	77% (60-89%)	1.6 (0.64-3)	5.8% (3.5-8.7%)	6.6% (3.8-10%)	69 (39-100)	3800 (2200-5600)
Togo	2000	Male	40 to 44	24% (12-42%)	76% (58-88%)	1.7 (0.73-3.3)	6.7% (3.9-10%)	7.7% (4.5-12%)	82 (47-120)	3900 (2300-5800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	2000	Male	45 to 49	25% (12-41%)	75% (59-88%)	1.8 (0.71-3.6)	7.8% (4.5-12%)	9.2% (5.2-14%)	97 (53-150)	4100 (2300-6100)
Togo	2000	Male	50 to 54	25% (12-45%)	75% (55-88%)	1.8 (0.69-3.8)	7.9% (4.4-12%)	9.4% (5.2-14%)	99 (53-160)	3700 (2000-5700)
Togo	2000	Male	55 to 59	25% (12-43%)	75% (57-88%)	1.8 (0.64-3.6)	7.7% (4.1-12%)	9.2% (4.8-14%)	100 (53-160)	3300 (1700-5100)
Togo	2000	Male	60 to 64	25% (12-42%)	75% (58-88%)	1.6 (0.55-3.6)	6.2% (2.9-10%)	7.3% (3.3-12%)	88 (39-150)	2400 (1100-4000)
Togo	2000	Male	65 to 69	25% (12-41%)	75% (59-88%)	1.3 (0.46-2.8)	5.2% (2.1-9.1%)	6.1% (2.5-11%)	76 (31-130)	1700 (720-3000)
Togo	2000	Male	70 to 74	23% (12-39%)	77% (61-88%)	1 (0.33-2.4)	4.1% (1.4-8.4%)	4.7% (1.6-9.9%)	58 (20-120)	1100 (360-2200)
Togo	2000	Male	75 to 79	21% (10-37%)	79% (63-90%)	0.76 (0.21-1.9)	2.9% (0.85-6.5%)	3.3% (0.95-7.5%)	33 (9.2-74)	470 (130-1000)
Togo	2000	Male	80 to 84	19% (8.7-35%)	81% (65-91%)	0.63 (0.17-1.5)	2.4% (0.77-5.6%)	2.8% (0.84-6.5%)	14 (4.3-33)	150 (48-340)
Togo	2000	Male	85 to 89	18% (8.7-33%)	82% (67-91%)	0.55 (0.15-1.4)	2% (0.58-4.8%)	2.3% (0.62-5.5%)	6.8 (1.9-17)	51 (15-120)
Togo	2000	Male	90 to 94	20% (9.3-35%)	80% (65-91%)	0.52 (0.12-1.4)	1.5% (0.31-4.2%)	1.7% (0.28-4.9%)	1.9 (0.32-5.6)	10 (2.2-29)
Togo	2000	Male	95 plus	23% (9.5-45%)	77% (55-91%)	0.55 (0.11-1.6)	1.6% (0.18-4.8%)	1.8% (0.073-5.6%)	0.48 (0.02-1.5)	1.7 (0.19-5.2)
Togo	2005	Female	All Ages	14% (10-19%)	86% (81-90%)	0.52 (0.38-0.67)	1.1% (0.71-1.6%)	2% (1-3.3%)	550 (280-910)	20000 (13000-29000)
Togo	2005	Female	15 to 19	13% (5.9-24%)	87% (76-94%)	0.43 (0.14-0.9)	1.1% (0.54-1.9%)	1.1% (0.29-2.4%)	5.4 (1.5-12)	690 (350-1300)
Togo	2005	Female	20 to 24	16% (7.4-29%)	84% (71-93%)	0.68 (0.27-1.3)	1.9% (1-3.1%)	1.6% (0.54-3.2%)	12 (4-23)	1500 (810-2300)
Togo	2005	Female	25 to 29	19% (8.4-33%)	81% (67-92%)	0.85 (0.34-1.5)	1.6% (0.86-2.6%)	1.4% (0.53-2.7%)	16 (6.7-30)	1600 (900-2500)
Togo	2005	Female	30 to 34	21% (8.8-37%)	79% (63-91%)	0.99 (0.41-1.9)	1.6% (0.91-2.6%)	1.6% (0.7-2.9%)	22 (11-38)	1600 (950-2700)
Togo	2005	Female	35 to 39	24% (11-41%)	76% (59-89%)	1.1 (0.44-2)	1.8% (0.96-3%)	1.9% (0.84-3.4%)	25 (12-43)	1500 (860-2500)
Togo	2005	Female	40 to 44	25% (13-43%)	75% (57-87%)	1.2 (0.48-2.3)	2.4% (1.2-4%)	2.7% (1.3-4.7%)	32 (15-56)	1700 (880-2800)
Togo	2005	Female	45 to 49	26% (11-43%)	74% (57-89%)	1.2 (0.45-2.4)	3.1% (1.5-5.3%)	3.8% (1.8-6.7%)	40 (18-72)	1800 (870-3100)
Togo	2005	Female	50 to 54	26% (12-46%)	74% (54-88%)	1.3 (0.49-2.5)	4.1% (2-6.8%)	5.1% (2.3-8.7%)	53 (23-92)	2000 (920-3400)
Togo	2005	Female	55 to 59	27% (13-47%)	73% (53-87%)	1.4 (0.49-2.8)	4.8% (2-8.6%)	6.1% (2.4-11%)	66 (25-120)	2100 (820-3900)
Togo	2005	Female	60 to 64	28% (14-48%)	72% (52-86%)	1.4 (0.5-3.1)	4.5% (1.7-8.1%)	5.6% (2.1-10%)	68 (23-130)	1900 (650-3500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	2005	Female	65 to 69	29% (15-47%)	71% (53-85%)	1.3 (0.46-2.7)	4.5% (1.5-8.7%)	5.4% (1.8-11%)	74 (23-150)	1700 (550-3400)
Togo	2005	Female	70 to 74	27% (14-43%)	73% (57-86%)	1 (0.34-2.3)	3.7% (0.98-8.3%)	4.5% (1.1-9.7%)	62 (15-140)	1100 (280-2600)
Togo	2005	Female	75 to 79	23% (11-40%)	77% (60-89%)	0.81 (0.24-2)	2.8% (0.55-6.7%)	3.3% (0.62-8%)	38 (6.9-98)	540 (100-1400)
Togo	2005	Female	80 to 84	21% (9.4-38%)	79% (62-91%)	0.71 (0.19-1.6)	2.3% (0.37-5.9%)	2.7% (0.39-7.1%)	18 (2.7-46)	190 (31-470)
Togo	2005	Female	85 to 89	22% (9.8-38%)	78% (62-90%)	0.73 (0.2-1.6)	2% (0.19-5.6%)	2.4% (0.16-6.6%)	10 (0.69-29)	78 (6.5-220)
Togo	2005	Female	90 to 94	25% (11-42%)	75% (58-89%)	0.8 (0.21-2)	1.9% (0.0043-5.7%)	2.2% (-0.067-6.7%)	4.1 (-0.13-13)	21 (0.05-65)
Togo	2005	Female	95 plus	29% (12-53%)	71% (47-88%)	0.93 (0.21-2.4)	1.9% (-0.24-6%)	2.2% (-0.42-7%)	1.1 (-0.21-3.6)	3.9 (-0.49-12)
Togo	2005	Male	All Ages	11% (8.4-16%)	89% (84-92%)	0.62 (0.47-0.79)	2% (1.5-2.6%)	3.2% (2.2-4.3%)	970 (670-1300)	40000 (31000-51000)
Togo	2005	Male	15 to 19	8.8% (4.2-16%)	91% (84-96%)	0.36 (0.12-0.78)	1.3% (0.67-2.4%)	1.4% (0.48-2.9%)	9.7 (3.3-20)	1000 (480-1800)
Togo	2005	Male	20 to 24	14% (6.1-24%)	86% (76-94%)	0.73 (0.28-1.4)	3.2% (1.7-5.2%)	3.2% (1.3-5.8%)	33 (14-60)	2800 (1500-4700)
Togo	2005	Male	25 to 29	17% (8.4-30%)	83% (70-92%)	1.1 (0.44-2)	4.1% (2.2-6.5%)	4.3% (2-7.5%)	56 (26-95)	4100 (2200-6500)
Togo	2005	Male	30 to 34	20% (9.5-36%)	80% (64-91%)	1.4 (0.62-2.6)	4.5% (2.7-7%)	4.8% (2.6-7.9%)	72 (41-110)	4500 (2800-6900)
Togo	2005	Male	35 to 39	22% (11-39%)	78% (61-89%)	1.6 (0.64-3)	5.1% (3-7.7%)	5.7% (3.1-8.7%)	81 (46-120)	4400 (2600-6600)
Togo	2005	Male	40 to 44	23% (12-40%)	77% (60-88%)	1.7 (0.67-3.2)	5.8% (3.4-8.8%)	6.6% (3.8-10%)	93 (51-140)	4500 (2500-6600)
Togo	2005	Male	45 to 49	24% (11-40%)	76% (60-89%)	1.8 (0.65-3.5)	7% (3.8-10%)	8.1% (4.3-12%)	110 (58-160)	4600 (2500-6900)
Togo	2005	Male	50 to 54	25% (12-44%)	75% (56-88%)	1.8 (0.67-3.6)	7.3% (3.9-11%)	8.5% (4.6-13%)	110 (58-170)	4000 (2200-6200)
Togo	2005	Male	55 to 59	25% (12-43%)	75% (57-88%)	1.7 (0.59-3.5)	7.4% (3.7-12%)	8.7% (4.3-13%)	110 (55-180)	3600 (1800-5700)
Togo	2005	Male	60 to 64	25% (12-42%)	75% (58-88%)	1.6 (0.53-3.4)	6% (2.7-9.8%)	6.9% (3-11%)	98 (44-160)	2700 (1200-4400)
Togo	2005	Male	65 to 69	24% (12-41%)	76% (59-88%)	1.3 (0.43-2.7)	5% (2-8.6%)	5.8% (2.4-10%)	82 (33-140)	1900 (750-3300)
Togo	2005	Male	70 to 74	23% (12-39%)	77% (61-88%)	0.98 (0.32-2.3)	3.8% (1.3-8.1%)	4.4% (1.5-9.3%)	62 (21-130)	1100 (380-2400)
Togo	2005	Male	75 to 79	21% (10-37%)	79% (63-90%)	0.75 (0.2-1.9)	2.7% (0.75-6.3%)	3.1% (0.84-7.3%)	34 (9.1-80)	480 (130-1100)
Togo	2005	Male	80 to 84	19% (8.4-33%)	81% (67-92%)	0.62 (0.17-1.4)	2.3% (0.74-5.4%)	2.6% (0.82-6.3%)	15 (4.6-35)	160 (50-370)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	2005	Male	85 to 89	18% (8.4-32%)	82% (68-92%)	0.53 (0.15-1.3)	1.9% (0.55-4.6%)	2.1% (0.57-5.3%)	6.9 (1.8-17)	52 (15-130)
Togo	2005	Male	90 to 94	19% (9.1-34%)	81% (66-91%)	0.52 (0.12-1.4)	1.4% (0.29-4.1%)	1.6% (0.25-4.8%)	2 (0.32-6.1)	11 (2.1-31)
Togo	2005	Male	95 plus	23% (9.2-44%)	77% (56-91%)	0.55 (0.1-1.6)	1.5% (0.18-4.6%)	1.7% (0.034-5.4%)	0.53 (0.011-1.7)	1.9 (0.23-6)
Togo	2010	Female	All Ages	14% (10-19%)	86% (81-90%)	0.54 (0.37-0.74)	1.3% (0.79-1.8%)	2.2% (1.1-3.6%)	600 (300-980)	22000 (14000-32000)
Togo	2010	Female	15 to 19	13% (6.1-23%)	87% (77-94%)	0.45 (0.15-0.98)	1% (0.52-1.8%)	1% (0.26-2.2%)	5.6 (1.5-12)	730 (360-1300)
Togo	2010	Female	20 to 24	17% (7.7-30%)	83% (70-92%)	0.72 (0.27-1.4)	2.2% (1.2-3.6%)	2% (0.69-4%)	14 (4.5-27)	1600 (890-2600)
Togo	2010	Female	25 to 29	19% (8.2-34%)	81% (66-92%)	0.88 (0.36-1.7)	2% (1.1-3.2%)	1.8% (0.75-3.5%)	18 (8-33)	1800 (1000-2800)
Togo	2010	Female	30 to 34	21% (9.2-38%)	79% (62-91%)	1 (0.42-1.9)	1.9% (1.1-3%)	1.9% (0.9-3.3%)	25 (13-43)	1900 (1100-3000)
Togo	2010	Female	35 to 39	23% (11-39%)	77% (61-89%)	1.1 (0.45-2.1)	2% (1.1-3.3%)	2.1% (0.99-3.7%)	29 (14-49)	1800 (1000-2900)
Togo	2010	Female	40 to 44	25% (13-42%)	75% (58-87%)	1.2 (0.46-2.3)	2.5% (1.3-4.1%)	3% (1.4-4.9%)	38 (18-62)	2000 (1000-3100)
Togo	2010	Female	45 to 49	26% (12-43%)	74% (57-88%)	1.2 (0.46-2.5)	3.3% (1.6-5.4%)	4.1% (1.9-6.9%)	46 (21-80)	2100 (1000-3500)
Togo	2010	Female	50 to 54	26% (12-46%)	74% (54-88%)	1.3 (0.49-2.7)	4.2% (1.9-7.2%)	5.3% (2.4-9.2%)	59 (25-110)	2200 (980-4000)
Togo	2010	Female	55 to 59	26% (13-46%)	74% (54-87%)	1.4 (0.49-3)	4.9% (2-8.8%)	6.3% (2.5-11%)	73 (28-140)	2400 (930-4400)
Togo	2010	Female	60 to 64	28% (14-47%)	72% (53-86%)	1.5 (0.5-3.1)	4.6% (1.7-8.2%)	5.7% (2.1-10%)	75 (27-140)	2100 (750-3800)
Togo	2010	Female	65 to 69	29% (14-47%)	71% (53-86%)	1.3 (0.45-2.8)	4.5% (1.5-8.5%)	5.5% (1.8-10%)	80 (25-160)	1800 (580-3600)
Togo	2010	Female	70 to 74	26% (14-43%)	74% (57-86%)	1.1 (0.33-2.5)	3.7% (0.94-8.4%)	4.4% (1.1-10%)	66 (16-160)	1200 (300-2900)
Togo	2010	Female	75 to 79	23% (11-40%)	77% (60-89%)	0.81 (0.23-2)	2.7% (0.49-6.9%)	3.2% (0.52-8.1%)	40 (6.8-110)	570 (100-1500)
Togo	2010	Female	80 to 84	21% (9.4-37%)	79% (63-91%)	0.72 (0.21-1.7)	2.2% (0.39-5.8%)	2.6% (0.41-6.9%)	19 (2.7-50)	200 (31-510)
Togo	2010	Female	85 to 89	21% (10-38%)	79% (62-90%)	0.73 (0.19-1.7)	2% (0.15-5.8%)	2.3% (0.1-6.8%)	11 (0.51-33)	83 (6.1-240)
Togo	2010	Female	90 to 94	24% (11-43%)	76% (57-89%)	0.8 (0.2-1.9)	1.9% (-0.045-5.6%)	2.2% (-0.13-6.8%)	4.4 (-0.25-14)	23 (-0.55-68)
Togo	2010	Female	95 plus	29% (12-53%)	71% (47-88%)	0.93 (0.21-2.5)	1.9% (-0.23-6.1%)	2.2% (-0.42-7.1%)	1.3 (-0.25-4.1)	4.5 (-0.56-14)
Togo	2010	Male	All Ages	12% (8.6-16%)	88% (84-91%)	0.65 (0.46-0.88)	2.3% (1.7-3%)	3.6% (2.4-4.8%)	1100 (740-1500)	46000 (34000-59000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	2010	Male	15 to 19	8.9% (4.2-16%)	91% (84-96%)	0.37 (0.12-0.82)	1.3% (0.66-2.4%)	1.3% (0.48-2.8%)	10 (3.6-22)	1100 (520-1900)
Togo	2010	Male	20 to 24	14% (6.1-25%)	86% (75-94%)	0.75 (0.27-1.5)	3.5% (1.9-5.7%)	3.6% (1.5-6.5%)	37 (15-66)	3200 (1700-5200)
Togo	2010	Male	25 to 29	17% (7.9-31%)	83% (69-92%)	1.1 (0.43-2.1)	4.8% (2.6-7.5%)	5.2% (2.4-8.8%)	65 (31-110)	4700 (2500-7500)
Togo	2010	Male	30 to 34	20% (8.9-36%)	80% (64-91%)	1.4 (0.57-2.6)	5.3% (3-8.1%)	5.8% (3.1-9.3%)	87 (47-140)	5500 (3300-8300)
Togo	2010	Male	35 to 39	23% (10-39%)	77% (61-90%)	1.6 (0.61-3.1)	5.6% (3.3-8.5%)	6.3% (3.5-9.8%)	99 (54-150)	5500 (3100-8200)
Togo	2010	Male	40 to 44	23% (12-40%)	77% (60-88%)	1.7 (0.64-3.4)	6.3% (3.5-9.6%)	7.2% (3.9-11%)	110 (59-170)	5300 (2900-8200)
Togo	2010	Male	45 to 49	24% (11-41%)	76% (59-89%)	1.8 (0.65-3.7)	7.5% (4.1-11%)	8.7% (4.6-13%)	120 (65-190)	5300 (2800-8000)
Togo	2010	Male	50 to 54	25% (11-43%)	75% (57-89%)	1.8 (0.66-4)	7.5% (4-12%)	8.8% (4.6-14%)	120 (63-190)	4500 (2400-7100)
Togo	2010	Male	55 to 59	25% (13-43%)	75% (57-87%)	1.8 (0.58-3.8)	7.6% (3.7-12%)	8.9% (4.3-14%)	120 (60-200)	4000 (2000-6400)
Togo	2010	Male	60 to 64	25% (12-43%)	75% (57-88%)	1.6 (0.52-3.5)	6.1% (2.7-10%)	7.1% (3.1-12%)	110 (47-180)	3000 (1300-5000)
Togo	2010	Male	65 to 69	24% (12-41%)	76% (59-88%)	1.3 (0.41-2.9)	5% (2-8.8%)	5.8% (2.3-10%)	92 (36-170)	2100 (830-3700)
Togo	2010	Male	70 to 74	23% (11-38%)	77% (62-89%)	1 (0.31-2.5)	3.8% (1.2-8.2%)	4.4% (1.4-9.5%)	68 (22-140)	1200 (400-2600)
Togo	2010	Male	75 to 79	21% (9.9-37%)	79% (63-90%)	0.76 (0.21-1.9)	2.7% (0.68-6.3%)	3% (0.75-7.3%)	36 (9.2-87)	510 (140-1200)
Togo	2010	Male	80 to 84	19% (8.5-34%)	81% (66-92%)	0.62 (0.18-1.5)	2.2% (0.69-5.2%)	2.5% (0.74-6%)	16 (4.7-38)	170 (52-400)
Togo	2010	Male	85 to 89	18% (8.6-32%)	82% (68-91%)	0.53 (0.14-1.3)	1.8% (0.53-4.4%)	2% (0.54-5.1%)	7.2 (1.9-18)	55 (16-140)
Togo	2010	Male	90 to 94	19% (8.5-34%)	81% (66-92%)	0.51 (0.11-1.3)	1.3% (0.26-3.8%)	1.5% (0.21-4.4%)	2.1 (0.28-6.1)	11 (2.2-32)
Togo	2010	Male	95 plus	23% (9.2-44%)	77% (56-91%)	0.53 (0.098-1.6)	1.4% (0.15-4.6%)	1.5% (0.027-5.2%)	0.56 (0.0098-1.9)	2.1 (0.21-6.6)
Togo	2016	Female	All Ages	14% (11-20%)	86% (80-89%)	0.61 (0.35-0.98)	1.6% (0.94-2.4%)	2.9% (1.2-4.8%)	720 (300-1200)	26000 (15000-39000)
Togo	2016	Female	15 to 19	13% (6-23%)	87% (77-94%)	0.48 (0.15-1.1)	1% (0.52-1.8%)	0.91% (0.22-2%)	5.5 (1.3-13)	790 (390-1400)
Togo	2016	Female	20 to 24	16% (7.3-29%)	84% (71-93%)	0.78 (0.24-1.7)	2.4% (1.4-3.8%)	2.1% (0.63-4.2%)	13 (3.7-25)	1700 (930-2700)
Togo	2016	Female	25 to 29	19% (8.2-33%)	81% (67-92%)	0.96 (0.35-2)	2.6% (1.5-4.2%)	2.6% (1-5.2%)	19 (7.7-36)	1900 (1100-3100)
Togo	2016	Female	30 to 34	21% (9.5-37%)	79% (63-90%)	1.1 (0.41-2.3)	2.6% (1.5-4.1%)	2.8% (1.3-5.2%)	27 (13-47)	2100 (1200-3300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	2016	Female	35 to 39	24% (11-41%)	76% (59-89%)	1.2 (0.44-2.5)	2.7% (1.5-4.4%)	3.3% (1.5-5.6%)	35 (16-59)	2200 (1200-3500)
Togo	2016	Female	40 to 44	25% (13-43%)	75% (57-87%)	1.3 (0.45-2.6)	3.4% (1.7-5.4%)	4.4% (2-7.3%)	46 (21-77)	2400 (1200-3800)
Togo	2016	Female	45 to 49	26% (12-42%)	74% (58-88%)	1.3 (0.45-3)	4.2% (1.9-7.1%)	5.6% (2.5-9.5%)	57 (25-100)	2500 (1200-4400)
Togo	2016	Female	50 to 54	26% (12-45%)	74% (55-88%)	1.4 (0.47-3.1)	4.8% (2.1-8.1%)	6.4% (2.6-11%)	71 (29-120)	2700 (1200-4700)
Togo	2016	Female	55 to 59	27% (13-45%)	73% (55-87%)	1.6 (0.47-3.6)	5.3% (2-9.3%)	6.9% (2.6-12%)	85 (31-160)	2800 (1000-5000)
Togo	2016	Female	60 to 64	28% (14-47%)	72% (53-86%)	1.6 (0.49-3.8)	4.8% (1.6-8.7%)	5.9% (2.1-11%)	87 (27-160)	2400 (780-4400)
Togo	2016	Female	65 to 69	29% (15-47%)	71% (53-85%)	1.4 (0.42-3.3)	4.7% (1.4-9.3%)	5.7% (1.6-11%)	97 (27-200)	2200 (630-4500)
Togo	2016	Female	70 to 74	26% (14-43%)	74% (57-86%)	1.2 (0.33-2.7)	3.9% (0.83-8.6%)	4.6% (0.92-10%)	81 (17-190)	1500 (320-3300)
Togo	2016	Female	75 to 79	23% (11-39%)	77% (61-89%)	0.9 (0.24-2.4)	2.9% (0.38-7.3%)	3.4% (0.43-8.6%)	52 (6.3-130)	740 (99-1900)
Togo	2016	Female	80 to 84	21% (9.3-37%)	79% (63-91%)	0.79 (0.21-2)	2.4% (0.26-6.3%)	2.8% (0.26-7.4%)	24 (2.2-66)	260 (28-680)
Togo	2016	Female	85 to 89	21% (10-38%)	79% (62-90%)	0.8 (0.2-2)	2.2% (0.099-6.1%)	2.5% (0.061-7.2%)	14 (0.34-40)	110 (4.6-290)
Togo	2016	Female	90 to 94	25% (12-43%)	75% (57-88%)	0.89 (0.2-2.2)	2.1% (-0.12-5.9%)	2.4% (-0.22-7%)	5.8 (-0.54-17)	30 (-1.7-86)
Togo	2016	Female	95 plus	30% (12-53%)	70% (47-88%)	1 (0.2-2.8)	2.1% (-0.26-6.6%)	2.4% (-0.46-7.5%)	1.7 (-0.33-5.4)	5.9 (-0.73-18)
Togo	2016	Male	All Ages	12% (8.6-16%)	88% (84-91%)	0.73 (0.42-1.2)	2.6% (1.8-3.5%)	4% (2.4-5.5%)	1200 (730-1700)	50000 (34000-67000)
Togo	2016	Male	15 to 19	8.7% (4.2-16%)	91% (84-96%)	0.39 (0.12-0.93)	1.3% (0.62-2.3%)	1.2% (0.39-2.6%)	11 (3.4-22)	1200 (550-2100)
Togo	2016	Male	20 to 24	13% (6.3-24%)	87% (76-94%)	0.8 (0.26-1.8)	3.4% (1.8-5.6%)	3.5% (1.3-6.3%)	35 (13-65)	3200 (1600-5300)
Togo	2016	Male	25 to 29	17% (8-31%)	83% (69-92%)	1.2 (0.42-2.6)	5% (2.7-7.9%)	5.5% (2.5-9.3%)	64 (29-110)	4800 (2500-7700)
Togo	2016	Male	30 to 34	20% (9.1-36%)	80% (64-91%)	1.5 (0.57-3.1)	5.9% (3.3-9.1%)	6.6% (3.4-11%)	91 (47-150)	5900 (3500-9000)
Togo	2016	Male	35 to 39	23% (11-39%)	77% (61-89%)	1.7 (0.61-3.4)	6.1% (3.5-9.3%)	7% (3.8-11%)	110 (58-170)	6200 (3500-9400)
Togo	2016	Male	40 to 44	24% (12-41%)	76% (59-88%)	1.9 (0.66-4)	6.6% (3.5-9.9%)	7.7% (4-11%)	130 (64-200)	6100 (3300-9400)
Togo	2016	Male	45 to 49	24% (11-40%)	76% (60-89%)	2 (0.65-4.6)	7.7% (4.2-12%)	9.1% (4.8-14%)	140 (72-220)	5900 (3200-9200)
Togo	2016	Male	50 to 54	25% (11-43%)	75% (57-89%)	2 (0.65-4.7)	7.4% (3.8-11%)	8.8% (4.4-14%)	130 (64-210)	4900 (2500-7600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Togo	2016	Male	55 to 59	25% (12-42%)	75% (58-88%)	2 (0.59-4.4)	7.4% (3.6-12%)	8.7% (4.2-14%)	130 (61-210)	4200 (2000-6800)
Togo	2016	Male	60 to 64	25% (12-42%)	75% (58-88%)	1.8 (0.53-4.2)	5.9% (2.7-10%)	7% (3-12%)	110 (50-200)	3100 (1400-5400)
Togo	2016	Male	65 to 69	24% (12-42%)	76% (58-88%)	1.5 (0.43-3.3)	4.9% (1.9-8.6%)	5.8% (2.2-10%)	98 (38-180)	2300 (890-4000)
Togo	2016	Male	70 to 74	23% (12-39%)	77% (61-88%)	1.1 (0.3-2.6)	3.8% (1.1-7.7%)	4.4% (1.2-9.2%)	76 (21-160)	1400 (400-2900)
Togo	2016	Male	75 to 79	21% (10-36%)	79% (64-90%)	0.83 (0.21-2.2)	2.7% (0.64-6.3%)	3.1% (0.71-7.4%)	42 (9.5-100)	600 (140-1400)
Togo	2016	Male	80 to 84	19% (8.6-33%)	81% (67-91%)	0.67 (0.17-1.7)	2.3% (0.59-5.3%)	2.6% (0.63-6.2%)	19 (4.6-43)	200 (52-460)
Togo	2016	Male	85 to 89	18% (8.2-32%)	82% (68-92%)	0.58 (0.14-1.5)	1.8% (0.41-4.7%)	2.1% (0.4-5.6%)	8.4 (1.7-22)	64 (14-160)
Togo	2016	Male	90 to 94	19% (9.1-34%)	81% (66-91%)	0.56 (0.12-1.5)	1.4% (0.22-4.2%)	1.6% (0.16-4.9%)	2.5 (0.26-7.6)	13 (2.1-39)
Togo	2016	Male	95 plus	23% (9.7-43%)	77% (57-90%)	0.6 (0.1-1.7)	1.5% (0.1-4.7%)	1.6% (-0.016-5.5%)	0.65 (-0.0062-2.2)	2.4 (0.16-7.5)
American Samoa	1990	Female	All Ages	18% (12-25%)	82% (75-88%)	0.35 (0.18-0.61)	0.52% (0.1-1.2%)	0.38% (-0.37-1.6%)	0.32 (-0.3-1.4)	26 (4.9-60)
American Samoa	1990	Female	15 to 19	19% (8.2-35%)	81% (65-92%)	0.4 (0.13-0.86)	0.93% (0.46-1.8%)	1.5% (0.45-3.7%)	0.019 (0.0058-0.047)	2.5 (1.2-4.8)
American Samoa	1990	Female	20 to 24	23% (9.7-41%)	77% (59-90%)	0.63 (0.23-1.4)	1.6% (0.89-2.7%)	2.1% (0.66-4.7%)	0.034 (0.01-0.075)	4.9 (2.6-8.7)
American Samoa	1990	Female	25 to 29	25% (10-44%)	75% (56-90%)	0.65 (0.23-1.3)	1.6% (0.98-2.8%)	2.3% (0.95-4.7%)	0.042 (0.017-0.088)	5.2 (3-8.9)
American Samoa	1990	Female	30 to 34	25% (10-45%)	75% (55-90%)	0.67 (0.24-1.4)	1.4% (0.73-2.5%)	2% (0.72-4.3%)	0.039 (0.014-0.087)	4.1 (2.1-7.5)
American Samoa	1990	Female	35 to 39	26% (10-44%)	74% (56-90%)	0.69 (0.25-1.5)	1.3% (0.57-2.8%)	2.1% (0.75-4.7%)	0.049 (0.017-0.11)	3.7 (1.6-7.7)
American Samoa	1990	Female	40 to 44	25% (12-43%)	75% (57-88%)	0.64 (0.23-1.4)	1.1% (0.27-2.7%)	1.9% (0.46-4.5%)	0.052 (0.011-0.13)	3 (0.7-7.3)
American Samoa	1990	Female	45 to 49	25% (11-42%)	75% (58-89%)	0.55 (0.17-1.2)	0.75% (-0.088-2.2%)	1.4% (0.12-3.6%)	0.042 (0.0037-0.11)	1.8 (-0.21-5.2)
American Samoa	1990	Female	50 to 54	24% (11-43%)	76% (57-89%)	0.49 (0.15-1.1)	0.42% (-0.51-1.9%)	0.9% (-0.33-3%)	0.036 (-0.014-0.12)	1.1 (-1.4-4.8)
American Samoa	1990	Female	55 to 59	24% (11-43%)	76% (57-89%)	0.44 (0.14-1)	0.24% (-0.76-1.7%)	0.64% (-0.56-2.7%)	0.028 (-0.024-0.12)	0.53 (-1.8-3.8)
American Samoa	1990	Female	60 to 64	24% (11-44%)	76% (56-89%)	0.42 (0.13-1)	-0.073% (-1.1-1.4%)	0.17% (-1-2.1%)	0.01 (-0.065-0.13)	-0.19 (-2.8-3.5)
American Samoa	1990	Female	65 to 69	25% (11-42%)	75% (58-89%)	0.42 (0.13-0.98)	-0.1% (-1.2-1.2%)	0.11% (-1.2-1.8%)	0.0083 (-0.084-0.13)	-0.25 (-2.8-2.8)
American Samoa	1990	Female	70 to 74	25% (11-41%)	75% (59-89%)	0.41 (0.13-0.93)	-0.23% (-1.2-1.1%)	-0.12% (-1.3-1.5%)	-0.0084 (-0.087-0.1)	-0.39 (-2.1-1.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	1990	Female	75 to 79	24% (10-43%)	76% (57-90%)	0.39 (0.11-1)	-0.099% (-1.1-1.3%)	0.013% (-1.1-1.7%)	0.0008 (-0.072-0.11)	-0.12 (-1.3-1.5)
American Samoa	1990	Female	80 to 84	23% (9.3-43%)	77% (57-91%)	0.37 (0.11-0.94)	-0.18% (-1.1-1.1%)	-0.13% (-1.2-1.4%)	-0.013 (-0.12-0.13)	-0.23 (-1.4-1.5)
American Samoa	1990	Female	85 to 89	24% (10-43%)	76% (57-90%)	0.38 (0.11-0.9)	-0.25% (-1.2-0.96%)	-0.24% (-1.3-1.1%)	-0.016 (-0.091-0.077)	-0.15 (-0.74-0.58)
American Samoa	1990	Female	90 to 94	27% (12-47%)	73% (53-88%)	0.43 (0.13-1.1)	-0.23% (-1.1-1.1%)	-0.23% (-1.3-1.3%)	-0.0065 (-0.035-0.036)	-0.039 (-0.19-0.2)
American Samoa	1990	Female	95 plus	32% (13-54%)	68% (46-87%)	0.5 (0.15-1.3)	-0.38% (-1.6-1.2%)	-0.47% (-1.8-1.2%)	-0.0038 (-0.015-0.0099)	-0.012 (-0.051-0.038)
American Samoa	1990	Male	All Ages	38% (30-48%)	62% (52-70%)	0.79 (0.43-1.3)	2.2% (1.1-3.9%)	2% (0.68-4.2%)	3 (0.99-6.3)	170 (79-300)
American Samoa	1990	Male	15 to 19	35% (20-53%)	65% (47-80%)	0.49 (0.19-1)	1.8% (0.95-3.3%)	2.4% (0.93-4.6%)	0.082 (0.032-0.16)	7.5 (3.8-13)
American Samoa	1990	Male	20 to 24	53% (34-71%)	47% (29-66%)	1.3 (0.6-2.4)	4.9% (2.6-8.2%)	6.1% (2.8-11%)	0.31 (0.14-0.56)	25 (13-42)
American Samoa	1990	Male	25 to 29	61% (41-77%)	39% (23-59%)	1.9 (0.93-3.4)	6.1% (3.4-10%)	7.6% (3.9-14%)	0.39 (0.19-0.7)	29 (16-49)
American Samoa	1990	Male	30 to 34	62% (41-79%)	38% (21-59%)	2 (0.91-3.6)	5.4% (2.7-9.4%)	6.6% (3-12%)	0.38 (0.16-0.72)	26 (13-46)
American Samoa	1990	Male	35 to 39	60% (42-77%)	40% (23-58%)	1.8 (0.78-3.4)	4.6% (2.1-8.9%)	5.7% (2.2-11%)	0.32 (0.12-0.67)	20 (9-38)
American Samoa	1990	Male	40 to 44	57% (39-75%)	43% (25-61%)	1.5 (0.64-3)	3.5% (1.1-7.4%)	4.1% (1.1-9.3%)	0.28 (0.068-0.66)	16 (5.2-34)
American Samoa	1990	Male	45 to 49	54% (32-72%)	46% (28-68%)	1.3 (0.55-2.7)	2.8% (0.78-6.5%)	3.2% (0.59-7.7%)	0.26 (0.044-0.63)	13 (3.6-30)
American Samoa	1990	Male	50 to 54	51% (32-70%)	49% (30-68%)	1.1 (0.44-2.2)	2.4% (0.52-5.7%)	2.6% (0.36-6.9%)	0.27 (0.04-0.72)	12 (2.6-30)
American Samoa	1990	Male	55 to 59	48% (31-68%)	52% (32-69%)	0.88 (0.35-1.8)	1.5% (0.059-4.4%)	1.6% (-0.076-5.2%)	0.18 (-0.0084-0.56)	6.8 (0.25-20)
American Samoa	1990	Male	60 to 64	46% (27-66%)	54% (34-73%)	0.74 (0.29-1.6)	1.2% (-0.065-3.8%)	1.3% (-0.17-4.2%)	0.16 (-0.021-0.54)	5 (-0.27-16)
American Samoa	1990	Male	65 to 69	44% (26-63%)	56% (37-74%)	0.67 (0.26-1.3)	0.92% (-0.13-2.9%)	1% (-0.18-3.3%)	0.13 (-0.023-0.43)	3.3 (-0.44-10)
American Samoa	1990	Male	70 to 74	41% (24-59%)	59% (41-76%)	0.59 (0.24-1.2)	0.69% (-0.25-2.8%)	0.75% (-0.3-3.1%)	0.099 (-0.042-0.41)	2 (-0.7-7.9)
American Samoa	1990	Male	75 to 79	38% (22-57%)	62% (43-78%)	0.5 (0.19-1.1)	0.52% (-0.21-2.2%)	0.55% (-0.3-2.4%)	0.067 (-0.037-0.28)	1.1 (-0.45-4.3)
American Samoa	1990	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.44 (0.17-0.95)	0.46% (-0.13-1.7%)	0.47% (-0.19-1.9%)	0.046 (-0.018-0.18)	0.55 (-0.16-2)
American Samoa	1990	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.42 (0.17-0.87)	0.32% (-0.24-1.4%)	0.32% (-0.33-1.5%)	0.015 (-0.016-0.072)	0.14 (-0.1-0.58)
American Samoa	1990	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.43 (0.17-0.92)	0.21% (-0.32-1.2%)	0.18% (-0.42-1.2%)	0.0026 (-0.0059-0.018)	0.018 (-0.028-0.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	1990	Male	95 plus	38% (20-59%)	62% (41-80%)	0.47 (0.17-1)	0.14% (-0.51-1.4%)	0.044% (-0.7-1.4%)	0.00012 (-0.002-0.004)	0.0016 (-0.0058-0.016)
American Samoa	1995	Female	All Ages	18% (12-25%)	82% (75-88%)	0.35 (0.18-0.6)	0.53% (0.043-1.3%)	0.36% (-0.45-1.6%)	0.36 (-0.45-1.5)	30 (2.5-71)
American Samoa	1995	Female	15 to 19	19% (8.8-33%)	81% (67-91%)	0.4 (0.14-0.9)	0.95% (0.47-1.8%)	1.6% (0.47-3.8%)	0.021 (0.0064-0.051)	2.7 (1.3-5.1)
American Samoa	1995	Female	20 to 24	23% (10-41%)	77% (59-90%)	0.64 (0.23-1.3)	1.6% (0.9-2.8%)	2.1% (0.7-4.7%)	0.036 (0.012-0.082)	5 (2.7-8.8)
American Samoa	1995	Female	25 to 29	24% (10-42%)	76% (58-90%)	0.65 (0.24-1.4)	1.7% (0.98-3%)	2.3% (0.96-4.8%)	0.05 (0.02-0.1)	5.6 (3.2-9.9)
American Samoa	1995	Female	30 to 34	25% (9.8-46%)	75% (54-90%)	0.68 (0.23-1.4)	1.5% (0.71-2.7%)	2% (0.67-4.3%)	0.053 (0.017-0.11)	5.1 (2.4-9.5)
American Samoa	1995	Female	35 to 39	26% (11-44%)	74% (56-89%)	0.7 (0.25-1.5)	1.4% (0.57-2.9%)	2.1% (0.73-4.6%)	0.072 (0.024-0.16)	5.1 (2.1-10)
American Samoa	1995	Female	40 to 44	25% (12-42%)	75% (58-88%)	0.64 (0.24-1.4)	1.1% (0.23-2.8%)	1.8% (0.37-4.4%)	0.072 (0.014-0.18)	4.1 (0.8-9.8)
American Samoa	1995	Female	45 to 49	24% (10-42%)	76% (58-90%)	0.55 (0.16-1.2)	0.73% (-0.2-2.2%)	1.3% (-0.011-3.6%)	0.058 (-0.00052-0.17)	2.5 (-0.73-7.7)
American Samoa	1995	Female	50 to 54	24% (10-42%)	76% (58-90%)	0.48 (0.15-1.1)	0.37% (-0.7-1.9%)	0.78% (-0.56-2.8%)	0.043 (-0.031-0.16)	1.2 (-2.3-6.4)
American Samoa	1995	Female	55 to 59	23% (11-41%)	77% (59-89%)	0.44 (0.14-1)	0.18% (-0.93-1.6%)	0.53% (-0.72-2.5%)	0.03 (-0.04-0.14)	0.5 (-2.5-4.4)
American Samoa	1995	Female	60 to 64	23% (10-42%)	77% (58-90%)	0.42 (0.13-1)	-0.14% (-1.3-1.3%)	0.074% (-1.2-1.8%)	0.0055 (-0.09-0.14)	-0.41 (-3.8-3.8)
American Samoa	1995	Female	65 to 69	25% (11-44%)	75% (56-89%)	0.43 (0.13-1)	-0.18% (-1.4-1.2%)	0.009% (-1.4-1.7%)	0.0007 (-0.11-0.14)	-0.47 (-3.5-3)
American Samoa	1995	Female	70 to 74	25% (12-43%)	75% (57-88%)	0.41 (0.12-0.94)	-0.31% (-1.4-1.1%)	-0.22% (-1.5-1.6%)	-0.021 (-0.14-0.13)	-0.7 (-3.1-2.4)
American Samoa	1995	Female	75 to 79	24% (10-42%)	76% (58-90%)	0.38 (0.11-0.96)	-0.15% (-1.1-1.2%)	-0.055% (-1.2-1.5%)	-0.0042 (-0.087-0.11)	-0.21 (-1.5-1.6)
American Samoa	1995	Female	80 to 84	23% (9.9-43%)	77% (57-90%)	0.37 (0.11-0.93)	-0.23% (-1.1-0.98%)	-0.19% (-1.2-1.2%)	-0.02 (-0.13-0.14)	-0.31 (-1.5-1.4)
American Samoa	1995	Female	85 to 89	24% (10-42%)	76% (58-90%)	0.38 (0.12-0.9)	-0.3% (-1.2-0.84%)	-0.31% (-1.4-1%)	-0.025 (-0.11-0.079)	-0.22 (-0.92-0.61)
American Samoa	1995	Female	90 to 94	27% (12-47%)	73% (53-88%)	0.43 (0.13-1)	-0.28% (-1.2-1%)	-0.29% (-1.3-1.1%)	-0.01 (-0.046-0.041)	-0.06 (-0.26-0.21)
American Samoa	1995	Female	95 plus	32% (14-55%)	68% (45-86%)	0.51 (0.15-1.2)	-0.45% (-1.7-1.2%)	-0.55% (-2-1.3%)	-0.0053 (-0.019-0.013)	-0.017 (-0.066-0.048)
American Samoa	1995	Male	All Ages	38% (30-47%)	62% (53-70%)	0.8 (0.45-1.4)	2.2% (1.1-4.1%)	2.1% (0.72-4.5%)	3.2 (1.1-6.7)	180 (85-330)
American Samoa	1995	Male	15 to 19	35% (20-53%)	65% (47-80%)	0.5 (0.19-1)	1.9% (0.94-3.3%)	2.4% (0.97-4.8%)	0.087 (0.035-0.17)	8 (4-14)
American Samoa	1995	Male	20 to 24	53% (34-70%)	47% (30-66%)	1.4 (0.62-2.6)	5% (2.6-8.6%)	6.2% (2.8-11%)	0.3 (0.14-0.56)	25 (13-43)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	1995	Male	25 to 29	60% (41-77%)	40% (23-59%)	1.9 (0.93-3.4)	6.1% (3.3-10%)	7.8% (3.8-14%)	0.39 (0.19-0.71)	29 (16-51)
American Samoa	1995	Male	30 to 34	61% (40-78%)	39% (22-60%)	2 (0.9-3.6)	5.4% (2.8-9.4%)	6.6% (3.1-12%)	0.38 (0.16-0.72)	27 (13-48)
American Samoa	1995	Male	35 to 39	60% (42-77%)	40% (23-58%)	1.8 (0.81-3.4)	4.6% (2.2-8.6%)	5.7% (2.3-11%)	0.36 (0.14-0.75)	23 (10-44)
American Samoa	1995	Male	40 to 44	58% (40-75%)	42% (25-60%)	1.6 (0.66-3)	3.5% (1.2-7.5%)	4.1% (1.2-9.3%)	0.31 (0.081-0.73)	18 (6-39)
American Samoa	1995	Male	45 to 49	53% (32-72%)	47% (28-68%)	1.3 (0.52-2.7)	2.8% (0.81-6.4%)	3.2% (0.73-7.8%)	0.29 (0.062-0.69)	15 (4.1-33)
American Samoa	1995	Male	50 to 54	50% (30-69%)	50% (31-70%)	1.1 (0.43-2.2)	2.4% (0.53-5.7%)	2.7% (0.36-6.8%)	0.29 (0.041-0.76)	13 (3-31)
American Samoa	1995	Male	55 to 59	48% (30-68%)	52% (32-70%)	0.9 (0.34-1.9)	1.5% (0.05-4.3%)	1.7% (-0.085-5%)	0.2 (-0.0098-0.62)	7.7 (0.25-22)
American Samoa	1995	Male	60 to 64	46% (27-66%)	54% (34-73%)	0.75 (0.29-1.7)	1.2% (-0.082-3.9%)	1.3% (-0.17-4.4%)	0.19 (-0.024-0.63)	5.7 (-0.37-19)
American Samoa	1995	Male	65 to 69	45% (26-63%)	55% (37-74%)	0.68 (0.27-1.4)	0.94% (-0.14-3.1%)	1% (-0.19-3.6%)	0.14 (-0.027-0.5)	3.5 (-0.5-12)
American Samoa	1995	Male	70 to 74	42% (25-60%)	58% (40-75%)	0.6 (0.24-1.2)	0.69% (-0.27-2.7%)	0.75% (-0.32-3%)	0.096 (-0.042-0.39)	1.9 (-0.72-7.6)
American Samoa	1995	Male	75 to 79	38% (22-56%)	62% (44-78%)	0.51 (0.19-1.1)	0.53% (-0.22-2.2%)	0.55% (-0.26-2.3%)	0.062 (-0.029-0.27)	1 (-0.4-4.1)
American Samoa	1995	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.45 (0.18-0.94)	0.47% (-0.14-1.7%)	0.48% (-0.19-1.8%)	0.041 (-0.015-0.16)	0.49 (-0.15-1.8)
American Samoa	1995	Male	85 to 89	34% (19-50%)	66% (50-81%)	0.42 (0.17-0.87)	0.33% (-0.23-1.3%)	0.33% (-0.32-1.5%)	0.017 (-0.016-0.074)	0.15 (-0.1-0.59)
American Samoa	1995	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.44 (0.17-0.94)	0.22% (-0.32-1.2%)	0.19% (-0.43-1.2%)	0.0033 (-0.0075-0.021)	0.023 (-0.033-0.12)
American Samoa	1995	Male	95 plus	39% (21-60%)	61% (40-79%)	0.48 (0.17-1.1)	0.16% (-0.54-1.5%)	0.064% (-0.73-1.6%)	0.00021 (-0.0025-0.0053)	0.0021 (-0.0072-0.02)
American Samoa	2000	Female	All Ages	18% (13-25%)	82% (75-87%)	0.36 (0.18-0.61)	0.51% (-0.021-1.3%)	0.32% (-0.52-1.6%)	0.36 (-0.57-1.7)	31 (-1.2-79)
American Samoa	2000	Female	15 to 19	19% (8.3-33%)	81% (67-92%)	0.42 (0.14-0.92)	0.96% (0.46-1.9%)	1.6% (0.46-3.6%)	0.021 (0.0063-0.051)	2.7 (1.3-5.4)
American Samoa	2000	Female	20 to 24	23% (9.9-40%)	77% (60-90%)	0.65 (0.23-1.4)	1.6% (0.86-2.7%)	2.2% (0.71-4.7%)	0.035 (0.011-0.076)	4.8 (2.5-8.3)
American Samoa	2000	Female	25 to 29	24% (10-43%)	76% (57-90%)	0.67 (0.24-1.4)	1.7% (0.97-2.9%)	2.3% (0.93-4.7%)	0.05 (0.02-0.1)	5.6 (3.1-10)
American Samoa	2000	Female	30 to 34	25% (9.9-45%)	75% (55-90%)	0.69 (0.24-1.5)	1.4% (0.7-2.8%)	2% (0.71-4.3%)	0.061 (0.02-0.14)	5.8 (2.6-11)
American Samoa	2000	Female	35 to 39	26% (10-45%)	74% (55-90%)	0.72 (0.27-1.5)	1.4% (0.55-2.9%)	2.1% (0.76-4.5%)	0.09 (0.03-0.2)	6.4 (2.5-14)
American Samoa	2000	Female	40 to 44	25% (12-43%)	75% (57-88%)	0.67 (0.24-1.4)	1.1% (0.13-2.9%)	1.8% (0.34-4.5%)	0.09 (0.017-0.23)	5 (0.59-12)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	2000	Female	45 to 49	25% (11-42%)	75% (58-89%)	0.57 (0.19-1.2)	0.68% (-0.31-2.2%)	1.2% (-0.084-3.5%)	0.073 (-0.0043-0.2)	3 (-1.3-9.3)
American Samoa	2000	Female	50 to 54	24% (11-42%)	76% (58-89%)	0.5 (0.16-1.1)	0.3% (-0.82-1.9%)	0.71% (-0.68-2.8%)	0.047 (-0.048-0.19)	1.2 (-3.4-7.3)
American Samoa	2000	Female	55 to 59	24% (11-42%)	76% (58-89%)	0.46 (0.14-1.1)	0.093% (-1.1-1.5%)	0.43% (-0.91-2.3%)	0.028 (-0.058-0.15)	0.29 (-3.4-4.8)
American Samoa	2000	Female	60 to 64	24% (11-43%)	76% (57-89%)	0.43 (0.13-1)	-0.21% (-1.5-1.2%)	-0.011% (-1.4-1.7%)	-0.00099 (-0.12-0.14)	-0.69 (-4.8-3.9)
American Samoa	2000	Female	65 to 69	25% (12-42%)	75% (58-88%)	0.44 (0.14-1)	-0.26% (-1.4-1.1%)	-0.093% (-1.5-1.6%)	-0.0081 (-0.13-0.14)	-0.71 (-4-3.1)
American Samoa	2000	Female	70 to 74	25% (12-41%)	75% (59-88%)	0.42 (0.13-0.96)	-0.38% (-1.4-1%)	-0.32% (-1.6-1.3%)	-0.036 (-0.18-0.15)	-1.1 (-4.1-2.8)
American Samoa	2000	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.39 (0.12-0.94)	-0.22% (-1.3-1.1%)	-0.14% (-1.3-1.4%)	-0.011 (-0.1-0.11)	-0.32 (-1.9-1.6)
American Samoa	2000	Female	80 to 84	23% (9.5-42%)	77% (58-91%)	0.37 (0.12-0.94)	-0.28% (-1.3-0.94%)	-0.27% (-1.4-1.2%)	-0.029 (-0.15-0.12)	-0.4 (-1.8-1.3)
American Samoa	2000	Female	85 to 89	23% (10-41%)	77% (59-90%)	0.38 (0.12-0.91)	-0.33% (-1.3-0.9%)	-0.36% (-1.5-1%)	-0.032 (-0.13-0.097)	-0.27 (-1.1-0.73)
American Samoa	2000	Female	90 to 94	27% (12-46%)	73% (54-88%)	0.44 (0.13-1)	-0.31% (-1.2-1%)	-0.35% (-1.4-1.2%)	-0.015 (-0.063-0.054)	-0.087 (-0.35-0.29)
American Samoa	2000	Female	95 plus	31% (13-55%)	69% (45-87%)	0.52 (0.14-1.3)	-0.48% (-1.8-1.2%)	-0.6% (-2.1-1.2%)	-0.0076 (-0.027-0.016)	-0.024 (-0.092-0.058)
American Samoa	2000	Male	All Ages	38% (30-47%)	62% (53-70%)	0.83 (0.45-1.4)	2.2% (1.1-4%)	2.1% (0.69-4.3%)	3.1 (1-6.4)	180 (86-330)
American Samoa	2000	Male	15 to 19	35% (20-52%)	65% (48-80%)	0.52 (0.2-1)	1.9% (0.95-3.4%)	2.5% (0.98-4.9%)	0.092 (0.037-0.18)	8.6 (4.2-15)
American Samoa	2000	Male	20 to 24	52% (33-70%)	48% (30-67%)	1.4 (0.63-2.7)	5% (2.6-8.7%)	6.4% (2.8-12%)	0.3 (0.13-0.55)	24 (12-42)
American Samoa	2000	Male	25 to 29	60% (40-76%)	40% (24-60%)	1.9 (0.93-3.5)	6.1% (3.3-10%)	7.8% (3.8-14%)	0.38 (0.18-0.66)	29 (16-48)
American Samoa	2000	Male	30 to 34	61% (40-78%)	39% (22-60%)	2 (0.98-3.7)	5.2% (2.7-9%)	6.5% (2.9-12%)	0.37 (0.16-0.69)	27 (14-47)
American Samoa	2000	Male	35 to 39	60% (42-76%)	40% (24-58%)	1.8 (0.82-3.4)	4.5% (2.1-8.5%)	5.7% (2.3-11%)	0.38 (0.15-0.77)	25 (11-47)
American Samoa	2000	Male	40 to 44	57% (40-74%)	43% (26-60%)	1.6 (0.68-3.1)	3.4% (1.2-7.2%)	4.1% (1.2-9%)	0.32 (0.087-0.76)	19 (6.5-41)
American Samoa	2000	Male	45 to 49	53% (32-71%)	47% (29-68%)	1.4 (0.55-2.7)	2.8% (0.83-6.1%)	3.2% (0.63-7.4%)	0.3 (0.062-0.73)	16 (4.5-35)
American Samoa	2000	Male	50 to 54	50% (31-68%)	50% (32-69%)	1.1 (0.45-2.2)	2.3% (0.54-5.4%)	2.6% (0.37-6.3%)	0.3 (0.042-0.73)	13 (2.9-31)
American Samoa	2000	Male	55 to 59	48% (31-66%)	52% (34-69%)	0.93 (0.35-1.9)	1.5% (-0.024-4.4%)	1.7% (-0.17-5.2%)	0.21 (-0.022-0.65)	8.3 (-0.13-24)
American Samoa	2000	Male	60 to 64	45% (26-65%)	55% (35-74%)	0.77 (0.29-1.7)	1.2% (-0.094-3.9%)	1.3% (-0.19-4.6%)	0.19 (-0.029-0.65)	6 (-0.5-20)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	2000	Male	65 to 69	44% (25-63%)	56% (37-75%)	0.7 (0.28-1.5)	0.88% (-0.21-3.2%)	0.97% (-0.33-3.6%)	0.14 (-0.045-0.49)	3.5 (-0.86-12)
American Samoa	2000	Male	70 to 74	41% (24-59%)	59% (41-76%)	0.61 (0.25-1.3)	0.65% (-0.29-2.7%)	0.69% (-0.35-3%)	0.083 (-0.043-0.35)	1.7 (-0.78-7.1)
American Samoa	2000	Male	75 to 79	37% (22-56%)	63% (44-78%)	0.52 (0.19-1.1)	0.49% (-0.26-2.2%)	0.5% (-0.34-2.3%)	0.05 (-0.035-0.23)	0.84 (-0.45-3.6)
American Samoa	2000	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.46 (0.19-0.97)	0.43% (-0.18-1.6%)	0.43% (-0.26-1.8%)	0.031 (-0.019-0.13)	0.38 (-0.16-1.5)
American Samoa	2000	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.44 (0.17-0.92)	0.29% (-0.29-1.3%)	0.27% (-0.39-1.4%)	0.014 (-0.02-0.076)	0.14 (-0.14-0.65)
American Samoa	2000	Male	90 to 94	35% (22-52%)	65% (48-78%)	0.45 (0.16-0.95)	0.19% (-0.37-1.1%)	0.14% (-0.5-1.2%)	0.003 (-0.011-0.025)	0.024 (-0.047-0.15)
American Samoa	2000	Male	95 plus	38% (20-59%)	62% (41-80%)	0.49 (0.16-1.1)	0.094% (-0.59-1.3%)	-0.023% (-0.82-1.3%)	-0.00011 (-0.0039-0.0062)	0.0018 (-0.011-0.025)
American Samoa	2005	Female	All Ages	17% (12-24%)	83% (76-88%)	0.37 (0.19-0.65)	0.55% (-0.037-1.5%)	0.38% (-0.53-1.8%)	0.47 (-0.66-2.3)	37 (-2.5-100)
American Samoa	2005	Female	15 to 19	18% (8-31%)	82% (69-92%)	0.42 (0.14-0.96)	0.96% (0.46-1.9%)	1.6% (0.46-3.7%)	0.022 (0.0066-0.053)	2.8 (1.3-5.6)
American Samoa	2005	Female	20 to 24	22% (9.2-40%)	78% (60-91%)	0.68 (0.23-1.6)	1.7% (0.9-3%)	2.2% (0.71-5%)	0.033 (0.011-0.075)	4.6 (2.4-8.5)
American Samoa	2005	Female	25 to 29	23% (9.4-42%)	77% (58-91%)	0.69 (0.24-1.5)	1.7% (0.96-3.1%)	2.3% (0.89-5%)	0.048 (0.019-0.11)	5.6 (3-10)
American Samoa	2005	Female	30 to 34	24% (8.9-42%)	76% (58-91%)	0.71 (0.24-1.5)	1.5% (0.74-2.8%)	2% (0.68-4.4%)	0.071 (0.023-0.16)	6.8 (3.3-13)
American Samoa	2005	Female	35 to 39	25% (9.6-42%)	75% (58-90%)	0.74 (0.27-1.5)	1.4% (0.57-3%)	2.1% (0.74-4.7%)	0.11 (0.037-0.26)	8.2 (3.2-18)
American Samoa	2005	Female	40 to 44	24% (11-40%)	76% (60-89%)	0.68 (0.25-1.4)	1.1% (0.13-2.9%)	1.8% (0.3-4.4%)	0.11 (0.018-0.28)	6.3 (0.71-16)
American Samoa	2005	Female	45 to 49	23% (10-42%)	77% (58-90%)	0.58 (0.19-1.2)	0.71% (-0.29-2.2%)	1.3% (-0.055-3.4%)	0.098 (-0.004-0.28)	4.1 (-1.6-13)
American Samoa	2005	Female	50 to 54	23% (9.8-41%)	77% (59-90%)	0.51 (0.16-1.1)	0.32% (-0.81-2%)	0.73% (-0.65-3%)	0.059 (-0.053-0.25)	1.6 (-3.9-10)
American Samoa	2005	Female	55 to 59	22% (9.8-42%)	78% (58-90%)	0.46 (0.14-1)	0.12% (-1.1-1.8%)	0.45% (-0.84-2.5%)	0.034 (-0.065-0.19)	0.45 (-4-6.4)
American Samoa	2005	Female	60 to 64	23% (9.7-41%)	77% (59-90%)	0.44 (0.13-1.1)	-0.16% (-1.4-1.4%)	0.034% (-1.4-1.9%)	0.0032 (-0.13-0.18)	-0.6 (-5.3-5.2)
American Samoa	2005	Female	65 to 69	24% (10-42%)	76% (58-90%)	0.45 (0.14-1)	-0.21% (-1.5-1.4%)	-0.042% (-1.6-1.8%)	-0.0036 (-0.14-0.17)	-0.59 (-4.6-3.8)
American Samoa	2005	Female	70 to 74	24% (10-40%)	76% (60-90%)	0.43 (0.13-1.1)	-0.29% (-1.4-1.2%)	-0.22% (-1.5-1.5%)	-0.031 (-0.22-0.23)	-1 (-5-4.3)
American Samoa	2005	Female	75 to 79	22% (9.6-40%)	78% (60-90%)	0.4 (0.11-1)	-0.16% (-1.3-1.1%)	-0.089% (-1.4-1.4%)	-0.0074 (-0.12-0.12)	-0.25 (-2-1.9)
American Samoa	2005	Female	80 to 84	22% (9.3-41%)	78% (59-91%)	0.39 (0.11-0.94)	-0.22% (-1.3-1%)	-0.2% (-1.4-1.3%)	-0.023 (-0.16-0.15)	-0.33 (-1.9-1.6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	2005	Female	85 to 89	23% (9.5-41%)	77% (59-91%)	0.41 (0.12-1)	-0.29% (-1.4-1%)	-0.32% (-1.5-1.2%)	-0.03 (-0.15-0.11)	-0.25 (-1.2-0.86)
American Samoa	2005	Female	90 to 94	26% (12-45%)	74% (55-88%)	0.47 (0.14-1.1)	-0.28% (-1.3-1.2%)	-0.32% (-1.5-1.3%)	-0.016 (-0.074-0.066)	-0.088 (-0.41-0.37)
American Samoa	2005	Female	95 plus	31% (13-53%)	69% (47-87%)	0.54 (0.15-1.3)	-0.45% (-1.8-1.3%)	-0.58% (-2.1-1.3%)	-0.0095 (-0.035-0.022)	-0.03 (-0.12-0.084)
American Samoa	2005	Male	All Ages	37% (29-46%)	63% (54-71%)	0.86 (0.48-1.5)	2.4% (1.1-4.4%)	2.2% (0.75-4.7%)	3.6 (1.2-7.8)	210 (98-400)
American Samoa	2005	Male	15 to 19	33% (19-51%)	67% (49-81%)	0.53 (0.19-1.1)	1.9% (0.95-3.4%)	2.5% (0.95-4.9%)	0.097 (0.037-0.19)	9.1 (4.4-16)
American Samoa	2005	Male	20 to 24	51% (33-69%)	49% (31-67%)	1.4 (0.64-2.8)	5.2% (2.7-9%)	6.5% (3-12%)	0.3 (0.14-0.57)	25 (13-43)
American Samoa	2005	Male	25 to 29	59% (40-76%)	41% (24-60%)	2 (0.97-3.7)	5.6% (3-9.4%)	6.7% (3.2-12%)	0.41 (0.19-0.74)	32 (17-54)
American Samoa	2005	Male	30 to 34	60% (39-77%)	40% (23-61%)	2.2 (1-4)	5.5% (2.8-9.3%)	6.7% (3.1-12%)	0.42 (0.19-0.79)	31 (16-54)
American Samoa	2005	Male	35 to 39	59% (40-76%)	41% (24-60%)	1.9 (0.89-3.6)	4.7% (2.2-8.5%)	5.9% (2.5-11%)	0.48 (0.19-0.96)	31 (14-59)
American Samoa	2005	Male	40 to 44	56% (38-73%)	44% (27-62%)	1.6 (0.7-3.1)	3.5% (1.2-7.2%)	4.2% (1.2-9%)	0.39 (0.11-0.87)	23 (8.1-48)
American Samoa	2005	Male	45 to 49	52% (31-70%)	48% (30-69%)	1.4 (0.58-2.7)	2.7% (0.65-5.8%)	3% (0.65-6.8%)	0.36 (0.069-0.82)	19 (4.5-41)
American Samoa	2005	Male	50 to 54	49% (30-68%)	51% (32-70%)	1.1 (0.45-2.2)	2.4% (0.46-5.4%)	2.7% (0.38-6.4%)	0.34 (0.046-0.87)	16 (2.9-37)
American Samoa	2005	Male	55 to 59	47% (29-66%)	53% (34-71%)	0.94 (0.36-2)	1.6% (0.075-4.3%)	1.7% (-0.038-4.9%)	0.26 (-0.0059-0.74)	10 (0.45-27)
American Samoa	2005	Male	60 to 64	44% (26-65%)	56% (35-74%)	0.79 (0.3-1.7)	1.2% (-0.097-3.9%)	1.3% (-0.21-4.4%)	0.23 (-0.036-0.76)	7.2 (-0.59-23)
American Samoa	2005	Male	65 to 69	42% (24-61%)	58% (39-76%)	0.71 (0.29-1.6)	0.95% (-0.18-3.4%)	1% (-0.25-3.8%)	0.16 (-0.038-0.57)	4.2 (-0.76-14)
American Samoa	2005	Male	70 to 74	40% (23-57%)	60% (43-77%)	0.62 (0.24-1.4)	0.68% (-0.23-2.6%)	0.73% (-0.29-2.9%)	0.087 (-0.035-0.34)	1.8 (-0.61-7)
American Samoa	2005	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.52 (0.19-1.1)	0.52% (-0.22-2.1%)	0.53% (-0.3-2.3%)	0.051 (-0.028-0.21)	0.86 (-0.36-3.3)
American Samoa	2005	Male	80 to 84	33% (18-51%)	67% (49-82%)	0.46 (0.17-0.93)	0.46% (-0.15-1.7%)	0.45% (-0.24-1.8%)	0.03 (-0.016-0.12)	0.38 (-0.12-1.4)
American Samoa	2005	Male	85 to 89	32% (18-48%)	68% (52-82%)	0.44 (0.17-0.9)	0.32% (-0.26-1.5%)	0.3% (-0.36-1.6%)	0.015 (-0.018-0.079)	0.14 (-0.12-0.65)
American Samoa	2005	Male	90 to 94	33% (19-50%)	67% (50-81%)	0.45 (0.16-0.95)	0.22% (-0.31-1.2%)	0.18% (-0.4-1.3%)	0.0043 (-0.0099-0.032)	0.033 (-0.047-0.19)
American Samoa	2005	Male	95 plus	36% (19-57%)	64% (43-81%)	0.49 (0.16-1.1)	0.15% (-0.53-1.4%)	0.037% (-0.76-1.4%)	0.00024 (-0.0047-0.0093)	0.0039 (-0.014-0.037)
American Samoa	2010	Female	All Ages	18% (12-24%)	82% (76-88%)	0.4 (0.21-0.7)	0.63% (-0.029-1.7%)	0.51% (-0.47-2.1%)	0.69 (-0.65-2.9)	48 (-2.1-120)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	2010	Female	15 to 19	18% (8.1-33%)	82% (67-92%)	0.46 (0.14-1.1)	1% (0.47-2%)	1.7% (0.47-4%)	0.023 (0.0065-0.057)	3 (1.3-5.9)
American Samoa	2010	Female	20 to 24	22% (9-39%)	78% (61-91%)	0.71 (0.24-1.7)	1.7% (0.9-3.1%)	2.3% (0.72-5.5%)	0.032 (0.0098-0.077)	4.5 (2.3-8.2)
American Samoa	2010	Female	25 to 29	23% (9.2-43%)	77% (57-91%)	0.72 (0.25-1.5)	1.8% (0.96-3.1%)	2.3% (0.92-5%)	0.048 (0.018-0.1)	5.6 (3-10)
American Samoa	2010	Female	30 to 34	24% (9-44%)	76% (56-91%)	0.74 (0.24-1.6)	1.6% (0.73-3%)	2.1% (0.7-4.7%)	0.082 (0.027-0.18)	7.9 (3.8-15)
American Samoa	2010	Female	35 to 39	24% (9.1-42%)	76% (58-91%)	0.76 (0.27-1.6)	1.5% (0.6-3%)	2.2% (0.79-4.7%)	0.15 (0.049-0.32)	11 (4.2-22)
American Samoa	2010	Female	40 to 44	23% (11-41%)	77% (59-89%)	0.7 (0.26-1.6)	1.2% (0.2-3%)	1.9% (0.31-4.6%)	0.15 (0.025-0.38)	8.4 (1.3-22)
American Samoa	2010	Female	45 to 49	23% (9.7-41%)	77% (59-90%)	0.61 (0.21-1.4)	0.83% (-0.24-2.6%)	1.4% (-0.016-4%)	0.14 (-0.0015-0.39)	6.1 (-1.6-19)
American Samoa	2010	Female	50 to 54	23% (10-41%)	77% (59-90%)	0.54 (0.17-1.2)	0.43% (-0.7-2.2%)	0.87% (-0.54-3.2%)	0.085 (-0.051-0.31)	2.6 (-4.2-13)
American Samoa	2010	Female	55 to 59	22% (10-41%)	78% (59-90%)	0.48 (0.16-1.1)	0.21% (-0.97-1.8%)	0.57% (-0.72-2.6%)	0.048 (-0.061-0.23)	0.88 (-4-7.9)
American Samoa	2010	Female	60 to 64	22% (10-40%)	78% (60-90%)	0.46 (0.15-1.1)	-0.1% (-1.4-1.5%)	0.11% (-1.4-2.1%)	0.011 (-0.14-0.22)	-0.42 (-5.9-6.5)
American Samoa	2010	Female	65 to 69	23% (11-41%)	77% (59-89%)	0.47 (0.14-1.1)	-0.12% (-1.5-1.6%)	0.072% (-1.5-2.1%)	0.0071 (-0.14-0.2)	-0.34 (-4.5-5)
American Samoa	2010	Female	70 to 74	23% (10-40%)	77% (60-90%)	0.45 (0.13-1.1)	-0.19% (-1.3-1.4%)	-0.094% (-1.4-1.9%)	-0.017 (-0.25-0.33)	-0.86 (-5.9-6.4)
American Samoa	2010	Female	75 to 79	22% (9.2-40%)	78% (60-91%)	0.42 (0.12-0.99)	-0.08% (-1.3-1.4%)	0.0085% (-1.3-1.7%)	0.001 (-0.12-0.16)	-0.13 (-2.1-2.4)
American Samoa	2010	Female	80 to 84	21% (8.6-40%)	79% (60-91%)	0.4 (0.11-0.97)	-0.15% (-1.3-1.3%)	-0.12% (-1.4-1.6%)	-0.015 (-0.16-0.19)	-0.24 (-2-2)
American Samoa	2010	Female	85 to 89	22% (9.6-40%)	78% (60-90%)	0.42 (0.13-1)	-0.21% (-1.3-1.3%)	-0.23% (-1.5-1.5%)	-0.023 (-0.15-0.15)	-0.2 (-1.2-1.2)
American Samoa	2010	Female	90 to 94	25% (11-44%)	75% (56-89%)	0.47 (0.15-1.1)	-0.21% (-1.3-1.3%)	-0.24% (-1.4-1.5%)	-0.013 (-0.079-0.078)	-0.07 (-0.43-0.42)
American Samoa	2010	Female	95 plus	30% (13-52%)	70% (48-87%)	0.55 (0.16-1.4)	-0.35% (-1.8-1.5%)	-0.47% (-2.1-1.5%)	-0.0089 (-0.039-0.029)	-0.027 (-0.13-0.11)
American Samoa	2010	Male	All Ages	37% (29-46%)	63% (54-71%)	0.91 (0.49-1.6)	2.6% (1.2-4.6%)	2.5% (0.84-5.1%)	4 (1.4-8.5)	230 (110-420)
American Samoa	2010	Male	15 to 19	33% (18-51%)	67% (49-82%)	0.54 (0.19-1.1)	1.9% (0.97-3.5%)	2.5% (0.98-5.1%)	0.1 (0.038-0.2)	9.5 (4.7-17)
American Samoa	2010	Male	20 to 24	51% (32-68%)	49% (32-68%)	1.5 (0.65-3)	5.3% (2.8-9.2%)	6.7% (3.1-12%)	0.29 (0.13-0.54)	24 (12-43)
American Samoa	2010	Male	25 to 29	58% (38-75%)	42% (25-62%)	2.1 (0.98-3.9)	6.4% (3.5-10%)	8.2% (4-14%)	0.42 (0.2-0.73)	33 (18-54)
American Samoa	2010	Male	30 to 34	59% (38-76%)	41% (24-62%)	2.2 (1-4.1)	5.5% (2.8-9.4%)	6.9% (3.1-12%)	0.44 (0.2-0.81)	33 (17-57)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	2010	Male	35 to 39	58% (39-76%)	42% (24-61%)	1.9 (0.91-3.6)	4.7% (2.2-8.4%)	5.9% (2.6-11%)	0.55 (0.23-1.1)	37 (17-65)
American Samoa	2010	Male	40 to 44	55% (38-74%)	45% (26-62%)	1.7 (0.77-3.2)	3.6% (1.3-7.3%)	4.3% (1.4-9%)	0.45 (0.14-0.98)	27 (9.9-56)
American Samoa	2010	Male	45 to 49	52% (31-71%)	48% (29-69%)	1.4 (0.61-2.8)	3% (0.86-6.4%)	3.4% (0.84-7.7%)	0.41 (0.1-0.94)	22 (6.2-47)
American Samoa	2010	Male	50 to 54	49% (30-68%)	51% (32-70%)	1.2 (0.48-2.3)	2.5% (0.57-5.8%)	2.9% (0.46-6.9%)	0.4 (0.066-0.96)	18 (3.9-42)
American Samoa	2010	Male	55 to 59	46% (29-66%)	54% (34-71%)	0.98 (0.37-1.9)	1.7% (0.14-4.5%)	1.9% (0.011-5.1%)	0.31 (0.0017-0.9)	12 (1.1-33)
American Samoa	2010	Male	60 to 64	44% (25-63%)	56% (37-75%)	0.81 (0.31-1.8)	1.3% (-0.036-3.8%)	1.5% (-0.1-4.4%)	0.28 (-0.021-0.87)	8.8 (-0.24-26)
American Samoa	2010	Male	65 to 69	42% (24-61%)	58% (39-76%)	0.74 (0.29-1.6)	1.1% (-0.046-3.6%)	1.2% (-0.093-4.1%)	0.2 (-0.015-0.69)	5.1 (-0.21-17)
American Samoa	2010	Male	70 to 74	40% (23-58%)	60% (42-77%)	0.64 (0.23-1.4)	0.78% (-0.16-2.7%)	0.84% (-0.21-3%)	0.096 (-0.025-0.35)	2 (-0.41-7.1)
American Samoa	2010	Male	75 to 79	36% (21-54%)	64% (46-79%)	0.54 (0.2-1.2)	0.59% (-0.17-2.2%)	0.61% (-0.26-2.4%)	0.053 (-0.022-0.22)	0.9 (-0.27-3.4)
American Samoa	2010	Male	80 to 84	33% (19-51%)	67% (49-81%)	0.48 (0.19-0.98)	0.51% (-0.12-1.9%)	0.52% (-0.2-2%)	0.031 (-0.012-0.12)	0.39 (-0.091-1.4)
American Samoa	2010	Male	85 to 89	32% (18-48%)	68% (52-82%)	0.45 (0.17-0.93)	0.37% (-0.23-1.7%)	0.35% (-0.33-1.8%)	0.017 (-0.016-0.084)	0.16 (-0.099-0.71)
American Samoa	2010	Male	90 to 94	33% (19-50%)	67% (50-81%)	0.46 (0.16-1)	0.26% (-0.27-1.4%)	0.22% (-0.37-1.4%)	0.0052 (-0.009-0.034)	0.039 (-0.04-0.2)
American Samoa	2010	Male	95 plus	36% (19-57%)	64% (43-81%)	0.5 (0.16-1.2)	0.21% (-0.45-1.5%)	0.094% (-0.67-1.6%)	0.0069 (-0.005-0.012)	0.062 (-0.014-0.047)
American Samoa	2016	Female	All Ages	18% (13-25%)	82% (75-87%)	0.41 (0.22-0.73)	0.67% (-0.0099-1.7%)	0.55% (-0.47-2.2%)	0.9 (-0.75-3.6)	60 (-0.79-150)
American Samoa	2016	Female	15 to 19	18% (7.7-33%)	82% (67-92%)	0.46 (0.15-1.1)	1% (0.48-2%)	1.7% (0.51-4.1%)	0.025 (0.0072-0.06)	3.2 (1.5-6.4)
American Samoa	2016	Female	20 to 24	22% (8.9-39%)	78% (61-91%)	0.73 (0.23-1.7)	1.8% (0.97-3.2%)	2.4% (0.76-5.5%)	0.032 (0.0094-0.079)	4.4 (2.2-8.2)
American Samoa	2016	Female	25 to 29	23% (9-41%)	77% (59-91%)	0.74 (0.24-1.6)	1.8% (1-3.2%)	2.4% (0.96-5.2%)	0.05 (0.019-0.11)	5.8 (3.1-11)
American Samoa	2016	Female	30 to 34	24% (8.7-45%)	76% (55-91%)	0.76 (0.25-1.6)	1.6% (0.77-3%)	2.2% (0.72-5.1%)	0.1 (0.033-0.23)	9.7 (4.5-19)
American Samoa	2016	Female	35 to 39	25% (9.7-43%)	75% (57-90%)	0.78 (0.28-1.7)	1.6% (0.67-3.2%)	2.3% (0.87-5%)	0.21 (0.071-0.46)	15 (5.9-31)
American Samoa	2016	Female	40 to 44	24% (12-41%)	76% (59-88%)	0.72 (0.28-1.5)	1.3% (0.21-3%)	1.9% (0.34-4.7%)	0.2 (0.034-0.49)	12 (1.9-28)
American Samoa	2016	Female	45 to 49	23% (9.9-41%)	77% (59-90%)	0.61 (0.21-1.3)	0.83% (-0.21-2.6%)	1.4% (-0.021-3.8%)	0.2 (-0.0026-0.55)	8.6 (-2.2-27)
American Samoa	2016	Female	50 to 54	23% (10-42%)	77% (58-90%)	0.55 (0.18-1.2)	0.43% (-0.79-2.3%)	0.85% (-0.58-3.4%)	0.11 (-0.08-0.46)	3.3 (-6-19)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	2016	Female	55 to 59	22% (10-41%)	78% (59-90%)	0.49 (0.16-1.1)	0.22% (-0.99-1.9%)	0.57% (-0.85-2.7%)	0.06 (-0.088-0.3)	1.1 (-5.1-9.8)
American Samoa	2016	Female	60 to 64	23% (10-41%)	77% (59-90%)	0.47 (0.16-1.1)	-0.087% (-1.4-1.6%)	0.11% (-1.4-2.3%)	0.014 (-0.18-0.29)	-0.41 (-7.4-7.9)
American Samoa	2016	Female	65 to 69	24% (11-42%)	76% (58-89%)	0.49 (0.15-1.1)	-0.11% (-1.5-1.6%)	0.064% (-1.5-2.2%)	0.0068 (-0.17-0.23)	-0.38 (-5.5-5.5)
American Samoa	2016	Female	70 to 74	24% (11-41%)	76% (59-89%)	0.47 (0.14-1.1)	-0.2% (-1.4-1.4%)	-0.11% (-1.5-1.8%)	-0.028 (-0.4-0.46)	-1.2 (-8.9-9)
American Samoa	2016	Female	75 to 79	22% (9.8-40%)	78% (60-90%)	0.43 (0.13-1)	-0.081% (-1.2-1.4%)	0.0007% (-1.3-1.7%)	-0.00022 (-0.13-0.18)	-0.15 (-2.3-2.5)
American Samoa	2016	Female	80 to 84	22% (9.3-40%)	78% (60-91%)	0.41 (0.12-0.99)	-0.15% (-1.3-1.4%)	-0.13% (-1.4-1.7%)	-0.018 (-0.19-0.24)	-0.27 (-2.3-2.5)
American Samoa	2016	Female	85 to 89	23% (9.9-40%)	77% (60-90%)	0.43 (0.13-1.1)	-0.21% (-1.3-1.4%)	-0.24% (-1.5-1.6%)	-0.027 (-0.17-0.17)	-0.22 (-1.3-1.4)
American Samoa	2016	Female	90 to 94	26% (11-46%)	74% (54-89%)	0.5 (0.15-1.2)	-0.21% (-1.3-1.5%)	-0.25% (-1.5-1.7%)	-0.016 (-0.092-0.1)	-0.08 (-0.49-0.56)
American Samoa	2016	Female	95 plus	31% (12-53%)	69% (47-88%)	0.59 (0.17-1.5)	-0.39% (-1.9-1.7%)	-0.51% (-2.3-1.8%)	-0.011 (-0.048-0.038)	-0.032 (-0.16-0.14)
American Samoa	2016	Male	All Ages	37% (30-47%)	63% (53-70%)	0.95 (0.5-1.6)	2.7% (1.3-4.8%)	2.7% (0.93-5.2%)	4.8 (1.6-10)	270 (120-500)
American Samoa	2016	Male	15 to 19	33% (19-51%)	67% (49-81%)	0.57 (0.2-1.2)	1.9% (0.95-3.5%)	2.6% (1-5.3%)	0.1 (0.039-0.21)	9.9 (4.9-18)
American Samoa	2016	Male	20 to 24	51% (32-69%)	49% (31-68%)	1.6 (0.68-3.1)	5.4% (2.8-9.3%)	6.9% (3-13%)	0.28 (0.12-0.53)	24 (12-41)
American Samoa	2016	Male	25 to 29	59% (38-75%)	41% (25-62%)	2.2 (1-4.1)	6.4% (3.4-10%)	8.2% (3.8-14%)	0.43 (0.19-0.76)	34 (18-57)
American Samoa	2016	Male	30 to 34	60% (38-77%)	40% (23-62%)	2.3 (1.1-4.3)	5.6% (2.9-9.3%)	7% (3.2-13%)	0.49 (0.2-0.94)	37 (18-66)
American Samoa	2016	Male	35 to 39	58% (40-76%)	42% (24-60%)	2 (0.99-3.7)	4.9% (2.3-8.4%)	6.1% (2.7-11%)	0.71 (0.28-1.4)	47 (22-85)
American Samoa	2016	Male	40 to 44	56% (37-74%)	44% (26-63%)	1.7 (0.77-3.3)	3.7% (1.4-7%)	4.4% (1.5-9%)	0.57 (0.18-1.2)	35 (13-68)
American Samoa	2016	Male	45 to 49	51% (30-70%)	49% (30-70%)	1.5 (0.63-2.8)	3.1% (1-6.4%)	3.5% (0.98-7.6%)	0.52 (0.14-1.2)	28 (8.9-60)
American Samoa	2016	Male	50 to 54	48% (29-68%)	52% (32-71%)	1.2 (0.49-2.5)	2.7% (0.7-6%)	3.1% (0.64-7.2%)	0.49 (0.11-1.2)	22 (5.7-52)
American Samoa	2016	Male	55 to 59	46% (28-66%)	54% (34-72%)	1 (0.39-2.1)	1.8% (0.16-4.8%)	2% (0.094-5.4%)	0.41 (0.018-1.1)	16 (1.4-42)
American Samoa	2016	Male	60 to 64	44% (26-65%)	56% (35-74%)	0.83 (0.33-1.8)	1.4% (-0.0083-4%)	1.5% (-0.076-4.6%)	0.35 (-0.018-1.1)	11 (-0.069-33)
American Samoa	2016	Male	65 to 69	42% (25-61%)	58% (39-75%)	0.74 (0.29-1.6)	1.1% (-0.0074-3.5%)	1.3% (-0.05-4%)	0.23 (-0.0094-0.75)	6 (-0.039-19)
American Samoa	2016	Male	70 to 74	39% (23-57%)	61% (43-77%)	0.66 (0.25-1.4)	0.83% (-0.13-2.9%)	0.9% (-0.17-3.2%)	0.1 (-0.02-0.37)	2.2 (-0.34-7.5)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
American Samoa	2016	Male	75 to 79	36% (21-55%)	64% (45-79%)	0.56 (0.21-1.2)	0.63% (-0.14-2.3%)	0.65% (-0.23-2.6%)	0.054 (-0.019-0.21)	0.91 (-0.21-3.4)
American Samoa	2016	Male	80 to 84	33% (18-51%)	67% (49-82%)	0.5 (0.19-1)	0.57% (-0.092-2%)	0.57% (-0.17-2.1%)	0.032 (-0.0091-0.12)	0.4 (-0.067-1.4)
American Samoa	2016	Male	85 to 89	31% (17-47%)	69% (53-83%)	0.47 (0.18-0.98)	0.42% (-0.2-1.7%)	0.4% (-0.29-1.9%)	0.018 (-0.013-0.084)	0.17 (-0.085-0.7)
American Samoa	2016	Male	90 to 94	33% (20-51%)	67% (49-80%)	0.49 (0.18-1.1)	0.31% (-0.24-1.5%)	0.26% (-0.36-1.6%)	0.0063 (-0.0085-0.038)	0.045 (-0.036-0.22)
American Samoa	2016	Male	95 plus	37% (19-58%)	63% (42-81%)	0.54 (0.18-1.3)	0.26% (-0.45-1.8%)	0.14% (-0.66-1.9%)	0.0011 (-0.0051-0.015)	0.008 (-0.014-0.058)
Bermuda	1990	Female	All Ages	32% (26-39%)	68% (61-74%)	2.6 (2.3-2.9)	5.6% (4.1-7.4%)	7.4% (4.8-11%)	15 (9.7-22)	380 (280-510)
Bermuda	1990	Female	15 to 19	37% (24-53%)	63% (47-76%)	3 (1.6-4.9)	4.1% (2.6-5.9%)	10% (6-14%)	0.14 (0.082-0.2)	13 (8.3-19)
Bermuda	1990	Female	20 to 24	43% (28-60%)	57% (40-72%)	4.5 (2.6-6.9)	4.7% (3.4-6.1%)	12% (8.4-16%)	0.12 (0.083-0.16)	15 (10-20)
Bermuda	1990	Female	25 to 29	43% (27-60%)	57% (40-73%)	4.7 (2.8-7.1)	5.9% (4.4-7.7%)	12% (8.3-15%)	0.24 (0.17-0.32)	21 (16-27)
Bermuda	1990	Female	30 to 34	41% (25-59%)	59% (41-75%)	4.5 (2.6-7)	6.5% (4.7-8.7%)	12% (8.2-16%)	0.31 (0.21-0.43)	22 (16-30)
Bermuda	1990	Female	35 to 39	40% (26-58%)	60% (42-74%)	4.2 (2.4-6.7)	5.9% (4.1-8.2%)	11% (7.7-16%)	0.28 (0.19-0.42)	19 (13-26)
Bermuda	1990	Female	40 to 44	39% (25-57%)	61% (43-75%)	3.9 (2.1-6.4)	7% (4.8-10%)	13% (8.4-18%)	0.42 (0.27-0.63)	23 (15-33)
Bermuda	1990	Female	45 to 49	37% (22-54%)	63% (46-78%)	3.8 (1.9-6.5)	7.3% (4.7-11%)	12% (8-18%)	0.51 (0.33-0.77)	24 (16-36)
Bermuda	1990	Female	50 to 54	34% (20-50%)	66% (50-80%)	3.4 (1.8-5.8)	8.1% (5.1-12%)	12% (7.5-17%)	0.79 (0.5-1.2)	31 (20-47)
Bermuda	1990	Female	55 to 59	30% (18-46%)	70% (54-82%)	2.9 (1.5-5)	8.1% (4.9-12%)	11% (6.6-16%)	1.1 (0.66-1.6)	36 (22-55)
Bermuda	1990	Female	60 to 64	26% (16-41%)	74% (59-84%)	2.5 (1.2-4.4)	7.4% (4.2-12%)	9.2% (5.2-14%)	1.3 (0.75-2.2)	38 (21-61)
Bermuda	1990	Female	65 to 69	24% (15-37%)	76% (63-85%)	2.1 (1.1-3.5)	6.9% (4-11%)	8.4% (4.9-13%)	1.5 (0.86-2.4)	35 (20-56)
Bermuda	1990	Female	70 to 74	22% (13-35%)	78% (65-87%)	1.8 (0.87-3.2)	6.1% (3.1-10%)	7.2% (3.7-12%)	1.7 (0.84-2.8)	31 (16-52)
Bermuda	1990	Female	75 to 79	19% (11-32%)	81% (68-89%)	1.5 (0.68-2.8)	5.4% (2.4-9.4%)	6.2% (2.7-11%)	1.8 (0.79-3.2)	26 (11-46)
Bermuda	1990	Female	80 to 84	18% (9.8-29%)	82% (71-90%)	1.4 (0.68-2.5)	5.5% (2.5-9.4%)	6.1% (2.7-11%)	2.2 (1-3.9)	23 (11-40)
Bermuda	1990	Female	85 to 89	18% (10-28%)	82% (72-90%)	1.5 (0.67-2.6)	5.4% (2.4-9.2%)	6% (2.7-10%)	1.7 (0.72-2.9)	12 (5.6-21)
Bermuda	1990	Female	90 to 94	20% (11-31%)	80% (69-89%)	1.7 (0.75-3.1)	5.6% (2.4-10%)	6.3% (2.6-11%)	0.79 (0.32-1.5)	4 (1.7-7.3)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	1990	Female	95 plus	23% (12-40%)	77% (60-88%)	2 (0.73-4.1)	6% (2-11%)	6.7% (2.1-13%)	0.25 (0.077-0.48)	0.87 (0.28-1.6)
Bermuda	1990	Male	All Ages	54% (46-63%)	46% (37-54%)	5.8 (5.3-6.3)	12% (9.3-15%)	13% (9.5-16%)	31 (23-41)	1000 (800-1300)
Bermuda	1990	Male	15 to 19	55% (37-72%)	45% (28-63%)	4.5 (2.5-7.1)	9.1% (5.8-13%)	16% (9.7-23%)	0.36 (0.22-0.53)	32 (21-46)
Bermuda	1990	Male	20 to 24	66% (48-80%)	34% (20-52%)	8.6 (5.5-12)	16% (11-21%)	23% (15-31%)	0.85 (0.54-1.2)	70 (48-92)
Bermuda	1990	Male	25 to 29	69% (50-82%)	31% (18-50%)	10 (6.8-14)	16% (11-20%)	22% (14-29%)	0.95 (0.63-1.3)	73 (52-93)
Bermuda	1990	Male	30 to 34	70% (50-83%)	30% (17-50%)	11 (6.9-15)	15% (11-19%)	20% (14-26%)	1 (0.74-1.4)	70 (51-89)
Bermuda	1990	Male	35 to 39	69% (53-83%)	31% (17-47%)	10 (6.4-15)	16% (12-20%)	19% (15-24%)	1.7 (1.2-2.2)	95 (72-120)
Bermuda	1990	Male	40 to 44	68% (52-82%)	32% (18-48%)	9.7 (5.6-15)	15% (12-19%)	19% (14-24%)	1.7 (1.3-2.1)	85 (64-110)
Bermuda	1990	Male	45 to 49	66% (46-81%)	34% (19-54%)	9.1 (5.1-15)	15% (12-19%)	18% (14-23%)	2 (1.5-2.5)	88 (66-110)
Bermuda	1990	Male	50 to 54	65% (46-81%)	35% (19-54%)	8.4 (4.8-14)	15% (11-19%)	17% (12-22%)	2.6 (1.9-3.5)	100 (74-130)
Bermuda	1990	Male	55 to 59	63% (44-79%)	37% (21-56%)	7.7 (4.2-12)	14% (10-19%)	16% (12-21%)	2.9 (2.1-3.9)	96 (69-130)
Bermuda	1990	Male	60 to 64	61% (40-78%)	39% (22-60%)	6.9 (3.6-12)	13% (8.5-17%)	14% (9.5-20%)	3.4 (2.3-4.8)	96 (65-130)
Bermuda	1990	Male	65 to 69	59% (40-76%)	41% (24-60%)	6.1 (3.3-9.9)	12% (7.9-17%)	13% (8.7-19%)	3.5 (2.3-5.1)	82 (53-120)
Bermuda	1990	Male	70 to 74	55% (37-71%)	45% (29-63%)	5 (2.6-8.6)	10% (6.4-15%)	11% (7-16%)	3.3 (2-4.7)	61 (39-88)
Bermuda	1990	Male	75 to 79	50% (31-69%)	50% (31-69%)	4.2 (2-7.1)	8.6% (5.5-13%)	9.5% (5.9-14%)	3 (1.8-4.3)	43 (26-62)
Bermuda	1990	Male	80 to 84	45% (27-62%)	55% (38-73%)	3.5 (1.7-5.6)	7.5% (4.5-11%)	8.3% (5-13%)	2.2 (1.3-3.2)	23 (14-35)
Bermuda	1990	Male	85 to 89	40% (24-56%)	60% (44-76%)	3.1 (1.5-5.2)	6.7% (4.1-10%)	7.4% (4.4-12%)	1.2 (0.73-1.9)	9.5 (5.7-14)
Bermuda	1990	Male	90 to 94	40% (25-57%)	60% (43-75%)	3.1 (1.5-5.6)	6.3% (3.7-9.7%)	6.9% (3.8-11%)	0.44 (0.25-0.71)	2.4 (1.4-3.7)
Bermuda	1990	Male	95 plus	44% (25-64%)	56% (36-75%)	3.4 (1.4-6.4)	6.6% (3.3-11%)	7.2% (3.4-12%)	0.1 (0.049-0.17)	0.37 (0.19-0.6)
Bermuda	1995	Female	All Ages	31% (25-38%)	69% (62-75%)	2.8 (2.5-3.1)	5.5% (4-7.4%)	7.5% (4.9-11%)	16 (11-23)	390 (280-520)
Bermuda	1995	Female	15 to 19	37% (24-53%)	63% (47-76%)	3.2 (1.6-5.1)	3.8% (2.4-5.4%)	9.8% (5.8-14%)	0.11 (0.066-0.16)	11 (7.1-16)
Bermuda	1995	Female	20 to 24	43% (27-59%)	57% (41-73%)	4.8 (2.7-7.2)	3.9% (2.8-5.1%)	9.2% (6.4-12%)	0.07 (0.047-0.099)	11 (7.9-15)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	1995	Female	25 to 29	42% (26-59%)	58% (41-74%)	4.9 (3-7.2)	4.7% (3.6-6.1%)	9.2% (6.6-12%)	0.14 (0.098-0.2)	15 (11-21)
Bermuda	1995	Female	30 to 34	40% (24-58%)	60% (42-76%)	4.7 (2.9-7.1)	5.8% (4.1-7.9%)	11% (7.2-15%)	0.31 (0.2-0.45)	23 (16-32)
Bermuda	1995	Female	35 to 39	40% (25-57%)	60% (43-75%)	4.4 (2.5-7)	6.2% (4.2-8.8%)	11% (7.1-16%)	0.37 (0.23-0.53)	23 (16-33)
Bermuda	1995	Female	40 to 44	38% (24-56%)	62% (44-76%)	4.1 (2.2-6.7)	6.9% (4.6-9.9%)	12% (7.8-18%)	0.49 (0.31-0.74)	27 (17-39)
Bermuda	1995	Female	45 to 49	36% (22-53%)	64% (47-78%)	4 (2.1-6.9)	7.3% (4.6-11%)	12% (7.5-18%)	0.57 (0.35-0.88)	27 (17-41)
Bermuda	1995	Female	50 to 54	33% (20-49%)	67% (51-80%)	3.6 (1.9-6)	7.8% (4.9-12%)	11% (7.2-17%)	0.78 (0.48-1.2)	31 (20-47)
Bermuda	1995	Female	55 to 59	29% (17-45%)	71% (55-83%)	3 (1.5-5.1)	7.7% (4.7-11%)	10% (6.4-16%)	1 (0.63-1.6)	36 (22-55)
Bermuda	1995	Female	60 to 64	26% (15-41%)	74% (59-85%)	2.5 (1.3-4.5)	7.1% (4.2-11%)	9% (5.3-14%)	1.2 (0.72-2)	36 (21-56)
Bermuda	1995	Female	65 to 69	23% (14-37%)	77% (63-86%)	2.2 (1.2-3.6)	6.9% (4.1-11%)	8.5% (5.1-13%)	1.6 (0.95-2.4)	37 (22-57)
Bermuda	1995	Female	70 to 74	21% (13-34%)	79% (66-87%)	1.9 (0.94-3.4)	6.2% (3.4-10%)	7.4% (4.1-12%)	1.8 (0.97-2.9)	33 (18-54)
Bermuda	1995	Female	75 to 79	19% (11-31%)	81% (69-89%)	1.6 (0.74-3)	5.5% (2.7-9.2%)	6.4% (3.1-11%)	1.8 (0.86-3.2)	27 (12-45)
Bermuda	1995	Female	80 to 84	17% (9.8-27%)	83% (73-90%)	1.5 (0.73-2.6)	5.6% (2.7-9.2%)	6.3% (3-11%)	2.3 (1.1-3.9)	24 (12-41)
Bermuda	1995	Female	85 to 89	17% (9.9-27%)	83% (73-90%)	1.6 (0.72-2.8)	5.7% (2.8-9.7%)	6.3% (3-11%)	2.2 (1.1-3.9)	17 (8-29)
Bermuda	1995	Female	90 to 94	19% (11-30%)	81% (70-89%)	1.8 (0.77-3.3)	5.8% (2.7-10%)	6.5% (2.9-12%)	1.1 (0.49-2)	5.7 (2.5-10)
Bermuda	1995	Female	95 plus	23% (12-39%)	77% (61-88%)	2.1 (0.75-4.3)	6.2% (2.3-12%)	6.9% (2.4-13%)	0.39 (0.14-0.74)	1.4 (0.5-2.5)
Bermuda	1995	Male	All Ages	54% (45-62%)	46% (38-55%)	6.1 (5.7-6.6)	11% (8.4-13%)	12% (8.8-15%)	28 (21-37)	890 (690-1100)
Bermuda	1995	Male	15 to 19	54% (36-71%)	46% (29-64%)	4.7 (2.6-7.5)	8% (5.4-11%)	15% (9.4-22%)	0.23 (0.14-0.34)	23 (15-32)
Bermuda	1995	Male	20 to 24	65% (47-80%)	35% (20-53%)	9 (5.7-13)	14% (10-18%)	20% (13-27%)	0.6 (0.38-0.82)	53 (37-70)
Bermuda	1995	Male	25 to 29	68% (49-82%)	32% (18-51%)	11 (7.2-15)	12% (9-16%)	17% (11-23%)	0.58 (0.37-0.79)	50 (35-65)
Bermuda	1995	Male	30 to 34	69% (49-83%)	31% (17-51%)	11 (7.4-16)	13% (9.4-16%)	17% (12-21%)	0.88 (0.62-1.2)	62 (46-80)
Bermuda	1995	Male	35 to 39	69% (52-83%)	31% (17-48%)	11 (6.7-16)	13% (10-17%)	18% (14-22%)	1.1 (0.79-1.4)	65 (48-83)
Bermuda	1995	Male	40 to 44	68% (51-82%)	32% (18-49%)	10 (5.8-16)	13% (10-17%)	17% (13-21%)	1.3 (1-1.7)	70 (54-89)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	1995	Male	45 to 49	66% (45-81%)	34% (19-55%)	9.6 (5.4-16)	14% (11-18%)	17% (13-22%)	1.8 (1.3-2.4)	82 (60-110)
Bermuda	1995	Male	50 to 54	64% (44-80%)	36% (20-56%)	8.8 (4.9-14)	13% (9.8-17%)	16% (12-21%)	2.1 (1.5-2.8)	83 (61-110)
Bermuda	1995	Male	55 to 59	62% (44-78%)	38% (22-56%)	8 (4.3-13)	13% (9.8-18%)	16% (11-20%)	2.9 (2-3.9)	96 (69-130)
Bermuda	1995	Male	60 to 64	60% (40-78%)	40% (22-60%)	7.2 (3.9-12)	12% (8.4-16%)	14% (9.4-19%)	2.9 (2-4.1)	83 (58-120)
Bermuda	1995	Male	65 to 69	58% (39-76%)	42% (24-61%)	6.3 (3.5-10)	12% (7.7-16%)	13% (8.7-18%)	3.3 (2.2-4.7)	77 (52-110)
Bermuda	1995	Male	70 to 74	54% (36-71%)	46% (29-64%)	5.2 (2.8-9)	9.7% (6.3-14%)	11% (6.9-16%)	3.2 (2.1-4.6)	60 (39-86)
Bermuda	1995	Male	75 to 79	49% (31-68%)	51% (32-69%)	4.3 (2.1-7.4)	8.4% (5.4-12%)	9.3% (5.8-13%)	2.9 (1.8-4.3)	43 (27-61)
Bermuda	1995	Male	80 to 84	43% (26-62%)	57% (38-74%)	3.6 (1.8-5.9)	7.3% (4.3-11%)	8.1% (4.7-12%)	2.4 (1.4-3.5)	25 (15-38)
Bermuda	1995	Male	85 to 89	39% (23-56%)	61% (44-77%)	3.1 (1.6-5.4)	6.5% (3.9-9.9%)	7.2% (4.2-11%)	1.3 (0.78-2.1)	10 (6.3-16)
Bermuda	1995	Male	90 to 94	39% (24-56%)	61% (44-76%)	3.2 (1.5-5.8)	6.4% (3.8-9.7%)	7% (4-11%)	0.69 (0.39-1.1)	3.6 (2.1-5.6)
Bermuda	1995	Male	95 plus	43% (24-63%)	57% (37-76%)	3.5 (1.4-6.6)	6.7% (3.6-11%)	7.2% (3.7-12%)	0.18 (0.093-0.3)	0.65 (0.34-1)
Bermuda	2000	Female	All Ages	30% (24-37%)	70% (63-76%)	2.8 (2.5-3.1)	5% (3.6-6.7%)	6.9% (4.5-9.8%)	16 (10-23)	350 (250-480)
Bermuda	2000	Female	15 to 19	35% (23-52%)	65% (48-77%)	3.2 (1.7-5.3)	3.3% (2.1-4.6%)	8.9% (5.3-13%)	0.088 (0.052-0.13)	9.6 (6.2-14)
Bermuda	2000	Female	20 to 24	41% (26-58%)	59% (42-74%)	4.8 (2.7-7.5)	3.4% (2.4-4.4%)	6.4% (4-8.9%)	0.045 (0.028-0.065)	8.9 (6.2-12)
Bermuda	2000	Female	25 to 29	41% (25-57%)	59% (43-75%)	5 (3-7.5)	3.8% (2.9-5%)	6.5% (4.4-8.8%)	0.085 (0.057-0.12)	12 (8.5-16)
Bermuda	2000	Female	30 to 34	39% (23-56%)	61% (44-77%)	4.8 (2.8-7.2)	3.9% (2.8-5.3%)	6.4% (4.1-9.3%)	0.15 (0.09-0.22)	15 (10-20)
Bermuda	2000	Female	35 to 39	38% (24-55%)	62% (45-76%)	4.4 (2.6-6.9)	4.2% (2.8-5.9%)	7% (4.3-10%)	0.23 (0.13-0.35)	17 (11-25)
Bermuda	2000	Female	40 to 44	37% (23-55%)	63% (45-77%)	4.2 (2.2-6.6)	5.4% (3.5-7.8%)	9.3% (5.8-14%)	0.36 (0.22-0.56)	21 (13-31)
Bermuda	2000	Female	45 to 49	35% (21-52%)	65% (48-79%)	4.1 (2.1-6.9)	6% (3.7-9.1%)	9.9% (6.2-15%)	0.48 (0.29-0.75)	24 (15-36)
Bermuda	2000	Female	50 to 54	32% (19-48%)	68% (52-81%)	3.6 (2-5.9)	6.8% (4.2-10%)	10% (6.4-16%)	0.68 (0.41-1)	28 (17-43)
Bermuda	2000	Female	55 to 59	28% (16-43%)	72% (57-84%)	3 (1.5-5.1)	7% (4.3-11%)	9.7% (5.9-15%)	0.95 (0.59-1.5)	33 (20-50)
Bermuda	2000	Female	60 to 64	24% (14-39%)	76% (61-86%)	2.6 (1.3-4.6)	6.6% (3.9-10%)	8.5% (5-13%)	1.3 (0.74-2)	36 (22-57)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	2000	Female	65 to 69	22% (14-36%)	78% (64-86%)	2.2 (1.2-3.7)	6.4% (3.8-9.9%)	7.9% (4.7-12%)	1.5 (0.88-2.3)	35 (21-54)
Bermuda	2000	Female	70 to 74	20% (12-32%)	80% (68-88%)	1.9 (0.93-3.4)	5.8% (3.3-9.1%)	7% (4-11%)	1.7 (0.93-2.7)	31 (18-50)
Bermuda	2000	Female	75 to 79	18% (10-29%)	82% (71-90%)	1.6 (0.74-3)	5.2% (2.7-8.4%)	6.1% (3.1-10%)	1.8 (0.89-3)	26 (13-43)
Bermuda	2000	Female	80 to 84	16% (9.2-27%)	84% (73-91%)	1.5 (0.74-2.7)	5.4% (2.8-8.8%)	6.1% (3.1-10%)	2.5 (1.2-4.2)	26 (13-44)
Bermuda	2000	Female	85 to 89	16% (9.3-26%)	84% (74-91%)	1.5 (0.71-2.8)	5.4% (2.8-9.1%)	6% (3-10%)	2.3 (1.2-4)	17 (8.7-29)
Bermuda	2000	Female	90 to 94	18% (11-29%)	82% (71-89%)	1.7 (0.76-3.2)	5.6% (2.7-9.5%)	6.2% (3-11%)	1.2 (0.59-2.2)	6.4 (3.1-11)
Bermuda	2000	Female	95 plus	22% (11-37%)	78% (63-89%)	2.1 (0.74-4.2)	6% (2.4-11%)	6.6% (2.7-12%)	0.47 (0.19-0.87)	1.6 (0.66-3)
Bermuda	2000	Male	All Ages	53% (44-61%)	47% (39-56%)	6.4 (5.9-6.8)	9% (7.1-11%)	10% (7.7-13%)	26 (20-34)	750 (580-940)
Bermuda	2000	Male	15 to 19	52% (35-70%)	48% (30-65%)	4.8 (2.6-7.9)	6.1% (4.1-8.4%)	13% (7.6-18%)	0.14 (0.08-0.2)	16 (11-23)
Bermuda	2000	Male	20 to 24	64% (45-79%)	36% (21-55%)	9.3 (5.7-14)	11% (8.2-14%)	16% (10-22%)	0.34 (0.21-0.47)	35 (25-47)
Bermuda	2000	Male	25 to 29	67% (48-81%)	33% (19-52%)	11 (7.3-16)	8.2% (6.1-11%)	8.8% (4.7-13%)	0.22 (0.11-0.34)	28 (19-39)
Bermuda	2000	Male	30 to 34	68% (48-82%)	32% (18-52%)	12 (7.5-17)	9% (6.6-12%)	11% (7-15%)	0.51 (0.33-0.75)	42 (30-57)
Bermuda	2000	Male	35 to 39	68% (50-82%)	32% (18-50%)	11 (7-16)	7.6% (5.5-9.7%)	9% (5.8-12%)	0.4 (0.24-0.56)	33 (24-44)
Bermuda	2000	Male	40 to 44	66% (50-81%)	34% (19-50%)	10 (5.9-16)	8.9% (6.7-11%)	11% (8-14%)	0.73 (0.5-0.98)	43 (32-57)
Bermuda	2000	Male	45 to 49	64% (44-80%)	36% (20-56%)	9.8 (5.6-16)	12% (8.8-15%)	14% (10-17%)	1.7 (1.2-2.2)	79 (58-100)
Bermuda	2000	Male	50 to 54	63% (44-79%)	37% (21-56%)	8.9 (5-15)	12% (8.8-15%)	14% (11-19%)	1.6 (1.2-2.1)	66 (49-86)
Bermuda	2000	Male	55 to 59	60% (42-78%)	40% (22-58%)	8.1 (4.4-13)	12% (9-16%)	14% (11-19%)	2.7 (1.9-3.6)	91 (65-120)
Bermuda	2000	Male	60 to 64	58% (38-77%)	42% (23-62%)	7.3 (3.8-12)	11% (7.7-15%)	13% (9-17%)	2.6 (1.8-3.6)	75 (53-100)
Bermuda	2000	Male	65 to 69	57% (37-74%)	43% (26-63%)	6.4 (3.5-10)	11% (7.3-15%)	12% (8.2-17%)	3.4 (2.3-4.7)	80 (54-110)
Bermuda	2000	Male	70 to 74	53% (35-70%)	47% (30-65%)	5.3 (2.8-8.8)	9.3% (6-13%)	10% (6.7-15%)	3.5 (2.2-4.9)	65 (43-92)
Bermuda	2000	Male	75 to 79	48% (30-67%)	52% (33-70%)	4.3 (2.1-7.4)	8% (5.2-11%)	8.8% (5.6-13%)	3.1 (2-4.5)	45 (29-64)
Bermuda	2000	Male	80 to 84	42% (25-60%)	58% (40-75%)	3.6 (1.8-5.9)	7% (4.3-10%)	7.7% (4.7-11%)	2.6 (1.6-3.8)	28 (17-41)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	2000	Male	85 to 89	38% (22-54%)	62% (46-78%)	3.2 (1.6-5.5)	6.3% (3.8-9.5%)	6.9% (4.1-11%)	1.5 (0.89-2.3)	12 (7-18)
Bermuda	2000	Male	90 to 94	38% (24-55%)	62% (45-76%)	3.2 (1.5-5.8)	6.3% (3.8-9.6%)	6.8% (3.9-11%)	0.92 (0.54-1.4)	4.9 (2.9-7.5)
Bermuda	2000	Male	95 plus	41% (23-61%)	59% (39-77%)	3.6 (1.4-6.7)	6.7% (3.7-11%)	7.2% (3.8-12%)	0.26 (0.14-0.43)	0.95 (0.52-1.5)
Bermuda	2005	Female	All Ages	29% (24-36%)	71% (64-76%)	2.7 (2.4-3)	4.5% (3.3-6.1%)	6.6% (4.3-9.3%)	13 (8.8-19)	300 (220-410)
Bermuda	2005	Female	15 to 19	35% (22-51%)	65% (49-78%)	3.1 (1.6-5.1)	2.9% (1.9-4.1%)	9.1% (5.5-13%)	0.07 (0.041-0.1)	8.4 (5.3-12)
Bermuda	2005	Female	20 to 24	40% (25-57%)	60% (43-75%)	4.6 (2.5-7.2)	3.2% (2.3-4.3%)	6.4% (4-9.2%)	0.038 (0.022-0.057)	8.8 (6-12)
Bermuda	2005	Female	25 to 29	40% (24-57%)	60% (43-76%)	4.7 (2.8-7.3)	3.5% (2.6-4.6%)	6.2% (4-8.5%)	0.056 (0.034-0.08)	9.5 (6.7-13)
Bermuda	2005	Female	30 to 34	38% (23-55%)	62% (45-77%)	4.5 (2.6-6.9)	4% (2.9-5.3%)	7.7% (5.1-11%)	0.12 (0.079-0.18)	13 (9-18)
Bermuda	2005	Female	35 to 39	37% (23-55%)	63% (45-77%)	4.2 (2.5-6.6)	3.9% (2.7-5.5%)	7.8% (4.9-11%)	0.16 (0.099-0.25)	14 (9.6-20)
Bermuda	2005	Female	40 to 44	36% (23-55%)	64% (45-77%)	4 (2.1-6.2)	4.8% (3.2-7.1%)	9.8% (6.2-15%)	0.32 (0.2-0.49)	20 (13-30)
Bermuda	2005	Female	45 to 49	34% (20-50%)	66% (50-80%)	3.8 (2-6.4)	5.4% (3.4-8.2%)	10% (6.3-16%)	0.38 (0.23-0.61)	20 (13-30)
Bermuda	2005	Female	50 to 54	31% (18-47%)	69% (53-82%)	3.4 (1.8-5.6)	6.1% (3.7-9.3%)	10% (6.3-16%)	0.58 (0.36-0.89)	25 (16-38)
Bermuda	2005	Female	55 to 59	27% (16-43%)	73% (57-84%)	2.9 (1.5-5)	6.3% (3.8-9.6%)	9.5% (5.8-14%)	0.77 (0.47-1.2)	27 (17-42)
Bermuda	2005	Female	60 to 64	24% (14-39%)	76% (61-86%)	2.4 (1.2-4.4)	6.1% (3.6-9.3%)	8.2% (5-13%)	1 (0.61-1.6)	30 (18-47)
Bermuda	2005	Female	65 to 69	22% (13-34%)	78% (66-87%)	2.1 (1.1-3.6)	5.8% (3.4-8.8%)	7.6% (4.6-11%)	1.2 (0.73-1.9)	30 (18-46)
Bermuda	2005	Female	70 to 74	20% (12-31%)	80% (69-88%)	1.8 (0.89-3.2)	5.2% (2.9-8.1%)	6.6% (3.8-10%)	1.3 (0.73-2.1)	25 (14-40)
Bermuda	2005	Female	75 to 79	17% (10-28%)	83% (72-90%)	1.5 (0.68-2.8)	4.7% (2.4-7.7%)	5.8% (3-9.3%)	1.4 (0.72-2.4)	21 (11-35)
Bermuda	2005	Female	80 to 84	16% (9-26%)	84% (74-91%)	1.4 (0.7-2.5)	5% (2.5-8.1%)	5.8% (2.9-9.4%)	2.2 (1.1-3.6)	23 (12-38)
Bermuda	2005	Female	85 to 89	16% (9.3-25%)	84% (75-91%)	1.5 (0.69-2.6)	5.1% (2.6-8.4%)	5.7% (2.9-9.4%)	2.1 (1.1-3.5)	16 (8.1-26)
Bermuda	2005	Female	90 to 94	18% (10-28%)	82% (72-90%)	1.7 (0.74-3)	5.2% (2.6-8.8%)	5.9% (2.8-10%)	1.2 (0.56-2)	6.1 (3-10)
Bermuda	2005	Female	95 plus	21% (11-36%)	79% (64-89%)	2 (0.72-4)	5.7% (2.3-10%)	6.4% (2.4-12%)	0.49 (0.19-0.91)	1.7 (0.67-3.1)
Bermuda	2005	Male	All Ages	52% (44-61%)	48% (39-56%)	6.2 (5.7-6.6)	9.5% (7.6-12%)	11% (8.3-13%)	24 (18-30)	730 (580-910)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	2005	Male	15 to 19	52% (34-69%)	48% (31-66%)	4.6 (2.5-7.4)	6.4% (4.3-8.8%)	15% (9.2-22%)	0.14 (0.088-0.22)	17 (12-24)
Bermuda	2005	Male	20 to 24	63% (44-79%)	37% (21-56%)	8.8 (5.3-13)	14% (9.8-18%)	21% (13-29%)	0.54 (0.34-0.76)	49 (34-65)
Bermuda	2005	Male	25 to 29	66% (48-81%)	34% (19-52%)	11 (6.8-15)	13% (9.1-16%)	19% (12-26%)	0.46 (0.29-0.63)	42 (29-54)
Bermuda	2005	Male	30 to 34	67% (47-82%)	33% (18-53%)	11 (7-16)	10% (7.7-13%)	15% (10-20%)	0.44 (0.29-0.6)	38 (28-50)
Bermuda	2005	Male	35 to 39	67% (49-82%)	33% (18-51%)	11 (6.7-15)	11% (8-14%)	17% (12-21%)	0.59 (0.42-0.78)	43 (31-54)
Bermuda	2005	Male	40 to 44	66% (49-81%)	34% (19-51%)	10 (5.6-15)	12% (9.2-15%)	17% (13-21%)	1.2 (0.87-1.5)	65 (49-83)
Bermuda	2005	Male	45 to 49	64% (43-79%)	36% (21-57%)	9.4 (5.4-15)	12% (9.2-15%)	16% (12-20%)	1.4 (1-1.7)	65 (49-84)
Bermuda	2005	Male	50 to 54	62% (42-78%)	38% (22-58%)	8.5 (4.7-13)	12% (9-15%)	15% (12-20%)	1.7 (1.3-2.3)	71 (53-92)
Bermuda	2005	Male	55 to 59	60% (42-77%)	40% (23-58%)	7.7 (4.2-13)	12% (9-16%)	15% (11-19%)	2.4 (1.8-3.2)	82 (61-110)
Bermuda	2005	Male	60 to 64	58% (37-76%)	42% (24-63%)	6.9 (3.7-12)	11% (7.7-14%)	13% (9.2-17%)	2.2 (1.6-2.9)	65 (46-87)
Bermuda	2005	Male	65 to 69	56% (36-74%)	44% (26-64%)	6.1 (3.3-9.8)	10% (7.2-14%)	12% (8.5-16%)	2.5 (1.7-3.4)	60 (42-81)
Bermuda	2005	Male	70 to 74	52% (33-70%)	48% (30-67%)	5.1 (2.7-8.6)	8.9% (6-12%)	10% (7-14%)	2.8 (1.9-3.8)	53 (36-73)
Bermuda	2005	Male	75 to 79	47% (28-66%)	53% (34-72%)	4.1 (2-7.1)	7.7% (5-11%)	8.7% (5.6-12%)	2.8 (1.8-3.9)	41 (26-58)
Bermuda	2005	Male	80 to 84	41% (24-59%)	59% (41-76%)	3.4 (1.7-5.7)	6.7% (4.2-9.8%)	7.5% (4.6-11%)	2.2 (1.4-3.3)	24 (15-36)
Bermuda	2005	Male	85 to 89	37% (21-53%)	63% (47-79%)	3 (1.5-5.1)	5.8% (3.6-8.8%)	6.6% (4-10%)	1.2 (0.78-1.9)	9.9 (6.4-15)
Bermuda	2005	Male	90 to 94	37% (23-54%)	63% (46-77%)	3 (1.5-5.6)	5.9% (3.6-9%)	6.4% (3.8-9.9%)	0.88 (0.52-1.4)	4.7 (2.8-7.1)
Bermuda	2005	Male	95 plus	40% (22-61%)	60% (39-78%)	3.3 (1.3-6.4)	6.3% (3.4-10%)	6.8% (3.6-11%)	0.27 (0.14-0.44)	0.97 (0.53-1.5)
Bermuda	2010	Female	All Ages	29% (23-35%)	71% (65-77%)	2.4 (2.1-2.7)	4% (2.9-5.4%)	5.8% (3.8-8.2%)	11 (6.9-15)	260 (190-340)
Bermuda	2010	Female	15 to 19	34% (22-50%)	66% (50-78%)	2.7 (1.4-4.5)	2.6% (1.7-3.8%)	8.7% (4.9-13%)	0.056 (0.031-0.088)	7.3 (4.6-11)
Bermuda	2010	Female	20 to 24	40% (25-56%)	60% (44-75%)	4.1 (2.3-6.4)	3.4% (2.5-4.5%)	7.5% (4.8-11%)	0.044 (0.027-0.067)	9.5 (6.4-14)
Bermuda	2010	Female	25 to 29	39% (24-56%)	61% (44-76%)	4.2 (2.4-6.6)	3.7% (2.7-4.8%)	7.3% (4.8-10%)	0.063 (0.039-0.091)	10 (7.1-14)
Bermuda	2010	Female	30 to 34	38% (23-54%)	62% (46-77%)	4 (2.3-6.2)	3.8% (2.7-5.2%)	7.9% (5.2-11%)	0.1 (0.062-0.15)	11 (8-16)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	2010	Female	35 to 39	37% (23-54%)	63% (46-77%)	3.7 (2.1-5.8)	3.7% (2.6-5.2%)	8.1% (5.2-12%)	0.14 (0.085-0.22)	13 (8.7-18)
Bermuda	2010	Female	40 to 44	35% (22-54%)	65% (46-78%)	3.5 (1.9-5.6)	4.3% (2.8-6.2%)	9.2% (5.8-14%)	0.25 (0.15-0.38)	17 (11-25)
Bermuda	2010	Female	45 to 49	34% (19-50%)	66% (50-81%)	3.3 (1.7-5.8)	4.7% (2.9-7.2%)	9.5% (5.9-15%)	0.35 (0.21-0.57)	20 (12-30)
Bermuda	2010	Female	50 to 54	30% (18-46%)	70% (54-82%)	3 (1.6-5)	5.3% (3.3-8.1%)	9.4% (5.8-15%)	0.49 (0.3-0.77)	22 (14-34)
Bermuda	2010	Female	55 to 59	26% (16-42%)	74% (58-84%)	2.5 (1.3-4.3)	5.4% (3.2-8.2%)	8.6% (5.3-13%)	0.67 (0.4-1)	25 (15-38)
Bermuda	2010	Female	60 to 64	23% (14-37%)	77% (63-86%)	2.1 (1-3.7)	5.2% (3-8%)	7.4% (4.4-11%)	0.79 (0.46-1.2)	24 (14-38)
Bermuda	2010	Female	65 to 69	21% (13-34%)	79% (66-87%)	1.9 (0.98-3.2)	4.9% (2.9-7.6%)	6.8% (4.1-10%)	0.91 (0.53-1.4)	23 (13-36)
Bermuda	2010	Female	70 to 74	19% (12-31%)	81% (69-88%)	1.6 (0.79-2.9)	4.4% (2.4-7.1%)	5.8% (3.2-9.2%)	1 (0.56-1.7)	20 (11-33)
Bermuda	2010	Female	75 to 79	17% (9.8-28%)	83% (72-90%)	1.3 (0.61-2.4)	4% (2-6.5%)	5% (2.5-8.2%)	1.1 (0.55-1.8)	17 (8.3-28)
Bermuda	2010	Female	80 to 84	16% (8.5-26%)	84% (74-92%)	1.3 (0.59-2.2)	4.2% (2.1-6.7%)	5% (2.5-8.1%)	1.6 (0.75-2.6)	17 (8.3-29)
Bermuda	2010	Female	85 to 89	16% (8.9-25%)	84% (75-91%)	1.3 (0.6-2.3)	4.2% (2.1-7%)	4.9% (2.4-8.1%)	1.6 (0.74-2.6)	12 (5.8-20)
Bermuda	2010	Female	90 to 94	17% (10-28%)	83% (72-90%)	1.5 (0.64-2.7)	4.4% (1.9-7.6%)	5.1% (2.2-9.1%)	0.99 (0.43-1.8)	5.2 (2.3-9.1)
Bermuda	2010	Female	95 plus	21% (10-35%)	79% (65-90%)	1.8 (0.63-3.6)	4.8% (1.8-8.7%)	5.5% (1.9-10%)	0.43 (0.15-0.82)	1.5 (0.55-2.8)
Bermuda	2010	Male	All Ages	52% (43-60%)	48% (40-57%)	5.5 (5-6)	9.4% (7.6-11%)	10% (8.2-13%)	23 (18-29)	730 (590-900)
Bermuda	2010	Male	15 to 19	51% (33-69%)	49% (31-67%)	4 (2.2-6.4)	6.8% (4.4-9.5%)	15% (8.5-22%)	0.18 (0.1-0.27)	20 (13-27)
Bermuda	2010	Male	20 to 24	62% (43-78%)	38% (22-57%)	7.8 (4.7-11)	14% (10-19%)	21% (13-30%)	0.64 (0.4-0.91)	57 (38-76)
Bermuda	2010	Male	25 to 29	65% (46-81%)	35% (19-54%)	9.4 (5.9-13)	14% (10-18%)	21% (13-28%)	0.69 (0.44-0.94)	56 (40-74)
Bermuda	2010	Male	30 to 34	67% (46-82%)	33% (18-54%)	9.9 (6.1-14)	11% (8.1-14%)	17% (11-22%)	0.43 (0.29-0.57)	36 (26-47)
Bermuda	2010	Male	35 to 39	66% (49-81%)	34% (19-51%)	9.3 (5.8-14)	11% (8.2-14%)	16% (12-21%)	0.6 (0.43-0.78)	43 (32-55)
Bermuda	2010	Male	40 to 44	65% (48-80%)	35% (20-52%)	8.9 (5.1-13)	12% (8.9-15%)	16% (13-21%)	1 (0.78-1.3)	59 (45-76)
Bermuda	2010	Male	45 to 49	63% (42-79%)	37% (21-58%)	8.2 (4.7-13)	11% (8.7-14%)	16% (12-19%)	1.3 (1-1.7)	67 (51-85)
Bermuda	2010	Male	50 to 54	61% (41-78%)	39% (22-59%)	7.5 (4.2-12)	11% (8.4-14%)	15% (11-18%)	1.4 (1.1-1.9)	61 (47-80)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	2010	Male	55 to 59	59% (40-76%)	41% (24-60%)	6.8 (3.6-11)	12% (8.8-15%)	14% (11-18%)	2.4 (1.8-3.1)	83 (62-110)
Bermuda	2010	Male	60 to 64	57% (37-76%)	43% (24-63%)	6.1 (3.2-11)	10% (7.5-14%)	12% (9.1-16%)	2.2 (1.6-3)	65 (47-87)
Bermuda	2010	Male	65 to 69	55% (36-73%)	45% (27-64%)	5.4 (2.9-8.7)	9.7% (6.8-13%)	12% (8.4-15%)	2.3 (1.6-3.1)	57 (39-76)
Bermuda	2010	Male	70 to 74	51% (32-68%)	49% (32-68%)	4.4 (2.3-7.6)	8.4% (5.7-12%)	9.8% (6.6-14%)	2.5 (1.7-3.6)	49 (33-69)
Bermuda	2010	Male	75 to 79	46% (28-66%)	54% (34-72%)	3.6 (1.8-6.3)	7.2% (4.6-10%)	8.1% (5.2-11%)	2.6 (1.7-3.8)	39 (25-56)
Bermuda	2010	Male	80 to 84	41% (24-59%)	59% (41-76%)	3 (1.5-5)	6.3% (4-9%)	7.1% (4.5-10%)	2.2 (1.4-3.3)	25 (16-36)
Bermuda	2010	Male	85 to 89	36% (21-53%)	64% (47-79%)	2.6 (1.3-4.5)	5.6% (3.4-8.4%)	6.2% (3.8-9.6%)	1.4 (0.84-2.1)	11 (6.7-17)
Bermuda	2010	Male	90 to 94	36% (22-53%)	64% (47-78%)	2.7 (1.3-4.9)	5.6% (3.3-8.4%)	6.1% (3.5-9.3%)	0.92 (0.53-1.4)	4.9 (3-7.5)
Bermuda	2010	Male	95 plus	39% (22-60%)	61% (40-78%)	2.9 (1.2-5.7)	5.8% (3-9.4%)	6.3% (3.1-10%)	0.28 (0.14-0.46)	1 (0.52-1.6)
Bermuda	2016	Female	All Ages	28% (22-34%)	72% (66-78%)	2.3 (1.8-2.9)	3.5% (2.6-4.8%)	5.3% (3.5-7.5%)	8.7 (5.5-13)	230 (160-310)
Bermuda	2016	Female	15 to 19	33% (21-49%)	67% (51-79%)	2.6 (1.2-4.4)	2.4% (1.5-3.6%)	8.2% (4.4-12%)	0.05 (0.026-0.08)	6.9 (4.3-9.9)
Bermuda	2016	Female	20 to 24	38% (23-55%)	62% (45-77%)	3.8 (2-6.3)	3.3% (2.4-4.5%)	7.6% (4.8-11%)	0.042 (0.025-0.065)	9.5 (6.3-13)
Bermuda	2016	Female	25 to 29	38% (22-55%)	62% (45-78%)	3.9 (2.1-6.3)	3.5% (2.6-4.6%)	7% (4.5-9.7%)	0.06 (0.038-0.09)	11 (7.6-15)
Bermuda	2016	Female	30 to 34	36% (22-54%)	64% (46-78%)	3.8 (2.1-6)	3.6% (2.6-4.8%)	7.4% (4.9-11%)	0.091 (0.056-0.14)	12 (8.2-16)
Bermuda	2016	Female	35 to 39	35% (22-52%)	65% (48-78%)	3.5 (1.9-5.6)	3.4% (2.4-4.7%)	7.4% (4.7-11%)	0.11 (0.063-0.17)	11 (7.3-15)
Bermuda	2016	Female	40 to 44	34% (21-52%)	66% (48-79%)	3.3 (1.7-5.6)	3.8% (2.5-5.6%)	8.6% (5.4-13%)	0.2 (0.11-0.33)	15 (9.1-22)
Bermuda	2016	Female	45 to 49	32% (18-48%)	68% (52-82%)	3.2 (1.6-5.6)	4.3% (2.7-6.7%)	8.9% (5.4-14%)	0.32 (0.19-0.54)	19 (11-29)
Bermuda	2016	Female	50 to 54	29% (17-45%)	71% (55-83%)	2.8 (1.4-4.9)	4.7% (2.9-7.4%)	8.7% (5.4-13%)	0.52 (0.29-0.82)	24 (15-38)
Bermuda	2016	Female	55 to 59	25% (15-40%)	75% (60-85%)	2.4 (1.1-4.1)	4.6% (2.8-7.1%)	7.8% (4.8-12%)	0.58 (0.34-0.91)	23 (13-35)
Bermuda	2016	Female	60 to 64	22% (13-37%)	78% (63-87%)	2 (0.95-3.6)	4.3% (2.6-6.8%)	6.7% (4-10%)	0.67 (0.38-1.1)	21 (12-34)
Bermuda	2016	Female	65 to 69	20% (12-33%)	80% (67-88%)	1.7 (0.88-3)	4.1% (2.4-6.5%)	6.1% (3.6-9.3%)	0.72 (0.4-1.1)	19 (11-30)
Bermuda	2016	Female	70 to 74	18% (11-29%)	82% (71-89%)	1.5 (0.7-2.8)	3.7% (1.9-6.2%)	5.2% (2.8-8.5%)	0.79 (0.41-1.3)	16 (8.5-27)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	2016	Female	75 to 79	16% (9.2-27%)	84% (73-91%)	1.3 (0.56-2.3)	3.4% (1.7-5.8%)	4.5% (2.2-7.8%)	0.92 (0.44-1.6)	15 (6.8-25)
Bermuda	2016	Female	80 to 84	15% (8.1-25%)	85% (75-92%)	1.2 (0.54-2.1)	3.4% (1.7-5.7%)	4.4% (2.1-7.2%)	1.1 (0.53-1.9)	13 (6-22)
Bermuda	2016	Female	85 to 89	15% (8.4-23%)	85% (77-92%)	1.2 (0.53-2.2)	3.5% (1.7-6%)	4.3% (2-7.4%)	1.2 (0.54-2.1)	9.4 (4.3-16)
Bermuda	2016	Female	90 to 94	17% (9.4-26%)	83% (74-91%)	1.4 (0.58-2.7)	3.8% (1.7-6.6%)	4.5% (1.9-8%)	0.9 (0.38-1.6)	4.8 (2.1-8.5)
Bermuda	2016	Female	95 plus	20% (9.9-34%)	80% (66-90%)	1.6 (0.56-3.4)	4.2% (1.5-7.8%)	4.9% (1.6-9.4%)	0.42 (0.14-0.83)	1.5 (0.53-2.8)
Bermuda	2016	Male	All Ages	51% (43-60%)	49% (40-57%)	5.4 (4.2-6.8)	8.7% (7-11%)	9.6% (7.7-12%)	23 (17-29)	720 (570-910)
Bermuda	2016	Male	15 to 19	50% (32-68%)	50% (32-68%)	3.8 (1.9-6.5)	5.8% (3.9-8%)	14% (7.9-21%)	0.12 (0.067-0.19)	16 (10-22)
Bermuda	2016	Male	20 to 24	61% (42-77%)	39% (23-58%)	7.4 (4.1-11)	13% (9.2-16%)	20% (13-28%)	0.46 (0.28-0.66)	46 (32-63)
Bermuda	2016	Male	25 to 29	64% (44-80%)	36% (20-56%)	9 (5.3-14)	12% (9.2-16%)	19% (12-26%)	0.51 (0.31-0.73)	48 (34-64)
Bermuda	2016	Male	30 to 34	65% (45-81%)	35% (19-55%)	9.5 (5.6-14)	11% (7.9-14%)	16% (11-22%)	0.44 (0.28-0.63)	40 (28-53)
Bermuda	2016	Male	35 to 39	65% (48-80%)	35% (20-52%)	8.9 (5.3-14)	9.8% (7.6-13%)	15% (11-19%)	0.46 (0.31-0.65)	36 (26-48)
Bermuda	2016	Male	40 to 44	64% (47-79%)	36% (21-53%)	8.4 (4.5-13)	11% (8.1-13%)	15% (12-19%)	0.85 (0.59-1.1)	51 (37-68)
Bermuda	2016	Male	45 to 49	62% (40-78%)	38% (22-60%)	7.8 (4.1-13)	11% (8.1-14%)	15% (11-19%)	1.3 (0.91-1.8)	65 (48-86)
Bermuda	2016	Male	50 to 54	60% (41-77%)	40% (23-59%)	7.1 (3.7-12)	10% (7.8-13%)	14% (11-18%)	1.7 (1.2-2.3)	72 (53-97)
Bermuda	2016	Male	55 to 59	58% (39-76%)	42% (24-61%)	6.4 (3.3-11)	11% (8-14%)	14% (10-18%)	2.4 (1.7-3.3)	85 (61-120)
Bermuda	2016	Male	60 to 64	56% (35-74%)	44% (26-65%)	5.8 (3-10)	9.5% (6.8-13%)	12% (8.5-16%)	2.3 (1.6-3.3)	70 (49-97)
Bermuda	2016	Male	65 to 69	54% (35-72%)	46% (28-65%)	5.1 (2.7-8.7)	9.1% (6.4-12%)	11% (7.9-15%)	2.4 (1.7-3.4)	60 (42-82)
Bermuda	2016	Male	70 to 74	50% (32-67%)	50% (33-68%)	4.2 (2.1-7.4)	7.8% (5.3-11%)	9.2% (6.3-13%)	2.5 (1.7-3.6)	50 (33-70)
Bermuda	2016	Male	75 to 79	45% (27-64%)	55% (36-73%)	3.4 (1.6-6)	6.6% (4.3-9.4%)	7.6% (4.8-11%)	2.6 (1.7-3.8)	40 (25-57)
Bermuda	2016	Male	80 to 84	39% (22-58%)	61% (42-78%)	2.8 (1.4-4.9)	5.8% (3.7-8.5%)	6.6% (4.2-9.7%)	2.1 (1.3-3.1)	23 (15-34)
Bermuda	2016	Male	85 to 89	35% (20-51%)	65% (49-80%)	2.5 (1.2-4.5)	5.1% (3.2-7.8%)	5.8% (3.5-9%)	1.3 (0.79-2)	11 (6.5-16)
Bermuda	2016	Male	90 to 94	35% (21-52%)	65% (48-79%)	2.5 (1.1-4.7)	5.2% (2.9-7.8%)	5.7% (3.1-8.7%)	0.96 (0.54-1.5)	5.2 (3-7.8)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Bermuda	2016	Male	95 plus	38% (20-59%)	62% (41-80%)	2.7 (1-5.4)	5.4% (2.6-8.9%)	5.9% (2.6-9.9%)	0.31 (0.13-0.51)	1.1 (0.52-1.8)
Greenland	1990	Female	All Ages	74% (67-80%)	26% (20-33%)	1.8 (1.1-2.8)	5.1% (2.8-8.3%)	5% (1.9-9.4%)	9.1 (3.5-18)	450 (240-720)
Greenland	1990	Female	15 to 19	64% (48-79%)	36% (21-52%)	1.6 (0.75-2.8)	7.2% (1.9-15%)	11% (2.3-22%)	0.4 (0.091-0.85)	30 (8.1-62)
Greenland	1990	Female	20 to 24	80% (66-90%)	20% (10-34%)	3 (1.6-5.1)	9.8% (4.2-17%)	19% (6.3-33%)	0.71 (0.24-1.3)	55 (24-95)
Greenland	1990	Female	25 to 29	84% (74-91%)	16% (8.6-26%)	3 (1.7-5.1)	10% (4.5-17%)	19% (7.1-33%)	0.98 (0.38-1.7)	68 (31-110)
Greenland	1990	Female	30 to 34	86% (77-93%)	14% (6.9-23%)	2.9 (1.6-4.8)	8.8% (4.2-14%)	17% (6.7-28%)	0.69 (0.29-1.2)	45 (21-74)
Greenland	1990	Female	35 to 39	87% (78-94%)	13% (6.3-22%)	2.6 (1.4-4.5)	8.5% (4.6-13%)	15% (7.2-24%)	0.63 (0.3-1.1)	38 (21-61)
Greenland	1990	Female	40 to 44	87% (78-93%)	13% (6.6-22%)	2.5 (1.3-4.4)	9.9% (5.7-15%)	15% (7.8-23%)	0.86 (0.46-1.4)	44 (25-72)
Greenland	1990	Female	45 to 49	86% (76-93%)	14% (6.9-24%)	2.4 (1.3-4)	9.2% (5.6-14%)	13% (7.9-20%)	0.93 (0.54-1.5)	43 (26-66)
Greenland	1990	Female	50 to 54	85% (75-92%)	15% (8-25%)	2.2 (1.1-3.8)	8.2% (4.7-13%)	11% (6.4-18%)	0.86 (0.48-1.4)	34 (19-55)
Greenland	1990	Female	55 to 59	83% (71-91%)	17% (8.8-29%)	2 (1-3.4)	7% (3.6-12%)	8.6% (4.4-14%)	1.3 (0.65-2.2)	42 (21-72)
Greenland	1990	Female	60 to 64	80% (67-90%)	20% (9.9-33%)	1.8 (0.92-3.3)	4.8% (1.6-9.4%)	5.8% (2-11%)	0.99 (0.34-1.9)	27 (9.2-55)
Greenland	1990	Female	65 to 69	77% (62-88%)	23% (12-38%)	1.6 (0.82-2.8)	3% (0.28-7.2%)	3.6% (0.37-8.3%)	0.58 (0.061-1.4)	13 (1.2-32)
Greenland	1990	Female	70 to 74	74% (60-85%)	26% (15-40%)	1.4 (0.66-2.6)	1.6% (-0.97-6.2%)	1.9% (-1-6.9%)	0.37 (-0.2-1.4)	6.5 (-3.9-26)
Greenland	1990	Female	75 to 79	72% (56-84%)	28% (16-44%)	1.3 (0.62-2.4)	0.32% (-2.4-4.8%)	0.38% (-2.7-5.3%)	0.076 (-0.53-1)	1 (-7.5-15)
Greenland	1990	Female	80 to 84	69% (51-82%)	31% (18-49%)	1.3 (0.61-2.3)	-0.8% (-4-4.1%)	-0.87% (-4.2-4.2%)	-0.17 (-0.83-0.82)	-1.7 (-8.5-8.6)
Greenland	1990	Female	85 to 89	67% (49-80%)	33% (20-51%)	1.2 (0.6-2.3)	-0.96% (-3.8-3.4%)	-1.1% (-4.2-3.5%)	-0.095 (-0.36-0.31)	-0.66 (-2.6-2.4)
Greenland	1990	Female	90 to 94	67% (51-80%)	33% (20-49%)	1.3 (0.63-2.5)	-1% (-4.4-3.6%)	-1.2% (-4.8-3.8%)	-0.033 (-0.13-0.1)	-0.16 (-0.65-0.54)
Greenland	1990	Female	95 plus	69% (51-84%)	31% (16-49%)	1.4 (0.62-2.7)	-1.4% (-5-4.7%)	-1.6% (-5.7-4.9%)	-0.0094 (-0.033-0.028)	-0.03 (-0.11-0.1)
Greenland	1990	Male	All Ages	82% (76-87%)	18% (13-24%)	2.2 (1.4-3.3)	11% (6.1-16%)	10% (5.3-16%)	30 (15-47)	1600 (880-2500)
Greenland	1990	Male	15 to 19	68% (52-82%)	32% (18-48%)	1.7 (0.78-3.1)	11% (2.9-22%)	12% (2.8-24%)	1.9 (0.41-3.6)	130 (34-260)
Greenland	1990	Male	20 to 24	84% (74-91%)	16% (8.5-26%)	3.4 (1.8-5.8)	21% (9.2-34%)	23% (9.1-39%)	5.7 (2.2-9.7)	400 (170-660)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	1990	Male	25 to 29	89% (82-94%)	11% (5.9-18%)	3.6 (2-5.9)	20% (9.8-32%)	23% (9.5-37%)	4.4 (1.9-7)	300 (140-460)
Greenland	1990	Male	30 to 34	92% (85-96%)	8.2% (4-15%)	3.4 (1.9-5.6)	19% (9.8-28%)	22% (10-34%)	3.3 (1.5-5.2)	210 (110-310)
Greenland	1990	Male	35 to 39	92% (87-96%)	7.5% (3.6-13%)	3 (1.6-5.2)	16% (9.7-24%)	20% (11-31%)	2.4 (1.3-3.8)	140 (80-210)
Greenland	1990	Male	40 to 44	93% (87-96%)	7.3% (3.5-13%)	2.8 (1.4-4.9)	14% (8.4-22%)	18% (9.3-28%)	2.5 (1.3-4)	130 (71-200)
Greenland	1990	Male	45 to 49	93% (85-96%)	7.4% (3.5-15%)	2.7 (1.4-4.5)	12% (7.6-19%)	16% (8.9-25%)	1.9 (1-3.1)	89 (54-140)
Greenland	1990	Male	50 to 54	92% (84-96%)	7.8% (3.7-16%)	2.5 (1.3-4.3)	10% (5.6-17%)	12% (6-20%)	2.1 (1.1-3.7)	82 (45-140)
Greenland	1990	Male	55 to 59	92% (84-96%)	8.5% (4-16%)	2.3 (1.2-3.9)	8.2% (4.5-14%)	9.2% (4.8-16%)	1.9 (0.99-3.3)	64 (35-110)
Greenland	1990	Male	60 to 64	91% (82-96%)	9.4% (4.2-18%)	2.1 (1.1-3.7)	6% (2.6-11%)	6.5% (2.6-12%)	1.8 (0.7-3.4)	50 (21-94)
Greenland	1990	Male	65 to 69	89% (81-95%)	11% (4.7-19%)	1.9 (1-3.3)	4.1% (1-8.5%)	4.3% (1-9%)	1 (0.24-2.2)	24 (6-50)
Greenland	1990	Male	70 to 74	88% (80-94%)	12% (5.8-20%)	1.6 (0.79-3.1)	2.4% (-0.082-6.5%)	2.5% (-0.2-6.9%)	0.62 (-0.049-1.7)	12 (-0.41-32)
Greenland	1990	Male	75 to 79	87% (78-94%)	13% (6.3-22%)	1.5 (0.7-2.8)	1.4% (-0.84-5.6%)	1.4% (-1-5.7%)	0.31 (-0.24-1.3)	4.7 (-3-19)
Greenland	1990	Male	80 to 84	86% (75-93%)	14% (6.9-25%)	1.4 (0.72-2.5)	0.75% (-1.8-4.6%)	0.68% (-1.9-4.7%)	0.063 (-0.18-0.44)	0.76 (-1.8-4.7)
Greenland	1990	Male	85 to 89	85% (74-92%)	15% (8.3-26%)	1.4 (0.69-2.4)	0.39% (-1.7-3.6%)	0.25% (-2-3.6%)	0.01 (-0.081-0.14)	0.12 (-0.55-1.1)
Greenland	1990	Male	90 to 94	85% (75-92%)	15% (7.8-25%)	1.4 (0.68-2.6)	-0.23% (-2.4-3%)	-0.48% (-2.9-3%)	-0.0068 (-0.041-0.042)	-0.018 (-0.19-0.23)
Greenland	1990	Male	95 plus	85% (73-93%)	15% (6.8-27%)	1.4 (0.64-2.7)	-0.82% (-3.3-2.9%)	-1.2% (-4-2.7%)	-0.0041 (-0.014-0.0091)	-0.01 (-0.043-0.037)
Greenland	1995	Female	All Ages	74% (67-80%)	26% (20-33%)	1.7 (1-2.6)	5.3% (2.9-8.5%)	5% (1.9-9.4%)	8.8 (3.4-17)	420 (230-690)
Greenland	1995	Female	15 to 19	65% (49-79%)	35% (21-51%)	1.5 (0.7-2.8)	7% (1.7-14%)	11% (2-23%)	0.35 (0.062-0.75)	26 (6-55)
Greenland	1995	Female	20 to 24	80% (66-89%)	20% (11-34%)	2.9 (1.5-4.9)	9.6% (4-17%)	19% (6-34%)	0.37 (0.12-0.67)	29 (11-49)
Greenland	1995	Female	25 to 29	84% (74-91%)	16% (8.8-26%)	3 (1.6-5)	10% (4.4-17%)	20% (7.2-34%)	0.84 (0.27-1.5)	59 (25-98)
Greenland	1995	Female	30 to 34	86% (77-93%)	14% (6.9-23%)	2.9 (1.6-4.8)	8.8% (4.2-14%)	17% (6.8-29%)	0.81 (0.31-1.4)	53 (25-87)
Greenland	1995	Female	35 to 39	87% (78-94%)	13% (6.4-22%)	2.6 (1.4-4.3)	8.3% (4.3-13%)	15% (6.9-24%)	0.69 (0.31-1.2)	42 (22-67)
Greenland	1995	Female	40 to 44	87% (79-93%)	13% (6.5-21%)	2.4 (1.3-4.2)	9.5% (5.6-14%)	15% (7.9-23%)	0.93 (0.5-1.5)	48 (27-74)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	1995	Female	45 to 49	86% (76-93%)	14% (7-24%)	2.4 (1.3-4)	9% (5.4-13%)	14% (8-21%)	0.78 (0.44-1.2)	36 (21-55)
Greenland	1995	Female	50 to 54	85% (74-92%)	15% (8.1-26%)	2.2 (1.1-3.8)	8.1% (4.8-13%)	11% (6.5-17%)	1 (0.58-1.6)	39 (23-62)
Greenland	1995	Female	55 to 59	83% (71-91%)	17% (8.8-29%)	2 (1-3.4)	7% (3.7-11%)	8.6% (4.5-14%)	1.2 (0.62-1.9)	39 (21-64)
Greenland	1995	Female	60 to 64	80% (67-90%)	20% (10-33%)	1.7 (0.89-3.1)	4.7% (1.8-8.7%)	5.8% (2.3-10%)	0.97 (0.39-1.8)	27 (11-50)
Greenland	1995	Female	65 to 69	77% (62-88%)	23% (12-38%)	1.5 (0.81-2.7)	2.9% (0.35-6.8%)	3.5% (0.49-7.7%)	0.74 (0.1-1.7)	17 (2.1-39)
Greenland	1995	Female	70 to 74	74% (60-85%)	26% (15-40%)	1.4 (0.64-2.4)	1.5% (-1-5.7%)	1.8% (-1.1-6.5%)	0.35 (-0.2-1.3)	6.1 (-4.1-23)
Greenland	1995	Female	75 to 79	72% (56-84%)	28% (16-44%)	1.2 (0.59-2.4)	0.24% (-2.4-4.5%)	0.29% (-2.6-5%)	0.064 (-0.58-1.1)	0.83 (-7.9-15)
Greenland	1995	Female	80 to 84	69% (51-82%)	31% (18-49%)	1.2 (0.59-2.2)	-0.79% (-3.8-3.9%)	-0.87% (-4-3.9%)	-0.15 (-0.68-0.67)	-1.5 (-7-7)
Greenland	1995	Female	85 to 89	67% (49-80%)	33% (20-51%)	1.2 (0.58-2.2)	-0.95% (-3.7-3.1%)	-1.1% (-4-3.2%)	-0.089 (-0.33-0.26)	-0.62 (-2.4-2)
Greenland	1995	Female	90 to 94	67% (52-80%)	33% (20-48%)	1.3 (0.61-2.4)	-1% (-4.1-3.5%)	-1.2% (-4.6-3.7%)	-0.035 (-0.13-0.11)	-0.17 (-0.67-0.59)
Greenland	1995	Female	95 plus	68% (51-84%)	32% (16-49%)	1.4 (0.6-2.7)	-1.4% (-5-4.3%)	-1.6% (-5.5-4.5%)	-0.012 (-0.042-0.034)	-0.039 (-0.14-0.12)
Greenland	1995	Male	All Ages	81% (76-86%)	19% (14-24%)	2.1 (1.3-3.1)	10% (5.9-15%)	9.2% (5-14%)	23 (13-37)	1200 (700-1800)
Greenland	1995	Male	15 to 19	68% (52-81%)	32% (19-48%)	1.6 (0.75-2.9)	11% (2.7-20%)	12% (2.6-23%)	1.2 (0.26-2.4)	90 (23-170)
Greenland	1995	Male	20 to 24	84% (74-91%)	16% (8.6-26%)	3.3 (1.7-5.5)	20% (8.7-33%)	23% (8.5-38%)	2.4 (0.88-4.1)	170 (70-280)
Greenland	1995	Male	25 to 29	89% (82-94%)	11% (5.9-18%)	3.5 (1.9-5.8)	20% (10-30%)	23% (10-37%)	3.3 (1.4-5.4)	230 (110-350)
Greenland	1995	Male	30 to 34	92% (85-96%)	8.3% (4-15%)	3.3 (1.9-5.6)	18% (9.7-26%)	21% (10-34%)	2.9 (1.4-4.7)	190 (100-290)
Greenland	1995	Male	35 to 39	92% (87-96%)	7.5% (3.6-13%)	3 (1.6-5)	15% (9.6-23%)	20% (11-30%)	2.1 (1.1-3.2)	120 (77-180)
Greenland	1995	Male	40 to 44	93% (87-97%)	7.3% (3.5-13%)	2.7 (1.4-4.7)	14% (8.5-21%)	17% (9.6-27%)	2 (1.1-3.2)	110 (62-160)
Greenland	1995	Male	45 to 49	93% (85-97%)	7.4% (3.5-15%)	2.6 (1.4-4.4)	12% (7.6-18%)	16% (9.2-25%)	1.4 (0.85-2.3)	70 (45-110)
Greenland	1995	Male	50 to 54	92% (84-96%)	7.9% (3.8-16%)	2.4 (1.2-4.1)	9.8% (5.7-16%)	12% (6.3-19%)	2.2 (1.2-3.7)	87 (51-140)
Greenland	1995	Male	55 to 59	91% (84-96%)	8.5% (4.1-16%)	2.2 (1.2-3.8)	8.1% (4.8-13%)	9.4% (5.3-16%)	1.7 (0.96-2.9)	58 (34-96)
Greenland	1995	Male	60 to 64	91% (82-96%)	9.5% (4.2-18%)	2 (1-3.5)	6% (2.8-11%)	6.6% (3-12%)	1.7 (0.75-3)	48 (22-85)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	1995	Male	65 to 69	89% (81-95%)	11% (4.6-19%)	1.9 (0.98-3.2)	4.1% (1.5-8%)	4.4% (1.4-8.7%)	1.3 (0.42-2.5)	30 (11-58)
Greenland	1995	Male	70 to 74	88% (80-94%)	12% (5.8-20%)	1.6 (0.77-2.9)	2.5% (0.038-6.2%)	2.6% (-0.1-6.6%)	0.6 (-0.024-1.6)	11 (0.17-29)
Greenland	1995	Male	75 to 79	87% (78-94%)	13% (6.3-22%)	1.4 (0.68-2.7)	1.4% (-0.69-5.1%)	1.4% (-0.89-5.3%)	0.3 (-0.19-1.1)	4.5 (-2.2-16)
Greenland	1995	Male	80 to 84	86% (76-93%)	14% (6.9-24%)	1.4 (0.69-2.4)	0.69% (-1.7-4.2%)	0.63% (-1.9-4.3%)	0.057 (-0.18-0.39)	0.69 (-1.7-4.2)
Greenland	1995	Male	85 to 89	84% (74-92%)	16% (8.4-26%)	1.3 (0.66-2.3)	0.28% (-1.8-3.5%)	0.16% (-2.1-3.5%)	0.0062 (-0.081-0.14)	0.088 (-0.56-1.1)
Greenland	1995	Male	90 to 94	85% (75-92%)	15% (7.8-25%)	1.4 (0.65-2.5)	-0.34% (-2.5-2.8%)	-0.57% (-2.9-2.7%)	-0.0085 (-0.043-0.04)	-0.028 (-0.21-0.23)
Greenland	1995	Male	95 plus	85% (73-93%)	15% (6.6-27%)	1.4 (0.6-2.7)	-0.94% (-3.4-2.3%)	-1.3% (-4.1-2.2%)	-0.005 (-0.015-0.0083)	-0.014 (-0.049-0.033)
Greenland	2000	Female	All Ages	74% (67-80%)	26% (20-33%)	1.7 (1-2.6)	5.9% (3.3-9.3%)	5.4% (2.4-10%)	10 (4.3-18)	460 (260-740)
Greenland	2000	Female	15 to 19	65% (49-79%)	35% (21-51%)	1.5 (0.71-2.8)	7.1% (1.7-15%)	11% (2.1-23%)	0.41 (0.077-0.84)	31 (7.4-61)
Greenland	2000	Female	20 to 24	80% (67-89%)	20% (11-33%)	2.9 (1.4-5)	11% (4.4-18%)	19% (6.1-34%)	0.49 (0.16-0.88)	37 (15-63)
Greenland	2000	Female	25 to 29	84% (74-91%)	16% (8.7-26%)	3 (1.6-5.2)	12% (5.5-19%)	21% (8.1-35%)	0.64 (0.24-1.1)	43 (19-71)
Greenland	2000	Female	30 to 34	86% (77-93%)	14% (6.9-23%)	2.9 (1.6-4.9)	9.4% (4.4-16%)	18% (7.3-30%)	0.86 (0.35-1.5)	55 (26-91)
Greenland	2000	Female	35 to 39	87% (78-94%)	13% (6.4-22%)	2.6 (1.4-4.3)	8.1% (4.4-13%)	15% (7.2-24%)	0.81 (0.38-1.3)	50 (27-79)
Greenland	2000	Female	40 to 44	87% (79-94%)	13% (6.3-21%)	2.4 (1.3-4.2)	9.3% (5.8-14%)	15% (8.8-23%)	0.97 (0.56-1.5)	51 (31-77)
Greenland	2000	Female	45 to 49	86% (76-93%)	14% (6.9-24%)	2.3 (1.3-4)	9.5% (6.2-14%)	14% (8.8-21%)	1.1 (0.68-1.7)	50 (32-74)
Greenland	2000	Female	50 to 54	85% (74-92%)	15% (8.2-26%)	2.2 (1.1-3.8)	9.1% (5.7-14%)	12% (7.4-18%)	1.2 (0.73-1.9)	47 (29-72)
Greenland	2000	Female	55 to 59	83% (72-91%)	17% (8.9-28%)	2 (1-3.5)	7.2% (4.1-11%)	9% (5.1-14%)	1.3 (0.75-2.1)	44 (25-71)
Greenland	2000	Female	60 to 64	80% (68-90%)	20% (10-32%)	1.8 (0.87-3.1)	5.2% (2.3-9.4%)	6.3% (2.8-11%)	1 (0.45-1.9)	29 (13-53)
Greenland	2000	Female	65 to 69	77% (63-88%)	23% (12-37%)	1.6 (0.79-2.7)	3.3% (0.74-7%)	3.9% (0.96-8.2%)	0.88 (0.22-1.9)	20 (4.5-44)
Greenland	2000	Female	70 to 74	74% (59-85%)	26% (15-41%)	1.4 (0.66-2.4)	1.7% (-0.72-5.6%)	2% (-0.78-6.5%)	0.52 (-0.2-1.6)	9.2 (-3.9-30)
Greenland	2000	Female	75 to 79	72% (56-84%)	28% (16-44%)	1.3 (0.59-2.3)	0.25% (-2.2-4.3%)	0.29% (-2.5-4.7%)	0.067 (-0.56-1)	0.92 (-8-15)
Greenland	2000	Female	80 to 84	69% (52-82%)	31% (18-48%)	1.2 (0.6-2.2)	-0.57% (-3.5-3.8%)	-0.64% (-3.7-4%)	-0.092 (-0.52-0.57)	-0.89 (-5.3-6)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	2000	Female	85 to 89	67% (50-80%)	33% (20-50%)	1.2 (0.58-2.2)	-0.89% (-3.7-3.3%)	-1% (-4-3.4%)	-0.1 (-0.4-0.35)	-0.69 (-3-2.6)
Greenland	2000	Female	90 to 94	67% (52-81%)	33% (19-48%)	1.3 (0.62-2.3)	-0.85% (-3.8-3.6%)	-1% (-4.3-3.8%)	-0.032 (-0.13-0.12)	-0.15 (-0.67-0.63)
Greenland	2000	Female	95 plus	69% (50-84%)	31% (16-50%)	1.4 (0.6-2.6)	-1.2% (-4.8-4.6%)	-1.5% (-5.3-4.8%)	-0.013 (-0.048-0.042)	-0.04 (-0.16-0.15)
Greenland	2000	Male	All Ages	81% (76-87%)	19% (13-24%)	2 (1.2-3.1)	10% (6.3-15%)	9.5% (5.5-15%)	24 (13-37)	1200 (710-1700)
Greenland	2000	Male	15 to 19	68% (53-82%)	32% (18-47%)	1.6 (0.78-3)	11% (2.8-20%)	12% (2.7-24%)	1.1 (0.23-2.2)	82 (21-160)
Greenland	2000	Male	20 to 24	84% (73-92%)	16% (8.4-27%)	3.4 (1.7-5.9)	21% (9.1-34%)	24% (8.8-39%)	2.2 (0.85-3.7)	160 (69-260)
Greenland	2000	Male	25 to 29	89% (82-94%)	11% (6.2-18%)	3.6 (1.9-6)	20% (11-30%)	24% (11-38%)	1.7 (0.77-2.7)	120 (62-180)
Greenland	2000	Male	30 to 34	92% (85-96%)	8.4% (4.2-15%)	3.4 (1.8-5.8)	18% (10-27%)	23% (11-36%)	2.5 (1.3-4)	160 (94-250)
Greenland	2000	Male	35 to 39	92% (86-96%)	7.6% (3.6-14%)	3 (1.6-5)	16% (9.9-23%)	21% (11-31%)	2.1 (1.1-3.3)	130 (80-190)
Greenland	2000	Male	40 to 44	93% (87-97%)	7.3% (3.4-13%)	2.7 (1.5-4.7)	15% (8.9-21%)	19% (10-28%)	2.4 (1.4-3.6)	130 (78-180)
Greenland	2000	Male	45 to 49	93% (85-97%)	7.3% (3.4-15%)	2.6 (1.4-4.4)	13% (8.6-19%)	17% (10-25%)	2.1 (1.3-3.2)	97 (62-140)
Greenland	2000	Male	50 to 54	92% (84-96%)	7.7% (3.8-16%)	2.5 (1.2-4.1)	11% (6.8-17%)	13% (7.9-21%)	2.1 (1.2-3.4)	84 (52-130)
Greenland	2000	Male	55 to 59	92% (84-96%)	8.5% (4-16%)	2.3 (1.2-3.9)	8.9% (5.5-14%)	10% (6.2-16%)	2.5 (1.5-4.1)	85 (51-130)
Greenland	2000	Male	60 to 64	91% (82-96%)	9.4% (4.3-18%)	2.1 (1-3.7)	6.6% (3.5-11%)	7.4% (3.8-13%)	2 (1-3.6)	57 (29-98)
Greenland	2000	Male	65 to 69	89% (81-95%)	11% (4.7-19%)	1.9 (0.95-3.2)	4.8% (2-8.7%)	5.3% (2.2-9.6%)	1.5 (0.61-2.8)	35 (16-64)
Greenland	2000	Male	70 to 74	88% (80-94%)	12% (5.8-20%)	1.6 (0.81-2.9)	2.9% (0.56-6.6%)	3.1% (0.49-7%)	0.98 (0.15-2.2)	18 (3.4-42)
Greenland	2000	Male	75 to 79	87% (78-94%)	13% (6.4-22%)	1.5 (0.7-2.7)	1.7% (-0.37-5.1%)	1.8% (-0.57-5.4%)	0.38 (-0.12-1.2)	5.6 (-1.2-17)
Greenland	2000	Male	80 to 84	86% (76-93%)	14% (6.9-24%)	1.4 (0.72-2.4)	0.91% (-1.3-4.5%)	0.86% (-1.4-4.6%)	0.066 (-0.11-0.35)	0.77 (-1.1-3.8)
Greenland	2000	Male	85 to 89	85% (75-92%)	15% (8.5-25%)	1.3 (0.67-2.4)	0.28% (-1.9-3.7%)	0.18% (-2.2-3.7%)	0.0093 (-0.11-0.19)	0.12 (-0.79-1.5)
Greenland	2000	Male	90 to 94	85% (76-92%)	15% (7.8-24%)	1.4 (0.65-2.5)	-0.17% (-2.3-2.9%)	-0.38% (-2.7-2.9%)	-0.0057 (-0.041-0.044)	-0.014 (-0.19-0.25)
Greenland	2000	Male	95 plus	85% (73-94%)	15% (6.4-27%)	1.4 (0.62-2.8)	-0.81% (-3.3-2.4%)	-1.2% (-3.9-2.3%)	-0.0047 (-0.016-0.0095)	-0.012 (-0.05-0.037)
Greenland	2005	Female	All Ages	74% (67-80%)	26% (20-33%)	1.7 (1.1-2.7)	6% (3.6-9%)	5.8% (3-9.7%)	11 (5.4-19)	480 (290-720)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	2005	Female	15 to 19	65% (49-79%)	35% (21-51%)	1.6 (0.75-2.8)	7.1% (1.7-14%)	11% (2.1-23%)	0.44 (0.083-0.93)	33 (8.1-68)
Greenland	2005	Female	20 to 24	80% (67-89%)	20% (11-33%)	3 (1.5-5.2)	11% (4.5-20%)	20% (6-35%)	0.64 (0.19-1.1)	47 (18-81)
Greenland	2005	Female	25 to 29	84% (74-91%)	16% (8.9-26%)	3.1 (1.7-5.4)	9.5% (4.6-15%)	21% (8.1-36%)	0.36 (0.13-0.6)	27 (12-42)
Greenland	2005	Female	30 to 34	86% (77-93%)	14% (7.1-23%)	3 (1.6-5.2)	11% (5-17%)	18% (7.4-30%)	0.66 (0.26-1.1)	41 (19-66)
Greenland	2005	Female	35 to 39	87% (78-93%)	13% (7.1-22%)	2.7 (1.5-4.4)	7.8% (4.5-12%)	16% (8.2-25%)	0.64 (0.33-1)	40 (23-60)
Greenland	2005	Female	40 to 44	87% (79-93%)	13% (6.9-21%)	2.5 (1.4-4.3)	9% (5.8-13%)	16% (9.9-24%)	1 (0.62-1.5)	55 (35-79)
Greenland	2005	Female	45 to 49	86% (75-93%)	14% (7-25%)	2.4 (1.4-4.1)	10% (6.9-15%)	15% (10-22%)	1.4 (0.92-2.1)	64 (44-92)
Greenland	2005	Female	50 to 54	85% (74-92%)	15% (8.2-26%)	2.3 (1.2-3.9)	9.8% (6.4-14%)	13% (8.5-19%)	1.6 (1-2.4)	62 (40-92)
Greenland	2005	Female	55 to 59	83% (71-91%)	17% (8.9-29%)	2 (1.1-3.4)	7.8% (4.7-12%)	10% (6.2-15%)	1.2 (0.73-1.9)	41 (25-62)
Greenland	2005	Female	60 to 64	80% (67-90%)	20% (9.9-33%)	1.8 (0.87-3.1)	5.6% (2.9-9.4%)	6.9% (3.7-11%)	1.2 (0.67-2)	35 (18-59)
Greenland	2005	Female	65 to 69	77% (62-88%)	23% (12-38%)	1.6 (0.82-2.8)	3.7% (1.5-6.9%)	4.5% (1.8-8.1%)	0.93 (0.37-1.7)	21 (8.1-39)
Greenland	2005	Female	70 to 74	74% (60-85%)	26% (15-40%)	1.4 (0.7-2.5)	2.1% (-0.23-5.5%)	2.4% (-0.18-6.2%)	0.67 (-0.052-1.7)	12 (-1.3-32)
Greenland	2005	Female	75 to 79	72% (57-84%)	28% (16-43%)	1.3 (0.63-2.3)	0.62% (-1.6-4.4%)	0.71% (-1.8-4.7%)	0.21 (-0.55-1.4)	2.9 (-7.8-20)
Greenland	2005	Female	80 to 84	69% (52-82%)	31% (18-48%)	1.3 (0.63-2.3)	-0.22% (-3-3.9%)	-0.27% (-3.2-4%)	-0.037 (-0.43-0.55)	-0.33 (-4.4-5.8)
Greenland	2005	Female	85 to 89	67% (50-80%)	33% (20-50%)	1.3 (0.62-2.2)	-0.57% (-3.3-3.8%)	-0.67% (-3.6-3.9%)	-0.06 (-0.32-0.35)	-0.4 (-2.3-2.6)
Greenland	2005	Female	90 to 94	67% (52-80%)	33% (20-48%)	1.4 (0.64-2.4)	-0.51% (-3.6-3.9%)	-0.66% (-3.9-4.2%)	-0.02 (-0.12-0.12)	-0.087 (-0.59-0.66)
Greenland	2005	Female	95 plus	69% (50-84%)	31% (16-50%)	1.5 (0.64-2.7)	-0.83% (-4.4-4.8%)	-1.1% (-5-5.1%)	-0.0094 (-0.044-0.045)	-0.028 (-0.15-0.16)
Greenland	2005	Male	All Ages	82% (76-87%)	18% (13-24%)	2.1 (1.3-3.2)	11% (6.4-15%)	9.7% (5.8-15%)	25 (15-38)	1200 (700-1700)
Greenland	2005	Male	15 to 19	68% (53-82%)	32% (18-47%)	1.7 (0.82-3.1)	10% (2.8-20%)	13% (2.6-25%)	0.98 (0.2-2)	74 (20-140)
Greenland	2005	Male	20 to 24	85% (74-91%)	15% (8.5-26%)	3.5 (1.8-6)	22% (9.4-35%)	24% (9.3-41%)	2.9 (1.1-5)	200 (85-340)
Greenland	2005	Male	25 to 29	89% (82-94%)	11% (6-18%)	3.7 (1.9-6.4)	21% (11-31%)	25% (11-40%)	1.4 (0.6-2.2)	100 (51-150)
Greenland	2005	Male	30 to 34	92% (85-96%)	8.2% (4-15%)	3.5 (1.9-6.2)	18% (10-27%)	24% (11-37%)	1.3 (0.61-2)	87 (49-130)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	2005	Male	35 to 39	92% (86-96%)	7.5% (3.7-14%)	3.1 (1.7-5.1)	17% (10-24%)	22% (12-33%)	2.3 (1.2-3.4)	130 (82-190)
Greenland	2005	Male	40 to 44	93% (87-97%)	7.3% (3.5-13%)	2.8 (1.6-4.9)	15% (9.4-22%)	19% (11-29%)	2.6 (1.5-4)	140 (86-200)
Greenland	2005	Male	45 to 49	93% (86-97%)	7.4% (3.5-14%)	2.7 (1.6-4.5)	14% (8.9-20%)	18% (11-26%)	2.5 (1.5-3.6)	110 (74-160)
Greenland	2005	Male	50 to 54	92% (85-96%)	7.8% (3.7-15%)	2.6 (1.4-4.3)	11% (7.4-17%)	14% (8.9-23%)	2 (1.3-3.2)	83 (54-130)
Greenland	2005	Male	55 to 59	92% (84-96%)	8.5% (4-16%)	2.3 (1.2-4)	9.7% (6.2-15%)	12% (7.3-18%)	2.7 (1.7-4.2)	92 (59-140)
Greenland	2005	Male	60 to 64	91% (82-96%)	9.4% (4.2-18%)	2.1 (1.1-3.6)	7% (3.9-12%)	8.1% (4.4-14%)	2.3 (1.2-3.9)	66 (36-110)
Greenland	2005	Male	65 to 69	90% (81-95%)	10% (4.7-19%)	1.9 (1-3.3)	5.2% (2.3-9.1%)	5.7% (2.5-10%)	1.8 (0.76-3.2)	42 (18-74)
Greenland	2005	Male	70 to 74	88% (81-94%)	12% (5.8-19%)	1.7 (0.86-2.9)	3.3% (0.8-7%)	3.5% (0.77-7.6%)	1.2 (0.28-2.7)	23 (5.8-50)
Greenland	2005	Male	75 to 79	87% (78-94%)	13% (6.2-22%)	1.5 (0.77-2.7)	1.8% (-0.34-5.3%)	1.9% (-0.46-5.7%)	0.55 (-0.14-1.6)	8.1 (-1.6-23)
Greenland	2005	Male	80 to 84	86% (75-93%)	14% (6.8-25%)	1.4 (0.74-2.5)	1.1% (-1.1-4.4%)	1% (-1.3-4.6%)	0.087 (-0.11-0.38)	0.98 (-1-4)
Greenland	2005	Male	85 to 89	85% (74-92%)	15% (8.3-26%)	1.4 (0.7-2.4)	0.62% (-1.4-3.9%)	0.55% (-1.6-4.1%)	0.019 (-0.054-0.14)	0.17 (-0.36-1.1)
Greenland	2005	Male	90 to 94	85% (75-92%)	15% (7.7-25%)	1.4 (0.7-2.6)	0.18% (-1.8-3.3%)	-0.0061% (-2.2-3.4%)	-0.000086 (-0.027-0.041)	0.012 (-0.12-0.23)
Greenland	2005	Male	95 plus	85% (72-93%)	15% (6.8-28%)	1.4 (0.66-2.8)	-0.41% (-2.8-3%)	-0.75% (-3.5-3%)	-0.0023 (-0.011-0.0091)	-0.0048 (-0.033-0.035)
Greenland	2010	Female	All Ages	75% (68-80%)	25% (20-32%)	1.9 (1.2-2.8)	6% (3.7-8.9%)	6.3% (3.6-10%)	11 (6-17)	450 (280-670)
Greenland	2010	Female	15 to 19	64% (47-79%)	36% (21-53%)	1.6 (0.76-2.9)	7.5% (1.8-15%)	12% (2.1-24%)	0.53 (0.092-1.2)	40 (9-83)
Greenland	2010	Female	20 to 24	80% (65-89%)	20% (11-35%)	3.1 (1.6-5.4)	10% (4.2-17%)	20% (6.4-35%)	0.52 (0.17-0.94)	40 (17-69)
Greenland	2010	Female	25 to 29	84% (73-91%)	16% (8.7-27%)	3.2 (1.7-5.5)	7.5% (3.9-12%)	22% (8.9-36%)	0.25 (0.1-0.43)	20 (10-32)
Greenland	2010	Female	30 to 34	86% (77-93%)	14% (6.9-23%)	3.1 (1.7-5.5)	9.4% (4.4-15%)	19% (7.7-31%)	0.47 (0.19-0.79)	30 (14-49)
Greenland	2010	Female	35 to 39	87% (78-94%)	13% (6.5-22%)	2.8 (1.5-4.7)	8% (4.6-12%)	17% (8.6-26%)	0.38 (0.2-0.61)	23 (14-37)
Greenland	2010	Female	40 to 44	87% (78-93%)	13% (6.6-22%)	2.6 (1.4-4.4)	8.4% (5.6-12%)	17% (10-24%)	0.81 (0.48-1.2)	45 (29-66)
Greenland	2010	Female	45 to 49	86% (76-93%)	14% (6.8-24%)	2.5 (1.4-4.2)	9.8% (6.7-14%)	15% (10-21%)	1.7 (1.1-2.4)	76 (51-110)
Greenland	2010	Female	50 to 54	85% (74-92%)	15% (8.3-26%)	2.3 (1.2-3.9)	8.6% (5.8-12%)	13% (8.5-18%)	1.3 (0.87-1.9)	53 (36-77)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	2010	Female	55 to 59	83% (71-91%)	17% (9-29%)	2.1 (1.1-3.5)	7.5% (4.6-11%)	9.9% (6.3-15%)	1.4 (0.89-2.2)	48 (30-73)
Greenland	2010	Female	60 to 64	80% (67-90%)	20% (10-33%)	1.9 (0.95-3.2)	5.5% (3-9%)	7% (4-11%)	1 (0.58-1.7)	29 (16-48)
Greenland	2010	Female	65 to 69	77% (62-88%)	23% (12-38%)	1.7 (0.86-2.8)	3.8% (1.6-7%)	4.7% (2.1-8%)	1.1 (0.49-1.9)	25 (11-45)
Greenland	2010	Female	70 to 74	74% (59-85%)	26% (15-41%)	1.4 (0.71-2.6)	2.3% (0.12-5.7%)	2.7% (0.22-6.5%)	0.67 (0.053-1.6)	12 (0.59-30)
Greenland	2010	Female	75 to 79	71% (55-84%)	29% (16-45%)	1.3 (0.63-2.4)	0.98% (-1.1-4.3%)	1.1% (-1.2-4.8%)	0.32 (-0.35-1.3)	4.5 (-5.1-19)
Greenland	2010	Female	80 to 84	68% (50-82%)	32% (18-50%)	1.3 (0.64-2.3)	0.38% (-1.9-3.7%)	0.4% (-2.1-4%)	0.035 (-0.18-0.37)	0.39 (-1.8-4)
Greenland	2010	Female	85 to 89	65% (48-79%)	35% (21-52%)	1.3 (0.63-2.3)	0.11% (-2.2-3.6%)	0.049% (-2.4-3.8%)	0.0022 (-0.11-0.17)	0.039 (-0.75-1.3)
Greenland	2010	Female	90 to 94	66% (52-79%)	34% (21-48%)	1.4 (0.66-2.5)	0.031% (-2.6-3.9%)	-0.087% (-3-4.2%)	-0.0014 (-0.05-0.071)	0.0032 (-0.25-0.38)
Greenland	2010	Female	95 plus	68% (50-84%)	32% (16-50%)	1.5 (0.67-2.9)	-0.2% (-3.6-5%)	-0.41% (-4.2-5.5%)	-0.0023 (-0.024-0.03)	-0.0043 (-0.077-0.11)
Greenland	2010	Male	All Ages	82% (77-87%)	18% (13-23%)	2.2 (1.4-3.3)	10% (6.5-14%)	9.6% (5.8-14%)	23 (14-36)	1100 (680-1600)
Greenland	2010	Male	15 to 19	68% (52-81%)	32% (19-48%)	1.8 (0.86-3.3)	10% (3.1-20%)	13% (3-26%)	0.88 (0.19-1.8)	67 (19-130)
Greenland	2010	Male	20 to 24	84% (73-91%)	16% (8.6-27%)	3.6 (1.9-6.4)	21% (9.4-34%)	25% (9.4-41%)	2.3 (0.88-3.9)	170 (73-270)
Greenland	2010	Male	25 to 29	89% (82-94%)	11% (6.1-18%)	3.8 (2.1-6.4)	21% (11-31%)	25% (11-40%)	1.7 (0.76-2.7)	120 (61-180)
Greenland	2010	Male	30 to 34	92% (85-96%)	8.4% (4.2-15%)	3.6 (2-6.4)	18% (10-26%)	24% (12-38%)	1.1 (0.56-1.7)	76 (43-110)
Greenland	2010	Male	35 to 39	92% (86-96%)	7.6% (3.6-14%)	3.2 (1.8-5.3)	16% (10-23%)	23% (13-34%)	1.2 (0.69-1.8)	74 (47-110)
Greenland	2010	Male	40 to 44	93% (86-97%)	7.3% (3.4-14%)	2.9 (1.6-4.9)	14% (9.3-21%)	20% (12-31%)	1.9 (1.1-2.9)	100 (64-150)
Greenland	2010	Male	45 to 49	93% (85-97%)	7.3% (3.4-15%)	2.8 (1.6-4.6)	14% (9-20%)	18% (11-27%)	2.7 (1.6-4.1)	120 (80-190)
Greenland	2010	Male	50 to 54	92% (85-96%)	7.8% (3.8-15%)	2.6 (1.4-4.5)	11% (7.4-17%)	15% (9.2-23%)	2.5 (1.5-3.9)	100 (66-160)
Greenland	2010	Male	55 to 59	91% (84-96%)	8.5% (4-16%)	2.4 (1.3-4)	9.7% (6.4-14%)	12% (7.7-18%)	2.7 (1.7-4.2)	93 (61-140)
Greenland	2010	Male	60 to 64	90% (82-96%)	9.5% (4.4-18%)	2.2 (1.2-3.8)	7.2% (4.1-12%)	8.6% (4.9-14%)	2.1 (1.2-3.4)	60 (34-99)
Greenland	2010	Male	65 to 69	89% (81-95%)	11% (4.6-19%)	2 (1.1-3.3)	5.5% (2.8-8.9%)	6.2% (3.1-10%)	2.3 (1.1-3.8)	53 (26-88)
Greenland	2010	Male	70 to 74	88% (80-94%)	12% (5.9-20%)	1.7 (0.89-3.1)	3.5% (0.98-7%)	3.8% (0.99-7.7%)	1.3 (0.32-2.7)	24 (6.6-50)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	2010	Male	75 to 79	87% (78-94%)	13% (6.5-22%)	1.6 (0.81-2.8)	2% (-0.28-5.4%)	2.1% (-0.38-5.8%)	0.73 (-0.14-2.1)	11 (-1.5-30)
Greenland	2010	Male	80 to 84	86% (75-93%)	14% (6.9-25%)	1.5 (0.77-2.6)	1.5% (-0.42-4.4%)	1.5% (-0.59-4.7%)	0.08 (-0.033-0.25)	0.92 (-0.26-2.8)
Greenland	2010	Male	85 to 89	85% (75-91%)	15% (8.5-25%)	1.4 (0.76-2.5)	1% (-0.79-3.8%)	0.94% (-1.1-4%)	0.023 (-0.026-0.097)	0.2 (-0.16-0.77)
Greenland	2010	Male	90 to 94	85% (76-92%)	15% (7.9-24%)	1.5 (0.74-2.6)	0.53% (-1.4-3.6%)	0.35% (-1.9-3.8%)	0.003 (-0.015-0.031)	0.025 (-0.068-0.18)
Greenland	2010	Male	95 plus	85% (73-94%)	15% (6.5-27%)	1.5 (0.69-2.8)	-0.029% (-2.3-3.1%)	-0.37% (-3-3.1%)	-0.00088 (-0.007-0.0073)	-0.00024 (-0.021-0.028)
Greenland	2016	Female	All Ages	75% (68-81%)	25% (19-32%)	2 (1.2-3.1)	5.5% (3.5-8.2%)	6.2% (3.7-9.7%)	9.2 (4.9-15)	380 (220-590)
Greenland	2016	Female	15 to 19	63% (47-78%)	37% (22-53%)	1.6 (0.78-2.9)	7% (1.8-14%)	12% (2.4-24%)	0.37 (0.073-0.8)	28 (7.1-58)
Greenland	2016	Female	20 to 24	80% (67-89%)	20% (11-33%)	3.2 (1.6-5.5)	9.2% (4.1-16%)	20% (7-35%)	0.41 (0.14-0.74)	33 (14-54)
Greenland	2016	Female	25 to 29	84% (74-92%)	16% (8.2-26%)	3.3 (1.9-5.5)	6.4% (3.5-9.9%)	21% (8.8-35%)	0.22 (0.087-0.39)	20 (11-31)
Greenland	2016	Female	30 to 34	86% (77-93%)	14% (7.1-23%)	3.2 (1.8-5.4)	8.3% (4.2-13%)	18% (8.3-31%)	0.44 (0.19-0.79)	30 (15-50)
Greenland	2016	Female	35 to 39	87% (77-93%)	13% (6.8-23%)	2.9 (1.6-5)	7.2% (4.3-11%)	17% (8.9-25%)	0.31 (0.15-0.53)	20 (11-32)
Greenland	2016	Female	40 to 44	87% (77-93%)	13% (7.1-23%)	2.7 (1.5-4.6)	7.9% (5.2-11%)	17% (11-25%)	0.34 (0.2-0.53)	20 (13-29)
Greenland	2016	Female	45 to 49	86% (75-93%)	14% (7.1-25%)	2.6 (1.4-4.4)	9.4% (6.3-13%)	16% (11-22%)	1.2 (0.73-1.9)	56 (35-84)
Greenland	2016	Female	50 to 54	85% (73-92%)	15% (8.1-27%)	2.4 (1.3-4)	7.7% (5-11%)	12% (8.2-18%)	1.3 (0.79-2.2)	56 (34-86)
Greenland	2016	Female	55 to 59	83% (70-91%)	17% (9.2-30%)	2.1 (1.2-3.6)	6.8% (4.2-10%)	9.6% (6.1-14%)	1.4 (0.83-2.2)	49 (28-76)
Greenland	2016	Female	60 to 64	80% (66-90%)	20% (10-34%)	1.9 (1-3.3)	5.2% (2.9-8.4%)	6.9% (4-11%)	1.2 (0.64-2)	34 (18-57)
Greenland	2016	Female	65 to 69	77% (62-88%)	23% (12-38%)	1.7 (0.9-2.9)	3.8% (1.7-6.8%)	4.8% (2.3-8.3%)	0.76 (0.35-1.4)	17 (7.5-33)
Greenland	2016	Female	70 to 74	74% (59-85%)	26% (15-41%)	1.5 (0.71-2.6)	2.3% (0.43-5.6%)	2.9% (0.67-6.6%)	0.77 (0.17-1.8)	14 (2.5-32)
Greenland	2016	Female	75 to 79	70% (54-83%)	30% (17-46%)	1.3 (0.63-2.4)	1.2% (-0.73-4.4%)	1.5% (-0.78-5.2%)	0.31 (-0.16-1.1)	4.3 (-2.6-16)
Greenland	2016	Female	80 to 84	68% (50-82%)	32% (18-50%)	1.3 (0.67-2.3)	0.58% (-1.6-4.1%)	0.64% (-1.8-4.4%)	0.041 (-0.12-0.29)	0.44 (-1.3-3.1)
Greenland	2016	Female	85 to 89	65% (50-79%)	35% (21-50%)	1.3 (0.65-2.4)	0.27% (-1.9-3.7%)	0.24% (-2.3-4%)	0.012 (-0.12-0.21)	0.12 (-0.86-1.7)
Greenland	2016	Female	90 to 94	66% (51-80%)	34% (20-49%)	1.4 (0.69-2.6)	0.17% (-2.3-4%)	0.078% (-2.8-4.3%)	0.0016 (-0.054-0.088)	0.02 (-0.27-0.48)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Greenland	2016	Female	95 plus	69% (50-83%)	31% (17-50%)	1.5 (0.69-3)	-0.048% (-3.3-5%)	-0.24% (-3.8-5.4%)	-0.0013 (-0.02-0.029)	-0.00099 (-0.067-0.1)
Greenland	2016	Male	All Ages	83% (77-88%)	17% (12-23%)	2.3 (1.4-3.6)	9.9% (6.5-14%)	9.2% (5.7-14%)	24 (13-38)	1000 (610-1600)
Greenland	2016	Male	15 to 19	67% (51-81%)	33% (19-49%)	1.8 (0.86-3.2)	10% (3-20%)	13% (3.1-26%)	0.69 (0.16-1.5)	53 (15-100)
Greenland	2016	Male	20 to 24	84% (73-92%)	16% (8.3-27%)	3.7 (2-6.4)	21% (9.6-33%)	25% (10-42%)	2.1 (0.78-3.7)	150 (65-260)
Greenland	2016	Male	25 to 29	89% (82-94%)	11% (6-18%)	3.9 (2.2-6.6)	20% (11-30%)	26% (12-41%)	1.7 (0.74-2.8)	120 (62-190)
Greenland	2016	Male	30 to 34	92% (85-96%)	8.5% (4.2-15%)	3.8 (2.1-6.4)	18% (10-27%)	25% (12-38%)	1.3 (0.6-2.1)	86 (47-130)
Greenland	2016	Male	35 to 39	92% (86-96%)	7.6% (3.7-14%)	3.3 (1.8-5.7)	16% (10-23%)	23% (14-35%)	1 (0.53-1.6)	62 (37-94)
Greenland	2016	Male	40 to 44	93% (87-97%)	7.3% (3.5-13%)	3 (1.7-5.2)	14% (9.2-21%)	21% (13-33%)	1 (0.56-1.7)	56 (34-87)
Greenland	2016	Male	45 to 49	93% (85-97%)	7.5% (3.4-15%)	2.9 (1.6-4.7)	14% (9.2-21%)	19% (12-28%)	2.4 (1.3-3.8)	110 (66-170)
Greenland	2016	Male	50 to 54	92% (84-96%)	8% (3.9-16%)	2.7 (1.5-4.5)	11% (7.3-17%)	15% (9.5-23%)	2.7 (1.5-4.8)	110 (66-180)
Greenland	2016	Male	55 to 59	91% (84-96%)	8.7% (4.1-16%)	2.5 (1.4-4.1)	9.5% (6.3-15%)	12% (7.7-19%)	3.3 (1.9-5.5)	110 (67-180)
Greenland	2016	Male	60 to 64	90% (82-96%)	9.6% (4.2-18%)	2.3 (1.2-3.9)	7.1% (4.2-12%)	8.8% (5.1-15%)	2.5 (1.3-4.3)	71 (40-120)
Greenland	2016	Male	65 to 69	89% (81-95%)	11% (4.8-19%)	2 (1.1-3.4)	5.5% (2.8-9.4%)	6.3% (3.2-11%)	2.1 (0.95-3.7)	49 (23-85)
Greenland	2016	Male	70 to 74	88% (79-94%)	12% (6.1-21%)	1.8 (0.9-3.1)	3.5% (1.1-7.4%)	3.9% (1.1-8.3%)	1.9 (0.54-4.1)	36 (10-76)
Greenland	2016	Male	75 to 79	87% (78-94%)	13% (6.3-22%)	1.6 (0.77-2.9)	2.1% (-0.062-5.4%)	2.2% (-0.13-5.7%)	0.76 (-0.056-2)	11 (-0.37-29)
Greenland	2016	Male	80 to 84	86% (75-93%)	14% (7.3-25%)	1.5 (0.77-2.6)	1.5% (-0.32-4.5%)	1.6% (-0.45-4.8%)	0.075 (-0.021-0.23)	0.86 (-0.17-2.5)
Greenland	2016	Male	85 to 89	84% (74-91%)	16% (8.6-26%)	1.5 (0.78-2.6)	1.1% (-0.72-4%)	1% (-0.99-4.2%)	0.028 (-0.026-0.11)	0.24 (-0.16-0.9)
Greenland	2016	Male	90 to 94	85% (75-92%)	15% (8-25%)	1.5 (0.79-2.7)	0.58% (-1.4-3.7%)	0.42% (-1.7-3.8%)	0.0045 (-0.019-0.041)	0.036 (-0.086-0.23)
Greenland	2016	Male	95 plus	85% (73-94%)	15% (6.5-27%)	1.5 (0.75-2.8)	0.022% (-2.3-3.7%)	-0.31% (-3-3.6%)	-0.00071 (-0.0068-0.0081)	0.00017 (-0.02-0.03)
Guam	1990	Female	All Ages	36% (31-43%)	64% (57-69%)	0.31 (0.08-0.65)	0.28% (-0.075-0.82%)	-0.14% (-0.82-0.62%)	-0.3 (-1.8-1.3)	36 (-9.7-99)
Guam	1990	Female	15 to 19	39% (27-54%)	61% (46-73%)	0.45 (0.11-1)	1.1% (0.37-2.3%)	2.1% (0.15-5.4%)	0.057 (0.0041-0.15)	6.6 (2.3-13)
Guam	1990	Female	20 to 24	46% (32-61%)	54% (39-68%)	0.66 (0.16-1.4)	1.8% (0.83-3.3%)	2.9% (0.48-6.7%)	0.1 (0.017-0.23)	13 (5.6-23)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	1990	Female	25 to 29	46% (33-62%)	54% (38-67%)	0.53 (0.14-1.2)	1.6% (0.87-2.7%)	2.5% (0.72-5.2%)	0.11 (0.03-0.23)	13 (6.9-22)
Guam	1990	Female	30 to 34	45% (30-61%)	55% (39-70%)	0.48 (0.11-1.1)	1.1% (0.56-2%)	1.5% (0.37-3.6%)	0.078 (0.018-0.2)	9.1 (4.3-17)
Guam	1990	Female	35 to 39	44% (30-60%)	56% (40-70%)	0.51 (0.12-1.1)	1% (0.45-2.2%)	1.7% (0.53-4.1%)	0.11 (0.033-0.27)	8.4 (3.6-18)
Guam	1990	Female	40 to 44	42% (29-59%)	58% (41-71%)	0.47 (0.11-1.1)	0.69% (0.052-1.7%)	1.2% (0.13-3.3%)	0.096 (0.01-0.27)	5.6 (0.45-14)
Guam	1990	Female	45 to 49	42% (27-59%)	58% (41-73%)	0.38 (0.096-0.84)	0.21% (-0.41-0.99%)	0.52% (-0.43-1.9%)	0.042 (-0.035-0.16)	1.4 (-2.8-6.6)
Guam	1990	Female	50 to 54	42% (28-58%)	58% (42-72%)	0.32 (0.081-0.72)	-0.16% (-0.93-0.52%)	0.022% (-0.99-1.1%)	0.0026 (-0.12-0.13)	-1.2 (-7.1-3.9)
Guam	1990	Female	55 to 59	39% (26-56%)	61% (44-74%)	0.27 (0.065-0.59)	-0.27% (-1.1-0.39%)	-0.093% (-1.1-0.76%)	-0.014 (-0.17-0.12)	-2.2 (-8.2-3.1)
Guam	1990	Female	60 to 64	37% (24-54%)	63% (46-76%)	0.23 (0.055-0.53)	-0.49% (-1.5-0.22%)	-0.43% (-1.6-0.44%)	-0.086 (-0.32-0.088)	-4 (-12-1.9)
Guam	1990	Female	65 to 69	38% (26-52%)	62% (48-74%)	0.23 (0.06-0.51)	-0.49% (-1.3-0.24%)	-0.41% (-1.5-0.48%)	-0.086 (-0.31-0.1)	-3.4 (-9.4-1.7)
Guam	1990	Female	70 to 74	39% (27-54%)	61% (46-73%)	0.22 (0.054-0.49)	-0.76% (-1.7--0.015%)	-0.82% (-2-0.068%)	-0.18 (-0.43-0.016)	-4.3 (-9.2--0.092)
Guam	1990	Female	75 to 79	39% (26-54%)	61% (46-74%)	0.22 (0.052-0.5)	-0.59% (-1.5-0.13%)	-0.59% (-1.7-0.29%)	-0.13 (-0.36-0.062)	-2.4 (-5.9-0.49)
Guam	1990	Female	80 to 84	38% (25-54%)	62% (46-75%)	0.22 (0.052-0.5)	-0.78% (-1.7--0.069%)	-0.86% (-2--0.051%)	-0.17 (-0.39--0.011)	-2.1 (-4.6--0.19)
Guam	1990	Female	85 to 89	39% (26-53%)	61% (47-74%)	0.23 (0.054-0.52)	-0.79% (-1.8--0.012%)	-0.89% (-2.1-0.033%)	-0.14 (-0.33-0.005)	-1.1 (-2.6--0.016)
Guam	1990	Female	90 to 94	41% (28-54%)	59% (46-72%)	0.25 (0.059-0.58)	-0.73% (-1.6--0.021%)	-0.82% (-1.8-0.0035%)	-0.06 (-0.13-0.00026)	-0.33 (-0.71--0.01)
Guam	1990	Female	95 plus	44% (28-61%)	56% (39-72%)	0.29 (0.067-0.69)	-0.76% (-1.9-0.12%)	-0.89% (-2.2-0.18%)	-0.018 (-0.044-0.0034)	-0.061 (-0.15-0.0092)
Guam	1990	Male	All Ages	54% (46-62%)	46% (38-54%)	0.69 (0.19-1.4)	2.6% (1.2-4.9%)	2.2% (0.74-4.6%)	7.8 (2.7-16)	470 (220-890)
Guam	1990	Male	15 to 19	47% (30-64%)	53% (36-70%)	0.53 (0.14-1.2)	2.4% (0.83-5%)	3.3% (0.67-7.6%)	0.27 (0.054-0.64)	24 (8.4-51)
Guam	1990	Male	20 to 24	64% (46-79%)	36% (21-54%)	1.2 (0.33-2.7)	5.9% (2.1-12%)	7.5% (1.7-16%)	1.1 (0.27-2.4)	91 (33-180)
Guam	1990	Male	25 to 29	70% (53-82%)	30% (18-47%)	1.3 (0.33-2.7)	6% (2.6-11%)	7.5% (2.1-15%)	1.1 (0.3-2.1)	82 (35-150)
Guam	1990	Male	30 to 34	71% (52-84%)	29% (16-48%)	1.2 (0.32-2.7)	5.2% (2.4-9.6%)	6.4% (2.2-13%)	0.92 (0.29-1.9)	67 (31-130)
Guam	1990	Male	35 to 39	69% (52-83%)	31% (17-48%)	1.1 (0.27-2.5)	4.8% (2.4-8.6%)	6% (2.4-12%)	0.91 (0.37-1.8)	59 (30-110)
Guam	1990	Male	40 to 44	67% (50-82%)	33% (18-50%)	1 (0.25-2.3)	3.8% (1.9-7.1%)	4.5% (1.9-9.2%)	0.8 (0.33-1.6)	47 (23-89)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	1990	Male	45 to 49	64% (43-80%)	36% (20-57%)	0.91 (0.24-2)	3.1% (1.4-5.8%)	3.6% (1.3-7.2%)	0.64 (0.23-1.3)	33 (15-62)
Guam	1990	Male	50 to 54	62% (43-79%)	38% (21-57%)	0.75 (0.19-1.7)	2.5% (1.1-5%)	2.8% (0.92-6.1%)	0.64 (0.21-1.3)	29 (12-56)
Guam	1990	Male	55 to 59	61% (42-78%)	39% (22-58%)	0.58 (0.14-1.2)	1.4% (0.26-3.3%)	1.4% (0.056-3.8%)	0.43 (0.017-1.1)	17 (3.4-40)
Guam	1990	Male	60 to 64	58% (38-77%)	42% (23-62%)	0.49 (0.12-1.1)	0.83% (-0.1-2.4%)	0.85% (-0.3-2.7%)	0.34 (-0.12-1.1)	11 (-1.4-33)
Guam	1990	Male	65 to 69	56% (37-74%)	44% (26-63%)	0.46 (0.11-1)	0.68% (-0.16-2.2%)	0.73% (-0.28-2.4%)	0.31 (-0.12-1)	7.9 (-1.7-25)
Guam	1990	Male	70 to 74	54% (35-70%)	46% (30-65%)	0.42 (0.096-0.95)	0.41% (-0.37-1.7%)	0.42% (-0.48-1.9%)	0.13 (-0.15-0.57)	2.7 (-2.3-11)
Guam	1990	Male	75 to 79	50% (32-69%)	50% (31-68%)	0.37 (0.087-0.86)	0.33% (-0.38-1.5%)	0.32% (-0.52-1.6%)	0.081 (-0.13-0.42)	1.4 (-1.6-6.4)
Guam	1990	Male	80 to 84	47% (28-65%)	53% (35-72%)	0.34 (0.08-0.78)	0.36% (-0.3-1.3%)	0.36% (-0.41-1.5%)	0.071 (-0.079-0.28)	0.87 (-0.72-3.2)
Guam	1990	Male	85 to 89	45% (29-62%)	55% (38-71%)	0.32 (0.074-0.77)	0.16% (-0.48-1%)	0.13% (-0.61-1.1%)	0.013 (-0.061-0.12)	0.15 (-0.42-0.93)
Guam	1990	Male	90 to 94	47% (31-64%)	53% (36-69%)	0.34 (0.078-0.77)	0.13% (-0.5-0.88%)	0.086% (-0.65-0.92%)	0.0028 (-0.022-0.03)	0.026 (-0.097-0.18)
Guam	1990	Male	95 plus	50% (30-70%)	50% (30-70%)	0.38 (0.086-0.94)	-0.11% (-0.88-0.87%)	-0.26% (-1.2-0.81%)	-0.0019 (-0.0088-0.0061)	-0.0032 (-0.027-0.026)
Guam	1995	Female	All Ages	36% (31-43%)	64% (57-69%)	0.32 (0.071-0.63)	0.23% (-0.17-0.75%)	-0.22% (-0.93-0.53%)	-0.6 (-2.5-1.4)	33 (-23-110)
Guam	1995	Female	15 to 19	39% (28-54%)	61% (46-72%)	0.46 (0.098-1)	1.1% (0.4-2.3%)	2.2% (0.17-5.4%)	0.061 (0.0048-0.15)	6.9 (2.5-14)
Guam	1995	Female	20 to 24	46% (32-61%)	54% (39-68%)	0.67 (0.14-1.4)	1.8% (0.83-3.2%)	2.9% (0.5-6.6%)	0.1 (0.017-0.23)	13 (5.9-23)
Guam	1995	Female	25 to 29	46% (32-62%)	54% (38-68%)	0.55 (0.11-1.2)	1.6% (0.84-2.7%)	2.5% (0.74-5.3%)	0.12 (0.033-0.25)	13 (6.9-23)
Guam	1995	Female	30 to 34	45% (31-61%)	55% (39-69%)	0.5 (0.1-1.1)	1.1% (0.54-2%)	1.5% (0.37-3.7%)	0.095 (0.022-0.24)	11 (4.8-20)
Guam	1995	Female	35 to 39	44% (29-60%)	56% (40-71%)	0.52 (0.11-1.2)	1% (0.41-2.2%)	1.7% (0.53-4%)	0.13 (0.04-0.33)	10 (3.8-21)
Guam	1995	Female	40 to 44	42% (29-59%)	58% (41-71%)	0.48 (0.11-1)	0.65% (-0.0035-1.8%)	1.2% (0.087-3.1%)	0.12 (0.0083-0.32)	6.6 (-0.037-18)
Guam	1995	Female	45 to 49	42% (27-59%)	58% (41-73%)	0.39 (0.084-0.83)	0.16% (-0.52-0.95%)	0.45% (-0.52-1.8%)	0.055 (-0.064-0.22)	1.6 (-5.3-9.3)
Guam	1995	Female	50 to 54	42% (28-59%)	58% (41-72%)	0.33 (0.069-0.72)	-0.22% (-1.1-0.53%)	-0.043% (-1.1-1.1%)	-0.0059 (-0.15-0.14)	-1.8 (-8.9-4.4)
Guam	1995	Female	55 to 59	39% (26-55%)	61% (45-74%)	0.28 (0.06-0.61)	-0.34% (-1.1-0.39%)	-0.17% (-1.2-0.77%)	-0.032 (-0.22-0.14)	-3.2 (-11-3.7)
Guam	1995	Female	60 to 64	37% (24-54%)	63% (46-76%)	0.24 (0.049-0.51)	-0.56% (-1.5-0.21%)	-0.51% (-1.7-0.4%)	-0.12 (-0.42-0.098)	-5.5 (-15-2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	1995	Female	65 to 69	38% (26-54%)	62% (46-74%)	0.24 (0.05-0.52)	-0.58% (-1.6-0.24%)	-0.52% (-1.7-0.44%)	-0.15 (-0.48-0.12)	-5.4 (-14-2.1)
Guam	1995	Female	70 to 74	40% (27-55%)	60% (45-73%)	0.23 (0.052-0.5)	-0.85% (-1.9--0.024%)	-0.93% (-2.2-0.047%)	-0.26 (-0.63-0.012)	-6 (-13--0.16)
Guam	1995	Female	75 to 79	39% (27-55%)	61% (45-73%)	0.22 (0.047-0.52)	-0.63% (-1.5-0.14%)	-0.63% (-1.7-0.29%)	-0.17 (-0.48-0.079)	-3.2 (-7.8-0.75)
Guam	1995	Female	80 to 84	38% (26-54%)	62% (46-74%)	0.23 (0.047-0.51)	-0.81% (-1.8--0.04%)	-0.9% (-2--0.00095%)	-0.23 (-0.52--0.00022)	-2.8 (-6--0.14)
Guam	1995	Female	85 to 89	39% (26-53%)	61% (47-74%)	0.24 (0.049-0.52)	-0.83% (-1.8-0.022%)	-0.93% (-2.2-0.016%)	-0.19 (-0.42-0.0032)	-1.5 (-3.4--0.041)
Guam	1995	Female	90 to 94	41% (28-55%)	59% (45-72%)	0.26 (0.056-0.62)	-0.77% (-1.6--0.04%)	-0.87% (-1.9--0.02%)	-0.088 (-0.19--0.0021)	-0.48 (-1-0.026)
Guam	1995	Female	95 plus	44% (28-61%)	56% (39-72%)	0.29 (0.058-0.69)	-0.77% (-1.9-0.14%)	-0.91% (-2.3-0.14%)	-0.028 (-0.069-0.0041)	-0.095 (-0.23-0.018)
Guam	1995	Male	All Ages	53% (45-61%)	47% (39-55%)	0.69 (0.15-1.4)	2.6% (1.2-4.7%)	2.1% (0.75-4.5%)	8.7 (3.1-19)	520 (240-980)
Guam	1995	Male	15 to 19	47% (29-64%)	53% (36-71%)	0.53 (0.11-1.2)	2.5% (0.77-5.2%)	3.4% (0.63-7.9%)	0.26 (0.049-0.63)	23 (7.2-49)
Guam	1995	Male	20 to 24	63% (45-78%)	37% (22-55%)	1.2 (0.26-2.6)	6% (2-12%)	7.6% (1.7-16%)	1 (0.23-2.1)	78 (26-150)
Guam	1995	Male	25 to 29	69% (52-82%)	31% (18-48%)	1.3 (0.3-2.7)	6% (2.4-11%)	7.5% (2.1-15%)	1.2 (0.33-2.5)	94 (38-170)
Guam	1995	Male	30 to 34	70% (52-84%)	30% (16-48%)	1.3 (0.3-2.7)	5.2% (2.3-9.6%)	6.3% (2-13%)	1.1 (0.34-2.2)	78 (35-150)
Guam	1995	Male	35 to 39	69% (52-83%)	31% (17-48%)	1.2 (0.24-2.6)	4.7% (2.4-8.5%)	5.9% (2.4-11%)	1 (0.41-2)	67 (34-120)
Guam	1995	Male	40 to 44	67% (51-82%)	33% (18-49%)	1.1 (0.22-2.3)	3.8% (1.9-7%)	4.5% (1.9-8.9%)	0.95 (0.4-1.9)	55 (27-100)
Guam	1995	Male	45 to 49	64% (43-79%)	36% (21-57%)	0.92 (0.2-2)	3.1% (1.5-5.6%)	3.5% (1.4-7%)	0.8 (0.3-1.6)	42 (20-78)
Guam	1995	Male	50 to 54	62% (42-78%)	38% (22-58%)	0.76 (0.17-1.7)	2.6% (1.2-5.2%)	2.9% (1.1-6.4%)	0.68 (0.24-1.4)	31 (14-61)
Guam	1995	Male	55 to 59	60% (42-78%)	40% (22-58%)	0.59 (0.13-1.3)	1.4% (0.34-3.2%)	1.5% (0.076-3.8%)	0.48 (0.024-1.2)	19 (4.5-43)
Guam	1995	Male	60 to 64	58% (38-76%)	42% (24-62%)	0.49 (0.1-1.1)	0.91% (0.006-2.3%)	0.96% (-0.2-2.7%)	0.34 (-0.071-0.95)	11 (0.076-29)
Guam	1995	Male	65 to 69	56% (37-74%)	44% (26-63%)	0.46 (0.09-1)	0.72% (-0.1-2.1%)	0.77% (-0.2-2.5%)	0.41 (-0.11-1.3)	11 (-1.5-31)
Guam	1995	Male	70 to 74	54% (35-71%)	46% (29-65%)	0.42 (0.089-0.96)	0.45% (-0.38-1.7%)	0.47% (-0.5-2%)	0.22 (-0.24-0.92)	4.7 (-3.7-18)
Guam	1995	Male	75 to 79	50% (33-68%)	50% (32-67%)	0.38 (0.074-0.84)	0.37% (-0.4-1.5%)	0.36% (-0.52-1.7%)	0.11 (-0.15-0.48)	1.8 (-2-7.5)
Guam	1995	Male	80 to 84	47% (29-65%)	53% (35-71%)	0.34 (0.07-0.77)	0.39% (-0.26-1.4%)	0.4% (-0.38-1.5%)	0.097 (-0.09-0.37)	1.2 (-0.79-4.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	1995	Male	85 to 89	45% (28-62%)	55% (38-72%)	0.33 (0.069-0.74)	0.18% (-0.45-1%)	0.15% (-0.6-1.1%)	0.022 (-0.088-0.16)	0.24 (-0.58-1.3)
Guam	1995	Male	90 to 94	46% (31-64%)	54% (36-69%)	0.34 (0.075-0.86)	0.15% (-0.46-0.95%)	0.11% (-0.61-1%)	0.0055 (-0.032-0.052)	0.046 (-0.14-0.29)
Guam	1995	Male	95 plus	50% (30-69%)	50% (31-70%)	0.38 (0.074-0.93)	-0.098% (-0.83-0.92%)	-0.24% (-1.1-0.88%)	-0.0026 (-0.012-0.0091)	-0.0042 (-0.035-0.038)
Guam	2000	Female	All Ages	33% (29-38%)	67% (62-71%)	0.32 (0.099-0.61)	0.11% (-0.26-0.61%)	-0.43% (-1.2-0.35%)	-1.3 (-3.5-1.1)	17 (-42-96)
Guam	2000	Female	15 to 19	37% (26-51%)	63% (49-74%)	0.49 (0.15-1)	1.1% (0.4-2.1%)	2.1% (0.24-5.4%)	0.062 (0.0069-0.15)	7.2 (2.7-14)
Guam	2000	Female	20 to 24	43% (32-57%)	57% (43-68%)	0.71 (0.2-1.4)	1.7% (0.84-3.1%)	2.9% (0.59-6.4%)	0.095 (0.019-0.21)	12 (5.9-22)
Guam	2000	Female	25 to 29	42% (32-56%)	58% (44-68%)	0.57 (0.17-1.1)	1.5% (0.83-2.5%)	2.3% (0.67-4.7%)	0.098 (0.027-0.21)	12 (6.6-21)
Guam	2000	Female	30 to 34	41% (31-55%)	59% (45-69%)	0.51 (0.16-1)	1.1% (0.49-1.9%)	1.4% (0.31-3.3%)	0.078 (0.017-0.19)	9.9 (4.6-18)
Guam	2000	Female	35 to 39	39% (29-53%)	61% (47-71%)	0.52 (0.15-1.1)	0.96% (0.4-1.9%)	1.5% (0.44-3.6%)	0.13 (0.034-0.28)	10 (4.3-20)
Guam	2000	Female	40 to 44	34% (24-47%)	66% (53-76%)	0.43 (0.12-0.91)	0.62% (0.068-1.5%)	1.1% (0.12-2.6%)	0.11 (0.011-0.28)	6.8 (0.76-16)
Guam	2000	Female	45 to 49	34% (24-49%)	66% (51-76%)	0.35 (0.11-0.7)	0.17% (-0.39-0.86%)	0.41% (-0.45-1.6%)	0.056 (-0.064-0.22)	1.9 (-4.5-9.9)
Guam	2000	Female	50 to 54	38% (28-51%)	62% (49-72%)	0.33 (0.1-0.67)	-0.27% (-0.99-0.44%)	-0.12% (-1.1-0.96%)	-0.021 (-0.2-0.17)	-3.3 (-12-5.3)
Guam	2000	Female	55 to 59	37% (27-50%)	63% (50-73%)	0.29 (0.093-0.6)	-0.48% (-1.3-0.22%)	-0.35% (-1.4-0.62%)	-0.065 (-0.26-0.11)	-4.7 (-12-2.2)
Guam	2000	Female	60 to 64	35% (24-50%)	65% (50-76%)	0.25 (0.078-0.52)	-0.72% (-1.6-0.015%)	-0.72% (-1.9-0.19%)	-0.2 (-0.52-0.052)	-8.3 (-19-0.18)
Guam	2000	Female	65 to 69	36% (24-50%)	64% (50-76%)	0.25 (0.075-0.53)	-0.72% (-1.6--0.0037%)	-0.7% (-1.8-0.21%)	-0.23 (-0.61-0.066)	-7.8 (-18--0.038)
Guam	2000	Female	70 to 74	36% (25-51%)	64% (49-75%)	0.23 (0.069-0.5)	-0.93% (-1.9--0.15%)	-1% (-2.2--0.1%)	-0.32 (-0.7-0.031)	-7.3 (-15--1.2)
Guam	2000	Female	75 to 79	36% (25-52%)	64% (48-75%)	0.23 (0.071-0.49)	-0.75% (-1.6--0.051%)	-0.8% (-1.8-0.023%)	-0.27 (-0.63-0.0082)	-4.8 (-11--0.34)
Guam	2000	Female	80 to 84	36% (24-51%)	64% (49-76%)	0.24 (0.069-0.5)	-0.91% (-1.8--0.17%)	-1% (-2.1--0.15%)	-0.38 (-0.78--0.054)	-4.5 (-9--0.82)
Guam	2000	Female	85 to 89	36% (24-50%)	64% (50-76%)	0.25 (0.07-0.54)	-0.93% (-1.9--0.16%)	-1.1% (-2.3--0.15%)	-0.29 (-0.61--0.038)	-2.3 (-4.9--0.38)
Guam	2000	Female	90 to 94	38% (26-52%)	62% (48-74%)	0.28 (0.078-0.62)	-0.88% (-1.7--0.14%)	-1% (-2--0.14%)	-0.14 (-0.28--0.02)	-0.78 (-1.5--0.12)
Guam	2000	Female	95 plus	41% (26-58%)	59% (42-74%)	0.31 (0.082-0.72)	-0.88% (-2-0.032%)	-1% (-2.4-0.056%)	-0.049 (-0.11-0.0027)	-0.17 (-0.38-0.006)
Guam	2000	Male	All Ages	58% (51-66%)	42% (34-49%)	0.83 (0.27-1.6)	2.4% (1-4.9%)	1.8% (0.36-4.2%)	8.3 (1.7-20)	520 (220-1000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	2000	Male	15 to 19	53% (37-69%)	47% (31-63%)	0.71 (0.21-1.5)	3% (0.93-6%)	4.3% (0.82-9.5%)	0.33 (0.063-0.73)	28 (8.6-58)
Guam	2000	Male	20 to 24	69% (53-81%)	31% (19-47%)	1.6 (0.5-3.2)	7% (2.6-13%)	9.4% (2.8-18%)	1 (0.28-2)	78 (30-150)
Guam	2000	Male	25 to 29	73% (61-84%)	27% (16-39%)	1.6 (0.52-3.2)	6.5% (2.8-12%)	8.5% (2.7-17%)	1.1 (0.32-2.1)	80 (34-140)
Guam	2000	Male	30 to 34	74% (61-85%)	26% (15-39%)	1.5 (0.51-3)	5.4% (2.4-9.9%)	6.7% (2.2-13%)	1 (0.32-2.1)	75 (32-140)
Guam	2000	Male	35 to 39	74% (61-85%)	26% (15-39%)	1.4 (0.43-2.9)	4.8% (2.3-8.6%)	6.1% (2.3-12%)	1.1 (0.42-2.2)	75 (36-140)
Guam	2000	Male	40 to 44	72% (59-83%)	28% (17-41%)	1.3 (0.4-2.6)	3.8% (1.8-7.3%)	4.6% (1.7-9.3%)	0.97 (0.35-2)	57 (26-110)
Guam	2000	Male	45 to 49	71% (55-82%)	29% (18-45%)	1.2 (0.36-2.4)	3% (1.1-6.1%)	3.5% (1.1-7.7%)	0.91 (0.26-2)	48 (19-98)
Guam	2000	Male	50 to 54	71% (56-83%)	29% (17-44%)	1.1 (0.34-2.1)	2.3% (0.55-5.4%)	2.6% (0.23-6.3%)	0.85 (0.075-2.2)	39 (9.5-91)
Guam	2000	Male	55 to 59	68% (53-81%)	32% (19-47%)	0.81 (0.25-1.6)	1% (-0.35-3.1%)	0.99% (-0.71-3.5%)	0.34 (-0.24-1.2)	15 (-5-45)
Guam	2000	Male	60 to 64	62% (42-78%)	38% (22-58%)	0.61 (0.19-1.3)	0.61% (-0.53-2.2%)	0.58% (-0.86-2.5%)	0.25 (-0.38-1.1)	9.2 (-8.3-34)
Guam	2000	Male	65 to 69	59% (40-76%)	41% (24-60%)	0.56 (0.17-1.1)	0.54% (-0.43-2%)	0.55% (-0.64-2.3%)	0.27 (-0.31-1.1)	7.4 (-5.8-28)
Guam	2000	Male	70 to 74	57% (39-73%)	43% (27-61%)	0.52 (0.15-1.1)	0.24% (-0.72-1.6%)	0.21% (-0.97-1.7%)	0.1 (-0.5-0.9)	2.7 (-8.3-18)
Guam	2000	Male	75 to 79	53% (35-71%)	47% (29-65%)	0.46 (0.14-0.97)	0.11% (-0.8-1.3%)	0.046% (-1-1.4%)	0.022 (-0.5-0.65)	0.93 (-6.3-11)
Guam	2000	Male	80 to 84	50% (31-69%)	50% (31-69%)	0.42 (0.13-0.89)	0.15% (-0.62-1.2%)	0.1% (-0.75-1.3%)	0.041 (-0.31-0.5)	0.76 (-3-5.8)
Guam	2000	Male	85 to 89	48% (31-66%)	52% (34-69%)	0.41 (0.11-0.9)	0.015% (-0.72-0.96%)	-0.06% (-0.94-1%)	-0.012 (-0.19-0.21)	0.027 (-1.3-1.7)
Guam	2000	Male	90 to 94	50% (34-68%)	50% (32-66%)	0.42 (0.11-0.97)	0.009% (-0.67-0.84%)	-0.066% (-0.88-0.86%)	-0.0055 (-0.072-0.071)	0.0047 (-0.34-0.42)
Guam	2000	Male	95 plus	53% (32-72%)	47% (28-68%)	0.46 (0.12-1.1)	-0.24% (-1.1-0.82%)	-0.43% (-1.4-0.77%)	-0.0079 (-0.027-0.014)	-0.018 (-0.08-0.061)
Guam	2005	Female	All Ages	30% (26-35%)	70% (65-74%)	0.33 (0.12-0.64)	0.036% (-0.36-0.54%)	-0.52% (-1.2-0.21%)	-2 (-4.7-0.77)	6.5 (-60-93)
Guam	2005	Female	15 to 19	35% (25-49%)	65% (51-75%)	0.53 (0.18-1.1)	1.1% (0.42-2.1%)	2.2% (0.32-5%)	0.07 (0.0095-0.16)	8.1 (3.1-16)
Guam	2005	Female	20 to 24	40% (31-53%)	60% (47-69%)	0.74 (0.25-1.5)	1.8% (0.88-3.2%)	2.9% (0.68-6.4%)	0.095 (0.023-0.21)	12 (6-23)
Guam	2005	Female	25 to 29	40% (31-53%)	60% (47-69%)	0.61 (0.22-1.2)	1.5% (0.84-2.6%)	2.3% (0.64-4.7%)	0.084 (0.025-0.18)	11 (5.8-19)
Guam	2005	Female	30 to 34	39% (30-52%)	61% (48-70%)	0.54 (0.2-1.1)	1.1% (0.5-2%)	1.4% (0.31-3.4%)	0.074 (0.017-0.18)	9.4 (4.2-17)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	2005	Female	35 to 39	35% (27-49%)	65% (51-73%)	0.53 (0.19-1.1)	0.95% (0.41-1.9%)	1.5% (0.43-3.5%)	0.11 (0.031-0.27)	9.7 (4.1-20)
Guam	2005	Female	40 to 44	29% (21-41%)	71% (59-79%)	0.41 (0.13-0.83)	0.62% (0.1-1.5%)	0.99% (0.12-2.6%)	0.11 (0.013-0.29)	7.6 (1.2-18)
Guam	2005	Female	45 to 49	29% (20-42%)	71% (58-80%)	0.33 (0.11-0.67)	0.22% (-0.28-0.87%)	0.45% (-0.34-1.6%)	0.067 (-0.051-0.23)	2.7 (-3.6-11)
Guam	2005	Female	50 to 54	34% (25-47%)	66% (53-75%)	0.33 (0.11-0.68)	-0.27% (-0.95-0.38%)	-0.13% (-1-0.93%)	-0.026 (-0.21-0.18)	-3.6 (-13-5.1)
Guam	2005	Female	55 to 59	34% (25-48%)	66% (52-75%)	0.3 (0.1-0.6)	-0.53% (-1.2-0.13%)	-0.43% (-1.4-0.48%)	-0.11 (-0.36-0.13)	-7.3 (-17-1.8)
Guam	2005	Female	60 to 64	31% (21-46%)	69% (54-79%)	0.26 (0.084-0.52)	-0.7% (-1.6--0.034%)	-0.7% (-1.8-0.14%)	-0.19 (-0.48-0.036)	-7.7 (-17--0.38)
Guam	2005	Female	65 to 69	32% (22-47%)	68% (53-78%)	0.25 (0.081-0.52)	-0.72% (-1.6--0.053%)	-0.73% (-1.8-0.13%)	-0.28 (-0.7-0.05)	-9.2 (-20--0.68)
Guam	2005	Female	70 to 74	33% (22-49%)	67% (51-78%)	0.24 (0.081-0.52)	-0.91% (-1.8--0.2%)	-1% (-2.1--0.16%)	-0.46 (-0.97--0.071)	-11 (-21--2.4)
Guam	2005	Female	75 to 79	33% (22-48%)	67% (52-78%)	0.24 (0.08-0.5)	-0.77% (-1.6--0.086%)	-0.83% (-1.9--0.018%)	-0.34 (-0.78--0.0077)	-6 (-13--0.69)
Guam	2005	Female	80 to 84	32% (21-47%)	68% (53-79%)	0.24 (0.081-0.5)	-0.9% (-1.8--0.22%)	-1% (-2.1--0.22%)	-0.47 (-0.96--0.1)	-5.5 (-11--1.4)
Guam	2005	Female	85 to 89	33% (21-45%)	67% (55-79%)	0.26 (0.081-0.54)	-0.96% (-1.9--0.22%)	-1.1% (-2.2--0.23%)	-0.46 (-0.93--0.093)	-3.6 (-7.1--0.82)
Guam	2005	Female	90 to 94	34% (23-48%)	66% (52-77%)	0.28 (0.087-0.64)	-0.9% (-1.7--0.16%)	-1% (-2--0.18%)	-0.21 (-0.4--0.036)	-1.1 (-2.1--0.19)
Guam	2005	Female	95 plus	37% (23-55%)	63% (45-77%)	0.32 (0.091-0.74)	-0.9% (-2--0.00031%)	-1.1% (-2.4--0.055%)	-0.073 (-0.16--0.0037)	-0.24 (-0.53--0.000083)
Guam	2005	Male	All Ages	58% (51-65%)	42% (35-49%)	0.94 (0.33-1.8)	2.4% (0.94-5.1%)	1.7% (0.17-4.5%)	9.5 (0.95-26)	580 (220-1200)
Guam	2005	Male	15 to 19	55% (40-69%)	45% (31-60%)	0.85 (0.3-1.7)	3.4% (1.1-6.6%)	4.9% (1.3-10%)	0.43 (0.11-0.88)	35 (12-70)
Guam	2005	Male	20 to 24	70% (56-82%)	30% (18-44%)	1.9 (0.66-3.8)	7.4% (2.9-13%)	9.7% (3.1-19%)	1.1 (0.35-2.2)	87 (35-160)
Guam	2005	Male	25 to 29	74% (61-84%)	26% (16-39%)	1.8 (0.67-3.7)	7.2% (3.1-13%)	9.4% (3.2-18%)	1.1 (0.35-2)	79 (35-140)
Guam	2005	Male	30 to 34	74% (61-85%)	26% (15-39%)	1.8 (0.63-3.5)	5.7% (2.6-11%)	7.2% (2.7-14%)	1 (0.34-2)	72 (32-140)
Guam	2005	Male	35 to 39	72% (60-84%)	28% (16-40%)	1.6 (0.56-3.2)	5.1% (2.4-9.5%)	6.5% (2.5-13%)	1.2 (0.44-2.4)	76 (36-140)
Guam	2005	Male	40 to 44	71% (59-82%)	29% (18-41%)	1.5 (0.5-2.9)	4% (1.7-7.9%)	4.8% (1.7-10%)	1.2 (0.42-2.6)	70 (30-140)
Guam	2005	Male	45 to 49	69% (53-81%)	31% (19-47%)	1.3 (0.46-2.6)	3.4% (1.4-6.9%)	3.9% (1.3-8.8%)	1.2 (0.38-2.6)	60 (24-130)
Guam	2005	Male	50 to 54	69% (53-81%)	31% (19-47%)	1.2 (0.4-2.3)	2.5% (0.67-6.1%)	2.8% (0.36-7.1%)	1.1 (0.14-2.8)	49 (12-110)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	2005	Male	55 to 59	68% (53-81%)	32% (19-47%)	0.94 (0.34-1.9)	1.2% (-0.3-3.8%)	1.2% (-0.72-4.3%)	0.56 (-0.35-2.1)	24 (-6.2-79)
Guam	2005	Male	60 to 64	66% (48-81%)	34% (19-52%)	0.78 (0.27-1.6)	0.55% (-0.77-2.6%)	0.49% (-1.1-3%)	0.23 (-0.56-1.4)	8.9 (-13-44)
Guam	2005	Male	65 to 69	61% (42-78%)	39% (22-58%)	0.68 (0.22-1.4)	0.54% (-0.6-2.5%)	0.54% (-0.81-2.8%)	0.32 (-0.49-1.7)	9 (-10-42)
Guam	2005	Male	70 to 74	56% (37-72%)	44% (28-63%)	0.59 (0.2-1.2)	0.27% (-0.75-2%)	0.23% (-1-2.2%)	0.14 (-0.64-1.4)	3.7 (-11-27)
Guam	2005	Male	75 to 79	52% (34-70%)	48% (30-66%)	0.52 (0.17-1.1)	0.14% (-0.82-1.5%)	0.065% (-1.1-1.7%)	0.037 (-0.64-0.95)	1.4 (-8.2-15)
Guam	2005	Male	80 to 84	49% (31-67%)	51% (33-69%)	0.47 (0.16-0.97)	0.14% (-0.68-1.3%)	0.075% (-0.88-1.4%)	0.038 (-0.47-0.73)	0.89 (-4.4-8.6)
Guam	2005	Male	85 to 89	47% (30-64%)	53% (36-70%)	0.45 (0.15-0.99)	-0.0014% (-0.8-1.1%)	-0.09% (-1-1.2%)	-0.03 (-0.34-0.4)	-0.0018 (-2.3-3.3)
Guam	2005	Male	90 to 94	49% (33-66%)	51% (34-67%)	0.47 (0.14-1.1)	-0.024% (-0.74-0.92%)	-0.12% (-0.96-0.94%)	-0.013 (-0.1-0.1)	-0.015 (-0.47-0.6)
Guam	2005	Male	95 plus	52% (31-72%)	48% (28-69%)	0.51 (0.15-1.2)	-0.21% (-1.1-0.95%)	-0.4% (-1.5-0.86%)	-0.011 (-0.041-0.025)	-0.023 (-0.12-0.11)
Guam	2010	Female	All Ages	31% (27-36%)	69% (64-73%)	0.35 (0.12-0.7)	-0.012% (-0.43-0.56%)	-0.62% (-1.4-0.18%)	-3.1 (-6.8-0.87)	-2.3 (-8.7-110)
Guam	2010	Female	15 to 19	36% (26-50%)	64% (50-74%)	0.56 (0.2-1.2)	1.1% (0.37-2.3%)	1.8% (0.26-4.4%)	0.083 (0.012-0.2)	9.1 (3.1-18)
Guam	2010	Female	20 to 24	40% (30-53%)	60% (47-70%)	0.75 (0.26-1.5)	1.8% (0.89-3.3%)	3% (0.64-6.9%)	0.12 (0.026-0.27)	15 (6.7-27)
Guam	2010	Female	25 to 29	40% (29-54%)	60% (46-71%)	0.61 (0.21-1.3)	1.6% (0.84-2.7%)	2.3% (0.73-4.8%)	0.099 (0.03-0.21)	12 (6.1-20)
Guam	2010	Female	30 to 34	39% (29-53%)	61% (47-71%)	0.56 (0.19-1.1)	1.1% (0.48-2.1%)	1.4% (0.31-3.5%)	0.078 (0.017-0.2)	9 (3.9-17)
Guam	2010	Female	35 to 39	38% (29-52%)	62% (48-71%)	0.59 (0.2-1.2)	1% (0.39-2.3%)	1.7% (0.46-3.9%)	0.14 (0.04-0.34)	11 (3.9-24)
Guam	2010	Female	40 to 44	36% (27-51%)	64% (49-73%)	0.53 (0.18-1.1)	0.65% (-0.039-1.8%)	1.1% (0.024-3.1%)	0.14 (0.0034-0.4)	8.2 (-0.47-23)
Guam	2010	Female	45 to 49	33% (24-47%)	67% (53-76%)	0.39 (0.14-0.81)	0.22% (-0.39-1.1%)	0.49% (-0.38-1.9%)	0.094 (-0.073-0.36)	3.2 (-5.8-16)
Guam	2010	Female	50 to 54	34% (24-47%)	66% (53-76%)	0.34 (0.12-0.68)	-0.24% (-0.91-0.49%)	-0.11% (-1-1%)	-0.026 (-0.25-0.24)	-3.6 (-14-7.6)
Guam	2010	Female	55 to 59	31% (22-43%)	69% (57-78%)	0.28 (0.093-0.59)	-0.39% (-1.1-0.24%)	-0.29% (-1.2-0.57%)	-0.093 (-0.39-0.18)	-6.3 (-18-3.8)
Guam	2010	Female	60 to 64	27% (18-41%)	73% (59-82%)	0.22 (0.074-0.46)	-0.51% (-1.3-0.056%)	-0.5% (-1.5-0.25%)	-0.2 (-0.6-0.1)	-8.4 (-21-0.93)
Guam	2010	Female	65 to 69	31% (21-44%)	69% (56-79%)	0.24 (0.08-0.51)	-0.66% (-1.5-0.0012%)	-0.66% (-1.7-0.17%)	-0.26 (-0.67-0.067)	-8.4 (-19-0.015)
Guam	2010	Female	70 to 74	35% (23-50%)	65% (50-77%)	0.26 (0.087-0.55)	-1% (-2--0.23%)	-1.1% (-2.3--0.18%)	-0.66 (-1.4-0.1)	-15 (-29--3.3)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	2010	Female	75 to 79	35% (24-51%)	65% (49-76%)	0.27 (0.089-0.56)	-0.94% (-1.9--0.18%)	-1% (-2.2--0.11%)	-0.68 (-1.5--0.075)	-12 (-23--2.1)
Guam	2010	Female	80 to 84	35% (23-50%)	65% (50-77%)	0.27 (0.093-0.58)	-1% (-2--0.28%)	-1.2% (-2.4--0.3%)	-0.8 (-1.6--0.2)	-9.3 (-18--2.5)
Guam	2010	Female	85 to 89	36% (24-49%)	64% (51-76%)	0.29 (0.097-0.62)	-1.1% (-2.2--0.28%)	-1.3% (-2.5--0.32%)	-0.64 (-1.3--0.16)	-5.1 (-9.8--1.3)
Guam	2010	Female	90 to 94	37% (25-51%)	63% (49-75%)	0.32 (0.098-0.71)	-1.1% (-2.1--0.28%)	-1.2% (-2.4--0.3%)	-0.36 (-0.69--0.086)	-1.9 (-3.6--0.49)
Guam	2010	Female	95 plus	40% (26-57%)	60% (43-74%)	0.36 (0.11-0.83)	-1.1% (-2.3--0.086%)	-1.3% (-2.8--0.12%)	-0.12 (-0.25--0.011)	-0.39 (-0.84--0.031)
Guam	2010	Male	All Ages	57% (50-65%)	43% (35-50%)	0.97 (0.34-1.9)	2.5% (0.9-5.1%)	1.7% (0.17-4.4%)	12 (1.2-31)	680 (260-1500)
Guam	2010	Male	15 to 19	52% (37-67%)	48% (33-63%)	0.84 (0.28-1.7)	3.4% (1.2-6.7%)	4.9% (1.1-11%)	0.48 (0.11-1.1)	40 (14-81)
Guam	2010	Male	20 to 24	69% (54-81%)	31% (19-46%)	1.9 (0.69-3.9)	7.7% (2.9-14%)	10% (3.3-20%)	1.4 (0.42-2.7)	100 (38-190)
Guam	2010	Male	25 to 29	74% (61-84%)	26% (16-39%)	2 (0.67-3.9)	7.6% (3.3-14%)	9.9% (3.5-19%)	1.2 (0.41-2.3)	87 (38-160)
Guam	2010	Male	30 to 34	73% (59-84%)	27% (16-41%)	1.8 (0.62-3.5)	6% (2.6-11%)	7.4% (2.5-15%)	1 (0.34-2.2)	73 (32-140)
Guam	2010	Male	35 to 39	72% (58-84%)	28% (16-42%)	1.7 (0.55-3.2)	5.4% (2.4-10%)	6.7% (2.6-13%)	1.3 (0.49-2.6)	80 (37-150)
Guam	2010	Male	40 to 44	73% (59-84%)	27% (16-41%)	1.6 (0.54-3.1)	4.3% (1.8-9%)	5.2% (1.8-11%)	1.4 (0.48-3.1)	81 (34-170)
Guam	2010	Male	45 to 49	69% (53-81%)	31% (19-47%)	1.4 (0.51-2.8)	3.7% (1.4-9.7%)	4.2% (1.4-9.1%)	1.7 (0.53-3.9)	84 (33-180)
Guam	2010	Male	50 to 54	67% (51-80%)	33% (20-49%)	1.2 (0.41-2.4)	2.8% (0.86-6.5%)	3.1% (0.63-7.6%)	1.5 (0.31-3.7)	65 (19-150)
Guam	2010	Male	55 to 59	66% (50-79%)	34% (21-50%)	0.95 (0.34-1.9)	1.4% (-0.21-4.1%)	1.4% (-0.59-4.6%)	0.83 (-0.36-2.9)	33 (-4.8-100)
Guam	2010	Male	60 to 64	64% (47-79%)	36% (21-53%)	0.79 (0.27-1.6)	0.6% (-0.75-3%)	0.54% (-1.1-3.3%)	0.39 (-0.83-2.4)	14 (-19-72)
Guam	2010	Male	65 to 69	59% (42-75%)	41% (25-58%)	0.69 (0.24-1.4)	0.57% (-0.61-2.6%)	0.57% (-0.85-2.9%)	0.38 (-0.56-2)	11 (-12-49)
Guam	2010	Male	70 to 74	54% (36-70%)	46% (30-64%)	0.57 (0.19-1.2)	0.32% (-0.7-2.1%)	0.29% (-0.89-2.3%)	0.23 (-0.7-1.8)	5.6 (-12-35)
Guam	2010	Male	75 to 79	50% (33-67%)	50% (33-67%)	0.51 (0.16-1.1)	0.16% (-0.8-1.6%)	0.084% (-1-1.7%)	0.064 (-0.81-1.3)	2 (-10-21)
Guam	2010	Male	80 to 84	47% (29-65%)	53% (35-71%)	0.46 (0.15-0.99)	0.17% (-0.61-1.3%)	0.11% (-0.81-1.4%)	0.07 (-0.53-0.9)	1.3 (-5-11)
Guam	2010	Male	85 to 89	45% (28-62%)	55% (38-72%)	0.44 (0.14-0.97)	0.026% (-0.74-1%)	-0.059% (-0.93-1.1%)	-0.025 (-0.4-0.47)	0.096 (-2.7-3.9)
Guam	2010	Male	90 to 94	46% (31-64%)	54% (36-69%)	0.45 (0.14-1.1)	-0.016% (-0.73-0.93%)	-0.11% (-0.98-0.9%)	-0.02 (-0.17-0.16)	-0.017 (-0.77-1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	2010	Male	95 plus	49% (29-69%)	51% (31-71%)	0.49 (0.14-1.1)	-0.19% (-1.1-0.94%)	-0.38% (-1.4-0.83%)	-0.013 (-0.052-0.03)	-0.027 (-0.15-0.13)
Guam	2016	Female	All Ages	33% (29-38%)	67% (62-71%)	0.37 (0.13-0.76)	-0.1% (-0.6-0.52%)	-0.76% (-1.6-0.096%)	-4.6 (-10-0.63)	-23 (-130-120)
Guam	2016	Female	15 to 19	37% (27-51%)	63% (49-73%)	0.59 (0.2-1.3)	1.2% (0.4-2.4%)	2.4% (0.31-6.1%)	0.086 (0.01-0.22)	9.6 (3.3-20)
Guam	2016	Female	20 to 24	41% (31-54%)	59% (46-69%)	0.79 (0.26-1.7)	1.9% (0.9-3.6%)	3.1% (0.69-7.2%)	0.13 (0.026-0.31)	16 (7.6-32)
Guam	2016	Female	25 to 29	40% (29-54%)	60% (46-71%)	0.62 (0.2-1.2)	1.6% (0.83-2.8%)	2.3% (0.71-4.9%)	0.11 (0.033-0.25)	14 (7.2-25)
Guam	2016	Female	30 to 34	40% (30-54%)	60% (46-70%)	0.57 (0.2-1.2)	1.1% (0.48-2.1%)	1.4% (0.3-3.6%)	0.084 (0.016-0.22)	9.8 (4.1-20)
Guam	2016	Female	35 to 39	42% (32-57%)	58% (43-68%)	0.65 (0.22-1.3)	1% (0.34-2.4%)	1.7% (0.43-4.2%)	0.14 (0.03-0.36)	10 (3.2-24)
Guam	2016	Female	40 to 44	41% (32-56%)	59% (44-68%)	0.62 (0.21-1.3)	0.61% (-0.17-2%)	1.1% (-0.22-3.6%)	0.14 (-0.026-0.44)	7.8 (-2-26)
Guam	2016	Female	45 to 49	37% (27-51%)	63% (49-73%)	0.45 (0.16-0.92)	0.12% (-0.61-1.1%)	0.4% (-0.69-2%)	0.076 (-0.13-0.38)	1.9 (-9.3-17)
Guam	2016	Female	50 to 54	35% (25-49%)	65% (51-75%)	0.36 (0.12-0.74)	-0.28% (-1.1-0.5%)	-0.16% (-1.2-1%)	-0.046 (-0.37-0.29)	-5.3 (-20-9.3)
Guam	2016	Female	55 to 59	30% (21-44%)	70% (56-79%)	0.27 (0.088-0.59)	-0.37% (-1.1-0.25%)	-0.28% (-1.3-0.58%)	-0.1 (-0.46-0.22)	-7 (-20-4.8)
Guam	2016	Female	60 to 64	26% (18-40%)	74% (60-82%)	0.22 (0.073-0.46)	-0.48% (-1.2-0.094%)	-0.47% (-1.4-0.24%)	-0.22 (-0.68-0.11)	-9.4 (-24-1.9)
Guam	2016	Female	65 to 69	31% (21-46%)	69% (54-79%)	0.25 (0.082-0.52)	-0.67% (-1.6-0.016%)	-0.68% (-1.8-0.17%)	-0.41 (-1.1-0.11)	-13 (-31-0.32)
Guam	2016	Female	70 to 74	37% (25-52%)	63% (48-75%)	0.28 (0.091-0.6)	-1.1% (-2.2--0.26%)	-1.2% (-2.5--0.24%)	-0.75 (-1.6--0.15)	-17 (-34--4.3)
Guam	2016	Female	75 to 79	38% (25-54%)	62% (46-75%)	0.3 (0.097-0.65)	-1% (-2.1--0.21%)	-1.2% (-2.5--0.15%)	-0.92 (-2--0.13)	-15 (-31--3.3)
Guam	2016	Female	80 to 84	38% (25-54%)	62% (46-75%)	0.3 (0.097-0.67)	-1.2% (-2.3--0.35%)	-1.3% (-2.7--0.31%)	-1.3 (-2.6--0.31)	-15 (-29--4.6)
Guam	2016	Female	85 to 89	38% (26-52%)	62% (48-74%)	0.32 (0.098-0.69)	-1.2% (-2.4--0.34%)	-1.4% (-2.9--0.36%)	-1 (-2.1--0.27)	-8.2 (-16--2.3)
Guam	2016	Female	90 to 94	40% (27-54%)	60% (46-73%)	0.35 (0.1-0.8)	-1.2% (-2.2--0.26%)	-1.4% (-2.6--0.29%)	-0.46 (-0.88--0.099)	-2.4 (-4.6--0.56)
Guam	2016	Female	95 plus	43% (27-60%)	57% (40-73%)	0.4 (0.12-0.92)	-1.2% (-2.5--0.087%)	-1.5% (-3-0.14%)	-0.19 (-0.4-0.019)	-0.63 (-1.3--0.043)
Guam	2016	Male	All Ages	56% (49-64%)	44% (36-51%)	0.98 (0.34-2)	2.4% (0.84-5.3%)	1.6% (0.16-4.5%)	14 (1.4-38)	760 (260-1700)
Guam	2016	Male	15 to 19	47% (32-63%)	53% (37-68%)	0.75 (0.24-1.6)	3.1% (1.1-6.4%)	4.5% (1.1-9.9%)	0.47 (0.12-1)	39 (14-80)
Guam	2016	Male	20 to 24	66% (51-79%)	34% (21-49%)	1.8 (0.63-3.7)	7.8% (3-15%)	10% (3.4-21%)	1.5 (0.5-3.1)	120 (45-220)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Guam	2016	Male	25 to 29	73% (60-84%)	27% (16-40%)	2 (0.65-3.9)	7.5% (3-14%)	9.7% (3.1-19%)	1.4 (0.46-2.7)	100 (42-190)
Guam	2016	Male	30 to 34	72% (58-83%)	28% (17-42%)	1.8 (0.59-3.8)	6% (2.6-11%)	7.4% (2.5-15%)	1.1 (0.35-2.3)	77 (32-150)
Guam	2016	Male	35 to 39	72% (59-84%)	28% (16-41%)	1.7 (0.57-3.5)	5.5% (2.5-10%)	6.8% (2.7-14%)	1.2 (0.44-2.5)	77 (35-150)
Guam	2016	Male	40 to 44	73% (59-84%)	27% (16-41%)	1.6 (0.54-3.3)	4.5% (1.9-9%)	5.3% (1.8-11%)	1.4 (0.45-3.1)	80 (32-160)
Guam	2016	Male	45 to 49	69% (52-82%)	31% (18-48%)	1.4 (0.48-3)	3.8% (1.4-8.2%)	4.4% (1.3-9.9%)	1.7 (0.49-4)	87 (33-190)
Guam	2016	Male	50 to 54	67% (51-80%)	33% (20-49%)	1.2 (0.41-2.5)	2.9% (0.8-7%)	3.2% (0.6-8.2%)	1.9 (0.35-4.8)	84 (23-200)
Guam	2016	Male	55 to 59	64% (48-78%)	36% (22-52%)	0.92 (0.3-1.9)	1.5% (-0.17-4.3%)	1.5% (-0.48-4.9%)	1.1 (-0.36-3.5)	43 (-5.5-130)
Guam	2016	Male	60 to 64	62% (44-78%)	38% (22-56%)	0.75 (0.25-1.6)	0.72% (-0.58-3.1%)	0.68% (-0.89-3.4%)	0.59 (-0.78-2.9)	21 (-17-92)
Guam	2016	Male	65 to 69	59% (42-75%)	41% (25-58%)	0.68 (0.23-1.4)	0.66% (-0.5-2.7%)	0.67% (-0.7-3%)	0.68 (-0.75-3.1)	18 (-15-75)
Guam	2016	Male	70 to 74	53% (35-69%)	47% (31-65%)	0.56 (0.18-1.2)	0.39% (-0.61-2.1%)	0.37% (-0.82-2.4%)	0.34 (-0.75-2.2)	8 (-12-42)
Guam	2016	Male	75 to 79	48% (30-66%)	52% (34-70%)	0.49 (0.15-1.1)	0.25% (-0.73-1.7%)	0.19% (-0.95-1.9%)	0.18 (-0.87-1.8)	3.9 (-11-28)
Guam	2016	Male	80 to 84	45% (27-63%)	55% (37-73%)	0.44 (0.14-0.97)	0.24% (-0.53-1.5%)	0.19% (-0.67-1.7%)	0.17 (-0.59-1.4)	2.5 (-5.6-16)
Guam	2016	Male	85 to 89	43% (26-60%)	57% (40-74%)	0.43 (0.13-0.94)	0.092% (-0.65-1.4%)	0.019% (-0.85-1.4%)	0.0093 (-0.44-0.74)	0.41 (-2.9-6.2)
Guam	2016	Male	90 to 94	45% (29-62%)	55% (38-71%)	0.45 (0.13-1)	0.036% (-0.68-0.99%)	-0.051% (-0.92-0.98%)	-0.011 (-0.2-0.21)	0.048 (-0.92-1.3)
Guam	2016	Male	95 plus	48% (28-68%)	52% (32-72%)	0.49 (0.13-1.1)	-0.13% (-1-1%)	-0.31% (-1.4-0.96%)	-0.019 (-0.085-0.06)	-0.032 (-0.25-0.25)
Northern Mariana Islands	1990	Female	All Ages	21% (13-30%)	79% (70-87%)	0.25 (0.033-0.58)	0.83% (0.45-1.7%)	0.74% (0.15-2.1%)	0.4 (0.078-1.1)	33 (17-67)
Northern Mariana Islands	1990	Female	15 to 19	20% (8.6-36%)	80% (64-91%)	0.24 (0.026-0.67)	0.79% (0.34-1.6%)	1.1% (0.14-3.1%)	0.0088 (0.0011-0.028)	1.4 (0.56-3.1)
Northern Mariana Islands	1990	Female	20 to 24	24% (10-44%)	76% (56-90%)	0.37 (0.043-0.96)	1.4% (0.77-2.6%)	1.6% (0.28-4%)	0.042 (0.007-0.12)	6.9 (3.5-13)
Northern Mariana Islands	1990	Female	25 to 29	26% (11-45%)	74% (55-89%)	0.38 (0.046-1)	1.5% (0.87-2.6%)	1.8% (0.53-4.1%)	0.05 (0.015-0.12)	6.5 (3.7-12)
Northern Mariana Islands	1990	Female	30 to 34	26% (11-48%)	74% (52-89%)	0.39 (0.049-0.99)	1.3% (0.72-2.5%)	1.7% (0.57-3.8%)	0.057 (0.018-0.14)	5.9 (3.2-11)
Northern Mariana Islands	1990	Female	35 to 39	27% (12-46%)	73% (54-88%)	0.41 (0.05-1.1)	1.3% (0.64-2.7%)	1.8% (0.63-4.2%)	0.064 (0.021-0.16)	4.8 (2.3-10)
Northern Mariana Islands	1990	Female	40 to 44	26% (12-44%)	74% (56-88%)	0.38 (0.043-1)	1.2% (0.5-2.6%)	1.7% (0.64-4.2%)	0.058 (0.019-0.15)	3.3 (1.3-7.9)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	1990	Female	45 to 49	26% (11-45%)	74% (55-89%)	0.33 (0.031-1)	0.87% (0.25-2.4%)	1.3% (0.44-3.6%)	0.035 (0.011-0.1)	1.6 (0.44-4.6)
Northern Mariana Islands	1990	Female	50 to 54	26% (12-46%)	74% (54-88%)	0.3 (0.025-0.91)	0.62% (-0.059-2%)	1% (0.095-3.1%)	0.026 (0.0023-0.085)	0.95 (-0.082-3.2)
Northern Mariana Islands	1990	Female	55 to 59	25% (11-45%)	75% (55-89%)	0.26 (0.023-0.83)	0.47% (-0.27-1.9%)	0.8% (-0.12-2.7%)	0.025 (-0.0035-0.083)	0.69 (-0.37-2.8)
Northern Mariana Islands	1990	Female	60 to 64	25% (12-45%)	75% (55-88%)	0.25 (0.026-0.78)	0.19% (-0.58-1.3%)	0.41% (-0.45-1.9%)	0.012 (-0.013-0.055)	0.22 (-0.65-1.5)
Northern Mariana Islands	1990	Female	65 to 69	26% (12-44%)	74% (56-88%)	0.25 (0.028-0.75)	0.17% (-0.59-1.4%)	0.37% (-0.53-2%)	0.013 (-0.019-0.074)	0.18 (-0.65-1.6)
Northern Mariana Islands	1990	Female	70 to 74	26% (12-43%)	74% (57-88%)	0.24 (0.027-0.71)	0.027% (-0.61-1.2%)	0.14% (-0.62-1.6%)	0.0054 (-0.024-0.06)	0.026 (-0.59-1.1)
Northern Mariana Islands	1990	Female	75 to 79	25% (11-43%)	75% (57-89%)	0.22 (0.022-0.65)	0.11% (-0.53-1.1%)	0.23% (-0.51-1.5%)	0.011 (-0.026-0.071)	0.1 (-0.47-1)
Northern Mariana Islands	1990	Female	80 to 84	24% (11-43%)	76% (57-89%)	0.21 (0.021-0.59)	-0.055% (-0.69-0.94%)	-0.019% (-0.75-1.1%)	-0.00081 (-0.034-0.054)	-0.032 (-0.41-0.56)
Northern Mariana Islands	1990	Female	85 to 89	25% (11-43%)	75% (57-89%)	0.22 (0.023-0.65)	-0.086% (-0.8-0.83%)	-0.068% (-0.89-0.99%)	-0.002 (-0.026-0.031)	-0.023 (-0.21-0.23)
Northern Mariana Islands	1990	Female	90 to 94	28% (13-49%)	72% (51-87%)	0.25 (0.024-0.73)	-0.12% (-0.75-0.75%)	-0.11% (-0.83-0.84%)	-0.0014 (-0.01-0.011)	-0.0086 (-0.057-0.059)
Northern Mariana Islands	1990	Female	95 plus	33% (14-56%)	67% (44-86%)	0.29 (0.031-0.85)	-0.034% (-0.99-1%)	-0.051% (-1.2-1.1%)	-0.00015 (-0.0034-0.0033)	-0.00039 (-0.012-0.012)
Northern Mariana Islands	1990	Male	All Ages	47% (38-57%)	53% (43-62%)	0.61 (0.084-1.3)	3.3% (2-5.7%)	3.6% (1.8-6.8%)	2.8 (1.3-5.4)	170 (97-300)
Northern Mariana Islands	1990	Male	15 to 19	37% (22-54%)	63% (46-78%)	0.28 (0.033-0.75)	1.8% (0.88-3.4%)	2.4% (0.78-5.3%)	0.027 (0.0082-0.062)	2.8 (1.3-5.6)
Northern Mariana Islands	1990	Male	20 to 24	55% (36-73%)	45% (27-64%)	0.75 (0.09-1.8)	4.4% (2-8.1%)	5.4% (1.4-11%)	0.14 (0.035-0.3)	13 (5.5-24)
Northern Mariana Islands	1990	Male	25 to 29	62% (43-78%)	38% (22-57%)	1 (0.13-2.4)	5.5% (2.7-9.7%)	6.9% (2.4-14%)	0.34 (0.11-0.67)	29 (14-52)
Northern Mariana Islands	1990	Male	30 to 34	63% (42-80%)	37% (20-58%)	1.1 (0.15-2.6)	5.4% (2.9-9.3%)	6.7% (2.7-13%)	0.42 (0.15-0.84)	33 (17-58)
Northern Mariana Islands	1990	Male	35 to 39	62% (44-78%)	38% (22-56%)	0.96 (0.13-2.3)	5.1% (3-8.6%)	6.6% (3.2-12%)	0.43 (0.18-0.83)	29 (16-50)
Northern Mariana Islands	1990	Male	40 to 44	59% (41-76%)	41% (24-59%)	0.84 (0.1-2.1)	4.4% (2.6-7.5%)	5.5% (2.9-9.9%)	0.4 (0.18-0.76)	24 (13-43)
Northern Mariana Islands	1990	Male	45 to 49	55% (34-74%)	45% (26-66%)	0.73 (0.078-2)	4.1% (2.5-7.1%)	4.9% (2.7-9.2%)	0.35 (0.18-0.68)	18 (10-33)
Northern Mariana Islands	1990	Male	50 to 54	52% (33-72%)	48% (28-67%)	0.61 (0.066-1.7)	3.5% (2.1-6.5%)	4.2% (2.2-8.3%)	0.29 (0.14-0.58)	13 (6.8-24)
Northern Mariana Islands	1990	Male	55 to 59	50% (32-70%)	50% (30-68%)	0.49 (0.055-1.4)	2.2% (1.2-4.7%)	2.5% (1.2-5.6%)	0.14 (0.06-0.32)	5.3 (2.6-12)
Northern Mariana Islands	1990	Male	60 to 64	48% (27-68%)	52% (32-73%)	0.41 (0.052-1.2)	1.5% (0.71-3.7%)	1.7% (0.71-4.4%)	0.081 (0.033-0.2)	2.6 (1.2-6.1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	1990	Male	65 to 69	46% (27-66%)	54% (34-73%)	0.38 (0.046-1.1)	1.3% (0.58-3.2%)	1.5% (0.65-3.8%)	0.081 (0.032-0.21)	2 (0.87-5.1)
Northern Mariana Islands	1990	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.33 (0.038-0.93)	1% (0.39-2.6%)	1.1% (0.4-3%)	0.04 (0.014-0.11)	0.81 (0.3-2.1)
Northern Mariana Islands	1990	Male	75 to 79	39% (23-58%)	61% (42-77%)	0.29 (0.034-0.78)	0.8% (0.27-2.1%)	0.87% (0.28-2.4%)	0.029 (0.0092-0.078)	0.47 (0.16-1.2)
Northern Mariana Islands	1990	Male	80 to 84	37% (21-55%)	63% (45-79%)	0.25 (0.03-0.66)	0.87% (0.4-1.9%)	0.97% (0.42-2.2%)	0.021 (0.0091-0.048)	0.24 (0.11-0.54)
Northern Mariana Islands	1990	Male	85 to 89	35% (21-52%)	65% (48-79%)	0.24 (0.026-0.67)	0.54% (0.12-1.5%)	0.56% (0.089-1.6%)	0.0069 (0.0011-0.02)	0.059 (0.013-0.17)
Northern Mariana Islands	1990	Male	90 to 94	37% (22-54%)	63% (46-78%)	0.25 (0.026-0.65)	0.54% (0.14-1.4%)	0.57% (0.13-1.5%)	0.0031 (0.0007-0.0082)	0.018 (0.0048-0.046)
Northern Mariana Islands	1990	Male	95 plus	40% (22-62%)	60% (38-78%)	0.28 (0.031-0.77)	0.31% (-0.13-1.2%)	0.24% (-0.29-1.2%)	0.00041 (-0.00049-0.0021)	0.0022 (-0.001-0.0085)
Northern Mariana Islands	1995	Female	All Ages	20% (13-29%)	80% (71-87%)	0.27 (0.036-0.66)	0.89% (0.49-1.7%)	0.85% (0.25-2.3%)	0.55 (0.15-1.6)	46 (24-91)
Northern Mariana Islands	1995	Female	15 to 19	19% (9-36%)	81% (64-91%)	0.25 (0.031-0.68)	0.81% (0.36-1.6%)	1.2% (0.18-3.4%)	0.009 (0.0013-0.028)	1.4 (0.63-3.1)
Northern Mariana Islands	1995	Female	20 to 24	24% (10-41%)	76% (59-90%)	0.38 (0.042-1)	1.4% (0.78-2.5%)	1.6% (0.3-4.1%)	0.04 (0.0076-0.11)	7.3 (3.8-13)
Northern Mariana Islands	1995	Female	25 to 29	25% (10-43%)	75% (57-90%)	0.39 (0.047-1.1)	1.5% (0.9-2.6%)	1.8% (0.56-4.2%)	0.074 (0.022-0.18)	11 (6.2-19)
Northern Mariana Islands	1995	Female	30 to 34	26% (9.9-46%)	74% (54-90%)	0.41 (0.048-1.1)	1.4% (0.76-2.5%)	1.7% (0.59-4%)	0.074 (0.024-0.18)	8.4 (4.5-15)
Northern Mariana Islands	1995	Female	35 to 39	27% (11-45%)	73% (55-89%)	0.43 (0.054-1.1)	1.3% (0.64-2.6%)	1.9% (0.66-4.2%)	0.093 (0.031-0.22)	7.3 (3.5-15)
Northern Mariana Islands	1995	Female	40 to 44	26% (12-44%)	74% (56-88%)	0.41 (0.048-1.1)	1.2% (0.53-2.5%)	1.8% (0.72-4.2%)	0.084 (0.031-0.2)	5 (2.2-11)
Northern Mariana Islands	1995	Female	45 to 49	26% (11-45%)	74% (55-89%)	0.35 (0.039-1)	0.87% (0.25-2.3%)	1.4% (0.47-3.6%)	0.063 (0.02-0.18)	3 (0.83-8.5)
Northern Mariana Islands	1995	Female	50 to 54	25% (11-45%)	75% (55-89%)	0.31 (0.032-0.97)	0.62% (-0.066-2.1%)	1.1% (0.14-3.2%)	0.035 (0.0047-0.12)	1.3 (-0.13-4.5)
Northern Mariana Islands	1995	Female	55 to 59	24% (11-44%)	76% (56-89%)	0.28 (0.026-0.88)	0.47% (-0.25-1.8%)	0.85% (-0.042-2.8%)	0.029 (-0.0015-0.096)	0.81 (-0.43-3.2)
Northern Mariana Islands	1995	Female	60 to 64	25% (11-45%)	75% (55-89%)	0.27 (0.023-0.86)	0.22% (-0.56-1.5%)	0.47% (-0.45-2.2%)	0.016 (-0.016-0.076)	0.31 (-0.77-2)
Northern Mariana Islands	1995	Female	65 to 69	26% (12-45%)	74% (55-88%)	0.27 (0.028-0.88)	0.19% (-0.58-1.6%)	0.42% (-0.49-2.3%)	0.015 (-0.018-0.078)	0.21 (-0.67-1.8)
Northern Mariana Islands	1995	Female	70 to 74	26% (11-44%)	74% (56-89%)	0.25 (0.027-0.76)	0.05% (-0.62-1.3%)	0.18% (-0.61-1.7%)	0.0075 (-0.024-0.075)	0.053 (-0.65-1.3)
Northern Mariana Islands	1995	Female	75 to 79	24% (10-43%)	76% (57-90%)	0.23 (0.024-0.67)	0.15% (-0.49-1.2%)	0.29% (-0.46-1.7%)	0.013 (-0.02-0.072)	0.13 (-0.39-1)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	1995	Female	80 to 84	23% (11-42%)	77% (58-89%)	0.21 (0.022-0.6)	-0.0072% (-0.63-1%)	0.039% (-0.67-1.3%)	0.002 (-0.031-0.061)	-0.0039 (-0.39-0.68)
Northern Mariana Islands	1995	Female	85 to 89	24% (11-42%)	76% (58-89%)	0.22 (0.024-0.64)	-0.032% (-0.71-0.95%)	-0.0041% (-0.75-1.2%)	-0.00012 (-0.031-0.044)	-0.011 (-0.26-0.33)
Northern Mariana Islands	1995	Female	90 to 94	28% (12-47%)	72% (53-88%)	0.25 (0.026-0.74)	-0.065% (-0.66-0.93%)	-0.058% (-0.71-1.1%)	-0.00099 (-0.012-0.019)	-0.007 (-0.069-0.098)
Northern Mariana Islands	1995	Female	95 plus	32% (13-54%)	68% (46-87%)	0.3 (0.031-0.88)	0.052% (-0.9-1.2%)	0.041% (-1-1.3%)	0.00019 (-0.0047-0.0058)	0.00095 (-0.016-0.021)
Northern Mariana Islands	1995	Male	All Ages	45% (36-55%)	55% (45-64%)	0.61 (0.078-1.4)	3.3% (2-5.6%)	3.7% (2-6.7%)	3.2 (1.6-6.3)	200 (120-350)
Northern Mariana Islands	1995	Male	15 to 19	36% (21-54%)	64% (46-79%)	0.29 (0.035-0.8)	1.8% (0.87-3.4%)	2.5% (0.83-5.4%)	0.031 (0.0098-0.073)	3.4 (1.6-6.8)
Northern Mariana Islands	1995	Male	20 to 24	54% (35-71%)	46% (29-65%)	0.78 (0.095-2)	4.4% (2-8%)	5.6% (1.5-11%)	0.15 (0.041-0.32)	14 (6.2-27)
Northern Mariana Islands	1995	Male	25 to 29	61% (42-78%)	39% (22-58%)	1.1 (0.14-2.6)	5.5% (2.8-9.3%)	7.1% (2.5-14%)	0.34 (0.12-0.69)	31 (15-54)
Northern Mariana Islands	1995	Male	30 to 34	62% (42-79%)	38% (21-58%)	1.2 (0.16-2.9)	5.4% (3-8.9%)	7% (2.8-13%)	0.43 (0.16-0.86)	36 (18-63)
Northern Mariana Islands	1995	Male	35 to 39	61% (42-77%)	39% (23-58%)	1 (0.12-2.4)	5.2% (3.1-8.6%)	6.9% (3.4-12%)	0.52 (0.23-0.99)	37 (20-64)
Northern Mariana Islands	1995	Male	40 to 44	58% (41-76%)	42% (24-59%)	0.88 (0.11-2.1)	4.5% (2.8-7.6%)	5.8% (3.2-10%)	0.46 (0.22-0.88)	28 (16-50)
Northern Mariana Islands	1995	Male	45 to 49	55% (33-73%)	45% (27-67%)	0.76 (0.089-2)	4.2% (2.6-7.1%)	5.2% (3-9.2%)	0.45 (0.23-0.83)	24 (14-42)
Northern Mariana Islands	1995	Male	50 to 54	52% (32-71%)	48% (29-68%)	0.63 (0.071-1.7)	3.7% (2.2-6.5%)	4.5% (2.5-8.3%)	0.32 (0.16-0.63)	14 (8-27)
Northern Mariana Islands	1995	Male	55 to 59	49% (31-69%)	51% (31-69%)	0.51 (0.053-1.4)	2.3% (1.3-4.7%)	2.7% (1.4-5.8%)	0.18 (0.085-0.42)	7.2 (3.7-15)
Northern Mariana Islands	1995	Male	60 to 64	47% (28-67%)	53% (33-72%)	0.43 (0.043-1.3)	1.6% (0.79-3.7%)	1.8% (0.83-4.4%)	0.11 (0.046-0.28)	3.5 (1.6-8.4)
Northern Mariana Islands	1995	Male	65 to 69	45% (27-64%)	55% (36-73%)	0.4 (0.044-1.1)	1.4% (0.65-3.5%)	1.7% (0.73-4.1%)	0.088 (0.037-0.22)	2.2 (1-5.5)
Northern Mariana Islands	1995	Male	70 to 74	43% (26-61%)	57% (39-74%)	0.35 (0.046-0.96)	1.1% (0.43-2.8%)	1.2% (0.46-3.3%)	0.058 (0.022-0.15)	1.2 (0.47-3.1)
Northern Mariana Islands	1995	Male	75 to 79	39% (23-58%)	61% (42-77%)	0.3 (0.038-0.84)	0.87% (0.35-2.3%)	0.96% (0.34-2.7%)	0.033 (0.011-0.091)	0.53 (0.21-1.4)
Northern Mariana Islands	1995	Male	80 to 84	36% (20-54%)	64% (46-80%)	0.27 (0.033-0.72)	0.92% (0.43-2.2%)	1% (0.47-2.5%)	0.03 (0.013-0.073)	0.34 (0.16-0.83)
Northern Mariana Islands	1995	Male	85 to 89	34% (20-51%)	66% (49-80%)	0.25 (0.03-0.71)	0.59% (0.17-1.8%)	0.62% (0.14-2%)	0.012 (0.0025-0.039)	0.1 (0.028-0.31)
Northern Mariana Islands	1995	Male	90 to 94	36% (22-53%)	64% (47-78%)	0.26 (0.03-0.74)	0.58% (0.17-1.7%)	0.63% (0.17-1.8%)	0.0047 (0.0012-0.014)	0.027 (0.0082-0.076)
Northern Mariana Islands	1995	Male	95 plus	40% (22-60%)	60% (40-78%)	0.3 (0.033-0.82)	0.36% (-0.083-1.5%)	0.29% (-0.23-1.5%)	0.00058 (-0.00046-0.003)	0.0029 (-0.00068-0.012)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	2000	Female	All Ages	20% (13-29%)	80% (71-87%)	0.34 (0.1-0.76)	0.95% (0.48-1.9%)	0.92% (0.19-2.5%)	0.65 (0.12-1.7)	56 (28-110)
Northern Mariana Islands	2000	Female	15 to 19	19% (8.6-35%)	81% (65-91%)	0.32 (0.067-0.81)	0.89% (0.44-1.7%)	1.5% (0.34-3.7%)	0.012 (0.0026-0.034)	1.8 (0.86-3.7)
Northern Mariana Islands	2000	Female	20 to 24	24% (10-42%)	76% (58-90%)	0.5 (0.11-1.2)	1.6% (0.85-2.8%)	2.1% (0.51-4.9%)	0.053 (0.013-0.14)	9.2 (4.8-16)
Northern Mariana Islands	2000	Female	25 to 29	25% (11-44%)	75% (56-89%)	0.51 (0.12-1.3)	1.6% (0.96-2.9%)	2.2% (0.77-4.9%)	0.1 (0.036-0.23)	14 (7.8-25)
Northern Mariana Islands	2000	Female	30 to 34	26% (10-45%)	74% (55-90%)	0.52 (0.12-1.3)	1.5% (0.76-2.7%)	2% (0.73-4.5%)	0.092 (0.032-0.21)	10 (5-18)
Northern Mariana Islands	2000	Female	35 to 39	27% (11-46%)	73% (54-89%)	0.54 (0.13-1.2)	1.4% (0.66-2.6%)	2.1% (0.77-4.4%)	0.11 (0.038-0.25)	8.7 (4-17)
Northern Mariana Islands	2000	Female	40 to 44	26% (13-44%)	74% (56-87%)	0.51 (0.11-1.3)	1.2% (0.45-2.6%)	2% (0.76-4.4%)	0.096 (0.035-0.22)	5.7 (2.1-12)
Northern Mariana Islands	2000	Female	45 to 49	26% (11-43%)	74% (57-89%)	0.44 (0.094-1.1)	0.85% (0.1-2.2%)	1.5% (0.41-3.8%)	0.083 (0.021-0.23)	3.8 (0.46-10)
Northern Mariana Islands	2000	Female	50 to 54	25% (12-43%)	75% (57-88%)	0.39 (0.077-1.1)	0.57% (-0.22-2.1%)	1.1% (-0.048-3.5%)	0.043 (-0.0018-0.14)	1.4 (-0.53-5.6)
Northern Mariana Islands	2000	Female	55 to 59	24% (11-43%)	76% (57-89%)	0.35 (0.064-0.97)	0.38% (-0.43-1.8%)	0.81% (-0.16-2.7%)	0.029 (-0.0058-0.1)	0.7 (-0.77-3.3)
Northern Mariana Islands	2000	Female	60 to 64	24% (11-43%)	76% (57-89%)	0.33 (0.066-0.93)	0.1% (-0.75-1.4%)	0.38% (-0.64-2.2%)	0.016 (-0.025-0.091)	0.18 (-1.3-2.4)
Northern Mariana Islands	2000	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.33 (0.067-0.95)	0.057% (-0.81-1.5%)	0.3% (-0.78-2.3%)	0.011 (-0.029-0.082)	0.071 (-0.99-1.9)
Northern Mariana Islands	2000	Female	70 to 74	25% (11-42%)	75% (58-89%)	0.31 (0.061-0.81)	-0.075% (-0.92-1.4%)	0.056% (-0.9-2%)	0.0026 (-0.039-0.087)	-0.082 (-1-1.7)
Northern Mariana Islands	2000	Female	75 to 79	24% (10-41%)	76% (59-90%)	0.29 (0.056-0.79)	0.06% (-0.76-1.3%)	0.2% (-0.73-1.7%)	0.0081 (-0.03-0.068)	0.047 (-0.59-1)
Northern Mariana Islands	2000	Female	80 to 84	24% (9.6-43%)	76% (57-90%)	0.28 (0.052-0.73)	-0.086% (-0.9-1.2%)	-0.038% (-0.98-1.4%)	-0.0017 (-0.047-0.07)	-0.055 (-0.59-0.74)
Northern Mariana Islands	2000	Female	85 to 89	25% (11-44%)	75% (56-89%)	0.3 (0.059-0.77)	-0.12% (-0.99-1.1%)	-0.1% (-1.1-1.4%)	-0.0047 (-0.052-0.062)	-0.052 (-0.44-0.48)
Northern Mariana Islands	2000	Female	90 to 94	28% (13-47%)	72% (53-87%)	0.34 (0.062-0.89)	-0.13% (-0.86-1.2%)	-0.13% (-0.95-1.3%)	-0.0033 (-0.024-0.034)	-0.021 (-0.13-0.19)
Northern Mariana Islands	2000	Female	95 plus	33% (14-56%)	67% (44-86%)	0.4 (0.074-1.1)	-0.043% (-1.1-1.5%)	-0.071% (-1.3-1.6%)	-0.00051 (-0.0094-0.012)	-0.0012 (-0.032-0.044)
Northern Mariana Islands	2000	Male	All Ages	45% (37-55%)	55% (45-63%)	0.81 (0.25-1.6)	3.8% (2.3-6.2%)	4.4% (2.3-7.8%)	4.2 (2.1-7.7)	260 (150-420)
Northern Mariana Islands	2000	Male	15 to 19	36% (21-54%)	64% (46-79%)	0.39 (0.093-1)	2% (1-3.6%)	3.1% (1.2-6.4%)	0.04 (0.015-0.085)	4.2 (2.1-7.8)
Northern Mariana Islands	2000	Male	20 to 24	54% (35-71%)	46% (29-65%)	1.1 (0.28-2.4)	5.1% (2.6-9%)	7.1% (2.5-14%)	0.2 (0.068-0.4)	18 (8.9-33)
Northern Mariana Islands	2000	Male	25 to 29	61% (41-77%)	39% (23-59%)	1.4 (0.43-3.1)	6.2% (3.4-10%)	8.7% (3.6-16%)	0.46 (0.18-0.89)	40 (21-68)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	2000	Male	30 to 34	62% (41-78%)	38% (22-59%)	1.5 (0.45-3.2)	6% (3.4-9.6%)	8.4% (3.8-15%)	0.54 (0.23-1)	44 (24-73)
Northern Mariana Islands	2000	Male	35 to 39	61% (43-77%)	39% (23-57%)	1.3 (0.35-2.8)	5.7% (3.3-9.3%)	8% (4.1-13%)	0.65 (0.3-1.2)	45 (25-75)
Northern Mariana Islands	2000	Male	40 to 44	58% (41-76%)	42% (24-59%)	1.2 (0.29-2.5)	4.9% (2.9-8.2%)	6.6% (3.6-11%)	0.56 (0.27-1)	34 (19-59)
Northern Mariana Islands	2000	Male	45 to 49	55% (33-73%)	45% (27-67%)	1 (0.23-2.4)	4.5% (2.7-7.7%)	5.9% (3.3-10%)	0.63 (0.31-1.2)	33 (19-58)
Northern Mariana Islands	2000	Male	50 to 54	52% (32-69%)	48% (31-68%)	0.84 (0.19-2)	4% (2.3-7.2%)	5% (2.8-9.4%)	0.44 (0.22-0.87)	20 (11-37)
Northern Mariana Islands	2000	Male	55 to 59	49% (31-68%)	51% (32-69%)	0.68 (0.15-1.7)	2.5% (1.3-5.2%)	3% (1.4-6.6%)	0.22 (0.096-0.5)	8.5 (4.1-19)
Northern Mariana Islands	2000	Male	60 to 64	47% (28-67%)	53% (33-72%)	0.57 (0.13-1.5)	1.7% (0.7-4.2%)	2.1% (0.78-5.1%)	0.15 (0.054-0.38)	4.7 (1.9-12)
Northern Mariana Islands	2000	Male	65 to 69	45% (26-64%)	55% (36-74%)	0.51 (0.13-1.3)	1.5% (0.61-3.8%)	1.8% (0.69-4.6%)	0.1 (0.037-0.26)	2.6 (1-6.7)
Northern Mariana Islands	2000	Male	70 to 74	42% (25-60%)	58% (40-75%)	0.45 (0.11-1.1)	1.1% (0.37-3%)	1.4% (0.42-3.5%)	0.07 (0.021-0.19)	1.4 (0.45-3.8)
Northern Mariana Islands	2000	Male	75 to 79	39% (23-57%)	61% (43-77%)	0.39 (0.096-0.94)	0.95% (0.3-2.5%)	1.1% (0.3-2.9%)	0.034 (0.01-0.096)	0.56 (0.17-1.5)
Northern Mariana Islands	2000	Male	80 to 84	36% (20-55%)	64% (45-80%)	0.35 (0.086-0.83)	1% (0.42-2.5%)	1.2% (0.47-2.8%)	0.034 (0.013-0.082)	0.39 (0.16-0.93)
Northern Mariana Islands	2000	Male	85 to 89	35% (20-52%)	65% (48-80%)	0.33 (0.08-0.8)	0.65% (0.12-2%)	0.69% (0.082-2.2%)	0.016 (0.0019-0.052)	0.14 (0.024-0.43)
Northern Mariana Islands	2000	Male	90 to 94	36% (22-54%)	64% (46-78%)	0.34 (0.076-0.87)	0.62% (0.13-1.8%)	0.66% (0.095-2%)	0.0073 (0.001-0.022)	0.042 (0.0091-0.12)
Northern Mariana Islands	2000	Male	95 plus	40% (22-60%)	60% (40-78%)	0.38 (0.076-1)	0.41% (-0.15-1.8%)	0.33% (-0.33-1.9%)	0.00086 (-0.00083-0.0048)	0.0044 (-0.0016-0.019)
Northern Mariana Islands	2005	Female	All Ages	20% (12-29%)	80% (71-88%)	0.42 (0.17-0.85)	1.1% (0.54-2%)	1.1% (0.19-2.7%)	0.87 (0.16-2.3)	72 (35-140)
Northern Mariana Islands	2005	Female	15 to 19	19% (8.5-34%)	81% (66-91%)	0.39 (0.1-0.99)	0.98% (0.5-1.9%)	1.8% (0.51-4.3%)	0.016 (0.0045-0.04)	2.3 (1.1-4.6)
Northern Mariana Islands	2005	Female	20 to 24	23% (9.8-41%)	77% (59-90%)	0.61 (0.16-1.4)	1.8% (0.98-3%)	2.5% (0.75-5.6%)	0.073 (0.021-0.17)	12 (6.2-21)
Northern Mariana Islands	2005	Female	25 to 29	24% (10-42%)	76% (58-90%)	0.62 (0.18-1.5)	1.8% (1.1-3%)	2.5% (0.98-5.3%)	0.14 (0.05-0.3)	18 (11-31)
Northern Mariana Islands	2005	Female	30 to 34	25% (9.6-45%)	75% (55-90%)	0.64 (0.2-1.5)	1.6% (0.88-2.9%)	2.4% (0.93-5.2%)	0.12 (0.046-0.27)	13 (6.6-23)
Northern Mariana Islands	2005	Female	35 to 39	25% (11-44%)	75% (56-89%)	0.66 (0.2-1.4)	1.5% (0.73-3.1%)	2.4% (0.95-5.1%)	0.15 (0.053-0.33)	11 (5.2-22)
Northern Mariana Islands	2005	Female	40 to 44	25% (12-42%)	75% (58-88%)	0.6 (0.17-1.4)	1.3% (0.47-2.8%)	2.2% (0.78-4.7%)	0.12 (0.041-0.26)	7 (2.6-15)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	2005	Female	45 to 49	24% (11-41%)	76% (59-89%)	0.52 (0.13-1.3)	0.93% (0.099-2.5%)	1.6% (0.42-4.1%)	0.12 (0.031-0.31)	5.5 (0.59-15)
Northern Mariana Islands	2005	Female	50 to 54	24% (11-42%)	76% (58-89%)	0.46 (0.12-1.2)	0.6% (-0.38-2.3%)	1.1% (-0.097-3.6%)	0.058 (-0.0053-0.19)	2 (-1.3-7.9)
Northern Mariana Islands	2005	Female	55 to 59	23% (11-42%)	77% (58-89%)	0.42 (0.1-1.1)	0.4% (-0.55-2%)	0.84% (-0.3-3%)	0.033 (-0.012-0.12)	0.83 (-1.1-4)
Northern Mariana Islands	2005	Female	60 to 64	23% (10-42%)	77% (58-90%)	0.4 (0.095-1.1)	0.14% (-0.87-1.6%)	0.45% (-0.7-2.4%)	0.023 (-0.036-0.12)	0.31 (-1.8-3.5)
Northern Mariana Islands	2005	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.41 (0.092-1)	0.09% (-0.95-1.7%)	0.36% (-0.89-2.3%)	0.014 (-0.035-0.091)	0.12 (-1.3-2.1)
Northern Mariana Islands	2005	Female	70 to 74	24% (11-41%)	76% (59-89%)	0.38 (0.091-0.94)	-0.05% (-0.96-1.5%)	0.091% (-0.99-1.9%)	0.0046 (-0.05-0.1)	-0.063 (-1.3-2)
Northern Mariana Islands	2005	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.36 (0.083-0.91)	0.086% (-0.86-1.4%)	0.23% (-0.82-1.9%)	0.0091 (-0.033-0.071)	0.066 (-0.65-1)
Northern Mariana Islands	2005	Female	80 to 84	23% (9.2-43%)	77% (57-91%)	0.35 (0.079-0.86)	-0.032% (-1.1-3%)	0.028% (-1.1-1.6%)	0.0015 (-0.057-0.079)	-0.022 (-0.72-0.85)
Northern Mariana Islands	2005	Female	85 to 89	24% (11-43%)	76% (57-89%)	0.37 (0.087-0.9)	-0.076% (-1.1-3%)	-0.057% (-1.2-1.6%)	-0.003 (-0.059-0.08)	-0.037 (-0.48-0.64)
Northern Mariana Islands	2005	Female	90 to 94	27% (12-47%)	73% (53-88%)	0.42 (0.095-1)	-0.075% (-0.91-1.3%)	-0.076% (-1.1-5%)	-0.0026 (-0.034-0.052)	-0.016 (-0.19-0.28)
Northern Mariana Islands	2005	Female	95 plus	32% (13-55%)	68% (45-87%)	0.5 (0.11-1.3)	-0.074% (-1.4-1.7%)	-0.13% (-1.6-1.9%)	-0.0015 (-0.02-0.023)	-0.0035 (-0.065-0.083)
Northern Mariana Islands	2005	Male	All Ages	45% (36-54%)	55% (46-64%)	1 (0.4-1.9)	4.2% (2.5-6.8%)	4.8% (2.5-8.4%)	5.6 (2.7-10)	340 (190-570)
Northern Mariana Islands	2005	Male	15 to 19	35% (20-52%)	65% (48-80%)	0.47 (0.14-1.1)	2.1% (1.1-4%)	3.5% (1.3-7.2%)	0.05 (0.018-0.11)	5.2 (2.6-9.6)
Northern Mariana Islands	2005	Male	20 to 24	53% (34-71%)	47% (29-66%)	1.3 (0.42-2.7)	5.7% (2.9-9.7%)	8.2% (3.1-16%)	0.26 (0.1-0.51)	23 (12-41)
Northern Mariana Islands	2005	Male	25 to 29	60% (41-77%)	40% (23-59%)	1.8 (0.67-3.5)	6.4% (3.5-10%)	8.6% (3.7-15%)	0.65 (0.27-1.2)	56 (29-91)
Northern Mariana Islands	2005	Male	30 to 34	61% (40-78%)	39% (22-60%)	1.9 (0.71-3.7)	6.3% (3.4-10%)	8.4% (3.7-14%)	0.74 (0.32-1.3)	58 (31-94)
Northern Mariana Islands	2005	Male	35 to 39	60% (41-76%)	40% (24-59%)	1.6 (0.56-3.2)	6.2% (3.6-10%)	8.7% (4.3-15%)	0.86 (0.4-1.6)	59 (32-97)
Northern Mariana Islands	2005	Male	40 to 44	57% (40-75%)	43% (25-60%)	1.4 (0.47-3)	5.3% (3-8.9%)	7.1% (3.7-12%)	0.71 (0.33-1.3)	43 (23-74)
Northern Mariana Islands	2005	Male	45 to 49	53% (32-72%)	47% (28-68%)	1.3 (0.38-2.8)	4.8% (2.7-8.3%)	6.1% (3.2-11%)	0.92 (0.45-1.7)	48 (26-85)
Northern Mariana Islands	2005	Male	50 to 54	50% (31-69%)	50% (31-69%)	1 (0.3-2.3)	4.2% (2.2-7.7%)	5.2% (2.6-10%)	0.63 (0.3-1.3)	28 (14-54)
Northern Mariana Islands	2005	Male	55 to 59	48% (30-67%)	52% (33-70%)	0.83 (0.25-2)	2.6% (1.2-5.7%)	3.1% (1.3-6.9%)	0.27 (0.1-0.61)	10 (4.5-23)
Northern Mariana Islands	2005	Male	60 to 64	46% (27-66%)	54% (34-73%)	0.69 (0.2-1.7)	1.9% (0.71-4.6%)	2.2% (0.77-5.6%)	0.21 (0.071-0.55)	6.8 (2.5-17)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	2005	Male	65 to 69	44% (25-62%)	56% (38-75%)	0.63 (0.19-1.5)	1.7% (0.57-4.3%)	2% (0.66-4.9%)	0.13 (0.039-0.32)	3.2 (1.1-8.1)
Northern Mariana Islands	2005	Male	70 to 74	41% (24-58%)	59% (42-76%)	0.55 (0.18-1.3)	1.3% (0.34-3.4%)	1.5% (0.39-4%)	0.092 (0.023-0.24)	1.9 (0.46-4.9)
Northern Mariana Islands	2005	Male	75 to 79	37% (22-56%)	63% (44-78%)	0.47 (0.15-1.1)	1.1% (0.3-2.8%)	1.2% (0.31-3.2%)	0.039 (0.0099-0.1)	0.64 (0.18-1.7)
Northern Mariana Islands	2005	Male	80 to 84	34% (19-53%)	66% (47-81%)	0.41 (0.12-0.95)	1.1% (0.42-2.5%)	1.3% (0.45-2.9%)	0.04 (0.014-0.09)	0.46 (0.17-1.1)
Northern Mariana Islands	2005	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.39 (0.11-0.89)	0.75% (0.13-2.1%)	0.8% (0.098-2.4%)	0.022 (0.0027-0.066)	0.19 (0.033-0.54)
Northern Mariana Islands	2005	Male	90 to 94	34% (21-51%)	66% (49-79%)	0.4 (0.12-0.93)	0.68% (0.14-1.9%)	0.71% (0.093-2.1%)	0.011 (0.0014-0.032)	0.065 (0.013-0.19)
Northern Mariana Islands	2005	Male	95 plus	38% (20-59%)	62% (41-80%)	0.44 (0.11-1.1)	0.52% (-0.094-2%)	0.43% (-0.29-2.1%)	0.0018 (-0.0012-0.009)	0.0091 (-0.0017-0.036)
Northern Mariana Islands	2010	Female	All Ages	20% (13-29%)	80% (71-87%)	0.47 (0.2-0.86)	1.2% (0.59-2.2%)	1.2% (0.28-3.1%)	1.2 (0.25-2.9)	92 (45-170)
Northern Mariana Islands	2010	Female	15 to 19	18% (8.3-33%)	82% (67-92%)	0.42 (0.12-1)	0.99% (0.52-1.9%)	1.9% (0.63-4.5%)	0.018 (0.0056-0.044)	2.7 (1.4-5.3)
Northern Mariana Islands	2010	Female	20 to 24	23% (9.3-41%)	77% (59-91%)	0.64 (0.19-1.5)	1.8% (0.99-2.9%)	2.6% (0.86-5.7%)	0.082 (0.026-0.19)	14 (7.4-23)
Northern Mariana Islands	2010	Female	25 to 29	24% (9.7-42%)	76% (58-90%)	0.66 (0.22-1.4)	1.8% (1.1-3%)	2.5% (0.96-5.1%)	0.16 (0.065-0.34)	23 (13-38)
Northern Mariana Islands	2010	Female	30 to 34	25% (9.4-44%)	75% (56-91%)	0.69 (0.23-1.5)	1.7% (0.96-2.9%)	2.5% (1.1-5%)	0.14 (0.06-0.3)	15 (8.3-27)
Northern Mariana Islands	2010	Female	35 to 39	25% (11-44%)	75% (56-89%)	0.72 (0.24-1.6)	1.7% (0.8-3.3%)	2.6% (1.1-5.5%)	0.19 (0.071-0.4)	14 (6.5-28)
Northern Mariana Islands	2010	Female	40 to 44	25% (12-42%)	75% (58-88%)	0.67 (0.21-1.5)	1.4% (0.54-3%)	2.4% (0.86-5%)	0.15 (0.052-0.33)	8.8 (3.3-19)
Northern Mariana Islands	2010	Female	45 to 49	25% (11-43%)	75% (57-89%)	0.59 (0.17-1.4)	1.1% (0.18-2.8%)	1.9% (0.52-4.6%)	0.18 (0.051-0.48)	8.5 (1.5-23)
Northern Mariana Islands	2010	Female	50 to 54	25% (11-44%)	75% (56-89%)	0.52 (0.15-1.3)	0.73% (-0.32-2.6%)	1.3% (-0.042-4%)	0.09 (-0.0028-0.28)	3.2 (-1.5-12)
Northern Mariana Islands	2010	Female	55 to 59	24% (11-44%)	76% (56-89%)	0.48 (0.13-1.2)	0.51% (-0.6-2.2%)	1% (-0.37-3.2%)	0.045 (-0.017-0.16)	1.2 (-1.4-5.3)
Northern Mariana Islands	2010	Female	60 to 64	24% (11-43%)	76% (57-89%)	0.46 (0.13-1.2)	0.22% (-0.86-1.9%)	0.56% (-0.67-2.7%)	0.037 (-0.048-0.18)	0.61 (-2.5-5.5)
Northern Mariana Islands	2010	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.46 (0.12-1.1)	0.18% (-0.95-1.9%)	0.48% (-0.84-2.8%)	0.021 (-0.038-0.12)	0.26 (-1.4-2.8)
Northern Mariana Islands	2010	Female	70 to 74	24% (11-42%)	76% (58-89%)	0.43 (0.11-1.1)	0.04% (-0.93-1.7%)	0.21% (-0.95-2.4%)	0.012 (-0.056-0.15)	0.062 (-1.4-2.8)
Northern Mariana Islands	2010	Female	75 to 79	23% (10-42%)	77% (58-90%)	0.4 (0.1-0.97)	0.18% (-0.81-1.7%)	0.34% (-0.76-2.1%)	0.014 (-0.029-0.082)	0.13 (-0.61-1.3)
Northern Mariana Islands	2010	Female	80 to 84	23% (9.4-43%)	77% (57-91%)	0.39 (0.097-0.99)	0.04% (-0.93-1.4%)	0.11% (-0.97-1.8%)	0.0066 (-0.057-0.11)	0.032 (-0.73-1.2)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	2010	Female	85 to 89	24% (11-44%)	76% (56-89%)	0.41 (0.1-1)	-0.0031% (-1-1.5%)	0.025% (-1.1-1.7%)	0.0016 (-0.068-0.11)	-0.0014 (-0.57-0.84)
Northern Mariana Islands	2010	Female	90 to 94	28% (13-47%)	72% (53-87%)	0.47 (0.12-1.1)	-0.018% (-0.95-1.6%)	-0.018% (-1.1-1.8%)	-0.00073 (-0.044-0.075)	-0.0047 (-0.24-0.42)
Northern Mariana Islands	2010	Female	95 plus	33% (14-54%)	67% (46-86%)	0.55 (0.13-1.3)	0.021% (-1.3-1.9%)	-0.025% (-1.6-2%)	-0.00043 (-0.028-0.036)	0.0015 (-0.092-0.13)
Northern Mariana Islands	2010	Male	All Ages	45% (36-55%)	55% (45-64%)	1.1 (0.51-2)	4.8% (2.9-7.6%)	5.6% (3-9.3%)	8 (4.1-14)	470 (270-770)
Northern Mariana Islands	2010	Male	15 to 19	35% (20-53%)	65% (47-80%)	0.51 (0.16-1.2)	2.2% (1.2-3.8%)	3.8% (1.6-7.5%)	0.058 (0.024-0.12)	6.1 (3.4-11)
Northern Mariana Islands	2010	Male	20 to 24	53% (34-70%)	47% (30-66%)	1.4 (0.52-2.8)	5.9% (3.2-10%)	8.8% (3.8-16%)	0.32 (0.14-0.62)	29 (15-50)
Northern Mariana Islands	2010	Male	25 to 29	60% (40-76%)	40% (24-60%)	1.9 (0.81-3.8)	7.3% (4.2-11%)	11% (5.1-18%)	0.92 (0.4-1.6)	76 (42-120)
Northern Mariana Islands	2010	Male	30 to 34	61% (40-78%)	39% (22-60%)	2 (0.86-3.8)	7.2% (4.1-11%)	10% (5-17%)	1 (0.46-1.8)	77 (42-120)
Northern Mariana Islands	2010	Male	35 to 39	60% (41-77%)	40% (23-59%)	1.8 (0.71-3.5)	6.7% (3.8-11%)	9.4% (4.8-16%)	1.2 (0.56-2.1)	79 (42-130)
Northern Mariana Islands	2010	Male	40 to 44	57% (39-75%)	43% (25-61%)	1.6 (0.58-3.3)	5.8% (3.3-9.7%)	7.7% (4-13%)	0.98 (0.46-1.8)	57 (30-98)
Northern Mariana Islands	2010	Male	45 to 49	54% (33-72%)	46% (28-67%)	1.4 (0.49-2.9)	5.2% (2.9-8.9%)	6.6% (3.6-12%)	1.5 (0.72-2.8)	75 (40-130)
Northern Mariana Islands	2010	Male	50 to 54	51% (32-69%)	49% (31-68%)	1.1 (0.38-2.4)	4.5% (2.4-8.2%)	5.6% (2.8-10%)	0.98 (0.44-1.9)	43 (20-81)
Northern Mariana Islands	2010	Male	55 to 59	48% (30-68%)	52% (32-70%)	0.92 (0.3-2.1)	2.8% (1.2-6%)	3.3% (1.3-7.1%)	0.35 (0.13-0.79)	13 (5.6-29)
Northern Mariana Islands	2010	Male	60 to 64	46% (26-66%)	54% (34-74%)	0.77 (0.24-1.8)	2.1% (0.69-5.1%)	2.4% (0.72-6.1%)	0.32 (0.097-0.83)	10 (3.3-25)
Northern Mariana Islands	2010	Male	65 to 69	44% (26-63%)	56% (37-74%)	0.69 (0.24-1.5)	1.9% (0.57-4.5%)	2.2% (0.68-5.3%)	0.16 (0.049-0.4)	4.1 (1.2-9.7)
Northern Mariana Islands	2010	Male	70 to 74	41% (25-59%)	59% (41-75%)	0.6 (0.21-1.3)	1.4% (0.35-3.7%)	1.7% (0.4-4.3%)	0.12 (0.029-0.32)	2.4 (0.58-6.4)
Northern Mariana Islands	2010	Male	75 to 79	38% (23-56%)	62% (44-77%)	0.51 (0.17-1.2)	1.2% (0.34-3.1%)	1.4% (0.34-3.6%)	0.044 (0.011-0.11)	0.71 (0.19-1.8)
Northern Mariana Islands	2010	Male	80 to 84	35% (20-53%)	65% (47-80%)	0.46 (0.15-0.98)	1.2% (0.44-2.8%)	1.4% (0.5-3.2%)	0.049 (0.017-0.11)	0.56 (0.2-1.3)
Northern Mariana Islands	2010	Male	85 to 89	33% (19-50%)	67% (50-81%)	0.43 (0.14-0.96)	0.83% (0.15-2.3%)	0.89% (0.12-2.5%)	0.028 (0.0035-0.083)	0.24 (0.044-0.68)
Northern Mariana Islands	2010	Male	90 to 94	35% (21-52%)	65% (48-79%)	0.44 (0.14-0.97)	0.73% (0.15-2%)	0.76% (0.1-2.2%)	0.015 (0.002-0.042)	0.087 (0.017-0.24)
Northern Mariana Islands	2010	Male	95 plus	38% (20-59%)	62% (41-80%)	0.48 (0.13-1.1)	0.59% (-0.095-2.2%)	0.5% (-0.28-2.2%)	0.0031 (-0.0018-0.013)	0.015 (-0.0025-0.054)
Northern Mariana Islands	2016	Female	All Ages	21% (13-30%)	79% (70-87%)	0.49 (0.22-0.89)	1.2% (0.6-2.2%)	1.3% (0.29-3.2%)	1.5 (0.3-3.9)	120 (57-220)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	2016	Female	15 to 19	19% (8.9-33%)	81% (67-91%)	0.44 (0.14-1)	1% (0.52-1.9%)	2% (0.65-4.6%)	0.024 (0.0074-0.056)	3.4 (1.7-6.5)
Northern Mariana Islands	2016	Female	20 to 24	23% (9.7-40%)	77% (60-90%)	0.68 (0.21-1.5)	1.8% (1-3%)	2.7% (0.89-5.8%)	0.11 (0.034-0.24)	17 (9.5-28)
Northern Mariana Islands	2016	Female	25 to 29	24% (10-43%)	76% (57-90%)	0.69 (0.23-1.4)	1.8% (1.1-3%)	2.6% (1-5.2%)	0.22 (0.082-0.47)	30 (17-52)
Northern Mariana Islands	2016	Female	30 to 34	25% (9-45%)	75% (55-91%)	0.7 (0.24-1.5)	1.7% (0.96-2.9%)	2.6% (1.1-5.2%)	0.18 (0.07-0.37)	19 (10-34)
Northern Mariana Islands	2016	Female	35 to 39	26% (10-45%)	74% (55-90%)	0.74 (0.27-1.6)	1.7% (0.84-3.3%)	2.7% (1.1-5.7%)	0.23 (0.086-0.5)	18 (8.2-34)
Northern Mariana Islands	2016	Female	40 to 44	26% (12-43%)	74% (57-88%)	0.7 (0.23-1.5)	1.5% (0.55-3.2%)	2.5% (0.87-5.3%)	0.18 (0.058-0.4)	11 (3.9-23)
Northern Mariana Islands	2016	Female	45 to 49	25% (11-43%)	75% (57-89%)	0.6 (0.18-1.4)	1.1% (0.15-3%)	1.8% (0.46-4.7%)	0.26 (0.063-0.68)	12 (1.6-33)
Northern Mariana Islands	2016	Female	50 to 54	24% (11-43%)	76% (57-89%)	0.53 (0.15-1.3)	0.73% (-0.3-2.6%)	1.3% (-0.099-4.1%)	0.12 (-0.0083-0.39)	4.4 (-1.7-16)
Northern Mariana Islands	2016	Female	55 to 59	24% (11-43%)	76% (57-89%)	0.47 (0.13-1.2)	0.51% (-0.56-2.1%)	0.97% (-0.35-3.3%)	0.051 (-0.019-0.17)	1.4 (-1.5-6)
Northern Mariana Islands	2016	Female	60 to 64	24% (11-43%)	76% (57-89%)	0.46 (0.13-1.1)	0.21% (-0.84-1.8%)	0.52% (-0.74-2.6%)	0.047 (-0.069-0.25)	0.8 (-3.5-7.3)
Northern Mariana Islands	2016	Female	65 to 69	25% (11-43%)	75% (57-89%)	0.47 (0.12-1.1)	0.16% (-1-2%)	0.43% (-0.98-2.7%)	0.022 (-0.05-0.13)	0.27 (-1.7-3.2)
Northern Mariana Islands	2016	Female	70 to 74	25% (11-42%)	75% (58-89%)	0.45 (0.12-1.1)	0.024% (-1-1.9%)	0.18% (-1.1-2.5%)	0.012 (-0.077-0.17)	0.046 (-1.8-3.5)
Northern Mariana Islands	2016	Female	75 to 79	24% (11-42%)	76% (58-89%)	0.42 (0.12-1.1)	0.18% (-0.79-1.8%)	0.34% (-0.81-2.3%)	0.013 (-0.031-0.088)	0.14 (-0.61-1.4)
Northern Mariana Islands	2016	Female	80 to 84	23% (10-41%)	77% (59-90%)	0.4 (0.1-0.99)	0.043% (-0.89-1.6%)	0.11% (-0.98-2%)	0.0074 (-0.067-0.14)	0.041 (-0.83-1.5)
Northern Mariana Islands	2016	Female	85 to 89	24% (11-43%)	76% (57-89%)	0.42 (0.11-1)	0.0046% (-0.95-1.8%)	0.024% (-1.1-2%)	0.0018 (-0.08-0.14)	0.003 (-0.65-1.1)
Northern Mariana Islands	2016	Female	90 to 94	28% (12-47%)	72% (53-88%)	0.48 (0.12-1.1)	-0.015% (-1-1.7%)	-0.025% (-1.1-2%)	-0.0013 (-0.056-0.1)	-0.0049 (-0.3-0.54)
Northern Mariana Islands	2016	Female	95 plus	33% (13-55%)	67% (45-87%)	0.56 (0.13-1.4)	0.005% (-1.4-2%)	-0.052% (-1.8-2.2%)	-0.0011 (-0.038-0.048)	0.00047 (-0.13-0.17)
Northern Mariana Islands	2016	Male	All Ages	46% (37-56%)	54% (44-63%)	1.2 (0.56-2.1)	4.9% (2.9-7.6%)	5.7% (3-9.4%)	11 (5.1-19)	620 (350-1000)
Northern Mariana Islands	2016	Male	15 to 19	35% (20-53%)	65% (47-80%)	0.53 (0.19-1.2)	2.2% (1.2-4.1%)	3.9% (1.7-8%)	0.075 (0.032-0.16)	7.8 (4.2-14)
Northern Mariana Islands	2016	Male	20 to 24	53% (34-71%)	47% (29-66%)	1.4 (0.58-2.8)	6.1% (3.3-10%)	9% (4.2-16%)	0.42 (0.19-0.79)	37 (20-62)
Northern Mariana Islands	2016	Male	25 to 29	60% (40-76%)	40% (24-60%)	2 (0.88-3.8)	7.3% (4.2-12%)	11% (5-18%)	1.2 (0.52-2.2)	100 (54-160)
Northern Mariana Islands	2016	Male	30 to 34	61% (40-78%)	39% (22-60%)	2.1 (0.88-3.9)	7.3% (4.1-11%)	10% (4.9-17%)	1.3 (0.55-2.3)	96 (52-160)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Northern Mariana Islands	2016	Male	35 to 39	60% (41-78%)	40% (22-59%)	1.9 (0.79-3.8)	6.9% (4-11%)	9.5% (4.9-16%)	1.5 (0.7-2.8)	100 (54-170)
Northern Mariana Islands	2016	Male	40 to 44	58% (39-76%)	42% (24-61%)	1.6 (0.61-3.4)	5.9% (3.2-9.9%)	7.7% (4-14%)	1.2 (0.53-2.3)	70 (36-130)
Northern Mariana Islands	2016	Male	45 to 49	53% (32-72%)	47% (28-68%)	1.4 (0.49-2.9)	5.2% (2.9-8.9%)	6.6% (3.4-12%)	2.2 (1-4)	110 (58-190)
Northern Mariana Islands	2016	Male	50 to 54	50% (31-69%)	50% (31-69%)	1.2 (0.4-2.5)	4.6% (2.3-8.2%)	5.6% (2.7-10%)	1.4 (0.6-2.7)	61 (28-110)
Northern Mariana Islands	2016	Male	55 to 59	48% (30-68%)	52% (32-70%)	0.94 (0.31-2.1)	2.8% (1.1-6%)	3.3% (1.2-7.1%)	0.42 (0.14-0.94)	16 (6.4-35)
Northern Mariana Islands	2016	Male	60 to 64	46% (27-66%)	54% (34-73%)	0.77 (0.27-1.8)	2.1% (0.68-4.9%)	2.4% (0.72-5.8%)	0.45 (0.13-1.1)	14 (4.7-33)
Northern Mariana Islands	2016	Male	65 to 69	44% (26-63%)	56% (37-74%)	0.69 (0.25-1.5)	1.9% (0.63-4.3%)	2.2% (0.68-5.1%)	0.19 (0.061-0.45)	4.8 (1.6-11)
Northern Mariana Islands	2016	Male	70 to 74	41% (25-59%)	59% (41-75%)	0.61 (0.21-1.4)	1.5% (0.36-4%)	1.7% (0.4-4.6%)	0.15 (0.035-0.4)	3 (0.71-8.2)
Northern Mariana Islands	2016	Male	75 to 79	38% (22-57%)	62% (43-78%)	0.53 (0.18-1.2)	1.3% (0.37-3.3%)	1.4% (0.37-3.7%)	0.046 (0.012-0.12)	0.75 (0.21-1.9)
Northern Mariana Islands	2016	Male	80 to 84	35% (19-53%)	65% (47-81%)	0.47 (0.16-1)	1.3% (0.47-3%)	1.4% (0.52-3.3%)	0.059 (0.021-0.14)	0.68 (0.25-1.6)
Northern Mariana Islands	2016	Male	85 to 89	33% (19-49%)	67% (51-81%)	0.45 (0.15-1)	0.87% (0.16-2.5%)	0.92% (0.12-2.7%)	0.035 (0.0046-0.1)	0.3 (0.054-0.82)
Northern Mariana Islands	2016	Male	90 to 94	35% (21-53%)	65% (47-79%)	0.46 (0.15-1)	0.78% (0.16-2.2%)	0.8% (0.095-2.4%)	0.019 (0.0022-0.056)	0.11 (0.022-0.31)
Northern Mariana Islands	2016	Male	95 plus	39% (20-59%)	61% (41-80%)	0.51 (0.14-1.2)	0.64% (-0.1-2.2%)	0.53% (-0.33-2.3%)	0.0044 (-0.0027-0.019)	0.021 (-0.0036-0.077)
Puerto Rico	1990	Female	All Ages	35% (29-42%)	65% (58-71%)	1.2 (0.77-1.8)	2.3% (1.3-3.6%)	2.2% (0.88-4.2%)	240 (93-450)	9600 (5700-15000)
Puerto Rico	1990	Female	15 to 19	42% (28-58%)	58% (42-72%)	1.5 (0.85-2.6)	2.6% (1.7-3.9%)	6.9% (4-11%)	4.1 (2.4-6.5)	450 (290-660)
Puerto Rico	1990	Female	20 to 24	48% (32-64%)	52% (36-68%)	2.3 (1.3-3.6)	4.4% (3-6.2%)	8.3% (4.9-13%)	7.6 (4.4-12)	820 (540-1200)
Puerto Rico	1990	Female	25 to 29	47% (31-64%)	53% (36-69%)	2.1 (1.2-3.5)	4.3% (2.9-6.1%)	6.8% (4.1-10%)	9.4 (5.5-14)	900 (600-1300)
Puerto Rico	1990	Female	30 to 34	46% (29-63%)	54% (37-71%)	1.9 (1.1-3.2)	4% (2.7-5.8%)	6.6% (4.3-9.8%)	11 (6.9-16)	910 (610-1300)
Puerto Rico	1990	Female	35 to 39	45% (30-62%)	55% (38-70%)	1.9 (1-3.1)	3.9% (2.7-5.6%)	7% (4.7-10%)	13 (8.4-19)	910 (600-1300)
Puerto Rico	1990	Female	40 to 44	43% (28-62%)	57% (38-72%)	1.8 (0.95-2.9)	4.1% (2.7-6%)	7.8% (5.2-12%)	17 (11-25)	1000 (670-1500)
Puerto Rico	1990	Female	45 to 49	42% (26-59%)	58% (41-74%)	1.6 (0.86-2.7)	3.7% (2.3-5.9%)	7% (4.6-11%)	18 (11-27)	880 (550-1300)
Puerto Rico	1990	Female	50 to 54	38% (24-55%)	62% (45-76%)	1.4 (0.74-2.3)	3.4% (2-5.5%)	6% (3.6-9.2%)	19 (11-30)	810 (460-1300)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	1990	Female	55 to 59	33% (20-50%)	67% (50-80%)	1.2 (0.59-2.1)	3% (1.5-5.2%)	4.7% (2.6-7.8%)	21 (12-35)	740 (390-1300)
Puerto Rico	1990	Female	60 to 64	30% (18-46%)	70% (54-82%)	1 (0.51-1.8)	2.3% (0.88-4.4%)	3.4% (1.4-6.2%)	22 (9.1-40)	640 (250-1200)
Puerto Rico	1990	Female	65 to 69	27% (17-42%)	73% (58-83%)	0.89 (0.45-1.6)	1.8% (0.43-3.7%)	2.5% (0.72-4.9%)	23 (6.5-45)	560 (140-1100)
Puerto Rico	1990	Female	70 to 74	25% (16-37%)	75% (63-84%)	0.75 (0.38-1.4)	1.4% (0.21-3.1%)	1.8% (0.3-4.1%)	21 (3.6-48)	400 (65-940)
Puerto Rico	1990	Female	75 to 79	22% (13-34%)	78% (66-87%)	0.65 (0.32-1.2)	0.98% (-0.097-2.4%)	1.2% (-0.11-3.1%)	18 (-1.6-46)	280 (-28-700)
Puerto Rico	1990	Female	80 to 84	20% (11-32%)	80% (68-89%)	0.59 (0.28-1)	0.83% (-0.025-2.2%)	1% (-0.091-2.7%)	15 (-1.4-41)	170 (-5.7-460)
Puerto Rico	1990	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.58 (0.28-1)	0.69% (-0.21-2.1%)	0.76% (-0.37-2.4%)	11 (-5-33)	90 (-28-270)
Puerto Rico	1990	Female	90 to 94	22% (13-34%)	78% (66-87%)	0.65 (0.31-1.2)	0.67% (-0.38-2.3%)	0.66% (-0.63-2.6%)	5.7 (-5.5-22)	36 (-21-120)
Puerto Rico	1990	Female	95 plus	25% (14-41%)	75% (59-86%)	0.75 (0.31-1.5)	0.65% (-0.72-2.8%)	0.56% (-1.1-3.1%)	1.4 (-2.9-8)	6.8 (-7.9-29)
Puerto Rico	1990	Male	All Ages	57% (49-65%)	43% (35-51%)	2.7 (1.8-4.1)	10% (7.4-13%)	9.9% (6.6-14%)	1500 (1000-2100)	60000 (43000-77000)
Puerto Rico	1990	Male	15 to 19	57% (40-73%)	43% (27-60%)	2.5 (1.4-4.1)	8.9% (5.6-13%)	13% (7.7-21%)	28 (16-44)	2300 (1500-3600)
Puerto Rico	1990	Male	20 to 24	67% (50-81%)	33% (19-50%)	4.6 (2.7-7.2)	17% (11-24%)	22% (14-32%)	70 (43-100)	5300 (3600-7400)
Puerto Rico	1990	Male	25 to 29	71% (53-83%)	29% (17-47%)	4.6 (2.8-7.3)	15% (10-20%)	17% (11-24%)	76 (49-110)	5500 (3800-7500)
Puerto Rico	1990	Male	30 to 34	71% (53-84%)	29% (16-47%)	4.7 (2.8-7.7)	13% (9.2-17%)	15% (10-20%)	82 (55-110)	5300 (3800-7200)
Puerto Rico	1990	Male	35 to 39	71% (55-84%)	29% (16-45%)	5 (2.9-7.8)	15% (11-18%)	17% (13-22%)	110 (78-140)	6200 (4700-8000)
Puerto Rico	1990	Male	40 to 44	70% (55-84%)	30% (16-45%)	4.7 (2.7-7.4)	16% (13-20%)	20% (15-24%)	120 (94-160)	6500 (5000-8000)
Puerto Rico	1990	Male	45 to 49	68% (49-82%)	32% (18-51%)	4 (2.4-6.3)	16% (12-20%)	20% (15-24%)	120 (93-150)	5700 (4300-7200)
Puerto Rico	1990	Male	50 to 54	67% (48-82%)	33% (18-52%)	3.6 (2-5.6)	14% (10-18%)	18% (13-23%)	120 (90-160)	4900 (3700-6400)
Puerto Rico	1990	Male	55 to 59	65% (48-81%)	35% (19-52%)	3.4 (1.9-5.5)	13% (8.9-17%)	16% (11-21%)	130 (91-180)	4600 (3200-6200)
Puerto Rico	1990	Male	60 to 64	63% (44-80%)	37% (20-56%)	3.1 (1.7-5.1)	11% (6.8-15%)	13% (8.1-18%)	150 (92-200)	4200 (2700-5900)
Puerto Rico	1990	Male	65 to 69	62% (42-78%)	38% (22-58%)	2.6 (1.4-4.4)	8.8% (5.3-13%)	10% (6.2-15%)	150 (90-220)	3600 (2200-5300)
Puerto Rico	1990	Male	70 to 74	58% (40-74%)	42% (26-60%)	2.2 (1.2-3.7)	6.4% (3.5-10%)	7.4% (4-12%)	120 (66-200)	2400 (1300-3900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	1990	Male	75 to 79	53% (34-71%)	47% (29-66%)	1.8 (0.91-3.1)	4.9% (2.4-8.4%)	5.6% (2.6-9.8%)	100 (49-180)	1600 (760-2700)
Puerto Rico	1990	Male	80 to 84	47% (29-66%)	53% (34-71%)	1.6 (0.79-2.9)	3.9% (1.7-6.9%)	4.4% (1.9-8%)	71 (32-130)	820 (380-1400)
Puerto Rico	1990	Male	85 to 89	43% (26-60%)	57% (40-74%)	1.5 (0.7-2.6)	3.2% (1.2-6.1%)	3.6% (1.2-6.9%)	47 (16-91)	380 (150-720)
Puerto Rico	1990	Male	90 to 94	43% (28-61%)	57% (39-72%)	1.5 (0.77-2.8)	3% (0.82-6.1%)	3.2% (0.69-6.9%)	21 (4.4-45)	120 (32-240)
Puerto Rico	1990	Male	95 plus	46% (27-67%)	54% (33-73%)	1.7 (0.79-3.2)	3.2% (0.59-6.9%)	3.3% (0.25-7.7%)	5 (0.38-12)	20 (3.6-42)
Puerto Rico	1995	Female	All Ages	32% (26-39%)	68% (61-74%)	1.1 (0.7-1.6)	2.2% (1.3-3.4%)	2.3% (1-4.1%)	270 (120-500)	10000 (6000-16000)
Puerto Rico	1995	Female	15 to 19	39% (26-56%)	61% (44-74%)	1.5 (0.85-2.6)	2.6% (1.6-4%)	6.5% (3.6-11%)	4.2 (2.3-6.7)	450 (290-680)
Puerto Rico	1995	Female	20 to 24	44% (29-60%)	56% (40-71%)	2.2 (1.2-3.5)	4% (2.6-5.6%)	7.4% (4.3-12%)	6.3 (3.7-10)	730 (490-1100)
Puerto Rico	1995	Female	25 to 29	43% (28-60%)	57% (40-72%)	1.9 (1.1-3.1)	3.6% (2.4-5.1%)	5.2% (3-8%)	6.8 (3.9-10)	720 (490-1000)
Puerto Rico	1995	Female	30 to 34	40% (25-57%)	60% (43-75%)	1.7 (0.92-2.9)	3.3% (2.2-4.8%)	4.7% (2.9-7.3%)	9.2 (5.7-14)	820 (550-1200)
Puerto Rico	1995	Female	35 to 39	38% (25-56%)	62% (44-75%)	1.6 (0.87-2.7)	3.4% (2.3-4.9%)	5.3% (3.4-8%)	13 (8.1-20)	930 (620-1300)
Puerto Rico	1995	Female	40 to 44	38% (24-56%)	62% (44-76%)	1.5 (0.82-2.6)	3.4% (2.3-5.1%)	6% (4-8.8%)	15 (10-23)	940 (620-1400)
Puerto Rico	1995	Female	45 to 49	36% (23-52%)	64% (48-77%)	1.4 (0.76-2.3)	3.4% (2.1-5.2%)	6% (3.9-9%)	19 (13-30)	980 (620-1500)
Puerto Rico	1995	Female	50 to 54	33% (20-49%)	67% (51-80%)	1.2 (0.64-2.1)	3.2% (1.9-5%)	5.5% (3.5-8.2%)	22 (14-33)	920 (560-1400)
Puerto Rico	1995	Female	55 to 59	29% (18-45%)	71% (55-82%)	1.1 (0.54-1.9)	2.7% (1.4-4.5%)	4.3% (2.4-7%)	21 (12-35)	760 (400-1300)
Puerto Rico	1995	Female	60 to 64	27% (16-42%)	73% (58-84%)	0.95 (0.48-1.7)	2.3% (1-4.3%)	3.4% (1.7-6.2%)	25 (12-44)	730 (330-1400)
Puerto Rico	1995	Female	65 to 69	26% (17-40%)	74% (60-83%)	0.88 (0.43-1.5)	2% (0.73-3.7%)	2.8% (1.2-5.1%)	27 (12-50)	650 (250-1200)
Puerto Rico	1995	Female	70 to 74	24% (15-38%)	76% (62-85%)	0.76 (0.38-1.4)	1.5% (0.43-3.2%)	2.1% (0.65-4.3%)	27 (8.5-57)	520 (150-1100)
Puerto Rico	1995	Female	75 to 79	21% (13-34%)	79% (66-87%)	0.66 (0.33-1.2)	1.2% (0.19-2.7%)	1.5% (0.29-3.6%)	24 (4.4-55)	350 (58-830)
Puerto Rico	1995	Female	80 to 84	20% (12-31%)	80% (69-88%)	0.6 (0.28-1.1)	1% (0.14-2.4%)	1.3% (0.19-3%)	23 (3.4-55)	260 (37-610)
Puerto Rico	1995	Female	85 to 89	20% (12-30%)	80% (70-88%)	0.59 (0.27-1.1)	0.86% (-0.0034-2.2%)	1% (-0.089-2.6%)	16 (-1.5-41)	140 (-0.54-330)
Puerto Rico	1995	Female	90 to 94	21% (13-33%)	79% (67-87%)	0.66 (0.32-1.2)	0.88% (-0.16-2.5%)	0.93% (-0.31-2.9%)	10 (-3.5-32)	61 (-11-180)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	1995	Female	95 plus	25% (14-40%)	75% (60-86%)	0.77 (0.31-1.5)	0.85% (-0.52-2.9%)	0.82% (-0.8-3.2%)	3.4 (-3.3-13)	14 (-8.5-48)
Puerto Rico	1995	Male	All Ages	57% (50-65%)	43% (35-50%)	3.1 (2-4.6)	10% (7.5-13%)	10% (6.8-13%)	1800 (1200-2400)	68000 (49000-87000)
Puerto Rico	1995	Male	15 to 19	58% (40-74%)	42% (26-60%)	2.7 (1.6-4.5)	10% (6.3-15%)	14% (8.2-22%)	42 (24-67)	3300 (2000-5100)
Puerto Rico	1995	Male	20 to 24	69% (53-82%)	31% (18-47%)	5.3 (3.2-8)	19% (12-26%)	23% (15-33%)	100 (65-150)	7400 (5000-10000)
Puerto Rico	1995	Male	25 to 29	72% (57-84%)	28% (16-43%)	5.3 (3.3-8.2)	16% (11-21%)	19% (13-27%)	84 (54-120)	5900 (4000-8100)
Puerto Rico	1995	Male	30 to 34	72% (57-84%)	28% (16-43%)	5.3 (3.2-8.2)	12% (8.5-16%)	13% (9.1-18%)	77 (51-110)	5200 (3600-6900)
Puerto Rico	1995	Male	35 to 39	71% (58-84%)	29% (16-42%)	5.4 (3.3-8.3)	13% (9.5-15%)	14% (11-18%)	97 (69-130)	5800 (4300-7400)
Puerto Rico	1995	Male	40 to 44	70% (55-82%)	30% (18-45%)	5.1 (3-7.9)	14% (11-17%)	17% (13-20%)	120 (92-150)	6400 (4900-8000)
Puerto Rico	1995	Male	45 to 49	68% (52-81%)	32% (19-48%)	4.4 (2.6-6.8)	15% (12-18%)	18% (14-22%)	140 (110-180)	6800 (5300-8400)
Puerto Rico	1995	Male	50 to 54	68% (52-82%)	32% (18-48%)	4 (2.4-6.3)	15% (11-18%)	18% (14-23%)	160 (120-210)	6600 (4900-8300)
Puerto Rico	1995	Male	55 to 59	65% (50-80%)	35% (20-50%)	3.6 (2.2-5.9)	14% (9.7-18%)	17% (12-21%)	170 (120-220)	5800 (4100-7500)
Puerto Rico	1995	Male	60 to 64	61% (42-78%)	39% (22-58%)	3.1 (1.8-5.1)	11% (7.4-15%)	13% (8.8-18%)	170 (110-230)	4900 (3200-6700)
Puerto Rico	1995	Male	65 to 69	58% (39-76%)	42% (24-61%)	2.6 (1.4-4.3)	8.9% (5.5-13%)	11% (6.6-15%)	160 (98-230)	3800 (2400-5500)
Puerto Rico	1995	Male	70 to 74	54% (36-71%)	46% (29-64%)	2.1 (1.1-3.8)	6.6% (3.7-10%)	7.7% (4.3-12%)	150 (81-240)	2800 (1600-4600)
Puerto Rico	1995	Male	75 to 79	49% (31-68%)	51% (32-69%)	1.8 (0.94-3.1)	5.1% (2.7-8.5%)	5.9% (3.1-9.9%)	120 (59-190)	1700 (910-2900)
Puerto Rico	1995	Male	80 to 84	44% (26-61%)	56% (39-74%)	1.6 (0.77-2.7)	4.2% (2.1-7.1%)	4.8% (2.3-8.2%)	95 (46-160)	1100 (530-1800)
Puerto Rico	1995	Male	85 to 89	39% (24-55%)	61% (45-76%)	1.4 (0.69-2.5)	3.4% (1.6-6%)	3.8% (1.7-6.9%)	59 (26-110)	480 (220-850)
Puerto Rico	1995	Male	90 to 94	40% (25-56%)	60% (44-75%)	1.5 (0.76-2.7)	3.1% (1.2-5.9%)	3.4% (1.1-6.7%)	28 (9.3-55)	160 (59-300)
Puerto Rico	1995	Male	95 plus	43% (24-63%)	57% (37-76%)	1.7 (0.76-3)	3.3% (1-6.6%)	3.5% (0.8-7.4%)	8.1 (1.8-17)	32 (9.7-61)
Puerto Rico	2000	Female	All Ages	28% (23-34%)	72% (66-77%)	1 (0.63-1.5)	2% (1.2-3.2%)	2.2% (1-3.8%)	270 (120-470)	9400 (5700-15000)
Puerto Rico	2000	Female	15 to 19	38% (26-54%)	62% (46-74%)	1.5 (0.85-2.5)	2.4% (1.5-3.6%)	6.3% (3.6-11%)	3.3 (1.8-5.4)	380 (240-570)
Puerto Rico	2000	Female	20 to 24	41% (27-55%)	59% (45-73%)	2 (1.2-3.2)	3.7% (2.5-5.3%)	7.2% (4.2-12%)	5.2 (3-8.4)	660 (440-940)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	2000	Female	25 to 29	37% (25-53%)	63% (47-75%)	1.7 (0.94-2.7)	3.4% (2.4-4.7%)	5.7% (3.4-8.6%)	5.1 (3-7.9)	620 (430-870)
Puerto Rico	2000	Female	30 to 34	34% (22-49%)	66% (51-78%)	1.5 (0.84-2.4)	3.1% (2.1-4.4%)	4.9% (3.2-7.6%)	6.7 (4.2-10)	650 (440-960)
Puerto Rico	2000	Female	35 to 39	35% (24-50%)	65% (50-76%)	1.5 (0.84-2.4)	3.1% (2.1-4.4%)	5.3% (3.5-7.9%)	9.7 (6.4-14)	760 (520-1100)
Puerto Rico	2000	Female	40 to 44	34% (23-50%)	66% (50-77%)	1.4 (0.8-2.3)	3.3% (2.1-4.9%)	6.3% (4.2-9%)	14 (9.4-21)	890 (580-1300)
Puerto Rico	2000	Female	45 to 49	32% (21-47%)	68% (53-79%)	1.3 (0.71-2)	3.2% (1.9-4.9%)	5.8% (3.7-8.7%)	18 (11-27)	920 (570-1400)
Puerto Rico	2000	Female	50 to 54	28% (18-41%)	72% (59-82%)	1 (0.59-1.7)	2.8% (1.7-4.6%)	5% (3.1-7.5%)	21 (14-33)	920 (560-1500)
Puerto Rico	2000	Female	55 to 59	23% (15-36%)	77% (64-85%)	0.84 (0.44-1.5)	2.4% (1.3-4.1%)	3.9% (2.3-6.3%)	22 (13-35)	780 (440-1300)
Puerto Rico	2000	Female	60 to 64	22% (13-35%)	78% (65-87%)	0.76 (0.4-1.3)	2.1% (1-3.8%)	3.2% (1.7-5.5%)	23 (12-40)	700 (340-1200)
Puerto Rico	2000	Female	65 to 69	22% (14-34%)	78% (66-86%)	0.75 (0.39-1.3)	1.9% (0.85-3.4%)	2.7% (1.4-4.8%)	28 (14-49)	660 (300-1200)
Puerto Rico	2000	Female	70 to 74	21% (13-33%)	79% (67-87%)	0.67 (0.35-1.2)	1.5% (0.51-3%)	2.1% (0.79-4%)	27 (10-51)	520 (180-1000)
Puerto Rico	2000	Female	75 to 79	18% (11-29%)	82% (71-89%)	0.59 (0.29-1.1)	1.2% (0.33-2.6%)	1.6% (0.51-3.4%)	27 (8.6-56)	400 (110-860)
Puerto Rico	2000	Female	80 to 84	17% (10-28%)	83% (72-90%)	0.54 (0.27-0.95)	1% (0.24-2.2%)	1.3% (0.33-2.8%)	25 (6-52)	280 (66-590)
Puerto Rico	2000	Female	85 to 89	17% (10-26%)	83% (74-90%)	0.53 (0.24-0.97)	0.91% (0.13-2.1%)	1.1% (0.13-2.5%)	20 (2.3-45)	160 (24-370)
Puerto Rico	2000	Female	90 to 94	19% (11-29%)	81% (71-89%)	0.58 (0.29-1.1)	0.91% (0.026-2.3%)	1% (-0.08-2.7%)	12 (-0.95-32)	71 (2.1-180)
Puerto Rico	2000	Female	95 plus	21% (12-36%)	79% (64-88%)	0.68 (0.29-1.3)	0.93% (-0.17-2.6%)	0.95% (-0.44-3.1%)	4.3 (-2-14)	18 (-3.2-50)
Puerto Rico	2000	Male	All Ages	59% (52-67%)	41% (33-48%)	3.3 (2.2-4.9)	11% (7.9-14%)	11% (7.2-14%)	1700 (1200-2300)	64000 (47000-83000)
Puerto Rico	2000	Male	15 to 19	57% (42-73%)	43% (27-58%)	2.7 (1.6-4.3)	9.5% (6-14%)	14% (8.4-22%)	31 (18-49)	2500 (1600-3800)
Puerto Rico	2000	Male	20 to 24	71% (57-83%)	29% (17-43%)	5.6 (3.4-8.5)	18% (13-25%)	23% (15-34%)	89 (59-130)	6600 (4600-9300)
Puerto Rico	2000	Male	25 to 29	74% (61-84%)	26% (16-39%)	5.6 (3.5-8.4)	16% (12-22%)	21% (14-29%)	72 (48-100)	5200 (3800-7100)
Puerto Rico	2000	Male	30 to 34	72% (59-84%)	28% (16-41%)	5.4 (3.4-8.2)	14% (10-18%)	17% (12-24%)	64 (45-88)	4500 (3200-5900)
Puerto Rico	2000	Male	35 to 39	73% (61-84%)	27% (16-39%)	5.8 (3.6-8.8)	14% (10-17%)	17% (13-22%)	84 (61-110)	5200 (3900-6500)
Puerto Rico	2000	Male	40 to 44	72% (59-83%)	28% (17-41%)	5.6 (3.3-8.6)	14% (11-18%)	18% (14-22%)	100 (79-130)	5600 (4400-6900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	2000	Male	45 to 49	71% (55-82%)	29% (18-45%)	4.8 (2.9-7.2)	15% (12-19%)	19% (15-23%)	130 (99-160)	6200 (4800-7700)
Puerto Rico	2000	Male	50 to 54	70% (55-83%)	30% (17-45%)	4.3 (2.5-6.7)	15% (11-19%)	19% (14-24%)	170 (120-210)	6800 (5000-8600)
Puerto Rico	2000	Male	55 to 59	69% (53-81%)	31% (19-47%)	4.1 (2.5-6.4)	14% (9.9-18%)	17% (12-23%)	180 (130-230)	6200 (4500-8100)
Puerto Rico	2000	Male	60 to 64	64% (48-79%)	36% (21-52%)	3.5 (2.1-5.6)	11% (7.7-15%)	14% (9.5-19%)	170 (120-230)	5100 (3500-6900)
Puerto Rico	2000	Male	65 to 69	60% (42-76%)	40% (24-58%)	2.8 (1.6-4.6)	9.1% (5.7-13%)	11% (7-16%)	160 (100-230)	4000 (2500-5600)
Puerto Rico	2000	Male	70 to 74	56% (38-72%)	44% (28-62%)	2.3 (1.3-4)	6.7% (3.7-11%)	8.1% (4.4-13%)	140 (75-220)	2700 (1500-4300)
Puerto Rico	2000	Male	75 to 79	51% (32-70%)	49% (30-68%)	1.9 (1-3.3)	5.3% (2.7-8.6%)	6.2% (3-10%)	120 (60-200)	1800 (930-3000)
Puerto Rico	2000	Male	80 to 84	45% (28-63%)	55% (37-72%)	1.7 (0.81-2.8)	4.3% (2.1-7.1%)	5% (2.5-8.3%)	95 (46-160)	1100 (540-1800)
Puerto Rico	2000	Male	85 to 89	41% (24-57%)	59% (43-76%)	1.5 (0.74-2.6)	3.7% (1.7-6.3%)	4.2% (1.9-7.3%)	67 (30-120)	540 (260-930)
Puerto Rico	2000	Male	90 to 94	41% (26-59%)	59% (41-74%)	1.6 (0.81-2.9)	3.3% (1.3-6.1%)	3.7% (1.4-7%)	32 (12-60)	180 (74-340)
Puerto Rico	2000	Male	95 plus	44% (25-65%)	56% (35-75%)	1.8 (0.81-3.2)	3.6% (1.2-6.7%)	3.9% (1-7.7%)	10 (2.8-20)	40 (14-76)
Puerto Rico	2005	Female	All Ages	29% (25-34%)	71% (66-75%)	1.1 (0.69-1.7)	2.1% (1.3-3.3%)	2.2% (1.2-3.7%)	280 (150-480)	9500 (5700-15000)
Puerto Rico	2005	Female	15 to 19	38% (26-54%)	62% (46-74%)	1.5 (0.88-2.6)	2.1% (1.4-3.2%)	6.4% (3.7-11%)	2.4 (1.4-4)	310 (200-470)
Puerto Rico	2005	Female	20 to 24	42% (30-56%)	58% (44-70%)	2.2 (1.3-3.5)	3.8% (2.6-5.5%)	7.7% (4.5-12%)	5.2 (3-8.3)	650 (430-930)
Puerto Rico	2005	Female	25 to 29	40% (29-56%)	60% (44-71%)	1.9 (1.1-3)	3.5% (2.4-5%)	6.5% (3.8-10%)	4.9 (2.9-7.6)	600 (400-880)
Puerto Rico	2005	Female	30 to 34	37% (26-52%)	63% (48-74%)	1.7 (1-2.7)	3.1% (2.1-4.3%)	5.5% (3.5-8.4%)	5.5 (3.4-8.3)	580 (390-850)
Puerto Rico	2005	Female	35 to 39	36% (26-51%)	64% (49-74%)	1.6 (0.94-2.7)	3% (2-4.3%)	5.5% (3.7-8.2%)	8.3 (5.4-12)	680 (460-1000)
Puerto Rico	2005	Female	40 to 44	35% (25-51%)	65% (49-75%)	1.5 (0.87-2.5)	3.1% (2-4.8%)	6.1% (4-9.1%)	13 (8.3-19)	810 (530-1200)
Puerto Rico	2005	Female	45 to 49	34% (24-47%)	66% (53-76%)	1.4 (0.8-2.3)	3.2% (2-5.1%)	6% (3.9-9.1%)	18 (11-27)	920 (560-1400)
Puerto Rico	2005	Female	50 to 54	30% (21-43%)	70% (57-79%)	1.2 (0.69-1.9)	2.9% (1.7-4.7%)	5.2% (3.2-8%)	21 (13-33)	920 (520-1500)
Puerto Rico	2005	Female	55 to 59	25% (17-37%)	75% (63-83%)	0.96 (0.54-1.7)	2.6% (1.4-4.5%)	4.4% (2.6-7%)	25 (15-42)	910 (510-1600)
Puerto Rico	2005	Female	60 to 64	22% (15-35%)	78% (65-85%)	0.82 (0.46-1.4)	2.2% (1.1-4.2%)	3.4% (1.8-6%)	27 (14-48)	810 (400-1500)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	2005	Female	65 to 69	21% (14-32%)	79% (68-86%)	0.73 (0.4-1.2)	1.9% (0.94-3.5%)	2.9% (1.6-5%)	29 (15-50)	690 (340-1300)
Puerto Rico	2005	Female	70 to 74	18% (12-26%)	82% (74-88%)	0.59 (0.31-1)	1.6% (0.66-2.9%)	2.2% (1.1-4%)	29 (14-52)	560 (250-1000)
Puerto Rico	2005	Female	75 to 79	14% (9.1-22%)	86% (78-91%)	0.46 (0.24-0.82)	1.2% (0.54-2.3%)	1.7% (0.78-3.1%)	27 (13-50)	410 (180-780)
Puerto Rico	2005	Female	80 to 84	13% (7.6-21%)	87% (79-92%)	0.42 (0.21-0.75)	1% (0.41-2.1%)	1.3% (0.56-2.6%)	27 (11-54)	310 (120-600)
Puerto Rico	2005	Female	85 to 89	13% (7.8-20%)	87% (80-92%)	0.42 (0.2-0.76)	0.88% (0.3-1.9%)	1.1% (0.35-2.3%)	21 (6.7-45)	170 (60-360)
Puerto Rico	2005	Female	90 to 94	14% (8.5-21%)	86% (79-91%)	0.45 (0.22-0.81)	0.89% (0.19-2%)	1% (0.17-2.4%)	14 (2.3-34)	84 (18-190)
Puerto Rico	2005	Female	95 plus	16% (8.8-28%)	84% (72-91%)	0.52 (0.22-0.96)	0.88% (0.04-2.3%)	0.92% (-0.11-2.6%)	5.5 (-0.69-16)	23 (1-58)
Puerto Rico	2005	Male	All Ages	59% (52-66%)	41% (34-48%)	3.4 (2.2-5.1)	10% (7.5-13%)	10% (6.7-14%)	1600 (1100-2200)	59000 (43000-76000)
Puerto Rico	2005	Male	15 to 19	57% (41-72%)	43% (28-59%)	2.8 (1.6-4.3)	8.9% (5.6-13%)	14% (8.4-22%)	24 (14-36)	2000 (1300-2900)
Puerto Rico	2005	Male	20 to 24	71% (56-82%)	29% (18-44%)	5.7 (3.5-8.7)	18% (13-26%)	24% (16-35%)	86 (57-120)	6400 (4400-9000)
Puerto Rico	2005	Male	25 to 29	73% (60-84%)	27% (16-40%)	5.7 (3.5-8.6)	17% (12-23%)	22% (15-32%)	75 (50-110)	5400 (3800-7400)
Puerto Rico	2005	Male	30 to 34	71% (58-83%)	29% (17-42%)	5.4 (3.4-8.2)	14% (10-19%)	19% (13-26%)	59 (40-83)	4100 (3000-5500)
Puerto Rico	2005	Male	35 to 39	73% (60-84%)	27% (16-40%)	5.9 (3.6-9.1)	13% (10-17%)	18% (13-24%)	66 (49-88)	4200 (3200-5300)
Puerto Rico	2005	Male	40 to 44	72% (60-83%)	28% (17-40%)	5.7 (3.4-8.7)	14% (11-17%)	18% (14-23%)	83 (62-100)	4600 (3500-5700)
Puerto Rico	2005	Male	45 to 49	69% (54-81%)	31% (19-46%)	4.7 (2.9-7.3)	14% (10-17%)	18% (13-22%)	110 (80-130)	5200 (3900-6400)
Puerto Rico	2005	Male	50 to 54	68% (52-81%)	32% (19-48%)	4.2 (2.5-6.6)	14% (10-17%)	18% (13-22%)	140 (100-180)	5800 (4400-7400)
Puerto Rico	2005	Male	55 to 59	67% (51-80%)	33% (20-49%)	4 (2.4-6.3)	13% (9.1-17%)	16% (12-21%)	170 (120-220)	6100 (4300-7900)
Puerto Rico	2005	Male	60 to 64	63% (47-78%)	37% (22-53%)	3.5 (2.1-5.6)	11% (7.1-15%)	13% (8.8-18%)	170 (110-230)	5100 (3400-6900)
Puerto Rico	2005	Male	65 to 69	60% (43-75%)	40% (25-57%)	2.9 (1.7-4.7)	8.8% (5.5-12%)	11% (6.7-15%)	160 (99-220)	3900 (2500-5600)
Puerto Rico	2005	Male	70 to 74	58% (42-73%)	42% (27-58%)	2.5 (1.4-4.1)	6.8% (3.7-10%)	8.2% (4.4-12%)	140 (73-210)	2700 (1500-4200)
Puerto Rico	2005	Male	75 to 79	52% (35-69%)	48% (31-65%)	2 (1.1-3.3)	5.3% (2.6-8.7%)	6.2% (3-10%)	120 (57-190)	1800 (890-2900)
Puerto Rico	2005	Male	80 to 84	44% (27-62%)	56% (38-73%)	1.7 (0.87-2.8)	4.3% (2.1-7.2%)	5% (2.4-8.4%)	98 (48-160)	1100 (550-1800)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	2005	Male	85 to 89	39% (24-55%)	61% (45-76%)	1.5 (0.75-2.6)	3.6% (1.7-6.2%)	4.2% (1.9-7.2%)	67 (30-120)	550 (260-950)
Puerto Rico	2005	Male	90 to 94	40% (26-56%)	60% (44-74%)	1.6 (0.83-2.7)	3.5% (1.4-6.3%)	3.9% (1.5-7.2%)	40 (15-74)	220 (92-400)
Puerto Rico	2005	Male	95 plus	43% (25-63%)	57% (37-75%)	1.8 (0.8-3.2)	3.7% (1.3-7.1%)	4.1% (1.2-8.1%)	14 (4.2-28)	55 (19-100)
Puerto Rico	2010	Female	All Ages	30% (26-35%)	70% (65-74%)	1.2 (0.72-1.8)	2% (1.3-3.3%)	2.1% (1.2-3.6%)	270 (150-470)	9200 (5600-14000)
Puerto Rico	2010	Female	15 to 19	38% (27-54%)	62% (46-73%)	1.6 (0.86-2.6)	2% (1.3-3%)	6.5% (3.8-11%)	1.9 (1.1-3.2)	280 (180-440)
Puerto Rico	2010	Female	20 to 24	45% (33-58%)	55% (42-67%)	2.4 (1.4-3.8)	3.8% (2.6-5.5%)	8.3% (4.8-13%)	4.3 (2.4-6.8)	580 (380-850)
Puerto Rico	2010	Female	25 to 29	45% (34-60%)	55% (40-66%)	2.2 (1.3-3.5)	3.7% (2.5-5.3%)	7.5% (4.5-12%)	4.6 (2.6-7.1)	580 (380-840)
Puerto Rico	2010	Female	30 to 34	42% (31-57%)	58% (43-69%)	2 (1.2-3.2)	3.3% (2.2-4.8%)	6.4% (4-9.6%)	5.6 (3.4-8.5)	600 (400-880)
Puerto Rico	2010	Female	35 to 39	38% (28-53%)	62% (47-72%)	1.7 (1-2.9)	3.1% (2.1-4.5%)	6.2% (4-9.2%)	7.9 (5-12)	650 (430-970)
Puerto Rico	2010	Female	40 to 44	35% (25-51%)	65% (49-75%)	1.5 (0.89-2.5)	3% (1.9-4.7%)	6.2% (4-9.3%)	11 (7.4-17)	750 (470-1100)
Puerto Rico	2010	Female	45 to 49	35% (25-48%)	65% (52-75%)	1.5 (0.85-2.4)	3.1% (1.8-5.1%)	6% (3.8-9.2%)	17 (10-26)	880 (510-1400)
Puerto Rico	2010	Female	50 to 54	31% (22-43%)	69% (57-78%)	1.2 (0.68-2)	2.9% (1.6-4.9%)	5.2% (3.1-8.3%)	21 (12-33)	920 (510-1500)
Puerto Rico	2010	Female	55 to 59	26% (18-37%)	74% (63-82%)	0.98 (0.55-1.7)	2.6% (1.4-4.5%)	4.4% (2.6-7.3%)	24 (14-41)	890 (490-1500)
Puerto Rico	2010	Female	60 to 64	22% (15-34%)	78% (66-85%)	0.8 (0.43-1.4)	2.1% (1-3.8%)	3.4% (1.8-5.7%)	27 (14-46)	820 (410-1500)
Puerto Rico	2010	Female	65 to 69	18% (13-28%)	82% (72-87%)	0.64 (0.35-1.1)	1.9% (0.99-3.3%)	2.8% (1.6-4.7%)	28 (16-47)	690 (370-1200)
Puerto Rico	2010	Female	70 to 74	16% (11-24%)	84% (76-89%)	0.53 (0.28-0.91)	1.5% (0.67-2.7%)	2.2% (1.1-3.7%)	27 (13-47)	520 (240-940)
Puerto Rico	2010	Female	75 to 79	13% (8.4-19%)	87% (81-92%)	0.42 (0.21-0.76)	1.2% (0.55-2.2%)	1.6% (0.84-3%)	27 (13-48)	400 (180-750)
Puerto Rico	2010	Female	80 to 84	11% (6.5-18%)	89% (82-93%)	0.36 (0.18-0.68)	0.95% (0.42-1.9%)	1.3% (0.6-2.4%)	25 (12-48)	280 (120-550)
Puerto Rico	2010	Female	85 to 89	11% (6.6-17%)	89% (83-93%)	0.36 (0.17-0.64)	0.84% (0.35-1.7%)	1% (0.41-2.1%)	22 (8.8-44)	180 (75-360)
Puerto Rico	2010	Female	90 to 94	12% (7.4-18%)	88% (82-93%)	0.39 (0.19-0.67)	0.84% (0.28-1.8%)	0.96% (0.26-2.1%)	15 (4.2-34)	91 (30-200)
Puerto Rico	2010	Female	95 plus	14% (7.6-24%)	86% (76-92%)	0.44 (0.19-0.84)	0.85% (0.19-2%)	0.88% (0.04-2.4%)	6.1 (0.28-17)	26 (5.9-63)
Puerto Rico	2010	Male	All Ages	59% (52-67%)	41% (33-48%)	3.5 (2.2-5.2)	10% (7.2-13%)	9.9% (6.5-13%)	1500 (1000-2100)	57000 (40000-74000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	2010	Male	15 to 19	56% (40-71%)	44% (29-60%)	2.7 (1.6-4.4)	8.8% (5.6-14%)	14% (8.4-23%)	22 (13-35)	1900 (1200-2800)
Puerto Rico	2010	Male	20 to 24	71% (56-82%)	29% (18-44%)	5.7 (3.4-8.7)	19% (13-27%)	25% (16-37%)	82 (53-120)	6100 (4100-8700)
Puerto Rico	2010	Male	25 to 29	75% (62-85%)	25% (15-38%)	5.9 (3.6-8.8)	18% (13-25%)	24% (16-35%)	81 (54-120)	5600 (3900-7900)
Puerto Rico	2010	Male	30 to 34	74% (61-85%)	26% (15-39%)	5.7 (3.5-8.7)	15% (11-21%)	21% (15-30%)	67 (46-93)	4500 (3200-6100)
Puerto Rico	2010	Male	35 to 39	72% (59-84%)	28% (16-41%)	5.7 (3.6-8.9)	13% (10-17%)	20% (14-26%)	57 (42-77)	3700 (2700-4800)
Puerto Rico	2010	Male	40 to 44	71% (58-82%)	29% (18-42%)	5.5 (3.4-8.6)	13% (10-17%)	19% (14-24%)	72 (53-91)	4100 (3100-5100)
Puerto Rico	2010	Male	45 to 49	69% (53-81%)	31% (19-47%)	4.7 (2.8-7.3)	13% (9.5-17%)	17% (13-22%)	91 (66-120)	4500 (3300-5700)
Puerto Rico	2010	Male	50 to 54	66% (50-79%)	34% (21-50%)	4 (2.4-6.3)	13% (9.1-16%)	17% (12-21%)	120 (88-160)	5300 (3700-6800)
Puerto Rico	2010	Male	55 to 59	65% (49-78%)	35% (22-51%)	3.8 (2.3-6)	12% (8.3-16%)	16% (11-20%)	150 (100-200)	5400 (3700-7000)
Puerto Rico	2010	Male	60 to 64	62% (45-77%)	38% (23-55%)	3.4 (1.9-5.4)	10% (6.8-14%)	13% (8.6-17%)	170 (110-230)	5100 (3400-6900)
Puerto Rico	2010	Male	65 to 69	61% (45-76%)	39% (24-55%)	3 (1.7-4.8)	8.8% (5.4-12%)	11% (6.8-15%)	170 (100-230)	4100 (2500-5800)
Puerto Rico	2010	Male	70 to 74	58% (42-72%)	42% (28-58%)	2.5 (1.4-4)	6.7% (3.6-10%)	8.1% (4.4-12%)	140 (74-210)	2700 (1500-4100)
Puerto Rico	2010	Male	75 to 79	51% (34-68%)	49% (32-66%)	2 (1.1-3.3)	5.1% (2.5-8.3%)	6% (3-9.7%)	110 (56-180)	1800 (880-2800)
Puerto Rico	2010	Male	80 to 84	43% (26-60%)	57% (40-74%)	1.6 (0.87-2.9)	4.1% (2.1-7%)	4.8% (2.4-8.3%)	93 (46-160)	1100 (540-1800)
Puerto Rico	2010	Male	85 to 89	38% (24-53%)	62% (47-76%)	1.5 (0.73-2.5)	3.5% (1.7-6%)	4.1% (1.8-7.1%)	70 (32-120)	570 (270-980)
Puerto Rico	2010	Male	90 to 94	39% (25-55%)	61% (45-75%)	1.5 (0.77-2.7)	3.4% (1.4-6%)	3.8% (1.4-7%)	40 (15-73)	230 (96-400)
Puerto Rico	2010	Male	95 plus	42% (24-62%)	58% (38-76%)	1.7 (0.77-3.2)	3.6% (1.3-6.6%)	4% (1.2-7.7%)	14 (4.3-27)	54 (20-98)
Puerto Rico	2016	Female	All Ages	30% (26-36%)	70% (64-74%)	1.2 (0.71-1.9)	2.1% (1.2-3.5%)	2.3% (1.1-4.1%)	320 (150-570)	9700 (5400-15000)
Puerto Rico	2016	Female	15 to 19	39% (27-55%)	61% (45-73%)	1.7 (0.85-2.8)	2.1% (1.3-3.2%)	6.9% (3.6-11%)	1.7 (0.89-2.9)	250 (150-380)
Puerto Rico	2016	Female	20 to 24	45% (32-59%)	55% (41-68%)	2.5 (1.4-4.2)	3.9% (2.5-5.4%)	8.9% (4.8-14%)	4.2 (2.2-6.7)	570 (360-820)
Puerto Rico	2016	Female	25 to 29	45% (32-61%)	55% (39-68%)	2.3 (1.2-3.8)	3.7% (2.4-5.2%)	7.7% (4.2-12%)	4.3 (2.2-7.1)	560 (350-810)
Puerto Rico	2016	Female	30 to 34	43% (30-59%)	57% (41-70%)	2.1 (1.1-3.6)	3.3% (2.1-4.9%)	7.1% (4.2-11%)	5.4 (3.1-8.5)	580 (370-860)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	2016	Female	35 to 39	39% (28-56%)	61% (44-72%)	1.9 (1-3.2)	3.2% (2.1-4.9%)	6.8% (4.2-10%)	8 (4.8-12)	670 (420-990)
Puerto Rico	2016	Female	40 to 44	36% (24-52%)	64% (48-76%)	1.6 (0.89-2.7)	3.1% (1.9-4.8%)	6.6% (4.1-9.7%)	11 (6.4-17)	730 (430-1100)
Puerto Rico	2016	Female	45 to 49	34% (23-49%)	66% (51-77%)	1.5 (0.81-2.6)	3.2% (1.9-5.3%)	6.4% (3.9-10%)	17 (9.9-26)	880 (510-1400)
Puerto Rico	2016	Female	50 to 54	29% (19-43%)	71% (57-81%)	1.2 (0.62-2.1)	3% (1.6-5.1%)	5.5% (3.2-8.9%)	21 (12-35)	940 (510-1600)
Puerto Rico	2016	Female	55 to 59	24% (16-36%)	76% (64-84%)	0.97 (0.47-1.7)	2.7% (1.4-4.7%)	4.6% (2.6-7.7%)	25 (14-42)	910 (480-1600)
Puerto Rico	2016	Female	60 to 64	21% (14-33%)	79% (67-86%)	0.8 (0.39-1.4)	2.2% (1.1-4.1%)	3.6% (1.9-6.3%)	27 (15-48)	840 (420-1500)
Puerto Rico	2016	Female	65 to 69	18% (11-28%)	82% (72-89%)	0.64 (0.32-1.1)	2% (0.98-3.5%)	3% (1.6-5.1%)	33 (18-56)	830 (410-1400)
Puerto Rico	2016	Female	70 to 74	16% (9.5-26%)	84% (74-91%)	0.56 (0.27-1)	1.6% (0.76-3.1%)	2.4% (1.2-4.3%)	33 (17-61)	640 (310-1200)
Puerto Rico	2016	Female	75 to 79	15% (8.8-24%)	85% (76-91%)	0.5 (0.23-1)	1.4% (0.55-2.7%)	1.9% (0.83-3.6%)	32 (14-65)	490 (190-1000)
Puerto Rico	2016	Female	80 to 84	14% (7.7-23%)	86% (77-92%)	0.46 (0.22-0.88)	1.1% (0.41-2.4%)	1.5% (0.56-3.1%)	32 (12-66)	360 (130-790)
Puerto Rico	2016	Female	85 to 89	14% (7.6-21%)	86% (79-92%)	0.46 (0.21-0.87)	0.99% (0.32-2.2%)	1.2% (0.37-2.8%)	28 (8.7-64)	230 (74-510)
Puerto Rico	2016	Female	90 to 94	15% (8.6-23%)	85% (77-91%)	0.5 (0.23-0.94)	1% (0.26-2.5%)	1.2% (0.24-3%)	23 (4.7-56)	130 (34-320)
Puerto Rico	2016	Female	95 plus	17% (9.2-30%)	83% (70-91%)	0.57 (0.22-1.1)	1.1% (0.17-2.7%)	1.1% (0.004-3.3%)	10 (0.039-29)	42 (7-110)
Puerto Rico	2016	Male	All Ages	59% (52-66%)	41% (34-48%)	3.6 (2.1-5.7)	10% (7-13%)	9.8% (6.3-13%)	1600 (960-2200)	55000 (37000-73000)
Puerto Rico	2016	Male	15 to 19	56% (39-72%)	44% (28-61%)	2.8 (1.5-4.6)	8.4% (5.2-13%)	15% (8.1-23%)	18 (9.6-29)	1500 (940-2300)
Puerto Rico	2016	Male	20 to 24	70% (55-82%)	30% (18-45%)	5.8 (3.4-9.5)	18% (12-26%)	25% (15-36%)	73 (45-110)	5500 (3600-7900)
Puerto Rico	2016	Male	25 to 29	75% (63-86%)	25% (14-37%)	6.2 (3.5-9.9)	18% (12-24%)	24% (15-34%)	68 (43-98)	4900 (3200-6700)
Puerto Rico	2016	Male	30 to 34	75% (62-86%)	25% (14-38%)	6.1 (3.4-9.8)	15% (11-20%)	22% (14-30%)	57 (38-81)	4000 (2700-5400)
Puerto Rico	2016	Male	35 to 39	70% (56-82%)	30% (18-44%)	5.7 (3.2-9.3)	13% (9.8-17%)	19% (14-26%)	56 (38-75)	3600 (2600-4700)
Puerto Rico	2016	Male	40 to 44	69% (55-81%)	31% (19-45%)	5.4 (3.1-8.9)	13% (9.9-17%)	19% (14-24%)	67 (47-88)	3800 (2800-4800)
Puerto Rico	2016	Male	45 to 49	69% (52-81%)	31% (19-48%)	4.8 (2.7-7.8)	13% (9.9-17%)	18% (13-23%)	91 (65-120)	4500 (3200-5800)
Puerto Rico	2016	Male	50 to 54	65% (49-78%)	35% (22-51%)	4 (2.2-6.7)	13% (9-17%)	17% (12-22%)	120 (84-160)	5100 (3500-6600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Puerto Rico	2016	Male	55 to 59	63% (47-77%)	37% (23-53%)	3.8 (2-6.3)	12% (8.4-16%)	16% (11-20%)	160 (110-210)	5600 (3800-7300)
Puerto Rico	2016	Male	60 to 64	60% (44-76%)	40% (24-56%)	3.4 (1.8-5.7)	10% (6.7-14%)	13% (8.6-18%)	160 (100-220)	4900 (3200-6700)
Puerto Rico	2016	Male	65 to 69	61% (45-76%)	39% (24-55%)	3.1 (1.7-5.1)	9.2% (5.4-13%)	11% (7-16%)	190 (110-270)	4700 (2700-6600)
Puerto Rico	2016	Male	70 to 74	58% (41-73%)	42% (27-59%)	2.6 (1.3-4.5)	6.9% (3.7-11%)	8.4% (4.5-13%)	160 (82-240)	3100 (1600-4700)
Puerto Rico	2016	Male	75 to 79	49% (30-67%)	51% (33-70%)	1.9 (0.94-3.5)	5.1% (2.5-8.5%)	6.1% (3-10%)	120 (57-210)	1900 (890-3100)
Puerto Rico	2016	Male	80 to 84	43% (26-61%)	57% (39-74%)	1.7 (0.81-3.1)	4.2% (2-7.1%)	5% (2.4-8.4%)	100 (47-170)	1200 (550-1900)
Puerto Rico	2016	Male	85 to 89	38% (24-54%)	62% (46-76%)	1.5 (0.69-2.7)	3.7% (1.7-6.3%)	4.3% (1.9-7.4%)	76 (34-130)	620 (290-1100)
Puerto Rico	2016	Male	90 to 94	38% (24-55%)	62% (45-76%)	1.6 (0.73-2.9)	3.5% (1.4-6.2%)	4% (1.5-7.2%)	47 (18-85)	270 (110-470)
Puerto Rico	2016	Male	95 plus	41% (24-62%)	59% (38-76%)	1.8 (0.71-3.4)	3.7% (1.3-6.7%)	4.1% (1.2-7.9%)	18 (5.3-35)	70 (25-130)
Virgin Islands, U.S.	1990	Female	All Ages	47% (41-54%)	53% (46-59%)	1.5 (0.95-2.2)	2.9% (1.6-4.7%)	3.4% (1.2-6.5%)	9.1 (3.3-17)	360 (190-590)
Virgin Islands, U.S.	1990	Female	15 to 19	54% (40-69%)	46% (31-60%)	1.9 (1.1-3)	3.2% (2.1-4.7%)	7.6% (4.6-11%)	0.18 (0.11-0.27)	18 (12-27)
Virgin Islands, U.S.	1990	Female	20 to 24	58% (45-71%)	42% (29-55%)	2.5 (1.6-3.9)	4.8% (3.4-6.5%)	9.6% (6.3-14%)	0.21 (0.13-0.3)	22 (15-31)
Virgin Islands, U.S.	1990	Female	25 to 29	57% (44-72%)	43% (28-56%)	2.4 (1.4-3.7)	4.6% (3.3-6.3%)	8.7% (5.8-12%)	0.24 (0.15-0.35)	23 (17-33)
Virgin Islands, U.S.	1990	Female	30 to 34	56% (42-71%)	44% (29-58%)	2.3 (1.4-3.6)	4.5% (3.1-6.1%)	8.6% (5.8-12%)	0.32 (0.2-0.47)	26 (18-36)
Virgin Islands, U.S.	1990	Female	35 to 39	56% (42-71%)	44% (29-58%)	2.2 (1.3-3.4)	4.4% (3.1-6.1%)	8.5% (6-12%)	0.45 (0.31-0.65)	31 (21-44)
Virgin Islands, U.S.	1990	Female	40 to 44	55% (41-72%)	45% (28-59%)	2.2 (1.3-3.4)	4.7% (3.1-6.9%)	8.7% (5.9-12%)	0.74 (0.48-1.1)	42 (27-61)
Virgin Islands, U.S.	1990	Female	45 to 49	54% (38-69%)	46% (31-62%)	2.1 (1.3-3.3)	4.9% (3-7.5%)	8.3% (5.3-12%)	0.95 (0.58-1.5)	45 (27-70)
Virgin Islands, U.S.	1990	Female	50 to 54	51% (36-67%)	49% (33-64%)	2 (1.1-3.1)	4.8% (2.6-7.7%)	7.4% (4.1-12%)	1.1 (0.61-1.8)	46 (24-76)
Virgin Islands, U.S.	1990	Female	55 to 59	47% (32-64%)	53% (36-68%)	1.8 (0.98-2.9)	4.2% (1.9-7.2%)	6.1% (2.8-10%)	0.92 (0.43-1.6)	32 (14-56)
Virgin Islands, U.S.	1990	Female	60 to 64	44% (29-61%)	56% (39-71%)	1.5 (0.82-2.5)	3.3% (1-6.4%)	4.5% (1.5-8.3%)	0.93 (0.31-1.8)	27 (8.7-52)
Virgin Islands, U.S.	1990	Female	65 to 69	41% (28-57%)	59% (43-72%)	1.4 (0.76-2.2)	2.5% (0.32-5.5%)	3.3% (0.51-7.3%)	0.79 (0.12-1.8)	19 (2.4-42)
Virgin Islands, U.S.	1990	Female	70 to 74	38% (25-53%)	62% (47-75%)	1.2 (0.62-2)	1.7% (-0.25-4.5%)	2.2% (-0.29-5.6%)	0.67 (-0.092-1.7)	13 (-1.8-33)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	1990	Female	75 to 79	34% (22-51%)	66% (49-78%)	1 (0.54-1.7)	1.3% (-0.45-3.8%)	1.6% (-0.52-4.6%)	0.56 (-0.18-1.6)	8.3 (-2.8-24)
Virgin Islands, U.S.	1990	Female	80 to 84	32% (20-46%)	68% (54-80%)	0.98 (0.5-1.6)	1.3% (-0.54-3.6%)	1.4% (-0.69-4.3%)	0.43 (-0.21-1.3)	4.8 (-2-14)
Virgin Islands, U.S.	1990	Female	85 to 89	31% (21-44%)	69% (56-79%)	1 (0.51-1.8)	1.3% (-0.6-3.7%)	1.4% (-0.83-4.3%)	0.33 (-0.2-1)	2.7 (-1.2-8)
Virgin Islands, U.S.	1990	Female	90 to 94	34% (22-48%)	66% (52-78%)	1.1 (0.6-2)	1.4% (-0.7-4.2%)	1.4% (-0.98-4.7%)	0.16 (-0.11-0.51)	0.89 (-0.44-2.7)
Virgin Islands, U.S.	1990	Female	95 plus	38% (24-56%)	62% (44-76%)	1.3 (0.67-2.3)	1.6% (-0.7-5%)	1.6% (-1.1-5.8%)	0.053 (-0.036-0.19)	0.21 (-0.092-0.66)
Virgin Islands, U.S.	1990	Male	All Ages	60% (52-68%)	40% (32-48%)	2.4 (1.6-3.5)	9.3% (6.7-12%)	10% (6.7-14%)	36 (24-52)	1500 (1000-2000)
Virgin Islands, U.S.	1990	Male	15 to 19	62% (45-76%)	38% (24-55%)	2.2 (1.3-3.5)	8.3% (5.7-11%)	11% (7.3-16%)	0.79 (0.49-1.2)	68 (46-98)
Virgin Islands, U.S.	1990	Male	20 to 24	71% (54-84%)	29% (16-46%)	4 (2.5-6.2)	16% (11-21%)	19% (13-27%)	1.4 (0.89-2)	110 (79-160)
Virgin Islands, U.S.	1990	Male	25 to 29	73% (57-85%)	27% (15-43%)	4.2 (2.6-6.2)	16% (11-20%)	19% (13-27%)	1.4 (0.93-2.1)	110 (80-160)
Virgin Islands, U.S.	1990	Male	30 to 34	74% (58-86%)	26% (14-42%)	4.1 (2.6-6.4)	15% (11-19%)	19% (13-26%)	1.6 (1-2.3)	120 (81-150)
Virgin Islands, U.S.	1990	Male	35 to 39	74% (59-86%)	26% (14-41%)	3.9 (2.4-6)	15% (11-19%)	19% (14-25%)	1.8 (1.2-2.6)	120 (82-150)
Virgin Islands, U.S.	1990	Male	40 to 44	73% (59-85%)	27% (15-41%)	3.8 (2.3-5.8)	15% (11-19%)	19% (14-25%)	3 (2.1-4.1)	160 (120-220)
Virgin Islands, U.S.	1990	Male	45 to 49	72% (55-85%)	28% (15-45%)	3.7 (2.2-5.7)	14% (10-19%)	18% (12-24%)	3.9 (2.7-5.4)	190 (130-250)
Virgin Islands, U.S.	1990	Male	50 to 54	70% (54-84%)	30% (16-46%)	3.4 (2.1-5.4)	13% (9.3-18%)	16% (11-22%)	4.1 (2.8-5.8)	170 (120-230)
Virgin Islands, U.S.	1990	Male	55 to 59	69% (52-83%)	31% (17-48%)	3 (1.8-4.8)	11% (7.5-16%)	14% (8.7-19%)	3.5 (2.2-5.1)	120 (79-170)
Virgin Islands, U.S.	1990	Male	60 to 64	67% (48-83%)	33% (17-52%)	2.8 (1.6-4.4)	9.3% (5.7-14%)	11% (6.4-16%)	4 (2.3-6.1)	120 (69-180)
Virgin Islands, U.S.	1990	Male	65 to 69	66% (48-81%)	34% (19-52%)	2.7 (1.6-4.3)	8.5% (4.9-13%)	9.7% (5.5-15%)	3.7 (2-5.7)	88 (50-140)
Virgin Islands, U.S.	1990	Male	70 to 74	63% (45-77%)	37% (23-55%)	2.5 (1.3-4.1)	6.9% (3.5-11%)	7.7% (3.8-13%)	3.2 (1.6-5.4)	62 (31-100)
Virgin Islands, U.S.	1990	Male	75 to 79	58% (39-75%)	42% (25-61%)	2.1 (1.1-3.4)	5.4% (2.4-9.2%)	6% (2.6-10%)	2.3 (1-4.1)	35 (15-59)
Virgin Islands, U.S.	1990	Male	80 to 84	53% (34-70%)	47% (30-66%)	1.7 (0.86-2.8)	3.9% (1.5-7.3%)	4.2% (1.6-8.2%)	1.1 (0.39-2.1)	12 (4.7-23)
Virgin Islands, U.S.	1990	Male	85 to 89	48% (31-65%)	52% (35-69%)	1.4 (0.69-2.4)	2.8% (1.1-6%)	3% (0.99-6.5%)	0.45 (0.15-0.99)	3.6 (1.4-7.6)
Virgin Islands, U.S.	1990	Male	90 to 94	48% (33-66%)	52% (34-67%)	1.2 (0.63-2.2)	1.9% (0.46-4.5%)	1.9% (0.17-4.9%)	0.095 (0.0088-0.25)	0.57 (0.14-1.4)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	1990	Male	95 plus	51% (32-71%)	49% (29-68%)	1.2 (0.54-2.1)	1.3% (-0.36-3.9%)	1.1% (-1-4.1%)	0.011 (-0.01-0.043)	0.054 (-0.015-0.17)
Virgin Islands, U.S.	1995	Female	All Ages	47% (42-54%)	53% (46-58%)	1.5 (0.97-2.3)	2.9% (1.6-4.9%)	3.3% (1.1-6.5%)	10 (3.4-20)	390 (200-650)
Virgin Islands, U.S.	1995	Female	15 to 19	54% (40-69%)	46% (31-60%)	1.9 (1.1-3.1)	3.2% (2-4.6%)	7.2% (4.4-11%)	0.15 (0.092-0.24)	15 (9.6-22)
Virgin Islands, U.S.	1995	Female	20 to 24	58% (45-71%)	42% (29-55%)	2.6 (1.6-4.1)	4.5% (3.2-6.3%)	8.8% (5.6-13%)	0.22 (0.14-0.31)	23 (16-32)
Virgin Islands, U.S.	1995	Female	25 to 29	57% (44-72%)	43% (28-56%)	2.4 (1.5-3.9)	4.3% (3-5.9%)	7.8% (4.4-11%)	0.21 (0.12-0.31)	21 (14-30)
Virgin Islands, U.S.	1995	Female	30 to 34	56% (41-71%)	44% (29-59%)	2.3 (1.3-3.6)	4.1% (2.9-5.7%)	7.8% (4.9-11%)	0.31 (0.2-0.46)	27 (18-37)
Virgin Islands, U.S.	1995	Female	35 to 39	56% (42-71%)	44% (29-58%)	2.2 (1.3-3.4)	4.1% (2.8-5.9%)	7.9% (5.3-11%)	0.44 (0.28-0.64)	30 (20-43)
Virgin Islands, U.S.	1995	Female	40 to 44	55% (41-71%)	45% (29-59%)	2.2 (1.3-3.5)	4.5% (3-6.5%)	8.3% (5.5-12%)	0.68 (0.44-1)	39 (25-57)
Virgin Islands, U.S.	1995	Female	45 to 49	54% (38-70%)	46% (30-62%)	2.2 (1.2-3.4)	4.8% (2.9-7.5%)	8.1% (5.1-12%)	1.1 (0.66-1.6)	51 (30-80)
Virgin Islands, U.S.	1995	Female	50 to 54	51% (36-66%)	49% (34-64%)	2 (1.2-3.1)	4.7% (2.5-7.7%)	7.3% (4.1-12%)	1.3 (0.7-2)	52 (28-84)
Virgin Islands, U.S.	1995	Female	55 to 59	47% (33-64%)	53% (36-67%)	1.8 (0.98-2.9)	4.1% (1.8-7.2%)	6% (2.9-10%)	1.3 (0.61-2.2)	45 (20-78)
Virgin Islands, U.S.	1995	Female	60 to 64	43% (29-60%)	57% (40-71%)	1.6 (0.83-2.6)	3.3% (1-6.4%)	4.5% (1.5-8.7%)	0.98 (0.32-1.9)	29 (8.9-57)
Virgin Islands, U.S.	1995	Female	65 to 69	40% (28-56%)	60% (44-72%)	1.4 (0.74-2.3)	2.5% (0.35-5.5%)	3.3% (0.56-7%)	1 (0.17-2.1)	24 (3.4-51)
Virgin Islands, U.S.	1995	Female	70 to 74	37% (25-53%)	63% (47-75%)	1.2 (0.61-2)	1.8% (-0.22-4.5%)	2.3% (-0.23-5.7%)	0.74 (-0.072-1.9)	14 (-1.7-36)
Virgin Islands, U.S.	1995	Female	75 to 79	34% (22-49%)	66% (51-78%)	1 (0.52-1.7)	1.4% (-0.35-3.9%)	1.7% (-0.51-4.9%)	0.67 (-0.2-1.9)	9.9 (-2.6-28)
Virgin Islands, U.S.	1995	Female	80 to 84	32% (20-46%)	68% (54-80%)	0.99 (0.51-1.7)	1.3% (-0.47-3.8%)	1.5% (-0.64-4.5%)	0.65 (-0.27-1.9)	7.2 (-2.5-21)
Virgin Islands, U.S.	1995	Female	85 to 89	31% (20-45%)	69% (55-80%)	1 (0.52-1.8)	1.3% (-0.51-3.9%)	1.5% (-0.75-4.5%)	0.45 (-0.23-1.4)	3.6 (-1.4-10)
Virgin Islands, U.S.	1995	Female	90 to 94	34% (22-47%)	66% (53-78%)	1.1 (0.58-2)	1.4% (-0.6-4.5%)	1.5% (-0.93-5.2%)	0.24 (-0.15-0.85)	1.4 (-0.59-4.5)
Virgin Islands, U.S.	1995	Female	95 plus	38% (23-57%)	62% (43-77%)	1.3 (0.67-2.4)	1.7% (-0.7-5.5%)	1.7% (-1.2-6.2%)	0.084 (-0.057-0.3)	0.33 (-0.14-1.1)
Virgin Islands, U.S.	1995	Male	All Ages	61% (53-69%)	39% (31-47%)	2.5 (1.6-3.8)	9.6% (6.7-13%)	10% (6.4-14%)	42 (26-60)	1700 (1100-2200)
Virgin Islands, U.S.	1995	Male	15 to 19	61% (45-76%)	39% (24-55%)	2.2 (1.3-3.6)	8.3% (5.6-11%)	11% (7-16%)	0.7 (0.44-1)	59 (41-84)
Virgin Islands, U.S.	1995	Male	20 to 24	70% (54-83%)	30% (17-46%)	4.1 (2.5-6.2)	16% (11-21%)	19% (12-26%)	1.6 (0.99-2.2)	130 (88-170)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	1995	Male	25 to 29	73% (57-85%)	27% (15-43%)	4.3 (2.6-6.6)	15% (11-20%)	18% (12-26%)	1.4 (0.87-2)	110 (75-150)
Virgin Islands, U.S.	1995	Male	30 to 34	74% (58-86%)	26% (14-42%)	4.2 (2.6-6.6)	14% (9.7-19%)	17% (11-24%)	1.8 (1-2.6)	130 (83-170)
Virgin Islands, U.S.	1995	Male	35 to 39	74% (59-86%)	26% (14-41%)	4 (2.4-6.3)	14% (9.8-18%)	18% (11-24%)	2.1 (1.3-3)	130 (89-180)
Virgin Islands, U.S.	1995	Male	40 to 44	73% (59-85%)	27% (15-41%)	3.9 (2.3-6.2)	14% (10-19%)	18% (13-24%)	2.6 (1.7-3.7)	140 (98-190)
Virgin Islands, U.S.	1995	Male	45 to 49	72% (53-85%)	28% (15-47%)	3.8 (2.3-6)	14% (9.9-18%)	17% (12-23%)	4.3 (2.8-5.9)	200 (140-270)
Virgin Islands, U.S.	1995	Male	50 to 54	70% (54-84%)	30% (16-46%)	3.5 (2-5.4)	13% (9.1-18%)	16% (11-21%)	5.4 (3.6-7.5)	220 (150-300)
Virgin Islands, U.S.	1995	Male	55 to 59	69% (52-83%)	31% (17-48%)	3.1 (1.8-5.1)	11% (7.5-16%)	14% (8.6-19%)	5.1 (3.2-7.3)	180 (110-250)
Virgin Islands, U.S.	1995	Male	60 to 64	67% (48-83%)	33% (17-52%)	2.9 (1.6-4.5)	9.4% (5.5-14%)	11% (6.2-16%)	4.1 (2.3-6.1)	120 (70-170)
Virgin Islands, U.S.	1995	Male	65 to 69	66% (47-81%)	34% (19-53%)	2.7 (1.6-4.5)	8.7% (4.8-13%)	9.9% (5.4-15%)	4.8 (2.6-7.5)	120 (63-180)
Virgin Islands, U.S.	1995	Male	70 to 74	63% (45-77%)	37% (23-55%)	2.5 (1.3-4.2)	7% (3.5-11%)	7.8% (3.7-13%)	3.5 (1.6-5.7)	67 (33-110)
Virgin Islands, U.S.	1995	Male	75 to 79	58% (39-76%)	42% (24-61%)	2.1 (1.1-3.5)	5.5% (2.4-9.3%)	6.1% (2.6-11%)	2.7 (1.1-4.7)	41 (18-70)
Virgin Islands, U.S.	1995	Male	80 to 84	52% (33-70%)	48% (30-67%)	1.7 (0.86-2.9)	3.9% (1.6-7.4%)	4.3% (1.6-8.2%)	1.6 (0.59-3)	18 (7.1-33)
Virgin Islands, U.S.	1995	Male	85 to 89	48% (31-64%)	52% (36-69%)	1.4 (0.69-2.4)	2.9% (1.1-6%)	3.1% (1-6.5%)	0.65 (0.22-1.4)	5.2 (2-11)
Virgin Islands, U.S.	1995	Male	90 to 94	48% (32-65%)	52% (35-68%)	1.2 (0.63-2.2)	2% (0.43-5%)	2% (0.22-5.5%)	0.16 (0.017-0.44)	0.97 (0.21-2.4)
Virgin Islands, U.S.	1995	Male	95 plus	51% (31-70%)	49% (30-69%)	1.2 (0.55-2.1)	1.4% (-0.3-4.1%)	1.2% (-0.81-4.4%)	0.021 (-0.014-0.078)	0.1 (-0.021-0.3)
Virgin Islands, U.S.	2000	Female	All Ages	43% (38-49%)	57% (51-62%)	1.3 (0.87-2)	2.7% (1.3-4.5%)	2.9% (0.86-5.8%)	11 (3.2-21)	370 (180-630)
Virgin Islands, U.S.	2000	Female	15 to 19	50% (37-65%)	50% (35-63%)	1.8 (1-2.9)	2.9% (1.9-4.3%)	6.9% (4.3-11%)	0.14 (0.088-0.23)	15 (9.5-22)
Virgin Islands, U.S.	2000	Female	20 to 24	52% (39-65%)	48% (35-61%)	2.2 (1.3-3.4)	4.1% (2.8-5.7%)	7.9% (4.6-12%)	0.13 (0.078-0.21)	15 (11-22)
Virgin Islands, U.S.	2000	Female	25 to 29	52% (41-67%)	48% (33-59%)	2.2 (1.3-3.4)	3.8% (2.5-5.2%)	6.6% (2.9-10%)	0.15 (0.068-0.23)	17 (11-25)
Virgin Islands, U.S.	2000	Female	30 to 34	53% (41-66%)	47% (34-59%)	2.1 (1.3-3.3)	3.7% (2.4-5.2%)	7% (3.7-10%)	0.23 (0.13-0.35)	21 (14-30)
Virgin Islands, U.S.	2000	Female	35 to 39	51% (40-66%)	49% (34-60%)	2 (1.2-3.1)	3.7% (2.5-5.2%)	7% (4.4-9.7%)	0.37 (0.23-0.52)	27 (17-38)
Virgin Islands, U.S.	2000	Female	40 to 44	48% (37-63%)	52% (37-63%)	1.8 (1.1-2.9)	3.8% (2.5-5.5%)	7.1% (4.7-10%)	0.55 (0.35-0.81)	33 (21-48)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	2000	Female	45 to 49	47% (36-62%)	53% (38-64%)	1.8 (1.1-2.8)	4.2% (2.5-6.4%)	7.1% (4.4-11%)	0.87 (0.52-1.3)	42 (25-64)
Virgin Islands, U.S.	2000	Female	50 to 54	46% (34-59%)	54% (41-66%)	1.7 (1-2.7)	4.1% (2.3-6.7%)	6.5% (3.8-10%)	1.3 (0.73-2)	53 (29-86)
Virgin Islands, U.S.	2000	Female	55 to 59	42% (32-57%)	58% (43-68%)	1.6 (0.94-2.4)	3.7% (1.8-6.4%)	5.4% (2.8-9.1%)	1.4 (0.69-2.3)	48 (22-82)
Virgin Islands, U.S.	2000	Female	60 to 64	43% (32-58%)	57% (42-68%)	1.5 (0.89-2.4)	3% (0.94-5.9%)	4.2% (1.4-8%)	1.2 (0.38-2.3)	34 (11-67)
Virgin Islands, U.S.	2000	Female	65 to 69	39% (28-55%)	61% (45-72%)	1.3 (0.73-2.2)	2.3% (0.25-5%)	3.1% (0.53-6.7%)	1.1 (0.17-2.2)	25 (2.7-53)
Virgin Islands, U.S.	2000	Female	70 to 74	34% (23-49%)	66% (51-77%)	1.1 (0.56-1.9)	1.6% (-0.11-4%)	2.1% (-0.14-5.2%)	0.75 (-0.053-1.9)	14 (-0.98-37)
Virgin Islands, U.S.	2000	Female	75 to 79	31% (21-45%)	69% (55-79%)	0.94 (0.5-1.6)	1.3% (-0.26-3.6%)	1.6% (-0.34-4.5%)	0.7 (-0.15-2)	10 (-2.1-29)
Virgin Islands, U.S.	2000	Female	80 to 84	29% (19-43%)	71% (57-81%)	0.9 (0.46-1.5)	1.2% (-0.35-3.5%)	1.4% (-0.52-4.2%)	0.73 (-0.27-2.2)	8.3 (-2.4-24)
Virgin Islands, U.S.	2000	Female	85 to 89	29% (18-41%)	71% (59-82%)	0.92 (0.48-1.6)	1.2% (-0.36-3.6%)	1.4% (-0.55-4.1%)	0.64 (-0.25-1.9)	5.2 (-1.5-15)
Virgin Islands, U.S.	2000	Female	90 to 94	31% (20-44%)	69% (56-80%)	1 (0.54-1.8)	1.4% (-0.4-4%)	1.4% (-0.65-4.5%)	0.33 (-0.14-1)	1.9 (-0.55-5.5)
Virgin Islands, U.S.	2000	Female	95 plus	35% (21-53%)	65% (47-79%)	1.2 (0.6-2.1)	1.6% (-0.47-5%)	1.6% (-0.81-5.6%)	0.13 (-0.065-0.44)	0.51 (-0.15-1.6)
Virgin Islands, U.S.	2000	Male	All Ages	64% (57-71%)	36% (29-43%)	2.8 (1.8-4.1)	10% (6.9-14%)	10% (6.4-15%)	50 (31-70)	1800 (1200-2500)
Virgin Islands, U.S.	2000	Male	15 to 19	63% (47-78%)	37% (22-53%)	2.4 (1.4-3.8)	8.8% (5.8-12%)	7.2% (7.6-18%)	0.76 (0.46-1.1)	64 (43-90)
Virgin Islands, U.S.	2000	Male	20 to 24	73% (59-84%)	27% (16-41%)	4.5 (2.8-6.7)	17% (12-22%)	20% (14-29%)	1.2 (0.79-1.7)	98 (69-130)
Virgin Islands, U.S.	2000	Male	25 to 29	76% (64-86%)	24% (14-36%)	4.7 (2.9-7)	16% (11-21%)	19% (12-27%)	1.3 (0.77-1.9)	98 (67-140)
Virgin Islands, U.S.	2000	Male	30 to 34	78% (66-88%)	22% (12-34%)	4.7 (2.9-7.4)	15% (10-20%)	19% (11-26%)	1.6 (0.99-2.4)	120 (78-160)
Virgin Islands, U.S.	2000	Male	35 to 39	78% (66-88%)	22% (12-34%)	4.5 (2.8-6.9)	14% (9.9-19%)	18% (11-25%)	2.2 (1.3-3.2)	140 (92-190)
Virgin Islands, U.S.	2000	Male	40 to 44	76% (64-86%)	24% (14-36%)	4.3 (2.6-6.6)	15% (10-19%)	18% (12-25%)	3 (1.9-4.2)	170 (110-220)
Virgin Islands, U.S.	2000	Male	45 to 49	74% (60-85%)	26% (15-40%)	4.2 (2.6-6.4)	15% (10-19%)	18% (12-24%)	4 (2.7-5.4)	190 (130-250)
Virgin Islands, U.S.	2000	Male	50 to 54	74% (59-85%)	26% (15-41%)	3.9 (2.4-6)	14% (9.6-18%)	17% (11-22%)	6.1 (4.1-8.3)	250 (170-330)
Virgin Islands, U.S.	2000	Male	55 to 59	71% (57-83%)	29% (17-43%)	3.4 (2.1-5.2)	12% (7.7-16%)	14% (8.9-19%)	6.9 (4.3-9.6)	240 (150-330)
Virgin Islands, U.S.	2000	Male	60 to 64	70% (54-84%)	30% (16-46%)	3.2 (1.8-4.8)	9.8% (5.6-14%)	11% (6.3-17%)	5.8 (3.3-8.6)	170 (98-250)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	2000	Male	65 to 69	69% (52-83%)	31% (17-48%)	3.1 (1.7-4.8)	9.1% (4.9-14%)	10% (5.6-16%)	5.4 (2.9-8.3)	130 (71-200)
Virgin Islands, U.S.	2000	Male	70 to 74	66% (49-80%)	34% (20-51%)	2.8 (1.5-4.5)	7.4% (3.6-12%)	8.4% (4-14%)	4.7 (2.2-7.7)	90 (43-140)
Virgin Islands, U.S.	2000	Male	75 to 79	61% (43-77%)	39% (23-57%)	2.4 (1.3-4)	5.8% (2.5-10%)	6.5% (2.7-11%)	3.1 (1.3-5.4)	46 (20-79)
Virgin Islands, U.S.	2000	Male	80 to 84	55% (36-73%)	45% (27-64%)	1.9 (0.96-3.2)	4.2% (1.7-8%)	4.6% (1.7-8.9%)	2.1 (0.77-4)	24 (9.1-45)
Virgin Islands, U.S.	2000	Male	85 to 89	50% (33-67%)	50% (33-67%)	1.5 (0.76-2.7)	3.1% (1.1-6.3%)	3.3% (1.1-6.9%)	1.1 (0.34-2.2)	8.7 (3.1-18)
Virgin Islands, U.S.	2000	Male	90 to 94	51% (35-69%)	49% (31-65%)	1.4 (0.72-2.5)	2.1% (0.35-5.1%)	2.1% (0.12-5.7%)	0.26 (0.015-0.7)	1.6 (0.26-3.9)
Virgin Islands, U.S.	2000	Male	95 plus	54% (34-73%)	46% (27-66%)	1.3 (0.61-2.4)	1.6% (-0.29-4.6%)	1.4% (-0.82-5%)	0.041 (-0.025-0.15)	0.19 (-0.034-0.55)
Virgin Islands, U.S.	2005	Female	All Ages	42% (37-48%)	58% (52-63%)	1.4 (0.88-2)	2.7% (1.3-4.5%)	2.9% (0.87-5.6%)	12 (3.6-24)	390 (190-650)
Virgin Islands, U.S.	2005	Female	15 to 19	48% (36-64%)	52% (36-64%)	1.7 (1-2.8)	2.9% (1.9-4.3%)	7% (4.2-11%)	0.13 (0.074-0.2)	13 (8.4-19)
Virgin Islands, U.S.	2005	Female	20 to 24	50% (38-63%)	50% (37-62%)	2.2 (1.3-3.3)	3.9% (2.6-5.5%)	7.5% (3.6-11%)	0.15 (0.074-0.24)	18 (12-26)
Virgin Islands, U.S.	2005	Female	25 to 29	51% (40-66%)	49% (34-60%)	2.1 (1.3-3.3)	3.6% (2.2-5.1%)	6.1% (1.1-9.7%)	0.12 (0.023-0.2)	15 (8.6-21)
Virgin Islands, U.S.	2005	Female	30 to 34	50% (39-64%)	50% (36-61%)	2 (1.3-3.1)	3.4% (2-4.9%)	6.2% (2-9.5%)	0.19 (0.066-0.29)	17 (10-26)
Virgin Islands, U.S.	2005	Female	35 to 39	50% (40-64%)	50% (36-60%)	2 (1.2-3)	3.5% (2.3-5%)	6.7% (3.9-9.3%)	0.32 (0.19-0.48)	23 (15-34)
Virgin Islands, U.S.	2005	Female	40 to 44	47% (36-62%)	53% (38-64%)	1.8 (1.1-2.8)	3.7% (2.4-5.5%)	7% (4.6-9.8%)	0.54 (0.34-0.79)	32 (20-47)
Virgin Islands, U.S.	2005	Female	45 to 49	47% (36-61%)	53% (39-64%)	1.9 (1.1-2.9)	4.1% (2.5-6.1%)	7.1% (4.5-10%)	0.85 (0.54-1.2)	42 (25-61)
Virgin Islands, U.S.	2005	Female	50 to 54	46% (35-59%)	54% (41-65%)	1.8 (1.1-2.8)	4.2% (2.4-6.8%)	6.8% (4-11%)	1.2 (0.72-1.9)	50 (28-80)
Virgin Islands, U.S.	2005	Female	55 to 59	41% (31-55%)	59% (45-69%)	1.5 (0.95-2.4)	3.6% (1.8-6.2%)	5.4% (2.9-8.9%)	1.6 (0.83-2.5)	54 (27-92)
Virgin Islands, U.S.	2005	Female	60 to 64	42% (32-57%)	58% (43-68%)	1.5 (0.91-2.4)	3.1% (1-5.8%)	4.3% (1.6-7.9%)	1.6 (0.58-2.8)	45 (15-85)
Virgin Islands, U.S.	2005	Female	65 to 69	38% (29-53%)	62% (47-71%)	1.3 (0.77-2.1)	2.4% (0.47-5%)	3.2% (0.67-6.5%)	1.2 (0.26-2.5)	30 (5.7-61)
Virgin Islands, U.S.	2005	Female	70 to 74	33% (24-46%)	67% (54-76%)	1 (0.57-1.7)	1.7% (0.082-3.9%)	2.2% (0.082-4.9%)	0.96 (0.035-2.1)	18 (0.98-42)
Virgin Islands, U.S.	2005	Female	75 to 79	33% (24-46%)	67% (54-76%)	1 (0.57-1.6)	1.4% (-0.28-3.6%)	1.7% (-0.33-4.5%)	0.75 (-0.14-2)	11 (-2.2-30)
Virgin Islands, U.S.	2005	Female	80 to 84	31% (21-44%)	69% (56-79%)	0.99 (0.52-1.7)	1.4% (-0.34-3.7%)	1.6% (-0.45-4.4%)	0.99 (-0.27-2.7)	11 (-2.8-29)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	2005	Female	85 to 89	30% (19-41%)	70% (59-81%)	0.97 (0.52-1.6)	1.3% (-0.26-3.8%)	1.5% (-0.44-4.4%)	0.84 (-0.25-2.5)	6.8 (-1.3-19)
Virgin Islands, U.S.	2005	Female	90 to 94	32% (21-45%)	68% (55-79%)	1.1 (0.58-1.9)	1.5% (-0.25-4.3%)	1.6% (-0.48-4.9%)	0.55 (-0.16-1.7)	3.1 (-0.52-9.1)
Virgin Islands, U.S.	2005	Female	95 plus	36% (23-53%)	64% (47-77%)	1.3 (0.63-2.3)	1.7% (-0.37-5.3%)	1.8% (-0.72-5.9%)	0.19 (-0.08-0.65)	0.76 (-0.16-2.3)
Virgin Islands, U.S.	2005	Male	All Ages	65% (58-72%)	35% (28-42%)	2.9 (1.9-4.3)	10% (7-14%)	10% (6.3-15%)	57 (35-82)	2000 (1400-2700)
Virgin Islands, U.S.	2005	Male	15 to 19	62% (46-76%)	38% (24-54%)	2.3 (1.4-3.7)	8.8% (5.7-12%)	13% (7.4-18%)	0.64 (0.37-0.93)	53 (35-75)
Virgin Islands, U.S.	2005	Male	20 to 24	73% (58-84%)	27% (16-42%)	4.5 (2.8-6.7)	15% (11-21%)	18% (11-26%)	1.5 (0.93-2.1)	120 (83-170)
Virgin Islands, U.S.	2005	Male	25 to 29	77% (65-86%)	23% (14-35%)	4.8 (3-7.2)	16% (9.9-22%)	20% (10-29%)	1.2 (0.63-1.8)	94 (59-130)
Virgin Islands, U.S.	2005	Male	30 to 34	78% (66-87%)	22% (13-34%)	4.7 (3-7.2)	15% (9.4-20%)	18% (10-27%)	1.4 (0.78-2.1)	100 (63-140)
Virgin Islands, U.S.	2005	Male	35 to 39	77% (66-87%)	23% (13-34%)	4.5 (2.8-6.8)	14% (9.8-19%)	18% (12-25%)	2.1 (1.3-2.9)	130 (85-180)
Virgin Islands, U.S.	2005	Male	40 to 44	76% (65-86%)	24% (14-35%)	4.4 (2.8-6.6)	15% (10-19%)	19% (12-25%)	3 (1.9-4.1)	160 (110-220)
Virgin Islands, U.S.	2005	Male	45 to 49	73% (59-84%)	27% (16-41%)	4.2 (2.6-6.3)	15% (10-19%)	18% (12-24%)	4.3 (3-5.9)	210 (150-270)
Virgin Islands, U.S.	2005	Male	50 to 54	73% (60-85%)	27% (15-40%)	4 (2.5-6)	14% (9.8-18%)	17% (12-22%)	5.5 (3.8-7.3)	220 (160-290)
Virgin Islands, U.S.	2005	Male	55 to 59	72% (58-84%)	28% (16-42%)	3.6 (2.2-5.5)	12% (8-17%)	15% (9.5-20%)	7.8 (5-11)	270 (180-360)
Virgin Islands, U.S.	2005	Male	60 to 64	70% (54-83%)	30% (17-46%)	3.2 (1.9-5)	10% (5.9-15%)	12% (6.7-17%)	7.9 (4.5-12)	230 (140-340)
Virgin Islands, U.S.	2005	Male	65 to 69	68% (52-81%)	32% (19-48%)	3 (1.8-4.7)	9.2% (5-14%)	11% (5.8-16%)	6.9 (3.8-10)	170 (93-240)
Virgin Islands, U.S.	2005	Male	70 to 74	66% (50-79%)	34% (21-50%)	2.8 (1.6-4.5)	7.6% (3.8-12%)	8.6% (4.2-14%)	5.4 (2.6-8.5)	110 (54-160)
Virgin Islands, U.S.	2005	Male	75 to 79	67% (51-80%)	33% (20-49%)	2.8 (1.6-4.4)	6.4% (2.7-11%)	7.2% (2.9-12%)	4.4 (1.8-7.4)	67 (29-110)
Virgin Islands, U.S.	2005	Male	80 to 84	61% (44-76%)	39% (24-56%)	2.2 (1.2-3.6)	4.8% (1.8-8.9%)	5.3% (1.9-10%)	2.9 (1.1-5.5)	33 (12-60)
Virgin Islands, U.S.	2005	Male	85 to 89	52% (35-68%)	48% (32-65%)	1.6 (0.86-2.7)	3.3% (1.2-6.4%)	3.5% (1.1-7.1%)	1.3 (0.4-2.6)	10 (3.7-20)
Virgin Islands, U.S.	2005	Male	90 to 94	53% (37-69%)	47% (31-63%)	1.5 (0.79-2.5)	2.3% (0.39-5.2%)	2.3% (0.11-5.8%)	0.4 (0.02-1)	2.4 (0.42-5.4)
Virgin Islands, U.S.	2005	Male	95 plus	55% (37-73%)	45% (27-63%)	1.4 (0.67-2.5)	1.7% (-0.25-5%)	1.5% (-0.79-5.4%)	0.064 (-0.032-0.22)	0.29 (-0.042-0.85)
Virgin Islands, U.S.	2010	Female	All Ages	41% (36-47%)	59% (53-64%)	1.4 (0.94-2.2)	2.7% (1.4-4.8%)	3% (0.92-6.2%)	14 (4.3-29)	420 (200-730)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	2010	Female	15 to 19	48% (36-64%)	52% (36-64%)	1.8 (1.1-2.9)	2.9% (1.8-4.3%)	7% (4.1-11%)	0.12 (0.069-0.19)	12 (7.7-18)
Virgin Islands, U.S.	2010	Female	20 to 24	50% (39-64%)	50% (36-61%)	2.3 (1.4-3.5)	4% (2.6-5.7%)	7.6% (3.3-12%)	0.12 (0.055-0.19)	15 (9.1-21)
Virgin Islands, U.S.	2010	Female	25 to 29	51% (40-65%)	49% (35-60%)	2.2 (1.4-3.4)	3.4% (1.9-5%)	5.5% (0.84-9.7%)	0.11 (0.017-0.2)	14 (7.6-21)
Virgin Islands, U.S.	2010	Female	30 to 34	49% (38-63%)	51% (37-62%)	2 (1.3-3.1)	3.2% (1.8-4.8%)	5.7% (0.94-9.2%)	0.16 (0.025-0.26)	16 (8.4-23)
Virgin Islands, U.S.	2010	Female	35 to 39	49% (38-64%)	51% (36-62%)	2 (1.2-3)	3.4% (2-5%)	6.3% (3-9.1%)	0.27 (0.12-0.4)	20 (12-30)
Virgin Islands, U.S.	2010	Female	40 to 44	47% (36-62%)	53% (38-64%)	1.9 (1.2-3)	3.7% (2.5-5.6%)	7.1% (4.6-10%)	0.51 (0.31-0.76)	30 (19-44)
Virgin Islands, U.S.	2010	Female	45 to 49	47% (36-61%)	53% (39-64%)	1.9 (1.2-3)	4.1% (2.6-6.3%)	7.3% (4.7-11%)	0.82 (0.5-1.2)	41 (25-63)
Virgin Islands, U.S.	2010	Female	50 to 54	46% (35-59%)	54% (41-65%)	1.9 (1.1-3)	4.3% (2.5-6.9%)	7% (4.2-11%)	1.2 (0.75-1.9)	52 (29-84)
Virgin Islands, U.S.	2010	Female	55 to 59	41% (31-55%)	59% (45-69%)	1.6 (0.95-2.6)	3.7% (1.9-6.3%)	5.6% (3-9.4%)	1.5 (0.77-2.4)	51 (26-88)
Virgin Islands, U.S.	2010	Female	60 to 64	40% (31-55%)	60% (45-69%)	1.5 (0.92-2.5)	3.1% (1.2-5.9%)	4.4% (1.8-8.2%)	1.8 (0.76-3.4)	54 (21-100)
Virgin Islands, U.S.	2010	Female	65 to 69	38% (29-52%)	62% (48-71%)	1.3 (0.8-2.2)	2.5% (0.62-5.3%)	3.4% (0.91-7%)	1.7 (0.45-3.5)	41 (9.6-88)
Virgin Islands, U.S.	2010	Female	70 to 74	33% (24-46%)	67% (54-76%)	1.1 (0.6-1.8)	1.8% (0.3-4.3%)	2.4% (0.46-5.5%)	1.2 (0.24-2.8)	24 (3.8-56)
Virgin Islands, U.S.	2010	Female	75 to 79	33% (23-47%)	67% (53-77%)	1.1 (0.6-1.7)	1.5% (-0.11-4%)	1.9% (-0.14-4.9%)	1.1 (-0.079-2.7)	16 (-1.2-41)
Virgin Islands, U.S.	2010	Female	80 to 84	32% (22-46%)	68% (54-78%)	1.1 (0.57-1.8)	1.6% (-0.096-4.3%)	1.9% (-0.15-5.1%)	1.3 (-0.1-3.4)	14 (-0.87-38)
Virgin Islands, U.S.	2010	Female	85 to 89	31% (20-43%)	69% (57-80%)	1 (0.55-1.7)	1.6% (-0.075-4.2%)	1.8% (-0.21-4.8%)	1.2 (-0.14-3.1)	9.5 (-0.45-24)
Virgin Islands, U.S.	2010	Female	90 to 94	33% (22-46%)	67% (54-78%)	1.2 (0.6-2)	1.8% (-0.071-4.6%)	1.9% (-0.29-5.3%)	0.78 (-0.12-2.2)	4.4 (-0.18-12)
Virgin Islands, U.S.	2010	Female	95 plus	37% (23-55%)	63% (45-77%)	1.4 (0.67-2.4)	2.1% (-0.14-5.6%)	2.2% (-0.46-6.2%)	0.35 (-0.073-1)	1.3 (-0.087-3.6)
Virgin Islands, U.S.	2010	Male	All Ages	65% (58-72%)	35% (28-42%)	3.1 (2.1-4.7)	11% (7.2-14%)	11% (6.5-15%)	68 (42-98)	2200 (1500-3000)
Virgin Islands, U.S.	2010	Male	15 to 19	61% (45-76%)	39% (24-55%)	2.4 (1.4-3.8)	8.8% (5.6-12%)	13% (7.5-19%)	0.64 (0.37-0.94)	54 (35-75)
Virgin Islands, U.S.	2010	Male	20 to 24	73% (58-84%)	27% (16-42%)	4.8 (3-7.2)	17% (12-23%)	21% (14-31%)	1.2 (0.76-1.7)	94 (65-130)
Virgin Islands, U.S.	2010	Male	25 to 29	77% (66-87%)	23% (13-34%)	5.2 (3.3-7.9)	16% (8.2-22%)	20% (7.8-29%)	1.2 (0.45-1.8)	91 (46-130)
Virgin Islands, U.S.	2010	Male	30 to 34	77% (65-87%)	23% (13-35%)	5 (3.2-7.7)	15% (8.3-21%)	18% (8.4-27%)	1.4 (0.62-2.1)	96 (52-140)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	2010	Male	35 to 39	77% (65-87%)	23% (13-35%)	4.8 (3-7.1)	15% (9.7-19%)	19% (11-26%)	1.9 (1.2-2.8)	120 (79-160)
Virgin Islands, U.S.	2010	Male	40 to 44	77% (65-86%)	23% (14-35%)	4.8 (3-7.2)	15% (11-20%)	19% (14-26%)	3.1 (2.1-4.3)	170 (120-230)
Virgin Islands, U.S.	2010	Male	45 to 49	73% (58-84%)	27% (16-42%)	4.4 (2.8-6.7)	15% (11-19%)	19% (13-24%)	4.6 (3.2-6.3)	210 (150-290)
Virgin Islands, U.S.	2010	Male	50 to 54	73% (59-85%)	27% (15-41%)	4.2 (2.7-6.5)	15% (10-19%)	18% (12-23%)	6.6 (4.7-9)	270 (190-360)
Virgin Islands, U.S.	2010	Male	55 to 59	73% (59-84%)	27% (16-41%)	3.8 (2.4-5.9)	13% (8.7-18%)	16% (10-21%)	7.6 (5-10)	260 (170-360)
Virgin Islands, U.S.	2010	Male	60 to 64	70% (54-83%)	30% (17-46%)	3.4 (2.1-5.3)	11% (6.5-15%)	12% (7.3-18%)	9.3 (5.6-13)	270 (160-390)
Virgin Islands, U.S.	2010	Male	65 to 69	68% (52-82%)	32% (18-48%)	3.2 (1.9-5)	9.8% (5.5-14%)	11% (6.2-17%)	10 (5.9-16)	250 (140-370)
Virgin Islands, U.S.	2010	Male	70 to 74	66% (51-79%)	34% (21-49%)	3.1 (1.8-5)	8.3% (4.4-12%)	9.4% (4.9-14%)	7.8 (4-12)	150 (80-230)
Virgin Islands, U.S.	2010	Male	75 to 79	67% (51-80%)	33% (20-49%)	3 (1.8-4.9)	7.1% (3.3-11%)	7.9% (3.5-13%)	5.9 (2.6-9.6)	89 (42-140)
Virgin Islands, U.S.	2010	Male	80 to 84	61% (42-76%)	39% (24-58%)	2.4 (1.3-3.9)	5.4% (2.2-9.7%)	5.9% (2.3-11%)	4.2 (1.6-7.6)	46 (19-82)
Virgin Islands, U.S.	2010	Male	85 to 89	52% (34-68%)	48% (32-66%)	1.7 (0.9-2.9)	3.7% (1.4-7.1%)	4% (1.4-7.9%)	1.8 (0.63-3.6)	15 (5.4-28)
Virgin Islands, U.S.	2010	Male	90 to 94	52% (36-69%)	48% (31-64%)	1.5 (0.81-2.7)	2.6% (0.62-5.7%)	2.7% (0.41-6.2%)	0.52 (0.079-1.2)	3 (0.72-6.6)
Virgin Islands, U.S.	2010	Male	95 plus	54% (35-73%)	46% (27-65%)	1.4 (0.66-2.6)	2% (-0.016-5.3%)	1.9% (-0.49-5.7%)	0.11 (-0.028-0.34)	0.47 (-0.0035-1.3)
Virgin Islands, U.S.	2016	Female	All Ages	46% (40-52%)	54% (48-60%)	1.7 (1.1-2.4)	2.8% (1.3-5%)	3.1% (0.95-6.2%)	17 (5.2-35)	450 (200-820)
Virgin Islands, U.S.	2016	Female	15 to 19	53% (40-68%)	47% (32-60%)	2.1 (1.2-3.3)	2.9% (1.9-4.4%)	7.6% (4.3-12%)	0.098 (0.054-0.16)	10 (6.4-16)
Virgin Islands, U.S.	2016	Female	20 to 24	57% (43-71%)	43% (29-57%)	2.8 (1.6-4.2)	4.1% (2.6-5.9%)	8.2% (3.1-13%)	0.14 (0.051-0.22)	17 (10-24)
Virgin Islands, U.S.	2016	Female	25 to 29	57% (43-72%)	43% (28-57%)	2.6 (1.6-3.9)	3.4% (1.8-5.2%)	5.5% (0.83-10%)	0.091 (0.013-0.18)	12 (6.1-19)
Virgin Islands, U.S.	2016	Female	30 to 34	56% (41-71%)	44% (29-59%)	2.5 (1.5-3.7)	3.2% (1.6-5%)	5.5% (0.83-9.9%)	0.13 (0.018-0.24)	14 (6.4-22)
Virgin Islands, U.S.	2016	Female	35 to 39	55% (42-70%)	45% (30-58%)	2.4 (1.5-3.7)	3.4% (1.7-5.3%)	6.5% (1.4-10%)	0.23 (0.051-0.4)	18 (8.7-28)
Virgin Islands, U.S.	2016	Female	40 to 44	54% (41-71%)	46% (29-59%)	2.4 (1.4-3.7)	4% (2.5-6.1%)	7.7% (4.5-11%)	0.44 (0.24-0.7)	27 (16-41)
Virgin Islands, U.S.	2016	Female	45 to 49	53% (37-69%)	47% (31-63%)	2.3 (1.4-3.6)	4.4% (2.6-6.9%)	7.9% (4.8-12%)	0.81 (0.44-1.3)	40 (22-64)
Virgin Islands, U.S.	2016	Female	50 to 54	50% (35-66%)	50% (34-65%)	2.2 (1.2-3.4)	4.5% (2.5-7.5%)	7.6% (4.6-12%)	1.2 (0.68-2)	51 (29-85)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	2016	Female	55 to 59	46% (31-63%)	54% (37-69%)	1.9 (1-2.9)	4.1% (2-7%)	6.3% (3.4-10%)	1.6 (0.83-2.7)	57 (28-98)
Virgin Islands, U.S.	2016	Female	60 to 64	42% (28-60%)	58% (40-72%)	1.7 (0.95-2.7)	3.3% (1.3-6.1%)	4.8% (2.1-8.6%)	1.8 (0.74-3.2)	52 (20-97)
Virgin Islands, U.S.	2016	Female	65 to 69	40% (27-55%)	60% (45-73%)	1.5 (0.8-2.4)	2.7% (0.59-5.6%)	3.7% (0.95-7.4%)	2.1 (0.57-4.4)	52 (12-110)
Virgin Islands, U.S.	2016	Female	70 to 74	37% (25-53%)	63% (47-75%)	1.2 (0.68-2.1)	2% (0.23-4.7%)	2.7% (0.35-6.1%)	2 (0.27-4.5)	38 (4.4-89)
Virgin Islands, U.S.	2016	Female	75 to 79	33% (21-49%)	67% (51-79%)	1.1 (0.6-1.9)	1.7% (0.015-4.1%)	2.1% (-0.0087-5.2%)	1.6 (-0.0067-4)	24 (0.2-62)
Virgin Islands, U.S.	2016	Female	80 to 84	31% (20-45%)	69% (55-80%)	1.1 (0.56-1.9)	1.7% (0.026-4.2%)	2% (-0.01-5.1%)	1.8 (-0.0095-4.7)	21 (0.32-53)
Virgin Islands, U.S.	2016	Female	85 to 89	31% (20-44%)	69% (56-80%)	1.1 (0.55-1.9)	1.7% (-0.045-4.4%)	1.9% (-0.2-5.1%)	1.5 (-0.15-3.8)	12 (-0.29-30)
Virgin Islands, U.S.	2016	Female	90 to 94	33% (22-47%)	67% (53-78%)	1.2 (0.63-2.1)	1.9% (-0.12-4.9%)	2.1% (-0.34-5.7%)	0.98 (-0.16-2.7)	5.5 (-0.34-14)
Virgin Islands, U.S.	2016	Female	95 plus	37% (23-56%)	63% (44-77%)	1.4 (0.67-2.6)	2.2% (-0.068-6.2%)	2.4% (-0.4-7.2%)	0.46 (-0.079-1.4)	1.8 (-0.057-5)
Virgin Islands, U.S.	2016	Male	All Ages	62% (54-69%)	38% (31-46%)	3 (2-4.5)	9.9% (6.9-13%)	9.7% (5.8-13%)	71 (43-100)	2100 (1400-2900)
Virgin Islands, U.S.	2016	Male	15 to 19	61% (44-76%)	39% (24-56%)	2.6 (1.5-4.1)	8.6% (5.6-12%)	13% (7.7-19%)	0.56 (0.32-0.82)	47 (30-67)
Virgin Islands, U.S.	2016	Male	20 to 24	70% (53-83%)	30% (17-47%)	4.8 (2.9-7.4)	16% (11-22%)	20% (13-30%)	1.3 (0.77-2)	110 (68-150)
Virgin Islands, U.S.	2016	Male	25 to 29	73% (56-85%)	27% (15-44%)	4.9 (3-7.4)	14% (6-21%)	18% (3.4-28%)	0.92 (0.18-1.5)	73 (31-110)
Virgin Islands, U.S.	2016	Male	30 to 34	74% (57-85%)	26% (15-43%)	4.9 (3-7.4)	13% (4.9-19%)	17% (2.6-26%)	0.96 (0.15-1.6)	70 (25-100)
Virgin Islands, U.S.	2016	Male	35 to 39	74% (58-86%)	26% (14-42%)	4.7 (2.8-6.9)	14% (7.4-19%)	17% (7.3-25%)	1.4 (0.55-2.1)	89 (47-130)
Virgin Islands, U.S.	2016	Male	40 to 44	73% (57-85%)	27% (15-43%)	4.6 (2.7-7)	15% (10-19%)	19% (12-25%)	2.3 (1.4-3.4)	130 (83-180)
Virgin Islands, U.S.	2016	Male	45 to 49	71% (53-84%)	29% (16-47%)	4.4 (2.7-6.7)	15% (11-19%)	18% (13-24%)	3.9 (2.6-5.6)	190 (130-250)
Virgin Islands, U.S.	2016	Male	50 to 54	70% (52-83%)	30% (17-48%)	4.1 (2.4-6.3)	14% (10-18%)	17% (12-22%)	5.8 (3.8-8.1)	240 (160-320)
Virgin Islands, U.S.	2016	Male	55 to 59	68% (51-82%)	32% (18-49%)	3.6 (2.1-5.6)	13% (8.6-17%)	15% (10-20%)	7.9 (5.1-11)	270 (180-370)
Virgin Islands, U.S.	2016	Male	60 to 64	66% (47-82%)	34% (18-53%)	3.3 (1.9-5.3)	10% (6.6-15%)	12% (7.6-17%)	8.1 (4.9-12)	240 (140-340)
Virgin Islands, U.S.	2016	Male	65 to 69	65% (47-81%)	35% (19-53%)	3.2 (1.8-5.1)	9.8% (5.9-14%)	11% (6.7-16%)	11 (6.3-16)	260 (150-390)
Virgin Islands, U.S.	2016	Male	70 to 74	62% (44-77%)	38% (23-56%)	2.9 (1.7-4.7)	8% (4.4-12%)	9.2% (4.8-14%)	11 (5.9-17)	210 (120-330)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Virgin Islands, U.S.	2016	Male	75 to 79	57% (38-75%)	43% (25-62%)	2.4 (1.3-4.1)	6.3% (3.1-10%)	7.1% (3.4-11%)	7.5 (3.5-12)	110 (53-180)
Virgin Islands, U.S.	2016	Male	80 to 84	51% (33-69%)	49% (31-67%)	1.9 (1-3.4)	4.8% (2-8.2%)	5.3% (2.1-9.3%)	4.9 (2.1-8.7)	55 (24-95)
Virgin Islands, U.S.	2016	Male	85 to 89	47% (29-64%)	53% (36-71%)	1.6 (0.75-2.7)	3.6% (1.4-6.9%)	3.9% (1.4-7.8%)	2.3 (0.85-4.5)	19 (7.5-35)
Virgin Islands, U.S.	2016	Male	90 to 94	47% (31-64%)	53% (36-69%)	1.4 (0.7-2.5)	2.6% (0.76-5.5%)	2.7% (0.56-6.1%)	0.7 (0.15-1.6)	4.1 (1.2-8.7)
Virgin Islands, U.S.	2016	Male	95 plus	49% (30-69%)	51% (31-70%)	1.3 (0.58-2.5)	2.1% (0.17-5.3%)	2% (-0.28-5.8%)	0.13 (-0.019-0.38)	0.56 (0.045-1.4)
South Sudan	1990	Female	All Ages	33% (27-39%)	67% (61-73%)	0.12 (0.054-0.25)	0.11% (0.029-0.19%)	0.098% (-0.079-0.26%)	46 (-38-120)	3800 (1000-6700)
South Sudan	1990	Female	15 to 19	35% (22-50%)	65% (50-78%)	0.16 (0.055-0.35)	0.31% (0.093-0.59%)	0.14% (-0.18-0.54%)	1 (-1.4-4.4)	270 (77-540)
South Sudan	1990	Female	20 to 24	41% (26-57%)	59% (43-74%)	0.21 (0.078-0.5)	0.55% (0.24-0.95%)	0.16% (-0.23-0.63%)	1.5 (-2-5.8)	510 (210-900)
South Sudan	1990	Female	25 to 29	43% (28-59%)	57% (41-72%)	0.26 (0.092-0.59)	0.67% (0.26-1.1%)	0.33% (-0.24-0.98%)	2.9 (-1.9-8.3)	530 (200-890)
South Sudan	1990	Female	30 to 34	45% (30-62%)	55% (38-70%)	0.27 (0.095-0.63)	0.68% (0.25-1.2%)	0.45% (-0.15-1.2%)	3.6 (-1.3-9.3)	470 (170-850)
South Sudan	1990	Female	35 to 39	48% (33-63%)	52% (37-67%)	0.25 (0.094-0.56)	0.65% (0.23-1.1%)	0.55% (-0.036-1.3%)	4.3 (-0.27-9.8)	400 (140-690)
South Sudan	1990	Female	40 to 44	50% (35-65%)	50% (35-65%)	0.25 (0.088-0.55)	0.67% (0.29-1.1%)	0.71% (0.18-1.3%)	6.3 (1.5-12)	410 (180-680)
South Sudan	1990	Female	45 to 49	49% (32-66%)	51% (34-68%)	0.26 (0.093-0.58)	0.62% (0.18-1.1%)	0.71% (0.095-1.4%)	7.1 (0.97-15)	370 (110-680)
South Sudan	1990	Female	50 to 54	48% (32-65%)	52% (35-68%)	0.26 (0.09-0.6)	0.64% (0.11-1.3%)	0.79% (0.067-1.7%)	9 (0.76-19)	370 (59-710)
South Sudan	1990	Female	55 to 59	46% (31-63%)	54% (37-69%)	0.27 (0.089-0.63)	0.36% (-0.17-0.99%)	0.46% (-0.22-1.3%)	6.1 (-3-18)	200 (-97-550)
South Sudan	1990	Female	60 to 64	47% (31-64%)	53% (36-69%)	0.27 (0.085-0.63)	0.28% (-0.31-0.8%)	0.37% (-0.35-1.1%)	5.6 (-5.7-16)	140 (-150-440)
South Sudan	1990	Female	65 to 69	48% (33-64%)	52% (36-67%)	0.25 (0.078-0.58)	0.22% (-0.4-0.72%)	0.26% (-0.47-0.89%)	4.2 (-7.6-15)	100 (-170-340)
South Sudan	1990	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.22 (0.067-0.48)	-0.055% (-0.77-0.47%)	-0.1% (-0.93-0.5%)	-1.7 (-15-8.4)	-19 (-270-160)
South Sudan	1990	Female	75 to 79	44% (28-61%)	56% (39-72%)	0.19 (0.058-0.42)	-0.043% (-0.83-0.48%)	-0.054% (-0.98-0.57%)	-0.67 (-13-7.3)	-9 (-180-100)
South Sudan	1990	Female	80 to 84	43% (27-60%)	57% (40-73%)	0.17 (0.056-0.41)	-0.19% (-0.92-0.25%)	-0.22% (-1.1-0.3%)	-1.6 (-7.7-2.2)	-17 (-80-23)
South Sudan	1990	Female	85 to 89	43% (28-59%)	57% (41-72%)	0.17 (0.051-0.38)	-0.23% (-0.83-0.15%)	-0.27% (-0.98-0.17%)	-1.1 (-4-0.69)	-8.1 (-29-5.3)
South Sudan	1990	Female	90 to 94	45% (29-59%)	55% (41-71%)	0.16 (0.053-0.39)	-0.31% (-1-0.1%)	-0.41% (-1.3-0.095%)	-0.59 (-1.9-0.14)	-2.8 (-9.2-0.93)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	1990	Female	95 plus	47% (29-64%)	53% (36-71%)	0.16 (0.048-0.4)	-0.057% (-0.74-0.36%)	-0.13% (-0.95-0.37%)	-0.057 (-0.41-0.16)	-0.099 (-1.3-0.64)
South Sudan	1990	Male	All Ages	41% (35-48%)	59% (52-65%)	0.15 (0.065-0.3)	0.25% (0.1-0.42%)	0.27% (0.0081-0.55%)	140 (4.4-290)	10000 (4000-17000)
South Sudan	1990	Male	15 to 19	41% (29-57%)	59% (43-71%)	0.18 (0.065-0.41)	0.68% (0.37-1.1%)	0.28% (-0.032-0.83%)	2.6 (-0.33-7.3)	670 (360-1100)
South Sudan	1990	Male	20 to 24	52% (38-67%)	48% (33-62%)	0.24 (0.09-0.55)	1.1% (0.57-1.9%)	0.35% (-0.17-1.1%)	4.9 (-2.3-15)	1400 (670-2300)
South Sudan	1990	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.29 (0.11-0.66)	1.3% (0.57-2.2%)	0.56% (-0.32-1.7%)	7 (-3.7-20)	1300 (560-2200)
South Sudan	1990	Male	30 to 34	58% (43-72%)	42% (28-57%)	0.33 (0.12-0.73)	1.3% (0.36-2.5%)	0.72% (-0.48-2.2%)	8.1 (-5.4-24)	1100 (290-2000)
South Sudan	1990	Male	35 to 39	60% (46-74%)	40% (26-54%)	0.35 (0.13-0.75)	1.3% (0.33-2.6%)	1% (-0.28-2.6%)	10 (-3.1-25)	930 (220-1700)
South Sudan	1990	Male	40 to 44	60% (47-75%)	40% (25-53%)	0.36 (0.14-0.78)	1.4% (0.49-2.6%)	1.3% (0.11-2.8%)	15 (1.2-32)	1000 (340-1900)
South Sudan	1990	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.36 (0.13-0.82)	1.5% (0.53-2.6%)	1.5% (0.21-3.1%)	18 (2.6-39)	1000 (340-1900)
South Sudan	1990	Male	50 to 54	61% (45-75%)	39% (25-55%)	0.35 (0.13-0.82)	1.5% (0.41-2.6%)	1.6% (0.18-3.1%)	19 (2.3-40)	880 (240-1600)
South Sudan	1990	Male	55 to 59	60% (46-75%)	40% (25-54%)	0.34 (0.12-0.79)	1.3% (0.3-2.3%)	1.4% (0.1-2.7%)	18 (1.3-36)	680 (160-1300)
South Sudan	1990	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.31 (0.11-0.74)	1.1% (0.33-2%)	1.2% (0.21-2.4%)	17 (2.7-33)	540 (150-980)
South Sudan	1990	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.26 (0.088-0.59)	0.64% (0.042-1.2%)	0.68% (-0.077-1.4%)	9.7 (-0.95-21)	250 (16-500)
South Sudan	1990	Male	70 to 74	56% (41-70%)	44% (30-59%)	0.21 (0.069-0.48)	0.47% (-0.11-0.95%)	0.52% (-0.18-1.1%)	7 (-2.6-15)	140 (-32-290)
South Sudan	1990	Male	75 to 79	55% (41-70%)	45% (30-59%)	0.18 (0.058-0.4)	0.39% (-0.13-0.8%)	0.43% (-0.21-0.95%)	4.4 (-2.2-9.7)	67 (-22-140)
South Sudan	1990	Male	80 to 84	53% (36-69%)	47% (31-64%)	0.16 (0.055-0.37)	0.25% (-0.16-0.56%)	0.25% (-0.24-0.61%)	1.3 (-1.3-3.2)	16 (-9.9-37)
South Sudan	1990	Male	85 to 89	51% (35-65%)	49% (35-65%)	0.14 (0.047-0.32)	0.25% (-0.11-0.5%)	0.22% (-0.2-0.54%)	0.66 (-0.6-1.6)	6.4 (-2.9-13)
South Sudan	1990	Male	90 to 94	51% (36-66%)	49% (34-64%)	0.13 (0.044-0.3)	0.16% (-0.21-0.41%)	0.092% (-0.39-0.39%)	0.1 (-0.43-0.43)	1 (-1.5-2.7)
South Sudan	1990	Male	95 plus	53% (34-71%)	47% (29-66%)	0.12 (0.037-0.29)	0.24% (-0.15-0.5%)	0.15% (-0.34-0.5%)	0.046 (-0.1-0.15)	0.3 (-0.18-0.63)
South Sudan	1995	Female	All Ages	32% (27-39%)	68% (61-73%)	0.12 (0.05-0.25)	0.12% (0.028-0.22%)	0.11% (-0.12-0.3%)	44 (-46-120)	3600 (790-6500)
South Sudan	1995	Female	15 to 19	35% (22-50%)	65% (50-78%)	0.15 (0.05-0.34)	0.33% (0.072-0.66%)	0.16% (-0.25-0.62%)	0.95 (-1.5-3.9)	250 (54-490)
South Sudan	1995	Female	20 to 24	41% (26-57%)	59% (43-74%)	0.21 (0.074-0.47)	0.56% (0.21-0.95%)	0.17% (-0.29-0.68%)	1.4 (-2.4-5.7)	470 (170-820)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	1995	Female	25 to 29	43% (28-59%)	57% (41-72%)	0.25 (0.089-0.57)	0.64% (0.22-1.2%)	0.31% (-0.28-0.97%)	2.7 (-2.3-8.3)	520 (170-920)
South Sudan	1995	Female	30 to 34	45% (30-61%)	55% (39-70%)	0.26 (0.095-0.61)	0.63% (0.21-1.1%)	0.4% (-0.18-1.1%)	3.5 (-1.6-9.5)	460 (150-840)
South Sudan	1995	Female	35 to 39	48% (33-63%)	52% (37-67%)	0.25 (0.092-0.54)	0.62% (0.19-1.2%)	0.52% (-0.13-1.3%)	4.4 (-1.1-11)	400 (120-730)
South Sudan	1995	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.25 (0.086-0.56)	0.65% (0.25-1.1%)	0.68% (0.088-1.3%)	5.4 (0.62-11)	350 (130-590)
South Sudan	1995	Female	45 to 49	49% (32-66%)	51% (34-68%)	0.25 (0.093-0.58)	0.61% (0.15-1.1%)	0.69% (0.052-1.4%)	6.6 (0.43-14)	340 (81-650)
South Sudan	1995	Female	50 to 54	48% (32-64%)	52% (36-68%)	0.26 (0.089-0.62)	0.64% (0.052-1.3%)	0.78% (0.0051-1.7%)	9.1 (0.055-20)	370 (27-760)
South Sudan	1995	Female	55 to 59	46% (31-63%)	54% (37-69%)	0.26 (0.086-0.64)	0.35% (-0.24-0.96%)	0.44% (-0.34-1.3%)	5.8 (-4.1-19)	190 (-130-580)
South Sudan	1995	Female	60 to 64	47% (31-65%)	53% (35-69%)	0.26 (0.087-0.61)	0.29% (-0.29-0.86%)	0.38% (-0.34-1.1%)	5.9 (-5.1-17)	150 (-150-460)
South Sudan	1995	Female	65 to 69	48% (33-64%)	52% (36-67%)	0.24 (0.081-0.55)	0.24% (-0.41-0.78%)	0.28% (-0.48-0.95%)	4.7 (-8.2-16)	110 (-180-370)
South Sudan	1995	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.21 (0.067-0.49)	-0.055% (-0.77-0.48%)	-0.1% (-0.95-0.57%)	-1.7 (-16-9)	-19 (-290-170)
South Sudan	1995	Female	75 to 79	45% (29-61%)	55% (39-71%)	0.19 (0.06-0.44)	-0.038% (-0.86-0.52%)	-0.047% (-1-0.62%)	-0.63 (-14-8)	-8.4 (-190-120)
South Sudan	1995	Female	80 to 84	44% (28-61%)	56% (39-72%)	0.17 (0.059-0.4)	-0.21% (-0.99-0.28%)	-0.24% (-1.2-0.33%)	-2 (-9.5-2.7)	-21 (-100-28)
South Sudan	1995	Female	85 to 89	44% (28-58%)	56% (42-72%)	0.17 (0.053-0.4)	-0.26% (-0.96-0.17%)	-0.31% (-1.1-0.2%)	-1.5 (-5.5-1)	-11 (-41-7.6)
South Sudan	1995	Female	90 to 94	45% (29-60%)	55% (40-71%)	0.16 (0.054-0.38)	-0.31% (-1-0.11%)	-0.41% (-1.2-0.1%)	-0.75 (-2.3-0.19)	-3.5 (-11-1.2)
South Sudan	1995	Female	95 plus	47% (30-65%)	53% (35-70%)	0.16 (0.049-0.37)	-0.05% (-0.73-0.38%)	-0.12% (-0.97-0.41%)	-0.063 (-0.5-0.21)	-0.1 (-1.5-0.81)
South Sudan	1995	Male	All Ages	40% (34-48%)	60% (52-66%)	0.15 (0.059-0.31)	0.29% (0.099-0.5%)	0.33% (-0.0041-0.72%)	150 (-2-320)	9900 (3400-17000)
South Sudan	1995	Male	15 to 19	42% (29-58%)	58% (42-71%)	0.18 (0.061-0.41)	0.83% (0.44-1.3%)	0.39% (-0.078-1.1%)	2.6 (-0.45-7.4)	630 (320-1000)
South Sudan	1995	Male	20 to 24	52% (38-66%)	48% (34-62%)	0.24 (0.084-0.53)	1.4% (0.64-2.4%)	0.52% (-0.33-1.6%)	5.1 (-3.4-16)	1300 (570-2100)
South Sudan	1995	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.29 (0.11-0.63)	1.5% (0.55-2.6%)	0.69% (-0.49-2%)	7.5 (-5-22)	1300 (490-2300)
South Sudan	1995	Male	30 to 34	58% (42-73%)	42% (27-58%)	0.32 (0.12-0.69)	1.3% (0.28-2.5%)	0.75% (-0.6-2.2%)	8.7 (-6.7-26)	1100 (230-2100)
South Sudan	1995	Male	35 to 39	60% (46-74%)	40% (26-54%)	0.34 (0.13-0.71)	1.4% (0.31-2.7%)	1.1% (-0.34-2.7%)	11 (-3.3-31)	990 (210-2000)
South Sudan	1995	Male	40 to 44	60% (46-76%)	40% (24-54%)	0.35 (0.13-0.77)	1.4% (0.37-2.5%)	1.3% (-0.053-2.8%)	13 (-0.45-31)	850 (230-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	1995	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.35 (0.14-0.8)	1.5% (0.43-2.7%)	1.5% (0.095-3.1%)	18 (1.2-39)	940 (270-1800)
South Sudan	1995	Male	50 to 54	61% (46-76%)	39% (24-54%)	0.34 (0.12-0.81)	1.5% (0.33-2.6%)	1.6% (0.094-3.1%)	20 (1.3-42)	890 (200-1700)
South Sudan	1995	Male	55 to 59	61% (45-75%)	39% (25-55%)	0.34 (0.12-0.79)	1.3% (0.22-2.3%)	1.4% (0.0096-2.8%)	18 (0.11-38)	690 (120-1300)
South Sudan	1995	Male	60 to 64	60% (44-75%)	40% (25-56%)	0.31 (0.11-0.72)	1.2% (0.3-2.1%)	1.3% (0.19-2.5%)	18 (2.4-36)	560 (130-1000)
South Sudan	1995	Male	65 to 69	58% (43-75%)	42% (25-57%)	0.26 (0.086-0.57)	0.71% (0.017-1.4%)	0.77% (-0.089-1.7%)	11 (-0.98-23)	280 (7.1-550)
South Sudan	1995	Male	70 to 74	56% (42-72%)	44% (28-58%)	0.21 (0.069-0.46)	0.5% (-0.12-1%)	0.56% (-0.2-1.2%)	7.7 (-2.2-17)	150 (-33-310)
South Sudan	1995	Male	75 to 79	55% (41-71%)	45% (29-59%)	0.17 (0.058-0.39)	0.44% (-0.13-0.9%)	0.49% (-0.19-1.1%)	5.1 (-2.1-11)	78 (-24-160)
South Sudan	1995	Male	80 to 84	53% (36-70%)	47% (30-64%)	0.16 (0.056-0.35)	0.29% (-0.16-0.64%)	0.29% (-0.25-0.7%)	1.7 (-1.4-4)	20 (-11-45)
South Sudan	1995	Male	85 to 89	51% (35-65%)	49% (35-65%)	0.14 (0.048-0.31)	0.3% (-0.097-0.58%)	0.29% (-0.18-0.63%)	1 (-0.64-2.3)	9.2 (-2.9-18)
South Sudan	1995	Male	90 to 94	51% (36-66%)	49% (34-64%)	0.13 (0.044-0.29)	0.17% (-0.19-0.42%)	0.12% (-0.34-0.43%)	0.16 (-0.46-0.58)	1.4 (-1.6-3.5)
South Sudan	1995	Male	95 plus	53% (34-72%)	47% (28-66%)	0.12 (0.036-0.28)	0.28% (-0.12-0.55%)	0.21% (-0.29-0.55%)	0.071 (-0.1-0.19)	0.4 (-0.17-0.8)
South Sudan	2000	Female	All Ages	32% (27-39%)	68% (61-73%)	0.12 (0.05-0.25)	0.13% (-0.00036-0.26%)	0.11% (-0.17-0.37%)	48 (-76-170)	4100 (-11-8300)
South Sudan	2000	Female	15 to 19	34% (22-50%)	66% (50-78%)	0.15 (0.05-0.34)	0.33% (0.054-0.63%)	0.13% (-0.31-0.64%)	0.91 (-2.1-4.4)	280 (48-560)
South Sudan	2000	Female	20 to 24	41% (26-57%)	59% (43-74%)	0.2 (0.07-0.46)	0.56% (0.17-1%)	0.14% (-0.36-0.69%)	1.4 (-3.4-6.3)	550 (160-1000)
South Sudan	2000	Female	25 to 29	43% (28-59%)	57% (41-72%)	0.25 (0.086-0.56)	0.56% (0.15-1%)	0.23% (-0.3-0.83%)	2.8 (-3.3-9.8)	590 (160-1100)
South Sudan	2000	Female	30 to 34	45% (30-61%)	55% (39-70%)	0.26 (0.091-0.6)	0.54% (0.12-1%)	0.3% (-0.24-0.9%)	3.8 (-3.1-12)	540 (130-1000)
South Sudan	2000	Female	35 to 39	48% (33-63%)	52% (37-67%)	0.24 (0.084-0.55)	0.52% (0.061-1%)	0.38% (-0.25-1.1%)	4.9 (-3.1-14)	470 (60-930)
South Sudan	2000	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.24 (0.08-0.54)	0.56% (0.15-1.1%)	0.54% (-0.036-1.2%)	6.3 (-0.45-14)	420 (110-790)
South Sudan	2000	Female	45 to 49	49% (32-65%)	51% (35-68%)	0.25 (0.088-0.56)	0.53% (-0.015-1.1%)	0.57% (-0.17-1.4%)	6.4 (-1.7-16)	340 (-9.3-710)
South Sudan	2000	Female	50 to 54	47% (32-65%)	53% (35-68%)	0.25 (0.088-0.57)	0.57% (-0.14-1.3%)	0.68% (-0.26-1.7%)	9.6 (-4.1-25)	390 (-96-910)
South Sudan	2000	Female	55 to 59	46% (31-63%)	54% (37-69%)	0.26 (0.089-0.61)	0.32% (-0.38-1.1%)	0.4% (-0.53-1.4%)	6.8 (-8.9-24)	220 (-280-730)
South Sudan	2000	Female	60 to 64	47% (31-63%)	53% (37-69%)	0.26 (0.086-0.59)	0.28% (-0.36-0.93%)	0.36% (-0.45-1.2%)	6.9 (-8.3-24)	180 (-230-610)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	2000	Female	65 to 69	48% (33-64%)	52% (36-67%)	0.24 (0.079-0.54)	0.24% (-0.45-0.84%)	0.29% (-0.51-1%)	6.2 (-11-23)	140 (-250-510)
South Sudan	2000	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.21 (0.066-0.51)	-0.051% (-0.85-0.56%)	-0.094% (-1-0.65%)	-2 (-23-14)	-23 (-400-250)
South Sudan	2000	Female	75 to 79	45% (29-61%)	55% (39-71%)	0.18 (0.057-0.44)	-0.024% (-0.89-0.56%)	-0.028% (-1-0.65%)	-0.49 (-18-11)	-6.9 (-260-160)
South Sudan	2000	Female	80 to 84	44% (27-61%)	56% (39-73%)	0.17 (0.054-0.39)	-0.19% (-0.96-0.29%)	-0.22% (-1.1-0.36%)	-2.3 (-12-3.7)	-25 (-120-38)
South Sudan	2000	Female	85 to 89	43% (28-58%)	57% (42-72%)	0.16 (0.05-0.39)	-0.23% (-0.91-0.2%)	-0.28% (-1.1-0.24%)	-1.7 (-6.4-1.4)	-12 (-47-10)
South Sudan	2000	Female	90 to 94	45% (29-60%)	55% (40-71%)	0.16 (0.05-0.38)	-0.29% (-1-0.14%)	-0.38% (-1.2-0.14%)	-0.95 (-3.1-0.34)	-4.4 (-15-2)
South Sudan	2000	Female	95 plus	47% (29-65%)	53% (35-71%)	0.15 (0.046-0.37)	-0.0061% (-0.67-0.44%)	-0.076% (-0.9-0.45%)	-0.054 (-0.62-0.32)	-0.016 (-1.9-1.3)
South Sudan	2000	Male	All Ages	41% (35-48%)	59% (52-65%)	0.14 (0.06-0.31)	0.33% (0.091-0.59%)	0.35% (-0.078-0.82%)	170 (-38-410)	12000 (3200-21000)
South Sudan	2000	Male	15 to 19	41% (29-57%)	59% (43-71%)	0.18 (0.06-0.41)	0.82% (0.4-1.4%)	0.34% (-0.16-1.1%)	2.8 (-1.4-8.8)	750 (350-1300)
South Sudan	2000	Male	20 to 24	52% (39-66%)	48% (34-61%)	0.24 (0.084-0.52)	1.4% (0.58-2.4%)	0.47% (-0.42-1.6%)	5.7 (-4.9-20)	1600 (640-2800)
South Sudan	2000	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.28 (0.099-0.62)	1.4% (0.48-2.5%)	0.59% (-0.53-1.8%)	8.4 (-6.8-27)	1600 (530-2900)
South Sudan	2000	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.32 (0.11-0.69)	1.2% (0.16-2.4%)	0.66% (-0.65-2.1%)	10 (-10-34)	1400 (180-2800)
South Sudan	2000	Male	35 to 39	59% (45-74%)	41% (26-55%)	0.34 (0.12-0.72)	1.2% (0.18-2.4%)	0.85% (-0.46-2.5%)	14 (-7.5-41)	1200 (180-2600)
South Sudan	2000	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.34 (0.12-0.76)	1.2% (0.21-2.5%)	1.1% (-0.19-2.7%)	16 (-3.1-41)	1100 (180-2200)
South Sudan	2000	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.34 (0.13-0.77)	1.3% (0.22-2.5%)	1.3% (-0.11-2.8%)	17 (-1.7-40)	930 (160-1900)
South Sudan	2000	Male	50 to 54	61% (45-76%)	39% (24-55%)	0.34 (0.12-0.77)	1.4% (0.14-2.7%)	1.4% (-0.2-3.1%)	22 (-3-51)	990 (93-2000)
South Sudan	2000	Male	55 to 59	61% (45-75%)	39% (25-55%)	0.33 (0.12-0.76)	1.2% (0.038-2.4%)	1.3% (-0.22-2.9%)	22 (-3.7-50)	840 (26-1700)
South Sudan	2000	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.3 (0.11-0.7)	1.2% (0.21-2.2%)	1.2% (0.009-2.6%)	22 (0.16-48)	700 (110-1400)
South Sudan	2000	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.25 (0.085-0.56)	0.68% (-0.1-1.5%)	0.73% (-0.24-1.8%)	13 (-3.7-32)	340 (-51-760)
South Sudan	2000	Male	70 to 74	56% (42-70%)	44% (30-58%)	0.2 (0.07-0.49)	0.49% (-0.19-1.1%)	0.54% (-0.28-1.3%)	9.5 (-4.9-24)	190 (-71-440)
South Sudan	2000	Male	75 to 79	55% (40-70%)	45% (30-60%)	0.17 (0.056-0.38)	0.44% (-0.16-0.95%)	0.48% (-0.24-1.1%)	6.8 (-3.3-16)	100 (-36-230)
South Sudan	2000	Male	80 to 84	53% (36-70%)	47% (30-64%)	0.15 (0.054-0.34)	0.29% (-0.19-0.71%)	0.3% (-0.27-0.79%)	2.3 (-2-6)	28 (-17-66)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	2000	Male	85 to 89	51% (35-65%)	49% (35-65%)	0.13 (0.047-0.32)	0.3% (-0.12-0.62%)	0.3% (-0.21-0.67%)	1.3 (-0.91-2.9)	12 (-4.6-24)
South Sudan	2000	Male	90 to 94	51% (36-66%)	49% (34-64%)	0.12 (0.043-0.31)	0.19% (-0.18-0.45%)	0.14% (-0.32-0.47%)	0.26 (-0.57-0.84)	2.2 (-2-5.1)
South Sudan	2000	Male	95 plus	53% (33-71%)	47% (29-67%)	0.12 (0.034-0.28)	0.31% (-0.12-0.6%)	0.24% (-0.32-0.6%)	0.12 (-0.15-0.29)	0.61 (-0.24-1.2)
South Sudan	2005	Female	All Ages	32% (27-39%)	68% (61-73%)	0.12 (0.049-0.26)	0.14% (-0.012-0.3%)	0.11% (-0.23-0.43%)	53 (-110-220)	4800 (-370-10000)
South Sudan	2005	Female	15 to 19	34% (22-50%)	66% (50-78%)	0.16 (0.051-0.36)	0.37% (0.043-0.75%)	0.15% (-0.36-0.78%)	1.2 (-2.7-6.1)	360 (40-730)
South Sudan	2005	Female	20 to 24	41% (26-57%)	59% (43-74%)	0.22 (0.07-0.5)	0.63% (0.21-1.1%)	0.17% (-0.43-0.78%)	1.7 (-4.1-7.8)	660 (200-1200)
South Sudan	2005	Female	25 to 29	43% (27-60%)	57% (40-73%)	0.26 (0.091-0.58)	0.6% (0.16-1.1%)	0.25% (-0.32-0.89%)	3.3 (-4.1-11)	710 (190-1300)
South Sudan	2005	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.27 (0.095-0.63)	0.52% (0.12-0.98%)	0.28% (-0.27-0.91%)	4.4 (-4.4-14)	640 (140-1200)
South Sudan	2005	Female	35 to 39	48% (32-64%)	52% (36-68%)	0.25 (0.088-0.57)	0.5% (0.022-1%)	0.35% (-0.29-1.1%)	5.8 (-4.6-17)	570 (29-1200)
South Sudan	2005	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.25 (0.083-0.58)	0.51% (0.056-1%)	0.46% (-0.13-1.1%)	7.4 (-2.1-18)	510 (53-990)
South Sudan	2005	Female	45 to 49	49% (32-66%)	51% (34-68%)	0.26 (0.089-0.59)	0.51% (-0.095-1.1%)	0.54% (-0.25-1.4%)	7.9 (-3.7-20)	420 (-78-930)
South Sudan	2005	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.26 (0.094-0.62)	0.55% (-0.25-1.4%)	0.66% (-0.4-1.8%)	9.7 (-6.2-27)	400 (-180-1000)
South Sudan	2005	Female	55 to 59	46% (31-62%)	54% (38-69%)	0.27 (0.093-0.64)	0.31% (-0.52-1.2%)	0.39% (-0.72-1.6%)	7.2 (-13-30)	230 (-390-920)
South Sudan	2005	Female	60 to 64	47% (30-63%)	53% (37-70%)	0.27 (0.092-0.67)	0.25% (-0.52-1%)	0.34% (-0.61-1.4%)	7.5 (-13-31)	190 (-380-770)
South Sudan	2005	Female	65 to 69	48% (32-64%)	52% (36-68%)	0.25 (0.082-0.6)	0.23% (-0.56-0.92%)	0.28% (-0.7-1.2%)	7 (-17-29)	160 (-370-640)
South Sudan	2005	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.22 (0.069-0.57)	-0.051% (-0.87-0.61%)	-0.096% (-1.1-0.7%)	-2.4 (-28-18)	-28 (-480-340)
South Sudan	2005	Female	75 to 79	45% (28-61%)	55% (39-72%)	0.19 (0.058-0.49)	-0.031% (-0.95-0.6%)	-0.035% (-1.1-0.71%)	-0.73 (-24-16)	-11 (-340-220)
South Sudan	2005	Female	80 to 84	43% (27-61%)	57% (39-73%)	0.18 (0.054-0.43)	-0.2% (-0.98-0.35%)	-0.23% (-1.1-0.42%)	-3.4 (-17-6.3)	-37 (-180-64)
South Sudan	2005	Female	85 to 89	43% (27-58%)	57% (42-73%)	0.17 (0.052-0.43)	-0.24% (-0.98-0.22%)	-0.29% (-1.2-0.26%)	-2 (-7.9-1.9)	-15 (-60-14)
South Sudan	2005	Female	90 to 94	45% (29-60%)	55% (40-71%)	0.17 (0.051-0.39)	-0.3% (-1.1-0.17%)	-0.4% (-1.3-0.2%)	-1.1 (-3.5-0.52)	-5 (-17-2.9)
South Sudan	2005	Female	95 plus	47% (30-65%)	53% (35-70%)	0.16 (0.045-0.42)	-0.0015% (-0.67-0.46%)	-0.067% (-0.9-0.49%)	-0.058 (-0.79-0.43)	-0.0013 (-2.5-1.6)
South Sudan	2005	Male	All Ages	40% (34-48%)	60% (52-66%)	0.15 (0.059-0.32)	0.38% (0.079-0.68%)	0.4% (-0.12-0.93%)	210 (-60-510)	14000 (3100-26000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	2005	Male	15 to 19	41% (28-57%)	59% (43-72%)	0.18 (0.062-0.42)	0.96% (0.43-1.6%)	0.44% (-0.23-1.4%)	3.6 (-1.7-11)	950 (420-1600)
South Sudan	2005	Male	20 to 24	52% (38-66%)	48% (34-62%)	0.24 (0.08-0.57)	1.8% (0.75-3.1%)	0.64% (-0.68-2.2%)	6.9 (-7-24)	1900 (740-3300)
South Sudan	2005	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.29 (0.11-0.67)	1.7% (0.51-3%)	0.77% (-0.74-2.4%)	11 (-10-34)	1900 (540-3500)
South Sudan	2005	Male	30 to 34	58% (42-72%)	42% (28-58%)	0.33 (0.12-0.73)	1.4% (0.17-2.6%)	0.76% (-0.7-2.4%)	13 (-13-41)	1700 (230-3400)
South Sudan	2005	Male	35 to 39	59% (46-74%)	41% (26-54%)	0.35 (0.12-0.79)	1.3% (0.18-2.7%)	0.94% (-0.51-2.7%)	17 (-9.5-50)	1600 (200-3300)
South Sudan	2005	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.36 (0.12-0.84)	1.3% (0.17-2.6%)	1.1% (-0.21-2.8%)	21 (-4-53)	1400 (180-2800)
South Sudan	2005	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.35 (0.12-0.79)	1.3% (0.15-2.6%)	1.3% (-0.25-3.1%)	23 (-4.6-53)	1200 (130-2400)
South Sudan	2005	Male	50 to 54	61% (45-75%)	39% (25-55%)	0.35 (0.13-0.84)	1.4% (-0.069-2.8%)	1.4% (-0.51-3.3%)	22 (-7.2-53)	1000 (-58-2100)
South Sudan	2005	Male	55 to 59	60% (45-75%)	40% (25-55%)	0.34 (0.12-0.81)	1.3% (-0.013-2.7%)	1.3% (-0.35-3.2%)	25 (-6.2-61)	950 (-8.1-2100)
South Sudan	2005	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.31 (0.11-0.78)	1.2% (0.084-2.3%)	1.3% (-0.095-2.7%)	28 (-1.9-58)	860 (63-1700)
South Sudan	2005	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.26 (0.087-0.62)	0.7% (-0.14-1.6%)	0.75% (-0.31-1.8%)	17 (-6.1-41)	430 (-85-970)
South Sudan	2005	Male	70 to 74	56% (41-70%)	44% (30-59%)	0.21 (0.071-0.52)	0.5% (-0.29-1.2%)	0.55% (-0.38-1.4%)	12 (-8.6-31)	230 (-130-560)
South Sudan	2005	Male	75 to 79	54% (40-70%)	46% (30-60%)	0.17 (0.056-0.43)	0.45% (-0.23-1%)	0.49% (-0.34-1.2%)	8.7 (-5.8-21)	130 (-66-300)
South Sudan	2005	Male	80 to 84	52% (36-69%)	48% (31-64%)	0.16 (0.051-0.38)	0.3% (-0.25-0.79%)	0.31% (-0.36-0.88%)	3.3 (-4-9.5)	41 (-34-110)
South Sudan	2005	Male	85 to 89	50% (35-64%)	50% (36-65%)	0.14 (0.045-0.36)	0.31% (-0.17-0.66%)	0.3% (-0.29-0.73%)	1.6 (-1.6-3.9)	14 (-7.7-31)
South Sudan	2005	Male	90 to 94	51% (35-66%)	49% (34-65%)	0.13 (0.042-0.31)	0.2% (-0.21-0.51%)	0.15% (-0.34-0.52%)	0.31 (-0.71-1.1)	2.6 (-2.7-6.3)
South Sudan	2005	Male	95 plus	52% (32-71%)	48% (29-68%)	0.12 (0.035-0.31)	0.32% (-0.15-0.67%)	0.25% (-0.35-0.68%)	0.15 (-0.21-0.41)	0.78 (-0.38-1.6)
South Sudan	2010	Female	All Ages	33% (27-39%)	67% (61-73%)	0.13 (0.054-0.28)	0.16% (-0.027-0.34%)	0.11% (-0.28-0.47%)	63 (-160-280)	6000 (-1000-13000)
South Sudan	2010	Female	15 to 19	35% (22-50%)	65% (50-78%)	0.17 (0.051-0.38)	0.4% (0.029-0.78%)	0.18% (-0.39-0.86%)	1.7 (-4-7.7)	480 (32-980)
South Sudan	2010	Female	20 to 24	41% (26-57%)	59% (43-74%)	0.23 (0.071-0.52)	0.7% (0.2-1.2%)	0.2% (-0.46-0.89%)	2.3 (-5.4-10)	880 (250-1600)
South Sudan	2010	Female	25 to 29	43% (27-59%)	57% (41-73%)	0.27 (0.092-0.6)	0.67% (0.18-1.2%)	0.29% (-0.35-1%)	4.3 (-5.6-15)	890 (260-1600)
South Sudan	2010	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.28 (0.099-0.66)	0.57% (0.12-1.1%)	0.31% (-0.26-1.1%)	5.7 (-4.8-19)	810 (170-1600)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	2010	Female	35 to 39	48% (33-63%)	52% (37-67%)	0.27 (0.09-0.6)	0.51% (0.00098-1.1%)	0.36% (-0.3-1.1%)	7.1 (-5.9-23)	700 (1.5-1500)
South Sudan	2010	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.26 (0.085-0.59)	0.54% (0.025-1.1%)	0.49% (-0.16-1.2%)	9.6 (-3.3-24)	660 (33-1300)
South Sudan	2010	Female	45 to 49	49% (32-65%)	51% (35-68%)	0.27 (0.091-0.63)	0.52% (-0.091-1.2%)	0.55% (-0.27-1.5%)	10 (-4.9-29)	550 (-96-1300)
South Sudan	2010	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.27 (0.095-0.64)	0.59% (-0.28-1.6%)	0.7% (-0.41-2%)	13 (-8.8-38)	530 (-270-1400)
South Sudan	2010	Female	55 to 59	46% (31-62%)	54% (38-69%)	0.28 (0.096-0.66)	0.32% (-0.59-1.4%)	0.41% (-0.8-1.9%)	7.9 (-16-35)	250 (-480-1100)
South Sudan	2010	Female	60 to 64	47% (31-64%)	53% (36-69%)	0.28 (0.097-0.66)	0.25% (-0.56-1.1%)	0.34% (-0.68-1.5%)	8.4 (-18-36)	210 (-500-920)
South Sudan	2010	Female	65 to 69	48% (33-64%)	52% (36-67%)	0.26 (0.091-0.62)	0.22% (-0.59-0.94%)	0.26% (-0.71-1.2%)	7.9 (-22-37)	180 (-520-800)
South Sudan	2010	Female	70 to 74	47% (33-62%)	53% (38-67%)	0.23 (0.075-0.57)	-0.082% (-0.94-0.62%)	-0.13% (-1.2-0.73%)	-4 (-36-23)	-56 (-640-410)
South Sudan	2010	Female	75 to 79	45% (30-61%)	55% (39-70%)	0.2 (0.065-0.51)	-0.056% (-1-0.62%)	-0.062% (-1.2-0.74%)	-1.7 (-35-21)	-26 (-490-290)
South Sudan	2010	Female	80 to 84	44% (27-60%)	56% (40-73%)	0.19 (0.061-0.47)	-0.23% (-1-0.35%)	-0.26% (-1.2-0.41%)	-5.3 (-24-8.6)	-58 (-260-87)
South Sudan	2010	Female	85 to 89	43% (27-58%)	57% (42-73%)	0.18 (0.054-0.44)	-0.27% (-1-0.24%)	-0.32% (-1.2-0.3%)	-3.1 (-12-3)	-23 (-88-21)
South Sudan	2010	Female	90 to 94	45% (30-59%)	55% (41-70%)	0.17 (0.054-0.41)	-0.33% (-1.1-0.16%)	-0.43% (-1.4-0.17%)	-1.4 (-4.4-0.51)	-6.5 (-22-3.3)
South Sudan	2010	Female	95 plus	47% (29-65%)	53% (35-71%)	0.17 (0.047-0.43)	-0.01% (-0.7-0.48%)	-0.083% (-0.93-0.54%)	-0.075 (-0.83-0.49)	-0.033 (-2.5-1.8)
South Sudan	2010	Male	All Ages	40% (34-48%)	60% (52-66%)	0.16 (0.063-0.33)	0.44% (0.092-0.8%)	0.45% (-0.14-1.1%)	280 (-79-660)	19000 (4300-34000)
South Sudan	2010	Male	15 to 19	40% (27-55%)	60% (45-73%)	0.19 (0.058-0.44)	1% (0.46-1.7%)	0.47% (-0.26-1.5%)	5 (-2.9-15)	1300 (550-2200)
South Sudan	2010	Male	20 to 24	51% (37-66%)	49% (34-63%)	0.25 (0.081-0.57)	1.9% (0.77-3.3%)	0.73% (-0.74-2.4%)	9.9 (-10-35)	2500 (920-4500)
South Sudan	2010	Male	25 to 29	55% (39-70%)	45% (30-61%)	0.3 (0.11-0.69)	1.9% (0.52-3.4%)	0.92% (-0.92-2.9%)	14 (-14-46)	2500 (690-4500)
South Sudan	2010	Male	30 to 34	58% (43-73%)	42% (27-57%)	0.34 (0.13-0.77)	1.6% (0.26-3.1%)	0.93% (-0.82-2.8%)	17 (-16-55)	2200 (360-4300)
South Sudan	2010	Male	35 to 39	59% (45-74%)	41% (26-55%)	0.37 (0.13-0.83)	1.4% (0.18-2.9%)	1.1% (-0.59-2.9%)	23 (-12-63)	2000 (250-4100)
South Sudan	2010	Male	40 to 44	60% (46-75%)	40% (25-54%)	0.38 (0.12-0.86)	1.4% (0.2-2.8%)	1.3% (-0.25-3.1%)	29 (-5.9-71)	1900 (260-3700)
South Sudan	2010	Male	45 to 49	60% (45-75%)	40% (25-55%)	0.37 (0.13-0.83)	1.4% (0.13-2.9%)	1.4% (-0.24-3.2%)	31 (-5.6-71)	1600 (150-3300)
South Sudan	2010	Male	50 to 54	60% (45-74%)	40% (26-55%)	0.37 (0.13-0.83)	1.5% (-0.095-3.1%)	1.5% (-0.47-3.6%)	31 (-10-74)	1400 (-86-2900)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	2010	Male	55 to 59	60% (45-75%)	40% (25-55%)	0.36 (0.13-0.81)	1.4% (-0.098-3%)	1.4% (-0.45-3.6%)	28 (-9-69)	1100 (-78-2300)
South Sudan	2010	Male	60 to 64	59% (43-76%)	41% (24-57%)	0.32 (0.11-0.73)	1.3% (0.15-2.5%)	1.4% (-0.072-2.9%)	34 (-1.9-72)	1000 (110-2100)
South Sudan	2010	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.27 (0.096-0.62)	0.73% (-0.2-1.6%)	0.78% (-0.35-1.9%)	22 (-11-54)	560 (-150-1300)
South Sudan	2010	Male	70 to 74	56% (41-70%)	44% (30-59%)	0.22 (0.076-0.54)	0.51% (-0.29-1.2%)	0.57% (-0.41-1.5%)	16 (-11-40)	310 (-170-750)
South Sudan	2010	Male	75 to 79	54% (39-70%)	46% (30-61%)	0.18 (0.064-0.46)	0.45% (-0.32-1.1%)	0.5% (-0.43-1.2%)	12 (-10-29)	170 (-110-410)
South Sudan	2010	Male	80 to 84	52% (35-69%)	48% (31-65%)	0.16 (0.057-0.39)	0.31% (-0.27-0.82%)	0.31% (-0.39-0.93%)	4.6 (-5.7-14)	57 (-51-150)
South Sudan	2010	Male	85 to 89	50% (34-64%)	50% (36-66%)	0.14 (0.049-0.34)	0.31% (-0.18-0.71%)	0.3% (-0.31-0.78%)	2.2 (-2.2-5.8)	20 (-12-46)
South Sudan	2010	Male	90 to 94	50% (35-67%)	50% (33-65%)	0.13 (0.045-0.32)	0.2% (-0.23-0.53%)	0.15% (-0.37-0.55%)	0.37 (-0.95-1.4)	3.1 (-3.4-8.3)
South Sudan	2010	Male	95 plus	52% (33-71%)	48% (29-67%)	0.13 (0.035-0.33)	0.32% (-0.18-0.73%)	0.24% (-0.38-0.75%)	0.16 (-0.25-0.49)	0.85 (-0.47-2)
South Sudan	2016	Female	All Ages	33% (27-39%)	67% (61-73%)	0.14 (0.057-0.3)	0.18% (-0.032-0.41%)	0.12% (-0.34-0.57%)	80 (-240-400)	8100 (-1500-18000)
South Sudan	2016	Female	15 to 19	34% (22-51%)	66% (49-78%)	0.17 (0.057-0.4)	0.41% (0.05-0.8%)	0.19% (-0.39-0.84%)	2.2 (-4.7-10)	620 (68-1300)
South Sudan	2016	Female	20 to 24	41% (26-57%)	59% (43-74%)	0.24 (0.079-0.52)	0.76% (0.25-1.3%)	0.23% (-0.47-0.99%)	3.2 (-6.9-14)	1200 (400-2100)
South Sudan	2016	Female	25 to 29	43% (28-59%)	57% (41-72%)	0.28 (0.1-0.63)	0.77% (0.25-1.4%)	0.35% (-0.36-1.2%)	6 (-6.6-20)	1200 (400-2300)
South Sudan	2016	Female	30 to 34	45% (29-61%)	55% (39-71%)	0.3 (0.1-0.7)	0.68% (0.11-1.3%)	0.4% (-0.34-1.3%)	7.8 (-6.6-24)	1100 (170-2100)
South Sudan	2016	Female	35 to 39	48% (32-63%)	52% (37-68%)	0.28 (0.097-0.67)	0.62% (0.034-1.3%)	0.47% (-0.32-1.4%)	9.8 (-6.3-29)	940 (50-2000)
South Sudan	2016	Female	40 to 44	49% (35-65%)	51% (35-65%)	0.28 (0.096-0.66)	0.64% (0.072-1.3%)	0.61% (-0.13-1.5%)	13 (-2.7-32)	880 (92-1700)
South Sudan	2016	Female	45 to 49	49% (32-64%)	51% (36-68%)	0.29 (0.098-0.64)	0.62% (-0.14-1.4%)	0.67% (-0.37-1.9%)	15 (-7.6-42)	780 (-160-1900)
South Sudan	2016	Female	50 to 54	47% (32-64%)	53% (36-68%)	0.29 (0.1-0.66)	0.66% (-0.35-1.8%)	0.79% (-0.5-2.3%)	19 (-12-56)	760 (-400-2100)
South Sudan	2016	Female	55 to 59	46% (31-62%)	54% (38-69%)	0.29 (0.11-0.7)	0.35% (-0.71-1.5%)	0.45% (-0.98-2%)	12 (-25-52)	370 (-780-1600)
South Sudan	2016	Female	60 to 64	47% (30-63%)	53% (37-70%)	0.29 (0.1-0.68)	0.26% (-0.65-1.1%)	0.36% (-0.79-1.6%)	10 (-21-42)	250 (-630-1100)
South Sudan	2016	Female	65 to 69	48% (33-64%)	52% (36-67%)	0.27 (0.096-0.65)	0.21% (-0.72-1%)	0.26% (-0.88-1.2%)	9.1 (-31-47)	210 (-700-1000)
South Sudan	2016	Female	70 to 74	47% (32-62%)	53% (38-68%)	0.24 (0.082-0.57)	-0.1% (-1.1-0.68%)	-0.15% (-1.3-0.78%)	-6.4 (-54-33)	-90 (-970-620)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	2016	Female	75 to 79	45% (29-61%)	55% (39-71%)	0.21 (0.067-0.51)	-0.082% (-1.1-0.61%)	-0.093% (-1.3-0.72%)	-3.7 (-50-29)	-53 (-720-410)
South Sudan	2016	Female	80 to 84	44% (27-61%)	56% (39-73%)	0.2 (0.059-0.46)	-0.27% (-1.1-0.35%)	-0.3% (-1.3-0.42%)	-8.9 (-37-12)	-97 (-390-120)
South Sudan	2016	Female	85 to 89	43% (28-58%)	57% (42-72%)	0.19 (0.059-0.46)	-0.3% (-1.1-0.24%)	-0.36% (-1.3-0.29%)	-5.4 (-19-4.5)	-40 (-140-31)
South Sudan	2016	Female	90 to 94	45% (29-60%)	55% (40-71%)	0.18 (0.055-0.45)	-0.38% (-1.2-0.15%)	-0.48% (-1.5-0.15%)	-2.4 (-7.3-0.72)	-11 (-36-4.4)
South Sudan	2016	Female	95 plus	47% (29-65%)	53% (35-71%)	0.18 (0.05-0.46)	-0.033% (-0.75-0.5%)	-0.11% (-1-0.55%)	-0.13 (-1.1-0.64)	-0.15 (-3.5-2.4)
South Sudan	2016	Male	All Ages	41% (34-48%)	59% (52-66%)	0.17 (0.068-0.36)	0.52% (0.099-1%)	0.53% (-0.17-1.3%)	390 (-130-940)	26000 (5000-49000)
South Sudan	2016	Male	15 to 19	41% (28-56%)	59% (44-72%)	0.2 (0.065-0.45)	1% (0.48-1.8%)	0.5% (-0.27-1.5%)	7.1 (-3.6-22)	1700 (740-2900)
South Sudan	2016	Male	20 to 24	52% (37-66%)	48% (34-63%)	0.26 (0.089-0.58)	2% (0.75-3.5%)	0.8% (-0.86-2.6%)	15 (-14-49)	3500 (1200-6100)
South Sudan	2016	Male	25 to 29	56% (40-70%)	44% (30-60%)	0.32 (0.12-0.72)	2.1% (0.58-3.8%)	1.1% (-1-3.4%)	21 (-20-69)	3500 (940-6600)
South Sudan	2016	Male	30 to 34	58% (42-73%)	42% (27-58%)	0.36 (0.13-0.85)	1.8% (0.22-3.6%)	1.1% (-0.94-3.4%)	25 (-22-77)	3000 (360-6200)
South Sudan	2016	Male	35 to 39	59% (46-75%)	41% (25-54%)	0.38 (0.14-0.88)	1.7% (0.069-3.5%)	1.3% (-0.71-3.6%)	32 (-16-89)	2700 (120-5600)
South Sudan	2016	Male	40 to 44	60% (45-75%)	40% (25-55%)	0.39 (0.14-0.96)	1.6% (0.13-3.3%)	1.5% (-0.39-3.7%)	39 (-9.2-97)	2500 (190-5300)
South Sudan	2016	Male	45 to 49	61% (45-76%)	39% (24-55%)	0.39 (0.13-0.87)	1.7% (0.13-3.3%)	1.7% (-0.28-3.8%)	45 (-7.3-110)	2300 (180-4800)
South Sudan	2016	Male	50 to 54	61% (45-75%)	39% (25-55%)	0.38 (0.14-0.88)	1.6% (-0.072-3.4%)	1.7% (-0.49-4%)	45 (-12-110)	2000 (-79-4300)
South Sudan	2016	Male	55 to 59	60% (45-75%)	40% (25-55%)	0.38 (0.14-0.86)	1.5% (-0.18-3.3%)	1.6% (-0.57-3.9%)	41 (-15-100)	1600 (-180-3500)
South Sudan	2016	Male	60 to 64	59% (44-75%)	41% (25-56%)	0.34 (0.12-0.77)	1.3% (0.097-2.6%)	1.5% (-0.085-3%)	40 (-2.5-84)	1200 (87-2400)
South Sudan	2016	Male	65 to 69	58% (43-73%)	42% (27-57%)	0.28 (0.1-0.65)	0.76% (-0.21-1.7%)	0.81% (-0.38-2%)	27 (-12-66)	680 (-180-1500)
South Sudan	2016	Male	70 to 74	56% (41-70%)	44% (30-59%)	0.23 (0.081-0.53)	0.54% (-0.36-1.4%)	0.6% (-0.51-1.6%)	22 (-18-58)	430 (-290-1100)
South Sudan	2016	Male	75 to 79	54% (40-71%)	46% (29-60%)	0.19 (0.067-0.43)	0.48% (-0.31-1.1%)	0.53% (-0.42-1.3%)	17 (-13-42)	250 (-160-620)
South Sudan	2016	Male	80 to 84	52% (35-69%)	48% (31-65%)	0.17 (0.06-0.39)	0.32% (-0.32-0.87%)	0.32% (-0.43-1%)	6.8 (-9.4-21)	84 (-79-240)
South Sudan	2016	Male	85 to 89	50% (35-66%)	50% (34-65%)	0.15 (0.05-0.36)	0.32% (-0.22-0.78%)	0.31% (-0.35-0.87%)	3.3 (-3.7-9.2)	30 (-21-73)
South Sudan	2016	Male	90 to 94	51% (35-66%)	49% (34-65%)	0.14 (0.045-0.34)	0.21% (-0.28-0.57%)	0.15% (-0.44-0.6%)	0.55 (-1.6-2.2)	4.6 (-6.1-13)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
South Sudan	2016	Male	95 plus	52% (32-71%)	48% (29-68%)	0.13 (0.037-0.33)	0.32% (-0.25-0.77%)	0.25% (-0.48-0.81%)	0.21 (-0.41-0.68)	1.1 (-0.87-2.6)
Sudan	1990	Female	All Ages	11% (6.8-17%)	89% (83-93%)	0.52 (0.37-0.69)	1.3% (0.65-1.9%)	2.4% (0.91-4.5%)	2100 (810-3900)	76000 (39000-120000)
Sudan	1990	Female	15 to 19	9.9% (3.3-21%)	90% (79-97%)	0.41 (0.13-0.91)	1.3% (0.41-2.9%)	2% (0.51-4.8%)	40 (10-94)	3300 (990-7100)
Sudan	1990	Female	20 to 24	13% (4-26%)	87% (74-96%)	0.66 (0.21-1.4)	2.5% (0.88-4.9%)	3.6% (1-7.6%)	81 (22-170)	6200 (2200-12000)
Sudan	1990	Female	25 to 29	14% (4.5-29%)	86% (71-96%)	0.81 (0.25-1.6)	3% (1.1-5.7%)	4.6% (1.5-9.2%)	95 (31-190)	6600 (2600-13000)
Sudan	1990	Female	30 to 34	16% (4.2-33%)	84% (67-96%)	0.96 (0.31-1.8)	3.5% (1.4-6.4%)	5.5% (1.9-11%)	110 (38-220)	6900 (2700-13000)
Sudan	1990	Female	35 to 39	18% (5.2-35%)	82% (65-95%)	1.1 (0.35-2.1)	3.6% (1.4-6.9%)	5.6% (1.8-11%)	120 (40-240)	6600 (2400-13000)
Sudan	1990	Female	40 to 44	19% (7-35%)	81% (65-93%)	1.3 (0.44-2.4)	4.1% (1.5-7.6%)	6.4% (2.1-12%)	140 (46-260)	6700 (2400-13000)
Sudan	1990	Female	45 to 49	19% (7-36%)	81% (64-93%)	1.3 (0.45-2.8)	4.4% (1.6-8.7%)	6.8% (2.3-13%)	160 (52-310)	6700 (2300-13000)
Sudan	1990	Female	50 to 54	20% (8.1-39%)	80% (61-92%)	1.4 (0.45-3)	4.8% (1.4-9.6%)	6.9% (2-14%)	190 (54-390)	7100 (2100-14000)
Sudan	1990	Female	55 to 59	21% (8.8-41%)	79% (59-91%)	1.4 (0.41-3)	4.9% (1.1-10%)	6.7% (1.5-14%)	220 (51-460)	7100 (1700-15000)
Sudan	1990	Female	60 to 64	22% (9.1-41%)	78% (59-91%)	1.3 (0.37-3.1)	4.6% (0.6-9.9%)	6% (0.76-13%)	240 (31-530)	6600 (870-15000)
Sudan	1990	Female	65 to 69	23% (10-40%)	77% (60-90%)	1.2 (0.34-2.6)	3.8% (0.33-8.9%)	4.8% (0.43-11%)	230 (20-550)	5100 (460-12000)
Sudan	1990	Female	70 to 74	21% (9.2-38%)	79% (62-91%)	0.94 (0.25-2.3)	2.9% (-0.21-8.9%)	3.6% (-0.25-11%)	190 (-13-560)	3400 (-230-10000)
Sudan	1990	Female	75 to 79	19% (7.5-37%)	81% (63-93%)	0.77 (0.17-2.1)	2.3% (-0.34-7.6%)	2.8% (-0.4-9.1%)	130 (-19-440)	1800 (-280-6200)
Sudan	1990	Female	80 to 84	17% (6.1-38%)	83% (62-94%)	0.7 (0.15-2)	2.4% (-0.16-7.8%)	2.9% (-0.21-9.7%)	100 (-7.6-360)	1100 (-73-3700)
Sudan	1990	Female	85 to 89	18% (6.1-39%)	82% (61-94%)	0.71 (0.14-2.1)	2.3% (-0.17-8.1%)	2.7% (-0.23-9.6%)	60 (-4.9-210)	440 (-32-1500)
Sudan	1990	Female	90 to 94	21% (6.7-43%)	79% (57-93%)	0.8 (0.15-2.3)	2.1% (-0.42-7.6%)	2.5% (-0.54-9.4%)	21 (-4.5-82)	110 (-21-400)
Sudan	1990	Female	95 plus	25% (7.6-50%)	75% (50-92%)	0.94 (0.16-2.8)	2.1% (-0.85-8.3%)	2.5% (-1-10%)	5.7 (-2.4-23)	19 (-7.5-76)
Sudan	1990	Male	All Ages	8.7% (5.9-12%)	91% (88-94%)	0.58 (0.44-0.75)	1.3% (0.9-1.8%)	2.2% (1.4-3.3%)	2100 (1300-3000)	85000 (60000-120000)
Sudan	1990	Male	15 to 19	7.2% (3.1-14%)	93% (86-97%)	0.29 (0.097-0.64)	1.1% (0.55-2.2%)	1.5% (0.54-3.2%)	29 (11-63)	2700 (1300-5400)
Sudan	1990	Male	20 to 24	11% (4.6-22%)	89% (78-95%)	0.66 (0.24-1.3)	2.9% (1.4-5.1%)	3.6% (1.4-6.9%)	93 (37-180)	7400 (3600-13000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	1990	Male	25 to 29	14% (5.4-26%)	86% (74-95%)	1 (0.41-2)	4.2% (2-7.2%)	5.6% (2.4-10%)	130 (57-250)	9300 (4500-16000)
Sudan	1990	Male	30 to 34	15% (5.9-30%)	85% (70-94%)	1.3 (0.54-2.4)	4.9% (2.5-8.2%)	6.6% (3.2-11%)	160 (74-270)	9700 (5000-16000)
Sudan	1990	Male	35 to 39	17% (6.7-31%)	83% (69-93%)	1.5 (0.59-2.8)	5.3% (2.7-8.7%)	7.2% (3.4-12%)	180 (83-300)	9800 (5000-16000)
Sudan	1990	Male	40 to 44	17% (7.9-31%)	83% (69-92%)	1.6 (0.62-3.2)	5% (2.5-8.5%)	6.8% (3.2-12%)	170 (81-300)	8700 (4300-14000)
Sudan	1990	Male	45 to 49	17% (6.6-31%)	83% (69-93%)	1.6 (0.53-3.4)	5.1% (2.3-8.6%)	6.8% (2.8-12%)	190 (80-320)	8100 (3600-14000)
Sudan	1990	Male	50 to 54	16% (7-32%)	84% (68-93%)	1.6 (0.55-3.5)	4.6% (2-7.9%)	5.9% (2.5-10%)	180 (77-320)	7000 (3000-12000)
Sudan	1990	Male	55 to 59	16% (7-29%)	84% (71-93%)	1.5 (0.51-3.2)	4.5% (2-8%)	5.7% (2.5-10%)	200 (86-350)	6400 (2900-11000)
Sudan	1990	Male	60 to 64	15% (6.5-28%)	85% (72-94%)	1.4 (0.42-3)	4.1% (1.6-7.7%)	5.1% (2-9.5%)	200 (80-380)	5700 (2200-11000)
Sudan	1990	Male	65 to 69	14% (6.4-26%)	86% (74-94%)	1.1 (0.36-2.4)	3.6% (1.3-7%)	4.4% (1.5-8.7%)	190 (67-370)	4400 (1600-8400)
Sudan	1990	Male	70 to 74	13% (6.1-24%)	87% (76-94%)	0.87 (0.26-2.1)	2.9% (0.71-6.4%)	3.5% (0.83-7.8%)	160 (37-360)	3000 (710-6600)
Sudan	1990	Male	75 to 79	11% (4.9-22%)	89% (78-95%)	0.67 (0.17-1.7)	2.2% (0.37-5.2%)	2.5% (0.41-6.2%)	100 (17-260)	1500 (250-3600)
Sudan	1990	Male	80 to 84	10% (4.1-20%)	90% (80-96%)	0.55 (0.14-1.4)	1.9% (0.38-4.6%)	2.3% (0.44-5.4%)	65 (13-160)	680 (140-1600)
Sudan	1990	Male	85 to 89	9.8% (4-19%)	90% (81-96%)	0.5 (0.14-1.3)	1.6% (0.28-4%)	1.8% (0.31-4.6%)	29 (5-75)	220 (40-560)
Sudan	1990	Male	90 to 94	11% (4.5-22%)	89% (78-95%)	0.53 (0.13-1.4)	1.3% (0.16-3.8%)	1.5% (0.13-4.5%)	9.8 (0.82-28)	51 (6.3-140)
Sudan	1990	Male	95 plus	14% (5-29%)	86% (71-95%)	0.6 (0.12-1.7)	1.3% (0.0043-4.1%)	1.5% (-0.072-4.8%)	2.3 (-0.11-7.2)	8 (0.025-25)
Sudan	1995	Female	All Ages	11% (6.6-17%)	89% (83-93%)	0.63 (0.49-0.79)	1.6% (0.93-2.4%)	3.1% (1.4-5.2%)	3000 (1400-5200)	110000 (63000-160000)
Sudan	1995	Female	15 to 19	9.9% (3-22%)	90% (78-97%)	0.5 (0.14-1.1)	1.5% (0.52-3.2%)	2.5% (0.66-5.5%)	54 (15-120)	4400 (1500-9300)
Sudan	1995	Female	20 to 24	13% (4-28%)	87% (72-96%)	0.81 (0.26-1.7)	2.8% (1-5.4%)	4.3% (1.2-9%)	110 (32-230)	8300 (3200-16000)
Sudan	1995	Female	25 to 29	14% (3.8-30%)	86% (70-96%)	0.98 (0.29-1.9)	3.3% (1.3-6%)	5.3% (1.7-10%)	130 (41-250)	8900 (3400-16000)
Sudan	1995	Female	30 to 34	16% (4.1-32%)	84% (68-96%)	1.2 (0.35-2.2)	3.8% (1.5-6.8%)	6.2% (2.2-11%)	150 (51-270)	9200 (3700-16000)
Sudan	1995	Female	35 to 39	18% (5-34%)	82% (66-95%)	1.3 (0.44-2.6)	4% (1.6-7.1%)	6.3% (2.3-12%)	160 (55-290)	8700 (3300-15000)
Sudan	1995	Female	40 to 44	18% (6.7-35%)	82% (65-93%)	1.5 (0.54-2.8)	4.5% (1.9-8.2%)	7.3% (2.8-13%)	190 (72-340)	9100 (3700-16000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	1995	Female	45 to 49	19% (6.9-36%)	81% (64-93%)	1.6 (0.56-3.4)	5% (1.9-9.3%)	7.8% (2.9-14%)	220 (81-410)	9400 (3600-17000)
Sudan	1995	Female	50 to 54	20% (7.8-38%)	80% (62-92%)	1.7 (0.56-3.6)	5.5% (1.9-10%)	7.9% (2.7-15%)	270 (88-510)	9900 (3300-19000)
Sudan	1995	Female	55 to 59	21% (8.2-40%)	79% (60-92%)	1.7 (0.5-3.7)	5.6% (1.7-11%)	7.7% (2.4-15%)	310 (96-600)	9900 (3100-19000)
Sudan	1995	Female	60 to 64	22% (8.2-41%)	78% (59-92%)	1.6 (0.43-3.7)	5.3% (1.1-11%)	6.9% (1.4-14%)	340 (68-700)	9300 (1900-19000)
Sudan	1995	Female	65 to 69	22% (9.5-40%)	78% (60-91%)	1.4 (0.38-3)	4.6% (0.81-10%)	5.9% (1-13%)	340 (56-760)	7700 (1300-17000)
Sudan	1995	Female	70 to 74	21% (8.4-38%)	79% (62-92%)	1.1 (0.27-2.8)	3.6% (0.077-9.4%)	4.5% (0.079-12%)	280 (5-760)	5200 (110-13000)
Sudan	1995	Female	75 to 79	19% (6.8-38%)	81% (62-93%)	0.92 (0.19-2.6)	2.9% (-0.16-8.6%)	3.5% (-0.18-10%)	200 (-10-620)	2900 (-150-8600)
Sudan	1995	Female	80 to 84	17% (6.3-38%)	83% (62-94%)	0.84 (0.18-2.5)	3% (0.06-9.1%)	3.6% (0.071-11%)	160 (3-490)	1700 (31-5000)
Sudan	1995	Female	85 to 89	18% (6.2-39%)	82% (61-94%)	0.86 (0.18-2.5)	2.9% (0.032-9.3%)	3.5% (0.037-11%)	94 (1-300)	680 (7.1-2100)
Sudan	1995	Female	90 to 94	21% (7-44%)	79% (56-93%)	0.98 (0.19-2.9)	2.7% (-0.15-8.9%)	3.3% (-0.21-11%)	34 (-2.1-110)	170 (-8.9-560)
Sudan	1995	Female	95 plus	26% (8-50%)	74% (50-92%)	1.2 (0.19-3.6)	2.8% (-0.5-9.7%)	3.3% (-0.65-12%)	9.6 (-1.8-34)	32 (-5.5-110)
Sudan	1995	Male	All Ages	8.6% (5.9-12%)	91% (88-94%)	0.69 (0.54-0.87)	1.5% (1.1-2%)	2.6% (1.8-3.7%)	2800 (1900-3900)	110000 (84000-150000)
Sudan	1995	Male	15 to 19	7% (2.9-14%)	93% (86-97%)	0.34 (0.11-0.73)	1.3% (0.61-2.6%)	1.8% (0.64-3.9%)	39 (14-88)	3600 (1700-7300)
Sudan	1995	Male	20 to 24	11% (4.2-21%)	89% (79-96%)	0.77 (0.28-1.5)	3.4% (1.6-5.8%)	4.4% (1.7-8.2%)	130 (52-240)	10000 (4900-18000)
Sudan	1995	Male	25 to 29	14% (5.5-26%)	86% (74-94%)	1.2 (0.51-2.2)	4.7% (2.4-7.8%)	6.4% (2.9-11%)	180 (84-320)	13000 (6600-21000)
Sudan	1995	Male	30 to 34	15% (6.1-29%)	85% (71-94%)	1.5 (0.66-2.7)	5.3% (2.8-8.8%)	7.3% (3.7-13%)	210 (110-360)	13000 (6900-21000)
Sudan	1995	Male	35 to 39	17% (6.7-31%)	83% (69-93%)	1.8 (0.7-3.2)	5.7% (3.1-9.3%)	7.9% (4-13%)	230 (120-390)	13000 (6800-21000)
Sudan	1995	Male	40 to 44	17% (7.7-30%)	83% (70-92%)	1.9 (0.74-3.6)	5.5% (2.8-8.8%)	7.4% (3.6-12%)	230 (110-380)	11000 (5900-18000)
Sudan	1995	Male	45 to 49	17% (6.8-31%)	83% (69-93%)	1.9 (0.63-4)	5.5% (2.7-9%)	7.3% (3.4-12%)	250 (120-420)	11000 (5200-18000)
Sudan	1995	Male	50 to 54	16% (7-31%)	84% (69-93%)	1.9 (0.64-3.9)	5% (2.4-8.3%)	6.5% (3.1-11%)	240 (110-410)	9200 (4400-15000)
Sudan	1995	Male	55 to 59	15% (7-29%)	85% (71-93%)	1.8 (0.62-3.8)	4.9% (2.4-8.4%)	6.2% (3.1-11%)	260 (130-450)	8500 (4200-15000)
Sudan	1995	Male	60 to 64	15% (6.5-28%)	85% (72-93%)	1.6 (0.5-3.5)	4.5% (2-8.4%)	5.6% (2.5-10%)	270 (120-500)	7400 (3400-14000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	1995	Male	65 to 69	14% (6.4-27%)	86% (73-94%)	1.3 (0.43-2.8)	4% (1.6-7.3%)	4.9% (2-8.9%)	260 (100-470)	5900 (2400-11000)
Sudan	1995	Male	70 to 74	13% (5.7-23%)	87% (77-94%)	0.99 (0.29-2.3)	3.3% (0.92-6.6%)	3.9% (1.1-8%)	220 (60-450)	4000 (1100-8200)
Sudan	1995	Male	75 to 79	11% (4.7-21%)	89% (79-95%)	0.75 (0.19-1.9)	2.4% (0.52-5.4%)	2.8% (0.6-6.4%)	140 (31-320)	2000 (430-4500)
Sudan	1995	Male	80 to 84	9.9% (4-20%)	90% (80-96%)	0.63 (0.17-1.6)	2.2% (0.55-5%)	2.6% (0.65-5.8%)	91 (22-200)	960 (240-2100)
Sudan	1995	Male	85 to 89	9.7% (3.8-19%)	90% (81-96%)	0.58 (0.16-1.5)	1.8% (0.39-4.4%)	2.2% (0.42-5.2%)	44 (8.5-110)	330 (70-780)
Sudan	1995	Male	90 to 94	11% (4.4-21%)	89% (79-96%)	0.61 (0.16-1.6)	1.6% (0.26-4.3%)	1.9% (0.26-5.1%)	14 (2-38)	72 (12-190)
Sudan	1995	Male	95 plus	13% (5.1-28%)	87% (72-95%)	0.71 (0.14-2)	1.6% (0.067-4.6%)	1.8% (-0.0044-5.3%)	3.4 (-0.0083-10)	12 (0.5-35)
Sudan	2000	Female	All Ages	11% (6.5-17%)	89% (83-94%)	0.63 (0.5-0.78)	1.5% (0.92-2.3%)	3% (1.4-5%)	3100 (1400-5200)	110000 (63000-160000)
Sudan	2000	Female	15 to 19	9.7% (3.1-21%)	90% (79-97%)	0.49 (0.14-1.1)	1.3% (0.48-2.8%)	2.2% (0.6-4.8%)	47 (13-100)	4000 (1400-8400)
Sudan	2000	Female	20 to 24	12% (3.7-27%)	88% (73-96%)	0.78 (0.24-1.6)	2.4% (0.98-4.6%)	3.7% (1.1-7.7%)	96 (29-200)	7700 (3100-15000)
Sudan	2000	Female	25 to 29	14% (4.1-29%)	86% (71-96%)	0.95 (0.3-1.9)	2.9% (1.2-5.3%)	4.7% (1.6-9%)	120 (40-230)	8500 (3500-15000)
Sudan	2000	Female	30 to 34	16% (4.3-32%)	84% (68-96%)	1.1 (0.39-2.1)	3.4% (1.4-6%)	5.6% (2.1-10%)	140 (52-260)	9000 (3800-16000)
Sudan	2000	Female	35 to 39	18% (5.1-34%)	82% (66-95%)	1.3 (0.46-2.5)	3.6% (1.5-6.1%)	5.9% (2.2-10%)	150 (59-270)	8600 (3600-14000)
Sudan	2000	Female	40 to 44	19% (6.5-35%)	81% (65-93%)	1.5 (0.55-2.9)	4.2% (1.7-7.5%)	6.8% (2.6-12%)	180 (71-330)	9100 (3700-16000)
Sudan	2000	Female	45 to 49	19% (6.7-36%)	81% (64-93%)	1.6 (0.56-3.4)	4.5% (1.9-8.2%)	7.2% (2.8-13%)	210 (86-390)	9300 (3900-17000)
Sudan	2000	Female	50 to 54	20% (8.2-38%)	80% (62-92%)	1.7 (0.57-3.5)	5.1% (1.9-9.5%)	7.5% (2.6-14%)	270 (94-500)	10000 (3600-19000)
Sudan	2000	Female	55 to 59	21% (8.8-40%)	79% (60-91%)	1.7 (0.54-3.8)	5.4% (1.7-10%)	7.6% (2.4-15%)	320 (100-620)	10000 (3300-20000)
Sudan	2000	Female	60 to 64	22% (9-42%)	78% (58-91%)	1.7 (0.46-3.9)	5.1% (1.1-11%)	6.7% (1.4-14%)	350 (74-740)	9600 (2100-20000)
Sudan	2000	Female	65 to 69	22% (9.7-40%)	78% (60-90%)	1.4 (0.4-3.2)	4.5% (0.79-9.9%)	5.7% (1-13%)	350 (63-770)	8000 (1400-17000)
Sudan	2000	Female	70 to 74	21% (9.2-38%)	79% (62-91%)	1.1 (0.29-2.8)	3.4% (0.07-8.8%)	4.2% (0.077-11%)	290 (5.4-750)	5300 (110-14000)
Sudan	2000	Female	75 to 79	19% (7.4-38%)	81% (62-93%)	0.93 (0.2-2.6)	2.7% (-0.25-8.1%)	3.3% (-0.31-9.9%)	210 (-21-630)	2900 (-270-8900)
Sudan	2000	Female	80 to 84	17% (6.1-39%)	83% (61-94%)	0.85 (0.17-2.6)	2.8% (0.028-8.6%)	3.3% (0.019-10%)	170 (0.92-520)	1700 (17-5400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	2000	Female	85 to 89	18% (6.2-39%)	82% (61-94%)	0.85 (0.18-2.6)	2.7% (-0.0056-8.5%)	3.2% (-0.02-10%)	96 (-0.63-310)	700 (-1.4-2200)
Sudan	2000	Female	90 to 94	20% (6.9-42%)	80% (58-93%)	0.96 (0.19-2.9)	2.5% (-0.19-8.3%)	3% (-0.25-10%)	35 (-3-120)	180 (-13-590)
Sudan	2000	Female	95 plus	25% (7.6-49%)	75% (51-92%)	1.1 (0.19-3.5)	2.5% (-0.54-8.7%)	3% (-0.73-11%)	9.7 (-2.4-35)	33 (-7.2-110)
Sudan	2000	Male	All Ages	8.5% (5.9-12%)	91% (88-94%)	0.69 (0.55-0.85)	1.6% (1.1-2.1%)	2.6% (1.8-3.7%)	3000 (2000-4100)	120000 (90000-160000)
Sudan	2000	Male	15 to 19	7% (2.9-13%)	93% (87-97%)	0.33 (0.11-0.72)	1.3% (0.62-2.4%)	1.7% (0.63-3.7%)	40 (15-91)	3800 (1800-7400)
Sudan	2000	Male	20 to 24	11% (4.1-21%)	89% (79-96%)	0.75 (0.27-1.5)	3.1% (1.5-5.4%)	4% (1.6-7.4%)	130 (54-250)	11000 (5200-19000)
Sudan	2000	Male	25 to 29	13% (5.3-25%)	87% (75-95%)	1.2 (0.5-2.2)	4.4% (2.3-7.4%)	6% (2.7-10%)	190 (89-340)	14000 (7100-23000)
Sudan	2000	Male	30 to 34	15% (5.8-30%)	85% (70-94%)	1.5 (0.66-2.7)	4.9% (2.7-8.2%)	6.9% (3.4-12%)	220 (110-380)	14000 (7500-23000)
Sudan	2000	Male	35 to 39	17% (6.7-30%)	83% (70-93%)	1.8 (0.72-3.2)	5.4% (2.9-8.6%)	7.4% (3.8-12%)	250 (120-410)	14000 (7400-22000)
Sudan	2000	Male	40 to 44	17% (7.8-30%)	83% (70-92%)	1.9 (0.74-3.6)	5.2% (2.8-8.5%)	7.1% (3.6-12%)	240 (120-400)	12000 (6200-19000)
Sudan	2000	Male	45 to 49	17% (6.7-30%)	83% (70-93%)	2 (0.63-4)	5.2% (2.6-8.6%)	7.1% (3.3-12%)	260 (130-430)	11000 (5600-18000)
Sudan	2000	Male	50 to 54	16% (6.7-31%)	84% (69-93%)	1.9 (0.63-3.9)	4.8% (2.4-7.9%)	6.3% (3.1-10%)	260 (120-430)	9900 (4900-16000)
Sudan	2000	Male	55 to 59	15% (6.5-29%)	85% (71-94%)	1.8 (0.6-3.9)	4.7% (2.4-8%)	6% (2.9-10%)	270 (130-470)	9000 (4500-15000)
Sudan	2000	Male	60 to 64	15% (6.1-28%)	85% (72-94%)	1.6 (0.5-3.6)	4.3% (1.9-7.8%)	5.3% (2.3-9.9%)	290 (130-530)	7900 (3500-15000)
Sudan	2000	Male	65 to 69	14% (6.3-26%)	86% (74-94%)	1.3 (0.42-2.7)	3.8% (1.5-6.8%)	4.7% (1.9-8.5%)	270 (110-490)	6300 (2600-11000)
Sudan	2000	Male	70 to 74	13% (6-22%)	87% (78-94%)	0.99 (0.29-2.4)	3.1% (0.92-6.3%)	3.7% (1.1-7.7%)	230 (68-480)	4300 (1300-8800)
Sudan	2000	Male	75 to 79	11% (4.8-21%)	89% (79-95%)	0.76 (0.18-2)	2.3% (0.49-5.4%)	2.7% (0.55-6.3%)	150 (31-350)	2200 (470-5000)
Sudan	2000	Male	80 to 84	9.9% (4-20%)	90% (80-96%)	0.64 (0.16-1.6)	2.1% (0.53-4.8%)	2.5% (0.61-5.6%)	100 (24-230)	1100 (260-2400)
Sudan	2000	Male	85 to 89	9.7% (3.8-19%)	90% (81-96%)	0.59 (0.16-1.6)	1.8% (0.37-4.2%)	2.1% (0.39-5%)	50 (9.6-120)	370 (77-890)
Sudan	2000	Male	90 to 94	11% (4.4-21%)	89% (79-96%)	0.63 (0.16-1.7)	1.5% (0.23-4%)	1.8% (0.22-4.8%)	17 (2.1-45)	88 (13-230)
Sudan	2000	Male	95 plus	13% (4.9-29%)	87% (71-95%)	0.72 (0.14-2.1)	1.5% (0.055-4.5%)	1.7% (-0.025-5.2%)	3.8 (-0.056-12)	14 (0.49-40)
Sudan	2005	Female	All Ages	11% (6.6-16%)	89% (84-93%)	0.6 (0.47-0.75)	1.5% (0.86-2.2%)	2.8% (1.2-4.8%)	2800 (1200-4700)	98000 (59000-140000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	2005	Female	15 to 19	9.6% (3.2-21%)	90% (79-97%)	0.47 (0.14-1)	1.2% (0.47-2.3%)	1.9% (0.59-4.2%)	39 (12-85)	3500 (1400-7100)
Sudan	2005	Female	20 to 24	12% (3.7-27%)	88% (73-96%)	0.75 (0.24-1.6)	2.1% (0.9-4%)	3.4% (1.1-7%)	83 (26-170)	6900 (2900-13000)
Sudan	2005	Female	25 to 29	14% (4-29%)	86% (71-96%)	0.9 (0.28-1.8)	2.6% (1.1-4.5%)	4.2% (1.5-8.4%)	100 (35-200)	7700 (3400-14000)
Sudan	2005	Female	30 to 34	15% (4-33%)	85% (67-96%)	1.1 (0.36-2)	2.9% (1.3-5.1%)	5% (1.9-9.4%)	120 (46-230)	8200 (3700-14000)
Sudan	2005	Female	35 to 39	17% (4.6-33%)	83% (67-95%)	1.3 (0.43-2.3)	3.1% (1.3-5.3%)	5.3% (2.9-2%)	140 (51-250)	8100 (3400-14000)
Sudan	2005	Female	40 to 44	18% (6.3-34%)	82% (66-94%)	1.4 (0.51-2.6)	3.6% (1.5-6.4%)	6% (2.3-11%)	170 (65-300)	8500 (3500-15000)
Sudan	2005	Female	45 to 49	18% (6.5-34%)	82% (66-93%)	1.5 (0.5-3.1)	3.8% (1.5-7.1%)	6.2% (2.4-12%)	190 (76-360)	8500 (3500-16000)
Sudan	2005	Female	50 to 54	19% (7.5-36%)	81% (64-92%)	1.6 (0.5-3.3)	4.4% (1.4-8.3%)	6.6% (2.2-12%)	250 (79-470)	9500 (3100-18000)
Sudan	2005	Female	55 to 59	20% (8.1-40%)	80% (60-92%)	1.6 (0.48-3.6)	4.8% (1.4-9.3%)	6.7% (1.9-13%)	300 (84-600)	9800 (2800-20000)
Sudan	2005	Female	60 to 64	21% (8.7-41%)	79% (59-91%)	1.6 (0.43-3.7)	4.4% (0.88-9.6%)	6% (1.1-13%)	320 (61-690)	9000 (1800-19000)
Sudan	2005	Female	65 to 69	22% (9.4-39%)	78% (61-91%)	1.4 (0.39-3)	3.9% (0.62-8.7%)	5.1% (0.79-11%)	320 (50-720)	7500 (1100-16000)
Sudan	2005	Female	70 to 74	21% (8.4-37%)	79% (63-92%)	1.1 (0.27-2.7)	3% (-0.024-8%)	3.8% (-0.053-10%)	270 (-3.9-730)	5000 (-39-13000)
Sudan	2005	Female	75 to 79	19% (7-38%)	81% (62-93%)	0.9 (0.19-2.6)	2.3% (-0.29-7.4%)	2.9% (-0.36-9.1%)	200 (-25-620)	2800 (-340-8900)
Sudan	2005	Female	80 to 84	17% (6-38%)	83% (62-94%)	0.82 (0.17-2.4)	2.4% (-0.033-7.5%)	2.9% (-0.032-9%)	160 (-1.8-500)	1600 (-22-5100)
Sudan	2005	Female	85 to 89	18% (5.9-38%)	82% (62-94%)	0.81 (0.16-2.4)	2.2% (-0.091-7.5%)	2.7% (-0.12-9.2%)	96 (-4.2-330)	710 (-28-2400)
Sudan	2005	Female	90 to 94	20% (6.9-43%)	80% (57-93%)	0.92 (0.17-2.8)	2.1% (-0.27-7.4%)	2.5% (-0.38-9.2%)	36 (-5.2-130)	180 (-22-630)
Sudan	2005	Female	95 plus	25% (7.4-49%)	75% (51-93%)	1.1 (0.19-3.4)	2.2% (-0.67-8%)	2.5% (-0.86-9.6%)	9.8 (-3.3-37)	33 (-10-120)
Sudan	2005	Male	All Ages	8.6% (5.9-12%)	91% (88-94%)	0.67 (0.54-0.82)	1.6% (1.2-2.1%)	2.7% (1.8-3.7%)	3000 (2000-4100)	120000 (92000-160000)
Sudan	2005	Male	15 to 19	6.9% (2.8-14%)	93% (86-97%)	0.32 (0.098-0.71)	1.2% (0.62-2.4%)	1.6% (0.62-3.6%)	39 (15-88)	3800 (1900-7500)
Sudan	2005	Male	20 to 24	11% (4.2-21%)	89% (79-96%)	0.72 (0.27-1.4)	3% (1.6-5.2%)	3.9% (1.6-7.2%)	120 (51-230)	10000 (5400-18000)
Sudan	2005	Male	25 to 29	13% (5.4-25%)	87% (75-95%)	1.1 (0.47-2.1)	4.2% (2.2-7%)	5.8% (2.6-11%)	190 (83-330)	14000 (6900-23000)
Sudan	2005	Male	30 to 34	15% (5.6-29%)	85% (71-94%)	1.5 (0.62-2.6)	4.6% (2.5-7.6%)	6.6% (3.3-11%)	220 (110-380)	14000 (7700-23000)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	2005	Male	35 to 39	17% (6.4-30%)	83% (70-94%)	1.7 (0.67-3)	5% (2.7-8%)	7% (3.5-12%)	250 (120-400)	14000 (7400-22000)
Sudan	2005	Male	40 to 44	17% (7.5-30%)	83% (70-92%)	1.8 (0.7-3.4)	4.8% (2.6-7.9%)	6.7% (3.4-11%)	240 (120-410)	12000 (6500-20000)
Sudan	2005	Male	45 to 49	17% (6.9-31%)	83% (69-93%)	1.8 (0.6-3.7)	4.9% (2.4-8%)	6.6% (3.2-11%)	260 (120-420)	11000 (5600-18000)
Sudan	2005	Male	50 to 54	16% (6.8-31%)	84% (69-93%)	1.8 (0.62-3.8)	4.5% (2.2-7.5%)	5.9% (2.9-10%)	260 (130-440)	10000 (5100-17000)
Sudan	2005	Male	55 to 59	15% (6.5-29%)	85% (71-93%)	1.7 (0.56-3.6)	4.4% (2.1-7.6%)	5.7% (2.8-9.7%)	280 (140-500)	9400 (4700-16000)
Sudan	2005	Male	60 to 64	15% (6.2-28%)	85% (72-94%)	1.6 (0.47-3.4)	4% (1.7-7.2%)	5% (2.1-9.1%)	290 (120-530)	8100 (3500-15000)
Sudan	2005	Male	65 to 69	14% (6.3-26%)	86% (74-94%)	1.3 (0.4-2.6)	3.6% (1.4-6.4%)	4.4% (1.8-7.9%)	280 (110-510)	6500 (2700-12000)
Sudan	2005	Male	70 to 74	13% (5.7-23%)	87% (77-94%)	0.95 (0.28-2.3)	2.9% (0.8-5.9%)	3.4% (0.95-7.2%)	230 (64-480)	4300 (1200-8900)
Sudan	2005	Male	75 to 79	11% (4.7-20%)	89% (80-95%)	0.72 (0.18-1.9)	2.1% (0.44-4.8%)	2.5% (0.51-5.7%)	150 (31-350)	2200 (460-5000)
Sudan	2005	Male	80 to 84	9.7% (3.8-19%)	90% (81-96%)	0.6 (0.15-1.5)	1.9% (0.46-4.4%)	2.2% (0.51-5.1%)	100 (24-230)	1100 (260-2500)
Sudan	2005	Male	85 to 89	9.5% (3.9-19%)	90% (81-96%)	0.55 (0.15-1.4)	1.6% (0.31-3.9%)	1.8% (0.33-4.5%)	53 (9.8-130)	400 (78-990)
Sudan	2005	Male	90 to 94	11% (4.3-21%)	89% (79-96%)	0.58 (0.15-1.5)	1.4% (0.16-3.7%)	1.6% (0.13-4.2%)	18 (1.6-49)	96 (11-250)
Sudan	2005	Male	95 plus	13% (4.9-28%)	87% (72-95%)	0.67 (0.13-1.9)	1.3% (-0.0045-4.2%)	1.5% (-0.1-4.8%)	4.3 (-0.29-14)	15 (-0.051-49)
Sudan	2010	Female	All Ages	11% (6.9-17%)	89% (83-93%)	0.65 (0.51-0.84)	1.5% (0.88-2.2%)	2.9% (1.3-4.8%)	2800 (1300-4700)	100000 (62000-140000)
Sudan	2010	Female	15 to 19	9.6% (3.2-21%)	90% (79-97%)	0.49 (0.15-1.1)	1.1% (0.47-2.3%)	2% (0.57-4.3%)	39 (12-87)	3700 (1500-7400)
Sudan	2010	Female	20 to 24	12% (3.8-26%)	88% (74-96%)	0.78 (0.24-1.6)	2% (0.92-3.8%)	3.3% (1.1-6.7%)	78 (26-160)	6800 (3100-13000)
Sudan	2010	Female	25 to 29	14% (4-29%)	86% (71-96%)	0.96 (0.3-1.9)	2.4% (1.1-4.3%)	4.2% (1.4-8.1%)	99 (33-190)	7700 (3600-14000)
Sudan	2010	Female	30 to 34	16% (4.2-32%)	84% (68-96%)	1.1 (0.37-2.1)	2.8% (1.3-4.7%)	5% (1.9-8.9%)	120 (47-220)	8500 (3800-14000)
Sudan	2010	Female	35 to 39	18% (4.7-34%)	82% (66-95%)	1.3 (0.42-2.5)	3% (1.3-5.3%)	5.3% (2-9.5%)	150 (56-270)	8800 (3900-15000)
Sudan	2010	Female	40 to 44	18% (6.1-35%)	82% (65-94%)	1.5 (0.53-2.8)	3.4% (1.4-6%)	5.9% (2.3-10%)	180 (69-310)	9100 (3800-16000)
Sudan	2010	Female	45 to 49	19% (7-35%)	81% (65-93%)	1.6 (0.54-3.3)	3.7% (1.5-6.7%)	6.2% (2.4-11%)	200 (81-370)	9200 (3800-17000)
Sudan	2010	Female	50 to 54	20% (8.2-38%)	80% (62-92%)	1.7 (0.56-3.5)	4.2% (1.5-7.8%)	6.6% (2.3-12%)	250 (87-480)	9800 (3300-18000)


Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	2010	Female	55 to 59	21% (8.6-40%)	79% (60-91%)	1.7 (0.51-3.7)	4.5% (1.3-8.8%)	6.7% (2-13%)	300 (85-580)	10000 (3000-19000)
Sudan	2010	Female	60 to 64	22% (9-42%)	78% (58-91%)	1.6 (0.46-3.9)	4.3% (0.83-9.1%)	5.9% (1.1-13%)	320 (61-680)	9000 (1700-19000)
Sudan	2010	Female	65 to 69	22% (9.9-40%)	78% (60-90%)	1.4 (0.43-3.2)	3.8% (0.64-8.3%)	5.1% (0.85-11%)	320 (53-720)	7500 (1300-16000)
Sudan	2010	Female	70 to 74	21% (9.3-37%)	79% (63-91%)	1.1 (0.3-2.8)	2.9% (-0.047-7.6%)	3.7% (-0.036-9.8%)	270 (-2.5-700)	5000 (-81-13000)
Sudan	2010	Female	75 to 79	18% (7.2-37%)	82% (63-93%)	0.91 (0.2-2.4)	2.2% (-0.28-6.7%)	2.7% (-0.37-8.4%)	190 (-27-600)	2800 (-380-8600)
Sudan	2010	Female	80 to 84	17% (5.8-36%)	83% (64-94%)	0.82 (0.17-2.5)	2.2% (-0.029-6.9%)	2.7% (-0.05-8.4%)	160 (-3-500)	1600 (-22-5200)
Sudan	2010	Female	85 to 89	17% (6-38%)	83% (62-94%)	0.83 (0.16-2.5)	2.2% (-0.091-7.4%)	2.6% (-0.16-8.9%)	100 (-6.1-350)	770 (-33-2600)
Sudan	2010	Female	90 to 94	20% (6.8-42%)	80% (58-93%)	0.94 (0.18-2.9)	2.1% (-0.27-7.1%)	2.5% (-0.39-8.7%)	43 (-6.7-150)	220 (-30-770)
Sudan	2010	Female	95 plus	24% (7.7-49%)	76% (51-92%)	1.1 (0.19-3.6)	2.1% (-0.63-7.9%)	2.5% (-0.79-9.5%)	12 (-3.8-44)	41 (-11-150)
Sudan	2010	Male	All Ages	8.6% (5.8-12%)	91% (88-94%)	0.7 (0.54-0.87)	1.7% (1.2-2.2%)	2.7% (1.8-3.8%)	3200 (2100-4400)	130000 (98000-170000)
Sudan	2010	Male	15 to 19	6.9% (2.9-13%)	93% (87-97%)	0.32 (0.096-0.72)	1.3% (0.63-2.5%)	1.7% (0.62-4%)	44 (16-100)	4300 (2100-8800)
Sudan	2010	Male	20 to 24	11% (4.2-20%)	89% (80-96%)	0.73 (0.25-1.5)	3% (1.5-5.2%)	3.9% (1.6-7.5%)	130 (52-240)	11000 (5500-19000)
Sudan	2010	Male	25 to 29	13% (5.1-25%)	87% (75-95%)	1.1 (0.47-2.2)	4.1% (2.1-6.7%)	5.7% (2.5-10%)	190 (86-340)	14000 (7300-24000)
Sudan	2010	Male	30 to 34	15% (5.6-29%)	85% (71-94%)	1.5 (0.61-2.6)	4.5% (2.4-7.3%)	6.5% (3.2-11%)	230 (110-390)	15000 (8200-24000)
Sudan	2010	Male	35 to 39	16% (6.3-30%)	84% (70-94%)	1.7 (0.68-3)	4.8% (2.6-7.7%)	6.9% (3.4-11%)	260 (130-430)	15000 (8100-24000)
Sudan	2010	Male	40 to 44	17% (7.5-30%)	83% (70-93%)	1.8 (0.7-3.4)	4.6% (2.5-7.7%)	6.5% (3.3-11%)	260 (130-430)	13000 (7100-21000)
Sudan	2010	Male	45 to 49	16% (6.4-30%)	84% (70-94%)	1.8 (0.61-3.7)	4.7% (2.4-7.5%)	6.4% (3.1-11%)	270 (130-440)	12000 (6200-19000)
Sudan	2010	Male	50 to 54	16% (6.8-30%)	84% (70-93%)	1.8 (0.61-3.7)	4.3% (2.2-7.1%)	5.8% (2.9-9.6%)	270 (130-460)	11000 (5400-17000)
Sudan	2010	Male	55 to 59	15% (6.7-28%)	85% (72-93%)	1.8 (0.59-3.5)	4.2% (2.2-7.3%)	5.6% (2.8-9.4%)	290 (150-500)	9700 (5000-17000)
Sudan	2010	Male	60 to 64	14% (6-28%)	86% (72-94%)	1.6 (0.5-3.6)	3.9% (1.8-6.9%)	4.9% (2.3-8.9%)	300 (140-540)	8600 (4000-15000)
Sudan	2010	Male	65 to 69	14% (6.2-26%)	86% (74-94%)	1.3 (0.41-2.8)	3.5% (1.4-6.3%)	4.3% (1.8-7.7%)	300 (120-530)	6900 (2900-12000)
Sudan	2010	Male	70 to 74	13% (5.8-22%)	87% (78-94%)	0.98 (0.28-2.3)	2.8% (0.85-5.6%)	3.4% (1-6.9%)	250 (73-500)	4600 (1400-9400)

Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	2010	Male	75 to 79	11% (4.7-21%)	89% (79-95%)	0.74 (0.19-2)	2% (0.43-4.6%)	2.4% (0.51-5.5%)	170 (35-380)	2400 (520-5500)
Sudan	2010	Male	80 to 84	9.8% (3.9-20%)	90% (80-96%)	0.63 (0.16-1.6)	1.9% (0.42-4.3%)	2.2% (0.48-4.9%)	110 (25-260)	1200 (280-2700)
Sudan	2010	Male	85 to 89	9.6% (3.7-20%)	90% (80-96%)	0.58 (0.16-1.6)	1.6% (0.32-3.9%)	1.8% (0.34-4.6%)	61 (12-150)	460 (95-1100)
Sudan	2010	Male	90 to 94	11% (4.3-21%)	89% (79-96%)	0.6 (0.15-1.6)	1.4% (0.17-3.6%)	1.6% (0.14-4.3%)	23 (2-63)	120 (16-320)
Sudan	2010	Male	95 plus	13% (5-28%)	87% (72-95%)	0.69 (0.14-2)	1.3% (-0.009-4.2%)	1.5% (-0.11-4.8%)	5.5 (-0.39-18)	20 (-0.14-61)
Sudan	2016	Female	All Ages	12% (7.2-18%)	88% (82-93%)	0.73 (0.45-1.1)	1.6% (0.86-2.6%)	3.1% (1.1-5.6%)	3000 (1100-5400)	110000 (58000-160000)
Sudan	2016	Female	15 to 19	10% (3.1-22%)	90% (78-97%)	0.54 (0.15-1.2)	1.1% (0.45-2.2%)	2.1% (0.58-4.7%)	39 (11-87)	3800 (1500-7500)
Sudan	2016	Female	20 to 24	13% (4-28%)	87% (72-96%)	0.86 (0.26-1.9)	2% (0.91-3.7%)	3.4% (1.1-7.3%)	75 (23-160)	6900 (3100-13000)
Sudan	2016	Female	25 to 29	14% (4.3-30%)	86% (70-96%)	1 (0.29-2.3)	2.3% (1.1-4.3%)	4.2% (1.4-8.6%)	93 (32-190)	7600 (3600-14000)
Sudan	2016	Female	30 to 34	16% (4.4-33%)	84% (67-96%)	1.2 (0.36-2.5)	2.6% (1.2-4.6%)	4.9% (1.8-9.3%)	120 (44-220)	8400 (3800-15000)
Sudan	2016	Female	35 to 39	17% (4.7-33%)	83% (67-95%)	1.4 (0.41-2.8)	2.8% (1.1-5%)	5.2% (1.8-9.7%)	150 (51-280)	9000 (3600-16000)
Sudan	2016	Female	40 to 44	18% (6.5-34%)	82% (66-94%)	1.6 (0.49-3.2)	3.2% (1.3-5.7%)	5.9% (2.3-10%)	190 (75-340)	9900 (4200-17000)
Sudan	2016	Female	45 to 49	19% (6.6-36%)	81% (64-93%)	1.7 (0.54-3.5)	3.5% (1.3-6.6%)	6.1% (2.2-11%)	220 (78-420)	10000 (3800-19000)
Sudan	2016	Female	50 to 54	20% (6.9-39%)	80% (61-93%)	1.7 (0.52-3.8)	4% (1.2-7.6%)	6.4% (2.1-12%)	270 (85-520)	11000 (3400-21000)
Sudan	2016	Female	55 to 59	20% (7.5-40%)	80% (60-93%)	1.7 (0.48-3.9)	4.2% (1.2-8.3%)	6.4% (1.8-13%)	310 (85-620)	11000 (3000-21000)
Sudan	2016	Female	60 to 64	22% (8.4-40%)	78% (60-92%)	1.7 (0.46-4.2)	4% (0.7-8.6%)	5.6% (1-12%)	340 (61-720)	9600 (1700-20000)
Sudan	2016	Female	65 to 69	22% (9.3-40%)	78% (60-91%)	1.5 (0.41-3.4)	3.7% (0.45-8.1%)	5% (0.7-11%)	340 (48-770)	8000 (1000-18000)
Sudan	2016	Female	70 to 74	21% (8.9-38%)	79% (62-91%)	1.2 (0.3-3.1)	2.9% (-0.064-7.5%)	3.8% (-0.06-9.7%)	290 (-4.5-770)	5400 (-120-14000)
Sudan	2016	Female	75 to 79	19% (7.6-39%)	81% (61-92%)	1 (0.21-2.8)	2.2% (-0.32-6.6%)	2.8% (-0.39-8.4%)	220 (-31-650)	3200 (-460-9400)
Sudan	2016	Female	80 to 84	18% (6-39%)	82% (61-94%)	0.93 (0.17-2.9)	2.3% (-0.083-7.3%)	2.8% (-0.13-9.1%)	180 (-8.1-590)	1900 (-67-6200)
Sudan	2016	Female	85 to 89	18% (6.3-40%)	82% (60-94%)	0.93 (0.17-2.8)	2.3% (-0.17-7.7%)	2.7% (-0.23-9.2%)	130 (-11-440)	970 (-72-3300)
Sudan	2016	Female	90 to 94	21% (7-43%)	79% (57-93%)	1 (0.19-3.3)	2.2% (-0.32-7.3%)	2.6% (-0.45-8.8%)	57 (-9.7-190)	290 (-44-960)


Location	Year	Sex	Age	Prevalence of current drinkers	Prevalence of abstainers	Population average of standard drinks daily	Percent of total attributable deaths	Percent of total attributable DALYs	Attributable deaths	Attributable DALYs
Sudan	2016	Female	95 plus	25% (7.9-51%)	75% (49-92%)	1.2 (0.21-4)	2.3% (-0.68-7.9%)	2.7% (-0.85-9.6%)	17 (-5.7-62)	60 (-18-210)
Sudan	2016	Male	All Ages	9.1% (6.2-13%)	91% (87-94%)	0.78 (0.49-1.2)	1.9% (1.3-2.7%)	2.9% (1.8-4.2%)	3500 (2200-4900)	140000 (100000-200000)
Sudan	2016	Male	15 to 19	7% (2.8-14%)	93% (86-97%)	0.35 (0.1-0.82)	1.3% (0.62-2.7%)	1.8% (0.59-4.3%)	50 (17-120)	5000 (2400-10000)
Sudan	2016	Male	20 to 24	11% (4.2-20%)	89% (80-96%)	0.78 (0.27-1.7)	3% (1.5-5.3%)	3.9% (1.5-7.8%)	140 (57-280)	12000 (6300-22000)
Sudan	2016	Male	25 to 29	13% (5.3-25%)	87% (75-95%)	1.2 (0.44-2.5)	4% (2-6.7%)	5.6% (2.4-10%)	210 (85-370)	15000 (7700-26000)
Sudan	2016	Male	30 to 34	15% (5.8-29%)	85% (71-94%)	1.6 (0.6-3.1)	4.4% (2.4-7.3%)	6.4% (3-11%)	240 (110-420)	16000 (8400-26000)
Sudan	2016	Male	35 to 39	17% (6.5-30%)	83% (70-93%)	1.8 (0.62-3.6)	4.7% (2.4-7.6%)	6.8% (3.2-11%)	280 (140-480)	17000 (8600-27000)
Sudan	2016	Male	40 to 44	17% (7.5-30%)	83% (70-92%)	1.9 (0.65-3.9)	4.5% (2.3-7.7%)	6.5% (3.1-11%)	290 (140-500)	15000 (7900-25000)
Sudan	2016	Male	45 to 49	17% (6.6-30%)	83% (70-93%)	2 (0.6-4.3)	4.5% (2.2-7.3%)	6.4% (2.9-11%)	310 (140-500)	14000 (6900-22000)
Sudan	2016	Male	50 to 54	16% (6.6-30%)	84% (70-93%)	1.9 (0.61-4.2)	4.2% (2-6.9%)	5.7% (2.6-9.4%)	300 (140-500)	12000 (5800-20000)
Sudan	2016	Male	55 to 59	15% (6.6-29%)	85% (71-93%)	1.8 (0.56-4)	4.1% (2-7%)	5.4% (2.6-9.3%)	320 (150-550)	11000 (5300-18000)
Sudan	2016	Male	60 to 64	14% (6.1-28%)	86% (72-94%)	1.6 (0.49-3.8)	3.7% (1.6-6.6%)	4.7% (2-8.6%)	320 (140-600)	9200 (4100-17000)
Sudan	2016	Male	65 to 69	14% (6-25%)	86% (75-94%)	1.3 (0.39-3)	3.3% (1.3-5.9%)	4.1% (1.6-7.4%)	320 (120-570)	7500 (3000-14000)
Sudan	2016	Male	70 to 74	12% (5.6-22%)	88% (78-94%)	1 (0.27-2.4)	2.6% (0.74-5.1%)	3.2% (0.88-6.2%)	270 (73-520)	5000 (1400-9800)
Sudan	2016	Male	75 to 79	11% (4.5-21%)	89% (79-96%)	0.76 (0.18-2)	1.9% (0.39-4.3%)	2.3% (0.45-5.1%)	180 (35-410)	2600 (530-5900)
Sudan	2016	Male	80 to 84	9.5% (3.8-19%)	90% (81-96%)	0.64 (0.14-1.6)	1.8% (0.38-4%)	2.1% (0.43-4.7%)	120 (26-290)	1300 (290-3100)
Sudan	2016	Male	85 to 89	9.4% (3.8-18%)	91% (82-96%)	0.59 (0.15-1.5)	1.5% (0.26-3.7%)	1.7% (0.27-4.3%)	71 (11-180)	550 (94-1300)
Sudan	2016	Male	90 to 94	11% (4.2-22%)	89% (78-96%)	0.63 (0.15-1.7)	1.4% (0.1-3.5%)	1.6% (0.061-4.2%)	28 (1.1-77)	150 (11-400)
Sudan	2016	Male	95 plus	13% (4.9-27%)	87% (73-95%)	0.73 (0.13-2.2)	1.3% (-0.023-4.1%)	1.5% (-0.15-4.7%)	7.4 (-0.77-23)	27 (-0.46-81)

Supplemental Results Figure 1: Attributable DALY rate disaggregated by cause by location, sex, and age in 2016


Attributable DALY rate disaggregated by cause, China, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, North Korea, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Taiwan, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Indonesia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Laos, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Malaysia, by sex and age in 2016.




Cause

- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
- Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Transport injuries
- Unintentional injuries
 - Self-harm
 - Interpersonal violence

Attributable DALY rate disaggregated by cause, Maldives, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Mauritius, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Philippines, by sex and age in 2016.


Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Alcohol use disorders
 - Lip and oral cavity cancer
 - Diabetes mellitus
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Sport injuries


Attributable DALY rate disaggregated by cause, Sri Lanka, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Seychelles, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Thailand, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Timor-Leste, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Vietnam, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Federated States of Micronesia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Fiji, by sex and age in 2016.

Female

Male




Cause

- Tuberculosis
- Lower respiratory infections
- Esophageal cancer
- Liver cancer
- Larynx cancer
- Breast cancer
- Colon and rectum cancer
- Lip and oral cavity cancer
- Pharynx and nasopharynx cancer
- Ischaemic heart disease
- Ischaemic stroke
- Hemorrhagic stroke
- Hypertensive heart disease
- Atrial fibrillation and flutter
- Cirrhosis and other chronic liver diseases
- Pancreatitis
- Epilepsy
- Alcohol use disorders
- Diabetes mellitus
- Unintentional injuries
- Self-harm
- Interpersonal violence

Attributable DALY rate disaggregated by cause, Kiribati, by sex and age in 2016.

Female

Male




- Cause**
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - 2015 sport injuries
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence

Attributable DALY rate disaggregated by cause, Marshall Islands, by sex and age in 2016.

Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - 2016 sport injuries

Attributable DALY rate disaggregated by cause, Papua New Guinea, by sex and age in 2016.

Female


Male



Attributable DALY rate disaggregated by cause, Samoa, by sex and age in 2016.


Female

Male



- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - 2018 sport injuries


Attributable DALY rate disaggregated by cause, Solomon Islands, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Tonga, by sex and age in 2016.


Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - 2020 sport injuries


Attributable DALY rate disaggregated by cause, Vanuatu, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Armenia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Azerbaijan, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Georgia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Kazakhstan, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Kyrgyzstan, by sex and age in 2016.


Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - 2026 sport injuries


Attributable DALY rate disaggregated by cause, Mongolia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Tajikistan, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Turkmenistan, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Uzbekistan, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Albania, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Bosnia and Herzegovina, by sex and age in 2016.

Female


Male




Cause

- Tuberculosis
- Ischaemic stroke
- Unintentional injuries
- Lower respiratory infections
- Hemorrhagic stroke
- Self-harm
- Esophageal cancer
- Hypertensive heart disease
- Interpersonal violence
- Liver cancer
- Atrial fibrillation and flutter
- Pharynx and nasopharynx cancer
- Larynx cancer
- Cirrhosis and other chronic liver diseases
- 2012
- Breast cancer
- Pancreatitis
- Transport injuries
- Colon and rectum cancer
- Epilepsy
- 2012
- Lip and oral cavity cancer
- Alcohol use disorders
- 2012
- Pharynx and nasopharynx cancer
- 2012
- Ischaemic heart disease
- 2012


Attributable DALY rate disaggregated by cause, Bulgaria, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Croatia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Czech Republic, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Hungary, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Macedonia, by sex and age in 2016.


Female

Male



- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - Sport injuries


Attributable DALY rate disaggregated by cause, Montenegro, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Poland, by sex and age in 2016.

Female

Male




Cause

- | | | |
|---|---|--|
| <ul style="list-style-type: none"> ■ Tuberculosis ■ Lower respiratory infections ■ Esophageal cancer ■ Liver cancer ■ Larynx cancer ■ Breast cancer ■ Colon and rectum cancer ■ Lip and oral cavity cancer ■ Pharynx and nasopharynx cancer ■ Ischaemic heart disease | <ul style="list-style-type: none"> ■ Ischaemic stroke ■ Hemorrhagic stroke ■ Hypertensive heart disease ■ Atrial fibrillation and flutter ■ Cirrhosis and other chronic liver diseases ■ Pancreatitis ■ Epilepsy ■ Alcohol use disorders ■ Diabetes mellitus ■ Unintentional injuries | <ul style="list-style-type: none"> ■ Self-harm ■ Interpersonal violence ■ Sport injuries |
|---|---|--|

Attributable DALY rate disaggregated by cause, Romania, by sex and age in 2016.

Female

Male




- Cause**
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - 2040 sport injuries
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence


Attributable DALY rate disaggregated by cause, Serbia, by sex and age in 2016.

Female


Male




Attributable DALY rate disaggregated by cause, Slovakia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Slovenia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Belarus, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Estonia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Latvia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Lithuania, by sex and age in 2016.

Female


Male



Cause

- Tuberculosis
- Ischaemic stroke
- Unintentional injuries
- Lower respiratory infections
- Hemorrhagic stroke
- Self-harm
- Esophageal cancer
- Hypertensive heart disease
- Interpersonal violence
- Liver cancer
- Atrial fibrillation and flutter
- Larynx cancer
- Cirrhosis and other chronic liver diseases
- Breast cancer
- Pancreatitis
- Colon and rectum cancer
- Epilepsy
- Lip and oral cavity cancer
- Alcohol use disorders
- Pharynx and nasopharynx cancer
- Diabetes mellitus
- Ischaemic heart disease
- Sport injuries


Attributable DALY rate disaggregated by cause, Moldova, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Russia, by sex and age in 2016.

Female


Male



Attributable DALY rate disaggregated by cause, Ukraine, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Brunei, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Japan, by sex and age in 2016.

Female


Male




Cause

- | | | |
|---|---|---|
| ■ Tuberculosis | ■ Ischaemic stroke | ■ Unintentional injuries |
| ■ Lower respiratory infections | ■ Hemorrhagic stroke | ■ Self-harm |
| ■ Esophageal cancer | ■ Hypertensive heart disease | ■ Interpersonal violence |
| ■ Liver cancer | ■ Atrial fibrillation and flutter | |
| ■ Larynx cancer | ■ Pancreatitis | |
| ■ Breast cancer | ■ Cirrhosis and other chronic liver diseases | |
| ■ Colon and rectum cancer | ■ Epilepsy | |
| ■ Lip and oral cavity cancer | ■ Alcohol use disorders | |
| ■ Pharynx and nasopharynx cancer | ■ Diabetes mellitus | |
| ■ Ischaemic heart disease | ■ 2012 sport injuries | |


Attributable DALY rate disaggregated by cause, South Korea, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Singapore, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Australia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, New Zealand, by sex and age in 2016.

Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - Sport injuries


Attributable DALY rate disaggregated by cause, Andorra, by sex and age in 2016.

Female


Male




Attributable DALY rate disaggregated by cause, Austria, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Belgium, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Cyprus, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Finland, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, France, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Germany, by sex and age in 2016.

Female

Male




- Cause
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence

Attributable DALY rate disaggregated by cause, Greece, by sex and age in 2016.

Female

Male




Cause

- | | | |
|---|---|---|
| ■ Tuberculosis | ■ Ischaemic stroke | ■ Unintentional injuries |
| ■ Lower respiratory infections | ■ Hemorrhagic stroke | ■ Self-harm |
| ■ Esophageal cancer | ■ Hypertensive heart disease | ■ Interpersonal violence |
| ■ Liver cancer | ■ Atrial fibrillation and flutter | |
| ■ Larynx cancer | ■ Cirrhosis and other chronic liver diseases | |
| ■ Breast cancer | ■ Pancreatitis | |
| ■ Colon and rectum cancer | ■ Epilepsy | |
| ■ Lip and oral cavity cancer | ■ Alcohol use disorders | |
| ■ Pharynx and nasopharynx cancer | ■ Diabetes mellitus | |
| ■ Ischaemic heart disease | ■ Sport injuries | |

Attributable DALY rate disaggregated by cause, Greenland, by sex and age in 2016.

Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - Sport injuries

Attributable DALY rate disaggregated by cause, Iceland, by sex and age in 2016.

Female

Male




- Cause
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence
 - Sport injuries

Attributable DALY rate disaggregated by cause, Ireland, by sex and age in 2016.


Female

Male



- Cause
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence
 - Transport injuries


Attributable DALY rate disaggregated by cause, Israel, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Italy, by sex and age in 2016.

Female


Male




Attributable DALY rate disaggregated by cause, Luxembourg, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Malta, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Netherlands, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Norway, by sex and age in 2016.

Female

Male




Attributable DALY rate disaggregated by cause, Portugal, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Spain, by sex and age in 2016.

Female


Male



Attributable DALY rate disaggregated by cause, Sweden, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Switzerland, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, United Kingdom, by sex and age in 2016.


Female

Male



- Cause
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Alcohol use disorders
 - Lip and oral cavity cancer
 - Diabetes mellitus
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - 2019 sport injuries


Attributable DALY rate disaggregated by cause, Argentina, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Chile, by sex and age in 2016.

Female

Male




- Cause
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - 2011 sport injuries

Attributable DALY rate disaggregated by cause, Uruguay, by sex and age in 2016.

Female


Male



Cause

- | | | |
|---|---|---|
| ■ Tuberculosis | ■ Ischaemic stroke | ■ Unintentional injuries |
| ■ Lower respiratory infections | ■ Hemorrhagic stroke | ■ Self-harm |
| ■ Esophageal cancer | ■ Hypertensive heart disease | ■ Interpersonal violence |
| ■ Liver cancer | ■ Atrial fibrillation and flutter | |
| ■ Larynx cancer | ■ Cirrhosis and other chronic liver diseases | |
| ■ Breast cancer | ■ Pancreatitis | |
| ■ Colon and rectum cancer | ■ Epilepsy | |
| ■ Lip and oral cavity cancer | ■ Alcohol use disorders | |
| ■ Pharynx and nasopharynx cancer | ■ Diabetes mellitus | |
| ■ Ischaemic heart disease | ■ 2012 sport injuries | |


Attributable DALY rate disaggregated by cause, Canada, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, United States, by sex and age in 2016.

Female


Male



Cause

- | | | |
|---|---|---|
| ■ Tuberculosis | ■ Ischaemic stroke | ■ Unintentional injuries |
| ■ Lower respiratory infections | ■ Hemorrhagic stroke | ■ Self-harm |
| ■ Esophageal cancer | ■ Hypertensive heart disease | ■ Interpersonal violence |
| ■ Liver cancer | ■ Atrial fibrillation and flutter | |
| ■ Larynx cancer | ■ Cirrhosis and other chronic liver diseases | |
| ■ Breast cancer | ■ Pancreatitis | |
| ■ Colon and rectum cancer | ■ Epilepsy | |
| ■ Lip and oral cavity cancer | ■ Alcohol use disorders | |
| ■ Pharynx and nasopharynx cancer | ■ Diabetes mellitus | |
| ■ Ischaemic heart disease | ■ 2014 sport injuries | |


Attributable DALY rate disaggregated by cause, Antigua and Barbuda, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, The Bahamas, by sex and age in 2016.

Female


Male




Cause

2016

Attributable DALY rate disaggregated by cause, Barbados, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Belize, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Bermuda, by sex and age in 2016.

Female


Male



Attributable DALY rate disaggregated by cause, Cuba, by sex and age in 2016.

Female


Male




Cause

- | | | |
|---|---|---|
| ■ Tuberculosis | ■ Ischaemic stroke | ■ Unintentional injuries |
| ■ Lower respiratory infections | ■ Hemorrhagic stroke | ■ Self-harm |
| ■ Esophageal cancer | ■ Hypertensive heart disease | ■ Interpersonal violence |
| ■ Liver cancer | ■ Atrial fibrillation and flutter | |
| ■ Larynx cancer | ■ Cirrhosis and other chronic liver diseases | |
| ■ Breast cancer | ■ Pancreatitis | |
| ■ Colon and rectum cancer | ■ Epilepsy | |
| ■ Lip and oral cavity cancer | ■ Alcohol use disorders | |
| ■ Pharynx and nasopharynx cancer | ■ Diabetes mellitus | |
| ■ Ischaemic heart disease | ■ Transport injuries | |


Attributable DALY rate disaggregated by cause, Dominica, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Dominican Republic, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Grenada, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Guyana, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Haiti, by sex and age in 2016.

Female

Male




Attributable DALY rate disaggregated by cause, Jamaica, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Saint Lucia, by sex and age in 2016.

Female

Male




Attributable DALY rate disaggregated by cause, Saint Vincent and the Grenadines, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Suriname, by sex and age in 2016.

Female

Male




Attributable DALY rate disaggregated by cause, Trinidad and Tobago, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Bolivia, by sex and age in 2016.

Female

Male




- | | | |
|---|---|--|
| <ul style="list-style-type: none"> ■ Tuberculosis ■ Lower respiratory infections ■ Esophageal cancer ■ Liver cancer ■ Larynx cancer ■ Breast cancer ■ Colon and rectum cancer ■ Lip and oral cavity cancer ■ Pharynx and nasopharynx cancer ■ Ischaemic heart disease | <ul style="list-style-type: none"> ■ Ischaemic stroke ■ Hemorrhagic stroke ■ Hypertensive heart disease ■ Atrial fibrillation and flutter ■ Cirrhosis and other chronic liver diseases ■ Pancreatitis ■ Epilepsy ■ Alcohol use disorders ■ Diabetes mellitus ■ Transport injuries | <ul style="list-style-type: none"> ■ Unintentional injuries ■ Self-harm ■ Interpersonal violence |
|---|---|--|

Attributable DALY rate disaggregated by cause, Ecuador, by sex and age in 2016.

Female


Male




Attributable DALY rate disaggregated by cause, Peru, by sex and age in 2016.

Female


Male




Attributable DALY rate disaggregated by cause, Colombia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Costa Rica, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, El Salvador, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Guatemala, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Honduras, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Mexico, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Nicaragua, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Panama, by sex and age in 2016.

Female

Male




- Cause
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Transport injuries
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence


Attributable DALY rate disaggregated by cause, Venezuela, by sex and age in 2016.

Female


Male




Attributable DALY rate disaggregated by cause, Brazil, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Paraguay, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Afghanistan, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Algeria, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Bahrain, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Egypt, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Iran, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Iraq, by sex and age in 2016.

Female

Male




- Cause**
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence
 - Sport injuries

Attributable DALY rate disaggregated by cause, Jordan, by sex and age in 2016.

Female

Male




Cause


Attributable DALY rate disaggregated by cause, Kuwait, by sex and age in 2016.

Female

Male




Attributable DALY rate disaggregated by cause, Lebanon, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Libya, by sex and age in 2016.


Female

Male



- | | | |
|---|---|--|
| <ul style="list-style-type: none"> ■ Tuberculosis ■ Lower respiratory infections ■ Esophageal cancer ■ Liver cancer ■ Larynx cancer ■ Breast cancer ■ Colon and rectum cancer ■ Lip and oral cavity cancer ■ Pharynx and nasopharynx cancer ■ Ischaemic heart disease | <ul style="list-style-type: none"> ■ Ischaemic stroke ■ Hemorrhagic stroke ■ Hypertensive heart disease ■ Atrial fibrillation and flutter ■ Cirrhosis and other chronic liver diseases ■ Pancreatitis ■ Epilepsy ■ Alcohol use disorders ■ Diabetes mellitus ■ Transport injuries | <ul style="list-style-type: none"> ■ Unintentional injuries ■ Self-harm ■ Interpersonal violence |
|---|---|--|


Attributable DALY rate disaggregated by cause, Palestine, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Oman, by sex and age in 2016.

Female

Male




Attributable DALY rate disaggregated by cause, Qatar, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Sudan, by sex and age in 2016.


Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - Transport injuries
 - Interpersonal violence


Attributable DALY rate disaggregated by cause, Saudi Arabia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Syria, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Tunisia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Turkey, by sex and age in 2016.

Female


Male



Attributable DALY rate disaggregated by cause, United Arab Emirates, by sex and age in 2016.

Female


Male



Cause

- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
- Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Transport injuries
- Unintentional injuries
 - Self-harm
 - Interpersonal violence


Attributable DALY rate disaggregated by cause, Yemen, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Bangladesh, by sex and age in 2016.

Female

Male




Cause

- Tuberculosis
- Lower respiratory infections
- Esophageal cancer
- Liver cancer
- Larynx cancer
- Breast cancer
- Colon and rectum cancer
- Lip and oral cavity cancer
- Pharynx and nasopharynx cancer
- Ischaemic heart disease
- Ischaemic stroke
- Hemorrhagic stroke
- Hypertensive heart disease
- Atrial fibrillation and flutter
- Cirrhosis and other chronic liver diseases
- Pancreatitis
- Epilepsy
- Alcohol use disorders
- Diabetes mellitus
- Transport injuries
- Unintentional injuries
- Self-harm
- Interpersonal violence

Attributable DALY rate disaggregated by cause, Bhutan, by sex and age in 2016.

Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Atrial fibrillation and flutter
 - Interpersonal violence
 - Liver cancer
 - Hypertensive heart disease
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - Transport injuries

Attributable DALY rate disaggregated by cause, India, by sex and age in 2016.

Female

Male




- Cause
- Tuberculosis
 - Lower respiratory infections
 - Esophageal cancer
 - Liver cancer
 - Larynx cancer
 - Breast cancer
 - Colon and rectum cancer
 - Lip and oral cavity cancer
 - Pharynx and nasopharynx cancer
 - Ischaemic heart disease
 - Ischaemic stroke
 - Hemorrhagic stroke
 - Hypertensive heart disease
 - Atrial fibrillation and flutter
 - Cirrhosis and other chronic liver diseases
 - Pancreatitis
 - Epilepsy
 - Alcohol use disorders
 - Diabetes mellitus
 - Unintentional injuries
 - Self-harm
 - Interpersonal violence
 - Sport injuries

Attributable DALY rate disaggregated by cause, Nepal, by sex and age in 2016.


Female

Male



- | | | |
|---|---|--|
| <ul style="list-style-type: none"> ■ Tuberculosis ■ Lower respiratory infections ■ Esophageal cancer ■ Liver cancer ■ Larynx cancer ■ Breast cancer ■ Colon and rectum cancer ■ Lip and oral cavity cancer ■ Pharynx and nasopharynx cancer ■ Ischaemic heart disease | <ul style="list-style-type: none"> ■ Ischaemic stroke ■ Hemorrhagic stroke ■ Hypertensive heart disease ■ Atrial fibrillation and flutter ■ Cirrhosis and other chronic liver diseases ■ Pancreatitis ■ Epilepsy ■ Alcohol use disorders ■ Diabetes mellitus ■ Transport injuries | <ul style="list-style-type: none"> ■ Unintentional injuries ■ Self-harm ■ Interpersonal violence |
|---|---|--|


Attributable DALY rate disaggregated by cause, Pakistan, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Angola, by sex and age in 2016.

Female

Male




- Cause
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Interpersonal violence
 - Liver cancer
 - Atrial fibrillation and flutter
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - Traumatic transport injuries

Attributable DALY rate disaggregated by cause, Congo, by sex and age in 2016.


Female

Male




- | | | |
|---|---|---|
| <ul style="list-style-type: none"> ■ Tuberculosis ■ Lower respiratory infections ■ Esophageal cancer ■ Liver cancer ■ Larynx cancer ■ Breast cancer ■ Colon and rectum cancer ■ Lip and oral cavity cancer ■ Pharynx and nasopharynx cancer ■ Ischaemic heart disease | <ul style="list-style-type: none"> ■ Ischaemic stroke ■ Hemorrhagic stroke ■ Hypertensive heart disease ■ Atrial fibrillation and flutter ■ Cirrhosis and other chronic liver diseases ■ Pancreatitis ■ Epilepsy ■ Alcohol use disorders ■ Diabetes mellitus ■ Unintentional injuries | <ul style="list-style-type: none"> ■ Self-harm ■ Interpersonal violence |
|---|---|---|


Attributable DALY rate disaggregated by cause, Democratic Republic of the Congo, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Equatorial Guinea, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Gabon, by sex and age in 2016.




Cause

2146


Attributable DALY rate disaggregated by cause, Burundi, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Comoros, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Djibouti, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Eritrea, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Ethiopia, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Kenya, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Madagascar, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Malawi, by sex and age in 2016.


Female

Male



- | | |
|---|--|
| <p>Cause</p> <ul style="list-style-type: none"> Tuberculosis Lower respiratory infections Esophageal cancer Liver cancer Larynx cancer Breast cancer Colon and rectum cancer Lip and oral cavity cancer Pharynx and nasopharynx cancer Ischaemic heart disease | <ul style="list-style-type: none"> Ischaemic stroke Hemorrhagic stroke Hypertensive heart disease Atrial fibrillation and flutter Cirrhosis and other chronic liver diseases Pancreatitis Epilepsy Alcohol use disorders Diabetes mellitus Unintentional injuries Self-harm Interpersonal violence Sport injuries |
|---|--|


Attributable DALY rate disaggregated by cause, Mozambique, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Rwanda, by sex and age in 2016.


Female

Male




- Cause**
- Tuberculosis
 - Ischaemic stroke
 - Unintentional injuries
 - Lower respiratory infections
 - Hemorrhagic stroke
 - Self-harm
 - Esophageal cancer
 - Hypertensive heart disease
 - Liver cancer
 - Atrial fibrillation and flutter
 - Interpersonal violence
 - Larynx cancer
 - Cirrhosis and other chronic liver diseases
 - Breast cancer
 - Pancreatitis
 - Colon and rectum cancer
 - Epilepsy
 - Lip and oral cavity cancer
 - Alcohol use disorders
 - Pharynx and nasopharynx cancer
 - Diabetes mellitus
 - Ischaemic heart disease
 - Sport injuries


Attributable DALY rate disaggregated by cause, Somalia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, South Sudan, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Tanzania, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Zambia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Botswana, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Lesotho, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, South Africa, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Swaziland, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Benin, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Burkina Faso, by sex and age in 2016.


Female

Male



- | | | |
|---|---|---|
| <p>Cause</p> <ul style="list-style-type: none"> Tuberculosis Lower respiratory infections Esophageal cancer Liver cancer Larynx cancer Breast cancer Colon and rectum cancer Lip and oral cavity cancer Pharynx and nasopharynx cancer Ischaemic heart disease | <ul style="list-style-type: none"> Ischaemic stroke Hemorrhagic stroke Hypertensive heart disease Atrial fibrillation and flutter Cirrhosis and other chronic liver diseases Pancreatitis Epilepsy Alcohol use disorders Diabetes mellitus Sport injuries | <ul style="list-style-type: none"> Unintentional injuries Self-harm Interpersonal violence |
|---|---|---|


Attributable DALY rate disaggregated by cause, Cameroon, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Cape Verde, by sex and age in 2016.

Female

Male




- | | | |
|---|---|--|
| <ul style="list-style-type: none"> ■ Tuberculosis ■ Lower respiratory infections ■ Esophageal cancer ■ Liver cancer ■ Larynx cancer ■ Breast cancer ■ Colon and rectum cancer ■ Lip and oral cavity cancer ■ Pharynx and nasopharynx cancer ■ Ischaemic heart disease | <ul style="list-style-type: none"> ■ Ischaemic stroke ■ Hemorrhagic stroke ■ Hypertensive heart disease ■ Atrial fibrillation and flutter ■ Cirrhosis and other chronic liver diseases ■ Pancreatitis ■ Epilepsy ■ Alcohol use disorders ■ Diabetes mellitus ■ Transport injuries | <ul style="list-style-type: none"> ■ Unintentional injuries ■ Self-harm ■ Interpersonal violence |
|---|---|--|

Attributable DALY rate disaggregated by cause, Chad, by sex and age in 2016.


Female

Male




- | | | |
|---|--|--|
| <ul style="list-style-type: none"> ■ Tuberculosis ■ Lower respiratory infections ■ Esophageal cancer ■ Liver cancer ■ Larynx cancer ■ Breast cancer ■ Colon and rectum cancer ■ Lip and oral cavity cancer ■ Pharynx and nasopharynx cancer ■ Ischaemic heart disease | <ul style="list-style-type: none"> ■ Ischaemic stroke ■ Hemorrhagic stroke ■ Hypertensive heart disease ■ Atrial fibrillation and flutter ■ Cirrhosis and other chronic liver diseases ■ Pancreatitis ■ Epilepsy ■ Alcohol use disorders ■ Diabetes mellitus ■ Sports injuries | <ul style="list-style-type: none"> ■ Unintentional injuries ■ Self-harm ■ Interpersonal violence |
|---|--|--|


Attributable DALY rate disaggregated by cause, Cote d'Ivoire, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, The Gambia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Guinea, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Guinea-Bissau, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Liberia, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Mali, by sex and age in 2016.

Female


Male




Attributable DALY rate disaggregated by cause, Mauritania, by sex and age in 2016.




Attributable DALY rate disaggregated by cause, Niger, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Nigeria, by sex and age in 2016.



Attributable DALY rate disaggregated by cause, Sao Tome and Principe, by sex and age in 2016.



Cause

2183

