

Anti-inflammatory activity of a polymeric proanthocyanidin from *Serjania schiedeana*

David Osvaldo Salinas-Sánchez^{1,2}, Enrique Jiménez-Ferrer¹, Verónica Sánchez-Sánchez^{1,3}, Alejandro Zamilpa¹, Manasés González-Cortázar¹, Tortoriello J¹ and Maribel Herrera-Ruiz^{1,*}

¹ Biomedical Research Center of the South (IMSS), Argentina 1, Col. Centro, Xochitepec 62790, Morelos, Mexico; E-Mails: enriqueferrer_mx@yahoo.com (E.J.-F.); azamilpa_2000@yahoo.com.mx (A.Z.); gmanases@hotmail.com (M.G.-C.)

² Biodiversity and Conservation Research Center. (UAEM), Av. Universidad 1001, Col. Chamilpa, Cuernavaca 62209, Morelos, Mexico; E-Mail: davidos@uaem.mx (D.O.S.-S.)

³ Faculty of Biological Sciences (FBC), UAEM, Av. Universidad 1001, Col. Chamilpa, 62209 Cuernavaca, Morelos, Mexico; E-mail: vEra1o@outlook.es (V.S.S)

* Correspondence: cibus_herj@yahoo.com.mx; Tel.: +52-777-361-215-5




Fig. 1S ¹H NMR (CD₃OD, 400 MHz) Compound ETP.




Fig. 2S ^{13}C NMR (CD_3OD , 100 MHz) Compound ETP.




Fig 3S. ^1H – ^1H COSY (CD_3OD , 400 MHz) Compound ETP.




Fig4S. gHSQC (CD₃OD, 400 MHz) Compound ETP




Fig. 5S. HMBC (CD₃OD, 400 MHz) Compound ETP




Fig 6S. Mass spectra Compound ETP




Fig 7S. HPLC Analysis of Fraction F14-9