

Supplementary Info File

In-vivo shift of the microbiota in oral biofilm in response to frequent sucrose consumption

Annette Carola Anderson^{1**}, Michael Rothballer²⁺, Markus Jörg Altenburger¹, Johan Peter Woelber¹, Lamprini Karygianni³, Ilias Lagkouvardos⁴, Elmar Hellwig¹, Ali Al-Ahmad¹

¹Department of Operative Dentistry and Periodontology, Faculty of Medicine, Albert-Ludwigs-University, Freiburg, Germany

²Institute of Network Biology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany

³Clinic of Preventive Dentistry, Periodontology and Cariology, Center of Dental medicine, University of Zürich, Switzerland

⁴ZIEL - Institute for Food and Health, Core Facility Microbiome/NGS, Technical University of Munich, Freising, Germany

*ACA and MR contributed equally to this work

Address correspondence to:

Dr. Annette Anderson
Department of Operative Dentistry and Periodontology,
Hugstetter Straße 55
79106 Freiburg
Germany
annette.anderson@uniklinik-freiburg.de

Figure S1: Relative abundances

Relative abundances (%) of the different phyla found in all samples from the 11 study participants.




Figure S5: Abundances of *Streptococcus* spp.

Relative abundances of the different *Streptococcus* species of all samples from the 11 study participants. *Streptococcus* spp. refers to those OTUs which could not be unambiguously assigned to one of the oral *Streptococcus* species.

