

Table S2, related to Figure3. Primers used for analyzing changes in gene expression in mouse liver or fat tissues

	Forward	Reverse
Cyclophilin A	TGGAGAGCACCAAGACAGACA	TGCCGGAGTCGACAATGAT
Casp2	CACCCTCTTCAAGCTTTTGG	CGAAAACTCT TGGAGCTG
uPA	CTTCCCACTACCTTGGCTGG	CCAGCCAAGGTAGTGGGAAG
TNF α	ACGGCATGGATCTCAAAGAC	AGATAGCAAATCGGCTGACG
Adrp30	GCACTGGCAAGTTCTACTGCA A	GTAGGTGAAGAGAACGGCCTT GT
C/ebp β	ACGACTTCCTCTCCGACCTCT	CGAGGCTCACGTAACCGTAGT
PPAR γ	TGAAAGAAGCGGTGAACCACT G	TGGCATCTCTGTGTCAACCAT
UCP1	ACTGCCACACCTCCAGTCATT	CTTTGCCTCACTCAGGATTGG
α SMA	G TTCAGTGGTGCCTCTGTCA	ACTGGGACGACATGGAAAAG
Tgfb1	GGAGAGCCCTGGATACCAAC	AAGTTGGCATGGTAGCCCTT