

Supplementary materials: Factors Secreted by Cancer-Associated Fibroblasts that Sustain Cancer Stem Properties in Head and Neck Squamous Carcinoma Cells as Potential Therapeutic Targets

Saúl Álvarez-Teijeiro, Cristina García-Inclán, M. Ángeles Villaronga, Pedro Casado, Francisco Hermida-Prado, Rocío Granda-Díaz, Juan P. Rodrigo, Fernando Calvo, Nagore del-Río-Ibisate, Alberto Gandarillas, Francisco Moris, Mario Hermsen, Pedro Cutillas and Juana M. García-Pedrero

Table S1. Primers used for real-time RT-PCR (5'→3').

Genes	Forward	Reverse
<i>ABCG2</i>	ATGGATTTACGGCTTIGCAG	TCTTCGCCAGTACATGTTGC
<i>ALDH1</i>	AACAGTGTGGGTGAATTGCT	GGAAACCGTACTCTCCCAGT
<i>BMI1</i>	CTGGAGAAGGAATGGTCCAC	CACTTCCAGTCTCCAGCA
<i>CD133</i>	CAGTGAGAAAGTGGCATCGT	TCCTATGCCAAACCAAACA
<i>CD44</i>	GATCATCTTGGCATCCCTCT	TGAGTCCACTTGGCTTTCTG
<i>NANOG</i>	TGCTTATTCAGGACAGCCCT	TCTGGTCTTCTGTTTCTTACT
<i>NESTIN</i>	CAGCGTTGGAACAGAGGTTG	GCTGGCACAGGTGTCTCAAG
<i>NOTCH1</i>	AGACGGCATCAACACGGCCTTC	GTGTAGCTGTCCACGCAGTCCG
<i>OCT4</i>	GGCAACCTGGAGAATTTGTT	ACTCGGACCACATCCTTCTC
<i>RPL19</i>	GCGGAAGGGTACAGCCAAT	GCAGCCGGCGCAA
<i>SOX2</i>	TCAGGAGTTGTCAAGGCAGAGAAG	CTCAGTCCTAGTCTTAAAGAGGCAGC