VIROLOGICA SINICA

Electronic Supplementary Material

HearNPV Pseudotyped with PIF1, 2, and 3 from MabrNPV: Infectivity and Complex Stability

George Alliwa Makalliwa[1,2,3], Xi Wang [1,3], Huanyu Zhang[1,3], Nan Zhang[1,3], Cheng Chen[1,3], Jiang Li[1], Fei Deng[1], Hualin Wang[1], Manli Wang[1✉], Zhihong Hu[1✉]

1. State Key Laboratory of Virology, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan 430071, China;
2. Jomo Kenyatta University of Agriculture and Technology, P.O. Box 62000 – 00200, Nairobi, Kenya;
3. University of the Chinese Academy of Sciences, Beijing 100049, China

Supplementary Table S1. Primers used in this study.

Primer	Sequence (5′–3′)	Purpose
p*Hapif1*-F	TCC<u>CCCGGG</u>TTGTTTGGCTGTGTGTGTATCGTC	Amplifying *Mapif1* promoter region
p*Hapif1*-R	CCG<u>CTCGAG</u>GCATACATGTTACTTTATTCAACTTAG	
p*Hapif2*-F	TCC<u>CCCGGG</u>ACATGTTTGCAAGACCACATAAAACTGTAC	Amplifying *Hapif2* promoter region
p*Hapif2*-R	CCG<u>CTCGAG</u>CGTTTGATACTTAACAATTG	
p*Hapif3*-F	TCC<u>CCCGGG</u>AGACATTTTTGATATAAAATGATATTATATTATTAAAAATATAAAAATATGATGCAATAA	Amplifying *Hapif3* promoter region
p*Hapif3*-R	CCG<u>CTCGAG</u>GATGGTACGCTTGCTTATTCAG	
Mbpif1-F	CCG<u>CTCGAG</u>ATGTATTTAATCGTAGCTGCCGTTCTAC	Amplifying *Mbpif1* coding region
Mbpif1-FLAG-R	CATG<u>CCATGG</u>TTA*CTTGTCATCGTCGTCCTTGTAGTC*TGTTACCGAATAGTTTGCGTACGTCTTTAGAATGTCA	
MbPIF2-F	CCG<u>CTCGAG</u>ATGATCAATGCTGCATTGGGATTTATC	Amplifying *Mbpif2* coding region
MbPIF2-FLAG-R	CATG<u>CCATGG</u>TTA*CTTGTCATCGTCGTCCTTGTAGTC*AGTCTCGGGCTGTGGTCGCGGATATCCCAACGTAACTCTGTCGCGAATGTCCATC	
Mbpif3-F	CCG<u>CTCGAG</u>ATGTCGCCGACTACCATTT	Amplifying *Mbpif3* coding region
Mbpif3-FLAG-R	CATG<u>CCATGG</u>TTA*CTTGTCATCGTCGTCCTTGTAGTC*CACGACAGTAGTGTAGATCTTGCTGTACACATTTGCGGCTTG	