

Group	Total	MDR	Pyuria	Antibiotic use	LUTS	Chronic infection	Uncomplicated infection	Complicated infection	Pyelonephritis
All samples	78	45 (58%)	42 (54%)	36 (46%)	26 (33%)	34 (44%)	9 (12%)	63 (81%)	5 (6%)
<i>BF+</i>	41	14 (34%)*	23 (56%)	16 (39%)	14 (34%)	15 (37%)	6 (15%)	34 (83%)	1 (2%)
<i>BF-</i>	37	31 (84%)*	19 (51%)	20 (54%)	12 (32%)	19 (51%)	3 (8%)	29 (78%)	4 (11%)
Chronic	34	21 (62%)	18 (53%)	26 (76%)	13 (38%)	34 (100%)	3 (9%)	28 (82%)	3 (9%)
<i>BF+</i>	15	5 (33%)*	9 (60%)	12 (80%)	6 (40%)	15 (100%)	2 (13%)	12 (80%)	1 (7%)
<i>BF-</i>	19	16 (84%)*	9 (47%)	14 (74%)	7 (37%)	19 (100%)	1 (5%)	16 (84%)	2 (11%)
Symptomatic	26	16 (62%)	15 (58%)	14 (54%)	26 (100%)	13 (50%)	5 (19%)	21 (81%)	0 (0%)
<i>BF+</i>	14	5 (36%)	6 (43%)	6 (43%)	14 (100%)	6 (43%)	3 (21%)	11 (79%)	0 (0%)
<i>BF-</i>	12	11 (92%)	9 (75%)	8 (67%)	12 (100%)	7 (58%)	2 (17%)	10 (83%)	0 (0%)
Asymptomatic	40	22 (55%)	26 (65%)	18 (45%)	0 (0%)	16 (40%)	4 (10%)	31 (78%)	5 (12%)
<i>BF+</i>	23	6 (26%)*	17 (74%)	9 (39%)	0 (0%)	8 (35%)	3 (13%)	19 (83%)	1 (4%)
<i>BF-</i>	17	16 (94%)*	9 (53%)	9 (53%)	0 (0%)	8 (47%)	1 (6%)	12 (71%)	4 (24%)