

Evaluation of a community intervention to reduce exposure of young children living in farmworker households to organophosphate pesticides

Authors

William C. Griffith, Ph.D.<sup>a,b</sup>, Eric M. Vigoren, MS<sup>a,b</sup>, Marissa N. Smith, MS<sup>a,b\*</sup>, Tomomi Workman, BS<sup>a,b\*</sup>, Beti Thompson, Ph.D.<sup>c</sup>, Gloria D. Coronado, Ph.D.<sup>d</sup>, Elaine M. Faustman, Ph.D.<sup>a,b</sup>

<sup>a</sup>Department of Environmental and Occupational Health Sciences, School of Public Health, University of Washington, Seattle, WA

<sup>b</sup>Washington Institute for Risk Analysis & Risk Communication, University of Washington, Seattle, WA

<sup>c</sup>Cancer Prevention Program, Fred Hutchinson Cancer Research Center, Seattle, WA

<sup>d</sup>Kaiser Permanente Center for Health Research, Portland, OR

Correspondence to:

Elaine M. Faustman

Department of Environmental and Occupational Health Sciences

School of Public Health

University of Washington

4225 Roosevelt Way NE Suite 100

Seattle, WA 98105

Fax: (206) 685-4696

E-mail: faustman@uw.edu

Running Title: Intervention reducing child exposure

[This publication was made possible by grants PO1 ES009601 from NIEHS, and EPA RD-83170901, RD826886, *RD83451401*, RD-83273301.]

Supplemental Information

Table S1: Total Active Ingredients Applied in Washington State in 1999, 2001 and 2003.  
Data are shown in pounds.

	1999	2001	2003
azinphos-methyl-apple	309,300	241,400	289,200
Phosmet-apple	46,000	138,300	70,700
Malathion-apple	22,300	1,600	Not Available
azinphos-methyl-pear	33,000	22,800	28,700
Phosmet-pear	11,500	29,900	18,800
Malathion-pear	Not available	Not available	Not available
Total	422,100	434,000	407,400