

S2 Table. Groups for preliminary analysis of MARCH1 SNPs and type 2 diabetes, grouped by race/ethnicity and genotyping platform

	White, not Hispanic (n=6892)								Black, not Hispanic (n=1596)			
	Vanderbilt ^a /660W ^b (n=139)		Northwestern/660W (n=1134)		HPFS/Affy (n=2397)		NHS/Affy (n=3222)		Vanderbilt/1M (n=1354)		Northwestern/1M (n=242)	
	Cases	Controls	Cases	Controls	Cases	Controls	Cases	Controls	Cases	Controls	Cases	Controls
Cases (%)	99 (71.2)	40	530 (46.7)	604	1116 (46.6)	1281	1466 (45.5)	1756	598 (44.2)	756	153 (63.2)	89
Females (%)	45 (45.5)	22 (55.0)	233 (44.0)	337 (55.8)	0 (0.0)	0 (0.0)	1466 (100.0)	1756 (100.0)	373 (62.4)	501 (66.3)	97 (63.4)	68 (76.4)
Family history of diabetes												
No (%)	47 (47.5)	28 (70.0)	0	0	635 (56.9)	1009 (78.8)	741 (50.5)	1370 (78.0)	128 (21.4)	479 (63.4)	0	0
Yes (%)	48 (48.5)	5 (12.5)	460 (86.8)	0	481 (43.1)	272 (21.2)	725 (49.5)	386 (22.0)	456 (76.3)	204 (27.0)	143 (93.5)	0
Unknown/missing (%)	4 (4.0)	7 (17.5)	70 (13.2)	604 (100.0)	0	0	0	0	14 (2.3)	73 (9.7)	10 (6.5)	89 (100.0)

HPFS, Health Professional Follow-up Study; NHS, Nurses' Health Study

^aIndicates study

^bIndicates genotyping platform