

1 Assessing scale-wise similarity of curves with a thick pen:
2 as illustrated through comparisons of spectral irradiance

3 Saara M. Hartikainen¹, Agnieszka Jach², Aurea Grané³,

T. Matthew Robson^{1,*}

4 ¹*University of Helsinki, Viikki Plant Science Centre, PB 65, Helsinki, Finland;* ²*Department of*
5 *Finance and Statistics, Hanken School of Economics, PB 479, Helsinki, Finland;* ³*Department of*
6 *Statistics, Universidad Carlos III de Madrid, C/ Madrid, 126, Getafe, Spain*

7 **Appendix: Graphs**

*Correspondence author. E-mail: matthew.robson@helsinki.fi




Figure A.1: TPMA for spectral irradiance from positions sunfleck and full-sun, for a given stand and for a given date.




Figure A.2: TPMA for spectral irradiance from positions sunfleck and full-sun, averaged over wavelengths, for a given stand and for a given date.




Figure A.3: TPMA for spectral irradiance from positions semi-shade and full-sun, for a given stand and for a given date. No data for semi-shade in Quercus.




Figure A.4: TPMA for spectral irradiance from positions semi-shade and full-sun, averaged over wavelengths, for a given stand and for a given date.




Figure A.5: TPMA for spectral irradiance from positions shade and full-sun, for a given stand and for a given date.




Figure A.6: TPMA for spectral irradiance from positions shade and full-sun, averaged over wavelengths, for a given stand and for a given date.