

The increase of triterpene saponin production induced by trans-anethole in hairy root cultures of *Panax quinquefolium*

Ewa Kochan ^{1,*}, Piotr Szymczyk ¹, Łukasz Kuźma ², Grażyna Szymańska ¹, Anna Wajs-Bonikowska ³, Radosław Bonikowski ³, Monika Sienkiewicz ⁴

¹ Pharmaceutical Biotechnology Department, Medical University of Lodz, Muszyńskiego 1, Lodz 90-151, Poland; ewa.kochan@umed.lodz.pl (E.K.); piotr.szymczyk@umed.lodz.pl (P.S.); grazyna.szymanska@umed.lodz.pl (G.S.)

² Department of Biology and Pharmaceutical Botany, Medical University of Lodz, Muszyńskiego 1, Lodz 90-151, Poland; lukasz.kuzma@umed.lodz.pl

³ Institute of General Food Chemistry, Biotechnology and Food Science, Lodz University of Technology, Stefanowskiego St. 4/10, 90-924 Lodz, Poland; anna.wajs-bonikowska@p.lodz.pl (A.W.-B.); radoslaw.bonikowski@p.lodz.pl (R.B.)

⁴ Department of Allergology and Respiratory Rehabilitation, 2nd Chair of Otolaryngology Medical University of Lodz, Poland; monika.sienkiewicz@umed.lodz.pl

* correspondence: ewa.kochan@umed.lodz.pl




Figure S1. Schematic representation of the possible responses of cells to elicitation. R: receptor; PL: phospholipase; MAPKs: mitogen activated protein kinases; ROS: reactive oxygen species; RNS: reactive nitrogen species; TF: transcription factors [according to 9].