

Evaluating Marine Cyanobacteria as a Source for CNS Receptor Ligands

Andrea L. Rague, Stacy-Ann J. Parker, and Kevin J. Tidgewell

Department of Medicinal Chemistry, Mylan School of Pharmacy, Duquesne University,
Pittsburgh, PA 15282, United States

Contents of Supporting Information

Contents	page
Figure S1. ^1H NMR spectrum of compound 2 in CDCl_3 at 500 MHz	2
Figure S2. COSY spectrum of compound 2 in CDCl_3 at 500 MHz	3
Figure S3. HSQC spectrum of compound 2 in CDCl_3 at 500 MHz	4
Figure S4. HMBC spectrum of compound 2 in CDCl_3 at 500 MHz	5
Figure S5. GCMS spectrum of 2	6




Figure S1. ¹H NMR spectrum of compound 2 in CDCl₃ at 500 MHz




Figure S2. COSY spectrum of compound 2 in CDCl₃ at 500 MHz




Figure S3. HSQC spectrum of compound 2 in CDCl₃ at 500 MHz




Figure S4. HMBC spectrum of compound 2 in CDCl₃ at 500 MHz




Figure S5. GC-MS chromatogram and mass spectrum of compound 2. (A) TIC, total ion chromatogram, of 1 and (B) mass spectrum of compound 2.