

SUPPLEMENTAL MATERIAL

Table S1. Temporal trends in clinical outcomes in patients who survived the index ACS hospitalization.

	2008	2010	2013	2016	P trend
Entire cohort					
N	1673	1686	1632	1695	
30 day MACE	172 (10.3)	139 (8.2)	144 (8.8)	123 (7.3)	0.006
30 day mortality	29 (1.7)	34 (2.0)	29 (1.8)	23 (1.4)	0.334
1 year mortality	92 (5.6)	99 (5.9)	103 (6.4)	103 (6.2)	0.387
Low-risk					
N	639	597	566	605	
30 day MACE	39 (6.1)	34 (5.7)	43 (7.6)	46 (7.6)	0.1
30 day mortality	5 (0.8)	2 (0.3)	3 (0.5)	5 (0.8)	0.8
1 year mortality	6 (1.0)	7 (1.2)	11 (2.0)	10 (1.7)	0.1
Intermediate-risk					
N	429	457	433	453	
30 day MACE	35 (8.2)	31 (6.8)	36 (8.3)	28 (6.2)	0.4
30 day mortality	4 (0.9)	3 (0.7)	6 (1.4)	5 (1.1)	0.5
1 year mortality	13 (3.1)	15 (3.3)	20 (4.7)	16 (3.6)	0.4
High-risk					
N	605	632	633	637	
30 day MACE	98 (16.2)	74 (11.7)	65 (10.3)	49 (7.7)	<0.001
30 day mortality	20 (3.3)	29 (4.6)	20 (3.2)	13 (2.1)	0.1
1 year mortality	73 (12.1)	77 (12.2)	72 (11.6)	77 (12.3)	0.9

Values are presented as n (%).

DAPT: Dual antiplatelet therapy; MACE: Death/Unstable angina pectoris/MI/CVA/Stent thrombosis/Urgent revascularization.