

Supplementary Online Content

Lee S, Kim BJ, Lee YB, Lee WS. Hair Regrowth Outcomes of Contact Immunotherapy for Patients With Alopecia Areata: A Systematic Review and Meta-analysis. *JAMA Dermatol.* Published online August 01, 2018. doi:10.1001/jamadermatol.2018.2312

Supplement. eText 1. Search strategy

eFigure 1. Literature search

eTable 1. Definition of therapeutic regrowth

eTable 2. Study characteristics and quality assessment

eTable 3. Sensitivity analysis

eTable 4. Treatment-related adverse effects

eReferences

This supplementary material has been provided by the authors to give readers additional information about their work.

eText 1. Search strategy

MEDLINE

(“areata”[title] OR “totalis”[title] OR “universalis”[title]) AND
 (“sensitizer”[title] OR “sensitization”[title] OR “immunotherapy”[title] OR
 “Diphenylcyclopropenone”[title] OR “diphenylcyclopropenone”[title] OR “diphencyprone”[title] OR
 “Squaric acid dibutyl ester”[title] OR “squaric”[title])

Embase

(‘areata’:ti OR ‘totalis’:ti OR ‘universalis’:ti) AND
 (‘sensitizer’:ti OR ‘sensitization’:ti OR ‘immunotherapy’:ti OR
 ‘Diphenylcyclopropenone’:ti OR ‘diphenylcyclopropenone’:ti OR ‘diphencyprone’:ti OR
 ‘Squaric acid dibutyl ester’:ti OR ‘squaric’:ti)


Cochrane Library

#1: ‘areata’:ti OR ‘totalis’:ti OR ‘universalis’:ti

#2: ‘sensitizer’:ti OR ‘sensitization’:ti OR ‘immunotherapy’:ti OR
 ‘Diphenylcyclopropenone’:ti OR ‘diphenylcyclopropenone’:ti OR ‘diphencyprone’:ti OR
 ‘Squaric acid dibutyl ester’:ti OR ‘squaric’:ti

Final query: #1 AND #2

eFigure 1. Literature search



PRISMA flow diagram of literature search and study selection for the systematic review of studies on contact immunotherapy for patients with alopecia areata.

eTable 1. Definition of therapeutic regrowth

Grade	Quantitative criteria ^a	Qualitative criteria
Any regrowth	<10% regrowth	Any or little regrowth Hull and Norris ²⁹ grade I (vellus hair) Any improvement in Hull and Norris grade No clear definition of therapeutic regrowth
Minor regrowth	10–59% regrowth	Partial, some, good, or moderate regrowth Cosmetically unacceptable regrowth Hull and Norris grade II (sparse terminal hair)
Major regrowth	60–89% regrowth	Diffuse or persistent regrowth Cosmetically acceptable regrowth Hull and Norris grade III (terminal hair with patchy alopecia)
Complete regrowth	90–100% regrowth	Complete or excellent regrowth Hull and Norris grade IV (terminal hair all over the head)

^aOnly studies using quantitative criteria for therapeutic regrowth were included in the sensitivity analysis (see eTable 3).

eTable 2. Study characteristics and quality assessment

Study name	Study setting	Sensitizer	Study population	Total, n	Female, n	Age, y	AT/AU, n	Follow-up, m	Any regrrowth	Minor regrrowth	Major regrrowth	Complete regrowth	SA LT ≥ 50	Duration ≥ 1 y	Atopic disease	Nail involvement	Adverse effects	Recurrence rate
Thuangtong et al ³⁰	Thailand, 2017	Diphenylcyclopropenone	Recalcitrant AT/AU	20	12	35	20	2.7	-	-	O	-	O	-	-	-	O	-
Tiwary et al ³¹	India, 2016	Diphenylcyclopropenone, squaric acid dibutyl ester	AA (>20%)	24	10	NA	0	6	-	O	O	-	O	-	-	-	O	-
Lamb et al ³²	UK, 2016	Diphenylcyclopropenone	AA	133	99	35	53	NA	O	-	-	O	O	O	O	O	O	-
Durdur et al ³³	Turkey, 2015	Diphenylcyclopropenone	Chronic, extensive, recalcitrant AA	22	9	17	9	12	-	O	O	-	O	-			O	O
Chiang et al ³⁴	US, 2015	Diphenylcyclopropenone	AA	50	36	37.5	39	36	-	O	-	-	O	-			O	O
Luk et al ³⁵	China, 2013	Diphenylcyclopropenone	AA	29	15	28.9	15	NA	O	-	O	O	O	O	O	O	O	O
Khoury et al ³⁶	Lebanon, 2013	Diphenylcyclopropenone	AA	34	15	24.72	14	16	-	-	-	-	O	-	-	-	O	O
Dhurat et al ³⁷	India, 2013	Diphenylcyclopropenone	AA with at least two patches	29	NA	NA	0	6	O	-	-	-	O	-	-	-	-	-

Study name	Study setting	Sensitizer	Study population	Total, n	Female, n	Age, y	AT/AU, n	Follow-up, m	Any regrrowth	Minor regrrowth	Major regrrowth	Complete regrowth	SA LT ≥ 50	DURATION ≥ 1 y	Atopic disease	Nail involvement	Adverse effects	Recurrence rate
Ohlmeier et al ³⁸	Germany, 2012	Diphenylcyclopropenone	AA	135	95	36	24	20	O	-	-	-	O	-	-	-	-	O
Khan et al ³⁹	UK, 2011	Diphenylcyclopropenone	AA	60	43	NA	22	10.5	O	O	-	-	O	-	-	-	O	O
Bhat et al ⁴⁰	UK, 2011	Diphenylcyclopropenone	Recalcitrant AA	29	23	NA	14	12	-	O	-	-	O	-	-	-	-	O
Avgerinou et al ⁴¹	Greece, 2007	Diphenylcyclopropenone	Extensive or chronic (>25% or >1 year)	54	15	27.2	6	6.1	-	-	-	-	O	-	-	-	O	O
Sotiridis et al ⁴²	Greece, 2006	Diphenylcyclopropenone	Severe AA (>50%)	38	19	24.5	16	12	-	O	O	O	O	O	O	-	O	O
Ajith et al ⁴³	India, 2006	Squaric acid dibutyl ester	Recalcitrant AA	60	32	25	20	3.2	-	O	-	-	O	-	O	O	O	O
Kwon et al ⁴⁴	Korea, 2005	Squaric acid dibutyl ester	AA (>25%)	22	12	27.8	7	6	O	-	-	-	O	-	-	-	O	O
Dall'oglio et al ⁴⁵	Italy, 2005	Squaric acid dibutyl ester	Recalcitrant AA (>40%)	54	26	25	23	8	-	-	O	O	O	-	-	-	-	O
Mun et al ⁴⁶	Korea, 2004	Diphenylcyclopropenone	Recalcitrant AA	36	23	NA	10	NA	O	-	O	O	O	-	O	O	O	-

Study name	Study setting	Sensitizer	Study population	Total, n	Female, n	Age, y	AT/AU, n	Follow-up, m	Any regrrowth	Minor regrrowth	Major regrrowth	Complete regrowth	SA LT ≥ 50	DURATION ≥ 1 y	Atopic disease	Nail involvement	Adverse effects	Recurrence rate
			(>25%)															
Lee et al ⁴⁷	Korea, 2004	Diphenylcyclopropenone	Extensive AA (>50%)	39	17	25	16	5.8	-	-	O	O	O	O	-	-	O	O
Wisman et al ⁴⁸	Canada, 2001	Diphenylcyclopropenone	AA (>25%)	148	91	36.3	35	7.2	-	O	-	-	O	-	-	-	O	O
Pardasani et al ⁴⁹	US, 2001	Squaric acid dibutyl ester	AA	36	22	27.2	8	7	O	-	-	-	O	-	-	O	-	-
Pericin et al ⁵⁰	Switzerland, 1998	Diphenylcyclopropenone	Severe AA (>40%)	68	36	NA	52	15.8	O	-	O	O	O	-	-	-	O	O
Iijima et al ¹³	Japan, 1997	Squaric acid dibutyl ester	AA (>25%)	48	32	NA	14	6	O	-	-	-	O	O	O	-	-	-
Amer et al ⁵¹	Egypt, 1997	Diphenylcyclopropenone	AA	49	21	NA	27	NA	O	O	O	O	O	-	-	-	O	-
Weise et al ⁵²	Germany, 1996	Diphenylcyclopropenone	Severe AA	105	78	32	62	NA	-	O	-	-	O	-	O	O	-	-
Tosti et al ⁵³	Italy, 1996	Squaric acid dibutyl ester	Pediatric severe AA	33	13	9.5	33	NA	O	-	O	O	O	-	O	O	O	O
Schutteelaar et al ²⁴	Netherlands, 1996	Diphenylcyclopropenone	Pediatric AA	25	14	10.7	15	10.8	O	-	O	O	O	O	O	-	-	-
Micali	Italy,	Squaric acid	Recal	14	77	23	15	12	-	-	-	-	O	-	-	-	-	-

Study name	Study setting	Sensitizer	Study population	Total, n	Female, n	Age, y	AT/AU, n	Follow-up, m	Any regrrowth	Minor regrrowth	Major regrrowth	Complete regrowth	SA LT ≥ 50	DURATION ≥ 1 y	Atopic disease	Nail involvement	Adverse effects	Recurrence rate
et al ⁵⁴	1996	dibutyl ester	citrant AA	4														
van Baar et al ⁵⁵	Netherlands, 1995	Diphenylcyclopropenone	Severe AA	13 9	73	NA	85	43	-	-	O	O	O	-	-	-	O	-
Orechchia et al ⁵⁶	Italy, 1994	Squaric acid dibutyl ester	Pediatric recalcitrant AA (>75%)	28	16	10	11	NA	O	O	-	-	O	O	O	-	O	O
Shapiro et al ⁵⁷	Canada, 1993	Diphenylcyclopropenone	AA (>50%)	7	4	32.7	4	NA	-	O	-	-	O	O	O	O	-	O
van der Steen et al ⁵⁸	Netherlands, 1992	Diphenylcyclopropenone	AA (>40%)	13 9	NA	NA	85	NA	-	O	-	-	O	O	-	O	-	-
Hull et al ²¹	UK, 1991	Diphenylcyclopropenone	Pediatric severe AA	12	4	11.2	8	7.3	O	-	O	O	O	O	O	-	O	O
Hull et al ⁵⁹	UK, 1991	Diphenylcyclopropenone	AA (>6 months)	78	61	NA	45	8	-	-	O	O	O	-	-	-	-	-
Chun et al ⁶⁰	Korea, 1991	Diphenylcyclopropenone	Recalcitrant AA	20	6	20.1	10	5	O	-	O	O	O	O	-	-	O	-
Berth-Jones	UK, 1990	Diphenylcyclopropenone	AT (>1 year)	11	6	39	11	6	O	-	-	-	O	O	-	-	-	-

Study name	Study setting	Sensitizer	Study population	Total, n	Female, n	Age, y	AT/AU, n	Follow-up, m	Any regrrowth	Minor regrrowth	Major regrrowth	Complete regrowth	SA LT ≥ 50	DURATION ≥ 1 y	Atopic disease	Nail involvement	Adverse effects	Recurrence rate
et al ⁶¹																		
Monk et al ⁶²	UK, 1989	Diphenylcyclopropenone	Extensive AA (>50%, >5 years)	18	10	26.9	14	NA	-	O	-	-	O	O	-	-	O	-
Hull et al ⁶³	UK, 1988	Diphenylcyclopropenone	Extensive AA (>1 year)	28	25	NA	13	8	-	-	-	-	O	O	-	-	-	-
Hatzis K et al ¹⁶	Greece, 1988	Diphenylcyclopropenone	Extensive AA	45	19	25.4	22	NA	O	O	-	-	O	-	-	-	O	-
Caserio et al ⁶⁴	US, 1987	Squaric acid dibutyl ester	Recalcitrant AA	14	11	16.5	12	7.6	-	O	-	-	O	-	-	-	O	-
Valsecchi et al ⁶⁵	Italy, 1986	Squaric acid dibutyl ester	Recalcitrant AA	45	NA	NA	23	NA	-	-	-	-	O	-	-	-	-	O
Valsecchi et al ⁶⁶	Italy, 1985	Squaric acid dibutyl ester	AA (>30%)	38	17	NA	20	NA	-	O	-	-	O	-	-	-	O	-
Orechchia et al ¹²	Italy, 1985	Diphenylcyclopropenone	AA (>50%)	26	11	NA	17	5.1	O	-	O	O	O	O	O	O	O	O
Case et al ⁶⁷	US, 1984	Squaric acid dibutyl ester	AA (>30%)	21	11	34	10	3	-	O	O	O	O	-	-	-	O	O
Giannetti et al ⁶⁸	Italy, 1983	Squaric acid dibutyl ester	AA	26	11	20	6	3.6	-	O	O	O	O	O	O	-	O	O

Study name	Study setting	Sensitizer	Study population	Total, n	Female, n	Age, y	AT/AU, n	Follow-up, m	Any regrowth	Minor regrowth	Major regrowth	Complete regrowth	SA LT ≥ 50	DURATION ≥ 1 y	Atopic disease	Nail involvement	Adverse effects	Recurrence rate
Flowers et al ⁶⁹	US, 1982	Squaric acid dibutyl ester	AA (>30%)	8	5	33.9	5	2.3	O	-	O	O	O	O	-	-	O	-

AT, alopecia totalis; AU, alopecia universalis; AA, alopecia areata; SALT, The Severity of Alopecia Tool; NA, not available

eTable 3. Sensitivity analysis

Original analysis (both quantitative and qualitative studies)								
AA subtype	Any regrowth		Minor regrowth		Major regrowth		Complete regrowth	
	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)
PA	74.6 (67.9–80.3)	15 (459)	64.9 (57.7–71.6)	20 (409)	56.1 (45.9–65.8)	18 (430)	42.6 (29.9–56.3)	17 (371)
AT/AU	54.5 (46.2–62.6)	16 (285)	45.0 (35.9–54.4)	20 (357)	28.7 (24.1–33.8)	18 (405)	24.9 (18.2–33.0)	18 (443)
Overall	65.5 (59.1–71.3)	16 (744)	55.5 (49.0–61.9)	22 (766)	39.4 (31.8–47.6)	19 (835)	32.3 (25.3–40.2)	18 (814)
Sensitivity analysis (quantitative studies only)								
AA subtype	-		10–59% regrowth		60–89% regrowth		90–100% regrowth	
	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)	Response rate (%) (95% CI [%])	No. of studies, (No. of patients)
PA	-	-	65.1 (56.5–72.7)	11 (275)	52.2 (23.2–79.8)	4 (41)	43.5 (26.4–62.4)	7 (112)
AT/AU	-	-	52.0 (37.5–66.1)	9 (184)	23.3 (15.4–33.8)	5 (83)	27.7 (20.4–36.4)	7 (126)
Overall	-	-	59.8 (52.0–67.2)	11 (459)	39.1 (27.5–52.2)	5 (124)	34.4 (21.8–49.7)	7 (238)

AA, alopecia areata; 95% CI, 95% confidential interval; PA, patchy alopecia; AT, alopecia totalis; AU, alopecia universalis

eTable 4. Treatment-related adverse effects

Adverse effects	Incidence rate (%) (95% confidence interval [%])	No. of studies	No. of patients
Severe eczema	30.8 (22.7–40.3)	13	1220
Lymphadenopathy	25.7 (18.2–35.0)	24	1213
Generalized eczema	15.8 (10.1–23.8)	17	814
Hyperpigmentation	12.7 (4.7–30.2)	5	304
Influenza-like symptoms	11.1 (5.2–21.9)	7	239
Contact urticaria	9.0 (5.2–15.2)	7	295
Folliculitis	7.1 (3.6–13.6)	4	116
Hypopigmentation	5.5 (3.0–9.7)	11	549
Erythema multiforme-like eruption	1.7 (0.6–4.4)	3	246

eReferences

29. Cotellessa C, Peris K, Caracciolo E, Mordenti C, Chimenti S. The use of topical diphenylcyclopropenone for the treatment of extensive alopecia areata. *J Am Acad Dermatol.* 2001;44(1):73-76.
30. Thuangtong R, Varothai S, Triwongwaranat D, Rujitharanawong C. Multi-concentration level patch test guided diphenyl cyclopropenone (Diphenylcyclopropenone) treatment in alopecia totalis or alopecia universalis. *J Med Assoc Thai.* 2017;100(1):86-92.
31. Tiwary AK, Mishra DK, Chaudhary SS. Comparative study of efficacy and safety of topical squaric acid dibutylester and diphenylcyclopropenone for the treatment of alopecia areata. *N Am J Med Sci.* 2016;8(6):237-242.
32. Lamb RC, Young D, Holmes S. Retrospective review of dipencyprone in the treatment of alopecia areata. *Clin Exp Dermatol.* 2016;41(4):352-358.
33. Durdu M, Ozcan D, Baba M, Seckin D. Efficacy and safety of diphenylcyclopropenone alone or in combination with anthralin in the treatment of chronic extensive alopecia areata: a retrospective case series. *J Am Acad Dermatol.* 2015;72(4):640-650.
34. Chiang KS, Mesinkovska NA, Piliang MP, Bergfeld WF. Clinical efficacy of diphenylcyclopropenone in alopecia areata: retrospective data analysis of 50 patients. *J Investig Dermatol Symp Proc.* 2015;17(2):50-55.
35. Luk NM, Chiu LS, Lee KC, et al. Efficacy and safety of diphenylcyclopropenone among Chinese patients with steroid resistant and extensive alopecia areata. *J Eur Acad Dermatol Venereol.* 2013;27(3):e400-405.
36. El Khoury J, Abd-el-Baki J, Succariah F, Abbas O, Kibbi AG, Kurban M. Topical immunomodulation with diphenylcyclopropenone for alopecia areata: the Lebanese experience. *Int J Dermatol.* 2013;52(12):1551-1556.
37. Dhurat R, Pund P, Ghate S. Dipencyprone (Diphenylcyclopropenone) in treatment of patchy alopecia areata (AA). *J Investig Dermatol.* 2013;133(5):1392.
38. Ohlmeier MC, Traupe H, Luger TA, Bohm M. Topical immunotherapy with diphenylcyclopropenone of patients with alopecia areata--a large retrospective study on 142 patients with a self-controlled design. *J Eur Acad Dermatol Venereol.* 2012;26(4):503-507.
39. Khan M, Gach JE. A retrospective study of dipencyprone therapy for alopecia areata in 60 patients. *Br J Dermatol.* 2011;165(Suppl. 1):45.
40. Bhat A, Sripathy K, Wahie S, Carr M. Efficacy and cost-efficiency of dipencyprone for alopecia areata. *Br J Dermatol.* 2011;165:43-44.
41. Avgerinou G, Gregoriou S, Rigopoulos D, Stratigos A, Kalogeromitros D, Katsambas A. Alopecia areata: topical immunotherapy treatment with dipencyprone. *J Eur Acad Dermatol Venereol.* 2008;22(3):320-323.
42. Sotiriadis D, Patsatsi A, Lazaridou E, Kastanis A, Vakirlis E, Chrysomallis F. Topical immunotherapy with diphenylcyclopropenone in the treatment of chronic extensive alopecia areata. *Clin Exp Dermatol.* 2007;32(1):48-51.
43. Ajith C, Gupta S, Kanwar AJ. Efficacy and safety of the topical sensitizer squaric acid dibutyl ester in alopecia areata and factors influencing the outcome. *J Drugs Dermatol.* 2006;5(3):262-266.
44. Kwon KS, Kim MB, Kwon YW, Park HJ, Oh CK, Jang HS. Squaric acid dibutyl ester (squaric acid dibutylester) immunotherapy in alopecia areata. *Korean J Dermatol.* 2005;43(5):630-637.
45. Dall'oglio F, Nasca MR, Musumeci ML, et al. Topical immunomodulator therapy with squaric acid dibutylester (Squaric acid dibutyl ester) is effective treatment for severe alopecia areata (AA): results of an open-label, paired-comparison, clinical trial. *J Dermatolog Treat.* 2005;16(1):10-14.
46. Mun JH, Ihm CW. Diphenylcyclopropenone (diphenylcyclopropenone) immunotherapy in severe alopecia areata - Clinical study of 36 cases. *Korean J Dermatol.* 2004;42(6):710-717.
47. Lee SG, Shin JH, Choi GS. Therapeutic effect of topical diphenylcyclopropenone (Diphenylcyclopropenone) for the treatment of extensive alopecia areata. *Korean J Dermatol.* 2004;42(9):1130-1137.
48. Wiseman MC, Shapiro J, MacDonald N, Lui H. Predictive model for immunotherapy of alopecia areata with dipencyprone. *Arch Dermatol.* 2001;137(8):1063-1068.
49. Pardasani AG, Turner E, McMichael AJ. Squaric acid dibutylester: indications for use and efficacy in alopecia areata. *Arch Dermatol.* 2001;137(7):970-972.
50. Pericin M, Trueb RM. Topical immunotherapy of severe alopecia areata with diphenylcyclopropenone: evaluation of 68 cases. *Dermatology (Basel, Switzerland).* 1998;196(4):418-

51. Amer M, Metwalli M, Tosson Z. Successful treatment of alopecia areata and alopecia totalis with diphencyprone. *J Eur Acad Dermatol Venereol*. 1997;9(1):83-84.
52. Weise K, Kretzschmar L, John SM, Hamm H. Topical immunotherapy in alopecia areata: anamnestic and clinical criteria of prognostic significance. *Dermatology (Basel, Switzerland)*. 1996;192(2):129-133.
53. Tosti A, Guidetti MS, Bardazzi F, Misciali C. Long-term results of topical immunotherapy in children with alopecia totalis or alopecia universalis. *J Am Acad Dermatol*. 1996;35(2 Pt 1):199-201.
54. Micali G, Cicero RL, Nasca MR, Sapuppo A. Treatment of alopecia areata with squaric acid dibutylester. *Int J Dermatol*. 1996;35(1):52-56.
55. van Baar HM, van der Vleuten CJ, van de Kerkhof PC. Dapsone versus topical immunotherapy in alopecia areata. *Br J Dermatol*. 1995;133(2):270-274.
56. Orecchia G, Malagoli P, Santagostino L. Treatment of severe alopecia areata with squaric acid dibutylester in pediatric patients. *Pediatr Dermatol*. 1994;11(1):65-68.
57. Shapiro J, Tan J, Ho V, Abbott F, Tron V. Treatment of chronic severe alopecia areata with topical diphenylcyclopropenone and 5% minoxidil: a clinical and immunopathologic evaluation. *J Am Acad Dermatol*. 1993;29(5 Pt 1):729-735.
58. van der Steen PH, Boezeman JB, Happé R. Topical immunotherapy for alopecia areata: re-evaluation of 139 cases after an additional follow-up period of 19 months. *Dermatology (Basel, Switzerland)*. 1992;184(3):198-201.
59. Hull SM, Cunliffe WJ. Successful treatment of alopecia areata using the contact allergen diphencyprone. *Br J Dermatol*. 1991;124(2):212-213.
60. Chun DK, Eun HC, Lee YS. Treatment of alopecia areata with diphencyprone. *Korean J Dermatol*. 1991;29(3):407-413.
61. Berth-Jones J, Hutchinson PE. Treatment of alopecia totalis with a combination of inosine pranobex and diphencyprone compared to each treatment alone. *Clin Exp Dermatol*. 1991;16(3):172-175.
62. Monk B. Induction of hair growth in alopecia totalis with diphencyprone sensitization. *Clin Exp Dermatol*. 1989;14(2):154-157.
63. Hull SM, Norris JF. Diphenyprone in the treatment of long-standing alopecia areata. *Br J Dermatol*. 1988;119(3):367-374.
64. Caserio RJ. Treatment of alopecia areata with squaric acid dibutylester. *Arch Dermatol*. 1987;123(8):1036-1041.
65. Valsecchi R, Cainelli T, Foiadelli L, Rossi A. Topical immunotherapy of alopecia areata. A follow-up study. *Acta Derm Venereol*. 1986;66(3):269-272.
66. Valsecchi R, Cainelli T, Tornaghi A, et al. Squaric acid dibutylester treatment of alopecia areata. *Clin Exp Dermatol*. 1985;10(3):233-238.
67. Case PC, Mitchell AJ, Swanson NA, Vanderveen EE, Ellis CN, Headington JT. Topical therapy of alopecia areata with squaric acid dibutylester. *J Am Acad Dermatol*. 1984;10(3):447-450.
68. Giannetti A, Orecchia G. Clinical experience on the treatment of alopecia areata with squaric acid dibutyl ester. *Dermatologica*. 1983;167(5):280-282.
69. Flowers FP, Slazinski L, Fenske NA, Pullara TJ. Topical squaric acid dibutylester therapy for alopecia areata. *Cutis*. 1982;30(6):733-736.