

Supplemental Table 1. Epidemiological characteristics among 4,283 white and 615 black women with endometrial carcinoma in the NRG Oncology/GOG 210 Study			
	Race, n (%) <sup>1</sup>		
	White	Black	p <sup>2</sup>
	n=4,283	n=615	
<b>Age at diagnosis</b>			0.66
<54	846 (19.8)	118 (19.2)	
55-59	806 (18.8)	99 (16.1)	
60-64	786 (18.4)	117 (19.0)	
65-69	673 (15.7)	122 (19.8)	
≥70	1172 (27.4)	159 (25.9)	
Median (interquartile range)	63 (56, 70)	63 (57, 70)	
<b>Education</b>			<.0001
Less than high school	415 (9.7)	137 (22.3)	
High school/GED	1403 (32.8)	158 (25.7)	
Some college/technical school	1186 (27.7)	189 (30.7)	
College graduate/beyond	1268 (29.6)	123 (20.0)	
<b>Income</b>			<.0001
< \$20,000	817 (19.1)	258 (42.0)	
\$20,000-\$39,999	940 (22.0)	130 (21.1)	
\$40,000-\$69,999	1019 (23.8)	102 (16.6)	
≥\$70,000	1009 (23.6)	53 (8.6)	
<b>Age at menarche</b>			0.006
≤11	901 (21.0)	138 (22.4)	
12	1256 (29.3)	147 (23.9)	
13	1131 (26.4)	139 (22.6)	
≥14	884 (20.6)	151 (24.6)	
<b>Parity</b>			<.0001
Nulliparous	810 (18.9)	71 (11.5)	
1-2	1797 (42.0)	200 (32.5)	
≥3	1542 (36.0)	310 (50.4)	
<b>BMI 1 year ago, kg/m<sup>2</sup></b>			<.0001
Normal (< 25 kg/m <sup>2</sup> )	944 (22.0)	33 (5.4)	
Overweight (25-30 kg/m <sup>2</sup> )	951 (22.2)	86 (14.0)	
Obese (≥ 30 kg/m <sup>2</sup> )	1956 (45.7)	356 (57.9)	
<b>Smoking status</b>			0.07
Never	2714 (63.4)	413 (67.2)	
Current	282 (6.6)	32 (5.2)	
Former	1191 (27.8)	149 (24.2)	
<b>Diabetes</b>			<.0001
Never	3266 (76.3)	391 (63.6)	
Ever	824 (19.2)	177 (28.8)	

<b>Oral contraceptive use</b>			0.09
Never	1831 (42.8)	232 (37.7)	
Ever	2341 (54.7)	346 (56.3)	
<b>Menopausal status</b>			0.77
Premenopausal	372 (8.7)	50 (8.1)	
Postmenopausal	3818 (89.1)	537 (87.3)	
<b>Menopausal hormone use</b>			<.0001
Never	2882 (67.3)	527 (85.7)	
Estrogen only	329 (7.7)	38 (6.2)	
Progestin only	181 (4.2)	12 (2.0)	
Estrogen + progestin	849 (19.8)	34 (5.5)	
<b>Tamoxifen use</b>			0.17
Never	3933 (91.8)	550 (89.4)	
Ever	199 (4.7)	20 (3.3)	
<b>History of breast cancer</b>			0.22
Never	3857 (90.1)	551 (89.6)	
Ever	331 (7.7)	38 (6.2)	
1 Numbers may not sum to total because of missing data			
2 P value was from two-sided $\chi^2$ test among women with non-missing data for the variable			
3 Among patients for whom the procedure was performed			