

Supplemental Data Table S2. Frequency of *TCR* clonality detected using fragment analysis at initial diagnosis according to three age groups of T-ALL patients

	Total (N=51)	Children (<12 yr; N=16)	Adolescents (12–18 yr; N=9)	Adults (>18 yr; N=26)
None, N (%)	13 (25.5)	1 (7.7)	4 (30.8)	8 (61.5)
<i>TCR</i> β , N (%)	2 (3.9)	1 (50.0)		1 (50.0)
<i>TCR</i> γ , N (%)	11 (21.6)	5 (45.5)		6 (54.5)
<i>TCR</i> δ , N (%)	5 (9.8)	2 (40.0)		3 (60.0)
<i>TCR</i> $\beta+TCR$ γ , N (%)	11 (21.6)	6 (54.5)	3 (27.3)	2 (18.2)
<i>TCR</i> $\beta+TCR$ δ , N (%)	0 (0.0)			
<i>TCR</i> $\gamma+TCR$ δ , N (%)	4 (7.8)			4 (100)
<i>TCR</i> $\beta+TCR$ $\gamma+TCR$ δ , N (%)	5 (9.8)	1 (20.0)	2 (40.0)	2 (40.0)
At least one <i>TCR</i>	38 (74.5)	15 (93.8)	5 (55.6)	18 (69.2)

Abbreviations: *TCR*, T-cell receptor; T-ALL, T-acute lymphoblastic leukemia.