

Three-dimensional structure of the orbicularis retaining ligament: an anatomical study using micro-computed tomography

Jehoon O[#], Hyun-Jin Kwon[#], You-Jin Choi, Tae-Hyeon Cho and Hun-Mu Yang

#co-1st authors. J.O. and H.J.K. contributed equally to the manuscript.

Department of Anatomy, Yonsei University College of Medicine, Seoul, Republic of Korea

Correspondence to:

Hun-Mu Yang, DDS, PhD

Department of Anatomy, Yonsei University College of Medicine

Address: 50-1 Yonsei-ro, Seodaemun-gu, Seoul, 03722, Republic of Korea

Telephone: +82-2-2228-1649

E-mail: yanghm@yuhs.ac

Authors information

1. Jehoon O, MFA

Department of Anatomy, Yonsei University College of Medicine

Address: 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, South Korea

E-mail: jhoon81@yuhs.ac

2. Hyun-Jin Kwon, PhD Student

Department of Anatomy, Yonsei University College of Medicine

Address: 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, South Korea

E-mail: hjkwon128@yuhs.ac

3. You-Jin Choi, PhD

Department of Anatomy, Yonsei University College of Medicine

Address: 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, South Korea

E-mail: CYJ7797@yuhs.ac

4. Tae-Hyeon Cho, BS

Department of Anatomy, Yonsei University College of Medicine

Address: 50-1 Yonsei-ro, Seodaemun-gu, Seoul 03722, South Korea


E-mail: CHOTH@yuhs.ac

Supplementary Video S1

Three-dimensional structure of the orbicularis retaining ligament

Video showing the three-dimensional structure of the orbicularis retaining ligament (ORL) reconstructed from micro-computed tomography images.


Supplementary Figure S1



Modified Verhoeff Van Gieson staining images

Modified Verhoeff Van Gieson staining images of the medial (a), central (b) and lateral (c) portions of the ORL. S, sagittal; A, anterior.

Supplementary Figure S2



Immunofluorescence of the ORL

Immunofluorescence of the ORL for elastin (a, blue), collagen type I (b, green) and actin (c, red). (d) Merged image.