

SUPPLEMENTARY TABLE S1. DEMOGRAPHICS AND BASELINE CHARACTERISTICS

	<i>At risk (A)</i> n = 300	<i>Diagnosed (D)</i> n = 871	<i>Hysterectomy (H)</i> n = 272
Age, mean ± SD, years	33 ± 8.8	41 ± 8.3 ^A	51 ± 6.6 ^{AD}
Age group, %			
18–24 years	18	3 ^A	0 ^{AD}
25–34 years	41	18 ^A	2 ^{AD}
35–44 years	30	43 ^A	12 ^{AD}
45–54 years	10	33 ^A	50 ^{AD}
55–64 years	1	2	36 ^{AD}
Race and ethnicity, %			
White	40	40	54 ^{AD}
Black	19	30 ^A	30 ^A
Hispanic	31	22	10 ^{AD}
Other	4	1 ^A	3
≥2 races, non-Hispanic	6	7	3 ^D
Education, bachelor's degree or higher, %	16	40 ^A	27 ^{AD}
Income, mean ± SD, in thousands (US dollar)	50 ± 41	71 ± 45 ^A	76 ± 45 ^A
Age of first period, mean ± SD, years	12.1 ± 1.8	12.1 ± 1.7	12.3 ± 1.4
Comorbidities ^a in ≥5% of total population, %			
Depression	39	27 ^A	25 ^A
Anxiety	35	26 ^A	18 ^{AD}
High blood pressure	19	21	37 ^{AD}
Ovarian cysts	15	31 ^A	27 ^A
Obesity	21	17	24 ^D
High cholesterol	14	13	33 ^{AD}
Endometriosis	4	18 ^A	20 ^A
Polyps	2	13 ^A	17 ^A
Diabetes/elevated glucose levels	11	7	17 ^D
Infertility	3	7 ^A	4
Polycystic ovary syndrome	4	7 ^A	3 ^D

^aIndicates formal diagnosis by a healthcare provider.

Letters denote statistical difference at 90% confidence between cohorts, where A indicates a significant difference versus at-risk cohort and D indicates a significant difference versus diagnosed cohort.

The *n* values reflect the unweighted base; mean values and percentages are based on the weighted base. Percentages have been rounded to the nearest whole number.