

SUPPLEMENTARY TABLE S2. UTERINE FIBROID SYMPTOM AND HEALTH-RELATED
QUALITY OF LIFE SYMPTOM SEVERITY ITEMS

<i>Question, % of women^{a,b}</i>	<i>At risk n = 300</i>	<i>Diagnosed n = 871</i>	<i>General population^c n = 5961</i>
Feeling fatigued	83%	44%	25%
Heavy bleeding during menstrual period	78%	43%	17%
Feeling tightness or pressure in pelvic area	64%	34%	15%
Passing blood clots during menstrual period	62%	38%	17%
Frequent urination during daytime hours	55%	28%	11%
Fluctuation in the duration of menstrual period compared to previous cycles	49%	27%	11%
Fluctuation in length of monthly cycle compared to previous cycles	45%	26%	11%
Frequent nighttime urination	47%	24%	8%

^aEach question asks how much distress a participant has experienced from each symptom during the previous month; response options ranged from 1 (“Not At All”) to 5 (“A Very Great Deal”).

^bValues reported indicate percentage of women reporting a 4 (“A Great Deal”) or 5 (“A Very Great Deal”) of distress during past month.

^cGeneral population represents women who completed the Screener, had not undergone a hysterectomy, and were premenopausal.

No statistical analyses were performed between the cohorts for item-level data. The *n* values reflect the unweighted base; percentages are based on the weighted base.