Appendix 1. Hispanic American participation in prospective cohort studies in the U.S., by size (largest to smallest)

	Authors,			D 1.4		Biological
Cohort	Year	Focus	Location	Population	# subjects	samples (n)
Kaiser Permanente						
Southern California					920 034	
(KPSC) Children's	Koebnick C,		Southern	Multi-ethnic, children	(approx. 50%	
health study	et al., 2007	Childhood obesity	California	aged 2-19	Hispanic)	None
				Latino, African-		
	Henderson			American, Japanese-		Blood, Urine
The Multiethnic	BE, et		Hawaii,	American, Native	215 000 (47	(~15 427 Latino
Cohort	al.,1993	Diet and cancer	California	Hawaiian, NHW	438 Latino)	samples)*
				Non-Hispanic whites,		
		Current and former		African-American,		
California	Bernstein L,	public school teachers		Hispanic, Asian and	133 479 (~5%	Blood, Saliva
Teacher's Study	et al., 2002-	and administrators	California	Pacific Island women	Hispanic)	(~2000 samples)*
California Men's	Caan, et al.,			Hispanics, Blacks,	84 170 (11	• /
Health Study	2000	Prostate cancer	California	whites, aged 45 to 69	428 Latinos)	None
				Multi-ethnic, over age	>37 000	
Health and	Weir DR, et	Health, retirement,		50 (oversampling of	(approx. 9%	Blood, Saliva
Retirement Study	al., 1992	aging	National	Hispanics)	Hispanic)	(~1405 samples)*
,	,	Role of genetic,			1 /	Blood, saliva,
		acculturation and				and urine (56%,
	Chow W, Wu	sociodemogrpahic				63% and 43% of
Mano A Mano	X, et al,	factors in cancer and				all participants,
cohort study	2001-	chronic disease	Houston, Texas	Mexican-Americans	25 483 MAs	respectively)
Diabetes Study of						
Northern California	Moffet HH,		Northern	Multi-ethnic, aged 30-	20 188 (3717	
(DISTANCE)	et al., 2005	Diabetes	California	75	Latinos)	None
(BISTITIVEE)	Ct u1., 2003	Prevalence and	Cumomu	7.5	Euthiosy	110110
Hispanic		development of disease				
Community Health	Aviles-Santa	in Hispanics/Latinos,	Bronx, Chicago,			
Study/ Study of	L, et al.,	role of acculturation,	Miami, San	Hispanic/Latinos,		Blood, urine
Latinos	2008-	risk factors	Diego	aged 18 to 74 years	16 415	(15 643 samples)
Latifics	2000-	TISK 10001S	Diego	4504 10 to 14 years	10 713	Blood, urine
	Victor RG,			Blacks, non-Hispanic		(~475
	Hobbs HH,		Dallas County,	whites, Hispanics,	6101 (approx.	participants aged
Dallas Heart Study	2000	Cardiovascular health	Texas	aged 18 to 65	14% Hispanic)	30-65)*
Danas Heart Study	4000	Carulovasculai licalili	1 CAAS	aged 10 to 03	14/0 Hispailie)	30-03)

Northern	Elkind M, et		Northern			
Manhattan Study	al.,1993	Stroke	Manhattan	Hispanic, black, white	4400	Blood
	Stern M,			Mexican Americans		
	Lorenzo C, et	D' 1 1		and non-Hispanic		
San Antonio Heart	al., 1979,	Diabetes and	G	whites, aged 25-64	2201 3.64	D1 1
Study	1988	Cardiovascular Disease	San Antonio	years	3301 MAs	Blood
Hispanic						
Established			A			
Populations for the Epidemiology		Physical and health	Arizona, California,			
Study of the Elderly	Markides KS,	conditions in older	Colorado, New	Mexican-Americans,		
(HEPESE)	et al.,1993	Mexican-Americans	Mexico, Texas	aged 65 and older	3050	None
Life After Cancer	Caan B,	MCAICAII-AIIICHCAIIS	Wickico, Texas	Multi-ethnic,	3030	None
Epidemiology	Sternfeld B,		Northern	diagnosed with breast	2321 (6%	
Study (LACE)	et al.,2000	Breast cancer survivors	California, Utah	cancer, aged 18-79	Hispanic)	None
Lovelace Smokers	Tesfaigzi Y,	Breast carreer survivors	Cumoma, Ctan	Hispanic, NHW,	2300 (369	DNA (307
Cohort	2000	COPD, lung cancer	New Mexico	smokers aged 40-75	Hispanic)	samples)
	Fisher-Hoch	, <u>C</u>		<u> </u>	1 /	1 /
Cameron County	SP, et al.,			Mexican Americans,		Blood (1847
Hispanic Cohort	2003	Obesity, diabetes	Texas	aged 18 and older	2000	participants)
				Patients with AD and		_
				mild cognitive		
Texas Harris				impairment; large %		
Alzheimer's	O'Bryant SE,			Mexican-American		
Research Study	et al., 2009	Alzheimer's disease	Texas	participants	2000	Blood
_		Minority health,				
Boston Area		urologic				D1 1/500/ 0.11
Community Health	Piccolo RS,	signs/symptoms, type 2	.	a called a decident	107611	Blood (68% of all
survey	et al., 2002	diabetes	Boston	Multi-ethnic	1876 Hispanic	subjects), DNA
Sacramento Area	Gonzalez			M . A .		
Latino Study on	HM, et al., 1998	Mortality, dementia	California	Mexican Americans, aged 60 or older	1789	None
Aging	1998	Mortanty, dementia	NY, Baltimore,	aged 60 of older	1/89	None
			Chicago, LA,			
Multi-ethnic Study	Olson J, et		Minnesota,	Multi-ethnic (22%		Blood, urine
of Atherosclerosis	al., 2000	Cardiovascular disease	North Carolina	Hispanic), aged 45-84	1496	(1496 samples)
Latina Gestational	Chasan-	The state of the s	- A Committee of the Co	Puerto Rican,	2.12.0	(1.50 Sampies)
Diabetes Study/	Taber L, et	Gestational diabetes	Massachusetts	Dominican, pregnant	1231	Blood
	, -•			, F		

Proyecto Buena	al.,2000,			women, aged 16-40		
Salud	2006			_		
Viva la Familia	Butte NF, et					_
Study	al., 2000	Childhood obesity	Texas	Hispanic children	1030	Blood (1030)
	Bergman RN,			African American,		_
	Wagenknecht	Insulin Resistance,		Hispanic, non-		
IRAS/ IRAS	LE, et al.,	Atherosclerosis,		Hispanic white, aged		
Family Study	1992, 1999	Visceral Adiposity	multi-center	18-81	1000	Blood, urine
SUNSHINE cohort						
(follow up of						
SHINE 4-Corners						
Breast Cancer	Sedjo RL, et	Breast cancer, women's	Arizona,	Non-Hispanic whites,	650 (204	
Study)	al., 2007	health	Colorado	Hispanics	Hispanics)	None
Project FRONTIER	O'Bryant SE,			Multiethnic rural		_
(Cochran County	Lambert M,	Aging, cognitive		adults aged 40 and	519 (235	
Aging Study)	et al., 2008	health, rural health	Texas	above	Hispanics)	Blood
San Antonio				Mexican Americans,		
Longitudinal Study	Gonzalez H,			European Americans,		
of Aging	et al., 1992-	Mortality, Aging	San Antonio	aged 65 to 80	394 MAs	None
				Mexican American,		_
Hispanic Chronic	Lash JP, et			Puerto Rican, other		
Renal Insufficiency	al., 2005-	Chronic Kidney		Latin American, aged		Blood, urine (327
Cohort Study	2008	Disease	Chicago	21-74 with CKD	327 MAs	samples each)

Note. * indicates approximate amounts based on reported percentages of Hispanic or Latino in the cohort