

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Identifying causes for socioeconomic differences in the treatment of coronary heart disease: a qualitative study from the perspective of elderly patients

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-024151
Article Type:	Research
Date Submitted by the Author:	11-May-2018
Complete List of Authors:	Schröder, Sara Lena; Martin Luther University Halle-Wittenberg, Germany, Institute of Medical Sociology Fink, Astrid; Martin Luther University Halle-Wittenberg, Institute of Medical Sociology Richter, Matthias; Martin Luther University Halle-Wittenberg, Institute of Medical Sociology
Keywords:	health services research, socioeconomic status, Coronary heart disease < CARDIOLOGY, QUALITATIVE RESEARCH, healthcare inequalities, health services accessibility

SCHOLARONE™
Manuscripts

Peer Review Only

1
2
3 **Identifying causes for socioeconomic differences in the treatment of coronary**
4 **heart disease: a qualitative study from the perspective of elderly patients**
5
6
7

8
9 Sara L. Schröder, Astrid Fink (PhD), Matthias Richter (PhD)
10

11
12 Institute of Medical Sociology, Martin Luther University Halle-Wittenberg, Halle (Saale),
13 Germany
14
15

16
17
18
19 **Corresponding author:**
20

21 Sara Lena Schröder

22 Institute of Medical Sociology (IMS)

23 Martin Luther University Halle-Wittenberg

24 Magdeburger Str. 8

25 06112 Halle (Saale)

26 Germany
27

28 Email: sara.schroeder@medizin.uni-halle.de
29
30

31
32
33
34
35
36 Word count: 4087
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives - Socioeconomic status (SES) is associated with treatment of coronary heart disease (CHD) to the disadvantage of deprived patients, but little is known regarding the reasons for socioeconomic inequalities in healthcare. This qualitative study provides an in-depth insight into socioeconomic differences in patient's experiences with treatment of CHD to understand the underlying causes for socioeconomic differences.

Design - Longitudinal qualitative study using in-depth semi-structured interviews to explore patient's experiences with treatment. We analysed the transcripts of the records according to qualitative content analysis, and identified differences between high and low SES by comparing and contrasting the narratives.

Setting - University hospital in Halle (Saale), Germany.

Participants - 41 elderly patients (aged 59-80) who suffered CHD.

Results - Three major themes characterize the socioeconomic differences in the patient's experiences with treatment: (1) information: patients with high SES had greater knowledge about treatment and could use medical records as sources of information; (2) illness perception: patients with low SES focused on improving symptoms and survival, while patients with high SES focused on physical performance and disease management; and (3) perceived role in healthcare: patients with low SES tended to delegate responsibility to healthcare professionals.

Conclusions - Information, the patient's perceived role in health care, and illness perception may mediate the association between SES and treatment of CHD. These factors should be considered in quantitative studies to better understand the disparities in treatment and mortality. We suggest that improving patient-physician-communication and patient knowledge could change their understanding of CHD and their perceived role in healthcare and reduce inequalities in CHD treatment.

Trial registration number - DRKS00007839.

1
2
3
4
5 **Keywords:** coronary heart disease, socioeconomic status, healthcare inequalities, qualitative
6 research, health services accessibility
7
8
9

10 **Strengths and Limitations**

- 11 - Less is known about the underlying causes of socioeconomic inequalities in CHD
12 treatment.
13
- 14 - The longitudinal design enables us to gather information of the patients experiences along
15 the entire pathway of treatment of CHD.
16
- 17 - We undertook a comparative analysis to explore differences in patient's perspectives and
18 experiences with treatment.
19
- 20 - As most patients were treated according to guidelines, we cannot conclude if there are
21 socioeconomic differences in treatment of CHD in Germany.
22
- 23 - Only minor issues in each phase of treatment were found to differ according to SES, but
24 altogether these show a consistent image of differences in the patient's perspectives with
25 CHD treatment.
26
27
28
29
30
31
32

33 **Funding**

34 This work was supported by the Wilhelm Roux Programme of the Medical Faculty of Martin
35 Luther University, Halle-Wittenberg with grant number FKZ: 28/40.
36
37
38
39

40 **Competing Interests**

41 The authors declare that they have no competing interests.
42
43
44
45

46 **Authors contribution:**

47 SLS wrote the initial draft of this manuscript, collected data and analysed the transcripts. MR
48 and AF conceptualized and designed the project, helped interpreting the data, and critically
49 reviewed the manuscript. All authors have read and approved the final version of the manuscript.
50
51
52
53
54
55
56
57
58
59
60

Data Sharing Statement

Participants of this study were guaranteed that only the study research team would have access to the interviews and transcripts thus data are not available for sharing.

Ethics approval

Ethical Review Committee of the Medical Faculty of Martin Luther University, Halle-Wittenberg

For peer review only

Introduction

Cardiovascular diseases remain the leading cause of death worldwide.[1–3] Treatment of coronary heart disease (CHD), including revascularization, rehabilitation and long-term medication, aims to improve disease-related quality of life, including exercise capacity, to prevent further cardiac events and reduce mortality. Socioeconomic inequalities in the incidence and mortality of CHD, to the disadvantage of patients with low socioeconomic status (SES), have been well explored.[4–6] Therefore, patients of low SES have a greater need for treatment. Access to treatment is aimed to be on the basis of need rather than of gender, residence or SES, and guidelines have been established to reduce inequalities in healthcare.[7] But several studies showed strong socioeconomic differences in CHD treatment, to the disadvantage of patients with low SES.[8]

How inequalities in healthcare arise is complex and largely unknown. As socioeconomic differences in treatment are provable independent of the underlying healthcare system, they may be attributable to reasons other than financial affordability of healthcare costs.[8] This is particularly the case in countries such as Germany, where individuals have statutory health insurance and very low out-of-pocket payments. In cardiac rehabilitation, a recent study found that comorbidities, self-efficacy, anxiety and depression, cohabitation, commute, disease severity or type of treatment do not significantly mediate the association of SES with attendance and participation.[9] But generally health literacy and communication between physician and patient are discussed to explain healthcare inequalities, as they are strongly associated with SES.[10–12] Although there is no clear evidence regarding whether an increase in treatment rates reduced socioeconomic disparities in treatment access,[13, 14] studies found that socioeconomic inequalities in CHD treatment can be partly explained by different distances to hospitals with on-site cardiac facilities.[15, 16]

Using quantitative data, these studies remain at a descriptive level, and cannot explain the underlying causes and determinants of socioeconomic inequalities in CHD treatment and how these factors lead to inequalities in healthcare. Taking the patients' individual needs and perceptions into account is helpful to reveal new explanatory approaches from the patient's perspective and explore how socioeconomic differences arise. Qualitative research aims to

1
2
3 understand healthcare interactions from the patient's perspective and can help identify the
4 underlying causes. To our knowledge, there are only two qualitative study exploring
5 socioeconomic differences in the treatment of CHD. Manderbacka analysed differences in
6 healthcare encounters and found that doctor-centred decision making was more common in low-
7 SES patients.[17] Pedersen et al. found that concerning barriers to cardiac rehabilitation, only
8 low SES patients felt excluded due to their divergent health beliefs.[18] To gather scientific
9 evidence on the underlying causes for socioeconomic differences, we aimed to identify
10 socioeconomic differences in the patient's perspective and their experiences with the entire
11 pathway of CHD treatment as these might be possible factors and mechanisms that may lead to
12 inequalities in CHD treatment.
13
14
15
16
17
18
19
20
21

22 **Material and methods**

23 **Study design**

24
25 We conducted an exploratory qualitative longitudinal study of elderly CHD patients to examine
26 socioeconomic differences in access, utilization and quality of treatment.[19] A purposive
27 sampling strategy was used to select patients with a confirmed diagnosis of CHD at a university
28 hospital in Germany. Between November 2014 and April 2015, a study nurse contacted patients
29 aged 59-80 who were hospitalized with CHD, informed them through a flyer and explained the
30 purpose of the study. We approached 96 eligible patients during hospitalization and asked them
31 to attend an interview to share their experiences with CHD treatment. Participation was
32 voluntary, the patients received no incentives, and 35 patients refused to participate. Maximum
33 variation was used to assure that men and women with different SES and different CHD
34 severities were represented to cover a broad spectrum of treatment experience. Patients were
35 enrolled in the study after providing written informed consent until theoretical saturation was
36 reached. In total, 48 interviews were conducted in the hospital. After 6 months, we contacted the
37 participants again and asked them to participate in a follow-up interview. Nine patients refused
38 to participate due to their disease burden, death, lack of interest or failure to respond to the
39 invitation letter. Thirty-nine follow-up interviews were conducted between June and October
40 2015. The study was approved by the Ethics Review Committee of the Medical Faculty at the
41 Martin Luther University, Halle-Wittenberg.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Data collection

We conducted in-depth interviews to explore patients' perspectives on their heart disease and their personal experiences with the treatment they received. SLS conducted two preliminary interviews that were used to further develop the semi-structured interview guides (S1 pdf). We began the baseline interviews with a broad general question about the medical history. More detailed questions were asked individually based on each patient's narrative. The patient's basic sociodemographic data were obtained with a short standardized questionnaire following the baseline interview, and the interviewer subsequently wrote a field note for each interview.

Baseline interviews took place in a private room of the hospital, and follow-up interviews were held either at the patient's home or at the medical facility, depending on the patient's choice. Two employees of the Institute of Medical Sociology (IMS) conducted the interviews face-to-face: SLS (female) is a research associate and an economist who has conducted qualitative health research since 2009 and conducted most of the interviews; Nils Bormann (male) is a medical student assistant and an economist who was trained prior to conducting the interviews. The interviewers introduced themselves as members of the IMS not working in the department of cardiology. Each baseline interview included only the patient and one or both interviewers. In some follow-up interviews, family members were present at the patient's request. The interviews lasted an average of 35 minutes at baseline and 42 minutes at follow-up and were audiotaped with the interviewee's permission.

SES was defined by educational level and occupation based on the German epidemiological standards.[20] Education was measured by level of schooling and academic qualifications and was classified on an 8-point scale. Occupation was measured by the last occupational group the patient belonged to and classified according to job autonomy on a 5-point scale.[21] A composite index of SES was derived using a sum score of both items, which can range between 2 and 13. Index values between 2 and 7 were rated "low SES", and values between 8 and 13 were rated "high SES".

Data analysis

1
2
3 SLS analysed the pseudonymized interview transcripts inductively with codes and themes
4 derived from the data. Five transcripts were double-coded by two additional members of the
5 qualitative research working group at the IMS, who also participated in the discussion and
6 evaluation of the data. The working group was composed of researchers with different
7 backgrounds, namely, sociology, health care research, economics, nursing and educational
8 science. In the present analysis, we excluded 7 patients whose treatment experiences had been
9 overshadowed by other heart diseases with little relation to CHD throughout the narratives. Of
10 the remaining 41 baseline interviews included in this analysis, 17 patients had been diagnosed
11 with CHD up to one year prior to the baseline interview. Therefore, we included their follow-up
12 interviews, as they had not gone through the entire treatment pathway at baseline. The narratives
13 were coded inductively and analysed in accordance with qualitative content analysis.[22] We
14 identified socioeconomic differences by comparing and contrasting the patients' narratives and
15 identified SES-specific patient perspectives and treatment experiences. We used MAXQDA 11
16 software to assist with the data management and analyses.
17
18
19
20
21
22
23
24
25
26
27
28

29 **Patient Involvement**

30 By exploring patient's experiences with treatment in this study with an exploratory design, the
31 patient's preferences and priorities led the data collection during the interviews and as well due
32 to an inductive approach throughout data analysis. The participants did not provide feedback on
33 their transcripts or the findings, but interested patients received a summary of the main results.
34
35
36
37
38

39 **Results**

40 This analysis was based on the transcripts of 41 baseline and 17 follow-up interviews. The
41 baseline characteristics of the 41 patients are shown in Table 1. The mean patient age was 69.7
42 years, and the study population at baseline comprised 15 (37%) women and 26 (63%) men. The
43 low-SES group comprised 23 (56%) patients, and the high-SES group comprised 18 (44%)
44 patients.
45
46
47
48
49
50

51 Treatment of CHD was reported as a complex process with the involvement of many providers.
52 The majority of patients reported treatment according to guidelines. After being diagnosed with
53 CHD through coronary angiography (CAG), coronary arteries were widened with PCI, CABG or
54
55
56
57
58
59

1
2
3 ballooning, and the hospital organized the referral to inpatient CR. Patients generally reported
4 that they regularly visit their general practitioner (GP), who prescribes the medication as
5 recommended from the hospital. Furthermore, patients had regular appointments with a resident
6 cardiologist. Only a few patients (from both SES groups) reported having problems with
7 treatment at the hospital, secondary preventive drugs, attending regular follow-ups or
8 participating in CR. However, we found differences according to SES in the patients' individual
9 perceptions and perspectives on access, utilization and quality of treatment. Pseudonymized
10 quotes reflective of the themes are presented for patients of high and low SES at various
11 treatment stages.
12
13
14
15
16
17
18
19

20 **Treatment at the hospital**

21 Most patients reported a history of revascularization with balloon dilatation, stents, or coronary
22 artery bypass graft surgery, and some patients reported that a second CAG was scheduled and
23 conducted in a timely manner, e.g., to protect the patient's circulatory system. However, only
24 low-SES patients reported that this second CAG was performed to implant stents following a
25 diagnostic-only CAG without mentioning any reason. Whereas many patients of both low and
26 high SES rated it important for hospital staff to be friendly, attentive and caring, predominantly
27 low-SES patients mentioned aspects of being taken seriously either positively (e.g., taking
28 preferences seriously, being asked how one feels) or negatively (e.g., discussions about but not
29 with oneself, no conversations at all). While assessing the quality of treatment, some low-SES
30 patients complained about being discharged with open wounds after CABG. Generally, while
31 assessing the quality of treatment, low-SES patients tended to focus on symptom improvement,
32 eliminating the need for visits to the doctor or improving survival, whereas high-SES patients
33 predominantly aimed to improve physical performance.
34
35
36
37
38
39
40
41
42
43
44
45

46 “Well, but then it came up with the heart and I myself had the feeling that it is probably
47 the only thing that really gets you back on your feet. I was really feeling sluggish. So. I
48 did indeed make an effort to try to walk again quickly and to/ I really did, but somehow
49 in the end the energy was lacking.” (Mr. Lehmann, 63 years, high-SES)
50
51
52
53
54
55
56
57
58
59
60

1
2
3 “Well, in the hope that everything stays the same. Let’s say that you are well. Well,
4 getting around with it quite well, with no complications arising. As I said, that one may
5 reach a slightly older age, not just until retirement, but perhaps even a bit longer.” (Mr.
6 Lange, 60 years, low-SES)
7
8
9

10
11 Knowledge of different treatment options became apparent mainly in patients with high SES.
12 However, an information deficit was found in some narratives, e.g., patients reporting that they
13 did not know the outcome of medical investigations, were unaware of the treatment that had been
14 performed or rated the medical reports incomprehensible. An information deficit became
15 apparent mainly in low-SES patients, although this was not always experienced with a perceived
16 greater need for information.
17
18
19
20
21
22
23

24 “For this there are, that’s at least what has been explained to me, the clogged coronary
25 arteries can be expanded by implanting stents, if you know what these are, incorporating
26 these small tubes that eliminate the narrowed areas. Or if that is not feasible, which can
27 happen as well that a vessel ruptures or something like that, then (...) it is done surgically.
28 Then bypasses are produced from other parts of the body and incorporated.” (Mr.
29 Wagner, 75 years, high-SES)
30
31
32
33
34
35

36 “Yes, it has been explained to me, but I didn’t get it. First of all, I sometimes didn’t
37 understand it, because he expressed himself in such a professional manner. The nurses
38 already started laughing, because I/ and then I thought: ‘No, you better give up. You
39 better read through it next time’” (Mrs. Schubert, 80 years, low-SES)
40
41
42
43
44

45 We found paternalistic and shared decision making in both SES groups, but paternalistic
46 decisions were found to be more unquestioned in low-SES patients, whereas high-SES patients
47 understood the reasons why decisions were made. When decisions have been made jointly, we
48 found that high-SES patients were involved more actively in decision-making, whereas low-SES
49 patients saw themselves as only responsible for responding (agree or disagree) to a
50 recommendation given.
51
52
53
54
55
56
57
58
59
60

1
2
3 “With this I entirely relied on the physicians. In fact, I didn’t really thought about it.
4 When they said: ‘This’, then we do this. They do ask indeed: ‘Mr. Zimmermann, do you
5 agree?’ I mean, when I go to the hospital, I let myself be treated, but this is not meant to
6 be negative in any way. But I rely on the physicians. [...] They are the professionals. I
7 would never pretend to be the wise guy, as I know people who do so, not physicians, but
8 private persons, who dictate people what to do.” (Mr. Zimmermann, 76 years, low-SES)
9
10
11
12
13
14

15 “Today quite a few medical exams have been conducted with me. Now you have to wait,
16 but I guess it won’t change anything about this final decision. The requirement to me to
17 decide this.” (Mr. Hartmann, 69 years, high-SES)
18
19
20
21

22 **Cardiac rehabilitation**

23 Some patients assessed the time between hospital discharge and beginning of inpatient CR. In
24 doing so, only high-SES patients focused on individual demands for disease management,
25 whereas only low-SES patients reported they could not participate in therapeutic CR treatments
26 due to their recent coronary artery bypass surgery. Additionally, we found that while assessing
27 the quality of CR, high-SES patients tended to place an emphasis on physical performance
28 improvement, as well as increased knowledge about necessary behaviour changes and their heart
29 disease in general, whereas in the narratives of low-SES patients, the conduction of regular
30 examinations, such as daily blood pressure measurement, was more important.
31
32
33
34
35
36
37
38

39 “Over there I primarily learned through lectures and, as I said before, through meetings
40 about what the issue of a heart attack means, how it emerges and so on. This improved
41 much of my understanding. Previously, before you have something like that you don’t
42 care about it. So, concerning this matter one was educated quite well.” (Mr. Jung, 67
43 years, high-SES)
44
45
46
47
48

49 “Indeed, in rehabilitation it was great. Well, there was/ every morning you had to go to
50 the presentation, weighing, measuring blood pressure and stuff like that.” (Mrs. Koch, 62
51 years, low-SES)
52
53
54
55
56
57
58
59
60

Treatment with drugs

Although specific drugs, agents or trade names were rarely mentioned throughout the narratives, access to and utilization of any secondary prevention drugs were fairly high in both SES groups. However, we found that with respect to medication preferences, some low-SES patients highlighted the importance that involved physicians appropriately coordinate the selection of prescribed drugs, while some stated they generally did not like to take drugs. However, one high-SES patient reported that he prefers to take combination medication.

“Or, for example, if there are changes in medication, it is very important. With prothrombin time and everything; you need to be always in good hands. Not that one says this way and the other one says that way. That’s bad.” (Mr. Köhler, 66 years, low-SES)

“I have to take many drugs; they upset the stomach and everything. I had to take 17 or 18 different pills. Well, and then I recognized, I read about it, there is a patch and so on and using that you can come off of 5 drugs at once. The patches are indeed very expensive and the doctor immediately said ‘no and no and no’ and so on. Now I have this patch and I am feeling fine.” (Mr. Richter, 66 years, high-SES)

Ambulatory aftercare

Some patients in the low-SES group mentioned undergoing regular check-ups every month with their GP to assess prothrombin time (Quick-test). While rating the quality of GPs, mainly low-SES patients mentioned the importance of thorough medical examinations and doctor availability. Some mainly low-SES patients complained about the GP examining or referring patients only on demand or the GP not helping them make an appointment with a cardiologist. Some predominantly low-SES patients reported that they see themselves as responsible for retention of their medical record. However, some patients, mainly with high SES, mentioned obtaining or inquiring for a recommendation from the GP regarding a particular specialist or hospital.

“But, I wasn’t correctly/ At first I didn’t know something like that existed. Then I asked my doctor. I say: ‘Don’t one has to go to any follow-up?’ - ‘Well, I could forward you

1
2
3 there, but you better try to make an appointment yourself’.” (Mrs. Körtig, 59 years, low-
4 SES)
5
6
7

8 “In one consultation it was like, ‘To whom should I go?’ And then he said, ‘I have a
9 number of colleagues here that I can recommend to you’ And then I say, ‘Alright, then
10 make an appointment for me’ And that’s how I came to Mrs. Dr. Alpha.” (Mr. Winkler,
11 72 years, high-SES)
12
13
14
15

16
17 In addition, the patient’s experiences with access to and utilization of regular check-ups at a
18 resident cardiologist varied according to SES. Some low-SES patients perceived appointments
19 with specialists as generally unnecessary or expected instructions from physicians about the
20 necessity of appointments due to their own uncertainty. Some mainly high-SES patients used
21 medical reports as sources of information for follow-up care.
22
23
24
25

26
27 “As I said, I had to wait a very long time for this 24-hour ECG. Then it was done. She
28 [cardiologist] wrote her report, my GP got it and nothing else came of it. [...] I mean, if
29 there would have been something acute, surely something would have come up, but it
30 still was, that it is not” (Mr. Köhler, 66 years, low-SES)
31
32
33
34
35

36 “I guess that I will have to go to my GP for follow up and he does further. [...] That is my
37 current state. But I am not sure yet, because I have not received the medical report yet. I
38 am still waiting for it.” (Mr. Schäfer, 71 years, high-SES)
39
40
41
42

43 **Underlying causes of socioeconomic differences in CHD treatment**

44 While comparing the patients’ experiences across the healthcare sectors, 3 major themes (Fig 1)
45 emerged to characterize differences and may be the underlying reasons for socioeconomic
46 inequalities in CHD treatment, namely, information, a patient’s individual illness perception and
47 a their perceived role in health care.
48
49
50
51

52
53 First, the narratives suggest that low-SES patients are less informed about their treatment than
54 high-SES patients, which became apparent, e.g., through their lack of knowledge regarding the
55
56
57
58
59

1
2
3 treatment that was performed, as well as their lack of knowledge about the reasons for or
4 outcomes of medical investigations. However, high-SES patients knew about different treatment
5 options. Whereas high-SES patients used medical reports as sources of information, low-SES
6 patients missed conversations at the hospital or rated the reports as incomprehensible. While
7 most patients rated the healthcare system (especially hospitals) as overloaded, some made the
8 connection that patients are given far too little information and are required to actively request
9 information because of excessive work demands on health workers due to the healthcare system
10 economization.
11
12
13
14
15
16
17

18
19 “And as well, someone who is cognitively not that fit anymore, one does not understand
20 at all what you are told. And that is sometimes not so nice. Well, but probably that’s just
21 our system, that is/ which is not working, I think”. (Mrs. Koch, 62 years, low-SES)
22
23
24

25
26 Second, we found differences in patients’ illness perceptions. Whereas low-SES patients seemed
27 to understand CHD through putting emphasis on symptoms and medical parameters measured
28 through medical examinations, high-SES patients focused on physical performance and disease
29 management, which requires knowledge about CHD.
30
31
32
33

34
35 Finally, we found differences in the patient's perceived role in their health care. Low-SES
36 patients tended to delegate responsibility for treatment, which became apparent, e.g., by the
37 patients not questioning the reasons for what and why something (diagnostic-only CAG,
38 discharge with open wounds) is done, assuming healthcare professionals responsible that patients
39 are being taken seriously, not perceiving a need to obtain information although lacking
40 information, not being as involved in decision making, relying on physicians to coordinate care
41 (referrals, medication), seeing themselves as responsible only for retention of medical records
42 and having a tendency to have general views rather than opinions. However, high-SES patients
43 seem to feel more responsible for treatment, e.g., focusing more strongly on disease
44 management, having increased healthcare knowledge, making informed choices or actively
45 involving their GP.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

In this study, we found 3 major themes where the patients' experiences with CHD treatment differed between patients of high and low SES: information, the patient's individual perception of CHD and their perceived role in health care. Regarding information patients with high SES had greater knowledge about treatment and could use medical records as sources of information. Regarding illness perception patients with low SES focused on improving symptoms and survival, while patients with high SES focused on physical performance and disease management. Regarding the patient's perceived role in healthcare patients with low SES tended to delegate responsibility to healthcare professionals.

First, we found that low-SES patients seem to lack knowledge about treatment compared with high-SES patients, get less information and have problems understanding the information provided to them. This is in accordance with previous studies, which found that patients of low SES are given less information in healthcare,[10, 23] and are therefore less informed, e.g., about glaucoma treatment[24]. Studies found that lack of information and knowledge results in low utilization, especially in low-SES patients.[25–27] Studies showed that patients are given little information by health professionals and therefore wide information gaps exist for all CHD patients.[28, 29] Second, illness perception differs as we found that low-SES patients seem to focus more strongly on symptoms, survival and good results of physical check-ups in their illness perception, while high-SES patients tend to focus on physical performance and disease management. The latter reveals more knowledge about CHD, and studies found that lack of knowledge regarding CHD is associated with both low SES and adherence.[30–34] Thereby we confirm the findings of another study that low SES, but not high-SES, patients felt excluded due to their divergent health beliefs.[18] This becomes particular relevant as focusing on and improving physical performance are important for reducing cardiovascular mortality and inequalities in AMI mortality, independently of inequalities in access to healthcare.[35] Finally, the perceived role in health care differed according to SES: high-SES patients tended to assume responsibility for treatment, whereas low-SES patients showed a tendency to hand over responsibility. This latter finding was characterized by “putting up with everything needed” and expecting to be told that “everything is fine”, with no further need to see a physician. Different responsibilities are becoming apparent through decision making between physician and patient,

1
2
3 as a study found that high-SES patients perceive their involvement as participating in the
4 decision making after exploring other options, whereas low-SES patients see themselves only
5 being responsible for agreeing or disagreeing with the recommendation made.[36] Therefore we
6 confirm another study, which found that inequalities might be caused by the fact that shared
7 decision making is more common in high-SES patients.[17] Previous studies showed that
8 engaged patients who actively participate in treatment have good compliance.[37]
9

10
11
12
13
14
15 Although we provide novel insights into the underlying causes of socioeconomic differences in
16 treatment, there are several limitations. We recruited elderly patients in a university hospital
17 setting in a large city in the eastern part of Germany. Therefore, these results may not reflect the
18 experiences of younger patients or patients living in other regions. Importantly, data were
19 generated in a country with statutory health insurance. Therefore, financial concerns may better
20 explain socioeconomic differences in treatment in other countries without universal healthcare
21 systems. Additionally, only a few patients reported problems with access to and utilization of
22 treatment, and throughout the narratives, only minor perspectives were found to differ according
23 to SES. However, consideration of these aspects together led us to an understanding of the
24 underlying causes of socioeconomic inequalities, based on the words of the patients studied.
25 Finally, especially when assessing physician-patient communication, it has to be considered that
26 we only analysed the patients' perspectives and viewpoints.
27
28
29
30
31
32
33
34
35
36

37
38 Given that CHD is the leading cause of death worldwide, the generation of hypotheses regarding
39 the potential factors and mechanisms underlying the socioeconomic differences in CHD
40 treatment is essential for explaining and reducing these inequalities. The level of information
41 and knowledge, perceived role in health care and assumption of responsibility, as well as
42 individual illness perception should be considered in further quantitative studies as mediating
43 factors between SES and treatment. We conclude that improvement in patient-provider
44 communication might be the key to reducing healthcare inequalities, as it may increase
45 knowledge about treatment, improve the understanding of CHD and empower patients to assume
46 responsibility in treatment, which can improve engagement in treatment and utilization.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgments

We acknowledge the patients who participated in this study and shared their healthcare experiences with us. We thank the staff at the Department of Cardiology at the University Hospital in Halle for their assistance with screening patients for this stud. Finally we thank Niels Bormann (NB) for assisting with data collection, and the members of the qualitative research working group at the IMS, who helped with coding, discussing and evaluating the data.

For peer review only

References

- 1 Nichols M, Townsend N, Scarborough P, et al. Cardiovascular disease in Europe: epidemiological update. *Eur Heart J* 2013;34(39):3028–34.
- 2 GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385(9963):117–71.
- 3 Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart Disease and Stroke Statistics-2017 Update: A Report From the American Heart Association. *CIRCULATION* 2017;135(10):e146-e603.
- 4 Mackenbach JP, Cavelaars AE, Kunst AE, et al. Socioeconomic inequalities in cardiovascular disease mortality; an international study. *Eur Heart J* 2000;21(14):1141–51.
- 5 Havranek EP, Mujahid MS, Barr DA, et al. Social Determinants of Risk and Outcomes for Cardiovascular Disease: A Scientific Statement From the American Heart Association. *CIRCULATION* 2015;132(9):873–98.
- 6 Baldi I, Costa G, Foltran F, et al. Effect of educational attainment on incidence and mortality for ischemic heart and cerebrovascular diseases: a systematic review and trend estimation. *Int J Cardiol* 2013;168(5):4959–63.
- 7 Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease. *J Am Coll Cardiol* 2012;60(24):e44-e164.
- 8 Schröder SL, Richter M, Schröder J, et al. Socioeconomic inequalities in access to treatment for coronary heart disease: A systematic review. *Int J Cardiol* 2016;219:70–78.
- 9 Pedersen M, Egerod I, Overgaard D, et al. Social inequality in phase II cardiac rehabilitation attendance: The impact of potential mediators. *European journal of cardiovascular nursing* 2018;17(4):345–55.
- 10 Verlinde E, Laender N de, Maesschalck S de, et al. The social gradient in doctor-patient communication. *Int J Equity Health* 2012;11:12.
- 11 Ghisi GLdM, Chaves GSdS, Britto RR, et al. Health literacy and coronary artery disease: A systematic review. *Patient Educ Couns* 2018;101(2):177–84.

- 1
2
3 12 Sørensen K, Pelikan JM, Röthlin F, et al. Health literacy in Europe: comparative results of
4 the European health literacy survey (HLS-EU). *Eur J Public Health* 2015;25(6):1053–58.
5
6 13 Alter DA, Naylor CD, Austin P, et al. Effects of Socioeconomic Status on Access to
7 Invasive Cardiac Procedures and on Mortality after Acute Myocardial Infarction. *N Engl J*
8 *Med* 1999;341(18):1359–67.
9
10 14 Lemstra ME, Alsabbagh W, Rajakumar RJ, et al. Neighbourhood income and cardiac
11 rehabilitation access as determinants of nonattendance and noncompletion. *Can J Cardiol*
12 2013;29(12):1599–603.
13
14 15 Perelman J, Shmueli A, McDonald KM, et al. Inequality in treatment use among elderly
15 patients with acute myocardial infarction: USA, Belgium and Quebec. *BMC Health Serv*
16 *Res* 2009;9(1):130.
17
18 16 Philbin EF, McCullough PA, DiSalvo TG, et al. Socioeconomic status is an important
19 determinant of the use of invasive procedures after acute myocardial infarction in New
20 York State. *CIRCULATION* 2000;102(19 Suppl 3):15.
21
22 17 Manderbacka K. Exploring gender and socioeconomic differences in treatment of coronary
23 heart disease. *Eur J Public Health* 2005;15(6):634–39.
24
25 18 Pedersen M, Overgaard D, Andersen I, et al. Experience of exclusion: A framework
26 analysis of socioeconomic factors affecting cardiac rehabilitation participation among
27 patients with acute coronary syndrome. *European journal of cardiovascular nursing*
28 2017;16(8):715–23.
29
30 19 Schröder SL, Fink A, Schumann N, et al. How socioeconomic inequalities impact
31 pathways of care for coronary artery disease among elderly patients: Study protocol for a
32 qualitative longitudinal study. *BMJ Open* 2015;5(11):e008060.
33
34 20 Jöckel K, Babitsch B, Bellach B, et al. Messung und Quantifizierung soziographischer
35 Merkmale in epidemiologischen Studien. In: Ahrens W, Bellach B-M, Jöckel K-H,
36 eds. *Messung soziodemographischer Merkmale in der Epidemiologie*. München: MMV
37 Medizin Verlag, 1998:7-38.
38
39 21 Hoffmeyer-Zlotnik J. "Stellung im Beruf" als Ersatz für eine Berufsklassifikation zur
40 Ermittlung von sozialem Prestige. In: *ZUMA Nachrichten* 27 (2003), 53, pp. 114-127:114–
41 27.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 22 Schreier M. Qualitative content analysis in practice. Los Angeles, London, New Dehli,
4 Singapore, Washington DC: Sage 2012.
5
6 23 Willems S, Maesschalck S de, Deveugele M, et al. Socio-economic status of the patient
7 and doctor-patient communication: does it make a difference? *Patient Educ Couns*
8 2005;56(2):139–46.
9
10 24 Hoevenaars JGMM, Schouten JSAG, van den Borne B, et al. Socioeconomic differences in
11 glaucoma patients' knowledge, need for information and expectations of treatments. *Acta*
12 *Ophthalmol Scand* 2006;84(1):84–91.
13
14 25 Czarny MJ, Nathan AS, Yeh RW, et al. Adherence to dual antiplatelet therapy after
15 coronary stenting: a systematic review. *CLINICAL CARDIOLOGY* 2014;37(8):505–13.
16
17 26 Gordon K, Smith F, Dhillon S. Effective chronic disease management: Patients'
18 perspectives on medication-related problems. *Patient Educ Couns* 2007;65(3):407–15.
19
20 27 Dankner R, Drory Y, Geulayov G, et al. A controlled intervention to increase participation
21 in cardiac rehabilitation. *Eur J Prev Cardiol* 2015;22(9):1121–28.
22
23 28 Nolan MT, McKee G. Is Knowledge Level of Coronary Heart Disease and Risk Factors
24 Among Post-Percutaneous Coronary Intervention Patients Adequate? *J Cardiovasc Nurs*
25 2016;31(3):E1-9.
26
27 29 Nunes SRT, Rego G, Nunes R. Right or duty of information: A Habermasian perspective.
28 *Nurs Ethics* 2016;23(1):36–47.
29
30 30 Al Hamarneh YN, Crealey GE, McElnay JC. Coronary heart disease: Health knowledge
31 and behaviour. *Int J Clin Pharm* 2011;33(1):111–23.
32
33 31 Kayaniyil S, Ardern CI, Winstanley J, et al. Degree and correlates of cardiac knowledge
34 and awareness among cardiac inpatients. *Patient Educ Couns* 2009;75(1):99–107.
35
36 32 Tchicaya A, Braun M, Lorentz N, et al. Social inequality in awareness of cardiovascular
37 risk factors in patients undergoing coronary angiography. *Eur J Prev Cardiol*
38 2013;20(5):872–79.
39
40 33 Fennessy MM, DeVon HA, Ryan C, et al. Changing illness perceptions and adherence to
41 dual antiplatelet therapy in patients with stable coronary disease. *J Cardiovasc Nurs*
42 2013;28(6):573–83.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 34 Kucukarslan SN. A review of published studies of patients' illness perceptions and
4 medication adherence: lessons learned and future directions. *Res Social Adm Pharm*
5 2012;8(5):371–82.
6
7
8 35 Alter DA, Franklin B, Ko DT, et al. Socioeconomic status, functional recovery, and long-
9 term mortality among patients surviving acute myocardial infarction. *PLoS One*
10 2014;8(6):e65130.
11
12 36 Smith SK, Dixon A, Trevena L, et al. Exploring patient involvement in healthcare decision
13 making across different education and functional health literacy groups. *Soc Sci Med*
14 2009;69(12):1805–12.
15
16 37 Kambhampati S, Ashvetiya T, Stone NJ, et al. Shared Decision-Making and Patient
17 Empowerment in Preventive Cardiology. *Curr Cardiol Rep* 2016;18(5):49.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1. Characteristics of the patients at baseline (n = 41).

Characteristics	Overall patients	Low-SES patients	High-SES patients
Total	41	23	18
Age, mean	69.7	69.7	69.6
Male	26	10	16
Female	15	13	2
Publicly insured	41	23	18
One-vessel disease	13	8	5
Two-vessel disease	11	6	5
Three-vessel disease	17	9	8
CHD diagnosed for up to 1 year	20	13	7
CHD diagnosed longer than 1 year ago	21	10	11

CHD, coronary heart disease; SES, socioeconomic status

Figure legends

Fig 1. Three themes characterizing socioeconomic differences in the patients' experiences with coronary heart disease treatment.

Supporting information

S1 Appendix. Interview guides.




Fig 1. Three themes characterizing socioeconomic differences in the patients' experiences with coronary heart disease treatment.

174x164mm (150 x 150 DPI)

Interview Guide Baseline

Key question	Concrete supplementary questions
Please begin by telling me about the medical history of your heart disease starting from the first symptoms until this hospital stay.	<ul style="list-style-type: none"> - Have you ever been to a general practitioner/ cardiologist/ in a rehabilitation clinic before because of your heart disease? - How was your disease diagnosed? - Why are you currently in hospital?
Which positive and negative experiences have you made, concerning the care you have received?	<ul style="list-style-type: none"> - Can you think of any other positive or negative experience you have made with the hospital or office-based physicians? - Was there anything you were particularly satisfied with? - Was there anything you were unsatisfied with? - Have you experienced any problems with the further processing of the treatment? - How have you experienced talking with your physicians? - How was a decision reached about what treatment you were to receive?
How would you describe the quality of your treatment?	<ul style="list-style-type: none"> - How satisfied are you with your treatment? - Do you think that everyone in Germany is able to receive good treatment? Why do you think that is?
How do you manage your heart disease right now?	<ul style="list-style-type: none"> - What will happen next? - What medical care do you expect to receive in the next months? - What hopes do you have for your health in the future? - Is there something else you would like to tell me?

Questions aimed at maintaining the conversational flow:

- Please tell me exactly how things went with...
- Could you perhaps give me a few more details?
- What happened next? / And after that?
- What else comes into your mind?
- What do you associate with ...?

Interview Guide Follow Up

Key question	Concrete supplementary questions
Please begin by telling me how the treatment of your heart disease has progressed after our last conversation.	<ul style="list-style-type: none"> - Which doctors have you seen meanwhile, e.g. general practitioner (GP)/ cardiologist? - Have you been in a rehabilitation clinic or do you attend a heart training group? - What exactly has been done by each of the physicians (GP, cardiologist, hospital, rehabilitation)? <ul style="list-style-type: none"> - Which tasks of medical care have been taken care of by which doctor? - Who prescribes you heart drugs?
What positive and negative experiences have you made, concerning the treatment and care you have received?	<ul style="list-style-type: none"> - Can you think of any other experiences you have had with the GP/ cardiologist/ rehabilitation clinic? - Was there anything you were particularly satisfied or not satisfied with? - How does the cooperation between GP, specialists and the doctors at the hospital work? - Have you experienced any problems with the postoperative management and any further treatment after you have been discharged from hospital? - Have you had to actively arrange yourself to the further treatment? - Have you ever changed your GP or cardiologist and what have been the particular reasons?
Last time we talked about your expectations of your medical care – to what extent were they fulfilled?	<ul style="list-style-type: none"> - Have your expectations changed during the course of treatment?
What kind of influence does your heart disease have on your everyday life?	<ul style="list-style-type: none"> - What kind of heart disease/s do you have? - How far is your everyday life constrained by your heart disease, and which tasks can't be managed by yourself anymore? - Who assists you in coping with the disease? - What do you personally contribute to a better health?
What will happen next?	<ul style="list-style-type: none"> - What hopes do you have for your future health? - Is there something else you would like to tell me?

Questions aimed at maintaining the conversational flow:

- Please tell me exactly how things went with...
- Could you perhaps give me a few more details?
- What happened next? / And after that?
- What else comes into your mind?
- What do you associate with ...?

Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist

<i>No. Item</i>	<i>Guide questions/description</i>	<i>Page</i>
Domain 1: Research team and reflexivity		
<i>Personal Characteristics</i>		
1. Interviewer/facilitator	Which author/s conducted the interview or focus group?	7
2. Credentials	What were the researcher's credentials? E.g. PhD, MD	1/7
3. Occupation	What was their occupation at the time of the study?	1/7
4. Gender	Was the researcher male or female?	7
5. Experience and training	What experience or training did the researcher have?	7
<i>Relationship with participants</i>		
6. Relationship established	Was a relationship established prior to study commencement?	6-7
7. Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	7
8. Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? e.g. Bias, assumptions, reasons and interests in the research topic	7
Domain 2: study design		
<i>Theoretical framework</i>		
9. Methodological orientation and Theory	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	8
<i>Participant selection</i>		
10. Sampling	How were participants selected? e.g. purposive, convenience, consecutive, snowball	6
11. Method of approach	How were participants approached? e.g. face-to-face, telephone, mail, email	6
12. Sample size	How many participants were in the study?	6/8
13. Non-participation	How many people refused to participate or dropped out? Reasons?	6
<i>Setting</i>		
14. Setting of data collection	Where was the data collected? e.g. home, clinic, workplace	7
15. Presence of non-participants	Was anyone else present besides the participants and researchers?	7
16. Description of sample	What are the important characteristics of the sample? e.g. demographic data, date	8/ Table 1
<i>Data collection</i>		
17. Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	7/ Appendix S1
18. Repeat interviews	Were repeat interviews carried out? If yes, how many?	6-7
19. Audio/visual recording	Did the research use audio or visual recording to collect the data?	7
20. Field notes	Were field notes made during and/or after the interview or focus group?	7
21. Duration	What was the duration of the interviews or focus group?	

<i>No. Item</i>	<i>Guide questions/description</i>	<i>Page</i>
22. Data saturation	Was data saturation discussed?	6
23. Transcripts returned	Were transcripts returned to participants for comment and/or correction?	8
Domain 3: analysis and findings		
<i>Data analysis</i>		
24. Number of data coders	How many data coders coded the data?	8
25. Description of the coding tree	Did authors provide a description of the coding tree?	No
26. Derivation of themes	Were themes identified in advance or derived from the data?	8
27. Software	What software, if applicable, was used to manage the data?	8
28. Participant checking	Did participants provide feedback on the findings?	8
<i>Reporting</i>		
29. Quotations presented	Were participant quotations presented to illustrate the themes / findings? Was each quotation identified? e.g. participant number	9-14
30. Data and findings consistent	Was there consistency between the data presented and the findings?	yes
31. Clarity of major themes	Were major themes clearly presented in the findings?	13-14
32. Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	9-13

BMJ Open

Socioeconomic differences in experiences with treatment of coronary heart disease: a qualitative study from the perspective of elderly patients

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-024151.R1
Article Type:	Research
Date Submitted by the Author:	26-Jul-2018
Complete List of Authors:	Schröder, Sara Lena; Martin Luther University Halle-Wittenberg, Germany, Institute of Medical Sociology Fink, Astrid; Martin Luther University Halle-Wittenberg, Institute of Medical Sociology Richter, Matthias; Martin Luther University Halle-Wittenberg, Institute of Medical Sociology
Primary Subject Heading:	Health services research
Secondary Subject Heading:	Cardiovascular medicine, Public health, Qualitative research
Keywords:	health services research, socioeconomic status, Coronary heart disease < CARDIOLOGY, QUALITATIVE RESEARCH, healthcare inequalities, health services accessibility

SCHOLARONE™
Manuscripts

1
2
3 **Socioeconomic differences in experiences with treatment of coronary heart disease: a**
4
5 **qualitative study from the perspective of elderly patients**
6
7

8
9
10 Sara L. Schröder¹, Astrid Fink¹ (PhD), Matthias Richter¹ (PhD)
11
12

13
14 ¹Institute of Medical Sociology, Martin Luther University Halle-Wittenberg, Halle (Saale),
15
16 Germany
17
18

19
20
21 **Corresponding author:**
22

23 Sara Lena Schröder
24

25
26 Institute of Medical Sociology (IMS)
27

28 Martin Luther University Halle-Wittenberg
29

30
31 Magdeburger Str. 8
32

33 06112 Halle (Saale)
34

35 Germany
36

37 Email: sara.schroeder@medizin.uni-halle.de
38
39

40
41
42 Word count: **4761**
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives: This qualitative study aims to analyse socioeconomic differences in patients' experiences along the treatment pathway for coronary heart disease (CHD).

Design: A longitudinal qualitative study using in-depth semi-structured interviews to explore patients' experiences with treatment was conducted. We analysed the transcripts of the records according to qualitative content analysis and identified differences between patients with lower and higher socioeconomic status (SES) by comparing and contrasting the narratives.

Setting: The University Hospital in Halle (Saale), Germany.

Participants: Forty-one elderly patients (aged 59-80) who suffered from CHD.

Results: From various patient's experiences along the pathway of care which were found to differ according to SES we derived three major themes: (1) information: patients with higher SES had greater knowledge about treatment and could use medical records as sources of information; (2) illness perception: patients with lower SES focused on improving symptoms and survival, while patients with higher SES focused on physical performance and disease management; and (3) perceived role in healthcare: patients with lower SES tended to delegate responsibility to healthcare professionals.

Conclusions: Differences in the patient's knowledge about treatment, their perceived role in healthcare, and illness perception can be the factors and mechanisms that contribute to explain

1
2
3 socioeconomic inequalities in the treatment of CHD. These factors should be considered in
4
5 quantitative studies to better understand the disparities in treatment and mortality. We suggest
6
7 that improving patient-physician-communication and patient knowledge can change the patient's
8
9 understanding of CHD as well as their perceived role in healthcare and reduce inequalities in
10
11 understanding of CHD as well as their perceived role in healthcare and reduce inequalities in
12
13 CHD treatment.
14
15
16

17 **Trial registration number:** DRKS00007839.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5 **Keywords:** coronary heart disease, socioeconomic status, healthcare inequalities, qualitative
6
7
8 research, health services accessibility
9

10 **Strengths and Limitations**

- 11 - The longitudinal design enabled us to collect information regarding the patient's
12 experiences along the treatment pathway for coronary heart disease from a first
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- We performed a comparative analysis to explore differences between patients with lower
and higher socioeconomic status regarding their perspectives and experiences with
treatment.
- Generalizability in this study might be limited as we only recruited elderly patients in a
large city in the eastern part of Germany; therefore, the results may not reflect the
experiences of younger patients with coronary heart disease or those in countries without
universal healthcare systems.

36 **Funding**

This work was supported by the Wilhelm Roux Programme of the Medical Faculty of
Martin Luther University, Halle-Wittenberg with grant number FKZ: 28/40.

43 **Competing Interests**

The authors declare that they have no competing interests.

47 **Author Contributions:**

SLS wrote the initial draft of this manuscript, collected data and analysed the transcripts.
MR and AF conceptualized and designed the project, helped interpreting the data, and critically
reviewed the manuscript. All authors have read and approved the final version of the manuscript.

Data Sharing Statement

Participants of this study were guaranteed that only the study research team would have access to the interviews and transcripts thus data are not available for sharing.

Ethics Approval

Approval for the study was obtained from the Ethical Review Committee of the Medical Faculty of Martin Luther University, Halle-Wittenberg (No. 2014-95).

For peer review only

Introduction

Coronary heart disease (CHD) remains the leading cause of death worldwide.¹⁻³ Treatment of CHD includes revascularization, rehabilitation and long-term medication and aims to improve disease-related quality of life, including exercise capacity, to prevent further cardiac events and reduce mortality. Socioeconomic inequalities in the incidence and mortality of CHD, to the disadvantage of patients with low socioeconomic status (SES), have been well explored.⁴⁻⁶ Therefore, patients of low SES have a greater need for treatment. Access to treatment is aimed to be on the basis of need rather than of gender, residence or SES, and guidelines note the importance of reducing inequalities in healthcare.⁷ However, several studies showed strong socioeconomic differences in CHD treatment, to the disadvantage of patients with low SES.⁸

How inequalities in healthcare arise is complex and largely unknown. As socioeconomic differences in treatment are provable regardless of the underlying healthcare system, they may be attributable to reasons other than financial affordability of healthcare costs.⁸ This is particularly the case in countries such as Germany, where individuals have statutory health insurance and very low out-of-pocket payments. In cardiac rehabilitation, a recent study found that comorbidities, self-efficacy, anxiety and depression, cohabitation, commute, disease severity or type of treatment do not significantly mediate the association of SES with attendance and participation.⁹ However, health literacy and communication between physician and patient are discussed to generally explain healthcare inequalities, as they are strongly associated with SES.¹⁰⁻¹² Although there is no clear evidence regarding whether an increase in treatment rates reduces socioeconomic disparities in access to treatment,^{13, 14} Perelman et al. found that socioeconomic inequalities in CHD treatment can be partly explained by different distances to hospitals with on-site cardiac facilities.¹⁵

1
2
3 Therefore, the underlying causes and determinants of socioeconomic inequalities in CHD
4 treatment are still not fully understood or analysed. Taking the patients' individual needs and
5 perceptions into account is helpful to reveal new explanatory approaches from the patient's
6 perspective and explore how socioeconomic differences might arise. Qualitative research aims to
7 understand healthcare interactions from the patient's perspective and can help to identify the
8 mechanisms that lead to inequalities in healthcare. To our knowledge, there are only two
9 qualitative studies exploring socioeconomic differences in the treatment of CHD. Manderbacka
10 analysed differences in healthcare encounters and found that doctor-centred decision making was
11 more common in lower-SES patients.¹⁶ Pedersen et al. found that concerning barriers to cardiac
12 rehabilitation, only lower-SES patients felt excluded due to their divergent health beliefs.¹⁷ To
13 gather scientific evidence on the possible factors and mechanisms of socioeconomic inequalities
14 in CHD treatment, we aimed to identify socioeconomic differences in the patient's perspective
15 and their experiences with the treatment pathway for CHD in all sectors from therapy to
16 aftercare.

35 **Material and methods**

37 **Study design**

39
40 Based on the methodology of grounded theory, we conducted an exploratory qualitative
41 longitudinal study of elderly CHD patients to examine socioeconomic differences in access,
42 utilization and quality of treatment.¹⁸ A purposive sampling strategy was used to select patients
43 with a confirmed diagnosis of CHD at a university hospital in Germany. Additionally, the
44 patients had to fulfil the inclusion criteria of being between 60 and 80 years and having one
45 additional principal or secondary diagnosis of stable angina pectoris, acute coronary syndrome or
46 cardiac arrhythmia. Patients were excluded from the study if they had insufficient language skills
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 to conduct an interview in German language or were moribund. Between November 2014 and
4
5 April 2015, a study nurse contacted patients who were hospitalized in the department of
6
7 cardiology, informed them through a flyer and explained the purpose of the study. We
8
9 approached 96 eligible patients and asked them to attend an interview to share their experiences
10
11 with CHD treatment. Participation was voluntary, the patients received no incentives, and 61
12
13 patients agreed to participate. Of those, we gradually chose 48 for an interview in accordance
14
15 with maximum variation to assure that men and women with different SES and different CHD
16
17 severities (number of atherosclerotic altered vessels, type of revascularization needed, and
18
19 manifestations of angina pectoris or myocardial infarction) were represented who covered a
20
21 broad spectrum of treatment experience. To achieve maximum variation, we also included one
22
23 59-year-old patient because of relevant experiences. Patients were only enrolled in the study after
24
25 providing written informed consent. After 48 interviews theoretical saturation in terms of
26
27 experiences with diagnosis and treatment of different CHD severities was reached. We stopped
28
29 recruiting new interviewees when the experiences of new interviews were very similar to the
30
31 narratives reported earlier by other patients, and no new categories emerged during inductive
32
33 coding. After 6 months, we contacted the 48 participants again and asked them to participate in a
34
35 follow-up interview. Eight patients refused to participate due to their disease burden, lack of
36
37 interest, or failure to respond to the invitation letter, and one patient was no longer alive. The
38
39 thirty-nine follow-up interviews were conducted between June and October 2015. The study was
40
41 approved by the Ethics Review Committee of the Medical Faculty at the Martin Luther
42
43 University, Halle-Wittenberg (No. 2014-95).

51 **Data collection**

52
53
54
55
56
57
58
59
60

1
2
3 We conducted semi-structured interviews to explore the patients' perspectives on their
4 heart disease and their personal experiences with the treatment they received. The interview
5 guide (S1 Appendix) was developed according to Helfferich with key questions that evoke
6 narrations, which were derived from the state of research.¹⁹ To cover the same topics within each
7 interview, these key questions were supplemented by specific questions, which were only asked
8 if not mentioned by the patient himself. The key and supplementary questions were developed by
9 a process of collecting, checking, sorting, and subsuming topics and questions of relevance in
10 accordance with the research question and aim. SLS conducted two preliminary interviews that
11 were used to further develop the semi-structured interview guides. We began the baseline
12 interviews with a broad general question about the medical history. More detailed questions were
13 asked individually based on each patient's narrative. All Interviews were conducted and analysed
14 in German language. For the Quotations and interview guide presented in this manuscript we
15 conducted a double-blind translation from German to English that was checked by a third person.
16 The patient's basic sociodemographic data were obtained with a short, standardized
17 questionnaire following the baseline interview, and the interviewer subsequently wrote a field
18 note for each interview.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Baseline interviews were conducted in a private room of the hospital, and follow-up interviews were held either at the patient's home or at the medical facility, depending on the patient's choice. Two employees of the Institute of Medical Sociology (IMS) conducted the interviews face-to-face: SLS (female) is a research associate and an economist who has conducted qualitative health research since 2009 and conducted most of the interviews; Nils Bormann (male) is a medical student assistant and an economist who was trained prior to conducting the interviews. The interviewers introduced themselves as members of the IMS, not

1
2
3 working in the department of cardiology. Each baseline interview included only the patient and
4 one or both interviewers. In some follow up interviews, family members were present at the
5 patient's request. The interviews lasted an average of 35 minutes at baseline and 42 minutes at
6 follow-up and were audiotaped with the interviewee's permission.
7
8
9

10
11
12 SES was defined by educational level and occupation based on the German
13 epidemiological standards.²⁰ Education was measured by level of schooling and academic
14 qualifications and was classified on an 8-point scale. Occupation was measured by the last
15 occupational group the patient belonged to and classified according to job autonomy on a 5-point
16 scale.²¹ A composite index of SES was derived using a sum score of both items, which ranged
17 between 2 and 13. Index values between 2 and 7 were rated "lower SES", and values between 8
18 and 13 were rated "higher SES".
19
20
21
22
23
24
25
26
27

28 **Data analysis**

29
30
31 After constructing the main categories that originated from the research questions, SLS
32 coded the pseudonymised interview transcripts in a data-driven approach (inductively) sentence
33 by sentence with codes and themes derived from the data in accordance with qualitative content
34 analysis.²² The coding started simultaneously with the conducting of the interviews. Five
35 contrasting transcripts were double-coded by two additional members of the qualitative research
36 working group at the IMS, who also participated in the discussion and evaluation of the data. The
37 working group was composed of researchers with different backgrounds, namely, sociology,
38 healthcare research, economics, nursing and educational science. The emerging codes were
39 organized into subcategories with the existing main categories. Afterwards, we identified
40 socioeconomic differences by comparing and contrasting the codes of the frame between patients
41 with lower and higher SES. Codes that were found only in one group constituted the origin of
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 comparing and contrasting the narratives to identify SES-specific patient perspectives and
4
5 experiences in each phase of treatment. Building on these results, we constructed overall major
6
7 themes, which characterized the differences across the continuum of care. We used MAXQDA
8
9
10 11 software to assist with the data management and analyses. In the present analysis, we
11
12 excluded 7 patients because neither CHD nor any CHD-indicative treatment was mentioned
13
14 throughout their interviews as the narratives of treatment experiences had been overshadowed by
15
16 other heart diseases. Additionally, to the 41 baseline interviews, we included the narratives of
17
18 those 17 follow-up interviews in the current analysis, which provided important additional
19
20 information to answer the research question. In total, we analysed 58 interviews from 41
21
22 patients.
23
24

25 26 **Patient involvement**

27
28 By exploring patients' experiences with treatment in this study through an exploratory
29
30 design, the patients' preferences and priorities led the data collection during the interviews and
31
32 due to an inductive data-driven approach as well, throughout data analysis. The participants did
33
34 not provide feedback on their transcripts or the findings.
35
36

37 38 **Results**

39
40 This analysis was based on the transcripts of 58 interviews (41 baseline and 17 follow-up) with
41
42 41 patients. The baseline characteristics of the 41 patients are shown in Table 1. Throughout the
43
44 narratives, treatment of CHD was reported as a complex process with the involvement of many
45
46 providers. The majority of the patients reported utilizing treatment, e.g., revascularization at the
47
48 hospital; inpatient cardiac rehabilitation; visiting their general practitioner (GP) regularly, who
49
50 prescribes the medication as recommended from the hospital; and having regular appointments
51
52 with a resident cardiologist. Only a few patients (from both SES groups) reported having
53
54
55
56
57
58
59
60

1
2
3 problems with access to, utilization of or quality of treatment. However, we found differences
4
5 according to SES in the patients' individual perceptions and perspectives on treatment, which are
6
7 described separately hereafter subdivided into four sections: treatment at the hospital,
8
9 rehabilitation, treatment with drugs and ambulatory aftercare. Quotations reflective of the
10
11 differences are presented for patients of lower and higher SES at the various stages of treatment.
12
13 Following, three major themes, which were derived from the sector-specific differences, are
14
15 depicted: "information", the "patient's individual perception of CHD" and the "perceived role in
16
17 healthcare". These themes might characterize the factors and mechanisms of socioeconomic
18
19 differences in treatment across the continuum of care.
20
21
22

23 24 **Treatment at the hospital**

25
26 With regard to revascularization, some patients reported that a second coronary
27
28 angiography (CAG) was scheduled and conducted within a few days for different reasons.
29
30 However, only lower-SES patients did not scrutinise or mention the reason when a diagnostic-
31
32 only CAG was followed by a second CAG with stent implantation. Additionally, whereas many
33
34 patients of both lower and higher SES rated it important for hospital staff to be friendly, attentive
35
36 and caring, predominantly lower-SES patients mentioned aspects of being taken seriously either
37
38 positively (e.g., taking preferences seriously, being asked how one feels) or negatively (e.g.,
39
40 discussions about but not with oneself, no conversations at all). While assessing the quality of
41
42 treatment, some lower-SES patients complained about being discharged with open wounds after
43
44 coronary artery bypass graft surgery. Generally, while assessing the quality of treatment, lower-
45
46 SES patients tended to focus on symptom improvement, eliminating the need for visits to the
47
48 doctor or improving survival, whereas higher-SES patients predominantly aimed to improve
49
50 physical performance.
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5 “Well, in the hope that everything stays the same. Let’s say that you are well. Well,
6 getting around with it quite well, with no complications arising. As I said, that one may reach a
7 slightly older age, not just until retirement, but perhaps even a bit longer.” (Mr. Lange, 60 years,
8 lower SES)
9
10
11
12
13

14 “Well, but then it came up with the heart and I myself had the feeling that it is probably
15 the only thing that really gets you back on your feet. I was really feeling sluggish. So I did
16 indeed make an effort to try to walk again quickly and to/ I really did, but somehow in the end
17 the energy was lacking.” (Mr. Lehmann, 63 years, higher SES)
18
19
20
21
22
23

24 Knowledge of different treatment options became apparent mainly in patients with high
25 SES. However, an information deficit was found in some narratives, e.g., patients reporting that
26 they did not know the outcome of medical examinations; they were unaware of the treatment that
27 had been performed; or they rated the medical reports incomprehensible. An information deficit
28 became apparent mainly in lower-SES patients, although this was not always experienced with a
29 perceived greater need for information.
30
31
32
33
34
35
36
37

38 “Yes, it has been explained to me, but I didn’t get it. First of all, I sometimes didn’t
39 understand it, because he expressed himself in such a professional manner. The nurses already
40 started laughing, because I/ and then I thought: ‘No, you better give up. You better read through
41 it next time’” (Mrs. Schubert, 80 years, lower SES)
42
43
44
45
46

47 “For this there are, that’s at least what has been explained to me, the clogged coronary
48 arteries can be expanded by implanting stents, if you know what these are, incorporating these
49 small tubes that eliminate the narrowed areas. Or if that is not feasible, which can happen as well
50
51
52
53
54
55
56
57
58
59
60

1
2
3 that a vessel ruptures or something like that, then (...) it is done surgically. Then bypasses are
4 produced from other parts of the body and incorporated.” (Mr. Wagner, 75 years, higher SES)
5
6

7 We found paternalistic and shared decision making in both SES groups, but paternalistic
8 decisions were found to be more unquestioned in lower-SES patients, whereas higher-SES
9 patients understood the reasons why decisions were made. When decisions were made jointly,
10 we found that higher-SES patients were involved more actively in decision making, whereas
11 lower-SES patients thought of themselves as being only responsible for responding (agree or
12 disagree) to a recommendation given.
13
14
15
16
17
18
19
20

21 “With this I entirely relied on the physicians. In fact, I didn’t really thought about it.
22 When they said: ‘This’, then we do this. They do ask indeed: ‘Mr. Zimmermann, do you agree?’
23 I mean, when I go to the hospital, I let myself be treated, but this is not meant to be negative in
24 any way. But I rely on the physicians. [...] They are the professionals. I would never pretend to
25 be the wise guy, as I know people who do so, not physicians, but private persons, who dictate
26 people what to do.” (Mr. Zimmermann, 76 years, lower SES)
27
28
29
30
31
32
33
34

35 “Today, quite a few medical exams have been conducted with me. Now you have to wait,
36 but I guess it won’t change anything about this final decision. The demand placed on me to
37 decide this.” (Mr. Hartmann, 69 years, higher SES)
38
39
40
41

42 **Cardiac rehabilitation**

43 Some patients reported whether the time interval between their discharge from hospital
44 and the beginning of inpatient cardiac rehabilitation was either too long or too short for their
45 individual needs. In doing so, only lower-SES patients reported they could not participate in
46 therapeutic cardiac rehabilitation treatments due to their recent coronary artery bypass surgery,
47 whereas only higher-SES patients focused their reasons on individual demands for disease
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 management. Additionally, we found that while assessing the quality of cardiac rehabilitation, in
4 the narratives of lower-SES patients, performing regular examinations, such as daily blood
5 pressure measurement or electrocardiograms, was more important, whereas higher-SES patients
6 tended to place an emphasis on physical performance improvement, as well as increased
7 knowledge about necessary behaviour changes and their heart disease in general.
8
9
10
11
12
13

14 “Indeed, in rehabilitation it was great. Well, there was/ every morning you had to go to
15 the presentation, weighing, measuring blood pressure and stuff like that.” (Mrs. Koch, 62 years,
16 lower SES)
17
18
19
20

21 “Over there I primarily learned through lectures and, as I said before, through meetings
22 about what the issue of a heart attack means, how it emerges and so on. This improved much of
23 my understanding. Previously, before you have something like that you don’t care about it. So,
24 concerning this matter one was educated quite well.” (Mr. Jung, 67 years, higher SES)
25
26
27
28
29
30

31 **Treatment with drugs**

32
33 Although specific drugs, agents or trade names were rarely mentioned throughout the
34 narratives, access to and utilization of any medications were fairly high in both SES groups.
35 However, we found that with respect to medication preferences, some lower-SES patients
36 highlighted the importance that the involved physicians should appropriately coordinate the
37 selection of prescribed drugs, while some stated they generally did not like to take drugs.
38
39
40
41
42
43
44
45 However, one higher-SES patient reported that he prefers to take combination medication.
46

47 “Or, for example, if there are changes in medication, it is very important. With
48 prothrombin time and everything; you need to be always in good hands. Not that one says this
49 way and the other one says that way. That’s bad.” (Mr. Köhler, 66 years, lower SES)
50
51
52
53
54
55
56
57
58
59
60

1
2
3 “I have to take many drugs; they upset the stomach and everything. I had to take 17 or 18
4 different pills. Well, and then I recognized, I read about it, there is a patch and so on and using
5 that you can come off of 5 drugs at once. The patches are indeed very expensive and the doctor
6 immediately said ‘no and no and no’ and so on. Now I have this patch and I am feeling fine.”
7
8
9

10
11
12 (Mr. Richter, 66 years, higher SES)
13

14 **Ambulatory aftercare**

15
16
17 Some patients in the lower-SES group mentioned undergoing regular check-ups every
18 month with their GP to assess prothrombin time (Quick-test). While rating the quality of GPs,
19 mainly lower-SES patients mentioned the importance of thorough medical examinations and
20 doctor availability. Some mainly lower-SES patients complained about the GP examining or
21 referring patients only on demand or the GP not helping them make an appointment with a
22 cardiologist. Some predominantly lower-SES patients reported that they see themselves as
23 responsible for retention of their medical record. However, some patients, mainly with higher
24 SES, mentioned obtaining or inquiring for a recommendation from the GP regarding a particular
25 specialist or hospital.
26
27
28
29
30
31
32
33
34
35
36

37
38 “However, I wasn’t correctly/ At first I didn’t know something like that existed. Then, I
39 asked my doctor. I say: ‘Doesn’t one have to go to any follow up?’ - ‘Well, I could forward you
40 there, but you better try to make an appointment yourself’.” (Mrs. Körtig, 59 years, lower SES)
41
42
43

44
45 “In one consultation it was like, ‘To whom should I go?’ And then he said, ‘I have a
46 number of colleagues here that I can recommend to you’ And then I say, ‘Alright, then make an
47 appointment for me’, and that’s how I came to Mrs. Dr. Alpha.” (Mr. Winkler, 72 years, higher
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 In addition, the patient's experiences with access to and utilization of regular check-ups at
4 a resident cardiologist varied according to SES. Some lower-SES patients perceived
5 appointments with specialists as generally unnecessary or expected instructions from physicians
6 about the necessity of appointments due to their own uncertainty. Some mainly higher-SES
7 patients used medical reports as sources of information for the need of follow-up care.
8
9
10
11
12
13

14 "As I said, I had to wait a very long time for this 24-hour ambulatory electrocardiogram.
15 Then it was done. She [cardiologist] wrote her report, my GP got it and nothing else came of it.
16 [...] I mean, if there would have been something acute, surely something would have come up,
17 but it was still, that it is not/" (Mr. Köhler, 66 years, lower SES)
18
19
20
21
22
23

24 "I guess that I will have to go to my GP for follow up and he does further. [...] That is my
25 current state. But I am not sure yet, because I have not received the medical report yet. I am still
26 waiting for it." (Mr. Schäfer, 71 years, higher SES)
27
28
29
30

31 **Factors and mechanisms of socioeconomic differences in CHD treatment**

32
33 While comparing the differences in the patients' experiences according to SES across the
34 pathway of care, three major themes (Fig. 1) emerged to characterize differences and may be the
35 factors and mechanisms for socioeconomic inequalities in CHD treatment, namely, information,
36 patient's individual illness perception and their perceived role in healthcare.
37
38
39
40
41

42 First, the narratives suggest that lower-SES patients are less informed about their
43 treatment than higher-SES patients, which became apparent, e.g., through their lack of
44 knowledge regarding the treatment that was performed, as well as their lack of knowledge about
45 the reasons for or outcomes of medical examinations. However, higher-SES patients knew about
46 different treatment options. Whereas lower-SES patients reported they missed conversations at
47 the hospital or rated the reports as incomprehensible, higher-SES patients explained that they use
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 medical reports as sources of information. While most patients rated the healthcare system
4
5 (especially hospitals) as overloaded, some made the connection that patients are given far too
6
7 little information and are required to actively request information because of excessive work
8
9 demands on health workers due to the healthcare system economization.
10

11
12 “And as well, someone who is cognitively not that fit anymore, one does not understand
13
14 at all what you are told. And that is sometimes not so nice. Well, but probably that’s just our
15
16 system, that is/ which is not working, I think”. (Mrs. Koch, 62 years, lower SES)
17
18

19
20 “You need to directly address the physicians; they do not have the time as well. You can
21
22 notice that during the doctor’s visits. Well, then it is clear. But otherwise you get an answer to
23
24 any question; pleasant, polite, courteous.” (Mr. Schäfer, 71 years, higher SES)
25

26
27 Second, we found differences in patients’ illness perceptions. Whereas lower-SES
28
29 patients seemed to understand CHD through putting emphasis on symptoms and medical
30
31 parameters measured through medical examinations, higher-SES patients focused on physical
32
33 performance and disease management, which requires knowledge about CHD.
34

35
36 Finally, we found differences in the patient's perceived role in their healthcare. Lower-
37
38 SES patients tended to delegate responsibility for treatment, which became apparent, e.g., by the
39
40 patients not questioning the reasons for the decisions made, assuming that healthcare
41
42 professionals are responsible for patients being taken seriously, not perceiving a need to obtain
43
44 information even when lacking information, being less involved in decision making, relying on
45
46 physicians to coordinate care (referrals, medication), viewing themselves as only responsible for
47
48 the retention of medical records, and having a tendency to have general views rather than
49
50 opinions. Contrastingly, higher-SES patients seem to feel more responsible for treatment, e.g.,
51
52
53
54
55
56
57
58
59
60

1
2
3 focusing more strongly on disease management, having increased healthcare knowledge, making
4 informed choices or actively involving their GP.
5
6

7 **Discussion**

8
9
10 In this study, we elaborated three major themes from the patients' experiences with CHD
11 treatment that were found to differ between patients of lower and higher SES: information,
12 the patient's individual perception of CHD and their perceived role in healthcare. Regarding
13 information, patients with higher SES had greater knowledge about treatment and could use
14 medical records as sources of information. Regarding illness perception, patients with lower SES
15 focused on improving symptoms and survival, while patients with higher SES focused on
16 physical performance and disease management. Regarding the patient's perceived role in
17 healthcare, patients with lower SES tended to delegate responsibility to healthcare professionals.
18
19
20
21
22
23
24
25
26
27

28 First, we found that lower-SES patients seem to lack knowledge about treatment
29 compared with higher-SES patients, obtain less information and have problems understanding
30 the information provided to them. This is in accordance with previous studies, which found that
31 knowledge about treatment was a barrier of utilization and access for lower-SES patients, e.g. to
32 optimal oesophageal cancer care or reproductive healthcare.^{23,24} Especially in low-SES patients
33 it is well known that lack of information and knowledge results in low utilization.^{25,26} This is
34 probably due to the fact that patients of lower SES are generally given less information in
35 healthcare.¹⁰ Especially with regard to CHD, studies showed that patients are given little
36 information by health professionals, and therefore, wide information gaps exist for all CHD
37 patients.^{27,28} Second, illness perception differs, as we found that lower-SES patients seem to
38 focus more strongly on symptoms, survival and good results of physical check-ups in their
39 illness perception, while higher-SES patients tend to focus on physical performance and disease
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 management. The latter reveals more knowledge about CHD, and studies found that lack of
4
5 knowledge regarding CHD is associated with both low SES and adherence.^{29–32} Therefore, we
6
7 confirm the findings of another study that only lower SES patients felt excluded from cardiac
8
9 rehabilitation due to their divergent health beliefs.¹⁷ This becomes particularly relevant as
10
11 focusing on and improving physical performance are important for reducing cardiovascular
12
13 mortality and inequalities in AMI mortality, independently of inequalities in access to
14
15 healthcare.³³ Finally, the perceived role in healthcare differed according to SES: higher-SES
16
17 patients tended to assume responsibility for treatment, whereas lower-SES patients showed a
18
19 tendency to relinquish responsibility. This latter finding was characterized by “putting up with
20
21 everything needed” and expecting to be told that “everything is fine”, with no further need to see
22
23 a physician. Different responsibilities are becoming apparent through the decision making
24
25 process between the physician and the patient, as a study found that high-SES patients perceive
26
27 their involvement as participating in the decision making after exploring other options, whereas
28
29 low-SES patients see themselves only being responsible for agreeing or disagreeing with the
30
31 recommendation made.³⁴ Therefore, we confirm another study, which found that inequalities
32
33 exist in the way that shared decision making is more common in high-SES patients.¹⁶ However,
34
35 the patient’s assumed role is likely influenced by the physician, as a review found that physicians
36
37 reported lower levels of facilitating shared decision making with patients who were rated to be
38
39 unable or unwilling to participate in decision making from the physicians perspective or who
40
41 showed limited awareness about their condition.³⁵
42
43
44
45
46
47
48

49 Although we provide novel insights into the possible factors and mechanisms of
50
51 socioeconomic differences in treatment, the study has several limitations. The generalizability
52
53 might be limited as we only recruited elderly patients in the eastern part of Germany. Therefore,
54
55
56
57
58
59

1
2
3 these results may not reflect the experiences of younger patients or those living in other regions.
4
5 Importantly, data were generated in a country with statutory health insurance and financial
6
7 concerns may better explain socioeconomic differences in treatment in other countries without
8
9 universal healthcare systems. There might be transferability of some of our findings to settings of
10
11 CHD treatment with similar organizational and economic contexts. We increased the rigour of
12
13 the study by taking the consolidation criteria for reporting qualitative research (COREQ) into
14
15 account during the research process.³⁶ Reflexivity was increased by using content analysis with a
16
17 coding frame as well as coding the interviews and discussing the results with different
18
19 researchers of a qualitative working group, and reporting on the process of interpretation within
20
21 the results section of the manuscript. Only a few patients reported problems with access to and
22
23 utilization of treatment. Throughout the narratives, only some experiences and viewpoints on
24
25 treatment were found to differ according to SES. However, consideration of these aspects
26
27 together led us to generate three data-driven major themes, which may elucidate the possible
28
29 mechanisms of socioeconomic inequalities in treatment. But we did not find differences in
30
31 treatment according to SES. Especially when assessing the patient's knowledge, information,
32
33 illness perception and delegation of responsibility, it needs to be considered that we only
34
35 analysed the patients' perspectives and viewpoints independently of the information that were
36
37 given through communication by healthcare staff. Finally, as we operationalised SES
38
39 dichotomously based on education and occupation, we simplified the continuum of social status.
40
41 We did not collect data on income, as income presents a sensitive personal issue, especially in
42
43 longitudinal studies, and for older people income is a less appropriate indicator for SES because
44
45 it is highly affected by retirement and difficult to collect reliable information.³⁷
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Given that CHD is the leading cause of death worldwide, the generation of hypotheses
4 regarding the potential factors underlying the socioeconomic differences in CHD treatment is
5 essential for explaining and reducing these inequalities. The level of information and knowledge,
6 individual illness perception and perceived role in healthcare can be the mechanisms that
7 contribute to explain socioeconomic inequalities in CHD treatment and should be considered in
8 further quantitative studies as mediating factors between SES and treatment. We conclude that
9 improvement in patient-provider communication might be the key to reducing healthcare
10 inequalities, as it may increase knowledge about treatment, improve the understanding of CHD,
11 and empower patients to assume responsibility in treatment, which can improve engagement in
12 treatment and utilization.
13
14
15
16
17
18
19
20
21
22
23
24

25 26 **Acknowledgments**

27
28 We acknowledge the patients who participated in this study and shared their healthcare
29 experiences with us. We thank the staff at the Department of Cardiology at the University
30 Hospital in Halle for their assistance with screening patients for this study. Finally, we thank
31 Niels Bormann for assisting with data collection, and the members of the qualitative research
32 working group at the IMS, who helped with coding, discussing and evaluating the data.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- 1 Nichols M, Townsend N, Scarborough P, et al. Cardiovascular disease in Europe: epidemiological update. *Eur Heart J* 2013;34:3028–34.
- 2 GBD. Mortality and causes of death collaborators. Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2013;2015(9963):117–71.
- 3 Benjamin EJ, Blaha MJ, Chiuve SE, et al. Heart disease and stroke Statistics-2017 update: A report From the American Heart Association. *Circulation* 2017;135:e146–e603.
- 4 Khaing W, Vallibhakara SA, Attia J, et al. Effects of education and income on cardiovascular outcomes: A systematic review and meta-analysis. *Eur J Prev Cardiol* 2017;24:1032–42.
- 5 Havranek EP, Mujahid MS, Barr DA, et al. Social determinants of risk and outcomes for cardiovascular disease: A scientific statement From the American Heart Association. *Circulation* 2015;132:873–98.
- 6 Mestral C de, Stringhini S. Socioeconomic status and cardiovascular disease; an update. *Curr Cardiol Rep* 2017;19:115.
- 7 Fihn SD, Gardin JM, Abrams J, et al. 2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease. *J Am Coll Cardiol* 2012;60(24):e44–e164.
- 8 Schröder SL, Richter M, Schröder J, et al. Socioeconomic inequalities in access to treatment for coronary heart disease: A systematic review. *Int J Cardiol* 2016;219:70–8.

- 1
2
3 9 Pedersen M, Egerod I, Overgaard D, et al. Social inequality in phase II cardiac
4 rehabilitation attendance: the impact of potential mediators. *Eur J Cardiovasc Nurs*
5 2018;17:345–55.
6
7
8
9
10 10 Verlinde E, Laender N de, Maesschalck S de, et al. The social gradient in doctor-patient
11 communication. *Int J Equity Health* 2012;11:12.
12
13
14 11 Ghisi GLM, Chaves GSDS, Britto RR, et al. Health literacy and coronary artery disease:
15 A systematic review. *Patient Educ Couns* 2018;101:177–84.
16
17
18
19 12 Sørensen K, Pelikan JM, Röthlin F, et al. Health literacy in Europe: comparative results
20 of the European health literacy survey (HLS-EU). *Eur J Public Health* 2015;25:1053–58.
21
22
23 13 Hetemaa T, Keskimäki I, Manderbacka K, et al. How did the recent increase in the
24 supply of coronary operations in Finland affect socioeconomic and gender equity in their
25 use? *J Epidemiol Commun Health* 2003;57:178–85.
26
27
28
29
30 14 Lemstra ME, Alsabbagh W, Rajakumar RJ, et al. Neighbourhood income and cardiac
31 rehabilitation access as determinants of nonattendance and noncompletion. *Can J Cardiol*
32 2013;29:1599–603.
33
34
35
36
37 15 Perelman J, Shmueli A, McDonald KM, et al. Inequality in treatment use among elderly
38 patients with acute myocardial infarction: USA, Belgium and Quebec. *BMC Health Serv*
39 *Res* 2009;9:130.
40
41
42
43
44 16 Manderbacka K. Exploring gender and socioeconomic differences in treatment of
45 coronary heart disease. *Eur J Public Health* 2005;15:634–39.
46
47
48
49 17 Pedersen M, Overgaard D, Andersen I, et al. Experience of exclusion: A framework
50 analysis of socioeconomic factors affecting cardiac rehabilitation participation among
51 patients with acute coronary syndrome. *Eur J Cardiovasc Nurs* 2017;16:715–23.
52
53
54
55
56
57
58
59
60

- 1
2
3 18 Schröder SL, Fink A, Schumann N, et al. How socioeconomic inequalities impact
4 pathways of care for coronary artery disease among elderly patients: study protocol for a
5 qualitative longitudinal study. *BMJ Open* 2015;5:e008060.
6
7
8
9
10 19 Helfferich C. *Die Qualität qualitativer Daten: Manual für die Durchführung qualitativer*
11 *interviews*. 3rd ed. Wiesbaden: VS Verlag für Sozialwissenschaften 2009.
12
13
14 20 Destatis. Demografische standards. 2017.
15
16 [https://www.destatis.de/DE/Methoden/DemografischeRegionaleStandards/Demografisch](https://www.destatis.de/DE/Methoden/DemografischeRegionaleStandards/DemografischeStandardsInfo.html)
17 [eStandardsInfo.html](https://www.destatis.de/DE/Methoden/DemografischeRegionaleStandards/DemografischeStandardsInfo.html) (accessed 16 Jul 2018).
18
19
20
21 21 Hoffmeyer-Zlotnik J. "Stellung im Beruf" als Ersatz für eine Berufsklassifikation zur
22 Ermittlung von sozialem Prestige. In: *ZUMA Nachrichten* 27 2003;53:114-127:114–27.
23
24
25 22 Schreier M. *Qualitative content analysis in practice*. Los Angeles. London, New Dehli,
26 Singapore, Washington DC: Sage 2012.
27
28
29
30 23 Zimmerman MS. Information Poverty and Reproductive Healthcare; Assessing the
31 Reasons for Inequity between Income Groups. *Soc Work Public Health* 2017;32:210–
32 221.
33
34
35
36
37 24 Lineback CM, Mervak CM, Revels S'sL, et al. Barriers to Accessing Optimal Esophageal
38 Cancer Care for Socioeconomically Disadvantaged Patients. *Ann Thorac Surg*
39 2017;103:416–421.
40
41
42
43
44 25 Czarny MJ, Nathan AS, Yeh RW, et al. Adherence to dual antiplatelet therapy after
45 coronary stenting: a systematic review. *Clin Cardiol* 2014;37:505–13.
46
47
48
49 26 Dankner R, Drory Y, Geulayov G, et al. A controlled intervention to increase
50 participation in cardiac rehabilitation. *Eur J Prev Cardiol* 2015;22:1121–28.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 27 Nolan MT, McKee G. Is knowledge level of coronary heart disease and risk factors
4 Among post-percutaneous coronary intervention patients adequate? *J Cardiovasc Nurs*
5 2016;31:E1-E9.
6
7
8
9
10 28 Nunes SRT, Rego G, Nunes R. Right or duty of information: A Habermasian perspective.
11 *Nurs Eth* 2016;23:36–47.
12
13
14 29 Kayaniyil S, Ardern CI, Winstanley J, et al. Degree and correlates of cardiac knowledge
15 and awareness among cardiac inpatients. *Patient Educ Couns* 2009;75:99–107.
16
17
18
19 30 Tchicaya A, Lorentz N, Demarest S, et al. Persistence of socioeconomic inequalities in
20 the knowledge of cardiovascular risk factors five years after coronary angiography. *Eur J*
21 *Cardiovasc Nurs* 2018;17(2):136–147.
22
23
24
25
26 31 Fennessy MM, Devon HA, Ryan C, et al. Changing illness perceptions and adherence to
27 dual antiplatelet therapy in patients with stable coronary disease. *J Cardiovasc Nurs*
28 2013;28:573–83.
29
30
31
32
33 32 Kucukarslan SN. A review of published studies of patients' illness perceptions and
34 medication adherence: lessons learned and future directions. *Res Social Adm Pharm*
35 2012;8:371–82.
36
37
38
39
40 33 Alter DA, Franklin B, Ko DT, et al. Socioeconomic status, functional recovery, and long-
41 term mortality among patients surviving acute myocardial infarction. *PLOS ONE*
42 2014;8:e65130.
43
44
45
46
47 34 Smith SK, Dixon A, Trevena L, et al. Exploring patient involvement in healthcare
48 decision making across different education and functional health literacy groups. *Soc Sci*
49 *Med* 2009;69:1805–12.
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 35 Pollard S, Bansback N, Bryan S. Physician attitudes toward shared decision making: A
4 systematic review. *Patient Educ Couns* 2015;98:1046-57.
5
6
7
8 36 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research
9 (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care*
10 2007;19:349–57.
11
12
13
14 37 Galobardes B, Shaw M, Lawlor DA, et al. Indicators of socioeconomic position (part 1).
15 *J Epidemiol Community Health* 2006;60(1):7–12.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 **Tables**
4
5
6
7
8
9

10 **Table 1.** Characteristics of the patients at baseline (n = 41).
11
12
13

Characteristics	Number in overall patients	Number in lower-SES patients	Number in higher-SES patients
Total	41	23	18
Age, the mean	69.7	69.7	69.6
Male	26	10	16
Female	15	13	2
Publicly insured	41	23	18
One-vessel disease	13	8	5
Two-vessel disease	11	6	5
Three-vessel disease	17	9	8
CHD diagnosed for up to 1 year	20	13	7
CHD diagnosed longer than 1 year ago	21	10	11

CHD, coronary heart disease; SES, socioeconomic status

1
2
3 **Figure legends**
4

5 **Fig 1.** Three major themes derived from socioeconomic differences in patients' experiences with
6 coronary heart disease treatment.
7
8
9

10
11
12 **Supporting information**
13

14 S1 Appendix. Interview guides.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig 1. Three major themes derived from socioeconomic differences in patients' experiences with coronary heart disease treatment.

190x254mm (300 x 300 DPI)

Interview Guide Baseline

Key question	Concrete supplementary questions
Please begin by telling me about the medical history of your heart disease starting from the first symptoms until this hospital stay.	<ul style="list-style-type: none"> - Have you ever been to a general practitioner/ cardiologist/ in a rehabilitation clinic before because of your heart disease? - How was your disease diagnosed? - Why are you currently in hospital?
Which positive and negative experiences have you made, concerning the care you have received?	<ul style="list-style-type: none"> - Can you think of any other positive or negative experience you have made with the hospital or office-based physicians? - Was there anything you were particularly satisfied with? - Was there anything you were unsatisfied with? - Have you experienced any problems with the further processing of the treatment? - How have you experienced talking with your physicians? - How was a decision reached about what treatment you were to receive?
How would you describe the quality of your treatment?	<ul style="list-style-type: none"> - How satisfied are you with your treatment? - Do you think that everyone in Germany is able to receive good treatment? Why do you think that is?
How do you manage your heart disease right now?	<ul style="list-style-type: none"> - What will happen next? - What medical care do you expect to receive in the next months? - What hopes do you have for your health in the future? - Is there something else you would like to tell me?

Questions aimed at maintaining the conversational flow:

- Please tell me exactly how things went with...
- Could you perhaps give me a few more details?
- What happened next? / And after that?
- What else comes into your mind?
- What do you associate with ...?

Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist

<i>No. Item</i>	<i>Guide questions/description</i>	<i>Page</i>
Domain 1: Research team and reflexivity		
<i>Personal Characteristics</i>		
1. Interviewer/facilitator	Which author/s conducted the interview or focus group?	9
2. Credentials	What were the researcher's credentials? E.g. PhD, MD	1/9
3. Occupation	What was their occupation at the time of the study?	1/9
4. Gender	Was the researcher male or female?	9
5. Experience and training	What experience or training did the researcher have?	9
<i>Relationship with participants</i>		
6. Relationship established	Was a relationship established prior to study commencement?	9-10
7. Participant knowledge of the interviewer	What did the participants know about the researcher? e.g. personal goals, reasons for doing the research	9-10
8. Interviewer characteristics	What characteristics were reported about the interviewer/facilitator? e.g. Bias, assumptions, reasons and interests in the research topic	9-10
Domain 2: study design		
<i>Theoretical framework</i>		
9. Methodological orientation and Theory	What methodological orientation was stated to underpin the study? e.g. grounded theory, discourse analysis, ethnography, phenomenology, content analysis	7
<i>Participant selection</i>		
10. Sampling	How were participants selected? e.g. purposive, convenience, consecutive, snowball	7
11. Method of approach	How were participants approached? e.g. face-to-face, telephone, mail, email	8
12. Sample size	How many participants were in the study?	8/11
13. Non-participation	How many people refused to participate or dropped out? Reasons?	8
<i>Setting</i>		
14. Setting of data collection	Where was the data collected? e.g. home, clinic, workplace	9
15. Presence of non-participants	Was anyone else present besides the participants and researchers?	10
16. Description of sample	What are the important characteristics of the sample? e.g. demographic data, date	Table 1
<i>Data collection</i>		
17. Interview guide	Were questions, prompts, guides provided by the authors? Was it pilot tested?	9/ S1 Appendix
18. Repeat interviews	Were repeat interviews carried out? If yes, how many?	8
19. Audio/visual recording	Did the research use audio or visual recording to collect the data?	10
20. Field notes	Were field notes made during and/or after the interview or focus group?	9
21. Duration	What was the duration of the interviews or focus group?	10

<i>No. Item</i>	<i>Guide questions/description</i>	<i>Page</i>
22. Data saturation	Was data saturation discussed?	8
23. Transcripts returned	Were transcripts returned to participants for comment and/or correction?	11
Domain 3: analysis and findings		
<i>Data analysis</i>		
24. Number of data coders	How many data coders coded the data?	10
25. Description of the coding tree	Did authors provide a description of the coding tree?	No
26. Derivation of themes	Were themes identified in advance or derived from the data?	10
27. Software	What software, if applicable, was used to manage the data?	11
28. Participant checking	Did participants provide feedback on the findings?	11
<i>Reporting</i>		
29. Quotations presented	Were participant quotations presented to illustrate the themes / findings? Was each quotation identified? e.g. participant number	13-18
30. Data and findings consistent	Was there consistency between the data presented and the findings?	yes
31. Clarity of major themes	Were major themes clearly presented in the findings?	17-19/ Figure 1
32. Clarity of minor themes	Is there a description of diverse cases or discussion of minor themes?	11-17