

Table 3 supplementary. Prevalences of chronic pain and of other chronic diseases by sex, age and disabling condition.

Variables		Condition		Disabling Condition (Subpopulation with the condition/s)		Disabling Condition (Total population)		
		Prevalence	95% Confidence Interval	Prevalence ^b	95% Confidence Interval	Prevalence ^c	95% Confidence Interval	
Chronic Pain ^a	Total	17.2%	16.3-18.12	66.9%	64.14-69.55	11.5%	10.73-12.28	
	Sex (p<0.001)	Women	23.9%	22.5-25.4	68.2%	64.9-71.3	16.3%	15.1-17.6
		Men	10.3%	9.3-11.4	63.9%	58.6-68.8	6.6%	5.8-7.5
	Age groups (p<0.001)	16-44 yrs	11.4%	10.4-12.5	48.2%	43.4-53.1	5.5%	4.8-6.3
		45-64	18.2%	16.5-20	70.4%	65.3-75	12.8%	11.4-14.4
+65		32.3%	29.7-35	82.7%	78.6-86.1	26.7%	24.2-29.2	
At least one Chronic Disease (besides Chronic Pain)	Total	45.9%	44.67-47.08	59.7%	58-61.47	29.5%	28.45-30.66	
	Sex (p<0.001)	Women	52.9%	51.2-54.6	64.5%	62.2-66.7	36.8%	35.2-38.5
		Men	38.6%	37-40.3	53.1%	50.3-55.8	22.1%	20.7-23.5
	Age groups (p<0.001)	16-44 yrs	23.7%	22.3-25.1	43.3%	40-46.7	13%	11.9-14.2
		45-64	56.9%	54.6-59.1	56.6%	53.6-59.5	34%	31.9-36.2
+65		92.1%	90.4-93.5	74.9%	72.3-77.4	69.7%	67.1-72.3	

^a Chronic Pains: 'migraine/headache/chronic cephalalgia /frequent headache', 'angina/chest pain', 'back pain, neck pain, shoulder pain, waist pain, cervical/low back pain' or 'menstrual pain'.

^b Percentage of disabling population within that population with the corresponding condition (chronic pain or at least one chronic disease).

^c Percentage of population with both disabling and the corresponding condition (chronic pain or at least one chronic disease).