

Evaluation of the adequacy of drug prescriptions in patients with chronic kidney disease: results from the CKD-REIN cohort


S.M. Laville, M. Metzger, B. Stengel, C. Jacquelinet, C. Combe, D. Fouque, M. Laville, L. Frimat, C. Ayav, E. Speyer, B.M. Robinson, Z.A. Massy, S. Liabeuf

British Journal of Clinical Pharmacology

Corresponding author:



Bénédicte Stengel
Director, Inserm U1018, Team 5
UPS-UVSQ, CESP, Centre for Epidemiology and Population Health
EpReC Team, Renal and Cardiovascular Epidemiology
16, avenue P. Vaillant Couturier
94 807 Villejuif cedex
France
Phone: + 33 1 4559 5039
E-mail: benedicte.stengel@inserm.fr

a. Number of inappropriate prescriptions per patient (n=3011)



b. Number of inappropriately high dose prescriptions per patient (n=3011)

c. Number of contraindicated prescriptions per patient (n=3011)



Legend

- No drug
- One drug
- Two drugs
- Three drugs or more

Supplementary Figure 1: Appropriateness of prescriptions, expressed as the number of inappropriate (i.e. contraindicated or inappropriately high dose drug), inappropriately high dose or contraindicated drugs per patient (n=3011 patients receiving at least one medication). a. Number of inappropriate prescriptions per patient. b. Number of inappropriately high dose prescriptions per patient. c. Number of contraindicated prescriptions per patient.