

Online Data Supplement for:

**Predicting 30-day mortality in patients with Sepsis; an exploratory analysis of process of care and patient characteristics**

Miriam Sanderson<sup>1</sup> [mzymis@exmail.nottingham.ac.uk](mailto:mzymis@exmail.nottingham.ac.uk)

Dr Marc Chikhani<sup>2,3</sup> [marc.chikhani@nottingham.ac.uk](mailto:marc.chikhani@nottingham.ac.uk)

Esme Blyth<sup>4</sup> [eblyth1@sheffield.ac.uk](mailto:eblyth1@sheffield.ac.uk)

Sally Wood<sup>2</sup> [sally.wood@nuh.nhs.uk](mailto:sally.wood@nuh.nhs.uk)

Prof Iain K Moppett<sup>2,3</sup> [iain.moppett@nottingham.ac.uk](mailto:iain.moppett@nottingham.ac.uk)

Dr Tricia McKeever<sup>5</sup> [tricia.mckeever@nottingham.ac.uk](mailto:tricia.mckeever@nottingham.ac.uk)

Dr Mark J R Simmonds<sup>2</sup> [mark.simmonds@nuh.nhs.uk](mailto:mark.simmonds@nuh.nhs.uk)

1. School of Medicine, University of Nottingham

2. Nottingham University Hospitals NHS Trust

3. Division of Clinical Neuroscience, Faculty of Medicine and Health Sciences,  
University of Nottingham

4. Sheffield Teaching Hospitals NHS Foundation Trust

5. Epidemiology and Public Health, Faculty of Medicine and Health Sciences,  
University of Nottingham

All correspondence to: [Marc.chikhani@nottingham.ac.uk](mailto:Marc.chikhani@nottingham.ac.uk)

---

**Table E1 Age and sex of patients with severe sepsis and univariate association with 30 day mortality**

Variable	Dead (N=119) % (n)	Alive (N=336) % (n)	OR <sup>a</sup>	95%CI <sup>b</sup>	P-value
Age (years) mean (SD <sup>c</sup> )	12.1 (70.3)	17.4 (61.8)			<0.001
<b>Sex</b>					
Female N(%)	25.9 (50)	74.1 (143)	1.00		
Male N(%)	26.3 (69)	73.7 (193)	1.02	0.67-1.56	0.92

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval

<sup>c</sup> Standard Deviation

**Table E2 Symptoms of systemic inflammatory response syndrome and univariate association with 30 day mortality**

SIRS* Criteria	Dead (N=119) % (n)	Alive (N=336) % (n)	OR <sup>a</sup>	95%CI <sup>b</sup>	P-value
Fever - Core temperature >38.3°C	17.8 (48)	82.2 (221)	0.35	0.23-0.55	<0.001
Hypothermia - Core temperature <36°C	51.5 (24)	48.9 (23)	3.44	1.83-6.45	<0.001
Heart Rate >90/min	26.6 (102)	73.4 (282)	1.15	0.64-2.07	0.65
Respiratory Rate >25/min	27.1 (71)	72.8 (190))	1.14	0.74-1.74	0.56
Altered mental status	36.8 (32)	63.2 (55)	1.88	1.14-3.10	0.01
Oedema	30.8 (4)	69.2 (9)	1.26	0.38-4.19	0.75
Blood Glucose >6.6mmol/L	31 (22)	69 (49)	1.33	0.76-2.31	0.31
Leukocytosis - White blood cells>12x10 <sup>9</sup> /L	26.2 (66)	173.8 (86)	1.00	0.66-1.53	0.98
Leukopenia - White blood cells<4x10 <sup>9</sup> /L	29.5 (18)	70.5 (43)	1.21	0.67-2.20	0.52
C-Reactive Protein >30mg/L	26.5 (81)	73.5 (225)	1.05	0.67-1.65	0.83

\*Systemic inflammatory response syndrome

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval

**Table E3 Symptoms of organ dysfunction or tissue hypoperfusion and univariate association with 30 day mortality**

<b>Criteria for organ dysfunction or tissue hypoperfusion</b>	<b>Dead (N=119) % (n)</b>	<b>Alive (N=336) % (n)</b>	<b>OR<sup>a</sup></b>	<b>95%CI<sup>b</sup></b>	<b>p-value</b>
Saturations (<92% room air) or (PaO <sub>2</sub> /FiO <sub>2</sub> <sup>c</sup> <40)	27.1 (62)	72.9 (167)	1.10	0.72-1.67	0.65
Urine Output <30ml per hour for 2 hours	26.7 (23)	73.3 (63)	1.04	0.61-1.77	0.89
Creatinine >175µmol/L or increased >45µmol/L	29.4 (50)	70.6 (120)	1.30	0.85-2.00	0.22
Coagulation Abnormalities <sup>d</sup>	50 (7)	50 (7)	2.94	1.00-8.61	0.04
Thrombocytopenia (<100x10 <sup>9</sup> /L)	48.3 (14)	52.7 (15)	2.85	1.32-6.15	0.01
Hyperbilirubinaemia (>35mmol/L)	35.0 (14)	65.0 (26)	1.59	0.80-3.16	0.18
Ileus	25.0 (1)	75.0 (3)	0.94	0.10-9.13	1.00
Hyperlactaemia (>2mmol/L)	29.0 (83)	71.0 (203)	1.51	0.96-2.37	0.07
Capillary refill >5 seconds	37.5 (6)	62.5 (10)	1.73	0.61-4.88	0.29
Mottling	60.0 (9)	40.0 (6)	4.50	1.55-13.08	<0.001
Arterial Hypotension <sup>e</sup>	25.5 (76)	74.5 (222)	0.91	0.59-1.41	0.66

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval

<sup>c</sup> Partial Pressure of Oxygen in arterial blood/Fraction of inspired oxygen

<sup>d</sup> International Normalised Ratio>1.5 or Activated Pro-Thrombin Time >60seconds

<sup>e</sup> Systolic blood pressure <90mmHg, mean arterial pressure <70mmHg or Systolic blood pressure decrease >40mmHg

**Table E4 Locations of Early Care and univariate association with 30 day mortality**

<b>Admissions and Early Treatment locations</b>	<b>Dead (N=119) %(n)</b>	<b>Alive (N=336) %(n)</b>	<b>OR<sup>a</sup></b>	<b>95%CI<sup>b</sup></b>	<b>P-value</b>
<b>Admission Source</b>					
Emergency Department	24.6 (71)	75.4 (218)	1.00		
Elective	25 (5)	75 (15)	1.02	0.36-2.92	
General Practitioner	29.5 (43)	70.6 (103)	1.28	0.82-2.00	0.28
<b>GP referral?</b>					
Yes	29.5 (43)	70.6 (103)	1.27	0.82-1.99	0.27

No	24.6 (76)	75.4 (233)	1.00		
<b>Speciality at Time Zero</b>					
Medical	28.0 (106)	72.0 (273)	1.00		
Surgical	18.2 (12)	81.8 (54)	0.57	0.29-1.11	0.124
Other	10.0 (1)	90.0 (9)	0.29	0.04-2.29	
<b>Surgical ward at Time Zero?</b>					
Yes	18.2 (12)	81.8 (54)	0.59	0.30-1.14	0.111
No	27.5 (107)	72.5 (282)	1.00		

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval

**Table E5 Sepsis source and control and univariate association with 30 day mortality**

	<b>Dead (N=119) %(n)</b>	<b>Alive (N=336) %(n)</b>	<b>OR<sup>a</sup></b>	<b>95%CI<sup>b</sup></b>	<b>P- value</b>
<b>Provisional Sepsis Source at 6Hrs<sup>c</sup></b>					
Identified N(%)	25.4 (107)	74.6 (315)	0.59	0.28-1.25	0.17
Not identified N(%)	36.4 (12)	63.7 (21)	1.00		
<b>Location</b>					
Lungs N(%)	28.8 (66)	71.2 (163)	1.32	0.87-2.01	0.19
Liver N(%)	30.2 (13)	69.8 (30)	1.25	0.63-2.49	0.52
Unperforated bowel N(%)	0 (0)	100 (4)			0.58
Peritoneum N(%)	28.6 (4)	71.4 (10)	1.13	0.35-3.69	0.77
Urinary tract N(%)	20.0 (14)	80.0 (56)	0.67	0.36-1.25	0.20
Intracranial N(%)	33.3 (1)	66.8 (2)	1.42	0.13-15.8	1.00
Heart N(%)	33.3 (1)	66.7 (2)	1.42	0.13-15.8	1.00
Skin N(%)	11.4 (4)	88.6 (31)	0.34	0.12-0.99	0.04
Bone N(%)	22.2 (2)	78.8 (7)	0.80	0.16-3.92	1.00
Ear nose throat N(%)	0 (0)	100 (2)			1.00
Gynaecological N(%)	16.7 (1)	83.3 (5)	0.56	0.06-4.85	1.00
Line N(%)	0 (0)	100 (3)			0.57
<b>Imaging</b>					
Appropriate Imaging N(%)	25.5 (109)	74.5 (319)	0.58	0.26-1.31	0.19
No appropriate Imaging N(%)	37.0 (10)	63.0 (17)	1.00		
<b>Time to imaging - median (IQR*)<sup>d</sup></b>	0.92(0.07-2.78)	0.60 (0.00-2.50)			0.59
<b>Source Control</b>					
<b>Source control appropriate immediately?</b>					
Yes N(%)	18.9 (10)	81.1 (43)	0.63	0.30-1.29	0.20
No N(%)	72.9 (293)	27.1 (109)	1.00		
<b>When appropriate, source control within 6 Hours?</b>					
Yes N(%)	22.2 (6)	77.8 (21)	1.57	0.39-6.37	0.44
No N(%)	15.4 (4)	84.6 (22)	0.62	0.23-1.69	

\*Interquartile Range

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval

<sup>c</sup> 3 patients with missing data

<sup>d</sup> 4 patients with missing data

**Table E6 Patient escalation and univariate association with 30 day mortality**

<b>Patient Escalation</b>	<b>Dead (N=119) %(n)</b>	<b>Alive (N=336) %(n)</b>	<b>OR<sup>a</sup></b>	<b>95%CI<sup>b</sup></b>	<b>P- value</b>
<b>Early Warning Score<sup>c</sup></b>					
0-2 "stable" and 3 "At risk" N(%)	29.6 (8)	70.4 (19)	1.00		
4-5 "Doctor Review" N(%)	28.7 (27)	71.3 (67)	0.96	0.37-2.46	
6+ "Critical Care Outreach Team aware" N(%)	25.4 (84)	74.6 (247)	0.81	0.34-1.92	0.46
<b>Appropriate escalation?</b>					
Yes N(%)	74.4 (325)	25.6 (112)	0.40	0.13-1.23	0.12
No N(%)	46.2 (6)	53.9 (7)	1.00		
N/A N(%)	20.0 (1)	80.0 (4)	0.29	0.02-3.91	
<b>Time to escalation - median (IQR*)</b>	0.0(-0.12- 0.0)	0.0 (-0.02-0.0)			0.96
<b>Critical Care Outreach Team called within 1 Hour?<sup>d</sup></b>					
Yes N(%)	30.4 (21)	69.6 (48)	1.23	0.69-2.19	0.62
No N(%)	26.2 (76)	73.8 (214)	1.00		
Out of Hours	23.7 (22)	76.3 (71)	0.87	0.51-1.51	
<b>Time to Doctor - median (IQR*)</b>	0.17(0.00- 0.55)	0.17 (0.00- 0.50)			0.99
<b>Grade of doctor</b>					
F1 N(%)	12.2 (5)	87.8 (36)	0.49	0.15-1.58	0.15
F2/ST1/ST2 N(%)	28.7 (55)	71.4 (137)	1.42	0.68-2.99	
ST3+/SPR N(%)	27.9 (48)	72.1 (124)	1.37	0.65-2.91	
Consultant N(%)	22.0 (11)	78.0 (39)	1.00		
<b>Critical Care Outreach Team review after Initial diagnosis?</b>					
Yes N(%)	25.6 (33)	74.4 (96)	0.96	0.60-1.53	0.86
No N(%)	26.4 (86)	73.6 (240)	1.00		
<b>Time to Critical Care Outreach Team - median (IQR)</b>	4.08(1.20- 11.00)	5.08(1.76- 0.17)			0.53
<b>Time to Senior Doctor - median (IQR*)<sup>e</sup></b>	0.75(1.30- 2.15)	0.75 (0.13- 2.07)			0.84
<b>No senior doctor seen N(%)</b>	0 (0)	100(1)			

<b>Considered for Intensive Therapy Unit (ITU)?</b>					
Yes N(%)	26.2 (119)	73.8 (335)			
No N(%)	100 (1)	0 (0)			1.00
<b>Time to Intensive Therapy Unit assessment - median (IQR*)<sup>f</sup></b>	3.67(2.00-8.00)	3.56 (2.13-7.39)			0.98
<b>Time to Intensive Therapy Unit Admission - median (IQR*)</b>	7.00(4.05-13.25)	6.80 (4.00-11.13)			0.65
<b>Time to Intensive Therapy Unit Assessment Level 3 - median (IQR*)<sup>g</sup></b>	17.93(6.00-20.91)	15.88(12.40-31.05)			0.57
<b>Time to Intensive Therapy Unit Level 3 - median (IQR*)</b>	21.33(6.00-29.33)	17.83(12.75-31.48)			0.83

\*Interquartile Range

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval

<sup>c</sup> 3 patients with missing data

<sup>d</sup> 3 patients with missing data

<sup>e</sup> 1 patient with missing data

<sup>f</sup> 5 patients with missing data

<sup>g</sup> 5 patients with missing data

**Table E7 Blood cultures and antibiotics and univariate association with 30 day mortality**

<b>Variable</b>	<b>Dead (N=119) % (n)</b>	<b>Alive (N=336) % (n)</b>	<b>OR<sup>a</sup></b>	<b>95%CI<sup>b</sup></b>	<b>P-value</b>
<b>Blood cultures within 6 Hours of time zero?</b>					
1 set N(%)	25.7 (70)	74.3 (202)	0.81	0.50-1.31	0.19
2+ set N(%)	21.2 (14)	78.8 (52)	0.63	0.31-1.29	
No N(%)	29.9 (35)	70.1 (82)	1.00		
<b>Blood culture before Time Zero (if no to previous)</b>					
Yes N(%)	30.1 (22)	69.9 (51)	1.03	0.45-2.34	0.33
No N(%)	29.6 (13)	70.5 (31)	1.00		
<b>Time to Blood Culture - median</b>					



<b>(IQR*)</b>					
Before Time Zero	26.5 (36)	73.5 (100)	1.00		0.73
0-4 Hours	24.6 (58)	75.4 (178)	0.91	0.56-1.47	
4+ Hours	31.8 (14)	68.2 (30)	1.30	0.62-2.73	
<b>Blood culture before antibiotics?</b>					
Yes N(%)	25.0 (61)	75.0 (183)	0.88	1.58-1.34	0.55
No N(%)	27.5 (58)	72.5 (153)	1.00		
<b>Antibiotic after Time Zero</b>					
Yes N(%)	26.2 (119)	73.9 (336)			
<b>Antibiotic adhere local guidelines</b>					
Yes N(%)	26.3 (119)	73.7 (336)			1.00
No N(%)	0 (0)	100 (2)			
<b>Antibiotics before Time Zero</b>					
Yes N(%)	31.0 (27)	69.0 (60)	1.35	0.81-2.26	0.25
No N(%)	25.0 (92)	75.0 (276)	1.00		
<b>Time to antibiotic - median (IQR*)</b>	0.56 (0-1.4)	0.5 (0.07-1.0)			0.69

\*Interquartile Range

<sup>a</sup>Odds Ratio

<sup>b</sup> 95% Confidence Interval

**Table E8 Lactate, the hypotensive patient, and univariate association with 30 day mortality**

<b>Variable</b>	<b>Dead (N=119) %(n)</b>	<b>Alive (N=336) %(n)</b>	<b>OR<sup>1</sup></b>	<b>95%CI<sup>2</sup></b>	<b>P- value</b>
<b>Lactate Measured?</b>					
Yes N(%)	25.7 (114)	74.3 (330)	0.41	0.12-1.39	0.14
No N(%)	45.5 (5)	54.6 (6)	1.00		
<b>Lactate Value - median (IQR*)</b>	3.30(2.20- 5.40)	2.55 (1.50- 4.10)			<0.001
<b>Time to lactate - median (IQR*)<sup>3</sup></b>	0.25 (0- 0.93)	0.38 (0.0- 1.28)			0.16
<b>Hypotension or hyperlactate?</b>					
Yes N(%)	27.2 (98)	72.9 (263)	1.30	0.76-2.22	0.35
No N(%)	22.3 (21)	77.7 (73)	1.00		
<b>Crystalloid or colloid?</b>					
Yes N(%)	27.1 (95)	72.9 (255)	0.99	0.26-3.83	0.65
No N(%)	27.3 (3)	72.7 (8)	1.00		
N/A N(%)	22.3 (21)	77.7 (73)	0.77	0.19-3.15	
<b>Time to fluid - median (IQR*)</b>	0.50(0.17- 1.17)	0.50(0.22- 1.03)			0.60
<b>Fail fluid challenge<sup>a</sup> or new renal impairment?<sup>a</sup></b>					
Yes N(%)	26.9 (93)	73.1 (253)	1.17	0.71-1.94	0.53
No N(%)	23.9 (26)	76.2 (83)	1.00		
<b>Urine Catheter?</b>					
Yes N(%)	27.0 (90)	73.0 (243)	1.23	0.33-4.59	0.82
No N(%)	23.1 (3)	76.9 (10)	1.00		
N/A N(%)	23.9 (26)	76.2 (83)	1.04	0.27-4.08	
<b>Fluid balance chart completed?</b>					
Yes N(%)	27.1 (88)	72.9 (237)	1.19	0.42-3.35	0.67
No N(%)	23.8 (5)	76.2 (16)	1.00		
N/A N(%)	23.9 (26)	76.2 (83)	1.00	0.33-3.01	

<sup>a</sup> Persistent systolic blood pressure <90mmHg or mean arterial pressure <70mmHg despite fluid

resuscitation

\*Interquartile Range

<sup>1</sup> Odds Ratio

<sup>2</sup> 95% Confidence Interval

<sup>3</sup>2 patients with missing data

**Table E9 Interventions for a patient with septic shock, and univariate association with 30 day mortality**

<b>Variable</b>	<b>Dead (N=119) %(n)</b>	<b>Alive (N=336) %(n)</b>	<b>OR<sup>1</sup></b>	<b>95%CI<sup>2</sup></b>	<b>P-value</b>
<b>Fluid refractory<sup>a</sup> sepsis?</b>					
Yes N(%)	23.9 (59)	76.1 (188)	0.60	0.36-1.00	0.09
No N(%)	34.3 (34)	65.7 (65)	1.00		
N/A N(%)	23.9 (26)	76.1 (83)	0.60	0.33-1.1	
<b>Central Venous Catheter placed?</b>					
Yes N(%)	27.0 (90)	73.0 (243)	1.05	0.28-3.94	0.63
No N(%)	23.1 (3)	76.9 (10)	1.00		
N/A N(%)	23.9 (26)	76.2 (83)	1.35	0.36-5.08	
<b>Time to Central Venous Catheter - median (IQR*)</b>	6.91(3.58-12.4)	5.95 (3.32-12.5)			0.64
<b>Vasopressors?<sup>3</sup></b>					
Yes N(%)	21.1 (54)	72.9 (145)	2.89	0.97-8.56	0.19
No N(%)	11.4 (4)	88.6 (31)	1.00		
N/A N(%)	28.2 (58)	71.8 (148)	3.04	1.03-8.99	
<b>Time to Vasopressors - median (IQR*)</b>	7.75(4.65-15.55)	7.33(4.25-13.75)			0.75
<b>Remain hypotensive<sup>a?4</sup></b>					
Yes N(%)	54.6 (12)	45.5 (10)	3.80	1.53-9.40	0.01
No N(%)	24.0 (43)	76.0 (136)	1.00		
N/A N(%)	26.4 (63)	73.6 (176)	1.13	0.72-1.77	
<b>Time to inotropes - median (IQR*)</b>	9.25(4.00-19.35)	7.75 (5.67-14.55)			0.62
<b>If low Haemoglobin (&lt;7.0g/dl), blood transfused?</b>					
Yes N(%)	44.4 (4)	55.6 (5)	2.30	0.61-8.76	0.25
N/A N(%)	25.8 (115)	74.2 (331)	1.00		

<sup>a</sup> Persistent systolic blood pressure <90mmHg or mean arterial pressure <70mmHg despite fluid resuscitation

\*Interquartile Range

<sup>1</sup> Odds Ratio

<sup>2</sup> 95% Confidence Interval

<sup>3</sup> 15 patients with missing data

<sup>4</sup> 15 patients with missing data

**Table E10 Interventions within allotted time-frame and univariate association with 30 day mortality**

<b>Variable</b>	<b>Dead (N=119) %(n)</b>	<b>Alive (N=336) %(n)</b>	<b>OR<sup>a</sup></b>	<b>95%CI<sup>b</sup></b>	<b>P- value</b>
<b>Lactate measured within 6 Hours?</b>					
Yes N(%)	25.7 (110)	74.3 (318)	0.69	0.30-1.59	0.38
No N(%)	33.3 (9)	66.7 (18)	1.00		
<b>Blood Culture within 6 Hours?</b>					
Yes N(%)	25.6 (105)	74.5 (306)	0.74	0.38-1.44	0.37
No N(%)	31.8 (14)	68.2 (30)	1.00		
<b>Antibiotics within 1 Hour?</b>					
Yes N(%)	24.5 (84)	75.5 (259)	0.71	0.45-1.14	0.16
No N(%)	31.3 (35)	68.8 (77)	1.00		
<b>Fluid resuscitation within 6 Hours?</b>					
Yes N(%)	26.6 (93)	73.4 (257)	0.63	0.18-2.22	0.58
No N(%)	36.4 (4)	26.6 (7)	1.00		
N/A N(%)	23.4 (22)	76.6 (72)	0.53	0.14-2.00	
<b>Urine Catheter within 6 Hours?<sup>c</sup></b>					
Yes N(%)	27.2 (89)	72.8 (238)	1.50	0.41-5.43	0.88
No N(%)	20.0 (3)	80.0 (12)	1.00		
N/A N(%)	23.9 (26)	76.2 (86)	1.25	0.33-4.78	
Missing N(%)	25.0 (1)	75.0 (3)	1.33	0.10-19.20	
<b>Fluid Balance chart completed within 6 Hours?<sup>d</sup></b>					
Yes N(%)	26.6 (87)	73.4 (240)	1.27	0.41-3.96	0.91
No N(%)	22.2 (4)	77.8 (14)	1.00		
N/A N(%)	25.0 (26)	75.0 (78)	1.17	0.35-3.86	
<b>Central Venous Catheter within 6 Hours?</b>					
Yes N(%)	22.7 (29)	77.3 (99)	0.77	0.43-1.39	0.88
No N(%)	27.0 (30)	73.0 (81)			
N/A N(%)	27.8 (60)	72.2 (156)	1.01		
<b>Vasopressors within 6 Hours?</b>					
Yes N(%)	23.4 (22)	76.6 (72)	0.83	0.44-1.54	0.95
No N(%)	27.0 (34)	73.0 (92)			
N/A N(%)	26.8 (63)	73.2 (172)	0.99	0.61-1.62	

<b>Inotropes within 6 Hours?<sup>e</sup></b>					
Yes N(%)	41.7 (5)	58.3 (7)	0.79	0.18-3.37	0.035
No N(%)	47.6 (10)	52.4 (11)	1.00		
N/A N(%)	24.9 (104)	75.1 (313)	0.36	0.15-0.89	
<b>Blood Transfusion within 6 Hours?<sup>f</sup></b>					
Yes N(%)	33.3 (2)	66.7 (4)	1.00	0.09-11.02	0.29
No N(%)	33.3 (2)	66.7 (4)	1.00		
N/A N(%)	26.6 (114)	73.4 (315)	0.72	0.13-4.00	

\*Interquartile Range

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval

<sup>c</sup> 4 patients missing data

<sup>d</sup> 6 patients missing data

<sup>e</sup> 5 patients missing data

<sup>f</sup> 14 patients missing data

**Table E11 Hospital-acquired sepsis and resource issues, and univariate association with 30 day mortality**

<b>Variable</b>	<b>Dead (N=119) %(n)</b>	<b>Alive (N=336) %(n)</b>	<b>OR<sup>a</sup></b>	<b>95%CI<sup>b</sup></b>	<b>P- value</b>
<b>Hospital-acquired sepsis?</b>					
Yes N(%)	35.8 (34)	64.2 (61)	1.80	1.11-2.94	0.016
No N(%)	23.6 (85)	76.4 (275)			
<b>Resource issues?</b>					
Yes N(%)	23.6 (17)	76.4 (55)	0.85	0.47-1.53	0.59
No N(%)	26.6 (102)	73.4 (281)	1.00		

<sup>a</sup> Odds Ratio

<sup>b</sup> 95% Confidence Interval