

Year	Primary prevention		Secondary prevention	
	Crude n=688,813	Adjusted* n=219,254	Crude n=347,025	Adjusted* n=114,325
	Mean (95% CI)	Mean (95% CI)	Mean (95% CI)	Mean (95% CI)
2007	35.1 (34.7–35.6)	42.2 (40.7–43.7)	74.2 (73.6–74.7)	65.6 (63.5–67.6)
2008	39.5 (39.1–39.9)	46.3 (44.8–47.7)	76.4 (76.0–76.9)	69.4 (67.5–71.2)
2009	43.4 (43.1–43.8)	49.8 (48.4–51.3)	78.1 (77.7–78.5)	72.0 (70.3–73.7)
2010	45.6 (45.2–45.9)	52.4 (51.0–53.9)	78.4 (78.0–78.8)	73.8 (72.1–75.5)
2011	47.8 (47.5–48.1)	52.7 (51.3–54.1)	78.9 (78.5–79.2)	74.4 (72.7–76.1)
2012	48.7 (48.4–49.0)	52.7 (51.3–54.2)	77.8 n(77.4–78.2)	72.9 (71.1–74.6)
2013	49.2 (48.9–49.6)	52.7 (51.2–54.2)	76.6 (76.2–77.0)	71.6 (69.7–73.4)
2014	50.2 (49.9–50.6)	52.1 (50.6–53.6)	76.0 (75.6–76.4)	71.1 (69.2–72.9)

* Adjusted for year, county council, type of care, sex, age, HbA1c, eGFR, diabetes duration, diabetes medications, antihypertensives, antiplatelets, blood pressure, microalbuminuria, macroalbuminuria, BMI, physical activity, smoking and cholesterol levels.