

	Primary prevention		Secondary prevention	
	Crude n=751,864	Adjusted* n=237,751	Crude n=437,618	Adjusted* n=140,513
Year	Mean (95% CI)	Mean (95% CI)	Mean (95% CI)	Mean (95% CI)
2007	34.6 (34.2–35.0)	36.1 (34.7–37.5)	70.7 (70.2–71.2)	58.7 (56.8–60.6)
2008	38.9 (38.5–39.3)	40.2 (38.9–41.6)	72.8 (72.4–73.3)	63.2 (61.4–64.9)
2009	42.7 (42.3–43.0)	43.7 (42.4–45.1)	74.7 (74.3–75.1)	66.3 (64.6–68.0)
2010	44.7 (44.4–45.0)	46.3 (44.9–47.7)	74.8 (74.5–75.1)	68.0 (66.4–69.7)
2011	46.7 (46.4–47.0)	46.6 (45.2–48.0)	75.2 (74.9–75.5)	69.3 (67.7–70.9)
2012	47.4 (47.1–47.7)	46.6 (45.2–48.0)	73.7 (73.3–74.0)	68.1 (66.4–69.7)
2013	47.6 (47.3–47.9)	46.6 (45.2–48.0)	72.0 (71.6–72.3)	67.1 (65.4–68.8)
2014	48.3 (48.0–48.7)	46.1 (44.7–47.5)	71.1 (70.7–71.5)	66.1 (64.4–67.8)

\* Adjusted for year, county council, type of care, sex, age, HbA1c, eGFR, diabetes duration, diabetes medications, antihypertensives, antiplatelets, blood pressure, microalbuminuria, macroalbuminuria, BMI, physical activity, smoking and cholesterol levels.