

Supplementary Data

Synthesis of Disaccharides Containing 6-deoxy- α -L-Talose as Potential Heparan Sulfate Mimetics

Jon K. Fairweather¹, **Ligong Liu**^{1,2}, **Tomislav Karoli**^{1,2} and **Vito Ferro**^{1,3*}

¹ Drug Design Group, Progen Pharmaceuticals Ltd, Brisbane QLD 4076, Australia



² Current address: The University of Queensland, Institute for Molecular Bioscience, Brisbane, QLD 4072, Australia



³ Current address: The University of Queensland, School of Chemistry and Molecular Biosciences, Brisbane, QLD 4072, Australia


* Author to whom correspondence should be addressed; E-Mail: v.ferro@uq.edu.au; Tel.: +61-7-3346-9598; Fax: +61-7-3365-4299.

Contents


¹ H NMR (400 MHz) spectrum for compound 12	S3
¹ H NMR (400 MHz) spectrum for compound 20	S4
¹³ C NMR (100 MHz) spectrum for compound 20	S5
¹ H NMR (400 MHz) spectrum for compound 14	S6
¹ H NMR (400 MHz) spectrum for compound 22	S7
¹³ C NMR (100 MHz) spectrum for compound 22	S8
¹ H NMR (200 MHz) spectrum for compound 23	S9
¹ H NMR (400 MHz) spectrum for compound 25	S10
¹ H NMR (400 MHz) spectrum for compound 27	S11
¹ H NMR (400 MHz) spectrum for compound 26	S12
¹ H NMR (400 MHz) spectrum for compound 30	S13
¹³ C NMR (100 MHz) spectrum for compound 30	S14
¹ H NMR (400 MHz) spectrum for compound 31	S15
¹ H NMR (400 MHz) spectrum for compound 32	S16











20








14















S9





S10



S11





S13

