

Supplemental Information

**miR-30 Family Reduction Maintains Self-Renewal
and Promotes Tumorigenesis in NSCLC-Initiating
Cells by Targeting Oncogene TM4SF1**

Yu-Shui Ma, Fei Yu, Xiao-Ming Zhong, Gai-Xia Lu, Xian-Ling Cong, Shao-Bo Xue, Wen-Ting Xie, Li-Kun Hou, Li-Juan Pang, Wei Wu, Wei Zhang, Le-Le Cong, Tie Liu, Hui-Deng Long, Ran Sun, Hong-Yan Sun, Zhong-Wei Lv, Chun-Yan Wu, and Da Fu

Supplementary Figure and Legends




Fig. S1. Clinical significance of MiR-30c/a and TM4SF1 in NSCLC. (A)

Kaplan-Meier survival analysis to evaluate the OS and PFS of miR-30c expression in NSCLC patients. (B) Kaplan-Meier survival analysis to evaluate the OS and PFS of miR-30a expression in NSCLC patients. (C) Kaplan-Meier survival analysis to evaluate the OS and PFS of TM4SF1 expression in NSCLC patients. (D) Kaplan-Meier survival analysis to evaluate the OS and PFS of miR-30c/a combined with TM4SF1 expression in NSCLC.