

New Phytologist Supporting Information

Article title: **Arbuscular mycorrhizal fungi reduce nitrous oxide emissions from N₂O hotspots**

Authors: Kate Storer, Aisha Coggan, Phil Ineson, Angela Hodge

Article acceptance date: 26th October 2017.

The following Supporting Information is available for this article:

Table S1. Mean plant biomass parameters from AMF and non-AMF treatments in Experiment 1 ± standard error of the mean ($n = 12$). There were no significant differences between treatments for any of the parameters ($P > 0.05$ in each case) as determined using two-way ANOVAs.

	AMF	Non-AMF
Leaf DW (g)	1.22 ^a ± 0.05	1.16 ^a ± 0.07
Stalk DW (g)	2.08 ^b ± 0.08	1.97 ^b ± 0.11
Tassel DW (g)	0.03 ^c ± 0.01	0.02 ^c ± 0.003
Total shoot DW (g)	3.89 ^d ± 0.32	3.82 ^d ± 0.33
Root DW (g)	1.24 ^e ± 0.29	1.68 ^e ± 0.45
Total plant DW (g)	5.13 ^f ± 0.41	5.50 ^f ± 0.57
Root weight ratio	0.23 ^g ± 0.03	0.27 ^g ± 0.05

Mean values with the same letters are not significantly different at $P = 0.05$.