

Age-related response to an acute innate immune challenge in mice: proteomics reveals a telomere maintenance-related cost.

François Criscuolo^{1‡}, Gabriele Sorci², Margaux Behaim-Delarbre¹, Sandrine Zahn¹, Bruno Faivre² and Fabrice Bertile^{1‡}

¹Université de Strasbourg, CNRS, IPHC UMR 7178, 67000 Strasbourg, France.

²Biogéosciences, CNRS UMR 6282, Université Bourgogne Franche-Comté, Dijon, France.

[‡]Authors for correspondence (francois.criscuolo@iphc.cnrs.fr, fbertile@unistra.fr)

Electronic Supplementary Material 3 : Body mass and telomere length raw data

Table S3

Variables	Group size	Mean values	Confidence interval 95%
(a) Body mass loss (g)			
Old LPS	7	-2.6 ± 0.3	-3.3 ; -2.0
Old PBS	6	-2.1 ± 0.3	-2.8 ; -1.4
Young LPS	6	-2.8 ± 0.3	-3.5 ; -2.1
Young PBS	6	-1.0 ± 0.3	-1.7 ; -0.3
(b) Telomere length (T/S ratio)			
Old LPS	6	0.892 ± 0.104	0.672 ; 1.111
Old PBS	6	0.825 ± 0.127	0.556 ; 1.094
Young LPS	6	1.008 ± 0.114	0.767 ; 1.249
Young PBS	6	1.256 ± 0.115	1.015 ; 1.497

Descriptive statistics on body mass loss and final telomere length measured in splenocytes of old (12 months) and young (1 month) mice challenged with LPS or PBS.