

**SUPPLEMENTAL TABLE**

**Supplemental Table 1. Marginal structural model\* results for the association of incident atrial fibrillation with subsequent incident cardiovascular events in adults with chronic kidney disease.**

<i>Outcome</i>	<i>Estimated HR (95% CI)</i>	<i>p-value</i>
<b>Incident heart failure</b>		
<b>No Incident AF</b>	Ref	
<b>Incident AF</b>	4.80 ( 3.51 - 6.56 )	<.0001
<b>Incident MI</b>		
<b>No Incident AF</b>	Ref	
<b>Incident AF</b>	3.10 ( 2.11 - 4.54 )	<.0001
<b>Incident stroke</b>		
<b>No Incident AF</b>	Ref	
<b>Incident AF</b>	1.75 ( 0.99 - 3.09 )	0.02
<b>Death</b>		
<b>No Incident AF</b>	Ref	
<b>Incident AF</b>	3.69 ( 2.93 - 4.64 )	<.0001
<b>Composite of death and incident CVD</b>		
<b>No Incident AF</b>	Ref	
<b>Incident AF</b>	3.68 ( 2.91 - 4.66 )	<.0001

\*Adjusted for age, sex, race/ethnicity, clinical site, diabetes, hypertension, any history of cardiovascular disease, tobacco use, systolic blood pressure, BMI, eGFR, proteinuria, BMI, LDL cholesterol, HDL cholesterol, ACEi/ARB use,  $\beta$ -blocker use and diuretic use