

Supplementary table 2

Secondary outcome measures in males

	Donepezil means (SEM)	Placebo means (SEM)	Treatment effect*				<i>p</i>
			means	(95% confidence interval)	
Psychosis							
Cx in total PPQ	-0.13 (0.24)	0.65 (0.30)	-0.78	(-1.53 - -0.027)	0.042
Sleepiness							
Cx in Epworth Sleep Scale	0.59 (0.66)	2.67 (0.70)	-2.09	(-3.97 - -0.21)	0.029
Cognitive function							
Cx in scores of MMSE	0.03 (0.30)	-0.35 (0.31)	0.38	(-0.46 - 1.21)	0.384
Cx in WMS-1 (auditory memory)	14.7 (2.11)	8.85 (1.77)	5.85	(0.46 - 11.25)	0.034
Cx in WMS-2 (visual memory)	4.57 (1.04)	4.10 (1.08)	0.47	(-2.46 - 3.39)	0.755
Cx in WMS-4 (attention)	2.09 (1.37)	3.38 (1.52)	-1.28	(-5.28 - 2.72)	0.529
Cx in FAB	0.03 (0.34)	0.87 (0.35)	-0.85	(-1.81 - 0.11)	0.083

*Effect size was estimated using GEE (working matrix M-dependent)

Cx: changes, SEM: standard error of mean

Higher scores in PPQ, and Epworth Sleep Scale are worse.

Higher scores in MMSE, WMS, and FAB are better.