

In vitro Chondrocyte Responses in Mg-doped Wollastonite/Hydrogel Composite Scaffolds for Osteochondral Interface Regeneration

Xinning Yu^{1,2,4}, Tengfei Zhao^{1,2}, Yiyi Qi^{1,2}, Jianyang Luo^{1,2}, Jinghua Fang^{1,2,4}, Xianyan Yang³, Xiaonan Liu^{1,2}, Tengjing Xu^{1,2}, Quanming Yang^{1,2}, Zhongru Gou³, Xuesong Dai^{1,2*}

¹Department of Orthopaedic Surgery, Second Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou 310009, China.

²Orthopaedics Research Institute, Zhejiang University, Hangzhou, 310009, China.

³Bio-nanomaterials and Regenerative Medicine Research Division, Zhejiang-California International NanoSystems Institute, Zhejiang University, Hangzhou 310058, China.

⁴Department of Orthopaedic Surgery, Hangzhou Mingzhou Hospital, Hangzhou 311215, China.

Correspondence should be addressed to Xuesong Dai (E-mail: daixshz@zju.edu.cn)

Figure S1




Figure S2




Figure S3




Figure S4




Figure S5




Figure S6

