Supplementary Figure

Actein inhibits the proliferation and adhesion of human breast cancer cells and suppresses migration *in vivo*

Xiao-Xiao Wu^{1,#}, Grace Gar-Lee Yue^{2,3,#}, Jin-Run Dong^{4,#}, Christopher Wai-Kei Lam⁵, Chun-Kwok Wong^{1,2,3,6,*}, Ming-Hua Qiu^{4,*}, Clara Bik-San Lau^{2,3,*}

¹Department of Chemical Pathology; ²Institute of Chinese Medicine; ³State Key Laboratory of Phytochemistry and Plant Resources in West China, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong SAR, China; ⁴State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China; ⁵State Key Laboratory of Quality Research in Chinese Medicines, Macau Institute for Applied Research in Medicine and Health, Macau University of Science and Technology, Taipa, Macau; ⁶Li Dak Sum Yip Yio Chin R & D Centre for Chinese Medicine, The Chinese University of Hong Kong, Hong Kong SAR, China.

*Corresponding authors

Clara Bik-San LAU, Institute of Chinese Medicine, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong. Tel: +852 3943 6109, Fax: +852 2603 5248, E-mail: claralau@cuhk.edu.hk

Ming-Hua QIU, State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, China. Tel: +86 871 6522 3257, Fax: +86 871 6522 3255, E-mail: mhchiu@mail.kib.ac.cn

Chun-Kwok WONG, Department of Chemical Pathology, Prince of Wales Hospital, Shatin, New Territories, Hong Kong. Tel: +852 3505 2964, Fax: +852 2636 5090, E-mail: ck-wong@cuhk.edu.hk

These authors contributed equally to this work.




Figure S1: Survival rate of embryos after treated with or without actein for the following 5 days. Results were expressed as ratio of alive embryos per group in treated and vehicle-treated control of 4 independent experiments.