

Article: Energy crops affecting farmland birds in Central Europe: insights from a miscanthus-dominated landscape

Journal: *Biologia*

Authors: Jan M. Kaczmarek*, Tadeusz Mizera & Piotr Tryjanowski

*Corresponding author; Institute of Zoology, Poznań University of Life Sciences, Wojska Polskiego 71C, 60-625 Poznań, Poland; email: kaczmara@up.poznan.pl

Online Resource 1. Land cover in the study plots (i.e. buffers with a 150 m radius around observation points). Values for each visit were summed to provide a standardized estimate of land cover in each season. Differences in values between the seasons are a consequence of varying number of visits to study plots as well as agricultural activity.

COVER TYPE	EARLY SPRING		BREEDING SEASON		SUMMER		AUTUMN		WINTER		TOTAL	
	Area [ha]	%	Area [ha]	%	Area [ha]	%	Area [ha]	%	Area [ha]	%	Area [ha]	%
Cereals	39.89	23.48	137.11	26.89	45.37	12.45	37.12	18.48	59.31	23.41	318.80	21.27
Low marginal vegetation	15.43	9.08	45.70	8.97	32.73	8.98	17.56	8.75	21.79	8.60	133.21	8.89
Meadows	20.79	12.24	59.52	11.68	39.29	10.78	19.80	9.86	30.10	11.88	169.50	11.31
Miscanthus	53.06	31.23	153.21	30.05	100.93	27.68	51.89	25.83	74.54	29.42	433.63	28.94
Orchards	3.36	1.97	10.03	1.97	6.33	1.73	4.82	2.40	4.91	1.94	29.45	1.97
Other crops	9.77	5.75	52.70	10.34	24.94	6.84	17.58	8.76	18.60	7.34	123.59	8.25
Shrubs	1.25	0.73	3.70	0.72	2.52	0.69	1.32	0.66	1.59	0.63	10.38	0.69
Stubble/ploughed fields	17.24	10.14	20.29	3.98	93.90	25.76	39.84	19.84	28.61	11.29	199.88	13.34
Trees	9.16	5.39	27.55	5.40	18.58	5.09	10.91	5.43	13.90	5.49	80.10	5.35
SUM	169.93	100.00	509.80	100.00	364.59	100.00	200.83	100.00	253.35	100.00	1498.50	100.00