

Supplemental Table 1: Baseline demographics among mild ischemic stroke patients enrolled in study versus those excluded.

	Enrolled (n=229)	Excluded (n=16)	p valu e
Demographics			
Age (SD), y	64.9 (15.0)	62.5 (20.6)	0.47
Male, n (%)	126 (55.0%)	7 (43.8%)	0.27
Black, n (%)	68 (29.7%)	4 (25.0%)	0.47
Private Insurance, n (%)	84 (36.7%)	9 (56.3%)	0.10
mRS score at baseline, median (IQR)	0 (0-3)	0 (0-3)	0.38
Comorbid medical conditions, n (%)			
Diabetes mellitus	53 (23.1%)	2 (12.5%)	0.26
Hypercholesterolemia	162 (70.7%)	13 (82.3%)	0.28
Atrial fibrillation/Flutter	20 (8.7%)	2 (12.5%)	0.43
Cardiac disease	42 (18.3%)	3 (1.3%)	0.59
Prior stroke	40 (17.5%)	4 (25.0%)	0.32
Chronic kidney disease	15 (6.6%)	0 (0%)	0.35
Hypertension	172 (75.1%)	13 (81.2%)	0.42
Peripheral artery disease	14 (6.1%)	1 (6.3%)	0.65
Smoking within past 6 months	44 (19.2%)	3 (18.8%)	0.63 2
Clinical characteristics			
Initial NIHSS score, median (IQR)	1 (0-5)	1 (0-5)	0.18
TOAST subtype, n (%)			0.43
Cardioembolic	44 (19.2%)	1 (6.3%)	
Large artery atherosclerosis	41 (17.9%)	4 (25.0%)	

Small artery disease	37 (16.2%)	2 (12.5%)
Other	28 (12.2%)	4 (25.0%)
Cryptogenic	79 (34.5%)	5 (31.3%)

Abbreviations: SD - standard deviation, IQR - interquartile range, mRS - modified Rankin Score, NIHSS – National Institutes of Health Stroke Scale, TOAST - Trial of Org 10172 in Acute Stroke Treatment