

Cell, Volume 172

Supplemental Information

Western Diet Triggers NLRP3-Dependent

Innate Immune Reprogramming

Anette Christ, Patrick Günther, Mario A.R. Lauterbach, Peter Duewell, Debjani Biswas, Karin Pelka, Claus J. Scholz, Marije Oosting, Kristian Haendler, Kevin Baßler, Kathrin Klee, Jonas Schulte-Schrepping, Thomas Ulas, Simone J.C.F.M. Moorlag, Vinod Kumar, Min Hi Park, Leo A.B. Joosten, Laszlo A. Groh, Niels P. Riksen, Terje Espevik, Andreas Schlitzer, Yang Li, Michael L. Fitzgerald, Mihai G. Netea, Joachim L. Schultze, and Eicke Latz

Supplemental Tables

Table 1. Serum cytokine levels. Female *Ldlr^{-/-}* mice were fed either chow diet (CD), western diet (WD) for 4 weeks or WD for 4 weeks followed by CD for 4 weeks (WD>CD) and serum cytokines were measured. pg/ml = picogram per milliliter

Cytokine (pg/ml)	CD	WD	WD>CD
IL-1 α	12.15	107.45	54.45
IL-1 β	554.00	2288.98	472.31
IL-2	41.22	247.23	39.90
IL-3	13.36	102.23	13.62
IL-4	7.47	42.08	7.95
IL-5	60.22	156.47	30.42
IL-6	22.81	62.56	16.20
IL-9	318.81	802.64	302.35
IL-10	87.03	385.39	78.13
IL-12p40	264.02	211.49	253.39
IL-12p70	157.05	627.88	185.46
IL-13	368.36	1530.54	477.75
IL-17	52.85	199.10	69.98
G-CSF	159.48	272.95	130.05
GM-CSF	212.68	561.09	211.18
IFN γ	19.02	93.98	18.69
CXCL1	163.69	140.47	87.91
MCP-1	436.50	1222.79	338.51
MIP-1 α	18.58	37.06	17.17
MIP-1 β	58.84	209.35	51.37
CCL5	76.34	85.01	89.40
TNF- α	1965.22	3002.10	2207.60
IL-18	1498.37	3184.53	1618.29
VEGF	22.63	115.80	22.64