

## Brief Report

### **ARGONAUT-I: Activity of cefiderocol (S-649266), a siderophore cephalosporin, against Gram-negative bacteria including carbapenem resistant nonfermenters and Enterobacteriaceae with defined extended-spectrum $\beta$ -lactamases and carbapenemases**

Michael R. Jacobs\*, MD PhD<sup>1</sup>, Ayman M. Abdelhamed, MD PhD<sup>2</sup>, Caryn E. Good, MA/MPH<sup>3</sup>, Daniel D. Rhoads, MD<sup>2</sup>, Kristine M. Hujer, BS<sup>4,5</sup>, Andrea M. Hujer, BS<sup>4,5</sup>, T. Nicholas Domitrovic, MS<sup>4,5</sup>, Susan D. Rudin, BS<sup>4,5</sup>, Sandra S. Richter, MD<sup>6</sup>, David van Duin, MD/PhD<sup>7</sup>, Barry N. Kreiswirth, PhD<sup>8</sup>, Chris Greco, PhD<sup>9</sup>, Derrick E. Fouts, PhD<sup>9</sup>, Robert A. Bonomo, MD\*<sup>4,5,10</sup> on behalf of the Antibacterial Resistance Leadership Group (ARLG)

#### **Supplemental material**

Figure S1. Structure of cefiderocol (S-649266).

Table S1. Interpretative breakpoints used.



**Table S1. Interpretative breakpoints used.**

Agent	Range tested (mg/L)	Interpretative breakpoints (mg/L) <sup>a,b</sup>			
		Enterobacteriaceae	<i>Acinetobacter spp.</i>	<i>Pseudomonas aeruginosa</i>	<i>Stenotrophomonas maltophilia</i>
Cefiderocol	0.03-64	- <sup>c</sup>	-	-	-
Amikacin	4-64	≤16/32/>32	≤16/32/>32	≤16/32/>32	-
Ciprofloxacin	0.24-4	≤1/2/>2	≤1/2/>2	≤1/2/>2	-
Colistin	0.5-8	≤2/-/>2 <sup>d</sup>	≤2/-/>2	≤2/-/>2	≤2/-/>2 <sup>d</sup>
Tigecycline	0.25-4	≤2/4/>4 <sup>e</sup>	-	-	-
Aztreonam	0.5-32	≤4/8/>8	-	≤8/16/>16	-
Ceftolozane-tazobactam <sup>f</sup>	0.03-64	≤2/4/>4	-	≤4/8/>8	-
Cefepime	0.5-16	≤2/4-8 <sup>g</sup> />8	-	≤8/16/>16	-
Ceftazidime	0.03-64	≤4/8/>8	≤8/16/>16	≤8/16/>16	≤8/16/>16
Ceftazidime-avibactam <sup>f</sup>	0.03-64	≤8/-/>8	≤8/-/>8 <sup>d</sup>	≤8/-/>8	≤8/-/> 8 <sup>d</sup>
Meropenem	0.03-64	≤1/2/>2	≤2/4/>4	≤2/4/>4	-

<sup>a</sup> Breakpoints shown as susceptible/intermediate/resistant values unless otherwise indicated

<sup>b</sup> CLSI breakpoints (1) shown unless otherwise indicated

<sup>c</sup> -, no interpretative breakpoints available; provisional susceptible breakpoint of ≤4 mg/L used for cefiderocol

<sup>d</sup> Breakpoint for *P. aeruginosa* used

<sup>e</sup> FDA breakpoint

<sup>f</sup> Tazobactam and avibactam tested in fixed concentration of 4 mg/L with ceftolozane and ceftazidime, respectively

<sup>g</sup> Susceptible, dose-dependent

## **References**

1. Clinical and Laboratory Standards Institute. 2018. M100-S28. Performance Standards for Antimicrobial Susceptibility Testing, 28th Edition. Clinical and Laboratory Standards Institute, Wayne, PA.