

	REAL-WORLD (N=46)	PALOMA-3 (N=345)
Most common haematological AEs (all grades)	neutropenia (93%), leukopenia (89%), thrombocytopenia (48%), anaemia (46%)	neutropenia (81%), leukopenia (50%), thrombocytopenia (21%), anaemia (28%)
Most common non-haematological AEs (all grades)	fatigue (59%), decreased appetite (37%), infections (35%), nausea (35%), increased ASAT (28%), dizziness (26%)	infections (41%), fatigue (39%), nausea (32%), headache (23%), diarrhoea (21%), constipation (19%), alopecia (17%)
Grade 3-4 haematological AEs	neutropenia (63%), leukopenia (33%)	neutropenia (65%), leukopenia (28%)
Grade 3-4 non-haematological AEs (>2.5%)	pneumonia (n=3, 7%), pleural effusion (n=3, 7%), hypertension (n=2, 4%), ascites (n=2, 4%), increased ASAT (n=2, 4%)	no non-haematological AEs >2.5%
Serious AEs (all causalities)	occurred in 13% of patients (n=6)	occurred in 13% of patients (n=44)
Febrile neutropenia	did not occur	was uncommon (n=3, 0.9%)
<i>Note.</i> PALOMA-3 data were adapted from Cristofanilli et al. (2016) and Verma et al. (2016)		