

Table S1

Drivers	Strategy (n's)			Cell body location							Projections															
	I	II	III	SNC	RR	VTA	IF	CLi	DR/PG	CPr	CPi	CPc	CPt	ACB core	ACB msh	ACB lsh	OT	CEA c	CEA m	CEA I	BLA	IA	BST ov	LS	PFC	ENT
Dat-ires-Cre ¹	3			✓	✓	✓	✓	✓	✓	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
		2 (SNC/ VTA)		✓	■	✓	✓	■	■	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Aldh1a1-CreER ^{T2}		2		✓	✓	✓	✓	✓	✓	+	+	+	+	+	+	+	+	+	+	+	+	+	+	#	#	+
		6 (SNC)		✓	■	■	■	■	■	+	+	+	+/	-	+	+/	+/	-	-	-	-	-	-	+	-	-
		6 (VTA)		■	■	✓	✓	✓	■	-	-	-	-	-	+	+/	+/	-	-	-	-	-	-	+	-	-
Adcyap1-Cre ²		3		✓	■	✓	✓	✓	■	+	+	+	+/	-	+	+/	+/	-	-	-	-	-	-	+	-	-
		3 (Frepe)		✓	✓	✓	✓	✓	■	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#
Calb1-Cre		3 (SNC)		✓	■	✓	■	■	■	+	+	+	+	+/	-	-	-	+/	-	-	+/	+/	-	-	+/	+/
		3		✓	✓	✓	✓	✓	✓	✓	+	+	+	+	+/	+/	+/	+/	-	+	+/	+/	-	-	-	-
Cck-Cre ³		2 (Frepe & Ai65)		✓	■	✓	✓	✓	✓	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#
		2 (VTA)		■	■	✓	✓	■	■	+	+/	-	-	+	+	+	+	-	-	-	+	-	-	+	+	+
Ndnf-dgCre ⁴		3		✓	✓	✓	■	■	■	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#
		3		✓	✓	✓	■	■	■	+	+	+	+/	+	-	+/	+/	-	-	-	-	-	-	-	-	-
Sox6-FSF-Cre		5 (Frepe)		✓	✓	✓	■	■	■	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#
		2 (VTA/ SNC)		✓	■	✓	■	■	■	+	+	+/	-	+	-	+	+/	-	-	-	-	-	-	-	-	-
		2 (SNC)		✓	■	■	■	■	■	+	+	+	+/	+	-	+/	+/	-	-	-	-	-	-	-	-	-
Vglut2-Cre ⁵		3 (Frepe)		✓	✓	✓	✓	✓	✓	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#	#
		5 (VTA)		■	■	✓	✓	■	■	-	-	-	-	+/	+	+/	+	-	-	-	+	-	-	+	+	+/
		3 (SNC)		✓	■	■	■	■	■	-	+/	+/	+	-	-	-	-	+	-	-	-	-	-	-	-	-
Vip-Cre ³		3		■	■	■	■	■	■	✓	-	-	-	-	-	-	-	-	+/	+	-	-	+	-	-	-
		3		■	■	■	■	■	■	✓	-	-	-	-	-	-	-	-	+/	+	-	-	+	-	-	-

n = the number of mice used in independent experiment.

+ = dense projections; +/- = fibers present; - = fibers absent; # = not analyzed due to recombination outside midbrain; ## = not analyzed due to the lack of recombination .

References

1. Bäckman, C. M., Malik, N., Zhang, Y., Shan, L., Grinberg, A., Hoffer, B. J., et al. (2006). Characterization of a mouse strain expressing Cre recombinase from the 3' untranslated region of the dopamine transporter locus. *Genesis (New York, NY : 2000)*, 44(8), 383–390.
2. Krashes, M. J., Shah, B. P., Madara, J. C., Olson, D. P., Strochlic, D. E., Garfield, A. S., et al. (2014). An excitatory paraventricular nucleus to AgRP neuron circuit that drives hunger. *Nature*, 507(7491), 238–242.
3. Taniguchi, H., He, M., Wu, P., Kim, S., Paik, R., Sugino, K., et al. (2011). A resource of cre driver lines for genetic targeting of GABAergic neurons in cerebral cortex. *Neuron*, 71(6), 995–1013.
4. Tasic, B., Menon, V., Nguyen, T. N., Kim, T. K., Jarsky, T., Yao, Z., et al. (2016). Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nature Neuroscience*, 19(2), 335–346.
5. Vong, L., Ye, C., Yang, Z., Choi, B., Chua, S., & Lowell, B. B. (2011). Leptin action on GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. *Neuron*, 71(1), 142–154.

Table S2

DA neuron subtypes	Preferred Strategy	Mouse drivers
Aldh1a1+/Sox6+ (SNC)	Viral injection (CreON,FlpON) in SNC	Aldh1a1-CreER ^{T2} ; Th-2A-Flpo
Calb1+/Sox6+ (SNC)	Viral injection (CreON,FlpON) in SNC	Calb1-Cre; Th-2A-Flpo
Vglut2+ (SNC)	Viral injection (CreON,FlpON) in SNC	Vglut2-Cre; Th-2A-Flpo
Sox6+/Calb1+ (VTA)	Viral injection (CreON,FlpON) in VTA	Sox6-FSF-Cre; Th-2A-Flpo
Vglut2+/Otx2+ (VTA)	Viral injection (CreON,FlpON) in VTA	Vglut2-Cre; Th-2A-Flpo
Aldh1a1+/Otx2+ (VTA)	Viral injection (CreON,FlpON) in VTA	Aldh1a1-CreER ^{T2} ; Th-2A-Flpo
Vip+ (DR/PAG)	Transgenic reporter (e.g. Ai65)	Vip-Cre; Th-2A-Flpo