

Supplementary Table SI Bisphenol and phthalate urinary concentrations (n = 1233 women).

	Median (IQR) (ng/ml)	Percentage of values below the LOD
Total bisphenols ^a	9.22 (3.51, 20.28)	
Bisphenol A (BPA)	1.65 (0.72, 3.58)	21.7
Bisphenol S (BPS)	0.35 (0.17, 1.03)	31.5
Bisphenol F (BPF)	0.57 (0.30, 1.32)	60.0
Phthalic acid (PA)	56.99 (30.62, 124.05)	0.3
Low molecular weight (LMW) metabolites ^a	1087.42 (423.91, 2925.28)	
Monomethylphthalate (mMP)	5.43 (2.75, 9.75)	0.2
Monoethylphthalate (mEP)	137.85 (41.59, 484.50)	0.1
Mono-isobutylphthalate (mIBP)	20.69 (9.44, 45.33)	0.2
Mono-n-butylphthalate (mBP)	15.58 (6.89, 31.00)	0.7
High molecular weight (HMW) metabolites ^a	219.66 (110.84, 405.51)	
<i>Di-2-ethylhexylphthalate (DEHP) metabolites^a</i>	171.33 (87.61, 323.30)	
Mono-(2-ethyl-5-carboxypentyl)phthalate (mECP)	15.96 (8.08, 31.60)	0.2
Mono-(2-ethyl-5-hydroxyhexyl)phthalate (mEHHP)	11.76 (5.67, 22.92)	0.2
Mono-(2-ethyl-5-oxohexyl)phthalate (mEOHP)	7.66 (3.46, 15.48)	–
Mono-[(2-carboxymethyl)hexyl]phthalate (mCMHP)	13.90 (7.55, 26.26)	0.1
<i>Di-n-octylphthalate (DNOP)^a</i>	5.72 (3.10, 11.03)	
Mono(3-carboxypropyl)phthalate (mCPP)	1.44 (0.78, 2.77)	–
<i>Other high molecular weight metabolites</i>		
Monobenzylphthalate (mBzP)	6.55 (3.06, 12.81)	8.3
Mono-hexylphthalate (mHxP)	0.33 (0.16, 0.63)	24.0
Mono-2-heptylphthalate (mHpP)	1.09 (0.60, 2.29)	36.7

LOD = limit of detection.

^aGroups are molar concentrations in nmol/L with non-detectable levels of separate metabolites imputed as LOD/sqrt(2). Separate metabolites are included only if <80% of values was below the LOD.